

Computer Algebra Independent Integration Tests

Summer 2023 edition

6-Hyperbolic-functions/6.4-Hyperbolic-cotangent/176-6.4.7-d-
hyper- \hat{m} -a+b-c-coth- \hat{n} - \hat{p}

Nasser M. Abbasi

September 5, 2023

Compiled on September 5, 2023 at 3:09pm

Contents

1	Introduction	3
2	detailed summary tables of results	21
3	Listing of integrals	39
4	Appendix	363

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	8
1.4	Performance based on number of rules Rubi used	10
1.5	Performance based on number of steps Rubi used	11
1.6	Solved integrals histogram based on leaf size of result	12
1.7	Solved integrals histogram based on CPU time used	13
1.8	Leaf size vs. CPU time used	14
1.9	list of integrals with no known antiderivative	15
1.10	List of integrals solved by CAS but has no known antiderivative	15
1.11	list of integrals solved by CAS but failed verification	15
1.12	Timing	16
1.13	Verification	16
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [53]. This is test number [176].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (53)	0.00 (0)
Mathematica	100.00 (53)	0.00 (0)
Fricas	100.00 (53)	0.00 (0)
Maple	81.13 (43)	18.87 (10)
Giac	66.04 (35)	33.96 (18)
Mupad	60.38 (32)	39.62 (21)
Maxima	30.19 (16)	69.81 (37)
Sympy	13.21 (7)	86.79 (46)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	66.038	15.094	18.868	0.000
Maple	33.962	45.283	1.887	18.868
Giac	7.547	47.170	11.321	33.962
Maxima	7.547	16.981	5.660	69.811
Fricas	3.774	90.566	5.660	0.000
Sympy	1.887	11.321	0.000	86.792
Mupad	0.000	60.377	0.000	39.623

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Maple	10	100.00	0.00	0.00
Giac	18	22.22	0.00	77.78
Mupad	21	0.00	100.00	0.00
Maxima	37	100.00	0.00	0.00
Sympy	46	93.48	6.52	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Rubi	0.07
Maple	0.19
Maxima	0.29
Fricas	0.40
Giac	0.47
Mathematica	0.80
Mupad	2.44
Sympy	3.84

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Rubi	63.92	1.00	56.00	1.00
Mathematica	110.08	1.72	65.00	1.05
Maple	186.91	3.01	137.00	2.74
Maxima	209.25	2.63	64.50	2.18
Mupad	249.41	2.21	44.00	1.00
Sympy	292.71	3.96	253.00	4.35
Giac	316.83	4.81	241.00	4.06
Fricas	3314.85	40.97	2111.00	37.57

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {37, 42, 44, 47, 49}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
```

```
x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in *Rubi Table file*

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2023
Design-vide

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	22
2.2	Detailed conclusion table per each integral for all CAS systems	25
2.3	Detailed conclusion table specific for Rubi results	36

2.1 List of integrals sorted by grade for each CAS

Rubi	22
Mma	22
Maple	23
Fricas	23
Maxima	23
Giac	24
Mupad	24
Sympy	24

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 13, 14, 15, 16, 18, 19, 20, 22, 23, 24, 25, 26, 28, 29, 31, 32, 34, 36, 38, 48, 49, 50, 51, 52, 53 }

B grade { 10, 12, 17, 27, 30, 33, 35, 39 }

C grade { 21, 37, 40, 41, 42, 43, 44, 45, 46, 47 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 50, 51, 52 }

B grade { 16, 17, 18, 19, 22, 23, 24, 25, 28, 29, 30, 31, 32, 33, 34, 35, 38, 39, 40, 43, 44, 45, 48, 49 }

C grade { 53 }

F normal fail { 20, 21, 26, 27, 36, 37, 41, 42, 46, 47 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 11, 15 }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 30, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 50, 51, 52, 53 }

C grade { 29, 31, 49 }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maxima

A grade { 5, 11, 13, 15 }

B grade { 1, 2, 3, 4, 6, 7, 8, 9, 50 }

C grade { 10, 12, 14 }

F normal fail { 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 51, 52, 53 }

F(-1) timeout fail { }

F(-2) exception fail { }

Giac

A grade { 5, 11, 15, 50 }

B grade { 1, 2, 3, 4, 6, 7, 8, 9, 13, 19, 20, 26, 28, 30, 34, 35, 38, 39, 40, 42, 43, 44, 45, 47, 48 }

C grade { 10, 12, 14, 29, 31, 49 }

F normal fail { 27, 51, 52, 53 }

F(-1) timedout fail { }

F(-2) exception fail { 16, 17, 18, 21, 22, 23, 24, 25, 32, 33, 36, 37, 41, 46 }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 18, 22, 24, 28, 29, 30, 32, 34, 35, 38, 40, 43, 45, 48, 49, 50 }

C grade { }

F normal fail { }

F(-1) timedout fail { 17, 19, 20, 21, 23, 25, 26, 27, 31, 33, 36, 37, 39, 41, 42, 44, 46, 47, 51, 52, 53 }

F(-2) exception fail { }

Sympy

A grade { 9 }

B grade { 1, 2, 3, 4, 5, 50 }

C grade { }

F normal fail { 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 51, 52, 53 }

F(-1) timedout fail { 6, 7, 8 }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	160	160	231	153	624	2111	639	721	158
N.S.	1	1.00	1.44	0.96	3.90	13.19	3.99	4.51	0.99
time (sec)	N/A	0.066	3.219	0.343	0.216	0.270	14.402	0.330	2.114

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	110	127	112	410	1164	478	447	111
N.S.	1	1.00	1.15	1.02	3.73	10.58	4.35	4.06	1.01
time (sec)	N/A	0.050	2.254	0.245	0.206	0.254	5.832	0.316	2.045

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	74	100	75	239	557	332	241	72
N.S.	1	1.00	1.35	1.01	3.23	7.53	4.49	3.26	0.97
time (sec)	N/A	0.036	1.508	0.177	0.203	0.267	2.329	0.291	2.006

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	65	46	114	197	211	103	41
N.S.	1	1.00	1.51	1.07	2.65	4.58	4.91	2.40	0.95
time (sec)	N/A	0.025	0.494	0.119	0.194	0.262	1.086	0.277	0.125

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	47	71	56	488	253	65	37
N.S.	1	1.00	1.02	1.54	1.22	10.61	5.50	1.41	0.80
time (sec)	N/A	0.056	0.080	0.130	0.291	0.280	2.489	0.276	0.122

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	89	97	103	207	1952	0	198	110
N.S.	1	1.00	1.09	1.16	2.33	21.93	0.00	2.22	1.24
time (sec)	N/A	0.074	0.626	0.211	0.320	0.327	0.000	0.293	2.313

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	142	142	147	156	509	7508	0	409	2719
N.S.	1	1.00	1.04	1.10	3.58	52.87	0.00	2.88	19.15
time (sec)	N/A	0.118	0.349	0.306	0.386	0.382	0.000	0.325	2.712

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	201	201	203	219	925	20031	0	750	3685
N.S.	1	1.00	1.01	1.09	4.60	99.66	0.00	3.73	18.33
time (sec)	N/A	0.200	0.598	0.485	0.506	0.521	0.000	0.357	2.810

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	27	38	70	34	38	15
N.S.	1	1.00	1.00	1.42	2.00	3.68	1.79	2.00	0.79
time (sec)	N/A	0.038	0.115	0.088	0.283	0.265	0.274	0.268	0.114

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	C	B	F	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	30	4	19	45	0	26	3
N.S.	1	1.00	10.00	1.33	6.33	15.00	0.00	8.67	1.00
time (sec)	N/A	0.014	0.053	0.188	0.277	0.253	0.000	0.284	1.905

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	28	15	17	17	0	23	14
N.S.	1	1.00	2.00	1.07	1.21	1.21	0.00	1.64	1.00
time (sec)	N/A	0.017	0.048	0.109	0.279	0.255	0.000	0.291	2.089

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	C	B	F	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	51	21	49	279	0	60	20
N.S.	1	1.00	2.12	0.88	2.04	11.62	0.00	2.50	0.83
time (sec)	N/A	0.016	0.102	0.095	0.302	0.282	0.000	0.274	1.946

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	49	28	46	211	0	52	27
N.S.	1	1.00	1.58	0.90	1.48	6.81	0.00	1.68	0.87
time (sec)	N/A	0.017	0.087	0.143	0.301	0.259	0.000	0.268	1.917

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	B	F	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	14	11	40	0	24	18
N.S.	1	1.00	1.00	1.08	0.85	3.08	0.00	1.85	1.38
time (sec)	N/A	0.015	0.042	0.127	0.280	0.261	0.000	0.271	2.092

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	11	12	11	2	0	18	15
N.S.	1	1.00	1.00	1.09	1.00	0.18	0.00	1.64	1.36
time (sec)	N/A	0.016	0.040	0.059	0.276	0.255	0.000	0.283	1.969

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	63	60	253	0	2367	0	0	66
N.S.	1	1.00	0.95	4.02	0.00	37.57	0.00	0.00	1.05
time (sec)	N/A	0.098	0.200	0.211	0.000	0.363	0.000	0.000	3.902

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	85	191	276	0	4877	0	0	0
N.S.	1	1.00	2.25	3.25	0.00	57.38	0.00	0.00	0.00
time (sec)	N/A	0.092	0.836	0.140	0.000	0.434	0.000	0.000	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	44	44	238	0	1551	0	0	51
N.S.	1	1.00	1.00	5.41	0.00	35.25	0.00	0.00	1.16
time (sec)	N/A	0.056	0.035	0.094	0.000	0.313	0.000	0.000	2.335

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	60	60	82	238	0	3455	0	262	0
N.S.	1	1.00	1.37	3.97	0.00	57.58	0.00	4.37	0.00
time (sec)	N/A	0.034	0.329	0.139	0.000	0.369	0.000	0.477	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	B	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	56	56	56	0	0	3479	0	259	0
N.S.	1	1.00	1.00	0.00	0.00	62.12	0.00	4.62	0.00
time (sec)	N/A	0.078	0.038	0.000	0.000	0.380	0.000	0.512	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	48	48	42	0	0	1539	0	0	0
N.S.	1	1.00	0.88	0.00	0.00	32.06	0.00	0.00	0.00
time (sec)	N/A	0.063	0.101	0.000	0.000	0.324	0.000	0.000	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	82	82	86	488	0	4940	0	0	112
N.S.	1	1.00	1.05	5.95	0.00	60.24	0.00	0.00	1.37
time (sec)	N/A	0.104	0.437	0.093	0.000	0.470	0.000	0.000	9.988

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	123	123	219	529	0	10286	0	0	0
N.S.	1	1.00	1.78	4.30	0.00	83.63	0.00	0.00	0.00
time (sec)	N/A	0.169	0.951	0.098	0.000	0.597	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	63	59	473	0	2362	0	0	64
N.S.	1	1.00	0.94	7.51	0.00	37.49	0.00	0.00	1.02
time (sec)	N/A	0.071	0.164	0.074	0.000	0.373	0.000	0.000	4.154

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	111	473	0	5037	0	0	0
N.S.	1	1.00	1.26	5.38	0.00	57.24	0.00	0.00	0.00
time (sec)	N/A	0.060	0.578	0.125	0.000	0.429	0.000	0.000	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	B	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	71	71	71	0	0	3949	0	470	0
N.S.	1	1.00	1.00	0.00	0.00	55.62	0.00	6.62	0.00
time (sec)	N/A	0.101	0.076	0.000	0.000	0.426	0.000	3.809	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	77	77	180	0	0	4025	0	0	0
N.S.	1	1.00	2.34	0.00	0.00	52.27	0.00	0.00	0.00
time (sec)	N/A	0.088	0.468	0.000	0.000	0.417	0.000	0.000	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	60	97	0	683	0	119	68
N.S.	1	1.00	1.94	3.13	0.00	22.03	0.00	3.84	2.19
time (sec)	N/A	0.020	0.120	0.229	0.000	0.261	0.000	0.282	2.038

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	45	45	62	142	0	226	0	124	43
N.S.	1	1.00	1.38	3.16	0.00	5.02	0.00	2.76	0.96
time (sec)	N/A	0.024	0.082	0.203	0.000	0.255	0.000	0.302	2.014

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	50	50	116	158	0	1043	0	265	78
N.S.	1	1.00	2.32	3.16	0.00	20.86	0.00	5.30	1.56
time (sec)	N/A	0.028	0.415	0.103	0.000	0.276	0.000	0.287	2.120

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	67	118	211	0	361	0	285	0
N.S.	1	1.00	1.76	3.15	0.00	5.39	0.00	4.25	0.00
time (sec)	N/A	0.036	0.264	0.107	0.000	0.262	0.000	0.309	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	47	129	0	1576	0	0	39
N.S.	1	1.00	1.00	2.74	0.00	33.53	0.00	0.00	0.83
time (sec)	N/A	0.082	0.099	0.117	0.000	0.315	0.000	0.000	2.388

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	60	60	134	137	0	3513	0	0	0
N.S.	1	1.00	2.23	2.28	0.00	58.55	0.00	0.00	0.00
time (sec)	N/A	0.071	0.238	0.118	0.000	0.394	0.000	0.000	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	29	114	0	1298	0	196	23
N.S.	1	1.00	1.00	3.93	0.00	44.76	0.00	6.76	0.79
time (sec)	N/A	0.050	0.021	0.155	0.000	0.296	0.000	0.419	2.306

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	77	114	0	1357	0	196	25
N.S.	1	1.00	2.48	3.68	0.00	43.77	0.00	6.32	0.81
time (sec)	N/A	0.021	0.137	0.137	0.000	0.298	0.000	0.388	2.223

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	56	56	56	0	0	3397	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	60.66	0.00	0.00	0.00
time (sec)	N/A	0.077	0.045	0.000	0.000	0.386	0.000	0.000	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	51	51	127	0	0	1621	0	0	0
N.S.	1	1.00	2.49	0.00	0.00	31.78	0.00	0.00	0.00
time (sec)	N/A	0.072	0.774	0.000	0.000	0.325	0.000	0.000	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	52	52	52	287	0	2541	0	359	45
N.S.	1	1.00	1.00	5.52	0.00	48.87	0.00	6.90	0.87
time (sec)	N/A	0.090	0.164	0.097	0.000	0.358	0.000	0.459	2.769

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	53	109	289	0	2279	0	363	0
N.S.	1	1.00	2.06	5.45	0.00	43.00	0.00	6.85	0.00
time (sec)	N/A	0.077	1.107	0.100	0.000	0.351	0.000	0.459	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	49	49	41	273	0	2299	0	364	41
N.S.	1	1.00	0.84	5.57	0.00	46.92	0.00	7.43	0.84
time (sec)	N/A	0.066	0.047	0.075	0.000	0.355	0.000	0.471	2.613

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	78	78	70	0	0	6991	0	0	0
N.S.	1	1.00	0.90	0.00	0.00	89.63	0.00	0.00	0.00
time (sec)	N/A	0.108	0.063	0.000	0.000	0.545	0.000	0.000	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	F	B	F	B	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	85	85	260	0	0	3931	0	540	0
N.S.	1	1.00	3.06	0.00	0.00	46.25	0.00	6.35	0.00
time (sec)	N/A	0.114	7.926	0.000	0.000	0.506	0.000	0.599	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	74	63	435	0	6560	0	951	82
N.S.	1	1.00	0.85	5.88	0.00	88.65	0.00	12.85	1.11
time (sec)	N/A	0.108	0.119	0.105	0.000	0.689	0.000	0.596	4.250

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	B	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	215	454	0	6591	0	952	0
N.S.	1	1.00	2.44	5.16	0.00	74.90	0.00	10.82	0.00
time (sec)	N/A	0.092	7.067	0.104	0.000	0.682	0.000	0.551	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	43	420	0	5736	0	909	76
N.S.	1	1.00	0.61	6.00	0.00	81.94	0.00	12.99	1.09
time (sec)	N/A	0.077	0.048	0.081	0.000	0.630	0.000	0.575	4.332

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	108	108	73	0	0	19199	0	0	0
N.S.	1	1.00	0.68	0.00	0.00	177.77	0.00	0.00	0.00
time (sec)	N/A	0.151	0.072	0.000	0.000	1.242	0.000	0.000	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	F	B	F	B	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	131	131	1350	0	0	10729	0	1133	0
N.S.	1	1.00	10.31	0.00	0.00	81.90	0.00	8.65	0.00
time (sec)	N/A	0.168	8.544	0.000	0.000	1.296	0.000	0.769	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	62	0	547	0	69	63
N.S.	1	1.00	1.00	2.48	0.00	21.88	0.00	2.76	2.52
time (sec)	N/A	0.013	0.071	0.200	0.000	0.253	0.000	0.265	2.096

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	C	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	49	66	0	175	0	73	22
N.S.	1	1.00	1.81	2.44	0.00	6.48	0.00	2.70	0.81
time (sec)	N/A	0.014	0.079	0.198	0.000	0.263	0.000	0.267	2.189

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	40	41	73	95	102	25	38
N.S.	1	1.00	1.05	1.08	1.92	2.50	2.68	0.66	1.00
time (sec)	N/A	0.046	0.093	0.099	0.289	0.273	0.500	0.263	0.106

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	89	86	116	0	5172	0	0	0
N.S.	1	1.00	0.97	1.30	0.00	58.11	0.00	0.00	0.00
time (sec)	N/A	0.102	0.170	0.815	0.000	0.402	0.000	0.000	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	40	40	40	37	0	1290	0	0	0
N.S.	1	1.00	1.00	0.92	0.00	32.25	0.00	0.00	0.00
time (sec)	N/A	0.056	0.021	0.577	0.000	0.406	0.000	0.000	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	74	73	431	0	3938	0	0	0
N.S.	1	1.00	0.99	5.82	0.00	53.22	0.00	0.00	0.00
time (sec)	N/A	0.089	0.528	0.670	0.000	0.495	0.000	0.000	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [50] had the largest ratio of [.6250000000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	4	3	1.00	14	0.214
2	A	4	3	1.00	14	0.214
3	A	4	3	1.00	14	0.214
4	A	4	3	1.00	14	0.214
5	A	3	3	1.00	14	0.214
6	A	5	5	1.00	14	0.357
7	A	6	6	1.00	14	0.429
8	A	7	6	1.00	14	0.429
9	A	3	3	1.00	10	0.300
10	A	3	3	1.00	12	0.250
11	A	4	4	1.00	10	0.400
12	A	4	4	1.00	12	0.333
13	A	5	5	1.00	10	0.500
14	A	3	3	1.00	12	0.250
15	A	3	3	1.00	10	0.300
16	A	6	6	1.00	17	0.353
17	A	7	6	1.00	17	0.353
18	A	5	5	1.00	15	0.333
19	A	6	5	1.00	12	0.417
20	A	7	5	1.00	15	0.333
21	A	5	5	1.00	17	0.294
22	A	7	6	1.00	17	0.353
23	A	8	7	1.00	17	0.412
24	A	6	5	1.00	15	0.333

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
25	A	7	6	1.00	12	0.500
26	A	8	6	1.00	15	0.400
27	A	7	6	1.00	17	0.353
28	A	5	5	1.00	10	0.500
29	A	6	5	1.00	12	0.417
30	A	6	6	1.00	10	0.600
31	A	7	6	1.00	12	0.500
32	A	5	5	1.00	17	0.294
33	A	6	5	1.00	17	0.294
34	A	4	4	1.00	15	0.267
35	A	3	3	1.00	12	0.250
36	A	7	5	1.00	15	0.333
37	A	5	5	1.00	17	0.294
38	A	5	5	1.00	17	0.294
39	A	4	4	1.00	17	0.235
40	A	5	5	1.00	15	0.333
41	A	8	6	1.00	15	0.400
42	A	6	6	1.00	17	0.353
43	A	6	6	1.00	17	0.353
44	A	6	6	1.00	17	0.353
45	A	6	5	1.00	15	0.333
46	A	9	7	1.00	15	0.467
47	A	7	7	1.00	17	0.412
48	A	3	3	1.00	10	0.300
49	A	3	3	1.00	12	0.250
50	A	6	5	1.00	8	0.625
51	A	8	7	1.00	15	0.467
52	A	4	4	1.00	15	0.267
53	A	6	6	1.00	15	0.400

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int (a + b \coth^2(c + dx))^5 dx \dots\dots\dots$	42
3.2	$\int (a + b \coth^2(c + dx))^4 dx \dots\dots\dots$	50
3.3	$\int (a + b \coth^2(c + dx))^3 dx \dots\dots\dots$	57
3.4	$\int (a + b \coth^2(c + dx))^2 dx \dots\dots\dots$	63
3.5	$\int \frac{1}{a+b \coth^2(c+dx)} dx \dots\dots\dots$	68
3.6	$\int \frac{1}{(a+b \coth^2(c+dx))^2} dx \dots\dots\dots$	73
3.7	$\int \frac{1}{(a+b \coth^2(c+dx))^3} dx \dots\dots\dots$	79
3.8	$\int \frac{1}{(a+b \coth^2(c+dx))^4} dx \dots\dots\dots$	87
3.9	$\int \frac{1}{1-2 \coth^2(x)} dx \dots\dots\dots$	96
3.10	$\int \sqrt{1 - \coth^2(x)} dx \dots\dots\dots$	100
3.11	$\int \sqrt{-1 + \coth^2(x)} dx \dots\dots\dots$	104
3.12	$\int (1 - \coth^2(x))^{3/2} dx \dots\dots\dots$	108
3.13	$\int (-1 + \coth^2(x))^{3/2} dx \dots\dots\dots$	113
3.14	$\int \frac{1}{\sqrt{1-\coth^2(x)}} dx \dots\dots\dots$	118
3.15	$\int \frac{1}{\sqrt{-1+\coth^2(x)}} dx \dots\dots\dots$	122
3.16	$\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx \dots\dots\dots$	126
3.17	$\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx \dots\dots\dots$	133
3.18	$\int \coth(x) \sqrt{a + b \coth^2(x)} dx \dots\dots\dots$	141
3.19	$\int \sqrt{a + b \coth^2(x)} dx \dots\dots\dots$	147
3.20	$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx \dots\dots\dots$	154

3.21	$\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx$	161
3.22	$\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx$	166
3.23	$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx$	174
3.24	$\int \coth(x) (a + b \coth^2(x))^{3/2} dx$	180
3.25	$\int (a + b \coth^2(x))^{3/2} dx$	187
3.26	$\int (a + b \coth^2(x))^{3/2} \tanh(x) dx$	192
3.27	$\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx$	200
3.28	$\int \sqrt{1 + \coth^2(x)} dx$	207
3.29	$\int \sqrt{-1 - \coth^2(x)} dx$	212
3.30	$\int (1 + \coth^2(x))^{3/2} dx$	217
3.31	$\int (-1 - \coth^2(x))^{3/2} dx$	223
3.32	$\int \frac{\coth^3(x)}{\sqrt{a+b \coth^2(x)}} dx$	229
3.33	$\int \frac{\coth^2(x)}{\sqrt{a+b \coth^2(x)}} dx$	235
3.34	$\int \frac{\coth(x)}{\sqrt{a+b \coth^2(x)}} dx$	242
3.35	$\int \frac{1}{\sqrt{a+b \coth^2(x)}} dx$	247
3.36	$\int \frac{\tanh(x)}{\sqrt{a+b \coth^2(x)}} dx$	252
3.37	$\int \frac{\tanh^2(x)}{\sqrt{a+b \coth^2(x)}} dx$	258
3.38	$\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{3/2}} dx$	264
3.39	$\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{3/2}} dx$	271
3.40	$\int \frac{\coth(x)}{(a+b \coth^2(x))^{3/2}} dx$	278
3.41	$\int \frac{\tanh(x)}{(a+b \coth^2(x))^{3/2}} dx$	285
3.42	$\int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{3/2}} dx$	290
3.43	$\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{5/2}} dx$	299
3.44	$\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{5/2}} dx$	305
3.45	$\int \frac{\coth(x)}{(a+b \coth^2(x))^{5/2}} dx$	311
3.46	$\int \frac{\tanh(x)}{(a+b \coth^2(x))^{5/2}} dx$	317
3.47	$\int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{5/2}} dx$	322
3.48	$\int \frac{1}{\sqrt{1+\coth^2(x)}} dx$	330
3.49	$\int \frac{1}{\sqrt{-1-\coth^2(x)}} dx$	335

3.50	$\int \frac{1}{1+\coth^3(x)} dx$	340
3.51	$\int \coth(x) \sqrt{a + b \coth^4(x)} dx$	345
3.52	$\int \frac{\coth(x)}{\sqrt{a+b \coth^4(x)}} dx$	351
3.53	$\int \frac{\coth(x)}{(a+b \coth^4(x))^{3/2}} dx$	356

3.1 $\int (a + b \coth^2(c + dx))^5 dx$

Optimal result	42
Rubi [A] (verified)	42
Mathematica [A] (verified)	44
Maple [A] (verified)	44
Fricas [B] (verification not implemented)	45
Sympy [B] (verification not implemented)	46
Maxima [B] (verification not implemented)	47
Giac [B] (verification not implemented)	48
Mupad [B] (verification not implemented)	49

Optimal result

Integrand size = 14, antiderivative size = 160

$$\int (a + b \coth^2(c + dx))^5 dx = (a + b)^5 x - \frac{b(5a^4 + 10a^3b + 10a^2b^2 + 5ab^3 + b^4) \coth(c + dx)}{d} - \frac{b^2(10a^3 + 10a^2b + 5ab^2 + b^3) \coth^3(c + dx)}{3d} - \frac{b^3(10a^2 + 5ab + b^2) \coth^5(c + dx)}{5d} - \frac{b^4(5a + b) \coth^7(c + dx)}{7d} - \frac{b^5 \coth^9(c + dx)}{9d}$$

[Out] (a+b)^5*x-b*(5*a^4+10*a^3*b+10*a^2*b^2+5*a*b^3+b^4)*coth(d*x+c)/d-1/3*b^2*(10*a^3+10*a^2*b+5*a*b^2+b^3)*coth(d*x+c)^3/d-1/5*b^3*(10*a^2+5*a*b+b^2)*coth(d*x+c)^5/d-1/7*b^4*(5*a+b)*coth(d*x+c)^7/d-1/9*b^5*coth(d*x+c)^9/d

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3742, 398, 212}

$$\int (a + b \coth^2(c + dx))^5 dx = -\frac{b^3(10a^2 + 5ab + b^2) \coth^5(c + dx)}{5d} - \frac{b^2(10a^3 + 10a^2b + 5ab^2 + b^3) \coth^3(c + dx)}{3d} - \frac{b(5a^4 + 10a^3b + 10a^2b^2 + 5ab^3 + b^4) \coth(c + dx)}{d} - \frac{b^4(5a + b) \coth^7(c + dx)}{7d} + x(a + b)^5 - \frac{b^5 \coth^9(c + dx)}{9d}$$

[In] Int[(a + b*Coth[c + d*x]^2)^5, x]

[Out] (a + b)^5*x - (b*(5*a^4 + 10*a^3*b + 10*a^2*b^2 + 5*a*b^3 + b^4)*Coth[c + d*x])/d - (b^2*(10*a^3 + 10*a^2*b + 5*a*b^2 + b^3)*Coth[c + d*x]^3)/(3*d) - (b^3*(10*a^2 + 5*a*b + b^2)*Coth[c + d*x]^5)/(5*d) - (b^4*(5*a + b)*Coth[c + d*x]^7)/(7*d) - (b^5*Coth[c + d*x]^9)/(9*d)

Rule 212

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 398

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rule 3742

Int[((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int \frac{(a+bx^2)^5}{1-x^2} dx, x, \coth(c+dx)\right)}{d} \\
 &= \frac{\text{Subst}\left(\int \left(-b(5a^4 + 10a^3b + 10a^2b^2 + 5ab^3 + b^4) - b^2(10a^3 + 10a^2b + 5ab^2 + b^3)x^2 - b^3(10a^2 + 5ab + b^2)x^4 - b^4(5a + b)x^6 - b^5x^8\right) dx, x, \coth(c+dx)\right)}{d} \\
 &= -\frac{b(5a^4 + 10a^3b + 10a^2b^2 + 5ab^3 + b^4) \coth(c+dx)}{d} \\
 &\quad - \frac{b^2(10a^3 + 10a^2b + 5ab^2 + b^3) \coth^3(c+dx)}{3d} \\
 &\quad - \frac{b^3(10a^2 + 5ab + b^2) \coth^5(c+dx)}{5d} - \frac{b^4(5a + b) \coth^7(c+dx)}{7d} \\
 &\quad - \frac{b^5 \coth^9(c+dx)}{9d} + \frac{(a+b)^5 \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \coth(c+dx)\right)}{d}
 \end{aligned}$$

$$= (a+b)^5 x - \frac{b(5a^4 + 10a^3b + 10a^2b^2 + 5ab^3 + b^4) \coth(c+dx)}{d} - \frac{b^2(10a^3 + 10a^2b + 5ab^2 + b^3) \coth^3(c+dx)}{3d} - \frac{b^3(10a^2 + 5ab + b^2) \coth^5(c+dx)}{5d} - \frac{b^4(5a+b) \coth^7(c+dx)}{7d} - \frac{b^5 \coth^9(c+dx)}{9d}$$

Mathematica [A] (verified)

Time = 3.22 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.44

$$\int (a + b \coth^2(c + dx))^5 dx = \frac{b^5 \coth^9(c + dx) \left(35 + 45 \tanh^2(c + dx) + 63 \tanh^4(c + dx) + 105 \tanh^6(c + dx) + 315 \tanh^8(c + dx) + \dots \right)}{\dots}$$

[In] Integrate[(a + b*Coth[c + d*x]^2)^5,x]

[Out] -1/315*(b^5*Coth[c + d*x]^9*(35 + 45*Tanh[c + d*x]^2 + 63*Tanh[c + d*x]^4 + 105*Tanh[c + d*x]^6 + 315*Tanh[c + d*x]^8 + (1575*a^4*Tanh[c + d*x]^8)/b^4 - (315*(a + b)^5*ArcTanh[Sqrt[Tanh[c + d*x]^2]]*Tanh[c + d*x]^10)/(b^5*Sqrt[Tanh[c + d*x]^2]) + (1050*a^3*Tanh[c + d*x]^6*(1 + 3*Tanh[c + d*x]^2))/b^3 + (210*a^2*Tanh[c + d*x]^4*(3 + 5*Tanh[c + d*x]^2 + 15*Tanh[c + d*x]^4))/b^2 + (15*a*Tanh[c + d*x]^2*(15 + 21*Tanh[c + d*x]^2 + 35*Tanh[c + d*x]^4 + 105*Tanh[c + d*x]^6))/b)/d

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.96

method	result
parallelrisc	$\frac{-35b^5 \coth(dx+c)^9 + (-225ab^4 - 45b^5) \coth(dx+c)^7 + (-630a^2b^3 - 315ab^4 - 63b^5) \coth(dx+c)^5 + (-1050a^3b^2 - 1050a^2b^3 - 1050ab^4 - 1050b^5) \coth(dx+c)^3 + (-315a^4b - 315a^3b^2 - 315a^2b^3 - 315ab^4 - 315b^5) \coth(dx+c) + 315a^5}{315d}$
derivativedivides	$\frac{-5a^4b \coth(dx+c) - 10a^3b^2 \coth(dx+c) - 10a^2b^3 \coth(dx+c) - 5ab^4 \coth(dx+c) - \frac{5ab^4 \coth(dx+c)^7}{7} - 2a^2b^3 \coth(dx+c)^5 - 1050a^3b^2 \coth(dx+c)^3 - 1050a^2b^3 \coth(dx+c) - 1050ab^4 \coth(dx+c) - 1050b^5}{d}$
default	$\frac{-5a^4b \coth(dx+c) - 10a^3b^2 \coth(dx+c) - 10a^2b^3 \coth(dx+c) - 5ab^4 \coth(dx+c) - \frac{5ab^4 \coth(dx+c)^7}{7} - 2a^2b^3 \coth(dx+c)^5 - 1050a^3b^2 \coth(dx+c)^3 - 1050a^2b^3 \coth(dx+c) - 1050ab^4 \coth(dx+c) - 1050b^5}{d}$
parts	$a^5x + \frac{b^5 \left(-\frac{\coth(dx+c)^9}{9} - \frac{\coth(dx+c)^7}{7} - \frac{\coth(dx+c)^5}{5} - \frac{\coth(dx+c)^3}{3} - \coth(dx+c) - \frac{\ln(\coth(dx+c)-1)}{2} + \frac{\ln(\coth(dx+c)+1)}{2} \right)}{d}$
risc	$a^5x + 5ba^4x + 10a^3b^2x + 10b^3a^2x + 5ab^4x + b^5x - \frac{2b(-31500a^3be^{2dx+2c} - 34020a^2b^2e^{2dx+2c} - 17400ab^3e^{2dx+2c} - 3150b^4e^{2dx+2c} - 3150b^5e^{2dx+2c})}{d}$

[In] int((a+coth(d*x+c)^2*b)^5,x,method=_RETURNVERBOSE)

```
[Out] 1/315*(-35*b^5*coth(d*x+c)^9+(-225*a*b^4-45*b^5)*coth(d*x+c)^7+(-630*a^2*b^3-315*a*b^4-63*b^5)*coth(d*x+c)^5+(-1050*a^3*b^2-1050*a^2*b^3-525*a*b^4-105*b^5)*coth(d*x+c)^3-1575*(a^4+2*a^3*b+2*a^2*b^2+a*b^3+1/5*b^4)*b*coth(d*x+c)+315*d*x*(a+b)^5)/d
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2111 vs. 2(152) = 304.

Time = 0.27 (sec) , antiderivative size = 2111, normalized size of antiderivative = 13.19

$$\int (a + b \coth^2(c + dx))^5 dx = \text{Too large to display}$$

```
[In] integrate((a+b*coth(d*x+c)^2)^5,x, algorithm="fricas")
```

```
[Out] -1/315*((1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5)*cosh(d*x + c)^9 + 9*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5)*cosh(d*x + c)*sinh(d*x + c)^8 - (1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x)*sinh(d*x + c)^9 - 9*(1225*a^4*b + 2800*a^3*b^2 + 2730*a^2*b^3 + 1240*a*b^4 + 213*b^5)*cosh(d*x + c)^7 + 9*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x - 4*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x)*cosh(d*x + c)^2)*sinh(d*x + c)^7 + 21*(4*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5)*cosh(d*x + c)^3 - 3*(1225*a^4*b + 2800*a^3*b^2 + 2730*a^2*b^3 + 1240*a*b^4 + 213*b^5)*cosh(d*x + c))*sinh(d*x + c)^6 + 36*(875*a^4*b + 1750*a^3*b^2 + 1680*a^2*b^3 + 890*a*b^4 + 213*b^5)*cosh(d*x + c)^5 - 9*(6300*a^4*b + 16800*a^3*b^2 + 19320*a^2*b^3 + 10560*a*b^4 + 2252*b^5 + 14*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x)*cosh(d*x + c)^4 + 1260*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x - 21*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x)*cosh(d*x + c)^2)*sinh(d*x + c)^5 + 9*(14*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5)*cosh(d*x + c)^5 - 35*(1225*a^4*b + 2800*a^3*b^2 + 2730*a^2*b^3 + 1240*a*b^4 + 213*b^5)*cosh(d*x + c)^3 + 20*(875*a^4*b + 1750*a^3*b^2 + 1680*a^2*b^3 + 890*a*b^4 + 213*b^5)*cosh(d*x + c))*sinh(d*x + c)^4 - 84*(525*a^4*b + 950*a^3*b^2 + 980*a^2*b^3 + 490*a*b^4 + 63*b^5)*cosh(d*x + c)^3 - 3*(28*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x)*cosh(d*x + c)^6 - 44100*a^4*b - 117600*a^3*b^2 - 135240*a^2*b^3 - 73920*a*b^4 - 15764*b^5 - 105*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x)*cosh(d*x + c)^4 - 8820*(a^5 + 5*a^4*b
```

+ 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x + 120*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x)*cosh(d*x + c)^2)*sinh(d*x + c)^3 + 9*(4*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5)*cosh(d*x + c)^7 - 21*(1225*a^4*b + 2800*a^3*b^2 + 2730*a^2*b^3 + 1240*a*b^4 + 213*b^5)*cosh(d*x + c)^5 + 40*(875*a^4*b + 1750*a^3*b^2 + 1680*a^2*b^3 + 890*a*b^4 + 213*b^5)*cosh(d*x + c)^3 - 28*(525*a^4*b + 950*a^3*b^2 + 980*a^2*b^3 + 490*a*b^4 + 63*b^5)*cosh(d*x + c))*sinh(d*x + c)^2 + 126*(175*a^4*b + 300*a^3*b^2 + 330*a^2*b^3 + 140*a*b^4 + 63*b^5)*cosh(d*x + c) - 9*((1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x)*cosh(d*x + c)^8 - 7*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x)*cosh(d*x + c)^6 + 22050*a^4*b + 58800*a^3*b^2 + 67620*a^2*b^3 + 36960*a*b^4 + 7882*b^5 + 20*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x)*cosh(d*x + c)^4 + 4410*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x - 28*(1575*a^4*b + 4200*a^3*b^2 + 4830*a^2*b^3 + 2640*a*b^4 + 563*b^5 + 315*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x)*cosh(d*x + c)^2)*sinh(d*x + c))/(d*sinh(d*x + c)^9 + 9*(4*d*cosh(d*x + c)^2 - d)*sinh(d*x + c)^7 + 9*(14*d*cosh(d*x + c)^4 - 21*d*cosh(d*x + c)^2 + 4*d)*sinh(d*x + c)^5 + 3*(28*d*cosh(d*x + c)^6 - 105*d*cosh(d*x + c)^4 + 120*d*cosh(d*x + c)^2 - 28*d)*sinh(d*x + c)^3 + 9*(d*cosh(d*x + c)^8 - 7*d*cosh(d*x + c)^6 + 20*d*cosh(d*x + c)^4 - 28*d*cosh(d*x + c)^2 + 14*d)*sinh(d*x + c))

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 639 vs. $2(148) = 296$.

Time = 14.40 (sec) , antiderivative size = 639, normalized size of antiderivative = 3.99

$$\int (a + b \coth^2(c + dx))^5 dx$$

$$= \begin{cases} x(a + b \coth^2(c))^5 \\ -\frac{a^5 \log(-e^{-dx})}{d} - \frac{5a^4 b \log(-e^{-dx}) \coth^2(dx + \log(-e^{-dx}))}{d} - \frac{10a^3 b^2 \log(-e^{-dx}) \coth^4(dx + \log(-e^{-dx}))}{d} - \frac{10a^2 b^3 \log(-e^{-dx}) \coth^6(dx + \log(-e^{-dx}))}{d} \\ a^5 x + 5a^4 b x \coth^2(dx + \log(e^{-dx})) + 10a^3 b^2 x \coth^4(dx + \log(e^{-dx})) + 10a^2 b^3 x \coth^6(dx + \log(e^{-dx})) \\ a^5 x + 5a^4 b x - \frac{5a^4 b}{d \tanh(c+dx)} + 10a^3 b^2 x - \frac{10a^3 b^2}{d \tanh(c+dx)} - \frac{10a^3 b^2}{3d \tanh^3(c+dx)} + 10a^2 b^3 x - \frac{10a^2 b^3}{d \tanh(c+dx)} - \frac{10a^2 b^3}{3d \tanh^3(c+dx)} \end{cases}$$

[In] integrate((a+b*coth(d*x+c)**2)**5,x)

[Out] Piecewise((x*(a + b*coth(c)**2)**5, Eq(d, 0)), (-a**5*log(-exp(-d*x))/d - 5*a**4*b*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**2/d - 10*a**3*b**2*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**4/d - 10*a**2*b**3*log(-exp(-d*x)

```

)*coth(d*x + log(-exp(-d*x)))*6/d - 5*a*b**4*log(-exp(-d*x))*coth(d*x + lo
g(-exp(-d*x)))*8/d - b**5*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))*10/
d, Eq(c, log(-exp(-d*x))), (a**5*x + 5*a**4*b*x*coth(d*x + log(exp(-d*x)))
**2 + 10*a**3*b**2*x*coth(d*x + log(exp(-d*x)))**4 + 10*a**2*b**3*x*coth(d*
x + log(exp(-d*x)))**6 + 5*a*b**4*x*coth(d*x + log(exp(-d*x)))**8 + b**5*x*
coth(d*x + log(exp(-d*x)))**10, Eq(c, log(exp(-d*x))), (a**5*x + 5*a**4*b*
x - 5*a**4*b/(d*tanh(c + d*x)) + 10*a**3*b**2*x - 10*a**3*b**2/(d*tanh(c +
d*x)) - 10*a**3*b**2/(3*d*tanh(c + d*x)**3) + 10*a**2*b**3*x - 10*a**2*b**3
/(d*tanh(c + d*x)) - 10*a**2*b**3/(3*d*tanh(c + d*x)**3) - 2*a**2*b**3/(d*t
anh(c + d*x)**5) + 5*a*b**4*x - 5*a*b**4/(d*tanh(c + d*x)) - 5*a*b**4/(3*d*
tanh(c + d*x)**3) - a*b**4/(d*tanh(c + d*x)**5) - 5*a*b**4/(7*d*tanh(c + d*
x)**7) + b**5*x - b**5/(d*tanh(c + d*x)) - b**5/(3*d*tanh(c + d*x)**3) - b*
**5/(5*d*tanh(c + d*x)**5) - b**5/(7*d*tanh(c + d*x)**7) - b**5/(9*d*tanh(c
+ d*x)**9), True))

```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 624 vs. $2(152) = 304$.

Time = 0.22 (sec) , antiderivative size = 624, normalized size of antiderivative = 3.90

$$\begin{aligned}
& \int (a + b \coth^2(c + dx))^5 dx \\
&= \frac{1}{315} b^5 \left(315x + \frac{315c}{d} - \frac{2(3492e^{(-2dx-2c)} - 13968e^{(-4dx-4c)} + 26292e^{(-6dx-6c)} - 39438e^{(-8dx-8c)} + 31500e^{(-10dx-10c)} - 21000e^{(-12dx-12c)} + 6300e^{(-14dx-14c)} - 1575e^{(-16dx-16c)} - 563)}{d(9e^{(-2dx-2c)} - 36e^{(-4dx-4c)} + 84e^{(-6dx-6c)} - 126e^{(-8dx-8c)} + 126e^{(-10dx-10c)} - 84e^{(-12dx-12c)} + 36e^{(-14dx-14c)} - 9e^{(-16dx-16c)} + e^{(-18dx-18c)} - 1)} \right) \\
&+ \frac{1}{21} ab^4 \left(105x + \frac{105c}{d} - \frac{8(203e^{(-2dx-2c)} - 609e^{(-4dx-4c)} + 770e^{(-6dx-6c)} - 770e^{(-8dx-8c)} + 315e^{(-10dx-10c)} - 563)}{d(7e^{(-2dx-2c)} - 21e^{(-4dx-4c)} + 35e^{(-6dx-6c)} - 35e^{(-8dx-8c)} + 21e^{(-10dx-10c)} - 7e^{(-12dx-12c)} + 7e^{(-14dx-14c)} - 7e^{(-16dx-16c)} + 7e^{(-18dx-18c)} - 1)} \right) \\
&+ \frac{2}{3} a^2 b^3 \left(15x + \frac{15c}{d} - \frac{2(70e^{(-2dx-2c)} - 140e^{(-4dx-4c)} + 90e^{(-6dx-6c)} - 45e^{(-8dx-8c)} - 23)}{d(5e^{(-2dx-2c)} - 10e^{(-4dx-4c)} + 10e^{(-6dx-6c)} - 5e^{(-8dx-8c)} + e^{(-10dx-10c)} - 1)} \right) \\
&+ \frac{10}{3} a^3 b^2 \left(3x + \frac{3c}{d} - \frac{4(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} - 2)}{d(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} + e^{(-6dx-6c)} - 1)} \right) \\
&+ 5a^4 b \left(x + \frac{c}{d} + \frac{2}{d(e^{(-2dx-2c)} - 1)} \right) + a^5 x
\end{aligned}$$

[In] integrate((a+b*coth(d*x+c)^2)^5,x, algorithm="maxima")

```

[Out] 1/315*b^5*(315*x + 315*c/d - 2*(3492*e^(-2*d*x - 2*c) - 13968*e^(-4*d*x - 4
*c) + 26292*e^(-6*d*x - 6*c) - 39438*e^(-8*d*x - 8*c) + 31500*e^(-10*d*x -
10*c) - 21000*e^(-12*d*x - 12*c) + 6300*e^(-14*d*x - 14*c) - 1575*e^(-16*d*
x - 16*c) - 563)/(d*(9*e^(-2*d*x - 2*c) - 36*e^(-4*d*x - 4*c) + 84*e^(-6*d*
x - 6*c) - 126*e^(-8*d*x - 8*c) + 126*e^(-10*d*x - 10*c) - 84*e^(-12*d*x -
12*c) + 36*e^(-14*d*x - 14*c) - 9*e^(-16*d*x - 16*c) + e^(-18*d*x - 18*c) -
1))) + 1/21*a*b^4*(105*x + 105*c/d - 8*(203*e^(-2*d*x - 2*c) - 609*e^(-4*d

```


Mupad [B] (verification not implemented)

Time = 2.11 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.99

$$\int (a + b \coth^2(c + dx))^5 dx = x(a + b)^5 - \frac{\coth(c + dx)^3 (10 a^3 b^2 + 10 a^2 b^3 + 5 a b^4 + b^5)}{3 d} - \frac{\coth(c + dx)^5 (10 a^2 b^3 + 5 a b^4 + b^5)}{5 d} - \frac{\coth(c + dx)^7 (b^5 + 5 a b^4)}{7 d} - \frac{b^5 \coth(c + dx)^9}{9 d} - \frac{b \coth(c + dx) (5 a^4 + 10 a^3 b + 10 a^2 b^2 + 5 a b^3 + b^4)}{d}$$

```
[In] int((a + b*coth(c + d*x)^2)^5,x)
```

```
[Out] x*(a + b)^5 - (coth(c + d*x)^3*(5*a*b^4 + b^5 + 10*a^2*b^3 + 10*a^3*b^2))/(3*d) - (coth(c + d*x)^5*(5*a*b^4 + b^5 + 10*a^2*b^3))/(5*d) - (coth(c + d*x)^7*(5*a*b^4 + b^5))/(7*d) - (b^5*coth(c + d*x)^9)/(9*d) - (b*coth(c + d*x)*(5*a*b^3 + 10*a^3*b + 5*a^4 + b^4 + 10*a^2*b^2))/d
```

3.2 $\int (a + b \coth^2(c + dx))^4 dx$

Optimal result	50
Rubi [A] (verified)	50
Mathematica [A] (verified)	52
Maple [A] (verified)	52
Fricas [B] (verification not implemented)	53
Sympy [B] (verification not implemented)	54
Maxima [B] (verification not implemented)	55
Giac [B] (verification not implemented)	55
Mupad [B] (verification not implemented)	56

Optimal result

Integrand size = 14, antiderivative size = 110

$$\int (a + b \coth^2(c + dx))^4 dx = (a + b)^4 x - \frac{b(2a + b)(2a^2 + 2ab + b^2) \coth(c + dx)}{d} - \frac{b^2(6a^2 + 4ab + b^2) \coth^3(c + dx)}{3d} - \frac{b^3(4a + b) \coth^5(c + dx)}{5d} - \frac{b^4 \coth^7(c + dx)}{7d}$$

[Out] (a+b)^4*x-b*(2*a+b)*(2*a^2+2*a*b+b^2)*coth(d*x+c)/d-1/3*b^2*(6*a^2+4*a*b+b^2)*coth(d*x+c)^3/d-1/5*b^3*(4*a+b)*coth(d*x+c)^5/d-1/7*b^4*coth(d*x+c)^7/d

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3742, 398, 212}

$$\int (a + b \coth^2(c + dx))^4 dx = -\frac{b^2(6a^2 + 4ab + b^2) \coth^3(c + dx)}{3d} - \frac{b(2a + b)(2a^2 + 2ab + b^2) \coth(c + dx)}{d} - \frac{b^3(4a + b) \coth^5(c + dx)}{5d} + x(a + b)^4 - \frac{b^4 \coth^7(c + dx)}{7d}$$

[In] Int[(a + b*Coth[c + d*x]^2)^4, x]

[Out] (a + b)^4*x - (b*(2*a + b)*(2*a^2 + 2*a*b + b^2)*Coth[c + d*x])/d - (b^2*(6*a^2 + 4*a*b + b^2)*Coth[c + d*x]^3)/(3*d) - (b^3*(4*a + b)*Coth[c + d*x]^5)/(5*d) - (b^4*Coth[c + d*x]^7)/(7*d)

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 398

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :=
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(
ff*x)^n)^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Subst}\left(\int \frac{(a+bx^2)^4}{1-x^2} dx, x, \coth(c+dx)\right)}{d} \\
&= \frac{\text{Subst}\left(\int \left(-b(2a+b)(2a^2+2ab+b^2) - b^2(6a^2+4ab+b^2)x^2 - b^3(4a+b)x^4 - b^4x^6 + \frac{(a+b)^4}{1-x^2}\right) dx, x, \coth(c+dx)\right)}{d} \\
&= -\frac{b(2a+b)(2a^2+2ab+b^2)\coth(c+dx)}{d} - \frac{b^2(6a^2+4ab+b^2)\coth^3(c+dx)}{3d} \\
&\quad - \frac{b^3(4a+b)\coth^5(c+dx)}{5d} - \frac{b^4\coth^7(c+dx)}{7d} + \frac{(a+b)^4\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \coth(c+dx)\right)}{d} \\
&= (a+b)^4x - \frac{b(2a+b)(2a^2+2ab+b^2)\coth(c+dx)}{d} \\
&\quad - \frac{b^2(6a^2+4ab+b^2)\coth^3(c+dx)}{3d} - \frac{b^3(4a+b)\coth^5(c+dx)}{5d} - \frac{b^4\coth^7(c+dx)}{7d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 2.25 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.15

$$\int (a + b \coth^2(c + dx))^4 dx = \frac{\coth(c + dx) \left(b(105(4a^3 + 6a^2b + 4ab^2 + b^3)) + 35b(6a^2 + 4ab + b^2) \coth^2(c + dx) + 21b^2(4a + b) \coth^4(c + dx) + 15b^3 \coth^6(c + dx) - 105(a + b)^4 \operatorname{ArcTanh}[\operatorname{Sqrt}[\operatorname{Tanh}^2(c + dx)]] \operatorname{Sqrt}[\operatorname{Tanh}^2(c + dx)] \right)}{105d}$$

`[In] Integrate[(a + b*Coth[c + d*x]^2)^4,x]`

```
[Out] -1/105*(Coth[c + d*x]*(b*(105*(4*a^3 + 6*a^2*b + 4*a*b^2 + b^3) + 35*b*(6*a^2 + 4*a*b + b^2)*Coth[c + d*x]^2 + 21*b^2*(4*a + b)*Coth[c + d*x]^4 + 15*b^3*Coth[c + d*x]^6) - 105*(a + b)^4*ArcTanh[Sqrt[Tanh[c + d*x]^2]]*Sqrt[Tanh[c + d*x]^2]))/d
```

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.02

method	result
parallelrisch	$\frac{-15b^4 \coth(dx+c)^7 + (-84ab^3 - 21b^4) \coth(dx+c)^5 + (-210a^2b^2 - 140ab^3 - 35b^4) \coth(dx+c)^3 + (-420a^3b - 630a^2b^2 - 420a^3b^3 - 105b^4) \coth(dx+c) + 105(a+b)^4 \operatorname{ArcTanh}[\operatorname{Sqrt}[\operatorname{Tanh}^2(dx+c)]] \operatorname{Sqrt}[\operatorname{Tanh}^2(dx+c)]}{105d}$
derivativedivides	$\frac{-4 \coth(dx+c)a^3b - 6a^2b^2 \coth(dx+c) - 4ab^3 \coth(dx+c) - \frac{4ab^3 \coth(dx+c)^5}{5} - 2a^2b^2 \coth(dx+c)^3 - \frac{4 \coth(dx+c)^3 ab^3}{3} - b^4 \coth(dx+c)}{d}$
default	$\frac{-4 \coth(dx+c)a^3b - 6a^2b^2 \coth(dx+c) - 4ab^3 \coth(dx+c) - \frac{4ab^3 \coth(dx+c)^5}{5} - 2a^2b^2 \coth(dx+c)^3 - \frac{4 \coth(dx+c)^3 ab^3}{3} - b^4 \coth(dx+c)}{d}$
parts	$x a^4 + \frac{b^4 \left(-\frac{\coth(dx+c)^7}{7} - \frac{\coth(dx+c)^5}{5} - \frac{\coth(dx+c)^3}{3} - \coth(dx+c) - \frac{\ln(\coth(dx+c)-1)}{2} + \frac{\ln(\coth(dx+c)+1)}{2} \right)}{d} + \frac{4ab^3 \left(-\frac{\coth(dx+c)^6}{6} - \frac{\coth(dx+c)^4}{4} - \frac{\coth(dx+c)^2}{2} - \coth(dx+c) \right)}{d}$
risch	$x a^4 + 4b a^3 x + 6a^2 b^2 x + 4a b^3 x + b^4 x - \frac{8b(161a^2b^2 + 105b^3e^{12dx+12c} + 105a^3 + 609e^{4dx+4c}b^3 + 315a^2b e^{12dx+12c})}{105d}$

`[In] int((a+coth(d*x+c)^2*b)^4,x,method=_RETURNVERBOSE)`

```
[Out] 1/105*(-15*b^4*coth(d*x+c)^7+(-84*a*b^3-21*b^4)*coth(d*x+c)^5+(-210*a^2*b^2-140*a*b^3-35*b^4)*coth(d*x+c)^3+(-420*a^3*b-630*a^2*b^2-420*a*b^3-105*b^4)*coth(d*x+c)+105*d*x*(a+b)^4)/d
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1164 vs. $2(104) = 208$.

Time = 0.25 (sec) , antiderivative size = 1164, normalized size of antiderivative = 10.58

$$\int (a + b \operatorname{coth}^2(c + dx))^4 dx = \text{Too large to display}$$

[In] integrate((a+b*coth(d*x+c)^2)^4,x, algorithm="fricas")

[Out]
$$\begin{aligned} & -1/105*(4*(105*a^3*b + 210*a^2*b^2 + 161*a*b^3 + 44*b^4)*\cosh(d*x + c)^7 + \\ & 28*(105*a^3*b + 210*a^2*b^2 + 161*a*b^3 + 44*b^4)*\cosh(d*x + c)*\sinh(d*x + \\ & c)^6 - (420*a^3*b + 840*a^2*b^2 + 644*a*b^3 + 176*b^4 + 105*(a^4 + 4*a^3*b \\ & + 6*a^2*b^2 + 4*a*b^3 + b^4)*d*x)*\sinh(d*x + c)^7 - 28*(75*a^3*b + 120*a^2* \\ & b^2 + 71*a*b^3 + 14*b^4)*\cosh(d*x + c)^5 + 7*(420*a^3*b + 840*a^2*b^2 + 644 \\ & *a*b^3 + 176*b^4 + 105*(a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a*b^3 + b^4)*d*x - 3* \\ & (420*a^3*b + 840*a^2*b^2 + 644*a*b^3 + 176*b^4 + 105*(a^4 + 4*a^3*b + 6*a^2 \\ & *b^2 + 4*a*b^3 + b^4)*d*x)*\cosh(d*x + c)^2*\sinh(d*x + c)^5 + 140*((105*a^3 \\ & *b + 210*a^2*b^2 + 161*a*b^3 + 44*b^4)*\cosh(d*x + c)^3 - (75*a^3*b + 120*a^ \\ & 2*b^2 + 71*a*b^3 + 14*b^4)*\cosh(d*x + c))*\sinh(d*x + c)^4 + 84*(45*a^3*b + \\ & 60*a^2*b^2 + 41*a*b^3 + 14*b^4)*\cosh(d*x + c)^3 - 7*(5*(420*a^3*b + 840*a^2 \\ & *b^2 + 644*a*b^3 + 176*b^4 + 105*(a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a*b^3 + b^4 \\ &)*d*x)*\cosh(d*x + c)^4 + 1260*a^3*b + 2520*a^2*b^2 + 1932*a*b^3 + 528*b^4 + \\ & 315*(a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a*b^3 + b^4)*d*x - 10*(420*a^3*b + 840* \\ & a^2*b^2 + 644*a*b^3 + 176*b^4 + 105*(a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a*b^3 + \\ & b^4)*d*x)*\cosh(d*x + c)^2*\sinh(d*x + c)^3 + 28*(3*(105*a^3*b + 210*a^2*b^2 \\ & + 161*a*b^3 + 44*b^4)*\cosh(d*x + c)^5 - 10*(75*a^3*b + 120*a^2*b^2 + 71*a* \\ & b^3 + 14*b^4)*\cosh(d*x + c)^3 + 9*(45*a^3*b + 60*a^2*b^2 + 41*a*b^3 + 14*b^ \\ & 4)*\cosh(d*x + c))*\sinh(d*x + c)^2 - 420*(5*a^3*b + 6*a^2*b^2 + 5*a*b^3)*\cos \\ & h(d*x + c) - 7*((420*a^3*b + 840*a^2*b^2 + 644*a*b^3 + 176*b^4 + 105*(a^4 + \\ & 4*a^3*b + 6*a^2*b^2 + 4*a*b^3 + b^4)*d*x)*\cosh(d*x + c)^6 - 5*(420*a^3*b + \\ & 840*a^2*b^2 + 644*a*b^3 + 176*b^4 + 105*(a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a*b \\ & ^3 + b^4)*d*x)*\cosh(d*x + c)^4 - 2100*a^3*b - 4200*a^2*b^2 - 3220*a*b^3 - 8 \\ & 80*b^4 - 525*(a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a*b^3 + b^4)*d*x + 9*(420*a^3*b \\ & + 840*a^2*b^2 + 644*a*b^3 + 176*b^4 + 105*(a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a \\ & *b^3 + b^4)*d*x)*\cosh(d*x + c)^2*\sinh(d*x + c))/(d*\sinh(d*x + c)^7 + 7*(3* \\ & d*\cosh(d*x + c)^2 - d)*\sinh(d*x + c)^5 + 7*(5*d*\cosh(d*x + c)^4 - 10*d*\cosh \\ & (d*x + c)^2 + 3*d)*\sinh(d*x + c)^3 + 7*(d*\cosh(d*x + c)^6 - 5*d*\cosh(d*x + \\ & c)^4 + 9*d*\cosh(d*x + c)^2 - 5*d)*\sinh(d*x + c)) \end{aligned}$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 478 vs. $2(99) = 198$.

Time = 5.83 (sec) , antiderivative size = 478, normalized size of antiderivative = 4.35

$$\int (a + b \coth^2(c + dx))^4 dx$$

$$= \begin{cases} x(a + b \coth^2(c))^4 \\ -\frac{a^4 \log(-e^{-dx})}{d} - \frac{4a^3 b \log(-e^{-dx}) \coth^2(dx + \log(-e^{-dx}))}{d} - \frac{6a^2 b^2 \log(-e^{-dx}) \coth^4(dx + \log(-e^{-dx}))}{d} - \frac{4ab^3 \log(-e^{-dx}) \coth^6(dx + \log(-e^{-dx}))}{d} \\ a^4 x + 4a^3 b x \coth^2(dx + \log(e^{-dx})) + 6a^2 b^2 x \coth^4(dx + \log(e^{-dx})) + 4ab^3 x \coth^6(dx + \log(e^{-dx})) + \\ a^4 x + 4a^3 b x - \frac{4a^3 b}{d \tanh(c+dx)} + 6a^2 b^2 x - \frac{6a^2 b^2}{d \tanh(c+dx)} - \frac{2a^2 b^2}{d \tanh^3(c+dx)} + 4ab^3 x - \frac{4ab^3}{d \tanh(c+dx)} - \frac{4ab^3}{3d \tanh^3(c+dx)} - \frac{b^4}{5d \tanh^5(c+dx)} \end{cases}$$

[In] integrate((a+b*coth(d*x+c)**2)**4,x)

[Out] Piecewise((x*(a + b*coth(c)**2)**4, Eq(d, 0)), (-a**4*log(-exp(-d*x))/d - 4*a**3*b*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**2/d - 6*a**2*b**2*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**4/d - 4*a*b**3*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**6/d - b**4*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**8/d, Eq(c, log(-exp(-d*x)))), (a**4*x + 4*a**3*b*x*coth(d*x + log(exp(-d*x)))**2 + 6*a**2*b**2*x*coth(d*x + log(exp(-d*x)))**4 + 4*a*b**3*x*coth(d*x + log(exp(-d*x)))**6 + b**4*x*coth(d*x + log(exp(-d*x)))**8, Eq(c, log(exp(-d*x)))), (a**4*x + 4*a**3*b*x - 4*a**3*b/(d*tanh(c + d*x)) + 6*a**2*b**2*x - 6*a**2*b**2/(d*tanh(c + d*x)) - 2*a**2*b**2/(d*tanh(c + d*x)**3) + 4*a*b**3*x - 4*a*b**3/(d*tanh(c + d*x)) - 4*a*b**3/(3*d*tanh(c + d*x)**3) - 4*a*b**3/(5*d*tanh(c + d*x)**5) + b**4*x - b**4/(d*tanh(c + d*x)) - b**4/(3*d*tanh(c + d*x)**3) - b**4/(5*d*tanh(c + d*x)**5) - b**4/(7*d*tanh(c + d*x)**7), True))

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 410 vs. $2(104) = 208$.

Time = 0.21 (sec) , antiderivative size = 410, normalized size of antiderivative = 3.73

$$\int (a + b \coth^2(c + dx))^4 dx$$

$$= \frac{1}{105} b^4 \left(105x + \frac{105c}{d} - \frac{8(203e^{(-2dx-2c)} - 609e^{(-4dx-4c)} + 770e^{(-6dx-6c)} - 770e^{(-8dx-8c)} + 315e^{(-10dx-10c)} - 105e^{(-12dx-12c)} - 44)}{d(7e^{(-2dx-2c)} - 21e^{(-4dx-4c)} + 35e^{(-6dx-6c)} - 35e^{(-8dx-8c)} + 21e^{(-10dx-10c)} - 7e^{(-12dx-12c)} + e^{(-14dx-14c)} - 1)} \right)$$

$$+ \frac{4}{15} ab^3 \left(15x + \frac{15c}{d} - \frac{2(70e^{(-2dx-2c)} - 140e^{(-4dx-4c)} + 90e^{(-6dx-6c)} - 45e^{(-8dx-8c)} - 23)}{d(5e^{(-2dx-2c)} - 10e^{(-4dx-4c)} + 10e^{(-6dx-6c)} - 5e^{(-8dx-8c)} + e^{(-10dx-10c)} - 1)} \right)$$

$$+ 2a^2b^2 \left(3x + \frac{3c}{d} - \frac{4(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} - 2)}{d(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} + e^{(-6dx-6c)} - 1)} \right)$$

$$+ 4a^3b \left(x + \frac{c}{d} + \frac{2}{d(e^{(-2dx-2c)} - 1)} \right) + a^4x$$

[In] integrate((a+b*coth(d*x+c)^2)^4,x, algorithm="maxima")

[Out] 1/105*b^4*(105*x + 105*c/d - 8*(203*e^(-2*d*x - 2*c) - 609*e^(-4*d*x - 4*c) + 770*e^(-6*d*x - 6*c) - 770*e^(-8*d*x - 8*c) + 315*e^(-10*d*x - 10*c) - 105*e^(-12*d*x - 12*c) - 44)/(d*(7*e^(-2*d*x - 2*c) - 21*e^(-4*d*x - 4*c) + 35*e^(-6*d*x - 6*c) - 35*e^(-8*d*x - 8*c) + 21*e^(-10*d*x - 10*c) - 7*e^(-12*d*x - 12*c) + e^(-14*d*x - 14*c) - 1))) + 4/15*a*b^3*(15*x + 15*c/d - 2*(70*e^(-2*d*x - 2*c) - 140*e^(-4*d*x - 4*c) + 90*e^(-6*d*x - 6*c) - 45*e^(-8*d*x - 8*c) - 23)/(d*(5*e^(-2*d*x - 2*c) - 10*e^(-4*d*x - 4*c) + 10*e^(-6*d*x - 6*c) - 5*e^(-8*d*x - 8*c) + e^(-10*d*x - 10*c) - 1))) + 2*a^2*b^2*(3*x + 3*c/d - 4*(3*e^(-2*d*x - 2*c) - 3*e^(-4*d*x - 4*c) - 2)/(d*(3*e^(-2*d*x - 2*c) - 3*e^(-4*d*x - 4*c) + e^(-6*d*x - 6*c) - 1))) + 4*a^3*b*(x + c/d + 2/(d*(e^(-2*d*x - 2*c) - 1))) + a^4*x

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 447 vs. $2(104) = 208$.

Time = 0.32 (sec) , antiderivative size = 447, normalized size of antiderivative = 4.06

$$\int (a + b \coth^2(c + dx))^4 dx$$

$$= \frac{105(a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4)(dx + c) - \frac{8(105a^3be^{(12dx+12c)} + 315a^2b^2e^{(12dx+12c)} + 315ab^3e^{(12dx+12c)} + 105b^4e^{(12dx+12c)} - 105a^3be^{(-12dx-12c)} - 315a^2b^2e^{(-12dx-12c)} - 315ab^3e^{(-12dx-12c)} - 105b^4e^{(-12dx-12c)} - 44)}{d(7e^{(-2dx-2c)} - 21e^{(-4dx-4c)} + 35e^{(-6dx-6c)} - 35e^{(-8dx-8c)} + 21e^{(-10dx-10c)} - 7e^{(-12dx-12c)} + e^{(-14dx-14c)} - 1)}}{d} + \frac{4(15a^3b^3(15x + 15c/d - 2(70e^{(-2dx-2c)} - 140e^{(-4dx-4c)} + 90e^{(-6dx-6c)} - 45e^{(-8dx-8c)} - 23)/(d(5e^{(-2dx-2c)} - 10e^{(-4dx-4c)} + 10e^{(-6dx-6c)} - 5e^{(-8dx-8c)} + e^{(-10dx-10c)} - 1))) + 2a^2b^2(3x + 3c/d - 4(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} - 2)/(d(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} + e^{(-6dx-6c)} - 1))) + 4a^3b(x + c/d + 2/(d(e^{(-2dx-2c)} - 1))) + a^4x)}{15}$$

[In] integrate((a+b*coth(d*x+c)^2)^4,x, algorithm="giac")

```
[Out] 1/105*(105*(a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a*b^3 + b^4)*(d*x + c) - 8*(105*a^3*b*e^(12*d*x + 12*c) + 315*a^2*b^2*e^(12*d*x + 12*c) + 315*a*b^3*e^(12*d*x + 12*c) + 105*b^4*e^(12*d*x + 12*c) - 630*a^3*b*e^(10*d*x + 10*c) - 1575*a^2*b^2*e^(10*d*x + 10*c) - 1260*a*b^3*e^(10*d*x + 10*c) - 315*b^4*e^(10*d*x + 10*c) + 1575*a^3*b*e^(8*d*x + 8*c) + 3360*a^2*b^2*e^(8*d*x + 8*c) + 2555*a*b^3*e^(8*d*x + 8*c) + 770*b^4*e^(8*d*x + 8*c) - 2100*a^3*b*e^(6*d*x + 6*c) - 3990*a^2*b^2*e^(6*d*x + 6*c) - 3080*a*b^3*e^(6*d*x + 6*c) - 770*b^4*e^(6*d*x + 6*c) + 1575*a^3*b*e^(4*d*x + 4*c) + 2835*a^2*b^2*e^(4*d*x + 4*c) + 2121*a*b^3*e^(4*d*x + 4*c) + 609*b^4*e^(4*d*x + 4*c) - 630*a^3*b*e^(2*d*x + 2*c) - 1155*a^2*b^2*e^(2*d*x + 2*c) - 812*a*b^3*e^(2*d*x + 2*c) - 203*b^4*e^(2*d*x + 2*c) + 105*a^3*b + 210*a^2*b^2 + 161*a*b^3 + 44*b^4)/(e^(2*d*x + 2*c) - 1)^7)/d
```

Mupad [B] (verification not implemented)

Time = 2.05 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.01

$$\int (a + b \coth^2(c + dx))^4 dx = x(a + b)^4 - \frac{\coth(c + dx)^3 (6a^2b^2 + 4ab^3 + b^4)}{3d} - \frac{\coth(c + dx)^5 (b^4 + 4ab^3)}{5d} - \frac{b^4 \coth(c + dx)^7}{7d} - \frac{b \coth(c + dx) (4a^3 + 6a^2b + 4ab^2 + b^3)}{d}$$

```
[In] int((a + b*coth(c + d*x)^2)^4,x)
```

```
[Out] x*(a + b)^4 - (coth(c + d*x)^3*(4*a*b^3 + b^4 + 6*a^2*b^2))/(3*d) - (coth(c + d*x)^5*(4*a*b^3 + b^4))/(5*d) - (b^4*coth(c + d*x)^7)/(7*d) - (b*coth(c + d*x)*(4*a*b^2 + 6*a^2*b + 4*a^3 + b^3))/d
```


3.3 $\int (a + b \coth^2(c + dx))^3 dx$

Optimal result	57
Rubi [A] (verified)	57
Mathematica [A] (verified)	58
Maple [A] (verified)	59
Fricas [B] (verification not implemented)	59
Sympy [B] (verification not implemented)	60
Maxima [B] (verification not implemented)	60
Giac [B] (verification not implemented)	61
Mupad [B] (verification not implemented)	62

Optimal result

Integrand size = 14, antiderivative size = 74

$$\int (a + b \coth^2(c + dx))^3 dx = (a + b)^3 x - \frac{b(3a^2 + 3ab + b^2) \coth(c + dx)}{d} - \frac{b^2(3a + b) \coth^3(c + dx)}{3d} - \frac{b^3 \coth^5(c + dx)}{5d}$$

[Out] (a+b)^3*x-b*(3*a^2+3*a*b+b^2)*coth(d*x+c)/d-1/3*b^2*(3*a+b)*coth(d*x+c)^3/d-1/5*b^3*coth(d*x+c)^5/d

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3742, 398, 212}

$$\int (a + b \coth^2(c + dx))^3 dx = -\frac{b(3a^2 + 3ab + b^2) \coth(c + dx)}{d} - \frac{b^2(3a + b) \coth^3(c + dx)}{3d} + x(a + b)^3 - \frac{b^3 \coth^5(c + dx)}{5d}$$

[In] Int[(a + b*Coth[c + d*x]^2)^3,x]

[Out] (a + b)^3*x - (b*(3*a^2 + 3*a*b + b^2)*Coth[c + d*x])/d - (b^2*(3*a + b)*Coth[c + d*x]^3)/(3*d) - (b^3*Coth[c + d*x]^5)/(5*d)

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt

$Q[a, 0] \parallel LtQ[b, 0]$

Rule 398

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :>
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(
ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int \frac{(a+bx^2)^3}{1-x^2} dx, x, \coth(c+dx)\right)}{d} \\
 &= \frac{\text{Subst}\left(\int \left(-b(3a^2+3ab+b^2) - b^2(3a+b)x^2 - b^3x^4 + \frac{(a+b)^3}{1-x^2}\right) dx, x, \coth(c+dx)\right)}{d} \\
 &= -\frac{b(3a^2+3ab+b^2)\coth(c+dx)}{d} - \frac{b^2(3a+b)\coth^3(c+dx)}{3d} \\
 &\quad - \frac{b^3\coth^5(c+dx)}{5d} + \frac{(a+b)^3\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \coth(c+dx)\right)}{d} \\
 &= (a+b)^3x - \frac{b(3a^2+3ab+b^2)\coth(c+dx)}{d} - \frac{b^2(3a+b)\coth^3(c+dx)}{3d} - \frac{b^3\coth^5(c+dx)}{5d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 1.51 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.35

$$\begin{aligned}
 &\int (a + b \coth^2(c + dx))^3 dx \\
 &= -\frac{b \coth(c + dx) (15(3a^2 + 3ab + b^2) + 5b(3a + b) \coth^2(c + dx) + 3b^2 \coth^4(c + dx))}{15d} \\
 &\quad + \frac{(a + b)^3 \operatorname{arctanh}\left(\sqrt{\tanh^2(c + dx)}\right) \tanh(c + dx)}{d \sqrt{\tanh^2(c + dx)}}
 \end{aligned}$$

[In] Integrate[(a + b*Coth[c + d*x]^2)^3,x]

[Out] $-1/15*(b*\text{Coth}[c + d*x]*(15*(3*a^2 + 3*a*b + b^2) + 5*b*(3*a + b)*\text{Coth}[c + d*x]^2 + 3*b^2*\text{Coth}[c + d*x]^4))/d + ((a + b)^3*\text{ArcTanh}[\text{Sqrt}[\text{Tanh}[c + d*x]^2]])*\text{Tanh}[c + d*x]/(d*\text{Sqrt}[\text{Tanh}[c + d*x]^2])$

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.01

method	result
parallelrisch	$\frac{-3b^3 \coth(dx+c)^5 + (-15ab^2 - 5b^3) \coth(dx+c)^3 + (-45a^2b - 45ab^2 - 15b^3) \coth(dx+c) + 15dx(a+b)^3}{15d}$
derivativedivides	$\frac{-3a^2b \coth(dx+c) - 3 \coth(dx+c) a b^2 - a b^2 \coth(dx+c)^3 - \frac{b^3 \coth(dx+c)^3}{3} - b^3 \coth(dx+c) - \frac{b^3 \coth(dx+c)^5}{5} + \frac{(a^3 + 3a^2b + 3ab^2 + b^3) \coth(dx+c)}{d}}{d}$
default	$\frac{-3a^2b \coth(dx+c) - 3 \coth(dx+c) a b^2 - a b^2 \coth(dx+c)^3 - \frac{b^3 \coth(dx+c)^3}{3} - b^3 \coth(dx+c) - \frac{b^3 \coth(dx+c)^5}{5} + \frac{(a^3 + 3a^2b + 3ab^2 + b^3) \coth(dx+c)}{d}}{d}$
parts	$a^3x + \frac{b^3 \left(-\frac{\coth(dx+c)^5}{5} - \frac{\coth(dx+c)^3}{3} - \coth(dx+c) - \frac{\ln(\coth(dx+c)-1) + \ln(\coth(dx+c)+1)}{2} \right)}{d} + \frac{3ab^2 \left(-\frac{\coth(dx+c)^3}{3} - \coth(dx+c) \right)}{d}$
risch	$a^3x + 3ba^2x + 3a^2bx + b^3x - \frac{2b(45a^2e^{8dx+8c} + 90abe^{8dx+8c} + 45b^2e^{8dx+8c} - 180a^2e^{6dx+6c} - 270abe^{6dx+6c} - 90b^3e^{6dx+6c})}{d}$

[In] int((a+coth(d*x+c)^2*b)^3,x,method=_RETURNVERBOSE)

[Out] $1/15*(-3*b^3*\text{coth}(d*x+c)^5 + (-15*a*b^2 - 5*b^3)*\text{coth}(d*x+c)^3 + (-45*a^2*b - 45*a*b^2 - 15*b^3)*\text{coth}(d*x+c) + 15*d*x*(a+b)^3)/d$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 557 vs. $2(70) = 140$.

Time = 0.27 (sec) , antiderivative size = 557, normalized size of antiderivative = 7.53

$$\int (a + b \coth^2(c + dx))^3 dx = \frac{(45a^2b + 60ab^2 + 23b^3) \cosh(dx+c)^5 + 5(45a^2b + 60ab^2 + 23b^3) \cosh(dx+c) \sinh(dx+c)^4 - (45a^2b + 60ab^2 + 23b^3) \sinh(dx+c)^5}{d}$$

[In] integrate((a+b*coth(d*x+c)^2)^3,x, algorithm="fricas")

[Out] $-1/15*((45*a^2*b + 60*a*b^2 + 23*b^3)*\cosh(d*x + c)^5 + 5*(45*a^2*b + 60*a*b^2 + 23*b^3)*\cosh(d*x + c)*\sinh(d*x + c)^4 - (45*a^2*b + 60*a*b^2 + 23*b^3)*\sinh(d*x + c)^5 - 5*(27*a^2*b + 24*a*b^2 + 5*b^3)*\cosh(d*x + c)^3 + 5*(45*a^2*b + 60*a*b^2 + 23*b^3 + 15*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*x - 2*(45*a^2*b + 60*a*b^2 + 23*b^3 + 15*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*x)*\cosh(d*x + c)^2*\sinh(d*x + c)^3 + 5*(27*a^2*b + 24*a*b^2 + 5*b^3)*\sinh(d*x + c)^3)$

$(45a^2b + 60ab^2 + 23b^3) \cosh(dx + c)^3 - 3(27a^2b + 24ab^2 + 5b^3) \cosh(dx + c) \sinh(dx + c)^2 + 10(9a^2b + 6ab^2 + 5b^3) \cosh(dx + c) - 5((45a^2b + 60ab^2 + 23b^3 + 15(a^3 + 3a^2b + 3ab^2 + b^3)dx) \cosh(dx + c)^4 + 90a^2b + 120ab^2 + 46b^3 + 30(a^3 + 3a^2b + 3ab^2 + b^3)dx - 3(45a^2b + 60ab^2 + 23b^3 + 15(a^3 + 3a^2b + 3ab^2 + b^3)dx) \cosh(dx + c)^2 \sinh(dx + c)) / (d \sinh(dx + c)^5 + 5(2d \cosh(dx + c)^2 - d) \sinh(dx + c)^3 + 5(d \cosh(dx + c)^4 - 3d \cosh(dx + c)^2 + 2d) \sinh(dx + c))$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 332 vs. $2(65) = 130$.

Time = 2.33 (sec) , antiderivative size = 332, normalized size of antiderivative = 4.49

$$\int (a + b \coth^2(c + dx))^3 dx$$

$$= \begin{cases} x(a + b \coth^2(c))^3 \\ -\frac{a^3 \log(-e^{-dx})}{d} - \frac{3a^2 b \log(-e^{-dx}) \coth^2(dx + \log(-e^{-dx}))}{d} - \frac{3ab^2 \log(-e^{-dx}) \coth^4(dx + \log(-e^{-dx}))}{d} - \frac{b^3 \log(-e^{-dx}) \coth^6(dx + \log(-e^{-dx}))}{d} \\ a^3 x + 3a^2 b x \coth^2(dx + \log(e^{-dx})) + 3ab^2 x \coth^4(dx + \log(e^{-dx})) + b^3 x \coth^6(dx + \log(e^{-dx})) \\ a^3 x + 3a^2 b x - \frac{3a^2 b}{d \tanh(c+dx)} + 3ab^2 x - \frac{3ab^2}{d \tanh(c+dx)} - \frac{ab^2}{d \tanh^3(c+dx)} + b^3 x - \frac{b^3}{d \tanh(c+dx)} - \frac{b^3}{3d \tanh^3(c+dx)} - \frac{b^3}{5d \tanh^5(c+dx)} \end{cases}$$

[In] integrate((a+b*coth(d*x+c)**2)**3,x)

[Out] Piecewise((x*(a + b*coth(c)**2)**3, Eq(d, 0)), (-a**3*log(-exp(-d*x))/d - 3*a**2*b*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**2/d - 3*a*b**2*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**4/d - b**3*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**6/d, Eq(c, log(-exp(-d*x)))), (a**3*x + 3*a**2*b*x*coth(d*x + log(exp(-d*x)))**2 + 3*a*b**2*x*coth(d*x + log(exp(-d*x)))**4 + b**3*x*coth(d*x + log(exp(-d*x)))**6, Eq(c, log(exp(-d*x)))), (a**3*x + 3*a**2*b*x - 3*a**2*b/(d*tanh(c + d*x)) + 3*a*b**2*x - 3*a*b**2/(d*tanh(c + d*x)) - a*b**2/(d*tanh(c + d*x)**3) + b**3*x - b**3/(d*tanh(c + d*x)) - b**3/(3*d*tanh(c + d*x)**3) - b**3/(5*d*tanh(c + d*x)**5), True))

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 239 vs. $2(70) = 140$.

Mupad [B] (verification not implemented)

Time = 2.01 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.97

$$\int (a + b \coth^2(c + dx))^3 dx = x(a + b)^3 - \frac{\coth(c + dx)^3 (b^3 + 3ab^2)}{3d} - \frac{b^3 \coth(c + dx)^5}{5d} - \frac{b \coth(c + dx) (3a^2 + 3ab + b^2)}{d}$$

[In] int((a + b*coth(c + d*x)^2)^3,x)

[Out] x*(a + b)^3 - (coth(c + d*x)^3*(3*a*b^2 + b^3))/(3*d) - (b^3*coth(c + d*x)^5)/(5*d) - (b*coth(c + d*x)*(3*a*b + 3*a^2 + b^2))/d

3.4 $\int (a + b \coth^2(c + dx))^2 dx$

Optimal result	63
Rubi [A] (verified)	63
Mathematica [A] (verified)	64
Maple [A] (verified)	65
Fricas [B] (verification not implemented)	65
Sympy [B] (verification not implemented)	66
Maxima [B] (verification not implemented)	66
Giac [B] (verification not implemented)	67
Mupad [B] (verification not implemented)	67

Optimal result

Integrand size = 14, antiderivative size = 43

$$\int (a + b \coth^2(c + dx))^2 dx = (a + b)^2 x - \frac{b(2a + b) \coth(c + dx)}{d} - \frac{b^2 \coth^3(c + dx)}{3d}$$

[Out] (a+b)^2*x-b*(2*a+b)*coth(d*x+c)/d-1/3*b^2*coth(d*x+c)^3/d

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3742, 398, 212}

$$\int (a + b \coth^2(c + dx))^2 dx = -\frac{b(2a + b) \coth(c + dx)}{d} + x(a + b)^2 - \frac{b^2 \coth^3(c + dx)}{3d}$$

[In] Int[(a + b*Coth[c + d*x]^2)^2,x]

[Out] (a + b)^2*x - (b*(2*a + b)*Coth[c + d*x])/d - (b^2*Coth[c + d*x]^3)/(3*d)

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 398

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0]

0] && GeQ[p, -q]

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :>
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(
ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int \frac{(a+bx^2)^2}{1-x^2} dx, x, \coth(c+dx)\right)}{d} \\
 &= \frac{\text{Subst}\left(\int \left(-b(2a+b) - b^2x^2 + \frac{(a+b)^2}{1-x^2}\right) dx, x, \coth(c+dx)\right)}{d} \\
 &= -\frac{b(2a+b)\coth(c+dx)}{d} - \frac{b^2\coth^3(c+dx)}{3d} + \frac{(a+b)^2\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \coth(c+dx)\right)}{d} \\
 &= (a+b)^2x - \frac{b(2a+b)\coth(c+dx)}{d} - \frac{b^2\coth^3(c+dx)}{3d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.51

$$\int (a + b \coth^2(c + dx))^2 dx = \frac{\coth(c + dx) \left(b(6a + 3b + b \coth^2(c + dx)) - 3(a + b)^2 \operatorname{arctanh}\left(\sqrt{\tanh^2(c + dx)}\right) \sqrt{\tanh^2(c + dx)} \right)}{3d}$$

[In] Integrate[(a + b*Coth[c + d*x]^2)^2,x]

[Out] -1/3*(Coth[c + d*x]*(b*(6*a + 3*b + b*Coth[c + d*x]^2) - 3*(a + b)^2*ArcTan h[Sqrt[Tanh[c + d*x]^2]]*Sqrt[Tanh[c + d*x]^2]))/d

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.07

method	result
parallelrisch	$\frac{-b^2 \coth(dx+c)^3 + (-6ab-3b^2) \coth(dx+c) + 3dx(a+b)^2}{3d}$
derivativedivides	$\frac{-\frac{b^2 \coth(dx+c)^3}{3} - 2 \coth(dx+c)ab - \coth(dx+c)b^2 - \frac{(a^2+2ab+b^2) \ln(\coth(dx+c)-1)}{2} + \frac{(a^2+2ab+b^2) \ln(\coth(dx+c)+1)}{2}}{d}$
default	$\frac{-\frac{b^2 \coth(dx+c)^3}{3} - 2 \coth(dx+c)ab - \coth(dx+c)b^2 - \frac{(a^2+2ab+b^2) \ln(\coth(dx+c)-1)}{2} + \frac{(a^2+2ab+b^2) \ln(\coth(dx+c)+1)}{2}}{d}$
risch	$a^2x + 2abx + b^2x - \frac{4b(3ae^{4dx+4c} + 3be^{4dx+4c} - 6e^{2dx+2c}a - 3be^{2dx+2c} + 3a+2b)}{3d(e^{2dx+2c}-1)^3}$
parts	$a^2x + \frac{b^2 \left(-\frac{\coth(dx+c)^3}{3} - \coth(dx+c) - \frac{\ln(\coth(dx+c)-1)}{2} + \frac{\ln(\coth(dx+c)+1)}{2} \right)}{d} + \frac{2ab \left(-\coth(dx+c) - \frac{\ln(\coth(dx+c)-1)}{2} \right)}{d}$

```
[In] int((a+coth(d*x+c)^2*b)^2,x,method=_RETURNVERBOSE)
```

```
[Out] 1/3*(-b^2*coth(d*x+c)^3+(-6*a*b-3*b^2)*coth(d*x+c)+3*d*x*(a+b)^2)/d
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 197 vs. 2(41) = 82.

Time = 0.26 (sec) , antiderivative size = 197, normalized size of antiderivative = 4.58

$$\int (a + b \coth^2(c + dx))^2 dx = \frac{2(3ab + 2b^2) \cosh(dx + c)^3 + 6(3ab + 2b^2) \cosh(dx + c) \sinh(dx + c)^2 - (3(a^2 + 2ab + b^2)dx + 6ab)}{3(d \sinh(dx + c))}$$

```
[In] integrate((a+b*coth(d*x+c)^2)^2,x, algorithm="fricas")
```

```
[Out] -1/3*(2*(3*a*b + 2*b^2)*cosh(d*x + c)^3 + 6*(3*a*b + 2*b^2)*cosh(d*x + c)*sinh(d*x + c)^2 - (3*(a^2 + 2*a*b + b^2)*d*x + 6*a*b + 4*b^2)*sinh(d*x + c)^3 - 6*a*b*cosh(d*x + c) + 3*(3*(a^2 + 2*a*b + b^2)*d*x - (3*(a^2 + 2*a*b + b^2)*d*x + 6*a*b + 4*b^2)*cosh(d*x + c)^2 + 6*a*b + 4*b^2)*sinh(d*x + c))/(d*sinh(d*x + c)^3 + 3*(d*cosh(d*x + c)^2 - d)*sinh(d*x + c))
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 211 vs. $2(36) = 72$.

Time = 1.09 (sec) , antiderivative size = 211, normalized size of antiderivative = 4.91

$$\int (a + b \coth^2(c + dx))^2 dx$$

$$= \begin{cases} x(a + b \coth^2(c))^2 & \text{for } d = 0 \\ -\frac{a^2 \log(-e^{-dx})}{d} - \frac{2ab \log(-e^{-dx}) \coth^2(dx + \log(-e^{-dx}))}{d} - \frac{b^2 \log(-e^{-dx}) \coth^4(dx + \log(-e^{-dx}))}{d} & \text{for } c = \log(-e^{-dx}) \\ a^2x + 2abx \coth^2(dx + \log(e^{-dx})) + b^2x \coth^4(dx + \log(e^{-dx})) & \text{for } c = \log(e^{-dx}) \\ a^2x + 2abx - \frac{2ab}{d \tanh(c+dx)} + b^2x - \frac{b^2}{d \tanh(c+dx)} - \frac{b^2}{3d \tanh^3(c+dx)} & \text{otherwise} \end{cases}$$

[In] integrate((a+b*coth(d*x+c)**2)**2,x)

[Out] Piecewise((x*(a + b*coth(c)**2)**2, Eq(d, 0)), (-a**2*log(-exp(-d*x))/d - 2*a*b*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**2/d - b**2*log(-exp(-d*x))*coth(d*x + log(-exp(-d*x)))**4/d, Eq(c, log(-exp(-d*x)))), (a**2*x + 2*a*b*x*coth(d*x + log(exp(-d*x)))**2 + b**2*x*coth(d*x + log(exp(-d*x)))**4, Eq(c, log(exp(-d*x)))), (a**2*x + 2*a*b*x - 2*a*b/(d*tanh(c + d*x)) + b**2*x - b**2/(d*tanh(c + d*x)) - b**2/(3*d*tanh(c + d*x)**3), True))

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 114 vs. $2(41) = 82$.

Time = 0.19 (sec) , antiderivative size = 114, normalized size of antiderivative = 2.65

$$\int (a + b \coth^2(c + dx))^2 dx$$

$$= \frac{1}{3} b^2 \left(3x + \frac{3c}{d} - \frac{4(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} - 2)}{d(3e^{(-2dx-2c)} - 3e^{(-4dx-4c)} + e^{(-6dx-6c)} - 1)} \right) + 2ab \left(x + \frac{c}{d} + \frac{2}{d(e^{(-2dx-2c)} - 1)} \right) + a^2x$$

[In] integrate((a+b*coth(d*x+c)^2)^2,x, algorithm="maxima")

[Out] 1/3*b^2*(3*x + 3*c/d - 4*(3*e^(-2*d*x - 2*c) - 3*e^(-4*d*x - 4*c) - 2)/(d*(3*e^(-2*d*x - 2*c) - 3*e^(-4*d*x - 4*c) + e^(-6*d*x - 6*c) - 1))) + 2*a*b*(x + c/d + 2/(d*(e^(-2*d*x - 2*c) - 1))) + a^2*x

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 103 vs. $2(41) = 82$.

Time = 0.28 (sec) , antiderivative size = 103, normalized size of antiderivative = 2.40

$$\int (a + b \coth^2(c + dx))^2 dx$$

$$= \frac{3(a^2 + 2ab + b^2)(dx + c) - \frac{4(3abe^{(4dx+4c)} + 3b^2e^{(4dx+4c)} - 6abe^{(2dx+2c)} - 3b^2e^{(2dx+2c)} + 3ab + 2b^2)}{(e^{(2dx+2c)} - 1)^3}}{3d}$$

[In] integrate((a+b*coth(d*x+c)^2)^2,x, algorithm="giac")

[Out] $\frac{1}{3} * (3 * (a^2 + 2 * a * b + b^2) * (d * x + c) - 4 * (3 * a * b * e^{(4 * d * x + 4 * c)} + 3 * b^2 * e^{(4 * d * x + 4 * c)} - 6 * a * b * e^{(2 * d * x + 2 * c)} - 3 * b^2 * e^{(2 * d * x + 2 * c)} + 3 * a * b + 2 * b^2) / (e^{(2 * d * x + 2 * c)} - 1)^3) / d$

Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.95

$$\int (a + b \coth^2(c + dx))^2 dx = x(a + b)^2 - \frac{b^2 \coth(c + dx)^3}{3d} - \frac{b \coth(c + dx)(2a + b)}{d}$$

[In] int((a + b*coth(c + d*x)^2)^2,x)

[Out] $x * (a + b)^2 - (b^2 * \coth(c + d * x)^3) / (3 * d) - (b * \coth(c + d * x) * (2 * a + b)) / d$

3.5 $\int \frac{1}{a+b \coth^2(c+dx)} dx$

Optimal result	68
Rubi [A] (verified)	68
Mathematica [A] (verified)	69
Maple [A] (verified)	69
Fricas [B] (verification not implemented)	70
Sympy [B] (verification not implemented)	71
Maxima [A] (verification not implemented)	71
Giac [A] (verification not implemented)	72
Mupad [B] (verification not implemented)	72

Optimal result

Integrand size = 14, antiderivative size = 46

$$\int \frac{1}{a+b \coth^2(c+dx)} dx = \frac{x}{a+b} - \frac{\sqrt{b} \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{\sqrt{a}(a+b)d}$$

[Out] $x/(a+b) - \arctan(a^{1/2} \tanh(dx+c)/b^{1/2}) * b^{1/2} / (a+b) / d / a^{1/2}$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3741, 3756, 211}

$$\int \frac{1}{a+b \coth^2(c+dx)} dx = \frac{x}{a+b} - \frac{\sqrt{b} \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{\sqrt{ad}(a+b)}$$

[In] `Int[(a + b*Coth[c + d*x]^2)^(-1), x]`

[Out] $x/(a+b) - (\text{Sqrt}[b] * \text{ArcTan}[(\text{Sqrt}[a] * \text{Tanh}[c + d*x]) / \text{Sqrt}[b]]) / (\text{Sqrt}[a] * (a + b) * d)$

Rule 211

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

Rule 3741

`Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^(-1), x_Symbol] := Simp[x/(a - b), x] - Dist[b/(a - b), Int[Sec[e + f*x]^2/(a + b*Tan[e + f*x]^2), x], x]`

```
;/ FreeQ[{a, b, e, f}, x] && NeQ[a, b]
```

Rule 3756

```
Int[sec[(e_.) + (f_.)*(x_.)]^(m_)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_.)])^(n_))^(p_.), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/(c^(m - 1)*f), Subst[Int[(c^2 + ff^2*x^2)^(m/2 - 1)*(a + b*(ff*x)^n)^p, x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && IntegerQ[m/2] && (IntegerQ[n, p] || IGtQ[m, 0] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{x}{a+b} - \frac{b \int \frac{\operatorname{csch}^2(c+dx)}{a+b \operatorname{coth}^2(c+dx)} dx}{a+b} \\ &= \frac{x}{a+b} + \frac{b \operatorname{Subst}\left(\int \frac{1}{a+bx^2} dx, x, \operatorname{coth}(c+dx)\right)}{(a+b)d} \\ &= \frac{x}{a+b} - \frac{\sqrt{b} \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{\sqrt{a}(a+b)d} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.02

$$\int \frac{1}{a+b \operatorname{coth}^2(c+dx)} dx = \frac{-\frac{\sqrt{b} \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{\sqrt{a}} + \operatorname{arctanh}(\tanh(c+dx))}{(a+b)d}$$

```
[In] Integrate[(a + b*Coth[c + d*x]^2)^(-1),x]
```

```
[Out] (-((Sqrt[b]*ArcTan[(Sqrt[a]*Tanh[c + d*x])/Sqrt[b]])/Sqrt[a]) + ArcTanh[Tanh[c + d*x]])/((a + b)*d)
```

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.54

method	result	size
derivativedivides	$\frac{b \arctan\left(\frac{b \coth(dx+c)}{\sqrt{ab}}\right)}{(a+b)\sqrt{ab}} - \frac{\ln(\coth(dx+c)-1)}{2a+2b} + \frac{\ln(\coth(dx+c)+1)}{2a+2b}$	71
default	$\frac{b \arctan\left(\frac{b \coth(dx+c)}{\sqrt{ab}}\right)}{(a+b)\sqrt{ab}} - \frac{\ln(\coth(dx+c)-1)}{2a+2b} + \frac{\ln(\coth(dx+c)+1)}{2a+2b}$	71
risch	$\frac{x}{a+b} + \frac{\sqrt{-ab} \ln\left(\frac{e^{2dx+2c} - 2\sqrt{-ab} + a - b}{a+b}\right)}{2a(a+b)d} - \frac{\sqrt{-ab} \ln\left(\frac{e^{2dx+2c} + 2\sqrt{-ab} - a + b}{a+b}\right)}{2a(a+b)d}$	108

[In] `int(1/(a+coth(d*x+c)^2*b),x,method=_RETURNVERBOSE)`

[Out] `1/d*(b/(a+b)/(a*b)^(1/2)*arctan(b*coth(d*x+c)/(a*b)^(1/2))-1/(2*a+2*b)*ln(c
oth(d*x+c)-1)+1/(2*a+2*b)*ln(coth(d*x+c)+1))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 81 vs. 2(38) = 76.

Time = 0.28 (sec) , antiderivative size = 488, normalized size of antiderivative = 10.61

$$\int \frac{1}{a + b \coth^2(c + dx)} dx$$

$$= \left[\frac{2 dx + \sqrt{-\frac{b}{a}} \log\left(\frac{(a^2 + 2ab + b^2) \cosh(dx+c)^4 + 4(a^2 + 2ab + b^2) \cosh(dx+c) \sinh(dx+c)^3 + (a^2 + 2ab + b^2) \sinh(dx+c)^4 - 2(a^2 - b^2) \cosh(dx+c) \sinh(dx+c)^2 + (a^2 - b^2) \sinh(dx+c)^2}{(a+b) \cosh(dx+c)^4 + 4(a+b) \cosh(dx+c) \sinh(dx+c)^3 + (a+b) \sinh(dx+c)^4}\right)}{(a+b) \cosh(dx+c)^4 + 4(a+b) \cosh(dx+c) \sinh(dx+c)^3 + (a+b) \sinh(dx+c)^4} \right]$$

[In] `integrate(1/(a+b*coth(d*x+c)^2),x, algorithm="fricas")`

[Out] `[1/2*(2*d*x + sqrt(-b/a)*log(((a^2 + 2*a*b + b^2)*cosh(d*x + c)^4 + 4*(a^2 + 2*a*b + b^2)*cosh(d*x + c)*sinh(d*x + c)^3 + (a^2 + 2*a*b + b^2)*sinh(d*x + c)^4 - 2*(a^2 - b^2)*cosh(d*x + c)^2 + 2*(3*(a^2 + 2*a*b + b^2)*cosh(d*x + c)^2 - a^2 + b^2)*sinh(d*x + c)^2 + a^2 - 6*a*b + b^2 + 4*((a^2 + 2*a*b + b^2)*cosh(d*x + c)^3 - (a^2 - b^2)*cosh(d*x + c))*sinh(d*x + c) - 4*((a^2 + a*b)*cosh(d*x + c)^2 + 2*(a^2 + a*b)*cosh(d*x + c)*sinh(d*x + c) + (a^2 + a*b)*sinh(d*x + c)^2 - a^2 + a*b)*sqrt(-b/a)))/((a + b)*cosh(d*x + c)^4 + 4*(a + b)*cosh(d*x + c)*sinh(d*x + c)^3 + (a + b)*sinh(d*x + c)^4 - 2*(a - b)*cosh(d*x + c)^2 + 2*(3*(a + b)*cosh(d*x + c)^2 - a + b)*sinh(d*x + c)^2 + 4*((a + b)*cosh(d*x + c)^3 - (a - b)*cosh(d*x + c))*sinh(d*x + c) + a + b)))/((a + b)*d), (d*x - sqrt(b/a)*arctan(1/2*((a + b)*cosh(d*x + c)^2 + 2*(a + b)*cosh(d*x + c)*sinh(d*x + c) + (a + b)*sinh(d*x + c)^2 - a + b)*sqrt(b/a/b)))/((a + b)*d)]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 253 vs. 2(37) = 74.

Time = 2.49 (sec) , antiderivative size = 253, normalized size of antiderivative = 5.50

$$\int \frac{1}{a + b \coth^2(c + dx)} dx$$

$$= \begin{cases} \frac{\infty x}{\coth^2(c)} & \text{for } a = 0 \wedge b = 0 \wedge d = 0 \\ \frac{x - \frac{\tanh(c+dx)}{d}}{b} & \text{for } a = 0 \\ -\frac{dx \tanh^2(c+dx)}{2bd \tanh^2(c+dx) - 2bd} + \frac{dx}{2bd \tanh^2(c+dx) - 2bd} - \frac{\tanh(c+dx)}{2bd \tanh^2(c+dx) - 2bd} & \text{for } a = -b \\ \frac{x}{a} & \text{for } b = 0 \\ \frac{x}{a + b \coth^2(c)} & \text{for } d = 0 \\ \frac{2adx\sqrt{-\frac{b}{a}}}{2a^2d\sqrt{-\frac{b}{a}} + 2abd\sqrt{-\frac{b}{a}}} - \frac{b \log\left(-\sqrt{-\frac{b}{a}} + \tanh(c+dx)\right)}{2a^2d\sqrt{-\frac{b}{a}} + 2abd\sqrt{-\frac{b}{a}}} + \frac{b \log\left(\sqrt{-\frac{b}{a}} + \tanh(c+dx)\right)}{2a^2d\sqrt{-\frac{b}{a}} + 2abd\sqrt{-\frac{b}{a}}} & \text{otherwise} \end{cases}$$

[In] integrate(1/(a+b*coth(d*x+c)**2),x)

[Out] Piecewise((zoo*x/coth(c)**2, Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), ((x - tanh(c + d*x)/d)/b, Eq(a, 0)), (-d*x*tanh(c + d*x)**2/(2*b*d*tanh(c + d*x)**2 - 2*b*d) + d*x/(2*b*d*tanh(c + d*x)**2 - 2*b*d) - tanh(c + d*x)/(2*b*d*tanh(c + d*x)**2 - 2*b*d), Eq(a, -b)), (x/a, Eq(b, 0)), (x/(a + b*coth(c)**2), Eq(d, 0)), (2*a*d*x*sqrt(-b/a)/(2*a**2*d*sqrt(-b/a) + 2*a*b*d*sqrt(-b/a)) - b*log(-sqrt(-b/a) + tanh(c + d*x))/(2*a**2*d*sqrt(-b/a) + 2*a*b*d*sqrt(-b/a)) + b*log(sqrt(-b/a) + tanh(c + d*x))/(2*a**2*d*sqrt(-b/a) + 2*a*b*d*sqrt(-b/a)), True))

Maxima [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.22

$$\int \frac{1}{a + b \coth^2(c + dx)} dx = \frac{b \arctan\left(\frac{(a+b)e^{(-2dx-2c)-a+b}}{2\sqrt{ab}}\right)}{\sqrt{ab}(a+b)d} + \frac{dx + c}{(a+b)d}$$

[In] integrate(1/(a+b*coth(d*x+c)^2),x, algorithm="maxima")

[Out] b*arctan(1/2*((a + b)*e^(-2*d*x - 2*c) - a + b)/sqrt(a*b))/(sqrt(a*b)*(a + b)*d) + (d*x + c)/((a + b)*d)

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.41

$$\int \frac{1}{a + b \coth^2(c + dx)} dx = -\frac{\frac{b \arctan\left(\frac{ae^{(2dx+2c)} + be^{(2dx+2c)} - a + b}{2\sqrt{ab}}\right)}{\sqrt{ab}(a+b)} - \frac{dx+c}{a+b}}{d}$$

[In] integrate(1/(a+b*coth(d*x+c)^2),x, algorithm="giac")

[Out] -(b*arctan(1/2*(a*e^(2*d*x + 2*c) + b*e^(2*d*x + 2*c) - a + b)/sqrt(a*b)))/(sqrt(a*b)*(a + b)) - (d*x + c)/(a + b))/d

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\int \frac{1}{a + b \coth^2(c + dx)} dx = \frac{x}{a + b} + \frac{b \operatorname{atan}\left(\frac{b \coth(c + dx)}{\sqrt{ab}}\right)}{d \sqrt{ab} (a + b)}$$

[In] int(1/(a + b*coth(c + d*x)^2),x)

[Out] x/(a + b) + (b*atan((b*coth(c + d*x))/(a*b)^(1/2)))/(d*(a*b)^(1/2)*(a + b))

3.6 $\int \frac{1}{(a+b \coth^2(c+dx))^2} dx$

Optimal result	73
Rubi [A] (verified)	73
Mathematica [A] (verified)	75
Maple [A] (verified)	75
Fricas [B] (verification not implemented)	76
Sympy [F(-1)]	77
Maxima [B] (verification not implemented)	77
Giac [B] (verification not implemented)	78
Mupad [B] (verification not implemented)	78

Optimal result

Integrand size = 14, antiderivative size = 89

$$\int \frac{1}{(a+b \coth^2(c+dx))^2} dx = \frac{x}{(a+b)^2} - \frac{\sqrt{b}(3a+b) \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{2a^{3/2}(a+b)^2 d} + \frac{b \coth(c+dx)}{2a(a+b)d(a+b \coth^2(c+dx))}$$

[Out] $x/(a+b)^2 + 1/2*b*\coth(d*x+c)/a/(a+b)/d/(a+b*\coth(d*x+c)^2) - 1/2*(3*a+b)*\arctan(\sqrt{a}*\tanh(d*x+c)/\sqrt{b})/b^{1/2}/a^{3/2}/(a+b)^2/d$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {3742, 425, 536, 212, 211}

$$\int \frac{1}{(a+b \coth^2(c+dx))^2} dx = -\frac{\sqrt{b}(3a+b) \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{2a^{3/2}d(a+b)^2} + \frac{b \coth(c+dx)}{2ad(a+b)(a+b \coth^2(c+dx))} + \frac{x}{(a+b)^2}$$

[In] $\text{Int}[(a + b*\text{Coth}[c + d*x]^2)^{-2}, x]$

[Out] $x/(a+b)^2 - (\text{Sqrt}[b]*(3*a+b)*\text{ArcTan}[(\text{Sqrt}[a]*\text{Tanh}[c+d*x])/ \text{Sqrt}[b]])/(2*a^{3/2}*(a+b)^2*d) + (b*\text{Coth}[c+d*x])/(2*a*(a+b)*d*(a+b*\text{Coth}[c+d*x]^2))$

Rule 211

$\text{Int}[(a_ + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] \text{ ; FreeQ}\{a, b\}, x\} \ \&\& \ \text{PosQ}[a/b]$

Rule 212

$\text{Int}[(a_ + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] \text{ ; FreeQ}\{a, b\}, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 425

$\text{Int}[(a_ + (b_.)*(x_)^{n_})^{p_}*((c_ + (d_.)*(x_)^{n_})^{q_}), x_Symbol] \rightarrow \text{Simp}[(-b)*x*(a + b*x^n)^{p+1}*((c + d*x^n)^{q+1}/(a*n*(p+1)*(b*c - a*d))], x] + \text{Dist}[1/(a*n*(p+1)*(b*c - a*d)), \text{Int}[(a + b*x^n)^{p+1}*(c + d*x^n)^q*\text{Simp}[b*c + n*(p+1)*(b*c - a*d) + d*b*(n*(p+q+2) + 1)*x^n, x], x], x] \text{ ; FreeQ}\{a, b, c, d, n, q\}, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ !(\ !\text{IntegerQ}[p] \ \&\& \ \text{IntegerQ}[q] \ \&\& \ \text{LtQ}[q, -1]) \ \&\& \ \text{IntBinomialQ}[a, b, c, d, n, p, q, x]$

Rule 536

$\text{Int}[(e_ + (f_.)*(x_)^{n_})/((a_ + (b_.)*(x_)^{n_})*((c_ + (d_.)*(x_)^{n_}))), x_Symbol] \rightarrow \text{Dist}[(b*e - a*f)/(b*c - a*d), \text{Int}[1/(a + b*x^n), x], x] - \text{Dist}[(d*e - c*f)/(b*c - a*d), \text{Int}[1/(c + d*x^n), x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, n\}, x\}$

Rule 3742

$\text{Int}[(a_ + (b_.)*((c_)*\tan[(e_ + (f_.)*(x_)]))^{n_})^{p_}, x_Symbol] \rightarrow \text{With}\{\{ff = \text{FreeFactors}[\text{Tan}[e + f*x], x]\}, \text{Dist}[c*(ff/f), \text{Subst}[\text{Int}[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(\text{Tan}[e + f*x]/ff)], x]\} \text{ ; FreeQ}\{a, b, c, e, f, n, p\}, x\} \ \&\& \ (\text{IntegersQ}[n, p] \ || \ \text{IGtQ}[p, 0] \ || \ \text{EqQ}[n^2, 4] \ || \ \text{EqQ}[n^2, 16])$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\text{Subst}\left(\int \frac{1}{(1-x^2)(a+bx^2)^2} dx, x, \coth(c+dx)\right)}{d} \\ &= \frac{b \coth(c+dx)}{2a(a+b)d(a+b \coth^2(c+dx))} - \frac{\text{Subst}\left(\int \frac{b-2(a+b)+bx^2}{(1-x^2)(a+bx^2)} dx, x, \coth(c+dx)\right)}{2a(a+b)d} \end{aligned}$$

$$\begin{aligned}
&= \frac{b \coth(c + dx)}{2a(a + b)d(a + b \coth^2(c + dx))} + \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \coth(c + dx)\right)}{(a + b)^2 d} \\
&\quad + \frac{(b(3a + b))\text{Subst}\left(\int \frac{1}{a+bx^2} dx, x, \coth(c + dx)\right)}{2a(a + b)^2 d} \\
&= \frac{x}{(a + b)^2} - \frac{\sqrt{b}(3a + b) \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{2a^{3/2}(a + b)^2 d} + \frac{b \coth(c + dx)}{2a(a + b)d(a + b \coth^2(c + dx))}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.63 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.09

$$\begin{aligned}
&\int \frac{1}{(a + b \coth^2(c + dx))^2} dx \\
&= \frac{\frac{\sqrt{b}(3a+b) \arctan\left(\frac{\sqrt{b} \coth(c+dx)}{\sqrt{a}}\right)}{a^{3/2}} + \frac{b(a+b) \coth(c+dx)}{a(a+b \coth^2(c+dx))} - \log(1 - \coth(c + dx)) + \log(1 + \coth(c + dx))}{2(a + b)^2 d}
\end{aligned}$$

[In] Integrate[(a + b*Coth[c + d*x]^2)^(-2), x]

[Out] ((Sqrt[b]*(3*a + b)*ArcTan[(Sqrt[b]*Coth[c + d*x])/Sqrt[a]])/a^(3/2) + (b*(a + b)*Coth[c + d*x])/(a*(a + b*Coth[c + d*x]^2)) - Log[1 - Coth[c + d*x]] + Log[1 + Coth[c + d*x]])/(2*(a + b)^2*d)

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.16

method	result
derivativedivides	$ \frac{-\frac{\ln(\coth(dx+c)-1)}{2(a+b)^2} + \frac{\ln(\coth(dx+c)+1)}{2(a+b)^2} + \frac{b\left(\frac{(a+b)\coth(dx+c)}{2a(a+\coth(dx+c)^2b)} + \frac{(3a+b)\arctan\left(\frac{b\coth(dx+c)}{\sqrt{ab}}\right)}{2a\sqrt{ab}}\right)}{(a+b)^2}}{d} $
default	$ \frac{-\frac{\ln(\coth(dx+c)-1)}{2(a+b)^2} + \frac{\ln(\coth(dx+c)+1)}{2(a+b)^2} + \frac{b\left(\frac{(a+b)\coth(dx+c)}{2a(a+\coth(dx+c)^2b)} + \frac{(3a+b)\arctan\left(\frac{b\coth(dx+c)}{\sqrt{ab}}\right)}{2a\sqrt{ab}}\right)}{(a+b)^2}}{d} $
risch	$ \frac{x}{a^2+2ab+b^2} + \frac{b(e^{2dx+2c}a-b)e^{2dx+2c}-a-b}{da(a+b)^2(ae^{4dx+4c}+be^{4dx+4c}-2e^{2dx+2c}a+2be^{2dx+2c}+a+b)} + \frac{3\sqrt{-ab}\ln\left(\frac{e^{2dx+2c}-2\sqrt{-ab}+a-b}{a+b}\right)}{4a(a+b)^2d} $

[In] int(1/(a+coth(d*x+c)^2*b)^2,x,method=_RETURNVERBOSE)

[Out] 1/d*(-1/2/(a+b)^2*ln(coth(d*x+c)-1)+1/2/(a+b)^2*ln(coth(d*x+c)+1)+b/(a+b)^2*(1/2*(a+b)/a*coth(d*x+c)/(a+coth(d*x+c)^2*b)+1/2*(3*a+b)/a/(a*b)^(1/2)*arctan(b*coth(d*x+c)/(a*b)^(1/2))))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 814 vs. 2(77) = 154.

Time = 0.33 (sec) , antiderivative size = 1952, normalized size of antiderivative = 21.93

$$\int \frac{1}{(a + b \coth^2(c + dx))^2} dx = \text{Too large to display}$$

[In] integrate(1/(a+b*coth(d*x+c)^2)^2,x, algorithm="fricas")

[Out] [1/4*(4*(a^2 + a*b)*d*x*cosh(d*x + c)^4 + 16*(a^2 + a*b)*d*x*cosh(d*x + c)*sinh(d*x + c)^3 + 4*(a^2 + a*b)*d*x*sinh(d*x + c)^4 + 4*(a^2 + a*b)*d*x - 4*(2*(a^2 - a*b)*d*x - a*b + b^2)*cosh(d*x + c)^2 + 4*(6*(a^2 + a*b)*d*x*cosh(d*x + c)^2 - 2*(a^2 - a*b)*d*x + a*b - b^2)*sinh(d*x + c)^2 + ((3*a^2 + 4*a*b + b^2)*cosh(d*x + c)^4 + 4*(3*a^2 + 4*a*b + b^2)*cosh(d*x + c)*sinh(d*x + c)^3 + (3*a^2 + 4*a*b + b^2)*sinh(d*x + c)^4 - 2*(3*a^2 - 2*a*b - b^2)*cosh(d*x + c)^2 + 2*(3*(3*a^2 + 4*a*b + b^2)*cosh(d*x + c)^2 - 3*a^2 + 2*a*b + b^2)*sinh(d*x + c)^2 + 3*a^2 + 4*a*b + b^2 + 4*((3*a^2 + 4*a*b + b^2)*cosh(d*x + c)^3 - (3*a^2 - 2*a*b - b^2)*cosh(d*x + c))*sinh(d*x + c))*sqrt(-b/a)*log(((a^2 + 2*a*b + b^2)*cosh(d*x + c)^4 + 4*(a^2 + 2*a*b + b^2)*cosh(d*x + c)*sinh(d*x + c)^3 + (a^2 + 2*a*b + b^2)*sinh(d*x + c)^4 - 2*(a^2 - b^2)*cosh(d*x + c)^2 + 2*(3*(a^2 + 2*a*b + b^2)*cosh(d*x + c)^2 - a^2 + b^2)*sinh(d*x + c)^2 + a^2 - 6*a*b + b^2 + 4*((a^2 + 2*a*b + b^2)*cosh(d*x + c)^3 - (a^2 - b^2)*cosh(d*x + c))*sinh(d*x + c) - 4*((a^2 + a*b)*cosh(d*x + c)^2 + 2*(a^2 + a*b)*cosh(d*x + c)*sinh(d*x + c) + (a^2 + a*b)*sinh(d*x + c)^2 - a^2 + a*b)*sqrt(-b/a))/((a + b)*cosh(d*x + c)^4 + 4*(a + b)*cosh(d*x + c)*sinh(d*x + c)^3 + (a + b)*sinh(d*x + c)^4 - 2*(a - b)*cosh(d*x + c)^2 + 2*(3*(a + b)*cosh(d*x + c)^2 - a + b)*sinh(d*x + c)^2 + 4*((a + b)*cosh(d*x + c)^3 - (a - b)*cosh(d*x + c))*sinh(d*x + c) + a + b) - 4*a*b - 4*b^2 + 8*(2*(a^2 + a*b)*d*x*cosh(d*x + c)^3 - (2*(a^2 - a*b)*d*x - a*b + b^2)*cosh(d*x + c))*sinh(d*x + c))/((a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*d*cosh(d*x + c)^4 + 4*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*d*cosh(d*x + c)*sinh(d*x + c)^3 + (a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*d*sinh(d*x + c)^4 - 2*(a^4 + a^3*b - a^2*b^2 - a*b^3)*d*cosh(d*x + c)^2 + 2*(3*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*d*cosh(d*x + c)^2 - (a^4 + a^3*b - a^2*b^2 - a*b^3)*d)*sinh(d*x + c)^2 + (a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*d + 4*((a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*d*cosh(d*x + c)^3 - (a^4 + a^3*b - a^2*b^2 - a*b^3)*d*cosh(d*x + c))*sinh(d*x + c)), 1/2*(2*(a^2 + a*b)*d*x*cosh(d*x + c)^4 + 8*(a^2 + a*b)*d*x*cosh(d*x + c)*sinh(d*x + c)^3 + 2*(a^2 + a*b)*d*x*sinh(d*x + c)^4 + 2*(a^2 + a*b)*d*x - 2*(2*(a^2 - a*b)*d*x - a*b + b^2)*cosh(d*x + c)^2 + 2*(6*(a^2 + a*b)*d*x*cosh(d*x + c)^2 - 2*(a^2 - a*b)*d*x + a*b - b^2)*sinh(d*x + c)^2 - ((3*a^2 + 4*a*b + b^2)*cosh(d*x + c)^4 + 4*(3*a^2 + 4*a*b + b^2)*cosh(d*x + c)*sinh(d*x + c)^3 + (3*a^2 + 4*a*b + b^2)*sinh(d*x + c)^4 - 2*(3*a^2 - 2*a*b - b^2)*cosh(d*x + c)^2 + 2*(3*(3*a^2 + 4*a*b + b^2)*cosh(d*x + c)^2 - 3*a^2 + 2*a*b + b^2)*sinh(d*x + c)^2 + 3*a^2 + 4*a*b + b^2 + 4*((3*a^2

+ 4*a*b + b^2)*cosh(d*x + c)^3 - (3*a^2 - 2*a*b - b^2)*cosh(d*x + c)*sinh(d*x + c))*sqrt(b/a)*arctan(1/2*((a + b)*cosh(d*x + c)^2 + 2*(a + b)*cosh(d*x + c)*sinh(d*x + c) + (a + b)*sinh(d*x + c)^2 - a + b)*sqrt(b/a)/b) - 2*a*b - 2*b^2 + 4*(2*(a^2 + a*b)*d*x*cosh(d*x + c)^3 - (2*(a^2 - a*b)*d*x - a*b + b^2)*cosh(d*x + c))*sinh(d*x + c))/((a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*d*cosh(d*x + c)^4 + 4*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*d*cosh(d*x + c)*sinh(d*x + c)^3 + (a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*d*sinh(d*x + c)^4 - 2*(a^4 + a^3*b - a^2*b^2 - a*b^3)*d*cosh(d*x + c)^2 + 2*(3*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*d*cosh(d*x + c)^2 - (a^4 + a^3*b - a^2*b^2 - a*b^3)*d)*sinh(d*x + c)^2 + (a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*d + 4*((a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*d*cosh(d*x + c)^3 - (a^4 + a^3*b - a^2*b^2 - a*b^3)*d*cosh(d*x + c))*sinh(d*x + c))]

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \coth^2(c + dx))^2} dx = \text{Timed out}$$

[In] integrate(1/(a+b*coth(d*x+c)**2)**2,x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 207 vs. 2(77) = 154.

Time = 0.32 (sec) , antiderivative size = 207, normalized size of antiderivative = 2.33

$$\int \frac{1}{(a + b \coth^2(c + dx))^2} dx = \frac{(3ab + b^2) \arctan\left(\frac{(a+b)e^{(-2dx-2c)} - a+b}{2\sqrt{ab}}\right)}{2(a^3 + 2a^2b + ab^2)\sqrt{abd}} + \frac{ab + b^2 - (ab - b^2)e^{(-2dx-2c)}}{(a^4 + 3a^3b + 3a^2b^2 + ab^3 - 2(a^4 + a^3b - a^2b^2 - ab^3)e^{(-2dx-2c)} + (a^4 + 3a^3b + 3a^2b^2 + ab^3)e^{(-4dx-4c)})} + \frac{dx + c}{(a^2 + 2ab + b^2)d}$$

[In] integrate(1/(a+b*coth(d*x+c)^2)^2,x, algorithm="maxima")

[Out] 1/2*(3*a*b + b^2)*arctan(1/2*((a + b)*e^(-2*d*x - 2*c) - a + b)/sqrt(a*b))/((a^3 + 2*a^2*b + a*b^2)*sqrt(a*b)*d) + (a*b + b^2 - (a*b - b^2)*e^(-2*d*x - 2*c))/((a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3 - 2*(a^4 + a^3*b - a^2*b^2 - a*b^3)*e^(-2*d*x - 2*c) + (a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*e^(-4*d*x - 4*c))*d) + (d*x + c)/((a^2 + 2*a*b + b^2)*d)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 198 vs. 2(77) = 154.

Time = 0.29 (sec) , antiderivative size = 198, normalized size of antiderivative = 2.22

$$\int \frac{1}{(a + b \coth^2(c + dx))^2} dx = \frac{(3ab + b^2) \arctan\left(\frac{ae^{(2dx+2c)} + be^{(2dx+2c)} - a + b}{2\sqrt{ab}}\right)}{(a^3 + 2a^2b + ab^2)\sqrt{ab}} - \frac{2(dx+c)}{a^2 + 2ab + b^2} - \frac{2(ab e^{(2dx+2c)} - b^2 e^{(2dx+2c)} - ab - b^2)}{(a^3 + 2a^2b + ab^2)(ae^{(4dx+4c)} + be^{(4dx+4c)} - 2ae^{(2dx+2c)} + 2be^{(2dx+2c)})}$$

[In] integrate(1/(a+b*coth(d*x+c)^2)^2,x, algorithm="giac")

[Out] -1/2*((3*a*b + b^2)*arctan(1/2*(a*e^(2*d*x + 2*c) + b*e^(2*d*x + 2*c) - a + b)/sqrt(a*b))/((a^3 + 2*a^2*b + a*b^2)*sqrt(a*b)) - 2*(d*x + c)/(a^2 + 2*a*b + b^2) - 2*(a*b*e^(2*d*x + 2*c) - b^2*e^(2*d*x + 2*c) - a*b - b^2)/((a^3 + 2*a^2*b + a*b^2)*(a*e^(4*d*x + 4*c) + b*e^(4*d*x + 4*c) - 2*a*e^(2*d*x + 2*c) + 2*b*e^(2*d*x + 2*c) + a + b))/d

Mupad [B] (verification not implemented)

Time = 2.31 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.24

$$\int \frac{1}{(a + b \coth^2(c + dx))^2} dx = \frac{\frac{ax}{(a+b)^2} + \frac{bx \coth(c+dx)^2}{(a+b)^2} + \frac{b \coth(c+dx)}{2ad(a+b)}}{b \coth(c + dx)^2 + a} + \frac{\operatorname{atan}\left(\frac{b \coth(c+dx)}{\sqrt{ab}}\right) (b^2 + 3ab)}{\sqrt{ab} (2a^3d + ab(4ad + 2bd))}$$

[In] int(1/(a + b*coth(c + d*x)^2)^2,x)

[Out] ((a*x)/(a + b)^2 + (b*x*coth(c + d*x)^2)/(a + b)^2 + (b*coth(c + d*x))/(2*a*d*(a + b)))/(a + b*coth(c + d*x)^2) + (atan((b*coth(c + d*x))/(a*b)^(1/2)))*(3*a*b + b^2))/((a*b)^(1/2)*(2*a^3*d + a*b*(4*a*d + 2*b*d)))

3.7 $\int \frac{1}{(a+b \coth^2(c+dx))^3} dx$

Optimal result	79
Rubi [A] (verified)	79
Mathematica [A] (verified)	81
Maple [A] (verified)	82
Fricas [B] (verification not implemented)	82
Sympy [F(-1)]	83
Maxima [B] (verification not implemented)	83
Giac [B] (verification not implemented)	84
Mupad [B] (verification not implemented)	84

Optimal result

Integrand size = 14, antiderivative size = 142

$$\int \frac{1}{(a+b \coth^2(c+dx))^3} dx = \frac{x}{(a+b)^3} - \frac{\sqrt{b}(15a^2+10ab+3b^2) \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{8a^{5/2}(a+b)^3 d}$$

$$+ \frac{b \coth(c+dx)}{4a(a+b)d (a+b \coth^2(c+dx))^2}$$

$$+ \frac{b(7a+3b) \coth(c+dx)}{8a^2(a+b)^2 d (a+b \coth^2(c+dx))}$$

[Out] x/(a+b)^3+1/4*b*coth(d*x+c)/a/(a+b)/d/(a+b*coth(d*x+c)^2)^2+1/8*b*(7*a+3*b)*coth(d*x+c)/a^2/(a+b)^2/d/(a+b*coth(d*x+c)^2)-1/8*(15*a^2+10*a*b+3*b^2)*arctan(a^(1/2)*tanh(d*x+c)/b^(1/2))*b^(1/2)/a^(5/2)/(a+b)^3/d

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3742, 425, 541, 536, 212, 211}

$$\int \frac{1}{(a+b \coth^2(c+dx))^3} dx = \frac{b(7a+3b) \coth(c+dx)}{8a^2 d (a+b)^2 (a+b \coth^2(c+dx))}$$

$$- \frac{\sqrt{b}(15a^2+10ab+3b^2) \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{8a^{5/2} d (a+b)^3}$$

$$+ \frac{b \coth(c+dx)}{4ad(a+b) (a+b \coth^2(c+dx))^2} + \frac{x}{(a+b)^3}$$

[In] Int[(a + b*Coth[c + d*x]^2)^(-3), x]

[Out] x/(a + b)^3 - (Sqrt[b]*(15*a^2 + 10*a*b + 3*b^2)*ArcTan[(Sqrt[a]*Tanh[c + d*x])/Sqrt[b]])/(8*a^(5/2)*(a + b)^3*d) + (b*Coth[c + d*x])/(4*a*(a + b)*d*(a + b*Coth[c + d*x]^2)^2) + (b*(7*a + 3*b)*Coth[c + d*x])/(8*a^2*(a + b)^2*d*(a + b*Coth[c + d*x]^2))

Rule 211

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 425

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-b)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(p + 1)*(b*c - a*d))), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 536

Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 541

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Simp[(-b*e - a*f)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(b*c - a*d)*(p + 1))), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rule 3742

Int[((a_) + (b_.)*((c_.)*tan[e_.] + (f_.)*(x_)))^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(

$\text{ff*x})^n)^p/(c^2 + \text{ff}^2*x^2), x], x, c*(\text{Tan}[e + f*x]/\text{ff}), x]] /; \text{FreeQ}\{a, b, c, e, f, n, p\}, x] \&\& (\text{IntegersQ}[n, p] \parallel \text{IGtQ}[p, 0] \parallel \text{EqQ}[n^2, 4] \parallel \text{E} \\ \text{qQ}[n^2, 16])$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int \frac{1}{(1-x^2)(a+bx^2)^3} dx, x, \coth(c+dx)\right)}{d} \\
 &= \frac{b \coth(c+dx)}{4a(a+b)d (a+b \coth^2(c+dx))^2} - \frac{\text{Subst}\left(\int \frac{b-4(a+b)+3bx^2}{(1-x^2)(a+bx^2)^2} dx, x, \coth(c+dx)\right)}{4a(a+b)d} \\
 &= \frac{b \coth(c+dx)}{4a(a+b)d (a+b \coth^2(c+dx))^2} + \frac{b(7a+3b) \coth(c+dx)}{8a^2(a+b)^2d (a+b \coth^2(c+dx))} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{8a^2+7ab+3b^2-b(7a+3b)x^2}{(1-x^2)(a+bx^2)} dx, x, \coth(c+dx)\right)}{8a^2(a+b)^2d} \\
 &= \frac{b \coth(c+dx)}{4a(a+b)d (a+b \coth^2(c+dx))^2} + \frac{b(7a+3b) \coth(c+dx)}{8a^2(a+b)^2d (a+b \coth^2(c+dx))} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \coth(c+dx)\right)}{(a+b)^3d} \\
 &\quad + \frac{(b(15a^2+10ab+3b^2)) \text{Subst}\left(\int \frac{1}{a+bx^2} dx, x, \coth(c+dx)\right)}{8a^2(a+b)^3d} \\
 &= \frac{x}{(a+b)^3} - \frac{\sqrt{b}(15a^2+10ab+3b^2) \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{8a^{5/2}(a+b)^3d} \\
 &\quad + \frac{b \coth(c+dx)}{4a(a+b)d (a+b \coth^2(c+dx))^2} + \frac{b(7a+3b) \coth(c+dx)}{8a^2(a+b)^2d (a+b \coth^2(c+dx))}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.04

$$\begin{aligned}
 &\int \frac{1}{(a+b \coth^2(c+dx))^3} dx \\
 &= \frac{\sqrt{b}(15a^2+10ab+3b^2) \arctan\left(\frac{\sqrt{b} \coth(c+dx)}{\sqrt{a}}\right)}{a^{5/2}} + \frac{2b(a+b)^2 \coth(c+dx)}{a(a+b \coth^2(c+dx))^2} + \frac{b(a+b)(7a+3b) \coth(c+dx)}{a^2(a+b \coth^2(c+dx))} - 4 \log(1 - \coth(c+dx)) + \dots \\
 &\hspace{15em} 8(a+b)^3d
 \end{aligned}$$

[In] Integrate[(a + b*Coth[c + d*x]^2)^(-3), x]

```
[Out] ((Sqrt[b]*(15*a^2 + 10*a*b + 3*b^2)*ArcTan[(Sqrt[b]*Coth[c + d*x])/Sqrt[a]]
)/a^(5/2) + (2*b*(a + b)^2*Coth[c + d*x])/(a*(a + b*Coth[c + d*x]^2)^2) + (
b*(a + b)*(7*a + 3*b)*Coth[c + d*x])/(a^2*(a + b*Coth[c + d*x]^2)) - 4*Log[
1 - Coth[c + d*x]] + 4*Log[1 + Coth[c + d*x]])/(8*(a + b)^3*d)
```

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.10

method	result
derivativedivides	$\frac{\frac{\ln(\coth(dx+c)+1)}{2(a+b)^3} - \frac{\ln(\coth(dx+c)-1)}{2(a+b)^3} + \frac{b \left(\frac{b(7a^2+10ab+3b^2)\coth(dx+c)^3}{8a^2} + \frac{(9a^2+14ab+5b^2)\coth(dx+c)}{8a} + \frac{(15a^2+10ab+3b^2)\arctan\left(\frac{\sqrt{b}\coth(dx+c)}{\sqrt{a}}\right)}{8a^2\sqrt{b}} \right)}{(a+\coth(dx+c)^2b)^2}}{(a+b)^3}}{d}$
default	$\frac{\frac{\ln(\coth(dx+c)+1)}{2(a+b)^3} - \frac{\ln(\coth(dx+c)-1)}{2(a+b)^3} + \frac{b \left(\frac{b(7a^2+10ab+3b^2)\coth(dx+c)^3}{8a^2} + \frac{(9a^2+14ab+5b^2)\coth(dx+c)}{8a} + \frac{(15a^2+10ab+3b^2)\arctan\left(\frac{\sqrt{b}\coth(dx+c)}{\sqrt{a}}\right)}{8a^2\sqrt{b}} \right)}{(a+\coth(dx+c)^2b)^2}}{(a+b)^3}}{d}$
risch	$\frac{x}{a^3+3a^2b+3ab^2+b^3} + \frac{b(9a^3e^{6dx+6c}-a^2be^{6dx+6c}-13ab^2e^{6dx+6c}-3e^{6dx+6c}b^3-27a^3e^{4dx+4c}+9a^2be^{4dx+4c}-21ab^2e^{4dx+4c}-21a^2be^{2dx+2c}-21ab^2e^{2dx+2c})}{4(ae^{4dx+4c}+be^{4dx+4c}-2e^{2dx+2c})}$

```
[In] int(1/(a+coth(d*x+c)^2*b)^3,x,method=_RETURNVERBOSE)
```

```
[Out] 1/d*(1/2/(a+b)^3*ln(coth(d*x+c)+1)-1/2/(a+b)^3*ln(coth(d*x+c)-1)+b/(a+b)^3*
((1/8*b*(7*a^2+10*a*b+3*b^2)/a^2*coth(d*x+c)^3+1/8*(9*a^2+14*a*b+5*b^2)/a*c
oth(d*x+c))/(a+coth(d*x+c)^2*b)^2+1/8*(15*a^2+10*a*b+3*b^2)/a^2/(a*b)^(1/2)
*arctan(b*coth(d*x+c)/(a*b)^(1/2))))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3592 vs. $2(128) = 256$.

Time = 0.38 (sec) , antiderivative size = 7508, normalized size of antiderivative = 52.87

$$\int \frac{1}{(a + b \coth^2(c + dx))^3} dx = \text{Too large to display}$$

```
[In] integrate(1/(a+b*coth(d*x+c)^2)^3,x, algorithm="fricas")
```

```
[Out] Too large to include
```

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \coth^2(c + dx))^3} dx = \text{Timed out}$$

[In] integrate(1/(a+b*coth(d*x+c)**2)**3,x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 509 vs. 2(128) = 256.

Time = 0.39 (sec) , antiderivative size = 509, normalized size of antiderivative = 3.58

$$\int \frac{1}{(a + b \coth^2(c + dx))^3} dx = \frac{(15 a^2 b + 10 a b^2 + 3 b^3) \arctan\left(\frac{(a+b)e^{(-2 dx - 2c)} - a + b}{2\sqrt{ab}}\right)}{8(a^5 + 3 a^4 b + 3 a^3 b^2 + a^2 b^3)\sqrt{abd}}$$

$$+ \frac{9 a^3 b + 21 a^2 b^2 + 15 a b^3 + 3 b^4 - (27 a^3 b + 13 a^2 b^2 - 23 a^3 b^3 - 9 b^4)e^{(-2 dx - 2c)} + 3(9 a^3 b - 3 a^2 b^2 + 7 a^3 b^3 + 3 b^4)e^{(-4 dx - 4c)} - (9 a^3 b - a^2 b^2 - 13 a^3 b^3 - 3 b^4)e^{(-6 dx - 6c)}}{4(a^7 + 5 a^6 b + 10 a^5 b^2 + 10 a^4 b^3 + 5 a^3 b^4 + a^2 b^5 - 4(a^7 + 3 a^6 b + 2 a^5 b^2 - 2 a^4 b^3 - 3 a^3 b^4 - a^2 b^5)e^{(-2 dx - 2c)} + 2(3 a^7 + 7 a^6 b + 6 a^5 b^2 + 6 a^4 b^3 + 7 a^3 b^4 + 3 a^2 b^5)e^{(-4 dx - 4c)} - 4(a^7 + 3 a^6 b + 2 a^5 b^2 - 2 a^4 b^3 - 3 a^3 b^4 - a^2 b^5)e^{(-6 dx - 6c)} + (a^7 + 5 a^6 b + 10 a^5 b^2 + 10 a^4 b^3 + 5 a^3 b^4 + a^2 b^5)e^{(-8 dx - 8c)})d}$$

$$+ \frac{dx + c}{(a^3 + 3 a^2 b + 3 a b^2 + b^3)d}$$

[In] integrate(1/(a+b*coth(d*x+c)^2)^3,x, algorithm="maxima")

```
[Out] 1/8*(15*a^2*b + 10*a*b^2 + 3*b^3)*arctan(1/2*((a + b)*e^(-2*d*x - 2*c) - a + b)/sqrt(a*b))/((a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3)*sqrt(a*b)*d) + 1/4*(9*a^3*b + 21*a^2*b^2 + 15*a*b^3 + 3*b^4 - (27*a^3*b + 13*a^2*b^2 - 23*a^3*b^3 - 9*b^4)*e^(-2*d*x - 2*c) + 3*(9*a^3*b - 3*a^2*b^2 + 7*a^3*b^3 + 3*b^4)*e^(-4*d*x - 4*c) - (9*a^3*b - a^2*b^2 - 13*a^3*b^3 - 3*b^4)*e^(-6*d*x - 6*c))/((a^7 + 5*a^6*b + 10*a^5*b^2 + 10*a^4*b^3 + 5*a^3*b^4 + a^2*b^5 - 4*(a^7 + 3*a^6*b + 2*a^5*b^2 - 2*a^4*b^3 - 3*a^3*b^4 - a^2*b^5)*e^(-2*d*x - 2*c) + 2*(3*a^7 + 7*a^6*b + 6*a^5*b^2 + 6*a^4*b^3 + 7*a^3*b^4 + 3*a^2*b^5)*e^(-4*d*x - 4*c) - 4*(a^7 + 3*a^6*b + 2*a^5*b^2 - 2*a^4*b^3 - 3*a^3*b^4 - a^2*b^5)*e^(-6*d*x - 6*c) + (a^7 + 5*a^6*b + 10*a^5*b^2 + 10*a^4*b^3 + 5*a^3*b^4 + a^2*b^5)*e^(-8*d*x - 8*c))*d) + (d*x + c)/((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 409 vs. 2(128) = 256.

Time = 0.32 (sec) , antiderivative size = 409, normalized size of antiderivative = 2.88

$$\int \frac{1}{(a + b \coth^2(c + dx))^3} dx = \frac{(15a^2b + 10ab^2 + 3b^3) \arctan\left(\frac{ae^{(2dx+2c)} + be^{(2dx+2c)} - a + b}{2\sqrt{ab}}\right)}{(a^5 + 3a^4b + 3a^3b^2 + a^2b^3)\sqrt{ab}} - \frac{8(dx+c)}{a^3 + 3a^2b + 3ab^2 + b^3} - \frac{2(9a^3be^{(6dx+6c)} - a^2b^2e^{(6dx+6c)} - 13ab^3e^{(6dx+6c)})}{(a^5 + 3a^4b + 3a^3b^2 + a^2b^3)\sqrt{ab}}$$

[In] integrate(1/(a+b*coth(d*x+c)^2)^3,x, algorithm="giac")

[Out]
$$-1/8*((15*a^2*b + 10*a*b^2 + 3*b^3)*\arctan(1/2*(a*e^{(2*d*x + 2*c)} + b*e^{(2*d*x + 2*c)} - a + b)/\sqrt{a*b}))/((a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3)*\sqrt{a*b}) - 8*(d*x + c)/(a^3 + 3*a^2*b + 3*a*b^2 + b^3) - 2*(9*a^3*b*e^{(6*d*x + 6*c)} - a^2*b^2*e^{(6*d*x + 6*c)} - 13*a*b^3*e^{(6*d*x + 6*c)} - 27*a^3*b*e^{(4*d*x + 4*c)} + 9*a^2*b^2*e^{(4*d*x + 4*c)} - 21*a*b^3*e^{(4*d*x + 4*c)} - 9*b^4*e^{(4*d*x + 4*c)} + 27*a^3*b*e^{(2*d*x + 2*c)} + 13*a^2*b^2*e^{(2*d*x + 2*c)} - 23*a*b^3*e^{(2*d*x + 2*c)} - 9*b^4*e^{(2*d*x + 2*c)} - 9*a^3*b - 21*a^2*b^2 - 15*a*b^3 - 3*b^4)/(a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3)*(a*e^{(4*d*x + 4*c)} + b*e^{(4*d*x + 4*c)} - 2*a*e^{(2*d*x + 2*c)} + 2*b*e^{(2*d*x + 2*c)} + a + b)^2)/d$$

Mupad [B] (verification not implemented)

Time = 2.71 (sec) , antiderivative size = 2719, normalized size of antiderivative = 19.15

$$\int \frac{1}{(a + b \coth^2(c + dx))^3} dx = \text{Too large to display}$$

[In] int(1/(a + b*coth(c + d*x)^2)^3,x)

[Out]
$$\log(\coth(c + d*x) + 1)/(2*a^3*d + 2*b^3*d + 6*a*b^2*d + 6*a^2*b*d) + ((\coth(c + d*x)*(9*a*b + 5*b^2))/(8*a*(2*a*b + a^2 + b^2)) + (b*\coth(c + d*x)^3*(7*a*b + 3*b^2))/(8*a*(a*b^2 + 2*a^2*b + a^3)))/(a^2*d + b^2*d*\coth(c + d*x)^4 + 2*a*b*d*\coth(c + d*x)^2) - \log(\coth(c + d*x) - 1)/(2*d*(a + b)^3) - (a \tan(\frac{(-a^5*b)^{1/2}*(\coth(c + d*x)*(60*a*b^6 + 9*b^7 + 190*a^2*b^5 + 300*a^3*b^4 + 289*a^4*b^3))}{(32*(a^8*d^2 + 4*a^7*b*d^2 + a^4*b^4*d^2 + 4*a^5*b^3*d^2 + 6*a^6*b^2*d^2))} + ((96*a^2*b^10*d^2 + 800*a^3*b^9*d^2 + 3040*a^4*b^8*d^2 + 6816*a^5*b^7*d^2 + 9760*a^6*b^6*d^2 + 9056*a^7*b^5*d^2 + 5280*a^8*b^4*d^2 + 1760*a^9*b^3*d^2 + 256*a^10*b^2*d^2)/(64*(a^10*d^3 + 6*a^9*b*d^3 + a^4*b^6*d^3 + 6*a^5*b^5*d^3 + 15*a^6*b^4*d^3 + 20*a^7*b^3*d^3 + 15*a^8*b^2*d^3)) - (\coth(c + d*x)*(-a^5*b)^{1/2}*(10*a*b + 15*a^2 + 3*b^2)*(256*a^4*$$

$$\frac{(b^3d + 3a^6b^2d + 3a^7bd)(10ab + 15a^2 + 3b^2)}{(16(a^8d + a^5b^3d + 3a^6b^2d + 3a^7bd))(-a^5b)^{1/2}(10ab + 15a^2 + 3b^2)i} / (8(a^8d + a^5b^3d + 3a^6b^2d + 3a^7bd))$$

3.8 $\int \frac{1}{(a+b \coth^2(c+dx))^4} dx$

Optimal result	87
Rubi [A] (verified)	88
Mathematica [A] (verified)	90
Maple [A] (verified)	90
Fricas [B] (verification not implemented)	91
Sympy [F(-1)]	91
Maxima [B] (verification not implemented)	92
Giac [B] (verification not implemented)	93
Mupad [B] (verification not implemented)	93

Optimal result

Integrand size = 14, antiderivative size = 201

$$\int \frac{1}{(a+b \coth^2(c+dx))^4} dx = \frac{x}{(a+b)^4} - \frac{\sqrt{b}(35a^3 + 35a^2b + 21ab^2 + 5b^3) \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{16a^{7/2}(a+b)^4d} + \frac{b \coth(c+dx)}{6a(a+b)d (a+b \coth^2(c+dx))^3} + \frac{b(11a+5b) \coth(c+dx)}{24a^2(a+b)^2d (a+b \coth^2(c+dx))^2} + \frac{b(19a^2+16ab+5b^2) \coth(c+dx)}{16a^3(a+b)^3d (a+b \coth^2(c+dx))}$$

```
[Out] x/(a+b)^4+1/6*b*coth(d*x+c)/a/(a+b)/d/(a+b*coth(d*x+c)^2)^3+1/24*b*(11*a+5*
b)*coth(d*x+c)/a^2/(a+b)^2/d/(a+b*coth(d*x+c)^2)^2+1/16*b*(19*a^2+16*a*b+5*
b^2)*coth(d*x+c)/a^3/(a+b)^3/d/(a+b*coth(d*x+c)^2)-1/16*(35*a^3+35*a^2*b+21
*a*b^2+5*b^3)*arctan(a^(1/2)*tanh(d*x+c)/b^(1/2))*b^(1/2)/a^(7/2)/(a+b)^4/d
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3742, 425, 541, 536, 212, 211}

$$\int \frac{1}{(a + b \coth^2(c + dx))^4} dx = \frac{b(11a + 5b) \coth(c + dx)}{24a^2 d(a + b)^2 (a + b \coth^2(c + dx))^2} + \frac{b(19a^2 + 16ab + 5b^2) \coth(c + dx)}{16a^3 d(a + b)^3 (a + b \coth^2(c + dx))} - \frac{\sqrt{b}(35a^3 + 35a^2b + 21ab^2 + 5b^3) \arctan\left(\frac{\sqrt{a} \tanh(c + dx)}{\sqrt{b}}\right)}{16a^{7/2} d(a + b)^4} + \frac{b \coth(c + dx)}{6ad(a + b) (a + b \coth^2(c + dx))^3} + \frac{x}{(a + b)^4}$$

[In] Int[(a + b*Coth[c + d*x]^2)^(-4), x]

[Out] x/(a + b)^4 - (Sqrt[b]*(35*a^3 + 35*a^2*b + 21*a*b^2 + 5*b^3)*ArcTan[(Sqrt[a]*Tanh[c + d*x])/Sqrt[b]])/(16*a^(7/2)*(a + b)^4*d) + (b*Coth[c + d*x])/(6*a*(a + b)*d*(a + b*Coth[c + d*x]^2)^3) + (b*(11*a + 5*b)*Coth[c + d*x])/(2*4*a^2*(a + b)^2*d*(a + b*Coth[c + d*x]^2)^2) + (b*(19*a^2 + 16*a*b + 5*b^2)*Coth[c + d*x])/(16*a^3*(a + b)^3*d*(a + b*Coth[c + d*x]^2))

Rule 211

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 425

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-b)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(p + 1)*(b*c - a*d))), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 536

Int[((e_) + (f_)*(x_)^(n_))/(((a_) + (b_)*(x_)^(n_))*((c_) + (d_)*(x_)^(n_))), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 541

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] := Simp[(-b*e - a*f)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(b*c - a*d)*(p + 1))), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rule 3742

Int[((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int \frac{1}{(1-x^2)(a+bx^2)^4} dx, x, \coth(c+dx)\right)}{d} \\
 &= \frac{b \coth(c+dx)}{6a(a+b)d (a+b \coth^2(c+dx))^3} - \frac{\text{Subst}\left(\int \frac{b-6(a+b)+5bx^2}{(1-x^2)(a+bx^2)^3} dx, x, \coth(c+dx)\right)}{6a(a+b)d} \\
 &= \frac{b \coth(c+dx)}{6a(a+b)d (a+b \coth^2(c+dx))^3} + \frac{b(11a+5b) \coth(c+dx)}{24a^2(a+b)^2d (a+b \coth^2(c+dx))^2} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{3(8a^2+11ab+5b^2)-3b(11a+5b)x^2}{(1-x^2)(a+bx^2)^2} dx, x, \coth(c+dx)\right)}{24a^2(a+b)^2d} \\
 &= \frac{b \coth(c+dx)}{6a(a+b)d (a+b \coth^2(c+dx))^3} + \frac{b(11a+5b) \coth(c+dx)}{24a^2(a+b)^2d (a+b \coth^2(c+dx))^2} \\
 &\quad + \frac{b(19a^2+16ab+5b^2) \coth(c+dx)}{16a^3(a+b)^3d (a+b \coth^2(c+dx))} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{-3(16a^3+19a^2b+16ab^2+5b^3)+3b(19a^2+16ab+5b^2)x^2}{(1-x^2)(a+bx^2)} dx, x, \coth(c+dx)\right)}{48a^3(a+b)^3d}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{b \coth(c+dx)}{6a(a+b)d(a+b \coth^2(c+dx))^3} + \frac{b(11a+5b) \coth(c+dx)}{24a^2(a+b)^2d(a+b \coth^2(c+dx))^2} \\
&\quad + \frac{b(19a^2+16ab+5b^2) \coth(c+dx)}{16a^3(a+b)^3d(a+b \coth^2(c+dx))} + \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \coth(c+dx)\right)}{(a+b)^4d} \\
&\quad + \frac{(b(35a^3+35a^2b+21ab^2+5b^3)) \text{Subst}\left(\int \frac{1}{a+bx^2} dx, x, \coth(c+dx)\right)}{16a^3(a+b)^4d} \\
&= \frac{x}{(a+b)^4} - \frac{\sqrt{b}(35a^3+35a^2b+21ab^2+5b^3) \arctan\left(\frac{\sqrt{a} \tanh(c+dx)}{\sqrt{b}}\right)}{16a^{7/2}(a+b)^4d} \\
&\quad + \frac{b \coth(c+dx)}{6a(a+b)d(a+b \coth^2(c+dx))^3} + \frac{b(11a+5b) \coth(c+dx)}{24a^2(a+b)^2d(a+b \coth^2(c+dx))^2} \\
&\quad + \frac{b(19a^2+16ab+5b^2) \coth(c+dx)}{16a^3(a+b)^3d(a+b \coth^2(c+dx))}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.60 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.01

$$\int \frac{1}{(a+b \coth^2(c+dx))^4} dx$$

$$= \frac{3\sqrt{b}(35a^3+35a^2b+21ab^2+5b^3) \arctan\left(\frac{\sqrt{b} \coth(c+dx)}{\sqrt{a}}\right)}{a^{7/2}} + \frac{8b(a+b)^3 \coth(c+dx)}{a(a+b \coth^2(c+dx))^3} + \frac{2b(a+b)^2(11a+5b) \coth(c+dx)}{a^2(a+b \coth^2(c+dx))^2} + \frac{3b(a+b)(19a^2+16ab+5b^2) \coth(c+dx)}{a^3(a+b \coth^2(c+dx))} + \frac{48(a+b)^4d}{48(a+b)^4d}$$

[In] Integrate[(a + b*Coth[c + d*x]^2)^(-4), x]

[Out] ((3*sqrt[b]*(35*a^3 + 35*a^2*b + 21*a*b^2 + 5*b^3)*ArcTan[(sqrt[b]*Coth[c + d*x])/sqrt[a]])/a^(7/2) + (8*b*(a + b)^3*Coth[c + d*x])/(a*(a + b*Coth[c + d*x]^2)^3) + (2*b*(a + b)^2*(11*a + 5*b)*Coth[c + d*x])/(a^2*(a + b*Coth[c + d*x]^2)^2) + (3*b*(a + b)*(19*a^2 + 16*a*b + 5*b^2)*Coth[c + d*x])/(a^3*(a + b*Coth[c + d*x]^2)) - 24*Log[1 - Coth[c + d*x]] + 24*Log[1 + Coth[c + d*x]])/(48*(a + b)^4*d)

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.09

method	result
derivativedivides	$\frac{-\frac{\ln(\coth(dx+c)-1)}{2(a+b)^4} + \frac{\ln(\coth(dx+c)+1)}{2(a+b)^4} + b \left(\frac{b^2(19a^3+35a^2b+21ab^2+5b^3)\coth(dx+c)^5}{16a^3} + \frac{b(17a^3+33a^2b+21ab^2+5b^3)\coth(dx+c)}{(a+\coth(dx+c)^2b)^3} \right)}{d}$
default	$\frac{-\frac{\ln(\coth(dx+c)-1)}{2(a+b)^4} + \frac{\ln(\coth(dx+c)+1)}{2(a+b)^4} + b \left(\frac{b^2(19a^3+35a^2b+21ab^2+5b^3)\coth(dx+c)^5}{16a^3} + \frac{b(17a^3+33a^2b+21ab^2+5b^3)\coth(dx+c)}{(a+\coth(dx+c)^2b)^3} \right)}{d}$
risch	Expression too large to display

[In] `int(1/(a+coth(d*x+c)^2*b)^4,x,method=_RETURNVERBOSE)`

[Out] $1/d * (-1/2/(a+b)^4 * \ln(\coth(d*x+c)-1) + 1/2/(a+b)^4 * \ln(\coth(d*x+c)+1) + 1/(a+b)^4 * b * ((1/16*b^2*(19*a^3+35*a^2*b+21*a*b^2+5*b^3)/a^3*\coth(d*x+c)^5 + 1/6*b*(17*a^3+33*a^2*b+21*a*b^2+5*b^3)/a^2*\coth(d*x+c)^3 + 1/16*(29*a^3+61*a^2*b+43*a*b^2+11*b^3)/a*\coth(d*x+c))/(a+\coth(d*x+c)^2*b)^3 + 1/16*(35*a^3+35*a^2*b+21*a*b^2+5*b^3)/a^3/(a*b)^(1/2)*\arctan(b*\coth(d*x+c)/(a*b)^(1/2))))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 9853 vs. $2(185) = 370$.

Time = 0.52 (sec) , antiderivative size = 20031, normalized size of antiderivative = 99.66

$$\int \frac{1}{(a + b \coth^2(c + dx))^4} dx = \text{Too large to display}$$

[In] `integrate(1/(a+b*coth(d*x+c)^2)^4,x, algorithm="fricas")`

[Out] Too large to include

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \coth^2(c + dx))^4} dx = \text{Timed out}$$

[In] `integrate(1/(a+b*coth(d*x+c)**2)**4,x)`

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 925 vs. 2(185) = 370.

Time = 0.51 (sec) , antiderivative size = 925, normalized size of antiderivative = 4.60

$$\int \frac{1}{(a + b \coth^2(c + dx))^4} dx = \frac{(35 a^3 b + 35 a^2 b^2 + 21 a b^3 + 5 b^4) \arctan\left(\frac{(a+b)e^{(-2 dx - 2 c)} - a + b}{2\sqrt{ab}}\right)}{16 (a^7 + 4 a^6 b + 6 a^5 b^2 + 4 a^4 b^3 + a^3 b^4) \sqrt{abd}}$$

$$+ \frac{87 a^5 b + 319 a^4 b^2 + 450 a^3 b^3 + 306 a^2 b^4 + 103 a b^5 + 15 b^6 - 3(145 a^5 b + 267 a^4 b^2 + 34 a^3 b^3 - 178 a^2 b^4 - 115 a b^5 - 25 b^6) e^{(-2 dx - 2 c)} + 6(145 a^5 b + 93 a^4 b^2 - 6 a^3 b^3 + 106 a^2 b^4 + 85 a b^5 + 25 b^6) e^{(-4 dx - 4 c)} - 2(435 a^5 b + 29 a^4 b^2 + 162 a^3 b^3 - 306 a^2 b^4 - 245 a b^5 - 75 b^6) e^{(-6 dx - 6 c)} + 3(145 a^5 b + 17 a^4 b^2 - 58 a^3 b^3 + 150 a^2 b^4 + 105 a b^5 + 25 b^6) e^{(-8 dx - 8 c)} - 3(29 a^5 b + 23 a^4 b^2 - 62 a^3 b^3 - 82 a^2 b^4 - 31 a b^5 - 5 b^6) e^{(-10 dx - 10 c)}}{24 (a^{10} + 7 a^9 b + 21 a^8 b^2 + 35 a^7 b^3 + 35 a^6 b^4 + 21 a^5 b^5 + 7 a^4 b^6 + a^3 b^7 - 6 (a^{10} + 5 a^9 b + 9 a^8 b^2 + 5 a^7 b^3 + 35 a^6 b^4 + 21 a^5 b^5 + 7 a^4 b^6 + a^3 b^7) e^{(-2 dx - 2 c)} + 3(5 a^{10} + 19 a^9 b + 25 a^8 b^2 + 15 a^7 b^3 + 15 a^6 b^4 + 25 a^5 b^5 + 19 a^4 b^6 + 5 a^3 b^7) e^{(-4 dx - 4 c)} - 4(5 a^{10} + 17 a^9 b + 21 a^8 b^2 + 9 a^7 b^3 - 9 a^6 b^4 - 21 a^5 b^5 - 17 a^4 b^6 - 5 a^3 b^7) e^{(-6 dx - 6 c)} + 3(5 a^{10} + 19 a^9 b + 25 a^8 b^2 + 15 a^7 b^3 + 15 a^6 b^4 + 25 a^5 b^5 + 19 a^4 b^6 + 5 a^3 b^7) e^{(-8 dx - 8 c)} - 6(a^{10} + 5 a^9 b + 9 a^8 b^2 + 5 a^7 b^3 - 5 a^6 b^4 - 9 a^5 b^5 - 5 a^4 b^6 - a^3 b^7) e^{(-10 dx - 10 c)} + (a^{10} + 7 a^9 b + 21 a^8 b^2 + 35 a^7 b^3 + 35 a^6 b^4 + 21 a^5 b^5 + 7 a^4 b^6 + a^3 b^7) e^{(-12 dx - 12 c)}} d$$

[In] integrate(1/(a+b*coth(d*x+c)^2)^4,x, algorithm="maxima")

[Out] 1/16*(35*a^3*b + 35*a^2*b^2 + 21*a*b^3 + 5*b^4)*arctan(1/2*((a + b)*e^(-2*d*x - 2*c) - a + b)/sqrt(a*b))/((a^7 + 4*a^6*b + 6*a^5*b^2 + 4*a^4*b^3 + a^3*b^4)*sqrt(a*b)*d) + 1/24*(87*a^5*b + 319*a^4*b^2 + 450*a^3*b^3 + 306*a^2*b^4 + 103*a*b^5 + 15*b^6 - 3*(145*a^5*b + 267*a^4*b^2 + 34*a^3*b^3 - 178*a^2*b^4 - 115*a*b^5 - 25*b^6)*e^(-2*d*x - 2*c) + 6*(145*a^5*b + 93*a^4*b^2 - 6*a^3*b^3 + 106*a^2*b^4 + 85*a*b^5 + 25*b^6)*e^(-4*d*x - 4*c) - 2*(435*a^5*b + 29*a^4*b^2 + 162*a^3*b^3 - 306*a^2*b^4 - 245*a*b^5 - 75*b^6)*e^(-6*d*x - 6*c) + 3*(145*a^5*b + 17*a^4*b^2 - 58*a^3*b^3 + 150*a^2*b^4 + 105*a*b^5 + 25*b^6)*e^(-8*d*x - 8*c) - 3*(29*a^5*b + 23*a^4*b^2 - 62*a^3*b^3 - 82*a^2*b^4 - 31*a*b^5 - 5*b^6)*e^(-10*d*x - 10*c))/((a^10 + 7*a^9*b + 21*a^8*b^2 + 35*a^7*b^3 + 35*a^6*b^4 + 21*a^5*b^5 + 7*a^4*b^6 + a^3*b^7 - 6*(a^10 + 5*a^9*b + 9*a^8*b^2 + 5*a^7*b^3 - 5*a^6*b^4 - 9*a^5*b^5 - 5*a^4*b^6 - a^3*b^7)*e^(-2*d*x - 2*c) + 3*(5*a^10 + 19*a^9*b + 25*a^8*b^2 + 15*a^7*b^3 + 15*a^6*b^4 + 25*a^5*b^5 + 19*a^4*b^6 + 5*a^3*b^7)*e^(-4*d*x - 4*c) - 4*(5*a^10 + 17*a^9*b + 21*a^8*b^2 + 9*a^7*b^3 - 9*a^6*b^4 - 21*a^5*b^5 - 17*a^4*b^6 - 5*a^3*b^7)*e^(-6*d*x - 6*c) + 3*(5*a^10 + 19*a^9*b + 25*a^8*b^2 + 15*a^7*b^3 + 15*a^6*b^4 + 25*a^5*b^5 + 19*a^4*b^6 + 5*a^3*b^7)*e^(-8*d*x - 8*c) - 6*(a^10 + 5*a^9*b + 9*a^8*b^2 + 5*a^7*b^3 - 5*a^6*b^4 - 9*a^5*b^5 - 5*a^4*b^6 - a^3*b^7)*e^(-10*d*x - 10*c) + (a^10 + 7*a^9*b + 21*a^8*b^2 + 35*a^7*b^3 + 35*a^6*b^4 + 21*a^5*b^5 + 7*a^4*b^6 + a^3*b^7)*e^(-12*d*x - 12*c))*d) + (d*x + c)/((a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a*b^3 + b^4)*d)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 750 vs. $2(185) = 370$.

Time = 0.36 (sec) , antiderivative size = 750, normalized size of antiderivative = 3.73

$$\int \frac{1}{(a + b \coth^2(c + dx))^4} dx = \frac{3(35a^3b + 35a^2b^2 + 21ab^3 + 5b^4) \arctan\left(\frac{ae^{(2dx+2c)} + be^{(2dx+2c)} - a + b}{2\sqrt{ab}}\right)}{(a^7 + 4a^6b + 6a^5b^2 + 4a^4b^3 + a^3b^4)\sqrt{ab}} - \frac{48(dx+c)}{a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4} - \frac{2(87a^5be^{(10dx+10c)} + 69a^4b^2e^{(10dx+10c)} + \dots)}{\dots}$$

[In] integrate(1/(a+b*coth(d*x+c)^2)^4,x, algorithm="giac")

[Out]
$$-1/48*(3*(35*a^3*b + 35*a^2*b^2 + 21*a*b^3 + 5*b^4)*\arctan(1/2*(a*e^{(2*d*x + 2*c)} + b*e^{(2*d*x + 2*c)} - a + b)/\sqrt{a*b}))/((a^7 + 4*a^6*b + 6*a^5*b^2 + 4*a^4*b^3 + a^3*b^4)*\sqrt{a*b}) - 48*(d*x + c)/(a^4 + 4*a^3*b + 6*a^2*b^2 + 4*a*b^3 + b^4) - 2*(87*a^5*b*e^{(10*d*x + 10*c)} + 69*a^4*b^2*e^{(10*d*x + 10*c)} - 186*a^3*b^3*e^{(10*d*x + 10*c)} - 246*a^2*b^4*e^{(10*d*x + 10*c)} - 93*a*b^5*e^{(10*d*x + 10*c)} - 15*b^6*e^{(10*d*x + 10*c)} - 435*a^5*b*e^{(8*d*x + 8*c)} - 51*a^4*b^2*e^{(8*d*x + 8*c)} + 174*a^3*b^3*e^{(8*d*x + 8*c)} - 450*a^2*b^4*e^{(8*d*x + 8*c)} - 315*a*b^5*e^{(8*d*x + 8*c)} - 75*b^6*e^{(8*d*x + 8*c)} + 870*a^5*b*e^{(6*d*x + 6*c)} + 58*a^4*b^2*e^{(6*d*x + 6*c)} + 324*a^3*b^3*e^{(6*d*x + 6*c)} - 612*a^2*b^4*e^{(6*d*x + 6*c)} - 490*a*b^5*e^{(6*d*x + 6*c)} - 150*b^6*e^{(6*d*x + 6*c)} - 870*a^5*b*e^{(4*d*x + 4*c)} - 558*a^4*b^2*e^{(4*d*x + 4*c)} + 36*a^3*b^3*e^{(4*d*x + 4*c)} - 636*a^2*b^4*e^{(4*d*x + 4*c)} - 510*a*b^5*e^{(4*d*x + 4*c)} - 150*b^6*e^{(4*d*x + 4*c)} + 435*a^5*b*e^{(2*d*x + 2*c)} + 801*a^4*b^2*e^{(2*d*x + 2*c)} + 102*a^3*b^3*e^{(2*d*x + 2*c)} - 534*a^2*b^4*e^{(2*d*x + 2*c)} - 345*a*b^5*e^{(2*d*x + 2*c)} - 75*b^6*e^{(2*d*x + 2*c)} - 87*a^5*b - 319*a^4*b^2 - 450*a^3*b^3 - 306*a^2*b^4 - 103*a*b^5 - 15*b^6)/((a^7 + 4*a^6*b + 6*a^5*b^2 + 4*a^4*b^3 + a^3*b^4)*(a*e^{(4*d*x + 4*c)} + b*e^{(4*d*x + 4*c)} - 2*a*e^{(2*d*x + 2*c)} + 2*b*e^{(2*d*x + 2*c)} + a + b)^3))/d$$

Mupad [B] (verification not implemented)

Time = 2.81 (sec) , antiderivative size = 3685, normalized size of antiderivative = 18.33

$$\int \frac{1}{(a + b \coth^2(c + dx))^4} dx = \text{Too large to display}$$

[In] int(1/(a + b*coth(c + d*x)^2)^4,x)

[Out]
$$\log(\coth(c + d*x) + 1)/(2*a^4*d + 2*b^4*d + 12*a^2*b^2*d + 8*a*b^3*d + 8*a^3*b*d) + ((\coth(c + d*x)^3*(16*a*b^3 + 5*b^4 + 17*a^2*b^2))/(6*a^2*(3*a*b^2 + 3*a^2*b + a^3 + b^3)) + (\coth(c + d*x)*(32*a*b^2 + 29*a^2*b + 11*b^3)))/(\dots)$$

$$\begin{aligned}
& 16a*(3a*b^2 + 3a^2*b + a^3 + b^3) + (b^2*\coth(c + d*x)^5*(16a*b^2 + 19 \\
& a^2*b + 5b^3))/(16a^2*(a*b^3 + 3a^3*b + a^4 + 3a^2*b^2))/(a^3*d + b^3 \\
& *d*\coth(c + d*x)^6 + 3a^2*b*d*\coth(c + d*x)^2 + 3a*b^2*d*\coth(c + d*x)^4) \\
& - \log(\coth(c + d*x) - 1)/(2*d*(a + b)^4) - (\operatorname{atan}(\frac{(-a^7*b)^{1/2}*((\coth(c + \\
& d*x)*(210a*b^8 + 25b^9 + 791a^2*b^7 + 1820a^3*b^6 + 2695a^4*b^5 + 2 \\
& 450a^5*b^4 + 1481a^6*b^3))}{(128*(a^{12}*d^2 + 6a^{11}*b*d^2 + a^6*b^6*d^2 + \\
& 6a^7*b^5*d^2 + 15a^8*b^4*d^2 + 20a^9*b^3*d^2 + 15a^{10}*b^2*d^2))} + (((5 \\
& a^3*b^{13}*d^2)/4 + 14a^4*b^{12}*d^2 + (287a^5*b^{11}*d^2)/4 + 224a^6*b^{10}*d^2 \\
& + (953a^7*b^9*d^2)/2 + 728a^8*b^8*d^2 + (1631a^9*b^7*d^2)/2 + 668a^{10} \\
& *b^6*d^2 + (1561a^{11}*b^5*d^2)/4 + 154a^{12}*b^4*d^2 + (147a^{13}*b^3*d^2)/4 \\
& + 4a^{14}*b^2*d^2)/(a^{15}*d^3 + 9a^{14}*b*d^3 + a^6*b^9*d^3 + 9a^7*b^8*d^3 + \\
& 36a^8*b^7*d^3 + 84a^9*b^6*d^3 + 126a^{10}*b^5*d^3 + 126a^{11}*b^4*d^3 + 84a^{12} \\
& *b^3*d^3 + 36a^{13}*b^2*d^3) - (\coth(c + d*x)*(-a^7*b)^{1/2}*(21a*b^2 + \\
& 35a^2*b + 35a^3 + 5b^3)*(1024a^6*b^{11}*d^2 + 7168a^7*b^{10}*d^2 + 20480a^8 \\
& *b^9*d^2 + 28672a^9*b^8*d^2 + 14336a^{10}*b^7*d^2 - 14336a^{11}*b^6*d^2 - \\
& 28672a^{12}*b^5*d^2 - 20480a^{13}*b^4*d^2 - 7168a^{14}*b^3*d^2 - 1024a^{15}*b^2 \\
& *d^2))/(4096*(a^{11}*d + a^7*b^4*d + 4a^8*b^3*d + 6a^9*b^2*d + 4a^{10}*b*d) \\
& *(a^{12}*d^2 + 6a^{11}*b*d^2 + a^6*b^6*d^2 + 6a^7*b^5*d^2 + 15a^8*b^4*d^2 + \\
& 20a^9*b^3*d^2 + 15a^{10}*b^2*d^2)))*(-a^7*b)^{1/2}*(21a*b^2 + 35a^2*b + 3 \\
& 5a^3 + 5b^3))/(32*(a^{11}*d + a^7*b^4*d + 4a^8*b^3*d + 6a^9*b^2*d + 4a^{10} \\
& *b*d)))*(21a*b^2 + 35a^2*b + 35a^3 + 5b^3)*i)/(32*(a^{11}*d + a^7*b^4*d \\
& + 4a^8*b^3*d + 6a^9*b^2*d + 4a^{10}*b*d)) + ((-a^7*b)^{1/2}*((\coth(c + d* \\
& x)*(210a*b^8 + 25b^9 + 791a^2*b^7 + 1820a^3*b^6 + 2695a^4*b^5 + 2450a^5 \\
& *b^4 + 1481a^6*b^3)))/(128*(a^{12}*d^2 + 6a^{11}*b*d^2 + a^6*b^6*d^2 + 6a^7 \\
& *b^5*d^2 + 15a^8*b^4*d^2 + 20a^9*b^3*d^2 + 15a^{10}*b^2*d^2)) - (((5a^3* \\
& b^{13}*d^2)/4 + 14a^4*b^{12}*d^2 + (287a^5*b^{11}*d^2)/4 + 224a^6*b^{10}*d^2 + (\\
& 953a^7*b^9*d^2)/2 + 728a^8*b^8*d^2 + (1631a^9*b^7*d^2)/2 + 668a^{10}*b^6* \\
& d^2 + (1561a^{11}*b^5*d^2)/4 + 154a^{12}*b^4*d^2 + (147a^{13}*b^3*d^2)/4 + 4a^ \\
& ^{14}*b^2*d^2)/(a^{15}*d^3 + 9a^{14}*b*d^3 + a^6*b^9*d^3 + 9a^7*b^8*d^3 + 36a^8 \\
& *b^7*d^3 + 84a^9*b^6*d^3 + 126a^{10}*b^5*d^3 + 126a^{11}*b^4*d^3 + 84a^{12} \\
& *b^3*d^3 + 36a^{13}*b^2*d^3) + (\coth(c + d*x)*(-a^7*b)^{1/2}*(21a*b^2 + 35a^ \\
& ^2*b + 35a^3 + 5b^3)*(1024a^6*b^{11}*d^2 + 7168a^7*b^{10}*d^2 + 20480a^8*b^ \\
& ^9*d^2 + 28672a^9*b^8*d^2 + 14336a^{10}*b^7*d^2 - 14336a^{11}*b^6*d^2 - 2867 \\
& 2a^{12}*b^5*d^2 - 20480a^{13}*b^4*d^2 - 7168a^{14}*b^3*d^2 - 1024a^{15}*b^2*d^2 \\
&))/(4096*(a^{11}*d + a^7*b^4*d + 4a^8*b^3*d + 6a^9*b^2*d + 4a^{10}*b*d)*(a^{12} \\
& *d^2 + 6a^{11}*b*d^2 + a^6*b^6*d^2 + 6a^7*b^5*d^2 + 15a^8*b^4*d^2 + 20a^9 \\
& *b^3*d^2 + 15a^{10}*b^2*d^2)))*(-a^7*b)^{1/2}*(21a*b^2 + 35a^2*b + 35a^3 \\
& + 5b^3))/(32*(a^{11}*d + a^7*b^4*d + 4a^8*b^3*d + 6a^9*b^2*d + 4a^{10}*b*d) \\
&))*(21a*b^2 + 35a^2*b + 35a^3 + 5b^3)*i)/(32*(a^{11}*d + a^7*b^4*d + 4a^ \\
& ^8*b^3*d + 6a^9*b^2*d + 4a^{10}*b*d)))/(((185a*b^7)/128 + (25*b^8)/128 + \\
& (303a^2*b^6)/64 + (567a^3*b^5)/64 + (1225a^4*b^4)/128 + (665a^5*b^3)/12 \\
& 8)/(a^{15}*d^3 + 9a^{14}*b*d^3 + a^6*b^9*d^3 + 9a^7*b^8*d^3 + 36a^8*b^7*d^3 \\
& + 84a^9*b^6*d^3 + 126a^{10}*b^5*d^3 + 126a^{11}*b^4*d^3 + 84a^{12}*b^3*d^3 + \\
& 36a^{13}*b^2*d^3) + ((-a^7*b)^{1/2}*((\coth(c + d*x)*(210a*b^8 + 25b^9 + 79 \\
& 1a^2*b^7 + 1820a^3*b^6 + 2695a^4*b^5 + 2450a^5*b^4 + 1481a^6*b^3))/(12
\end{aligned}$$

$$\begin{aligned}
& 8*(a^{12}d^2 + 6a^{11}b*d^2 + a^6*b^6*d^2 + 6a^7*b^5*d^2 + 15a^8*b^4*d^2 + \\
& 20a^9*b^3*d^2 + 15a^{10}*b^2*d^2)) + (((5a^3*b^{13}d^2)/4 + 14a^4*b^{12}d^2 \\
& ^2 + (287a^5*b^{11}d^2)/4 + 224a^6*b^{10}d^2 + (953a^7*b^9d^2)/2 + 728a^8*b^8d^2 \\
& + (1631a^9*b^7d^2)/2 + 668a^{10}*b^6d^2 + (1561a^{11}*b^5d^2)/4 \\
& + 154a^{12}*b^4d^2 + (147a^{13}*b^3d^2)/4 + 4a^{14}*b^2d^2)/(a^{15}d^3 + 9a^{14}*b*d^3 \\
& + a^6*b^9d^3 + 9a^7*b^8d^3 + 36a^8*b^7d^3 + 84a^9*b^6d^3 + 126a^{10}*b^5d^3 \\
& + 126a^{11}*b^4d^3 + 84a^{12}*b^3d^3 + 36a^{13}*b^2d^3) \\
& - (\coth(c + d*x)*(-a^7*b)^{(1/2)}*(21*a*b^2 + 35*a^2*b + 35*a^3 + 5*b^3)*(102 \\
& 4*a^6*b^{11}d^2 + 7168*a^7*b^{10}d^2 + 20480*a^8*b^9d^2 + 28672*a^9*b^8d^2 \\
& + 14336*a^{10}*b^7d^2 - 14336*a^{11}*b^6d^2 - 28672*a^{12}*b^5d^2 - 20480*a^{13} \\
& *b^4d^2 - 7168*a^{14}*b^3d^2 - 1024*a^{15}*b^2d^2))/(4096*(a^{11}d + a^7*b^4* \\
& d + 4*a^8*b^3*d + 6*a^9*b^2*d + 4*a^{10}*b*d))*(a^{12}d^2 + 6a^{11}*b*d^2 + a^6* \\
& b^6*d^2 + 6a^7*b^5*d^2 + 15a^8*b^4*d^2 + 20a^9*b^3*d^2 + 15a^{10}*b^2*d^2 \\
&))*(-a^7*b)^{(1/2)}*(21*a*b^2 + 35*a^2*b + 35*a^3 + 5*b^3))/(32*(a^{11}d + a^7* \\
& b^4*d + 4*a^8*b^3*d + 6*a^9*b^2*d + 4*a^{10}*b*d))*(21*a*b^2 + 35*a^2*b + \\
& 35*a^3 + 5*b^3))/(32*(a^{11}d + a^7*b^4*d + 4*a^8*b^3*d + 6*a^9*b^2*d + 4*a^ \\
& 10*b*d)) - ((-a^7*b)^{(1/2)}*(\coth(c + d*x)*(210*a*b^8 + 25*b^9 + 791*a^2*b^7 \\
& + 1820*a^3*b^6 + 2695*a^4*b^5 + 2450*a^5*b^4 + 1481*a^6*b^3))/(128*(a^{12} \\
& d^2 + 6a^{11}*b*d^2 + a^6*b^6*d^2 + 6a^7*b^5*d^2 + 15a^8*b^4*d^2 + 20a^9* \\
& b^3*d^2 + 15a^{10}*b^2*d^2)) - (((5a^3*b^{13}d^2)/4 + 14a^4*b^{12}d^2 + (28 \\
& 7a^5*b^{11}d^2)/4 + 224a^6*b^{10}d^2 + (953a^7*b^9d^2)/2 + 728a^8*b^8d^2 \\
& + (1631a^9*b^7d^2)/2 + 668a^{10}*b^6d^2 + (1561a^{11}*b^5d^2)/4 + 154a^{12}*b^4d^2 \\
& + (147a^{13}*b^3d^2)/4 + 4a^{14}*b^2d^2)/(a^{15}d^3 + 9a^{14}*b*d^3 + a^6*b^9d^3 \\
& + 9a^7*b^8d^3 + 36a^8*b^7d^3 + 84a^9*b^6d^3 + 126a^{10}*b^5d^3 + 126a^{11}*b^4d^3 \\
& + 84a^{12}*b^3d^3 + 36a^{13}*b^2d^3) + (\coth(c + d*x)*(-a^7*b)^{(1/2)}*(21*a*b^2 + 35*a^2*b + 35*a^3 + 5*b^3)*(1024*a^6*b^{11}d^2 + 7168*a^7*b^{10}d^2 + 20480*a^8*b^9d^2 + 28672*a^9*b^8d^2 + 14336*a^{10}*b^7d^2 - 14336*a^{11}*b^6d^2 - 28672*a^{12}*b^5d^2 - 20480*a^{13}*b^4d^2 - 7168*a^{14}*b^3d^2 - 1024*a^{15}*b^2d^2))/(4096*(a^{11}d + a^7*b^4*d + 4*a^8*b^3*d + 6*a^9*b^2*d + 4*a^{10}*b*d))*(a^{12}d^2 + 6a^{11}*b*d^2 + a^6*b^6*d^2 + 6a^7*b^5*d^2 + 15a^8*b^4*d^2 + 20a^9*b^3*d^2 + 15a^{10}*b^2*d^2))*(-a^7*b)^{(1/2)}*(21*a*b^2 + 35*a^2*b + 35*a^3 + 5*b^3))/(32*(a^{11}d + a^7*b^4*d + 4*a^8*b^3*d + 6*a^9*b^2*d + 4*a^{10}*b*d))*(21*a*b^2 + 35*a^2*b + 35*a^3 + 5*b^3))/(32*(a^{11}d + a^7*b^4*d + 4*a^8*b^3*d + 6*a^9*b^2*d + 4*a^{10}*b*d)))*(-a^7*b)^{(1/2)}*(21*a*b^2 + 35*a^2*b + 35*a^3 + 5*b^3)*1i)/(16*(a^{11}d + a^7*b^4*d + 4*a^8*b^3*d + 6*a^9*b^2*d + 4*a^{10}*b*d))
\end{aligned}$$

3.9 $\int \frac{1}{1-2 \coth^2(x)} dx$

Optimal result	96
Rubi [A] (verified)	96
Mathematica [A] (verified)	97
Maple [A] (verified)	97
Fricas [B] (verification not implemented)	98
Sympy [A] (verification not implemented)	98
Maxima [B] (verification not implemented)	98
Giac [B] (verification not implemented)	99
Mupad [B] (verification not implemented)	99

Optimal result

Integrand size = 10, antiderivative size = 19

$$\int \frac{1}{1-2 \coth^2(x)} dx = -x + \sqrt{2} \operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)$$

[Out] -x+arctanh(1/2*2^(1/2)*tanh(x))*2^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3741, 3756, 212}

$$\int \frac{1}{1-2 \coth^2(x)} dx = \sqrt{2} \operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right) - x$$

[In] Int[(1 - 2*Coth[x]^2)^(-1), x]

[Out] -x + Sqrt[2]*ArcTanh[Tanh[x]/Sqrt[2]]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3741

Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^(-1), x_Symbol] := Simp[x/(a - b), x] - Dist[b/(a - b), Int[Sec[e + f*x]^2/(a + b*Tan[e + f*x]^2), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a, b]

Rule 3756

```
Int[sec[(e_.) + (f_.)*(x_.)]^(m_.)*((a_.) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.))^(p_.), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/(c^(m - 1)*f), Subst[Int[(c^2 + ff^2*x^2)^(m/2 - 1)*(a + b*(ff*x)^n)^p, x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && IntegerQ[m/2] && (IntegersQ[n, p] || IGtQ[m, 0] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])
```

Rubi steps

$$\begin{aligned} \text{integral} &= -x - 2 \int \frac{\operatorname{csch}^2(x)}{1 - 2 \operatorname{coth}^2(x)} dx \\ &= -x + 2 \operatorname{Subst} \left(\int \frac{1}{1 - 2x^2} dx, x, \operatorname{coth}(x) \right) \\ &= -x + \sqrt{2} \operatorname{arctanh} \left(\frac{\operatorname{tanh}(x)}{\sqrt{2}} \right) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 - 2 \operatorname{coth}^2(x)} dx = -x + \sqrt{2} \operatorname{arctanh} \left(\frac{\operatorname{tanh}(x)}{\sqrt{2}} \right)$$

[In] Integrate[(1 - 2*Coth[x]^2)^(-1), x]

[Out] -x + Sqrt[2]*ArcTanh[Tanh[x]/Sqrt[2]]

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.42

method	result	size
derivativedivides	$-\frac{\ln(1+\operatorname{coth}(x))}{2} + \frac{\ln(\operatorname{coth}(x)-1)}{2} + \sqrt{2} \operatorname{arctanh}(\operatorname{coth}(x) \sqrt{2})$	27
default	$-\frac{\ln(1+\operatorname{coth}(x))}{2} + \frac{\ln(\operatorname{coth}(x)-1)}{2} + \sqrt{2} \operatorname{arctanh}(\operatorname{coth}(x) \sqrt{2})$	27
risch	$-x + \frac{\sqrt{2} \ln(e^{2x}+3-2\sqrt{2})}{2} - \frac{\sqrt{2} \ln(e^{2x}+3+2\sqrt{2})}{2}$	39

[In] int(1/(1-2*coth(x)^2), x, method=_RETURNVERBOSE)

[Out] -1/2*ln(1+coth(x))+1/2*ln(coth(x)-1)+2^(1/2)*arctanh(coth(x)*2^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 70 vs. $2(16) = 32$.

Time = 0.27 (sec) , antiderivative size = 70, normalized size of antiderivative = 3.68

$$\int \frac{1}{1 - 2 \coth^2(x)} dx$$

$$= \frac{1}{2} \sqrt{2} \log \left(-\frac{3(2\sqrt{2} - 3) \cosh(x)^2 - 4(3\sqrt{2} - 4) \cosh(x) \sinh(x) + 3(2\sqrt{2} - 3) \sinh(x)^2 + 2\sqrt{2} - 3}{\cosh(x)^2 + \sinh(x)^2 + 3} \right) - x$$

[In] integrate(1/(1-2*coth(x)^2),x, algorithm="fricas")

[Out] 1/2*sqrt(2)*log(-(3*(2*sqrt(2) - 3)*cosh(x)^2 - 4*(3*sqrt(2) - 4)*cosh(x)*sinh(x) + 3*(2*sqrt(2) - 3)*sinh(x)^2 + 2*sqrt(2) - 3)/(cosh(x)^2 + sinh(x)^2 + 3)) - x

Sympy [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.79

$$\int \frac{1}{1 - 2 \coth^2(x)} dx = -x - \frac{\sqrt{2} \log(\tanh(x) - \sqrt{2})}{2} + \frac{\sqrt{2} \log(\tanh(x) + \sqrt{2})}{2}$$

[In] integrate(1/(1-2*coth(x)**2),x)

[Out] -x - sqrt(2)*log(tanh(x) - sqrt(2))/2 + sqrt(2)*log(tanh(x) + sqrt(2))/2

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 38 vs. $2(16) = 32$.

Time = 0.28 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.00

$$\int \frac{1}{1 - 2 \coth^2(x)} dx = -\frac{1}{2} \sqrt{2} \log \left(-\frac{2\sqrt{2} - e^{(-2x)} - 3}{2\sqrt{2} + e^{(-2x)} + 3} \right) - x$$

[In] integrate(1/(1-2*coth(x)^2),x, algorithm="maxima")

[Out] -1/2*sqrt(2)*log(-(2*sqrt(2) - e^(-2*x) - 3)/(2*sqrt(2) + e^(-2*x) + 3)) - x

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 38 vs. $2(16) = 32$.

Time = 0.27 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.00

$$\int \frac{1}{1 - 2 \coth^2(x)} dx = \frac{1}{2} \sqrt{2} \log \left(-\frac{2\sqrt{2} - e^{(2x)} - 3}{2\sqrt{2} + e^{(2x)} + 3} \right) - x$$

[In] integrate(1/(1-2*coth(x)^2),x, algorithm="giac")

[Out] 1/2*sqrt(2)*log(-(2*sqrt(2) - e^(2*x) - 3)/(2*sqrt(2) + e^(2*x) + 3)) - x

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{1 - 2 \coth^2(x)} dx = \sqrt{2} \operatorname{atanh}(\sqrt{2} \coth(x)) - x$$

[In] int(-1/(2*coth(x)^2 - 1),x)

[Out] 2^(1/2)*atanh(2^(1/2)*coth(x)) - x

3.10 $\int \sqrt{1 - \coth^2(x)} dx$

Optimal result	100
Rubi [A] (verified)	100
Mathematica [B] (verified)	101
Maple [A] (verified)	101
Fricas [B] (verification not implemented)	102
Sympy [F]	102
Maxima [C] (verification not implemented)	102
Giac [C] (verification not implemented)	102
Mupad [B] (verification not implemented)	103

Optimal result

Integrand size = 12, antiderivative size = 3

$$\int \sqrt{1 - \coth^2(x)} dx = \arcsin(\coth(x))$$

[Out] arcsin(coth(x))

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3738, 4207, 222}

$$\int \sqrt{1 - \coth^2(x)} dx = \arcsin(\coth(x))$$

[In] Int[Sqrt[1 - Coth[x]^2], x]

[Out] ArcSin[Coth[x]]

Rule 222

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

Rule 3738

Int[(u_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)]^2)^p], x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]

Rule 4207

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] :> With[{ff = FreeFac
tors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[Int[(b + b*ff^2*x^2)^(p - 1),
x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \sqrt{-\operatorname{csch}^2(x)} dx \\ &= \operatorname{Subst}\left(\int \frac{1}{\sqrt{1-x^2}} dx, x, \operatorname{coth}(x)\right) \\ &= \arcsin(\operatorname{coth}(x)) \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 30 vs. $2(3) = 6$.

Time = 0.05 (sec) , antiderivative size = 30, normalized size of antiderivative = 10.00

$$\int \sqrt{1 - \operatorname{coth}^2(x)} dx = \sqrt{-\operatorname{csch}^2(x)} \left(-\log\left(\cosh\left(\frac{x}{2}\right)\right) + \log\left(\sinh\left(\frac{x}{2}\right)\right) \right) \sinh(x)$$

```
[In] Integrate[Sqrt[1 - Coth[x]^2], x]
```

```
[Out] Sqrt[-Csch[x]^2]*(-Log[Cosh[x/2]] + Log[Sinh[x/2]])*Sinh[x]
```

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
derivativedivides	$\arcsin(\operatorname{coth}(x))$	4
default	$\arcsin(\operatorname{coth}(x))$	4
risch	$\sqrt{-\frac{e^{2x}}{(e^{2x}-1)^2}} e^{-x}(e^{2x}-1) \ln(e^x-1) - \sqrt{-\frac{e^{2x}}{(e^{2x}-1)^2}} e^{-x}(e^{2x}-1) \ln(e^x+1)$	67

```
[In] int((1-coth(x)^2)^(1/2), x, method=_RETURNVERBOSE)
```

```
[Out] arcsin(coth(x))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 45 vs. $2(3) = 6$.

Time = 0.25 (sec) , antiderivative size = 45, normalized size of antiderivative = 15.00

$$\int \sqrt{1 - \coth^2(x)} dx = -2 \arctan \left(\frac{\sqrt{-\frac{e^{(2x)}}{e^{(4x)} - 2e^{(2x)} + 1}} (e^{(2x)} - 1)}{\cosh(x) e^x + e^x \sinh(x)} \right)$$

[In] integrate((1-coth(x)^2)^(1/2),x, algorithm="fricas")

[Out] -2*arctan(sqrt(-e^(2*x)/(e^(4*x) - 2*e^(2*x) + 1))*(e^(2*x) - 1)/(cosh(x)*e^x + e^x*sinh(x)))

Sympy [F]

$$\int \sqrt{1 - \coth^2(x)} dx = \int \sqrt{1 - \coth^2(x)} dx$$

[In] integrate((1-coth(x)**2)**(1/2),x)

[Out] Integral(sqrt(1 - coth(x)**2), x)

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 6.33

$$\int \sqrt{1 - \coth^2(x)} dx = i \log(e^{-x} + 1) - i \log(e^{-x} - 1)$$

[In] integrate((1-coth(x)^2)^(1/2),x, algorithm="maxima")

[Out] I*log(e^(-x) + 1) - I*log(e^(-x) - 1)

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 8.67

$$\int \sqrt{1 - \coth^2(x)} dx = (i \log(e^x + 1) - i \log(|e^x - 1|)) \operatorname{sgn}(-e^{(2x)} + 1)$$

[In] integrate((1-coth(x)^2)^(1/2),x, algorithm="giac")

[Out] (I*log(e^x + 1) - I*log(abs(e^x - 1)))*sgn(-e^(2*x) + 1)

Mupad [B] (verification not implemented)

Time = 1.90 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \sqrt{1 - \coth^2(x)} dx = \operatorname{asin}(\coth(x))$$

[In] `int((1 - coth(x)^2)^(1/2),x)`

[Out] `asin(coth(x))`

3.11 $\int \sqrt{-1 + \coth^2(x)} dx$

Optimal result	104
Rubi [A] (verified)	104
Mathematica [A] (verified)	105
Maple [A] (verified)	105
Fricas [A] (verification not implemented)	106
Sympy [F]	106
Maxima [A] (verification not implemented)	106
Giac [A] (verification not implemented)	107
Mupad [B] (verification not implemented)	107

Optimal result

Integrand size = 10, antiderivative size = 14

$$\int \sqrt{-1 + \coth^2(x)} dx = -\operatorname{arctanh}\left(\frac{\coth(x)}{\sqrt{\operatorname{csch}^2(x)}}\right)$$

[Out] $-\operatorname{arctanh}(\coth(x)/(\operatorname{csch}(x)^2)^{(1/2)})$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3738, 4207, 223, 212}

$$\int \sqrt{-1 + \coth^2(x)} dx = -\operatorname{arctanh}\left(\frac{\coth(x)}{\sqrt{\operatorname{csch}^2(x)}}\right)$$

[In] $\operatorname{Int}[\operatorname{Sqrt}[-1 + \operatorname{Coth}[x]^2], x]$

[Out] $-\operatorname{ArcTanh}[\operatorname{Coth}[x]/\operatorname{Sqrt}[\operatorname{Csch}[x]^2]]$

Rule 212

$\operatorname{Int}[(a_ + (b_ \cdot (x_)^2)^{-1}), x_Symbol] := \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2] \cdot \operatorname{Rt}[-b, 2])) \cdot \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2] \cdot (x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 223

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_ + (b_ \cdot (x_)^2)], x_Symbol] := \operatorname{Subst}[\operatorname{Int}[1/(1 - b \cdot x^2), x], x, x/\operatorname{Sqrt}[a + b \cdot x^2]] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a, 0]$

Rule 3738

```
Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]
```

Rule 4207

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \sqrt{\operatorname{csch}^2(x)} dx \\
 &= -\operatorname{Subst}\left(\int \frac{1}{\sqrt{-1+x^2}} dx, x, \operatorname{coth}(x)\right) \\
 &= -\operatorname{Subst}\left(\int \frac{1}{1-x^2} dx, x, \frac{\operatorname{coth}(x)}{\sqrt{\operatorname{csch}^2(x)}}\right) \\
 &= -\operatorname{arctanh}\left(\frac{\operatorname{coth}(x)}{\sqrt{\operatorname{csch}^2(x)}}\right)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 28, normalized size of antiderivative = 2.00

$$\int \sqrt{-1 + \operatorname{coth}^2(x)} dx = \sqrt{\operatorname{csch}^2(x)} \left(-\log\left(\cosh\left(\frac{x}{2}\right)\right) + \log\left(\sinh\left(\frac{x}{2}\right)\right) \right) \sinh(x)$$

```
[In] Integrate[Sqrt[-1 + Coth[x]^2], x]
```

```
[Out] Sqrt[Csch[x]^2]*(-Log[Cosh[x/2]] + Log[Sinh[x/2]])*Sinh[x]
```

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

method	result	size
derivativedivides	$-\ln\left(\coth(x) + \sqrt{\coth(x)^2 - 1}\right)$	15
default	$-\ln\left(\coth(x) + \sqrt{\coth(x)^2 - 1}\right)$	15
risch	$\sqrt{\frac{e^{2x}}{(e^{2x}-1)^2}} e^{-x}(e^{2x}-1)\ln(e^x-1) - \sqrt{\frac{e^{2x}}{(e^{2x}-1)^2}} e^{-x}(e^{2x}-1)\ln(e^x+1)$	65

```
[In] int((coth(x)^2-1)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -ln(coth(x)+(coth(x)^2-1)^(1/2))
```

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.21

$$\int \sqrt{-1 + \coth^2(x)} dx = -\log(\cosh(x) + \sinh(x) + 1) + \log(\cosh(x) + \sinh(x) - 1)$$

```
[In] integrate((-1+coth(x)^2)^(1/2),x, algorithm="fricas")
```

```
[Out] -log(cosh(x) + sinh(x) + 1) + log(cosh(x) + sinh(x) - 1)
```

Sympy [F]

$$\int \sqrt{-1 + \coth^2(x)} dx = \int \sqrt{\coth^2(x) - 1} dx$$

```
[In] integrate((-1+coth(x)**2)**(1/2),x)
```

```
[Out] Integral(sqrt(coth(x)**2 - 1), x)
```

Maxima [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.21

$$\int \sqrt{-1 + \coth^2(x)} dx = \log(e^{-x} + 1) - \log(e^{-x} - 1)$$

```
[In] integrate((-1+coth(x)^2)^(1/2),x, algorithm="maxima")
```

```
[Out] log(e^(-x) + 1) - log(e^(-x) - 1)
```

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.64

$$\int \sqrt{-1 + \coth^2(x)} dx = -(\log(e^x + 1) - \log(|e^x - 1|)) \operatorname{sgn}(e^{(2x)} - 1)$$

[In] integrate((-1+coth(x)^2)^(1/2),x, algorithm="giac")

[Out] -(log(e^x + 1) - log(abs(e^x - 1)))*sgn(e^(2*x) - 1)

Mupad [B] (verification not implemented)

Time = 2.09 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \sqrt{-1 + \coth^2(x)} dx = -\ln \left(\coth(x) + \sqrt{\coth(x)^2 - 1} \right)$$

[In] int((coth(x)^2 - 1)^(1/2),x)

[Out] -log(coth(x) + (coth(x)^2 - 1)^(1/2))

3.12 $\int (1 - \coth^2(x))^{3/2} dx$

Optimal result	108
Rubi [A] (verified)	108
Mathematica [B] (verified)	109
Maple [A] (verified)	110
Fricas [B] (verification not implemented)	110
Sympy [F]	111
Maxima [C] (verification not implemented)	111
Giac [C] (verification not implemented)	111
Mupad [B] (verification not implemented)	112

Optimal result

Integrand size = 12, antiderivative size = 24

$$\int (1 - \coth^2(x))^{3/2} dx = \frac{1}{2} \arcsin(\coth(x)) + \frac{1}{2} \coth(x) \sqrt{-\operatorname{csch}^2(x)}$$

[Out] 1/2*arcsin(coth(x))+1/2*coth(x)*(-csch(x)^2)^(1/2)

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3738, 4207, 201, 222}

$$\int (1 - \coth^2(x))^{3/2} dx = \frac{1}{2} \arcsin(\coth(x)) + \frac{1}{2} \coth(x) \sqrt{-\operatorname{csch}^2(x)}$$

[In] Int[(1 - Coth[x]^2)^(3/2), x]

[Out] ArcSin[Coth[x]]/2 + (Coth[x]*Sqrt[-Csch[x]^2])/2

Rule 201

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p
+ 1)), x] + Dist[a*n*(p/(n*p + 1)), Int[(a + b*x^n)^(p - 1), x], x] /; Free
Q[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] &&
IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n],
Denominator[p]])
```

Rule 222

```
Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt
[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

Rule 3738

```
Int[(u_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)]^2)^p, x_Symbol] := Int[A
ctivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ
[a, b]
```

Rule 4207

```
Int[((b_)*sec[(e_) + (f_)*(x_)]^2)^p, x_Symbol] := With[{ff = FreeFac
tors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[Int[(b + b*ff^2*x^2)^(p - 1),
x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int (-\operatorname{csch}^2(x))^{3/2} dx \\
 &= \operatorname{Subst}\left(\int \sqrt{1-x^2} dx, x, \operatorname{coth}(x)\right) \\
 &= \frac{1}{2} \operatorname{coth}(x) \sqrt{-\operatorname{csch}^2(x)} + \frac{1}{2} \operatorname{Subst}\left(\int \frac{1}{\sqrt{1-x^2}} dx, x, \operatorname{coth}(x)\right) \\
 &= \frac{1}{2} \arcsin(\operatorname{coth}(x)) + \frac{1}{2} \operatorname{coth}(x) \sqrt{-\operatorname{csch}^2(x)}
 \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 51 vs. $2(24) = 48$.

Time = 0.10 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.12

$$\begin{aligned}
 \int (1 - \operatorname{coth}^2(x))^{3/2} dx &= \frac{1}{8} \sqrt{-\operatorname{csch}^2(x)} \left(\operatorname{csch}^2\left(\frac{x}{2}\right) \right. \\
 &\quad \left. - 4 \log\left(\cosh\left(\frac{x}{2}\right)\right) + 4 \log\left(\sinh\left(\frac{x}{2}\right)\right) + \operatorname{sech}^2\left(\frac{x}{2}\right) \right) \sinh(x)
 \end{aligned}$$

```
[In] Integrate[(1 - Coth[x]^2)^(3/2), x]
```

```
[Out] (Sqrt[-Csch[x]^2]*(Csch[x/2]^2 - 4*Log[Cosh[x/2]] + 4*Log[Sinh[x/2]] + Sech
[x/2]^2)*Sinh[x])/8
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{\coth(x)\sqrt{1-\coth(x)^2}}{2} + \frac{\arcsin(\coth(x))}{2}$	21
default	$\frac{\coth(x)\sqrt{1-\coth(x)^2}}{2} + \frac{\arcsin(\coth(x))}{2}$	21
risch	$\frac{\sqrt{-\frac{e^{2x}}{(e^{2x}-1)^2}}(1+e^{2x})}{e^{2x}-1} - \frac{\sqrt{-\frac{e^{2x}}{(e^{2x}-1)^2}}e^{-x}(e^{2x}-1)\ln(e^x+1)}{2} + \frac{\sqrt{-\frac{e^{2x}}{(e^{2x}-1)^2}}e^{-x}(e^{2x}-1)\ln(e^x-1)}{2}$	99

```
[In] int((1-coth(x)^2)^(3/2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/2*coth(x)*(1-coth(x)^2)^(1/2)+1/2*arcsin(coth(x))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 279 vs. 2(18) = 36.

Time = 0.28 (sec) , antiderivative size = 279, normalized size of antiderivative = 11.62

$$\int (1 - \coth^2(x))^{3/2} dx =$$

$$(4 \cosh(x) e^x \sinh(x)^3 + e^x \sinh(x)^4 + 2(3 \cosh(x)^2 - 1)e^x \sinh(x)^2 + 4(\cosh(x)^3 - \cosh(x))e^x \sinh(x))$$

```
[In] integrate((1-coth(x)^2)^(3/2),x, algorithm="fricas")
```

```
[Out] -((4*cosh(x)*e^x*sinh(x)^3 + e^x*sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*e^x*sinh(x)^2 + 4*(cosh(x)^3 - cosh(x))*e^x*sinh(x) + (cosh(x)^4 - 2*cosh(x)^2 + 1)*e^x*arctan(sqrt(-e^(2*x))/(e^(4*x) - 2*e^(2*x) + 1))*(e^(2*x) - 1)/(cosh(x)*e^x + e^x*sinh(x))) - ((e^(2*x) - 1)*sinh(x)^3 - cosh(x)^3 + 3*(cosh(x)*e^(2*x) - cosh(x))*sinh(x)^2 + (cosh(x)^3 + cosh(x))*e^(2*x) - (3*cosh(x)^2 - (3*cosh(x)^2 + 1)*e^(2*x) + 1)*sinh(x) - cosh(x))*sqrt(-e^(2*x)/(e^(4*x) - 2*e^(2*x) + 1)))/(4*cosh(x)*e^x*sinh(x)^3 + e^x*sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*e^x*sinh(x)^2 + 4*(cosh(x)^3 - cosh(x))*e^x*sinh(x) + (cosh(x)^4 - 2*cosh(x)^2 + 1)*e^x)
```

Sympy [F]

$$\int (1 - \coth^2(x))^{3/2} dx = \int (1 - \coth^2(x))^{\frac{3}{2}} dx$$

[In] integrate((1-coth(x)**2)**(3/2),x)

[Out] Integral((1 - coth(x)**2)**(3/2), x)

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 49, normalized size of antiderivative = 2.04

$$\int (1 - \coth^2(x))^{3/2} dx = \frac{i e^{(-x)} + i e^{(-3x)}}{2 e^{(-2x)} - e^{(-4x)} - 1} + \frac{1}{2} i \log(e^{(-x)} + 1) - \frac{1}{2} i \log(e^{(-x)} - 1)$$

[In] integrate((1-coth(x)^2)^(3/2),x, algorithm="maxima")

[Out] (I*e^(-x) + I*e^(-3*x))/(2*e^(-2*x) - e^(-4*x) - 1) + 1/2*I*log(e^(-x) + 1) - 1/2*I*log(e^(-x) - 1)

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 60, normalized size of antiderivative = 2.50

$$\int (1 - \coth^2(x))^{3/2} dx = -\frac{1}{4} \left(\frac{4(i e^{(-x)} + i e^x)}{(e^{(-x)} + e^x)^2 - 4} - i \log(e^{(-x)} + e^x + 2) + i \log(e^{(-x)} + e^x - 2) \right) \operatorname{sgn}(-e^{(2x)} + 1)$$

[In] integrate((1-coth(x)^2)^(3/2),x, algorithm="giac")

[Out] -1/4*(4*(I*e^(-x) + I*e^x)/((e^(-x) + e^x)^2 - 4) - I*log(e^(-x) + e^x + 2) + I*log(e^(-x) + e^x - 2))*sgn(-e^(2*x) + 1)

Mupad [B] (verification not implemented)

Time = 1.95 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int (1 - \operatorname{coth}^2(x))^{3/2} dx = \frac{\operatorname{asin}(\operatorname{coth}(x))}{2} + \frac{\operatorname{coth}(x) \sqrt{1 - \operatorname{coth}(x)^2}}{2}$$

[In] `int((1 - coth(x)^2)^(3/2),x)`

[Out] `asin(coth(x))/2 + (coth(x)*(1 - coth(x)^2)^(1/2))/2`

3.13 $\int (-1 + \coth^2(x))^{3/2} dx$

Optimal result	113
Rubi [A] (verified)	113
Mathematica [A] (verified)	115
Maple [A] (verified)	115
Fricas [B] (verification not implemented)	115
Sympy [F]	116
Maxima [A] (verification not implemented)	116
Giac [B] (verification not implemented)	116
Mupad [B] (verification not implemented)	117

Optimal result

Integrand size = 10, antiderivative size = 31

$$\int (-1 + \coth^2(x))^{3/2} dx = \frac{1}{2} \operatorname{arctanh}\left(\frac{\coth(x)}{\sqrt{\operatorname{csch}^2(x)}}\right) - \frac{1}{2} \coth(x) \sqrt{\operatorname{csch}^2(x)}$$

[Out] $1/2*\operatorname{arctanh}(\coth(x)/(\operatorname{csch}(x)^2)^{(1/2)})-1/2*\coth(x)*(\operatorname{csch}(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3738, 4207, 201, 223, 212}

$$\int (-1 + \coth^2(x))^{3/2} dx = \frac{1}{2} \operatorname{arctanh}\left(\frac{\coth(x)}{\sqrt{\operatorname{csch}^2(x)}}\right) - \frac{1}{2} \coth(x) \sqrt{\operatorname{csch}^2(x)}$$

[In] $\operatorname{Int}[(-1 + \operatorname{Coth}[x]^2)^{(3/2)}, x]$

[Out] $\operatorname{ArcTanh}[\operatorname{Coth}[x]/\operatorname{Sqrt}[\operatorname{Csch}[x]^2]]/2 - (\operatorname{Coth}[x]*\operatorname{Sqrt}[\operatorname{Csch}[x]^2])/2$

Rule 201

$\operatorname{Int}[(a + b*x^n)^p, x_Symbol] \rightarrow \operatorname{Simp}[x*(a + b*x^n)^p/(n*p + 1), x] + \operatorname{Dist}[a*n*(p/(n*p + 1)), \operatorname{Int}[(a + b*x^n)^{p-1}, x], x] /;$ Free Q[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 223

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x],
x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

Rule 3738

```
Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^p, x_Symbol] := Int[A
ctivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ
[a, b]
```

Rule 4207

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^p, x_Symbol] := With[{ff = FreeFac
tors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[Int[(b + b*ff^2*x^2)^(p - 1),
x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \operatorname{csch}^2(x)^{3/2} dx \\
&= -\operatorname{Subst}\left(\int \sqrt{-1+x^2} dx, x, \operatorname{coth}(x)\right) \\
&= -\frac{1}{2} \operatorname{coth}(x) \sqrt{\operatorname{csch}^2(x)} + \frac{1}{2} \operatorname{Subst}\left(\int \frac{1}{\sqrt{-1+x^2}} dx, x, \operatorname{coth}(x)\right) \\
&= -\frac{1}{2} \operatorname{coth}(x) \sqrt{\operatorname{csch}^2(x)} + \frac{1}{2} \operatorname{Subst}\left(\int \frac{1}{1-x^2} dx, x, \frac{\operatorname{coth}(x)}{\sqrt{\operatorname{csch}^2(x)}}\right) \\
&= \frac{1}{2} \operatorname{arctanh}\left(\frac{\operatorname{coth}(x)}{\sqrt{\operatorname{csch}^2(x)}}\right) - \frac{1}{2} \operatorname{coth}(x) \sqrt{\operatorname{csch}^2(x)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.58

$$\int (-1 + \coth^2(x))^{3/2} dx = -\frac{1}{8}\sqrt{\operatorname{csch}^2(x)}\left(\operatorname{csch}^2\left(\frac{x}{2}\right) - 4\log\left(\cosh\left(\frac{x}{2}\right)\right) + 4\log\left(\sinh\left(\frac{x}{2}\right)\right) + \operatorname{sech}^2\left(\frac{x}{2}\right)\right)\sinh(x)$$

[In] Integrate[(-1 + Coth[x]^2)^(3/2), x]

[Out] -1/8*(Sqrt[Csch[x]^2]*(Csch[x/2]^2 - 4*Log[Cosh[x/2]] + 4*Log[Sinh[x/2]] + Sech[x/2]^2)*Sinh[x])

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

method	result	size
derivativedivides	$-\frac{\coth(x)\sqrt{\coth(x)^2-1}}{2} + \frac{\ln\left(\coth(x)+\sqrt{\coth(x)^2-1}\right)}{2}$	28
default	$-\frac{\coth(x)\sqrt{\coth(x)^2-1}}{2} + \frac{\ln\left(\coth(x)+\sqrt{\coth(x)^2-1}\right)}{2}$	28
risch	$-\frac{\sqrt{\frac{e^{2x}}{(e^{2x}-1)^2}}(1+e^{2x})}{e^{2x}-1} + \frac{\sqrt{\frac{e^{2x}}{(e^{2x}-1)^2}}e^{-x}(e^{2x}-1)\ln(e^x+1)}{2} - \frac{\sqrt{\frac{e^{2x}}{(e^{2x}-1)^2}}e^{-x}(e^{2x}-1)\ln(e^x-1)}{2}$	97

[In] int((coth(x)^2-1)^(3/2), x, method=_RETURNVERBOSE)

[Out] -1/2*coth(x)*(coth(x)^2-1)^(1/2)+1/2*ln(coth(x)+(coth(x)^2-1)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 211 vs. 2(23) = 46.

Time = 0.26 (sec) , antiderivative size = 211, normalized size of antiderivative = 6.81

$$\int (-1 + \coth^2(x))^{3/2} dx = \frac{2 \cosh(x)^3 + 6 \cosh(x) \sinh(x)^2 + 2 \sinh(x)^3 - (\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4) + 2(3 \cosh(x) \sinh(x)^2 - 2 \cosh(x)^2 + 4 \sinh(x)^2)}{8}$$

[In] integrate((-1+coth(x)^2)^(3/2), x, algorithm="fricas")

[Out] -1/2*(2*cosh(x)^3 + 6*cosh(x)*sinh(x)^2 + 2*sinh(x)^3 - (cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*sinh(x)^2 - 2*cosh(x)^2 + 4

```

*(cosh(x)^3 - cosh(x))*sinh(x) + 1)*log(cosh(x) + sinh(x) + 1) + (cosh(x)^4
+ 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*sinh(x)^2 - 2*cosh
(x)^2 + 4*(cosh(x)^3 - cosh(x))*sinh(x) + 1)*log(cosh(x) + sinh(x) - 1) + 2
*(3*cosh(x)^2 + 1)*sinh(x) + 2*cosh(x))/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 +
sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*sinh(x)^2 - 2*cosh(x)^2 + 4*(cosh(x)^3 - co
sh(x))*sinh(x) + 1)

```

Sympy [F]

$$\int (-1 + \coth^2(x))^{3/2} dx = \int (\coth^2(x) - 1)^{\frac{3}{2}} dx$$

```
[In] integrate((-1+coth(x)**2)**(3/2),x)
```

```
[Out] Integral((coth(x)**2 - 1)**(3/2), x)
```

Maxima [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.48

$$\int (-1 + \coth^2(x))^{3/2} dx = -\frac{e^{(-x)} + e^{(-3x)}}{2e^{(-2x)} - e^{(-4x)} - 1} - \frac{1}{2} \log(e^{(-x)} + 1) + \frac{1}{2} \log(e^{(-x)} - 1)$$

```
[In] integrate((-1+coth(x)^2)^(3/2),x, algorithm="maxima")
```

```
[Out] -(e^(-x) + e^(-3*x))/(2*e^(-2*x) - e^(-4*x) - 1) - 1/2*log(e^(-x) + 1) + 1/
2*log(e^(-x) - 1)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 52 vs. 2(23) = 46.

Time = 0.27 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.68

$$\int (-1 + \coth^2(x))^{3/2} dx = -\frac{1}{4} \left(\frac{4(e^{(-x)} + e^x)}{(e^{(-x)} + e^x)^2 - 4} - \log(e^{(-x)} + e^x + 2) + \log(e^{(-x)} + e^x - 2) \right) \operatorname{sgn}(e^{(2x)} - 1)$$

```
[In] integrate((-1+coth(x)^2)^(3/2),x, algorithm="giac")
```

```
[Out] -1/4*(4*(e^(-x) + e^x)/((e^(-x) + e^x)^2 - 4) - log(e^(-x) + e^x + 2) + log
(e^(-x) + e^x - 2))*sgn(e^(2*x) - 1)
```

Mupad [B] (verification not implemented)

Time = 1.92 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

$$\int (-1 + \coth^2(x))^{3/2} dx = \frac{\ln\left(\coth(x) + \sqrt{\coth(x)^2 - 1}\right)}{2} - \frac{\coth(x) \sqrt{\coth(x)^2 - 1}}{2}$$

[In] int((coth(x)^2 - 1)^(3/2),x)

[Out] log(coth(x) + (coth(x)^2 - 1)^(1/2))/2 - (coth(x)*(coth(x)^2 - 1)^(1/2))/2

$$3.14 \quad \int \frac{1}{\sqrt{1-\coth^2(x)}} dx$$

Optimal result	118
Rubi [A] (verified)	118
Mathematica [A] (verified)	119
Maple [A] (verified)	119
Fricas [B] (verification not implemented)	120
Sympy [F]	120
Maxima [C] (verification not implemented)	120
Giac [C] (verification not implemented)	121
Mupad [B] (verification not implemented)	121

Optimal result

Integrand size = 12, antiderivative size = 13

$$\int \frac{1}{\sqrt{1-\coth^2(x)}} dx = \frac{\coth(x)}{\sqrt{-\operatorname{csch}^2(x)}}$$

[Out] $\coth(x)/(-\operatorname{csch}(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3738, 4207, 197}

$$\int \frac{1}{\sqrt{1-\coth^2(x)}} dx = \frac{\coth(x)}{\sqrt{-\operatorname{csch}^2(x)}}$$

[In] `Int[1/Sqrt[1 - Coth[x]^2], x]`

[Out] `Coth[x]/Sqrt[-Csch[x]^2]`

Rule 197

`Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1) / a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

Rule 3738

`Int[(u_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^2)^(p_), x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ`

[a, b]

Rule 4207

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] :> With[{ff = FreeFac
tors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[Int[(b + b*ff^2*x^2)^(p - 1),
x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \frac{1}{\sqrt{-\operatorname{csch}^2(x)}} dx \\ &= \operatorname{Subst}\left(\int \frac{1}{(1-x^2)^{3/2}} dx, x, \operatorname{coth}(x)\right) \\ &= \frac{\operatorname{coth}(x)}{\sqrt{-\operatorname{csch}^2(x)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1 - \operatorname{coth}^2(x)}} dx = \frac{\operatorname{coth}(x)}{\sqrt{-\operatorname{csch}^2(x)}}$$

[In] Integrate[1/Sqrt[1 - Coth[x]^2],x]

[Out] Coth[x]/Sqrt[-Csch[x]^2]

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.08

method	result	size
derivativedivides	$\frac{\operatorname{coth}(x)}{\sqrt{1 - \operatorname{coth}^2(x)}}$	14
default	$\frac{\operatorname{coth}(x)}{\sqrt{1 - \operatorname{coth}^2(x)}}$	14
risch	$\frac{e^{2x}}{2\sqrt{-\frac{e^{2x}}{(e^{2x}-1)^2}}(e^{2x}-1)} + \frac{1}{2(e^{2x}-1)\sqrt{-\frac{e^{2x}}{(e^{2x}-1)^2}}}$	58

[In] int(1/(1-coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out] $1/(1-\coth(x)^2)^{(1/2)}*\coth(x)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 40 vs. $2(11) = 22$.

Time = 0.26 (sec) , antiderivative size = 40, normalized size of antiderivative = 3.08

$$\int \frac{1}{\sqrt{1 - \coth^2(x)}} dx = -(\cosh(x) e^{2x} - \cosh(x)) \sqrt{-\frac{e^{2x}}{e^{4x} - 2e^{2x} + 1}} e^{(-x)}$$

[In] `integrate(1/(1-coth(x)^2)^(1/2),x, algorithm="fricas")`

[Out] `-(cosh(x)*e^(2*x) - cosh(x))*sqrt(-e^(2*x)/(e^(4*x) - 2*e^(2*x) + 1))*e^(-x)`

Sympy [F]

$$\int \frac{1}{\sqrt{1 - \coth^2(x)}} dx = \int \frac{1}{\sqrt{1 - \coth^2(x)}} dx$$

[In] `integrate(1/(1-coth(x)**2)**(1/2),x)`

[Out] `Integral(1/sqrt(1 - coth(x)**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1}{\sqrt{1 - \coth^2(x)}} dx = \frac{1}{2}i e^{(-x)} + \frac{1}{2}i e^x$$

[In] `integrate(1/(1-coth(x)^2)^(1/2),x, algorithm="maxima")`

[Out] `1/2*I*e^(-x) + 1/2*I*e^x`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.85

$$\int \frac{1}{\sqrt{1 - \coth^2(x)}} dx = -\frac{-i e^{(-x)} - i e^x}{2 \operatorname{sgn}(-e^{(2x)} + 1)}$$

[In] integrate(1/(1-coth(x)^2)^(1/2),x, algorithm="giac")

[Out] -1/2*(-I*e^(-x) - I*e^x)/sgn(-e^(2*x) + 1)

Mupad [B] (verification not implemented)

Time = 2.09 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.38

$$\int \frac{1}{\sqrt{1 - \coth^2(x)}} dx = -\cosh(x) \sinh(x) \sqrt{-\frac{1}{\cosh(x)^2 - 1}}$$

[In] int(1/(1 - coth(x)^2)^(1/2),x)

[Out] -cosh(x)*sinh(x)*(-1/(cosh(x)^2 - 1))^(1/2)

$$3.15 \quad \int \frac{1}{\sqrt{-1 + \coth^2(x)}} dx$$

Optimal result	122
Rubi [A] (verified)	122
Mathematica [A] (verified)	123
Maple [A] (verified)	123
Fricas [A] (verification not implemented)	124
Sympy [F]	124
Maxima [A] (verification not implemented)	124
Giac [A] (verification not implemented)	125
Mupad [B] (verification not implemented)	125

Optimal result

Integrand size = 10, antiderivative size = 11

$$\int \frac{1}{\sqrt{-1 + \coth^2(x)}} dx = \frac{\coth(x)}{\sqrt{\operatorname{csch}^2(x)}}$$

[Out] $\coth(x)/(\operatorname{csch}(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3738, 4207, 197}

$$\int \frac{1}{\sqrt{-1 + \coth^2(x)}} dx = \frac{\coth(x)}{\sqrt{\operatorname{csch}^2(x)}}$$

[In] `Int[1/Sqrt[-1 + Coth[x]^2], x]`

[Out] `Coth[x]/Sqrt[Csch[x]^2]`

Rule 197

`Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1) / a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

Rule 3738

`Int[(u_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)]^2)^(p_), x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ`

[a, b]

Rule 4207

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \frac{1}{\sqrt{\operatorname{csch}^2(x)}} dx \\ &= -\operatorname{Subst}\left(\int \frac{1}{(-1 + x^2)^{3/2}} dx, x, \operatorname{coth}(x)\right) \\ &= \frac{\operatorname{coth}(x)}{\sqrt{\operatorname{csch}^2(x)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{-1 + \operatorname{coth}^2(x)}} dx = \frac{\operatorname{coth}(x)}{\sqrt{\operatorname{csch}^2(x)}}$$

[In] Integrate[1/Sqrt[-1 + Coth[x]^2],x]

[Out] Coth[x]/Sqrt[Csch[x]^2]

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
derivativedivides	$\frac{\operatorname{coth}(x)}{\sqrt{\operatorname{coth}(x)^2 - 1}}$	12
default	$\frac{\operatorname{coth}(x)}{\sqrt{\operatorname{coth}(x)^2 - 1}}$	12
risch	$\frac{e^{2x}}{2\sqrt{\frac{e^{2x}}{(e^{2x}-1)^2}}(e^{2x}-1)} + \frac{1}{2(e^{2x}-1)\sqrt{\frac{e^{2x}}{(e^{2x}-1)^2}}}$	56

[In] int(1/(coth(x)^2-1)^(1/2),x,method=_RETURNVERBOSE)

[Out] $\coth(x)/(\coth(x)^2-1)^{(1/2)}$

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.18

$$\int \frac{1}{\sqrt{-1 + \coth^2(x)}} dx = \cosh(x)$$

[In] `integrate(1/(-1+coth(x)^2)^(1/2),x, algorithm="fricas")`

[Out] $\cosh(x)$

Sympy [F]

$$\int \frac{1}{\sqrt{-1 + \coth^2(x)}} dx = \int \frac{1}{\sqrt{\coth^2(x) - 1}} dx$$

[In] `integrate(1/(-1+coth(x)**2)**(1/2),x)`

[Out] `Integral(1/sqrt(coth(x)**2 - 1), x)`

Maxima [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{-1 + \coth^2(x)}} dx = -\frac{1}{2} e^{(-x)} - \frac{1}{2} e^x$$

[In] `integrate(1/(-1+coth(x)^2)^(1/2),x, algorithm="maxima")`

[Out] $-1/2*e^{(-x)} - 1/2*e^x$

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.64

$$\int \frac{1}{\sqrt{-1 + \coth^2(x)}} dx = \frac{e^{(-x)} + e^x}{2 \operatorname{sgn}(e^{(2x)} - 1)}$$

[In] integrate(1/(-1+coth(x)^2)^(1/2),x, algorithm="giac")

[Out] 1/2*(e^(-x) + e^x)/sgn(e^(2*x) - 1)

Mupad [B] (verification not implemented)

Time = 1.97 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.36

$$\int \frac{1}{\sqrt{-1 + \coth^2(x)}} dx = \cosh(x) \sinh(x) \sqrt{\frac{1}{\cosh(x)^2 - 1}}$$

[In] int(1/(coth(x)^2 - 1)^(1/2),x)

[Out] cosh(x)*sinh(x)*(1/(cosh(x)^2 - 1))^(1/2)

3.16 $\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx$

Optimal result	126
Rubi [A] (verified)	126
Mathematica [A] (verified)	128
Maple [B] (verified)	129
Fricas [B] (verification not implemented)	129
Sympy [F]	131
Maxima [F]	131
Giac [F(-2)]	131
Mupad [B] (verification not implemented)	132

Optimal result

Integrand size = 17, antiderivative size = 63

$$\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx = \sqrt{a+b} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a+b}} \right) - \sqrt{a + b \coth^2(x)} - \frac{(a + b \coth^2(x))^{3/2}}{3b}$$

[Out] $-1/3*(a+b*\coth(x)^2)^{(3/2)}/b+\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2)}/(a+b)^{(1/2)})*(a+b)^{(1/2)}-(a+b*\coth(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3751, 457, 81, 52, 65, 214}

$$\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx = \sqrt{a+b} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a+b}} \right) - \frac{(a + b \coth^2(x))^{3/2}}{3b} - \sqrt{a + b \coth^2(x)}$$

[In] $\operatorname{Int}[\operatorname{Coth}[x]^3 \operatorname{Sqrt}[a + b \operatorname{Coth}[x]^2], x]$

[Out] $\operatorname{Sqrt}[a + b] \operatorname{ArcTanh}[\operatorname{Sqrt}[a + b \operatorname{Coth}[x]^2] / \operatorname{Sqrt}[a + b]] - \operatorname{Sqrt}[a + b \operatorname{Coth}[x]^2] - (a + b \operatorname{Coth}[x]^2)^{(3/2)} / (3*b)$

Rule 52

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(
b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 214

```
Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 457

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_
.), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p
*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[
b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{x^3\sqrt{a+bx^2}}{1-x^2} dx, x, \coth(x)\right) \\
&= \frac{1}{2}\text{Subst}\left(\int \frac{x\sqrt{a+bx}}{1-x} dx, x, \coth^2(x)\right) \\
&= -\frac{(a+b\coth^2(x))^{3/2}}{3b} + \frac{1}{2}\text{Subst}\left(\int \frac{\sqrt{a+bx}}{1-x} dx, x, \coth^2(x)\right) \\
&= -\sqrt{a+b\coth^2(x)} - \frac{(a+b\coth^2(x))^{3/2}}{3b} \\
&\quad + \frac{1}{2}(a+b)\text{Subst}\left(\int \frac{1}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x)\right) \\
&= -\sqrt{a+b\coth^2(x)} - \frac{(a+b\coth^2(x))^{3/2}}{3b} + \frac{(a+b)\text{Subst}\left(\int \frac{1}{1+\frac{a}{b}-\frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)}\right)}{b} \\
&= \sqrt{a+b}\text{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right) - \sqrt{a+b\coth^2(x)} - \frac{(a+b\coth^2(x))^{3/2}}{3b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.95

$$\begin{aligned}
\int \coth^3(x)\sqrt{a+b\coth^2(x)} dx &= \sqrt{a+b}\text{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right) \\
&\quad - \frac{\sqrt{a+b\coth^2(x)}(a+3b+b\coth^2(x))}{3b}
\end{aligned}$$

[In] Integrate[Coth[x]^3*Sqrt[a + b*Coth[x]^2],x]

[Out] Sqrt[a + b]*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]] - (Sqrt[a + b*Coth[x]^2]*(a + 3*b + b*Coth[x]^2))/(3*b)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 252 vs. 2(51) = 102.

Time = 0.21 (sec) , antiderivative size = 253, normalized size of antiderivative = 4.02

method	result
derivativedivides	$-\frac{(a+b \operatorname{coth}(x)^2)^{\frac{3}{2}}}{3b} - \frac{\sqrt{b(\operatorname{coth}(x)-1)^2+2b(\operatorname{coth}(x)-1)+a+b}}{2} - \frac{\sqrt{b} \ln\left(\frac{b(\operatorname{coth}(x)-1)+b}{\sqrt{b}} + \sqrt{b(\operatorname{coth}(x)-1)^2+2b(\operatorname{coth}(x)-1)+a+b}\right)}{2}$
default	$-\frac{(a+b \operatorname{coth}(x)^2)^{\frac{3}{2}}}{3b} - \frac{\sqrt{b(\operatorname{coth}(x)-1)^2+2b(\operatorname{coth}(x)-1)+a+b}}{2} - \frac{\sqrt{b} \ln\left(\frac{b(\operatorname{coth}(x)-1)+b}{\sqrt{b}} + \sqrt{b(\operatorname{coth}(x)-1)^2+2b(\operatorname{coth}(x)-1)+a+b}\right)}{2}$

[In] `int(coth(x)^3*(a+b*coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/3*(a+b*\operatorname{coth}(x)^2)^{(3/2)}/b-1/2*(b*(\operatorname{coth}(x)-1)^2+2*b*(\operatorname{coth}(x)-1)+a+b)^{(1/2)}-1/2*b^{(1/2)}*\ln((b*(\operatorname{coth}(x)-1)+b)/b^{(1/2)}+(b*(\operatorname{coth}(x)-1)^2+2*b*(\operatorname{coth}(x)-1)+a+b)^{(1/2)})+1/2*(a+b)^{(1/2)}*\ln((2*a+2*b+2*b*(\operatorname{coth}(x)-1)+2*(a+b)^{(1/2)}*(b*(\operatorname{coth}(x)-1)^2+2*b*(\operatorname{coth}(x)-1)+a+b)^{(1/2)})/(\operatorname{coth}(x)-1))-1/2*(b*(1+\operatorname{coth}(x))^2-2*b*(1+\operatorname{coth}(x))+a+b)^{(1/2)}+1/2*b^{(1/2)}*\ln((b*(1+\operatorname{coth}(x))-b)/b^{(1/2)}+(b*(1+\operatorname{coth}(x))^2-2*b*(1+\operatorname{coth}(x))+a+b)^{(1/2)})+1/2*(a+b)^{(1/2)}*\ln((2*a+2*b-2*b*(1+\operatorname{coth}(x))+2*(a+b)^{(1/2)}*(b*(1+\operatorname{coth}(x))^2-2*b*(1+\operatorname{coth}(x))+a+b)^{(1/2)})/(1+\operatorname{coth}(x))))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 903 vs. 2(51) = 102.

Time = 0.36 (sec) , antiderivative size = 2367, normalized size of antiderivative = 37.57

$$\int \operatorname{coth}^3(x) \sqrt{a + b \operatorname{coth}^2(x)} dx = \text{Too large to display}$$

[In] `integrate(coth(x)^3*(a+b*coth(x)^2)^(1/2),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & [1/12*(3*(b*\cosh(x)^6 + 6*b*\cosh(x)*\sinh(x)^5 + b*\sinh(x)^6 - 3*b*\cosh(x)^4 \\ & + 3*(5*b*\cosh(x)^2 - b)*\sinh(x)^4 + 4*(5*b*\cosh(x)^3 - 3*b*\cosh(x))*\sinh(x) \\ &)^3 + 3*b*\cosh(x)^2 + 3*(5*b*\cosh(x)^4 - 6*b*\cosh(x)^2 + b)*\sinh(x)^2 + 6*(\\ & b*\cosh(x)^5 - 2*b*\cosh(x)^3 + b*\cosh(x))*\sinh(x) - b)*\sqrt{a + b}*\log(-((a^ \\ & 3 + a^2*b)*\cosh(x)^8 + 8*(a^3 + a^2*b)*\cosh(x)*\sinh(x)^7 + (a^3 + a^2*b)*\si \\ & nh(x)^8 - 2*(2*a^3 + a^2*b)*\cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b) \\ &)*\cosh(x)^2)*\sinh(x)^6 + 4*(14*(a^3 + a^2*b)*\cosh(x)^3 - 3*(2*a^3 + a^2*b)*\c \\ & osh(x))*\sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*\cosh(x)^4 + (70*(a^3 + \\ & a^2*b)*\cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*\cosh(\\ & x)^2)*\sinh(x)^4 + 4*(14*(a^3 + a^2*b)*\cosh(x)^5 - 10*(2*a^3 + a^2*b)*\cosh(x) \\ &)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*\cosh(x))*\sinh(x)^3 + a^3 + 3*a^2*b + \\ & 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*\cosh(x)^2 + 2*(14*(a^3 + a^2*b)*\c \end{aligned}$$

$$\begin{aligned}
& \text{osh}(x)^6 - 15*(2*a^3 + a^2*b)*\text{cosh}(x)^4 - 2*a^3 - 3*a^2*b + b^3 + 3*(6*a^3 \\
& + 4*a^2*b - a*b^2 + b^3)*\text{cosh}(x)^2*\sinh(x)^2 + \sqrt{2}*(a^2*\text{cosh}(x)^6 + 6* \\
& a^2*\text{cosh}(x)*\sinh(x)^5 + a^2*\sinh(x)^6 - 3*a^2*\text{cosh}(x)^4 + 3*(5*a^2*\text{cosh}(x)^ \\
& 2 - a^2)*\sinh(x)^4 + 4*(5*a^2*\text{cosh}(x)^3 - 3*a^2*\text{cosh}(x))*\sinh(x)^3 + (3*a^2 \\
& + 2*a*b - b^2)*\text{cosh}(x)^2 + (15*a^2*\text{cosh}(x)^4 - 18*a^2*\text{cosh}(x)^2 + 3*a^2 + \\
& 2*a*b - b^2)*\sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*\text{cosh}(x)^5 - 6*a^2*\text{cosh} \\
& h(x)^3 + (3*a^2 + 2*a*b - b^2)*\text{cosh}(x))*\sinh(x))*\sqrt{a + b}*\sqrt{((a + b)* \\
& \text{cosh}(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\text{cosh}(x)^2 - 2*\text{cosh}(x)*\sinh(x) + \sinh \\
& h(x)^2))} + 4*(2*(a^3 + a^2*b)*\text{cosh}(x)^7 - 3*(2*a^3 + a^2*b)*\text{cosh}(x)^5 + (6* \\
& a^3 + 4*a^2*b - a*b^2 + b^3)*\text{cosh}(x)^3 - (2*a^3 + 3*a^2*b - b^3)*\text{cosh}(x))*\sinh(x) \\
&)/(\text{cosh}(x)^6 + 6*\text{cosh}(x)^5*\sinh(x) + 15*\text{cosh}(x)^4*\sinh(x)^2 + 20*\text{cosh} \\
& (x)^3*\sinh(x)^3 + 15*\text{cosh}(x)^2*\sinh(x)^4 + 6*\text{cosh}(x)*\sinh(x)^5 + \sinh(x)^6) \\
&) + 3*(b*\text{cosh}(x)^6 + 6*b*\text{cosh}(x)*\sinh(x)^5 + b*\sinh(x)^6 - 3*b*\text{cosh}(x)^4 + \\
& 3*(5*b*\text{cosh}(x)^2 - b)*\sinh(x)^4 + 4*(5*b*\text{cosh}(x)^3 - 3*b*\text{cosh}(x))*\sinh(x)^3 \\
& + 3*b*\text{cosh}(x)^2 + 3*(5*b*\text{cosh}(x)^4 - 6*b*\text{cosh}(x)^2 + b)*\sinh(x)^2 + 6*(b*\text{c} \\
& osh(x)^5 - 2*b*\text{cosh}(x)^3 + b*\text{cosh}(x))*\sinh(x) - b)*\sqrt{a + b}*\log(((a + b) \\
& *\text{cosh}(x)^4 + 4*(a + b)*\text{cosh}(x)*\sinh(x)^3 + (a + b)*\sinh(x)^4 + 2*b*\text{cosh}(x)^ \\
& 2 + 2*(3*(a + b)*\text{cosh}(x)^2 + b)*\sinh(x)^2 + \sqrt{2}*(\text{cosh}(x)^2 + 2*\text{cosh}(x)* \\
& \sinh(x) + \sinh(x)^2 + 1))*\sqrt{a + b}*\sqrt{((a + b)*\text{cosh}(x)^2 + (a + b)*\sinh \\
& (x)^2 - a + b)/(\text{cosh}(x)^2 - 2*\text{cosh}(x)*\sinh(x) + \sinh(x)^2))} + 4*((a + b)*\text{c} \\
& osh(x)^3 + b*\text{cosh}(x))*\sinh(x) + a + b)/(\text{cosh}(x)^2 + 2*\text{cosh}(x)*\sinh(x) + \sinh \\
& (x)^2)) - 4*\sqrt{2}*((a + 4*b)*\text{cosh}(x)^4 + 4*(a + 4*b)*\text{cosh}(x)*\sinh(x)^3 + \\
& (a + 4*b)*\sinh(x)^4 - 2*(a + 2*b)*\text{cosh}(x)^2 + 2*(3*(a + 4*b)*\text{cosh}(x)^2 - a \\
& - 2*b)*\sinh(x)^2 + 4*((a + 4*b)*\text{cosh}(x)^3 - (a + 2*b)*\text{cosh}(x))*\sinh(x) + a \\
& + 4*b)*\sqrt{((a + b)*\text{cosh}(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\text{cosh}(x)^2 - 2* \\
& \text{cosh}(x)*\sinh(x) + \sinh(x)^2))}/(b*\text{cosh}(x)^6 + 6*b*\text{cosh}(x)*\sinh(x)^5 + b*\sinh \\
& h(x)^6 - 3*b*\text{cosh}(x)^4 + 3*(5*b*\text{cosh}(x)^2 - b)*\sinh(x)^4 + 4*(5*b*\text{cosh}(x)^3 \\
& - 3*b*\text{cosh}(x))*\sinh(x)^3 + 3*b*\text{cosh}(x)^2 + 3*(5*b*\text{cosh}(x)^4 - 6*b*\text{cosh}(x)^ \\
& 2 + b)*\sinh(x)^2 + 6*(b*\text{cosh}(x)^5 - 2*b*\text{cosh}(x)^3 + b*\text{cosh}(x))*\sinh(x) - b) \\
& , -1/6*(3*(b*\text{cosh}(x)^6 + 6*b*\text{cosh}(x)*\sinh(x)^5 + b*\sinh(x)^6 - 3*b*\text{cosh}(x)^ \\
& 4 + 3*(5*b*\text{cosh}(x)^2 - b)*\sinh(x)^4 + 4*(5*b*\text{cosh}(x)^3 - 3*b*\text{cosh}(x))*\sinh \\
& (x)^3 + 3*b*\text{cosh}(x)^2 + 3*(5*b*\text{cosh}(x)^4 - 6*b*\text{cosh}(x)^2 + b)*\sinh(x)^2 + 6* \\
& (b*\text{cosh}(x)^5 - 2*b*\text{cosh}(x)^3 + b*\text{cosh}(x))*\sinh(x) - b)*\sqrt{-a - b}*\arctan(\\
& \sqrt{2}*(a*\text{cosh}(x)^2 + 2*a*\text{cosh}(x)*\sinh(x) + a*\sinh(x)^2 - a - b)*\sqrt{-a - \\
& b}*\sqrt{((a + b)*\text{cosh}(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\text{cosh}(x)^2 - 2*\text{cosh} \\
& h(x)*\sinh(x) + \sinh(x)^2))}/((a^2 + a*b)*\text{cosh}(x)^4 + 4*(a^2 + a*b)*\text{cosh}(x)*\sinh \\
& (x)^3 + (a^2 + a*b)*\sinh(x)^4 - (2*a^2 + a*b - b^2)*\text{cosh}(x)^2 + (6*(a^2 \\
& + a*b)*\text{cosh}(x)^2 - 2*a^2 - a*b + b^2)*\sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(2* \\
& (a^2 + a*b)*\text{cosh}(x)^3 - (2*a^2 + a*b - b^2)*\text{cosh}(x))*\sinh(x)) + 3*(b*\text{cosh} \\
& (x)^6 + 6*b*\text{cosh}(x)*\sinh(x)^5 + b*\sinh(x)^6 - 3*b*\text{cosh}(x)^4 + 3*(5*b*\text{cosh}(x) \\
& ^2 - b)*\sinh(x)^4 + 4*(5*b*\text{cosh}(x)^3 - 3*b*\text{cosh}(x))*\sinh(x)^3 + 3*b*\text{cosh}(x) \\
& ^2 + 3*(5*b*\text{cosh}(x)^4 - 6*b*\text{cosh}(x)^2 + b)*\sinh(x)^2 + 6*(b*\text{cosh}(x)^5 - 2*b \\
& *\text{cosh}(x)^3 + b*\text{cosh}(x))*\sinh(x) - b)*\sqrt{-a - b}*\arctan(\sqrt{2}*(\text{cosh}(x)^2 \\
& + 2*\text{cosh}(x)*\sinh(x) + \sinh(x)^2 + 1))*\sqrt{-a - b}*\sqrt{((a + b)*\text{cosh}(x)^2 \\
& + (a + b)*\sinh(x)^2 - a + b)/(\text{cosh}(x)^2 - 2*\text{cosh}(x)*\sinh(x) + \sinh(x)^2))}/(
\end{aligned}$$

```
(a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a
- b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cos
h(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)) + 2*sqrt(2)*((a + 4*b)*cosh(x)^
4 + 4*(a + 4*b)*cosh(x)*sinh(x)^3 + (a + 4*b)*sinh(x)^4 - 2*(a + 2*b)*cosh(
x)^2 + 2*(3*(a + 4*b)*cosh(x)^2 - a - 2*b)*sinh(x)^2 + 4*((a + 4*b)*cosh(x)
^3 - (a + 2*b)*cosh(x))*sinh(x) + a + 4*b)*sqrt(((a + b)*cosh(x)^2 + (a + b
)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/(b*cosh(
x)^6 + 6*b*cosh(x)*sinh(x)^5 + b*sinh(x)^6 - 3*b*cosh(x)^4 + 3*(5*b*cosh(x)
^2 - b)*sinh(x)^4 + 4*(5*b*cosh(x)^3 - 3*b*cosh(x))*sinh(x)^3 + 3*b*cosh(x)
^2 + 3*(5*b*cosh(x)^4 - 6*b*cosh(x)^2 + b)*sinh(x)^2 + 6*(b*cosh(x)^5 - 2*b
*cosh(x)^3 + b*cosh(x))*sinh(x) - b)]
```

Sympy [F]

$$\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx = \int \sqrt{a + b \coth^2(x)} \coth^3(x) dx$$

```
[In] integrate(coth(x)**3*(a+b*coth(x)**2)**(1/2),x)
```

```
[Out] Integral(sqrt(a + b*coth(x)**2)*coth(x)**3, x)
```

Maxima [F]

$$\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx = \int \sqrt{b \coth^2(x) + a} \coth^3(x) dx$$

```
[In] integrate(coth(x)^3*(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*coth(x)^2 + a)*coth(x)^3, x)
```

Giac [F(-2)]

Exception generated.

$$\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx = \text{Exception raised: TypeError}$$

```
[In] integrate(coth(x)^3*(a+b*coth(x)^2)^(1/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type
```

Mupad [B] (verification not implemented)

Time = 3.90 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.05

$$\int \coth^3(x) \sqrt{a + b \coth^2(x)} dx = -\sqrt{b \coth^2(x) + a} - \frac{(b \coth^2(x) + a)^{3/2}}{3b} - 2 \operatorname{atan} \left(\frac{2 \sqrt{b \coth^2(x) + a} \sqrt{-\frac{a}{4} - \frac{b}{4}}}{a + b} \right) \sqrt{-\frac{a}{4} - \frac{b}{4}}$$

[In] int(coth(x)^3*(a + b*coth(x)^2)^(1/2),x)

[Out] - (a + b*coth(x)^2)^(1/2) - (a + b*coth(x)^2)^(3/2)/(3*b) - 2*atan((2*(a + b*coth(x)^2)^(1/2)*(- a/4 - b/4)^(1/2))/(a + b))*(- a/4 - b/4)^(1/2)

3.17 $\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx$

Optimal result	133
Rubi [A] (verified)	133
Mathematica [B] (verified)	135
Maple [B] (verified)	136
Fricas [B] (verification not implemented)	136
Sympy [F]	140
Maxima [F]	140
Giac [F(-2)]	140
Mupad [F(-1)]	140

Optimal result

Integrand size = 17, antiderivative size = 85

$$\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx = -\frac{(a + 2b) \operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right)}{2\sqrt{b}} + \sqrt{a + b} \operatorname{arctanh}\left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right) - \frac{1}{2} \coth(x) \sqrt{a + b \coth^2(x)}$$

[Out] $-1/2*(a+2*b)*\operatorname{arctanh}(\coth(x)*b^{(1/2)}/(a+b*\coth(x)^2)^{(1/2)})/b^{(1/2)}+\operatorname{arctanh}(\coth(x)*(a+b)^{(1/2)}/(a+b*\coth(x)^2)^{(1/2)})*(a+b)^{(1/2)}-1/2*\coth(x)*(a+b*\coth(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3751, 489, 537, 223, 212, 385}

$$\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx = -\frac{(a + 2b) \operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right)}{2\sqrt{b}} + \sqrt{a + b} \operatorname{arctanh}\left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right) - \frac{1}{2} \coth(x) \sqrt{a + b \coth^2(x)}$$

[In] Int[Coth[x]^2*Sqrt[a + b*Coth[x]^2],x]

[Out] $-1/2*((a + 2*b)*\text{ArcTanh}[(\text{Sqrt}[b]*\text{Coth}[x])/\text{Sqrt}[a + b*\text{Coth}[x]^2]])/\text{Sqrt}[b] + \text{Sqrt}[a + b]*\text{ArcTanh}[(\text{Sqrt}[a + b]*\text{Coth}[x])/\text{Sqrt}[a + b*\text{Coth}[x]^2]] - (\text{Coth}[x]*\text{Sqrt}[a + b*\text{Coth}[x]^2])/2$

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 223

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 489

Int[((e_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[e^(n - 1)*(e*x)^(m - n + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^q/(b*(m + n*(p + q) + 1))), x] - Dist[e^n/(b*(m + n*(p + q) + 1)), Int[(e*x)^(m - n)*(a + b*x^n)^p*(c + d*x^n)^(q - 1)*Simp[a*c*(m - n + 1) + (a*d*(m - n + 1) - n*q*(b*c - a*d))*x^n, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && GtQ[q, 0] && GtQ[m - n + 1, 0] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

Rule 537

Int[((e_) + (f_.)*(x_)^(n_))/(((a_) + (b_.)*(x_)^(n_))*Sqrt[(c_) + (d_.)*(x_)^(n_)]), x_Symbol] := Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 3751

Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration

alQ[n]))

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{x^2\sqrt{a+bx^2}}{1-x^2} dx, x, \coth(x)\right) \\
&= -\frac{1}{2}\coth(x)\sqrt{a+b\coth^2(x)} + \frac{1}{2}\text{Subst}\left(\int \frac{a+(a+2b)x^2}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right) \\
&= -\frac{1}{2}\coth(x)\sqrt{a+b\coth^2(x)} + \frac{1}{2}(-a-2b)\text{Subst}\left(\int \frac{1}{\sqrt{a+bx^2}} dx, x, \coth(x)\right) \\
&\quad + (a+b)\text{Subst}\left(\int \frac{1}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right) \\
&= -\frac{1}{2}\coth(x)\sqrt{a+b\coth^2(x)} \\
&\quad + \frac{1}{2}(-a-2b)\text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{\coth(x)}{\sqrt{a+b\coth^2(x)}}\right) \\
&\quad + (a+b)\text{Subst}\left(\int \frac{1}{1-(a+b)x^2} dx, x, \frac{\coth(x)}{\sqrt{a+b\coth^2(x)}}\right) \\
&= -\frac{(a+2b)\text{arctanh}\left(\frac{\sqrt{b}\coth(x)}{\sqrt{a+b\coth^2(x)}}\right)}{2\sqrt{b}} \\
&\quad + \sqrt{a+b}\text{arctanh}\left(\frac{\sqrt{a+b}\coth(x)}{\sqrt{a+b\coth^2(x)}}\right) - \frac{1}{2}\coth(x)\sqrt{a+b\coth^2(x)}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 191 vs. 2(85) = 170.

Time = 0.84 (sec) , antiderivative size = 191, normalized size of antiderivative = 2.25

$$\int \coth^2(x)\sqrt{a+b\coth^2(x)} dx = \frac{\sqrt{(-a+b+(a+b)\cosh(2x))\text{csch}^2(x)}\left(\sqrt{2}\sqrt{a+b}(a+2b)\text{arctanh}\left(\frac{\sqrt{2}\sqrt{b}\cosh(x)}{\sqrt{-a+b+(a+b)\cosh(2x)}}\right) + \sqrt{b}\left(-2\sqrt{2}\right)\right)}{2\sqrt{2}\sqrt{b}\sqrt{a+b}\sqrt{-a+b}}$$

`[In] Integrate[Coth[x]^2*Sqrt[a + b*Coth[x]^2], x]`

```
[Out] -1/2*(Sqrt[(-a + b + (a + b)*Cosh[2*x])*Csch[x]^2]*(Sqrt[2]*Sqrt[a + b]*(a + 2*b)*ArcTanh[(Sqrt[2]*Sqrt[b]*Cosh[x])/Sqrt[-a + b + (a + b)*Cosh[2*x]]] + Sqrt[b]*(-2*Sqrt[2]*(a + b)*ArcTanh[(Sqrt[2]*Sqrt[a + b]*Cosh[x])/Sqrt[-a + b + (a + b)*Cosh[2*x]]] + Sqrt[a + b]*Sqrt[-a + b + (a + b)*Cosh[2*x])*Csch[x]*Csch[x]))*Sinh[x])/(Sqrt[2]*Sqrt[b]*Sqrt[a + b]*Sqrt[-a + b + (a + b)*Cosh[2*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 275 vs. 2(67) = 134.

Time = 0.14 (sec) , antiderivative size = 276, normalized size of antiderivative = 3.25

method	result
derivativedivides	$-\frac{\coth(x)\sqrt{a+b\coth(x)^2}}{2} - \frac{a \ln\left(\sqrt{b}\coth(x) + \sqrt{a+b\coth(x)^2}\right)}{2\sqrt{b}} - \frac{\sqrt{b(\coth(x)-1)^2 + 2b(\coth(x)-1) + a + b}}{2} - \frac{\sqrt{b} \ln}{\dots}$
default	$-\frac{\coth(x)\sqrt{a+b\coth(x)^2}}{2} - \frac{a \ln\left(\sqrt{b}\coth(x) + \sqrt{a+b\coth(x)^2}\right)}{2\sqrt{b}} - \frac{\sqrt{b(\coth(x)-1)^2 + 2b(\coth(x)-1) + a + b}}{2} - \frac{\sqrt{b} \ln}{\dots}$

```
[In] int(coth(x)^2*(a+b*coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -1/2*coth(x)*(a+b*coth(x)^2)^(1/2)-1/2*a/b^(1/2)*ln(b^(1/2)*coth(x)+(a+b*coth(x)^2)^(1/2))-1/2*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)-1/2*b^(1/2)*ln((b*(coth(x)-1)+b)/b^(1/2)+(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))+1/2*(a+b)^(1/2)*ln((2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(1/2)*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(coth(x)-1))+1/2*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)-1/2*b^(1/2)*ln((b*(1+coth(x))-b)/b^(1/2)+(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))-1/2*(a+b)^(1/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b)^(1/2)*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))/(1+coth(x)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 907 vs. 2(67) = 134.

Time = 0.43 (sec) , antiderivative size = 4877, normalized size of antiderivative = 57.38

$$\int \coth^2(x)\sqrt{a+b\coth^2(x)} dx = \text{Too large to display}$$

```
[In] integrate(coth(x)^2*(a+b*coth(x)^2)^(1/2),x, algorithm="fricas")
```

```
[Out] [1/4*((b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 - 2*b*cosh(x)^2 + 2*(3*b*cosh(x)^2 - b)*sinh(x)^2 + 4*(b*cosh(x)^3 - b*cosh(x))*sinh(x) + b)*sqrt(a + b)*log(((a*b^2 + b^3)*cosh(x)^8 + 8*(a*b^2 + b^3)*cosh(x)*sinh(x)^7 + (a*b^2 + b^3)*sinh(x)^8 + 2*(a*b^2 + 2*b^3)*cosh(x)^6 + 2*(a*b^2 + 2*b^3)*sinh(x)^6 + 2*(a*b^2 + 2*b^3)*cosh(x)^4 + 2*(a*b^2 + 2*b^3)*sinh(x)^4 + 2*(a*b^2 + 2*b^3)*cosh(x)^2 + 2*(a*b^2 + 2*b^3)*sinh(x)^2 + a*b^2 + b^3))
```


$$\begin{aligned}
& 3 + 14*(a*b^2 + b^3)*\cosh(x)^2*\sinh(x)^6 + 4*(14*(a*b^2 + b^3)*\cosh(x)^3 + \\
& 3*(a*b^2 + 2*b^3)*\cosh(x))*\sinh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*\cos \\
& h(x)^4 + (70*(a*b^2 + b^3)*\cosh(x)^4 + a^3 - a^2*b + 4*a*b^2 + 6*b^3 + 30*(\\
& a*b^2 + 2*b^3)*\cosh(x)^2)*\sinh(x)^4 + 4*(14*(a*b^2 + b^3)*\cosh(x)^5 + 10*(a \\
& *b^2 + 2*b^3)*\cosh(x)^3 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*\cosh(x))*\sinh(x)^ \\
& 3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 - 3*a*b^2 - 2*b^3)*\cosh(x)^2 + 2 \\
& *(14*(a*b^2 + b^3)*\cosh(x)^6 + 15*(a*b^2 + 2*b^3)*\cosh(x)^4 - a^3 + 3*a*b^2 \\
& + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 + 6*b^3)*\cosh(x)^2)*\sinh(x)^2 + \sqrt{2} \\
& *(b^2*\cosh(x)^6 + 6*b^2*\cosh(x)*\sinh(x)^5 + b^2*\sinh(x)^6 + 3*b^2*\cosh(x)^4 \\
& + 3*(5*b^2*\cosh(x)^2 + b^2)*\sinh(x)^4 + 4*(5*b^2*\cosh(x)^3 + 3*b^2*\cosh(x) \\
&)*\sinh(x)^3 - (a^2 - 2*a*b - 3*b^2)*\cosh(x)^2 + (15*b^2*\cosh(x)^4 + 18*b^2* \\
& \cosh(x)^2 - a^2 + 2*a*b + 3*b^2)*\sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(3*b^2*c \\
& osh(x)^5 + 6*b^2*\cosh(x)^3 - (a^2 - 2*a*b - 3*b^2)*\cosh(x))*\sinh(x))*\sqrt{a \\
& + b}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*c \\
& osh(x)*\sinh(x) + \sinh(x)^2))} + 4*(2*(a*b^2 + b^3)*\cosh(x)^7 + 3*(a*b^2 + 2* \\
& b^3)*\cosh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*\cosh(x)^3 - (a^3 - 3*a*b^2 \\
& - 2*b^3)*\cosh(x))*\sinh(x))/(\cosh(x)^6 + 6*\cosh(x)^5*\sinh(x) + 15*\cosh(x)^4 \\
& *\sinh(x)^2 + 20*\cosh(x)^3*\sinh(x)^3 + 15*\cosh(x)^2*\sinh(x)^4 + 6*\cosh(x)*\si \\
& nh(x)^5 + \sinh(x)^6)) + ((a + 2*b)*\cosh(x)^4 + 4*(a + 2*b)*\cosh(x)*\sinh(x)^ \\
& 3 + (a + 2*b)*\sinh(x)^4 - 2*(a + 2*b)*\cosh(x)^2 + 2*(3*(a + 2*b)*\cosh(x)^2 \\
& - a - 2*b)*\sinh(x)^2 + 4*((a + 2*b)*\cosh(x)^3 - (a + 2*b)*\cosh(x))*\sinh(x) \\
& + a + 2*b)*\sqrt{b}*\log(-((a + 2*b)*\cosh(x)^4 + 4*(a + 2*b)*\cosh(x)*\sinh(x)^ \\
& 3 + (a + 2*b)*\sinh(x)^4 - 2*(a - 2*b)*\cosh(x)^2 + 2*(3*(a + 2*b)*\cosh(x)^2 \\
& - a + 2*b)*\sinh(x)^2 - 2*\sqrt{2}*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 \\
& + 1)*\sqrt{b}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x) \\
& ^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2))} + 4*((a + 2*b)*\cosh(x)^3 - (a - 2*b)*c \\
& osh(x))*\sinh(x) + a + 2*b)/(\cosh(x)^4 + 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4 + 2 \\
& *(3*\cosh(x)^2 - 1)*\sinh(x)^2 - 2*\cosh(x)^2 + 4*(\cosh(x)^3 - \cosh(x))*\sinh(x) \\
&) + 1)) + (b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 - 2*b*\cosh(x)^ \\
& 2 + 2*(3*b*\cosh(x)^2 - b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 - b*\cosh(x))*\sinh(x) + \\
& b)*\sqrt{a + b}*\log(-((a + b)*\cosh(x)^4 + 4*(a + b)*\cosh(x)*\sinh(x)^3 + (a \\
& + b)*\sinh(x)^4 - 2*a*\cosh(x)^2 + 2*(3*(a + b)*\cosh(x)^2 - a)*\sinh(x)^2 + sq \\
& rt(2)*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 - 1)*\sqrt{a + b}*\sqrt{((a \\
& + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) \\
& + \sinh(x)^2))} + 4*((a + b)*\cosh(x)^3 - a*\cosh(x))*\sinh(x) + a + b)/(\cosh(x) \\
& ^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2)) - 2*\sqrt{2}*(b*\cosh(x)^2 + 2*b*\cosh(x) \\
& *\sinh(x) + b*\sinh(x)^2 + b)*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a \\
& + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)))/(b*\cosh(x)^4 + 4*b*\cosh \\
& (x)*\sinh(x)^3 + b*\sinh(x)^4 - 2*b*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 - b)*\sinh(x) \\
& ^2 + 4*(b*\cosh(x)^3 - b*\cosh(x))*\sinh(x) + b), 1/4*(2*((a + 2*b)*\cosh(x)^4 \\
& + 4*(a + 2*b)*\cosh(x)*\sinh(x)^3 + (a + 2*b)*\sinh(x)^4 - 2*(a + 2*b)*\cosh(x) \\
& ^2 + 2*(3*(a + 2*b)*\cosh(x)^2 - a - 2*b)*\sinh(x)^2 + 4*((a + 2*b)*\cosh(x)^3 \\
& - (a + 2*b)*\cosh(x))*\sinh(x) + a + 2*b)*\sqrt{-b}*\arctan(\sqrt{2}*(\cosh(x)^2 \\
& + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 + 1)*\sqrt{-b}*\sqrt{((a + b)*\cosh(x)^2 + (a \\
& + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)))/(a +
\end{aligned}$$

$$\begin{aligned}
& b) \cosh(x)^4 + 4(a+b)\cosh(x)\sinh(x)^3 + (a+b)\sinh(x)^4 - 2(a-b) \\
& * \cosh(x)^2 + 2(3(a+b)\cosh(x)^2 - a+b)\sinh(x)^2 + 4((a+b)\cosh(x) \\
& ^3 - (a-b)\cosh(x))\sinh(x) + a+b) + (b\cosh(x)^4 + 4b\cosh(x)\sinh(x) \\
&)^3 + b\sinh(x)^4 - 2b\cosh(x)^2 + 2(3b\cosh(x)^2 - b)\sinh(x)^2 + 4(b* \\
& \cosh(x)^3 - b\cosh(x))\sinh(x) + b)\sqrt{a+b}\log(((a*b^2 + b^3)\cosh(x)^ \\
& 8 + 8(a*b^2 + b^3)\cosh(x)\sinh(x)^7 + (a*b^2 + b^3)\sinh(x)^8 + 2(a*b^2 \\
& + 2*b^3)\cosh(x)^6 + 2(a*b^2 + 2*b^3 + 14*(a*b^2 + b^3)\cosh(x)^2)\sinh(x) \\
& ^6 + 4*(14*(a*b^2 + b^3)\cosh(x)^3 + 3*(a*b^2 + 2*b^3)\cosh(x))\sinh(x)^5 + \\
& (a^3 - a^2*b + 4*a*b^2 + 6*b^3)\cosh(x)^4 + (70*(a*b^2 + b^3)\cosh(x)^4 + \\
& a^3 - a^2*b + 4*a*b^2 + 6*b^3 + 30*(a*b^2 + 2*b^3)\cosh(x)^2)\sinh(x)^4 + 4 \\
& *(14*(a*b^2 + b^3)\cosh(x)^5 + 10*(a*b^2 + 2*b^3)\cosh(x)^3 + (a^3 - a^2*b \\
& + 4*a*b^2 + 6*b^3)\cosh(x))\sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(\\
& a^3 - 3*a*b^2 - 2*b^3)\cosh(x)^2 + 2*(14*(a*b^2 + b^3)\cosh(x)^6 + 15*(a*b^ \\
& 2 + 2*b^3)\cosh(x)^4 - a^3 + 3*a*b^2 + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 + 6 \\
& *b^3)\cosh(x)^2)\sinh(x)^2 + \sqrt{2}*(b^2*\cosh(x)^6 + 6*b^2*\cosh(x)\sinh(x) \\
& ^5 + b^2*\sinh(x)^6 + 3*b^2*\cosh(x)^4 + 3*(5*b^2*\cosh(x)^2 + b^2)\sinh(x)^4 \\
& + 4*(5*b^2*\cosh(x)^3 + 3*b^2*\cosh(x))\sinh(x)^3 - (a^2 - 2*a*b - 3*b^2)*\cos \\
& h(x)^2 + (15*b^2*\cosh(x)^4 + 18*b^2*\cosh(x)^2 - a^2 + 2*a*b + 3*b^2)\sinh(x) \\
&)^2 + a^2 + 2*a*b + b^2 + 2*(3*b^2*\cosh(x)^5 + 6*b^2*\cosh(x)^3 - (a^2 - 2*a \\
& *b - 3*b^2)\cosh(x))\sinh(x))\sqrt{a+b}\sqrt{((a+b)\cosh(x)^2 + (a+b) \\
& *\sinh(x)^2 - a+b)/(\cosh(x)^2 - 2*\cosh(x)\sinh(x) + \sinh(x)^2))} + 4*(2*(a* \\
& b^2 + b^3)\cosh(x)^7 + 3*(a*b^2 + 2*b^3)\cosh(x)^5 + (a^3 - a^2*b + 4*a*b^2 \\
& + 6*b^3)\cosh(x)^3 - (a^3 - 3*a*b^2 - 2*b^3)\cosh(x))\sinh(x))/(\cosh(x)^6 \\
& + 6*\cosh(x)^5*\sinh(x) + 15*\cosh(x)^4*\sinh(x)^2 + 20*\cosh(x)^3*\sinh(x)^3 + 1 \\
& 5*\cosh(x)^2*\sinh(x)^4 + 6*\cosh(x)\sinh(x)^5 + \sinh(x)^6) + (b*\cosh(x)^4 + \\
& 4*b*\cosh(x)\sinh(x)^3 + b*\sinh(x)^4 - 2*b*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 - b) \\
& *\sinh(x)^2 + 4*(b*\cosh(x)^3 - b*\cosh(x))\sinh(x) + b)\sqrt{a+b}\log(-((a \\
& + b)\cosh(x)^4 + 4*(a+b)\cosh(x)\sinh(x)^3 + (a+b)\sinh(x)^4 - 2*a*\cosh \\
& (x)^2 + 2*(3*(a+b)\cosh(x)^2 - a)\sinh(x)^2 + \sqrt{2}*(\cosh(x)^2 + 2*\cosh \\
& (x)\sinh(x) + \sinh(x)^2 - 1))\sqrt{a+b}\sqrt{((a+b)\cosh(x)^2 + (a+b)* \\
& \sinh(x)^2 - a+b)/(\cosh(x)^2 - 2*\cosh(x)\sinh(x) + \sinh(x)^2))} + 4*((a+b) \\
&)\cosh(x)^3 - a*\cosh(x))\sinh(x) + a+b)/(\cosh(x)^2 + 2*\cosh(x)\sinh(x) + \\
& \sinh(x)^2)) - 2*\sqrt{2}*(b*\cosh(x)^2 + 2*b*\cosh(x)\sinh(x) + b*\sinh(x)^2 + \\
& b)\sqrt{((a+b)\cosh(x)^2 + (a+b)\sinh(x)^2 - a+b)/(\cosh(x)^2 - 2*\cosh \\
& (x)\sinh(x) + \sinh(x)^2))}/(b*\cosh(x)^4 + 4*b*\cosh(x)\sinh(x)^3 + b*\sinh(x) \\
& ^4 - 2*b*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 - b)\sinh(x)^2 + 4*(b*\cosh(x)^3 - b*c \\
& osh(x))\sinh(x) + b), -1/4*(2*(b*\cosh(x)^4 + 4*b*\cosh(x)\sinh(x)^3 + b*\sinh \\
& (x)^4 - 2*b*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 - b)\sinh(x)^2 + 4*(b*\cosh(x)^3 - \\
& b*\cosh(x))\sinh(x) + b)\sqrt{-a-b}\arctan(\sqrt{2}*(b*\cosh(x)^2 + 2*b*\cosh \\
& (x)\sinh(x) + b*\sinh(x)^2 + a+b)\sqrt{-a-b}\sqrt{((a+b)\cosh(x)^2 + (\\
& a+b)\sinh(x)^2 - a+b)/(\cosh(x)^2 - 2*\cosh(x)\sinh(x) + \sinh(x)^2))}/((a* \\
& b + b^2)\cosh(x)^4 + 4*(a*b + b^2)\cosh(x)\sinh(x)^3 + (a*b + b^2)\sinh(x)^ \\
& 4 - (a^2 - a*b - 2*b^2)\cosh(x)^2 + (6*(a*b + b^2)\cosh(x)^2 - a^2 + a*b + \\
& 2*b^2)\sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(2*(a*b + b^2)\cosh(x)^3 - (a^2 - \\
& a*b - 2*b^2)\cosh(x))\sinh(x))) + 2*(b*\cosh(x)^4 + 4*b*\cosh(x)\sinh(x)^3 +
\end{aligned}$$

$\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)) / (b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 - 2*b*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 - b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 - b*\cosh(x))*\sinh(x) + b)$

Sympy [F]

$$\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx = \int \sqrt{a + b \coth^2(x)} \coth^2(x) dx$$

[In] integrate(coth(x)**2*(a+b*coth(x)**2)**(1/2),x)

[Out] Integral(sqrt(a + b*coth(x)**2)*coth(x)**2, x)

Maxima [F]

$$\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx = \int \sqrt{b \coth(x)^2 + a} \coth(x)^2 dx$$

[In] integrate(coth(x)^2*(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*coth(x)^2 + a)*coth(x)^2, x)

Giac [F(-2)]

Exception generated.

$$\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx = \text{Exception raised: TypeError}$$

[In] integrate(coth(x)^2*(a+b*coth(x)^2)^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx);;OUTPUT>Error: Bad Argument Type

Mupad [F(-1)]

Timed out.

$$\int \coth^2(x) \sqrt{a + b \coth^2(x)} dx = \int \coth(x)^2 \sqrt{b \coth(x)^2 + a} dx$$

[In] int(coth(x)^2*(a + b*coth(x)^2)^(1/2),x)

[Out] int(coth(x)^2*(a + b*coth(x)^2)^(1/2), x)

3.18 $\int \coth(x) \sqrt{a + b \coth^2(x)} dx$

Optimal result	141
Rubi [A] (verified)	141
Mathematica [A] (verified)	143
Maple [B] (verified)	143
Fricas [B] (verification not implemented)	144
Sympy [F]	145
Maxima [F]	145
Giac [F(-2)]	145
Mupad [B] (verification not implemented)	146

Optimal result

Integrand size = 15, antiderivative size = 44

$$\int \coth(x) \sqrt{a + b \coth^2(x)} dx = \sqrt{a + b} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right) - \sqrt{a + b \coth^2(x)}$$

[Out] $\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2)/(a+b)^{(1/2)})}*(a+b)^{(1/2)}-(a+b*\coth(x)^2)^{(1/2)})$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3751, 455, 52, 65, 214}

$$\int \coth(x) \sqrt{a + b \coth^2(x)} dx = \sqrt{a + b} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right) - \sqrt{a + b \coth^2(x)}$$

[In] $\operatorname{Int}[\operatorname{Coth}[x]*\operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2], x]$

[Out] $\operatorname{Sqrt}[a + b]*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2]/\operatorname{Sqrt}[a + b]] - \operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2]$

Rule 52

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_. + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)*((c + d*x)^n/(b*(m + n + 1))}, x] + \operatorname{Dist}[n*((b*c - a*d)/(b*(m + n + 1))], \operatorname{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d\}, x \ \&\& \operatorname{NeQ}[b*c - a*d, 0] \ \&\& \operatorname{GtQ}[n, 0] \ \&\& \operatorname{NeQ}[m + n + 1, 0] \ \&\& \operatorname{!}(\operatorname{IGtQ}$

`[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0])) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 65

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 214

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Rule 455

`Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Dist[1/n, Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n + 1, 0]`

Rule 3751

`Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst} \left(\int \frac{x\sqrt{a+bx^2}}{1-x^2} dx, x, \coth(x) \right) \\
 &= \frac{1}{2} \text{Subst} \left(\int \frac{\sqrt{a+bx}}{1-x} dx, x, \coth^2(x) \right) \\
 &= -\sqrt{a+b\coth^2(x)} + \frac{1}{2}(a+b) \text{Subst} \left(\int \frac{1}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x) \right) \\
 &= -\sqrt{a+b\coth^2(x)} + \frac{(a+b) \text{Subst} \left(\int \frac{1}{1+\frac{a}{b}-\frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)} \right)}{b}
 \end{aligned}$$

$$= \sqrt{a+b} \operatorname{arctanh} \left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}} \right) - \sqrt{a+b \coth^2(x)}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00

$$\int \coth(x) \sqrt{a+b \coth^2(x)} dx = \sqrt{a+b} \operatorname{arctanh} \left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}} \right) - \sqrt{a+b \coth^2(x)}$$

[In] Integrate[Coth[x]*Sqrt[a + b*Coth[x]^2],x]

[Out] Sqrt[a + b]*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]] - Sqrt[a + b*Coth[x]^2]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 237 vs. 2(36) = 72.

Time = 0.09 (sec) , antiderivative size = 238, normalized size of antiderivative = 5.41

method	result
derivativedivides	$-\frac{\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{2} + \frac{\sqrt{b} \ln\left(\frac{b(1+\coth(x))-b}{\sqrt{b}} + \sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}\right)}{2} + \dots$
default	$-\frac{\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{2} + \frac{\sqrt{b} \ln\left(\frac{b(1+\coth(x))-b}{\sqrt{b}} + \sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}\right)}{2} + \dots$

[In] int(coth(x)*(a+b*coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out]
$$-1/2*(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^(1/2)+1/2*b^(1/2)*\ln((b*(1+\coth(x))-b)/b^(1/2)+(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^(1/2))+1/2*(a+b)^(1/2)*\ln((2*a+2*b-2*b*(1+\coth(x))+2*(a+b)^(1/2)*(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^(1/2))/(1+\coth(x)))-1/2*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^(1/2)-1/2*b^(1/2)*\ln((b*(\coth(x)-1)+b)/b^(1/2)+(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^(1/2))+1/2*(a+b)^(1/2)*\ln((2*a+2*b+2*b*(\coth(x)-1)+2*(a+b)^(1/2)*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^(1/2)))/(\coth(x)-1))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 495 vs. $2(36) = 72$.

Time = 0.31 (sec) , antiderivative size = 1551, normalized size of antiderivative = 35.25

$$\int \coth(x) \sqrt{a + b \coth^2(x)} dx = \text{Too large to display}$$

[In] integrate(coth(x)*(a+b*coth(x)^2)^(1/2),x, algorithm="fricas")

[Out] $\frac{1}{4} \left((\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1) \sqrt{a+b} \log\left(-\left(a^3 + a^2 b\right) \cosh(x)^8 + 8 \left(a^3 + a^2 b\right) \cosh(x) \sinh(x)^7 + \left(a^3 + a^2 b\right) \sinh(x)^8 - 2 \left(2 a^3 + a^2 b\right) \cosh(x)^6 - 2 \left(2 a^3 + a^2 b - 14 \left(a^3 + a^2 b\right) \cosh(x)^2\right) \sinh(x)^6 + 4 \left(14 \left(a^3 + a^2 b\right) \cosh(x)^3 - 3 \left(2 a^3 + a^2 b\right) \cosh(x) \sinh(x)^5 + \left(6 a^3 + 4 a^2 b - a b^2 + b^3\right) \cosh(x)^4 + \left(70 \left(a^3 + a^2 b\right) \cosh(x)^4 + 6 a^3 + 4 a^2 b - a b^2 + b^3 - 30 \left(2 a^3 + a^2 b\right) \cosh(x)^2\right) \sinh(x)^4 + 4 \left(14 \left(a^3 + a^2 b\right) \cosh(x)^5 - 10 \left(2 a^3 + a^2 b\right) \cosh(x)^3 + \left(6 a^3 + 4 a^2 b - a b^2 + b^3\right) \cosh(x) \sinh(x)^3 + a^3 + 3 a^2 b + 3 a b^2 + b^3 - 2 \left(2 a^3 + 3 a^2 b - b^3\right) \cosh(x)^2 + 2 \left(14 \left(a^3 + a^2 b\right) \cosh(x)^6 - 15 \left(2 a^3 + a^2 b\right) \cosh(x)^4 - 2 a^3 - 3 a^2 b + b^3 + 3 \left(6 a^3 + 4 a^2 b - a b^2 + b^3\right) \cosh(x)^2 \sinh(x)^2 + \sqrt{2} \left(a^2 \cosh(x)^6 + 6 a^2 \cosh(x) \sinh(x)^5 + a^2 \sinh(x)^6 - 3 a^2 \cosh(x)^4 + 3 \left(5 a^2 \cosh(x)^2 - a^2\right) \sinh(x)^4 + 4 \left(5 a^2 \cosh(x)^3 - 3 a^2 \cosh(x)\right) \sinh(x)^3 + \left(3 a^2 + 2 a b - b^2\right) \cosh(x)^2 + \left(15 a^2 \cosh(x)^4 - 18 a^2 \cosh(x)^2 + 3 a^2 + 2 a b - b^2\right) \sinh(x)^2 - a^2 - 2 a b - b^2 + 2 \left(3 a^2 \cosh(x)^5 - 6 a^2 \cosh(x)^3 + \left(3 a^2 + 2 a b - b^2\right) \cosh(x)\right) \sinh(x)\right) \sqrt{a+b} \sqrt{\left(\left(a+b\right) \cosh(x)^2 + \left(a+b\right) \sinh(x)^2 - a+b\right) / \left(\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2\right)} + 4 \left(2 \left(a^3 + a^2 b\right) \cosh(x)^7 - 3 \left(2 a^3 + a^2 b\right) \cosh(x)^5 + \left(6 a^3 + 4 a^2 b - a b^2 + b^3\right) \cosh(x)^3 - \left(2 a^3 + 3 a^2 b - b^3\right) \cosh(x) \sinh(x)\right) / \left(\cosh(x)^6 + 6 \cosh(x)^5 \sinh(x) + 15 \cosh(x)^4 \sinh(x)^2 + 20 \cosh(x)^3 \sinh(x)^3 + 15 \cosh(x)^2 \sinh(x)^4 + 6 \cosh(x) \sinh(x)^5 + \sinh(x)^6\right) + \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1\right) \sqrt{a+b} \log\left(\left(a+b\right) \cosh(x)^4 + 4 \left(a+b\right) \cosh(x) \sinh(x)^3 + \left(a+b\right) \sinh(x)^4 + 2 b \cosh(x)^2 + 2 \left(3 \left(a+b\right) \cosh(x)^2 + b\right) \sinh(x)^2 + \sqrt{2} \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1\right) \sqrt{a+b} \sqrt{\left(\left(a+b\right) \cosh(x)^2 + \left(a+b\right) \sinh(x)^2 - a+b\right) / \left(\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2\right)} + 4 \left(\left(a+b\right) \cosh(x)^3 + b \cosh(x)\right) \sinh(x) + a+b\right) / \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2\right) - 4 \sqrt{2} \sqrt{\left(\left(a+b\right) \cosh(x)^2 + \left(a+b\right) \sinh(x)^2 - a+b\right) / \left(\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2\right)} / \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1\right), -1/2 \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1\right) \sqrt{-a-b} \arctan\left(\sqrt{2} \left(a \cosh(x)^2 + 2 a \cosh(x) \sinh(x) + a \sinh(x)^2 - a - b\right) \sqrt{-a-b} \sqrt{\left(\left(a+b\right) \cosh(x)^2 + \left(a+b\right) \sinh(x)^2 - a+b\right) / \left(\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2\right)} / \left(\left(a^2 + a b\right) \cosh(x)^4 + 4 \left(a^2 + a b\right) \cosh(x) \sinh(x)^3 + \left(a^2 + a b\right) \sinh(x)^4 - \left(2 a^2 + a b - b^2\right) \cosh(x)^2 + \left(6 \left(a^2 + a b\right) \cosh(x)^2 - 2 a^2 - a b + b^2\right) \sinh(x)^2 + a^2 + 2\right)$


```
*a*b + b^2 + 2*(2*(a^2 + a*b)*cosh(x)^3 - (2*a^2 + a*b - b^2)*cosh(x))*sinh(x)) + (cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(-a - b)*arctan(sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(-a - b)*sqrt((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)) + 2*sqrt(2)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)]
```

Sympy [F]

$$\int \coth(x) \sqrt{a + b \coth^2(x)} dx = \int \sqrt{a + b \coth^2(x)} \coth(x) dx$$

```
[In] integrate(coth(x)*(a+b*coth(x)**2)**(1/2),x)
```

```
[Out] Integral(sqrt(a + b*coth(x)**2)*coth(x), x)
```

Maxima [F]

$$\int \coth(x) \sqrt{a + b \coth^2(x)} dx = \int \sqrt{b \coth(x)^2 + a} \coth(x) dx$$

```
[In] integrate(coth(x)*(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*coth(x)^2 + a)*coth(x), x)
```

Giac [F(-2)]

Exception generated.

$$\int \coth(x) \sqrt{a + b \coth^2(x)} dx = \text{Exception raised: TypeError}$$

```
[In] integrate(coth(x)*(a+b*coth(x)^2)^(1/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type
```

Mupad [B] (verification not implemented)

Time = 2.34 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.16

$$\int \coth(x) \sqrt{a + b \coth^2(x)} dx = -\sqrt{b \coth(x)^2 + a} - 2 \operatorname{atan} \left(\frac{2 \sqrt{b \coth(x)^2 + a} \sqrt{-\frac{a}{4} - \frac{b}{4}}}{a + b} \right) \sqrt{-\frac{a}{4} - \frac{b}{4}}$$

`[In] int(coth(x)*(a + b*coth(x)^2)^(1/2),x)`

```
[Out] - (a + b*coth(x)^2)^(1/2) - 2*atan((2*(a + b*coth(x)^2)^(1/2)*(- a/4 - b/4)^(1/2))/(a + b))*(- a/4 - b/4)^(1/2)
```

3.19 $\int \sqrt{a + b \coth^2(x)} dx$

Optimal result	147
Rubi [A] (verified)	147
Mathematica [A] (verified)	149
Maple [B] (verified)	149
Fricas [B] (verification not implemented)	150
Sympy [F]	152
Maxima [F]	152
Giac [B] (verification not implemented)	153
Mupad [F(-1)]	153

Optimal result

Integrand size = 12, antiderivative size = 60

$$\int \sqrt{a + b \coth^2(x)} dx = -\sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) + \sqrt{a + b} \operatorname{arctanh} \left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right)$$

[Out] $-\operatorname{arctanh}(\coth(x) \cdot b^{(1/2)} / (a + b \cdot \coth(x)^2)^{(1/2)}) \cdot b^{(1/2)} + \operatorname{arctanh}(\coth(x) \cdot (a + b)^{(1/2)} / (a + b \cdot \coth(x)^2)^{(1/2)}) \cdot (a + b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3742, 399, 223, 212, 385}

$$\int \sqrt{a + b \coth^2(x)} dx = \sqrt{a + b} \operatorname{arctanh} \left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) - \sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right)$$

[In] `Int[Sqrt[a + b*Coth[x]^2],x]`

[Out] $-(\operatorname{Sqrt}[b] \cdot \operatorname{ArcTanh}[(\operatorname{Sqrt}[b] \cdot \operatorname{Coth}[x]) / \operatorname{Sqrt}[a + b \cdot \operatorname{Coth}[x]^2]]) + \operatorname{Sqrt}[a + b] \cdot \operatorname{ArcTanh}[(\operatorname{Sqrt}[a + b] \cdot \operatorname{Coth}[x]) / \operatorname{Sqrt}[a + b \cdot \operatorname{Coth}[x]^2]]$

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 223

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x],
x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Su
bst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b
, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 399

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Di
st[b/d, Int[(a + b*x^n)^(p - 1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^
n)^(p - 1)/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*
d, 0] && EqQ[n*(p - 1) + 1, 0] && IntegerQ[n]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :=
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(
ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst} \left(\int \frac{\sqrt{a + bx^2}}{1 - x^2} dx, x, \coth(x) \right) \\ &= - \left((-a - b) \text{Subst} \left(\int \frac{1}{(1 - x^2) \sqrt{a + bx^2}} dx, x, \coth(x) \right) \right) \\ &\quad - b \text{Subst} \left(\int \frac{1}{\sqrt{a + bx^2}} dx, x, \coth(x) \right) \end{aligned}$$

$$\begin{aligned}
&= - \left((-a - b) \text{Subst} \left(\int \frac{1}{1 - (a + b)x^2} dx, x, \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} \right) \right) \\
&\quad - b \text{Subst} \left(\int \frac{1}{1 - bx^2} dx, x, \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} \right) \\
&= -\sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) + \sqrt{a + b} \operatorname{arctanh} \left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.37

$$\begin{aligned}
\int \sqrt{a + b \coth^2(x)} dx &= \sqrt{-a - b} \arctan \left(\frac{\coth(x) \sqrt{a + b \coth^2(x)} - \sqrt{b} \operatorname{csch}^2(x)}{\sqrt{-a - b}} \right) \\
&\quad + \sqrt{b} \log \left(-\sqrt{b} \coth(x) + \sqrt{a + b \coth^2(x)} \right)
\end{aligned}$$

[In] Integrate[Sqrt[a + b*Coth[x]^2],x]

[Out] Sqrt[-a - b]*ArcTan[(Coth[x]*Sqrt[a + b*Coth[x]^2] - Sqrt[b]*Csch[x]^2)/Sqrt[-a - b]] + Sqrt[b]*Log[-(Sqrt[b]*Coth[x]) + Sqrt[a + b*Coth[x]^2]]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 237 vs. 2(48) = 96.

Time = 0.14 (sec) , antiderivative size = 238, normalized size of antiderivative = 3.97

method	result
derivativedivides	$\frac{\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{2} - \frac{\sqrt{b} \ln\left(\frac{b(1+\coth(x))-b}{\sqrt{b}} + \sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}\right)}{2} - \frac{\sqrt{a+b}}{2}$
default	$\frac{\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{2} - \frac{\sqrt{b} \ln\left(\frac{b(1+\coth(x))-b}{\sqrt{b}} + \sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}\right)}{2} - \frac{\sqrt{a+b}}{2}$

[In] int((a+b*coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/2*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)-1/2*b^(1/2)*ln((b*(1+coth(x))-b)/b^(1/2)+(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))-1/2*(a+b)^(1/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b)^(1/2)*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))/(1+coth(x)))-1/2*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)-1/2

$$2*b^{(1/2)}*\ln((b*(\coth(x)-1)+b)/b^{(1/2)}+(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{(1/2)})+1/2*(a+b)^{(1/2)}*\ln((2*a+2*b+2*b*(\coth(x)-1)+2*(a+b)^{(1/2)}*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{(1/2)})/(\coth(x)-1))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 552 vs. $2(48) = 96$.

Time = 0.37 (sec) , antiderivative size = 3455, normalized size of antiderivative = 57.58

$$\int \sqrt{a + b \coth^2(x)} dx = \text{Too large to display}$$

[In] integrate((a+b*coth(x)^2)^(1/2),x, algorithm="fricas")

[Out] [1/4*sqrt(a + b)*log(((a*b^2 + b^3)*cosh(x)^8 + 8*(a*b^2 + b^3)*cosh(x)*sinh(x)^7 + (a*b^2 + b^3)*sinh(x)^8 + 2*(a*b^2 + 2*b^3)*cosh(x)^6 + 2*(a*b^2 + 2*b^3 + 14*(a*b^2 + b^3)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a*b^2 + b^3)*cosh(x)^3 + 3*(a*b^2 + 2*b^3)*cosh(x))*sinh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^4 + (70*(a*b^2 + b^3)*cosh(x)^4 + a^3 - a^2*b + 4*a*b^2 + 6*b^3 + 30*(a*b^2 + 2*b^3)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a*b^2 + b^3)*cosh(x)^5 + 10*(a*b^2 + 2*b^3)*cosh(x)^3 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 - 3*a*b^2 - 2*b^3)*cosh(x)^2 + 2*(14*(a*b^2 + b^3)*cosh(x)^6 + 15*(a*b^2 + 2*b^3)*cosh(x)^4 - a^3 + 3*a*b^2 + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(b^2*cosh(x)^6 + 6*b^2*cosh(x)*sinh(x)^5 + b^2*sinh(x)^6 + 3*b^2*cosh(x)^4 + 3*(5*b^2*cosh(x)^2 + b^2)*sinh(x)^4 + 4*(5*b^2*cosh(x)^3 + 3*b^2*cosh(x))*sinh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x)^2 + (15*b^2*cosh(x)^4 + 18*b^2*cosh(x)^2 - a^2 + 2*a*b + 3*b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(3*b^2*cosh(x)^5 + 6*b^2*cosh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a*b^2 + b^3)*cosh(x)^7 + 3*(a*b^2 + 2*b^3)*cosh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^3 - (a^3 - 3*a*b^2 - 2*b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6)) + 1/2*sqrt(b)*log(-((a + 2*b)*cosh(x)^4 + 4*(a + 2*b)*cosh(x)*sinh(x)^3 + (a + 2*b)*sinh(x)^4 - 2*(a - 2*b)*cosh(x)^2 + 2*(3*(a + 2*b)*cosh(x)^2 - a + 2*b)*sinh(x)^2 - 2*sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*((a + 2*b)*cosh(x)^3 - (a - 2*b)*cosh(x))*sinh(x) + a + 2*b)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*sinh(x)^2 - 2*cosh(x)^2 + 4*(cosh(x)^3 - cosh(x))*sinh(x) + 1)) + 1/4*sqrt(a + b)*log(-((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*a*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a)*sinh(x)^2 + sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)

$$\begin{aligned}
& /(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)) + 4*((a + b)*\cosh(x)^3 - a*\cosh(x))*\sinh(x) + a + b)/(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2)), \sqrt{-b}*\arctan(\sqrt{2}*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 + 1)*\sqrt{-b} \\
& *\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)})/((a + b)*\cosh(x)^4 + 4*(a + b)*\cosh(x)*\sinh(x)^3 + \\
& (a + b)*\sinh(x)^4 - 2*(a - b)*\cosh(x)^2 + 2*(3*(a + b)*\cosh(x)^2 - a + b)*\sinh(x)^2 + 4*((a + b)*\cosh(x)^3 - (a - b)*\cosh(x))*\sinh(x) + a + b)) + 1/4* \\
& \sqrt{a + b}*\log(((a*b^2 + b^3)*\cosh(x)^8 + 8*(a*b^2 + b^3)*\cosh(x)*\sinh(x)^7 + (a*b^2 + b^3)*\sinh(x)^8 + 2*(a*b^2 + 2*b^3)*\cosh(x)^6 + 2*(a*b^2 + 2*b^3 \\
& + 14*(a*b^2 + b^3)*\cosh(x)^2)*\sinh(x)^6 + 4*(14*(a*b^2 + b^3)*\cosh(x)^3 + 3*(a*b^2 + 2*b^3)*\cosh(x))*\sinh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*\cosh(x)^4 + (70*(a*b^2 + b^3)*\cosh(x)^4 + a^3 - a^2*b + 4*a*b^2 + 6*b^3 + 30*(a*b^2 + 2*b^3)*\cosh(x)^2)*\sinh(x)^4 + 4*(14*(a*b^2 + b^3)*\cosh(x)^5 + 10*(a*b^2 + 2*b^3)*\cosh(x)^3 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*\cosh(x))*\sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 - 3*a*b^2 - 2*b^3)*\cosh(x)^2 + 2*(14*(a*b^2 + b^3)*\cosh(x)^6 + 15*(a*b^2 + 2*b^3)*\cosh(x)^4 - a^3 + 3*a*b^2 + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 + 6*b^3)*\cosh(x)^2)*\sinh(x)^2 + \sqrt{2}*(b^2*\cosh(x)^6 + 6*b^2*\cosh(x)*\sinh(x)^5 + b^2*\sinh(x)^6 + 3*b^2*\cosh(x)^4 + 3*(5*b^2*\cosh(x)^2 + b^2)*\sinh(x)^4 + 4*(5*b^2*\cosh(x)^3 + 3*b^2*\cosh(x))*\sinh(x)^3 - (a^2 - 2*a*b - 3*b^2)*\cosh(x)^2 + (15*b^2*\cosh(x)^4 + 18*b^2*\cosh(x)^2 - a^2 + 2*a*b + 3*b^2)*\sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(3*b^2*\cosh(x)^5 + 6*b^2*\cosh(x)^3 - (a^2 - 2*a*b - 3*b^2)*\cosh(x))*\sinh(x))*\sqrt{a + b}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2))} + 4*(2*(a*b^2 + b^3)*\cosh(x)^7 + 3*(a*b^2 + 2*b^3)*\cosh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*\cosh(x)^3 - (a^3 - 3*a*b^2 - 2*b^3)*\cosh(x))*\sinh(x))/(\cosh(x)^6 + 6*\cosh(x)^5*\sinh(x) + 15*\cosh(x)^4*\sinh(x)^2 + 20*\cosh(x)^3*\sinh(x)^3 + 15*\cosh(x)^2*\sinh(x)^4 + 6*\cosh(x)*\sinh(x)^5 + \sinh(x)^6)) + 1/4*\sqrt{a + b}*\log(-((a + b)*\cosh(x)^4 + 4*(a + b)*\cosh(x)*\sinh(x)^3 + (a + b)*\sinh(x)^4 - 2*a*\cosh(x)^2 + 2*(3*(a + b)*\cosh(x)^2 - a)*\sinh(x)^2 + \sqrt{2}*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 - 1)*\sqrt{a + b}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2))} + 4*((a + b)*\cosh(x)^3 - a*\cosh(x))*\sinh(x) + a + b)/(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2)), -1/2*\sqrt{-a - b}*\arctan(\sqrt{2}*(b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 + a + b)*\sqrt{-a - b}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)})/((a*b + b^2)*\cosh(x)^4 + 4*(a*b + b^2)*\cosh(x)*\sinh(x)^3 + (a*b + b^2)*\sinh(x)^4 - (a^2 - a*b - 2*b^2)*\cosh(x)^2 + (6*(a*b + b^2)*\cosh(x)^2 - a^2 + a*b + 2*b^2)*\sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(2*(a*b + b^2)*\cosh(x)^3 - (a^2 - a*b - 2*b^2)*\cosh(x))*\sinh(x))) - 1/2*\sqrt{-a - b}*\arctan(\sqrt{2}*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 - 1)*\sqrt{-a - b}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)})/((a + b)*\cosh(x)^4 + 4*(a + b)*\cosh(x)*\sinh(x)^3 + (a + b)*\sinh(x)^4 - 2*(a - b)*\cosh(x)^2 + 2*(3*(a + b)*\cosh(x)^2 - a + b)*\sinh(x)^2 + 4*((a + b)*\cosh(x)^3 - (a - b)*\cosh(x))*\sinh(x) + a + b)) + 1/2*\sqrt{b}*\log(-((a + 2*b)*\cosh(x)^4 + 4*(a + 2*b)*\cosh(x)*
\end{aligned}$$

```

sinh(x)^3 + (a + 2*b)*sinh(x)^4 - 2*(a - 2*b)*cosh(x)^2 + 2*(3*(a + 2*b)*co
sh(x)^2 - a + 2*b)*sinh(x)^2 - 2*sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + s
inh(x)^2 + 1)*sqrt(b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/
(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*((a + 2*b)*cosh(x)^3 - (a
- 2*b)*cosh(x))*sinh(x) + a + 2*b)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(
x)^4 + 2*(3*cosh(x)^2 - 1)*sinh(x)^2 - 2*cosh(x)^2 + 4*(cosh(x)^3 - cosh(x)
)*sinh(x) + 1)), -1/2*sqrt(-a - b)*arctan(sqrt(2)*(b*cosh(x)^2 + 2*b*cosh(x)
)*sinh(x) + b*sinh(x)^2 + a + b)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a
+ b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2))/((a*b
+ b^2)*cosh(x)^4 + 4*(a*b + b^2)*cosh(x)*sinh(x)^3 + (a*b + b^2)*sinh(x)^4
- (a^2 - a*b - 2*b^2)*cosh(x)^2 + (6*(a*b + b^2)*cosh(x)^2 - a^2 + a*b + 2*
b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(2*(a*b + b^2)*cosh(x)^3 - (a^2 - a*
b - 2*b^2)*cosh(x))*sinh(x)) - 1/2*sqrt(-a - b)*arctan(sqrt(2)*(cosh(x)^2
+ 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 +
(a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((
a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a -
b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh
(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)) + sqrt(-b)*arctan(sqrt(2)*(cosh(
x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(-b)*sqrt(((a + b)*cosh(x)^2
+ (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((
a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a
- b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cos
h(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b))]

```

Sympy [F]

$$\int \sqrt{a + b \coth^2(x)} dx = \int \sqrt{a + b \coth^2(x)} dx$$

```
[In] integrate((a+b*coth(x)**2)**(1/2),x)
```

```
[Out] Integral(sqrt(a + b*coth(x)**2), x)
```

Maxima [F]

$$\int \sqrt{a + b \coth^2(x)} dx = \int \sqrt{b \coth^2(x) + a} dx$$

```
[In] integrate((a+b*coth(x)^2)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*coth(x)^2 + a), x)
```


Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 262 vs. 2(48) = 96.

Time = 0.48 (sec) , antiderivative size = 262, normalized size of antiderivative = 4.37

$$\int \sqrt{a + b \coth^2(x)} dx$$

$$= \frac{1}{2} \left(\frac{4 b \arctan \left(-\frac{\sqrt{a+be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b} - \sqrt{a+b}}{2\sqrt{-b}} \right)}{\sqrt{-b}} - \sqrt{a+b} \log \left(\left| \left(\sqrt{a+be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b} - \sqrt{a+b} \right) \right. \right. \right. \right. \\ \left. \left. \left. \left. - 1 \right) \right| \right) \right)$$

[In] integrate((a+b*coth(x)^2)^(1/2),x, algorithm="giac")

[Out] 1/2*(4*b*arctan(-1/2*(sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - sqrt(a + b))/sqrt(-b))/sqrt(-b) - sqrt(a + b)*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*(a + b) - sqrt(a + b)*(a - b))) + sqrt(a + b)*log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) + sqrt(a + b))) - sqrt(a + b)*log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - sqrt(a + b))))*sgn(e^(2*x) - 1)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \coth^2(x)} dx = \int \sqrt{b \coth^2(x) + a} dx$$

[In] int((a + b*coth(x)^2)^(1/2),x)

[Out] int((a + b*coth(x)^2)^(1/2), x)

3.20 $\int \sqrt{a + b \coth^2(x)} \tanh(x) dx$

Optimal result	154
Rubi [A] (verified)	154
Mathematica [A] (verified)	156
Maple [F]	156
Fricas [B] (verification not implemented)	157
Sympy [F]	159
Maxima [F]	159
Giac [B] (verification not implemented)	160
Mupad [F(-1)]	160

Optimal result

Integrand size = 15, antiderivative size = 56

$$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx = -\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a}}\right) + \sqrt{a + b} \operatorname{arctanh}\left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}}\right)$$

[Out] $-\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2)}/a^{(1/2)})*a^{(1/2)}+\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2)}/(a+b)^{(1/2)})*(a+b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3751, 457, 85, 65, 214}

$$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx = \sqrt{a + b} \operatorname{arctanh}\left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}}\right) - \sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a}}\right)$$

[In] `Int[Sqrt[a + b*Coth[x]^2]*Tanh[x], x]`

[Out] $-(\operatorname{Sqrt}[a]*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2]/\operatorname{Sqrt}[a]]) + \operatorname{Sqrt}[a + b]*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2]/\operatorname{Sqrt}[a + b]]$

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 85

```
Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))),
x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[(e + f*x)^(p - 1)/(a + b*x),
x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[(e + f*x)^(p - 1)/(c + d*x), x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[0, p, 1]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 457

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.
), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p
*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[
b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst}\left(\int \frac{\sqrt{a+bx^2}}{x(1-x^2)} dx, x, \coth(x)\right) \\ &= \frac{1}{2} \text{Subst}\left(\int \frac{\sqrt{a+bx}}{(1-x)x} dx, x, \coth^2(x)\right) \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2}a \text{Subst} \left(\int \frac{1}{x\sqrt{a+bx}} dx, x, \coth^2(x) \right) \\
&\quad + \frac{1}{2}(a+b) \text{Subst} \left(\int \frac{1}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x) \right) \\
&= \frac{a \text{Subst} \left(\int \frac{1}{-\frac{a}{b} + \frac{x^2}{b}} dx, x, \sqrt{a+b \coth^2(x)} \right)}{b} + \frac{(a+b) \text{Subst} \left(\int \frac{1}{1 + \frac{a}{b} - \frac{x^2}{b}} dx, x, \sqrt{a+b \coth^2(x)} \right)}{b} \\
&= -\sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a}} \right) + \sqrt{a+b} \operatorname{arctanh} \left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00

$$\begin{aligned}
\int \sqrt{a+b \coth^2(x)} \tanh(x) dx &= -\sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a}} \right) \\
&\quad + \sqrt{a+b} \operatorname{arctanh} \left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}} \right)
\end{aligned}$$

[In] Integrate[Sqrt[a + b*Coth[x]^2]*Tanh[x],x]

[Out] -(Sqrt[a]*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a]]) + Sqrt[a + b]*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]]

Maple [F]

$$\int \sqrt{a+b \coth(x)^2} \tanh(x) dx$$

[In] int((a+b*coth(x)^2)^(1/2)*tanh(x),x)

[Out] int((a+b*coth(x)^2)^(1/2)*tanh(x),x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 557 vs. 2(44) = 88.

Time = 0.38 (sec) , antiderivative size = 3479, normalized size of antiderivative = 62.12

$$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx = \text{Too large to display}$$

[In] integrate((a+b*coth(x)^2)^(1/2)*tanh(x),x, algorithm="fricas")

[Out] [1/4*sqrt(a + b)*log(-(a^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^4 + (70*(a^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*cosh(x)^2 + 2*(14*(a^3 + a^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*cosh(x)^4 - 2*a^3 - 3*a^2*b + b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(a^2*cosh(x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*sinh(x)^6 - 3*a^2*cosh(x)^4 + 3*(5*a^2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^2*cosh(x)^3 - 3*a^2*cosh(x))*sinh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 + (15*a^2*cosh(x)^4 - 18*a^2*cosh(x)^2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*cosh(x)^5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a^3 + a^2*b)*cosh(x)^7 - 3*(2*a^3 + a^2*b)*cosh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^3 - (2*a^3 + 3*a^2*b - b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6) + 1/2*sqrt(a)*log(-(2*a + b)*cosh(x)^4 + 4*(2*a + b)*cosh(x)*sinh(x)^3 + (2*a + b)*sinh(x)^4 - 2*(2*a - b)*cosh(x)^2 + 2*(3*(2*a + b)*cosh(x)^2 - 2*a + b)*sinh(x)^2 - 2*sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(a)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*((2*a + b)*cosh(x)^3 - (2*a - b)*cosh(x))*sinh(x) + 2*a + b)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 + 1)*sinh(x)^2 + 2*cosh(x)^2 + 4*(cosh(x)^3 + cosh(x))*sinh(x) + 1) + 1/4*sqrt(a + b)*log(((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 + 2*b*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 + b)*sinh(x)^2 + sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*((a + b)*cosh(x)^3 + b*cosh(x))*sinh(x) + a + b)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)), sqrt(-a)*arctan(sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(-a)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)

$$\begin{aligned}
&)*\sinh(x) + \sinh(x)^2)/((a + b)*\cosh(x)^4 + 4*(a + b)*\cosh(x)*\sinh(x)^3 + \\
& (a + b)*\sinh(x)^4 - 2*(a - b)*\cosh(x)^2 + 2*(3*(a + b)*\cosh(x)^2 - a + b)*\sinh(x)^2 + 4*((a + b)*\cosh(x)^3 - (a - b)*\cosh(x))*\sinh(x) + a + b)) + 1/4* \\
& \sqrt{a + b}*\log(-((a^3 + a^2*b)*\cosh(x)^8 + 8*(a^3 + a^2*b)*\cosh(x)*\sinh(x) \\
& ^7 + (a^3 + a^2*b)*\sinh(x)^8 - 2*(2*a^3 + a^2*b)*\cosh(x)^6 - 2*(2*a^3 + a^2 \\
& *b - 14*(a^3 + a^2*b)*\cosh(x)^2)*\sinh(x)^6 + 4*(14*(a^3 + a^2*b)*\cosh(x)^3 \\
& - 3*(2*a^3 + a^2*b)*\cosh(x))*\sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*\cosh(x)^4 + (70*(a^3 + a^2*b)*\cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30* \\
& (2*a^3 + a^2*b)*\cosh(x)^2)*\sinh(x)^4 + 4*(14*(a^3 + a^2*b)*\cosh(x)^5 - 10*(\\
& 2*a^3 + a^2*b)*\cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*\cosh(x))*\sinh(x) \\
& ^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*\cosh(x)^2 + \\
& 2*(14*(a^3 + a^2*b)*\cosh(x)^6 - 15*(2*a^3 + a^2*b)*\cosh(x)^4 - 2*a^3 - 3*a^2 \\
& *b + b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*\cosh(x)^2)*\sinh(x)^2 + \sqrt{2} \\
&)*(a^2*\cosh(x)^6 + 6*a^2*\cosh(x)*\sinh(x)^5 + a^2*\sinh(x)^6 - 3*a^2*\cosh(x)^4 \\
& + 3*(5*a^2*\cosh(x)^2 - a^2)*\sinh(x)^4 + 4*(5*a^2*\cosh(x)^3 - 3*a^2*\cosh(x) \\
&))*\sinh(x)^3 + (3*a^2 + 2*a*b - b^2)*\cosh(x)^2 + (15*a^2*\cosh(x)^4 - 18*a^2 \\
& *\cosh(x)^2 + 3*a^2 + 2*a*b - b^2)*\sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2* \\
& \cosh(x)^5 - 6*a^2*\cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*\cosh(x))*\sinh(x))*\sqrt{ \\
& a + b}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2* \\
& \cosh(x)*\sinh(x) + \sinh(x)^2)) + 4*(2*(a^3 + a^2*b)*\cosh(x)^7 - 3*(2*a^3 + a \\
& ^2*b)*\cosh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*\cosh(x)^3 - (2*a^3 + 3*a^2 \\
& *b - b^3)*\cosh(x))*\sinh(x))/(\cosh(x)^6 + 6*\cosh(x)^5*\sinh(x) + 15*\cosh(x)^4 \\
& *4*\sinh(x)^2 + 20*\cosh(x)^3*\sinh(x)^3 + 15*\cosh(x)^2*\sinh(x)^4 + 6*\cosh(x)*\sinh(x)^5 + \sinh(x)^6)) + 1/4*\sqrt{a + b}*\log(((a + b)*\cosh(x)^4 + 4*(a + b) \\
& *\cosh(x)*\sinh(x)^3 + (a + b)*\sinh(x)^4 + 2*b*\cosh(x)^2 + 2*(3*(a + b)*\cosh(x)^2 + b)*\sinh(x)^2 + \sqrt{2}*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 + \\
& 1)*\sqrt{a + b}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x) \\
&)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)) + 4*((a + b)*\cosh(x)^3 + b*\cosh(x))*\sinh(x) + a + b)/(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2)), -1/2*\sqrt{-a \\
& - b}*\arctan(\sqrt{2}*(a*\cosh(x)^2 + 2*a*\cosh(x)*\sinh(x) + a*\sinh(x)^2 - a - \\
& b)*\sqrt{-a - b}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x) \\
&)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)))/((a^2 + a*b)*\cosh(x)^4 + 4*(a^2 + a* \\
& b)*\cosh(x)*\sinh(x)^3 + (a^2 + a*b)*\sinh(x)^4 - (2*a^2 + a*b - b^2)*\cosh(x)^2 \\
& + (6*(a^2 + a*b)*\cosh(x)^2 - 2*a^2 - a*b + b^2)*\sinh(x)^2 + a^2 + 2*a*b + \\
& b^2 + 2*(2*(a^2 + a*b)*\cosh(x)^3 - (2*a^2 + a*b - b^2)*\cosh(x))*\sinh(x)) \\
& - 1/2*\sqrt{-a - b}*\arctan(\sqrt{2}*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 \\
& + 1)*\sqrt{-a - b}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(c \\
& osh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)))/((a + b)*\cosh(x)^4 + 4*(a + b)*c \\
& osh(x)*\sinh(x)^3 + (a + b)*\sinh(x)^4 - 2*(a - b)*\cosh(x)^2 + 2*(3*(a + b)*c \\
& osh(x)^2 - a + b)*\sinh(x)^2 + 4*((a + b)*\cosh(x)^3 - (a - b)*\cosh(x))*\sinh(x) \\
& + a + b)) + 1/2*\sqrt{a}*\log(-((2*a + b)*\cosh(x)^4 + 4*(2*a + b)*\cosh(x)* \\
& \sinh(x)^3 + (2*a + b)*\sinh(x)^4 - 2*(2*a - b)*\cosh(x)^2 + 2*(3*(2*a + b)*co \\
& sh(x)^2 - 2*a + b)*\sinh(x)^2 - 2*\sqrt{2}*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + s \\
& inhx)^2 - 1)*\sqrt{a}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/ \\
& (\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)) + 4*((2*a + b)*\cosh(x)^3 - (2*
\end{aligned}$$

```

a - b)*cosh(x))*sinh(x) + 2*a + b)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 + 1)*sinh(x)^2 + 2*cosh(x)^2 + 4*(cosh(x)^3 + cosh(x))*sinh(x) + 1)), sqrt(-a)*arctan(sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(-a)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)) - 1/2*sqrt(-a - b)*arctan(sqrt(2)*(a*cosh(x)^2 + 2*a*cosh(x))*sinh(x) + a*sinh(x)^2 - a - b)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a^2 + a*b)*cosh(x)^4 + 4*(a^2 + a*b)*cosh(x)*sinh(x)^3 + (a^2 + a*b)*sinh(x)^4 - (2*a^2 + a*b - b^2)*cosh(x)^2 + (6*(a^2 + a*b)*cosh(x)^2 - 2*a^2 - a*b + b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(2*(a^2 + a*b)*cosh(x)^3 - (2*a^2 + a*b - b^2)*cosh(x))*sinh(x))) - 1/2*sqrt(-a - b)*arctan(sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b))]

```

Sympy [F]

$$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx = \int \sqrt{a + b \coth^2(x)} \tanh(x) dx$$

```
[In] integrate((a+b*coth(x)**2)**(1/2)*tanh(x),x)
```

```
[Out] Integral(sqrt(a + b*coth(x)**2)*tanh(x), x)
```

Maxima [F]

$$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx = \int \sqrt{b \coth(x)^2 + a} \tanh(x) dx$$

```
[In] integrate((a+b*coth(x)^2)^(1/2)*tanh(x),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*coth(x)^2 + a)*tanh(x), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 259 vs. 2(44) = 88.

Time = 0.51 (sec) , antiderivative size = 259, normalized size of antiderivative = 4.62

$$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx = -\frac{1}{2} \left(\frac{4a \arctan\left(-\frac{\sqrt{a+be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b + \sqrt{a+b}}}{2\sqrt{-a}}\right)}{\sqrt{-a}} + \sqrt{a+b} \log\left(\left|\left(\sqrt{a+be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b + \sqrt{a+b}}\right) - 1\right|\right) \right)$$

[In] integrate((a+b*coth(x)^2)^(1/2)*tanh(x),x, algorithm="giac")

[Out] -1/2*(4*a*arctan(-1/2*(sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) + sqrt(a + b))/sqrt(-a))/sqrt(-a) + sqrt(a + b)*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*(a + b) - sqrt(a + b)*(a - b))) + sqrt(a + b)*log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) + sqrt(a + b))) - sqrt(a + b)*log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - sqrt(a + b))))*sgn(e^(2*x) - 1)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \coth^2(x)} \tanh(x) dx = \int \tanh(x) \sqrt{b \coth^2(x) + a} dx$$

[In] int(tanh(x)*(a + b*coth(x)^2)^(1/2),x)

[Out] int(tanh(x)*(a + b*coth(x)^2)^(1/2), x)

3.21 $\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx$

Optimal result	161
Rubi [A] (verified)	161
Mathematica [C] (verified)	163
Maple [F]	163
Fricas [B] (verification not implemented)	163
Sympy [F]	165
Maxima [F]	165
Giac [F(-2)]	165
Mupad [F(-1)]	165

Optimal result

Integrand size = 17, antiderivative size = 48

$$\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx = \sqrt{a + b} \operatorname{arctanh} \left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) - \sqrt{a + b \coth^2(x)} \tanh(x)$$

[Out] $\operatorname{arctanh}(\coth(x) * (a+b)^{(1/2)} / (a+b * \coth(x)^2)^{(1/2)}) * (a+b)^{(1/2)} - (a+b * \coth(x)^2)^{(1/2)} * \tanh(x)$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3751, 486, 12, 385, 212}

$$\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx = \sqrt{a + b} \operatorname{arctanh} \left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) - \tanh(x) \sqrt{a + b \coth^2(x)}$$

[In] $\operatorname{Int}[\operatorname{Sqrt}[a + b * \operatorname{Coth}[x]^2] * \operatorname{Tanh}[x]^2, x]$

[Out] $\operatorname{Sqrt}[a + b] * \operatorname{ArcTanh}[(\operatorname{Sqrt}[a + b] * \operatorname{Coth}[x]) / \operatorname{Sqrt}[a + b * \operatorname{Coth}[x]^2]] - \operatorname{Sqrt}[a + b * \operatorname{Coth}[x]^2] * \operatorname{Tanh}[x]$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 385

Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 486

Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[(e*x)^(m + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^q/(a*e*(m + 1))), x] - Dist[1/(a*e^n*(m + 1)), Int[(e*x)^(m + n)*(a + b*x^n)^p*(c + d*x^n)^(q - 1)*Simp[c*b*(m + 1) + n*(b*c*(p + 1) + a*d*q) + d*(b*(m + 1) + b*n*(p + q + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[0, q, 1] && LtQ[m, -1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

Rule 3751

Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{\sqrt{a+bx^2}}{x^2(1-x^2)} dx, x, \coth(x)\right) \\
 &= -\sqrt{a+b\coth^2(x)} \tanh(x) + \text{Subst}\left(\int \frac{a+b}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right) \\
 &= -\sqrt{a+b\coth^2(x)} \tanh(x) + (a+b)\text{Subst}\left(\int \frac{1}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right)
 \end{aligned}$$

$$\begin{aligned}
&= -\sqrt{a + b \coth^2(x)} \tanh(x) + (a+b) \text{Subst} \left(\int \frac{1}{1 - (a+b)x^2} dx, x, \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} \right) \\
&= \sqrt{a + b} \arctan \left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) - \sqrt{a + b \coth^2(x)} \tanh(x)
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.10 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.88

$$\begin{aligned}
&\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx \\
&= -\sqrt{a + b \coth^2(x)} \text{Hypergeometric2F1} \left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{(a+b) \coth^2(x)}{a + b \coth^2(x)} \right) \tanh(x)
\end{aligned}$$

[In] Integrate[Sqrt[a + b*Coth[x]^2]*Tanh[x]^2,x]

[Out] -(Sqrt[a + b*Coth[x]^2]*Hypergeometric2F1[-1/2, 1, 1/2, ((a + b)*Coth[x]^2)/(a + b*Coth[x]^2)]*Tanh[x])

Maple [F]

$$\int \sqrt{a + b \coth(x)^2} \tanh(x)^2 dx$$

[In] int((a+b*coth(x)^2)^(1/2)*tanh(x)^2,x)

[Out] int((a+b*coth(x)^2)^(1/2)*tanh(x)^2,x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 490 vs. 2(40) = 80.

Time = 0.32 (sec) , antiderivative size = 1539, normalized size of antiderivative = 32.06

$$\int \sqrt{a + b \coth^2(x)} \tanh^2(x) dx = \text{Too large to display}$$

[In] integrate((a+b*coth(x)^2)^(1/2)*tanh(x)^2,x, algorithm="fricas")

[Out] [1/4*((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(a + b)*log((a*b^2 + b^3)*cosh(x)^8 + 8*(a*b^2 + b^3)*cosh(x)*sinh(x)^7 + (a*b^2 + b^3)*sinh(x)^8 + 2*(a*b^2 + 2*b^3)*cosh(x)^6 + 2*(a*b^2 + 2*b^3 + 14*(a*b^2 + b^3)*

$$\begin{aligned}
& \cosh(x)^2 * \sinh(x)^6 + 4 * (14 * (a * b^2 + b^3) * \cosh(x)^3 + 3 * (a * b^2 + 2 * b^3) * \cosh(x)) * \sinh(x)^5 + (a^3 - a^2 * b + 4 * a * b^2 + 6 * b^3) * \cosh(x)^4 + (70 * (a * b^2 + b^3) * \cosh(x)^4 + a^3 - a^2 * b + 4 * a * b^2 + 6 * b^3 + 30 * (a * b^2 + 2 * b^3) * \cosh(x)^2) * \sinh(x)^4 + 4 * (14 * (a * b^2 + b^3) * \cosh(x)^5 + 10 * (a * b^2 + 2 * b^3) * \cosh(x)^3 + (a^3 - a^2 * b + 4 * a * b^2 + 6 * b^3) * \cosh(x)) * \sinh(x)^3 + a^3 + 3 * a^2 * b + 3 * a * b^2 + b^3 - 2 * (a^3 - 3 * a * b^2 - 2 * b^3) * \cosh(x)^2 + 2 * (14 * (a * b^2 + b^3) * \cosh(x)^6 + 15 * (a * b^2 + 2 * b^3) * \cosh(x)^4 - a^3 + 3 * a * b^2 + 2 * b^3 + 3 * (a^3 - a^2 * b + 4 * a * b^2 + 6 * b^3) * \cosh(x)^2) * \sinh(x)^2 + \sqrt{2} * (b^2 * \cosh(x)^6 + 6 * b^2 * \cosh(x) * \sinh(x)^5 + b^2 * \sinh(x)^6 + 3 * b^2 * \cosh(x)^4 + 3 * (5 * b^2 * \cosh(x)^2 + b^2) * \sinh(x)^4 + 4 * (5 * b^2 * \cosh(x)^3 + 3 * b^2 * \cosh(x)) * \sinh(x)^3 - (a^2 - 2 * a * b - 3 * b^2) * \cosh(x)^2 + (15 * b^2 * \cosh(x)^4 + 18 * b^2 * \cosh(x)^2 - a^2 + 2 * a * b + 3 * b^2) * \sinh(x)^2 + a^2 + 2 * a * b + b^2 + 2 * (3 * b^2 * \cosh(x)^5 + 6 * b^2 * \cosh(x)^3 - (a^2 - 2 * a * b - 3 * b^2) * \cosh(x)) * \sinh(x)) * \sqrt{a + b} * \sqrt{((a + b) * \cosh(x)^2 + (a + b) * \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 * \cosh(x) * \sinh(x) + \sinh(x)^2))} + 4 * (2 * (a * b^2 + b^3) * \cosh(x)^7 + 3 * (a * b^2 + 2 * b^3) * \cosh(x)^5 + (a^3 - a^2 * b + 4 * a * b^2 + 6 * b^3) * \cosh(x)^3 - (a^3 - 3 * a * b^2 - 2 * b^3) * \cosh(x)) * \sinh(x) / (\cosh(x)^6 + 6 * \cosh(x)^5 * \sinh(x) + 15 * \cosh(x)^4 * \sinh(x)^2 + 20 * \cosh(x)^3 * \sinh(x)^3 + 15 * \cosh(x)^2 * \sinh(x)^4 + 6 * \cosh(x) * \sinh(x)^5 + \sinh(x)^6) + (\cosh(x)^2 + 2 * \cosh(x) * \sinh(x) + \sinh(x)^2 + 1) * \sqrt{a + b} * \log(-((a + b) * \cosh(x)^4 + 4 * (a + b) * \cosh(x) * \sinh(x)^3 + (a + b) * \sinh(x)^4 - 2 * a * \cosh(x)^2 + 2 * (3 * (a + b) * \cosh(x)^2 - a) * \sinh(x)^2 + \sqrt{2} * (\cosh(x)^2 + 2 * \cosh(x) * \sinh(x) + \sinh(x)^2 - 1) * \sqrt{a + b} * \sqrt{((a + b) * \cosh(x)^2 + (a + b) * \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 * \cosh(x) * \sinh(x) + \sinh(x)^2))} + 4 * ((a + b) * \cosh(x)^3 - a * \cosh(x)) * \sinh(x) + a + b) / (\cosh(x)^2 + 2 * \cosh(x) * \sinh(x) + \sinh(x)^2)) - 4 * \sqrt{2} * \sqrt{((a + b) * \cosh(x)^2 + (a + b) * \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 * \cosh(x) * \sinh(x) + \sinh(x)^2))} / (\cosh(x)^2 + 2 * \cosh(x) * \sinh(x) + \sinh(x)^2 + 1), -1/2 * ((\cosh(x)^2 + 2 * \cosh(x) * \sinh(x) + \sinh(x)^2 + 1) * \sqrt{-a - b} * \arctan(\sqrt{2} * (b * \cosh(x)^2 + 2 * b * \cosh(x) * \sinh(x) + b * \sinh(x)^2 + a + b) * \sqrt{-a - b} * \sqrt{((a + b) * \cosh(x)^2 + (a + b) * \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 * \cosh(x) * \sinh(x) + \sinh(x)^2)}) / ((a * b + b^2) * \cosh(x)^4 + 4 * (a * b + b^2) * \cosh(x) * \sinh(x)^3 + (a * b + b^2) * \sinh(x)^4 - (a^2 - a * b - 2 * b^2) * \cosh(x)^2 + (6 * (a * b + b^2) * \cosh(x)^2 - a^2 + a * b + 2 * b^2) * \sinh(x)^2 + a^2 + 2 * a * b + b^2 + 2 * (2 * (a * b + b^2) * \cosh(x)^3 - (a^2 - a * b - 2 * b^2) * \cosh(x)) * \sinh(x))) + (\cosh(x)^2 + 2 * \cosh(x) * \sinh(x) + \sinh(x)^2 + 1) * \sqrt{-a - b} * \arctan(\sqrt{2} * (\cosh(x)^2 + 2 * \cosh(x) * \sinh(x) + \sinh(x)^2 - 1) * \sqrt{-a - b} * \sqrt{((a + b) * \cosh(x)^2 + (a + b) * \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 * \cosh(x) * \sinh(x) + \sinh(x)^2)}) / ((a + b) * \cosh(x)^4 + 4 * (a + b) * \cosh(x) * \sinh(x)^3 + (a + b) * \sinh(x)^4 - 2 * (a - b) * \cosh(x)^2 + 2 * (3 * (a + b) * \cosh(x)^2 - a + b) * \sinh(x)^2 + 4 * ((a + b) * \cosh(x)^3 - (a - b) * \cosh(x)) * \sinh(x) + a + b)) + 2 * \sqrt{2} * \sqrt{((a + b) * \cosh(x)^2 + (a + b) * \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 * \cosh(x) * \sinh(x) + \sinh(x)^2))} / (\cosh(x)^2 + 2 * \cosh(x) * \sinh(x) + \sinh(x)^2 + 1)]
\end{aligned}$$

Sympy [F]

$$\int \sqrt{a + b \coth^2(x) \tanh^2(x)} dx = \int \sqrt{a + b \coth^2(x) \tanh^2(x)} dx$$

[In] `integrate((a+b*coth(x)**2)**(1/2)*tanh(x)**2,x)`

[Out] `Integral(sqrt(a + b*coth(x)**2)*tanh(x)**2, x)`

Maxima [F]

$$\int \sqrt{a + b \coth^2(x) \tanh^2(x)} dx = \int \sqrt{b \coth(x)^2 + a} \tanh(x)^2 dx$$

[In] `integrate((a+b*coth(x)^2)^(1/2)*tanh(x)^2,x, algorithm="maxima")`

[Out] `integrate(sqrt(b*coth(x)^2 + a)*tanh(x)^2, x)`

Giac [F(-2)]

Exception generated.

$$\int \sqrt{a + b \coth^2(x) \tanh^2(x)} dx = \text{Exception raised: TypeError}$$

[In] `integrate((a+b*coth(x)^2)^(1/2)*tanh(x)^2,x, algorithm="giac")`

[Out] `Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \coth^2(x) \tanh^2(x)} dx = \int \tanh(x)^2 \sqrt{b \coth(x)^2 + a} dx$$

[In] `int(tanh(x)^2*(a + b*coth(x)^2)^(1/2),x)`

[Out] `int(tanh(x)^2*(a + b*coth(x)^2)^(1/2), x)`

3.22 $\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx$

Optimal result	166
Rubi [A] (verified)	166
Mathematica [A] (verified)	168
Maple [B] (verified)	169
Fricas [B] (verification not implemented)	169
Sympy [F]	173
Maxima [F]	173
Giac [F(-2)]	173
Mupad [B] (verification not implemented)	173

Optimal result

Integrand size = 17, antiderivative size = 82

$$\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx = (a + b)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right) - (a + b) \sqrt{a + b \coth^2(x)} - \frac{1}{3} (a + b \coth^2(x))^{3/2} - \frac{(a + b \coth^2(x))^{5/2}}{5b}$$

[Out] $(a+b)^{(3/2)} * \operatorname{arctanh}((a+b * \coth(x)^2)^{(1/2)} / (a+b)^{(1/2)}) - 1/3 * (a+b * \coth(x)^2)^{(3/2)} - 1/5 * (a+b * \coth(x)^2)^{(5/2)} / b - (a+b) * (a+b * \coth(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3751, 457, 81, 52, 65, 214}

$$\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx = (a + b)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right) - \frac{(a + b \coth^2(x))^{5/2}}{5b} - \frac{1}{3} (a + b \coth^2(x))^{3/2} - (a + b) \sqrt{a + b \coth^2(x)}$$

[In] $\operatorname{Int}[\operatorname{Coth}[x]^3 * (a + b * \operatorname{Coth}[x]^2)^{(3/2)}, x]$

[Out] $(a + b)^{(3/2)} * \operatorname{ArcTanh}[\operatorname{Sqrt}[a + b * \operatorname{Coth}[x]^2] / \operatorname{Sqrt}[a + b]] - (a + b) * \operatorname{Sqrt}[a + b * \operatorname{Coth}[x]^2] - (a + b * \operatorname{Coth}[x]^2)^{(3/2)} / 3 - (a + b * \operatorname{Coth}[x]^2)^{(5/2)} / (5 * b)$

Rule 52

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(
b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 214

```
Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 457

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_
.), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p
*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[
b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{x^3(a+bx^2)^{3/2}}{1-x^2} dx, x, \coth(x)\right) \\
&= \frac{1}{2}\text{Subst}\left(\int \frac{x(a+bx)^{3/2}}{1-x} dx, x, \coth^2(x)\right) \\
&= -\frac{(a+b\coth^2(x))^{5/2}}{5b} + \frac{1}{2}\text{Subst}\left(\int \frac{(a+bx)^{3/2}}{1-x} dx, x, \coth^2(x)\right) \\
&= -\frac{1}{3}(a+b\coth^2(x))^{3/2} - \frac{(a+b\coth^2(x))^{5/2}}{5b} + \frac{1}{2}(a+b)\text{Subst}\left(\int \frac{\sqrt{a+bx}}{1-x} dx, x, \coth^2(x)\right) \\
&= -\left((a+b)\sqrt{a+b\coth^2(x)}\right) - \frac{1}{3}(a+b\coth^2(x))^{3/2} - \frac{(a+b\coth^2(x))^{5/2}}{5b} \\
&\quad + \frac{1}{2}(a+b)^2\text{Subst}\left(\int \frac{1}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x)\right) \\
&= -\left((a+b)\sqrt{a+b\coth^2(x)}\right) - \frac{1}{3}(a+b\coth^2(x))^{3/2} - \frac{(a+b\coth^2(x))^{5/2}}{5b} \\
&\quad + \frac{(a+b)^2\text{Subst}\left(\int \frac{1}{1+\frac{a}{b}-\frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)}\right)}{b} \\
&= (a+b)^{3/2}\text{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right) - (a+b)\sqrt{a+b\coth^2(x)} \\
&\quad - \frac{1}{3}(a+b\coth^2(x))^{3/2} - \frac{(a+b\coth^2(x))^{5/2}}{5b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.05

$$\int \coth^3(x) (a+b\coth^2(x))^{3/2} dx = (a+b)^{3/2}\text{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right) - \frac{\sqrt{a+b\coth^2(x)}(3a^2+20ab+15b^2+b(6a+5b)\coth^2(x)+3b^2\coth^4(x))}{15b}$$

[In] Integrate[Coth[x]^3*(a+b*Coth[x]^2)^(3/2),x]

[Out] (a+b)^(3/2)*ArcTanh[Sqrt[a+b*Coth[x]^2]/Sqrt[a+b]] - (Sqrt[a+b*Coth[x]^2]*(3*a^2+20*a*b+15*b^2+b*(6*a+5*b)*Coth[x]^2+3*b^2*Coth[x]^4))/(15*b)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 487 vs. $2(66) = 132$.

Time = 0.09 (sec) , antiderivative size = 488, normalized size of antiderivative = 5.95

method	result
derivativedivides	$-\frac{(a+b \coth(x)^2)^{\frac{5}{2}}}{5b} - \frac{(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{\frac{3}{2}}}{6} - b \left(\frac{(2b(\coth(x)-1)+2b)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}{4b} \right)$
default	$-\frac{(a+b \coth(x)^2)^{\frac{5}{2}}}{5b} - \frac{(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{\frac{3}{2}}}{6} - b \left(\frac{(2b(\coth(x)-1)+2b)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}{4b} \right)$

[In] `int(coth(x)^3*(a+b*coth(x)^2)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/5*(a+b*\coth(x)^2)^{5/2}/b-1/6*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{3/2}-1/2*b*(1/4*(2*b*(\coth(x)-1)+2*b)/b*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{1/2}+1/8*(4*(a+b)*b-4*b^2)/b^{3/2}*\ln((b*(\coth(x)-1)+b)/b^{1/2}+(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{1/2}))-1/2*(a+b)*((b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{1/2}+b^{1/2})*\ln((b*(\coth(x)-1)+b)/b^{1/2}+(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{1/2}))-1/2*(a+b)*\ln((2*a+2*b+2*b*(\coth(x)-1)+2*(a+b)^{1/2})*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{1/2})/(\coth(x)-1))-1/6*(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{3/2}+1/2*b*(1/4*(2*b*(1+\coth(x))-2*b)/b*(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{1/2}+1/8*(4*(a+b)*b-4*b^2)/b^{3/2}*\ln((b*(1+\coth(x))-b)/b^{1/2}+(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{1/2}))-1/2*(a+b)*((b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{1/2}-b^{1/2})*\ln((b*(1+\coth(x))-b)/b^{1/2}+(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{1/2}))-1/2*(a+b)*\ln((2*a+2*b-2*b*(1+\coth(x))+2*(a+b)^{1/2})*(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{1/2})/(1+\coth(x)))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2154 vs. $2(66) = 132$.

Time = 0.47 (sec) , antiderivative size = 4940, normalized size of antiderivative = 60.24

$$\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx = \text{Too large to display}$$

[In] `integrate(coth(x)^3*(a+b*coth(x)^2)^(3/2),x, algorithm="fricas")`

[Out]
$$[1/60*(15*((a*b + b^2)*\cosh(x)^{10} + 10*(a*b + b^2)*\cosh(x)*\sinh(x)^9 + (a*b + b^2)*\sinh(x)^{10} - 5*(a*b + b^2)*\cosh(x)^8 + 5*(9*(a*b + b^2)*\cosh(x)^2 - a*b - b^2)*\sinh(x)^8 + 40*(3*(a*b + b^2)*\cosh(x)^3 - (a*b + b^2)*\cosh(x))*\sinh(x)^7 + 10*(a*b + b^2)*\cosh(x)^6 + 10*(21*(a*b + b^2)*\cosh(x)^4 - 14*(a*b + b^2)*\cosh(x)^2 + a*b + b^2)*\sinh(x)^6 + 4*(63*(a*b + b^2)*\cosh(x)^5 -$$

$$\begin{aligned}
& 70*(a*b + b^2)*\cosh(x)^3 + 15*(a*b + b^2)*\cosh(x))*\sinh(x)^5 - 10*(a*b + b^2)*\cosh(x)^4 + 10*(21*(a*b + b^2)*\cosh(x)^6 - 35*(a*b + b^2)*\cosh(x)^4 + 15*(a*b + b^2)*\cosh(x)^2 - a*b - b^2)*\sinh(x)^4 + 40*(3*(a*b + b^2)*\cosh(x)^7 - 7*(a*b + b^2)*\cosh(x)^5 + 5*(a*b + b^2)*\cosh(x)^3 - (a*b + b^2)*\cosh(x))*\sinh(x)^3 + 5*(a*b + b^2)*\cosh(x)^2 + 5*(9*(a*b + b^2)*\cosh(x)^8 - 28*(a*b + b^2)*\cosh(x)^6 + 30*(a*b + b^2)*\cosh(x)^4 - 12*(a*b + b^2)*\cosh(x)^2 + a*b + b^2)*\sinh(x)^2 - a*b - b^2 + 10*((a*b + b^2)*\cosh(x)^9 - 4*(a*b + b^2)*\cosh(x)^7 + 6*(a*b + b^2)*\cosh(x)^5 - 4*(a*b + b^2)*\cosh(x)^3 + (a*b + b^2)*\cosh(x))*\sinh(x))*\sqrt{a + b}*\log(-((a^3 + a^2*b)*\cosh(x)^8 + 8*(a^3 + a^2*b)*\cosh(x)*\sinh(x)^7 + (a^3 + a^2*b)*\sinh(x)^8 - 2*(2*a^3 + a^2*b)*\cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b)*\cosh(x)^2)*\sinh(x)^6 + 4*(14*(a^3 + a^2*b)*\cosh(x)^3 - 3*(2*a^3 + a^2*b)*\cosh(x))*\sinh(x)^5 + (6*a^3 + 4*a^2*b*b - a*b^2 + b^3)*\cosh(x)^4 + (70*(a^3 + a^2*b)*\cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*\cosh(x)^2)*\sinh(x)^4 + 4*(14*(a^3 + a^2*b)*\cosh(x)^5 - 10*(2*a^3 + a^2*b)*\cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*\cosh(x))*\sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*\cosh(x)^2 + 2*(14*(a^3 + a^2*b)*\cosh(x)^6 - 15*(2*a^3 + a^2*b)*\cosh(x)^4 - 2*a^3 - 3*a^2*b + b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*\cosh(x)^2)*\sinh(x)^2 + \sqrt{2}*(a^2*\cosh(x)^6 + 6*a^2*\cosh(x)*\sinh(x)^5 + a^2*\sinh(x)^6 - 3*a^2*\cosh(x)^4 + 3*(5*a^2*\cosh(x)^2 - a^2)*\sinh(x)^4 + 4*(5*a^2*\cosh(x)^3 - 3*a^2*\cosh(x))*\sinh(x)^3 + (3*a^2 + 2*a*b - b^2)*\cosh(x)^2 + (15*a^2*\cosh(x)^4 - 18*a^2*\cosh(x)^2 + 3*a^2 + 2*a*b - b^2)*\sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*\cosh(x)^5 - 6*a^2*\cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*\cosh(x))*\sinh(x))*\sqrt{a + b}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2))} + 4*(2*(a^3 + a^2*b)*\cosh(x)^7 - 3*(2*a^3 + a^2*b)*\cosh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*\cosh(x)^3 - (2*a^3 + 3*a^2*b - b^3)*\cosh(x))*\sinh(x))/(\cosh(x)^6 + 6*\cosh(x)^5*\sinh(x) + 15*\cosh(x)^4*\sinh(x)^2 + 20*\cosh(x)^3*\sinh(x)^3 + 15*\cosh(x)^2*\sinh(x)^4 + 6*\cosh(x)*\sinh(x)^5 + \sinh(x)^6)) + 15*((a*b + b^2)*\cosh(x)^10 + 10*(a*b + b^2)*\cosh(x)*\sinh(x)^9 + (a*b + b^2)*\sinh(x)^10 - 5*(a*b + b^2)*\cosh(x)^8 + 5*(9*(a*b + b^2)*\cosh(x)^2 - a*b - b^2)*\sinh(x)^8 + 40*(3*(a*b + b^2)*\cosh(x)^3 - (a*b + b^2)*\cosh(x))*\sinh(x)^7 + 10*(a*b + b^2)*\cosh(x)^6 + 10*(21*(a*b + b^2)*\cosh(x)^4 - 14*(a*b + b^2)*\cosh(x)^2 + a*b + b^2)*\sinh(x)^6 + 4*(63*(a*b + b^2)*\cosh(x)^5 - 70*(a*b + b^2)*\cosh(x)^3 + 15*(a*b + b^2)*\cosh(x))*\sinh(x)^5 - 10*(a*b + b^2)*\cosh(x)^4 + 10*(21*(a*b + b^2)*\cosh(x)^6 - 35*(a*b + b^2)*\cosh(x)^4 + 15*(a*b + b^2)*\cosh(x)^2 - a*b - b^2)*\sinh(x)^4 + 40*(3*(a*b + b^2)*\cosh(x)^7 - 7*(a*b + b^2)*\cosh(x)^5 + 5*(a*b + b^2)*\cosh(x)^3 - (a*b + b^2)*\cosh(x))*\sinh(x)^3 + 5*(a*b + b^2)*\cosh(x)^2 + 5*(9*(a*b + b^2)*\cosh(x)^8 - 28*(a*b + b^2)*\cosh(x)^6 + 30*(a*b + b^2)*\cosh(x)^4 - 12*(a*b + b^2)*\cosh(x)^2 + a*b + b^2)*\sinh(x)^2 - a*b - b^2 + 10*((a*b + b^2)*\cosh(x)^9 - 4*(a*b + b^2)*\cosh(x)^7 + 6*(a*b + b^2)*\cosh(x)^5 - 4*(a*b + b^2)*\cosh(x)^3 + (a*b + b^2)*\cosh(x))*\sinh(x))*\sqrt{a + b}*\log(((a + b)*\cosh(x)^4 + 4*(a + b)*\cosh(x)*\sinh(x)^3 + (a + b)*\sinh(x)^4 + 2*b*\cosh(x)^2 + 2*(3*(a + b)*\cosh(x)^2 + b)*\sinh(x)^2 + \sqrt{2}*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 + 1))*\sqrt{a + b}*\sqrt{((a + b)*\cosh(x)^2 + (a +
\end{aligned}$$

$$\begin{aligned}
& b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2)) + 4 * ((a + b) * \cosh(x)^3 + b * \cosh(x) * \sinh(x) + a + b) / (\cosh(x)^2 + 2 \cosh(x) * \sinh(x) + \sinh(x)^2)) - 4 * \sqrt{2} * ((3 * a^2 + 26 * a * b + 23 * b^2) * \cosh(x)^8 + 8 * (3 * a^2 + 26 * a * b + 23 * b^2) * \cosh(x) * \sinh(x)^7 + (3 * a^2 + 26 * a * b + 23 * b^2) * \sinh(x)^8 - 4 * (3 * a^2 + 20 * a * b + 12 * b^2) * \cosh(x)^6 + 4 * (7 * (3 * a^2 + 26 * a * b + 23 * b^2) * \cosh(x)^2 - 3 * a^2 - 20 * a * b - 12 * b^2) * \sinh(x)^6 + 8 * (7 * (3 * a^2 + 26 * a * b + 23 * b^2) * \cosh(x)^3 - 3 * (3 * a^2 + 20 * a * b + 12 * b^2) * \cosh(x)) * \sinh(x)^5 + 2 * (9 * a^2 + 54 * a * b + 49 * b^2) * \cosh(x)^4 + 2 * (35 * (3 * a^2 + 26 * a * b + 23 * b^2) * \cosh(x)^4 - 30 * (3 * a^2 + 20 * a * b + 12 * b^2) * \cosh(x)^2 + 9 * a^2 + 54 * a * b + 49 * b^2) * \sinh(x)^4 + 8 * (7 * (3 * a^2 + 26 * a * b + 23 * b^2) * \cosh(x)^5 - 10 * (3 * a^2 + 20 * a * b + 12 * b^2) * \cosh(x)^3 + (9 * a^2 + 54 * a * b + 49 * b^2) * \cosh(x)) * \sinh(x)^3 - 4 * (3 * a^2 + 20 * a * b + 12 * b^2) * \cosh(x)^2 + 4 * (7 * (3 * a^2 + 26 * a * b + 23 * b^2) * \cosh(x)^6 - 15 * (3 * a^2 + 20 * a * b + 12 * b^2) * \cosh(x)^4 + 3 * (9 * a^2 + 54 * a * b + 49 * b^2) * \cosh(x)^2 - 3 * a^2 - 20 * a * b - 12 * b^2) * \sinh(x)^2 + 3 * a^2 + 26 * a * b + 23 * b^2 + 8 * ((3 * a^2 + 26 * a * b + 23 * b^2) * \cosh(x)^7 - 3 * (3 * a^2 + 20 * a * b + 12 * b^2) * \cosh(x)^5 + (9 * a^2 + 54 * a * b + 49 * b^2) * \cosh(x)^3 - (3 * a^2 + 20 * a * b + 12 * b^2) * \cosh(x)) * \sinh(x)) * \sqrt{2} * ((a + b) * \cosh(x)^2 + (a + b) * \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) * \sinh(x) + \sinh(x)^2)) / (b * \cosh(x)^10 + 10 * b * \cosh(x) * \sinh(x)^9 + b * \sinh(x)^10 - 5 * b * \cosh(x)^8 + 5 * (9 * b * \cosh(x)^2 - b) * \sinh(x)^8 + 40 * (3 * b * \cosh(x)^3 - b * \cosh(x)) * \sinh(x)^7 + 10 * b * \cosh(x)^6 + 10 * (21 * b * \cosh(x)^4 - 14 * b * \cosh(x)^2 + b) * \sinh(x)^6 + 4 * (63 * b * \cosh(x)^5 - 70 * b * \cosh(x)^3 + 15 * b * \cosh(x)) * \sinh(x)^5 - 10 * b * \cosh(x)^4 + 10 * (21 * b * \cosh(x)^6 - 35 * b * \cosh(x)^4 + 15 * b * \cosh(x)^2 - b) * \sinh(x)^4 + 40 * (3 * b * \cosh(x)^7 - 7 * b * \cosh(x)^5 + 5 * b * \cosh(x)^3 - b * \cosh(x)) * \sinh(x)^3 + 5 * b * \cosh(x)^2 + 5 * (9 * b * \cosh(x)^8 - 28 * b * \cosh(x)^6 + 30 * b * \cosh(x)^4 - 12 * b * \cosh(x)^2 + b) * \sinh(x)^2 + 10 * (b * \cosh(x)^9 - 4 * b * \cosh(x)^7 + 6 * b * \cosh(x)^5 - 4 * b * \cosh(x)^3 + b * \cosh(x)) * \sinh(x) - b), -1/30 * (15 * ((a * b + b^2) * \cosh(x)^10 + 10 * (a * b + b^2) * \cosh(x) * \sinh(x)^9 + (a * b + b^2) * \sinh(x)^10 - 5 * (a * b + b^2) * \cosh(x)^8 + 5 * (9 * (a * b + b^2) * \cosh(x)^2 - a * b - b^2) * \sinh(x))^8 + 40 * (3 * (a * b + b^2) * \cosh(x)^3 - (a * b + b^2) * \cosh(x)) * \sinh(x)^7 + 10 * (a * b + b^2) * \cosh(x)^6 + 10 * (21 * (a * b + b^2) * \cosh(x)^4 - 14 * (a * b + b^2) * \cosh(x)^2 + a * b + b^2) * \sinh(x)^6 + 4 * (63 * (a * b + b^2) * \cosh(x)^5 - 70 * (a * b + b^2) * \cosh(x)^3 + 15 * (a * b + b^2) * \cosh(x)) * \sinh(x)^5 - 10 * (a * b + b^2) * \cosh(x)^4 + 10 * (21 * (a * b + b^2) * \cosh(x)^6 - 35 * (a * b + b^2) * \cosh(x)^4 + 15 * (a * b + b^2) * \cosh(x)^2 - a * b - b^2) * \sinh(x)^4 + 40 * (3 * (a * b + b^2) * \cosh(x)^7 - 7 * (a * b + b^2) * \cosh(x)^5 + 5 * (a * b + b^2) * \cosh(x)^3 - (a * b + b^2) * \cosh(x)) * \sinh(x)^3 + 5 * (a * b + b^2) * \cosh(x)^2 + 5 * (9 * (a * b + b^2) * \cosh(x)^8 - 28 * (a * b + b^2) * \cosh(x)^6 + 30 * (a * b + b^2) * \cosh(x)^4 - 12 * (a * b + b^2) * \cosh(x)^2 + a * b + b^2) * \sinh(x)^2 - a * b - b^2 + 10 * ((a * b + b^2) * \cosh(x)^9 - 4 * (a * b + b^2) * \cosh(x)^7 + 6 * (a * b + b^2) * \cosh(x)^5 - 4 * (a * b + b^2) * \cosh(x)^3 + (a * b + b^2) * \cosh(x)) * \sinh(x)) * \sqrt{-a - b} * \arctan(\sqrt{2} * (a * \cosh(x)^2 + 2 * a * \cosh(x) * \sinh(x) + a * \sinh(x))^2 - a - b) * \sqrt{-a - b} * \sqrt{((a + b) * \cosh(x)^2 + (a + b) * \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 * \cosh(x) * \sinh(x) + \sinh(x)^2)) / ((a^2 + a * b) * \cosh(x)^4 + 4 * (a^2 + a * b) * \cosh(x) * \sinh(x)^3 + (a^2 + a * b) * \sinh(x)^4 - (2 * a^2 + a * b - b^2) * \cosh(x)^2 + (6 * (a^2 + a * b) * \cosh(x)^2 - 2 * a^2 - a * b + b^2) * \sinh(x)^2 + a^2 + 2 * a * b + b^2 + 2 * (2 * (a^2 + a * b) * \cosh(x)^3 - (2 * a^2 + a * b - b^2) * \cosh(x)) *
\end{aligned}$$

$$\begin{aligned}
& \sinh(x)) + 15*((a*b + b^2)*\cosh(x)^{10} + 10*(a*b + b^2)*\cosh(x)*\sinh(x)^9 + \\
& (a*b + b^2)*\sinh(x)^{10} - 5*(a*b + b^2)*\cosh(x)^8 + 5*(9*(a*b + b^2)*\cosh(x) \\
&)^2 - a*b - b^2)*\sinh(x)^8 + 40*(3*(a*b + b^2)*\cosh(x)^3 - (a*b + b^2)*\cosh \\
& (x))*\sinh(x)^7 + 10*(a*b + b^2)*\cosh(x)^6 + 10*(21*(a*b + b^2)*\cosh(x)^4 - \\
& 14*(a*b + b^2)*\cosh(x)^2 + a*b + b^2)*\sinh(x)^6 + 4*(63*(a*b + b^2)*\cosh(x) \\
& ^5 - 70*(a*b + b^2)*\cosh(x)^3 + 15*(a*b + b^2)*\cosh(x))*\sinh(x)^5 - 10*(a*b \\
& + b^2)*\cosh(x)^4 + 10*(21*(a*b + b^2)*\cosh(x)^6 - 35*(a*b + b^2)*\cosh(x)^4 \\
& + 15*(a*b + b^2)*\cosh(x)^2 - a*b - b^2)*\sinh(x)^4 + 40*(3*(a*b + b^2)*\cosh \\
& (x)^7 - 7*(a*b + b^2)*\cosh(x)^5 + 5*(a*b + b^2)*\cosh(x)^3 - (a*b + b^2)*\cos \\
& h(x))*\sinh(x)^3 + 5*(a*b + b^2)*\cosh(x)^2 + 5*(9*(a*b + b^2)*\cosh(x)^8 - 28 \\
& *(a*b + b^2)*\cosh(x)^6 + 30*(a*b + b^2)*\cosh(x)^4 - 12*(a*b + b^2)*\cosh(x)^2 \\
& + a*b + b^2)*\sinh(x)^2 - a*b - b^2 + 10*((a*b + b^2)*\cosh(x)^9 - 4*(a*b + \\
& b^2)*\cosh(x)^7 + 6*(a*b + b^2)*\cosh(x)^5 - 4*(a*b + b^2)*\cosh(x)^3 + (a*b \\
& + b^2)*\cosh(x))*\sinh(x))*\sqrt{-a - b}*\arctan(\sqrt{2}*\sqrt{-a - b}*\sqrt{((a \\
& + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) \\
& + \sinh(x)^2)))/((a + b)*\cosh(x)^2 + 2*(a + b)*\cosh(x)*\sinh(x) + (a + b)*\sinh \\
& (x)^2 + a + b)) + 2*\sqrt{2}*((3*a^2 + 26*a*b + 23*b^2)*\cosh(x)^8 + 8*(3*a^2 \\
& + 26*a*b + 23*b^2)*\cosh(x)*\sinh(x)^7 + (3*a^2 + 26*a*b + 23*b^2)*\sinh(x)^8 \\
& - 4*(3*a^2 + 20*a*b + 12*b^2)*\cosh(x)^6 + 4*(7*(3*a^2 + 26*a*b + 23*b^2)*\c \\
& osh(x)^2 - 3*a^2 - 20*a*b - 12*b^2)*\sinh(x)^6 + 8*(7*(3*a^2 + 26*a*b + 23*b \\
& ^2)*\cosh(x)^3 - 3*(3*a^2 + 20*a*b + 12*b^2)*\cosh(x))*\sinh(x)^5 + 2*(9*a^2 + \\
& 54*a*b + 49*b^2)*\cosh(x)^4 + 2*(35*(3*a^2 + 26*a*b + 23*b^2)*\cosh(x)^4 - 3 \\
& 0*(3*a^2 + 20*a*b + 12*b^2)*\cosh(x)^2 + 9*a^2 + 54*a*b + 49*b^2)*\sinh(x)^4 \\
& + 8*(7*(3*a^2 + 26*a*b + 23*b^2)*\cosh(x)^5 - 10*(3*a^2 + 20*a*b + 12*b^2)*\c \\
& osh(x)^3 + (9*a^2 + 54*a*b + 49*b^2)*\cosh(x))*\sinh(x)^3 - 4*(3*a^2 + 20*a*b \\
& + 12*b^2)*\cosh(x)^2 + 4*(7*(3*a^2 + 26*a*b + 23*b^2)*\cosh(x)^6 - 15*(3*a^2 \\
& + 20*a*b + 12*b^2)*\cosh(x)^4 + 3*(9*a^2 + 54*a*b + 49*b^2)*\cosh(x)^2 - 3*a \\
& ^2 - 20*a*b - 12*b^2)*\sinh(x)^2 + 3*a^2 + 26*a*b + 23*b^2 + 8*((3*a^2 + 26* \\
& a*b + 23*b^2)*\cosh(x)^7 - 3*(3*a^2 + 20*a*b + 12*b^2)*\cosh(x)^5 + (9*a^2 + \\
& 54*a*b + 49*b^2)*\cosh(x)^3 - (3*a^2 + 20*a*b + 12*b^2)*\cosh(x))*\sinh(x))*\sq \\
& rt(((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\s \\
& inh(x) + \sinh(x)^2)))/(b*\cosh(x)^{10} + 10*b*\cosh(x)*\sinh(x)^9 + b*\sinh(x)^{10} \\
& - 5*b*\cosh(x)^8 + 5*(9*b*\cosh(x)^2 - b)*\sinh(x)^8 + 40*(3*b*\cosh(x)^3 - b* \\
& \cosh(x))*\sinh(x)^7 + 10*b*\cosh(x)^6 + 10*(21*b*\cosh(x)^4 - 14*b*\cosh(x)^2 + \\
& b)*\sinh(x)^6 + 4*(63*b*\cosh(x)^5 - 70*b*\cosh(x)^3 + 15*b*\cosh(x))*\sinh(x)^ \\
& 5 - 10*b*\cosh(x)^4 + 10*(21*b*\cosh(x)^6 - 35*b*\cosh(x)^4 + 15*b*\cosh(x)^2 - \\
& b)*\sinh(x)^4 + 40*(3*b*\cosh(x)^7 - 7*b*\cosh(x)^5 + 5*b*\cosh(x)^3 - b*\cosh \\
& (x))*\sinh(x)^3 + 5*b*\cosh(x)^2 + 5*(9*b*\cosh(x)^8 - 28*b*\cosh(x)^6 + 30*b*\c \\
& osh(x)^4 - 12*b*\cosh(x)^2 + b)*\sinh(x)^2 + 10*(b*\cosh(x)^9 - 4*b*\cosh(x)^7 + \\
& 6*b*\cosh(x)^5 - 4*b*\cosh(x)^3 + b*\cosh(x))*\sinh(x) - b]
\end{aligned}$$

Sympy [F]

$$\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx = \int (a + b \coth^2(x))^{\frac{3}{2}} \coth^3(x) dx$$

[In] integrate(coth(x)**3*(a+b*coth(x)**2)**(3/2),x)

[Out] Integral((a + b*coth(x)**2)**(3/2)*coth(x)**3, x)

Maxima [F]

$$\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx = \int (b \coth(x)^2 + a)^{\frac{3}{2}} \coth(x)^3 dx$$

[In] integrate(coth(x)^3*(a+b*coth(x)^2)^(3/2),x, algorithm="maxima")

[Out] integrate((b*coth(x)^2 + a)^(3/2)*coth(x)^3, x)

Giac [F(-2)]

Exception generated.

$$\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx = \text{Exception raised: TypeError}$$

[In] integrate(coth(x)^3*(a+b*coth(x)^2)^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);OUTPUT:Error: Bad Argument Type

Mupad [B] (verification not implemented)

Time = 9.99 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.37

$$\int \coth^3(x) (a + b \coth^2(x))^{3/2} dx = -\frac{(b \coth(x)^2 + a)^{5/2}}{5b} - \left(\frac{a+b}{3b} - \frac{a}{3b}\right) (b \coth(x)^2 + a)^{3/2} \\ - (a+b) \left(\frac{a+b}{b} - \frac{a}{b}\right) \sqrt{b \coth(x)^2 + a} - \operatorname{atan}\left(\frac{(a+b)^{3/2} \sqrt{b \coth(x)^2 + a} \operatorname{li}}{a^2 + 2ab + b^2}\right) (a+b)^{3/2} \operatorname{li}$$

[In] int(coth(x)^3*(a + b*coth(x)^2)^(3/2),x)

[Out] - (a + b*coth(x)^2)^(5/2)/(5*b) - ((a + b)/(3*b) - a/(3*b))*(a + b*coth(x)^2)^(3/2) - atan(((a + b)^(3/2)*(a + b*coth(x)^2)^(1/2)*1i)/(2*a*b + a^2 + b^2))*(a + b)^(3/2)*1i - (a + b)*((a + b)/b - a/b)*(a + b*coth(x)^2)^(1/2)

3.23 $\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx$

Optimal result	174
Rubi [A] (verified)	174
Mathematica [A] (verified)	177
Maple [B] (verified)	177
Fricas [B] (verification not implemented)	178
Sympy [F]	178
Maxima [F]	179
Giac [F(-2)]	179
Mupad [F(-1)]	179

Optimal result

Integrand size = 17, antiderivative size = 123

$$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx = -\frac{(3a^2 + 12ab + 8b^2) \operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right)}{8\sqrt{b}}$$

$$+ (a + b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right)$$

$$- \frac{1}{8}(5a + 4b) \coth(x) \sqrt{a + b \coth^2(x)} - \frac{1}{4} b \coth^3(x) \sqrt{a + b \coth^2(x)}$$

[Out] (a+b)^(3/2)*arctanh(coth(x)*(a+b)^(1/2)/(a+b*coth(x)^2)^(1/2))-1/8*(3*a^2+12*a*b+8*b^2)*arctanh(coth(x)*b^(1/2)/(a+b*coth(x)^2)^(1/2))/b^(1/2)-1/8*(5*a+4*b)*coth(x)*(a+b*coth(x)^2)^(1/2)-1/4*b*coth(x)^3*(a+b*coth(x)^2)^(1/2)

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.412$, Rules used = {3751, 488, 596, 537, 223, 212, 385}

$$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx = -\frac{(3a^2 + 12ab + 8b^2) \operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right)}{8\sqrt{b}}$$

$$+ (a + b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right)$$

$$- \frac{1}{8}(5a + 4b) \coth(x) \sqrt{a + b \coth^2(x)} - \frac{1}{4} b \coth^3(x) \sqrt{a + b \coth^2(x)}$$

[In] Int[Coth[x]^2*(a + b*Coth[x]^2)^(3/2), x]

[Out]
$$-1/8*((3*a^2 + 12*a*b + 8*b^2)*\text{ArcTanh}[(\text{Sqrt}[b]*\text{Coth}[x])/\text{Sqrt}[a + b*\text{Coth}[x]^2]])/\text{Sqrt}[b] + (a + b)^{(3/2)}*\text{ArcTanh}[(\text{Sqrt}[a + b]*\text{Coth}[x])/\text{Sqrt}[a + b*\text{Coth}[x]^2]] - ((5*a + 4*b)*\text{Coth}[x]*\text{Sqrt}[a + b*\text{Coth}[x]^2])/8 - (b*\text{Coth}[x]^3*\text{Sqrt}[a + b*\text{Coth}[x]^2])/4$$

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 223

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 385

Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 488

Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q - 1)/(b*e*(m + n*(p + q) + 1))), x] + Dist[1/(b*(m + n*(p + q) + 1)), Int[(e*x)^m*(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp[c*((c*b - a*d)*(m + 1) + c*b*n*(p + q)) + (d*(c*b - a*d)*(m + 1) + d*n*(q - 1)*(b*c - a*d) + c*b*d*n*(p + q))*x^n, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

Rule 537

Int[((e_) + (f_)*(x_)^(n_))/(((a_) + (b_)*(x_)^(n_))*Sqrt[(c_) + (d_)*(x_)^(n_)]), x_Symbol] := Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 596

Int[((g_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] := Simp[f*g^(n - 1)*(g*x)^(m - n + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(b*d*(m + n*(p + q + 1) + 1))), x] - Dist[g^n/(b*d*(m + n*(p + q + 1) + 1)), Int[(g*x)^(m - n)*(a +

```

b*x^n)^p*(c + d*x^n)^q*Simp[a*f*c*(m - n + 1) + (a*f*d*(m + n*q + 1) + b*(f
*c*(m + n*p + 1) - e*d*(m + n*(p + q + 1) + 1))*x^n, x], x] /; FreeQ[{
a, b, c, d, e, f, g, p, q}, x] && IGtQ[n, 0] && GtQ[m, n - 1]

```

Rule 3751

```

Int[((d_)*tan[(e_.) + (f_)*(x_)]^(m_))*((a_) + (b_)*((c_)*tan[(e_.) +
(f_)*(x_)]^(n_))^(p_)), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst} \left(\int \frac{x^2(a + bx^2)^{3/2}}{1 - x^2} dx, x, \coth(x) \right) \\
&= -\frac{1}{4}b \coth^3(x) \sqrt{a + b \coth^2(x)} - \frac{1}{4} \text{Subst} \left(\int \frac{x^2(-a(4a + 3b) - b(5a + 4b)x^2)}{(1 - x^2) \sqrt{a + bx^2}} dx, x, \coth(x) \right) \\
&= -\frac{1}{8}(5a + 4b) \coth(x) \sqrt{a + b \coth^2(x)} - \frac{1}{4}b \coth^3(x) \sqrt{a + b \coth^2(x)} \\
&\quad - \frac{\text{Subst} \left(\int \frac{-ab(5a+4b) - b(3a^2+12ab+8b^2)x^2}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x) \right)}{8b} \\
&= -\frac{1}{8}(5a + 4b) \coth(x) \sqrt{a + b \coth^2(x)} - \frac{1}{4}b \coth^3(x) \sqrt{a + b \coth^2(x)} \\
&\quad + (a + b)^2 \text{Subst} \left(\int \frac{1}{(1 - x^2) \sqrt{a + bx^2}} dx, x, \coth(x) \right) \\
&\quad - \frac{1}{8}(3a^2 + 12ab + 8b^2) \text{Subst} \left(\int \frac{1}{\sqrt{a + bx^2}} dx, x, \coth(x) \right) \\
&= -\frac{1}{8}(5a + 4b) \coth(x) \sqrt{a + b \coth^2(x)} - \frac{1}{4}b \coth^3(x) \sqrt{a + b \coth^2(x)} \\
&\quad + (a + b)^2 \text{Subst} \left(\int \frac{1}{1 - (a + b)x^2} dx, x, \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} \right) \\
&\quad - \frac{1}{8}(3a^2 + 12ab + 8b^2) \text{Subst} \left(\int \frac{1}{1 - bx^2} dx, x, \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} \right)
\end{aligned}$$

$$\begin{aligned}
&= -\frac{(3a^2 + 12ab + 8b^2) \operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{8\sqrt{b}} \\
&\quad + (a+b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right) - \frac{1}{8}(5a+4b) \coth(x) \sqrt{a+b \coth^2(x)} \\
&\quad\quad\quad - \frac{1}{4} b \coth^3(x) \sqrt{a+b \coth^2(x)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.95 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.78

$$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx = \frac{\sqrt{(-a+b+(a+b)\cosh(2x))\operatorname{csch}^2(x)} \left(-\sqrt{2}\sqrt{a+b}(3a^2+12ab+8b^2) \operatorname{arctanh}\left(\frac{\sqrt{b}\coth(x)}{\sqrt{a+b\coth^2(x)}}\right) \right)}{\dots}$$

[In] Integrate[Coth[x]^2*(a + b*Coth[x]^2)^(3/2), x]

[Out] (Sqrt[(-a + b + (a + b)*Cosh[2*x])*Csch[x]^2]*(-(Sqrt[2]*Sqrt[a + b]*(3*a^2 + 12*a*b + 8*b^2)*ArcTanh[(Sqrt[2]*Sqrt[b]*Cosh[x])/Sqrt[-a + b + (a + b)*Cosh[2*x]]) + Sqrt[b]*(8*Sqrt[2]*(a + b)^2*ArcTanh[(Sqrt[2]*Sqrt[a + b]*Cosh[x])/Sqrt[-a + b + (a + b)*Cosh[2*x]]) - Sqrt[a + b]*Sqrt[-a + b + (a + b)*Cosh[2*x])*Coth[x]*Csch[x]*(5*a + 6*b + 2*b*Csch[x]^2)))*Sinh[x])/(8*Sqrt[2]*Sqrt[b]*Sqrt[a + b]*Sqrt[-a + b + (a + b)*Cosh[2*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 528 vs. 2(101) = 202.

Time = 0.10 (sec) , antiderivative size = 529, normalized size of antiderivative = 4.30

method	result
derivativedivides	$-\frac{\coth(x)(a+b\coth(x)^2)^{3/2}}{4} - \frac{3a\left(\frac{\coth(x)\sqrt{a+b\coth(x)^2}}{2} + \frac{a\ln(\sqrt{b}\coth(x)+\sqrt{a+b\coth(x)^2})}{2\sqrt{b}}\right)}{4} - \frac{(b(\coth(x)-1)^2+2b)}{\dots}$
default	$-\frac{\coth(x)(a+b\coth(x)^2)^{3/2}}{4} - \frac{3a\left(\frac{\coth(x)\sqrt{a+b\coth(x)^2}}{2} + \frac{a\ln(\sqrt{b}\coth(x)+\sqrt{a+b\coth(x)^2})}{2\sqrt{b}}\right)}{4} - \frac{(b(\coth(x)-1)^2+2b)}{\dots}$

[In] int(coth(x)^2*(a+b*coth(x)^2)^(3/2), x, method=_RETURNVERBOSE)

[Out] -1/4*coth(x)*(a+b*coth(x)^2)^(3/2)-3/4*a*(1/2*coth(x)*(a+b*coth(x)^2)^(1/2)+1/2*a/b^(1/2)*ln(b^(1/2)*coth(x)+(a+b*coth(x)^2)^(1/2)))-1/6*(b*(coth(x)-1

$$\begin{aligned} &)^2+2*b*(\coth(x)-1)+a+b)^{(3/2)}-1/2*b*(1/4*(2*b*(\coth(x)-1)+2*b)/b*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{(1/2)}+1/8*(4*(a+b)*b-4*b^2)/b^{(3/2)}*\ln((b*(\coth(x)-1)+b)/b^{(1/2)}+(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{(1/2)}))-1/2*(a+b)* \\ &((b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{(1/2)}+b^{(1/2)}*\ln((b*(\coth(x)-1)+b)/b^{(1/2)}+(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{(1/2)}))-(a+b)^{(1/2)}*\ln((2*a+2*b+2*b*(\coth(x)-1)+2*(a+b)^{(1/2)}*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{(1/2)})/(\coth(x)-1))))+1/6*(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{(3/2)}-1/2*b*(1/4*(2*b*(1+\coth(x))-2*b)/b*(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{(1/2)}+1/8*(4*(a+b)*b-4*b^2)/b^{(3/2)}*\ln((b*(1+\coth(x))-b)/b^{(1/2)}+(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{(1/2)}))+1/2*(a+b)*((b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{(1/2)}-b^{(1/2)}*\ln((b*(1+\coth(x))-b)/b^{(1/2)}+(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{(1/2)}))-(a+b)^{(1/2)}*\ln((2*a+2*b-2*b*(1+\coth(x))+2*(a+b)^{(1/2)}*(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{(1/2)})/(1+\coth(x)))) \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2260 vs. $2(101) = 202$.

Time = 0.60 (sec) , antiderivative size = 10286, normalized size of antiderivative = 83.63

$$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx = \text{Too large to display}$$

[In] integrate(coth(x)^2*(a+b*coth(x)^2)^(3/2),x, algorithm="fricas")

[Out] Too large to include

Sympy [F]

$$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx = \int (a + b \coth^2(x))^{\frac{3}{2}} \coth^2(x) dx$$

[In] integrate(coth(x)**2*(a+b*coth(x)**2)**(3/2),x)

[Out] Integral((a + b*coth(x)**2)**(3/2)*coth(x)**2, x)

Maxima [F]

$$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx = \int (b \coth(x)^2 + a)^{\frac{3}{2}} \coth(x)^2 dx$$

[In] `integrate(coth(x)^2*(a+b*coth(x)^2)^(3/2),x, algorithm="maxima")`

[Out] `integrate((b*coth(x)^2 + a)^(3/2)*coth(x)^2, x)`

Giac [F(-2)]

Exception generated.

$$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx = \text{Exception raised: TypeError}$$

[In] `integrate(coth(x)^2*(a+b*coth(x)^2)^(3/2),x, algorithm="giac")`

[Out] `Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type`

Mupad [F(-1)]

Timed out.

$$\int \coth^2(x) (a + b \coth^2(x))^{3/2} dx = \int \coth(x)^2 (b \coth(x)^2 + a)^{3/2} dx$$

[In] `int(coth(x)^2*(a + b*coth(x)^2)^(3/2),x)`

[Out] `int(coth(x)^2*(a + b*coth(x)^2)^(3/2), x)`

3.24 $\int \coth(x) (a + b \coth^2(x))^{3/2} dx$

Optimal result	180
Rubi [A] (verified)	180
Mathematica [A] (verified)	182
Maple [B] (verified)	182
Fricas [B] (verification not implemented)	183
Sympy [F]	185
Maxima [F]	185
Giac [F(-2)]	185
Mupad [B] (verification not implemented)	186

Optimal result

Integrand size = 15, antiderivative size = 63

$$\int \coth(x) (a + b \coth^2(x))^{3/2} dx = (a + b)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right) - (a + b) \sqrt{a + b \coth^2(x)} - \frac{1}{3} (a + b \coth^2(x))^{3/2}$$

[Out] (a+b)^(3/2)*arctanh((a+b*coth(x)^2)^(1/2)/(a+b)^(1/2))-1/3*(a+b*coth(x)^2)^(3/2)-(a+b)*(a+b*coth(x)^2)^(1/2)

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3751, 455, 52, 65, 214}

$$\int \coth(x) (a + b \coth^2(x))^{3/2} dx = (a + b)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}} \right) - (a + b) \sqrt{a + b \coth^2(x)} - \frac{1}{3} (a + b \coth^2(x))^{3/2}$$

[In] Int[Coth[x]*(a + b*Coth[x]^2)^(3/2),x]

[Out] (a + b)^(3/2)*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]] - (a + b)*Sqrt[a + b*Coth[x]^2] - (a + b*Coth[x]^2)^(3/2)/3

Rule 52

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*(b*c - a*d)/(
b*(m + n + 1)), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 455

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.
), x_Symbol] := Dist[1/n, Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x
] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n +
1, 0]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{x(a + bx^2)^{3/2}}{1 - x^2} dx, x, \coth(x)\right) \\
&= \frac{1}{2} \text{Subst}\left(\int \frac{(a + bx)^{3/2}}{1 - x} dx, x, \coth^2(x)\right) \\
&= -\frac{1}{3}(a + b \coth^2(x))^{3/2} + \frac{1}{2}(a + b) \text{Subst}\left(\int \frac{\sqrt{a + bx}}{1 - x} dx, x, \coth^2(x)\right)
\end{aligned}$$

$$\begin{aligned}
&= -\left((a+b)\sqrt{a+b\coth^2(x)}\right) - \frac{1}{3}(a+b\coth^2(x))^{3/2} \\
&\quad + \frac{1}{2}(a+b)^2 \text{Subst}\left(\int \frac{1}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x)\right) \\
&= -\left((a+b)\sqrt{a+b\coth^2(x)}\right) - \frac{1}{3}(a+b\coth^2(x))^{3/2} \\
&\quad + \frac{(a+b)^2 \text{Subst}\left(\int \frac{1}{1+\frac{a}{b}-\frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)}\right)}{b} \\
&= (a+b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right) - (a+b)\sqrt{a+b\coth^2(x)} - \frac{1}{3}(a+b\coth^2(x))^{3/2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.94

$$\begin{aligned}
\int \coth(x) (a+b\coth^2(x))^{3/2} dx &= (a+b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right) \\
&\quad - \frac{1}{3}\sqrt{a+b\coth^2(x)}(4a+3b+b\coth^2(x))
\end{aligned}$$

[In] Integrate[Coth[x]*(a + b*Coth[x]^2)^(3/2), x]

[Out] (a + b)^(3/2)*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]] - (Sqrt[a + b*Coth[x]^2]*(4*a + 3*b + b*Coth[x]^2))/3

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 472 vs. 2(51) = 102.

Time = 0.07 (sec) , antiderivative size = 473, normalized size of antiderivative = 7.51

method	result
derivativedivides	$-\frac{(b(1+\coth(x))^2-2b(1+\coth(x))+a+b)^{\frac{3}{2}}}{6} + \frac{b\left(\frac{(2b(1+\coth(x))-2b)\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{4b} + \frac{(4(a+b)b-4b^2)}{2}\right)}{2}$
default	$-\frac{(b(1+\coth(x))^2-2b(1+\coth(x))+a+b)^{\frac{3}{2}}}{6} + \frac{b\left(\frac{(2b(1+\coth(x))-2b)\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{4b} + \frac{(4(a+b)b-4b^2)}{2}\right)}{2}$

[In] int(coth(x)*(a+b*coth(x)^2)^(3/2), x, method=_RETURNVERBOSE)

```
[Out] -1/6*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(3/2)+1/2*b*(1/4*(2*b*(1+coth(x))
)-2*b)/b*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)+1/8*(4*(a+b)*b-4*b^2)/
b^(3/2)*ln((b*(1+coth(x))-b)/b^(1/2)+(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(
1/2))-1/2*(a+b)*((b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)-b^(1/2)*ln((
b*(1+coth(x))-b)/b^(1/2)+(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))-(a+b)
^(1/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b)^(1/2)*(b*(1+coth(x))^2-2*b*(1+co
th(x))+a+b)^(1/2))/(1+coth(x))))-1/6*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(
3/2)-1/2*b*(1/4*(2*b*(coth(x)-1)+2*b)/b*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a
+b)^(1/2)+1/8*(4*(a+b)*b-4*b^2)/b^(3/2)*ln((b*(coth(x)-1)+b)/b^(1/2)+(b*(co
th(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))-1/2*(a+b)*((b*(coth(x)-1)^2+2*b*(co
th(x)-1)+a+b)^(1/2)+b^(1/2)*ln((b*(coth(x)-1)+b)/b^(1/2)+(b*(coth(x)-1)^2+2
*b*(coth(x)-1)+a+b)^(1/2))-(a+b)^(1/2)*ln((2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(
1/2)*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(coth(x)-1)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 865 vs. 2(51) = 102.

Time = 0.37 (sec) , antiderivative size = 2362, normalized size of antiderivative = 37.49

$$\int \coth(x) (a + b \coth^2(x))^{3/2} dx = \text{Too large to display}$$

```
[In] integrate(coth(x)*(a+b*coth(x)^2)^(3/2),x, algorithm="fricas")
```

```
[Out] [1/12*(3*((a + b)*cosh(x)^6 + 6*(a + b)*cosh(x)*sinh(x)^5 + (a + b)*sinh(x)
^6 - 3*(a + b)*cosh(x)^4 + 3*(5*(a + b)*cosh(x)^2 - a - b)*sinh(x)^4 + 4*(5
*(a + b)*cosh(x)^3 - 3*(a + b)*cosh(x))*sinh(x)^3 + 3*(a + b)*cosh(x)^2 + 3
*(5*(a + b)*cosh(x)^4 - 6*(a + b)*cosh(x)^2 + a + b)*sinh(x)^2 + 6*((a + b)
*cosh(x)^5 - 2*(a + b)*cosh(x)^3 + (a + b)*cosh(x))*sinh(x) - a - b)*sqrt(a
+ b)*log(-((a^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (
a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 1
4*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2
*a^3 + a^2*b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^
4 + (70*(a^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3
+ a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3
+ a^2*b)*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))*sinh(x)^3 + a
^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*cosh(x)^2 + 2*(14*
(a^3 + a^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*cosh(x)^4 - 2*a^3 - 3*a^2*b +
b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(a^2
*cosh(x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*sinh(x)^6 - 3*a^2*cosh(x)^4 + 3*
(5*a^2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^2*cosh(x)^3 - 3*a^2*cosh(x))*sin
h(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 + (15*a^2*cosh(x)^4 - 18*a^2*cosh(
x)^2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*cosh(x)
)^5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x))*sinh(x))*sqrt(a + b)
*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x
```

$$\begin{aligned}
&)*\sinh(x) + \sinh(x)^2)) + 4*(2*(a^3 + a^2*b)*\cosh(x)^7 - 3*(2*a^3 + a^2*b)* \\
& \cosh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*\cosh(x)^3 - (2*a^3 + 3*a^2*b - \\
& b^3)*\cosh(x))*\sinh(x))/(\cosh(x)^6 + 6*\cosh(x)^5*\sinh(x) + 15*\cosh(x)^4*\sinh \\
& (x)^2 + 20*\cosh(x)^3*\sinh(x)^3 + 15*\cosh(x)^2*\sinh(x)^4 + 6*\cosh(x)*\sinh(x) \\
& ^5 + \sinh(x)^6)) + 3*((a + b)*\cosh(x)^6 + 6*(a + b)*\cosh(x)*\sinh(x)^5 + (a \\
& + b)*\sinh(x)^6 - 3*(a + b)*\cosh(x)^4 + 3*(5*(a + b)*\cosh(x)^2 - a - b)*\sinh \\
& (x)^4 + 4*(5*(a + b)*\cosh(x)^3 - 3*(a + b)*\cosh(x))*\sinh(x)^3 + 3*(a + b)*\c \\
& osh(x)^2 + 3*(5*(a + b)*\cosh(x)^4 - 6*(a + b)*\cosh(x)^2 + a + b)*\sinh(x)^2 \\
& + 6*((a + b)*\cosh(x)^5 - 2*(a + b)*\cosh(x)^3 + (a + b)*\cosh(x))*\sinh(x) - a \\
& - b)*\sqrt{a + b}*\log(((a + b)*\cosh(x)^4 + 4*(a + b)*\cosh(x)*\sinh(x)^3 + (a \\
& + b)*\sinh(x)^4 + 2*b*\cosh(x)^2 + 2*(3*(a + b)*\cosh(x)^2 + b)*\sinh(x)^2 + s \\
& qrt(2)*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 + 1))*\sqrt{a + b}*\sqrt{((a \\
& + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) \\
& + \sinh(x)^2)) + 4*((a + b)*\cosh(x)^3 + b*\cosh(x))*\sinh(x) + a + b)/(\cosh(x) \\
&)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2)) - 16*\sqrt{2}*((a + b)*\cosh(x)^4 + 4*(\\
& a + b)*\cosh(x)*\sinh(x)^3 + (a + b)*\sinh(x)^4 - (2*a + b)*\cosh(x)^2 + (6*(a \\
& + b)*\cosh(x)^2 - 2*a - b)*\sinh(x)^2 + 2*(2*(a + b)*\cosh(x)^3 - (2*a + b)*\c \\
& osh(x))*\sinh(x) + a + b)*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b \\
&)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)))/(\cosh(x)^6 + 6*\cosh(x)*\sinh \\
& (x)^5 + \sinh(x)^6 + 3*(5*\cosh(x)^2 - 1)*\sinh(x)^4 - 3*\cosh(x)^4 + 4*(5*\cosh \\
& (x)^3 - 3*\cosh(x))*\sinh(x)^3 + 3*(5*\cosh(x)^4 - 6*\cosh(x)^2 + 1)*\sinh(x)^2 \\
& + 3*\cosh(x)^2 + 6*(\cosh(x)^5 - 2*\cosh(x)^3 + \cosh(x))*\sinh(x) - 1), -1/6*(3 \\
& *((a + b)*\cosh(x)^6 + 6*(a + b)*\cosh(x)*\sinh(x)^5 + (a + b)*\sinh(x)^6 - 3*(\\
& a + b)*\cosh(x)^4 + 3*(5*(a + b)*\cosh(x)^2 - a - b)*\sinh(x)^4 + 4*(5*(a + b) \\
& *\cosh(x)^3 - 3*(a + b)*\cosh(x))*\sinh(x)^3 + 3*(a + b)*\cosh(x)^2 + 3*(5*(a + \\
& b)*\cosh(x)^4 - 6*(a + b)*\cosh(x)^2 + a + b)*\sinh(x)^2 + 6*((a + b)*\cosh(x) \\
& ^5 - 2*(a + b)*\cosh(x)^3 + (a + b)*\cosh(x))*\sinh(x) - a - b)*\sqrt{-a - b}*a \\
& rctan(\sqrt{2}*(a*\cosh(x)^2 + 2*a*\cosh(x)*\sinh(x) + a*\sinh(x)^2 - a - b)*\sqrt{ \\
& t(-a - b)*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - \\
& 2*\cosh(x)*\sinh(x) + \sinh(x)^2)))/((a^2 + a*b)*\cosh(x)^4 + 4*(a^2 + a*b)*\c \\
& osh(x)*\sinh(x)^3 + (a^2 + a*b)*\sinh(x)^4 - (2*a^2 + a*b - b^2)*\cosh(x)^2 + (6 \\
& *(a^2 + a*b)*\cosh(x)^2 - 2*a^2 - a*b + b^2)*\sinh(x)^2 + a^2 + 2*a*b + b^2 + \\
& 2*(2*(a^2 + a*b)*\cosh(x)^3 - (2*a^2 + a*b - b^2)*\cosh(x))*\sinh(x))) + 3*((\\
& a + b)*\cosh(x)^6 + 6*(a + b)*\cosh(x)*\sinh(x)^5 + (a + b)*\sinh(x)^6 - 3*(a + \\
& b)*\cosh(x)^4 + 3*(5*(a + b)*\cosh(x)^2 - a - b)*\sinh(x)^4 + 4*(5*(a + b)*\c \\
& osh(x)^3 - 3*(a + b)*\cosh(x))*\sinh(x)^3 + 3*(a + b)*\cosh(x)^2 + 3*(5*(a + b) \\
& *\cosh(x)^4 - 6*(a + b)*\cosh(x)^2 + a + b)*\sinh(x)^2 + 6*((a + b)*\cosh(x)^5 \\
& - 2*(a + b)*\cosh(x)^3 + (a + b)*\cosh(x))*\sinh(x) - a - b)*\sqrt{-a - b}*\arct \\
& an(\sqrt{2}*\sqrt{-a - b}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b) \\
&)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)))/((a + b)*\cosh(x)^2 + 2*(a + \\
& b)*\cosh(x)*\sinh(x) + (a + b)*\sinh(x)^2 + a + b)) + 8*\sqrt{2}*((a + b)*\cosh \\
& (x)^4 + 4*(a + b)*\cosh(x)*\sinh(x)^3 + (a + b)*\sinh(x)^4 - (2*a + b)*\cosh(x)^ \\
& 2 + (6*(a + b)*\cosh(x)^2 - 2*a - b)*\sinh(x)^2 + 2*(2*(a + b)*\cosh(x)^3 - (2 \\
& *a + b)*\cosh(x))*\sinh(x) + a + b)*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x) \\
& ^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)))/(\cosh(x)^6 + 6*\c \\
\end{aligned}$$


```
sh(x)*sinh(x)^5 + sinh(x)^6 + 3*(5*cosh(x)^2 - 1)*sinh(x)^4 - 3*cosh(x)^4 +
  4*(5*cosh(x)^3 - 3*cosh(x))*sinh(x)^3 + 3*(5*cosh(x)^4 - 6*cosh(x)^2 + 1)*
sinh(x)^2 + 3*cosh(x)^2 + 6*(cosh(x)^5 - 2*cosh(x)^3 + cosh(x))*sinh(x) - 1
)]
```

Sympy [F]

$$\int \coth(x) (a + b \coth^2(x))^{3/2} dx = \int (a + b \coth^2(x))^{\frac{3}{2}} \coth(x) dx$$

```
[In] integrate(coth(x)*(a+b*coth(x)**2)**(3/2),x)
```

```
[Out] Integral((a + b*coth(x)**2)**(3/2)*coth(x), x)
```

Maxima [F]

$$\int \coth(x) (a + b \coth^2(x))^{3/2} dx = \int (b \coth(x)^2 + a)^{\frac{3}{2}} \coth(x) dx$$

```
[In] integrate(coth(x)*(a+b*coth(x)^2)^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((b*coth(x)^2 + a)^(3/2)*coth(x), x)
```

Giac [F(-2)]

Exception generated.

$$\int \coth(x) (a + b \coth^2(x))^{3/2} dx = \text{Exception raised: TypeError}$$

```
[In] integrate(coth(x)*(a+b*coth(x)^2)^(3/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type
```

Mupad [B] (verification not implemented)

Time = 4.15 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.02

$$\int \coth(x) (a + b \coth^2(x))^{3/2} dx = \operatorname{atanh}\left(\frac{(a+b)^{3/2} \sqrt{b \coth(x)^2 + a}}{a^2 + 2ab + b^2}\right) (a+b)^{3/2} - (a+b) \sqrt{b \coth(x)^2 + a} - \frac{(b \coth(x)^2 + a)^{3/2}}{3}$$

`[In] int(coth(x)*(a + b*coth(x)^2)^(3/2),x)`
`[Out] atanh(((a + b)^(3/2)*(a + b*coth(x)^2)^(1/2))/(2*a*b + a^2 + b^2))*(a + b)^(3/2) - (a + b)*(a + b*coth(x)^2)^(1/2) - (a + b*coth(x)^2)^(3/2)/3`

3.25 $\int (a + b \coth^2(x))^{3/2} dx$

Optimal result	187
Rubi [A] (verified)	187
Mathematica [A] (verified)	189
Maple [B] (verified)	190
Fricas [B] (verification not implemented)	190
Sympy [F]	191
Maxima [F]	191
Giac [F(-2)]	191
Mupad [F(-1)]	191

Optimal result

Integrand size = 12, antiderivative size = 88

$$\int (a + b \coth^2(x))^{3/2} dx = -\frac{1}{2}\sqrt{b}(3a + 2b)\operatorname{arctanh}\left(\frac{\sqrt{b}\coth(x)}{\sqrt{a + b\coth^2(x)}}\right) + (a + b)^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{a + b}\coth(x)}{\sqrt{a + b\coth^2(x)}}\right) - \frac{1}{2}b\coth(x)\sqrt{a + b\coth^2(x)}$$

[Out] (a+b)^(3/2)*arctanh(coth(x)*(a+b)^(1/2)/(a+b*coth(x)^2)^(1/2))-1/2*(3*a+2*b)*arctanh(coth(x)*b^(1/2)/(a+b*coth(x)^2)^(1/2))*b^(1/2)-1/2*b*coth(x)*(a+b*coth(x)^2)^(1/2)

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3742, 427, 537, 223, 212, 385}

$$\int (a + b \coth^2(x))^{3/2} dx = (a + b)^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{a + b}\coth(x)}{\sqrt{a + b\coth^2(x)}}\right) - \frac{1}{2}\sqrt{b}(3a + 2b)\operatorname{arctanh}\left(\frac{\sqrt{b}\coth(x)}{\sqrt{a + b\coth^2(x)}}\right) - \frac{1}{2}b\coth(x)\sqrt{a + b\coth^2(x)}$$

[In] Int[(a + b*Coth[x]^2)^(3/2),x]

[Out] $-1/2*(\text{Sqrt}[b]*(3*a + 2*b)*\text{ArcTanh}[(\text{Sqrt}[b]*\text{Coth}[x])/\text{Sqrt}[a + b*\text{Coth}[x]^2]]) + (a + b)^{(3/2)}*\text{ArcTanh}[(\text{Sqrt}[a + b]*\text{Coth}[x])/\text{Sqrt}[a + b*\text{Coth}[x]^2]] - (b*\text{Coth}[x]*\text{Sqrt}[a + b*\text{Coth}[x]^2])/2$

Rule 212

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 223

$\text{Int}[1/\text{Sqrt}[(a_ + (b_)*(x_)^2), x_Symbol] \rightarrow \text{Subst}[\text{Int}[1/(1 - b*x^2), x], x, x/\text{Sqrt}[a + b*x^2]] /; \text{FreeQ}\{a, b\}, x \ \&\& \ !\text{GtQ}[a, 0]$

Rule 385

$\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}/((c_ + (d_)*(x_)^{(n_)}), x_Symbol] \rightarrow \text{Subst}[\text{Int}[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^{(1/n)}] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[n*p + 1, 0] \ \&\& \ \text{IntegerQ}[n]$

Rule 427

$\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)*((c_ + (d_)*(x_)^{(n_)})^{(q_)}), x_Symbol] \rightarrow \text{Simp}[d*x*(a + b*x^n)^{(p+1)}*((c + d*x^n)^{(q-1)}/(b*(n*(p+q) + 1))), x] + \text{Dist}[1/(b*(n*(p+q) + 1)), \text{Int}[(a + b*x^n)^p*(c + d*x^n)^{(q-2)}*\text{Simp}[c*(b*c*(n*(p+q) + 1) - a*d) + d*(b*c*(n*(p+2*q-1) + 1) - a*d*(n*(q-1) + 1))*x^n, x], x] /; \text{FreeQ}\{a, b, c, d, n, p\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{GtQ}[q, 1] \ \&\& \ \text{NeQ}[n*(p+q) + 1, 0] \ \&\& \ !\text{IGtQ}[p, 1] \ \&\& \ \text{IntBinomialQ}[a, b, c, d, n, p, q, x]$

Rule 537

$\text{Int}[(e_ + (f_)*(x_)^{(n_)})/((a_ + (b_)*(x_)^{(n_)})*\text{Sqrt}[(c_ + (d_)*(x_)^{(n_)})]), x_Symbol] \rightarrow \text{Dist}[f/b, \text{Int}[1/\text{Sqrt}[c + d*x^n], x], x] + \text{Dist}[(b*e - a*f)/b, \text{Int}[1/((a + b*x^n)*\text{Sqrt}[c + d*x^n]), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x]$

Rule 3742

$\text{Int}[(a_ + (b_)*((c_)*\text{tan}[e_ + (f_)*(x_)])^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{With}\{\{ff = \text{FreeFactors}[\text{Tan}[e + f*x], x]\}, \text{Dist}[c*(ff/f), \text{Subst}[\text{Int}[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(\text{Tan}[e + f*x]/ff)], x] /; \text{FreeQ}\{a, b, c, e, f, n, p\}, x \ \&\& \ (\text{IntegersQ}[n, p] \ || \ \text{IGtQ}[p, 0] \ || \ \text{EqQ}[n^2, 4] \ || \ \text{EqQ}[n^2, 16])$

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{(a + bx^2)^{3/2}}{1 - x^2} dx, x, \coth(x)\right) \\
&= -\frac{1}{2}b \coth(x) \sqrt{a + b \coth^2(x)} - \frac{1}{2} \text{Subst}\left(\int \frac{-a(2a + b) - b(3a + 2b)x^2}{(1 - x^2) \sqrt{a + bx^2}} dx, x, \coth(x)\right) \\
&= -\frac{1}{2}b \coth(x) \sqrt{a + b \coth^2(x)} + (a + b)^2 \text{Subst}\left(\int \frac{1}{(1 - x^2) \sqrt{a + bx^2}} dx, x, \coth(x)\right) \\
&\quad - \frac{1}{2}(b(3a + 2b)) \text{Subst}\left(\int \frac{1}{\sqrt{a + bx^2}} dx, x, \coth(x)\right) \\
&= -\frac{1}{2}b \coth(x) \sqrt{a + b \coth^2(x)} \\
&\quad + (a + b)^2 \text{Subst}\left(\int \frac{1}{1 - (a + b)x^2} dx, x, \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}}\right) \\
&\quad - \frac{1}{2}(b(3a + 2b)) \text{Subst}\left(\int \frac{1}{1 - bx^2} dx, x, \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}}\right) \\
&= -\frac{1}{2}\sqrt{b}(3a + 2b) \operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right) \\
&\quad + (a + b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right) - \frac{1}{2}b \coth(x) \sqrt{a + b \coth^2(x)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.58 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.26

$$\int (a + b \coth^2(x))^{3/2} dx = \frac{1}{2} \left(-2(-a - b)^{3/2} \arctan\left(\frac{\coth(x) \sqrt{a + b \coth^2(x)} - \sqrt{b} \operatorname{csch}^2(x)}{\sqrt{-a - b}}\right) - b \coth(x) \sqrt{a + b \coth^2(x)} \right)$$

[In] Integrate[(a + b*Coth[x]^2)^(3/2), x]

[Out] (-2*(-a - b)^(3/2)*ArcTan[(Coth[x]*Sqrt[a + b*Coth[x]^2] - Sqrt[b]*Csch[x]^2)/Sqrt[-a - b]] - b*Coth[x]*Sqrt[a + b*Coth[x]^2] + Sqrt[b]*(3*a + 2*b)*Log[-(Sqrt[b]*Coth[x]) + Sqrt[a + b*Coth[x]^2]])/2

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 472 vs. 2(70) = 140.

Time = 0.12 (sec) , antiderivative size = 473, normalized size of antiderivative = 5.38

method	result
derivativedivides	$-\frac{(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{\frac{3}{2}}}{6} - \frac{b \left(\frac{(2b(\coth(x)-1)+2b)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}{4b} + \frac{(4(a+b)b-4b^2)}{2} \right)}{2}$
default	$-\frac{(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{\frac{3}{2}}}{6} - \frac{b \left(\frac{(2b(\coth(x)-1)+2b)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}{4b} + \frac{(4(a+b)b-4b^2)}{2} \right)}{2}$

[In] int((a+b*coth(x)^2)^(3/2),x,method=_RETURNVERBOSE)

[Out]
$$-1/6*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{(3/2)}-1/2*b*(1/4*(2*b*(\coth(x)-1)+2*b)/b*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{(1/2)}+1/8*(4*(a+b)*b-4*b^2)/b^{(3/2)}*\ln((b*(\coth(x)-1)+b)/b^{(1/2)}+(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{(1/2)}))-1/2*(a+b)*((b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{(1/2)}+b^{(1/2)}*\ln((b*(\coth(x)-1)+b)/b^{(1/2)}+(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{(1/2)})-(a+b)^{(1/2)}*\ln((2*a+2*b+2*b*(\coth(x)-1)+2*(a+b)^{(1/2)}*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{(1/2)})/(\coth(x)-1))))+1/6*(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{(3/2)}-1/2*b*(1/4*(2*b*(1+\coth(x))-2*b)/b*(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{(1/2)}+1/8*(4*(a+b)*b-4*b^2)/b^{(3/2)}*\ln((b*(1+\coth(x))-b)/b^{(1/2)}+(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{(1/2)}))+1/2*(a+b)*((b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{(1/2)}-b^{(1/2)}*\ln((b*(1+\coth(x))-b)/b^{(1/2)}+(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{(1/2)})-(a+b)^{(1/2)}*\ln((2*a+2*b-2*b*(1+\coth(x))+2*(a+b)^{(1/2)}*(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{(1/2)})/(1+\coth(x))))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 947 vs. 2(70) = 140.

Time = 0.43 (sec) , antiderivative size = 5037, normalized size of antiderivative = 57.24

$$\int (a + b \coth^2(x))^{3/2} dx = \text{Too large to display}$$

[In] integrate((a+b*coth(x)^2)^(3/2),x, algorithm="fricas")

[Out] Too large to include

Sympy [F]

$$\int (a + b \coth^2(x))^{3/2} dx = \int (a + b \coth^2(x))^{\frac{3}{2}} dx$$

```
[In] integrate((a+b*coth(x)**2)**(3/2),x)
```

```
[Out] Integral((a + b*coth(x)**2)**(3/2), x)
```

Maxima [F]

$$\int (a + b \coth^2(x))^{3/2} dx = \int (b \coth(x)^2 + a)^{\frac{3}{2}} dx$$

```
[In] integrate((a+b*coth(x)^2)^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((b*coth(x)^2 + a)^(3/2), x)
```

Giac [F(-2)]

Exception generated.

$$\int (a + b \coth^2(x))^{3/2} dx = \text{Exception raised: TypeError}$$

```
[In] integrate((a+b*coth(x)^2)^(3/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type
```

Mupad [F(-1)]

Timed out.

$$\int (a + b \coth^2(x))^{3/2} dx = \int (b \coth(x)^2 + a)^{3/2} dx$$

```
[In] int((a + b*coth(x)^2)^(3/2),x)
```

```
[Out] int((a + b*coth(x)^2)^(3/2), x)
```

3.26 $\int (a + b \coth^2(x))^{3/2} \tanh(x) dx$

Optimal result	192
Rubi [A] (verified)	192
Mathematica [A] (verified)	194
Maple [F]	195
Fricas [B] (verification not implemented)	195
Sympy [F]	198
Maxima [F]	198
Giac [B] (verification not implemented)	198
Mupad [F(-1)]	199

Optimal result

Integrand size = 15, antiderivative size = 71

$$\int (a + b \coth^2(x))^{3/2} \tanh(x) dx = -a^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a}}\right) + (a + b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}}\right) - b\sqrt{a + b \coth^2(x)}$$

[Out] $-a^{(3/2)}*\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2)}/a^{(1/2)})+(a+b)^{(3/2)}*\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2)}/(a+b)^{(1/2)})-b*(a+b*\coth(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3751, 457, 86, 162, 65, 214}

$$\int (a + b \coth^2(x))^{3/2} \tanh(x) dx = a^{3/2} \left(-\operatorname{arctanh}\left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a}}\right) \right) + (a + b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \coth^2(x)}}{\sqrt{a + b}}\right) - b\sqrt{a + b \coth^2(x)}$$

[In] $\operatorname{Int}[(a + b*\operatorname{Coth}[x]^2)^{(3/2)}*\operatorname{Tanh}[x], x]$

[Out] $-(a^{(3/2)}*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2]/\operatorname{Sqrt}[a]]) + (a + b)^{(3/2)}*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2]/\operatorname{Sqrt}[a + b]] - b*\operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2]$

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 86

```
Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))),
x_Symbol] := Simp[f*(e + f*x)^(p - 1)/(b*d*(p - 1)), x] + Dist[1/(b*d), I
nt[(b*d*e^2 - a*c*f^2 + f*(2*b*d*e - b*c*f - a*d*f)*x)*((e + f*x)^(p - 2)/(
(a + b*x)*(c + d*x))), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 1]
```

Rule 162

```
Int[(((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)))/(((a_.) + (b_.)*(x_))*
((c_.) + (d_.)*(x_))), x_Symbol] := Dist[(b*g - a*h)/(b*c - a*d), Int[(e +
f*x)^p/(a + b*x), x], x] - Dist[(d*g - c*h)/(b*c - a*d), Int[(e + f*x)^p/(c
+ d*x), x], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x]
```

Rule 214

```
Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 457

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_
), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p
*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[
b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\text{integral} = \text{Subst} \left(\int \frac{(a + bx^2)^{3/2}}{x(1 - x^2)} dx, x, \coth(x) \right)$$

$$\begin{aligned}
&= \frac{1}{2} \text{Subst} \left(\int \frac{(a+bx)^{3/2}}{(1-x)x} dx, x, \coth^2(x) \right) \\
&= -b\sqrt{a+b\coth^2(x)} - \frac{1}{2} \text{Subst} \left(\int \frac{-a^2 + (-2a-b)bx}{(1-x)x\sqrt{a+bx}} dx, x, \coth^2(x) \right) \\
&= -b\sqrt{a+b\coth^2(x)} + \frac{1}{2} a^2 \text{Subst} \left(\int \frac{1}{x\sqrt{a+bx}} dx, x, \coth^2(x) \right) \\
&\quad + \frac{1}{2} (a+b)^2 \text{Subst} \left(\int \frac{1}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x) \right) \\
&= -b\sqrt{a+b\coth^2(x)} + \frac{a^2 \text{Subst} \left(\int \frac{1}{-\frac{a}{b} + \frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)} \right)}{b} \\
&\quad + \frac{(a+b)^2 \text{Subst} \left(\int \frac{1}{1 + \frac{a}{b} - \frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)} \right)}{b} \\
&= -a^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a}} \right) \\
&\quad + (a+b)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}} \right) - b\sqrt{a+b\coth^2(x)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00

$$\begin{aligned}
\int (a+b\coth^2(x))^{3/2} \tanh(x) dx &= -a^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a}} \right) \\
&+ (a+b)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}} \right) - b\sqrt{a+b\coth^2(x)}
\end{aligned}$$

[In] Integrate[(a + b*Coth[x]^2)^(3/2)*Tanh[x], x]

[Out] -(a^(3/2)*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a]]) + (a + b)^(3/2)*ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]] - b*Sqrt[a + b*Coth[x]^2]

Maple [F]

$$\int (a + b \coth(x)^2)^{\frac{3}{2}} \tanh(x) dx$$

[In] int((a+b*coth(x)^2)^(3/2)*tanh(x),x)

[Out] int((a+b*coth(x)^2)^(3/2)*tanh(x),x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 641 vs. 2(57) = 114.

Time = 0.43 (sec) , antiderivative size = 3949, normalized size of antiderivative = 55.62

$$\int (a + b \coth^2(x))^{3/2} \tanh(x) dx = \text{Too large to display}$$

[In] integrate((a+b*coth(x)^2)^(3/2)*tanh(x),x, algorithm="fricas")

[Out] [1/4*(((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 - a - b)*sqrt(a + b)*log(-((a^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^4 + (70*(a^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*cosh(x)^2 + 2*(14*(a^3 + a^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*cosh(x)^4 - 2*a^3 - 3*a^2*b + b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(a^2*cosh(x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*sinh(x)^6 - 3*a^2*cosh(x)^4 + 3*(5*a^2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^2*cosh(x)^3 - 3*a^2*cosh(x))*sinh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 + (15*a^2*cosh(x)^4 - 18*a^2*cosh(x)^2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*cosh(x)^5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a^3 + a^2*b)*cosh(x)^7 - 3*(2*a^3 + a^2*b)*cosh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^3 - (2*a^3 + 3*a^2*b - b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6) + 2*(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 - a)*sqrt(a)*log(-((2*a + b)*cosh(x)^4 + 4*(2*a + b)*cosh(x)*sinh(x))^3 + (2*a + b)*sinh(x)^4 - 2*(2*a - b)*cosh(x)^2 + 2*(3*(2*a + b)*cosh(x)^2 - 2*a + b)*sinh(x)^2 - 2*sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(a)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*((2*a + b)*cosh(x)^3 - (2*a - b)

$$\begin{aligned}
& * \cosh(x) \sinh(x) + 2a + b) / (\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + \\
& 2(3 \cosh(x)^2 + 1) \sinh(x)^2 + 2 \cosh(x)^2 + 4(\cosh(x)^3 + \cosh(x)) \sinh(x) \\
& (x) + 1)) + ((a + b) \cosh(x)^2 + 2(a + b) \cosh(x) \sinh(x) + (a + b) \sinh(x) \\
&)^2 - a - b) \sqrt{a + b} \log(((a + b) \cosh(x)^4 + 4(a + b) \cosh(x) \sinh(x) \\
&)^3 + (a + b) \sinh(x)^4 + 2b \cosh(x)^2 + 2(3(a + b) \cosh(x)^2 + b) \sinh(x) \\
&)^2 + \sqrt{2}(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1) \sqrt{a + b} \sqrt{ \\
& } \sqrt{((a + b) \cosh(x)^2 + (a + b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) * \\
& \sinh(x) + \sinh(x)^2)} + 4((a + b) \cosh(x)^3 + b \cosh(x)) \sinh(x) + a + b) / \\
& (\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2)) - 4 \sqrt{2} b \sqrt{((a + b) \cosh(x) \\
&)^2 + (a + b) \sinh(x)^2 - a + b} / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x) \\
&)^2)) / (\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1), 1/4(4(a \cosh(x)^2 \\
& + 2a \cosh(x) \sinh(x) + a \sinh(x)^2 - a) \sqrt{-a} \arctan(\sqrt{2}(\cosh(x) \\
&)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1) \sqrt{-a}) \sqrt{((a + b) \cosh(x)^2 + \\
& (a + b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2)} / ((a \\
& + b) \cosh(x)^4 + 4(a + b) \cosh(x) \sinh(x)^3 + (a + b) \sinh(x)^4 - 2(a - \\
& b) \cosh(x)^2 + 2(3(a + b) \cosh(x)^2 - a + b) \sinh(x)^2 + 4((a + b) \cosh(x) \\
&)^3 - (a - b) \cosh(x)) \sinh(x) + a + b)) + ((a + b) \cosh(x)^2 + 2(a + b) \\
& \cosh(x) \sinh(x) + (a + b) \sinh(x)^2 - a - b) \sqrt{a + b} \log(-((a^3 + a^2 b) \\
&) \cosh(x)^8 + 8(a^3 + a^2 b) \cosh(x) \sinh(x)^7 + (a^3 + a^2 b) \sinh(x)^8 - \\
& 2(2a^3 + a^2 b) \cosh(x)^6 - 2(2a^3 + a^2 b - 14(a^3 + a^2 b) \cosh(x)^2) \sinh(x)^6 + 4(14(a^3 + a^2 b) \cosh(x)^3 - 3(2a^3 + a^2 b) \cosh(x)) \sinh(x)^5 + (6a^3 + 4a^2 b - a b^2 + b^3) \cosh(x)^4 + (70(a^3 + a^2 b) \cosh(x)^4 + 6a^3 + 4a^2 b - a b^2 + b^3 - 30(2a^3 + a^2 b) \cosh(x)^2) \sinh(x)^4 + 4(14(a^3 + a^2 b) \cosh(x)^5 - 10(2a^3 + a^2 b) \cosh(x)^3 + (6a^3 + 4a^2 b - a b^2 + b^3) \cosh(x)) \sinh(x)^3 + a^3 + 3a^2 b + 3a b^2 + b^3 - 2(2a^3 + 3a^2 b - b^3) \cosh(x)^2 + 2(14(a^3 + a^2 b) \cosh(x)^6 - 15(2a^3 + a^2 b) \cosh(x)^4 - 2a^3 - 3a^2 b + b^3 + 3(6a^3 + 4a^2 b - a b^2 + b^3) \cosh(x)^2) \sinh(x)^2 + \sqrt{2}(a^2 \cosh(x)^6 + 6a^2 \cosh(x) \sinh(x)^5 + a^2 \sinh(x)^6 - 3a^2 \cosh(x)^4 + 3(5a^2 \cosh(x)^2 - a^2) \sinh(x)^4 + 4(5a^2 \cosh(x)^3 - 3a^2 \cosh(x)) \sinh(x)^3 + (3a^2 + 2a b - b^2) \cosh(x)^2 + (15a^2 \cosh(x)^4 - 18a^2 \cosh(x)^2 + 3a^2 + 2a b - b^2) \sinh(x)^2 - a^2 - 2a b - b^2 + 2(3a^2 \cosh(x)^5 - 6a^2 \cosh(x)^3 + (3a^2 + 2a b - b^2) \cosh(x)) \sinh(x)) \sqrt{a + b} \sqrt{((a + b) \cosh(x)^2 + (a + b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2)} + 4(2(a^3 + a^2 b) \cosh(x)^7 - 3(2a^3 + a^2 b) \cosh(x)^5 + (6a^3 + 4a^2 b - a b^2 + b^3) \cosh(x)^3 - (2a^3 + 3a^2 b - b^3) \cosh(x)) \sinh(x)) / (\cosh(x)^6 + 6 \cosh(x)^5 \sinh(x) + 15 \cosh(x)^4 \sinh(x)^2 + 20 \cosh(x)^3 \sinh(x)^3 + 15 \cosh(x)^2 \sinh(x)^4 + 6 \cosh(x) \sinh(x)^5 + \sinh(x)^6)) + ((a + b) \cosh(x)^2 + 2(a + b) \cosh(x) \sinh(x) + (a + b) \sinh(x)^2 - a - b) \sqrt{a + b} \log(((a + b) \cosh(x)^4 + 4(a + b) \cosh(x) \sinh(x)^3 + (a + b) \sinh(x)^4 + 2b \cosh(x)^2 + 2(3(a + b) \cosh(x)^2 + b) \sinh(x)^2 + \sqrt{2}(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1) \sqrt{a + b} \sqrt{((a + b) \cosh(x)^2 + (a + b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2)} + 4((a + b) \cosh(x)^3 + b \cosh(x)) \sinh(x) + a + b) / (\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2)) - 4 \sqrt{2} b \sqrt{((a + b) \cosh(x)^2 + (a + b)
\end{aligned}$$

$$\begin{aligned}
& *sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2))/((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1), -1/2*(((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 - a - b)*sqrt(-a - b)*arctan(sqrt(2)*(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 - a - b)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a^2 + a*b)*cosh(x)^4 + 4*(a^2 + a*b)*cosh(x)*sinh(x)^3 + (a^2 + a*b)*sinh(x)^4 - (2*a^2 + a*b - b^2)*cosh(x)^2 + (6*(a^2 + a*b)*cosh(x)^2 - 2*a^2 - a*b + b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(2*(a^2 + a*b)*cosh(x)^3 - (2*a^2 + a*b - b^2)*cosh(x))*sinh(x))) + ((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 - a - b)*sqrt(-a - b)*arctan(sqrt(2)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a + b)) - (a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 - a)*sqrt(a)*log(-((2*a + b)*cosh(x)^4 + 4*(2*a + b)*cosh(x)*sinh(x)^3 + (2*a + b)*sinh(x)^4 - 2*(2*a - b)*cosh(x)^2 + 2*(3*(2*a + b)*cosh(x)^2 - 2*a + b)*sinh(x)^2 - 2*sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(a)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*((2*a + b)*cosh(x)^3 - (2*a - b)*cosh(x))*sinh(x) + 2*a + b)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 + 1)*sinh(x)^2 + 2*cosh(x)^2 + 4*(cosh(x)^3 + cosh(x))*sinh(x) + 1)) + 2*sqrt(2)*b*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1), 1/2*(2*(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 - a)*sqrt(-a)*arctan(sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(-a)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)) - ((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 - a - b)*sqrt(-a - b)*arctan(sqrt(2)*(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 - a - b)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a^2 + a*b)*cosh(x)^4 + 4*(a^2 + a*b)*cosh(x)*sinh(x)^3 + (a^2 + a*b)*sinh(x)^4 - (2*a^2 + a*b - b^2)*cosh(x)^2 + (6*(a^2 + a*b)*cosh(x)^2 - 2*a^2 - a*b + b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(2*(a^2 + a*b)*cosh(x)^3 - (2*a^2 + a*b - b^2)*cosh(x))*sinh(x))) - ((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 - a - b)*sqrt(-a - b)*arctan(sqrt(2)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a + b)) - 2*sqrt(2)*b*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1))]
\end{aligned}$$

Sympy [F]

$$\int (a + b \coth^2(x))^{3/2} \tanh(x) dx = \int (a + b \coth^2(x))^{\frac{3}{2}} \tanh(x) dx$$

[In] integrate((a+b*coth(x)**2)**(3/2)*tanh(x),x)

[Out] Integral((a + b*coth(x)**2)**(3/2)*tanh(x), x)

Maxima [F]

$$\int (a + b \coth^2(x))^{3/2} \tanh(x) dx = \int (b \coth(x)^2 + a)^{\frac{3}{2}} \tanh(x) dx$$

[In] integrate((a+b*coth(x)^2)^(3/2)*tanh(x),x, algorithm="maxima")

[Out] integrate((b*coth(x)^2 + a)^(3/2)*tanh(x), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 470 vs. 2(57) = 114.

Time = 3.81 (sec) , antiderivative size = 470, normalized size of antiderivative = 6.62

$$\begin{aligned} & \int (a + b \coth^2(x))^{3/2} \tanh(x) dx = \\ & \frac{2a^2 \arctan\left(-\frac{\sqrt{a+b}e^{2x} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b + \sqrt{a+b}}}{2\sqrt{-a}}\right) \operatorname{sgn}(e^{2x} - 1)}{\sqrt{-a}} \\ & + \frac{1}{2} (a + b)^{\frac{3}{2}} \log\left(\left|\sqrt{a + be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b + \sqrt{a + b}}\right|\right) \operatorname{sgn}(e^{2x} - 1) \\ & - \frac{1}{2} (a + b)^{\frac{3}{2}} \log\left(\left|-\sqrt{a + be^{2x}} + \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b + \sqrt{a + b}}\right|\right) \operatorname{sgn}(e^{2x} - 1) \\ & - \frac{(a^2 + 2ab + b^2) \log\left(\left|-\left(\sqrt{a + be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b}\right)(a + b) + \sqrt{a + b}(a - b)\right|\right)}{2\sqrt{a + b}} \\ & + \frac{4\left(\left(\sqrt{a + be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b}\right)b^2 \operatorname{sgn}(e^{2x} - 1) + \sqrt{a + b}\right)}{\left(\sqrt{a + be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b}\right)^2 - 2\left(\sqrt{a + be^{2x}} - \sqrt{ae^{4x} + be^{4x} - 2ae^{2x} + 2be^{2x} + a + b}\right)} \end{aligned}$$

[In] integrate((a+b*coth(x)^2)^(3/2)*tanh(x),x, algorithm="giac")

```
[Out] -2*a^2*arctan(-1/2*(sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*
e^(2*x) + 2*b*e^(2*x) + a + b) + sqrt(a + b))/sqrt(-a))*sgn(e^(2*x) - 1)/sq
rt(-a) + 1/2*(a + b)^(3/2)*log(abs(sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b
*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) + sqrt(a + b)))*sgn(e^(2*x) -
1) - 1/2*(a + b)^(3/2)*log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e
^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) + sqrt(a + b)))*sgn(e^(2*x) - 1
) - 1/2*(a^2 + 2*a*b + b^2)*log(abs(-(sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x)
+ b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*(a + b) + sqrt(a + b)*(a
- b)))*sgn(e^(2*x) - 1)/sqrt(a + b) + 4*((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4
*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*b^2*sgn(e^(2*x) - 1)
+ sqrt(a + b)*b^2*sgn(e^(2*x) - 1))/((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x)
+ b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))^2 - 2*(sqrt(a + b)*e^(2*x)
) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*sqrt(a
+ b) + a - 3*b)
```

Mupad [F(-1)]

Timed out.

$$\int (a + b \coth^2(x))^{3/2} \tanh(x) dx = \int \tanh(x) (b \coth(x)^2 + a)^{3/2} dx$$

```
[In] int(tanh(x)*(a + b*coth(x)^2)^(3/2),x)
```

```
[Out] int(tanh(x)*(a + b*coth(x)^2)^(3/2), x)
```

3.27 $\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx$

Optimal result	200
Rubi [A] (verified)	200
Mathematica [B] (verified)	202
Maple [F]	203
Fricas [B] (verification not implemented)	203
Sympy [F]	206
Maxima [F]	206
Giac [F]	206
Mupad [F(-1)]	206

Optimal result

Integrand size = 17, antiderivative size = 77

$$\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx = -b^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right) + (a + b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right) - a \sqrt{a + b \coth^2(x)} \tanh(x)$$

[Out] $-b^{(3/2)} \operatorname{arctanh}(\coth(x) * b^{(1/2)} / (a + b * \coth(x)^2)^{(1/2)}) + (a + b)^{(3/2)} \operatorname{arctanh}(\coth(x) * (a + b)^{(1/2)} / (a + b * \coth(x)^2)^{(1/2)}) - a * (a + b * \coth(x)^2)^{(1/2)} * \tanh(x)$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3751, 485, 537, 223, 212, 385}

$$\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx = b^{3/2} \left(-\operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right) \right) + (a + b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}}\right) - a \tanh(x) \sqrt{a + b \coth^2(x)}$$

[In] $\operatorname{Int}[(a + b * \operatorname{Coth}[x]^2)^{(3/2)} * \operatorname{Tanh}[x]^2, x]$

[Out] $-(b^{3/2} \operatorname{ArcTanh}[\operatorname{Sqrt}[b] \operatorname{Coth}[x]] / \operatorname{Sqrt}[a + b \operatorname{Coth}[x]^2]) + (a + b)^{3/2} \operatorname{ArcTanh}[\operatorname{Sqrt}[a + b] \operatorname{Coth}[x]] / \operatorname{Sqrt}[a + b \operatorname{Coth}[x]^2] - a \operatorname{Sqrt}[a + b \operatorname{Coth}[x]^2] \operatorname{Tanh}[x]$

Rule 212

$\operatorname{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1 / (\operatorname{Rt}[a, 2] \operatorname{Rt}[-b, 2])) \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2] \cdot (x / \operatorname{Rt}[a, 2])], x] /;$ $\operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b] \ \&\& \ (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 223

$\operatorname{Int}[1 / \operatorname{Sqrt}[(a + (b \cdot x)^2)], x_Symbol] \rightarrow \operatorname{Subst}[\operatorname{Int}[1 / (1 - b \cdot x^2), x], x, x / \operatorname{Sqrt}[a + b \cdot x^2]] /;$ $\operatorname{FreeQ}\{a, b, x\} \ \&\& \ !\operatorname{GtQ}[a, 0]$

Rule 385

$\operatorname{Int}[(a + (b \cdot x)^n)^p / ((c + (d \cdot x)^n)), x_Symbol] \rightarrow \operatorname{Subst}[\operatorname{Int}[1 / (c - (b \cdot c - a \cdot d) \cdot x^n), x], x, x / (a + b \cdot x^n)^{1/n}] /;$ $\operatorname{FreeQ}\{a, b, c, d, x\} \ \&\& \ \operatorname{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \operatorname{EqQ}[n \cdot p + 1, 0] \ \&\& \ \operatorname{IntegerQ}[n]$

Rule 485

$\operatorname{Int}[(e \cdot x)^m \cdot (a + (b \cdot x)^n)^p \cdot (c + (d \cdot x)^n)^q, x_Symbol] \rightarrow \operatorname{Simp}[c \cdot (e \cdot x)^{m+1} \cdot (a + b \cdot x^n)^{p+1} \cdot (c + d \cdot x^n)^{q-1} / (a \cdot e^{m+1}), x] - \operatorname{Dist}[1 / (a \cdot e^{m+1}), \operatorname{Int}[(e \cdot x)^{m+n} \cdot (a + b \cdot x^n)^p \cdot (c + d \cdot x^n)^{q-2} \cdot \operatorname{Simp}[c \cdot (c \cdot b - a \cdot d) \cdot (m+1) + c \cdot n \cdot (b \cdot c \cdot (p+1) + a \cdot d \cdot (q-1)) + d \cdot ((c \cdot b - a \cdot d) \cdot (m+1) + c \cdot b \cdot n \cdot (p+q)) \cdot x^n, x], x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, e, p, x\} \ \&\& \ \operatorname{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \operatorname{IGtQ}[n, 0] \ \&\& \ \operatorname{GtQ}[q, 1] \ \&\& \ \operatorname{LtQ}[m, -1] \ \&\& \ \operatorname{IntBinomialQ}[a, b, c, d, e, m, n, p, q, x]$

Rule 537

$\operatorname{Int}[(e + (f \cdot x)^n) / ((a + (b \cdot x)^n) \operatorname{Sqrt}[c + (d \cdot x)^n]), x_Symbol] \rightarrow \operatorname{Dist}[f/b, \operatorname{Int}[1 / \operatorname{Sqrt}[c + d \cdot x^n], x], x] + \operatorname{Dist}[(b \cdot e - a \cdot f) / b, \operatorname{Int}[1 / ((a + b \cdot x^n) \operatorname{Sqrt}[c + d \cdot x^n]), x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, e, f, n, x\}$

Rule 3751

$\operatorname{Int}[(d \cdot \tan[e + (f \cdot x)])^m \cdot (a + (b \cdot (c \cdot \tan[e + (f \cdot x)]))^n)^p, x_Symbol] \rightarrow \operatorname{With}\{ff = \operatorname{FreeFactors}[\operatorname{Tan}[e + f \cdot x], x]\}, \operatorname{Dist}[c \cdot (ff/f), \operatorname{Subst}[\operatorname{Int}[(d \cdot ff \cdot (x/c))^m \cdot (a + b \cdot (ff \cdot x)^n)^p / (c^2 + ff^2 \cdot x^2), x], x, c \cdot (\operatorname{Tan}[e + f \cdot x] / ff)], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, e, f, m, n, p, x\} \ \&\& \ (\operatorname{IGtQ}[p, 0] \ || \ \operatorname{EqQ}[n, 2] \ || \ \operatorname{EqQ}[n, 4] \ || \ (\operatorname{IntegerQ}[p] \ \&\& \ \operatorname{RationalQ}[n]))$

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst} \left(\int \frac{(a + bx^2)^{3/2}}{x^2(1-x^2)} dx, x, \coth(x) \right) \\
&= -a\sqrt{a + b \coth^2(x)} \tanh(x) + \text{Subst} \left(\int \frac{a(a + 2b) + b^2x^2}{(1-x^2)\sqrt{a + bx^2}} dx, x, \coth(x) \right) \\
&= -a\sqrt{a + b \coth^2(x)} \tanh(x) - b^2 \text{Subst} \left(\int \frac{1}{\sqrt{a + bx^2}} dx, x, \coth(x) \right) \\
&\quad + (a + b)^2 \text{Subst} \left(\int \frac{1}{(1-x^2)\sqrt{a + bx^2}} dx, x, \coth(x) \right) \\
&= -a\sqrt{a + b \coth^2(x)} \tanh(x) - b^2 \text{Subst} \left(\int \frac{1}{1 - bx^2} dx, x, \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} \right) \\
&\quad + (a + b)^2 \text{Subst} \left(\int \frac{1}{1 - (a + b)x^2} dx, x, \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} \right) \\
&= -b^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) \\
&\quad + (a + b)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a + b} \coth(x)}{\sqrt{a + b \coth^2(x)}} \right) - a\sqrt{a + b \coth^2(x)} \tanh(x)
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 180 vs. 2(77) = 154.

Time = 0.47 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.34

$$\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx = \frac{\left(-\sqrt{2}b^{3/2}\sqrt{a+b} \operatorname{arctanh} \left(\frac{\sqrt{2}\sqrt{b} \cosh(x)}{\sqrt{-a+b+(a+b) \cosh(2x)}} \right) \cosh(x) + \sqrt{2}(a+b)^2 \operatorname{arctanh} \left(\frac{\sqrt{a+b} \cosh(x)}{\sqrt{-a+b+(a+b) \cosh(2x)}} \right) \right)}{\sqrt{2}\sqrt{a+b}\sqrt{-a+b+(a+b) \cosh(2x)}}$$

[In] Integrate[(a + b*Coth[x]^2)^(3/2)*Tanh[x]^2,x]

[Out] ((-(Sqrt[2]*b^(3/2)*Sqrt[a + b]*ArcTanh[(Sqrt[2]*Sqrt[b]*Cosh[x])/Sqrt[-a + b + (a + b)*Cosh[2*x]])*Cosh[x]) + Sqrt[2]*(a + b)^2*ArcTanh[(Sqrt[2]*Sqrt[a + b]*Cosh[x])/Sqrt[-a + b + (a + b)*Cosh[2*x]])*Cosh[x] - a*Sqrt[a + b]*Sqrt[-a + b + (a + b)*Cosh[2*x]])*Sqrt[(-a + b + (a + b)*Cosh[2*x])*Csch[x]^2]*Tanh[x])/(Sqrt[2]*Sqrt[a + b]*Sqrt[-a + b + (a + b)*Cosh[2*x]])

Maple [F]

$$\int (a + b \coth(x)^2)^{\frac{3}{2}} \tanh(x)^2 dx$$

[In] int((a+b*coth(x)^2)^(3/2)*tanh(x)^2,x)

[Out] int((a+b*coth(x)^2)^(3/2)*tanh(x)^2,x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 695 vs. 2(63) = 126.

Time = 0.42 (sec) , antiderivative size = 4025, normalized size of antiderivative = 52.27

$$\int (a + b \coth^2(x))^{\frac{3}{2}} \tanh^2(x) dx = \text{Too large to display}$$

[In] integrate((a+b*coth(x)^2)^(3/2)*tanh(x)^2,x, algorithm="fricas")

[Out] [1/4*(((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a + b)*sqrt(a + b)*log(((a*b^2 + b^3)*cosh(x)^8 + 8*(a*b^2 + b^3)*cosh(x)*sinh(x)^7 + (a*b^2 + b^3)*sinh(x)^8 + 2*(a*b^2 + 2*b^3)*cosh(x)^6 + 2*(a*b^2 + 2*b^3 + 14*(a*b^2 + b^3)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a*b^2 + b^3)*cosh(x)^3 + 3*(a*b^2 + 2*b^3)*cosh(x))*sinh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^4 + (70*(a*b^2 + b^3)*cosh(x)^4 + a^3 - a^2*b + 4*a*b^2 + 6*b^3 + 30*(a*b^2 + 2*b^3)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a*b^2 + b^3)*cosh(x)^5 + 10*(a*b^2 + 2*b^3)*cosh(x)^3 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 - 3*a*b^2 - 2*b^3)*cosh(x)^2 + 2*(14*(a*b^2 + b^3)*cosh(x)^6 + 15*(a*b^2 + 2*b^3)*cosh(x)^4 - a^3 + 3*a*b^2 + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(b^2*cosh(x)^6 + 6*b^2*cosh(x)*sinh(x)^5 + b^2*sinh(x)^6 + 3*b^2*cosh(x)^4 + 3*(5*b^2*cosh(x)^2 + b^2)*sinh(x)^4 + 4*(5*b^2*cosh(x)^3 + 3*b^2*cosh(x))*sinh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x)^2 + (15*b^2*cosh(x)^4 + 18*b^2*cosh(x)^2 - a^2 + 2*a*b + 3*b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(3*b^2*cosh(x)^5 + 6*b^2*cosh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a*b^2 + b^3)*cosh(x)^7 + 3*(a*b^2 + 2*b^3)*cosh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^3 - (a^3 - 3*a*b^2 - 2*b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6) + 2*(b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + b)*sqrt(b)*log(-(a + 2*b)*cosh(x)^4 + 4*(a + 2*b)*cosh(x)*sinh(x)^3 + (a + 2*b)*sinh(x)^4 - 2*(a - 2*b)*cosh(x)^2 + 2*(3*(a + 2*b)*cosh(x)^2 - a + 2*b)*sinh(x)^2 - 2*sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*((a + 2*b)*cosh(x)^3 - (a - 2*b)*

$$\begin{aligned}
& \cosh(x) \cdot \sinh(x) + a + 2b) / (\cosh(x)^4 + 4 \cdot \cosh(x) \cdot \sinh(x)^3 + \sinh(x)^4 + \\
& 2 \cdot (3 \cdot \cosh(x)^2 - 1) \cdot \sinh(x)^2 - 2 \cdot \cosh(x)^2 + 4 \cdot (\cosh(x)^3 - \cosh(x)) \cdot \sinh(x) \\
& + 1)) + ((a + b) \cdot \cosh(x)^2 + 2 \cdot (a + b) \cdot \cosh(x) \cdot \sinh(x) + (a + b) \cdot \sinh(x) \\
& ^2 + a + b) \cdot \sqrt{a + b} \cdot \log(-((a + b) \cdot \cosh(x)^4 + 4 \cdot (a + b) \cdot \cosh(x) \cdot \sinh(x) \\
& ^3 + (a + b) \cdot \sinh(x)^4 - 2 \cdot a \cdot \cosh(x)^2 + 2 \cdot (3 \cdot (a + b) \cdot \cosh(x)^2 - a) \cdot \sinh(x) \\
&)^2 + \sqrt{2} \cdot (\cosh(x)^2 + 2 \cdot \cosh(x) \cdot \sinh(x) + \sinh(x)^2 - 1) \cdot \sqrt{a + b} \cdot \sqrt{ \\
& \sqrt{((a + b) \cdot \cosh(x)^2 + (a + b) \cdot \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cdot \cosh(x) \cdot \\
& \sinh(x) + \sinh(x)^2)} + 4 \cdot ((a + b) \cdot \cosh(x)^3 - a \cdot \cosh(x)) \cdot \sinh(x) + a + b) / \\
& (\cosh(x)^2 + 2 \cdot \cosh(x) \cdot \sinh(x) + \sinh(x)^2)) - 4 \cdot \sqrt{2} \cdot a \cdot \sqrt{((a + b) \cdot \cosh(x) \\
& ^2 + (a + b) \cdot \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cdot \cosh(x) \cdot \sinh(x) + \sinh(x) \\
& ^2))} / (\cosh(x)^2 + 2 \cdot \cosh(x) \cdot \sinh(x) + \sinh(x)^2 + 1), 1/4 \cdot (4 \cdot (b \cdot \cosh(x) \\
& ^2 + 2 \cdot b \cdot \cosh(x) \cdot \sinh(x) + b \cdot \sinh(x)^2 + b) \cdot \sqrt{-b} \cdot \arctan(\sqrt{2} \cdot (\cosh(x) \\
& ^2 + 2 \cdot \cosh(x) \cdot \sinh(x) + \sinh(x)^2 + 1) \cdot \sqrt{-b}) \cdot \sqrt{((a + b) \cdot \cosh(x)^2 + \\
& (a + b) \cdot \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cdot \cosh(x) \cdot \sinh(x) + \sinh(x)^2)} / ((a \\
& + b) \cdot \cosh(x)^4 + 4 \cdot (a + b) \cdot \cosh(x) \cdot \sinh(x)^3 + (a + b) \cdot \sinh(x)^4 - 2 \cdot (a - \\
& b) \cdot \cosh(x)^2 + 2 \cdot (3 \cdot (a + b) \cdot \cosh(x)^2 - a + b) \cdot \sinh(x)^2 + 4 \cdot ((a + b) \cdot \cosh(x) \\
& ^3 - (a - b) \cdot \cosh(x)) \cdot \sinh(x) + a + b)) + ((a + b) \cdot \cosh(x)^2 + 2 \cdot (a + b) \cdot \\
& \cosh(x) \cdot \sinh(x) + (a + b) \cdot \sinh(x)^2 + a + b) \cdot \sqrt{a + b} \cdot \log(((a \cdot b^2 + b^3) \\
& \cdot \cosh(x)^8 + 8 \cdot (a \cdot b^2 + b^3) \cdot \cosh(x) \cdot \sinh(x)^7 + (a \cdot b^2 + b^3) \cdot \sinh(x)^8 + \\
& 2 \cdot (a \cdot b^2 + 2 \cdot b^3) \cdot \cosh(x)^6 + 2 \cdot (a \cdot b^2 + 2 \cdot b^3 + 14 \cdot (a \cdot b^2 + b^3) \cdot \cosh(x)^2) \\
&) \cdot \sinh(x)^6 + 4 \cdot (14 \cdot (a \cdot b^2 + b^3) \cdot \cosh(x)^3 + 3 \cdot (a \cdot b^2 + 2 \cdot b^3) \cdot \cosh(x)) \cdot \sinh(x) \\
& ^5 + (a^3 - a^2 \cdot b + 4 \cdot a \cdot b^2 + 6 \cdot b^3) \cdot \cosh(x)^4 + (70 \cdot (a \cdot b^2 + b^3) \cdot \cosh(x) \\
& ^4 + a^3 - a^2 \cdot b + 4 \cdot a \cdot b^2 + 6 \cdot b^3 + 30 \cdot (a \cdot b^2 + 2 \cdot b^3) \cdot \cosh(x)^2) \cdot \sinh(x) \\
& ^4 + 4 \cdot (14 \cdot (a \cdot b^2 + b^3) \cdot \cosh(x)^5 + 10 \cdot (a \cdot b^2 + 2 \cdot b^3) \cdot \cosh(x)^3 + (a^3 \\
& - a^2 \cdot b + 4 \cdot a \cdot b^2 + 6 \cdot b^3) \cdot \cosh(x)) \cdot \sinh(x)^3 + a^3 + 3 \cdot a^2 \cdot b + 3 \cdot a \cdot b^2 + \\
& b^3 - 2 \cdot (a^3 - 3 \cdot a \cdot b^2 - 2 \cdot b^3) \cdot \cosh(x)^2 + 2 \cdot (14 \cdot (a \cdot b^2 + b^3) \cdot \cosh(x)^6 + \\
& 15 \cdot (a \cdot b^2 + 2 \cdot b^3) \cdot \cosh(x)^4 - a^3 + 3 \cdot a \cdot b^2 + 2 \cdot b^3 + 3 \cdot (a^3 - a^2 \cdot b + 4 \cdot \\
& a \cdot b^2 + 6 \cdot b^3) \cdot \cosh(x)^2) \cdot \sinh(x)^2 + \sqrt{2} \cdot (b^2 \cdot \cosh(x)^6 + 6 \cdot b^2 \cdot \cosh(x) \\
&) \cdot \sinh(x)^5 + b^2 \cdot \sinh(x)^6 + 3 \cdot b^2 \cdot \cosh(x)^4 + 3 \cdot (5 \cdot b^2 \cdot \cosh(x)^2 + b^2) \cdot \sinh(x) \\
& ^4 + 4 \cdot (5 \cdot b^2 \cdot \cosh(x)^3 + 3 \cdot b^2 \cdot \cosh(x)) \cdot \sinh(x)^3 - (a^2 - 2 \cdot a \cdot b - 3 \\
& \cdot b^2) \cdot \cosh(x)^2 + (15 \cdot b^2 \cdot \cosh(x)^4 + 18 \cdot b^2 \cdot \cosh(x)^2 - a^2 + 2 \cdot a \cdot b + 3 \cdot b^2) \\
& \cdot \sinh(x)^2 + a^2 + 2 \cdot a \cdot b + b^2 + 2 \cdot (3 \cdot b^2 \cdot \cosh(x)^5 + 6 \cdot b^2 \cdot \cosh(x)^3 - (\\
& a^2 - 2 \cdot a \cdot b - 3 \cdot b^2) \cdot \cosh(x)) \cdot \sinh(x)) \cdot \sqrt{a + b} \cdot \sqrt{((a + b) \cdot \cosh(x)^2 \\
& + (a + b) \cdot \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cdot \cosh(x) \cdot \sinh(x) + \sinh(x)^2)} + \\
& 4 \cdot (2 \cdot (a \cdot b^2 + b^3) \cdot \cosh(x)^7 + 3 \cdot (a \cdot b^2 + 2 \cdot b^3) \cdot \cosh(x)^5 + (a^3 - a^2 \cdot b \\
& + 4 \cdot a \cdot b^2 + 6 \cdot b^3) \cdot \cosh(x)^3 - (a^3 - 3 \cdot a \cdot b^2 - 2 \cdot b^3) \cdot \cosh(x)) \cdot \sinh(x)) / (\cosh(x) \\
& ^6 + 6 \cdot \cosh(x)^5 \cdot \sinh(x) + 15 \cdot \cosh(x)^4 \cdot \sinh(x)^2 + 20 \cdot \cosh(x)^3 \cdot \sinh(x) \\
& ^3 + 15 \cdot \cosh(x)^2 \cdot \sinh(x)^4 + 6 \cdot \cosh(x) \cdot \sinh(x)^5 + \sinh(x)^6)) + ((a + \\
& b) \cdot \cosh(x)^2 + 2 \cdot (a + b) \cdot \cosh(x) \cdot \sinh(x) + (a + b) \cdot \sinh(x)^2 + a + b) \cdot \sqrt{a + b} \cdot \log(-((a + b) \cdot \cosh(x) \\
& ^4 + 4 \cdot (a + b) \cdot \cosh(x) \cdot \sinh(x)^3 + (a + b) \cdot \sinh(x)^4 - 2 \cdot a \cdot \cosh(x)^2 + 2 \cdot (3 \cdot (a + b) \cdot \cosh(x)^2 - a) \cdot \sinh(x) \\
&)^2 + \sqrt{2} \cdot (\cosh(x)^2 + 2 \cdot \cosh(x) \cdot \sinh(x) + \sinh(x)^2 - 1) \cdot \sqrt{a + b} \cdot \sqrt{((a + b) \cdot \cosh(x) \\
& ^2 + (a + b) \cdot \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cdot \cosh(x) \cdot \sinh(x) + \sinh(x) \\
& ^2)} + 4 \cdot ((a + b) \cdot \cosh(x)^3 - a \cdot \cosh(x)) \cdot \sinh(x) + a + b) / (\cosh(x)^2 + 2 \cdot \cosh(x) \cdot \sinh(x) \\
& + \sinh(x)^2)) - 4 \cdot \sqrt{2} \cdot a \cdot \sqrt{((a + b) \cdot \cosh(x)^2 + (a + b)
\end{aligned}$$

$$\begin{aligned}
& *sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2))/((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1), -1/2*(((a + b)*cosh(x)^2 + 2*(a + b) \\
&)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a + b)*sqrt(-a - b)*arctan(sqrt(2)* \\
& (b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + a + b)*sqrt(-a - b)*sqrt \\
& (((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh \\
& h(x) + sinh(x)^2))/((a*b + b^2)*cosh(x)^4 + 4*(a*b + b^2)*cosh(x)*sinh(x)^3 \\
& + (a*b + b^2)*sinh(x)^4 - (a^2 - a*b - 2*b^2)*cosh(x)^2 + (6*(a*b + b^2)*c \\
& osh(x)^2 - a^2 + a*b + 2*b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(2*(a*b + b \\
& ^2)*cosh(x)^3 - (a^2 - a*b - 2*b^2)*cosh(x))*sinh(x))) + ((a + b)*cosh(x)^2 \\
& + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a + b)*sqrt(-a - b)*arct \\
& an(sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(-a - b)*sqrt \\
& t(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*si \\
& nh(x) + sinh(x)^2))/((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + \\
& b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(\\
& x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)) - (b*cosh(\\
& x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + b)*sqrt(b)*log(-((a + 2*b)*cosh(\\
& x)^4 + 4*(a + 2*b)*cosh(x)*sinh(x)^3 + (a + 2*b)*sinh(x)^4 - 2*(a - 2*b)*co \\
& sh(x)^2 + 2*(3*(a + 2*b)*cosh(x)^2 - a + 2*b)*sinh(x)^2 - 2*sqrt(2)*(cosh(x) \\
&)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(b)*sqrt(((a + b)*cosh(x)^2 + \\
& (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4 \\
& *((a + 2*b)*cosh(x)^3 - (a - 2*b)*cosh(x))*sinh(x) + a + 2*b)/(cosh(x)^4 + \\
& 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*sinh(x)^2 - 2*cosh(x) \\
& ^2 + 4*(cosh(x)^3 - cosh(x))*sinh(x) + 1)) + 2*sqrt(2)*a*sqrt(((a + b)*cosh \\
& (x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x) \\
& ^2)))/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1), -1/2*(((a + b)*cosh(\\
& x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a + b)*sqrt(-a - b)* \\
& arctan(sqrt(2)*(b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + a + b)*sq \\
& rt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 \\
& - 2*cosh(x)*sinh(x) + sinh(x)^2))/((a*b + b^2)*cosh(x)^4 + 4*(a*b + b^2)*co \\
& sh(x)*sinh(x)^3 + (a*b + b^2)*sinh(x)^4 - (a^2 - a*b - 2*b^2)*cosh(x)^2 + (\\
& 6*(a*b + b^2)*cosh(x)^2 - a^2 + a*b + 2*b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 \\
& + 2*(2*(a*b + b^2)*cosh(x)^3 - (a^2 - a*b - 2*b^2)*cosh(x))*sinh(x))) + ((a \\
& + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a + b)*sq \\
& rt(-a - b)*arctan(sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*s \\
& qrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 \\
& - 2*cosh(x)*sinh(x) + sinh(x)^2))/((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*s \\
& inh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 \\
& - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + \\
& b)) - 2*(b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + b)*sqrt(-b)*arc \\
& tan(sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(-b)*sqrt((\\
& (a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(\\
& x) + sinh(x)^2))/((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b) \\
& *sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^ \\
& 2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)) + 2*sqrt(2)*a \\
& *sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)
\end{aligned}$$

) $\sinh(x) + \sinh(x)^2$))/($\cosh(x)^2 + 2\cosh(x)\sinh(x) + \sinh(x)^2 + 1$)]

Sympy [F]

$$\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx = \int (a + b \coth^2(x))^{\frac{3}{2}} \tanh^2(x) dx$$

[In] integrate((a+b*coth(x)**2)**(3/2)*tanh(x)**2,x)

[Out] Integral((a + b*coth(x)**2)**(3/2)*tanh(x)**2, x)

Maxima [F]

$$\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx = \int (b \coth(x)^2 + a)^{\frac{3}{2}} \tanh(x)^2 dx$$

[In] integrate((a+b*coth(x)^2)^(3/2)*tanh(x)^2,x, algorithm="maxima")

[Out] integrate((b*coth(x)^2 + a)^(3/2)*tanh(x)^2, x)

Giac [F]

$$\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx = \int (b \coth(x)^2 + a)^{\frac{3}{2}} \tanh(x)^2 dx$$

[In] integrate((a+b*coth(x)^2)^(3/2)*tanh(x)^2,x, algorithm="giac")

[Out] sage0*x

Mupad [F(-1)]

Timed out.

$$\int (a + b \coth^2(x))^{3/2} \tanh^2(x) dx = \int \tanh(x)^2 (b \coth(x)^2 + a)^{3/2} dx$$

[In] int(tanh(x)^2*(a + b*coth(x)^2)^(3/2),x)

[Out] int(tanh(x)^2*(a + b*coth(x)^2)^(3/2), x)

3.28 $\int \sqrt{1 + \coth^2(x)} dx$

Optimal result	207
Rubi [A] (verified)	207
Mathematica [A] (verified)	209
Maple [B] (verified)	209
Fricas [B] (verification not implemented)	209
Sympy [F]	210
Maxima [F]	210
Giac [B] (verification not implemented)	211
Mupad [B] (verification not implemented)	211

Optimal result

Integrand size = 10, antiderivative size = 31

$$\int \sqrt{1 + \coth^2(x)} dx = -\operatorname{arcsinh}(\coth(x)) + \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2} \coth(x)}{\sqrt{1 + \coth^2(x)}}\right)$$

[Out] $-\operatorname{arcsinh}(\coth(x)) + \operatorname{arctanh}(\coth(x) * 2^{(1/2)} / (1 + \coth(x)^2)^{(1/2)}) * 2^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3742, 399, 221, 385, 212}

$$\int \sqrt{1 + \coth^2(x)} dx = \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2} \coth(x)}{\sqrt{\coth^2(x) + 1}}\right) - \operatorname{arcsinh}(\coth(x))$$

[In] $\operatorname{Int}[\operatorname{Sqrt}[1 + \operatorname{Coth}[x]^2], x]$

[Out] $-\operatorname{ArcSinh}[\operatorname{Coth}[x]] + \operatorname{Sqrt}[2] * \operatorname{ArcTanh}[(\operatorname{Sqrt}[2] * \operatorname{Coth}[x]) / \operatorname{Sqrt}[1 + \operatorname{Coth}[x]^2]]$

Rule 212

$\operatorname{Int}[(a_+ + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1 / (\operatorname{Rt}[a, 2] * \operatorname{Rt}[-b, 2])) * \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2] * (x / \operatorname{Rt}[a, 2])], x] /;$ FreeQ[{a, b}, x] && NegQ[a/b] && (Gt Q[a, 0] || LtQ[b, 0])

Rule 221

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt
[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Su
bst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b
, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 399

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Di
st[b/d, Int[(a + b*x^n)^(p - 1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^
n)^(p - 1)/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*
d, 0] && EqQ[n*(p - 1) + 1, 0] && IntegerQ[n]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :=
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*
ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{\sqrt{1+x^2}}{1-x^2} dx, x, \coth(x)\right) \\
&= 2\text{Subst}\left(\int \frac{1}{(1-x^2)\sqrt{1+x^2}} dx, x, \coth(x)\right) - \text{Subst}\left(\int \frac{1}{\sqrt{1+x^2}} dx, x, \coth(x)\right) \\
&= -\text{arcsinh}(\coth(x)) + 2\text{Subst}\left(\int \frac{1}{1-2x^2} dx, x, \frac{\coth(x)}{\sqrt{1+\coth^2(x)}}\right) \\
&= -\text{arcsinh}(\coth(x)) + \sqrt{2}\text{arctanh}\left(\frac{\sqrt{2}\coth(x)}{\sqrt{1+\coth^2(x)}}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.94

$$\int \sqrt{1 + \coth^2(x)} dx$$

$$= \frac{\sqrt{1 + \coth^2(x)} \left(-\operatorname{arctanh} \left(\frac{\cosh(x)}{\sqrt{\cosh(2x)}} \right) + \sqrt{2} \log \left(\sqrt{2} \cosh(x) + \sqrt{\cosh(2x)} \right) \right) \sinh(x)}{\sqrt{\cosh(2x)}}$$

[In] Integrate[Sqrt[1 + Coth[x]^2], x]

[Out] (Sqrt[1 + Coth[x]^2]*(-ArcTanh[Cosh[x]/Sqrt[Cosh[2*x]]] + Sqrt[2]*Log[Sqrt[2]*Cosh[x] + Sqrt[Cosh[2*x]]])*Sinh[x])/Sqrt[Cosh[2*x]]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 96 vs. 2(25) = 50.

Time = 0.23 (sec) , antiderivative size = 97, normalized size of antiderivative = 3.13

method	result
derivativedivides	$-\frac{\sqrt{(\coth(x)-1)^2+2\coth(x)}}{2} - \operatorname{arcsinh}(\coth(x)) + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(2+2\coth(x))\sqrt{2}}{4\sqrt{(\coth(x)-1)^2+2\coth(x)}}\right)}{2} + \frac{\sqrt{1+\coth(x)}}{2}$
default	$-\frac{\sqrt{(\coth(x)-1)^2+2\coth(x)}}{2} - \operatorname{arcsinh}(\coth(x)) + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(2+2\coth(x))\sqrt{2}}{4\sqrt{(\coth(x)-1)^2+2\coth(x)}}\right)}{2} + \frac{\sqrt{1+\coth(x)}}{2}$

[In] int((1+coth(x)^2)^(1/2), x, method=_RETURNVERBOSE)

[Out] -1/2*((coth(x)-1)^2+2*coth(x))^(1/2)-arcsinh(coth(x))+1/2*2^(1/2)*arctanh(1/4*(2+2*coth(x))*2^(1/2)/((coth(x)-1)^2+2*coth(x))^(1/2))+1/2*((1+coth(x))^2-2*coth(x))^(1/2)-1/2*2^(1/2)*arctanh(1/4*(2-2*coth(x))*2^(1/2)/((1+coth(x))^2-2*coth(x))^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 683 vs. 2(25) = 50.

Time = 0.26 (sec) , antiderivative size = 683, normalized size of antiderivative = 22.03

$$\int \sqrt{1 + \coth^2(x)} dx = \text{Too large to display}$$

[In] integrate((1+coth(x)^2)^(1/2), x, algorithm="fricas")

```
[Out] 1/4*sqrt(2)*log(2*(cosh(x)^8 + 8*cosh(x)*sinh(x)^7 + sinh(x)^8 + (28*cosh(x)^2 + 3)*sinh(x)^6 + 3*cosh(x)^6 + 2*(28*cosh(x)^3 + 9*cosh(x))*sinh(x)^5 + 5*(14*cosh(x)^4 + 9*cosh(x)^2 + 1)*sinh(x)^4 + 5*cosh(x)^4 + 4*(14*cosh(x)^5 + 15*cosh(x)^3 + 5*cosh(x))*sinh(x)^3 + (28*cosh(x)^6 + 45*cosh(x)^4 + 30*cosh(x)^2 + 4)*sinh(x)^2 + 4*cosh(x)^2 + 2*(4*cosh(x)^7 + 9*cosh(x)^5 + 10*cosh(x)^3 + 4*cosh(x))*sinh(x) + (sqrt(2)*cosh(x)^6 + 6*sqrt(2)*cosh(x)*sinh(x)^5 + sqrt(2)*sinh(x)^6 + 3*(5*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x)^4 + 3*sqrt(2)*cosh(x)^4 + 4*(5*sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x))*sinh(x)^3 + (15*sqrt(2)*cosh(x)^4 + 18*sqrt(2)*cosh(x)^2 + 4*sqrt(2))*sinh(x)^2 + 4*sqrt(2)*cosh(x)^2 + 2*(3*sqrt(2)*cosh(x)^5 + 6*sqrt(2)*cosh(x)^3 + 4*sqrt(2)*cosh(x))*sinh(x) + 4*sqrt(2))*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4)/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6)) + 1/4*sqrt(2)*log(-2*(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 - 1)*sinh(x)^2 - cosh(x)^2 + 2*(2*cosh(x)^3 - cosh(x))*sinh(x) + (sqrt(2)*cosh(x)^2 + 2*sqrt(2)*cosh(x)*sinh(x) + sqrt(2)*sinh(x)^2 - sqrt(2))*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 1)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)) - 1/2*log((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 2*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 1)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)) + 1/2*log((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 2*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 1)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2))
```

Sympy [F]

$$\int \sqrt{1 + \coth^2(x)} dx = \int \sqrt{\coth^2(x) + 1} dx$$

```
[In] integrate((1+coth(x)**2)**(1/2),x)
```

```
[Out] Integral(sqrt(coth(x)**2 + 1), x)
```

Maxima [F]

$$\int \sqrt{1 + \coth^2(x)} dx = \int \sqrt{\coth(x)^2 + 1} dx$$

```
[In] integrate((1+coth(x)^2)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(coth(x)^2 + 1), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs. 2(25) = 50.

Time = 0.28 (sec) , antiderivative size = 119, normalized size of antiderivative = 3.84

$$\int \sqrt{1 + \coth^2(x)} dx$$

$$= \frac{1}{2} \sqrt{2} \left(\sqrt{2} \log \left(\frac{|-2\sqrt{2} + 2\sqrt{e^{4x} + 1} - 2e^{2x} + 2|}{2(\sqrt{2} + \sqrt{e^{4x} + 1} - e^{2x} + 1)} \right) + \log(\sqrt{e^{4x} + 1} - e^{2x} + 1) - \log(\sqrt{e^{4x} + 1} + e^{2x} + 1) \right) - 1)$$

[In] integrate((1+coth(x)^2)^(1/2),x, algorithm="giac")

[Out] 1/2*sqrt(2)*(sqrt(2)*log(1/2*abs(-2*sqrt(2) + 2*sqrt(e^(4*x) + 1) - 2*e^(2*x) + 2)/(sqrt(2) + sqrt(e^(4*x) + 1) - e^(2*x) + 1)) + log(sqrt(e^(4*x) + 1) - e^(2*x) + 1) - log(sqrt(e^(4*x) + 1) + e^(2*x) + 1))*sgn(e^(2*x) - 1)

Mupad [B] (verification not implemented)

Time = 2.04 (sec) , antiderivative size = 68, normalized size of antiderivative = 2.19

$$\int \sqrt{1 + \coth^2(x)} dx = \frac{\sqrt{2} \left(\ln \left(\coth(x) + \sqrt{2} \sqrt{\coth(x)^2 + 1} + 1 \right) - \ln(\coth(x) - 1) \right)}{2} - \operatorname{asinh}(\coth(x)) + \frac{\sqrt{2} \left(\ln(\coth(x) + 1) - \ln \left(\sqrt{2} \sqrt{\coth(x)^2 + 1} - \coth(x) + 1 \right) \right)}{2}$$

[In] int((coth(x)^2 + 1)^(1/2),x)

[Out] (2^(1/2)*(log(coth(x) + 2^(1/2)*(coth(x)^2 + 1)^(1/2) + 1) - log(coth(x) - 1)))/2 - asinh(coth(x)) + (2^(1/2)*(log(coth(x) + 1) - log(2^(1/2)*(coth(x)^2 + 1)^(1/2) - coth(x) + 1)))/2

3.29 $\int \sqrt{-1 - \coth^2(x)} dx$

Optimal result	212
Rubi [A] (verified)	212
Mathematica [A] (verified)	214
Maple [B] (verified)	214
Fricas [C] (verification not implemented)	215
Sympy [F]	215
Maxima [F]	216
Giac [C] (verification not implemented)	216
Mupad [B] (verification not implemented)	216

Optimal result

Integrand size = 12, antiderivative size = 45

$$\int \sqrt{-1 - \coth^2(x)} dx = \arctan\left(\frac{\coth(x)}{\sqrt{-1 - \coth^2(x)}}\right) - \sqrt{2} \arctan\left(\frac{\sqrt{2} \coth(x)}{\sqrt{-1 - \coth^2(x)}}\right)$$

[Out] $\arctan(\coth(x)/(-1-\coth(x)^2)^{(1/2)})-\arctan(\coth(x)*2^{(1/2)/(-1-\coth(x)^2)^{(1/2)})*2^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3742, 399, 223, 209, 385}

$$\int \sqrt{-1 - \coth^2(x)} dx = \arctan\left(\frac{\coth(x)}{\sqrt{-\coth^2(x) - 1}}\right) - \sqrt{2} \arctan\left(\frac{\sqrt{2} \coth(x)}{\sqrt{-\coth^2(x) - 1}}\right)$$

[In] $\text{Int}[\text{Sqrt}[-1 - \text{Coth}[x]^2], x]$

[Out] $\text{ArcTan}[\text{Coth}[x]/\text{Sqrt}[-1 - \text{Coth}[x]^2]] - \text{Sqrt}[2]*\text{ArcTan}[(\text{Sqrt}[2]*\text{Coth}[x])/\text{Sqrt}[-1 - \text{Coth}[x]^2]]$

Rule 209

$\text{Int}[(a_+ + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /;$ $\text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rule 223

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x],
x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Su
bst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b
, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 399

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Di
st[b/d, Int[(a + b*x^n)^(p - 1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^
n)^(p - 1)/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*
d, 0] && EqQ[n*(p - 1) + 1, 0] && IntegerQ[n]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :=
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(
ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst} \left(\int \frac{\sqrt{-1-x^2}}{1-x^2} dx, x, \coth(x) \right) \\
&= - \left(2 \text{Subst} \left(\int \frac{1}{\sqrt{-1-x^2}(1-x^2)} dx, x, \coth(x) \right) \right) + \text{Subst} \left(\int \frac{1}{\sqrt{-1-x^2}} dx, x, \coth(x) \right) \\
&= - \left(2 \text{Subst} \left(\int \frac{1}{1+2x^2} dx, x, \frac{\coth(x)}{\sqrt{-1-\coth^2(x)}} \right) \right) \\
&\quad + \text{Subst} \left(\int \frac{1}{1+x^2} dx, x, \frac{\coth(x)}{\sqrt{-1-\coth^2(x)}} \right) \\
&= \arctan \left(\frac{\coth(x)}{\sqrt{-1-\coth^2(x)}} \right) - \sqrt{2} \arctan \left(\frac{\sqrt{2} \coth(x)}{\sqrt{-1-\coth^2(x)}} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.38

$$\int \sqrt{-1 - \coth^2(x)} dx$$

$$= \frac{\sqrt{-1 - \coth^2(x)} \left(-\operatorname{arctanh}\left(\frac{\cosh(x)}{\sqrt{\cosh(2x)}}\right) + \sqrt{2} \log\left(\sqrt{2} \cosh(x) + \sqrt{\cosh(2x)}\right) \right) \sinh(x)}{\sqrt{\cosh(2x)}}$$

[In] Integrate[Sqrt[-1 - Coth[x]^2], x]

[Out] (Sqrt[-1 - Coth[x]^2]*(-ArcTanh[Cosh[x]/Sqrt[Cosh[2*x]]] + Sqrt[2]*Log[Sqrt[2]*Cosh[x] + Sqrt[Cosh[2*x]]])*Sinh[x])/Sqrt[Cosh[2*x]]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(37) = 74.

Time = 0.20 (sec) , antiderivative size = 142, normalized size of antiderivative = 3.16

method	result
derivativedivides	$\frac{\sqrt{-(1+\coth(x))^2+2\coth(x)}}{2} + \frac{\arctan\left(\frac{\coth(x)}{\sqrt{-(1+\coth(x))^2+2\coth(x)}}\right)}{2} - \frac{\sqrt{2} \arctan\left(\frac{(-2+2\coth(x))\sqrt{2}}{4\sqrt{-(1+\coth(x))^2+2\coth(x)}}\right)}{2} - \frac{\sqrt{2} \arctan\left(\frac{(-2+2\coth(x))\sqrt{2}}{4\sqrt{-(1+\coth(x))^2+2\coth(x)}}\right)}{2}$
default	$\frac{\sqrt{-(1+\coth(x))^2+2\coth(x)}}{2} + \frac{\arctan\left(\frac{\coth(x)}{\sqrt{-(1+\coth(x))^2+2\coth(x)}}\right)}{2} - \frac{\sqrt{2} \arctan\left(\frac{(-2+2\coth(x))\sqrt{2}}{4\sqrt{-(1+\coth(x))^2+2\coth(x)}}\right)}{2} - \frac{\sqrt{2} \arctan\left(\frac{(-2+2\coth(x))\sqrt{2}}{4\sqrt{-(1+\coth(x))^2+2\coth(x)}}\right)}{2}$

[In] int((-1-coth(x)^2)^(1/2), x, method=_RETURNVERBOSE)

[Out] 1/2*(-(1+coth(x))^2+2*coth(x))^(1/2)+1/2*arctan(coth(x)/(-(1+coth(x))^2+2*coth(x))^(1/2))-1/2*2^(1/2)*arctan(1/4*(-2+2*coth(x))*2^(1/2)/(-(1+coth(x))^2+2*coth(x))^(1/2))-1/2*(-(coth(x)-1)^2-2*coth(x))^(1/2)+1/2*arctan(coth(x)/(-(coth(x)-1)^2-2*coth(x))^(1/2))+1/2*2^(1/2)*arctan(1/4*(-2-2*coth(x))*2^(1/2)/(-(coth(x)-1)^2-2*coth(x))^(1/2))

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 226, normalized size of antiderivative = 5.02

$$\begin{aligned}
 & \int \sqrt{-1 - \coth^2(x)} dx \\
 &= -\frac{1}{4} \sqrt{-2} \log \left(-\left(\sqrt{-2} \sqrt{-2 e^{4x} - 2} + 2 e^{2x} - 2 \right) e^{(-2x)} \right) \\
 &+ \frac{1}{4} \sqrt{-2} \log \left(\left(\sqrt{-2} \sqrt{-2 e^{4x} - 2} - 2 e^{2x} + 2 \right) e^{(-2x)} \right) \\
 &+ \frac{1}{4} \sqrt{-2} \log \left(-2 \left(\sqrt{-2 e^{4x} - 2} (e^{2x} + 2) + \sqrt{-2} e^{4x} + \sqrt{-2} e^{2x} + 2 \sqrt{-2} \right) e^{(-4x)} \right) \\
 &- \frac{1}{4} \sqrt{-2} \log \left(-2 \left(\sqrt{-2 e^{4x} - 2} (e^{2x} + 2) - \sqrt{-2} e^{4x} - \sqrt{-2} e^{2x} - 2 \sqrt{-2} \right) e^{(-4x)} \right) \\
 &+ \frac{1}{2} i \log \left(-4 \left(i \sqrt{-2 e^{4x} - 2} + e^{2x} + 1 \right) e^{(-2x)} \right) \\
 &- \frac{1}{2} i \log \left(-4 \left(-i \sqrt{-2 e^{4x} - 2} + e^{2x} + 1 \right) e^{(-2x)} \right)
 \end{aligned}$$

[In] integrate((-1-coth(x)^2)^(1/2),x, algorithm="fricas")

[Out] -1/4*sqrt(-2)*log(-(sqrt(-2)*sqrt(-2*e^(4*x) - 2) + 2*e^(2*x) - 2)*e^(-2*x)) + 1/4*sqrt(-2)*log((sqrt(-2)*sqrt(-2*e^(4*x) - 2) - 2*e^(2*x) + 2)*e^(-2*x)) + 1/4*sqrt(-2)*log(-2*(sqrt(-2*e^(4*x) - 2)*(e^(2*x) + 2) + sqrt(-2)*e^(4*x) + sqrt(-2)*e^(2*x) + 2*sqrt(-2))*e^(-4*x)) - 1/4*sqrt(-2)*log(-2*(sqrt(-2*e^(4*x) - 2)*(e^(2*x) + 2) - sqrt(-2)*e^(4*x) - sqrt(-2)*e^(2*x) - 2*sqrt(-2))*e^(-4*x)) + 1/2*I*log(-4*(I*sqrt(-2*e^(4*x) - 2) + e^(2*x) + 1)*e^(-2*x)) - 1/2*I*log(-4*(-I*sqrt(-2*e^(4*x) - 2) + e^(2*x) + 1)*e^(-2*x))

Sympy [F]

$$\int \sqrt{-1 - \coth^2(x)} dx = \int \sqrt{-\coth^2(x) - 1} dx$$

[In] integrate((-1-coth(x)**2)**(1/2),x)

[Out] Integral(sqrt(-coth(x)**2 - 1), x)

Maxima [F]

$$\int \sqrt{-1 - \coth^2(x)} dx = \int \sqrt{-\coth(x)^2 - 1} dx$$

[In] integrate((-1-coth(x)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-coth(x)^2 - 1), x)

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 124, normalized size of antiderivative = 2.76

$$\int \sqrt{-1 - \coth^2(x)} dx =$$

$$-\frac{1}{2} \sqrt{2} \left(i \sqrt{2} \log \left(\frac{|-2\sqrt{2} + 2\sqrt{e^{4x} + 1} - 2e^{2x} + 2|}{2(\sqrt{2} + \sqrt{e^{4x} + 1} - e^{2x} + 1)} \right) + i \log(\sqrt{e^{4x} + 1} - e^{2x} + 1) - i \log(\sqrt{e^{4x} + 1} + e^{2x} + 1) \right)$$

[In] integrate((-1-coth(x)^2)^(1/2),x, algorithm="giac")

[Out] -1/2*sqrt(2)*(I*sqrt(2)*log(1/2*abs(-2*sqrt(2) + 2*sqrt(e^(4*x) + 1) - 2*e^(2*x) + 2)/(sqrt(2) + sqrt(e^(4*x) + 1) - e^(2*x) + 1)) + I*log(sqrt(e^(4*x) + 1) - e^(2*x) + 1) - I*log(sqrt(e^(4*x) + 1) + e^(2*x) + 1)) - I*log(-sqrt(e^(4*x) + 1) + e^(2*x) + 1))*sgn(-e^(2*x) + 1)

Mupad [B] (verification not implemented)

Time = 2.01 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.96

$$\int \sqrt{-1 - \coth^2(x)} dx = -\sqrt{2} \operatorname{atan} \left(\frac{\sqrt{2} \coth(x)}{\sqrt{-\coth(x)^2 - 1}} \right) - \ln \left(\coth(x) - \sqrt{-\coth(x)^2 - 1} \right) \operatorname{li}$$

[In] int((- coth(x)^2 - 1)^(1/2),x)

[Out] - log(coth(x) - (- coth(x)^2 - 1)^(1/2)*1i)*1i - 2^(1/2)*atan((2^(1/2)*coth(x))/(- coth(x)^2 - 1)^(1/2))

3.30 $\int (1 + \coth^2(x))^{3/2} dx$

Optimal result	217
Rubi [A] (verified)	217
Mathematica [B] (verified)	219
Maple [B] (verified)	219
Fricas [B] (verification not implemented)	220
Sympy [F]	221
Maxima [F]	221
Giac [B] (verification not implemented)	222
Mupad [B] (verification not implemented)	222

Optimal result

Integrand size = 10, antiderivative size = 50

$$\int (1 + \coth^2(x))^{3/2} dx = -\frac{5}{2} \operatorname{arcsinh}(\coth(x)) + 2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2} \coth(x)}{\sqrt{1 + \coth^2(x)}}\right) - \frac{1}{2} \coth(x) \sqrt{1 + \coth^2(x)}$$

[Out] $-5/2*\operatorname{arcsinh}(\coth(x))+2*\operatorname{arctanh}(\coth(x)*2^{(1/2)}/(1+\coth(x)^2)^{(1/2}))*2^{(1/2)}-1/2*\coth(x)*(1+\coth(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3742, 427, 537, 221, 385, 212}

$$\int (1 + \coth^2(x))^{3/2} dx = -\frac{5}{2} \operatorname{arcsinh}(\coth(x)) + 2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2} \coth(x)}{\sqrt{\coth^2(x) + 1}}\right) - \frac{1}{2} \coth(x) \sqrt{\coth^2(x) + 1}$$

[In] $\operatorname{Int}[(1 + \operatorname{Coth}[x]^2)^{(3/2)}, x]$

[Out] $(-5*\operatorname{ArcSinh}[\operatorname{Coth}[x]])/2 + 2*\operatorname{Sqrt}[2]*\operatorname{ArcTanh}[(\operatorname{Sqrt}[2]*\operatorname{Coth}[x])/\operatorname{Sqrt}[1 + \operatorname{Coth}[x]^2]] - (\operatorname{Coth}[x]*\operatorname{Sqrt}[1 + \operatorname{Coth}[x]^2])/2$

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 221

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt
[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Su
bst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b
, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 427

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[d*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q - 1)/(b*(n*(p + q) + 1))),
x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp
[c*(b*c*(n*(p + q) + 1) - a*d) + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q -
1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d,
0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a
, b, c, d, n, p, q, x]
```

Rule 537

```
Int[((e_) + (f_.)*(x_)^(n_))/(((a_) + (b_.)*(x_)^(n_))*Sqrt[(c_) + (d_.)*(x
_)^(n_)]), x_Symbol] := Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e
- a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d
, e, f, n}, x]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :=
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(
ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\text{integral} = \text{Subst}\left(\int \frac{(1+x^2)^{3/2}}{1-x^2} dx, x, \coth(x)\right)$$

$$\begin{aligned}
&= -\frac{1}{2} \coth(x) \sqrt{1 + \coth^2(x)} - \frac{1}{2} \text{Subst} \left(\int \frac{-3 - 5x^2}{(1 - x^2) \sqrt{1 + x^2}} dx, x, \coth(x) \right) \\
&= -\frac{1}{2} \coth(x) \sqrt{1 + \coth^2(x)} - \frac{5}{2} \text{Subst} \left(\int \frac{1}{\sqrt{1 + x^2}} dx, x, \coth(x) \right) \\
&\quad + 4 \text{Subst} \left(\int \frac{1}{(1 - x^2) \sqrt{1 + x^2}} dx, x, \coth(x) \right) \\
&= -\frac{5}{2} \text{arcsinh}(\coth(x)) - \frac{1}{2} \coth(x) \sqrt{1 + \coth^2(x)} \\
&\quad + 4 \text{Subst} \left(\int \frac{1}{1 - 2x^2} dx, x, \frac{\coth(x)}{\sqrt{1 + \coth^2(x)}} \right) \\
&= -\frac{5}{2} \text{arcsinh}(\coth(x)) + 2\sqrt{2} \text{arctanh} \left(\frac{\sqrt{2} \coth(x)}{\sqrt{1 + \coth^2(x)}} \right) - \frac{1}{2} \coth(x) \sqrt{1 + \coth^2(x)}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 116 vs. $2(50) = 100$.

Time = 0.42 (sec) , antiderivative size = 116, normalized size of antiderivative = 2.32

$$\int (1 + \coth^2(x))^{3/2} dx = -\frac{1}{8} (1 + \coth^2(x))^{3/2} \text{sech}^2(2x) \left(16 \text{arctanh} \left(\frac{\cosh(x)}{\sqrt{\cosh(2x)}} \right) \sqrt{\cosh(2x)} \sinh^3(x) + 4 \left(\arctan \left(\frac{\cosh(x)}{\sqrt{-\cosh(2x)}} \right) \sqrt{-\cosh(2x)} \right) \right)$$

[In] Integrate[(1 + Coth[x]^2)^(3/2), x]

[Out] -1/8*((1 + Coth[x]^2)^(3/2)*Sech[2*x]^2*(16*ArcTanh[Cosh[x]/Sqrt[Cosh[2*x]]]*Sqrt[Cosh[2*x]]*Sinh[x]^3 + 4*(ArcTan[Cosh[x]/Sqrt[-Cosh[2*x]]]*Sqrt[-Cosh[2*x]] - 4*Sqrt[2]*Sqrt[Cosh[2*x]]*Log[Sqrt[2]*Cosh[x] + Sqrt[Cosh[2*x]]])*Sinh[x]^3 + Sinh[4*x]))

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 157 vs. $2(38) = 76$.

Time = 0.10 (sec) , antiderivative size = 158, normalized size of antiderivative = 3.16

method	result
derivativedivides	$\frac{\left((1+\coth(x))^2-2\coth(x)\right)^{\frac{3}{2}}}{6} - \frac{\coth(x)\sqrt{(1+\coth(x))^2-2\coth(x)}}{4} - \frac{5\operatorname{arcsinh}(\coth(x))}{2} + \sqrt{(1+\coth(x))^2}$
default	$\frac{\left((1+\coth(x))^2-2\coth(x)\right)^{\frac{3}{2}}}{6} - \frac{\coth(x)\sqrt{(1+\coth(x))^2-2\coth(x)}}{4} - \frac{5\operatorname{arcsinh}(\coth(x))}{2} + \sqrt{(1+\coth(x))^2}$

```
[In] int((1+coth(x)^2)^(3/2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/6*((1+coth(x))^2-2*coth(x))^(3/2)-1/4*coth(x)*((1+coth(x))^2-2*coth(x))^(1/2)-5/2*arcsinh(coth(x))+((1+coth(x))^2-2*coth(x))^(1/2)-2^(1/2)*arctanh(1/4*(2-2*coth(x))*2^(1/2)/((1+coth(x))^2-2*coth(x))^(1/2))-1/6*((coth(x)-1)^2+2*coth(x))^(3/2)-1/4*coth(x)*((coth(x)-1)^2+2*coth(x))^(1/2)-((coth(x)-1)^2+2*coth(x))^(1/2)+2^(1/2)*arctanh(1/4*(2+2*coth(x))*2^(1/2)/((coth(x)-1)^2+2*coth(x))^(1/2)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1043 vs. 2(38) = 76.

Time = 0.28 (sec) , antiderivative size = 1043, normalized size of antiderivative = 20.86

$$\int (1 + \coth^2(x))^{3/2} dx = \text{Too large to display}$$

```
[In] integrate((1+coth(x)^2)^(3/2),x, algorithm="fricas")
```

```
[Out] 1/4*(2*(sqrt(2)*cosh(x)^4 + 4*sqrt(2)*cosh(x)*sinh(x)^3 + sqrt(2)*sinh(x)^4 + 2*(3*sqrt(2)*cosh(x)^2 - sqrt(2))*sinh(x)^2 - 2*sqrt(2)*cosh(x)^2 + 4*(sqrt(2)*cosh(x)^3 - sqrt(2)*cosh(x))*sinh(x) + sqrt(2))*log(2*(cosh(x)^8 + 8*cosh(x)*sinh(x)^7 + sinh(x)^8 + (28*cosh(x)^2 + 3)*sinh(x)^6 + 3*cosh(x)^6 + 2*(28*cosh(x)^3 + 9*cosh(x))*sinh(x)^5 + 5*(14*cosh(x)^4 + 9*cosh(x)^2 + 1)*sinh(x)^4 + 5*cosh(x)^4 + 4*(14*cosh(x)^5 + 15*cosh(x)^3 + 5*cosh(x))*sinh(x)^3 + (28*cosh(x)^6 + 45*cosh(x)^4 + 30*cosh(x)^2 + 4)*sinh(x)^2 + 4*cosh(x)^2 + 2*(4*cosh(x)^7 + 9*cosh(x)^5 + 10*cosh(x)^3 + 4*cosh(x))*sinh(x) + (sqrt(2)*cosh(x)^6 + 6*sqrt(2)*cosh(x)*sinh(x)^5 + sqrt(2)*sinh(x)^6 + 3*(5*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x)^4 + 3*sqrt(2)*cosh(x)^4 + 4*(5*sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x))*sinh(x)^3 + (15*sqrt(2)*cosh(x)^4 + 18*sqrt(2)*cosh(x)^2 + 4*sqrt(2))*sinh(x)^2 + 4*sqrt(2)*cosh(x)^2 + 2*(3*sqrt(2)*cosh(x)^5 + 6*sqrt(2)*cosh(x)^3 + 4*sqrt(2)*cosh(x))*sinh(x) + 4*sqrt(2))*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4)/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6)) + 2*(sqrt(2)*cosh(x)^4 + 4*sqrt(2)*cosh(x)*sinh(x)^3 + sqrt(2)*sinh(x)^4 + 2*(3*sqrt(2)*cosh(x)^2 - sqrt(2))*sinh(x)^2 - 2*sqrt(2)*cosh(x)^2 + 4*(sqrt
```

```
(2)*cosh(x)^3 - sqrt(2)*cosh(x)*sinh(x) + sqrt(2))*log(-2*(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 - 1)*sinh(x)^2 - cosh(x)^2 + 2*(2*cosh(x)^3 - cosh(x))*sinh(x) + (sqrt(2)*cosh(x)^2 + 2*sqrt(2)*cosh(x)*sinh(x) + sqrt(2)*sinh(x)^2 - sqrt(2))*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 1)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)) - 5*(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*sinh(x)^2 - 2*cosh(x)^2 + 4*(cosh(x)^3 - cosh(x))*sinh(x) + 1)*log((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 2*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 1)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)) + 5*(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*sinh(x)^2 - 2*cosh(x)^2 + 4*(cosh(x)^3 - cosh(x))*sinh(x) + 1)*log((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 2*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 1)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)) - 4*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*sinh(x)^2 - 2*cosh(x)^2 + 4*(cosh(x)^3 - cosh(x))*sinh(x) + 1)
```

Sympy [F]

$$\int (1 + \coth^2(x))^{3/2} dx = \int (\coth^2(x) + 1)^{\frac{3}{2}} dx$$

```
[In] integrate((1+coth(x)**2)**(3/2),x)
```

```
[Out] Integral((coth(x)**2 + 1)**(3/2), x)
```

Maxima [F]

$$\int (1 + \coth^2(x))^{3/2} dx = \int (\coth(x)^2 + 1)^{\frac{3}{2}} dx$$

```
[In] integrate((1+coth(x)^2)^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((coth(x)^2 + 1)^(3/2), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 265 vs. 2(38) = 76.

Time = 0.29 (sec) , antiderivative size = 265, normalized size of antiderivative = 5.30

$$\int (1 + \coth^2(x))^{3/2} dx = \frac{1}{4} \sqrt{2} \left(5 \sqrt{2} \log \left(\frac{|-2\sqrt{2} + 2\sqrt{e^{4x} + 1} - 2e^{2x} + 2|}{2(\sqrt{2} + \sqrt{e^{4x} + 1} - e^{2x} + 1)} \right) \operatorname{sgn}(e^{2x} - 1) + 4 \log \left(\sqrt{e^{4x} + 1} \right) \right)$$

[In] integrate((1+coth(x)^2)^(3/2),x, algorithm="giac")

[Out] 1/4*sqrt(2)*(5*sqrt(2)*log(1/2*abs(-2*sqrt(2) + 2*sqrt(e^(4*x) + 1) - 2*e^(2*x) + 2)/(sqrt(2) + sqrt(e^(4*x) + 1) - e^(2*x) + 1))*sgn(e^(2*x) - 1) + 4*log(sqrt(e^(4*x) + 1) - e^(2*x) + 1)*sgn(e^(2*x) - 1) - 4*log(sqrt(e^(4*x) + 1) - e^(2*x))*sgn(e^(2*x) - 1) - 4*log(-sqrt(e^(4*x) + 1) + e^(2*x) + 1)*sgn(e^(2*x) - 1) - 4*(3*(sqrt(e^(4*x) + 1) - e^(2*x))^3*sgn(e^(2*x) - 1) + (sqrt(e^(4*x) + 1) - e^(2*x))^2*sgn(e^(2*x) - 1) - (sqrt(e^(4*x) + 1) - e^(2*x))*sgn(e^(2*x) - 1) + sgn(e^(2*x) - 1))/((sqrt(e^(4*x) + 1) - e^(2*x))^2 + 2*sqrt(e^(4*x) + 1) - 2*e^(2*x) - 1)^2)

Mupad [B] (verification not implemented)

Time = 2.12 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.56

$$\int (1 + \coth^2(x))^{3/2} dx = \sqrt{2} \left(\ln \left(\coth(x) + \sqrt{2} \sqrt{\coth(x)^2 + 1} + 1 \right) - \ln(\coth(x) - 1) \right) - \frac{\coth(x) \sqrt{\coth(x)^2 + 1}}{2} - \frac{5 \operatorname{asinh}(\coth(x))}{2} + \sqrt{2} \left(\ln(\coth(x) + 1) - \ln \left(\sqrt{2} \sqrt{\coth(x)^2 + 1} - \coth(x) + 1 \right) \right)$$

[In] int((coth(x)^2 + 1)^(3/2),x)

[Out] 2^(1/2)*(log(coth(x) + 2^(1/2)*(coth(x)^2 + 1)^(1/2) + 1) - log(coth(x) - 1)) - (coth(x)*(coth(x)^2 + 1)^(1/2))/2 - (5*asinh(coth(x)))/2 + 2^(1/2)*(log(coth(x) + 1) - log(2^(1/2)*(coth(x)^2 + 1)^(1/2) - coth(x) + 1))

3.31 $\int (-1 - \coth^2(x))^{3/2} dx$

Optimal result	223
Rubi [A] (verified)	223
Mathematica [A] (verified)	225
Maple [B] (verified)	226
Fricas [C] (verification not implemented)	226
Sympy [F]	227
Maxima [F]	227
Giac [C] (verification not implemented)	227
Mupad [F(-1)]	228

Optimal result

Integrand size = 12, antiderivative size = 67

$$\int (-1 - \coth^2(x))^{3/2} dx = -\frac{5}{2} \arctan\left(\frac{\coth(x)}{\sqrt{-1 - \coth^2(x)}}\right) + 2\sqrt{2} \arctan\left(\frac{\sqrt{2} \coth(x)}{\sqrt{-1 - \coth^2(x)}}\right) + \frac{1}{2} \coth(x) \sqrt{-1 - \coth^2(x)}$$

[Out] $-5/2*\arctan(\coth(x)/(-1-\coth(x)^2)^{(1/2)})+2*\arctan(\coth(x)*2^{(1/2)/(-1-\coth(x)^2)^{(1/2)})*2^{(1/2)}+1/2*\coth(x)*(-1-\coth(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3742, 427, 537, 223, 209, 385}

$$\int (-1 - \coth^2(x))^{3/2} dx = -\frac{5}{2} \arctan\left(\frac{\coth(x)}{\sqrt{-\coth^2(x) - 1}}\right) + 2\sqrt{2} \arctan\left(\frac{\sqrt{2} \coth(x)}{\sqrt{-\coth^2(x) - 1}}\right) + \frac{1}{2} \coth(x) \sqrt{-\coth^2(x) - 1}$$

[In] $\text{Int}[(-1 - \text{Coth}[x]^2)^{(3/2)}, x]$

[Out] $(-5*\text{ArcTan}[\text{Coth}[x]/\text{Sqrt}[-1 - \text{Coth}[x]^2]])/2 + 2*\text{Sqrt}[2]*\text{ArcTan}[(\text{Sqrt}[2]*\text{Coth}[x])/\text{Sqrt}[-1 - \text{Coth}[x]^2]] + (\text{Coth}[x]*\text{Sqrt}[-1 - \text{Coth}[x]^2])/2$

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 223

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 427

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[d*x*(a + b*x^n)^(p+1)*((c + d*x^n)^(q-1)/(b*(n*(p+q)+1))), x] + Dist[1/(b*(n*(p+q)+1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q-2)*Simp[c*(b*c*(n*(p+q)+1) - a*d) + d*(b*c*(n*(p+2*q-1)+1) - a*d*(n*(q-1)+1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[n*(p+q)+1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]
```

Rule 537

```
Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*Sqrt[(c_) + (d_.)*(x_)^(n_)], x_Symbol] := Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])
```

Rubi steps

$$\text{integral} = \text{Subst}\left(\int \frac{(-1-x^2)^{3/2}}{1-x^2} dx, x, \text{coth}(x)\right)$$

$$\begin{aligned}
&= \frac{1}{2} \coth(x) \sqrt{-1 - \coth^2(x)} - \frac{1}{2} \text{Subst} \left(\int \frac{-3 - 5x^2}{\sqrt{-1 - x^2} (1 - x^2)} dx, x, \coth(x) \right) \\
&= \frac{1}{2} \coth(x) \sqrt{-1 - \coth^2(x)} - \frac{5}{2} \text{Subst} \left(\int \frac{1}{\sqrt{-1 - x^2}} dx, x, \coth(x) \right) \\
&\quad + 4 \text{Subst} \left(\int \frac{1}{\sqrt{-1 - x^2} (1 - x^2)} dx, x, \coth(x) \right) \\
&= \frac{1}{2} \coth(x) \sqrt{-1 - \coth^2(x)} - \frac{5}{2} \text{Subst} \left(\int \frac{1}{1 + x^2} dx, x, \frac{\coth(x)}{\sqrt{-1 - \coth^2(x)}} \right) \\
&\quad + 4 \text{Subst} \left(\int \frac{1}{1 + 2x^2} dx, x, \frac{\coth(x)}{\sqrt{-1 - \coth^2(x)}} \right) \\
&= -\frac{5}{2} \arctan \left(\frac{\coth(x)}{\sqrt{-1 - \coth^2(x)}} \right) \\
&\quad + 2\sqrt{2} \arctan \left(\frac{\sqrt{2} \coth(x)}{\sqrt{-1 - \coth^2(x)}} \right) + \frac{1}{2} \coth(x) \sqrt{-1 - \coth^2(x)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.76

$$\int (-1 - \coth^2(x))^{3/2} dx = -\frac{1}{8}(-1 - \coth^2(x))^{3/2} \text{sech}^2(2x) \left(16 \text{arctanh} \left(\frac{\cosh(x)}{\sqrt{\cosh(2x)}} \right) \sqrt{\cosh(2x)} \sinh^3(x) + 4 \left(\arctan \left(\frac{\cosh(x)}{\sqrt{-\cosh(2x)}} \right) \sqrt{-\cosh(2x)} \right) \right)$$

[In] Integrate[(-1 - Coth[x]^2)^(3/2), x]

[Out] -1/8*((-1 - Coth[x]^2)^(3/2)*Sech[2*x]^2*(16*ArcTanh[Cosh[x]/Sqrt[Cosh[2*x]]]*Sqrt[Cosh[2*x]]*Sinh[x]^3 + 4*(ArcTan[Cosh[x]/Sqrt[-Cosh[2*x]]]*Sqrt[-Cosh[2*x]] - 4*Sqrt[2]*Sqrt[Cosh[2*x]]*Log[Sqrt[2]*Cosh[x] + Sqrt[Cosh[2*x]]])*Sinh[x]^3 + Sinh[4*x]))

$$+ 2) - \sqrt{-2} * e^{(4*x)} - \sqrt{-2} * e^{(2*x)} - 2 * \sqrt{-2}) * e^{(-4*x)} + 2 * \sqrt{-2} * e^{(4*x)} - 2) * (e^{(2*x)} + 1)) / (e^{(4*x)} - 2 * e^{(2*x)} + 1)$$

Sympy [F]

$$\int (-1 - \coth^2(x))^{3/2} dx = \int (-\coth^2(x) - 1)^{\frac{3}{2}} dx$$

[In] integrate((-1-coth(x)**2)**(3/2),x)

[Out] Integral((-coth(x)**2 - 1)**(3/2), x)

Maxima [F]

$$\int (-1 - \coth^2(x))^{3/2} dx = \int (-\coth(x)^2 - 1)^{\frac{3}{2}} dx$$

[In] integrate((-1-coth(x)^2)^(3/2),x, algorithm="maxima")

[Out] integrate((-coth(x)^2 - 1)^(3/2), x)

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 285, normalized size of antiderivative = 4.25

$$\int (-1 - \coth^2(x))^{3/2} dx = -\frac{1}{4} \sqrt{2} \left(-5i \sqrt{2} \log \left(\frac{|-2\sqrt{2} + 2\sqrt{e^{(4x)} + 1} - 2e^{(2x)} + 2|}{2(\sqrt{2} + \sqrt{e^{(4x)} + 1} - e^{(2x)} + 1)} \right) \operatorname{sgn}(-e^{(2x)} + 1) - 4i \log \left(\sqrt{e^{(4x)} + 1} - e^{(2x)} \right) \right)$$

[In] integrate((-1-coth(x)^2)^(3/2),x, algorithm="giac")

[Out]
$$-1/4 * \sqrt{2} * (-5 * I * \sqrt{2} * \log(1/2 * \operatorname{abs}(-2 * \sqrt{2} + 2 * \sqrt{e^{(4*x)} + 1} - 2 * e^{(2*x)} + 2) / (\sqrt{2} + \sqrt{e^{(4*x)} + 1} - e^{(2*x)} + 1)) * \operatorname{sgn}(-e^{(2*x)} + 1) - 4 * I * \log(\sqrt{e^{(4*x)} + 1} - e^{(2*x)}) * \operatorname{sgn}(-e^{(2*x)} + 1) + 4 * I * \log(\sqrt{e^{(4*x)} + 1} - e^{(2*x)}) * \operatorname{sgn}(-e^{(2*x)} + 1) + 4 * I * \log(-\sqrt{e^{(4*x)} + 1} - e^{(2*x)}) * \operatorname{sgn}(-e^{(2*x)} + 1) + 4 * (3 * I * (\sqrt{e^{(4*x)} + 1} - e^{(2*x)})^3 * \operatorname{sgn}(-e^{(2*x)} + 1) + I * (\sqrt{e^{(4*x)} + 1} - e^{(2*x)})^2 * \operatorname{sgn}(-e^{(2*x)} + 1) + (-I * \sqrt{e^{(4*x)} + 1} + I * e^{(2*x)}) * \operatorname{sgn}(-e^{(2*x)} + 1) + I * \operatorname{sgn}(-e^{(2*x)} + 1)) / ((\sqrt{e^{(4*x)} + 1} - e^{(2*x)})^2 + 2 * \sqrt{e^{(4*x)} + 1} - 2 * e^{(2*x)} - 1)^2)$$

Mupad [F(-1)]

Timed out.

$$\int (-1 - \coth^2(x))^{3/2} dx = \int (-\coth(x)^2 - 1)^{3/2} dx$$

```
[In] int((- coth(x)^2 - 1)^(3/2),x)
```

```
[Out] int((- coth(x)^2 - 1)^(3/2), x)
```

$$3.32 \quad \int \frac{\coth^3(x)}{\sqrt{a+b \coth^2(x)}} dx$$

Optimal result	229
Rubi [A] (verified)	229
Mathematica [A] (verified)	231
Maple [B] (verified)	231
Fricas [B] (verification not implemented)	232
Sympy [F]	233
Maxima [F]	233
Giac [F(-2)]	233
Mupad [B] (verification not implemented)	234

Optimal result

Integrand size = 17, antiderivative size = 47

$$\int \frac{\coth^3(x)}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}} - \frac{\sqrt{a+b \coth^2(x)}}{b}$$

[Out] $\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2)/(a+b)^{(1/2)})}/(a+b)^{(1/2)}-(a+b*\coth(x)^2)^{(1/2)}/b$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3751, 457, 81, 65, 214}

$$\int \frac{\coth^3(x)}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}} - \frac{\sqrt{a+b \coth^2(x)}}{b}$$

[In] $\operatorname{Int}[\operatorname{Coth}[x]^3/\operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2], x]$

[Out] $\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2]/\operatorname{Sqrt}[a + b]]/\operatorname{Sqrt}[a + b] - \operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2]/b$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] \rightarrow \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) +$

$d*(x^p/b)^n, x, (a + b*x)^{(1/p)}, x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 81

$\text{Int}[(a_.) + (b_.)*(x_.)]*((c_.) + (d_.)*(x_.)^{(n_.)})*((e_.) + (f_.)*(x_.)^{(p_.)}), x_Symbol] \rightarrow \text{Simp}[b*(c + d*x)^{(n + 1)}*(e + f*x)^{(p + 1)}/(d*f*(n + p + 2)), x] + \text{Dist}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1))]/(d*f*(n + p + 2)), \text{Int}[(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \&\& \text{NeQ}[n + p + 2, 0]$

Rule 214

$\text{Int}[(a_.) + (b_.)*(x_.)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{NegQ}[a/b]$

Rule 457

$\text{Int}(x_.)^{(m_.)}*((a_.) + (b_.)*(x_.)^{(n_.)})^{(p_.)}*((c_.) + (d_.)*(x_.)^{(n_.)})^{(q_.)}), x_Symbol] \rightarrow \text{Dist}[1/n, \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)}*(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; \text{FreeQ}\{a, b, c, d, m, n, p, q\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]$

Rule 3751

$\text{Int}[(d_.)*\tan[(e_.) + (f_.)*(x_.)])^{(m_.)}*((a_.) + (b_.)*((c_.)*\tan[(e_.) + (f_.)*(x_.)])^{(n_.)})^{(p_.)}), x_Symbol] \rightarrow \text{With}\{\text{ff} = \text{FreeFactors}[\text{Tan}[e + f*x], x]\}, \text{Dist}[c*(\text{ff}/f), \text{Subst}[\text{Int}[(d*\text{ff}*(x/c))^{(m_*)}*((a + b*(\text{ff}*x)^n)^p/(c^2 + \text{ff}^2*x^2)], x], x, c*(\text{Tan}[e + f*x]/\text{ff})], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x] \&\& (\text{IGtQ}[p, 0] \|\| \text{EqQ}[n, 2] \|\| \text{EqQ}[n, 4] \|\| (\text{IntegerQ}[p] \&\& \text{RationalQ}[n]))$

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst}\left(\int \frac{x^3}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right) \\ &= \frac{1}{2}\text{Subst}\left(\int \frac{x}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x)\right) \\ &= -\frac{\sqrt{a+b\coth^2(x)}}{b} + \frac{1}{2}\text{Subst}\left(\int \frac{1}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x)\right) \\ &= -\frac{\sqrt{a+b\coth^2(x)}}{b} + \frac{\text{Subst}\left(\int \frac{1}{1+\frac{a}{b}-\frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)}\right)}{b} \end{aligned}$$

$$= \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}} - \frac{\sqrt{a+b\coth^2(x)}}{b}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00

$$\int \frac{\coth^3(x)}{\sqrt{a+b\coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}} - \frac{\sqrt{a+b\coth^2(x)}}{b}$$

[In] Integrate[Coth[x]^3/Sqrt[a + b*Coth[x]^2], x]

[Out] ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]]/Sqrt[a + b] - Sqrt[a + b*Coth[x]^2]/b

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 128 vs. 2(39) = 78.

Time = 0.12 (sec) , antiderivative size = 129, normalized size of antiderivative = 2.74

method	result
derivativedivides	$-\frac{\sqrt{a+b\coth(x)^2}}{b} + \frac{\ln\left(\frac{2a+2b+2b(\coth(x)-1)+2\sqrt{a+b}\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}{\coth(x)-1}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))}{\coth(x)+1}\right)}{2\sqrt{a+b}}$
default	$-\frac{\sqrt{a+b\coth(x)^2}}{b} + \frac{\ln\left(\frac{2a+2b+2b(\coth(x)-1)+2\sqrt{a+b}\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}{\coth(x)-1}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))}{\coth(x)+1}\right)}{2\sqrt{a+b}}$

[In] int(coth(x)^3/(a+b*coth(x)^2)^(1/2), x, method=_RETURNVERBOSE)

[Out] $-(a+b\coth(x)^2)^{1/2}/b+1/2/(a+b)^{1/2}*\ln((2*a+2*b+2*b*(\coth(x)-1)+2*(a+b)^{1/2}*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{1/2})/(\coth(x)-1))+1/2/(a+b)^{1/2}*\ln((2*a+2*b-2*b*(1+\coth(x))+2*(a+b)^{1/2}*(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{1/2})/(1+\coth(x)))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 472 vs. 2(39) = 78.

Time = 0.31 (sec) , antiderivative size = 1576, normalized size of antiderivative = 33.53

$$\int \frac{\coth^3(x)}{\sqrt{a + b \coth^2(x)}} dx = \text{Too large to display}$$

[In] integrate(coth(x)^3/(a+b*coth(x)^2)^(1/2),x, algorithm="fricas")

[Out] [1/4*((b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 - b)*sqrt(a + b)*log(-(a^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^4 + (70*(a^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*cosh(x)^2 + 2*(14*(a^3 + a^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*cosh(x)^4 - 2*a^3 - 3*a^2*b + b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(a^2*cosh(x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*sinh(x)^6 - 3*a^2*cosh(x)^4 + 3*(5*a^2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^2*cosh(x)^3 - 3*a^2*cosh(x))*sinh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 + (15*a^2*cosh(x)^4 - 18*a^2*cosh(x)^2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*cosh(x)^5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a^3 + a^2*b)*cosh(x)^7 - 3*(2*a^3 + a^2*b)*cosh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^3 - (2*a^3 + 3*a^2*b - b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6) + (b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 - b)*sqrt(a + b)*log(((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 + 2*b*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 + b)*sinh(x)^2 + sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*((a + b)*cosh(x)^3 + b*cosh(x))*sinh(x) + a + b)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)) - 4*sqrt(2)*(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a*b + b^2)*cosh(x)^2 + 2*(a*b + b^2)*cosh(x)*sinh(x) + (a*b + b^2)*sinh(x)^2 - a*b - b^2), -1/2*((b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 - b)*sqrt(-a - b)*arctan(sqrt(2)*(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 - a - b)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a^2 + a*b)*cosh(x)^4 + 4*(a^2 + a


```
*b)*cosh(x)*sinh(x)^3 + (a^2 + a*b)*sinh(x)^4 - (2*a^2 + a*b - b^2)*cosh(x)
^2 + (6*(a^2 + a*b)*cosh(x)^2 - 2*a^2 - a*b + b^2)*sinh(x)^2 + a^2 + 2*a*b
+ b^2 + 2*(2*(a^2 + a*b)*cosh(x)^3 - (2*a^2 + a*b - b^2)*cosh(x))*sinh(x))
+ (b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 - b)*sqrt(-a - b)*arcta
n(sqrt(2)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)
/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2))/((a + b)*cosh(x)^2 + 2*(a + b
)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a + b)) + 2*sqrt(2)*(a + b)*sqrt(((
a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x)
) + sinh(x)^2)))/((a*b + b^2)*cosh(x)^2 + 2*(a*b + b^2)*cosh(x)*sinh(x) + (
a*b + b^2)*sinh(x)^2 - a*b - b^2)]
```

Sympy [F]

$$\int \frac{\coth^3(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\coth^3(x)}{\sqrt{a + b \coth^2(x)}} dx$$

```
[In] integrate(coth(x)**3/(a+b*coth(x)**2)**(1/2),x)
```

```
[Out] Integral(coth(x)**3/sqrt(a + b*coth(x)**2), x)
```

Maxima [F]

$$\int \frac{\coth^3(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\coth(x)^3}{\sqrt{b \coth(x)^2 + a}} dx$$

```
[In] integrate(coth(x)^3/(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(coth(x)^3/sqrt(b*coth(x)^2 + a), x)
```

Giac [F(-2)]

Exception generated.

$$\int \frac{\coth^3(x)}{\sqrt{a + b \coth^2(x)}} dx = \text{Exception raised: TypeError}$$

```
[In] integrate(coth(x)^3/(a+b*coth(x)^2)^(1/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type
```

Mupad [B] (verification not implemented)

Time = 2.39 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.83

$$\int \frac{\coth^3(x)}{\sqrt{a + b \coth^2(x)}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \coth(x)^2 + a}}{\sqrt{a+b}}\right)}{\sqrt{a+b}} - \frac{\sqrt{b \coth(x)^2 + a}}{b}$$

[In] `int(coth(x)^3/(a + b*coth(x)^2)^(1/2),x)`

[Out] `atanh((a + b*coth(x)^2)^(1/2)/(a + b)^(1/2))/(a + b)^(1/2) - (a + b*coth(x)^2)^(1/2)/b`

3.33 $\int \frac{\coth^2(x)}{\sqrt{a+b \coth^2(x)}} dx$

Optimal result	235
Rubi [A] (verified)	235
Mathematica [B] (verified)	237
Maple [B] (verified)	237
Fricas [B] (verification not implemented)	238
Sympy [F]	240
Maxima [F]	240
Giac [F(-2)]	241
Mupad [F(-1)]	241

Optimal result

Integrand size = 17, antiderivative size = 60

$$\int \frac{\coth^2(x)}{\sqrt{a+b \coth^2(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{\sqrt{b}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{\sqrt{a+b}}$$

[Out] $-\operatorname{arctanh}(\coth(x)*b^{(1/2)/(a+b*\coth(x)^2)^{(1/2)})/b^{(1/2)}+\operatorname{arctanh}(\coth(x)*(a+b)^{(1/2)/(a+b*\coth(x)^2)^{(1/2)})/(a+b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3751, 494, 223, 212, 385}

$$\int \frac{\coth^2(x)}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{\sqrt{a+b}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{\sqrt{b}}$$

[In] $\text{Int}[\text{Coth}[x]^2/\text{Sqrt}[a + b*\text{Coth}[x]^2], x]$

[Out] $-(\text{ArcTanh}[(\text{Sqrt}[b]*\text{Coth}[x])/(\text{Sqrt}[a + b*\text{Coth}[x]^2])]/\text{Sqrt}[b]) + \text{ArcTanh}[(\text{Sqrt}[a + b]*\text{Coth}[x])/(\text{Sqrt}[a + b*\text{Coth}[x]^2])]/\text{Sqrt}[a + b]$

Rule 212

$\text{Int}[(a_1 + (b_1*x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{Gt}$

Q[a, 0] || LtQ[b, 0])

Rule 223

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 494

Int[(((e_.)*(x_))^(m_)*((c_) + (d_.)*(x_)^(n_))^(q_.))/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Dist[e^n/b, Int[(e*x)^(m - n)*(c + d*x^n)^q, x], x] - Dist[a*(e^n/b), Int[(e*x)^(m - n)*((c + d*x^n)^q/(a + b*x^n)), x], x] /; FreeQ[{a, b, c, d, e, m, q}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LeQ[n, m, 2*n - 1] && IntBinomialQ[a, b, c, d, e, m, n, -1, q, x]

Rule 3751

Int[(((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst} \left(\int \frac{x^2}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x) \right) \\
 &= -\text{Subst} \left(\int \frac{1}{\sqrt{a+bx^2}} dx, x, \coth(x) \right) + \text{Subst} \left(\int \frac{1}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x) \right) \\
 &= -\text{Subst} \left(\int \frac{1}{1-bx^2} dx, x, \frac{\coth(x)}{\sqrt{a+b\coth^2(x)}} \right) \\
 &\quad + \text{Subst} \left(\int \frac{1}{1-(a+b)x^2} dx, x, \frac{\coth(x)}{\sqrt{a+b\coth^2(x)}} \right)
 \end{aligned}$$

$$= -\frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\coth(x)}{\sqrt{a+b\coth^2(x)}}\right)}{\sqrt{b}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b}\coth(x)}{\sqrt{a+b\coth^2(x)}}\right)}{\sqrt{a+b}}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 134 vs. $2(60) = 120$.

Time = 0.24 (sec) , antiderivative size = 134, normalized size of antiderivative = 2.23

$$\int \frac{\coth^2(x)}{\sqrt{a+b\coth^2(x)}} dx$$

$$= \frac{\left(-\sqrt{a+b}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{b}\cosh(x)}{\sqrt{-a+b+(a+b)\cosh(2x)}}\right) + \sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{a+b}\cosh(x)}{\sqrt{-a+b+(a+b)\cosh(2x)}}\right)\right) \sqrt{-a+b+(a+b)\cosh(2x)}}{\sqrt{b}\sqrt{a+b}\sqrt{-a+b+(a+b)\cosh(2x)}}$$

[In] Integrate[Coth[x]^2/Sqrt[a + b*Coth[x]^2], x]

[Out] ((-(Sqrt[a + b]*ArcTanh[(Sqrt[2]*Sqrt[b]*Cosh[x])/Sqrt[-a + b + (a + b)*Cosh[2*x]]) + Sqrt[b]*ArcTanh[(Sqrt[2]*Sqrt[a + b]*Cosh[x])/Sqrt[-a + b + (a + b)*Cosh[2*x]])*Sqrt[(-a + b + (a + b)*Cosh[2*x])*Csch[x]^2]*Sinh[x])/(Sqrt[b]*Sqrt[a + b]*Sqrt[-a + b + (a + b)*Cosh[2*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 136 vs. $2(48) = 96$.

Time = 0.12 (sec) , antiderivative size = 137, normalized size of antiderivative = 2.28

method	result
derivativedivides	$-\frac{\ln\left(\sqrt{b}\coth(x)+\sqrt{a+b\coth^2(x)}\right)}{\sqrt{b}} - \frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))+2\sqrt{a+b}\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{1+\coth(x)}}{2\sqrt{a+b}}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))+2\sqrt{a+b}\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{1+\coth(x)}}{2\sqrt{a+b}}\right)}{2\sqrt{a+b}}$
default	$-\frac{\ln\left(\sqrt{b}\coth(x)+\sqrt{a+b\coth^2(x)}\right)}{\sqrt{b}} - \frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))+2\sqrt{a+b}\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{1+\coth(x)}}{2\sqrt{a+b}}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))+2\sqrt{a+b}\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{1+\coth(x)}}{2\sqrt{a+b}}\right)}{2\sqrt{a+b}}$

[In] int(coth(x)^2/(a+b*coth(x)^2)^(1/2), x, method=_RETURNVERBOSE)

[Out] $-\ln(b^{1/2}\coth(x)+(a+b\coth(x)^2)^{1/2})/b^{1/2}-1/2/(a+b)^{1/2}\ln((2*a+2*b-2*b*(1+\coth(x))+2*(a+b)^{1/2}*(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{1/2})/(1+\coth(x)))+1/2/(a+b)^{1/2}\ln((2*a+2*b+2*b*(\coth(x)-1)+2*(a+b)^{1/2}*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{1/2})/(\coth(x)-1))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 567 vs. 2(48) = 96.

Time = 0.39 (sec) , antiderivative size = 3513, normalized size of antiderivative = 58.55

$$\int \frac{\coth^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \text{Too large to display}$$

[In] integrate(coth(x)^2/(a+b*coth(x)^2)^(1/2),x, algorithm="fricas")

[Out] [1/4*(sqrt(a + b)*b*log(((a*b^2 + b^3)*cosh(x)^8 + 8*(a*b^2 + b^3)*cosh(x)*sinh(x)^7 + (a*b^2 + b^3)*sinh(x)^8 + 2*(a*b^2 + 2*b^3)*cosh(x)^6 + 2*(a*b^2 + 2*b^3 + 14*(a*b^2 + b^3)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a*b^2 + b^3)*cosh(x)^3 + 3*(a*b^2 + 2*b^3)*cosh(x))*sinh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^4 + (70*(a*b^2 + b^3)*cosh(x)^4 + a^3 - a^2*b + 4*a*b^2 + 6*b^3 + 30*(a*b^2 + 2*b^3)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a*b^2 + b^3)*cosh(x)^5 + 10*(a*b^2 + 2*b^3)*cosh(x)^3 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 - 3*a*b^2 - 2*b^3)*cosh(x)^2 + 2*(14*(a*b^2 + b^3)*cosh(x)^6 + 15*(a*b^2 + 2*b^3)*cosh(x)^4 - a^3 + 3*a*b^2 + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(b^2*cosh(x)^6 + 6*b^2*cosh(x)*sinh(x)^5 + b^2*sinh(x)^6 + 3*b^2*cosh(x)^4 + 3*(5*b^2*cosh(x)^2 + b^2)*sinh(x)^4 + 4*(5*b^2*cosh(x)^3 + 3*b^2*cosh(x))*sinh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x)^2 + (15*b^2*cosh(x)^4 + 18*b^2*cosh(x)^2 - a^2 + 2*a*b + 3*b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(3*b^2*cosh(x)^5 + 6*b^2*cosh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a*b^2 + b^3)*cosh(x)^7 + 3*(a*b^2 + 2*b^3)*cosh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^3 - (a^3 - 3*a*b^2 - 2*b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6)) + 2*(a + b)*sqrt(b)*log(-((a + 2*b)*cosh(x)^4 + 4*(a + 2*b)*cosh(x)*sinh(x)^3 + (a + 2*b)*sinh(x)^4 - 2*(a - 2*b)*cosh(x)^2 + 2*(3*(a + 2*b)*cosh(x)^2 - a + 2*b)*sinh(x)^2 - 2*sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*((a + 2*b)*cosh(x)^3 - (a - 2*b)*cosh(x))*sinh(x) + a + 2*b)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*sinh(x)^2 - 2*cosh(x)^2 + 4*(cosh(x)^3 - cosh(x))*sinh(x) + 1)) + sqrt(a + b)*b*log(-((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*a*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a)*sinh(x)^2 + sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*((a + b)*cosh(x)^3 - a*cosh(x))*sinh(x) + a + b)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)))/(a*b + b^2), 1/4*(4*(a + b)*sqrt(-b)*arctan(sqrt(2)*(cosh(x)^2 + 2*cosh(x)

$$\begin{aligned}
&) * \sinh(x) + \sinh(x)^2 + 1) * \sqrt{-b} * \sqrt{((a + b) * \cosh(x)^2 + (a + b) * \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 * \cosh(x) * \sinh(x) + \sinh(x)^2))} / ((a + b) * \cosh(x)^4 + 4 * (a + b) * \cosh(x) * \sinh(x)^3 + (a + b) * \sinh(x)^4 - 2 * (a - b) * \cosh(x)^2 + 2 * (3 * (a + b) * \cosh(x)^2 - a + b) * \sinh(x)^2 + 4 * ((a + b) * \cosh(x)^3 - (a - b) * \cosh(x)) * \sinh(x) + a + b)) + \sqrt{a + b} * b * \log(((a * b^2 + b^3) * \cosh(x)^8 + 8 * (a * b^2 + b^3) * \cosh(x) * \sinh(x)^7 + (a * b^2 + b^3) * \sinh(x)^8 + 2 * (a * b^2 + 2 * b^3) * \cosh(x)^6 + 2 * (a * b^2 + 2 * b^3 + 14 * (a * b^2 + b^3) * \cosh(x)^2) * \sinh(x)^6 + 4 * (14 * (a * b^2 + b^3) * \cosh(x)^3 + 3 * (a * b^2 + 2 * b^3) * \cosh(x)) * \sinh(x)^5 + (a^3 - a^2 * b + 4 * a * b^2 + 6 * b^3) * \cosh(x)^4 + (70 * (a * b^2 + b^3) * \cosh(x)^4 + a^3 - a^2 * b + 4 * a * b^2 + 6 * b^3 + 30 * (a * b^2 + 2 * b^3) * \cosh(x)^2) * \sinh(x)^4 + 4 * (14 * (a * b^2 + b^3) * \cosh(x)^5 + 10 * (a * b^2 + 2 * b^3) * \cosh(x)^3 + (a^3 - a^2 * b + 4 * a * b^2 + 6 * b^3) * \cosh(x)) * \sinh(x)^3 + a^3 + 3 * a^2 * b + 3 * a * b^2 + b^3 - 2 * (a^3 - 3 * a * b^2 - 2 * b^3) * \cosh(x)^2 + 2 * (14 * (a * b^2 + b^3) * \cosh(x)^6 + 15 * (a * b^2 + 2 * b^3) * \cosh(x)^4 - a^3 + 3 * a * b^2 + 2 * b^3 + 3 * (a^3 - a^2 * b + 4 * a * b^2 + 6 * b^3) * \cosh(x)^2) * \sinh(x)^2 + \sqrt{2} * (b^2 * \cosh(x)^6 + 6 * b^2 * \cosh(x) * \sinh(x)^5 + b^2 * \sinh(x)^6 + 3 * b^2 * \cosh(x)^4 + 3 * (5 * b^2 * \cosh(x)^2 + b^2) * \sinh(x)^4 + 4 * (5 * b^2 * \cosh(x)^3 + 3 * b^2 * \cosh(x)) * \sinh(x)^3 - (a^2 - 2 * a * b - 3 * b^2) * \cosh(x)^2 + (15 * b^2 * \cosh(x)^4 + 18 * b^2 * \cosh(x)^2 - a^2 + 2 * a * b + 3 * b^2) * \sinh(x)^2 + a^2 + 2 * a * b + b^2 + 2 * (3 * b^2 * \cosh(x)^5 + 6 * b^2 * \cosh(x)^3 - (a^2 - 2 * a * b - 3 * b^2) * \cosh(x)) * \sinh(x)) * \sqrt{a + b} * \sqrt{((a + b) * \cosh(x)^2 + (a + b) * \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 * \cosh(x) * \sinh(x) + \sinh(x)^2))} + 4 * (2 * (a * b^2 + b^3) * \cosh(x)^7 + 3 * (a * b^2 + 2 * b^3) * \cosh(x)^5 + (a^3 - a^2 * b + 4 * a * b^2 + 6 * b^3) * \cosh(x)^3 - (a^3 - 3 * a * b^2 - 2 * b^3) * \cosh(x)) * \sinh(x)) / (\cosh(x)^6 + 6 * \cosh(x)^5 * \sinh(x) + 15 * \cosh(x)^4 * \sinh(x)^2 + 20 * \cosh(x)^3 * \sinh(x)^3 + 15 * \cosh(x)^2 * \sinh(x)^4 + 6 * \cosh(x) * \sinh(x)^5 + \sinh(x)^6)) + \sqrt{a + b} * b * \log(-((a + b) * \cosh(x)^4 + 4 * (a + b) * \cosh(x) * \sinh(x)^3 + (a + b) * \sinh(x)^4 - 2 * a * \cosh(x)^2 + 2 * (3 * (a + b) * \cosh(x)^2 - a) * \sinh(x)^2 + \sqrt{2} * (\cosh(x)^2 + 2 * \cosh(x) * \sinh(x) + \sinh(x)^2 - 1) * \sqrt{a + b} * \sqrt{((a + b) * \cosh(x)^2 + (a + b) * \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 * \cosh(x) * \sinh(x) + \sinh(x)^2))} + 4 * ((a + b) * \cosh(x)^3 - a * \cosh(x)) * \sinh(x) + a + b) / (\cosh(x)^2 + 2 * \cosh(x) * \sinh(x) + \sinh(x)^2))) / (a * b + b^2), -1/2 * (\sqrt{-a - b} * b * \arctan(\sqrt{2} * (b * \cosh(x)^2 + 2 * b * \cosh(x) * \sinh(x) + b * \sinh(x)^2 + a + b) * \sqrt{-a - b} * \sqrt{((a + b) * \cosh(x)^2 + (a + b) * \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 * \cosh(x) * \sinh(x) + \sinh(x)^2))} / ((a * b + b^2) * \cosh(x)^4 + 4 * (a * b + b^2) * \cosh(x) * \sinh(x)^3 + (a * b + b^2) * \sinh(x)^4 - (a^2 - a * b - 2 * b^2) * \cosh(x)^2 + (6 * (a * b + b^2) * \cosh(x)^2 - a^2 + a * b + 2 * b^2) * \sinh(x)^2 + a^2 + 2 * a * b + b^2 + 2 * (2 * (a * b + b^2) * \cosh(x)^3 - (a^2 - a * b - 2 * b^2) * \cosh(x)) * \sinh(x))) + \sqrt{-a - b} * b * \arctan(\sqrt{2} * (\cosh(x)^2 + 2 * \cosh(x) * \sinh(x) + \sinh(x)^2 - 1) * \sqrt{-a - b} * \sqrt{((a + b) * \cosh(x)^2 + (a + b) * \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 * \cosh(x) * \sinh(x) + \sinh(x)^2))} / ((a + b) * \cosh(x)^4 + 4 * (a + b) * \cosh(x) * \sinh(x)^3 + (a + b) * \sinh(x)^4 - 2 * (a - b) * \cosh(x)^2 + 2 * (3 * (a + b) * \cosh(x)^2 - a + b) * \sinh(x)^2 + 4 * ((a + b) * \cosh(x)^3 - (a - b) * \cosh(x)) * \sinh(x) + a + b)) - (a + b) * \sqrt{b} * \log(-((a + 2 * b) * \cosh(x)^4 + 4 * (a + 2 * b) * \cosh(x) * \sinh(x)^3 + (a + 2 * b) * \sinh(x)^4 - 2 * (a - 2 * b) * \cosh(x)^2 + 2 * (3 * (a + 2 * b) * \cosh(x)^2 - a + 2 * b) * \sinh(x)^2 - 2 * \sqrt{2} * (\cosh(x)^2 + 2 * \cosh(x) * \sinh(x) + \sinh(x)^2 + 1) * \sqrt{b} * \sqrt{
\end{aligned}$$

```

(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*((a + 2*b)*cosh(x)^3 - (a - 2*b)*cosh(x))*sinh(x) + a + 2*b)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*sinh(x)^2 - 2*cosh(x)^2 + 4*(cosh(x)^3 - cosh(x))*sinh(x) + 1)))/(a*b + b^2), -1/2*(sqrt(-a - b)*b*arctan(sqrt(2)*(b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + a + b)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a*b + b^2)*cosh(x)^4 + 4*(a*b + b^2)*cosh(x)*sinh(x)^3 + (a*b + b^2)*sinh(x)^4 - (a^2 - a*b - 2*b^2)*cosh(x)^2 + (6*(a*b + b^2)*cosh(x)^2 - a^2 + a*b + 2*b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(2*(a*b + b^2)*cosh(x)^3 - (a^2 - a*b - 2*b^2)*cosh(x))*sinh(x))) + sqrt(-a - b)*b*arctan(sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)) - 2*(a + b)*sqrt(-b)*arctan(sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(-b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)))/(a*b + b^2)]

```

Sympy [F]

$$\int \frac{\coth^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\coth^2(x)}{\sqrt{a + b \coth^2(x)}} dx$$

```
[In] integrate(coth(x)**2/(a+b*coth(x)**2)**(1/2),x)
```

```
[Out] Integral(coth(x)**2/sqrt(a + b*coth(x)**2), x)
```

Maxima [F]

$$\int \frac{\coth^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\coth(x)^2}{\sqrt{b \coth(x)^2 + a}} dx$$

```
[In] integrate(coth(x)^2/(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(coth(x)^2/sqrt(b*coth(x)^2 + a), x)
```


Giac [F(-2)]

Exception generated.

$$\int \frac{\coth^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \text{Exception raised: TypeError}$$

[In] integrate(coth(x)^2/(a+b*coth(x)^2)^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type

Mupad [F(-1)]

Timed out.

$$\int \frac{\coth^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\coth(x)^2}{\sqrt{b \coth(x)^2 + a}} dx$$

[In] int(coth(x)^2/(a + b*coth(x)^2)^(1/2),x)

[Out] int(coth(x)^2/(a + b*coth(x)^2)^(1/2), x)

$$3.34 \quad \int \frac{\coth(x)}{\sqrt{a+b \coth^2(x)}} dx$$

Optimal result	242
Rubi [A] (verified)	242
Mathematica [A] (verified)	244
Maple [B] (verified)	244
Fricas [B] (verification not implemented)	244
Sympy [F]	245
Maxima [F]	246
Giac [B] (verification not implemented)	246
Mupad [B] (verification not implemented)	246

Optimal result

Integrand size = 15, antiderivative size = 29

$$\int \frac{\coth(x)}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}}$$

[Out] $\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2)/(a+b)^{(1/2)})/(a+b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3751, 455, 65, 214}

$$\int \frac{\coth(x)}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}}$$

[In] $\operatorname{Int}[\operatorname{Coth}[x]/\operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2], x]$

[Out] $\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2]/\operatorname{Sqrt}[a + b]]/\operatorname{Sqrt}[a + b]$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] :> \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1) - 1)*(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}], x]] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{LeQ}[-1, n, 0] \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Den}$

ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 455

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_.), x_Symbol] := Dist[1/n, Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n + 1, 0]

Rule 3751

Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{x}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right) \\
 &= \frac{1}{2}\text{Subst}\left(\int \frac{1}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x)\right) \\
 &= \frac{\text{Subst}\left(\int \frac{1}{1+\frac{a}{b}-\frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)}\right)}{b} \\
 &= \frac{\text{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}}$$

[In] Integrate[Coth[x]/Sqrt[a + b*Coth[x]^2], x]

[Out] ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]]/Sqrt[a + b]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 113 vs. 2(23) = 46.

Time = 0.16 (sec) , antiderivative size = 114, normalized size of antiderivative = 3.93

method	result
derivativedivides	$\frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))+2\sqrt{a+b}\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{1+\coth(x)}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\frac{2a+2b+2b(\coth(x)-1)+2\sqrt{a+b}\sqrt{b(\coth(x)-1)^2-2b(\coth(x)-1)+a+b}}{\coth(x)-1}\right)}{2\sqrt{a+b}}$
default	$\frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))+2\sqrt{a+b}\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{1+\coth(x)}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\frac{2a+2b+2b(\coth(x)-1)+2\sqrt{a+b}\sqrt{b(\coth(x)-1)^2-2b(\coth(x)-1)+a+b}}{\coth(x)-1}\right)}{2\sqrt{a+b}}$

[In] int(coth(x)/(a+b*coth(x)^2)^(1/2), x, method=_RETURNVERBOSE)

[Out] 1/2/(a+b)^(1/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b)^(1/2)*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))/(1+coth(x)))+1/2/(a+b)^(1/2)*ln((2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(1/2)*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(coth(x)-1))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 333 vs. 2(23) = 46.

Time = 0.30 (sec) , antiderivative size = 1298, normalized size of antiderivative = 44.76

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} dx = \text{Too large to display}$$

[In] integrate(coth(x)/(a+b*coth(x)^2)^(1/2), x, algorithm="fricas")

[Out] [1/4*(sqrt(a + b)*log(-(a^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3

```

+ a^2*b - 14*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh
(x)^3 - 3*(2*a^3 + a^2*b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b
^3)*cosh(x)^4 + (70*(a^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3
- 30*(2*a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5
- 10*(2*a^3 + a^2*b)*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))*s
inh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*cosh(x
)^2 + 2*(14*(a^3 + a^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*cosh(x)^4 - 2*a^3
- 3*a^2*b + b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 +
sqrt(2)*(a^2*cosh(x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*sinh(x)^6 - 3*a^2*co
sh(x)^4 + 3*(5*a^2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^2*cosh(x)^3 - 3*a^2*
cosh(x))*sinh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 + (15*a^2*cosh(x)^4 -
18*a^2*cosh(x)^2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(
3*a^2*cosh(x)^5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x))*sinh(x))
*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^
2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a^3 + a^2*b)*cosh(x)^7 - 3*(2*a
^3 + a^2*b)*cosh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^3 - (2*a^3
+ 3*a^2*b - b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*co
sh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cos
h(x)*sinh(x)^5 + sinh(x)^6)) + sqrt(a + b)*log(((a + b)*cosh(x)^4 + 4*(a +
b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 + 2*b*cosh(x)^2 + 2*(3*(a + b)*cos
h(x)^2 + b)*sinh(x)^2 + sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2
+ 1)*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh
(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*((a + b)*cosh(x)^3 + b*cosh(x))
*sinh(x) + a + b)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)))/(a + b), -1
/2*(sqrt(-a - b)*arctan(sqrt(2)*(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh
(x)^2 - a - b)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a
+ b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a^2 + a*b)*cosh(x)^4 +
4*(a^2 + a*b)*cosh(x)*sinh(x)^3 + (a^2 + a*b)*sinh(x)^4 - (2*a^2 + a*b - b
^2)*cosh(x)^2 + (6*(a^2 + a*b)*cosh(x)^2 - 2*a^2 - a*b + b^2)*sinh(x)^2 + a
^2 + 2*a*b + b^2 + 2*(2*(a^2 + a*b)*cosh(x)^3 - (2*a^2 + a*b - b^2)*cosh(x)
)*sinh(x))) + sqrt(-a - b)*arctan(sqrt(2)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)
)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2
)))/((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a +
b)))/(a + b)]

```

Sympy [F]

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} dx$$

[In] integrate(coth(x)/(a+b*coth(x)**2)**(1/2),x)

[Out] Integral(coth(x)/sqrt(a + b*coth(x)**2), x)

Maxima [F]

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\coth(x)}{\sqrt{b \coth^2(x) + a}} dx$$

[In] integrate(coth(x)/(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(coth(x)/sqrt(b*coth(x)^2 + a), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 196 vs. 2(23) = 46.

Time = 0.42 (sec) , antiderivative size = 196, normalized size of antiderivative = 6.76

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} dx = \frac{\log\left(\left|-\left(\sqrt{a+be^{2x}}-\sqrt{ae^{4x}+be^{4x}-2ae^{2x}+2be^{2x}+a+b}\right)(a+b)+\sqrt{a+b}(a-b)\right|\right)}{\sqrt{a+b}} + \frac{\log\left(\left|-\sqrt{a+be^{2x}}+\sqrt{ae^{4x}+be^{4x}-2ae^{2x}+2be^{2x}+a+b}\right|\right)}{\sqrt{a+b}}$$

$$- \frac{2 \operatorname{sgn}(e^{2x} - 1)}{2 \operatorname{sgn}(e^{2x} - 1)}$$

[In] integrate(coth(x)/(a+b*coth(x)^2)^(1/2),x, algorithm="giac")

[Out] -1/2*(log(abs(-sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*(a + b) + sqrt(a + b)*(a - b)))/sqrt(a + b) + log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) + sqrt(a + b)))/sqrt(a + b) - log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - sqrt(a + b)))/sqrt(a + b))/sgn(e^(2*x) - 1)

Mupad [B] (verification not implemented)

Time = 2.31 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^2(x)}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \coth^2(x) + a}}{\sqrt{a + b}}\right)}{\sqrt{a + b}}$$

[In] int(coth(x)/(a + b*coth(x)^2)^(1/2),x)

[Out] atanh((a + b*coth(x)^2)^(1/2)/(a + b)^(1/2))/(a + b)^(1/2)

$$3.35 \quad \int \frac{1}{\sqrt{a+b \coth^2(x)}} dx$$

Optimal result	247
Rubi [A] (verified)	247
Mathematica [B] (verified)	248
Maple [B] (verified)	249
Fricas [B] (verification not implemented)	249
Sympy [F]	250
Maxima [F]	250
Giac [B] (verification not implemented)	251
Mupad [B] (verification not implemented)	251

Optimal result

Integrand size = 12, antiderivative size = 31

$$\int \frac{1}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{\sqrt{a+b}}$$

[Out] $\operatorname{arctanh}(\coth(x) \cdot (a+b)^{(1/2)} / (a+b \cdot \coth(x)^2)^{(1/2)}) / (a+b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3742, 385, 212}

$$\int \frac{1}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{\sqrt{a+b}}$$

[In] $\operatorname{Int}[1/\operatorname{Sqrt}[a + b \cdot \operatorname{Coth}[x]^2], x]$

[Out] $\operatorname{ArcTanh}[(\operatorname{Sqrt}[a + b] \cdot \operatorname{Coth}[x]) / \operatorname{Sqrt}[a + b \cdot \operatorname{Coth}[x]^2]] / \operatorname{Sqrt}[a + b]$

Rule 212

$\operatorname{Int}[(a_+ + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2] \cdot \operatorname{Rt}[-b, 2])) \cdot \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2] \cdot (x/\operatorname{Rt}[a, 2])], x] /;$ $\operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b] \ \&\& \ (\operatorname{Gt} Q[a, 0] \ || \ \operatorname{Lt} Q[b, 0])$

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Su
bst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b
, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :=
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(
ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst} \left(\int \frac{1}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x) \right) \\ &= \text{Subst} \left(\int \frac{1}{1-(a+b)x^2} dx, x, \frac{\coth(x)}{\sqrt{a+b\coth^2(x)}} \right) \\ &= \frac{\text{arctanh} \left(\frac{\sqrt{a+b}\coth(x)}{\sqrt{a+b\coth^2(x)}} \right)}{\sqrt{a+b}} \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 77 vs. $2(31) = 62$.

Time = 0.14 (sec) , antiderivative size = 77, normalized size of antiderivative = 2.48

$$\int \frac{1}{\sqrt{a+b\coth^2(x)}} dx = \frac{\text{arctanh} \left(\frac{\sqrt{\frac{(a+b)\coth^2(x)}{a}}}{\sqrt{1+\frac{b\coth^2(x)}{a}}} \right) \coth(x) \sqrt{1+\frac{b\coth^2(x)}{a}}}{\sqrt{\frac{(a+b)\coth^2(x)}{a}} \sqrt{a+b\coth^2(x)}}$$

```
[In] Integrate[1/Sqrt[a + b*Coth[x]^2], x]
```

```
[Out] (ArcTanh[Sqrt[((a + b)*Coth[x]^2)/a]/Sqrt[1 + (b*Coth[x]^2)/a]]*Coth[x]*Sqr
t[1 + (b*Coth[x]^2)/a])/(Sqrt[((a + b)*Coth[x]^2)/a]*Sqrt[a + b*Coth[x]^2])
```


Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 113 vs. $2(25) = 50$.

Time = 0.14 (sec) , antiderivative size = 114, normalized size of antiderivative = 3.68

method	result
derivativedivides	$-\frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))+2\sqrt{a+b}\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{1+\coth(x)}}{2\sqrt{a+b}}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\frac{2a+2b+2b(\coth(x)-1)+2\sqrt{a+b}\sqrt{b(\coth(x)-1)}}{\coth(x)-1}}{2\sqrt{a+b}}\right)}{2\sqrt{a+b}}$
default	$-\frac{\ln\left(\frac{2a+2b-2b(1+\coth(x))+2\sqrt{a+b}\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}{1+\coth(x)}}{2\sqrt{a+b}}\right)}{2\sqrt{a+b}} + \frac{\ln\left(\frac{2a+2b+2b(\coth(x)-1)+2\sqrt{a+b}\sqrt{b(\coth(x)-1)}}{\coth(x)-1}}{2\sqrt{a+b}}\right)}{2\sqrt{a+b}}$

[In] `int(1/(a+b*coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/2/(a+b)^{(1/2)}*\ln((2*a+2*b-2*b*(1+\coth(x))+2*(a+b)^{(1/2)}*(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^{(1/2)})/(1+\coth(x)))+1/2/(a+b)^{(1/2)}*\ln((2*a+2*b+2*b*(\coth(x)-1)+2*(a+b)^{(1/2)}*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{(1/2)})/(\coth(x)-1))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 399 vs. $2(25) = 50$.

Time = 0.30 (sec) , antiderivative size = 1357, normalized size of antiderivative = 43.77

$$\int \frac{1}{\sqrt{a + b \coth^2(x)}} dx = \text{Too large to display}$$

[In] `integrate(1/(a+b*coth(x)^2)^(1/2),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & [1/4*(\sqrt{a + b})*\log(((a*b^2 + b^3)*\cosh(x)^8 + 8*(a*b^2 + b^3)*\cosh(x)*\sinh(x)^7 + (a*b^2 + b^3)*\sinh(x)^8 + 2*(a*b^2 + 2*b^3)*\cosh(x)^6 + 2*(a*b^2 + 2*b^3 + 14*(a*b^2 + b^3)*\cosh(x)^2)*\sinh(x)^6 + 4*(14*(a*b^2 + b^3)*\cosh(x)^3 + 3*(a*b^2 + 2*b^3)*\cosh(x))*\sinh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*\cosh(x)^4 + (70*(a*b^2 + b^3)*\cosh(x)^4 + a^3 - a^2*b + 4*a*b^2 + 6*b^3 + 30*(a*b^2 + 2*b^3)*\cosh(x)^2)*\sinh(x)^4 + 4*(14*(a*b^2 + b^3)*\cosh(x)^5 + 10*(a*b^2 + 2*b^3)*\cosh(x)^3 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*\cosh(x))*\sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 - 3*a*b^2 - 2*b^3)*\cosh(x)^2 + 2*(14*(a*b^2 + b^3)*\cosh(x)^6 + 15*(a*b^2 + 2*b^3)*\cosh(x)^4 - a^3 + 3*a*b^2 + 2*b^3 + 3*(a^3 - a^2*b + 4*a*b^2 + 6*b^3)*\cosh(x)^2)*\sinh(x)^2 + \sqrt{2}*(b^2*\cosh(x)^6 + 6*b^2*\cosh(x)*\sinh(x)^5 + b^2*\sinh(x)^6 + 3*b^2*\cosh(x)^4 + 3*(5*b^2*\cosh(x)^2 + b^2)*\sinh(x)^4 + 4*(5*b^2*\cosh(x)^3 + 3*b^2*\cosh(x))*\sinh(x)^3 - (a^2 - 2*a*b - 3*b^2)*\cosh(x)^2 + (15*b^2*\cosh(x)^4 + 18*b^2*\cosh(x)^2 - a^2 + 2*a*b + 3*b^2)*\sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(3*b^2*\cosh(x)^5 + 6*b^2*\cosh(x)^3 - (a^2 - 2*a*b - 3*b^2)*\cosh(x))*\sinh(x))* \end{aligned}$$

```

sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2
- 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a*b^2 + b^3)*cosh(x)^7 + 3*(a*b^
2 + 2*b^3)*cosh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^3 - (a^3 - 3
*a*b^2 - 2*b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cos
h(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh
(x)*sinh(x)^5 + sinh(x)^6)) + sqrt(a + b)*log(-((a + b)*cosh(x)^4 + 4*(a +
b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*a*cosh(x)^2 + 2*(3*(a + b)*cos
h(x)^2 - a)*sinh(x)^2 + sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2
- 1)*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh
(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*((a + b)*cosh(x)^3 - a*cosh(x))
*sinh(x) + a + b)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)))/(a + b), -1
/2*(sqrt(-a - b)*arctan(sqrt(2)*(b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh
(x)^2 + a + b)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a
+ b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a*b + b^2)*cosh(x)^4 +
4*(a*b + b^2)*cosh(x)*sinh(x)^3 + (a*b + b^2)*sinh(x)^4 - (a^2 - a*b - 2*b
^2)*cosh(x)^2 + (6*(a*b + b^2)*cosh(x)^2 - a^2 + a*b + 2*b^2)*sinh(x)^2 + a
^2 + 2*a*b + b^2 + 2*(2*(a*b + b^2)*cosh(x)^3 - (a^2 - a*b - 2*b^2)*cosh(x)
)*sinh(x))) + sqrt(-a - b)*arctan(sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) +
sinh(x)^2 - 1)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a
+ b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a + b)*cosh(x)^4 + 4*(
a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(
a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x)
))*sinh(x) + a + b)))/(a + b)]

```

Sympy [F]

$$\int \frac{1}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{1}{\sqrt{a + b \coth^2(x)}} dx$$

```
[In] integrate(1/(a+b*coth(x)**2)**(1/2),x)
```

```
[Out] Integral(1/sqrt(a + b*coth(x)**2), x)
```

Maxima [F]

$$\int \frac{1}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{1}{\sqrt{b \coth(x)^2 + a}} dx$$

```
[In] integrate(1/(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(1/sqrt(b*coth(x)^2 + a), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 196 vs. 2(25) = 50.

Time = 0.39 (sec) , antiderivative size = 196, normalized size of antiderivative = 6.32

$$\int \frac{1}{\sqrt{a + b \coth^2(x)}} dx = \frac{\log\left(\left|-\left(\sqrt{a+b}e^{2x}-\sqrt{ae^{4x}+be^{4x}-2ae^{2x}+2be^{2x}+a+b}\right)(a+b)+\sqrt{a+b}(a-b)\right|\right)}{\sqrt{a+b}} - \frac{\log\left(\left|-\sqrt{a+b}e^{2x}+\sqrt{ae^{4x}+be^{4x}-2ae^{2x}+2be^{2x}+a+b}\right|\right)}{2 \operatorname{sgn}(e^{2x}-1)}$$

[In] integrate(1/(a+b*coth(x)^2)^(1/2),x, algorithm="giac")

[Out] -1/2*(log(abs(-sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*(a + b) + sqrt(a + b)*(a - b)))/sqrt(a + b) - 1/2*log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) + sqrt(a + b)))/sqrt(a + b) + log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - sqrt(a + b)))/sqrt(a + b))/sgn(e^(2*x) - 1)

Mupad [B] (verification not implemented)

Time = 2.22 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.81

$$\int \frac{1}{\sqrt{a + b \coth^2(x)}} dx = \frac{\operatorname{atanh}\left(\frac{\coth(x) \sqrt{a+b}}{\sqrt{b \coth^2(x) + a}}\right)}{\sqrt{a+b}}$$

[In] int(1/(a + b*coth(x)^2)^(1/2),x)

[Out] atanh((coth(x)*(a + b)^(1/2))/(a + b*coth(x)^2)^(1/2))/(a + b)^(1/2)

$$3.36 \quad \int \frac{\tanh(x)}{\sqrt{a+b \coth^2(x)}} dx$$

Optimal result	252
Rubi [A] (verified)	252
Mathematica [A] (verified)	254
Maple [F]	254
Fricas [B] (verification not implemented)	254
Sympy [F]	257
Maxima [F]	257
Giac [F(-2)]	257
Mupad [F(-1)]	257

Optimal result

Integrand size = 15, antiderivative size = 56

$$\int \frac{\tanh(x)}{\sqrt{a+b \coth^2(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}}$$

[Out] $-\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2)}/a^{(1/2)})/a^{(1/2)}+\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2)}/(a+b)^{(1/2)})/(a+b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3751, 457, 88, 65, 214}

$$\int \frac{\tanh(x)}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}}$$

[In] `Int[Tanh[x]/Sqrt[a + b*Coth[x]^2],x]`

[Out] $-(\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2]/\operatorname{Sqrt}[a]]/\operatorname{Sqrt}[a]) + \operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2]/\operatorname{Sqrt}[a + b]]/\operatorname{Sqrt}[a + b]$

Rule 65

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +`

$d*(x^p/b)^n, x], x, (a + b*x)^{1/p}], x]] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 88

$\text{Int}[(e_.) + (f_.)*(x_.)^p]/((a_.) + (b_.)*(x_.)*((c_.) + (d_.)*(x_))), x_Symbol] \rightarrow \text{Dist}[b/(b*c - a*d), \text{Int}[(e + f*x)^p/(a + b*x), x], x] - \text{Dist}[d/(b*c - a*d), \text{Int}[(e + f*x)^p/(c + d*x), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, p\}, x] \&\& \text{!IntegerQ}[p]$

Rule 214

$\text{Int}[(a_.) + (b_.)*(x_.)^2]^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{NegQ}[a/b]$

Rule 457

$\text{Int}[(x_.)^{m_.*((a_.) + (b_.)*(x_.)^{n_})^p*((c_.) + (d_.)*(x_.)^{n_})^q}, x_Symbol] \rightarrow \text{Dist}[1/n, \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)*(a + b*x)^p*(c + d*x)^q}, x], x, x^n], x] /; \text{FreeQ}\{a, b, c, d, m, n, p, q\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]$

Rule 3751

$\text{Int}[(d_.)*\tan[(e_.) + (f_.)*(x_.)]^{m_.*((a_.) + (b_.)*((c_.)*\tan[(e_.) + (f_.)*(x_.)]^{n_})^p}, x_Symbol] \rightarrow \text{With}\{\text{ff} = \text{FreeFactors}[\text{Tan}[e + f*x], x]\}, \text{Dist}[c*(\text{ff}/f), \text{Subst}[\text{Int}[(d*\text{ff}*(x/c))^m*((a + b*(\text{ff}*x)^n)^p/(c^2 + \text{ff}^2*x^2)], x], x, c*(\text{Tan}[e + f*x]/\text{ff})], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x] \&\& (\text{IGtQ}[p, 0] \parallel \text{EqQ}[n, 2] \parallel \text{EqQ}[n, 4] \parallel (\text{IntegerQ}[p] \&\& \text{RationalQ}[n]))$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{1}{x(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right) \\
 &= \frac{1}{2}\text{Subst}\left(\int \frac{1}{(1-x)x\sqrt{a+bx}} dx, x, \coth^2(x)\right) \\
 &= \frac{1}{2}\text{Subst}\left(\int \frac{1}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x)\right) + \frac{1}{2}\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx}} dx, x, \coth^2(x)\right) \\
 &= \frac{\text{Subst}\left(\int \frac{1}{1+\frac{a-x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)}\right)}{b} + \frac{\text{Subst}\left(\int \frac{1}{-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)}\right)}{b}
 \end{aligned}$$

$$= -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(x)}{\sqrt{a+b\coth^2(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right)}{\sqrt{a+b}}$$

[In] Integrate[Tanh[x]/Sqrt[a + b*Coth[x]^2],x]

[Out] -(ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a]]/Sqrt[a]) + ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]]/Sqrt[a + b]

Maple [F]

$$\int \frac{\tanh(x)}{\sqrt{a+b\coth(x)^2}} dx$$

[In] int(tanh(x)/(a+b*coth(x)^2)^(1/2),x)

[Out] int(tanh(x)/(a+b*coth(x)^2)^(1/2),x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 503 vs. 2(44) = 88.

Time = 0.39 (sec) , antiderivative size = 3397, normalized size of antiderivative = 60.66

$$\int \frac{\tanh(x)}{\sqrt{a+b\coth^2(x)}} dx = \text{Too large to display}$$

[In] integrate(tanh(x)/(a+b*coth(x)^2)^(1/2),x, algorithm="fricas")

[Out] [1/4*(sqrt(a + b)*a*log(-(a^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^4 + (70*(a^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))

$$\begin{aligned}
& * \sinh(x)^3 + a^3 + 3a^2b + 3ab^2 + b^3 - 2(2a^3 + 3a^2b - b^3) \cosh(x)^2 + 2(14(a^3 + a^2b) \cosh(x)^6 - 15(2a^3 + a^2b) \cosh(x)^4 - 2a^3 - 3a^2b + b^3 + 3(6a^3 + 4a^2b - ab^2 + b^3) \cosh(x)^2) \sinh(x)^2 \\
& + \sqrt{2}(a^2 \cosh(x)^6 + 6a^2 \cosh(x) \sinh(x)^5 + a^2 \sinh(x)^6 - 3a^2 \cosh(x)^4 + 3(5a^2 \cosh(x)^2 - a^2) \sinh(x)^4 + 4(5a^2 \cosh(x)^3 - 3a^2 \cosh(x)) \sinh(x)^3 + (3a^2 + 2ab - b^2) \cosh(x)^2 + (15a^2 \cosh(x)^4 - 18a^2 \cosh(x)^2 + 3a^2 + 2ab - b^2) \sinh(x)^2 - a^2 - 2ab - b^2 + 2 \\
& * (3a^2 \cosh(x)^5 - 6a^2 \cosh(x)^3 + (3a^2 + 2ab - b^2) \cosh(x)) \sinh(x)) \sqrt{a+b} \sqrt{((a+b) \cosh(x)^2 + (a+b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2))} + 4(2(a^3 + a^2b) \cosh(x)^7 - 3(2a^3 + a^2b) \cosh(x)^5 + (6a^3 + 4a^2b - ab^2 + b^3) \cosh(x)^3 - (2a^3 + 3a^2b - b^3) \cosh(x)) \sinh(x) / (\cosh(x)^6 + 6 \cosh(x)^5 \sinh(x) + 15 \cosh(x)^4 \sinh(x)^2 + 20 \cosh(x)^3 \sinh(x)^3 + 15 \cosh(x)^2 \sinh(x)^4 + 6 \cosh(x) \sinh(x)^5 + \sinh(x)^6) + 2(a+b) \sqrt{a} \log(-((2a+b) \cosh(x)^4 + 4(2a+b) \cosh(x) \sinh(x)^3 + (2a+b) \sinh(x)^4 - 2(2a-b) \cosh(x)^2 + 2(3(2a+b) \cosh(x)^2 - 2a + b) \sinh(x)^2 - 2\sqrt{2}(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1) \sqrt{a} \sqrt{((a+b) \cosh(x)^2 + (a+b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2))} + 4((2a+b) \cosh(x)^3 - (2a-b) \cosh(x)) \sinh(x) + 2a + b) / (\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + 2(3 \cosh(x)^2 + 1) \sinh(x)^2 + 2 \cosh(x)^2 + 4(\cosh(x)^3 + \cosh(x)) \sinh(x) + 1)) + \sqrt{a+b} a \log(((a+b) \cosh(x)^4 + 4(a+b) \cosh(x) \sinh(x)^3 + (a+b) \sinh(x)^4 + 2b \cosh(x)^2 + 2(3(a+b) \cosh(x)^2 + b) \sinh(x)^2 + \sqrt{2}(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1) \sqrt{a+b} \sqrt{((a+b) \cosh(x)^2 + (a+b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2))} + 4((a+b) \cosh(x)^3 + b \cosh(x)) \sinh(x) + a + b) / (\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2)) / (a^2 + ab), 1/4(4\sqrt{-a}(a+b) \arctan(\sqrt{2}(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1) \sqrt{-a} \sqrt{((a+b) \cosh(x)^2 + (a+b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2))} / ((a+b) \cosh(x)^4 + 4(a+b) \cosh(x) \sinh(x)^3 + (a+b) \sinh(x)^4 - 2(a-b) \cosh(x)^2 + 2(3(a+b) \cosh(x)^2 - a + b) \sinh(x)^2 + 4((a+b) \cosh(x)^3 - (a-b) \cosh(x)) \sinh(x) + a + b)) + \sqrt{a+b} a \log(-((a^3 + a^2b) \cosh(x)^8 + 8(a^3 + a^2b) \cosh(x) \sinh(x)^7 + (a^3 + a^2b) \sinh(x)^8 - 2(2a^3 + a^2b) \cosh(x)^6 - 2(2a^3 + a^2b - 14(a^3 + a^2b) \cosh(x)^2) \sinh(x)^6 + 4(14(a^3 + a^2b) \cosh(x)^3 - 3(2a^3 + a^2b) \cosh(x)) \sinh(x)^5 + (6a^3 + 4a^2b - ab^2 + b^3) \cosh(x)^4 + (70(a^3 + a^2b) \cosh(x)^4 + 6a^3 + 4a^2b - ab^2 + b^3 - 30(2a^3 + a^2b) \cosh(x)^2) \sinh(x)^4 + 4(14(a^3 + a^2b) \cosh(x)^5 - 10(2a^3 + a^2b) \cosh(x)^3 + (6a^3 + 4a^2b - ab^2 + b^3) \cosh(x)) \sinh(x)^3 + a^3 + 3a^2b + 3ab^2 + b^3 - 2(2a^3 + 3a^2b - b^3) \cosh(x)^2 + 2(14(a^3 + a^2b) \cosh(x)^6 - 15(2a^3 + a^2b) \cosh(x)^4 - 2a^3 - 3a^2b + b^3 + 3(6a^3 + 4a^2b - ab^2 + b^3) \cosh(x)^2) \sinh(x)^2 + \sqrt{2}(a^2 \cosh(x)^6 + 6a^2 \cosh(x) \sinh(x)^5 + a^2 \sinh(x)^6 - 3a^2 \cosh(x)^4 + 3(5a^2 \cosh(x)^2 - a^2) \sinh(x)^4 + 4(5a^2 \cosh(x)^3 - 3a^2 \cosh(x)) \sinh(x)^3 + (3a^2 + 2ab - b^2) \cosh(x)^2 + (15a^2 \cosh(x)^4 - 18a^2 \cosh(x)^2 + 3a^2 + 2ab - b^2) \sinh(x)^2
\end{aligned}$$

$$\begin{aligned}
& 2 - a^2 - 2ab - b^2 + 2(3a^2 \cosh(x)^5 - 6a^2 \cosh(x)^3 + (3a^2 + 2a \\
& * b - b^2) \cosh(x) \sinh(x)) \sqrt{a+b} \sqrt{((a+b) \cosh(x)^2 + (a+b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2)} + 4(2(a^3 \\
& + a^2 b) \cosh(x)^7 - 3(2a^3 + a^2 b) \cosh(x)^5 + (6a^3 + 4a^2 b - ab^2 \\
& + b^3) \cosh(x)^3 - (2a^3 + 3a^2 b - b^3) \cosh(x) \sinh(x)) / (\cosh(x)^6 + \\
& 6 \cosh(x)^5 \sinh(x) + 15 \cosh(x)^4 \sinh(x)^2 + 20 \cosh(x)^3 \sinh(x)^3 + 15 \cosh(x)^2 \sinh(x)^4 + 6 \cosh(x) \sinh(x)^5 + \sinh(x)^6) + \sqrt{a+b} a \log \\
& (((a+b) \cosh(x)^4 + 4(a+b) \cosh(x) \sinh(x)^3 + (a+b) \sinh(x)^4 + 2b \\
& * \cosh(x)^2 + 2(3(a+b) \cosh(x)^2 + b) \sinh(x)^2 + \sqrt{2}(\cosh(x)^2 + 2 \\
& * \cosh(x) \sinh(x) + \sinh(x)^2 + 1)) \sqrt{a+b} \sqrt{((a+b) \cosh(x)^2 + (a \\
& + b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2)} + 4((\\
& a+b) \cosh(x)^3 + b \cosh(x) \sinh(x) + a + b) / (\cosh(x)^2 + 2 \cosh(x) \sinh(x) \\
& + \sinh(x)^2))) / (a^2 + ab), -1/2(a \sqrt{-a-b}) \arctan(\sqrt{2}(a \cosh(x)^2 + 2a \cosh(x) \sinh(x) + a \sinh(x)^2 - a - b) \sqrt{-a-b}) \sqrt{((a+b) \cosh(x)^2 + (a+b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2)}) / ((a^2 + ab) \cosh(x)^4 + 4(a^2 + ab) \cosh(x) \sinh(x)^3 + (a^2 + ab) \sinh(x)^4 - (2a^2 + ab - b^2) \cosh(x)^2 + (6(a^2 + ab) \cosh(x)^2 - 2a^2 - ab + b^2) \sinh(x)^2 + a^2 + 2ab + b^2 + 2(2(a^2 + ab) \cosh(x)^3 - (2a^2 + ab - b^2) \cosh(x) \sinh(x))) + a \sqrt{-a-b}) \arctan(\sqrt{2} \sqrt{-a-b} \sqrt{((a+b) \cosh(x)^2 + (a+b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2)}) / ((a+b) \cosh(x)^2 + 2(a+b) \cosh(x) \sinh(x) + (a+b) \sinh(x)^2 + a + b)) - (a+b) \sqrt{a} \log(-((2a+b) \cosh(x)^4 + 4(2a+b) \cosh(x) \sinh(x)^3 + (2a+b) \sinh(x)^4 - 2(2a-b) \cosh(x)^2 + 2(3(2a+b) \cosh(x)^2 - 2a+b) \sinh(x)^2 - 2\sqrt{2}(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1)) \sqrt{a} \sqrt{((a+b) \cosh(x)^2 + (a+b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2)}) + 4(((2a+b) \cosh(x)^3 - (2a-b) \cosh(x) \sinh(x) + 2a+b) / (\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + 2(3 \cosh(x)^2 + 1) \sinh(x)^2 + 2 \cosh(x)^2 + 4(\cosh(x)^3 + \cosh(x) \sinh(x) + 1)))) / (a^2 + ab), 1/2(2 \sqrt{2} \sqrt{-a}(a+b) \arctan(\sqrt{2}(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1) \sqrt{-a}) \sqrt{((a+b) \cosh(x)^2 + (a+b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2)}) / ((a+b) \cosh(x)^4 + 4(a+b) \cosh(x) \sinh(x)^3 + (a+b) \sinh(x)^4 - 2(a-b) \cosh(x)^2 + 2(3(a+b) \cosh(x)^2 - a+b) \sinh(x)^2 + 4((a+b) \cosh(x)^3 - (a-b) \cosh(x) \sinh(x) + a+b)) - a \sqrt{-a-b}) \arctan(\sqrt{2}(a \cosh(x)^2 + 2a \cosh(x) \sinh(x) + a \sinh(x)^2 - a - b) \sqrt{-a-b}) \sqrt{((a+b) \cosh(x)^2 + (a+b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2)}) / ((a^2 + ab) \cosh(x)^4 + 4(a^2 + ab) \cosh(x) \sinh(x)^3 + (a^2 + ab) \sinh(x)^4 - (2a^2 + ab - b^2) \cosh(x)^2 + (6(a^2 + ab) \cosh(x)^2 - 2a^2 - ab + b^2) \sinh(x)^2 + a^2 + 2ab + b^2 + 2(2(a^2 + ab) \cosh(x)^3 - (2a^2 + ab - b^2) \cosh(x) \sinh(x))) - a \sqrt{-a-b}) \arctan(\sqrt{2} \sqrt{-a-b} \sqrt{((a+b) \cosh(x)^2 + (a+b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2)}) / ((a+b) \cosh(x)^2 + 2(a+b) \cosh(x) \sinh(x) + (a+b) \sinh(x)^2 + a + b))) / (a^2 + ab)]
\end{aligned}$$

Sympy [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{a + b \coth^2(x)}} dx$$

[In] integrate(tanh(x)/(a+b*coth(x)**2)**(1/2),x)

[Out] Integral(tanh(x)/sqrt(a + b*coth(x)**2), x)

Maxima [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{b \coth(x)^2 + a}} dx$$

[In] integrate(tanh(x)/(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(tanh(x)/sqrt(b*coth(x)^2 + a), x)

Giac [F(-2)]

Exception generated.

$$\int \frac{\tanh(x)}{\sqrt{a + b \coth^2(x)}} dx = \text{Exception raised: TypeError}$$

[In] integrate(tanh(x)/(a+b*coth(x)^2)^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx)::OUTPUT:Error: Bad Argument Type

Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{b \coth(x)^2 + a}} dx$$

[In] int(tanh(x)/(a + b*coth(x)^2)^(1/2),x)

[Out] int(tanh(x)/(a + b*coth(x)^2)^(1/2), x)

$$3.37 \quad \int \frac{\tanh^2(x)}{\sqrt{a+b \coth^2(x)}} dx$$

Optimal result	258
Rubi [A] (verified)	258
Mathematica [C] (warning: unable to verify)	260
Maple [F]	260
Fricas [B] (verification not implemented)	260
Sympy [F]	262
Maxima [F]	262
Giac [F(-2)]	262
Mupad [F(-1)]	263

Optimal result

Integrand size = 17, antiderivative size = 51

$$\int \frac{\tanh^2(x)}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{\sqrt{a+b}} - \frac{\sqrt{a+b \coth^2(x)} \tanh(x)}{a}$$

[Out] $\operatorname{arctanh}(\coth(x) \cdot (a+b)^{(1/2)} / (a+b \cdot \coth(x)^2)^{(1/2)}) / (a+b)^{(1/2)} - (a+b \cdot \coth(x)^2)^{(1/2)} \cdot \tanh(x) / a$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3751, 491, 12, 385, 212}

$$\int \frac{\tanh^2(x)}{\sqrt{a+b \coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{\sqrt{a+b}} - \frac{\tanh(x) \sqrt{a+b \coth^2(x)}}{a}$$

[In] `Int[Tanh[x]^2/Sqrt[a + b*Coth[x]^2],x]`

[Out] `ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]]/Sqrt[a + b] - (Sqrt[a + b*Coth[x]^2]*Tanh[x])/a`

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]`

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 385

Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 491

Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[(e*x)^(m + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*c*e*(m + 1))), x] - Dist[1/(a*c*e^n*(m + 1)), Int[(e*x)^(m + n)*(a + b*x^n)^p*(c + d*x^n)^q*Simp[(b*c + a*d)*(m + n + 1) + n*(b*c*p + a*d*q) + b*d*(m + n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

Rule 3751

Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{1}{x^2(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right) \\
 &= -\frac{\sqrt{a+b\coth^2(x)\tanh(x)}}{a} + \frac{\text{Subst}\left(\int \frac{a}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right)}{a} \\
 &= -\frac{\sqrt{a+b\coth^2(x)\tanh(x)}}{a} + \text{Subst}\left(\int \frac{1}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right) \\
 &= -\frac{\sqrt{a+b\coth^2(x)\tanh(x)}}{a} + \text{Subst}\left(\int \frac{1}{1-(a+b)x^2} dx, x, \frac{\coth(x)}{\sqrt{a+b\coth^2(x)}}\right)
 \end{aligned}$$

$$= \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b}\coth(x)}{\sqrt{a+b\coth^2(x)}}\right)}{\sqrt{a+b}} - \frac{\sqrt{a+b\coth^2(x)}\tanh(x)}{a}$$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.77 (sec) , antiderivative size = 127, normalized size of antiderivative = 2.49

$$\int \frac{\tanh^2(x)}{\sqrt{a+b\coth^2(x)}} dx$$

$$= \frac{\left(1 + \frac{b\coth^2(x)}{a}\right) \sinh^2(x) \left(\frac{4(a+b)\cosh^2(x)(a+b\coth^2(x)) \operatorname{Hypergeometric2F1}\left(2, 2, \frac{5}{2}, \frac{(a+b)\cosh^2(x)}{a}\right)}{3a^2} + \frac{\arcsin\left(\sqrt{\frac{(a+b)\cosh^2(x)}{a}}\right)(a+b\coth^2(x))}{a\sqrt{-\frac{(a+b)\cosh^2(x)(a+b\coth^2(x))}{a^2}}}\right)}{\sqrt{a+b\coth^2(x)}}$$

[In] Integrate[Tanh[x]^2/Sqrt[a + b*Coth[x]^2], x]

[Out] ((1 + (b*Coth[x]^2)/a)*Sinh[x]^2*((4*(a + b)*Cosh[x]^2*(a + b*Coth[x]^2)*Hypergeometric2F1[2, 2, 5/2, ((a + b)*Cosh[x]^2)/a])/(3*a^2) + (ArcSin[Sqrt[(a + b)*Cosh[x]^2/a]]*(a + 2*b*Coth[x]^2))/(a*Sqrt[-((a + b)*Cosh[x]^2*(a + b*Coth[x]^2)*Sinh[x]^2/a^2)]))*Tanh[x])/Sqrt[a + b*Coth[x]^2]

Maple [F]

$$\int \frac{\tanh(x)^2}{\sqrt{a+b\coth(x)^2}} dx$$

[In] int(tanh(x)^2/(a+b*coth(x)^2)^(1/2), x)

[Out] int(tanh(x)^2/(a+b*coth(x)^2)^(1/2), x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 531 vs. 2(43) = 86.

Time = 0.33 (sec) , antiderivative size = 1621, normalized size of antiderivative = 31.78

$$\int \frac{\tanh^2(x)}{\sqrt{a+b\coth^2(x)}} dx = \text{Too large to display}$$

[In] integrate(tanh(x)^2/(a+b*coth(x)^2)^(1/2), x, algorithm="fricas")

```
[Out] [1/4*((a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 + a)*sqrt(a + b)*log
(((a*b^2 + b^3)*cosh(x)^8 + 8*(a*b^2 + b^3)*cosh(x)*sinh(x)^7 + (a*b^2 + b^
3)*sinh(x)^8 + 2*(a*b^2 + 2*b^3)*cosh(x)^6 + 2*(a*b^2 + 2*b^3 + 14*(a*b^2 +
b^3)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a*b^2 + b^3)*cosh(x)^3 + 3*(a*b^2 + 2*b
^3)*cosh(x))*sinh(x)^5 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^4 + (70*(a
*b^2 + b^3)*cosh(x)^4 + a^3 - a^2*b + 4*a*b^2 + 6*b^3 + 30*(a*b^2 + 2*b^3)*
cosh(x)^2)*sinh(x)^4 + 4*(14*(a*b^2 + b^3)*cosh(x)^5 + 10*(a*b^2 + 2*b^3)*c
osh(x)^3 + (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2
*b + 3*a*b^2 + b^3 - 2*(a^3 - 3*a*b^2 - 2*b^3)*cosh(x)^2 + 2*(14*(a*b^2 + b
^3)*cosh(x)^6 + 15*(a*b^2 + 2*b^3)*cosh(x)^4 - a^3 + 3*a*b^2 + 2*b^3 + 3*(a
^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(b^2*cosh(x)^6
+ 6*b^2*cosh(x)*sinh(x)^5 + b^2*sinh(x)^6 + 3*b^2*cosh(x)^4 + 3*(5*b^2*cos
h(x)^2 + b^2)*sinh(x)^4 + 4*(5*b^2*cosh(x)^3 + 3*b^2*cosh(x))*sinh(x)^3 - (
a^2 - 2*a*b - 3*b^2)*cosh(x)^2 + (15*b^2*cosh(x)^4 + 18*b^2*cosh(x)^2 - a^2
+ 2*a*b + 3*b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(3*b^2*cosh(x)^5 + 6*b^
2*cosh(x)^3 - (a^2 - 2*a*b - 3*b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a
+ b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x)
+ sinh(x)^2)) + 4*(2*(a*b^2 + b^3)*cosh(x)^7 + 3*(a*b^2 + 2*b^3)*cosh(x)^5
+ (a^3 - a^2*b + 4*a*b^2 + 6*b^3)*cosh(x)^3 - (a^3 - 3*a*b^2 - 2*b^3)*cosh(
x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20
*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(
x)^6)) + (a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 + a)*sqrt(a + b)*
log(-((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 -
2*a*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a)*sinh(x)^2 + sqrt(2)*(cosh(x)^2
+ 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(a + b))*sqrt(((a + b)*cosh(x)^2 +
(a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) +
4*((a + b)*cosh(x)^3 - a*cosh(x))*sinh(x) + a + b)/(cosh(x)^2 + 2*cosh(x)*s
inh(x) + sinh(x)^2)) - 4*sqrt(2)*(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*
sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a^2 + a*
b)*cosh(x)^2 + 2*(a^2 + a*b)*cosh(x)*sinh(x) + (a^2 + a*b)*sinh(x)^2 + a^2
+ a*b), -1/2*((a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 + a)*sqrt(-a
- b)*arctan(sqrt(2)*(b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + a +
b)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh
(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a*b + b^2)*cosh(x)^4 + 4*(a*b + b
^2)*cosh(x)*sinh(x)^3 + (a*b + b^2)*sinh(x)^4 - (a^2 - a*b - 2*b^2)*cosh(x)
^2 + (6*(a*b + b^2)*cosh(x)^2 - a^2 + a*b + 2*b^2)*sinh(x)^2 + a^2 + 2*a*b
+ b^2 + 2*(2*(a*b + b^2)*cosh(x)^3 - (a^2 - a*b - 2*b^2)*cosh(x))*sinh(x))
+ (a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 + a)*sqrt(-a - b)*arcta
n(sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(-a - b)*sqrt
(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sin
h(x) + sinh(x)^2)))/((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a +
b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)
)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)) + 2*sqrt(2)
*(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 -
2*cosh(x)*sinh(x) + sinh(x)^2)))/((a^2 + a*b)*cosh(x)^2 + 2*(a^2 + a*b)*cos
```

$h(x)*\sinh(x) + (a^2 + a*b)*\sinh(x)^2 + a^2 + a*b]$

Sympy [F]

$$\int \frac{\tanh^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\tanh^2(x)}{\sqrt{a + b \coth^2(x)}} dx$$

[In] `integrate(tanh(x)**2/(a+b*coth(x)**2)**(1/2),x)`

[Out] `Integral(tanh(x)**2/sqrt(a + b*coth(x)**2), x)`

Maxima [F]

$$\int \frac{\tanh^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\tanh(x)^2}{\sqrt{b \coth(x)^2 + a}} dx$$

[In] `integrate(tanh(x)^2/(a+b*coth(x)^2)^(1/2),x, algorithm="maxima")`

[Out] `integrate(tanh(x)^2/sqrt(b*coth(x)^2 + a), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{\tanh^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \text{Exception raised: TypeError}$$

[In] `integrate(tanh(x)^2/(a+b*coth(x)^2)^(1/2),x, algorithm="giac")`

[Out] `Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type`

Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh^2(x)}{\sqrt{a + b \coth^2(x)}} dx = \int \frac{\tanh(x)^2}{\sqrt{b \coth(x)^2 + a}} dx$$

```
[In] int(tanh(x)^2/(a + b*coth(x)^2)^(1/2), x)
```

```
[Out] int(tanh(x)^2/(a + b*coth(x)^2)^(1/2), x)
```

$$3.38 \quad \int \frac{\coth^3(x)}{(a+b \coth^2(x))^{3/2}} dx$$

Optimal result	264
Rubi [A] (verified)	264
Mathematica [A] (verified)	266
Maple [B] (verified)	266
Fricas [B] (verification not implemented)	267
Sympy [F]	269
Maxima [F]	269
Giac [B] (verification not implemented)	269
Mupad [B] (verification not implemented)	270

Optimal result

Integrand size = 17, antiderivative size = 52

$$\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} + \frac{a}{b(a+b)\sqrt{a+b \coth^2(x)}}$$

[Out] $\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2)/(a+b)^{(1/2)})/(a+b)^{(3/2)}+a/b/(a+b)/(a+b*\coth(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3751, 457, 79, 65, 214}

$$\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} + \frac{a}{b(a+b)\sqrt{a+b \coth^2(x)}}$$

[In] $\operatorname{Int}[\operatorname{Coth}[x]^3/(a+b*\operatorname{Coth}[x]^2)^{(3/2)},x]$

[Out] $\operatorname{ArcTanh}[\operatorname{Sqrt}[a+b*\operatorname{Coth}[x]^2]/\operatorname{Sqrt}[a+b]]/(a+b)^{(3/2)}+a/(b*(a+b)*\operatorname{Sqrt}[a+b*\operatorname{Coth}[x]^2])$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) +$

$d*(x^p/b)^n, x, (a + b*x)^{1/p}, x] /; \text{FreeQ}\{a, b, c, d, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 79

$\text{Int}[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^{(n_.)}*((e_.) + (f_.)*(x_))^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[(-b*e - a*f)*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)})/(f*(p + 1)*(c*f - d*e)), x] - \text{Dist}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)), \text{Int}[(c + d*x)^n*(e + f*x)^{(p + 1)}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x\} \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ (\ !\text{LtQ}[n, -1] \ || \ \text{IntegerQ}[p] \ || \ \!(\text{IntegerQ}[n] \ || \ \!(\text{EqQ}[e, 0] \ || \ \!(\text{EqQ}[c, 0] \ || \ \text{LtQ}[p, n]))))$

Rule 214

$\text{Int}[(a_.) + (b_.)*(x_)^2]^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x\} \ \&\& \ \text{NegQ}[a/b]$

Rule 457

$\text{Int}[(x_)^{(m_.)}*((a_.) + (b_.)*(x_)^{(n_.)})^{(p_.)}*((c_.) + (d_.)*(x_)^{(n_.)})^{(q_.)}, x_Symbol] \rightarrow \text{Dist}[1/n, \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)*(a + b*x)^p*(c + d*x)^q}, x], x, x^n], x] /; \text{FreeQ}\{a, b, c, d, m, n, p, q\}, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]$

Rule 3751

$\text{Int}[(d_.)*\text{tan}[(e_.) + (f_.)*(x_)]]^{(m_.)}*((a_.) + (b_.)*((c_.)*\text{tan}[(e_.) + (f_.)*(x_)])^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \text{With}\{\text{ff} = \text{FreeFactors}[\text{Tan}[e + f*x], x]\}, \text{Dist}[c*(\text{ff}/f), \text{Subst}[\text{Int}[(d*\text{ff}*(x/c))^m*((a + b*(\text{ff}*x)^n)^p/(c^2 + \text{ff}^2*x^2)], x], x, c*(\text{Tan}[e + f*x]/\text{ff}), x] /; \text{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x\} \ \&\& \ (\text{IGtQ}[p, 0] \ || \ \text{EqQ}[n, 2] \ || \ \text{EqQ}[n, 4] \ || \ (\text{IntegerQ}[p] \ \&\& \ \text{RationalQ}[n]))$

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst}\left(\int \frac{x^3}{(1-x^2)(a+bx^2)^{3/2}} dx, x, \coth(x)\right) \\ &= \frac{1}{2} \text{Subst}\left(\int \frac{x}{(1-x)(a+bx)^{3/2}} dx, x, \coth^2(x)\right) \\ &= \frac{a}{b(a+b)\sqrt{a+b\coth^2(x)}} + \frac{\text{Subst}\left(\int \frac{1}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x)\right)}{2(a+b)} \end{aligned}$$

$$= \frac{a}{b(a+b)\sqrt{a+b\coth^2(x)}} + \frac{\text{Subst}\left(\int \frac{1}{1+\frac{a}{b}-\frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)}\right)}{b(a+b)}$$

$$= \frac{\text{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} + \frac{a}{b(a+b)\sqrt{a+b\coth^2(x)}}$$

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.00

$$\int \frac{\coth^3(x)}{(a+b\coth^2(x))^{3/2}} dx = \frac{\text{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} + \frac{a}{b(a+b)\sqrt{a+b\coth^2(x)}}$$

[In] Integrate[Coth[x]^3/(a + b*Coth[x]^2)^(3/2), x]

[Out] ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]]/(a + b)^(3/2) + a/(b*(a + b)*Sqrt[a + b*Coth[x]^2])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 286 vs. 2(44) = 88.

Time = 0.10 (sec) , antiderivative size = 287, normalized size of antiderivative = 5.52

method	result
derivativedivides	$\frac{1}{b\sqrt{a+b\coth(x)^2}} - \frac{1}{2(a+b)\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}} - \frac{b(2b(1+\coth(x))-2b)}{(a+b)(4(a+b)b-4b^2)\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}$
default	$\frac{1}{b\sqrt{a+b\coth(x)^2}} - \frac{1}{2(a+b)\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}} - \frac{b(2b(1+\coth(x))-2b)}{(a+b)(4(a+b)b-4b^2)\sqrt{b(1+\coth(x))^2-2b(1+\coth(x))+a+b}}$

[In] int(coth(x)^3/(a+b*coth(x)^2)^(3/2), x, method=_RETURNVERBOSE)

[Out] 1/b/(a+b*coth(x)^2)^(1/2)-1/2/(a+b)/(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)-b/(a+b)*(2*b*(1+coth(x))-2*b)/(4*(a+b)*b-4*b^2)/(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)+1/2/(a+b)^(3/2)*ln((2*a+2*b-2*b*(1+coth(x))+2*(a+b)^(1/2)*(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2))/(1+coth(x)))-1/2/(a+b)/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)+b/(a+b)*(2*b*(coth(x)-1)+2*b)/(4*(a+b)*b-4*b^2)/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)+1/2/(a+b)^(3/2)*

$\ln((2*a+2*b+2*b*(\coth(x)-1)+2*(a+b)^{(1/2)}*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^{(1/2)))/(\coth(x)-1))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 990 vs. 2(44) = 88.

Time = 0.36 (sec) , antiderivative size = 2541, normalized size of antiderivative = 48.87

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{3/2}} dx = \text{Too large to display}$$

[In] integrate(coth(x)^3/(a+b*coth(x)^2)^(3/2),x, algorithm="fricas")

[Out] [1/4*(((a*b + b^2)*cosh(x)^4 + 4*(a*b + b^2)*cosh(x)*sinh(x)^3 + (a*b + b^2)*sinh(x)^4 - 2*(a*b - b^2)*cosh(x)^2 + 2*(3*(a*b + b^2)*cosh(x)^2 - a*b + b^2)*sinh(x)^2 + a*b + b^2 + 4*((a*b + b^2)*cosh(x)^3 - (a*b - b^2)*cosh(x))*sinh(x))*sqrt(a + b)*log(-((a^3 + a^2*b)*cosh(x)^8 + 8*(a^3 + a^2*b)*cosh(x)*sinh(x)^7 + (a^3 + a^2*b)*sinh(x)^8 - 2*(2*a^3 + a^2*b)*cosh(x)^6 - 2*(2*a^3 + a^2*b - 14*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^6 + 4*(14*(a^3 + a^2*b)*cosh(x)^3 - 3*(2*a^3 + a^2*b)*cosh(x))*sinh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^4 + (70*(a^3 + a^2*b)*cosh(x)^4 + 6*a^3 + 4*a^2*b - a*b^2 + b^3 - 30*(2*a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(14*(a^3 + a^2*b)*cosh(x)^5 - 10*(2*a^3 + a^2*b)*cosh(x)^3 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x))*sinh(x)^3 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(2*a^3 + 3*a^2*b - b^3)*cosh(x)^2 + 2*(14*(a^3 + a^2*b)*cosh(x)^6 - 15*(2*a^3 + a^2*b)*cosh(x)^4 - 2*a^3 - 3*a^2*b + b^3 + 3*(6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + sqrt(2)*(a^2*cosh(x)^6 + 6*a^2*cosh(x)*sinh(x)^5 + a^2*sinh(x)^6 - 3*a^2*cosh(x)^4 + 3*(5*a^2*cosh(x)^2 - a^2)*sinh(x)^4 + 4*(5*a^2*cosh(x)^3 - 3*a^2*cosh(x))*sinh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x)^2 + (15*a^2*cosh(x)^4 - 18*a^2*cosh(x)^2 + 3*a^2 + 2*a*b - b^2)*sinh(x)^2 - a^2 - 2*a*b - b^2 + 2*(3*a^2*cosh(x)^5 - 6*a^2*cosh(x)^3 + (3*a^2 + 2*a*b - b^2)*cosh(x))*sinh(x))*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*(2*(a^3 + a^2*b)*cosh(x)^7 - 3*(2*a^3 + a^2*b)*cosh(x)^5 + (6*a^3 + 4*a^2*b - a*b^2 + b^3)*cosh(x)^3 - (2*a^3 + 3*a^2*b - b^3)*cosh(x))*sinh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6)) + ((a*b + b^2)*cosh(x)^4 + 4*(a*b + b^2)*cosh(x)*sinh(x)^3 + (a*b + b^2)*sinh(x)^4 - 2*(a*b - b^2)*cosh(x)^2 + 2*(3*(a*b + b^2)*cosh(x)^2 - a*b + b^2)*sinh(x)^2 + a*b + b^2 + 4*((a*b + b^2)*cosh(x)^3 - (a*b - b^2)*cosh(x))*sinh(x))*sqrt(a + b)*log(((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 + 2*b*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 + b)*sinh(x)^2 + sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 4*((a + b)*cosh(x)^3 +

$$\begin{aligned}
& b \cosh(x) \sinh(x) + a + b) / (\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2) + \\
& 4 \sqrt{2} * ((a^2 + a*b) \cosh(x)^2 + 2*(a^2 + a*b) \cosh(x) \sinh(x) + (a^2 + a \\
& *b) \sinh(x)^2 - a^2 - a*b) \sqrt{((a + b) \cosh(x)^2 + (a + b) \sinh(x)^2 - a \\
& + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2))} / ((a^3*b + 3*a^2*b^2 + 3* \\
& a*b^3 + b^4) \cosh(x)^4 + 4*(a^3*b + 3*a^2*b^2 + 3*a*b^3 + b^4) \cosh(x) \sinh \\
& (x)^3 + (a^3*b + 3*a^2*b^2 + 3*a*b^3 + b^4) \sinh(x)^4 + a^3*b + 3*a^2*b^2 + \\
& 3*a*b^3 + b^4 - 2*(a^3*b + a^2*b^2 - a*b^3 - b^4) \cosh(x)^2 - 2*(a^3*b + a \\
& ^2*b^2 - a*b^3 - b^4 - 3*(a^3*b + 3*a^2*b^2 + 3*a*b^3 + b^4) \cosh(x)^2) \sin \\
& h(x)^2 + 4*((a^3*b + 3*a^2*b^2 + 3*a*b^3 + b^4) \cosh(x)^3 - (a^3*b + a^2*b^ \\
& 2 - a*b^3 - b^4) \cosh(x)) \sinh(x)), -1/2 * (((a*b + b^2) \cosh(x)^4 + 4*(a*b + \\
& b^2) \cosh(x) \sinh(x)^3 + (a*b + b^2) \sinh(x)^4 - 2*(a*b - b^2) \cosh(x)^2 + \\
& 2*(3*(a*b + b^2) \cosh(x)^2 - a*b + b^2) \sinh(x)^2 + a*b + b^2 + 4*((a*b + \\
& b^2) \cosh(x)^3 - (a*b - b^2) \cosh(x)) \sinh(x)) \sqrt{-a - b} \arctan(\sqrt{2} * \\
& (a \cosh(x)^2 + 2*a \cosh(x) \sinh(x) + a \sinh(x)^2 - a - b) \sqrt{-a - b} \sqrt{ \\
& ((a + b) \cosh(x)^2 + (a + b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sin \\
& h(x) + \sinh(x)^2)}) / ((a^2 + a*b) \cosh(x)^4 + 4*(a^2 + a*b) \cosh(x) \sinh(x)^3 \\
& + (a^2 + a*b) \sinh(x)^4 - (2*a^2 + a*b - b^2) \cosh(x)^2 + (6*(a^2 + a*b) \c \\
& osh(x)^2 - 2*a^2 - a*b + b^2) \sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(2*(a^2 + a \\
& *b) \cosh(x)^3 - (2*a^2 + a*b - b^2) \cosh(x)) \sinh(x))) + ((a*b + b^2) \cosh(\\
& x)^4 + 4*(a*b + b^2) \cosh(x) \sinh(x)^3 + (a*b + b^2) \sinh(x)^4 - 2*(a*b - b \\
& ^2) \cosh(x)^2 + 2*(3*(a*b + b^2) \cosh(x)^2 - a*b + b^2) \sinh(x)^2 + a*b + b \\
& ^2 + 4*((a*b + b^2) \cosh(x)^3 - (a*b - b^2) \cosh(x)) \sinh(x)) \sqrt{-a - b} * \\
& \arctan(\sqrt{2} * (\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1) \sqrt{-a - b} \\
& \sqrt{((a + b) \cosh(x)^2 + (a + b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \\
&) \sinh(x) + \sinh(x)^2)}) / ((a + b) \cosh(x)^4 + 4*(a + b) \cosh(x) \sinh(x)^3 + \\
& (a + b) \sinh(x)^4 - 2*(a - b) \cosh(x)^2 + 2*(3*(a + b) \cosh(x)^2 - a + b) \sin \\
& h(x)^2 + 4*((a + b) \cosh(x)^3 - (a - b) \cosh(x)) \sinh(x) + a + b)) - 2 \sqrt{2} * ((a^2 + a*b) \cosh(x)^2 + 2*(a^2 + a*b) \cosh(x) \sinh(x) + (a^2 + a*b) \sinh(x)^2 - a^2 - a*b) \sqrt{((a + b) \cosh(x)^2 + (a + b) \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2))} / ((a^3*b + 3*a^2*b^2 + 3*a*b^3 + b^4) \cosh(x)^4 + 4*(a^3*b + 3*a^2*b^2 + 3*a*b^3 + b^4) \cosh(x) \sinh(x)^3 + (a^3*b + 3*a^2*b^2 + 3*a*b^3 + b^4) \sinh(x)^4 + a^3*b + 3*a^2*b^2 + 3*a*b^3 + b^4 - 2*(a^3*b + a^2*b^2 - a*b^3 - b^4) \cosh(x)^2 - 2*(a^3*b + a^2*b^2 - a*b^3 - b^4 - 3*(a^3*b + 3*a^2*b^2 + 3*a*b^3 + b^4) \cosh(x)^2) \sinh(x)^2 + 4*((a^3*b + 3*a^2*b^2 + 3*a*b^3 + b^4) \cosh(x)^3 - (a^3*b + a^2*b^2 - a*b^3 - b^4) \cosh(x)) \sinh(x)]
\end{aligned}$$

Sympy [F]

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\coth^3(x)}{(a + b \coth^2(x))^{\frac{3}{2}}} dx$$

[In] integrate(coth(x)**3/(a+b*coth(x)**2)**(3/2), x)

[Out] Integral(coth(x)**3/(a + b*coth(x)**2)**(3/2), x)

Maxima [F]

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\coth(x)^3}{(b \coth(x)^2 + a)^{\frac{3}{2}}} dx$$

[In] integrate(coth(x)^3/(a+b*coth(x)^2)^(3/2), x, algorithm="maxima")

[Out] integrate(coth(x)^3/(b*coth(x)^2 + a)^(3/2), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 359 vs. 2(44) = 88.

Time = 0.46 (sec) , antiderivative size = 359, normalized size of antiderivative = 6.90

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{3/2}} dx = \frac{\frac{(a^3+a^2b)e^{(2x)}}{a^3\operatorname{sgn}(e^{(2x)}-1)+2a^2b^2\operatorname{sgn}(e^{(2x)}-1)+ab^3\operatorname{sgn}(e^{(2x)}-1)} - \frac{a^3+a^2b}{a^3\operatorname{sgn}(e^{(2x)}-1)+2a^2b^2\operatorname{sgn}(e^{(2x)}-1)+ab^3\operatorname{sgn}(e^{(2x)}-1)}}{\sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}}$$

$$- \frac{\log\left(\left|\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)\sqrt{a + b} - a + b\right|\right)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

$$- \frac{\log\left(\left|\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)\sqrt{a + b} - a - b\right|\right)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

$$+ \frac{\log\left(\left|-\sqrt{a + be^{(2x)}} + \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b} - \sqrt{a + b}\right|\right)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

[In] integrate(coth(x)^3/(a+b*coth(x)^2)^(3/2), x, algorithm="giac")

[Out] ((a^3 + a^2*b)*e^(2*x)/(a^3*b*sgn(e^(2*x) - 1) + 2*a^2*b^2*sgn(e^(2*x) - 1) + a*b^3*sgn(e^(2*x) - 1)) - (a^3 + a^2*b)/(a^3*b*sgn(e^(2*x) - 1) + 2*a^2*b^2*sgn(e^(2*x) - 1) + a*b^3*sgn(e^(2*x) - 1)))/sqrt(a*e^(4*x) + b*e^(4*x))

```

- 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - 1/2*log(abs((sqrt(a + b)*e^(2*x) - s
qrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*sqrt(a + b)
- a + b))/((a + b)^(3/2)*sgn(e^(2*x) - 1)) - 1/2*log(abs((sqrt(a + b)*e^(2
*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*sqrt
(a + b) - a - b))/((a + b)^(3/2)*sgn(e^(2*x) - 1)) + 1/2*log(abs(-sqrt(a +
b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b
) - sqrt(a + b)))/((a + b)^(3/2)*sgn(e^(2*x) - 1))

```

Mupad [B] (verification not implemented)

Time = 2.77 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.87

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{3/2}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \coth(x)^2 + a}}{\sqrt{a+b}}\right)}{(a + b)^{3/2}} + \frac{a}{(b^2 + a b) \sqrt{b \coth(x)^2 + a}}$$

```
[In] int(coth(x)^3/(a + b*coth(x)^2)^(3/2),x)
```

```
[Out] atanh((a + b*coth(x)^2)^(1/2)/(a + b)^(1/2))/(a + b)^(3/2) + a/((a*b + b^2)
*(a + b*coth(x)^2)^(1/2))
```

3.39 $\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{3/2}} dx$

Optimal result	271
Rubi [A] (verified)	271
Mathematica [B] (verified)	273
Maple [B] (verified)	273
Fricas [B] (verification not implemented)	274
Sympy [F]	275
Maxima [F]	276
Giac [B] (verification not implemented)	276
Mupad [F(-1)]	277

Optimal result

Integrand size = 17, antiderivative size = 53

$$\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{(a+b)^{3/2}} - \frac{\coth(x)}{(a+b)\sqrt{a+b \coth^2(x)}}$$

[Out] $\operatorname{arctanh}(\coth(x) \cdot (a+b)^{1/2} / (a+b \coth(x)^2)^{1/2}) / (a+b)^{3/2} - \coth(x) / (a+b \coth(x)^2)^{1/2}$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3751, 482, 385, 212}

$$\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{(a+b)^{3/2}} - \frac{\coth(x)}{(a+b)\sqrt{a+b \coth^2(x)}}$$

[In] $\operatorname{Int}[\operatorname{Coth}[x]^2 / (a + b \operatorname{Coth}[x]^2)^{3/2}, x]$

[Out] $\operatorname{ArcTanh}[\operatorname{Sqrt}[a+b] \operatorname{Coth}[x] / \operatorname{Sqrt}[a+b \operatorname{Coth}[x]^2]] / (a+b)^{3/2} - \operatorname{Coth}[x] / ((a+b) \operatorname{Sqrt}[a+b \operatorname{Coth}[x]^2])$

Rule 212

$\operatorname{Int}[(a_+ + (b_-) \cdot (x_-)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1 / (\operatorname{Rt}[a, 2] \operatorname{Rt}[-b, 2])) * \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2] \cdot (x / \operatorname{Rt}[a, 2])], x] / ; \operatorname{FreeQ}\{a, b, x\} \ \&\& \operatorname{NegQ}[a/b] \ \&\& \operatorname{Gt}$

Q[a, 0] || LtQ[b, 0])

Rule 385

Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 482

Int[((e_)*(x_)^(m_))*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[e^(n - 1)*(e*x)^(m - n + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(n*(b*c - a*d)*(p + 1))), x] - Dist[e^n/(n*(b*c - a*d)*(p + 1)), Int[(e*x)^(m - n)*(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(m - n + 1) + d*(m + n*(p + q + 1) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[p, -1] && GeQ[n, m - n + 1] && GtQ[m - n + 1, 0] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

Rule 3751

Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{x^2}{(1-x^2)(a+bx^2)^{3/2}} dx, x, \coth(x)\right) \\
 &= -\frac{\coth(x)}{(a+b)\sqrt{a+b\coth^2(x)}} + \frac{\text{Subst}\left(\int \frac{1}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right)}{a+b} \\
 &= -\frac{\coth(x)}{(a+b)\sqrt{a+b\coth^2(x)}} + \frac{\text{Subst}\left(\int \frac{1}{1-(a+b)x^2} dx, x, \frac{\coth(x)}{\sqrt{a+b\coth^2(x)}}\right)}{a+b} \\
 &= \frac{\arctanh\left(\frac{\sqrt{a+b}\coth(x)}{\sqrt{a+b\coth^2(x)}}\right)}{(a+b)^{3/2}} - \frac{\coth(x)}{(a+b)\sqrt{a+b\coth^2(x)}}
 \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 109 vs. $2(53) = 106$.

Time = 1.11 (sec) , antiderivative size = 109, normalized size of antiderivative = 2.06

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \frac{-2(a + b) \coth(x) + \frac{\operatorname{arctanh}\left(\frac{\sqrt{\frac{(a+b) \coth^2(x)}{a}}}{\sqrt{1 + \frac{b \coth^2(x)}{a}}}\right) (-a + b + (a+b) \cosh(2x)) \sqrt{\frac{(a+b) \coth^2(x)}{a}} \operatorname{csch}(2x)}{\sqrt{1 + \frac{b \coth^2(x)}{a}}}}{2(a + b)^2 \sqrt{a + b \coth^2(x)}}$$

[In] Integrate[Coth[x]^2/(a + b*Coth[x]^2)^(3/2), x]

[Out] $(-2*(a + b)*\operatorname{Coth}[x] + (\operatorname{ArcTanh}[\operatorname{Sqrt}[(a + b)*\operatorname{Coth}[x]^2/a]/\operatorname{Sqrt}[1 + (b*\operatorname{Coth}[x]^2)/a]])*(-a + b + (a + b)*\operatorname{Cosh}[2*x])*\operatorname{Sqrt}[(a + b)*\operatorname{Coth}[x]^2/a]*\operatorname{Csch}[x]*\operatorname{Sech}[x])/\operatorname{Sqrt}[1 + (b*\operatorname{Coth}[x]^2)/a])/(2*(a + b)^2*\operatorname{Sqrt}[a + b*\operatorname{Coth}[x]^2])$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 288 vs. $2(45) = 90$.

Time = 0.10 (sec) , antiderivative size = 289, normalized size of antiderivative = 5.45

method	result
derivativedivides	$-\frac{\coth(x)}{a\sqrt{a+b\coth(x)^2}} - \frac{1}{2(a+b)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}} + \frac{b(2b(\coth(x)-1)+2b)}{(a+b)(4(a+b)b-4b^2)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}$
default	$-\frac{\coth(x)}{a\sqrt{a+b\coth(x)^2}} - \frac{1}{2(a+b)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}} + \frac{b(2b(\coth(x)-1)+2b)}{(a+b)(4(a+b)b-4b^2)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}}$

[In] int(coth(x)^2/(a+b*coth(x)^2)^(3/2), x, method=_RETURNVERBOSE)

[Out] $-\operatorname{coth}(x)/a/(a+b*\operatorname{coth}(x)^2)^{(1/2)}-1/2/(a+b)/(b*(\operatorname{coth}(x)-1)^2+2*b*(\operatorname{coth}(x)-1)+a+b)^{(1/2)}+b/(a+b)*(2*b*(\operatorname{coth}(x)-1)+2*b)/(4*(a+b)*b-4*b^2)/(b*(\operatorname{coth}(x)-1)^2+2*b*(\operatorname{coth}(x)-1)+a+b)^{(1/2)}+1/2/(a+b)^{(3/2)}*\ln((2*a+2*b+2*b*(\operatorname{coth}(x)-1)+2*(a+b)^{(1/2)}*(b*(\operatorname{coth}(x)-1)^2+2*b*(\operatorname{coth}(x)-1)+a+b)^{(1/2)})/(\operatorname{coth}(x)-1))+1/2/(a+b)/(b*(1+\operatorname{coth}(x))^2-2*b*(1+\operatorname{coth}(x))+a+b)^{(1/2)}+b/(a+b)*(2*b*(1+\operatorname{coth}(x))-2*b)/(4*(a+b)*b-4*b^2)/(b*(1+\operatorname{coth}(x))^2-2*b*(1+\operatorname{coth}(x))+a+b)^{(1/2)}-1/2/(a+b)^{(3/2)}*\ln((2*a+2*b-2*b*(1+\operatorname{coth}(x))+2*(a+b)^{(1/2)}*(b*(1+\operatorname{coth}(x))^2-2*b*(1+\operatorname{coth}(x))+a+b)^{(1/2)})/(1+\operatorname{coth}(x)))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 860 vs. $2(45) = 90$.

Time = 0.35 (sec) , antiderivative size = 2279, normalized size of antiderivative = 43.00

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \text{Too large to display}$$

[In] integrate(coth(x)^2/(a+b*coth(x)^2)^(3/2),x, algorithm="fricas")

[Out] $\frac{1}{4} \left((a+b) \cosh(x)^4 + 4(a+b) \cosh(x) \sinh(x)^3 + (a+b) \sinh(x)^4 - 2(a-b) \cosh(x)^2 + 2(3(a+b) \cosh(x)^2 - a+b) \sinh(x)^2 + 4((a+b) \cosh(x)^3 - (a-b) \cosh(x)) \sinh(x) + a+b \right) \sqrt{a+b} \log\left((a^2 b^2 + b^3) \cosh(x)^8 + 8(a^2 b^2 + b^3) \cosh(x) \sinh(x)^7 + (a^2 b^2 + b^3) \sinh(x)^8 + 2(a^2 b^2 + 2b^3) \cosh(x)^6 + 2(a^2 b^2 + 2b^3 + 14(a^2 b^2 + b^3) \cosh(x)^2) \sinh(x)^6 + 4(14(a^2 b^2 + b^3) \cosh(x)^3 + 3(a^2 b^2 + 2b^3) \cosh(x)) \sinh(x)^5 + (a^3 - a^2 b + 4a^2 b^2 + 6b^3) \cosh(x)^4 + (70(a^2 b^2 + b^3) \cosh(x)^4 + a^3 - a^2 b + 4a^2 b^2 + 6b^3 + 30(a^2 b^2 + 2b^3) \cosh(x)^2) \sinh(x)^4 + 4(14(a^2 b^2 + b^3) \cosh(x)^5 + 10(a^2 b^2 + 2b^3) \cosh(x)^3 + (a^3 - a^2 b + 4a^2 b^2 + 6b^3) \cosh(x)) \sinh(x)^3 + a^3 + 3a^2 b + 3a^2 b^2 + b^3 - 2(a^3 - 3a^2 b^2 - 2b^3) \cosh(x)^2 + 2(14(a^2 b^2 + b^3) \cosh(x))^6 + 15(a^2 b^2 + 2b^3) \cosh(x)^4 - a^3 + 3a^2 b^2 + 2b^3 + 3(a^3 - a^2 b + 4a^2 b^2 + 6b^3) \cosh(x)^2 \sinh(x)^2 + \sqrt{2} (b^2 \cosh(x)^6 + 6b^2 \cosh(x) \sinh(x)^5 + b^2 \sinh(x)^6 + 3b^2 \cosh(x)^4 + 3(5b^2 \cosh(x)^2 + b^2) \sinh(x)^4 + 4(5b^2 \cosh(x)^3 + 3b^2 \cosh(x)) \sinh(x)^3 - (a^2 - 2ab - 3b^2) \cosh(x)^2 + (15b^2 \cosh(x)^4 + 18b^2 \cosh(x)^2 - a^2 + 2ab + 3b^2) \sinh(x)^2 + a^2 + 2ab + b^2 + 2(3b^2 \cosh(x)^5 + 6b^2 \cosh(x)^3 - (a^2 - 2ab - 3b^2) \cosh(x)) \sinh(x) \right) \sqrt{a+b} \sqrt{\left((a+b) \cosh(x)^2 + (a+b) \sinh(x)^2 - a+b \right) / \left(\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2 \right)} + 4(2(a^2 b^2 + b^3) \cosh(x)^7 + 3(a^2 b^2 + 2b^3) \cosh(x)^5 + (a^3 - a^2 b + 4a^2 b^2 + 6b^3) \cosh(x)^3 - (a^3 - 3a^2 b^2 - 2b^3) \cosh(x)) \sinh(x) / \left(\cosh(x)^6 + 6 \cosh(x)^5 \sinh(x) + 15 \cosh(x)^4 \sinh(x)^2 + 20 \cosh(x)^3 \sinh(x)^3 + 15 \cosh(x)^2 \sinh(x)^4 + 6 \cosh(x) \sinh(x)^5 + \sinh(x)^6 \right) + \left((a+b) \cosh(x)^4 + 4(a+b) \cosh(x) \sinh(x)^3 + (a+b) \sinh(x)^4 - 2(a-b) \cosh(x)^2 + 2(3(a+b) \cosh(x)^2 - a+b) \sinh(x)^2 + 4((a+b) \cosh(x)^3 - (a-b) \cosh(x)) \sinh(x) + a+b \right) \sqrt{a+b} \log\left(-((a+b) \cosh(x))^4 + 4(a+b) \cosh(x) \sinh(x)^3 + (a+b) \sinh(x)^4 - 2a \cosh(x)^2 + 2(3(a+b) \cosh(x)^2 - a) \sinh(x)^2 + \sqrt{2} (\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1) \sqrt{a+b} \sqrt{\left((a+b) \cosh(x)^2 + (a+b) \sinh(x)^2 - a+b \right) / \left(\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2 \right)} + 4((a+b) \cosh(x)^3 - a \cosh(x)) \sinh(x) + a+b \right) / \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 \right) - 4 \sqrt{2} \left((a+b) \cosh(x)^2 + 2(a+b) \cosh(x) \sinh(x) + (a+b) \sinh(x)^2 + a+b \right) \sqrt{\left((a+b) \cosh(x)^2 + (a+b) \sinh(x)^2 - a+b \right) / \left(\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2 \right)} \right) / \left((a^3 + 3a^2 b + 3a^2 b^2 + b^3) \cosh(x)^4 + 4(a^3 + 3a^2 b + 3a^2 b^2 + b^3) \cosh(x) \sinh(x)^3 + (a^3 + 3a^2 b \right.$

```

b + 3*a*b^2 + b^3)*sinh(x)^4 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 + a^2
*b - a*b^2 - b^3)*cosh(x)^2 - 2*(a^3 + a^2*b - a*b^2 - b^3 - 3*(a^3 + 3*a^2
*b + 3*a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + 4*((a^3 + 3*a^2*b + 3*a*b^2 + b^
3)*cosh(x)^3 - (a^3 + a^2*b - a*b^2 - b^3)*cosh(x))*sinh(x)), -1/2*((a + b
)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*c
osh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3
- (a - b)*cosh(x))*sinh(x) + a + b)*sqrt(-a - b)*arctan(sqrt(2)*(b*cosh(x)
^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + a + b)*sqrt(-a - b)*sqrt(((a + b)*
cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sin
h(x)^2)))/((a*b + b^2)*cosh(x)^4 + 4*(a*b + b^2)*cosh(x)*sinh(x)^3 + (a*b +
b^2)*sinh(x)^4 - (a^2 - a*b - 2*b^2)*cosh(x)^2 + (6*(a*b + b^2)*cosh(x)^2 -
a^2 + a*b + 2*b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(2*(a*b + b^2)*cosh(x
)^3 - (a^2 - a*b - 2*b^2)*cosh(x))*sinh(x))) + ((a + b)*cosh(x)^4 + 4*(a +
b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a +
b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*s
inh(x) + a + b)*sqrt(-a - b)*arctan(sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x)
+ sinh(x)^2 - 1)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 -
a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a + b)*cosh(x)^4 + 4
*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3
*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh
(x))*sinh(x) + a + b)) + 2*sqrt(2)*((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*s
inh(x) + (a + b)*sinh(x)^2 + a + b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(
x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a^3 + 3*a^2*b
+ 3*a*b^2 + b^3)*cosh(x)^4 + 4*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*cosh(x)*sin
h(x)^3 + (a^3 + 3*a^2*b + 3*a*b^2 + b^3)*sinh(x)^4 + a^3 + 3*a^2*b + 3*a*b^
2 + b^3 - 2*(a^3 + a^2*b - a*b^2 - b^3)*cosh(x)^2 - 2*(a^3 + a^2*b - a*b^2
- b^3 - 3*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + 4*((a^3 +
3*a^2*b + 3*a*b^2 + b^3)*cosh(x)^3 - (a^3 + a^2*b - a*b^2 - b^3)*cosh(x))*s
inh(x))]

```

Sympy [F]

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\coth^2(x)}{(a + b \coth^2(x))^{\frac{3}{2}}} dx$$

[In] integrate(coth(x)**2/(a+b*coth(x)**2)**(3/2), x)

[Out] Integral(coth(x)**2/(a + b*coth(x)**2)**(3/2), x)

Maxima [F]

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\coth(x)^2}{(b \coth(x)^2 + a)^{3/2}} dx$$

[In] integrate(coth(x)^2/(a+b*coth(x)^2)^(3/2),x, algorithm="maxima")

[Out] integrate(coth(x)^2/(b*coth(x)^2 + a)^(3/2), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 363 vs. 2(45) = 90.

Time = 0.46 (sec) , antiderivative size = 363, normalized size of antiderivative = 6.85

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \frac{\frac{(a^2b+ab^2)e^{(2x)}}{a^3\operatorname{sgn}(e^{(2x)}-1)+2a^2b^2\operatorname{sgn}(e^{(2x)}-1)+ab^3\operatorname{sgn}(e^{(2x)}-1)} + \frac{a^2b+ab^2}{a^3\operatorname{sgn}(e^{(2x)}-1)+2a^2b^2\operatorname{sgn}(e^{(2x)}-1)+ab^3\operatorname{sgn}(e^{(2x)}-1)}}{\sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}} \log \left(\left| \left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b} \right) \sqrt{a + b} - a + b \right| \right) - \frac{2(a + b)^{\frac{3}{2}} \operatorname{sgn}(e^{(2x)} - 1)}{\log \left(\left| \left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b} \right) \sqrt{a + b} - a - b \right| \right)} + \frac{2(a + b)^{\frac{3}{2}} \operatorname{sgn}(e^{(2x)} - 1)}{\log \left(\left| -\sqrt{a + be^{(2x)}} + \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b} - \sqrt{a + b} \right| \right)}$$

[In] integrate(coth(x)^2/(a+b*coth(x)^2)^(3/2),x, algorithm="giac")

[Out] -((a^2*b + a*b^2)*e^(2*x)/(a^3*b*sgn(e^(2*x) - 1) + 2*a^2*b^2*sgn(e^(2*x) - 1) + a*b^3*sgn(e^(2*x) - 1)) + (a^2*b + a*b^2)/(a^3*b*sgn(e^(2*x) - 1) + 2*a^2*b^2*sgn(e^(2*x) - 1) + a*b^3*sgn(e^(2*x) - 1)))/sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - 1/2*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*sqrt(a + b) - a + b))/((a + b)^(3/2)*sgn(e^(2*x) - 1)) + 1/2*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*sqrt(a + b) - a - b))/((a + b)^(3/2)*sgn(e^(2*x) - 1)) - 1/2*log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - sqrt(a + b)))/((a + b)^(3/2)*sgn(e^(2*x) - 1))

Mupad [F(-1)]

Timed out.

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\coth(x)^2}{(b \coth(x)^2 + a)^{3/2}} dx$$

```
[In] int(coth(x)^2/(a + b*coth(x)^2)^(3/2), x)
```

```
[Out] int(coth(x)^2/(a + b*coth(x)^2)^(3/2), x)
```

$$3.40 \quad \int \frac{\coth(x)}{(a+b \coth^2(x))^{3/2}} dx$$

Optimal result	278
Rubi [A] (verified)	278
Mathematica [C] (verified)	280
Maple [B] (verified)	280
Fricas [B] (verification not implemented)	281
Sympy [F]	282
Maxima [F]	283
Giac [B] (verification not implemented)	283
Mupad [B] (verification not implemented)	284

Optimal result

Integrand size = 15, antiderivative size = 49

$$\int \frac{\coth(x)}{(a+b \coth^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} - \frac{1}{(a+b)\sqrt{a+b \coth^2(x)}}$$

[Out] $\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2)/(a+b)^{(1/2)})/(a+b)^{(3/2)}-1/(a+b)/(a+b*\coth(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3751, 455, 53, 65, 214}

$$\int \frac{\coth(x)}{(a+b \coth^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} - \frac{1}{(a+b)\sqrt{a+b \coth^2(x)}}$$

[In] $\operatorname{Int}[\operatorname{Coth}[x]/(a+b*\operatorname{Coth}[x]^2)^{(3/2)}, x]$

[Out] $\operatorname{ArcTanh}[\operatorname{Sqrt}[a+b*\operatorname{Coth}[x]^2]/\operatorname{Sqrt}[a+b]]/(a+b)^{(3/2)}-1/((a+b)*\operatorname{Sqrt}[a+b*\operatorname{Coth}[x]^2])$

Rule 53

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a+b*x)^{(m+1)*((c+d*x)^{(n+1))/(b*c-a*d)*(m+1))}, x] - \operatorname{Dist}[d*(($

```

m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && !(LtQ
[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && I
ntLinearQ[a, b, c, d, m, n, x]

```

Rule 65

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

```

Rule 214

```

Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

```

Rule 455

```

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.
), x_Symbol] := Dist[1/n, Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x
] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n +
1, 0]

```

Rule 3751

```

Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{x}{(1-x^2)(a+bx^2)^{3/2}} dx, x, \coth(x)\right) \\
&= \frac{1}{2} \text{Subst}\left(\int \frac{1}{(1-x)(a+bx)^{3/2}} dx, x, \coth^2(x)\right) \\
&= -\frac{1}{(a+b)\sqrt{a+b\coth^2(x)}} + \frac{\text{Subst}\left(\int \frac{1}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x)\right)}{2(a+b)}
\end{aligned}$$

$$= -\frac{1}{(a+b)\sqrt{a+b\coth^2(x)}} + \frac{\text{Subst}\left(\int \frac{1}{1+\frac{a}{b}-\frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)}\right)}{b(a+b)}$$

$$= \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} - \frac{1}{(a+b)\sqrt{a+b\coth^2(x)}}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.05 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.84

$$\int \frac{\coth(x)}{(a+b\coth^2(x))^{3/2}} dx = -\frac{\operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b\coth^2(x)}{a+b}\right)}{(a+b)\sqrt{a+b\coth^2(x)}}$$

[In] Integrate[Coth[x]/(a + b*Coth[x]^2)^(3/2), x]

[Out] -(Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Coth[x]^2)/(a + b)]/((a + b)*Sqrt[a + b*Coth[x]^2]))

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 272 vs. 2(41) = 82.

Time = 0.08 (sec) , antiderivative size = 273, normalized size of antiderivative = 5.57

method	result
derivativedivides	$-\frac{1}{2(a+b)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}} + \frac{b(2b(\coth(x)-1)+2b)}{(a+b)(4(a+b)b-4b^2)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}} + \dots$
default	$-\frac{1}{2(a+b)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}} + \frac{b(2b(\coth(x)-1)+2b)}{(a+b)(4(a+b)b-4b^2)\sqrt{b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b}} + \dots$

[In] int(coth(x)/(a+b*coth(x)^2)^(3/2), x, method=_RETURNVERBOSE)

[Out] -1/2/(a+b)/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)+b/(a+b)*(2*b*(coth(x)-1)+2*b)/(4*(a+b)*b-4*b^2)/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2)+1/2/(a+b)^(3/2)*ln((2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(1/2)*(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(coth(x)-1))-1/2/(a+b)/(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)-b/(a+b)*(2*b*(1+coth(x))-2*b)/(4*(a+b)*b-4*b^2)/(b*(1+coth(x))^2-2*b*(1+coth(x))+a+b)^(1/2)

$(x)^{-2-2*b*(1+\coth(x))+a+b}^{(1/2)+1/2/(a+b)^{(3/2)}*\ln((2*a+2*b-2*b*(1+\coth(x)))+2*(a+b)^{(1/2)}*(b*(1+\coth(x))^{-2-2*b*(1+\coth(x))+a+b}^{(1/2)})/(1+\coth(x)))}$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 869 vs. $2(41) = 82$.

Time = 0.35 (sec) , antiderivative size = 2299, normalized size of antiderivative = 46.92

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{3/2}} dx = \text{Too large to display}$$

[In] integrate(coth(x)/(a+b*coth(x)^2)^(3/2),x, algorithm="fricas")

[Out] $\frac{1}{4}(((a + b)\cosh(x)^4 + 4(a + b)\cosh(x)\sinh(x)^3 + (a + b)\sinh(x)^4 - 2(a - b)\cosh(x)^2 + 2(3(a + b)\cosh(x)^2 - a + b)\sinh(x)^2 + 4((a + b)\cosh(x)^3 - (a - b)\cosh(x))\sinh(x) + a + b)\sqrt{a + b}\log(-((a^3 + a^2b)\cosh(x)^8 + 8(a^3 + a^2b)\cosh(x)\sinh(x)^7 + (a^3 + a^2b)\sinh(x)^8 - 2(2a^3 + a^2b)\cosh(x)^6 - 2(2a^3 + a^2b - 14(a^3 + a^2b)\cosh(x)^2)\sinh(x)^6 + 4(14(a^3 + a^2b)\cosh(x)^3 - 3(2a^3 + a^2b)\cosh(x))\sinh(x)^5 + (6a^3 + 4a^2b - ab^2 + b^3)\cosh(x)^4 + (70(a^3 + a^2b)\cosh(x)^4 + 6a^3 + 4a^2b - ab^2 + b^3 - 30(2a^3 + a^2b)\cosh(x)^2)\sinh(x)^4 + 4(14(a^3 + a^2b)\cosh(x)^5 - 10(2a^3 + a^2b)\cosh(x)^3 + (6a^3 + 4a^2b - ab^2 + b^3)\cosh(x))\sinh(x)^3 + a^3 + 3a^2b + 3ab^2 + b^3 - 2(2a^3 + 3a^2b - b^3)\cosh(x)^2 + 2(14(a^3 + a^2b)\cosh(x)^6 - 15(2a^3 + a^2b)\cosh(x)^4 - 2a^3 - 3a^2b + b^3 + 3(6a^3 + 4a^2b - ab^2 + b^3)\cosh(x)^2)\sinh(x)^2 + \sqrt{2}(a^2\cosh(x)^6 + 6a^2\cosh(x)\sinh(x)^5 + a^2\sinh(x)^6 - 3a^2\cosh(x)^4 + 3(5a^2\cosh(x)^2 - a^2)\sinh(x)^4 + 4(5a^2\cosh(x)^3 - 3a^2\cosh(x))\sinh(x)^3 + (3a^2 + 2ab - b^2)\cosh(x)^2 + (15a^2\cosh(x)^4 - 18a^2\cosh(x)^2 + 3a^2 + 2ab - b^2)\sinh(x)^2 - a^2 - 2ab - b^2 + 2(3a^2\cosh(x)^5 - 6a^2\cosh(x)^3 + (3a^2 + 2ab - b^2)\cosh(x))\sinh(x))\sqrt{a + b}\sqrt{((a + b)\cosh(x)^2 + (a + b)\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2\cosh(x)\sinh(x) + \sinh(x)^2))} + 4(2(a^3 + a^2b)\cosh(x)^7 - 3(2a^3 + a^2b)\cosh(x)^5 + (6a^3 + 4a^2b - ab^2 + b^3)\cosh(x)^3 - (2a^3 + 3a^2b - b^3)\cosh(x))\sinh(x)/(\cosh(x)^6 + 6\cosh(x)^5\sinh(x) + 15\cosh(x)^4\sinh(x)^2 + 20\cosh(x)^3\sinh(x)^3 + 15\cosh(x)^2\sinh(x)^4 + 6\cosh(x)\sinh(x)^5 + \sinh(x)^6)) + ((a + b)\cosh(x)^4 + 4(a + b)\cosh(x)\sinh(x)^3 + (a + b)\sinh(x)^4 - 2(a - b)\cosh(x)^2 + 2(3(a + b)\cosh(x)^2 - a + b)\sinh(x)^2 + 4((a + b)\cosh(x)^3 - (a - b)\cosh(x))\sinh(x) + a + b)\sqrt{a + b}\log(((a + b)\cosh(x)^4 + 4(a + b)\cosh(x)\sinh(x)^3 + (a + b)\sinh(x)^4 + 2b\cosh(x)^2 + 2(3(a + b)\cosh(x)^2 + b)\sinh(x)^2 + \sqrt{2}(\cosh(x)^2 + 2\cosh(x)\sinh(x) + \sinh(x)^2 + 1)\sqrt{a + b}\sqrt{((a + b)\cosh(x)^2 + (a + b)\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2\cosh(x)\sinh(x) + \sinh(x)^2))} + 4((a + b)\cosh(x)^3 + b\cosh(x))\sinh(x) + a + b)/(\cosh(x)^2 + 2\cosh(x)\sinh(x) + \sinh(x)^2))$

```

- 4*sqrt(2)*((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 - a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*cosh(x)^4 + 4*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*cosh(x)*sinh(x)^3 + (a^3 + 3*a^2*b + 3*a*b^2 + b^3)*sinh(x)^4 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 + a^2*b - a*b^2 - b^3)*cosh(x)^2 - 2*(a^3 + a^2*b - a*b^2 - b^3 - 3*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + 4*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*cosh(x)^3 - (a^3 + a^2*b - a*b^2 - b^3)*cosh(x))*sinh(x)), -1/2*((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)*sqrt(-a - b)*arctan(sqrt(2)*(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 - a - b)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a^2 + a*b)*cosh(x)^4 + 4*(a^2 + a*b)*cosh(x)*sinh(x)^3 + (a^2 + a*b)*sinh(x)^4 - (2*a^2 + a*b - b^2)*cosh(x)^2 + (6*(a^2 + a*b)*cosh(x)^2 - 2*a^2 - a*b + b^2)*sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(2*(a^2 + a*b)*cosh(x)^3 - (2*a^2 + a*b - b^2)*cosh(x))*sinh(x))) + ((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)*sqrt(-a - b)*arctan(sqrt(2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)) + 2*sqrt(2)*((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 - a - b)*sqrt(((a + b)*cosh(x)^2 + (a + b)*sinh(x)^2 - a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*cosh(x)^4 + 4*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*cosh(x)*sinh(x)^3 + (a^3 + 3*a^2*b + 3*a*b^2 + b^3)*sinh(x)^4 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 + a^2*b - a*b^2 - b^3)*cosh(x)^2 - 2*(a^3 + a^2*b - a*b^2 - b^3 - 3*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*cosh(x)^2)*sinh(x)^2 + 4*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*cosh(x)^3 - (a^3 + a^2*b - a*b^2 - b^3)*cosh(x))*sinh(x))]

```

Sympy [F]

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\coth(x)}{(a + b \coth^2(x))^{\frac{3}{2}}} dx$$

[In] integrate(coth(x)/(a+b*coth(x)**2)**(3/2),x)

[Out] Integral(coth(x)/(a + b*coth(x)**2)**(3/2), x)

Maxima [F]

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\coth(x)}{(b \coth(x)^2 + a)^{3/2}} dx$$

[In] integrate(coth(x)/(a+b*coth(x)^2)^(3/2),x, algorithm="maxima")

[Out] integrate(coth(x)/(b*coth(x)^2 + a)^(3/2), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 364 vs. 2(41) = 82.

Time = 0.47 (sec) , antiderivative size = 364, normalized size of antiderivative = 7.43

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{3/2}} dx =$$

$$\frac{(a^2b+ab^2)e^{(2x)}}{a^3\operatorname{sgn}(e^{(2x)}-1)+2a^2b^2\operatorname{sgn}(e^{(2x)}-1)+ab^3\operatorname{sgn}(e^{(2x)}-1)} - \frac{a^2b+ab^2}{a^3\operatorname{sgn}(e^{(2x)}-1)+2a^2b^2\operatorname{sgn}(e^{(2x)}-1)+ab^3\operatorname{sgn}(e^{(2x)}-1)}$$

$$\frac{\sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}}{\log\left(\left|\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)\sqrt{a + b} - a + b\right|\right)}$$

$$\frac{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}{\log\left(\left|\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)\sqrt{a + b} - a - b\right|\right)}$$

$$\frac{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}{\log\left(\left|-\sqrt{a + be^{(2x)}} + \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b} - \sqrt{a + b}\right|\right)}$$

$$+ \frac{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

[In] integrate(coth(x)/(a+b*coth(x)^2)^(3/2),x, algorithm="giac")

[Out] -((a^2*b + a*b^2)*e^(2*x)/(a^3*b*sgn(e^(2*x) - 1) + 2*a^2*b^2*sgn(e^(2*x) - 1) + a*b^3*sgn(e^(2*x) - 1)) - (a^2*b + a*b^2)/(a^3*b*sgn(e^(2*x) - 1) + 2*a^2*b^2*sgn(e^(2*x) - 1) + a*b^3*sgn(e^(2*x) - 1)))/sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - 1/2*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*sqrt(a + b) - a + b))/((a + b)^(3/2)*sgn(e^(2*x) - 1)) - 1/2*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*sqrt(a + b) - a - b))/((a + b)^(3/2)*sgn(e^(2*x) - 1)) + 1/2*log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - sqrt(a + b)))/((a + b)^(3/2)*sgn(e^(2*x) - 1))

Mupad [B] (verification not implemented)

Time = 2.61 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.84

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{3/2}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \coth(x)^2 + a}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} - \frac{1}{(a+b) \sqrt{b \coth(x)^2 + a}}$$

[In] `int(coth(x)/(a + b*coth(x)^2)^(3/2),x)`

[Out] `atanh((a + b*coth(x)^2)^(1/2)/(a + b)^(1/2))/(a + b)^(3/2) - 1/((a + b)*(a + b*coth(x)^2)^(1/2))`

$$3.41 \quad \int \frac{\tanh(x)}{(a+b \coth^2(x))^{3/2}} dx$$

Optimal result	285
Rubi [A] (verified)	285
Mathematica [C] (verified)	287
Maple [F]	288
Fricas [B] (verification not implemented)	288
Sympy [F]	288
Maxima [F]	288
Giac [F(-2)]	289
Mupad [F(-1)]	289

Optimal result

Integrand size = 15, antiderivative size = 78

$$\int \frac{\tanh(x)}{(a+b \coth^2(x))^{3/2}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a}}\right)}{a^{3/2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} + \frac{b}{a(a+b)\sqrt{a+b \coth^2(x)}}$$

[Out] $-\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2))/a^{(1/2)})/a^{(3/2)}+\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2))/(a+b)^{(1/2)})/(a+b)^{(3/2)}+b/a/(a+b)/(a+b*\coth(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3751, 457, 87, 162, 65, 214}

$$\int \frac{\tanh(x)}{(a+b \coth^2(x))^{3/2}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a}}\right)}{a^{3/2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{3/2}} + \frac{b}{a(a+b)\sqrt{a+b \coth^2(x)}}$$

[In] Int[Tanh[x]/(a + b*Coth[x]^2)^(3/2), x]

[Out] -(ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a]]/a^(3/2)) + ArcTanh[Sqrt[a + b*Coth[x]^2]/Sqrt[a + b]]/(a + b)^(3/2) + b/(a*(a + b)*Sqrt[a + b*Coth[x]^2])

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 87

Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[f*(e + f*x)^(p + 1)/((p + 1)*(b*e - a*f)*(d*e - c*f)), x] + Dist[1/((b*e - a*f)*(d*e - c*f)), Int[(b*d*e - b*c*f - a*d*f - b*d*f*x)*(e + f*x)^(p + 1)/((a + b*x)*(c + d*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[p, -1]

Rule 162

Int[(((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)))/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Dist[(b*g - a*h)/(b*c - a*d), Int[(e + f*x)^p/(a + b*x), x], x] - Dist[(d*g - c*h)/(b*c - a*d), Int[(e + f*x)^p/(c + d*x), x], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x]

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 457

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]

Rule 3751

Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration

alQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst} \left(\int \frac{1}{x(1-x^2)(a+bx^2)^{3/2}} dx, x, \coth(x) \right) \\
 &= \frac{1}{2} \text{Subst} \left(\int \frac{1}{(1-x)x(a+bx)^{3/2}} dx, x, \coth^2(x) \right) \\
 &= \frac{b}{a(a+b)\sqrt{a+b\coth^2(x)}} - \frac{\text{Subst} \left(\int \frac{-a-b+bx}{(1-x)x\sqrt{a+bx}} dx, x, \coth^2(x) \right)}{2a(a+b)} \\
 &= \frac{b}{a(a+b)\sqrt{a+b\coth^2(x)}} + \frac{\text{Subst} \left(\int \frac{1}{x\sqrt{a+bx}} dx, x, \coth^2(x) \right)}{2a} \\
 &\quad + \frac{\text{Subst} \left(\int \frac{1}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x) \right)}{2(a+b)} \\
 &= \frac{b}{a(a+b)\sqrt{a+b\coth^2(x)}} + \frac{\text{Subst} \left(\int \frac{1}{-\frac{a}{b} + \frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)} \right)}{ab} \\
 &\quad + \frac{\text{Subst} \left(\int \frac{1}{1 + \frac{a}{b} - \frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)} \right)}{b(a+b)} \\
 &= -\frac{\text{arctanh} \left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a}} \right)}{a^{3/2}} + \frac{\text{arctanh} \left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}} \right)}{(a+b)^{3/2}} + \frac{b}{a(a+b)\sqrt{a+b\coth^2(x)}}
 \end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.06 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.90

$$\int \frac{\tanh(x)}{(a+b\coth^2(x))^{3/2}} dx = \frac{-a \text{Hypergeometric2F1} \left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b\coth^2(x)}{a+b} \right) + (a+b) \text{Hypergeometric2F1} \left(-\frac{1}{2}, 1, \frac{1}{2}, 1 + \frac{b\coth^2(x)}{a} \right)}{a(a+b)\sqrt{a+b\coth^2(x)}}$$

[In] Integrate[Tanh[x]/(a + b*Coth[x]^2)^(3/2), x]

[Out] $(-a \text{Hypergeometric2F1}[-1/2, 1, 1/2, (a + b \text{Coth}[x]^2)/(a + b)]) + (a + b) \text{Hypergeometric2F1}[-1/2, 1, 1/2, 1 + (b \text{Coth}[x]^2)/a]) / (a(a + b) \text{Sqrt}[a + b \text{Coth}[x]^2])$

Maple [F]

$$\int \frac{\tanh(x)}{(a + b \coth(x)^2)^{\frac{3}{2}}} dx$$

[In] `int(tanh(x)/(a+b*coth(x)^2)^(3/2),x)`

[Out] `int(tanh(x)/(a+b*coth(x)^2)^(3/2),x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1437 vs. $2(64) = 128$.

Time = 0.55 (sec) , antiderivative size = 6991, normalized size of antiderivative = 89.63

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{3/2}} dx = \text{Too large to display}$$

[In] `integrate(tanh(x)/(a+b*coth(x)^2)^(3/2),x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\tanh(x)}{(a + b \coth^2(x))^{\frac{3}{2}}} dx$$

[In] `integrate(tanh(x)/(a+b*coth(x)**2)**(3/2),x)`

[Out] `Integral(tanh(x)/(a + b*coth(x)**2)**(3/2), x)`

Maxima [F]

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\tanh(x)}{(b \coth(x)^2 + a)^{\frac{3}{2}}} dx$$

[In] `integrate(tanh(x)/(a+b*coth(x)^2)^(3/2),x, algorithm="maxima")`

[Out] `integrate(tanh(x)/(b*coth(x)^2 + a)^(3/2), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{3/2}} dx = \text{Exception raised: TypeError}$$

[In] integrate(tanh(x)/(a+b*coth(x)^2)^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type

Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\tanh(x)}{(b \coth(x)^2 + a)^{3/2}} dx$$

[In] int(tanh(x)/(a + b*coth(x)^2)^(3/2),x)

[Out] int(tanh(x)/(a + b*coth(x)^2)^(3/2), x)

$$3.42 \quad \int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{3/2}} dx$$

Optimal result	290
Rubi [A] (verified)	290
Mathematica [C] (warning: unable to verify)	293
Maple [F]	293
Fricas [B] (verification not implemented)	293
Sympy [F]	296
Maxima [F]	296
Giac [B] (verification not implemented)	297
Mupad [F(-1)]	298

Optimal result

Integrand size = 17, antiderivative size = 85

$$\int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{(a+b)^{3/2}} + \frac{b \tanh(x)}{a(a+b)\sqrt{a+b \coth^2(x)}} - \frac{(a+2b)\sqrt{a+b \coth^2(x)} \tanh(x)}{a^2(a+b)}$$

[Out] $\operatorname{arctanh}(\coth(x) \cdot (a+b)^{(1/2)} / (a+b \cdot \coth(x)^2)^{(1/2)}) / (a+b)^{(3/2)} + b \cdot \tanh(x) / a / (a+b) / (a+b \cdot \coth(x)^2)^{(1/2)} - (a+2 \cdot b) \cdot (a+b \cdot \coth(x)^2)^{(1/2)} \cdot \tanh(x) / a^2 / (a+b)$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3751, 483, 597, 12, 385, 212}

$$\int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{3/2}} dx = -\frac{(a+2b) \tanh(x) \sqrt{a+b \coth^2(x)}}{a^2(a+b)} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{(a+b)^{3/2}} + \frac{b \tanh(x)}{a(a+b)\sqrt{a+b \coth^2(x)}}$$

[In] $\operatorname{Int}[\operatorname{Tanh}[x]^2 / (a + b \cdot \operatorname{Coth}[x]^2)^{(3/2)}, x]$

[Out] ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]]/(a + b)^(3/2) + (b*Tanh[x])/(a*(a + b)*Sqrt[a + b*Coth[x]^2]) - ((a + 2*b)*Sqrt[a + b*Coth[x]^2]*Tanh[x])/(a^2*(a + b))

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 385

Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 483

Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-b)*(e*x)^(m + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*e*n*(b*c - a*d)*(p + 1))), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(e*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*b*(m + 1) + n*(b*c - a*d)*(p + 1) + d*b*(m + n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, m, q}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[p, -1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

Rule 597

Int[((g_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] := Simp[e*(g*x)^(m + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*c*g*(m + 1))), x] + Dist[1/(a*c*g^n*(m + 1)), Int[(g*x)^(m + n)*(a + b*x^n)^p*(c + d*x^n)^q*Simp[a*f*c*(m + 1) - e*(b*c + a*d)*(m + n + 1) - e*n*(b*c*p + a*d*q) - b*e*d*(m + n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p, q}, x] && IGtQ[n, 0] && LtQ[m, -1]

Rule 3751

Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n}

, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{1}{x^2(1-x^2)(a+bx^2)^{3/2}} dx, x, \coth(x)\right) \\
&= \frac{b \tanh(x)}{a(a+b)\sqrt{a+b \coth^2(x)}} - \frac{\text{Subst}\left(\int \frac{-a-2b+2bx^2}{x^2(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right)}{a(a+b)} \\
&= \frac{b \tanh(x)}{a(a+b)\sqrt{a+b \coth^2(x)}} - \frac{(a+2b)\sqrt{a+b \coth^2(x)} \tanh(x)}{a^2(a+b)} \\
&\quad + \frac{\text{Subst}\left(\int \frac{a^2}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right)}{a^2(a+b)} \\
&= \frac{b \tanh(x)}{a(a+b)\sqrt{a+b \coth^2(x)}} - \frac{(a+2b)\sqrt{a+b \coth^2(x)} \tanh(x)}{a^2(a+b)} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right)}{a+b} \\
&= \frac{b \tanh(x)}{a(a+b)\sqrt{a+b \coth^2(x)}} - \frac{(a+2b)\sqrt{a+b \coth^2(x)} \tanh(x)}{a^2(a+b)} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{1-(a+b)x^2} dx, x, \frac{\coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{a+b} \\
&= \frac{\text{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{(a+b)^{3/2}} + \frac{b \tanh(x)}{a(a+b)\sqrt{a+b \coth^2(x)}} - \frac{(a+2b)\sqrt{a+b \coth^2(x)} \tanh(x)}{a^2(a+b)}
\end{aligned}$$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 7.93 (sec) , antiderivative size = 260, normalized size of antiderivative = 3.06

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \frac{\sinh^2(x) \tanh(x) \left(\frac{8(a+b) \cosh^2(x)(2a^2+5ab \coth^2(x)+3b^2 \coth^4(x)) \operatorname{Hypergeometric2F1}\left(2, 2, \frac{7}{2}, \frac{(a+b) \cosh^2(x)}{a}\right)}{15a^3} - \left(8(a+b) \left((-1)a \coth(x) - I b \coth(x)^3 \right)^2 \operatorname{HypergeometricPFQ}\left[\{2, 2, 2\}, \{1, 7/2\}, \frac{(a+b) \cosh^2(x)}{a} \frac{\sinh^2(x)}{(15a^3)} - \frac{(3a^2 + 12ab \coth^2(x) + 8b^2 \coth^4(x)) (\operatorname{ArcSin}[\sqrt{\frac{(a+b) \cosh^2(x)}{a}}] (-a - b \coth(x)^2) - a \operatorname{Csch}[x]^2 \sqrt{-((a+b) \cosh^2(x) (a + b \coth(x)^2) \sinh^2(x) / a^2)})}]{\frac{(a+b) \cosh^2(x) (a + b \coth(x)^2) \sinh^2(x)}{a^2}} \right)}{a \sqrt{a + b \coth(x)^2}} \right)}{a \sqrt{a + b \coth(x)^2}}$$

[In] Integrate[Tanh[x]^2/(a + b*Coth[x]^2)^(3/2), x]

[Out] (Sinh[x]^2*Tanh[x]*((8*(a + b)*Cosh[x]^2*(2*a^2 + 5*a*b*Coth[x]^2 + 3*b^2*Coth[x]^4)*Hypergeometric2F1[2, 2, 7/2, ((a + b)*Cosh[x]^2)/a])/(15*a^3) - (8*(a + b)*((-I)*a*Coth[x] - I*b*Coth[x]^3)^2*HypergeometricPFQ[{2, 2, 2}, {1, 7/2}, ((a + b)*Cosh[x]^2)/a]*Sinh[x]^2)/(15*a^3) - ((3*a^2 + 12*a*b*Coth[x]^2 + 8*b^2*Coth[x]^4)*(ArcSin[Sqrt[((a + b)*Cosh[x]^2)/a]]*(-a - b*Coth[x]^2) - a*Csch[x]^2*sqrt[-((a + b)*Cosh[x]^2*(a + b*Coth[x]^2)*Sinh[x]^2)/a^2]))*Tanh[x]^2)/(a^2*(a + b)*sqrt[-((a + b)*Cosh[x]^2*(a + b*Coth[x]^2)*Sinh[x]^2)/a^2])))/(a*sqrt[a + b*Coth[x]^2])

Maple [F]

$$\int \frac{\tanh(x)^2}{(a + b \coth(x)^2)^{3/2}} dx$$

[In] int(tanh(x)^2/(a+b*coth(x)^2)^(3/2), x)

[Out] int(tanh(x)^2/(a+b*coth(x)^2)^(3/2), x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1686 vs. 2(75) = 150.

Time = 0.51 (sec) , antiderivative size = 3931, normalized size of antiderivative = 46.25

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \text{Too large to display}$$

[In] integrate(tanh(x)^2/(a+b*coth(x)^2)^(3/2), x, algorithm="fricas")

[Out] [1/4*((a^3 + a^2*b)*cosh(x)^6 + 6*(a^3 + a^2*b)*cosh(x)*sinh(x)^5 + (a^3 + a^2*b)*sinh(x)^6 - (a^3 - 3*a^2*b)*cosh(x)^4 - (a^3 - 3*a^2*b - 15*(a^3 + a^2*b)*cosh(x)^2)*sinh(x)^4 + 4*(5*(a^3 + a^2*b)*cosh(x)^3 - (a^3 - 3*a^2*b

$$\begin{aligned}
&) * \cosh(x) * \sinh(x)^3 + a^3 + a^2 * b - (a^3 - 3 * a^2 * b) * \cosh(x)^2 + (15 * (a^3 + \\
& a^2 * b) * \cosh(x)^4 - a^3 + 3 * a^2 * b - 6 * (a^3 - 3 * a^2 * b) * \cosh(x)^2) * \sinh(x)^2 \\
& + 2 * (3 * (a^3 + a^2 * b) * \cosh(x)^5 - 2 * (a^3 - 3 * a^2 * b) * \cosh(x)^3 - (a^3 - 3 * a^2 * \\
& b) * \cosh(x)) * \sinh(x) * \sqrt{a + b} * \log(((a * b^2 + b^3) * \cosh(x)^8 + 8 * (a * b^2 + \\
& b^3) * \cosh(x) * \sinh(x)^7 + (a * b^2 + b^3) * \sinh(x)^8 + 2 * (a * b^2 + 2 * b^3) * \cosh(x) \\
& x)^6 + 2 * (a * b^2 + 2 * b^3 + 14 * (a * b^2 + b^3) * \cosh(x)^2) * \sinh(x)^6 + 4 * (14 * (a * \\
& b^2 + b^3) * \cosh(x)^3 + 3 * (a * b^2 + 2 * b^3) * \cosh(x)) * \sinh(x)^5 + (a^3 - a^2 * b \\
& + 4 * a * b^2 + 6 * b^3) * \cosh(x)^4 + (70 * (a * b^2 + b^3) * \cosh(x)^4 + a^3 - a^2 * b + \\
& 4 * a * b^2 + 6 * b^3 + 30 * (a * b^2 + 2 * b^3) * \cosh(x)^2) * \sinh(x)^4 + 4 * (14 * (a * b^2 + \\
& b^3) * \cosh(x)^5 + 10 * (a * b^2 + 2 * b^3) * \cosh(x)^3 + (a^3 - a^2 * b + 4 * a * b^2 + 6 * \\
& b^3) * \cosh(x)) * \sinh(x)^3 + a^3 + 3 * a^2 * b + 3 * a * b^2 + b^3 - 2 * (a^3 - 3 * a * b^2 \\
& - 2 * b^3) * \cosh(x)^2 + 2 * (14 * (a * b^2 + b^3) * \cosh(x)^6 + 15 * (a * b^2 + 2 * b^3) * \cosh(x) \\
& h(x)^4 - a^3 + 3 * a * b^2 + 2 * b^3 + 3 * (a^3 - a^2 * b + 4 * a * b^2 + 6 * b^3) * \cosh(x)^2) * \sinh(x)^2 + \sqrt{2} * (b^2 * \cosh(x)^6 + 6 * b^2 * \cosh(x) * \sinh(x)^5 + b^2 * \sinh(x) \\
& x)^6 + 3 * b^2 * \cosh(x)^4 + 3 * (5 * b^2 * \cosh(x)^2 + b^2) * \sinh(x)^4 + 4 * (5 * b^2 * \cosh(x)^3 + 3 * b^2 * \cosh(x)) * \sinh(x)^3 - (a^2 - 2 * a * b - 3 * b^2) * \cosh(x)^2 + (15 * b \\
& ^2 * \cosh(x)^4 + 18 * b^2 * \cosh(x)^2 - a^2 + 2 * a * b + 3 * b^2) * \sinh(x)^2 + a^2 + 2 * \\
& a * b + b^2 + 2 * (3 * b^2 * \cosh(x)^5 + 6 * b^2 * \cosh(x)^3 - (a^2 - 2 * a * b - 3 * b^2) * \cosh(x)) * \sinh(x) * \sqrt{a + b} * \sqrt{((a + b) * \cosh(x)^2 + (a + b) * \sinh(x)^2 - a \\
& + b) / (\cosh(x)^2 - 2 * \cosh(x) * \sinh(x) + \sinh(x)^2)} + 4 * (2 * (a * b^2 + b^3) * \cosh(x) \\
& h(x)^7 + 3 * (a * b^2 + 2 * b^3) * \cosh(x)^5 + (a^3 - a^2 * b + 4 * a * b^2 + 6 * b^3) * \cosh(x) \\
& x)^3 - (a^3 - 3 * a * b^2 - 2 * b^3) * \cosh(x)) * \sinh(x) / (\cosh(x)^6 + 6 * \cosh(x)^5 * \\
& \sinh(x) + 15 * \cosh(x)^4 * \sinh(x)^2 + 20 * \cosh(x)^3 * \sinh(x)^3 + 15 * \cosh(x)^2 * \sinh(x) \\
& h(x)^4 + 6 * \cosh(x) * \sinh(x)^5 + \sinh(x)^6)) + ((a^3 + a^2 * b) * \cosh(x)^6 + 6 * \\
& (a^3 + a^2 * b) * \cosh(x) * \sinh(x)^5 + (a^3 + a^2 * b) * \sinh(x)^6 - (a^3 - 3 * a^2 * b) \\
& * \cosh(x)^4 - (a^3 - 3 * a^2 * b - 15 * (a^3 + a^2 * b) * \cosh(x)^2) * \sinh(x)^4 + 4 * (5 * \\
& (a^3 + a^2 * b) * \cosh(x)^3 - (a^3 - 3 * a^2 * b) * \cosh(x)) * \sinh(x)^3 + a^3 + a^2 * b \\
& - (a^3 - 3 * a^2 * b) * \cosh(x)^2 + (15 * (a^3 + a^2 * b) * \cosh(x)^4 - a^3 + 3 * a^2 * b - \\
& 6 * (a^3 - 3 * a^2 * b) * \cosh(x)^2) * \sinh(x)^2 + 2 * (3 * (a^3 + a^2 * b) * \cosh(x)^5 - 2 * \\
& (a^3 - 3 * a^2 * b) * \cosh(x)^3 - (a^3 - 3 * a^2 * b) * \cosh(x)) * \sinh(x) * \sqrt{a + b} * \log(-((a + b) * \cosh(x)^4 + 4 * (a + b) * \cosh(x) * \sinh(x)^3 + (a + b) * \sinh(x)^4 - \\
& 2 * a * \cosh(x)^2 + 2 * (3 * (a + b) * \cosh(x)^2 - a) * \sinh(x)^2 + \sqrt{2} * (\cosh(x)^2 \\
& + 2 * \cosh(x) * \sinh(x) + \sinh(x)^2 - 1) * \sqrt{a + b} * \sqrt{((a + b) * \cosh(x)^2 + \\
& (a + b) * \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 * \cosh(x) * \sinh(x) + \sinh(x)^2)} + 4 \\
& * ((a + b) * \cosh(x)^3 - a * \cosh(x)) * \sinh(x) + a + b) / (\cosh(x)^2 + 2 * \cosh(x) * \sinh(x) + \sinh(x)^2)) - 4 * \sqrt{2} * ((a^3 + 3 * a^2 * b + 4 * a * b^2 + 2 * b^3) * \cosh(x)^4 \\
& + 4 * (a^3 + 3 * a^2 * b + 4 * a * b^2 + 2 * b^3) * \cosh(x) * \sinh(x)^3 + (a^3 + 3 * a^2 * b \\
& + 4 * a * b^2 + 2 * b^3) * \sinh(x)^4 + a^3 + 3 * a^2 * b + 4 * a * b^2 + 2 * b^3 - 2 * (a^3 + a \\
& ^2 * b - 2 * a * b^2 - 2 * b^3) * \cosh(x)^2 - 2 * (a^3 + a^2 * b - 2 * a * b^2 - 2 * b^3 - 3 * (a \\
& ^3 + 3 * a^2 * b + 4 * a * b^2 + 2 * b^3) * \cosh(x)^2) * \sinh(x)^2 + 4 * ((a^3 + 3 * a^2 * b + \\
& 4 * a * b^2 + 2 * b^3) * \cosh(x)^3 - (a^3 + a^2 * b - 2 * a * b^2 - 2 * b^3) * \cosh(x)) * \sinh(x) \\
& x) * \sqrt{((a + b) * \cosh(x)^2 + (a + b) * \sinh(x)^2 - a + b) / (\cosh(x)^2 - 2 * \cosh(x) * \sinh(x) + \sinh(x)^2))} / ((a^5 + 3 * a^4 * b + 3 * a^3 * b^2 + a^2 * b^3) * \cosh(x)^6 \\
& + 6 * (a^5 + 3 * a^4 * b + 3 * a^3 * b^2 + a^2 * b^3) * \cosh(x) * \sinh(x)^5 + (a^5 + 3 * a^4 * \\
& 4 * b + 3 * a^3 * b^2 + a^2 * b^3) * \sinh(x)^6 + a^5 + 3 * a^4 * b + 3 * a^3 * b^2 + a^2 * b^3
\end{aligned}$$

$$\begin{aligned}
& - (a^5 - a^4*b - 5*a^3*b^2 - 3*a^2*b^3)*\cosh(x)^4 - (a^5 - a^4*b - 5*a^3*b^2 - 3*a^2*b^3 - 15*(a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3)*\cosh(x)^2)*\sinh(x)^4 \\
& + 4*(5*(a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3)*\cosh(x)^3 - (a^5 - a^4*b - 5*a^3*b^2 - 3*a^2*b^3)*\cosh(x))*\sinh(x)^3 - (a^5 - a^4*b - 5*a^3*b^2 - 3*a^2*b^3)*\cosh(x)^2 \\
& - (a^5 - a^4*b - 5*a^3*b^2 - 3*a^2*b^3 - 15*(a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3)*\cosh(x)^4 + 6*(a^5 - a^4*b - 5*a^3*b^2 - 3*a^2*b^3)*\cosh(x)^2)*\sinh(x)^2 \\
& + 2*(3*(a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3)*\cosh(x)^5 - 2*(a^5 - a^4*b - 5*a^3*b^2 - 3*a^2*b^3)*\cosh(x)^3 - (a^5 - a^4*b - 5*a^3*b^2 - 3*a^2*b^3)*\cosh(x))*\sinh(x)), \\
& -1/2*((a^3 + a^2*b)*\cosh(x)^6 + 6*(a^3 + a^2*b)*\cosh(x)*\sinh(x)^5 + (a^3 + a^2*b)*\sinh(x)^6 - (a^3 - 3*a^2*b)*\cosh(x)^4 - (a^3 - 3*a^2*b - 15*(a^3 + a^2*b)*\cosh(x)^2)*\sinh(x)^4 \\
& + 4*(5*(a^3 + a^2*b)*\cosh(x)^3 - (a^3 - 3*a^2*b)*\cosh(x))*\sinh(x)^3 + a^3 + a^2*b - (a^3 - 3*a^2*b)*\cosh(x)^2 + (15*(a^3 + a^2*b)*\cosh(x)^4 - a^3 + 3*a^2*b - 6*(a^3 - 3*a^2*b)*\cosh(x)^2)*\sinh(x)^2 \\
& + 2*(3*(a^3 + a^2*b)*\cosh(x)^5 - 2*(a^3 - 3*a^2*b)*\cosh(x)^3 - (a^3 - 3*a^2*b)*\cosh(x))*\sinh(x))*\sqrt{-a - b}*\arctan(\sqrt{2}*(b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 + a + b)*\sqrt{-a - b}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)})/((a*b + b^2)*\cosh(x)^4 + 4*(a*b + b^2)*\cosh(x)*\sinh(x)^3 + (a*b + b^2)*\sinh(x)^4 - (a^2 - a*b - 2*b^2)*\cosh(x)^2 + (6*(a*b + b^2)*\cosh(x)^2 - a^2 + a*b + 2*b^2)*\sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(2*(a*b + b^2)*\cosh(x)^3 - (a^2 - a*b - 2*b^2)*\cosh(x))*\sinh(x))) + ((a^3 + a^2*b)*\cosh(x)^6 + 6*(a^3 + a^2*b)*\cosh(x)*\sinh(x)^5 + (a^3 + a^2*b)*\sinh(x)^6 - (a^3 - 3*a^2*b)*\cosh(x)^4 - (a^3 - 3*a^2*b - 15*(a^3 + a^2*b)*\cosh(x)^2)*\sinh(x)^4 + 4*(5*(a^3 + a^2*b)*\cosh(x)^3 - (a^3 - 3*a^2*b)*\cosh(x))*\sinh(x)^3 + a^3 + a^2*b - (a^3 - 3*a^2*b)*\cosh(x)^2 + (15*(a^3 + a^2*b)*\cosh(x)^4 - a^3 + 3*a^2*b - 6*(a^3 - 3*a^2*b)*\cosh(x)^2)*\sinh(x)^2 + 2*(3*(a^3 + a^2*b)*\cosh(x)^5 - 2*(a^3 - 3*a^2*b)*\cosh(x)^3 - (a^3 - 3*a^2*b)*\cosh(x))*\sinh(x))*\sqrt{-a - b}*\arctan(\sqrt{2}*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 - 1)*\sqrt{-a - b}*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)})/((a + b)*\cosh(x)^4 + 4*(a + b)*\cosh(x)*\sinh(x)^3 + (a + b)*\sinh(x)^4 - 2*(a - b)*\cosh(x)^2 + 2*(3*(a + b)*\cosh(x)^2 - a + b)*\sinh(x)^2 + 4*((a + b)*\cosh(x)^3 - (a - b)*\cosh(x))*\sinh(x) + a + b)) + 2*\sqrt{2}*((a^3 + 3*a^2*b + 4*a*b^2 + 2*b^3)*\cosh(x)^4 + 4*(a^3 + 3*a^2*b + 4*a*b^2 + 2*b^3)*\cosh(x)*\sinh(x)^3 + (a^3 + 3*a^2*b + 4*a*b^2 + 2*b^3)*\sinh(x)^4 + a^3 + 3*a^2*b + 4*a*b^2 + 2*b^3 - 2*(a^3 + a^2*b - 2*a*b^2 - 2*b^3)*\cosh(x)^2 - 2*(a^3 + a^2*b - 2*a*b^2 - 2*b^3 - 3*(a^3 + 3*a^2*b + 4*a*b^2 + 2*b^3)*\cosh(x)^2)*\sinh(x)^2 + 4*((a^3 + 3*a^2*b + 4*a*b^2 + 2*b^3)*\cosh(x)^3 - (a^3 + a^2*b - 2*a*b^2 - 2*b^3)*\cosh(x))*\sinh(x))*\sqrt{((a + b)*\cosh(x)^2 + (a + b)*\sinh(x)^2 - a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)})/((a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3)*\cosh(x)^6 + 6*(a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3)*\cosh(x)*\sinh(x)^5 + (a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3)*\sinh(x)^6 + a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3 - (a^5 - a^4*b - 5*a^3*b^2 - 3*a^2*b^3)*\cosh(x)^4 - (a^5 - a^4*b - 5*a^3*b^2 - 3*a^2*b^3 - 15*(a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3)*\cosh(x)^2)*\sinh(x)^4 + 4*(5*(a^5 + 3*a^4*b + 3*a^3*b^2 + a^2*b^3)*\cosh(x)^3 - (a^5 - a^4*b - 5*
\end{aligned}$$

$a^3b^2 - 3a^2b^3) \cosh(x) \sinh(x)^3 - (a^5 - a^4b - 5a^3b^2 - 3a^2b^3) \cosh(x)^2 - (a^5 - a^4b - 5a^3b^2 - 3a^2b^3 - 15(a^5 + 3a^4b + 3a^3b^2 + a^2b^3) \cosh(x)^4 + 6(a^5 - a^4b - 5a^3b^2 - 3a^2b^3) \cosh(x)^2) \sinh(x)^2 + 2(3(a^5 + 3a^4b + 3a^3b^2 + a^2b^3) \cosh(x)^5 - 2(a^5 - a^4b - 5a^3b^2 - 3a^2b^3) \cosh(x)^3 - (a^5 - a^4b - 5a^3b^2 - 3a^2b^3) \cosh(x) \sinh(x))]$

Sympy [F]

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{\frac{3}{2}}} dx$$

[In] integrate(tanh(x)**2/(a+b*coth(x)**2)**(3/2),x)

[Out] Integral(tanh(x)**2/(a + b*coth(x)**2)**(3/2), x)

Maxima [F]

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\tanh(x)^2}{(b \coth(x)^2 + a)^{\frac{3}{2}}} dx$$

[In] integrate(tanh(x)^2/(a+b*coth(x)^2)^(3/2),x, algorithm="maxima")

[Out] integrate(tanh(x)^2/(b*coth(x)^2 + a)^(3/2), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 540 vs. 2(75) = 150.

Time = 0.60 (sec) , antiderivative size = 540, normalized size of antiderivative = 6.35

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{3/2}} dx =$$

$$\frac{\frac{(a^2b^3+ab^4)e^{(2x)}}{a^5b\operatorname{sgn}(e^{(2x)}-1)+2a^4b^2\operatorname{sgn}(e^{(2x)}-1)+a^3b^3\operatorname{sgn}(e^{(2x)}-1)} + \frac{a^2b^3+ab^4}{a^5b\operatorname{sgn}(e^{(2x)}-1)+2a^4b^2\operatorname{sgn}(e^{(2x)}-1)+a^3b^3\operatorname{sgn}(e^{(2x)}-1)}}{\sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}}$$

$$\frac{\log\left(\left|\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)\sqrt{a + b} - a + b\right|\right)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

$$+ \frac{\log\left(\left|\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)\sqrt{a + b} - a - b\right|\right)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

$$\frac{\log\left(\left|-\sqrt{a + be^{(2x)}} + \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b} - \sqrt{a + b}\right|\right)}{2(a + b)^{\frac{3}{2}}\operatorname{sgn}(e^{(2x)} - 1)}$$

$$\frac{4\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)}{\left(\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)^2 + 2\left(\sqrt{a + be^{(2x)}} - \sqrt{ae^{(4x)} + be^{(4x)} - 2ae^{(2x)} + 2be^{(2x)} + a + b}\right)\sqrt{a + b} - 3a + b\right)}$$

[In] integrate(tanh(x)^2/(a+b*coth(x)^2)^(3/2),x, algorithm="giac")

[Out] -((a^2*b^3 + a*b^4)*e^(2*x)/(a^5*b*sgn(e^(2*x) - 1) + 2*a^4*b^2*sgn(e^(2*x) - 1) + a^3*b^3*sgn(e^(2*x) - 1)) + (a^2*b^3 + a*b^4)/(a^5*b*sgn(e^(2*x) - 1) + 2*a^4*b^2*sgn(e^(2*x) - 1) + a^3*b^3*sgn(e^(2*x) - 1)))/sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - 1/2*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*sqrt(a + b) - a + b))/((a + b)^(3/2)*sgn(e^(2*x) - 1)) + 1/2*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*sqrt(a + b) - a - b))/((a + b)^(3/2)*sgn(e^(2*x) - 1)) - 1/2*log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - sqrt(a + b)))/((a + b)^(3/2)*sgn(e^(2*x) - 1)) - 4*(sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b) - sqrt(a + b))/(((sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))^2 + 2*(sqrt(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*sqrt(a + b) - 3*a + b)*a*sgn(e^(2*x) - 1))

Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{3/2}} dx = \int \frac{\tanh(x)^2}{(b \coth(x)^2 + a)^{3/2}} dx$$

```
[In] int(tanh(x)^2/(a + b*coth(x)^2)^(3/2),x)
```

```
[Out] int(tanh(x)^2/(a + b*coth(x)^2)^(3/2), x)
```

3.43 $\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{5/2}} dx$

Optimal result	299
Rubi [A] (verified)	299
Mathematica [C] (verified)	301
Maple [B] (verified)	302
Fricas [B] (verification not implemented)	302
Sympy [F]	303
Maxima [F]	303
Giac [B] (verification not implemented)	303
Mupad [B] (verification not implemented)	304

Optimal result

Integrand size = 17, antiderivative size = 74

$$\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{5/2}} + \frac{a}{3b(a+b)(a+b \coth^2(x))^{3/2}} - \frac{1}{(a+b)^2 \sqrt{a+b \coth^2(x)}}$$

[Out] $\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2)}/(a+b)^{(1/2)})/(a+b)^{(5/2)}+1/3*a/b/(a+b)/(a+b*\coth(x)^2)^{(3/2)}-1/(a+b)^2/(a+b*\coth(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3751, 457, 79, 53, 65, 214}

$$\int \frac{\coth^3(x)}{(a+b \coth^2(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{5/2}} + \frac{a}{3b(a+b)(a+b \coth^2(x))^{3/2}} - \frac{1}{(a+b)^2 \sqrt{a+b \coth^2(x)}}$$

[In] $\operatorname{Int}[\operatorname{Coth}[x]^3/(a+b*\operatorname{Coth}[x]^2)^{(5/2)},x]$

[Out] $\operatorname{ArcTanh}[\operatorname{Sqrt}[a+b*\operatorname{Coth}[x]^2]/\operatorname{Sqrt}[a+b]]/(a+b)^{(5/2)}+a/(3*b*(a+b)*(a+b*\operatorname{Coth}[x]^2)^{(3/2)})-1/((a+b)^2*\operatorname{Sqrt}[a+b*\operatorname{Coth}[x]^2])$

Rule 53

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((
m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && !(LtQ[
n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && I
ntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[
b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 79

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/
(f*(p + 1)*(c*f - d*e))), x] - Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c
*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)), Int[(c + d*x)^n*(e + f*x)^(p + 1), x]
, x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && ( !LtQ[n, -1] || I
ntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))
))
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 457

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q.
_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p
*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[
b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
```

alQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{x^3}{(1-x^2)(a+bx^2)^{5/2}} dx, x, \coth(x)\right) \\
 &= \frac{1}{2} \text{Subst}\left(\int \frac{x}{(1-x)(a+bx)^{5/2}} dx, x, \coth^2(x)\right) \\
 &= \frac{a}{3b(a+b)(a+b\coth^2(x))^{3/2}} + \frac{\text{Subst}\left(\int \frac{1}{(1-x)(a+bx)^{3/2}} dx, x, \coth^2(x)\right)}{2(a+b)} \\
 &= \frac{a}{3b(a+b)(a+b\coth^2(x))^{3/2}} - \frac{1}{(a+b)^2\sqrt{a+b\coth^2(x)}} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{1}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x)\right)}{2(a+b)^2} \\
 &= \frac{a}{3b(a+b)(a+b\coth^2(x))^{3/2}} - \frac{1}{(a+b)^2\sqrt{a+b\coth^2(x)}} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{1}{1+\frac{a}{b}-\frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)}\right)}{b(a+b)^2} \\
 &= \frac{\text{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{5/2}} + \frac{a}{3b(a+b)(a+b\coth^2(x))^{3/2}} - \frac{1}{(a+b)^2\sqrt{a+b\coth^2(x)}}
 \end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.12 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.85

$$\int \frac{\coth^3(x)}{(a+b\coth^2(x))^{5/2}} dx = \frac{a(a+b) - 3b(a+b\coth^2(x)) \text{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b\coth^2(x)}{a+b}\right)}{3b(a+b)^2(a+b\coth^2(x))^{3/2}}$$

[In] Integrate[Coth[x]^3/(a + b*Coth[x]^2)^(5/2), x]

[Out] (a*(a + b) - 3*b*(a + b*Coth[x]^2)*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Coth[x]^2)/(a + b)])/(3*b*(a + b)^2*(a + b*Coth[x]^2)^(3/2))

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 434 vs. $2(62) = 124$.

Time = 0.10 (sec) , antiderivative size = 435, normalized size of antiderivative = 5.88

method	result
derivativedivides	$\frac{1}{3b(a+b\coth(x)^2)^{3/2}} - \frac{1}{6(a+b)(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{3/2}} + \frac{b\coth(x)}{6(a+b)a(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{3/2}}$
default	$\frac{1}{3b(a+b\coth(x)^2)^{3/2}} - \frac{1}{6(a+b)(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{3/2}} + \frac{b\coth(x)}{6(a+b)a(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{3/2}}$

[In] `int(coth(x)^3/(a+b*coth(x)^2)^(5/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{3} \frac{b}{(a+b\coth(x)^2)^{3/2}} - \frac{1}{6} \frac{1}{(a+b)(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{3/2}} + \frac{1}{6} \frac{b\coth(x)}{(a+b)a(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{3/2}}$
 $+ \frac{1}{3} \frac{b}{(a+b)a^2(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{1/2}} \coth(x) - \frac{1}{2} \frac{1}{(a+b)^2} \frac{1}{(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{1/2}} + \frac{1}{2} \frac{1}{(a+b)^2} \frac{1}{a(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{1/2}} * b\coth(x) + \frac{1}{2} \frac{1}{(a+b)^{5/2}} * \ln\left(\frac{(2a+2b+2b(\coth(x)-1)+2(a+b)^{1/2})(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{1/2}}{(a+b)(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{1/2}}\right) - \frac{1}{6} \frac{1}{(a+b)(b(1+\coth(x))^2+2b(1+\coth(x))+a+b)^{3/2}} - \frac{1}{6} \frac{b}{(a+b)a} \frac{1}{(b(1+\coth(x))^2+2b(1+\coth(x))+a+b)^{3/2}} \coth(x) - \frac{1}{3} \frac{b}{(a+b)a^2} \frac{1}{(b(1+\coth(x))^2+2b(1+\coth(x))+a+b)^{1/2}} \coth(x) - \frac{1}{2} \frac{1}{(a+b)^2} \frac{1}{(b(1+\coth(x))^2+2b(1+\coth(x))+a+b)^{1/2}} - \frac{1}{2} \frac{1}{(a+b)^2} \frac{1}{a(b(1+\coth(x))^2+2b(1+\coth(x))+a+b)^{1/2}} * b\coth(x) + \frac{1}{2} \frac{1}{(a+b)^{5/2}} * \ln\left(\frac{(2a+2b-2b(1+\coth(x))+2(a+b)^{1/2})(b(1+\coth(x))^2+2b(1+\coth(x))+a+b)^{1/2}}{(1+\coth(x))(b(1+\coth(x))^2+2b(1+\coth(x))+a+b)^{1/2}}\right)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2964 vs. $2(62) = 124$.

Time = 0.69 (sec) , antiderivative size = 6560, normalized size of antiderivative = 88.65

$$\int \frac{\coth^3(x)}{(a+b\coth^2(x))^{5/2}} dx = \text{Too large to display}$$

[In] `integrate(coth(x)^3/(a+b*coth(x)^2)^(5/2),x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\coth^3(x)}{(a + b \coth^2(x))^{\frac{5}{2}}} dx$$

[In] integrate(coth(x)**3/(a+b*coth(x)**2)**(5/2),x)

[Out] Integral(coth(x)**3/(a + b*coth(x)**2)**(5/2), x)

Maxima [F]

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\coth(x)^3}{(b \coth(x)^2 + a)^{\frac{5}{2}}} dx$$

[In] integrate(coth(x)^3/(a+b*coth(x)^2)^(5/2),x, algorithm="maxima")

[Out] integrate(coth(x)^3/(b*coth(x)^2 + a)^(5/2), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 951 vs. 2(62) = 124.

Time = 0.60 (sec) , antiderivative size = 951, normalized size of antiderivative = 12.85

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{5/2}} dx = \text{Too large to display}$$

[In] integrate(coth(x)^3/(a+b*coth(x)^2)^(5/2),x, algorithm="giac")

[Out] 1/3*(((a^8*b*sgn(e^(2*x) - 1) + 2*a^7*b^2*sgn(e^(2*x) - 1) - 5*a^6*b^3*sgn(e^(2*x) - 1) - 20*a^5*b^4*sgn(e^(2*x) - 1) - 25*a^4*b^5*sgn(e^(2*x) - 1) - 14*a^3*b^6*sgn(e^(2*x) - 1) - 3*a^2*b^7*sgn(e^(2*x) - 1))*e^(2*x)/(a^8*b^2 + 6*a^7*b^3 + 15*a^6*b^4 + 20*a^5*b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*b^8) - 3*(a^8*b*sgn(e^(2*x) - 1) + 2*a^7*b^2*sgn(e^(2*x) - 1) - a^6*b^3*sgn(e^(2*x) - 1) - 4*a^5*b^4*sgn(e^(2*x) - 1) - a^4*b^5*sgn(e^(2*x) - 1) + 2*a^3*b^6*sgn(e^(2*x) - 1) + a^2*b^7*sgn(e^(2*x) - 1)))/(a^8*b^2 + 6*a^7*b^3 + 15*a^6*b^4 + 20*a^5*b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*b^8))*e^(2*x) + 3*(a^8*b*sgn(e^(2*x) - 1) + 2*a^7*b^2*sgn(e^(2*x) - 1) - a^6*b^3*sgn(e^(2*x) - 1) - 4*a^5*b^4*sgn(e^(2*x) - 1) - a^4*b^5*sgn(e^(2*x) - 1) + 2*a^3*b^6*sgn(e^(2*x) - 1) + a^2*b^7*sgn(e^(2*x) - 1)))/(a^8*b^2 + 6*a^7*b^3 + 15*a^6*b^4 + 20*a^5*b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*b^8))*e^(2*x) - (a^8*b*sgn(e^(2*x) - 1) + 2*a^7*b^2*sgn(e^(2*x) - 1) - 5*a^6*b^3*sgn(e^(2*x) - 1) - 20*a^5*b^4*

```

sgn(e^(2*x) - 1) - 25*a^4*b^5*sgn(e^(2*x) - 1) - 14*a^3*b^6*sgn(e^(2*x) - 1)
) - 3*a^2*b^7*sgn(e^(2*x) - 1))/(a^8*b^2 + 6*a^7*b^3 + 15*a^6*b^4 + 20*a^5*
b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*b^8))/(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*
x) + 2*b*e^(2*x) + a + b)^(3/2) - 1/2*log(abs((sqrt(a + b)*e^(2*x) - sqrt(a
*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) + a + b))*sqrt(a + b) - a
+ b))/((a^2 + 2*a*b + b^2)*sqrt(a + b)*sgn(e^(2*x) - 1)) - 1/2*log(abs((sqr
t(a + b)*e^(2*x) - sqrt(a*e^(4*x) + b*e^(4*x) - 2*a*e^(2*x) + 2*b*e^(2*x) +
a + b))*sqrt(a + b) - a - b))/((a^2 + 2*a*b + b^2)*sqrt(a + b)*sgn(e^(2*x)
- 1)) + 1/2*log(abs(-sqrt(a + b)*e^(2*x) + sqrt(a*e^(4*x) + b*e^(4*x) - 2*
a*e^(2*x) + 2*b*e^(2*x) + a + b) - sqrt(a + b)))/((a^2 + 2*a*b + b^2)*sqrt(
a + b)*sgn(e^(2*x) - 1))

```

Mupad [B] (verification not implemented)

Time = 4.25 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.11

$$\int \frac{\coth^3(x)}{(a + b \coth^2(x))^{5/2}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \coth(x)^2 + a} (2a^2 + 4ab + 2b^2)}{2(a+b)^{5/2}}\right)}{(a+b)^{5/2}} + \frac{\frac{a}{3(a+b)} - \frac{b(b \coth(x)^2 + a)}{(a+b)^2}}{b(b \coth(x)^2 + a)^{3/2}}$$

[In] int(coth(x)^3/(a + b*coth(x)^2)^(5/2),x)

[Out] atanh(((a + b*coth(x)^2)^(1/2)*(4*a*b + 2*a^2 + 2*b^2))/(2*(a + b)^(5/2)))/
(a + b)^(5/2) + (a/(3*(a + b)) - (b*(a + b*coth(x)^2))/(a + b)^2)/(b*(a + b
*coth(x)^2)^(3/2))

$$3.44 \quad \int \frac{\coth^2(x)}{(a+b \coth^2(x))^{5/2}} dx$$

Optimal result	305
Rubi [A] (verified)	305
Mathematica [C] (warning: unable to verify)	307
Maple [B] (verified)	308
Fricas [B] (verification not implemented)	309
Sympy [F]	309
Maxima [F]	309
Giac [B] (verification not implemented)	310
Mupad [F(-1)]	310

Optimal result

Integrand size = 17, antiderivative size = 88

$$\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{(a+b)^{5/2}} - \frac{\coth(x)}{3(a+b)(a+b \coth^2(x))^{3/2}} - \frac{(2a-b) \coth(x)}{3a(a+b)^2 \sqrt{a+b \coth^2(x)}}$$

[Out] $\operatorname{arctanh}(\coth(x) \cdot (a+b)^{1/2} / (a+b \cdot \coth(x)^2)^{1/2}) / (a+b)^{5/2} - 1/3 \cdot \coth(x) / (a+b) / (a+b \cdot \coth(x)^2)^{3/2} - 1/3 \cdot (2 \cdot a - b) \cdot \coth(x) / a / (a+b)^2 / (a+b \cdot \coth(x)^2)^{1/2}$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3751, 482, 541, 12, 385, 212}

$$\int \frac{\coth^2(x)}{(a+b \coth^2(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{(a+b)^{5/2}} - \frac{(2a-b) \coth(x)}{3a(a+b)^2 \sqrt{a+b \coth^2(x)}} - \frac{\coth(x)}{3(a+b)(a+b \coth^2(x))^{3/2}}$$

[In] $\text{Int}[\text{Coth}[x]^2 / (a + b \cdot \text{Coth}[x]^2)^{5/2}, x]$

```
[Out] ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]]/(a + b)^(5/2) - Coth[x]
]/(3*(a + b)*(a + b*Coth[x]^2)^(3/2)) - ((2*a - b)*Coth[x])/(3*a*(a + b)^2*
Sqrt[a + b*Coth[x]^2])
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Su
bst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b
, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 482

```
Int[((e_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_
))^ (q_), x_Symbol] := Simp[e^(n - 1)*(e*x)^(m - n + 1)*(a + b*x^n)^(p + 1)*
((c + d*x^n)^(q + 1)/(n*(b*c - a*d)*(p + 1))), x] - Dist[e^n/(n*(b*c - a*d)
*(p + 1)), Int[(e*x)^(m - n)*(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(m -
n + 1) + d*(m + n*(p + q + 1) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e,
q}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[p, -1] && GeQ[n, m - n +
1] && GtQ[m - n + 1, 0] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]
```

Rule 541

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f
_.)*(x_)^(n_)), x_Symbol] := Simp[(- (b*e - a*f))*x*(a + b*x^n)^(p + 1)*((c
+ d*x^n)^(q + 1)/(a*n*(b*c - a*d)*(p + 1))), x] + Dist[1/(a*n*(b*c - a*d)*(
p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*
c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x] /; Fre
eQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
```

alQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{x^2}{(1-x^2)(a+bx^2)^{5/2}} dx, x, \coth(x)\right) \\
 &= -\frac{\coth(x)}{3(a+b)(a+b\coth^2(x))^{3/2}} + \frac{\text{Subst}\left(\int \frac{1+2x^2}{(1-x^2)(a+bx^2)^{3/2}} dx, x, \coth(x)\right)}{3(a+b)} \\
 &= -\frac{\coth(x)}{3(a+b)(a+b\coth^2(x))^{3/2}} - \frac{(2a-b)\coth(x)}{3a(a+b)^2\sqrt{a+b\coth^2(x)}} \\
 &\quad - \frac{\text{Subst}\left(\int -\frac{3a}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right)}{3a(a+b)^2} \\
 &= -\frac{\coth(x)}{3(a+b)(a+b\coth^2(x))^{3/2}} - \frac{(2a-b)\coth(x)}{3a(a+b)^2\sqrt{a+b\coth^2(x)}} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{1}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right)}{(a+b)^2} \\
 &= -\frac{\coth(x)}{3(a+b)(a+b\coth^2(x))^{3/2}} - \frac{(2a-b)\coth(x)}{3a(a+b)^2\sqrt{a+b\coth^2(x)}} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{1}{1-(a+b)x^2} dx, x, \frac{\coth(x)}{\sqrt{a+b\coth^2(x)}}\right)}{(a+b)^2} \\
 &= \frac{\text{arctanh}\left(\frac{\sqrt{a+b}\coth(x)}{\sqrt{a+b\coth^2(x)}}\right)}{(a+b)^{5/2}} - \frac{\coth(x)}{3(a+b)(a+b\coth^2(x))^{3/2}} - \frac{(2a-b)\coth(x)}{3a(a+b)^2\sqrt{a+b\coth^2(x)}}
 \end{aligned}$$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 7.07 (sec) , antiderivative size = 215, normalized size of antiderivative = 2.44

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{5/2}} dx =$$

$$\frac{\cosh^2(x) \coth(x) \left(\frac{4(a+b) \cosh^2(x) (a+b \coth^2(x)) \operatorname{Hypergeometric2F1}\left(2, 2, \frac{9}{2}, \frac{(a+b) \cosh^2(x)}{a}\right)}{35a^2} - \frac{(-5a - 2b \coth^2(x)) \left(3 \arcsin\left(\sqrt{\frac{(a+b) \cosh^2(x)}{a}}\right) \right)}{3a^2 \sqrt{a + b \coth^2(x)} \left(1 + \right)} \right)}{3a^2 \sqrt{a + b \coth^2(x)} \left(1 + \right)}$$

[In] Integrate[Coth[x]^2/(a + b*Coth[x]^2)^(5/2), x]

[Out] -1/3*(Cosh[x]^2*Coth[x]*((4*(a + b)*Cosh[x]^2*(a + b*Coth[x]^2)*Hypergeomet
ric2F1[2, 2, 9/2, ((a + b)*Cosh[x]^2)/a])/(35*a^2) - ((-5*a - 2*b*Coth[x]^2
)*(3*ArcSin[Sqrt[((a + b)*Cosh[x]^2)/a]]*(a + b*Coth[x]^2)^2 - a*(-4*b*Coth
[x]^2 + a*(-3 - Coth[x]^2))*Csch[x]^2*Sqrt[-(((a + b)*Cosh[x]^2*(a + b*Coth
[x]^2)*Sinh[x]^2)/a^2]))*Tanh[x]^4)/(3*a*(a + b)^2*Sqrt[-(((a + b)*Cosh[x]^2
*(a + b*Coth[x]^2)*Sinh[x]^2)/a^2)])))/(a^2*Sqrt[a + b*Coth[x]^2]*(1 + (b*
Coth[x]^2)/a))

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 453 vs. 2(74) = 148.

Time = 0.10 (sec) , antiderivative size = 454, normalized size of antiderivative = 5.16

method	result
derivativedivides	$-\frac{\coth(x)}{3a(a+b \coth(x)^2)^{\frac{3}{2}}} - \frac{2 \coth(x)}{3a^2 \sqrt{a+b \coth(x)^2}} - \frac{1}{6(a+b)(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{\frac{3}{2}}} + \frac{1}{6(a+b)a(b \coth(x)-1)}$
default	$-\frac{\coth(x)}{3a(a+b \coth(x)^2)^{\frac{3}{2}}} - \frac{2 \coth(x)}{3a^2 \sqrt{a+b \coth(x)^2}} - \frac{1}{6(a+b)(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b)^{\frac{3}{2}}} + \frac{1}{6(a+b)a(b \coth(x)-1)}$

[In] int(coth(x)^2/(a+b*coth(x)^2)^(5/2), x, method=_RETURNVERBOSE)

[Out] -1/3*coth(x)/a/(a+b*coth(x)^2)^(3/2)-2/3/a^2*coth(x)/(a+b*coth(x)^2)^(1/2)-
1/6/(a+b)/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(3/2)+1/6*b/(a+b)/a/(b*(cot
h(x)-1)^2+2*b*(coth(x)-1)+a+b)^(3/2)*coth(x)+1/3*b/(a+b)/a^2/(b*(coth(x)-1
^2+2*b*(coth(x)-1)+a+b)^(1/2)*coth(x)-1/2/(a+b)^2/(b*(coth(x)-1)^2+2*b*(cot
h(x)-1)+a+b)^(1/2)+1/2/(a+b)^2/a/(b*(coth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2
) *b*coth(x)+1/2/(a+b)^(5/2)*ln((2*a+2*b+2*b*(coth(x)-1)+2*(a+b)^(1/2)*(b*(c
oth(x)-1)^2+2*b*(coth(x)-1)+a+b)^(1/2))/(coth(x)-1))+1/6/(a+b)/(b*(1+coth(x
)^2-2*b*(1+coth(x))+a+b)^(3/2)+1/6*b/(a+b)/a/(b*(1+coth(x))^2-2*b*(1+coth(x)

$x)) + a + b)^{3/2} \coth(x) + 1/3 * b / (a + b) / a^2 / (b * (1 + \coth(x))^2 - 2 * b * (1 + \coth(x)) + a + b)^{1/2} * \coth(x) + 1/2 / (a + b)^2 / (b * (1 + \coth(x))^2 - 2 * b * (1 + \coth(x)) + a + b)^{1/2} + 1/2 / (a + b)^2 / a / (b * (1 + \coth(x))^2 - 2 * b * (1 + \coth(x)) + a + b)^{1/2} * b * \coth(x) - 1/2 / (a + b)^{5/2} * \ln((2 * a + 2 * b - 2 * b * (1 + \coth(x)) + 2 * (a + b)^{1/2} * (b * (1 + \coth(x))^2 - 2 * b * (1 + \coth(x)) + a + b)^{1/2}) / (1 + \coth(x)))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3016 vs. $2(74) = 148$.

Time = 0.68 (sec) , antiderivative size = 6591, normalized size of antiderivative = 74.90

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \text{Too large to display}$$

[In] integrate(coth(x)^2/(a+b*coth(x)^2)^(5/2),x, algorithm="fricas")

[Out] Too large to include

Sympy [F]

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\coth^2(x)}{(a + b \coth^2(x))^{5/2}} dx$$

[In] integrate(coth(x)**2/(a+b*coth(x)**2)**(5/2),x)

[Out] Integral(coth(x)**2/(a + b*coth(x)**2)**(5/2), x)

Maxima [F]

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\coth(x)^2}{(b \coth(x)^2 + a)^{5/2}} dx$$

[In] integrate(coth(x)^2/(a+b*coth(x)^2)^(5/2),x, algorithm="maxima")

[Out] integrate(coth(x)^2/(b*coth(x)^2 + a)^(5/2), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 952 vs. $2(74) = 148$.

Time = 0.55 (sec) , antiderivative size = 952, normalized size of antiderivative = 10.82

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \text{Too large to display}$$

[In] integrate(coth(x)^2/(a+b*coth(x)^2)^(5/2),x, algorithm="giac")

[Out]
$$-1/3 * (((3*a^7*b^2*\text{sgn}(e^{2*x}) - 1) + 14*a^6*b^3*\text{sgn}(e^{2*x}) - 1) + 25*a^5*b^4*\text{sgn}(e^{2*x}) - 1) + 20*a^4*b^5*\text{sgn}(e^{2*x}) - 1) + 5*a^3*b^6*\text{sgn}(e^{2*x}) - 1) - 2*a^2*b^7*\text{sgn}(e^{2*x}) - 1) - a*b^8*\text{sgn}(e^{2*x}) - 1)) * e^{2*x} / (a^8*b^2 + 6*a^7*b^3 + 15*a^6*b^4 + 20*a^5*b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*b^8) - 3*(a^7*b^2*\text{sgn}(e^{2*x}) - 1) + 2*a^6*b^3*\text{sgn}(e^{2*x}) - 1) - a^5*b^4*\text{sgn}(e^{2*x}) - 1) - 4*a^4*b^5*\text{sgn}(e^{2*x}) - 1) - a^3*b^6*\text{sgn}(e^{2*x}) - 1) + 2*a^2*b^7*\text{sgn}(e^{2*x}) - 1) + a*b^8*\text{sgn}(e^{2*x}) - 1)) / (a^8*b^2 + 6*a^7*b^3 + 15*a^6*b^4 + 20*a^5*b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*b^8)) * e^{2*x} - 3*(a^7*b^2*\text{sgn}(e^{2*x}) - 1) + 2*a^6*b^3*\text{sgn}(e^{2*x}) - 1) - a^5*b^4*\text{sgn}(e^{2*x}) - 1) - 4*a^4*b^5*\text{sgn}(e^{2*x}) - 1) - a^3*b^6*\text{sgn}(e^{2*x}) - 1) + 2*a^2*b^7*\text{sgn}(e^{2*x}) - 1) + a*b^8*\text{sgn}(e^{2*x}) - 1)) / (a^8*b^2 + 6*a^7*b^3 + 15*a^6*b^4 + 20*a^5*b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*b^8)) * e^{2*x} + (3*a^7*b^2*\text{sgn}(e^{2*x}) - 1) + 14*a^6*b^3*\text{sgn}(e^{2*x}) - 1) + 25*a^5*b^4*\text{sgn}(e^{2*x}) - 1) + 20*a^4*b^5*\text{sgn}(e^{2*x}) - 1) + 5*a^3*b^6*\text{sgn}(e^{2*x}) - 1) - 2*a^2*b^7*\text{sgn}(e^{2*x}) - 1) - a*b^8*\text{sgn}(e^{2*x}) - 1)) / (a^8*b^2 + 6*a^7*b^3 + 15*a^6*b^4 + 20*a^5*b^5 + 15*a^4*b^6 + 6*a^3*b^7 + a^2*b^8)) / (a*e^{4*x} + b*e^{4*x} - 2*a*e^{2*x} + 2*b*e^{2*x} + a + b) * sqrt(a + b) - a + b) / ((a^2 + 2*a*b + b^2)*sqrt(a + b)*sgn(e^{2*x}) - 1) + 1/2*log(abs((sqrt(a + b)*e^{2*x} - sqrt(a*e^{4*x} + b*e^{4*x} - 2*a*e^{2*x} + 2*b*e^{2*x} + a + b)) * sqrt(a + b) - a + b)) / ((a^2 + 2*a*b + b^2)*sqrt(a + b)*sgn(e^{2*x}) - 1)) - 1/2*log(abs(-sqrt(a + b)*e^{2*x} + sqrt(a*e^{4*x} + b*e^{4*x} - 2*a*e^{2*x} + 2*b*e^{2*x} + a + b) - sqrt(a + b))) / ((a^2 + 2*a*b + b^2)*sqrt(a + b)*sgn(e^{2*x}) - 1))$$

Mupad [F(-1)]

Timed out.

$$\int \frac{\coth^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\coth(x)^2}{(b \coth(x)^2 + a)^{5/2}} dx$$

[In] int(coth(x)^2/(a + b*coth(x)^2)^(5/2),x)

[Out] int(coth(x)^2/(a + b*coth(x)^2)^(5/2), x)

$$3.45 \quad \int \frac{\coth(x)}{(a+b \coth^2(x))^{5/2}} dx$$

Optimal result	311
Rubi [A] (verified)	311
Mathematica [C] (verified)	313
Maple [B] (verified)	313
Fricas [B] (verification not implemented)	314
Sympy [F]	315
Maxima [F]	315
Giac [B] (verification not implemented)	315
Mupad [B] (verification not implemented)	316

Optimal result

Integrand size = 15, antiderivative size = 70

$$\int \frac{\coth(x)}{(a+b \coth^2(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{5/2}} - \frac{1}{3(a+b)(a+b \coth^2(x))^{3/2}} - \frac{1}{(a+b)^2 \sqrt{a+b \coth^2(x)}}$$

[Out] $\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2)}/(a+b)^{(1/2)})/(a+b)^{(5/2)}-1/3/(a+b)/(a+b*\coth(x)^2)^{(3/2)}-1/(a+b)^2/(a+b*\coth(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3751, 455, 53, 65, 214}

$$\int \frac{\coth(x)}{(a+b \coth^2(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{5/2}} - \frac{1}{(a+b)^2 \sqrt{a+b \coth^2(x)}} - \frac{1}{3(a+b)(a+b \coth^2(x))^{3/2}}$$

[In] $\operatorname{Int}[\operatorname{Coth}[x]/(a+b*\operatorname{Coth}[x]^2)^{(5/2)},x]$

[Out] $\operatorname{ArcTanh}[\operatorname{Sqrt}[a+b*\operatorname{Coth}[x]^2]/\operatorname{Sqrt}[a+b]]/(a+b)^{(5/2)}-1/(3*(a+b)*(a+b*\operatorname{Coth}[x]^2)^{(3/2)})-1/((a+b)^2*\operatorname{Sqrt}[a+b*\operatorname{Coth}[x]^2])$

Rule 53

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((
m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x]
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && !(LtQ
[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && I
ntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 455

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.
), x_Symbol] := Dist[1/n, Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x
] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n +
1, 0]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst}\left(\int \frac{x}{(1-x^2)(a+bx^2)^{5/2}} dx, x, \coth(x)\right) \\ &= \frac{1}{2} \text{Subst}\left(\int \frac{1}{(1-x)(a+bx)^{5/2}} dx, x, \coth^2(x)\right) \end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{3(a+b)(a+b\coth^2(x))^{3/2}} + \frac{\text{Subst}\left(\int \frac{1}{(1-x)(a+bx)^{3/2}} dx, x, \coth^2(x)\right)}{2(a+b)} \\
&= -\frac{1}{3(a+b)(a+b\coth^2(x))^{3/2}} - \frac{1}{(a+b)^2\sqrt{a+b\coth^2(x)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x)\right)}{2(a+b)^2} \\
&= -\frac{1}{3(a+b)(a+b\coth^2(x))^{3/2}} - \frac{1}{(a+b)^2\sqrt{a+b\coth^2(x)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{1+\frac{a}{b}-\frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)}\right)}{b(a+b)^2} \\
&= \frac{\text{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{5/2}} - \frac{1}{3(a+b)(a+b\coth^2(x))^{3/2}} - \frac{1}{(a+b)^2\sqrt{a+b\coth^2(x)}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.05 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.61

$$\int \frac{\coth(x)}{(a+b\coth^2(x))^{5/2}} dx = -\frac{\text{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{a+b\coth^2(x)}{a+b}\right)}{3(a+b)(a+b\coth^2(x))^{3/2}}$$

[In] Integrate[Coth[x]/(a + b*Coth[x]^2)^(5/2), x]

[Out] -1/3*Hypergeometric2F1[-3/2, 1, -1/2, (a + b*Coth[x]^2)/(a + b)]/((a + b)*(a + b*Coth[x]^2)^(3/2))

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 419 vs. 2(58) = 116.

Time = 0.08 (sec) , antiderivative size = 420, normalized size of antiderivative = 6.00

method	result
derivativedivides	$-\frac{1}{6(a+b)\left(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b\right)^{\frac{3}{2}}} + \frac{b \coth(x)}{6(a+b)a\left(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b\right)^{\frac{3}{2}}} + \frac{1}{3(a+b)a^2}$
default	$-\frac{1}{6(a+b)\left(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b\right)^{\frac{3}{2}}} + \frac{b \coth(x)}{6(a+b)a\left(b(\coth(x)-1)^2+2b(\coth(x)-1)+a+b\right)^{\frac{3}{2}}} + \frac{1}{3(a+b)a^2}$

[In] `int(coth(x)/(a+b*coth(x)^2)^(5/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/6/(a+b)/(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^(3/2)+1/6*b/(a+b)/a/(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^(3/2)*\coth(x)+1/3*b/(a+b)/a^2/(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^(1/2)*\coth(x)-1/2/(a+b)^2/(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^(1/2)+1/2/(a+b)^2/a/(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^(1/2)*b*\coth(x)+1/2/(a+b)^(5/2)*\ln((2*a+2*b+2*b*(\coth(x)-1)+2*(a+b)^(1/2)*(b*(\coth(x)-1)^2+2*b*(\coth(x)-1)+a+b)^(1/2)))/(\coth(x)-1))-1/6/(a+b)/(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^(3/2)-1/6*b/(a+b)/a/(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^(3/2)*\coth(x)-1/3*b/(a+b)/a^2/(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^(1/2)*\coth(x)-1/2/(a+b)^2/(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^(1/2)-1/2/(a+b)^2/a/(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^(1/2)*b*\coth(x)+1/2/(a+b)^(5/2)*\ln((2*a+2*b-2*b*(1+\coth(x))+2*(a+b)^(1/2)*(b*(1+\coth(x))^2-2*b*(1+\coth(x))+a+b)^(1/2)))/(1+\coth(x)))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2552 vs. $2(58) = 116$.

Time = 0.63 (sec) , antiderivative size = 5736, normalized size of antiderivative = 81.94

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{5/2}} dx = \text{Too large to display}$$

[In] `integrate(coth(x)/(a+b*coth(x)^2)^(5/2),x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\coth(x)}{(a + b \coth^2(x))^{\frac{5}{2}}} dx$$

[In] integrate(coth(x)/(a+b*coth(x)**2)**(5/2), x)

[Out] Integral(coth(x)/(a + b*coth(x)**2)**(5/2), x)

Maxima [F]

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\coth(x)}{(b \coth(x)^2 + a)^{\frac{5}{2}}} dx$$

[In] integrate(coth(x)/(a+b*coth(x)^2)^(5/2), x, algorithm="maxima")

[Out] integrate(coth(x)/(b*coth(x)^2 + a)^(5/2), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 909 vs. $2(58) = 116$.

Time = 0.58 (sec) , antiderivative size = 909, normalized size of antiderivative = 12.99

$$\int \frac{\coth(x)}{(a + b \coth^2(x))^{5/2}} dx = \text{Too large to display}$$

[In] integrate(coth(x)/(a+b*coth(x)^2)^(5/2), x, algorithm="giac")

[Out]
$$\begin{aligned} & -4/3 * (((a^7 * b^2 + 5 * a^6 * b^3 + 10 * a^5 * b^4 + 10 * a^4 * b^5 + 5 * a^3 * b^6 + a^2 * b^7) * e^{(2 * x)} / (a^8 * b^2 * \operatorname{sgn}(e^{(2 * x)} - 1) + 6 * a^7 * b^3 * \operatorname{sgn}(e^{(2 * x)} - 1) + 15 * a^6 * b^4 * \operatorname{sgn}(e^{(2 * x)} - 1) + 20 * a^5 * b^5 * \operatorname{sgn}(e^{(2 * x)} - 1) + 15 * a^4 * b^6 * \operatorname{sgn}(e^{(2 * x)} - 1) + 6 * a^3 * b^7 * \operatorname{sgn}(e^{(2 * x)} - 1) + a^2 * b^8 * \operatorname{sgn}(e^{(2 * x)} - 1)) - 3 * (a^7 * b^2 + 4 * a^6 * b^3 + 6 * a^5 * b^4 + 4 * a^4 * b^5 + a^3 * b^6) / (a^8 * b^2 * \operatorname{sgn}(e^{(2 * x)} - 1) + 6 * a^7 * b^3 * \operatorname{sgn}(e^{(2 * x)} - 1) + 15 * a^6 * b^4 * \operatorname{sgn}(e^{(2 * x)} - 1) + 20 * a^5 * b^5 * \operatorname{sgn}(e^{(2 * x)} - 1) + 15 * a^4 * b^6 * \operatorname{sgn}(e^{(2 * x)} - 1) + 6 * a^3 * b^7 * \operatorname{sgn}(e^{(2 * x)} - 1) + a^2 * b^8 * \operatorname{sgn}(e^{(2 * x)} - 1))) * e^{(2 * x)} + 3 * (a^7 * b^2 + 4 * a^6 * b^3 + 6 * a^5 * b^4 + 4 * a^4 * b^5 + a^3 * b^6) / (a^8 * b^2 * \operatorname{sgn}(e^{(2 * x)} - 1) + 6 * a^7 * b^3 * \operatorname{sgn}(e^{(2 * x)} - 1) + 15 * a^6 * b^4 * \operatorname{sgn}(e^{(2 * x)} - 1) + 20 * a^5 * b^5 * \operatorname{sgn}(e^{(2 * x)} - 1) + 15 * a^4 * b^6 * \operatorname{sgn}(e^{(2 * x)} - 1) + 6 * a^3 * b^7 * \operatorname{sgn}(e^{(2 * x)} - 1) + a^2 * b^8 * \operatorname{sgn}(e^{(2 * x)} - 1))) * e^{(2 * x)} - (a^7 * b^2 + 5 * a^6 * b^3 + 10 * a^5 * b^4 + 10 * a^4 * b^5 + 5 * a^3 * b^6 + a^2 * b^7) / (a^8 * b^2 * \operatorname{sgn}(e^{(2 * x)} - 1) + 6 * a^7 * b^3 * \operatorname{sgn}(e^{(2 * x)} - 1) + 15 * a^6 * b^4 * \operatorname{sgn}(e^{(2 * x)} - 1) + 20 * a^5 * b^5 * \operatorname{sgn}(e^{(2 * x)} - 1) + 15 * a^4 * b^6 * \operatorname{sgn}(e^{(2 * x)} - 1) + 6 * a^3 * b^7 * \operatorname{sgn}(e^{(2 * x)} - 1) + a^2 * b^8 * \operatorname{sgn}(e^{(2 * x)} - 1)) \end{aligned}$$

$$\begin{aligned} & \left(e^{2x} - 1 \right) + 20a^5b^5 \operatorname{sgn}(e^{2x} - 1) + 15a^4b^6 \operatorname{sgn}(e^{2x} - 1) + 6 \\ & a^3b^7 \operatorname{sgn}(e^{2x} - 1) + a^2b^8 \operatorname{sgn}(e^{2x} - 1) \Big) / (a e^{4x} + b e^{4x} \\ & - 2a e^{2x} + 2b e^{2x} + a + b)^{3/2} - 1/2 \log(\operatorname{abs}(\sqrt{a+b} e^{2x} \\ & - \sqrt{a e^{4x} + b e^{4x} - 2a e^{2x} + 2b e^{2x} + a + b}) \sqrt{a+b} \\ & - a + b) / ((a^2 + 2ab + b^2) \sqrt{a+b} \operatorname{sgn}(e^{2x} - 1)) - 1/ \\ & 2 \log(\operatorname{abs}(\sqrt{a+b} e^{2x} - \sqrt{a e^{4x} + b e^{4x} - 2a e^{2x} + 2b e^{2x} + a + b}) \sqrt{a+b} \\ & - a - b) / ((a^2 + 2ab + b^2) \sqrt{a+b} \operatorname{sgn}(e^{2x} - 1)) + 1/2 \log(\operatorname{abs}(-\sqrt{a+b} e^{2x} \\ & + \sqrt{a e^{4x} + b e^{4x} - 2a e^{2x} + 2b e^{2x} + a + b} - \sqrt{a+b})) / ((a^2 + 2ab \\ & + b^2) \sqrt{a+b} \operatorname{sgn}(e^{2x} - 1)) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 4.33 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.09

$$\int \frac{\operatorname{coth}(x)}{(a + b \operatorname{coth}^2(x))^{5/2}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \operatorname{coth}(x)^2 + a(2a^2 + 4ab + 2b^2)}}{2(a+b)^{5/2}}\right)}{(a+b)^{5/2}} - \frac{\frac{1}{3(a+b)} + \frac{b \operatorname{coth}(x)^2 + a}{(a+b)^2}}{(b \operatorname{coth}(x)^2 + a)^{3/2}}$$

[In] int(coth(x)/(a + b*coth(x)^2)^(5/2),x)

[Out] atanh(((a + b*coth(x)^2)^(1/2)*(4*a*b + 2*a^2 + 2*b^2))/(2*(a + b)^(5/2)))/
(a + b)^(5/2) - (1/(3*(a + b)) + (a + b*coth(x)^2)/(a + b)^2)/(a + b*coth(x)
)^2)^(3/2)

$$3.46 \quad \int \frac{\tanh(x)}{(a+b \coth^2(x))^{5/2}} dx$$

Optimal result	317
Rubi [A] (verified)	317
Mathematica [C] (verified)	320
Maple [F]	320
Fricas [B] (verification not implemented)	320
Sympy [F]	321
Maxima [F]	321
Giac [F(-2)]	321
Mupad [F(-1)]	321

Optimal result

Integrand size = 15, antiderivative size = 108

$$\int \frac{\tanh(x)}{(a+b \coth^2(x))^{5/2}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a}}\right)}{a^{5/2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{5/2}} + \frac{b}{3a(a+b)(a+b \coth^2(x))^{3/2}} + \frac{b(2a+b)}{a^2(a+b)^2 \sqrt{a+b \coth^2(x)}}$$

[Out] $-\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2))/a^{(1/2)})/a^{(5/2)}+\operatorname{arctanh}((a+b*\coth(x)^2)^{(1/2))/(a+b)^{(1/2)})/(a+b)^{(5/2)}+1/3*b/a/(a+b)/(a+b*\coth(x)^2)^{(3/2)}+b*(2*a+b)/a^2/(a+b)^2/(a+b*\coth(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3751, 457, 87, 157, 162, 65, 214}

$$\int \frac{\tanh(x)}{(a+b \coth^2(x))^{5/2}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a}}\right)}{a^{5/2}} + \frac{b(2a+b)}{a^2(a+b)^2 \sqrt{a+b \coth^2(x)}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{5/2}} + \frac{b}{3a(a+b)(a+b \coth^2(x))^{3/2}}$$

[In] $\operatorname{Int}[\operatorname{Tanh}[x]/(a+b*\operatorname{Coth}[x]^2)^{(5/2)},x]$

[Out] $-(\text{ArcTanh}[\text{Sqrt}[a + b \cdot \text{Coth}[x]^2]/\text{Sqrt}[a]]/a^{(5/2)}) + \text{ArcTanh}[\text{Sqrt}[a + b \cdot \text{Coth}[x]^2]/\text{Sqrt}[a + b]]/(a + b)^{(5/2)} + b/(3 \cdot a \cdot (a + b) \cdot (a + b \cdot \text{Coth}[x]^2)^{(3/2)}) + (b \cdot (2 \cdot a + b))/(a^2 \cdot (a + b)^2 \cdot \text{Sqrt}[a + b \cdot \text{Coth}[x]^2])$

Rule 65

$\text{Int}[(a_. + (b_.)(x_))^{(m_)} \cdot ((c_.) + (d_.)(x_))^{(n_)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p \cdot (m + 1) - 1)} \cdot (c - a \cdot (d/b) + d \cdot (x^p/b))^{(n)}, x], x, (a + b \cdot x)^{(1/p)}], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 87

$\text{Int}[(e_. + (f_.)(x_))^{(p_)} / (((a_.) + (b_.)(x_)) \cdot ((c_.) + (d_.)(x_))), x_Symbol] \rightarrow \text{Simp}[f \cdot ((e + f \cdot x)^{(p + 1)} / ((p + 1) \cdot (b \cdot e - a \cdot f) \cdot (d \cdot e - c \cdot f))), x] + \text{Dist}[1 / ((b \cdot e - a \cdot f) \cdot (d \cdot e - c \cdot f)), \text{Int}[(b \cdot d \cdot e - b \cdot c \cdot f - a \cdot d \cdot f - b \cdot d \cdot f \cdot x) \cdot ((e + f \cdot x)^{(p + 1)} / ((a + b \cdot x) \cdot (c + d \cdot x))), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{LtQ}[p, -1]$

Rule 157

$\text{Int}[(a_. + (b_.)(x_))^{(m_)} \cdot ((c_.) + (d_.)(x_))^{(n_)} \cdot ((e_.) + (f_.)(x_))^{(p_)} \cdot ((g_.) + (h_.)(x_)), x_Symbol] \rightarrow \text{Simp}[(b \cdot g - a \cdot h) \cdot (a + b \cdot x)^{(m + 1)} \cdot (c + d \cdot x)^{(n + 1)} \cdot ((e + f \cdot x)^{(p + 1)} / ((m + 1) \cdot (b \cdot c - a \cdot d) \cdot (b \cdot e - a \cdot f))), x] + \text{Dist}[1 / ((m + 1) \cdot (b \cdot c - a \cdot d) \cdot (b \cdot e - a \cdot f)), \text{Int}[(a + b \cdot x)^{(m + 1)} \cdot (c + d \cdot x)^n \cdot (e + f \cdot x)^p \cdot \text{Simp}[(a \cdot d \cdot f \cdot g - b \cdot (d \cdot e + c \cdot f) \cdot g + b \cdot c \cdot e \cdot h) \cdot (m + 1) - (b \cdot g - a \cdot h) \cdot (d \cdot e \cdot (n + 1) + c \cdot f \cdot (p + 1)) - d \cdot f \cdot (b \cdot g - a \cdot h) \cdot (m + n + p + 3) \cdot x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, h, n, p\}, x] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntegersQ}[2 \cdot m, 2 \cdot n, 2 \cdot p]$

Rule 162

$\text{Int}[(e_. + (f_.)(x_))^{(p_)} \cdot ((g_.) + (h_.)(x_)) / (((a_.) + (b_.)(x_)) \cdot ((c_.) + (d_.)(x_))), x_Symbol] \rightarrow \text{Dist}[(b \cdot g - a \cdot h) / (b \cdot c - a \cdot d), \text{Int}[(e + f \cdot x)^p / (a + b \cdot x), x], x] - \text{Dist}[(d \cdot g - c \cdot h) / (b \cdot c - a \cdot d), \text{Int}[(e + f \cdot x)^p / (c + d \cdot x), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, h\}, x]$

Rule 214

$\text{Int}[(a_. + (b_.)(x_)^2)^{(-1)}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) \cdot \text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$

Rule 457

$\text{Int}[(x_)^{(m_)} \cdot ((a_.) + (b_.)(x_)^{(n_)})^{(p_)} \cdot ((c_.) + (d_.)(x_)^{(n_)})^{(q_.)}, x_Symbol] \rightarrow \text{Dist}[1/n, \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)} \cdot (a + b \cdot x)^p$

$*(c + d*x)^q, x], x, x^n], x] /; \text{FreeQ}[\{a, b, c, d, m, n, p, q\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]$

Rule 3751

$\text{Int}[\{(d \cdot) \cdot \tan[(e \cdot) + (f \cdot) \cdot (x \cdot)]\}^{(m \cdot)} \cdot \{(a \cdot) + (b \cdot) \cdot \{(c \cdot) \cdot \tan[(e \cdot) + (f \cdot) \cdot (x \cdot)]\}^{(n \cdot)}\}^{(p \cdot)}, x_Symbol] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Tan}[e + f \cdot x], x]\}, \text{Dist}[c \cdot (ff/f), \text{Subst}[\text{Int}[(d \cdot ff \cdot (x/c))^{m \cdot} \cdot \{(a + b \cdot (ff \cdot x)^n\}^p / (c^2 + ff^2 \cdot x^2)}, x], x, c \cdot (\text{Tan}[e + f \cdot x] / ff)], x]] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ (\text{IGtQ}[p, 0] \ || \ \text{EqQ}[n, 2] \ || \ \text{EqQ}[n, 4] \ || \ (\text{IntegerQ}[p] \ \&\& \ \text{RationalQ}[n]))$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{1}{x(1-x^2)(a+bx^2)^{5/2}} dx, x, \coth(x)\right) \\
 &= \frac{1}{2} \text{Subst}\left(\int \frac{1}{(1-x)x(a+bx)^{5/2}} dx, x, \coth^2(x)\right) \\
 &= \frac{b}{3a(a+b)(a+b\coth^2(x))^{3/2}} - \frac{\text{Subst}\left(\int \frac{-a-b+bx}{(1-x)x(a+bx)^{3/2}} dx, x, \coth^2(x)\right)}{2a(a+b)} \\
 &= \frac{b}{3a(a+b)(a+b\coth^2(x))^{3/2}} + \frac{b(2a+b)}{a^2(a+b)^2\sqrt{a+b\coth^2(x)}} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{-\frac{1}{2}(a+b)^2 + \frac{1}{2}b(2a+b)x}{(1-x)x\sqrt{a+bx}} dx, x, \coth^2(x)\right)}{a^2(a+b)^2} \\
 &= \frac{b}{3a(a+b)(a+b\coth^2(x))^{3/2}} + \frac{b(2a+b)}{a^2(a+b)^2\sqrt{a+b\coth^2(x)}} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx}} dx, x, \coth^2(x)\right)}{2a^2} + \frac{\text{Subst}\left(\int \frac{1}{(1-x)\sqrt{a+bx}} dx, x, \coth^2(x)\right)}{2(a+b)^2} \\
 &= \frac{b}{3a(a+b)(a+b\coth^2(x))^{3/2}} + \frac{b(2a+b)}{a^2(a+b)^2\sqrt{a+b\coth^2(x)}} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{1}{-\frac{a}{b} + \frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)}\right)}{a^2b} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{1}{1 + \frac{a}{b} - \frac{x^2}{b}} dx, x, \sqrt{a+b\coth^2(x)}\right)}{b(a+b)^2}
 \end{aligned}$$

$$= -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a}}\right)}{a^{5/2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b\coth^2(x)}}{\sqrt{a+b}}\right)}{(a+b)^{5/2}} \\ + \frac{b}{3a(a+b)(a+b\coth^2(x))^{3/2}} + \frac{b(2a+b)}{a^2(a+b)^2\sqrt{a+b\coth^2(x)}}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.07 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.68

$$\int \frac{\tanh(x)}{(a+b\coth^2(x))^{5/2}} dx = \frac{-a \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{a+b\coth^2(x)}{a+b}\right) + (a+b) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, 1 + \frac{b\coth^2(x)}{a}\right)}{3a(a+b)(a+b\coth^2(x))^{3/2}}$$

[In] Integrate[Tanh[x]/(a + b*Coth[x]^2)^(5/2),x]

[Out] $(-a \operatorname{Hypergeometric2F1}[-3/2, 1, -1/2, (a + b \operatorname{Coth}[x]^2)/(a + b)]) + (a + b) \operatorname{Hypergeometric2F1}[-3/2, 1, -1/2, 1 + (b \operatorname{Coth}[x]^2)/a]) / (3a(a + b)(a + b \operatorname{Coth}[x]^2)^{3/2})$

Maple [F]

$$\int \frac{\tanh(x)}{(a+b\coth(x)^2)^{5/2}} dx$$

[In] int(tanh(x)/(a+b*coth(x)^2)^(5/2),x)

[Out] int(tanh(x)/(a+b*coth(x)^2)^(5/2),x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4452 vs. 2(90) = 180.

Time = 1.24 (sec) , antiderivative size = 19199, normalized size of antiderivative = 177.77

$$\int \frac{\tanh(x)}{(a+b\coth^2(x))^{5/2}} dx = \text{Too large to display}$$

[In] integrate(tanh(x)/(a+b*coth(x)^2)^(5/2),x, algorithm="fricas")

[Out] Too large to include

Sympy [F]

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\tanh(x)}{(a + b \coth^2(x))^{5/2}} dx$$

[In] integrate(tanh(x)/(a+b*coth(x)**2)**(5/2), x)

[Out] Integral(tanh(x)/(a + b*coth(x)**2)**(5/2), x)

Maxima [F]

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\tanh(x)}{(b \coth(x)^2 + a)^{5/2}} dx$$

[In] integrate(tanh(x)/(a+b*coth(x)^2)^(5/2), x, algorithm="maxima")

[Out] integrate(tanh(x)/(b*coth(x)^2 + a)^(5/2), x)

Giac [F(-2)]

Exception generated.

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{5/2}} dx = \text{Exception raised: TypeError}$$

[In] integrate(tanh(x)/(a+b*coth(x)^2)^(5/2), x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type

Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\tanh(x)}{(b \coth(x)^2 + a)^{5/2}} dx$$

[In] int(tanh(x)/(a + b*coth(x)^2)^(5/2), x)

[Out] int(tanh(x)/(a + b*coth(x)^2)^(5/2), x)

$$3.47 \quad \int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{5/2}} dx$$

Optimal result	322
Rubi [A] (verified)	322
Mathematica [C] (warning: unable to verify)	325
Maple [F]	327
Fricas [B] (verification not implemented)	327
Sympy [F]	327
Maxima [F]	327
Giac [B] (verification not implemented)	328
Mupad [F(-1)]	329

Optimal result

Integrand size = 17, antiderivative size = 131

$$\int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{(a+b)^{5/2}} + \frac{b \tanh(x)}{3a(a+b)(a+b \coth^2(x))^{3/2}}$$

$$+ \frac{b(7a+4b) \tanh(x)}{3a^2(a+b)^2 \sqrt{a+b \coth^2(x)}} - \frac{(3a+2b)(a+4b) \sqrt{a+b \coth^2(x)} \tanh(x)}{3a^3(a+b)^2}$$

[Out] $\operatorname{arctanh}(\coth(x) \cdot (a+b)^{1/2} / (a+b \coth(x)^2)^{1/2}) / (a+b)^{5/2} + 1/3 \cdot b \cdot \tanh(x) / a / (a+b) / (a+b \coth(x)^2)^{3/2} + 1/3 \cdot b \cdot (7a+4b) \cdot \tanh(x) / a^2 / (a+b)^2 / (a+b \coth(x)^2)^{1/2} - 1/3 \cdot (3a+2b) \cdot (a+4b) \cdot (a+b \coth(x)^2)^{1/2} \cdot \tanh(x) / a^3 / (a+b)^2$

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.412$, Rules used = {3751, 483, 593, 597, 12, 385, 212}

$$\int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{5/2}} dx = -\frac{(3a+2b)(a+4b) \tanh(x) \sqrt{a+b \coth^2(x)}}{3a^3(a+b)^2}$$

$$+ \frac{b(7a+4b) \tanh(x)}{3a^2(a+b)^2 \sqrt{a+b \coth^2(x)}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{(a+b)^{5/2}} + \frac{b \tanh(x)}{3a(a+b)(a+b \coth^2(x))^{3/2}}$$

[In] Int[Tanh[x]^2/(a + b*Coth[x]^2)^(5/2), x]

[Out] ArcTanh[(Sqrt[a + b]*Coth[x])/Sqrt[a + b*Coth[x]^2]]/(a + b)^(5/2) + (b*Tanh[x])/(3*a*(a + b)*(a + b*Coth[x]^2)^(3/2)) + (b*(7*a + 4*b)*Tanh[x])/(3*a^2*(a + b)^2*Sqrt[a + b*Coth[x]^2]) - ((3*a + 2*b)*(a + 4*b)*Sqrt[a + b*Coth[x]^2]*Tanh[x])/(3*a^3*(a + b)^2)

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 483

Int[((e_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-b)*(e*x)^(m + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*e*n*(b*c - a*d)*(p + 1))), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(e*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*b*(m + 1) + n*(b*c - a*d)*(p + 1) + d*b*(m + n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, m, q}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[p, -1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

Rule 593

Int[((g_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Simp[(-b*e - a*f)*(g*x)^(m + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*g*n*(b*c - a*d)*(p + 1))), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(g*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f)*(m + 1) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(m + n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, q}, x] && IGtQ[n, 0] && LtQ[p, -1]

Rule 597

Int[((g_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Simp[e*(g*x)^(m + 1)*(a + b

```
*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*c*g^(m + 1))), x] + Dist[1/(a*c*g^n*(
m + 1)), Int[(g*x)^(m + n)*(a + b*x^n)^p*(c + d*x^n)^q*Simp[a*f*c*(m + 1) -
e*(b*c + a*d)*(m + n + 1) - e*n*(b*c*p + a*d*q) - b*e*d*(m + n*(p + q + 2)
+ 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p, q}, x] && IGtQ[n, 0
] && LtQ[m, -1]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_.) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{1}{x^2(1-x^2)(a+bx^2)^{5/2}} dx, x, \coth(x)\right) \\
&= \frac{b \tanh(x)}{3a(a+b)(a+b \coth^2(x))^{3/2}} - \frac{\text{Subst}\left(\int \frac{-3a-4b+4bx^2}{x^2(1-x^2)(a+bx^2)^{3/2}} dx, x, \coth(x)\right)}{3a(a+b)} \\
&= \frac{b \tanh(x)}{3a(a+b)(a+b \coth^2(x))^{3/2}} + \frac{b(7a+4b) \tanh(x)}{3a^2(a+b)^2 \sqrt{a+b \coth^2(x)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{(3a+2b)(a+4b)-2b(7a+4b)x^2}{x^2(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right)}{3a^2(a+b)^2} \\
&= \frac{b \tanh(x)}{3a(a+b)(a+b \coth^2(x))^{3/2}} + \frac{b(7a+4b) \tanh(x)}{3a^2(a+b)^2 \sqrt{a+b \coth^2(x)}} \\
&\quad - \frac{(3a+2b)(a+4b) \sqrt{a+b \coth^2(x)} \tanh(x)}{3a^3(a+b)^2} \\
&\quad - \frac{\text{Subst}\left(\int -\frac{3a^3}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right)}{3a^3(a+b)^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b \tanh(x)}{3a(a+b) (a+b \coth^2(x))^{3/2}} + \frac{b(7a+4b) \tanh(x)}{3a^2(a+b)^2 \sqrt{a+b \coth^2(x)}} \\
&\quad - \frac{(3a+2b)(a+4b) \sqrt{a+b \coth^2(x)} \tanh(x)}{3a^3(a+b)^2} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{(1-x^2)\sqrt{a+bx^2}} dx, x, \coth(x)\right)}{(a+b)^2} \\
&= \frac{b \tanh(x)}{3a(a+b) (a+b \coth^2(x))^{3/2}} + \frac{b(7a+4b) \tanh(x)}{3a^2(a+b)^2 \sqrt{a+b \coth^2(x)}} \\
&\quad - \frac{(3a+2b)(a+4b) \sqrt{a+b \coth^2(x)} \tanh(x)}{3a^3(a+b)^2} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{1-(a+b)x^2} dx, x, \frac{\coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{(a+b)^2} \\
&= \frac{\text{arctanh}\left(\frac{\sqrt{a+b} \coth(x)}{\sqrt{a+b \coth^2(x)}}\right)}{(a+b)^{5/2}} + \frac{b \tanh(x)}{3a(a+b) (a+b \coth^2(x))^{3/2}} \\
&\quad + \frac{b(7a+4b) \tanh(x)}{3a^2(a+b)^2 \sqrt{a+b \coth^2(x)}} - \frac{(3a+2b)(a+4b) \sqrt{a+b \coth^2(x)} \tanh(x)}{3a^3(a+b)^2}
\end{aligned}$$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 8.54 (sec) , antiderivative size = 1350, normalized size of antiderivative = 10.31

$$\int \frac{\tanh^2(x)}{(a+b \coth^2(x))^{5/2}} dx = \frac{\sinh^2(x)}{3a^2(a+b)^2} \left(\frac{16b^3(-i \coth(x) + i \coth^3(x))^2}{a(a+b)^2} + \frac{40b \text{CSch}^2(x)}{a+b} + \frac{160b^2 \coth^2(x) \text{CSch}^2(x)}{3a(a+b)} + \frac{64b^3 \coth^4(x)}{3a^2} \right)$$

[In] Integrate[Tanh[x]^2/(a + b*Coth[x]^2)^(5/2), x]

[Out] (Sinh[x]^2*((16*b^3*((-I)*Coth[x] + I*Coth[x]^3)^2)/(a*(a + b)^2) + (40*b*Csch[x]^2)/(a + b) + (160*b^2*Coth[x]^2*Csch[x]^2)/(3*a*(a + b)) + (64*b^3*Coth[x]^4*Csch[x]^2)/(3*a^2*(a + b)) - (40*b^2*Csch[x]^4)/(a + b)^2 + (92*(a + b)*Cosh[x]^2*Hypergeometric2F1[2, 2, 9/2, ((a + b)*Cosh[x]^2)/a])/(105*a) + (124*b*(a + b)*Cosh[x]^2*Coth[x]^2*Hypergeometric2F1[2, 2, 9/2, ((a + b)*Cosh[x]^2)/a])/(35*a^2) + (152*b^2*(a + b)*Cosh[x]^2*Coth[x]^4*Hypergeome

$$\begin{aligned}
& \text{tric2F1}[2, 2, 9/2, ((a + b) \cdot \text{Cosh}[x]^2)/a]/(35 \cdot a^3) + (176 \cdot b^3 \cdot (a + b) \cdot \text{Cosh}[x]^2 \cdot \text{Coth}[x]^6 \cdot \text{Hypergeometric2F1}[2, 2, 9/2, ((a + b) \cdot \text{Cosh}[x]^2)/a]/(105 \cdot a^4) \\
& + (24 \cdot (a + b) \cdot \text{Cosh}[x]^2 \cdot \text{HypergeometricPFQ}[\{2, 2, 2\}, \{1, 9/2\}, ((a + b) \cdot \text{Cosh}[x]^2)/a]/(35 \cdot a) + (16 \cdot b \cdot (a + b) \cdot \text{Cosh}[x]^2 \cdot \text{Coth}[x]^2 \cdot \text{HypergeometricPFQ}[\{2, 2, 2\}, \{1, 9/2\}, ((a + b) \cdot \text{Cosh}[x]^2)/a]/(7 \cdot a^2) \\
& + (88 \cdot b^2 \cdot (a + b) \cdot \text{Cosh}[x]^2 \cdot \text{Coth}[x]^4 \cdot \text{HypergeometricPFQ}[\{2, 2, 2\}, \{1, 9/2\}, ((a + b) \cdot \text{Cosh}[x]^2)/a]/(35 \cdot a^3) + (32 \cdot b^3 \cdot (a + b) \cdot \text{Cosh}[x]^2 \cdot \text{Coth}[x]^6 \cdot \text{HypergeometricPFQ}[\{2, 2, 2\}, \{1, 9/2\}, ((a + b) \cdot \text{Cosh}[x]^2)/a]/(35 \cdot a^4) \\
& + (16 \cdot (a + b) \cdot \text{Cosh}[x]^2 \cdot \text{HypergeometricPFQ}[\{2, 2, 2, 2\}, \{1, 1, 9/2\}, ((a + b) \cdot \text{Cosh}[x]^2)/a]/(105 \cdot a) + (16 \cdot b \cdot (a + b) \cdot \text{Cosh}[x]^2 \cdot \text{Coth}[x]^2 \cdot \text{HypergeometricPFQ}[\{2, 2, 2, 2\}, \{1, 1, 9/2\}, ((a + b) \cdot \text{Cosh}[x]^2)/a]/(35 \cdot a^2) \\
& + (16 \cdot b^2 \cdot (a + b) \cdot \text{Cosh}[x]^2 \cdot \text{Coth}[x]^4 \cdot \text{HypergeometricPFQ}[\{2, 2, 2, 2\}, \{1, 1, 9/2\}, ((a + b) \cdot \text{Cosh}[x]^2)/a]/(35 \cdot a^3) + (16 \cdot b^3 \cdot (a + b) \cdot \text{Cosh}[x]^2 \cdot \text{Coth}[x]^6 \cdot \text{HypergeometricPFQ}[\{2, 2, 2, 2\}, \{1, 1, 9/2\}, ((a + b) \cdot \text{Cosh}[x]^2)/a]/(105 \cdot a^4) \\
& + (20 \cdot a \cdot \text{Sech}[x]^2)/(3 \cdot (a + b)) - (30 \cdot a \cdot b \cdot \text{Csch}[x]^2 \cdot \text{Sech}[x]^2)/(a + b)^2 - (5 \cdot a^2 \cdot \text{Sech}[x]^4)/(a + b)^2 \\
& + (5 \cdot \text{ArcSin}[\text{Sqrt}[(a + b) \cdot \text{Cosh}[x]^2)/a])/((a + b) \cdot \text{Cosh}[x]^2/a)^{(5/2)} \cdot \text{Sqrt}[-((a + b) \cdot \text{Coth}[x]^2 \cdot \text{Sinh}[x]^2)/a]) + (30 \cdot b \cdot \text{ArcSin}[\text{Sqrt}[(a + b) \cdot \text{Cosh}[x]^2)/a] \cdot \text{Coth}[x]^2)/(a \cdot ((a + b) \cdot \text{Cosh}[x]^2/a)^{(5/2)} \cdot \text{Sqrt}[-((a + b) \cdot \text{Coth}[x]^2 \cdot \text{Sinh}[x]^2)/a]) \\
& + (40 \cdot b^2 \cdot \text{ArcSin}[\text{Sqrt}[(a + b) \cdot \text{Cosh}[x]^2)/a] \cdot \text{Coth}[x]^4)/(a^2 \cdot ((a + b) \cdot \text{Cosh}[x]^2/a)^{(5/2)} \cdot \text{Sqrt}[-((a + b) \cdot \text{Coth}[x]^2 \cdot \text{Sinh}[x]^2)/a]) \\
& + (16 \cdot b^3 \cdot \text{ArcSin}[\text{Sqrt}[(a + b) \cdot \text{Cosh}[x]^2)/a] \cdot \text{Coth}[x]^6)/(a^3 \cdot ((a + b) \cdot \text{Cosh}[x]^2/a)^{(5/2)} \cdot \text{Sqrt}[-((a + b) \cdot \text{Coth}[x]^2 \cdot \text{Sinh}[x]^2)/a]) + (5 \cdot \text{ArcSin}[\text{Sqrt}[(a + b) \cdot \text{Cosh}[x]^2)/a])/ \text{Sqrt}[-((a + b) \cdot \text{Cosh}[x]^2 \cdot (a + b) \cdot \text{Coth}[x]^2 \cdot \text{Sinh}[x]^2)/a^2] \\
& + (30 \cdot b \cdot \text{ArcSin}[\text{Sqrt}[(a + b) \cdot \text{Cosh}[x]^2)/a] \cdot \text{Coth}[x]^2)/(a \cdot \text{Sqrt}[-((a + b) \cdot \text{Cosh}[x]^2 \cdot (a + b) \cdot \text{Coth}[x]^2 \cdot \text{Sinh}[x]^2)/a^2]) + (40 \cdot b^2 \cdot \text{ArcSin}[\text{Sqrt}[(a + b) \cdot \text{Cosh}[x]^2)/a] \cdot \text{Coth}[x]^4)/(a^2 \cdot \text{Sqrt}[-((a + b) \cdot \text{Cosh}[x]^2 \cdot (a + b) \cdot \text{Coth}[x]^2 \cdot \text{Sinh}[x]^2)/a^2]) \\
& + (16 \cdot b^3 \cdot \text{ArcSin}[\text{Sqrt}[(a + b) \cdot \text{Cosh}[x]^2)/a] \cdot \text{Coth}[x]^6)/(a^3 \cdot \text{Sqrt}[-((a + b) \cdot \text{Cosh}[x]^2 \cdot (a + b) \cdot \text{Coth}[x]^2 \cdot \text{Sinh}[x]^2)/a^2]) - (60 \cdot b \cdot \text{ArcSin}[\text{Sqrt}[(a + b) \cdot \text{Cosh}[x]^2)/a] \cdot \text{Csch}[x]^2)/((a + b) \cdot \text{Sqrt}[-((a + b) \cdot \text{Cosh}[x]^2 \cdot (a + b) \cdot \text{Coth}[x]^2 \cdot \text{Sinh}[x]^2)/a^2]) \\
& - (80 \cdot b^2 \cdot \text{ArcSin}[\text{Sqrt}[(a + b) \cdot \text{Cosh}[x]^2)/a] \cdot \text{Coth}[x]^2 \cdot \text{Csch}[x]^2)/(a \cdot (a + b) \cdot \text{Sqrt}[-((a + b) \cdot \text{Cosh}[x]^2 \cdot (a + b) \cdot \text{Coth}[x]^2 \cdot \text{Sinh}[x]^2)/a^2]) - (32 \cdot b^3 \cdot \text{ArcSin}[\text{Sqrt}[(a + b) \cdot \text{Cosh}[x]^2)/a] \cdot \text{Coth}[x]^4 \cdot \text{Csch}[x]^2)/(a^2 \cdot (a + b) \cdot \text{Sqrt}[-((a + b) \cdot \text{Cosh}[x]^2 \cdot (a + b) \cdot \text{Coth}[x]^2 \cdot \text{Sinh}[x]^2)/a^2]) \\
& - (10 \cdot a \cdot \text{ArcSin}[\text{Sqrt}[(a + b) \cdot \text{Cosh}[x]^2)/a] \cdot \text{Sech}[x]^2)/((a + b) \cdot \text{Sqrt}[-((a + b) \cdot \text{Cosh}[x]^2 \cdot (a + b) \cdot \text{Coth}[x]^2 \cdot \text{Sinh}[x]^2)/a^2])) \cdot \text{Tanh}[x]/(a^2 \cdot \text{Sqrt}[a + b \cdot \text{Coth}[x]^2] \cdot (1 + (b \cdot \text{Coth}[x]^2)/a))
\end{aligned}$$

Maple [F]

$$\int \frac{\tanh(x)^2}{(a + b \coth(x)^2)^{\frac{5}{2}}} dx$$

[In] int(tanh(x)^2/(a+b*coth(x)^2)^(5/2),x)

[Out] int(tanh(x)^2/(a+b*coth(x)^2)^(5/2),x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5085 vs. 2(113) = 226.

Time = 1.30 (sec) , antiderivative size = 10729, normalized size of antiderivative = 81.90

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{\frac{5}{2}}} dx = \text{Too large to display}$$

[In] integrate(tanh(x)^2/(a+b*coth(x)^2)^(5/2),x, algorithm="fricas")

[Out] Too large to include

Sympy [F]

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{\frac{5}{2}}} dx = \int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{\frac{5}{2}}} dx$$

[In] integrate(tanh(x)**2/(a+b*coth(x)**2)**(5/2),x)

[Out] Integral(tanh(x)**2/(a + b*coth(x)**2)**(5/2), x)

Maxima [F]

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{\frac{5}{2}}} dx = \int \frac{\tanh(x)^2}{(b \coth(x)^2 + a)^{\frac{5}{2}}} dx$$

[In] integrate(tanh(x)^2/(a+b*coth(x)^2)^(5/2),x, algorithm="maxima")

[Out] integrate(tanh(x)^2/(b*coth(x)^2 + a)^(5/2), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1133 vs. $2(113) = 226$.

Time = 0.77 (sec) , antiderivative size = 1133, normalized size of antiderivative = 8.65

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \text{Too large to display}$$

[In] integrate(tanh(x)^2/(a+b*coth(x)^2)^(5/2),x, algorithm="giac")

[Out]
$$-1/3 * (((9*a^{13}*b^4 + 50*a^{12}*b^5 + 115*a^{11}*b^6 + 140*a^{10}*b^7 + 95*a^9*b^8 + 34*a^8*b^9 + 5*a^7*b^{10}) * e^{(2*x)} / (a^{16}*b^2*sgn(e^{(2*x)} - 1) + 6*a^{15}*b^3*sgn(e^{(2*x)} - 1) + 15*a^{14}*b^4*sgn(e^{(2*x)} - 1) + 20*a^{13}*b^5*sgn(e^{(2*x)} - 1) + 15*a^{12}*b^6*sgn(e^{(2*x)} - 1) + 6*a^{11}*b^7*sgn(e^{(2*x)} - 1) + a^{10}*b^8*sgn(e^{(2*x)} - 1)) - 3*(3*a^{13}*b^4 + 6*a^{12}*b^5 - 11*a^{11}*b^6 - 44*a^{10}*b^7 - 51*a^9*b^8 - 26*a^8*b^9 - 5*a^7*b^{10}) / (a^{16}*b^2*sgn(e^{(2*x)} - 1) + 6*a^{15}*b^3*sgn(e^{(2*x)} - 1) + 15*a^{14}*b^4*sgn(e^{(2*x)} - 1) + 20*a^{13}*b^5*sgn(e^{(2*x)} - 1) + 15*a^{12}*b^6*sgn(e^{(2*x)} - 1) + 6*a^{11}*b^7*sgn(e^{(2*x)} - 1) + a^{10}*b^8*sgn(e^{(2*x)} - 1))) * e^{(2*x)} - 3*(3*a^{13}*b^4 + 6*a^{12}*b^5 - 11*a^{11}*b^6 - 44*a^{10}*b^7 - 51*a^9*b^8 - 26*a^8*b^9 - 5*a^7*b^{10}) / (a^{16}*b^2*sgn(e^{(2*x)} - 1) + 6*a^{15}*b^3*sgn(e^{(2*x)} - 1) + 15*a^{14}*b^4*sgn(e^{(2*x)} - 1) + 20*a^{13}*b^5*sgn(e^{(2*x)} - 1) + 15*a^{12}*b^6*sgn(e^{(2*x)} - 1) + 6*a^{11}*b^7*sgn(e^{(2*x)} - 1) + a^{10}*b^8*sgn(e^{(2*x)} - 1))) * e^{(2*x)} + (9*a^{13}*b^4 + 50*a^{12}*b^5 + 115*a^{11}*b^6 + 140*a^{10}*b^7 + 95*a^9*b^8 + 34*a^8*b^9 + 5*a^7*b^{10}) / (a^{16}*b^2*sgn(e^{(2*x)} - 1) + 6*a^{15}*b^3*sgn(e^{(2*x)} - 1) + 15*a^{14}*b^4*sgn(e^{(2*x)} - 1) + 20*a^{13}*b^5*sgn(e^{(2*x)} - 1) + 15*a^{12}*b^6*sgn(e^{(2*x)} - 1) + 6*a^{11}*b^7*sgn(e^{(2*x)} - 1) + a^{10}*b^8*sgn(e^{(2*x)} - 1))) / (a*e^{(4*x)} + b*e^{(4*x)} - 2*a*e^{(2*x)} + 2*b*e^{(2*x)} + a + b)^{(3/2)} - 1/2*log(abs((sqrt(a + b)*e^{(2*x)} - sqrt(a*e^{(4*x)} + b*e^{(4*x)} - 2*a*e^{(2*x)} + 2*b*e^{(2*x)} + a + b)) * sqrt(a + b) - a + b)) / ((a^2 + 2*a*b + b^2)*sqrt(a + b)*sgn(e^{(2*x)} - 1)) + 1/2*log(abs((sqrt(a + b)*e^{(2*x)} - sqrt(a*e^{(4*x)} + b*e^{(4*x)} - 2*a*e^{(2*x)} + 2*b*e^{(2*x)} + a + b)) * sqrt(a + b) - a - b)) / ((a^2 + 2*a*b + b^2)*sqrt(a + b)*sgn(e^{(2*x)} - 1)) - 1/2*log(abs(-sqrt(a + b)*e^{(2*x)} + sqrt(a*e^{(4*x)} + b*e^{(4*x)} - 2*a*e^{(2*x)} + 2*b*e^{(2*x)} + a + b) - sqrt(a + b))) / ((a^2 + 2*a*b + b^2)*sqrt(a + b)*sgn(e^{(2*x)} - 1)) - 4*(sqrt(a + b)*e^{(2*x)} - sqrt(a*e^{(4*x)} + b*e^{(4*x)} - 2*a*e^{(2*x)} + 2*b*e^{(2*x)} + a + b) - sqrt(a + b)) / ((sqrt(a + b)*e^{(2*x)} - sqrt(a*e^{(4*x)} + b*e^{(4*x)} - 2*a*e^{(2*x)} + 2*b*e^{(2*x)} + a + b))^2 + 2*(sqrt(a + b)*e^{(2*x)} - sqrt(a*e^{(4*x)} + b*e^{(4*x)} - 2*a*e^{(2*x)} + 2*b*e^{(2*x)} + a + b)) * sqrt(a + b) - 3*a + b) * a^2 * sgn(e^{(2*x)} - 1))$$

Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh^2(x)}{(a + b \coth^2(x))^{5/2}} dx = \int \frac{\tanh(x)^2}{(b \coth(x)^2 + a)^{5/2}} dx$$

```
[In] int(tanh(x)^2/(a + b*coth(x)^2)^(5/2), x)
```

```
[Out] int(tanh(x)^2/(a + b*coth(x)^2)^(5/2), x)
```

$$3.48 \quad \int \frac{1}{\sqrt{1+\coth^2(x)}} dx$$

Optimal result	330
Rubi [A] (verified)	330
Mathematica [A] (verified)	331
Maple [B] (verified)	332
Fricas [B] (verification not implemented)	332
Sympy [F]	333
Maxima [F]	333
Giac [B] (verification not implemented)	333
Mupad [B] (verification not implemented)	334

Optimal result

Integrand size = 10, antiderivative size = 25

$$\int \frac{1}{\sqrt{1+\coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}\coth(x)}{\sqrt{1+\coth^2(x)}}\right)}{\sqrt{2}}$$

[Out] 1/2*arctanh(coth(x)*2^(1/2)/(1+coth(x)^2)^(1/2))*2^(1/2)

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3742, 385, 212}

$$\int \frac{1}{\sqrt{1+\coth^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}\coth(x)}{\sqrt{\coth^2(x)+1}}\right)}{\sqrt{2}}$$

[In] Int[1/Sqrt[1 + Coth[x]^2],x]

[Out] ArcTanh[(Sqrt[2]*Coth[x])/Sqrt[1 + Coth[x]^2]]/Sqrt[2]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst} \left(\int \frac{1}{(1-x^2)\sqrt{1+x^2}} dx, x, \coth(x) \right) \\ &= \text{Subst} \left(\int \frac{1}{1-2x^2} dx, x, \frac{\coth(x)}{\sqrt{1+\coth^2(x)}} \right) \\ &= \frac{\operatorname{arctanh} \left(\frac{\sqrt{2}\coth(x)}{\sqrt{1+\coth^2(x)}} \right)}{\sqrt{2}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1+\coth^2(x)}} dx = \frac{\operatorname{arctanh} \left(\frac{\sqrt{2}\coth(x)}{\sqrt{1+\coth^2(x)}} \right)}{\sqrt{2}}$$

```
[In] Integrate[1/Sqrt[1 + Coth[x]^2], x]
```

```
[Out] ArcTanh[(Sqrt[2]*Coth[x])/Sqrt[1 + Coth[x]^2]]/Sqrt[2]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs. $2(20) = 40$.

Time = 0.20 (sec) , antiderivative size = 62, normalized size of antiderivative = 2.48

method	result	size
derivativedivides	$-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(2-2\coth(x))\sqrt{2}}{4\sqrt{(1+\coth(x))^2-2\coth(x)}}\right)}{4} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(2+2\coth(x))\sqrt{2}}{4\sqrt{(\coth(x)-1)^2+2\coth(x)}}\right)}{4}$	62
default	$-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(2-2\coth(x))\sqrt{2}}{4\sqrt{(1+\coth(x))^2-2\coth(x)}}\right)}{4} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(2+2\coth(x))\sqrt{2}}{4\sqrt{(\coth(x)-1)^2+2\coth(x)}}\right)}{4}$	62

[In] `int(1/(1+coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/4*2^{(1/2)}*\operatorname{arctanh}(1/4*(2-2*\coth(x))*2^{(1/2)}/((1+\coth(x))^2-2*\coth(x))^{(1/2)})+1/4*2^{(1/2)}*\operatorname{arctanh}(1/4*(2+2*\coth(x))*2^{(1/2)}/((\coth(x)-1)^2+2*\coth(x))^{(1/2)})$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 547 vs. $2(20) = 40$.

Time = 0.25 (sec) , antiderivative size = 547, normalized size of antiderivative = 21.88

$$\int \frac{1}{\sqrt{1 + \coth^2(x)}} dx = \text{Too large to display}$$

[In] `integrate(1/(1+coth(x)^2)^(1/2),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & 1/8*\sqrt{2}*\log(2*(\cosh(x)^8 + 8*\cosh(x)*\sinh(x)^7 + \sinh(x)^8 + (28*\cosh(x)^2 + 3)*\sinh(x)^6 + 3*\cosh(x)^6 + 2*(28*\cosh(x)^3 + 9*\cosh(x))*\sinh(x)^5 + \\ & 5*(14*\cosh(x)^4 + 9*\cosh(x)^2 + 1)*\sinh(x)^4 + 5*\cosh(x)^4 + 4*(14*\cosh(x)^5 + 15*\cosh(x)^3 + 5*\cosh(x))*\sinh(x)^3 + (28*\cosh(x)^6 + 45*\cosh(x)^4 + 30*\cosh(x)^2 + 4)*\sinh(x)^2 + 4*\cosh(x)^2 + 2*(4*\cosh(x)^7 + 9*\cosh(x)^5 + 10*\cosh(x)^3 + 4*\cosh(x))*\sinh(x) + (\sqrt{2}*\cosh(x)^6 + 6*\sqrt{2}*\cosh(x)*\sinh(x)^5 + \sqrt{2}*\sinh(x)^6 + 3*(5*\sqrt{2}*\cosh(x)^2 + \sqrt{2})*\sinh(x)^4 + 3*\sqrt{2}*\cosh(x)^4 + 4*(5*\sqrt{2}*\cosh(x)^3 + 3*\sqrt{2}*\cosh(x))*\sinh(x)^3 + (15*\sqrt{2}*\cosh(x)^4 + 18*\sqrt{2}*\cosh(x)^2 + 4*\sqrt{2})*\sinh(x)^2 + 4*\sqrt{2}*\cosh(x)^2 + 2*(3*\sqrt{2}*\cosh(x)^5 + 6*\sqrt{2}*\cosh(x)^3 + 4*\sqrt{2}*\cosh(x))*\sinh(x) + 4*\sqrt{2})*\sqrt{((\cosh(x)^2 + \sinh(x)^2)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)) + 4}/(\cosh(x)^6 + 6*\cosh(x)^5*\sinh(x) + 15*\cosh(x)^4*\sinh(x)^2 + 20*\cosh(x)^3*\sinh(x)^3 + 15*\cosh(x)^2*\sinh(x)^4 + 6*\cosh(x)*\sinh(x)^5 + \sinh(x)^6)) + 1/8*\sqrt{2}*\log(-2*(\cosh(x)^4 + 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4 + (6*\cosh(x)^2 - 1)*\sinh(x)^2 - \cosh(x)^2 + 2*(2*\cosh(x)^3 - \cosh(x))*\sinh(x) + (\sqrt{2}*\cosh(x)^2 + 2*\sqrt{2}*\cosh(x)*\sinh(x) \end{aligned}$$

+ sqrt(2)*sinh(x)^2 - sqrt(2))*sqrt((cosh(x)^2 + sinh(x)^2)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2) + 1)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2))

Sympy [F]

$$\int \frac{1}{\sqrt{1 + \coth^2(x)}} dx = \int \frac{1}{\sqrt{\coth^2(x) + 1}} dx$$

[In] integrate(1/(1+coth(x)**2)**(1/2),x)

[Out] Integral(1/sqrt(coth(x)**2 + 1), x)

Maxima [F]

$$\int \frac{1}{\sqrt{1 + \coth^2(x)}} dx = \int \frac{1}{\sqrt{\coth(x)^2 + 1}} dx$$

[In] integrate(1/(1+coth(x)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(coth(x)^2 + 1), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 69 vs. 2(20) = 40.

Time = 0.26 (sec) , antiderivative size = 69, normalized size of antiderivative = 2.76

$$\int \frac{1}{\sqrt{1 + \coth^2(x)}} dx = \frac{\sqrt{2} \left(\log \left(\sqrt{e^{4x} + 1} - e^{2x} + 1 \right) - \log \left(\sqrt{e^{4x} + 1} - e^{2x} \right) - \log \left(-\sqrt{e^{4x} + 1} + e^{2x} + 1 \right) \right)}{4 \operatorname{sgn}(e^{2x} - 1)}$$

[In] integrate(1/(1+coth(x)^2)^(1/2),x, algorithm="giac")

[Out] 1/4*sqrt(2)*(log(sqrt(e^(4*x) + 1) - e^(2*x) + 1) - log(sqrt(e^(4*x) + 1) - e^(2*x)) - log(-sqrt(e^(4*x) + 1) + e^(2*x) + 1))/sgn(e^(2*x) - 1)

Mupad [B] (verification not implemented)

Time = 2.10 (sec) , antiderivative size = 63, normalized size of antiderivative = 2.52

$$\int \frac{1}{\sqrt{1 + \coth^2(x)}} dx = \frac{\sqrt{2} \left(\ln \left(\coth(x) + \sqrt{2} \sqrt{\coth(x)^2 + 1} + 1 \right) - \ln(\coth(x) - 1) \right)}{4} + \frac{\sqrt{2} \left(\ln(\coth(x) + 1) - \ln \left(\sqrt{2} \sqrt{\coth(x)^2 + 1} - \coth(x) + 1 \right) \right)}{4}$$

`[In] int(1/(coth(x)^2 + 1)^(1/2),x)`

```
[Out] (2^(1/2)*(log(coth(x) + 2^(1/2)*(coth(x)^2 + 1)^(1/2) + 1) - log(coth(x) - 1)))/4 + (2^(1/2)*(log(coth(x) + 1) - log(2^(1/2)*(coth(x)^2 + 1)^(1/2) - coth(x) + 1)))/4
```

$$3.49 \quad \int \frac{1}{\sqrt{-1 - \coth^2(x)}} dx$$

Optimal result	335
Rubi [A] (verified)	335
Mathematica [A] (warning: unable to verify)	336
Maple [B] (verified)	337
Fricas [C] (verification not implemented)	337
Sympy [F]	338
Maxima [F]	338
Giac [C] (verification not implemented)	338
Mupad [B] (verification not implemented)	339

Optimal result

Integrand size = 12, antiderivative size = 27

$$\int \frac{1}{\sqrt{-1 - \coth^2(x)}} dx = \frac{\arctan\left(\frac{\sqrt{2} \coth(x)}{\sqrt{-1 - \coth^2(x)}}\right)}{\sqrt{2}}$$

[Out] 1/2*arctan(coth(x)*2^(1/2)/(-1-coth(x)^2)^(1/2))*2^(1/2)

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3742, 385, 209}

$$\int \frac{1}{\sqrt{-1 - \coth^2(x)}} dx = \frac{\arctan\left(\frac{\sqrt{2} \coth(x)}{\sqrt{-\coth^2(x) - 1}}\right)}{\sqrt{2}}$$

[In] Int[1/Sqrt[-1 - Coth[x]^2],x]

[Out] ArcTan[(Sqrt[2]*Coth[x])/Sqrt[-1 - Coth[x]^2]]/Sqrt[2]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst} \left(\int \frac{1}{\sqrt{-1-x^2}(1-x^2)} dx, x, \coth(x) \right) \\ &= \text{Subst} \left(\int \frac{1}{1+2x^2} dx, x, \frac{\coth(x)}{\sqrt{-1-\coth^2(x)}} \right) \\ &= \frac{\arctan \left(\frac{\sqrt{2}\coth(x)}{\sqrt{-1-\coth^2(x)}} \right)}{\sqrt{2}} \end{aligned}$$

Mathematica [A] (warning: unable to verify)

Time = 0.08 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.81

$$\int \frac{1}{\sqrt{-1-\coth^2(x)}} dx = \frac{\text{arcsinh} \left(\frac{\sqrt{2}\coth(x)}{\sqrt{1-\coth^2(x)}} \right) \sqrt{1+\coth^2(x)}}{\sqrt{2}\sqrt{-1-\coth^2(x)}}$$

```
[In] Integrate[1/Sqrt[-1 - Coth[x]^2], x]
```

```
[Out] (ArcSinh[(Sqrt[2]*Coth[x])/Sqrt[1 - Coth[x]^2]]*Sqrt[1 + Coth[x]^2])/(Sqrt[2]*Sqrt[-1 - Coth[x]^2])
```


Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. $2(22) = 44$.

Time = 0.20 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.44

method	result	size
derivativedivides	$-\frac{\sqrt{2} \arctan\left(\frac{(-2-2 \coth(x))\sqrt{2}}{4\sqrt{-(\coth(x)-1)^2-2 \coth(x)}}\right)}{4} + \frac{\sqrt{2} \arctan\left(\frac{(-2+2 \coth(x))\sqrt{2}}{4\sqrt{-(1+\coth(x))^2+2 \coth(x)}}\right)}{4}$	66
default	$-\frac{\sqrt{2} \arctan\left(\frac{(-2-2 \coth(x))\sqrt{2}}{4\sqrt{-(\coth(x)-1)^2-2 \coth(x)}}\right)}{4} + \frac{\sqrt{2} \arctan\left(\frac{(-2+2 \coth(x))\sqrt{2}}{4\sqrt{-(1+\coth(x))^2+2 \coth(x)}}\right)}{4}$	66

[In] `int(1/(-1-coth(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/4*2^{(1/2)}*\arctan(1/4*(-2-2*\coth(x))*2^{(1/2)/(-(\coth(x)-1)^2-2*\coth(x))^{(1/2)})+1/4*2^{(1/2)}*\arctan(1/4*(-2+2*\coth(x))*2^{(1/2)/(-(1+\coth(x))^2+2*\coth(x))^{(1/2)})}$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 175, normalized size of antiderivative = 6.48

$$\begin{aligned} & \int \frac{1}{\sqrt{-1 - \coth^2(x)}} dx \\ &= \frac{1}{8}i\sqrt{2} \log\left(\frac{1}{2}\left(i\sqrt{2}\sqrt{-2e^{4x}-2} + 2e^{2x}-2\right)e^{(-2x)}\right) \\ & \quad - \frac{1}{8}i\sqrt{2} \log\left(\frac{1}{2}\left(-i\sqrt{2}\sqrt{-2e^{4x}-2} + 2e^{2x}-2\right)e^{(-2x)}\right) \\ & \quad - \frac{1}{8}i\sqrt{2} \log\left(\left(\sqrt{-2e^{4x}-2}(e^{2x}+2) + i\sqrt{2}e^{4x} + i\sqrt{2}e^{2x} + 2i\sqrt{2}\right)e^{(-4x)}\right) \\ & \quad + \frac{1}{8}i\sqrt{2} \log\left(\left(\sqrt{-2e^{4x}-2}(e^{2x}+2) - i\sqrt{2}e^{4x} - i\sqrt{2}e^{2x} - 2i\sqrt{2}\right)e^{(-4x)}\right) \end{aligned}$$

[In] `integrate(1/(-1-coth(x)^2)^(1/2),x, algorithm="fricas")`

[Out]
$$1/8*I*\sqrt{2}*\log(1/2*(I*\sqrt{2}*\sqrt{-2*e^{4*x}-2} + 2*e^{2*x}-2)*e^{(-2*x)}) - 1/8*I*\sqrt{2}*\log(1/2*(-I*\sqrt{2}*\sqrt{-2*e^{4*x}-2} + 2*e^{2*x}-2)*e^{(-2*x)}) - 1/8*I*\sqrt{2}*\log((\sqrt{-2*e^{4*x}-2}*(e^{2*x}+2) + I*\sqrt{2}*e^{4*x} + I*\sqrt{2}*e^{2*x} + 2*I*\sqrt{2})*e^{(-4*x)}) + 1/8*I*\sqrt{2}*\log((\sqrt{-2*e^{4*x}-2}*(e^{2*x}+2) - I*\sqrt{2}*e^{4*x} - I*\sqrt{2}*e^{2*x} - 2*I*\sqrt{2})*e^{(-4*x)})}$$

Sympy [F]

$$\int \frac{1}{\sqrt{-1 - \coth^2(x)}} dx = \int \frac{1}{\sqrt{-\coth^2(x) - 1}} dx$$

```
[In] integrate(1/(-1-coth(x)**2)**(1/2),x)
```

```
[Out] Integral(1/sqrt(-coth(x)**2 - 1), x)
```

Maxima [F]

$$\int \frac{1}{\sqrt{-1 - \coth^2(x)}} dx = \int \frac{1}{\sqrt{-\coth^2(x) - 1}} dx$$

```
[In] integrate(1/(-1-coth(x)^2)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(1/sqrt(-coth(x)^2 - 1), x)
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.70

$$\int \frac{1}{\sqrt{-1 - \coth^2(x)}} dx = \frac{\sqrt{2} \left(-i \log \left(\sqrt{e^{(4x)} + 1} - e^{(2x)} + 1 \right) + i \log \left(\sqrt{e^{(4x)} + 1} - e^{(2x)} \right) + i \log \left(-\sqrt{e^{(4x)} + 1} + e^{(2x)} + 1 \right) \right)}{4 \operatorname{sgn}(-e^{(2x)} + 1)}$$

```
[In] integrate(1/(-1-coth(x)^2)^(1/2),x, algorithm="giac")
```

```
[Out] -1/4*sqrt(2)*(-I*log(sqrt(e^(4*x) + 1) - e^(2*x) + 1) + I*log(sqrt(e^(4*x) + 1) - e^(2*x)) + I*log(-sqrt(e^(4*x) + 1) + e^(2*x) + 1))/sgn(-e^(2*x) + 1)
```

Mupad [B] (verification not implemented)

Time = 2.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int \frac{1}{\sqrt{-1 - \coth^2(x)}} dx = \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2} \coth(x)}{\sqrt{-\coth(x)^2 - 1}}\right)}{2}$$

[In] `int(1/(- coth(x)^2 - 1)^(1/2),x)`

[Out] `(2^(1/2)*atan((2^(1/2)*coth(x))/(- coth(x)^2 - 1)^(1/2)))/2`

3.50 $\int \frac{1}{1+\coth^3(x)} dx$

Optimal result	340
Rubi [A] (verified)	340
Mathematica [A] (verified)	342
Maple [A] (verified)	342
Fricas [B] (verification not implemented)	342
Sympy [B] (verification not implemented)	343
Maxima [B] (verification not implemented)	343
Giac [A] (verification not implemented)	344
Mupad [B] (verification not implemented)	344

Optimal result

Integrand size = 8, antiderivative size = 38

$$\int \frac{1}{1+\coth^3(x)} dx = \frac{x}{2} - \frac{2 \arctan\left(\frac{1-2\coth(x)}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{1}{6(1+\coth(x))}$$

[Out] 1/2*x-1/6/(1+coth(x))-2/9*arctan(1/3*(1-2*coth(x))*3^(1/2))*3^(1/2)

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3742, 2099, 213, 632, 210}

$$\int \frac{1}{1+\coth^3(x)} dx = -\frac{2 \arctan\left(\frac{1-2\coth(x)}{\sqrt{3}}\right)}{3\sqrt{3}} + \frac{x}{2} - \frac{1}{6(\coth(x)+1)}$$

[In] Int[(1 + Coth[x]^3)^(-1), x]

[Out] x/2 - (2*ArcTan[(1 - 2*Coth[x])/Sqrt[3]])/(3*Sqrt[3]) - 1/(6*(1 + Coth[x]))

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] &&

(LtQ[a, 0] || GtQ[b, 0])

Rule 632

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 2099

Int[(P_)^(p_)*(Q_)^(q_), x_Symbol] := With[{PP = Factor[P]}, Int[ExpandIntegrand[PP^p*Q^q, x], x] /; !SumQ[NonfreeFactors[PP, x]]] /; FreeQ[q, x] && PolyQ[P, x] && PolyQ[Q, x] && IntegerQ[p] && NeQ[P, x]

Rule 3742

Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegerQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{1}{(1-x^2)(1+x^3)} dx, x, \coth(x)\right) \\
 &= \text{Subst}\left(\int \left(\frac{1}{6(1+x)^2} - \frac{1}{2(-1+x^2)} + \frac{1}{3(1-x+x^2)}\right) dx, x, \coth(x)\right) \\
 &= -\frac{1}{6(1+\coth(x))} + \frac{1}{3}\text{Subst}\left(\int \frac{1}{1-x+x^2} dx, x, \coth(x)\right) \\
 &\quad - \frac{1}{2}\text{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \coth(x)\right) \\
 &= \frac{x}{2} - \frac{1}{6(1+\coth(x))} - \frac{2}{3}\text{Subst}\left(\int \frac{1}{-3-x^2} dx, x, -1+2\coth(x)\right) \\
 &= \frac{x}{2} - \frac{2\arctan\left(\frac{1-2\coth(x)}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{1}{6(1+\coth(x))}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.05

$$\int \frac{1}{1 + \coth^3(x)} dx = \frac{2 \arctan\left(\frac{1-2\tanh(x)}{\sqrt{3}}\right)}{3\sqrt{3}} + \frac{1}{2} \operatorname{arctanh}(\tanh(x)) + \frac{1}{6(1 + \tanh(x))}$$

[In] Integrate[(1 + Coth[x]^3)^(-1), x]

[Out] (2*ArcTan[(1 - 2*Tanh[x])/Sqrt[3]])/(3*Sqrt[3]) + ArcTanh[Tanh[x]]/2 + 1/(6*(1 + Tanh[x]))

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.08

method	result	size
derivativedivides	$-\frac{1}{6(1+\coth(x))} + \frac{\ln(1+\coth(x))}{4} - \frac{\ln(\coth(x)-1)}{4} + \frac{2\sqrt{3} \arctan\left(\frac{(2\coth(x)-1)\sqrt{3}}{3}\right)}{9}$	41
default	$-\frac{1}{6(1+\coth(x))} + \frac{\ln(1+\coth(x))}{4} - \frac{\ln(\coth(x)-1)}{4} + \frac{2\sqrt{3} \arctan\left(\frac{(2\coth(x)-1)\sqrt{3}}{3}\right)}{9}$	41
risch	$\frac{x}{2} + \frac{e^{-2x}}{12} + \frac{i\sqrt{3} \ln(e^{2x}-i\sqrt{3})}{9} - \frac{i\sqrt{3} \ln(e^{2x}+i\sqrt{3})}{9}$	47

[In] int(1/(1+coth(x)^3),x,method=_RETURNVERBOSE)

[Out] -1/6/(1+coth(x))+1/4*ln(1+coth(x))-1/4*ln(coth(x)-1)+2/9*3^(1/2)*arctan(1/3*(2*coth(x)-1)*3^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs. 2(29) = 58.

Time = 0.27 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.50

$$\int \frac{1}{1 + \coth^3(x)} dx = \frac{18x \cosh(x)^2 + 36x \cosh(x) \sinh(x) + 18x \sinh(x)^2 + 8(\sqrt{3} \cosh(x)^2 + 2\sqrt{3} \cosh(x) \sinh(x) + \sqrt{3} \sinh(x)^2) \arctan\left(\frac{-1/3(\sqrt{3} \cosh(x) + \sqrt{3} \sinh(x))}{\cosh(x) - \sinh(x)}\right) + 3}{36(\cosh(x)^2 + 2\cosh(x) \sinh(x) + \sinh(x)^2)}$$

[In] integrate(1/(1+coth(x)^3),x, algorithm="fricas")

[Out] 1/36*(18*x*cosh(x)^2 + 36*x*cosh(x)*sinh(x) + 18*x*sinh(x)^2 + 8*(sqrt(3)*cosh(x)^2 + 2*sqrt(3)*cosh(x)*sinh(x) + sqrt(3)*sinh(x)^2)*arctan(-1/3*(sqrt(3)*cosh(x) + sqrt(3)*sinh(x))/(cosh(x) - sinh(x))) + 3)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 102 vs. $2(36) = 72$.

Time = 0.50 (sec) , antiderivative size = 102, normalized size of antiderivative = 2.68

$$\int \frac{1}{1 + \coth^3(x)} dx = \frac{9x \tanh(x)}{18 \tanh(x) + 18} + \frac{9x}{18 \tanh(x) + 18} - \frac{4\sqrt{3} \tanh(x) \operatorname{atan}\left(\frac{2\sqrt{3} \tanh(x)}{3} - \frac{\sqrt{3}}{3}\right)}{18 \tanh(x) + 18} - \frac{4\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3} \tanh(x)}{3} - \frac{\sqrt{3}}{3}\right)}{18 \tanh(x) + 18} + \frac{3}{18 \tanh(x) + 18}$$

[In] integrate(1/(1+coth(x)**3),x)

[Out] $9*x*\tanh(x)/(18*\tanh(x) + 18) + 9*x/(18*\tanh(x) + 18) - 4*\sqrt{3}*\tanh(x)*\operatorname{atan}(2*\sqrt{3}*\tanh(x)/3 - \sqrt{3}/3)/(18*\tanh(x) + 18) - 4*\sqrt{3}*\operatorname{atan}(2*\sqrt{3}*\tanh(x)/3 - \sqrt{3}/3)/(18*\tanh(x) + 18) + 3/(18*\tanh(x) + 18)$

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. $2(29) = 58$.

Time = 0.29 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.92

$$\int \frac{1}{1 + \coth^3(x)} dx = -\frac{2}{9} \sqrt{3} \arctan\left(\frac{1}{6} \cdot 3^{\frac{3}{4}} \sqrt{2} \left(2\sqrt{3}e^{-x} + 3^{\frac{1}{4}} \sqrt{2}\right)\right) + \frac{2}{9} \sqrt{3} \arctan\left(\frac{1}{6} \cdot 3^{\frac{3}{4}} \sqrt{2} \left(2\sqrt{3}e^{-x} - 3^{\frac{1}{4}} \sqrt{2}\right)\right) + \frac{1}{2} x + \frac{1}{12} e^{(-2x)}$$

[In] integrate(1/(1+coth(x)^3),x, algorithm="maxima")

[Out] $-2/9*\sqrt{3}*\arctan(1/6*3^{(3/4)}*\sqrt{2}*(2*\sqrt{3}*e^{-x} + 3^{(1/4)}*\sqrt{2})) + 2/9*\sqrt{3}*\arctan(1/6*3^{(3/4)}*\sqrt{2}*(2*\sqrt{3}*e^{-x} - 3^{(1/4)}*\sqrt{2})) + 1/2*x + 1/12*e^{(-2*x)}$

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.66

$$\int \frac{1}{1 + \coth^3(x)} dx = -\frac{2}{9} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3} e^{2x}\right) + \frac{1}{2} x + \frac{1}{12} e^{(-2x)}$$

[In] integrate(1/(1+coth(x)^3),x, algorithm="giac")

[Out] -2/9*sqrt(3)*arctan(1/3*sqrt(3)*e^(2*x)) + 1/2*x + 1/12*e^(-2*x)

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \coth^3(x)} dx = \frac{\frac{x}{2} + \frac{\coth(x)}{6} + \frac{x \coth(x)}{2}}{\coth(x) + 1} + \frac{2\sqrt{3} \operatorname{atan}\left(\frac{\sqrt{3}(2\coth(x)-1)}{3}\right)}{9}$$

[In] int(1/(coth(x)^3 + 1),x)

[Out] (x/2 + coth(x)/6 + (x*coth(x))/2)/(coth(x) + 1) + (2*3^(1/2)*atan((3^(1/2)*(2*coth(x) - 1))/3))/9

3.51 $\int \coth(x) \sqrt{a + b \coth^4(x)} dx$

Optimal result	345
Rubi [A] (verified)	345
Mathematica [A] (verified)	348
Maple [A] (verified)	348
Fricas [B] (verification not implemented)	349
Sympy [F]	349
Maxima [F]	349
Giac [F]	349
Mupad [F(-1)]	350

Optimal result

Integrand size = 15, antiderivative size = 89

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx = -\frac{1}{2} \sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{b} \coth^2(x)}{\sqrt{a + b \coth^4(x)}} \right) + \frac{1}{2} \sqrt{a + b} \operatorname{arctanh} \left(\frac{a + b \coth^2(x)}{\sqrt{a + b} \sqrt{a + b \coth^4(x)}} \right) - \frac{1}{2} \sqrt{a + b \coth^4(x)}$$

[Out] $-1/2*\operatorname{arctanh}(\coth(x)^2*b^{(1/2)}/(a+b*\coth(x)^4)^{(1/2)})*b^{(1/2)}+1/2*\operatorname{arctanh}((a+b*\coth(x)^2)/(a+b)^{(1/2)}/(a+b*\coth(x)^4)^{(1/2)})*(a+b)^{(1/2)}-1/2*(a+b*\coth(x)^4)^{(1/2)}$

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3751, 1262, 749, 858, 223, 212, 739}

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx = -\frac{1}{2} \sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{b} \coth^2(x)}{\sqrt{a + b \coth^4(x)}} \right) + \frac{1}{2} \sqrt{a + b} \operatorname{arctanh} \left(\frac{a + b \coth^2(x)}{\sqrt{a + b} \sqrt{a + b \coth^4(x)}} \right) - \frac{1}{2} \sqrt{a + b \coth^4(x)}$$

[In] Int[Coth[x]*Sqrt[a + b*Coth[x]^4], x]

[Out] $-\frac{1}{2}(\sqrt{b} \operatorname{ArcTanh}[\frac{\sqrt{b} \operatorname{Coth}[x]^2}{\sqrt{a + b \operatorname{Coth}[x]^4}}]) + (\sqrt{a + b} \operatorname{ArcTanh}[\frac{a + b \operatorname{Coth}[x]^2}{\sqrt{a + b} \sqrt{a + b \operatorname{Coth}[x]^4}}]))/2 - \operatorname{Sqrt}[a + b \operatorname{Coth}[x]^4]/2$

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 223

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 739

Int[1/(((d_) + (e_.)*(x_))*Sqrt[(a_) + (c_.)*(x_)^2]), x_Symbol] := -Subst[Int[1/(c*d^2 + a*e^2 - x^2), x], x, (a*e - c*d*x)/Sqrt[a + c*x^2]] /; FreeQ[{a, c, d, e}, x]

Rule 749

Int[((d_) + (e_.)*(x_))^(m_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + c*x^2)^p/(e*(m + 2*p + 1))), x] + Dist[2*(p/(e*(m + 2*p + 1))), Int[(d + e*x)^m*Simp[a*e - c*d*x, x]*(a + c*x^2)^(p - 1), x], x] /; FreeQ[{a, c, d, e, m}, x] && NeQ[c*d^2 + a*e^2, 0] && GtQ[p, 0] && NeQ[m + 2*p + 1, 0] && (!RationalQ[m] || LtQ[m, 1]) && !ILtQ[m + 2*p, 0] && IntQuadraticQ[a, 0, c, d, e, m, p, x]

Rule 858

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g, m, p}, x] && NeQ[c*d^2 + a*e^2, 0] && !IGtQ[m, 0]

Rule 1262

Int[(x_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_.), x_Symbol] := Dist[1/2, Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, c, d, e, p, q}, x]

Rule 3751

Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_.))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],

`x]], Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))`

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst} \left(\int \frac{x\sqrt{a+bx^4}}{1-x^2} dx, x, \coth(x) \right) \\
&= \frac{1}{2} \text{Subst} \left(\int \frac{\sqrt{a+bx^2}}{1-x} dx, x, \coth^2(x) \right) \\
&= -\frac{1}{2} \sqrt{a+b\coth^4(x)} - \frac{1}{2} \text{Subst} \left(\int \frac{-a-bx}{(1-x)\sqrt{a+bx^2}} dx, x, \coth^2(x) \right) \\
&= -\frac{1}{2} \sqrt{a+b\coth^4(x)} - \frac{1}{2}(-a-b) \text{Subst} \left(\int \frac{1}{(1-x)\sqrt{a+bx^2}} dx, x, \coth^2(x) \right) \\
&\quad - \frac{1}{2} b \text{Subst} \left(\int \frac{1}{\sqrt{a+bx^2}} dx, x, \coth^2(x) \right) \\
&= -\frac{1}{2} \sqrt{a+b\coth^4(x)} - \frac{1}{2} b \text{Subst} \left(\int \frac{1}{1-bx^2} dx, x, \frac{\coth^2(x)}{\sqrt{a+b\coth^4(x)}} \right) \\
&\quad - \frac{1}{2}(a+b) \text{Subst} \left(\int \frac{1}{a+b-x^2} dx, x, \frac{-a-b\coth^2(x)}{\sqrt{a+b\coth^4(x)}} \right) \\
&= -\frac{1}{2} \sqrt{b} \text{arctanh} \left(\frac{\sqrt{b}\coth^2(x)}{\sqrt{a+b\coth^4(x)}} \right) \\
&\quad + \frac{1}{2} \sqrt{a+b} \text{arctanh} \left(\frac{a+b\coth^2(x)}{\sqrt{a+b}\sqrt{a+b\coth^4(x)}} \right) - \frac{1}{2} \sqrt{a+b\coth^4(x)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.97

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx = \frac{1}{2} \left(-\sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{b} \coth^2(x)}{\sqrt{a + b \coth^4(x)}} \right) + \sqrt{a + b} \operatorname{arctanh} \left(\frac{a + b \coth^2(x)}{\sqrt{a + b} \sqrt{a + b \coth^4(x)}} \right) - \sqrt{a + b \coth^4(x)} \right)$$

[In] Integrate[Coth[x]*Sqrt[a + b*Coth[x]^4],x]

[Out] $(-\sqrt{b} \operatorname{ArcTanh}[(\sqrt{b} \operatorname{Coth}[x]^2)/\sqrt{a + b \operatorname{Coth}[x]^4}]) + \sqrt{a + b} \operatorname{ArcTanh}[(a + b \operatorname{Coth}[x]^2)/(\sqrt{a + b} \sqrt{a + b \operatorname{Coth}[x]^4})] - \sqrt{a + b \operatorname{Coth}[x]^4})/2$

Maple [A] (verified)

Time = 0.82 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.30

method	result
derivativedivides	$-\frac{\sqrt{a+b \coth(x)^4}}{2} - \frac{\sqrt{b} \ln(2\sqrt{b} \coth(x)^2 + 2\sqrt{a+b \coth(x)^4})}{2} + \frac{b \operatorname{arctanh}\left(\frac{2b \coth(x)^2 + 2a}{2\sqrt{a+b} \sqrt{a+b \coth(x)^4}}\right)}{2\sqrt{a+b}} + \frac{a \operatorname{arctanh}\left(\frac{a + b \coth^2(x)}{\sqrt{a+b} \sqrt{a+b \coth^4(x)}}\right)}{2}$
default	$-\frac{\sqrt{a+b \coth(x)^4}}{2} - \frac{\sqrt{b} \ln(2\sqrt{b} \coth(x)^2 + 2\sqrt{a+b \coth(x)^4})}{2} + \frac{b \operatorname{arctanh}\left(\frac{2b \coth(x)^2 + 2a}{2\sqrt{a+b} \sqrt{a+b \coth(x)^4}}\right)}{2\sqrt{a+b}} + \frac{a \operatorname{arctanh}\left(\frac{a + b \coth^2(x)}{\sqrt{a+b} \sqrt{a+b \coth^4(x)}}\right)}{2}$

[In] int(coth(x)*(a+b*coth(x)^4)^(1/2),x,method=_RETURNVERBOSE)

[Out] $-1/2*(a+b*\coth(x)^4)^(1/2)-1/2*b^(1/2)*\ln(2*b^(1/2)*\coth(x)^2+2*(a+b*\coth(x)^4)^(1/2))+1/2*b/(a+b)^(1/2)*\operatorname{arctanh}(1/2*(2*b*\coth(x)^2+2*a)/(a+b)^(1/2)/(a+b*\coth(x)^4)^(1/2))+1/2*a/(a+b)^(1/2)*\operatorname{arctanh}(1/2*(2*b*\coth(x)^2+2*a)/(a+b)^(1/2)/(a+b*\coth(x)^4)^(1/2))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1053 vs. 2(69) = 138.
 Time = 0.40 (sec) , antiderivative size = 5172, normalized size of antiderivative = 58.11

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx = \text{Too large to display}$$

[In] `integrate(coth(x)*(a+b*coth(x)^4)^(1/2),x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx = \int \sqrt{a + b \coth^4(x)} \coth(x) dx$$

[In] `integrate(coth(x)*(a+b*coth(x)**4)**(1/2),x)`

[Out] `Integral(sqrt(a + b*coth(x)**4)*coth(x), x)`

Maxima [F]

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx = \int \sqrt{b \coth^4(x) + a} \coth(x) dx$$

[In] `integrate(coth(x)*(a+b*coth(x)^4)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(b*coth(x)^4 + a)*coth(x), x)`

Giac [F]

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx = \int \sqrt{b \coth^4(x) + a} \coth(x) dx$$

[In] `integrate(coth(x)*(a+b*coth(x)^4)^(1/2),x, algorithm="giac")`

[Out] `integrate(sqrt(b*coth(x)^4 + a)*coth(x), x)`

Mupad [F(-1)]

Timed out.

$$\int \coth(x) \sqrt{a + b \coth^4(x)} dx = \int \coth(x) \sqrt{b \coth^4(x) + a} dx$$

```
[In] int(coth(x)*(a + b*coth(x)^4)^(1/2),x)
```

```
[Out] int(coth(x)*(a + b*coth(x)^4)^(1/2), x)
```

$$3.52 \quad \int \frac{\coth(x)}{\sqrt{a+b \coth^4(x)}} dx$$

Optimal result	351
Rubi [A] (verified)	351
Mathematica [A] (verified)	353
Maple [A] (verified)	353
Fricas [B] (verification not implemented)	353
Sympy [F]	354
Maxima [F]	355
Giac [F]	355
Mupad [F(-1)]	355

Optimal result

Integrand size = 15, antiderivative size = 40

$$\int \frac{\coth(x)}{\sqrt{a+b \coth^4(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{a+b \coth^2(x)}{\sqrt{a+b} \sqrt{a+b \coth^4(x)}}\right)}{2\sqrt{a+b}}$$

[Out] 1/2*arctanh((a+b*coth(x)^2)/(a+b)^(1/2)/(a+b*coth(x)^4)^(1/2))/(a+b)^(1/2)

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3751, 1262, 739, 212}

$$\int \frac{\coth(x)}{\sqrt{a+b \coth^4(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{a+b \coth^2(x)}{\sqrt{a+b} \sqrt{a+b \coth^4(x)}}\right)}{2\sqrt{a+b}}$$

[In] Int[Coth[x]/Sqrt[a + b*Coth[x]^4], x]

[Out] ArcTanh[(a + b*Coth[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Coth[x]^4])]/(2*Sqrt[a + b])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt

$Q[a, 0] \parallel LtQ[b, 0]$)

Rule 739

```
Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (c_)*(x_)^2]), x_Symbol] := -Subst[
Int[1/(c*d^2 + a*e^2 - x^2), x], x, (a*e - c*d*x)/Sqrt[a + c*x^2]] /; FreeQ
[{a, c, d, e}, x]
```

Rule 1262

```
Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol]
:= Dist[1/2, Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ
[{a, c, d, e, p, q}, x]
```

Rule 3751

```
Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) +
(f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst} \left(\int \frac{x}{(1-x^2)\sqrt{a+bx^4}} dx, x, \coth(x) \right) \\
&= \frac{1}{2} \text{Subst} \left(\int \frac{1}{(1-x)\sqrt{a+bx^2}} dx, x, \coth^2(x) \right) \\
&= - \left(\frac{1}{2} \text{Subst} \left(\int \frac{1}{a+b-x^2} dx, x, \frac{-a-b\coth^2(x)}{\sqrt{a+b\coth^4(x)}} \right) \right) \\
&= \frac{\text{arctanh} \left(\frac{a+b\coth^2(x)}{\sqrt{a+b}\sqrt{a+b\coth^4(x)}} \right)}{2\sqrt{a+b}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^4(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{a + b \coth^2(x)}{\sqrt{a+b}\sqrt{a + b \coth^4(x)}}\right)}{2\sqrt{a+b}}$$

[In] Integrate[Coth[x]/Sqrt[a + b*Coth[x]^4],x]

[Out] ArcTanh[(a + b*Coth[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Coth[x]^4])]/(2*Sqrt[a + b])

Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.92

method	result	size
derivativedivides	$\frac{\operatorname{arctanh}\left(\frac{2b \coth(x)^2 + 2a}{2\sqrt{a+b}\sqrt{a + b \coth(x)^4}}\right)}{2\sqrt{a+b}}$	37
default	$\frac{\operatorname{arctanh}\left(\frac{2b \coth(x)^2 + 2a}{2\sqrt{a+b}\sqrt{a + b \coth(x)^4}}\right)}{2\sqrt{a+b}}$	37

[In] int(coth(x)/(a+b*coth(x)^4)^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/2/(a+b)^(1/2)*arctanh(1/2*(2*b*coth(x)^2+2*a)/(a+b)^(1/2)/(a+b*coth(x)^4)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 626 vs. 2(32) = 64.

Time = 0.41 (sec) , antiderivative size = 1290, normalized size of antiderivative = 32.25

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^4(x)}} dx = \text{Too large to display}$$

[In] integrate(coth(x)/(a+b*coth(x)^4)^(1/2),x, algorithm="fricas")

[Out] [1/4*log(((a^2 + 2*a*b + b^2)*cosh(x)^8 + 8*(a^2 + 2*a*b + b^2)*cosh(x)*sinh(x)^7 + (a^2 + 2*a*b + b^2)*sinh(x)^8 - 4*(a^2 - b^2)*cosh(x)^6 + 4*(7*(a^2 + 2*a*b + b^2)*cosh(x)^2 - a^2 + b^2)*sinh(x)^6 + 8*(7*(a^2 + 2*a*b + b^2)*cosh(x)^3 - 3*(a^2 - b^2)*cosh(x))*sinh(x)^5 + 2*(3*a^2 + 2*a*b + 3*b^2)*

```

cosh(x)^4 + 2*(35*(a^2 + 2*a*b + b^2)*cosh(x)^4 - 30*(a^2 - b^2)*cosh(x)^2
+ 3*a^2 + 2*a*b + 3*b^2)*sinh(x)^4 + 8*(7*(a^2 + 2*a*b + b^2)*cosh(x)^5 - 1
0*(a^2 - b^2)*cosh(x)^3 + (3*a^2 + 2*a*b + 3*b^2)*cosh(x))*sinh(x)^3 - 4*(a
^2 - b^2)*cosh(x)^2 + 4*(7*(a^2 + 2*a*b + b^2)*cosh(x)^6 - 15*(a^2 - b^2)*c
osh(x)^4 + 3*(3*a^2 + 2*a*b + 3*b^2)*cosh(x)^2 - a^2 + b^2)*sinh(x)^2 + sqr
t(2)*((a + b)*cosh(x)^4 + 4*(a + b)*cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 -
2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a +
b)*cosh(x)^3 - (a - b)*cosh(x))*sinh(x) + a + b)*sqrt(a + b)*sqrt(((a + b)*
cosh(x)^4 + (a + b)*sinh(x)^4 - 4*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^
2 - 2*a + 2*b)*sinh(x)^2 + 3*a + 3*b)/(cosh(x)^4 - 4*cosh(x)^3*sinh(x) + 6*
cosh(x)^2*sinh(x)^2 - 4*cosh(x)*sinh(x)^3 + sinh(x)^4)) + a^2 + 2*a*b + b^2
+ 8*((a^2 + 2*a*b + b^2)*cosh(x)^7 - 3*(a^2 - b^2)*cosh(x)^5 + (3*a^2 + 2*
a*b + 3*b^2)*cosh(x)^3 - (a^2 - b^2)*cosh(x))*sinh(x))/(cosh(x)^4 + 4*cosh(
x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4))/sq
rt(a + b), -1/2*sqrt(-a - b)*arctan(sqrt(2)*((a + b)*cosh(x)^4 + 4*(a + b)*
cosh(x)*sinh(x)^3 + (a + b)*sinh(x)^4 - 2*(a - b)*cosh(x)^2 + 2*(3*(a + b)*
cosh(x)^2 - a + b)*sinh(x)^2 + 4*((a + b)*cosh(x)^3 - (a - b)*cosh(x))*sinh
(x) + a + b)*sqrt(-a - b)*sqrt(((a + b)*cosh(x)^4 + (a + b)*sinh(x)^4 - 4*(
a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - 2*a + 2*b)*sinh(x)^2 + 3*a + 3*
b)/(cosh(x)^4 - 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 - 4*cosh(x)*sin
h(x)^3 + sinh(x)^4))/((a^2 + 2*a*b + b^2)*cosh(x)^8 + 8*(a^2 + 2*a*b + b^2)
*cosh(x)*sinh(x)^7 + (a^2 + 2*a*b + b^2)*sinh(x)^8 - 4*(a^2 - b^2)*cosh(x)^
6 + 4*(7*(a^2 + 2*a*b + b^2)*cosh(x)^2 - a^2 + b^2)*sinh(x)^6 + 8*(7*(a^2 +
2*a*b + b^2)*cosh(x)^3 - 3*(a^2 - b^2)*cosh(x))*sinh(x)^5 + 6*(a^2 + 2*a*b
+ b^2)*cosh(x)^4 + 2*(35*(a^2 + 2*a*b + b^2)*cosh(x)^4 - 30*(a^2 - b^2)*co
sh(x)^2 + 3*a^2 + 6*a*b + 3*b^2)*sinh(x)^4 + 8*(7*(a^2 + 2*a*b + b^2)*cosh(
x)^5 - 10*(a^2 - b^2)*cosh(x)^3 + 3*(a^2 + 2*a*b + b^2)*cosh(x))*sinh(x)^3
- 4*(a^2 - b^2)*cosh(x)^2 + 4*(7*(a^2 + 2*a*b + b^2)*cosh(x)^6 - 15*(a^2 -
b^2)*cosh(x)^4 + 9*(a^2 + 2*a*b + b^2)*cosh(x)^2 - a^2 + b^2)*sinh(x)^2 + a
^2 + 2*a*b + b^2 + 8*((a^2 + 2*a*b + b^2)*cosh(x)^7 - 3*(a^2 - b^2)*cosh(x)
^5 + 3*(a^2 + 2*a*b + b^2)*cosh(x)^3 - (a^2 - b^2)*cosh(x))*sinh(x)))/(a +
b)]

```

Sympy [F]

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^4(x)}} dx = \int \frac{\coth(x)}{\sqrt{a + b \coth^4(x)}} dx$$

[In] integrate(coth(x)/(a+b*coth(x)**4)**(1/2),x)

[Out] Integral(coth(x)/sqrt(a + b*coth(x)**4), x)

Maxima [F]

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^4(x)}} dx = \int \frac{\coth(x)}{\sqrt{b \coth(x)^4 + a}} dx$$

[In] integrate(coth(x)/(a+b*coth(x)^4)^(1/2),x, algorithm="maxima")

[Out] integrate(coth(x)/sqrt(b*coth(x)^4 + a), x)

Giac [F]

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^4(x)}} dx = \int \frac{\coth(x)}{\sqrt{b \coth(x)^4 + a}} dx$$

[In] integrate(coth(x)/(a+b*coth(x)^4)^(1/2),x, algorithm="giac")

[Out] integrate(coth(x)/sqrt(b*coth(x)^4 + a), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\coth(x)}{\sqrt{a + b \coth^4(x)}} dx = \int \frac{\coth(x)}{\sqrt{b \coth(x)^4 + a}} dx$$

[In] int(coth(x)/(a + b*coth(x)^4)^(1/2),x)

[Out] int(coth(x)/(a + b*coth(x)^4)^(1/2), x)

3.53 $\int \frac{\coth(x)}{(a+b\coth^4(x))^{3/2}} dx$

Optimal result	356
Rubi [A] (verified)	356
Mathematica [A] (verified)	358
Maple [C] (verified)	358
Fricas [B] (verification not implemented)	359
Sympy [F]	361
Maxima [F]	362
Giac [F]	362
Mupad [F(-1)]	362

Optimal result

Integrand size = 15, antiderivative size = 74

$$\int \frac{\coth(x)}{(a+b\coth^4(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{a+b\coth^2(x)}{\sqrt{a+b}\sqrt{a+b\coth^4(x)}}\right)}{2(a+b)^{3/2}} - \frac{a-b\coth^2(x)}{2a(a+b)\sqrt{a+b\coth^4(x)}}$$

[Out] $1/2*\operatorname{arctanh}((a+b*\coth(x)^2)/(a+b)^{(1/2)/(a+b*\coth(x)^4)^{(1/2))}/(a+b)^{(3/2)+1/2*(-a+b*\coth(x)^2)/a/(a+b)/(a+b*\coth(x)^4)^{(1/2)}$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3751, 1262, 755, 12, 739, 212}

$$\int \frac{\coth(x)}{(a+b\coth^4(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{a+b\coth^2(x)}{\sqrt{a+b}\sqrt{a+b\coth^4(x)}}\right)}{2(a+b)^{3/2}} - \frac{a-b\coth^2(x)}{2a(a+b)\sqrt{a+b\coth^4(x)}}$$

[In] $\operatorname{Int}[\operatorname{Coth}[x]/(a+b*\operatorname{Coth}[x]^4)^{(3/2)}, x]$

[Out] $\operatorname{ArcTanh}[(a+b*\operatorname{Coth}[x]^2)/(\operatorname{Sqrt}[a+b]*\operatorname{Sqrt}[a+b*\operatorname{Coth}[x]^4])]/(2*(a+b)^{(3/2)}) - (a-b*\operatorname{Coth}[x]^2)/(2*a*(a+b)*\operatorname{Sqrt}[a+b*\operatorname{Coth}[x]^4])$

Rule 12

$\operatorname{Int}[(a_*)*(u_), x_Symbol] \rightarrow \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \&\& \operatorname{!Match} Q[u, (b_*)*(v_)] /; \operatorname{FreeQ}[b, x]$

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 739

Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (c_)*(x_)^2]), x_Symbol] := -Subst[Int[1/(c*d^2 + a*e^2 - x^2), x], x, (a*e - c*d*x)/Sqrt[a + c*x^2]] /; FreeQ[{a, c, d, e}, x]

Rule 755

Int[((d_) + (e_)*(x_))^(m_)*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(-(d + e*x)^(m + 1))*(a*e + c*d*x)*((a + c*x^2)^(p + 1)/(2*a*(p + 1)*(c*d^2 + a*e^2))), x] + Dist[1/(2*a*(p + 1)*(c*d^2 + a*e^2)), Int[(d + e*x)^m*Sim p[c*d^2*(2*p + 3) + a*e^2*(m + 2*p + 3) + c*e*d*(m + 2*p + 4)*x, x]*(a + c*x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e, m}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && IntQuadraticQ[a, 0, c, d, e, m, p, x]

Rule 1262

Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Dist[1/2, Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, c, d, e, p, q}, x]

Rule 3751

Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst} \left(\int \frac{x}{(1-x^2)(a+bx^4)^{3/2}} dx, x, \coth(x) \right) \\
 &= \frac{1}{2} \text{Subst} \left(\int \frac{1}{(1-x)(a+bx^2)^{3/2}} dx, x, \coth^2(x) \right) \\
 &= -\frac{a-b\coth^2(x)}{2a(a+b)\sqrt{a+b\coth^4(x)}} + \frac{\text{Subst} \left(\int \frac{a}{(1-x)\sqrt{a+bx^2}} dx, x, \coth^2(x) \right)}{2a(a+b)}
 \end{aligned}$$

$$\begin{aligned}
 &= -\frac{a - b \operatorname{coth}^2(x)}{2a(a + b)\sqrt{a + b \operatorname{coth}^4(x)}} + \frac{\operatorname{Subst}\left(\int \frac{1}{(1-x)\sqrt{a+bx^2}} dx, x, \operatorname{coth}^2(x)\right)}{2(a + b)} \\
 &= -\frac{a - b \operatorname{coth}^2(x)}{2a(a + b)\sqrt{a + b \operatorname{coth}^4(x)}} - \frac{\operatorname{Subst}\left(\int \frac{1}{a+b-x^2} dx, x, \frac{-a-b \operatorname{coth}^2(x)}{\sqrt{a+b \operatorname{coth}^4(x)}}\right)}{2(a + b)} \\
 &= \frac{\operatorname{arctanh}\left(\frac{a+b \operatorname{coth}^2(x)}{\sqrt{a+b}\sqrt{a+b \operatorname{coth}^4(x)}}\right)}{2(a + b)^{3/2}} - \frac{a - b \operatorname{coth}^2(x)}{2a(a + b)\sqrt{a + b \operatorname{coth}^4(x)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.53 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.99

$$\int \frac{\operatorname{coth}(x)}{(a + b \operatorname{coth}^4(x))^{3/2}} dx = \frac{1}{2} \left(\frac{\operatorname{arctanh}\left(\frac{a+b \operatorname{coth}^2(x)}{\sqrt{a+b}\sqrt{a+b \operatorname{coth}^4(x)}}\right)}{(a + b)^{3/2}} - \frac{a - b \operatorname{coth}^2(x)}{a(a + b)\sqrt{a + b \operatorname{coth}^4(x)}} \right)$$

[In] Integrate[Coth[x]/(a + b*Coth[x]^4)^(3/2), x]

[Out] (ArcTanh[(a + b*Coth[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Coth[x]^4])]/(a + b)^(3/2) - (a - b*Coth[x]^2)/(a*(a + b)*Sqrt[a + b*Coth[x]^4]))/2

Maple [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 0.67 (sec) , antiderivative size = 431, normalized size of antiderivative = 5.82

method	result
derivativedivides	$ \frac{b\left(-\frac{\operatorname{coth}(x)^3}{4a(a+b)} + \frac{\operatorname{coth}(x)^2}{4a(a+b)} - \frac{\operatorname{coth}(x)}{4a(a+b)} - \frac{1}{4(a+b)b}\right)}{\sqrt{\left(\operatorname{coth}(x)^4 + \frac{a}{b}\right)b}} - \frac{\operatorname{arctanh}\left(\frac{2b \operatorname{coth}(x)^2 + 2a}{2\sqrt{a+b}\sqrt{a+b \operatorname{coth}(x)^4}}\right) + \frac{\sqrt{1 - \frac{i\sqrt{b} \operatorname{coth}(x)^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{b} \operatorname{coth}(x)^2}{\sqrt{a}}}}{2\sqrt{a+b}}}{2(a+b)} $
default	$ \frac{b\left(-\frac{\operatorname{coth}(x)^3}{4a(a+b)} + \frac{\operatorname{coth}(x)^2}{4a(a+b)} - \frac{\operatorname{coth}(x)}{4a(a+b)} - \frac{1}{4(a+b)b}\right)}{\sqrt{\left(\operatorname{coth}(x)^4 + \frac{a}{b}\right)b}} - \frac{\operatorname{arctanh}\left(\frac{2b \operatorname{coth}(x)^2 + 2a}{2\sqrt{a+b}\sqrt{a+b \operatorname{coth}(x)^4}}\right) + \frac{\sqrt{1 - \frac{i\sqrt{b} \operatorname{coth}(x)^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{b} \operatorname{coth}(x)^2}{\sqrt{a}}}}{2\sqrt{a+b}}}{2(a+b)} $

[In] `int(coth(x)/(a+b*coth(x)^4)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$b*(-1/4/a/(a+b)*\coth(x)^3+1/4/a/(a+b)*\coth(x)^2-1/4/a/(a+b)*\coth(x)-1/4/(a+b)/b)/((\coth(x)^4+a/b)*b)^{(1/2)}-1/2/(a+b)*(-1/2/(a+b))^{(1/2)}*\operatorname{arctanh}(1/2*(2*b*\coth(x)^2+2*a)/(a+b))^{(1/2)}/(a+b*\coth(x)^4)^{(1/2)}+1/(I/a^{(1/2)}*b^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*b^{(1/2)}*\coth(x)^2)^{(1/2)}*(1+I/a^{(1/2)}*b^{(1/2)}*\coth(x)^2)^{(1/2)}/(a+b*\coth(x)^4)^{(1/2)}*\operatorname{EllipticPi}(\coth(x)*(I/a^{(1/2)}*b^{(1/2)})^{(1/2)},-I*a^{(1/2)}/b^{(1/2)},(-I/a^{(1/2)}*b^{(1/2)})^{(1/2)}/(I/a^{(1/2)}*b^{(1/2)})^{(1/2)})))+b*(1/4/a/(a+b)*\coth(x)^3+1/4/a/(a+b)*\coth(x)^2+1/4/a/(a+b)*\coth(x)-1/4/(a+b)/b)/((\coth(x)^4+a/b)*b)^{(1/2)}-1/2/(a+b)*(-1/2/(a+b))^{(1/2)}*\operatorname{arctanh}(1/2*(2*b*\coth(x)^2+2*a)/(a+b))^{(1/2)}/(a+b*\coth(x)^4)^{(1/2)}-1/(I/a^{(1/2)}*b^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*b^{(1/2)}*\coth(x)^2)^{(1/2)}*(1+I/a^{(1/2)}*b^{(1/2)}*\coth(x)^2)^{(1/2)}/(a+b*\coth(x)^4)^{(1/2)}*\operatorname{EllipticPi}(\coth(x)*(I/a^{(1/2)}*b^{(1/2)})^{(1/2)},-I*a^{(1/2)}/b^{(1/2)},(-I/a^{(1/2)}*b^{(1/2)})^{(1/2)}/(I/a^{(1/2)}*b^{(1/2)})^{(1/2)}))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1947 vs. 2(63) = 126.

Time = 0.50 (sec) , antiderivative size = 3938, normalized size of antiderivative = 53.22

$$\int \frac{\coth(x)}{(a + b \coth^4(x))^{3/2}} dx = \text{Too large to display}$$

[In] `integrate(coth(x)/(a+b*coth(x)^4)^(3/2),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & [1/4*((a^2 + a*b)*\cosh(x)^8 + 8*(a^2 + a*b)*\cosh(x)*\sinh(x)^7 + (a^2 + a*b) \\ &)*\sinh(x)^8 - 4*(a^2 - a*b)*\cosh(x)^6 + 4*(7*(a^2 + a*b)*\cosh(x)^2 - a^2 + \\ & a*b)*\sinh(x)^6 + 8*(7*(a^2 + a*b)*\cosh(x)^3 - 3*(a^2 - a*b)*\cosh(x))*\sinh(x) \\ &)^5 + 6*(a^2 + a*b)*\cosh(x)^4 + 2*(35*(a^2 + a*b)*\cosh(x)^4 - 30*(a^2 - a*b) \\ &)*\cosh(x)^2 + 3*a^2 + 3*a*b)*\sinh(x)^4 + 8*(7*(a^2 + a*b)*\cosh(x)^5 - 10*(a \\ & ^2 - a*b)*\cosh(x)^3 + 3*(a^2 + a*b)*\cosh(x))*\sinh(x)^3 - 4*(a^2 - a*b)*\cosh \\ & (x)^2 + 4*(7*(a^2 + a*b)*\cosh(x)^6 - 15*(a^2 - a*b)*\cosh(x)^4 + 9*(a^2 + a* \\ & b)*\cosh(x)^2 - a^2 + a*b)*\sinh(x)^2 + a^2 + a*b + 8*((a^2 + a*b)*\cosh(x)^7 \\ & - 3*(a^2 - a*b)*\cosh(x)^5 + 3*(a^2 + a*b)*\cosh(x)^3 - (a^2 - a*b)*\cosh(x))* \\ & \sinh(x))*\sqrt{a + b}*\log(((a^2 + 2*a*b + b^2)*\cosh(x)^8 + 8*(a^2 + 2*a*b + \\ & b^2)*\cosh(x)*\sinh(x)^7 + (a^2 + 2*a*b + b^2)*\sinh(x)^8 - 4*(a^2 - b^2)*\cosh \\ & (x)^6 + 4*(7*(a^2 + 2*a*b + b^2)*\cosh(x)^2 - a^2 + b^2)*\sinh(x)^6 + 8*(7*(a \\ & ^2 + 2*a*b + b^2)*\cosh(x)^3 - 3*(a^2 - b^2)*\cosh(x))*\sinh(x)^5 + 2*(3*a^2 + \\ & 2*a*b + 3*b^2)*\cosh(x)^4 + 2*(35*(a^2 + 2*a*b + b^2)*\cosh(x)^4 - 30*(a^2 - \\ & b^2)*\cosh(x)^2 + 3*a^2 + 2*a*b + 3*b^2)*\sinh(x)^4 + 8*(7*(a^2 + 2*a*b + b^ \\ & 2)*\cosh(x)^5 - 10*(a^2 - b^2)*\cosh(x)^3 + (3*a^2 + 2*a*b + 3*b^2)*\cosh(x))* \\ & \sinh(x)^3 - 4*(a^2 - b^2)*\cosh(x)^2 + 4*(7*(a^2 + 2*a*b + b^2)*\cosh(x)^6 - \\ & 15*(a^2 - b^2)*\cosh(x)^4 + 3*(3*a^2 + 2*a*b + 3*b^2)*\cosh(x)^2 - a^2 + b^2) \\ &)*\sinh(x)^2 + \sqrt{2}*((a + b)*\cosh(x)^4 + 4*(a + b)*\cosh(x)*\sinh(x)^3 + (a \\ & + b)*\sinh(x)^4 - 2*(a - b)*\cosh(x)^2 + 2*(3*(a + b)*\cosh(x)^2 - a + b)*\sinh \end{aligned}$$

$$\begin{aligned}
& (x)^2 + 4*((a + b)*\cosh(x)^3 - (a - b)*\cosh(x))*\sinh(x) + a + b)*\sqrt{a + b} \\
&)*\sqrt{((a + b)*\cosh(x)^4 + (a + b)*\sinh(x)^4 - 4*(a - b)*\cosh(x)^2 + 2*(3* \\
& (a + b)*\cosh(x)^2 - 2*a + 2*b)*\sinh(x)^2 + 3*a + 3*b)/(\cosh(x)^4 - 4*\cosh(x) \\
&)^3*\sinh(x) + 6*\cosh(x)^2*\sinh(x)^2 - 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4)) + a \\
& ^2 + 2*a*b + b^2 + 8*((a^2 + 2*a*b + b^2)*\cosh(x)^7 - 3*(a^2 - b^2)*\cosh(x) \\
& ^5 + (3*a^2 + 2*a*b + 3*b^2)*\cosh(x)^3 - (a^2 - b^2)*\cosh(x))*\sinh(x))/(\cos \\
& h(x)^4 + 4*\cosh(x)^3*\sinh(x) + 6*\cosh(x)^2*\sinh(x)^2 + 4*\cosh(x)*\sinh(x)^3 \\
& + \sinh(x)^4)) - 2*\sqrt{2}*((a^2 - b^2)*\cosh(x)^4 + 4*(a^2 - b^2)*\cosh(x)*\si \\
& nh(x)^3 + (a^2 - b^2)*\sinh(x)^4 - 2*(a^2 + 2*a*b + b^2)*\cosh(x)^2 + 2*(3*(a \\
& ^2 - b^2)*\cosh(x)^2 - a^2 - 2*a*b - b^2)*\sinh(x)^2 + a^2 - b^2 + 4*((a^2 - \\
& b^2)*\cosh(x)^3 - (a^2 + 2*a*b + b^2)*\cosh(x))*\sinh(x))*\sqrt{((a + b)*\cosh(x) \\
&)^4 + (a + b)*\sinh(x)^4 - 4*(a - b)*\cosh(x)^2 + 2*(3*(a + b)*\cosh(x)^2 - 2* \\
& a + 2*b)*\sinh(x)^2 + 3*a + 3*b)/(\cosh(x)^4 - 4*\cosh(x)^3*\sinh(x) + 6*\cosh(x) \\
&)^2*\sinh(x)^2 - 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4)))/((a^4 + 3*a^3*b + 3*a^2*b^2 \\
& + a*b^3)*\cosh(x)^8 + 8*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*\cosh(x)*\sinh \\
& (x)^7 + (a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*\sinh(x)^8 - 4*(a^4 + a^3*b - a^2 \\
& *b^2 - a*b^3)*\cosh(x)^6 - 4*(a^4 + a^3*b - a^2*b^2 - a*b^3 - 7*(a^4 + 3*a^3 \\
& *b + 3*a^2*b^2 + a*b^3)*\cosh(x)^2)*\sinh(x)^6 + 8*(7*(a^4 + 3*a^3*b + 3*a^2 \\
& *b^2 + a*b^3)*\cosh(x)^3 - 3*(a^4 + a^3*b - a^2*b^2 - a*b^3)*\cosh(x))*\sinh(x) \\
&)^5 + 6*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*\cosh(x)^4 + 2*(35*(a^4 + 3*a^3*b \\
& + 3*a^2*b^2 + a*b^3)*\cosh(x)^4 + 3*a^4 + 9*a^3*b + 9*a^2*b^2 + 3*a*b^3 - \\
& 30*(a^4 + a^3*b - a^2*b^2 - a*b^3)*\cosh(x)^2)*\sinh(x)^4 + a^4 + 3*a^3*b + 3 \\
& *a^2*b^2 + a*b^3 + 8*(7*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*\cosh(x)^5 - 10* \\
& (a^4 + a^3*b - a^2*b^2 - a*b^3)*\cosh(x)^3 + 3*(a^4 + 3*a^3*b + 3*a^2*b^2 + \\
& a*b^3)*\cosh(x))*\sinh(x)^3 - 4*(a^4 + a^3*b - a^2*b^2 - a*b^3)*\cosh(x)^2 + 4 \\
& *(7*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*\cosh(x)^6 - 15*(a^4 + a^3*b - a^2*b \\
& ^2 - a*b^3)*\cosh(x)^4 - a^4 - a^3*b + a^2*b^2 + a*b^3 + 9*(a^4 + 3*a^3*b + \\
& 3*a^2*b^2 + a*b^3)*\cosh(x)^2)*\sinh(x)^2 + 8*((a^4 + 3*a^3*b + 3*a^2*b^2 + a \\
& *b^3)*\cosh(x)^7 - 3*(a^4 + a^3*b - a^2*b^2 - a*b^3)*\cosh(x)^5 + 3*(a^4 + 3* \\
& a^3*b + 3*a^2*b^2 + a*b^3)*\cosh(x)^3 - (a^4 + a^3*b - a^2*b^2 - a*b^3)*\cosh \\
& (x))*\sinh(x)), -1/2*((a^2 + a*b)*\cosh(x)^8 + 8*(a^2 + a*b)*\cosh(x)*\sinh(x) \\
& ^7 + (a^2 + a*b)*\sinh(x)^8 - 4*(a^2 - a*b)*\cosh(x)^6 + 4*(7*(a^2 + a*b)*\cos \\
& h(x)^2 - a^2 + a*b)*\sinh(x)^6 + 8*(7*(a^2 + a*b)*\cosh(x)^3 - 3*(a^2 - a*b)* \\
& \cosh(x))*\sinh(x)^5 + 6*(a^2 + a*b)*\cosh(x)^4 + 2*(35*(a^2 + a*b)*\cosh(x)^4 \\
& - 30*(a^2 - a*b)*\cosh(x)^2 + 3*a^2 + 3*a*b)*\sinh(x)^4 + 8*(7*(a^2 + a*b)*\cos \\
& sh(x)^5 - 10*(a^2 - a*b)*\cosh(x)^3 + 3*(a^2 + a*b)*\cosh(x))*\sinh(x)^3 - 4*(\\
& a^2 - a*b)*\cosh(x)^2 + 4*(7*(a^2 + a*b)*\cosh(x)^6 - 15*(a^2 - a*b)*\cosh(x)^ \\
& 4 + 9*(a^2 + a*b)*\cosh(x)^2 - a^2 + a*b)*\sinh(x)^2 + a^2 + a*b + 8*((a^2 + \\
& a*b)*\cosh(x)^7 - 3*(a^2 - a*b)*\cosh(x)^5 + 3*(a^2 + a*b)*\cosh(x)^3 - (a^2 - \\
& a*b)*\cosh(x))*\sinh(x))*\sqrt{-a - b}*\arctan(\sqrt{2}*((a + b)*\cosh(x)^4 + 4* \\
& (a + b)*\cosh(x)*\sinh(x)^3 + (a + b)*\sinh(x)^4 - 2*(a - b)*\cosh(x)^2 + 2*(3* \\
& (a + b)*\cosh(x)^2 - a + b)*\sinh(x)^2 + 4*((a + b)*\cosh(x)^3 - (a - b)*\cosh(\\
& x))*\sinh(x) + a + b)*\sqrt{-a - b}*\sqrt{((a + b)*\cosh(x)^4 + (a + b)*\sinh(x) \\
&)^4 - 4*(a - b)*\cosh(x)^2 + 2*(3*(a + b)*\cosh(x)^2 - 2*a + 2*b)*\sinh(x)^2 + \\
& 3*a + 3*b)/(\cosh(x)^4 - 4*\cosh(x)^3*\sinh(x) + 6*\cosh(x)^2*\sinh(x)^2 - 4*\cos
\end{aligned}$$


```

h(x)*sinh(x)^3 + sinh(x)^4))/((a^2 + 2*a*b + b^2)*cosh(x)^8 + 8*(a^2 + 2*a*
b + b^2)*cosh(x)*sinh(x)^7 + (a^2 + 2*a*b + b^2)*sinh(x)^8 - 4*(a^2 - b^2)*
cosh(x)^6 + 4*(7*(a^2 + 2*a*b + b^2)*cosh(x)^2 - a^2 + b^2)*sinh(x)^6 + 8*(
7*(a^2 + 2*a*b + b^2)*cosh(x)^3 - 3*(a^2 - b^2)*cosh(x))*sinh(x)^5 + 6*(a^2
+ 2*a*b + b^2)*cosh(x)^4 + 2*(35*(a^2 + 2*a*b + b^2)*cosh(x)^4 - 30*(a^2 -
b^2)*cosh(x)^2 + 3*a^2 + 6*a*b + 3*b^2)*sinh(x)^4 + 8*(7*(a^2 + 2*a*b + b^
2)*cosh(x)^5 - 10*(a^2 - b^2)*cosh(x)^3 + 3*(a^2 + 2*a*b + b^2)*cosh(x))*si
nh(x)^3 - 4*(a^2 - b^2)*cosh(x)^2 + 4*(7*(a^2 + 2*a*b + b^2)*cosh(x)^6 - 15
*(a^2 - b^2)*cosh(x)^4 + 9*(a^2 + 2*a*b + b^2)*cosh(x)^2 - a^2 + b^2)*sinh(
x)^2 + a^2 + 2*a*b + b^2 + 8*((a^2 + 2*a*b + b^2)*cosh(x)^7 - 3*(a^2 - b^2)
*cosh(x)^5 + 3*(a^2 + 2*a*b + b^2)*cosh(x)^3 - (a^2 - b^2)*cosh(x))*sinh(x)
)) + sqrt(2)*((a^2 - b^2)*cosh(x)^4 + 4*(a^2 - b^2)*cosh(x)*sinh(x)^3 + (a^
2 - b^2)*sinh(x)^4 - 2*(a^2 + 2*a*b + b^2)*cosh(x)^2 + 2*(3*(a^2 - b^2)*cos
h(x)^2 - a^2 - 2*a*b - b^2)*sinh(x)^2 + a^2 - b^2 + 4*((a^2 - b^2)*cosh(x)^
3 - (a^2 + 2*a*b + b^2)*cosh(x))*sinh(x))*sqrt(((a + b)*cosh(x)^4 + (a + b)
*sinh(x)^4 - 4*(a - b)*cosh(x)^2 + 2*(3*(a + b)*cosh(x)^2 - 2*a + 2*b)*sinh
(x)^2 + 3*a + 3*b)/(cosh(x)^4 - 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2
- 4*cosh(x)*sinh(x)^3 + sinh(x)^4)))/((a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*
cosh(x)^8 + 8*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)*sinh(x)^7 + (a^4
+ 3*a^3*b + 3*a^2*b^2 + a*b^3)*sinh(x)^8 - 4*(a^4 + a^3*b - a^2*b^2 - a*b^3
)*cosh(x)^6 - 4*(a^4 + a^3*b - a^2*b^2 - a*b^3 - 7*(a^4 + 3*a^3*b + 3*a^2*b
^2 + a*b^3)*cosh(x)^2)*sinh(x)^6 + 8*(7*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)
*cosh(x)^3 - 3*(a^4 + a^3*b - a^2*b^2 - a*b^3)*cosh(x))*sinh(x)^5 + 6*(a^4
+ 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)^4 + 2*(35*(a^4 + 3*a^3*b + 3*a^2*b^2
+ a*b^3)*cosh(x)^4 + 3*a^4 + 9*a^3*b + 9*a^2*b^2 + 3*a*b^3 - 30*(a^4 + a^3
*b - a^2*b^2 - a*b^3)*cosh(x)^2)*sinh(x)^4 + a^4 + 3*a^3*b + 3*a^2*b^2 + a*
b^3 + 8*(7*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)^5 - 10*(a^4 + a^3*b
- a^2*b^2 - a*b^3)*cosh(x)^3 + 3*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)
))*sinh(x)^3 - 4*(a^4 + a^3*b - a^2*b^2 - a*b^3)*cosh(x)^2 + 4*(7*(a^4 + 3*
a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)^6 - 15*(a^4 + a^3*b - a^2*b^2 - a*b^3)*c
osh(x)^4 - a^4 - a^3*b + a^2*b^2 + a*b^3 + 9*(a^4 + 3*a^3*b + 3*a^2*b^2 + a
*b^3)*cosh(x)^2)*sinh(x)^2 + 8*((a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)
^7 - 3*(a^4 + a^3*b - a^2*b^2 - a*b^3)*cosh(x)^5 + 3*(a^4 + 3*a^3*b + 3*a^2
*b^2 + a*b^3)*cosh(x)^3 - (a^4 + a^3*b - a^2*b^2 - a*b^3)*cosh(x))*sinh(x))
]

```

Sympy [F]

$$\int \frac{\coth(x)}{(a + b \coth^4(x))^{3/2}} dx = \int \frac{\coth(x)}{(a + b \coth^4(x))^{\frac{3}{2}}} dx$$

[In] integrate(coth(x)/(a+b*coth(x)**4)**(3/2),x)

[Out] Integral(coth(x)/(a + b*coth(x)**4)**(3/2), x)

Maxima [F]

$$\int \frac{\coth(x)}{(a + b \coth^4(x))^{3/2}} dx = \int \frac{\coth(x)}{(b \coth^4(x) + a)^{3/2}} dx$$

[In] integrate(coth(x)/(a+b*coth(x)^4)^(3/2),x, algorithm="maxima")

[Out] integrate(coth(x)/(b*coth(x)^4 + a)^(3/2), x)

Giac [F]

$$\int \frac{\coth(x)}{(a + b \coth^4(x))^{3/2}} dx = \int \frac{\coth(x)}{(b \coth^4(x) + a)^{3/2}} dx$$

[In] integrate(coth(x)/(a+b*coth(x)^4)^(3/2),x, algorithm="giac")

[Out] integrate(coth(x)/(b*coth(x)^4 + a)^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\coth(x)}{(a + b \coth^4(x))^{3/2}} dx = \int \frac{\coth(x)}{(b \coth^4(x) + a)^{3/2}} dx$$

[In] int(coth(x)/(a + b*coth(x)^4)^(3/2),x)

[Out] int(coth(x)/(a + b*coth(x)^4)^(3/2), x)

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 363

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*      Small rewrite of logic in main function to make it*)
(*      match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
```

```

(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCo
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count is
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"
    ]
  ]
  ,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<>
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

  finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)

```

```

(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```

```

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3, ExpnType[expn[[1]]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
            If[Head[expn]===RootSum,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,

```

```

    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result, optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);

```

```

#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 ("
```

```

                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_c
    end if
  else #result contains complex but optimal is not
    if debug then
      print("result contains complex but optimal is not");
    fi;
    return "C","Result contains complex when optimal does not.";
  fi;
else # result do not contain complex
  # this assumes optimal do not as well. No check is needed here.
  if debug then
    print("result do not contain complex, this assumes optimal do not as well");
  fi;
  if leaf_count_result<=2*leaf_count_optimal then
    if debug then
      print("leaf_count_result<=2*leaf_count_optimal");
    fi;
    return "A"," ";
  else
    if debug then
      print("leaf_count_result>2*leaf_count_optimal");
    fi;
    return "B",cat("Leaf count of result is larger than twice the leaf count of opt
                                convert(leaf_count_result,string)," $ vs. $2(",
                                convert(leaf_count_optimal,string)," )=",convert(2*leaf_coun
    fi;
  fi;
else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
                convert(ExpnType_result,string)," vs. order ",
                convert(ExpnType_optimal,string),".");
fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

```



```

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`) or type(expn,'*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else

```

```

9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u), u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnTy
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+' or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)+str(leaf_count_optimal)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)+str(ExpnType_optimal)

```

```

# print("Before returning. grade=", grade, " grade_annotation=", grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

# Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
# Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
# June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
# July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    # print("Enter tree_size, expr is ", expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: # isinstance(expr, Pow):
        if expr.operands()[1] == 1/2: # expr.args[1] == Rational(1,2):
            if debug: print("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception,AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```



```

    return 1
  else:
    return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
  else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. " + str(leaf_c

else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_result)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```