

Computer Algebra Independent Integration Tests

Summer 2023 edition

7-Inverse-hyperbolic-functions/7.5-Inverse-hyperbolic-secant/201-
7.5.2-Inverse-hyperbolic-secant-functions

Nasser M. Abbasi

September 5, 2023

Compiled on September 5, 2023 at 2:01pm

Contents

1	Introduction	3
2	detailed summary tables of results	21
3	Listing of integrals	51
4	Appendix	609

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	8
1.4	Performance based on number of rules Rubi used	10
1.5	Performance based on number of steps Rubi used	11
1.6	Solved integrals histogram based on leaf size of result	12
1.7	Solved integrals histogram based on CPU time used	13
1.8	Leaf size vs. CPU time used	14
1.9	list of integrals with no known antiderivative	15
1.10	List of integrals solved by CAS but has no known antiderivative	15
1.11	list of integrals solved by CAS but failed verification	15
1.12	Timing	16
1.13	Verification	16
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [100]. This is test number [201].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (100)	0.00 (0)
Mathematica	99.00 (99)	1.00 (1)
Maple	83.00 (83)	17.00 (17)
Fricas	73.00 (73)	27.00 (27)
Mupad	56.00 (56)	44.00 (44)
Maxima	21.00 (21)	79.00 (79)
Giac	4.00 (4)	96.00 (96)
Sympy	3.00 (3)	97.00 (97)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	97.000	2.000	1.000	0.000
Mathematica	65.000	9.000	25.000	1.000
Maple	60.000	2.000	21.000	17.000
Fricas	57.000	16.000	0.000	27.000
Maxima	21.000	0.000	0.000	79.000
Sympy	3.000	0.000	0.000	97.000
Giac	0.000	4.000	0.000	96.000
Mupad	0.000	56.000	0.000	44.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of

error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	1	100.00	0.00	0.00
Maple	17	100.00	0.00	0.00
Fricas	27	77.78	0.00	22.22
Mupad	44	0.00	100.00	0.00
Maxima	79	93.67	0.00	6.33
Giac	96	89.58	0.00	10.42
Sympy	97	98.97	1.03	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Rubi	0.16
Maxima	0.22
Fricas	0.25
Maple	0.47
Giac	0.66
Mathematica	1.11
Sympy	3.21
Mupad	13.15

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Maxima	49.76	0.58	46.00	0.67
Sympy	78.33	1.23	82.00	0.95
Rubi	135.47	1.06	107.00	1.00
Fricas	136.37	1.55	87.00	1.05
Maple	177.34	1.83	104.00	1.15
Giac	194.75	2.59	197.50	2.62
Mathematica	222.89	1.49	101.00	1.09
Mupad	329.71	2.76	88.50	1.74

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {47, 53}

Mathematica {18, 57, 59, 60}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
```

```
x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2023
Design-vide

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	22
2.2	Detailed conclusion table per each integral for all CAS systems	25
2.3	Detailed conclusion table specific for Rubi results	46

2.1 List of integrals sorted by grade for each CAS

Rubi	22
Mma	22
Maple	23
Fricas	23
Maxima	23
Giac	24
Mupad	24
Sympy	24

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 98, 99, 100 }

B grade { 38, 97 }

C grade { 39 }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Mma

A grade { 8, 9, 10, 11, 12, 15, 16, 17, 20, 21, 22, 24, 25, 26, 27, 28, 30, 32, 34, 39, 40, 41, 42, 43, 44, 47, 51, 53, 55, 56, 57, 58, 59, 60, 61, 62, 64, 66, 68, 69, 70, 71, 72, 73, 74, 75, 76, 78, 80, 82, 83, 84, 85, 86, 87, 88, 89, 91, 93, 94, 95, 96, 98, 99, 100 }

B grade { 4, 6, 7, 23, 29, 31, 36, 38, 97 }

C grade { 1, 2, 3, 5, 13, 14, 18, 33, 35, 37, 45, 46, 48, 49, 50, 52, 54, 63, 65, 67, 77, 79, 81, 90, 92 }

F normal fail { 19 }

F(-1) timeout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 9, 10, 11, 13, 14, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 31, 32, 34, 36, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 52, 53, 54, 66, 68, 70, 71, 72, 73, 74, 75, 82, 83, 84, 85, 86, 87, 89, 91, 93, 95, 97, 98, 99 }

B grade { 6, 80 }

C grade { 5, 7, 8, 33, 35, 37, 51, 55, 63, 65, 67, 69, 76, 77, 78, 79, 81, 90, 92, 94, 96 }

F normal fail { 12, 15, 16, 17, 18, 19, 30, 56, 57, 58, 59, 60, 61, 62, 64, 88, 100 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 20, 21, 22, 23, 25, 26, 27, 28, 32, 33, 34, 35, 37, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 54, 55, 63, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 89, 91, 94, 96, 97 }

B grade { 2, 3, 4, 6, 7, 8, 36, 38, 51, 53, 90, 92, 93, 95, 99, 100 }

C grade { }

F normal fail { 5, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 24, 30, 52, 56, 57, 58, 59, 88, 98 }

F(-1) timedout fail { }

F(-2) exception fail { 29, 31, 60, 61, 62, 64 }

Maxima

A grade { 4, 20, 21, 22, 23, 25, 26, 27, 28, 32, 34, 39, 41, 43, 47, 66, 71, 73, 75, 99, 100 }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 24, 29, 30, 31, 33, 35, 36, 37, 38, 40, 42, 44, 45, 46, 48, 49, 50, 51, 52, 53, 54, 55, 58, 59, 63, 64, 65, 67, 68, 69, 70, 72, 74, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98 }

F(-1) timedout fail { }

F(-2) exception fail { 56, 57, 60, 61, 62 }

Giac**A grade** { }**B grade** { 45, 47, 49, 53 }**C grade** { }**F normal fail** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 52, 54, 56, 58, 59, 60, 61, 62, 63, 64, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 }**F(-1) timedout fail** { }**F(-2) exception fail** { 32, 33, 34, 46, 48, 50, 51, 55, 57, 65 }**Mupad****A grade** { }**B grade** { 4, 25, 28, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 47, 49, 51, 53, 55, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 89, 90, 91, 92, 93, 94, 95, 96, 97, 99, 100 }**C grade** { }**F normal fail** { }**F(-1) timedout fail** { 1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 26, 27, 29, 30, 31, 46, 48, 50, 52, 54, 56, 57, 58, 59, 60, 61, 62, 63, 64, 88, 98 }**F(-2) exception fail** { }**Sympy****A grade** { 38, 55, 70 }**B grade** { }**C grade** { }**F normal fail** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 }**F(-1) timedout fail** { 100 }**F(-2) exception fail** { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	203	203	225	250	0	345	0	0	0
N.S.	1	1.00	1.11	1.23	0.00	1.70	0.00	0.00	0.00
time (sec)	N/A	0.119	0.552	0.842	0.000	0.312	0.000	0.000	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	153	153	200	189	0	327	0	0	0
N.S.	1	1.00	1.31	1.24	0.00	2.14	0.00	0.00	0.00
time (sec)	N/A	0.075	0.249	0.734	0.000	0.295	0.000	0.000	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	107	176	111	0	308	0	0	0
N.S.	1	1.00	1.64	1.04	0.00	2.88	0.00	0.00	0.00
time (sec)	N/A	0.047	0.184	0.723	0.000	0.282	0.000	0.000	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	44	97	44	31	253	0	0	43
N.S.	1	1.00	2.20	1.00	0.70	5.75	0.00	0.00	0.98
time (sec)	N/A	0.042	0.276	0.413	0.203	0.268	0.000	0.000	4.983

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	170	170	332	882	0	0	0	0	0
N.S.	1	1.00	1.95	5.19	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.219	0.314	0.878	0.000	0.000	0.000	0.000	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	244	171	0	651	0	0	0
N.S.	1	1.00	3.49	2.44	0.00	9.30	0.00	0.00	0.00
time (sec)	N/A	0.083	0.269	0.947	0.000	0.301	0.000	0.000	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	C	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	133	133	315	370	0	865	0	0	0
N.S.	1	1.00	2.37	2.78	0.00	6.50	0.00	0.00	0.00
time (sec)	N/A	0.181	0.897	0.961	0.000	0.320	0.000	0.000	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	197	197	368	608	0	987	0	0	0
N.S.	1	1.00	1.87	3.09	0.00	5.01	0.00	0.00	0.00
time (sec)	N/A	0.244	0.360	0.974	0.000	0.331	0.000	0.000	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	279	279	305	599	0	0	0	0	0
N.S.	1	1.00	1.09	2.15	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.193	2.278	0.870	0.000	0.000	0.000	0.000	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	149	149	172	331	0	0	0	0	0
N.S.	1	1.00	1.15	2.22	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.118	0.523	0.776	0.000	0.000	0.000	0.000	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	80	105	192	0	0	0	0	0
N.S.	1	1.00	1.31	2.40	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.057	0.240	0.545	0.000	0.000	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	274	274	280	0	0	0	0	0	0
N.S.	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.363	0.336	0.000	0.000	0.000	0.000	0.000	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	224	224	678	362	0	0	0	0	0
N.S.	1	1.00	3.03	1.62	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.307	2.832	0.922	0.000	0.000	0.000	0.000	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	965	965	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.049	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	164	164	84	54	58	57	0	0	0
N.S.	1	1.00	0.51	0.33	0.35	0.35	0.00	0.00	0.00
time (sec)	N/A	0.022	0.050	0.252	0.210	0.255	0.000	0.000	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	126	126	72	49	46	52	0	0	0
N.S.	1	1.00	0.57	0.39	0.37	0.41	0.00	0.00	0.00
time (sec)	N/A	0.019	0.037	0.221	0.212	0.250	0.000	0.000	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	56	42	34	45	0	0	0
N.S.	1	1.00	0.64	0.48	0.39	0.51	0.00	0.00	0.00
time (sec)	N/A	0.013	0.030	0.231	0.215	0.260	0.000	0.000	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	118	36	19	39	0	0	0
N.S.	1	1.00	2.74	0.84	0.44	0.91	0.00	0.00	0.00
time (sec)	N/A	0.005	0.127	0.230	0.204	0.254	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	45	65	0	0	0	0	0
N.S.	1	1.00	0.98	1.41	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.071	0.058	0.468	0.000	0.000	0.000	0.000	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	111	64	65	45	0	0	40
N.S.	1	1.00	1.13	0.65	0.66	0.46	0.00	0.00	0.41
time (sec)	N/A	0.016	0.083	0.253	0.219	0.262	0.000	0.000	4.901

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	136	136	125	79	92	54	0	0	0
N.S.	1	1.00	0.92	0.58	0.68	0.40	0.00	0.00	0.00
time (sec)	N/A	0.020	0.123	0.230	0.210	0.259	0.000	0.000	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	172	172	140	91	116	60	0	0	0
N.S.	1	1.00	0.81	0.53	0.67	0.35	0.00	0.00	0.00
time (sec)	N/A	0.025	0.137	0.232	0.204	0.262	0.000	0.000	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	25	20	16	22	0	0	17
N.S.	1	1.00	1.19	0.95	0.76	1.05	0.00	0.00	0.81
time (sec)	N/A	0.005	0.022	0.230	0.224	0.254	0.000	0.000	0.070

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	61	219	111	0	0	0	0	0
N.S.	1	1.00	3.59	1.82	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.078	1.086	1.066	0.000	0.000	0.000	0.000	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	54	54	49	0	0	0	0	0	0
N.S.	1	1.00	0.91	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.073	0.059	0.000	0.000	0.000	0.000	0.000	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	77	77	249	135	0	0	0	0	0
N.S.	1	1.00	3.23	1.75	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.066	1.859	1.099	0.000	0.000	0.000	0.000	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	64	83	65	64	47	65	0	0	75
N.S.	1	1.30	1.02	1.00	0.73	1.02	0.00	0.00	1.17
time (sec)	N/A	0.025	0.099	0.047	0.240	0.245	0.000	0.000	4.127

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	A	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	84	84	97	118	0	95	0	0	521
N.S.	1	1.00	1.15	1.40	0.00	1.13	0.00	0.00	6.20
time (sec)	N/A	0.025	0.143	0.045	0.000	0.250	0.000	0.000	16.438

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	52	48	54	38	54	0	0	55
N.S.	1	1.37	1.26	1.42	1.00	1.42	0.00	0.00	1.45
time (sec)	N/A	0.019	0.068	0.049	0.222	0.254	0.000	0.000	5.112

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	53	75	92	0	79	0	0	303
N.S.	1	1.00	1.42	1.74	0.00	1.49	0.00	0.00	5.72
time (sec)	N/A	0.014	0.067	0.048	0.000	0.253	0.000	0.000	10.972

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	39	79	80	0	115	0	0	182
N.S.	1	1.62	3.29	3.33	0.00	4.79	0.00	0.00	7.58
time (sec)	N/A	0.082	0.053	0.103	0.000	0.259	0.000	0.000	6.311

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	48	64	75	92	0	77	0	0	184
N.S.	1	1.33	1.56	1.92	0.00	1.60	0.00	0.00	3.83
time (sec)	N/A	0.021	0.056	0.048	0.000	0.270	0.000	0.000	6.371

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	B	A	F	B	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	99	93	91	0	128	71	0	71
N.S.	1	2.83	2.66	2.60	0.00	3.66	2.03	0.00	2.03
time (sec)	N/A	0.030	0.066	0.046	0.000	0.254	2.665	0.000	4.878

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	55	84	43	53	43	52	0	0	58
N.S.	1	1.53	0.78	0.96	0.78	0.95	0.00	0.00	1.05
time (sec)	N/A	0.027	0.054	0.043	0.224	0.262	0.000	0.000	4.587

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	132	132	110	110	0	138	0	0	602
N.S.	1	1.00	0.83	0.83	0.00	1.05	0.00	0.00	4.56
time (sec)	N/A	0.042	0.087	0.046	0.000	0.246	0.000	0.000	19.502

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	115	115	60	63	51	60	0	0	76
N.S.	1	1.00	0.52	0.55	0.44	0.52	0.00	0.00	0.66
time (sec)	N/A	0.036	0.073	0.048	0.218	0.250	0.000	0.000	4.644

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	163	163	129	132	0	148	0	0	878
N.S.	1	1.00	0.79	0.81	0.00	0.91	0.00	0.00	5.39
time (sec)	N/A	0.054	0.133	0.053	0.000	0.260	0.000	0.000	41.119

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	146	146	76	71	60	69	0	0	95
N.S.	1	1.00	0.52	0.49	0.41	0.47	0.00	0.00	0.65
time (sec)	N/A	0.051	0.098	0.056	0.231	0.248	0.000	0.000	4.862

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	194	194	145	152	0	156	0	0	1155
N.S.	1	1.00	0.75	0.78	0.00	0.80	0.00	0.00	5.95
time (sec)	N/A	0.074	0.137	0.060	0.000	0.258	0.000	0.000	43.823

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	111	111	111	137	0	116	0	205	521
N.S.	1	1.00	1.00	1.23	0.00	1.05	0.00	1.85	4.69
time (sec)	N/A	0.044	0.183	0.201	0.000	0.252	0.000	0.320	18.815

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	115	115	143	114	0	79	0	0	0
N.S.	1	1.00	1.24	0.99	0.00	0.69	0.00	0.00	0.00
time (sec)	N/A	0.037	0.350	1.070	0.000	0.079	0.000	0.000	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	58	71	56	60	42	60	0	190	57
N.S.	1	1.22	0.97	1.03	0.72	1.03	0.00	3.28	0.98
time (sec)	N/A	0.028	0.110	0.052	0.226	0.257	0.000	0.300	5.472

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	112	112	140	136	0	91	0	0	0
N.S.	1	1.00	1.25	1.21	0.00	0.81	0.00	0.00	0.00
time (sec)	N/A	0.049	0.460	0.659	0.000	0.097	0.000	0.000	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	63	92	112	0	102	0	132	306
N.S.	1	1.00	1.46	1.78	0.00	1.62	0.00	2.10	4.86
time (sec)	N/A	0.027	0.103	0.098	0.000	0.262	0.000	0.297	10.881

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	67	112	102	0	62	0	0	0
N.S.	1	1.00	1.67	1.52	0.00	0.93	0.00	0.00	0.00
time (sec)	N/A	0.021	0.273	0.576	0.000	0.078	0.000	0.000	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	68	68	100	127	0	133	0	0	182
N.S.	1	1.00	1.47	1.87	0.00	1.96	0.00	0.00	2.68
time (sec)	N/A	0.030	0.069	0.110	0.000	0.247	0.000	0.000	7.093

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	147	147	135	132	0	0	0	0	0
N.S.	1	1.00	0.92	0.90	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.051	0.328	0.770	0.000	0.000	0.000	0.000	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	93	22	103	0	102	0	252	185
N.S.	1	1.16	0.28	1.29	0.00	1.28	0.00	3.15	2.31
time (sec)	N/A	0.040	0.086	0.106	0.000	0.254	0.000	1.726	7.400

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	109	109	139	0	0	0	0	0	0
N.S.	1	1.00	1.28	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.060	0.713	0.000	0.000	0.000	0.000	0.000	0.000

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F(-2)	F(-2)	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	133	133	186	0	0	0	0	0	0
N.S.	1	1.00	1.40	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.065	4.331	0.000	0.000	0.000	0.000	0.000	0.000

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F(-2)	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	119	119	159	0	0	0	0	0	0
N.S.	1	1.00	1.34	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.042	0.327	0.000	0.000	0.000	0.000	0.000	0.000

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F(-2)	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	105	105	164	0	0	0	0	0	0
N.S.	1	1.00	1.56	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.043	0.251	0.000	0.000	0.000	0.000	0.000	0.000

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	87	106	96	116	0	102	0	0	0
N.S.	1	1.22	1.10	1.33	0.00	1.17	0.00	0.00	0.00
time (sec)	N/A	0.047	0.152	1.171	0.000	0.285	0.000	0.000	0.000

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	107	107	166	0	0	0	0	0	0
N.S.	1	1.00	1.55	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.047	0.243	0.000	0.000	0.000	0.000	0.000	0.000

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	A	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	203	203	105	136	0	103	0	0	808
N.S.	1	1.00	0.52	0.67	0.00	0.51	0.00	0.00	3.98
time (sec)	N/A	0.530	0.159	0.060	0.000	0.262	0.000	0.000	25.515

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	117	52	72	42	62	0	0	63
N.S.	1	1.00	0.44	0.62	0.36	0.53	0.00	0.00	0.54
time (sec)	N/A	0.400	0.075	0.051	0.224	0.244	0.000	0.000	4.678

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	169	169	86	105	0	87	0	0	420
N.S.	1	1.00	0.51	0.62	0.00	0.51	0.00	0.00	2.49
time (sec)	N/A	0.338	0.078	0.054	0.000	0.249	0.000	0.000	12.778

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	85	89	98	0	124	0	0	56
N.S.	1	1.00	1.05	1.15	0.00	1.46	0.00	0.00	0.66
time (sec)	N/A	0.314	0.069	0.045	0.000	0.251	0.000	0.000	7.010

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	57	57	75	111	0	85	0	0	162
N.S.	1	1.00	1.32	1.95	0.00	1.49	0.00	0.00	2.84
time (sec)	N/A	0.131	0.107	0.055	0.000	0.272	0.000	0.000	7.778

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	86	86	86	110	0	138	82	0	323
N.S.	1	1.00	1.00	1.28	0.00	1.60	0.95	0.00	3.76
time (sec)	N/A	0.317	0.067	0.047	0.000	0.248	2.889	0.000	13.879

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	57	57	52	73	46	61	0	0	67
N.S.	1	1.00	0.91	1.28	0.81	1.07	0.00	0.00	1.18
time (sec)	N/A	0.275	0.073	0.046	0.220	0.253	0.000	0.000	4.762

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	147	147	121	131	0	146	0	0	885
N.S.	1	1.00	0.82	0.89	0.00	0.99	0.00	0.00	6.02
time (sec)	N/A	0.333	0.129	0.056	0.000	0.257	0.000	0.000	46.860

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	183	183	69	84	56	69	0	0	86
N.S.	1	1.00	0.38	0.46	0.31	0.38	0.00	0.00	0.47
time (sec)	N/A	0.378	0.092	0.050	0.212	0.269	0.000	0.000	5.425

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	267	267	137	153	0	156	0	0	2480
N.S.	1	1.00	0.51	0.57	0.00	0.58	0.00	0.00	9.29
time (sec)	N/A	0.414	0.157	0.062	0.000	0.255	0.000	0.000	68.991

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	301	301	85	92	65	78	0	0	105
N.S.	1	1.00	0.28	0.31	0.22	0.26	0.00	0.00	0.35
time (sec)	N/A	0.412	0.108	0.056	0.213	0.248	0.000	0.000	5.340

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	147	147	65	531	0	65	0	0	73
N.S.	1	1.00	0.44	3.61	0.00	0.44	0.00	0.00	0.50
time (sec)	N/A	0.438	0.096	0.351	0.000	0.250	0.000	0.000	5.348

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	163	163	97	120	0	95	0	0	795
N.S.	1	1.00	0.60	0.74	0.00	0.58	0.00	0.00	4.88
time (sec)	N/A	0.382	0.128	0.918	0.000	0.251	0.000	0.000	25.409

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	75	75	48	269	0	54	0	0	57
N.S.	1	1.00	0.64	3.59	0.00	0.72	0.00	0.00	0.76
time (sec)	N/A	0.348	0.069	0.286	0.000	0.248	0.000	0.000	4.887

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	94	94	75	94	0	79	0	0	407
N.S.	1	1.00	0.80	1.00	0.00	0.84	0.00	0.00	4.33
time (sec)	N/A	0.202	0.079	0.876	0.000	0.263	0.000	0.000	13.356

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	65	72	2612	0	115	0	0	47
N.S.	1	1.00	1.11	40.18	0.00	1.77	0.00	0.00	0.72
time (sec)	N/A	0.108	0.045	0.282	0.000	0.256	0.000	0.000	7.361

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	74	96	0	76	0	0	184
N.S.	1	1.00	1.61	2.09	0.00	1.65	0.00	0.00	4.00
time (sec)	N/A	0.282	0.044	0.894	0.000	0.278	0.000	0.000	7.480

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	92	96	0	128	0	0	323
N.S.	1	1.00	1.28	1.33	0.00	1.78	0.00	0.00	4.49
time (sec)	N/A	0.274	0.063	0.886	0.000	0.257	0.000	0.000	16.763

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	116	43	58	0	52	0	0	58
N.S.	1	1.00	0.37	0.50	0.00	0.45	0.00	0.00	0.50
time (sec)	N/A	0.313	0.059	0.914	0.000	0.256	0.000	0.000	5.728

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	69	74	0	69	0	0	76
N.S.	1	1.00	0.78	0.84	0.00	0.78	0.00	0.00	0.86
time (sec)	N/A	0.131	0.220	0.691	0.000	0.244	0.000	0.000	5.423

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	75	75	110	122	0	91	0	0	340
N.S.	1	1.00	1.47	1.63	0.00	1.21	0.00	0.00	4.53
time (sec)	N/A	0.117	0.199	0.682	0.000	0.255	0.000	0.000	11.191

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	45	45	44	52	0	49	0	0	44
N.S.	1	1.00	0.98	1.16	0.00	1.09	0.00	0.00	0.98
time (sec)	N/A	0.097	0.135	0.697	0.000	0.264	0.000	0.000	5.245

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	37	68	97	0	53	0	0	84
N.S.	1	1.00	1.84	2.62	0.00	1.43	0.00	0.00	2.27
time (sec)	N/A	0.061	0.200	0.694	0.000	0.252	0.000	0.000	6.105

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	71	71	73	82	0	92	0	0	59
N.S.	1	1.00	1.03	1.15	0.00	1.30	0.00	0.00	0.83
time (sec)	N/A	0.095	0.165	0.704	0.000	0.242	0.000	0.000	5.954

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	59	65	0	62	0	0	37
N.S.	1	1.00	1.40	1.55	0.00	1.48	0.00	0.00	0.88
time (sec)	N/A	0.110	0.213	0.722	0.000	0.264	0.000	0.000	5.083

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	108	108	108	119	0	156	0	0	331
N.S.	1	1.00	1.00	1.10	0.00	1.44	0.00	0.00	3.06
time (sec)	N/A	0.128	0.216	0.729	0.000	0.249	0.000	0.000	17.892

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	85	90	90	0	89	0	0	75
N.S.	1	1.00	1.06	1.06	0.00	1.05	0.00	0.00	0.88
time (sec)	N/A	0.124	0.323	0.731	0.000	0.251	0.000	0.000	5.484

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	B	A	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	26	28	36	0	35	0	0	76
N.S.	1	2.17	2.33	3.00	0.00	2.92	0.00	0.00	6.33
time (sec)	N/A	0.821	0.441	0.734	0.000	0.247	0.000	0.000	5.778

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	61	52	111	0	0	0	0	0
N.S.	1	1.00	0.85	1.82	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.087	0.101	0.783	0.000	0.000	0.000	0.000	0.000

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	57	57	106	53	38	283	0	0	56
N.S.	1	1.00	1.86	0.93	0.67	4.96	0.00	0.00	0.98
time (sec)	N/A	0.081	0.174	0.185	0.199	0.271	0.000	0.000	5.585

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	A	B	F(-1)	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	58	58	106	0	40	385	0	0	54
N.S.	1	1.00	1.83	0.00	0.69	6.64	0.00	0.00	0.93
time (sec)	N/A	0.093	0.218	0.000	0.196	0.304	0.000	0.000	5.009

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. The column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [15] had the largest ratio of [1.1999999999999996]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	8	7	1.00	10	0.700
2	A	7	6	1.00	10	0.600
3	A	6	6	1.00	8	0.750
4	A	4	4	1.00	6	0.667
5	A	14	8	1.00	10	0.800
6	A	5	5	1.00	10	0.500
7	A	7	7	1.00	10	0.700
8	A	8	8	1.00	10	0.800
9	A	17	9	1.00	12	0.750
10	A	11	8	1.00	10	0.800
11	A	8	6	1.00	8	0.750
12	A	17	9	1.00	12	0.750
13	A	12	8	1.00	12	0.667
14	A	23	11	1.00	12	0.917
15	A	16	12	1.00	10	1.200
16	A	10	7	1.00	8	0.875
17	A	20	10	1.00	12	0.833
18	A	14	9	1.00	12	0.750
19	A	32	13	1.00	12	1.083
20	A	4	3	1.00	10	0.300
21	A	4	3	1.00	10	0.300
22	A	4	3	1.00	8	0.375
23	A	3	3	1.00	6	0.500
24	A	7	6	1.00	10	0.600

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
25	A	5	5	1.00	10	0.500
26	A	6	5	1.00	10	0.500
27	A	7	5	1.00	10	0.500
28	A	3	3	1.00	4	0.750
29	A	7	6	1.00	10	0.600
30	A	7	6	1.00	10	0.600
31	A	7	7	1.00	10	0.700
32	A	5	5	1.30	10	0.500
33	A	5	5	1.00	10	0.500
34	A	3	3	1.37	10	0.300
35	A	4	4	1.00	8	0.500
36	A	3	3	1.62	6	0.500
37	A	5	5	1.33	10	0.500
38	B	6	6	2.83	10	0.600
39	C	5	5	1.53	10	0.500
40	A	8	6	1.00	10	0.600
41	A	7	5	1.00	10	0.500
42	A	10	6	1.00	10	0.600
43	A	9	5	1.00	10	0.500
44	A	12	6	1.00	10	0.600
45	A	6	6	1.00	12	0.500
46	A	5	5	1.00	12	0.417
47	A	4	4	1.22	12	0.333
48	A	7	7	1.00	12	0.583
49	A	5	5	1.00	12	0.417
50	A	4	4	1.00	12	0.333
51	A	6	6	1.00	10	0.600
52	A	8	8	1.00	8	1.000
53	A	5	5	1.16	12	0.417
54	A	5	5	1.00	12	0.417
55	A	7	7	1.00	12	0.583
56	A	4	4	1.00	12	0.333
57	A	4	4	1.00	12	0.333
58	A	4	4	1.00	10	0.400
59	A	5	5	1.00	12	0.417

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
60	A	4	4	1.00	12	0.333
61	A	4	4	1.00	10	0.400
62	A	4	4	1.00	8	0.500
63	A	6	6	1.22	12	0.500
64	A	4	4	1.00	12	0.333
65	A	9	6	1.00	12	0.500
66	A	8	6	1.00	12	0.500
67	A	7	6	1.00	12	0.500
68	A	8	5	1.00	10	0.500
69	A	7	5	1.00	8	0.625
70	A	5	3	1.00	12	0.250
71	A	4	2	1.00	12	0.167
72	A	5	3	1.00	12	0.250
73	A	4	2	1.00	12	0.167
74	A	5	3	1.00	12	0.250
75	A	4	2	1.00	12	0.167
76	A	8	5	1.00	12	0.417
77	A	7	5	1.00	12	0.417
78	A	6	5	1.00	12	0.417
79	A	5	4	1.00	10	0.400
80	A	6	4	1.00	8	0.500
81	A	5	3	1.00	12	0.250
82	A	5	3	1.00	12	0.250
83	A	4	2	1.00	12	0.167
84	A	5	3	1.00	12	0.250
85	A	4	2	1.00	12	0.167
86	A	5	3	1.00	12	0.250
87	A	4	2	1.00	12	0.167
88	A	5	4	1.00	24	0.167
89	A	8	7	1.00	22	0.318
90	A	7	7	1.00	22	0.318
91	A	4	4	1.00	22	0.182
92	A	5	5	1.00	20	0.250
93	A	8	8	1.00	19	0.421
94	A	5	5	1.00	22	0.227

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
95	A	9	8	1.00	22	0.364
96	A	8	7	1.00	22	0.318
97	B	7	5	2.17	25	0.200
98	A	8	8	1.00	19	0.421
99	A	5	5	1.00	12	0.417
100	A	5	5	1.00	14	0.357

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^3 \operatorname{sech}^{-1}(a + bx) dx$	55
3.2	$\int x^2 \operatorname{sech}^{-1}(a + bx) dx$	61
3.3	$\int x \operatorname{sech}^{-1}(a + bx) dx$	67
3.4	$\int \operatorname{sech}^{-1}(a + bx) dx$	72
3.5	$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x} dx$	76
3.6	$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^2} dx$	84
3.7	$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^3} dx$	89
3.8	$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^4} dx$	96
3.9	$\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx$	104
3.10	$\int x \operatorname{sech}^{-1}(a + bx)^2 dx$	113
3.11	$\int \operatorname{sech}^{-1}(a + bx)^2 dx$	120
3.12	$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} dx$	125
3.13	$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx$	134
3.14	$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^3} dx$	141
3.15	$\int x \operatorname{sech}^{-1}(a + bx)^3 dx$	154
3.16	$\int \operatorname{sech}^{-1}(a + bx)^3 dx$	163
3.17	$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx$	169
3.18	$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^2} dx$	181
3.19	$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^3} dx$	189
3.20	$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx$	204
3.21	$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx$	209
3.22	$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx$	213
3.23	$\int \operatorname{sech}^{-1}(\sqrt{x}) dx$	217

3.24	$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx$	221
3.25	$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx$	226
3.26	$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx$	231
3.27	$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx$	236
3.28	$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx$	242
3.29	$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx$	246
3.30	$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx$	251
3.31	$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx$	255
3.32	$\int e^{\operatorname{sech}^{-1}(ax)} x^4 dx$	260
3.33	$\int e^{\operatorname{sech}^{-1}(ax)} x^3 dx$	265
3.34	$\int e^{\operatorname{sech}^{-1}(ax)} x^2 dx$	270
3.35	$\int e^{\operatorname{sech}^{-1}(ax)} x dx$	274
3.36	$\int e^{\operatorname{sech}^{-1}(ax)} dx$	278
3.37	$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x} dx$	282
3.38	$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^2} dx$	287
3.39	$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^3} dx$	292
3.40	$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^4} dx$	296
3.41	$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^5} dx$	302
3.42	$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^6} dx$	307
3.43	$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^7} dx$	313
3.44	$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^8} dx$	319
3.45	$\int e^{\operatorname{sech}^{-1}(ax^2)} x^7 dx$	326
3.46	$\int e^{\operatorname{sech}^{-1}(ax^2)} x^6 dx$	332
3.47	$\int e^{\operatorname{sech}^{-1}(ax^2)} x^5 dx$	337
3.48	$\int e^{\operatorname{sech}^{-1}(ax^2)} x^4 dx$	341
3.49	$\int e^{\operatorname{sech}^{-1}(ax^2)} x^3 dx$	346
3.50	$\int e^{\operatorname{sech}^{-1}(ax^2)} x^2 dx$	351
3.51	$\int e^{\operatorname{sech}^{-1}(ax^2)} x dx$	355
3.52	$\int e^{\operatorname{sech}^{-1}(ax^2)} dx$	360
3.53	$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x} dx$	366
3.54	$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^2} dx$	371
3.55	$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^3} dx$	376
3.56	$\int e^{\operatorname{sech}^{-1}(ax^3)} x^m dx$	382
3.57	$\int e^{\operatorname{sech}^{-1}(ax^2)} x^m dx$	387
3.58	$\int e^{\operatorname{sech}^{-1}(ax)} x^m dx$	392
3.59	$\int e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} x^m dx$	396

3.60	$\int e^{\operatorname{sech}^{-1}(ax^p)} x^m dx$	400
3.61	$\int e^{\operatorname{sech}^{-1}(ax^p)} x dx$	405
3.62	$\int e^{\operatorname{sech}^{-1}(ax^p)} dx$	410
3.63	$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x} dx$	415
3.64	$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x^2} dx$	420
3.65	$\int e^{2\operatorname{sech}^{-1}(ax)} x^4 dx$	424
3.66	$\int e^{2\operatorname{sech}^{-1}(ax)} x^3 dx$	431
3.67	$\int e^{2\operatorname{sech}^{-1}(ax)} x^2 dx$	436
3.68	$\int e^{2\operatorname{sech}^{-1}(ax)} x dx$	442
3.69	$\int e^{2\operatorname{sech}^{-1}(ax)} dx$	447
3.70	$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x} dx$	452
3.71	$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^2} dx$	456
3.72	$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^3} dx$	460
3.73	$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^4} dx$	466
3.74	$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^5} dx$	471
3.75	$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^6} dx$	478
3.76	$\int e^{-\operatorname{sech}^{-1}(ax)} x^4 dx$	483
3.77	$\int e^{-\operatorname{sech}^{-1}(ax)} x^3 dx$	488
3.78	$\int e^{-\operatorname{sech}^{-1}(ax)} x^2 dx$	494
3.79	$\int e^{-\operatorname{sech}^{-1}(ax)} x dx$	499
3.80	$\int e^{-\operatorname{sech}^{-1}(ax)} dx$	504
3.81	$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x} dx$	510
3.82	$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^2} dx$	514
3.83	$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^3} dx$	519
3.84	$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^4} dx$	523
3.85	$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^5} dx$	529
3.86	$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^6} dx$	533
3.87	$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^7} dx$	540
3.88	$\int \frac{e^{\operatorname{sech}^{-1}(cx)} (dx)^m}{1-c^2x^2} dx$	545
3.89	$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^4}{1-c^2x^2} dx$	550
3.90	$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^3}{1-c^2x^2} dx$	555
3.91	$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^2}{1-c^2x^2} dx$	560
3.92	$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x}{1-c^2x^2} dx$	564
3.93	$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{1-c^2x^2} dx$	568
3.94	$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x(1-c^2x^2)} dx$	573

3.95	$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^2(1-c^2x^2)} dx \dots\dots\dots$	578
3.96	$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^3(1-c^2x^2)} dx \dots\dots\dots$	584
3.97	$\int \frac{x(-1+ae^{\operatorname{sech}^{-1}(ax)})}{1-a^2x^2} dx \dots\dots\dots$	589
3.98	$\int \frac{\operatorname{sech}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx \dots\dots\dots$	594
3.99	$\int x^3 \operatorname{sech}^{-1}(a+bx^4) dx \dots\dots\dots$	599
3.100	$\int x^{-1+n} \operatorname{sech}^{-1}(a+bx^n) dx \dots\dots\dots$	604

3.1 $\int x^3 \operatorname{sech}^{-1}(a + bx) dx$

Optimal result	55
Rubi [A] (verified)	56
Mathematica [C] (verified)	58
Maple [A] (verified)	59
Fricas [A] (verification not implemented)	59
Sympy [F]	60
Maxima [F]	60
Giac [F]	60
Mupad [F(-1)]	60

Optimal result

Integrand size = 10, antiderivative size = 203

$$\int x^3 \operatorname{sech}^{-1}(a + bx) dx = -\frac{(2 + 17a^2) \sqrt{\frac{1-a-bx}{1+a+bx}}(1 + a + bx)}{12b^4} - \frac{x^2 \sqrt{\frac{1-a-bx}{1+a+bx}}(1 + a + bx)}{12b^2}$$

$$+ \frac{a(a + bx) \sqrt{\frac{1-a-bx}{1+a+bx}}(1 + a + bx)}{3b^4} - \frac{a^4 \operatorname{sech}^{-1}(a + bx)}{4b^4}$$

$$+ \frac{1}{4} x^4 \operatorname{sech}^{-1}(a + bx) + \frac{a(1 + 2a^2) \arctan\left(\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{a+bx}\right)}{2b^4}$$

```
[Out] -1/4*a^4*arcsech(b*x+a)/b^4+1/4*x^4*arcsech(b*x+a)+1/2*a*(2*a^2+1)*arctan((
b*x+a+1)*((-b*x-a+1)/(b*x+a+1))^(1/2)/(b*x+a))/b^4-1/12*(17*a^2+2)*(b*x+a+1
)*((-b*x-a+1)/(b*x+a+1))^(1/2)/b^4-1/12*x^2*(b*x+a+1)*((-b*x-a+1)/(b*x+a+1)
)^(1/2)/b^2+1/3*a*(b*x+a)*(b*x+a+1)*((-b*x-a+1)/(b*x+a+1))^(1/2)/b^4
```

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {6456, 5576, 3867, 4133, 3855, 3852, 8}

$$\int x^3 \operatorname{sech}^{-1}(a + bx) dx = -\frac{a^4 \operatorname{sech}^{-1}(a + bx)}{4b^4} + \frac{(2a^2 + 1) a \arctan\left(\frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a+bx}\right)}{2b^4} - \frac{(17a^2 + 2) \sqrt{\frac{-a-bx+1}{a+bx+1}}(a + bx + 1)}{12b^4} + \frac{a(a + bx) \sqrt{\frac{-a-bx+1}{a+bx+1}}(a + bx + 1)}{3b^4} - \frac{x^2 \sqrt{\frac{-a-bx+1}{a+bx+1}}(a + bx + 1)}{12b^2} + \frac{1}{4} x^4 \operatorname{sech}^{-1}(a + bx)$$

[In] Int[x^3*ArcSech[a + b*x],x]

[Out] -1/12*((2 + 17*a^2)*Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x))/b^4 - (x^2*Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x))/(12*b^2) + (a*(a + b*x)*Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x))/(3*b^4) - (a^4*ArcSech[a + b*x])/(4*b^4) + (x^4*ArcSech[a + b*x])/4 + (a*(1 + 2*a^2)*ArcTan[(Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x))/(a + b*x)])/(2*b^4)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3852

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3867

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_), x_Symbol] := Simp[(-b^2)*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n - 2)/(d*(n - 1))), x] + Dist[1/(n - 1), Int[(a + b*Csc[c + d*x])^(n - 3)*Simp[a^3*(n - 1) + (b*(b^2*(n - 2) + 3*a^2*(n - 1)))*Csc[c + d*x] + (a*b^2*(3*n - 4))*Csc[c + d*x]^2, x], x], x] /;

FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 2] && IntegerQ[2*n]

Rule 4133

Int[((A_.) + csc[(e_.) + (f_.)*(x_)])*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.) * (csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :> Simp[(-b)*C*Csc[e + f*x]*(Cot[e + f*x]/(2*f)), x] + Dist[1/2, Int[Simp[2*A*a + (2*B*a + b*(2*A + C))*Csc[e + f*x] + 2*(a*C + B*b)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x]

Rule 5576

Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sech[(c_.) + (d_.)*(x_)]^(n_.)*Tanh[(c_.) + (d_.)*(x_)]), x_Symbol] :> Simp[(-(e + f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rule 6456

Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int x \text{sech}(x)(-a + \text{sech}(x))^3 \tanh(x) dx, x, \text{sech}^{-1}(a + bx)\right)}{b^4} \\
 &= \frac{1}{4}x^4 \text{sech}^{-1}(a + bx) - \frac{\text{Subst}\left(\int (-a + \text{sech}(x))^4 dx, x, \text{sech}^{-1}(a + bx)\right)}{4b^4} \\
 &= -\frac{x^2 \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{12b^2} + \frac{1}{4}x^4 \text{sech}^{-1}(a + bx) \\
 &\quad - \frac{\text{Subst}\left(\int (-a + \text{sech}(x))(-3a^3 + (2 + 9a^2)\text{sech}(x) - 8a\text{sech}^2(x)) dx, x, \text{sech}^{-1}(a + bx)\right)}{12b^4} \\
 &= -\frac{x^2 \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{12b^2} + \frac{a(a+bx) \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{3b^4} + \frac{1}{4}x^4 \text{sech}^{-1}(a + bx) \\
 &\quad - \frac{\text{Subst}\left(\int (6a^4 - 12a(1 + 2a^2)\text{sech}(x) + 2(2 + 17a^2)\text{sech}^2(x)) dx, x, \text{sech}^{-1}(a + bx)\right)}{24b^4}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{x^2 \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{12b^2} + \frac{a(a+bx) \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{3b^4} - \frac{a^4 \operatorname{sech}^{-1}(a+bx)}{4b^4} \\
&+ \frac{1}{4} x^4 \operatorname{sech}^{-1}(a+bx) + \frac{(a(1+2a^2)) \operatorname{Subst}(\int \operatorname{sech}(x) dx, x, \operatorname{sech}^{-1}(a+bx))}{2b^4} \\
&- \frac{(2+17a^2) \operatorname{Subst}(\int \operatorname{sech}^2(x) dx, x, \operatorname{sech}^{-1}(a+bx))}{12b^4} \\
&= -\frac{x^2 \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{12b^2} + \frac{a(a+bx) \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{3b^4} - \frac{a^4 \operatorname{sech}^{-1}(a+bx)}{4b^4} \\
&+ \frac{1}{4} x^4 \operatorname{sech}^{-1}(a+bx) + \frac{a(1+2a^2) \arctan\left(\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{a+bx}\right)}{2b^4} \\
&- \frac{(i(2+17a^2)) \operatorname{Subst}(\int 1 dx, x, -i \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx))}{12b^4} \\
&= -\frac{(2+17a^2) \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{12b^4} - \frac{x^2 \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{12b^2} \\
&+ \frac{a(a+bx) \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{3b^4} - \frac{a^4 \operatorname{sech}^{-1}(a+bx)}{4b^4} \\
&+ \frac{1}{4} x^4 \operatorname{sech}^{-1}(a+bx) + \frac{a(1+2a^2) \arctan\left(\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{a+bx}\right)}{2b^4}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.55 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.11

$$\int x^3 \operatorname{sech}^{-1}(a+bx) dx =$$

$$\sqrt{-\frac{-1+a+bx}{1+a+bx}}(2+2a+13a^2+13a^3+(2-4a+9a^2)bx+(1-3a)b^2x^2+b^3x^3)-3b^4x^4\operatorname{sech}^{-1}(a+bx)-$$

[In] Integrate[x^3*ArcSech[a + b*x],x]

[Out] -1/12*(Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(2 + 2*a + 13*a^2 + 13*a^3 + (2 - 4*a + 9*a^2)*b*x + (1 - 3*a)*b^2*x^2 + b^3*x^3) - 3*b^4*x^4*ArcSech[a + b*x] - 3*a^4*Log[a + b*x] + 3*a^4*Log[1 + Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]] + a*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))] + b*x*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]) + (6*I)*a*(1 + 2*a^2)*Log[(-2*I)*(a + b*x) + 2*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]]*(1 + a + b*x))/b^4

Maple [A] (verified)

Time = 0.84 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.23

method	result
derivativedivides	$\frac{\operatorname{arcsech}(bx+a)a^4 - \operatorname{arcsech}(bx+a)a^3(bx+a) + \frac{3}{2}\operatorname{arcsech}(bx+a)a^2(bx+a)^2 - \operatorname{arcsech}(bx+a)a(bx+a)^3 + \frac{\operatorname{arcsech}(bx+a)(bx+a)^4}{4}}{1}$
default	$\frac{\operatorname{arcsech}(bx+a)a^4 - \operatorname{arcsech}(bx+a)a^3(bx+a) + \frac{3}{2}\operatorname{arcsech}(bx+a)a^2(bx+a)^2 - \operatorname{arcsech}(bx+a)a(bx+a)^3 + \frac{\operatorname{arcsech}(bx+a)(bx+a)^4}{4}}{1}$
parts	$\frac{x^4 \operatorname{arcsech}(bx+a)}{4} - \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \left(3 \operatorname{csgn}(b) \operatorname{arctanh}\left(\frac{1}{\sqrt{-b^2x^2-2abx-a^2+1}}\right) a^4 + \operatorname{csgn}(b)b^2x^2\sqrt{-b^2x^2-2abx-a^2+1} \right)}{1}$

[In] `int(x^3*arcsech(b*x+a),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{b^4} \left(\frac{1}{4} \operatorname{arcsech}(bx+a) a^4 - \operatorname{arcsech}(bx+a) a^3 (bx+a) + \frac{3}{2} \operatorname{arcsech}(bx+a) a^2 (bx+a)^2 - \operatorname{arcsech}(bx+a) a (bx+a)^3 + \frac{\operatorname{arcsech}(bx+a) (bx+a)^4}{4} \right. \\ \left. - \frac{1}{12} \left(-\frac{bx+a-1}{bx+a} \right)^{1/2} (bx+a) \left(\frac{bx+a+1}{bx+a} \right)^{1/2} \left(3a^4 \operatorname{arctanh}\left(\frac{1}{\sqrt{1-(bx+a)^2}}\right) + 12a^3 \operatorname{arcsin}(bx+a) + 18a^2 \sqrt{1-(bx+a)^2} - 6a \sqrt{1-(bx+a)^2} \right. \right. \\ \left. \left. + (1-(bx+a)^2)^{1/2} (bx+a)^2 + 6a \operatorname{arcsin}(bx+a) + 2 \sqrt{1-(bx+a)^2} \right) \sqrt{1-(bx+a)^2} \right)$$

Fricas [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 345, normalized size of antiderivative = 1.70

$$\int x^3 \operatorname{sech}^{-1}(a+bx) dx$$

$$= \frac{6b^4x^4 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}+1}}{bx+a}\right) - 3a^4 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}+1}}{x}\right) + 3a^4 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}+1}}{x}\right)}{1}$$

[In] `integrate(x^3*arcsech(b*x+a),x, algorithm="fricas")`

[Out]
$$\frac{1}{24} \left(6b^4x^4 \log\left(\frac{(bx+a)\sqrt{-(b^2x^2+2abx+a^2-1)/(b^2x^2+2abx+a^2)+1}}{bx+a}\right) - 3a^4 \log\left(\frac{(bx+a)\sqrt{-(b^2x^2+2abx+a^2-1)/(b^2x^2+2abx+a^2)+1}}{x}\right) + 3a^4 \log\left(\frac{(bx+a)\sqrt{-(b^2x^2+2abx+a^2-1)/(b^2x^2+2abx+a^2)+1}}{x}\right) \right. \\ \left. + 12(2a^3+a) \operatorname{arctan}\left(\frac{(b^2x^2+2abx+a^2)\sqrt{-(b^2x^2+2abx+a^2-1)/(b^2x^2+2abx+a^2)+1}}{(b^2x^2+2abx+a^2-1)}\right) - 2(b^3x^3 - 3ab^2x^2 + 13a^3 + (9a^2+2)b^2x + 2a) \sqrt{-(b^2x^2+2abx+a^2-1)/(b^2x^2+2abx+a^2)+1} \right) / b^4$$

Sympy [F]

$$\int x^3 \operatorname{sech}^{-1}(a + bx) dx = \int x^3 \operatorname{arsech}(a + bx) dx$$

[In] integrate(x**3*asech(b*x+a),x)

[Out] Integral(x**3*asech(a + b*x), x)

Maxima [F]

$$\int x^3 \operatorname{sech}^{-1}(a + bx) dx = \int x^3 \operatorname{arsech}(bx + a) dx$$

[In] integrate(x^3*arcsech(b*x+a),x, algorithm="maxima")

[Out] 1/8*(2*b^4*x^4*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a) - 2*b^4*x^4*log(b*x + a) - b^2*x^2 + 6*a*b*x - (a^4 + 4*a^3 + 6*a^2 + 4*a + 1)*log(b*x + a + 1) - 2*(b^4*x^4 - a^4)*log(b*x + a) - (a^4 - 4*a^3 + 6*a^2 - 4*a + 1)*log(-b*x - a + 1))/b^4 + integrate(1/4*(b^2*x^5 + a*b*x^4)/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^(1/2*log(b*x + a + 1) + 1/2*log(-b*x - a + 1)) - 1), x)

Giac [F]

$$\int x^3 \operatorname{sech}^{-1}(a + bx) dx = \int x^3 \operatorname{arsech}(bx + a) dx$$

[In] integrate(x^3*arcsech(b*x+a),x, algorithm="giac")

[Out] integrate(x^3*arcsech(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int x^3 \operatorname{sech}^{-1}(a + bx) dx = \int x^3 \operatorname{acosh}\left(\frac{1}{a + bx}\right) dx$$

[In] int(x^3*acosh(1/(a + b*x)),x)

[Out] int(x^3*acosh(1/(a + b*x)), x)

3.2 $\int x^2 \operatorname{sech}^{-1}(a + bx) dx$

Optimal result	61
Rubi [A] (verified)	61
Mathematica [C] (verified)	63
Maple [A] (verified)	64
Fricas [B] (verification not implemented)	64
Sympy [F]	65
Maxima [F]	65
Giac [F]	65
Mupad [F(-1)]	66

Optimal result

Integrand size = 10, antiderivative size = 153

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx = \frac{5a \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6b^3} - \frac{x \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6b^2} + \frac{a^3 \operatorname{sech}^{-1}(a+bx)}{3b^3} + \frac{1}{3} x^3 \operatorname{sech}^{-1}(a+bx) - \frac{(1+6a^2) \arctan\left(\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{a+bx}\right)}{6b^3}$$

[Out] $1/3*a^3*\operatorname{arcsech}(b*x+a)/b^3+1/3*x^3*\operatorname{arcsech}(b*x+a)-1/6*(6*a^2+1)*\arctan((b*x+a+1)*((-b*x-a+1)/(b*x+a+1))^{(1/2)/(b*x+a)})/b^3+5/6*a*(b*x+a+1)*((-b*x-a+1)/(b*x+a+1))^{(1/2)}/b^3-1/6*x*(b*x+a+1)*((-b*x-a+1)/(b*x+a+1))^{(1/2)}/b^2$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6456, 5576, 3867, 3855, 3852, 8}

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx = \frac{a^3 \operatorname{sech}^{-1}(a + bx)}{3b^3} - \frac{(6a^2 + 1) \arctan\left(\frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a+bx}\right)}{6b^3} + \frac{5a \sqrt{\frac{-a-bx+1}{a+bx+1}}(a + bx + 1)}{6b^3} - \frac{x \sqrt{\frac{-a-bx+1}{a+bx+1}}(a + bx + 1)}{6b^2} + \frac{1}{3} x^3 \operatorname{sech}^{-1}(a + bx)$$

[In] Int[x^2*ArcSech[a + b*x],x]

[Out] (5*a*Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x))/(6*b^3) - (x*Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x))/(6*b^2) + (a^3*ArcSech[a + b*x])/(3*b^3) + (x^3*ArcSech[a + b*x])/3 - ((1 + 6*a^2)*ArcTan[(Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x))/(a + b*x])/(6*b^3)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3852

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3867

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_), x_Symbol] := Simp[(-b^2)*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n - 2)/(d*(n - 1))), x] + Dist[1/(n - 1), Int[(a + b*Csc[c + d*x])^(n - 3)*Simp[a^3*(n - 1) + (b*(b^2*(n - 2) + 3*a^2*(n - 1)))*Csc[c + d*x] + (a*b^2*(3*n - 4))*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 2] && IntegerQ[2*n]

Rule 5576

Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sech[(c_.) + (d_.)*(x_)]^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rule 6456

Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int x \operatorname{sech}(x)(-a + \operatorname{sech}(x))^2 \tanh(x) dx, x, \operatorname{sech}^{-1}(a + bx)\right)}{b^3} \\
&= \frac{1}{3}x^3 \operatorname{sech}^{-1}(a + bx) - \frac{\text{Subst}\left(\int (-a + \operatorname{sech}(x))^3 dx, x, \operatorname{sech}^{-1}(a + bx)\right)}{3b^3} \\
&= -\frac{x \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6b^2} + \frac{1}{3}x^3 \operatorname{sech}^{-1}(a + bx) \\
&\quad - \frac{\text{Subst}\left(\int (-2a^3 + (1+6a^2) \operatorname{sech}(x) - 5a \operatorname{sech}^2(x)) dx, x, \operatorname{sech}^{-1}(a + bx)\right)}{6b^3} \\
&= -\frac{x \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6b^2} + \frac{a^3 \operatorname{sech}^{-1}(a + bx)}{3b^3} + \frac{1}{3}x^3 \operatorname{sech}^{-1}(a + bx) \\
&\quad + \frac{(5a) \text{Subst}\left(\int \operatorname{sech}^2(x) dx, x, \operatorname{sech}^{-1}(a + bx)\right)}{6b^3} \\
&\quad - \frac{(1+6a^2) \text{Subst}\left(\int \operatorname{sech}(x) dx, x, \operatorname{sech}^{-1}(a + bx)\right)}{6b^3} \\
&= -\frac{x \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6b^2} + \frac{a^3 \operatorname{sech}^{-1}(a + bx)}{3b^3} \\
&\quad + \frac{1}{3}x^3 \operatorname{sech}^{-1}(a + bx) - \frac{(1+6a^2) \arctan\left(\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{a+bx}\right)}{6b^3} \\
&\quad + \frac{(5ia) \text{Subst}\left(\int 1 dx, x, -i \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\right)}{6b^3} \\
&= \frac{5a \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6b^3} - \frac{x \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6b^2} + \frac{a^3 \operatorname{sech}^{-1}(a + bx)}{3b^3} \\
&\quad + \frac{1}{3}x^3 \operatorname{sech}^{-1}(a + bx) - \frac{(1+6a^2) \arctan\left(\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{a+bx}\right)}{6b^3}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.31

$$\begin{aligned}
&\int x^2 \operatorname{sech}^{-1}(a + bx) dx \\
&= \frac{\sqrt{-\frac{-1+a+bx}{1+a+bx}}(5a^2 - bx(1+bx) + a(5+4bx)) + 2b^3 x^3 \operatorname{sech}^{-1}(a + bx) - 2a^3 \log(a + bx) + 2a^3 \log\left(1 + \sqrt{-\frac{-1+a+bx}{1+a+bx}}\right)}{6b^3}
\end{aligned}$$

[In] Integrate[x^2*ArcSech[a + b*x],x]

[Out] (Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(5*a^2 - b*x*(1 + b*x) + a*(5 + 4*b*x)) + 2*b^3*x^3*ArcSech[a + b*x] - 2*a^3*Log[a + b*x] + 2*a^3*Log[1 + Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]] + a*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))] + b*x*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]] + I*(1 + 6*a^2)*Log[(-2*I)*(a + b*x) + 2*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(1 + a + b*x)]/(6*b^3)

Maple [A] (verified)

Time = 0.73 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.24

method	result
derivativedivides	$\frac{-\frac{\operatorname{arcsech}(bx+a)a^3}{3} + \operatorname{arcsech}(bx+a)a^2(bx+a) - \operatorname{arcsech}(bx+a)a(bx+a)^2 + \frac{\operatorname{arcsech}(bx+a)(bx+a)^3}{3} + \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}}}{b^3}}{b^3}$
default	$\frac{-\frac{\operatorname{arcsech}(bx+a)a^3}{3} + \operatorname{arcsech}(bx+a)a^2(bx+a) - \operatorname{arcsech}(bx+a)a(bx+a)^2 + \frac{\operatorname{arcsech}(bx+a)(bx+a)^3}{3} + \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}}}{b^3}}{b^3}$
parts	$\frac{x^3 \operatorname{arcsech}(bx+a)}{3} + \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \left(2 \operatorname{arctanh}\left(\frac{1}{\sqrt{-b^2x^2-2abx-a^2+1}}\right) \operatorname{csgn}(b)a^3 - \sqrt{-b^2x^2-2abx-a^2+1} \right)}{b^3}$

[In] int(x^2*arcsech(b*x+a),x,method=_RETURNVERBOSE)

[Out] 1/b^3*(-1/3*arcsech(b*x+a)*a^3+arcsech(b*x+a)*a^2*(b*x+a)-arcsech(b*x+a)*a*(b*x+a)^2+1/3*arcsech(b*x+a)*(b*x+a)^3+1/6*(-(b*x+a-1)/(b*x+a))^(1/2)*(b*x+a)*((b*x+a+1)/(b*x+a))^(1/2)*(2*a^3*arctanh(1/(1-(b*x+a)^2))^(1/2))+6*a^2*arcsin(b*x+a)+6*a*(1-(b*x+a)^2)^(1/2)-(b*x+a)*(1-(b*x+a)^2)^(1/2)+arcsin(b*x+a))/(1-(b*x+a)^2)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 327 vs. 2(131) = 262.

Time = 0.30 (sec) , antiderivative size = 327, normalized size of antiderivative = 2.14

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx = \frac{2b^3x^3 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}+1}{bx+a}\right) + a^3 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}+1}{x}\right) - a^3 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}}{x}\right)}{6b^3}$$

[In] integrate(x^2*arcsech(b*x+a),x, algorithm="fricas")


```
[Out] 1/6*(2*b^3*x^3*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)) + a^3*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/x) - a^3*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) - 1)/x) - (6*a^2 + 1)*arctan((b^2*x^2 + 2*a*b*x + a^2)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)))/(b^2*x^2 + 2*a*b*x + a^2 - 1)) - (b^2*x^2 - 4*a*b*x - 5*a^2)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)))/b^3
```

Sympy [F]

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx = \int x^2 \operatorname{arsech}(a + bx) dx$$

```
[In] integrate(x**2*asech(b*x+a),x)
```

```
[Out] Integral(x**2*asech(a + b*x), x)
```

Maxima [F]

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx = \int x^2 \operatorname{arsech}(bx + a) dx$$

```
[In] integrate(x^2*arcsech(b*x+a),x, algorithm="maxima")
```

```
[Out] 1/6*(2*b^3*x^3*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a) - 2*b^3*x^3*log(b*x + a) - 2*b*x + (a^3 + 3*a^2 + 3*a + 1)*log(b*x + a + 1) - 2*(b^3*x^3 + a^3)*log(b*x + a) + (a^3 - 3*a^2 + 3*a - 1)*log(-b*x - a + 1))/b^3 + integrate(1/3*(b^2*x^4 + a*b*x^3)/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^(1/2*log(b*x + a + 1) + 1/2*log(-b*x - a + 1)) - 1), x)
```

Giac [F]

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx = \int x^2 \operatorname{arsech}(bx + a) dx$$

```
[In] integrate(x^2*arcsech(b*x+a),x, algorithm="giac")
```

```
[Out] integrate(x^2*arcsech(b*x + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{sech}^{-1}(a + bx) dx = \int x^2 \operatorname{acosh}\left(\frac{1}{a + bx}\right) dx$$

```
[In] int(x^2*acosh(1/(a + b*x)),x)
```

```
[Out] int(x^2*acosh(1/(a + b*x)), x)
```

3.3 $\int x \operatorname{sech}^{-1}(a + bx) dx$

Optimal result	67
Rubi [A] (verified)	67
Mathematica [C] (verified)	69
Maple [A] (verified)	69
Fricas [B] (verification not implemented)	70
Sympy [F]	70
Maxima [F]	71
Giac [F]	71
Mupad [F(-1)]	71

Optimal result

Integrand size = 8, antiderivative size = 107

$$\int x \operatorname{sech}^{-1}(a + bx) dx = -\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{2b^2} - \frac{a^2 \operatorname{sech}^{-1}(a+bx)}{2b^2} + \frac{1}{2}x^2 \operatorname{sech}^{-1}(a+bx) + \frac{a \arctan\left(\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{a+bx}\right)}{b^2}$$

[Out] $-1/2*a^2*\operatorname{arcsech}(b*x+a)/b^2+1/2*x^2*\operatorname{arcsech}(b*x+a)+a*\arctan((b*x+a+1)*((-b*x-a+1)/(b*x+a+1))^{(1/2)/(b*x+a)})/b^2-1/2*(b*x+a+1)*((-b*x-a+1)/(b*x+a+1))^{(1/2)/b^2}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6456, 5576, 3858, 3855, 3852, 8}

$$\int x \operatorname{sech}^{-1}(a + bx) dx = -\frac{a^2 \operatorname{sech}^{-1}(a + bx)}{2b^2} + \frac{a \arctan\left(\frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{a+bx}\right)}{b^2} - \frac{\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{2b^2} + \frac{1}{2}x^2 \operatorname{sech}^{-1}(a + bx)$$

[In] $\operatorname{Int}[x*\operatorname{ArcSech}[a + b*x], x]$

[Out] $-1/2*(\operatorname{Sqrt}[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x))/b^2 - (a^2*\operatorname{ArcSech}[a + b*x])/(2*b^2) + (x^2*\operatorname{ArcSech}[a + b*x])/2 + (a*\operatorname{ArcTan}[(\operatorname{Sqrt}[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x))/(a + b*x)])/b^2$

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 3852

`Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

Rule 3855

`Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

Rule 3858

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^2, x_Symbol] := Simp[a^2*x, x] + (Dist[2*a*b, Int[Csc[c + d*x], x], x] + Dist[b^2, Int[Csc[c + d*x]^2, x], x]) /; FreeQ[{a, b, c, d}, x]`

Rule 5576

`Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sech[(c_.) + (d_.)*(x_)])^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

Rule 6456

`Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int x \text{sech}(x)(-a + \text{sech}(x)) \tanh(x) dx, x, \text{sech}^{-1}(a + bx)\right)}{b^2} \\
 &= \frac{1}{2}x^2 \text{sech}^{-1}(a + bx) - \frac{\text{Subst}\left(\int (-a + \text{sech}(x))^2 dx, x, \text{sech}^{-1}(a + bx)\right)}{2b^2} \\
 &= -\frac{a^2 \text{sech}^{-1}(a + bx)}{2b^2} + \frac{1}{2}x^2 \text{sech}^{-1}(a + bx) - \frac{\text{Subst}\left(\int \text{sech}^2(x) dx, x, \text{sech}^{-1}(a + bx)\right)}{2b^2} \\
 &\quad + \frac{a \text{Subst}\left(\int \text{sech}(x) dx, x, \text{sech}^{-1}(a + bx)\right)}{b^2}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{a^2 \operatorname{sech}^{-1}(a+bx)}{2b^2} + \frac{1}{2}x^2 \operatorname{sech}^{-1}(a+bx) + \frac{a \arctan\left(\frac{\sqrt{\frac{1-a-bx}{1+bx}}(1+a+bx)}{a+bx}\right)}{b^2} \\
&\quad - \frac{i \operatorname{Subst}\left(\int 1 dx, x, -i\sqrt{\frac{1-a-bx}{1+bx}}(1+a+bx)\right)}{2b^2} \\
&= -\frac{\sqrt{\frac{1-a-bx}{1+bx}}(1+a+bx)}{2b^2} - \frac{a^2 \operatorname{sech}^{-1}(a+bx)}{2b^2} \\
&\quad + \frac{1}{2}x^2 \operatorname{sech}^{-1}(a+bx) + \frac{a \arctan\left(\frac{\sqrt{\frac{1-a-bx}{1+bx}}(1+a+bx)}{a+bx}\right)}{b^2}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.18 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.64

$$\int x \operatorname{sech}^{-1}(a+bx) dx = \frac{-\sqrt{-\frac{-1+a+bx}{1+bx}}(1+a+bx) + b^2 x^2 \operatorname{sech}^{-1}(a+bx) + a^2 \log(a+bx) - a^2 \log\left(1 + \sqrt{-\frac{-1+a+bx}{1+bx}}\right) + a\sqrt{-\frac{-1+a+bx}{1+bx}}}{2b^2}$$

[In] Integrate[x*ArcSech[a + b*x],x]

[Out] $(-\sqrt{-\frac{-1+a+bx}{1+bx}}(1+a+bx) + b^2 x^2 \operatorname{ArcSech}[a + b*x] + a^2 \operatorname{Log}[a + b*x] - a^2 \operatorname{Log}[1 + \sqrt{-\frac{-1+a+bx}{1+bx}}]) + a \sqrt{-\frac{-1+a+bx}{1+bx}} + b*x \sqrt{-\frac{-1+a+bx}{1+bx}} - (2*I)*a \operatorname{Log}[(-2*I)*(a + b*x) + 2*\sqrt{-\frac{-1+a+bx}{1+bx}}](1 + a + b*x)]/(2*b^2)$

Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.04

method	result
derivativedivides	$-\operatorname{arcsech}(bx+a)a(bx+a) + \frac{\operatorname{arcsech}(bx+a)(bx+a)^2}{2} - \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}}(2a \arcsin(bx+a) + \sqrt{1-(bx+a)^2})}{2\sqrt{1-(bx+a)^2}}$
default	$-\operatorname{arcsech}(bx+a)a(bx+a) + \frac{\operatorname{arcsech}(bx+a)(bx+a)^2}{2} - \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}}(2a \arcsin(bx+a) + \sqrt{1-(bx+a)^2})}{2\sqrt{1-(bx+a)^2}}$
parts	$\frac{x^2 \operatorname{arcsech}(bx+a)}{2} - \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}}\left(\operatorname{csgn}(b) \operatorname{arctanh}\left(\frac{1}{\sqrt{-b^2x^2-2abx-a^2+1}}\right)\right)a^2 + \operatorname{csgn}(b)\sqrt{-b^2x^2-2abx-a^2+1}}{2b^2\sqrt{-b^2x^2-2abx-a^2+1}}$

[In] `int(x*arcsech(b*x+a),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{b^2}(-\operatorname{arcsech}(bx+a)a(bx+a)+\frac{1}{2}\operatorname{arcsech}(bx+a)(bx+a)^2-\frac{1}{2}(-(bx+a-1)/(bx+a))^{1/2}(bx+a)((bx+a+1)/(bx+a))^{1/2})(2a\arcsin(bx+a)+(1-(bx+a)^2)^{1/2}))/((1-(bx+a)^2)^{1/2})$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 308 vs. $2(93) = 186$.

Time = 0.28 (sec) , antiderivative size = 308, normalized size of antiderivative = 2.88

$$\int x \operatorname{sech}^{-1}(a + bx) dx$$

$$= \frac{2b^2x^2 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}+1}{bx+a}\right) - a^2 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}}{x}\right) + a^2 \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}}{x}\right)}{4b^2}$$

[In] `integrate(x*arcsech(b*x+a),x, algorithm="fricas")`

[Out] $\frac{1}{4}(2b^2x^2 \log(((bx+a)\sqrt{-(b^2x^2+2abx+a^2-1)/(b^2x^2+2abx+a^2)}+1)/(bx+a)) - a^2 \log(((bx+a)\sqrt{-(b^2x^2+2abx+a^2-1)/(b^2x^2+2abx+a^2)}-1)/x) + a^2 \log(((bx+a)\sqrt{-(b^2x^2+2abx+a^2-1)/(b^2x^2+2abx+a^2)}+1)/x) + 4a \operatorname{arctan}((b^2x^2+2abx+a^2)\sqrt{-(b^2x^2+2abx+a^2-1)/(b^2x^2+2abx+a^2)})/(b^2x^2+2abx+a^2-1)) - 2(bx+a)\sqrt{-(b^2x^2+2abx+a^2-1)/(b^2x^2+2abx+a^2)})/b^2$

Sympy [F]

$$\int x \operatorname{sech}^{-1}(a + bx) dx = \int x \operatorname{asech}(a + bx) dx$$

[In] `integrate(x*asech(b*x+a),x)`

[Out] `Integral(x*asech(a + b*x), x)`

Maxima [F]

$$\int x \operatorname{sech}^{-1}(a + bx) dx = \int x \operatorname{arsech}(bx + a) dx$$

[In] integrate(x*arcsech(b*x+a),x, algorithm="maxima")

[Out] 1/4*(2*b^2*x^2*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a) - 2*b^2*x^2*log(b*x + a) - (a^2 + 2*a + 1)*log(b*x + a + 1) - 2*(b^2*x^2 - a^2)*log(b*x + a) - (a^2 - 2*a + 1)*log(-b*x - a + 1))/b^2 + integrate(1/2*(b^2*x^3 + a*b*x^2)/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^(1/2*log(b*x + a + 1) + 1/2*log(-b*x - a + 1)) - 1), x)

Giac [F]

$$\int x \operatorname{sech}^{-1}(a + bx) dx = \int x \operatorname{arsech}(bx + a) dx$$

[In] integrate(x*arcsech(b*x+a),x, algorithm="giac")

[Out] integrate(x*arcsech(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{sech}^{-1}(a + bx) dx = \int x \operatorname{acosh}\left(\frac{1}{a + bx}\right) dx$$

[In] int(x*acosh(1/(a + b*x)),x)

[Out] int(x*acosh(1/(a + b*x)), x)

3.4 $\int \operatorname{sech}^{-1}(a + bx) dx$

Optimal result	72
Rubi [A] (verified)	72
Mathematica [B] (verified)	73
Maple [A] (verified)	74
Fricas [B] (verification not implemented)	74
Sympy [F]	75
Maxima [A] (verification not implemented)	75
Giac [F]	75
Mupad [B] (verification not implemented)	75

Optimal result

Integrand size = 6, antiderivative size = 44

$$\int \operatorname{sech}^{-1}(a + bx) dx = \frac{(a + bx)\operatorname{sech}^{-1}(a + bx)}{b} - \frac{2 \arctan\left(\sqrt{\frac{1-a-bx}{1+a+bx}}\right)}{b}$$

[Out] (b*x+a)*arcsech(b*x+a)/b-2*arctan(((b*x+a+1)/(b*x+a+1))^(1/2))/b

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {6448, 1983, 12, 209}

$$\int \operatorname{sech}^{-1}(a + bx) dx = \frac{(a + bx)\operatorname{sech}^{-1}(a + bx)}{b} - \frac{2 \arctan\left(\sqrt{\frac{-a-bx+1}{a+bx+1}}\right)}{b}$$

[In] Int[ArcSech[a + b*x], x]

[Out] ((a + b*x)*ArcSech[a + b*x])/b - (2*ArcTan[Sqrt[(1 - a - b*x)/(1 + a + b*x)]])/b

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a

, 0] || GtQ[b, 0])

Rule 1983

```
Int[(u_)^(r_)*(((e_)*((a_) + (b_)*(x_)^(n_.)))/((c_) + (d_)*(x_)^(n_.)
))^ (p_), x_Symbol] := With[{q = Denominator[p]}, Dist[q*e*((b*c - a*d)/n),
Subst[Int[SimplifyIntegrand[x^(q*(p + 1) - 1)*((-a)*e + c*x^q)^(1/n - 1)/(
b*e - d*x^q)^(1/n + 1)]*(u /. x -> ((-a)*e + c*x^q)^(1/n)/(b*e - d*x^q)^(1/
n))^r, x], x], x, (e*((a + b*x^n)/(c + d*x^n)))^(1/q)], x]] /; FreeQ[{a, b,
c, d, e}, x] && PolynomialQ[u, x] && FractionQ[p] && IntegerQ[1/n] && Inte
gerQ[r]
```

Rule 6448

```
Int[ArcSech[(c_) + (d_)*(x_)], x_Symbol] := Simp[(c + d*x)*(ArcSech[c + d*
x]/d), x] + Int[Sqrt[(1 - c - d*x)/(1 + c + d*x)]/(1 - c - d*x), x] /; Free
Q[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(a + bx)\operatorname{sech}^{-1}(a + bx)}{b} + \int \frac{\sqrt{\frac{1-a-bx}{1+a+bx}}}{1-a-bx} dx \\
&= \frac{(a + bx)\operatorname{sech}^{-1}(a + bx)}{b} - (4b)\operatorname{Subst}\left(\int \frac{1}{2b^2(1+x^2)} dx, x, \sqrt{\frac{1-a-bx}{1+a+bx}}\right) \\
&= \frac{(a + bx)\operatorname{sech}^{-1}(a + bx)}{b} - \frac{2\operatorname{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\frac{1-a-bx}{1+a+bx}}\right)}{b} \\
&= \frac{(a + bx)\operatorname{sech}^{-1}(a + bx)}{b} - \frac{2\arctan\left(\sqrt{\frac{1-a-bx}{1+a+bx}}\right)}{b}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 97 vs. 2(44) = 88.

Time = 0.28 (sec) , antiderivative size = 97, normalized size of antiderivative = 2.20

$$\begin{aligned}
\int \operatorname{sech}^{-1}(a + bx) dx &= x\operatorname{sech}^{-1}(a + bx) \\
&\quad - \frac{2\sqrt{-\frac{1+a+bx}{1+a+bx}}\left(-a\arctan\left(\sqrt{\frac{-1+a+bx}{1+a+bx}}\right) + \operatorname{arctanh}\left(\sqrt{\frac{-1+a+bx}{1+a+bx}}\right)\right)}{b\sqrt{\frac{-1+a+bx}{1+a+bx}}}
\end{aligned}$$

[In] Integrate[ArcSech[a + b*x], x]

[Out] x*ArcSech[a + b*x] - (2*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(-(a*ArcTan[Sqrt[-(-1 + a + b*x)/(1 + a + b*x)]])) + ArcTanh[Sqrt[-(-1 + a + b*x)/(1 + a + b*x)]])/(b*Sqrt[-(-1 + a + b*x)/(1 + a + b*x)])

Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00

method	result
derivativeldivides	$\frac{(bx+a) \operatorname{arcsech}(bx+a) - \arctan\left(\sqrt{\frac{1}{bx+a}-1} \sqrt{\frac{1}{bx+a}+1}\right)}{b}$
default	$\frac{(bx+a) \operatorname{arcsech}(bx+a) - \arctan\left(\sqrt{\frac{1}{bx+a}-1} \sqrt{\frac{1}{bx+a}+1}\right)}{b}$
parts	$x \operatorname{arcsech}(bx+a) + \frac{\sqrt{-\frac{bx+a-1}{bx+a}} (bx+a) \sqrt{\frac{bx+a+1}{bx+a}} \left(\operatorname{csgn}(b) \operatorname{arctanh}\left(\frac{1}{\sqrt{-b^2x^2-2abx-a^2+1}}\right) a + \arctan\left(\frac{\operatorname{csgn}(b)}{\sqrt{-(bx+a)}}\right)\right)}{b\sqrt{-b^2x^2-2abx-a^2+1}}$

[In] int(arcsech(b*x+a), x, method=_RETURNVERBOSE)

[Out] 1/b*((b*x+a)*arcsech(b*x+a)-arctan((1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 253 vs. 2(40) = 80.

Time = 0.27 (sec) , antiderivative size = 253, normalized size of antiderivative = 5.75

$$\int \operatorname{sech}^{-1}(a + bx) dx$$

$$= \frac{2bx \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}+1}}{bx+a}\right) + a \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}+1}}{x}\right) - a \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}-1}}{x}\right)}{2b}$$

[In] integrate(arcsech(b*x+a), x, algorithm="fricas")

[Out] 1/2*(2*b*x*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)) + a*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/x) - a*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) - 1)/x) - 2*arctan((b^2*x^2 + 2*a*b*x + a^2)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)))/(b^2*x^2 + 2*a*b*x + a^2 - 1))/b

Sympy [F]

$$\int \operatorname{sech}^{-1}(a + bx) dx = \int \operatorname{arsech}(a + bx) dx$$

[In] integrate(arsech(b*x+a), x)

[Out] Integral(arsech(a + b*x), x)

Maxima [A] (verification not implemented)

none

Time = 0.20 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.70

$$\int \operatorname{sech}^{-1}(a + bx) dx = \frac{(bx + a) \operatorname{arsech}(bx + a) - \arctan\left(\sqrt{\frac{1}{(bx+a)^2} - 1}\right)}{b}$$

[In] integrate(arcsech(b*x+a), x, algorithm="maxima")

[Out] ((b*x + a)*arcsech(b*x + a) - arctan(sqrt(1/(b*x + a)^2 - 1)))/b

Giac [F]

$$\int \operatorname{sech}^{-1}(a + bx) dx = \int \operatorname{arsech}(bx + a) dx$$

[In] integrate(arcsech(b*x+a), x, algorithm="giac")

[Out] integrate(arcsech(b*x + a), x)

Mupad [B] (verification not implemented)

Time = 4.98 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.98

$$\int \operatorname{sech}^{-1}(a + bx) dx = \frac{\operatorname{atan}\left(\frac{1}{\sqrt{\frac{1}{a+bx}-1}\sqrt{\frac{1}{a+bx}+1}}\right) + \operatorname{acosh}\left(\frac{1}{a+bx}\right)(a + bx)}{b}$$

[In] int(acosh(1/(a + b*x)), x)

[Out] (atan(1/((1/(a + b*x) - 1)^(1/2)*(1/(a + b*x) + 1)^(1/2))) + acosh(1/(a + b*x)))*(a + b*x)/b

3.5 $\int \frac{\operatorname{sech}^{-1}(a+bx)}{x} dx$

Optimal result	76
Rubi [A] (verified)	77
Mathematica [C] (verified)	80
Maple [C] (verified)	81
Fricas [F]	82
Sympy [F]	82
Maxima [F]	82
Giac [F]	82
Mupad [F(-1)]	83

Optimal result

Integrand size = 10, antiderivative size = 170

$$\begin{aligned} \int \frac{\operatorname{sech}^{-1}(a+bx)}{x} dx &= \operatorname{sech}^{-1}(a+bx) \log \left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\ &\quad + \operatorname{sech}^{-1}(a+bx) \log \left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\ &\quad - \operatorname{sech}^{-1}(a+bx) \log \left(1 + e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\ &\quad + \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) + \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\ &\quad - \frac{1}{2} \operatorname{PolyLog} \left(2, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) \end{aligned}$$

```
[Out] -arcsech(b*x+a)*ln(1+(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)
+arcsech(b*x+a)*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/
(1-(-a^2+1)^(1/2)))+arcsech(b*x+a)*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1
/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))-1/2*polylog(2,-(1/(b*x+a)+(1/(b*x+a)
-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)+polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2
)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))+polylog(2,a*(1/(b*x+a)+(1/(b*x+a)
-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {6456, 5714, 5689, 3799, 2221, 2317, 2438, 5681}

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x} dx = \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) + \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) \\ + \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) \\ + \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) \\ - \frac{1}{2} \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\ - \operatorname{sech}^{-1}(a+bx) \log\left(e^{2\operatorname{sech}^{-1}(a+bx)} + 1\right)$$

[In] Int[ArcSech[a + b*x]/x, x]

[Out] ArcSech[a + b*x]*Log[1 - (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])] + ArcSech[a + b*x]*Log[1 - (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])] - ArcSech[a + b*x]*Log[1 + E^(2*ArcSech[a + b*x])] + PolyLog[2, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])] + PolyLog[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])] - PolyLog[2, -E^(2*ArcSech[a + b*x])]/2

Rule 2221

Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2317

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_)*((d_) + (e_)*(x_))^(n_)]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3799

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x
_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(
c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x))), x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

Rule 5681

```
Int[(((e_.) + (f_.)*(x_))^(m_.)*Sinh[(c_.) + (d_.)*(x_)])/(Cosh[(c_.) + (d_
.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[-(e + f*x)^(m + 1)/(b*f*(m + 1)),
x] + (Int[(e + f*x)^m*(E^(c + d*x)/(a - Rt[a^2 - b^2, 2] + b*E^(c + d*x)))
, x] + Int[(e + f*x)^m*(E^(c + d*x)/(a + Rt[a^2 - b^2, 2] + b*E^(c + d*x)))
, x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[a^2 - b^2, 0]
```

Rule 5689

```
Int[(((e_.) + (f_.)*(x_))^(m_.)*Tanh[(c_.) + (d_.)*(x_)]^(n_.))/(Cosh[(c_.)
+ (d_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Dist[1/a, Int[(e + f*x)^m*Tanh[c
+ d*x]^n, x], x] - Dist[b/a, Int[(e + f*x)^m*Sinh[c + d*x]*(Tanh[c + d*x]^
(n - 1)/(a + b*Cosh[c + d*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && I
GtQ[m, 0] && IGtQ[n, 0]
```

Rule 5714

```
Int[(((e_.) + (f_.)*(x_))^(m_.)*(F_)[(c_.) + (d_.)*(x_)]^(n_.)*(G_)[(c_.) +
(d_.)*(x_)]^(p_.))/(a_ + (b_.)*Sech[(c_.) + (d_.)*(x_)]), x_Symbol] := I
nt[(e + f*x)^m*Cosh[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Cosh[c + d*x
])), x] /; FreeQ[{a, b, c, d, e, f}, x] && HyperbolicQ[F] && HyperbolicQ[G]
&& IntegersQ[m, n, p]
```

Rule 6456

```
Int[((a_.) + ArcSech[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Ta
nh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\text{Subst}\left(\int \frac{x \operatorname{sech}(x) \tanh(x)}{-a + \operatorname{sech}(x)} dx, x, \operatorname{sech}^{-1}(a + bx)\right) \\ &= -\text{Subst}\left(\int \frac{x \tanh(x)}{1 - a \cosh(x)} dx, x, \operatorname{sech}^{-1}(a + bx)\right) \end{aligned}$$

$$\begin{aligned}
&= -\left(a\text{Subst}\left(\int \frac{x \sinh(x)}{1 - a \cosh(x)} dx, x, \text{sech}^{-1}(a + bx)\right) \right) \\
&\quad - \text{Subst}\left(\int x \tanh(x) dx, x, \text{sech}^{-1}(a + bx)\right) \\
&= -\left(2\text{Subst}\left(\int \frac{e^{2x} x}{1 + e^{2x}} dx, x, \text{sech}^{-1}(a + bx)\right) \right) \\
&\quad - a\text{Subst}\left(\int \frac{e^x x}{1 - \sqrt{1 - a^2} - ae^x} dx, x, \text{sech}^{-1}(a + bx)\right) \\
&\quad - a\text{Subst}\left(\int \frac{e^x x}{1 + \sqrt{1 - a^2} - ae^x} dx, x, \text{sech}^{-1}(a + bx)\right) \\
&= \text{sech}^{-1}(a + bx) \log\left(1 - \frac{ae^{\text{sech}^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}}\right) + \text{sech}^{-1}(a + bx) \log\left(1 - \frac{ae^{\text{sech}^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}}\right) \\
&\quad - \text{sech}^{-1}(a + bx) \log\left(1 + e^{2\text{sech}^{-1}(a+bx)}\right) \\
&\quad - \text{Subst}\left(\int \log\left(1 - \frac{ae^x}{1 - \sqrt{1 - a^2}}\right) dx, x, \text{sech}^{-1}(a + bx)\right) \\
&\quad - \text{Subst}\left(\int \log\left(1 - \frac{ae^x}{1 + \sqrt{1 - a^2}}\right) dx, x, \text{sech}^{-1}(a + bx)\right) \\
&\quad + \text{Subst}\left(\int \log(1 + e^{2x}) dx, x, \text{sech}^{-1}(a + bx)\right) \\
&= \text{sech}^{-1}(a + bx) \log\left(1 - \frac{ae^{\text{sech}^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}}\right) + \text{sech}^{-1}(a + bx) \log\left(1 - \frac{ae^{\text{sech}^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}}\right) \\
&\quad - \text{sech}^{-1}(a + bx) \log\left(1 + e^{2\text{sech}^{-1}(a+bx)}\right) \\
&\quad + \frac{1}{2}\text{Subst}\left(\int \frac{\log(1 + x)}{x} dx, x, e^{2\text{sech}^{-1}(a+bx)}\right) \\
&\quad - \text{Subst}\left(\int \frac{\log\left(1 - \frac{ax}{1 - \sqrt{1 - a^2}}\right)}{x} dx, x, e^{\text{sech}^{-1}(a+bx)}\right) \\
&\quad - \text{Subst}\left(\int \frac{\log\left(1 - \frac{ax}{1 + \sqrt{1 - a^2}}\right)}{x} dx, x, e^{\text{sech}^{-1}(a+bx)}\right) \\
&= \text{sech}^{-1}(a + bx) \log\left(1 - \frac{ae^{\text{sech}^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}}\right) + \text{sech}^{-1}(a + bx) \log\left(1 - \frac{ae^{\text{sech}^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}}\right) \\
&\quad - \text{sech}^{-1}(a + bx) \log\left(1 + e^{2\text{sech}^{-1}(a+bx)}\right) + \text{PolyLog}\left(2, \frac{ae^{\text{sech}^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}}\right) \\
&\quad + \text{PolyLog}\left(2, \frac{ae^{\text{sech}^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}}\right) - \frac{1}{2}\text{PolyLog}\left(2, -e^{2\text{sech}^{-1}(a+bx)}\right)
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 332, normalized size of antiderivative = 1.95

$$\begin{aligned}
\int \frac{\operatorname{sech}^{-1}(a+bx)}{x} dx = & -4i \arcsin\left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}}\right) \operatorname{arctanh}\left(\frac{(1+a) \tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a^2}}\right) \\
& - \operatorname{sech}^{-1}(a+bx) \log\left(1 + e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\
& + \operatorname{sech}^{-1}(a+bx) \log\left(1 + \frac{(-1 + \sqrt{1-a^2}) e^{-\operatorname{sech}^{-1}(a+bx)}}{a}\right) \\
& + 2i \arcsin\left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}}\right) \log\left(1 + \frac{(-1 + \sqrt{1-a^2}) e^{-\operatorname{sech}^{-1}(a+bx)}}{a}\right) \\
& + \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{(1 + \sqrt{1-a^2}) e^{-\operatorname{sech}^{-1}(a+bx)}}{a}\right) \\
& - 2i \arcsin\left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}}\right) \log\left(1 - \frac{(1 + \sqrt{1-a^2}) e^{-\operatorname{sech}^{-1}(a+bx)}}{a}\right) \\
& + \frac{1}{2} \operatorname{PolyLog}\left(2, -e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\
& - \operatorname{PolyLog}\left(2, -\frac{(-1 + \sqrt{1-a^2}) e^{-\operatorname{sech}^{-1}(a+bx)}}{a}\right) \\
& - \operatorname{PolyLog}\left(2, \frac{(1 + \sqrt{1-a^2}) e^{-\operatorname{sech}^{-1}(a+bx)}}{a}\right)
\end{aligned}$$

[In] Integrate[ArcSech[a + b*x]/x,x]

[Out] (-4*I)*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*ArcTanh[(((1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[1 - a^2]] - ArcSech[a + b*x]*Log[1 + E^(-2*ArcSech[a + b*x])] + ArcSech[a + b*x]*Log[1 + (-1 + Sqrt[1 - a^2])/(a*E^ArcSech[a + b*x])] + (2*I)*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*Log[1 + (-1 + Sqrt[1 - a^2])/(a*E^ArcSech[a + b*x])] + ArcSech[a + b*x]*Log[1 - (1 + Sqrt[1 - a^2])/(a*E^ArcSech[a + b*x])] - (2*I)*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*Log[1 - (1 + Sqrt[1 - a^2])/(a*E^ArcSech[a + b*x])] + PolyLog[2, -E^(-2*ArcSech[a + b*x])]/2 - PolyLog[2, -((-1 + Sqrt[1 - a^2])/(a*E^ArcSech[a + b*x]))] - PolyLog[2, (1 + Sqrt[1 - a^2])/(a*E^ArcSech[a + b*x])]

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.88 (sec) , antiderivative size = 882, normalized size of antiderivative = 5.19

method	result
derivativedivides	$\frac{\operatorname{arcsech}(bx+a) \ln\left(\frac{a\left(\frac{1}{bx+a} + \sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right) + \sqrt{-a^2+1}-1}{-1+\sqrt{-a^2+1}}\right)}{2} + \frac{\operatorname{arcsech}(bx+a) \ln\left(\frac{-a\left(\frac{1}{bx+a} + \sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)}{1+\sqrt{-a^2+1}}\right)}{2}$
default	$\frac{\operatorname{arcsech}(bx+a) \ln\left(\frac{a\left(\frac{1}{bx+a} + \sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right) + \sqrt{-a^2+1}-1}{-1+\sqrt{-a^2+1}}\right)}{2} + \frac{\operatorname{arcsech}(bx+a) \ln\left(\frac{-a\left(\frac{1}{bx+a} + \sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)}{1+\sqrt{-a^2+1}}\right)}{2}$

[In] `int(arcsech(b*x+a)/x,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{2} \operatorname{arcsech}(b*x+a) \ln\left(\left(a \cdot \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} \cdot \left(\frac{1}{(b*x+a)} + 1\right)^{1/2}\right) + (-a^2+1)^{1/2} - 1\right) / \left(-1 + (-a^2+1)^{1/2}\right)\right) + \frac{1}{2} \operatorname{arcsech}(b*x+a) \ln\left(\left(-a \cdot \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} \cdot \left(\frac{1}{(b*x+a)} + 1\right)^{1/2}\right) + (-a^2+1)^{1/2} + 1\right) / \left(1 + (-a^2+1)^{1/2}\right)\right) + \frac{1}{2} \cdot (-a^2+1)^{1/2} / (a^2-1) \cdot \operatorname{arcsech}(b*x+a) \ln\left(\left(a \cdot \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} \cdot \left(\frac{1}{(b*x+a)} + 1\right)^{1/2}\right) + (-a^2+1)^{1/2} - 1\right) / \left(-1 + (-a^2+1)^{1/2}\right)\right) - \frac{1}{2} \cdot (-a^2+1)^{1/2} / (a^2-1) \cdot \operatorname{arcsech}(b*x+a) \ln\left(\left(-a \cdot \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} \cdot \left(\frac{1}{(b*x+a)} + 1\right)^{1/2}\right) + (-a^2+1)^{1/2} + 1\right) / \left(1 + (-a^2+1)^{1/2}\right)\right) + \operatorname{dilog}\left(\left(a \cdot \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} \cdot \left(\frac{1}{(b*x+a)} + 1\right)^{1/2}\right) + (-a^2+1)^{1/2} - 1\right) / \left(-1 + (-a^2+1)^{1/2}\right)\right) + \operatorname{dilog}\left(\left(-a \cdot \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} \cdot \left(\frac{1}{(b*x+a)} + 1\right)^{1/2}\right) + (-a^2+1)^{1/2} + 1\right) / \left(1 + (-a^2+1)^{1/2}\right)\right) - \operatorname{arcsech}(b*x+a) \ln\left(1 + I \cdot \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} \cdot \left(\frac{1}{(b*x+a)} + 1\right)^{1/2}\right)\right) - \operatorname{arcsech}(b*x+a) \ln\left(1 - I \cdot \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} \cdot \left(\frac{1}{(b*x+a)} + 1\right)^{1/2}\right)\right) - \operatorname{dilog}\left(1 + I \cdot \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} \cdot \left(\frac{1}{(b*x+a)} + 1\right)^{1/2}\right)\right) - \operatorname{dilog}\left(1 - I \cdot \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} \cdot \left(\frac{1}{(b*x+a)} + 1\right)^{1/2}\right)\right) + \frac{1}{2} \cdot (a^2 + (-a^2+1)^{1/2} - 1) \cdot \operatorname{arcsech}(b*x+a) \cdot \left(\ln\left(\left(-a \cdot \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} \cdot \left(\frac{1}{(b*x+a)} + 1\right)^{1/2}\right) + (-a^2+1)^{1/2} + 1\right) / \left(1 + (-a^2+1)^{1/2}\right)\right) + a^2 \cdot \ln\left(\left(a \cdot \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} \cdot \left(\frac{1}{(b*x+a)} + 1\right)^{1/2}\right) + (-a^2+1)^{1/2} - 1\right) / \left(-1 + (-a^2+1)^{1/2}\right)\right) + a^2 - 2 \cdot \ln\left(\left(a \cdot \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} \cdot \left(\frac{1}{(b*x+a)} + 1\right)^{1/2}\right) + (-a^2+1)^{1/2} - 1\right) / \left(-1 + (-a^2+1)^{1/2}\right)\right) - 2 \cdot \ln\left(\left(-a \cdot \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} \cdot \left(\frac{1}{(b*x+a)} + 1\right)^{1/2}\right) + (-a^2+1)^{1/2} + 1\right) / \left(1 + (-a^2+1)^{1/2}\right)\right) + (-a^2+1)^{1/2} / (a^2-1)$

Fricas [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arosech}(bx + a)}{x} dx$$

[In] integrate(arcsech(b*x+a)/x,x, algorithm="fricas")

[Out] integral(arcsech(b*x + a)/x, x)

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{asech}(a + bx)}{x} dx$$

[In] integrate(asech(b*x+a)/x,x)

[Out] Integral(asech(a + b*x)/x, x)

Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arosech}(bx + a)}{x} dx$$

[In] integrate(arcsech(b*x+a)/x,x, algorithm="maxima")

[Out] integrate(arcsech(b*x + a)/x, x)

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arosech}(bx + a)}{x} dx$$

[In] integrate(arcsech(b*x+a)/x,x, algorithm="giac")

[Out] integrate(arcsech(b*x + a)/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)}{x} dx$$

```
[In] int(acosh(1/(a + b*x))/x,x)
```

```
[Out] int(acosh(1/(a + b*x))/x, x)
```

3.6 $\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^2} dx$

Optimal result	84
Rubi [A] (verified)	84
Mathematica [B] (verified)	86
Maple [B] (verified)	86
Fricas [B] (verification not implemented)	87
Sympy [F]	88
Maxima [F]	88
Giac [F]	88
Mupad [F(-1)]	88

Optimal result

Integrand size = 10, antiderivative size = 70

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^2} dx = -\frac{b\operatorname{sech}^{-1}(a+bx)}{a} - \frac{\operatorname{sech}^{-1}(a+bx)}{x} + \frac{2b\operatorname{arctanh}\left(\frac{\sqrt{1+a}\tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{a\sqrt{1-a^2}}$$

[Out] $-b*\operatorname{arcsech}(b*x+a)/a-\operatorname{arcsech}(b*x+a)/x+2*b*\operatorname{arctanh}((1+a)^{(1/2)}*\tanh(1/2*\operatorname{arcsech}(b*x+a))/(1-a)^{(1/2}))/a/(-a^2+1)^{(1/2)}$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6456, 5576, 3868, 2738, 214}

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^2} dx = \frac{2b\operatorname{arctanh}\left(\frac{\sqrt{a+1}\tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{a\sqrt{1-a^2}} - \frac{b\operatorname{sech}^{-1}(a+bx)}{a} - \frac{\operatorname{sech}^{-1}(a+bx)}{x}$$

[In] $\operatorname{Int}[\operatorname{ArcSech}[a + b*x]/x^2, x]$

[Out] $-((b*\operatorname{ArcSech}[a + b*x])/a) - \operatorname{ArcSech}[a + b*x]/x + (2*b*\operatorname{ArcTanh}[(\operatorname{Sqrt}[1 + a]*\operatorname{Tanh}[\operatorname{ArcSech}[a + b*x]/2])/(\operatorname{Sqrt}[1 - a])])/(a*\operatorname{Sqrt}[1 - a^2])$

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 2738

Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[2*(e/d), Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]

Rule 3868

Int[(csc[(c_) + (d_)*(x_)]*(b_) + (a_))^(-1), x_Symbol] := Simp[x/a, x] - Dist[1/a, Int[1/(1 + (a/b)*Sin[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]

Rule 5576

Int[((e_) + (f_)*(x_))^(m_)*Sech[(c_) + (d_)*(x_)]*((a_) + (b_)*Sech[(c_) + (d_)*(x_)])^(n_)*Tanh[(c_) + (d_)*(x_)], x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rule 6456

Int[((a_) + ArcSech[(c_) + (d_)*(x_)]*(b_))^(p_)*((e_) + (f_)*(x_))^(m_), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\left(b\text{Subst}\left(\int \frac{x\text{sech}(x)\tanh(x)}{(-a + \text{sech}(x))^2} dx, x, \text{sech}^{-1}(a + bx)\right)\right) \\
 &= -\frac{\text{sech}^{-1}(a + bx)}{x} + b\text{Subst}\left(\int \frac{1}{-a + \text{sech}(x)} dx, x, \text{sech}^{-1}(a + bx)\right) \\
 &= -\frac{b\text{sech}^{-1}(a + bx)}{a} - \frac{\text{sech}^{-1}(a + bx)}{x} + \frac{b\text{Subst}\left(\int \frac{1}{1 - a\cosh(x)} dx, x, \text{sech}^{-1}(a + bx)\right)}{a} \\
 &= -\frac{b\text{sech}^{-1}(a + bx)}{a} - \frac{\text{sech}^{-1}(a + bx)}{x} \\
 &\quad + \frac{(2b)\text{Subst}\left(\int \frac{1}{1 - a - (1+a)x^2} dx, x, \tanh\left(\frac{1}{2}\text{sech}^{-1}(a + bx)\right)\right)}{a}
 \end{aligned}$$

$$= -\frac{b \operatorname{sech}^{-1}(a+bx)}{a} - \frac{\operatorname{sech}^{-1}(a+bx)}{x} + \frac{2b \operatorname{arctanh}\left(\frac{\sqrt{1+a} \tanh\left(\frac{1}{2} \operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{a\sqrt{1-a^2}}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 244 vs. $2(70) = 140$.

Time = 0.27 (sec) , antiderivative size = 244, normalized size of antiderivative = 3.49

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^2} dx = -\frac{\operatorname{sech}^{-1}(a+bx)}{x} + \frac{b\left(-\log(x) + \sqrt{1-a^2} \log(a+bx) - \sqrt{1-a^2} \log\left(1 + \sqrt{-\frac{-1+a+bx}{1+a+bx}} + a\sqrt{-\frac{-1+a+bx}{1+a+bx}} + bx\sqrt{-\frac{-1+a+bx}{1+a+bx}}\right)\right)}{a\sqrt{1-a^2}}$$

[In] Integrate[ArcSech[a + b*x]/x^2,x]

[Out] $-(\operatorname{ArcSech}[a + b*x]/x) + (b*(-\operatorname{Log}[x] + \operatorname{Sqrt}[1 - a^2]*\operatorname{Log}[a + b*x] - \operatorname{Sqrt}[1 - a^2]*\operatorname{Log}[1 + \operatorname{Sqrt}[-(1 + a + b*x)/(1 + a + b*x)]] + a*\operatorname{Sqrt}[-(1 + a + b*x)/(1 + a + b*x)]] + b*x*\operatorname{Sqrt}[-(1 + a + b*x)/(1 + a + b*x)]] + \operatorname{Log}[1 - a^2 - a*b*x + \operatorname{Sqrt}[1 - a^2]*\operatorname{Sqrt}[-(1 + a + b*x)/(1 + a + b*x)]] + a*\operatorname{Sqrt}[1 - a^2]*\operatorname{Sqrt}[-(1 + a + b*x)/(1 + a + b*x)]] + \operatorname{Sqrt}[1 - a^2]*b*x*\operatorname{Sqrt}[-(1 + a + b*x)/(1 + a + b*x)]])/(a*\operatorname{Sqrt}[1 - a^2])$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 170 vs. $2(62) = 124$.

Time = 0.95 (sec) , antiderivative size = 171, normalized size of antiderivative = 2.44

method	result
derivativedivides	$b \left(-\frac{\operatorname{arcsech}(bx+a)}{bx} - \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{1-(bx+a)^2}}\right) a^2 + \sqrt{-a^2+1} \ln\left(\frac{2\sqrt{-a^2+1}\sqrt{1-(bx+a)}}{bx}\right)}{\sqrt{1-(bx+a)^2} a(-1+a)(1+a)} \right)}{\sqrt{1-(bx+a)^2} a(-1+a)(1+a)} \right)$
default	$b \left(-\frac{\operatorname{arcsech}(bx+a)}{bx} - \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{1-(bx+a)^2}}\right) a^2 + \sqrt{-a^2+1} \ln\left(\frac{2\sqrt{-a^2+1}\sqrt{1-(bx+a)}}{bx}\right)}{\sqrt{1-(bx+a)^2} a(-1+a)(1+a)} \right)}{\sqrt{1-(bx+a)^2} a(-1+a)(1+a)} \right)$
parts	$-\frac{\operatorname{arcsech}(bx+a)}{x} - \frac{b\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{-b^2x^2-2abx-a^2+1}}\right) a^2 + \sqrt{-a^2+1} \ln\left(\frac{-2a^2+2-2abx+2bx^2}{\sqrt{-b^2x^2-2abx-a^2+1}(1+a)}\right)}{\sqrt{-b^2x^2-2abx-a^2+1}(1+a)} \right)}{\sqrt{-b^2x^2-2abx-a^2+1}(1+a)}$

[In] int(arcsech(b*x+a)/x^2,x,method=_RETURNVERBOSE)

[Out] $b*(-1/b/x*\operatorname{arcsech}(b*x+a)-(-(b*x+a-1)/(b*x+a))^{1/2}*(b*x+a)*((b*x+a+1)/(b*x+a))^{1/2}*(\operatorname{arctanh}(1/(1-(b*x+a)^2)^{1/2}))*a^2+(-a^2+1)^{1/2}*ln(2*((-a^2+1)^{1/2}*(1-(b*x+a)^2)^{1/2}-(b*x+a)*a+1)/b/x)-\operatorname{arctanh}(1/(1-(b*x+a)^2)^{1/2}))/((1-(b*x+a)^2)^{1/2})/a/(-1+a)/(1+a))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 320 vs. $2(62) = 124$.

Time = 0.30 (sec) , antiderivative size = 651, normalized size of antiderivative = 9.30

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^2} dx$$

$$= \frac{\left[(a^2-1)bx \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}+1}{x}\right) - (a^2-1)bx \log\left(\frac{(bx+a)\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}-1}{x}\right) + \sqrt{-a^2+1}bx \right]}{2(a^3 - \dots)}$$

[In] `integrate(arcsech(b*x+a)/x^2,x, algorithm="fricas")`

[Out] $[-1/2*((a^2-1)*b*x*\log(((b*x+a)*\sqrt{-(b^2*x^2+2*a*b*x+a^2-1)/(b^2*x^2+2*a*b*x+a^2)}+1)/x) - (a^2-1)*b*x*\log(((b*x+a)*\sqrt{-(b^2*x^2+2*a*b*x+a^2-1)/(b^2*x^2+2*a*b*x+a^2)}-1)/x) + \sqrt{-a^2+1})*b*x*\log(((2*a^2-1)*b^2*x^2+2*a^4+4*(a^3-a)*b*x-4*a^2-2*(a*b^2*x^2+a^3+(2*a^2-1)*b*x-a)*\sqrt{-a^2+1}*\sqrt{-(b^2*x^2+2*a*b*x+a^2-1)/(b^2*x^2+2*a*b*x+a^2)}+2)/x^2) + 2*(a^3-a)*\log(((b*x+a)*\sqrt{-(b^2*x^2+2*a*b*x+a^2-1)/(b^2*x^2+2*a*b*x+a^2)}+1)/(b*x+a)))/((a^3-a)*x), -1/2*((a^2-1)*b*x*\log(((b*x+a)*\sqrt{-(b^2*x^2+2*a*b*x+a^2-1)/(b^2*x^2+2*a*b*x+a^2)}+1)/x) - (a^2-1)*b*x*\log(((b*x+a)*\sqrt{-(b^2*x^2+2*a*b*x+a^2-1)/(b^2*x^2+2*a*b*x+a^2)}-1)/x) + 2*\sqrt{a^2-1}*b*x*\operatorname{arctan}((a*b^2*x^2+a^3+(2*a^2-1)*b*x-a)*\sqrt{a^2-1}*\sqrt{-(b^2*x^2+2*a*b*x+a^2-1)/(b^2*x^2+2*a*b*x+a^2)})/((a^2-1)*b^2*x^2+a^4+2*(a^3-a)*b*x-2*a^2+1)) + 2*(a^3-a)*\log(((b*x+a)*\sqrt{-(b^2*x^2+2*a*b*x+a^2-1)/(b^2*x^2+2*a*b*x+a^2)}+1)/(b*x+a)))/((a^3-a)*x)]$

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{arsech}(a + bx)}{x^2} dx$$

[In] integrate(arsech(b*x+a)/x**2,x)

[Out] Integral(arsech(a + b*x)/x**2, x)

Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{arsech}(bx + a)}{x^2} dx$$

[In] integrate(arcsech(b*x+a)/x^2,x, algorithm="maxima")

[Out] b*log(x)/(a^3 - a) - 1/2*((a^2*b - a*b)*x*log(b*x + a + 1) + (a^2*b + a*b)*x*log(-b*x - a + 1) + 2*(a^3 - a)*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a) - 2*(a^3 + (a^2*b - b)*x - a)*log(b*x + a) - 2*(a^3 - a)*log(b*x + a))/((a^3 - a)*x) - integrate((b^2*x + a*b)/(b^2*x^3 + 2*a*b*x^2 + (a^2 - 1)*x + (b^2*x^3 + 2*a*b*x^2 + (a^2 - 1)*x)*e^(1/2*log(b*x + a + 1) + 1/2*log(-b*x - a + 1))), x)

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{arsech}(bx + a)}{x^2} dx$$

[In] integrate(arcsech(b*x+a)/x^2,x, algorithm="giac")

[Out] integrate(arcsech(b*x + a)/x^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)}{x^2} dx$$

[In] int(acosh(1/(a + b*x))/x^2,x)

[Out] int(acosh(1/(a + b*x))/x^2, x)

3.7 $\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^3} dx$

Optimal result	89
Rubi [A] (verified)	89
Mathematica [B] (verified)	92
Maple [C] (verified)	93
Fricas [B] (verification not implemented)	93
Sympy [F]	94
Maxima [F]	94
Giac [F]	95
Mupad [F(-1)]	95

Optimal result

Integrand size = 10, antiderivative size = 133

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^3} dx = \frac{b\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{2a(1-a^2)x} + \frac{b^2\operatorname{sech}^{-1}(a+bx)}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)}{2x^2} - \frac{(1-2a^2)b^2\operatorname{arctanh}\left(\frac{\sqrt{1+a}\tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{a^2(1-a^2)^{3/2}}$$

[Out] $1/2*b^2*\operatorname{arcsech}(b*x+a)/a^2-1/2*\operatorname{arcsech}(b*x+a)/x^2-(-2*a^2+1)*b^2*\operatorname{arctanh}((1+a)^{(1/2)}*\tanh(1/2*\operatorname{arcsech}(b*x+a))/(1-a)^{(1/2)})/a^2/(-a^2+1)^{(3/2)}+1/2*b*(b*x+a+1)*((-b*x-a+1)/(b*x+a+1))^{(1/2)}/a/(-a^2+1)/x$

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {6456, 5576, 3870, 4004, 3916, 2738, 214}

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^3} dx = -\frac{(1-2a^2)b^2\operatorname{arctanh}\left(\frac{\sqrt{a+1}\tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{a^2(1-a^2)^{3/2}} + \frac{b^2\operatorname{sech}^{-1}(a+bx)}{2a^2} + \frac{b\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{2a(1-a^2)x} - \frac{\operatorname{sech}^{-1}(a+bx)}{2x^2}$$

[In] Int[ArcSech[a + b*x]/x^3,x]

[Out] $(b\sqrt{(1-a-bx)/(1+a+bx)}(1+a+bx))/(2a(1-a^2)x) + (b^2\text{ArcSech}[a+bx])/(2a^2) - \text{ArcSech}[a+bx]/(2x^2) - ((1-2a^2)b^2\text{ArcTanh}[\sqrt{1+a}\text{Tanh}[\text{ArcSech}[a+bx]/2]]/\sqrt{1-a})/(a^2(1-a^2)^{(3/2)})$

Rule 214

$\text{Int}[(a_.) + (b_.)x^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b, x\} \&\& \text{NegQ}[a/b]$

Rule 2738

$\text{Int}[(a_.) + (b_.)\sin[\pi/2 + (c_.) + (d_.)x])^{-1}, x_Symbol] \rightarrow \text{With}\{e = \text{FreeFactors}[\text{Tan}[(c + dx)/2], x], \text{Dist}[2(e/d), \text{Subst}[\text{Int}[1/(a + b + (a - b)e^2x^2), x], x, \text{Tan}[(c + dx)/2]/e], x]\} /; \text{FreeQ}\{a, b, c, d, x\} \&\& \text{NeQ}[a^2 - b^2, 0]$

Rule 3870

$\text{Int}[(\text{csc}[(c_.) + (d_.)x] + (b_.) + (a_.)^n), x_Symbol] \rightarrow \text{Simp}[b^2\text{Cot}[c + dx]*(a + b\text{Csc}[c + dx])^{n+1}/(a*d*(n+1)*(a^2 - b^2)), x] + \text{Dist}[1/(a*(n+1)*(a^2 - b^2)), \text{Int}[(a + b\text{Csc}[c + dx])^{n+1}*\text{Simp}[(a^2 - b^2)*(n+1) - a*b*(n+1)*\text{Csc}[c + dx] + b^2*(n+2)*\text{Csc}[c + dx]^2, x], x], x] /; \text{FreeQ}\{a, b, c, d, x\} \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{LtQ}[n, -1] \&\& \text{IntegerQ}[2*n]$

Rule 3916

$\text{Int}[\text{csc}[(e_.) + (f_.)x]/(\text{csc}[(e_.) + (f_.)x] + (b_.) + (a_)), x_Symbol] \rightarrow \text{Dist}[1/b, \text{Int}[1/(1 + (a/b)\text{Sin}[e + fx]), x], x] /; \text{FreeQ}\{a, b, e, f, x\} \&\& \text{NeQ}[a^2 - b^2, 0]$

Rule 4004

$\text{Int}[(\text{csc}[(e_.) + (f_.)x] + (d_.) + (c_))/(\text{csc}[(e_.) + (f_.)x] + (b_.) + (a_)), x_Symbol] \rightarrow \text{Simp}[c*(x/a), x] - \text{Dist}[(b*c - a*d)/a, \text{Int}[\text{Csc}[e + fx]/(a + b\text{Csc}[e + fx]), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, x\} \&\& \text{NeQ}[b*c - a*d, 0]$

Rule 5576

$\text{Int}[(e_.) + (f_.)x)^{m_.*\text{Sech}[(c_.) + (d_.)x] * ((a_.) + (b_.)\text{Sech}[(c_.) + (d_.)x])^{n_.*\text{Tanh}[(c_.) + (d_.)x]}, x_Symbol] \rightarrow \text{Simp}[(-e + fx)^m * ((a + b\text{Sech}[c + dx])^{n+1}/(b*d*(n+1))), x] + \text{Dist}[f*(m/(b*d*(n+1))), \text{Int}[(e + fx)^{m-1} * (a + b\text{Sech}[c + dx])^{n+1}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n, x\} \&\& \text{IGtQ}[m, 0] \&\& \text{NeQ}[n, -1]$

Rule 6456

Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol] :> Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rubi steps

$$\begin{aligned}
\text{integral} &= -\left(b^2 \text{Subst}\left(\int \frac{x \operatorname{sech}(x) \tanh(x)}{(-a + \operatorname{sech}(x))^3} dx, x, \operatorname{sech}^{-1}(a + bx)\right)\right) \\
&= -\frac{\operatorname{sech}^{-1}(a + bx)}{2x^2} + \frac{1}{2} b^2 \text{Subst}\left(\int \frac{1}{(-a + \operatorname{sech}(x))^2} dx, x, \operatorname{sech}^{-1}(a + bx)\right) \\
&= \frac{b\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{2a(1-a^2)x} - \frac{\operatorname{sech}^{-1}(a+bx)}{2x^2} - \frac{b^2 \text{Subst}\left(\int \frac{1-a^2-a\operatorname{sech}(x)}{-a+\operatorname{sech}(x)} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{2a(1-a^2)} \\
&= \frac{b\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{2a(1-a^2)x} + \frac{b^2 \operatorname{sech}^{-1}(a+bx)}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)}{2x^2} \\
&\quad - \frac{((1-2a^2)b^2) \text{Subst}\left(\int \frac{\operatorname{sech}(x)}{-a+\operatorname{sech}(x)} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{2a^2(1-a^2)} \\
&= \frac{b\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{2a(1-a^2)x} + \frac{b^2 \operatorname{sech}^{-1}(a+bx)}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)}{2x^2} \\
&\quad - \frac{((1-2a^2)b^2) \text{Subst}\left(\int \frac{1}{1-a\cosh(x)} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{2a^2(1-a^2)} \\
&= \frac{b\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{2a(1-a^2)x} + \frac{b^2 \operatorname{sech}^{-1}(a+bx)}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)}{2x^2} \\
&\quad - \frac{((1-2a^2)b^2) \text{Subst}\left(\int \frac{1}{1-a-(1+a)x^2} dx, x, \tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)\right)}{a^2(1-a^2)} \\
&= \frac{b\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{2a(1-a^2)x} + \frac{b^2 \operatorname{sech}^{-1}(a+bx)}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)}{2x^2} \\
&\quad - \frac{(1-2a^2)b^2 \operatorname{arctanh}\left(\frac{\sqrt{1+a}\tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{a^2(1-a^2)^{3/2}}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 315 vs. $2(133) = 266$.

Time = 0.90 (sec) , antiderivative size = 315, normalized size of antiderivative = 2.37

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^3} dx$$

$$= \frac{1}{2} \left(-\frac{b\sqrt{-\frac{-1+a+bx}{1+a+bx}}(1+a+bx)}{(-1+a)a(1+a)x} - \frac{\operatorname{sech}^{-1}(a+bx)}{x^2} - \frac{(-1+2a^2)b^2 \log(x)}{a^2(1-a^2)^{3/2}} \right.$$

$$\left. - \frac{b^2 \log(a+bx)}{a^2} + \frac{b^2 \log\left(1 + \sqrt{-\frac{-1+a+bx}{1+a+bx}} + a\sqrt{-\frac{-1+a+bx}{1+a+bx}} + bx\sqrt{-\frac{-1+a+bx}{1+a+bx}}\right)}{a^2} \right.$$

$$\left. + \frac{(-1+2a^2)b^2 \log\left(1 - a^2 - abx + \sqrt{1-a^2}\sqrt{-\frac{-1+a+bx}{1+a+bx}} + a\sqrt{1-a^2}\sqrt{-\frac{-1+a+bx}{1+a+bx}} + \sqrt{1-a^2}bx\sqrt{-\frac{-1+a+bx}{1+a+bx}}\right)}{a^2(1-a^2)^{3/2}} \right)$$

[In] Integrate[ArcSech[a + b*x]/x^3,x]

[Out] $\left(-\left(\frac{b\sqrt{-\left(-1+a+bx\right)/\left(1+a+bx\right)}}{\left(-1+a\right)a\left(1+a\right)x}\right) - \frac{\operatorname{ArcSech}\left[a+bx\right]}{x^2} - \frac{\left(-1+2a^2\right)b^2\operatorname{Log}\left[x\right]}{a^2\left(1-a^2\right)^{3/2}} - \frac{b^2\operatorname{Log}\left[a+bx\right]}{a^2} + \frac{b^2\operatorname{Log}\left[1+\sqrt{-\left(-1+a+bx\right)/\left(1+a+bx\right)} + a\sqrt{-\left(-1+a+bx\right)/\left(1+a+bx\right)} + bx\sqrt{-\left(-1+a+bx\right)/\left(1+a+bx\right)}\right]}{a^2} + \frac{\left(-1+2a^2\right)b^2\operatorname{Log}\left[1-a^2-abx+\sqrt{1-a^2}\sqrt{-\left(-1+a+bx\right)/\left(1+a+bx\right)} + a\sqrt{1-a^2}\sqrt{-\left(-1+a+bx\right)/\left(1+a+bx\right)} + \sqrt{1-a^2}bx\sqrt{-\left(-1+a+bx\right)/\left(1+a+bx\right)}\right]}{a^2\left(1-a^2\right)^{3/2}}\right)/2$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.96 (sec) , antiderivative size = 370, normalized size of antiderivative = 2.78

method	result
parts	$-\frac{\operatorname{arcsech}(bx+a)}{2x^2} - \frac{b\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \operatorname{csgn}(b)^2 \left(-\operatorname{arctanh}\left(\frac{1}{\sqrt{-b^2x^2-2abx-a^2+1}}\right)\right) a^4 bx - 2\sqrt{-a^2+1} \ln\left(\frac{1}{\sqrt{-b^2x^2-2abx-a^2+1}}\right)}{2x^2}$
derivativedivides	$b^2 \left(-\frac{\operatorname{arcsech}(bx+a)}{2b^2x^2} - \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{1-(bx+a)^2}}\right)\right) a^5 - \operatorname{arctanh}\left(\frac{1}{\sqrt{1-(bx+a)^2}}\right) a^4 (bx+a)}{2b^2x^2} \right)$
default	$b^2 \left(-\frac{\operatorname{arcsech}(bx+a)}{2b^2x^2} - \frac{\sqrt{-\frac{bx+a-1}{bx+a}}(bx+a)\sqrt{\frac{bx+a+1}{bx+a}} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{1-(bx+a)^2}}\right)\right) a^5 - \operatorname{arctanh}\left(\frac{1}{\sqrt{1-(bx+a)^2}}\right) a^4 (bx+a)}{2b^2x^2} \right)$

[In] `int(arcsech(b*x+a)/x^3,x,method=_RETURNVERBOSE)`

[Out]
$$-1/2*\operatorname{arcsech}(b*x+a)/x^2 - 1/2*b*(-(b*x+a-1)/(b*x+a))^{1/2}*(b*x+a)*((b*x+a+1)/(b*x+a))^{1/2}* \operatorname{csgn}(b)^2*(-\operatorname{arctanh}(1/(-b^2*x^2-2*a*b*x-a^2+1)^{1/2}))*a^4*b*x - 2*(-a^2+1)^{1/2}*\ln(2*(-a*b*x+(-a^2+1)^{1/2}*(-b^2*x^2-2*a*b*x-a^2+1)^{1/2}-a^2+1)/x)*a^2*b*x+2*\operatorname{arctanh}(1/(-b^2*x^2-2*a*b*x-a^2+1)^{1/2}))*a^2*b*x+a^3*(-b^2*x^2-2*a*b*x-a^2+1)^{1/2}+(-a^2+1)^{1/2}*\ln(2*(-a*b*x+(-a^2+1)^{1/2}*(-b^2*x^2-2*a*b*x-a^2+1)^{1/2}-a^2+1)/x)*b*x-\operatorname{arctanh}(1/(-b^2*x^2-2*a*b*x-a^2+1)^{1/2}))*b*x-(-b^2*x^2-2*a*b*x-a^2+1)^{1/2}*a/(-b^2*x^2-2*a*b*x-a^2+1)^{1/2}/(1+a)/(-1+a)/a^2/(a^2-1)/x$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 427 vs. 2(112) = 224.

Time = 0.32 (sec) , antiderivative size = 865, normalized size of antiderivative = 6.50

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^3} dx$$

$$= \left[\frac{(2a^2-1)\sqrt{-a^2+1}b^2x^2 \log\left(\frac{(2a^2-1)b^2x^2+2a^4+4(a^3-a)bx-4a^2+2(ab^2x^2+a^3+(2a^2-1)bx-a)\sqrt{-a^2+1}\sqrt{-\frac{b^2x^2+2abx+a^2}{b^2x^2+2abx+a^2}}}{x^2}\right)}{x^2} \right]$$

[In] `integrate(arcsech(b*x+a)/x^3,x, algorithm="fricas")`

```
[Out] [-1/4*((2*a^2 - 1)*sqrt(-a^2 + 1)*b^2*x^2*log(((2*a^2 - 1)*b^2*x^2 + 2*a^4 + 4*(a^3 - a)*b*x - 4*a^2 + 2*(a*b^2*x^2 + a^3 + (2*a^2 - 1)*b*x - a)*sqrt(-a^2 + 1)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 2)/x^2) - (a^4 - 2*a^2 + 1)*b^2*x^2*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/x) + (a^4 - 2*a^2 + 1)*b^2*x^2*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) - 1)/x) + 2*(a^6 - 2*a^4 + a^2)*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)) + 2*((a^3 - a)*b^2*x^2 + (a^4 - a^2)*b*x)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)))/((a^6 - 2*a^4 + a^2)*x^2), 1/4*(2*(2*a^2 - 1)*sqrt(a^2 - 1)*b^2*x^2*arctan((a*b^2*x^2 + a^3 + (2*a^2 - 1)*b*x - a)*sqrt(a^2 - 1)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)))/((a^2 - 1)*b^2*x^2 + a^4 + 2*(a^3 - a)*b*x - 2*a^2 + 1)) + (a^4 - 2*a^2 + 1)*b^2*x^2*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/x) - (a^4 - 2*a^2 + 1)*b^2*x^2*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) - 1)/x) - 2*(a^6 - 2*a^4 + a^2)*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)) - 2*((a^3 - a)*b^2*x^2 + (a^4 - a^2)*b*x)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)))/((a^6 - 2*a^4 + a^2)*x^2)]
```

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{asech}(a + bx)}{x^3} dx$$

```
[In] integrate(asech(b*x+a)/x**3,x)
```

```
[Out] Integral(asech(a + b*x)/x**3, x)
```

Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{arsech}(bx + a)}{x^3} dx$$

```
[In] integrate(arcsech(b*x+a)/x^3,x, algorithm="maxima")
```

```
[Out] -1/2*(3*a^2*b^2 - b^2)*log(x)/(a^6 - 2*a^4 + a^2) + 1/4*((a^4*b^2 - 2*a^3*b^2 + a^2*b^2)*x^2*log(b*x + a + 1) + (a^4*b^2 + 2*a^3*b^2 + a^2*b^2)*x^2*log(-b*x - a + 1) - 2*(a^3*b - a*b)*x - 2*(a^6 - 2*a^4 + a^2)*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a) + 2*(a^6 - 2*a^4 - (a^4*b^2 - 2*a^2*b^2 + b^2)*x^2 + a^2)*log(b*x + a) + 2*(a^6 - 2*a^4 + a^2)*log(b*x + a))/((a^6 - 2*a^4 + a^2)*x^2) - integrate(1/2*(b^2*x + a*b)/(b^2*x^4 + 2*a*b*x^3 + (a^2 - 1)*x^2 + (b^2*x^4 + 2*a*b*x^3 + (a^2 - 1)*x^2)*e^(1/2*log(b*x + a + 1) + 1/2*log(-b*x - a + 1))), x)
```

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{ar} \operatorname{sech}(bx + a)}{x^3} dx$$

[In] integrate(arcsech(b*x+a)/x^3,x, algorithm="giac")

[Out] integrate(arcsech(b*x + a)/x^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)}{x^3} dx$$

[In] int(acosh(1/(a + b*x))/x^3,x)

[Out] int(acosh(1/(a + b*x))/x^3, x)

3.8 $\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^4} dx$

Optimal result	96
Rubi [A] (verified)	97
Mathematica [A] (verified)	100
Maple [C] (verified)	100
Fricas [B] (verification not implemented)	101
Sympy [F]	102
Maxima [F]	102
Giac [F]	103
Mupad [F(-1)]	103

Optimal result

Integrand size = 10, antiderivative size = 197

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^4} dx = \frac{b\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6a(1-a^2)x^2} - \frac{(2-5a^2)b^2\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6a^2(1-a^2)^2x}$$

$$- \frac{b^3\operatorname{sech}^{-1}(a+bx)}{3a^3} - \frac{\operatorname{sech}^{-1}(a+bx)}{3x^3}$$

$$+ \frac{(2-5a^2+6a^4)b^3\operatorname{arctanh}\left(\frac{\sqrt{1+a}\tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{3a^3(1-a^2)^{5/2}}$$

```
[Out] -1/3*b^3*arcsech(b*x+a)/a^3-1/3*arcsech(b*x+a)/x^3+1/3*(6*a^4-5*a^2+2)*b^3*
arctanh((1+a)^(1/2)*tanh(1/2*arcsech(b*x+a))/(1-a)^(1/2))/a^3/(-a^2+1)^(5/2
)+1/6*b*(b*x+a+1)*((-b*x-a+1)/(b*x+a+1))^(1/2)/a/(-a^2+1)/x^2-1/6*(-5*a^2+2
)*b^2*(b*x+a+1)*((-b*x-a+1)/(b*x+a+1))^(1/2)/a^2/(-a^2+1)^2/x
```


Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {6456, 5576, 3870, 4145, 4004, 3916, 2738, 214}

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^4} dx = -\frac{b^3 \operatorname{sech}^{-1}(a+bx)}{3a^3} - \frac{(2-5a^2)b^2 \sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{6a^2(1-a^2)^2 x} + \frac{b \sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)}{6a(1-a^2)x^2} + \frac{(6a^4-5a^2+2)b^3 \operatorname{arctanh}\left(\frac{\sqrt{a+1} \tanh\left(\frac{1}{2} \operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{3a^3(1-a^2)^{5/2}} - \frac{\operatorname{sech}^{-1}(a+bx)}{3x^3}$$

[In] Int[ArcSech[a + b*x]/x^4,x]

[Out] (b*Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x))/(6*a*(1 - a^2)*x^2) - (2 - 5*a^2)*b^2*Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x)/(6*a^2*(1 - a^2)^2*x) - (b^3*ArcSech[a + b*x])/(3*a^3) - ArcSech[a + b*x]/(3*x^3) + ((2 - 5*a^2 + 6*a^4)*b^3*ArcTanh[(Sqrt[1 + a]*Tanh[ArcSech[a + b*x]/2])/Sqrt[1 - a]])/(3*a^3*(1 - a^2)^(5/2))

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 2738

Int[((a_) + (b_)*sin[Pi/2 + (c_.) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[2*(e/d), Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]

Rule 3870

Int[(csc[(c_.) + (d_)*(x_)])*(b_.) + (a_))^(n_), x_Symbol] := Simp[b^2*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n + 1)/(a*d*(n + 1)*(a^2 - b^2))), x] + Dist[1/(a*(n + 1)*(a^2 - b^2)), Int[(a + b*Csc[c + d*x])^(n + 1)*Simp[(a^2 - b^2)*(n + 1) - a*b*(n + 1)*Csc[c + d*x] + b^2*(n + 2)*Csc[c + d*x]^2, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3916

```
Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol]
:> Dist[1/b, Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x]
&& NeQ[a^2 - b^2, 0]
```

Rule 4004

```
Int[(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol]
:> Simp[c*(x/a), x] - Dist[(b*c - a*d)/a, Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x]
&& NeQ[b*c - a*d, 0]
```

Rule 4145

```
Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.))
*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_), x_Symbol]
:> Simp[(A*b^2 - a*b*B + a^2*C)*Cot[e + f*x]*((a + b*Csc[e + f*x])^(m + 1)/(a*f*(m + 1)*(a^2 - b^2))), x]
+ Dist[1/(a*(m + 1)*(a^2 - b^2)), Int[(a + b*Csc[e + f*x])^(m + 1)*Simp[A*(a^2 - b^2)*(m + 1) - a*(A*b - a*B + b*C)*(m + 1)*Csc[e + f*x] + (A*b^2 - a*b*B + a^2*C)*(m + 2)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x]
&& NeQ[a^2 - b^2, 0] && LtQ[m, -1]
```

Rule 5576

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sech[(c_.) + (d_.)*(x_)])^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol]
:> Simp[(-(e + f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]
&& IGtQ[m, 0] && NeQ[n, -1]
```

Rule 6456

```
Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol]
:> Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x]
&& IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(b^3 \text{Subst}\left(\int \frac{x \operatorname{sech}(x) \tanh(x)}{(-a + \operatorname{sech}(x))^4} dx, x, \operatorname{sech}^{-1}(a + bx)\right)\right) \\ &= -\frac{\operatorname{sech}^{-1}(a + bx)}{3x^3} + \frac{1}{3}b^3 \text{Subst}\left(\int \frac{1}{(-a + \operatorname{sech}(x))^3} dx, x, \operatorname{sech}^{-1}(a + bx)\right) \end{aligned}$$

$$\begin{aligned}
&= \frac{b\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6a(1-a^2)x^2} - \frac{\operatorname{sech}^{-1}(a+bx)}{3x^3} \\
&\quad - \frac{b^3 \operatorname{Subst}\left(\int \frac{2(1-a^2)-2a\operatorname{sech}(x)-\operatorname{sech}^2(x)}{(-a+\operatorname{sech}(x))^2} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{6a(1-a^2)} \\
&= \frac{b\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6a(1-a^2)x^2} - \frac{(2-5a^2)b^2\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6a^2(1-a^2)^2x} - \frac{\operatorname{sech}^{-1}(a+bx)}{3x^3} \\
&\quad + \frac{b^3 \operatorname{Subst}\left(\int \frac{2(1-a^2)^2-a(1-4a^2)\operatorname{sech}(x)}{-a+\operatorname{sech}(x)} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{6a^2(1-a^2)^2} \\
&= \frac{b\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6a(1-a^2)x^2} - \frac{(2-5a^2)b^2\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6a^2(1-a^2)^2x} - \frac{b^3\operatorname{sech}^{-1}(a+bx)}{3a^3} \\
&\quad - \frac{\operatorname{sech}^{-1}(a+bx)}{3x^3} + \frac{((2-5a^2+6a^4)b^3) \operatorname{Subst}\left(\int \frac{\operatorname{sech}(x)}{-a+\operatorname{sech}(x)} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{6a^3(1-a^2)^2} \\
&= \frac{b\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6a(1-a^2)x^2} - \frac{(2-5a^2)b^2\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6a^2(1-a^2)^2x} - \frac{b^3\operatorname{sech}^{-1}(a+bx)}{3a^3} \\
&\quad - \frac{\operatorname{sech}^{-1}(a+bx)}{3x^3} + \frac{((2-5a^2+6a^4)b^3) \operatorname{Subst}\left(\int \frac{1}{1-a\cosh(x)} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{6a^3(1-a^2)^2} \\
&= \frac{b\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6a(1-a^2)x^2} - \frac{(2-5a^2)b^2\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6a^2(1-a^2)^2x} \\
&\quad - \frac{b^3\operatorname{sech}^{-1}(a+bx)}{3a^3} - \frac{\operatorname{sech}^{-1}(a+bx)}{3x^3} \\
&\quad + \frac{((2-5a^2+6a^4)b^3) \operatorname{Subst}\left(\int \frac{1}{1-a-(1+a)x^2} dx, x, \tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)\right)}{3a^3(1-a^2)^2} \\
&= \frac{b\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6a(1-a^2)x^2} - \frac{(2-5a^2)b^2\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)}{6a^2(1-a^2)^2x} - \frac{b^3\operatorname{sech}^{-1}(a+bx)}{3a^3} \\
&\quad - \frac{\operatorname{sech}^{-1}(a+bx)}{3x^3} + \frac{(2-5a^2+6a^4)b^3 \operatorname{arctanh}\left(\frac{\sqrt{1+a}\tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{3a^3(1-a^2)^{5/2}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 368, normalized size of antiderivative = 1.87

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^4} dx$$

$$= \frac{1}{6} \left(\frac{b \sqrt{-\frac{-1+a+bx}{1+a+bx}} (a - a^4 - abx - 2bx(1 + bx) + a^3(-1 + 4bx) + a^2(1 + 5bx + 5b^2x^2))}{(-1 + a)^2 a^2 (1 + a)^2 x^2} \right.$$

$$- \frac{2 \operatorname{sech}^{-1}(a + bx)}{x^3} - \frac{(2 - 5a^2 + 6a^4) b^3 \log(x)}{a^3 (1 - a^2)^{5/2}} + \frac{2b^3 \log(a + bx)}{a^3}$$

$$- \frac{2b^3 \log \left(1 + \sqrt{-\frac{-1+a+bx}{1+a+bx}} + a \sqrt{-\frac{-1+a+bx}{1+a+bx}} + bx \sqrt{-\frac{-1+a+bx}{1+a+bx}} \right)}{a^3}$$

$$+ \frac{(2 - 5a^2 + 6a^4) b^3 \log \left(1 - a^2 - abx + \sqrt{1 - a^2} \sqrt{-\frac{-1+a+bx}{1+a+bx}} + a \sqrt{1 - a^2} \sqrt{-\frac{-1+a+bx}{1+a+bx}} + \sqrt{1 - a^2} bx \sqrt{-\frac{-1+a+bx}{1+a+bx}} \right)}{a^3 (1 - a^2)^{5/2}}$$

`[In] Integrate[ArcSech[a + b*x]/x^4,x]`

```
[Out] ((b*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(a - a^4 - a*b*x - 2*b*x*(1 + b*x)
) + a^3*(-1 + 4*b*x) + a^2*(1 + 5*b*x + 5*b^2*x^2)))/((-1 + a)^2*a^2*(1 + a
)^2*x^2) - (2*ArcSech[a + b*x])/x^3 - ((2 - 5*a^2 + 6*a^4)*b^3*Log[x])/(a^3
*(1 - a^2)^(5/2)) + (2*b^3*Log[a + b*x])/a^3 - (2*b^3*Log[1 + Sqrt[-((-1 +
a + b*x)/(1 + a + b*x))]] + a*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))] + b*x*Sqr
rt[-((-1 + a + b*x)/(1 + a + b*x))])/a^3 + ((2 - 5*a^2 + 6*a^4)*b^3*Log[1
- a^2 - a*b*x + Sqrt[1 - a^2]*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))] + a*Sqr
t[1 - a^2]*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))] + Sqrt[1 - a^2]*b*x*Sqrt[-
((-1 + a + b*x)/(1 + a + b*x))])/a^3*(1 - a^2)^(5/2))/6
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.97 (sec) , antiderivative size = 608, normalized size of antiderivative = 3.09

method	result
parts	$-\frac{\operatorname{arcsech}(bx+a)}{3x^3} - \frac{b \sqrt{-\frac{bx+a-1}{bx+a}} (bx+a) \sqrt{\frac{bx+a+1}{bx+a}} \operatorname{csgn}(b)^2 \left(2 \operatorname{arctanh} \left(\frac{1}{\sqrt{-b^2x^2 - 2abx - a^2 + 1}} \right) a^6 b^2 x^2 + 6 \sqrt{-a^2 + 1} \ln \right)}{3x^3}$
derivativedivides	Expression too large to display
default	Expression too large to display

`[In] int(arcsech(b*x+a)/x^4,x,method=_RETURNVERBOSE)`

```
[Out] -1/3*arcsech(b*x+a)/x^3-1/6*b*(-(b*x+a-1)/(b*x+a))^(1/2)*(b*x+a)*((b*x+a+1)
/(b*x+a))^(1/2)*csgn(b)^2*(2*arctanh(1/(-b^2*x^2-2*a*b*x-a^2+1)^(1/2))*a^6*
b^2*x^2+6*(-a^2+1)^(1/2)*ln(2*(-a*b*x+(-a^2+1)^(1/2)*(-b^2*x^2-2*a*b*x-a^2+
1)^(1/2)-a^2+1)/x)*a^4*b^2*x^2-6*arctanh(1/(-b^2*x^2-2*a*b*x-a^2+1)^(1/2))*
a^4*b^2*x^2-5*(-a^2+1)^(1/2)*ln(2*(-a*b*x+(-a^2+1)^(1/2)*(-b^2*x^2-2*a*b*x-
a^2+1)^(1/2)-a^2+1)/x)*a^2*b^2*x^2-5*a^5*b*x*(-b^2*x^2-2*a*b*x-a^2+1)^(1/2)
+6*arctanh(1/(-b^2*x^2-2*a*b*x-a^2+1)^(1/2))*a^2*b^2*x^2+a^6*(-b^2*x^2-2*a*
b*x-a^2+1)^(1/2)+2*(-a^2+1)^(1/2)*ln(2*(-a*b*x+(-a^2+1)^(1/2)*(-b^2*x^2-2*a
*b*x-a^2+1)^(1/2)-a^2+1)/x)*b^2*x^2+7*(-b^2*x^2-2*a*b*x-a^2+1)^(1/2)*a^3*b*
x-2*arctanh(1/(-b^2*x^2-2*a*b*x-a^2+1)^(1/2))*b^2*x^2-2*(-b^2*x^2-2*a*b*x-a
^2+1)^(1/2)*a^4-2*(-b^2*x^2-2*a*b*x-a^2+1)^(1/2)*a*b*x+(-b^2*x^2-2*a*b*x-a
^2+1)^(1/2)*a^2)/x^2/(a^2-1)^2/(-1+a)/(1+a)/a^3/(-b^2*x^2-2*a*b*x-a^2+1)^(1/
2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 487 vs. 2(167) = 334.

Time = 0.33 (sec) , antiderivative size = 987, normalized size of antiderivative = 5.01

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{x^4} dx$$

$$= \frac{(6a^4 - 5a^2 + 2)\sqrt{-a^2 + 1}b^3x^3 \log\left(\frac{(2a^2-1)b^2x^2+2a^4+4(a^3-a)bx-4a^2-2(ab^2x^2+a^3+(2a^2-1)bx-a)\sqrt{-a^2+1}\sqrt{-\frac{b^2x^2}{b^2x^2+2abx+a^2}}}{x^2}\right)}{(6a^4 - 5a^2 + 2)\sqrt{a^2 - 1}b^3x^3 \arctan\left(\frac{(ab^2x^2+a^3+(2a^2-1)bx-a)\sqrt{a^2-1}\sqrt{-\frac{b^2x^2+2abx+a^2-1}{b^2x^2+2abx+a^2}}}{(a^2-1)b^2x^2+a^4+2(a^3-a)bx-2a^2+1}\right)} + (a^6 - 3a^4 + 3a^2)$$

```
[In] integrate(arcsech(b*x+a)/x^4,x, algorithm="fricas")
```

```
[Out] [-1/12*((6*a^4 - 5*a^2 + 2)*sqrt(-a^2 + 1)*b^3*x^3*log(((2*a^2 - 1)*b^2*x^2
+ 2*a^4 + 4*(a^3 - a)*b*x - 4*a^2 - 2*(a*b^2*x^2 + a^3 + (2*a^2 - 1)*b*x -
a)*sqrt(-a^2 + 1)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x +
a^2)) + 2)/x^2) + 2*(a^6 - 3*a^4 + 3*a^2 - 1)*b^3*x^3*log(((b*x + a)*sqrt(
-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/x) - 2*(a^6
- 3*a^4 + 3*a^2 - 1)*b^3*x^3*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2
- 1)/(b^2*x^2 + 2*a*b*x + a^2)) - 1)/x) + 4*(a^9 - 3*a^7 + 3*a^5 - a^3)*log
(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2))
```

```
+ 1)/(b*x + a)) - 2*((5*a^5 - 7*a^3 + 2*a)*b^3*x^3 + (4*a^6 - 5*a^4 + a^2)*
b^2*x^2 - (a^7 - 2*a^5 + a^3)*b*x)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2
*x^2 + 2*a*b*x + a^2)))/((a^9 - 3*a^7 + 3*a^5 - a^3)*x^3), -1/6*((6*a^4 - 5
*a^2 + 2)*sqrt(a^2 - 1)*b^3*x^3*arctan((a*b^2*x^2 + a^3 + (2*a^2 - 1)*b*x -
a)*sqrt(a^2 - 1)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x +
a^2)))/((a^2 - 1)*b^2*x^2 + a^4 + 2*(a^3 - a)*b*x - 2*a^2 + 1)) + (a^6 - 3*a
^4 + 3*a^2 - 1)*b^3*x^3*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/
(b^2*x^2 + 2*a*b*x + a^2)) + 1)/x) - (a^6 - 3*a^4 + 3*a^2 - 1)*b^3*x^3*log(
((b*x + a)*sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) -
1)/x) + 2*(a^9 - 3*a^7 + 3*a^5 - a^3)*log(((b*x + a)*sqrt(-(b^2*x^2 + 2*a*
b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)) - ((5*a^5 - 7*a^3
+ 2*a)*b^3*x^3 + (4*a^6 - 5*a^4 + a^2)*b^2*x^2 - (a^7 - 2*a^5 + a^3)*b*x)*
sqrt(-(b^2*x^2 + 2*a*b*x + a^2 - 1)/(b^2*x^2 + 2*a*b*x + a^2)))/((a^9 - 3*a
^7 + 3*a^5 - a^3)*x^3)]
```

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^4} dx = \int \frac{\operatorname{asech}(a + bx)}{x^4} dx$$

```
[In] integrate(asech(b*x+a)/x**4,x)
```

```
[Out] Integral(asech(a + b*x)/x**4, x)
```

Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^4} dx = \int \frac{\operatorname{arsech}(bx + a)}{x^4} dx$$

```
[In] integrate(arcsech(b*x+a)/x^4,x, algorithm="maxima")
```

```
[Out] 1/3*(6*a^4*b^3 - 3*a^2*b^3 + b^3)*log(x)/(a^9 - 3*a^7 + 3*a^5 - a^3) - 1/6*
((a^6*b^3 - 3*a^5*b^3 + 3*a^4*b^3 - a^3*b^3)*x^3*log(b*x + a + 1) + (a^6*b^
3 + 3*a^5*b^3 + 3*a^4*b^3 + a^3*b^3)*x^3*log(-b*x - a + 1) - 2*(3*a^5*b^2 -
4*a^3*b^2 + a*b^2)*x^2 + (a^6*b - 2*a^4*b + a^2*b)*x + 2*(a^9 - 3*a^7 + 3*
a^5 - a^3)*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)
*sqrt(-b*x - a + 1)*a + b*x + a) - 2*(a^9 - 3*a^7 + 3*a^5 + (a^6*b^3 - 3*a^
4*b^3 + 3*a^2*b^3 - b^3)*x^3 - a^3)*log(b*x + a) - 2*(a^9 - 3*a^7 + 3*a^5 -
a^3)*log(b*x + a))/((a^9 - 3*a^7 + 3*a^5 - a^3)*x^3) - integrate(1/3*(b^2*x
+ a*b)/(b^2*x^5 + 2*a*b*x^4 + (a^2 - 1)*x^3 + (b^2*x^5 + 2*a*b*x^4 + (a^2
- 1)*x^3)*e^(1/2*log(b*x + a + 1) + 1/2*log(-b*x - a + 1))), x)
```

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^4} dx = \int \frac{\operatorname{arosech}(bx + a)}{x^4} dx$$

[In] integrate(arcsech(b*x+a)/x^4,x, algorithm="giac")

[Out] integrate(arcsech(b*x + a)/x^4, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{x^4} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)}{x^4} dx$$

[In] int(acosh(1/(a + b*x))/x^4,x)

[Out] int(acosh(1/(a + b*x))/x^4, x)

3.9 $\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx$

Optimal result	104
Rubi [A] (verified)	105
Mathematica [A] (verified)	109
Maple [A] (verified)	110
Fricas [F]	110
Sympy [F]	111
Maxima [F]	111
Giac [F]	111
Mupad [F(-1)]	112

Optimal result

Integrand size = 12, antiderivative size = 279

$$\begin{aligned}
 \int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx = & -\frac{x}{3b^2} + \frac{2a\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)}{b^3} \\
 & - \frac{(a+bx)\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)}{3b^3} \\
 & + \frac{a^3 \operatorname{sech}^{-1}(a+bx)^2}{3b^3} + \frac{1}{3}x^3 \operatorname{sech}^{-1}(a+bx)^2 \\
 & - \frac{2 \operatorname{sech}^{-1}(a+bx) \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3} \\
 & - \frac{4a^2 \operatorname{sech}^{-1}(a+bx) \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3} \\
 & + \frac{2a \log(a+bx)}{b^3} + \frac{i \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3} \\
 & + \frac{2ia^2 \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3} \\
 & - \frac{i \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3} - \frac{2ia^2 \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3}
 \end{aligned}$$

[Out] $-1/3*x/b^2+1/3*a^3*\operatorname{arcsech}(b*x+a)^2/b^3+1/3*x^3*\operatorname{arcsech}(b*x+a)^2-2/3*\operatorname{arcsech}(b*x+a)*\arctan(1/(b*x+a)+(1/(b*x+a)-1)^{(1/2)}*(1/(b*x+a)+1)^{(1/2)})/b^3-4*a^2*\operatorname{arcsech}(b*x+a)*\arctan(1/(b*x+a)+(1/(b*x+a)-1)^{(1/2)}*(1/(b*x+a)+1)^{(1/2)})/b^3+2*a*\ln(b*x+a)/b^3+1/3*I*\operatorname{polylog}(2, -I*(1/(b*x+a)+(1/(b*x+a)-1)^{(1/2)}*(1/(b*x+a)+1)^{(1/2)}))/b^3+2*I*a^2*\operatorname{polylog}(2, -I*(1/(b*x+a)+(1/(b*x+a)-1)^{(1/2)}*(1/(b*x+a)+1)^{(1/2)}))/b^3-1/3*I*\operatorname{polylog}(2, I*(1/(b*x+a)+(1/(b*x+a)-1)^{(1/2)}*(1/(b*x+a)+1)^{(1/2)}))/b^3-2*I*a^2*\operatorname{polylog}(2, I*(1/(b*x+a)+(1/(b*x+a)-1)^{(1/2)}*(1/(b*x+a)+1)^{(1/2)}))/b^3$

$(1/(b*x+a)+1)^{(1/2)})/b^3-2*I*a^2*polylog(2,I*(1/(b*x+a)+(1/(b*x+a)-1)^{(1/2)})*(1/(b*x+a)+1)^{(1/2)})/b^3+2*a*(b*x+a+1)*arcsech(b*x+a)*((-b*x-a+1)/(b*x+a+1))^{(1/2)}/b^3-1/3*(b*x+a)*(b*x+a+1)*arcsech(b*x+a)*((-b*x-a+1)/(b*x+a+1))^{(1/2)}/b^3$

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6456, 5576, 4275, 4265, 2317, 2438, 4269, 3556, 4270}

$$\int x^2 \operatorname{sech}^{-1}(a+bx)^2 dx = \frac{a^3 \operatorname{sech}^{-1}(a+bx)^2}{3b^3} - \frac{4a^2 \operatorname{sech}^{-1}(a+bx) \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3} + \frac{2ia^2 \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3} - \frac{2ia^2 \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3} - \frac{2 \operatorname{sech}^{-1}(a+bx) \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3} + \frac{i \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3} - \frac{i \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3} + \frac{2a \log(a+bx)}{b^3} + \frac{2a \sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1) \operatorname{sech}^{-1}(a+bx)}{b^3} - \frac{(a+bx) \sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1) \operatorname{sech}^{-1}(a+bx)}{3b^3} + \frac{1}{3}x^3 \operatorname{sech}^{-1}(a+bx)^2 - \frac{x}{3b^2}$$

[In] Int[x^2*ArcSech[a + b*x]^2,x]

[Out] $-1/3*x/b^2 + (2*a*\sqrt{(1 - a - b*x)/(1 + a + b*x)}*(1 + a + b*x)*\operatorname{ArcSech}[a + b*x])/b^3 - ((a + b*x)*\sqrt{(1 - a - b*x)/(1 + a + b*x)}*(1 + a + b*x)*\operatorname{ArcSech}[a + b*x])/(3*b^3) + (a^3*\operatorname{ArcSech}[a + b*x]^2)/(3*b^3) + (x^3*\operatorname{ArcSech}[a + b*x]^2)/3 - (2*\operatorname{ArcSech}[a + b*x]*\operatorname{ArcTan}[E^{\operatorname{ArcSech}[a + b*x]}])/(3*b^3) - (4*a^2*\operatorname{ArcSech}[a + b*x]*\operatorname{ArcTan}[E^{\operatorname{ArcSech}[a + b*x]}])/b^3 + (2*a*\log[a + b*x])/b^3 + ((1/3)*\operatorname{PolyLog}[2, (-1)*E^{\operatorname{ArcSech}[a + b*x]}])/b^3 + ((2*I)*a^2*\operatorname{PolyLog}[2, (-1)*E^{\operatorname{ArcSech}[a + b*x]}])/b^3 - ((1/3)*\operatorname{PolyLog}[2, I*E^{\operatorname{ArcSech}[a + b*x]}])/b^3 - ((2*I)*a^2*\operatorname{PolyLog}[2, I*E^{\operatorname{ArcSech}[a + b*x]}])/b^3$

Rule 2317

Int[Log[(a_) + (b_.)*((F_)^(e_.)*((c_.) + (d_.)*(x_)))]^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))]

)^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 4265

Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_)^(m_.)), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Dist[d*(m/(f*fz*I)), Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Dist[d*(m/(f*fz*I)), Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]

Rule 4269

Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_)^(m_.)), x_Symbol] := Simp[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

Rule 4270

Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^n*((c_.) + (d_.)*(x_)), x_Symbol] := Simp[(-b^2)*(c + d*x)*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))), x] + (Dist[b^2*((n - 2)/(n - 1)), Int[(c + d*x)*(b*Csc[e + f*x])^(n - 2), x], x] - Simp[b^2*d*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2]

Rule 4275

Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^n*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]

Rule 5576

Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sech[(c_.) + (d_.)*(x_)])^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b

d(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /
; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rule 6456

Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int x^2 \text{sech}(x)(-a + \text{sech}(x))^2 \tanh(x) dx, x, \text{sech}^{-1}(a + bx)\right)}{b^3} \\
 &= \frac{1}{3}x^3 \text{sech}^{-1}(a + bx)^2 - \frac{2\text{Subst}\left(\int x(-a + \text{sech}(x))^3 dx, x, \text{sech}^{-1}(a + bx)\right)}{3b^3} \\
 &= \frac{1}{3}x^3 \text{sech}^{-1}(a + bx)^2 \\
 &\quad - \frac{2\text{Subst}\left(\int (-a^3x + 3a^2x\text{sech}(x) - 3ax\text{sech}^2(x) + x\text{sech}^3(x)) dx, x, \text{sech}^{-1}(a + bx)\right)}{3b^3} \\
 &= \frac{a^3 \text{sech}^{-1}(a + bx)^2}{3b^3} + \frac{1}{3}x^3 \text{sech}^{-1}(a + bx)^2 \\
 &\quad - \frac{2\text{Subst}\left(\int x\text{sech}^3(x) dx, x, \text{sech}^{-1}(a + bx)\right)}{3b^3} \\
 &\quad + \frac{(2a)\text{Subst}\left(\int x\text{sech}^2(x) dx, x, \text{sech}^{-1}(a + bx)\right)}{b^3} \\
 &\quad - \frac{(2a^2)\text{Subst}\left(\int x\text{sech}(x) dx, x, \text{sech}^{-1}(a + bx)\right)}{b^3} \\
 &= -\frac{x}{3b^2} + \frac{2a\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\text{sech}^{-1}(a+bx)}{b^3} \\
 &\quad - \frac{(a+bx)\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\text{sech}^{-1}(a+bx)}{3b^3} + \frac{a^3 \text{sech}^{-1}(a+bx)^2}{3b^3} \\
 &\quad + \frac{1}{3}x^3 \text{sech}^{-1}(a+bx)^2 - \frac{4a^2 \text{sech}^{-1}(a+bx) \arctan\left(e^{\text{sech}^{-1}(a+bx)}\right)}{b^3} \\
 &\quad - \frac{\text{Subst}\left(\int x\text{sech}(x) dx, x, \text{sech}^{-1}(a+bx)\right)}{3b^3} \\
 &\quad - \frac{(2a)\text{Subst}\left(\int \tanh(x) dx, x, \text{sech}^{-1}(a+bx)\right)}{b^3} \\
 &\quad + \frac{(2ia^2)\text{Subst}\left(\int \log(1-ie^x) dx, x, \text{sech}^{-1}(a+bx)\right)}{b^3} \\
 &\quad - \frac{(2ia^2)\text{Subst}\left(\int \log(1+ie^x) dx, x, \text{sech}^{-1}(a+bx)\right)}{b^3}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{x}{3b^2} + \frac{2a\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)}{b^3} \\
&\quad - \frac{(a+bx)\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)}{3b^3} + \frac{a^3\operatorname{sech}^{-1}(a+bx)^2}{3b^3} \\
&\quad + \frac{1}{3}x^3\operatorname{sech}^{-1}(a+bx)^2 - \frac{2\operatorname{sech}^{-1}(a+bx)\arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3} \\
&\quad - \frac{4a^2\operatorname{sech}^{-1}(a+bx)\arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3} + \frac{2a\log(a+bx)}{b^3} \\
&\quad + \frac{i\operatorname{Subst}\left(\int \log(1-ie^x) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{3b^3} \\
&\quad - \frac{i\operatorname{Subst}\left(\int \log(1+ie^x) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{3b^3} \\
&\quad + \frac{(2ia^2)\operatorname{Subst}\left(\int \frac{\log(1-ix)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3} \\
&\quad - \frac{(2ia^2)\operatorname{Subst}\left(\int \frac{\log(1+ix)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3} \\
&= -\frac{x}{3b^2} + \frac{2a\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)}{b^3} \\
&\quad - \frac{(a+bx)\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)}{3b^3} + \frac{a^3\operatorname{sech}^{-1}(a+bx)^2}{3b^3} \\
&\quad + \frac{1}{3}x^3\operatorname{sech}^{-1}(a+bx)^2 - \frac{2\operatorname{sech}^{-1}(a+bx)\arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3} \\
&\quad - \frac{4a^2\operatorname{sech}^{-1}(a+bx)\arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3} + \frac{2a\log(a+bx)}{b^3} \\
&\quad + \frac{2ia^2\operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3} - \frac{2ia^2\operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3} \\
&\quad + \frac{i\operatorname{Subst}\left(\int \frac{\log(1-ix)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3} - \frac{i\operatorname{Subst}\left(\int \frac{\log(1+ix)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{x}{3b^2} + \frac{2a\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)}{b^3} \\
&\quad - \frac{(a+bx)\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)}{3b^3} + \frac{a^3\operatorname{sech}^{-1}(a+bx)^2}{3b^3} \\
&\quad + \frac{1}{3}x^3\operatorname{sech}^{-1}(a+bx)^2 - \frac{2\operatorname{sech}^{-1}(a+bx)\arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3} \\
&\quad - \frac{4a^2\operatorname{sech}^{-1}(a+bx)\arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3} + \frac{2a\log(a+bx)}{b^3} \\
&\quad + \frac{i\operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3} + \frac{2ia^2\operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3} \\
&\quad - \frac{i\operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{3b^3} - \frac{2ia^2\operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 2.28 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.09

$$\int x^2\operatorname{sech}^{-1}(a+bx)^2 dx = \frac{2(a+bx)\sqrt{-\frac{-1+a+bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx) + 6a(a+bx)^2\operatorname{sech}^{-1}(a+bx)^2 - 2(a+bx)^3\operatorname{sech}^{-1}(a+bx)^2}{b^3}$$

[In] Integrate[x^2*ArcSech[a + b*x]^2,x]

[Out]
$$\begin{aligned}
&-1/6*(2*(a + b*x)*\operatorname{Sqrt}[-((-1 + a + b*x)/(1 + a + b*x))]*(1 + a + b*x)*\operatorname{ArcSech}[a + b*x] \\
&+ 6*a*(a + b*x)^2*\operatorname{ArcSech}[a + b*x]^2 - 2*(a + b*x)^3*\operatorname{ArcSech}[a + b*x]^2 \\
&+ 2*(a + b*x - 6*a*\operatorname{Sqrt}[-((-1 + a + b*x)/(1 + a + b*x))]*(1 + a + b*x)*\operatorname{ArcSech}[a + b*x] \\
&- 3*a^2*(a + b*x)*\operatorname{ArcSech}[a + b*x]^2) + 12*a*\operatorname{Log}[(a + b*x)^{-1}] - (1 + 6*a^2)*(Pi*\operatorname{Log}[1 - I*E^{\operatorname{ArcSech}[a + b*x]}] - (2*I)*\operatorname{ArcSech}[a + b*x]*\operatorname{Log}[1 - I*E^{\operatorname{ArcSech}[a + b*x]}] - Pi*\operatorname{Log}[1 + I*E^{\operatorname{ArcSech}[a + b*x]}] \\
&+ (2*I)*\operatorname{ArcSech}[a + b*x]*\operatorname{Log}[1 + I*E^{\operatorname{ArcSech}[a + b*x]}] - Pi*\operatorname{Log}[\operatorname{Cot}[(Pi + (2*I)*\operatorname{ArcSech}[a + b*x])/4]]) + (2*I)*\operatorname{PolyLog}[2, (-I)*E^{\operatorname{ArcSech}[a + b*x]}] - (2*I)*\operatorname{PolyLog}[2, I*E^{\operatorname{ArcSech}[a + b*x]}]))/b^3
\end{aligned}$$

Maple [A] (verified)

Time = 0.87 (sec) , antiderivative size = 599, normalized size of antiderivative = 2.15

method	result
derivativedivides	$\frac{\operatorname{arcsech}(bx+a)^2 a^2 (bx+a) - \operatorname{arcsech}(bx+a)^2 a (bx+a)^2 + \frac{\operatorname{arcsech}(bx+a)^2 (bx+a)^3}{3} + 2 \operatorname{arcsech}(bx+a) \sqrt{-\frac{bx+a-1}{bx+a}} \sqrt{\frac{bx+a+1}{bx+a}} a}{\dots}$
default	$\operatorname{arcsech}(bx+a)^2 a^2 (bx+a) - \operatorname{arcsech}(bx+a)^2 a (bx+a)^2 + \frac{\operatorname{arcsech}(bx+a)^2 (bx+a)^3}{3} + 2 \operatorname{arcsech}(bx+a) \sqrt{-\frac{bx+a-1}{bx+a}} \sqrt{\frac{bx+a+1}{bx+a}} a$

[In] int(x^2*arcsech(b*x+a)^2,x,method=_RETURNVERBOSE)

```
[Out] 1/b^3*(arcsech(b*x+a)^2*a^2*(b*x+a)-arcsech(b*x+a)^2*a*(b*x+a)^2+1/3*arcsech(b*x+a)^2*(b*x+a)^3+2*arcsech(b*x+a)*(-(b*x+a-1)/(b*x+a))^(1/2)*((b*x+a+1)/(b*x+a))^(1/2)*a*(b*x+a)-1/3*arcsech(b*x+a)*(-(b*x+a-1)/(b*x+a))^(1/2)*((b*x+a+1)/(b*x+a))^(1/2)*(b*x+a)^2-2*a*arcsech(b*x+a)-1/3*b*x-1/3*a+1/3*I*arcsech(b*x+a)*ln(1+I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))))-1/3*I*dilog(1-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))))-2*I*a^2*dilog(1-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))))+2*I*a^2*dilog(1+I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))))-2*ln(1+(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))^2)*a+4*a*ln(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))-1/3*I*arcsech(b*x+a)*ln(1-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))))-2*I*a^2*arcsech(b*x+a)*ln(1-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))))+2*I*a^2*arcsech(b*x+a)*ln(1+I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))))+1/3*I*dilog(1+I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))))
```

Fricas [F]

$$\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx = \int x^2 \operatorname{ar} \operatorname{sech}(bx + a)^2 dx$$

[In] integrate(x^2*arcsech(b*x+a)^2,x, algorithm="fricas")

[Out] integral(x^2*arcsech(b*x + a)^2, x)

Sympy [F]

$$\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx = \int x^2 \operatorname{asech}^2(a + bx) dx$$

```
[In] integrate(x**2*asech(b*x+a)**2,x)
```

```
[Out] Integral(x**2*asech(a + b*x)**2, x)
```

Maxima [F]

$$\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx = \int x^2 \operatorname{arsech}(bx + a)^2 dx$$

```
[In] integrate(x^2*arcsech(b*x+a)^2,x, algorithm="maxima")
```

```
[Out] 1/3*x^3*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^2 - integrate(-2/3*(6*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^2 + 6*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2)*log(b*x + a)^2 - (b^3*x^5 + 2*a*b^2*x^4 + (a^2*b - b)*x^3 + 6*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2)*log(b*x + a) + (3*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2)*sqrt(b*x + a + 1)*log(b*x + a) + (2*b^3*x^5 + 4*a*b^2*x^4 + (2*a^2*b - b)*x^3 + 3*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2)*log(b*x + a))*sqrt(b*x + a + 1))*sqrt(-b*x - a + 1))*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1) + (3*a^2*b - b)*x - a), x)
```

Giac [F]

$$\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx = \int x^2 \operatorname{arsech}(bx + a)^2 dx$$

```
[In] integrate(x^2*arcsech(b*x+a)^2,x, algorithm="giac")
```

```
[Out] integrate(x^2*arcsech(b*x + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{sech}^{-1}(a + bx)^2 dx = \int x^2 \operatorname{acosh}\left(\frac{1}{a + bx}\right)^2 dx$$

```
[In] int(x^2*acosh(1/(a + b*x))^2,x)
```

```
[Out] int(x^2*acosh(1/(a + b*x))^2, x)
```


3.10 $\int x \operatorname{sech}^{-1}(a + bx)^2 dx$

Optimal result	113
Rubi [A] (verified)	113
Mathematica [A] (verified)	116
Maple [A] (verified)	117
Fricas [F]	117
Sympy [F]	118
Maxima [F]	118
Giac [F]	118
Mupad [F(-1)]	119

Optimal result

Integrand size = 10, antiderivative size = 149

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx = -\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)}{b^2} - \frac{a^2 \operatorname{sech}^{-1}(a+bx)^2}{2b^2}$$

$$+ \frac{1}{2}x^2 \operatorname{sech}^{-1}(a+bx)^2 + \frac{4a \operatorname{sech}^{-1}(a+bx) \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2}$$

$$- \frac{\log(a+bx)}{b^2} - \frac{2ia \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2}$$

$$+ \frac{2ia \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2}$$

```
[Out] -1/2*a^2*arcsech(b*x+a)^2/b^2+1/2*x^2*arcsech(b*x+a)^2+4*a*arcsech(b*x+a)*a
rctan(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/b^2-ln(b*x+a)/b^2-
2*I*a*polylog(2,-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b^2
+2*I*a*polylog(2,I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b^2
-(b*x+a+1)*arcsech(b*x+a)*((-b*x-a+1)/(b*x+a+1))^(1/2)/b^2
```

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used

= {6456, 5576, 4275, 4265, 2317, 2438, 4269, 3556}

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx = -\frac{a^2 \operatorname{sech}^{-1}(a + bx)^2}{2b^2} + \frac{4a \operatorname{sech}^{-1}(a + bx) \arctan\left(e^{\operatorname{sech}^{-1}(a + bx)}\right)}{b^2}$$

$$- \frac{2ia \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b^2}$$

$$+ \frac{2ia \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b^2} - \frac{\log(a + bx)}{b^2}$$

$$- \frac{\sqrt{\frac{-a - bx + 1}{a + bx + 1}}(a + bx + 1) \operatorname{sech}^{-1}(a + bx)}{b^2} + \frac{1}{2}x^2 \operatorname{sech}^{-1}(a + bx)^2$$

[In] Int[x*ArcSech[a + b*x]^2,x]

[Out] -((Sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x)*ArcSech[a + b*x])/b^2) - (a^2*ArcSech[a + b*x]^2)/(2*b^2) + (x^2*ArcSech[a + b*x]^2)/2 + (4*a*ArcSech[a + b*x]*ArcTan[E^ArcSech[a + b*x]])/b^2 - Log[a + b*x]/b^2 - ((2*I)*a*PolyLog[2, (-I)*E^ArcSech[a + b*x]])/b^2 + ((2*I)*a*PolyLog[2, I*E^ArcSech[a + b*x]])/b^2

Rule 2317

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3556

Int[tan[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 4265

Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Dist[d*(m/(f*fz*I)), Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Dist[d*(m/(f*fz*I)), Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]

Rule 4269

```
Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp
[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*
Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 4275

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)
, x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 5576

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sech[
(c_.) + (d_.)*(x_)])^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-(e
+ f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b
*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /
; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 6456

```
Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Ta
nh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int x^2 \text{sech}(x)(-a + \text{sech}(x)) \tanh(x) dx, x, \text{sech}^{-1}(a + bx)\right)}{b^2} \\
&= \frac{1}{2}x^2 \text{sech}^{-1}(a + bx)^2 - \frac{\text{Subst}\left(\int x(-a + \text{sech}(x))^2 dx, x, \text{sech}^{-1}(a + bx)\right)}{b^2} \\
&= \frac{1}{2}x^2 \text{sech}^{-1}(a + bx)^2 - \frac{\text{Subst}\left(\int (a^2x - 2ax \text{sech}(x) + x \text{sech}^2(x)) dx, x, \text{sech}^{-1}(a + bx)\right)}{b^2} \\
&= -\frac{a^2 \text{sech}^{-1}(a + bx)^2}{2b^2} + \frac{1}{2}x^2 \text{sech}^{-1}(a + bx)^2 \\
&\quad - \frac{\text{Subst}\left(\int x \text{sech}^2(x) dx, x, \text{sech}^{-1}(a + bx)\right)}{b^2} \\
&\quad + \frac{(2a)\text{Subst}\left(\int x \text{sech}(x) dx, x, \text{sech}^{-1}(a + bx)\right)}{b^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)}{b^2} - \frac{a^2\operatorname{sech}^{-1}(a+bx)^2}{2b^2} \\
&\quad + \frac{1}{2}x^2\operatorname{sech}^{-1}(a+bx)^2 + \frac{4a\operatorname{sech}^{-1}(a+bx)\arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad + \frac{\operatorname{Subst}\left(\int \tanh(x) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{b^2} \\
&\quad - \frac{(2ia)\operatorname{Subst}\left(\int \log(1-ie^x) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{b^2} \\
&\quad + \frac{(2ia)\operatorname{Subst}\left(\int \log(1+ie^x) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{b^2} \\
&= -\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)}{b^2} - \frac{a^2\operatorname{sech}^{-1}(a+bx)^2}{2b^2} \\
&\quad + \frac{1}{2}x^2\operatorname{sech}^{-1}(a+bx)^2 + \frac{4a\operatorname{sech}^{-1}(a+bx)\arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad - \frac{\log(a+bx)}{b^2} - \frac{(2ia)\operatorname{Subst}\left(\int \frac{\log(1-ix)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad + \frac{(2ia)\operatorname{Subst}\left(\int \frac{\log(1+ix)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&= -\frac{\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)}{b^2} - \frac{a^2\operatorname{sech}^{-1}(a+bx)^2}{2b^2} \\
&\quad + \frac{1}{2}x^2\operatorname{sech}^{-1}(a+bx)^2 + \frac{4a\operatorname{sech}^{-1}(a+bx)\arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} - \frac{\log(a+bx)}{b^2} \\
&\quad - \frac{2ia\operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} + \frac{2ia\operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.52 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.15

$$\begin{aligned}
&\int x\operatorname{sech}^{-1}(a+bx)^2 dx \\
&= \frac{-2\sqrt{-\frac{-1+a+bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx) - 2a(a+bx)\operatorname{sech}^{-1}(a+bx)^2 + (a+bx)^2\operatorname{sech}^{-1}(a+bx)^2 - 4ia}{b^2}
\end{aligned}$$

[In] Integrate[x*ArcSech[a + b*x]^2,x]

[Out] (-2*Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(1 + a + b*x)*ArcSech[a + b*x] - 2*a*(a + b*x)*ArcSech[a + b*x]^2 + (a + b*x)^2*ArcSech[a + b*x]^2 - (4*I)*a

`*ArcSech[a + b*x]*(Log[1 - I/E^ArcSech[a + b*x]] - Log[1 + I/E^ArcSech[a + b*x]]) + 2*Log[(a + b*x)^(-1)] - (4*I)*a*(PolyLog[2, (-I)/E^ArcSech[a + b*x]] - PolyLog[2, I/E^ArcSech[a + b*x]])/(2*b^2)`

Maple [A] (verified)

Time = 0.78 (sec) , antiderivative size = 331, normalized size of antiderivative = 2.22

method	result
derivativedivides	$-\frac{\operatorname{arcsech}(bx+a)\left(2\operatorname{arcsech}(bx+a)a(bx+a)-\operatorname{arcsech}(bx+a)(bx+a)^2+2\sqrt{-\frac{bx+a-1}{bx+a}}\sqrt{\frac{bx+a+1}{bx+a}}(bx+a)-2\right)}{2}-2\ln\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}}-\sqrt{\frac{1}{bx+a}}\right)$
default	$-\frac{\operatorname{arcsech}(bx+a)\left(2\operatorname{arcsech}(bx+a)a(bx+a)-\operatorname{arcsech}(bx+a)(bx+a)^2+2\sqrt{-\frac{bx+a-1}{bx+a}}\sqrt{\frac{bx+a+1}{bx+a}}(bx+a)-2\right)}{2}-2\ln\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}}-\sqrt{\frac{1}{bx+a}}\right)$

[In] `int(x*arcsech(b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{b^2}\left(-\frac{1}{2}\operatorname{arcsech}(bx+a)\left(2\operatorname{arcsech}(bx+a)a(bx+a)-\operatorname{arcsech}(bx+a)(bx+a)^2+2\sqrt{-\frac{bx+a-1}{bx+a}}\sqrt{\frac{bx+a+1}{bx+a}}(bx+a)-2\right)+2\ln\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}}-\sqrt{\frac{1}{bx+a}}\right)\right)+\frac{1}{b^2}\left(-\frac{1}{2}\operatorname{arcsech}(bx+a)\left(2\operatorname{arcsech}(bx+a)a(bx+a)-\operatorname{arcsech}(bx+a)(bx+a)^2+2\sqrt{-\frac{bx+a-1}{bx+a}}\sqrt{\frac{bx+a+1}{bx+a}}(bx+a)-2\right)+2\ln\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}}-\sqrt{\frac{1}{bx+a}}\right)\right)+2Ia\operatorname{arcsech}(bx+a)\ln\left(1+I\sqrt{\frac{1}{bx+a}}+I\sqrt{\frac{1}{bx+a}}\right)+2Ia\operatorname{arcsech}(bx+a)\ln\left(1-I\sqrt{\frac{1}{bx+a}}-I\sqrt{\frac{1}{bx+a}}\right)+2Ia\operatorname{dilog}\left(1+I\sqrt{\frac{1}{bx+a}}+I\sqrt{\frac{1}{bx+a}}\right)+2Ia\operatorname{dilog}\left(1-I\sqrt{\frac{1}{bx+a}}-I\sqrt{\frac{1}{bx+a}}\right)\right)$$

Fricas [F]

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx = \int x \operatorname{ar} \operatorname{sech}(bx + a)^2 dx$$

[In] `integrate(x*arcsech(b*x+a)^2,x, algorithm="fricas")`

[Out] `integral(x*arcsech(b*x + a)^2, x)`

Sympy [F]

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx = \int x \operatorname{asech}^2(a + bx) dx$$

```
[In] integrate(x*asech(b*x+a)**2,x)
```

```
[Out] Integral(x*asech(a + b*x)**2, x)
```

Maxima [F]

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx = \int x \operatorname{arsech}(bx + a)^2 dx$$

```
[In] integrate(x*arcsech(b*x+a)^2,x, algorithm="maxima")
```

```
[Out] 1/2*x^2*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^2 - integrate(-(4*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^2 + 4*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*log(b*x + a)^2 - (b^3*x^4 + 2*a*b^2*x^3 + (a^2*b - b)*x^2 + 4*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*log(b*x + a) + (2*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*sqrt(b*x + a + 1)*log(b*x + a) + (2*b^3*x^4 + 4*a*b^2*x^3 + (2*a^2*b - b)*x^2 + 2*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*log(b*x + a))*sqrt(b*x + a + 1))*sqrt(-b*x - a + 1))*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1) + (3*a^2*b - b)*x - a), x)
```

Giac [F]

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx = \int x \operatorname{arsech}(bx + a)^2 dx$$

```
[In] integrate(x*arcsech(b*x+a)^2,x, algorithm="giac")
```

```
[Out] integrate(x*arcsech(b*x + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{sech}^{-1}(a + bx)^2 dx = \int x \operatorname{acosh}\left(\frac{1}{a + bx}\right)^2 dx$$

```
[In] int(x*acosh(1/(a + b*x))^2,x)
```

```
[Out] int(x*acosh(1/(a + b*x))^2, x)
```

3.11 $\int \operatorname{sech}^{-1}(a + bx)^2 dx$

Optimal result	120
Rubi [A] (verified)	120
Mathematica [A] (verified)	122
Maple [A] (verified)	123
Fricas [F]	123
Sympy [F]	123
Maxima [F]	124
Giac [F]	124
Mupad [F(-1)]	124

Optimal result

Integrand size = 8, antiderivative size = 80

$$\int \operatorname{sech}^{-1}(a + bx)^2 dx = \frac{(a + bx)\operatorname{sech}^{-1}(a + bx)^2}{b} - \frac{4\operatorname{sech}^{-1}(a + bx) \arctan\left(e^{\operatorname{sech}^{-1}(a + bx)}\right)}{b} + \frac{2i \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b} - \frac{2i \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b}$$

[Out] (b*x+a)*arcsech(b*x+a)^2/b-4*arcsech(b*x+a)*arctan(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/b+2*I*polylog(2,-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b-2*I*polylog(2,I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6450, 6414, 5526, 4265, 2317, 2438}

$$\int \operatorname{sech}^{-1}(a + bx)^2 dx = -\frac{4\operatorname{sech}^{-1}(a + bx) \arctan\left(e^{\operatorname{sech}^{-1}(a + bx)}\right)}{b} + \frac{2i \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b} - \frac{2i \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b} + \frac{(a + bx)\operatorname{sech}^{-1}(a + bx)^2}{b}$$

[In] Int[ArcSech[a + b*x]^2,x]

[Out] $((a + b*x)*\text{ArcSech}[a + b*x]^2)/b - (4*\text{ArcSech}[a + b*x]*\text{ArcTan}[E^{\text{ArcSech}[a + b*x]}])/b + ((2*I)*\text{PolyLog}[2, (-I)*E^{\text{ArcSech}[a + b*x]}])/b - ((2*I)*\text{PolyLog}[2, I*E^{\text{ArcSech}[a + b*x]}])/b$

Rule 2317

$\text{Int}[\text{Log}[(a_) + (b_)*(F_)^{((e_)*((c_) + (d_)*(x_)))^{(n_)}], x_Symbol] \rightarrow \text{Dist}[1/(d*e*n*\text{Log}[F]), \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^{(e*(c + d*x)})^n], x] \text{ ; FreeQ}\{F, a, b, c, d, e, n\}, x] \ \&\& \ \text{GtQ}[a, 0]$

Rule 2438

$\text{Int}[\text{Log}[(c_)*((d_) + (e_)*(x_)^{(n_)})]/(x_), x_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n]/n, x] \text{ ; FreeQ}\{c, d, e, n\}, x] \ \&\& \ \text{EqQ}[c*d, 1]$

Rule 4265

$\text{Int}[\text{csc}[(e_) + \text{Pi}*(k_) + (\text{Complex}[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^{(m_)}, x_Symbol] \rightarrow \text{Simp}[-2*(c + d*x)^m*(\text{ArcTanh}[E^{((-I)*e + f*fz*x)/E^{(I*k*\text{Pi})}}]/(f*fz*I)), x] + (-\text{Dist}[d*(m/(f*fz*I)), \text{Int}[(c + d*x)^{m-1}*\text{Log}[1 - E^{((-I)*e + f*fz*x)/E^{(I*k*\text{Pi})}}], x], x] + \text{Dist}[d*(m/(f*fz*I)), \text{Int}[(c + d*x)^{m-1}*\text{Log}[1 + E^{((-I)*e + f*fz*x)/E^{(I*k*\text{Pi})}}], x], x]) \text{ ; FreeQ}\{c, d, e, f, fz\}, x] \ \&\& \ \text{IntegerQ}[2*k] \ \&\& \ \text{IGtQ}[m, 0]$

Rule 5526

$\text{Int}[(x_)^{(m_)}*\text{Sech}[(a_) + (b_)*(x_)^{(n_)}]^{(p_)}*\text{Tanh}[(a_) + (b_)*(x_)^{(n_)}]^{(q_)}, x_Symbol] \rightarrow \text{Simp}[-(x^{(m-n+1)})*(\text{Sech}[a + b*x^n]^p/(b^n*p)), x] + \text{Dist}[(m-n+1)/(b^n*p), \text{Int}[x^{(m-n)}*\text{Sech}[a + b*x^n]^p, x], x] \text{ ; FreeQ}\{a, b, p\}, x] \ \&\& \ \text{RationalQ}[m] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ \text{GeQ}[m-n, 0] \ \&\& \ \text{EqQ}[q, 1]$

Rule 6414

$\text{Int}[(a_) + \text{ArcSech}[(c_)*(x_)]*(b_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[-c^{(-1)}, \text{Subst}[\text{Int}[(a + b*x)^n*\text{Sech}[x]*\text{Tanh}[x], x], x, \text{ArcSech}[c*x]], x] \text{ ; FreeQ}\{a, b, c, n\}, x] \ \&\& \ \text{IGtQ}[n, 0]$

Rule 6450

$\text{Int}[(a_) + \text{ArcSech}[(c_) + (d_)*(x_)]*(b_)]^{(p_)}, x_Symbol] \rightarrow \text{Dist}[1/d, \text{Subst}[\text{Int}[(a + b*\text{ArcSech}[x])^p, x], x, c + d*x], x] \text{ ; FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{IGtQ}[p, 0]$

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Subst}\left(\int \text{sech}^{-1}(x)^2 dx, x, a + bx\right)}{b} \\
&= -\frac{\text{Subst}\left(\int x^2 \text{sech}(x) \tanh(x) dx, x, \text{sech}^{-1}(a + bx)\right)}{b} \\
&= \frac{(a + bx)\text{sech}^{-1}(a + bx)^2}{b} - \frac{2\text{Subst}\left(\int x \text{sech}(x) dx, x, \text{sech}^{-1}(a + bx)\right)}{b} \\
&= \frac{(a + bx)\text{sech}^{-1}(a + bx)^2}{b} - \frac{4\text{sech}^{-1}(a + bx) \arctan\left(e^{\text{sech}^{-1}(a + bx)}\right)}{b} \\
&\quad + \frac{(2i)\text{Subst}\left(\int \log(1 - ie^x) dx, x, \text{sech}^{-1}(a + bx)\right)}{b} \\
&\quad - \frac{(2i)\text{Subst}\left(\int \log(1 + ie^x) dx, x, \text{sech}^{-1}(a + bx)\right)}{b} \\
&= \frac{(a + bx)\text{sech}^{-1}(a + bx)^2}{b} - \frac{4\text{sech}^{-1}(a + bx) \arctan\left(e^{\text{sech}^{-1}(a + bx)}\right)}{b} \\
&\quad + \frac{(2i)\text{Subst}\left(\int \frac{\log(1 - ix)}{x} dx, x, e^{\text{sech}^{-1}(a + bx)}\right)}{b} \\
&\quad - \frac{(2i)\text{Subst}\left(\int \frac{\log(1 + ix)}{x} dx, x, e^{\text{sech}^{-1}(a + bx)}\right)}{b} \\
&= \frac{(a + bx)\text{sech}^{-1}(a + bx)^2}{b} - \frac{4\text{sech}^{-1}(a + bx) \arctan\left(e^{\text{sech}^{-1}(a + bx)}\right)}{b} \\
&\quad + \frac{2i \text{PolyLog}\left(2, -ie^{\text{sech}^{-1}(a + bx)}\right)}{b} - \frac{2i \text{PolyLog}\left(2, ie^{\text{sech}^{-1}(a + bx)}\right)}{b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.31

$$\begin{aligned}
&\int \text{sech}^{-1}(a + bx)^2 dx \\
&= \frac{i\left(\text{sech}^{-1}(a + bx)\left(-i(a + bx)\text{sech}^{-1}(a + bx) + 2 \log\left(1 - ie^{-\text{sech}^{-1}(a + bx)}\right) - 2 \log\left(1 + ie^{-\text{sech}^{-1}(a + bx)}\right)\right)\right)}{b} +
\end{aligned}$$

[In] Integrate[ArcSech[a + b*x]^2,x]

[Out] (I*(ArcSech[a + b*x]*((-I)*(a + b*x)*ArcSech[a + b*x] + 2*Log[1 - I/E^ArcSech[a + b*x]] - 2*Log[1 + I/E^ArcSech[a + b*x]])) + 2*PolyLog[2, (-I)/E^ArcSech[a + b*x]] - 2*PolyLog[2, I/E^ArcSech[a + b*x]]))/b

Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 192, normalized size of antiderivative = 2.40

method	result
derivativedivides	$\frac{\operatorname{arcsech}(bx+a)^2(bx+a)+2i \operatorname{arcsech}(bx+a) \ln\left(1+i\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)-2i \operatorname{arcsech}(bx+a) \ln\left(1-i\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)}{\operatorname{arcsech}(bx+a)^2(bx+a)+2i \operatorname{arcsech}(bx+a) \ln\left(1+i\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)-2i \operatorname{arcsech}(bx+a) \ln\left(1-i\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)}$
default	$\frac{\operatorname{arcsech}(bx+a)^2(bx+a)+2i \operatorname{arcsech}(bx+a) \ln\left(1+i\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)-2i \operatorname{arcsech}(bx+a) \ln\left(1-i\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)}{\operatorname{arcsech}(bx+a)^2(bx+a)+2i \operatorname{arcsech}(bx+a) \ln\left(1+i\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)-2i \operatorname{arcsech}(bx+a) \ln\left(1-i\left(\frac{1}{bx+a}+\sqrt{\frac{1}{bx+a}-1}\sqrt{\frac{1}{bx+a}+1}\right)\right)}$

```
[In] int(arcsech(b*x+a)^2,x,method=_RETURNVERBOSE)
```

```
[Out] 1/b*(arcsech(b*x+a)^2*(b*x+a)+2*I*arcsech(b*x+a)*ln(1+I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))-2*I*arcsech(b*x+a)*ln(1-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))+2*I*dilog(1+I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))-2*I*dilog(1-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))))
```

Fricas [F]

$$\int \operatorname{sech}^{-1}(a + bx)^2 dx = \int \operatorname{arsech}(bx + a)^2 dx$$

```
[In] integrate(arcsech(b*x+a)^2,x, algorithm="fricas")
```

```
[Out] integral(arcsech(b*x + a)^2, x)
```

Sympy [F]

$$\int \operatorname{sech}^{-1}(a + bx)^2 dx = \int \operatorname{asech}^2(a + bx) dx$$

```
[In] integrate(asech(b*x+a)**2,x)
```

```
[Out] Integral(asech(a + b*x)**2, x)
```

Maxima [F]

$$\int \operatorname{sech}^{-1}(a + bx)^2 dx = \int \operatorname{arsech}(bx + a)^2 dx$$

[In] integrate(arcsech(b*x+a)^2,x, algorithm="maxima")

[Out] x*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^2 - integrate(-2*(2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^2 + 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a)^2 - (b^3*x^3 + 2*a*b^2*x^2 + (a^2*b - b)*x + 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a) + ((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*log(b*x + a) + (2*b^3*x^3 + 4*a*b^2*x^2 + (2*a^2*b - b)*x + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a))*sqrt(b*x + a + 1))*sqrt(-b*x - a + 1))*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1) + (3*a^2*b - b)*x - a), x)

Giac [F]

$$\int \operatorname{sech}^{-1}(a + bx)^2 dx = \int \operatorname{arsech}(bx + a)^2 dx$$

[In] integrate(arcsech(b*x+a)^2,x, algorithm="giac")

[Out] integrate(arcsech(b*x + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \operatorname{sech}^{-1}(a + bx)^2 dx = \int \operatorname{acosh}\left(\frac{1}{a + bx}\right)^2 dx$$

[In] int(acosh(1/(a + b*x))^2,x)

[Out] int(acosh(1/(a + b*x))^2, x)

3.12 $\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} dx$

Optimal result	125
Rubi [A] (verified)	126
Mathematica [A] (verified)	131
Maple [F]	132
Fricas [F]	132
Sympy [F]	132
Maxima [F]	132
Giac [F]	133
Mupad [F(-1)]	133

Optimal result

Integrand size = 12, antiderivative size = 274

$$\begin{aligned}
 \int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} dx &= \operatorname{sech}^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
 &\quad + \operatorname{sech}^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
 &\quad - \operatorname{sech}^{-1}(a+bx)^2 \log \left(1 + e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
 &\quad + 2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
 &\quad + 2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
 &\quad - \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left(2, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
 &\quad - 2 \operatorname{PolyLog} \left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) - 2 \operatorname{PolyLog} \left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
 &\quad + \frac{1}{2} \operatorname{PolyLog} \left(3, -e^{2\operatorname{sech}^{-1}(a+bx)} \right)
 \end{aligned}$$

```

[Out] -arcsech(b*x+a)^2*ln(1+(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))^
2)+arcsech(b*x+a)^2*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/
2)))/(1-(-a^2+1)^(1/2))+arcsech(b*x+a)^2*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1
/2)*(1/(b*x+a)+1)^(1/2)))/(1+(-a^2+1)^(1/2))-arcsech(b*x+a)*polylog(2,-(1/(
b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)+2*arcsech(b*x+a)*polylog
(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1-(-a^2+1)^(1/2))
)+2*arcsech(b*x+a)*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)

```

$$\begin{aligned} & \frac{1}{2} \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} \left(\frac{1}{1+(-a^2+1)^{1/2}} \right) + \frac{1}{2} \operatorname{polylog}\left(3, -\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} * \right. \\ & \left. \frac{1}{(b*x+a)+1} \right)^{1/2} - 2 \operatorname{polylog}\left(3, a * \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} * \right. \right. \\ & \left. \left. \frac{1}{(b*x+a)+1} \right)^{1/2} \right) / \left(1 - (-a^2+1)^{1/2}\right) - 2 \operatorname{polylog}\left(3, a * \left(\frac{1}{(b*x+a)} + \left(\frac{1}{(b*x+a)} - 1\right)^{1/2} * \right. \right. \\ & \left. \left. \frac{1}{(b*x+a)+1} \right)^{1/2} \right) / \left(1 + (-a^2+1)^{1/2}\right) \end{aligned}$$

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 274, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6456, 5714, 5689, 3799, 2221, 2611, 2320, 6724, 5681}

$$\begin{aligned} \int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} dx &= 2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\ &+ 2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2} + 1}\right) \\ &- 2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) - 2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2} + 1}\right) \\ &+ \operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\ &+ \operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2} + 1}\right) \\ &- \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\ &+ \frac{1}{2} \operatorname{PolyLog}\left(3, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\ &- \operatorname{sech}^{-1}(a+bx)^2 \log\left(e^{2\operatorname{sech}^{-1}(a+bx)} + 1\right) \end{aligned}$$

[In] Int[ArcSech[a + b*x]^2/x, x]

[Out] ArcSech[a + b*x]^2*Log[1 - (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])] + ArcSech[a + b*x]^2*Log[1 - (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])] - ArcSech[a + b*x]^2*Log[1 + E^(2*ArcSech[a + b*x])] + 2*ArcSech[a + b*x]*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])] + 2*ArcSech[a + b*x]*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])] - ArcSech[a + b*x]*PolyLog[2, -E^(2*ArcSech[a + b*x])] - 2*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])] - 2*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])] + PolyLog[3, -E^(2*ArcSech[a + b*x])]/2

Rule 2221

Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di

st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2611

Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.))] * ((f_.) + (g_.)*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3799

Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x))), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

Rule 5681

Int[(((e_.) + (f_.)*(x_))^(m_.)*Sinh[(c_.) + (d_.)*(x_)])/(Cosh[(c_.) + (d_.)*(x_)]*(b_.) + (a_.), x_Symbol] := Simp[-(e + f*x)^(m + 1)/(b*f*(m + 1)), x] + (Int[(e + f*x)^m*(E^(c + d*x))/(a - Rt[a^2 - b^2, 2] + b*E^(c + d*x)), x] + Int[(e + f*x)^m*(E^(c + d*x))/(a + Rt[a^2 - b^2, 2] + b*E^(c + d*x)), x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[a^2 - b^2, 0]

Rule 5689

Int[(((e_.) + (f_.)*(x_))^(m_.)*Tanh[(c_.) + (d_.)*(x_)])^(n_.)/(Cosh[(c_.) + (d_.)*(x_)]*(b_.) + (a_.), x_Symbol] := Dist[1/a, Int[(e + f*x)^m*Tanh[c + d*x]^n, x], x] - Dist[b/a, Int[(e + f*x)^m*Sinh[c + d*x]*(Tanh[c + d*x]^(n - 1)/(a + b*Cosh[c + d*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[n, 0]

Rule 5714

Int[(((e_.) + (f_.)*(x_))^(m_.)*(F_) [(c_.) + (d_.)*(x_)]^(n_.)*(G_) [(c_.) + (d_.)*(x_)]^(p_.))/(a_.) + (b_.)*Sech[(c_.) + (d_.)*(x_)], x_Symbol] := I

```
nt[(e + f*x)^m*Cosh[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Cosh[c + d*x
])), x] /; FreeQ[{a, b, c, d, e, f}, x] && HyperbolicQ[F] && HyperbolicQ[G]
&& IntegersQ[m, n, p]
```

Rule 6456

```
Int[((a_.) + ArcSech[(c_) + (d_.)*(x_)])*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Ta
nh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{x^2 \operatorname{sech}(x) \tanh(x)}{-a + \operatorname{sech}(x)} dx, x, \operatorname{sech}^{-1}(a + bx)\right) \\
&= -\text{Subst}\left(\int \frac{x^2 \tanh(x)}{1 - a \cosh(x)} dx, x, \operatorname{sech}^{-1}(a + bx)\right) \\
&= -\left(a \text{Subst}\left(\int \frac{x^2 \sinh(x)}{1 - a \cosh(x)} dx, x, \operatorname{sech}^{-1}(a + bx)\right)\right) \\
&\quad - \text{Subst}\left(\int x^2 \tanh(x) dx, x, \operatorname{sech}^{-1}(a + bx)\right) \\
&= -\left(2 \text{Subst}\left(\int \frac{e^{2x} x^2}{1 + e^{2x}} dx, x, \operatorname{sech}^{-1}(a + bx)\right)\right) \\
&\quad - a \text{Subst}\left(\int \frac{e^x x^2}{1 - \sqrt{1 - a^2} - a e^x} dx, x, \operatorname{sech}^{-1}(a + bx)\right) \\
&\quad - a \text{Subst}\left(\int \frac{e^x x^2}{1 + \sqrt{1 - a^2} - a e^x} dx, x, \operatorname{sech}^{-1}(a + bx)\right)
\end{aligned}$$

$$\begin{aligned}
&= \operatorname{sech}^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) + \operatorname{sech}^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&\quad - \operatorname{sech}^{-1}(a+bx)^2 \log \left(1 + e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
&\quad - 2\operatorname{Subst} \left(\int x \log \left(1 - \frac{ae^x}{1 - \sqrt{1-a^2}} \right) dx, x, \operatorname{sech}^{-1}(a+bx) \right) \\
&\quad - 2\operatorname{Subst} \left(\int x \log \left(1 - \frac{ae^x}{1 + \sqrt{1-a^2}} \right) dx, x, \operatorname{sech}^{-1}(a+bx) \right) \\
&\quad + 2\operatorname{Subst} \left(\int x \log (1 + e^{2x}) dx, x, \operatorname{sech}^{-1}(a+bx) \right) \\
&= \operatorname{sech}^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) + \operatorname{sech}^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&\quad - \operatorname{sech}^{-1}(a+bx)^2 \log \left(1 + e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
&\quad + 2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
&\quad + 2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&\quad - \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left(2, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
&\quad - 2\operatorname{Subst} \left(\int \operatorname{PolyLog} \left(2, \frac{ae^x}{1 - \sqrt{1-a^2}} \right) dx, x, \operatorname{sech}^{-1}(a+bx) \right) \\
&\quad - 2\operatorname{Subst} \left(\int \operatorname{PolyLog} \left(2, \frac{ae^x}{1 + \sqrt{1-a^2}} \right) dx, x, \operatorname{sech}^{-1}(a+bx) \right) \\
&\quad + \operatorname{Subst} \left(\int \operatorname{PolyLog} (2, -e^{2x}) dx, x, \operatorname{sech}^{-1}(a+bx) \right)
\end{aligned}$$

$$\begin{aligned}
&= \operatorname{sech}^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) + \operatorname{sech}^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&\quad - \operatorname{sech}^{-1}(a+bx)^2 \log \left(1 + e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
&\quad + 2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
&\quad + 2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&\quad - \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left(2, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
&\quad + \frac{1}{2} \operatorname{Subst} \left(\int \frac{\operatorname{PolyLog}(2, -x)}{x} dx, x, e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
&\quad - 2 \operatorname{Subst} \left(\int \frac{\operatorname{PolyLog} \left(2, \frac{ax}{1-\sqrt{1-a^2}} \right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)} \right) \\
&\quad - 2 \operatorname{Subst} \left(\int \frac{\operatorname{PolyLog} \left(2, \frac{ax}{1+\sqrt{1-a^2}} \right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)} \right) \\
&= \operatorname{sech}^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) + \operatorname{sech}^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&\quad - \operatorname{sech}^{-1}(a+bx)^2 \log \left(1 + e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
&\quad + 2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
&\quad + 2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&\quad - \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left(2, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) - 2 \operatorname{PolyLog} \left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
&\quad - 2 \operatorname{PolyLog} \left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) + \frac{1}{2} \operatorname{PolyLog} \left(3, -e^{2\operatorname{sech}^{-1}(a+bx)} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.34 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.02

$$\begin{aligned}
\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} dx = & -\frac{2}{3}\operatorname{sech}^{-1}(a+bx)^3 - \operatorname{sech}^{-1}(a+bx)^2 \log\left(1 + e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\
& + \operatorname{sech}^{-1}(a+bx)^2 \log\left(1 + \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}}\right) \\
& + \operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
& + \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\
& + 2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}}\right) \\
& + 2\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
& + \frac{1}{2} \operatorname{PolyLog}\left(3, -e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\
& - 2 \operatorname{PolyLog}\left(3, -\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}}\right) - 2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right)
\end{aligned}$$

[In] Integrate[ArcSech[a + b*x]^2/x,x]

```

[Out] (-2*ArcSech[a + b*x]^3)/3 - ArcSech[a + b*x]^2*Log[1 + E^(-2*ArcSech[a + b*
x])] + ArcSech[a + b*x]^2*Log[1 + (a*E^ArcSech[a + b*x])/(-1 + Sqrt[1 - a^2
])] + ArcSech[a + b*x]^2*Log[1 - (a*E^ArcSech[a + b*x]/(1 + Sqrt[1 - a^2])
] + ArcSech[a + b*x]*PolyLog[2, -E^(-2*ArcSech[a + b*x])] + 2*ArcSech[a + b
*x]*PolyLog[2, -((a*E^ArcSech[a + b*x])/(-1 + Sqrt[1 - a^2]))] + 2*ArcSech[
a + b*x]*PolyLog[2, (a*E^ArcSech[a + b*x]/(1 + Sqrt[1 - a^2])] + PolyLog[3
, -E^(-2*ArcSech[a + b*x])/2 - 2*PolyLog[3, -((a*E^ArcSech[a + b*x])/(-1 +
Sqrt[1 - a^2]))] - 2*PolyLog[3, (a*E^ArcSech[a + b*x]/(1 + Sqrt[1 - a^2])
]

```

Maple [F]

$$\int \frac{\operatorname{arcsech}(bx+a)^2}{x} dx$$

```
[In] int(arcsech(b*x+a)^2/x,x)
```

```
[Out] int(arcsech(b*x+a)^2/x,x)
```

Fricas [F]

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} dx = \int \frac{\operatorname{arsech}(bx+a)^2}{x} dx$$

```
[In] integrate(arcsech(b*x+a)^2/x,x, algorithm="fricas")
```

```
[Out] integral(arcsech(b*x + a)^2/x, x)
```

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} dx = \int \frac{\operatorname{asech}^2(a+bx)}{x} dx$$

```
[In] integrate(asech(b*x+a)**2/x,x)
```

```
[Out] Integral(asech(a + b*x)**2/x, x)
```

Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} dx = \int \frac{\operatorname{arsech}(bx+a)^2}{x} dx$$

```
[In] integrate(arcsech(b*x+a)^2/x,x, algorithm="maxima")
```

```
[Out] integrate(arcsech(b*x + a)^2/x, x)
```

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{ar} \operatorname{sech}(bx + a)^2}{x} dx$$

[In] integrate(arcsech(b*x+a)^2/x,x, algorithm="giac")

[Out] integrate(arcsech(b*x + a)^2/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)^2}{x} dx$$

[In] int(acosh(1/(a + b*x))^2/x,x)

[Out] int(acosh(1/(a + b*x))^2/x, x)

3.13 $\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx$

Optimal result	134
Rubi [A] (verified)	135
Mathematica [C] (verified)	138
Maple [A] (verified)	139
Fricas [F]	139
Sympy [F]	139
Maxima [F]	140
Giac [F]	140
Mupad [F(-1)]	140

Optimal result

Integrand size = 12, antiderivative size = 224

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx = -\frac{b\operatorname{sech}^{-1}(a+bx)^2}{a} - \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} + \frac{2b\operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{2b\operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{2b \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{2b \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}}$$

```
[Out] -b*arcsech(b*x+a)^2/a-arcsech(b*x+a)^2/x+2*b*arcsech(b*x+a)*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-2*b*arcsech(b*x+a)*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+2*b*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-2*b*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)
```

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {6456, 5576, 4276, 3401, 2296, 2221, 2317, 2438}

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx = \frac{2b \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{2b \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a\sqrt{1-a^2}}$$

$$+ \frac{2b \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}}$$

$$- \frac{2b \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a\sqrt{1-a^2}}$$

$$- \frac{b \operatorname{sech}^{-1}(a+bx)^2}{a} - \frac{\operatorname{sech}^{-1}(a+bx)^2}{x}$$

[In] Int[ArcSech[a + b*x]^2/x^2,x]

[Out] -((b*ArcSech[a + b*x]^2)/a) - ArcSech[a + b*x]^2/x + (2*b*ArcSech[a + b*x]*Log[1 - (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) - (2*b*ArcSech[a + b*x]*Log[1 - (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) + (2*b*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) - (2*b*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2])

Rule 2221

Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2296

Int[(F_)^(u)*((f_) + (g_)*(x_))^(m_)]/((a_) + (b_)*(F_)^(u) + (c_)*(F_)^(v)), x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[2*(c/q), Int[(f + g*x)^m*(F^u/(b - q + 2*c*F^u)), x], x] - Dist[2*(c/q), Int[(f + g*x)^m*(F^u/(b + q + 2*c*F^u)), x], x]] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v, 2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]

Rule 2317

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))]

)^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_.)*((d_.) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3401

Int[((c_.) + (d_.)*(x_)^(m_.))/((a_.) + (b_.)*sin[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]), x_Symbol] := Dist[2, Int[((c + d*x)^m*(E^((-I)*e + f*fz*x)/(b + (2*a*E^((-I)*e + f*fz*x))/E^(I*Pi*(k - 1/2)) - (b*E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi))))/E^(I*Pi*(k - 1/2)), x], x] /; FreeQ[{a, b, c, d, e, f, fz}, x] && IntegerQ[2*k] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]

Rule 4276

Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)^(n_.))*((c_.) + (d_.)*(x_)^(m_.)), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Sin[e + f*x]^n)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGtQ[m, 0]

Rule 5576

Int[((e_.) + (f_.)*(x_)^(m_.))*Sech[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sech[(c_.) + (d_.)*(x_)]^(n_.)*Tanh[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[(-e + f*x)^m*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rule 6456

Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_)^(m_.)), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\left(b\text{Subst}\left(\int \frac{x^2 \operatorname{sech}(x) \tanh(x)}{(-a + \operatorname{sech}(x))^2} dx, x, \operatorname{sech}^{-1}(a + bx)\right)\right) \\
 &= -\frac{\operatorname{sech}^{-1}(a + bx)^2}{x} + (2b)\text{Subst}\left(\int \frac{x}{-a + \operatorname{sech}(x)} dx, x, \operatorname{sech}^{-1}(a + bx)\right) \\
 &= -\frac{\operatorname{sech}^{-1}(a + bx)^2}{x} + (2b)\text{Subst}\left(\int \left(-\frac{x}{a} + \frac{x}{a(1 - a \cosh(x))}\right) dx, x, \operatorname{sech}^{-1}(a + bx)\right)
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{b\operatorname{sech}^{-1}(a+bx)^2}{a} - \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} + \frac{(2b)\operatorname{Subst}\left(\int \frac{x}{1-a\cosh(x)} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a} \\
&= -\frac{b\operatorname{sech}^{-1}(a+bx)^2}{a} - \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} + \frac{(4b)\operatorname{Subst}\left(\int \frac{e^x x}{-a+2e^x-ae^{2x}} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a} \\
&= -\frac{b\operatorname{sech}^{-1}(a+bx)^2}{a} - \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} \\
&\quad - \frac{(4b)\operatorname{Subst}\left(\int \frac{e^x x}{2-2\sqrt{1-a^2}-2ae^x} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a^2}} \\
&\quad + \frac{(4b)\operatorname{Subst}\left(\int \frac{e^x x}{2+2\sqrt{1-a^2}-2ae^x} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{1-a^2}} \\
&= -\frac{b\operatorname{sech}^{-1}(a+bx)^2}{a} - \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} + \frac{2b\operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{2b\operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{(2b)\operatorname{Subst}\left(\int \log\left(1 - \frac{2ae^x}{2-2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a\sqrt{1-a^2}} \\
&\quad + \frac{(2b)\operatorname{Subst}\left(\int \log\left(1 - \frac{2ae^x}{2+2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a\sqrt{1-a^2}} \\
&= -\frac{b\operatorname{sech}^{-1}(a+bx)^2}{a} - \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} + \frac{2b\operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{2b\operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{(2b)\operatorname{Subst}\left(\int \frac{\log\left(1 - \frac{2ax}{2-2\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{a\sqrt{1-a^2}} \\
&\quad + \frac{(2b)\operatorname{Subst}\left(\int \frac{\log\left(1 - \frac{2ax}{2+2\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{a\sqrt{1-a^2}}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{b\operatorname{sech}^{-1}(a+bx)^2}{a} - \frac{\operatorname{sech}^{-1}(a+bx)^2}{x} + \frac{2b\operatorname{sech}^{-1}(a+bx)\log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{2b\operatorname{sech}^{-1}(a+bx)\log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad + \frac{2b\operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{2b\operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.83 (sec) , antiderivative size = 678, normalized size of antiderivative = 3.03

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx$$

$$= \frac{-(a+bx)\operatorname{sech}^{-1}(a+bx)^2}{x} + \frac{2b\left(2\operatorname{sech}^{-1}(a+bx)\arctan\left(\frac{(-1+a)\coth\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{-1+a^2}}\right) - 2i\arccos\left(\frac{1}{a}\right)\arctan\left(\frac{(1+a)\tanh\left(\frac{1}{2}\operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{-1+a^2}}\right)\right)}{\sqrt{-1+a^2}}$$

[In] Integrate[ArcSech[a + b*x]^2/x^2,x]

[Out] (-((a + b*x)*ArcSech[a + b*x]^2)/x) + (2*b*(2*ArcSech[a + b*x]*ArcTan[(-1 + a)*Coth[ArcSech[a + b*x]/2]]/Sqrt[-1 + a^2]] - (2*I)*ArcCos[a^(-1)]*ArcTan[(((1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]] + (ArcCos[a^(-1)] + 2*(ArcTan[(-1 + a)*Coth[ArcSech[a + b*x]/2]]/Sqrt[-1 + a^2]] + ArcTan[(((1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]]))*Log[Sqrt[-1 + a^2]/(Sqrt[2]*Sqrt[a]*E^(ArcSech[a + b*x]/2)*Sqrt[-((b*x)/(a + b*x))])] + (ArcCos[a^(-1)] - 2*(ArcTan[(-1 + a)*Coth[ArcSech[a + b*x]/2]]/Sqrt[-1 + a^2]] + ArcTan[(((1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]]))*Log[(Sqrt[-1 + a^2]*E^(ArcSech[a + b*x]/2))/(Sqrt[2]*Sqrt[a]*Sqrt[-((b*x)/(a + b*x))])] - (ArcCos[a^(-1)] + 2*ArcTan[(((1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]])*Log[-(((-1 + a)*(1 + a - I*Sqrt[-1 + a^2])*(-1 + Tanh[ArcSech[a + b*x]/2]))/(a*(-1 + a + I*Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2])))] - (ArcCos[a^(-1)] - 2*ArcTan[(((1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]])*Log[(-1 + a)*(1 + a + I*Sqrt[-1 + a^2])*(1 + Tanh[ArcSech[a + b*x]/2])]/(a*(-1 + a + I*Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2]))] + I*(PolyLog[2, ((-1 - I*Sqrt[-1 + a^2])*(-1 + a - I*Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2]))/(a*(-1 + a + I*Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2]))] - PolyLog[2, ((I + Sqrt[-1 + a^2])*(-1 + a - I*Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2]))/(a*((-I)*(-1 + a) + Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2])))]/Sqrt[-1 + a^2])/a

Maple [A] (verified)

Time = 0.92 (sec) , antiderivative size = 362, normalized size of antiderivative = 1.62

method	result
derivativedivides	$b \left(-\frac{(bx+a) \operatorname{arcsech}(bx+a)^2}{abx} + \frac{2\sqrt{-a^2+1} \operatorname{arcsech}(bx+a) \ln \left(\frac{-a \left(\frac{1}{bx+a} + \sqrt{\frac{1}{bx+a}-1} \sqrt{\frac{1}{bx+a}+1} \right) + \sqrt{-a^2+1+1}}{1+\sqrt{-a^2+1}} \right)}{a(a^2-1)} \right)$
default	$b \left(-\frac{(bx+a) \operatorname{arcsech}(bx+a)^2}{abx} + \frac{2\sqrt{-a^2+1} \operatorname{arcsech}(bx+a) \ln \left(\frac{-a \left(\frac{1}{bx+a} + \sqrt{\frac{1}{bx+a}-1} \sqrt{\frac{1}{bx+a}+1} \right) + \sqrt{-a^2+1+1}}{1+\sqrt{-a^2+1}} \right)}{a(a^2-1)} \right)$

```
[In] int(arcsech(b*x+a)^2/x^2,x,method=_RETURNVERBOSE)
```

```
[Out] b*(-(b*x+a)*arcsech(b*x+a)^2/a/b/x+2*(-a^2+1)^(1/2)/a/(a^2-1)*arcsech(b*x+a)
)*ln((-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))+(-a^2+1)^(1/2)
+1)/(1+(-a^2+1)^(1/2)))-2*(-a^2+1)^(1/2)/a/(a^2-1)*arcsech(b*x+a)*ln((a*(1/
(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))+(-a^2+1)^(1/2)-1)/(-1+(-a^
2+1)^(1/2)))+2*(-a^2+1)^(1/2)/a/(a^2-1)*dilog((-a*(1/(b*x+a)+(1/(b*x+a)-1)^(
1/2)*(1/(b*x+a)+1)^(1/2)))+(-a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))-2*(-a^2+1)
^(1/2)/a/(a^2-1)*dilog((a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)
)))+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2)))
```

Fricas [F]

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx = \int \frac{\operatorname{arsech}(bx+a)^2}{x^2} dx$$

```
[In] integrate(arcsech(b*x+a)^2/x^2,x, algorithm="fricas")
```

```
[Out] integral(arcsech(b*x + a)^2/x^2, x)
```

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx = \int \frac{\operatorname{asech}^2(a+bx)}{x^2} dx$$

```
[In] integrate(asech(b*x+a)**2/x**2,x)
```

```
[Out] Integral(asech(a + b*x)**2/x**2, x)
```

Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx = \int \frac{\operatorname{arosech}(bx+a)^2}{x^2} dx$$

[In] integrate(arcsech(b*x+a)^2/x^2,x, algorithm="maxima")

[Out] -log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^2/x - integrate(-2*(2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^2 + 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a)^2 + (b^3*x^3 + 2*a*b^2*x^2 + (a^2*b - b)*x - 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a) - ((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*log(b*x + a) - (2*b^3*x^3 + 4*a*b^2*x^2 + (2*a^2*b - b)*x - (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a))*sqrt(b*x + a + 1))*sqrt(-b*x - a + 1))*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a))/(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2 + (b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)), x)

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx = \int \frac{\operatorname{arosech}(bx+a)^2}{x^2} dx$$

[In] integrate(arcsech(b*x+a)^2/x^2,x, algorithm="giac")

[Out] integrate(arcsech(b*x + a)^2/x^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^2} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)^2}{x^2} dx$$

[In] int(acosh(1/(a + b*x))^2/x^2,x)

[Out] int(acosh(1/(a + b*x))^2/x^2, x)

3.14 $\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^3} dx$

Optimal result	141
Rubi [A] (verified)	142
Mathematica [C] (verified)	149
Maple [A] (verified)	151
Fricas [F]	152
Sympy [F]	152
Maxima [F]	152
Giac [F]	153
Mupad [F(-1)]	153

Optimal result

Integrand size = 12, antiderivative size = 537

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^3} dx = \frac{b^2 \sqrt{\frac{1-a-bx}{1+a+bx}} (1+a+bx) \operatorname{sech}^{-1}(a+bx)}{a(1-a^2)(a+bx) \left(1 - \frac{a}{a+bx}\right)} + \frac{b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2}$$

$$- \frac{\operatorname{sech}^{-1}(a+bx)^2}{2x^2} + \frac{b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}}$$

$$- \frac{2b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}}$$

$$- \frac{b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}}$$

$$+ \frac{2b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} + \frac{b^2 \log\left(\frac{x}{a+bx}\right)}{a^2(1-a^2)}$$

$$+ \frac{b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} - \frac{2b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}}$$

$$- \frac{b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} + \frac{2b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}}$$

[Out] $1/2*b^2*\operatorname{arcsech}(b*x+a)^2/a^2-1/2*\operatorname{arcsech}(b*x+a)^2/x^2+b^2*\ln(x/(b*x+a))/a^2$
 $/(-a^2+1)+b^2*\operatorname{arcsech}(b*x+a)*\ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1-(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(3/2)-b^2*\operatorname{arcsech}(b*x+a)*\ln(1$
 $-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/$
 $a^2/(-a^2+1)^(3/2)+b^2*\operatorname{polylog}(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)$

$$\begin{aligned} &)+1)^{(1/2)})/(1-(-a^2+1)^{(1/2)})))/a^2/(-a^2+1)^{(3/2)}-b^2*\text{polylog}(2,a*(1/(b*x+a)+1/(b*x+a)-1)^{(1/2)}*(1/(b*x+a)+1)^{(1/2)}))/(1+(-a^2+1)^{(1/2)})))/a^2/(-a^2+1)^{(3/2)}-2*b^2*\text{arcsech}(b*x+a)*\ln(1-a*(1/(b*x+a)+1/(b*x+a)-1)^{(1/2)}*(1/(b*x+a)+1)^{(1/2)}))/(1-(-a^2+1)^{(1/2)})))/a^2/(-a^2+1)^{(1/2)}+2*b^2*\text{arcsech}(b*x+a)*\ln(1-a*(1/(b*x+a)+1/(b*x+a)-1)^{(1/2)}*(1/(b*x+a)+1)^{(1/2)}))/(1+(-a^2+1)^{(1/2)})))/a^2/(-a^2+1)^{(1/2)}-2*b^2*\text{polylog}(2,a*(1/(b*x+a)+1/(b*x+a)-1)^{(1/2)}*(1/(b*x+a)+1)^{(1/2)}))/(1-(-a^2+1)^{(1/2)})))/a^2/(-a^2+1)^{(1/2)}+2*b^2*\text{polylog}(2,a*(1/(b*x+a)+1/(b*x+a)-1)^{(1/2)}*(1/(b*x+a)+1)^{(1/2)}))/(1+(-a^2+1)^{(1/2)})))/a^2/(-a^2+1)^{(1/2)}+b^2*(b*x+a+1)*\text{arcsech}(b*x+a)*((-b*x-a+1)/(b*x+a+1))^{(1/2)}/a/(-a^2+1)/(b*x+a)/(1-a/(b*x+a)) \end{aligned}$$

Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 537, normalized size of antiderivative = 1.00, number of steps used = 23, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$, Rules used = {6456, 5576, 4276, 3405, 3401, 2296, 2221, 2317, 2438, 2747, 31}

$$\begin{aligned} \int \frac{\text{sech}^{-1}(a+bx)^2}{x^3} dx = & -\frac{2b^2 \text{PolyLog}\left(2, \frac{ae^{\text{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2\sqrt{1-a^2}} + \frac{b^2 \text{PolyLog}\left(2, \frac{ae^{\text{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\ & + \frac{2b^2 \text{PolyLog}\left(2, \frac{ae^{\text{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a^2\sqrt{1-a^2}} - \frac{b^2 \text{PolyLog}\left(2, \frac{ae^{\text{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a^2(1-a^2)^{3/2}} \\ & + \frac{b^2 \log\left(\frac{x}{a+bx}\right)}{a^2(1-a^2)} + \frac{b^2 \text{sech}^{-1}(a+bx)^2}{2a^2} \\ & + \frac{b^2 \sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\text{sech}^{-1}(a+bx)}{a(1-a^2)(a+bx)\left(1-\frac{a}{a+bx}\right)} \\ & - \frac{2b^2 \text{sech}^{-1}(a+bx) \log\left(1-\frac{ae^{\text{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2\sqrt{1-a^2}} \\ & + \frac{b^2 \text{sech}^{-1}(a+bx) \log\left(1-\frac{ae^{\text{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\ & + \frac{2b^2 \text{sech}^{-1}(a+bx) \log\left(1-\frac{ae^{\text{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a^2\sqrt{1-a^2}} \\ & - \frac{b^2 \text{sech}^{-1}(a+bx) \log\left(1-\frac{ae^{\text{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a^2(1-a^2)^{3/2}} - \frac{\text{sech}^{-1}(a+bx)^2}{2x^2} \end{aligned}$$

[In] Int[ArcSech[a + b*x]^2/x^3, x]

[Out] (b^2*sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x)*ArcSech[a + b*x])/(a*(1 - a^2)*(a + b*x)*(1 - a/(a + b*x))) + (b^2*ArcSech[a + b*x]^2)/(2*a^2) -

```
ArcSech[a + b*x]^2/(2*x^2) + (b^2*ArcSech[a + b*x]*Log[1 - (a*E^ArcSech[a +
b*x])/(1 - Sqrt[1 - a^2])])/(a^2*(1 - a^2)^(3/2)) - (2*b^2*ArcSech[a + b*x]
)*Log[1 - (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/(a^2*Sqrt[1 - a^2])
- (b^2*ArcSech[a + b*x]*Log[1 - (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])
)/(a^2*(1 - a^2)^(3/2)) + (2*b^2*ArcSech[a + b*x]*Log[1 - (a*E^ArcSech[a +
b*x])/(1 + Sqrt[1 - a^2])])/(a^2*Sqrt[1 - a^2]) + (b^2*Log[x/(a + b*x)]/(a
^2*(1 - a^2)) + (b^2*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])
)/(a^2*(1 - a^2)^(3/2)) - (2*b^2*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 - Sqr
t[1 - a^2])])/(a^2*Sqrt[1 - a^2]) - (b^2*PolyLog[2, (a*E^ArcSech[a + b*x])/
(1 + Sqrt[1 - a^2])])/(a^2*(1 - a^2)^(3/2)) + (2*b^2*PolyLog[2, (a*E^ArcSec
h[a + b*x])/(1 + Sqrt[1 - a^2])])/(a^2*Sqrt[1 - a^2])
```

Rule 31

```
Int[((a_) + (b_.)*(x_))^-1, x_Symbol] := Simp[Log[RemoveContent[a + b*x,
x]]/b, x] /; FreeQ[{a, b}, x]
```

Rule 2221

```
Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/
((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)
))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2296

```
Int[((F_)^(u_.)*((f_.) + (g_.)*(x_))^(m_.))/((a_.) + (b_.)*(F_)^(u_) + (c_.)
*(F_)^(v_)), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[2*(c/q), Int[
(f + g*x)^m*(F^u/(b - q + 2*c*F^u)), x], x] - Dist[2*(c/q), Int[(f + g*x)^m
*(F^u/(b + q + 2*c*F^u)), x], x]] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v,
2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol]
:= Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x))
)^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*((d_) + (e_.)*(x_))^(n_.)]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2747

```
Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m
_.)), x_Symbol] := Dist[1/(b^p*f), Subst[Int[(a + x)^m*(b^2 - x^2)^(p - 1)/
```

2), x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && NeQ[a^2 - b^2, 0]

Rule 3401

Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*sin[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]), x_Symbol] := Dist[2, Int[((c + d*x)^m*(E^((-I)*e + f*fz*x)/(b + (2*a*E^((-I)*e + f*fz*x))/E^(I*Pi*(k - 1/2)) - (b*E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi))))/E^(I*Pi*(k - 1/2)), x], x] /; FreeQ[{a, b, c, d, e, f, fz}, x] && IntegerQ[2*k] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]

Rule 3405

Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[b*(c + d*x)^m*(Cos[e + f*x]/(f*(a^2 - b^2)*(a + b*Sin[e + f*x]))], x] + (Dist[a/(a^2 - b^2), Int[(c + d*x)^m/(a + b*Sin[e + f*x]), x], x] - Dist[b*d*(m/(f*(a^2 - b^2))), Int[(c + d*x)^(m - 1)*(Cos[e + f*x]/(a + b*Sin[e + f*x])), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]

Rule 4276

Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Sin[e + f*x])^n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGtQ[m, 0]

Rule 5576

Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_) + (b_.)*Sech[(c_.) + (d_.)*(x_)]^(n_.)*Tanh[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rule 6456

Int[((a_.) + ArcSech[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rubi steps

$$\text{integral} = -\left(b^2 \text{Subst}\left(\int \frac{x^2 \text{sech}(x) \tanh(x)}{(-a + \text{sech}(x))^3} dx, x, \text{sech}^{-1}(a + bx)\right)\right)$$

$$\begin{aligned}
&= -\frac{\operatorname{sech}^{-1}(a+bx)^2}{2x^2} + b^2 \operatorname{Subst} \left(\int \frac{x}{(-a + \operatorname{sech}(x))^2} dx, x, \operatorname{sech}^{-1}(a+bx) \right) \\
&= -\frac{\operatorname{sech}^{-1}(a+bx)^2}{2x^2} + b^2 \operatorname{Subst} \left(\int \left(\frac{x}{a^2} + \frac{x}{a^2(-1+a \cosh(x))^2} \right. \right. \\
&\quad \left. \left. + \frac{2x}{a^2(-1+a \cosh(x))} \right) dx, x, \operatorname{sech}^{-1}(a+bx) \right) \\
&= \frac{b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)^2}{2x^2} \\
&\quad + \frac{b^2 \operatorname{Subst} \left(\int \frac{x}{(-1+a \cosh(x))^2} dx, x, \operatorname{sech}^{-1}(a+bx) \right)}{a^2} \\
&\quad + \frac{(2b^2) \operatorname{Subst} \left(\int \frac{x}{-1+a \cosh(x)} dx, x, \operatorname{sech}^{-1}(a+bx) \right)}{a^2} \\
&= \frac{b^2 \sqrt{\frac{1-a-bx}{1+a+bx}} (1+a+bx) \operatorname{sech}^{-1}(a+bx)}{a(1-a^2)(a+bx) \left(1 - \frac{a}{a+bx}\right)} + \frac{b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2} \\
&\quad - \frac{\operatorname{sech}^{-1}(a+bx)^2}{2x^2} + \frac{(4b^2) \operatorname{Subst} \left(\int \frac{e^x x}{a-2e^x+ae^{2x}} dx, x, \operatorname{sech}^{-1}(a+bx) \right)}{a^2} \\
&\quad - \frac{b^2 \operatorname{Subst} \left(\int \frac{x}{-1+a \cosh(x)} dx, x, \operatorname{sech}^{-1}(a+bx) \right)}{a^2(1-a^2)} \\
&\quad + \frac{b^2 \operatorname{Subst} \left(\int \frac{\sinh(x)}{-1+a \cosh(x)} dx, x, \operatorname{sech}^{-1}(a+bx) \right)}{a(1-a^2)} \\
&= \frac{b^2 \sqrt{\frac{1-a-bx}{1+a+bx}} (1+a+bx) \operatorname{sech}^{-1}(a+bx)}{a(1-a^2)(a+bx) \left(1 - \frac{a}{a+bx}\right)} + \frac{b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)^2}{2x^2} \\
&\quad + \frac{b^2 \operatorname{Subst} \left(\int \frac{1}{-1+x} dx, x, \frac{a}{a+bx} \right)}{a^2(1-a^2)} - \frac{(2b^2) \operatorname{Subst} \left(\int \frac{e^x x}{a-2e^x+ae^{2x}} dx, x, \operatorname{sech}^{-1}(a+bx) \right)}{a^2(1-a^2)} \\
&\quad + \frac{(4b^2) \operatorname{Subst} \left(\int \frac{e^x x}{-2-2\sqrt{1-a^2}+2ae^x} dx, x, \operatorname{sech}^{-1}(a+bx) \right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{(4b^2) \operatorname{Subst} \left(\int \frac{e^x x}{-2+2\sqrt{1-a^2}+2ae^x} dx, x, \operatorname{sech}^{-1}(a+bx) \right)}{a\sqrt{1-a^2}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^2 \sqrt{\frac{1-a-bx}{1+a+bx}} (1+a+bx) \operatorname{sech}^{-1}(a+bx)}{a(1-a^2)(a+bx) \left(1 - \frac{a}{a+bx}\right)} + \frac{b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2} \\
&\quad - \frac{\operatorname{sech}^{-1}(a+bx)^2}{2x^2} - \frac{2b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&\quad + \frac{2b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} + \frac{b^2 \log\left(\frac{x}{a+bx}\right)}{a^2(1-a^2)} \\
&\quad - \frac{(2b^2) \operatorname{Subst}\left(\int \frac{e^x x}{-2-2\sqrt{1-a^2}+2ae^x} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a(1-a^2)^{3/2}} \\
&\quad + \frac{(2b^2) \operatorname{Subst}\left(\int \frac{e^x x}{-2+2\sqrt{1-a^2}+2ae^x} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a(1-a^2)^{3/2}} \\
&\quad - \frac{(2b^2) \operatorname{Subst}\left(\int \log\left(1 + \frac{2ae^x}{-2-2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2 \sqrt{1-a^2}} \\
&\quad + \frac{(2b^2) \operatorname{Subst}\left(\int \log\left(1 + \frac{2ae^x}{-2+2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2 \sqrt{1-a^2}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^2 \sqrt{\frac{1-a-bx}{1+a+bx}} (1+a+bx) \operatorname{sech}^{-1}(a+bx)}{a(1-a^2)(a+bx)\left(1-\frac{a}{a+bx}\right)} + \frac{b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2} \\
&\quad - \frac{\operatorname{sech}^{-1}(a+bx)^2}{2x^2} + \frac{b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
&\quad - \frac{2b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&\quad - \frac{b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
&\quad + \frac{2b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} + \frac{b^2 \log\left(\frac{x}{a+bx}\right)}{a^2(1-a^2)} \\
&\quad + \frac{b^2 \operatorname{Subst}\left(\int \log\left(1 + \frac{2ae^x}{-2-2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2(1-a^2)^{3/2}} \\
&\quad - \frac{b^2 \operatorname{Subst}\left(\int \log\left(1 + \frac{2ae^x}{-2+2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2(1-a^2)^{3/2}} \\
&\quad - \frac{(2b^2) \operatorname{Subst}\left(\int \frac{\log\left(1 + \frac{2ax}{-2-2\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{a^2 \sqrt{1-a^2}} \\
&\quad + \frac{(2b^2) \operatorname{Subst}\left(\int \frac{\log\left(1 + \frac{2ax}{-2+2\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{a^2 \sqrt{1-a^2}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^2 \sqrt{\frac{1-a-bx}{1+a+bx}} (1+a+bx) \operatorname{sech}^{-1}(a+bx)}{a(1-a^2)(a+bx) \left(1 - \frac{a}{a+bx}\right)} + \frac{b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2} \\
&- \frac{\operatorname{sech}^{-1}(a+bx)^2}{2x^2} + \frac{b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
&- \frac{2b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&- \frac{b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
&+ \frac{2b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} + \frac{b^2 \log\left(\frac{x}{a+bx}\right)}{a^2(1-a^2)} \\
&- \frac{2b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} + \frac{2b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{b^2 \operatorname{Subst}\left(\int \frac{\log\left(1 + \frac{2ax}{-2-2\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{a^2(1-a^2)^{3/2}} \\
&- \frac{b^2 \operatorname{Subst}\left(\int \frac{\log\left(1 + \frac{2ax}{-2+2\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{a^2(1-a^2)^{3/2}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^2 \sqrt{\frac{1-a-bx}{1+a+bx}} (1+a+bx) \operatorname{sech}^{-1}(a+bx)}{a(1-a^2)(a+bx) \left(1 - \frac{a}{a+bx}\right)} + \frac{b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2} \\
&\quad - \frac{\operatorname{sech}^{-1}(a+bx)^2}{2x^2} + \frac{b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
&\quad - \frac{2b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&\quad - \frac{b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
&\quad + \frac{2b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} + \frac{b^2 \log\left(\frac{x}{a+bx}\right)}{a^2(1-a^2)} \\
&\quad + \frac{b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} - \frac{2b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&\quad - \frac{b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} + \frac{2b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 8.00 (sec) , antiderivative size = 1439, normalized size of antiderivative = 2.68

$$\begin{aligned}
\int \frac{\operatorname{sech}^{-1}(a+bx)^2}{x^3} dx &= -\frac{(a+bx)^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2 x^2} \\
&+ \frac{b \operatorname{sech}^{-1}(a+bx) \left(-a \sqrt{-\frac{-1+a+bx}{1+a+bx}} (1+a+bx) + (-1+a^2)(a+bx) \operatorname{sech}^{-1}(a+bx) \right)}{(-1+a)a^2(1+a)x} \\
&+ \frac{b^2 \log\left(\frac{bx}{a+bx}\right)}{a^2 - a^4} \\
&- \frac{2b^2 \left(2 \operatorname{sech}^{-1}(a+bx) \arctan\left(\frac{(-1+a) \coth\left(\frac{1}{2} \operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{-1+a^2}}\right) - 2i \arccos\left(\frac{1}{a}\right) \arctan\left(\frac{(1+a) \tanh\left(\frac{1}{2} \operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{-1+a^2}}\right) \right)}{1} \\
&+ \frac{b^2 \left(2 \operatorname{sech}^{-1}(a+bx) \arctan\left(\frac{(-1+a) \coth\left(\frac{1}{2} \operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{-1+a^2}}\right) - 2i \arccos\left(\frac{1}{a}\right) \arctan\left(\frac{(1+a) \tanh\left(\frac{1}{2} \operatorname{sech}^{-1}(a+bx)\right)}{\sqrt{-1+a^2}}\right) \right)}{1}
\end{aligned}$$

[In] Integrate[ArcSech[a + b*x]^2/x^3, x]

```

[Out] -1/2*((a + b*x)^2*ArcSech[a + b*x]^2)/(a^2*x^2) + (b*ArcSech[a + b*x]*(-a*
Sqrt[-((-1 + a + b*x)/(1 + a + b*x))]*(1 + a + b*x)) + (-1 + a^2)*(a + b*x)
*ArcSech[a + b*x]))/((-1 + a)*a^2*(1 + a)*x) + (b^2*Log[(b*x)/(a + b*x)])/(
a^2 - a^4) - (2*b^2*(2*ArcSech[a + b*x]*ArcTan[((-1 + a)*Coth[ArcSech[a + b
*x]/2]))/Sqrt[-1 + a^2]] - (2*I)*ArcCos[a^(-1)]*ArcTan[((1 + a)*Tanh[ArcSech
[a + b*x]/2])/Sqrt[-1 + a^2]] + (ArcCos[a^(-1)] + 2*(ArcTan[((-1 + a)*Coth[
ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]] + ArcTan[((1 + a)*Tanh[ArcSech[a + b*x
]/2])/Sqrt[-1 + a^2]]))*Log[Sqrt[-1 + a^2]/(Sqrt[2]*Sqrt[a]*E^(ArcSech[a +
b*x]/2)*Sqrt[-((b*x)/(a + b*x))])] + (ArcCos[a^(-1)] - 2*(ArcTan[((-1 + a)*
Coth[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]] + ArcTan[((1 + a)*Tanh[ArcSech[a
+ b*x]/2])/Sqrt[-1 + a^2]]))*Log[(Sqrt[-1 + a^2]*E^(ArcSech[a + b*x]/2))/(S
qrt[2]*Sqrt[a]*Sqrt[-((b*x)/(a + b*x))])] - (ArcCos[a^(-1)] + 2*ArcTan[((1
+ a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]])*Log[-(((1 + a)*(1 + a - I*
Sqrt[-1 + a^2])*(-1 + Tanh[ArcSech[a + b*x]/2]))/(a*(-1 + a + I*Sqrt[-1 + a
^2]*Tanh[ArcSech[a + b*x]/2])))] - (ArcCos[a^(-1)] - 2*ArcTan[((1 + a)*Tanh
[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]])*Log[((-1 + a)*(1 + a + I*Sqrt[-1 + a
^2])*(1 + Tanh[ArcSech[a + b*x]/2]))/(a*(-1 + a + I*Sqrt[-1 + a^2]*Tanh[Arc
Sech[a + b*x]/2]))] + I*(PolyLog[2, ((-1 - I*Sqrt[-1 + a^2])*(-1 + a - I*Sq
rt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2]))/(a*(-1 + a + I*Sqrt[-1 + a^2]*Tanh[
ArcSech[a + b*x]/2]))] - PolyLog[2, ((I + Sqrt[-1 + a^2])*(-1 + a - I*Sqrt[
-1 + a^2]*Tanh[ArcSech[a + b*x]/2]))/(a*((-I)*(-1 + a) + Sqrt[-1 + a^2]*Tan
h[ArcSech[a + b*x]/2])))])))/(-1 + a^2)^(3/2) + (b^2*(2*ArcSech[a + b*x]*Arc
Tan[((-1 + a)*Coth[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]] - (2*I)*ArcCos[a^(-
1)]*ArcTan[((1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]] + (ArcCos[a^(-
1)] + 2*(ArcTan[((-1 + a)*Coth[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]] + ArcT
an[((1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]]))*Log[Sqrt[-1 + a^2]/
(Sqrt[2]*Sqrt[a]*E^(ArcSech[a + b*x]/2)*Sqrt[-((b*x)/(a + b*x))])] + (ArcCo
s[a^(-1)] - 2*(ArcTan[((-1 + a)*Coth[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]] +
ArcTan[((1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]]))*Log[(Sqrt[-1 +
a^2]*E^(ArcSech[a + b*x]/2))/(Sqrt[2]*Sqrt[a]*Sqrt[-((b*x)/(a + b*x))])] -
(ArcCos[a^(-1)] + 2*ArcTan[((1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^
2]])*Log[-(((1 + a)*(1 + a - I*Sqrt[-1 + a^2])*(-1 + Tanh[ArcSech[a + b*x]
/2]))/(a*(-1 + a + I*Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2])))] - (ArcCos[
a^(-1)] - 2*ArcTan[((1 + a)*Tanh[ArcSech[a + b*x]/2])/Sqrt[-1 + a^2]])*Log[
((-1 + a)*(1 + a + I*Sqrt[-1 + a^2])*(1 + Tanh[ArcSech[a + b*x]/2]))/(a*(-1
+ a + I*Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2]))] + I*(PolyLog[2, ((-1 -
I*Sqrt[-1 + a^2])*(-1 + a - I*Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2]))/(a*
(-1 + a + I*Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2]))] - PolyLog[2, ((I + S
qrt[-1 + a^2])*(-1 + a - I*Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2]))/(a*((-
I)*(-1 + a) + Sqrt[-1 + a^2]*Tanh[ArcSech[a + b*x]/2])))])))/(-1 + a^2)^(3/2))

```

Maple [A] (verified)

Time = 1.21 (sec) , antiderivative size = 982, normalized size of antiderivative = 1.83

method	result
derivativedivides	$b^2 \left(-\frac{\operatorname{arcsech}(bx+a) \left(2 \operatorname{arcsech}(bx+a) a^3 (bx+a) - \operatorname{arcsech}(bx+a) a^2 (bx+a)^2 - 2 \sqrt{-\frac{bx+a-1}{bx+a}} \sqrt{\frac{bx+a+1}{bx+a}} a^2 (bx+a) + 2 \sqrt{\frac{bx+a-1}{bx+a}} \sqrt{\frac{bx+a+1}{bx+a}} a^2 (bx+a) \right)}{2a^2(a^2)}$
default	$b^2 \left(-\frac{\operatorname{arcsech}(bx+a) \left(2 \operatorname{arcsech}(bx+a) a^3 (bx+a) - \operatorname{arcsech}(bx+a) a^2 (bx+a)^2 - 2 \sqrt{-\frac{bx+a-1}{bx+a}} \sqrt{\frac{bx+a+1}{bx+a}} a^2 (bx+a) + 2 \sqrt{\frac{bx+a-1}{bx+a}} \sqrt{\frac{bx+a+1}{bx+a}} a^2 (bx+a) \right)}{2a^2(a^2)}$

[In] `int(arcsech(b*x+a)^2/x^3,x,method=_RETURNVERBOSE)`

[Out]
$$b^2 \cdot \left(-\frac{1}{2} \operatorname{arcsech}(bx+a) \cdot \left(2 \operatorname{arcsech}(bx+a) \cdot a^3 (bx+a) - \operatorname{arcsech}(bx+a) \cdot a^2 (bx+a)^2 - 2 \cdot \left(-\frac{bx+a-1}{bx+a} \right)^{1/2} \cdot \left(\frac{bx+a+1}{bx+a} \right)^{1/2} \cdot a^2 (bx+a) + 2 \cdot \left(-\frac{bx+a-1}{bx+a} \right)^{1/2} \cdot \left(\frac{bx+a+1}{bx+a} \right)^{1/2} \cdot a \cdot (bx+a)^2 - 2 \operatorname{arcsech}(bx+a) \cdot a \cdot (bx+a) + \operatorname{arcsech}(bx+a) \cdot (bx+a)^2 + 2 \cdot a^2 - 4 \cdot (bx+a) \cdot a + 2 \cdot (bx+a)^2 \right) / a^2 / (a^2 - 1) / b^2 / x^2 + 2/a^2 / (a^2 - 1) \cdot \ln\left(\frac{1}{bx+a} + \left(\frac{1}{bx+a} - 1 \right)^{1/2} \cdot \left(\frac{1}{bx+a} + 1 \right)^{1/2} \right) - 1/a^2 / (a^2 - 1) \cdot \ln\left(a \cdot \left(\frac{1}{bx+a} + \left(\frac{1}{bx+a} - 1 \right)^{1/2} \cdot \left(\frac{1}{bx+a} + 1 \right)^{1/2} \right)^2 + a - 2 / (bx+a) - 2 \cdot \left(\frac{1}{bx+a} - 1 \right)^{1/2} \cdot \left(\frac{1}{bx+a} + 1 \right)^{1/2} \right) + (-a^2 + 1)^{1/2} / a^2 / (a^2 - 1)^2 \cdot \operatorname{arcsech}(bx+a) \cdot \ln\left(\left(-a \cdot \left(\frac{1}{bx+a} + \left(\frac{1}{bx+a} - 1 \right)^{1/2} \cdot \left(\frac{1}{bx+a} + 1 \right)^{1/2} \right) + (-a^2 + 1)^{1/2} + 1 \right) / \left(1 + (-a^2 + 1)^{1/2} \right) \right) - (-a^2 + 1)^{1/2} / a^2 / (a^2 - 1)^2 \cdot \operatorname{arcsech}(bx+a) \cdot \ln\left(\left(a \cdot \left(\frac{1}{bx+a} + \left(\frac{1}{bx+a} - 1 \right)^{1/2} \cdot \left(\frac{1}{bx+a} + 1 \right)^{1/2} \right) + (-a^2 + 1)^{1/2} - 1 \right) / \left(-1 + (-a^2 + 1)^{1/2} \right) \right) + (-a^2 + 1)^{1/2} / a^2 / (a^2 - 1)^2 \cdot \operatorname{dilog}\left(\left(-a \cdot \left(\frac{1}{bx+a} + \left(\frac{1}{bx+a} - 1 \right)^{1/2} \cdot \left(\frac{1}{bx+a} + 1 \right)^{1/2} \right) + (-a^2 + 1)^{1/2} + 1 \right) / \left(1 + (-a^2 + 1)^{1/2} \right) \right) - (-a^2 + 1)^{1/2} / a^2 / (a^2 - 1)^2 \cdot \operatorname{dilog}\left(\left(a \cdot \left(\frac{1}{bx+a} + \left(\frac{1}{bx+a} - 1 \right)^{1/2} \cdot \left(\frac{1}{bx+a} + 1 \right)^{1/2} \right) + (-a^2 + 1)^{1/2} - 1 \right) / \left(-1 + (-a^2 + 1)^{1/2} \right) \right) + 2 \cdot \left(-a^2 + 1 \right)^{1/2} / (a^2 - 1)^2 \cdot \operatorname{arcsech}(bx+a) \cdot \ln\left(\left(-a \cdot \left(\frac{1}{bx+a} + \left(\frac{1}{bx+a} - 1 \right)^{1/2} \cdot \left(\frac{1}{bx+a} + 1 \right)^{1/2} \right) + (-a^2 + 1)^{1/2} + 1 \right) / \left(1 + (-a^2 + 1)^{1/2} \right) \right) + 2 \cdot \left(-a^2 + 1 \right)^{1/2} / (a^2 - 1)^2 \cdot \operatorname{arcsech}(bx+a) \cdot \ln\left(\left(a \cdot \left(\frac{1}{bx+a} + \left(\frac{1}{bx+a} - 1 \right)^{1/2} \cdot \left(\frac{1}{bx+a} + 1 \right)^{1/2} \right) + (-a^2 + 1)^{1/2} - 1 \right) / \left(-1 + (-a^2 + 1)^{1/2} \right) \right) - 2 \cdot \left(-a^2 + 1 \right)^{1/2} / (a^2 - 1)^2 \cdot \operatorname{dilog}\left(\left(-a \cdot \left(\frac{1}{bx+a} + \left(\frac{1}{bx+a} - 1 \right)^{1/2} \cdot \left(\frac{1}{bx+a} + 1 \right)^{1/2} \right) + (-a^2 + 1)^{1/2} + 1 \right) / \left(1 + (-a^2 + 1)^{1/2} \right) \right) + \left(-a^2 + 1 \right)^{1/2} + 1 \right) / \left(1 + (-a^2 + 1)^{1/2} \right) \right) + 2 \cdot \left(-a^2 + 1 \right)^{1/2} / (a^2 - 1)^2 \cdot \operatorname{dilog}\left(\left(a \cdot \left(\frac{1}{bx+a} + \left(\frac{1}{bx+a} - 1 \right)^{1/2} \cdot \left(\frac{1}{bx+a} + 1 \right)^{1/2} \right) + (-a^2 + 1)^{1/2} - 1 \right) / \left(-1 + (-a^2 + 1)^{1/2} \right) \right) \right)$$

Fricas [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{arsech}(bx + a)^2}{x^3} dx$$

[In] integrate(arcsech(b*x+a)^2/x^3,x, algorithm="fricas")

[Out] integral(arcsech(b*x + a)^2/x^3, x)

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{asech}^2(a + bx)}{x^3} dx$$

[In] integrate(asech(b*x+a)**2/x**3,x)

[Out] Integral(asech(a + b*x)**2/x**3, x)

Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{arsech}(bx + a)^2}{x^3} dx$$

[In] integrate(arcsech(b*x+a)^2/x^3,x, algorithm="maxima")

[Out]
$$-1/2*\log(\sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1})*b*x + \sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1}*a + b*x + a)^2/x^2 - \operatorname{integrate}(-4*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*\sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1}*\log(b*x + a)^2 + 4*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*\log(b*x + a)^2 + (b^3*x^3 + 2*a*b^2*x^2 + (a^2*b - b)*x - 4*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*\log(b*x + a) - (2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*\sqrt{b*x + a + 1}*\log(b*x + a) - (2*b^3*x^3 + 4*a*b^2*x^2 + (2*a^2*b - b)*x - 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*\log(b*x + a))*\sqrt{b*x + a + 1})*\sqrt{-b*x - a + 1})*\log(\sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1})*b*x + \sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1}*a + b*x + a))/ (b^3*x^6 + 3*a*b^2*x^5 + (3*a^2*b - b)*x^4 + (a^3 - a)*x^3 + (b^3*x^6 + 3*a*b^2*x^5 + (3*a^2*b - b)*x^4 + (a^3 - a)*x^3)*\sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1}), x)$$

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{ar} \operatorname{sech}(bx + a)^2}{x^3} dx$$

[In] integrate(arcsech(b*x+a)^2/x^3,x, algorithm="giac")

[Out] integrate(arcsech(b*x + a)^2/x^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)^2}{x^3} dx$$

[In] int(acosh(1/(a + b*x))^2/x^3,x)

[Out] int(acosh(1/(a + b*x))^2/x^3, x)

3.15 $\int x \operatorname{sech}^{-1}(a + bx)^3 dx$

Optimal result	154
Rubi [A] (verified)	155
Mathematica [A] (verified)	160
Maple [F]	161
Fricas [F]	161
Sympy [F]	161
Maxima [F]	161
Giac [F]	162
Mupad [F(-1)]	162

Optimal result

Integrand size = 10, antiderivative size = 260

$$\begin{aligned}
 \int x \operatorname{sech}^{-1}(a + bx)^3 dx = & -\frac{3 \operatorname{sech}^{-1}(a + bx)^2}{2b^2} - \frac{3 \sqrt{\frac{1-a-bx}{1+a+bx}} (1+a+bx) \operatorname{sech}^{-1}(a + bx)^2}{2b^2} \\
 & - \frac{a^2 \operatorname{sech}^{-1}(a + bx)^3}{2b^2} + \frac{1}{2} x^2 \operatorname{sech}^{-1}(a + bx)^3 \\
 & + \frac{6a \operatorname{sech}^{-1}(a + bx)^2 \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
 & + \frac{3 \operatorname{sech}^{-1}(a + bx) \log\left(1 + e^{2 \operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
 & - \frac{6ia \operatorname{sech}^{-1}(a + bx) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
 & + \frac{6ia \operatorname{sech}^{-1}(a + bx) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
 & + \frac{3 \operatorname{PolyLog}\left(2, -e^{2 \operatorname{sech}^{-1}(a+bx)}\right)}{2b^2} \\
 & + \frac{6ia \operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
 & - \frac{6ia \operatorname{PolyLog}\left(3, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2}
 \end{aligned}$$

[Out] $-3/2*\operatorname{arcsech}(b*x+a)^2/b^2-1/2*a^2*\operatorname{arcsech}(b*x+a)^3/b^2+1/2*x^2*\operatorname{arcsech}(b*x+a)^3+6*a*\operatorname{arcsech}(b*x+a)^2*\arctan(1/(b*x+a)+(1/(b*x+a)-1)^{(1/2)}*(1/(b*x+a)+1)^{(1/2)})/b^2+3*\operatorname{arcsech}(b*x+a)*\ln(1+(1/(b*x+a)+(1/(b*x+a)-1)^{(1/2)}*(1/(b*x+a)+1)^{(1/2)}))^2/b^2-6*I*a*\operatorname{arcsech}(b*x+a)*\operatorname{polylog}(2,-I*(1/(b*x+a)+(1/(b*x+a)-1)^{(1/2)}*(1/(b*x+a)+1)^{(1/2)}))$

$$\begin{aligned} & 1)^{(1/2)} * (1/(b*x+a)+1)^{(1/2)}) / b^2 + 6*I*a*arcsech(b*x+a)*polylog(2, I*(1/(b*x+a) \\ & + (1/(b*x+a)-1)^{(1/2)} * (1/(b*x+a)+1)^{(1/2)}) / b^2 + 3/2*polylog(2, -(1/(b*x+a) \\ & + (1/(b*x+a)-1)^{(1/2)} * (1/(b*x+a)+1)^{(1/2)})^2) / b^2 + 6*I*a*polylog(3, -I*(1/(b*x \\ & + a) + (1/(b*x+a)-1)^{(1/2)} * (1/(b*x+a)+1)^{(1/2)}) / b^2 - 6*I*a*polylog(3, I*(1/(b*x \\ & + a) + (1/(b*x+a)-1)^{(1/2)} * (1/(b*x+a)+1)^{(1/2)}) / b^2 - 3/2*(b*x+a+1)*arcsech(b*x \\ & + a)^2 * ((-b*x-a+1)/(b*x+a+1))^{(1/2)} / b^2 \end{aligned}$$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 1.200$, Rules used = {6456, 5576, 4275, 4265, 2611, 2320, 6724, 4269, 3799, 2221, 2317, 2438}

$$\begin{aligned} \int x \operatorname{sech}^{-1}(a+bx)^3 dx = & -\frac{a^2 \operatorname{sech}^{-1}(a+bx)^3}{2b^2} + \frac{6a \operatorname{sech}^{-1}(a+bx)^2 \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\ & - \frac{6ia \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\ & + \frac{6ia \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\ & + \frac{3 \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right)}{2b^2} \\ & + \frac{6ia \operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\ & - \frac{6ia \operatorname{PolyLog}\left(3, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\ & - \frac{3\sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1)\operatorname{sech}^{-1}(a+bx)^2}{2b^2} - \frac{3\operatorname{sech}^{-1}(a+bx)^2}{2b^2} \\ & + \frac{3\operatorname{sech}^{-1}(a+bx) \log\left(e^{2\operatorname{sech}^{-1}(a+bx)} + 1\right)}{b^2} + \frac{1}{2}x^2 \operatorname{sech}^{-1}(a+bx)^3 \end{aligned}$$

[In] Int[x*ArcSech[a + b*x]^3, x]

[Out] $(-3*\operatorname{ArcSech}[a + b*x]^2)/(2*b^2) - (3*\sqrt{[(1 - a - b*x)/(1 + a + b*x)]}*(1 + a + b*x)*\operatorname{ArcSech}[a + b*x]^2)/(2*b^2) - (a^2*\operatorname{ArcSech}[a + b*x]^3)/(2*b^2) + (x^2*\operatorname{ArcSech}[a + b*x]^3)/2 + (6*a*\operatorname{ArcSech}[a + b*x]^2*\operatorname{ArcTan}[E^{\operatorname{ArcSech}[a + b*x]}])/b^2 + (3*\operatorname{ArcSech}[a + b*x]*\operatorname{Log}[1 + E^{(2*\operatorname{ArcSech}[a + b*x])}])/b^2 - ((6*I)*a*\operatorname{ArcSech}[a + b*x]*\operatorname{PolyLog}[2, (-I)*E^{\operatorname{ArcSech}[a + b*x]}])/b^2 + ((6*I)*a*\operatorname{ArcSech}[a + b*x]*\operatorname{PolyLog}[2, I*E^{\operatorname{ArcSech}[a + b*x]}])/b^2 + (3*\operatorname{PolyLog}[2, -E^{(2*\operatorname{ArcSech}[a + b*x])}])/(2*b^2) + ((6*I)*a*\operatorname{PolyLog}[3, (-I)*E^{\operatorname{ArcSech}[a + b*x]}])/b^2 - ((6*I)*a*\operatorname{PolyLog}[3, I*E^{\operatorname{ArcSech}[a + b*x]}])/b^2$

Rule 2221

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)
))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol]
:= Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*
(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2611

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)
*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m
- 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3799

```
Int[(((c_) + (d_)*(x_))^(m_))*tan[(e_) + (Complex[0, fz_])*(f_)*(x_)], x
_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c
+ d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x))))], x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

Rule 4265

```
Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_
))^m, x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(
I*k*Pi)]/(f*fz*I)), x] + (-Dist[d*(m/(f*fz*I)), Int[(c + d*x)^(m - 1)*Log[1
```

- $E^{(-I)*e + f*fz*x}/E^{(I*k*Pi)}$, x], x] + Dist[d*(m/(f*fz*I)), Int[(c + d*x)^(m - 1)*Log[1 + E^{(-I)*e + f*fz*x}/E^{(I*k*Pi)}], x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]

Rule 4269

Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

Rule 4275

Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]

Rule 5576

Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sech[(c_.) + (d_.)*(x_)])^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rule 6456

Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Tanh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rule 6724

Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int x^3 \text{sech}(x)(-a + \text{sech}(x)) \tanh(x) dx, x, \text{sech}^{-1}(a + bx)\right)}{b^2} \\ &= \frac{1}{2}x^2 \text{sech}^{-1}(a + bx)^3 - \frac{3\text{Subst}\left(\int x^2(-a + \text{sech}(x))^2 dx, x, \text{sech}^{-1}(a + bx)\right)}{2b^2} \\ &= \frac{1}{2}x^2 \text{sech}^{-1}(a + bx)^3 - \frac{3\text{Subst}\left(\int (a^2x^2 - 2ax^2 \text{sech}(x) + x^2 \text{sech}^2(x)) dx, x, \text{sech}^{-1}(a + bx)\right)}{2b^2} \end{aligned}$$

$$\begin{aligned}
&= -\frac{a^2 \operatorname{sech}^{-1}(a+bx)^3}{2b^2} + \frac{1}{2} x^2 \operatorname{sech}^{-1}(a+bx)^3 \\
&\quad - \frac{3 \operatorname{Subst}\left(\int x^2 \operatorname{sech}^2(x) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{2b^2} \\
&\quad + \frac{(3a) \operatorname{Subst}\left(\int x^2 \operatorname{sech}(x) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{b^2} \\
&= -\frac{3\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx) \operatorname{sech}^{-1}(a+bx)^2}{2b^2} - \frac{a^2 \operatorname{sech}^{-1}(a+bx)^3}{2b^2} \\
&\quad + \frac{1}{2} x^2 \operatorname{sech}^{-1}(a+bx)^3 + \frac{6a \operatorname{sech}^{-1}(a+bx)^2 \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad + \frac{3 \operatorname{Subst}\left(\int x \tanh(x) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{b^2} \\
&\quad - \frac{(6ia) \operatorname{Subst}\left(\int x \log(1-ie^x) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{b^2} \\
&\quad + \frac{(6ia) \operatorname{Subst}\left(\int x \log(1+ie^x) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{b^2} \\
&= -\frac{3 \operatorname{sech}^{-1}(a+bx)^2}{2b^2} - \frac{3\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx) \operatorname{sech}^{-1}(a+bx)^2}{2b^2} - \frac{a^2 \operatorname{sech}^{-1}(a+bx)^3}{2b^2} \\
&\quad + \frac{1}{2} x^2 \operatorname{sech}^{-1}(a+bx)^3 + \frac{6a \operatorname{sech}^{-1}(a+bx)^2 \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad - \frac{6ia \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad + \frac{6ia \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad + \frac{6 \operatorname{Subst}\left(\int \frac{e^{2x} x}{1+e^{2x}} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{b^2} \\
&\quad + \frac{(6ia) \operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, -ie^x\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{b^2} \\
&\quad - \frac{(6ia) \operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, ie^x\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{b^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3\operatorname{sech}^{-1}(a+bx)^2}{2b^2} - \frac{3\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)^2}{2b^2} - \frac{a^2\operatorname{sech}^{-1}(a+bx)^3}{2b^2} \\
&\quad + \frac{1}{2}x^2\operatorname{sech}^{-1}(a+bx)^3 + \frac{6a\operatorname{sech}^{-1}(a+bx)^2 \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad + \frac{3\operatorname{sech}^{-1}(a+bx) \log\left(1+e^{2\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad - \frac{6ia\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad + \frac{6ia\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad - \frac{3\operatorname{Subst}\left(\int \log(1+e^{2x}) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{b^2} \\
&\quad + \frac{(6ia)\operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(2, -ix)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad - \frac{(6ia)\operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(2, ix)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&= -\frac{3\operatorname{sech}^{-1}(a+bx)^2}{2b^2} - \frac{3\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)^2}{2b^2} - \frac{a^2\operatorname{sech}^{-1}(a+bx)^3}{2b^2} \\
&\quad + \frac{1}{2}x^2\operatorname{sech}^{-1}(a+bx)^3 + \frac{6a\operatorname{sech}^{-1}(a+bx)^2 \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad + \frac{3\operatorname{sech}^{-1}(a+bx) \log\left(1+e^{2\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad - \frac{6ia\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad + \frac{6ia\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} + \frac{6ia \operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&\quad - \frac{6ia \operatorname{PolyLog}\left(3, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} - \frac{3\operatorname{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2\operatorname{sech}^{-1}(a+bx)}\right)}{2b^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3\operatorname{sech}^{-1}(a+bx)^2}{2b^2} - \frac{3\sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)^2}{2b^2} - \frac{a^2\operatorname{sech}^{-1}(a+bx)^3}{2b^2} \\
&+ \frac{1}{2}x^2\operatorname{sech}^{-1}(a+bx)^3 + \frac{6a\operatorname{sech}^{-1}(a+bx)^2\arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&+ \frac{3\operatorname{sech}^{-1}(a+bx)\log\left(1+e^{2\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&- \frac{6ia\operatorname{sech}^{-1}(a+bx)\operatorname{PolyLog}\left(2,-ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} \\
&+ \frac{6ia\operatorname{sech}^{-1}(a+bx)\operatorname{PolyLog}\left(2,ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} + \frac{3\operatorname{PolyLog}\left(2,-e^{2\operatorname{sech}^{-1}(a+bx)}\right)}{2b^2} \\
&+ \frac{6ia\operatorname{PolyLog}\left(3,-ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2} - \frac{6ia\operatorname{PolyLog}\left(3,ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.65 (sec) , antiderivative size = 254, normalized size of antiderivative = 0.98

$$\int x\operatorname{sech}^{-1}(a+bx)^3 dx$$

$$= \frac{-3\sqrt{-\frac{-1+a+bx}{1+a+bx}}(1+a+bx)\operatorname{sech}^{-1}(a+bx)^2 - 2a(a+bx)\operatorname{sech}^{-1}(a+bx)^3 + (a+bx)^2\operatorname{sech}^{-1}(a+bx)^3 + 3\operatorname{sech}^{-1}(a+bx)^3}{2b^2}$$

[In] Integrate[x*ArcSech[a + b*x]^3,x]

[Out] $(-3\sqrt{-((-1+a+b*x)/(1+a+b*x))}*(1+a+b*x)*\operatorname{ArcSech}[a+b*x]^2 - 2*a*(a+b*x)*\operatorname{ArcSech}[a+b*x]^3 + (a+b*x)^2*\operatorname{ArcSech}[a+b*x]^3 + 3*\operatorname{ArcSech}[a+b*x]*(\operatorname{ArcSech}[a+b*x] + 2*\log[1+E^{(-2*\operatorname{ArcSech}[a+b*x])}])) - 3*\operatorname{PolyLog}[2,-E^{(-2*\operatorname{ArcSech}[a+b*x])}] + (6*I)*a*(-(\operatorname{ArcSech}[a+b*x]^2*(\log[1-I/E^{\operatorname{ArcSech}[a+b*x]}] - \log[1+I/E^{\operatorname{ArcSech}[a+b*x]}])) - 2*\operatorname{ArcSech}[a+b*x]*(\operatorname{PolyLog}[2,(-I)/E^{\operatorname{ArcSech}[a+b*x]}] - \operatorname{PolyLog}[2,I/E^{\operatorname{ArcSech}[a+b*x]}]) - 2*\operatorname{PolyLog}[3,(-I)/E^{\operatorname{ArcSech}[a+b*x]}] + 2*\operatorname{PolyLog}[3,I/E^{\operatorname{ArcSech}[a+b*x]}])))/(2*b^2)$

Maple [F]

$$\int x \operatorname{arcsech}(bx + a)^3 dx$$

```
[In] int(x*arcsech(b*x+a)^3,x)
```

```
[Out] int(x*arcsech(b*x+a)^3,x)
```

Fricas [F]

$$\int x \operatorname{sech}^{-1}(a + bx)^3 dx = \int x \operatorname{arsech}(bx + a)^3 dx$$

```
[In] integrate(x*arcsech(b*x+a)^3,x, algorithm="fricas")
```

```
[Out] integral(x*arcsech(b*x + a)^3, x)
```

Sympy [F]

$$\int x \operatorname{sech}^{-1}(a + bx)^3 dx = \int x \operatorname{asech}^3(a + bx) dx$$

```
[In] integrate(x*asech(b*x+a)**3,x)
```

```
[Out] Integral(x*asech(a + b*x)**3, x)
```

Maxima [F]

$$\int x \operatorname{sech}^{-1}(a + bx)^3 dx = \int x \operatorname{arsech}(bx + a)^3 dx$$

```
[In] integrate(x*arcsech(b*x+a)^3,x, algorithm="maxima")
```

```
[Out] 1/2*x^2*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^3 - integrate(1/2*(16*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^3 + 16*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*log(b*x + a)^3 + 3*(b^3*x^4 + 2*a*b^2*x^3 + (a^2*b - b)*x^2 + 4*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*log(b*x + a) + (2*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*sqrt(b*x + a + 1)*log(b*x + a) + (2*b^3*x^4 + 4*a*b^2*x^3 + (2*a^2*b - b)*x^2 + 2*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*log(b*x + a))*sqrt(b*x + a + 1)*sqrt
```

$(-b*x - a + 1))*\log(\sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1}*b*x + \sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1}*a + b*x + a)^2 - 24*((b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*\sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1}*\log(b*x + a)^2 + (b^3*x^4 + 3*a*b^2*x^3 + (3*a^2*b - b)*x^2 + (a^3 - a)*x)*\log(b*x + a)^2)*\log(\sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1}*b*x + \sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1}*a + b*x + a))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*\sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1} + (3*a^2*b - b)*x - a), x)$

Giac [F]

$$\int x \operatorname{sech}^{-1}(a + bx)^3 dx = \int x \operatorname{ar} \operatorname{sech}(bx + a)^3 dx$$

[In] integrate(x*arcsech(b*x+a)^3,x, algorithm="giac")

[Out] integrate(x*arcsech(b*x + a)^3, x)

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{sech}^{-1}(a + bx)^3 dx = \int x \operatorname{acosh}\left(\frac{1}{a + bx}\right)^3 dx$$

[In] int(x*acosh(1/(a + b*x))^3,x)

[Out] int(x*acosh(1/(a + b*x))^3, x)

3.16 $\int \operatorname{sech}^{-1}(a + bx)^3 dx$

Optimal result	163
Rubi [A] (verified)	164
Mathematica [A] (verified)	167
Maple [F]	167
Fricas [F]	167
Sympy [F]	167
Maxima [F]	168
Giac [F]	168
Mupad [F(-1)]	168

Optimal result

Integrand size = 8, antiderivative size = 136

$$\int \operatorname{sech}^{-1}(a + bx)^3 dx = \frac{(a + bx)\operatorname{sech}^{-1}(a + bx)^3}{b} - \frac{6\operatorname{sech}^{-1}(a + bx)^2 \arctan\left(e^{\operatorname{sech}^{-1}(a + bx)}\right)}{b}$$

$$+ \frac{6i\operatorname{sech}^{-1}(a + bx) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b}$$

$$- \frac{6i\operatorname{sech}^{-1}(a + bx) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b}$$

$$- \frac{6i \operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b} + \frac{6i \operatorname{PolyLog}\left(3, ie^{\operatorname{sech}^{-1}(a + bx)}\right)}{b}$$

```
[Out] (b*x+a)*arcsech(b*x+a)^3/b-6*arcsech(b*x+a)^2*arctan(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/b+6*I*arcsech(b*x+a)*polylog(2,-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b-6*I*arcsech(b*x+a)*polylog(2,I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b-6*I*polylog(3,-I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b+6*I*polylog(3,I*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/b
```

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {6450, 6414, 5526, 4265, 2611, 2320, 6724}

$$\int \operatorname{sech}^{-1}(a + bx)^3 dx = -\frac{6\operatorname{sech}^{-1}(a + bx)^2 \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} + \frac{6i\operatorname{sech}^{-1}(a + bx) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} - \frac{6i\operatorname{sech}^{-1}(a + bx) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} - \frac{6i \operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} + \frac{6i \operatorname{PolyLog}\left(3, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} + \frac{(a + bx)\operatorname{sech}^{-1}(a + bx)^3}{b}$$

[In] Int[ArcSech[a + b*x]^3,x]

[Out] ((a + b*x)*ArcSech[a + b*x]^3)/b - (6*ArcSech[a + b*x]^2*ArcTan[E^ArcSech[a + b*x]])/b + ((6*I)*ArcSech[a + b*x]*PolyLog[2, (-I)*E^ArcSech[a + b*x]])/b - ((6*I)*ArcSech[a + b*x]*PolyLog[2, I*E^ArcSech[a + b*x]])/b - ((6*I)*PolyLog[3, (-I)*E^ArcSech[a + b*x]])/b + ((6*I)*PolyLog[3, I*E^ArcSech[a + b*x]])/b

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2611

Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(- (f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x))))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x))))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 4265

Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/E^((

$I*k*\text{Pi}]/(f*fz*I)), x] + (-\text{Dist}[d*(m/(f*fz*I)), \text{Int}[(c + d*x)^{(m-1)}*\text{Log}[1 - E^{((-I)*e + f*fz*x)/E^{(I*k*\text{Pi})}}], x], x] + \text{Dist}[d*(m/(f*fz*I)), \text{Int}[(c + d*x)^{(m-1)}*\text{Log}[1 + E^{((-I)*e + f*fz*x)/E^{(I*k*\text{Pi})}}], x], x]) /; \text{FreeQ}\{c, d, e, f, fz\}, x] \&\& \text{IntegerQ}[2*k] \&\& \text{IGtQ}[m, 0]$

Rule 5526

$\text{Int}[(x_)^{(m_*)}*\text{Sech}[(a_*) + (b_*)*(x_)^{(n_*)}]^{(p_*)}*\text{Tanh}[(a_*) + (b_*)*(x_)^{(n_*)}]^{(q_*)}, x_Symbol] :> \text{Simp}[(-x^{(m-n+1)})*(\text{Sech}[a + b*x^n]^p/(b*n*p)), x] + \text{Dist}[(m-n+1)/(b*n*p), \text{Int}[x^{(m-n)}*\text{Sech}[a + b*x^n]^p, x], x] /; \text{FreeQ}\{a, b, p\}, x] \&\& \text{RationalQ}[m] \&\& \text{IntegerQ}[n] \&\& \text{GeQ}[m-n, 0] \&\& \text{EqQ}[q, 1]$

Rule 6414

$\text{Int}[(a_*) + \text{ArcSech}[(c_*)*(x_)]*(b_*)^{(n_*)}, x_Symbol] :> \text{Dist}[-c^{(-1)}, \text{Subst}[\text{Int}[(a + b*x)^n*\text{Sech}[x]*\text{Tanh}[x], x], x, \text{ArcSech}[c*x]], x] /; \text{FreeQ}\{a, b, c, n\}, x] \&\& \text{IGtQ}[n, 0]$

Rule 6450

$\text{Int}[(a_*) + \text{ArcSech}[(c_*) + (d_*)*(x_)]*(b_*)^{(p_*)}, x_Symbol] :> \text{Dist}[1/d, \text{Subst}[\text{Int}[(a + b*\text{ArcSech}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{IGtQ}[p, 0]$

Rule 6724

$\text{Int}[\text{PolyLog}[n_*, (c_*)*((a_*) + (b_*)*(x_))^{(p_*)}]/((d_*) + (e_*)*(x_)), x_Symbol] :> \text{Simp}[\text{PolyLog}[n + 1, c*(a + b*x)^p/(e*p), x] /; \text{FreeQ}\{a, b, c, d, e, n, p\}, x] \&\& \text{EqQ}[b*d, a*e]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}(\int \text{sech}^{-1}(x)^3 dx, x, a + bx)}{b} \\
 &= -\frac{\text{Subst}(\int x^3 \text{sech}(x) \tanh(x) dx, x, \text{sech}^{-1}(a + bx))}{b} \\
 &= \frac{(a + bx)\text{sech}^{-1}(a + bx)^3}{b} - \frac{3\text{Subst}(\int x^2 \text{sech}(x) dx, x, \text{sech}^{-1}(a + bx))}{b} \\
 &= \frac{(a + bx)\text{sech}^{-1}(a + bx)^3}{b} - \frac{6\text{sech}^{-1}(a + bx)^2 \arctan(e^{\text{sech}^{-1}(a + bx)})}{b} \\
 &\quad + \frac{(6i)\text{Subst}(\int x \log(1 - ie^x) dx, x, \text{sech}^{-1}(a + bx))}{b} \\
 &\quad - \frac{(6i)\text{Subst}(\int x \log(1 + ie^x) dx, x, \text{sech}^{-1}(a + bx))}{b}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{(a+bx)\operatorname{sech}^{-1}(a+bx)^3}{b} - \frac{6\operatorname{sech}^{-1}(a+bx)^2 \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} \\
&+ \frac{6i\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} \\
&- \frac{6i\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} \\
&- \frac{(6i)\operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, -ie^x\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{b} \\
&+ \frac{(6i)\operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, ie^x\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{b} \\
&= \frac{(a+bx)\operatorname{sech}^{-1}(a+bx)^3}{b} - \frac{6\operatorname{sech}^{-1}(a+bx)^2 \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} \\
&+ \frac{6i\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} \\
&- \frac{6i\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} \\
&- \frac{(6i)\operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(2, -ix)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} \\
&+ \frac{(6i)\operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(2, ix)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} \\
&= \frac{(a+bx)\operatorname{sech}^{-1}(a+bx)^3}{b} - \frac{6\operatorname{sech}^{-1}(a+bx)^2 \arctan\left(e^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} \\
&+ \frac{6i\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} \\
&- \frac{6i\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} \\
&- \frac{6i \operatorname{PolyLog}\left(3, -ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b} + \frac{6i \operatorname{PolyLog}\left(3, ie^{\operatorname{sech}^{-1}(a+bx)}\right)}{b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.12

$$\int \operatorname{sech}^{-1}(a+bx)^3 dx = \frac{(a+bx)\operatorname{sech}^{-1}(a+bx)^3}{b} - \frac{3i\left(-\operatorname{sech}^{-1}(a+bx)^2\left(\log\left(1-ie^{-\operatorname{sech}^{-1}(a+bx)}\right) - \log\left(1+ie^{-\operatorname{sech}^{-1}(a+bx)}\right)\right) - 2\operatorname{sech}^{-1}(a+bx)\left(\operatorname{PolyLog}\right)}{b}$$

[In] Integrate[ArcSech[a + b*x]^3,x]

[Out] ((a + b*x)*ArcSech[a + b*x]^3)/b - ((3*I)*(-(ArcSech[a + b*x]^2*(Log[1 - I/E^ArcSech[a + b*x]] - Log[1 + I/E^ArcSech[a + b*x]])) - 2*ArcSech[a + b*x]*(PolyLog[2, (-I)/E^ArcSech[a + b*x]] - PolyLog[2, I/E^ArcSech[a + b*x]])) - 2*(PolyLog[3, (-I)/E^ArcSech[a + b*x]] - PolyLog[3, I/E^ArcSech[a + b*x]]))/b

Maple [F]

$$\int \operatorname{arcsech}(bx+a)^3 dx$$

[In] int(arcsech(b*x+a)^3,x)

[Out] int(arcsech(b*x+a)^3,x)

Fricas [F]

$$\int \operatorname{sech}^{-1}(a+bx)^3 dx = \int \operatorname{arsech}(bx+a)^3 dx$$

[In] integrate(arcsech(b*x+a)^3,x, algorithm="fricas")

[Out] integral(arcsech(b*x + a)^3, x)

Sympy [F]

$$\int \operatorname{sech}^{-1}(a+bx)^3 dx = \int \operatorname{asech}^3(a+bx) dx$$

[In] integrate(asech(b*x+a)**3,x)

[Out] Integral(asech(a + b*x)**3, x)

Maxima [F]

$$\int \operatorname{sech}^{-1}(a + bx)^3 dx = \int \operatorname{arsech}(bx + a)^3 dx$$

[In] integrate(arcsech(b*x+a)^3,x, algorithm="maxima")

[Out] x*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^3 - integrate((8*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^3 + 8*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a)^3 + 3*(b^3*x^3 + 2*a*b^2*x^2 + (a^2*b - b)*x + 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a) + ((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*log(b*x + a) + (2*b^3*x^3 + 4*a*b^2*x^2 + (2*a^2*b - b)*x + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a))*sqrt(b*x + a + 1))*sqrt(-b*x - a + 1)*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^2 - 12*((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^2 + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a)^2)*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1) + (3*a^2*b - b)*x - a), x)

Giac [F]

$$\int \operatorname{sech}^{-1}(a + bx)^3 dx = \int \operatorname{arsech}(bx + a)^3 dx$$

[In] integrate(arcsech(b*x+a)^3,x, algorithm="giac")

[Out] integrate(arcsech(b*x + a)^3, x)

Mupad [F(-1)]

Timed out.

$$\int \operatorname{sech}^{-1}(a + bx)^3 dx = \int \operatorname{acosh}\left(\frac{1}{a + bx}\right)^3 dx$$

[In] int(acosh(1/(a + b*x))^3,x)

[Out] int(acosh(1/(a + b*x))^3, x)

3.17 $\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx$

Optimal result	169
Rubi [A] (verified)	170
Mathematica [A] (verified)	178
Maple [F]	179
Fricas [F]	179
Sympy [F]	179
Maxima [F]	179
Giac [F]	180
Mupad [F(-1)]	180

Optimal result

Integrand size = 12, antiderivative size = 378

$$\begin{aligned}
 \int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx &= \operatorname{sech}^{-1}(a+bx)^3 \log \left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
 &\quad + \operatorname{sech}^{-1}(a+bx)^3 \log \left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
 &\quad - \operatorname{sech}^{-1}(a+bx)^3 \log \left(1 + e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
 &\quad + 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
 &\quad + 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
 &\quad - \frac{3}{2} \operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog} \left(2, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
 &\quad - 6\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
 &\quad - 6\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
 &\quad + \frac{3}{2} \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog} \left(3, -e^{2\operatorname{sech}^{-1}(a+bx)} \right) \\
 &\quad + 6 \operatorname{PolyLog} \left(4, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) + 6 \operatorname{PolyLog} \left(4, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
 &\quad - \frac{3}{4} \operatorname{PolyLog} \left(4, -e^{2\operatorname{sech}^{-1}(a+bx)} \right)
 \end{aligned}$$

```
[Out] -arcsech(b*x+a)^3*ln(1+(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)+arcsech(b*x+a)^3*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1-(-a^2+1)^(1/2))+arcsech(b*x+a)^3*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1+(-a^2+1)^(1/2))-3/2*arcsech(b*x+a)^2*polylog(2,-(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)+3*arcsech(b*x+a)^2*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1-(-a^2+1)^(1/2))+3*arcsech(b*x+a)^2*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1+(-a^2+1)^(1/2))+3/2*arcsech(b*x+a)*polylog(3,-(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)-6*arcsech(b*x+a)*polylog(3,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1-(-a^2+1)^(1/2))-6*arcsech(b*x+a)*polylog(3,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1+(-a^2+1)^(1/2))-3/4*polylog(4,-(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)+6*polylog(4,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1-(-a^2+1)^(1/2))+6*polylog(4,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)))/(1+(-a^2+1)^(1/2))
```

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 378, normalized size of antiderivative = 1.00, number of steps used = 20, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules

used = {6456, 5714, 5689, 3799, 2221, 2611, 6744, 2320, 6724, 5681}

$$\begin{aligned}
 \int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx &= 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) \\
 &+ 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) \\
 &- 6\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) \\
 &- 6\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) \\
 &+ 6 \operatorname{PolyLog}\left(4, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) + 6 \operatorname{PolyLog}\left(4, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) \\
 &+ \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) \\
 &+ \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) \\
 &- \frac{3}{2} \operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\
 &+ \frac{3}{2} \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\
 &- \frac{3}{4} \operatorname{PolyLog}\left(4, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\
 &- \operatorname{sech}^{-1}(a+bx)^3 \log\left(e^{2\operatorname{sech}^{-1}(a+bx)} + 1\right)
 \end{aligned}$$

[In] Int[ArcSech[a + b*x]^3/x,x]

[Out] ArcSech[a + b*x]^3*Log[1 - (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])] + ArcSech[a + b*x]^3*Log[1 - (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])] - ArcSech[a + b*x]^3*Log[1 + E^(2*ArcSech[a + b*x])] + 3*ArcSech[a + b*x]^2*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])] + 3*ArcSech[a + b*x]^2*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])] - (3*ArcSech[a + b*x]^2*PolyLog[2, -E^(2*ArcSech[a + b*x])])/2 - 6*ArcSech[a + b*x]*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])] - 6*ArcSech[a + b*x]*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])] + (3*ArcSech[a + b*x]*PolyLog[3, -E^(2*ArcSech[a + b*x])])/2 + 6*PolyLog[4, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])] + 6*PolyLog[4, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])] - (3*PolyLog[4, -E^(2*ArcSech[a + b*x])])/4

Rule 2221

Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] :> Simp

```

[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

```

Rule 2320

```

Int[u_, x_Symbol] :=> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

```

Rule 2611

```

Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] :=> Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

```

Rule 3799

```

Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] :=> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x))), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

```

Rule 5681

```

Int[(((e_.) + (f_.)*(x_))^(m_.)*Sinh[(c_.) + (d_.)*(x_)])/(Cosh[(c_.) + (d_.)*(x_)]*(b_.) + (a_)), x_Symbol] :=> Simp[-(e + f*x)^(m + 1)/(b*f*(m + 1)), x] + (Int[(e + f*x)^m*(E^(c + d*x)/(a - Rt[a^2 - b^2, 2] + b*E^(c + d*x))), x] + Int[(e + f*x)^m*(E^(c + d*x)/(a + Rt[a^2 - b^2, 2] + b*E^(c + d*x))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[a^2 - b^2, 0]

```

Rule 5689

```

Int[(((e_.) + (f_.)*(x_))^(m_.)*Tanh[(c_.) + (d_.)*(x_)])^(n_.)/(Cosh[(c_.) + (d_.)*(x_)]*(b_.) + (a_)), x_Symbol] :=> Dist[1/a, Int[(e + f*x)^m*Tanh[c + d*x]^n, x], x] - Dist[b/a, Int[(e + f*x)^m*Sinh[c + d*x]*(Tanh[c + d*x])^(n - 1)/(a + b*Cosh[c + d*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[n, 0]

```

Rule 5714

```

Int[(((e_.) + (f_.)*(x_))^(m_.)*(F_) [(c_.) + (d_.)*(x_)]^(n_.)*(G_) [(c_.) + (d_.)*(x_)]^(p_.))/(a_ + (b_.)*Sech[(c_.) + (d_.)*(x_)]), x_Symbol] :=> I

```

```

nt[(e + f*x)^m*Cosh[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Cosh[c + d*x
]))], x] /; FreeQ[{a, b, c, d, e, f}, x] && HyperbolicQ[F] && HyperbolicQ[G]
&& IntegersQ[m, n, p]

```

Rule 6456

```

Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(
m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Ta
nh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

```

Rule 6724

```

Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_.))^(p_.)]/((d_.) + (e_.)*(x_.)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]

```

Rule 6744

```

Int[((e_.) + (f_.)*(x_.))^(m_.)*PolyLog[n_, (d_.)*(F_)^((c_.)*((a_.) + (b_.
)*(x_.)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^
(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c,
d, e, f, n, p}, x] && GtQ[m, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{x^3 \operatorname{sech}(x) \tanh(x)}{-a + \operatorname{sech}(x)} dx, x, \operatorname{sech}^{-1}(a + bx)\right) \\
&= -\text{Subst}\left(\int \frac{x^3 \tanh(x)}{1 - a \cosh(x)} dx, x, \operatorname{sech}^{-1}(a + bx)\right) \\
&= -\left(a \text{Subst}\left(\int \frac{x^3 \sinh(x)}{1 - a \cosh(x)} dx, x, \operatorname{sech}^{-1}(a + bx)\right)\right) \\
&\quad - \text{Subst}\left(\int x^3 \tanh(x) dx, x, \operatorname{sech}^{-1}(a + bx)\right) \\
&= -\left(2 \text{Subst}\left(\int \frac{e^{2x} x^3}{1 + e^{2x}} dx, x, \operatorname{sech}^{-1}(a + bx)\right)\right) \\
&\quad - a \text{Subst}\left(\int \frac{e^x x^3}{1 - \sqrt{1 - a^2} - a e^x} dx, x, \operatorname{sech}^{-1}(a + bx)\right) \\
&\quad - a \text{Subst}\left(\int \frac{e^x x^3}{1 + \sqrt{1 - a^2} - a e^x} dx, x, \operatorname{sech}^{-1}(a + bx)\right)
\end{aligned}$$

$$\begin{aligned}
&= \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) + \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad - \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 + e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\
&\quad - 3\operatorname{Subst}\left(\int x^2 \log\left(1 - \frac{ae^x}{1 - \sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right) \\
&\quad - 3\operatorname{Subst}\left(\int x^2 \log\left(1 - \frac{ae^x}{1 + \sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right) \\
&\quad + 3\operatorname{Subst}\left(\int x^2 \log(1 + e^{2x}) dx, x, \operatorname{sech}^{-1}(a+bx)\right) \\
&= \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) + \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad - \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 + e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\
&\quad + 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\
&\quad + 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad - \frac{3}{2}\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\
&\quad + 3\operatorname{Subst}\left(\int x \operatorname{PolyLog}(2, -e^{2x}) dx, x, \operatorname{sech}^{-1}(a+bx)\right) \\
&\quad - 6\operatorname{Subst}\left(\int x \operatorname{PolyLog}\left(2, \frac{ae^x}{1 - \sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right) \\
&\quad - 6\operatorname{Subst}\left(\int x \operatorname{PolyLog}\left(2, \frac{ae^x}{1 + \sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)
\end{aligned}$$

$$\begin{aligned}
&= \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) + \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad - \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 + e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\
&\quad + 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\
&\quad + 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad - \frac{3}{2}\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\
&\quad - 6\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\
&\quad - 6\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad + \frac{3}{2}\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\
&\quad - \frac{3}{2}\operatorname{Subst}\left(\int \operatorname{PolyLog}\left(3, -e^{2x}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right) \\
&\quad + 6\operatorname{Subst}\left(\int \operatorname{PolyLog}\left(3, \frac{ae^x}{1 - \sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right) \\
&\quad + 6\operatorname{Subst}\left(\int \operatorname{PolyLog}\left(3, \frac{ae^x}{1 + \sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)
\end{aligned}$$

$$\begin{aligned}
&= \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) + \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad - \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 + e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\
&\quad + 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\
&\quad + 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad - \frac{3}{2}\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\
&\quad - 6\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\
&\quad - 6\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad + \frac{3}{2}\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\
&\quad - \frac{3}{4}\operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(3, -x)}{x} dx, x, e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\
&\quad + 6\operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(3, \frac{ax}{1-\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right) \\
&\quad + 6\operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(3, \frac{ax}{1+\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)
\end{aligned}$$

$$\begin{aligned}
&= \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) + \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad - \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 + e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\
&\quad + 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\
&\quad + 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad - \frac{3}{2}\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) \\
&\quad - 6\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\
&\quad - 6\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad + \frac{3}{2}\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, -e^{2\operatorname{sech}^{-1}(a+bx)}\right) + 6 \operatorname{PolyLog}\left(4, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\
&\quad + 6 \operatorname{PolyLog}\left(4, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) - \frac{3}{4} \operatorname{PolyLog}\left(4, -e^{2\operatorname{sech}^{-1}(a+bx)}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 384, normalized size of antiderivative = 1.02

$$\begin{aligned}
\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx = & -\frac{1}{2}\operatorname{sech}^{-1}(a+bx)^4 - \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 + e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\
& + \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 + \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}}\right) \\
& + \operatorname{sech}^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
& + \frac{3}{2}\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, -e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\
& + 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, -\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}}\right) \\
& + 3\operatorname{sech}^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
& + \frac{3}{2}\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, -e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\
& - 6\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, -\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}}\right) \\
& - 6\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
& + \frac{3}{4} \operatorname{PolyLog}\left(4, -e^{-2\operatorname{sech}^{-1}(a+bx)}\right) \\
& + 6 \operatorname{PolyLog}\left(4, -\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}}\right) + 6 \operatorname{PolyLog}\left(4, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right)
\end{aligned}$$

```
[In] Integrate[ArcSech[a + b*x]^3/x,x]
```

```
[Out] -1/2*ArcSech[a + b*x]^4 - ArcSech[a + b*x]^3*Log[1 + E^(-2*ArcSech[a + b*x])
] + ArcSech[a + b*x]^3*Log[1 + (a*E^ArcSech[a + b*x])/(-1 + Sqrt[1 - a^2])
] + ArcSech[a + b*x]^3*Log[1 - (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])]
+ (3*ArcSech[a + b*x]^2*PolyLog[2, -E^(-2*ArcSech[a + b*x])])/2 + 3*ArcSech
[a + b*x]^2*PolyLog[2, -((a*E^ArcSech[a + b*x])/(-1 + Sqrt[1 - a^2]))] + 3*
ArcSech[a + b*x]^2*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])] +
(3*ArcSech[a + b*x]*PolyLog[3, -E^(-2*ArcSech[a + b*x])])/2 - 6*ArcSech[a
+ b*x]*PolyLog[3, -((a*E^ArcSech[a + b*x])/(-1 + Sqrt[1 - a^2]))] - 6*ArcSe
ch[a + b*x]*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])] + (3*Pol
yLog[4, -E^(-2*ArcSech[a + b*x])])/4 + 6*PolyLog[4, -((a*E^ArcSech[a + b*x]
)/(-1 + Sqrt[1 - a^2]))] + 6*PolyLog[4, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1
- a^2])]
```

Maple [F]

$$\int \frac{\operatorname{arcsech}(bx+a)^3}{x} dx$$

[In] int(arcsech(b*x+a)^3/x,x)

[Out] int(arcsech(b*x+a)^3/x,x)

Fricas [F]

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx = \int \frac{\operatorname{arsech}(bx+a)^3}{x} dx$$

[In] integrate(arcsech(b*x+a)^3/x,x, algorithm="fricas")

[Out] integral(arcsech(b*x + a)^3/x, x)

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx = \int \frac{\operatorname{asech}^3(a+bx)}{x} dx$$

[In] integrate(asech(b*x+a)**3/x,x)

[Out] Integral(asech(a + b*x)**3/x, x)

Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} dx = \int \frac{\operatorname{arsech}(bx+a)^3}{x} dx$$

[In] integrate(arcsech(b*x+a)^3/x,x, algorithm="maxima")

[Out] integrate(arcsech(b*x + a)^3/x, x)

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x} dx = \int \frac{\operatorname{ar} \operatorname{sech}(bx + a)^3}{x} dx$$

[In] integrate(arcsech(b*x+a)^3/x,x, algorithm="giac")

[Out] integrate(arcsech(b*x + a)^3/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)^3}{x} dx$$

[In] int(acosh(1/(a + b*x))^3/x,x)

[Out] int(acosh(1/(a + b*x))^3/x, x)

3.18 $\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^2} dx$

Optimal result	181
Rubi [A] (verified)	182
Mathematica [C] (warning: unable to verify)	186
Maple [F]	186
Fricas [F]	186
Sympy [F]	187
Maxima [F]	187
Giac [F]	187
Mupad [F(-1)]	188

Optimal result

Integrand size = 12, antiderivative size = 330

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^2} dx = -\frac{b\operatorname{sech}^{-1}(a+bx)^3}{a} - \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} + \frac{3b\operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{3b\operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{6b\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{6b\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{6b \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{6b \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}}$$

```
[Out] -b*arcsech(b*x+a)^3/a-arcsech(b*x+a)^3/x+3*b*arcsech(b*x+a)^2*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-3*b*arcsech(b*x+a)^2*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+6*b*arcsech(b*x+a)*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-6*b*arcsech(b*x+a)*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-6*b*polylog(3,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+6*b*polylog(3,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)
```

Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 330, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6456, 5576, 4276, 3401, 2296, 2221, 2611, 2320, 6724}

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^2} dx = \frac{6b\operatorname{sech}^{-1}(a+bx)\operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{6b\operatorname{sech}^{-1}(a+bx)\operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a\sqrt{1-a^2}} - \frac{6b\operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{6b\operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a\sqrt{1-a^2}} + \frac{3b\operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{3b\operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a\sqrt{1-a^2}} - \frac{b\operatorname{sech}^{-1}(a+bx)^3}{a} - \frac{\operatorname{sech}^{-1}(a+bx)^3}{x}$$

[In] Int[ArcSech[a + b*x]^3/x^2,x]

[Out] -((b*ArcSech[a + b*x]^3)/a) - ArcSech[a + b*x]^3/x + (3*b*ArcSech[a + b*x]^2*Log[1 - (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) - (3*b*ArcSech[a + b*x]^2*Log[1 - (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) + (6*b*ArcSech[a + b*x]*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) - (6*b*ArcSech[a + b*x]*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) - (6*b*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) + (6*b*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2])

Rule 2221

Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] :> Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2296

Int[(((F_)^(u_)*((f_) + (g_)*(x_))^(m_))/((a_) + (b_)*(F_)^(u_) + (c_)*(F_)^(v_)), x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[2*(c/q), Int[

```
(f + g*x)^m*(F^u/(b - q + 2*c*F^u)), x], x] - Dist[2*(c/q), Int[(f + g*x)^m
*(F^u/(b + q + 2*c*F^u)), x], x]] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v,
2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*
(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*(f_.) + (g_.)
*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m
- 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3401

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_.) + (b_.)*sin[(e_.) + Pi*(k_.) + (Comple
x[0, fz_])*(f_.)*(x_)]), x_Symbol] := Dist[2, Int[((c + d*x)^m*(E^((-I)*e +
f*fz*x)/(b + (2*a*E^((-I)*e + f*fz*x))/E^(I*Pi*(k - 1/2)) - (b*E^(2*((-I)*
e + f*fz*x))/E^(2*I*k*Pi)))/E^(I*Pi*(k - 1/2)), x], x] /; FreeQ[{a, b, c,
d, e, f, fz}, x] && IntegerQ[2*k] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]
```

Rule 4276

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)
, x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Si
n[e + f*x]^n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGt
Q[m, 0]
```

Rule 5576

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sech[
(c_.) + (d_.)*(x_)]^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-e
+ f*x)^m*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b
*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /
; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 6456

```
Int[((a_.) + ArcSech[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Ta
```

nh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rule 6724

Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\left(b\text{Subst}\left(\int \frac{x^3 \text{sech}(x) \tanh(x)}{(-a + \text{sech}(x))^2} dx, x, \text{sech}^{-1}(a + bx)\right)\right) \\
 &= -\frac{\text{sech}^{-1}(a + bx)^3}{x} + (3b)\text{Subst}\left(\int \frac{x^2}{-a + \text{sech}(x)} dx, x, \text{sech}^{-1}(a + bx)\right) \\
 &= -\frac{\text{sech}^{-1}(a + bx)^3}{x} + (3b)\text{Subst}\left(\int \left(-\frac{x^2}{a} + \frac{x^2}{a(1 - a \cosh(x))}\right) dx, x, \text{sech}^{-1}(a + bx)\right) \\
 &= -\frac{b\text{sech}^{-1}(a + bx)^3}{a} - \frac{\text{sech}^{-1}(a + bx)^3}{x} + \frac{(3b)\text{Subst}\left(\int \frac{x^2}{1 - a \cosh(x)} dx, x, \text{sech}^{-1}(a + bx)\right)}{a} \\
 &= -\frac{b\text{sech}^{-1}(a + bx)^3}{a} - \frac{\text{sech}^{-1}(a + bx)^3}{x} + \frac{(6b)\text{Subst}\left(\int \frac{e^x x^2}{-a + 2e^x - ae^{2x}} dx, x, \text{sech}^{-1}(a + bx)\right)}{a} \\
 &= -\frac{b\text{sech}^{-1}(a + bx)^3}{a} - \frac{\text{sech}^{-1}(a + bx)^3}{x} \\
 &\quad - \frac{(6b)\text{Subst}\left(\int \frac{e^x x^2}{2 - 2\sqrt{1 - a^2} - 2ae^x} dx, x, \text{sech}^{-1}(a + bx)\right)}{\sqrt{1 - a^2}} \\
 &\quad + \frac{(6b)\text{Subst}\left(\int \frac{e^x x^2}{2 + 2\sqrt{1 - a^2} - 2ae^x} dx, x, \text{sech}^{-1}(a + bx)\right)}{\sqrt{1 - a^2}} \\
 &= -\frac{b\text{sech}^{-1}(a + bx)^3}{a} - \frac{\text{sech}^{-1}(a + bx)^3}{x} + \frac{3b\text{sech}^{-1}(a + bx)^2 \log\left(1 - \frac{ae^{\text{sech}^{-1}(a + bx)}}{1 - \sqrt{1 - a^2}}\right)}{a\sqrt{1 - a^2}} \\
 &\quad - \frac{3b\text{sech}^{-1}(a + bx)^2 \log\left(1 - \frac{ae^{\text{sech}^{-1}(a + bx)}}{1 + \sqrt{1 - a^2}}\right)}{a\sqrt{1 - a^2}} \\
 &\quad - \frac{(6b)\text{Subst}\left(\int x \log\left(1 - \frac{2ae^x}{2 - 2\sqrt{1 - a^2}}\right) dx, x, \text{sech}^{-1}(a + bx)\right)}{a\sqrt{1 - a^2}} \\
 &\quad + \frac{(6b)\text{Subst}\left(\int x \log\left(1 - \frac{2ae^x}{2 + 2\sqrt{1 - a^2}}\right) dx, x, \text{sech}^{-1}(a + bx)\right)}{a\sqrt{1 - a^2}}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{b\operatorname{sech}^{-1}(a+bx)^3}{a} - \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} + \frac{3b\operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{3b\operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad + \frac{6b\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{6b\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{(6b)\operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, \frac{2ae^x}{2-2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a\sqrt{1-a^2}} \\
&\quad + \frac{(6b)\operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, \frac{2ae^x}{2+2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a\sqrt{1-a^2}} \\
&= -\frac{b\operatorname{sech}^{-1}(a+bx)^3}{a} - \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} + \frac{3b\operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{3b\operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad + \frac{6b\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{6b\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{(6b)\operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(2, \frac{ax}{1-\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{a\sqrt{1-a^2}} \\
&\quad + \frac{(6b)\operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(2, \frac{ax}{1+\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{a\sqrt{1-a^2}}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{b\operatorname{sech}^{-1}(a+bx)^3}{a} - \frac{\operatorname{sech}^{-1}(a+bx)^3}{x} + \frac{3b\operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{3b\operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad + \frac{6b\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{6b\operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{6b \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{6b \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}}
\end{aligned}$$

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 69.42 (sec) , antiderivative size = 8527, normalized size of antiderivative = 25.84

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^2} dx = \text{Result too large to show}$$

[In] Integrate[ArcSech[a + b*x]^3/x^2,x]

[Out] Result too large to show

Maple [F]

$$\int \frac{\operatorname{arcsech}(bx+a)^3}{x^2} dx$$

[In] int(arcsech(b*x+a)^3/x^2,x)

[Out] int(arcsech(b*x+a)^3/x^2,x)

Fricas [F]

$$\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^2} dx = \int \frac{\operatorname{arosech}(bx+a)^3}{x^2} dx$$

[In] integrate(arcsech(b*x+a)^3/x^2,x, algorithm="fricas")

[Out] integral(arcsech(b*x + a)^3/x^2, x)

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{asech}^3(a + bx)}{x^2} dx$$

```
[In] integrate(asech(b*x+a)**3/x**2,x)
```

```
[Out] Integral(asech(a + b*x)**3/x**2, x)
```

Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{arsech}(bx + a)^3}{x^2} dx$$

```
[In] integrate(arcsech(b*x+a)^3/x^2,x, algorithm="maxima")
```

```
[Out] -log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^3/x - integrate((8*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^3 + 8*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a)^3 - 3*(b^3*x^3 + 2*a*b^2*x^2 + (a^2*b - b)*x - 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a) - ((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*log(b*x + a) - (2*b^3*x^3 + 4*a*b^2*x^2 + (2*a^2*b - b)*x - (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a))*sqrt(b*x + a + 1))*sqrt(-b*x - a + 1))*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)^2 - 12*((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*log(b*x + a)^2 + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*log(b*x + a)^2)*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a))/(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2 + (b^3*x^5 + 3*a*b^2*x^4 + (3*a^2*b - b)*x^3 + (a^3 - a)*x^2)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)), x)
```

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{arsech}(bx + a)^3}{x^2} dx$$

```
[In] integrate(arcsech(b*x+a)^3/x^2,x, algorithm="giac")
```

```
[Out] integrate(arcsech(b*x + a)^3/x^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)^3}{x^2} dx$$

```
[In] int(acosh(1/(a + b*x))^3/x^2,x)
```

```
[Out] int(acosh(1/(a + b*x))^3/x^2, x)
```

$$3.19 \quad \int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^3} dx$$

Optimal result	190
Rubi [A] (verified)	191
Mathematica [F]	202
Maple [F]	202
Fricas [F]	202
Sympy [F]	202
Maxima [F]	203
Giac [F]	203
Mupad [F(-1)]	203

Optimal result

Integrand size = 12, antiderivative size = 965

$$\begin{aligned}
\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^3} dx = & -\frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2(1-a^2)} + \frac{3b^2 \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx) \operatorname{sech}^{-1}(a+bx)^2}{2a(1-a^2)(a+bx)\left(1-\frac{a}{a+bx}\right)} \\
& + \frac{b^2 \operatorname{sech}^{-1}(a+bx)^3}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)^3}{2x^2} \\
& + \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
& + \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{2a^2(1-a^2)^{3/2}} \\
& - \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2\sqrt{1-a^2}} \\
& + \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
& - \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{2a^2(1-a^2)^{3/2}} \\
& + \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2\sqrt{1-a^2}} \\
& + \frac{3b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
& + \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
& - \frac{6b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2\sqrt{1-a^2}} \\
& + \frac{3b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
& - \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
& + \frac{6b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2\sqrt{1-a^2}} \\
& - \frac{3b^2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} + \frac{6b^2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2\sqrt{1-a^2}} \\
& + \frac{3b^2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} - \frac{6b^2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2\sqrt{1-a^2}}
\end{aligned}$$

```
[Out] -3/2*b^2*arcsech(b*x+a)^2/a^2/(-a^2+1)+1/2*b^2*arcsech(b*x+a)^3/a^2-1/2*arc
sech(b*x+a)^3/x^2+3*b^2*arcsech(b*x+a)*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)
)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a^2/(-a^2+1)+3/2*b^2*arcsech(b*x
+a)^2*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)
^(1/2)))/a^2/(-a^2+1)^(3/2)+3*b^2*arcsech(b*x+a)*ln(1-a*(1/(b*x+a)+(1/(b*x
+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a^2/(-a^2+1)-3/2*b^2*
arcsech(b*x+a)^2*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2)
)/(1+(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(3/2)+3*b^2*polylog(2,a*(1/(b*x+a)+(1/(b*
x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a^2/(-a^2+1)+3*b^2*a
rcsech(b*x+a)*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2
)))/(1-(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(3/2)+3*b^2*polylog(2,a*(1/(b*x+a)+(1/(
b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a^2/(-a^2+1)-3*b^2
*arcsech(b*x+a)*polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1
/2)))/(1+(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(3/2)-3*b^2*polylog(3,a*(1/(b*x+a)+(1
/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(3/
2)+3*b^2*polylog(3,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1
+(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(3/2)-3*b^2*arcsech(b*x+a)^2*ln(1-a*(1/(b*x+
a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)^(1/2)))/a^2/(-a^2+1)
^(1/2)+3*b^2*arcsech(b*x+a)^2*ln(1-a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*
x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(1/2)-6*b^2*arcsech(b*x+a)*
polylog(2,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-(-a^2+1)
^(1/2)))/a^2/(-a^2+1)^(1/2)+6*b^2*arcsech(b*x+a)*polylog(2,a*(1/(b*x+a)+(1/(
b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(1/2
)+6*b^2*polylog(3,a*(1/(b*x+a)+(1/(b*x+a)-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1-
(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(1/2)-6*b^2*polylog(3,a*(1/(b*x+a)+(1/(b*x+a)
-1)^(1/2)*(1/(b*x+a)+1)^(1/2))/(1+(-a^2+1)^(1/2)))/a^2/(-a^2+1)^(1/2)+3/2*b
^2*(b*x+a+1)*arcsech(b*x+a)^2*((-b*x-a+1)/(b*x+a+1))^(1/2)/a/(-a^2+1)/(b*x+
a)/(1-a/(b*x+a))
```

Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 965, normalized size of antiderivative = 1.00,
number of steps used = 32, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 1.083$, Rules

used = {6456, 5576, 4276, 3405, 3401, 2296, 2221, 2611, 2320, 6724, 5681, 2317, 2438}

$$\begin{aligned}
\int \frac{\operatorname{sech}^{-1}(a+bx)^3}{x^3} dx = & \frac{b^2 \operatorname{sech}^{-1}(a+bx)^3}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)^3}{2x^2} - \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2(1-a^2)} \\
& - \frac{3b^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) \operatorname{sech}^{-1}(a+bx)^2}{a^2 \sqrt{1-a^2}} \\
& + \frac{3b^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) \operatorname{sech}^{-1}(a+bx)^2}{2a^2(1-a^2)^{3/2}} \\
& + \frac{3b^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) \operatorname{sech}^{-1}(a+bx)^2}{a^2 \sqrt{1-a^2}} \\
& - \frac{3b^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) \operatorname{sech}^{-1}(a+bx)^2}{2a^2(1-a^2)^{3/2}} \\
& + \frac{3b^2 \sqrt{\frac{-a-bx+1}{a+bx+1}}(a+bx+1) \operatorname{sech}^{-1}(a+bx)^2}{2a(1-a^2)(a+bx)\left(1 - \frac{a}{a+bx}\right)} \\
& + \frac{3b^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) \operatorname{sech}^{-1}(a+bx)}{a^2(1-a^2)} \\
& + \frac{3b^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) \operatorname{sech}^{-1}(a+bx)}{a^2(1-a^2)} \\
& - \frac{6b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) \operatorname{sech}^{-1}(a+bx)}{a^2 \sqrt{1-a^2}} \\
& + \frac{3b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) \operatorname{sech}^{-1}(a+bx)}{a^2(1-a^2)^{3/2}} \\
& + \frac{6b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) \operatorname{sech}^{-1}(a+bx)}{a^2 \sqrt{1-a^2}} \\
& - \frac{3b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) \operatorname{sech}^{-1}(a+bx)}{a^2(1-a^2)^{3/2}} \\
& + \frac{3b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)} + \frac{3b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a^2(1-a^2)} \\
& + \frac{6b^2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} - \frac{3b^2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
& - \frac{6b^2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a^2 \sqrt{1-a^2}} + \frac{3b^2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a^2(1-a^2)^{3/2}}
\end{aligned}$$

[In] Int[ArcSech[a + b*x]^3/x^3,x]

[Out]
$$\begin{aligned} & (-3*b^2*ArcSech[a + b*x]^2)/(2*a^2*(1 - a^2)) + (3*b^2*sqrt[(1 - a - b*x)/(1 + a + b*x)]*(1 + a + b*x)*ArcSech[a + b*x]^2)/(2*a*(1 - a^2)*(a + b*x)*(1 - a/(a + b*x))) + (b^2*ArcSech[a + b*x]^3)/(2*a^2) - ArcSech[a + b*x]^3/(2*x^2) + (3*b^2*ArcSech[a + b*x]*Log[1 - (a*E^ArcSech[a + b*x])]/(1 - Sqrt[1 - a^2]))/(a^2*(1 - a^2)) + (3*b^2*ArcSech[a + b*x]^2*Log[1 - (a*E^ArcSech[a + b*x])]/(1 - Sqrt[1 - a^2]))/(2*a^2*(1 - a^2)^(3/2)) - (3*b^2*ArcSech[a + b*x]^2*Log[1 - (a*E^ArcSech[a + b*x])]/(1 - Sqrt[1 - a^2]))/(a^2*sqrt[1 - a^2]) + (3*b^2*ArcSech[a + b*x]*Log[1 - (a*E^ArcSech[a + b*x])]/(1 + Sqrt[1 - a^2]))/(a^2*(1 - a^2)) - (3*b^2*ArcSech[a + b*x]^2*Log[1 - (a*E^ArcSech[a + b*x])]/(1 + Sqrt[1 - a^2]))/(2*a^2*(1 - a^2)^(3/2)) + (3*b^2*ArcSech[a + b*x]^2*Log[1 - (a*E^ArcSech[a + b*x])]/(1 + Sqrt[1 - a^2]))/(a^2*sqrt[1 - a^2]) + (3*b^2*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/(a^2*(1 - a^2)) + (3*b^2*ArcSech[a + b*x]*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/(a^2*(1 - a^2)^(3/2)) - (6*b^2*ArcSech[a + b*x]*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/(a^2*sqrt[1 - a^2]) + (3*b^2*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])/(a^2*(1 - a^2)) - (3*b^2*ArcSech[a + b*x]*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])/(a^2*(1 - a^2)^(3/2)) + (6*b^2*ArcSech[a + b*x]*PolyLog[2, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])/(a^2*sqrt[1 - a^2]) - (3*b^2*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/(a^2*(1 - a^2)^(3/2)) + (6*b^2*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 - Sqrt[1 - a^2])])/(a^2*sqrt[1 - a^2]) + (3*b^2*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])/(a^2*(1 - a^2)^(3/2)) - (6*b^2*PolyLog[3, (a*E^ArcSech[a + b*x])/(1 + Sqrt[1 - a^2])])/(a^2*sqrt[1 - a^2]) \end{aligned}$$

Rule 2221

Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2296

Int[((F_)^(u_)*((f_) + (g_)*(x_))^(m_))/((a_) + (b_)*(F_)^(u_) + (c_)*(F_)^(v_)), x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[2*(c/q), Int[(f + g*x)^m*(F^u/(b - q + 2*c*F^u)), x], x] - Dist[2*(c/q), Int[(f + g*x)^m*(F^u/(b + q + 2*c*F^u)), x], x]] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v, 2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]

Rule 2317

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))]

)^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2438

Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 2611

Int[Log[1 + (e_.)*((F_)^(c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3401

Int[((c_.) + (d_.)*(x_)^(m_.))/((a_) + (b_.)*sin[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]), x_Symbol] := Dist[2, Int[((c + d*x)^m*(E^((-I)*e + f*fz*x)/(b + (2*a*E^((-I)*e + f*fz*x))/E^(I*Pi*(k - 1/2)) - (b*E^(2*((-I)*e + f*fz*x))/E^(2*I*k*Pi))))/E^(I*Pi*(k - 1/2)), x], x] /; FreeQ[{a, b, c, d, e, f, fz}, x] && IntegerQ[2*k] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]

Rule 3405

Int[((c_.) + (d_.)*(x_)^(m_.))/((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[b*(c + d*x)^m*(Cos[e + f*x]/(f*(a^2 - b^2)*(a + b*Sin[e + f*x]))], x] + (Dist[a/(a^2 - b^2), Int[(c + d*x)^m/(a + b*Sin[e + f*x]), x], x] - Dist[b*d*(m/(f*(a^2 - b^2))), Int[(c + d*x)^(m - 1)*(Cos[e + f*x]/(a + b*Sin[e + f*x])), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]

Rule 4276

Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Sin[e + f*x])^n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGtQ[m, 0]

Rule 5576

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sech[(c_.) + (d_.)*(x_)]*((a_) + (b_.)*Sech[
(c_.) + (d_.)*(x_)]^(n_.)*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(-(e
+ f*x)^m)*((a + b*Sech[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b
*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sech[c + d*x])^(n + 1), x], x] /
; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5681

```
Int[(((e_.) + (f_.)*(x_))^(m_.)*Sinh[(c_.) + (d_.)*(x_)]/(Cosh[(c_.) + (d_
.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[-(e + f*x)^(m + 1)/(b*f*(m + 1)),
x] + (Int[(e + f*x)^m*(E^(c + d*x)/(a - Rt[a^2 - b^2, 2] + b*E^(c + d*x)))
, x] + Int[(e + f*x)^m*(E^(c + d*x)/(a + Rt[a^2 - b^2, 2] + b*E^(c + d*x)))
, x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[a^2 - b^2, 0]
```

Rule 6456

```
Int[((a_.) + ArcSech[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Sech[x]*Ta
nh[x]*(d*e - c*f + f*Sech[x])^m, x], x, ArcSech[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\left(b^2 \text{Subst}\left(\int \frac{x^3 \text{sech}(x) \tanh(x)}{(-a + \text{sech}(x))^3} dx, x, \text{sech}^{-1}(a + bx)\right)\right) \\
&= -\frac{\text{sech}^{-1}(a + bx)^3}{2x^2} + \frac{1}{2}(3b^2) \text{Subst}\left(\int \frac{x^2}{(-a + \text{sech}(x))^2} dx, x, \text{sech}^{-1}(a + bx)\right) \\
&= -\frac{\text{sech}^{-1}(a + bx)^3}{2x^2} + \frac{1}{2}(3b^2) \text{Subst}\left(\int \left(\frac{x^2}{a^2} + \frac{x^2}{a^2(-1 + a \cosh(x))^2}\right.\right. \\
&\quad \left.\left. + \frac{2x^2}{a^2(-1 + a \cosh(x))}\right) dx, x, \text{sech}^{-1}(a + bx)\right)
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^2 \operatorname{sech}^{-1}(a+bx)^3}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)^3}{2x^2} \\
&\quad + \frac{(3b^2) \operatorname{Subst}\left(\int \frac{x^2}{(-1+a \cosh(x))^2} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{2a^2} \\
&\quad + \frac{(3b^2) \operatorname{Subst}\left(\int \frac{x^2}{-1+a \cosh(x)} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2} \\
&= \frac{3b^2 \sqrt{\frac{1-a-bx}{1+a+bx}} (1+a+bx) \operatorname{sech}^{-1}(a+bx)^2}{2a(1-a^2)(a+bx)\left(1-\frac{a}{a+bx}\right)} + \frac{b^2 \operatorname{sech}^{-1}(a+bx)^3}{2a^2} \\
&\quad - \frac{\operatorname{sech}^{-1}(a+bx)^3}{2x^2} + \frac{(6b^2) \operatorname{Subst}\left(\int \frac{e^x x^2}{a-2e^x+ae^{2x}} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2} \\
&\quad - \frac{(3b^2) \operatorname{Subst}\left(\int \frac{x^2}{-1+a \cosh(x)} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{2a^2(1-a^2)} \\
&\quad + \frac{(3b^2) \operatorname{Subst}\left(\int \frac{x \sinh(x)}{-1+a \cosh(x)} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a(1-a^2)} \\
&= -\frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2(1-a^2)} + \frac{3b^2 \sqrt{\frac{1-a-bx}{1+a+bx}} (1+a+bx) \operatorname{sech}^{-1}(a+bx)^2}{2a(1-a^2)(a+bx)\left(1-\frac{a}{a+bx}\right)} \\
&\quad + \frac{b^2 \operatorname{sech}^{-1}(a+bx)^3}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)^3}{2x^2} \\
&\quad - \frac{(3b^2) \operatorname{Subst}\left(\int \frac{e^x x^2}{a-2e^x+ae^{2x}} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2(1-a^2)} \\
&\quad + \frac{(3b^2) \operatorname{Subst}\left(\int \frac{e^x x}{-1-\sqrt{1-a^2}+ae^x} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a(1-a^2)} \\
&\quad + \frac{(3b^2) \operatorname{Subst}\left(\int \frac{e^x x}{-1+\sqrt{1-a^2}+ae^x} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a(1-a^2)} \\
&\quad + \frac{(6b^2) \operatorname{Subst}\left(\int \frac{e^x x^2}{-2-2\sqrt{1-a^2}+2ae^x} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{(6b^2) \operatorname{Subst}\left(\int \frac{e^x x^2}{-2+2\sqrt{1-a^2}+2ae^x} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a\sqrt{1-a^2}}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2(1-a^2)} + \frac{3b^2 \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx) \operatorname{sech}^{-1}(a+bx)^2}{2a(1-a^2)(a+bx)\left(1-\frac{a}{a+bx}\right)} \\
&+ \frac{b^2 \operatorname{sech}^{-1}(a+bx)^3}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)^3}{2x^2} + \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
&- \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&- \frac{(3b^2) \operatorname{Subst}\left(\int \frac{e^x x^2}{-2-2\sqrt{1-a^2}+2ae^x} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a(1-a^2)^{3/2}} \\
&+ \frac{(3b^2) \operatorname{Subst}\left(\int \frac{e^x x^2}{-2+2\sqrt{1-a^2}+2ae^x} dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a(1-a^2)^{3/2}} \\
&- \frac{(3b^2) \operatorname{Subst}\left(\int \log\left(1 + \frac{ae^x}{-1-\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2(1-a^2)} \\
&- \frac{(3b^2) \operatorname{Subst}\left(\int \log\left(1 + \frac{ae^x}{-1+\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2(1-a^2)} \\
&- \frac{(6b^2) \operatorname{Subst}\left(\int x \log\left(1 + \frac{2ae^x}{-2-2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{(6b^2) \operatorname{Subst}\left(\int x \log\left(1 + \frac{2ae^x}{-2+2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2 \sqrt{1-a^2}}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2(1-a^2)} + \frac{3b^2 \sqrt{\frac{1-a-bx}{1+a+bx}} (1+a+bx) \operatorname{sech}^{-1}(a+bx)^2}{2a(1-a^2)(a+bx)\left(1-\frac{a}{a+bx}\right)} \\
&+ \frac{b^2 \operatorname{sech}^{-1}(a+bx)^3}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)^3}{2x^2} + \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{2a^2(1-a^2)^{3/2}} \\
&- \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
&- \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{2a^2(1-a^2)^{3/2}} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&- \frac{6b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{6b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{(3b^2) \operatorname{Subst}\left(\int x \log\left(1+\frac{2ae^x}{-2-2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2(1-a^2)^{3/2}} \\
&- \frac{(3b^2) \operatorname{Subst}\left(\int x \log\left(1+\frac{2ae^x}{-2+2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2(1-a^2)^{3/2}} \\
&- \frac{(3b^2) \operatorname{Subst}\left(\int \frac{\log\left(1+\frac{ax}{-1-\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{a^2(1-a^2)} \\
&- \frac{(3b^2) \operatorname{Subst}\left(\int \frac{\log\left(1+\frac{ax}{-1+\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{a^2(1-a^2)} \\
&- \frac{(6b^2) \operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, -\frac{2ae^x}{-2-2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{(6b^2) \operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, -\frac{2ae^x}{-2+2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2 \sqrt{1-a^2}}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2(1-a^2)} + \frac{3b^2 \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx) \operatorname{sech}^{-1}(a+bx)^2}{2a(1-a^2)(a+bx)\left(1-\frac{a}{a+bx}\right)} \\
&+ \frac{b^2 \operatorname{sech}^{-1}(a+bx)^3}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)^3}{2x^2} + \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{2a^2(1-a^2)^{3/2}} \\
&- \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
&- \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{2a^2(1-a^2)^{3/2}} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} + \frac{3b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
&- \frac{6b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{3b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)} - \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
&+ \frac{6b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{(3b^2) \operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, -\frac{2ae^x}{-2-2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2(1-a^2)^{3/2}} \\
&- \frac{(3b^2) \operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, -\frac{2ae^x}{-2+2\sqrt{1-a^2}}\right) dx, x, \operatorname{sech}^{-1}(a+bx)\right)}{a^2(1-a^2)^{3/2}} \\
&+ \frac{(6b^2) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(2, \frac{ax}{1-\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{a^2 \sqrt{1-a^2}} \\
&- \frac{(6b^2) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(2, \frac{ax}{1+\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{a^2 \sqrt{1-a^2}}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2(1-a^2)} + \frac{3b^2 \sqrt{\frac{1-a-bx}{1+a+bx}} (1+a+bx) \operatorname{sech}^{-1}(a+bx)^2}{2a(1-a^2)(a+bx)\left(1-\frac{a}{a+bx}\right)} \\
&+ \frac{b^2 \operatorname{sech}^{-1}(a+bx)^3}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)^3}{2x^2} + \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{2a^2(1-a^2)^{3/2}} \\
&- \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
&- \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{2a^2(1-a^2)^{3/2}} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} + \frac{3b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
&- \frac{6b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{3b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)} - \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
&+ \frac{6b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{6b^2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} - \frac{6b^2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&- \frac{(3b^2) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(2, \frac{ax}{1-\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{a^2(1-a^2)^{3/2}} \\
&+ \frac{(3b^2) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(2, \frac{ax}{1+\sqrt{1-a^2}}\right)}{x} dx, x, e^{\operatorname{sech}^{-1}(a+bx)}\right)}{a^2(1-a^2)^{3/2}}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2}{2a^2(1-a^2)} + \frac{3b^2 \sqrt{\frac{1-a-bx}{1+a+bx}}(1+a+bx) \operatorname{sech}^{-1}(a+bx)^2}{2a(1-a^2)(a+bx)\left(1-\frac{a}{a+bx}\right)} \\
&+ \frac{b^2 \operatorname{sech}^{-1}(a+bx)^3}{2a^2} - \frac{\operatorname{sech}^{-1}(a+bx)^3}{2x^2} + \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{2a^2(1-a^2)^{3/2}} \\
&- \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
&- \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{2a^2(1-a^2)^{3/2}} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx)^2 \log\left(1-\frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} + \frac{3b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)} \\
&+ \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
&- \frac{6b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{3b^2 \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)} - \frac{3b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} \\
&+ \frac{6b^2 \operatorname{sech}^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&- \frac{3b^2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} + \frac{6b^2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}} \\
&+ \frac{3b^2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} - \frac{6b^2 \operatorname{PolyLog}\left(3, \frac{ae^{\operatorname{sech}^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a^2 \sqrt{1-a^2}}
\end{aligned}$$

Mathematica [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^3} dx = \int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^3} dx$$

```
[In] Integrate[ArcSech[a + b*x]^3/x^3,x]
```

```
[Out] Integrate[ArcSech[a + b*x]^3/x^3, x]
```

Maple [F]

$$\int \frac{\operatorname{arcsech}(bx + a)^3}{x^3} dx$$

```
[In] int(arcsech(b*x+a)^3/x^3,x)
```

```
[Out] int(arcsech(b*x+a)^3/x^3,x)
```

Fricas [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^3} dx = \int \frac{\operatorname{arsech}(bx + a)^3}{x^3} dx$$

```
[In] integrate(arcsech(b*x+a)^3/x^3,x, algorithm="fricas")
```

```
[Out] integral(arcsech(b*x + a)^3/x^3, x)
```

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^3} dx = \int \frac{\operatorname{asech}^3(a + bx)}{x^3} dx$$

```
[In] integrate(asech(b*x+a)**3/x**3,x)
```

```
[Out] Integral(asech(a + b*x)**3/x**3, x)
```

Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^3} dx = \int \frac{\operatorname{arosech}(bx + a)^3}{x^3} dx$$

[In] integrate(arcsech(b*x+a)^3/x^3,x, algorithm="maxima")

[Out] $-1/2*\log(\sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1})*b*x + \sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1}*a + b*x + a)^3/x^2 - \operatorname{integrate}(1/2*(16*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*\sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1}*\log(b*x + a)^3 + 16*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*\log(b*x + a)^3 - 3*(b^3*x^3 + 2*a*b^2*x^2 + (a^2*b - b)*x - 4*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*\log(b*x + a) - (2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*\sqrt{b*x + a + 1}*\log(b*x + a) - (2*b^3*x^3 + 4*a*b^2*x^2 + (2*a^2*b - b)*x - 2*(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*\log(b*x + a))*\sqrt{b*x + a + 1})*\sqrt{-b*x - a + 1})*\log(\sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1})*b*x + \sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1}*a + b*x + a)^2 - 24*((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*\sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1}*\log(b*x + a)^2 + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2*b - b)*x - a)*\log(b*x + a)^2)*\log(\sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1})*b*x + \sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1}*a + b*x + a))/(b^3*x^6 + 3*a*b^2*x^5 + (3*a^2*b - b)*x^4 + (a^3 - a)*x^3 + (b^3*x^6 + 3*a*b^2*x^5 + (3*a^2*b - b)*x^4 + (a^3 - a)*x^3)*\sqrt{b*x + a + 1}*\sqrt{-b*x - a + 1}), x)$

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^3} dx = \int \frac{\operatorname{arosech}(bx + a)^3}{x^3} dx$$

[In] integrate(arcsech(b*x+a)^3/x^3,x, algorithm="giac")

[Out] integrate(arcsech(b*x + a)^3/x^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)^3}{x^3} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)^3}{x^3} dx$$

[In] int(acosh(1/(a + b*x))^3/x^3,x)

[Out] int(acosh(1/(a + b*x))^3/x^3, x)

3.20 $\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx$

Optimal result	204
Rubi [A] (verified)	204
Mathematica [A] (verified)	206
Maple [A] (verified)	206
Fricas [A] (verification not implemented)	206
Sympy [F]	207
Maxima [A] (verification not implemented)	207
Giac [F]	207
Mupad [F(-1)]	208

Optimal result

Integrand size = 10, antiderivative size = 164

$$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{1-x}{4\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} + \frac{(1-x)^2}{4\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}}$$

$$-\frac{3(1-x)^3}{20\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}}$$

$$+\frac{(1-x)^4}{28\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} + \frac{1}{4}x^4 \operatorname{sech}^{-1}(\sqrt{x})$$

[Out] $1/4*x^4*\operatorname{arcsech}(x^{(1/2)})+1/4*(-1+x)/x^{(1/2)/(-1+1/x^{(1/2)})^{(1/2)/(1+1/x^{(1/2)})^{(1/2)}}+1/4*(1-x)^2/x^{(1/2)/(-1+1/x^{(1/2)})^{(1/2)/(1+1/x^{(1/2)})^{(1/2)}}-3/20*(1-x)^3/x^{(1/2)/(-1+1/x^{(1/2)})^{(1/2)/(1+1/x^{(1/2)})^{(1/2)}}+1/28*(1-x)^4/x^{(1/2)/(-1+1/x^{(1/2)})^{(1/2)/(1+1/x^{(1/2)})^{(1/2)}}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {6480, 12, 45}

$$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx = \frac{1}{4}x^4 \operatorname{sech}^{-1}(\sqrt{x}) + \frac{(1-x)^4}{28\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}}$$

$$-\frac{3(1-x)^3}{20\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}}$$

$$+\frac{(1-x)^2}{4\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}} - \frac{1-x}{4\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}}$$

[In] Int[x^3*ArcSech[Sqrt[x]],x]

[Out] $-1/4*(1-x)/(\text{Sqrt}[-1+1/\text{Sqrt}[x]]*\text{Sqrt}[1+1/\text{Sqrt}[x]]*\text{Sqrt}[x]) + (1-x)^2/(4*\text{Sqrt}[-1+1/\text{Sqrt}[x]]*\text{Sqrt}[1+1/\text{Sqrt}[x]]*\text{Sqrt}[x]) - (3*(1-x)^3)/(20*\text{Sqrt}[-1+1/\text{Sqrt}[x]]*\text{Sqrt}[1+1/\text{Sqrt}[x]]*\text{Sqrt}[x]) + (1-x)^4/(28*\text{Sqrt}[-1+1/\text{Sqrt}[x]]*\text{Sqrt}[1+1/\text{Sqrt}[x]]*\text{Sqrt}[x]) + (x^4*\text{ArcSech}[\text{Sqrt}[x]])/4$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 6480

Int[((a_.) + ArcSech[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcSech[u])/(d*(m + 1))), x] + Dist[b*(Sqrt[1 - u^2]/(d*(m + 1)*u*Sqrt[-1 + 1/u]*Sqrt[1 + 1/u])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[1 - u^2])), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{4}x^4\text{sech}^{-1}(\sqrt{x}) + \frac{\sqrt{1-x} \int \frac{x^3}{2\sqrt{1-x}} dx}{4\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} \\
 &= \frac{1}{4}x^4\text{sech}^{-1}(\sqrt{x}) + \frac{\sqrt{1-x} \int \frac{x^3}{\sqrt{1-x}} dx}{8\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} \\
 &= \frac{1}{4}x^4\text{sech}^{-1}(\sqrt{x}) + \frac{\sqrt{1-x} \int \left(\frac{1}{\sqrt{1-x}} - 3\sqrt{1-x} + 3(1-x)^{3/2} - (1-x)^{5/2} \right) dx}{8\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} \\
 &= -\frac{1-x}{4\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} + \frac{(1-x)^2}{4\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} \\
 &\quad - \frac{3(1-x)^3}{20\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} + \frac{(1-x)^4}{28\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} + \frac{1}{4}x^4\text{sech}^{-1}(\sqrt{x})
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.51

$$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{1}{140} \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}} (16 + 16\sqrt{x} + 8x + 8x^{3/2} + 6x^2 + 6x^{5/2} + 5x^3 + 5x^{7/2}) + \frac{1}{4} x^4 \operatorname{sech}^{-1}(\sqrt{x})$$

[In] Integrate[x^3*ArcSech[Sqrt[x]],x]

[Out] -1/140*(Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])])*(16 + 16*Sqrt[x] + 8*x + 8*x^(3/2) + 6*x^2 + 6*x^(5/2) + 5*x^3 + 5*x^(7/2))) + (x^4*ArcSech[Sqrt[x]])/4

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.33

method	result	size
derivativedivides	$\frac{x^4 \operatorname{arcsech}(\sqrt{x})}{4} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}} (5x^3+6x^2+8x+16)}{140}$	54
default	$\frac{x^4 \operatorname{arcsech}(\sqrt{x})}{4} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}} (5x^3+6x^2+8x+16)}{140}$	54
parts	$\frac{x^4 \operatorname{arcsech}(\sqrt{x})}{4} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}} (5x^3+6x^2+8x+16)}{140}$	54

[In] int(x^3*arcsech(x^(1/2)),x,method=_RETURNVERBOSE)

[Out] 1/4*x^4*arcsech(x^(1/2))-1/140*(-(x^(1/2)-1)/x^(1/2))^(1/2)*x^(1/2)*((x^(1/2)+1)/x^(1/2))^(1/2)*(5*x^3+6*x^2+8*x+16)

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.35

$$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx = \frac{1}{4} x^4 \log \left(\frac{x \sqrt{-\frac{x-1}{x}} + \sqrt{x}}{x} \right) - \frac{1}{140} (5x^3 + 6x^2 + 8x + 16) \sqrt{x} \sqrt{-\frac{x-1}{x}}$$

[In] integrate(x^3*arcsech(x^(1/2)),x, algorithm="fricas")

[Out] $\frac{1}{4}x^4 \log\left(\frac{x\sqrt{-x-1} + \sqrt{x}}{x}\right) - \frac{1}{140}(5x^3 + 6x^2 + 8x + 16)\sqrt{x}\sqrt{-x-1}$

Sympy [F]

$$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x^3 \operatorname{asech}(\sqrt{x}) dx$$

[In] `integrate(x**3*asech(x**(1/2)),x)`

[Out] `Integral(x**3*asech(sqrt(x)), x)`

Maxima [A] (verification not implemented)

none

Time = 0.21 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.35

$$\begin{aligned} \int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx &= \frac{1}{28} x^{\frac{7}{2}} \left(\frac{1}{x} - 1\right)^{\frac{7}{2}} - \frac{3}{20} x^{\frac{5}{2}} \left(\frac{1}{x} - 1\right)^{\frac{5}{2}} \\ &\quad + \frac{1}{4} x^4 \operatorname{arsech}(\sqrt{x}) + \frac{1}{4} x^{\frac{3}{2}} \left(\frac{1}{x} - 1\right)^{\frac{3}{2}} - \frac{1}{4} \sqrt{x} \sqrt{\frac{1}{x} - 1} \end{aligned}$$

[In] `integrate(x^3*arcsech(x^(1/2)),x, algorithm="maxima")`

[Out] $\frac{1}{28}x^{7/2}(1/x - 1)^{7/2} - \frac{3}{20}x^{5/2}(1/x - 1)^{5/2} + \frac{1}{4}x^4 \operatorname{arcsech}(\sqrt{x}) + \frac{1}{4}x^{3/2}(1/x - 1)^{3/2} - \frac{1}{4}\sqrt{x}\sqrt{1/x - 1}$

Giac [F]

$$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x^3 \operatorname{arsech}(\sqrt{x}) dx$$

[In] `integrate(x^3*arcsech(x^(1/2)),x, algorithm="giac")`

[Out] `integrate(x^3*arcsech(sqrt(x)), x)`

Mupad [F(-1)]

Timed out.

$$\int x^3 \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x^3 \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right) dx$$

```
[In] int(x^3*acosh(1/x^(1/2)),x)
```

```
[Out] int(x^3*acosh(1/x^(1/2)), x)
```


3.21 $\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx$

Optimal result	209
Rubi [A] (verified)	209
Mathematica [A] (verified)	211
Maple [A] (verified)	211
Fricas [A] (verification not implemented)	211
Sympy [F]	212
Maxima [A] (verification not implemented)	212
Giac [F]	212
Mupad [F(-1)]	212

Optimal result

Integrand size = 10, antiderivative size = 126

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{1-x}{3\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} + \frac{2(1-x)^2}{9\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} - \frac{(1-x)^3}{15\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} + \frac{1}{3}x^3 \operatorname{sech}^{-1}(\sqrt{x})$$

[Out] $\frac{1}{3}x^3 \operatorname{arcsech}(x^{1/2}) + \frac{1}{3}(-1+x)/x^{1/2}/(-1+1/x^{1/2})^{1/2}/(1+1/x^{1/2})^{1/2} + \frac{2}{9}(1-x)^2/x^{1/2}/(-1+1/x^{1/2})^{1/2}/(1+1/x^{1/2})^{1/2} - \frac{1}{15}(1-x)^3/x^{1/2}/(-1+1/x^{1/2})^{1/2}/(1+1/x^{1/2})^{1/2}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {6480, 12, 45}

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = \frac{1}{3}x^3 \operatorname{sech}^{-1}(\sqrt{x}) - \frac{(1-x)^3}{15\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}} + \frac{2(1-x)^2}{9\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}} - \frac{1-x}{3\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}}$$

[In] `Int[x^2*ArcSech[Sqrt[x]],x]`

[Out] $-\frac{1}{3}(1-x)/(\sqrt{-1+1/\sqrt{x}}*\sqrt{1+1/\sqrt{x}}*\sqrt{x}) + \frac{2*(1-x)^2}{(9*\sqrt{-1+1/\sqrt{x}}*\sqrt{1+1/\sqrt{x}}*\sqrt{x})} - \frac{(1-x)^3}{(15*\sqrt{-1+1/\sqrt{x}}*\sqrt{1+1/\sqrt{x}}*\sqrt{x})} + \frac{(x^3*\operatorname{ArcSech}[\sqrt{x}])}{3}$

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 45

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && Le
Q[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

Rule 6480

```
Int[((a_.) + ArcSech[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Si
mp[(c + d*x)^(m + 1)*((a + b*ArcSech[u])/(d*(m + 1))), x] + Dist[b*(Sqrt[1
- u^2]/(d*(m + 1)*u*Sqrt[-1 + 1/u]*Sqrt[1 + 1/u])), Int[SimplifyIntegrand[(
c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[1 - u^2])), x], x], x] /; FreeQ[{a, b, c,
d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c
+ d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{3}x^3 \operatorname{sech}^{-1}(\sqrt{x}) + \frac{\sqrt{1-x} \int \frac{x^2}{2\sqrt{1-x}} dx}{3\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} \\
&= \frac{1}{3}x^3 \operatorname{sech}^{-1}(\sqrt{x}) + \frac{\sqrt{1-x} \int \frac{x^2}{\sqrt{1-x}} dx}{6\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} \\
&= \frac{1}{3}x^3 \operatorname{sech}^{-1}(\sqrt{x}) + \frac{\sqrt{1-x} \int \left(\frac{1}{\sqrt{1-x}} - 2\sqrt{1-x} + (1-x)^{3/2} \right) dx}{6\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} \\
&= -\frac{1-x}{3\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} + \frac{2(1-x)^2}{9\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} \\
&\quad - \frac{(1-x)^3}{15\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} + \frac{1}{3}x^3 \operatorname{sech}^{-1}(\sqrt{x})
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.57

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{1}{45} \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}} (8+8\sqrt{x}+4x+4x^{3/2}+3x^2+3x^{5/2}) + \frac{1}{3} x^3 \operatorname{sech}^{-1}(\sqrt{x})$$

`[In] Integrate[x^2*ArcSech[Sqrt[x]],x]`

```
[Out] -1/45*(Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])])*(8 + 8*Sqrt[x] + 4*x + 4*x^(3/2) +
3*x^2 + 3*x^(5/2))) + (x^3*ArcSech[Sqrt[x]])/3
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.39

method	result	size
derivativedivides	$\frac{x^3 \operatorname{arcsech}(\sqrt{x})}{3} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}} (3x^2+4x+8)}{45}$	49
default	$\frac{x^3 \operatorname{arcsech}(\sqrt{x})}{3} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}} (3x^2+4x+8)}{45}$	49
parts	$\frac{x^3 \operatorname{arcsech}(\sqrt{x})}{3} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}} (3x^2+4x+8)}{45}$	49

`[In] int(x^2*arcsech(x^(1/2)),x,method=_RETURNVERBOSE)`

```
[Out] 1/3*x^3*arcsech(x^(1/2))-1/45*(-(x^(1/2)-1)/x^(1/2))^(1/2)*x^(1/2)*((x^(1/2)
)+1)/x^(1/2))^(1/2)*(3*x^2+4*x+8)
```

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.41

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = \frac{1}{3} x^3 \log \left(\frac{x \sqrt{-\frac{x-1}{x}} + \sqrt{x}}{x} \right) - \frac{1}{45} (3x^2 + 4x + 8) \sqrt{x} \sqrt{-\frac{x-1}{x}}$$

`[In] integrate(x^2*arcsech(x^(1/2)),x, algorithm="fricas")`

```
[Out] 1/3*x^3*log((x*sqrt(-(x - 1)/x) + sqrt(x))/x) - 1/45*(3*x^2 + 4*x + 8)*sqrt
(x)*sqrt(-(x - 1)/x)
```

Sympy [F]

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x^2 \operatorname{asech}(\sqrt{x}) dx$$

[In] integrate(x**2*asech(x**(1/2)),x)

[Out] Integral(x**2*asech(sqrt(x)), x)

Maxima [A] (verification not implemented)

none

Time = 0.21 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.37

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{1}{15} x^{\frac{5}{2}} \left(\frac{1}{x} - 1\right)^{\frac{5}{2}} + \frac{1}{3} x^3 \operatorname{arsech}(\sqrt{x}) + \frac{2}{9} x^{\frac{3}{2}} \left(\frac{1}{x} - 1\right)^{\frac{3}{2}} - \frac{1}{3} \sqrt{x} \sqrt{\frac{1}{x} - 1}$$

[In] integrate(x^2*arcsech(x^(1/2)),x, algorithm="maxima")

[Out] -1/15*x^(5/2)*(1/x - 1)^(5/2) + 1/3*x^3*arcsech(sqrt(x)) + 2/9*x^(3/2)*(1/x - 1)^(3/2) - 1/3*sqrt(x)*sqrt(1/x - 1)

Giac [F]

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x^2 \operatorname{arsech}(\sqrt{x}) dx$$

[In] integrate(x^2*arcsech(x^(1/2)),x, algorithm="giac")

[Out] integrate(x^2*arcsech(sqrt(x)), x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x^2 \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right) dx$$

[In] int(x^2*acosh(1/x^(1/2)),x)

[Out] int(x^2*acosh(1/x^(1/2)), x)

3.22 $\int x \operatorname{sech}^{-1}(\sqrt{x}) dx$

Optimal result	213
Rubi [A] (verified)	213
Mathematica [A] (verified)	214
Maple [A] (verified)	215
Fricas [A] (verification not implemented)	215
Sympy [F]	215
Maxima [A] (verification not implemented)	216
Giac [F]	216
Mupad [F(-1)]	216

Optimal result

Integrand size = 8, antiderivative size = 88

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{1-x}{2\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} + \frac{(1-x)^2}{6\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} + \frac{1}{2}x^2 \operatorname{sech}^{-1}(\sqrt{x})$$

[Out] $\frac{1}{2}x^2 \operatorname{arcsech}(x^{1/2}) + \frac{1}{2}(-1+x)/x^{1/2}/(-1+1/x^{1/2})^{1/2}/(1+1/x^{1/2})^{1/2} + \frac{1}{6}(1-x)^2/x^{1/2}/(-1+1/x^{1/2})^{1/2}/(1+1/x^{1/2})^{1/2}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6480, 12, 45}

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = \frac{1}{2}x^2 \operatorname{sech}^{-1}(\sqrt{x}) + \frac{(1-x)^2}{6\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}} - \frac{1-x}{2\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}}$$

[In] `Int[x*ArcSech[Sqrt[x]],x]`

[Out] $-1/2*(1-x)/(\operatorname{Sqrt}[-1+1/\operatorname{Sqrt}[x]]*\operatorname{Sqrt}[1+1/\operatorname{Sqrt}[x]]*\operatorname{Sqrt}[x]) + (1-x)^2/(6*\operatorname{Sqrt}[-1+1/\operatorname{Sqrt}[x]]*\operatorname{Sqrt}[1+1/\operatorname{Sqrt}[x]]*\operatorname{Sqrt}[x]) + (x^2*\operatorname{ArcSech}[\operatorname{Sqrt}[x]])/2$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 45

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && ( !IntegerQ[n] || (EqQ[c, 0] && Le
Q[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

Rule 6480

```
Int[((a_.) + ArcSech[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Si
mp[(c + d*x)^(m + 1)*((a + b*ArcSech[u])/(d*(m + 1))), x] + Dist[b*Sqrt[1
- u^2]/(d*(m + 1)*u*Sqrt[-1 + 1/u]*Sqrt[1 + 1/u]), Int[SimplifyIntegrand[(
c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[1 - u^2])), x], x], x] /; FreeQ[{a, b, c,
d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c
+ d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{2}x^2 \operatorname{sech}^{-1}(\sqrt{x}) + \frac{\sqrt{1-x} \int \frac{x}{2\sqrt{1-x}} dx}{2\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} \\
&= \frac{1}{2}x^2 \operatorname{sech}^{-1}(\sqrt{x}) + \frac{\sqrt{1-x} \int \frac{x}{\sqrt{1-x}} dx}{4\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} \\
&= \frac{1}{2}x^2 \operatorname{sech}^{-1}(\sqrt{x}) + \frac{\sqrt{1-x} \int \left(\frac{1}{\sqrt{1-x}} - \sqrt{1-x} \right) dx}{4\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} \\
&= -\frac{1-x}{2\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} + \frac{(1-x)^2}{6\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}} + \frac{1}{2}x^2 \operatorname{sech}^{-1}(\sqrt{x})
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.64

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{1}{6} \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}} (2 + 2\sqrt{x} + x + x^{3/2}) + \frac{1}{2} x^2 \operatorname{sech}^{-1}(\sqrt{x})$$

```
[In] Integrate[x*ArcSech[Sqrt[x]], x]
```

```
[Out] -1/6*(Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]*(2 + 2*Sqrt[x] + x + x^(3/2))) + (x
^2*ArcSech[Sqrt[x]])/2
```

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.48

method	result	size
derivativedivides	$\frac{x^2 \operatorname{arcsech}(\sqrt{x})}{2} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}(x+2)}{6}$	42
default	$\frac{x^2 \operatorname{arcsech}(\sqrt{x})}{2} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}(x+2)}{6}$	42
parts	$\frac{x^2 \operatorname{arcsech}(\sqrt{x})}{2} - \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}(x+2)}{6}$	42

[In] `int(x*arcsech(x^(1/2)),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{2}x^2 \operatorname{arcsech}(x^{1/2}) - \frac{1}{6}(-x^{1/2}-1)/x^{1/2} \sqrt{x} \sqrt{(x^{1/2}+1)/x^{1/2}}(x+2)$

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.51

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = \frac{1}{2} x^2 \log \left(\frac{x \sqrt{-\frac{x-1}{x}} + \sqrt{x}}{x} \right) - \frac{1}{6} (x+2) \sqrt{x} \sqrt{-\frac{x-1}{x}}$$

[In] `integrate(x*arcsech(x^(1/2)),x, algorithm="fricas")`

[Out] $\frac{1}{2}x^2 \log((x \sqrt{-(x-1)/x} + \sqrt{x})/x) - \frac{1}{6}(x+2) \sqrt{x} \sqrt{-(x-1)/x}$

Sympy [F]

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x \operatorname{asech}(\sqrt{x}) dx$$

[In] `integrate(x*asech(x**(1/2)),x)`

[Out] `Integral(x*asech(sqrt(x)), x)`

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.39

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = \frac{1}{6} x^{\frac{3}{2}} \left(\frac{1}{x} - 1 \right)^{\frac{3}{2}} + \frac{1}{2} x^2 \operatorname{ar} \operatorname{sech}(\sqrt{x}) - \frac{1}{2} \sqrt{x} \sqrt{\frac{1}{x} - 1}$$

[In] integrate(x*arcsech(x^(1/2)),x, algorithm="maxima")

[Out] 1/6*x^(3/2)*(1/x - 1)^(3/2) + 1/2*x^2*arcsech(sqrt(x)) - 1/2*sqrt(x)*sqrt(1/x - 1)

Giac [F]

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x \operatorname{ar} \operatorname{sech}(\sqrt{x}) dx$$

[In] integrate(x*arcsech(x^(1/2)),x, algorithm="giac")

[Out] integrate(x*arcsech(sqrt(x)), x)

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{sech}^{-1}(\sqrt{x}) dx = \int x \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right) dx$$

[In] int(x*acosh(1/x^(1/2)),x)

[Out] int(x*acosh(1/x^(1/2)), x)

3.23 $\int \operatorname{sech}^{-1}(\sqrt{x}) dx$

Optimal result	217
Rubi [A] (verified)	217
Mathematica [B] (verified)	218
Maple [A] (verified)	218
Fricas [A] (verification not implemented)	219
Sympy [F]	219
Maxima [A] (verification not implemented)	220
Giac [F]	220
Mupad [F(-1)]	220

Optimal result

Integrand size = 6, antiderivative size = 43

$$\int \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{1-x}{\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} + x\operatorname{sech}^{-1}(\sqrt{x})$$

[Out] $x*\operatorname{arcsech}(x^{(1/2)})+(-1+x)/x^{(1/2)/(-1+1/x^{(1/2)})^{(1/2)/(1+1/x^{(1/2)})^{(1/2)}}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6478, 12, 32}

$$\int \operatorname{sech}^{-1}(\sqrt{x}) dx = x\operatorname{sech}^{-1}(\sqrt{x}) - \frac{1-x}{\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}}$$

[In] Int[ArcSech[Sqrt[x]],x]

[Out] $-((1-x)/(\operatorname{Sqrt}[-1+1/\operatorname{Sqrt}[x]]*\operatorname{Sqrt}[1+1/\operatorname{Sqrt}[x]]*\operatorname{Sqrt}[x]))+x*\operatorname{ArcSech}[\operatorname{Sqrt}[x]]$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 32

Int[((a_.) + (b_.)*(x_))^(m_), x_Symbol] := Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]

Rule 6478

```
Int[ArcSech[u_], x_Symbol] := Simp[x*ArcSech[u], x] + Dist[Sqrt[1 - u^2]/(u
*Sqrt[-1 + 1/u]*Sqrt[1 + 1/u]), Int[SimplifyIntegrand[x*(D[u, x]/(u*Sqrt[1
- u^2])), x], x], x] /; InverseFunctionFreeQ[u, x] && !FunctionOfExponenti
alQ[u, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= x \operatorname{sech}^{-1}(\sqrt{x}) + \frac{\sqrt{1-x} \int \frac{1}{2\sqrt{1-x}} dx}{\sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}}} \sqrt{x}} \\ &= x \operatorname{sech}^{-1}(\sqrt{x}) + \frac{\sqrt{1-x} \int \frac{1}{\sqrt{1-x}} dx}{2\sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}}} \sqrt{x}} \\ &= -\frac{1-x}{\sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}}} \sqrt{x}} + x \operatorname{sech}^{-1}(\sqrt{x}) \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 118 vs. $2(43) = 86$.

Time = 0.13 (sec) , antiderivative size = 118, normalized size of antiderivative = 2.74

$$\int \operatorname{sech}^{-1}(\sqrt{x}) dx = -\frac{2(-1 + \sqrt{1 - \sqrt{x}})^2 (-1 + \sqrt{1 + \sqrt{x}})^2 \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}} \sqrt{1 + \sqrt{x}}}{(-2 + \sqrt{1 - \sqrt{x}} + \sqrt{1 + \sqrt{x}})^2 \sqrt{1 - \sqrt{x}}} + x \operatorname{sech}^{-1}(\sqrt{x})$$

```
[In] Integrate[ArcSech[Sqrt[x]], x]
```

```
[Out] (-2*(-1 + Sqrt[1 - Sqrt[x]])^2*(-1 + Sqrt[1 + Sqrt[x]])^2*Sqrt[(1 - Sqrt[x]
)/(1 + Sqrt[x])]*Sqrt[1 + Sqrt[x]])/((-2 + Sqrt[1 - Sqrt[x]] + Sqrt[1 + Sqr
t[x]])^2*Sqrt[1 - Sqrt[x]]) + x*ArcSech[Sqrt[x]]
```

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$x \operatorname{arcsech}(\sqrt{x}) - \sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}$	36
default	$x \operatorname{arcsech}(\sqrt{x}) - \sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}$	36
parts	$x \operatorname{arcsech}(\sqrt{x}) - \sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}} \sqrt{x} \sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}$	36

```
[In] int(arcsech(x^(1/2)),x,method=_RETURNVERBOSE)
```

```
[Out] x*arcsech(x^(1/2))-(-(x^(1/2)-1)/x^(1/2))^(1/2)*x^(1/2)*((x^(1/2)+1)/x^(1/2))^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.91

$$\int \operatorname{sech}^{-1}(\sqrt{x}) dx = x \log \left(\frac{x \sqrt{-\frac{x-1}{x}} + \sqrt{x}}{x} \right) - \sqrt{x} \sqrt{-\frac{x-1}{x}}$$

```
[In] integrate(arcsech(x^(1/2)),x, algorithm="fricas")
```

```
[Out] x*log((x*sqrt(-(x - 1)/x) + sqrt(x))/x) - sqrt(x)*sqrt(-(x - 1)/x)
```

Sympy [F]

$$\int \operatorname{sech}^{-1}(\sqrt{x}) dx = \int \operatorname{asech}(\sqrt{x}) dx$$

```
[In] integrate(asech(x**(1/2)),x)
```

```
[Out] Integral(asech(sqrt(x)), x)
```

Maxima [A] (verification not implemented)

none

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.44

$$\int \operatorname{sech}^{-1}(\sqrt{x}) \, dx = x \operatorname{arsech}(\sqrt{x}) - \sqrt{x} \sqrt{\frac{1}{x} - 1}$$

[In] integrate(arcsech(x^(1/2)),x, algorithm="maxima")

[Out] x*arcsech(sqrt(x)) - sqrt(x)*sqrt(1/x - 1)

Giac [F]

$$\int \operatorname{sech}^{-1}(\sqrt{x}) \, dx = \int \operatorname{arsech}(\sqrt{x}) \, dx$$

[In] integrate(arcsech(x^(1/2)),x, algorithm="giac")

[Out] integrate(arcsech(sqrt(x)), x)

Mupad [F(-1)]

Timed out.

$$\int \operatorname{sech}^{-1}(\sqrt{x}) \, dx = \int \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right) \, dx$$

[In] int(acosh(1/x^(1/2)),x)

[Out] int(acosh(1/x^(1/2)), x)

3.24 $\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx$

Optimal result	221
Rubi [A] (verified)	221
Mathematica [A] (verified)	223
Maple [A] (verified)	223
Fricas [F]	224
Sympy [F]	224
Maxima [F]	224
Giac [F]	224
Mupad [F(-1)]	225

Optimal result

Integrand size = 10, antiderivative size = 46

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx = \operatorname{sech}^{-1}(\sqrt{x})^2 - 2\operatorname{sech}^{-1}(\sqrt{x}) \log\left(1 + e^{2\operatorname{sech}^{-1}(\sqrt{x})}\right) - \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(\sqrt{x})}\right)$$

[Out] $\operatorname{arcsech}(x^{(1/2)})^2 - 2*\operatorname{arcsech}(x^{(1/2)})*\ln(1+(1/x^{(1/2)}+(-1+1/x^{(1/2)})^{(1/2)})*(1+1/x^{(1/2)})^{(1/2)})^2) - \operatorname{polylog}(2, -(1/x^{(1/2)}+(-1+1/x^{(1/2)})^{(1/2)})*(1+1/x^{(1/2)})^{(1/2)})^2)$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6416, 5882, 3799, 2221, 2317, 2438}

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx = -\operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(\sqrt{x})}\right) + \operatorname{sech}^{-1}(\sqrt{x})^2 - 2\operatorname{sech}^{-1}(\sqrt{x}) \log\left(e^{2\operatorname{sech}^{-1}(\sqrt{x})} + 1\right)$$

[In] $\operatorname{Int}[\operatorname{ArcSech}[\operatorname{Sqrt}[x]]/x, x]$

[Out] $\operatorname{ArcSech}[\operatorname{Sqrt}[x]]^2 - 2*\operatorname{ArcSech}[\operatorname{Sqrt}[x]]*\operatorname{Log}[1 + E^{(2*\operatorname{ArcSech}[\operatorname{Sqrt}[x]])}] - \operatorname{PolyLog}[2, -E^{(2*\operatorname{ArcSech}[\operatorname{Sqrt}[x]])}]$

Rule 2221

$\operatorname{Int}[(((F_.)^{((g_.)*((e_.) + (f_.)*(x_)))})^{(n_.)}*((c_.) + (d_.)*(x_))^{(m_.)})/((a_.) + (b_.)*((F_.)^{((g_.)*((e_.) + (f_.)*(x_)))})^{(n_.)}), x_Symbol] \rightarrow \operatorname{Simp}$

```

[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

```

Rule 2317

```

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

```

Rule 2438

```

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

```

Rule 3799

```

Int[((c_) + (d_)*(x_)^(m_))*tan[(e_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol]
:> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

```

Rule 5882

```

Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)/(x_), x_Symbol] :> Dist[1/b, Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]

```

Rule 6416

```

Int[((a_) + ArcSech[(c_)*(x_)])*(b_)/(x_), x_Symbol] :> -Subst[Int[(a + b*ArcCosh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= 2\text{Subst}\left(\int \frac{\text{sech}^{-1}(x)}{x} dx, x, \sqrt{x}\right) \\
&= -\left(2\text{Subst}\left(\int \frac{\text{arccosh}(x)}{x} dx, x, \frac{1}{\sqrt{x}}\right)\right) \\
&= -\left(2\text{Subst}\left(\int x \tanh(x) dx, x, \text{arccosh}\left(\frac{1}{\sqrt{x}}\right)\right)\right) \\
&= \text{arccosh}\left(\frac{1}{\sqrt{x}}\right)^2 - 4\text{Subst}\left(\int \frac{e^{2x}x}{1 + e^{2x}} dx, x, \text{arccosh}\left(\frac{1}{\sqrt{x}}\right)\right)
\end{aligned}$$

$$\begin{aligned}
&= \operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)^2 - 2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right) \log\left(1 + e^{2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)}\right) \\
&\quad + 2\operatorname{Subst}\left(\int \log(1 + e^{2x}) dx, x, \operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)\right) \\
&= \operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)^2 - 2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right) \log\left(1 + e^{2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)}\right) \\
&\quad + \operatorname{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)}\right) \\
&= \operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)^2 - 2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right) \log\left(1 + e^{2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)}\right) - \operatorname{PolyLog}\left(2, -e^{2\operatorname{arccosh}\left(\frac{1}{\sqrt{x}}\right)}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.98

$$\begin{aligned}
\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx &= -\operatorname{sech}^{-1}(\sqrt{x}) \left(\operatorname{sech}^{-1}(\sqrt{x}) + 2 \log\left(1 + e^{-2\operatorname{sech}^{-1}(\sqrt{x})}\right) \right) \\
&\quad + \operatorname{PolyLog}\left(2, -e^{-2\operatorname{sech}^{-1}(\sqrt{x})}\right)
\end{aligned}$$

[In] Integrate[ArcSech[Sqrt[x]]/x,x]

[Out] -(ArcSech[Sqrt[x]]*(ArcSech[Sqrt[x]] + 2*Log[1 + E^(-2*ArcSech[Sqrt[x]])])) + PolyLog[2, -E^(-2*ArcSech[Sqrt[x]])]

Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.41

method	result
derivativedivides	$\operatorname{arcsech}(\sqrt{x})^2 - 2 \operatorname{arcsech}(\sqrt{x}) \ln\left(1 + \left(\frac{1}{\sqrt{x}} + \sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\right)^2\right) - \operatorname{polylog}\left(2, -\left(\frac{1}{\sqrt{x}} + \sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\right)^2\right)$
default	$\operatorname{arcsech}(\sqrt{x})^2 - 2 \operatorname{arcsech}(\sqrt{x}) \ln\left(1 + \left(\frac{1}{\sqrt{x}} + \sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\right)^2\right) - \operatorname{polylog}\left(2, -\left(\frac{1}{\sqrt{x}} + \sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\right)^2\right)$

[In] int(arcsech(x^(1/2))/x,x,method=_RETURNVERBOSE)

[Out] arcsech(x^(1/2))^2-2*arcsech(x^(1/2))*ln(1+(1/x^(1/2)+(-1+1/x^(1/2))^(1/2))*(1+1/x^(1/2))^(1/2))^2)-polylog(2,-(1/x^(1/2)+(-1+1/x^(1/2))^(1/2))*(1+1/x^(1/2))^(1/2))^2)

Fricas [F]

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arosech}(\sqrt{x})}{x} dx$$

[In] integrate(arcsech(x^(1/2))/x,x, algorithm="fricas")

[Out] integral(arcsech(sqrt(x))/x, x)

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{asech}(\sqrt{x})}{x} dx$$

[In] integrate(asech(x**(1/2))/x,x)

[Out] Integral(asech(sqrt(x))/x, x)

Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arosech}(\sqrt{x})}{x} dx$$

[In] integrate(arcsech(x^(1/2))/x,x, algorithm="maxima")

[Out] -1/4*log(x)^2 + log(x)*log(sqrt(sqrt(x) + 1)*sqrt(-sqrt(x) + 1) + 1) - log(sqrt(x) + 1)*log(sqrt(x)) - log(sqrt(x))*log(-sqrt(x) + 1) - dilog(-sqrt(x)) - dilog(sqrt(x)) + integrate(1/2*log(x)/((x - 1)*e^(1/2*log(sqrt(x) + 1) + 1/2*log(-sqrt(x) + 1)) + x - 1), x)

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arosech}(\sqrt{x})}{x} dx$$

[In] integrate(arcsech(x^(1/2))/x,x, algorithm="giac")

[Out] integrate(arcsech(sqrt(x))/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right)}{x} dx$$

```
[In] int(acosh(1/x^(1/2))/x,x)
```

```
[Out] int(acosh(1/x^(1/2))/x, x)
```

3.25 $\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx$

Optimal result	226
Rubi [A] (verified)	226
Mathematica [A] (verified)	228
Maple [A] (verified)	228
Fricas [A] (verification not implemented)	229
Sympy [F]	229
Maxima [A] (verification not implemented)	229
Giac [F]	230
Mupad [B] (verification not implemented)	230

Optimal result

Integrand size = 10, antiderivative size = 98

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx = \frac{1-x}{2\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}x^{3/2}}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} + \frac{\sqrt{1-x}\operatorname{arctanh}(\sqrt{1-x})}{2\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}}$$

[Out] $-\operatorname{arcsech}(x^{(1/2)})/x+1/2*(1-x)/x^{(3/2)/(-1+1/x^{(1/2)})^{(1/2)/(1+1/x^{(1/2)})^{(1/2)+1/2*\operatorname{arctanh}((1-x)^{(1/2)})*(1-x)^{(1/2)/x^{(1/2)/(-1+1/x^{(1/2)})^{(1/2)/(1+1/x^{(1/2)})^{(1/2)}}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6480, 12, 44, 65, 212}

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{1-x}\operatorname{arctanh}(\sqrt{1-x})}{2\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}} + \frac{1-x}{2\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}x^{3/2}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x}$$

[In] `Int[ArcSech[Sqrt[x]]/x^2,x]`

[Out] $(1-x)/(2*\operatorname{Sqrt}[-1+1/\operatorname{Sqrt}[x]]*\operatorname{Sqrt}[1+1/\operatorname{Sqrt}[x]]*x^{(3/2)}) - \operatorname{ArcSech}[\operatorname{Sqrt}[x]]/x + (\operatorname{Sqrt}[1-x]*\operatorname{ArcTanh}[\operatorname{Sqrt}[1-x]])/(2*\operatorname{Sqrt}[-1+1/\operatorname{Sqrt}[x]]*\operatorname{Sqrt}[1+1/\operatorname{Sqrt}[x]]*\operatorname{Sqrt}[x])$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]`

Rule 44

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((
m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !Int
egerQ[n] && LtQ[n, 0]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 6480

```
Int[((a_.) + ArcSech[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Si
mp[(c + d*x)^(m + 1)*((a + b*ArcSech[u])/(d*(m + 1))), x] + Dist[b*(Sqrt[1
- u^2]/(d*(m + 1)*u*Sqrt[-1 + 1/u]*Sqrt[1 + 1/u])), Int[SimplifyIntegrand[(
c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[1 - u^2])), x], x], x] /; FreeQ[{a, b, c,
d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c
+ d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} - \frac{\sqrt{1-x} \int \frac{1}{2\sqrt{1-xx^2}} dx}{\sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}}} \sqrt{x}} \\
&= -\frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} - \frac{\sqrt{1-x} \int \frac{1}{\sqrt{1-xx^2}} dx}{2\sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}}} \sqrt{x}} \\
&= \frac{1-x}{2\sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}}} x^{3/2}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} - \frac{\sqrt{1-x} \int \frac{1}{\sqrt{1-xx^2}} dx}{4\sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}}} \sqrt{x}} \\
&= \frac{1-x}{2\sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}}} x^{3/2}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} + \frac{\sqrt{1-x} \operatorname{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{1-x}\right)}{2\sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}}} \sqrt{x}}
\end{aligned}$$

$$= \frac{1-x}{2\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}x^{3/2}}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x} + \frac{\sqrt{1-x}\operatorname{arctanh}(\sqrt{1-x})}{2\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx$$

$$= \frac{\sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}(1+\sqrt{x}) - 2\operatorname{sech}^{-1}(\sqrt{x}) + x \log\left(1 + \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}\sqrt{x}\right) - \frac{1}{2}x \log(x)}{2x}$$

[In] Integrate[ArcSech[Sqrt[x]]/x^2,x]

[Out] (Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]*(1 + Sqrt[x]) - 2*ArcSech[Sqrt[x]] + x*Log[1 + Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]] + Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])])*Sqrt[x] - (x*Log[x])/2)/(2*x)

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.65

method	result	size
derivativedivides	$-\frac{\operatorname{arcsech}(\sqrt{x})}{x} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(\operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x+\sqrt{1-x}\right)}{2\sqrt{x}\sqrt{1-x}}$	64
default	$-\frac{\operatorname{arcsech}(\sqrt{x})}{x} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(\operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x+\sqrt{1-x}\right)}{2\sqrt{x}\sqrt{1-x}}$	64
parts	$-\frac{\operatorname{arcsech}(\sqrt{x})}{x} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(\operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x+\sqrt{1-x}\right)}{2\sqrt{x}\sqrt{1-x}}$	64

[In] int(arcsech(x^(1/2))/x^2,x,method=_RETURNVERBOSE)

[Out] -arcsech(x^(1/2))/x+1/2*(-(x^(1/2)-1)/x^(1/2))^(1/2)/x^(1/2)*((x^(1/2)+1)/x^(1/2))^(1/2)*(arctanh(1/(1-x)^(1/2))*x+(1-x)^(1/2))/(1-x)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.46

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx = \frac{(x-2) \log\left(\frac{x\sqrt{-\frac{x-1}{x}} + \sqrt{x}}{x}\right) + \sqrt{x}\sqrt{-\frac{x-1}{x}}}{2x}$$

[In] integrate(arcsech(x^(1/2))/x^2,x, algorithm="fricas")

[Out] 1/2*((x - 2)*log((x*sqrt(-(x - 1)/x) + sqrt(x))/x) + sqrt(x)*sqrt(-(x - 1)/x))/x

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx = \int \frac{\operatorname{asech}(\sqrt{x})}{x^2} dx$$

[In] integrate(asech(x**(1/2))/x**2,x)

[Out] Integral(asech(sqrt(x))/x**2, x)

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.66

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx = -\frac{\sqrt{x}\sqrt{\frac{1}{x}-1}}{2\left(x\left(\frac{1}{x}-1\right)-1\right)} - \frac{\operatorname{arsech}(\sqrt{x})}{x} + \frac{1}{4} \log\left(\sqrt{x}\sqrt{\frac{1}{x}-1}+1\right) - \frac{1}{4} \log\left(\sqrt{x}\sqrt{\frac{1}{x}-1}-1\right)$$

[In] integrate(arcsech(x^(1/2))/x^2,x, algorithm="maxima")

[Out] -1/2*sqrt(x)*sqrt(1/x - 1)/(x*(1/x - 1) - 1) - arcsech(sqrt(x))/x + 1/4*log(sqrt(x)*sqrt(1/x - 1) + 1) - 1/4*log(sqrt(x)*sqrt(1/x - 1) - 1)

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx = \int \frac{\operatorname{arosech}(\sqrt{x})}{x^2} dx$$

[In] integrate(arcsech(x^(1/2))/x^2,x, algorithm="giac")

[Out] integrate(arcsech(sqrt(x))/x^2, x)

Mupad [B] (verification not implemented)

Time = 4.90 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.41

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{\frac{1}{\sqrt{x}} - 1} \sqrt{\frac{1}{\sqrt{x}} + 1}}{2\sqrt{x}} - \frac{2 \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right) \left(\frac{1}{2\sqrt{x}} - \frac{\sqrt{x}}{4}\right)}{\sqrt{x}}$$

[In] int(acosh(1/x^(1/2))/x^2,x)

[Out] ((1/x^(1/2) - 1)^(1/2)*(1/x^(1/2) + 1)^(1/2))/(2*x^(1/2)) - (2*acosh(1/x^(1/2))*(1/(2*x^(1/2)) - x^(1/2)/4))/x^(1/2)

3.26 $\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx$

Optimal result	231
Rubi [A] (verified)	231
Mathematica [A] (verified)	233
Maple [A] (verified)	234
Fricas [A] (verification not implemented)	234
Sympy [F]	234
Maxima [A] (verification not implemented)	235
Giac [F]	235
Mupad [F(-1)]	235

Optimal result

Integrand size = 10, antiderivative size = 136

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx = \frac{1-x}{8\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}x^{5/2}}} + \frac{3(1-x)}{16\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}x^{3/2}}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{2x^2} + \frac{3\sqrt{1-x}\operatorname{arctanh}(\sqrt{1-x})}{16\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}}$$

[Out] $-1/2*\operatorname{arcsech}(x^{(1/2)})/x^2+1/8*(1-x)/x^{(5/2)/(-1+1/x^{(1/2)})^{(1/2)/(1+1/x^{(1/2)})^{(1/2)}}+3/16*(1-x)/x^{(3/2)/(-1+1/x^{(1/2)})^{(1/2)/(1+1/x^{(1/2)})^{(1/2)}}+3/16*\operatorname{arctanh}((1-x)^{(1/2)})*(1-x)^{(1/2)/x^{(1/2)/(-1+1/x^{(1/2)})^{(1/2)/(1+1/x^{(1/2)})^{(1/2)}}}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6480, 12, 44, 65, 212}

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx = \frac{3\sqrt{1-x}\operatorname{arctanh}(\sqrt{1-x})}{16\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}} + \frac{3(1-x)}{16\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}x^{3/2}} + \frac{1-x}{8\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}x^{5/2}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{2x^2}$$

[In] Int[ArcSech[Sqrt[x]]/x^3,x]

```
[Out] (1 - x)/(8*Sqrt[-1 + 1/Sqrt[x]]*Sqrt[1 + 1/Sqrt[x]]*x^(5/2)) + (3*(1 - x))/
(16*Sqrt[-1 + 1/Sqrt[x]]*Sqrt[1 + 1/Sqrt[x]]*x^(3/2)) - ArcSech[Sqrt[x]]/(2
*x^2) + (3*Sqrt[1 - x]*ArcTanh[Sqrt[1 - x]])/(16*Sqrt[-1 + 1/Sqrt[x]]*Sqrt[
1 + 1/Sqrt[x]]*Sqrt[x])
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 44

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((
m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !Int
egerQ[n] && LtQ[n, 0]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 6480

```
Int[((a_.) + ArcSech[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Si
mp[(c + d*x)^(m + 1)*((a + b*ArcSech[u])/(d*(m + 1))), x] + Dist[b*(Sqrt[1
- u^2]/(d*(m + 1)*u*Sqrt[-1 + 1/u]*Sqrt[1 + 1/u])), Int[SimplifyIntegrand[(
c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[1 - u^2])), x], x], x] /; FreeQ[{a, b, c,
d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c
+ d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]
```

Rubi steps

$$\text{integral} = -\frac{\operatorname{sech}^{-1}(\sqrt{x})}{2x^2} - \frac{\sqrt{1-x} \int \frac{1}{2\sqrt{1-xx^3}} dx}{2\sqrt{-1 + \frac{1}{\sqrt{x}}} \sqrt{1 + \frac{1}{\sqrt{x}} \sqrt{x}}}$$

$$\begin{aligned}
&= -\frac{\operatorname{sech}^{-1}(\sqrt{x})}{2x^2} - \frac{\sqrt{1-x} \int \frac{1}{\sqrt{1-xx^3}} dx}{4\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} \\
&= \frac{1-x}{8\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{5/2}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{2x^2} - \frac{(3\sqrt{1-x}) \int \frac{1}{\sqrt{1-xx^2}} dx}{16\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} \\
&= \frac{1-x}{8\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{5/2}} + \frac{3(1-x)}{16\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{3/2}} \\
&\quad - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{2x^2} - \frac{(3\sqrt{1-x}) \int \frac{1}{\sqrt{1-xx}} dx}{32\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} \\
&= \frac{1-x}{8\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{5/2}} + \frac{3(1-x)}{16\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{3/2}} \\
&\quad - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{2x^2} + \frac{(3\sqrt{1-x}) \operatorname{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{1-x}\right)}{16\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} \\
&= \frac{1-x}{8\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{5/2}} + \frac{3(1-x)}{16\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{3/2}} \\
&\quad - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{2x^2} + \frac{3\sqrt{1-x} \operatorname{arctanh}(\sqrt{1-x})}{16\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.92

$$\begin{aligned}
\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx &= \frac{1}{16} \left(\frac{\sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}(2+2\sqrt{x}+3x+3x^{3/2})}{x^2} - \frac{8\operatorname{sech}^{-1}(\sqrt{x})}{x^2} \right) \\
&\quad + 3 \log \left(1 + \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}} + \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}\sqrt{x} \right) - \frac{3 \log(x)}{2}
\end{aligned}$$

[In] Integrate[ArcSech[Sqrt[x]]/x^3,x]

[Out] ((Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]*(2 + 2*Sqrt[x] + 3*x + 3*x^(3/2)))/x^2 - (8*ArcSech[Sqrt[x]])/x^2 + 3*Log[1 + Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]] + Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]*Sqrt[x]] - (3*Log[x])/2)/16

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.58

method	result	size
derivativedivides	$-\frac{\operatorname{arcsech}(\sqrt{x})}{2x^2} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(3\operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x^2+3\sqrt{1-x}x+2\sqrt{1-x}\right)}{16x^{\frac{3}{2}}\sqrt{1-x}}$	79
default	$-\frac{\operatorname{arcsech}(\sqrt{x})}{2x^2} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(3\operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x^2+3\sqrt{1-x}x+2\sqrt{1-x}\right)}{16x^{\frac{3}{2}}\sqrt{1-x}}$	79
parts	$-\frac{\operatorname{arcsech}(\sqrt{x})}{2x^2} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(3\operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x^2+3\sqrt{1-x}x+2\sqrt{1-x}\right)}{16x^{\frac{3}{2}}\sqrt{1-x}}$	79

[In] int(arcsech(x^(1/2))/x^3,x,method=_RETURNVERBOSE)

```
[Out] -1/2*arcsech(x^(1/2))/x^2+1/16*(-(x^(1/2)-1)/x^(1/2))^(1/2)/x^(3/2)*((x^(1/2)+1)/x^(1/2))^(1/2)*(3*arctanh(1/(1-x)^(1/2)))*x^2+3*(1-x)^(1/2)*x+2*(1-x)^(1/2))/(1-x)^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.40

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx = \frac{(3x+2)\sqrt{x}\sqrt{-\frac{x-1}{x}} + (3x^2-8)\log\left(\frac{x\sqrt{-\frac{x-1}{x}}+\sqrt{x}}{x}\right)}{16x^2}$$

[In] integrate(arcsech(x^(1/2))/x^3,x, algorithm="fricas")

```
[Out] 1/16*((3*x + 2)*sqrt(x)*sqrt(-(x - 1)/x) + (3*x^2 - 8)*log((x*sqrt(-(x - 1)/x) + sqrt(x))/x))/x^2
```

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\operatorname{asech}(\sqrt{x})}{x^3} dx$$

[In] integrate(asech(x**(1/2))/x**3,x)

[Out] Integral(asech(sqrt(x))/x**3, x)

Maxima [A] (verification not implemented)

none

Time = 0.21 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.68

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx = -\frac{3x^{\frac{3}{2}}\left(\frac{1}{x}-1\right)^{\frac{3}{2}} - 5\sqrt{x}\sqrt{\frac{1}{x}-1}}{16\left(x^2\left(\frac{1}{x}-1\right)^2 - 2x\left(\frac{1}{x}-1\right) + 1\right)} - \frac{\operatorname{arsech}(\sqrt{x})}{2x^2} + \frac{3}{32} \log\left(\sqrt{x}\sqrt{\frac{1}{x}-1} + 1\right) - \frac{3}{32} \log\left(\sqrt{x}\sqrt{\frac{1}{x}-1} - 1\right)$$

[In] integrate(arcsech(x^(1/2))/x^3,x, algorithm="maxima")

```
[Out] -1/16*(3*x^(3/2)*(1/x - 1)^(3/2) - 5*sqrt(x)*sqrt(1/x - 1))/(x^2*(1/x - 1)^2 - 2*x*(1/x - 1) + 1) - 1/2*arcsech(sqrt(x))/x^2 + 3/32*log(sqrt(x)*sqrt(1/x - 1) + 1) - 3/32*log(sqrt(x)*sqrt(1/x - 1) - 1)
```

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\operatorname{arsech}(\sqrt{x})}{x^3} dx$$

[In] integrate(arcsech(x^(1/2))/x^3,x, algorithm="giac")

[Out] integrate(arcsech(sqrt(x))/x^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right)}{x^3} dx$$

[In] int(acosh(1/x^(1/2))/x^3,x)

[Out] int(acosh(1/x^(1/2))/x^3, x)

3.27 $\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx$

Optimal result	236
Rubi [A] (verified)	236
Mathematica [A] (verified)	239
Maple [A] (verified)	239
Fricas [A] (verification not implemented)	240
Sympy [F]	240
Maxima [A] (verification not implemented)	240
Giac [F]	241
Mupad [F(-1)]	241

Optimal result

Integrand size = 10, antiderivative size = 172

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx = \frac{1-x}{18\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{7/2}} + \frac{5(1-x)}{72\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{5/2}}$$

$$+ \frac{5(1-x)}{48\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{3/2}}$$

$$- \frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3} + \frac{5\sqrt{1-x}\operatorname{arctanh}(\sqrt{1-x})}{48\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}}$$

[Out] $-1/3*\operatorname{arcsech}(x^{(1/2)})/x^3+1/18*(1-x)/x^{(7/2)/(-1+1/x^{(1/2)})^{(1/2)/(1+1/x^{(1/2)})^{(1/2)}}+5/72*(1-x)/x^{(5/2)/(-1+1/x^{(1/2)})^{(1/2)/(1+1/x^{(1/2)})^{(1/2)}}+5/48*(1-x)/x^{(3/2)/(-1+1/x^{(1/2)})^{(1/2)/(1+1/x^{(1/2)})^{(1/2)}}+5/48*\operatorname{arctanh}((1-x)^{(1/2))*(1-x)^{(1/2)}/x^{(1/2)/(-1+1/x^{(1/2)})^{(1/2)/(1+1/x^{(1/2)})^{(1/2)}})$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6480, 12, 44, 65, 212}

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx = \frac{5\sqrt{1-x}\operatorname{arctanh}(\sqrt{1-x})}{48\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}\sqrt{x}} + \frac{5(1-x)}{48\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}x^{3/2}}$$

$$+ \frac{5(1-x)}{72\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}x^{5/2}} + \frac{1-x}{18\sqrt{\frac{1}{\sqrt{x}}-1}\sqrt{\frac{1}{\sqrt{x}}+1}x^{7/2}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3}$$

[In] Int[ArcSech[Sqrt[x]]/x^4,x]

[Out] (1 - x)/(18*Sqrt[-1 + 1/Sqrt[x]]*Sqrt[1 + 1/Sqrt[x]]*x^(7/2)) + (5*(1 - x))/(72*Sqrt[-1 + 1/Sqrt[x]]*Sqrt[1 + 1/Sqrt[x]]*x^(5/2)) + (5*(1 - x))/(48*Sqrt[-1 + 1/Sqrt[x]]*Sqrt[1 + 1/Sqrt[x]]*x^(3/2)) - ArcSech[Sqrt[x]]/(3*x^3) + (5*Sqrt[1 - x]*ArcTanh[Sqrt[1 - x]])/(48*Sqrt[-1 + 1/Sqrt[x]]*Sqrt[1 + 1/Sqrt[x]]*Sqrt[x])

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 44

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !IntegerQ[n] && LtQ[n, 0]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 6480

Int[((a_.) + ArcSech[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcSech[u])/(d*(m + 1))), x] + Dist[b*(Sqrt[1 - u^2]/(d*(m + 1)*u*Sqrt[-1 + 1/u]*Sqrt[1 + 1/u])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[1 - u^2])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]

Rubi steps

$$\text{integral} = -\frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3} - \frac{\sqrt{1-x} \int \frac{1}{2\sqrt{1-xx^4}} dx}{3\sqrt{-1 + \frac{1}{\sqrt{x}}}\sqrt{1 + \frac{1}{\sqrt{x}}}\sqrt{x}}$$

$$\begin{aligned}
&= -\frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3} - \frac{\sqrt{1-x} \int \frac{1}{\sqrt{1-xx^4}} dx}{6\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} \\
&= \frac{1-x}{18\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{7/2}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3} - \frac{(5\sqrt{1-x}) \int \frac{1}{\sqrt{1-xx^3}} dx}{36\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} \\
&= \frac{1-x}{18\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{7/2}} + \frac{5(1-x)}{72\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{5/2}} \\
&\quad - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3} - \frac{(5\sqrt{1-x}) \int \frac{1}{\sqrt{1-xx^2}} dx}{48\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} \\
&= \frac{1-x}{18\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{7/2}} + \frac{5(1-x)}{72\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{5/2}} \\
&\quad + \frac{5(1-x)}{48\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{3/2}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3} - \frac{(5\sqrt{1-x}) \int \frac{1}{\sqrt{1-xx}} dx}{96\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} \\
&= \frac{1-x}{18\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{7/2}} + \frac{5(1-x)}{72\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{5/2}} \\
&\quad + \frac{5(1-x)}{48\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{3/2}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3} \\
&\quad + \frac{(5\sqrt{1-x}) \operatorname{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{1-x}\right)}{48\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}} \\
&= \frac{1-x}{18\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{7/2}} + \frac{5(1-x)}{72\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{5/2}} \\
&\quad + \frac{5(1-x)}{48\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}x^{3/2}} - \frac{\operatorname{sech}^{-1}(\sqrt{x})}{3x^3} + \frac{5\sqrt{1-x} \operatorname{arctanh}(\sqrt{1-x})}{48\sqrt{-1+\frac{1}{\sqrt{x}}}\sqrt{1+\frac{1}{\sqrt{x}}}\sqrt{x}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.81

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx$$

$$= \frac{\sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}(8 + 8\sqrt{x} + 10x + 10x^{3/2} + 15x^2 + 15x^{5/2}) - 48\operatorname{sech}^{-1}(\sqrt{x}) + 15x^3 \log\left(1 + \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}\right) + \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}}}{144x^3}$$

`[In] Integrate[ArcSech[Sqrt[x]]/x^4,x]`

```
[Out] (Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])])*(8 + 8*Sqrt[x] + 10*x + 10*x^(3/2) + 15*x^2 + 15*x^(5/2)) - 48*ArcSech[Sqrt[x]] + 15*x^3*Log[1 + Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])] + Sqrt[(1 - Sqrt[x])/(1 + Sqrt[x])]*Sqrt[x]] - (15*x^3*Log[x])/2)/(144*x^3)
```

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.53

method	result	size
derivativedivides	$-\frac{\operatorname{arcsech}(\sqrt{x})}{3x^3} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(15 \operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x^3 + 15\sqrt{1-x}x^2 + 10\sqrt{1-x}x + 8\sqrt{1-x}\right)}{144x^{\frac{5}{2}}\sqrt{1-x}}$	91
default	$-\frac{\operatorname{arcsech}(\sqrt{x})}{3x^3} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(15 \operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x^3 + 15\sqrt{1-x}x^2 + 10\sqrt{1-x}x + 8\sqrt{1-x}\right)}{144x^{\frac{5}{2}}\sqrt{1-x}}$	91
parts	$-\frac{\operatorname{arcsech}(\sqrt{x})}{3x^3} + \frac{\sqrt{-\frac{\sqrt{x}-1}{\sqrt{x}}}\sqrt{\frac{\sqrt{x}+1}{\sqrt{x}}}\left(15 \operatorname{arctanh}\left(\frac{1}{\sqrt{1-x}}\right)x^3 + 15\sqrt{1-x}x^2 + 10\sqrt{1-x}x + 8\sqrt{1-x}\right)}{144x^{\frac{5}{2}}\sqrt{1-x}}$	91

`[In] int(arcsech(x^(1/2))/x^4,x,method=_RETURNVERBOSE)`

```
[Out] -1/3*arcsech(x^(1/2))/x^3+1/144*(-(x^(1/2)-1)/x^(1/2))^(1/2)/x^(5/2)*((x^(1/2)+1)/x^(1/2))^(1/2)*(15*arctanh(1/(1-x)^(1/2))*x^3+15*(1-x)^(1/2)*x^2+10*(1-x)^(1/2)*x+8*(1-x)^(1/2))/(1-x)^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.35

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx = \frac{(15x^2 + 10x + 8)\sqrt{x}\sqrt{-\frac{x-1}{x}} + 3(5x^3 - 16)\log\left(\frac{x\sqrt{-\frac{x-1}{x}} + \sqrt{x}}{x}\right)}{144x^3}$$

[In] integrate(arcsech(x^(1/2))/x^4,x, algorithm="fricas")

[Out] 1/144*((15*x^2 + 10*x + 8)*sqrt(x)*sqrt(-(x - 1)/x) + 3*(5*x^3 - 16)*log((x *sqrt(-(x - 1)/x) + sqrt(x))/x))/x^3

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\operatorname{asech}(\sqrt{x})}{x^4} dx$$

[In] integrate(asech(x**(1/2))/x**4,x)

[Out] Integral(asech(sqrt(x))/x**4, x)

Maxima [A] (verification not implemented)

none

Time = 0.20 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.67

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx = -\frac{15x^{\frac{5}{2}}\left(\frac{1}{x}-1\right)^{\frac{5}{2}} - 40x^{\frac{3}{2}}\left(\frac{1}{x}-1\right)^{\frac{3}{2}} + 33\sqrt{x}\sqrt{\frac{1}{x}-1}}{144\left(x^3\left(\frac{1}{x}-1\right)^3 - 3x^2\left(\frac{1}{x}-1\right)^2 + 3x\left(\frac{1}{x}-1\right) - 1\right)} - \frac{\operatorname{arosech}(\sqrt{x})}{3x^3} + \frac{5}{96}\log\left(\sqrt{x}\sqrt{\frac{1}{x}-1} + 1\right) - \frac{5}{96}\log\left(\sqrt{x}\sqrt{\frac{1}{x}-1} - 1\right)$$

[In] integrate(arcsech(x^(1/2))/x^4,x, algorithm="maxima")

[Out] -1/144*(15*x^(5/2)*(1/x - 1)^(5/2) - 40*x^(3/2)*(1/x - 1)^(3/2) + 33*sqrt(x)*sqrt(1/x - 1))/(x^3*(1/x - 1)^3 - 3*x^2*(1/x - 1)^2 + 3*x*(1/x - 1) - 1) - 1/3*arcsech(sqrt(x))/x^3 + 5/96*log(sqrt(x)*sqrt(1/x - 1) + 1) - 5/96*log(sqrt(x)*sqrt(1/x - 1) - 1)

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\operatorname{ar} \operatorname{sech}(\sqrt{x})}{x^4} dx$$

[In] integrate(arcsech(x^(1/2))/x^4,x, algorithm="giac")

[Out] integrate(arcsech(sqrt(x))/x^4, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right)}{x^4} dx$$

[In] int(acosh(1/x^(1/2))/x^4,x)

[Out] int(acosh(1/x^(1/2))/x^4, x)

3.28 $\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx$

Optimal result	242
Rubi [A] (verified)	242
Mathematica [A] (verified)	243
Maple [A] (verified)	243
Fricas [A] (verification not implemented)	244
Sympy [F]	244
Maxima [A] (verification not implemented)	244
Giac [F]	244
Mupad [B] (verification not implemented)	245

Optimal result

Integrand size = 4, antiderivative size = 21

$$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx = -\sqrt{-1+x}\sqrt{1+x} + x \operatorname{arccosh}(x)$$

[Out] $x \operatorname{arccosh}(x) - (-1+x)^{(1/2)} * (1+x)^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6462, 5879, 75}

$$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx = x \operatorname{arccosh}(x) - \sqrt{x-1}\sqrt{x+1}$$

[In] $\operatorname{Int}[\operatorname{ArcSech}[x^{-1}], x]$

[Out] $-(\operatorname{Sqrt}[-1+x] * \operatorname{Sqrt}[1+x]) + x * \operatorname{ArcCosh}[x]$

Rule 75

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]
```

Rule 5879

```
Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.), x_Symbol] :> Simp[x*(a + b*ArcCosh[c*x])^n, x] - Dist[b*c*n, Int[x*((a + b*ArcCosh[c*x])^(n - 1))/(Sqrt[
```

$1 + c*x]*\text{Sqrt}[-1 + c*x]))$, $x]$, $x]$ /; $\text{FreeQ}[\{a, b, c\}, x]$ && $\text{GtQ}[n, 0]$

Rule 6462

$\text{Int}[\text{ArcSech}[(c_.) / ((a_.) + (b_.) * (x_)^{(n_.)})]^{(m_.)} * (u_.), x_Symbol] \rightarrow \text{Int}[u * \text{ArcCosh}[a/c + b * (x^n/c)]^m, x]$ /; $\text{FreeQ}[\{a, b, c, n, m\}, x]$

Rubi steps

$$\begin{aligned} \text{integral} &= \int \text{arccosh}(x) dx \\ &= x \text{arccosh}(x) - \int \frac{x}{\sqrt{-1+x}\sqrt{1+x}} dx \\ &= -\sqrt{-1+x}\sqrt{1+x} + x \text{arccosh}(x) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.19

$$\int \text{sech}^{-1}\left(\frac{1}{x}\right) dx = -\sqrt{\frac{-1+x}{1+x}}(1+x) + x \text{sech}^{-1}\left(\frac{1}{x}\right)$$

[In] $\text{Integrate}[\text{ArcSech}[x^{(-1)}], x]$

[Out] $-(\text{Sqrt}[(-1 + x)/(1 + x)] * (1 + x)) + x * \text{ArcSech}[x^{(-1)}]$

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.95

method	result	size
parts	$x \text{arcsech}\left(\frac{1}{x}\right) - \sqrt{x-1}\sqrt{1+x}$	20
derivativedivides	$x \text{arcsech}\left(\frac{1}{x}\right) - \sqrt{-\left(-1 + \frac{1}{x}\right)x} \sqrt{\left(1 + \frac{1}{x}\right)x}$	29
default	$x \text{arcsech}\left(\frac{1}{x}\right) - \sqrt{-\left(-1 + \frac{1}{x}\right)x} \sqrt{\left(1 + \frac{1}{x}\right)x}$	29

[In] $\text{int}(\text{arcsech}(1/x), x, \text{method}=_RETURNVERBOSE)$

[Out] $x * \text{arcsech}(1/x) - (x-1)^{(1/2)} * (1+x)^{(1/2)}$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx = x \log\left(x + \sqrt{x^2 - 1}\right) - \sqrt{x^2 - 1}$$

[In] integrate(arcsech(1/x),x, algorithm="fricas")

[Out] x*log(x + sqrt(x^2 - 1)) - sqrt(x^2 - 1)

Sympy [F]

$$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx = \int \operatorname{asech}\left(\frac{1}{x}\right) dx$$

[In] integrate(asech(1/x),x)

[Out] Integral(asech(1/x), x)

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.76

$$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx = x \operatorname{arsech}\left(\frac{1}{x}\right) - \sqrt{x^2 - 1}$$

[In] integrate(arcsech(1/x),x, algorithm="maxima")

[Out] x*arcsech(1/x) - sqrt(x^2 - 1)

Giac [F]

$$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx = \int \operatorname{arsech}\left(\frac{1}{x}\right) dx$$

[In] integrate(arcsech(1/x),x, algorithm="giac")

[Out] integrate(arcsech(1/x), x)

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \operatorname{sech}^{-1}\left(\frac{1}{x}\right) dx = x \operatorname{acosh}(x) - \sqrt{x-1} \sqrt{x+1}$$

[In] int(acosh(x),x)

[Out] x*acosh(x) - (x - 1)^(1/2)*(x + 1)^(1/2)

3.29 $\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx$

Optimal result	246
Rubi [A] (verified)	246
Mathematica [B] (verified)	248
Maple [A] (verified)	249
Fricas [F(-2)]	249
Sympy [F]	249
Maxima [F]	250
Giac [F]	250
Mupad [F(-1)]	250

Optimal result

Integrand size = 10, antiderivative size = 61

$$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx = \frac{\operatorname{sech}^{-1}(ax^n)^2}{2n} - \frac{\operatorname{sech}^{-1}(ax^n) \log\left(1 + e^{2\operatorname{sech}^{-1}(ax^n)}\right)}{n} - \frac{\operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(ax^n)}\right)}{2n}$$

[Out] $1/2*\operatorname{arcsech}(a*x^n)^2/n - \operatorname{arcsech}(a*x^n)*\ln(1+(1/a/(x^n)+(1/a/(x^n)-1)^{(1/2))*1/a/(x^n)+1)^{(1/2)})/n - 1/2*\operatorname{polylog}(2, -(1/a/(x^n)+(1/a/(x^n)-1)^{(1/2))*(1/a/(x^n)+1)^{(1/2)})/n$

Rubi [A] (verified)

Time = 0.08 (sec), antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6416, 5882, 3799, 2221, 2317, 2438}

$$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx = -\frac{\operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(ax^n)}\right)}{2n} + \frac{\operatorname{sech}^{-1}(ax^n)^2}{2n} - \frac{\operatorname{sech}^{-1}(ax^n) \log\left(e^{2\operatorname{sech}^{-1}(ax^n)} + 1\right)}{n}$$

[In] $\operatorname{Int}[\operatorname{ArcSech}[a*x^n]/x, x]$

[Out] $\operatorname{ArcSech}[a*x^n]^2/(2*n) - (\operatorname{ArcSech}[a*x^n]*\operatorname{Log}[1 + E^{(2*\operatorname{ArcSech}[a*x^n])}])/n - \operatorname{PolyLog}[2, -E^{(2*\operatorname{ArcSech}[a*x^n])}]/(2*n)$

Rule 2221

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] :> Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)
))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*(c_) + (d_)*(x_)))^(n_)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3799

```
Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_]*(f_)*(x_)]], x
_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(
c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))], x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

Rule 5882

```
Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)/(x_), x_Symbol] :> Dist[1/b,
Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a,
b, c}, x] && IGtQ[n, 0]
```

Rule 6416

```
Int[((a_) + ArcSech[(c_)*(x_)]*(b_))/(x_), x_Symbol] :> -Subst[Int[(a +
b*ArcCosh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]
```

Rubi steps

$$\text{integral} = \frac{\text{Subst}\left(\int \frac{\text{sech}^{-1}(ax)}{x} dx, x, x^n\right)}{n}$$

$$= -\frac{\text{Subst}\left(\int \frac{\text{arccosh}\left(\frac{x}{a}\right)}{x} dx, x, x^{-n}\right)}{n}$$

$$\begin{aligned}
&= -\frac{\text{Subst}\left(\int x \tanh(x) dx, x, \text{arccosh}\left(\frac{x^{-n}}{a}\right)\right)}{n} \\
&= \frac{\text{arccosh}\left(\frac{x^{-n}}{a}\right)^2}{2n} - \frac{2\text{Subst}\left(\int \frac{e^{2x} x}{1+e^{2x}} dx, x, \text{arccosh}\left(\frac{x^{-n}}{a}\right)\right)}{n} \\
&= \frac{\text{arccosh}\left(\frac{x^{-n}}{a}\right)^2}{2n} - \frac{\text{arccosh}\left(\frac{x^{-n}}{a}\right) \log\left(1 + e^{2\text{arccosh}\left(\frac{x^{-n}}{a}\right)}\right)}{n} \\
&\quad + \frac{\text{Subst}\left(\int \log(1 + e^{2x}) dx, x, \text{arccosh}\left(\frac{x^{-n}}{a}\right)\right)}{n} \\
&= \frac{\text{arccosh}\left(\frac{x^{-n}}{a}\right)^2}{2n} - \frac{\text{arccosh}\left(\frac{x^{-n}}{a}\right) \log\left(1 + e^{2\text{arccosh}\left(\frac{x^{-n}}{a}\right)}\right)}{n} \\
&\quad + \frac{\text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2\text{arccosh}\left(\frac{x^{-n}}{a}\right)}\right)}{2n} \\
&= \frac{\text{arccosh}\left(\frac{x^{-n}}{a}\right)^2}{2n} - \frac{\text{arccosh}\left(\frac{x^{-n}}{a}\right) \log\left(1 + e^{2\text{arccosh}\left(\frac{x^{-n}}{a}\right)}\right)}{n} - \frac{\text{PolyLog}\left(2, -e^{2\text{arccosh}\left(\frac{x^{-n}}{a}\right)}\right)}{2n}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 219 vs. $2(61) = 122$.

Time = 1.09 (sec) , antiderivative size = 219, normalized size of antiderivative = 3.59

$$\int \frac{\text{sech}^{-1}(ax^n)}{x} dx = \text{sech}^{-1}(ax^n) \log(x) + \frac{\sqrt{\frac{1-ax^n}{1+ax^n}} (4\sqrt{-1+a^2x^{2n}} \arctan(\sqrt{-1+a^2x^{2n}}) (2n \log(x) - \log(a^2x^{2n})) + \sqrt{1-a^2x^{2n}} (\log^2(a^2x^{2n}) - 4 \log(a^2x^{2n})))}{8(n - a^n x^n)}$$

[In] Integrate[ArcSech[a*x^n]/x,x]

[Out] ArcSech[a*x^n]*Log[x] + (Sqrt[(1 - a*x^n)/(1 + a*x^n)]*(4*Sqrt[-1 + a^2*x^(2*n)]*ArcTan[Sqrt[-1 + a^2*x^(2*n)]]*(2*n*Log[x] - Log[a^2*x^(2*n)]) + Sqrt[1 - a^2*x^(2*n)]*(Log[a^2*x^(2*n)]^2 - 4*Log[a^2*x^(2*n)]*Log[(1 + Sqrt[1 - a^2*x^(2*n)])]/2] + 2*Log[(1 + Sqrt[1 - a^2*x^(2*n)])]/2]^2 - 4*PolyLog[2, 1/2 - Sqrt[1 - a^2*x^(2*n)]]/2)))/(8*(n - a^n*x^n))

Maple [A] (verified)

Time = 1.07 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.82

method	result
derivativedivides	$\frac{\frac{\operatorname{arcsech}(ax^n)^2}{2} - \operatorname{arcsech}(ax^n) \ln\left(1 + \left(\frac{x^{-n}}{a} + \sqrt{\frac{x^{-n}}{a} - 1} \sqrt{\frac{x^{-n}}{a} + 1}\right)^2\right)}{n} - \frac{\operatorname{polylog}\left(2, -\left(\frac{x^{-n}}{a} + \sqrt{\frac{x^{-n}}{a} - 1} \sqrt{\frac{x^{-n}}{a} + 1}\right)^2\right)}{2}}$
default	$\frac{\frac{\operatorname{arcsech}(ax^n)^2}{2} - \operatorname{arcsech}(ax^n) \ln\left(1 + \left(\frac{x^{-n}}{a} + \sqrt{\frac{x^{-n}}{a} - 1} \sqrt{\frac{x^{-n}}{a} + 1}\right)^2\right)}{n} - \frac{\operatorname{polylog}\left(2, -\left(\frac{x^{-n}}{a} + \sqrt{\frac{x^{-n}}{a} - 1} \sqrt{\frac{x^{-n}}{a} + 1}\right)^2\right)}{2}}$

[In] int(arcsech(a*x^n)/x,x,method=_RETURNVERBOSE)

[Out] 1/n*(1/2*arcsech(a*x^n)^2-arcsech(a*x^n)*ln(1+(1/a/(x^n)+(1/a/(x^n)-1)^(1/2))*(1/a/(x^n)+1)^(1/2))^2)-1/2*polylog(2,-(1/a/(x^n)+(1/a/(x^n)-1)^(1/2))*(1/a/(x^n)+1)^(1/2))^2))

Fricas [F(-2)]

Exception generated.

$$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx = \text{Exception raised: TypeError}$$

[In] integrate(arcsech(a*x^n)/x,x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{asech}(ax^n)}{x} dx$$

[In] integrate(asech(a*x**n)/x,x)

[Out] Integral(asech(a*x**n)/x, x)

Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{arosech}(ax^n)}{x} dx$$

[In] integrate(arcsech(a*x^n)/x,x, algorithm="maxima")

[Out] a^2*n*integrate(x^(2*n)*log(x)/(a^2*x*x^(2*n) + (a^2*x*x^(2*n) - x)*sqrt(a*x^n + 1)*sqrt(-a*x^n + 1) - x), x) + n*integrate(1/2*log(x)/(a*x*x^n + x), x) - n*integrate(1/2*log(x)/(a*x*x^n - x), x) + log(sqrt(a*x^n + 1)*sqrt(-a*x^n + 1) + 1)*log(x) - log(a)*log(x) - log(x)*log(x^n)

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{arosech}(ax^n)}{x} dx$$

[In] integrate(arcsech(a*x^n)/x,x, algorithm="giac")

[Out] integrate(arcsech(a*x^n)/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{ax^n}\right)}{x} dx$$

[In] int(acosh(1/(a*x^n))/x,x)

[Out] int(acosh(1/(a*x^n))/x, x)

3.30 $\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx$

Optimal result	251
Rubi [A] (verified)	251
Mathematica [A] (verified)	253
Maple [F]	253
Fricas [F]	253
Sympy [F]	254
Maxima [F]	254
Giac [F]	254
Mupad [F(-1)]	254

Optimal result

Integrand size = 10, antiderivative size = 54

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = \frac{1}{10} \operatorname{sech}^{-1}(ax^5)^2 - \frac{1}{5} \operatorname{sech}^{-1}(ax^5) \log\left(1 + e^{2\operatorname{sech}^{-1}(ax^5)}\right) - \frac{1}{10} \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(ax^5)}\right)$$

[Out] 1/10*arcsech(a*x^5)^2-1/5*arcsech(a*x^5)*ln(1+(1/a/x^5+(1/a/x^5-1)^(1/2))*(1/a/x^5+1)^(1/2))^2)-1/10*polylog(2,-(1/a/x^5+(1/a/x^5-1)^(1/2))*(1/a/x^5+1)^(1/2))^2)

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6416, 5882, 3799, 2221, 2317, 2438}

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = -\frac{1}{10} \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(ax^5)}\right) + \frac{1}{10} \operatorname{sech}^{-1}(ax^5)^2 - \frac{1}{5} \operatorname{sech}^{-1}(ax^5) \log\left(e^{2\operatorname{sech}^{-1}(ax^5)} + 1\right)$$

[In] Int[ArcSech[a*x^5]/x,x]

[Out] ArcSech[a*x^5]^2/10 - (ArcSech[a*x^5]*Log[1 + E^(2*ArcSech[a*x^5])])/5 - PolyLog[2, -E^(2*ArcSech[a*x^5])]/10

Rule 2221

Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] :> Simp

```
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^n)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_)*((d_) + (e_)*(x_)^n)]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3799

```
Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol]
:> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

Rule 5882

```
Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)/(x_), x_Symbol] :> Dist[1/b, Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]
```

Rule 6416

```
Int[((a_) + ArcSech[(c_)*(x_)]*(b_))/(x_), x_Symbol] :> -Subst[Int[(a + b*ArcCosh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{5} \text{Subst} \left(\int \frac{\text{sech}^{-1}(ax)}{x} dx, x, x^5 \right) \\
 &= - \left(\frac{1}{5} \text{Subst} \left(\int \frac{\text{arccosh}\left(\frac{x}{a}\right)}{x} dx, x, \frac{1}{x^5} \right) \right) \\
 &= - \left(\frac{1}{5} \text{Subst} \left(\int x \tanh(x) dx, x, \text{sech}^{-1}(ax^5) \right) \right) \\
 &= \frac{1}{10} \text{sech}^{-1}(ax^5)^2 - \frac{2}{5} \text{Subst} \left(\int \frac{e^{2x} x}{1 + e^{2x}} dx, x, \text{sech}^{-1}(ax^5) \right)
 \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{10} \operatorname{sech}^{-1}(ax^5)^2 - \frac{1}{5} \operatorname{sech}^{-1}(ax^5) \log\left(1 + e^{2\operatorname{sech}^{-1}(ax^5)}\right) \\
&\quad + \frac{1}{5} \operatorname{Subst}\left(\int \log(1 + e^{2x}) dx, x, \operatorname{sech}^{-1}(ax^5)\right) \\
&= \frac{1}{10} \operatorname{sech}^{-1}(ax^5)^2 - \frac{1}{5} \operatorname{sech}^{-1}(ax^5) \log\left(1 + e^{2\operatorname{sech}^{-1}(ax^5)}\right) \\
&\quad + \frac{1}{10} \operatorname{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2\operatorname{sech}^{-1}(ax^5)}\right) \\
&= \frac{1}{10} \operatorname{sech}^{-1}(ax^5)^2 - \frac{1}{5} \operatorname{sech}^{-1}(ax^5) \log\left(1 + e^{2\operatorname{sech}^{-1}(ax^5)}\right) - \frac{1}{10} \operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(ax^5)}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.91

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = \frac{1}{10} \left(-\operatorname{sech}^{-1}(ax^5) \left(\operatorname{sech}^{-1}(ax^5) + 2 \log\left(1 + e^{-2\operatorname{sech}^{-1}(ax^5)}\right) \right) + \operatorname{PolyLog}\left(2, -e^{-2\operatorname{sech}^{-1}(ax^5)}\right) \right)$$

[In] Integrate[ArcSech[a*x^5]/x,x]

[Out] $(-\operatorname{ArcSech}[a*x^5]*(\operatorname{ArcSech}[a*x^5] + 2*\operatorname{Log}[1 + E^{(-2*\operatorname{ArcSech}[a*x^5])}])) + \operatorname{PolyLog}[2, -E^{(-2*\operatorname{ArcSech}[a*x^5])}])/10$

Maple [F]

$$\int \frac{\operatorname{arcsech}(ax^5)}{x} dx$$

[In] int(arcsech(a*x^5)/x,x)

[Out] int(arcsech(a*x^5)/x,x)

Fricas [F]

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arsech}(ax^5)}{x} dx$$

[In] integrate(arcsech(a*x^5)/x,x, algorithm="fricas")

[Out] integral(arcsech(a*x^5)/x, x)

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arsech}(ax^5)}{x} dx$$

[In] integrate(arsech(a*x**5)/x,x)

[Out] Integral(arsech(a*x**5)/x, x)

Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arsech}(ax^5)}{x} dx$$

[In] integrate(arcsech(a*x^5)/x,x, algorithm="maxima")

[Out] integrate(arcsech(a*x^5)/x, x)

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arsech}(ax^5)}{x} dx$$

[In] integrate(arcsech(a*x^5)/x,x, algorithm="giac")

[Out] integrate(arcsech(a*x^5)/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{ax^5}\right)}{x} dx$$

[In] int(acosh(1/(a*x^5))/x,x)

[Out] int(acosh(1/(a*x^5))/x, x)

3.31 $\int \operatorname{sech}^{-1}(ce^{a+bx}) dx$

Optimal result	255
Rubi [A] (verified)	255
Mathematica [B] (verified)	257
Maple [A] (verified)	258
Fricas [F(-2)]	258
Sympy [F]	259
Maxima [F]	259
Giac [F]	259
Mupad [F(-1)]	259

Optimal result

Integrand size = 10, antiderivative size = 77

$$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx = \frac{\operatorname{sech}^{-1}(ce^{a+bx})^2}{2b} - \frac{\operatorname{sech}^{-1}(ce^{a+bx}) \log\left(1 + e^{2\operatorname{sech}^{-1}(ce^{a+bx})}\right)}{b} - \frac{\operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(ce^{a+bx})}\right)}{2b}$$

[Out] $1/2*\operatorname{arcsech}(c*\exp(b*x+a))^2/b - \operatorname{arcsech}(c*\exp(b*x+a))*\ln(1+(1/c/\exp(b*x+a)+(1/c/\exp(b*x+a)-1)^{(1/2)*(1/c/\exp(b*x+a)+1)^{(1/2)})^2)/b - 1/2*\operatorname{polylog}(2, -(1/c/\exp(b*x+a)+(1/c/\exp(b*x+a)-1)^{(1/2)*(1/c/\exp(b*x+a)+1)^{(1/2)})^2)/b$

Rubi [A] (verified)

Time = 0.07 (sec), antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {2320, 6416, 5882, 3799, 2221, 2317, 2438}

$$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx = -\frac{\operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(ce^{a+bx})}\right)}{2b} + \frac{\operatorname{sech}^{-1}(ce^{a+bx})^2}{2b} - \frac{\operatorname{sech}^{-1}(ce^{a+bx}) \log\left(e^{2\operatorname{sech}^{-1}(ce^{a+bx})} + 1\right)}{b}$$

[In] $\operatorname{Int}[\operatorname{ArcSech}[c*E^{(a + b*x)}], x]$

[Out] $\operatorname{ArcSech}[c*E^{(a + b*x)}]^2/(2*b) - (\operatorname{ArcSech}[c*E^{(a + b*x)}]*\operatorname{Log}[1 + E^{(2*\operatorname{ArcSech}[c*E^{(a + b*x)})}]])/b - \operatorname{PolyLog}[2, -E^{(2*\operatorname{ArcSech}[c*E^{(a + b*x)})}]]/(2*b)$

Rule 2221

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)
))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol]
:= Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*
(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3799

```
Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_])*(f_)*(x_)], x
_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(
c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x))))], x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

Rule 5882

```
Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)/(x_), x_Symbol] := Dist[1/b,
Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a,
b, c}, x] && IGtQ[n, 0]
```

Rule 6416

```
Int[((a_) + ArcSech[(c_)*(x_)])*(b_)^(n_)/(x_), x_Symbol] := -Subst[Int[(a +
b*ArcCosh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]
```


Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int \frac{\text{sech}^{-1}(cx)}{x} dx, x, e^{a+bx}\right)}{b} \\
 &= -\frac{\text{Subst}\left(\int \frac{\text{arccosh}\left(\frac{x}{c}\right)}{x} dx, x, e^{-a-bx}\right)}{b} \\
 &= -\frac{\text{Subst}\left(\int x \tanh(x) dx, x, \text{arccosh}\left(\frac{e^{-a-bx}}{c}\right)\right)}{b} \\
 &= \frac{\text{arccosh}\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} - \frac{2\text{Subst}\left(\int \frac{e^{2x}x}{1+e^{2x}} dx, x, \text{arccosh}\left(\frac{e^{-a-bx}}{c}\right)\right)}{b} \\
 &= \frac{\text{arccosh}\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} - \frac{\text{arccosh}\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 + e^{2\text{arccosh}\left(\frac{e^{-a-bx}}{c}\right)}\right)}{b} \\
 &\quad + \frac{\text{Subst}\left(\int \log(1 + e^{2x}) dx, x, \text{arccosh}\left(\frac{e^{-a-bx}}{c}\right)\right)}{b} \\
 &= \frac{\text{arccosh}\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} - \frac{\text{arccosh}\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 + e^{2\text{arccosh}\left(\frac{e^{-a-bx}}{c}\right)}\right)}{b} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2\text{arccosh}\left(\frac{e^{-a-bx}}{c}\right)}\right)}{2b} \\
 &= \frac{\text{arccosh}\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} - \frac{\text{arccosh}\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 + e^{2\text{arccosh}\left(\frac{e^{-a-bx}}{c}\right)}\right)}{b} \\
 &\quad - \frac{\text{PolyLog}\left(2, -e^{2\text{arccosh}\left(\frac{e^{-a-bx}}{c}\right)}\right)}{2b}
 \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 249 vs. 2(77) = 154.

Time = 1.86 (sec) , antiderivative size = 249, normalized size of antiderivative = 3.23

$$\begin{aligned}
 \int \text{sech}^{-1}(ce^{a+bx}) dx &= x \text{sech}^{-1}(ce^{a+bx}) \\
 &\quad - \frac{\sqrt{\frac{1-ce^{a+bx}}{1+ce^{a+bx}}} \sqrt{1+ce^{a+bx}} \left(\arctanh\left(\sqrt{1-c^2e^{2(a+bx)}}\right) (8bx - 4 \log(c^2e^{2(a+bx)})) - \log^2(c^2e^{2(a+bx)}) + 4 \log\right)}{2b}
 \end{aligned}$$

```
[In] Integrate[ArcSech[c*E^(a + b*x)],x]
```

```
[Out] x*ArcSech[c*E^(a + b*x)] - (Sqrt[(1 - c*E^(a + b*x))/(1 + c*E^(a + b*x))]*Sqrt[1 + c*E^(a + b*x)]*(ArcTanh[Sqrt[1 - c^2*E^(2*(a + b*x))]]*(8*b*x - 4*Log[c^2*E^(2*(a + b*x))]) - Log[c^2*E^(2*(a + b*x))]^2 + 4*Log[c^2*E^(2*(a + b*x))]*Log[(1 + Sqrt[1 - c^2*E^(2*(a + b*x))])/2] - 2*Log[(1 + Sqrt[1 - c^2*E^(2*(a + b*x))])/2]^2 + 4*PolyLog[2, (1 - Sqrt[1 - c^2*E^(2*(a + b*x))])/2]))/(8*b*Sqrt[1 - c*E^(a + b*x)])
```

Maple [A] (verified)

Time = 1.10 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.75

method	result
derivativedivides	$\frac{\frac{\operatorname{arcsech}\left(\frac{e^{bx+a}}{c}\right)^2}{2} - \operatorname{arcsech}\left(\frac{e^{bx+a}}{c}\right) \ln\left(1 + \left(\frac{e^{-bx-a}}{c} + \sqrt{\frac{e^{-bx-a}}{c} - 1} \sqrt{\frac{e^{-bx-a}}{c} + 1}\right)^2\right)}{b} - \frac{\operatorname{polylog}\left(2, -\left(\frac{e^{-bx-a}}{c} + \sqrt{\frac{e^{-bx-a}}{c} + 1}\right)\right)}{2}}$
default	$\frac{\frac{\operatorname{arcsech}\left(\frac{e^{bx+a}}{c}\right)^2}{2} - \operatorname{arcsech}\left(\frac{e^{bx+a}}{c}\right) \ln\left(1 + \left(\frac{e^{-bx-a}}{c} + \sqrt{\frac{e^{-bx-a}}{c} - 1} \sqrt{\frac{e^{-bx-a}}{c} + 1}\right)^2\right)}{b} - \frac{\operatorname{polylog}\left(2, -\left(\frac{e^{-bx-a}}{c} + \sqrt{\frac{e^{-bx-a}}{c} + 1}\right)\right)}{2}}$

```
[In] int(arcsech(exp(b*x+a)*c),x,method=_RETURNVERBOSE)
```

```
[Out] 1/b*(1/2*arcsech(exp(b*x+a)*c)^2-arcsech(exp(b*x+a)*c)*ln(1+(1/c/exp(b*x+a)+(1/c/exp(b*x+a)-1)^(1/2)*(1/c/exp(b*x+a)+1)^(1/2))^2)-1/2*polylog(2,-(1/c/exp(b*x+a)+(1/c/exp(b*x+a)-1)^(1/2)*(1/c/exp(b*x+a)+1)^(1/2))^2))
```

Fricas [F(-2)]

Exception generated.

$$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx = \text{Exception raised: TypeError}$$

```
[In] integrate(arcsech(c*exp(b*x+a)),x, algorithm="fricas")
```

```
[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)
```

Sympy [F]

$$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx = \int \operatorname{asech}(ce^{a+bx}) dx$$

[In] integrate(asech(c*exp(b*x+a)),x)

[Out] Integral(asech(c*exp(a + b*x)), x)

Maxima [F]

$$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx = \int \operatorname{arsech}(ce^{(bx+a)}) dx$$

[In] integrate(arcsech(c*exp(b*x+a)),x, algorithm="maxima")

[Out] b*c^2*integrate(x*e^(2*b*x + 2*a)/(c^2*e^(2*b*x + 2*a) + (c^2*e^(2*b*x + 2*a) - 1)*e^(1/2*log(c*e^(b*x + a) + 1) + 1/2*log(-c*e^(b*x + a) + 1)) - 1), x) - 1/2*b*x^2 - (a + log(c))*x + x*log(sqrt(c*e^(b*x + a) + 1)*sqrt(-c*e^(b*x + a) + 1) + 1) - 1/2*(b*x*log(c*e^(b*x + a) + 1) + dilog(-c*e^(b*x + a)))/b - 1/2*(b*x*log(-c*e^(b*x + a) + 1) + dilog(c*e^(b*x + a)))/b

Giac [F]

$$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx = \int \operatorname{arsech}(ce^{(bx+a)}) dx$$

[In] integrate(arcsech(c*exp(b*x+a)),x, algorithm="giac")

[Out] integrate(arcsech(c*e^(b*x + a)), x)

Mupad [F(-1)]

Timed out.

$$\int \operatorname{sech}^{-1}(ce^{a+bx}) dx = \int \operatorname{acosh}\left(\frac{e^{-a-bx}}{c}\right) dx$$

[In] int(acosh(exp(- a - b*x)/c),x)

[Out] int(acosh(exp(- a - b*x)/c), x)

3.32 $\int e^{\operatorname{sech}^{-1}(ax)} x^4 dx$

Optimal result	260
Rubi [A] (verified)	260
Mathematica [A] (verified)	262
Maple [A] (verified)	262
Fricas [A] (verification not implemented)	262
Sympy [F]	263
Maxima [A] (verification not implemented)	263
Giac [F(-2)]	263
Mupad [B] (verification not implemented)	264

Optimal result

Integrand size = 10, antiderivative size = 64

$$\int e^{\operatorname{sech}^{-1}(ax)} x^4 dx = -\frac{2e^{\operatorname{sech}^{-1}(ax)} x}{15a^4} + \frac{x^2}{15a^3} - \frac{e^{\operatorname{sech}^{-1}(ax)} x^3}{15a^2} + \frac{x^4}{20a} + \frac{1}{5} e^{\operatorname{sech}^{-1}(ax)} x^5$$

[Out] $-2/15*(1/a/x+(1/a/x-1)^{(1/2)}*(1+1/a/x)^{(1/2)})*x/a^4+1/15*x^2/a^3-1/15*(1/a/x+(1/a/x-1)^{(1/2)}*(1+1/a/x)^{(1/2)})*x^3/a^2+1/20*x^4/a+1/5*(1/a/x+(1/a/x-1)^{(1/2)}*(1+1/a/x)^{(1/2)})*x^5$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.30, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6470, 30, 102, 12, 75}

$$\int e^{\operatorname{sech}^{-1}(ax)} x^4 dx = -\frac{2\sqrt{1-ax}}{15a^5\sqrt{\frac{1}{ax+1}}} - \frac{x^2\sqrt{1-ax}}{15a^3\sqrt{\frac{1}{ax+1}}} + \frac{1}{5}x^5e^{\operatorname{sech}^{-1}(ax)} + \frac{x^4}{20a}$$

[In] `Int[E^ArcSech[a*x]*x^4,x]`

[Out] $x^4/(20*a) + (E^{\operatorname{ArcSech}[a*x]}*x^5)/5 - (2*\sqrt{1-ax})/(15*a^5*\sqrt{[(1+ax)^{-1}]}) - (x^2*\sqrt{1-ax})/(15*a^3*\sqrt{[(1+ax)^{-1}]})$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]`

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 75

Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]

Rule 102

Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Dist[1/(d*f*(m + n + p + 1)), Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]

Rule 6470

Int[E^ArcSech[(a_)*(x_)^(p_)]*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*(E^ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] + Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, m, p}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{5}e^{\text{sech}^{-1}(ax)}x^5 + \frac{\int x^3 dx}{5a} + \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{x^3}{\sqrt{1-ax}\sqrt{1+ax}} dx}{5a} \\
 &= \frac{x^4}{20a} + \frac{1}{5}e^{\text{sech}^{-1}(ax)}x^5 - \frac{x^2\sqrt{1-ax}}{15a^3\sqrt{\frac{1}{1+ax}}} - \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int -\frac{2x}{\sqrt{1-ax}\sqrt{1+ax}} dx}{15a^3} \\
 &= \frac{x^4}{20a} + \frac{1}{5}e^{\text{sech}^{-1}(ax)}x^5 - \frac{x^2\sqrt{1-ax}}{15a^3\sqrt{\frac{1}{1+ax}}} + \frac{\left(2\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{x}{\sqrt{1-ax}\sqrt{1+ax}} dx}{15a^3} \\
 &= \frac{x^4}{20a} + \frac{1}{5}e^{\text{sech}^{-1}(ax)}x^5 - \frac{2\sqrt{1-ax}}{15a^5\sqrt{\frac{1}{1+ax}}} - \frac{x^2\sqrt{1-ax}}{15a^3\sqrt{\frac{1}{1+ax}}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.02

$$\int e^{\operatorname{sech}^{-1}(ax)} x^4 dx = \frac{15a^4 x^4 + 4\sqrt{\frac{1-ax}{1+ax}}(1+ax)^2(-2+2ax-3a^2x^2+3a^3x^3)}{60a^5}$$

[In] Integrate[E^ArcSech[a*x]*x^4,x]

[Out] (15*a^4*x^4 + 4*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2*(-2 + 2*a*x - 3*a^2*x^2 + 3*a^3*x^3))/(60*a^5)

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{\sqrt{\frac{ax+1}{ax}} x \sqrt{-\frac{ax-1}{ax}} (a^2x^2-1)(3a^2x^2+2)}{15a^4} + \frac{x^4}{4a}$	64

[In] int((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x^4,x,method=_RETURNVERBOSE)

[Out] 1/15*((a*x+1)/a/x)^(1/2)*x*(-(a*x-1)/a/x)^(1/2)*(a^2*x^2-1)*(3*a^2*x^2+2)/a^4+1/4*x^4/a

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.02

$$\int e^{\operatorname{sech}^{-1}(ax)} x^4 dx = \frac{15a^3x^4 + 4(3a^4x^5 - a^2x^3 - 2x)\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}}{60a^4}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x^4,x, algorithm="fricas")

[Out] 1/60*(15*a^3*x^4 + 4*(3*a^4*x^5 - a^2*x^3 - 2*x)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)))/a^4

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax)} x^4 dx = \frac{\int x^3 dx + \int ax^4 \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} dx}{a}$$

```
[In] integrate((1/a/x+(1/a/x-1)**(1/2))*(1+1/a/x)**(1/2))*x**4,x)
```

```
[Out] (Integral(x**3, x) + Integral(a*x**4*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)), x))/a
```

Maxima [A] (verification not implemented)

```
none
```

```
Time = 0.24 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.73
```

$$\int e^{\operatorname{sech}^{-1}(ax)} x^4 dx = \frac{x^4}{4a} + \frac{(3a^4x^4 - a^2x^2 - 2)\sqrt{ax+1}\sqrt{-ax+1}}{15a^5}$$

```
[In] integrate((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))*x^4,x, algorithm="maxima")
```

```
[Out] 1/4*x^4/a + 1/15*(3*a^4*x^4 - a^2*x^2 - 2)*sqrt(a*x + 1)*sqrt(-a*x + 1)/a^5
```

Giac [F(-2)]

```
Exception generated.
```

$$\int e^{\operatorname{sech}^{-1}(ax)} x^4 dx = \text{Exception raised: TypeError}$$

```
[In] integrate((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))*x^4,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to roun
ding error%%{1,[0,3,2,2,0,0]}%%+%%{1,[0,2,0,1,1,1]}%% / %%{1,[0,0,2,3,
0,0]}%%
```

Mupad [B] (verification not implemented)

Time = 4.13 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.17

$$\int e^{\operatorname{sech}^{-1}(ax)} x^4 dx = \frac{x^4}{4a} - \sqrt{\frac{1}{ax} - 1} \left(\frac{2x \sqrt{\frac{1}{ax} + 1}}{15a^4} - \frac{x^5 \sqrt{\frac{1}{ax} + 1}}{5} + \frac{x^3 \sqrt{\frac{1}{ax} + 1}}{15a^2} \right)$$

[In] int(x^4*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x)),x)

[Out] x^4/(4*a) - (1/(a*x) - 1)^(1/2)*((2*x*(1/(a*x) + 1)^(1/2))/(15*a^4) - (x^5*(1/(a*x) + 1)^(1/2))/5 + (x^3*(1/(a*x) + 1)^(1/2))/(15*a^2))

3.33 $\int e^{\operatorname{sech}^{-1}(ax)} x^3 dx$

Optimal result	265
Rubi [A] (verified)	265
Mathematica [C] (verified)	267
Maple [C] (verified)	267
Fricas [A] (verification not implemented)	267
Sympy [F]	268
Maxima [F]	268
Giac [F(-2)]	268
Mupad [B] (verification not implemented)	269

Optimal result

Integrand size = 10, antiderivative size = 84

$$\int e^{\operatorname{sech}^{-1}(ax)} x^3 dx = \frac{x^3}{12a} + \frac{1}{4} e^{\operatorname{sech}^{-1}(ax)} x^4 - \frac{x\sqrt{1-ax}}{8a^3 \sqrt{\frac{1}{1+ax}}} + \frac{\sqrt{\frac{1}{1+ax}} \sqrt{1+ax} \arcsin(ax)}{8a^4}$$

[Out] 1/12*x^3/a+1/4*(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x^4-1/8*x*(-a*x+1)^(1/2)/a^3/(1/(a*x+1))^(1/2)+1/8*arcsin(a*x)*(1/(a*x+1))^(1/2)*(a*x+1)^(1/2)/a^4

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6470, 30, 92, 41, 222}

$$\int e^{\operatorname{sech}^{-1}(ax)} x^3 dx = \frac{\sqrt{\frac{1}{ax+1}} \sqrt{ax+1} \arcsin(ax)}{8a^4} - \frac{x\sqrt{1-ax}}{8a^3 \sqrt{\frac{1}{ax+1}}} + \frac{1}{4} x^4 e^{\operatorname{sech}^{-1}(ax)} + \frac{x^3}{12a}$$

[In] Int[E^ArcSech[a*x]*x^3,x]

[Out] x^3/(12*a) + (E^ArcSech[a*x]*x^4)/4 - (x*Sqrt[1 - a*x])/(8*a^3*Sqrt[(1 + a*x)^(-1)]) + (Sqrt[(1 + a*x)^(-1)]*Sqrt[1 + a*x]*ArcSin[a*x])/(8*a^4)

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 41

Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))

Rule 92

Int[((a_) + (b_)*(x_))^(p_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(m_), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]

Rule 222

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

Rule 6470

Int[E^ArcSech[(a_)*(x_)^(p_)]*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*(E^ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] + Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, m, p}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{4}e^{\text{sech}^{-1}(ax)}x^4 + \frac{\int x^2 dx}{4a} + \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{x^2}{\sqrt{1-ax}\sqrt{1+ax}} dx}{4a} \\
 &= \frac{x^3}{12a} + \frac{1}{4}e^{\text{sech}^{-1}(ax)}x^4 - \frac{x\sqrt{1-ax}}{8a^3\sqrt{\frac{1}{1+ax}}} + \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{\sqrt{1-ax}\sqrt{1+ax}} dx}{8a^3} \\
 &= \frac{x^3}{12a} + \frac{1}{4}e^{\text{sech}^{-1}(ax)}x^4 - \frac{x\sqrt{1-ax}}{8a^3\sqrt{\frac{1}{1+ax}}} + \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{\sqrt{1-a^2x^2}} dx}{8a^3} \\
 &= \frac{x^3}{12a} + \frac{1}{4}e^{\text{sech}^{-1}(ax)}x^4 - \frac{x\sqrt{1-ax}}{8a^3\sqrt{\frac{1}{1+ax}}} + \frac{\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \arcsin(ax)}{8a^4}
 \end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.15

$$\int e^{\operatorname{sech}^{-1}(ax)} x^3 dx = \frac{8a^3 x^3 - 3a \sqrt{\frac{1-ax}{1+ax}} (x + ax^2 - 2a^2 x^3 - 2a^3 x^4) + 3i \log \left(-2iax + 2 \sqrt{\frac{1-ax}{1+ax}} (1 + ax) \right)}{24a^4}$$

[In] Integrate[E^ArcSech[a*x]*x^3,x]

[Out] (8*a^3*x^3 - 3*a*Sqrt[(1 - a*x)/(1 + a*x)]*(x + a*x^2 - 2*a^2*x^3 - 2*a^3*x^4) + (3*I)*Log[(-2*I)*a*x + 2*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)]/(24*a^4)

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.04 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.40

method	result	size
default	$\frac{\sqrt{\frac{ax+1}{ax}} x \sqrt{-\frac{ax-1}{ax}} \left(2 \operatorname{csgn}(a) a^3 x^3 \sqrt{-a^2 x^2 + 1} - \sqrt{-a^2 x^2 + 1} x \operatorname{csgn}(a) a + \arctan \left(\frac{\operatorname{csgn}(a) ax}{\sqrt{-a^2 x^2 + 1}} \right) \right) \operatorname{csgn}(a)}{8 \sqrt{-a^2 x^2 + 1} a^3} + \frac{x^3}{3a}$	118

[In] int((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))*x^3,x,method=_RETURNVERBOSE)

[Out] 1/8*((a*x+1)/a/x)^(1/2)*x*(-(a*x-1)/a/x)^(1/2)*(2*csgn(a)*a^3*x^3*(-a^2*x^2+1)^(1/2)-(a^2*x^2+1)^(1/2)*x*csgn(a)*a+arctan(csgn(a)*a*x/(-a^2*x^2+1)^(1/2)))*csgn(a)/(-a^2*x^2+1)^(1/2)/a^3+1/3*x^3/a

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.13

$$\int e^{\operatorname{sech}^{-1}(ax)} x^3 dx = \frac{8a^3 x^3 + 3(2a^4 x^4 - a^2 x^2) \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} - 3 \arctan \left(\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} \right)}{24a^4}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))*x^3,x, algorithm="fricas")

[Out] 1/24*(8*a^3*x^3 + 3*(2*a^4*x^4 - a^2*x^2)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 3*arctan(sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x))))/a^4

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax)} x^3 dx = \frac{\int x^2 dx + \int ax^3 \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} dx}{a}$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))*x**3,x)

[Out] (Integral(x**2, x) + Integral(a*x**3*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)), x))/a

Maxima [F]

$$\int e^{\operatorname{sech}^{-1}(ax)} x^3 dx = \int x^3 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}} \right) dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x^3,x, algorithm="maxima")

[Out] 1/3*x^3/a + integrate(sqrt(a*x + 1)*sqrt(-a*x + 1)*x^2, x)/a

Giac [F(-2)]

Exception generated.

$$\int e^{\operatorname{sech}^{-1}(ax)} x^3 dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x^3,x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Unable to divide, perhaps due to rounding error%%{1,[0,2,2,2,0,0]}%%+%%{1,[0,1,0,1,1,1]}%% / %%{1,[0,0,2,3,0,0]}%%

Mupad [B] (verification not implemented)

Time = 16.44 (sec) , antiderivative size = 521, normalized size of antiderivative = 6.20

$$\int e^{\operatorname{sech}^{-1}(ax)} x^3 dx = \frac{\ln\left(\frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^2}{\left(\sqrt{\frac{1}{ax}+1-1}\right)^2} + 1\right) \operatorname{li}}{8a^4} - \frac{\frac{\operatorname{li}}{1024a^4} + \frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^2 \operatorname{li}}{128a^4 \left(\sqrt{\frac{1}{ax}+1-1}\right)^2} + \frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^4 11i}{512a^4 \left(\sqrt{\frac{1}{ax}+1-1}\right)^4} + \frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^6 7i}{256a^4 \left(\sqrt{\frac{1}{ax}+1-1}\right)^6} - \frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^8 239i}{1024a^4 \left(\sqrt{\frac{1}{ax}+1-1}\right)^8} + \frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^{10}}{256a^4 \left(\sqrt{\frac{1}{ax}+1-1}\right)^{10}}}{\frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^4}{\left(\sqrt{\frac{1}{ax}+1-1}\right)^4} + \frac{4\left(\sqrt{\frac{1}{ax}-1-i}\right)^6}{\left(\sqrt{\frac{1}{ax}+1-1}\right)^6} + \frac{6\left(\sqrt{\frac{1}{ax}-1-i}\right)^8}{\left(\sqrt{\frac{1}{ax}+1-1}\right)^8} + \frac{4\left(\sqrt{\frac{1}{ax}-1-i}\right)^{10}}{\left(\sqrt{\frac{1}{ax}+1-1}\right)^{10}} + \frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^{12}}{\left(\sqrt{\frac{1}{ax}+1-1}\right)^{12}}} - \frac{\ln\left(\frac{\sqrt{\frac{1}{ax}-1-i}}{\sqrt{\frac{1}{ax}+1-1}}\right) \operatorname{li}}{8a^4} + \frac{x^3}{3a} - \frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^2 \operatorname{li}}{256a^4 \left(\sqrt{\frac{1}{ax}+1-1}\right)^2} - \frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^4 \operatorname{li}}{1024a^4 \left(\sqrt{\frac{1}{ax}+1-1}\right)^4}$$

[In] `int(x^3*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x)),x)`

[Out] `(log(((1/(a*x) - 1)^(1/2) - 1i)^2/((1/(a*x) + 1)^(1/2) - 1)^2 + 1)*1i)/(8*a^4) - (1i/(1024*a^4) + (((1/(a*x) - 1)^(1/2) - 1i)^2*1i)/(128*a^4*((1/(a*x) + 1)^(1/2) - 1)^2) + (((1/(a*x) - 1)^(1/2) - 1i)^4*11i)/(512*a^4*((1/(a*x) + 1)^(1/2) - 1)^4) + (((1/(a*x) - 1)^(1/2) - 1i)^6*7i)/(256*a^4*((1/(a*x) + 1)^(1/2) - 1)^6) - (((1/(a*x) - 1)^(1/2) - 1i)^8*239i)/(1024*a^4*((1/(a*x) + 1)^(1/2) - 1)^8) + (((1/(a*x) - 1)^(1/2) - 1i)^10*1i)/(256*a^4*((1/(a*x) + 1)^(1/2) - 1)^10))/(((1/(a*x) - 1)^(1/2) - 1i)^4/((1/(a*x) + 1)^(1/2) - 1)^4 + (4*((1/(a*x) - 1)^(1/2) - 1i)^6)/((1/(a*x) + 1)^(1/2) - 1)^6 + (6*((1/(a*x) - 1)^(1/2) - 1i)^8)/((1/(a*x) + 1)^(1/2) - 1)^8 + (4*((1/(a*x) - 1)^(1/2) - 1i)^10)/((1/(a*x) + 1)^(1/2) - 1)^10 + ((1/(a*x) - 1)^(1/2) - 1i)^12/((1/(a*x) + 1)^(1/2) - 1)^12) - (log(((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1))*1i)/(8*a^4) + x^3/(3*a) - (((1/(a*x) - 1)^(1/2) - 1i)^2*1i)/(256*a^4*((1/(a*x) + 1)^(1/2) - 1)^2) - (((1/(a*x) - 1)^(1/2) - 1i)^4*1i)/(1024*a^4*((1/(a*x) + 1)^(1/2) - 1)^4)`

3.34 $\int e^{\operatorname{sech}^{-1}(ax)} x^2 dx$

Optimal result	270
Rubi [A] (verified)	270
Mathematica [A] (verified)	271
Maple [A] (verified)	271
Fricas [A] (verification not implemented)	272
Sympy [F]	272
Maxima [A] (verification not implemented)	272
Giac [F(-2)]	273
Mupad [B] (verification not implemented)	273

Optimal result

Integrand size = 10, antiderivative size = 38

$$\int e^{\operatorname{sech}^{-1}(ax)} x^2 dx = -\frac{e^{\operatorname{sech}^{-1}(ax)} x}{3a^2} + \frac{x^2}{6a} + \frac{1}{3} e^{\operatorname{sech}^{-1}(ax)} x^3$$

[Out] $-1/3*(1/a/x+(1/a/x-1)^{(1/2)}*(1+1/a/x)^{(1/2)})*x/a^2+1/6*x^2/a+1/3*(1/a/x+(1/a/x-1)^{(1/2)}*(1+1/a/x)^{(1/2)})*x^3$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.37, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {6470, 30, 75}

$$\int e^{\operatorname{sech}^{-1}(ax)} x^2 dx = -\frac{\sqrt{1-ax}}{3a^3 \sqrt{\frac{1}{ax+1}}} + \frac{1}{3} x^3 e^{\operatorname{sech}^{-1}(ax)} + \frac{x^2}{6a}$$

[In] `Int[E^ArcSech[a*x]*x^2,x]`

[Out] $x^2/(6*a) + (E^{\operatorname{ArcSech}[a*x]}*x^3)/3 - \operatorname{Sqrt}[1 - a*x]/(3*a^3*\operatorname{Sqrt}[(1 + a*x)^{-1}])$

Rule 30

`Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]`

Rule 75

`Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/(d*f*(n + p + 1))`

2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]

Rule 6470

Int[E^ArcSech[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*(E^ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] + Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, m, p}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{3} e^{\operatorname{sech}^{-1}(ax)} x^3 + \frac{\int x dx}{3a} + \frac{\left(\sqrt{\frac{1}{1+ax}} \sqrt{1+ax}\right) \int \frac{x}{\sqrt{1-ax}\sqrt{1+ax}} dx}{3a} \\ &= \frac{x^2}{6a} + \frac{1}{3} e^{\operatorname{sech}^{-1}(ax)} x^3 - \frac{\sqrt{1-ax}}{3a^3 \sqrt{\frac{1}{1+ax}}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.26

$$\int e^{\operatorname{sech}^{-1}(ax)} x^2 dx = \frac{3a^2 x^2 + 2(-1 + ax) \sqrt{\frac{1-ax}{1+ax}} (1 + ax)^2}{6a^3}$$

[In] Integrate[E^ArcSech[a*x]*x^2,x]

[Out] (3*a^2*x^2 + 2*(-1 + a*x)*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2)/(6*a^3)

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.42

method	result	size
default	$\frac{\sqrt{\frac{ax+1}{ax}} x \sqrt{-\frac{ax-1}{ax}} (a^2 x^2 - 1)}{3a^2} + \frac{x^2}{2a}$	54

[In] int((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))*x^2,x,method=_RETURNVERBOSE)

[Out] 1/3*((a*x+1)/a/x)^(1/2)*x*(-(a*x-1)/a/x)^(1/2)*(a^2*x^2-1)/a^2+1/2*x^2/a

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.42

$$\int e^{\operatorname{sech}^{-1}(ax)} x^2 dx = \frac{3ax^2 + 2(a^2x^3 - x)\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}}{6a^2}$$

```
[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x^2,x, algorithm="fricas")
```

```
[Out] 1/6*(3*a*x^2 + 2*(a^2*x^3 - x)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)))/a^2
```

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax)} x^2 dx = \frac{\int x dx + \int ax^2 \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} dx}{a}$$

```
[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))*x**2,x)
```

```
[Out] (Integral(x, x) + Integral(a*x**2*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)), x))/a
```

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int e^{\operatorname{sech}^{-1}(ax)} x^2 dx = \frac{x^2}{2a} + \frac{(a^2x^2 - 1)\sqrt{ax + 1}\sqrt{-ax + 1}}{3a^3}$$

```
[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x^2,x, algorithm="maxima")
```

```
[Out] 1/2*x^2/a + 1/3*(a^2*x^2 - 1)*sqrt(a*x + 1)*sqrt(-a*x + 1)/a^3
```


Giac [F(-2)]

Exception generated.

$$\int e^{\operatorname{sech}^{-1}(ax)} x^2 dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))*x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
 UT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{1,[0,1,2,2,0,0]%%}+%%{1,[0,0,0,1,1,1]%%} / %%{1,[0,0,2,3,0,0]%%}

Mupad [B] (verification not implemented)

Time = 5.11 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.45

$$\int e^{\operatorname{sech}^{-1}(ax)} x^2 dx = \sqrt{\frac{1}{ax} - 1} \left(\frac{x^3 \sqrt{\frac{1}{ax} + 1}}{3} - \frac{x \sqrt{\frac{1}{ax} + 1}}{3a^2} \right) + \frac{x^2}{2a}$$

[In] int(x^2*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x)),x)

[Out] (1/(a*x) - 1)^(1/2)*((x^3*(1/(a*x) + 1)^(1/2))/3 - (x*(1/(a*x) + 1)^(1/2))/(3*a^2)) + x^2/(2*a)

3.35 $\int e^{\operatorname{sech}^{-1}(ax)} x dx$

Optimal result	274
Rubi [A] (verified)	274
Mathematica [C] (verified)	275
Maple [C] (verified)	276
Fricas [A] (verification not implemented)	276
Sympy [F]	276
Maxima [F]	277
Giac [F]	277
Mupad [B] (verification not implemented)	277

Optimal result

Integrand size = 8, antiderivative size = 53

$$\int e^{\operatorname{sech}^{-1}(ax)} x dx = \frac{x}{2a} + \frac{1}{2} e^{\operatorname{sech}^{-1}(ax)} x^2 + \frac{\sqrt{\frac{1}{1+ax}} \sqrt{1+ax} \arcsin(ax)}{2a^2}$$

[Out] $1/2*x/a+1/2*(1/a/x+(1/a/x-1)^{(1/2)}*(1+1/a/x)^{(1/2)})*x^2+1/2*\arcsin(a*x)*(1/(a*x+1))^{(1/2)}*(a*x+1)^{(1/2)}/a^2$

Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6470, 8, 41, 222}

$$\int e^{\operatorname{sech}^{-1}(ax)} x dx = \frac{\sqrt{\frac{1}{ax+1}} \sqrt{ax+1} \arcsin(ax)}{2a^2} + \frac{1}{2} x^2 e^{\operatorname{sech}^{-1}(ax)} + \frac{x}{2a}$$

[In] Int[E^ArcSech[a*x]*x,x]

[Out] $x/(2*a) + (E^{\operatorname{ArcSech}[a*x]}*x^2)/2 + (\operatorname{Sqrt}[(1 + a*x)^{-1}]*\operatorname{Sqrt}[1 + a*x]*\operatorname{ArcSin}[a*x])/(2*a^2)$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 41

Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(m_.), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (

IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))

Rule 222

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

Rule 6470

Int[E^ArcSech[(a_)*(x_)^(p_)]*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*(E^ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] + Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, m, p}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{2}e^{\text{sech}^{-1}(ax)}x^2 + \frac{\int 1 dx}{2a} + \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{\sqrt{1-ax}\sqrt{1+ax}} dx}{2a} \\ &= \frac{x}{2a} + \frac{1}{2}e^{\text{sech}^{-1}(ax)}x^2 + \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{\sqrt{1-a^2x^2}} dx}{2a} \\ &= \frac{x}{2a} + \frac{1}{2}e^{\text{sech}^{-1}(ax)}x^2 + \frac{\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \arcsin(ax)}{2a^2} \end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.42

$$\int e^{\text{sech}^{-1}(ax)}x dx = \frac{2ax + ax\sqrt{\frac{1-ax}{1+ax}}(1+ax) + i \log\left(-2iax + 2\sqrt{\frac{1-ax}{1+ax}}(1+ax)\right)}{2a^2}$$

[In] Integrate[E^ArcSech[a*x]*x,x]

[Out] (2*a*x + a*x*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x) + I*Log[(-2*I)*a*x + 2*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)])/(2*a^2)

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.05 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.74

method	result	size
default	$\frac{\sqrt{\frac{ax+1}{ax}} x \sqrt{-\frac{ax-1}{ax}} \left(\sqrt{-a^2x^2+1} x \operatorname{csgn}(a) a + \arctan\left(\frac{\operatorname{csgn}(a)ax}{\sqrt{-a^2x^2+1}}\right) \right) \operatorname{csgn}(a)}{2\sqrt{-a^2x^2+1} a} + \frac{x}{a}$	92

[In] `int((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{2} * \left(\frac{a*x+1}{a/x} \right)^{1/2} * x * \left(-\frac{a*x-1}{a/x} \right)^{1/2} * \left(\sqrt{-a^2*x^2+1} \right)^{1/2} * x * \operatorname{csgn}(a) * a + \arctan\left(\operatorname{csgn}(a) * a * x / \left(-a^2*x^2+1 \right)^{1/2}\right) / \left(-a^2*x^2+1 \right)^{1/2} * \operatorname{csgn}(a) / a + x / a$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.49

$$\int e^{\operatorname{sech}^{-1}(ax)} x dx = \frac{a^2 x^2 \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} + 2ax - \arctan\left(\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}}\right)}{2a^2}$$

[In] `integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x,x, algorithm="fricas")`

[Out] $\frac{1}{2} * (a^2 * x^2 * \operatorname{sqrt}((a*x + 1)/(a*x)) * \operatorname{sqrt}(-(a*x - 1)/(a*x)) + 2*a*x - \operatorname{arctan}(\operatorname{sqrt}((a*x + 1)/(a*x)) * \operatorname{sqrt}(-(a*x - 1)/(a*x)))) / a^2$

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax)} x dx = \frac{\int 1 dx + \int ax \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} dx}{a}$$

[In] `integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))*x,x)`

[Out] $(\operatorname{Integral}(1, x) + \operatorname{Integral}(a*x*\operatorname{sqrt}(-1 + 1/(a*x))*\operatorname{sqrt}(1 + 1/(a*x)), x))/a$

Maxima [F]

$$\int e^{\operatorname{sech}^{-1}(ax)} x dx = \int x \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} \right) dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x,x, algorithm="maxima")

[Out] x/a + integrate(sqrt(a*x + 1)*sqrt(-a*x + 1), x)/a

Giac [F]

$$\int e^{\operatorname{sech}^{-1}(ax)} x dx = \int x \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} \right) dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x,x, algorithm="giac")

[Out] integrate(x*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x)), x)

Mupad [B] (verification not implemented)

Time = 10.97 (sec) , antiderivative size = 303, normalized size of antiderivative = 5.72

$$\begin{aligned} \int e^{\operatorname{sech}^{-1}(ax)} x dx = & \frac{\ln \left(\frac{\left(\sqrt{\frac{1}{ax} - 1 - i} \right)^2}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^2} + 1 \right) \operatorname{li}}{2a^2} - \frac{\ln \left(\frac{\sqrt{\frac{1}{ax} - 1 - i}}{\sqrt{\frac{1}{ax} + 1 - 1}} \right) \operatorname{li}}{2a^2} \\ & + \frac{\frac{\operatorname{li}}{32a^2} + \frac{\left(\sqrt{\frac{1}{ax} - 1 - i} \right)^2 \operatorname{li}}{16a^2 \left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^2} - \frac{\left(\sqrt{\frac{1}{ax} - 1 - i} \right)^4 15i}{32a^2 \left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^4}}{\frac{\left(\sqrt{\frac{1}{ax} - 1 - i} \right)^2}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^2} + \frac{2 \left(\sqrt{\frac{1}{ax} - 1 - i} \right)^4}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^4} + \frac{\left(\sqrt{\frac{1}{ax} - 1 - i} \right)^6}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^6}} \\ & + \frac{x}{a} + \frac{\left(\sqrt{\frac{1}{ax} - 1 - i} \right)^2 \operatorname{li}}{32a^2 \left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^2} \end{aligned}$$

[In] int(x*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x)),x)

[Out] (log(((1/(a*x) - 1)^(1/2) - 1i)^2/((1/(a*x) + 1)^(1/2) - 1)^2 + 1)*1i)/(2*a^2) - (log(((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1))*1i)/(2*a^2) + (1i/(32*a^2) + (((1/(a*x) - 1)^(1/2) - 1i)^2*1i)/(16*a^2*((1/(a*x) + 1)^(1/2) - 1)^2) - (((1/(a*x) - 1)^(1/2) - 1i)^4*15i)/(32*a^2*((1/(a*x) + 1)^(1/2) - 1)^4))/(((1/(a*x) - 1)^(1/2) - 1i)^2/((1/(a*x) + 1)^(1/2) - 1)^2 + (2*((1/(a*x) - 1)^(1/2) - 1i)^4)/((1/(a*x) + 1)^(1/2) - 1)^4 + ((1/(a*x) - 1)^(1/2) - 1i)^6/((1/(a*x) + 1)^(1/2) - 1)^6) + x/a + (((1/(a*x) - 1)^(1/2) - 1i)^2*1i)/(32*a^2*((1/(a*x) + 1)^(1/2) - 1)^2)

3.36 $\int e^{\operatorname{sech}^{-1}(ax)} dx$

Optimal result	278
Rubi [A] (verified)	278
Mathematica [B] (verified)	279
Maple [A] (verified)	280
Fricas [B] (verification not implemented)	280
Sympy [F]	280
Maxima [F]	281
Giac [F]	281
Mupad [B] (verification not implemented)	281

Optimal result

Integrand size = 6, antiderivative size = 24

$$\int e^{\operatorname{sech}^{-1}(ax)} dx = e^{\operatorname{sech}^{-1}(ax)} x - \frac{\operatorname{sech}^{-1}(ax)}{a} + \frac{\log(x)}{a}$$

[Out] $(1/a/x + (1/a/x - 1)^{1/2} * (1 + 1/a/x)^{1/2}) * x - \operatorname{arcsech}(a*x)/a + \ln(x)/a$

Rubi [A] (verified)

Time = 0.08 (sec), antiderivative size = 39, normalized size of antiderivative = 1.62, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6464, 1984, 214}

$$\int e^{\operatorname{sech}^{-1}(ax)} dx = -\frac{2 \operatorname{arctanh}\left(\sqrt{\frac{1-ax}{ax+1}}\right)}{a} + \frac{\log(x)}{a} + x e^{\operatorname{sech}^{-1}(ax)}$$

[In] `Int[E^ArcSech[a*x],x]`

[Out] `E^ArcSech[a*x]*x - (2*ArcTanh[Sqrt[(1 - a*x)/(1 + a*x)]])/a + Log[x]/a`

Rule 214

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Rule 1984

`Int[(u_)^(r_.)*(x_)^(m_.)*(((e_.)*((a_.) + (b_.)*(x_)^(n_.)))/((c_) + (d_.)*(x_)^(n_.)))^(p_), x_Symbol] := With[{q = Denominator[p]}, Dist[q*e*((b*c - a*d)/n), Subst[Int[SimplifyIntegrand[x^(q*(p + 1) - 1)*(((a)*e + c*x^q)^`

```
((m + 1)/n - 1)/(b*e - d*x^q)^((m + 1)/n + 1))*(u /. x -> ((-a)*e + c*x^q)^
(1/n)/(b*e - d*x^q)^(1/n))^r, x], x], x, (e*((a + b*x^n)/(c + d*x^n)))^(1/q
)], x]] /; FreeQ[{a, b, c, d, e}, x] && PolynomialQ[u, x] && FractionQ[p] &
& IntegerQ[1/n] && IntegersQ[m, r]
```

Rule 6464

```
Int[E^ArcSech[(a_.)*(x_)], x_Symbol] :> Simp[x*E^ArcSech[a*x], x] + (Dist[1
/a, Int[(1/(x*(1 - a*x)))*Sqrt[(1 - a*x)/(1 + a*x)], x], x] + Simp[Log[x]/a
, x]) /; FreeQ[a, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= e^{\operatorname{sech}^{-1}(ax)}x + \frac{\log(x)}{a} + \frac{\int \frac{\sqrt{\frac{1-ax}{1+ax}}}{x(1-ax)} dx}{a} \\ &= e^{\operatorname{sech}^{-1}(ax)}x + \frac{\log(x)}{a} - 4\operatorname{Subst}\left(\int \frac{1}{2a - 2ax^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right) \\ &= e^{\operatorname{sech}^{-1}(ax)}x - \frac{2\operatorname{arctanh}\left(\sqrt{\frac{1-ax}{1+ax}}\right)}{a} + \frac{\log(x)}{a} \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 79 vs. $2(24) = 48$.

Time = 0.05 (sec) , antiderivative size = 79, normalized size of antiderivative = 3.29

$$\int e^{\operatorname{sech}^{-1}(ax)} dx = \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax) + 2\log(ax) - \log\left(1 + \sqrt{\frac{1-ax}{1+ax}} + ax\sqrt{\frac{1-ax}{1+ax}}\right)}{a}$$

```
[In] Integrate[E^ArcSech[a*x], x]
```

```
[Out] (Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x) + 2*Log[a*x] - Log[1 + Sqrt[(1 - a*x)/
(1 + a*x)] + a*x*Sqrt[(1 - a*x)/(1 + a*x)]])/a
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 80, normalized size of antiderivative = 3.33

method	result	size
default	$\frac{\ln(x)}{a} - \frac{\sqrt{-\frac{ax-1}{ax}} x \sqrt{\frac{ax+1}{ax}} \left(-\sqrt{-a^2x^2+1} + \operatorname{arctanh}\left(\frac{1}{\sqrt{-a^2x^2+1}}\right) \right)}{\sqrt{-a^2x^2+1}}$	80

[In] int(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2),x,method=_RETURNVERBOSE)

[Out] ln(x)/a-(-(a*x-1)/a/x)^(1/2)*x*((a*x+1)/a/x)^(1/2)*(-(-a^2*x^2+1)^(1/2)+arc tanh(1/(-a^2*x^2+1)^(1/2)))/(-a^2*x^2+1)^(1/2)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 115 vs. 2(49) = 98.

Time = 0.26 (sec) , antiderivative size = 115, normalized size of antiderivative = 4.79

$$\int e^{\operatorname{sech}^{-1}(ax)} dx = \frac{2ax\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} - \log\left(ax\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} + 1\right) + \log\left(ax\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} - 1\right) + 2\log(x)}{2a}$$

[In] integrate(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2),x, algorithm="fricas")

[Out] 1/2*(2*a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + 1) + log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 1) + 2*log(x))/a

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax)} dx = \frac{\int \frac{1}{x} dx + \int a\sqrt{-1 + \frac{1}{ax}}\sqrt{1 + \frac{1}{ax}} dx}{a}$$

[In] integrate(1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2),x)

[Out] (Integral(1/x, x) + Integral(a*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)), x))/a

Maxima [F]

$$\int e^{\operatorname{sech}^{-1}(ax)} dx = \int \sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} dx$$

[In] integrate(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x), x)

Giac [F]

$$\int e^{\operatorname{sech}^{-1}(ax)} dx = \int \sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} dx$$

[In] integrate(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x), x)

Mupad [B] (verification not implemented)

Time = 6.31 (sec) , antiderivative size = 182, normalized size of antiderivative = 7.58

$$\int e^{\operatorname{sech}^{-1}(ax)} dx = \frac{\ln(x)}{a} - \frac{4 \operatorname{atanh}\left(\frac{\sqrt{\frac{1}{ax}-1-i}}{\sqrt{\frac{1}{ax}+1-1}}\right)}{a} + \frac{\frac{5\left(\sqrt{\frac{1}{ax}-1-i}\right)^2}{\left(\sqrt{\frac{1}{ax}+1-1}\right)^2} + 1}{\frac{4a\left(\sqrt{\frac{1}{ax}-1-i}\right)}{\sqrt{\frac{1}{ax}+1-1}} + \frac{4a\left(\sqrt{\frac{1}{ax}-1-i}\right)^3}{\left(\sqrt{\frac{1}{ax}+1-1}\right)^3}} + \frac{\sqrt{\frac{1}{ax}-1-i}}{4a\left(\sqrt{\frac{1}{ax}+1-1}\right)}$$

[In] int((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x),x)

[Out] log(x)/a - (4*atanh(((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1)))/a + ((5*((1/(a*x) - 1)^(1/2) - 1i)^2)/((1/(a*x) + 1)^(1/2) - 1)^2 + 1)/((4*a*((1/(a*x) - 1)^(1/2) - 1i))/((1/(a*x) + 1)^(1/2) - 1) + (4*a*((1/(a*x) - 1)^(1/2) - 1i)^3)/((1/(a*x) + 1)^(1/2) - 1)^3 + ((1/(a*x) - 1)^(1/2) - 1i)/(4*a*((1/(a*x) + 1)^(1/2) - 1)))

3.37 $\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x} dx$

Optimal result	282
Rubi [A] (verified)	282
Mathematica [C] (verified)	284
Maple [C] (verified)	284
Fricas [A] (verification not implemented)	284
Sympy [F]	285
Maxima [F]	285
Giac [F]	285
Mupad [B] (verification not implemented)	285

Optimal result

Integrand size = 10, antiderivative size = 48

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x} dx = -\frac{2}{1 - \sqrt{\frac{1-ax}{1+ax}}} + 2 \arctan \left(\sqrt{\frac{1-ax}{1+ax}} \right)$$

[Out] 2*arctan(((−a*x+1)/(a*x+1))^(1/2))-2/(1-((−a*x+1)/(a*x+1))^(1/2))

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.33, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6469, 99, 12, 41, 222}

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x} dx = -\sqrt{\frac{1}{ax+1}} \sqrt{ax+1} \arcsin(ax) - \frac{\sqrt{1-ax}}{ax \sqrt{\frac{1}{ax+1}}} - \frac{1}{ax}$$

[In] Int[E^ArcSech[a*x]/x,x]

[Out] -(1/(a*x)) - Sqrt[1 - a*x]/(a*x*Sqrt[(1 + a*x)^(-1)]) - Sqrt[(1 + a*x)^(-1)]*Sqrt[1 + a*x]*ArcSin[a*x]

Rule 12

Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 41

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(m_), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))
```

Rule 99

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p/(b*(m + 1)), x] - Dist[1/(b*(m + 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^(p - 1)*Simp[d*e*n + c*f*p + d*f*(n + p)*x, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[m, -1] && GtQ[n, 0] && GtQ[p, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])
```

Rule 222

```
Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

Rule 6469

```
Int[E^ArcSech[(a_)*(x_)^(p_)]/(x_), x_Symbol] := -Simp[(a*p*x^p)^(-1), x] + Dist[(Sqrt[1 + a*x^p]/a)*Sqrt[1/(1 + a*x^p)], Int[Sqrt[1 + a*x^p]*(Sqrt[1 - a*x^p]/x^(p + 1)), x], x] /; FreeQ[{a, p}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{1}{ax} + \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{\sqrt{1-ax}\sqrt{1+ax}}{x^2} dx}{a} \\
 &= -\frac{1}{ax} - \frac{\sqrt{1-ax}}{ax\sqrt{\frac{1}{1+ax}}} - \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{a^2}{\sqrt{1-ax}\sqrt{1+ax}} dx}{a} \\
 &= -\frac{1}{ax} - \frac{\sqrt{1-ax}}{ax\sqrt{\frac{1}{1+ax}}} - \left(a\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{\sqrt{1-ax}\sqrt{1+ax}} dx \\
 &= -\frac{1}{ax} - \frac{\sqrt{1-ax}}{ax\sqrt{\frac{1}{1+ax}}} - \left(a\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{\sqrt{1-a^2x^2}} dx \\
 &= -\frac{1}{ax} - \frac{\sqrt{1-ax}}{ax\sqrt{\frac{1}{1+ax}}} - \sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \arcsin(ax)
 \end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.06 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.56

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x} dx = -\frac{1}{ax} + \left(-1 - \frac{1}{ax}\right) \sqrt{\frac{1-ax}{1+ax}} - i \log \left(-2iax + 2\sqrt{\frac{1-ax}{1+ax}}(1+ax)\right)$$

[In] Integrate[E^ArcSech[a*x]/x,x]

[Out] -(1/(a*x)) + (-1 - 1/(a*x))*Sqrt[(1 - a*x)/(1 + a*x)] - I*Log[(-2*I)*a*x + 2*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)]

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.05 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.92

method	result	size
default	$-\frac{\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} \left(\arctan\left(\frac{\operatorname{csgn}(a)ax}{\sqrt{-a^2x^2+1}}\right) ax + \operatorname{csgn}(a)\sqrt{-a^2x^2+1} \right) \operatorname{csgn}(a)}{\sqrt{-a^2x^2+1}} - \frac{1}{ax}$	92

[In] int((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x,x,method=_RETURNVERBOSE)

[Out] -((a*x+1)/a/x)^(1/2)*(-(a*x-1)/a/x)^(1/2)*(arctan(csgn(a)*a*x/(-a^2*x^2+1)^(1/2))*a*x+csgn(a)*(-a^2*x^2+1)^(1/2))*csgn(a)/(-a^2*x^2+1)^(1/2)-1/a/x

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.60

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x} dx = -\frac{ax \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} - ax \arctan\left(\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}}\right) + 1}{ax}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x,x, algorithm="fricas")

[Out] -(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - a*x*arctan(sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x))) + 1)/(a*x)

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x} dx = \int \frac{1}{x^2} dx + \int \frac{a\sqrt{-1+\frac{1}{ax}}\sqrt{1+\frac{1}{ax}}}{x} dx$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))/x,x)

[Out] (Integral(x**(-2), x) + Integral(a*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x))/x, x))/a

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x} dx = \int \frac{\sqrt{\frac{1}{ax} + 1}\sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}}}{x} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x,x, algorithm="maxima")

[Out] integrate(sqrt(a*x + 1)*sqrt(-a*x + 1)/x^2, x)/a - 1/(a*x)

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x} dx = \int \frac{\sqrt{\frac{1}{ax} + 1}\sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}}}{x} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))/x, x)

Mupad [B] (verification not implemented)

Time = 6.37 (sec) , antiderivative size = 184, normalized size of antiderivative = 3.83

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x} dx = -\ln \left(\frac{\left(\sqrt{\frac{1}{ax} - 1 - i} \right)^2}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^2} + 1 \right) \operatorname{li} + \ln \left(\frac{\sqrt{\frac{1}{ax} - 1 - i}}{\sqrt{\frac{1}{ax} + 1 - 1}} \right) \operatorname{li} - \frac{1}{ax} + \frac{\left(\sqrt{\frac{1}{ax} - 1 - i} \right)^2 8i}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^2 \left(1 + \frac{\left(\sqrt{\frac{1}{ax} - 1 - i} \right)^4}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^4} - \frac{2 \left(\sqrt{\frac{1}{ax} - 1 - i} \right)^2}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^2} \right)}$$

[In] `int(((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))/x,x)`

[Out] `log(((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1))*1i - log(((1/(a*x) - 1)^(1/2) - 1i)^2/((1/(a*x) + 1)^(1/2) - 1)^2 + 1)*1i - 1/(a*x) + (((1/(a*x) - 1)^(1/2) - 1i)^2*8i)/(((1/(a*x) + 1)^(1/2) - 1)^2*((1/(a*x) - 1)^(1/2) - 1i)^4/((1/(a*x) + 1)^(1/2) - 1)^4 - (2*((1/(a*x) - 1)^(1/2) - 1i)^2)/((1/(a*x) + 1)^(1/2) - 1)^2 + 1))`

3.38 $\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^2} dx$

Optimal result	287
Rubi [B] (verified)	287
Mathematica [B] (verified)	289
Maple [A] (verified)	289
Fricas [B] (verification not implemented)	290
Sympy [A] (verification not implemented)	290
Maxima [F]	290
Giac [F]	291
Mupad [B] (verification not implemented)	291

Optimal result

Integrand size = 10, antiderivative size = 35

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^2} dx = -\frac{e^{\operatorname{sech}^{-1}(ax)}}{2x} + a \operatorname{arctanh}\left(\sqrt{\frac{1-ax}{1+ax}}\right)$$

[Out] $-1/2*(1/a/x+(1/a/x-1)^{(1/2)}*(1+1/a/x)^{(1/2)})/x+a*\operatorname{arctanh}(((-a*x+1)/(a*x+1))^{(1/2)})$

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 99 vs. $2(35) = 70$.

Time = 0.03 (sec), antiderivative size = 99, normalized size of antiderivative = 2.83, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6470, 30, 105, 12, 94, 214}

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^2} dx = \frac{1}{2}a\sqrt{\frac{1}{ax+1}}\sqrt{ax+1}\operatorname{arctanh}\left(\sqrt{1-ax}\sqrt{ax+1}\right) + \frac{\sqrt{1-ax}}{2ax^2\sqrt{\frac{1}{ax+1}}} + \frac{1}{2ax^2} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{x}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcSech}[a*x]}/x^2, x]$

[Out] $1/(2*a*x^2) - E^{\operatorname{ArcSech}[a*x]}/x + \operatorname{Sqrt}[1 - a*x]/(2*a*x^2*\operatorname{Sqrt}[(1 + a*x)^{-1}]) + (a*\operatorname{Sqrt}[(1 + a*x)^{-1}]*\operatorname{Sqrt}[1 + a*x]*\operatorname{ArcTanh}[\operatorname{Sqrt}[1 - a*x]*\operatorname{Sqrt}[1 + a*x]])/2$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 94

Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))), x_Symbol] := Dist[b*f, Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]

Rule 105

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 6470

Int[E^ArcSech[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*(E^ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] + Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, m, p}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{e^{\operatorname{sech}^{-1}(ax)}}{x} - \frac{\int \frac{1}{x^3} dx}{a} - \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{x^3\sqrt{1-ax}\sqrt{1+ax}} dx}{a} \\ &= \frac{1}{2ax^2} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{x} + \frac{\sqrt{1-ax}}{2ax^2\sqrt{\frac{1}{1+ax}}} - \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{a^2}{x\sqrt{1-ax}\sqrt{1+ax}} dx}{2a} \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2ax^2} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{x} + \frac{\sqrt{1-ax}}{2ax^2\sqrt{\frac{1}{1+ax}}} - \frac{1}{2} \left(a\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int \frac{1}{x\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{2ax^2} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{x} + \frac{\sqrt{1-ax}}{2ax^2\sqrt{\frac{1}{1+ax}}} \\
&\quad + \frac{1}{2} \left(a^2\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \operatorname{Subst} \left(\int \frac{1}{a-ax^2} dx, x, \sqrt{1-ax}\sqrt{1+ax} \right) \\
&= \frac{1}{2ax^2} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{x} + \frac{\sqrt{1-ax}}{2ax^2\sqrt{\frac{1}{1+ax}}} + \frac{1}{2} a\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \operatorname{arctanh} \left(\sqrt{1-ax}\sqrt{1+ax} \right)
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 93 vs. $2(35) = 70$.

Time = 0.07 (sec) , antiderivative size = 93, normalized size of antiderivative = 2.66

$$\begin{aligned}
\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^2} dx &= \frac{1}{2} \left(-\frac{1}{ax^2} - \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)}{ax^2} - a \log(x) \right. \\
&\quad \left. + a \log \left(1 + \sqrt{\frac{1-ax}{1+ax}} + ax\sqrt{\frac{1-ax}{1+ax}} \right) \right)
\end{aligned}$$

[In] Integrate[E^ArcSech[a*x]/x^2,x]

[Out] $(-(1/(a*x^2)) - (\operatorname{Sqrt}[(1 - a*x)/(1 + a*x)]*(1 + a*x))/(a*x^2) - a*\operatorname{Log}[x] + a*\operatorname{Log}[1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)] + a*x*\operatorname{Sqrt}[(1 - a*x)/(1 + a*x)]])/2$

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 91, normalized size of antiderivative = 2.60

method	result	size
default	$\frac{\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}\left(a^2x^2\operatorname{arctanh}\left(\frac{1}{\sqrt{-a^2x^2+1}}\right)-\sqrt{-a^2x^2+1}\right)}{2x\sqrt{-a^2x^2+1}} - \frac{1}{2ax^2}$	91

[In] int((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))/x^2,x,method=_RETURNVERBOSE)

[Out] $1/2*((a*x+1)/a/x)^(1/2)/x*(-(a*x-1)/a/x)^(1/2)*(a^2*x^2*\operatorname{arctanh}(1/(-a^2*x^2+1)^(1/2))-(-a^2*x^2+1)^(1/2))/(-a^2*x^2+1)^(1/2)-1/2/a/x^2$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 128 vs. $2(56) = 112$.

Time = 0.25 (sec) , antiderivative size = 128, normalized size of antiderivative = 3.66

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^2} dx = \frac{a^2 x^2 \log\left(ax \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} + 1\right) - a^2 x^2 \log\left(ax \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} - 1\right) - 2ax \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} - 2}{4ax^2}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^2,x, algorithm="fricas")

[Out] 1/4*(a^2*x^2*log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + 1) - a^2*x^2*log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 1) - 2*a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 2)/(a*x^2)

Sympy [A] (verification not implemented)

Time = 2.66 (sec) , antiderivative size = 71, normalized size of antiderivative = 2.03

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^2} dx = -a \left(2\sqrt{-1 + \frac{1}{ax}} \left(\frac{\left(1 + \frac{1}{ax}\right)^{\frac{3}{2}}}{4} - \frac{\sqrt{1 + \frac{1}{ax}}}{4} \right) - \log \left(2\sqrt{-1 + \frac{1}{ax}} + 2\sqrt{1 + \frac{1}{ax}} \right) \right) - \frac{1}{2ax^2}$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))/x**2,x)

[Out] -a*(2*sqrt(-1 + 1/(a*x))*((1 + 1/(a*x))**(3/2)/4 - sqrt(1 + 1/(a*x))/4) - 1*log(2*sqrt(-1 + 1/(a*x)) + 2*sqrt(1 + 1/(a*x)))) - 1/(2*a*x**2)

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^2} dx = \int \frac{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}}{x^2} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^2,x, algorithm="maxima")

[Out] integrate(sqrt(a*x + 1)*sqrt(-a*x + 1)/x^3, x)/a - 1/2/(a*x^2)

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^2} dx = \int \frac{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}}{x^2} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^2,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))/x^2, x)

Mupad [B] (verification not implemented)

Time = 4.88 (sec) , antiderivative size = 71, normalized size of antiderivative = 2.03

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^2} dx = \frac{a \ln \left(\sqrt{\frac{1}{ax} - 1} \sqrt{\frac{1}{ax} + 1} + \frac{1}{ax} \right)}{2} - \frac{1}{2ax^2} - \frac{\sqrt{\frac{1}{ax} - 1} \sqrt{\frac{1}{ax} + 1}}{2x}$$

[In] int(((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))/x^2,x)

[Out] (a*log((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x)))/2 - 1/(2*a*x^2) - ((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2))/(2*x)

3.39 $\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^3} dx$

Optimal result	292
Rubi [C] (verified)	292
Mathematica [A] (verified)	294
Maple [A] (verified)	294
Fricas [A] (verification not implemented)	294
Sympy [F]	295
Maxima [A] (verification not implemented)	295
Giac [F]	295
Mupad [B] (verification not implemented)	295

Optimal result

Integrand size = 10, antiderivative size = 55

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^3} dx = -\frac{1}{3ax^3} - \frac{8a^2 \left(\frac{1-ax}{1+ax}\right)^{3/2}}{3 \left(1 - \frac{1-ax}{1+ax}\right)^3}$$

[Out] $-1/3/a/x^3-8/3*a^2*((-a*x+1)/(a*x+1))^{(3/2)}/(1+(a*x-1)/(a*x+1))^3$

Rubi [C] (verified)

Result contains higher order function than in optimal. Order 3 vs. order 2 in optimal.

Time = 0.03 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.53, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6470, 30, 105, 12, 97}

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^3} dx = \frac{\sqrt{1-ax}}{6ax^3 \sqrt{\frac{1}{ax+1}}} + \frac{1}{6ax^3} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{2x^2} + \frac{a\sqrt{1-ax}}{3x \sqrt{\frac{1}{ax+1}}}$$

[In] `Int[E^ArcSech[a*x]/x^3,x]`

[Out] $1/(6*a*x^3) - E^{\operatorname{ArcSech}[a*x]}/(2*x^2) + \operatorname{Sqrt}[1 - a*x]/(6*a*x^3*\operatorname{Sqrt}[(1 + a*x)^{-1}]) + (a*\operatorname{Sqrt}[1 - a*x])/(3*x*\operatorname{Sqrt}[(1 + a*x)^{-1}])$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 30

`Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]`

Rule 97

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && EqQ[a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1), 0] && NeQ[m, -1]`

Rule 105

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])`

Rule 6470

`Int[E^ArcSech[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*(E^ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] + Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, m, p}, x] && NeQ[m, -1]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{e^{\operatorname{sech}^{-1}(ax)}}{2x^2} - \frac{\int \frac{1}{x^4} dx}{2a} - \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{x^4\sqrt{1-ax}\sqrt{1+ax}} dx}{2a} \\
 &= \frac{1}{6ax^3} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{2x^2} + \frac{\sqrt{1-ax}}{6ax^3\sqrt{\frac{1}{1+ax}}} + \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int -\frac{2a^2}{x^2\sqrt{1-ax}\sqrt{1+ax}} dx}{6a} \\
 &= \frac{1}{6ax^3} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{2x^2} + \frac{\sqrt{1-ax}}{6ax^3\sqrt{\frac{1}{1+ax}}} - \frac{1}{3} \left(a\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{x^2\sqrt{1-ax}\sqrt{1+ax}} dx \\
 &= \frac{1}{6ax^3} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{2x^2} + \frac{\sqrt{1-ax}}{6ax^3\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{3x\sqrt{\frac{1}{1+ax}}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.78

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^3} dx = \frac{-1 + (-1 + ax) \sqrt{\frac{1-ax}{1+ax}} (1 + ax)^2}{3ax^3}$$

[In] Integrate[E^ArcSech[a*x]/x^3,x]

[Out] (-1 + (-1 + a*x)*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2)/(3*a*x^3)

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.96

method	result	size
default	$\frac{\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} (a^2 x^2 - 1)}{3x^2} - \frac{1}{3ax^3}$	53

[In] int((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^3,x,method=_RETURNVERBOSE)

[Out] 1/3*((a*x+1)/a/x)^(1/2)/x^2*(-(a*x-1)/a/x)^(1/2)*(a^2*x^2-1)-1/3/a/x^3

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.95

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^3} dx = \frac{(a^3 x^3 - ax) \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} - 1}{3ax^3}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^3,x, algorithm="fricas")

[Out] 1/3*((a^3*x^3 - a*x)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 1)/(a*x^3)

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^3} dx = \frac{\int \frac{1}{x^4} dx + \int \frac{a\sqrt{-1+\frac{1}{ax}}\sqrt{1+\frac{1}{ax}}}{x^3} dx}{a}$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))/x**3,x)

[Out] (Integral(x**(-4), x) + Integral(a*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x))/x**3, x))/a

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.78

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^3} dx = \frac{(a^2x^3 - x)\sqrt{ax + 1}\sqrt{-ax + 1}}{3ax^4} - \frac{1}{3ax^3}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^3,x, algorithm="maxima")

[Out] 1/3*(a^2*x^3 - x)*sqrt(a*x + 1)*sqrt(-a*x + 1)/(a*x^4) - 1/3/(a*x^3)

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^3} dx = \int \frac{\sqrt{\frac{1}{ax} + 1}\sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}}{x^3} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^3,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))/x^3, x)

Mupad [B] (verification not implemented)

Time = 4.59 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.05

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^3} dx = -\frac{1}{3ax^3} - \frac{\left(\frac{\sqrt{\frac{1}{ax}+1}}{3} - \frac{a^2x^2\sqrt{\frac{1}{ax}+1}}{3}\right)\sqrt{\frac{1}{ax}-1}}{x^2}$$

[In] int(((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))/x^3,x)

[Out] - 1/(3*a*x^3) - (((1/(a*x) + 1)^(1/2)/3 - (a^2*x^2*(1/(a*x) + 1)^(1/2))/3)*(1/(a*x) - 1)^(1/2))/x^2

3.40 $\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^4} dx$

Optimal result	296
Rubi [A] (verified)	296
Mathematica [A] (verified)	298
Maple [A] (verified)	299
Fricas [A] (verification not implemented)	299
Sympy [F]	299
Maxima [F]	300
Giac [F]	300
Mupad [B] (verification not implemented)	300

Optimal result

Integrand size = 10, antiderivative size = 132

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^4} dx = \frac{1}{12ax^4} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{3x^3} + \frac{\sqrt{1-ax}}{12ax^4\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{8x^2\sqrt{\frac{1}{1+ax}}} + \frac{1}{8}a^3\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\operatorname{arctanh}\left(\sqrt{1-ax}\sqrt{1+ax}\right)$$

[Out] 1/12/a/x^4-1/3*(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^3+1/12*(-a*x+1)^(1/2)/a/x^4/(1/(a*x+1))^(1/2)+1/8*a*(-a*x+1)^(1/2)/x^2/(1/(a*x+1))^(1/2)+1/8*a^3*arctanh((-a*x+1)^(1/2)*(a*x+1)^(1/2))*(1/(a*x+1))^(1/2)*(a*x+1)^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6470, 30, 105, 12, 94, 214}

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^4} dx = \frac{1}{8}a^3\sqrt{\frac{1}{ax+1}}\sqrt{ax+1}\operatorname{arctanh}\left(\sqrt{1-ax}\sqrt{ax+1}\right) + \frac{\sqrt{1-ax}}{12ax^4\sqrt{\frac{1}{ax+1}}} + \frac{1}{12ax^4} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{3x^3} + \frac{a\sqrt{1-ax}}{8x^2\sqrt{\frac{1}{ax+1}}}$$

[In] Int[E^ArcSech[a*x]/x^4,x]

[Out] 1/(12*a*x^4) - E^ArcSech[a*x]/(3*x^3) + Sqrt[1 - a*x]/(12*a*x^4*Sqrt[(1 + a*x)^(-1)]) + (a*Sqrt[1 - a*x])/(8*x^2*Sqrt[(1 + a*x)^(-1)]) + (a^3*Sqrt[(1 + a*x)^(-1)]*Sqrt[1 + a*x]*ArcTanh[Sqrt[1 - a*x]*Sqrt[1 + a*x]])/8

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 30

```
Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && N
eQ[m, -1]
```

Rule 94

```
Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]*((e_) + (f_)*(x_
))), x_Symbol] := Dist[b*f, Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x],
x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[
2*b*d*e - f*(b*c + a*d), 0]
```

Rule 105

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_
))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x
)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a
*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*
(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x,
x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (Integer
Q[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])
```

Rule 214

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 6470

```
Int[E^ArcSech[(a_)*(x_)^(p_)]*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*(E^
ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] +
Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sq
rt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x)) /; FreeQ[{a, m, p}, x] && NeQ[m, -1
]
```

Rubi steps

$$\text{integral} = -\frac{e^{\text{sech}^{-1}(ax)}}{3x^3} - \frac{\int \frac{1}{x^5} dx}{3a} - \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{x^5\sqrt{1-ax}\sqrt{1+ax}} dx}{3a}$$

$$\begin{aligned}
&= \frac{1}{12ax^4} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{3x^3} + \frac{\sqrt{1-ax}}{12ax^4\sqrt{\frac{1}{1+ax}}} + \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int -\frac{3a^2}{x^3\sqrt{1-ax}\sqrt{1+ax}} dx}{12a} \\
&= \frac{1}{12ax^4} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{3x^3} + \frac{\sqrt{1-ax}}{12ax^4\sqrt{\frac{1}{1+ax}}} - \frac{1}{4} \left(a\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int \frac{1}{x^3\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{12ax^4} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{3x^3} + \frac{\sqrt{1-ax}}{12ax^4\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{8x^2\sqrt{\frac{1}{1+ax}}} \\
&\quad - \frac{1}{8} \left(a\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int \frac{a^2}{x\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{12ax^4} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{3x^3} + \frac{\sqrt{1-ax}}{12ax^4\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{8x^2\sqrt{\frac{1}{1+ax}}} \\
&\quad - \frac{1}{8} \left(a^3\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int \frac{1}{x\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{12ax^4} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{3x^3} + \frac{\sqrt{1-ax}}{12ax^4\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{8x^2\sqrt{\frac{1}{1+ax}}} \\
&\quad + \frac{1}{8} \left(a^4\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \operatorname{Subst} \left(\int \frac{1}{a-ax^2} dx, x, \sqrt{1-ax}\sqrt{1+ax} \right) \\
&= \frac{1}{12ax^4} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{3x^3} + \frac{\sqrt{1-ax}}{12ax^4\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{8x^2\sqrt{\frac{1}{1+ax}}} \\
&\quad + \frac{1}{8} a^3 \sqrt{\frac{1}{1+ax}} \sqrt{1+ax} \operatorname{arctanh} \left(\sqrt{1-ax}\sqrt{1+ax} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.83

$$\begin{aligned}
&\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^4} dx \\
&= \frac{-2 + \sqrt{\frac{1-ax}{1+ax}}(-2 - 2ax + a^2x^2 + a^3x^3) - a^4x^4 \log(x) + a^4x^4 \log\left(1 + \sqrt{\frac{1-ax}{1+ax}} + ax\sqrt{\frac{1-ax}{1+ax}}\right)}{8ax^4}
\end{aligned}$$

[In] Integrate[E^ArcSech[a*x]/x^4,x]

[Out] (-2 + Sqrt[(1 - a*x)/(1 + a*x)]*(-2 - 2*a*x + a^2*x^2 + a^3*x^3) - a^4*x^4*Log[x] + a^4*x^4*Log[1 + Sqrt[(1 - a*x)/(1 + a*x)] + a*x*Sqrt[(1 - a*x)/(1 + a*x)]])/(8*a*x^4)

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{-a^2x^2+1}}\right) a^4x^4 + a^2x^2\sqrt{-a^2x^2+1} - 2\sqrt{-a^2x^2+1} \right)}{8x^3\sqrt{-a^2x^2+1}} - \frac{1}{4ax^4}$	110

[In] int((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))/x^4,x,method=_RETURNVERBOSE)

[Out] $1/8*((a*x+1)/a/x)^{(1/2)}/x^3*(-(a*x-1)/a/x)^{(1/2)}*(\operatorname{arctanh}(1/(-a^2*x^2+1)^{(1/2)}))*a^4*x^4+a^2*x^2*(-a^2*x^2+1)^{(1/2)}-2*(-a^2*x^2+1)^{(1/2)})/(-a^2*x^2+1)^{(1/2)}-1/4/a/x^4$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.05

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^4} dx$$

$$= \frac{a^4x^4 \log\left(ax\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}+1\right) - a^4x^4 \log\left(ax\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}-1\right) + 2(a^3x^3 - 2ax)\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}}{16ax^4}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))/x^4,x, algorithm="fricas")

[Out] $1/16*(a^4*x^4*\log(a*x*\sqrt{(a*x+1)/(a*x)})*\sqrt{-(a*x-1)/(a*x)}+1) - a^4*x^4*\log(a*x*\sqrt{(a*x+1)/(a*x)})*\sqrt{-(a*x-1)/(a*x)}-1) + 2*(a^3*x^3 - 2*a*x)*\sqrt{(a*x+1)/(a*x)}*\sqrt{-(a*x-1)/(a*x)} - 4)/(a*x^4)$

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^4} dx = \int \frac{1}{x^5} dx + \int \frac{a\sqrt{-1+\frac{1}{ax}}\sqrt{1+\frac{1}{ax}}}{x^4} dx$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2))*(1+1/a/x)**(1/2))/x**4,x)

[Out] (Integral(x**(-5), x) + Integral(a*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x))/x**4, x))/a

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^4} dx = \int \frac{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}}{x^4} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^4,x, algorithm="maxima")

[Out] integrate(sqrt(a*x + 1)*sqrt(-a*x + 1)/x^5, x)/a - 1/4/(a*x^4)

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^4} dx = \int \frac{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}}{x^4} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^4,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))/x^4, x)

Mupad [B] (verification not implemented)

Time = 19.50 (sec) , antiderivative size = 602, normalized size of antiderivative = 4.56

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^4} dx = \frac{a^3 \operatorname{atanh}\left(\frac{\sqrt{\frac{1}{ax}-1-i}}{\sqrt{\frac{1}{ax}+1-1}}\right)}{2} - \frac{35 a^3 \left(\sqrt{\frac{1}{ax}-1-i}\right)^3}{2 \left(\sqrt{\frac{1}{ax}+1-1}\right)^3} + \frac{273 a^3 \left(\sqrt{\frac{1}{ax}-1-i}\right)^5}{2 \left(\sqrt{\frac{1}{ax}+1-1}\right)^5} + \frac{715 a^3 \left(\sqrt{\frac{1}{ax}-1-i}\right)^7}{2 \left(\sqrt{\frac{1}{ax}+1-1}\right)^7} + \frac{715 a^3 \left(\sqrt{\frac{1}{ax}-1-i}\right)^9}{2 \left(\sqrt{\frac{1}{ax}+1-1}\right)^9} + \frac{273 a^3 \left(\sqrt{\frac{1}{ax}-1-i}\right)^{11}}{2 \left(\sqrt{\frac{1}{ax}+1-1}\right)^{11}} + \frac{35 a^3 \left(\sqrt{\frac{1}{ax}-1-i}\right)^{13}}{2 \left(\sqrt{\frac{1}{ax}+1-1}\right)^{13}} - \frac{1}{4 a x^4}$$

[In] int(((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))/x^4,x)

[Out] (a^3*atanh(((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1)))/2 - ((35*a^3*((1/(a*x) - 1)^(1/2) - 1i)^3)/(2*((1/(a*x) + 1)^(1/2) - 1)^3) + (273*a^3*((1/(a*x) - 1)^(1/2) - 1i)^5)/(2*((1/(a*x) + 1)^(1/2) - 1)^5) + (715*a^3*((1/(a*x) - 1)^(1/2) - 1i)^7)/(2*((1/(a*x) + 1)^(1/2) - 1)^7) + (715*a^3*((1/(a*x) - 1)^(1/2) - 1i)^9)/(2*((1/(a*x) + 1)^(1/2) - 1)^9) + (273*a^3*((1/

$$\begin{aligned}
& (a*x - 1)^{(1/2) - 1i} / (2*((1/(a*x) + 1)^{(1/2) - 1})^{11}) + (35*a^3*((1/(a*x) - 1)^{(1/2) - 1i})^{13}) / (2*((1/(a*x) + 1)^{(1/2) - 1})^{13}) + (a^3*((1/(a*x) - 1)^{(1/2) - 1i})^{15}) / (2*((1/(a*x) + 1)^{(1/2) - 1})^{15}) + (a^3*((1/(a*x) - 1)^{(1/2) - 1i})) / (2*((1/(a*x) + 1)^{(1/2) - 1})) / ((28*((1/(a*x) - 1)^{(1/2) - 1i})^4) / ((1/(a*x) + 1)^{(1/2) - 1})^4 - (8*((1/(a*x) - 1)^{(1/2) - 1i})^2) / ((1/(a*x) + 1)^{(1/2) - 1})^2 - (56*((1/(a*x) - 1)^{(1/2) - 1i})^6) / ((1/(a*x) + 1)^{(1/2) - 1})^6 + (70*((1/(a*x) - 1)^{(1/2) - 1i})^8) / ((1/(a*x) + 1)^{(1/2) - 1})^8 - (56*((1/(a*x) - 1)^{(1/2) - 1i})^{10}) / ((1/(a*x) + 1)^{(1/2) - 1})^{10} + (28*((1/(a*x) - 1)^{(1/2) - 1i})^{12}) / ((1/(a*x) + 1)^{(1/2) - 1})^{12} - (8*((1/(a*x) - 1)^{(1/2) - 1i})^{14}) / ((1/(a*x) + 1)^{(1/2) - 1})^{14} + ((1/(a*x) - 1)^{(1/2) - 1i})^{16} / ((1/(a*x) + 1)^{(1/2) - 1})^{16} + 1) - 1/(4*a*x^4)
\end{aligned}$$

3.41 $\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^5} dx$

Optimal result	302
Rubi [A] (verified)	302
Mathematica [A] (verified)	304
Maple [A] (verified)	304
Fricas [A] (verification not implemented)	305
Sympy [F]	305
Maxima [A] (verification not implemented)	305
Giac [F]	306
Mupad [B] (verification not implemented)	306

Optimal result

Integrand size = 10, antiderivative size = 115

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^5} dx = \frac{1}{20ax^5} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{4x^4} + \frac{\sqrt{1-ax}}{20ax^5\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{15x^3\sqrt{\frac{1}{1+ax}}} + \frac{2a^3\sqrt{1-ax}}{15x\sqrt{\frac{1}{1+ax}}}$$

[Out] $1/20/a/x^5 - 1/4*(1/a/x + (1/a/x - 1)^{(1/2)}*(1 + 1/a/x)^{(1/2)})/x^4 + 1/20*(-a*x + 1)^{(1/2)}/a/x^5/(1/(a*x + 1))^{(1/2)} + 1/15*a*(-a*x + 1)^{(1/2)}/x^3/(1/(a*x + 1))^{(1/2)} + 2/15*a^3*(-a*x + 1)^{(1/2)}/x/(1/(a*x + 1))^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6470, 30, 105, 12, 97}

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^5} dx = \frac{2a^3\sqrt{1-ax}}{15x\sqrt{\frac{1}{ax+1}}} + \frac{\sqrt{1-ax}}{20ax^5\sqrt{\frac{1}{ax+1}}} + \frac{1}{20ax^5} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{4x^4} + \frac{a\sqrt{1-ax}}{15x^3\sqrt{\frac{1}{ax+1}}}$$

[In] Int[E^ArcSech[a*x]/x^5, x]

[Out] $1/(20*a*x^5) - E^{\operatorname{ArcSech}[a*x]}/(4*x^4) + \operatorname{Sqrt}[1 - a*x]/(20*a*x^5*\operatorname{Sqrt}[(1 + a*x)^{-1}]) + (a*\operatorname{Sqrt}[1 - a*x])/(15*x^3*\operatorname{Sqrt}[(1 + a*x)^{-1}]) + (2*a^3*\operatorname{Sqrt}[1 - a*x])/(15*x*\operatorname{Sqrt}[(1 + a*x)^{-1}])$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 30

`Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]`

Rule 97

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && EqQ[a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1), 0] && NeQ[m, -1]`

Rule 105

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])`

Rule 6470

`Int[E^ArcSech[(a_.)*(x_)^(p_.)]*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*(E^ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] + Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, m, p}, x] && NeQ[m, -1]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{e^{\operatorname{sech}^{-1}(ax)}}{4x^4} - \frac{\int \frac{1}{x^6} dx}{4a} - \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{x^6\sqrt{1-ax}\sqrt{1+ax}} dx}{4a} \\
 &= \frac{1}{20ax^5} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{4x^4} + \frac{\sqrt{1-ax}}{20ax^5\sqrt{\frac{1}{1+ax}}} + \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int -\frac{4a^2}{x^4\sqrt{1-ax}\sqrt{1+ax}} dx}{20a} \\
 &= \frac{1}{20ax^5} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{4x^4} + \frac{\sqrt{1-ax}}{20ax^5\sqrt{\frac{1}{1+ax}}} - \frac{1}{5} \left(a\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{x^4\sqrt{1-ax}\sqrt{1+ax}} dx
 \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{20ax^5} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{4x^4} + \frac{\sqrt{1-ax}}{20ax^5\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{15x^3\sqrt{\frac{1}{1+ax}}} \\
&\quad + \frac{1}{15} \left(a\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int -\frac{2a^2}{x^2\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{20ax^5} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{4x^4} + \frac{\sqrt{1-ax}}{20ax^5\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{15x^3\sqrt{\frac{1}{1+ax}}} \\
&\quad - \frac{1}{15} \left(2a^3\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int \frac{1}{x^2\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{20ax^5} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{4x^4} + \frac{\sqrt{1-ax}}{20ax^5\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{15x^3\sqrt{\frac{1}{1+ax}}} + \frac{2a^3\sqrt{1-ax}}{15x\sqrt{\frac{1}{1+ax}}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.52

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^5} dx = \frac{-3 + \sqrt{\frac{1-ax}{1+ax}}(1+ax)^2(-3 + 3ax - 2a^2x^2 + 2a^3x^3)}{15ax^5}$$

[In] Integrate[E^ArcSech[a*x]/x^5,x]

[Out] (-3 + Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2*(-3 + 3*a*x - 2*a^2*x^2 + 2*a^3*x^3))/(15*a*x^5)

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.55

method	result	size
default	$\frac{\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}(a^2x^2-1)(2a^2x^2+3)}{15x^4} - \frac{1}{5ax^5}$	63

[In] int((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^5,x,method=_RETURNVERBOSE)

[Out] 1/15*((a*x+1)/a/x)^(1/2)/x^4*(-(a*x-1)/a/x)^(1/2)*(a^2*x^2-1)*(2*a^2*x^2+3)-1/5/a/x^5

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.52

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^5} dx = \frac{(2a^5x^5 + a^3x^3 - 3ax)\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} - 3}{15ax^5}$$

```
[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^5,x, algorithm="fricas")
```

```
[Out] 1/15*((2*a^5*x^5 + a^3*x^3 - 3*a*x)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 3)/(a*x^5)
```

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^5} dx = \frac{\int \frac{1}{x^6} dx + \int \frac{a\sqrt{-1+\frac{1}{ax}}\sqrt{1+\frac{1}{ax}}}{x^5} dx}{a}$$

```
[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))/x**5,x)
```

```
[Out] (Integral(x**(-6), x) + Integral(a*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x))/x**5, x))/a
```

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.44

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^5} dx = \frac{(2a^4x^5 + a^2x^3 - 3x)\sqrt{ax+1}\sqrt{-ax+1}}{15ax^6} - \frac{1}{5ax^5}$$

```
[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^5,x, algorithm="maxima")
```

```
[Out] 1/15*(2*a^4*x^5 + a^2*x^3 - 3*x)*sqrt(a*x + 1)*sqrt(-a*x + 1)/(a*x^6) - 1/5/(a*x^5)
```

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^5} dx = \int \frac{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}}{x^5} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^5,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))/x^5, x)

Mupad [B] (verification not implemented)

Time = 4.64 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.66

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^5} dx = \frac{\sqrt{\frac{1}{ax} - 1} \left(\frac{a^2 x^2 \sqrt{\frac{1}{ax} + 1}}{15} - \frac{\sqrt{\frac{1}{ax} + 1}}{5} + \frac{2 a^4 x^4 \sqrt{\frac{1}{ax} + 1}}{15} \right)}{x^4} - \frac{1}{5 a x^5}$$

[In] int(((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))/x^5,x)

[Out] ((1/(a*x) - 1)^(1/2)*((a^2*x^2*(1/(a*x) + 1)^(1/2))/15 - (1/(a*x) + 1)^(1/2)/5 + (2*a^4*x^4*(1/(a*x) + 1)^(1/2))/15))/x^4 - 1/(5*a*x^5)

3.42 $\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^6} dx$

Optimal result	307
Rubi [A] (verified)	307
Mathematica [A] (verified)	310
Maple [A] (verified)	310
Fricas [A] (verification not implemented)	310
Sympy [F]	311
Maxima [F]	311
Giac [F]	311
Mupad [B] (verification not implemented)	312

Optimal result

Integrand size = 10, antiderivative size = 163

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^6} dx = \frac{1}{30ax^6} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{5x^5} + \frac{\sqrt{1-ax}}{30ax^6\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{24x^4\sqrt{\frac{1}{1+ax}}} + \frac{a^3\sqrt{1-ax}}{16x^2\sqrt{\frac{1}{1+ax}}} + \frac{1}{16}a^5\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\operatorname{arctanh}\left(\sqrt{1-ax}\sqrt{1+ax}\right)$$

[Out] 1/30/a/x^6-1/5*(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^5+1/30*(-a*x+1)^(1/2)/a/x^6/(1/(a*x+1))^(1/2)+1/24*a*(-a*x+1)^(1/2)/x^4/(1/(a*x+1))^(1/2)+1/16*a^3*(-a*x+1)^(1/2)/x^2/(1/(a*x+1))^(1/2)+1/16*a^5*arctanh((-a*x+1)^(1/2)*(a*x+1)^(1/2))*(1/(a*x+1))^(1/2)*(a*x+1)^(1/2)

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6470, 30, 105, 12, 94, 214}

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^6} dx = \frac{1}{16}a^5\sqrt{\frac{1}{ax+1}}\sqrt{ax+1}\operatorname{arctanh}\left(\sqrt{1-ax}\sqrt{ax+1}\right) + \frac{a^3\sqrt{1-ax}}{16x^2\sqrt{\frac{1}{ax+1}}} + \frac{\sqrt{1-ax}}{30ax^6\sqrt{\frac{1}{ax+1}}} + \frac{1}{30ax^6} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{5x^5} + \frac{a\sqrt{1-ax}}{24x^4\sqrt{\frac{1}{ax+1}}}$$

[In] Int[E^ArcSech[a*x]/x^6,x]

[Out] 1/(30*a*x^6) - E^ArcSech[a*x]/(5*x^5) + Sqrt[1 - a*x]/(30*a*x^6*Sqrt[(1 + a*x)^(-1)]) + (a*Sqrt[1 - a*x])/(24*x^4*Sqrt[(1 + a*x)^(-1)]) + (a^3*Sqrt[1

$- a*x]/(16*x^2*\text{Sqrt}[(1 + a*x)^{-1}]) + (a^5*\text{Sqrt}[(1 + a*x)^{-1}]*\text{Sqrt}[1 + a*x]*\text{ArcTanh}[\text{Sqrt}[1 - a*x]*\text{Sqrt}[1 + a*x]])/16$

Rule 12

$\text{Int}[(a_*)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{Match}[\text{Q}[u, (b_*)*(v_)] /; \text{FreeQ}[b, x]]$

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}/(m + 1), x] /; \text{FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 94

$\text{Int}[1/(\text{Sqrt}[(a_.) + (b_.)*(x_)]*\text{Sqrt}[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))), x_Symbol] \rightarrow \text{Dist}[b*f, \text{Subst}[\text{Int}[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, \text{Sqrt}[a + b*x]*\text{Sqrt}[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{EqQ}[2*b*d*e - f*(b*c + a*d), 0]$

Rule 105

$\text{Int}[(a_.) + (b_.)*(x_)]^{(m_.)}*((c_.) + (d_.)*(x_))^{(n_.)}*((e_.) + (f_.)*(x_))^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[b*(a + b*x)^{(m + 1)}*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)}/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + \text{Dist}[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), \text{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^n*(e + f*x)^p*\text{Simp}[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \text{ILtQ}[m, -1] \ \&\& \ (\text{IntegerQ}[n] \ || \ \text{IntegersQ}[2*n, 2*p] \ || \ \text{ILtQ}[m + n + p + 3, 0])$

Rule 214

$\text{Int}[(a_.) + (b_.)*(x_)^2]^{(-1)}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$

Rule 6470

$\text{Int}[E^{\text{ArcSech}[(a_.)*(x_)^{(p_.)}]}*(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}*(E^{\text{ArcSech}[a*x^p]/(m + 1)}), x] + (\text{Dist}[p/(a*(m + 1)), \text{Int}[x^{(m - p)}, x], x] + \text{Dist}[p*(\text{Sqrt}[1 + a*x^p]/(a*(m + 1)))*\text{Sqrt}[1/(1 + a*x^p)], \text{Int}[x^{(m - p)}/(\text{Sqrt}[1 + a*x^p]*\text{Sqrt}[1 - a*x^p]), x], x]) /; \text{FreeQ}[\{a, m, p\}, x] \ \&\& \ \text{NeQ}[m, -1]$

Rubi steps

$$\text{integral} = -\frac{e^{\text{sech}^{-1}(ax)}}{5x^5} - \frac{\int \frac{1}{x^7} dx}{5a} - \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{x^7\sqrt{1-ax}\sqrt{1+ax}} dx}{5a}$$

$$\begin{aligned}
&= \frac{1}{30ax^6} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{5x^5} + \frac{\sqrt{1-ax}}{30ax^6\sqrt{\frac{1}{1+ax}}} + \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int -\frac{5a^2}{x^5\sqrt{1-ax}\sqrt{1+ax}} dx}{30a} \\
&= \frac{1}{30ax^6} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{5x^5} + \frac{\sqrt{1-ax}}{30ax^6\sqrt{\frac{1}{1+ax}}} - \frac{1}{6} \left(a\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int \frac{1}{x^5\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{30ax^6} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{5x^5} + \frac{\sqrt{1-ax}}{30ax^6\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{24x^4\sqrt{\frac{1}{1+ax}}} \\
&\quad + \frac{1}{24} \left(a\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int -\frac{3a^2}{x^3\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{30ax^6} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{5x^5} + \frac{\sqrt{1-ax}}{30ax^6\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{24x^4\sqrt{\frac{1}{1+ax}}} \\
&\quad - \frac{1}{8} \left(a^3\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int \frac{1}{x^3\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{30ax^6} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{5x^5} + \frac{\sqrt{1-ax}}{30ax^6\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{24x^4\sqrt{\frac{1}{1+ax}}} + \frac{a^3\sqrt{1-ax}}{16x^2\sqrt{\frac{1}{1+ax}}} \\
&\quad - \frac{1}{16} \left(a^3\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int \frac{a^2}{x\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{30ax^6} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{5x^5} + \frac{\sqrt{1-ax}}{30ax^6\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{24x^4\sqrt{\frac{1}{1+ax}}} + \frac{a^3\sqrt{1-ax}}{16x^2\sqrt{\frac{1}{1+ax}}} \\
&\quad - \frac{1}{16} \left(a^5\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int \frac{1}{x\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{30ax^6} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{5x^5} + \frac{\sqrt{1-ax}}{30ax^6\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{24x^4\sqrt{\frac{1}{1+ax}}} + \frac{a^3\sqrt{1-ax}}{16x^2\sqrt{\frac{1}{1+ax}}} \\
&\quad + \frac{1}{16} \left(a^6\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \operatorname{Subst} \left(\int \frac{1}{a-ax^2} dx, x, \sqrt{1-ax}\sqrt{1+ax} \right) \\
&= \frac{1}{30ax^6} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{5x^5} + \frac{\sqrt{1-ax}}{30ax^6\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{24x^4\sqrt{\frac{1}{1+ax}}} + \frac{a^3\sqrt{1-ax}}{16x^2\sqrt{\frac{1}{1+ax}}} \\
&\quad + \frac{1}{16} a^5 \sqrt{\frac{1}{1+ax}} \sqrt{1+ax} \operatorname{arctanh} \left(\sqrt{1-ax}\sqrt{1+ax} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.79

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^6} dx$$

$$= \frac{-8 + \sqrt{\frac{1-ax}{1+ax}}(-8 - 8ax + 2a^2x^2 + 2a^3x^3 + 3a^4x^4 + 3a^5x^5) - 3a^6x^6 \log(x) + 3a^6x^6 \log\left(1 + \sqrt{\frac{1-ax}{1+ax}} + ax\sqrt{\frac{1-ax}{1+ax}}\right)}{48ax^6}$$

`[In] Integrate[E^ArcSech[a*x]/x^6,x]`

```
[Out] (-8 + Sqrt[(1 - a*x)/(1 + a*x)]*(-8 - 8*a*x + 2*a^2*x^2 + 2*a^3*x^3 + 3*a^4*x^4 + 3*a^5*x^5) - 3*a^6*x^6*Log[x] + 3*a^6*x^6*Log[1 + Sqrt[(1 - a*x)/(1 + a*x)]] + a*x*Sqrt[(1 - a*x)/(1 + a*x)])/(48*a*x^6)
```

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.81

method	result	size
default	$\frac{\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} \left(3 \operatorname{arctanh}\left(\frac{1}{\sqrt{-a^2x^2+1}}\right) a^6x^6 + 3\sqrt{-a^2x^2+1} a^4x^4 + 2a^2x^2\sqrt{-a^2x^2+1} - 8\sqrt{-a^2x^2+1} \right)}{48x^5\sqrt{-a^2x^2+1}} - \frac{1}{6ax^6}$	132

`[In] int((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^6,x,method=_RETURNVERBOSE)`

```
[Out] 1/48*((a*x+1)/a/x)^(1/2)/x^5*(-(a*x-1)/a/x)^(1/2)*(3*arctanh(1/(-a^2*x^2+1)^(1/2))*a^6*x^6+3*(-a^2*x^2+1)^(1/2)*a^4*x^4+2*a^2*x^2*(-a^2*x^2+1)^(1/2)-8*(-a^2*x^2+1)^(1/2))/(-a^2*x^2+1)^(1/2)-1/6/a/x^6
```

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.91

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^6} dx$$

$$= \frac{3a^6x^6 \log\left(ax\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} + 1\right) - 3a^6x^6 \log\left(ax\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} - 1\right) + 2(3a^5x^5 + 2a^3x^3 - 8ax)\sqrt{\frac{ax+1}{ax}}}{96ax^6}$$

`[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^6,x, algorithm="fricas")`

```
[Out] 1/96*(3*a^6*x^6*log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + 1) - 3*a^6*x^6*log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 1) + 2*(3
```

$*a^5*x^5 + 2*a^3*x^3 - 8*a*x)*\text{sqrt}((a*x + 1)/(a*x))*\text{sqrt}(-(a*x - 1)/(a*x)) - 16)/(a*x^6)$

Sympy [F]

$$\int \frac{e^{\text{sech}^{-1}(ax)}}{x^6} dx = \int \frac{1}{x^7} dx + \int \frac{a\sqrt{-1+\frac{1}{ax}}\sqrt{1+\frac{1}{ax}}}{x^6} dx$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))/x**6,x)

[Out] (Integral(x**(-7), x) + Integral(a*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x))/x**6, x))/a

Maxima [F]

$$\int \frac{e^{\text{sech}^{-1}(ax)}}{x^6} dx = \int \frac{\sqrt{\frac{1}{ax} + 1}\sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}}}{x^6} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^6,x, algorithm="maxima")

[Out] integrate(sqrt(a*x + 1)*sqrt(-a*x + 1)/x^7, x)/a - 1/6/(a*x^6)

Giac [F]

$$\int \frac{e^{\text{sech}^{-1}(ax)}}{x^6} dx = \int \frac{\sqrt{\frac{1}{ax} + 1}\sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}}}{x^6} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^6,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))/x^6, x)

Mupad [B] (verification not implemented)

Time = 41.12 (sec) , antiderivative size = 878, normalized size of antiderivative = 5.39

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^6} dx$$

$$= \frac{35 a^5 \left(\sqrt{\frac{1}{ax}-1-i}\right)^3}{12 \left(\sqrt{\frac{1}{ax}+1-1}\right)^3} + \frac{757 a^5 \left(\sqrt{\frac{1}{ax}-1-i}\right)^5}{4 \left(\sqrt{\frac{1}{ax}+1-1}\right)^5} + \frac{7339 a^5 \left(\sqrt{\frac{1}{ax}-1-i}\right)^7}{4 \left(\sqrt{\frac{1}{ax}+1-1}\right)^7} + \frac{41929 a^5 \left(\sqrt{\frac{1}{ax}-1-i}\right)^9}{6 \left(\sqrt{\frac{1}{ax}+1-1}\right)^9} + \frac{25661 a^5 \left(\sqrt{\frac{1}{ax}-1-i}\right)^{11}}{2 \left(\sqrt{\frac{1}{ax}+1-1}\right)^{11}} + \frac{25661 a^5 \left(\sqrt{\frac{1}{ax}-1-i}\right)^{13}}{2 \left(\sqrt{\frac{1}{ax}+1-1}\right)^{13}} + \frac{41929 a^5 \left(\sqrt{\frac{1}{ax}-1-i}\right)^{15}}{6 \left(\sqrt{\frac{1}{ax}+1-1}\right)^{15}} + \frac{7339 a^5 \left(\sqrt{\frac{1}{ax}-1-i}\right)^{17}}{4 \left(\sqrt{\frac{1}{ax}+1-1}\right)^{17}} + \frac{757 a^5 \left(\sqrt{\frac{1}{ax}-1-i}\right)^{19}}{4 \left(\sqrt{\frac{1}{ax}+1-1}\right)^{19}} + \frac{35 a^5 \left(\sqrt{\frac{1}{ax}-1-i}\right)^{21}}{12 \left(\sqrt{\frac{1}{ax}+1-1}\right)^{21}} - \frac{a^5 \left(\sqrt{\frac{1}{ax}-1-i}\right)^{23}}{4 \left(\sqrt{\frac{1}{ax}+1-1}\right)^{23}} - \frac{a^5 \left(\sqrt{\frac{1}{ax}-1-i}\right)}{4 \left(\sqrt{\frac{1}{ax}+1-1}\right)^4} - \frac{1}{6 a x^6}$$

[In] int(((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))/x^6,x)

[Out] ((35*a^5*((1/(a*x) - 1)^(1/2) - 1i)^3)/(12*((1/(a*x) + 1)^(1/2) - 1)^3) + (757*a^5*((1/(a*x) - 1)^(1/2) - 1i)^5)/(4*((1/(a*x) + 1)^(1/2) - 1)^5) + (7339*a^5*((1/(a*x) - 1)^(1/2) - 1i)^7)/(4*((1/(a*x) + 1)^(1/2) - 1)^7) + (41929*a^5*((1/(a*x) - 1)^(1/2) - 1i)^9)/(6*((1/(a*x) + 1)^(1/2) - 1)^9) + (25661*a^5*((1/(a*x) - 1)^(1/2) - 1i)^11)/(2*((1/(a*x) + 1)^(1/2) - 1)^11) + (25661*a^5*((1/(a*x) - 1)^(1/2) - 1i)^13)/(2*((1/(a*x) + 1)^(1/2) - 1)^13) + (41929*a^5*((1/(a*x) - 1)^(1/2) - 1i)^15)/(6*((1/(a*x) + 1)^(1/2) - 1)^15) + (7339*a^5*((1/(a*x) - 1)^(1/2) - 1i)^17)/(4*((1/(a*x) + 1)^(1/2) - 1)^17) + (757*a^5*((1/(a*x) - 1)^(1/2) - 1i)^19)/(4*((1/(a*x) + 1)^(1/2) - 1)^19) + (35*a^5*((1/(a*x) - 1)^(1/2) - 1i)^21)/(12*((1/(a*x) + 1)^(1/2) - 1)^21) - (a^5*((1/(a*x) - 1)^(1/2) - 1i)^23)/(4*((1/(a*x) + 1)^(1/2) - 1)^23) - (a^5*((1/(a*x) - 1)^(1/2) - 1i))/(4*((1/(a*x) + 1)^(1/2) - 1)))/((66*((1/(a*x) - 1)^(1/2) - 1i)^4)/((1/(a*x) + 1)^(1/2) - 1)^4 - (12*((1/(a*x) - 1)^(1/2) - 1i)^2)/((1/(a*x) + 1)^(1/2) - 1)^2 - (220*((1/(a*x) - 1)^(1/2) - 1i)^6)/((1/(a*x) + 1)^(1/2) - 1)^6 + (495*((1/(a*x) - 1)^(1/2) - 1i)^8)/((1/(a*x) + 1)^(1/2) - 1)^8 - (792*((1/(a*x) - 1)^(1/2) - 1i)^10)/((1/(a*x) + 1)^(1/2) - 1)^10 + (924*((1/(a*x) - 1)^(1/2) - 1i)^12)/((1/(a*x) + 1)^(1/2) - 1)^12 - (792*((1/(a*x) - 1)^(1/2) - 1i)^14)/((1/(a*x) + 1)^(1/2) - 1)^14 + (495*((1/(a*x) - 1)^(1/2) - 1i)^16)/((1/(a*x) + 1)^(1/2) - 1)^16 - (220*((1/(a*x) - 1)^(1/2) - 1i)^18)/((1/(a*x) + 1)^(1/2) - 1)^18 + (66*((1/(a*x) - 1)^(1/2) - 1i)^20)/((1/(a*x) + 1)^(1/2) - 1)^20 - (12*((1/(a*x) - 1)^(1/2) - 1i)^22)/((1/(a*x) + 1)^(1/2) - 1)^22 + ((1/(a*x) - 1)^(1/2) - 1i)^24)/((1/(a*x) + 1)^(1/2) - 1)^24 + 1) + (a^5*atanh(((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1))))/4 - 1/(6*a*x^6)

3.43 $\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^7} dx$

Optimal result	313
Rubi [A] (verified)	313
Mathematica [A] (verified)	316
Maple [A] (verified)	316
Fricas [A] (verification not implemented)	316
Sympy [F]	317
Maxima [A] (verification not implemented)	317
Giac [F]	317
Mupad [B] (verification not implemented)	318

Optimal result

Integrand size = 10, antiderivative size = 146

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^7} dx = \frac{1}{42ax^7} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{6x^6} + \frac{\sqrt{1-ax}}{42ax^7\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{35x^5\sqrt{\frac{1}{1+ax}}} + \frac{4a^3\sqrt{1-ax}}{105x^3\sqrt{\frac{1}{1+ax}}} + \frac{8a^5\sqrt{1-ax}}{105x\sqrt{\frac{1}{1+ax}}}$$

[Out] $1/42/a/x^7-1/6*(1/a/x+(1/a/x-1)^{(1/2)}*(1+1/a/x)^{(1/2)})/x^6+1/42*(-a*x+1)^{(1/2)}/a/x^7/(1/(a*x+1))^{(1/2)}+1/35*a*(-a*x+1)^{(1/2)}/x^5/(1/(a*x+1))^{(1/2)}+4/105*a^3*(-a*x+1)^{(1/2)}/x^3/(1/(a*x+1))^{(1/2)}+8/105*a^5*(-a*x+1)^{(1/2)}/x/(1/(a*x+1))^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6470, 30, 105, 12, 97}

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^7} dx = \frac{8a^5\sqrt{1-ax}}{105x\sqrt{\frac{1}{ax+1}}} + \frac{4a^3\sqrt{1-ax}}{105x^3\sqrt{\frac{1}{ax+1}}} + \frac{\sqrt{1-ax}}{42ax^7\sqrt{\frac{1}{ax+1}}} + \frac{1}{42ax^7} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{6x^6} + \frac{a\sqrt{1-ax}}{35x^5\sqrt{\frac{1}{ax+1}}}$$

[In] Int[E^ArcSech[a*x]/x^7,x]

```
[Out] 1/(42*a*x^7) - E^ArcSech[a*x]/(6*x^6) + Sqrt[1 - a*x]/(42*a*x^7*Sqrt[(1 + a
*x)^(-1)]) + (a*Sqrt[1 - a*x]/(35*x^5*Sqrt[(1 + a*x)^(-1)])) + (4*a^3*Sqrt[
1 - a*x]/(105*x^3*Sqrt[(1 + a*x)^(-1)])) + (8*a^5*Sqrt[1 - a*x]/(105*x*Sqr
t[(1 + a*x)^(-1)])
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 30

```
Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && N
eQ[m, -1]
```

Rule 97

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_., x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x
)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] /; FreeQ[{a, b, c, d, e, f
, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && EqQ[a*d*f*(m + 1) + b*
c*f*(n + 1) + b*d*e*(p + 1), 0] && NeQ[m, -1]
```

Rule 105

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_., x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x
)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a
*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*
(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x,
x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (Integer
Q[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])
```

Rule 6470

```
Int[E^ArcSech[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*(E^
ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] +
Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sq
rt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, m, p}, x] && NeQ[m, -1
]
```

Rubi steps

$$\text{integral} = -\frac{e^{\operatorname{sech}^{-1}(ax)}}{6x^6} - \frac{\int \frac{1}{x^8} dx}{6a} - \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{x^8\sqrt{1-ax}\sqrt{1+ax}} dx}{6a}$$

$$\begin{aligned}
&= \frac{1}{42ax^7} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{6x^6} + \frac{\sqrt{1-ax}}{42ax^7\sqrt{\frac{1}{1+ax}}} + \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int -\frac{6a^2}{x^6\sqrt{1-ax}\sqrt{1+ax}} dx}{42a} \\
&= \frac{1}{42ax^7} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{6x^6} + \frac{\sqrt{1-ax}}{42ax^7\sqrt{\frac{1}{1+ax}}} - \frac{1}{7} \left(a\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int \frac{1}{x^6\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{42ax^7} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{6x^6} + \frac{\sqrt{1-ax}}{42ax^7\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{35x^5\sqrt{\frac{1}{1+ax}}} \\
&\quad + \frac{1}{35} \left(a\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int -\frac{4a^2}{x^4\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{42ax^7} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{6x^6} + \frac{\sqrt{1-ax}}{42ax^7\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{35x^5\sqrt{\frac{1}{1+ax}}} \\
&\quad - \frac{1}{35} \left(4a^3\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int \frac{1}{x^4\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{42ax^7} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{6x^6} + \frac{\sqrt{1-ax}}{42ax^7\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{35x^5\sqrt{\frac{1}{1+ax}}} + \frac{4a^3\sqrt{1-ax}}{105x^3\sqrt{\frac{1}{1+ax}}} \\
&\quad + \frac{1}{105} \left(4a^3\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int -\frac{2a^2}{x^2\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{42ax^7} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{6x^6} + \frac{\sqrt{1-ax}}{42ax^7\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{35x^5\sqrt{\frac{1}{1+ax}}} + \frac{4a^3\sqrt{1-ax}}{105x^3\sqrt{\frac{1}{1+ax}}} \\
&\quad - \frac{1}{105} \left(8a^5\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int \frac{1}{x^2\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{42ax^7} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{6x^6} + \frac{\sqrt{1-ax}}{42ax^7\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{35x^5\sqrt{\frac{1}{1+ax}}} + \frac{4a^3\sqrt{1-ax}}{105x^3\sqrt{\frac{1}{1+ax}}} + \frac{8a^5\sqrt{1-ax}}{105x\sqrt{\frac{1}{1+ax}}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.52

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^7} dx = \frac{-15 + \sqrt{\frac{1-ax}{1+ax}}(1+ax)^2(-15 + 15ax - 12a^2x^2 + 12a^3x^3 - 8a^4x^4 + 8a^5x^5)}{105ax^7}$$

[In] Integrate[E^ArcSech[a*x]/x^7,x]

[Out] (-15 + Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2*(-15 + 15*a*x - 12*a^2*x^2 + 12*a^3*x^3 - 8*a^4*x^4 + 8*a^5*x^5))/(105*a*x^7)

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.49

method	result	size
default	$\frac{\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} (a^2x^2-1)(8a^4x^4+12a^2x^2+15)}{105x^6} - \frac{1}{7ax^7}$	71

[In] int((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^7,x,method=_RETURNVERBOSE)

[Out] 1/105*((a*x+1)/a/x)^(1/2)/x^6*(-(a*x-1)/a/x)^(1/2)*(a^2*x^2-1)*(8*a^4*x^4+12*a^2*x^2+15)-1/7/a/x^7

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.47

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^7} dx = \frac{(8a^7x^7 + 4a^5x^5 + 3a^3x^3 - 15ax)\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} - 15}{105ax^7}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^7,x, algorithm="fricas")

[Out] 1/105*((8*a^7*x^7 + 4*a^5*x^5 + 3*a^3*x^3 - 15*a*x)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 15)/(a*x^7)

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^7} dx = \int \frac{1}{x^8} dx + \int \frac{a\sqrt{-1+\frac{1}{ax}}\sqrt{1+\frac{1}{ax}}}{x^7} dx$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2))*(1+1/a/x)**(1/2))/x**7,x)

[Out] (Integral(x**(-8), x) + Integral(a*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x))/x**7, x))/a

Maxima [A] (verification not implemented)

none

Time = 0.23 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.41

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^7} dx = \frac{(8a^6x^7 + 4a^4x^5 + 3a^2x^3 - 15x)\sqrt{ax+1}\sqrt{-ax+1}}{105ax^8} - \frac{1}{7ax^7}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))/x^7,x, algorithm="maxima")

[Out] 1/105*(8*a^6*x^7 + 4*a^4*x^5 + 3*a^2*x^3 - 15*x)*sqrt(a*x + 1)*sqrt(-a*x + 1)/(a*x^8) - 1/7/(a*x^7)

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^7} dx = \int \frac{\sqrt{\frac{1}{ax} + 1}\sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}}{x^7} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))/x^7,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))/x^7, x)

Mupad [B] (verification not implemented)

Time = 4.86 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.65

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^7} dx = \frac{\sqrt{\frac{1}{ax} - 1} \left(\frac{a^2 x^2 \sqrt{\frac{1}{ax} + 1}}{35} - \frac{\sqrt{\frac{1}{ax} + 1}}{7} + \frac{4a^4 x^4 \sqrt{\frac{1}{ax} + 1}}{105} + \frac{8a^6 x^6 \sqrt{\frac{1}{ax} + 1}}{105} \right)}{x^6} - \frac{1}{7ax^7}$$

[In] int(((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))/x^7,x)

[Out] ((1/(a*x) - 1)^(1/2)*((a^2*x^2*(1/(a*x) + 1)^(1/2))/35 - (1/(a*x) + 1)^(1/2)/7 + (4*a^4*x^4*(1/(a*x) + 1)^(1/2))/105 + (8*a^6*x^6*(1/(a*x) + 1)^(1/2))/105))/x^6 - 1/(7*a*x^7)

3.44 $\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^8} dx$

Optimal result	319
Rubi [A] (verified)	319
Mathematica [A] (verified)	322
Maple [A] (verified)	322
Fricas [A] (verification not implemented)	323
Sympy [F]	323
Maxima [F]	324
Giac [F]	324
Mupad [B] (verification not implemented)	324

Optimal result

Integrand size = 10, antiderivative size = 194

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^8} dx = \frac{1}{56ax^8} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{7x^7} + \frac{\sqrt{1-ax}}{56ax^8\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{48x^6\sqrt{\frac{1}{1+ax}}} + \frac{5a^3\sqrt{1-ax}}{192x^4\sqrt{\frac{1}{1+ax}}} \\ + \frac{5a^5\sqrt{1-ax}}{128x^2\sqrt{\frac{1}{1+ax}}} + \frac{5}{128}a^7\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\operatorname{arctanh}\left(\sqrt{1-ax}\sqrt{1+ax}\right)$$

[Out] $1/56/a/x^8-1/7*(1/a/x+(1/a/x-1)^{(1/2)}*(1+1/a/x)^{(1/2)})/x^7+1/56*(-a*x+1)^{(1/2)}/a/x^8/(1/(a*x+1))^{(1/2)}+1/48*a*(-a*x+1)^{(1/2)}/x^6/(1/(a*x+1))^{(1/2)}+5/192*a^3*(-a*x+1)^{(1/2)}/x^4/(1/(a*x+1))^{(1/2)}+5/128*a^5*(-a*x+1)^{(1/2)}/x^2/(1/(a*x+1))^{(1/2)}+5/128*a^7*\operatorname{arctanh}((-a*x+1)^{(1/2)}*(a*x+1)^{(1/2)})*(1/(a*x+1))^{(1/2)}*(a*x+1)^{(1/2)}$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6470, 30, 105, 12, 94, 214}

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^8} dx = \frac{5}{128}a^7\sqrt{\frac{1}{ax+1}}\sqrt{ax+1}\operatorname{arctanh}\left(\sqrt{1-ax}\sqrt{ax+1}\right) + \frac{5a^5\sqrt{1-ax}}{128x^2\sqrt{\frac{1}{ax+1}}} \\ + \frac{5a^3\sqrt{1-ax}}{192x^4\sqrt{\frac{1}{ax+1}}} + \frac{\sqrt{1-ax}}{56ax^8\sqrt{\frac{1}{ax+1}}} + \frac{1}{56ax^8} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{7x^7} + \frac{a\sqrt{1-ax}}{48x^6\sqrt{\frac{1}{ax+1}}}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcSech}[a*x]}/x^8, x]$

```
[Out] 1/(56*a*x^8) - E^ArcSech[a*x]/(7*x^7) + Sqrt[1 - a*x]/(56*a*x^8*Sqrt[(1 + a*x)^(-1)]) + (a*Sqrt[1 - a*x])/(48*x^6*Sqrt[(1 + a*x)^(-1)]) + (5*a^3*Sqrt[1 - a*x])/(192*x^4*Sqrt[(1 + a*x)^(-1)]) + (5*a^5*Sqrt[1 - a*x])/(128*x^2*Sqrt[(1 + a*x)^(-1)]) + (5*a^7*Sqrt[(1 + a*x)^(-1)]*Sqrt[1 + a*x]*ArcTanh[Sqrt[1 - a*x]*Sqrt[1 + a*x]])/128
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 30

```
Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]
```

Rule 94

```
Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))), x_Symbol] := Dist[b*f, Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]
```

Rule 105

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 6470

```
Int[E^ArcSech[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*(E^ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] + Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, m, p}, x] && NeQ[m, -1]
```


Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{e^{\operatorname{sech}^{-1}(ax)}}{7x^7} - \frac{\int \frac{1}{x^9} dx}{7a} - \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{x^9\sqrt{1-ax}\sqrt{1+ax}} dx}{7a} \\
&= \frac{1}{56ax^8} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{7x^7} + \frac{\sqrt{1-ax}}{56ax^8\sqrt{\frac{1}{1+ax}}} + \frac{\left(\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int -\frac{7a^2}{x^7\sqrt{1-ax}\sqrt{1+ax}} dx}{56a} \\
&= \frac{1}{56ax^8} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{7x^7} + \frac{\sqrt{1-ax}}{56ax^8\sqrt{\frac{1}{1+ax}}} - \frac{1}{8} \left(a\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{x^7\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{56ax^8} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{7x^7} + \frac{\sqrt{1-ax}}{56ax^8\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{48x^6\sqrt{\frac{1}{1+ax}}} \\
&\quad + \frac{1}{48} \left(a\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int -\frac{5a^2}{x^5\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{56ax^8} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{7x^7} + \frac{\sqrt{1-ax}}{56ax^8\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{48x^6\sqrt{\frac{1}{1+ax}}} \\
&\quad - \frac{1}{48} \left(5a^3\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{x^5\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{56ax^8} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{7x^7} + \frac{\sqrt{1-ax}}{56ax^8\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{48x^6\sqrt{\frac{1}{1+ax}}} + \frac{5a^3\sqrt{1-ax}}{192x^4\sqrt{\frac{1}{1+ax}}} \\
&\quad + \frac{1}{192} \left(5a^3\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int -\frac{3a^2}{x^3\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{56ax^8} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{7x^7} + \frac{\sqrt{1-ax}}{56ax^8\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{48x^6\sqrt{\frac{1}{1+ax}}} + \frac{5a^3\sqrt{1-ax}}{192x^4\sqrt{\frac{1}{1+ax}}} \\
&\quad - \frac{1}{64} \left(5a^5\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{1}{x^3\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{56ax^8} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{7x^7} + \frac{\sqrt{1-ax}}{56ax^8\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{48x^6\sqrt{\frac{1}{1+ax}}} + \frac{5a^3\sqrt{1-ax}}{192x^4\sqrt{\frac{1}{1+ax}}} \\
&\quad + \frac{5a^5\sqrt{1-ax}}{128x^2\sqrt{\frac{1}{1+ax}}} - \frac{1}{128} \left(5a^5\sqrt{\frac{1}{1+ax}}\sqrt{1+ax}\right) \int \frac{a^2}{x\sqrt{1-ax}\sqrt{1+ax}} dx
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{56ax^8} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{7x^7} + \frac{\sqrt{1-ax}}{56ax^8\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{48x^6\sqrt{\frac{1}{1+ax}}} + \frac{5a^3\sqrt{1-ax}}{192x^4\sqrt{\frac{1}{1+ax}}} \\
&\quad + \frac{5a^5\sqrt{1-ax}}{128x^2\sqrt{\frac{1}{1+ax}}} - \frac{1}{128} \left(5a^7\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \int \frac{1}{x\sqrt{1-ax}\sqrt{1+ax}} dx \\
&= \frac{1}{56ax^8} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{7x^7} + \frac{\sqrt{1-ax}}{56ax^8\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{48x^6\sqrt{\frac{1}{1+ax}}} + \frac{5a^3\sqrt{1-ax}}{192x^4\sqrt{\frac{1}{1+ax}}} + \frac{5a^5\sqrt{1-ax}}{128x^2\sqrt{\frac{1}{1+ax}}} \\
&\quad + \frac{1}{128} \left(5a^8\sqrt{\frac{1}{1+ax}}\sqrt{1+ax} \right) \operatorname{Subst} \left(\int \frac{1}{a-ax^2} dx, x, \sqrt{1-ax}\sqrt{1+ax} \right) \\
&= \frac{1}{56ax^8} - \frac{e^{\operatorname{sech}^{-1}(ax)}}{7x^7} + \frac{\sqrt{1-ax}}{56ax^8\sqrt{\frac{1}{1+ax}}} + \frac{a\sqrt{1-ax}}{48x^6\sqrt{\frac{1}{1+ax}}} + \frac{5a^3\sqrt{1-ax}}{192x^4\sqrt{\frac{1}{1+ax}}} \\
&\quad + \frac{5a^5\sqrt{1-ax}}{128x^2\sqrt{\frac{1}{1+ax}}} + \frac{5}{128} a^7 \sqrt{\frac{1}{1+ax}} \sqrt{1+ax} \operatorname{arctanh} \left(\sqrt{1-ax}\sqrt{1+ax} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.75

$$\begin{aligned}
&\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^8} dx \\
&= \frac{-48 + \sqrt{\frac{1-ax}{1+ax}} (-48 - 48ax + 8a^2x^2 + 8a^3x^3 + 10a^4x^4 + 10a^5x^5 + 15a^6x^6 + 15a^7x^7) - 15a^8x^8 \log(x) + 15a^8x^8 \operatorname{Log}\left[1 + \sqrt{\frac{1-ax}{1+ax}}\right] + a^8x^8 \operatorname{Log}\left[1 + \sqrt{\frac{1-ax}{1+ax}}\right]}{384ax^8}
\end{aligned}$$

[In] Integrate[E^ArcSech[a*x]/x^8,x]

[Out] (-48 + Sqrt[(1 - a*x)/(1 + a*x)]*(-48 - 48*a*x + 8*a^2*x^2 + 8*a^3*x^3 + 10*a^4*x^4 + 10*a^5*x^5 + 15*a^6*x^6 + 15*a^7*x^7) - 15*a^8*x^8*Log[x] + 15*a^8*x^8*Log[1 + Sqrt[(1 - a*x)/(1 + a*x)]] + a*x*Sqrt[(1 - a*x)/(1 + a*x)])/(384*a*x^8)

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.78

method	result
default	$\frac{\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} \left(15 \operatorname{arctanh} \left(\frac{1}{\sqrt{-a^2x^2+1}} \right) a^8x^8 + 15\sqrt{-a^2x^2+1} a^6x^6 + 10\sqrt{-a^2x^2+1} a^4x^4 + 8a^2x^2\sqrt{-a^2x^2+1} - 48\sqrt{-a^2x^2+1} \right)}{384x^7\sqrt{-a^2x^2+1}}$

```
[In] int((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^8,x,method=_RETURNVERBOSE)
[Out] 1/384*((a*x+1)/a/x)^(1/2)/x^7*(-(a*x-1)/a/x)^(1/2)*(15*arctanh(1/(-a^2*x^2+
1)^(1/2))*a^8*x^8+15*(-a^2*x^2+1)^(1/2)*a^6*x^6+10*(-a^2*x^2+1)^(1/2)*a^4*x
^4+8*a^2*x^2*(-a^2*x^2+1)^(1/2)-48*(-a^2*x^2+1)^(1/2))/(-a^2*x^2+1)^(1/2)-1
/8/a/x^8
```

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.80

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^8} dx$$

$$= \frac{15 a^8 x^8 \log \left(ax \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} + 1 \right) - 15 a^8 x^8 \log \left(ax \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} - 1 \right) + 2 (15 a^7 x^7 + 10 a^5 x^5 + 8 a^3 x^3 - 48 a x) \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} - 96}{768 a x^8}$$

```
[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^8,x, algorithm="fricas"
)
```

```
[Out] 1/768*(15*a^8*x^8*log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + 1)
- 15*a^8*x^8*log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 1) + 2
*(15*a^7*x^7 + 10*a^5*x^5 + 8*a^3*x^3 - 48*a*x)*sqrt((a*x + 1)/(a*x))*sqrt(
-(a*x - 1)/(a*x)) - 96)/(a*x^8)
```

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^8} dx = \frac{\int \frac{1}{x^9} dx + \int \frac{a \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}}}{x^8} dx}{a}$$

```
[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))/x**8,x)
```

```
[Out] (Integral(x**(-9), x) + Integral(a*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x))/x**
8, x))/a
```

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^8} dx = \int \frac{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}}{x^8} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^8,x, algorithm="maxima")

[Out] integrate(sqrt(a*x + 1)*sqrt(-a*x + 1)/x^9, x)/a - 1/8/(a*x^8)

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^8} dx = \int \frac{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}}{x^8} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^8,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))/x^8, x)

Mupad [B] (verification not implemented)

Time = 43.82 (sec) , antiderivative size = 1155, normalized size of antiderivative = 5.95

$$\int \frac{e^{\operatorname{sech}^{-1}(ax)}}{x^8} dx = \text{Too large to display}$$

[In] int(((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))/x^8,x)

[Out] (5*a^7*atanh(((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1)))/32 - ((1723*a^7*((1/(a*x) - 1)^(1/2) - 1i)^5)/(96*((1/(a*x) + 1)^(1/2) - 1)^5) - (235*a^7*((1/(a*x) - 1)^(1/2) - 1i)^3)/(96*((1/(a*x) + 1)^(1/2) - 1)^3) + (72283*a^7*((1/(a*x) - 1)^(1/2) - 1i)^7)/(32*((1/(a*x) + 1)^(1/2) - 1)^7) + (848801*a^7*((1/(a*x) - 1)^(1/2) - 1i)^9)/(32*((1/(a*x) + 1)^(1/2) - 1)^9) + (4181067*a^7*((1/(a*x) - 1)^(1/2) - 1i)^11)/(32*((1/(a*x) + 1)^(1/2) - 1)^11) + (10994181*a^7*((1/(a*x) - 1)^(1/2) - 1i)^13)/(32*((1/(a*x) + 1)^(1/2) - 1)^13) + (17457599*a^7*((1/(a*x) - 1)^(1/2) - 1i)^15)/(32*((1/(a*x) + 1)^(1/2) - 1)^15) + (17457599*a^7*((1/(a*x) - 1)^(1/2) - 1i)^17)/(32*((1/(a*x) + 1)^(1/2) - 1)^17) + (10994181*a^7*((1/(a*x) - 1)^(1/2) - 1i)^19)/(32*((1/(a*x) + 1)^(1/2) - 1)^19) + (4181067*a^7*((1/(a*x) - 1)^(1/2) - 1i)^21)/(32*((1/(a*x) + 1)^(1/2) - 1)^21) + (848801*a^7*((1/(a*x) - 1)^(1/2) - 1i)^23)/(32*((1/(a*x) + 1)^(1/2) - 1)^23) + (72283*a^7*((1/(a*x) - 1)^(1/2) - 1i)^25)/(32*((1/(a*x) + 1)^(1/2) - 1)^25) + (1723*a^7*((1/(a*x) - 1)^(1/2) -

$$\begin{aligned}
& 1i)^{27}/(96*((1/(a*x) + 1)^{(1/2)} - 1)^{27}) - (235*a^7*((1/(a*x) - 1)^{(1/2)} - \\
& 1i)^{29})/(96*((1/(a*x) + 1)^{(1/2)} - 1)^{29}) + (5*a^7*((1/(a*x) - 1)^{(1/2)} - \\
& 1i)^{31})/(32*((1/(a*x) + 1)^{(1/2)} - 1)^{31}) + (5*a^7*((1/(a*x) - 1)^{(1/2)} - 1 \\
& i))/((32*((1/(a*x) + 1)^{(1/2)} - 1)))/((120*((1/(a*x) - 1)^{(1/2)} - 1i)^4)/((1 \\
& /((a*x) + 1)^{(1/2)} - 1)^4 - (16*((1/(a*x) - 1)^{(1/2)} - 1i)^2)/((1/(a*x) + 1) \\
& ^{(1/2)} - 1)^2 - (560*((1/(a*x) - 1)^{(1/2)} - 1i)^6)/((1/(a*x) + 1)^{(1/2)} - 1 \\
&)^6 + (1820*((1/(a*x) - 1)^{(1/2)} - 1i)^8)/((1/(a*x) + 1)^{(1/2)} - 1)^8 - (43 \\
& 68*((1/(a*x) - 1)^{(1/2)} - 1i)^{10})/((1/(a*x) + 1)^{(1/2)} - 1)^{10} + (8008*((1/ \\
& (a*x) - 1)^{(1/2)} - 1i)^{12})/((1/(a*x) + 1)^{(1/2)} - 1)^{12} - (11440*((1/(a*x) \\
& - 1)^{(1/2)} - 1i)^{14})/((1/(a*x) + 1)^{(1/2)} - 1)^{14} + (12870*((1/(a*x) - 1)^{(\\
& 1/2)} - 1i)^{16})/((1/(a*x) + 1)^{(1/2)} - 1)^{16} - (11440*((1/(a*x) - 1)^{(1/2)} - \\
& 1i)^{18})/((1/(a*x) + 1)^{(1/2)} - 1)^{18} + (8008*((1/(a*x) - 1)^{(1/2)} - 1i)^{20} \\
&)/((1/(a*x) + 1)^{(1/2)} - 1)^{20} - (4368*((1/(a*x) - 1)^{(1/2)} - 1i)^{22})/((1/(\\
& a*x) + 1)^{(1/2)} - 1)^{22} + (1820*((1/(a*x) - 1)^{(1/2)} - 1i)^{24})/((1/(a*x) + \\
& 1)^{(1/2)} - 1)^{24} - (560*((1/(a*x) - 1)^{(1/2)} - 1i)^{26})/((1/(a*x) + 1)^{(1/2)} \\
& - 1)^{26} + (120*((1/(a*x) - 1)^{(1/2)} - 1i)^{28})/((1/(a*x) + 1)^{(1/2)} - 1)^{28} \\
& - (16*((1/(a*x) - 1)^{(1/2)} - 1i)^{30})/((1/(a*x) + 1)^{(1/2)} - 1)^{30} + ((1/(a \\
& *x) - 1)^{(1/2)} - 1i)^{32}/((1/(a*x) + 1)^{(1/2)} - 1)^{32} + 1) - 1/(8*a*x^8)
\end{aligned}$$

3.45 $\int e^{\operatorname{sech}^{-1}(ax^2)} x^7 dx$

Optimal result	326
Rubi [A] (verified)	326
Mathematica [C] (verified)	328
Maple [A] (verified)	328
Fricas [A] (verification not implemented)	329
Sympy [F]	329
Maxima [F]	329
Giac [B] (verification not implemented)	330
Mupad [B] (verification not implemented)	330

Optimal result

Integrand size = 12, antiderivative size = 111

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^7 dx = \frac{x^6}{24a} + \frac{1}{8} e^{\operatorname{sech}^{-1}(ax^2)} x^8 - \frac{x^2 \sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \sqrt{1-a^2x^4}}{16a^3} + \frac{\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \arcsin(ax^2)}{16a^4}$$

[Out] 1/24*x^6/a+1/8*(1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^8+1/16*arcsin(a*x^2)*(1/(a*x^2+1))^(1/2)*(a*x^2+1)^(1/2)/a^4-1/16*x^2*(1/(a*x^2+1))^(1/2)*(a*x^2+1)^(1/2)*(-a^2*x^4+1)^(1/2)/a^3

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6470, 30, 265, 281, 327, 222}

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^7 dx = \frac{\sqrt{\frac{1}{ax^2+1}} \sqrt{ax^2+1} \arcsin(ax^2)}{16a^4} - \frac{x^2 \sqrt{\frac{1}{ax^2+1}} \sqrt{ax^2+1} \sqrt{1-a^2x^4}}{16a^3} + \frac{x^6}{24a} + \frac{1}{8} x^8 e^{\operatorname{sech}^{-1}(ax^2)}$$

[In] Int[E^ArcSech[a*x^2]*x^7,x]

[Out] x^6/(24*a) + (E^ArcSech[a*x^2]*x^8)/8 - (x^2*sqrt[(1 + a*x^2)^(-1)]*sqrt[1 + a*x^2]*sqrt[1 - a^2*x^4])/(16*a^3) + (sqrt[(1 + a*x^2)^(-1)]*sqrt[1 + a*x^2]*ArcSin[a*x^2])/(16*a^4)

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 222

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

Rule 265

Int[((c_)*(x_))^(m_)*((a1_) + (b1_)*(x_)^(n_))^(p_)*((a2_) + (b2_)*(x_)^(n_))^(p_), x_Symbol] := Int[(c*x)^(m*(a1*a2 + b1*b2*x^(2*n)))^p, x] /; FreeQ[{a1, b1, a2, b2, c, m, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))

Rule 281

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

Rule 327

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 6470

Int[E^ArcSech[(a_)*(x_)^(p_)]*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*(E^ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] + Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x)) /; FreeQ[{a, m, p}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{8} e^{\text{sech}^{-1}(ax^2)} x^8 + \frac{\int x^5 dx}{4a} + \frac{\left(\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \int \frac{x^5}{\sqrt{1-ax^2} \sqrt{1+ax^2}} dx}{4a} \\ &= \frac{x^6}{24a} + \frac{1}{8} e^{\text{sech}^{-1}(ax^2)} x^8 + \frac{\left(\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \int \frac{x^5}{\sqrt{1-a^2x^4}} dx}{4a} \end{aligned}$$

$$\begin{aligned}
&= \frac{x^6}{24a} + \frac{1}{8} e^{\operatorname{sech}^{-1}(ax^2)} x^8 + \frac{\left(\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \operatorname{Subst}\left(\int \frac{x^2}{\sqrt{1-a^2x^2}} dx, x, x^2\right)}{8a} \\
&= \frac{x^6}{24a} + \frac{1}{8} e^{\operatorname{sech}^{-1}(ax^2)} x^8 - \frac{x^2 \sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \sqrt{1-a^2x^4}}{16a^3} \\
&\quad + \frac{\left(\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \operatorname{Subst}\left(\int \frac{1}{\sqrt{1-a^2x^2}} dx, x, x^2\right)}{16a^3} \\
&= \frac{x^6}{24a} + \frac{1}{8} e^{\operatorname{sech}^{-1}(ax^2)} x^8 - \frac{x^2 \sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \sqrt{1-a^2x^4}}{16a^3} + \frac{\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \arcsin(ax^2)}{16a^4}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.18 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.00

$$\begin{aligned}
&\int e^{\operatorname{sech}^{-1}(ax^2)} x^7 dx \\
&= \frac{8a^3x^6 - 3a\sqrt{\frac{1-ax^2}{1+ax^2}}(x^2 + ax^4 - 2a^2x^6 - 2a^3x^8) + 3i \log\left(-2iax^2 + 2\sqrt{\frac{1-ax^2}{1+ax^2}}(1+ax^2)\right)}{48a^4}
\end{aligned}$$

[In] Integrate[E^ArcSech[a*x^2]*x^7,x]

[Out] (8*a^3*x^6 - 3*a*Sqrt[(1 - a*x^2)/(1 + a*x^2)]*(x^2 + a*x^4 - 2*a^2*x^6 - 2*a^3*x^8) + (3*I)*Log[(-2*I)*a*x^2 + 2*Sqrt[(1 - a*x^2)/(1 + a*x^2)]*(1 + a*x^2)])/(48*a^4)

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.23

method	result	size
default	$\frac{\sqrt{-\frac{ax^2-1}{ax^2}} x^2 \sqrt{\frac{ax^2+1}{ax^2}} \left(2x^6 \sqrt{-\frac{x^4a^2-1}{a^2}} a^4 - x^2 \sqrt{-\frac{x^4a^2-1}{a^2}} a^2 + \arctan\left(\frac{x^2}{\sqrt{-\frac{x^4a^2-1}{a^2}}}\right) \right)}{16\sqrt{-\frac{x^4a^2-1}{a^2}} a^4} + \frac{x^6}{6a}$	137

[In] int((1/a/x^2+(1/a/x^2-1)^(1/2))*(1/a/x^2+1)^(1/2))*x^7,x,method=_RETURNVERBOSE)

[Out] 1/16*(-(a*x^2-1)/a/x^2)^(1/2)*x^2*((a*x^2+1)/a/x^2)^(1/2)*(2*x^6*(-(a^2*x^4-1)/a^2)^(1/2)*a^4-x^2*(-(a^2*x^4-1)/a^2)^(1/2)*a^2+arctan(x^2/(-(a^2*x^4-1)/a^2)^(1/2)))/(-(a^2*x^4-1)/a^2)^(1/2)/a^4+1/6/a*x^6

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.05

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^7 dx$$

$$= \frac{8a^3x^6 + 3(2a^4x^8 - a^2x^4)\sqrt{\frac{ax^2+1}{ax^2}}\sqrt{-\frac{ax^2-1}{ax^2}} - 6\arctan\left(\frac{ax^2\sqrt{\frac{ax^2+1}{ax^2}}\sqrt{-\frac{ax^2-1}{ax^2}}-1}{ax^2}\right)}{48a^4}$$

```
[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^7,x, algorithm="fricas")
```

```
[Out] 1/48*(8*a^3*x^6 + 3*(2*a^4*x^8 - a^2*x^4)*sqrt((a*x^2 + 1)/(a*x^2))*sqrt(-(a*x^2 - 1)/(a*x^2)) - 6*arctan((a*x^2*sqrt((a*x^2 + 1)/(a*x^2))*sqrt(-(a*x^2 - 1)/(a*x^2)) - 1)/(a*x^2)))/a^4
```

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^7 dx = \frac{\int x^5 dx + \int ax^7 \sqrt{-1 + \frac{1}{ax^2}} \sqrt{1 + \frac{1}{ax^2}} dx}{a}$$

```
[In] integrate((1/a/x**2+(1/a/x**2-1)**(1/2)*(1/a/x**2+1)**(1/2))*x**7,x)
```

```
[Out] (Integral(x**5, x) + Integral(a*x**7*sqrt(-1 + 1/(a*x**2))*sqrt(1 + 1/(a*x**2)), x))/a
```

Maxima [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^7 dx = \int x^7 \left(\sqrt{\frac{1}{ax^2} + 1} \sqrt{\frac{1}{ax^2} - 1} + \frac{1}{ax^2} \right) dx$$

```
[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^7,x, algorithm="maxima")
```

```
[Out] 1/6*x^6/a + integrate(sqrt(a*x^2 + 1)*sqrt(-a*x^2 + 1)*x^5, x)/a
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 205 vs. 2(76) = 152.

Time = 0.32 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.85

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^7 dx$$

$$= \frac{8a^2x^6 + 4\sqrt{a^2x^2+a}\sqrt{-a^2x^2+a}\left((a^2x^2+a)\left(\frac{2(a^2x^2+a)}{a^4} - \frac{7}{a^3}\right) + \frac{9}{a^2}\right) + \left(\sqrt{a^2x^2+a}\sqrt{-a^2x^2+a}\left((a^2x^2+a)\left(\frac{2(a^2x^2+a)}{a^4} - \frac{7}{a^3}\right) + \frac{9}{a^2}\right) + \frac{9}{a^2}\right)}{48a^3}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^7,x, algorithm="giac")

[Out] 1/48*(8*a^2*x^6 + 4*sqrt(a^2*x^2 + a)*sqrt(-a^2*x^2 + a)*((a^2*x^2 + a)*(2*(a^2*x^2 + a)/a^4 - 7/a^3) + 9/a^2) + (sqrt(a^2*x^2 + a)*sqrt(-a^2*x^2 + a)*((a^2*x^2 + a)*(2*(a^2*x^2 + a)*(3*(a^2*x^2 + a)/a^6 - 13/a^5) + 43/a^4) - 39/a^3) - 18*arcsin(1/2*sqrt(2)*sqrt(a^2*x^2 + a)/sqrt(a))/a^2)*a + 24*arcsin(1/2*sqrt(2)*sqrt(a^2*x^2 + a)/sqrt(a))/a/a^3

Mupad [B] (verification not implemented)

Time = 18.82 (sec) , antiderivative size = 521, normalized size of antiderivative = 4.69

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^7 dx = \frac{\ln\left(\frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^2}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^2} + 1\right) \operatorname{li}}{16a^4}$$

$$- \frac{\frac{\operatorname{li}}{2048a^4} + \frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^2 \operatorname{li}}{256a^4 \left(\sqrt{\frac{1}{ax^2}+1-1}\right)^2} + \frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^4 11i}{1024a^4 \left(\sqrt{\frac{1}{ax^2}+1-1}\right)^4} + \frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^6 7i}{512a^4 \left(\sqrt{\frac{1}{ax^2}+1-1}\right)^6} - \frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^8 239i}{2048a^4 \left(\sqrt{\frac{1}{ax^2}+1-1}\right)^8} + \frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^{10} 11i}{512a^4 \left(\sqrt{\frac{1}{ax^2}+1-1}\right)^{10}} + \frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^{12} 11i}{512a^4 \left(\sqrt{\frac{1}{ax^2}+1-1}\right)^{12}}}{\frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^4}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^4} + \frac{4\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^6}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^6} + \frac{6\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^8}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^8} + \frac{4\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^{10}}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^{10}} + \frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^{12}}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^{12}}}$$

$$- \frac{\ln\left(\frac{\sqrt{\frac{1}{ax^2}-1-i}}{\sqrt{\frac{1}{ax^2}+1-1}}\right) \operatorname{li}}{16a^4} + \frac{x^6}{6a} - \frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^2 \operatorname{li}}{512a^4 \left(\sqrt{\frac{1}{ax^2}+1-1}\right)^2} - \frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^4 \operatorname{li}}{2048a^4 \left(\sqrt{\frac{1}{ax^2}+1-1}\right)^4}$$

[In] int(x^7*((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2)),x)

[Out] (log(((1/(a*x^2) - 1)^(1/2) - 1i)^2/((1/(a*x^2) + 1)^(1/2) - 1)^2 + 1)*1i)/(16*a^4) - (1i/(2048*a^4) + (((1/(a*x^2) - 1)^(1/2) - 1i)^2*1i)/(256*a^4*((1/(a*x^2) + 1)^(1/2) - 1)^2) + (((1/(a*x^2) - 1)^(1/2) - 1i)^4*11i)/(1024*a^4*((1/(a*x^2) + 1)^(1/2) - 1)^4) + (((1/(a*x^2) - 1)^(1/2) - 1i)^6*7i)/(512*a^4*((1/(a*x^2) + 1)^(1/2) - 1)^6) - (((1/(a*x^2) - 1)^(1/2) - 1i)^8*239i)/(2048*a^4*((1/(a*x^2) + 1)^(1/2) - 1)^8) + (((1/(a*x^2) - 1)^(1/2) - 1i)^10*11i)/(512*a^4*((1/(a*x^2) + 1)^(1/2) - 1)^10) + (((1/(a*x^2) - 1)^(1/2) - 1i)^12*11i)/(512*a^4*((1/(a*x^2) + 1)^(1/2) - 1)^12))

$$\begin{aligned}
& 2*a^4*((1/(a*x^2) + 1)^{(1/2)} - 1)^6 - (((1/(a*x^2) - 1)^{(1/2)} - 1i)^8*239i \\
&)/(2048*a^4*((1/(a*x^2) + 1)^{(1/2)} - 1)^8) + (((1/(a*x^2) - 1)^{(1/2)} - 1i)^{10}*1i)/(512*a^4*((1/(a*x^2) + 1)^{(1/2)} - 1)^{10}))/(((1/(a*x^2) - 1)^{(1/2)} - 1i)^4/((1/(a*x^2) + 1)^{(1/2)} - 1)^4 + (4*((1/(a*x^2) - 1)^{(1/2)} - 1i)^6)/((1/(a*x^2) + 1)^{(1/2)} - 1)^6 + (6*((1/(a*x^2) - 1)^{(1/2)} - 1i)^8)/((1/(a*x^2) + 1)^{(1/2)} - 1)^8 + (4*((1/(a*x^2) - 1)^{(1/2)} - 1i)^{10})/((1/(a*x^2) + 1)^{(1/2)} - 1)^{10} + ((1/(a*x^2) - 1)^{(1/2)} - 1i)^{12}/((1/(a*x^2) + 1)^{(1/2)} - 1)^{12}) - (\log(((1/(a*x^2) - 1)^{(1/2)} - 1i)/((1/(a*x^2) + 1)^{(1/2)} - 1))*1i)/(16*a^4) + x^6/(6*a) - (((1/(a*x^2) - 1)^{(1/2)} - 1i)^2*1i)/(512*a^4*((1/(a*x^2) + 1)^{(1/2)} - 1)^2) - (((1/(a*x^2) - 1)^{(1/2)} - 1i)^4*1i)/(2048*a^4*((1/(a*x^2) + 1)^{(1/2)} - 1)^4)
\end{aligned}$$

3.46 $\int e^{\operatorname{sech}^{-1}(ax^2)} x^6 dx$

Optimal result	332
Rubi [A] (verified)	332
Mathematica [C] (verified)	334
Maple [A] (verified)	334
Fricas [A] (verification not implemented)	335
Sympy [F]	335
Maxima [F]	335
Giac [F(-2)]	336
Mupad [F(-1)]	336

Optimal result

Integrand size = 12, antiderivative size = 115

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^6 dx = \frac{2x^5}{35a} + \frac{1}{7} e^{\operatorname{sech}^{-1}(ax^2)} x^7 - \frac{2x \sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \sqrt{1-a^2x^4}}{21a^3} + \frac{2 \sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \operatorname{EllipticF}(\arcsin(\sqrt{ax}), -1)}{21a^{7/2}}$$

[Out] $2/35*x^5/a+1/7*(1/a/x^2+(1/a/x^2-1)^{(1/2)}*(1/a/x^2+1)^{(1/2}))*x^7+2/21*\operatorname{EllipticF}(x*a^{(1/2)},I)*(1/(a*x^2+1))^{(1/2)}*(a*x^2+1)^{(1/2)}/a^{(7/2)}-2/21*x*(1/(a*x^2+1))^{(1/2)}*(a*x^2+1)^{(1/2)}*(-a^2*x^4+1)^{(1/2)}/a^3$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {6470, 30, 265, 327, 227}

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^6 dx = \frac{2 \sqrt{\frac{1}{ax^2+1}} \sqrt{ax^2+1} \operatorname{EllipticF}(\arcsin(\sqrt{ax}), -1)}{21a^{7/2}} - \frac{2x \sqrt{\frac{1}{ax^2+1}} \sqrt{ax^2+1} \sqrt{1-a^2x^4}}{21a^3} + \frac{2x^5}{35a} + \frac{1}{7} x^7 e^{\operatorname{sech}^{-1}(ax^2)}$$

[In] Int[E^ArcSech[a*x^2]*x^6,x]

[Out] $(2*x^5)/(35*a) + (E^{\operatorname{ArcSech}[a*x^2]}*x^7)/7 - (2*x*\operatorname{Sqrt}[(1+a*x^2)^{-1}]*\operatorname{Sqrt}[1+a*x^2]*\operatorname{Sqrt}[1-a^2*x^4])/(21*a^3) + (2*\operatorname{Sqrt}[(1+a*x^2)^{-1}]*\operatorname{Sqrt}[1+a*x^2]*\operatorname{EllipticF}[\operatorname{ArcSin}[\operatorname{Sqrt}[a]*x], -1])/(21*a^{(7/2)})$

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 227

Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := Simp[EllipticF[ArcSin[Rt[-b, 4]*(x/Rt[a, 4])], -1]/(Rt[a, 4]*Rt[-b, 4]), x] /; FreeQ[{a, b}, x] && NegQ[b/a] && GtQ[a, 0]

Rule 265

Int[((c_)*(x_))^(m_)*((a1_) + (b1_)*(x_)^(n_))^(p_)*((a2_) + (b2_)*(x_)^(n_))^(p_), x_Symbol] := Int[(c*x)^(m*(a1*a2 + b1*b2*x^(2*n)))^p, x] /; FreeQ[{a1, b1, a2, b2, c, m, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))

Rule 327

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 6470

Int[E^ArcSech[(a_)*(x_)^(p_)]*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*(E^ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] + Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, m, p}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{7} e^{\text{sech}^{-1}(ax^2)} x^7 + \frac{2 \int x^4 dx}{7a} + \frac{\left(2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{x^4}{\sqrt{1-ax^2}\sqrt{1+ax^2}} dx}{7a} \\
 &= \frac{2x^5}{35a} + \frac{1}{7} e^{\text{sech}^{-1}(ax^2)} x^7 + \frac{\left(2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{x^4}{\sqrt{1-a^2x^4}} dx}{7a} \\
 &= \frac{2x^5}{35a} + \frac{1}{7} e^{\text{sech}^{-1}(ax^2)} x^7 - \frac{2x\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\sqrt{1-a^2x^4}}{21a^3} + \frac{\left(2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{1}{\sqrt{1-a^2x^4}} dx}{21a^3}
 \end{aligned}$$

$$= \frac{2x^5}{35a} + \frac{1}{7} e^{\operatorname{sech}^{-1}(ax^2)} x^7 - \frac{2x \sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \sqrt{1-a^2x^4}}{21a^3} + \frac{2 \sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \operatorname{EllipticF}(\arcsin(\sqrt{ax}), -1)}{21a^{7/2}}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.35 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.24

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^6 dx = \frac{2\sqrt{2} \sqrt{\frac{e^{\operatorname{sech}^{-1}(ax^2)}}{1+e^{2\operatorname{sech}^{-1}(ax^2)}}} x^5 \left(-5 - 17e^{2\operatorname{sech}^{-1}(ax^2)} - 67e^{4\operatorname{sech}^{-1}(ax^2)} + 5e^{6\operatorname{sech}^{-1}(ax^2)} + 5 \left(1 + e^{2\operatorname{sech}^{-1}(ax^2)} \right)^{7/2} \right)}{105a \left(1 + e^{2\operatorname{sech}^{-1}(ax^2)} \right)^3 (ax^2)^{5/2}}$$

[In] Integrate[E^ArcSech[a*x^2]*x^6,x]

[Out] (-2*Sqrt[2]*Sqrt[E^ArcSech[a*x^2]/(1 + E^(2*ArcSech[a*x^2]))]*x^5*(-5 - 17*E^(2*ArcSech[a*x^2]) - 67*E^(4*ArcSech[a*x^2]) + 5*E^(6*ArcSech[a*x^2]) + 5*(1 + E^(2*ArcSech[a*x^2]))^(7/2)*Hypergeometric2F1[1/4, 1/2, 5/4, -E^(2*ArcSech[a*x^2])]))/(105*a*(1 + E^(2*ArcSech[a*x^2]))^3*(a*x^2)^(5/2))

Maple [A] (verified)

Time = 1.07 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.99

method	result	size
default	$\frac{\sqrt{-\frac{ax^2-1}{ax^2}} x^2 \sqrt{\frac{ax^2+1}{ax^2}} \left(3a^{\frac{9}{2}} x^9 - 5a^{\frac{5}{2}} x^5 - 2 \operatorname{EllipticF}(x\sqrt{a}, i) \sqrt{-ax^2+1} \sqrt{ax^2+1} + 2x\sqrt{a} \right)}{21a^{\frac{5}{2}} (x^4 a^2 - 1)} + \frac{x^5}{5a}$	114

[In] int((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^6,x,method=_RETURNVERBOSE)

[Out] 1/21*(-(a*x^2-1)/a/x^2)^(1/2)*x^2*((a*x^2+1)/a/x^2)^(1/2)*(3*a^(9/2)*x^9-5*a^(5/2)*x^5-2*EllipticF(x*a^(1/2),I)*(-a*x^2+1)^(1/2)*(a*x^2+1)^(1/2)+2*x*a^(1/2))/a^(5/2)/(a^2*x^4-1)+1/5/a*x^5

Fricas [A] (verification not implemented)

none

Time = 0.08 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.69

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^6 dx = \frac{21 a^2 x^5 + 5 (3 a^3 x^7 - 2 a x^3) \sqrt{\frac{ax^2+1}{ax^2}} \sqrt{\frac{-ax^2-1}{ax^2}} + \frac{10i F(\arcsin(\frac{1}{\sqrt{ax}}) | -1)}{\sqrt{a}}}{105 a^3}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^6,x, algorithm="fricas")

[Out] 1/105*(21*a^2*x^5 + 5*(3*a^3*x^7 - 2*a*x^3)*sqrt((a*x^2 + 1)/(a*x^2))*sqrt(-(a*x^2 - 1)/(a*x^2)) + 10*I*elliptic_f(arcsin(1/(sqrt(a)*x)), -1)/sqrt(a))/a^3

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^6 dx = \frac{\int x^4 dx + \int ax^6 \sqrt{-1 + \frac{1}{ax^2}} \sqrt{1 + \frac{1}{ax^2}} dx}{a}$$

[In] integrate((1/a/x**2+(1/a/x**2-1)**(1/2)*(1/a/x**2+1)**(1/2))*x**6,x)

[Out] (Integral(x**4, x) + Integral(a*x**6*sqrt(-1 + 1/(a*x**2))*sqrt(1 + 1/(a*x**2)), x))/a

Maxima [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^6 dx = \int x^6 \left(\sqrt{\frac{1}{ax^2} + 1} \sqrt{\frac{1}{ax^2} - 1 + \frac{1}{ax^2}} \right) dx$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^6,x, algorithm="maxima")

[Out] 1/5*x^5/a + integrate(sqrt(a*x^2 + 1)*sqrt(-a*x^2 + 1)*x^4, x)/a

Giac [F(-2)]

Exception generated.

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^6 dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2))*(1/a/x^2+1)^(1/2))*x^6,x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx)::OUTPUT:Unable to divide, perhaps due to rounding error%%{1, [0,4,2,1,1,1]%%}+%%{1, [0,4,0,0,0,2]%%} / %%{1, [0,0,0,0,0,3]%%}

Mupad [F(-1)]

Timed out.

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^6 dx = \int x^6 \left(\sqrt{\frac{1}{ax^2} - 1} \sqrt{\frac{1}{ax^2} + 1} + \frac{1}{ax^2} \right) dx$$

[In] int(x^6*((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2)),x)

[Out] int(x^6*((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2)), x)

3.47 $\int e^{\operatorname{sech}^{-1}(ax^2)} x^5 dx$

Optimal result	337
Rubi [A] (warning: unable to verify)	337
Mathematica [A] (verified)	338
Maple [A] (verified)	339
Fricas [A] (verification not implemented)	339
Sympy [F]	339
Maxima [A] (verification not implemented)	340
Giac [B] (verification not implemented)	340
Mupad [B] (verification not implemented)	340

Optimal result

Integrand size = 12, antiderivative size = 58

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^5 dx = \frac{x^4}{12a} + \frac{1}{6} e^{\operatorname{sech}^{-1}(ax^2)} x^6 - \frac{\sqrt{1-ax^2}}{6a^3 \sqrt{\frac{1}{1+ax^2}}}$$

[Out] $1/12*x^4/a+1/6*(1/a/x^2+(1/a/x^2-1)^{(1/2)}*(1/a/x^2+1)^{(1/2)})*x^6-1/6*(-a*x^2+1)^{(1/2)}/a^3/(1/(a*x^2+1))^{(1/2)}$

Rubi [A] (warning: unable to verify)

Time = 0.03 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.22, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6470, 30, 265, 267}

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^5 dx = -\frac{\sqrt{\frac{1}{ax^2+1}} \sqrt{ax^2+1} \sqrt{1-a^2x^4}}{6a^3} + \frac{x^4}{12a} + \frac{1}{6} x^6 e^{\operatorname{sech}^{-1}(ax^2)}$$

[In] Int[E^ArcSech[a*x^2]*x^5,x]

[Out] $x^4/(12*a) + (E^{\operatorname{ArcSech}[a*x^2]}*x^6)/6 - (\operatorname{Sqrt}[(1 + a*x^2)^{-1}]*\operatorname{Sqrt}[1 + a*x^2]*\operatorname{Sqrt}[1 - a^2*x^4])/(6*a^3)$

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 265

```
Int[((c_.)*(x_))^(m_.)*((a1_) + (b1_.)*(x_)^(n_))^(p_)*((a2_) + (b2_.)*(x_)
^(n_))^(p_), x_Symbol] := Int[(c*x)^m*(a1*a2 + b1*b2*x^(2*n))^p, x] /; Free
Q[{a1, b1, a2, b2, c, m, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p]
|| (GtQ[a1, 0] && GtQ[a2, 0]))
```

Rule 267

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n)
^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] &&
NeQ[p, -1]
```

Rule 6470

```
Int[E^ArcSech[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*(E^
ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] +
Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqr
t[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x) /; FreeQ[{a, m, p}, x] && NeQ[m, -1
]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{6} e^{\operatorname{sech}^{-1}(ax^2)} x^6 + \frac{\int x^3 dx}{3a} + \frac{\left(\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \int \frac{x^3}{\sqrt{1-ax^2} \sqrt{1+ax^2}} dx}{3a} \\ &= \frac{x^4}{12a} + \frac{1}{6} e^{\operatorname{sech}^{-1}(ax^2)} x^6 + \frac{\left(\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \int \frac{x^3}{\sqrt{1-a^2x^4}} dx}{3a} \\ &= \frac{x^4}{12a} + \frac{1}{6} e^{\operatorname{sech}^{-1}(ax^2)} x^6 - \frac{\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \sqrt{1-a^2x^4}}{6a^3} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.97

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^5 dx = \frac{x^4}{4a} + \frac{(-1 + ax^2) \sqrt{\frac{1-ax^2}{1+ax^2}} (1 + ax^2)^2}{6a^3}$$

```
[In] Integrate[E^ArcSech[a*x^2]*x^5,x]
```

```
[Out] x^4/(4*a) + ((-1 + a*x^2)*Sqrt[(1 - a*x^2)/(1 + a*x^2)]*(1 + a*x^2)^2)/(6*a
^3)
```

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.03

method	result	size
default	$\frac{\sqrt{-\frac{ax^2-1}{ax^2}} x^2 \sqrt{\frac{ax^2+1}{ax^2}} (x^4 a^2 - 1)}{6a^2} + \frac{x^4}{4a}$	60

[In] int((1/a/x^2+(1/a/x^2-1)^(1/2))*(1/a/x^2+1)^(1/2))*x^5,x,method=_RETURNVERBOSE)

[Out] 1/6*(-(a*x^2-1)/a/x^2)^(1/2)*x^2*((a*x^2+1)/a/x^2)^(1/2)*(a^2*x^4-1)/a^2+1/4*x^4/a

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.03

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^5 dx = \frac{3ax^4 + 2(a^2x^6 - x^2)\sqrt{\frac{ax^2+1}{ax^2}}\sqrt{-\frac{ax^2-1}{ax^2}}}{12a^2}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2))*(1/a/x^2+1)^(1/2))*x^5,x, algorithm="fricas")

[Out] 1/12*(3*a*x^4 + 2*(a^2*x^6 - x^2)*sqrt((a*x^2 + 1)/(a*x^2))*sqrt(-(a*x^2 - 1)/(a*x^2)))/a^2

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^5 dx = \frac{\int x^3 dx + \int ax^5 \sqrt{-1 + \frac{1}{ax^2}} \sqrt{1 + \frac{1}{ax^2}} dx}{a}$$

[In] integrate((1/a/x**2+(1/a/x**2-1)**(1/2))*(1/a/x**2+1)**(1/2))*x**5,x)

[Out] (Integral(x**3, x) + Integral(a*x**5*sqrt(-1 + 1/(a*x**2))*sqrt(1 + 1/(a*x**2)), x))/a

Maxima [A] (verification not implemented)

none

Time = 0.23 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.72

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^5 dx = \frac{x^4}{4a} + \frac{(a^2x^4 - 1)\sqrt{ax^2 + 1}\sqrt{-ax^2 + 1}}{6a^3}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^5,x, algorithm="maxima")

[Out] 1/4*x^4/a + 1/6*(a^2*x^4 - 1)*sqrt(a*x^2 + 1)*sqrt(-a*x^2 + 1)/a^3

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 190 vs. 2(69) = 138.

Time = 0.30 (sec) , antiderivative size = 190, normalized size of antiderivative = 3.28

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^5 dx = \frac{\left(\sqrt{a^2x^2 + a}\sqrt{-a^2x^2 + a} \left((a^2x^2 + a) \left(\frac{2(a^2x^2 + a)}{a^4} - \frac{7}{a^3} \right) + \frac{9}{a^2} \right) + \frac{6 \arcsin\left(\frac{\sqrt{2}\sqrt{a^2x^2 + a}}{2\sqrt{a}}\right)}{a} \right) a - 3 \left(2a^2 \arcsin\left(\frac{\sqrt{2}\sqrt{a^2x^2 + a}}{2\sqrt{a}}\right) \right)}{12a^3}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^5,x, algorithm="giac")

[Out] 1/12*((sqrt(a^2*x^2 + a)*sqrt(-a^2*x^2 + a))*((a^2*x^2 + a)*(2*(a^2*x^2 + a)/a^4 - 7/a^3) + 9/a^2) + 6*arcsin(1/2*sqrt(2)*sqrt(a^2*x^2 + a)/sqrt(a))/a)*a - 3*(2*a^2*arcsin(1/2*sqrt(2)*sqrt(a^2*x^2 + a)/sqrt(a)) - sqrt(a^2*x^2 + a)*(a^2*x^2 - 2*a)*sqrt(-a^2*x^2 + a))/a^2 + 3*((a^2*x^2 + a)^2 - 2*(a^2*x^2 + a)*a)/a^2/a^3

Mupad [B] (verification not implemented)

Time = 5.47 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.98

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^5 dx = \sqrt{\frac{1}{ax^2} - 1} \left(\frac{x^6 \sqrt{\frac{1}{ax^2} + 1}}{6} - \frac{x^2 \sqrt{\frac{1}{ax^2} + 1}}{6a^2} \right) + \frac{x^4}{4a}$$

[In] int(x^5*((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2)),x)

[Out] (1/(a*x^2) - 1)^(1/2)*((x^6*(1/(a*x^2) + 1)^(1/2))/6 - (x^2*(1/(a*x^2) + 1)^(1/2))/(6*a^2)) + x^4/(4*a)

3.48 $\int e^{\operatorname{sech}^{-1}(ax^2)} x^4 dx$

Optimal result	341
Rubi [A] (verified)	341
Mathematica [C] (verified)	343
Maple [A] (verified)	344
Fricas [A] (verification not implemented)	344
Sympy [F]	344
Maxima [F]	345
Giac [F(-2)]	345
Mupad [F(-1)]	345

Optimal result

Integrand size = 12, antiderivative size = 112

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^4 dx = \frac{2x^3}{15a} + \frac{1}{5} e^{\operatorname{sech}^{-1}(ax^2)} x^5 + \frac{2\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} E(\arcsin(\sqrt{ax}) | -1)}{5a^{5/2}} - \frac{2\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \operatorname{EllipticF}(\arcsin(\sqrt{ax}), -1)}{5a^{5/2}}$$

[Out] 2/15*x^3/a+1/5*(1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^5+2/5*EllipticE(x*a^(1/2),I)*(1/(a*x^2+1))^(1/2)*(a*x^2+1)^(1/2)/a^(5/2)-2/5*EllipticF(x*a^(1/2),I)*(1/(a*x^2+1))^(1/2)*(a*x^2+1)^(1/2)/a^(5/2)

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {6470, 30, 265, 313, 227, 1213, 435}

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^4 dx = -\frac{2\sqrt{\frac{1}{ax^2+1}} \sqrt{ax^2+1} \operatorname{EllipticF}(\arcsin(\sqrt{ax}), -1)}{5a^{5/2}} + \frac{2\sqrt{\frac{1}{ax^2+1}} \sqrt{ax^2+1} E(\arcsin(\sqrt{ax}) | -1)}{5a^{5/2}} + \frac{2x^3}{15a} + \frac{1}{5} x^5 e^{\operatorname{sech}^{-1}(ax^2)}$$

[In] Int[E^ArcSech[a*x^2]*x^4,x]

[Out] (2*x^3)/(15*a) + (E^ArcSech[a*x^2]*x^5)/5 + (2*Sqrt[(1 + a*x^2)^(-1)]*Sqrt[1 + a*x^2]*EllipticE[ArcSin[Sqrt[a]*x], -1])/(5*a^(5/2)) - (2*Sqrt[(1 + a*x^2)^(-1)]*Sqrt[1 + a*x^2]*EllipticF[ArcSin[Sqrt[a]*x], -1])/(5*a^(5/2))

Rule 30

```
Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]
```

Rule 227

```
Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := Simp[EllipticF[ArcSin[Rt[-b, 4]*(x/Rt[a, 4])], -1]/(Rt[a, 4]*Rt[-b, 4]), x] /; FreeQ[{a, b}, x] && NegQ[b/a] && GtQ[a, 0]
```

Rule 265

```
Int[((c_)*(x_)^(m_))*((a1_) + (b1_)*(x_)^(n_))^(p_)*((a2_) + (b2_)*(x_)^(n_))^(p_), x_Symbol] := Int[(c*x)^m*(a1*a2 + b1*b2*x^(2*n))^p, x] /; FreeQ[{a1, b1, a2, b2, c, m, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))
```

Rule 313

```
Int[(x_)^2/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[-b/a, 2]}, Dist[-q^(-1), Int[1/Sqrt[a + b*x^4], x], x] + Dist[1/q, Int[(1 + q*x^2)/Sqrt[a + b*x^4], x], x]] /; FreeQ[{a, b}, x] && NegQ[b/a]
```

Rule 435

```
Int[Sqrt[(a_) + (b_)*(x_)^2]/Sqrt[(c_) + (d_)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]/(Sqrt[c]*Rt[-d/c, 2]))*EllipticE[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]
```

Rule 1213

```
Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] := Dist[d/Sqrt[a], Int[Sqrt[1 + e*(x^2/d)]/Sqrt[1 - e*(x^2/d)], x], x] /; FreeQ[{a, c, d, e}, x] && NegQ[c/a] && EqQ[c*d^2 + a*e^2, 0] && GtQ[a, 0]
```

Rule 6470

```
Int[E^ArcSech[(a_)*(x_)^(p_)]*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*(E^ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] + Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, m, p}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{5} e^{\operatorname{sech}^{-1}(ax^2)} x^5 + \frac{2 \int x^2 dx}{5a} + \frac{\left(2\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \int \frac{x^2}{\sqrt{1-ax^2}\sqrt{1+ax^2}} dx}{5a} \\
 &= \frac{2x^3}{15a} + \frac{1}{5} e^{\operatorname{sech}^{-1}(ax^2)} x^5 + \frac{\left(2\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \int \frac{x^2}{\sqrt{1-a^2x^4}} dx}{5a} \\
 &= \frac{2x^3}{15a} + \frac{1}{5} e^{\operatorname{sech}^{-1}(ax^2)} x^5 - \frac{\left(2\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \int \frac{1}{\sqrt{1-a^2x^4}} dx}{5a^2} \\
 &\quad + \frac{\left(2\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \int \frac{1+ax^2}{\sqrt{1-a^2x^4}} dx}{5a^2} \\
 &= \frac{2x^3}{15a} + \frac{1}{5} e^{\operatorname{sech}^{-1}(ax^2)} x^5 - \frac{2\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \operatorname{EllipticF}(\arcsin(\sqrt{ax}), -1)}{5a^{5/2}} \\
 &\quad + \frac{\left(2\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \int \frac{\sqrt{1+ax^2}}{\sqrt{1-a^2x^2}} dx}{5a^2} \\
 &= \frac{2x^3}{15a} + \frac{1}{5} e^{\operatorname{sech}^{-1}(ax^2)} x^5 + \frac{2\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} E(\arcsin(\sqrt{ax}) | -1)}{5a^{5/2}} \\
 &\quad - \frac{2\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \operatorname{EllipticF}(\arcsin(\sqrt{ax}), -1)}{5a^{5/2}}
 \end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.46 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.25

$$\begin{aligned}
 &\int e^{\operatorname{sech}^{-1}(ax^2)} x^4 dx \\
 &= \frac{1}{15} \left(\frac{5x^3}{a} + \frac{3\sqrt{\frac{1-ax^2}{1+ax^2}}(x^3 + ax^5)}{a} \right. \\
 &\quad \left. + \frac{6i\sqrt{\frac{1-ax^2}{1+ax^2}} \sqrt{1-a^2x^4} (E(i\operatorname{arcsinh}(\sqrt{-ax}) | -1) - \operatorname{EllipticF}(i\operatorname{arcsinh}(\sqrt{-ax}), -1))}{(-a)^{5/2}(-1+ax^2)} \right)
 \end{aligned}$$

[In] Integrate[E^ArcSech[a*x^2]*x^4,x]

[Out] ((5*x^3)/a + (3*Sqrt[(1 - a*x^2)/(1 + a*x^2)]*(x^3 + a*x^5))/a + ((6*I)*Sqrt[(1 - a*x^2)/(1 + a*x^2)]*Sqrt[1 - a^2*x^4]*(EllipticE[I*ArcSinh[Sqrt[-a]*x], -1] - EllipticF[I*ArcSinh[Sqrt[-a]*x], -1]))/((-a)^(5/2)*(-1 + a*x^2))/15

Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.21

method	result
default	$\frac{\sqrt{-\frac{ax^2-1}{ax^2}} x^2 \sqrt{\frac{ax^2+1}{ax^2}} \left(a^{\frac{7}{2}} x^7 - x^3 a^{\frac{3}{2}} + 2 \operatorname{EllipticF}(x\sqrt{a}, i) \sqrt{-ax^2+1} \sqrt{ax^2+1} - 2\sqrt{-ax^2+1} \sqrt{ax^2+1} \operatorname{EllipticE}(x\sqrt{a}, i) \right)}{5(x^4 a^2 - 1) a^{\frac{3}{2}}} + \frac{x^3}{3a}$

[In] int((1/a/x^2+(1/a/x^2-1)^(1/2))*(1/a/x^2+1)^(1/2))*x^4,x,method=_RETURNVERBOSE)

[Out] $\frac{1}{5} * (- (a*x^2-1)/a/x^2)^{(1/2)} * x^2 * ((a*x^2+1)/a/x^2)^{(1/2)} * (a^{(7/2)} * x^7 - x^3 * a^{(3/2)} + 2 * \operatorname{EllipticF}(x*a^{(1/2)}, I) * (-a*x^2+1)^{(1/2)} * (a*x^2+1)^{(1/2)} - 2 * (-a*x^2+1)^{(1/2)} * (a*x^2+1)^{(1/2)} * \operatorname{EllipticE}(x*a^{(1/2)}, I)) / (a^2 * x^4 - 1) / a^{(3/2)} + 1/3 * x^3/a$

Fricas [A] (verification not implemented)

none

Time = 0.10 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.81

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^4 dx = \frac{5a^2x^3 + 3(a^3x^5 - 2ax) \sqrt{\frac{ax^2+1}{ax^2}} \sqrt{-\frac{ax^2-1}{ax^2}} - \frac{6i E(\arcsin(\frac{1}{\sqrt{ax}}) | -1)}{\sqrt{a}} + \frac{6i F(\arcsin(\frac{1}{\sqrt{ax}}) | -1)}{\sqrt{a}}}{15a^3}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2))*(1/a/x^2+1)^(1/2))*x^4,x, algorithm="fricas")

[Out] $\frac{1}{15} * (5*a^2*x^3 + 3*(a^3*x^5 - 2*a*x) * \operatorname{sqrt}((a*x^2 + 1)/(a*x^2)) * \operatorname{sqrt}(-(a*x^2 - 1)/(a*x^2)) - 6*I * \operatorname{elliptic}_e(\arcsin(1/(\operatorname{sqrt}(a)*x)), -1)/\operatorname{sqrt}(a) + 6*I * \operatorname{elliptic}_f(\arcsin(1/(\operatorname{sqrt}(a)*x)), -1)/\operatorname{sqrt}(a)) / a^3$

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^4 dx = \frac{\int x^2 dx + \int ax^4 \sqrt{-1 + \frac{1}{ax^2}} \sqrt{1 + \frac{1}{ax^2}} dx}{a}$$

[In] integrate((1/a/x**2+(1/a/x**2-1)**(1/2))*(1/a/x**2+1)**(1/2))*x**4,x)

[Out] (Integral(x**2, x) + Integral(a*x**4*sqrt(-1 + 1/(a*x**2))*sqrt(1 + 1/(a*x**2)), x))/a

Maxima [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^4 dx = \int x^4 \left(\sqrt{\frac{1}{ax^2} + 1} \sqrt{\frac{1}{ax^2} - 1} + \frac{1}{ax^2} \right) dx$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^4,x, algorithm="maxima")

[Out] 1/3*x^3/a + integrate(sqrt(a*x^2 + 1)*sqrt(-a*x^2 + 1)*x^2, x)/a

Giac [F(-2)]

Exception generated.

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^4 dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^4,x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%[1, [0,2,2,1,1,1]%%]+%%[1, [0,2,0,0,0,2]%%] / %%[1, [0,0,0,0,0,3]%%

Mupad [F(-1)]

Timed out.

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^4 dx = \int x^4 \left(\sqrt{\frac{1}{ax^2} - 1} \sqrt{\frac{1}{ax^2} + 1} + \frac{1}{ax^2} \right) dx$$

[In] int(x^4*((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2)),x)

[Out] int(x^4*((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2)), x)

3.49 $\int e^{\operatorname{sech}^{-1}(ax^2)} x^3 dx$

Optimal result	346
Rubi [A] (verified)	346
Mathematica [C] (verified)	348
Maple [A] (verified)	348
Fricas [A] (verification not implemented)	348
Sympy [F]	349
Maxima [F]	349
Giac [B] (verification not implemented)	349
Mupad [B] (verification not implemented)	350

Optimal result

Integrand size = 12, antiderivative size = 63

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^3 dx = \frac{x^2}{4a} + \frac{1}{4} e^{\operatorname{sech}^{-1}(ax^2)} x^4 + \frac{\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \arcsin(ax^2)}{4a^2}$$

[Out] $1/4*x^2/a+1/4*(1/a/x^2+(1/a/x^2-1)^{(1/2)}*(1/a/x^2+1)^{(1/2}))*x^4+1/4*\arcsin(a*x^2)*(1/(a*x^2+1))^{(1/2)}*(a*x^2+1)^{(1/2)}/a^2$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {6470, 30, 265, 281, 222}

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^3 dx = \frac{\sqrt{\frac{1}{ax^2+1}} \sqrt{ax^2+1} \arcsin(ax^2)}{4a^2} + \frac{x^2}{4a} + \frac{1}{4} x^4 e^{\operatorname{sech}^{-1}(ax^2)}$$

[In] Int[E^ArcSech[a*x^2]*x^3,x]

[Out] $x^2/(4*a) + (E^{\operatorname{ArcSech}[a*x^2]}*x^4)/4 + (\operatorname{Sqrt}[(1 + a*x^2)^{-1}]*\operatorname{Sqrt}[1 + a*x^2]*\operatorname{ArcSin}[a*x^2])/(4*a^2)$

Rule 30

Int[(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 222

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

Rule 265

Int[((c_)*(x_))^(m_)*((a1_) + (b1_)*(x_)^(n_))^(p_)*((a2_) + (b2_)*(x_)^(n_))^(p_), x_Symbol] := Int[(c*x)^m*(a1*a2 + b1*b2*x^(2*n))^p, x] /; FreeQ[{a1, b1, a2, b2, c, m, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))

Rule 281

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

Rule 6470

Int[E^ArcSech[(a_)*(x_)^(p_)]*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*(E^ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] + Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, m, p}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{4} e^{\text{sech}^{-1}(ax^2)} x^4 + \frac{\int x dx}{2a} + \frac{\left(\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \int \frac{x}{\sqrt{1-ax^2} \sqrt{1+ax^2}} dx}{2a} \\
 &= \frac{x^2}{4a} + \frac{1}{4} e^{\text{sech}^{-1}(ax^2)} x^4 + \frac{\left(\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \int \frac{x}{\sqrt{1-a^2x^4}} dx}{2a} \\
 &= \frac{x^2}{4a} + \frac{1}{4} e^{\text{sech}^{-1}(ax^2)} x^4 + \frac{\left(\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \text{Subst}\left(\int \frac{1}{\sqrt{1-a^2x^2}} dx, x, x^2\right)}{4a} \\
 &= \frac{x^2}{4a} + \frac{1}{4} e^{\text{sech}^{-1}(ax^2)} x^4 + \frac{\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \arcsin(ax^2)}{4a^2}
 \end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.46

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^3 dx = \frac{2ax^2 + a\sqrt{\frac{1-ax^2}{1+ax^2}}(x^2 + ax^4) + i \log\left(-2iax^2 + 2\sqrt{\frac{1-ax^2}{1+ax^2}}(1 + ax^2)\right)}{4a^2}$$

[In] Integrate[E^ArcSech[a*x^2]*x^3,x]

[Out] (2*a*x^2 + a*Sqrt[(1 - a*x^2)/(1 + a*x^2)]*(x^2 + a*x^4) + I*Log[(-2*I)*a*x^2 + 2*Sqrt[(1 - a*x^2)/(1 + a*x^2)]*(1 + a*x^2)])/(4*a^2)

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.78

method	result	size
default	$\frac{\sqrt{-\frac{ax^2-1}{ax^2}} x^2 \sqrt{\frac{ax^2+1}{ax^2}} \left(x^2 \sqrt{-\frac{x^4 a^2-1}{a^2}} a^2 + \arctan\left(\frac{x^2}{\sqrt{-\frac{x^4 a^2-1}{a^2}}}\right) \right)}{4\sqrt{-\frac{x^4 a^2-1}{a^2}} a^2} + \frac{x^2}{2a}$	112

[In] int((1/a/x^2+(1/a/x^2-1)^(1/2))*(1/a/x^2+1)^(1/2))*x^3,x,method=_RETURNVERBOSE)

[Out] 1/4*(-(a*x^2-1)/a/x^2)^(1/2)*x^2*((a*x^2+1)/a/x^2)^(1/2)*(x^2*(-(a^2*x^4-1)/a^2)^(1/2)*a^2+arctan(x^2/(-(a^2*x^4-1)/a^2)^(1/2)))/(-(a^2*x^4-1)/a^2)^(1/2)/a^2+1/2*x^2/a

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.62

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^3 dx = \frac{a^2 x^4 \sqrt{\frac{ax^2+1}{ax^2}} \sqrt{-\frac{ax^2-1}{ax^2}} + 2ax^2 - 2 \arctan\left(\frac{ax^2 \sqrt{\frac{ax^2+1}{ax^2}} \sqrt{-\frac{ax^2-1}{ax^2}} - 1}{ax^2}\right)}{4a^2}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2))*(1/a/x^2+1)^(1/2))*x^3,x, algorithm="fricas")

[Out] 1/4*(a^2*x^4*sqrt((a*x^2 + 1)/(a*x^2))*sqrt(-(a*x^2 - 1)/(a*x^2)) + 2*a*x^2 - 2*arctan((a*x^2*sqrt((a*x^2 + 1)/(a*x^2))*sqrt(-(a*x^2 - 1)/(a*x^2)) - 1)/(a*x^2)))/a^2

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^3 dx = \frac{\int x dx + \int ax^3 \sqrt{-1 + \frac{1}{ax^2}} \sqrt{1 + \frac{1}{ax^2}} dx}{a}$$

[In] integrate((1/a/x**2+(1/a/x**2-1)**(1/2)*(1/a/x**2+1)**(1/2))*x**3,x)

[Out] (Integral(x, x) + Integral(a*x**3*sqrt(-1 + 1/(a*x**2))*sqrt(1 + 1/(a*x**2)), x))/a

Maxima [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^3 dx = \int x^3 \left(\sqrt{\frac{1}{ax^2} + 1} \sqrt{\frac{1}{ax^2} - 1} + \frac{1}{ax^2} \right) dx$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^3,x, algorithm="maxima")

[Out] 1/2*x^2/a + integrate(sqrt(a*x^2 + 1)*sqrt(-a*x^2 + 1)*x, x)/a

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 132 vs. 2(56) = 112.

Time = 0.30 (sec) , antiderivative size = 132, normalized size of antiderivative = 2.10

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^3 dx$$

$$= \frac{2a^2x^2 + 4a \arcsin\left(\frac{\sqrt{2}\sqrt{a^2x^2+a}}{2\sqrt{a}}\right) + 2\sqrt{a^2x^2+a}\sqrt{-a^2x^2+a} + 2a - \frac{2a^2 \arcsin\left(\frac{\sqrt{2}\sqrt{a^2x^2+a}}{2\sqrt{a}}\right) - \sqrt{a^2x^2+a}(a^2x^2-2a)}{a}}{4a^3}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^3,x, algorithm="giac")

[Out] 1/4*(2*a^2*x^2 + 4*a*arcsin(1/2*sqrt(2)*sqrt(a^2*x^2 + a)/sqrt(a)) + 2*sqrt(a^2*x^2 + a)*sqrt(-a^2*x^2 + a) + 2*a - (2*a^2*arcsin(1/2*sqrt(2)*sqrt(a^2*x^2 + a)/sqrt(a)) - sqrt(a^2*x^2 + a)*(a^2*x^2 - 2*a)*sqrt(-a^2*x^2 + a))/a)/a^3

Mupad [B] (verification not implemented)

Time = 10.88 (sec) , antiderivative size = 306, normalized size of antiderivative = 4.86

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^3 dx = \frac{\ln\left(\frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^2}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^2} + 1\right) \operatorname{li} - \ln\left(\frac{\sqrt{\frac{1}{ax^2}-1-i}}{\sqrt{\frac{1}{ax^2}+1-1}}\right) \operatorname{li}}{4a^2} + \frac{\frac{\operatorname{li}}{64a^2} + \frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^2 \operatorname{li}}{32a^2 \left(\sqrt{\frac{1}{ax^2}+1-1}\right)^2} - \frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^4 15i}{64a^2 \left(\sqrt{\frac{1}{ax^2}+1-1}\right)^4}}{\frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^2}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^2} + \frac{2\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^4}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^4} + \frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^6}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^6}} + \frac{x^2}{2a} + \frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^2 \operatorname{li}}{64a^2 \left(\sqrt{\frac{1}{ax^2}+1-1}\right)^2}$$

[In] int(x^3*((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2)),x)

```
[Out] (log(((1/(a*x^2) - 1)^(1/2) - 1i)^2/((1/(a*x^2) + 1)^(1/2) - 1)^2 + 1)*1i)/
(4*a^2) - (log(((1/(a*x^2) - 1)^(1/2) - 1i)/((1/(a*x^2) + 1)^(1/2) - 1))*1i
)/(4*a^2) + (1i/(64*a^2) + (((1/(a*x^2) - 1)^(1/2) - 1i)^2*1i)/(32*a^2*((1/
(a*x^2) + 1)^(1/2) - 1)^2) - (((1/(a*x^2) - 1)^(1/2) - 1i)^4*15i)/(64*a^2*(
(1/(a*x^2) + 1)^(1/2) - 1)^4))/(((1/(a*x^2) - 1)^(1/2) - 1i)^2/((1/(a*x^2)
+ 1)^(1/2) - 1)^2 + (2*((1/(a*x^2) - 1)^(1/2) - 1i)^4)/((1/(a*x^2) + 1)^(1/
2) - 1)^4 + ((1/(a*x^2) - 1)^(1/2) - 1i)^6/((1/(a*x^2) + 1)^(1/2) - 1)^6) +
x^2/(2*a) + (((1/(a*x^2) - 1)^(1/2) - 1i)^2*1i)/(64*a^2*((1/(a*x^2) + 1)^(
1/2) - 1)^2)
```

3.50 $\int e^{\operatorname{sech}^{-1}(ax^2)} x^2 dx$

Optimal result	351
Rubi [A] (verified)	351
Mathematica [C] (verified)	352
Maple [A] (verified)	353
Fricas [A] (verification not implemented)	353
Sympy [F]	353
Maxima [F]	354
Giac [F(-2)]	354
Mupad [F(-1)]	354

Optimal result

Integrand size = 12, antiderivative size = 67

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^2 dx = \frac{2x}{3a} + \frac{1}{3} e^{\operatorname{sech}^{-1}(ax^2)} x^3 + \frac{2\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \operatorname{EllipticF}(\arcsin(\sqrt{ax}), -1)}{3a^{3/2}}$$

[Out] $2/3*x/a+1/3*(1/a/x^2+(1/a/x^2-1)^{(1/2)}*(1/a/x^2+1)^{(1/2)})*x^3+2/3*\operatorname{EllipticF}(x*a^{(1/2)}, I)*(1/(a*x^2+1))^{(1/2)}*(a*x^2+1)^{(1/2)}/a^{(3/2)}$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6470, 8, 254, 227}

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^2 dx = \frac{2\sqrt{\frac{1}{ax^2+1}} \sqrt{ax^2+1} \operatorname{EllipticF}(\arcsin(\sqrt{ax}), -1)}{3a^{3/2}} + \frac{1}{3} x^3 e^{\operatorname{sech}^{-1}(ax^2)} + \frac{2x}{3a}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcSech}[a*x^2]}*x^2, x]$

[Out] $(2*x)/(3*a) + (E^{\operatorname{ArcSech}[a*x^2]}*x^3)/3 + (2*\operatorname{Sqrt}[(1 + a*x^2)^{-1}]*\operatorname{Sqrt}[1 + a*x^2]*\operatorname{EllipticF}[\operatorname{ArcSin}[\operatorname{Sqrt}[a]*x], -1])/(3*a^{(3/2)})$

Rule 8

$\operatorname{Int}[a_, x_Symbol] := \operatorname{Simp}[a*x, x] /; \operatorname{FreeQ}[a, x]$

Rule 227

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_) + (b_.)*(x_)^4], x_Symbol] := \operatorname{Simp}[\operatorname{EllipticF}[\operatorname{ArcSin}[\operatorname{Rt}[-b, 4]*(x/\operatorname{Rt}[a, 4])], -1]/(\operatorname{Rt}[a, 4]*\operatorname{Rt}[-b, 4]), x] /; \operatorname{FreeQ}\{a, b\}, x \&\& \operatorname{NegQ}[$

b/a] && GtQ[a, 0]

Rule 254

```
Int[((a1_.) + (b1_.)*(x_)^(n_))^(p_.)*((a2_.) + (b2_.)*(x_)^(n_))^(p_.), x_
Symbol] := Int[(a1*a2 + b1*b2*x^(2*n))^p, x] /; FreeQ[{a1, b1, a2, b2, n, p
}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]
))
```

Rule 6470

```
Int[E^ArcSech[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*(E^
ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] +
Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sq
rt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, m, p}, x] && NeQ[m, -1
]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{3} e^{\operatorname{sech}^{-1}(ax^2)} x^3 + \frac{2 \int 1 dx}{3a} + \frac{\left(2\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \int \frac{1}{\sqrt{1-ax^2}\sqrt{1+ax^2}} dx}{3a} \\ &= \frac{2x}{3a} + \frac{1}{3} e^{\operatorname{sech}^{-1}(ax^2)} x^3 + \frac{\left(2\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2}\right) \int \frac{1}{\sqrt{1-a^2x^4}} dx}{3a} \\ &= \frac{2x}{3a} + \frac{1}{3} e^{\operatorname{sech}^{-1}(ax^2)} x^3 + \frac{2\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \operatorname{EllipticF}\left(\arcsin(\sqrt{ax}), -1\right)}{3a^{3/2}} \end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.27 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.67

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^2 dx = \frac{2\sqrt{2}e^{-\operatorname{sech}^{-1}(ax^2)} \left(\frac{e^{\operatorname{sech}^{-1}(ax^2)}}{1+e^{2\operatorname{sech}^{-1}(ax^2)}}\right)^{3/2} x \left(-1 - 2e^{2\operatorname{sech}^{-1}(ax^2)} + \left(1 + e^{2\operatorname{sech}^{-1}(ax^2)}\right)^{3/2} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -E^{2\operatorname{sech}^{-1}(ax^2)}\right)\right)}{3a\sqrt{ax^2}}$$

[In] Integrate[E^ArcSech[a*x^2]*x^2,x]

```
[Out] (-2*Sqrt[2]*(E^ArcSech[a*x^2]/(1 + E^(2*ArcSech[a*x^2])))^(3/2)*x*(-1 - 2*E
^(2*ArcSech[a*x^2]) + (1 + E^(2*ArcSech[a*x^2]))^(3/2)*Hypergeometric2F1[1/
4, 1/2, 5/4, -E^(2*ArcSech[a*x^2])])/(3*a*E^ArcSech[a*x^2]*Sqrt[a*x^2])
```


Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.52

method	result	size
default	$\frac{\sqrt{-\frac{ax^2-1}{ax^2}} x^2 \sqrt{\frac{ax^2+1}{ax^2}} \left(a^{\frac{5}{2}} x^5 - 2 \operatorname{EllipticF}(x\sqrt{a}, i) \sqrt{-ax^2+1} \sqrt{ax^2+1} - x\sqrt{a} \right)}{3(x^4 a^2 - 1)\sqrt{a}} + \frac{x}{a}$	102

```
[In] int((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^2,x,method=_RETURNVERBOSE)
```

```
[Out] 1/3*(-(a*x^2-1)/a/x^2)^(1/2)*x^2*((a*x^2+1)/a/x^2)^(1/2)*(a^(5/2)*x^5-2*EllipticF(x*a^(1/2),I)*(-a*x^2+1)^(1/2)*(a*x^2+1)^(1/2)-x*a^(1/2))/(a^2*x^4-1)/a^(1/2)+x/a
```

Fricas [A] (verification not implemented)

none

Time = 0.08 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.93

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^2 dx = \frac{ax^3 \sqrt{\frac{ax^2+1}{ax^2}} \sqrt{-\frac{ax^2-1}{ax^2}} + 3x + \frac{2i F(\arcsin(\frac{1}{\sqrt{ax}}) | -1)}{\sqrt{a}}}{3a}$$

```
[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^2,x, algorithm="fricas")
```

```
[Out] 1/3*(a*x^3*sqrt((a*x^2 + 1)/(a*x^2))*sqrt(-(a*x^2 - 1)/(a*x^2)) + 3*x + 2*I*elliptic_f(arcsin(1/(sqrt(a)*x)), -1)/sqrt(a))/a
```

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^2 dx = \frac{\int 1 dx + \int ax^2 \sqrt{-1 + \frac{1}{ax^2}} \sqrt{1 + \frac{1}{ax^2}} dx}{a}$$

```
[In] integrate((1/a/x**2+(1/a/x**2-1)**(1/2)*(1/a/x**2+1)**(1/2))*x**2,x)
```

```
[Out] (Integral(1, x) + Integral(a*x**2*sqrt(-1 + 1/(a*x**2))*sqrt(1 + 1/(a*x**2)), x))/a
```

Maxima [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^2 dx = \int x^2 \left(\sqrt{\frac{1}{ax^2} + 1} \sqrt{\frac{1}{ax^2} - 1} + \frac{1}{ax^2} \right) dx$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^2,x, algorithm="maxima")

[Out] x/a + integrate(sqrt(a*x^2 + 1)*sqrt(-a*x^2 + 1), x)/a

Giac [F(-2)]

Exception generated.

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^2 dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Unable to divide, perhaps due to rounding error%%{1, [0,2,1,1,1]%%}+%%{1, [0,0,0,0,2]%%} / %%{1, [0,0,0,0,3]%%}%} Err

Mupad [F(-1)]

Timed out.

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^2 dx = \int x^2 \left(\sqrt{\frac{1}{ax^2} - 1} \sqrt{\frac{1}{ax^2} + 1} + \frac{1}{ax^2} \right) dx$$

[In] int(x^2*((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2)),x)

[Out] int(x^2*((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2)), x)

3.51 $\int e^{\operatorname{sech}^{-1}(ax^2)} x dx$

Optimal result	355
Rubi [A] (verified)	355
Mathematica [A] (verified)	357
Maple [C] (verified)	357
Fricas [B] (verification not implemented)	357
Sympy [F]	358
Maxima [F]	358
Giac [F(-2)]	358
Mupad [B] (verification not implemented)	359

Optimal result

Integrand size = 10, antiderivative size = 68

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x dx = \frac{1}{2} e^{\operatorname{sech}^{-1}(ax^2)} x^2 - \frac{\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \operatorname{arctanh}(\sqrt{1-a^2x^4})}{2a} + \frac{\log(x)}{a}$$

[Out] $1/2*(1/a/x^2+(1/a/x^2-1)^{(1/2)}*(1/a/x^2+1)^{(1/2)})*x^2+\ln(x)/a-1/2*\operatorname{arctanh}((-a^2*x^4+1)^{(1/2)}*(1/(a*x^2+1))^{(1/2)}*(a*x^2+1)^{(1/2)}/a$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6470, 29, 265, 272, 65, 214}

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x dx = -\frac{\sqrt{\frac{1}{ax^2+1}} \sqrt{ax^2+1} \operatorname{arctanh}(\sqrt{1-a^2x^4})}{2a} + \frac{1}{2} x^2 e^{\operatorname{sech}^{-1}(ax^2)} + \frac{\log(x)}{a}$$

[In] `Int[E^ArcSech[a*x^2]*x,x]`

[Out] $(E^{\operatorname{ArcSech}[a*x^2]}*x^2)/2 - (\operatorname{Sqrt}[(1+a*x^2)^{-1}]*\operatorname{Sqrt}[1+a*x^2]*\operatorname{ArcTanh}[\operatorname{Sqrt}[1-a^2*x^4]])/(2*a) + \operatorname{Log}[x]/a$

Rule 29

`Int[(x_)^(-1), x_Symbol] :> Simp[Log[x], x]`

Rule 65

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m+1)-1)*(c-a*(d/b) +`

$d*(x^p/b)^n, x], x, (a + b*x)^{(1/p)}, x]] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 214

$\text{Int}[(a + (b*x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{NegQ}[a/b]$

Rule 265

$\text{Int}[(c*x)^m*(a1 + (b1*x)^n)^p*(a2 + (b2*x)^n)^p, x_Symbol] \rightarrow \text{Int}[(c*x)^m*(a1*a2 + b1*b2*x^{2*n})^p, x] /; \text{FreeQ}\{a1, b1, a2, b2, c, m, n, p\}, x] \&\& \text{EqQ}[a2*b1 + a1*b2, 0] \&\& (\text{IntegerQ}[p] \vee (\text{GtQ}[a1, 0] \&\& \text{GtQ}[a2, 0]))$

Rule 272

$\text{Int}[x^m*(a + (b*x)^n)^p, x_Symbol] \rightarrow \text{Dist}[1/n, \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)*(a + b*x)^p}, x], x, x^n], x] /; \text{FreeQ}\{a, b, m, n, p\}, x] \&\& \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]$

Rule 6470

$\text{Int}[E^{\text{ArcSech}[(a*x)^p]}*(x)^m, x_Symbol] \rightarrow \text{Simp}[x^{m+1}*(E^{\text{ArcSech}[a*x^p]/(m+1)}, x] + (\text{Dist}[p/(a*(m+1)), \text{Int}[x^{m-p}, x], x] + \text{Dist}[p*(\text{Sqrt}[1 + a*x^p]/(a*(m+1)))*\text{Sqrt}[1/(1 + a*x^p)], \text{Int}[x^{m-p}/(\text{Sqrt}[1 + a*x^p]*\text{Sqrt}[1 - a*x^p]), x], x]) /; \text{FreeQ}\{a, m, p\}, x] \&\& \text{NeQ}[m, -1]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{2}e^{\text{sech}^{-1}(ax^2)}x^2 + \frac{\int \frac{1}{x} dx}{a} + \frac{\left(\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{1}{x\sqrt{1-ax^2}\sqrt{1+ax^2}} dx}{a} \\
 &= \frac{1}{2}e^{\text{sech}^{-1}(ax^2)}x^2 + \frac{\log(x)}{a} + \frac{\left(\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{1}{x\sqrt{1-a^2x^4}} dx}{a} \\
 &= \frac{1}{2}e^{\text{sech}^{-1}(ax^2)}x^2 + \frac{\log(x)}{a} + \frac{\left(\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \text{Subst}\left(\int \frac{1}{x\sqrt{1-a^2x}} dx, x, x^4\right)}{4a} \\
 &= \frac{1}{2}e^{\text{sech}^{-1}(ax^2)}x^2 + \frac{\log(x)}{a} - \frac{\left(\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \text{Subst}\left(\int \frac{1}{\frac{1}{a^2}-\frac{x^2}{a^2}} dx, x, \sqrt{1-a^2x^4}\right)}{2a^3} \\
 &= \frac{1}{2}e^{\text{sech}^{-1}(ax^2)}x^2 - \frac{\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\text{arctanh}(\sqrt{1-a^2x^4})}{2a} + \frac{\log(x)}{a}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.47

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x dx = \frac{\sqrt{\frac{1-ax^2}{1+ax^2}}(1+ax^2) + 2 \log(ax^2) - \log\left(1 + \sqrt{\frac{1-ax^2}{1+ax^2}} + ax^2 \sqrt{\frac{1-ax^2}{1+ax^2}}\right)}{2a}$$

[In] Integrate[E^ArcSech[a*x^2]*x,x]

[Out] (Sqrt[(1 - a*x^2)/(1 + a*x^2)]*(1 + a*x^2) + 2*Log[a*x^2] - Log[1 + Sqrt[(1 - a*x^2)/(1 + a*x^2)] + a*x^2*Sqrt[(1 - a*x^2)/(1 + a*x^2)]])/(2*a)

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.11 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.87

method	result	size
default	$\frac{\sqrt{-\frac{ax^2-1}{ax^2}} x^2 \sqrt{\frac{ax^2+1}{ax^2}} \left(\operatorname{csgn}\left(\frac{1}{a}\right) a \sqrt{-\frac{x^4 a^2 - 1}{a^2}} - \ln\left(\frac{2 \operatorname{csgn}\left(\frac{1}{a}\right) a \sqrt{-\frac{x^4 a^2 - 1}{a^2}} + 2}{a^2 x^2}\right) \right) \operatorname{csgn}\left(\frac{1}{a}\right)}{2a \sqrt{-\frac{x^4 a^2 - 1}{a^2}}} + \frac{\ln(x)}{a}$	127

[In] int((1/a/x^2+(1/a/x^2-1)^(1/2))*(1/a/x^2+1)^(1/2))*x,x,method=_RETURNVERBOSE)

[Out] 1/2*(-(a*x^2-1)/a/x^2)^(1/2)*x^2*((a*x^2+1)/a/x^2)^(1/2)*(csgn(1/a)*a*(-(a^2*x^4-1)/a^2)^(1/2)-ln(2*(csgn(1/a)*a*(-(a^2*x^4-1)/a^2)^(1/2)+1)/a^2/x^2))*csgn(1/a)/a/(-(a^2*x^4-1)/a^2)^(1/2)+ln(x)/a

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 133 vs. 2(61) = 122.

Time = 0.25 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.96

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x dx = \frac{2ax^2 \sqrt{\frac{ax^2+1}{ax^2}} \sqrt{-\frac{ax^2-1}{ax^2}} - \log\left(ax^2 \sqrt{\frac{ax^2+1}{ax^2}} \sqrt{-\frac{ax^2-1}{ax^2}} + 1\right) + \log\left(ax^2 \sqrt{\frac{ax^2+1}{ax^2}} \sqrt{-\frac{ax^2-1}{ax^2}} - 1\right) + 4 \log(x)}{4a}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2))*(1/a/x^2+1)^(1/2))*x,x, algorithm="fricas")

[Out] $\frac{1}{4} * (2 * a * x^2 * \sqrt{(a * x^2 + 1) / (a * x^2)}) * \sqrt{-(a * x^2 - 1) / (a * x^2)} - \log(a * x^2 * \sqrt{(a * x^2 + 1) / (a * x^2)}) * \sqrt{-(a * x^2 - 1) / (a * x^2)} + 1) + \log(a * x^2 * \sqrt{(a * x^2 + 1) / (a * x^2)}) * \sqrt{-(a * x^2 - 1) / (a * x^2)} - 1) + 4 * \log(x) / a$

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x dx = \frac{\int \frac{1}{x} dx + \int ax \sqrt{-1 + \frac{1}{ax^2}} \sqrt{1 + \frac{1}{ax^2}} dx}{a}$$

[In] `integrate((1/a/x**2+(1/a/x**2-1)**(1/2))*(1/a/x**2+1)**(1/2))*x,x)`

[Out] `(Integral(1/x, x) + Integral(a*x*sqrt(-1 + 1/(a*x**2))*sqrt(1 + 1/(a*x**2)), x))/a`

Maxima [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x dx = \int x \left(\sqrt{\frac{1}{ax^2} + 1} \sqrt{\frac{1}{ax^2} - 1 + \frac{1}{ax^2}} \right) dx$$

[In] `integrate((1/a/x^2+(1/a/x^2-1)^(1/2))*(1/a/x^2+1)^(1/2))*x,x, algorithm="maxima")`

[Out] `integrate(sqrt(a*x^2 + 1)*sqrt(-a*x^2 + 1)/x, x)/a + log(x)/a`

Giac [F(-2)]

Exception generated.

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x dx = \text{Exception raised: TypeError}$$

[In] `integrate((1/a/x^2+(1/a/x^2-1)^(1/2))*(1/a/x^2+1)^(1/2))*x,x, algorithm="giac")`

[Out] `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Limit: Max order reached or unable to make series expansion Error: Bad Argument Value`

Mupad [B] (verification not implemented)

Time = 7.09 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.68

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x dx = \frac{\ln(x)}{a} - \frac{2 \operatorname{atanh}\left(\frac{\sqrt{\frac{1}{ax^2}-1-i}}{\sqrt{\frac{1}{ax^2}+1-1}}\right)}{a} + \frac{\frac{5\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^2}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^2} + 1}{\frac{8a\left(\sqrt{\frac{1}{ax^2}-1-i}\right)}{\sqrt{\frac{1}{ax^2}+1-1}} + \frac{8a\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^3}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^3}} + \frac{\sqrt{\frac{1}{ax^2}-1-i}}{8a\left(\sqrt{\frac{1}{ax^2}+1-1}\right)}$$

[In] int(x*((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2)),x)

```
[Out] log(x)/a - (2*atanh(((1/(a*x^2) - 1)^(1/2) - 1i)/((1/(a*x^2) + 1)^(1/2) - 1
))) / a + ((5*((1/(a*x^2) - 1)^(1/2) - 1i)^2)/((1/(a*x^2) + 1)^(1/2) - 1)^2 +
1)/((8*a*((1/(a*x^2) - 1)^(1/2) - 1i))/((1/(a*x^2) + 1)^(1/2) - 1) + (8*a*
((1/(a*x^2) - 1)^(1/2) - 1i)^3)/((1/(a*x^2) + 1)^(1/2) - 1)^3) + ((1/(a*x^2
) - 1)^(1/2) - 1i)/(8*a*((1/(a*x^2) + 1)^(1/2) - 1))
```

3.52 $\int e^{\operatorname{sech}^{-1}(ax^2)} dx$

Optimal result	360
Rubi [A] (verified)	360
Mathematica [C] (verified)	363
Maple [A] (verified)	363
Fricas [F]	364
Sympy [F]	364
Maxima [F]	364
Giac [F]	364
Mupad [F(-1)]	365

Optimal result

Integrand size = 8, antiderivative size = 147

$$\int e^{\operatorname{sech}^{-1}(ax^2)} dx = -\frac{2}{ax} + e^{\operatorname{sech}^{-1}(ax^2)} x - \frac{2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\sqrt{1-a^2x^4}}{ax} - \frac{2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}E(\arcsin(\sqrt{ax})|-1)}{\sqrt{a}} + \frac{2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\operatorname{EllipticF}(\arcsin(\sqrt{ax}),-1)}{\sqrt{a}}$$

[Out] $-2/a/x+(1/a/x^2+(1/a/x^2-1)^{(1/2)}*(1/a/x^2+1)^{(1/2)}*x-2*\operatorname{EllipticE}(x*a^{(1/2)},I)*(1/(a*x^2+1))^{(1/2)}*(a*x^2+1)^{(1/2)}/a^{(1/2)}+2*\operatorname{EllipticF}(x*a^{(1/2)},I)*(1/(a*x^2+1))^{(1/2)}*(a*x^2+1)^{(1/2)}/a^{(1/2)}-2*(1/(a*x^2+1))^{(1/2)}*(a*x^2+1)^{(1/2)}*(-a^2*x^4+1)^{(1/2)}/a/x$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {6465, 30, 265, 331, 313, 227, 1213, 435}

$$\int e^{\operatorname{sech}^{-1}(ax^2)} dx = -\frac{2\sqrt{\frac{1}{ax^2+1}}\sqrt{ax^2+1}\sqrt{1-a^2x^4}}{ax} + \frac{2\sqrt{\frac{1}{ax^2+1}}\sqrt{ax^2+1}\operatorname{EllipticF}(\arcsin(\sqrt{ax}),-1)}{\sqrt{a}} - \frac{2\sqrt{\frac{1}{ax^2+1}}\sqrt{ax^2+1}E(\arcsin(\sqrt{ax})|-1)}{\sqrt{a}} + xe^{\operatorname{sech}^{-1}(ax^2)} - \frac{2}{ax}$$

[In] Int[E^ArcSech[a*x^2], x]

[Out] $-2/(a*x) + E^{\text{ArcSech}[a*x^2]}*x - (2*\sqrt{(1 + a*x^2)^{-1}}*\sqrt{1 + a*x^2}*\sqrt{1 - a^2*x^4})/(a*x) - (2*\sqrt{(1 + a*x^2)^{-1}}*\sqrt{1 + a*x^2}*\text{EllipticE}[\text{ArcSin}[\sqrt{a}*x], -1])/\sqrt{a} + (2*\sqrt{(1 + a*x^2)^{-1}}*\sqrt{1 + a*x^2}*\text{EllipticF}[\text{ArcSin}[\sqrt{a}*x], -1])/\sqrt{a}$

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NegQ[m, -1]

Rule 227

Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := Simp[EllipticF[ArcSin[Rt[-b, 4]*(x/Rt[a, 4])], -1]/(Rt[a, 4]*Rt[-b, 4]), x] /; FreeQ[{a, b}, x] && NegQ[b/a] && GtQ[a, 0]

Rule 265

Int[((c_)*(x_))^(m_)*((a1_) + (b1_)*(x_)^(n_))^(p_)*((a2_) + (b2_)*(x_)^(n_))^(p_), x_Symbol] := Int[(c*x)^m*(a1*a2 + b1*b2*x^(2*n))^p, x] /; FreeQ[{a1, b1, a2, b2, c, m, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))

Rule 313

Int[(x_)^2/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[-b/a, 2]}, Dist[-q^(-1), Int[1/Sqrt[a + b*x^4], x], x] + Dist[1/q, Int[(1 + q*x^2)/Sqrt[a + b*x^4], x], x] /; FreeQ[{a, b}, x] && NegQ[b/a]

Rule 331

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Dist[b*((m + n*(p + 1) + 1)/(a*c^n*(m + 1))), Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 435

Int[Sqrt[(a_) + (b_)*(x_)^2]/Sqrt[(c_) + (d_)*(x_)^2], x_Symbol] := Simp[(Sqrt[a]/(Sqrt[c]*Rt[-d/c, 2]))*EllipticE[ArcSin[Rt[-d/c, 2]*x], b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a, 0]

Rule 1213

Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] := Dist[d/Sqrt[a], Int[Sqrt[1 + e*(x^2/d)]/Sqrt[1 - e*(x^2/d)], x], x] /; FreeQ[{a, c,

d, e}, x] && NegQ[c/a] && EqQ[c*d^2 + a*e^2, 0] && GtQ[a, 0]

Rule 6465

Int[E^ArcSech[(a_.)*(x_)^(p_)], x_Symbol] := Simp[x*E^ArcSech[a*x^p], x] + (Dist[p/a, Int[1/x^p, x], x] + Dist[p*(Sqrt[1 + a*x^p]/a)*Sqrt[1/(1 + a*x^p)]], Int[1/(x^p*Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x) /; FreeQ[{a, p}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= e^{\operatorname{sech}^{-1}(ax^2)} x + \frac{2 \int \frac{1}{x^2} dx}{a} + \frac{\left(2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{1}{x^2\sqrt{1-ax^2}\sqrt{1+ax^2}} dx}{a} \\
 &= -\frac{2}{ax} + e^{\operatorname{sech}^{-1}(ax^2)} x + \frac{\left(2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{1}{x^2\sqrt{1-a^2x^4}} dx}{a} \\
 &= -\frac{2}{ax} + e^{\operatorname{sech}^{-1}(ax^2)} x - \frac{2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\sqrt{1-a^2x^4}}{ax} \\
 &\quad - \left(2a\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{x^2}{\sqrt{1-a^2x^4}} dx \\
 &= -\frac{2}{ax} + e^{\operatorname{sech}^{-1}(ax^2)} x - \frac{2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\sqrt{1-a^2x^4}}{ax} \\
 &\quad + \left(2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{1}{\sqrt{1-a^2x^4}} dx \\
 &\quad - \left(2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{1+ax^2}{\sqrt{1-a^2x^4}} dx \\
 &= -\frac{2}{ax} + e^{\operatorname{sech}^{-1}(ax^2)} x - \frac{2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\sqrt{1-a^2x^4}}{ax} \\
 &\quad + \frac{2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2} \operatorname{EllipticF}(\arcsin(\sqrt{ax}), -1)}{\sqrt{a}} \\
 &\quad - \left(2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{\sqrt{1+ax^2}}{\sqrt{1-ax^2}} dx
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{2}{ax} + e^{\operatorname{sech}^{-1}(ax^2)} x - \frac{2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\sqrt{1-a^2x^4}}{ax} \\
&\quad - \frac{2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}E(\arcsin(\sqrt{ax})|-1)}{\sqrt{a}} \\
&\quad + \frac{2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\operatorname{EllipticF}(\arcsin(\sqrt{ax}),-1)}{\sqrt{a}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.92

$$\begin{aligned}
&\int e^{\operatorname{sech}^{-1}(ax^2)} dx \\
&= -\frac{1}{ax} + \left(-\frac{1}{ax} - x\right) \sqrt{\frac{1-ax^2}{1+ax^2}} \\
&\quad - \frac{2i\sqrt{\frac{1-ax^2}{1+ax^2}}\sqrt{1-a^2x^4}(E(i\operatorname{arcsinh}(\sqrt{-ax})|-1) - \operatorname{EllipticF}(i\operatorname{arcsinh}(\sqrt{-ax}),-1))}{\sqrt{-a}(-1+ax^2)}
\end{aligned}$$

[In] Integrate[E^ArcSech[a*x^2],x]

[Out] $-(1/(a*x)) + (-1/(a*x) - x)*\operatorname{Sqrt}[(1 - a*x^2)/(1 + a*x^2)] - ((2*I)*\operatorname{Sqrt}[(1 - a*x^2)/(1 + a*x^2)]*\operatorname{Sqrt}[1 - a^2*x^4]*(\operatorname{EllipticE}[I*\operatorname{ArcSinh}[\operatorname{Sqrt}[-a]*x], -1) - \operatorname{EllipticF}[I*\operatorname{ArcSinh}[\operatorname{Sqrt}[-a]*x], -1]))/(\operatorname{Sqrt}[-a]*(-1 + a*x^2))$

Maple [A] (verified)

Time = 0.77 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.90

method	result
default	$-\frac{1}{ax} - \frac{\sqrt{-\frac{ax^2-1}{ax^2}}x\sqrt{\frac{ax^2+1}{ax^2}}(x^4a^2+2\sqrt{-ax^2+1}\sqrt{ax^2+1}x\operatorname{EllipticF}(x\sqrt{a},i)\sqrt{a}-2\sqrt{-ax^2+1}\sqrt{ax^2+1}x\operatorname{EllipticE}(x\sqrt{a},i)\sqrt{a}}{x^4a^2-1}$

[In] `int(1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $-1/a/x - (-a*x^2-1)/a/x^2)^{(1/2)}*x*((a*x^2+1)/a/x^2)^{(1/2)}*(x^4*a^2+2*(-a*x^2+1)^{(1/2)}*(a*x^2+1)^{(1/2)}*x*\operatorname{EllipticF}(x*a^{(1/2)},I)*a^{(1/2)}-2*(-a*x^2+1)^{(1/2)}*(a*x^2+1)^{(1/2)}*x*\operatorname{EllipticE}(x*a^{(1/2)},I)*a^{(1/2)}-1)/(a^2*x^4-1)$

Fricas [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} dx = \int \sqrt{\frac{1}{ax^2} + 1} \sqrt{\frac{1}{ax^2} - 1} + \frac{1}{ax^2} dx$$

[In] integrate(1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2),x, algorithm="fricas")

[Out] integral((a*x^2*sqrt((a*x^2 + 1)/(a*x^2))*sqrt(-(a*x^2 - 1)/(a*x^2)) + 1)/(a*x^2), x)

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} dx = \frac{\int \frac{1}{x^2} dx + \int a \sqrt{-1 + \frac{1}{ax^2}} \sqrt{1 + \frac{1}{ax^2}} dx}{a}$$

[In] integrate(1/a/x**2+(1/a/x**2-1)**(1/2)*(1/a/x**2+1)**(1/2),x)

[Out] (Integral(x**(-2), x) + Integral(a*sqrt(-1 + 1/(a*x**2))*sqrt(1 + 1/(a*x**2)), x))/a

Maxima [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} dx = \int \sqrt{\frac{1}{ax^2} + 1} \sqrt{\frac{1}{ax^2} - 1} + \frac{1}{ax^2} dx$$

[In] integrate(1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(a*x^2 + 1)*sqrt(-a*x^2 + 1)/x^2, x)/a - 1/(a*x)

Giac [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} dx = \int \sqrt{\frac{1}{ax^2} + 1} \sqrt{\frac{1}{ax^2} - 1} + \frac{1}{ax^2} dx$$

[In] integrate(1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(1/(a*x^2) + 1)*sqrt(1/(a*x^2) - 1) + 1/(a*x^2), x)

Mupad [F(-1)]

Timed out.

$$\int e^{\operatorname{sech}^{-1}(ax^2)} dx = \int \sqrt{\frac{1}{ax^2} - 1} \sqrt{\frac{1}{ax^2} + 1} + \frac{1}{ax^2} dx$$

```
[In] int((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2), x)
```

```
[Out] int((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2), x)
```

3.53 $\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x} dx$

Optimal result	366
Rubi [A] (warning: unable to verify)	366
Mathematica [A] (verified)	368
Maple [A] (verified)	368
Fricas [B] (verification not implemented)	368
Sympy [F]	369
Maxima [F]	369
Giac [B] (verification not implemented)	369
Mupad [B] (verification not implemented)	370

Optimal result

Integrand size = 12, antiderivative size = 80

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x} dx = -\frac{1}{2ax^2} - \frac{\sqrt{1-ax^2}}{2ax^2\sqrt{\frac{1}{1+ax^2}}} - \frac{1}{2}\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\arcsin(ax^2)$$

[Out] $-1/2/a/x^2-1/2*(-a*x^2+1)^{(1/2)}/a/x^2/(1/(a*x^2+1))^{(1/2)}-1/2*\arcsin(a*x^2)*$
 $(1/(a*x^2+1))^{(1/2)}*(a*x^2+1)^{(1/2)}$

Rubi [A] (warning: unable to verify)

Time = 0.04 (sec), antiderivative size = 93, normalized size of antiderivative = 1.16, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {6469, 265, 281, 283, 222}

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x} dx = -\frac{\sqrt{\frac{1}{ax^2+1}}\sqrt{ax^2+1}\sqrt{1-a^2x^4}}{2ax^2} - \frac{1}{2}\sqrt{\frac{1}{ax^2+1}}\sqrt{ax^2+1}\arcsin(ax^2) - \frac{1}{2ax^2}$$

[In] `Int[E^ArcSech[a*x^2]/x,x]`

[Out] $-1/2*1/(a*x^2) - (\operatorname{Sqrt}[(1+a*x^2)^{-1}]*\operatorname{Sqrt}[1+a*x^2]*\operatorname{Sqrt}[1-a^2*x^4])$
 $/ (2*a*x^2) - (\operatorname{Sqrt}[(1+a*x^2)^{-1}]*\operatorname{Sqrt}[1+a*x^2]*\operatorname{ArcSin}[a*x^2])/2$

Rule 222

`Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

Rule 265

```
Int[((c_.)*(x_))^(m_.)*((a1_) + (b1_.)*(x_)^(n_))^(p_)*((a2_) + (b2_.)*(x_)
^(n_))^(p_), x_Symbol] := Int[(c*x)^(m*(a1*a2 + b1*b2*x^(2*n)))^p, x] /; Free
Q[{a1, b1, a2, b2, c, m, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p]
|| (GtQ[a1, 0] && GtQ[a2, 0]))
```

Rule 281

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m
+ 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x
^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

Rule 283

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*
x)^(m + 1)*((a + b*x^n)^p/(c*(m + 1))), x] - Dist[b*n*(p/(c^n*(m + 1))), In
t[(c*x)^(m + n)*(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[
n, 0] && GtQ[p, 0] && LtQ[m, -1] && !ILtQ[(m + n*p + n + 1)/n, 0] && IntBi
nomialQ[a, b, c, n, m, p, x]
```

Rule 6469

```
Int[E^ArcSech[(a_.)*(x_)^(p_.)]/(x_), x_Symbol] := -Simp[(a*p*x^p)^(-1), x]
+ Dist[(Sqrt[1 + a*x^p]/a)*Sqrt[1/(1 + a*x^p)], Int[Sqrt[1 + a*x^p]*(Sqrt[
1 - a*x^p]/x^(p + 1)), x], x] /; FreeQ[{a, p}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{1}{2ax^2} + \frac{\left(\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{\sqrt{1-ax^2}\sqrt{1+ax^2}}{x^3} dx}{a} \\
&= -\frac{1}{2ax^2} + \frac{\left(\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{\sqrt{1-a^2x^4}}{x^3} dx}{a} \\
&= -\frac{1}{2ax^2} + \frac{\left(\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \text{Subst}\left(\int \frac{\sqrt{1-a^2x^2}}{x^2} dx, x, x^2\right)}{2a} \\
&= -\frac{1}{2ax^2} - \frac{\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\sqrt{1-a^2x^4}}{2ax^2} \\
&\quad - \frac{1}{2} \left(a\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2} \right) \text{Subst}\left(\int \frac{1}{\sqrt{1-a^2x^2}} dx, x, x^2\right) \\
&= -\frac{1}{2ax^2} - \frac{\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\sqrt{1-a^2x^4}}{2ax^2} - \frac{1}{2} \sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2} \arcsin(ax^2)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.28

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x} dx = -\frac{1}{2}e^{\operatorname{sech}^{-1}(ax^2)} + \arctan\left(e^{\operatorname{sech}^{-1}(ax^2)}\right)$$

[In] Integrate[E^ArcSech[a*x^2]/x,x]

[Out] -1/2*E^ArcSech[a*x^2] + ArcTan[E^ArcSech[a*x^2]]

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.29

method	result	size
default	$-\frac{\sqrt{-\frac{ax^2-1}{ax^2}} \sqrt{\frac{ax^2+1}{ax^2}} \left(\arctan\left(\frac{x^2}{\sqrt{-\frac{x^4a^2-1}{a^2}}}\right) x^2 + \sqrt{-\frac{x^4a^2-1}{a^2}} \right)}{2\sqrt{-\frac{x^4a^2-1}{a^2}}} - \frac{1}{2ax^2}$	103

[In] int((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))/x,x,method=_RETURNVERBOSE)

[Out] -1/2*(-(a*x^2-1)/a/x^2)^(1/2)*((a*x^2+1)/a/x^2)^(1/2)*(arctan(x^2/(-(a^2*x^4-1)/a^2)^(1/2))*x^2+(-(a^2*x^4-1)/a^2)^(1/2))/(-(a^2*x^4-1)/a^2)^(1/2)-1/2/a/x^2

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 102 vs. 2(44) = 88.

Time = 0.25 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.28

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x} dx = -\frac{ax^2 \sqrt{\frac{ax^2+1}{ax^2}} \sqrt{-\frac{ax^2-1}{ax^2}} - 2ax^2 \arctan\left(\frac{ax^2 \sqrt{\frac{ax^2+1}{ax^2}} \sqrt{-\frac{ax^2-1}{ax^2}} - 1}{ax^2}\right) + 1}{2ax^2}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))/x,x, algorithm="fricas")

[Out] -1/2*(a*x^2*sqrt((a*x^2 + 1)/(a*x^2))*sqrt(-(a*x^2 - 1)/(a*x^2)) - 2*a*x^2*arctan((a*x^2*sqrt((a*x^2 + 1)/(a*x^2))*sqrt(-(a*x^2 - 1)/(a*x^2)) - 1)/(a*x^2)) + 1)/(a*x^2)

SymPy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x} dx = \int \frac{1}{x^3} dx + \int \frac{a\sqrt{-1+\frac{1}{ax^2}}\sqrt{1+\frac{1}{ax^2}}}{a} dx$$

[In] integrate((1/a/x**2+(1/a/x**2-1)**(1/2)*(1/a/x**2+1)**(1/2))/x,x)

[Out] (Integral(x**(-3), x) + Integral(a*sqrt(-1 + 1/(a*x**2))*sqrt(1 + 1/(a*x**2))/x, x))/a

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x} dx = \int \frac{\sqrt{\frac{1}{ax^2} + 1}\sqrt{\frac{1}{ax^2} - 1 + \frac{1}{ax^2}}}{x} dx$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))/x,x, algorithm="maxima")

[Out] integrate(sqrt(a*x^2 + 1)*sqrt(-a*x^2 + 1)/x^3, x)/a - 1/2/(a*x^2)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 252 vs. 2(44) = 88.

Time = 1.73 (sec) , antiderivative size = 252, normalized size of antiderivative = 3.15

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x} dx = \frac{\left(\pi + 2 \arctan \left(\frac{\sqrt{a^2x^2+a} \left(\frac{(\sqrt{2}\sqrt{a}-\sqrt{-a^2x^2+a})^2}{a^2x^2+a} - 1 \right)}{2(\sqrt{2}\sqrt{a}-\sqrt{-a^2x^2+a})} \right) \right) a^3 + \frac{4a^3 \left(\frac{\sqrt{2}\sqrt{a}-\sqrt{-a^2x^2+a}}{\sqrt{a^2x^2+a}} - \frac{\sqrt{a^2x^2+a}}{\sqrt{2}\sqrt{a}-\sqrt{-a^2x^2+a}} \right)}{\left(\frac{\sqrt{2}\sqrt{a}-\sqrt{-a^2x^2+a}}{\sqrt{a^2x^2+a}} - \frac{\sqrt{a^2x^2+a}}{\sqrt{2}\sqrt{a}-\sqrt{-a^2x^2+a}} \right)^2} + \frac{a^2}{x^2}}{2a^3}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))/x,x, algorithm="giac")

[Out] -1/2*((pi + 2*arctan(1/2*sqrt(a^2*x^2 + a))*((sqrt(2)*sqrt(a) - sqrt(-a^2*x^2 + a))^2/(a^2*x^2 + a) - 1)/(sqrt(2)*sqrt(a) - sqrt(-a^2*x^2 + a))))*a^3 + 4*a^3*((sqrt(2)*sqrt(a) - sqrt(-a^2*x^2 + a))/sqrt(a^2*x^2 + a) - sqrt(a^2*x^2 + a)/(sqrt(2)*sqrt(a) - sqrt(-a^2*x^2 + a)))/(((sqrt(2)*sqrt(a) - sqrt(-a^2*x^2 + a))/sqrt(a^2*x^2 + a) - sqrt(a^2*x^2 + a)/(sqrt(2)*sqrt(a) - sqrt(-a^2*x^2 + a)))^2 - 4) + a^2/x^2)/a^3

Mupad [B] (verification not implemented)

Time = 7.40 (sec) , antiderivative size = 185, normalized size of antiderivative = 2.31

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x} dx = -\frac{\ln\left(\frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^2}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^2} + 1\right) \operatorname{li}}{2} + \frac{\ln\left(\frac{\sqrt{\frac{1}{ax^2}-1-i}}{\sqrt{\frac{1}{ax^2}+1-1}}\right) \operatorname{li}}{2} - \frac{1}{2ax^2}$$

$$+ \frac{\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^2 8i}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^2 \left(2 + \frac{2\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^4}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^4} - \frac{4\left(\sqrt{\frac{1}{ax^2}-1-i}\right)^2}{\left(\sqrt{\frac{1}{ax^2}+1-1}\right)^2}\right)}$$

[In] int(((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2))/x,x)

[Out] (log(((1/(a*x^2) - 1)^(1/2) - 1i)/((1/(a*x^2) + 1)^(1/2) - 1))*1i)/2 - (log(((1/(a*x^2) - 1)^(1/2) - 1i)^2/((1/(a*x^2) + 1)^(1/2) - 1)^2 + 1)*1i)/2 - 1/(2*a*x^2) + (((1/(a*x^2) - 1)^(1/2) - 1i)^2*8i)/(((1/(a*x^2) + 1)^(1/2) - 1)^2*((2*((1/(a*x^2) - 1)^(1/2) - 1i)^4)/((1/(a*x^2) + 1)^(1/2) - 1)^4 - (4*((1/(a*x^2) - 1)^(1/2) - 1i)^2)/((1/(a*x^2) + 1)^(1/2) - 1)^2 + 2)))

3.54 $\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^2} dx$

Optimal result	371
Rubi [A] (verified)	371
Mathematica [C] (verified)	373
Maple [A] (verified)	373
Fricas [A] (verification not implemented)	374
Sympy [F]	374
Maxima [F]	374
Giac [F]	375
Mupad [F(-1)]	375

Optimal result

Integrand size = 12, antiderivative size = 115

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^2} dx = \frac{2}{3ax^3} - \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x} + \frac{2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\sqrt{1-a^2x^4}}{3ax^3} - \frac{2}{3}\sqrt{a}\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\operatorname{EllipticF}(\arcsin(\sqrt{ax}), -1)$$

[Out] $2/3/a/x^3 - (1/a/x^2 + (1/a/x^2 - 1)^{(1/2)} * (1/a/x^2 + 1)^{(1/2)})/x - 2/3 * \operatorname{EllipticF}(x * a^{(1/2)}, I) * a^{(1/2)} * (1/(a * x^2 + 1))^{(1/2)} * (a * x^2 + 1)^{(1/2)} + 2/3 * (1/(a * x^2 + 1))^{(1/2)} * (a * x^2 + 1)^{(1/2)} * (-a^2 * x^4 + 1)^{(1/2)} / a / x^3$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {6470, 30, 265, 331, 227}

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^2} dx = \frac{2\sqrt{\frac{1}{ax^2+1}}\sqrt{ax^2+1}\sqrt{1-a^2x^4}}{3ax^3} - \frac{2}{3}\sqrt{a}\sqrt{\frac{1}{ax^2+1}}\sqrt{ax^2+1}\operatorname{EllipticF}(\arcsin(\sqrt{ax}), -1) + \frac{2}{3ax^3} - \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcSech}[a*x^2]}/x^2, x]$

[Out] $2/(3*a*x^3) - E^{\text{ArcSech}[a*x^2]}/x + (2*\text{Sqrt}[(1 + a*x^2)^{-1}]*\text{Sqrt}[1 + a*x^2]*\text{Sqrt}[1 - a^2*x^4])/(3*a*x^3) - (2*\text{Sqrt}[a]*\text{Sqrt}[(1 + a*x^2)^{-1}]*\text{Sqrt}[1 + a*x^2]*\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[a]*x], -1])/3$

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}/(m + 1), x] \text{ /; FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 227

$\text{Int}[1/\text{Sqrt}[(a_) + (b_.)*(x_)^4], x_Symbol] \rightarrow \text{Simp}[\text{EllipticF}[\text{ArcSin}[\text{Rt}[-b, 4]*(x/\text{Rt}[a, 4])], -1]/(\text{Rt}[a, 4]*\text{Rt}[-b, 4]), x] \text{ /; FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[b/a] \ \&\& \ \text{GtQ}[a, 0]$

Rule 265

$\text{Int}[((c_.)*(x_))^{(m_.)}*((a1_) + (b1_.)*(x_)^{(n_)})^{(p_)}*((a2_) + (b2_.)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Int}[(c*x)^m*(a1*a2 + b1*b2*x^{(2*n)})^p, x] \text{ /; FreeQ}[\{a1, b1, a2, b2, c, m, n, p\}, x] \ \&\& \ \text{EqQ}[a2*b1 + a1*b2, 0] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{GtQ}[a1, 0] \ \&\& \ \text{GtQ}[a2, 0]))$

Rule 331

$\text{Int}[((c_.)*(x_))^{(m_.)}*((a_) + (b_.)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(c*x)^{(m + 1)}*((a + b*x^n)^{(p + 1)}/(a*c*(m + 1))), x] - \text{Dist}[b*((m + n*(p + 1) + 1)/(a*c^n*(m + 1))), \text{Int}[(c*x)^{(m + n)}*(a + b*x^n)^p, x], x] \text{ /; FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 6470

$\text{Int}[E^{\text{ArcSech}[(a_.)*(x_)^{(p_.)}]}*(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}*(E^{\text{ArcSech}[a*x^p]/(m + 1)}), x] + (\text{Dist}[p/(a*(m + 1)), \text{Int}[x^{(m - p)}, x], x] + \text{Dist}[p*(\text{Sqrt}[1 + a*x^p]/(a*(m + 1)))*\text{Sqrt}[1/(1 + a*x^p)], \text{Int}[x^{(m - p)}/(\text{Sqrt}[1 + a*x^p]*\text{Sqrt}[1 - a*x^p]), x], x]) \text{ /; FreeQ}[\{a, m, p\}, x] \ \&\& \ \text{NeQ}[m, -1]$

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{e^{\text{sech}^{-1}(ax^2)}}{x} - \frac{2 \int \frac{1}{x^4} dx}{a} - \frac{\left(2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{1}{x^4\sqrt{1-ax^2}\sqrt{1+ax^2}} dx}{a} \\ &= \frac{2}{3ax^3} - \frac{e^{\text{sech}^{-1}(ax^2)}}{x} - \frac{\left(2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{1}{x^4\sqrt{1-a^2x^4}} dx}{a} \end{aligned}$$

$$\begin{aligned}
&= \frac{2}{3ax^3} - \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x} + \frac{2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\sqrt{1-a^2x^4}}{3ax^3} \\
&\quad - \frac{1}{3} \left(2a\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2} \right) \int \frac{1}{\sqrt{1-a^2x^4}} dx \\
&= \frac{2}{3ax^3} - \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x} + \frac{2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\sqrt{1-a^2x^4}}{3ax^3} \\
&\quad - \frac{2}{3}\sqrt{a}\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2} \operatorname{EllipticF}(\arcsin(\sqrt{ax}), -1)
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.23 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.88

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^2} dx = \frac{a\sqrt{1+e^{2\operatorname{sech}^{-1}(ax^2)}}\sqrt{\frac{e^{\operatorname{sech}^{-1}(ax^2)}}{2+2e^{2\operatorname{sech}^{-1}(ax^2)}}}x\left(\sqrt{1+e^{2\operatorname{sech}^{-1}(ax^2)}}-4\operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -e^{2\operatorname{sech}^{-1}(ax^2)}\right)\right)}{3\sqrt{ax^2}}$$

[In] Integrate[E^ArcSech[a*x^2]/x^2,x]

[Out] $-1/3*(a*\sqrt{1+E^{(2*ArcSech[a*x^2])}})*\sqrt{E^{ArcSech[a*x^2]}/(2+2*E^{(2*ArcSech[a*x^2])})}*x*(\sqrt{1+E^{(2*ArcSech[a*x^2])}})-4*\operatorname{Hypergeometric2F1}[1/4, 1/2, 5/4, -E^{(2*ArcSech[a*x^2])}])/ \sqrt{a*x^2}$

Maple [A] (verified)

Time = 0.90 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.90

method	result	size
default	$\frac{\sqrt{-\frac{ax^2-1}{ax^2}}\sqrt{\frac{ax^2+1}{ax^2}}\left(2\sqrt{-ax^2+1}\sqrt{ax^2+1}\operatorname{EllipticF}(x\sqrt{a}, i)x^3a^{\frac{3}{2}}-x^4a^2+1\right)}{3x(x^4a^2-1)} - \frac{1}{3ax^3}$	104

[In] int((1/a/x^2+(1/a/x^2-1)^(1/2))*(1/a/x^2+1)^(1/2))/x^2,x,method=_RETURNVERBOSE)

[Out] $1/3*(-(a*x^2-1)/a/x^2)^(1/2)/x*((a*x^2+1)/a/x^2)^(1/2)*(2*(-a*x^2+1)^(1/2)*(a*x^2+1)^(1/2)*\operatorname{EllipticF}(x*a^(1/2), I)*x^3*a^(3/2)-x^4*a^2+1)/(a^2*x^4-1)-1/3/a/x^3$

Fricas [A] (verification not implemented)

none

Time = 0.08 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.56

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^2} dx = -\frac{2 a^{\frac{3}{2}} x^3 F(\arcsin(\sqrt{ax}) \mid -1) + ax^2 \sqrt{\frac{ax^2+1}{ax^2}} \sqrt{-\frac{ax^2-1}{ax^2}} + 1}{3 ax^3}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))/x^2,x, algorithm="fricas")

[Out] -1/3*(2*a^(3/2)*x^3*elliptic_f(arcsin(sqrt(a)*x), -1) + a*x^2*sqrt((a*x^2 + 1)/(a*x^2))*sqrt(-(a*x^2 - 1)/(a*x^2)) + 1)/(a*x^3)

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^2} dx = \int \frac{1}{x^4} dx + \int \frac{a \sqrt{-1 + \frac{1}{ax^2}} \sqrt{1 + \frac{1}{ax^2}}}{x^2} dx$$

[In] integrate((1/a/x**2+(1/a/x**2-1)**(1/2)*(1/a/x**2+1)**(1/2))/x**2,x)

[Out] (Integral(x**(-4), x) + Integral(a*sqrt(-1 + 1/(a*x**2))*sqrt(1 + 1/(a*x**2))/x**2, x))/a

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^2} dx = \int \frac{\sqrt{\frac{1}{ax^2} + 1} \sqrt{\frac{1}{ax^2} - 1 + \frac{1}{ax^2}}}{x^2} dx$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))/x^2,x, algorithm="maxima")

[Out] integrate(sqrt(a*x^2 + 1)*sqrt(-a*x^2 + 1)/x^4, x)/a - 1/3/(a*x^3)

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^2} dx = \int \frac{\sqrt{\frac{1}{ax^2} + 1} \sqrt{\frac{1}{ax^2} - 1} + \frac{1}{ax^2}}{x^2} dx$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))/x^2,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x^2) + 1)*sqrt(1/(a*x^2) - 1) + 1/(a*x^2))/x^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^2} dx = \int \frac{\sqrt{\frac{1}{ax^2} - 1} \sqrt{\frac{1}{ax^2} + 1} + \frac{1}{ax^2}}{x^2} dx$$

[In] int(((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2))/x^2,x)

[Out] int(((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2))/x^2, x)

3.55 $\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^3} dx$

Optimal result	376
Rubi [A] (verified)	376
Mathematica [A] (verified)	378
Maple [C] (verified)	379
Fricas [A] (verification not implemented)	379
Sympy [A] (verification not implemented)	380
Maxima [F]	380
Giac [F(-2)]	380
Mupad [B] (verification not implemented)	381

Optimal result

Integrand size = 12, antiderivative size = 118

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^3} dx = \frac{1}{4ax^4} - \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{2x^2} + \frac{\sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \sqrt{1-a^2x^4}}{4ax^4} + \frac{1}{4}a \sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \operatorname{arctanh}(\sqrt{1-a^2x^4})$$

[Out] 1/4/a/x^4-1/2*(1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))/x^2+1/4*a*arctanh((-a^2*x^4+1)^(1/2))*(1/(a*x^2+1))^(1/2)*(a*x^2+1)^(1/2)+1/4*(1/(a*x^2+1))^(1/2)*(a*x^2+1)^(1/2)*(-a^2*x^4+1)^(1/2)/a/x^4

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {6470, 30, 265, 272, 44, 65, 214}

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^3} dx = \frac{1}{4}a \sqrt{\frac{1}{ax^2+1}} \sqrt{ax^2+1} \operatorname{arctanh}(\sqrt{1-a^2x^4}) + \frac{\sqrt{\frac{1}{ax^2+1}} \sqrt{ax^2+1} \sqrt{1-a^2x^4}}{4ax^4} + \frac{1}{4ax^4} - \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{2x^2}$$

[In] Int[E^ArcSech[a*x^2]/x^3,x]

[Out] 1/(4*a*x^4) - E^ArcSech[a*x^2]/(2*x^2) + (Sqrt[(1 + a*x^2)^(-1)]*Sqrt[1 + a*x^2]*Sqrt[1 - a^2*x^4])/(4*a*x^4) + (a*Sqrt[(1 + a*x^2)^(-1)]*Sqrt[1 + a*x^2]*ArcTanh[Sqrt[1 - a^2*x^4]])/4

Rule 30

$\text{Int}[(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}/(m+1), x] /; \text{FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 44

$\text{Int}[(a_ + (b_)(x_))^{(m_)}((c_ + (d_)(x_))^{(n_)}), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m+1)}((c + d*x)^{(n+1)})/((b*c - a*d)*(m+1)), x] - \text{Dist}[d*((m+n+2)/((b*c - a*d)*(m+1))), \text{Int}[(a + b*x)^{(m+1)}(c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{ILtQ}[m, -1] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ \text{LtQ}[n, 0]$

Rule 65

$\text{Int}[(a_ + (b_)(x_))^{(m_)}((c_ + (d_)(x_))^{(n_)}), x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m+1)-1)}(c - a*(d/b) + d*(x^{p/b})^n, x], x, (a + b*x)^{(1/p)}], x]] /; \text{FreeQ}\{a, b, c, d\}, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 214

$\text{Int}[(a_ + (b_)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x\} \ \&\& \ \text{NegQ}[a/b]$

Rule 265

$\text{Int}[(c_)(x_)^{(m_)}((a1_ + (b1_)(x_)^{(n_)})^{(p_)}((a2_ + (b2_)(x_)^{(n_)})^{(p_)}), x_Symbol] \rightarrow \text{Int}[(c*x)^m*(a1*a2 + b1*b2*x^{(2*n)})^p, x] /; \text{FreeQ}\{a1, b1, a2, b2, c, m, n, p\}, x\} \ \&\& \ \text{EqQ}[a2*b1 + a1*b2, 0] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{GtQ}[a1, 0] \ \&\& \ \text{GtQ}[a2, 0]))$

Rule 272

$\text{Int}[(x_)^{(m_)}((a_ + (b_)(x_)^{(n_)})^{(p_)}), x_Symbol] \rightarrow \text{Dist}[1/n, \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n] - 1)}(a + b*x)^p, x], x, x^n], x] /; \text{FreeQ}\{a, b, m, n, p\}, x\} \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$

Rule 6470

$\text{Int}[E^{\text{ArcSech}[a*(x_)^{(p_)}]}(x_)^{(m_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}(E^{\text{ArcSech}[a*x^p]/(m+1)}, x] + (\text{Dist}[p/(a*(m+1)), \text{Int}[x^{(m-p)}, x], x] + \text{Dist}[p*(\text{Sqrt}[1 + a*x^p]/(a*(m+1)))*\text{Sqrt}[1/(1 + a*x^p)], \text{Int}[x^{(m-p)}/(\text{Sqrt}[1 + a*x^p]*\text{Sqrt}[1 - a*x^p]), x], x]) /; \text{FreeQ}\{a, m, p\}, x\} \ \&\& \ \text{NeQ}[m, -1]$

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{e^{\operatorname{sech}^{-1}(ax^2)}}{2x^2} - \frac{\int \frac{1}{x^5} dx}{a} - \frac{\left(\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{1}{x^5\sqrt{1-ax^2}\sqrt{1+ax^2}} dx}{a} \\
&= \frac{1}{4ax^4} - \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{2x^2} - \frac{\left(\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{1}{x^5\sqrt{1-a^2x^4}} dx}{a} \\
&= \frac{1}{4ax^4} - \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{2x^2} - \frac{\left(\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \operatorname{Subst}\left(\int \frac{1}{x^2\sqrt{1-a^2x}} dx, x, x^4\right)}{4a} \\
&= \frac{1}{4ax^4} - \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{2x^2} + \frac{\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\sqrt{1-a^2x^4}}{4ax^4} \\
&\quad - \frac{1}{8} \left(a\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2} \right) \operatorname{Subst}\left(\int \frac{1}{x\sqrt{1-a^2x}} dx, x, x^4\right) \\
&= \frac{1}{4ax^4} - \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{2x^2} + \frac{\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\sqrt{1-a^2x^4}}{4ax^4} \\
&\quad + \frac{\left(\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \operatorname{Subst}\left(\int \frac{1}{\frac{1}{a^2} - \frac{x^2}{a^2}} dx, x, \sqrt{1-a^2x^4}\right)}{4a} \\
&= \frac{1}{4ax^4} - \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{2x^2} + \frac{\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\sqrt{1-a^2x^4}}{4ax^4} \\
&\quad + \frac{1}{4} a \sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \operatorname{arctanh}\left(\sqrt{1-a^2x^4}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.89

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^3} dx = -\frac{1}{x^4} + \frac{\sqrt{\frac{1-ax^2}{1+ax^2}}(1+ax^2)}{x^4} - \frac{a^2 \sqrt{\frac{1-ax^2}{1+ax^2}}(1+ax^2) \operatorname{arctan}\left(\sqrt{-1+a^2x^4}\right)}{4a\sqrt{-1+a^2x^4}}$$

[In] Integrate[E^ArcSech[a*x^2]/x^3,x]

[Out] -1/4*(x^(-4) + (Sqrt[(1 - a*x^2)/(1 + a*x^2)]*(1 + a*x^2))/x^4 - (a^2*Sqrt[(1 - a*x^2)/(1 + a*x^2)]*(1 + a*x^2)*ArcTan[Sqrt[-1 + a^2*x^4]]/Sqrt[-1 + a^2*x^4])/a

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.10 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.09

method	result	size
default	$\frac{\sqrt{-\frac{ax^2-1}{ax^2}} \sqrt{\frac{ax^2+1}{ax^2}} \left(\ln \left(\frac{2 \operatorname{csgn}\left(\frac{1}{a}\right) a \sqrt{-\frac{x^4 a^2-1}{a^2}+2}}{a^2 x^2} \right) x^4 a - \sqrt{-\frac{x^4 a^2-1}{a^2}} \operatorname{csgn}\left(\frac{1}{a}\right) \right) \operatorname{csgn}\left(\frac{1}{a}\right)}{4x^2 \sqrt{-\frac{x^4 a^2-1}{a^2}}} - \frac{1}{4ax^4}$	129

[In] int((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))/x^3,x,method=_RETURNVERBOSE)

[Out] 1/4*(-(a*x^2-1)/a/x^2)^(1/2)/x^2*((a*x^2+1)/a/x^2)^(1/2)*(ln(2*(csgn(1/a)*a*(-(a^2*x^4-1)/a^2)^(1/2)+1)/a^2/x^2)*x^4*a-(sqrt(-a^2*x^4-1)/a^2)^(1/2)*csgn(1/a))*csgn(1/a)/(-(a^2*x^4-1)/a^2)^(1/2)-1/4/a/x^4

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.24

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^3} dx$$

$$= \frac{a^2 x^4 \log \left(ax^2 \sqrt{\frac{ax^2+1}{ax^2}} \sqrt{-\frac{ax^2-1}{ax^2}} + 1 \right) - a^2 x^4 \log \left(ax^2 \sqrt{\frac{ax^2+1}{ax^2}} \sqrt{-\frac{ax^2-1}{ax^2}} - 1 \right) - 2 a x^2 \sqrt{\frac{ax^2+1}{ax^2}} \sqrt{-\frac{ax^2-1}{ax^2}}}{8 a x^4}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))/x^3,x, algorithm="fricas")

[Out] 1/8*(a^2*x^4*log(a*x^2*sqrt((a*x^2 + 1)/(a*x^2))*sqrt(-(a*x^2 - 1)/(a*x^2)) + 1) - a^2*x^4*log(a*x^2*sqrt((a*x^2 + 1)/(a*x^2))*sqrt(-(a*x^2 - 1)/(a*x^2)) - 1) - 2*a*x^2*sqrt((a*x^2 + 1)/(a*x^2))*sqrt(-(a*x^2 - 1)/(a*x^2)) - 2)/(a*x^4)

Sympy [A] (verification not implemented)

Time = 4.07 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.69

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^3} dx$$

$$= -\frac{a \left(2\sqrt{-1 + \frac{1}{ax^2}} \left(\frac{\left(1 + \frac{1}{ax^2}\right)^{\frac{3}{2}}}{4} - \frac{\sqrt{1 + \frac{1}{ax^2}}}{4} \right) - \log \left(2\sqrt{-1 + \frac{1}{ax^2}} + 2\sqrt{1 + \frac{1}{ax^2}} \right) \right)}{2} - \frac{1}{4ax^4}$$

[In] integrate((1/a/x**2+(1/a/x**2-1)**(1/2)*(1/a/x**2+1)**(1/2))/x**3,x)

[Out] -a*(2*sqrt(-1 + 1/(a*x**2))*((1 + 1/(a*x**2))**(3/2)/4 - sqrt(1 + 1/(a*x**2))/4) - log(2*sqrt(-1 + 1/(a*x**2)) + 2*sqrt(1 + 1/(a*x**2))))/2 - 1/(4*a*x**4)

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^3} dx = \int \frac{\sqrt{\frac{1}{ax^2} + 1} \sqrt{\frac{1}{ax^2} - 1 + \frac{1}{ax^2}}}{x^3} dx$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))/x^3,x, algorithm="maxima")

[Out] integrate(sqrt(a*x^2 + 1)*sqrt(-a*x^2 + 1)/x^5, x)/a - 1/4/(a*x^4)

Giac [F(-2)]

Exception generated.

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^3} dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))/x^3,x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Limit: Max order reached or unable to make series expansion Error: Bad Argument Value

Mupad [B] (verification not implemented)

Time = 5.19 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.60

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^2)}}{x^3} dx = \frac{a \ln \left(\sqrt{\frac{1}{ax^2} - 1} \sqrt{\frac{1}{ax^2} + 1} + \frac{1}{ax^2} \right)}{4} - \frac{1}{4ax^4} - \frac{\sqrt{\frac{1}{ax^2} - 1} \sqrt{\frac{1}{ax^2} + 1}}{4x^2}$$

[In] int(((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2))/x^3,x)

[Out] (a*log((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2)))/4 - 1/(4*a*x^4) - ((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2))/(4*x^2)

3.56 $\int e^{\operatorname{sech}^{-1}(ax^3)} x^m dx$

Optimal result	382
Rubi [A] (verified)	382
Mathematica [A] (verified)	384
Maple [F]	384
Fricas [F]	384
Sympy [F]	385
Maxima [F(-2)]	385
Giac [F]	385
Mupad [F(-1)]	386

Optimal result

Integrand size = 12, antiderivative size = 109

$$\int e^{\operatorname{sech}^{-1}(ax^3)} x^m dx = -\frac{3x^{-2+m}}{a(2+m-m^2)} + \frac{e^{\operatorname{sech}^{-1}(ax^3)} x^{1+m}}{1+m} - \frac{3x^{-2+m} \sqrt{\frac{1}{1+ax^3}} \sqrt{1+ax^3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(-2+m), \frac{4+m}{6}, a^2 x^6\right)}{a(2+m-m^2)}$$

[Out] $-3*x^{(-2+m)}/a/(-m^2+m+2)+(1/a/x^3+(1/a/x^3-1)^{(1/2)}*(1/a/x^3+1)^{(1/2)})*x^{(1+m)}/(1+m)-3*x^{(-2+m)}*hypergeom([1/2, -1/3+1/6*m], [2/3+1/6*m], a^2*x^6)*(1/(a*x^3+1))^{(1/2)}*(a*x^3+1)^{(1/2)}/a/(-m^2+m+2)$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6470, 30, 265, 371}

$$\int e^{\operatorname{sech}^{-1}(ax^3)} x^m dx = -\frac{3\sqrt{\frac{1}{ax^3+1}} \sqrt{ax^3+1} x^{m-2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m-2}{6}, \frac{m+4}{6}, a^2 x^6\right)}{a(-m^2+m+2)} - \frac{3x^{m-2}}{a(-m^2+m+2)} + \frac{x^{m+1} e^{\operatorname{sech}^{-1}(ax^3)}}{m+1}$$

[In] Int[E^ArcSech[a*x^3]*x^m,x]

[Out] $(-3*x^{(-2 + m)})/(a*(2 + m - m^2)) + (E^{\text{ArcSech}[a*x^3]}*x^{(1 + m)})/(1 + m) - (3*x^{(-2 + m)}*\text{Sqrt}[(1 + a*x^3)^{(-1)}]*\text{Sqrt}[1 + a*x^3]*\text{Hypergeometric2F1}[1/2, (-2 + m)/6, (4 + m)/6, a^2*x^6])/(a*(2 + m - m^2))$

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}/(m + 1), x] /; \text{FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 265

$\text{Int}[(c_.)*(x_)^{(m_.)}*((a1_.) + (b1_.)*(x_)^{(n_.)})^{(p_.)}*((a2_.) + (b2_.)*(x_)^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Int}[(c*x)^m*(a1*a2 + b1*b2*x^{(2*n)})^p, x] /; \text{FreeQ}\{a1, b1, a2, b2, c, m, n, p\}, x] \ \&\& \ \text{EqQ}[a2*b1 + a1*b2, 0] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{GtQ}[a1, 0] \ \&\& \ \text{GtQ}[a2, 0]))$

Rule 371

$\text{Int}[(c_.)*(x_)^{(m_.)}*((a_.) + (b_.)*(x_)^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m + 1)}/(c*(m + 1))) * \text{Hypergeometric2F1}[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}\{a, b, c, m, n, p\}, x] \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

Rule 6470

$\text{Int}[E^{\text{ArcSech}[a_.*(x_)^{(p_.)}]}*(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}*(E^{\text{ArcSech}[a*x^p]/(m + 1)}), x] + (\text{Dist}[p/(a*(m + 1)), \text{Int}[x^{(m - p)}, x], x] + \text{Dist}[p*(\text{Sqrt}[1 + a*x^p]/(a*(m + 1)))*\text{Sqrt}[1/(1 + a*x^p)], \text{Int}[x^{(m - p)}/(\text{Sqrt}[1 + a*x^p]*\text{Sqrt}[1 - a*x^p]), x], x]) /; \text{FreeQ}\{a, m, p\}, x] \ \&\& \ \text{NeQ}[m, -1]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{e^{\text{sech}^{-1}(ax^3)}x^{1+m}}{1+m} + \frac{3 \int x^{-3+m} dx}{a(1+m)} + \frac{\left(3\sqrt{\frac{1}{1+ax^3}}\sqrt{1+ax^3}\right) \int \frac{x^{-3+m}}{\sqrt{1-ax^3}\sqrt{1+ax^3}} dx}{a(1+m)} \\ &= -\frac{3x^{-2+m}}{a(2+m-m^2)} + \frac{e^{\text{sech}^{-1}(ax^3)}x^{1+m}}{1+m} + \frac{\left(3\sqrt{\frac{1}{1+ax^3}}\sqrt{1+ax^3}\right) \int \frac{x^{-3+m}}{\sqrt{1-a^2x^6}} dx}{a(1+m)} \\ &= -\frac{3x^{-2+m}}{a(2+m-m^2)} + \frac{e^{\text{sech}^{-1}(ax^3)}x^{1+m}}{1+m} \\ &\quad - \frac{3x^{-2+m}\sqrt{\frac{1}{1+ax^3}}\sqrt{1+ax^3}\text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(-2+m), \frac{4+m}{6}, a^2x^6\right)}{a(2+m-m^2)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.61 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.72

$$\int e^{\operatorname{sech}^{-1}(ax^3)} x^m dx$$

$$= \frac{2^{\frac{1+m}{3}} e^{\operatorname{sech}^{-1}(ax^3)} \left(\frac{e^{\operatorname{sech}^{-1}(ax^3)}}{1+e^{2\operatorname{sech}^{-1}(ax^3)}} \right)^{\frac{1+m}{3}} \left(1 + e^{2\operatorname{sech}^{-1}(ax^3)} \right)^{\frac{1+m}{3}} x^{1+m} (ax^3)^{\frac{1}{3}(-1-m)} \left((10+m) \operatorname{Hypergeometric2F1} \left[\frac{4+m}{6}, \frac{4+m}{3}, \frac{10+m}{6}, -E^{2\operatorname{sech}^{-1}(ax^3)} \right] - E^{2\operatorname{sech}^{-1}(ax^3)} (4+m) \operatorname{Hypergeometric2F1} \left[\frac{4+m}{3}, \frac{10+m}{6}, \frac{16+m}{6}, -E^{2\operatorname{sech}^{-1}(ax^3)} \right] \right)}{(4+m)(10+m)}$$

[In] Integrate[E^ArcSech[a*x^3]*x^m,x]

[Out] (2^((1+m)/3)*E^ArcSech[a*x^3]*(E^ArcSech[a*x^3]/(1+E^(2*ArcSech[a*x^3])))^((1+m)/3)*(1+E^(2*ArcSech[a*x^3]))^((1+m)/3)*x^(1+m)*(a*x^3)^((-1-m)/3)*((10+m)*Hypergeometric2F1[(4+m)/6,(4+m)/3,(10+m)/6,-E^(2*ArcSech[a*x^3])] - E^(2*ArcSech[a*x^3])*(4+m)*Hypergeometric2F1[(4+m)/3,(10+m)/6,(16+m)/6,-E^(2*ArcSech[a*x^3])])/(4+m*(10+m))

Maple [F]

$$\int \left(\frac{1}{ax^3} + \sqrt{\frac{1}{ax^3} - 1} \sqrt{\frac{1}{ax^3} + 1} \right) x^m dx$$

[In] int((1/a/x^3+(1/a/x^3-1)^(1/2)*(1/a/x^3+1)^(1/2))*x^m,x)

[Out] int((1/a/x^3+(1/a/x^3-1)^(1/2)*(1/a/x^3+1)^(1/2))*x^m,x)

Fricas [F]

$$\int e^{\operatorname{sech}^{-1}(ax^3)} x^m dx = \int x^m \left(\sqrt{\frac{1}{ax^3} + 1} \sqrt{\frac{1}{ax^3} - 1} + \frac{1}{ax^3} \right) dx$$

[In] integrate((1/a/x^3+(1/a/x^3-1)^(1/2)*(1/a/x^3+1)^(1/2))*x^m,x, algorithm="fricas")

[Out] integral((a*x^3*x^m*sqrt((a*x^3+1)/(a*x^3))*sqrt(-(a*x^3-1)/(a*x^3))+x^m)/(a*x^3), x)

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax^3)} x^m dx = \frac{\int \frac{x^m}{x^3} dx + \int ax^m \sqrt{-1 + \frac{1}{ax^3}} \sqrt{1 + \frac{1}{ax^3}} dx}{a}$$

[In] integrate((1/a/x**3+(1/a/x**3-1)**(1/2)*(1/a/x**3+1)**(1/2))*x**m,x)

[Out] (Integral(x**m/x**3, x) + Integral(a*x**m*sqrt(-1 + 1/(a*x**3))*sqrt(1 + 1/(a*x**3)), x))/a

Maxima [F(-2)]

Exception generated.

$$\int e^{\operatorname{sech}^{-1}(ax^3)} x^m dx = \text{Exception raised: ValueError}$$

[In] integrate((1/a/x^3+(1/a/x^3-1)^(1/2)*(1/a/x^3+1)^(1/2))*x^m,x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested a additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(m-3>0)', see 'assume?' for more details)Is

Giac [F]

$$\int e^{\operatorname{sech}^{-1}(ax^3)} x^m dx = \int x^m \left(\sqrt{\frac{1}{ax^3} + 1} \sqrt{\frac{1}{ax^3} - 1 + \frac{1}{ax^3}} \right) dx$$

[In] integrate((1/a/x^3+(1/a/x^3-1)^(1/2)*(1/a/x^3+1)^(1/2))*x^m,x, algorithm="giac")

[Out] integrate(x^m*(sqrt(1/(a*x^3) + 1)*sqrt(1/(a*x^3) - 1) + 1/(a*x^3)), x)

Mupad [F(-1)]

Timed out.

$$\int e^{\operatorname{sech}^{-1}(ax^3)} x^m dx = \int x^m \left(\sqrt{\frac{1}{ax^3} - 1} \sqrt{\frac{1}{ax^3} + 1} + \frac{1}{ax^3} \right) dx$$

```
[In] int(x^m*((1/(a*x^3) - 1)^(1/2)*(1/(a*x^3) + 1)^(1/2) + 1/(a*x^3)),x)
```

```
[Out] int(x^m*((1/(a*x^3) - 1)^(1/2)*(1/(a*x^3) + 1)^(1/2) + 1/(a*x^3)), x)
```

3.57 $\int e^{\operatorname{sech}^{-1}(ax^2)} x^m dx$

Optimal result	387
Rubi [A] (verified)	387
Mathematica [A] (warning: unable to verify)	389
Maple [F]	389
Fricas [F]	389
Sympy [F]	390
Maxima [F(-2)]	390
Giac [F(-2)]	390
Mupad [F(-1)]	391

Optimal result

Integrand size = 12, antiderivative size = 107

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^m dx = -\frac{2x^{-1+m}}{a(1-m^2)} + \frac{e^{\operatorname{sech}^{-1}(ax^2)} x^{1+m}}{1+m} - \frac{2x^{-1+m} \sqrt{\frac{1}{1+ax^2}} \sqrt{1+ax^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(-1+m), \frac{3+m}{4}, a^2 x^4\right)}{a(1-m^2)}$$

[Out] $-2*x^{(-1+m)}/a/(-m^2+1)+(1/a/x^2+(1/a/x^2-1)^{(1/2)}*(1/a/x^2+1)^{(1/2)})*x^{(1+m)}/(1+m)-2*x^{(-1+m)}*hypergeom([1/2, -1/4+1/4*m], [3/4+1/4*m], a^2*x^4)*(1/(a*x^2+1))^{(1/2)}*(a*x^2+1)^{(1/2)}/a/(-m^2+1)$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6470, 30, 265, 371}

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^m dx = -\frac{2\sqrt{\frac{1}{ax^2+1}} \sqrt{ax^2+1} x^{m-1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m-1}{4}, \frac{m+3}{4}, a^2 x^4\right)}{a(1-m^2)} - \frac{2x^{m-1}}{a(1-m^2)} + \frac{x^{m+1} e^{\operatorname{sech}^{-1}(ax^2)}}{m+1}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcSech}[a*x^2]}*x^m, x]$

[Out] $(-2x^{(-1+m)})/(a*(1-m^2)) + (E^{\text{ArcSech}[a*x^2]}*x^{(1+m)})/(1+m) - (2*x^{(-1+m)}*\text{Sqrt}[(1+a*x^2)^{-1}]*\text{Sqrt}[1+a*x^2]*\text{Hypergeometric2F1}[1/2, (-1+m)/4, (3+m)/4, a^2*x^4])/(a*(1-m^2))$

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}/(m+1), x] /;$ FreeQ[m, x] && NeQ[m, -1]

Rule 265

$\text{Int}[(c_*)(x_)^{(m_.)}*((a1_) + (b1_)*(x_)^{(n_)})^{(p_)}*((a2_) + (b2_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Int}[(c*x)^m*(a1*a2 + b1*b2*x^{(2*n)})^p, x] /;$ FreeQ[{a1, b1, a2, b2, c, m, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))

Rule 371

$\text{Int}[(c_*)(x_)^{(m_.)}*((a_) + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[a^p*((c*x)^{(m+1)})/(c*(m+1))*\text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /;$ FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 6470

$\text{Int}[E^{\text{ArcSech}[(a_)*(x_)^{(p_.)}]}*(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*(E^{\text{ArcSech}[a*x^p]}/(m+1)), x] + (\text{Dist}[p/(a*(m+1)), \text{Int}[x^{(m-p)}, x], x] + \text{Dist}[p*(\text{Sqrt}[1+a*x^p]/(a*(m+1)))*\text{Sqrt}[1/(1+a*x^p)], \text{Int}[x^{(m-p)}/(\text{Sqrt}[1+a*x^p]*\text{Sqrt}[1-a*x^p]), x], x]) /;$ FreeQ[{a, m, p}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{e^{\text{sech}^{-1}(ax^2)}x^{1+m}}{1+m} + \frac{2 \int x^{-2+m} dx}{a(1+m)} + \frac{\left(2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{x^{-2+m}}{\sqrt{1-ax^2}\sqrt{1+ax^2}} dx}{a(1+m)} \\ &= -\frac{2x^{-1+m}}{a(1-m^2)} + \frac{e^{\text{sech}^{-1}(ax^2)}x^{1+m}}{1+m} + \frac{\left(2\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\right) \int \frac{x^{-2+m}}{\sqrt{1-a^2x^4}} dx}{a(1+m)} \\ &= -\frac{2x^{-1+m}}{a(1-m^2)} + \frac{e^{\text{sech}^{-1}(ax^2)}x^{1+m}}{1+m} \\ &\quad - \frac{2x^{-1+m}\sqrt{\frac{1}{1+ax^2}}\sqrt{1+ax^2}\text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(-1+m), \frac{3+m}{4}, a^2x^4\right)}{a(1-m^2)} \end{aligned}$$

Mathematica [A] (warning: unable to verify)

Time = 2.08 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.49

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^m dx$$

$$= \frac{2^{\frac{1+m}{2}} e^{\operatorname{sech}^{-1}(ax^2)} \left(\frac{e^{\operatorname{sech}^{-1}(ax^2)}}{1+e^{2\operatorname{sech}^{-1}(ax^2)}} \right)^{\frac{1+m}{2}} x^{1+m} (ax^2)^{\frac{1}{2}(-1-m)} \left((7+m) \operatorname{Hypergeometric2F1} \left(1, \frac{1-m}{4}, \frac{7+m}{4}, -e^{2\operatorname{sech}^{-1}(ax^2)} \right) - E^{2\operatorname{sech}^{-1}(ax^2)} (3+m)(7+m) \right)}{(3+m)(7+m)}$$

`[In] Integrate[E^ArcSech[a*x^2]*x^m,x]`

```
[Out] (2^((1+m)/2)*E^ArcSech[a*x^2]*(E^ArcSech[a*x^2]/(1+E^(2*ArcSech[a*x^2])))^((1+m)/2)*x^(1+m)*(a*x^2)^((-1-m)/2)*((7+m)*Hypergeometric2F1[1,
(1-m)/4,(7+m)/4,-E^(2*ArcSech[a*x^2])]) - E^(2*ArcSech[a*x^2])*(3+m)*Hypergeometric2F1[1,(5-m)/4,(11+m)/4,-E^(2*ArcSech[a*x^2])])/(3+m)*(7+m))
```

Maple [F]

$$\int \left(\frac{1}{ax^2} + \sqrt{\frac{1}{ax^2} - 1} \sqrt{\frac{1}{ax^2} + 1} \right) x^m dx$$

`[In] int((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^m,x)``[Out] int((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^m,x)`**Fricas [F]**

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^m dx = \int x^m \left(\sqrt{\frac{1}{ax^2} + 1} \sqrt{\frac{1}{ax^2} - 1} + \frac{1}{ax^2} \right) dx$$

`[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^m,x, algorithm="fricas")`

```
[Out] integral((a*x^2*x^m*sqrt((a*x^2+1)/(a*x^2))*sqrt(-(a*x^2-1)/(a*x^2))+x^m)/(a*x^2), x)
```

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^m dx = \frac{\int \frac{x^m}{x^2} dx + \int ax^m \sqrt{-1 + \frac{1}{ax^2}} \sqrt{1 + \frac{1}{ax^2}} dx}{a}$$

[In] integrate((1/a/x**2+(1/a/x**2-1)**(1/2)*(1/a/x**2+1)**(1/2))*x**m,x)

[Out] (Integral(x**m/x**2, x) + Integral(a*x**m*sqrt(-1 + 1/(a*x**2))*sqrt(1 + 1/(a*x**2)), x))/a

Maxima [F(-2)]

Exception generated.

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^m dx = \text{Exception raised: ValueError}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^m,x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(m-2>0)', see 'assume?' for more details)Is

Giac [F(-2)]

Exception generated.

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^m dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/x^2+(1/a/x^2-1)^(1/2)*(1/a/x^2+1)^(1/2))*x^m,x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int e^{\operatorname{sech}^{-1}(ax^2)} x^m dx = \int x^m \left(\sqrt{\frac{1}{ax^2} - 1} \sqrt{\frac{1}{ax^2} + 1} + \frac{1}{ax^2} \right) dx$$

```
[In] int(x^m*((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2)),x)
```

```
[Out] int(x^m*((1/(a*x^2) - 1)^(1/2)*(1/(a*x^2) + 1)^(1/2) + 1/(a*x^2)), x)
```

3.58 $\int e^{\operatorname{sech}^{-1}(ax)} x^m dx$

Optimal result	392
Rubi [A] (verified)	392
Mathematica [A] (verified)	394
Maple [F]	394
Fricas [F]	394
Sympy [F]	395
Maxima [F]	395
Giac [F]	395
Mupad [F(-1)]	395

Optimal result

Integrand size = 10, antiderivative size = 91

$$\int e^{\operatorname{sech}^{-1}(ax)} x^m dx = \frac{x^m}{am(1+m)} + \frac{e^{\operatorname{sech}^{-1}(ax)} x^{1+m}}{1+m} + \frac{x^m \sqrt{\frac{1}{1+ax}} \sqrt{1+ax} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m}{2}, \frac{2+m}{2}, a^2 x^2\right)}{am(1+m)}$$

[Out] $x^m/a/m/(1+m)+(1/a/x+(1/a/x-1)^{(1/2)}*(1+1/a/x)^{(1/2)})*x^{(1+m)}/(1+m)+x^m*\operatorname{hypergeom}([1/2, 1/2*m], [1+1/2*m], a^2*x^2)*(1/(a*x+1))^{(1/2)}*(a*x+1)^{(1/2)}/a/m/(1+m)$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {6470, 30, 126, 371}

$$\int e^{\operatorname{sech}^{-1}(ax)} x^m dx = \frac{\sqrt{\frac{1}{ax+1}} \sqrt{ax+1} x^m \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m}{2}, \frac{m+2}{2}, a^2 x^2\right)}{am(m+1)} + \frac{x^{m+1} e^{\operatorname{sech}^{-1}(ax)}}{m+1} + \frac{x^m}{am(m+1)}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcSech}[a*x]}*x^m, x]$

[Out] $x^m/(a*m*(1+m)) + (E^{\operatorname{ArcSech}[a*x]}*x^{(1+m)})/(1+m) + (x^m*\operatorname{Sqrt}[(1+a*x)^{-1}]*\operatorname{Sqrt}[1+a*x]*\operatorname{Hypergeometric2F1}[1/2, m/2, (2+m)/2, a^2*x^2])/(a*m*(1+m))$

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 126

Int[((f_)*(x_))^(p_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[(a*c + b*d*x^2)^m*(f*x)^p, x] /; FreeQ[{a, b, c, d, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && GtQ[a, 0] && GtQ[c, 0]

Rule 371

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 6470

Int[E^ArcSech[(a_)*(x_)^(p_)]*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*(E^ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] + Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, m, p}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{e^{\operatorname{sech}^{-1}(ax)} x^{1+m}}{1+m} + \frac{\int x^{-1+m} dx}{a(1+m)} + \frac{\left(\sqrt{\frac{1}{1+ax}} \sqrt{1+ax}\right) \int \frac{x^{-1+m}}{\sqrt{1-ax}\sqrt{1+ax}} dx}{a(1+m)} \\
 &= \frac{x^m}{am(1+m)} + \frac{e^{\operatorname{sech}^{-1}(ax)} x^{1+m}}{1+m} + \frac{\left(\sqrt{\frac{1}{1+ax}} \sqrt{1+ax}\right) \int \frac{x^{-1+m}}{\sqrt{1-a^2x^2}} dx}{a(1+m)} \\
 &= \frac{x^m}{am(1+m)} + \frac{e^{\operatorname{sech}^{-1}(ax)} x^{1+m}}{1+m} + \frac{x^m \sqrt{\frac{1}{1+ax}} \sqrt{1+ax} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m}{2}, \frac{2+m}{2}, a^2x^2\right)}{am(1+m)}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.59

$$\int e^{\operatorname{sech}^{-1}(ax)} x^m dx = \frac{2^{1+m} e^{2\operatorname{sech}^{-1}(ax)} \left(\frac{e^{\operatorname{sech}^{-1}(ax)}}{1+e^{2\operatorname{sech}^{-1}(ax)}} \right)^m \left(1 + e^{2\operatorname{sech}^{-1}(ax)} \right)^m x^m (ax)^{-m} \left(-\left((4+m) \operatorname{Hypergeometric2F1} \left(1 + \frac{m}{2}, 2 + m, 2 + m/2, -E^{(2\operatorname{sech}^{-1}(ax))} \right) \right) + E^{(2\operatorname{sech}^{-1}(ax))} (2+m) \operatorname{Hypergeometric2F1} \left(2 + m/2, 2 + m, 3 + m/2, -E^{(2\operatorname{sech}^{-1}(ax))} \right) \right)}{a(2+m)(4+m)(ax)^m}$$

[In] Integrate[E^ArcSech[a*x]*x^m,x]

[Out] $-\left((2^{1+m} E^{(2\operatorname{ArcSech}[a*x])} (E^{\operatorname{ArcSech}[a*x]} / (1 + E^{(2\operatorname{ArcSech}[a*x])}))^m (1 + E^{(2\operatorname{ArcSech}[a*x])})^m x^m \left(-\left((4+m) \operatorname{Hypergeometric2F1} \left[1 + m/2, 2 + m, 2 + m/2, -E^{(2\operatorname{ArcSech}[a*x])} \right] \right) + E^{(2\operatorname{ArcSech}[a*x])} (2+m) \operatorname{Hypergeometric2F1} \left[2 + m/2, 2 + m, 3 + m/2, -E^{(2\operatorname{ArcSech}[a*x])} \right] \right) \right) / (a(2+m)(4+m)(a*x)^m)$

Maple [F]

$$\int \left(\frac{1}{ax} + \sqrt{\frac{1}{ax} - 1} \sqrt{1 + \frac{1}{ax}} \right) x^m dx$$

[In] int((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x^m,x)

[Out] int((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x^m,x)

Fricas [F]

$$\int e^{\operatorname{sech}^{-1}(ax)} x^m dx = \int x^m \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} \right) dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x^m,x, algorithm="fricas")

[Out] integral((a*x*x^m*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + x^m)/(a*x), x)

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax)} x^m dx = \frac{\int \frac{x^m}{x} dx + \int ax^m \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} dx}{a}$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))*x**m,x)

[Out] (Integral(x**m/x, x) + Integral(a*x**m*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)), x))/a

Maxima [F]

$$\int e^{\operatorname{sech}^{-1}(ax)} x^m dx = \int x^m \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} \right) dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x^m,x, algorithm="maxima")

[Out] integrate(sqrt(a*x + 1)*sqrt(-a*x + 1)*x^m/x, x)/a + x^m/(a*m)

Giac [F]

$$\int e^{\operatorname{sech}^{-1}(ax)} x^m dx = \int x^m \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} \right) dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x^m,x, algorithm="giac")

[Out] integrate(x^m*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x)), x)

Mupad [F(-1)]

Timed out.

$$\int e^{\operatorname{sech}^{-1}(ax)} x^m dx = \int x^m \left(\sqrt{\frac{1}{ax} - 1} \sqrt{\frac{1}{ax} + 1} + \frac{1}{ax} \right) dx$$

[In] int(x^m*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x)),x)

[Out] int(x^m*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x)), x)

3.59 $\int e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} x^m dx$

Optimal result	396
Rubi [A] (verified)	396
Mathematica [A] (warning: unable to verify)	398
Maple [F]	398
Fricas [F]	398
Sympy [F]	399
Maxima [F]	399
Giac [F]	399
Mupad [F(-1)]	399

Optimal result

Integrand size = 12, antiderivative size = 109

$$\int e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} x^m dx = \frac{e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} x^{1+m}}{1+m} - \frac{x^{2+m}}{a(2+3m+m^2)} - \frac{\sqrt{\frac{1}{1+\frac{a}{x}}}\sqrt{1+\frac{a}{x}}x^{2+m} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-2-m), -\frac{m}{2}, \frac{a^2}{x^2}\right)}{a(2+3m+m^2)}$$

[Out] (x/a+(-1+x/a)^(1/2)*(1+x/a)^(1/2))*x^(1+m)/(1+m)-x^(2+m)/a/(m^2+3*m+2)-x^(2+m)*hypergeom([1/2, -1-1/2*m], [-1/2*m], a^2/x^2)*(1/(1+a/x))^(1/2)*(1+a/x)^(1/2)/a/(m^2+3*m+2)

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {6470, 30, 265, 346, 371}

$$\int e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} x^m dx = -\frac{\sqrt{\frac{1}{\frac{a}{x}+1}}\sqrt{\frac{a}{x}+1}x^{m+2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-m-2), -\frac{m}{2}, \frac{a^2}{x^2}\right)}{a(m^2+3m+2)} - \frac{x^{m+2}}{a(m^2+3m+2)} + \frac{x^{m+1}e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)}}{m+1}$$

[In] Int[E^ArcSech[a/x]*x^m,x]

[Out] (E^ArcSech[a/x]*x^(1+m))/(1+m)-x^(2+m)/(a*(2+3*m+m^2))-(Sqrt[(1+a/x)^(-1)]*Sqrt[1+a/x]*x^(2+m)*Hypergeometric2F1[1/2, (-2-m)/2, -1/2*m, a^2/x^2])/(a*(2+3*m+m^2))

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 265

Int[((c_)*(x_))^(m_)*((a1_) + (b1_)*(x_)^(n_))^(p_)*((a2_) + (b2_)*(x_)^(n_))^(p_), x_Symbol] := Int[(c*x)^(m*(a1*a2 + b1*b2*x^(2*n)))^p, x] /; FreeQ[{a1, b1, a2, b2, c, m, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))

Rule 346

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[(-c^(-1))*(c*x)^(m + 1)*(1/x)^(m + 1), Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x], x] /; FreeQ[{a, b, c, m, p}, x] && ILtQ[n, 0] && !RationalQ[m]

Rule 371

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 6470

Int[E^ArcSech[(a_)*(x_)^(p_)]*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*(E^ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] + Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x)) /; FreeQ[{a, m, p}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} x^{1+m}}{1+m} - \frac{\int x^{1+m} dx}{a(1+m)} - \frac{\left(\sqrt{\frac{1}{1+\frac{a}{x}}}\sqrt{1+\frac{a}{x}}\right) \int \frac{x^{1+m}}{\sqrt{1-\frac{a}{x}}\sqrt{1+\frac{a}{x}}} dx}{a(1+m)} \\
 &= \frac{e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} x^{1+m}}{1+m} - \frac{x^{2+m}}{a(2+3m+m^2)} - \frac{\left(\sqrt{\frac{1}{1+\frac{a}{x}}}\sqrt{1+\frac{a}{x}}\right) \int \frac{x^{1+m}}{\sqrt{1-\frac{a^2}{x^2}}} dx}{a(1+m)} \\
 &= \frac{e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} x^{1+m}}{1+m} - \frac{x^{2+m}}{a(2+3m+m^2)} + \frac{\left(\sqrt{\frac{1}{1+\frac{a}{x}}}\sqrt{1+\frac{a}{x}}\left(\frac{1}{x}\right)^m x^m\right) \operatorname{Subst}\left(\int \frac{x^{-3-m}}{\sqrt{1-a^2x^2}} dx, x, \frac{1}{x}\right)}{a(1+m)}
 \end{aligned}$$

$$= \frac{e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} x^{1+m}}{1+m} - \frac{x^{2+m}}{a(2+3m+m^2)}$$

$$- \frac{\sqrt{\frac{1}{1+\frac{a}{x}}}\sqrt{1+\frac{a}{x}}x^{2+m} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-2-m), -\frac{m}{2}, \frac{a^2}{x^2}\right)}{a(2+3m+m^2)}$$

Mathematica [A] (warning: unable to verify)

Time = 0.71 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.28

$$\int e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} x^m dx =$$

$$\frac{2^{-1-m} a e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} \left(\frac{e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)}}{1+e^{2\operatorname{sech}^{-1}\left(\frac{a}{x}\right)}}\right)^{-1-m} \left(\frac{a}{x}\right)^m x^m \left(-\left((-2+m)\operatorname{Hypergeometric2F1}\left(1, 1+\frac{m}{2}, 1-\frac{m}{2}, -e^{2\operatorname{sech}^{-1}\left(\frac{a}{x}\right)}\right)\right) + \left(2\operatorname{Hypergeometric2F1}\left(1, 2+\frac{m}{2}, 2-m, -e^{2\operatorname{sech}^{-1}\left(\frac{a}{x}\right)}\right)\right)\right)}{(-2+m)m}$$

[In] Integrate[E^ArcSech[a/x]*x^m,x]

[Out] -((2^(-1-m)*a*E^ArcSech[a/x]*(E^ArcSech[a/x]/(1+E^(2*ArcSech[a/x]))))^(-1-m)*(a/x)^m*x^m*(-((-2+m)*Hypergeometric2F1[1, 1+m/2, 1-m/2, -E^(2*ArcSech[a/x])]) + E^(2*ArcSech[a/x])*m*Hypergeometric2F1[1, 2+m/2, 2-m/2, -E^(2*ArcSech[a/x])]))/((-2+m)*m)

Maple [F]

$$\int \left(\frac{x}{a} + \sqrt{-1 + \frac{x}{a}} \sqrt{1 + \frac{x}{a}}\right) x^m dx$$

[In] int((x/a+(-1+x/a)^(1/2)*(1+x/a)^(1/2))*x^m,x)

[Out] int((x/a+(-1+x/a)^(1/2)*(1+x/a)^(1/2))*x^m,x)

Fricas [F]

$$\int e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} x^m dx = \int x^m \left(\sqrt{\frac{x}{a} + 1} \sqrt{\frac{x}{a} - 1 + \frac{x}{a}}\right) dx$$

[In] integrate((x/a+(-1+x/a)^(1/2)*(1+x/a)^(1/2))*x^m,x, algorithm="fricas")

[Out] integral((a*x^m*sqrt((a+x)/a)*sqrt(-(a-x)/a) + x*x^m)/a, x)

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} x^m dx = \frac{\int x x^m dx + \int a x^m \sqrt{-1 + \frac{x}{a}} \sqrt{1 + \frac{x}{a}} dx}{a}$$

[In] integrate((x/a+(-1+x/a)**(1/2)*(1+x/a)**(1/2))*x**m,x)

[Out] (Integral(x**m, x) + Integral(a*x**m*sqrt(-1 + x/a)*sqrt(1 + x/a), x))/a

Maxima [F]

$$\int e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} x^m dx = \int x^m \left(\sqrt{\frac{x}{a} + 1} \sqrt{\frac{x}{a} - 1 + \frac{x}{a}} \right) dx$$

[In] integrate((x/a+(-1+x/a)^(1/2)*(1+x/a)^(1/2))*x^m,x, algorithm="maxima")

[Out] x^2*x^m/(a*(m + 2)) + integrate(sqrt(a + x)*sqrt(-a + x)*x^m, x)/a

Giac [F]

$$\int e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} x^m dx = \int x^m \left(\sqrt{\frac{x}{a} + 1} \sqrt{\frac{x}{a} - 1 + \frac{x}{a}} \right) dx$$

[In] integrate((x/a+(-1+x/a)^(1/2)*(1+x/a)^(1/2))*x^m,x, algorithm="giac")

[Out] integrate(x^m*(sqrt(x/a + 1)*sqrt(x/a - 1) + x/a), x)

Mupad [F(-1)]

Timed out.

$$\int e^{\operatorname{sech}^{-1}\left(\frac{a}{x}\right)} x^m dx = \int x^m \left(\sqrt{\frac{x}{a} - 1} \sqrt{\frac{x}{a} + 1 + \frac{x}{a}} \right) dx$$

[In] int(x^m*((x/a - 1)^(1/2)*(x/a + 1)^(1/2) + x/a),x)

[Out] int(x^m*((x/a - 1)^(1/2)*(x/a + 1)^(1/2) + x/a), x)

3.60 $\int e^{\operatorname{sech}^{-1}(ax^p)} x^m dx$

Optimal result	400
Rubi [A] (verified)	400
Mathematica [A] (warning: unable to verify)	402
Maple [F]	402
Fricas [F(-2)]	402
Sympy [F]	403
Maxima [F(-2)]	403
Giac [F]	403
Mupad [F(-1)]	404

Optimal result

Integrand size = 12, antiderivative size = 133

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x^m dx$$

$$= \frac{e^{\operatorname{sech}^{-1}(ax^p)} x^{1+m}}{1+m} + \frac{px^{1+m-p}}{a(1+m)(1+m-p)}$$

$$+ \frac{px^{1+m-p} \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m-p}{2p}, \frac{1+m+p}{2p}, a^2 x^{2p}\right)}{a(1+m)(1+m-p)}$$

[Out] $(1/a/(x^p)+(1/a/(x^p)-1)^{(1/2)}*(1/a/(x^p)+1)^{(1/2)})*x^{(1+m)/(1+m)+p*x^{(1+m-p)/a/(1+m)/(1+m-p)+p*x^{(1+m-p)*\operatorname{hypergeom}([1/2, 1/2*(1+m-p)/p], [1/2*(1+m+p)/p], a^2*x^{(2p)})*(1/(1+a*x^p))^{(1/2)}*(1+a*x^p)^{(1/2)/a/(1+m)/(1+m-p)}$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6470, 30, 265, 371}

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x^m dx$$

$$= \frac{p\sqrt{\frac{1}{ax^p+1}}\sqrt{ax^p+1}x^{m-p+1}\operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m-p+1}{2p}, \frac{m+p+1}{2p}, a^2 x^{2p}\right)}{a(m+1)(m-p+1)}$$

$$+ \frac{px^{m-p+1}}{a(m+1)(m-p+1)} + \frac{x^{m+1}e^{\operatorname{sech}^{-1}(ax^p)}}{m+1}$$

[In] Int[E^ArcSech[a*x^p]*x^m,x]

[Out] $(E^{\text{ArcSech}[a*x^p]*x^{(1+m)}})/(1+m) + (p*x^{(1+m-p)})/(a*(1+m)*(1+m-p)) + (p*x^{(1+m-p)}*\text{Sqrt}[(1+a*x^p)^{-1}]*\text{Sqrt}[1+a*x^p]*\text{Hypergeometric2F1}[1/2, (1+m-p)/(2*p), (1+m+p)/(2*p), a^2*x^{(2*p)}])/(a*(1+m)*(1+m-p))$

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}/(m+1), x] /; \text{FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 265

$\text{Int}[(c_)*(x_)^{(m_.)}*((a1_) + (b1_)*(x_)^{(n_)})^{(p_)}*((a2_) + (b2_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Int}[(c*x)^m*(a1*a2 + b1*b2*x^{(2*n)})^p, x] /; \text{FreeQ}[\{a1, b1, a2, b2, c, m, n, p\}, x] \ \&\& \ \text{EqQ}[a2*b1 + a1*b2, 0] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{GtQ}[a1, 0] \ \&\& \ \text{GtQ}[a2, 0]))$

Rule 371

$\text{Int}[(c_)*(x_)^{(m_.)}*((a_) + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1)}/(c*(m+1))) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}[\{a, b, c, m, n, p\}, x] \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

Rule 6470

$\text{Int}[E^{\text{ArcSech}[a_)*(x_)^{(p_.)}]*x^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*(E^{\text{ArcSech}[a*x^p]/(m+1)}), x] + (\text{Dist}[p/(a*(m+1)), \text{Int}[x^{(m-p)}, x], x] + \text{Dist}[p*(\text{Sqrt}[1+a*x^p]/(a*(m+1)))*\text{Sqrt}[1/(1+a*x^p)], \text{Int}[x^{(m-p)}/(\text{Sqrt}[1+a*x^p]*\text{Sqrt}[1-a*x^p]), x], x]) /; \text{FreeQ}[\{a, m, p\}, x] \ \&\& \ \text{NeQ}[m, -1]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{e^{\text{sech}^{-1}(ax^p)} x^{1+m}}{1+m} + \frac{p \int x^{m-p} dx}{a(1+m)} + \frac{\left(p \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p}\right) \int \frac{x^{m-p}}{\sqrt{1-ax^p} \sqrt{1+ax^p}} dx}{a(1+m)} \\ &= \frac{e^{\text{sech}^{-1}(ax^p)} x^{1+m}}{1+m} + \frac{px^{1+m-p}}{a(1+m)(1+m-p)} + \frac{\left(p \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p}\right) \int \frac{x^{m-p}}{\sqrt{1-a^2x^{2p}}} dx}{a(1+m)} \\ &= \frac{e^{\text{sech}^{-1}(ax^p)} x^{1+m}}{1+m} + \frac{px^{1+m-p}}{a(1+m)(1+m-p)} \\ &\quad + \frac{px^{1+m-p} \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m-p}{2p}, \frac{1+m+p}{2p}, a^2x^{2p}\right)}{a(1+m)(1+m-p)} \end{aligned}$$

Mathematica [A] (warning: unable to verify)

Time = 4.33 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.40

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x^m dx$$

$$= \frac{2^{\frac{1+m}{p}} e^{\operatorname{sech}^{-1}(ax^p)} \left(\frac{e^{\operatorname{sech}^{-1}(ax^p)}}{1+e^{2\operatorname{sech}^{-1}(ax^p)}} \right)^{\frac{1+m}{p}} x^{1+m} (ax^p)^{-\frac{1+m}{p}} \left(-e^{2\operatorname{sech}^{-1}(ax^p)} (1+m+p) \operatorname{Hypergeometric2F1} \left(1, -\frac{1+m}{2(1+m+p)}, \frac{1+m+p}{2(1+m+p)}, -e^{2\operatorname{sech}^{-1}(ax^p)} \right) + (1+m+3p) \operatorname{Hypergeometric2F1} \left(1, 1 - \frac{1+m+p}{2(1+m+p)}, \frac{1+m+3p}{2(1+m+p)}, -e^{2\operatorname{sech}^{-1}(ax^p)} \right) \right)}{(1+m+p)}$$

[In] Integrate[E^ArcSech[a*x^p]*x^m,x]

[Out] (2^((1+m)/p)*E^ArcSech[a*x^p]*(E^ArcSech[a*x^p]/(1+E^(2*ArcSech[a*x^p])))^((1+m)/p)*x^(1+m)*(-(E^(2*ArcSech[a*x^p]))*(1+m+p)*Hypergeometric2F1[1,-1/2*(1+m-3*p)/p,(1+m+5*p)/(2*p),-E^(2*ArcSech[a*x^p])])+(1+m+3*p)*Hypergeometric2F1[1,1-(1+m+p)/(2*p),(1+m+3*p)/(2*p),-E^(2*ArcSech[a*x^p])]))/((1+m+p)*(1+m+3*p)*(a*x^p)^((1+m)/p))

Maple [F]

$$\int \left(\frac{x^{-p}}{a} + \sqrt{\frac{x^{-p}}{a} - 1} \sqrt{\frac{x^{-p}}{a} + 1} \right) x^m dx$$

[In] int((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))*x^m,x)

[Out] int((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))*x^m,x)

Fricas [F(-2)]

Exception generated.

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x^m dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))*x^m,x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x^m dx = \frac{\int x^m x^{-p} dx + \int ax^m \sqrt{-1 + \frac{x^{-p}}{a}} \sqrt{1 + \frac{x^{-p}}{a}} dx}{a}$$

[In] integrate((1/a/(x**p)+(1/a/(x**p)-1)**(1/2)*(1/a/(x**p)+1)**(1/2))*x**m,x)

[Out] (Integral(x**m/x**p, x) + Integral(a*x**m*sqrt(-1 + 1/(a*x**p))*sqrt(1 + 1/(a*x**p)), x))/a

Maxima [F(-2)]

Exception generated.

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x^m dx = \text{Exception raised: ValueError}$$

[In] integrate((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))*x^m,x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested a dditional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(m-p>0)', see 'assume?' for more details)Is

Giac [F]

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x^m dx = \int x^m \left(\sqrt{\frac{1}{ax^p} + 1} \sqrt{\frac{1}{ax^p} - 1 + \frac{1}{ax^p}} \right) dx$$

[In] integrate((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))*x^m,x, algorithm="giac")

[Out] integrate(x^m*(sqrt(1/(a*x^p) + 1)*sqrt(1/(a*x^p) - 1) + 1/(a*x^p)), x)

Mupad [F(-1)]

Timed out.

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x^m dx = \int x^m \left(\sqrt{\frac{1}{ax^p} - 1} \sqrt{\frac{1}{ax^p} + 1} + \frac{1}{ax^p} \right) dx$$

```
[In] int(x^m*((1/(a*x^p) - 1)^(1/2)*(1/(a*x^p) + 1)^(1/2) + 1/(a*x^p)),x)
```

```
[Out] int(x^m*((1/(a*x^p) - 1)^(1/2)*(1/(a*x^p) + 1)^(1/2) + 1/(a*x^p)), x)
```

3.61 $\int e^{\operatorname{sech}^{-1}(ax^p)} x dx$

Optimal result	405
Rubi [A] (verified)	405
Mathematica [A] (verified)	407
Maple [F]	407
Fricas [F(-2)]	407
Sympy [F]	408
Maxima [F(-2)]	408
Giac [F]	408
Mupad [F(-1)]	409

Optimal result

Integrand size = 10, antiderivative size = 119

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x dx$$

$$= \frac{1}{2} e^{\operatorname{sech}^{-1}(ax^p)} x^2 + \frac{px^{2-p}}{2a(2-p)}$$

$$+ \frac{px^{2-p} \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(-1 + \frac{2}{p}\right), \frac{1}{2}\left(1 + \frac{2}{p}\right), a^2 x^{2p}\right)}{2a(2-p)}$$

[Out] 1/2*(1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))*x^2+1/2*p*x^(2-p)/a/(2-p)+1/2*p*x^(2-p)*hypergeom([1/2, -1/2+1/p], [1/2+1/p], a^2*x^(2*p))*(1/(1+a*x^p))^(1/2)*(1+a*x^p)^(1/2)/a/(2-p)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {6470, 30, 265, 371}

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x dx$$

$$= \frac{px^{2-p} \sqrt{\frac{1}{ax^p+1}} \sqrt{ax^p+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{2}{p}-1\right), \frac{1}{2}\left(1 + \frac{2}{p}\right), a^2 x^{2p}\right)}{2a(2-p)}$$

$$+ \frac{px^{2-p}}{2a(2-p)} + \frac{1}{2} x^2 e^{\operatorname{sech}^{-1}(ax^p)}$$

[In] Int[E^ArcSech[a*x^p]*x,x]

[Out] $(E^{\text{ArcSech}[a*x^p]*x^2)/2 + (p*x^{(2-p)})/(2*a*(2-p)) + (p*x^{(2-p)}*\text{Sqrt}[(1+a*x^p)^{-1}]*\text{Sqrt}[1+a*x^p]*\text{Hypergeometric2F1}[1/2, (-1+2/p)/2, (1+2/p)/2, a^2*x^{(2*p)}])/(2*a*(2-p))$

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}/(m+1), x] /;$ FreeQ[m, x] && NeQ[m, -1]

Rule 265

$\text{Int}[(c_)*(x_)^{(m_.)}*((a1_) + (b1_)*(x_)^{(n_.)})^{(p_.)}*((a2_) + (b2_)*(x_)^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Int}[(c*x)^m*(a1*a2 + b1*b2*x^{(2*n)})^p, x] /;$ FreeQ[{a1, b1, a2, b2, c, m, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))

Rule 371

$\text{Int}[(c_)*(x_)^{(m_.)}*((a_) + (b_)*(x_)^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[a^p*((c*x)^{(m+1)}/(c*(m+1)))*\text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /;$ FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 6470

$\text{Int}[E^{\text{ArcSech}[(a_)*(x_)^{(p_.)}]}*(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*(E^{\text{ArcSech}[a*x^p]/(m+1)}), x] + (\text{Dist}[p/(a*(m+1)), \text{Int}[x^{(m-p)}, x], x] + \text{Dist}[p*(\text{Sqrt}[1+a*x^p]/(a*(m+1)))*\text{Sqrt}[1/(1+a*x^p)], \text{Int}[x^{(m-p)}/(\text{Sqrt}[1+a*x^p]*\text{Sqrt}[1-a*x^p]), x], x]) /;$ FreeQ[{a, m, p}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{2} e^{\text{sech}^{-1}(ax^p)} x^2 + \frac{p \int x^{1-p} dx}{2a} + \frac{\left(p \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p}\right) \int \frac{x^{1-p}}{\sqrt{1-ax^p} \sqrt{1+ax^p}} dx}{2a} \\ &= \frac{1}{2} e^{\text{sech}^{-1}(ax^p)} x^2 + \frac{px^{2-p}}{2a(2-p)} + \frac{\left(p \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p}\right) \int \frac{x^{1-p}}{\sqrt{1-a^2x^{2p}}} dx}{2a} \\ &= \frac{1}{2} e^{\text{sech}^{-1}(ax^p)} x^2 + \frac{px^{2-p}}{2a(2-p)} \\ &\quad + \frac{px^{2-p} \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(-1+\frac{2}{p}\right), \frac{1}{2}\left(1+\frac{2}{p}\right), a^2x^{2p}\right)}{2a(2-p)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.34

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x dx$$

$$= \frac{x^{2-p} \left(-1 - \sqrt{\frac{1-ax^p}{1+ax^p}} - ax^p \sqrt{\frac{1-ax^p}{1+ax^p}} + \frac{a^2 p x^{2p} \sqrt{\frac{1-ax^p}{1+ax^p}} \sqrt{1-a^2 x^{2p}} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} + \frac{1}{p}, \frac{3}{2} + \frac{1}{p}, a^2 x^{2p}\right)}{(2+p)(-1+ax^p)} \right)}{a(-2+p)}$$

[In] Integrate[E^ArcSech[a*x^p]*x,x]

[Out] (x^(2 - p)*(-1 - Sqrt[(1 - a*x^p)/(1 + a*x^p)] - a*x^p*Sqrt[(1 - a*x^p)/(1 + a*x^p)] + (a^2*p*x^(2*p)*Sqrt[(1 - a*x^p)/(1 + a*x^p)]*Sqrt[1 - a^2*x^(2*p)])*Hypergeometric2F1[1/2, 1/2 + p^(-1), 3/2 + p^(-1), a^2*x^(2*p)])/((2 + p)*(-1 + a*x^p)))/(a*(-2 + p))

Maple [F]

$$\int \left(\frac{x^{-p}}{a} + \sqrt{\frac{x^{-p}}{a} - 1} \sqrt{\frac{x^{-p}}{a} + 1} \right) x dx$$

[In] int((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))*x,x)

[Out] int((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))*x,x)

Fricas [F(-2)]

Exception generated.

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))*x,x, algorithm m="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x dx = \frac{\int x x^{-p} dx + \int ax \sqrt{-1 + \frac{x^{-p}}{a}} \sqrt{1 + \frac{x^{-p}}{a}} dx}{a}$$

[In] integrate((1/a/(x**p)+(1/a/(x**p)-1)**(1/2)*(1/a/(x**p)+1)**(1/2))*x,x)

[Out] (Integral(x/x**p, x) + Integral(a*x*sqrt(-1 + 1/(a*x**p))*sqrt(1 + 1/(a*x**p)), x))/a

Maxima [F(-2)]

Exception generated.

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x dx = \text{Exception raised: ValueError}$$

[In] integrate((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))*x,x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(1-p>0)', see 'assume?' for more details)Is

Giac [F]

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x dx = \int x \left(\sqrt{\frac{1}{ax^p} + 1} \sqrt{\frac{1}{ax^p} - 1} + \frac{1}{ax^p} \right) dx$$

[In] integrate((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))*x,x, algorithm="giac")

[Out] integrate(x*(sqrt(1/(a*x^p) + 1)*sqrt(1/(a*x^p) - 1) + 1/(a*x^p)), x)

Mupad [F(-1)]

Timed out.

$$\int e^{\operatorname{sech}^{-1}(ax^p)} x dx = \int x \left(\sqrt{\frac{1}{ax^p} - 1} \sqrt{\frac{1}{ax^p} + 1} + \frac{1}{ax^p} \right) dx$$

[In] `int(x*((1/(a*x^p) - 1)^(1/2)*(1/(a*x^p) + 1)^(1/2) + 1/(a*x^p)),x)`

[Out] `int(x*((1/(a*x^p) - 1)^(1/2)*(1/(a*x^p) + 1)^(1/2) + 1/(a*x^p)), x)`

3.62 $\int e^{\operatorname{sech}^{-1}(ax^p)} dx$

Optimal result	410
Rubi [A] (verified)	410
Mathematica [A] (verified)	412
Maple [F]	412
Fricas [F(-2)]	412
Sympy [F]	413
Maxima [F(-2)]	413
Giac [F]	413
Mupad [F(-1)]	414

Optimal result

Integrand size = 8, antiderivative size = 105

$$\int e^{\operatorname{sech}^{-1}(ax^p)} dx = e^{\operatorname{sech}^{-1}(ax^p)} x + \frac{px^{1-p}}{a(1-p)} + \frac{px^{1-p} \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} \left(-1 + \frac{1}{p}\right), \frac{1+p}{2p}, a^2 x^{2p}\right)}{a(1-p)}$$

[Out] (1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))*x+p*x^(1-p)/a/(1-p)+p*x^(1-p)*hypergeom([1/2, -1/2+1/2/p], [1/2*(p+1)/p], a^2*x^(2*p))*(1/(1+a*x^p))^(1/2)*(1+a*x^p)^(1/2)/a/(1-p)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6465, 30, 265, 371}

$$\int e^{\operatorname{sech}^{-1}(ax^p)} dx = \frac{px^{1-p} \sqrt{\frac{1}{ax^p+1}} \sqrt{ax^p+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} \left(\frac{1}{p} - 1\right), \frac{p+1}{2p}, a^2 x^{2p}\right)}{a(1-p)} + \frac{px^{1-p}}{a(1-p)} + x e^{\operatorname{sech}^{-1}(ax^p)}$$

[In] Int[E^ArcSech[a*x^p], x]

[Out] E^ArcSech[a*x^p]*x + (p*x^(1-p))/(a*(1-p)) + (p*x^(1-p)*Sqrt[(1+a*x^p)^(-1)]*Sqrt[1+a*x^p]*Hypergeometric2F1[1/2, (-1+p^(-1))/2, (1+p)/(2*p), a^2*x^(2*p)])/(a*(1-p))

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 265

Int[((c_)*(x_))^(m_)*((a1_) + (b1_)*(x_)^(n_))^(p_)*((a2_) + (b2_)*(x_)^(n_))^(p_), x_Symbol] := Int[(c*x)^m*(a1*a2 + b1*b2*x^(2*n))^p, x] /; FreeQ[{a1, b1, a2, b2, c, m, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))

Rule 371

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1))) * Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 6465

Int[E^ArcSech[(a_)*(x_)^(p_)], x_Symbol] := Simp[x*E^ArcSech[a*x^p], x] + (Dist[p/a, Int[1/x^p, x], x] + Dist[p*(Sqrt[1 + a*x^p]/a)*Sqrt[1/(1 + a*x^p)], Int[1/(x^p*Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, p}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= e^{\text{sech}^{-1}(ax^p)} x + \frac{p \int x^{-p} dx}{a} + \frac{\left(p \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p}\right) \int \frac{x^{-p}}{\sqrt{1-ax^p}\sqrt{1+ax^p}} dx}{a} \\
 &= e^{\text{sech}^{-1}(ax^p)} x + \frac{px^{1-p}}{a(1-p)} + \frac{\left(p \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p}\right) \int \frac{x^{-p}}{\sqrt{1-a^2x^{2p}}} dx}{a} \\
 &= e^{\text{sech}^{-1}(ax^p)} x + \frac{px^{1-p}}{a(1-p)} \\
 &\quad + \frac{px^{1-p} \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(-1 + \frac{1}{p}\right), \frac{1+p}{2p}, a^2x^{2p}\right)}{a(1-p)}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.56

$$\int e^{\operatorname{sech}^{-1}(ax^p)} dx$$

$$= \frac{x^{1-p} \left(-1 - \sqrt{\frac{1-ax^p}{1+ax^p}} - ax^p \sqrt{\frac{1-ax^p}{1+ax^p}} + \frac{a^2 p x^{2p} \sqrt{\frac{1-ax^p}{1+ax^p}} \sqrt{1-a^2 x^{2p}} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+p}{2p}, \frac{1}{2}\left(3+\frac{1}{p}\right), a^2 x^{2p}\right)}{(1+p)(-1+ax^p)} \right)}{a(-1+p)}$$

[In] Integrate[E^ArcSech[a*x^p],x]

[Out] (x^(1 - p)*(-1 - Sqrt[(1 - a*x^p)/(1 + a*x^p)] - a*x^p*Sqrt[(1 - a*x^p)/(1 + a*x^p)] + (a^2*p*x^(2*p)*Sqrt[(1 - a*x^p)/(1 + a*x^p)]*Sqrt[1 - a^2*x^(2*p)])*Hypergeometric2F1[1/2, (1 + p)/(2*p), (3 + p^(-1))/2, a^2*x^(2*p)])/((1 + p)*(-1 + a*x^p)))/(a*(-1 + p))

Maple [F]

$$\int \left(\frac{x^{-p}}{a} + \sqrt{\frac{x^{-p}}{a} - 1} \sqrt{\frac{x^{-p}}{a} + 1} \right) dx$$

[In] int(1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2),x)

[Out] int(1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2),x)

Fricas [F(-2)]

Exception generated.

$$\int e^{\operatorname{sech}^{-1}(ax^p)} dx = \text{Exception raised: TypeError}$$

[In] integrate(1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

Sympy [F]

$$\int e^{\operatorname{sech}^{-1}(ax^p)} dx = \frac{\int x^{-p} dx + \int a \sqrt{-1 + \frac{x^{-p}}{a}} \sqrt{1 + \frac{x^{-p}}{a}} dx}{a}$$

[In] integrate(1/a/(x**p)+(1/a/(x**p)-1)**(1/2)*(1/a/(x**p)+1)**(1/2),x)

[Out] (Integral(x**(-p), x) + Integral(a*sqrt(-1 + 1/(a*x**p))*sqrt(1 + 1/(a*x**p)), x))/a

Maxima [F(-2)]

Exception generated.

$$\int e^{\operatorname{sech}^{-1}(ax^p)} dx = \text{Exception raised: ValueError}$$

[In] integrate(1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested a dditional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(-p>0)', see 'assume?' for more details)Is

Giac [F]

$$\int e^{\operatorname{sech}^{-1}(ax^p)} dx = \int \sqrt{\frac{1}{ax^p} + 1} \sqrt{\frac{1}{ax^p} - 1} + \frac{1}{ax^p} dx$$

[In] integrate(1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(1/(a*x^p) + 1)*sqrt(1/(a*x^p) - 1) + 1/(a*x^p), x)

Mupad [F(-1)]

Timed out.

$$\int e^{\operatorname{sech}^{-1}(ax^p)} dx = \int \sqrt{\frac{1}{ax^p} - 1} \sqrt{\frac{1}{ax^p} + 1} + \frac{1}{ax^p} dx$$

```
[In] int((1/(a*x^p) - 1)^(1/2)*(1/(a*x^p) + 1)^(1/2) + 1/(a*x^p), x)
```

```
[Out] int((1/(a*x^p) - 1)^(1/2)*(1/(a*x^p) + 1)^(1/2) + 1/(a*x^p), x)
```

3.63 $\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x} dx$

Optimal result	415
Rubi [A] (verified)	415
Mathematica [C] (verified)	417
Maple [C] (verified)	417
Fricas [A] (verification not implemented)	418
Sympy [F]	418
Maxima [F]	418
Giac [F]	419
Mupad [F(-1)]	419

Optimal result

Integrand size = 12, antiderivative size = 87

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x} dx = -\frac{x^{-p}}{ap} - \frac{x^{-p}\sqrt{1-ax^p}}{ap\sqrt{\frac{1}{1+ax^p}}} - \frac{\sqrt{\frac{1}{1+ax^p}}\sqrt{1+ax^p}\arcsin(ax^p)}{p}$$

[Out] $-1/a/p/(x^p) - (1-ax^p)^{(1/2)}/a/p/(x^p)/((1+(1+ax^p))^{(1/2)}) - \arcsin(ax^p) * (1/(1+ax^p))^{(1/2)} * (1+ax^p)^{(1/2)}/p$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.22, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6469, 265, 352, 248, 283, 222}

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x} dx = -\frac{x^{-p}\sqrt{\frac{1}{ax^p+1}}\sqrt{ax^p+1}\sqrt{1-a^2x^{2p}}}{ap} - \frac{x^{-p}}{ap} - \frac{\sqrt{\frac{1}{ax^p+1}}\sqrt{ax^p+1}\operatorname{csc}^{-1}\left(\frac{x^{-p}}{a}\right)}{p}$$

[In] Int[E^ArcSech[a*x^p]/x,x]

[Out] $-(1/(a*p*x^p)) - (\operatorname{Sqrt}[(1+a*x^p)^{-1}]*\operatorname{Sqrt}[1+a*x^p]*\operatorname{Sqrt}[1-a^2*x^{(2*p)}])/(a*p*x^p) - (\operatorname{Sqrt}[(1+a*x^p)^{-1}]*\operatorname{Sqrt}[1+a*x^p]*\operatorname{ArcCsc}[1/(a*x^p)]) / p$

Rule 222

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

Rule 248

$\text{Int}[(a_+) + (b_+)(x_+)^{(n_+)}]^{(p_+)}, x_Symbol] \text{ :> } -\text{Subst}[\text{Int}[(a + b/x^n)^p/x^2, x], x, 1/x] \text{ /; FreeQ}\{a, b, p\}, x\} \ \&\& \ \text{ILtQ}[n, 0]$

Rule 265

$\text{Int}[(c_+)(x_+)^{(m_+)}((a1_+) + (b1_+)(x_+)^{(n_+)})^{(p_+)}((a2_+) + (b2_+)(x_+)^{(n_+)})^{(p_+)}, x_Symbol] \text{ :> } \text{Int}[(c*x)^m*(a1*a2 + b1*b2*x^{(2*n)})^p, x] \text{ /; FreeQ}\{a1, b1, a2, b2, c, m, n, p\}, x\} \ \&\& \ \text{EqQ}[a2*b1 + a1*b2, 0] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{GtQ}[a1, 0] \ \&\& \ \text{GtQ}[a2, 0]))$

Rule 283

$\text{Int}[(c_+)(x_+)^{(m_+)}((a_+) + (b_+)(x_+)^{(n_+)})^{(p_+)}, x_Symbol] \text{ :> } \text{Simp}[(c*x)^{(m+1)}*((a + b*x^n)^p/(c^{n*(m+1)})), x] - \text{Dist}[b*n*(p/(c^n*(m+1))), \text{Int}[(c*x)^{(m+n)}*(a + b*x^n)^{(p-1)}, x], x] \text{ /; FreeQ}\{a, b, c\}, x\} \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ !\text{ILtQ}[(m+n*p+n+1)/n, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 352

$\text{Int}(x_+)^{(m_+)}((a_+) + (b_+)(x_+)^{(n_+)})^{(p_+)}, x_Symbol] \text{ :> } \text{Dist}[1/(m+1), \text{Subst}[\text{Int}[(a + b*x^{\text{Simplify}[n/(m+1)])^p, x], x, x^{(m+1)}], x] \text{ /; FreeQ}\{a, b, m, n, p\}, x\} \ \&\& \ \text{IntegerQ}[\text{Simplify}[n/(m+1)]] \ \&\& \ !\text{IntegerQ}[n]$

Rule 6469

$\text{Int}[E^{\text{ArcSech}}[(a_+)(x_+)^{(p_+)})]/(x_+), x_Symbol] \text{ :> } -\text{Simp}[(a*p*x^p)^{-1}, x] + \text{Dist}[(\text{Sqrt}[1 + a*x^p]/a)*\text{Sqrt}[1/(1 + a*x^p)], \text{Int}[\text{Sqrt}[1 + a*x^p]*(\text{Sqrt}[1 - a*x^p]/x^{(p+1)}), x], x] \text{ /; FreeQ}\{a, p\}, x]$

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{x^{-p}}{ap} + \frac{\left(\sqrt{\frac{1}{1+ax^p}}\sqrt{1+ax^p}\right) \int x^{-1-p}\sqrt{1-ax^p}\sqrt{1+ax^p} dx}{a} \\ &= -\frac{x^{-p}}{ap} + \frac{\left(\sqrt{\frac{1}{1+ax^p}}\sqrt{1+ax^p}\right) \int x^{-1-p}\sqrt{1-a^2x^{2p}} dx}{a} \\ &= -\frac{x^{-p}}{ap} - \frac{\left(\sqrt{\frac{1}{1+ax^p}}\sqrt{1+ax^p}\right) \text{Subst}\left(\int \sqrt{1-\frac{a^2}{x^2}} dx, x, x^{-p}\right)}{ap} \\ &= -\frac{x^{-p}}{ap} + \frac{\left(\sqrt{\frac{1}{1+ax^p}}\sqrt{1+ax^p}\right) \text{Subst}\left(\int \frac{\sqrt{1-a^2x^2}}{x^2} dx, x, x^p\right)}{ap} \end{aligned}$$

$$= \frac{x^{-p}}{ap} - \frac{x^{-p} \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p} \sqrt{1-a^2x^{2p}}}{ap} - \frac{\left(a \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p}\right) \text{Subst}\left(\int \frac{1}{\sqrt{1-a^2x^2}} dx, x, x^p\right)}{p}$$

$$= \frac{x^{-p}}{ap} - \frac{x^{-p} \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p} \sqrt{1-a^2x^{2p}}}{ap} - \frac{\sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p} \arcsin(ax^p)}{p}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.10

$$\int \frac{e^{\text{sech}^{-1}(ax^p)}}{x} dx = -\frac{i\left(-ix^{-p} - i(a+x^{-p})\sqrt{\frac{1-ax^p}{1+ax^p}} + a \log\left(-2iax^p + 2\sqrt{\frac{1-ax^p}{1+ax^p}}(1+ax^p)\right)\right)}{ap}$$

[In] Integrate[E^ArcSech[a*x^p]/x,x]

[Out] ((-I)*((-I)/x^p - I*(a + x^(-p))*Sqrt[(1 - a*x^p)/(1 + a*x^p)] + a*Log[(-2*I)*a*x^p + 2*Sqrt[(1 - a*x^p)/(1 + a*x^p)]*(1 + a*x^p)]))/(a*p)

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 1.17 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.33

method	result	size
derivativedivides	$\frac{\sqrt{-\frac{(ax^p-1)x^{-p}}{a}} \sqrt{\frac{(1+ax^p)x^{-p}}{a}} \left(\arctan\left(\frac{\text{csgn}(a)ax^p}{\sqrt{-a^2x^{2p}+1}}\right)ax^p + \sqrt{-a^2x^{2p}+1} \text{csgn}(a)\right) \text{csgn}(a) - \frac{x^{-p}}{a}}{\sqrt{-a^2x^{2p}+1}}}{p}$	116
default	$\frac{\sqrt{-\frac{(ax^p-1)x^{-p}}{a}} \sqrt{\frac{(1+ax^p)x^{-p}}{a}} \left(\arctan\left(\frac{\text{csgn}(a)ax^p}{\sqrt{-a^2x^{2p}+1}}\right)ax^p + \sqrt{-a^2x^{2p}+1} \text{csgn}(a)\right) \text{csgn}(a) - \frac{x^{-p}}{a}}{\sqrt{-a^2x^{2p}+1}}}{p}$	116

[In] int((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))/x,x,method=_RETURNV ERBOSE)

[Out] 1/p*(-(-(a*x^p-1)/a/(x^p))^(1/2)*((1+a*x^p)/a/(x^p))^(1/2)*(arctan(csgn(a)*a*x^p/(-(x^p)^2*a^2+1)^(1/2))*a*x^p+(-(x^p)^2*a^2+1)^(1/2)*csgn(a))*csgn(a)/(-(x^p)^2*a^2+1)^(1/2)-1/a/(x^p))

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.17

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x} dx = -\frac{ax^p \sqrt{\frac{ax^p+1}{ax^p}} \sqrt{-\frac{ax^p-1}{ax^p}} - ax^p \arctan\left(\sqrt{\frac{ax^p+1}{ax^p}} \sqrt{-\frac{ax^p-1}{ax^p}}\right) + 1}{apx^p}$$

[In] integrate((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))/x,x, algorithm="fricas")

[Out] -(a*x^p*sqrt((a*x^p + 1)/(a*x^p))*sqrt(-(a*x^p - 1)/(a*x^p)) - a*x^p*arctan(sqrt((a*x^p + 1)/(a*x^p))*sqrt(-(a*x^p - 1)/(a*x^p))) + 1)/(a*p*x^p)

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x} dx = \int \frac{x^{-p}}{x} dx + \int \frac{a\sqrt{-1+\frac{x^{-p}}{a}}\sqrt{1+\frac{x^{-p}}{a}}}{x} dx$$

[In] integrate((1/a/(x**p)+(1/a/(x**p)-1)**(1/2)*(1/a/(x**p)+1)**(1/2))/x,x)

[Out] (Integral(1/(x*x**p), x) + Integral(a*sqrt(-1 + 1/(a*x**p))*sqrt(1 + 1/(a*x**p))/x, x))/a

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x} dx = \int \frac{\sqrt{\frac{1}{ax^p} + 1} \sqrt{\frac{1}{ax^p} - 1} + \frac{1}{ax^p}}{x} dx$$

[In] integrate((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))/x,x, algorithm="maxima")

[Out] integrate(sqrt(a*x^p + 1)*sqrt(-a*x^p + 1)/(x*x^p), x)/a - 1/(a*p*x^p)

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x} dx = \int \frac{\sqrt{\frac{1}{ax^p} + 1} \sqrt{\frac{1}{ax^p} - 1} + \frac{1}{ax^p}}{x} dx$$

[In] integrate((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))/x,x, algorithm m="giac")

[Out] integrate((sqrt(1/(a*x^p) + 1)*sqrt(1/(a*x^p) - 1) + 1/(a*x^p))/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x} dx = \int \frac{\sqrt{\frac{1}{ax^p} - 1} \sqrt{\frac{1}{ax^p} + 1} + \frac{1}{ax^p}}{x} dx$$

[In] int(((1/(a*x^p) - 1)^(1/2)*(1/(a*x^p) + 1)^(1/2) + 1/(a*x^p))/x,x)

[Out] int(((1/(a*x^p) - 1)^(1/2)*(1/(a*x^p) + 1)^(1/2) + 1/(a*x^p))/x, x)

3.64 $\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x^2} dx$

Optimal result	420
Rubi [A] (verified)	420
Mathematica [A] (verified)	422
Maple [F]	422
Fricas [F(-2)]	422
Sympy [F]	423
Maxima [F]	423
Giac [F]	423
Mupad [F(-1)]	423

Optimal result

Integrand size = 12, antiderivative size = 107

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x^2} dx = -\frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x} + \frac{px^{-1-p}}{a(1+p)} + \frac{px^{-1-p} \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -\frac{1+p}{2p}, -\frac{1-p}{2p}, a^2x^{2p}\right)}{a(1+p)}$$

[Out] $-(1/a/(x^p)+(1/a/(x^p)-1)^{(1/2)}*(1/a/(x^p)+1)^{(1/2)})/x+p*x^{(-1-p)}/a/(p+1)+p*x^{(-1-p)}*hypergeom([1/2, 1/2*(-1-p)/p], [1/2*(-1+p)/p], a^2*x^{(2*p)})*(1/(1+a*x^p))^{(1/2)}*(1+a*x^p)^{(1/2)}/a/(p+1)$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6470, 30, 265, 371}

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x^2} dx = \frac{px^{-p-1} \sqrt{\frac{1}{ax^p+1}} \sqrt{ax^p+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -\frac{p+1}{2p}, -\frac{1-p}{2p}, a^2x^{2p}\right)}{a(p+1)} + \frac{px^{-p-1}}{a(p+1)} - \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcSech}[a*x^p]}/x^2, x]$

[Out] $-(E^{\operatorname{ArcSech}[a*x^p]}/x) + (p*x^{(-1-p)})/(a*(1+p)) + (p*x^{(-1-p)}*\operatorname{Sqrt}[(1+a*x^p)^{-1}]*\operatorname{Sqrt}[1+a*x^p]*\operatorname{Hypergeometric2F1}[1/2, -1/2*(1+p)/p, -1/2*(1-p)/p, a^2*x^{(2*p)}])/(a*(1+p))$

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 265

Int[((c_)*(x_))^(m_)*((a1_) + (b1_)*(x_)^(n_))^(p_)*((a2_) + (b2_)*(x_)^(n_))^(p_), x_Symbol] := Int[(c*x)^m*(a1*a2 + b1*b2*x^(2*n))^p, x] /; FreeQ[{a1, b1, a2, b2, c, m, n, p}, x] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[p] || (GtQ[a1, 0] && GtQ[a2, 0]))

Rule 371

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 6470

Int[E^ArcSech[(a_)*(x_)^(p_)]*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*(E^ArcSech[a*x^p]/(m + 1)), x] + (Dist[p/(a*(m + 1)), Int[x^(m - p), x], x] + Dist[p*(Sqrt[1 + a*x^p]/(a*(m + 1)))*Sqrt[1/(1 + a*x^p)], Int[x^(m - p)/(Sqrt[1 + a*x^p]*Sqrt[1 - a*x^p]), x], x]) /; FreeQ[{a, m, p}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x} - \frac{p \int x^{-2-p} dx}{a} - \frac{\left(p \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p}\right) \int \frac{x^{-2-p}}{\sqrt{1-ax^p}\sqrt{1+ax^p}} dx}{a} \\
 &= -\frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x} + \frac{px^{-1-p}}{a(1+p)} - \frac{\left(p \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p}\right) \int \frac{x^{-2-p}}{\sqrt{1-a^2x^{2p}}} dx}{a} \\
 &= -\frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x} + \frac{px^{-1-p}}{a(1+p)} \\
 &\quad + \frac{px^{-1-p} \sqrt{\frac{1}{1+ax^p}} \sqrt{1+ax^p} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -\frac{1+p}{2p}, -\frac{1-p}{2p}, a^2x^{2p}\right)}{a(1+p)}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.55

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x^2} dx$$

$$= \frac{x^{-1-p} \left(-1 - \sqrt{\frac{1-ax^p}{1+ax^p}} - ax^p \sqrt{\frac{1-ax^p}{1+ax^p}} + \frac{a^2 p x^{2p} \sqrt{\frac{1-ax^p}{1+ax^p}} \sqrt{1-a^2 x^{2p}} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{-1+p}{2p}, \frac{3}{2} - \frac{1}{2p}, a^2 x^{2p}\right)}{(-1+p)(-1+ax^p)} \right)}{a(1+p)}$$

[In] Integrate[E^ArcSech[a*x^p]/x^2,x]

[Out] (x^(-1 - p))*(-1 - Sqrt[(1 - a*x^p)/(1 + a*x^p)]) - a*x^p*Sqrt[(1 - a*x^p)/(1 + a*x^p)] + (a^2*p*x^(2*p)*Sqrt[(1 - a*x^p)/(1 + a*x^p)]*Sqrt[1 - a^2*x^(2*p)]*Hypergeometric2F1[1/2, (-1 + p)/(2*p), 3/2 - 1/(2*p), a^2*x^(2*p)])/((-1 + p)*(-1 + a*x^p)))/(a*(1 + p))

Maple [F]

$$\int \frac{\frac{x^{-p}}{a} + \sqrt{\frac{x^{-p}}{a} - 1} \sqrt{\frac{x^{-p}}{a} + 1}}{x^2} dx$$

[In] int((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))/x^2,x)

[Out] int((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))/x^2,x)

Fricas [F(-2)]

Exception generated.

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x^2} dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))/x^2,x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x^2} dx = \int \frac{x^{-p}}{x^2} dx + \int \frac{a\sqrt{-1+\frac{x^{-p}}{a}}\sqrt{1+\frac{x^{-p}}{a}}}{x^2} dx$$

[In] integrate((1/a/(x**p)+(1/a/(x**p)-1)**(1/2)*(1/a/(x**p)+1)**(1/2))/x**2,x)

[Out] (Integral(1/(x**2*x**p), x) + Integral(a*sqrt(-1 + 1/(a*x**p))*sqrt(1 + 1/(a*x**p))/x**2, x))/a

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x^2} dx = \int \frac{\sqrt{\frac{1}{ax^p} + 1}\sqrt{\frac{1}{ax^p} - 1 + \frac{1}{ax^p}}}{x^2} dx$$

[In] integrate((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))/x^2,x, algorithm="maxima")

[Out] integrate(sqrt(a*x^p + 1)*sqrt(-a*x^p + 1)/(x^2*x^p), x)/a - x^(-p - 1)/(a*(p + 1))

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x^2} dx = \int \frac{\sqrt{\frac{1}{ax^p} + 1}\sqrt{\frac{1}{ax^p} - 1 + \frac{1}{ax^p}}}{x^2} dx$$

[In] integrate((1/a/(x^p)+(1/a/(x^p)-1)^(1/2)*(1/a/(x^p)+1)^(1/2))/x^2,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x^p) + 1)*sqrt(1/(a*x^p) - 1) + 1/(a*x^p))/x^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\operatorname{sech}^{-1}(ax^p)}}{x^2} dx = \int \frac{\sqrt{\frac{1}{ax^p} - 1}\sqrt{\frac{1}{ax^p} + 1 + \frac{1}{ax^p}}}{x^2} dx$$

[In] int(((1/(a*x^p) - 1)^(1/2)*(1/(a*x^p) + 1)^(1/2) + 1/(a*x^p))/x^2,x)

[Out] int(((1/(a*x^p) - 1)^(1/2)*(1/(a*x^p) + 1)^(1/2) + 1/(a*x^p))/x^2, x)

3.65 $\int e^{2\operatorname{sech}^{-1}(ax)} x^4 dx$

Optimal result	424
Rubi [A] (verified)	424
Mathematica [C] (verified)	427
Maple [C] (verified)	427
Fricas [A] (verification not implemented)	428
Sympy [F]	428
Maxima [F]	428
Giac [F(-2)]	429
Mupad [B] (verification not implemented)	429

Optimal result

Integrand size = 12, antiderivative size = 203

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^4 dx = \frac{5\sqrt{\frac{1-ax}{1+ax}}(1+ax)^2}{4a^5} + \frac{(1-ax)(1+ax)^4}{5a^5} + \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^4(5-6\sqrt{\frac{1-ax}{1+ax}})}{10a^5} + \frac{(1+ax)(4-\sqrt{\frac{1-ax}{1+ax}})}{4a^5} - \frac{(1+ax)^3(4+45\sqrt{\frac{1-ax}{1+ax}})}{30a^5} - \frac{\arctan\left(\sqrt{\frac{1-ax}{1+ax}}\right)}{2a^5}$$

[Out] 1/5*(-a*x+1)*(a*x+1)^4/a^5-1/2*arctan(((a*x+1)/(a*x+1))^(1/2))/a^5+1/4*(a*x+1)*(4-((a*x+1)/(a*x+1))^(1/2))/a^5+5/4*(a*x+1)^2*((a*x+1)/(a*x+1))^(1/2)/a^5+1/10*(a*x+1)^4*(5-6*((a*x+1)/(a*x+1))^(1/2))*((a*x+1)/(a*x+1))^(1/2)/a^5-1/30*(a*x+1)^3*(4+45*((a*x+1)/(a*x+1))^(1/2))/a^5

Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6472, 1818, 1825, 1828, 653, 209}

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^4 dx = -\frac{\arctan\left(\sqrt{\frac{1-ax}{ax+1}}\right)}{2a^5} + \frac{(1-ax)(ax+1)^4}{5a^5} + \frac{\sqrt{\frac{1-ax}{ax+1}}(5-6\sqrt{\frac{1-ax}{ax+1}})(ax+1)^4}{10a^5} - \frac{(45\sqrt{\frac{1-ax}{ax+1}}+4)(ax+1)^3}{30a^5} + \frac{5\sqrt{\frac{1-ax}{ax+1}}(ax+1)^2}{4a^5} + \frac{(4-\sqrt{\frac{1-ax}{ax+1}})(ax+1)}{4a^5}$$

[In] Int[E^(2*ArcSech[a*x])*x^4,x]

[Out] (5*sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2)/(4*a^5) + ((1 - a*x)*(1 + a*x)^4)/(5*a^5) + (sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^4*(5 - 6*sqrt[(1 - a*x)/(1 + a*x)]))/(10*a^5) + ((1 + a*x)*(4 - sqrt[(1 - a*x)/(1 + a*x)]))/(4*a^5) - ((1 + a*x)^3*(4 + 45*sqrt[(1 - a*x)/(1 + a*x)]))/(30*a^5) - ArcTan[sqrt[(1 - a*x)/(1 + a*x)]]/(2*a^5)

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 653

Int[((d_) + (e_)*(x_))*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[((a*e - c*d*x)/(2*a*c*(p + 1))*(a + c*x^2)^(p + 1), x] + Dist[d*((2*p + 3)/(2*a*(p + 1))), Int[(a + c*x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e}, x] && LtQ[p, -1] && NeQ[p, -3/2]

Rule 1818

Int[(Pq_)*((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(c*x)^m*(a + b*x^2)^(p + 1)*((a*g - b*f*x)/(2*a*b*(p + 1))), x] + Dist[c/(2*a*b*(p + 1)), Int[(c*x)^(m - 1)*(a + b*x^2)^(p + 1)*ExpandToSum[2*a*b*(p + 1)*x*Q - a*g*m + b*f*(m + 2*p + 3)*x, x], x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x] && LtQ[p, -1] && GtQ[m, 0]

Rule 1825

Int[(Pq_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[x*PolynomialQuotient[Pq, x, x]*(a + b*x^2)^p, x] /; FreeQ[{a, b, p}, x] && PolyQ[Pq, x] && EqQ[Coeff[Pq, x, 0], 0] && !MatchQ[Pq, x^(m_)*(u_)] /; IntegerQ[m]

Rule 1828

Int[(Pq_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(a*g - b*f*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Dist[1/(2*a*(p + 1)), Int[(a + b*x^2)^(p + 1)*ExpandToSum[2*a*(p + 1)*Q + f*(2*p + 3), x], x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && LtQ[p, -1]

Rule 6472

Int[E^(ArcSech[u_]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && Integer Q[n]

Rubi steps

$$\begin{aligned}
\text{integral} &= \int x^4 \left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax} \right)^2 dx \\
&= -\frac{4\text{Subst}\left(\int \frac{(-1+x)^2 x(1+x)^6}{(1+x^2)^6} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a^5} \\
&= \frac{(1-ax)(1+ax)^4}{5a^5} + \frac{2\text{Subst}\left(\int \frac{-42x-40x^2+130x^3+80x^4-30x^5-40x^6-10x^7}{(1+x^2)^5} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{5a^5} \\
&= \frac{(1-ax)(1+ax)^4}{5a^5} + \frac{2\text{Subst}\left(\int \frac{x(-42-40x+130x^2+80x^3-30x^4-40x^5-10x^6)}{(1+x^2)^5} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{5a^5} \\
&= \frac{(1-ax)(1+ax)^4}{5a^5} + \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^4 \left(5 - 6\sqrt{\frac{1-ax}{1+ax}}\right)}{10a^5} \\
&\quad - \frac{\text{Subst}\left(\int \frac{160-48x-960x^2+160x^3+320x^4+80x^5}{(1+x^2)^4} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{20a^5} \\
&= \frac{(1-ax)(1+ax)^4}{5a^5} + \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^4 \left(5 - 6\sqrt{\frac{1-ax}{1+ax}}\right)}{10a^5} \\
&\quad - \frac{(1+ax)^3 \left(4 + 45\sqrt{\frac{1-ax}{1+ax}}\right)}{30a^5} + \frac{\text{Subst}\left(\int \frac{480-480x-1920x^2-480x^3}{(1+x^2)^3} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{120a^5} \\
&= \frac{5\sqrt{\frac{1-ax}{1+ax}}(1+ax)^2}{4a^5} + \frac{(1-ax)(1+ax)^4}{5a^5} + \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^4 \left(5 - 6\sqrt{\frac{1-ax}{1+ax}}\right)}{10a^5} \\
&\quad - \frac{(1+ax)^3 \left(4 + 45\sqrt{\frac{1-ax}{1+ax}}\right)}{30a^5} - \frac{\text{Subst}\left(\int \frac{480+1920x}{(1+x^2)^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{480a^5} \\
&= \frac{5\sqrt{\frac{1-ax}{1+ax}}(1+ax)^2}{4a^5} + \frac{(1-ax)(1+ax)^4}{5a^5} + \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^4 \left(5 - 6\sqrt{\frac{1-ax}{1+ax}}\right)}{10a^5} \\
&\quad + \frac{(1+ax) \left(4 - \sqrt{\frac{1-ax}{1+ax}}\right)}{4a^5} - \frac{(1+ax)^3 \left(4 + 45\sqrt{\frac{1-ax}{1+ax}}\right)}{30a^5} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{2a^5}
\end{aligned}$$

$$= \frac{5\sqrt{\frac{1-ax}{1+ax}}(1+ax)^2}{4a^5} + \frac{(1-ax)(1+ax)^4}{5a^5} + \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^4 \left(5 - 6\sqrt{\frac{1-ax}{1+ax}}\right)}{10a^5} \\ + \frac{(1+ax) \left(4 - \sqrt{\frac{1-ax}{1+ax}}\right)}{4a^5} - \frac{(1+ax)^3 \left(4 + 45\sqrt{\frac{1-ax}{1+ax}}\right)}{30a^5} - \frac{\arctan\left(\sqrt{\frac{1-ax}{1+ax}}\right)}{2a^5}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.16 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.52

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^4 dx \\ = \frac{40a^3x^3 - 12a^5x^5 - 15a\sqrt{\frac{1-ax}{1+ax}}(x + ax^2 - 2a^2x^3 - 2a^3x^4) + 15i \log\left(-2iax + 2\sqrt{\frac{1-ax}{1+ax}}(1+ax)\right)}{60a^5}$$

[In] Integrate[E^(2*ArcSech[a*x])*x^4,x]

[Out] (40*a^3*x^3 - 12*a^5*x^5 - 15*a*Sqrt[(1 - a*x)/(1 + a*x)]*(x + a*x^2 - 2*a^2*x^3 - 2*a^3*x^4) + (15*I)*Log[(-2*I)*a*x + 2*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)]/(60*a^5)

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.06 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.67

method	result
default	$\frac{-\frac{1}{5}a^2x^5 + \frac{1}{3}x^3}{a^2} + \frac{\sqrt{\frac{ax+1}{ax}} x \sqrt{-\frac{ax-1}{ax}} \left(2 \operatorname{csgn}(a) a^3 x^3 \sqrt{-a^2x^2+1} - \sqrt{-a^2x^2+1} x \operatorname{csgn}(a) a + \arctan\left(\frac{\operatorname{csgn}(a)ax}{\sqrt{-a^2x^2+1}}\right)\right) \operatorname{csgn}(a)}{4a^4\sqrt{-a^2x^2+1}} + \frac{x^3}{3a^2}$

[In] int((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))^2*x^4,x,method=_RETURNVERBOSE)

[Out] 1/a^2*(-1/5*a^2*x^5+1/3*x^3)+1/4/a^4*((a*x+1)/a/x)^(1/2)*x*(-(a*x-1)/a/x)^(1/2)*(2*csgn(a)*a^3*x^3*(-a^2*x^2+1)^(1/2)-(-a^2*x^2+1)^(1/2)*x*csgn(a)*a+a*rctan(csgn(a)*a*x/(-a^2*x^2+1)^(1/2)))*csgn(a)/(-a^2*x^2+1)^(1/2)+1/3*x^3/a^2

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.51

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^4 dx$$

$$= -\frac{12 a^5 x^5 - 40 a^3 x^3 - 15 (2 a^4 x^4 - a^2 x^2) \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} + 15 \arctan\left(\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}}\right)}{60 a^5}$$

```
[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2*x^4,x, algorithm="fricas")
```

```
[Out] -1/60*(12*a^5*x^5 - 40*a^3*x^3 - 15*(2*a^4*x^4 - a^2*x^2)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + 15*arctan(sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x))))/a^5
```

Sympy [F]

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^4 dx = \frac{\int 2x^2 dx + \int (-a^2 x^4) dx + \int 2ax^3 \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} dx}{a^2}$$

```
[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))**2*x**4,x)
```

```
[Out] (Integral(2*x**2, x) + Integral(-a**2*x**4, x) + Integral(2*a*x**3*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)), x))/a**2
```

Maxima [F]

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^4 dx = \int x^4 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} \right)^2 dx$$

```
[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2*x^4,x, algorithm="maxima")
```

```
[Out] 2/3*x^3/a^2 + 2*integrate(sqrt(a*x + 1)*sqrt(-a*x + 1)*x^2, x)/a^2 - integrate(x^4, x)
```

Giac [F(-2)]

Exception generated.

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^4 dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2*x^4,x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{-1,[0,4,0,6,0,0]%%}+%%{1,[0,2,4,4,0,0]%%}+%%{1,[0,2,0,4,0,0]%%}

Mupad [B] (verification not implemented)

Time = 25.51 (sec) , antiderivative size = 808, normalized size of antiderivative = 3.98

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^4 dx =$$

$$\frac{\frac{1i}{512a^5} - \frac{(\sqrt{\frac{1}{ax}-1-i})^2 3i}{64a^5 (\sqrt{\frac{1}{ax}+1-1})^2} - \frac{(\sqrt{\frac{1}{ax}-1-i})^4 53i}{256a^5 (\sqrt{\frac{1}{ax}+1-1})^4} + \frac{(\sqrt{\frac{1}{ax}-1-i})^6 87i}{128a^5 (\sqrt{\frac{1}{ax}+1-1})^6} + \frac{(\sqrt{\frac{1}{ax}-1-i})^8 657i}{512a^5 (\sqrt{\frac{1}{ax}+1-1})^8} + \frac{(\sqrt{\frac{1}{ax}-1-i})^{10} 121i}{128a^5 (\sqrt{\frac{1}{ax}+1-1})^{10}}}{\frac{(\sqrt{\frac{1}{ax}-1-i})^4}{(\sqrt{\frac{1}{ax}+1-1})^4} + \frac{4(\sqrt{\frac{1}{ax}-1-i})^6}{(\sqrt{\frac{1}{ax}+1-1})^6} + \frac{6(\sqrt{\frac{1}{ax}-1-i})^8}{(\sqrt{\frac{1}{ax}+1-1})^8} + \frac{4(\sqrt{\frac{1}{ax}-1-i})^{10}}{(\sqrt{\frac{1}{ax}+1-1})^{10}} + \frac{(\sqrt{\frac{1}{ax}-1-i})^{12}}{(\sqrt{\frac{1}{ax}+1-1})^{12}}}$$

$$- \frac{\frac{1i}{16a^5} + \frac{(\sqrt{\frac{1}{ax}-1-i})^2 1i}{8a^5 (\sqrt{\frac{1}{ax}+1-1})^2} - \frac{(\sqrt{\frac{1}{ax}-1-i})^4 15i}{16a^5 (\sqrt{\frac{1}{ax}+1-1})^4}}{\frac{(\sqrt{\frac{1}{ax}-1-i})^2}{(\sqrt{\frac{1}{ax}+1-1})^2} + \frac{2(\sqrt{\frac{1}{ax}-1-i})^4}{(\sqrt{\frac{1}{ax}+1-1})^4} + \frac{(\sqrt{\frac{1}{ax}-1-i})^6}{(\sqrt{\frac{1}{ax}+1-1})^6}} - \frac{x^5 \left(\frac{a^2}{5} - \frac{2}{3x^2} \right)}{a^2}$$

$$- \frac{\ln \left(\frac{a\sqrt{\frac{1}{ax}+1-\frac{1}{x}} + a\sqrt{\frac{1}{ax}-1-i}}{2a-2a\sqrt{\frac{1}{ax}+1+\frac{1}{x}}} \right) 3i}{4a^5} - \frac{\ln \left(\frac{\sqrt{\frac{1}{ax}-1-i}}{\sqrt{\frac{1}{ax}+1-1}} \right) 1i}{4a^5} + \frac{\ln \left(\frac{2a\sqrt{\frac{a+\frac{1}{x}}{a}-\frac{2}{x}} + a\sqrt{\frac{a-\frac{1}{x}}{a} 2i}}{2a+\frac{1}{x}-2a\sqrt{\frac{a+\frac{1}{x}}{a}}} \right) 1i}{a^5}$$

$$- \frac{(\sqrt{\frac{1}{ax}-1-i})^2 1i}{128a^5 (\sqrt{\frac{1}{ax}+1-1})^2} - \frac{(\sqrt{\frac{1}{ax}-1-i})^4 1i}{512a^5 (\sqrt{\frac{1}{ax}+1-1})^4}$$

[In] int(x^4*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))^2,x)

[Out] (log((a*(-(a - 1/x)/a)^(1/2)*2i - 2/x + 2*a*((a + 1/x)/a)^(1/2))/(2*a + 1/x - 2*a*((a + 1/x)/a)^(1/2)))*1i)/a^5 - (1i/(512*a^5) - (((1/(a*x) - 1)^(1/2) - 1i)^2*3i)/(64*a^5*((1/(a*x) + 1)^(1/2) - 1)^2) - (((1/(a*x) - 1)^(1/2)

$$\begin{aligned}
& - 1i)^4 * 53i) / (256 * a^5 * ((1/(a*x) + 1)^{1/2} - 1)^4) + (((1/(a*x) - 1)^{1/2} \\
& - 1i)^6 * 87i) / (128 * a^5 * ((1/(a*x) + 1)^{1/2} - 1)^6) + (((1/(a*x) - 1)^{1/2} \\
& - 1i)^8 * 657i) / (512 * a^5 * ((1/(a*x) + 1)^{1/2} - 1)^8) + (((1/(a*x) - 1)^{1/2} \\
& - 1i)^{10} * 121i) / (128 * a^5 * ((1/(a*x) + 1)^{1/2} - 1)^{10}) / (((1/(a*x) - 1)^{1/2} \\
& - 1i)^4 / ((1/(a*x) + 1)^{1/2} - 1)^4 + (4 * ((1/(a*x) - 1)^{1/2} - 1i)^6) / (\\
& (1/(a*x) + 1)^{1/2} - 1)^6 + (6 * ((1/(a*x) - 1)^{1/2} - 1i)^8) / ((1/(a*x) + 1 \\
&)^{1/2} - 1)^8 + (4 * ((1/(a*x) - 1)^{1/2} - 1i)^{10}) / ((1/(a*x) + 1)^{1/2} - 1 \\
&)^{10} + ((1/(a*x) - 1)^{1/2} - 1i)^{12} / ((1/(a*x) + 1)^{1/2} - 1)^{12} - (\log((\\
& (1/(a*x) - 1)^{1/2} - 1i) / ((1/(a*x) + 1)^{1/2} - 1)) * 1i) / (4 * a^5) - (1i / (16 * \\
& a^5) + (((1/(a*x) - 1)^{1/2} - 1i)^2 * 1i) / (8 * a^5 * ((1/(a*x) + 1)^{1/2} - 1)^2 \\
&) - (((1/(a*x) - 1)^{1/2} - 1i)^4 * 15i) / (16 * a^5 * ((1/(a*x) + 1)^{1/2} - 1)^4 \\
&) / (((1/(a*x) - 1)^{1/2} - 1i)^2 / ((1/(a*x) + 1)^{1/2} - 1)^2 + (2 * ((1/(a*x) \\
& - 1)^{1/2} - 1i)^4) / ((1/(a*x) + 1)^{1/2} - 1)^4 + ((1/(a*x) - 1)^{1/2} - 1i \\
&)^6 / ((1/(a*x) + 1)^{1/2} - 1)^6) - (\log((a * (1/(a*x) - 1)^{1/2} * 1i + a * (1/(a \\
& * x) + 1)^{1/2} - 1/x) / (2 * a - 2 * a * (1/(a*x) + 1)^{1/2} + 1/x)) * 3i) / (4 * a^5) - \\
& (((1/(a*x) - 1)^{1/2} - 1i)^2 * 1i) / (128 * a^5 * ((1/(a*x) + 1)^{1/2} - 1)^2) - (\\
& ((1/(a*x) - 1)^{1/2} - 1i)^4 * 1i) / (512 * a^5 * ((1/(a*x) + 1)^{1/2} - 1)^4) - (x \\
& ^5 * (a^{2/5} - 2 / (3 * x^2))) / a^2
\end{aligned}$$

3.66 $\int e^{2\operatorname{sech}^{-1}(ax)} x^3 dx$

Optimal result	431
Rubi [A] (verified)	431
Mathematica [A] (verified)	433
Maple [A] (verified)	434
Fricas [A] (verification not implemented)	434
Sympy [F]	434
Maxima [A] (verification not implemented)	435
Giac [F]	435
Mupad [B] (verification not implemented)	435

Optimal result

Integrand size = 12, antiderivative size = 117

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^3 dx = -\frac{x}{a^3} + \frac{(1-ax)(1+ax)^3}{4a^4} + \frac{(1+ax)^2 \left(3 - 8\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^4} + \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^3 \left(4 - 3\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^4}$$

[Out] $-x/a^3 + 1/4*(-a*x+1)*(a*x+1)^3/a^4 + 1/6*(a*x+1)^2*(3-8*((-a*x+1)/(a*x+1))^(1/2))/a^4 + 1/6*(a*x+1)^3*(4-3*((-a*x+1)/(a*x+1))^(1/2))*((-a*x+1)/(a*x+1))^(1/2)/a^4$

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6472, 1818, 1825, 1828, 12, 267}

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^3 dx = \frac{(1-ax)(ax+1)^3}{4a^4} + \frac{\sqrt{\frac{1-ax}{ax+1}} \left(4 - 3\sqrt{\frac{1-ax}{ax+1}}\right) (ax+1)^3}{6a^4} + \frac{\left(3 - 8\sqrt{\frac{1-ax}{ax+1}}\right) (ax+1)^2}{6a^4} - \frac{x}{a^3}$$

[In] $\text{Int}[E^{(2*\text{ArcSech}[a*x])}*x^3,x]$

[Out] $-(x/a^3) + ((1-a*x)*(1+a*x)^3)/(4*a^4) + ((1+a*x)^2*(3-8*\text{Sqrt}[(1-a*x)/(1+a*x]]))/(6*a^4) + (\text{Sqrt}[(1-a*x)/(1+a*x)]*(1+a*x)^3*(4-3*\text{Sqrt}[(1-a*x)/(1+a*x]]))/(6*a^4)$

Rule 12

```
Int[(a_)*(u_), x_Symbol] :=> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 267

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :=> Simp[(a + b*x^n)^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]
```

Rule 1818

```
Int[(Pq_)*((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :=> With[{Q = PolynomialQuotient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(c*x)^m*(a + b*x^2)^(p + 1)*((a*g - b*f*x)/(2*a*b*(p + 1))), x] + Dist[c/(2*a*b*(p + 1)), Int[(c*x)^(m - 1)*(a + b*x^2)^(p + 1)*ExpandToSum[2*a*b*(p + 1)*x*Q - a*g*m + b*f*(m + 2*p + 3)*x, x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x] && LtQ[p, -1] && GtQ[m, 0]
```

Rule 1825

```
Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :=> Int[x*PolynomialQuotient[Pq, x, x]*(a + b*x^2)^p, x] /; FreeQ[{a, b, p}, x] && PolyQ[Pq, x] && EqQ[Coeff[Pq, x, 0], 0] && !MatchQ[Pq, x^(m_.)*(u_.)] /; IntegerQ[m]
```

Rule 1828

```
Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :=> With[{Q = PolynomialQuotient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(a*g - b*f*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Dist[1/(2*a*(p + 1)), Int[(a + b*x^2)^(p + 1)*ExpandToSum[2*a*(p + 1)*Q + f*(2*p + 3), x], x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && LtQ[p, -1]
```

Rule 6472

```
Int[E^(ArcSech[u_]*(n_.))*(x_)^(m_.), x_Symbol] :=> Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && IntegerQ[n]
```

Rubi steps

$$\text{integral} = \int x^3 \left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax} \right)^2 dx$$

$$\begin{aligned}
&= \frac{4 \operatorname{Subst}\left(\int \frac{(-1+x)x(1+x)^5}{(1+x^2)^5} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a^4} \\
&= \frac{(1-ax)(1+ax)^3}{4a^4} - \frac{\operatorname{Subst}\left(\int \frac{24x+32x^2-32x^3-32x^4-8x^5}{(1+x^2)^4} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{2a^4} \\
&= \frac{(1-ax)(1+ax)^3}{4a^4} - \frac{\operatorname{Subst}\left(\int \frac{x(24+32x-32x^2-32x^3-8x^4)}{(1+x^2)^4} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{2a^4} \\
&= \frac{(1-ax)(1+ax)^3}{4a^4} + \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^3 \left(4 - 3\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^4} \\
&\quad + \frac{\operatorname{Subst}\left(\int \frac{-64-48x+192x^2+48x^3}{(1+x^2)^3} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{12a^4} \\
&= \frac{(1-ax)(1+ax)^3}{4a^4} + \frac{(1+ax)^2 \left(3 - 8\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^4} \\
&\quad + \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^3 \left(4 - 3\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^4} - \frac{\operatorname{Subst}\left(\int -\frac{192x}{(1+x^2)^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{48a^4} \\
&= \frac{(1-ax)(1+ax)^3}{4a^4} + \frac{(1+ax)^2 \left(3 - 8\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^4} \\
&\quad + \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^3 \left(4 - 3\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^4} + \frac{4 \operatorname{Subst}\left(\int \frac{x}{(1+x^2)^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a^4} \\
&= -\frac{x}{a^3} + \frac{(1-ax)(1+ax)^3}{4a^4} + \frac{(1+ax)^2 \left(3 - 8\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^4} + \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^3 \left(4 - 3\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^4}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.44

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^3 dx = \frac{x^2}{a^2} - \frac{x^4}{4} + \frac{2(-1+ax)\sqrt{\frac{1-ax}{1+ax}}(1+ax)^2}{3a^4}$$

[In] Integrate[E^(2*ArcSech[a*x])*x^3,x]

[Out] x^2/a^2 - x^4/4 + (2*(-1 + a*x)*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2)/(3*a^4)

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.62

method	result	size
default	$\frac{-\frac{1}{4}x^4a^2 + \frac{1}{2}x^2}{a^2} + \frac{2\sqrt{\frac{ax+1}{ax}}x\sqrt{-\frac{ax-1}{ax}}(a^2x^2-1)}{3a^3} + \frac{x^2}{2a^2}$	72

[In] int((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2*x^3,x,method=_RETURNVERBOSE)

[Out] 1/a^2*(-1/4*x^4*a^2+1/2*x^2)+2/3/a^3*((a*x+1)/a/x)^(1/2)*x*(-(a*x-1)/a/x)^(1/2)*(a^2*x^2-1)+1/2*x^2/a^2

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.53

$$\int e^{2\operatorname{sech}^{-1}(ax)}x^3 dx = -\frac{3a^3x^4 - 12ax^2 - 8(a^2x^3 - x)\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}}{12a^3}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2*x^3,x, algorithm="fricas")

[Out] -1/12*(3*a^3*x^4 - 12*a*x^2 - 8*(a^2*x^3 - x)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)))/a^3

Sympy [F]

$$\int e^{2\operatorname{sech}^{-1}(ax)}x^3 dx = \frac{\int 2x dx + \int (-a^2x^3) dx + \int 2ax^2\sqrt{-1 + \frac{1}{ax}}\sqrt{1 + \frac{1}{ax}} dx}{a^2}$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))**2*x**3,x)

[Out] (Integral(2*x, x) + Integral(-a**2*x**3, x) + Integral(2*a*x**2*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)), x))/a**2

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.36

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^3 dx = -\frac{1}{4}x^4 + \frac{x^2}{a^2} + \frac{2(a^2x^2 - 1)\sqrt{ax+1}\sqrt{-ax+1}}{3a^4}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2*x^3,x, algorithm="maxima")

[Out] -1/4*x^4 + x^2/a^2 + 2/3*(a^2*x^2 - 1)*sqrt(a*x + 1)*sqrt(-a*x + 1)/a^4

Giac [F]

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^3 dx = \int x^3 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} \right)^2 dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2*x^3,x, algorithm="giac")

[Out] integrate(x^3*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))^2, x)

Mupad [B] (verification not implemented)

Time = 4.68 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.54

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^3 dx = \frac{x^2}{a^2} - \frac{x^4}{4} - \sqrt{\frac{1}{ax} - 1} \left(\frac{2x\sqrt{\frac{1}{ax} + 1}}{3a^3} - \frac{2x^3\sqrt{\frac{1}{ax} + 1}}{3a} \right)$$

[In] int(x^3*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))^2,x)

[Out] x^2/a^2 - x^4/4 - (1/(a*x) - 1)^(1/2)*((2*x*(1/(a*x) + 1)^(1/2))/(3*a^3) - (2*x^3*(1/(a*x) + 1)^(1/2))/(3*a))

3.67 $\int e^{2\operatorname{sech}^{-1}(ax)} x^2 dx$

Optimal result	436
Rubi [A] (verified)	436
Mathematica [C] (verified)	439
Maple [C] (verified)	439
Fricas [A] (verification not implemented)	439
Sympy [F]	440
Maxima [F]	440
Giac [F]	440
Mupad [B] (verification not implemented)	441

Optimal result

Integrand size = 12, antiderivative size = 169

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^2 dx = \frac{(1+ax)\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)\left(1+\sqrt{\frac{1-ax}{1+ax}}\right)}{2a^3} - \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^2\left(1+\sqrt{\frac{1-ax}{1+ax}}\right)^3}{6a^3} + \frac{(1+ax)^3\left(1+\sqrt{\frac{1-ax}{1+ax}}\right)^4}{12a^3} - \frac{2\arctan\left(\sqrt{\frac{1-ax}{1+ax}}\right)}{a^3}$$

[Out] $-2*\arctan\left(\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)/a^3+1/2*(a*x+1)*(1-\left(\frac{-a*x+1}{a*x+1}\right)^{1/2})*\left(1+\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)/a^3-1/6*(a*x+1)^2*\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}*\left(1+\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)^3/a^3+1/12*(a*x+1)^3*\left(1+\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)^4/a^3$

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6472, 835, 12, 743, 737, 209}

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^2 dx = -\frac{2\arctan\left(\sqrt{\frac{1-ax}{ax+1}}\right)}{a^3} + \frac{(ax+1)^3\left(\sqrt{\frac{1-ax}{ax+1}}+1\right)^4}{12a^3} - \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)^2\left(\sqrt{\frac{1-ax}{ax+1}}+1\right)^3}{6a^3} + \frac{(ax+1)\left(1-\sqrt{\frac{1-ax}{ax+1}}\right)\left(\sqrt{\frac{1-ax}{ax+1}}+1\right)}{2a^3}$$

[In] Int[E^(2*ArcSech[a*x])*x^2,x]

[Out] ((1 + a*x)*(1 - Sqrt[(1 - a*x)/(1 + a*x)])*(1 + Sqrt[(1 - a*x)/(1 + a*x]])) / (2*a^3) - (Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2*(1 + Sqrt[(1 - a*x)/(1 + a*x]]))^3 / (6*a^3) + ((1 + a*x)^3*(1 + Sqrt[(1 - a*x)/(1 + a*x]]))^4 / (12*a^3) - (2*ArcTan[Sqrt[(1 - a*x)/(1 + a*x]]) / a^3

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 737

Int[((d_) + (e_)*(x_)^2)^m*((a_) + (c_)*(x_)^2)^p, x_Symbol] := Simp[(d + e*x)^(m - 1)*(a*e - c*d*x)*((a + c*x^2)^(p + 1)/(2*a*c*(p + 1))), x] + Dist[(2*p + 3)*((c*d^2 + a*e^2)/(2*a*c*(p + 1))), Int[(d + e*x)^(m - 2)*(a + c*x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && EqQ[m + 2*p + 2, 0] && LtQ[p, -1]

Rule 743

Int[((d_) + (e_)*(x_)^2)^m*((a_) + (c_)*(x_)^2)^p, x_Symbol] := Simp[-(d + e*x)^m*(2*c*x)*((a + c*x^2)^(p + 1)/(4*a*c*(p + 1))), x] - Dist[m*(2*c*d)/(4*a*c*(p + 1)), Int[(d + e*x)^(m - 1)*(a + c*x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e, m, p}, x] && NeQ[c*d^2 + a*e^2, 0] && EqQ[m + 2*p + 3, 0] && LtQ[p, -1]

Rule 835

Int[((d_) + (e_)*(x_)^2)^m*((f_) + (g_)*(x_)^2)^p, x_Symbol] := Simp[(d + e*x)^m*(a + c*x^2)^(p + 1)*((a*g - c*f*x)/(2*a*c*(p + 1))), x] - Dist[1/(2*a*c*(p + 1)), Int[(d + e*x)^(m - 1)*(a + c*x^2)^(p + 1)*Simp[a*e*g*m - c*d*f*(2*p + 3) - c*e*f*(m + 2*p + 3)*x, x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && GtQ[m, 0] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])

Rule 6472

Int[E^(ArcSech[u]*(n_))*(x_)^m, x_Symbol] := Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)]) + (1/u)*Sqrt[(1 - u)/(1 + u))]^n, x] /; FreeQ[m, x] && Integer

Q[n]

Rubi steps

$$\begin{aligned}
\text{integral} &= \int x^2 \left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax} \right)^2 dx \\
&= -\frac{4\text{Subst}\left(\int \frac{x(1+x)^4}{(1+x^2)^4} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a^3} \\
&= \frac{(1+ax)^3 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^4}{12a^3} - \frac{2\text{Subst}\left(\int \frac{4(1+x)^3}{(1+x^2)^3} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{3a^3} \\
&= \frac{(1+ax)^3 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^4}{12a^3} - \frac{8\text{Subst}\left(\int \frac{(1+x)^3}{(1+x^2)^3} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{3a^3} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^2 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^3}{6a^3} + \frac{(1+ax)^3 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^4}{12a^3} \\
&\quad - \frac{2\text{Subst}\left(\int \frac{(1+x)^2}{(1+x^2)^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a^3} \\
&= \frac{(1+ax) \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right) \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}{2a^3} - \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^2 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^3}{6a^3} \\
&\quad + \frac{(1+ax)^3 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^4}{12a^3} - \frac{2\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a^3} \\
&= \frac{(1+ax) \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right) \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}{2a^3} - \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^2 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^3}{6a^3} \\
&\quad + \frac{(1+ax)^3 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^4}{12a^3} - \frac{2 \arctan\left(\sqrt{\frac{1-ax}{1+ax}}\right)}{a^3}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.51

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^2 dx = \frac{2x}{a^2} - \frac{x^3}{3} + \sqrt{\frac{1-ax}{1+ax}} \left(\frac{x}{a^2} + \frac{x^2}{a} \right) + \frac{i \log \left(-2iax + 2\sqrt{\frac{1-ax}{1+ax}}(1+ax) \right)}{a^3}$$

[In] Integrate[E^(2*ArcSech[a*x])*x^2,x]

[Out] (2*x)/a^2 - x^3/3 + Sqrt[(1 - a*x)/(1 + a*x)]*(x/a^2 + x^2/a) + (I*Log[(-2*I)*a*x + 2*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)])/a^3

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.05 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.62

method	result	size
default	$-\frac{1}{3} \frac{a^2 x^3 + x}{a^2} + \frac{\sqrt{\frac{ax+1}{ax}} x \sqrt{-\frac{ax-1}{ax}} \left(\sqrt{-a^2 x^2 + 1} x \operatorname{csgn}(a) + \arctan \left(\frac{\operatorname{csgn}(a) ax}{\sqrt{-a^2 x^2 + 1}} \right) \right) \operatorname{csgn}(a)}{a^2 \sqrt{-a^2 x^2 + 1}} + \frac{x}{a^2}$	105

[In] int((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))^2*x^2,x,method=_RETURNVERBOSE)

[Out] 1/a^2*(-1/3*a^2*x^3+x)+1/a^2*((a*x+1)/a/x)^(1/2)*x*(-(a*x-1)/a/x)^(1/2)*((-a^2*x^2+1)^(1/2)*x*csgn(a)*a+arctan(csgn(a)*a*x/(-a^2*x^2+1)^(1/2)))/(-a^2*x^2+1)^(1/2)*csgn(a)+x/a^2

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.51

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^2 dx = -\frac{a^3 x^3 - 3a^2 x^2 \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} - 6ax + 3 \arctan \left(\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} \right)}{3a^3}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))^2*x^2,x, algorithm="fricas")

[Out] -1/3*(a^3*x^3 - 3*a^2*x^2*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 6*a*x + 3*arctan(sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x))))/a^3

Sympy [F]

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^2 dx = \frac{\int 2 dx + \int (-a^2 x^2) dx + \int 2ax \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} dx}{a^2}$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))**2*x**2,x)

[Out] (Integral(2, x) + Integral(-a**2*x**2, x) + Integral(2*a*x*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)), x))/a**2

Maxima [F]

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^2 dx = \int x^2 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} \right)^2 dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2*x^2,x, algorithm="maxima")

[Out] 2*x/a^2 + 2*integrate(sqrt(a*x + 1)*sqrt(-a*x + 1), x)/a^2 - integrate(x^2, x)

Giac [F]

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^2 dx = \int x^2 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} \right)^2 dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2*x^2,x, algorithm="giac")

[Out] integrate(x^2*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))^2, x)

Mupad [B] (verification not implemented)

Time = 12.78 (sec) , antiderivative size = 420, normalized size of antiderivative = 2.49

$$\int e^{2\operatorname{sech}^{-1}(ax)} x^2 dx = \frac{\frac{1i}{16a^3} + \frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^2 1i}{8a^3 \left(\sqrt{\frac{1}{ax}+1-1}\right)^2} - \frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^4 15i}{16a^3 \left(\sqrt{\frac{1}{ax}+1-1}\right)^4} - \frac{x^3 \left(\frac{a^2}{3} - \frac{2}{x^2}\right)}{a^2}}{\frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^2}{\left(\sqrt{\frac{1}{ax}+1-1}\right)^2} + \frac{2\left(\sqrt{\frac{1}{ax}-1-i}\right)^4}{\left(\sqrt{\frac{1}{ax}+1-1}\right)^4} + \frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^6}{\left(\sqrt{\frac{1}{ax}+1-1}\right)^6}} + \frac{\left(\ln\left(\frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^2}{\left(\sqrt{\frac{1}{ax}+1-1}\right)^2} + 1\right) - \ln\left(\frac{\sqrt{\frac{1}{ax}-1-i}}{\sqrt{\frac{1}{ax}+1-1}}\right)\right) 2i}{a^3} + \frac{\ln\left(\frac{\sqrt{\frac{1}{ax}-1-i}}{\sqrt{\frac{1}{ax}+1-1}}\right) 1i}{a^3} - \frac{\ln\left(\frac{2a\sqrt{\frac{a+\frac{1}{x}}{a}} - \frac{2}{x} + a\sqrt{-\frac{a-\frac{1}{x}}{a}} 2i}{2a+\frac{1}{x} - 2a\sqrt{\frac{a+\frac{1}{x}}{a}}}\right) 1i}{a^3} + \frac{\left(\sqrt{\frac{1}{ax}-1-i}\right)^2 1i}{16a^3 \left(\sqrt{\frac{1}{ax}+1-1}\right)^2}$$

[In] int(x^2*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))^2,x)

[Out] ((log(((1/(a*x) - 1)^(1/2) - 1i)^2/((1/(a*x) + 1)^(1/2) - 1)^2 + 1) - log((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1)))*2i)/a^3 + (log(((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1)))*1i)/a^3 + (1i/(16*a^3) + ((1/(a*x) - 1)^(1/2) - 1i)^2*1i)/(8*a^3*((1/(a*x) + 1)^(1/2) - 1)^2) - (((1/(a*x) - 1)^(1/2) - 1i)^4*15i)/(16*a^3*((1/(a*x) + 1)^(1/2) - 1)^4))/(((1/(a*x) - 1)^(1/2) - 1i)^2/((1/(a*x) + 1)^(1/2) - 1)^2 + (2*((1/(a*x) - 1)^(1/2) - 1i)^4)/((1/(a*x) + 1)^(1/2) - 1)^4 + ((1/(a*x) - 1)^(1/2) - 1i)^6/((1/(a*x) + 1)^(1/2) - 1)^6) - (log((a*(-(a - 1/x)/a)^(1/2)*2i - 2/x + 2*a*((a + 1/x)/a)^(1/2)))/(2*a + 1/x - 2*a*((a + 1/x)/a)^(1/2)))*1i)/a^3 + (((1/(a*x) - 1)^(1/2) - 1i)^2*1i)/(16*a^3*((1/(a*x) + 1)^(1/2) - 1)^2) - (x^3*(a^2/3 - 2/x^2))/a^2)

3.68 $\int e^{2\operatorname{sech}^{-1}(ax)} x dx$

Optimal result	442
Rubi [A] (verified)	442
Mathematica [A] (verified)	444
Maple [A] (verified)	444
Fricas [A] (verification not implemented)	445
Sympy [F]	445
Maxima [F]	445
Giac [F]	446
Mupad [B] (verification not implemented)	446

Optimal result

Integrand size = 10, antiderivative size = 85

$$\int e^{2\operatorname{sech}^{-1}(ax)} x dx = -\frac{(1+ax)^2}{2a^2} + \frac{(1+ax)\left(1+2\sqrt{\frac{1-ax}{1+ax}}\right)}{a^2} + \frac{2\log(1+ax)}{a^2} + \frac{4\log\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)}{a^2}$$

[Out] $-1/2*(a*x+1)^2/a^2+2*\ln(a*x+1)/a^2+4*\ln(1-((-a*x+1)/(a*x+1))^(1/2))/a^2+(a*x+1)*(1+2*((-a*x+1)/(a*x+1))^(1/2))/a^2$

Rubi [A] (verified)

Time = 0.31 (sec), antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6472, 1661, 1607, 815, 266}

$$\int e^{2\operatorname{sech}^{-1}(ax)} x dx = -\frac{(ax+1)^2}{2a^2} + \frac{\left(2\sqrt{\frac{1-ax}{ax+1}}+1\right)(ax+1)}{a^2} + \frac{2\log(ax+1)}{a^2} + \frac{4\log\left(1-\sqrt{\frac{1-ax}{ax+1}}\right)}{a^2}$$

[In] $\text{Int}[E^{(2*\text{ArcSech}[a*x])}*x,x]$

[Out] $-1/2*(1+a*x)^2/a^2 + ((1+a*x)*(1+2*\text{Sqrt}[(1-a*x)/(1+a*x]]))/a^2 + (2*\text{Log}[1+a*x])/a^2 + (4*\text{Log}[1-\text{Sqrt}[(1-a*x)/(1+a*x]]))/a^2$

Rule 266

Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]

Rule 815

Int[(((d_) + (e_)*(x_)^(m_))*((f_) + (g_)*(x_)))/((a_) + (c_)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)/(a + c*x^2)), x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[m]

Rule 1607

Int[(u_)*((a_)*(x_)^(p_) + (b_)*(x_)^(q_))^(n_), x_Symbol] := Int[u*x^(n*p)*(a + b*x^(q - p))^n, x] /; FreeQ[{a, b, p, q}, x] && IntegerQ[n] && PosQ[q - p]

Rule 1661

Int[(Pq_)*((d_) + (e_)*(x_)^(m_))*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[(d + e*x)^m*Pq, a + c*x^2, x], f = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + c*x^2, x], x, 0], g = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + c*x^2, x], x, 1]}, Simp[(a*g - c*f*x)*((a + c*x^2)^(p + 1)/(2*a*c*(p + 1))), x] + Dist[1/(2*a*c*(p + 1)), Int[(d + e*x)^m*(a + c*x^2)^(p + 1)*ExpandToSum[(2*a*c*(p + 1)*Q]/(d + e*x)^m + (c*f*(2*p + 3))/(d + e*x)^m, x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Pq, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && ILtQ[m, 0]

Rule 6472

Int[E^(ArcSech[u_]*(n_))*(x_)^(m_), x_Symbol] := Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && IntegerQ[n]

Rubi steps

$$\begin{aligned} \text{integral} &= \int x \left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax} \right)^2 dx \\ &= \frac{4 \text{Subst} \left(\int \frac{x(1+x)^3}{(-1+x)(1+x^2)^3} dx, x, \sqrt{\frac{1-ax}{1+ax}} \right)}{a^2} \\ &= -\frac{(1+ax)^2}{2a^2} - \frac{\text{Subst} \left(\int \frac{-12x-4x^2}{(-1+x)(1+x^2)^2} dx, x, \sqrt{\frac{1-ax}{1+ax}} \right)}{a^2} \\ &= -\frac{(1+ax)^2}{2a^2} - \frac{\text{Subst} \left(\int \frac{(-12-4x)x}{(-1+x)(1+x^2)^2} dx, x, \sqrt{\frac{1-ax}{1+ax}} \right)}{a^2} \end{aligned}$$

$$\begin{aligned}
&= -\frac{(1+ax)^2}{2a^2} + \frac{(1+ax)\left(1+2\sqrt{\frac{1-ax}{1+ax}}\right)}{a^2} + \frac{\text{Subst}\left(\int \frac{8+8x}{(-1+x)(1+x^2)} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{2a^2} \\
&= -\frac{(1+ax)^2}{2a^2} + \frac{(1+ax)\left(1+2\sqrt{\frac{1-ax}{1+ax}}\right)}{a^2} + \frac{\text{Subst}\left(\int \left(\frac{8}{-1+x} - \frac{8x}{1+x^2}\right) dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{2a^2} \\
&= -\frac{(1+ax)^2}{2a^2} + \frac{(1+ax)\left(1+2\sqrt{\frac{1-ax}{1+ax}}\right)}{a^2} + \frac{4\log\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)}{a^2} - \frac{4\text{Subst}\left(\int \frac{x}{1+x^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a^2} \\
&= -\frac{(1+ax)^2}{2a^2} + \frac{(1+ax)\left(1+2\sqrt{\frac{1-ax}{1+ax}}\right)}{a^2} + \frac{2\log(1+ax)}{a^2} + \frac{4\log\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)}{a^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.05

$$\int e^{2\text{sech}^{-1}(ax)} x dx = \frac{-a^2 x^2 + 4\sqrt{\frac{1-ax}{1+ax}}(1+ax) + 8\log(x) - 4\log\left(1 + \sqrt{\frac{1-ax}{1+ax}} + ax\sqrt{\frac{1-ax}{1+ax}}\right)}{2a^2}$$

[In] Integrate[E^(2*ArcSech[a*x])*x,x]

[Out] $(-(a^2 x^2) + 4\sqrt{(1-ax)/(1+ax)}(1+ax) + 8\text{Log}[x] - 4\text{Log}[1 + \sqrt{(1-ax)/(1+ax)} + ax\sqrt{(1-ax)/(1+ax)}])/(2a^2)$

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.15

method	result	size
default	$\frac{-a^2 x^2 + \ln(x)}{a^2} - \frac{2\sqrt{\frac{ax+1}{ax}} x \sqrt{-\frac{ax-1}{ax}} \left(-\sqrt{-a^2 x^2 + 1} + \text{arctanh}\left(\frac{1}{\sqrt{-a^2 x^2 + 1}}\right)\right)}{a\sqrt{-a^2 x^2 + 1}} + \frac{\ln(x)}{a^2}$	98

[In] int((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2*x,x,method=_RETURNVERBOSE)

[Out] $1/a^2*(-1/2*a^2*x^2+\ln(x))-2/a*((ax+1)/a/x)^(1/2)*x*(-(ax-1)/a/x)^(1/2)*(-(-a^2*x^2+1)^(1/2)+\text{arctanh}(1/(-a^2*x^2+1)^(1/2)))/(-a^2*x^2+1)^(1/2)+\ln(x)/a^2$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.46

$$\int e^{2\operatorname{sech}^{-1}(ax)} x dx = \frac{a^2 x^2 - 4ax \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} + 2 \log \left(ax \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} + 1 \right) - 2 \log \left(ax \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} - 1 \right) - 4 \log(x)}{2a^2}$$

```
[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2*x,x, algorithm="fricas")
```

```
[Out] -1/2*(a^2*x^2 - 4*a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + 2*log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + 1) - 2*log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 1) - 4*log(x))/a^2
```

Sympy [F]

$$\int e^{2\operatorname{sech}^{-1}(ax)} x dx = \frac{\int \frac{2}{x} dx + \int (-a^2 x) dx + \int 2a \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} dx}{a^2}$$

```
[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))**2*x,x)
```

```
[Out] (Integral(2/x, x) + Integral(-a**2*x, x) + Integral(2*a*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)), x))/a**2
```

Maxima [F]

$$\int e^{2\operatorname{sech}^{-1}(ax)} x dx = \int x \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} \right)^2 dx$$

```
[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2*x,x, algorithm="maxima")
```

```
[Out] integrate(x*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))^2, x)
```

Giac [F]

$$\int e^{2\operatorname{sech}^{-1}(ax)} x dx = \int x \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} \right)^2 dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2*x,x, algorithm="giac")

[Out] integrate(x*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))^2, x)

Mupad [B] (verification not implemented)

Time = 7.01 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.66

$$\int e^{2\operatorname{sech}^{-1}(ax)} x dx = \frac{2x \sqrt{\frac{1}{ax} - 1} \sqrt{\frac{1}{ax} + 1}}{a} - \frac{2 \operatorname{acosh}\left(\frac{1}{ax}\right)}{a^2} - \frac{x^2}{2} - \frac{2 \ln\left(\frac{1}{x}\right)}{a^2}$$

[In] int(x*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))^2,x)

[Out] (2*x*(1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2))/a - (2*acosh(1/(a*x)))/a^2 - x^2/2 - (2*log(1/x))/a^2

3.69 $\int e^{2\operatorname{sech}^{-1}(ax)} dx$

Optimal result	447
Rubi [A] (verified)	447
Mathematica [A] (verified)	449
Maple [C] (verified)	449
Fricas [A] (verification not implemented)	450
Sympy [F]	450
Maxima [F]	450
Giac [F]	451
Mupad [B] (verification not implemented)	451

Optimal result

Integrand size = 8, antiderivative size = 57

$$\int e^{2\operatorname{sech}^{-1}(ax)} dx = -x - \frac{4}{a \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} + \frac{4 \arctan\left(\sqrt{\frac{1-ax}{1+ax}}\right)}{a}$$

[Out] $-x+4*\arctan(((-a*x+1)/(a*x+1))^{(1/2)})/a-4/a/(1-((-a*x+1)/(a*x+1))^{(1/2)})$

Rubi [A] (verified)

Time = 0.13 (sec), antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {6467, 1661, 12, 815, 209}

$$\int e^{2\operatorname{sech}^{-1}(ax)} dx = \frac{4 \arctan\left(\sqrt{\frac{1-ax}{ax+1}}\right)}{a} - \frac{4}{a \left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)} - x$$

[In] $\text{Int}[E^{(2*\text{ArcSech}[a*x])}, x]$

[Out] $-x - 4/(a*(1 - \text{Sqrt}[(1 - a*x)/(1 + a*x)])) + (4*\text{ArcTan}[\text{Sqrt}[(1 - a*x)/(1 + a*x)])]/a$

Rule 12

$\text{Int}[(a_*)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{Match}[\text{Q}[u, (b_)*(v_)] /; \text{FreeQ}[b, x]]$

Rule 209

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 815

```
Int[(((d_) + (e_)*(x_)^(m_))*((f_) + (g_)*(x_)))/((a_) + (c_)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)/(a + c*x^2)), x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[m]
```

Rule 1661

```
Int[(Pq_)*((d_) + (e_)*(x_)^(m_))*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] :> With[{Q = PolynomialQuotient[(d + e*x)^m*Pq, a + c*x^2, x], f = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + c*x^2, x], x, 0], g = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + c*x^2, x], x, 1]}, Simp[(a*g - c*f*x)*((a + c*x^2)^(p + 1)/(2*a*c*(p + 1))), x] + Dist[1/(2*a*c*(p + 1)), Int[(d + e*x)^m*(a + c*x^2)^(p + 1)*ExpandToSum[(2*a*c*(p + 1)*Q)/(d + e*x)^m + (c*f*(2*p + 3))/(d + e*x)^m, x], x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Pq, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && ILtQ[m, 0]
```

Rule 6467

```
Int[E^(ArcSech[u_]*(n_.)), x_Symbol] := Int[(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax} \right)^2 dx \\
 &= -\frac{4\text{Subst}\left(\int \frac{x(1+x)^2}{(-1+x)^2(1+x^2)^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a} \\
 &= -x + \frac{2\text{Subst}\left(\int -\frac{4x}{(-1+x)^2(1+x^2)} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a} \\
 &= -x - \frac{8\text{Subst}\left(\int \frac{x}{(-1+x)^2(1+x^2)} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a} \\
 &= -x - \frac{8\text{Subst}\left(\int \left(\frac{1}{2(-1+x)^2} - \frac{1}{2(1+x^2)}\right) dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a}
 \end{aligned}$$

$$\begin{aligned}
&= -x - \frac{4}{a \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} + \frac{4 \text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a} \\
&= -x - \frac{4}{a \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} + \frac{4 \arctan\left(\sqrt{\frac{1-ax}{1+ax}}\right)}{a}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.32

$$\int e^{2\text{sech}^{-1}(ax)} dx = -\frac{2 + a^2x^2 + 2\sqrt{\frac{1-ax}{1+ax}}(1+ax) + 2ax \arctan\left(\frac{ax}{\sqrt{\frac{1-ax}{1+ax}}(1+ax)}\right)}{a^2x}$$

[In] Integrate[E^(2*ArcSech[a*x]),x]

[Out] -((2 + a^2*x^2 + 2*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x) + 2*a*x*ArcTan[(a*x)/(Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x))])/(a^2*x))

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.06 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.95

method	result	size
default	$ \frac{-a^2x - \frac{1}{x}}{a^2} - \frac{2\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}\left(\arctan\left(\frac{\text{csgn}(a)ax}{\sqrt{-a^2x^2+1}}\right)ax + \text{csgn}(a)\sqrt{-a^2x^2+1}\right)\text{csgn}(a)}{a\sqrt{-a^2x^2+1}} - \frac{1}{xa^2} $	111

[In] int((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))^2,x,method=_RETURNVERBOSE)

[Out] 1/a^2*(-a^2*x-1/x)-2/a*((a*x+1)/a/x)^(1/2)*(-(a*x-1)/a/x)^(1/2)*(arctan(csgn(a)*a*x/(-a^2*x^2+1)^(1/2))*a*x+csgn(a)*(-a^2*x^2+1)^(1/2))*csgn(a)/(-a^2*x^2+1)^(1/2)-1/x/a^2

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.49

$$\int e^{2\operatorname{sech}^{-1}(ax)} dx = -\frac{a^2 x^2 + 2ax\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} - 2ax \arctan\left(\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}\right) + 2}{a^2 x}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2,x, algorithm="fricas")

[Out] -(a^2*x^2 + 2*a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 2*a*x*arctan(sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x))) + 2)/(a^2*x)

Sympy [F]

$$\int e^{2\operatorname{sech}^{-1}(ax)} dx = \frac{\int (-a^2) dx + \int \frac{2}{x^2} dx + \int \frac{2a\sqrt{-1+\frac{1}{ax}}\sqrt{1+\frac{1}{ax}}}{x} dx}{a^2}$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))**2,x)

[Out] (Integral(-a**2, x) + Integral(2/x**2, x) + Integral(2*a*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x))/x, x))/a**2

Maxima [F]

$$\int e^{2\operatorname{sech}^{-1}(ax)} dx = \int \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} \right)^2 dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2,x, algorithm="maxima")

[Out] -x + 2*integrate(sqrt(a*x + 1)*sqrt(-a*x + 1)/x^2, x)/a^2 + integrate(x^(-2), x)/a^2 - 1/(a^2*x)

Giac [F]

$$\int e^{2\operatorname{sech}^{-1}(ax)} dx = \int \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} \right)^2 dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))^2, x)

Mupad [B] (verification not implemented)

Time = 7.78 (sec) , antiderivative size = 162, normalized size of antiderivative = 2.84

$$\int e^{2\operatorname{sech}^{-1}(ax)} dx = -x - \frac{\left(\ln \left(\frac{\left(\sqrt{\frac{1}{ax} - 1} - i \right)^2}{\left(\sqrt{\frac{1}{ax} + 1} - 1 \right)^2} + 1 \right) - \ln \left(\frac{\sqrt{\frac{1}{ax} - 1} - i}{\sqrt{\frac{1}{ax} + 1} - 1} \right) \right) 2i}{a} - \frac{2}{a^2 x} + \frac{\left(1 + \sqrt{-\frac{a - \frac{1}{x}}{a}} i \right)^2 \left(\sqrt{\frac{a + \frac{1}{x}}{a}} - 1 \right)^2 4i}{a \left(\sqrt{\frac{a + \frac{1}{x}}{a}} i + \sqrt{-\frac{a - \frac{1}{x}}{a}} - 2i \right)^2}$$

[In] int(((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))^2,x)

[Out] (((-(a - 1/x)/a)^(1/2)*1i + 1)^2*(((a + 1/x)/a)^(1/2) - 1)^2*4i)/(a*(((a + 1/x)/a)^(1/2)*1i + (-(a - 1/x)/a)^(1/2) - 2i)^2) - ((log(((1/(a*x) - 1)^(1/2) - 1i)^2/((1/(a*x) + 1)^(1/2) - 1)^2 + 1) - log(((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1))))*2i)/a - 2/(a^2*x) - x

3.70 $\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x} dx$

Optimal result	452
Rubi [A] (verified)	452
Mathematica [A] (verified)	453
Maple [A] (verified)	454
Fricas [A] (verification not implemented)	454
Sympy [A] (verification not implemented)	454
Maxima [F]	455
Giac [F]	455
Mupad [B] (verification not implemented)	455

Optimal result

Integrand size = 12, antiderivative size = 86

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x} dx = -\frac{2}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{2}{1 - \sqrt{\frac{1-ax}{1+ax}}} - \log(1+ax) - 2 \log\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)$$

[Out] $-\ln(a*x+1)-2*\ln(1-((-a*x+1)/(a*x+1))^(1/2))-2/(1-((-a*x+1)/(a*x+1))^(1/2))-2+2/(1-((-a*x+1)/(a*x+1))^(1/2))$

Rubi [A] (verified)

Time = 0.32 (sec), antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6472, 1643, 266}

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x} dx = \frac{2}{1 - \sqrt{\frac{1-ax}{ax+1}}} - \frac{2}{\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^2} - \log(ax+1) - 2 \log\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)$$

[In] $\text{Int}[E^{(2*\text{ArcSech}[a*x])}/x, x]$

[Out] $-2/(1 - \text{Sqrt}[(1 - a*x)/(1 + a*x)])^2 + 2/(1 - \text{Sqrt}[(1 - a*x)/(1 + a*x)]) - \text{Log}[1 + a*x] - 2*\text{Log}[1 - \text{Sqrt}[(1 - a*x)/(1 + a*x)]]$

Rule 266

$\text{Int}[(x_)^{(m_.)}/((a_) + (b_.)*(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^n, x]]/(b*n), x] /; \text{FreeQ}\{[a, b, m, n], x\} \&\& \text{EqQ}[m, n - 1]$

Rule 1643

```
Int[(Pq_)*((d_) + (e_)*(x_)^(m_))*((a_) + (c_)*(x_)^2)^(p_), x_Symbol]
:> Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + c*x^2)^p, x], x] /; FreeQ[{a, c,
d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

Rule 6472

```
Int[E^(ArcSech[u_]*(n_))*(x_)^(m_), x_Symbol] :> Int[x^m*(1/u + Sqrt[(1 -
u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && Integer
Q[n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \frac{\left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax}\right)^2}{x} dx \\
&= 4\text{Subst}\left(\int \frac{x(1+x)}{(-1+x)^3(1+x^2)} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right) \\
&= 4\text{Subst}\left(\int \left(\frac{1}{(-1+x)^3} + \frac{1}{2(-1+x)^2} - \frac{1}{2(-1+x)} + \frac{x}{2(1+x^2)}\right) dx, x, \sqrt{\frac{1-ax}{1+ax}}\right) \\
&= -\frac{2}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{2}{1 - \sqrt{\frac{1-ax}{1+ax}}} - 2\log\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right) \\
&\quad + 2\text{Subst}\left(\int \frac{x}{1+x^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right) \\
&= -\frac{2}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{2}{1 - \sqrt{\frac{1-ax}{1+ax}}} - \log(1+ax) - 2\log\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.00

$$\int \frac{e^{2\text{sech}^{-1}(ax)}}{x} dx = -\frac{1}{a^2x^2} - \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)}{a^2x^2} - 2\log(x) + \log\left(1 + \sqrt{\frac{1-ax}{1+ax}} + ax\sqrt{\frac{1-ax}{1+ax}}\right)$$

```
[In] Integrate[E^(2*ArcSech[a*x])/x, x]
```

```
[Out] -(1/(a^2*x^2)) - (Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x))/(a^2*x^2) - 2*Log[x]
+ Log[1 + Sqrt[(1 - a*x)/(1 + a*x)] + a*x*Sqrt[(1 - a*x)/(1 + a*x)]]
```

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.28

method	result	size
default	$\frac{-a^2 \ln(x) - \frac{1}{2x^2}}{a^2} + \frac{\sqrt{\frac{ax+1}{ax}} \sqrt{\frac{-ax-1}{ax}} \left(a^2 x^2 \operatorname{arctanh}\left(\frac{1}{\sqrt{-a^2 x^2 + 1}}\right) - \sqrt{-a^2 x^2 + 1} \right)}{ax \sqrt{-a^2 x^2 + 1}} - \frac{1}{2a^2 x^2}$	110

[In] int((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x,x,method=_RETURNVERBOSE)

[Out] $\frac{1}{a^2} * (-a^2 * \ln(x) - 1/2/x^2) + 1/a * ((a*x+1)/a/x)^{(1/2)}/x * (-a*x-1)/a/x)^{(1/2)} * (a^2*x^2*\operatorname{arctanh}(1/(\sqrt{-a^2*x^2+1})^{(1/2)}) - (\sqrt{-a^2*x^2+1})^{(1/2)}) / (\sqrt{-a^2*x^2+1})^{(1/2)} - 1/2/a^2/x^2$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.60

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x} dx$$

$$= \frac{a^2 x^2 \log\left(ax \sqrt{\frac{ax+1}{ax}} \sqrt{\frac{-ax-1}{ax}} + 1\right) - a^2 x^2 \log\left(ax \sqrt{\frac{ax+1}{ax}} \sqrt{\frac{-ax-1}{ax}} - 1\right) - 2 a^2 x^2 \log(x) - 2 ax \sqrt{\frac{ax+1}{ax}} \sqrt{\frac{-ax-1}{ax}}}{2 a^2 x^2}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x,x, algorithm="fricas")

[Out] $\frac{1}{2} * (a^2 * x^2 * \log(a*x*\sqrt{(a*x+1)/(a*x)}*\sqrt{-(a*x-1)/(a*x)}+1) - a^2 * x^2 * \log(a*x*\sqrt{(a*x+1)/(a*x)}*\sqrt{-(a*x-1)/(a*x)}-1) - 2*a^2*x^2*\log(x) - 2*a*x*\sqrt{(a*x+1)/(a*x)}*\sqrt{-(a*x-1)/(a*x)}-2)/(a^2*x^2)$

Sympy [A] (verification not implemented)

Time = 2.89 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.95

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x} dx$$

$$= \frac{-2a^2 \cdot \left(2\sqrt{-1 + \frac{1}{ax}} \left(\frac{(1 + \frac{1}{ax})^{\frac{3}{2}}}{4} - \frac{\sqrt{1 + \frac{1}{ax}}}{4} \right) - \log\left(2\sqrt{-1 + \frac{1}{ax}} + 2\sqrt{1 + \frac{1}{ax}} \right) \right) - a^2 \log(x) - \frac{1}{x^2}}{a^2}$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))**2/x,x)

[Out] $(-2*a**2*(2*\sqrt{-1+1/(a*x)}*((1+1/(a*x))**(3/2)/4 - \sqrt{1+1/(a*x)})/4 - \log(2*\sqrt{-1+1/(a*x)}+2*\sqrt{1+1/(a*x)})) - a**2*\log(x) - 1/x**2)/a**2$

Maxima [F]

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x} dx = \int \frac{\left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}\right)^2}{x} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x,x, algorithm="maxima")

[Out] 2*integrate(sqrt(a*x + 1)*sqrt(-a*x + 1)/x^3, x)/a^2 - 1/(a^2*x^2) - integrate(1/x, x)

Giac [F]

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x} dx = \int \frac{\left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}\right)^2}{x} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))^2/x, x)

Mupad [B] (verification not implemented)

Time = 13.88 (sec) , antiderivative size = 323, normalized size of antiderivative = 3.76

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x} dx = \ln\left(\frac{1}{x}\right) - 4 \operatorname{atanh}\left(\frac{\sqrt{\frac{1}{ax} - 1} - i}{\sqrt{\frac{1}{ax} + 1} - 1}\right) + 2 \operatorname{acosh}\left(\frac{1}{ax}\right) + \frac{28\left(\sqrt{\frac{1}{ax} - 1} - i\right)^3}{\left(\sqrt{\frac{1}{ax} + 1} - 1\right)^3} + \frac{28\left(\sqrt{\frac{1}{ax} - 1} - i\right)^5}{\left(\sqrt{\frac{1}{ax} + 1} - 1\right)^5} + \frac{4\left(\sqrt{\frac{1}{ax} - 1} - i\right)^7}{\left(\sqrt{\frac{1}{ax} + 1} - 1\right)^7} + \frac{4\left(\sqrt{\frac{1}{ax} - 1} - i\right)}{\sqrt{\frac{1}{ax} + 1} - 1} - \frac{1}{a^2 x^2} + \frac{6\left(\sqrt{\frac{1}{ax} - 1} - i\right)^4}{\left(\sqrt{\frac{1}{ax} + 1} - 1\right)^4} - \frac{4\left(\sqrt{\frac{1}{ax} - 1} - i\right)^6}{\left(\sqrt{\frac{1}{ax} + 1} - 1\right)^6} + \frac{\left(\sqrt{\frac{1}{ax} - 1} - i\right)^8}{\left(\sqrt{\frac{1}{ax} + 1} - 1\right)^8} - \frac{4\left(\sqrt{\frac{1}{ax} - 1} - i\right)^2}{\left(\sqrt{\frac{1}{ax} + 1} - 1\right)^2}$$

[In] int(((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))^2/x,x)

[Out] log(1/x) - 4*atanh(((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1)) + 2*acosh(1/(a*x)) + ((28*((1/(a*x) - 1)^(1/2) - 1i)^3)/((1/(a*x) + 1)^(1/2) - 1)^3 + (28*((1/(a*x) - 1)^(1/2) - 1i)^5)/((1/(a*x) + 1)^(1/2) - 1)^5 + (4*((1/(a*x) - 1)^(1/2) - 1i)^7)/((1/(a*x) + 1)^(1/2) - 1)^7 + (4*((1/(a*x) - 1)^(1/2) - 1i))/((1/(a*x) + 1)^(1/2) - 1))/((6*((1/(a*x) - 1)^(1/2) - 1i)^4)/((1/(a*x) + 1)^(1/2) - 1)^4 - (4*((1/(a*x) - 1)^(1/2) - 1i)^2)/((1/(a*x) + 1)^(1/2) - 1)^2 - (4*((1/(a*x) - 1)^(1/2) - 1i)^6)/((1/(a*x) + 1)^(1/2) - 1)^6 + ((1/(a*x) - 1)^(1/2) - 1i)^8/((1/(a*x) + 1)^(1/2) - 1)^8 + 1) - 1/(a^2*x^2)

3.71 $\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^2} dx$

Optimal result	456
Rubi [A] (verified)	456
Mathematica [A] (verified)	457
Maple [A] (verified)	457
Fricas [A] (verification not implemented)	458
Sympy [F]	458
Maxima [A] (verification not implemented)	458
Giac [F]	459
Mupad [B] (verification not implemented)	459

Optimal result

Integrand size = 12, antiderivative size = 57

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^2} dx = -\frac{4a}{3\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3} + \frac{2a}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2}$$

[Out] $-4/3*a/(1-((-a*x+1)/(a*x+1))^{(1/2)})^3+2*a/(1-((-a*x+1)/(a*x+1))^{(1/2)})^2$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6472, 45}

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^2} dx = \frac{2a}{\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^2} - \frac{4a}{3\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^3}$$

[In] $\text{Int}[E^{(2*\text{ArcSech}[a*x])}/x^2, x]$

[Out] $(-4*a)/(3*(1 - \text{Sqrt}[(1 - a*x)/(1 + a*x)])^3) + (2*a)/(1 - \text{Sqrt}[(1 - a*x)/(1 + a*x)])^2$

Rule 45

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{IGtQ}[m, 0] \&\& (!\text{IntegerQ}[n] || (\text{EqQ}[c, 0] \&\& \text{LeQ}[7*m + 4*n + 4, 0]) || \text{LtQ}[9*m + 5*(n + 1), 0] || \text{GtQ}[m + n + 2, 0])$

Rule 6472


```
Int[E^(ArcSech[u_]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && IntegerQ[n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \frac{\left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax}\right)^2}{x^2} dx \\ &= -\left((4a)\text{Subst}\left(\int \frac{x}{(-1+x)^4} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)\right) \\ &= -\left((4a)\text{Subst}\left(\int \left(\frac{1}{(-1+x)^4} + \frac{1}{(-1+x)^3}\right) dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)\right) \\ &= -\frac{4a}{3\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3} + \frac{2a}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.91

$$\int \frac{e^{2\text{sech}^{-1}(ax)}}{x^2} dx = \frac{-2 + 3a^2x^2 + 2(-1 + ax)\sqrt{\frac{1-ax}{1+ax}}(1 + ax)^2}{3a^2x^3}$$

```
[In] Integrate[E^(2*ArcSech[a*x])/x^2,x]
```

```
[Out] (-2 + 3*a^2*x^2 + 2*(-1 + a*x)*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2)/(3*a^2*x^3)
```

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.28

method	result	size
default	$-\frac{1}{3a^3} + \frac{a^2}{x} + 2\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} (a^2x^2-1) - \frac{1}{3a^2x^3}$	73

```
[In] int((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))^2/x^2,x,method=_RETURNVERBOSE)
```

```
[Out] 1/a^2*(-1/3/x^3+a^2/x)+2/3/a*((a*x+1)/a/x)^(1/2)/x^2*(-(a*x-1)/a/x)^(1/2)*(a^2*x^2-1)-1/3/a^2/x^3
```

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.07

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^2} dx = \frac{3a^2x^2 + 2(a^3x^3 - ax)\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} - 2}{3a^2x^3}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^2,x, algorithm="fricas")

[Out] 1/3*(3*a^2*x^2 + 2*(a^3*x^3 - a*x)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 2)/(a^2*x^3)

Sympy [F]

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^2} dx = \frac{\int \frac{2}{x^4} dx + \int \left(-\frac{a^2}{x^2}\right) dx + \int \frac{2a\sqrt{-1+\frac{1}{ax}}\sqrt{1+\frac{1}{ax}}}{x^3} dx}{a^2}$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))**2/x**2,x)

[Out] (Integral(2/x**4, x) + Integral(-a**2/x**2, x) + Integral(2*a*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x))/x**3, x))/a**2

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.81

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^2} dx = \frac{1}{x} + \frac{2(a^2x^3 - x)\sqrt{ax+1}\sqrt{-ax+1}}{3a^2x^4} - \frac{2}{3a^2x^3}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^2,x, algorithm="maxima")

[Out] 1/x + 2/3*(a^2*x^3 - x)*sqrt(a*x + 1)*sqrt(-a*x + 1)/(a^2*x^4) - 2/3/(a^2*x^3)

Giac [F]

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^2} dx = \int \frac{\left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}\right)^2}{x^2} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^2,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))^2/x^2, x)

Mupad [B] (verification not implemented)

Time = 4.76 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.18

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^2} dx = \frac{a^2 x^2 - \frac{2}{3}}{a^2 x^3} - \frac{\sqrt{\frac{1}{ax} - 1} \left(\frac{2\sqrt{\frac{1}{ax} + 1}}{3a} - \frac{2ax^2\sqrt{\frac{1}{ax} + 1}}{3} \right)}{x^2}$$

[In] int(((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))^2/x^2,x)

[Out] (a^2*x^2 - 2/3)/(a^2*x^3) - ((1/(a*x) - 1)^(1/2)*((2*(1/(a*x) + 1)^(1/2))/(3*a) - (2*a*x^2*(1/(a*x) + 1)^(1/2))/3))/x^2

3.72 $\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^3} dx$

Optimal result	460
Rubi [A] (verified)	460
Mathematica [A] (verified)	462
Maple [A] (verified)	462
Fricas [A] (verification not implemented)	463
Sympy [F]	463
Maxima [F]	463
Giac [F]	464
Mupad [B] (verification not implemented)	464

Optimal result

Integrand size = 12, antiderivative size = 147

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^3} dx = -\frac{a^2}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^4} + \frac{2a^2}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3} - \frac{3a^2}{2\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{a^2}{2\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} + \frac{1}{2}a^2 \operatorname{arctanh}\left(\sqrt{\frac{1-ax}{1+ax}}\right)$$

[Out] $1/2*a^2*\operatorname{arctanh}\left(\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)-a^2/\left(1-\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)^4+2*a^2/\left(1-\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)^3-3/2*a^2/\left(1-\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)^2+1/2*a^2/\left(1-\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)$

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6472, 1626, 213}

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^3} dx = \frac{1}{2}a^2 \operatorname{arctanh}\left(\sqrt{\frac{1-ax}{ax+1}}\right) + \frac{a^2}{2\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)} - \frac{3a^2}{2\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^2} + \frac{2a^2}{\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^3} - \frac{a^2}{\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^4}$$

[In] $\operatorname{Int}\left[E^{(2*\operatorname{ArcSech}[a*x])}/x^3,x\right]$

[Out] $-(a^2/(1 - \sqrt{(1 - ax)/(1 + ax)}))^4 + (2a^2)/(1 - \sqrt{(1 - ax)/(1 + ax)})^3 - (3a^2)/(2(1 - \sqrt{(1 - ax)/(1 + ax)})^2) + a^2/(2(1 - \sqrt{(1 - ax)/(1 + ax)})) + (a^2 \operatorname{ArcTanh}[\sqrt{(1 - ax)/(1 + ax)}])/2$

Rule 213

$\operatorname{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(-\operatorname{Rt}[-a, 2] \operatorname{Rt}[b, 2])^{-1}] \operatorname{ArcTanh}[\operatorname{Rt}[b, 2] \operatorname{Rt}[-a, 2] / x] / ; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{LtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 1626

$\operatorname{Int}[(P_x) \cdot ((a) + (b \cdot x)^m) \cdot ((c) + (d \cdot x)^n) \cdot ((e) + (f \cdot x)^p), x_Symbol] \rightarrow \operatorname{Int}[\operatorname{ExpandIntegrand}[P_x \cdot (a + b \cdot x)^m \cdot (c + d \cdot x)^n \cdot (e + f \cdot x)^p, x], x] / ; \operatorname{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x \ \&\& \operatorname{PolyQ}[P_x, x] \ \&\& \operatorname{IntegersQ}[m, n]$

Rule 6472

$\operatorname{Int}[E^{\operatorname{ArcSech}[u] \cdot (n)} \cdot (x)^m, x_Symbol] \rightarrow \operatorname{Int}[x^m \cdot (1/u + \sqrt{(1 - u)/(1 + u)}) + (1/u) \cdot \sqrt{(1 - u)/(1 + u)}]^n, x] / ; \operatorname{FreeQ}[m, x] \ \&\& \operatorname{IntegerQ}[n]$

Rubi steps

$$\begin{aligned} \text{integral} &= \int \frac{\left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \sqrt{\frac{1-ax}{1+ax}}\right)^2}{x^3} dx \\ &= (4a^2) \operatorname{Subst}\left(\int \frac{x(1+x^2)}{(-1+x)^5(1+x)} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right) \\ &= (4a^2) \operatorname{Subst}\left(\int \left(\frac{1}{(-1+x)^5} + \frac{3}{2(-1+x)^4} + \frac{3}{4(-1+x)^3} + \frac{1}{8(-1+x)^2} - \frac{1}{8(-1+x^2)}\right) dx, x, \sqrt{\frac{1-ax}{1+ax}}\right) \\ &= -\frac{a^2}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^4} + \frac{2a^2}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3} - \frac{3a^2}{2\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} \\ &\quad + \frac{a^2}{2\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} - \frac{1}{2} a^2 \operatorname{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right) \end{aligned}$$

$$= -\frac{a^2}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^4} + \frac{2a^2}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3} - \frac{3a^2}{2\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2}$$

$$+ \frac{a^2}{2\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} + \frac{1}{2}a^2 \operatorname{arctanh}\left(\sqrt{\frac{1-ax}{1+ax}}\right)$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.82

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^3} dx$$

$$= \frac{(1+ax)\left(-2+2ax-2\sqrt{\frac{1-ax}{1+ax}}+a^2x^2\sqrt{\frac{1-ax}{1+ax}}\right)}{x^4} - a^4 \log(x) + a^4 \log\left(1 + \sqrt{\frac{1-ax}{1+ax}} + ax\sqrt{\frac{1-ax}{1+ax}}\right)$$

$$= \frac{\hspace{15em}}{4a^2}$$

[In] Integrate[E^(2*ArcSech[a*x])/x^3,x]

[Out] (((1 + a*x)*(-2 + 2*a*x - 2*Sqrt[(1 - a*x)/(1 + a*x)] + a^2*x^2*Sqrt[(1 - a*x)/(1 + a*x)]))/x^4 - a^4*Log[x] + a^4*Log[1 + Sqrt[(1 - a*x)/(1 + a*x)] + a*x*Sqrt[(1 - a*x)/(1 + a*x)]])/(4*a^2)

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.89

method	result	size
default	$-\frac{1}{4a^4} + \frac{a^2}{2x^2} + \frac{\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} \left(\operatorname{arctanh}\left(\frac{1}{\sqrt{-a^2x^2+1}}\right) a^4x^4 + a^2x^2\sqrt{-a^2x^2+1} - 2\sqrt{-a^2x^2+1} \right)}{4ax^3\sqrt{-a^2x^2+1}} - \frac{1}{4a^2x^4}$	131

[In] int((1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))^2/x^3,x,method=_RETURNVERBOSE)

[Out] 1/a^2*(-1/4/x^4+1/2*a^2/x^2)+1/4/a*((a*x+1)/a/x)^(1/2)/x^3*(-(a*x-1)/a/x)^(1/2)*(arctanh(1/(-a^2*x^2+1)^(1/2))*a^4*x^4+a^2*x^2*(-a^2*x^2+1)^(1/2)-2*(-a^2*x^2+1)^(1/2))/(-a^2*x^2+1)^(1/2)-1/4/a^2/x^4

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.99

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^3} dx = \frac{a^4 x^4 \log\left(ax \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} + 1\right) - a^4 x^4 \log\left(ax \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} - 1\right) + 4a^2 x^2 + 2(a^3 x^3 - 2ax) \sqrt{\frac{ax+1}{ax}}}{8a^2 x^4}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^3,x, algorithm="fricas")

[Out] 1/8*(a^4*x^4*log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + 1) - a^4*x^4*log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 1) + 4*a^2*x^2 + 2*(a^3*x^3 - 2*a*x)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 4)/(a^2*x^4)

Sympy [F]

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^3} dx = \frac{\int \frac{2}{x^5} dx + \int \left(-\frac{a^2}{x^3}\right) dx + \int \frac{2a\sqrt{-1+\frac{1}{ax}}\sqrt{1+\frac{1}{ax}}}{x^4} dx}{a^2}$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))**2/x**3,x)

[Out] (Integral(2/x**5, x) + Integral(-a**2/x**3, x) + Integral(2*a*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x))/x**4, x))/a**2

Maxima [F]

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^3} dx = \int \frac{\left(\sqrt{\frac{1}{ax} + 1}\sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}}\right)^2}{x^3} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^3,x, algorithm="maxima")

[Out] 2*integrate(sqrt(a*x + 1)*sqrt(-a*x + 1)/x^5, x)/a^2 - 1/2/(a^2*x^4) - integrate(x^(-3), x)

Giac [F]

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^3} dx = \int \frac{\left(\sqrt{\frac{1}{ax} + 1}\sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}\right)^2}{x^3} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^3,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))^2/x^3, x)

Mupad [B] (verification not implemented)

Time = 46.86 (sec) , antiderivative size = 885, normalized size of antiderivative = 6.02

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^3} dx = a^2 \operatorname{atanh}\left(\frac{\sqrt{\frac{1}{ax} - 1 - i}}{\sqrt{\frac{1}{ax} + 1 - 1}}\right) - \frac{28a^2\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^3}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^3} + \frac{28a^2\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^5}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^5} + \frac{4a^2\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^7}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^7} + \frac{4a^2\left(\sqrt{\frac{1}{ax} - 1 - i}\right)}{\sqrt{\frac{1}{ax} + 1 - 1}} - \frac{1 + \frac{6\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^4}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^4} - \frac{4\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^6}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^6} + \frac{\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^8}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^8} - \frac{4\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^2}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^2}}{1 + \frac{28\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^4}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^4} - \frac{56\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^6}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^6} + \frac{70\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^8}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^8} - \frac{56\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^{10}}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^{10}} + \frac{28\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^{12}}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^{12}} - \frac{8\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^2}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^2}} + \frac{23a^2\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^3}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^3} + \frac{333a^2\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^5}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^5} + \frac{671a^2\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^7}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^7} + \frac{671a^2\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^9}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^9} + \frac{333a^2\left(\sqrt{\frac{1}{ax} - 1 - i}\right)^{11}}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)^{11}} + \frac{23a^2\left(\sqrt{\frac{1}{ax} - 1 - i}\right)}{\left(\sqrt{\frac{1}{ax} + 1 - 1}\right)} + \frac{1}{2x^2} - \frac{1}{2a^2x^4}$$

[In] int(((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))^2/x^3,x)

[Out] a^2*atanh(((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1)) - ((28*a^2*((1/(a*x) - 1)^(1/2) - 1i)^3)/((1/(a*x) + 1)^(1/2) - 1)^3 + (28*a^2*((1/(a*x) - 1)^(1/2) - 1i)^5)/((1/(a*x) + 1)^(1/2) - 1)^5 + (4*a^2*((1/(a*x) - 1)^(1/2) - 1i)^7)/((1/(a*x) + 1)^(1/2) - 1)^7 + (4*a^2*((1/(a*x) - 1)^(1/2) - 1i))/((1/(a*x) + 1)^(1/2) - 1))/((6*((1/(a*x) - 1)^(1/2) - 1i)^4)/((1/(a*x) + 1)^(1/2) - 1)^4 - (4*((1/(a*x) - 1)^(1/2) - 1i)^2)/((1/(a*x) + 1)^(1/2) - 1)^2 - (4*((1/(a*x) - 1)^(1/2) - 1i)^6)/((1/(a*x) + 1)^(1/2) - 1)^6 + ((1/(a*x) - 1)^(1/2) - 1i)^8/((1/(a*x) + 1)^(1/2) - 1)^8 + 1) - ((23*a^2*((1/(a*x) - 1)^(1/2) - 1i)^3)/((1/(a*x) + 1)^(1/2) - 1)^3 + (333*a^2*((1/(a*x) - 1)^(1/2) - 1i)^5)/((1/(a*x) + 1)^(1/2) - 1)^5 + (671*a^2*((1/(a*x) - 1)^(1/2) - 1i)^7)/((1/(a*x) + 1)^(1/2) - 1)^7 + (671*a^2*((1/(a*x) - 1)^(1/2) - 1i))/((1/(a*x) + 1)^(1/2) - 1))

$$\begin{aligned}
& 1i)^9)/((1/(a*x) + 1)^{(1/2)} - 1)^9 + (333*a^2*((1/(a*x) - 1)^{(1/2)} - 1i)^{11} \\
&)/((1/(a*x) + 1)^{(1/2)} - 1)^{11} + (23*a^2*((1/(a*x) - 1)^{(1/2)} - 1i)^{13})/((1 \\
& / (a*x) + 1)^{(1/2)} - 1)^{13} - (3*a^2*((1/(a*x) - 1)^{(1/2)} - 1i)^{15})/((1/(a*x) \\
& + 1)^{(1/2)} - 1)^{15} - (3*a^2*((1/(a*x) - 1)^{(1/2)} - 1i))/((1/(a*x) + 1)^{(1/ \\
& 2)} - 1))/((28*((1/(a*x) - 1)^{(1/2)} - 1i)^4)/((1/(a*x) + 1)^{(1/2)} - 1)^4 - (\\
& 8*((1/(a*x) - 1)^{(1/2)} - 1i)^2)/((1/(a*x) + 1)^{(1/2)} - 1)^2 - (56*((1/(a*x) \\
& - 1)^{(1/2)} - 1i)^6)/((1/(a*x) + 1)^{(1/2)} - 1)^6 + (70*((1/(a*x) - 1)^{(1/2) \\
& - 1i)^8)/((1/(a*x) + 1)^{(1/2)} - 1)^8 - (56*((1/(a*x) - 1)^{(1/2)} - 1i)^{10})/ \\
& ((1/(a*x) + 1)^{(1/2)} - 1)^{10} + (28*((1/(a*x) - 1)^{(1/2)} - 1i)^{12})/((1/(a*x) \\
& + 1)^{(1/2)} - 1)^{12} - (8*((1/(a*x) - 1)^{(1/2)} - 1i)^{14})/((1/(a*x) + 1)^{(1/2 \\
&) - 1)^{14} + ((1/(a*x) - 1)^{(1/2)} - 1i)^{16}/((1/(a*x) + 1)^{(1/2)} - 1)^{16} + 1) \\
& + 1/(2*x^2) - 1/(2*a^2*x^4)
\end{aligned}$$

3.73 $\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^4} dx$

Optimal result	466
Rubi [A] (verified)	466
Mathematica [A] (verified)	468
Maple [A] (verified)	468
Fricas [A] (verification not implemented)	468
Sympy [F]	469
Maxima [A] (verification not implemented)	469
Giac [F]	469
Mupad [B] (verification not implemented)	470

Optimal result

Integrand size = 12, antiderivative size = 183

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^4} dx = -\frac{4a^3}{5\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^5} + \frac{2a^3}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^4} - \frac{7a^3}{3\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3} \\ + \frac{3a^3}{2\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} - \frac{a^3}{4\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} - \frac{a^3}{4\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}$$

[Out] $-4/5*a^3/(1-((-a*x+1)/(a*x+1))^(1/2))^5+2*a^3/(1-((-a*x+1)/(a*x+1))^(1/2))^4-7/3*a^3/(1-((-a*x+1)/(a*x+1))^(1/2))^3+3/2*a^3/(1-((-a*x+1)/(a*x+1))^(1/2))^2-1/4*a^3/(1-((-a*x+1)/(a*x+1))^(1/2))-1/4*a^3/(1+((-a*x+1)/(a*x+1))^(1/2))$

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6472, 1626}

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^4} dx = -\frac{a^3}{4\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)} - \frac{a^3}{4\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)} + \frac{3a^3}{2\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^2} \\ - \frac{7a^3}{3\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^3} + \frac{2a^3}{\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^4} - \frac{4a^3}{5\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^5}$$

[In] $\text{Int}[E^{(2*\text{ArcSech}[a*x])/x^4}, x]$

```
[Out] (-4*a^3)/(5*(1 - Sqrt[(1 - a*x)/(1 + a*x)])^5) + (2*a^3)/(1 - Sqrt[(1 - a*x)/(1 + a*x)])^4 - (7*a^3)/(3*(1 - Sqrt[(1 - a*x)/(1 + a*x)])^3) + (3*a^3)/(2*(1 - Sqrt[(1 - a*x)/(1 + a*x)])^2) - a^3/(4*(1 - Sqrt[(1 - a*x)/(1 + a*x)])) - a^3/(4*(1 + Sqrt[(1 - a*x)/(1 + a*x)]))
```

Rule 1626

```
Int[(Px_)*((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && IntegersQ[m, n]
```

Rule 6472

```
Int[E^(ArcSech[u_]*(n_.))*(x_)^(m_.), x_Symbol] :> Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{\left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax}\right)^2}{x^4} dx \\
 &= -\left((4a)\text{Subst}\left(\int \frac{x(a+ax^2)^2}{(-1+x)^6(1+x)^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)\right) \\
 &= -\left((4a)\text{Subst}\left(\int \left(\frac{a^2}{(-1+x)^6} + \frac{2a^2}{(-1+x)^5} + \frac{7a^2}{4(-1+x)^4} + \frac{3a^2}{4(-1+x)^3} + \frac{a^2}{16(-1+x)^2} - \frac{a^2}{16(1+x)}\right) dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)\right) \\
 &= -\frac{4a^3}{5\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^5} + \frac{2a^3}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^4} - \frac{7a^3}{3\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3} \\
 &\quad + \frac{3a^3}{2\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} - \frac{a^3}{4\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} - \frac{a^3}{4\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.38

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^4} dx = \frac{-6 + 5a^2x^2 + 2\sqrt{\frac{1-ax}{1+ax}}(1+ax)^2(-3+3ax-2a^2x^2+2a^3x^3)}{15a^2x^5}$$

[In] Integrate[E^(2*ArcSech[a*x])/x^4,x]

[Out] (-6 + 5*a^2*x^2 + 2*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2*(-3 + 3*a*x - 2*a^2*x^2 + 2*a^3*x^3))/(15*a^2*x^5)

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.46

method	result	size
default	$\frac{a^2}{3x^3} - \frac{1}{5x^5} + \frac{2\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}(a^2x^2-1)(2a^2x^2+3)}{15ax^4} - \frac{1}{5a^2x^5}$	84

[In] int((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^4,x,method=_RETURNVERBOSE)

[Out] 1/a^2*(1/3*a^2/x^3-1/5/x^5)+2/15/a*((a*x+1)/a/x)^(1/2)/x^4*(-(a*x-1)/a/x)^(1/2)*(a^2*x^2-1)*(2*a^2*x^2+3)-1/5/a^2/x^5

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.38

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^4} dx = \frac{5a^2x^2 + 2(2a^5x^5 + a^3x^3 - 3ax)\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} - 6}{15a^2x^5}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^4,x, algorithm="fricas")

[Out] 1/15*(5*a^2*x^2 + 2*(2*a^5*x^5 + a^3*x^3 - 3*a*x)*sqrt((a*x + 1)/(a*x))*sqrt(- (a*x - 1)/(a*x)) - 6)/(a^2*x^5)

Sympy [F]

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^4} dx = \frac{\int \frac{2}{x^6} dx + \int \left(-\frac{a^2}{x^4}\right) dx + \int \frac{2a\sqrt{-1+\frac{1}{ax}}\sqrt{1+\frac{1}{ax}}}{x^5} dx}{a^2}$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))**2/x**4,x)

[Out] (Integral(2/x**6, x) + Integral(-a**2/x**4, x) + Integral(2*a*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x))/x**5, x))/a**2

Maxima [A] (verification not implemented)

none

Time = 0.21 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.31

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^4} dx = \frac{1}{3x^3} + \frac{2(2a^4x^5 + a^2x^3 - 3x)\sqrt{ax+1}\sqrt{-ax+1}}{15a^2x^6} - \frac{2}{5a^2x^5}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^4,x, algorithm="maxima")

[Out] 1/3/x^3 + 2/15*(2*a^4*x^5 + a^2*x^3 - 3*x)*sqrt(a*x + 1)*sqrt(-a*x + 1)/(a^2*x^6) - 2/5/(a^2*x^5)

Giac [F]

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^4} dx = \int \frac{\left(\sqrt{\frac{1}{ax}+1}\sqrt{\frac{1}{ax}-1+\frac{1}{ax}}\right)^2}{x^4} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^4,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))^2/x^4, x)

Mupad [B] (verification not implemented)

Time = 5.42 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.47

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^4} dx = \frac{\sqrt{\frac{1}{ax} - 1} \left(\frac{2ax^2 \sqrt{\frac{1}{ax} + 1}}{15} - \frac{2\sqrt{\frac{1}{ax} + 1}}{5a} + \frac{4a^3 x^4 \sqrt{\frac{1}{ax} + 1}}{15} \right)}{x^4} + \frac{\frac{a^2 x^2}{3} - \frac{2}{5}}{a^2 x^5}$$

[In] int(((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))^2/x^4,x)

[Out] ((1/(a*x) - 1)^(1/2)*((2*a*x^2*(1/(a*x) + 1)^(1/2))/15 - (2*(1/(a*x) + 1)^(1/2))/(5*a) + (4*a^3*x^4*(1/(a*x) + 1)^(1/2))/15))/x^4 + ((a^2*x^2)/3 - 2/5)/(a^2*x^5)

3.74 $\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^5} dx$

Optimal result	471
Rubi [A] (verified)	471
Mathematica [A] (verified)	473
Maple [A] (verified)	474
Fricas [A] (verification not implemented)	474
Sympy [F]	474
Maxima [F]	475
Giac [F]	475
Mupad [B] (verification not implemented)	475

Optimal result

Integrand size = 12, antiderivative size = 267

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^5} dx = -\frac{2a^4}{3\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^6} + \frac{2a^4}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^5} - \frac{3a^4}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^4}$$

$$+ \frac{8a^4}{3\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3} - \frac{11a^4}{8\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{3a^4}{8\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)}$$

$$- \frac{a^4}{8\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{a^4}{8\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)} + \frac{1}{4}a^4 \operatorname{arctanh}\left(\sqrt{\frac{1-ax}{1+ax}}\right)$$

```
[Out] 1/4*a^4*arctanh(((a*x+1)/(a*x+1))^(1/2))-2/3*a^4/(1-((a*x+1)/(a*x+1))^(1/2))^6+2*a^4/(1-((a*x+1)/(a*x+1))^(1/2))^5-3*a^4/(1-((a*x+1)/(a*x+1))^(1/2))^4+8/3*a^4/(1-((a*x+1)/(a*x+1))^(1/2))^3-11/8*a^4/(1-((a*x+1)/(a*x+1))^(1/2))^2+3/8*a^4/(1-((a*x+1)/(a*x+1))^(1/2))-1/8*a^4/(1+((a*x+1)/(a*x+1))^(1/2))^2+1/8*a^4/(1+((a*x+1)/(a*x+1))^(1/2))
```

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used

= {6472, 1626, 213}

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^5} dx = \frac{1}{4}a^4 \operatorname{arctanh}\left(\sqrt{\frac{1-ax}{ax+1}}\right) + \frac{3a^4}{8\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)} + \frac{a^4}{8\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)} - \frac{11a^4}{8\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^2} - \frac{a^4}{8\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^2} + \frac{8a^4}{3\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^3} - \frac{3a^4}{\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^4} + \frac{2a^4}{\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^5} - \frac{2a^4}{3\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^6}$$

[In] Int[E^(2*ArcSech[a*x])/x^5,x]

[Out] (-2*a^4)/(3*(1 - Sqrt[(1 - a*x)/(1 + a*x)])^6) + (2*a^4)/(1 - Sqrt[(1 - a*x)/(1 + a*x)])^5 - (3*a^4)/(1 - Sqrt[(1 - a*x)/(1 + a*x)])^4 + (8*a^4)/(3*(1 - Sqrt[(1 - a*x)/(1 + a*x)])^3) - (11*a^4)/(8*(1 - Sqrt[(1 - a*x)/(1 + a*x)]))^2 + (3*a^4)/(8*(1 - Sqrt[(1 - a*x)/(1 + a*x)])) - a^4/(8*(1 + Sqrt[(1 - a*x)/(1 + a*x)]))^2 + a^4/(8*(1 + Sqrt[(1 - a*x)/(1 + a*x)])) + (a^4*ArcTanh[Sqrt[(1 - a*x)/(1 + a*x)]])/4

Rule 213

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 1626

Int[(Px)*((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)), x_Symbol] := Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && IntegersQ[m, n]

Rule 6472

Int[E^(ArcSech[u_]*(n_))*(x_)^(m_), x_Symbol] := Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && IntegerQ[n]

Rubi steps

$$\text{integral} = \int \frac{\left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax}\right)^2}{x^5} dx$$

$$\begin{aligned}
&= (4a)\text{Subst}\left(\int \frac{x(a+ax^2)^3}{(-1+x)^7(1+x)^3} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right) \\
&= (4a)\text{Subst}\left(\int \left(\frac{a^3}{(-1+x)^7} + \frac{5a^3}{2(-1+x)^6} + \frac{3a^3}{(-1+x)^5} + \frac{2a^3}{(-1+x)^4} + \frac{11a^3}{16(-1+x)^3}\right.\right. \\
&\quad \left.\left. + \frac{3a^3}{32(-1+x)^2} + \frac{a^3}{16(1+x)^3} - \frac{a^3}{32(1+x)^2} - \frac{a^3}{16(-1+x)^2}\right) dx, x, \sqrt{\frac{1-ax}{1+ax}}\right) \\
&= -\frac{2a^4}{3\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)^6} + \frac{2a^4}{\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)^5} - \frac{3a^4}{\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)^4} + \frac{8a^4}{3\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)^3} \\
&\quad - \frac{11a^4}{8\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{3a^4}{8\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)} - \frac{a^4}{8\left(1+\sqrt{\frac{1-ax}{1+ax}}\right)^2} \\
&\quad + \frac{a^4}{8\left(1+\sqrt{\frac{1-ax}{1+ax}}\right)} - \frac{1}{4}a^4\text{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right) \\
&= -\frac{2a^4}{3\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)^6} + \frac{2a^4}{\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)^5} - \frac{3a^4}{\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)^4} \\
&\quad + \frac{8a^4}{3\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)^3} - \frac{11a^4}{8\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{3a^4}{8\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)} \\
&\quad - \frac{a^4}{8\left(1+\sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{a^4}{8\left(1+\sqrt{\frac{1-ax}{1+ax}}\right)} + \frac{1}{4}a^4\text{arctanh}\left(\sqrt{\frac{1-ax}{1+ax}}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.51

$$\int \frac{e^{2\text{sech}^{-1}(ax)}}{x^5} dx = \frac{-8 + 6a^2x^2 + \sqrt{\frac{1-ax}{1+ax}}(-8 - 8ax + 2a^2x^2 + 2a^3x^3 + 3a^4x^4 + 3a^5x^5) - 3a^6x^6 \log(x) + 3a^6x^6 \log\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}{24a^2x^6}$$

[In] Integrate[E^(2*ArcSech[a*x])/x^5,x]

[Out] (-8 + 6*a^2*x^2 + Sqrt[(1 - a*x)/(1 + a*x)]*(-8 - 8*a*x + 2*a^2*x^2 + 2*a^3*x^3 + 3*a^4*x^4 + 3*a^5*x^5) - 3*a^6*x^6*Log[x] + 3*a^6*x^6*Log[1 + Sqrt[(1 - a*x)/(1 + a*x)]] + a*x*Sqrt[(1 - a*x)/(1 + a*x)])/(24*a^2*x^6)

Maxima [F]

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^5} dx = \int \frac{\left(\sqrt{\frac{1}{ax} + 1}\sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}\right)^2}{x^5} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^5,x, algorithm="maxima")

[Out] 2*integrate(sqrt(a*x + 1)*sqrt(-a*x + 1)/x^7, x)/a^2 - 1/3/(a^2*x^6) - integrate(x^(-5), x)

Giac [F]

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^5} dx = \int \frac{\left(\sqrt{\frac{1}{ax} + 1}\sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}\right)^2}{x^5} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^5,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))^2/x^5, x)

Mupad [B] (verification not implemented)

Time = 68.99 (sec) , antiderivative size = 2480, normalized size of antiderivative = 9.29

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^5} dx = \text{Too large to display}$$

[In] int(((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))^2/x^5,x)

[Out] ((311*a^4*((1/(a*x) - 1)^(1/2) - 1i)^5)/(2*((1/(a*x) + 1)^(1/2) - 1)^5) - (175*a^4*((1/(a*x) - 1)^(1/2) - 1i)^3)/(6*((1/(a*x) + 1)^(1/2) - 1)^3) + (8361*a^4*((1/(a*x) - 1)^(1/2) - 1i)^7)/(2*((1/(a*x) + 1)^(1/2) - 1)^7) + (42259*a^4*((1/(a*x) - 1)^(1/2) - 1i)^9)/(3*((1/(a*x) + 1)^(1/2) - 1)^9) + (25295*a^4*((1/(a*x) - 1)^(1/2) - 1i)^11)/((1/(a*x) + 1)^(1/2) - 1)^11 + (25295*a^4*((1/(a*x) - 1)^(1/2) - 1i)^13)/((1/(a*x) + 1)^(1/2) - 1)^13 + (42259*a^4*((1/(a*x) - 1)^(1/2) - 1i)^15)/(3*((1/(a*x) + 1)^(1/2) - 1)^15) + (8361*a^4*((1/(a*x) - 1)^(1/2) - 1i)^17)/(2*((1/(a*x) + 1)^(1/2) - 1)^17) + (311*a^4*((1/(a*x) - 1)^(1/2) - 1i)^19)/(2*((1/(a*x) + 1)^(1/2) - 1)^19) - (175*a^4*((1/(a*x) - 1)^(1/2) - 1i)^21)/(6*((1/(a*x) + 1)^(1/2) - 1)^21) + (5*a^4*((1/(a*x) - 1)^(1/2) - 1i)^23)/(2*((1/(a*x) + 1)^(1/2) - 1)^23) + (5*a^4*

$$\begin{aligned}
& \left(\frac{(1/(a*x) - 1)^{(1/2)} - 1i}{2*((1/(a*x) + 1)^{(1/2)} - 1)} \right) / \left(\frac{66*((1/(a*x) - 1)^{(1/2)} - 1i)^4}{((1/(a*x) + 1)^{(1/2)} - 1)^4 - (12*((1/(a*x) - 1)^{(1/2)} - 1i)^2)} \right) / \left(\frac{220*((1/(a*x) - 1)^{(1/2)} - 1i)^6}{((1/(a*x) + 1)^{(1/2)} - 1)^6 + (495*((1/(a*x) - 1)^{(1/2)} - 1i)^8)} \right) / \left(\frac{(1/(a*x) + 1)^{(1/2)} - 1)^8 - (792*((1/(a*x) - 1)^{(1/2)} - 1i)^{10})}{((1/(a*x) + 1)^{(1/2)} - 1)^{10} + (924*((1/(a*x) - 1)^{(1/2)} - 1i)^{12})} \right) / \left(\frac{(1/(a*x) + 1)^{(1/2)} - 1)^{14} - (792*((1/(a*x) - 1)^{(1/2)} - 1i)^{14})}{((1/(a*x) + 1)^{(1/2)} - 1)^{14} + (495*((1/(a*x) - 1)^{(1/2)} - 1i)^{16})} \right) / \left(\frac{(1/(a*x) - 1)^{(1/2)} - 1i)^{18}}{((1/(a*x) + 1)^{(1/2)} - 1)^{18} + (66*((1/(a*x) - 1)^{(1/2)} - 1i)^{20})} \right) / \left(\frac{(1/(a*x) + 1)^{(1/2)} - 1)^{22} - (12*((1/(a*x) - 1)^{(1/2)} - 1i)^{22})}{((1/(a*x) + 1)^{(1/2)} - 1)^{22} + ((1/(a*x) - 1)^{(1/2)} - 1i)^{24}} \right) / \left(\frac{(1/(a*x) + 1)^{(1/2)} - 1)^{24} + 1}{(a^4 * \operatorname{atanh}(((1/(a*x) - 1)^{(1/2)} - 1i) / ((1/(a*x) + 1)^{(1/2)} - 1)))} \right) / 2 - \left(\frac{a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^6 * 4096i}{3 * ((1/(a*x) + 1)^{(1/2)} - 1)^6} \right) + \left(\frac{a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^8 * 8192i}{3 * ((1/(a*x) + 1)^{(1/2)} - 1)^8} \right) + \left(\frac{a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{10} * 24576i}{5 * ((1/(a*x) + 1)^{(1/2)} - 1)^{10}} \right) + \left(\frac{a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{12} * 8192i}{3 * ((1/(a*x) + 1)^{(1/2)} - 1)^{12}} \right) + \left(\frac{a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{14} * 4096i}{3 * ((1/(a*x) + 1)^{(1/2)} - 1)^{14}} \right) / \left(\frac{45 * ((1/(a*x) - 1)^{(1/2)} - 1i)^4}{((1/(a*x) + 1)^{(1/2)} - 1)^4 - (10 * ((1/(a*x) - 1)^{(1/2)} - 1i)^2)} \right) / \left(\frac{(1/(a*x) + 1)^{(1/2)} - 1)^2 - (120 * ((1/(a*x) - 1)^{(1/2)} - 1i)^6)}{((1/(a*x) + 1)^{(1/2)} - 1)^6 + (210 * ((1/(a*x) - 1)^{(1/2)} - 1i)^8)} \right) / \left(\frac{(1/(a*x) + 1)^{(1/2)} - 1)^8 - (252 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{10})}{((1/(a*x) + 1)^{(1/2)} - 1)^{10} + (210 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{12})} \right) / \left(\frac{(1/(a*x) + 1)^{(1/2)} - 1)^{12} - (120 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{14})}{((1/(a*x) + 1)^{(1/2)} - 1)^{14} + (45 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{16})} \right) / \left(\frac{(1/(a*x) + 1)^{(1/2)} - 1)^{16} - (10 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{18})}{((1/(a*x) + 1)^{(1/2)} - 1)^{18} + ((1/(a*x) - 1)^{(1/2)} - 1i)^{20}} \right) / \left(\frac{(1/(a*x) + 1)^{(1/2)} - 1)^{20} + 1}{(a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^6 * 20480i)} \right) / \left(\frac{(1/(a*x) + 1)^{(1/2)} - 1)^6 + (a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^8 * 40960i)}{((1/(a*x) + 1)^{(1/2)} - 1)^8 + (a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{10} * 73728i)} \right) / \left(\frac{(1/(a*x) + 1)^{(1/2)} - 1)^{10} + (a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{12} * 40960i)}{((1/(a*x) + 1)^{(1/2)} - 1)^{12} + (a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{14} * 20480i)} \right) / \left(\frac{(1/(a*x) + 1)^{(1/2)} - 1)^{14}}{(15 * ((45 * ((1/(a*x) - 1)^{(1/2)} - 1i)^4) / ((1/(a*x) + 1)^{(1/2)} - 1)^4 - (10 * ((1/(a*x) - 1)^{(1/2)} - 1i)^2) / ((1/(a*x) + 1)^{(1/2)} - 1)^2 - (120 * ((1/(a*x) - 1)^{(1/2)} - 1i)^6) / ((1/(a*x) + 1)^{(1/2)} - 1)^6 + (210 * ((1/(a*x) - 1)^{(1/2)} - 1i)^8) / ((1/(a*x) + 1)^{(1/2)} - 1)^8 - (252 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{10}) / ((1/(a*x) + 1)^{(1/2)} - 1)^{10} + (210 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{12}) / ((1/(a*x) + 1)^{(1/2)} - 1)^{12} - (120 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{14}) / ((1/(a*x) + 1)^{(1/2)} - 1)^{14} + (45 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{16}) / ((1/(a*x) + 1)^{(1/2)} - 1)^{16} - (10 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{18}) / ((1/(a*x) + 1)^{(1/2)} - 1)^{18} + ((1/(a*x) - 1)^{(1/2)} - 1i)^{20} / ((1/(a*x) + 1)^{(1/2)} - 1)^{20} + 1)} \right) + \left(\frac{23 * a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^3}{((1/(a*x) + 1)^{(1/2)} - 1)^3} \right) + \left(\frac{333 * a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^5}{((1/(a*x) + 1)^{(1/2)} - 1)^5} \right) + \left(\frac{671 * a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^7}{((1/(a*x) + 1)^{(1/2)} - 1)^7} \right) + \left(\frac{671 * a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^9}{((1/(a*x) + 1)^{(1/2)} - 1)^9} \right) + \left(\frac{333 * a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{11}}{((1/(a*x) + 1)^{(1/2)} - 1)^{11}} \right) + \left(\frac{23 * a^4 * ((1/(a*x) - 1)^{(1/2)} - 1i)^{13}}{((1/(a*x) + 1)^{(1/2)} - 1)^{13}} \right)
\end{aligned}$$

$$\begin{aligned}
& /((1/(a*x) + 1)^{(1/2)} - 1)^{13} - (3*a^4*((1/(a*x) - 1)^{(1/2)} - 1i)^{15})/((1/(a*x) + 1)^{(1/2)} - 1)^{15} - (3*a^4*((1/(a*x) - 1)^{(1/2)} - 1i))/((1/(a*x) + 1)^{(1/2)} - 1) \\
& /((28*((1/(a*x) - 1)^{(1/2)} - 1i)^4)/((1/(a*x) + 1)^{(1/2)} - 1)^4 - (8*((1/(a*x) - 1)^{(1/2)} - 1i)^2)/((1/(a*x) + 1)^{(1/2)} - 1)^2 - (56*((1/(a*x) - 1)^{(1/2)} - 1i)^6)/((1/(a*x) + 1)^{(1/2)} - 1)^6 + (70*((1/(a*x) - 1)^{(1/2)} - 1i)^8)/((1/(a*x) + 1)^{(1/2)} - 1)^8 - (56*((1/(a*x) - 1)^{(1/2)} - 1i)^{10})/((1/(a*x) + 1)^{(1/2)} - 1)^{10} + (28*((1/(a*x) - 1)^{(1/2)} - 1i)^{12})/((1/(a*x) + 1)^{(1/2)} - 1)^{12} - (8*((1/(a*x) - 1)^{(1/2)} - 1i)^{14})/((1/(a*x) + 1)^{(1/2)} - 1)^{14} + ((1/(a*x) - 1)^{(1/2)} - 1i)^{16}/((1/(a*x) + 1)^{(1/2)} - 1)^{16} \\
& + 1) + 1/(4*x^4) - 1/(3*a^2*x^6)
\end{aligned}$$

3.75 $\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^6} dx$

Optimal result	478
Rubi [A] (verified)	478
Mathematica [A] (verified)	480
Maple [A] (verified)	480
Fricas [A] (verification not implemented)	481
Sympy [F]	481
Maxima [A] (verification not implemented)	481
Giac [F]	482
Mupad [B] (verification not implemented)	482

Optimal result

Integrand size = 12, antiderivative size = 301

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^6} dx = -\frac{4a^5}{7\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^7} + \frac{2a^5}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^6} - \frac{18a^5}{5\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^5} + \frac{4a^5}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^4}$$

$$- \frac{35a^5}{12\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3} + \frac{11a^5}{8\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} - \frac{a^5}{4\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)}$$

$$- \frac{a^5}{12\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^3} + \frac{a^5}{8\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^2} - \frac{a^5}{4\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}$$

[Out] $-4/7*a^5/(1-((-a*x+1)/(a*x+1))^(1/2))^7+2*a^5/(1-((-a*x+1)/(a*x+1))^(1/2))^6-18/5*a^5/(1-((-a*x+1)/(a*x+1))^(1/2))^5+4*a^5/(1-((-a*x+1)/(a*x+1))^(1/2))^4-35/12*a^5/(1-((-a*x+1)/(a*x+1))^(1/2))^3+11/8*a^5/(1-((-a*x+1)/(a*x+1))^(1/2))^2-1/4*a^5/(1-((-a*x+1)/(a*x+1))^(1/2))-1/12*a^5/(1+((-a*x+1)/(a*x+1))^(1/2))^3+1/8*a^5/(1+((-a*x+1)/(a*x+1))^(1/2))^2-1/4*a^5/(1+((-a*x+1)/(a*x+1))^(1/2))$

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 301, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used

= {6472, 1626}

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^6} dx = -\frac{a^5}{4\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)} - \frac{a^5}{4\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)} + \frac{11a^5}{8\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^2}$$

$$+ \frac{a^5}{8\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^2} - \frac{35a^5}{12\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^3} - \frac{a^5}{12\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^3}$$

$$+ \frac{4a^5}{\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^4} - \frac{18a^5}{5\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^5} + \frac{2a^5}{\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^6} - \frac{4a^5}{7\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^7}$$

[In] Int[E^(2*ArcSech[a*x])/x^6,x]

[Out] (-4*a^5)/(7*(1 - Sqrt[(1 - a*x)/(1 + a*x)])^7) + (2*a^5)/(1 - Sqrt[(1 - a*x)/(1 + a*x)])^6 - (18*a^5)/(5*(1 - Sqrt[(1 - a*x)/(1 + a*x)])^5) + (4*a^5)/(1 - Sqrt[(1 - a*x)/(1 + a*x)])^4 - (35*a^5)/(12*(1 - Sqrt[(1 - a*x)/(1 + a*x)])^3) + (11*a^5)/(8*(1 - Sqrt[(1 - a*x)/(1 + a*x)])^2) - a^5/(4*(1 - Sqrt[(1 - a*x)/(1 + a*x)])) - a^5/(12*(1 + Sqrt[(1 - a*x)/(1 + a*x)])^3) + a^5/(8*(1 + Sqrt[(1 - a*x)/(1 + a*x)])^2) - a^5/(4*(1 + Sqrt[(1 - a*x)/(1 + a*x)]))

Rule 1626

Int[(Px_)*((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && IntegersQ[m, n]

Rule 6472

Int[E^(ArcSech[u_]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && IntegerQ[n]

Rubi steps

$$\text{integral} = \int \frac{\left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \sqrt{\frac{1-ax}{1+ax}}\right)^2}{x^6} dx$$

$$= -\left((4a)\text{Subst}\left(\int \frac{x(a+ax^2)^4}{(-1+x)^8(1+x)^4} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)\right)$$

$$= -\left((4a)\text{Subst}\left(\int \left(\frac{a^4}{(-1+x)^8} + \frac{3a^4}{(-1+x)^7} + \frac{9a^4}{2(-1+x)^6} + \frac{4a^4}{(-1+x)^5} + \frac{35a^4}{16(-1+x)^4} + \frac{11a^4}{16(-1+x)^3} + \frac{a^4}{16(-1+x)^2} + \frac{a^4}{16(-1+x)} + \frac{a^4}{16}\right) dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)\right)$$

$$\begin{aligned}
&= -\frac{4a^5}{7\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^7} + \frac{2a^5}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^6} - \frac{18a^5}{5\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^5} + \frac{4a^5}{\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^4} \\
&\quad - \frac{35a^5}{12\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3} + \frac{11a^5}{8\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} - \frac{a^5}{4\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} \\
&\quad - \frac{a^5}{12\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^3} + \frac{a^5}{8\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^2} - \frac{a^5}{4\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.28

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^6} dx = \frac{-30 + 21a^2x^2 + 2\sqrt{\frac{1-ax}{1+ax}}(1+ax)^2(-15 + 15ax - 12a^2x^2 + 12a^3x^3 - 8a^4x^4 + 8a^5x^5)}{105a^2x^7}$$

[In] Integrate[E^(2*ArcSech[a*x])/x^6,x]

[Out] (-30 + 21*a^2*x^2 + 2*sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2*(-15 + 15*a*x - 12*a^2*x^2 + 12*a^3*x^3 - 8*a^4*x^4 + 8*a^5*x^5))/(105*a^2*x^7)

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.31

method	result	size
default	$-\frac{1}{7x^7} + \frac{a^2}{5x^5} + \frac{2\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}(a^2x^2-1)(8a^4x^4+12a^2x^2+15)}{105ax^6} - \frac{1}{7a^2x^7}$	92

[In] int((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^6,x,method=_RETURNVERBOSE)

[Out] 1/a^2*(-1/7/x^7+1/5*a^2/x^5)+2/105/a*((a*x+1)/a/x)^(1/2)/x^6*(-(a*x-1)/a/x)^(1/2)*(a^2*x^2-1)*(8*a^4*x^4+12*a^2*x^2+15)-1/7/a^2/x^7

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.26

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^6} dx = \frac{21 a^2 x^2 + 2 (8 a^7 x^7 + 4 a^5 x^5 + 3 a^3 x^3 - 15 a x) \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} - 30}{105 a^2 x^7}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^6,x, algorithm="fricas")

[Out] 1/105*(21*a^2*x^2 + 2*(8*a^7*x^7 + 4*a^5*x^5 + 3*a^3*x^3 - 15*a*x)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 30)/(a^2*x^7)

Sympy [F]

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^6} dx = \frac{\int \frac{2}{x^8} dx + \int \left(-\frac{a^2}{x^6}\right) dx + \int \frac{2a\sqrt{-1+\frac{1}{ax}}\sqrt{1+\frac{1}{ax}}}{x^7} dx}{a^2}$$

[In] integrate((1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))**2/x**6,x)

[Out] (Integral(2/x**8, x) + Integral(-a**2/x**6, x) + Integral(2*a*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x))/x**7, x))/a**2

Maxima [A] (verification not implemented)

none

Time = 0.21 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.22

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^6} dx = \frac{1}{5 x^5} + \frac{2 (8 a^6 x^7 + 4 a^4 x^5 + 3 a^2 x^3 - 15 x) \sqrt{ax+1} \sqrt{-ax+1}}{105 a^2 x^8} - \frac{2}{7 a^2 x^7}$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^6,x, algorithm="maxima")

[Out] 1/5/x^5 + 2/105*(8*a^6*x^7 + 4*a^4*x^5 + 3*a^2*x^3 - 15*x)*sqrt(a*x + 1)*sqrt(-a*x + 1)/(a^2*x^8) - 2/7/(a^2*x^7)

Giac [F]

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^6} dx = \int \frac{\left(\sqrt{\frac{1}{ax} + 1}\sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}\right)^2}{x^6} dx$$

[In] integrate((1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))^2/x^6,x, algorithm="giac")

[Out] integrate((sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))^2/x^6, x)

Mupad [B] (verification not implemented)

Time = 5.34 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.35

$$\int \frac{e^{2\operatorname{sech}^{-1}(ax)}}{x^6} dx = \frac{\frac{a^2 x^2}{5} - \frac{2}{7}}{a^2 x^7} + \frac{\sqrt{\frac{1}{ax} - 1} \left(\frac{2 a x^2 \sqrt{\frac{1}{ax} + 1}}{35} - \frac{2 \sqrt{\frac{1}{ax} + 1}}{7 a} + \frac{8 a^3 x^4 \sqrt{\frac{1}{ax} + 1}}{105} + \frac{16 a^5 x^6 \sqrt{\frac{1}{ax} + 1}}{105} \right)}{x^6}$$

[In] int(((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))^2/x^6,x)

[Out] ((a^2*x^2)/5 - 2/7)/(a^2*x^7) + ((1/(a*x) - 1)^(1/2)*((2*a*x^2*(1/(a*x) + 1)^(1/2))/35 - (2*(1/(a*x) + 1)^(1/2))/(7*a) + (8*a^3*x^4*(1/(a*x) + 1)^(1/2))/105 + (16*a^5*x^6*(1/(a*x) + 1)^(1/2))/105))/x^6

3.76 $\int e^{-\operatorname{sech}^{-1}(ax)} x^4 dx$

Optimal result	483
Rubi [A] (verified)	483
Mathematica [A] (verified)	485
Maple [C] (verified)	486
Fricas [A] (verification not implemented)	486
Sympy [F]	487
Maxima [F]	487
Giac [F]	487
Mupad [B] (verification not implemented)	487

Optimal result

Integrand size = 12, antiderivative size = 147

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^4 dx = -\frac{x}{a^4} - \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^5}{5a^5} + \frac{(1+ax)^2 \left(9 + 4\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^5} + \frac{(1+ax)^4 \left(5 + 16\sqrt{\frac{1-ax}{1+ax}}\right)}{20a^5} - \frac{(1+ax)^3 \left(15 + 17\sqrt{\frac{1-ax}{1+ax}}\right)}{15a^5}$$

[Out] $-x/a^4 - 1/5*(a*x+1)^5*((-a*x+1)/(a*x+1))^{(1/2)}/a^5 + 1/6*(a*x+1)^2*(9+4*((-a*x+1)/(a*x+1))^{(1/2)})/a^5 + 1/20*(a*x+1)^4*(5+16*((-a*x+1)/(a*x+1))^{(1/2)})/a^5 - 1/15*(a*x+1)^3*(15+17*((-a*x+1)/(a*x+1))^{(1/2)})/a^5$

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {6472, 1818, 1828, 12, 267}

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^4 dx = -\frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)^5}{5a^5} + \frac{\left(16\sqrt{\frac{1-ax}{ax+1}} + 5\right)(ax+1)^4}{20a^5} - \frac{\left(17\sqrt{\frac{1-ax}{ax+1}} + 15\right)(ax+1)^3}{15a^5} + \frac{\left(4\sqrt{\frac{1-ax}{ax+1}} + 9\right)(ax+1)^2}{6a^5} - \frac{x}{a^4}$$

[In] $\text{Int}[x^4/E^{\text{ArcSech}[a*x]}, x]$

[Out] $-(x/a^4) - (\text{Sqrt}[(1 - a*x)/(1 + a*x)]*(1 + a*x)^5)/(5*a^5) + ((1 + a*x)^2*(9 + 4*\text{Sqrt}[(1 - a*x)/(1 + a*x)]))/(6*a^5) + ((1 + a*x)^4*(5 + 16*\text{Sqrt}[(1 -$

$a*x)/(1 + a*x]])))/(20*a^5) - ((1 + a*x)^3*(15 + 17*sqrt[(1 - a*x)/(1 + a*x]])))/(15*a^5)$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]`

Rule 267

`Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^p_, x_Symbol] := Simp[(a + b*x^n)^(p + 1)/(b*m*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]`

Rule 1818

`Int[(Pq_)*((c_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^p_), x_Symbol] := With[{Q = PolynomialQuotient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(c*x)^m*(a + b*x^2)^(p + 1)*((a*g - b*f*x)/(2*a*b*(p + 1))), x] + Dist[c/(2*a*b*(p + 1)), Int[(c*x)^(m - 1)*(a + b*x^2)^(p + 1)*ExpandToSum[2*a*b*(p + 1)*x*Q - a*g*m + b*f*(m + 2*p + 3)*x, x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x] && LtQ[p, -1] && GtQ[m, 0]`

Rule 1828

`Int[(Pq_)*((a_) + (b_.)*(x_)^2)^p_, x_Symbol] := With[{Q = PolynomialQuotient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(a*g - b*f*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Dist[1/(2*a*(p + 1)), Int[(a + b*x^2)^(p + 1)*ExpandToSum[2*a*(p + 1)*Q + f*(2*p + 3), x], x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && LtQ[p, -1]`

Rule 6472

`Int[E^(ArcSech[u_]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && IntegerQ[n]`

Rubi steps

$$\begin{aligned} \text{integral} &= \int \frac{x^4}{\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax}} dx \\ &= \frac{4 \text{Subst}\left(\int \frac{(-1+x)^5 x(1+x)^3}{(1+x^2)^6} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a^5} \end{aligned}$$

$$\begin{aligned}
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^5}{5a^5} - \frac{2\text{Subst}\left(\int \frac{-16+10x+140x^2-30x^3-80x^4+30x^5+20x^6-10x^7}{(1+x^2)^5} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{5a^5} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^5}{5a^5} + \frac{(1+ax)^4\left(5+16\sqrt{\frac{1-ax}{1+ax}}\right)}{20a^5} \\
&\quad + \frac{\text{Subst}\left(\int \frac{-128+560x+800x^2-320x^3-160x^4+80x^5}{(1+x^2)^4} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{20a^5} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^5}{5a^5} + \frac{(1+ax)^4\left(5+16\sqrt{\frac{1-ax}{1+ax}}\right)}{20a^5} - \frac{(1+ax)^3\left(15+17\sqrt{\frac{1-ax}{1+ax}}\right)}{15a^5} \\
&\quad - \frac{\text{Subst}\left(\int \frac{-320+2400x+960x^2-480x^3}{(1+x^2)^3} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{120a^5} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^5}{5a^5} + \frac{(1+ax)^2\left(9+4\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^5} + \frac{(1+ax)^4\left(5+16\sqrt{\frac{1-ax}{1+ax}}\right)}{20a^5} \\
&\quad - \frac{(1+ax)^3\left(15+17\sqrt{\frac{1-ax}{1+ax}}\right)}{15a^5} + \frac{\text{Subst}\left(\int \frac{1920x}{(1+x^2)^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{480a^5} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^5}{5a^5} + \frac{(1+ax)^2\left(9+4\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^5} + \frac{(1+ax)^4\left(5+16\sqrt{\frac{1-ax}{1+ax}}\right)}{20a^5} \\
&\quad - \frac{(1+ax)^3\left(15+17\sqrt{\frac{1-ax}{1+ax}}\right)}{15a^5} + \frac{4\text{Subst}\left(\int \frac{x}{(1+x^2)^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a^5} \\
&= -\frac{x}{a^4} - \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^5}{5a^5} + \frac{(1+ax)^2\left(9+4\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^5} \\
&\quad + \frac{(1+ax)^4\left(5+16\sqrt{\frac{1-ax}{1+ax}}\right)}{20a^5} - \frac{(1+ax)^3\left(15+17\sqrt{\frac{1-ax}{1+ax}}\right)}{15a^5}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.44

$$\int e^{-\text{sech}^{-1}(ax)} x^4 dx = \frac{15a^4 x^4 - 4\sqrt{\frac{1-ax}{1+ax}}(1+ax)^2(-2+2ax-3a^2x^2+3a^3x^3)}{60a^5}$$

[In] Integrate[x^4/E^ArcSech[a*x], x]

[Out] (15*a^4*x^4 - 4*sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2*(-2 + 2*a*x - 3*a^2*x^2 + 3*a^3*x^3))/(60*a^5)

Maple [C] (verified)

Result contains higher order function than in optimal. Order 3 vs. order 2.

Time = 0.35 (sec) , antiderivative size = 531, normalized size of antiderivative = 3.61

method	result
default	$(ax+1) \left(15 \left(-\frac{ax-1}{ax} \right)^{\frac{7}{2}} \left(\frac{ax+1}{ax} \right)^{\frac{5}{2}} x^{10} a^{10} + 30 \left(-\frac{ax-1}{ax} \right)^{\frac{7}{2}} \left(\frac{ax+1}{ax} \right)^{\frac{5}{2}} x^8 a^8 - 30 \left(-\frac{ax-1}{ax} \right)^{\frac{7}{2}} \left(\frac{ax+1}{ax} \right)^{\frac{3}{2}} a^8 x^8 + 30 \left(-\frac{ax-1}{ax} \right)^{\frac{7}{2}} \ln(a^2 x^2) \left(\frac{ax+1}{ax} \right)^{\frac{5}{2}} \right)$

[In] `int(x^4/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{60} \cdot \frac{(ax+1)^7}{x^7} \cdot \left(15 \left(-\frac{ax-1}{ax} \right)^{\frac{7}{2}} \left(\frac{ax+1}{ax} \right)^{\frac{5}{2}} x^{10} a^{10} + 30 \left(-\frac{ax-1}{ax} \right)^{\frac{7}{2}} \left(\frac{ax+1}{ax} \right)^{\frac{5}{2}} x^8 a^8 - 30 \left(-\frac{ax-1}{ax} \right)^{\frac{7}{2}} \left(\frac{ax+1}{ax} \right)^{\frac{3}{2}} a^8 x^8 + 30 \left(-\frac{ax-1}{ax} \right)^{\frac{7}{2}} \ln(a^2 x^2) \left(\frac{ax+1}{ax} \right)^{\frac{5}{2}} \right)$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.44

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^4 dx = \frac{15 a^3 x^4 - 4 (3 a^4 x^5 - a^2 x^3 - 2 x) \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}}}{60 a^4}$$

[In] `integrate(x^4/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x, algorithm="fricas")`

[Out] $\frac{1}{60} \cdot \frac{(15 a^3 x^4 - 4 (3 a^4 x^5 - a^2 x^3 - 2 x) \sqrt{(ax+1)/(ax)} \sqrt{-(ax-1)/(ax)})}{a^4}$

Sympy [F]

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^4 dx = a \int \frac{x^5}{ax \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} + 1} dx$$

[In] integrate(x**4/(1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2)),x)

[Out] a*Integral(x**5/(a*x*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)) + 1), x)

Maxima [F]

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^4 dx = \int \frac{x^4}{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}}} dx$$

[In] integrate(x^4/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x, algorithm="maxima")

[Out] integrate(x^4/(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x)), x)

Giac [F]

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^4 dx = \int \frac{x^4}{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}}} dx$$

[In] integrate(x^4/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x, algorithm="giac")

[Out] integrate(x^4/(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x)), x)

Mupad [B] (verification not implemented)

Time = 5.35 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.50

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^4 dx = \frac{x^4}{4a} + \frac{\sqrt{\frac{1}{ax} - 1} \left(\frac{2x}{15a^4} + \frac{2}{15a^5} - \frac{x^5}{5} - \frac{x^4}{5a} + \frac{x^3}{15a^2} + \frac{x^2}{15a^3} \right)}{\sqrt{\frac{1}{ax} + 1}}$$

[In] int(x^4/((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x)),x)

[Out] x^4/(4*a) + ((1/(a*x) - 1)^(1/2)*((2*x)/(15*a^4) + 2/(15*a^5) - x^5/5 - x^4/(5*a) + x^3/(15*a^2) + x^2/(15*a^3)))/(1/(a*x) + 1)^(1/2)

3.77 $\int e^{-\operatorname{sech}^{-1}(ax)} x^3 dx$

Optimal result	488
Rubi [A] (verified)	488
Mathematica [C] (verified)	490
Maple [C] (verified)	491
Fricas [A] (verification not implemented)	491
Sympy [F]	491
Maxima [F]	492
Giac [F]	492
Mupad [B] (verification not implemented)	492

Optimal result

Integrand size = 12, antiderivative size = 163

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^3 dx = -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^4}{4a^4} + \frac{(1+ax)\left(8 + \sqrt{\frac{1-ax}{1+ax}}\right)}{8a^4} - \frac{(1+ax)^2\left(8 + 5\sqrt{\frac{1-ax}{1+ax}}\right)}{8a^4} + \frac{(1+ax)^3\left(4 + 9\sqrt{\frac{1-ax}{1+ax}}\right)}{12a^4} + \frac{\arctan\left(\sqrt{\frac{1-ax}{1+ax}}\right)}{4a^4}$$

[Out] $\frac{1}{4} \arctan\left(\frac{(-ax+1)}{(ax+1)}\right)^{(1/2)} / a^4 - \frac{1}{4} (ax+1)^4 \left(\frac{(-ax+1)}{(ax+1)}\right)^{(1/2)} / a^4 + \frac{1}{8} (ax+1) \left(8 + \left(\frac{(-ax+1)}{(ax+1)}\right)^{(1/2)}\right) / a^4 - \frac{1}{8} (ax+1)^2 \left(8 + 5 \left(\frac{(-ax+1)}{(ax+1)}\right)^{(1/2)}\right) / a^4 + \frac{1}{12} (ax+1)^3 \left(4 + 9 \left(\frac{(-ax+1)}{(ax+1)}\right)^{(1/2)}\right) / a^4$

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {6472, 1818, 1828, 653, 209}

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^3 dx = \frac{\arctan\left(\sqrt{\frac{1-ax}{ax+1}}\right)}{4a^4} - \frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)^4}{4a^4} + \frac{\left(9\sqrt{\frac{1-ax}{ax+1}} + 4\right)(ax+1)^3}{12a^4} - \frac{\left(5\sqrt{\frac{1-ax}{ax+1}} + 8\right)(ax+1)^2}{8a^4} + \frac{\left(\sqrt{\frac{1-ax}{ax+1}} + 8\right)(ax+1)}{8a^4}$$

[In] Int[x^3/E^ArcSech[a*x],x]


```
[Out] -1/4*(Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^4)/a^4 + ((1 + a*x)*(8 + Sqrt[(1 - a*x)/(1 + a*x)]))/(8*a^4) - ((1 + a*x)^2*(8 + 5*Sqrt[(1 - a*x)/(1 + a*x)]))/(8*a^4) + ((1 + a*x)^3*(4 + 9*Sqrt[(1 - a*x)/(1 + a*x)]))/(12*a^4) + ArcTan[Sqrt[(1 - a*x)/(1 + a*x)]]/(4*a^4)
```

Rule 209

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 653

```
Int[((d_) + (e_)*(x_))*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[((a*e - c*d*x)/(2*a*c*(p + 1)))*(a + c*x^2)^(p + 1), x] + Dist[d*((2*p + 3)/(2*a*(p + 1))), Int[(a + c*x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e}, x] && LtQ[p, -1] && NeQ[p, -3/2]
```

Rule 1818

```
Int[(Pq_)*((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(c*x)^m*(a + b*x^2)^(p + 1)*((a*g - b*f*x)/(2*a*b*(p + 1))), x] + Dist[c/(2*a*b*(p + 1)), Int[(c*x)^(m - 1)*(a + b*x^2)^(p + 1)*ExpandToSum[2*a*b*(p + 1)*x*Q - a*g*m + b*f*(m + 2*p + 3)*x, x], x], x]] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x] && LtQ[p, -1] && GtQ[m, 0]
```

Rule 1828

```
Int[(Pq_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(a*g - b*f*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Dist[1/(2*a*(p + 1)), Int[(a + b*x^2)^(p + 1)*ExpandToSum[2*a*(p + 1)*Q + f*(2*p + 3), x], x], x]] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && LtQ[p, -1]
```

Rule 6472

```
Int[E^(ArcSech[u_]*(n_))*(x_)^(m_), x_Symbol] := Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && IntegerQ[n]
```

Rubi steps

$$\text{integral} = \int \frac{x^3}{\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax}} dx$$

$$\begin{aligned}
&= -\frac{4\text{Subst}\left(\int \frac{(-1+x)^4 x(1+x)^2}{(1+x^2)^5} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a^4} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^4}{4a^4} + \frac{\text{Subst}\left(\int \frac{8-8x-48x^2+16x^3+16x^4-8x^5}{(1+x^2)^4} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{2a^4} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^4}{4a^4} + \frac{(1+ax)^3\left(4+9\sqrt{\frac{1-ax}{1+ax}}\right)}{12a^4} - \frac{\text{Subst}\left(\int \frac{24-144x-96x^2+48x^3}{(1+x^2)^3} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{12a^4} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^4}{4a^4} - \frac{(1+ax)^2\left(8+5\sqrt{\frac{1-ax}{1+ax}}\right)}{8a^4} \\
&\quad + \frac{(1+ax)^3\left(4+9\sqrt{\frac{1-ax}{1+ax}}\right)}{12a^4} + \frac{\text{Subst}\left(\int \frac{24-192x}{(1+x^2)^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{48a^4} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^4}{4a^4} + \frac{(1+ax)\left(8+\sqrt{\frac{1-ax}{1+ax}}\right)}{8a^4} - \frac{(1+ax)^2\left(8+5\sqrt{\frac{1-ax}{1+ax}}\right)}{8a^4} \\
&\quad + \frac{(1+ax)^3\left(4+9\sqrt{\frac{1-ax}{1+ax}}\right)}{12a^4} + \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{4a^4} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^4}{4a^4} + \frac{(1+ax)\left(8+\sqrt{\frac{1-ax}{1+ax}}\right)}{8a^4} - \frac{(1+ax)^2\left(8+5\sqrt{\frac{1-ax}{1+ax}}\right)}{8a^4} \\
&\quad + \frac{(1+ax)^3\left(4+9\sqrt{\frac{1-ax}{1+ax}}\right)}{12a^4} + \frac{\arctan\left(\sqrt{\frac{1-ax}{1+ax}}\right)}{4a^4}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.60

$$\begin{aligned}
&\int e^{-\text{sech}^{-1}(ax)} x^3 dx \\
&= \frac{8a^3 x^3 + 3a\sqrt{\frac{1-ax}{1+ax}}(x + ax^2 - 2a^2 x^3 - 2a^3 x^4) - 3i \log\left(-2iax + 2\sqrt{\frac{1-ax}{1+ax}}(1+ax)\right)}{24a^4}
\end{aligned}$$

[In] Integrate[x^3/E^ArcSech[a*x],x]

[Out] (8*a^3*x^3 + 3*a*Sqrt[(1 - a*x)/(1 + a*x)]*(x + a*x^2 - 2*a^2*x^3 - 2*a^3*x^4) - (3*I)*Log[(-2*I)*a*x + 2*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)]/(24*a^4)

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.92 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.74

method	result	size
default	$a \left(\frac{x^3}{3a^2} - \frac{\sqrt{-\frac{ax-1}{ax}} x \sqrt{\frac{ax+1}{ax}} \left(2 \operatorname{csgn}(a) a^3 x^3 \sqrt{-a^2 x^2 + 1} - \sqrt{-a^2 x^2 + 1} x \operatorname{csgn}(a) a + \arctan \left(\frac{\operatorname{csgn}(a) ax}{\sqrt{-a^2 x^2 + 1}} \right) \right) \operatorname{csgn}(a)}{8a^4 \sqrt{-a^2 x^2 + 1}} \right)$	120

[In] `int(x^3/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x,method=_RETURNVERBOSE)`

[Out] `a*(1/3*x^3/a^2-1/8/a^4*(-(a*x-1)/a/x)^(1/2)*x*((a*x+1)/a/x)^(1/2)*(2*csgn(a)*a^3*x^3*(-a^2*x^2+1)^(1/2)-(-a^2*x^2+1)^(1/2)*x*csgn(a)*a+arctan(csgn(a)*a*x/(-a^2*x^2+1)^(1/2)))*csgn(a)/(-a^2*x^2+1)^(1/2))`

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.58

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^3 dx = \frac{8a^3 x^3 - 3(2a^4 x^4 - a^2 x^2) \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} + 3 \arctan \left(\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} \right)}{24a^4}$$

[In] `integrate(x^3/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x, algorithm="fricas")`

[Out] `1/24*(8*a^3*x^3 - 3*(2*a^4*x^4 - a^2*x^2)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + 3*arctan(sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x))))/a^4`

Sympy [F]

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^3 dx = a \int \frac{x^4}{ax \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} + 1} dx$$

[In] `integrate(x**3/(1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2)),x)`

[Out] `a*Integral(x**4/(a*x*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)) + 1), x)`

Maxima [F]

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^3 dx = \int \frac{x^3}{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}} dx$$

[In] integrate(x^3/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x, algorithm="maxima")

[Out] integrate(x^3/(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x)), x)

Giac [F]

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^3 dx = \int \frac{x^3}{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax}} dx$$

[In] integrate(x^3/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x, algorithm="giac")

[Out] integrate(x^3/(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x)), x)

Mupad [B] (verification not implemented)

Time = 25.41 (sec) , antiderivative size = 795, normalized size of antiderivative = 4.88

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^3 dx = \frac{\ln\left(\frac{a\sqrt{\frac{1}{ax}+1}-\frac{1}{x}+a\sqrt{\frac{1}{ax}-1}i}{2a-2a\sqrt{\frac{1}{ax}+1+\frac{1}{x}}}\right) 3i}{8a^4} + \frac{\frac{1i}{1024a^4} - \frac{(\sqrt{\frac{1}{ax}-1-i})^2 3i}{128a^4(\sqrt{\frac{1}{ax}+1-1})^2} - \frac{(\sqrt{\frac{1}{ax}-1-i})^4 53i}{512a^4(\sqrt{\frac{1}{ax}+1-1})^4} + \frac{(\sqrt{\frac{1}{ax}-1-i})^6 87i}{256a^4(\sqrt{\frac{1}{ax}+1-1})^6} + \frac{(\sqrt{\frac{1}{ax}-1-i})^8 657i}{1024a^4(\sqrt{\frac{1}{ax}+1-1})^8} + \frac{(\sqrt{\frac{1}{ax}-1-i})^{10} 121i}{256a^4(\sqrt{\frac{1}{ax}+1-1})^{10}}}{\frac{(\sqrt{\frac{1}{ax}-1-i})^4}{(\sqrt{\frac{1}{ax}+1-1})^4} + \frac{4(\sqrt{\frac{1}{ax}-1-i})^6}{(\sqrt{\frac{1}{ax}+1-1})^6} + \frac{6(\sqrt{\frac{1}{ax}-1-i})^8}{(\sqrt{\frac{1}{ax}+1-1})^8} + \frac{4(\sqrt{\frac{1}{ax}-1-i})^{10}}{(\sqrt{\frac{1}{ax}+1-1})^{10}} + \frac{(\sqrt{\frac{1}{ax}-1-i})^{12}}{(\sqrt{\frac{1}{ax}+1-1})^{12}}} + \frac{\ln\left(\frac{\sqrt{\frac{1}{ax}-1-i}}{\sqrt{\frac{1}{ax}+1-1}}\right) 1i}{8a^4} + \frac{1i}{32a^4} + \frac{(\sqrt{\frac{1}{ax}-1-i})^2 1i}{16a^4(\sqrt{\frac{1}{ax}+1-1})^2} - \frac{(\sqrt{\frac{1}{ax}-1-i})^4 15i}{32a^4(\sqrt{\frac{1}{ax}+1-1})^4}}{\frac{(\sqrt{\frac{1}{ax}-1-i})^2}{(\sqrt{\frac{1}{ax}+1-1})^2} + \frac{2(\sqrt{\frac{1}{ax}-1-i})^4}{(\sqrt{\frac{1}{ax}+1-1})^4} + \frac{(\sqrt{\frac{1}{ax}-1-i})^6}{(\sqrt{\frac{1}{ax}+1-1})^6}} - \frac{\ln\left(\frac{2a\sqrt{\frac{a+\frac{1}{x}}{a}-\frac{2}{x}}+a\sqrt{-\frac{a-\frac{1}{x}}{a}} 2i}{2a+\frac{1}{x}-2a\sqrt{\frac{a+\frac{1}{x}}{a}}}\right) 1i}{2a^4} + \frac{x^3}{3a} + \frac{(\sqrt{\frac{1}{ax}-1-i})^2 1i}{256a^4(\sqrt{\frac{1}{ax}+1-1})^2} + \frac{(\sqrt{\frac{1}{ax}-1-i})^4 1i}{1024a^4(\sqrt{\frac{1}{ax}+1-1})^4}$$

[In] $\text{int}(x^3/((1/(a*x) - 1)^{(1/2)}*(1/(a*x) + 1)^{(1/2)} + 1/(a*x)),x)$

[Out] $(\log((a*(1/(a*x) - 1)^{(1/2)}*1i + a*(1/(a*x) + 1)^{(1/2)} - 1/x)/(2*a - 2*a*(1/(a*x) + 1)^{(1/2)} + 1/x))*3i)/(8*a^4) + (1i/(1024*a^4) - (((1/(a*x) - 1)^{(1/2)} - 1i)^2*3i)/(128*a^4*((1/(a*x) + 1)^{(1/2)} - 1)^2) - (((1/(a*x) - 1)^{(1/2)} - 1i)^4*53i)/(512*a^4*((1/(a*x) + 1)^{(1/2)} - 1)^4) + (((1/(a*x) - 1)^{(1/2)} - 1i)^6*87i)/(256*a^4*((1/(a*x) + 1)^{(1/2)} - 1)^6) + (((1/(a*x) - 1)^{(1/2)} - 1i)^8*657i)/(1024*a^4*((1/(a*x) + 1)^{(1/2)} - 1)^8) + (((1/(a*x) - 1)^{(1/2)} - 1i)^{10}*121i)/(256*a^4*((1/(a*x) + 1)^{(1/2)} - 1)^{10}))/(((1/(a*x) - 1)^{(1/2)} - 1i)^4/((1/(a*x) + 1)^{(1/2)} - 1)^4 + (4*((1/(a*x) - 1)^{(1/2)} - 1i)^6)/((1/(a*x) + 1)^{(1/2)} - 1)^6 + (6*((1/(a*x) - 1)^{(1/2)} - 1i)^8)/((1/(a*x) + 1)^{(1/2)} - 1)^8 + (4*((1/(a*x) - 1)^{(1/2)} - 1i)^{10})/((1/(a*x) + 1)^{(1/2)} - 1)^{10} + ((1/(a*x) - 1)^{(1/2)} - 1i)^{12}/((1/(a*x) + 1)^{(1/2)} - 1)^{12}) + (\log(((1/(a*x) - 1)^{(1/2)} - 1i)/((1/(a*x) + 1)^{(1/2)} - 1))*1i)/(8*a^4) + (1i/(32*a^4) + (((1/(a*x) - 1)^{(1/2)} - 1i)^2*1i)/(16*a^4*((1/(a*x) + 1)^{(1/2)} - 1)^2) - (((1/(a*x) - 1)^{(1/2)} - 1i)^4*15i)/(32*a^4*((1/(a*x) + 1)^{(1/2)} - 1)^4))/(((1/(a*x) - 1)^{(1/2)} - 1i)^2/((1/(a*x) + 1)^{(1/2)} - 1)^2 + (2*((1/(a*x) - 1)^{(1/2)} - 1i)^4)/((1/(a*x) + 1)^{(1/2)} - 1)^4 + ((1/(a*x) - 1)^{(1/2)} - 1i)^6/((1/(a*x) + 1)^{(1/2)} - 1)^6) - (\log((a*(-(a - 1/x)/a)^{(1/2)}*2i - 2/x + 2*a*((a + 1/x)/a)^{(1/2)}))/(2*a + 1/x - 2*a*((a + 1/x)/a)^{(1/2}))*1i)/(2*a^4) + x^3/(3*a) + (((1/(a*x) - 1)^{(1/2)} - 1i)^2*1i)/(256*a^4*((1/(a*x) + 1)^{(1/2)} - 1)^2) + (((1/(a*x) - 1)^{(1/2)} - 1i)^4*1i)/(1024*a^4*((1/(a*x) + 1)^{(1/2)} - 1)^4)$

3.78 $\int e^{-\operatorname{sech}^{-1}(ax)} x^2 dx$

Optimal result	494
Rubi [A] (verified)	494
Mathematica [A] (verified)	496
Maple [C] (verified)	496
Fricas [A] (verification not implemented)	496
Sympy [F]	497
Maxima [F]	497
Giac [F]	497
Mupad [B] (verification not implemented)	498

Optimal result

Integrand size = 12, antiderivative size = 75

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^2 dx = -\frac{x}{a^2} - \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^3}{3a^3} + \frac{(1+ax)^2 \left(3 + 4\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^3}$$

[Out] $-x/a^2 - 1/3*(a*x+1)^3*((-a*x+1)/(a*x+1))^{(1/2)}/a^3 + 1/6*(a*x+1)^2*(3+4*((-a*x+1)/(a*x+1))^{(1/2)})/a^3$

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {6472, 1818, 1828, 12, 267}

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^2 dx = -\frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)^3}{3a^3} + \frac{\left(4\sqrt{\frac{1-ax}{ax+1}} + 3\right)(ax+1)^2}{6a^3} - \frac{x}{a^2}$$

[In] $\text{Int}[x^2/E^{\text{ArcSech}[a*x]}, x]$

[Out] $-(x/a^2) - (\text{Sqrt}[(1 - a*x)/(1 + a*x)]*(1 + a*x)^3)/(3*a^3) + ((1 + a*x)^2*(3 + 4*\text{Sqrt}[(1 - a*x)/(1 + a*x)]))/(6*a^3)$

Rule 12

$\text{Int}[(a_*)(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /;$ FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 267

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n)
^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] &&
NeQ[p, -1]
```

Rule 1818

```
Int[(Pq_)*((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[
{Q = PolynomialQuotient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq
, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x,
1]}, Simp[(c*x)^m*(a + b*x^2)^(p + 1)*((a*g - b*f*x)/(2*a*b*(p + 1))), x]
+ Dist[c/(2*a*b*(p + 1)), Int[(c*x)^(m - 1)*(a + b*x^2)^(p + 1)*ExpandToSum
[2*a*b*(p + 1)*x*Q - a*g*m + b*f*(m + 2*p + 3)*x, x], x]] /; FreeQ[{a,
b, c}, x] && PolyQ[Pq, x] && LtQ[p, -1] && GtQ[m, 0]
```

Rule 1828

```
Int[(Pq_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuot
ient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x,
0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(a*g - b
*f*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Dist[1/(2*a*(p + 1)), Int
[(a + b*x^2)^(p + 1)*ExpandToSum[2*a*(p + 1)*Q + f*(2*p + 3), x], x], x]] /
; FreeQ[{a, b}, x] && PolyQ[Pq, x] && LtQ[p, -1]
```

Rule 6472

```
Int[E^(ArcSech[u_]*(n_))*(x_)^(m_), x_Symbol] := Int[x^m*(1/u + Sqrt[(1 -
u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && Integer
Q[n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \frac{x^2}{\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{1-ax}}{ax}} dx \\
&= \frac{4 \text{Subst}\left(\int \frac{(-1+x)^3 x(1+x)}{(1+x^2)^4} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a^3} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^3}{3a^3} - \frac{2 \text{Subst}\left(\int \frac{-4+6x+12x^2-6x^3}{(1+x^2)^3} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{3a^3} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^3}{3a^3} + \frac{(1+ax)^2 \left(3 + 4\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^3} + \frac{\text{Subst}\left(\int \frac{24x}{(1+x^2)^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{6a^3} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^3}{3a^3} + \frac{(1+ax)^2 \left(3 + 4\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^3} + \frac{4 \text{Subst}\left(\int \frac{x}{(1+x^2)^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a^3}
\end{aligned}$$

$$= -\frac{x}{a^2} - \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)^3}{3a^3} + \frac{(1+ax)^2 \left(3 + 4\sqrt{\frac{1-ax}{1+ax}}\right)}{6a^3}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.64

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^2 dx = \frac{3a^2 x^2 - 2(-1+ax)\sqrt{\frac{1-ax}{1+ax}}(1+ax)^2}{6a^3}$$

[In] Integrate[x^2/E^ArcSech[a*x],x]

[Out] (3*a^2*x^2 - 2*(-1 + a*x)*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2)/(6*a^3)

Maple [C] (verified)

Result contains higher order function than in optimal. Order 3 vs. order 2.

Time = 0.29 (sec) , antiderivative size = 269, normalized size of antiderivative = 3.59

method	result
default	$\frac{(ax+1)\left(3a^6 x^6 \left(\frac{ax+1}{ax}\right)^{\frac{3}{2}} \left(-\frac{ax-1}{ax}\right)^{\frac{5}{2}} + 3\left(-\frac{ax-1}{ax}\right)^{\frac{5}{2}} \ln(a^2 x^2) \left(\frac{ax+1}{ax}\right)^{\frac{3}{2}} x^4 a^4 - 3\left(-\frac{ax-1}{ax}\right)^{\frac{5}{2}} \sqrt{\frac{ax+1}{ax}} \ln(a^2 x^2) a^4 x^4 + 2a^7 x^7 - 3x^3 \ln(a^2 x^2)\right)}{6x^5 a^8 \left(\frac{ax+1}{ax}\right)^{\frac{5}{2}} \left(-\frac{ax-1}{ax}\right)^{\frac{5}{2}}}$

[In] int(x^2/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x,method=_RETURNVERBOSE)

[Out] 1/6*(a*x+1)/x^5*(3*a^6*x^6*((a*x+1)/a/x)^(3/2)*(-(a*x-1)/a/x)^(5/2)+3*(-(a*x-1)/a/x)^(5/2)*ln(a^2*x^2)*((a*x+1)/a/x)^(3/2)*x^4*a^4-3*(-(a*x-1)/a/x)^(5/2)*((a*x+1)/a/x)^(1/2)*ln(a^2*x^2)*a^4*x^4+2*a^7*x^7-3*x^3*ln(a^2*x^2)*((a*x+1)/a/x)^(1/2)*(-(a*x-1)/a/x)^(5/2)*a^3-2*a^6*x^6-6*a^5*x^5+6*a^4*x^4+6*a^3*x^3-6*a^2*x^2-2*a*x+2)/a^8/((a*x+1)/a/x)^(5/2)/(-(a*x-1)/a/x)^(5/2)

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.72

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^2 dx = \frac{3ax^2 - 2(a^2x^3 - x)\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}}{6a^2}$$

[In] integrate(x^2/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x, algorithm="fricas")

[Out] $\frac{1}{6} \cdot (3ax^2 - 2(a^2x^3 - x) \sqrt{\frac{ax+1}{ax}} \sqrt{\frac{-ax-1}{ax}}) / a^2$

Sympy [F]

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^2 dx = a \int \frac{x^3}{ax \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} + 1} dx$$

[In] `integrate(x**2/(1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2)),x)`

[Out] `a*Integral(x**3/(a*x*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)) + 1), x)`

Maxima [F]

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^2 dx = \int \frac{x^2}{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}}} dx$$

[In] `integrate(x^2/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x, algorithm="maxima")`

[Out] `integrate(x^2/(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x)), x)`

Giac [F]

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^2 dx = \int \frac{x^2}{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}}} dx$$

[In] `integrate(x^2/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x, algorithm="giac")`

[Out] `integrate(x^2/(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x)), x)`

Mupad [B] (verification not implemented)

Time = 4.89 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.76

$$\int e^{-\operatorname{sech}^{-1}(ax)} x^2 dx = \frac{x^2}{2a} + \frac{\sqrt{\frac{1}{ax} - 1} \left(\frac{x}{3a^2} + \frac{1}{3a^3} - \frac{x^3}{3} - \frac{x^2}{3a} \right)}{\sqrt{\frac{1}{ax} + 1}}$$

[In] int(x^2/((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x)),x)

[Out] x^2/(2*a) + ((1/(a*x) - 1)^(1/2)*(x/(3*a^2) + 1/(3*a^3) - x^3/3 - x^2/(3*a)))/(1/(a*x) + 1)^(1/2)

3.79 $\int e^{-\operatorname{sech}^{-1}(ax)} x dx$

Optimal result	499
Rubi [A] (verified)	499
Mathematica [C] (verified)	501
Maple [C] (verified)	501
Fricas [A] (verification not implemented)	501
Sympy [F]	502
Maxima [F]	502
Giac [F]	502
Mupad [B] (verification not implemented)	503

Optimal result

Integrand size = 10, antiderivative size = 94

$$\int e^{-\operatorname{sech}^{-1}(ax)} x dx = \frac{(1+ax)^2 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2}{4a^2} + \frac{(1+ax) \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}{2a^2} + \frac{\arctan\left(\sqrt{\frac{1-ax}{1+ax}}\right)}{a^2}$$

[Out] $\arctan\left(\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)/a^2+1/4*(a*x+1)^2*(1-\left(\frac{-a*x+1}{a*x+1}\right)^{1/2})^2/a^2+1/2*(a*x+1)*(1+\left(\frac{-a*x+1}{a*x+1}\right)^{1/2})/a^2$

Rubi [A] (verified)

Time = 0.20 (sec), antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {6472, 833, 653, 209}

$$\int e^{-\operatorname{sech}^{-1}(ax)} x dx = \frac{\arctan\left(\sqrt{\frac{1-ax}{ax+1}}\right)}{a^2} + \frac{(ax+1)^2 \left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^2}{4a^2} + \frac{(ax+1) \left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)}{2a^2}$$

[In] $\text{Int}[x/E^{\text{ArcSech}[a*x]}, x]$

[Out] $\left(\frac{(1+a*x)^2*(1-\text{Sqrt}[(1-a*x)/(1+a*x)])^2}{4*a^2} + \frac{(1+a*x)*(1+\text{Sqrt}[(1-a*x)/(1+a*x)])}{2*a^2} + \text{ArcTan}[\text{Sqrt}[(1-a*x)/(1+a*x)]]/a^2\right)$

Rule 209

$\text{Int}[\left(\frac{a}{x} + \frac{b}{x^2}\right)^{-1}, x_Symbol] := \text{Simp}\left[\frac{1}{\text{Rt}[a, 2]*\text{Rt}[b, 2]}\right]*\text{ArcTan}\left[\frac{\text{Rt}[b, 2]*x}{\text{Rt}[a, 2]}\right], x] /;$ FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 653

Int[((d_) + (e_)*(x_))*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[((a*e - c*d*x)/(2*a*c*(p + 1)))*(a + c*x^2)^(p + 1), x] + Dist[d*((2*p + 3)/(2*a*(p + 1))), Int[(a + c*x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e}, x] && LtQ[p, -1] && NeQ[p, -3/2]

Rule 833

Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m - 1)*(a + c*x^2)^(p + 1)*((a*(e*f + d*g) - (c*d*f - a*e*g)*x)/(2*a*c*(p + 1))), x] - Dist[1/(2*a*c*(p + 1)), Int[(d + e*x)^(m - 2)*(a + c*x^2)^(p + 1)*Simp[a*e*(e*f*(m - 1) + d*g*m) - c*d^2*f*(2*p + 3) + e*(a*e*g*m - c*d*f*(m + 2*p + 2))*x, x], x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && GtQ[m, 1] && (EqQ[d, 0] || (EqQ[m, 2] && EqQ[p, -3] && RationalQ[a, c, d, e, f, g]) || !ILtQ[m + 2*p + 3, 0])

Rule 6472

Int[E^(ArcSech[u_]*(n_))*(x_)^(m_), x_Symbol] := Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && IntegerQ[n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{x}{\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax}} dx \\
 &= -\frac{4\text{Subst}\left(\int \frac{(-1+x)^2 x}{(1+x^2)^3} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a^2} \\
 &= \frac{(1+ax)^2 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2}{4a^2} - \frac{\text{Subst}\left(\int \frac{-2+2x}{(1+x^2)^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a^2} \\
 &= \frac{(1+ax)^2 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2}{4a^2} + \frac{(1+ax) \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}{2a^2} + \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a^2} \\
 &= \frac{(1+ax)^2 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2}{4a^2} + \frac{(1+ax) \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}{2a^2} + \frac{\arctan\left(\sqrt{\frac{1-ax}{1+ax}}\right)}{a^2}
 \end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.80

$$\int e^{-\operatorname{sech}^{-1}(ax)} x dx = -\frac{-2ax + ax\sqrt{\frac{1-ax}{1+ax}}(1+ax) + i \log\left(-2iax + 2\sqrt{\frac{1-ax}{1+ax}}(1+ax)\right)}{2a^2}$$

[In] Integrate[x/E^ArcSech[a*x],x]

[Out] $-1/2*(-2*a*x + a*x*\operatorname{Sqrt}[(1 - a*x)/(1 + a*x)]*(1 + a*x) + I*\operatorname{Log}[(-2*I)*a*x + 2*\operatorname{Sqrt}[(1 - a*x)/(1 + a*x)]*(1 + a*x)])/a^2$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.88 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.00

method	result	size
default	$a\left(\frac{x}{a^2} - \frac{\sqrt{\frac{ax+1}{ax}} x \sqrt{-\frac{ax-1}{ax}} \left(\sqrt{-a^2x^2+1} x \operatorname{csgn}(a) a + \arctan\left(\frac{\operatorname{csgn}(a) ax}{\sqrt{-a^2x^2+1}}\right)\right) \operatorname{csgn}(a)}{2a^2\sqrt{-a^2x^2+1}}\right)$	94

[In] int(x/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x,method=_RETURNVERBOSE)

[Out] $a*(x/a^2-1/2/a^2*((a*x+1)/a/x)^(1/2)*x*(-(a*x-1)/a/x)^(1/2)*((-a^2*x^2+1)^(1/2)*x*\operatorname{csgn}(a)*a+\arctan(\operatorname{csgn}(a)*a*x/(-a^2*x^2+1)^(1/2)))/(-a^2*x^2+1)^(1/2)*\operatorname{csgn}(a))$

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.84

$$\int e^{-\operatorname{sech}^{-1}(ax)} x dx = -\frac{a^2x^2\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} - 2ax - \arctan\left(\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}\right)}{2a^2}$$

[In] integrate(x/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x, algorithm="fricas")

[Out] $-1/2*(a^2*x^2*\operatorname{sqrt}((a*x + 1)/(a*x))*\operatorname{sqrt}(-(a*x - 1)/(a*x)) - 2*a*x - \arctan(\operatorname{sqrt}((a*x + 1)/(a*x))*\operatorname{sqrt}(-(a*x - 1)/(a*x))))/a^2$

Sympy [F]

$$\int e^{-\operatorname{sech}^{-1}(ax)} x dx = a \int \frac{x^2}{ax \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} + 1} dx$$

[In] integrate(x/(1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2)),x)

[Out] a*Integral(x**2/(a*x*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)) + 1), x)

Maxima [F]

$$\int e^{-\operatorname{sech}^{-1}(ax)} x dx = \int \frac{x}{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}}} dx$$

[In] integrate(x/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x, algorithm="maxima")

[Out] integrate(x/(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x)), x)

Giac [F]

$$\int e^{-\operatorname{sech}^{-1}(ax)} x dx = \int \frac{x}{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}}} dx$$

[In] integrate(x/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2)),x, algorithm="giac")

[Out] integrate(x/(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x)), x)

Mupad [B] (verification not implemented)

Time = 13.36 (sec) , antiderivative size = 407, normalized size of antiderivative = 4.33

$$\int e^{-\operatorname{sech}^{-1}(ax)} x dx = \frac{x}{a} - \frac{\ln\left(\frac{\sqrt{\frac{1}{ax}-1-i}}{\sqrt{\frac{1}{ax}+1-1}}\right) \operatorname{li}}{2a^2} - \frac{\frac{\operatorname{li}}{32a^2} + \frac{(\sqrt{\frac{1}{ax}-1-i})^2 \operatorname{li}}{16a^2(\sqrt{\frac{1}{ax}+1-1})^2} - \frac{(\sqrt{\frac{1}{ax}-1-i})^4 15i}{32a^2(\sqrt{\frac{1}{ax}+1-1})^4}}{\frac{(\sqrt{\frac{1}{ax}-1-i})^2}{(\sqrt{\frac{1}{ax}+1-1})^2} + \frac{2(\sqrt{\frac{1}{ax}-1-i})^4}{(\sqrt{\frac{1}{ax}+1-1})^4} + \frac{(\sqrt{\frac{1}{ax}-1-i})^6}{(\sqrt{\frac{1}{ax}+1-1})^6}}$$

$$- \frac{\left(\ln\left(\frac{(\sqrt{\frac{1}{ax}-1-i})^2}{(\sqrt{\frac{1}{ax}+1-1})^2} + 1\right) - \ln\left(\frac{\sqrt{\frac{1}{ax}-1-i}}{\sqrt{\frac{1}{ax}+1-1}}\right)\right) \operatorname{li}}{a^2}$$

$$+ \frac{\ln\left(\frac{2a\sqrt{\frac{a+\frac{1}{x}}{a}} - \frac{2}{x} + a\sqrt{-\frac{a-\frac{1}{x}}{a}} 2i}{2a+\frac{1}{x} - 2a\sqrt{\frac{a+\frac{1}{x}}{a}}}\right) \operatorname{li}}{2a^2} - \frac{(\sqrt{\frac{1}{ax}-1-i})^2 \operatorname{li}}{32a^2(\sqrt{\frac{1}{ax}+1-1})^2}$$

[In] int(x/((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x)),x)

[Out] x/a - (log(((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1))*1i)/(2*a^2) - (1i/(32*a^2) + (((1/(a*x) - 1)^(1/2) - 1i)^2*1i)/(16*a^2*((1/(a*x) + 1)^(1/2) - 1)^2) - (((1/(a*x) - 1)^(1/2) - 1i)^4*15i)/(32*a^2*((1/(a*x) + 1)^(1/2) - 1)^4))/(((1/(a*x) - 1)^(1/2) - 1i)^2/((1/(a*x) + 1)^(1/2) - 1)^2 + (2*((1/(a*x) - 1)^(1/2) - 1i)^4)/((1/(a*x) + 1)^(1/2) - 1)^4 + ((1/(a*x) - 1)^(1/2) - 1i)^6/((1/(a*x) + 1)^(1/2) - 1)^6) - ((log(((1/(a*x) - 1)^(1/2) - 1i)^2/((1/(a*x) + 1)^(1/2) - 1)^2 + 1) - log(((1/(a*x) - 1)^(1/2) - 1i)/(1/(a*x) + 1)^(1/2) - 1))*1i)/a^2 + (log((a*(-a - 1/x)/a)^(1/2)*2i - 2/x + 2*a*((a + 1/x)/a)^(1/2)))/(2*a + 1/x - 2*a*((a + 1/x)/a)^(1/2))*1i)/(2*a^2) - (((1/(a*x) - 1)^(1/2) - 1i)^2*1i)/(32*a^2*((1/(a*x) + 1)^(1/2) - 1)^2)

3.80 $\int e^{-\operatorname{sech}^{-1}(ax)} dx$

Optimal result	504
Rubi [A] (verified)	504
Mathematica [A] (verified)	506
Maple [B] (verified)	506
Fricas [A] (verification not implemented)	507
Sympy [F]	508
Maxima [F]	508
Giac [F]	508
Mupad [B] (verification not implemented)	509

Optimal result

Integrand size = 8, antiderivative size = 65

$$\int e^{-\operatorname{sech}^{-1}(ax)} dx = -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)}{a} + \frac{\log(1+ax)}{a} + \frac{2 \log\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}{a}$$

[Out] $\ln(a*x+1)/a+2*\ln(1+((-a*x+1)/(a*x+1))^(1/2))/a-(a*x+1)*((-a*x+1)/(a*x+1))^(1/2)/a$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6467, 1661, 815, 266}

$$\int e^{-\operatorname{sech}^{-1}(ax)} dx = -\frac{\sqrt{\frac{1-ax}{ax+1}}(ax+1)}{a} + \frac{\log(ax+1)}{a} + \frac{2 \log\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)}{a}$$

[In] $\text{Int}[E^{(-\text{ArcSech}[a*x])}, x]$

[Out] $-\left(\frac{\text{Sqrt}[(1 - a*x)/(1 + a*x)]*(1 + a*x)}{a}\right) + \text{Log}[1 + a*x]/a + \frac{2*\text{Log}[1 + \text{Sqrt}[(1 - a*x)/(1 + a*x)]]}{a}$

Rule 266

$\text{Int}[(x_)^{(m_.)}/((a_) + (b_.)*(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^n, x]]/(b*n), x] /; \text{FreeQ}\{a, b, m, n, x\} \ \&\& \ \text{EqQ}[m, n - 1]$

Rule 815


```
Int[(((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_)))/((a_.) + (c_.)*(x_)^2),
  x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)/(a + c*x^2)), x],
  x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[m]
```

Rule 1661

```
Int[(Pq_)*((d_) + (e_.)*(x_))^(m_.)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] :
> With[{Q = PolynomialQuotient[(d + e*x)^m*Pq, a + c*x^2, x], f = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + c*x^2, x], x, 0], g = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + c*x^2, x], x, 1]}, Simp[(a*g - c*f*x)*((a + c*x^2)^(p + 1)/(2*a*c*(p + 1))), x] + Dist[1/(2*a*c*(p + 1)), Int[(d + e*x)^m*(a + c*x^2)^(p + 1)*ExpandToSum[(2*a*c*(p + 1)*Q)/(d + e*x)^m + (c*f*(2*p + 3))/(d + e*x)^m, x], x]] /; FreeQ[{a, c, d, e}, x] && PolyQ[Pq, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && ILtQ[m, 0]
```

Rule 6467

```
Int[E^(ArcSech[u_]*(n_.)), x_Symbol] :> Int[(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \frac{1}{\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax}} dx \\
&= \frac{4 \text{Subst}\left(\int \frac{(-1+x)x}{(1+x)(1+x^2)^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)}{a} - \frac{2 \text{Subst}\left(\int \frac{-1+x}{(1+x)(1+x^2)} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)}{a} - \frac{2 \text{Subst}\left(\int \left(\frac{1}{-1-x} + \frac{x}{1+x^2}\right) dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)}{a} + \frac{2 \log\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}{a} - \frac{2 \text{Subst}\left(\int \frac{x}{1+x^2} dx, x, \sqrt{\frac{1-ax}{1+ax}}\right)}{a} \\
&= -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)}{a} + \frac{\log(1+ax)}{a} + \frac{2 \log\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}{a}
\end{aligned}$$

[Out] $-1/2*(2*a*x*\sqrt{(a*x + 1)/(a*x)}*\sqrt{-(a*x - 1)/(a*x)} - \log(a*x*\sqrt{(a*x + 1)/(a*x)}*\sqrt{-(a*x - 1)/(a*x)} + 1) + \log(a*x*\sqrt{(a*x + 1)/(a*x)}*\sqrt{-(a*x - 1)/(a*x)} - 1) - 2*\log(x))/a$

Sympy [F]

$$\int e^{-\operatorname{sech}^{-1}(ax)} dx = a \int \frac{x}{ax \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax} + 1}} dx$$

[In] `integrate(1/(1/a/x+(1/a/x-1)**(1/2))*(1+1/a/x)**(1/2)),x)`

[Out] `a*Integral(x/(a*x*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)) + 1), x)`

Maxima [F]

$$\int e^{-\operatorname{sech}^{-1}(ax)} dx = \int \frac{1}{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}}} dx$$

[In] `integrate(1/(1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2)),x, algorithm="maxima")`

[Out] `integrate(1/(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x)), x)`

Giac [F]

$$\int e^{-\operatorname{sech}^{-1}(ax)} dx = \int \frac{1}{\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}}} dx$$

[In] `integrate(1/(1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2)),x, algorithm="giac")`

[Out] `integrate(1/(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x)), x)`

Mupad [B] (verification not implemented)

Time = 7.36 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.72

$$\int e^{-\operatorname{sech}^{-1}(ax)} dx = \frac{\operatorname{acosh}\left(\frac{1}{ax}\right)}{a} - \frac{\ln\left(\frac{1}{x}\right)}{a} - x \sqrt{\frac{1}{ax} - 1} \sqrt{\frac{1}{ax} + 1}$$

[In] int(1/((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x)),x)

[Out] acosh(1/(a*x))/a - log(1/x)/a - x*(1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2)

3.81 $\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x} dx$

Optimal result	510
Rubi [A] (verified)	510
Mathematica [C] (verified)	511
Maple [C] (verified)	512
Fricas [A] (verification not implemented)	512
Sympy [F]	512
Maxima [F]	513
Giac [F]	513
Mupad [B] (verification not implemented)	513

Optimal result

Integrand size = 12, antiderivative size = 46

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x} dx = -\frac{2}{1 + \sqrt{\frac{1-ax}{1+ax}}} - 2 \arctan \left(\sqrt{\frac{1-ax}{1+ax}} \right)$$

[Out] $-2*\arctan(((-a*x+1)/(a*x+1))^{(1/2)})-2/(1+((-a*x+1)/(a*x+1))^{(1/2)})$

Rubi [A] (verified)

Time = 0.28 (sec), antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6472, 815, 209}

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x} dx = -2 \arctan \left(\sqrt{\frac{1-ax}{ax+1}} \right) - \frac{2}{\sqrt{\frac{1-ax}{ax+1}} + 1}$$

[In] `Int[1/(E^ArcSech[a*x]*x),x]`

[Out] `-2/(1 + Sqrt[(1 - a*x)/(1 + a*x)]) - 2*ArcTan[Sqrt[(1 - a*x)/(1 + a*x)]]`

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 815

`Int[(((d_) + (e_.)*(x_)^m)*((f_) + (g_.)*(x_)))/((a_) + (c_.)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)/(a + c*x^2)), x],`

$x] /; \text{FreeQ}\{a, c, d, e, f, g\}, x] \&\& \text{NeQ}[c*d^2 + a*e^2, 0] \&\& \text{IntegerQ}[m]$

Rule 6472

$\text{Int}[E^{\text{ArcSech}[u_]*(n_.)}*(x_)^{(m_.)}, x_Symbol] \text{ :> Int}[x^m*(1/u + \text{Sqrt}[(1 - u)/(1 + u)] + (1/u)*\text{Sqrt}[(1 - u)/(1 + u)])^n, x] /; \text{FreeQ}[m, x] \&\& \text{IntegerQ}[n]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{1}{x \left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax} \right)} dx \\
 &= - \left(4 \text{Subst} \left(\int \frac{x}{(1+x)^2(1+x^2)} dx, x, \sqrt{\frac{1-ax}{1+ax}} \right) \right) \\
 &= - \left(4 \text{Subst} \left(\int \left(-\frac{1}{2(1+x)^2} + \frac{1}{2(1+x^2)} \right) dx, x, \sqrt{\frac{1-ax}{1+ax}} \right) \right) \\
 &= -\frac{2}{1 + \sqrt{\frac{1-ax}{1+ax}}} - 2 \text{Subst} \left(\int \frac{1}{1+x^2} dx, x, \sqrt{\frac{1-ax}{1+ax}} \right) \\
 &= -\frac{2}{1 + \sqrt{\frac{1-ax}{1+ax}}} - 2 \arctan \left(\sqrt{\frac{1-ax}{1+ax}} \right)
 \end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.04 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.61

$$\int \frac{e^{-\text{sech}^{-1}(ax)}}{x} dx = -\frac{1}{ax} + \left(1 + \frac{1}{ax} \right) \sqrt{\frac{1-ax}{1+ax}} + i \log \left(-2iax + 2\sqrt{\frac{1-ax}{1+ax}}(1+ax) \right)$$

[In] Integrate[1/(E^ArcSech[a*x]*x), x]

[Out] -(1/(a*x)) + (1 + 1/(a*x))*Sqrt[(1 - a*x)/(1 + a*x)] + I*Log[(-2*I)*a*x + 2*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)]

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.89 (sec) , antiderivative size = 96, normalized size of antiderivative = 2.09

method	result	size
default	$a \left(-\frac{1}{x a^2} + \frac{\sqrt{-\frac{ax-1}{ax}} \sqrt{\frac{ax+1}{ax}} \left(\arctan\left(\frac{\operatorname{csgn}(a)ax}{\sqrt{-a^2x^2+1}}\right) ax + \operatorname{csgn}(a)\sqrt{-a^2x^2+1} \right) \operatorname{csgn}(a)}{a\sqrt{-a^2x^2+1}} \right)$	96

[In] int(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x,x,method=_RETURNVERBOSE)

[Out] a*(-1/x/a^2+1/a*(-(a*x-1)/a/x)^(1/2)*((a*x+1)/a/x)^(1/2)*(arctan(csgn(a)*a*x/(-a^2*x^2+1)^(1/2))*a*x+csgn(a)*(-a^2*x^2+1)^(1/2))*csgn(a)/(-a^2*x^2+1)^(1/2))

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.65

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x} dx = \frac{ax \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} - ax \arctan\left(\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}}\right) - 1}{ax}$$

[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x,x, algorithm="fricas")

[Out] (a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - a*x*arctan(sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x))) - 1)/(a*x)

Sympy [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x} dx = a \int \frac{1}{ax \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} + 1} dx$$

[In] integrate(1/(1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))/x,x)

[Out] a*Integral(1/(a*x*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)) + 1), x)

Maxima [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x} dx = \int \frac{1}{x \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}} \right)} dx$$

[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x,x, algorithm="maxima")

[Out] integrate(1/(x*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))), x)

Giac [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x} dx = \int \frac{1}{x \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}} \right)} dx$$

[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x,x, algorithm="giac")

[Out] integrate(1/(x*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))), x)

Mupad [B] (verification not implemented)

Time = 7.48 (sec) , antiderivative size = 184, normalized size of antiderivative = 4.00

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x} dx = \ln \left(\frac{\left(\sqrt{\frac{1}{ax} - 1 - i} \right)^2}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^2} + 1 \right) \operatorname{li} - \ln \left(\frac{\sqrt{\frac{1}{ax} - 1 - i}}{\sqrt{\frac{1}{ax} + 1 - 1}} \right) \operatorname{li} - \frac{1}{ax} - \frac{\left(\sqrt{\frac{1}{ax} - 1 - i} \right)^2 8i}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^2 \left(1 + \frac{\left(\sqrt{\frac{1}{ax} - 1 - i} \right)^4}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^4} - \frac{2 \left(\sqrt{\frac{1}{ax} - 1 - i} \right)^2}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^2} \right)}$$

[In] int(1/(x*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))),x)

[Out] log(((1/(a*x) - 1)^(1/2) - 1i)^2/((1/(a*x) + 1)^(1/2) - 1)^2 + 1)*1i - log(((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1))*1i - 1/(a*x) - (((1/(a*x) - 1)^(1/2) - 1i)^2*8i)/(((1/(a*x) + 1)^(1/2) - 1)^2*((1/(a*x) - 1)^(1/2) - 1i)^4/((1/(a*x) + 1)^(1/2) - 1)^4 - (2*((1/(a*x) - 1)^(1/2) - 1i)^2)/((1/(a*x) + 1)^(1/2) - 1)^2 + 1))

3.82 $\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^2} dx$

Optimal result	514
Rubi [A] (verified)	514
Mathematica [A] (verified)	515
Maple [A] (verified)	516
Fricas [A] (verification not implemented)	516
Sympy [F]	517
Maxima [F]	517
Giac [F]	517
Mupad [B] (verification not implemented)	517

Optimal result

Integrand size = 12, antiderivative size = 72

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^2} dx = -\frac{a}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{a}{1 + \sqrt{\frac{1-ax}{1+ax}}} - a \operatorname{arctanh}\left(\sqrt{\frac{1-ax}{1+ax}}\right)$$

[Out] $-a \operatorname{arctanh}\left(\left(\frac{-a*x+1}{a*x+1}\right)^{(1/2)}\right) - a / \left(1 + \left(\frac{-a*x+1}{a*x+1}\right)^{(1/2)}\right) + a / \left(1 + \left(\frac{-a*x+1}{a*x+1}\right)^{(1/2)}\right)$

Rubi [A] (verified)

Time = 0.27 (sec), antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6472, 78, 213}

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^2} dx = a \left(-\operatorname{arctanh}\left(\sqrt{\frac{1-ax}{ax+1}}\right) \right) + \frac{a}{\sqrt{\frac{1-ax}{ax+1}} + 1} - \frac{a}{\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^2}$$

[In] $\operatorname{Int}\left[1/\left(E^{\operatorname{ArcSech}[a*x]}*x^2\right), x\right]$

[Out] $-(a/(1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)]))^2 + a/(1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)]) - a \operatorname{ArcTanh}[\operatorname{Sqrt}[(1 - a*x)/(1 + a*x)]]$

Rule 78

$\operatorname{Int}[(a_. + (b_.)*(x_.))*((c_. + (d_.)*(x_.))^{(n_.)*((e_. + (f_.)*(x_.))^{(p_.)}, x_Symbol)] :> \operatorname{Int}[\operatorname{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, e, f, n\}, x\} \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& ((\operatorname{ILtQ}[n, 0] \&\& \operatorname{ILtQ}[p, 0]) \|\operator\| \operatorname{EqQ}[p, 1] \|\operator\| (\operatorname{IGtQ}[p, 0] \&\& (!\operatorname{IntegerQ}[n] \|\operator\| \operatorname{LeQ}[9*p +$

5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f]))))

Rule 213

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 6472

Int[E^(ArcSech[u_]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && IntegerQ[n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{1}{x^2 \left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax} \right)} dx \\
 &= (4a) \text{Subst} \left(\int \frac{x}{(-1+x)(1+x)^3} dx, x, \sqrt{\frac{1-ax}{1+ax}} \right) \\
 &= (4a) \text{Subst} \left(\int \left(\frac{1}{2(1+x)^3} - \frac{1}{4(1+x)^2} + \frac{1}{4(-1+x^2)} \right) dx, x, \sqrt{\frac{1-ax}{1+ax}} \right) \\
 &= -\frac{a}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{a}{1 + \sqrt{\frac{1-ax}{1+ax}}} + a \text{Subst} \left(\int \frac{1}{-1+x^2} dx, x, \sqrt{\frac{1-ax}{1+ax}} \right) \\
 &= -\frac{a}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{a}{1 + \sqrt{\frac{1-ax}{1+ax}}} - a \text{arctanh} \left(\sqrt{\frac{1-ax}{1+ax}} \right)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.28

$$\begin{aligned}
 \int \frac{e^{-\text{sech}^{-1}(ax)}}{x^2} dx &= \frac{1}{2} \left(-\frac{1}{ax^2} + \frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)}{ax^2} + a \log(x) \right. \\
 &\quad \left. - a \log \left(1 + \sqrt{\frac{1-ax}{1+ax}} + ax \sqrt{\frac{1-ax}{1+ax}} \right) \right)
 \end{aligned}$$

[In] Integrate[1/(E^ArcSech[a*x]*x^2),x]

[Out] $(-1/(a*x^2)) + (\text{Sqrt}[(1 - a*x)/(1 + a*x)]*(1 + a*x))/(a*x^2) + a*\text{Log}[x] - a*\text{Log}[1 + \text{Sqrt}[(1 - a*x)/(1 + a*x)]] + a*x*\text{Sqrt}[(1 - a*x)/(1 + a*x)])/2$

Maple [A] (verified)

Time = 0.89 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.33

method	result	size
default	$a \left(-\frac{1}{2a^2x^2} - \frac{\sqrt{-\frac{ax-1}{ax}} \sqrt{\frac{ax+1}{ax}} \left(a^2x^2 \operatorname{arctanh}\left(\frac{1}{\sqrt{-a^2x^2+1}}\right) - \sqrt{-a^2x^2+1} \right)}{2ax\sqrt{-a^2x^2+1}} \right)$	96

[In] int(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^2,x,method=_RETURNVERBOSE)

[Out] $a*(-1/2/a^2/x^2-1/2/a*(-(a*x-1)/a/x)^(1/2)/x*((a*x+1)/a/x)^(1/2)*(a^2*x^2*\operatorname{rctanh}(1/(-a^2*x^2+1)^(1/2))-(-a^2*x^2+1)^(1/2))/(-a^2*x^2+1)^(1/2))$

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.78

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^2} dx = \frac{a^2x^2 \log\left(ax\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} + 1\right) - a^2x^2 \log\left(ax\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} - 1\right) - 2ax\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} + 2}{4ax^2}$$

[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^2,x, algorithm="fricas")

[Out] $-1/4*(a^2*x^2*\log(a*x*\text{sqrt}((a*x + 1)/(a*x))*\text{sqrt}(-(a*x - 1)/(a*x)) + 1) - a^2*x^2*\log(a*x*\text{sqrt}((a*x + 1)/(a*x))*\text{sqrt}(-(a*x - 1)/(a*x)) - 1) - 2*a*x*\text{sqrt}((a*x + 1)/(a*x))*\text{sqrt}(-(a*x - 1)/(a*x)) + 2)/(a*x^2)$

Sympy [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^2} dx = a \int \frac{1}{ax^2 \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} + x} dx$$

[In] integrate(1/(1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))/x**2,x)

[Out] a*Integral(1/(a*x**2*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)) + x), x)

Maxima [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^2} dx = \int \frac{1}{x^2 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}} \right)} dx$$

[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^2,x, algorithm="maxima")

[Out] integrate(1/(x^2*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))), x)

Giac [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^2} dx = \int \frac{1}{x^2 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}} \right)} dx$$

[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^2,x, algorithm="giac")

[Out] integrate(1/(x^2*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))), x)

Mupad [B] (verification not implemented)

Time = 16.76 (sec) , antiderivative size = 323, normalized size of antiderivative = 4.49

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^2} dx = 2a \operatorname{atanh} \left(\frac{\sqrt{\frac{1}{ax} - 1 - i}}{\sqrt{\frac{1}{ax} + 1 - 1}} \right) - a \operatorname{acosh} \left(\frac{1}{ax} \right) - \frac{1}{2ax^2} \\ - \frac{a \left(\frac{14 \left(\sqrt{\frac{1}{ax} - 1 - i} \right)^3}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^3} + \frac{14 \left(\sqrt{\frac{1}{ax} - 1 - i} \right)^5}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^5} + \frac{2 \left(\sqrt{\frac{1}{ax} - 1 - i} \right)^7}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^7} + \frac{2 \left(\sqrt{\frac{1}{ax} - 1 - i} \right)}{\sqrt{\frac{1}{ax} + 1 - 1}} \right)}{1 + \frac{6 \left(\sqrt{\frac{1}{ax} - 1 - i} \right)^4}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^4} - \frac{4 \left(\sqrt{\frac{1}{ax} - 1 - i} \right)^6}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^6} + \frac{\left(\sqrt{\frac{1}{ax} - 1 - i} \right)^8}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^8} - \frac{4 \left(\sqrt{\frac{1}{ax} - 1 - i} \right)^2}{\left(\sqrt{\frac{1}{ax} + 1 - 1} \right)^2}}$$

[In] `int(1/(x^2*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))),x)`

[Out] $2*a*\operatorname{atanh}\left(\frac{(1/(a*x) - 1)^{1/2} - 1i}{(1/(a*x) + 1)^{1/2} - 1}\right) - a*\operatorname{acosh}\left(\frac{1}{a*x}\right) - \frac{1}{2*a*x^2} - \left(a*\frac{(14*(1/(a*x) - 1)^{1/2} - 1i)^3}{(1/(a*x) + 1)^{1/2} - 1} + \frac{14*(1/(a*x) - 1)^{1/2} - 1i}{(1/(a*x) + 1)^{1/2} - 1}\right)^5 + \frac{2*(1/(a*x) - 1)^{1/2} - 1i}{(1/(a*x) + 1)^{1/2} - 1} + \frac{2*(1/(a*x) - 1)^{1/2} - 1i}{(1/(a*x) + 1)^{1/2} - 1}\right) / \left(\frac{6*(1/(a*x) - 1)^{1/2} - 1i}{(1/(a*x) + 1)^{1/2} - 1}\right)^4 - \frac{4*(1/(a*x) - 1)^{1/2} - 1i}{(1/(a*x) + 1)^{1/2} - 1}\right)^2 / \left(\frac{1}{a*x} + 1\right)^{1/2} - 1)^2 - \frac{4*(1/(a*x) - 1)^{1/2} - 1i}{(1/(a*x) + 1)^{1/2} - 1}\right)^6 / \left(\frac{1}{a*x} + 1\right)^{1/2} - 1)^6 + \frac{(1/(a*x) - 1)^{1/2} - 1i}{(1/(a*x) + 1)^{1/2} - 1}\right)^8 / \left(\frac{1}{a*x} + 1\right)^{1/2} - 1)^8 + 1)$

3.83 $\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^3} dx$

Optimal result	519
Rubi [A] (verified)	519
Mathematica [A] (verified)	520
Maple [A] (verified)	520
Fricas [A] (verification not implemented)	521
Sympy [F]	521
Maxima [F]	521
Giac [F]	522
Mupad [B] (verification not implemented)	522

Optimal result

Integrand size = 12, antiderivative size = 116

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^3} dx = -\frac{a^2}{2\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} - \frac{2a^2}{3\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^3} + \frac{a^2}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^2} - \frac{a^2}{2\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}$$

[Out] $-1/2*a^2/(1-((-a*x+1)/(a*x+1))^(1/2))-2/3*a^2/(1+((-a*x+1)/(a*x+1))^(1/2))^3+3*a^2/(1+((-a*x+1)/(a*x+1))^(1/2))^2-1/2*a^2/(1+((-a*x+1)/(a*x+1))^(1/2))$

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6472, 1626}

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^3} dx = -\frac{a^2}{2\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)} - \frac{a^2}{2\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)} + \frac{a^2}{\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^2} - \frac{2a^2}{3\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^3}$$

[In] $\text{Int}[1/(E^{\text{ArcSech}[a*x]*x^3}), x]$

[Out] $-1/2*a^2/(1 - \text{Sqrt}[(1 - a*x)/(1 + a*x)]) - (2*a^2)/(3*(1 + \text{Sqrt}[(1 - a*x)/(1 + a*x]))^3) + a^2/(1 + \text{Sqrt}[(1 - a*x)/(1 + a*x)])^2 - a^2/(2*(1 + \text{Sqrt}[(1 - a*x)/(1 + a*x)]))$

Rule 1626

$\text{Int}[(P_x) * ((a_{\cdot}) + (b_{\cdot})*(x_{\cdot}))^{(m_{\cdot})} * ((c_{\cdot}) + (d_{\cdot})*(x_{\cdot}))^{(n_{\cdot})} * ((e_{\cdot}) + (f_{\cdot})*(x_{\cdot}))^{(p_{\cdot})}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[P_x*(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x] \&\& \text{PolyQ}[P_x,$

x] && IntegersQ[m, n]

Rule 6472

Int[E^(ArcSech[u_]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && IntegerQ[n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{1}{x^3 \left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax} \right)} dx \\
 &= - \left((4a^2) \text{Subst} \left(\int \frac{x(1+x^2)}{(-1+x)^2(1+x)^4} dx, x, \sqrt{\frac{1-ax}{1+ax}} \right) \right) \\
 &= - \left((4a^2) \text{Subst} \left(\int \left(\frac{1}{8(-1+x)^2} - \frac{1}{2(1+x)^4} + \frac{1}{2(1+x)^3} - \frac{1}{8(1+x)^2} \right) dx, x, \sqrt{\frac{1-ax}{1+ax}} \right) \right) \\
 &= - \frac{a^2}{2 \left(1 - \sqrt{\frac{1-ax}{1+ax}} \right)} - \frac{2a^2}{3 \left(1 + \sqrt{\frac{1-ax}{1+ax}} \right)^3} + \frac{a^2}{\left(1 + \sqrt{\frac{1-ax}{1+ax}} \right)^2} - \frac{a^2}{2 \left(1 + \sqrt{\frac{1-ax}{1+ax}} \right)}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.37

$$\int \frac{e^{-\text{sech}^{-1}(ax)}}{x^3} dx = - \frac{1 + (-1 + ax) \sqrt{\frac{1-ax}{1+ax}} (1 + ax)^2}{3ax^3}$$

[In] Integrate[1/(E^ArcSech[a*x]*x^3), x]

[Out] -1/3*(1 + (-1 + a*x)*Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2)/(a*x^3)

Maple [A] (verified)

Time = 0.91 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.50

method	result	size
default	$a \left(-\frac{1}{3a^2x^3} - \frac{\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} (a^2x^2-1)}{3ax^2} \right)$	58


```
[In] int(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^3,x,method=_RETURNVERBOSE)
[Out] a*(-1/3/a^2/x^3-1/3/a*((a*x+1)/a/x)^(1/2)/x^2*(-(a*x-1)/a/x)^(1/2)*(a^2*x^2-1))
```

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.45

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^3} dx = -\frac{(a^3x^3 - ax)\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} + 1}{3ax^3}$$

```
[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^3,x, algorithm="fricas")
```

```
[Out] -1/3*((a^3*x^3 - a*x)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + 1)/(a*x^3)
```

Sympy [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^3} dx = a \int \frac{1}{ax^3\sqrt{-1 + \frac{1}{ax}}\sqrt{1 + \frac{1}{ax}} + x^2} dx$$

```
[In] integrate(1/(1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))/x**3,x)
```

```
[Out] a*Integral(1/(a*x**3*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)) + x**2), x)
```

Maxima [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^3} dx = \int \frac{1}{x^3\left(\sqrt{\frac{1}{ax}} + 1\sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}}\right)} dx$$

```
[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^3,x, algorithm="maxima")
```

```
[Out] integrate(1/(x^3*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))), x)
```

Giac [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^3} dx = \int \frac{1}{x^3 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}} \right)} dx$$

[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^3,x, algorithm="giac")

[Out] integrate(1/(x^3*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))), x)

Mupad [B] (verification not implemented)

Time = 5.73 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.50

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^3} dx = \frac{\sqrt{\frac{1}{ax} - 1} \left(\frac{x}{3} - \frac{ax^2}{3} + \frac{1}{3a} - \frac{a^2 x^3}{3} \right)}{x^3 \sqrt{\frac{1}{ax} + 1}} - \frac{1}{3ax^3}$$

[In] int(1/(x^3*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))),x)

[Out] ((1/(a*x) - 1)^(1/2)*(x/3 - (a*x^2)/3 + 1/(3*a) - (a^2*x^3)/3))/(x^3*(1/(a*x) + 1)^(1/2)) - 1/(3*a*x^3)

3.84 $\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^4} dx$

Optimal result	523
Rubi [A] (verified)	523
Mathematica [A] (verified)	525
Maple [A] (verified)	525
Fricas [A] (verification not implemented)	526
Sympy [F]	526
Maxima [F]	526
Giac [F]	527
Mupad [B] (verification not implemented)	527

Optimal result

Integrand size = 12, antiderivative size = 200

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^4} dx = -\frac{a^3}{4 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{a^3}{4 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} - \frac{a^3}{2 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^4}$$

$$+ \frac{a^3}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^3} - \frac{a^3}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{a^3}{2 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}$$

$$- \frac{1}{4} a^3 \operatorname{arctanh}\left(\sqrt{\frac{1-ax}{1+ax}}\right)$$

[Out] $-1/4*a^3*\operatorname{arctanh}\left(\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)-1/4*a^3/\left(1-\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)^2+1/4*a^3/\left(1-\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)-1/2*a^3/\left(1+\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)^4+a^3/\left(1+\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)^3-a^3/\left(1+\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)^2+1/2*a^3/\left(1+\left(\frac{-a*x+1}{a*x+1}\right)^{1/2}\right)$

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used

= {6472, 1626, 213}

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^4} dx = -\frac{1}{4}a^3 \operatorname{arctanh}\left(\sqrt{\frac{1-ax}{ax+1}}\right) + \frac{a^3}{4\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)} + \frac{a^3}{2\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)} - \frac{a^3}{4\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^2} - \frac{a^3}{\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^2} + \frac{a^3}{\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^3} - \frac{a^3}{2\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^4}$$

[In] Int[1/(E^ArcSech[a*x]*x^4),x]

[Out] -1/4*a^3/(1 - Sqrt[(1 - a*x)/(1 + a*x)])^2 + a^3/(4*(1 - Sqrt[(1 - a*x)/(1 + a*x)])) - a^3/(2*(1 + Sqrt[(1 - a*x)/(1 + a*x)])^4) + a^3/(1 + Sqrt[(1 - a*x)/(1 + a*x)])^3 - a^3/(1 + Sqrt[(1 - a*x)/(1 + a*x)])^2 + a^3/(2*(1 + Sqrt[(1 - a*x)/(1 + a*x)])) - (a^3*ArcTanh[Sqrt[(1 - a*x)/(1 + a*x)]])/4

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 1626

Int[(Px_)*((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && IntegersQ[m, n]

Rule 6472

Int[E^(ArcSech[u_]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && IntegerQ[n]

Rubi steps

$$\begin{aligned} \text{integral} &= \int \frac{1}{x^4 \left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \frac{\sqrt{\frac{1-ax}{1+ax}}}{ax} \right)} dx \\ &= (4a) \operatorname{Subst} \left(\int \frac{x(a + ax^2)^2}{(-1 + x)^3(1 + x)^5} dx, x, \sqrt{\frac{1 - ax}{1 + ax}} \right) \end{aligned}$$

$$\begin{aligned}
&= (4a)\text{Subst}\left(\int\left(\frac{a^2}{8(-1+x)^3}+\frac{a^2}{16(-1+x)^2}+\frac{a^2}{2(1+x)^5}-\frac{3a^2}{4(1+x)^4}+\frac{a^2}{2(1+x)^3}\right.\right. \\
&\quad\left.\left.-\frac{a^2}{8(1+x)^2}+\frac{a^2}{16(-1+x^2)}\right)dx,x,\sqrt{\frac{1-ax}{1+ax}}\right) \\
&= -\frac{a^3}{4\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)^2}+\frac{a^3}{4\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)}-\frac{a^3}{2\left(1+\sqrt{\frac{1-ax}{1+ax}}\right)^4}+\frac{a^3}{\left(1+\sqrt{\frac{1-ax}{1+ax}}\right)^3} \\
&\quad -\frac{a^3}{\left(1+\sqrt{\frac{1-ax}{1+ax}}\right)^2}+\frac{a^3}{2\left(1+\sqrt{\frac{1-ax}{1+ax}}\right)}+\frac{1}{4}a^3\text{Subst}\left(\int\frac{1}{-1+x^2}dx,x,\sqrt{\frac{1-ax}{1+ax}}\right) \\
&= -\frac{a^3}{4\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)^2}+\frac{a^3}{4\left(1-\sqrt{\frac{1-ax}{1+ax}}\right)}-\frac{a^3}{2\left(1+\sqrt{\frac{1-ax}{1+ax}}\right)^4}+\frac{a^3}{\left(1+\sqrt{\frac{1-ax}{1+ax}}\right)^3} \\
&\quad -\frac{a^3}{\left(1+\sqrt{\frac{1-ax}{1+ax}}\right)^2}+\frac{a^3}{2\left(1+\sqrt{\frac{1-ax}{1+ax}}\right)}-\frac{1}{4}a^3\text{arctanh}\left(\sqrt{\frac{1-ax}{1+ax}}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.55

$$\int\frac{e^{-\text{sech}^{-1}(ax)}}{x^4}dx=
\frac{2+\sqrt{\frac{1-ax}{1+ax}}(-2-2ax+a^2x^2+a^3x^3)-a^4x^4\log(x)+a^4x^4\log\left(1+\sqrt{\frac{1-ax}{1+ax}}+ax\sqrt{\frac{1-ax}{1+ax}}\right)}{8ax^4}$$

[In] Integrate[1/(E^ArcSech[a*x]*x^4),x]

[Out] -1/8*(2+Sqrt[(1-a*x)/(1+a*x)]*(-2-2*a*x+a^2*x^2+a^3*x^3)-a^4*x^4*Log[x]+a^4*x^4*Log[1+Sqrt[(1-a*x)/(1+a*x)]+a*x*Sqrt[(1-a*x)/(1+a*x)]])/(a*x^4)

Maple [A] (verified)

Time = 0.93 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.58

method	result	size
default	$a\left(-\frac{1}{4a^2x^4}-\frac{\sqrt{-\frac{ax-1}{ax}}\sqrt{\frac{ax+1}{ax}}\left(\text{arctanh}\left(\frac{1}{\sqrt{-a^2x^2+1}}\right)a^4x^4+a^2x^2\sqrt{-a^2x^2+1}-2\sqrt{-a^2x^2+1}\right)}{8ax^3\sqrt{-a^2x^2+1}}\right)$	115

```
[In] int(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^4,x,method=_RETURNVERBOSE)
[Out] a*(-1/4/a^2/x^4-1/8/a*(-(a*x-1)/a/x)^(1/2)/x^3*((a*x+1)/a/x)^(1/2)*(arctanh
(1/(-a^2*x^2+1)^(1/2))*a^4*x^4+a^2*x^2*(-a^2*x^2+1)^(1/2)-2*(-a^2*x^2+1)^(1
/2))/(-a^2*x^2+1)^(1/2))
```

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.69

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^4} dx = \frac{a^4 x^4 \log\left(ax \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} + 1\right) - a^4 x^4 \log\left(ax \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} - 1\right) + 2(a^3 x^3 - 2ax) \sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}}}{16 ax^4}$$

```
[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^4,x, algorithm="fricas")
```

```
[Out] -1/16*(a^4*x^4*log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + 1) -
a^4*x^4*log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 1) + 2*(a^3*
x^3 - 2*a*x)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + 4)/(a*x^4)
```

Sympy [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^4} dx = a \int \frac{1}{ax^4 \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} + x^3} dx$$

```
[In] integrate(1/(1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))/x**4,x)
```

```
[Out] a*Integral(1/(a*x**4*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)) + x**3), x)
```

Maxima [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^4} dx = \int \frac{1}{x^4 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}} \right)} dx$$

```
[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^4,x, algorithm="maxima")
```

```
[Out] integrate(1/(x^4*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))), x)
```

Giac [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^4} dx = \int \frac{1}{x^4 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1} + \frac{1}{ax} \right)} dx$$

[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^4,x, algorithm="giac")

[Out] integrate(1/(x^4*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))), x)

Mupad [B] (verification not implemented)

Time = 50.94 (sec) , antiderivative size = 1511, normalized size of antiderivative = 7.56

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^4} dx = \text{Too large to display}$$

[In] int(1/(x^4*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))),x)

[Out] ((a^3*((1/(a*x) - 1)^(1/2) - 1i)^4*192i)/((1/(a*x) + 1)^(1/2) - 1)^4 + (a^3 * ((1/(a*x) - 1)^(1/2) - 1i)^6*128i)/((1/(a*x) + 1)^(1/2) - 1)^6 + (a^3*((1/(a*x) - 1)^(1/2) - 1i)^8*192i)/((1/(a*x) + 1)^(1/2) - 1)^8)/(3*((15*((1/(a*x) - 1)^(1/2) - 1i)^4)/((1/(a*x) + 1)^(1/2) - 1)^4 - (6*((1/(a*x) - 1)^(1/2) - 1i)^2)/((1/(a*x) + 1)^(1/2) - 1)^2 - (20*((1/(a*x) - 1)^(1/2) - 1i)^6)/((1/(a*x) + 1)^(1/2) - 1)^6 + (15*((1/(a*x) - 1)^(1/2) - 1i)^8)/((1/(a*x) + 1)^(1/2) - 1)^8 - (6*((1/(a*x) - 1)^(1/2) - 1i)^10)/((1/(a*x) + 1)^(1/2) - 1)^10 + ((1/(a*x) - 1)^(1/2) - 1i)^12/((1/(a*x) + 1)^(1/2) - 1)^12 + 1)) - ((a^3*((1/(a*x) - 1)^(1/2) - 1i)^4*64i)/((1/(a*x) + 1)^(1/2) - 1)^4 + (a^3 * ((1/(a*x) - 1)^(1/2) - 1i)^6*128i)/(3*((1/(a*x) + 1)^(1/2) - 1)^6) + (a^3 * ((1/(a*x) - 1)^(1/2) - 1i)^8*64i)/((1/(a*x) + 1)^(1/2) - 1)^8)/(15*((1/(a*x) - 1)^(1/2) - 1i)^4)/((1/(a*x) + 1)^(1/2) - 1)^4 - (6*((1/(a*x) - 1)^(1/2) - 1i)^2)/((1/(a*x) + 1)^(1/2) - 1)^2 - (20*((1/(a*x) - 1)^(1/2) - 1i)^6)/((1/(a*x) + 1)^(1/2) - 1)^6 + (15*((1/(a*x) - 1)^(1/2) - 1i)^8)/((1/(a*x) + 1)^(1/2) - 1)^8 - (6*((1/(a*x) - 1)^(1/2) - 1i)^10)/((1/(a*x) + 1)^(1/2) - 1)^10 + ((1/(a*x) - 1)^(1/2) - 1i)^12/((1/(a*x) + 1)^(1/2) - 1)^12 + 1) - (a^3*atanh(((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1))))/2 + ((14*a^3*((1/(a*x) - 1)^(1/2) - 1i)^3)/((1/(a*x) + 1)^(1/2) - 1)^3 + (14*a^3*((1/(a*x) - 1)^(1/2) - 1i)^5)/((1/(a*x) + 1)^(1/2) - 1)^5 + (2*a^3*((1/(a*x) - 1)^(1/2) - 1i)^7)/((1/(a*x) + 1)^(1/2) - 1)^7 + (2*a^3*((1/(a*x) - 1)^(1/2) - 1i))/((1/(a*x) + 1)^(1/2) - 1))/((6*((1/(a*x) - 1)^(1/2) - 1i)^4)/((1/(a*x) + 1)^(1/2) - 1)^4 - (4*((1/(a*x) - 1)^(1/2) - 1i)^2)/((1/(a*x) + 1)^(1/2) - 1)^2 - (4*((1/(a*x) - 1)^(1/2) - 1i)^6)/((1/(a*x) + 1)^(1/2) - 1)^6 + ((1/(a*x) - 1)^(1/2) - 1i)^8/((1/(a*x) + 1)^(1/2) - 1)^8 + 1) + ((23*a^3*(

$$\begin{aligned}
& \frac{(1/(a*x) - 1)^{(1/2) - 1i}^3}{2*((1/(a*x) + 1)^{(1/2) - 1)^3} + (333*a^3*((1/(a*x) - 1)^{(1/2) - 1i}^5)/2*((1/(a*x) + 1)^{(1/2) - 1)^5) + (671*a^3*((1/(a*x) - 1)^{(1/2) - 1i}^7)/2*((1/(a*x) + 1)^{(1/2) - 1)^7) + (671*a^3*((1/(a*x) - 1)^{(1/2) - 1i}^9)/2*((1/(a*x) + 1)^{(1/2) - 1)^9) + (333*a^3*((1/(a*x) - 1)^{(1/2) - 1i}^{11})/2*((1/(a*x) + 1)^{(1/2) - 1)^{11}) + (23*a^3*((1/(a*x) - 1)^{(1/2) - 1i}^{13})/2*((1/(a*x) + 1)^{(1/2) - 1)^{13}) - (3*a^3*((1/(a*x) - 1)^{(1/2) - 1i}^{15})/2*((1/(a*x) + 1)^{(1/2) - 1)^{15}) - (3*a^3*((1/(a*x) - 1)^{(1/2) - 1i}))/2*((1/(a*x) + 1)^{(1/2) - 1}))/((28*((1/(a*x) - 1)^{(1/2) - 1i}^4)/((1/(a*x) + 1)^{(1/2) - 1)^4 - (8*((1/(a*x) - 1)^{(1/2) - 1i}^2)/((1/(a*x) + 1)^{(1/2) - 1)^2 - (56*((1/(a*x) - 1)^{(1/2) - 1i}^6)/((1/(a*x) + 1)^{(1/2) - 1)^6 + (70*((1/(a*x) - 1)^{(1/2) - 1i}^8)/((1/(a*x) + 1)^{(1/2) - 1)^8 - (56*((1/(a*x) - 1)^{(1/2) - 1i}^{10})/((1/(a*x) + 1)^{(1/2) - 1)^{10} + (28*((1/(a*x) - 1)^{(1/2) - 1i}^{12})/((1/(a*x) + 1)^{(1/2) - 1)^{12} - (8*((1/(a*x) - 1)^{(1/2) - 1i}^{14})/((1/(a*x) + 1)^{(1/2) - 1)^{14} + ((1/(a*x) - 1)^{(1/2) - 1i}^{16})/((1/(a*x) + 1)^{(1/2) - 1)^{16} + 1) - 1/(4*a*x^4)
\end{aligned}$$

3.85 $\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^5} dx$

Optimal result	529
Rubi [A] (verified)	529
Mathematica [A] (verified)	531
Maple [A] (verified)	531
Fricas [A] (verification not implemented)	531
Sympy [F]	532
Maxima [F]	532
Giac [F]	532
Mupad [B] (verification not implemented)	532

Optimal result

Integrand size = 12, antiderivative size = 233

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^5} dx = -\frac{a^4}{6 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3} + \frac{a^4}{4 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} - \frac{3a^4}{8 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} - \frac{2a^4}{5 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^5} + \frac{a^4}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^4} - \frac{4a^4}{3 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^3} + \frac{a^4}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^2} - \frac{3a^4}{8 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}$$

```
[Out] -1/6*a^4/(1-((-a*x+1)/(a*x+1))^(1/2))^3+1/4*a^4/(1-((-a*x+1)/(a*x+1))^(1/2))^2-3/8*a^4/(1-((-a*x+1)/(a*x+1))^(1/2))-2/5*a^4/(1+((-a*x+1)/(a*x+1))^(1/2))^5+a^4/(1+((-a*x+1)/(a*x+1))^(1/2))^4-4/3*a^4/(1+((-a*x+1)/(a*x+1))^(1/2))^3+a^4/(1+((-a*x+1)/(a*x+1))^(1/2))^2-3/8*a^4/(1+((-a*x+1)/(a*x+1))^(1/2))
```

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used

= {6472, 1626}

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^5} dx = -\frac{3a^4}{8 \left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)} - \frac{3a^4}{8 \left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)} + \frac{a^4}{4 \left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^2}$$

$$+ \frac{a^4}{\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^2} - \frac{a^4}{6 \left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^3} - \frac{a^4}{3 \left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^3}$$

$$+ \frac{a^4}{\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^4} - \frac{2a^4}{5 \left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^5}$$

[In] Int[1/(E^ArcSech[a*x]*x^5),x]

[Out] -1/6*a^4/(1 - Sqrt[(1 - a*x)/(1 + a*x)])^3 + a^4/(4*(1 - Sqrt[(1 - a*x)/(1 + a*x)])^2) - (3*a^4)/(8*(1 - Sqrt[(1 - a*x)/(1 + a*x)])) - (2*a^4)/(5*(1 + Sqrt[(1 - a*x)/(1 + a*x)])^5) + a^4/(1 + Sqrt[(1 - a*x)/(1 + a*x)])^4 - (4*a^4)/(3*(1 + Sqrt[(1 - a*x)/(1 + a*x)])^3) + a^4/(1 + Sqrt[(1 - a*x)/(1 + a*x)])^2 - (3*a^4)/(8*(1 + Sqrt[(1 - a*x)/(1 + a*x)]))

Rule 1626

Int[(Px_)*((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && IntegersQ[m, n]

Rule 6472

Int[E^(ArcSech[u_]*(n_.)*(x_))^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && IntegerQ[n]

Rubi steps

$$\text{integral} = \int \frac{1}{x^5 \left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \sqrt{\frac{1-ax}{1+ax}} \frac{1}{ax} \right)} dx$$

$$= - \left((4a) \operatorname{Subst} \left(\int \frac{x(a + ax^2)^3}{(-1 + x)^4(1 + x)^6} dx, x, \sqrt{\frac{1 - ax}{1 + ax}} \right) \right)$$

$$= - \left((4a) \operatorname{Subst} \left(\int \left(\frac{a^3}{8(-1 + x)^4} + \frac{a^3}{8(-1 + x)^3} + \frac{3a^3}{32(-1 + x)^2} - \frac{a^3}{2(1 + x)^6} + \frac{a^3}{(1 + x)^5} - \frac{a^3}{(1 + x)^4} + \frac{a^3}{2(1 + x)^3} \right) dx, x, \sqrt{\frac{1 - ax}{1 + ax}} \right) \right)$$

$$= -\frac{a^4}{6\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3} + \frac{a^4}{4\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} - \frac{3a^4}{8\left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} - \frac{2a^4}{5\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^5}$$

$$+ \frac{a^4}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^4} - \frac{4a^4}{3\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^3} + \frac{a^4}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^2} - \frac{3a^4}{8\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.26

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^5} dx = -\frac{3 + \sqrt{\frac{1-ax}{1+ax}}(1+ax)^2(-3 + 3ax - 2a^2x^2 + 2a^3x^3)}{15ax^5}$$

[In] Integrate[1/(E^ArcSech[a*x]*x^5),x]

[Out] -1/15*(3 + Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2*(-3 + 3*a*x - 2*a^2*x^2 + 2*a^3*x^3))/(a*x^5)

Maple [A] (verified)

Time = 0.94 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.29

method	result	size
default	$a\left(-\frac{1}{5a^2x^5} - \frac{\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}(a^2x^2-1)(2a^2x^2+3)}{15ax^4}\right)$	68

[In] int(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^5,x,method=_RETURNVERBOSE)

[Out] a*(-1/5/a^2/x^5-1/15/a*((a*x+1)/a/x)^(1/2)/x^4*(-(a*x-1)/a/x)^(1/2)*(a^2*x^2-1)*(2*a^2*x^2+3))

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.26

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^5} dx = -\frac{(2a^5x^5 + a^3x^3 - 3ax)\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} + 3}{15ax^5}$$

[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^5,x, algorithm="fricas")

[Out] -1/15*((2*a^5*x^5 + a^3*x^3 - 3*a*x)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + 3)/(a*x^5)

Sympy [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^5} dx = a \int \frac{1}{ax^5 \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax} + x^4}} dx$$

[In] integrate(1/(1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))/x**5,x)

[Out] a*Integral(1/(a*x**5*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)) + x**4), x)

Maxima [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^5} dx = \int \frac{1}{x^5 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}} \right)} dx$$

[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^5,x, algorithm="maxima")

[Out] integrate(1/(x^5*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))), x)

Giac [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^5} dx = \int \frac{1}{x^5 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}} \right)} dx$$

[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^5,x, algorithm="giac")

[Out] integrate(1/(x^5*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))), x)

Mupad [B] (verification not implemented)

Time = 5.17 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.32

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^5} dx = -\frac{1}{5ax^5} - \frac{\sqrt{\frac{1}{ax} - 1} \left(\frac{ax^2}{15} - \frac{x}{5} - \frac{1}{5a} + \frac{a^2x^3}{15} + \frac{2a^3x^4}{15} + \frac{2a^4x^5}{15} \right)}{x^5 \sqrt{\frac{1}{ax} + 1}}$$

[In] int(1/(x^5*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))),x)

[Out] - 1/(5*a*x^5) - ((1/(a*x) - 1)^(1/2)*((a*x^2)/15 - x/5 - 1/(5*a) + (a^2*x^3)/15 + (2*a^3*x^4)/15 + (2*a^4*x^5)/15))/(x^5*(1/(a*x) + 1)^(1/2))

3.86 $\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^6} dx$

Optimal result	533
Rubi [A] (verified)	534
Mathematica [A] (verified)	536
Maple [A] (verified)	536
Fricas [A] (verification not implemented)	536
Sympy [F]	537
Maxima [F]	537
Giac [F]	537
Mupad [B] (verification not implemented)	538

Optimal result

Integrand size = 12, antiderivative size = 320

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^6} dx = -\frac{a^5}{8 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^4} + \frac{a^5}{4 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3} - \frac{3a^5}{8 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2}$$

$$+ \frac{a^5}{4 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} - \frac{a^5}{3 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^6} + \frac{a^5}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^5}$$

$$- \frac{13a^5}{8 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^4} + \frac{19a^5}{12 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^3} - \frac{a^5}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^2}$$

$$+ \frac{3a^5}{8 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)} - \frac{1}{8} a^5 \operatorname{arctanh}\left(\sqrt{\frac{1-ax}{1+ax}}\right)$$

```
[Out] -1/8*a^5*arctanh(((a*x+1)/(a*x+1))^(1/2))-1/8*a^5/(1-((a*x+1)/(a*x+1))^(1/2))^4+1/4*a^5/(1-((a*x+1)/(a*x+1))^(1/2))^3-3/8*a^5/(1-((a*x+1)/(a*x+1))^(1/2))^2+1/4*a^5/(1-((a*x+1)/(a*x+1))^(1/2))-1/3*a^5/(1+((a*x+1)/(a*x+1))^(1/2))^6+a^5/(1+((a*x+1)/(a*x+1))^(1/2))^5-13/8*a^5/(1+((a*x+1)/(a*x+1))^(1/2))^4+19/12*a^5/(1+((a*x+1)/(a*x+1))^(1/2))^3-a^5/(1+((a*x+1)/(a*x+1))^(1/2))^2+3/8*a^5/(1+((a*x+1)/(a*x+1))^(1/2))
```

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6472, 1626, 213}

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^6} dx = -\frac{1}{8}a^5 \operatorname{arctanh}\left(\sqrt{\frac{1-ax}{ax+1}}\right) + \frac{a^5}{4\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)} + \frac{3a^5}{8\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)} - \frac{3a^5}{8\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^2} - \frac{a^5}{\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^2} + \frac{a^5}{4\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^3} + \frac{19a^5}{12\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^3} - \frac{a^5}{8\left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^4} - \frac{13a^5}{8\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^4} + \frac{a^5}{\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^5} - \frac{a^5}{3\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^6}$$

[In] Int[1/(E^ArcSech[a*x]*x^6),x]

[Out] $-1/8*a^5/(1 - \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^4 + a^5/(4*(1 - \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^3) - (3*a^5)/(8*(1 - \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^2) + a^5/(4*(1 - \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])) - a^5/(3*(1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^6) + a^5/(1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^5 - (13*a^5)/(8*(1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^4) + (19*a^5)/(12*(1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^3) - a^5/(1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^2 + (3*a^5)/(8*(1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])) - (a^5*\operatorname{ArcTanh}[\operatorname{Sqrt}[(1 - a*x)/(1 + a*x)]])/8$

Rule 213

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 1626

Int[(Px_)*((a_) + (b_)*(x_)^2)^(m_)*((c_) + (d_)*(x_)^2)^(n_)*((e_) + (f_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && IntegersQ[m, n]

Rule 6472

Int[E^(ArcSech[u_]*(n_))*(x_)^(m_), x_Symbol] := Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && Integer

Q[n]

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \frac{1}{x^6 \left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \sqrt{\frac{1-ax}{1+ax}} \frac{1}{ax} \right)} dx \\
&= (4a) \text{Subst} \left(\int \frac{x(a+ax^2)^4}{(-1+x)^5(1+x)^7} dx, x, \sqrt{\frac{1-ax}{1+ax}} \right) \\
&= (4a) \text{Subst} \left(\int \left(\frac{a^4}{8(-1+x)^5} + \frac{3a^4}{16(-1+x)^4} + \frac{3a^4}{16(-1+x)^3} + \frac{a^4}{16(-1+x)^2} \right. \right. \\
&\quad \left. \left. + \frac{a^4}{2(1+x)^7} - \frac{5a^4}{4(1+x)^6} + \frac{13a^4}{8(1+x)^5} - \frac{19a^4}{16(1+x)^4} + \frac{a^4}{2(1+x)^3} - \frac{3a^4}{32(1+x)^2} \right. \right. \\
&\quad \left. \left. + \frac{a^4}{32(-1+x^2)} \right) dx, x, \sqrt{\frac{1-ax}{1+ax}} \right) \\
&= -\frac{a^5}{8 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^4} + \frac{a^5}{4 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3} - \frac{3a^5}{8 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{a^5}{4 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} \\
&\quad - \frac{a^5}{3 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^6} + \frac{a^5}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^5} - \frac{13a^5}{8 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^4} + \frac{19a^5}{12 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^3} \\
&\quad - \frac{a^5}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{3a^5}{8 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)} + \frac{1}{8} a^5 \text{Subst} \left(\int \frac{1}{-1+x^2} dx, x, \sqrt{\frac{1-ax}{1+ax}} \right) \\
&= -\frac{a^5}{8 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^4} + \frac{a^5}{4 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3} - \frac{3a^5}{8 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{a^5}{4 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} \\
&\quad - \frac{a^5}{3 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^6} + \frac{a^5}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^5} - \frac{13a^5}{8 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^4} + \frac{19a^5}{12 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^3} \\
&\quad - \frac{a^5}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^2} + \frac{3a^5}{8 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)} - \frac{1}{8} a^5 \text{arctanh} \left(\sqrt{\frac{1-ax}{1+ax}} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.40

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^6} dx = \frac{8 + \sqrt{\frac{1-ax}{1+ax}}(-8 - 8ax + 2a^2x^2 + 2a^3x^3 + 3a^4x^4 + 3a^5x^5) - 3a^6x^6 \log(x) + 3a^6x^6 \log\left(1 + \sqrt{\frac{1-ax}{1+ax}} + ax\sqrt{\frac{1-ax}{1+ax}}\right)}{48ax^6}$$

[In] Integrate[1/(E^ArcSech[a*x]*x^6),x]

[Out] $-1/48*(8 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)]*(-8 - 8*a*x + 2*a^2*x^2 + 2*a^3*x^3 + 3*a^4*x^4 + 3*a^5*x^5) - 3*a^6*x^6*\operatorname{Log}[x] + 3*a^6*x^6*\operatorname{Log}[1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)] + a*x*\operatorname{Sqrt}[(1 - a*x)/(1 + a*x)]])/(a*x^6)$

Maple [A] (verified)

Time = 1.01 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.43

method	result	size
default	$a \left(-\frac{1}{6a^2x^6} - \frac{\sqrt{-\frac{ax-1}{ax}} \sqrt{\frac{ax+1}{ax}} \left(3 \operatorname{arctanh}\left(\frac{1}{\sqrt{-a^2x^2+1}}\right) a^6x^6 + 3\sqrt{-a^2x^2+1}a^4x^4 + 2a^2x^2\sqrt{-a^2x^2+1} - 8\sqrt{-a^2x^2+1} \right)}{48ax^5\sqrt{-a^2x^2+1}} \right)$	137

[In] int(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^6,x,method=_RETURNVERBOSE)

[Out] $a*(-1/6/a^2/x^6 - 1/48/a*(-(a*x-1)/a/x)^(1/2)/x^5*((a*x+1)/a/x)^(1/2)*(3*\operatorname{arctanh}(1/(-a^2*x^2+1)^(1/2))*a^6*x^6 + 3*(-a^2*x^2+1)^(1/2)*a^4*x^4 + 2*a^2*x^2*(-a^2*x^2+1)^(1/2) - 8*(-a^2*x^2+1)^(1/2))/(-a^2*x^2+1)^(1/2))$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.46

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^6} dx = \frac{3a^6x^6 \log\left(ax\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} + 1\right) - 3a^6x^6 \log\left(ax\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} - 1\right) + 2(3a^5x^5 + 2a^3x^3 - 8ax)\sqrt{-\frac{ax-1}{ax}}}{96ax^6}$$

[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^6,x, algorithm="fricas")


```
[Out] -1/96*(3*a^6*x^6*log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + 1)
- 3*a^6*x^6*log(a*x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) - 1) + 2*(
3*a^5*x^5 + 2*a^3*x^3 - 8*a*x)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x))
+ 16)/(a*x^6)
```

Sympy [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^6} dx = a \int \frac{1}{ax^6 \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} + x^5} dx$$

```
[In] integrate(1/(1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))/x**6,x)
```

```
[Out] a*Integral(1/(a*x**6*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)) + x**5), x)
```

Maxima [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^6} dx = \int \frac{1}{x^6 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}} \right)} dx$$

```
[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^6,x, algorithm="maxim
a")
```

```
[Out] integrate(1/(x^6*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))), x)
```

Giac [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^6} dx = \int \frac{1}{x^6 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}} \right)} dx$$

```
[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^6,x, algorithm="giac"
)
```

```
[Out] integrate(1/(x^6*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))), x)
```

Mupad [B] (verification not implemented)

Time = 69.66 (sec) , antiderivative size = 2479, normalized size of antiderivative = 7.75

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^6} dx = \text{Too large to display}$$

[In] int(1/(x^6*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))),x)

[Out] ((a^5*((1/(a*x) - 1)^(1/2) - 1i)^6*10240i)/((1/(a*x) + 1)^(1/2) - 1)^6 + (a^5*((1/(a*x) - 1)^(1/2) - 1i)^8*20480i)/((1/(a*x) + 1)^(1/2) - 1)^8 + (a^5*((1/(a*x) - 1)^(1/2) - 1i)^10*36864i)/((1/(a*x) + 1)^(1/2) - 1)^10 + (a^5*((1/(a*x) - 1)^(1/2) - 1i)^12*20480i)/((1/(a*x) + 1)^(1/2) - 1)^12 + (a^5*((1/(a*x) - 1)^(1/2) - 1i)^14*10240i)/((1/(a*x) + 1)^(1/2) - 1)^14)/(15*((45*((1/(a*x) - 1)^(1/2) - 1i)^4)/((1/(a*x) + 1)^(1/2) - 1)^4 - (10*((1/(a*x) - 1)^(1/2) - 1i)^2)/((1/(a*x) + 1)^(1/2) - 1)^2 - (120*((1/(a*x) - 1)^(1/2) - 1i)^6)/((1/(a*x) + 1)^(1/2) - 1)^6 + (210*((1/(a*x) - 1)^(1/2) - 1i)^8)/((1/(a*x) + 1)^(1/2) - 1)^8 - (252*((1/(a*x) - 1)^(1/2) - 1i)^10)/((1/(a*x) + 1)^(1/2) - 1)^10 + (210*((1/(a*x) - 1)^(1/2) - 1i)^12)/((1/(a*x) + 1)^(1/2) - 1)^12 - (120*((1/(a*x) - 1)^(1/2) - 1i)^14)/((1/(a*x) + 1)^(1/2) - 1)^14 + (45*((1/(a*x) - 1)^(1/2) - 1i)^16)/((1/(a*x) + 1)^(1/2) - 1)^16 - (10*((1/(a*x) - 1)^(1/2) - 1i)^18)/((1/(a*x) + 1)^(1/2) - 1)^18 + ((1/(a*x) - 1)^(1/2) - 1i)^20)/((1/(a*x) + 1)^(1/2) - 1)^20 + 1)) - (a^5*atanh((1/(a*x) - 1)^(1/2) - 1i)/((1/(a*x) + 1)^(1/2) - 1)))/4 - ((a^5*((1/(a*x) - 1)^(1/2) - 1i)^6*2048i)/(3*((1/(a*x) + 1)^(1/2) - 1)^6) + (a^5*((1/(a*x) - 1)^(1/2) - 1i)^8*4096i)/(3*((1/(a*x) + 1)^(1/2) - 1)^8) + (a^5*((1/(a*x) - 1)^(1/2) - 1i)^10*12288i)/(5*((1/(a*x) + 1)^(1/2) - 1)^10) + (a^5*((1/(a*x) - 1)^(1/2) - 1i)^12*4096i)/(3*((1/(a*x) + 1)^(1/2) - 1)^12) + (a^5*((1/(a*x) - 1)^(1/2) - 1i)^14*2048i)/(3*((1/(a*x) + 1)^(1/2) - 1)^14))/((45*((1/(a*x) - 1)^(1/2) - 1i)^4)/((1/(a*x) + 1)^(1/2) - 1)^4 - (10*((1/(a*x) - 1)^(1/2) - 1i)^2)/((1/(a*x) + 1)^(1/2) - 1)^2 - (120*((1/(a*x) - 1)^(1/2) - 1i)^6)/((1/(a*x) + 1)^(1/2) - 1)^6 + (210*((1/(a*x) - 1)^(1/2) - 1i)^8)/((1/(a*x) + 1)^(1/2) - 1)^8 - (252*((1/(a*x) - 1)^(1/2) - 1i)^10)/((1/(a*x) + 1)^(1/2) - 1)^10 + (210*((1/(a*x) - 1)^(1/2) - 1i)^12)/((1/(a*x) + 1)^(1/2) - 1)^12 - (120*((1/(a*x) - 1)^(1/2) - 1i)^14)/((1/(a*x) + 1)^(1/2) - 1)^14 + (45*((1/(a*x) - 1)^(1/2) - 1i)^16)/((1/(a*x) + 1)^(1/2) - 1)^16 - (10*((1/(a*x) - 1)^(1/2) - 1i)^18)/((1/(a*x) + 1)^(1/2) - 1)^18 + ((1/(a*x) - 1)^(1/2) - 1i)^20)/((1/(a*x) + 1)^(1/2) - 1)^20 + 1) - ((311*a^5*((1/(a*x) - 1)^(1/2) - 1i)^5)/(4*((1/(a*x) + 1)^(1/2) - 1)^5) - (175*a^5*((1/(a*x) - 1)^(1/2) - 1i)^3))/(12*((1/(a*x) + 1)^(1/2) - 1)^3) + (8361*a^5*((1/(a*x) - 1)^(1/2) - 1i)^7)/(4*((1/(a*x) + 1)^(1/2) - 1)^7) + (42259*a^5*((1/(a*x) - 1)^(1/2) - 1i)^9)/(6*((1/(a*x) + 1)^(1/2) - 1)^9) + (25295*a^5*((1/(a*x) - 1)^(1/2) - 1i)^11)/(2*((1/(a*x) + 1)^(1/2) - 1)^11) + (25295*a^5*((1/(a*x) - 1)^(1/2) - 1i)^13)/(2*((1/(a*x) + 1)^(1/2) - 1)^13) + (42259*a^5*((1/(a*x) - 1)^(1/2) - 1i)^15)/(6*((1/(a*x) + 1)^(1/2) - 1)^15) + (8361*a^5*((1/(a*x) - 1)^(1/2) - 1i)^17)/(4*((1/(a*x) + 1)^(1/2) - 1)^17) + (175*a^5*((1/(a*x) - 1)^(1/2) - 1i)^19)/(12*((1/(a*x) + 1)^(1/2) - 1)^19) + (311*a^5*((1/(a*x) - 1)^(1/2) - 1i)^21)/(4*((1/(a*x) + 1)^(1/2) - 1)^21) + 1))

$$\begin{aligned}
& 1i)^{17} / (4 * ((1/(a*x) + 1)^{1/2} - 1)^{17}) + (311*a^5 * ((1/(a*x) - 1)^{1/2} - \\
& 1i)^{19}) / (4 * ((1/(a*x) + 1)^{1/2} - 1)^{19}) - (175*a^5 * ((1/(a*x) - 1)^{1/2} - \\
& 1i)^{21}) / (12 * ((1/(a*x) + 1)^{1/2} - 1)^{21}) + (5*a^5 * ((1/(a*x) - 1)^{1/2} - 1 \\
& i)^{23}) / (4 * ((1/(a*x) + 1)^{1/2} - 1)^{23}) + (5*a^5 * ((1/(a*x) - 1)^{1/2} - 1i) \\
&) / (4 * ((1/(a*x) + 1)^{1/2} - 1)) / ((66 * ((1/(a*x) - 1)^{1/2} - 1i)^4) / ((1/(a* \\
& x) + 1)^{1/2} - 1)^4 - (12 * ((1/(a*x) - 1)^{1/2} - 1i)^2) / ((1/(a*x) + 1)^{1/ \\
& 2} - 1)^2 - (220 * ((1/(a*x) - 1)^{1/2} - 1i)^6) / ((1/(a*x) + 1)^{1/2} - 1)^6 \\
& + (495 * ((1/(a*x) - 1)^{1/2} - 1i)^8) / ((1/(a*x) + 1)^{1/2} - 1)^8 - (792 * ((1 \\
& / (a*x) - 1)^{1/2} - 1i)^{10}) / ((1/(a*x) + 1)^{1/2} - 1)^{10} + (924 * ((1/(a*x) - \\
& 1)^{1/2} - 1i)^{12}) / ((1/(a*x) + 1)^{1/2} - 1)^{12} - (792 * ((1/(a*x) - 1)^{1/2} \\
&) - 1i)^{14}) / ((1/(a*x) + 1)^{1/2} - 1)^{14} + (495 * ((1/(a*x) - 1)^{1/2} - 1i)^{ \\
& 16}) / ((1/(a*x) + 1)^{1/2} - 1)^{16} - (220 * ((1/(a*x) - 1)^{1/2} - 1i)^{18}) / ((1/ \\
& (a*x) + 1)^{1/2} - 1)^{18} + (66 * ((1/(a*x) - 1)^{1/2} - 1i)^{20}) / ((1/(a*x) + 1 \\
&)^{1/2} - 1)^{20} - (12 * ((1/(a*x) - 1)^{1/2} - 1i)^{22}) / ((1/(a*x) + 1)^{1/2} - \\
& 1)^{22} + ((1/(a*x) - 1)^{1/2} - 1i)^{24} / ((1/(a*x) + 1)^{1/2} - 1)^{24} + 1) - \\
& ((23*a^5 * ((1/(a*x) - 1)^{1/2} - 1i)^3) / (2 * ((1/(a*x) + 1)^{1/2} - 1)^3) + (3 \\
& 33*a^5 * ((1/(a*x) - 1)^{1/2} - 1i)^5) / (2 * ((1/(a*x) + 1)^{1/2} - 1)^5) + (671 \\
& *a^5 * ((1/(a*x) - 1)^{1/2} - 1i)^7) / (2 * ((1/(a*x) + 1)^{1/2} - 1)^7) + (671*a \\
& ^5 * ((1/(a*x) - 1)^{1/2} - 1i)^9) / (2 * ((1/(a*x) + 1)^{1/2} - 1)^9) + (333*a^5 \\
& * ((1/(a*x) - 1)^{1/2} - 1i)^{11}) / (2 * ((1/(a*x) + 1)^{1/2} - 1)^{11}) + (23*a^5 * \\
& ((1/(a*x) - 1)^{1/2} - 1i)^{13}) / (2 * ((1/(a*x) + 1)^{1/2} - 1)^{13}) - (3*a^5 * ((\\
& 1/(a*x) - 1)^{1/2} - 1i)^{15}) / (2 * ((1/(a*x) + 1)^{1/2} - 1)^{15}) - (3*a^5 * ((1/ \\
& (a*x) - 1)^{1/2} - 1i)) / (2 * ((1/(a*x) + 1)^{1/2} - 1)) / ((28 * ((1/(a*x) - 1)^{ \\
& 1/2} - 1i)^4) / ((1/(a*x) + 1)^{1/2} - 1)^4 - (8 * ((1/(a*x) - 1)^{1/2} - 1i)^{ \\
& 2}) / ((1/(a*x) + 1)^{1/2} - 1)^2 - (56 * ((1/(a*x) - 1)^{1/2} - 1i)^6) / ((1/(a*x) \\
&) + 1)^{1/2} - 1)^6 + (70 * ((1/(a*x) - 1)^{1/2} - 1i)^8) / ((1/(a*x) + 1)^{1/2} \\
&) - 1)^8 - (56 * ((1/(a*x) - 1)^{1/2} - 1i)^{10}) / ((1/(a*x) + 1)^{1/2} - 1)^{10} \\
& + (28 * ((1/(a*x) - 1)^{1/2} - 1i)^{12}) / ((1/(a*x) + 1)^{1/2} - 1)^{12} - (8 * ((1/ \\
& (a*x) - 1)^{1/2} - 1i)^{14}) / ((1/(a*x) + 1)^{1/2} - 1)^{14} + ((1/(a*x) - 1)^{1 \\
& /2} - 1i)^{16} / ((1/(a*x) + 1)^{1/2} - 1)^{16} + 1) - 1 / (6*a*x^6)
\end{aligned}$$

3.87 $\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^7} dx$

Optimal result	540
Rubi [A] (verified)	541
Mathematica [A] (verified)	542
Maple [A] (verified)	543
Fricas [A] (verification not implemented)	543
Sympy [F]	543
Maxima [F]	544
Giac [F]	544
Mupad [B] (verification not implemented)	544

Optimal result

Integrand size = 12, antiderivative size = 353

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^7} dx = -\frac{a^6}{10 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^5} + \frac{a^6}{4 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^4} - \frac{5a^6}{12 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^3}$$

$$+ \frac{3a^6}{8 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)^2} - \frac{5a^6}{16 \left(1 - \sqrt{\frac{1-ax}{1+ax}}\right)} - \frac{2a^6}{7 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^7}$$

$$+ \frac{a^6}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^6} - \frac{19a^6}{10 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^5} + \frac{9a^6}{4 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^4}$$

$$- \frac{11a^6}{6 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^3} + \frac{a^6}{\left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)^2} - \frac{5a^6}{16 \left(1 + \sqrt{\frac{1-ax}{1+ax}}\right)}$$

```
[Out] -1/10*a^6/(1-((-a*x+1)/(a*x+1))^(1/2))^5+1/4*a^6/(1-((-a*x+1)/(a*x+1))^(1/2))^4-5/12*a^6/(1-((-a*x+1)/(a*x+1))^(1/2))^3+3/8*a^6/(1-((-a*x+1)/(a*x+1))^(1/2))^2-5/16*a^6/(1-((-a*x+1)/(a*x+1))^(1/2))-2/7*a^6/(1+((-a*x+1)/(a*x+1))^(1/2))^7+a^6/(1+((-a*x+1)/(a*x+1))^(1/2))^6-19/10*a^6/(1+((-a*x+1)/(a*x+1))^(1/2))^5+9/4*a^6/(1+((-a*x+1)/(a*x+1))^(1/2))^4-11/6*a^6/(1+((-a*x+1)/(a*x+1))^(1/2))^3+a^6/(1+((-a*x+1)/(a*x+1))^(1/2))^2-5/16*a^6/(1+((-a*x+1)/(a*x+1))^(1/2))
```

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 353, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6472, 1626}

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^7} dx = -\frac{5a^6}{16 \left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)} - \frac{5a^6}{16 \left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)} + \frac{3a^6}{8 \left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^2}$$

$$+ \frac{a^6}{\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^2} - \frac{5a^6}{12 \left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^3} - \frac{11a^6}{6 \left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^3}$$

$$+ \frac{a^6}{4 \left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^4} + \frac{9a^6}{4 \left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^4} - \frac{a^6}{10 \left(1 - \sqrt{\frac{1-ax}{ax+1}}\right)^5}$$

$$- \frac{a^6}{10 \left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^5} + \frac{a^6}{\left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^6} - \frac{a^6}{7 \left(\sqrt{\frac{1-ax}{ax+1}} + 1\right)^7}$$

[In] Int[1/(E^ArcSech[a*x]*x^7),x]

[Out] $-1/10*a^6/(1 - \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^5 + a^6/(4*(1 - \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^4) - (5*a^6)/(12*(1 - \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^3) + (3*a^6)/(8*(1 - \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^2) - (5*a^6)/(16*(1 - \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])) - (2*a^6)/(7*(1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^7) + a^6/(1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^6 - (19*a^6)/(10*(1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^5) + (9*a^6)/(4*(1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^4) - (11*a^6)/(6*(1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^3) + a^6/(1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)])^2 - (5*a^6)/(16*(1 + \operatorname{Sqrt}[(1 - a*x)/(1 + a*x)]))$

Rule 1626

Int[(Px_)*((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && IntegersQ[m, n]

Rule 6472

Int[E^(ArcSech[u_]*(n_.))*(x_)^(m_.), x_Symbol] :> Int[x^m*(1/u + Sqrt[(1 - u)/(1 + u)] + (1/u)*Sqrt[(1 - u)/(1 + u)])^n, x] /; FreeQ[m, x] && IntegerQ[n]

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \frac{1}{x^7 \left(\frac{1}{ax} + \sqrt{\frac{1-ax}{1+ax}} + \sqrt{\frac{1-ax}{1+ax}} \right)} dx \\
&= - \left((4a) \text{Subst} \left(\int \frac{x(a+ax^2)^5}{(-1+x)^6(1+x)^8} dx, x, \sqrt{\frac{1-ax}{1+ax}} \right) \right) \\
&= - \left((4a) \text{Subst} \left(\int \left(\frac{a^5}{8(-1+x)^6} + \frac{a^5}{4(-1+x)^5} + \frac{5a^5}{16(-1+x)^4} + \frac{3a^5}{16(-1+x)^3} + \frac{5a^5}{64(-1+x)^2} - \frac{a^5}{2(1+x)} \right) dx, x, \sqrt{\frac{1-ax}{1+ax}} \right) \right) \\
&= - \frac{a^6}{10 \left(1 - \sqrt{\frac{1-ax}{1+ax}} \right)^5} + \frac{a^6}{4 \left(1 - \sqrt{\frac{1-ax}{1+ax}} \right)^4} - \frac{5a^6}{12 \left(1 - \sqrt{\frac{1-ax}{1+ax}} \right)^3} + \frac{3a^6}{8 \left(1 - \sqrt{\frac{1-ax}{1+ax}} \right)^2} \\
&\quad - \frac{5a^6}{16 \left(1 - \sqrt{\frac{1-ax}{1+ax}} \right)} - \frac{2a^6}{7 \left(1 + \sqrt{\frac{1-ax}{1+ax}} \right)^7} + \frac{a^6}{\left(1 + \sqrt{\frac{1-ax}{1+ax}} \right)^6} - \frac{19a^6}{10 \left(1 + \sqrt{\frac{1-ax}{1+ax}} \right)^5} \\
&\quad + \frac{9a^6}{4 \left(1 + \sqrt{\frac{1-ax}{1+ax}} \right)^4} - \frac{11a^6}{6 \left(1 + \sqrt{\frac{1-ax}{1+ax}} \right)^3} + \frac{a^6}{\left(1 + \sqrt{\frac{1-ax}{1+ax}} \right)^2} - \frac{5a^6}{16 \left(1 + \sqrt{\frac{1-ax}{1+ax}} \right)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.22

$$\begin{aligned}
&\int \frac{e^{-\text{sech}^{-1}(ax)}}{x^7} dx \\
&= \frac{15 + \sqrt{\frac{1-ax}{1+ax}}(1+ax)^2(-15 + 15ax - 12a^2x^2 + 12a^3x^3 - 8a^4x^4 + 8a^5x^5)}{105ax^7}
\end{aligned}$$

[In] Integrate[1/(E^ArcSech[a*x]*x^7),x]

[Out] -1/105*(15 + Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x)^2*(-15 + 15*a*x - 12*a^2*x^2 + 12*a^3*x^3 - 8*a^4*x^4 + 8*a^5*x^5))/(a*x^7)

Maple [A] (verified)

Time = 0.94 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.22

method	result	size
default	$a \left(-\frac{1}{7a^2x^7} - \frac{\sqrt{\frac{ax+1}{ax}} \sqrt{-\frac{ax-1}{ax}} (a^2x^2-1)(8a^4x^4+12a^2x^2+15)}{105ax^6} \right)$	76

[In] `int(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^7,x,method=_RETURNVERBOSE)`

[Out] `a*(-1/7/a^2/x^7-1/105/a*((a*x+1)/a/x)^(1/2)/x^6*(-(a*x-1)/a/x)^(1/2)*(a^2*x^2-1)*(8*a^4*x^4+12*a^2*x^2+15))`

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.20

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^7} dx = -\frac{(8a^7x^7 + 4a^5x^5 + 3a^3x^3 - 15ax)\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}} + 15}{105ax^7}$$

[In] `integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^7,x, algorithm="fricas")`

[Out] `-1/105*((8*a^7*x^7 + 4*a^5*x^5 + 3*a^3*x^3 - 15*a*x)*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x)) + 15)/(a*x^7)`

Sympy [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^7} dx = a \int \frac{1}{ax^7 \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}} + x^6} dx$$

[In] `integrate(1/(1/a/x+(1/a/x-1)**(1/2)*(1+1/a/x)**(1/2))/x**7,x)`

[Out] `a*Integral(1/(a*x**7*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x)) + x**6), x)`

Maxima [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^7} dx = \int \frac{1}{x^7 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}} \right)} dx$$

[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^7,x, algorithm="maxima")

[Out] integrate(1/(x^7*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))), x)

Giac [F]

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^7} dx = \int \frac{1}{x^7 \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}} \right)} dx$$

[In] integrate(1/(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))/x^7,x, algorithm="giac")

[Out] integrate(1/(x^7*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x))), x)

Mupad [B] (verification not implemented)

Time = 5.23 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.26

$$\int \frac{e^{-\operatorname{sech}^{-1}(ax)}}{x^7} dx = -\frac{1}{7ax^7} \frac{\sqrt{\frac{1}{ax} - 1} \left(\frac{ax^2}{35} - \frac{x}{7} - \frac{1}{7a} + \frac{a^2x^3}{35} + \frac{4a^3x^4}{105} + \frac{4a^4x^5}{105} + \frac{8a^5x^6}{105} + \frac{8a^6x^7}{105} \right)}{x^7 \sqrt{\frac{1}{ax} + 1}}$$

[In] int(1/(x^7*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x))),x)

[Out] - 1/(7*a*x^7) - ((1/(a*x) - 1)^(1/2)*((a*x^2)/35 - x/7 - 1/(7*a) + (a^2*x^3)/35 + (4*a^3*x^4)/105 + (4*a^4*x^5)/105 + (8*a^5*x^6)/105 + (8*a^6*x^7)/105))/(x^7*(1/(a*x) + 1)^(1/2))

$$3.88 \quad \int \frac{e^{\operatorname{sech}^{-1}(cx)}(dx)^m}{1-c^2x^2} dx$$

Optimal result	545
Rubi [A] (verified)	545
Mathematica [A] (verified)	547
Maple [F]	547
Fricas [F]	547
Sympy [F]	548
Maxima [F]	548
Giac [F]	548
Mupad [F(-1)]	549

Optimal result

Integrand size = 24, antiderivative size = 89

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}(dx)^m}{1-c^2x^2} dx = \frac{(dx)^m \sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m}{2}, \frac{2+m}{2}, c^2x^2\right)}{cm} + \frac{(dx)^m \operatorname{Hypergeometric2F1}\left(1, \frac{m}{2}, \frac{2+m}{2}, c^2x^2\right)}{cm}$$

[Out] (d*x)^m*hypergeom([1, 1/2*m], [1+1/2*m], c^2*x^2)/c/m+(d*x)^m*hypergeom([1/2, 1/2*m], [1+1/2*m], c^2*x^2)*(1/(c*x+1))^(1/2)*(c*x+1)^(1/2)/c/m

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6476, 1972, 126, 371}

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}(dx)^m}{1-c^2x^2} dx = \frac{\sqrt{\frac{1}{cx+1}} \sqrt{cx+1} (dx)^m \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m}{2}, \frac{m+2}{2}, c^2x^2\right)}{cm} + \frac{(dx)^m \operatorname{Hypergeometric2F1}\left(1, \frac{m}{2}, \frac{m+2}{2}, c^2x^2\right)}{cm}$$

[In] Int[(E^ArcSech[c*x]*(d*x)^m)/(1 - c^2*x^2), x]

[Out] ((d*x)^m*sqrt[(1 + c*x)^(-1)]*sqrt[1 + c*x]*Hypergeometric2F1[1/2, m/2, (2 + m)/2, c^2*x^2])/(c*m) + ((d*x)^m*Hypergeometric2F1[1, m/2, (2 + m)/2, c^2*x^2])/(c*m)

Rule 126

Int[((f_)*(x_))^(p_)*((a_)+(b_)*(x_))^(m_)*((c_)+(d_)*(x_))^(n_), x_Symbol] :> Int[(a*c + b*d*x^2)^m*(f*x)^p, x] /; FreeQ[{a, b, c, d, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && GtQ[a, 0] && GtQ[c, 0]

Rule 371

Int[((c_)*(x_))^(m_)*((a_)+(b_)*(x_))^(n_))^(p_), x_Symbol] :> Simp[a^p*((c*x)^(m+1)/(c*(m+1)))*Hypergeometric2F1[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 1972

Int[(u_)*((c_)*((a_)+(b_)*(x_))^(n_))^(q_))^(p_), x_Symbol] :> Dist[Simp[(c*(a+b*x^n)^q)^p/(a+b*x^n)^(p*q)], Int[u*(a+b*x^n)^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && GeQ[a, 0]

Rule 6476

Int[(E^ArcSech[(c_)*(x_)]*((d_)*(x_))^(m_)))/((a_)+(b_)*(x_)^2), x_Symbol] :> Dist[d/(a*c), Int[(d*x)^(m-1)*(Sqrt[1/(1+c*x)]/Sqrt[1-c*x]), x], x] + Dist[d/c, Int[(d*x)^(m-1)/(a+b*x^2), x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b+a*c^2, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{d \int \frac{(dx)^{-1+m} \sqrt{\frac{1}{1+cx}}}{\sqrt{1-cx}} dx}{c} + \frac{d \int \frac{(dx)^{-1+m}}{1-c^2x^2} dx}{c} \\
 &= \frac{(dx)^m \text{Hypergeometric2F1}\left(1, \frac{m}{2}, \frac{2+m}{2}, c^2x^2\right)}{cm} + \frac{\left(d\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{(dx)^{-1+m}}{\sqrt{1-cx}\sqrt{1+cx}} dx}{c} \\
 &= \frac{(dx)^m \text{Hypergeometric2F1}\left(1, \frac{m}{2}, \frac{2+m}{2}, c^2x^2\right)}{cm} + \frac{\left(d\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{(dx)^{-1+m}}{\sqrt{1-c^2x^2}} dx}{c} \\
 &= \frac{(dx)^m \sqrt{\frac{1}{1+cx}}\sqrt{1+cx} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m}{2}, \frac{2+m}{2}, c^2x^2\right)}{cm} \\
 &\quad + \frac{(dx)^m \text{Hypergeometric2F1}\left(1, \frac{m}{2}, \frac{2+m}{2}, c^2x^2\right)}{cm}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.80 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.07

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} (dx)^m}{1 - c^2 x^2} dx$$

$$= \frac{(dx)^m \left(\frac{\sqrt{\frac{1-cx}{1+cx}} (1+cx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m}{2}, 1 + \frac{m}{2}, c^2 x^2\right)}{\sqrt{1-c^2 x^2}} + \operatorname{Hypergeometric2F1}\left(1, \frac{m}{2}, 1 + \frac{m}{2}, c^2 x^2\right) \right)}{cm}$$

[In] Integrate[(E^ArcSech[c*x]*(d*x)^m)/(1 - c^2*x^2), x]

[Out] ((d*x)^m*((Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)*Hypergeometric2F1[1/2, m/2, 1 + m/2, c^2*x^2])/Sqrt[1 - c^2*x^2] + Hypergeometric2F1[1, m/2, 1 + m/2, c^2*x^2]))/(c*m)

Maple [F]

$$\int \frac{\left(\frac{1}{cx} + \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}\right) (dx)^m}{-c^2 x^2 + 1} dx$$

[In] int((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*(d*x)^m/(-c^2*x^2+1), x)

[Out] int((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*(d*x)^m/(-c^2*x^2+1), x)

Fricas [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} (dx)^m}{1 - c^2 x^2} dx = \int -\frac{(dx)^m \left(\sqrt{\frac{1}{cx} + 1} \sqrt{\frac{1}{cx} - 1 + \frac{1}{cx}} \right)}{c^2 x^2 - 1} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*(d*x)^m/(-c^2*x^2+1), x, algorithm="fricas")

[Out] integral(-((d*x)^m*c*x*sqrt((c*x + 1)/(c*x))*sqrt(-(c*x - 1)/(c*x)) + (d*x)^m)/(c^3*x^3 - c*x), x)

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} (dx)^m}{1 - c^2 x^2} dx = - \int \frac{(dx)^m}{c^2 x^3 - x} dx + \int \frac{cx(dx)^m \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}}{c^2 x^3 - x} dx$$

[In] integrate((1/c/x+(-1+1/c/x)**(1/2)*(1+1/c/x)**(1/2))*(d*x)**m/(-c**2*x**2+1),x)

[Out] -(Integral((d*x)**m/(c**2*x**3 - x), x) + Integral(c*x*(d*x)**m*sqrt(-1 + 1/(c*x))*sqrt(1 + 1/(c*x))/(c**2*x**3 - x), x))/c

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} (dx)^m}{1 - c^2 x^2} dx = \int - \frac{(dx)^m \left(\sqrt{\frac{1}{cx} + 1} \sqrt{\frac{1}{cx} - 1 + \frac{1}{cx}} \right)}{c^2 x^2 - 1} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*(d*x)^m/(-c^2*x^2+1),x, algorithm="maxima")

[Out] -d^m*integrate(sqrt(c*x + 1)*sqrt(-c*x + 1)*x^m/(c^3*x^3 - c*x), x) - d^m*integrate(1/2*x^m/(c*x + 1), x) - d^m*integrate(1/2*x^m/(c*x - 1), x) + d^m*x^m/(c*m)

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} (dx)^m}{1 - c^2 x^2} dx = \int - \frac{(dx)^m \left(\sqrt{\frac{1}{cx} + 1} \sqrt{\frac{1}{cx} - 1 + \frac{1}{cx}} \right)}{c^2 x^2 - 1} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*(d*x)^m/(-c^2*x^2+1),x, algorithm="giac")

[Out] integrate(-(d*x)^m*(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(c^2*x^2 - 1), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} (dx)^m}{1 - c^2 x^2} dx = - \int \frac{\left(\sqrt{\frac{1}{cx} - 1} \sqrt{\frac{1}{cx} + 1} + \frac{1}{cx} \right) (dx)^m}{c^2 x^2 - 1} dx$$

```
[In] int(-(((1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2) + 1/(c*x))*(d*x)^m)/(c^2*x^2 - 1), x)
```

```
[Out] -int(((1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2) + 1/(c*x))*(d*x)^m)/(c^2*x^2 - 1), x)
```

$$3.89 \quad \int \frac{e^{\operatorname{sech}^{-1}(cx)} x^4}{1-c^2 x^2} dx$$

Optimal result	550
Rubi [A] (verified)	550
Mathematica [A] (verified)	552
Maple [A] (verified)	552
Fricas [A] (verification not implemented)	553
Sympy [F]	553
Maxima [F]	553
Giac [F]	554
Mupad [B] (verification not implemented)	554

Optimal result

Integrand size = 22, antiderivative size = 88

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^4}{1-c^2 x^2} dx = -\frac{x^2}{2c^3} - \frac{2\sqrt{1-cx}}{3c^5 \sqrt{\frac{1}{1+cx}}} - \frac{x^2 \sqrt{1-cx}}{3c^3 \sqrt{\frac{1}{1+cx}}} - \frac{\log(1-c^2 x^2)}{2c^5}$$

[Out] $-1/2*x^2/c^3 - 1/2*\ln(-c^2*x^2+1)/c^5 - 2/3*(-c*x+1)^{(1/2)}/c^5/(1/(c*x+1))^{(1/2)}$
 $-1/3*x^2*(-c*x+1)^{(1/2)}/c^3/(1/(c*x+1))^{(1/2)}$

Rubi [A] (verified)

Time = 0.13 (sec), antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.318$, Rules used = {6476, 1972, 102, 12, 75, 272, 45}

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^4}{1-c^2 x^2} dx = -\frac{2\sqrt{1-cx}}{3c^5 \sqrt{\frac{1}{cx+1}}} - \frac{x^2 \sqrt{1-cx}}{3c^3 \sqrt{\frac{1}{cx+1}}} - \frac{x^2}{2c^3} - \frac{\log(1-c^2 x^2)}{2c^5}$$

[In] `Int[(E^ArcSech[c*x]*x^4)/(1 - c^2*x^2),x]`

[Out] $-1/2*x^2/c^3 - (2*\text{Sqrt}[1 - c*x])/(3*c^5*\text{Sqrt}[(1 + c*x)^{-1}]) - (x^2*\text{Sqrt}[1 - c*x])/(3*c^3*\text{Sqrt}[(1 + c*x)^{-1}]) - \text{Log}[1 - c^2*x^2]/(2*c^5)$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 45

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && Le
Q[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

Rule 75

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ
[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]
```

Rule 102

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_
))^p_., x_Symbol] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x
)^(p + 1)/(d*f*(m + n + p + 1))), x] + Dist[1/(d*f*(m + n + p + 1)), Int[(a
+ b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b
*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1)) + b*(a*d*f*(2*m + n + p) - b*
(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p
}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 1972

```
Int[(u_.)*((c_.)*((a_.) + (b_.)*(x_)^(n_.))^(q_.))^(p_), x_Symbol] := Dist[S
imp[(c*(a + b*x^n)^q)^p/(a + b*x^n)^(p*q)], Int[u*(a + b*x^n)^(p*q), x], x]
/; FreeQ[{a, b, c, n, p, q}, x] && GeQ[a, 0]
```

Rule 6476

```
Int[(E^ArcSech[(c_.)*(x_)])*((d_.)*(x_))^(m_.)/((a_) + (b_.)*(x_)^2), x_Sym
bol] := Dist[d/(a*c), Int[(d*x)^(m - 1)*(Sqrt[1/(1 + c*x)]/Sqrt[1 - c*x]),
x], x] + Dist[d/c, Int[(d*x)^(m - 1)/(a + b*x^2), x], x] /; FreeQ[{a, b, c,
d, m}, x] && EqQ[b + a*c^2, 0]
```

Rubi steps

$$\text{integral} = \frac{\int \frac{x^3 \sqrt{\frac{1}{1+cx}}}{\sqrt{1-cx}} dx}{c} + \frac{\int \frac{x^3}{1-c^2x^2} dx}{c}$$

$$\begin{aligned}
&= \frac{\text{Subst}\left(\int \frac{x}{1-c^2x} dx, x, x^2\right)}{2c} + \frac{\left(\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{x^3}{\sqrt{1-cx}\sqrt{1+cx}} dx}{c} \\
&= -\frac{x^2\sqrt{1-cx}}{3c^3\sqrt{\frac{1}{1+cx}}} + \frac{\text{Subst}\left(\int \left(-\frac{1}{c^2} - \frac{1}{c^2(-1+c^2x)}\right) dx, x, x^2\right)}{2c} \\
&\quad - \frac{\left(\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int -\frac{2x}{\sqrt{1-cx}\sqrt{1+cx}} dx}{3c^3} \\
&= -\frac{x^2}{2c^3} - \frac{x^2\sqrt{1-cx}}{3c^3\sqrt{\frac{1}{1+cx}}} - \frac{\log(1-c^2x^2)}{2c^5} + \frac{\left(2\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{x}{\sqrt{1-cx}\sqrt{1+cx}} dx}{3c^3} \\
&= -\frac{x^2}{2c^3} - \frac{2\sqrt{1-cx}}{3c^5\sqrt{\frac{1}{1+cx}}} - \frac{x^2\sqrt{1-cx}}{3c^3\sqrt{\frac{1}{1+cx}}} - \frac{\log(1-c^2x^2)}{2c^5}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.78

$$\int \frac{e^{\text{sech}^{-1}(cx)} x^4}{1-c^2x^2} dx = -\frac{3c^2x^2 + 2\sqrt{\frac{1-cx}{1+cx}}(2+2cx+c^2x^2+c^3x^3) + 3\log(1-c^2x^2)}{6c^5}$$

[In] Integrate[(E^ArcSech[c*x]*x^4)/(1 - c^2*x^2), x]

[Out] -1/6*(3*c^2*x^2 + 2*Sqrt[(1 - c*x)/(1 + c*x)]*(2 + 2*c*x + c^2*x^2 + c^3*x^3) + 3*Log[1 - c^2*x^2])/c^5

Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.84

method	result	size
default	$-\frac{\sqrt{\frac{-cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}} (c^2x^2+2)}{3c^4} + \frac{-\frac{x^2}{2c^2} - \frac{\ln(c^2x^2-1)}{2c^4}}{c}$	74

[In] int((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x^4/(-c^2*x^2+1), x, method=_RETURNVERBOSE)

[Out] -1/3*(-(c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*(c^2*x^2+2)/c^4+1/c*(-1/2*x^2/c^2-1/2/c^4*ln(c^2*x^2-1))

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.78

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^4}{1 - c^2 x^2} dx = -\frac{3c^2 x^2 + 2(c^3 x^3 + 2cx) \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} + 3 \log(c^2 x^2 - 1)}{6c^5}$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x^4/(-c^2*x^2+1),x, algorithm="fricas")

[Out] -1/6*(3*c^2*x^2 + 2*(c^3*x^3 + 2*c*x)*sqrt((c*x + 1)/(c*x))*sqrt(-(c*x - 1)/(c*x)) + 3*log(c^2*x^2 - 1))/c^5

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^4}{1 - c^2 x^2} dx = -\frac{\int \frac{x^3}{c^2 x^2 - 1} dx + \int \frac{cx^4 \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}}{c^2 x^2 - 1} dx}{c}$$

[In] integrate((1/c/x+(-1+1/c/x)**(1/2)*(1+1/c/x)**(1/2))*x**4/(-c**2*x**2+1),x)

[Out] -(Integral(x**3/(c**2*x**2 - 1), x) + Integral(c*x**4*sqrt(-1 + 1/(c*x))*sqrt(1 + 1/(c*x))/(c**2*x**2 - 1), x))/c

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^4}{1 - c^2 x^2} dx = \int -\frac{x^4 \left(\sqrt{\frac{1}{cx} + 1} \sqrt{\frac{1}{cx} - 1} + \frac{1}{cx} \right)}{c^2 x^2 - 1} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x^4/(-c^2*x^2+1),x, algorithm="maxima")

[Out] -integrate(x, x)/c^3 - 1/2*log(c*x + 1)/c^5 - 1/2*log(c*x - 1)/c^5 - integrate(sqrt(c*x + 1)*sqrt(-c*x + 1)*x^3/(c^3*x^2 - c), x)

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^4}{1 - c^2 x^2} dx = \int -\frac{x^4 \left(\sqrt{\frac{1}{cx} + 1} \sqrt{\frac{1}{cx} - 1} + \frac{1}{cx} \right)}{c^2 x^2 - 1} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x^4/(-c^2*x^2+1),x, algorithm="giac")

[Out] integrate(-x^4*(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(c^2*x^2 - 1), x)

Mupad [B] (verification not implemented)

Time = 5.42 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.86

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^4}{1 - c^2 x^2} dx = -\frac{\ln(c^2 x^2 - 1) + c^2 x^2}{2 c^5} - x^3 \sqrt{\frac{1}{cx} - 1} \left(\frac{\sqrt{\frac{1}{cx} + 1}}{3 c^2} + \frac{2 \sqrt{\frac{1}{cx} + 1}}{3 c^4 x^2} \right)$$

[In] int(-(x^4*((1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2) + 1/(c*x)))/(c^2*x^2 - 1),x)

[Out] - (log(c^2*x^2 - 1) + c^2*x^2)/(2*c^5) - x^3*(1/(c*x) - 1)^(1/2)*((1/(c*x) + 1)^(1/2)/(3*c^2) + (2*(1/(c*x) + 1)^(1/2))/(3*c^4*x^2))

3.90 $\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^3}{1-c^2 x^2} dx$

Optimal result	555
Rubi [A] (verified)	555
Mathematica [C] (verified)	557
Maple [C] (verified)	557
Fricas [B] (verification not implemented)	558
Sympy [F]	558
Maxima [F]	558
Giac [F]	559
Mupad [B] (verification not implemented)	559

Optimal result

Integrand size = 22, antiderivative size = 75

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^3}{1-c^2 x^2} dx = -\frac{x}{c^3} - \frac{x\sqrt{1-cx}}{2c^3\sqrt{\frac{1}{1+cx}}} + \frac{\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\arcsin(cx)}{2c^4} + \frac{\operatorname{arctanh}(cx)}{c^4}$$

[Out] $-x/c^3 + \operatorname{arctanh}(c*x)/c^4 - 1/2*x*(-c*x+1)^{(1/2)}/c^3/(1/(c*x+1))^{(1/2)} + 1/2*\arcsin(c*x)*(1/(c*x+1))^{(1/2)}*(c*x+1)^{(1/2)}/c^4$

Rubi [A] (verified)

Time = 0.12 (sec), antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.318$, Rules used = {6476, 1972, 92, 41, 222, 327, 212}

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^3}{1-c^2 x^2} dx = \frac{\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\arcsin(cx)}{2c^4} + \frac{\operatorname{arctanh}(cx)}{c^4} - \frac{x\sqrt{1-cx}}{2c^3\sqrt{\frac{1}{cx+1}}} - \frac{x}{c^3}$$

[In] $\operatorname{Int}[(E^{\operatorname{ArcSech}[c*x]}*x^3)/(1-c^2*x^2), x]$

[Out] $-(x/c^3) - (x*\operatorname{Sqrt}[1-c*x])/(2*c^3*\operatorname{Sqrt}[(1+c*x)^{-1}]) + (\operatorname{Sqrt}[(1+c*x)^{-1}]*\operatorname{Sqrt}[1+c*x]*\operatorname{ArcSin}[c*x])/(2*c^4) + \operatorname{ArcTanh}[c*x]/c^4$

Rule 41

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+) + (d_+)*(x_+))^{(m_+)}, x_Symbol] \rightarrow \operatorname{Int}[(a*c + b*d*x^2)^m, x] /; \operatorname{FreeQ}\{a, b, c, d, m\}, x] \&\& \operatorname{EqQ}[b*c + a*d, 0] \&\& (\operatorname{IntegerQ}[m] \parallel (\operatorname{GtQ}[a, 0] \&\& \operatorname{GtQ}[c, 0]))$

Rule 92

```
Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.))*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 222

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

Rule 327

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 1972

```
Int[(u_.)*((c_.)*((a_.) + (b_.)*(x_)^(n_.))^(q_))^(p_), x_Symbol] := Dist[Simp[(c*(a + b*x^n)^q)^p/(a + b*x^n)^(p*q)], Int[u*(a + b*x^n)^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && GeQ[a, 0]
```

Rule 6476

```
Int[(E^ArcSech[(c_.)*(x_)])*((d_.)*(x_))^(m_.))/((a_) + (b_.)*(x_)^2), x_Symbol] := Dist[d/(a*c), Int[(d*x)^(m - 1)*(Sqrt[1/(1 + c*x)]/Sqrt[1 - c*x]), x], x] + Dist[d/c, Int[(d*x)^(m - 1)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + a*c^2, 0]
```

Rubi steps

$$\text{integral} = \frac{\int \frac{x^2 \sqrt{\frac{1}{1+cx}}}{\sqrt{1-cx}} dx}{c} + \frac{\int \frac{x^2}{1-c^2x^2} dx}{c}$$

$$\begin{aligned}
&= -\frac{x}{c^3} + \frac{\int \frac{1}{1-c^2x^2} dx}{c^3} + \frac{\left(\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{x^2}{\sqrt{1-cx}\sqrt{1+cx}} dx}{c} \\
&= -\frac{x}{c^3} - \frac{x\sqrt{1-cx}}{2c^3\sqrt{\frac{1}{1+cx}}} + \frac{\operatorname{arctanh}(cx)}{c^4} + \frac{\left(\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{1}{\sqrt{1-cx}\sqrt{1+cx}} dx}{2c^3} \\
&= -\frac{x}{c^3} - \frac{x\sqrt{1-cx}}{2c^3\sqrt{\frac{1}{1+cx}}} + \frac{\operatorname{arctanh}(cx)}{c^4} + \frac{\left(\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{1}{\sqrt{1-c^2x^2}} dx}{2c^3} \\
&= -\frac{x}{c^3} - \frac{x\sqrt{1-cx}}{2c^3\sqrt{\frac{1}{1+cx}}} + \frac{\sqrt{\frac{1}{1+cx}}\sqrt{1+cx} \arcsin(cx)}{2c^4} + \frac{\operatorname{arctanh}(cx)}{c^4}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.20 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.47

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^3}{1-c^2x^2} dx = \frac{2cx + cx\sqrt{\frac{1-cx}{1+cx}} + c^2x^2\sqrt{\frac{1-cx}{1+cx}} + \log(1-cx) - \log(1+cx) - i \log\left(-2icx + 2\sqrt{\frac{1-cx}{1+cx}}(1+cx)\right)}{2c^4}$$

[In] Integrate[(E^ArcSech[c*x]*x^3)/(1 - c^2*x^2), x]

[Out] -1/2*(2*c*x + c*x*Sqrt[(1 - c*x)/(1 + c*x)] + c^2*x^2*Sqrt[(1 - c*x)/(1 + c*x)]) + Log[1 - c*x] - Log[1 + c*x] - I*Log[(-2*I)*c*x + 2*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)]/c^4

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.68 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.63

method	result	size
default	$-\frac{\sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}} \left(x \sqrt{-c^2x^2+1} \operatorname{csgn}(c) c - \arctan\left(\frac{\operatorname{csgn}(c)cx}{\sqrt{-c^2x^2+1}}\right) \right) \operatorname{csgn}(c)}{2c^3\sqrt{-c^2x^2+1}} + \frac{-\frac{x}{c^2} + \frac{\ln(cx+1)}{2c^3} - \frac{\ln(cx-1)}{2c^3}}{c}$	122

[In] int((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x^3/(-c^2*x^2+1), x, method=_RETURNVERBOSE)

[Out] $-1/2*(-(c*x-1)/c/x)^{(1/2)}*x*((c*x+1)/c/x)^{(1/2)}/c^3*(x*(-c^2*x^2+1)^{(1/2)}*c$
 $\text{sgn}(c)*c-\arctan(\text{csgn}(c)*c*x/(-c^2*x^2+1)^{(1/2)})/(-c^2*x^2+1)^{(1/2)}*\text{csgn}(c)$
 $+1/c*(-x/c^2+1/2/c^3*\ln(c*x+1)-1/2/c^3*\ln(c*x-1))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 91 vs. $2(45) = 90$.

Time = 0.25 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.21

$$\int \frac{e^{\text{sech}^{-1}(cx)} x^3}{1 - c^2 x^2} dx$$

$$= -\frac{c^2 x^2 \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} + 2cx + \arctan\left(\sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}}\right) - \log(cx+1) + \log(cx-1)}{2c^4}$$

[In] `integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x^3/(-c^2*x^2+1),x, algorithm="fricas")`

[Out] $-1/2*(c^2*x^2*\text{sqrt}((c*x+1)/(c*x))*\text{sqrt}(-(c*x-1)/(c*x))+2*c*x+\arctan(\text{sqrt}((c*x+1)/(c*x))*\text{sqrt}(-(c*x-1)/(c*x)))-\log(c*x+1)+\log(c*x-1))/c^4$

Sympy [F]

$$\int \frac{e^{\text{sech}^{-1}(cx)} x^3}{1 - c^2 x^2} dx = -\frac{\int \frac{x^2}{c^2 x^2 - 1} dx + \int \frac{cx^3 \sqrt{-1 + \frac{1}{cx}} \sqrt{1 + \frac{1}{cx}}}{c^2 x^2 - 1} dx}{c}$$

[In] `integrate((1/c/x+(-1+1/c/x)**(1/2)*(1+1/c/x)**(1/2))*x**3/(-c**2*x**2+1),x)`

[Out] $-(\text{Integral}(x**2/(c**2*x**2-1),x)+\text{Integral}(c*x**3*\text{sqrt}(-1+1/(c*x))*\text{sqrt}(1+1/(c*x))/(c**2*x**2-1),x))/c$

Maxima [F]

$$\int \frac{e^{\text{sech}^{-1}(cx)} x^3}{1 - c^2 x^2} dx = \int -\frac{x^3 \left(\sqrt{\frac{1}{cx} + 1} \sqrt{\frac{1}{cx} - 1 + \frac{1}{cx}} \right)}{c^2 x^2 - 1} dx$$

[In] `integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x^3/(-c^2*x^2+1),x, algorithm="maxima")`

[Out] $-x/c^3 + 1/2*\log(c*x+1)/c^4 - 1/2*\log(c*x-1)/c^4 - \text{integrate}(\text{sqrt}(c*x+1)*\text{sqrt}(-c*x+1)*x^2/(c^3*x^2-c),x)$

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^3}{1 - c^2 x^2} dx = \int -\frac{x^3 \left(\sqrt{\frac{1}{cx} + 1} \sqrt{\frac{1}{cx} - 1} + \frac{1}{cx} \right)}{c^2 x^2 - 1} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x^3/(-c^2*x^2+1),x, algo rithm="giac")

[Out] integrate(-x^3*(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(c^2*x^2 - 1), x)

Mupad [B] (verification not implemented)

Time = 11.19 (sec) , antiderivative size = 340, normalized size of antiderivative = 4.53

$$\begin{aligned} \int \frac{e^{\operatorname{sech}^{-1}(cx)} x^3}{1 - c^2 x^2} dx &= \frac{\operatorname{atanh}(cx) - cx}{c^4} - \frac{\ln \left(\frac{\sqrt{\frac{1}{cx} - 1} - i}{\sqrt{\frac{1}{cx} + 1} - 1} \right) \operatorname{li}}{2c^4} \\ &\quad - \frac{\frac{\operatorname{li}}{32c^4} + \frac{\left(\sqrt{\frac{1}{cx} - 1} - i\right)^2 \operatorname{li}}{16c^4 \left(\sqrt{\frac{1}{cx} + 1} - 1\right)^2} - \frac{\left(\sqrt{\frac{1}{cx} - 1} - i\right)^4 15i}{32c^4 \left(\sqrt{\frac{1}{cx} + 1} - 1\right)^4}}{\frac{\left(\sqrt{\frac{1}{cx} - 1} - i\right)^2}{\left(\sqrt{\frac{1}{cx} + 1} - 1\right)^2} + \frac{2\left(\sqrt{\frac{1}{cx} - 1} - i\right)^4}{\left(\sqrt{\frac{1}{cx} + 1} - 1\right)^4} + \frac{\left(\sqrt{\frac{1}{cx} - 1} - i\right)^6}{\left(\sqrt{\frac{1}{cx} + 1} - 1\right)^6}} \\ &\quad + \frac{\ln \left(\frac{2c\sqrt{\frac{c+\frac{1}{x}}{c}} - \frac{2}{x} + c\sqrt{-\frac{c-\frac{1}{x}}{c}} 2i}{2c+\frac{1}{x} - 2c\sqrt{\frac{c+\frac{1}{x}}{c}}} \right) \operatorname{li}}{2c^4} - \frac{\left(\sqrt{\frac{1}{cx} - 1} - i\right)^2 \operatorname{li}}{32c^4 \left(\sqrt{\frac{1}{cx} + 1} - 1\right)^2} \end{aligned}$$

[In] int(-(x^3*((1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2) + 1/(c*x)))/(c^2*x^2 - 1),x)

[Out] (atanh(c*x) - c*x)/c^4 - (log(((1/(c*x) - 1)^(1/2) - 1i)/((1/(c*x) + 1)^(1/2) - 1))*1i)/(2*c^4) - (1i/(32*c^4) + (((1/(c*x) - 1)^(1/2) - 1i)^2*1i)/(16*c^4*((1/(c*x) + 1)^(1/2) - 1)^2) - (((1/(c*x) - 1)^(1/2) - 1i)^4*15i)/(32*c^4*((1/(c*x) + 1)^(1/2) - 1)^4))/(((1/(c*x) - 1)^(1/2) - 1i)^2/((1/(c*x) + 1)^(1/2) - 1)^2 + (2*((1/(c*x) - 1)^(1/2) - 1i)^4)/((1/(c*x) + 1)^(1/2) - 1)^4 + ((1/(c*x) - 1)^(1/2) - 1i)^6/((1/(c*x) + 1)^(1/2) - 1)^6) + (log((c*(-(c - 1/x)/c)^(1/2)*2i - 2/x + 2*c*((c + 1/x)/c)^(1/2))/(2*c + 1/x - 2*c*((c + 1/x)/c)^(1/2)))*1i)/(2*c^4) - (((1/(c*x) - 1)^(1/2) - 1i)^2*1i)/(32*c^4*((1/(c*x) + 1)^(1/2) - 1)^2)

3.91 $\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^2}{1-c^2 x^2} dx$

Optimal result	560
Rubi [A] (verified)	560
Mathematica [A] (verified)	561
Maple [A] (verified)	562
Fricas [A] (verification not implemented)	562
Sympy [F]	562
Maxima [F]	563
Giac [F]	563
Mupad [B] (verification not implemented)	563

Optimal result

Integrand size = 22, antiderivative size = 45

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^2}{1-c^2 x^2} dx = -\frac{\sqrt{1-cx}}{c^3 \sqrt{\frac{1}{1+cx}}} - \frac{\log(1-c^2 x^2)}{2c^3}$$

[Out] $-1/2*\ln(-c^2*x^2+1)/c^3-(-c*x+1)^{(1/2)}/c^3/(1/(c*x+1))^{(1/2)}$

Rubi [A] (verified)

Time = 0.10 (sec), antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {6476, 1972, 75, 266}

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^2}{1-c^2 x^2} dx = -\frac{\sqrt{1-cx}}{c^3 \sqrt{\frac{1}{cx+1}}} - \frac{\log(1-c^2 x^2)}{2c^3}$$

[In] $\text{Int}[(E^{\text{ArcSech}[c*x]}*x^2)/(1-c^2*x^2),x]$

[Out] $-(\text{Sqrt}[1-c*x]/(c^3*\text{Sqrt}[(1+c*x)^{-1}]))) - \text{Log}[1-c^2*x^2]/(2*c^3)$

Rule 75

$\text{Int}[(a_.) + (b_.)*(x_.)*((c_.) + (d_.)*(x_.)^{(n_.)}*((e_.) + (f_.)*(x_.)^{(p_.)}), x_Symbol] :> \text{Simp}[b*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)}/(d*f*(n + p + 2))), x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, n, p, x\}$ && $\text{NeQ}[n + p + 2, 0]$ && $\text{EqQ}[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]$

Rule 266


```
Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]
```

Rule 1972

```
Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Dist[Simp[(c*(a + b*x^n)^q)^p/(a + b*x^n)^(p*q)], Int[u*(a + b*x^n)^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && GeQ[a, 0]
```

Rule 6476

```
Int[(E^ArcSech[(c_)*(x_)])*((d_)*(x_)^(m_))/((a_) + (b_)*(x_)^2), x_Symbol] := Dist[d/(a*c), Int[(d*x)^(m - 1)*(Sqrt[1/(1 + c*x)]/Sqrt[1 - c*x]), x], x] + Dist[d/c, Int[(d*x)^(m - 1)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + a*c^2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \frac{x\sqrt{\frac{1}{1+cx}}}{\sqrt{1-cx}} dx}{c} + \frac{\int \frac{x}{1-c^2x^2} dx}{c} \\ &= -\frac{\log(1-c^2x^2)}{2c^3} + \frac{\left(\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{x}{\sqrt{1-cx}\sqrt{1+cx}} dx}{c} \\ &= -\frac{\sqrt{1-cx}}{c^3\sqrt{\frac{1}{1+cx}}} - \frac{\log(1-c^2x^2)}{2c^3} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.98

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^2}{1-c^2x^2} dx = -\frac{2\sqrt{\frac{1-cx}{1+cx}}(1+cx) + \log(1-c^2x^2)}{2c^3}$$

```
[In] Integrate[(E^ArcSech[c*x]*x^2)/(1 - c^2*x^2), x]
```

```
[Out] -1/2*(2*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x) + Log[1 - c^2*x^2])/c^3
```

Maple [A] (verified)

Time = 0.70 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.16

method	result	size
default	$-\frac{\sqrt{-\frac{cx-1}{cx}}x\sqrt{\frac{cx+1}{cx}}}{c^2} - \frac{\ln(c^2x^2-1)}{2c^3}$	52

[In] int((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x^2/(-c^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] $-\left(-\frac{cx-1}{cx}\right)^{1/2}x\left(\frac{cx+1}{cx}\right)^{1/2}/c^2-1/2/c^3*\ln(c^2*x^2-1)$

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.09

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}x^2}{1-c^2x^2} dx = -\frac{2cx\sqrt{\frac{cx+1}{cx}}\sqrt{-\frac{cx-1}{cx}} + \log(c^2x^2-1)}{2c^3}$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x^2/(-c^2*x^2+1),x, algorithm="fricas")

[Out] $-1/2*(2*c*x*\sqrt{(c*x + 1)/(c*x)}*\sqrt{-(c*x - 1)/(c*x)} + \log(c^2*x^2 - 1))/c^3$

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}x^2}{1-c^2x^2} dx = -\int \frac{x}{c^2x^2-1} dx + \int \frac{cx^2\sqrt{-1+\frac{1}{cx}}\sqrt{1+\frac{1}{cx}}}{c^2x^2-1} dx$$

[In] integrate((1/c/x+(-1+1/c/x)**(1/2)*(1+1/c/x)**(1/2))*x**2/(-c**2*x**2+1),x)

[Out] $-(\operatorname{Integral}(x/(c**2*x**2 - 1), x) + \operatorname{Integral}(c*x**2*\sqrt{-1 + 1/(c*x)}*\sqrt{1 + 1/(c*x)})/(c**2*x**2 - 1), x))/c$

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^2}{1 - c^2 x^2} dx = \int -\frac{x^2 \left(\sqrt{\frac{1}{cx} + 1} \sqrt{\frac{1}{cx} - 1} + \frac{1}{cx} \right)}{c^2 x^2 - 1} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x^2/(-c^2*x^2+1),x, algorithm="maxima")

[Out] -1/2*log(c*x + 1)/c^3 - 1/2*log(c*x - 1)/c^3 - integrate(sqrt(c*x + 1)*sqrt(-c*x + 1)*x/(c^3*x^2 - c), x)

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^2}{1 - c^2 x^2} dx = \int -\frac{x^2 \left(\sqrt{\frac{1}{cx} + 1} \sqrt{\frac{1}{cx} - 1} + \frac{1}{cx} \right)}{c^2 x^2 - 1} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x^2/(-c^2*x^2+1),x, algorithm="giac")

[Out] integrate(-x^2*(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(c^2*x^2 - 1), x)

Mupad [B] (verification not implemented)

Time = 5.24 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.98

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x^2}{1 - c^2 x^2} dx = -\frac{\ln(c^2 x^2 - 1)}{2 c^3} - \frac{x \sqrt{\frac{1}{cx} - 1} \sqrt{\frac{1}{cx} + 1}}{c^2}$$

[In] int(-(x^2*((1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2) + 1/(c*x)))/(c^2*x^2 - 1),x)

[Out] - log(c^2*x^2 - 1)/(2*c^3) - (x*(1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2))/c^2

3.92 $\int \frac{e^{\operatorname{sech}^{-1}(cx)x}}{1-c^2x^2} dx$

Optimal result	564
Rubi [A] (verified)	564
Mathematica [C] (verified)	565
Maple [C] (verified)	566
Fricas [B] (verification not implemented)	566
Sympy [F]	566
Maxima [F]	567
Giac [F]	567
Mupad [B] (verification not implemented)	567

Optimal result

Integrand size = 20, antiderivative size = 37

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)x}}{1-c^2x^2} dx = \frac{\sqrt{\frac{1}{1+cx}} \sqrt{1+cx} \arcsin(cx)}{c^2} + \frac{\operatorname{arctanh}(cx)}{c^2}$$

[Out] $\operatorname{arctanh}(c*x)/c^2 + \arcsin(c*x)*(1/(c*x+1))^{(1/2)}*(c*x+1)^{(1/2)}/c^2$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6476, 1972, 41, 222, 212}

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)x}}{1-c^2x^2} dx = \frac{\sqrt{\frac{1}{cx+1}} \sqrt{cx+1} \arcsin(cx)}{c^2} + \frac{\operatorname{arctanh}(cx)}{c^2}$$

[In] $\operatorname{Int}[(E^{\operatorname{ArcSech}[c*x]*x})/(1 - c^2*x^2), x]$

[Out] $(\operatorname{Sqrt}[(1 + c*x)^{-1}]*\operatorname{Sqrt}[1 + c*x]*\operatorname{ArcSin}[c*x])/c^2 + \operatorname{ArcTanh}[c*x]/c^2$

Rule 41

$\operatorname{Int}[(a_ + (b_)*(x_))^{(m_)}*((c_ + (d_)*(x_))^{(m_)}), x_Symbol] \rightarrow \operatorname{Int}[(a*c + b*d*x^2)^m, x] /; \operatorname{FreeQ}\{a, b, c, d, m\}, x] \ \&\& \operatorname{EqQ}[b*c + a*d, 0] \ \&\& (\operatorname{IntegerQ}[m] \ || \ (\operatorname{GtQ}[a, 0] \ \&\& \operatorname{GtQ}[c, 0]))$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}], x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ \&\& \operatorname{GtQ}[b, 0])$

Q[a, 0] || LtQ[b, 0])

Rule 222

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

Rule 1972

Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] := Dist[Simp[(c*(a + b*x^n)^q)^p/(a + b*x^n)^(p*q)], Int[u*(a + b*x^n)^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && GeQ[a, 0]

Rule 6476

Int[(E^ArcSech[(c_)*(x_)])*((d_)*(x_)^(m_)))/((a_) + (b_)*(x_)^2), x_Symbol] := Dist[d/(a*c), Int[(d*x)^(m-1)*(Sqrt[1/(1+c*x)]/Sqrt[1-c*x]), x], x] + Dist[d/c, Int[(d*x)^(m-1)/(a+b*x^2), x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + a*c^2, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int \frac{\sqrt{\frac{1}{1+cx}}}{\sqrt{1-cx}} dx}{c} + \frac{\int \frac{1}{1-c^2x^2} dx}{c} \\
 &= \frac{\operatorname{arctanh}(cx)}{c^2} + \frac{\left(\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{1}{\sqrt{1-cx}\sqrt{1+cx}} dx}{c} \\
 &= \frac{\operatorname{arctanh}(cx)}{c^2} + \frac{\left(\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{1}{\sqrt{1-c^2x^2}} dx}{c} \\
 &= \frac{\sqrt{\frac{1}{1+cx}}\sqrt{1+cx} \arcsin(cx)}{c^2} + \frac{\operatorname{arctanh}(cx)}{c^2}
 \end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.20 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.84

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x}{1-c^2x^2} dx = -\frac{\log(1-cx)}{2c^2} + \frac{\log(1+cx)}{2c^2} + \frac{i \log\left(-2icx + 2\sqrt{\frac{1-cx}{1+cx}}(1+cx)\right)}{c^2}$$

[In] Integrate[(E^ArcSech[c*x]*x)/(1 - c^2*x^2), x]

[Out] -1/2*Log[1 - c*x]/c^2 + Log[1 + c*x]/(2*c^2) + (I*Log[(-2*I)*c*x + 2*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x)])/c^2

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.69 (sec) , antiderivative size = 97, normalized size of antiderivative = 2.62

method	result	size
default	$\frac{\sqrt{-\frac{cx-1}{cx}} x \sqrt{\frac{cx+1}{cx}} \arctan\left(\frac{\operatorname{csgn}(c)cx}{\sqrt{-(cx-1)(cx+1)}}\right) \operatorname{csgn}(c)}{\sqrt{-c^2x^2+1}c} + \frac{\ln(cx+1)}{2c} - \frac{\ln(cx-1)}{2c}$	97

[In] `int((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x/(-c^2*x^2+1),x,method=_RETURNVERBOSE)`

[Out] $(-(c*x-1)/c/x)^{(1/2)}*x*((c*x+1)/c/x)^{(1/2)}*\arctan(\operatorname{csgn}(c)*c*x/(-(c*x-1)*(c*x+1))^{(1/2)})/(-c^2*x^2+1)^{(1/2)}*\operatorname{csgn}(c)/c+1/c*(1/2/c*\ln(c*x+1)-1/2/c*\ln(c*x-1))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. $2(17) = 34$.

Time = 0.25 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.43

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x}{1 - c^2 x^2} dx = -\frac{2 \arctan\left(\sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}}\right) - \log(cx+1) + \log(cx-1)}{2c^2}$$

[In] `integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x/(-c^2*x^2+1),x,algorithm="fricas")`

[Out] $-1/2*(2*\arctan(\sqrt{(c*x+1)/(c*x)})*\sqrt{-(c*x-1)/(c*x)}) - \log(c*x+1) + \log(c*x-1))/c^2$

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x}{1 - c^2 x^2} dx = -\int \frac{cx\sqrt{-1+\frac{1}{cx}}\sqrt{1+\frac{1}{cx}}}{c^2x^2-1} dx + \int \frac{1}{c^2x^2-1} dx$$

[In] `integrate((1/c/x+(-1+1/c/x)**(1/2)*(1+1/c/x)**(1/2))*x/(-c**2*x**2+1),x)`

[Out] $-(\operatorname{Integral}(c*x*\sqrt{-1+1/(c*x)})*\sqrt{1+1/(c*x)})/(c**2*x**2-1), x) + \operatorname{Integral}(1/(c**2*x**2-1), x)/c$

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x}{1 - c^2 x^2} dx = \int -\frac{x \left(\sqrt{\frac{1}{cx} + 1} \sqrt{\frac{1}{cx} - 1} + \frac{1}{cx} \right)}{c^2 x^2 - 1} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x/(-c^2*x^2+1),x, algorithm="maxima")

[Out] 1/2*log(c*x + 1)/c^2 - 1/2*log(c*x - 1)/c^2 - integrate(sqrt(c*x + 1)*sqrt(-c*x + 1)/(c^3*x^2 - c), x)

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x}{1 - c^2 x^2} dx = \int -\frac{x \left(\sqrt{\frac{1}{cx} + 1} \sqrt{\frac{1}{cx} - 1} + \frac{1}{cx} \right)}{c^2 x^2 - 1} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))*x/(-c^2*x^2+1),x, algorithm="giac")

[Out] integrate(-x*(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(c^2*x^2 - 1), x)

Mupad [B] (verification not implemented)

Time = 6.10 (sec) , antiderivative size = 84, normalized size of antiderivative = 2.27

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)} x}{1 - c^2 x^2} dx = \frac{\operatorname{atanh}(cx)}{c^2} + \frac{\left(\ln \left(\frac{\left(\sqrt{\frac{1}{cx} - 1} - i \right)^2}{\left(\sqrt{\frac{1}{cx} + 1} - 1 \right)^2} + 1 \right) - \ln \left(\frac{\sqrt{\frac{1}{cx} - 1} - i}{\sqrt{\frac{1}{cx} + 1} - 1} \right) \right) i}{c^2}$$

[In] int(-x*((1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2) + 1/(c*x)))/(c^2*x^2 - 1), x)

[Out] ((log(((1/(c*x) - 1)^(1/2) - 1i)^2/((1/(c*x) + 1)^(1/2) - 1)^2 + 1) - log((1/(c*x) - 1)^(1/2) - 1i)/((1/(c*x) + 1)^(1/2) - 1)))*1i)/c^2 + atanh(c*x)/c^2

3.93 $\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{1-c^2x^2} dx$

Optimal result	568
Rubi [A] (verified)	568
Mathematica [A] (verified)	570
Maple [A] (verified)	570
Fricas [B] (verification not implemented)	571
Sympy [F]	571
Maxima [F]	571
Giac [F]	572
Mupad [B] (verification not implemented)	572

Optimal result

Integrand size = 19, antiderivative size = 71

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{1-c^2x^2} dx = -\frac{\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\operatorname{arctanh}(\sqrt{1-cx}\sqrt{1+cx})}{c} + \frac{\log(x)}{c} - \frac{\log(1-c^2x^2)}{2c}$$

[Out] $\ln(x)/c-1/2*\ln(-c^2*x^2+1)/c-\operatorname{arctanh}((-c*x+1)^{(1/2)}*(c*x+1)^{(1/2))}*(1/(c*x+1))^{(1/2)}*(c*x+1)^{(1/2)}/c$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {6474, 1972, 94, 214, 272, 36, 29, 31}

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{1-c^2x^2} dx = -\frac{\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\operatorname{arctanh}(\sqrt{1-cx}\sqrt{cx+1})}{c} - \frac{\log(1-c^2x^2)}{2c} + \frac{\log(x)}{c}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcSech}[c*x]}/(1-c^2*x^2), x]$

[Out] $-((\operatorname{Sqrt}[(1+c*x)^{-1}]*\operatorname{Sqrt}[1+c*x]*\operatorname{ArcTanh}[\operatorname{Sqrt}[1-c*x]*\operatorname{Sqrt}[1+c*x]])/c) + \operatorname{Log}[x]/c - \operatorname{Log}[1-c^2*x^2]/(2*c)$

Rule 29

$\operatorname{Int}[(x_)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Log}[x], x]$

Rule 31

Int[((a_) + (b_)*(x_))⁽⁻¹⁾, x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 36

Int[1/(((a_) + (b_)*(x_))*((c_) + (d_)*(x_))), x_Symbol] := Dist[b/(b*c - a*d), Int[1/(a + b*x), x], x] - Dist[d/(b*c - a*d), Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]

Rule 94

Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]*((e_) + (f_)*(x_))), x_Symbol] := Dist[b*f, Subst[Int[1/(d*(b*e - a*f)² + b*f²*x²), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]

Rule 214

Int[((a_) + (b_)*(x_)²)⁽⁻¹⁾, x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 272

Int[(x_)^{(m_)*((a_) + (b_)*(x_)^{(n_))^(p_)}, x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, xⁿ], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]}

Rule 1972

Int[(u_)*((c_)*((a_) + (b_)*(x_)^{(n_))^(q_))^(p_)}, x_Symbol] := Dist[Simp[(c*(a + b*xⁿ)^q)^p/(a + b*xⁿ)^(p*q)], Int[u*(a + b*xⁿ)^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && GeQ[a, 0]

Rule 6474

Int[E^{ArcSech[(c_)*(x_)]}/((a_) + (b_)*(x_)²), x_Symbol] := Dist[1/(a*c), Int[Sqrt[1/(1 + c*x)]/(x*Sqrt[1 - c*x]), x], x] + Dist[1/c, Int[1/(x*(a + b*x²)), x], x] /; FreeQ[{a, b, c}, x] && EqQ[b + a*c², 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \frac{\sqrt{\frac{1}{1+cx}}}{x\sqrt{1-cx}} dx}{c} + \frac{\int \frac{1}{x(1-c^2x^2)} dx}{c} \\ &= \frac{\text{Subst}\left(\int \frac{1}{x(1-c^2x)} dx, x, x^2\right)}{2c} + \frac{\left(\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{1}{x\sqrt{1-cx}\sqrt{1+cx}} dx}{c} \end{aligned}$$

$$\begin{aligned}
&= \frac{\text{Subst}\left(\int \frac{1}{x} dx, x, x^2\right)}{2c} + \frac{1}{2}c\text{Subst}\left(\int \frac{1}{1-c^2x} dx, x, x^2\right) \\
&\quad - \left(\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right)\text{Subst}\left(\int \frac{1}{c-cx^2} dx, x, \sqrt{1-cx}\sqrt{1+cx}\right) \\
&= -\frac{\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\text{arctanh}\left(\sqrt{1-cx}\sqrt{1+cx}\right)}{c} + \frac{\log(x)}{c} - \frac{\log(1-c^2x^2)}{2c}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.03

$$\int \frac{e^{\text{sech}^{-1}(cx)}}{1-c^2x^2} dx = \frac{2\log(x)}{c} - \frac{\log(1-c^2x^2)}{2c} - \frac{\log\left(1 + \sqrt{\frac{1-cx}{1+cx}} + cx\sqrt{\frac{1-cx}{1+cx}}\right)}{c}$$

[In] Integrate[E^ArcSech[c*x]/(1 - c^2*x^2),x]

[Out] (2*Log[x])/c - Log[1 - c^2*x^2]/(2*c) - Log[1 + Sqrt[(1 - c*x)/(1 + c*x)] + c*x*Sqrt[(1 - c*x)/(1 + c*x)]]/c

Maple [A] (verified)

Time = 0.70 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.15

method	result	size
default	$-\frac{\sqrt{-\frac{cx-1}{cx}}x\sqrt{\frac{cx+1}{cx}}\text{arctanh}\left(\frac{1}{\sqrt{-c^2x^2+1}}\right)}{\sqrt{-c^2x^2+1}} + \frac{-\frac{\ln(cx+1)}{2}+\ln(x)-\frac{\ln(cx-1)}{2}}{c}$	82

[In] int((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(-c^2*x^2+1),x,method=_RETURNV
ERBOSE)

[Out] -((-c*x-1)/c/x)^(1/2)*x*((c*x+1)/c/x)^(1/2)*arctanh(1/(-c^2*x^2+1)^(1/2))/(
-c^2*x^2+1)^(1/2)+1/c*(-1/2*ln(c*x+1)+ln(x)-1/2*ln(c*x-1))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 92 vs. 2(45) = 90.

Time = 0.24 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.30

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{1 - c^2x^2} dx = \frac{\log(c^2x^2 - 1) + \log\left(cx\sqrt{\frac{cx+1}{cx}}\sqrt{-\frac{cx-1}{cx}} + 1\right) - \log\left(cx\sqrt{\frac{cx+1}{cx}}\sqrt{-\frac{cx-1}{cx}} - 1\right) - 2\log(x)}{2c}$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(-c^2*x^2+1),x, algorithm m="fricas")

[Out] -1/2*(log(c^2*x^2 - 1) + log(c*x*sqrt((c*x + 1)/(c*x))*sqrt(-(c*x - 1)/(c*x)) + 1) - log(c*x*sqrt((c*x + 1)/(c*x))*sqrt(-(c*x - 1)/(c*x)) - 1) - 2*log(x))/c

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{1 - c^2x^2} dx = -\frac{\int \frac{cx\sqrt{-1+\frac{1}{cx}}\sqrt{1+\frac{1}{cx}}}{c^2x^3-x} dx + \int \frac{1}{c^2x^3-x} dx}{c}$$

[In] integrate((1/c/x+(-1+1/c/x)**(1/2)*(1+1/c/x)**(1/2))/(-c**2*x**2+1),x)

[Out] -(Integral(c*x*sqrt(-1 + 1/(c*x))*sqrt(1 + 1/(c*x))/(c**2*x**3 - x), x) + Integral(1/(c**2*x**3 - x), x))/c

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{1 - c^2x^2} dx = \int -\frac{\sqrt{\frac{1}{cx} + 1}\sqrt{\frac{1}{cx} - 1 + \frac{1}{cx}}}{c^2x^2 - 1} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(-c^2*x^2+1),x, algorithm m="maxima")

[Out] integrate(1/x, x)/c - 1/2*log(c*x + 1)/c - 1/2*log(c*x - 1)/c - integrate(sqrt(c*x + 1)*sqrt(-c*x + 1)/(c^3*x^3 - c*x), x)

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{1 - c^2x^2} dx = \int -\frac{\sqrt{\frac{1}{cx} + 1}\sqrt{\frac{1}{cx} - 1} + \frac{1}{cx}}{c^2x^2 - 1} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/(-c^2*x^2+1),x, algorithm m="giac")

[Out] integrate(-(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/(c^2*x^2 - 1), x)

Mupad [B] (verification not implemented)

Time = 5.95 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.83

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{1 - c^2x^2} dx = \frac{\ln(x)}{c} - \frac{4 \operatorname{atanh}\left(\frac{\sqrt{\frac{1}{cx} - 1 - i}}{\sqrt{\frac{1}{cx} + 1 - 1}}\right)}{c} - \frac{\ln(3c^2x^2 - 3)}{2c}$$

[In] int(-((1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2) + 1/(c*x))/(c^2*x^2 - 1),x)

[Out] log(x)/c - (4*atanh(((1/(c*x) - 1)^(1/2) - 1i)/((1/(c*x) + 1)^(1/2) - 1)))/c - log(3*c^2*x^2 - 3)/(2*c)

3.94 $\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x(1-c^2x^2)} dx$

Optimal result	573
Rubi [A] (verified)	573
Mathematica [A] (verified)	575
Maple [C] (verified)	575
Fricas [A] (verification not implemented)	575
Sympy [F]	576
Maxima [F]	576
Giac [F]	576
Mupad [B] (verification not implemented)	577

Optimal result

Integrand size = 22, antiderivative size = 42

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x(1-c^2x^2)} dx = -\frac{1}{cx} - \frac{\sqrt{1-cx}}{cx\sqrt{\frac{1}{1+cx}}} + \operatorname{arctanh}(cx)$$

[Out] $-1/c/x + \operatorname{arctanh}(c*x) - (-c*x+1)^{(1/2)}/c/x/(1/(c*x+1))^{(1/2)}$

Rubi [A] (verified)

Time = 0.11 (sec), antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.227$, Rules used = {6476, 1972, 97, 331, 212}

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x(1-c^2x^2)} dx = \operatorname{arctanh}(cx) - \frac{\sqrt{1-cx}}{cx\sqrt{\frac{1}{cx+1}}} - \frac{1}{cx}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcSech}[c*x]}/(x*(1-c^2*x^2)), x]$

[Out] $-(1/(c*x)) - \operatorname{Sqrt}[1-c*x]/(c*x*\operatorname{Sqrt}[(1+c*x)^{-1}]) + \operatorname{ArcTanh}[c*x]$

Rule 97

$\operatorname{Int}[(a_.) + (b_.)*(x_.)^{(m_.)}*((c_.) + (d_.)*(x_.)^{(n_.)}*((e_.) + (f_.)*(x_.)^{(p_.)}), x_Symbol] \rightarrow \operatorname{Simp}[b*(a + b*x)^{(m+1)}*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1})/((m+1)*(b*c - a*d)*(b*e - a*f))), x] /;$ $\operatorname{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x$ && $\operatorname{EqQ}[\operatorname{Simplify}[m + n + p + 3], 0]$ && $\operatorname{EqQ}[a*d*f*(m+1) + b*c*f*(n+1) + b*d*e*(p+1), 0]$ && $\operatorname{NeQ}[m, -1]$

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 331

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x
)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Dist[b*((m + n*(p + 1)
+ 1)/(a*c^n*(m + 1)), Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a,
b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p,
x]
```

Rule 1972

```
Int[(u_.)*((c_.)*((a_.) + (b_.)*(x_)^(n_.))^(q_.))^(p_), x_Symbol] := Dist[S
imp[(c*(a + b*x^n)^q)^p/(a + b*x^n)^(p*q)], Int[u*(a + b*x^n)^(p*q), x], x]
/; FreeQ[{a, b, c, n, p, q}, x] && GeQ[a, 0]
```

Rule 6476

```
Int[(E^ArcSech[(c_.)*(x_)])*((d_.)*(x_)^(m_.))/((a_) + (b_.)*(x_)^2), x_Sym
bol] := Dist[d/(a*c), Int[(d*x)^(m - 1)*(Sqrt[1/(1 + c*x)]/Sqrt[1 - c*x]),
x], x] + Dist[d/c, Int[(d*x)^(m - 1)/(a + b*x^2), x], x] /; FreeQ[{a, b, c,
d, m}, x] && EqQ[b + a*c^2, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\int \frac{\sqrt{\frac{1}{1+cx}}}{x^2\sqrt{1-cx}} dx}{c} + \frac{\int \frac{1}{x^2(1-c^2x^2)} dx}{c} \\
&= -\frac{1}{cx} + c \int \frac{1}{1-c^2x^2} dx + \frac{\left(\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{1}{x^2\sqrt{1-cx}\sqrt{1+cx}} dx}{c} \\
&= -\frac{1}{cx} - \frac{\sqrt{1-cx}}{cx\sqrt{\frac{1}{1+cx}}} + \operatorname{arctanh}(cx)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.40

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x(1-c^2x^2)} dx = -\frac{1}{cx} - \left(1 + \frac{1}{cx}\right) \sqrt{\frac{1-cx}{1+cx}} - \frac{1}{2} \log(1-cx) + \frac{1}{2} \log(1+cx)$$

[In] Integrate[E^ArcSech[c*x]/(x*(1 - c^2*x^2)),x]

[Out] -(1/(c*x)) - (1 + 1/(c*x))*Sqrt[(1 - c*x)/(1 + c*x)] - Log[1 - c*x]/2 + Log[1 + c*x]/2

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.72 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.55

method	result	size
default	$-\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \operatorname{csgn}(c)^2 + \frac{\frac{c \ln(cx+1)}{2} - \frac{c \ln(cx-1)}{2} - \frac{1}{x}}{c}$	65

[In] int((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/x/(-c^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] -((-c*x-1)/c/x)^(1/2)*((c*x+1)/c/x)^(1/2)*csgn(c)^2+1/c*(1/2*c*ln(c*x+1)-1/2*c*ln(c*x-1)-1/x)

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.48

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x(1-c^2x^2)} dx = -\frac{2cx \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} - cx \log(cx+1) + cx \log(cx-1) + 2}{2cx}$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/x/(-c^2*x^2+1),x, algorithm="fricas")

[Out] -1/2*(2*c*x*sqrt((c*x + 1)/(c*x))*sqrt(-(c*x - 1)/(c*x)) - c*x*log(c*x + 1) + c*x*log(c*x - 1) + 2)/(c*x)

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x(1-c^2x^2)} dx = -\int \frac{cx\sqrt{-1+\frac{1}{cx}}\sqrt{1+\frac{1}{cx}}}{c^2x^4-x^2} dx + \int \frac{1}{c^2x^4-x^2} dx$$

[In] integrate((1/c/x+(-1+1/c/x)**(1/2)*(1+1/c/x)**(1/2))/x/(-c**2*x**2+1), x)

[Out] -(Integral(c*x*sqrt(-1 + 1/(c*x))*sqrt(1 + 1/(c*x))/(c**2*x**4 - x**2), x) + Integral(1/(c**2*x**4 - x**2), x))/c

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x(1-c^2x^2)} dx = \int -\frac{\sqrt{\frac{1}{cx}+1}\sqrt{\frac{1}{cx}-1+\frac{1}{cx}}}{(c^2x^2-1)x} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/x/(-c^2*x^2+1), x, algorithm="maxima")

[Out] integrate(x^(-2), x)/c - integrate(sqrt(c*x + 1)*sqrt(-c*x + 1)/(c^3*x^4 - c*x^2), x) + 1/2*log(c*x + 1) - 1/2*log(c*x - 1)

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x(1-c^2x^2)} dx = \int -\frac{\sqrt{\frac{1}{cx}+1}\sqrt{\frac{1}{cx}-1+\frac{1}{cx}}}{(c^2x^2-1)x} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/x/(-c^2*x^2+1), x, algorithm="giac")

[Out] integrate(-(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/((c^2*x^2 - 1)*x), x)

Mupad [B] (verification not implemented)

Time = 5.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x(1-c^2x^2)} dx = \operatorname{atanh}(cx) - \sqrt{\frac{1}{cx} - 1} \sqrt{\frac{1}{cx} + 1} - \frac{1}{cx}$$

[In] int(-((1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2) + 1/(c*x))/(x*(c^2*x^2 - 1)), x)

[Out] atanh(c*x) - (1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2) - 1/(c*x)

3.95 $\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^2(1-c^2x^2)} dx$

Optimal result	578
Rubi [A] (verified)	578
Mathematica [A] (verified)	581
Maple [A] (verified)	581
Fricas [B] (verification not implemented)	581
Sympy [F]	582
Maxima [F]	582
Giac [F]	582
Mupad [B] (verification not implemented)	583

Optimal result

Integrand size = 22, antiderivative size = 108

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^2(1-c^2x^2)} dx = -\frac{1}{2cx^2} - \frac{\sqrt{1-cx}}{2cx^2\sqrt{\frac{1}{1+cx}}} - \frac{1}{2}c\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\operatorname{arctanh}\left(\sqrt{1-cx}\sqrt{1+cx}\right) + c\log(x) - \frac{1}{2}c\log(1-c^2x^2)$$

[Out] $-1/2/c/x^2+c*\ln(x)-1/2*c*\ln(-c^2*x^2+1)-1/2*(-c*x+1)^{(1/2)}/c/x^2/(1/(c*x+1))^{(1/2)}-1/2*c*\operatorname{arctanh}((-c*x+1)^{(1/2)}*(c*x+1)^{(1/2)})*(1/(c*x+1))^{(1/2)}*(c*x+1)^{(1/2)}$

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {6476, 1972, 105, 12, 94, 214, 272, 46}

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^2(1-c^2x^2)} dx = -\frac{1}{2}c\sqrt{\frac{1}{cx+1}}\sqrt{cx+1}\operatorname{arctanh}\left(\sqrt{1-cx}\sqrt{cx+1}\right) - \frac{1}{2}c\log(1-c^2x^2) - \frac{\sqrt{1-cx}}{2cx^2\sqrt{\frac{1}{cx+1}}} - \frac{1}{2cx^2} + c\log(x)$$

[In] $\operatorname{Int}\left[\frac{E^{\operatorname{ArcSech}[c*x]}}{x^2*(1-c^2*x^2)}, x\right]$

[Out] $-1/2*1/(c*x^2) - \operatorname{Sqrt}[1-c*x]/(2*c*x^2*\operatorname{Sqrt}[(1+c*x)^{-1}]) - (c*\operatorname{Sqrt}[(1+c*x)^{-1}]*\operatorname{Sqrt}[1+c*x]*\operatorname{ArcTanh}[\operatorname{Sqrt}[1-c*x]*\operatorname{Sqrt}[1+c*x]])/2 + c*\operatorname{Log}[x] - (c*\operatorname{Log}[1-c^2*x^2])/2$

Rule 12

```
Int[(a_)*(u_), x_Symbol] :=> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 46

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] :=> Int[Ex
pandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] &&
NeQ[b*c - a*d, 0] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m +
n + 2, 0])
```

Rule 94

```
Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]*((e_) + (f_)*(x_
))), x_Symbol] :=> Dist[b*f, Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x],
x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[
2*b*d*e - f*(b*c + a*d), 0]
```

Rule 105

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_
))^(p_), x_Symbol] :=> Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x
)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a
*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*
(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x,
x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (Integer
Q[n] || IntegerQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])
```

Rule 214

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :=> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 272

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :=> Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 1972

```
Int[(u_)*((c_)*((a_) + (b_)*(x_)^(n_))^(q_))^(p_), x_Symbol] :=> Dist[S
imp[(c*(a + b*x^n)^q)^p/(a + b*x^n)^(p*q)], Int[u*(a + b*x^n)^(p*q), x], x]
/; FreeQ[{a, b, c, n, p, q}, x] && GeQ[a, 0]
```

Rule 6476

Int[(E^ArcSech[(c_.)*(x_)]*((d_.)*(x_))^(m_.))/((a_) + (b_.)*(x_)^2), x_Symbol] :> Dist[d/(a*c), Int[(d*x)^(m - 1)*(Sqrt[1/(1 + c*x)]/Sqrt[1 - c*x]), x], x] + Dist[d/c, Int[(d*x)^(m - 1)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + a*c^2, 0]

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\int \frac{\sqrt{\frac{1}{1+cx}}}{x^3\sqrt{1-cx}} dx}{c} + \frac{\int \frac{1}{x^3(1-c^2x^2)} dx}{c} \\
&= \frac{\text{Subst}\left(\int \frac{1}{x^2(1-c^2x)} dx, x, x^2\right)}{2c} + \frac{\left(\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{1}{x^3\sqrt{1-cx}\sqrt{1+cx}} dx}{c} \\
&= -\frac{\sqrt{1-cx}}{2cx^2\sqrt{\frac{1}{1+cx}}} + \frac{\text{Subst}\left(\int \left(\frac{1}{x^2} + \frac{c^2}{x} - \frac{c^4}{-1+c^2x}\right) dx, x, x^2\right)}{2c} \\
&\quad + \frac{\left(\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{c^2}{x\sqrt{1-cx}\sqrt{1+cx}} dx}{2c} \\
&= -\frac{1}{2cx^2} - \frac{\sqrt{1-cx}}{2cx^2\sqrt{\frac{1}{1+cx}}} + c \log(x) - \frac{1}{2}c \log(1 - c^2x^2) \\
&\quad + \frac{1}{2} \left(c\sqrt{\frac{1}{1+cx}}\sqrt{1+cx} \right) \int \frac{1}{x\sqrt{1-cx}\sqrt{1+cx}} dx \\
&= -\frac{1}{2cx^2} - \frac{\sqrt{1-cx}}{2cx^2\sqrt{\frac{1}{1+cx}}} + c \log(x) - \frac{1}{2}c \log(1 - c^2x^2) \\
&\quad - \frac{1}{2} \left(c^2\sqrt{\frac{1}{1+cx}}\sqrt{1+cx} \right) \text{Subst}\left(\int \frac{1}{c - cx^2} dx, x, \sqrt{1-cx}\sqrt{1+cx}\right) \\
&= -\frac{1}{2cx^2} - \frac{\sqrt{1-cx}}{2cx^2\sqrt{\frac{1}{1+cx}}} - \frac{1}{2}c\sqrt{\frac{1}{1+cx}}\sqrt{1+cx} \arctanh\left(\sqrt{1-cx}\sqrt{1+cx}\right) \\
&\quad + c \log(x) - \frac{1}{2}c \log(1 - c^2x^2)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.00

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^2(1-c^2x^2)} dx = \frac{1}{2} \left(-\frac{1}{cx^2} - \frac{\sqrt{\frac{1-cx}{1+cx}}(1+cx)}{cx^2} + 3c \log(x) - c \log(1-c^2x^2) - c \log \left(1 + \sqrt{\frac{1-cx}{1+cx}} + cx \sqrt{\frac{1-cx}{1+cx}} \right) \right)$$

[In] Integrate[E^ArcSech[c*x]/(x^2*(1 - c^2*x^2)), x]

[Out] $(-(1/(c*x^2)) - (\operatorname{Sqrt}[(1 - c*x)/(1 + c*x)]*(1 + c*x))/(c*x^2) + 3*c*\operatorname{Log}[x] - c*\operatorname{Log}[1 - c^2*x^2] - c*\operatorname{Log}[1 + \operatorname{Sqrt}[(1 - c*x)/(1 + c*x)] + c*x*\operatorname{Sqrt}[(1 - c*x)/(1 + c*x])])/2$

Maple [A] (verified)

Time = 0.73 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.10

method	result	size
default	$-\frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \left(\operatorname{arctanh} \left(\frac{1}{\sqrt{-c^2x^2+1}} \right) c^2x^2 + \sqrt{-c^2x^2+1} \right)}{2x\sqrt{-c^2x^2+1}} + \frac{-\frac{c^2 \ln(cx+1)}{2} - \frac{1}{2x^2} + c^2 \ln(x) - \frac{c^2 \ln(cx-1)}{2}}{c}$	119

[In] int((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/x^2/(-c^2*x^2+1), x, method=_RETURNVERBOSE)

[Out] $-1/2*(-(c*x-1)/c/x)^(1/2)/x*((c*x+1)/c/x)^(1/2)*(\operatorname{arctanh}(1/(-c^2*x^2+1)^(1/2))*c^2*x^2+(-c^2*x^2+1)^(1/2))/(-c^2*x^2+1)^(1/2)+1/c*(-1/2*c^2*\ln(c*x+1)-1/2/x^2+c^2*\ln(x)-1/2*c^2*\ln(c*x-1))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 156 vs. 2(70) = 140.

Time = 0.25 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.44

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^2(1-c^2x^2)} dx = \frac{2c^2x^2 \log(c^2x^2 - 1) + c^2x^2 \log \left(cx \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} + 1 \right) - c^2x^2 \log \left(cx \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} - 1 \right) - 4c^2x^2 \log \left(1 + \sqrt{\frac{1-cx}{1+cx}} + cx \sqrt{\frac{1-cx}{1+cx}} \right)}{4cx^2}$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/x^2/(-c^2*x^2+1), x, algorithm="fricas")

[Out] $-1/4*(2*c^2*x^2*\log(c^2*x^2 - 1) + c^2*x^2*\log(c*x*\sqrt{(c*x + 1)/(c*x)})*\sqrt{-1/(c*x - 1)/(c*x)} + 1) - c^2*x^2*\log(c*x*\sqrt{(c*x + 1)/(c*x)})*\sqrt{-1/(c*x - 1)/(c*x)} - 1) - 4*c^2*x^2*\log(x) + 2*c*x*\sqrt{(c*x + 1)/(c*x)}*\sqrt{-1/(c*x - 1)/(c*x)} + 2)/(c*x^2)$

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^2(1-c^2x^2)} dx = -\frac{\int \frac{cx\sqrt{-1+\frac{1}{cx}}\sqrt{1+\frac{1}{cx}}}{c^2x^5-x^3} dx + \int \frac{1}{c^2x^5-x^3} dx}{c}$$

[In] `integrate((1/c/x+(-1+1/c/x)**(1/2)*(1+1/c/x)**(1/2))/x**2/(-c**2*x**2+1),x)`

[Out] $-(\operatorname{Integral}(c*x*\sqrt{-1 + 1/(c*x)}*\sqrt{1 + 1/(c*x)})/(c**2*x**5 - x**3), x) + \operatorname{Integral}(1/(c**2*x**5 - x**3), x))/c$

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^2(1-c^2x^2)} dx = \int -\frac{\sqrt{\frac{1}{cx} + 1}\sqrt{\frac{1}{cx} - 1} + \frac{1}{cx}}{(c^2x^2 - 1)x^2} dx$$

[In] `integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/x^2/(-c^2*x^2+1),x, algorithm="maxima")`

[Out] $c*\operatorname{integrate}(1/x, x) - 1/2*c*\log(c*x + 1) - 1/2*c*\log(c*x - 1) + \operatorname{integrate}(x^{-3}, x)/c - \operatorname{integrate}(\sqrt{c*x + 1}*\sqrt{-c*x + 1}/(c^3*x^5 - c*x^3), x)$

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^2(1-c^2x^2)} dx = \int -\frac{\sqrt{\frac{1}{cx} + 1}\sqrt{\frac{1}{cx} - 1} + \frac{1}{cx}}{(c^2x^2 - 1)x^2} dx$$

[In] `integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/x^2/(-c^2*x^2+1),x, algorithm="giac")`

[Out] $\operatorname{integrate}(-(\sqrt{1/(c*x) + 1}*\sqrt{1/(c*x) - 1} + 1/(c*x)))/((c^2*x^2 - 1)*x^2), x)$

Mupad [B] (verification not implemented)

Time = 17.89 (sec) , antiderivative size = 331, normalized size of antiderivative = 3.06

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^2(1-c^2x^2)} dx = c \ln(x) + \frac{2c\left(\sqrt{\frac{1}{cx}-1-i}\right)}{\sqrt{\frac{1}{cx}+1-1}} + \frac{14c\left(\sqrt{\frac{1}{cx}-1-i}\right)^3}{\left(\sqrt{\frac{1}{cx}+1-1}\right)^3} + \frac{14c\left(\sqrt{\frac{1}{cx}-1-i}\right)^5}{\left(\sqrt{\frac{1}{cx}+1-1}\right)^5} + \frac{2c\left(\sqrt{\frac{1}{cx}-1-i}\right)^7}{\left(\sqrt{\frac{1}{cx}+1-1}\right)^7} + \frac{1 + \frac{6\left(\sqrt{\frac{1}{cx}-1-i}\right)^4}{\left(\sqrt{\frac{1}{cx}+1-1}\right)^4} - \frac{4\left(\sqrt{\frac{1}{cx}-1-i}\right)^6}{\left(\sqrt{\frac{1}{cx}+1-1}\right)^6} + \frac{\left(\sqrt{\frac{1}{cx}-1-i}\right)^8}{\left(\sqrt{\frac{1}{cx}+1-1}\right)^8} - \frac{4\left(\sqrt{\frac{1}{cx}-1-i}\right)^2}{\left(\sqrt{\frac{1}{cx}+1-1}\right)^2}}{1} - \frac{c \ln(c^2 x^2 - 1)}{2} - 2c \operatorname{atanh}\left(\frac{\sqrt{\frac{1}{cx}-1-i}}{\sqrt{\frac{1}{cx}+1-1}}\right) - \frac{1}{2cx^2}$$

[In] int(-((1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2) + 1/(c*x)))/(x^2*(c^2*x^2 - 1)),x)

[Out] ((2*c*((1/(c*x) - 1)^(1/2) - 1i))/((1/(c*x) + 1)^(1/2) - 1) + (14*c*((1/(c*x) - 1)^(1/2) - 1i)^3)/((1/(c*x) + 1)^(1/2) - 1)^3 + (14*c*((1/(c*x) - 1)^(1/2) - 1i)^5)/((1/(c*x) + 1)^(1/2) - 1)^5 + (2*c*((1/(c*x) - 1)^(1/2) - 1i)^7)/((1/(c*x) + 1)^(1/2) - 1)^7)/((6*((1/(c*x) - 1)^(1/2) - 1i)^4)/((1/(c*x) + 1)^(1/2) - 1)^4 - (4*((1/(c*x) - 1)^(1/2) - 1i)^2)/((1/(c*x) + 1)^(1/2) - 1)^2 - (4*((1/(c*x) - 1)^(1/2) - 1i)^6)/((1/(c*x) + 1)^(1/2) - 1)^6 + ((1/(c*x) - 1)^(1/2) - 1i)^8/((1/(c*x) + 1)^(1/2) - 1)^8 + 1) - 2*c*atanh((1/(c*x) - 1)^(1/2) - 1i)/((1/(c*x) + 1)^(1/2) - 1)) - (c*log(c^2*x^2 - 1))/2 + c*log(x) - 1/(2*c*x^2)

3.96 $\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^3(1-c^2x^2)} dx$

Optimal result	584
Rubi [A] (verified)	584
Mathematica [A] (verified)	586
Maple [C] (verified)	587
Fricas [A] (verification not implemented)	587
Sympy [F]	587
Maxima [F]	588
Giac [F]	588
Mupad [B] (verification not implemented)	588

Optimal result

Integrand size = 22, antiderivative size = 85

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^3(1-c^2x^2)} dx = -\frac{1}{3cx^3} - \frac{c}{x} - \frac{\sqrt{1-cx}}{3cx^3\sqrt{\frac{1}{1+cx}}} - \frac{2c\sqrt{1-cx}}{3x\sqrt{\frac{1}{1+cx}}} + c^2 \operatorname{arctanh}(cx)$$

[Out] $-1/3/c/x^3 - c/x + c^2 \operatorname{arctanh}(c*x) - 1/3*(-c*x+1)^{(1/2)}/c/x^3/(1/(c*x+1))^{(1/2)} - 2/3*c*(-c*x+1)^{(1/2)}/x/(1/(c*x+1))^{(1/2)}$

Rubi [A] (verified)

Time = 0.12 (sec), antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.318$, Rules used = {6476, 1972, 105, 12, 97, 331, 212}

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^3(1-c^2x^2)} dx = c^2 \operatorname{arctanh}(cx) - \frac{\sqrt{1-cx}}{3cx^3\sqrt{\frac{1}{cx+1}}} - \frac{1}{3cx^3} - \frac{2c\sqrt{1-cx}}{3x\sqrt{\frac{1}{cx+1}}} - \frac{c}{x}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcSech}[c*x]}/(x^3*(1-c^2*x^2)), x]$

[Out] $-1/3*1/(c*x^3) - c/x - \operatorname{Sqrt}[1-c*x]/(3*c*x^3*\operatorname{Sqrt}[(1+c*x)^{-1}]) - (2*c*\operatorname{Sqrt}[1-c*x])/(3*x*\operatorname{Sqrt}[(1+c*x)^{-1}]) + c^2*\operatorname{ArcTanh}[c*x]$

Rule 12

$\operatorname{Int}[(a_*)(u_), x_Symbol] := \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /;$ FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 97


```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && EqQ[a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1), 0] && NeQ[m, -1]
```

Rule 105

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 331

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Dist[b*(m + n*(p + 1) + 1)/(a*c^n*(m + 1)), Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 1972

```
Int[(u_.)*((c_.)*((a_.) + (b_.)*(x_)^(n_.))^(q_.))^(p_), x_Symbol] := Dist[Simp[(c*(a + b*x^n)^q)^p/(a + b*x^n)^(p*q)], Int[u*(a + b*x^n)^(p*q), x], x] /; FreeQ[{a, b, c, n, p, q}, x] && GeQ[a, 0]
```

Rule 6476

```
Int[(E^ArcSech[(c_.)*(x_)])*((d_.)*(x_))^(m_.))/((a_) + (b_.)*(x_)^2), x_Symbol] := Dist[d/(a*c), Int[(d*x)^(m - 1)*(Sqrt[1/(1 + c*x)]/Sqrt[1 - c*x]), x], x] + Dist[d/c, Int[(d*x)^(m - 1)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + a*c^2, 0]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int \frac{\sqrt{\frac{1}{1+cx}}}{x^4\sqrt{1-cx}} dx}{c} + \frac{\int \frac{1}{x^4(1-c^2x^2)} dx}{c} \\
 &= -\frac{1}{3cx^3} + c \int \frac{1}{x^2(1-c^2x^2)} dx + \frac{\left(\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{1}{x^4\sqrt{1-cx}\sqrt{1+cx}} dx}{c} \\
 &= -\frac{1}{3cx^3} - \frac{c}{x} - \frac{\sqrt{1-cx}}{3cx^3\sqrt{\frac{1}{1+cx}}} + c^3 \int \frac{1}{1-c^2x^2} dx - \frac{\left(\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int -\frac{2c^2}{x^2\sqrt{1-cx}\sqrt{1+cx}} dx}{3c} \\
 &= -\frac{1}{3cx^3} - \frac{c}{x} - \frac{\sqrt{1-cx}}{3cx^3\sqrt{\frac{1}{1+cx}}} + c^2 \operatorname{arctanh}(cx) \\
 &\quad + \frac{1}{3} \left(2c\sqrt{\frac{1}{1+cx}}\sqrt{1+cx}\right) \int \frac{1}{x^2\sqrt{1-cx}\sqrt{1+cx}} dx \\
 &= -\frac{1}{3cx^3} - \frac{c}{x} - \frac{\sqrt{1-cx}}{3cx^3\sqrt{\frac{1}{1+cx}}} - \frac{2c\sqrt{1-cx}}{3x\sqrt{\frac{1}{1+cx}}} + c^2 \operatorname{arctanh}(cx)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.06

$$\begin{aligned}
 &\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^3(1-c^2x^2)} dx \\
 &= -\frac{2 + 6c^2x^2 + 2\sqrt{\frac{1-cx}{1+cx}}(1+cx+2c^2x^2+2c^3x^3) + 3c^3x^3 \log(1-cx) - 3c^3x^3 \log(1+cx)}{6cx^3}
 \end{aligned}$$

[In] Integrate[E^ArcSech[c*x]/(x^3*(1 - c^2*x^2)),x]

[Out] -1/6*(2 + 6*c^2*x^2 + 2*Sqrt[(1 - c*x)/(1 + c*x)]*(1 + c*x + 2*c^2*x^2 + 2*c^3*x^3) + 3*c^3*x^3*Log[1 - c*x] - 3*c^3*x^3*Log[1 + c*x])/(c*x^3)

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.73 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.06

method	result	size
default	$-\frac{\sqrt{-\frac{cx-1}{cx}} \sqrt{\frac{cx+1}{cx}} \operatorname{csgn}(c)^2 (2c^2x^2+1)}{3x^2} + \frac{\frac{c^3 \ln(cx+1)}{2} - \frac{1}{3x^3} - \frac{c^2}{x} - \frac{c^3 \ln(cx-1)}{2}}{c}$	90

[In] int((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/x^3/(-c^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] -1/3*(-(c*x-1)/c/x)^(1/2)/x^2*((c*x+1)/c/x)^(1/2)*csgn(c)^2*(2*c^2*x^2+1)+1/c*(1/2*c^3*ln(c*x+1)-1/3/x^3-c^2/x-1/2*c^3*ln(c*x-1))

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.05

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^3(1-c^2x^2)} dx = \frac{3c^3x^3 \log(cx+1) - 3c^3x^3 \log(cx-1) - 6c^2x^2 - 2(2c^3x^3 + cx) \sqrt{\frac{cx+1}{cx}} \sqrt{-\frac{cx-1}{cx}} - 2}{6cx^3}$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/x^3/(-c^2*x^2+1),x,algorithm="fricas")

[Out] 1/6*(3*c^3*x^3*log(c*x + 1) - 3*c^3*x^3*log(c*x - 1) - 6*c^2*x^2 - 2*(2*c^3*x^3 + c*x)*sqrt((c*x + 1)/(c*x))*sqrt(-(c*x - 1)/(c*x)) - 2)/(c*x^3)

Sympy [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^3(1-c^2x^2)} dx = -\int \frac{cx \sqrt{-1+\frac{1}{cx}} \sqrt{1+\frac{1}{cx}}}{c^2x^6-x^4} dx + \int \frac{1}{c^2x^6-x^4} dx$$

[In] integrate((1/c/x+(-1+1/c/x)**(1/2)*(1+1/c/x)**(1/2))/x**3/(-c**2*x**2+1),x)

[Out] -(Integral(c*x*sqrt(-1 + 1/(c*x))*sqrt(1 + 1/(c*x))/(c**2*x**6 - x**4), x) + Integral(1/(c**2*x**6 - x**4), x))/c

Maxima [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^3(1-c^2x^2)} dx = \int -\frac{\sqrt{\frac{1}{cx}+1}\sqrt{\frac{1}{cx}-1}+\frac{1}{cx}}{(c^2x^2-1)x^3} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/x^3/(-c^2*x^2+1),x, algorithm="maxima")

[Out] 1/2*c^2*log(c*x + 1) - 1/2*c^2*log(c*x - 1) + c*integrate(x^(-2), x) + integrate(x^(-4), x)/c - integrate(sqrt(c*x + 1)*sqrt(-c*x + 1)/(c^3*x^6 - c*x^4), x)

Giac [F]

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^3(1-c^2x^2)} dx = \int -\frac{\sqrt{\frac{1}{cx}+1}\sqrt{\frac{1}{cx}-1}+\frac{1}{cx}}{(c^2x^2-1)x^3} dx$$

[In] integrate((1/c/x+(-1+1/c/x)^(1/2)*(1+1/c/x)^(1/2))/x^3/(-c^2*x^2+1),x, algorithm="giac")

[Out] integrate(-(sqrt(1/(c*x) + 1)*sqrt(1/(c*x) - 1) + 1/(c*x))/((c^2*x^2 - 1)*x^3), x)

Mupad [B] (verification not implemented)

Time = 5.48 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.88

$$\int \frac{e^{\operatorname{sech}^{-1}(cx)}}{x^3(1-c^2x^2)} dx = c^2 \operatorname{atanh}(cx) - \frac{\left(\frac{\sqrt{\frac{1}{cx}+1}}{3} + \frac{2c^2x^2\sqrt{\frac{1}{cx}+1}}{3}\right)\sqrt{\frac{1}{cx}-1}}{x^2} - \frac{c^2x^2 + \frac{1}{3}}{cx^3}$$

[In] int(-((1/(c*x) - 1)^(1/2)*(1/(c*x) + 1)^(1/2) + 1/(c*x))/(x^3*(c^2*x^2 - 1)),x)

[Out] c^2*atanh(c*x) - (((1/(c*x) + 1)^(1/2)/3 + (2*c^2*x^2*(1/(c*x) + 1)^(1/2))/3)*(1/(c*x) - 1)^(1/2))/x^2 - (c^2*x^2 + 1/3)/(c*x^3)

$$3.97 \quad \int \frac{x \left(-1 + a e^{\operatorname{sech}^{-1}(ax)x} \right)}{1 - a^2 x^2} dx$$

Optimal result	589
Rubi [B] (verified)	589
Mathematica [B] (verified)	591
Maple [A] (verified)	591
Fricas [A] (verification not implemented)	591
Sympy [F]	592
Maxima [F]	592
Giac [F]	592
Mupad [B] (verification not implemented)	593

Optimal result

Integrand size = 25, antiderivative size = 12

$$\int \frac{x \left(-1 + a e^{\operatorname{sech}^{-1}(ax)x} \right)}{1 - a^2 x^2} dx = -\frac{e^{\operatorname{sech}^{-1}(ax)x}}{a}$$

[Out] $-(1/a/x + (1/a/x - 1)^{(1/2)} * (1 + 1/a/x)^{(1/2)}) * x/a$

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 26 vs. $2(12) = 24$.

Time = 0.82 (sec) , antiderivative size = 26, normalized size of antiderivative = 2.17, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {6857, 266, 6476, 1972, 75}

$$\int \frac{x \left(-1 + a e^{\operatorname{sech}^{-1}(ax)x} \right)}{1 - a^2 x^2} dx = -\frac{\sqrt{1 - ax}}{a^2 \sqrt{\frac{1}{ax+1}}}$$

[In] $\text{Int}[(x*(-1 + a*E^{\text{ArcSech}[a*x]*x)})/(1 - a^2*x^2), x]$

[Out] $-(\text{Sqrt}[1 - a*x]/(a^2*\text{Sqrt}[(1 + a*x)^{-1}])))$

Rule 75

$\text{Int}[(a_. + (b_.)*(x_.))*((c_. + (d_.)*(x_.))^{(n_.)*((e_. + (f_.)*(x_.))^{(p_.)}, x_Symbol] :> \text{Simp}[b*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)}/(d*f*(n + p + 2))], x] /; \text{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \&\& \text{NeQ}[n + p + 2, 0] \&\& \text{EqQ}$

$[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]$

Rule 266

$\text{Int}[(x_)^{(m_.)}/((a_) + (b_.)*(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^n, x]]/(b*n), x] /; \text{FreeQ}\{a, b, m, n\}, x\} \&\& \text{EqQ}[m, n - 1]$

Rule 1972

$\text{Int}[(u_.)*((c_.)*((a_.) + (b_.)*(x_)^{(n_.)})^{(q_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[\text{Simp}[(c*(a + b*x^n)^q)^p/(a + b*x^n)^{(p*q)}], \text{Int}[u*(a + b*x^n)^{(p*q)}, x], x] /; \text{FreeQ}\{a, b, c, n, p, q\}, x\} \&\& \text{GeQ}[a, 0]$

Rule 6476

$\text{Int}[(E^{\text{ArcSech}[(c_.)*(x_)])*((d_.)*(x_)^{(m_.)})}/((a_) + (b_.)*(x_)^2), x_Symbol] \rightarrow \text{Dist}[d/(a*c), \text{Int}[(d*x)^{(m-1)}*(\text{Sqrt}[1/(1 + c*x)]/\text{Sqrt}[1 - c*x]), x], x] + \text{Dist}[d/c, \text{Int}[(d*x)^{(m-1)}/(a + b*x^2), x], x] /; \text{FreeQ}\{a, b, c, d, m\}, x\} \&\& \text{EqQ}[b + a*c^2, 0]$

Rule 6857

$\text{Int}[(u_)/((a_) + (b_.)*(x_)^{(n_.)}), x_Symbol] \rightarrow \text{With}\{v = \text{RationalFunctionExpand}[u/(a + b*x^n), x]\}, \text{Int}[v, x] /; \text{SumQ}[v] /; \text{FreeQ}\{a, b\}, x\} \&\& \text{IGtQ}[n, 0]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \left(\frac{x}{-1 + a^2 x^2} - \frac{a e^{\text{sech}^{-1}(ax)} x^2}{-1 + a^2 x^2} \right) dx \\
 &= - \left(a \int \frac{e^{\text{sech}^{-1}(ax)} x^2}{-1 + a^2 x^2} dx \right) + \int \frac{x}{-1 + a^2 x^2} dx \\
 &= \frac{\log(1 - a^2 x^2)}{2a^2} + \int \frac{x \sqrt{\frac{1}{1+ax}}}{\sqrt{1-ax}} dx - \int \frac{x}{-1 + a^2 x^2} dx \\
 &= \left(\sqrt{\frac{1}{1+ax}} \sqrt{1+ax} \right) \int \frac{x}{\sqrt{1-ax} \sqrt{1+ax}} dx \\
 &= -\frac{\sqrt{1-ax}}{a^2 \sqrt{\frac{1}{1+ax}}}
 \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 28 vs. $2(12) = 24$.

Time = 0.44 (sec) , antiderivative size = 28, normalized size of antiderivative = 2.33

$$\int \frac{x(-1 + ae^{\operatorname{sech}^{-1}(ax)x})}{1 - a^2x^2} dx = -\frac{\sqrt{\frac{1-ax}{1+ax}}(1+ax)}{a^2}$$

[In] Integrate[(x*(-1 + a*E^ArcSech[a*x]*x))/(1 - a^2*x^2), x]

[Out] -((Sqrt[(1 - a*x)/(1 + a*x)]*(1 + a*x))/a^2)

Maple [A] (verified)

Time = 0.73 (sec) , antiderivative size = 36, normalized size of antiderivative = 3.00

method	result	size
gospers	$-\frac{x\sqrt{-\frac{ax-1}{ax}}\sqrt{\frac{ax+1}{ax}}}{a}$	36
default	$-\frac{x\sqrt{-\frac{ax-1}{ax}}\sqrt{\frac{ax+1}{ax}}}{a}$	36
risch	$-\frac{x\sqrt{-\frac{ax-1}{ax}}\sqrt{\frac{ax+1}{ax}}}{a}$	36

[In] int(x*(-1+a*(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x)/(-a^2*x^2+1), x, method=_RETURNVERBOSE)

[Out] -1/a*x*(-(a*x-1)/a/x)^(1/2)*((a*x+1)/a/x)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.92

$$\int \frac{x(-1 + ae^{\operatorname{sech}^{-1}(ax)x})}{1 - a^2x^2} dx = -\frac{x\sqrt{\frac{ax+1}{ax}}\sqrt{-\frac{ax-1}{ax}}}{a}$$

[In] integrate(x*(-1+a*(1/a/x+(1/a/x-1)^(1/2)*(1+1/a/x)^(1/2))*x)/(-a^2*x^2+1), x, algorithm="fricas")

[Out] -x*sqrt((a*x + 1)/(a*x))*sqrt(-(a*x - 1)/(a*x))/a

SymPy [F]

$$\int \frac{x \left(-1 + a e^{\operatorname{sech}^{-1}(ax)x} \right)}{1 - a^2 x^2} dx = -a \int \frac{x^2 \sqrt{-1 + \frac{1}{ax}} \sqrt{1 + \frac{1}{ax}}}{a^2 x^2 - 1} dx$$

[In] integrate(x*(-1+a*(1/a/x+(1/a/x-1)**(1/2))*(1+1/a/x)**(1/2))*x)/(-a**2*x**2+1),x)

[Out] -a*Integral(x**2*sqrt(-1 + 1/(a*x))*sqrt(1 + 1/(a*x))/(a**2*x**2 - 1), x)

Maxima [F]

$$\int \frac{x \left(-1 + a e^{\operatorname{sech}^{-1}(ax)x} \right)}{1 - a^2 x^2} dx = \int -\frac{\left(ax \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}} \right) - 1 \right) x}{a^2 x^2 - 1} dx$$

[In] integrate(x*(-1+a*(1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))*x)/(-a^2*x^2+1),x, algorithm="maxima")

[Out] -integrate((a*x*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x)) - 1)*x/(a^2*x^2 - 1), x)

Giac [F]

$$\int \frac{x \left(-1 + a e^{\operatorname{sech}^{-1}(ax)x} \right)}{1 - a^2 x^2} dx = \int -\frac{\left(ax \left(\sqrt{\frac{1}{ax} + 1} \sqrt{\frac{1}{ax} - 1 + \frac{1}{ax}} \right) - 1 \right) x}{a^2 x^2 - 1} dx$$

[In] integrate(x*(-1+a*(1/a/x+(1/a/x-1)^(1/2))*(1+1/a/x)^(1/2))*x)/(-a^2*x^2+1),x, algorithm="giac")

[Out] integrate(-(a*x*(sqrt(1/(a*x) + 1)*sqrt(1/(a*x) - 1) + 1/(a*x)) - 1)*x/(a^2*x^2 - 1), x)

Mupad [B] (verification not implemented)

Time = 5.78 (sec) , antiderivative size = 76, normalized size of antiderivative = 6.33

$$\int \frac{x \left(-1 + a e^{\operatorname{sech}^{-1}(ax)x} \right)}{1 - a^2 x^2} dx = \frac{\ln\left(\frac{1}{x}\right)}{a^2} - \frac{\ln\left(a + \frac{1}{x}\right)}{2a^2} - \frac{\ln\left(\frac{1}{x} - a\right)}{2a^2} + \frac{\ln(a^2 x^2 - 1)}{2a^2} - \frac{x \sqrt{\frac{1}{ax} - 1} \sqrt{\frac{1}{ax} + 1}}{a}$$

```
[In] int(-(x*(a*x*((1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2) + 1/(a*x)) - 1))/(a^2*x^2 - 1),x)
```

```
[Out] log(1/x)/a^2 - log(a + 1/x)/(2*a^2) - log(1/x - a)/(2*a^2) + log(a^2*x^2 - 1)/(2*a^2) - (x*(1/(a*x) - 1)^(1/2)*(1/(a*x) + 1)^(1/2))/a
```

3.98 $\int \frac{\operatorname{sech}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx$

Optimal result	594
Rubi [A] (verified)	594
Mathematica [A] (verified)	597
Maple [A] (verified)	597
Fricas [F]	597
Sympy [F]	598
Maxima [F]	598
Giac [F]	598
Mupad [F(-1)]	598

Optimal result

Integrand size = 19, antiderivative size = 61

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \frac{\operatorname{sech}^{-1}(a+bx)^2}{2d} - \frac{\operatorname{sech}^{-1}(a+bx) \log\left(1 + e^{2\operatorname{sech}^{-1}(a+bx)}\right)}{d} - \frac{\operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right)}{2d}$$

[Out] 1/2*arcsech(b*x+a)^2/d-arcsech(b*x+a)*ln(1+(1/(b*x+a)+(1/(b*x+a)-1)^(1/2))*1/(b*x+a)+1)^(1/2))/d-1/2*polylog(2,-(1/(b*x+a)+(1/(b*x+a)-1)^(1/2))*1/(b*x+a)+1)^(1/2))/d

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {6454, 12, 6416, 5882, 3799, 2221, 2317, 2438}

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = -\frac{\operatorname{PolyLog}\left(2, -e^{2\operatorname{sech}^{-1}(a+bx)}\right)}{2d} + \frac{\operatorname{sech}^{-1}(a+bx)^2}{2d} - \frac{\operatorname{sech}^{-1}(a+bx) \log\left(e^{2\operatorname{sech}^{-1}(a+bx)} + 1\right)}{d}$$

[In] Int[ArcSech[a + b*x]/((a*d)/b + d*x), x]

[Out] ArcSech[a + b*x]^2/(2*d) - (ArcSech[a + b*x]*Log[1 + E^(2*ArcSech[a + b*x])])/d - PolyLog[2, -E^(2*ArcSech[a + b*x])]/(2*d)

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 2221

Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[(c + d*x)^m/(b*f*g*n*Log[F])*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2317

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3799

Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

Rule 5882

Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)/(x_), x_Symbol] := Dist[1/b, Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]

Rule 6416

Int[((a_) + ArcSech[(c_)*(x_)])*(b_)/(x_), x_Symbol] := -Subst[Int[(a + b*ArcCosh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]

Rule 6454

Int[((a_) + ArcSech[(c_) + (d_)*(x_)])*(b_)^(p_)*((e_) + (f_)*(x_))^(m_), x_Symbol] := Dist[1/d, Subst[Int[(f*(x/d))^m*(a + b*ArcSech[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] &

& IGtQ[p, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int \frac{b \operatorname{sech}^{-1}(x)}{dx} dx, x, a + bx\right)}{b} \\
 &= \frac{\text{Subst}\left(\int \frac{\operatorname{sech}^{-1}(x)}{x} dx, x, a + bx\right)}{d} \\
 &= -\frac{\text{Subst}\left(\int \frac{\operatorname{arccosh}(x)}{x} dx, x, \frac{1}{a+bx}\right)}{d} \\
 &= -\frac{\text{Subst}\left(\int x \tanh(x) dx, x, \operatorname{arccosh}\left(\frac{1}{a+bx}\right)\right)}{d} \\
 &= \frac{\operatorname{arccosh}\left(\frac{1}{a+bx}\right)^2}{2d} - \frac{2\text{Subst}\left(\int \frac{e^{2x} x}{1+e^{2x}} dx, x, \operatorname{arccosh}\left(\frac{1}{a+bx}\right)\right)}{d} \\
 &= \frac{\operatorname{arccosh}\left(\frac{1}{a+bx}\right)^2}{2d} - \frac{\operatorname{arccosh}\left(\frac{1}{a+bx}\right) \log\left(1 + e^{2\operatorname{arccosh}\left(\frac{1}{a+bx}\right)}\right)}{d} \\
 &\quad + \frac{\text{Subst}\left(\int \log(1 + e^{2x}) dx, x, \operatorname{arccosh}\left(\frac{1}{a+bx}\right)\right)}{d} \\
 &= \frac{\operatorname{arccosh}\left(\frac{1}{a+bx}\right)^2}{2d} - \frac{\operatorname{arccosh}\left(\frac{1}{a+bx}\right) \log\left(1 + e^{2\operatorname{arccosh}\left(\frac{1}{a+bx}\right)}\right)}{d} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2\operatorname{arccosh}\left(\frac{1}{a+bx}\right)}\right)}{2d} \\
 &= \frac{\operatorname{arccosh}\left(\frac{1}{a+bx}\right)^2}{2d} - \frac{\operatorname{arccosh}\left(\frac{1}{a+bx}\right) \log\left(1 + e^{2\operatorname{arccosh}\left(\frac{1}{a+bx}\right)}\right)}{d} - \frac{\operatorname{PolyLog}\left(2, -e^{2\operatorname{arccosh}\left(\frac{1}{a+bx}\right)}\right)}{2d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.85

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \frac{-\operatorname{sech}^{-1}(a+bx) \left(\operatorname{sech}^{-1}(a+bx) + 2 \log \left(1 + e^{-2\operatorname{sech}^{-1}(a+bx)} \right) \right) + \operatorname{PolyLog} \left(2, -e^{-2\operatorname{sech}^{-1}(a+bx)} \right)}{2d}$$

`[In] Integrate[ArcSech[a + b*x]/((a*d)/b + d*x), x]`

```
[Out] (-ArcSech[a + b*x]*(ArcSech[a + b*x] + 2*Log[1 + E^(-2*ArcSech[a + b*x])])
) + PolyLog[2, -E^(-2*ArcSech[a + b*x])])/(2*d)
```

Maple [A] (verified)

Time = 0.78 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.82

method	result
derivativedivides	$\frac{b \operatorname{arcsech}(bx+a)^2}{2d} - \frac{b \operatorname{arcsech}(bx+a) \ln \left(1 + \left(\frac{1}{bx+a} + \sqrt{\frac{1}{bx+a} - 1} \sqrt{\frac{1}{bx+a} + 1} \right)^2 \right)}{d} - \frac{b \operatorname{polylog} \left(2, - \left(\frac{1}{bx+a} + \sqrt{\frac{1}{bx+a} - 1} \sqrt{\frac{1}{bx+a} + 1} \right)^2 \right)}{2d}$
default	$\frac{b \operatorname{arcsech}(bx+a)^2}{2d} - \frac{b \operatorname{arcsech}(bx+a) \ln \left(1 + \left(\frac{1}{bx+a} + \sqrt{\frac{1}{bx+a} - 1} \sqrt{\frac{1}{bx+a} + 1} \right)^2 \right)}{d} - \frac{b \operatorname{polylog} \left(2, - \left(\frac{1}{bx+a} + \sqrt{\frac{1}{bx+a} - 1} \sqrt{\frac{1}{bx+a} + 1} \right)^2 \right)}{2d}$

`[In] int(arcsech(b*x+a)/(a*d/b+d*x), x, method=_RETURNVERBOSE)`

```
[Out] 1/b*(1/2*b/d*arcsech(b*x+a)^2-b/d*arcsech(b*x+a)*ln(1+(1/(b*x+a)+(1/(b*x+a)
-1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2)-1/2*b/d*polylog(2,-(1/(b*x+a)+(1/(b*x+a)-
1)^(1/2)*(1/(b*x+a)+1)^(1/2))^2))
```

Fricas [F]

$$\int \frac{\operatorname{sech}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \int \frac{\operatorname{arosech}(bx+a)}{dx + \frac{ad}{b}} dx$$

`[In] integrate(arcsech(b*x+a)/(a*d/b+d*x), x, algorithm="fricas")``[Out] integral(b*arcsech(b*x + a)/(b*d*x + a*d), x)`

Sympy [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \frac{b \int \frac{\operatorname{asech}(a+bx)}{a+bx} dx}{d}$$

[In] integrate(asech(b*x+a)/(a*d/b+d*x),x)

[Out] b*Integral(asech(a + b*x)/(a + b*x), x)/d

Maxima [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{arsech}(bx + a)}{dx + \frac{ad}{b}} dx$$

[In] integrate(arcsech(b*x+a)/(a*d/b+d*x),x, algorithm="maxima")

[Out] 1/2*(2*log(sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*b*x + sqrt(b*x + a + 1)*sqrt(-b*x - a + 1)*a + b*x + a)*log(b*x + a) - 3*log(b*x + a)^2)/d - 1/2*(log(b*x + a + 1)*log(b*x + a) + dilog(-b*x - a))/d - 1/2*(log(b*x + a)*log(-b*x - a + 1) + dilog(b*x + a))/d + integrate((b^2*x + a*b)*log(b*x + a)/(b^2*d*x^2 + 2*a*b*d*x + a^2*d + (b^2*d*x^2 + 2*a*b*d*x + a^2*d - d)*sqrt(b*x + a + 1)*sqrt(-b*x - a + 1) - d), x)

Giac [F]

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{arsech}(bx + a)}{dx + \frac{ad}{b}} dx$$

[In] integrate(arcsech(b*x+a)/(a*d/b+d*x),x, algorithm="giac")

[Out] integrate(arcsech(b*x + a)/(d*x + a*d/b), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{sech}^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{acosh}\left(\frac{1}{a+bx}\right)}{dx + \frac{ad}{b}} dx$$

[In] int(acosh(1/(a + b*x))/(d*x + (a*d)/b),x)

[Out] int(acosh(1/(a + b*x))/(d*x + (a*d)/b), x)

3.99 $\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx$

Optimal result	599
Rubi [A] (verified)	599
Mathematica [A] (verified)	601
Maple [A] (verified)	601
Fricas [B] (verification not implemented)	601
Sympy [F]	602
Maxima [A] (verification not implemented)	602
Giac [F]	602
Mupad [B] (verification not implemented)	603

Optimal result

Integrand size = 12, antiderivative size = 57

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx = \frac{(a + bx^4) \operatorname{sech}^{-1}(a + bx^4)}{4b} - \frac{\arctan\left(\sqrt{\frac{1-a-bx^4}{1+a+bx^4}}\right)}{2b}$$

[Out] $1/4*(b*x^4+a)*\operatorname{arcsech}(b*x^4+a)/b-1/2*\arctan(((-b*x^4-a+1)/(b*x^4+a+1))^{1/2})/b$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {6847, 6448, 1983, 12, 209}

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx = \frac{(a + bx^4) \operatorname{sech}^{-1}(a + bx^4)}{4b} - \frac{\arctan\left(\sqrt{\frac{-a-bx^4+1}{a+bx^4+1}}\right)}{2b}$$

[In] $\operatorname{Int}[x^3*\operatorname{ArcSech}[a + b*x^4], x]$

[Out] $((a + b*x^4)*\operatorname{ArcSech}[a + b*x^4])/(4*b) - \operatorname{ArcTan}[\operatorname{Sqrt}[(1 - a - b*x^4)/(1 + a + b*x^4)]]/(2*b)$

Rule 12

$\operatorname{Int}[(a_*)*(u_), x_Symbol] \rightarrow \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /;$ FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 1983

```
Int[(u_)^(r_.)*(((e_.)*((a_.) + (b_.)*(x_)^(n_.)))/((c_) + (d_.)*(x_)^(n_.)))^(p_), x_Symbol] := With[{q = Denominator[p]}, Dist[q*e*((b*c - a*d)/n), Subst[Int[SimplifyIntegrand[x^(q*(p + 1) - 1)*((-a)*e + c*x^q)^(1/n - 1)/(b*e - d*x^q)^(1/n + 1)*(u /. x -> ((-a)*e + c*x^q)^(1/n)/(b*e - d*x^q)^(1/n))]^r, x], x], x, (e*((a + b*x^n)/(c + d*x^n)))^(1/q)], x] /; FreeQ[{a, b, c, d, e}, x] && PolynomialQ[u, x] && FractionQ[p] && IntegerQ[1/n] && IntegerQ[r]
```

Rule 6448

```
Int[ArcSech[(c_) + (d_.)*(x_)], x_Symbol] := Simp[(c + d*x)*(ArcSech[c + d*x]/d), x] + Int[Sqrt[(1 - c - d*x)/(1 + c + d*x)]/(1 - c - d*x), x] /; FreeQ[{c, d}, x]
```

Rule 6847

```
Int[(u_)*(x_)^(m_.), x_Symbol] := Dist[1/(m + 1), Subst[Int[SubstFor[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^(m + 1), u, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{4} \text{Subst} \left(\int \text{sech}^{-1}(a + bx) dx, x, x^4 \right) \\
 &= \frac{(a + bx^4) \text{sech}^{-1}(a + bx^4)}{4b} + \frac{1}{4} \text{Subst} \left(\int \frac{\sqrt{\frac{1-a-bx}{1+a+bx}}}{1-a-bx} dx, x, x^4 \right) \\
 &= \frac{(a + bx^4) \text{sech}^{-1}(a + bx^4)}{4b} - b \text{Subst} \left(\int \frac{1}{2b^2(1+x^2)} dx, x, \sqrt{\frac{1-a-bx^4}{1+a+bx^4}} \right) \\
 &= \frac{(a + bx^4) \text{sech}^{-1}(a + bx^4)}{4b} - \frac{\text{Subst} \left(\int \frac{1}{1+x^2} dx, x, \sqrt{\frac{1-a-bx^4}{1+a+bx^4}} \right)}{2b} \\
 &= \frac{(a + bx^4) \text{sech}^{-1}(a + bx^4)}{4b} - \frac{\arctan \left(\sqrt{\frac{1-a-bx^4}{1+a+bx^4}} \right)}{2b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.86

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx$$

$$= \frac{(a + bx^4) \operatorname{sech}^{-1}(a + bx^4) + \frac{2\sqrt{-\frac{-1+a+bx^4}{1+a+bx^4}} \sqrt{1-(a+bx^4)^2} \arctan\left(\frac{\sqrt{1-(a+bx^4)^2}}{-1+a+bx^4}\right)}{-1+a+bx^4}}{4b}$$

`[In] Integrate[x^3*ArcSech[a + b*x^4],x]`

```
[Out] ((a + b*x^4)*ArcSech[a + b*x^4] + (2*Sqrt[-((-1 + a + b*x^4)/(1 + a + b*x^4))]*Sqrt[1 - (a + b*x^4)^2]*ArcTan[Sqrt[1 - (a + b*x^4)^2]/(-1 + a + b*x^4)])/(-1 + a + b*x^4))/(4*b)
```

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.93

method	result	size
derivativedivides	$\frac{(bx^4+a) \operatorname{arcsech}(bx^4+a) - \arctan\left(\sqrt{\frac{1}{bx^4+a}} - 1 \sqrt{\frac{1}{bx^4+a} + 1}\right)}{4b}$	53
default	$\frac{(bx^4+a) \operatorname{arcsech}(bx^4+a) - \arctan\left(\sqrt{\frac{1}{bx^4+a}} - 1 \sqrt{\frac{1}{bx^4+a} + 1}\right)}{4b}$	53

`[In] int(x^3*arcsech(b*x^4+a),x,method=_RETURNVERBOSE)`

```
[Out] 1/4/b*((b*x^4+a)*arcsech(b*x^4+a)-arctan((1/(b*x^4+a)-1)^(1/2)*(1/(b*x^4+a)+1)^(1/2)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 283 vs. 2(49) = 98.

Time = 0.27 (sec) , antiderivative size = 283, normalized size of antiderivative = 4.96

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx$$

$$= \frac{2bx^4 \log\left(\frac{(bx^4+a)\sqrt{-\frac{b^2x^8+2abx^4+a^2-1}{b^2x^8+2abx^4+a^2}+1}}{bx^4+a}\right) + a \log\left(\frac{(bx^4+a)\sqrt{-\frac{b^2x^8+2abx^4+a^2-1}{b^2x^8+2abx^4+a^2}+1}}{x^4}\right) - a \log\left(\frac{(bx^4+a)\sqrt{-\frac{b^2x^8+2abx^4+a^2-1}{b^2x^8+2abx^4+a^2}+1}}{x^4}\right)}{8b}$$

`[In] integrate(x^3*arcsech(b*x^4+a),x, algorithm="fricas")`

```
[Out] 1/8*(2*b*x^4*log(((b*x^4 + a)*sqrt(-(b^2*x^8 + 2*a*b*x^4 + a^2 - 1)/(b^2*x^8 + 2*a*b*x^4 + a^2)) + 1)/(b*x^4 + a)) + a*log(((b*x^4 + a)*sqrt(-(b^2*x^8 + 2*a*b*x^4 + a^2 - 1)/(b^2*x^8 + 2*a*b*x^4 + a^2)) + 1)/x^4) - a*log(((b*x^4 + a)*sqrt(-(b^2*x^8 + 2*a*b*x^4 + a^2 - 1)/(b^2*x^8 + 2*a*b*x^4 + a^2)) - 1)/x^4) - 2*arctan((b^2*x^8 + 2*a*b*x^4 + a^2)*sqrt(-(b^2*x^8 + 2*a*b*x^4 + a^2 - 1)/(b^2*x^8 + 2*a*b*x^4 + a^2)))/(b^2*x^8 + 2*a*b*x^4 + a^2 - 1)))/b
```

Sympy [F]

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx = \int x^3 \operatorname{asech}(a + bx^4) dx$$

```
[In] integrate(x**3*asech(b*x**4+a), x)
```

```
[Out] Integral(x**3*asech(a + b*x**4), x)
```

Maxima [A] (verification not implemented)

none

Time = 0.20 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.67

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx = \frac{(bx^4 + a) \operatorname{arsech}(bx^4 + a) - \arctan\left(\sqrt{\frac{1}{(bx^4 + a)^2} - 1}\right)}{4b}$$

```
[In] integrate(x^3*arcsech(b*x^4+a), x, algorithm="maxima")
```

```
[Out] 1/4*((b*x^4 + a)*arcsech(b*x^4 + a) - arctan(sqrt(1/(b*x^4 + a)^2 - 1)))/b
```

Giac [F]

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx = \int x^3 \operatorname{arsech}(bx^4 + a) dx$$

```
[In] integrate(x^3*arcsech(b*x^4+a), x, algorithm="giac")
```

```
[Out] integrate(x^3*arcsech(b*x^4 + a), x)
```

Mupad [B] (verification not implemented)

Time = 5.58 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.98

$$\int x^3 \operatorname{sech}^{-1}(a + bx^4) dx = \frac{\operatorname{atan}\left(\frac{1}{\sqrt{\frac{1}{bx^4+a}-1}\sqrt{\frac{1}{bx^4+a}+1}}\right)}{4b} + \frac{\operatorname{acosh}\left(\frac{1}{bx^4+a}\right)(bx^4+a)}{4b}$$

[In] int(x^3*acosh(1/(a + b*x^4)),x)

[Out] atan(1/((1/(a + b*x^4) - 1)^(1/2)*(1/(a + b*x^4) + 1)^(1/2)))/(4*b) + (acosh(1/(a + b*x^4))*(a + b*x^4))/(4*b)

3.100 $\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx$

Optimal result	604
Rubi [A] (verified)	604
Mathematica [A] (verified)	606
Maple [F]	606
Fricas [B] (verification not implemented)	606
Sympy [F(-1)]	607
Maxima [A] (verification not implemented)	607
Giac [F]	607
Mupad [B] (verification not implemented)	608

Optimal result

Integrand size = 14, antiderivative size = 58

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx = \frac{(a + bx^n) \operatorname{sech}^{-1}(a + bx^n)}{bn} - \frac{2 \arctan\left(\sqrt{\frac{1-a-bx^n}{1+a+bx^n}}\right)}{bn}$$

[Out] $(a+b*x^n)*\operatorname{arcsech}(a+b*x^n)/b/n-2*\arctan(((1-a-b*x^n)/(1+a+b*x^n))^{(1/2)})/b/n$

Rubi [A] (verified)

Time = 0.09 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {6847, 6448, 1983, 12, 209}

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx = \frac{(a + bx^n) \operatorname{sech}^{-1}(a + bx^n)}{bn} - \frac{2 \arctan\left(\sqrt{\frac{-a-bx^n+1}{a+bx^n+1}}\right)}{bn}$$

[In] `Int[x^(-1 + n)*ArcSech[a + b*x^n],x]`

[Out] `((a + b*x^n)*ArcSech[a + b*x^n])/(b*n) - (2*ArcTan[Sqrt[(1 - a - b*x^n)/(1 + a + b*x^n)]])/(b*n)`

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]`

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 1983

Int[(u_)^(r_)*(((e_)*((a_) + (b_)*(x_)^(n_))))/((c_) + (d_)*(x_)^(n_))^(p_), x_Symbol] := With[{q = Denominator[p]}, Dist[q*e*((b*c - a*d)/n), Subst[Int[SimplifyIntegrand[x^(q*(p + 1) - 1)*(((a)*e + c*x^q)^(1/n - 1)/(b*e - d*x^q)^(1/n + 1))*(u /. x -> ((a)*e + c*x^q)^(1/n)/(b*e - d*x^q)^(1/n))^r, x], x], x, (e*((a + b*x^n)/(c + d*x^n))^(1/q)], x]] /; FreeQ[{a, b, c, d, e}, x] && PolynomialQ[u, x] && FractionQ[p] && IntegerQ[1/n] && IntegerQ[r]

Rule 6448

Int[ArcSech[(c_) + (d_)*(x_)], x_Symbol] := Simp[(c + d*x)*(ArcSech[c + d*x]/d), x] + Int[Sqrt[(1 - c - d*x)/(1 + c + d*x)]/(1 - c - d*x), x] /; FreeQ[{c, d}, x]

Rule 6847

Int[(u_)*(x_)^(m_), x_Symbol] := Dist[1/(m + 1), Subst[Int[SubstFor[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^(m + 1), u, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int \text{sech}^{-1}(a + bx) dx, x, x^n\right)}{n} \\
 &= \frac{(a + bx^n) \text{sech}^{-1}(a + bx^n)}{bn} + \frac{\text{Subst}\left(\int \frac{\sqrt{\frac{1-a-bx}{1+a+bx}}}{1-a-bx} dx, x, x^n\right)}{n} \\
 &= \frac{(a + bx^n) \text{sech}^{-1}(a + bx^n)}{bn} - \frac{(4b)\text{Subst}\left(\int \frac{1}{2b^2(1+x^2)} dx, x, \sqrt{\frac{1-a-bx^n}{1+a+bx^n}}\right)}{n} \\
 &= \frac{(a + bx^n) \text{sech}^{-1}(a + bx^n)}{bn} - \frac{2\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\frac{1-a-bx^n}{1+a+bx^n}}\right)}{bn} \\
 &= \frac{(a + bx^n) \text{sech}^{-1}(a + bx^n)}{bn} - \frac{2 \arctan\left(\sqrt{\frac{1-a-bx^n}{1+a+bx^n}}\right)}{bn}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.83

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx$$

$$= \frac{(a + bx^n) \operatorname{sech}^{-1}(a + bx^n) + \frac{2\sqrt{-\frac{-1+a+bx^n}{1+a+bx^n}} \sqrt{1-(a+bx^n)^2} \arctan\left(\frac{\sqrt{1-(a+bx^n)^2}}{-1+a+bx^n}\right)}{-1+a+bx^n}}{bn}$$

[In] Integrate[x[^](-1 + n)*ArcSech[a + b*x[^]n], x]

[Out] ((a + b*x[^]n)*ArcSech[a + b*x[^]n] + (2*Sqrt[-((-1 + a + b*x[^]n)/(1 + a + b*x[^]n))]*Sqrt[1 - (a + b*x[^]n)^2]*ArcTan[Sqrt[1 - (a + b*x[^]n)^2]/(-1 + a + b*x[^]n)])/(-1 + a + b*x[^]n))/(b*n)

Maple [F]

$$\int x^{-1+n} \operatorname{arcsech}(a + bx^n) dx$$

[In] int(x[^](-1+n)*arcsech(a+b*x[^]n), x)

[Out] int(x[^](-1+n)*arcsech(a+b*x[^]n), x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 385 vs. 2(54) = 108.

Time = 0.30 (sec) , antiderivative size = 385, normalized size of antiderivative = 6.64

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx$$

$$= \frac{2(b \cosh(n \log(x)) + b \sinh(n \log(x))) \log\left(\frac{\sqrt{-\frac{2ab+(a^2+b^2-1)\cosh(n\log(x))-(a^2-b^2-1)\sinh(n\log(x))}{\cosh(n\log(x))-\sinh(n\log(x))}}+1}}{b \cosh(n \log(x)) + b \sinh(n \log(x)) + a}\right) + a \log\left(\frac{\sqrt{-\frac{2ab+(a^2+b^2-1)\cosh(n\log(x))-(a^2-b^2-1)\sinh(n\log(x))}{\cosh(n\log(x))-\sinh(n\log(x))}}+1}}{b \cosh(n \log(x)) + b \sinh(n \log(x)) + a}\right)}{bn}$$

[In] integrate(x[^](-1+n)*arcsech(a+b*x[^]n), x, algorithm="fricas")

[Out] 1/2*(2*(b*cosh(n*log(x)) + b*sinh(n*log(x)))*log((sqrt(-(2*a*b + (a^2 + b^2 - 1)*cosh(n*log(x)) - (a^2 - b^2 - 1)*sinh(n*log(x)))/(cosh(n*log(x)) - sinh(n*log(x)))) + 1)/(b*cosh(n*log(x)) + b*sinh(n*log(x)) + a)) + a*log((sqrt(-(2*a*b + (a^2 + b^2 - 1)*cosh(n*log(x)) - (a^2 - b^2 - 1)*sinh(n*log(x)))/(cosh(n*log(x)) - sinh(n*log(x)))) + 1)/(cosh(n*log(x)) + sinh(n*log(x)))) - a*log((sqrt(-(2*a*b + (a^2 + b^2 - 1)*cosh(n*log(x)) - (a^2 - b^2 - 1)*sinh(n*log(x)))/(cosh(n*log(x)) - sinh(n*log(x)))) + 1)/(cosh(n*log(x)) + sinh(n*log(x))))

$$\frac{\sinh(n \log(x)) / (\cosh(n \log(x)) - \sinh(n \log(x))) - 1}{(\cosh(n \log(x)) + \sinh(n \log(x))) - 2 \arctan((b \cosh(n \log(x)) + b \sinh(n \log(x)) + a) \sqrt{-(2ab + (a^2 + b^2 - 1) \cosh(n \log(x)) - (a^2 - b^2 - 1) \sinh(n \log(x)))})} / (\cosh(n \log(x)) - \sinh(n \log(x))) / (b^2 \cosh(n \log(x))^2 + b^2 \sinh(n \log(x))^2 + 2ab \cosh(n \log(x)) + a^2 + 2(b^2 \cosh(n \log(x)) + ab) \sinh(n \log(x)) - 1) / (bn)$$

Sympy [F(-1)]

Timed out.

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx = \text{Timed out}$$

[In] integrate(x**(-1+n)*asech(a+b*x**n),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.20 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.69

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx = \frac{(bx^n + a) \operatorname{ar} \operatorname{sech}(bx^n + a) - \arctan\left(\sqrt{\frac{1}{(bx^n + a)^2} - 1}\right)}{bn}$$

[In] integrate(x^(-1+n)*arcsech(a+b*x^n),x, algorithm="maxima")

[Out] ((b*x^n + a)*arcsech(b*x^n + a) - arctan(sqrt(1/(b*x^n + a)^2 - 1)))/(b*n)

Giac [F]

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx = \int x^{n-1} \operatorname{ar} \operatorname{sech}(bx^n + a) dx$$

[In] integrate(x^(-1+n)*arcsech(a+b*x^n),x, algorithm="giac")

[Out] integrate(x^(n - 1)*arcsech(b*x^n + a), x)

Mupad [B] (verification not implemented)

Time = 5.01 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.93

$$\int x^{-1+n} \operatorname{sech}^{-1}(a + bx^n) dx = \frac{\operatorname{atan}\left(\frac{1}{\sqrt{\frac{1}{a+bx^n}-1}\sqrt{\frac{1}{a+bx^n}+1}}\right) + \operatorname{acosh}\left(\frac{1}{a+bx^n}\right) (a + bx^n)}{bn}$$

[In] `int(x^(n - 1)*acosh(1/(a + b*x^n)),x)`

[Out] `(atan(1/(((1/(a + b*x^n) - 1)^(1/2))*(1/(a + b*x^n) + 1)^(1/2)))) + acosh(1/(a + b*x^n))*(a + b*x^n))/(b*n)`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 609

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*      Small rewrite of logic in main function to make it*)
(*      match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
```

```

(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCo
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count is
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"}
    ]
  ]
  ,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<>
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

  finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)

```

```

(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```

```

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3, ExpnType[expn[[1]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
            If[Head[expn]===RootSum,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,

```

```

    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result, optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);

```

```

#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 ("
```

```

                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_c
    end if
  else #result contains complex but optimal is not
    if debug then
      print("result contains complex but optimal is not");
    fi;
    return "C","Result contains complex when optimal does not.";
  fi;
else # result do not contain complex
  # this assumes optimal do not as well. No check is needed here.
  if debug then
    print("result do not contain complex, this assumes optimal do not as well");
  fi;
  if leaf_count_result<=2*leaf_count_optimal then
    if debug then
      print("leaf_count_result<=2*leaf_count_optimal");
    fi;
    return "A"," ";
  else
    if debug then
      print("leaf_count_result>2*leaf_count_optimal");
    fi;
    return "B",cat("Leaf count of result is larger than twice the leaf count of opt
                                convert(leaf_count_result,string)," $ vs. $2(",
                                convert(leaf_count_optimal,string)," )=",convert(2*leaf_count
    fi;
  fi;
else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
                convert(ExpnType_result,string)," vs. order ",
                convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

```

```

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`) or type(expn,'*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else

```

```

9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u), u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```


Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#           Port of original Maple grading function by
#           Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#           added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnTy
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

```

```

# print("Before returning. grade=", grade, " grade_annotation=", grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

# Dec 24, 2019. Nasser: Ported original Maple grading function by
# Albert Rich to use with Sagemath. This is used to
# grade Fricas, Giac and Maxima results.
# Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
# 'arctan2', 'floor', 'abs', 'log_integral'
# June 4, 2022 Made default grade_annotation "none" instead of "" due
# issue later when reading the file.
# July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    # print("Enter tree_size, expr is ", expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: # isinstance(expr, Pow):
        if expr.operands()[1] == 1/2: # expr.args[1] == Rational(1,2):
            if debug: print("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception,AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

    return 1
  else:
    return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
  else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. " + str(leaf_c

else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_result

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```