

Computer Algebra Independent Integration Tests

Summer 2023 edition

7-Inverse-hyperbolic-functions/7.6-Inverse-hyperbolic-cosecant/203-
7.6.2-Inverse-hyperbolic-cosecant-functions

Nasser M. Abbasi

September 5, 2023

Compiled on September 5, 2023 at 2:00pm

Contents

1	Introduction	3
2	detailed summary tables of results	21
3	Listing of integrals	43
4	Appendix	427

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	8
1.4	Performance based on number of rules Rubi used	10
1.5	Performance based on number of steps Rubi used	11
1.6	Solved integrals histogram based on leaf size of result	12
1.7	Solved integrals histogram based on CPU time used	13
1.8	Leaf size vs. CPU time used	14
1.9	list of integrals with no known antiderivative	15
1.10	List of integrals solved by CAS but has no known antiderivative	15
1.11	list of integrals solved by CAS but failed verification	15
1.12	Timing	16
1.13	Verification	16
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [71]. This is test number [203].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (71)	0.00 (0)
Mathematica	98.59 (70)	1.41 (1)
Maple	74.65 (53)	25.35 (18)
Fricas	74.65 (53)	25.35 (18)
Maxima	59.15 (42)	40.85 (29)
Mupad	57.75 (41)	42.25 (30)
Sympy	50.70 (36)	49.30 (35)
Giac	45.07 (32)	54.93 (39)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	74.648	8.451	15.493	1.408
Fricas	47.887	26.761	0.000	25.352
Maxima	45.070	14.085	0.000	40.845
Sympy	42.254	1.408	7.042	49.296
Maple	30.986	36.620	7.042	25.352
Giac	28.169	16.901	0.000	54.930
Mupad	0.000	57.746	0.000	42.254

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	1	100.00	0.00	0.00
Fricas	18	88.89	0.00	11.11
Maple	18	100.00	0.00	0.00
Maxima	29	93.10	0.00	6.90
Mupad	30	0.00	100.00	0.00
Sympy	35	91.43	8.57	0.00
Giac	39	84.62	0.00	15.38

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Rubi	0.13
Maxima	0.23
Fricas	0.24
Giac	0.28
Maple	0.33
Mathematica	0.83
Sympy	1.67
Mupad	5.15

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	45.49	0.94	42.00	0.93
Sympy	57.83	1.13	50.00	1.18
Maxima	69.24	1.31	59.50	1.21
Giac	78.41	1.65	73.00	1.49
Fricas	104.72	1.78	70.00	1.37
Rubi	104.72	1.01	61.00	1.00
Maple	111.94	1.96	111.00	1.65
Mathematica	256.67	1.31	54.00	1.02

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {7, 8, 12, 13}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
```

```
x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in *Rubi Table file*

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2023
Design-vide

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	22
2.2	Detailed conclusion table per each integral for all CAS systems	25
2.3	Detailed conclusion table specific for Rubi results	40

2.1 List of integrals sorted by grade for each CAS

Rubi	22
Mma	22
Maple	23
Fricas	23
Maxima	23
Giac	24
Mupad	24
Sympy	24

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 6, 14, 15, 16, 17, 18, 19, 20, 21, 22, 24, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 39, 41, 43, 45, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69 }

B grade { 5, 9, 10, 25, 70, 71 }

C grade { 4, 7, 8, 12, 13, 23, 38, 40, 42, 44, 46 }

F normal fail { 11 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 14, 15, 16, 17, 19, 20, 21, 22, 27, 28, 29, 34, 36, 47, 49, 50, 55, 57, 70 }

B grade { 5, 6, 30, 31, 32, 33, 35, 39, 41, 43, 45, 51, 52, 53, 54, 56, 58, 60, 61, 62, 63, 64, 65, 66, 67, 68 }

C grade { 38, 40, 42, 44, 46 }

F normal fail { 4, 7, 8, 9, 10, 11, 12, 13, 18, 23, 24, 25, 26, 37, 48, 59, 69, 71 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 14, 15, 16, 17, 19, 20, 21, 22, 27, 28, 29, 30, 32, 34, 36, 38, 39, 40, 44, 46, 47, 49, 50, 51, 53, 57, 58, 60, 61, 62, 63, 64, 66, 68 }

B grade { 1, 2, 3, 5, 6, 31, 33, 35, 41, 43, 45, 52, 54, 55, 56, 65, 67, 70, 71 }

C grade { }

F normal fail { 4, 7, 8, 9, 10, 11, 12, 13, 18, 24, 26, 37, 42, 48, 59, 69 }

F(-1) timeout fail { }

F(-2) exception fail { 23, 25 }

Maxima

A grade { 14, 15, 16, 17, 19, 20, 21, 22, 27, 28, 29, 30, 31, 32, 34, 36, 39, 43, 47, 49, 50, 51, 52, 53, 55, 57, 61, 63, 66, 68, 70, 71 }

B grade { 33, 35, 41, 45, 54, 56, 58, 60, 62, 64 }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 18, 23, 24, 25, 26, 38, 40, 42, 44, 46, 59, 65, 67, 69 }

F(-1) timeout fail { }

F(-2) exception fail { 37, 48 }

Giac

A grade { 27, 28, 29, 30, 31, 35, 39, 41, 43, 49, 51, 56, 58, 60, 61, 62, 63, 64, 66, 68 }

B grade { 33, 34, 36, 45, 47, 50, 52, 54, 55, 57, 65, 67 }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 38, 40, 42, 44, 46, 69, 70, 71 }

F(-1) timeout fail { }

F(-2) exception fail { 26, 32, 37, 48, 53, 59 }

Mupad

A grade { }

B grade { 17, 19, 22, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 39, 41, 42, 43, 44, 45, 47, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 60, 61, 62, 63, 64, 65, 66, 67, 68, 70, 71 }

C grade { }

F normal fail { }

F(-1) timeout fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 18, 20, 21, 23, 24, 25, 26, 37, 38, 40, 46, 48, 59, 69 }

F(-2) exception fail { }

Sympy

A grade { 22, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 39, 41, 43, 45, 47, 48, 49, 50, 51, 52, 53, 55, 56, 57, 58, 66, 68 }

B grade { 54 }

C grade { 38, 40, 42, 44, 46 }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 23, 24, 25, 59, 60, 61, 62, 63, 64, 65, 67, 69 }

F(-1) timeout fail { 13, 70, 71 }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	147	147	149	227	0	325	0	0	0
N.S.	1	1.00	1.01	1.54	0.00	2.21	0.00	0.00	0.00
time (sec)	N/A	0.127	0.293	0.408	0.000	0.279	0.000	0.000	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	110	129	170	0	306	0	0	0
N.S.	1	1.00	1.17	1.55	0.00	2.78	0.00	0.00	0.00
time (sec)	N/A	0.081	0.179	0.317	0.000	0.278	0.000	0.000	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	75	75	110	97	0	285	0	0	0
N.S.	1	1.00	1.47	1.29	0.00	3.80	0.00	0.00	0.00
time (sec)	N/A	0.046	0.096	0.332	0.000	0.260	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	54	54	50	0	0	0	0	0	0
N.S.	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.079	0.025	0.000	0.000	0.000	0.000	0.000	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	77	77	236	0	0	0	0	0	0
N.S.	1	1.00	3.06	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.074	0.706	0.000	0.000	0.000	0.000	0.000	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	A	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	52	52	54	0	0	0	65	0	0
N.S.	1	1.00	1.04	0.00	0.00	0.00	1.25	0.00	0.00
time (sec)	N/A	0.035	0.034	0.000	0.000	0.000	2.803	0.000	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	54	54	49	53	50	53	63	78	41
N.S.	1	1.00	0.91	0.98	0.93	0.98	1.17	1.44	0.76
time (sec)	N/A	0.021	0.033	0.108	0.225	0.252	1.282	0.277	4.856

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	75	75	76	112	107	79	73	69	61
N.S.	1	1.00	1.01	1.49	1.43	1.05	0.97	0.92	0.81
time (sec)	N/A	0.033	0.042	0.056	0.231	0.252	2.592	0.300	5.490

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	38	43	25	41	41	44	33
N.S.	1	1.00	1.23	1.39	0.81	1.32	1.32	1.42	1.06
time (sec)	N/A	0.016	0.030	0.053	0.223	0.248	1.119	0.264	5.046

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	47	85	78	64	29	52	39
N.S.	1	1.00	1.00	1.81	1.66	1.36	0.62	1.11	0.83
time (sec)	N/A	0.023	0.023	0.053	0.230	0.244	1.575	0.283	4.883

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	31	35	113	64	86	48	66	36
N.S.	1	1.29	1.46	4.71	2.67	3.58	2.00	2.75	1.50
time (sec)	N/A	0.011	0.016	0.104	0.216	0.242	0.759	0.271	4.895

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	A	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	42	109	54	64	41	0	34
N.S.	1	1.00	1.11	2.87	1.42	1.68	1.08	0.00	0.89
time (sec)	N/A	0.023	0.028	0.052	0.236	0.240	2.271	0.000	5.156

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	40	40	43	145	86	102	37	82	42
N.S.	1	1.00	1.08	3.62	2.15	2.55	0.92	2.05	1.05
time (sec)	N/A	0.022	0.020	0.052	0.230	0.248	0.886	0.268	5.629

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	37	42	25	47	48	69	42
N.S.	1	1.00	1.19	1.35	0.81	1.52	1.55	2.23	1.35
time (sec)	N/A	0.017	0.024	0.057	0.227	0.267	0.995	0.278	4.802

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	65	53	173	129	113	95	103	61
N.S.	1	1.00	0.82	2.66	1.98	1.74	1.46	1.58	0.94
time (sec)	N/A	0.029	0.033	0.056	0.231	0.260	1.035	0.292	5.619

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	51	51	46	52	41	58	58	124	61
N.S.	1	1.00	0.90	1.02	0.80	1.14	1.14	2.43	1.20
time (sec)	N/A	0.029	0.026	0.055	0.234	0.246	1.105	0.295	5.165

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F(-2)	F	A	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	59	59	55	0	0	0	71	0	0
N.S.	1	1.00	0.93	0.00	0.00	0.00	1.20	0.00	0.00
time (sec)	N/A	0.035	0.047	0.000	0.000	0.000	2.731	0.000	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	A	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	202	202	112	150	0	89	48	0	0
N.S.	1	1.00	0.55	0.74	0.00	0.44	0.24	0.00	0.00
time (sec)	N/A	0.094	0.174	0.957	0.000	0.088	1.170	0.000	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	52	52	53	94	81	70	36	57	42
N.S.	1	1.00	1.02	1.81	1.56	1.35	0.69	1.10	0.81
time (sec)	N/A	0.025	0.038	0.106	0.227	0.273	1.755	0.258	5.315

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	A	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	86	86	113	104	0	56	41	0	0
N.S.	1	1.00	1.31	1.21	0.00	0.65	0.48	0.00	0.00
time (sec)	N/A	0.037	0.164	0.542	0.000	0.076	1.033	0.000	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	40	40	42	116	71	88	58	61	43
N.S.	1	1.00	1.05	2.90	1.78	2.20	1.45	1.52	1.08
time (sec)	N/A	0.026	0.022	0.115	0.227	0.253	2.975	0.267	5.714

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	F	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	165	165	96	146	0	0	42	0	24
N.S.	1	1.00	0.58	0.88	0.00	0.00	0.25	0.00	0.15
time (sec)	N/A	0.060	0.106	0.709	0.000	0.000	0.499	0.000	5.158

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	22	86	54	74	54	66	36
N.S.	1	1.00	0.48	1.87	1.17	1.61	1.17	1.43	0.78
time (sec)	N/A	0.024	0.038	0.100	0.219	0.242	4.951	0.281	5.115

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	A	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	91	91	96	111	0	57	42	0	27
N.S.	1	1.00	1.05	1.22	0.00	0.63	0.46	0.00	0.30
time (sec)	N/A	0.032	0.110	0.795	0.000	0.074	1.052	0.000	5.265

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	24	114	92	101	42	76	42
N.S.	1	1.00	0.57	2.71	2.19	2.40	1.00	1.81	1.00
time (sec)	N/A	0.029	0.034	0.114	0.227	0.245	1.339	0.270	6.321

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	A	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	181	181	114	171	0	97	44	0	0
N.S.	1	1.00	0.63	0.94	0.00	0.54	0.24	0.00	0.00
time (sec)	N/A	0.071	0.146	0.928	0.000	0.084	1.190	0.000	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	39	42	25	49	49	71	44
N.S.	1	1.00	1.26	1.35	0.81	1.58	1.58	2.29	1.42
time (sec)	N/A	0.017	0.034	0.100	0.220	0.255	1.459	0.273	4.798

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F(-2)	F	A	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	64	64	57	0	0	0	76	0	0
N.S.	1	1.00	0.89	0.00	0.00	0.00	1.19	0.00	0.00
time (sec)	N/A	0.233	0.046	0.000	0.000	0.000	3.103	0.000	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	85	84	127	117	87	82	80	73
N.S.	1	1.00	0.99	1.49	1.38	1.02	0.96	0.94	0.86
time (sec)	N/A	0.190	0.044	0.117	0.214	0.249	2.615	0.304	4.842

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	44	59	32	49	51	66	40
N.S.	1	1.00	1.16	1.55	0.84	1.29	1.34	1.74	1.05
time (sec)	N/A	0.203	0.032	0.128	0.221	0.249	1.120	0.293	5.231

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	52	52	57	98	89	72	36	62	51
N.S.	1	1.00	1.10	1.88	1.71	1.38	0.69	1.19	0.98
time (sec)	N/A	0.170	0.025	0.053	0.206	0.252	1.609	0.303	5.063

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	44	129	75	99	63	82	52
N.S.	1	1.00	1.02	3.00	1.74	2.30	1.47	1.91	1.21
time (sec)	N/A	0.121	0.033	0.057	0.220	0.262	1.733	0.276	5.090

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	A	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	52	113	59	73	49	0	47
N.S.	1	1.00	1.11	2.40	1.26	1.55	1.04	0.00	1.00
time (sec)	N/A	0.057	0.029	0.058	0.210	0.249	1.708	0.000	4.916

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	39	164	93	112	87	118	44
N.S.	1	1.00	1.03	4.32	2.45	2.95	2.29	3.11	1.16
time (sec)	N/A	0.162	0.036	0.056	0.244	0.242	2.003	0.283	4.869

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	54	46	56	28	57	58	75	51
N.S.	1	1.59	1.35	1.65	0.82	1.68	1.71	2.21	1.50
time (sec)	N/A	0.145	0.029	0.059	0.229	0.245	1.189	0.303	4.832

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	73	73	73	189	139	121	109	112	68
N.S.	1	1.00	1.00	2.59	1.90	1.66	1.49	1.53	0.93
time (sec)	N/A	0.176	0.034	0.059	0.256	0.254	1.247	0.275	5.225

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	58	58	54	65	52	67	71	134	67
N.S.	1	1.00	0.93	1.12	0.90	1.16	1.22	2.31	1.16
time (sec)	N/A	0.170	0.034	0.058	0.220	0.253	1.262	0.316	5.913

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	96	96	74	222	180	131	117	140	89
N.S.	1	1.00	0.77	2.31	1.88	1.36	1.22	1.46	0.93
time (sec)	N/A	0.195	0.050	0.059	0.230	0.253	1.305	0.280	5.502

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	85	85	88	0	0	0	0	0	0
N.S.	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.085	0.454	0.000	0.000	0.000	0.000	0.000	0.000

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	92	92	85	172	162	90	0	89	79
N.S.	1	1.00	0.92	1.87	1.76	0.98	0.00	0.97	0.86
time (sec)	N/A	0.078	0.085	0.955	0.310	0.266	0.000	0.284	5.337

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	64	125	49	58	0	85	61
N.S.	1	1.00	0.89	1.74	0.68	0.81	0.00	1.18	0.85
time (sec)	N/A	0.068	0.058	0.862	0.237	0.255	0.000	0.272	5.216

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	59	54	138	107	68	0	61	51
N.S.	1	1.00	0.92	2.34	1.81	1.15	0.00	1.03	0.86
time (sec)	N/A	0.063	0.065	0.862	0.328	0.277	0.000	0.291	5.247

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	35	89	31	38	0	44	31
N.S.	1	1.00	0.97	2.47	0.86	1.06	0.00	1.22	0.86
time (sec)	N/A	0.049	0.039	0.851	0.259	0.245	0.000	0.290	5.377

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	38	85	61	38	0	34	21
N.S.	1	1.00	1.41	3.15	2.26	1.41	0.00	1.26	0.78
time (sec)	N/A	0.042	0.115	0.858	0.308	0.248	0.000	0.270	5.824

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	37	170	0	80	0	70	38
N.S.	1	1.00	1.12	5.15	0.00	2.42	0.00	2.12	1.15
time (sec)	N/A	0.036	0.069	0.881	0.000	0.247	0.000	0.286	5.409

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	30	157	34	41	42	43	29
N.S.	1	1.00	1.00	5.23	1.13	1.37	1.40	1.43	0.97
time (sec)	N/A	0.052	0.061	0.847	0.318	0.248	2.257	0.272	5.505

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	60	60	58	216	0	130	0	114	61
N.S.	1	1.00	0.97	3.60	0.00	2.17	0.00	1.90	1.02
time (sec)	N/A	0.075	0.076	0.863	0.000	0.247	0.000	0.279	5.396

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	61	54	197	56	70	102	82	57
N.S.	1	1.00	0.89	3.23	0.92	1.15	1.67	1.34	0.93
time (sec)	N/A	0.067	0.082	0.868	0.317	0.258	2.259	0.273	5.203

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	61	61	53	0	0	0	0	0	0
N.S.	1	1.00	0.87	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.073	0.049	0.000	0.000	0.000	0.000	0.000	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	95	52	57	266	0	0	42
N.S.	1	1.00	2.07	1.13	1.24	5.78	0.00	0.00	0.91
time (sec)	N/A	0.047	0.097	0.174	0.221	0.278	0.000	0.000	5.752

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F	A	B	F(-1)	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	46	46	93	0	60	334	0	0	40
N.S.	1	1.00	2.02	0.00	1.30	7.26	0.00	0.00	0.87
time (sec)	N/A	0.056	0.127	0.000	0.237	0.287	0.000	0.000	5.626

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. The column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [42] had the largest ratio of [.8750000000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	8	7	1.00	10	0.700
2	A	7	6	1.00	10	0.600
3	A	6	6	1.00	8	0.750
4	A	14	8	1.00	10	0.800
5	A	6	6	1.00	10	0.600
6	A	8	8	1.00	10	0.800
7	A	20	9	1.00	20	0.450
8	A	17	9	1.00	20	0.450
9	A	11	8	1.00	18	0.444
10	A	8	6	1.00	12	0.500
11	A	17	9	1.00	20	0.450
12	A	12	8	1.00	20	0.400
13	A	23	11	1.00	20	0.550
14	A	4	3	1.00	10	0.300
15	A	4	3	1.00	10	0.300
16	A	4	3	1.00	8	0.375
17	A	3	3	1.00	6	0.500
18	A	7	6	1.00	10	0.600
19	A	5	5	1.00	10	0.500
20	A	6	5	1.00	10	0.500
21	A	7	5	1.00	10	0.500
22	A	3	3	1.00	4	0.750
23	A	7	6	1.00	10	0.600
24	A	7	6	1.00	10	0.600

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
25	A	7	7	1.00	10	0.700
26	A	4	4	1.00	10	0.400
27	A	4	4	1.00	10	0.400
28	A	7	7	1.00	10	0.700
29	A	3	3	1.00	10	0.300
30	A	6	6	1.00	8	0.750
31	A	5	5	1.29	6	0.833
32	A	6	6	1.00	10	0.600
33	A	5	5	1.00	10	0.500
34	A	3	3	1.00	10	0.300
35	A	6	6	1.00	10	0.600
36	A	5	4	1.00	10	0.400
37	A	4	4	1.00	12	0.333
38	A	8	8	1.00	12	0.667
39	A	6	6	1.00	12	0.500
40	A	5	5	1.00	12	0.417
41	A	6	6	1.00	10	0.600
42	A	7	7	1.00	8	0.875
43	A	6	6	1.00	12	0.500
44	A	5	5	1.00	12	0.417
45	A	6	6	1.00	12	0.500
46	A	7	7	1.00	12	0.583
47	A	3	3	1.00	12	0.250
48	A	5	4	1.00	12	0.333
49	A	8	7	1.00	12	0.583
50	A	4	3	1.00	12	0.250
51	A	7	6	1.00	12	0.500
52	A	6	5	1.00	10	0.500
53	A	7	6	1.00	8	0.750
54	A	6	5	1.00	12	0.417
55	A	4	3	1.59	12	0.250
56	A	7	6	1.00	12	0.500
57	A	6	4	1.00	12	0.333
58	A	8	6	1.00	12	0.500
59	A	4	3	1.00	23	0.130

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
60	A	9	7	1.00	21	0.333
61	A	6	5	1.00	21	0.238
62	A	7	7	1.00	21	0.333
63	A	3	3	1.00	21	0.143
64	A	5	5	1.00	19	0.263
65	A	7	7	1.00	18	0.389
66	A	4	4	1.00	21	0.190
67	A	7	6	1.00	21	0.286
68	A	7	5	1.00	21	0.238
69	A	8	8	1.00	19	0.421
70	A	6	6	1.00	12	0.500
71	A	6	6	1.00	14	0.429

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^3 \operatorname{csch}^{-1}(a + bx) dx$	46
3.2	$\int x^2 \operatorname{csch}^{-1}(a + bx) dx$	52
3.3	$\int x \operatorname{csch}^{-1}(a + bx) dx$	58
3.4	$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x} dx$	63
3.5	$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^2} dx$	70
3.6	$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^3} dx$	75
3.7	$\int (e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$	82
3.8	$\int (e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$	95
3.9	$\int (e + fx) (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$	105
3.10	$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$	112
3.11	$\int \frac{(a+b \operatorname{csch}^{-1}(c+dx))^2}{e+fx} dx$	118
3.12	$\int \frac{(a+b \operatorname{csch}^{-1}(c+dx))^2}{(e+fx)^2} dx$	129
3.13	$\int \frac{(a+b \operatorname{csch}^{-1}(c+dx))^2}{(e+fx)^3} dx$	138
3.14	$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx$	153
3.15	$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx$	157
3.16	$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx$	161
3.17	$\int \operatorname{csch}^{-1}(\sqrt{x}) dx$	165
3.18	$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx$	169
3.19	$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx$	173
3.20	$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx$	178
3.21	$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx$	183

3.22	$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx$	188
3.23	$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx$	192
3.24	$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx$	197
3.25	$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx$	201
3.26	$\int e^{\operatorname{csch}^{-1}(ax)} x^m dx$	206
3.27	$\int e^{\operatorname{csch}^{-1}(ax)} x^4 dx$	210
3.28	$\int e^{\operatorname{csch}^{-1}(ax)} x^3 dx$	214
3.29	$\int e^{\operatorname{csch}^{-1}(ax)} x^2 dx$	219
3.30	$\int e^{\operatorname{csch}^{-1}(ax)} x dx$	223
3.31	$\int e^{\operatorname{csch}^{-1}(ax)} dx$	228
3.32	$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x} dx$	232
3.33	$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^2} dx$	237
3.34	$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^3} dx$	242
3.35	$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^4} dx$	246
3.36	$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^5} dx$	251
3.37	$\int e^{\operatorname{csch}^{-1}(ax^2)} x^m dx$	255
3.38	$\int e^{\operatorname{csch}^{-1}(ax^2)} x^4 dx$	259
3.39	$\int e^{\operatorname{csch}^{-1}(ax^2)} x^3 dx$	265
3.40	$\int e^{\operatorname{csch}^{-1}(ax^2)} x^2 dx$	270
3.41	$\int e^{\operatorname{csch}^{-1}(ax^2)} x dx$	275
3.42	$\int e^{\operatorname{csch}^{-1}(ax^2)} dx$	280
3.43	$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x} dx$	286
3.44	$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^2} dx$	291
3.45	$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^3} dx$	296
3.46	$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^4} dx$	301
3.47	$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^5} dx$	307
3.48	$\int e^{2\operatorname{csch}^{-1}(ax)} x^m dx$	311
3.49	$\int e^{2\operatorname{csch}^{-1}(ax)} x^4 dx$	315
3.50	$\int e^{2\operatorname{csch}^{-1}(ax)} x^3 dx$	320
3.51	$\int e^{2\operatorname{csch}^{-1}(ax)} x^2 dx$	324
3.52	$\int e^{2\operatorname{csch}^{-1}(ax)} x dx$	329
3.53	$\int e^{2\operatorname{csch}^{-1}(ax)} dx$	334
3.54	$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x} dx$	339
3.55	$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^2} dx$	344
3.56	$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^3} dx$	348
3.57	$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^4} dx$	354

3.58	$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^5} dx$	358
3.59	$\int \frac{e^{\operatorname{csch}^{-1}(cx)}(dx)^m}{1+c^2x^2} dx$	364
3.60	$\int \frac{e^{\operatorname{csch}^{-1}(cx)}x^5}{1+c^2x^2} dx$	368
3.61	$\int \frac{e^{\operatorname{csch}^{-1}(cx)}x^4}{1+c^2x^2} dx$	374
3.62	$\int \frac{e^{\operatorname{csch}^{-1}(cx)}x^3}{1+c^2x^2} dx$	379
3.63	$\int \frac{e^{\operatorname{csch}^{-1}(cx)}x^2}{1+c^2x^2} dx$	384
3.64	$\int \frac{e^{\operatorname{csch}^{-1}(cx)}x}{1+c^2x^2} dx$	388
3.65	$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{1+c^2x^2} dx$	393
3.66	$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x(1+c^2x^2)} dx$	398
3.67	$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^2(1+c^2x^2)} dx$	402
3.68	$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^3(1+c^2x^2)} dx$	407
3.69	$\int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{a}{b}+dx} dx$	412
3.70	$\int x^3 \operatorname{csch}^{-1}(a+bx^4) dx$	417
3.71	$\int x^{-1+n} \operatorname{csch}^{-1}(a+bx^n) dx$	422

3.1 $\int x^3 \operatorname{csch}^{-1}(a + bx) dx$

Optimal result	46
Rubi [A] (verified)	46
Mathematica [A] (verified)	49
Maple [A] (verified)	49
Fricas [B] (verification not implemented)	50
Sympy [F]	50
Maxima [F]	50
Giac [F]	51
Mupad [F(-1)]	51

Optimal result

Integrand size = 10, antiderivative size = 147

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx = -\frac{(2 - 17a^2)(a + bx)\sqrt{1 + \frac{1}{(a+bx)^2}}}{12b^4} + \frac{x^2(a + bx)\sqrt{1 + \frac{1}{(a+bx)^2}}}{12b^2}$$

$$- \frac{a(a + bx)^2\sqrt{1 + \frac{1}{(a+bx)^2}}}{3b^4} - \frac{a^4 \operatorname{csch}^{-1}(a + bx)}{4b^4}$$

$$+ \frac{1}{4}x^4 \operatorname{csch}^{-1}(a + bx) + \frac{a(1 - 2a^2) \operatorname{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx)^2}}\right)}{2b^4}$$

[Out] $-1/4*a^4*\operatorname{arccsch}(b*x+a)/b^4+1/4*x^4*\operatorname{arccsch}(b*x+a)+1/2*a*(-2*a^2+1)*\operatorname{arctanh}((1+1/(b*x+a)^2)^{(1/2)})/b^4-1/12*(-17*a^2+2)*(b*x+a)*(1+1/(b*x+a)^2)^{(1/2)}/b^4+1/12*x^2*(b*x+a)*(1+1/(b*x+a)^2)^{(1/2)}/b^2-1/3*a*(b*x+a)^2*(1+1/(b*x+a)^2)^{(1/2)}/b^4$

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {6457, 5577, 3867, 4133, 3855, 3852, 8}

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx = -\frac{a^4 \operatorname{csch}^{-1}(a + bx)}{4b^4} + \frac{(1 - 2a^2) a \operatorname{arctanh}\left(\sqrt{\frac{1}{(a+bx)^2} + 1}\right)}{2b^4}$$

$$- \frac{(2 - 17a^2)(a + bx)\sqrt{\frac{1}{(a+bx)^2} + 1}}{12b^4} - \frac{a(a + bx)^2\sqrt{\frac{1}{(a+bx)^2} + 1}}{3b^4}$$

$$+ \frac{x^2(a + bx)\sqrt{\frac{1}{(a+bx)^2} + 1}}{12b^2} + \frac{1}{4}x^4 \operatorname{csch}^{-1}(a + bx)$$

[In] Int[x^3*ArcCsch[a + b*x],x]

[Out]
$$-1/12*((2 - 17*a^2)*(a + b*x)*\text{Sqrt}[1 + (a + b*x)^{-2}])/b^4 + (x^2*(a + b*x)*\text{Sqrt}[1 + (a + b*x)^{-2}])/(12*b^2) - (a*(a + b*x)^2*\text{Sqrt}[1 + (a + b*x)^{-2}])/(3*b^4) - (a^4*\text{ArcCsch}[a + b*x])/(4*b^4) + (x^4*\text{ArcCsch}[a + b*x])/4 + (a*(1 - 2*a^2)*\text{ArcTanh}[\text{Sqrt}[1 + (a + b*x)^{-2}]])/(2*b^4)$$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3852

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3867

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_), x_Symbol] := Simp[(-b^2)*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n - 2)/(d*(n - 1))), x] + Dist[1/(n - 1), Int[(a + b*Csc[c + d*x])^(n - 3)*Simp[a^3*(n - 1) + (b*(b^2*(n - 2) + 3*a^2*(n - 1)))*Csc[c + d*x] + (a*b^2*(3*n - 4))*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 2] && IntegerQ[2*n]

Rule 4133

Int[((A_.) + csc[(e_.) + (f_.)*(x_)])*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.))*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[(-b)*C*Csc[e + f*x]*(Cot[e + f*x]/(2*f)), x] + Dist[1/2, Int[Simp[2*A*a + (2*B*a + b*(2*A + C))*Csc[e + f*x] + 2*(a*C + B*b)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x]

Rule 5577

Int[Coth[(c_.) + (d_.)*(x_)]*Csch[(c_.) + (d_.)*(x_)]*(Csch[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rule 6457

```

Int[((a_.) + ArcCsch[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csch[x]*Co
th[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int x \coth(x) \operatorname{csch}(x) (-a + \operatorname{csch}(x))^3 dx, x, \operatorname{csch}^{-1}(a + bx)\right)}{b^4} \\
&= \frac{1}{4} x^4 \operatorname{csch}^{-1}(a + bx) - \frac{\text{Subst}\left(\int (-a + \operatorname{csch}(x))^4 dx, x, \operatorname{csch}^{-1}(a + bx)\right)}{4b^4} \\
&= \frac{x^2(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}}{12b^2} + \frac{1}{4} x^4 \operatorname{csch}^{-1}(a + bx) \\
&\quad - \frac{\text{Subst}\left(\int (-a + \operatorname{csch}(x)) (-3a^3 - (2 - 9a^2) \operatorname{csch}(x) - 8a \operatorname{csch}^2(x)) dx, x, \operatorname{csch}^{-1}(a + bx)\right)}{12b^4} \\
&= \frac{x^2(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}}{12b^2} - \frac{a(a + bx)^2 \sqrt{1 + \frac{1}{(a+bx)^2}}}{3b^4} + \frac{1}{4} x^4 \operatorname{csch}^{-1}(a + bx) \\
&\quad - \frac{\text{Subst}\left(\int (6a^4 + 12a(1 - 2a^2) \operatorname{csch}(x) - 2(2 - 17a^2) \operatorname{csch}^2(x)) dx, x, \operatorname{csch}^{-1}(a + bx)\right)}{24b^4} \\
&= \frac{x^2(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}}{12b^2} - \frac{a(a + bx)^2 \sqrt{1 + \frac{1}{(a+bx)^2}}}{3b^4} - \frac{a^4 \operatorname{csch}^{-1}(a + bx)}{4b^4} \\
&\quad + \frac{1}{4} x^4 \operatorname{csch}^{-1}(a + bx) + \frac{(2 - 17a^2) \text{Subst}\left(\int \operatorname{csch}^2(x) dx, x, \operatorname{csch}^{-1}(a + bx)\right)}{12b^4} \\
&\quad - \frac{(a(1 - 2a^2)) \text{Subst}\left(\int \operatorname{csch}(x) dx, x, \operatorname{csch}^{-1}(a + bx)\right)}{2b^4} \\
&= \frac{x^2(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}}{12b^2} - \frac{a(a + bx)^2 \sqrt{1 + \frac{1}{(a+bx)^2}}}{3b^4} - \frac{a^4 \operatorname{csch}^{-1}(a + bx)}{4b^4} \\
&\quad + \frac{1}{4} x^4 \operatorname{csch}^{-1}(a + bx) + \frac{a(1 - 2a^2) \operatorname{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx)^2}}\right)}{2b^4} \\
&\quad - \frac{(i(2 - 17a^2)) \text{Subst}\left(\int 1 dx, x, -i(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}\right)}{12b^4} \\
&= -\frac{(2 - 17a^2)(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}}{12b^4} + \frac{x^2(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}}{12b^2} - \frac{a(a + bx)^2 \sqrt{1 + \frac{1}{(a+bx)^2}}}{3b^4} \\
&\quad - \frac{a^4 \operatorname{csch}^{-1}(a + bx)}{4b^4} + \frac{1}{4} x^4 \operatorname{csch}^{-1}(a + bx) + \frac{a(1 - 2a^2) \operatorname{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx)^2}}\right)}{2b^4}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.01

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx$$

$$= \frac{\sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}}(-2a + 13a^3 - 2bx + 9a^2bx - 3ab^2x^2 + b^3x^3) + 3b^4x^4 \operatorname{csch}^{-1}(a + bx) - 3a^4 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}{12b^4}$$

`[In] Integrate[x^3*ArcCsch[a + b*x], x]`

```
[Out] (Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]*(-2*a + 13*a^3 - 2*b*x + 9
*a^2*b*x - 3*a*b^2*x^2 + b^3*x^3) + 3*b^4*x^4*ArcCsch[a + b*x] - 3*a^4*ArcS
inh[(a + b*x)^(-1)] + 6*a*(1 - 2*a^2)*Log[(a + b*x)*(1 + Sqrt[(1 + a^2 + 2*
a*b*x + b^2*x^2)/(a + b*x)^2])])/(12*b^4)
```

Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.54

method	result
derivativedivides	$\frac{\operatorname{arccsch}(bx+a)a^4}{4} - \operatorname{arccsch}(bx+a)a^3(bx+a) + \frac{3 \operatorname{arccsch}(bx+a)a^2(bx+a)^2}{2} - \operatorname{arccsch}(bx+a)a(bx+a)^3 + \frac{\operatorname{arccsch}(bx+a)(bx+a)^4}{4}$
default	$\frac{\operatorname{arccsch}(bx+a)a^4}{4} - \operatorname{arccsch}(bx+a)a^3(bx+a) + \frac{3 \operatorname{arccsch}(bx+a)a^2(bx+a)^2}{2} - \operatorname{arccsch}(bx+a)a(bx+a)^3 + \frac{\operatorname{arccsch}(bx+a)(bx+a)^4}{4}$
parts	$\frac{x^4 \operatorname{arccsch}(bx+a)}{4} - \frac{\sqrt{b^2x^2+2abx+a^2+1} \left(3a^4 \operatorname{arctanh}\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2+1}}\right) \sqrt{b^2-x^2} \sqrt{b^2x^2+2abx+a^2+1} b^2 \sqrt{b^2+4a^2} \right)}{\sqrt{b^2x^2+2abx+a^2+1}}$

`[In] int(x^3*arccsch(b*x+a), x, method=_RETURNVERBOSE)`

```
[Out] 1/b^4*(1/4*arccsch(b*x+a)*a^4-arccsch(b*x+a)*a^3*(b*x+a)+3/2*arccsch(b*x+a)
*a^2*(b*x+a)^2-arccsch(b*x+a)*a*(b*x+a)^3+1/4*arccsch(b*x+a)*(b*x+a)^4-1/12
*((b*x+a)^2+1)^(1/2)*(3*a^4*arctanh(1/((b*x+a)^2+1)^(1/2))+12*a^3*arcsinh(b
*x+a)-18*a^2*((b*x+a)^2+1)^(1/2)+6*a*(b*x+a)*((b*x+a)^2+1)^(1/2)-(b*x+a)^2*
((b*x+a)^2+1)^(1/2)-6*a*arcsinh(b*x+a)+2*((b*x+a)^2+1)^(1/2))/(((b*x+a)^2+1
)/(b*x+a)^2)^(1/2)/(b*x+a))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 325 vs. 2(127) = 254.

Time = 0.28 (sec) , antiderivative size = 325, normalized size of antiderivative = 2.21

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx$$

$$= \frac{3b^4 x^4 \log\left(\frac{(bx+a)\sqrt{\frac{b^2 x^2 + 2abx + a^2 + 1}{b^2 x^2 + 2abx + a^2}} + 1}{bx+a}\right) - 3a^4 \log\left(-bx + (bx+a)\sqrt{\frac{b^2 x^2 + 2abx + a^2 + 1}{b^2 x^2 + 2abx + a^2}} - a + 1\right) + 3a^4 \log(-bx$$

[In] integrate(x^3*arccsch(b*x+a),x, algorithm="fricas")

[Out] 1/12*(3*b^4*x^4*log(((b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)) - 3*a^4*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a + 1) + 3*a^4*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a - 1) + 6*(2*a^3 - a)*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a) + (b^3*x^3 - 3*a*b^2*x^2 + 13*a^3 + (9*a^2 - 2)*b*x - 2*a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)))/b^4

Sympy [F]

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx = \int x^3 \operatorname{acsch}(a + bx) dx$$

[In] integrate(x**3*acsch(b*x+a),x)

[Out] Integral(x**3*acsch(a + b*x), x)

Maxima [F]

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx = \int x^3 \operatorname{arcsch}(bx + a) dx$$

[In] integrate(x^3*arccsch(b*x+a),x, algorithm="maxima")

[Out] -1/2*(-I*a^3 + I*a)*(log(I*(b^2*x + a*b)/b + 1) - log(-I*(b^2*x + a*b)/b + 1))/b^4 + 1/8*(2*b^4*x^4*log(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) + 1) + b^2*x^2 - 6*a*b*x - (a^4 - 6*a^2 + 1)*log(b^2*x^2 + 2*a*b*x + a^2 + 1) - 2*(b^4*x^4 - a^4)*log(b*x + a))/b^4 + integrate(1/4*(b^2*x^5 + a*b*x^4)/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 + 1)^(3/2) + 1), x)

Giac [F]

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx = \int x^3 \operatorname{arcsch}(bx + a) dx$$

[In] integrate(x^3*arccsch(b*x+a),x, algorithm="giac")

[Out] integrate(x^3*arccsch(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int x^3 \operatorname{csch}^{-1}(a + bx) dx = \int x^3 \operatorname{asinh}\left(\frac{1}{a + bx}\right) dx$$

[In] int(x^3*asinh(1/(a + b*x)),x)

[Out] int(x^3*asinh(1/(a + b*x)), x)

3.2 $\int x^2 \operatorname{csch}^{-1}(a + bx) dx$

Optimal result	52
Rubi [A] (verified)	52
Mathematica [A] (verified)	54
Maple [A] (verified)	55
Fricas [B] (verification not implemented)	55
Sympy [F]	56
Maxima [F]	56
Giac [F]	56
Mupad [F(-1)]	57

Optimal result

Integrand size = 10, antiderivative size = 110

$$\int x^2 \operatorname{csch}^{-1}(a + bx) dx = -\frac{5a(a + bx)\sqrt{1 + \frac{1}{(a+bx)^2}}}{6b^3} + \frac{x(a + bx)\sqrt{1 + \frac{1}{(a+bx)^2}}}{6b^2}$$

$$+ \frac{a^3 \operatorname{csch}^{-1}(a + bx)}{3b^3} + \frac{1}{3}x^3 \operatorname{csch}^{-1}(a + bx)$$

$$- \frac{(1 - 6a^2) \operatorname{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx)^2}}\right)}{6b^3}$$

[Out] $1/3*a^3*\operatorname{arccsch}(b*x+a)/b^3+1/3*x^3*\operatorname{arccsch}(b*x+a)-1/6*(-6*a^2+1)*\operatorname{arctanh}((1+1/(b*x+a)^2)^{(1/2)})/b^3-5/6*a*(b*x+a)*(1+1/(b*x+a)^2)^{(1/2)}/b^3+1/6*x*(b*x+a)*(1+1/(b*x+a)^2)^{(1/2)}/b^2$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6457, 5577, 3867, 3855, 3852, 8}

$$\int x^2 \operatorname{csch}^{-1}(a + bx) dx = \frac{a^3 \operatorname{csch}^{-1}(a + bx)}{3b^3} - \frac{(1 - 6a^2) \operatorname{arctanh}\left(\sqrt{\frac{1}{(a+bx)^2} + 1}\right)}{6b^3}$$

$$- \frac{5a(a + bx)\sqrt{\frac{1}{(a+bx)^2} + 1}}{6b^3}$$

$$+ \frac{x(a + bx)\sqrt{\frac{1}{(a+bx)^2} + 1}}{6b^2} + \frac{1}{3}x^3 \operatorname{csch}^{-1}(a + bx)$$

[In] Int[x^2*ArcCsch[a + b*x],x]

[Out] $(-5*a*(a + b*x)*\sqrt{1 + (a + b*x)^{-2}})/(6*b^3) + (x*(a + b*x)*\sqrt{1 + (a + b*x)^{-2}})/(6*b^2) + (a^3*\text{ArcCsch}[a + b*x])/(3*b^3) + (x^3*\text{ArcCsch}[a + b*x])/3 - ((1 - 6*a^2)*\text{ArcTanh}[\sqrt{1 + (a + b*x)^{-2}}])/(6*b^3)$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3852

Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3867

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^(n_), x_Symbol] := Simp[(-b^2)*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n - 2)/(d*(n - 1))), x] + Dist[1/(n - 1), Int[(a + b*Csc[c + d*x])^(n - 3)*Simp[a^3*(n - 1) + (b*(b^2*(n - 2) + 3*a^2*(n - 1)))*Csc[c + d*x] + (a*b^2*(3*n - 4))*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 2] && IntegerQ[2*n]

Rule 5577

Int[Coth[(c_.) + (d_.)*(x_.)]*Csch[(c_.) + (d_.)*(x_.)]*(Csch[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^(n_.)*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rule 6457

Int[((a_.) + ArcCsch[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csch[x]*Coth[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int x \coth(x) \text{csch}(x) (-a + \text{csch}(x))^2 dx, x, \text{csch}^{-1}(a + bx)\right)}{b^3} \\
&= \frac{1}{3} x^3 \text{csch}^{-1}(a + bx) - \frac{\text{Subst}\left(\int (-a + \text{csch}(x))^3 dx, x, \text{csch}^{-1}(a + bx)\right)}{3b^3} \\
&= \frac{x(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}}{6b^2} + \frac{1}{3} x^3 \text{csch}^{-1}(a + bx) \\
&\quad - \frac{\text{Subst}\left(\int (-2a^3 - (1 - 6a^2) \text{csch}(x) - 5a \text{csch}^2(x)) dx, x, \text{csch}^{-1}(a + bx)\right)}{6b^3} \\
&= \frac{x(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}}{6b^2} + \frac{a^3 \text{csch}^{-1}(a + bx)}{3b^3} + \frac{1}{3} x^3 \text{csch}^{-1}(a + bx) \\
&\quad + \frac{(5a) \text{Subst}\left(\int \text{csch}^2(x) dx, x, \text{csch}^{-1}(a + bx)\right)}{6b^3} \\
&\quad + \frac{(1 - 6a^2) \text{Subst}\left(\int \text{csch}(x) dx, x, \text{csch}^{-1}(a + bx)\right)}{6b^3} \\
&= \frac{x(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}}{6b^2} + \frac{a^3 \text{csch}^{-1}(a + bx)}{3b^3} \\
&\quad + \frac{1}{3} x^3 \text{csch}^{-1}(a + bx) - \frac{(1 - 6a^2) \text{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx)^2}}\right)}{6b^3} \\
&\quad - \frac{(5ia) \text{Subst}\left(\int 1 dx, x, -i(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}\right)}{6b^3} \\
&= -\frac{5a(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}}{6b^3} + \frac{x(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}}{6b^2} + \frac{a^3 \text{csch}^{-1}(a + bx)}{3b^3} \\
&\quad + \frac{1}{3} x^3 \text{csch}^{-1}(a + bx) - \frac{(1 - 6a^2) \text{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx)^2}}\right)}{6b^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.17

$$\begin{aligned}
&\int x^2 \text{csch}^{-1}(a + bx) dx \\
&= \frac{(-5a^2 - 4abx + b^2x^2) \sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}} + 2b^3x^3 \text{csch}^{-1}(a + bx) + 2a^3 \text{arcsinh}\left(\frac{1}{a+bx}\right) + (-1 + 6a^2) \log\left((a + \right.}{6b^3}
\end{aligned}$$

[In] Integrate[x^2*ArcCsch[a + b*x], x]

```
[Out] ((-5*a^2 - 4*a*b*x + b^2*x^2)*Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2] + 2*b^3*x^3*ArcCsch[a + b*x] + 2*a^3*ArcSinh[(a + b*x)^(-1)] + (-1 + 6*a^2)*Log[(a + b*x)*(1 + Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2])])/(6*b^3)
```

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.55

method	result
derivativedivides	$-\frac{\operatorname{arccsch}(bx+a)a^3}{3} + \operatorname{arccsch}(bx+a)a^2(bx+a) - \operatorname{arccsch}(bx+a)a(bx+a)^2 + \frac{\operatorname{arccsch}(bx+a)(bx+a)^3}{3} + \frac{\sqrt{(bx+a)^2+1} \left(2a^3 \operatorname{arctanh} \left(\frac{1}{\sqrt{b^2x^2+2abx+a^2+1}} \right) \right)}{b^3}$
default	$-\frac{\operatorname{arccsch}(bx+a)a^3}{3} + \operatorname{arccsch}(bx+a)a^2(bx+a) - \operatorname{arccsch}(bx+a)a(bx+a)^2 + \frac{\operatorname{arccsch}(bx+a)(bx+a)^3}{3} + \frac{\sqrt{(bx+a)^2+1} \left(2a^3 \operatorname{arctanh} \left(\frac{1}{\sqrt{b^2x^2+2abx+a^2+1}} \right) \right)}{b^3}$
parts	$\frac{x^3 \operatorname{arccsch}(bx+a)}{3} + \frac{\sqrt{b^2x^2+2abx+a^2+1} \left(2a^3 \operatorname{arctanh} \left(\frac{1}{\sqrt{b^2x^2+2abx+a^2+1}} \right) \right) \sqrt{b^2+6} \ln \left(\frac{b^2x + \sqrt{b^2x^2+2abx+a^2+1} \sqrt{b^2+6}}{\sqrt{b^2+6}} \right)}{6b^3 \sqrt{b^2x^2+2abx+a^2+1}}$

```
[In] int(x^2*arccsch(b*x+a),x,method=_RETURNVERBOSE)
```

```
[Out] 1/b^3*(-1/3*arccsch(b*x+a)*a^3+arccsch(b*x+a)*a^2*(b*x+a)-arccsch(b*x+a)*a*(b*x+a)^2+1/3*arccsch(b*x+a)*(b*x+a)^3+1/6*((b*x+a)^2+1)^(1/2)*(2*a^3*arctanh(1/((b*x+a)^2+1)^(1/2))+6*a^2*arcsinh(b*x+a)-6*a*((b*x+a)^2+1)^(1/2)+(b*x+a)*((b*x+a)^2+1)^(1/2)-arcsinh(b*x+a))/(((b*x+a)^2+1)/(b*x+a)^2)^(1/2)/(b*x+a))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 306 vs. 2(94) = 188.

Time = 0.28 (sec) , antiderivative size = 306, normalized size of antiderivative = 2.78

$$\int x^2 \operatorname{csch}^{-1}(a + bx) dx$$

$$= \frac{2b^3x^3 \log \left(\frac{(bx+a) \sqrt{\frac{b^2x^2+2abx+a^2+1}{b^2x^2+2abx+a^2}} + 1}{bx+a} \right) + 2a^3 \log \left(-bx + (bx+a) \sqrt{\frac{b^2x^2+2abx+a^2+1}{b^2x^2+2abx+a^2}} - a + 1 \right) - 2a^3 \log \left(-bx + (bx+a) \sqrt{\frac{b^2x^2+2abx+a^2+1}{b^2x^2+2abx+a^2}} + a + 1 \right)}{6b^3}$$

```
[In] integrate(x^2*arccsch(b*x+a),x, algorithm="fricas")
```

```
[Out] 1/6*(2*b^3*x^3*log(((b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)) + 2*a^3*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a + 1) - 2*a^3*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) + a + 1))/(6*b^3)
```

$$+ 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a + 1) - 2*a^3*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a - 1) - (6*a^2 - 1)*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a) + (b^2*x^2 - 4*a*b*x - 5*a^2)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)))/b^3$$

Sympy [F]

$$\int x^2 \operatorname{csch}^{-1}(a + bx) dx = \int x^2 \operatorname{acsch}(a + bx) dx$$

[In] integrate(x**2*acsch(b*x+a),x)

[Out] Integral(x**2*acsch(a + b*x), x)

Maxima [F]

$$\int x^2 \operatorname{csch}^{-1}(a + bx) dx = \int x^2 \operatorname{arcsch}(bx + a) dx$$

[In] integrate(x^2*arccsch(b*x+a),x, algorithm="maxima")

[Out] -1/6*(3*I*a^2 - I)*(log(I*(b^2*x + a*b)/b + 1) - log(-I*(b^2*x + a*b)/b + 1))/b^3 + 1/6*(2*b^3*x^3*log(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) + 1) + 2*b*x + (a^3 - 3*a)*log(b^2*x^2 + 2*a*b*x + a^2 + 1) - 2*(b^3*x^3 + a^3)*log(b*x + a))/b^3 + integrate(1/3*(b^2*x^4 + a*b*x^3)/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 + 1)^(3/2) + 1), x)

Giac [F]

$$\int x^2 \operatorname{csch}^{-1}(a + bx) dx = \int x^2 \operatorname{arcsch}(bx + a) dx$$

[In] integrate(x^2*arccsch(b*x+a),x, algorithm="giac")

[Out] integrate(x^2*arccsch(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{csch}^{-1}(a + bx) dx = \int x^2 \operatorname{asinh}\left(\frac{1}{a + bx}\right) dx$$

```
[In] int(x^2*asinh(1/(a + b*x)),x)
```

```
[Out] int(x^2*asinh(1/(a + b*x)), x)
```

3.3 $\int x \operatorname{csch}^{-1}(a + bx) dx$

Optimal result	58
Rubi [A] (verified)	58
Mathematica [A] (verified)	60
Maple [A] (verified)	60
Fricas [B] (verification not implemented)	61
Sympy [F]	61
Maxima [F]	61
Giac [F]	62
Mupad [F(-1)]	62

Optimal result

Integrand size = 8, antiderivative size = 75

$$\int x \operatorname{csch}^{-1}(a + bx) dx = \frac{(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}}{2b^2} - \frac{a^2 \operatorname{csch}^{-1}(a + bx)}{2b^2} + \frac{1}{2} x^2 \operatorname{csch}^{-1}(a + bx) - \frac{a \operatorname{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx)^2}}\right)}{b^2}$$

[Out] $-1/2*a^2*\operatorname{arccsch}(b*x+a)/b^2+1/2*x^2*\operatorname{arccsch}(b*x+a)-a*\operatorname{arctanh}((1+1/(b*x+a)^2)^{(1/2)})/b^2+1/2*(b*x+a)*(1+1/(b*x+a)^2)^{(1/2)}/b^2$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6457, 5577, 3858, 3855, 3852, 8}

$$\int x \operatorname{csch}^{-1}(a + bx) dx = -\frac{a^2 \operatorname{csch}^{-1}(a + bx)}{2b^2} - \frac{a \operatorname{arctanh}\left(\sqrt{\frac{1}{(a+bx)^2} + 1}\right)}{b^2} + \frac{(a + bx) \sqrt{\frac{1}{(a+bx)^2} + 1}}{2b^2} + \frac{1}{2} x^2 \operatorname{csch}^{-1}(a + bx)$$

[In] $\operatorname{Int}[x*\operatorname{ArcCsch}[a + b*x], x]$

[Out] $((a + b*x)*\operatorname{Sqrt}[1 + (a + b*x)^{-2}])/(2*b^2) - (a^2*\operatorname{ArcCsch}[a + b*x])/(2*b^2) + (x^2*\operatorname{ArcCsch}[a + b*x])/2 - (a*\operatorname{ArcTanh}[\operatorname{Sqrt}[1 + (a + b*x)^{-2}]])/b^2$

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 3852

`Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

Rule 3855

`Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

Rule 3858

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_))^(n_), x_Symbol] := Simp[a^2*x, x] + (Dist[2*a*b, Int[Csc[c + d*x], x], x] + Dist[b^2, Int[Csc[c + d*x]^2, x], x]) /; FreeQ[{a, b, c, d}, x]`

Rule 5577

`Int[Coth[(c_.) + (d_.)*(x_)]*Csch[(c_.) + (d_.)*(x_)]*(Csch[(c_.) + (d_.)*(x_)]*(b_.) + (a_))^(n_)*((e_.) + (f_.)*(x_))^(m_), x_Symbol] := Simp[(-(e + f*x)^m)*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

Rule 6457

`Int[((a_.) + ArcCsch[(c_.) + (d_.)*(x_)]*(b_.))^(p_)*((e_.) + (f_.)*(x_))^(m_), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csch[x]*Coth[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int x \coth(x) \operatorname{csch}(x) (-a + \operatorname{csch}(x)) dx, x, \operatorname{csch}^{-1}(a + bx)\right)}{b^2} \\
 &= \frac{1}{2} x^2 \operatorname{csch}^{-1}(a + bx) - \frac{\text{Subst}\left(\int (-a + \operatorname{csch}(x))^2 dx, x, \operatorname{csch}^{-1}(a + bx)\right)}{2b^2} \\
 &= -\frac{a^2 \operatorname{csch}^{-1}(a + bx)}{2b^2} + \frac{1}{2} x^2 \operatorname{csch}^{-1}(a + bx) - \frac{\text{Subst}\left(\int \operatorname{csch}^2(x) dx, x, \operatorname{csch}^{-1}(a + bx)\right)}{2b^2} \\
 &\quad + \frac{a \text{Subst}\left(\int \operatorname{csch}(x) dx, x, \operatorname{csch}^{-1}(a + bx)\right)}{b^2}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{a^2 \operatorname{csch}^{-1}(a+bx)}{2b^2} + \frac{1}{2}x^2 \operatorname{csch}^{-1}(a+bx) - \frac{a \operatorname{arctanh}\left(\sqrt{1+\frac{1}{(a+bx)^2}}\right)}{b^2} \\
&\quad + \frac{i \operatorname{Subst}\left(\int 1 dx, x, -i(a+bx)\sqrt{1+\frac{1}{(a+bx)^2}}\right)}{2b^2} \\
&= \frac{(a+bx)\sqrt{1+\frac{1}{(a+bx)^2}}}{2b^2} - \frac{a^2 \operatorname{csch}^{-1}(a+bx)}{2b^2} + \frac{1}{2}x^2 \operatorname{csch}^{-1}(a+bx) - \frac{a \operatorname{arctanh}\left(\sqrt{1+\frac{1}{(a+bx)^2}}\right)}{b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.47

$$\begin{aligned}
&\int x \operatorname{csch}^{-1}(a+bx) dx \\
&= \frac{(a+bx)\sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}} + b^2x^2 \operatorname{csch}^{-1}(a+bx) - a^2 \operatorname{arcsinh}\left(\frac{1}{a+bx}\right) - 2a \log\left((a+bx)\left(1+\sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}}\right)\right)}{2b^2}
\end{aligned}$$

[In] Integrate[x*ArcCsch[a + b*x], x]

[Out] ((a + b*x)*Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2] + b^2*x^2*ArcCsch[a + b*x] - a^2*ArcSinh[(a + b*x)^(-1)] - 2*a*Log[(a + b*x)*(1 + Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2])])/(2*b^2)

Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.29

method	result
derivativedivides	$-\frac{\operatorname{arccsch}(bx+a)a(bx+a) + \frac{\operatorname{arccsch}(bx+a)(bx+a)^2}{2} - \frac{\sqrt{(bx+a)^2+1} \left(2a \operatorname{arcsinh}(bx+a) - \sqrt{(bx+a)^2+1}\right)}{2(bx+a)\sqrt{\frac{(bx+a)^2+1}{(bx+a)^2}}}{b^2}$
default	$-\frac{\operatorname{arccsch}(bx+a)a(bx+a) + \frac{\operatorname{arccsch}(bx+a)(bx+a)^2}{2} - \frac{\sqrt{(bx+a)^2+1} \left(2a \operatorname{arcsinh}(bx+a) - \sqrt{(bx+a)^2+1}\right)}{2(bx+a)\sqrt{\frac{(bx+a)^2+1}{(bx+a)^2}}}{b^2}$
parts	$\frac{x^2 \operatorname{arccsch}(bx+a)}{2} + \frac{\sqrt{b^2x^2+2abx+a^2+1} \left(-a^2 \operatorname{arctanh}\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2+1}}\right)\sqrt{b^2}-2a \ln\left(\frac{b^2x+\sqrt{b^2x^2+2abx+a^2+1}\sqrt{b^2}}{\sqrt{b^2}}\right)\right)}{2b^2\sqrt{\frac{b^2x^2+2abx+a^2+1}{(bx+a)^2}}(bx+a)\sqrt{b^2}}$

[In] int(x*arccsch(b*x+a), x, method=_RETURNVERBOSE)

[Out] 1/b^2*(-arccsch(b*x+a)*a*(b*x+a)+1/2*arccsch(b*x+a)*(b*x+a)^2-1/2*((b*x+a)^2+1)^(1/2)*(2*a*arcsinh(b*x+a)-((b*x+a)^2+1)^(1/2)))/(b*x+a)/(((b*x+a)^2+1)/(b*x+a)^2)^(1/2)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 285 vs. 2(65) = 130.

Time = 0.26 (sec) , antiderivative size = 285, normalized size of antiderivative = 3.80

$$\int x \operatorname{csch}^{-1}(a + bx) dx$$

$$= \frac{b^2 x^2 \log\left(\frac{(bx+a)\sqrt{\frac{b^2 x^2 + 2abx + a^2 + 1}{b^2 x^2 + 2abx + a^2}} + 1}{bx+a}\right) - a^2 \log\left(-bx + (bx+a)\sqrt{\frac{b^2 x^2 + 2abx + a^2 + 1}{b^2 x^2 + 2abx + a^2}} - a + 1\right) + a^2 \log\left(-bx + (bx+a)\sqrt{\frac{b^2 x^2 + 2abx + a^2 + 1}{b^2 x^2 + 2abx + a^2}} + a + 1\right)}{b^2}$$

[In] integrate(x*arccsch(b*x+a),x, algorithm="fricas")

[Out] 1/2*(b^2*x^2*log(((b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) + 1)/(b*x + a)) - a^2*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a + 1) + a^2*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a - 1) + 2*a*log(-b*x + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)) - a) + (b*x + a)*sqrt((b^2*x^2 + 2*a*b*x + a^2 + 1)/(b^2*x^2 + 2*a*b*x + a^2)))/b^2

Sympy [F]

$$\int x \operatorname{csch}^{-1}(a + bx) dx = \int x \operatorname{acsch}(a + bx) dx$$

[In] integrate(x*acsch(b*x+a),x)

[Out] Integral(x*acsch(a + b*x), x)

Maxima [F]

$$\int x \operatorname{csch}^{-1}(a + bx) dx = \int x \operatorname{arcsch}(bx + a) dx$$

[In] integrate(x*arccsch(b*x+a),x, algorithm="maxima")

[Out] 1/2*I*a*(log(I*(b^2*x + a*b)/b + 1) - log(-I*(b^2*x + a*b)/b + 1))/b^2 + 1/4*(2*b^2*x^2*log(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) + 1) - (a^2 - 1)*log(b^2*x^2 + 2*a*b*x + a^2 + 1) - 2*(b^2*x^2 - a^2)*log(b*x + a))/b^2 + integrate(1/2*(b^2*x^3 + a*b*x^2)/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 + 1)^(3/2) + 1), x)

Giac [F]

$$\int x \operatorname{csch}^{-1}(a + bx) dx = \int x \operatorname{arcsch}(bx + a) dx$$

[In] integrate(x*arccsch(b*x+a),x, algorithm="giac")

[Out] integrate(x*arccsch(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{csch}^{-1}(a + bx) dx = \int x \operatorname{asinh}\left(\frac{1}{a + bx}\right) dx$$

[In] int(x*asinh(1/(a + b*x)),x)

[Out] int(x*asinh(1/(a + b*x)), x)

3.4 $\int \frac{\operatorname{csch}^{-1}(a+bx)}{x} dx$

Optimal result	63
Rubi [A] (verified)	64
Mathematica [C] (verified)	67
Maple [F]	68
Fricas [F]	68
Sympy [F]	68
Maxima [F]	69
Giac [F]	69
Mupad [F(-1)]	69

Optimal result

Integrand size = 10, antiderivative size = 162

$$\begin{aligned} \int \frac{\operatorname{csch}^{-1}(a+bx)}{x} dx &= \operatorname{csch}^{-1}(a+bx) \log \left(1 - \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{1 - \sqrt{1+a^2}} \right) \\ &\quad + \operatorname{csch}^{-1}(a+bx) \log \left(1 - \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{1 + \sqrt{1+a^2}} \right) \\ &\quad - \operatorname{csch}^{-1}(a+bx) \log \left(1 - e^{2\operatorname{csch}^{-1}(a+bx)} \right) \\ &\quad + \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{1 - \sqrt{1+a^2}} \right) + \operatorname{PolyLog} \left(2, \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{1 + \sqrt{1+a^2}} \right) \\ &\quad - \frac{1}{2} \operatorname{PolyLog} \left(2, e^{2\operatorname{csch}^{-1}(a+bx)} \right) \end{aligned}$$

```
[Out] -arccsch(b*x+a)*ln(1-(1/(b*x+a)+(1+1/(b*x+a)^2)^(1/2))^2)+arccsch(b*x+a)*ln
(1-a*(1/(b*x+a)+(1+1/(b*x+a)^2)^(1/2))/(1-(a^2+1)^(1/2)))+arccsch(b*x+a)*ln
(1-a*(1/(b*x+a)+(1+1/(b*x+a)^2)^(1/2))/(1+(a^2+1)^(1/2)))-1/2*polylog(2,(1/
(b*x+a)+(1+1/(b*x+a)^2)^(1/2))^2)+polylog(2,a*(1/(b*x+a)+(1+1/(b*x+a)^2)^(1
/2))/(1-(a^2+1)^(1/2)))+polylog(2,a*(1/(b*x+a)+(1+1/(b*x+a)^2)^(1/2))/(1+(a
^2+1)^(1/2)))
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {6457, 5715, 5688, 3797, 2221, 2317, 2438, 5680}

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x} dx = \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{1-\sqrt{a^2+1}}\right) + \operatorname{PolyLog}\left(2, \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{\sqrt{a^2+1}+1}\right) \\ + \operatorname{csch}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{1-\sqrt{a^2+1}}\right) \\ + \operatorname{csch}^{-1}(a+bx) \log\left(1 - \frac{ae^{\operatorname{csch}^{-1}(a+bx)}}{\sqrt{a^2+1}+1}\right) \\ - \frac{1}{2} \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(a+bx)}\right) \\ - \operatorname{csch}^{-1}(a+bx) \log\left(1 - e^{2\operatorname{csch}^{-1}(a+bx)}\right)$$

[In] Int[ArcCsch[a + b*x]/x,x]

[Out] ArcCsch[a + b*x]*Log[1 - (a*E^ArcCsch[a + b*x])/(1 - Sqrt[1 + a^2])] + ArcCsch[a + b*x]*Log[1 - (a*E^ArcCsch[a + b*x])/(1 + Sqrt[1 + a^2])] - ArcCsch[a + b*x]*Log[1 - E^(2*ArcCsch[a + b*x])] + PolyLog[2, (a*E^ArcCsch[a + b*x])/(1 - Sqrt[1 + a^2])] + PolyLog[2, (a*E^ArcCsch[a + b*x])/(1 + Sqrt[1 + a^2])] - PolyLog[2, E^(2*ArcCsch[a + b*x])]/2

Rule 2221

Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2317

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3797


```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_
.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist
[2*I, Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)
)/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && Int
egerQ[4*k] && IGtQ[m, 0]
```

Rule 5680

```
Int[(Cosh[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.))/((a_.) + (b_.)*Sin
h[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[-(e + f*x)^(m + 1)/(b*f*(m + 1)),
x] + (Int[(e + f*x)^m*(E^(c + d*x)/(a - Rt[a^2 + b^2, 2] + b*E^(c + d*x)))
, x] + Int[(e + f*x)^m*(E^(c + d*x)/(a + Rt[a^2 + b^2, 2] + b*E^(c + d*x)))
, x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[a^2 + b^2, 0]
```

Rule 5688

```
Int[(Coth[(c_.) + (d_.)*(x_)]^(n_.)*((e_.) + (f_.)*(x_))^(m_.))/((a_.) + (b_
.)*Sinh[(c_.) + (d_.)*(x_)]), x_Symbol] := Dist[1/a, Int[(e + f*x)^m*Coth[c
+ d*x]^n, x], x] - Dist[b/a, Int[(e + f*x)^m*Cosh[c + d*x]*(Coth[c + d*x]^
(n - 1)/(a + b*Sinh[c + d*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && I
GtQ[m, 0] && IGtQ[n, 0]
```

Rule 5715

```
Int[(((e_.) + (f_.)*(x_))^(m_.)*(F_)[(c_.) + (d_.)*(x_)]^(n_.)*(G_)[(c_.) +
(d_.)*(x_)]^(p_.))/(Csch[(c_.) + (d_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := I
nt[(e + f*x)^m*Sinh[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Sinh[c + d*x
])), x] /; FreeQ[{a, b, c, d, e, f}, x] && HyperbolicQ[F] && HyperbolicQ[G]
&& IntegersQ[m, n, p]
```

Rule 6457

```
Int[((a_.) + ArcCsch[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csch[x]*Co
th[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\text{Subst}\left(\int \frac{x \coth(x) \operatorname{csch}(x)}{-a + \operatorname{csch}(x)} dx, x, \operatorname{csch}^{-1}(a + bx)\right) \\ &= -\text{Subst}\left(\int \frac{x \coth(x)}{1 - a \sinh(x)} dx, x, \operatorname{csch}^{-1}(a + bx)\right) \end{aligned}$$

$$\begin{aligned}
&= -\left(a \text{Subst} \left(\int \frac{x \cosh(x)}{1 - a \sinh(x)} dx, x, \text{csch}^{-1}(a + bx) \right) \right) \\
&\quad - \text{Subst} \left(\int x \coth(x) dx, x, \text{csch}^{-1}(a + bx) \right) \\
&= 2 \text{Subst} \left(\int \frac{e^{2x} x}{1 - e^{2x}} dx, x, \text{csch}^{-1}(a + bx) \right) \\
&\quad - a \text{Subst} \left(\int \frac{e^x x}{1 - \sqrt{1 + a^2} - ae^x} dx, x, \text{csch}^{-1}(a + bx) \right) \\
&\quad - a \text{Subst} \left(\int \frac{e^x x}{1 + \sqrt{1 + a^2} - ae^x} dx, x, \text{csch}^{-1}(a + bx) \right) \\
&= \text{csch}^{-1}(a + bx) \log \left(1 - \frac{ae^{\text{csch}^{-1}(a+bx)}}{1 - \sqrt{1 + a^2}} \right) + \text{csch}^{-1}(a + bx) \log \left(1 - \frac{ae^{\text{csch}^{-1}(a+bx)}}{1 + \sqrt{1 + a^2}} \right) \\
&\quad - \text{csch}^{-1}(a + bx) \log \left(1 - e^{2\text{csch}^{-1}(a+bx)} \right) \\
&\quad - \text{Subst} \left(\int \log \left(1 - \frac{ae^x}{1 - \sqrt{1 + a^2}} \right) dx, x, \text{csch}^{-1}(a + bx) \right) \\
&\quad - \text{Subst} \left(\int \log \left(1 - \frac{ae^x}{1 + \sqrt{1 + a^2}} \right) dx, x, \text{csch}^{-1}(a + bx) \right) \\
&\quad + \text{Subst} \left(\int \log(1 - e^{2x}) dx, x, \text{csch}^{-1}(a + bx) \right) \\
&= \text{csch}^{-1}(a + bx) \log \left(1 - \frac{ae^{\text{csch}^{-1}(a+bx)}}{1 - \sqrt{1 + a^2}} \right) + \text{csch}^{-1}(a + bx) \log \left(1 - \frac{ae^{\text{csch}^{-1}(a+bx)}}{1 + \sqrt{1 + a^2}} \right) \\
&\quad - \text{csch}^{-1}(a + bx) \log \left(1 - e^{2\text{csch}^{-1}(a+bx)} \right) \\
&\quad + \frac{1}{2} \text{Subst} \left(\int \frac{\log(1 - x)}{x} dx, x, e^{2\text{csch}^{-1}(a+bx)} \right) \\
&\quad - \text{Subst} \left(\int \frac{\log \left(1 - \frac{ax}{1 - \sqrt{1 + a^2}} \right)}{x} dx, x, e^{\text{csch}^{-1}(a+bx)} \right) \\
&\quad - \text{Subst} \left(\int \frac{\log \left(1 - \frac{ax}{1 + \sqrt{1 + a^2}} \right)}{x} dx, x, e^{\text{csch}^{-1}(a+bx)} \right) \\
&= \text{csch}^{-1}(a + bx) \log \left(1 - \frac{ae^{\text{csch}^{-1}(a+bx)}}{1 - \sqrt{1 + a^2}} \right) + \text{csch}^{-1}(a + bx) \log \left(1 - \frac{ae^{\text{csch}^{-1}(a+bx)}}{1 + \sqrt{1 + a^2}} \right) \\
&\quad - \text{csch}^{-1}(a + bx) \log \left(1 - e^{2\text{csch}^{-1}(a+bx)} \right) + \text{PolyLog} \left(2, \frac{ae^{\text{csch}^{-1}(a+bx)}}{1 - \sqrt{1 + a^2}} \right) \\
&\quad + \text{PolyLog} \left(2, \frac{ae^{\text{csch}^{-1}(a+bx)}}{1 + \sqrt{1 + a^2}} \right) - \frac{1}{2} \text{PolyLog} \left(2, e^{2\text{csch}^{-1}(a+bx)} \right)
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.48 (sec) , antiderivative size = 427, normalized size of antiderivative = 2.64

$$\begin{aligned}
& \int \frac{\operatorname{csch}^{-1}(a+bx)}{x} dx \\
&= \frac{1}{8} \left(\pi^2 - 4i\pi \operatorname{csch}^{-1}(a+bx) - 8\operatorname{csch}^{-1}(a+bx)^2 \right. \\
&\quad - 32 \arcsin \left(\frac{\sqrt{\frac{-i+a}{a}}}{\sqrt{2}} \right) \arctan \left(\frac{(1-ia) \cot \left(\frac{1}{4}(\pi + 2i\operatorname{csch}^{-1}(a+bx)) \right)}{\sqrt{1+a^2}} \right) \\
&\quad - 8\operatorname{csch}^{-1}(a+bx) \log \left(1 - e^{-2\operatorname{csch}^{-1}(a+bx)} \right) + 4i\pi \log \left(1 - \frac{(-1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) \\
&\quad + 8\operatorname{csch}^{-1}(a+bx) \log \left(1 - \frac{(-1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) \\
&\quad + 16i \arcsin \left(\frac{\sqrt{\frac{-i+a}{a}}}{\sqrt{2}} \right) \log \left(1 - \frac{(-1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) \\
&\quad + 4i\pi \log \left(1 + \frac{(1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) \\
&\quad + 8\operatorname{csch}^{-1}(a+bx) \log \left(1 + \frac{(1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) \\
&\quad - 16i \arcsin \left(\frac{\sqrt{\frac{-i+a}{a}}}{\sqrt{2}} \right) \log \left(1 + \frac{(1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) - 4i\pi \log \left(\frac{bx}{a+bx} \right) \\
&\quad + 4 \operatorname{PolyLog} \left(2, e^{-2\operatorname{csch}^{-1}(a+bx)} \right) + 8 \operatorname{PolyLog} \left(2, \frac{(-1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) \\
&\quad \left. + 8 \operatorname{PolyLog} \left(2, -\frac{(1 + \sqrt{1+a^2}) e^{\operatorname{csch}^{-1}(a+bx)}}{a} \right) \right)
\end{aligned}$$

[In] Integrate[ArcCsch[a + b*x]/x,x]

[Out] (Pi^2 - (4*I)*Pi*ArcCsch[a + b*x] - 8*ArcCsch[a + b*x]^2 - 32*ArcSin[Sqrt[(-I + a)/a]/Sqrt[2]]*ArcTan[((1 - I*a)*Cot[(Pi + (2*I)*ArcCsch[a + b*x])/4])/Sqrt[1 + a^2]] - 8*ArcCsch[a + b*x]*Log[1 - E^(-2*ArcCsch[a + b*x])] + (4*I)*Pi*Log[1 - ((-1 + Sqrt[1 + a^2])*E^ArcCsch[a + b*x])/a] + 8*ArcCsch[a + b*x]*Log[1 - ((-1 + Sqrt[1 + a^2])*E^ArcCsch[a + b*x])/a] + (16*I)*ArcSin[S

```

qrt[(-I + a)/a]/Sqrt[2]]*Log[1 - ((-1 + Sqrt[1 + a^2])*E^ArcCsch[a + b*x])/
a] + (4*I)*Pi*Log[1 + ((1 + Sqrt[1 + a^2])*E^ArcCsch[a + b*x])/a] + 8*ArcCs
ch[a + b*x]*Log[1 + ((1 + Sqrt[1 + a^2])*E^ArcCsch[a + b*x])/a] - (16*I)*Ar
cSin[Sqrt[(-I + a)/a]/Sqrt[2]]*Log[1 + ((1 + Sqrt[1 + a^2])*E^ArcCsch[a + b
*x])/a] - (4*I)*Pi*Log[(b*x)/(a + b*x)] + 4*PolyLog[2, E^(-2*ArcCsch[a + b*
x])] + 8*PolyLog[2, ((-1 + Sqrt[1 + a^2])*E^ArcCsch[a + b*x])/a] + 8*PolyLo
g[2, -(((1 + Sqrt[1 + a^2])*E^ArcCsch[a + b*x])/a))]/8

```

Maple [F]

$$\int \frac{\operatorname{arccsch}(bx + a)}{x} dx$$

```
[In] int(arccsch(b*x+a)/x,x)
```

```
[Out] int(arccsch(b*x+a)/x,x)
```

Fricas [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arcsch}(bx + a)}{x} dx$$

```
[In] integrate(arccsch(b*x+a)/x,x, algorithm="fricas")
```

```
[Out] integral(arccsch(b*x + a)/x, x)
```

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{acsch}(a + bx)}{x} dx$$

```
[In] integrate(acsch(b*x+a)/x,x)
```

```
[Out] Integral(acsch(a + b*x)/x, x)
```

Maxima [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arcsch}(bx + a)}{x} dx$$

[In] integrate(arccsch(b*x+a)/x,x, algorithm="maxima")

[Out] integrate(arccsch(b*x + a)/x, x)

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arcsch}(bx + a)}{x} dx$$

[In] integrate(arccsch(b*x+a)/x,x, algorithm="giac")

[Out] integrate(arccsch(b*x + a)/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{a+bx}\right)}{x} dx$$

[In] int(asinh(1/(a + b*x))/x,x)

[Out] int(asinh(1/(a + b*x))/x, x)

3.5 $\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^2} dx$

Optimal result	70
Rubi [A] (verified)	70
Mathematica [B] (verified)	72
Maple [B] (verified)	72
Fricas [B] (verification not implemented)	73
Sympy [F]	74
Maxima [F]	74
Giac [F]	74
Mupad [F(-1)]	74

Optimal result

Integrand size = 10, antiderivative size = 63

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^2} dx = -\frac{b\operatorname{csch}^{-1}(a+bx)}{a} - \frac{\operatorname{csch}^{-1}(a+bx)}{x} + \frac{2b\operatorname{arctanh}\left(\frac{a+\tanh\left(\frac{1}{2}\operatorname{csch}^{-1}(a+bx)\right)}{\sqrt{1+a^2}}\right)}{a\sqrt{1+a^2}}$$

[Out] $-b*\operatorname{arccsch}(b*x+a)/a-\operatorname{arccsch}(b*x+a)/x+2*b*\operatorname{arctanh}((a+\tanh(1/2*\operatorname{arccsch}(b*x+a)))/\sqrt{a^2+1})/a/\sqrt{a^2+1}$

Rubi [A] (verified)

Time = 0.08 (sec), antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6457, 5577, 3868, 2739, 632, 212}

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^2} dx = \frac{2b\operatorname{arctanh}\left(\frac{\tanh\left(\frac{1}{2}\operatorname{csch}^{-1}(a+bx)\right)+a}{\sqrt{a^2+1}}\right)}{a\sqrt{a^2+1}} - \frac{b\operatorname{csch}^{-1}(a+bx)}{a} - \frac{\operatorname{csch}^{-1}(a+bx)}{x}$$

[In] $\operatorname{Int}[\operatorname{ArcCsch}[a+b*x]/x^2,x]$

[Out] $-(b*\operatorname{ArcCsch}[a+b*x])/a - \operatorname{ArcCsch}[a+b*x]/x + (2*b*\operatorname{ArcTanh}[(a+\operatorname{Tanh}[\operatorname{ArcCsch}[a+b*x]/2])/ \operatorname{Sqrt}[1+a^2]])/(a*\operatorname{Sqrt}[1+a^2])$

Rule 212

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 632

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[In
t[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 2739

```
Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = Fre
eFactors[Tan[(c + d*x)/2], x]}, Dist[2*(e/d), Subst[Int[1/(a + 2*b*e*x + a*
e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[
a^2 - b^2, 0]
```

Rule 3868

```
Int[(csc[(c_) + (d_)*(x_)]*(b_) + (a_))(-1), x_Symbol] := Simp[x/a, x]
- Dist[1/a, Int[1/(1 + (a/b)*Sin[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x
] && NeQ[a^2 - b^2, 0]
```

Rule 5577

```
Int[Coth[(c_) + (d_)*(x_)]*Csch[(c_) + (d_)*(x_)]*(Csch[(c_) + (d_)*(
x_)]*(b_) + (a_))^(n_)*((e_) + (f_)*(x_))^(m_), x_Symbol] := Simp[(-e
+ f*x)^m*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b
*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x] /
; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 6457

```
Int[((a_) + ArcCsch[(c_) + (d_)*(x_)]*(b_))^(p_)*((e_) + (f_)*(x_))^(
m_), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csch[x]*Co
th[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(b\text{Subst}\left(\int \frac{x \coth(x) \operatorname{csch}(x)}{(-a + \operatorname{csch}(x))^2} dx, x, \operatorname{csch}^{-1}(a + bx)\right)\right) \\ &= -\frac{\operatorname{csch}^{-1}(a + bx)}{x} + b\text{Subst}\left(\int \frac{1}{-a + \operatorname{csch}(x)} dx, x, \operatorname{csch}^{-1}(a + bx)\right) \end{aligned}$$

$$\begin{aligned}
&= -\frac{b\operatorname{csch}^{-1}(a+bx)}{a} - \frac{\operatorname{csch}^{-1}(a+bx)}{x} + \frac{b\operatorname{Subst}\left(\int \frac{1}{1-a\sinh(x)} dx, x, \operatorname{csch}^{-1}(a+bx)\right)}{a} \\
&= -\frac{b\operatorname{csch}^{-1}(a+bx)}{a} - \frac{\operatorname{csch}^{-1}(a+bx)}{x} + \frac{(2b)\operatorname{Subst}\left(\int \frac{1}{1-2ax-x^2} dx, x, \tanh\left(\frac{1}{2}\operatorname{csch}^{-1}(a+bx)\right)\right)}{a} \\
&= -\frac{b\operatorname{csch}^{-1}(a+bx)}{a} - \frac{\operatorname{csch}^{-1}(a+bx)}{x} \\
&\quad - \frac{(4b)\operatorname{Subst}\left(\int \frac{1}{4(1+a^2)-x^2} dx, x, -2a - 2\tanh\left(\frac{1}{2}\operatorname{csch}^{-1}(a+bx)\right)\right)}{a} \\
&= -\frac{b\operatorname{csch}^{-1}(a+bx)}{a} - \frac{\operatorname{csch}^{-1}(a+bx)}{x} + \frac{2b\operatorname{arctanh}\left(\frac{a+\tanh\left(\frac{1}{2}\operatorname{csch}^{-1}(a+bx)\right)}{\sqrt{1+a^2}}\right)}{a\sqrt{1+a^2}}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 141 vs. $2(63) = 126$.

Time = 0.17 (sec) , antiderivative size = 141, normalized size of antiderivative = 2.24

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^2} dx = -\frac{\operatorname{csch}^{-1}(a+bx)}{x} - \frac{b\left(\sqrt{1+a^2}\operatorname{arcsinh}\left(\frac{1}{a+bx}\right) + \log(x) - \log\left(1+a^2+abx+a\sqrt{1+a^2}\sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}} + \sqrt{1+a^2}bx\sqrt{\frac{1+a^2}{(a+bx)^2}}\right)\right)}{a\sqrt{1+a^2}}$$

[In] Integrate[ArcCsch[a + b*x]/x^2,x]

[Out] -(ArcCsch[a + b*x]/x) - (b*(Sqrt[1 + a^2]*ArcSinh[(a + b*x)^(-1)] + Log[x] - Log[1 + a^2 + a*b*x + a*Sqrt[1 + a^2]*Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2] + Sqrt[1 + a^2]*b*x*Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]]))/(a*Sqrt[1 + a^2])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 126 vs. $2(57) = 114$.

Time = 0.64 (sec) , antiderivative size = 127, normalized size of antiderivative = 2.02

method	result
derivativedivides	$b \left(-\frac{\operatorname{arccsch}(bx+a)}{bx} - \frac{\sqrt{(bx+a)^2+1} \left(\operatorname{arctanh} \left(\frac{1}{\sqrt{(bx+a)^2+1}} \right) \sqrt{a^2+1} - \ln \left(\frac{2\sqrt{a^2+1} \sqrt{(bx+a)^2+1+2(bx+a)a+2}}{bx} \right) \right)}{\sqrt{\frac{(bx+a)^2+1}{(bx+a)^2}} (bx+a)a\sqrt{a^2+1}} \right)$
default	$b \left(-\frac{\operatorname{arccsch}(bx+a)}{bx} - \frac{\sqrt{(bx+a)^2+1} \left(\operatorname{arctanh} \left(\frac{1}{\sqrt{(bx+a)^2+1}} \right) \sqrt{a^2+1} - \ln \left(\frac{2\sqrt{a^2+1} \sqrt{(bx+a)^2+1+2(bx+a)a+2}}{bx} \right) \right)}{\sqrt{\frac{(bx+a)^2+1}{(bx+a)^2}} (bx+a)a\sqrt{a^2+1}} \right)$
parts	$-\frac{\operatorname{arccsch}(bx+a)}{x} - \frac{b\sqrt{b^2x^2+2abx+a^2+1} \left(\operatorname{arctanh} \left(\frac{1}{\sqrt{b^2x^2+2abx+a^2+1}} \right) \sqrt{a^2+1} - \ln \left(\frac{2a^2+2+2abx+2\sqrt{a^2+1}\sqrt{b^2x^2+2abx+a^2+1}}{x} \right) \right)}{\sqrt{\frac{b^2x^2+2abx+a^2+1}{(bx+a)^2}} (bx+a)a\sqrt{a^2+1}}$

[In] `int(arccsch(b*x+a)/x^2,x,method=_RETURNVERBOSE)`

[Out]
$$b \cdot \left(-\frac{1}{b \cdot x} \operatorname{arccsch}(b \cdot x + a) - \left((b \cdot x + a)^2 + 1 \right)^{1/2} \cdot \left(\operatorname{arctanh} \left(\frac{1}{\left((b \cdot x + a)^2 + 1 \right)^{1/2}} \right) \sqrt{a^2 + 1} - \ln \left(\frac{2 \cdot \left(a^2 + 1 \right)^{1/2} \cdot \left((b \cdot x + a)^2 + 1 \right)^{1/2} + (b \cdot x + a) \cdot a + 1}{b \cdot x} \right) \right) / \left(\left((b \cdot x + a)^2 + 1 \right) / (b \cdot x + a)^2 \right)^{1/2} / (b \cdot x + a) / a / \left(a^2 + 1 \right)^{1/2} \right)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 343 vs. $2(57) = 114$.

Time = 0.27 (sec) , antiderivative size = 343, normalized size of antiderivative = 5.44

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^2} dx =$$

$$(a^2 + 1)bx \log \left(-bx + (bx + a) \sqrt{\frac{b^2x^2 + 2abx + a^2 + 1}{b^2x^2 + 2abx + a^2}} - a + 1 \right) - (a^2 + 1)bx \log \left(-bx + (bx + a) \sqrt{\frac{b^2x^2 + 2abx + a^2 + 1}{b^2x^2 + 2abx + a^2}} \right)$$

[In] `integrate(arccsch(b*x+a)/x^2,x, algorithm="fricas")`

[Out]
$$-\left((a^2 + 1) \cdot b \cdot x \cdot \log(-b \cdot x + (b \cdot x + a) \cdot \sqrt{(b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1) / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2)}) - a + 1 \right) - (a^2 + 1) \cdot b \cdot x \cdot \log(-b \cdot x + (b \cdot x + a) \cdot \sqrt{(b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1) / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2)}) - a - 1 - \sqrt{(a^2 + 1) \cdot b \cdot x} \cdot \log(-a^2 \cdot b \cdot x + a^3 + (a \cdot b \cdot x + a^2 + (a \cdot b \cdot x + a^2) \cdot \sqrt{(b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1) / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2)}) + 1) \cdot \sqrt{(a^2 + 1)} + (a^3 + (a^2 + 1) \cdot b \cdot x + a) \cdot \sqrt{(b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1) / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2)}) + a) / x + (a^3 + a) \cdot \log(((b \cdot x + a) \cdot \sqrt{(b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1) / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2)}) + 1) / (b \cdot x + a)) / ((a^3 + a) \cdot x)$$

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{acsch}(a + bx)}{x^2} dx$$

[In] integrate(acsch(b*x+a)/x**2,x)

[Out] Integral(acsch(a + b*x)/x**2, x)

Maxima [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{arcsch}(bx + a)}{x^2} dx$$

[In] integrate(arccsch(b*x+a)/x^2,x, algorithm="maxima")

[Out] $-1/2*I*b*(\log(I*(b^2*x + a*b)/b + 1) - \log(-I*(b^2*x + a*b)/b + 1))/(a^2 + 1) - b*\log(x)/(a^3 + a) - 1/2*(a^2*b*x*\log(b^2*x^2 + 2*a*b*x + a^2 + 1) - 2*(a^3 + (a^2*b + b)*x + a)*\log(b*x + a) + 2*(a^3 + a)*\log(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1} + 1))/((a^3 + a)*x) - \operatorname{integrate}((b^2*x + a*b)/(b^2*x^3 + 2*a*b*x^2 + (a^2 + 1)*x + (b^2*x^3 + 2*a*b*x^2 + (a^2 + 1)*x)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}), x)$

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{arcsch}(bx + a)}{x^2} dx$$

[In] integrate(arccsch(b*x+a)/x^2,x, algorithm="giac")

[Out] integrate(arccsch(b*x + a)/x^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{a+bx}\right)}{x^2} dx$$

[In] int(asinh(1/(a + b*x))/x^2,x)

[Out] int(asinh(1/(a + b*x))/x^2, x)

3.6 $\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^3} dx$

Optimal result	75
Rubi [A] (verified)	75
Mathematica [A] (verified)	78
Maple [B] (verified)	78
Fricas [B] (verification not implemented)	79
Sympy [F]	80
Maxima [F]	80
Giac [F]	80
Mupad [F(-1)]	81

Optimal result

Integrand size = 10, antiderivative size = 114

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^3} dx = \frac{b(a+bx)\sqrt{1+\frac{1}{(a+bx)^2}}}{2a(1+a^2)x} + \frac{b^2\operatorname{csch}^{-1}(a+bx)}{2a^2} - \frac{\operatorname{csch}^{-1}(a+bx)}{2x^2} - \frac{(1+2a^2)b^2\operatorname{arctanh}\left(\frac{a+\tanh\left(\frac{1}{2}\operatorname{csch}^{-1}(a+bx)\right)}{\sqrt{1+a^2}}\right)}{a^2(1+a^2)^{3/2}}$$

[Out] $1/2*b^2*\operatorname{arccsch}(b*x+a)/a^2-1/2*\operatorname{arccsch}(b*x+a)/x^2-(2*a^2+1)*b^2*\operatorname{arctanh}\left(\frac{a+\tanh(1/2*\operatorname{arccsch}(b*x+a))}{\sqrt{1+a^2}}\right)/a^2/(a^2+1)^{3/2}+1/2*b*(b*x+a)*(1+1/(b*x+a)^2)^{1/2}/a/(a^2+1)/x$

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {6457, 5577, 3870, 4004, 3916, 2739, 632, 212}

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^3} dx = -\frac{(2a^2+1)b^2\operatorname{arctanh}\left(\frac{\tanh\left(\frac{1}{2}\operatorname{csch}^{-1}(a+bx)\right)+a}{\sqrt{a^2+1}}\right)}{a^2(a^2+1)^{3/2}} + \frac{b^2\operatorname{csch}^{-1}(a+bx)}{2a^2} + \frac{b(a+bx)\sqrt{\frac{1}{(a+bx)^2}+1}}{2a(a^2+1)x} - \frac{\operatorname{csch}^{-1}(a+bx)}{2x^2}$$

[In] Int[ArcCsch[a + b*x]/x^3,x]

[Out] $(b*(a + b*x)*\text{Sqrt}[1 + (a + b*x)^{-2}])/(2*a*(1 + a^2)*x) + (b^2*\text{ArcCsch}[a + b*x])/(2*a^2) - \text{ArcCsch}[a + b*x]/(2*x^2) - ((1 + 2*a^2)*b^2*\text{ArcTanh}[(a + \text{Tanh}[\text{ArcCsch}[a + b*x]/2])/\text{Sqrt}[1 + a^2]])/(a^2*(1 + a^2)^{(3/2)})$

Rule 212

$\text{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 632

$\text{Int}[(a + (b \cdot x) + (c \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}\{a, b, c\}, x \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 2739

$\text{Int}[(a + (b \cdot \sin[c + d \cdot x])^2)^{-1}, x_Symbol] \rightarrow \text{With}\{e = \text{FreeFactors}[\text{Tan}[(c + d \cdot x)/2], x]\}, \text{Dist}[2*(e/d), \text{Subst}[\text{Int}[1/(a + 2*b*e*x + a*e^2*x^2), x], x, \text{Tan}[(c + d \cdot x)/2]/e], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$

Rule 3870

$\text{Int}[(\text{csc}[c + d \cdot x] + (b \cdot x)^n)^{-1}, x_Symbol] \rightarrow \text{Simp}[b^2*\text{Cot}[c + d \cdot x]*(a + b*\text{Csc}[c + d \cdot x])^{n+1}/(a*d*(n+1)*(a^2 - b^2)), x] + \text{Dist}[1/(a*(n+1)*(a^2 - b^2)), \text{Int}[(a + b*\text{Csc}[c + d \cdot x])^{n+1}*\text{Simp}[(a^2 - b^2)*(n+1) - a*b*(n+1)*\text{Csc}[c + d \cdot x] + b^2*(n+2)*\text{Csc}[c + d \cdot x]^2, x], x], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 3916

$\text{Int}[\text{csc}[e + f \cdot x]/(\text{csc}[e + f \cdot x] + (b \cdot x)^n), x_Symbol] \rightarrow \text{Dist}[1/b, \text{Int}[1/(1 + (a/b)*\text{Sin}[e + f \cdot x]), x], x] /; \text{FreeQ}\{a, b, e, f\}, x \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$

Rule 4004

$\text{Int}[(\text{csc}[e + f \cdot x] + (b \cdot x)^n)/(a + b*\text{Csc}[e + f \cdot x]), x_Symbol] \rightarrow \text{Simp}[c*(x/a), x] - \text{Dist}[(b*c - a*d)/a, \text{Int}[\text{Csc}[e + f \cdot x]/(a + b*\text{Csc}[e + f \cdot x]), x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0]$

Rule 5577

```

Int[Coth[(c_.) + (d_.)*(x_)]*Csch[(c_.) + (d_.)*(x_)]*(Csch[(c_.) + (d_.)*(
x_)]*(b_.) + (a_.))^(n_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(-e
+ f*x)^m)*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b
*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x] /
; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

```

Rule 6457

```

Int[((a_.) + ArcCsch[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csch[x]*Co
th[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\left(b^2 \text{Subst}\left(\int \frac{x \coth(x) \operatorname{csch}(x)}{(-a + \operatorname{csch}(x))^3} dx, x, \operatorname{csch}^{-1}(a + bx)\right)\right) \\
&= -\frac{\operatorname{csch}^{-1}(a + bx)}{2x^2} + \frac{1}{2} b^2 \text{Subst}\left(\int \frac{1}{(-a + \operatorname{csch}(x))^2} dx, x, \operatorname{csch}^{-1}(a + bx)\right) \\
&= \frac{b(a + bx) \sqrt{1 + \frac{1}{(a + bx)^2}}}{2a(1 + a^2)x} - \frac{\operatorname{csch}^{-1}(a + bx)}{2x^2} + \frac{b^2 \text{Subst}\left(\int \frac{-1 - a^2 - a \operatorname{csch}(x)}{-a + \operatorname{csch}(x)} dx, x, \operatorname{csch}^{-1}(a + bx)\right)}{2a(1 + a^2)} \\
&= \frac{b(a + bx) \sqrt{1 + \frac{1}{(a + bx)^2}}}{2a(1 + a^2)x} + \frac{b^2 \operatorname{csch}^{-1}(a + bx)}{2a^2} - \frac{\operatorname{csch}^{-1}(a + bx)}{2x^2} \\
&\quad - \frac{((1 + 2a^2)b^2) \text{Subst}\left(\int \frac{\operatorname{csch}(x)}{-a + \operatorname{csch}(x)} dx, x, \operatorname{csch}^{-1}(a + bx)\right)}{2a^2(1 + a^2)} \\
&= \frac{b(a + bx) \sqrt{1 + \frac{1}{(a + bx)^2}}}{2a(1 + a^2)x} + \frac{b^2 \operatorname{csch}^{-1}(a + bx)}{2a^2} - \frac{\operatorname{csch}^{-1}(a + bx)}{2x^2} \\
&\quad - \frac{((1 + 2a^2)b^2) \text{Subst}\left(\int \frac{1}{1 - a \sinh(x)} dx, x, \operatorname{csch}^{-1}(a + bx)\right)}{2a^2(1 + a^2)} \\
&= \frac{b(a + bx) \sqrt{1 + \frac{1}{(a + bx)^2}}}{2a(1 + a^2)x} + \frac{b^2 \operatorname{csch}^{-1}(a + bx)}{2a^2} - \frac{\operatorname{csch}^{-1}(a + bx)}{2x^2} \\
&\quad - \frac{((1 + 2a^2)b^2) \text{Subst}\left(\int \frac{1}{1 - 2ax - x^2} dx, x, \tanh\left(\frac{1}{2} \operatorname{csch}^{-1}(a + bx)\right)\right)}{a^2(1 + a^2)}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b(a+bx)\sqrt{1+\frac{1}{(a+bx)^2}}}{2a(1+a^2)x} + \frac{b^2\operatorname{csch}^{-1}(a+bx)}{2a^2} - \frac{\operatorname{csch}^{-1}(a+bx)}{2x^2} \\
&\quad + \frac{(2(1+2a^2)b^2)\operatorname{Subst}\left(\int \frac{1}{4(1+a^2)-x^2} dx, x, -2a-2\tanh\left(\frac{1}{2}\operatorname{csch}^{-1}(a+bx)\right)\right)}{a^2(1+a^2)} \\
&= \frac{b(a+bx)\sqrt{1+\frac{1}{(a+bx)^2}}}{2a(1+a^2)x} + \frac{b^2\operatorname{csch}^{-1}(a+bx)}{2a^2} - \frac{\operatorname{csch}^{-1}(a+bx)}{2x^2} \\
&\quad - \frac{(1+2a^2)b^2\operatorname{arctanh}\left(\frac{a+\tanh\left(\frac{1}{2}\operatorname{csch}^{-1}(a+bx)\right)}{\sqrt{1+a^2}}\right)}{a^2(1+a^2)^{3/2}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.47 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.93

$$\begin{aligned}
&\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^3} dx \\
&= \frac{1}{2} \left(\frac{b(a+bx)\sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}}}{a(1+a^2)x} - \frac{\operatorname{csch}^{-1}(a+bx)}{x^2} + \frac{b^2\operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}{a^2} + \frac{(1+2a^2)b^2\log(x)}{a^2(1+a^2)^{3/2}} \right. \\
&\quad \left. - \frac{(1+2a^2)b^2\log\left(1+a^2+abx+a\sqrt{1+a^2}\sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}}+\sqrt{1+a^2}bx\sqrt{\frac{1+a^2+2abx+b^2x^2}{(a+bx)^2}}\right)}{a^2(1+a^2)^{3/2}} \right)
\end{aligned}$$

[In] Integrate[ArcCsch[a + b*x]/x^3,x]

[Out] ((b*(a + b*x)*Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2])/(a*(1 + a^2)*x) - ArcCsch[a + b*x]/x^2 + (b^2*ArcSinh[(a + b*x)^(-1)]/a^2 + ((1 + 2*a^2)*b^2*Log[x])/(a^2*(1 + a^2)^(3/2)) - ((1 + 2*a^2)*b^2*Log[1 + a^2 + a*b*x + a*Sqrt[1 + a^2]*Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2] + Sqrt[1 + a^2]*b*x*Sqrt[(1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]])/(a^2*(1 + a^2)^(3/2)))/2

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 317 vs. 2(100) = 200.

Time = 0.66 (sec) , antiderivative size = 318, normalized size of antiderivative = 2.79

method	result
parts	$-\frac{\operatorname{arccsch}(bx+a)}{2x^2} + \frac{b\sqrt{b^2x^2+2abx+a^2+1} \left((a^2+1)^{\frac{3}{2}} \operatorname{arctanh}\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2+1}}\right) a^2bx - 2 \ln\left(\frac{2a^2+2+2abx+2\sqrt{a^2+1}}{x}\right) \right)}{2x^2}$
derivativedivides	$b^2 \left(-\frac{\operatorname{arccsch}(bx+a)}{2b^2x^2} + \frac{\sqrt{(bx+a)^2+1} \left(-\operatorname{arctanh}\left(\frac{1}{\sqrt{(bx+a)^2+1}}\right) (a^2+1)^{\frac{3}{2}} a^3 + \operatorname{arctanh}\left(\frac{1}{\sqrt{(bx+a)^2+1}}\right) (a^2+1)^{\frac{3}{2}} a^2 \right)}{2b^2x^2} \right)$
default	$b^2 \left(-\frac{\operatorname{arccsch}(bx+a)}{2b^2x^2} + \frac{\sqrt{(bx+a)^2+1} \left(-\operatorname{arctanh}\left(\frac{1}{\sqrt{(bx+a)^2+1}}\right) (a^2+1)^{\frac{3}{2}} a^3 + \operatorname{arctanh}\left(\frac{1}{\sqrt{(bx+a)^2+1}}\right) (a^2+1)^{\frac{3}{2}} a^2 \right)}{2b^2x^2} \right)$

[In] `int(arccsch(b*x+a)/x^3,x,method=_RETURNVERBOSE)`

[Out]
$$-1/2*\operatorname{arccsch}(b*x+a)/x^2+1/2*b*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}*((a^2+1)^{(3/2)}*\operatorname{arctanh}(1/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2}))*a^2*b*x-2*\ln(2*(a*b*x+(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+a^2+1)/x)*a^4*b*x+b*\operatorname{arctanh}(1/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2}))*x*(a^2+1)^{(3/2)}+(a^2+1)^{(3/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}*a-3*\ln(2*(a*b*x+(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+a^2+1)/x)*a^2*b*x-b*\ln(2*(a*b*x+(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+a^2+1)/x)*x)/((b^2*x^2+2*a*b*x+a^2+1)/(b*x+a)^2)^{(1/2)}/(b*x+a)/a^2/(a^2+1)^{(5/2)}/x$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 461 vs. $2(100) = 200$.

Time = 0.28 (sec) , antiderivative size = 461, normalized size of antiderivative = 4.04

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{x^3} dx$$

$$= \frac{(2a^2+1)\sqrt{a^2+1}b^2x^2 \log\left(-\frac{a^2bx+a^3-(abx+a^2+(abx+a^2)\sqrt{\frac{b^2x^2+2abx+a^2+1}{b^2x^2+2abx+a^2+1}}+1)}{x}\sqrt{a^2+1}+(a^3+(a^2+1)bx+a)\sqrt{\frac{b^2x^2+2abx+a^2+1}{b^2x^2+2abx+a^2+1}}\right)}{2x^2}$$

[In] `integrate(arccsch(b*x+a)/x^3,x, algorithm="fricas")`

[Out]
$$1/2*((2*a^2+1)*\sqrt{a^2+1}*b^2*x^2*\log(-(a^2*b*x+a^3-(a*b*x+a^2+(a*b*x+a^2)*\sqrt{(b^2*x^2+2*a*b*x+a^2+1)/(b^2*x^2+2*a*b*x+a^2+1)}+1))*\sqrt{a^2+1}+(a^3+(a^2+1)*b*x+a)*\sqrt{(b^2*x^2+2*a*b*x+a^2+1)/(b^2*x^2+2*a*b*x+a^2+1)})))/x+(a^4+2*a^2+1)*b^2*x^2*\log(-b*x+(b*x+a)*\sqrt{(b^2*x^2+2*a*b*x+a^2+1)/(b^2*x^2+2*a*b*x+a^2+1)})-a+1)-(a^4+2*a^2+1)*b^2*x^2*\log(-b*x+(b*x+a)*\sqrt{(b^2*x^2+2*a*b*x+a^2+1)/(b^2*x^2+2*a*b*x+a^2+1)})-a-1)+(a^3+a)*b^2*$$

$$x^2 - (a^6 + 2a^4 + a^2) \log\left(\frac{(bx + a) \sqrt{(b^2x^2 + 2abx + a^2 + 1)}}{(b^2x^2 + 2abx + a^2)} + 1\right) + \frac{(a^3 + a)b^2x^2 + (a^4 + a^2)bx}{(b^2x^2 + 2abx + a^2)} \sqrt{\frac{(b^2x^2 + 2abx + a^2 + 1)}{(b^2x^2 + 2abx + a^2)}} \Big/ ((a^6 + 2a^4 + a^2)x^2)$$

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{acsch}(a + bx)}{x^3} dx$$

[In] integrate(acsch(b*x+a)/x**3,x)

[Out] Integral(acsch(a + b*x)/x**3, x)

Maxima [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{arcsch}(bx + a)}{x^3} dx$$

[In] integrate(arccsch(b*x+a)/x^3,x, algorithm="maxima")

[Out] $\frac{1}{2} I a b^2 (\log(I (b^2 x + a b) / b + 1) - \log(-I (b^2 x + a b) / b + 1)) / (a^4 + 2 a^2 + 1) + \frac{1}{2} (3 a^2 b^2 + b^2) \log(x) / (a^6 + 2 a^4 + a^2) + \frac{1}{4} ((a^4 b^2 - a^2 b^2) x^2 \log(b^2 x^2 + 2 a b x + a^2 + 1) + 2 (a^3 b + a b) x + 2 (a^6 + 2 a^4 - (a^4 b^2 + 2 a^2 b^2 + b^2) x^2 + a^2) \log(b x + a) - 2 (a^6 + 2 a^4 + a^2) \log(\sqrt{b^2 x^2 + 2 a b x + a^2 + 1} + 1)) / ((a^6 + 2 a^4 + a^2) x^2) - \int (1/2 (b^2 x + a b) / (b^2 x^4 + 2 a b x^3 + (a^2 + 1) x^2 + (b^2 x^4 + 2 a b x^3 + (a^2 + 1) x^2) \sqrt{b^2 x^2 + 2 a b x + a^2 + 1}), x)$

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{arcsch}(bx + a)}{x^3} dx$$

[In] integrate(arccsch(b*x+a)/x^3,x, algorithm="giac")

[Out] integrate(arccsch(b*x + a)/x^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{a+bx}\right)}{x^3} dx$$

```
[In] int(asinh(1/(a + b*x))/x^3,x)
```

```
[Out] int(asinh(1/(a + b*x))/x^3, x)
```

3.7 $\int (e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$

Optimal result	82
Rubi [A] (verified)	83
Mathematica [C] (warning: unable to verify)	91
Maple [F]	92
Fricas [F]	93
Sympy [F]	93
Maxima [F]	93
Giac [F]	94
Mupad [F(-1)]	94

Optimal result

Integrand size = 20, antiderivative size = 501

$$\begin{aligned}
 & \int (e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx \\
 &= \frac{b^2 f^2 (de - cf)x}{d^3} + \frac{b^2 f^3 (c + dx)^2}{12d^4} - \frac{bf^3 (c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{3d^4} \\
 &+ \frac{3bf(de - cf)^2 (c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{d^4} \\
 &+ \frac{bf^2 (de - cf)(c + dx)^2 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{d^4} \\
 &+ \frac{bf^3 (c + dx)^3 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{6d^4} \\
 &- \frac{(de - cf)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4d^4 f} + \frac{(e + fx)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4f} \\
 &- \frac{2bf^2 (de - cf) (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
 &+ \frac{4b(de - cf)^3 (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} - \frac{b^2 f^3 \log(c + dx)}{3d^4} \\
 &+ \frac{3b^2 f (de - cf)^2 \log(c + dx)}{d^4} - \frac{b^2 f^2 (de - cf) \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
 &+ \frac{2b^2 (de - cf)^3 \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
 &+ \frac{b^2 f^2 (de - cf) \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} - \frac{2b^2 (de - cf)^3 \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4}
 \end{aligned}$$

```
[Out] b^2*f^2*(-c*f+d*e)*x/d^3+1/12*b^2*f^3*(d*x+c)^2/d^4-1/4*(-c*f+d*e)^4*(a+b*arccsch(d*x+c))^2/d^4/f+1/4*(f*x+e)^4*(a+b*arccsch(d*x+c))^2/f-2*b*f^2*(-c*f+d*e)*(a+b*arccsch(d*x+c))*arctanh(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))/d^4+4*b*(-c*f+d*e)^3*(a+b*arccsch(d*x+c))*arctanh(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))/d^4-1/3*b^2*f^3*ln(d*x+c)/d^4+3*b^2*f*(-c*f+d*e)^2*ln(d*x+c)/d^4-b^2*f^2*(-c*f+d*e)*polylog(2,-1/(d*x+c)-(1+1/(d*x+c)^2)^(1/2))/d^4+2*b^2*(-c*f+d*e)^3*polylog(2,-1/(d*x+c)-(1+1/(d*x+c)^2)^(1/2))/d^4+b^2*f^2*(-c*f+d*e)*polylog(2,1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))/d^4-2*b^2*(-c*f+d*e)^3*polylog(2,1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))/d^4-1/3*b*f^3*(d*x+c)*(a+b*arccsch(d*x+c))*(1+1/(d*x+c)^2)^(1/2)/d^4+3*b*f*(-c*f+d*e)^2*(d*x+c)*(a+b*arccsch(d*x+c))*(1+1/(d*x+c)^2)^(1/2)/d^4+b*f^2*(-c*f+d*e)*(d*x+c)^2*(a+b*arccsch(d*x+c))*(1+1/(d*x+c)^2)^(1/2)/d^4+1/6*b*f^3*(d*x+c)^3*(a+b*arccsch(d*x+c))*(1+1/(d*x+c)^2)^(1/2)/d^4
```

Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 501, normalized size of antiderivative = 1.00, number of steps used = 20, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.450$, Rules used

= {6457, 5577, 4275, 4267, 2317, 2438, 4269, 3556, 4270}

$$\begin{aligned}
& \int (e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx \\
&= -\frac{2bf^2(de - cf) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right) (a + b \operatorname{csch}^{-1}(c + dx))}{d^4} \\
&+ \frac{4b(de - cf)^3 \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right) (a + b \operatorname{csch}^{-1}(c + dx))}{d^4} \\
&+ \frac{bf^2(c + dx)^2 \sqrt{\frac{1}{(c+dx)^2} + 1} (de - cf) (a + b \operatorname{csch}^{-1}(c + dx))}{d^4} \\
&- \frac{(de - cf)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4d^4 f} \\
&+ \frac{3bf(c + dx) \sqrt{\frac{1}{(c+dx)^2} + 1} (de - cf)^2 (a + b \operatorname{csch}^{-1}(c + dx))}{d^4} \\
&+ \frac{bf^3(c + dx)^3 \sqrt{\frac{1}{(c+dx)^2} + 1} (a + b \operatorname{csch}^{-1}(c + dx))}{6d^4} \\
&- \frac{bf^3(c + dx) \sqrt{\frac{1}{(c+dx)^2} + 1} (a + b \operatorname{csch}^{-1}(c + dx))}{3d^4} + \frac{(e + fx)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4f} \\
&- \frac{b^2 f^2 (de - cf) \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} + \frac{b^2 f^2 (de - cf) \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&+ \frac{2b^2 (de - cf)^3 \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} - \frac{2b^2 (de - cf)^3 \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&+ \frac{3b^2 f (de - cf)^2 \log(c + dx)}{d^4} + \frac{b^2 f^3 (c + dx)^2}{12d^4} - \frac{b^2 f^3 \log(c + dx)}{3d^4} + \frac{b^2 f^2 x (de - cf)}{d^3}
\end{aligned}$$

[In] Int[(e + f*x)^3*(a + b*ArcCsch[c + d*x])^2,x]

[Out] (b^2*f^2*(d*e - c*f)*x)/d^3 + (b^2*f^3*(c + d*x)^2)/(12*d^4) - (b*f^3*(c + d*x)*Sqrt[1 + (c + d*x)^(-2)]*(a + b*ArcCsch[c + d*x]))/(3*d^4) + (3*b*f*(d*e - c*f)^2*(c + d*x)*Sqrt[1 + (c + d*x)^(-2)]*(a + b*ArcCsch[c + d*x]))/d^4 + (b*f^2*(d*e - c*f)*(c + d*x)^2*Sqrt[1 + (c + d*x)^(-2)]*(a + b*ArcCsch[c + d*x]))/d^4 + (b*f^3*(c + d*x)^3*Sqrt[1 + (c + d*x)^(-2)]*(a + b*ArcCsch[c + d*x]))/(6*d^4) - ((d*e - c*f)^4*(a + b*ArcCsch[c + d*x])^2)/(4*d^4*f) + ((e + f*x)^4*(a + b*ArcCsch[c + d*x])^2)/(4*f) - (2*b*f^2*(d*e - c*f)*(a + b*ArcCsch[c + d*x])*ArcTanh[E^ArcCsch[c + d*x]])/d^4 + (4*b*(d*e - c*f)^3*(a + b*ArcCsch[c + d*x])*ArcTanh[E^ArcCsch[c + d*x]])/d^4 - (b^2*f^3*Log[c + d*x])/(3*d^4) + (3*b^2*f*(d*e - c*f)^2*Log[c + d*x])/d^4 - (b^2*f^2*(d*e - c*f)*PolyLog[2, -E^ArcCsch[c + d*x]])/d^4 + (2*b^2*(d*e - c*f)^3*PolyLog[2, -E^ArcCsch[c + d*x]])/d^4 + (b^2*f^2*(d*e - c*f)*PolyLog[2, E^ArcCsch[c + d*x]])/d^4 - (2*b^2*(d*e - c*f)^3*PolyLog[2, E^ArcCsch[c + d*x]])/d^4

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x))
)^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3556

```
Int[tan[(c_) + (d_)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 4267

```
Int[csc[(e_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x
_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x]
+ (-Dist[d*(m/(f*fz*I)), Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x]
], x], x] + Dist[d*(m/(f*fz*I)), Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e +
f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

Rule 4269

```
Int[csc[(e_) + (f_)*(x_)]^2*((c_) + (d_)*(x_))^(m_), x_Symbol] :> Simp
[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*
Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 4270

```
Int[(csc[(e_) + (f_)*(x_)]*(b_))^(n_)*((c_) + (d_)*(x_)), x_Symbol] :>
Simp[(-b^2)*(c + d*x)*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))),
x] + (Dist[b^2*((n - 2)/(n - 1)), Int[(c + d*x)*(b*Csc[e + f*x])^(n - 2),
x], x] - Simp[b^2*d*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x]) /
; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2]
```

Rule 4275

```
Int[(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(n_)*((c_) + (d_)*(x_))^(m_)
, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 5577

```
Int[Coth[(c_) + (d_)*(x_)]*Csch[(c_) + (d_)*(x_)]*(Csch[(c_) + (d_)*(
x_)]*(b_) + (a_))^(n_)*((e_) + (f_)*(x_))^(m_), x_Symbol] :> Simp[(-e
```

```

+ f*x)^m)*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b
*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x] /
; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

```

Rule 6457

```

Int[((a_.) + ArcCsch[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] :> Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csch[x]*Co
th[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int (a + bx)^2 \coth(x) \operatorname{csch}(x) (de - cf + f \operatorname{csch}(x))^3 dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^4} \\
&= \frac{(e + fx)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4f} \\
&\quad - \frac{b \text{Subst}\left(\int (a + bx) (de - cf + f \operatorname{csch}(x))^4 dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{2d^4 f} \\
&= \frac{(e + fx)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4f} \\
&\quad - \frac{b \text{Subst}\left(\int \left(d^4 e^4 \left(1 + \frac{cf(-4d^3 e^3 + 6cd^2 e^2 f - 4c^2 def^2 + c^3 f^3)}{d^4 e^4}\right) (a + bx) + 4d^3 e^3 f \left(1 - \frac{cf(3d^2 e^2 - 3cdef + c^2 f^2)}{d^3 e^3}\right)\right) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^4} \\
&= -\frac{(de - cf)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4d^4 f} + \frac{(e + fx)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4f} \\
&\quad - \frac{(bf^3) \text{Subst}\left(\int (a + bx) \operatorname{csch}^4(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{2d^4} \\
&\quad - \frac{(2bf^2(de - cf)) \text{Subst}\left(\int (a + bx) \operatorname{csch}^3(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^4} \\
&\quad - \frac{(3bf(de - cf)^2) \text{Subst}\left(\int (a + bx) \operatorname{csch}^2(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^4} \\
&\quad - \frac{(2b(de - cf)^3) \text{Subst}\left(\int (a + bx) \operatorname{csch}(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^4}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^2 f^2 (de - cf)x}{d^3} + \frac{b^2 f^3 (c + dx)^2}{12d^4} \\
&+ \frac{3bf(de - cf)^2(c + dx)\sqrt{1 + \frac{1}{(c+dx)^2}}(a + b\operatorname{csch}^{-1}(c + dx))}{d^4} \\
&+ \frac{bf^2(de - cf)(c + dx)^2\sqrt{1 + \frac{1}{(c+dx)^2}}(a + b\operatorname{csch}^{-1}(c + dx))}{d^4} \\
&+ \frac{bf^3(c + dx)^3\sqrt{1 + \frac{1}{(c+dx)^2}}(a + b\operatorname{csch}^{-1}(c + dx))}{6d^4} \\
&- \frac{(de - cf)^4(a + b\operatorname{csch}^{-1}(c + dx))^2}{4d^4 f} + \frac{(e + fx)^4(a + b\operatorname{csch}^{-1}(c + dx))^2}{4f} \\
&+ \frac{4b(de - cf)^3(a + b\operatorname{csch}^{-1}(c + dx))\operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&+ \frac{(bf^3)\operatorname{Subst}\left(\int(a + bx)\operatorname{csch}^2(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{3d^4} \\
&+ \frac{(bf^2(de - cf))\operatorname{Subst}\left(\int(a + bx)\operatorname{csch}(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^4} \\
&- \frac{(3b^2 f(de - cf)^2)\operatorname{Subst}\left(\int\coth(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^4} \\
&+ \frac{(2b^2(de - cf)^3)\operatorname{Subst}\left(\int\log(1 - e^x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^4} \\
&- \frac{(2b^2(de - cf)^3)\operatorname{Subst}\left(\int\log(1 + e^x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^4}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^2 f^2 (de - cf)x}{d^3} + \frac{b^2 f^3 (c + dx)^2}{12d^4} - \frac{bf^3 (c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{3d^4} \\
&+ \frac{3bf(de - cf)^2 (c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{d^4} \\
&+ \frac{bf^2 (de - cf)(c + dx)^2 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{d^4} \\
&+ \frac{bf^3 (c + dx)^3 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{6d^4} \\
&- \frac{(de - cf)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4d^4 f} + \frac{(e + fx)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4f} \\
&- \frac{2bf^2 (de - cf) (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&+ \frac{4b(de - cf)^3 (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&+ \frac{3b^2 f (de - cf)^2 \log(c + dx)}{d^4} + \frac{(b^2 f^3) \operatorname{Subst}\left(\int \coth(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{3d^4} \\
&- \frac{(b^2 f^2 (de - cf)) \operatorname{Subst}\left(\int \log(1 - e^x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^4} \\
&+ \frac{(b^2 f^2 (de - cf)) \operatorname{Subst}\left(\int \log(1 + e^x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^4} \\
&+ \frac{(2b^2 (de - cf)^3) \operatorname{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&- \frac{(2b^2 (de - cf)^3) \operatorname{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^2 f^2 (de - cf)x}{d^3} + \frac{b^2 f^3 (c + dx)^2}{12d^4} - \frac{bf^3 (c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{3d^4} \\
&+ \frac{3bf^2 (de - cf)^2 (c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{d^4} \\
&+ \frac{bf^2 (de - cf)(c + dx)^2 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{d^4} \\
&+ \frac{bf^3 (c + dx)^3 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{6d^4} \\
&- \frac{(de - cf)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4d^4 f} + \frac{(e + fx)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4f} \\
&- \frac{2bf^2 (de - cf) (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&+ \frac{4b(de - cf)^3 (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} - \frac{b^2 f^3 \log(c + dx)}{3d^4} \\
&+ \frac{3b^2 f (de - cf)^2 \log(c + dx)}{d^4} + \frac{2b^2 (de - cf)^3 \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&- \frac{2b^2 (de - cf)^3 \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&- \frac{(b^2 f^2 (de - cf)) \operatorname{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&+ \frac{(b^2 f^2 (de - cf)) \operatorname{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^2 f^2 (de - cf)x}{d^3} + \frac{b^2 f^3 (c + dx)^2}{12d^4} - \frac{bf^3 (c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{3d^4} \\
&+ \frac{3bf(de - cf)^2 (c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{d^4} \\
&+ \frac{bf^2 (de - cf)(c + dx)^2 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{d^4} \\
&+ \frac{bf^3 (c + dx)^3 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{6d^4} \\
&- \frac{(de - cf)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4d^4 f} + \frac{(e + fx)^4 (a + b \operatorname{csch}^{-1}(c + dx))^2}{4f} \\
&- \frac{2bf^2 (de - cf) (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&+ \frac{4b(de - cf)^3 (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} - \frac{b^2 f^3 \log(c + dx)}{3d^4} \\
&+ \frac{3b^2 f (de - cf)^2 \log(c + dx)}{d^4} - \frac{b^2 f^2 (de - cf) \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&+ \frac{2b^2 (de - cf)^3 \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&+ \frac{b^2 f^2 (de - cf) \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4} \\
&- \frac{2b^2 (de - cf)^3 \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^4}
\end{aligned}$$

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 13.06 (sec) , antiderivative size = 1487, normalized size of antiderivative = 2.97

$$\begin{aligned}
& \int (e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = a^2 e^3 x + \frac{3}{2} a^2 e^2 f x^2 + a^2 e f^2 x^3 \\
& + \frac{1}{4} a^2 f^3 x^4 + \frac{1}{6} ab \left(3x(4e^3 + 6e^2 f x + 4e f^2 x^2 + f^3 x^3) \operatorname{csch}^{-1}(c + dx) \right. \\
& \left. + \frac{f(c + dx) \sqrt{\frac{1+c^2+2cdx+d^2x^2}{(c+dx)^2}} ((-2 + 13c^2) f^2 - 2cdf(15e + 2fx) + d^2(18e^2 + 6efx + f^2x^2)) - 3c(-4d^3e^3}{d} \right. \\
& \left. + \frac{b^2 e^3 \left(-\operatorname{csch}^{-1}(c + dx) \left((c + dx) \operatorname{csch}^{-1}(c + dx) - 2 \log \left(1 - e^{-\operatorname{csch}^{-1}(c + dx)} \right) + 2 \log \left(1 + e^{-\operatorname{csch}^{-1}(c + dx)} \right) \right) \right)}{d} \right. \\
& \left. + \frac{3b^2 d e^2 f x \left(\frac{(c+dx) \sqrt{1 + \frac{1}{(c+dx)^2}} \operatorname{csch}^{-1}(c+dx)}{d^2} + \frac{(c+dx)^2 \operatorname{csch}^{-1}(c+dx)^2}{2d^2} - \frac{c \operatorname{csch}^{-1}(c+dx)^2 \operatorname{coth}(\frac{1}{2} \operatorname{csch}^{-1}(c+dx))}{2d^2} - \frac{\log(\frac{c}{c+dx})}{d^2} \right)}{d} \right. \\
& \left. + \frac{b^2 e f^2 \left(2(-2 + 12c \operatorname{csch}^{-1}(c + dx) + \operatorname{csch}^{-1}(c + dx)^2 - 6c^2 \operatorname{csch}^{-1}(c + dx)^2) \operatorname{coth}(\frac{1}{2} \operatorname{csch}^{-1}(c + dx)) + 2 \right)}{d} \right. \\
& \left. + \frac{b^2 f^3 x^3 \left(-16(2 \operatorname{csch}^{-1}(c + dx) - 18c^2 \operatorname{csch}^{-1}(c + dx) + 6c^3 \operatorname{csch}^{-1}(c + dx)^2 - 3c(-2 + \operatorname{csch}^{-1}(c + dx))^2) \right)}{d} \right)
\end{aligned}$$

`[In] Integrate[(e + f*x)^3*(a + b*ArcCsch[c + d*x])^2,x]`

```

[Out] a^2*e^3*x + (3*a^2*e^2*f*x^2)/2 + a^2*e*f^2*x^3 + (a^2*f^3*x^4)/4 + (a*b*(3
*x*(4*e^3 + 6*e^2*f*x + 4*e*f^2*x^2 + f^3*x^3)*ArcCsch[c + d*x] + (f*(c + d
*x)*Sqrt[(1 + c^2 + 2*c*d*x + d^2*x^2)/(c + d*x)^2]*((-2 + 13*c^2)*f^2 - 2*
c*d*f*(15*e + 2*f*x) + d^2*(18*e^2 + 6*e*f*x + f^2*x^2)) - 3*c*(-4*d^3*e^3
+ 6*c*d^2*e^2*f - 4*c^2*d*e*f^2 + c^3*f^3)*ArcSinh[(c + d*x)^(-1)] + 6*(2*d
^3*e^3 - 6*c*d^2*e^2*f + (-1 + 6*c^2)*d*e*f^2 + c*(1 - 2*c^2)*f^3)*Log[(c +
d*x)*(1 + Sqrt[(1 + c^2 + 2*c*d*x + d^2*x^2)/(c + d*x)^2])]/d^4))/6 - (b^
2*e^3*(-(ArcCsch[c + d*x]*((c + d*x)*ArcCsch[c + d*x] - 2*Log[1 - E^(-ArcCs
ch[c + d*x])]) + 2*Log[1 + E^(-ArcCsch[c + d*x])])) + 2*PolyLog[2, -E^(-ArcC
sch[c + d*x])] - 2*PolyLog[2, E^(-ArcCsch[c + d*x])]))/d - (3*b^2*d*e^2*f*x
*(((c + d*x)*Sqrt[1 + (c + d*x)^(-2)]*ArcCsch[c + d*x])/d^2 + ((c + d*x)^2*
ArcCsch[c + d*x]^2)/(2*d^2) - (c*ArcCsch[c + d*x]^2*Coth[ArcCsch[c + d*x]/2
])/d^2 - Log[(c + d*x)^(-1)]/d^2 - ((2*I)*c*(I*ArcCsch[c + d*x]*(Log[1
- E^(-ArcCsch[c + d*x])]) - Log[1 + E^(-ArcCsch[c + d*x])])) + I*(PolyLog[2,
-E^(-ArcCsch[c + d*x])]) - PolyLog[2, E^(-ArcCsch[c + d*x])])))/d^2 + (c*Arc

```

```

Csch[c + d*x]^2*Tanh[ArcCsch[c + d*x]/2])/(2*d^2)))/((c + d*x)*(-1 + c/(c +
d*x))) - (b^2*e*f^2*(2*(-2 + 12*c*ArcCsch[c + d*x] + ArcCsch[c + d*x]^2 -
6*c^2*ArcCsch[c + d*x]^2)*Coth[ArcCsch[c + d*x]/2] + 2*ArcCsch[c + d*x]*(-1
+ 3*c*ArcCsch[c + d*x])*Csch[ArcCsch[c + d*x]/2]^2 - (ArcCsch[c + d*x]^2*C
sch[ArcCsch[c + d*x]/2]^4)/(2*(c + d*x)) - 48*c*(Log[1/((c + d*x)*Sqrt[1 +
(c + d*x)^(-2)]]) + Log[Sqrt[1 + (c + d*x)^(-2)]]) + 8*(-1 + 6*c^2)*(ArcCsc
h[c + d*x]*(Log[1 - E^(-ArcCsch[c + d*x]]) - Log[1 + E^(-ArcCsch[c + d*x]])
) + PolyLog[2, -E^(-ArcCsch[c + d*x]]) - PolyLog[2, E^(-ArcCsch[c + d*x]])
) - 2*ArcCsch[c + d*x]*(1 + 3*c*ArcCsch[c + d*x])*Sech[ArcCsch[c + d*x]/2]^2
- 8*(c + d*x)^3*ArcCsch[c + d*x]^2*Sinh[ArcCsch[c + d*x]/2]^4 + 2*(2 + 12*
c*ArcCsch[c + d*x] - ArcCsch[c + d*x]^2 + 6*c^2*ArcCsch[c + d*x]^2)*Tanh[Arc
Csch[c + d*x]/2]))/(8*d^3) - (b^2*f^3*x^3*(-16*(2*ArcCsch[c + d*x] - 18*c^
2*ArcCsch[c + d*x] + 6*c^3*ArcCsch[c + d*x]^2 - 3*c*(-2 + ArcCsch[c + d*x]^
2))*Coth[ArcCsch[c + d*x]/2] + 2*(2 - 24*c*ArcCsch[c + d*x] - 3*ArcCsch[c +
d*x]^2 + 36*c^2*ArcCsch[c + d*x]^2)*Csch[ArcCsch[c + d*x]/2]^2 + 3*ArcCsch
[c + d*x]^2*Csch[ArcCsch[c + d*x]/2]^4 - (2*ArcCsch[c + d*x]*(-1 + 6*c*ArcC
sch[c + d*x])*Csch[ArcCsch[c + d*x]/2]^4)/(c + d*x) - 64*(-1 + 9*c^2)*(Log[
1/((c + d*x)*Sqrt[1 + (c + d*x)^(-2)]]) + Log[Sqrt[1 + (c + d*x)^(-2)]]) +
192*c*(-1 + 2*c^2)*(ArcCsch[c + d*x]*(Log[1 - E^(-ArcCsch[c + d*x]]) - Log[
1 + E^(-ArcCsch[c + d*x]]) + PolyLog[2, -E^(-ArcCsch[c + d*x]]) - PolyLog[
2, E^(-ArcCsch[c + d*x]]) - 2*(2 + 24*c*ArcCsch[c + d*x] - 3*ArcCsch[c + d
*x]^2 + 36*c^2*ArcCsch[c + d*x]^2)*Sech[ArcCsch[c + d*x]/2]^2 + 3*ArcCsch[c
+ d*x]^2*Sech[ArcCsch[c + d*x]/2]^4 - 32*(c + d*x)^3*ArcCsch[c + d*x]*(1 +
6*c*ArcCsch[c + d*x])*Sinh[ArcCsch[c + d*x]/2]^4 + 16*(-2*ArcCsch[c + d*x]
+ 18*c^2*ArcCsch[c + d*x] + 6*c^3*ArcCsch[c + d*x]^2 - 3*c*(-2 + ArcCsch[c
+ d*x]^2))*Tanh[ArcCsch[c + d*x]/2]))/(192*d*(c + d*x)^3*(-1 + c/(c + d*x)
)^3)

```

Maple [F]

$$\int (fx + e)^3 (a + b \operatorname{arccsch}(dx + c))^2 dx$$

```
[In] int((f*x+e)^3*(a+b*arccsch(d*x+c))^2,x)
```

```
[Out] int((f*x+e)^3*(a+b*arccsch(d*x+c))^2,x)
```

Fricas [F]

$$\int (e + fx)^3 (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (fx + e)^3 (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

[In] integrate((f*x+e)^3*(a+b*arcsch(d*x+c))^2,x, algorithm="fricas")

[Out] integral(a^2*f^3*x^3 + 3*a^2*e*f^2*x^2 + 3*a^2*e^2*f*x + a^2*e^3 + (b^2*f^3*x^3 + 3*b^2*e*f^2*x^2 + 3*b^2*e^2*f*x + b^2*e^3)*arcsch(d*x + c)^2 + 2*(a*b*f^3*x^3 + 3*a*b*e*f^2*x^2 + 3*a*b*e^2*f*x + a*b*e^3)*arcsch(d*x + c), x)

Sympy [F]

$$\int (e + fx)^3 (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (a + b \operatorname{arcsch}(c + dx))^2 (e + fx)^3 dx$$

[In] integrate((f*x+e)**3*(a+b*arcsch(d*x+c))**2,x)

[Out] Integral((a + b*arcsch(c + d*x))**2*(e + f*x)**3, x)

Maxima [F]

$$\int (e + fx)^3 (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (fx + e)^3 (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

[In] integrate((f*x+e)^3*(a+b*arcsch(d*x+c))^2,x, algorithm="maxima")

[Out] 1/4*a^2*f^3*x^4 + a^2*e*f^2*x^3 + 3/2*a^2*e^2*f*x^2 + a^2*e^3*x + (2*(d*x + c)*arcsch(d*x + c) + log(sqrt(1/(d*x + c)^2 + 1) + 1) - log(sqrt(1/(d*x + c)^2 + 1) - 1))*a*b*e^3/d + 1/4*(b^2*f^3*x^4 + 4*b^2*e*f^2*x^3 + 6*b^2*e^2*f*x^2 + 4*b^2*e^3*x)*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1)^2 - integrate(-1/2*(2*(b^2*d^2*f^3*x^5 + b^2*c^2*e^3 + b^2*e^3 + (3*b^2*d^2*e*f^2 + 2*b^2*c*d*f^3)*x^4 + (6*b^2*c*d*e*f^2 + b^2*c^2*f^3 + (3*d^2*e^2*f + f^3)*b^2)*x^3 + (6*b^2*c*d*e^2*f + 3*b^2*c^2*e*f^2 + (d^2*e^3 + 3*e*f^2)*b^2)*x^2 + (2*b^2*c*d*e^3 + 3*b^2*c^2*e^2*f + 3*b^2*e^2*f)*x)*log(d*x + c)^2 - 4*(a*b*d^2*f^3*x^5 + (3*a*b*d^2*e*f^2 + 2*a*b*c*d*f^3)*x^4 + (6*a*b*c*d*e*f^2 + a*b*c^2*f^3 + (3*d^2*e^2*f + f^3)*a*b)*x^3 + 3*(2*a*b*c*d*e^2*f + a*b*c^2*e*f^2 + a*b*e*f^2)*x^2 + 3*(a*b*c^2*e^2*f + a*b*e^2*f)*x)*log(d*x + c) + (4*a*b*d^2*f^3*x^5 + 4*(3*a*b*d^2*e*f^2 + 2*a*b*c*d*f^3)*x^4 + 4*(6*a*b*c*d*e*f^2 + a*b*c^2*f^3 + (3*d^2*e^2*f + f^3)*a*b)*x^3 + 12*(2*a*b*c*d*e^2*f + a*b*c^2*e*f^2 + a*b*e*f^2)*x^2 + 12*(a*b*c^2*e^2*f + a*b*e^2*f)*x - 4*(b^2*d^2

$$\begin{aligned}
& 2*f^3*x^5 + b^2*c^2*e^3 + b^2*e^3 + (3*b^2*d^2*e*f^2 + 2*b^2*c*d*f^3)*x^4 + \\
& (6*b^2*c*d*e*f^2 + b^2*c^2*f^3 + (3*d^2*e^2*f + f^3)*b^2)*x^3 + (6*b^2*c*d \\
& *e^2*f + 3*b^2*c^2*e*f^2 + (d^2*e^3 + 3*e*f^2)*b^2)*x^2 + (2*b^2*c*d*e^3 + \\
& 3*b^2*c^2*e^2*f + 3*b^2*e^2*f)*x)*\log(d*x + c) + ((4*a*b*d^2*f^3 - b^2*d^2* \\
& f^3)*x^5 + (12*a*b*d^2*e*f^2 - 4*b^2*d^2*e*f^2 + (8*a*b*d*f^3 - b^2*d*f^3)* \\
& c)*x^4 - 2*(3*b^2*d^2*e^2*f - 2*a*b*c^2*f^3 - 2*(3*d^2*e^2*f + f^3)*a*b - 2 \\
& *(6*a*b*d*e*f^2 - b^2*d*e*f^2)*c)*x^3 - 2*(2*b^2*d^2*e^3 - 6*a*b*c^2*e*f^2 \\
& - 6*a*b*e*f^2 - 3*(4*a*b*d*e^2*f - b^2*d*e^2*f)*c)*x^2 - 4*(b^2*c*d*e^3 - 3 \\
& *a*b*c^2*e^2*f - 3*a*b*e^2*f)*x - 4*(b^2*d^2*f^3*x^5 + b^2*c^2*e^3 + b^2*e^3 \\
& + (3*b^2*d^2*e*f^2 + 2*b^2*c*d*f^3)*x^4 + (6*b^2*c*d*e*f^2 + b^2*c^2*f^3 \\
& + (3*d^2*e^2*f + f^3)*b^2)*x^3 + (6*b^2*c*d*e^2*f + 3*b^2*c^2*e*f^2 + (d^2* \\
& e^3 + 3*e*f^2)*b^2)*x^2 + (2*b^2*c*d*e^3 + 3*b^2*c^2*e^2*f + 3*b^2*e^2*f)*x \\
&)*\log(d*x + c))*\sqrt{d^2*x^2 + 2*c*d*x + c^2 + 1})*\log(\sqrt{d^2*x^2 + 2*c*d \\
& *x + c^2 + 1} + 1) + 2*\sqrt{d^2*x^2 + 2*c*d*x + c^2 + 1})*((b^2*d^2*f^3*x^5 \\
& + b^2*c^2*e^3 + b^2*e^3 + (3*b^2*d^2*e*f^2 + 2*b^2*c*d*f^3)*x^4 + (6*b^2*c* \\
& d*e*f^2 + b^2*c^2*f^3 + (3*d^2*e^2*f + f^3)*b^2)*x^3 + (6*b^2*c*d*e^2*f + 3 \\
& *b^2*c^2*e*f^2 + (d^2*e^3 + 3*e*f^2)*b^2)*x^2 + (2*b^2*c*d*e^3 + 3*b^2*c^2* \\
& e^2*f + 3*b^2*e^2*f)*x)*\log(d*x + c))^2 - 2*(a*b*d^2*f^3*x^5 + (3*a*b*d^2*e* \\
& f^2 + 2*a*b*c*d*f^3)*x^4 + (6*a*b*c*d*e*f^2 + a*b*c^2*f^3 + (3*d^2*e^2*f + \\
& f^3)*a*b)*x^3 + 3*(2*a*b*c*d*e^2*f + a*b*c^2*e*f^2 + a*b*e*f^2)*x^2 + 3*(a* \\
& b*c^2*e^2*f + a*b*e^2*f)*x)*\log(d*x + c))/(d^2*x^2 + 2*c*d*x + c^2 + (d^2* \\
& x^2 + 2*c*d*x + c^2 + 1)^{(3/2)} + 1), x)
\end{aligned}$$

Giac [F]

$$\int (e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \int (fx + e)^3 (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

[In] integrate((f*x+e)^3*(a+b*arccsch(d*x+c))^2,x, algorithm="giac")

[Out] integrate((f*x + e)^3*(b*arccsch(d*x + c) + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int (e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \int (e + fx)^3 \left(a + b \operatorname{asinh}\left(\frac{1}{c + dx}\right) \right)^2 dx$$

[In] int((e + f*x)^3*(a + b*asinh(1/(c + d*x)))^2,x)

[Out] int((e + f*x)^3*(a + b*asinh(1/(c + d*x)))^2, x)

3.8 $\int (e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$

Optimal result	95
Rubi [A] (verified)	96
Mathematica [C] (warning: unable to verify)	101
Maple [F]	102
Fricas [F]	102
Sympy [F]	102
Maxima [F]	103
Giac [F]	103
Mupad [F(-1)]	104

Optimal result

Integrand size = 20, antiderivative size = 351

$$\begin{aligned}
 & \int (e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx \\
 &= \frac{b^2 f^2 x}{3d^2} + \frac{2bf(de - cf)(c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{d^3} \\
 &+ \frac{bf^2(c + dx)^2 \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{3d^3} - \frac{(de - cf)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2}{3d^3 f} \\
 &+ \frac{(e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2}{3f} - \frac{2bf^2 (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}(e^{\operatorname{csch}^{-1}(c+dx)})}{3d^3} \\
 &+ \frac{4b(de - cf)^2 (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}(e^{\operatorname{csch}^{-1}(c+dx)})}{d^3} \\
 &+ \frac{2b^2 f(de - cf) \log(c + dx)}{d^3} - \frac{b^2 f^2 \operatorname{PolyLog}(2, -e^{\operatorname{csch}^{-1}(c+dx)})}{3d^3} \\
 &+ \frac{2b^2 (de - cf)^2 \operatorname{PolyLog}(2, -e^{\operatorname{csch}^{-1}(c+dx)})}{d^3} \\
 &+ \frac{b^2 f^2 \operatorname{PolyLog}(2, e^{\operatorname{csch}^{-1}(c+dx)})}{3d^3} - \frac{2b^2 (de - cf)^2 \operatorname{PolyLog}(2, e^{\operatorname{csch}^{-1}(c+dx)})}{d^3}
 \end{aligned}$$

```
[Out] 1/3*b^2*f^2*x/d^2-1/3*(-c*f+d*e)^3*(a+b*arccsch(d*x+c))^2/d^3/f+1/3*(f*x+e)
^3*(a+b*arccsch(d*x+c))^2/f-2/3*b*f^2*(a+b*arccsch(d*x+c))*arctanh(1/(d*x+c)
)+(1+1/(d*x+c)^2)^(1/2))/d^3+4*b*(-c*f+d*e)^2*(a+b*arccsch(d*x+c))*arctanh(
1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))/d^3+2*b^2*f*(-c*f+d*e)*ln(d*x+c)/d^3-1/3*b
^2*f^2*polylog(2,-1/(d*x+c)-(1+1/(d*x+c)^2)^(1/2))/d^3+2*b^2*(-c*f+d*e)^2*p
olylog(2,-1/(d*x+c)-(1+1/(d*x+c)^2)^(1/2))/d^3+1/3*b^2*f^2*polylog(2,1/(d*x
```

$$+c)+(1+1/(d*x+c)^2)^{(1/2)}/d^3-2*b^2*(-c*f+d*e)^2*\text{polylog}(2,1/(d*x+c)+(1+1/(d*x+c)^2)^{(1/2)})/d^3+2*b*f*(-c*f+d*e)*(d*x+c)*(a+b*\text{arccsch}(d*x+c))*(1+1/(d*x+c)^2)^{(1/2)}/d^3+1/3*b*f^2*(d*x+c)^2*(a+b*\text{arccsch}(d*x+c))*(1+1/(d*x+c)^2)^{(1/2)}/d^3$$

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 351, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.450$, Rules used = {6457, 5577, 4275, 4267, 2317, 2438, 4269, 3556, 4270}

$$\int (e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$$

$$= \frac{4b(de - cf)^2 \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right) (a + b \operatorname{csch}^{-1}(c + dx))}{d^3}$$

$$- \frac{2bf^2 \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right) (a + b \operatorname{csch}^{-1}(c + dx))}{3d^3} - \frac{(de - cf)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2}{3d^3 f}$$

$$+ \frac{2bf(c + dx) \sqrt{\frac{1}{(c+dx)^2} + 1} (de - cf) (a + b \operatorname{csch}^{-1}(c + dx))}{d^3}$$

$$+ \frac{bf^2(c + dx)^2 \sqrt{\frac{1}{(c+dx)^2} + 1} (a + b \operatorname{csch}^{-1}(c + dx))}{3d^3}$$

$$+ \frac{(e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2}{3f} + \frac{2b^2(de - cf)^2 \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^3}$$

$$- \frac{2b^2(de - cf)^2 \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^3} + \frac{2b^2 f (de - cf) \log(c + dx)}{d^3}$$

$$- \frac{b^2 f^2 \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{3d^3} + \frac{b^2 f^2 \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{3d^3} + \frac{b^2 f^2 x}{3d^2}$$

[In] Int[(e + f*x)^2*(a + b*ArcCsch[c + d*x])^2,x]

[Out] (b^2*f^2*x)/(3*d^2) + (2*b*f*(d*e - c*f)*(c + d*x)*Sqrt[1 + (c + d*x)^(-2)]*(a + b*ArcCsch[c + d*x]))/d^3 + (b*f^2*(c + d*x)^2*Sqrt[1 + (c + d*x)^(-2)]*(a + b*ArcCsch[c + d*x]))/(3*d^3) - ((d*e - c*f)^3*(a + b*ArcCsch[c + d*x])^2)/(3*d^3*f) + ((e + f*x)^3*(a + b*ArcCsch[c + d*x])^2)/(3*f) - (2*b*f^2*(a + b*ArcCsch[c + d*x])*ArcTanh[E^ArcCsch[c + d*x]])/(3*d^3) + (4*b*(d*e - c*f)^2*(a + b*ArcCsch[c + d*x])*ArcTanh[E^ArcCsch[c + d*x]])/d^3 + (2*b^2*f*(d*e - c*f)*Log[c + d*x])/d^3 - (b^2*f^2*PolyLog[2, -E^ArcCsch[c + d*x]])/(3*d^3) + (2*b^2*(d*e - c*f)^2*PolyLog[2, -E^ArcCsch[c + d*x]])/d^3 + (b^2*f^2*PolyLog[2, E^ArcCsch[c + d*x]])/(3*d^3) - (2*b^2*(d*e - c*f)^2*PolyLog[2, E^ArcCsch[c + d*x]])/d^3

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x))
)^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3556

```
Int[tan[(c_) + (d_)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 4267

```
Int[csc[(e_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x
_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x]
+ (-Dist[d*(m/(f*fz*I)), Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x]
], x], x] + Dist[d*(m/(f*fz*I)), Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e +
f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

Rule 4269

```
Int[csc[(e_) + (f_)*(x_)]^2*((c_) + (d_)*(x_))^(m_), x_Symbol] :> Simp
[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*
Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 4270

```
Int[(csc[(e_) + (f_)*(x_)]*(b_))^(n_)*((c_) + (d_)*(x_)), x_Symbol] :>
Simp[(-b^2)*(c + d*x)*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))),
x] + (Dist[b^2*((n - 2)/(n - 1)), Int[(c + d*x)*(b*Csc[e + f*x])^(n - 2),
x], x] - Simp[b^2*d*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x]) /
; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2]
```

Rule 4275

```
Int[(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(n_)*((c_) + (d_)*(x_))^(m_)
, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 5577

```
Int[Coth[(c_) + (d_)*(x_)]*Csch[(c_) + (d_)*(x_)]*(Csch[(c_) + (d_)*(
x_)]*(b_) + (a_))^(n_)*((e_) + (f_)*(x_))^(m_), x_Symbol] :> Simp[(-e
```

```

+ f*x)^m)*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b
*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x] /
; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

```

Rule 6457

```

Int[((a_.) + ArcCsch[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] :> Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csch[x]*Co
th[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int (a + bx)^2 \coth(x) \operatorname{csch}(x) (de - cf + f \operatorname{csch}(x))^2 dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^3} \\
&= \frac{(e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2}{3f} \\
&\quad - \frac{(2b) \text{Subst}\left(\int (a + bx) (de - cf + f \operatorname{csch}(x))^3 dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{3d^3 f} \\
&= \frac{(e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2}{3f} \\
&\quad - \frac{(2b) \text{Subst}\left(\int \left(d^3 e^3 \left(1 - \frac{cf(3d^2 e^2 - 3cdef + c^2 f^2)}{d^3 e^3}\right) (a + bx) + 3d^2 e^2 f \left(1 + \frac{cf(-2de + cf)}{d^2 e^2}\right) (a + bx) \operatorname{csch}(x)\right) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{3d^3 f} \\
&= -\frac{(de - cf)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2}{3d^3 f} + \frac{(e + fx)^3 (a + b \operatorname{csch}^{-1}(c + dx))^2}{3f} \\
&\quad - \frac{(2bf^2) \text{Subst}\left(\int (a + bx) \operatorname{csch}^3(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{3d^3} \\
&\quad - \frac{(2bf(de - cf)) \text{Subst}\left(\int (a + bx) \operatorname{csch}^2(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^3} \\
&\quad - \frac{(2b(de - cf)^2) \text{Subst}\left(\int (a + bx) \operatorname{csch}(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^2 f^2 x}{3d^2} + \frac{2bf(de - cf)(c + dx)\sqrt{1 + \frac{1}{(c+dx)^2}}(a + b\operatorname{csch}^{-1}(c + dx))}{d^3} \\
&\quad + \frac{bf^2(c + dx)^2\sqrt{1 + \frac{1}{(c+dx)^2}}(a + b\operatorname{csch}^{-1}(c + dx))}{3d^3} \\
&\quad - \frac{(de - cf)^3(a + b\operatorname{csch}^{-1}(c + dx))^2}{3d^3 f} + \frac{(e + fx)^3(a + b\operatorname{csch}^{-1}(c + dx))^2}{3f} \\
&\quad + \frac{4b(de - cf)^2(a + b\operatorname{csch}^{-1}(c + dx))\operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^3} \\
&\quad + \frac{(bf^2)\operatorname{Subst}\left(\int(a + bx)\operatorname{csch}(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{3d^3} \\
&\quad - \frac{(2b^2 f(de - cf))\operatorname{Subst}\left(\int\coth(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^3} \\
&\quad + \frac{(2b^2(de - cf)^2)\operatorname{Subst}\left(\int\log(1 - e^x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^3} \\
&\quad - \frac{(2b^2(de - cf)^2)\operatorname{Subst}\left(\int\log(1 + e^x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^3} \\
&= \frac{b^2 f^2 x}{3d^2} + \frac{2bf(de - cf)(c + dx)\sqrt{1 + \frac{1}{(c+dx)^2}}(a + b\operatorname{csch}^{-1}(c + dx))}{d^3} \\
&\quad + \frac{bf^2(c + dx)^2\sqrt{1 + \frac{1}{(c+dx)^2}}(a + b\operatorname{csch}^{-1}(c + dx))}{3d^3} \\
&\quad - \frac{(de - cf)^3(a + b\operatorname{csch}^{-1}(c + dx))^2}{3d^3 f} + \frac{(e + fx)^3(a + b\operatorname{csch}^{-1}(c + dx))^2}{3f} \\
&\quad - \frac{2bf^2(a + b\operatorname{csch}^{-1}(c + dx))\operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{3d^3} \\
&\quad + \frac{4b(de - cf)^2(a + b\operatorname{csch}^{-1}(c + dx))\operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^3} \\
&\quad + \frac{2b^2 f(de - cf)\log(c + dx)}{d^3} - \frac{(b^2 f^2)\operatorname{Subst}\left(\int\log(1 - e^x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{3d^3} \\
&\quad + \frac{(b^2 f^2)\operatorname{Subst}\left(\int\log(1 + e^x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{3d^3} \\
&\quad + \frac{(2b^2(de - cf)^2)\operatorname{Subst}\left(\int\frac{\log(1-x)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^3} \\
&\quad - \frac{(2b^2(de - cf)^2)\operatorname{Subst}\left(\int\frac{\log(1+x)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^2 f^2 x}{3d^2} + \frac{2bf(de - cf)(c + dx)\sqrt{1 + \frac{1}{(c+dx)^2}}(a + b\operatorname{csch}^{-1}(c + dx))}{d^3} \\
&+ \frac{bf^2(c + dx)^2\sqrt{1 + \frac{1}{(c+dx)^2}}(a + b\operatorname{csch}^{-1}(c + dx))}{3d^3} \\
&- \frac{(de - cf)^3(a + b\operatorname{csch}^{-1}(c + dx))^2}{3d^3 f} + \frac{(e + fx)^3(a + b\operatorname{csch}^{-1}(c + dx))^2}{3f} \\
&- \frac{2bf^2(a + b\operatorname{csch}^{-1}(c + dx))\operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{3d^3} \\
&+ \frac{4b(de - cf)^2(a + b\operatorname{csch}^{-1}(c + dx))\operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^3} \\
&+ \frac{2b^2 f(de - cf)\log(c + dx)}{d^3} + \frac{2b^2(de - cf)^2\operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^3} \\
&- \frac{2b^2(de - cf)^2\operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^3} \\
&- \frac{(b^2 f^2)\operatorname{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{3d^3} \\
&+ \frac{(b^2 f^2)\operatorname{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{3d^3} \\
&= \frac{b^2 f^2 x}{3d^2} + \frac{2bf(de - cf)(c + dx)\sqrt{1 + \frac{1}{(c+dx)^2}}(a + b\operatorname{csch}^{-1}(c + dx))}{d^3} \\
&+ \frac{bf^2(c + dx)^2\sqrt{1 + \frac{1}{(c+dx)^2}}(a + b\operatorname{csch}^{-1}(c + dx))}{3d^3} \\
&- \frac{(de - cf)^3(a + b\operatorname{csch}^{-1}(c + dx))^2}{3d^3 f} + \frac{(e + fx)^3(a + b\operatorname{csch}^{-1}(c + dx))^2}{3f} \\
&- \frac{2bf^2(a + b\operatorname{csch}^{-1}(c + dx))\operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{3d^3} \\
&+ \frac{4b(de - cf)^2(a + b\operatorname{csch}^{-1}(c + dx))\operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^3} \\
&+ \frac{2b^2 f(de - cf)\log(c + dx)}{d^3} - \frac{b^2 f^2\operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{3d^3} \\
&+ \frac{2b^2(de - cf)^2\operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^3} \\
&+ \frac{b^2 f^2\operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{3d^3} - \frac{2b^2(de - cf)^2\operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^3}
\end{aligned}$$

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 7.97 (sec) , antiderivative size = 893, normalized size of antiderivative = 2.54

$$\begin{aligned}
& \int (e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2 dx \\
&= a^2 e^2 x + a^2 e f x^2 + \frac{1}{3} a^2 f^2 x^3 + \frac{1}{3} ab \left(2x(3e^2 + 3efx + f^2 x^2) \operatorname{csch}^{-1}(c + dx) \right. \\
&\quad \left. + \frac{-f(c + dx) \sqrt{\frac{1+c^2+2cdx+d^2x^2}{(c+dx)^2}} (5cf - d(6e + fx)) + 2c(3d^2 e^2 - 3cdef + c^2 f^2) \operatorname{arcsinh}\left(\frac{1}{c+dx}\right) + (6d^2 e^2 -}{d^3} \right. \\
&\quad \left. b^2 e^2 \left(-\operatorname{csch}^{-1}(c + dx) \left((c + dx) \operatorname{csch}^{-1}(c + dx) - 2 \log \left(1 - e^{-\operatorname{csch}^{-1}(c+dx)} \right) + 2 \log \left(1 + e^{-\operatorname{csch}^{-1}(c+dx)} \right) \right) \right)}{d} \right. \\
&\quad \left. 2b^2 defx \left(\frac{(c+dx) \sqrt{1 + \frac{1}{(c+dx)^2}} \operatorname{csch}^{-1}(c+dx)}{d^2} + \frac{(c+dx)^2 \operatorname{csch}^{-1}(c+dx)^2}{2d^2} - \frac{c \operatorname{csch}^{-1}(c+dx)^2 \operatorname{coth}\left(\frac{1}{2} \operatorname{csch}^{-1}(c+dx)\right)}{2d^2} - \frac{\log\left(\frac{1}{c+dx}\right)}{d^2} \right) \right. \\
&\quad \left. b^2 f^2 \left(2(-2 + 12c \operatorname{csch}^{-1}(c + dx) + \operatorname{csch}^{-1}(c + dx)^2 - 6c^2 \operatorname{csch}^{-1}(c + dx)^2) \operatorname{coth}\left(\frac{1}{2} \operatorname{csch}^{-1}(c + dx)\right) + 2 \right) \right)
\end{aligned}$$

[In] Integrate[(e + f*x)^2*(a + b*ArcCsch[c + d*x])^2,x]

```

[Out] a^2*e^2*x + a^2*e*f*x^2 + (a^2*f^2*x^3)/3 + (a*b*(2*x*(3*e^2 + 3*e*f*x + f^2*x^2)*ArcCsch[c + d*x] + (-f*(c + d*x)*Sqrt[(1 + c^2 + 2*c*d*x + d^2*x^2)/(c + d*x)^2]*(5*c*f - d*(6*e + f*x))) + 2*c*(3*d^2*e^2 - 3*c*d*e*f + c^2*f^2)*ArcSinh[(c + d*x)^(-1)] + (6*d^2*e^2 - 12*c*d*e*f + (-1 + 6*c^2)*f^2)*Log[(c + d*x)*(1 + Sqrt[(1 + c^2 + 2*c*d*x + d^2*x^2)/(c + d*x)^2])]/d^3))/3 - (b^2*e^2*(-ArcCsch[c + d*x]*((c + d*x)*ArcCsch[c + d*x] - 2*Log[1 - E^(-ArcCsch[c + d*x])]) + 2*Log[1 + E^(-ArcCsch[c + d*x])])) + 2*PolyLog[2, -E^(-ArcCsch[c + d*x])] - 2*PolyLog[2, E^(-ArcCsch[c + d*x])])/d - (2*b^2*d*e*f*x*((c + d*x)*Sqrt[1 + (c + d*x)^(-2)]*ArcCsch[c + d*x])/d^2 + ((c + d*x)^2*ArcCsch[c + d*x]^2)/(2*d^2) - (c*ArcCsch[c + d*x]^2*Coth[ArcCsch[c + d*x]/2])/(2*d^2) - Log[(c + d*x)^(-1)]/d^2 - ((2*I)*c*(I*ArcCsch[c + d*x]*(Log[1 - E^(-ArcCsch[c + d*x])]) - Log[1 + E^(-ArcCsch[c + d*x])]) + I*(PolyLog[2, -E^(-ArcCsch[c + d*x])] - PolyLog[2, E^(-ArcCsch[c + d*x])])))/d^2 + (c*ArcCsch[c + d*x]^2*Tanh[ArcCsch[c + d*x]/2])/(2*d^2))/((c + d*x)*(-1 + c/(c + d*x))) - (b^2*f^2*(2*(-2 + 12*c*ArcCsch[c + d*x] + ArcCsch[c + d*x]^2 - 6*c^2*ArcCsch[c + d*x]^2)*Coth[ArcCsch[c + d*x]/2] + 2*ArcCsch[c + d*x]*(-1 + 3*c*ArcCsch[c + d*x])*Csch[ArcCsch[c + d*x]/2]^2 - (ArcCsch[c + d*x]^2*Csch[ArcCsch[c + d*x]/2]^4)/(2*(c + d*x)) - 48*c*(Log[1/((c + d*x)*Sqrt[1 + (c + d*x)^(-2)])]) + Log[Sqrt[1 + (c + d*x)^(-2)])]) + 8*(-1 + 6*c^2)*(Arc

```

```
Csch[c + d*x]*(Log[1 - E^(-ArcCsch[c + d*x])] - Log[1 + E^(-ArcCsch[c + d*x]
)]) + PolyLog[2, -E^(-ArcCsch[c + d*x])] - PolyLog[2, E^(-ArcCsch[c + d*x]
)]) - 2*ArcCsch[c + d*x]*(1 + 3*c*ArcCsch[c + d*x])*Sech[ArcCsch[c + d*x]/2
]^2 - 8*(c + d*x)^3*ArcCsch[c + d*x]^2*Sinh[ArcCsch[c + d*x]/2]^4 + 2*(2 +
12*c*ArcCsch[c + d*x] - ArcCsch[c + d*x]^2 + 6*c^2*ArcCsch[c + d*x]^2)*Tanh
[ArcCsch[c + d*x]/2]))/(24*d^3)
```

Maple [F]

$$\int (fx + e)^2 (a + b \operatorname{arccsch}(dx + c))^2 dx$$

```
[In] int((f*x+e)^2*(a+b*arccsch(d*x+c))^2,x)
```

```
[Out] int((f*x+e)^2*(a+b*arccsch(d*x+c))^2,x)
```

Fricas [F]

$$\int (e + fx)^2 (a + b \operatorname{bsch}^{-1}(c + dx))^2 dx = \int (fx + e)^2 (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

```
[In] integrate((f*x+e)^2*(a+b*arccsch(d*x+c))^2,x, algorithm="fricas")
```

```
[Out] integral(a^2*f^2*x^2 + 2*a^2*e*f*x + a^2*e^2 + (b^2*f^2*x^2 + 2*b^2*e*f*x +
b^2*e^2)*arccsch(d*x + c)^2 + 2*(a*b*f^2*x^2 + 2*a*b*e*f*x + a*b*e^2)*arcc
sch(d*x + c), x)
```

Sympy [F]

$$\int (e + fx)^2 (a + b \operatorname{bsch}^{-1}(c + dx))^2 dx = \int (a + b \operatorname{acsch}(c + dx))^2 (e + fx)^2 dx$$

```
[In] integrate((f*x+e)**2*(a+b*acsch(d*x+c))**2,x)
```

```
[Out] Integral((a + b*acsch(c + d*x))**2*(e + f*x)**2, x)
```

Maxima [F]

$$\int (e + fx)^2 (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (fx + e)^2 (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

[In] integrate((f*x+e)^2*(a+b*arccsch(d*x+c))^2,x, algorithm="maxima")

[Out] $\frac{1}{3}a^2f^2x^3 + a^2efx^2 + a^2e^2x + (2(dx + c)\operatorname{arccsch}(dx + c) + \log(\sqrt{1/(dx + c)^2 + 1} + 1) - \log(\sqrt{1/(dx + c)^2 + 1} - 1))ab^2e^2/d + \frac{1}{3}(b^2f^2x^3 + 3b^2efx^2 + 3b^2e^2x)\log(\sqrt{d^2x^2 + 2cdx + c^2 + 1} + 1)^2 - \int (-1/3(3(b^2d^2f^2x^4 + b^2c^2e^2 + b^2e^2 + 2(b^2d^2ef + b^2cd^2f^2))x^3 + (4b^2cd^2ef + b^2c^2f^2 + (d^2e^2 + f^2)b^2)x^2 + 2(b^2cd^2e^2 + b^2c^2ef + b^2ef)x)\log(dx + c)^2 - 6(ab^2d^2f^2x^4 + 2(ab^2d^2ef + ab^2cd^2f^2))x^3 + (4ab^2cd^2ef + ab^2c^2f^2 + ab^2f^2)x^2 + 2(ab^2c^2ef + ab^2ef)x)\log(dx + c) + 2(3ab^2d^2f^2x^4 + 6(ab^2d^2ef + ab^2cd^2f^2))x^3 + 3(4ab^2cd^2ef + ab^2c^2f^2 + ab^2f^2)x^2 + 6(ab^2c^2ef + ab^2ef)x - 3(b^2d^2f^2x^4 + b^2c^2e^2 + b^2e^2 + 2(b^2d^2ef + b^2cd^2f^2))x^3 + (4b^2cd^2ef + b^2c^2f^2 + (d^2e^2 + f^2)b^2)x^2 + 2(b^2cd^2e^2 + b^2c^2ef + b^2ef)x)\log(dx + c) + ((3ab^2d^2f^2 - b^2d^2f^2)x^4 + (6ab^2d^2ef - 3b^2d^2ef + (6ab^2d^2f^2 - b^2d^2f^2)c)x^3 - 3(b^2d^2e^2 - ab^2c^2f^2 - ab^2f^2 - (4ab^2d^2ef - b^2d^2ef)c)x^2 - 3(b^2cd^2e^2 - 2ab^2c^2ef - 2ab^2ef)x - 3(b^2d^2f^2x^4 + b^2c^2e^2 + b^2e^2 + 2(b^2d^2ef + b^2cd^2f^2))x^3 + (4b^2cd^2ef + b^2c^2f^2 + (d^2e^2 + f^2)b^2)x^2 + 2(b^2cd^2e^2 + b^2c^2ef + b^2ef)x)\log(dx + c))\sqrt{d^2x^2 + 2cdx + c^2 + 1})\log(\sqrt{d^2x^2 + 2cdx + c^2 + 1} + 1) + 3\sqrt{d^2x^2 + 2cdx + c^2 + 1}((b^2d^2f^2x^4 + b^2c^2e^2 + b^2e^2 + 2(b^2d^2ef + b^2cd^2f^2))x^3 + (4b^2cd^2ef + b^2c^2f^2 + (d^2e^2 + f^2)b^2)x^2 + 2(b^2cd^2e^2 + b^2c^2ef + b^2ef)x)\log(dx + c)^2 - 2(ab^2d^2f^2x^4 + 2(ab^2d^2ef + ab^2cd^2f^2))x^3 + (4ab^2cd^2ef + ab^2c^2f^2 + ab^2f^2)x^2 + 2(ab^2c^2ef + ab^2ef)x)\log(dx + c)))/(d^2x^2 + 2cdx + c^2 + (d^2x^2 + 2cdx + c^2 + 1)^{3/2} + 1), x)$

Giac [F]

$$\int (e + fx)^2 (a + b \operatorname{arcsch}^{-1}(c + dx))^2 dx = \int (fx + e)^2 (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

[In] integrate((f*x+e)^2*(a+b*arccsch(d*x+c))^2,x, algorithm="giac")

[Out] integrate((f*x + e)^2*(b*arccsch(d*x + c) + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int (e + fx)^2 (a + \operatorname{bcsch}^{-1}(c + dx))^2 dx = \int (e + fx)^2 \left(a + b \operatorname{asinh}\left(\frac{1}{c + dx}\right) \right)^2 dx$$

```
[In] int((e + f*x)^2*(a + b*asinh(1/(c + d*x)))^2,x)
```

```
[Out] int((e + f*x)^2*(a + b*asinh(1/(c + d*x)))^2, x)
```


3.9 $\int (e + fx) (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$

Optimal result	105
Rubi [A] (verified)	106
Mathematica [B] (verified)	109
Maple [F]	110
Fricas [F]	110
Sympy [F]	110
Maxima [F]	110
Giac [F]	111
Mupad [F(-1)]	111

Optimal result

Integrand size = 18, antiderivative size = 194

$$\begin{aligned}
 & \int (e + fx) (a + b \operatorname{csch}^{-1}(c + dx))^2 dx \\
 &= \frac{bf(c + dx) \sqrt{1 + \frac{1}{(c + dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{d^2} \\
 & - \frac{(de - cf)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2d^2 f} + \frac{(e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2f} \\
 & + \frac{4b(de - cf) (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c + dx)}\right)}{d^2} + \frac{b^2 f \log(c + dx)}{d^2} \\
 & + \frac{2b^2(de - cf) \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c + dx)}\right)}{d^2} - \frac{2b^2(de - cf) \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c + dx)}\right)}{d^2}
 \end{aligned}$$

```

[Out] -1/2*(-c*f+d*e)^2*(a+b*arccsch(d*x+c))^2/d^2/f+1/2*(f*x+e)^2*(a+b*arccsch(d
*x+c))^2/f+4*b*(-c*f+d*e)*(a+b*arccsch(d*x+c))*arctanh(1/(d*x+c)+(1+1/(d*x+
c)^2)^(1/2))/d^2+b^2*f*ln(d*x+c)/d^2+2*b^2*(-c*f+d*e)*polylog(2,-1/(d*x+c)-
(1+1/(d*x+c)^2)^(1/2))/d^2-2*b^2*(-c*f+d*e)*polylog(2,1/(d*x+c)+(1+1/(d*x+c
)^2)^(1/2))/d^2+b*f*(d*x+c)*(a+b*arccsch(d*x+c))*(1+1/(d*x+c)^2)^(1/2)/d^2

```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {6457, 5577, 4275, 4267, 2317, 2438, 4269, 3556}

$$\int (e + fx) (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$$

$$= \frac{4b(de - cf) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right) (a + b \operatorname{csch}^{-1}(c + dx))}{d^2}$$

$$- \frac{(de - cf)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2d^2 f} + \frac{bf(c + dx) \sqrt{\frac{1}{(c+dx)^2} + 1} (a + b \operatorname{csch}^{-1}(c + dx))}{d^2}$$

$$+ \frac{(e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2f} + \frac{2b^2(de - cf) \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^2}$$

$$- \frac{2b^2(de - cf) \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^2} + \frac{b^2 f \log(c + dx)}{d^2}$$

[In] Int[(e + f*x)*(a + b*ArcCsch[c + d*x])^2,x]

[Out] (b*f*(c + d*x)*Sqrt[1 + (c + d*x)^(-2)]*(a + b*ArcCsch[c + d*x]))/d^2 - ((d * e - c*f)^2*(a + b*ArcCsch[c + d*x])^2)/(2*d^2*f) + ((e + f*x)^2*(a + b*ArcCsch[c + d*x])^2)/(2*f) + (4*b*(d*e - c*f)*(a + b*ArcCsch[c + d*x])*ArcTanh[E^ArcCsch[c + d*x]])/d^2 + (b^2*f*Log[c + d*x])/d^2 + (2*b^2*(d*e - c*f)*PolyLog[2, -E^ArcCsch[c + d*x]])/d^2 - (2*b^2*(d*e - c*f)*PolyLog[2, E^ArcCsch[c + d*x]])/d^2

Rule 2317

Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 4267

```
Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x
_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x]
+ (-Dist[d*(m/(f*fz*I)), Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)
], x], x] + Dist[d*(m/(f*fz*I)), Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e +
f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

Rule 4269

```
Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp
[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*
Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 4275

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)
, x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 5577

```
Int[Coth[(c_.) + (d_.)*(x_)]*Csch[(c_.) + (d_.)*(x_)]*(Csch[(c_.) + (d_.)*(
x_)]*(b_.) + (a_.))^(n_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(-(e
+ f*x)^m)*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b
*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x] /
; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 6457

```
Int[((a_.) + ArcCsch[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csch[x]*Co
th[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a,
b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int (a + bx)^2 \coth(x) \operatorname{csch}(x) (de - cf + f \operatorname{csch}(x)) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^2} \\ &= \frac{(e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2f} \\ &\quad - \frac{b \text{Subst}\left(\int (a + bx) (de - cf + f \operatorname{csch}(x))^2 dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^2 f} \end{aligned}$$

$$\begin{aligned}
&= \frac{(e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2f} \\
&\quad - \frac{b \operatorname{Subst}\left(\int \left(d^2 e^2 \left(1 + \frac{cf(-2de+cf)}{d^2 e^2}\right) (a + bx) + 2def \left(1 - \frac{cf}{de}\right) (a + bx) \operatorname{csch}(x) + f^2 (a + bx) \operatorname{csch}^2(x)\right) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^2 f} \\
&= -\frac{(de - cf)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2d^2 f} + \frac{(e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2f} \\
&\quad - \frac{(bf) \operatorname{Subst}\left(\int (a + bx) \operatorname{csch}^2(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^2} \\
&\quad - \frac{(2b(de - cf)) \operatorname{Subst}\left(\int (a + bx) \operatorname{csch}(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^2} \\
&= \frac{bf(c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{d^2} \\
&\quad - \frac{(de - cf)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2d^2 f} + \frac{(e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2f} \\
&\quad + \frac{4b(de - cf) (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^2} \\
&\quad - \frac{(b^2 f) \operatorname{Subst}\left(\int \coth(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^2} \\
&\quad + \frac{(2b^2(de - cf)) \operatorname{Subst}\left(\int \log(1 - e^x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^2} \\
&\quad - \frac{(2b^2(de - cf)) \operatorname{Subst}\left(\int \log(1 + e^x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{d^2} \\
&= \frac{bf(c + dx) \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{d^2} \\
&\quad - \frac{(de - cf)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2d^2 f} + \frac{(e + fx)^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2f} \\
&\quad + \frac{4b(de - cf) (a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^2} \\
&\quad + \frac{b^2 f \log(c + dx)}{d^2} + \frac{(2b^2(de - cf)) \operatorname{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^2} \\
&\quad - \frac{(2b^2(de - cf)) \operatorname{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{bf(c+dx)\sqrt{1+\frac{1}{(c+dx)^2}}(a+b\operatorname{csch}^{-1}(c+dx))}{d^2} \\
&\quad - \frac{(de-cf)^2(a+b\operatorname{csch}^{-1}(c+dx))^2}{2d^2f} + \frac{(e+fx)^2(a+b\operatorname{csch}^{-1}(c+dx))^2}{2f} \\
&\quad + \frac{4b(de-cf)(a+b\operatorname{csch}^{-1}(c+dx))\operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^2} \\
&\quad + \frac{b^2f\log(c+dx)}{d^2} + \frac{2b^2(de-cf)\operatorname{PolyLog}\left(2,-e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^2} \\
&\quad - \frac{2b^2(de-cf)\operatorname{PolyLog}\left(2,e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d^2}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 427 vs. $2(194) = 388$.

Time = 2.96 (sec) , antiderivative size = 427, normalized size of antiderivative = 2.20

$$\int (e+fx)(a+b\operatorname{csch}^{-1}(c+dx))^2 dx$$

$$\begin{aligned}
&= \frac{2a^2(de-cf)(c+dx) + a^2f(c+dx)^2 + 2abf(c+dx)\left(\sqrt{1+\frac{1}{(c+dx)^2}} + (c+dx)\operatorname{csch}^{-1}(c+dx)\right) + 2b^2f\left(\right)}{d^2}
\end{aligned}$$

[In] Integrate[(e + f*x)*(a + b*ArcCsch[c + d*x])^2,x]

[Out] $(2a^2(de-cf)(c+dx) + a^2f(c+dx)^2 + 2abf(c+dx)\left(\sqrt{1+\frac{1}{(c+dx)^2}} + (c+dx)\operatorname{csch}^{-1}(c+dx)\right) + 2b^2f\left(\right)) / (2d^2)$

Maple [F]

$$\int (fx + e) (a + b \operatorname{arccsch}(dx + c))^2 dx$$

[In] `int((f*x+e)*(a+b*arccsch(d*x+c))^2,x)`

[Out] `int((f*x+e)*(a+b*arccsch(d*x+c))^2,x)`

Fricas [F]

$$\int (e + fx) (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \int (fx + e)(b \operatorname{arcsch}(dx + c) + a)^2 dx$$

[In] `integrate((f*x+e)*(a+b*arccsch(d*x+c))^2,x, algorithm="fricas")`

[Out] `integral(a^2*f*x + a^2*e + (b^2*f*x + b^2*e)*arccsch(d*x + c)^2 + 2*(a*b*f*x + a*b*e)*arccsch(d*x + c), x)`

Sympy [F]

$$\int (e + fx) (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \int (a + b \operatorname{acsch}(c + dx))^2 (e + fx) dx$$

[In] `integrate((f*x+e)*(a+b*acsch(d*x+c))**2,x)`

[Out] `Integral((a + b*acsch(c + d*x))**2*(e + f*x), x)`

Maxima [F]

$$\int (e + fx) (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \int (fx + e)(b \operatorname{arcsch}(dx + c) + a)^2 dx$$

[In] `integrate((f*x+e)*(a+b*arccsch(d*x+c))^2,x, algorithm="maxima")`

[Out] `1/2*a^2*f*x^2 + a^2*e*x + (2*(d*x + c)*arccsch(d*x + c) + log(sqrt(1/(d*x + c)^2 + 1) + 1) - log(sqrt(1/(d*x + c)^2 + 1) - 1))*a*b*e/d + 1/2*(b^2*f*x^2 + 2*b^2*e*x)*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1)^2 - integrate(-((b^2*d^2*f*x^3 + b^2*c^2*e + b^2*e + (b^2*d^2*e + 2*b^2*c*d*f)*x^2 + (2*b^2*c*d*e + b^2*c^2*f + b^2*f)*x)*log(d*x + c)^2 - 2*(a*b*d^2*f*x^3 + 2*a*b*c*d*f*x^2 + (a*b*c^2*f + a*b*f)*x)*log(d*x + c) + (2*a*b*d^2*f*x^3 + 4*a*b*c*d*f*x^2 + 2*(a*b*c^2*f + a*b*f)*x - 2*(b^2*d^2*f*x^3 + b^2*c^2*e + b^2*e + (b^2*d^2*e + 2*b^2*c*d*f)*x^2 + (2*b^2*c*d*e + b^2*c^2*f + b^2*f)*x)*log(d*x`

+ c) + sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1)*((2*a*b*d^2*f - b^2*d^2*f)*x^3 - (2*b^2*d^2*e - (4*a*b*d*f - b^2*d*f)*c)*x^2 - 2*(b^2*c*d*e - a*b*c^2*f - a*b*f)*x - 2*(b^2*d^2*f*x^3 + b^2*c^2*e + b^2*e + (b^2*d^2*e + 2*b^2*c*d*f)*x^2 + (2*b^2*c*d*e + b^2*c^2*f + b^2*f)*x)*log(d*x + c))*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1) + sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1)*((b^2*d^2*f*x^3 + b^2*c^2*e + b^2*e + (b^2*d^2*e + 2*b^2*c*d*f)*x^2 + (2*b^2*c*d*e + b^2*c^2*f + b^2*f)*x)*log(d*x + c)^2 - 2*(a*b*d^2*f*x^3 + 2*a*b*c*d*f*x^2 + (a*b*c^2*f + a*b*f)*x)*log(d*x + c)))/(d^2*x^2 + 2*c*d*x + c^2 + (d^2*x^2 + 2*c*d*x + c^2 + 1)^(3/2) + 1), x)

Giac [F]

$$\int (e + fx) (a + b \operatorname{arcsch}(c + dx))^2 dx = \int (fx + e) (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

[In] integrate((f*x+e)*(a+b*arcsch(d*x+c))^2,x, algorithm="giac")

[Out] integrate((f*x + e)*(b*arcsch(d*x + c) + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int (e + fx) (a + b \operatorname{arcsch}(c + dx))^2 dx = \int (e + fx) \left(a + b \operatorname{asinh}\left(\frac{1}{c + dx}\right) \right)^2 dx$$

[In] int((e + f*x)*(a + b*asinh(1/(c + d*x)))^2,x)

[Out] int((e + f*x)*(a + b*asinh(1/(c + d*x)))^2, x)

3.10 $\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$

Optimal result	112
Rubi [A] (verified)	112
Mathematica [B] (verified)	115
Maple [F]	115
Fricas [F]	115
Sympy [F]	116
Maxima [F]	116
Giac [F]	116
Mupad [F(-1)]	117

Optimal result

Integrand size = 12, antiderivative size = 85

$$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \frac{(c + dx) (a + b \operatorname{csch}^{-1}(c + dx))^2}{d} + \frac{4b(a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c + dx)}\right)}{d} + \frac{2b^2 \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c + dx)}\right)}{d} - \frac{2b^2 \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c + dx)}\right)}{d}$$

```
[Out] (d*x+c)*(a+b*arccsch(d*x+c))^2/d+4*b*(a+b*arccsch(d*x+c))*arctanh(1/(d*x+c)
+(1+1/(d*x+c)^2)^(1/2))/d+2*b^2*polylog(2,-1/(d*x+c)-(1+1/(d*x+c)^2)^(1/2))
/d-2*b^2*polylog(2,1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))/d
```

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6451,

6415, 5560, 4267, 2317, 2438}

$$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \frac{4b \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(c+dx)}\right) (a + b \operatorname{csch}^{-1}(c + dx))}{d} + \frac{(c + dx) (a + b \operatorname{csch}^{-1}(c + dx))^2}{d} + \frac{2b^2 \operatorname{PolyLog}\left(2, -e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d} - \frac{2b^2 \operatorname{PolyLog}\left(2, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{d}$$

[In] Int[(a + b*ArcCsch[c + d*x])^2,x]

[Out] ((c + d*x)*(a + b*ArcCsch[c + d*x])^2)/d + (4*b*(a + b*ArcCsch[c + d*x])*ArcTanh[E^ArcCsch[c + d*x]])/d + (2*b^2*PolyLog[2, -E^ArcCsch[c + d*x]])/d - (2*b^2*PolyLog[2, E^ArcCsch[c + d*x]])/d

Rule 2317

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 4267

Int[csc[(e_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Dist[d*(m/(f*fz*I)), Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Dist[d*(m/(f*fz*I)), Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

Rule 5560

Int[Coth[(a_) + (b_)*(x_)]^(p_)*Csch[(a_) + (b_)*(x_)]^(n_)*((c_) + (d_)*(x_))^(m_), x_Symbol] :> Simp[(-c + d*x)^m*(Csch[a + b*x]^n/(b*n)), x] + Dist[d*(m/(b*n)), Int[(c + d*x)^(m - 1)*Csch[a + b*x]^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[p, 1] && GtQ[m, 0]

Rule 6415

```
Int[((a_.) + ArcCsch[(c_.)*(x_.)]*(b_.))^n, x_Symbol] := Dist[-c^(-1), Subst[Int[(a + b*x)^n*Csch[x]*Coth[x], x], x, ArcCsch[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[n, 0]
```

Rule 6451

```
Int[((a_.) + ArcCsch[(c_) + (d_.)*(x_.)]*(b_.))^p, x_Symbol] := Dist[1/d, Subst[Int[(a + b*ArcCsch[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[p, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Subst}\left(\int (a + b\text{csch}^{-1}(x))^2 dx, x, c + dx\right)}{d} \\
&= -\frac{\text{Subst}\left(\int (a + bx)^2 \coth(x)\text{csch}(x) dx, x, \text{csch}^{-1}(c + dx)\right)}{d} \\
&= \frac{(c + dx)(a + b\text{csch}^{-1}(c + dx))^2}{d} - \frac{(2b)\text{Subst}\left(\int (a + bx)\text{csch}(x) dx, x, \text{csch}^{-1}(c + dx)\right)}{d} \\
&= \frac{(c + dx)(a + b\text{csch}^{-1}(c + dx))^2}{d} + \frac{4b(a + b\text{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\text{csch}^{-1}(c+dx)}\right)}{d} \\
&\quad + \frac{(2b^2)\text{Subst}\left(\int \log(1 - e^x) dx, x, \text{csch}^{-1}(c + dx)\right)}{d} \\
&\quad - \frac{(2b^2)\text{Subst}\left(\int \log(1 + e^x) dx, x, \text{csch}^{-1}(c + dx)\right)}{d} \\
&= \frac{(c + dx)(a + b\text{csch}^{-1}(c + dx))^2}{d} + \frac{4b(a + b\text{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\text{csch}^{-1}(c+dx)}\right)}{d} \\
&\quad + \frac{(2b^2)\text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{\text{csch}^{-1}(c+dx)}\right)}{d} - \frac{(2b^2)\text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{\text{csch}^{-1}(c+dx)}\right)}{d} \\
&= \frac{(c + dx)(a + b\text{csch}^{-1}(c + dx))^2}{d} + \frac{4b(a + b\text{csch}^{-1}(c + dx)) \operatorname{arctanh}\left(e^{\text{csch}^{-1}(c+dx)}\right)}{d} \\
&\quad + \frac{2b^2 \operatorname{PolyLog}\left(2, -e^{\text{csch}^{-1}(c+dx)}\right)}{d} - \frac{2b^2 \operatorname{PolyLog}\left(2, e^{\text{csch}^{-1}(c+dx)}\right)}{d}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 176 vs. $2(85) = 170$.

Time = 0.21 (sec) , antiderivative size = 176, normalized size of antiderivative = 2.07

$$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx$$

$$= \frac{a^2 c + a^2 dx + 2ab(c + dx) \operatorname{csch}^{-1}(c + dx) + b^2 c \operatorname{csch}^{-1}(c + dx)^2 + b^2 dx \operatorname{csch}^{-1}(c + dx)^2 - 2b^2 \operatorname{csch}^{-1}(c + dx) \operatorname{csch}^{-1}(c + dx)}{d}$$

[In] Integrate[(a + b*ArcCsch[c + d*x])^2,x]

[Out] (a^2*c + a^2*d*x + 2*a*b*(c + d*x)*ArcCsch[c + d*x] + b^2*c*ArcCsch[c + d*x]^2 + b^2*d*x*ArcCsch[c + d*x]^2 - 2*b^2*ArcCsch[c + d*x]*Log[1 - E^(-ArcCsch[c + d*x])] + 2*b^2*ArcCsch[c + d*x]*Log[1 + E^(-ArcCsch[c + d*x])] + 2*a*b*Log[Cosh[ArcCsch[c + d*x]/2]] - 2*a*b*Log[Sinh[ArcCsch[c + d*x]/2]] - 2*b^2*PolyLog[2, -E^(-ArcCsch[c + d*x])] + 2*b^2*PolyLog[2, E^(-ArcCsch[c + d*x])])/d

Maple [F]

$$\int (a + b \operatorname{arcsch}(dx + c))^2 dx$$

[In] int((a+b*arcsch(d*x+c))^2,x)

[Out] int((a+b*arcsch(d*x+c))^2,x)

Fricas [F]

$$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \int (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

[In] integrate((a+b*arcsch(d*x+c))^2,x, algorithm="fricas")

[Out] integral(b^2*arcsch(d*x + c)^2 + 2*a*b*arcsch(d*x + c) + a^2, x)

Sympy [F]

$$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \int (a + b \operatorname{acsch}(c + dx))^2 dx$$

```
[In] integrate((a+b*acsch(d*x+c))**2,x)
```

```
[Out] Integral((a + b*acsch(c + d*x))**2, x)
```

Maxima [F]

$$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \int (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

```
[In] integrate((a+b*arccsch(d*x+c))^2,x, algorithm="maxima")
```

```
[Out] (x*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1)^2 - integrate(-((d^2*x^2 + 2*
c*d*x + c^2 + 1)^(3/2)*log(d*x + c)^2 + (d^2*x^2 + 2*c*d*x + c^2 + 1)*log(d
*x + c)^2 - 2*((d^2*x^2 + 2*c*d*x + c^2 + 1)*log(d*x + c) + sqrt(d^2*x^2 +
2*c*d*x + c^2 + 1)*(d^2*x^2 + c*d*x + (d^2*x^2 + 2*c*d*x + c^2 + 1)*log(d*x
+ c)))*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1))/(d^2*x^2 + 2*c*d*x + c^
2 + (d^2*x^2 + 2*c*d*x + c^2 + 1)^(3/2) + 1), x))*b^2 + a^2*x + (2*(d*x + c
)*arccsch(d*x + c) + log(sqrt(1/(d*x + c)^2 + 1) + 1) - log(sqrt(1/(d*x + c
)^2 + 1) - 1))*a*b/d
```

Giac [F]

$$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \int (b \operatorname{arcsch}(dx + c) + a)^2 dx$$

```
[In] integrate((a+b*arccsch(d*x+c))^2,x, algorithm="giac")
```

```
[Out] integrate((b*arccsch(d*x + c) + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int (a + b \operatorname{csch}^{-1}(c + dx))^2 dx = \int \left(a + b \operatorname{asinh}\left(\frac{1}{c + dx}\right) \right)^2 dx$$

```
[In] int((a + b*asinh(1/(c + d*x)))^2,x)
```

```
[Out] int((a + b*asinh(1/(c + d*x)))^2, x)
```

3.11 $\int \frac{\left(a+b\operatorname{csch}^{-1}(c+dx)\right)^2}{e+fx} dx$

Optimal result	119
Rubi [A] (verified)	120
Mathematica [F]	126
Maple [F]	127
Fricas [F]	127
Sympy [F]	127
Maxima [F]	127
Giac [F]	128
Mupad [F(-1)]	128

Optimal result

Integrand size = 20, antiderivative size = 475

$$\begin{aligned}
 & \int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{e + fx} dx \\
 &= -\frac{(a + b \operatorname{csch}^{-1}(c + dx))^2 \log\left(1 - e^{2 \operatorname{csch}^{-1}(c + dx)}\right)}{f} \\
 &+ \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2 \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f} \\
 &+ \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2 \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f} \\
 &- \frac{b(a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, e^{2 \operatorname{csch}^{-1}(c + dx)}\right)}{f} \\
 &+ \frac{2b(a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f} \\
 &+ \frac{2b(a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f} \\
 &+ \frac{b^2 \operatorname{PolyLog}\left(3, e^{2 \operatorname{csch}^{-1}(c + dx)}\right)}{2f} - \frac{2b^2 \operatorname{PolyLog}\left(3, -\frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f} \\
 &- \frac{2b^2 \operatorname{PolyLog}\left(3, -\frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f}
 \end{aligned}$$

```

[Out] -(a+b*arccsch(d*x+c))^2*ln(1-(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))^2)/f+(a+b*arccsch(d*x+c))^2*ln(1+(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))*(-c*f+d*e)/(f-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/f+(a+b*arccsch(d*x+c))^2*ln(1+(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))*(-c*f+d*e)/(f+(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/f-b*(a+b*arccsch(d*x+c))*polylog(2,(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))^2)/f+2*b*(a+b*arccsch(d*x+c))*polylog(2,-(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))*(-c*f+d*e)/(f-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/f+2*b*(a+b*arccsch(d*x+c))*polylog(2,-(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))*(-c*f+d*e)/(f+(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/f+1/2*b^2*polylog(3,(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))^2)/f-2*b^2*polylog(3,-(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))*(-c*f+d*e)/(f-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/f-2*b^2*polylog(3,-(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))*(-c*f+d*e)/(f+(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/f

```

Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 475, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.450$, Rules used = {6457, 5715, 5688, 3797, 2221, 2611, 2320, 6724, 5680}

$$\begin{aligned}
& \int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{e + fx} dx \\
&= \frac{2b(a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f - \sqrt{d^2 e^2 - 2cdf e + (c^2+1)f^2}}\right)}{f} \\
&+ \frac{2b(a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f + \sqrt{d^2 e^2 - 2cdf e + (c^2+1)f^2}}\right)}{f} \\
&+ \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2 \log\left(\frac{(de-cf)e^{\operatorname{csch}^{-1}(c+dx)}}{f - \sqrt{(c^2+1)f^2 - 2cdf e + d^2 e^2}} + 1\right)}{f} \\
&+ \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2 \log\left(\frac{(de-cf)e^{\operatorname{csch}^{-1}(c+dx)}}{\sqrt{(c^2+1)f^2 - 2cdf e + d^2 e^2} + f} + 1\right)}{f} \\
&- \frac{b \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(c+dx)}\right) (a + b \operatorname{csch}^{-1}(c + dx))}{f} \\
&- \frac{\log\left(1 - e^{2\operatorname{csch}^{-1}(c+dx)}\right) (a + b \operatorname{csch}^{-1}(c + dx))^2}{f} \\
&- \frac{2b^2 \operatorname{PolyLog}\left(3, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f - \sqrt{d^2 e^2 - 2cdf e + (c^2+1)f^2}}\right)}{f} \\
&- \frac{2b^2 \operatorname{PolyLog}\left(3, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f + \sqrt{d^2 e^2 - 2cdf e + (c^2+1)f^2}}\right)}{f} + \frac{b^2 \operatorname{PolyLog}\left(3, e^{2\operatorname{csch}^{-1}(c+dx)}\right)}{2f}
\end{aligned}$$

[In] Int[(a + b*ArcSch[c + d*x])^2/(e + f*x),x]

[Out] -(((a + b*ArcSch[c + d*x])^2*Log[1 - E^(2*ArcSch[c + d*x])])/f) + ((a + b*ArcSch[c + d*x])^2*Log[1 + (E^ArcSch[c + d*x]*(d*e - c*f))/(f - Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]])/f) + ((a + b*ArcSch[c + d*x])^2*Log[1 + (E^ArcSch[c + d*x]*(d*e - c*f))/(f + Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]])/f) - (b*(a + b*ArcSch[c + d*x])*PolyLog[2, E^(2*ArcSch[c + d*x])])/f + (2*b*(a + b*ArcSch[c + d*x])*PolyLog[2, -(E^ArcSch[c + d*x]*(d*e - c*f))/(f - Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]])/f) + (2*b*(a + b*ArcSch[c + d*x])*PolyLog[2, -(E^ArcSch[c + d*x]*(d*e - c*f))/(f + Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]])/f) + (b^2*PolyLog[3, E^(2*ArcSch[c

$$\frac{+ d*x]]]/(2*f) - (2*b^2*PolyLog[3, -((E^{\text{ArcCsch}[c + d*x]}*(d*e - c*f))/(f - \text{Sqrt}[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]))]/f - (2*b^2*PolyLog[3, -((E^{\text{ArcCsch}[c + d*x]}*(d*e - c*f))/(f + \text{Sqrt}[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]))]/f)))/f$$

Rule 2221

$$\text{Int}[(((F_)^((g_)*(e_)+(f_)*(x_)))^{(n_)*((c_)+(d_)*(x_))^{(m_)}))/((a_)+(b_)*((F_)^((g_)*(e_)+(f_)*(x_)))^{(n_)}), x_Symbol] \rightarrow \text{Simp} [((c + d*x)^m/(b*f*g*n*\text{Log}[F]))*\text{Log}[1 + b*((F^(g*(e + f*x)))^n/a)], x] - \text{Dist}[d*(m/(b*f*g*n*\text{Log}[F])), \text{Int}[(c + d*x)^{(m-1)}*\text{Log}[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&\& \text{IGtQ}[m, 0]$$

Rule 2320

$$\text{Int}[u, x_Symbol] \rightarrow \text{With}[\{v = \text{FunctionOfExponential}[u, x]\}, \text{Dist}[v/D[v, x], \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x] /; \text{FunctionOfExponentialQ}[u, x] \&\& !\text{MatchQ}[u, (w_)*((a_)*(v_)^{(n_))^{(m_)} /; \text{FreeQ}[\{a, m, n\}, x] \&\& \text{IntegerQ}[m*n]] \&\& !\text{MatchQ}[u, E^{((c_)*(a_)+(b_)*x)}*(F_)[v_] /; \text{FreeQ}[\{a, b, c\}, x] \&\& \text{InverseFunctionQ}[F[x]]]$$

Rule 2611

$$\text{Int}[\text{Log}[1 + (e_)*((F_)^((c_)*((a_)+(b_)*(x_)))^{(n_)})*((f_)+(g_)*(x_))^{(m_)}], x_Symbol] \rightarrow \text{Simp}[(-(f + g*x)^m)*(\text{PolyLog}[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*\text{Log}[F])), x] + \text{Dist}[g*(m/(b*c*n*\text{Log}[F])), \text{Int}[(f + g*x)^{(m-1)}*\text{PolyLog}[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; \text{FreeQ}[\{F, a, b, c, e, f, g, n\}, x] \&\& \text{GtQ}[m, 0]$$

Rule 3797

$$\text{Int}[((c_)+(d_)*(x_))^{(m_)}*\text{tan}[(e_)+\text{Pi}*(k_)+(\text{Complex}[0, fz_])*(f_)*(x_)], x_Symbol] \rightarrow \text{Simp}[(-I)*((c + d*x)^{(m+1)}/(d*(m+1))), x] + \text{Dist}[2*I, \text{Int}[((c + d*x)^m*(E^{(2*((-I)*e + f*fz*x))}/(1 + E^{(2*((-I)*e + f*fz*x)})/E^{(2*I*k*Pi)})))/E^{(2*I*k*Pi)}, x], x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \&\& \text{IntegerQ}[4*k] \&\& \text{IGtQ}[m, 0]$$

Rule 5680

$$\text{Int}[(\text{Cosh}[(c_)+(d_)*(x_)]*((e_)+(f_)*(x_))^{(m_)}))/((a_)+(b_)*\text{Sin}h[(c_)+(d_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[-(e + f*x)^{(m+1)}/(b*f*(m+1)), x] + (\text{Int}[(e + f*x)^m*(E^{(c + d*x)})/(a - \text{Rt}[a^2 + b^2, 2] + b*E^{(c + d*x)}), x] + \text{Int}[(e + f*x)^m*(E^{(c + d*x)})/(a + \text{Rt}[a^2 + b^2, 2] + b*E^{(c + d*x)}), x]) /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{IGtQ}[m, 0] \&\& \text{NeQ}[a^2 + b^2, 0]$$

Rule 5688

```
Int[(Coth[(c_.) + (d_.)*(x_)]^(n_.)*((e_.) + (f_.)*(x_))^(m_.))/((a_) + (b_.)*Sinh[(c_.) + (d_.)*(x_)]), x_Symbol] :> Dist[1/a, Int[(e + f*x)^m*Coth[c + d*x]^n, x], x] - Dist[b/a, Int[(e + f*x)^m*Cosh[c + d*x]*(Coth[c + d*x]^(n - 1)/(a + b*Sinh[c + d*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 5715

```
Int[(((e_.) + (f_.)*(x_))^(m_.)*(F_)[(c_.) + (d_.)*(x_)]^(n_.)*(G_)[(c_.) + (d_.)*(x_)]^(p_.))/(Csch[(c_.) + (d_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :> Int[(e + f*x)^m*Sinh[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Sinh[c + d*x])), x] /; FreeQ[{a, b, c, d, e, f}, x] && HyperbolicQ[F] && HyperbolicQ[G] && IntegersQ[m, n, p]
```

Rule 6457

```
Int[((a_.) + ArcCsch[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csch[x]*Coth[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{(a + bx)^2 \coth(x) \operatorname{csch}(x)}{de - cf + f \operatorname{csch}(x)} dx, x, \operatorname{csch}^{-1}(c + dx)\right) \\
&= -\text{Subst}\left(\int \frac{(a + bx)^2 \coth(x)}{f + (de - cf) \sinh(x)} dx, x, \operatorname{csch}^{-1}(c + dx)\right) \\
&= -\frac{\text{Subst}\left(\int (a + bx)^2 \coth(x) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{f} \\
&\quad + \frac{(de - cf) \text{Subst}\left(\int \frac{(a + bx)^2 \cosh(x)}{f + (de - cf) \sinh(x)} dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{f}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2\text{Subst}\left(\int \frac{e^{2x}(a+bx)^2}{1-e^{2x}} dx, x, \text{csch}^{-1}(c+dx)\right)}{f} \\
&+ \frac{(de-cf)\text{Subst}\left(\int \frac{e^x(a+bx)^2}{f+e^x(de-cf)-\sqrt{d^2e^2-2cdef+f^2+c^2f^2}} dx, x, \text{csch}^{-1}(c+dx)\right)}{f} \\
&+ \frac{(de-cf)\text{Subst}\left(\int \frac{e^x(a+bx)^2}{f+e^x(de-cf)+\sqrt{d^2e^2-2cdef+f^2+c^2f^2}} dx, x, \text{csch}^{-1}(c+dx)\right)}{f} \\
&= -\frac{(a+b\text{csch}^{-1}(c+dx))^2 \log\left(1-e^{2\text{csch}^{-1}(c+dx)}\right)}{f} \\
&+ \frac{(a+b\text{csch}^{-1}(c+dx))^2 \log\left(1+\frac{e^{\text{csch}^{-1}(c+dx)}(de-cf)}{f-\sqrt{d^2e^2-2cdef+(1+c^2)f^2}}\right)}{f} \\
&+ \frac{(a+b\text{csch}^{-1}(c+dx))^2 \log\left(1+\frac{e^{\text{csch}^{-1}(c+dx)}(de-cf)}{f+\sqrt{d^2e^2-2cdef+(1+c^2)f^2}}\right)}{f} \\
&+ \frac{(2b)\text{Subst}\left(\int (a+bx) \log(1-e^{2x}) dx, x, \text{csch}^{-1}(c+dx)\right)}{f} \\
&- \frac{(2b)\text{Subst}\left(\int (a+bx) \log\left(1+\frac{e^x(de-cf)}{f-\sqrt{d^2e^2-2cdef+f^2+c^2f^2}}\right) dx, x, \text{csch}^{-1}(c+dx)\right)}{f} \\
&- \frac{(2b)\text{Subst}\left(\int (a+bx) \log\left(1+\frac{e^x(de-cf)}{f+\sqrt{d^2e^2-2cdef+f^2+c^2f^2}}\right) dx, x, \text{csch}^{-1}(c+dx)\right)}{f}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{(a + b\operatorname{csch}^{-1}(c + dx))^2 \log\left(1 - e^{2\operatorname{csch}^{-1}(c+dx)}\right)}{f} \\
&+ \frac{(a + b\operatorname{csch}^{-1}(c + dx))^2 \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f - \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{f} \\
&+ \frac{(a + b\operatorname{csch}^{-1}(c + dx))^2 \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f + \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{f} \\
&- \frac{b(a + b\operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(c+dx)}\right)}{f} \\
&+ \frac{2b(a + b\operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f - \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{f} \\
&+ \frac{2b(a + b\operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f + \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{f} \\
&+ \frac{b^2 \operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, e^{2x}\right) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{f} \\
&- \frac{(2b^2) \operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, -\frac{e^x(de-cf)}{f - \sqrt{d^2e^2 - 2cdef + f^2 + c^2f^2}}\right) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{f} \\
&- \frac{(2b^2) \operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, -\frac{e^x(de-cf)}{f + \sqrt{d^2e^2 - 2cdef + f^2 + c^2f^2}}\right) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{f}
\end{aligned}$$

$$\begin{aligned}
&= - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2 \log\left(1 - e^{2 \operatorname{csch}^{-1}(c + dx)}\right)}{f} \\
&+ \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2 \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f} \\
&+ \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2 \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f} \\
&- \frac{b(a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, e^{2 \operatorname{csch}^{-1}(c + dx)}\right)}{f} \\
&+ \frac{2b(a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f} \\
&+ \frac{2b(a + b \operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{f} \\
&+ \frac{b^2 \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(2, x)}{x} dx, x, e^{2 \operatorname{csch}^{-1}(c + dx)}\right)}{2f} \\
&- \frac{(2b^2) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(2, \frac{(de - cf)x}{-f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{x} dx, x, e^{\operatorname{csch}^{-1}(c + dx)}\right)}{f} \\
&- \frac{(2b^2) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(2, -\frac{(de - cf)x}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2)f^2}}\right)}{x} dx, x, e^{\operatorname{csch}^{-1}(c + dx)}\right)}{f}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{(a + b\operatorname{csch}^{-1}(c + dx))^2 \log\left(1 - e^{2\operatorname{csch}^{-1}(c+dx)}\right)}{f} \\
&+ \frac{(a + b\operatorname{csch}^{-1}(c + dx))^2 \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f - \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{f} \\
&+ \frac{(a + b\operatorname{csch}^{-1}(c + dx))^2 \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f + \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{f} \\
&- \frac{b(a + b\operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(c+dx)}\right)}{f} \\
&+ \frac{2b(a + b\operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f - \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{f} \\
&+ \frac{2b(a + b\operatorname{csch}^{-1}(c + dx)) \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f + \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{f} \\
&+ \frac{b^2 \operatorname{PolyLog}\left(3, e^{2\operatorname{csch}^{-1}(c+dx)}\right)}{2f} - \frac{2b^2 \operatorname{PolyLog}\left(3, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f - \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{f} \\
&- \frac{2b^2 \operatorname{PolyLog}\left(3, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f + \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{f}
\end{aligned}$$

Mathematica [F]

$$\int \frac{(a + b\operatorname{csch}^{-1}(c + dx))^2}{e + fx} dx = \int \frac{(a + b\operatorname{csch}^{-1}(c + dx))^2}{e + fx} dx$$

[In] Integrate[(a + b*ArcCsch[c + d*x])^2/(e + f*x), x]

[Out] Integrate[(a + b*ArcCsch[c + d*x])^2/(e + f*x), x]

Maple [F]

$$\int \frac{(a + b \operatorname{arccsch}(dx + c))^2}{fx + e} dx$$

[In] int((a+b*arccsch(d*x+c))^2/(f*x+e),x)

[Out] int((a+b*arccsch(d*x+c))^2/(f*x+e),x)

Fricas [F]

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{e + fx} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{fx + e} dx$$

[In] integrate((a+b*arccsch(d*x+c))^2/(f*x+e),x, algorithm="fricas")

[Out] integral((b^2*arccsch(d*x + c)^2 + 2*a*b*arccsch(d*x + c) + a^2)/(f*x + e), x)

Sympy [F]

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{e + fx} dx = \int \frac{(a + b \operatorname{acsch}(c + dx))^2}{e + fx} dx$$

[In] integrate((a+b*acsch(d*x+c))**2/(f*x+e),x)

[Out] Integral((a + b*acsch(c + d*x))**2/(e + f*x), x)

Maxima [F]

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{e + fx} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{fx + e} dx$$

[In] integrate((a+b*arccsch(d*x+c))^2/(f*x+e),x, algorithm="maxima")

[Out] a^2*log(f*x + e)/f + integrate(b^2*log(sqrt(1/(d*x + c)^2 + 1) + 1/(d*x + c))^2/(f*x + e) + 2*a*b*log(sqrt(1/(d*x + c)^2 + 1) + 1/(d*x + c))/(f*x + e), x)

Giac [F]

$$\int \frac{(a + b \operatorname{arcsch}^{-1}(c + dx))^2}{e + fx} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{fx + e} dx$$

[In] integrate((a+b*arccsch(d*x+c))^2/(f*x+e),x, algorithm="giac")

[Out] integrate((b*arccsch(d*x + c) + a)^2/(f*x + e), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arcsch}^{-1}(c + dx))^2}{e + fx} dx = \int \frac{(a + b \operatorname{asinh}(\frac{1}{c+dx}))^2}{e + fx} dx$$

[In] int((a + b*asinh(1/(c + d*x)))^2/(e + f*x),x)

[Out] int((a + b*asinh(1/(c + d*x)))^2/(e + f*x), x)

$$3.12 \quad \int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^2} dx$$

Optimal result	129
Rubi [A] (verified)	130
Mathematica [C] (warning: unable to verify)	134
Maple [F]	135
Fricas [F]	135
Sympy [F]	136
Maxima [F]	136
Giac [F]	136
Mupad [F(-1)]	137

Optimal result

Integrand size = 20, antiderivative size = 448

$$\begin{aligned} & \int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^2} dx \\ &= \frac{d(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(de - cf)} - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(e + fx)} \\ & \quad - \frac{2bd(a + b \operatorname{csch}^{-1}(c + dx)) \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f - \sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}} \\ & \quad + \frac{2bd(a + b \operatorname{csch}^{-1}(c + dx)) \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f + \sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}} \\ & \quad - \frac{2b^2d \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f - \sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}} \\ & \quad + \frac{2b^2d \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f + \sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}} \end{aligned}$$

```
[Out] d*(a+b*arccsch(d*x+c))^2/f/(-c*f+d*e)-(a+b*arccsch(d*x+c))^2/f/(f*x+e)-2*b*
d*(a+b*arccsch(d*x+c))*ln(1+(1/(d*x+c)+(1+1/(d*x+c)^2)^(1/2))*(-c*f+d*e)/(f
-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/(-c*f+d*e)/(d^2*e^2-2*c*d*e*f+(c^2
+1)*f^2)^(1/2)+2*b*d*(a+b*arccsch(d*x+c))*ln(1+(1/(d*x+c)+(1+1/(d*x+c)^2)^(
1/2))*(-c*f+d*e)/(f+(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)))/(-c*f+d*e)/(d^2
*e^2-2*c*d*e*f+(c^2+1)*f^2)^(1/2)-2*b^2*d*polylog(2,-(1/(d*x+c)+(1+1/(d*x+c
```

$$\begin{aligned} & \left. \right)^2)^{(1/2)} * (-c*f+d*e) / (f - (d^2*e^2 - 2*c*d*e*f + (c^2+1)*f^2)^{(1/2)}) / (-c*f+d*e) \\ & / (d^2*e^2 - 2*c*d*e*f + (c^2+1)*f^2)^{(1/2)} + 2*b^2*d*polylog(2, -(1/(d*x+c) + (1+1/ \\ & (d*x+c)^2)^{(1/2)}) * (-c*f+d*e) / (f + (d^2*e^2 - 2*c*d*e*f + (c^2+1)*f^2)^{(1/2)})) / (-c \\ & *f+d*e) / (d^2*e^2 - 2*c*d*e*f + (c^2+1)*f^2)^{(1/2)} \end{aligned}$$

Rubi [A] (verified)

Time = 0.77 (sec) , antiderivative size = 448, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {6457, 5577, 4276, 3403, 2296, 2221, 2317, 2438}

$$\begin{aligned} & \int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^2} dx \\ & = -\frac{2bd(a + b \operatorname{csch}^{-1}(c + dx)) \log\left(\frac{(de - cf)e^{\operatorname{csch}^{-1}(c + dx)}}{f - \sqrt{(c^2 + 1)f^2 - 2cdf + d^2e^2}} + 1\right)}{(de - cf)\sqrt{(c^2 + 1)f^2 - 2cdf + d^2e^2}} \\ & + \frac{2bd(a + b \operatorname{csch}^{-1}(c + dx)) \log\left(\frac{(de - cf)e^{\operatorname{csch}^{-1}(c + dx)}}{\sqrt{(c^2 + 1)f^2 - 2cdf + d^2e^2} + f} + 1\right)}{(de - cf)\sqrt{(c^2 + 1)f^2 - 2cdf + d^2e^2}} + \frac{d(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(de - cf)} \\ & - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(e + fx)} - \frac{2b^2d \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f - \sqrt{d^2e^2 - 2cdf + (c^2 + 1)f^2}}\right)}{(de - cf)\sqrt{(c^2 + 1)f^2 - 2cdf + d^2e^2}} \\ & + \frac{2b^2d \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c + dx)}(de - cf)}{f + \sqrt{d^2e^2 - 2cdf + (c^2 + 1)f^2}}\right)}{(de - cf)\sqrt{(c^2 + 1)f^2 - 2cdf + d^2e^2}} \end{aligned}$$

[In] Int[(a + b*ArcCsch[c + d*x])^2/(e + f*x)^2,x]

[Out] (d*(a + b*ArcCsch[c + d*x])^2)/(f*(d*e - c*f)) - (a + b*ArcCsch[c + d*x])^2/(f*(e + f*x)) - (2*b*d*(a + b*ArcCsch[c + d*x])*Log[1 + (E^ArcCsch[c + d*x]*(d*e - c*f))/(f - Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]])/((d*e - c*f)*Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]) + (2*b*d*(a + b*ArcCsch[c + d*x])*Log[1 + (E^ArcCsch[c + d*x]*(d*e - c*f))/(f + Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]])/((d*e - c*f)*Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]) - (2*b^2*d*PolyLog[2, -((E^ArcCsch[c + d*x]*(d*e - c*f))/(f - Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2])])/(d*e - c*f)*Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]) + (2*b^2*d*PolyLog[2, -((E^ArcCsch[c + d*x]*(d*e - c*f))/(f + Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2])])/(d*e - c*f)*Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2])

Rule 2221

Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)*((c_) + (d_)*(x_))^(m_)) / ((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] :> Simp

$$\left[\left((c + dx)^m / (bfg^n \log[F]) \right) \log[1 + b((F^{g(e+fx)})^n/a)], x \right] - \text{Dist}[d(m/(bfg^n \log[F])), \text{Int}[(c + dx)^{(m-1)} \log[1 + b((F^{g(e+fx)})^n/a)], x], x] /;$$

$$\text{FreeQ}\{F, a, b, c, d, e, f, g, n\}, x \} \ \&\& \ \text{IGtQ}[m, 0]$$

Rule 2296

$$\text{Int}[(F^u)((f_.) + (g_.)x)^{(m_.)} / ((a_.) + (b_.)F^u + (c_.)F^v)], x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[b^2 - 4ac, 2]\}, \text{Dist}[2(c/q), \text{Int}[(f + gx)^m (F^u/(b - q + 2cF^u)), x], x] - \text{Dist}[2(c/q), \text{Int}[(f + gx)^m (F^u/(b + q + 2cF^u)), x], x] /;$$

$$\text{FreeQ}\{F, a, b, c, f, g\}, x \} \ \&\& \ \text{EqQ}[v, 2u] \ \&\& \ \text{LinearQ}[u, x] \ \&\& \ \text{NeQ}[b^2 - 4ac, 0] \ \&\& \ \text{IGtQ}[m, 0]$$

Rule 2317

$$\text{Int}[\log[(a_.) + (b_.)((F^{(e_.)((c_.) + (d_.)x))})^{(n_.)})], x_Symbol] \rightarrow \text{Dist}[1/(de^n \log[F]), \text{Subst}[\text{Int}[\log[a + bx]/x, x], x, (F^{e(c+dx)})^n], x] /;$$

$$\text{FreeQ}\{F, a, b, c, d, e, n\}, x \} \ \&\& \ \text{GtQ}[a, 0]$$

Rule 2438

$$\text{Int}[\log[(c_.)((d_.) + (e_.)x)^{(n_.)})] / (x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)ex^n/n, x] /;$$

$$\text{FreeQ}\{c, d, e, n\}, x \} \ \&\& \ \text{EqQ}[c*d, 1]$$

Rule 3403

$$\text{Int}[(c_.) + (d_.)x)^{(m_.)} / ((a_.) + (b_.)\sin[e_.] + (\text{Complex}[0, fz])f_.)x)], x_Symbol] \rightarrow \text{Dist}[2, \text{Int}[(c + dx)^m (E^{(-I)e + f*fx}) / ((-I)b + 2aE^{(-I)e + f*fx} + I*bE^{2*((-I)e + f*fx)})], x], x] /;$$

$$\text{FreeQ}\{a, b, c, d, e, f, fz\}, x \} \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{IGtQ}[m, 0]$$

Rule 4276

$$\text{Int}[(\csc[e_.] + (f_.)x)(b_.) + (a_.)^{(n_.)}((c_.) + (d_.)x)^{(m_.)}], x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c + dx)^m, 1/(\sin[e + fx]^n/(b + a\sin[e + fx])^n)], x], x] /;$$

$$\text{FreeQ}\{a, b, c, d, e, f\}, x \} \ \&\& \ \text{ILtQ}[n, 0] \ \&\& \ \text{IGtQ}[m, 0]$$

Rule 5577

$$\text{Int}[\text{Coth}[(c_.) + (d_.)x] \text{Csch}[(c_.) + (d_.)x] (\text{Csch}[(c_.) + (d_.)x] (b_.) + (a_.)^{(n_.)}((e_.) + (f_.)x)^{(m_.)})], x_Symbol] \rightarrow \text{Simp}[(-e + fx)^m ((a + b\text{Csch}[c + dx])^{(n+1)} / (b*d*(n+1))), x] + \text{Dist}[f*(m/(b*d*(n+1))), \text{Int}[(e + fx)^{(m-1)} (a + b\text{Csch}[c + dx])^{(n+1)}], x], x] /;$$

$$\text{FreeQ}\{a, b, c, d, e, f, n\}, x \} \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{NeQ}[n, -1]$$

Rule 6457

```
Int[((a_.) + ArcCsch[(c_) + (d_.)*(x_)]*(b_.))^p_.)*((e_.) + (f_.)*(x_))^m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csch[x]*Coth[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= - \left(d \text{Subst} \left(\int \frac{(a + bx)^2 \coth(x) \operatorname{csch}(x)}{(de - cf + f \operatorname{csch}(x))^2} dx, x, \operatorname{csch}^{-1}(c + dx) \right) \right) \\
&= - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(e + fx)} + \frac{(2bd) \text{Subst} \left(\int \frac{a+bx}{de - cf + f \operatorname{csch}(x)} dx, x, \operatorname{csch}^{-1}(c + dx) \right)}{f} \\
&= - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(e + fx)} \\
&\quad + \frac{(2bd) \text{Subst} \left(\int \left(\frac{a+bx}{de - cf} + \frac{f(a+bx)}{(-de + cf)(f + de(1 - \frac{cf}{de}) \sinh(x))} \right) dx, x, \operatorname{csch}^{-1}(c + dx) \right)}{f} \\
&= \frac{d(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(de - cf)} - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(e + fx)} \\
&\quad - \frac{(2bd) \text{Subst} \left(\int \frac{a+bx}{f + de(1 - \frac{cf}{de}) \sinh(x)} dx, x, \operatorname{csch}^{-1}(c + dx) \right)}{de - cf} \\
&= \frac{d(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(de - cf)} - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(e + fx)} \\
&\quad - \frac{(4bd) \text{Subst} \left(\int \frac{e^x(a+bx)}{2e^x f - de(1 - \frac{cf}{de}) + dee^{2x}(1 - \frac{cf}{de})} dx, x, \operatorname{csch}^{-1}(c + dx) \right)}{de - cf} \\
&= \frac{d(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(de - cf)} - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(e + fx)} \\
&\quad - \frac{(4bd) \text{Subst} \left(\int \frac{e^x(a+bx)}{2f + 2de e^x(1 - \frac{cf}{de}) - 2\sqrt{d^2 e^2 - 2cdef + f^2 + c^2 f^2}} dx, x, \operatorname{csch}^{-1}(c + dx) \right)}{\sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&\quad + \frac{(4bd) \text{Subst} \left(\int \frac{e^x(a+bx)}{2f + 2de e^x(1 - \frac{cf}{de}) + 2\sqrt{d^2 e^2 - 2cdef + f^2 + c^2 f^2}} dx, x, \operatorname{csch}^{-1}(c + dx) \right)}{\sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{d(a + b\operatorname{csch}^{-1}(c + dx))^2}{f(de - cf)} - \frac{(a + b\operatorname{csch}^{-1}(c + dx))^2}{f(e + fx)} \\
&\quad - \frac{2bd(a + b\operatorname{csch}^{-1}(c + dx)) \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f - \sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}} \\
&\quad + \frac{2bd(a + b\operatorname{csch}^{-1}(c + dx)) \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f + \sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}} \\
&\quad + \frac{(2b^2d) \operatorname{Subst}\left(\int \log\left(1 + \frac{2dee^x(1-\frac{cf}{de})}{2f-2\sqrt{d^2e^2-2cdf+f^2+c^2f^2}}\right) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{(de - cf)\sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}} \\
&\quad + \frac{(2b^2d) \operatorname{Subst}\left(\int \log\left(1 + \frac{2dee^x(1-\frac{cf}{de})}{2f+2\sqrt{d^2e^2-2cdf+f^2+c^2f^2}}\right) dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{(de - cf)\sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}} \\
&= \frac{d(a + b\operatorname{csch}^{-1}(c + dx))^2}{f(de - cf)} - \frac{(a + b\operatorname{csch}^{-1}(c + dx))^2}{f(e + fx)} \\
&\quad - \frac{2bd(a + b\operatorname{csch}^{-1}(c + dx)) \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f - \sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}} \\
&\quad + \frac{2bd(a + b\operatorname{csch}^{-1}(c + dx)) \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de-cf)}{f + \sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}} \\
&\quad + \frac{(2b^2d) \operatorname{Subst}\left(\int \frac{\log\left(1 + \frac{2de(1-\frac{cf}{de})x}{2f-2\sqrt{d^2e^2-2cdf+f^2+c^2f^2}}\right)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}} \\
&\quad + \frac{(2b^2d) \operatorname{Subst}\left(\int \frac{\log\left(1 + \frac{2de(1-\frac{cf}{de})x}{2f+2\sqrt{d^2e^2-2cdf+f^2+c^2f^2}}\right)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdf + (1+c^2)f^2}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{d(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(de - cf)} - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{f(e + fx)} \\
&\quad - \frac{2bd(a + b \operatorname{csch}^{-1}(c + dx)) \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)(de-cf)}}{f - \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}} \\
&\quad + \frac{2bd(a + b \operatorname{csch}^{-1}(c + dx)) \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)(de-cf)}}{f + \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}} \\
&\quad - \frac{2b^2d \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)(de-cf)}}{f - \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}} \\
&\quad + \frac{2b^2d \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)(de-cf)}}{f + \sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}\right)}{(de - cf)\sqrt{d^2e^2 - 2cdef + (1+c^2)f^2}}
\end{aligned}$$

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 12.80 (sec) , antiderivative size = 2061, normalized size of antiderivative = 4.60

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^2} dx = \text{Result too large to show}$$

[In] Integrate[(a + b*ArcCsch[c + d*x])^2/(e + f*x)^2,x]

[Out] $-(a^2/(f*(e + f*x))) - (2*a*b*(c + d*x)^2*(f + (d*e - c*f)/(c + d*x))^2*(\operatorname{ArcCsch}[c + d*x]/(f + (d*e)/(c + d*x) - (c*f)/(c + d*x)) - (2*\operatorname{ArcTan}[(d*e - c*f - f*\operatorname{Tanh}[\operatorname{ArcCsch}[c + d*x]/2])/ \operatorname{Sqrt}[-(d^2*e^2) + 2*c*d*e*f - (1 + c^2)*f^2]])/\operatorname{Sqrt}[-(d^2*e^2) + 2*c*d*e*f - (1 + c^2)*f^2])/(d*(-(d*e) + c*f)*(e + f*x)^2) - (b^2*(c + d*x)^2*(f + (d*e - c*f)/(c + d*x))^2*(\operatorname{ArcCsch}[c + d*x]^2/((- (d*e) + c*f)*(f + (d*e)/(c + d*x) - (c*f)/(c + d*x))) + (2*((-I)*\operatorname{Pi}*\operatorname{ArcTanh}[(-(d*e) + c*f + f*\operatorname{Tanh}[\operatorname{ArcCsch}[c + d*x]/2])/ \operatorname{Sqrt}[f^2 + (d*e - c*f)^2]])/\operatorname{Sqrt}[f^2 + (d*e - c*f)^2] - (2*(\operatorname{Pi}/2 - I*\operatorname{ArcCsch}[c + d*x])*\operatorname{ArcTanh}[(f - I*(d*e - c*f))*\operatorname{Cot}[(\operatorname{Pi}/2 - I*\operatorname{ArcCsch}[c + d*x])/2])/ \operatorname{Sqrt}[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2]) - 2*\operatorname{ArcCos}[((-I)*f)/(d*e - c*f)]*\operatorname{ArcTanh}[(f - I*(d*e - c*f))*\operatorname{Tan}[(\operatorname{Pi}/2 - I*\operatorname{ArcCsch}[c + d*x])/2])/ \operatorname{Sqrt}[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2]) + (\operatorname{ArcCos}[((-I)*f)/(d*e - c*f)] - (2*I)*(\operatorname{ArcTanh}[(f - I*(d*e - c*f))*\operatorname{Cot}[(\operatorname{Pi}/2 - I*\operatorname{ArcCsch}[c + d*x])/2])/ \operatorname{Sqrt}[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2]) - \operatorname{ArcTanh}[(f - I*(d*e - c*f))*\operatorname{Tan}[(\operatorname{Pi}/2 - I*\operatorname{ArcCsch}[c + d*x])/2])/ \operatorname{Sqrt}[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2]))*\operatorname{Log}[\operatorname{Sqrt}[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2]/(\operatorname{Sqrt}[2]*E^((I/2)*(\operatorname{Pi}/2 - I*\operatorname{ArcCsch}[c + d*x]))*\operatorname{Sqrt}[(-I)*(d*e - c*f)]*\operatorname{Sqrt}[f + (d*e - c*f)/(c + d*x)])] + (\operatorname{ArcCos}[$

```

((-I)*f)/(d*e - c*f)] + (2*I)*(ArcTanh[((f - I*(d*e - c*f))*Cot[(Pi/2 - I*ArcCsSch[c + d*x])/2])/Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2]] - ArcTanh[(-f - I*(d*e - c*f))*Tan[(Pi/2 - I*ArcCsSch[c + d*x])/2])/Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2]])*Log[(E^((I/2)*(Pi/2 - I*ArcCsSch[c + d*x]))*Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2])/(Sqrt[2]*Sqrt[(-I)*(d*e - c*f)]*Sqrt[f + (d*e - c*f)/(c + d*x)])] - (ArcCos[(-I)*f)/(d*e - c*f)] + (2*I)*ArcTanh[(-f - I*(d*e - c*f))*Tan[(Pi/2 - I*ArcCsSch[c + d*x])/2])/Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2]]*Log[1 - (I*(f - I*Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2]))*(f - I*(d*e - c*f) - Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2])*Tan[(Pi/2 - I*ArcCsSch[c + d*x])/2])]/((d*e - c*f)*(f - I*(d*e - c*f) + Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2])*Tan[(Pi/2 - I*ArcCsSch[c + d*x])/2])]] + (-ArcCos[(-I)*f)/(d*e - c*f)] + (2*I)*ArcTanh[(-f - I*(d*e - c*f))*Tan[(Pi/2 - I*ArcCsSch[c + d*x])/2])/Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2]]*Log[1 - (I*(f + I*Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2]))*(f - I*(d*e - c*f) - Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2])*Tan[(Pi/2 - I*ArcCsSch[c + d*x])/2])]/((d*e - c*f)*(f - I*(d*e - c*f) + Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2])*Tan[(Pi/2 - I*ArcCsSch[c + d*x])/2])]] + I*(PolyLog[2, (I*(f - I*Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2]))*(f - I*(d*e - c*f) - Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2])*Tan[(Pi/2 - I*ArcCsSch[c + d*x])/2])]/((d*e - c*f)*(f - I*(d*e - c*f) + Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2])*Tan[(Pi/2 - I*ArcCsSch[c + d*x])/2])]] - PolyLog[2, (I*(f + I*Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2]))*(f - I*(d*e - c*f) - Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2])*Tan[(Pi/2 - I*ArcCsSch[c + d*x])/2])]/((d*e - c*f)*(f - I*(d*e - c*f) + Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2])*Tan[(Pi/2 - I*ArcCsSch[c + d*x])/2])]]))/Sqrt[-(d^2*e^2) + 2*c*d*e*f - f^2 - c^2*f^2])]/(d*e - c*f))/((d*(e + f*x)^2)

```

Maple [F]

$$\int \frac{(a + b \operatorname{arccsch}(dx + c))^2}{(fx + e)^2} dx$$

```
[In] int((a+b*arccsch(d*x+c))^2/(f*x+e)^2,x)
```

```
[Out] int((a+b*arccsch(d*x+c))^2/(f*x+e)^2,x)
```

Fricas [F]

$$\int \frac{(a + b \operatorname{bsch}^{-1}(c + dx))^2}{(e + fx)^2} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{(fx + e)^2} dx$$

```
[In] integrate((a+b*arccsch(d*x+c))^2/(f*x+e)^2,x, algorithm="fricas")
```

```
[Out] integral((b^2*arccsch(d*x + c)^2 + 2*a*b*arccsch(d*x + c) + a^2)/(f^2*x^2 + 2*e*f*x + e^2), x)
```

SymPy [F]

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^2} dx = \int \frac{(a + b \operatorname{acsch}(c + dx))^2}{(e + fx)^2} dx$$

```
[In] integrate((a+b*acsch(d*x+c))**2/(f*x+e)**2,x)
```

```
[Out] Integral((a + b*acsch(c + d*x))**2/(e + f*x)**2, x)
```

Maxima [F]

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^2} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{(fx + e)^2} dx$$

```
[In] integrate((a+b*arccsch(d*x+c))^2/(f*x+e)^2,x, algorithm="maxima")
```

```
[Out] -b^2*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1)^2/(f^2*x + e*f) - a^2/(f^2*x + e*f) - integrate(-((b^2*d^2*f*x^2 + 2*b^2*c*d*f*x + (c^2*f + f)*b^2)*log(d*x + c)^2 - 2*(a*b*d^2*f*x^2 + 2*a*b*c*d*f*x + (c^2*f + f)*a*b)*log(d*x + c) + 2*(a*b*d^2*f*x^2 + 2*a*b*c*d*f*x + (c^2*f + f)*a*b - (b^2*d^2*f*x^2 + 2*b^2*c*d*f*x + (c^2*f + f)*b^2)*log(d*x + c) + (b^2*c*d*e + (c^2*f + f)*a*b + (a*b*d^2*f + b^2*d^2*f)*x^2 + (2*a*b*c*d*f + (d^2*e + c*d*f)*b^2)*x - (b^2*d^2*f*x^2 + 2*b^2*c*d*f*x + (c^2*f + f)*b^2)*log(d*x + c))*sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1))*log(sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1) + 1) + sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1)*((b^2*d^2*f*x^2 + 2*b^2*c*d*f*x + (c^2*f + f)*b^2)*log(d*x + c)^2 - 2*(a*b*d^2*f*x^2 + 2*a*b*c*d*f*x + (c^2*f + f)*a*b)*log(d*x + c)))/(d^2*f^3*x^4 + c^2*e^2*f + 2*(d^2*e*f^2 + c*d*f^3)*x^3 + e^2*f + (d^2*e^2*f + 4*c*d*e*f^2 + c^2*f^3 + f^3)*x^2 + 2*(c*d*e^2*f + c^2*e*f^2 + e*f^2)*x + (d^2*f^3*x^4 + c^2*e^2*f + 2*(d^2*e*f^2 + c*d*f^3)*x^3 + e^2*f + (d^2*e^2*f + 4*c*d*e*f^2 + c^2*f^3 + f^3)*x^2 + 2*(c*d*e^2*f + c^2*e*f^2 + e*f^2)*x)*sqrt(d^2*x^2 + 2*c*d*x + c^2 + 1)), x)
```

Giac [F]

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^2} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{(fx + e)^2} dx$$

```
[In] integrate((a+b*arccsch(d*x+c))^2/(f*x+e)^2,x, algorithm="giac")
```

```
[Out] integrate((b*arccsch(d*x + c) + a)^2/(f*x + e)^2, x)
```


Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^2} dx = \int \frac{(a + b \operatorname{asinh}(\frac{1}{c+dx}))^2}{(e + fx)^2} dx$$

```
[In] int((a + b*asinh(1/(c + d*x)))^2/(e + f*x)^2,x)
```

```
[Out] int((a + b*asinh(1/(c + d*x)))^2/(e + f*x)^2, x)
```

3.13
$$\int \frac{\left(a+b\operatorname{csch}^{-1}(c+dx)\right)^2}{(e+fx)^3} dx$$

Optimal result	139
Rubi [A] (verified)	140
Mathematica [C] (warning: unable to verify)	150
Maple [F]	150
Fricas [F]	150
Sympy [F(-1)]	150
Maxima [F]	151
Giac [F]	151
Mupad [F(-1)]	152

Optimal result

Integrand size = 20, antiderivative size = 1024

$$\begin{aligned}
& \int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^3} dx \\
&= -\frac{bd^2 f \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{(de - cf) (d^2 e^2 - 2cdef + (1 + c^2) f^2) \left(f + \frac{de - cf}{c + dx}\right)} \\
&+ \frac{d^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2f(de - cf)^2} - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{2f(e + fx)^2} \\
&+ \frac{bd^2 f^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}\right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&- \frac{2bd^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}\right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&- \frac{bd^2 f^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}\right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&+ \frac{2bd^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log\left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}\right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&+ \frac{b^2 d^2 f \log\left(f + \frac{de - cf}{c + dx}\right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)} \\
&+ \frac{b^2 d^2 f^2 \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}\right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&- \frac{2b^2 d^2 \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}\right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&- \frac{b^2 d^2 f^2 \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}\right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&+ \frac{2b^2 d^2 \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}\right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}
\end{aligned}$$

[Out] 1/2*d^2*(a+b*arccsch(d*x+c))^2/f/(-c*f+d*e)^2-1/2*(a+b*arccsch(d*x+c))^2/f/(f*x+e)^2+b^2*d^2*f*ln(f+(-c*f+d*e)/(d*x+c))/(-c*f+d*e)^2/(d^2*e^2-2*c*d*e*

$$\begin{aligned}
& f+(c^2+1)*f^2)+b*d^2*f^2*(a+b*\arccsch(d*x+c))*\ln(1+(1/(d*x+c)+(1+1/(d*x+c)^2)^{1/2}))*(-c*f+d*e)/(f-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{1/2}))/(-c*f+d*e)^2/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{3/2}-b*d^2*f^2*(a+b*\arccsch(d*x+c))*\ln(1+(1/(d*x+c)+(1+1/(d*x+c)^2)^{1/2}))*(-c*f+d*e)/(f+(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{1/2}))/(-c*f+d*e)^2/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{3/2}+b^2*d^2*f^2*\text{polylog}(2,-(1/(d*x+c)+(1+1/(d*x+c)^2)^{1/2}))*(-c*f+d*e)/(f-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{1/2}))/(-c*f+d*e)^2/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{3/2}-b^2*d^2*f^2*\text{polylog}(2,-(1/(d*x+c)+(1+1/(d*x+c)^2)^{1/2}))*(-c*f+d*e)/(f+(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{1/2}))/(-c*f+d*e)^2/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{3/2}-2*b*d^2*(a+b*\arccsch(d*x+c))*\ln(1+(1/(d*x+c)+(1+1/(d*x+c)^2)^{1/2}))*(-c*f+d*e)/(f-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{1/2}))/(-c*f+d*e)^2/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{1/2}+2*b*d^2*(a+b*\arccsch(d*x+c))*\ln(1+(1/(d*x+c)+(1+1/(d*x+c)^2)^{1/2}))*(-c*f+d*e)/(f+(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{1/2}))/(-c*f+d*e)^2/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{1/2}-2*b^2*d^2*\text{polylog}(2,-(1/(d*x+c)+(1+1/(d*x+c)^2)^{1/2}))*(-c*f+d*e)/(f-(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{1/2}))/(-c*f+d*e)^2/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{1/2}+2*b^2*d^2*\text{polylog}(2,-(1/(d*x+c)+(1+1/(d*x+c)^2)^{1/2}))*(-c*f+d*e)/(f+(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{1/2}))/(-c*f+d*e)^2/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^{1/2}-b*d^2*f*(a+b*\arccsch(d*x+c))*(1+1/(d*x+c)^2)^{1/2}/(-c*f+d*e)/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)/(f+(-c*f+d*e)/(d*x+c))
\end{aligned}$$

Rubi [A] (verified)

Time = 1.62 (sec) , antiderivative size = 1024, normalized size of antiderivative = 1.00, number of steps used = 23, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.550$, Rules

used = {6457, 5577, 4276, 3403, 2296, 2221, 2317, 2438, 3405, 2747, 31}

$$\begin{aligned}
& \int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^3} dx \\
&= \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2 d^2}{2f(de - cf)^2} - \frac{bf \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx)) d^2}{(de - cf)(d^2 e^2 - 2cdf e + (c^2 + 1)f^2) \left(f + \frac{de - cf}{c + dx}\right)} \\
&\quad - \frac{2b(a + b \operatorname{csch}^{-1}(c + dx)) \log\left(\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdf e + (c^2 + 1)f^2}} + 1\right) d^2}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdf e + (c^2 + 1)f^2}} \\
&\quad + \frac{bf^2(a + b \operatorname{csch}^{-1}(c + dx)) \log\left(\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdf e + (c^2 + 1)f^2}} + 1\right) d^2}{(de - cf)^2 (d^2 e^2 - 2cdf e + (c^2 + 1)f^2)^{3/2}} \\
&\quad + \frac{2b(a + b \operatorname{csch}^{-1}(c + dx)) \log\left(\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdf e + (c^2 + 1)f^2}} + 1\right) d^2}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdf e + (c^2 + 1)f^2}} \\
&\quad - \frac{bf^2(a + b \operatorname{csch}^{-1}(c + dx)) \log\left(\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdf e + (c^2 + 1)f^2}} + 1\right) d^2}{(de - cf)^2 (d^2 e^2 - 2cdf e + (c^2 + 1)f^2)^{3/2}} \\
&\quad + \frac{b^2 f \log\left(f + \frac{de - cf}{c + dx}\right) d^2}{(de - cf)^2 (d^2 e^2 - 2cdf e + (c^2 + 1)f^2)} - \frac{2b^2 \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdf e + (c^2 + 1)f^2}}\right) d^2}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdf e + (c^2 + 1)f^2}} \\
&\quad + \frac{b^2 f^2 \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdf e + (c^2 + 1)f^2}}\right) d^2}{(de - cf)^2 (d^2 e^2 - 2cdf e + (c^2 + 1)f^2)^{3/2}} \\
&\quad + \frac{2b^2 \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdf e + (c^2 + 1)f^2}}\right) d^2}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdf e + (c^2 + 1)f^2}} \\
&\quad - \frac{b^2 f^2 \operatorname{PolyLog}\left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdf e + (c^2 + 1)f^2}}\right) d^2}{(de - cf)^2 (d^2 e^2 - 2cdf e + (c^2 + 1)f^2)^{3/2}} - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{2f(e + fx)^2}
\end{aligned}$$

[In] Int[(a + b*ArcCsch[c + d*x])^2/(e + f*x)^3,x]

[Out] -((b*d^2*f*Sqrt[1 + (c + d*x)^(-2)]*(a + b*ArcCsch[c + d*x]))/((d*e - c*f)*(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2)*(f + (d*e - c*f)/(c + d*x)))) + (d^2*(a + b*ArcCsch[c + d*x])^2)/(2*f*(d*e - c*f)^2) - (a + b*ArcCsch[c + d*x])^2/(2*f*(e + f*x)^2) + (b*d^2*f^2*(a + b*ArcCsch[c + d*x])*Log[1 + (E^ArcCsch[c + d*x]*(d*e - c*f))/(f - Sqrt[d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2]])/((d*e - c*f)^2*(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2)^(3/2)) - (2*b*d^2*(a + b*ArcCsch[c + d*x])*Log[1 + (E^ArcCsch[c + d*x]*(d*e - c*f))/(f - Sqrt[d^2*

$$\frac{e^2 - 2cd*ef + (1 + c^2)*f^2)}{(d*e - c*f)^2*\sqrt{d^2*e^2 - 2cd*ef + (1 + c^2)*f^2}} - \frac{(b*d^2*f^2*(a + b*\text{ArcCsch}[c + d*x])*\text{Log}[1 + (E^{\text{ArcCsch}[c + d*x]}*(d*e - c*f))]/(f + \sqrt{d^2*e^2 - 2cd*ef + (1 + c^2)*f^2})]}{(d*e - c*f)^2*(d^2*e^2 - 2cd*ef + (1 + c^2)*f^2)^{3/2}} + \frac{(2*b*d^2*(a + b*\text{ArcCsch}[c + d*x])*\text{Log}[1 + (E^{\text{ArcCsch}[c + d*x]}*(d*e - c*f))]/(f + \sqrt{d^2*e^2 - 2cd*ef + (1 + c^2)*f^2})]}{(d*e - c*f)^2*\sqrt{d^2*e^2 - 2cd*ef + (1 + c^2)*f^2}} + \frac{(b^2*d^2*f*\text{Log}[f + (d*e - c*f)/(c + d*x)])}{(d*e - c*f)^2*(d^2*e^2 - 2cd*ef + (1 + c^2)*f^2)} + \frac{(b^2*d^2*f^2*\text{PolyLog}[2, -(E^{\text{ArcCsch}[c + d*x]}*(d*e - c*f))]/(f - \sqrt{d^2*e^2 - 2cd*ef + (1 + c^2)*f^2})]}{(d*e - c*f)^2*(d^2*e^2 - 2cd*ef + (1 + c^2)*f^2)^{3/2}} - \frac{(2*b^2*d^2*\text{PolyLog}[2, -(E^{\text{ArcCsch}[c + d*x]}*(d*e - c*f))]/(f - \sqrt{d^2*e^2 - 2cd*ef + (1 + c^2)*f^2})]}{(d*e - c*f)^2*(d^2*e^2 - 2cd*ef + (1 + c^2)*f^2)^{3/2}} - \frac{(2*b^2*d^2*\text{PolyLog}[2, -(E^{\text{ArcCsch}[c + d*x]}*(d*e - c*f))]/(f + \sqrt{d^2*e^2 - 2cd*ef + (1 + c^2)*f^2})]}{(d*e - c*f)^2*(d^2*e^2 - 2cd*ef + (1 + c^2)*f^2)^{3/2}} + \frac{(2*b^2*d^2*\text{PolyLog}[2, -(E^{\text{ArcCsch}[c + d*x]}*(d*e - c*f))]/(f + \sqrt{d^2*e^2 - 2cd*ef + (1 + c^2)*f^2})]}{(d*e - c*f)^2*\sqrt{d^2*e^2 - 2cd*ef + (1 + c^2)*f^2}}$$
Rule 31

```
Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]
```

Rule 2221

```
Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2296

```
Int[((F_)^(u_)*((f_.) + (g_.)*(x_))^(m_.))/((a_.) + (b_.)*(F_)^(u_) + (c_.)*(F_)^(v_)), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[2*(c/q), Int[(f + g*x)^m*(F^u/(b - q + 2*c*F^u)), x], x] - Dist[2*(c/q), Int[(f + g*x)^m*(F^u/(b + q + 2*c*F^u)), x], x]] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v, 2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*(d_) + (e_.)*(x_)^(n_.)]/(x_), x_Symbol] := Simp[-PolyLog[2,
(-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2747

```
Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m
_.), x_Symbol] := Dist[1/(b^p*f), Subst[Int[(a + x)^m*(b^2 - x^2)^((p - 1)/
2), x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p
- 1)/2] && NeQ[a^2 - b^2, 0]
```

Rule 3403

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*sin[(e_.) + (Complex[0, fz_]*)
(f_.)*(x_)]), x_Symbol] := Dist[2, Int[(c + d*x)^m*(E^((-I)*e + f*fz*x)/((-
I)*b + 2*a*E^((-I)*e + f*fz*x) + I*b*E^(2*((-I)*e + f*fz*x)))]), x], x] /; F
reeQ[{a, b, c, d, e, f, fz}, x] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]
```

Rule 3405

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2, x_
Symbol] := Simp[b*(c + d*x)^m*(Cos[e + f*x]/(f*(a^2 - b^2)*(a + b*Sin[e + f
*x]))], x] + (Dist[a/(a^2 - b^2), Int[(c + d*x)^m/(a + b*Sin[e + f*x]), x],
x] - Dist[b*d*(m/(f*(a^2 - b^2))), Int[(c + d*x)^(m - 1)*(Cos[e + f*x]/(a
+ b*Sin[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 - b^
2, 0] && IGtQ[m, 0]
```

Rule 4276

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)
, x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Si
n[e + f*x])^n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGt
Q[m, 0]
```

Rule 5577

```
Int[Coth[(c_.) + (d_.)*(x_)]*Csch[(c_.) + (d_.)*(x_)]*(Csch[(c_.) + (d_.)*(
x_)]*(b_.) + (a_))^(n_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(-e
+ f*x)^m*((a + b*Csch[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b
*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csch[c + d*x])^(n + 1), x], x] /
; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 6457

```
Int[((a_.) + ArcCsch[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] := Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csch[x]*Co
th[x]*(d*e - c*f + f*Csch[x])^m, x], x, ArcCsch[c + d*x]], x] /; FreeQ[{a,
```

b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rubi steps

$$\begin{aligned}
\text{integral} &= - \left(d^2 \text{Subst} \left(\int \frac{(a+bx)^2 \coth(x) \operatorname{csch}(x)}{(de-cf+f \operatorname{csch}(x))^3} dx, x, \operatorname{csch}^{-1}(c+dx) \right) \right) \\
&= - \frac{(a+b \operatorname{csch}^{-1}(c+dx))^2}{2f(e+fx)^2} + \frac{(bd^2) \text{Subst} \left(\int \frac{a+bx}{(de-cf+f \operatorname{csch}(x))^2} dx, x, \operatorname{csch}^{-1}(c+dx) \right)}{f} \\
&= - \frac{(a+b \operatorname{csch}^{-1}(c+dx))^2}{2f(e+fx)^2} \\
&\quad + \frac{(bd^2) \text{Subst} \left(\int \left(\frac{a+bx}{(de-cf)^2} + \frac{2f(a+bx)}{(de-cf)^2 (-f-de(1-\frac{cf}{de}) \sinh(x))} + \frac{f^2(a+bx)}{(de-cf)^2 (f+de(1-\frac{cf}{de}) \sinh(x))^2} \right) dx, x, \operatorname{csch}^{-1}(c+dx) \right)}{f} \\
&= \frac{d^2(a+b \operatorname{csch}^{-1}(c+dx))^2}{2f(de-cf)^2} - \frac{(a+b \operatorname{csch}^{-1}(c+dx))^2}{2f(e+fx)^2} \\
&\quad + \frac{(2bd^2) \text{Subst} \left(\int \frac{a+bx}{-f-de(1-\frac{cf}{de}) \sinh(x)} dx, x, \operatorname{csch}^{-1}(c+dx) \right)}{(de-cf)^2} \\
&\quad + \frac{(bd^2 f) \text{Subst} \left(\int \frac{a+bx}{(f+de(1-\frac{cf}{de}) \sinh(x))^2} dx, x, \operatorname{csch}^{-1}(c+dx) \right)}{(de-cf)^2} \\
&= - \frac{bd^2 f \sqrt{1 + \frac{1}{(c+dx)^2}} (a+b \operatorname{csch}^{-1}(c+dx))}{(de-cf)(d^2 e^2 - 2cdef + (1+c^2)f^2)(f + \frac{de-cf}{c+dx})} \\
&\quad + \frac{d^2(a+b \operatorname{csch}^{-1}(c+dx))^2}{2f(de-cf)^2} - \frac{(a+b \operatorname{csch}^{-1}(c+dx))^2}{2f(e+fx)^2} \\
&\quad + \frac{(4bd^2) \text{Subst} \left(\int \frac{e^x(a+bx)}{-2e^x f + de(1-\frac{cf}{de}) - dee^{2x}(1-\frac{cf}{de})} dx, x, \operatorname{csch}^{-1}(c+dx) \right)}{(de-cf)^2} \\
&\quad + \frac{(bd^2 f^2) \text{Subst} \left(\int \frac{a+bx}{f+de(1-\frac{cf}{de}) \sinh(x)} dx, x, \operatorname{csch}^{-1}(c+dx) \right)}{(de-cf)^2 (d^2 e^2 - 2cdef + (1+c^2)f^2)} \\
&\quad + \frac{(b^2 d^2 f) \text{Subst} \left(\int \frac{\cosh(x)}{f+de(1-\frac{cf}{de}) \sinh(x)} dx, x, \operatorname{csch}^{-1}(c+dx) \right)}{(de-cf)(d^2 e^2 - 2cdef + (1+c^2)f^2)}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{bd^2 f \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{(de - cf) (d^2 e^2 - 2cdef + (1 + c^2) f^2) \left(f + \frac{de - cf}{c + dx}\right)} + \frac{d^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2f (de - cf)^2} \\
&\quad - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{2f (e + fx)^2} + \frac{(b^2 d^2 f) \operatorname{Subst}\left(\int \frac{1}{f + x} dx, x, \frac{de \left(1 - \frac{cf}{de}\right)}{c + dx}\right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)} \\
&\quad + \frac{(2bd^2 f^2) \operatorname{Subst}\left(\int \frac{e^x (a + bx)}{2e^x f - de \left(1 - \frac{cf}{de}\right) + de e^{2x} \left(1 - \frac{cf}{de}\right)} dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)} \\
&\quad - \frac{(4bd^2) \operatorname{Subst}\left(\int \frac{e^x (a + bx)}{-2f - 2de e^x \left(1 - \frac{cf}{de}\right) - 2\sqrt{d^2 e^2 - 2cdef + f^2 + c^2 f^2}} dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{(de - cf) \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&\quad + \frac{(4bd^2) \operatorname{Subst}\left(\int \frac{e^x (a + bx)}{-2f - 2de e^x \left(1 - \frac{cf}{de}\right) + 2\sqrt{d^2 e^2 - 2cdef + f^2 + c^2 f^2}} dx, x, \operatorname{csch}^{-1}(c + dx)\right)}{(de - cf) \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{bd^2 f \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{(de - cf) (d^2 e^2 - 2cdef + (1 + c^2) f^2) (f + \frac{de - cf}{c + dx})} \\
&+ \frac{d^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2f(de - cf)^2} - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{2f(e + fx)^2} \\
&- \frac{2bd^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log \left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&+ \frac{2bd^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log \left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&+ \frac{b^2 d^2 f \log \left(\frac{e + fx}{c + dx} \right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)} \\
&+ \frac{(2bd^2 f^2) \operatorname{Subst} \left(\int \frac{e^x (a + bx)}{2f + 2dee^x \left(1 - \frac{cf}{de} \right) - 2\sqrt{d^2 e^2 - 2cdef + f^2 + c^2 f^2}} dx, x, \operatorname{csch}^{-1}(c + dx) \right)}{(de - cf) (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&- \frac{(2bd^2 f^2) \operatorname{Subst} \left(\int \frac{e^x (a + bx)}{2f + 2dee^x \left(1 - \frac{cf}{de} \right) + 2\sqrt{d^2 e^2 - 2cdef + f^2 + c^2 f^2}} dx, x, \operatorname{csch}^{-1}(c + dx) \right)}{(de - cf) (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&- \frac{(2b^2 d^2) \operatorname{Subst} \left(\int \log \left(1 - \frac{2dee^x \left(1 - \frac{cf}{de} \right)}{-2f - 2\sqrt{d^2 e^2 - 2cdef + f^2 + c^2 f^2}} \right) dx, x, \operatorname{csch}^{-1}(c + dx) \right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&+ \frac{(2b^2 d^2) \operatorname{Subst} \left(\int \log \left(1 - \frac{2dee^x \left(1 - \frac{cf}{de} \right)}{-2f + 2\sqrt{d^2 e^2 - 2cdef + f^2 + c^2 f^2}} \right) dx, x, \operatorname{csch}^{-1}(c + dx) \right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{bd^2 f \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{(de - cf) (d^2 e^2 - 2cdef + (1 + c^2) f^2) \left(f + \frac{de - cf}{c + dx}\right)} \\
&+ \frac{d^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2f(de - cf)^2} - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{2f(e + fx)^2} \\
&+ \frac{bd^2 f^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log \left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}\right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&- \frac{2bd^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log \left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}\right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&- \frac{bd^2 f^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log \left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}\right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&+ \frac{2bd^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log \left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}\right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&+ \frac{b^2 d^2 f \log \left(\frac{e + fx}{c + dx}\right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)} \\
&- \frac{(b^2 d^2 f^2) \operatorname{Subst} \left(\int \log \left(1 + \frac{2de e^x \left(1 - \frac{cf}{de}\right)}{2f - 2\sqrt{d^2 e^2 - 2cdef + f^2 + c^2 f^2}}\right) dx, x, \operatorname{csch}^{-1}(c + dx) \right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&+ \frac{(b^2 d^2 f^2) \operatorname{Subst} \left(\int \log \left(1 + \frac{2de e^x \left(1 - \frac{cf}{de}\right)}{2f + 2\sqrt{d^2 e^2 - 2cdef + f^2 + c^2 f^2}}\right) dx, x, \operatorname{csch}^{-1}(c + dx) \right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&- \frac{(2b^2 d^2) \operatorname{Subst} \left(\int \frac{\log \left(1 - \frac{2de \left(1 - \frac{cf}{de}\right) x}{-2f - 2\sqrt{d^2 e^2 - 2cdef + f^2 + c^2 f^2}}\right)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)} \right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&+ \frac{(2b^2 d^2) \operatorname{Subst} \left(\int \frac{\log \left(1 - \frac{2de \left(1 - \frac{cf}{de}\right) x}{-2f + 2\sqrt{d^2 e^2 - 2cdef + f^2 + c^2 f^2}}\right)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)} \right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{bd^2 f \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{(de - cf) (d^2 e^2 - 2cdef + (1 + c^2) f^2) (f + \frac{de - cf}{c + dx})} \\
&+ \frac{d^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2f(de - cf)^2} - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{2f(e + fx)^2} \\
&+ \frac{bd^2 f^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log \left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&- \frac{2bd^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log \left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&- \frac{bd^2 f^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log \left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&+ \frac{2bd^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log \left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&+ \frac{b^2 d^2 f \log \left(\frac{e + fx}{c + dx} \right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)} \\
&- \frac{2b^2 d^2 \operatorname{PolyLog} \left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&+ \frac{2b^2 d^2 \operatorname{PolyLog} \left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&- \frac{(b^2 d^2 f^2) \operatorname{Subst} \left(\int \frac{\log \left(1 + \frac{2de(1 - \frac{cf}{de})x}{2f - 2\sqrt{d^2 e^2 - 2cdef + f^2 + c^2 f^2}} \right)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)} \right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&+ \frac{(b^2 d^2 f^2) \operatorname{Subst} \left(\int \frac{\log \left(1 + \frac{2de(1 - \frac{cf}{de})x}{2f + 2\sqrt{d^2 e^2 - 2cdef + f^2 + c^2 f^2}} \right)}{x} dx, x, e^{\operatorname{csch}^{-1}(c+dx)} \right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{bd^2 f \sqrt{1 + \frac{1}{(c+dx)^2}} (a + b \operatorname{csch}^{-1}(c + dx))}{(de - cf) (d^2 e^2 - 2cdef + (1 + c^2) f^2) (f + \frac{de - cf}{c + dx})} \\
&+ \frac{d^2 (a + b \operatorname{csch}^{-1}(c + dx))^2}{2f(de - cf)^2} - \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{2f(e + fx)^2} \\
&+ \frac{bd^2 f^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log \left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&- \frac{2bd^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log \left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&- \frac{bd^2 f^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log \left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&+ \frac{2bd^2 (a + b \operatorname{csch}^{-1}(c + dx)) \log \left(1 + \frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&+ \frac{b^2 d^2 f \log \left(\frac{e + fx}{c + dx} \right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)} \\
&+ \frac{b^2 d^2 f^2 \operatorname{PolyLog} \left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&- \frac{2b^2 d^2 \operatorname{PolyLog} \left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f - \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \\
&- \frac{b^2 d^2 f^2 \operatorname{PolyLog} \left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 (d^2 e^2 - 2cdef + (1 + c^2) f^2)^{3/2}} \\
&+ \frac{2b^2 d^2 \operatorname{PolyLog} \left(2, -\frac{e^{\operatorname{csch}^{-1}(c+dx)}(de - cf)}{f + \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}} \right)}{(de - cf)^2 \sqrt{d^2 e^2 - 2cdef + (1 + c^2) f^2}}
\end{aligned}$$

Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 13.37 (sec) , antiderivative size = 8350, normalized size of antiderivative = 8.15

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^3} dx = \text{Result too large to show}$$

[In] Integrate[(a + b*ArcCsch[c + d*x])^2/(e + f*x)^3,x]

[Out] Result too large to show

Maple [F]

$$\int \frac{(a + b \operatorname{arccsch}(dx + c))^2}{(fx + e)^3} dx$$

[In] int((a+b*arccsch(d*x+c))^2/(f*x+e)^3,x)

[Out] int((a+b*arccsch(d*x+c))^2/(f*x+e)^3,x)

Fricas [F]

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^3} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{(fx + e)^3} dx$$

[In] integrate((a+b*arccsch(d*x+c))^2/(f*x+e)^3,x, algorithm="fricas")

[Out] integral((b^2*arccsch(d*x + c)^2 + 2*a*b*arccsch(d*x + c) + a^2)/(f^3*x^3 + 3*e*f^2*x^2 + 3*e^2*f*x + e^3), x)

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^3} dx = \text{Timed out}$$

[In] integrate((a+b*acsch(d*x+c))**2/(f*x+e)**3,x)

[Out] Timed out

Maxima [F]

$$\int \frac{(a + b \operatorname{arcsch}^{-1}(c + dx))^2}{(e + fx)^3} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{(fx + e)^3} dx$$

[In] integrate((a+b*arccsch(d*x+c))^2/(f*x+e)^3,x, algorithm="maxima")

[Out] $-1/2*b^2*\log(\sqrt{d^2*x^2 + 2*c*d*x + c^2 + 1} + 1)^2/(f^3*x^2 + 2*e*f^2*x + e^2*f) - 1/2*a^2/(f^3*x^2 + 2*e*f^2*x + e^2*f) - \operatorname{integrate}(-((b^2*d^2*f*x^2 + 2*b^2*c*d*f*x + (c^2*f + f)*b^2)*\log(d*x + c)^2 - 2*(a*b*d^2*f*x^2 + 2*a*b*c*d*f*x + (c^2*f + f)*a*b)*\log(d*x + c) + (2*a*b*d^2*f*x^2 + 4*a*b*c*d*f*x + 2*(c^2*f + f)*a*b - 2*(b^2*d^2*f*x^2 + 2*b^2*c*d*f*x + (c^2*f + f)*b^2)*\log(d*x + c) + (b^2*c*d*e + 2*(c^2*f + f)*a*b + (2*a*b*d^2*f + b^2*d^2*f)*x^2 + (4*a*b*c*d*f + (d^2*e + c*d*f)*b^2)*x - 2*(b^2*d^2*f*x^2 + 2*b^2*c*d*f*x + (c^2*f + f)*b^2)*\log(d*x + c))*\sqrt{d^2*x^2 + 2*c*d*x + c^2 + 1})*\log(\sqrt{d^2*x^2 + 2*c*d*x + c^2 + 1} + 1) + \sqrt{d^2*x^2 + 2*c*d*x + c^2 + 1}*((b^2*d^2*f*x^2 + 2*b^2*c*d*f*x + (c^2*f + f)*b^2)*\log(d*x + c)^2 - 2*(a*b*d^2*f*x^2 + 2*a*b*c*d*f*x + (c^2*f + f)*a*b)*\log(d*x + c)))/(d^2*f^4*x^5 + c^2*e^3*f + (3*d^2*e*f^3 + 2*c*d*f^4)*x^4 + e^3*f + (3*d^2*e^2*f^2 + 6*c*d*e*f^3 + c^2*f^4 + f^4)*x^3 + (d^2*e^3*f + 6*c*d*e^2*f^2 + 3*c^2*e*f^3 + 3*e*f^3)*x^2 + (2*c*d*e^3*f + 3*c^2*e^2*f^2 + 3*e^2*f^2)*x + (d^2*f^4*x^5 + c^2*e^3*f + (3*d^2*e*f^3 + 2*c*d*f^4)*x^4 + e^3*f + (3*d^2*e^2*f^2 + 6*c*d*e*f^3 + c^2*f^4 + f^4)*x^3 + (d^2*e^3*f + 6*c*d*e^2*f^2 + 3*c^2*e*f^3 + 3*e*f^3)*x^2 + (2*c*d*e^3*f + 3*c^2*e^2*f^2 + 3*e^2*f^2)*x)*\sqrt{d^2*x^2 + 2*c*d*x + c^2 + 1}), x)$

Giac [F]

$$\int \frac{(a + b \operatorname{arcsch}^{-1}(c + dx))^2}{(e + fx)^3} dx = \int \frac{(b \operatorname{arcsch}(dx + c) + a)^2}{(fx + e)^3} dx$$

[In] integrate((a+b*arccsch(d*x+c))^2/(f*x+e)^3,x, algorithm="giac")

[Out] integrate((b*arccsch(d*x + c) + a)^2/(f*x + e)^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{csch}^{-1}(c + dx))^2}{(e + fx)^3} dx = \int \frac{(a + b \operatorname{asinh}(\frac{1}{c+dx}))^2}{(e + fx)^3} dx$$

```
[In] int((a + b*asinh(1/(c + d*x)))^2/(e + f*x)^3,x)
```

```
[Out] int((a + b*asinh(1/(c + d*x)))^2/(e + f*x)^3, x)
```


3.14 $\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx$

Optimal result	153
Rubi [A] (verified)	153
Mathematica [A] (verified)	154
Maple [A] (verified)	155
Fricas [A] (verification not implemented)	155
Sympy [F]	155
Maxima [A] (verification not implemented)	156
Giac [F]	156
Mupad [F(-1)]	156

Optimal result

Integrand size = 10, antiderivative size = 114

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = -\frac{\sqrt{-1-x}\sqrt{x}}{4\sqrt{-x}} - \frac{(-1-x)^{3/2}\sqrt{x}}{4\sqrt{-x}} - \frac{3(-1-x)^{5/2}\sqrt{x}}{20\sqrt{-x}} - \frac{(-1-x)^{7/2}\sqrt{x}}{28\sqrt{-x}} + \frac{1}{4}x^4 \operatorname{csch}^{-1}(\sqrt{x})$$

[Out] 1/4*x^4*arccsch(x^(1/2))-1/4*(-1-x)^(3/2)*x^(1/2)/(-x)^(1/2)-3/20*(-1-x)^(5/2)*x^(1/2)/(-x)^(1/2)-1/28*(-1-x)^(7/2)*x^(1/2)/(-x)^(1/2)-1/4*(-1-x)^(1/2)*x^(1/2)/(-x)^(1/2)

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {6481, 12, 45}

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{4}x^4 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{(-x-1)^{7/2}\sqrt{x}}{28\sqrt{-x}} - \frac{3(-x-1)^{5/2}\sqrt{x}}{20\sqrt{-x}} - \frac{(-x-1)^{3/2}\sqrt{x}}{4\sqrt{-x}} - \frac{\sqrt{-x-1}\sqrt{x}}{4\sqrt{-x}}$$

[In] Int[x^3*ArcCsch[Sqrt[x]],x]

[Out] -1/4*(Sqrt[-1-x]*Sqrt[x])/Sqrt[-x] - ((-1-x)^(3/2)*Sqrt[x])/(4*Sqrt[-x]) - (3*(-1-x)^(5/2)*Sqrt[x])/(20*Sqrt[-x]) - ((-1-x)^(7/2)*Sqrt[x])/(28*Sqrt[-x]) + (x^4*ArcCsch[Sqrt[x]])/4

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 45

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

Rule 6481

```
Int[((a_.) + ArcCsch[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcCsch[u])/(d*(m + 1))), x] - Dist[b*(u/(d*(m + 1)*Sqrt[-u^2])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[-1 - u^2])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{4}x^4 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \frac{x^3}{2\sqrt{-1-x}} dx}{4\sqrt{-x}} \\
 &= \frac{1}{4}x^4 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \frac{x^3}{\sqrt{-1-x}} dx}{8\sqrt{-x}} \\
 &= \frac{1}{4}x^4 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \left(-\frac{1}{\sqrt{-1-x}} - 3\sqrt{-1-x} - 3(-1-x)^{3/2} - (-1-x)^{5/2} \right) dx}{8\sqrt{-x}} \\
 &= -\frac{\sqrt{-1-x}\sqrt{x}}{4\sqrt{-x}} - \frac{(-1-x)^{3/2}\sqrt{x}}{4\sqrt{-x}} - \frac{3(-1-x)^{5/2}\sqrt{x}}{20\sqrt{-x}} - \frac{(-1-x)^{7/2}\sqrt{x}}{28\sqrt{-x}} + \frac{1}{4}x^4 \operatorname{csch}^{-1}(\sqrt{x})
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.41

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{140} \sqrt{1 + \frac{1}{x}} \sqrt{x} (-16 + 8x - 6x^2 + 5x^3) + \frac{1}{4} x^4 \operatorname{csch}^{-1}(\sqrt{x})$$

```
[In] Integrate[x^3*ArcCsch[Sqrt[x]], x]
```

```
[Out] (Sqrt[1 + x^(-1)]*Sqrt[x]*(-16 + 8*x - 6*x^2 + 5*x^3))/140 + (x^4*ArcCsch[Sqrt[x]])/4
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.35

method	result	size
parts	$\frac{x^4 \operatorname{arccsch}(\sqrt{x})}{4} + \frac{\sqrt{\frac{1+x}{x}} \sqrt{x} (5x^3 - 6x^2 + 8x - 16)}{140}$	40
derivativedivides	$\frac{x^4 \operatorname{arccsch}(\sqrt{x})}{4} + \frac{(1+x)(5x^3 - 6x^2 + 8x - 16)}{140\sqrt{\frac{1+x}{x}} \sqrt{x}}$	43
default	$\frac{x^4 \operatorname{arccsch}(\sqrt{x})}{4} + \frac{(1+x)(5x^3 - 6x^2 + 8x - 16)}{140\sqrt{\frac{1+x}{x}} \sqrt{x}}$	43

[In] `int(x^3*arccsch(x^(1/2)),x,method=_RETURNVERBOSE)`

[Out] `1/4*x^4*arccsch(x^(1/2))+1/140*((1+x)/x)^(1/2)*x^(1/2)*(5*x^3-6*x^2+8*x-16)`

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.48

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{4} x^4 \log \left(\frac{x \sqrt{\frac{x+1}{x}} + \sqrt{x}}{x} \right) + \frac{1}{140} (5x^3 - 6x^2 + 8x - 16) \sqrt{x} \sqrt{\frac{x+1}{x}}$$

[In] `integrate(x^3*arccsch(x^(1/2)),x, algorithm="fricas")`

[Out] `1/4*x^4*log((x*sqrt((x + 1)/x) + sqrt(x))/x) + 1/140*(5*x^3 - 6*x^2 + 8*x - 16)*sqrt(x)*sqrt((x + 1)/x)`

Sympy [F]

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = \int x^3 \operatorname{acsch}(\sqrt{x}) dx$$

[In] `integrate(x**3*acsch(x**(1/2)),x)`

[Out] `Integral(x**3*acsch(sqrt(x)), x)`

Maxima [A] (verification not implemented)

none

Time = 0.20 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.51

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{28} x^{\frac{7}{2}} \left(\frac{1}{x} + 1\right)^{\frac{7}{2}} - \frac{3}{20} x^{\frac{5}{2}} \left(\frac{1}{x} + 1\right)^{\frac{5}{2}} + \frac{1}{4} x^4 \operatorname{arcsch}(\sqrt{x}) + \frac{1}{4} x^{\frac{3}{2}} \left(\frac{1}{x} + 1\right)^{\frac{3}{2}} - \frac{1}{4} \sqrt{x} \sqrt{\frac{1}{x} + 1}$$

[In] integrate(x^3*arccsch(x^(1/2)),x, algorithm="maxima")

[Out] 1/28*x^(7/2)*(1/x + 1)^(7/2) - 3/20*x^(5/2)*(1/x + 1)^(5/2) + 1/4*x^4*arccsch(sqrt(x)) + 1/4*x^(3/2)*(1/x + 1)^(3/2) - 1/4*sqrt(x)*sqrt(1/x + 1)

Giac [F]

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = \int x^3 \operatorname{arcsch}(\sqrt{x}) dx$$

[In] integrate(x^3*arccsch(x^(1/2)),x, algorithm="giac")

[Out] integrate(x^3*arccsch(sqrt(x)), x)

Mupad [F(-1)]

Timed out.

$$\int x^3 \operatorname{csch}^{-1}(\sqrt{x}) dx = \int x^3 \operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right) dx$$

[In] int(x^3*asinh(1/x^(1/2)),x)

[Out] int(x^3*asinh(1/x^(1/2)), x)

3.15 $\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx$

Optimal result	157
Rubi [A] (verified)	157
Mathematica [A] (verified)	158
Maple [A] (verified)	159
Fricas [A] (verification not implemented)	159
Sympy [F]	159
Maxima [A] (verification not implemented)	160
Giac [F]	160
Mupad [F(-1)]	160

Optimal result

Integrand size = 10, antiderivative size = 89

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{\sqrt{-1-x}\sqrt{x}}{3\sqrt{-x}} + \frac{2(-1-x)^{3/2}\sqrt{x}}{9\sqrt{-x}} + \frac{(-1-x)^{5/2}\sqrt{x}}{15\sqrt{-x}} + \frac{1}{3}x^3 \operatorname{csch}^{-1}(\sqrt{x})$$

[Out] $\frac{1}{3}x^3 \operatorname{arccsch}(x^{1/2}) + \frac{2}{9}(-1-x)^{3/2}x^{1/2}/(-x)^{1/2} + \frac{1}{15}(-1-x)^{5/2}x^{1/2}/(-x)^{1/2} + \frac{1}{3}x^3 \operatorname{arccsch}(x^{1/2})$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {6481, 12, 45}

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{3}x^3 \operatorname{csch}^{-1}(\sqrt{x}) + \frac{(-x-1)^{5/2}\sqrt{x}}{15\sqrt{-x}} + \frac{2(-x-1)^{3/2}\sqrt{x}}{9\sqrt{-x}} + \frac{\sqrt{-x-1}\sqrt{x}}{3\sqrt{-x}}$$

[In] $\operatorname{Int}[x^2 \operatorname{ArcCsch}[\operatorname{Sqrt}[x]], x]$

[Out] $(\operatorname{Sqrt}[-1-x] \operatorname{Sqrt}[x]) / (3 \operatorname{Sqrt}[-x]) + (2(-1-x)^{3/2} \operatorname{Sqrt}[x]) / (9 \operatorname{Sqrt}[-x]) + ((-1-x)^{5/2} \operatorname{Sqrt}[x]) / (15 \operatorname{Sqrt}[-x]) + (x^3 \operatorname{ArcCsch}[\operatorname{Sqrt}[x]]) / 3$

Rule 12

$\operatorname{Int}[(a_*)(u_), x_Symbol] := \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /;$ FreeQ[a, x] && !MatchQ[u, (b_*)(v_)] /; FreeQ[b, x]

Rule 45

$\operatorname{Int}[(a_.) + (b_.)*(x_.)^{(m_.)}*((c_.) + (d_.)*(x_.)^{(n_.)}), x_Symbol] := \operatorname{Int}[\operatorname{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /;$ FreeQ[{a, b, c, d, n},

`x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

Rule 6481

```
Int[((a_.) + ArcCsch[u_]*(b_.))*((c_.) + (d_.)*(x_.))^(m_.), x_Symbol] := Si
mp[(c + d*x)^(m + 1)*((a + b*ArcCsch[u])/(d*(m + 1))), x] - Dist[b*(u/(d*(m
+ 1)*Sqrt[-u^2])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqr
t[-1 - u^2])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && Inv
erseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !Func
tionOfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{3}x^3 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \frac{x^2}{2\sqrt{-1-x}} dx}{3\sqrt{-x}} \\
 &= \frac{1}{3}x^3 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \frac{x^2}{\sqrt{-1-x}} dx}{6\sqrt{-x}} \\
 &= \frac{1}{3}x^3 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \left(\frac{1}{\sqrt{-1-x}} + 2\sqrt{-1-x} + (-1-x)^{3/2} \right) dx}{6\sqrt{-x}} \\
 &= \frac{\sqrt{-1-x}\sqrt{x}}{3\sqrt{-x}} + \frac{2(-1-x)^{3/2}\sqrt{x}}{9\sqrt{-x}} + \frac{(-1-x)^{5/2}\sqrt{x}}{15\sqrt{-x}} + \frac{1}{3}x^3 \operatorname{csch}^{-1}(\sqrt{x})
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.47

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{45} \sqrt{1 + \frac{1}{x}} \sqrt{x} (8 - 4x + 3x^2) + \frac{1}{3} x^3 \operatorname{csch}^{-1}(\sqrt{x})$$

`[In] Integrate[x^2*ArcCsch[Sqrt[x]], x]`

`[Out] (Sqrt[1 + x^(-1)]*Sqrt[x]*(8 - 4*x + 3*x^2))/45 + (x^3*ArcCsch[Sqrt[x]])/3`

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.39

method	result	size
parts	$\frac{x^3 \operatorname{arccsch}(\sqrt{x})}{3} + \frac{\sqrt{\frac{1+x}{x}} \sqrt{x} (3x^2 - 4x + 8)}{45}$	35
derivativedivides	$\frac{x^3 \operatorname{arccsch}(\sqrt{x})}{3} + \frac{(1+x)(3x^2 - 4x + 8)}{45 \sqrt{\frac{1+x}{x}} \sqrt{x}}$	38
default	$\frac{x^3 \operatorname{arccsch}(\sqrt{x})}{3} + \frac{(1+x)(3x^2 - 4x + 8)}{45 \sqrt{\frac{1+x}{x}} \sqrt{x}}$	38

[In] `int(x^2*arccsch(x^(1/2)),x,method=_RETURNVERBOSE)`

[Out] `1/3*x^3*arccsch(x^(1/2))+1/45*((1+x)/x)^(1/2)*x^(1/2)*(3*x^2-4*x+8)`

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.56

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{3} x^3 \log \left(\frac{x \sqrt{\frac{x+1}{x}} + \sqrt{x}}{x} \right) + \frac{1}{45} (3x^2 - 4x + 8) \sqrt{x} \sqrt{\frac{x+1}{x}}$$

[In] `integrate(x^2*arccsch(x^(1/2)),x, algorithm="fricas")`

[Out] `1/3*x^3*log((x*sqrt((x + 1)/x) + sqrt(x))/x) + 1/45*(3*x^2 - 4*x + 8)*sqrt(x)*sqrt((x + 1)/x)`

Sympy [F]

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx = \int x^2 \operatorname{acsch}(\sqrt{x}) dx$$

[In] `integrate(x**2*acsch(x**(1/2)),x)`

[Out] `Integral(x**2*acsch(sqrt(x)), x)`

Maxima [A] (verification not implemented)

none

Time = 0.19 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.52

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{15} x^{\frac{5}{2}} \left(\frac{1}{x} + 1\right)^{\frac{5}{2}} + \frac{1}{3} x^3 \operatorname{arcsch}(\sqrt{x}) - \frac{2}{9} x^{\frac{3}{2}} \left(\frac{1}{x} + 1\right)^{\frac{3}{2}} + \frac{1}{3} \sqrt{x} \sqrt{\frac{1}{x} + 1}$$

[In] integrate(x^2*arccsch(x^(1/2)),x, algorithm="maxima")

[Out] 1/15*x^(5/2)*(1/x + 1)^(5/2) + 1/3*x^3*arccsch(sqrt(x)) - 2/9*x^(3/2)*(1/x + 1)^(3/2) + 1/3*sqrt(x)*sqrt(1/x + 1)

Giac [F]

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx = \int x^2 \operatorname{arcsch}(\sqrt{x}) dx$$

[In] integrate(x^2*arccsch(x^(1/2)),x, algorithm="giac")

[Out] integrate(x^2*arccsch(sqrt(x)), x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{csch}^{-1}(\sqrt{x}) dx = \int x^2 \operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right) dx$$

[In] int(x^2*asinh(1/x^(1/2)),x)

[Out] int(x^2*asinh(1/x^(1/2)), x)

3.16 $\int x \operatorname{csch}^{-1}(\sqrt{x}) dx$

Optimal result	161
Rubi [A] (verified)	161
Mathematica [A] (verified)	162
Maple [A] (verified)	163
Fricas [A] (verification not implemented)	163
Sympy [F]	163
Maxima [A] (verification not implemented)	164
Giac [F]	164
Mupad [F(-1)]	164

Optimal result

Integrand size = 8, antiderivative size = 64

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx = -\frac{\sqrt{-1-x}\sqrt{x}}{2\sqrt{-x}} - \frac{(-1-x)^{3/2}\sqrt{x}}{6\sqrt{-x}} + \frac{1}{2}x^2 \operatorname{csch}^{-1}(\sqrt{x})$$

[Out] $1/2*x^2*\operatorname{arccsch}(x^{(1/2)})-1/6*(-1-x)^{(3/2)}*x^{(1/2)}/(-x)^{(1/2)}-1/2*(-1-x)^{(1/2)}*x^{(1/2)}/(-x)^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6481, 12, 45}

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{2}x^2 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{(-x-1)^{3/2}\sqrt{x}}{6\sqrt{-x}} - \frac{\sqrt{-x-1}\sqrt{x}}{2\sqrt{-x}}$$

[In] `Int[x*ArcCsch[Sqrt[x]],x]`

[Out] $-1/2*(\operatorname{Sqrt}[-1-x]*\operatorname{Sqrt}[x])/\operatorname{Sqrt}[-x] - ((-1-x)^{(3/2)}*\operatorname{Sqrt}[x])/(6*\operatorname{Sqrt}[-x]) + (x^2*\operatorname{ArcCsch}[\operatorname{Sqrt}[x]])/2$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 45

`Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},`

`x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

Rule 6481

```
Int[((a_.) + ArcCsch[u_]*(b_.))*((c_.) + (d_.)*(x_.))^(m_.), x_Symbol] := Si
mp[(c + d*x)^(m + 1)*((a + b*ArcCsch[u])/(d*(m + 1))), x] - Dist[b*(u/(d*(m
+ 1)*Sqrt[-u^2])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqr
t[-1 - u^2])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && Inv
erseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !Func
tionOfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{2}x^2 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \frac{x}{2\sqrt{-1-x}} dx}{2\sqrt{-x}} \\
 &= \frac{1}{2}x^2 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \frac{x}{\sqrt{-1-x}} dx}{4\sqrt{-x}} \\
 &= \frac{1}{2}x^2 \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \left(-\frac{1}{\sqrt{-1-x}} - \sqrt{-1-x} \right) dx}{4\sqrt{-x}} \\
 &= -\frac{\sqrt{-1-x}\sqrt{x}}{2\sqrt{-x}} - \frac{(-1-x)^{3/2}\sqrt{x}}{6\sqrt{-x}} + \frac{1}{2}x^2 \operatorname{csch}^{-1}(\sqrt{x})
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.55

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{6} \sqrt{1 + \frac{1}{x}} (-2 + x) \sqrt{x} + \frac{1}{2} x^2 \operatorname{csch}^{-1}(\sqrt{x})$$

[In] Integrate[x*ArcCsch[Sqrt[x]],x]

[Out] (Sqrt[1 + x^(-1)]*(-2 + x)*Sqrt[x])/6 + (x^2*ArcCsch[Sqrt[x]])/2

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.44

method	result	size
parts	$\frac{x^2 \operatorname{arccsch}(\sqrt{x})}{2} + \frac{\sqrt{\frac{1+x}{x}} \sqrt{x} (x-2)}{6}$	28
derivativedivides	$\frac{x^2 \operatorname{arccsch}(\sqrt{x})}{2} + \frac{(1+x)(x-2)}{6\sqrt{\frac{1+x}{x}} \sqrt{x}}$	31
default	$\frac{x^2 \operatorname{arccsch}(\sqrt{x})}{2} + \frac{(1+x)(x-2)}{6\sqrt{\frac{1+x}{x}} \sqrt{x}}$	31

[In] `int(x*arccsch(x^(1/2)),x,method=_RETURNVERBOSE)`

[Out] $1/2*x^2*\operatorname{arccsch}(x^{(1/2)})+1/6*((1+x)/x)^{(1/2)}*x^{(1/2)}*(x-2)$

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.67

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{2} x^2 \log\left(\frac{x\sqrt{\frac{x+1}{x}} + \sqrt{x}}{x}\right) + \frac{1}{6} (x-2)\sqrt{x}\sqrt{\frac{x+1}{x}}$$

[In] `integrate(x*arccsch(x^(1/2)),x, algorithm="fricas")`

[Out] $1/2*x^2*\log((x*\sqrt{(x+1)/x} + \sqrt{x})/x) + 1/6*(x-2)*\sqrt{x}*\sqrt{(x+1)/x}$

Sympy [F]

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx = \int x \operatorname{acsch}(\sqrt{x}) dx$$

[In] `integrate(x*acsch(x**(1/2)),x)`

[Out] `Integral(x*acsch(sqrt(x)), x)`

Maxima [A] (verification not implemented)

none

Time = 0.23 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.53

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{1}{6} x^{\frac{3}{2}} \left(\frac{1}{x} + 1 \right)^{\frac{3}{2}} + \frac{1}{2} x^2 \operatorname{arcsch}(\sqrt{x}) - \frac{1}{2} \sqrt{x} \sqrt{\frac{1}{x} + 1}$$

[In] integrate(x*arccsch(x^(1/2)),x, algorithm="maxima")

[Out] 1/6*x^(3/2)*(1/x + 1)^(3/2) + 1/2*x^2*arccsch(sqrt(x)) - 1/2*sqrt(x)*sqrt(1/x + 1)

Giac [F]

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx = \int x \operatorname{arcsch}(\sqrt{x}) dx$$

[In] integrate(x*arccsch(x^(1/2)),x, algorithm="giac")

[Out] integrate(x*arccsch(sqrt(x)), x)

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{csch}^{-1}(\sqrt{x}) dx = \int x \operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right) dx$$

[In] int(x*asinh(1/x^(1/2)),x)

[Out] int(x*asinh(1/x^(1/2)), x)

3.17 $\int \operatorname{csch}^{-1}(\sqrt{x}) dx$

Optimal result	165
Rubi [A] (verified)	165
Mathematica [A] (verified)	166
Maple [A] (verified)	166
Fricas [A] (verification not implemented)	167
Sympy [F]	167
Maxima [A] (verification not implemented)	167
Giac [F]	167
Mupad [B] (verification not implemented)	168

Optimal result

Integrand size = 6, antiderivative size = 31

$$\int \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{\sqrt{-1-x}\sqrt{x}}{\sqrt{-x}} + x \operatorname{csch}^{-1}(\sqrt{x})$$

[Out] $x \operatorname{arccsch}(x^{1/2}) + (-1-x)^{1/2} x^{1/2} / (-x)^{1/2}$

Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6479, 12, 32}

$$\int \operatorname{csch}^{-1}(\sqrt{x}) dx = \frac{\sqrt{-x-1}\sqrt{x}}{\sqrt{-x}} + x \operatorname{csch}^{-1}(\sqrt{x})$$

[In] `Int[ArcCsch[Sqrt[x]], x]`

[Out] `(Sqrt[-1 - x]*Sqrt[x])/Sqrt[-x] + x*ArcCsch[Sqrt[x]]`

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]`

Rule 32

`Int[((a_.) + (b_.)*(x_))^(m_), x_Symbol] := Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]`

Rule 6479

```
Int[ArcCsch[u_], x_Symbol] := Simp[x*ArcCsch[u], x] - Dist[u/Sqrt[-u^2], Int[SimplifyIntegrand[x*(D[u, x]/(u*Sqrt[-1 - u^2])), x], x], x] /; InverseFunctionFreeQ[u, x] && !FunctionOfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= x \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \frac{1}{2\sqrt{-1-x}} dx}{\sqrt{-x}} \\ &= x \operatorname{csch}^{-1}(\sqrt{x}) - \frac{\sqrt{x} \int \frac{1}{\sqrt{-1-x}} dx}{2\sqrt{-x}} \\ &= \frac{\sqrt{-1-x}\sqrt{x}}{\sqrt{-x}} + x \operatorname{csch}^{-1}(\sqrt{x}) \end{aligned}$$

Mathematica [A] (verified)

Time = 2.31 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.77

$$\int \operatorname{csch}^{-1}(\sqrt{x}) dx = \sqrt{1 + \frac{1}{x}} \sqrt{x} + x \operatorname{csch}^{-1}(\sqrt{x})$$

```
[In] Integrate[ArcCsch[Sqrt[x]], x]
```

```
[Out] Sqrt[1 + x^(-1)]*Sqrt[x] + x*ArcCsch[Sqrt[x]]
```

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.68

method	result	size
parts	$x \operatorname{arccsch}(\sqrt{x}) + \sqrt{\frac{1+x}{x}} \sqrt{x}$	21
derivativedivides	$x \operatorname{arccsch}(\sqrt{x}) + \frac{1+x}{\sqrt{\frac{1+x}{x}} \sqrt{x}}$	24
default	$x \operatorname{arccsch}(\sqrt{x}) + \frac{1+x}{\sqrt{\frac{1+x}{x}} \sqrt{x}}$	24

```
[In] int(arccsch(x^(1/2)), x, method=_RETURNVERBOSE)
```

```
[Out] x*arccsch(x^(1/2))+((1+x)/x)^(1/2)*x^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.16

$$\int \operatorname{csch}^{-1}(\sqrt{x}) \, dx = x \log \left(\frac{x\sqrt{\frac{x+1}{x}} + \sqrt{x}}{x} \right) + \sqrt{x}\sqrt{\frac{x+1}{x}}$$

[In] integrate(arccsch(x^(1/2)),x, algorithm="fricas")

[Out] x*log((x*sqrt((x + 1)/x) + sqrt(x))/x) + sqrt(x)*sqrt((x + 1)/x)

Sympy [F]

$$\int \operatorname{csch}^{-1}(\sqrt{x}) \, dx = \int \operatorname{acsch}(\sqrt{x}) \, dx$$

[In] integrate(acsch(x**(1/2)),x)

[Out] Integral(acsch(sqrt(x)), x)

Maxima [A] (verification not implemented)

none

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.58

$$\int \operatorname{csch}^{-1}(\sqrt{x}) \, dx = x \operatorname{arcsch}(\sqrt{x}) + \sqrt{x}\sqrt{\frac{1}{x} + 1}$$

[In] integrate(arccsch(x^(1/2)),x, algorithm="maxima")

[Out] x*arccsch(sqrt(x)) + sqrt(x)*sqrt(1/x + 1)

Giac [F]

$$\int \operatorname{csch}^{-1}(\sqrt{x}) \, dx = \int \operatorname{arcsch}(\sqrt{x}) \, dx$$

[In] integrate(arccsch(x^(1/2)),x, algorithm="giac")

[Out] integrate(arccsch(sqrt(x)), x)

Mupad [B] (verification not implemented)

Time = 5.44 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.58

$$\int \operatorname{csch}^{-1}(\sqrt{x}) \, dx = x \operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right) + \sqrt{x} \sqrt{\frac{1}{x} + 1}$$

[In] `int(asinh(1/x^(1/2)),x)`

[Out] `x*asinh(1/x^(1/2)) + x^(1/2)*(1/x + 1)^(1/2)`

3.18 $\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx$

Optimal result	169
Rubi [A] (verified)	169
Mathematica [A] (verified)	171
Maple [F]	171
Fricas [F]	171
Sympy [F]	172
Maxima [F]	172
Giac [F]	172
Mupad [F(-1)]	172

Optimal result

Integrand size = 10, antiderivative size = 46

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = \operatorname{csch}^{-1}(\sqrt{x})^2 - 2\operatorname{csch}^{-1}(\sqrt{x}) \log\left(1 - e^{2\operatorname{csch}^{-1}(\sqrt{x})}\right) - \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(\sqrt{x})}\right)$$

[Out] $\operatorname{arccsch}(x^{(1/2)})^2 - 2*\operatorname{arccsch}(x^{(1/2)})*\ln(1 - (1/x^{(1/2)} + (1+1/x)^{(1/2)})^2) - \operatorname{polylog}(2, (1/x^{(1/2)} + (1+1/x)^{(1/2)})^2)$

Rubi [A] (verified)

Time = 0.07 (sec), antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6417, 5775, 3797, 2221, 2317, 2438}

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = -\operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(\sqrt{x})}\right) + \operatorname{csch}^{-1}(\sqrt{x})^2 - 2\operatorname{csch}^{-1}(\sqrt{x}) \log\left(1 - e^{2\operatorname{csch}^{-1}(\sqrt{x})}\right)$$

[In] $\operatorname{Int}[\operatorname{ArcCsch}[\operatorname{Sqrt}[x]]/x, x]$

[Out] $\operatorname{ArcCsch}[\operatorname{Sqrt}[x]]^2 - 2*\operatorname{ArcCsch}[\operatorname{Sqrt}[x]]*\operatorname{Log}[1 - E^{(2*\operatorname{ArcCsch}[\operatorname{Sqrt}[x]])}] - \operatorname{PolyLog}[2, E^{(2*\operatorname{ArcCsch}[\operatorname{Sqrt}[x]])}]$

Rule 2221

$\operatorname{Int}[(((F_.)^{((g_.) * ((e_.) + (f_.) * (x_)))})^{(n_.) * ((c_.) + (d_.) * (x_))^{(m_.)}}) / ((a_.) + (b_.) * (F_.)^{((g_.) * ((e_.) + (f_.) * (x_)))})^{(n_.)}], x_Symbol] \rightarrow \operatorname{Simp}$

```

[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

```

Rule 2317

```

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

```

Rule 2438

```

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

```

Rule 3797

```

Int[((c_) + (d_)*(x_)^(m_))*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]

```

Rule 5775

```

Int[((a_) + ArcSinh[(c_)*(x_)])*(b_)^(n_)/(x_), x_Symbol] :> Dist[1/b, Subst[Int[x^n*Coth[-a/b + x/b], x], x, a + b*ArcSinh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]

```

Rule 6417

```

Int[((a_) + ArcCsch[(c_)*(x_)])*(b_)^(n_)/(x_), x_Symbol] :> -Subst[Int[(a + b*ArcSinh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= 2\text{Subst}\left(\int \frac{\text{csch}^{-1}(x)}{x} dx, x, \sqrt{x}\right) \\
&= -\left(2\text{Subst}\left(\int \frac{\text{arcsinh}(x)}{x} dx, x, \frac{1}{\sqrt{x}}\right)\right) \\
&= -\left(2\text{Subst}\left(\int x \coth(x) dx, x, \text{arcsinh}\left(\frac{1}{\sqrt{x}}\right)\right)\right) \\
&= \text{arcsinh}\left(\frac{1}{\sqrt{x}}\right)^2 + 4\text{Subst}\left(\int \frac{e^{2x}}{1 - e^{2x}} dx, x, \text{arcsinh}\left(\frac{1}{\sqrt{x}}\right)\right)
\end{aligned}$$

$$\begin{aligned}
&= \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)^2 - 2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) \log\left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)}\right) \\
&\quad + 2\operatorname{Subst}\left(\int \log(1 - e^{2x}) dx, x, \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)\right) \\
&= \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)^2 - 2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) \log\left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)}\right) \\
&\quad + \operatorname{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)}\right) \\
&= \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)^2 - 2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right) \log\left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)}\right) - \operatorname{PolyLog}\left(2, e^{2\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = \operatorname{csch}^{-1}(\sqrt{x}) \left(\operatorname{csch}^{-1}(\sqrt{x}) - 2 \log\left(1 - e^{2\operatorname{csch}^{-1}(\sqrt{x})}\right) \right) - \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(\sqrt{x})}\right)$$

[In] Integrate[ArcCsch[Sqrt[x]]/x,x]

[Out] ArcCsch[Sqrt[x]]*(ArcCsch[Sqrt[x]] - 2*Log[1 - E^(2*ArcCsch[Sqrt[x]])]) - PolyLog[2, E^(2*ArcCsch[Sqrt[x]])]

Maple [F]

$$\int \frac{\operatorname{arccsch}(\sqrt{x})}{x} dx$$

[In] int(arccsch(x^(1/2))/x,x)

[Out] int(arccsch(x^(1/2))/x,x)

Fricas [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arcsch}(\sqrt{x})}{x} dx$$

[In] integrate(arccsch(x^(1/2))/x,x, algorithm="fricas")

[Out] integral(arccsch(sqrt(x))/x, x)

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{acsch}(\sqrt{x})}{x} dx$$

[In] integrate(acsch(x**(1/2))/x,x)

[Out] Integral(acsch(sqrt(x))/x, x)

Maxima [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arcsch}(\sqrt{x})}{x} dx$$

[In] integrate(arccsch(x^(1/2))/x,x, algorithm="maxima")

[Out] integrate(arccsch(sqrt(x))/x, x)

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arcsch}(\sqrt{x})}{x} dx$$

[In] integrate(arccsch(x^(1/2))/x,x, algorithm="giac")

[Out] integrate(arccsch(sqrt(x))/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right)}{x} dx$$

[In] int(asinh(1/x^(1/2))/x,x)

[Out] int(asinh(1/x^(1/2))/x, x)

3.19 $\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx$

Optimal result	173
Rubi [A] (verified)	173
Mathematica [A] (verified)	175
Maple [A] (verified)	175
Fricas [A] (verification not implemented)	175
Sympy [F]	176
Maxima [A] (verification not implemented)	176
Giac [F]	176
Mupad [B] (verification not implemented)	177

Optimal result

Integrand size = 10, antiderivative size = 63

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{-1-x}}{2\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} - \frac{\sqrt{x} \arctan(\sqrt{-1-x})}{2\sqrt{-x}}$$

[Out] $-\operatorname{arccsch}(x^{(1/2)})/x+1/2*(-1-x)^{(1/2)/(-x)^{(1/2)}/x^{(1/2)}-1/2*\arctan((-1-x)^{(1/2))*x^{(1/2)/(-x)^{(1/2)}}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6481, 12, 44, 65, 210}

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = -\frac{\sqrt{x} \arctan(\sqrt{-x-1})}{2\sqrt{-x}} + \frac{\sqrt{-x-1}}{2\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x}$$

[In] $\operatorname{Int}[\operatorname{ArcCsch}[\operatorname{Sqrt}[x]]/x^2, x]$

[Out] $\operatorname{Sqrt}[-1-x]/(2*\operatorname{Sqrt}[-x]*\operatorname{Sqrt}[x]) - \operatorname{ArcCsch}[\operatorname{Sqrt}[x]]/x - (\operatorname{Sqrt}[x]*\operatorname{ArcTan}[\operatorname{Sqrt}[-1-x]])/(2*\operatorname{Sqrt}[-x])$

Rule 12

$\operatorname{Int}[(a_*)*(u_), x_Symbol] \rightarrow \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \&\& \operatorname{!Match} Q[u, (b_)*(v_)] /; \operatorname{FreeQ}[b, x]$

Rule 44

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((
m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x]
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !Int
egerQ[n] && LtQ[n, 0]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 6481

```
Int[((a_.) + ArcCsch[u_] * (b_.)) * ((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Si
mp[(c + d*x)^(m + 1)*((a + b*ArcCsch[u])/(d*(m + 1))), x] - Dist[b*(u/(d*(m
+ 1)*Sqrt[-u^2])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqr
t[-1 - u^2])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && Inv
erseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !Func
tionOfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} + \frac{\sqrt{x} \int \frac{1}{2\sqrt{-1-xx^2}} dx}{\sqrt{-x}} \\
&= -\frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} + \frac{\sqrt{x} \int \frac{1}{\sqrt{-1-xx^2}} dx}{2\sqrt{-x}} \\
&= \frac{\sqrt{-1-x}}{2\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} - \frac{\sqrt{x} \int \frac{1}{\sqrt{-1-xx}} dx}{4\sqrt{-x}} \\
&= \frac{\sqrt{-1-x}}{2\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} + \frac{\sqrt{x} \operatorname{Subst}\left(\int \frac{1}{-1-x^2} dx, x, \sqrt{-1-x}\right)}{2\sqrt{-x}} \\
&= \frac{\sqrt{-1-x}}{2\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} - \frac{\sqrt{x} \arctan(\sqrt{-1-x})}{2\sqrt{-x}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.67

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{\frac{1+x}{x}}}{2\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x} - \frac{1}{2} \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)$$

`[In] Integrate[ArcCsch[Sqrt[x]]/x^2,x]``[Out] Sqrt[(1 + x)/x]/(2*Sqrt[x]) - ArcCsch[Sqrt[x]]/x - ArcSinh[1/Sqrt[x]]/2`**Maple [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.71

method	result	size
derivativedivides	$-\frac{\operatorname{arccsch}(\sqrt{x})}{x} + \frac{\sqrt{1+x} \left(-\operatorname{arctanh}\left(\frac{1}{\sqrt{1+x}}\right)x + \sqrt{1+x} \right)}{2\sqrt{\frac{1+x}{x}} x^{\frac{3}{2}}}$	45
default	$-\frac{\operatorname{arccsch}(\sqrt{x})}{x} + \frac{\sqrt{1+x} \left(-\operatorname{arctanh}\left(\frac{1}{\sqrt{1+x}}\right)x + \sqrt{1+x} \right)}{2\sqrt{\frac{1+x}{x}} x^{\frac{3}{2}}}$	45
parts	$-\frac{\operatorname{arccsch}(\sqrt{x})}{x} + \frac{\sqrt{\frac{1+x}{x}} \sqrt{x} (\ln(\sqrt{1+x}-1)x - \ln(\sqrt{1+x}+1)x + 2\sqrt{1+x})}{4\sqrt{1+x} (\sqrt{1+x}-1)(\sqrt{1+x}+1)}$	77

`[In] int(arccsch(x^(1/2))/x^2,x,method=_RETURNVERBOSE)``[Out] -arccsch(x^(1/2))/x+1/2*(1+x)^(1/2)*(-arctanh(1/(1+x)^(1/2))*x+(1+x)^(1/2))
/((1+x)/x)^(1/2)/x^(3/2)`**Fricas [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.70

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = -\frac{(x+2) \log\left(\frac{x\sqrt{\frac{x+1}{x}} + \sqrt{x}}{x}\right) - \sqrt{x}\sqrt{\frac{x+1}{x}}}{2x}$$

`[In] integrate(arccsch(x^(1/2))/x^2,x, algorithm="fricas")``[Out] -1/2*((x + 2)*log((x*sqrt((x + 1)/x) + sqrt(x))/x) - sqrt(x)*sqrt((x + 1)/x
))/x`

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = \int \frac{\operatorname{acsch}(\sqrt{x})}{x^2} dx$$

[In] integrate(acsch(x**(1/2))/x**2,x)

[Out] Integral(acsch(sqrt(x))/x**2, x)

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.03

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{x}\sqrt{\frac{1}{x}+1}}{2\left(x\left(\frac{1}{x}+1\right)-1\right)} - \frac{\operatorname{arcsch}(\sqrt{x})}{x} - \frac{1}{4} \log\left(\sqrt{x}\sqrt{\frac{1}{x}+1}+1\right) + \frac{1}{4} \log\left(\sqrt{x}\sqrt{\frac{1}{x}+1}-1\right)$$

[In] integrate(arccsch(x^(1/2))/x^2,x, algorithm="maxima")

[Out] 1/2*sqrt(x)*sqrt(1/x + 1)/(x*(1/x + 1) - 1) - arccsch(sqrt(x))/x - 1/4*log(sqrt(x)*sqrt(1/x + 1) + 1) + 1/4*log(sqrt(x)*sqrt(1/x + 1) - 1)

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = \int \frac{\operatorname{arcsch}(\sqrt{x})}{x^2} dx$$

[In] integrate(arccsch(x^(1/2))/x^2,x, algorithm="giac")

[Out] integrate(arccsch(sqrt(x))/x^2, x)

Mupad [B] (verification not implemented)

Time = 4.84 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.52

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{\frac{1}{x} + 1}}{2\sqrt{x}} - \frac{2 \operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right) \left(\frac{1}{2\sqrt{x}} + \frac{\sqrt{x}}{4}\right)}{\sqrt{x}}$$

[In] `int(asinh(1/x^(1/2))/x^2,x)`

[Out] `(1/x + 1)^(1/2)/(2*x^(1/2)) - (2*asinh(1/x^(1/2))*(1/(2*x^(1/2)) + x^(1/2)/4))/x^(1/2)`

3.20 $\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx$

Optimal result	178
Rubi [A] (verified)	178
Mathematica [A] (verified)	180
Maple [A] (verified)	180
Fricas [A] (verification not implemented)	180
Sympy [F]	181
Maxima [A] (verification not implemented)	181
Giac [F]	181
Mupad [F(-1)]	182

Optimal result

Integrand size = 10, antiderivative size = 90

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = \frac{\sqrt{-1-x}}{8\sqrt{-x}x^{3/2}} - \frac{3\sqrt{-1-x}}{16\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{2x^2} + \frac{3\sqrt{x} \arctan(\sqrt{-1-x})}{16\sqrt{-x}}$$

[Out] $-1/2*\operatorname{arccsch}(x^{(1/2)})/x^2+1/8*(-1-x)^{(1/2)}/x^{(3/2)}/(-x)^{(1/2)}-3/16*(-1-x)^{(1/2)}/(-x)^{(1/2)}/x^{(1/2)}+3/16*\arctan((-1-x)^{(1/2)})*x^{(1/2)}/(-x)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6481, 12, 44, 65, 210}

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = \frac{3\sqrt{x} \arctan(\sqrt{-x-1})}{16\sqrt{-x}} + \frac{\sqrt{-x-1}}{8\sqrt{-x}x^{3/2}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{2x^2} - \frac{3\sqrt{-x-1}}{16\sqrt{-x}\sqrt{x}}$$

[In] $\operatorname{Int}[\operatorname{ArcCsch}[\operatorname{Sqrt}[x]]/x^3, x]$

[Out] $\operatorname{Sqrt}[-1-x]/(8*\operatorname{Sqrt}[-x]*x^{(3/2)}) - (3*\operatorname{Sqrt}[-1-x])/(16*\operatorname{Sqrt}[-x]*\operatorname{Sqrt}[x]) - \operatorname{ArcCsch}[\operatorname{Sqrt}[x]]/(2*x^2) + (3*\operatorname{Sqrt}[x]*\operatorname{ArcTan}[\operatorname{Sqrt}[-1-x]])/(16*\operatorname{Sqrt}[-x])$

Rule 12

$\operatorname{Int}[(a_*)*(u_), x_Symbol] \rightarrow \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{Match} Q[u, (b_*)*(v_)] /; \operatorname{FreeQ}[b, x]$

Rule 44

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((
m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x]
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !Int
egerQ[n] && LtQ[n, 0]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 6481

```
Int[((a_.) + ArcCsch[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Si
mp[(c + d*x)^(m + 1)*((a + b*ArcCsch[u])/(d*(m + 1))), x] - Dist[b*(u/(d*(m
+ 1)*Sqrt[-u^2])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqr
t[-1 - u^2])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && Inv
erseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !Func
tionOfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\operatorname{csch}^{-1}(\sqrt{x})}{2x^2} + \frac{\sqrt{x} \int \frac{1}{2\sqrt{-1-xx^3}} dx}{2\sqrt{-x}} \\
&= -\frac{\operatorname{csch}^{-1}(\sqrt{x})}{2x^2} + \frac{\sqrt{x} \int \frac{1}{\sqrt{-1-xx^3}} dx}{4\sqrt{-x}} \\
&= \frac{\sqrt{-1-x}}{8\sqrt{-xx^{3/2}}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{2x^2} - \frac{(3\sqrt{x}) \int \frac{1}{\sqrt{-1-xx^2}} dx}{16\sqrt{-x}} \\
&= \frac{\sqrt{-1-x}}{8\sqrt{-xx^{3/2}}} - \frac{3\sqrt{-1-x}}{16\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{2x^2} + \frac{(3\sqrt{x}) \int \frac{1}{\sqrt{-1-xx}} dx}{32\sqrt{-x}} \\
&= \frac{\sqrt{-1-x}}{8\sqrt{-xx^{3/2}}} - \frac{3\sqrt{-1-x}}{16\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{2x^2} - \frac{(3\sqrt{x}) \operatorname{Subst}\left(\int \frac{1}{-1-x^2} dx, x, \sqrt{-1-x}\right)}{16\sqrt{-x}}
\end{aligned}$$

$$= \frac{\sqrt{-1-x}}{8\sqrt{-x}x^{3/2}} - \frac{3\sqrt{-1-x}}{16\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{2x^2} + \frac{3\sqrt{x} \arctan(\sqrt{-1-x})}{16\sqrt{-x}}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.52

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = \frac{\sqrt{1+\frac{1}{x}}(2-3x)\sqrt{x} - 8\operatorname{csch}^{-1}(\sqrt{x}) + 3x^2 \operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)}{16x^2}$$

[In] Integrate[ArcCsch[Sqrt[x]]/x^3,x]

[Out] (Sqrt[1 + x^(-1)]*(2 - 3*x)*Sqrt[x] - 8*ArcCsch[Sqrt[x]] + 3*x^2*ArcSinh[1/Sqrt[x]])/(16*x^2)

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.63

method	result	size
derivativedivides	$-\frac{\operatorname{arccsch}(\sqrt{x})}{2x^2} + \frac{\sqrt{1+x} \left(3 \operatorname{arctanh}\left(\frac{1}{\sqrt{1+x}}\right)x^2 - 3\sqrt{1+x}x + 2\sqrt{1+x} \right)}{16\sqrt{\frac{1+x}{x}}x^{\frac{5}{2}}}$	57
default	$-\frac{\operatorname{arccsch}(\sqrt{x})}{2x^2} + \frac{\sqrt{1+x} \left(3 \operatorname{arctanh}\left(\frac{1}{\sqrt{1+x}}\right)x^2 - 3\sqrt{1+x}x + 2\sqrt{1+x} \right)}{16\sqrt{\frac{1+x}{x}}x^{\frac{5}{2}}}$	57
parts	$-\frac{\operatorname{arccsch}(\sqrt{x})}{2x^2} + \frac{\sqrt{\frac{1+x}{x}}\sqrt{x} \left(3 \ln(\sqrt{1+x}+1)x^2 - 3 \ln(\sqrt{1+x}-1)x^2 - 6\sqrt{1+x}x + 4\sqrt{1+x} \right)}{32\sqrt{1+x}(\sqrt{1+x}+1)^2(\sqrt{1+x}-1)^2}$	90

[In] int(arccsch(x^(1/2))/x^3,x,method=_RETURNVERBOSE)

[Out] -1/2*arccsch(x^(1/2))/x^2+1/16*(1+x)^(1/2)*(3*arctanh(1/(1+x)^(1/2))*x^2-3*(1+x)^(1/2)*x+2*(1+x)^(1/2))/((1+x)/x)^(1/2)/x^(5/2)

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.59

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = -\frac{(3x-2)\sqrt{x}\sqrt{\frac{x+1}{x}} - (3x^2-8) \log\left(\frac{x\sqrt{\frac{x+1}{x}}+\sqrt{x}}{x}\right)}{16x^2}$$

[In] integrate(arccsch(x^(1/2))/x^3,x, algorithm="fricas")

[Out] $-1/16*((3*x - 2)*\sqrt{x}*\sqrt{(x + 1)/x} - (3*x^2 - 8)*\log((x*\sqrt{(x + 1)/x} + \sqrt{x})/x))/x^2$

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\operatorname{acsch}(\sqrt{x})}{x^3} dx$$

[In] `integrate(acsch(x**(1/2))/x**3,x)`

[Out] `Integral(acsch(sqrt(x))/x**3, x)`

Maxima [A] (verification not implemented)

none

Time = 0.19 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.02

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = -\frac{3x^{\frac{3}{2}}\left(\frac{1}{x} + 1\right)^{\frac{3}{2}} - 5\sqrt{x}\sqrt{\frac{1}{x} + 1}}{16\left(x^2\left(\frac{1}{x} + 1\right)^2 - 2x\left(\frac{1}{x} + 1\right) + 1\right)} - \frac{\operatorname{arcsch}(\sqrt{x})}{2x^2} + \frac{3}{32} \log\left(\sqrt{x}\sqrt{\frac{1}{x} + 1} + 1\right) - \frac{3}{32} \log\left(\sqrt{x}\sqrt{\frac{1}{x} + 1} - 1\right)$$

[In] `integrate(arccsch(x^(1/2))/x^3,x, algorithm="maxima")`

[Out] $-1/16*(3*x^(3/2)*(1/x + 1)^(3/2) - 5*\sqrt{x}*\sqrt{1/x + 1})/(x^2*(1/x + 1)^2 - 2*x*(1/x + 1) + 1) - 1/2*arccsch(sqrt(x))/x^2 + 3/32*\log(sqrt(x)*sqrt(1/x + 1) + 1) - 3/32*\log(sqrt(x)*sqrt(1/x + 1) - 1)$

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\operatorname{arcsch}(\sqrt{x})}{x^3} dx$$

[In] `integrate(arccsch(x^(1/2))/x^3,x, algorithm="giac")`

[Out] `integrate(arccsch(sqrt(x))/x^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right)}{x^3} dx$$

```
[In] int(asinh(1/x^(1/2))/x^3,x)
```

```
[Out] int(asinh(1/x^(1/2))/x^3, x)
```

3.21 $\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx$

Optimal result	183
Rubi [A] (verified)	183
Mathematica [A] (verified)	185
Maple [A] (verified)	185
Fricas [A] (verification not implemented)	186
Sympy [F]	186
Maxima [A] (verification not implemented)	186
Giac [F]	187
Mupad [F(-1)]	187

Optimal result

Integrand size = 10, antiderivative size = 115

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = \frac{\sqrt{-1-x}}{18\sqrt{-x}x^{5/2}} - \frac{5\sqrt{-1-x}}{72\sqrt{-x}x^{3/2}} + \frac{5\sqrt{-1-x}}{48\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3} - \frac{5\sqrt{x} \arctan(\sqrt{-1-x})}{48\sqrt{-x}}$$

[Out] $-1/3*\operatorname{arccsch}(x^{(1/2)})/x^3+1/18*(-1-x)^{(1/2)}/x^{(5/2)}/(-x)^{(1/2)}-5/72*(-1-x)^{(1/2)}/x^{(3/2)}/(-x)^{(1/2)}+5/48*(-1-x)^{(1/2)}/(-x)^{(1/2)}/x^{(1/2)}-5/48*\arctan((-1-x)^{(1/2))*x^{(1/2)}/(-x)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6481, 12, 44, 65, 210}

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = -\frac{5\sqrt{x} \arctan(\sqrt{-x-1})}{48\sqrt{-x}} - \frac{5\sqrt{-x-1}}{72\sqrt{-x}x^{3/2}} + \frac{\sqrt{-x-1}}{18\sqrt{-x}x^{5/2}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3} + \frac{5\sqrt{-x-1}}{48\sqrt{-x}\sqrt{x}}$$

[In] $\operatorname{Int}[\operatorname{ArcCsch}[\operatorname{Sqrt}[x]]/x^4, x]$

[Out] $\operatorname{Sqrt}[-1-x]/(18*\operatorname{Sqrt}[-x]*x^{(5/2)}) - (5*\operatorname{Sqrt}[-1-x])/(72*\operatorname{Sqrt}[-x]*x^{(3/2)}) + (5*\operatorname{Sqrt}[-1-x])/(48*\operatorname{Sqrt}[-x]*\operatorname{Sqrt}[x]) - \operatorname{ArcCsch}[\operatorname{Sqrt}[x]]/(3*x^3) - (5*\operatorname{Sqrt}[-1-x])/(48*\operatorname{Sqrt}[-x]*\operatorname{Sqrt}[x])$

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 44

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((
m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !Int
egerQ[n] && LtQ[n, 0]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 6481

```
Int[((a_.) + ArcCsch[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Si
mp[(c + d*x)^(m + 1)*((a + b*ArcCsch[u])/(d*(m + 1))), x] - Dist[b*(u/(d*(m
+ 1)*Sqrt[-u^2])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqr
t[-1 - u^2])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && Inv
erseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !Func
tionOfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3} + \frac{\sqrt{x} \int \frac{1}{2\sqrt{-1-xx^4}} dx}{3\sqrt{-x}} \\
&= -\frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3} + \frac{\sqrt{x} \int \frac{1}{\sqrt{-1-xx^4}} dx}{6\sqrt{-x}} \\
&= \frac{\sqrt{-1-x}}{18\sqrt{-xx^{5/2}}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3} - \frac{(5\sqrt{x}) \int \frac{1}{\sqrt{-1-xx^3}} dx}{36\sqrt{-x}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\sqrt{-1-x}}{18\sqrt{-xx^{5/2}}} - \frac{5\sqrt{-1-x}}{72\sqrt{-xx^{3/2}}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3} + \frac{(5\sqrt{x}) \int \frac{1}{\sqrt{-1-xx^2}} dx}{48\sqrt{-x}} \\
&= \frac{\sqrt{-1-x}}{18\sqrt{-xx^{5/2}}} - \frac{5\sqrt{-1-x}}{72\sqrt{-xx^{3/2}}} + \frac{5\sqrt{-1-x}}{48\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3} - \frac{(5\sqrt{x}) \int \frac{1}{\sqrt{-1-xx}} dx}{96\sqrt{-x}} \\
&= \frac{\sqrt{-1-x}}{18\sqrt{-xx^{5/2}}} - \frac{5\sqrt{-1-x}}{72\sqrt{-xx^{3/2}}} + \frac{5\sqrt{-1-x}}{48\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3} \\
&\quad + \frac{(5\sqrt{x}) \operatorname{Subst}\left(\int \frac{1}{-1-x^2} dx, x, \sqrt{-1-x}\right)}{48\sqrt{-x}} \\
&= \frac{\sqrt{-1-x}}{18\sqrt{-xx^{5/2}}} - \frac{5\sqrt{-1-x}}{72\sqrt{-xx^{3/2}}} + \frac{5\sqrt{-1-x}}{48\sqrt{-x}\sqrt{x}} - \frac{\operatorname{csch}^{-1}(\sqrt{x})}{3x^3} - \frac{5\sqrt{x} \arctan(\sqrt{-1-x})}{48\sqrt{-x}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.45

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = \frac{\sqrt{1+\frac{1}{x}}\sqrt{x}(8-10x+15x^2) - 48\operatorname{csch}^{-1}(\sqrt{x}) - 15x^3\operatorname{arcsinh}\left(\frac{1}{\sqrt{x}}\right)}{144x^3}$$

[In] Integrate[ArcCsch[Sqrt[x]]/x^4,x]

[Out] (Sqrt[1+x^(-1)]*Sqrt[x]*(8-10*x+15*x^2)-48*ArcCsch[Sqrt[x]]-15*x^3*ArcSinh[1/Sqrt[x]])/(144*x^3)

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.58

method	result	size
derivativedivides	$-\frac{\operatorname{arccsch}(\sqrt{x})}{3x^3} + \frac{\sqrt{1+x}(-15\operatorname{arctanh}\left(\frac{1}{\sqrt{1+x}}\right)x^3+15\sqrt{1+x}x^2-10\sqrt{1+x}x+8\sqrt{1+x})}{144\sqrt{\frac{1+x}{x}}x^{\frac{7}{2}}}$	67
default	$-\frac{\operatorname{arccsch}(\sqrt{x})}{3x^3} + \frac{\sqrt{1+x}(-15\operatorname{arctanh}\left(\frac{1}{\sqrt{1+x}}\right)x^3+15\sqrt{1+x}x^2-10\sqrt{1+x}x+8\sqrt{1+x})}{144\sqrt{\frac{1+x}{x}}x^{\frac{7}{2}}}$	67
parts	$-\frac{\operatorname{arccsch}(\sqrt{x})}{3x^3} + \frac{\sqrt{\frac{1+x}{x}}\sqrt{x}(15\ln(\sqrt{1+x}-1)x^3-15\ln(\sqrt{1+x}+1)x^3+30\sqrt{1+x}x^2-20\sqrt{1+x}x+16\sqrt{1+x})}{288\sqrt{1+x}(\sqrt{1+x}-1)^3(\sqrt{1+x}+1)^3}$	100

[In] int(arccsch(x^(1/2))/x^4,x,method=_RETURNVERBOSE)

[Out] -1/3*arccsch(x^(1/2))/x^3+1/144*(1+x)^(1/2)*(-15*arctanh(1/(1+x)^(1/2))*x^3+15*(1+x)^(1/2)*x^2-10*(1+x)^(1/2)*x+8*(1+x)^(1/2))/((1+x)/x)^(1/2)/x^(7/2)

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.50

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = \frac{(15x^2 - 10x + 8)\sqrt{x}\sqrt{\frac{x+1}{x}} - 3(5x^3 + 16)\log\left(\frac{x\sqrt{\frac{x+1}{x}} + \sqrt{x}}{x}\right)}{144x^3}$$

[In] integrate(arccsch(x^(1/2))/x^4,x, algorithm="fricas")

[Out] 1/144*((15*x^2 - 10*x + 8)*sqrt(x)*sqrt((x + 1)/x) - 3*(5*x^3 + 16)*log((x*sqrt((x + 1)/x) + sqrt(x))/x))/x^3

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\operatorname{acsch}(\sqrt{x})}{x^4} dx$$

[In] integrate(acsch(x**(1/2))/x**4,x)

[Out] Integral(acsch(sqrt(x))/x**4, x)

Maxima [A] (verification not implemented)

none

Time = 0.21 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.01

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = \frac{15x^{\frac{5}{2}}\left(\frac{1}{x} + 1\right)^{\frac{5}{2}} - 40x^{\frac{3}{2}}\left(\frac{1}{x} + 1\right)^{\frac{3}{2}} + 33\sqrt{x}\sqrt{\frac{1}{x} + 1}}{144\left(x^3\left(\frac{1}{x} + 1\right)^3 - 3x^2\left(\frac{1}{x} + 1\right)^2 + 3x\left(\frac{1}{x} + 1\right) - 1\right)} - \frac{\operatorname{arcsch}(\sqrt{x})}{3x^3} - \frac{5}{96}\log\left(\sqrt{x}\sqrt{\frac{1}{x} + 1} + 1\right) + \frac{5}{96}\log\left(\sqrt{x}\sqrt{\frac{1}{x} + 1} - 1\right)$$

[In] integrate(arccsch(x^(1/2))/x^4,x, algorithm="maxima")

[Out] 1/144*(15*x^(5/2)*(1/x + 1)^(5/2) - 40*x^(3/2)*(1/x + 1)^(3/2) + 33*sqrt(x)*sqrt(1/x + 1))/(x^3*(1/x + 1)^3 - 3*x^2*(1/x + 1)^2 + 3*x*(1/x + 1) - 1) - 1/3*arccsch(sqrt(x))/x^3 - 5/96*log(sqrt(x)*sqrt(1/x + 1) + 1) + 5/96*log(sqrt(x)*sqrt(1/x + 1) - 1)

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\operatorname{arcsch}(\sqrt{x})}{x^4} dx$$

[In] integrate(arccsch(x^(1/2))/x^4,x, algorithm="giac")

[Out] integrate(arccsch(sqrt(x))/x^4, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{\sqrt{x}}\right)}{x^4} dx$$

[In] int(asinh(1/x^(1/2))/x^4,x)

[Out] int(asinh(1/x^(1/2))/x^4, x)

3.22 $\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx$

Optimal result	188
Rubi [A] (verified)	188
Mathematica [A] (verified)	189
Maple [A] (verified)	189
Fricas [A] (verification not implemented)	190
Sympy [A] (verification not implemented)	190
Maxima [A] (verification not implemented)	190
Giac [F]	190
Mupad [B] (verification not implemented)	191

Optimal result

Integrand size = 4, antiderivative size = 16

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = -\sqrt{1+x^2} + x \operatorname{arcsinh}(x)$$

[Out] $x*\operatorname{arcsinh}(x)-(x^2+1)^{(1/2)}$

Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6463, 5772, 267}

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = x \operatorname{arcsinh}(x) - \sqrt{x^2 + 1}$$

[In] $\operatorname{Int}[\operatorname{ArcCsch}[x^{-1}], x]$

[Out] $-\operatorname{Sqrt}[1 + x^2] + x*\operatorname{ArcSinh}[x]$

Rule 267

$\operatorname{Int}[(x_)^{(m_.)}*((a_) + (b_.)*(x_)^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x^n)^{(p+1)}/(b*n*(p+1)), x] /;$ $\operatorname{FreeQ}\{a, b, m, n, p\}, x$ && $\operatorname{EqQ}[m, n-1]$ && $\operatorname{NeQ}[p, -1]$

Rule 5772

$\operatorname{Int}[(a_. + \operatorname{ArcSinh}[c_.*(x_)])*(b_.)^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[x*(a + b*\operatorname{ArcSinh}[c*x])^n, x] - \operatorname{Dist}[b*c*n, \operatorname{Int}[x*((a + b*\operatorname{ArcSinh}[c*x])^{(n-1)})/\operatorname{Sqrt}[1 + c^2*x^2]], x, x] /;$ $\operatorname{FreeQ}\{a, b, c\}, x$ && $\operatorname{GtQ}[n, 0]$

Rule 6463

```
Int[ArcCsch[(c_.)/((a_.) + (b_.)*(x_)^(n_.))]^(m_.)*(u_.), x_Symbol] := Int
[u*ArcSinh[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \operatorname{arcsinh}(x) dx \\ &= x \operatorname{arcsinh}(x) - \int \frac{x}{\sqrt{1+x^2}} dx \\ &= -\sqrt{1+x^2} + x \operatorname{arcsinh}(x) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = -\sqrt{1+x^2} + x \operatorname{csch}^{-1}\left(\frac{1}{x}\right)$$

```
[In] Integrate[ArcCsch[x^(-1)],x]
```

```
[Out] -Sqrt[1 + x^2] + x*ArcCsch[x^(-1)]
```

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

method	result	size
parts	$x \operatorname{arccsch}\left(\frac{1}{x}\right) - \sqrt{x^2 + 1}$	17
derivativedivides	$x \operatorname{arccsch}\left(\frac{1}{x}\right) - \frac{x^2\left(\frac{1}{x^2}+1\right)}{\sqrt{\left(\frac{1}{x^2}+1\right)x^2}}$	29
default	$x \operatorname{arccsch}\left(\frac{1}{x}\right) - \frac{x^2\left(\frac{1}{x^2}+1\right)}{\sqrt{\left(\frac{1}{x^2}+1\right)x^2}}$	29

```
[In] int(arccsch(1/x),x,method=_RETURNVERBOSE)
```

```
[Out] x*arccsch(1/x)-(x^2+1)^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = x \log\left(x + \sqrt{x^2 + 1}\right) - \sqrt{x^2 + 1}$$

[In] integrate(arccsch(1/x),x, algorithm="fricas")

[Out] x*log(x + sqrt(x^2 + 1)) - sqrt(x^2 + 1)

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = x \operatorname{acsch}\left(\frac{1}{x}\right) - \sqrt{x^2 + 1}$$

[In] integrate(acsch(1/x),x)

[Out] x*acsch(1/x) - sqrt(x**2 + 1)

Maxima [A] (verification not implemented)

none

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = x \operatorname{arcsch}\left(\frac{1}{x}\right) - \sqrt{x^2 + 1}$$

[In] integrate(arccsch(1/x),x, algorithm="maxima")

[Out] x*arccsch(1/x) - sqrt(x^2 + 1)

Giac [F]

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = \int \operatorname{arcsch}\left(\frac{1}{x}\right) dx$$

[In] integrate(arccsch(1/x),x, algorithm="giac")

[Out] integrate(arccsch(1/x), x)

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \operatorname{csch}^{-1}\left(\frac{1}{x}\right) dx = x \operatorname{asinh}(x) - \sqrt{x^2 + 1}$$

[In] int(asinh(x),x)

[Out] x*asinh(x) - (x^2 + 1)^(1/2)

3.23 $\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx$

Optimal result	192
Rubi [A] (verified)	192
Mathematica [C] (verified)	194
Maple [F]	195
Fricas [F(-2)]	195
Sympy [F]	195
Maxima [F]	195
Giac [F]	196
Mupad [F(-1)]	196

Optimal result

Integrand size = 10, antiderivative size = 61

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = \frac{\operatorname{csch}^{-1}(ax^n)^2}{2n} - \frac{\operatorname{csch}^{-1}(ax^n) \log\left(1 - e^{2\operatorname{csch}^{-1}(ax^n)}\right)}{n} - \frac{\operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(ax^n)}\right)}{2n}$$

[Out] 1/2*arccsch(a*x^n)^2/n-arccsch(a*x^n)*ln(1-(1/a/(x^n)+(1+1/a^2/(x^n)^2)^(1/2))^2)/n-1/2*polylog(2,(1/a/(x^n)+(1+1/a^2/(x^n)^2)^(1/2))^2)/n

Rubi [A] (verified)

Time = 0.08 (sec), antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6417, 5775, 3797, 2221, 2317, 2438}

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = -\frac{\operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(ax^n)}\right)}{2n} + \frac{\operatorname{csch}^{-1}(ax^n)^2}{2n} - \frac{\operatorname{csch}^{-1}(ax^n) \log\left(1 - e^{2\operatorname{csch}^{-1}(ax^n)}\right)}{n}$$

[In] Int[ArcCsch[a*x^n]/x,x]

[Out] ArcCsch[a*x^n]^2/(2*n) - (ArcCsch[a*x^n]*Log[1 - E^(2*ArcCsch[a*x^n])])/n - PolyLog[2, E^(2*ArcCsch[a*x^n])]/(2*n)

Rule 2221


```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)
))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol]
:= Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x))
)^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3797

```
Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_
)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist
[2*I, Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)
)/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && Int
egerQ[4*k] && IGtQ[m, 0]
```

Rule 5775

```
Int[((a_) + ArcSinh[(c_)*(x_)]*(b_))^(n_)/(x_), x_Symbol] := Dist[1/b,
Subst[Int[x^n*Coth[-a/b + x/b], x], x, a + b*ArcSinh[c*x]], x] /; FreeQ[{a,
b, c}, x] && IGtQ[n, 0]
```

Rule 6417

```
Int[((a_) + ArcCsch[(c_)*(x_)]*(b_))/(x_), x_Symbol] := -Subst[Int[(a +
b*ArcSinh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\text{Subst}\left(\int \frac{\text{csch}^{-1}(ax)}{x} dx, x, x^n\right)}{n} \\ &= -\frac{\text{Subst}\left(\int \frac{\text{arcsinh}\left(\frac{x}{a}\right)}{x} dx, x, x^{-n}\right)}{n} \end{aligned}$$

$$\begin{aligned}
&= -\frac{\text{Subst}\left(\int x \coth(x) dx, x, \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)\right)}{n} \\
&= \frac{\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)^2}{2n} + \frac{2\text{Subst}\left(\int \frac{e^{2x}x}{1-e^{2x}} dx, x, \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)\right)}{n} \\
&= \frac{\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)^2}{2n} - \frac{\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right) \log\left(1 - e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)}\right)}{n} \\
&\quad + \frac{\text{Subst}\left(\int \log(1 - e^{2x}) dx, x, \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)\right)}{n} \\
&= \frac{\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)^2}{2n} - \frac{\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right) \log\left(1 - e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)}\right)}{n} \\
&\quad + \frac{\text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)}\right)}{2n} \\
&= \frac{\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)^2}{2n} - \frac{\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right) \log\left(1 - e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)}\right)}{n} - \frac{\operatorname{PolyLog}\left(2, e^{2\operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)}\right)}{2n}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.06 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.05

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = -\frac{x^{-n} {}_3F_2\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{3}{2}, -\frac{x^{-2n}}{a^2}\right)}{an} + \left(\operatorname{csch}^{-1}(ax^n) - \operatorname{arcsinh}\left(\frac{x^{-n}}{a}\right)\right) \log(x)$$

[In] Integrate[ArcCsch[a*x^n]/x,x]

[Out] -(HypergeometricPFQ[{1/2, 1/2, 1/2}, {3/2, 3/2}, -(1/(a^2*x^(2*n)))]/(a*n*x^n)) + (ArcCsch[a*x^n] - ArcSinh[1/(a*x^n)])*Log[x]

Maple [F]

$$\int \frac{\operatorname{arccsch}(ax^n)}{x} dx$$

[In] `int(arccsch(a*x^n)/x,x)`

[Out] `int(arccsch(a*x^n)/x,x)`

Fricas [F(-2)]

Exception generated.

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = \text{Exception raised: TypeError}$$

[In] `integrate(arccsch(a*x^n)/x,x, algorithm="fricas")`

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{acsch}(ax^n)}{x} dx$$

[In] `integrate(acsch(a*x**n)/x,x)`

[Out] `Integral(acsch(a*x**n)/x, x)`

Maxima [F]

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{arcsch}(ax^n)}{x} dx$$

[In] `integrate(arccsch(a*x^n)/x,x, algorithm="maxima")`

[Out] `a^2*n*integrate(x^(2*n)*log(x)/(a^2*x*x^(2*n) + (a^2*x*x^(2*n) + x)*sqrt(a^2*x^(2*n) + 1) + x), x) + n*integrate(log(x)/(a^2*x*x^(2*n) + x), x) - log(a)*log(x) - log(x)*log(x^n) + log(x)*log(sqrt(a^2*x^(2*n) + 1) + 1)`

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{arcsch}(ax^n)}{x} dx$$

[In] integrate(arccsch(a*x^n)/x,x, algorithm="giac")

[Out] integrate(arccsch(a*x^n)/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{ax^n}\right)}{x} dx$$

[In] int(asinh(1/(a*x^n))/x,x)

[Out] int(asinh(1/(a*x^n))/x, x)

3.24 $\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx$

Optimal result	197
Rubi [A] (verified)	197
Mathematica [A] (verified)	199
Maple [F]	199
Fricas [F]	199
Sympy [F]	200
Maxima [F]	200
Giac [F]	200
Mupad [F(-1)]	200

Optimal result

Integrand size = 10, antiderivative size = 54

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = \frac{1}{10} \operatorname{csch}^{-1}(ax^5)^2 - \frac{1}{5} \operatorname{csch}^{-1}(ax^5) \log\left(1 - e^{2\operatorname{csch}^{-1}(ax^5)}\right) - \frac{1}{10} \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(ax^5)}\right)$$

[Out] 1/10*arccsch(a*x^5)^2-1/5*arccsch(a*x^5)*ln(1-(1/a/x^5+(1+1/a^2/x^10)^(1/2))^2)-1/10*polylog(2,(1/a/x^5+(1+1/a^2/x^10)^(1/2))^2)

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6417, 5775, 3797, 2221, 2317, 2438}

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = -\frac{1}{10} \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(ax^5)}\right) + \frac{1}{10} \operatorname{csch}^{-1}(ax^5)^2 - \frac{1}{5} \operatorname{csch}^{-1}(ax^5) \log\left(1 - e^{2\operatorname{csch}^{-1}(ax^5)}\right)$$

[In] Int[ArcCsch[a*x^5]/x,x]

[Out] ArcCsch[a*x^5]^2/10 - (ArcCsch[a*x^5]*Log[1 - E^(2*ArcCsch[a*x^5])])/5 - PolyLog[2, E^(2*ArcCsch[a*x^5])]/10

Rule 2221

Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] :> Simp

```

[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

```

Rule 2317

```

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

```

Rule 2438

```

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

```

Rule 3797

```

Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] :> Simp[(-1)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[((c + d*x)^m*(E^(2*((-1)*e + f*fz*x)))/(1 + E^(2*((-1)*e + f*fz*x)))/E^(2*I*k*Pi))]/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]

```

Rule 5775

```

Int[((a_) + ArcSinh[(c_)*(x_)])*(b_)^(n_)/(x_), x_Symbol] :> Dist[1/b, Subst[Int[x^n*Coth[-a/b + x/b], x], x, a + b*ArcSinh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]

```

Rule 6417

```

Int[((a_) + ArcCsch[(c_)*(x_)])*(b_)/(x_), x_Symbol] :> -Subst[Int[(a + b*ArcSinh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{5} \text{Subst} \left(\int \frac{\text{csch}^{-1}(ax)}{x} dx, x, x^5 \right) \\
&= - \left(\frac{1}{5} \text{Subst} \left(\int \frac{\text{arcsinh}\left(\frac{x}{a}\right)}{x} dx, x, \frac{1}{x^5} \right) \right) \\
&= - \left(\frac{1}{5} \text{Subst} \left(\int x \coth(x) dx, x, \text{csch}^{-1}(ax^5) \right) \right) \\
&= \frac{1}{10} \text{csch}^{-1}(ax^5)^2 + \frac{2}{5} \text{Subst} \left(\int \frac{e^{2x}x}{1 - e^{2x}} dx, x, \text{csch}^{-1}(ax^5) \right)
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{10} \operatorname{csch}^{-1}(ax^5)^2 - \frac{1}{5} \operatorname{csch}^{-1}(ax^5) \log\left(1 - e^{2\operatorname{csch}^{-1}(ax^5)}\right) \\
&\quad + \frac{1}{5} \operatorname{Subst}\left(\int \log(1 - e^{2x}) dx, x, \operatorname{csch}^{-1}(ax^5)\right) \\
&= \frac{1}{10} \operatorname{csch}^{-1}(ax^5)^2 - \frac{1}{5} \operatorname{csch}^{-1}(ax^5) \log\left(1 - e^{2\operatorname{csch}^{-1}(ax^5)}\right) \\
&\quad + \frac{1}{10} \operatorname{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2\operatorname{csch}^{-1}(ax^5)}\right) \\
&= \frac{1}{10} \operatorname{csch}^{-1}(ax^5)^2 - \frac{1}{5} \operatorname{csch}^{-1}(ax^5) \log\left(1 - e^{2\operatorname{csch}^{-1}(ax^5)}\right) - \frac{1}{10} \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(ax^5)}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.93

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = \frac{1}{10} \left(\operatorname{csch}^{-1}(ax^5)^2 - 2\operatorname{csch}^{-1}(ax^5) \log\left(1 - e^{2\operatorname{csch}^{-1}(ax^5)}\right) - \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(ax^5)}\right) \right)$$

[In] Integrate[ArcCsch[a*x^5]/x,x]

[Out] (ArcCsch[a*x^5]^2 - 2*ArcCsch[a*x^5]*Log[1 - E^(2*ArcCsch[a*x^5])] - PolyLog[2, E^(2*ArcCsch[a*x^5])])/10

Maple [F]

$$\int \frac{\operatorname{arccsch}(ax^5)}{x} dx$$

[In] int(arccsch(a*x^5)/x,x)

[Out] int(arccsch(a*x^5)/x,x)

Fricas [F]

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arcsch}(ax^5)}{x} dx$$

[In] integrate(arccsch(a*x^5)/x,x, algorithm="fricas")

[Out] integral(arccsch(a*x^5)/x, x)

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{acsch}(ax^5)}{x} dx$$

```
[In] integrate(acsch(a*x**5)/x,x)
```

```
[Out] Integral(acsch(a*x**5)/x, x)
```

Maxima [F]

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arcsch}(ax^5)}{x} dx$$

```
[In] integrate(arccsch(a*x^5)/x,x, algorithm="maxima")
```

```
[Out] 5*a^2*integrate(x^9*log(x)/(a^2*x^10 + (a^2*x^10 + 1)^(3/2) + 1), x) - 1/2*
log(a^2*x^10 + 1)*log(x) - log(a)*log(x) - 5/2*log(x)^2 + log(x)*log(sqrt(a
^2*x^10 + 1) + 1) - 1/20*dilog(-a^2*x^10)
```

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arcsch}(ax^5)}{x} dx$$

```
[In] integrate(arccsch(a*x^5)/x,x, algorithm="giac")
```

```
[Out] integrate(arccsch(a*x^5)/x, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{ax^5}\right)}{x} dx$$

```
[In] int(asinh(1/(a*x^5))/x,x)
```

```
[Out] int(asinh(1/(a*x^5))/x, x)
```


3.25 $\int \operatorname{csch}^{-1}(ce^{a+bx}) dx$

Optimal result	201
Rubi [A] (verified)	201
Mathematica [B] (verified)	203
Maple [F]	204
Fricas [F(-2)]	204
Sympy [F]	204
Maxima [F]	205
Giac [F]	205
Mupad [F(-1)]	205

Optimal result

Integrand size = 10, antiderivative size = 77

$$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx = \frac{\operatorname{csch}^{-1}(ce^{a+bx})^2}{2b} - \frac{\operatorname{csch}^{-1}(ce^{a+bx}) \log(1 - e^{2\operatorname{csch}^{-1}(ce^{a+bx})})}{b} - \frac{\operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(ce^{a+bx})}\right)}{2b}$$

[Out] 1/2*arccsch(c*exp(b*x+a))^2/b-arccsch(c*exp(b*x+a))*ln(1-(1/c/exp(b*x+a)+(1+1/c^2/exp(b*x+a)^2)^(1/2))^2)/b-1/2*polylog(2,(1/c/exp(b*x+a)+(1+1/c^2/exp(b*x+a)^2)^(1/2))^2)/b

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {2320, 6417, 5775, 3797, 2221, 2317, 2438}

$$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx = -\frac{\operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(ce^{a+bx})}\right)}{2b} + \frac{\operatorname{csch}^{-1}(ce^{a+bx})^2}{2b} - \frac{\operatorname{csch}^{-1}(ce^{a+bx}) \log(1 - e^{2\operatorname{csch}^{-1}(ce^{a+bx})})}{b}$$

[In] Int[ArcCsch[c*E^(a + b*x)],x]

[Out] ArcCsch[c*E^(a + b*x)]^2/(2*b) - (ArcCsch[c*E^(a + b*x)]*Log[1 - E^(2*ArcCsch[c*E^(a + b*x)])])/b - PolyLog[2, E^(2*ArcCsch[c*E^(a + b*x)])]/(2*b)

Rule 2221

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)
))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol]
:= Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*
(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3797

```
Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_
)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist
[2*I, Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)
))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && Int
egerQ[4*k] && IGtQ[m, 0]
```

Rule 5775

```
Int[((a_) + ArcSinh[(c_)*(x_)]*(b_))^(n_)/(x_), x_Symbol] := Dist[1/b,
Subst[Int[x^n*Coth[-a/b + x/b], x], x, a + b*ArcSinh[c*x]], x] /; FreeQ[{a,
b, c}, x] && IGtQ[n, 0]
```

Rule 6417

```
Int[((a_) + ArcCsch[(c_)*(x_)]*(b_))/(x_), x_Symbol] := -Subst[Int[(a +
b*ArcSinh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Subst}\left(\int \frac{\text{csch}^{-1}(cx)}{x} dx, x, e^{a+bx}\right)}{b} \\
&= -\frac{\text{Subst}\left(\int \frac{\text{arcsinh}\left(\frac{x}{c}\right)}{x} dx, x, e^{-a-bx}\right)}{b} \\
&= -\frac{\text{Subst}\left(\int x \coth(x) dx, x, \text{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)\right)}{b} \\
&= \frac{\text{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} + \frac{2\text{Subst}\left(\int \frac{e^{2x}x}{1-e^{2x}} dx, x, \text{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)\right)}{b} \\
&= \frac{\text{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} - \frac{\text{arcsinh}\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 - e^{2\text{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}\right)}{b} \\
&\quad + \frac{\text{Subst}\left(\int \log(1 - e^{2x}) dx, x, \text{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)\right)}{b} \\
&= \frac{\text{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} - \frac{\text{arcsinh}\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 - e^{2\text{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}\right)}{b} \\
&\quad + \frac{\text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2\text{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}\right)}{2b} \\
&= \frac{\text{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} - \frac{\text{arcsinh}\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 - e^{2\text{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}\right)}{b} \\
&\quad - \frac{\text{PolyLog}\left(2, e^{2\text{arcsinh}\left(\frac{e^{-a-bx}}{c}\right)}\right)}{2b}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 236 vs. 2(77) = 154.

Time = 0.71 (sec) , antiderivative size = 236, normalized size of antiderivative = 3.06

$$\begin{aligned}
\int \text{csch}^{-1}(ce^{a+bx}) dx &= x \text{csch}^{-1}(ce^{a+bx}) \\
&\quad + \frac{e^{-a-bx} \sqrt{1 + c^2 e^{2(a+bx)}} \left(\log^2(-c^2 e^{2(a+bx)}) + \text{arctanh}\left(\sqrt{1 + c^2 e^{2(a+bx)}}\right) (-8bx + 4 \log(-c^2 e^{2(a+bx)})) \right)}{2}
\end{aligned}$$

```
[In] Integrate[ArcCsch[c*E^(a + b*x)],x]
```

```
[Out] x*ArcCsch[c*E^(a + b*x)] + (E^(-a - b*x)*Sqrt[1 + c^2*E^(2*(a + b*x))]*(Log
[-(c^2*E^(2*(a + b*x)))]^2 + ArcTanh[Sqrt[1 + c^2*E^(2*(a + b*x))]]*(-8*b*x
+ 4*Log[-(c^2*E^(2*(a + b*x)))])) - 4*Log[-(c^2*E^(2*(a + b*x)))]*Log[(1 +
Sqrt[1 + c^2*E^(2*(a + b*x))])/2] + 2*Log[(1 + Sqrt[1 + c^2*E^(2*(a + b*x))
])/2]^2 - 4*PolyLog[2, (1 - Sqrt[1 + c^2*E^(2*(a + b*x))])/2]))/(8*b*c*Sqrt
[1 + 1/(c^2*E^(2*(a + b*x)))]])
```

Maple [F]

$$\int \operatorname{arccsch}(e^{bx+a}c) dx$$

```
[In] int(arccsch(exp(b*x+a)*c),x)
```

```
[Out] int(arccsch(exp(b*x+a)*c),x)
```

Fricas [F(-2)]

Exception generated.

$$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx = \text{Exception raised: TypeError}$$

```
[In] integrate(arccsch(c*exp(b*x+a)),x, algorithm="fricas")
```

```
[Out] Exception raised: TypeError >> Error detected within library code: integ
rate: implementation incomplete (constant residues)
```

Sympy [F]

$$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx = \int \operatorname{acsch}(ce^{a+bx}) dx$$

```
[In] integrate(acsch(c*exp(b*x+a)),x)
```

```
[Out] Integral(acsch(c*exp(a + b*x)), x)
```

Maxima [F]

$$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx = \int \operatorname{arcsch}(ce^{(bx+a)}) dx$$

[In] integrate(arccsch(c*exp(b*x+a)),x, algorithm="maxima")

[Out] b*c^2*integrate(x*e^(2*b*x + 2*a)/(c^2*e^(2*b*x + 2*a) + (c^2*e^(2*b*x + 2*a) + 1)^(3/2) + 1), x) - 1/2*b*x^2 - (a + log(c))*x + x*log(sqrt(c^2*e^(2*b*x + 2*a) + 1) + 1) - 1/4*(2*b*x*log(c^2*e^(2*b*x + 2*a) + 1) + dilog(-c^2*e^(2*b*x + 2*a)))/b

Giac [F]

$$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx = \int \operatorname{arcsch}(ce^{(bx+a)}) dx$$

[In] integrate(arccsch(c*exp(b*x+a)),x, algorithm="giac")

[Out] integrate(arccsch(c*e^(b*x + a)), x)

Mupad [F(-1)]

Timed out.

$$\int \operatorname{csch}^{-1}(ce^{a+bx}) dx = \int \operatorname{asinh}\left(\frac{e^{-a-bx}}{c}\right) dx$$

[In] int(asinh(exp(- a - b*x)/c),x)

[Out] int(asinh(exp(- a - b*x)/c), x)

3.26 $\int e^{\operatorname{csch}^{-1}(ax)} x^m dx$

Optimal result	206
Rubi [A] (verified)	206
Mathematica [A] (verified)	207
Maple [F]	208
Fricas [F]	208
Sympy [A] (verification not implemented)	208
Maxima [F]	208
Giac [F(-2)]	209
Mupad [F(-1)]	209

Optimal result

Integrand size = 10, antiderivative size = 52

$$\int e^{\operatorname{csch}^{-1}(ax)} x^m dx = \frac{x^m}{am} + \frac{x^{1+m} \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, \frac{1}{2}(-1-m), \frac{1-m}{2}, -\frac{1}{a^2 x^2}\right)}{1+m}$$

[Out] $x^m/a/m+x^{(1+m)}*\operatorname{hypergeom}([-1/2, -1/2-1/2*m], [1/2-1/2*m], -1/a^2/x^2)/(1+m)$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {6471, 30, 346, 371}

$$\int e^{\operatorname{csch}^{-1}(ax)} x^m dx = \frac{x^{m+1} \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, \frac{1}{2}(-m-1), \frac{1-m}{2}, -\frac{1}{a^2 x^2}\right)}{m+1} + \frac{x^m}{am}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcCsch}[a*x]}*x^m, x]$

[Out] $x^m/(a*m) + (x^{(1+m)}*\operatorname{Hypergeometric2F1}[-1/2, (-1-m)/2, (1-m)/2, -(1/(a^2*x^2))])/(1+m)$

Rule 30

$\operatorname{Int}[(x_)^{(m_.)}, x_Symbol] := \operatorname{Simp}[x^{(m+1)}/(m+1), x] /; \operatorname{FreeQ}[m, x] \ \&\& \operatorname{NeQ}[m, -1]$

Rule 346

$\operatorname{Int}[((c_.)*(x_))^{(m_.)}*((a_.) + (b_.)*(x_)^{(n_.)})^{(p_.)}, x_Symbol] := \operatorname{Dist}[(-c^{(-1)})*(c*x)^{(m+1)}*(1/x)^{(m+1)}, \operatorname{Subst}[\operatorname{Int}[(a + b/x^n)^p/x^{(m+2)}, x], x$

, 1/x], x] /; FreeQ[{a, b, c, m, p}, x] && ILtQ[n, 0] && !RationalQ[m]

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 6471

Int[E^ArcCsch[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Dist[1/a, Int[x^(m - p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int x^{-1+m} dx}{a} + \int \sqrt{1 + \frac{1}{a^2 x^2}} x^m dx \\ &= \frac{x^m}{am} - \left(\left(\frac{1}{x} \right)^m x^m \right) \text{Subst} \left(\int x^{-2-m} \sqrt{1 + \frac{x^2}{a^2}} dx, x, \frac{1}{x} \right) \\ &= \frac{x^m}{am} + \frac{x^{1+m} \text{Hypergeometric2F1} \left(-\frac{1}{2}, \frac{1}{2}(-1 - m), \frac{1-m}{2}, -\frac{1}{a^2 x^2} \right)}{1 + m} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.04

$$\begin{aligned} &\int e^{\text{csch}^{-1}(ax)} x^m dx \\ &= \frac{x^m}{am} + \frac{x^{1+m} \text{Hypergeometric2F1} \left(-\frac{1}{2}, \frac{1}{2}(-1 - m), 1 + \frac{1}{2}(-1 - m), -\frac{1}{a^2 x^2} \right)}{1 + m} \end{aligned}$$

[In] Integrate[E^ArcCsch[a*x]*x^m,x]

[Out] x^m/(a*m) + (x^(1 + m)*Hypergeometric2F1[-1/2, (-1 - m)/2, 1 + (-1 - m)/2, -(1/(a^2*x^2))])/(1 + m)

Maple [F]

$$\int \left(\frac{1}{ax} + \sqrt{1 + \frac{1}{a^2x^2}} \right) x^m dx$$

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))*x^m,x)

[Out] int((1/a/x+(1+1/a^2/x^2)^(1/2))*x^m,x)

Fricas [F]

$$\int e^{\operatorname{csch}^{-1}(ax)} x^m dx = \int x^m \left(\sqrt{\frac{1}{a^2x^2} + 1} + \frac{1}{ax} \right) dx$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))*x^m,x, algorithm="fricas")

[Out] integral((a*x*x^m*sqrt((a^2*x^2 + 1)/(a^2*x^2)) + x^m)/(a*x), x)

Sympy [A] (verification not implemented)

Time = 2.80 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.25

$$\int e^{\operatorname{csch}^{-1}(ax)} x^m dx = -\frac{a^m a^{-m-1} x^m \Gamma\left(-\frac{m}{2}\right) {}_2F_1\left(\frac{-\frac{1}{2}, \frac{m}{2}}{\frac{m}{2} + 1} \middle| a^2 x^2 e^{i\pi}\right)}{2\Gamma\left(1 - \frac{m}{2}\right)} - \frac{\begin{cases} -\log(x) & \text{for } m = 0 \\ -\frac{x^m}{m} & \text{otherwise} \end{cases}}{a}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))*x**m,x)

[Out] -a**m*a**(-m - 1)*x**m*gamma(-m/2)*hyper((-1/2, m/2), (m/2 + 1,), a**2*x**2*exp_polar(I*pi))/(2*gamma(1 - m/2)) - Piecewise((-log(x), Eq(m, 0)), (-x**m/m, True))/a

Maxima [F]

$$\int e^{\operatorname{csch}^{-1}(ax)} x^m dx = \int x^m \left(\sqrt{\frac{1}{a^2x^2} + 1} + \frac{1}{ax} \right) dx$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))*x^m,x, algorithm="maxima")

[Out] integrate(sqrt(a^2*x^2 + 1)*x^m/x, x)/a + x^m/(a*m)

Giac [F(-2)]

Exception generated.

$$\int e^{\operatorname{csch}^{-1}(ax)} x^m dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))*x^m,x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int e^{\operatorname{csch}^{-1}(ax)} x^m dx = \int x^m \left(\sqrt{\frac{1}{a^2 x^2} + 1} + \frac{1}{ax} \right) dx$$

[In] int(x^m*((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x)),x)

[Out] int(x^m*((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x)), x)

3.27 $\int e^{\operatorname{csch}^{-1}(ax)} x^4 dx$

Optimal result	210
Rubi [A] (verified)	210
Mathematica [A] (verified)	211
Maple [A] (verified)	212
Fricas [A] (verification not implemented)	212
Sympy [A] (verification not implemented)	212
Maxima [A] (verification not implemented)	213
Giac [A] (verification not implemented)	213
Mupad [B] (verification not implemented)	213

Optimal result

Integrand size = 10, antiderivative size = 54

$$\int e^{\operatorname{csch}^{-1}(ax)} x^4 dx = -\frac{2\left(1 + \frac{1}{a^2 x^2}\right)^{3/2} x^3}{15a^2} + \frac{x^4}{4a} + \frac{1}{5} \left(1 + \frac{1}{a^2 x^2}\right)^{3/2} x^5$$

[Out] $-2/15*(1+1/a^2/x^2)^{(3/2)}*x^3/a^2+1/4*x^4/a+1/5*(1+1/a^2/x^2)^{(3/2)}*x^5$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {6471, 30, 277, 270}

$$\int e^{\operatorname{csch}^{-1}(ax)} x^4 dx = \frac{1}{5} x^5 \left(\frac{1}{a^2 x^2} + 1 \right)^{3/2} - \frac{2x^3 \left(\frac{1}{a^2 x^2} + 1 \right)^{3/2}}{15a^2} + \frac{x^4}{4a}$$

[In] `Int[E^ArcCsch[a*x]*x^4,x]`

[Out] $(-2*(1 + 1/(a^2*x^2))^{(3/2)}*x^3)/(15*a^2) + x^4/(4*a) + ((1 + 1/(a^2*x^2))^{(3/2)}*x^5)/5$

Rule 30

`Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]`

Rule 270

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, n,`

p}, x] && EqQ[(m + 1)/n + p + 1, 0] && NeQ[m, -1]

Rule 277

Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x^(m + 1)*((a + b*x^n)^(p + 1)/(a*(m + 1))), x] - Dist[b*((m + n*(p + 1) + 1)/(a*(m + 1))), Int[x^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, m, n, p}, x] && ILtQ[Simplify[(m + 1)/n + p + 1], 0] && NeQ[m, -1]

Rule 6471

Int[E^ArcCsch[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Dist[1/a, Int[x^(m - p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int x^3 dx}{a} + \int \sqrt{1 + \frac{1}{a^2 x^2}} x^4 dx \\ &= \frac{x^4}{4a} + \frac{1}{5} \left(1 + \frac{1}{a^2 x^2}\right)^{3/2} x^5 - \frac{2 \int \sqrt{1 + \frac{1}{a^2 x^2}} x^2 dx}{5a^2} \\ &= -\frac{2 \left(1 + \frac{1}{a^2 x^2}\right)^{3/2} x^3}{15a^2} + \frac{x^4}{4a} + \frac{1}{5} \left(1 + \frac{1}{a^2 x^2}\right)^{3/2} x^5 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.91

$$\int e^{\text{csch}^{-1}(ax)} x^4 dx = \frac{x^4}{4a} + \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x (-2 + a^2 x^2 + 3a^4 x^4)}{15a^4}$$

[In] Integrate[E^ArcCsch[a*x]*x^4, x]

[Out] x^4/(4*a) + (Sqrt[1 + 1/(a^2*x^2)]*x*(-2 + a^2*x^2 + 3*a^4*x^4))/(15*a^4)

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.98

method	result	size
default	$\frac{\sqrt{\frac{a^2x^2+1}{a^2x^2}} x(a^2x^2+1)(3a^2x^2-2)}{15a^4} + \frac{x^4}{4a}$	53
trager	$\frac{(x^3+x^2+x+1)(x-1)}{4} + \frac{(3a^4x^4+a^2x^2-2)x\sqrt{-\frac{a^2x^2-1}{a^2x^2}}}{15a^3}$	63

[In] `int((1/a/x+(1+1/a^2/x^2)^(1/2))*x^4,x,method=_RETURNVERBOSE)`

[Out] `1/15*((a^2*x^2+1)/a^2/x^2)^(1/2)*x*(a^2*x^2+1)/a^4*(3*a^2*x^2-2)+1/4*x^4/a`

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.98

$$\int e^{\operatorname{csch}^{-1}(ax)} x^4 dx = \frac{15 a^3 x^4 + 4 (3 a^4 x^5 + a^2 x^3 - 2 x) \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}}}{60 a^4}$$

[In] `integrate((1/a/x+(1+1/a^2/x^2)^(1/2))*x^4,x, algorithm="fricas")`

[Out] `1/60*(15*a^3*x^4 + 4*(3*a^4*x^5 + a^2*x^3 - 2*x)*sqrt((a^2*x^2 + 1)/(a^2*x^2)))/a^4`

Sympy [A] (verification not implemented)

Time = 1.28 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.17

$$\int e^{\operatorname{csch}^{-1}(ax)} x^4 dx = \frac{x^4 \sqrt{a^2 x^2 + 1}}{5a} + \frac{x^4}{4a} + \frac{x^2 \sqrt{a^2 x^2 + 1}}{15a^3} - \frac{2\sqrt{a^2 x^2 + 1}}{15a^5}$$

[In] `integrate((1/a/x+(1+1/a**2/x**2)**(1/2))*x**4,x)`

[Out] `x**4*sqrt(a**2*x**2 + 1)/(5*a) + x**4/(4*a) + x**2*sqrt(a**2*x**2 + 1)/(15*a**3) - 2*sqrt(a**2*x**2 + 1)/(15*a**5)`

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.93

$$\int e^{\operatorname{csch}^{-1}(ax)} x^4 dx = \frac{x^4}{4a} + \frac{3a^2 x^5 \left(\frac{1}{a^2 x^2} + 1\right)^{\frac{5}{2}} - 5x^3 \left(\frac{1}{a^2 x^2} + 1\right)^{\frac{3}{2}}}{15a^2}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))*x^4,x, algorithm="maxima")

[Out] 1/4*x^4/a + 1/15*(3*a^2*x^5*(1/(a^2*x^2) + 1)^(5/2) - 5*x^3*(1/(a^2*x^2) + 1)^(3/2))/a^2

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.44

$$\int e^{\operatorname{csch}^{-1}(ax)} x^4 dx = -\frac{a^2 x^2 + 1}{2a^5} + \frac{2|a|\operatorname{sgn}(x)}{15a^6} + \frac{12(a^2 x^2 + 1)^{\frac{5}{2}}|a|\operatorname{sgn}(x) - 20(a^2 x^2 + 1)^{\frac{3}{2}}|a|\operatorname{sgn}(x) + 15(a^2 x^2 + 1)^2 a}{60a^6}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))*x^4,x, algorithm="giac")

[Out] -1/2*(a^2*x^2 + 1)/a^5 + 2/15*abs(a)*sgn(x)/a^6 + 1/60*(12*(a^2*x^2 + 1)^(5/2)*abs(a)*sgn(x) - 20*(a^2*x^2 + 1)^(3/2)*abs(a)*sgn(x) + 15*(a^2*x^2 + 1)^2*a)/a^6

Mupad [B] (verification not implemented)

Time = 4.86 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.76

$$\int e^{\operatorname{csch}^{-1}(ax)} x^4 dx = \sqrt{\frac{1}{a^2 x^2} + 1} \left(\frac{x^5}{5} - \frac{2x}{15a^4} + \frac{x^3}{15a^2} \right) + \frac{x^4}{4a}$$

[In] int(x^4*((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x)),x)

[Out] (1/(a^2*x^2) + 1)^(1/2)*(x^5/5 - (2*x)/(15*a^4) + x^3/(15*a^2)) + x^4/(4*a)

3.28 $\int e^{\operatorname{csch}^{-1}(ax)} x^3 dx$

Optimal result	214
Rubi [A] (verified)	214
Mathematica [A] (verified)	216
Maple [A] (verified)	216
Fricas [A] (verification not implemented)	217
Sympy [A] (verification not implemented)	217
Maxima [A] (verification not implemented)	217
Giac [A] (verification not implemented)	218
Mupad [B] (verification not implemented)	218

Optimal result

Integrand size = 10, antiderivative size = 75

$$\int e^{\operatorname{csch}^{-1}(ax)} x^3 dx = \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x^2}{8a^2} + \frac{x^3}{3a} + \frac{1}{4} \sqrt{1 + \frac{1}{a^2 x^2}} x^4 - \frac{\operatorname{arctanh}\left(\sqrt{1 + \frac{1}{a^2 x^2}}\right)}{8a^4}$$

[Out] 1/3*x^3/a-1/8*arctanh((1+1/a^2/x^2)^(1/2))/a^4+1/8*x^2*(1+1/a^2/x^2)^(1/2)/a^2+1/4*x^4*(1+1/a^2/x^2)^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {6471, 30, 272, 43, 44, 65, 214}

$$\int e^{\operatorname{csch}^{-1}(ax)} x^3 dx = \frac{x^2 \sqrt{\frac{1}{a^2 x^2} + 1}}{8a^2} + \frac{1}{4} x^4 \sqrt{\frac{1}{a^2 x^2} + 1} - \frac{\operatorname{arctanh}\left(\sqrt{\frac{1}{a^2 x^2} + 1}\right)}{8a^4} + \frac{x^3}{3a}$$

[In] Int[E^ArcCsch[a*x]*x^3,x]

[Out] (Sqrt[1 + 1/(a^2*x^2)]*x^2)/(8*a^2) + x^3/(3*a) + (Sqrt[1 + 1/(a^2*x^2)]*x^4)/4 - ArcTanh[Sqrt[1 + 1/(a^2*x^2)]]/(8*a^4)

Rule 30

Int[(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 43

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Dist[d*(n/(b*(m + 1))), I
nt[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !IntegerQ[n] && GtQ[n, 0]
```

Rule 44

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((
m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x]
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !Int
egerQ[n] && LtQ[n, 0]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 6471

```
Int[E^ArcCsch[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Dist[1/a, Int[x^(m
- p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x
]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int x^2 dx}{a} + \int \sqrt{1 + \frac{1}{a^2 x^2}} x^3 dx \\ &= \frac{x^3}{3a} - \frac{1}{2} \text{Subst} \left(\int \frac{\sqrt{1 + \frac{x}{a^2}}}{x^3} dx, x, \frac{1}{x^2} \right) \end{aligned}$$

$$\begin{aligned}
&= \frac{x^3}{3a} + \frac{1}{4} \sqrt{1 + \frac{1}{a^2 x^2}} x^4 - \frac{\text{Subst}\left(\int \frac{1}{x^2 \sqrt{1 + \frac{x}{a^2}}} dx, x, \frac{1}{x^2}\right)}{8a^2} \\
&= \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x^2}{8a^2} + \frac{x^3}{3a} + \frac{1}{4} \sqrt{1 + \frac{1}{a^2 x^2}} x^4 + \frac{\text{Subst}\left(\int \frac{1}{x \sqrt{1 + \frac{x}{a^2}}} dx, x, \frac{1}{x^2}\right)}{16a^4} \\
&= \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x^2}{8a^2} + \frac{x^3}{3a} + \frac{1}{4} \sqrt{1 + \frac{1}{a^2 x^2}} x^4 + \frac{\text{Subst}\left(\int \frac{1}{-a^2 + a^2 x^2} dx, x, \sqrt{1 + \frac{1}{a^2 x^2}}\right)}{8a^2} \\
&= \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x^2}{8a^2} + \frac{x^3}{3a} + \frac{1}{4} \sqrt{1 + \frac{1}{a^2 x^2}} x^4 - \frac{\text{arctanh}\left(\sqrt{1 + \frac{1}{a^2 x^2}}\right)}{8a^4}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.01

$$\begin{aligned}
&\int e^{\text{csch}^{-1}(ax)} x^3 dx \\
&= \frac{a^2 x^2 \left(3 \sqrt{1 + \frac{1}{a^2 x^2}} + 8ax + 6a^2 \sqrt{1 + \frac{1}{a^2 x^2}} x^2 \right) - 3 \log \left(\left(1 + \sqrt{1 + \frac{1}{a^2 x^2}} \right) x \right)}{24a^4}
\end{aligned}$$

[In] Integrate[E^ArcCsch[a*x]*x^3,x]

[Out] (a^2*x^2*(3*sqrt[1 + 1/(a^2*x^2)] + 8*a*x + 6*a^2*sqrt[1 + 1/(a^2*x^2)]*x^2) - 3*Log[(1 + sqrt[1 + 1/(a^2*x^2)])*x])/(24*a^4)

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.49

method	result	size
default	$\frac{\sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} x \left(2x \left(\frac{a^2 x^2 + 1}{a^2} \right)^{\frac{3}{2}} a^4 - x \sqrt{\frac{a^2 x^2 + 1}{a^2}} a^2 - \ln \left(x + \sqrt{\frac{a^2 x^2 + 1}{a^2}} \right) \right)}{8 \sqrt{\frac{a^2 x^2 + 1}{a^2}} a^4} + \frac{x^3}{3a}$	112

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))*x^3,x,method=_RETURNVERBOSE)

[Out] 1/8*((a^2*x^2+1)/a^2/x^2)^(1/2)*x*(2*x*((a^2*x^2+1)/a^2)^(3/2)*a^4-x*((a^2*x^2+1)/a^2)^(1/2)*a^2-ln(x+((a^2*x^2+1)/a^2)^(1/2)))/((a^2*x^2+1)/a^2)^(1/2)/a^4+1/3*x^3/a

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.05

$$\int e^{\operatorname{csch}^{-1}(ax)} x^3 dx = \frac{8a^3x^3 + 3(2a^4x^4 + a^2x^2)\sqrt{\frac{a^2x^2+1}{a^2x^2}} + 3 \log\left(ax\sqrt{\frac{a^2x^2+1}{a^2x^2}} - ax\right)}{24a^4}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))*x^3,x, algorithm="fricas")

[Out] 1/24*(8*a^3*x^3 + 3*(2*a^4*x^4 + a^2*x^2)*sqrt((a^2*x^2 + 1)/(a^2*x^2)) + 3*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x))/a^4

Sympy [A] (verification not implemented)

Time = 2.59 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.97

$$\int e^{\operatorname{csch}^{-1}(ax)} x^3 dx = \frac{ax^5}{4\sqrt{a^2x^2+1}} + \frac{x^3}{3a} + \frac{3x^3}{8a\sqrt{a^2x^2+1}} + \frac{x}{8a^3\sqrt{a^2x^2+1}} - \frac{\operatorname{asinh}(ax)}{8a^4}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))*x**3,x)

[Out] a*x**5/(4*sqrt(a**2*x**2 + 1)) + x**3/(3*a) + 3*x**3/(8*a*sqrt(a**2*x**2 + 1)) + x/(8*a**3*sqrt(a**2*x**2 + 1)) - asinh(a*x)/(8*a**4)

Maxima [A] (verification not implemented)

none

Time = 0.23 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.43

$$\int e^{\operatorname{csch}^{-1}(ax)} x^3 dx = \frac{x^3}{3a} + \frac{\left(\frac{1}{a^2x^2} + 1\right)^{\frac{3}{2}} + \sqrt{\frac{1}{a^2x^2} + 1}}{8\left(a^4\left(\frac{1}{a^2x^2} + 1\right)^2 - 2a^4\left(\frac{1}{a^2x^2} + 1\right) + a^4\right)} - \frac{\log\left(\sqrt{\frac{1}{a^2x^2} + 1} + 1\right)}{16a^4} + \frac{\log\left(\sqrt{\frac{1}{a^2x^2} + 1} - 1\right)}{16a^4}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))*x^3,x, algorithm="maxima")

[Out] 1/3*x^3/a + 1/8*((1/(a^2*x^2) + 1)^(3/2) + sqrt(1/(a^2*x^2) + 1))/(a^4*(1/(a^2*x^2) + 1)^2 - 2*a^4*(1/(a^2*x^2) + 1) + a^4) - 1/16*log(sqrt(1/(a^2*x^2) + 1) + 1)/a^4 + 1/16*log(sqrt(1/(a^2*x^2) + 1) - 1)/a^4

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.92

$$\int e^{\operatorname{csch}^{-1}(ax)} x^3 dx = \frac{1}{8} \sqrt{a^2 x^2 + 1} \left(\frac{2x^2 |a| \operatorname{sgn}(x)}{a^2} + \frac{|a| \operatorname{sgn}(x)}{a^4} \right) x + \frac{x^3}{3a} + \frac{\log(-x|a| + \sqrt{a^2 x^2 + 1}) \operatorname{sgn}(x)}{8a^4}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))*x^3,x, algorithm="giac")

[Out] 1/8*sqrt(a^2*x^2 + 1)*(2*x^2*abs(a)*sgn(x)/a^2 + abs(a)*sgn(x)/a^4)*x + 1/3*x^3/a + 1/8*log(-x*abs(a) + sqrt(a^2*x^2 + 1))*sgn(x)/a^4

Mupad [B] (verification not implemented)

Time = 5.49 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.81

$$\int e^{\operatorname{csch}^{-1}(ax)} x^3 dx = \frac{x^4 \sqrt{\frac{1}{a^2 x^2} + 1}}{4} - \frac{\operatorname{atanh}\left(\sqrt{\frac{1}{a^2 x^2} + 1}\right)}{8a^4} + \frac{x^3}{3a} + \frac{x^2 \sqrt{\frac{1}{a^2 x^2} + 1}}{8a^2}$$

[In] int(x^3*((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x)),x)

[Out] (x^4*(1/(a^2*x^2) + 1)^(1/2))/4 - atanh((1/(a^2*x^2) + 1)^(1/2))/(8*a^4) + x^3/(3*a) + (x^2*(1/(a^2*x^2) + 1)^(1/2))/(8*a^2)

3.29 $\int e^{\operatorname{csch}^{-1}(ax)} x^2 dx$

Optimal result	219
Rubi [A] (verified)	219
Mathematica [A] (verified)	220
Maple [A] (verified)	220
Fricas [A] (verification not implemented)	221
Sympy [A] (verification not implemented)	221
Maxima [A] (verification not implemented)	221
Giac [A] (verification not implemented)	221
Mupad [B] (verification not implemented)	222

Optimal result

Integrand size = 10, antiderivative size = 31

$$\int e^{\operatorname{csch}^{-1}(ax)} x^2 dx = \frac{x^2}{2a} + \frac{1}{3} \left(1 + \frac{1}{a^2 x^2}\right)^{3/2} x^3$$

[Out] $1/2*x^2/a+1/3*(1+1/a^2/x^2)^(3/2)*x^3$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {6471, 30, 270}

$$\int e^{\operatorname{csch}^{-1}(ax)} x^2 dx = \frac{1}{3} x^3 \left(\frac{1}{a^2 x^2} + 1 \right)^{3/2} + \frac{x^2}{2a}$$

[In] `Int[E^ArcCsch[a*x]*x^2,x]`

[Out] $x^2/(2*a) + ((1 + 1/(a^2*x^2))^(3/2)*x^3)/3$

Rule 30

`Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]`

Rule 270

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, n, p}, x] && EqQ[(m + 1)/n + p + 1, 0] && NeQ[m, -1]`

Rule 6471

```
Int[E^ArcCsch[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Dist[1/a, Int[x^(m
- p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x
]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int x dx}{a} + \int \sqrt{1 + \frac{1}{a^2 x^2}} x^2 dx \\ &= \frac{x^2}{2a} + \frac{1}{3} \left(1 + \frac{1}{a^2 x^2}\right)^{3/2} x^3 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.23

$$\int e^{\text{csch}^{-1}(ax)} x^2 dx = \frac{3ax^2 + 2\sqrt{1 + \frac{1}{a^2 x^2}}(x + a^2 x^3)}{6a^2}$$

[In] Integrate[E^ArcCsch[a*x]*x^2,x]

[Out] (3*a*x^2 + 2*Sqrt[1 + 1/(a^2*x^2)]*(x + a^2*x^3))/(6*a^2)

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.39

method	result	size
default	$\frac{\sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} x (a^2 x^2 + 1)}{3a^2} + \frac{x^2}{2a}$	43
trager	$\frac{\frac{(x-1)(1+x)}{2} + \frac{(a^2 x^2 + 1)x \sqrt{-\frac{a^2 x^2 - 1}{a^2 x^2}}}{3a}}{a}$	49

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))*x^2,x,method=_RETURNVERBOSE)

[Out] 1/3*((a^2*x^2+1)/a^2/x^2)^(1/2)*x*(a^2*x^2+1)/a^2+1/2*x^2/a

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.32

$$\int e^{\operatorname{csch}^{-1}(ax)} x^2 dx = \frac{3ax^2 + 2(a^2x^3 + x)\sqrt{\frac{a^2x^2+1}{a^2x^2}}}{6a^2}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))*x^2,x, algorithm="fricas")

[Out] 1/6*(3*a*x^2 + 2*(a^2*x^3 + x)*sqrt((a^2*x^2 + 1)/(a^2*x^2)))/a^2

Sympy [A] (verification not implemented)

Time = 1.12 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.32

$$\int e^{\operatorname{csch}^{-1}(ax)} x^2 dx = \frac{x^2\sqrt{a^2x^2+1}}{3a} + \frac{x^2}{2a} + \frac{\sqrt{a^2x^2+1}}{3a^3}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))*x**2,x)

[Out] x**2*sqrt(a**2*x**2 + 1)/(3*a) + x**2/(2*a) + sqrt(a**2*x**2 + 1)/(3*a**3)

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.81

$$\int e^{\operatorname{csch}^{-1}(ax)} x^2 dx = \frac{1}{3} x^3 \left(\frac{1}{a^2 x^2} + 1 \right)^{\frac{3}{2}} + \frac{x^2}{2a}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))*x^2,x, algorithm="maxima")

[Out] 1/3*x^3*(1/(a^2*x^2) + 1)^(3/2) + 1/2*x^2/a

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.42

$$\int e^{\operatorname{csch}^{-1}(ax)} x^2 dx = \frac{(a^2x^2+1)^{\frac{3}{2}}|a\operatorname{sgn}(x)|}{3a^4} + \frac{a^2x^2+1}{2a^3} - \frac{|a\operatorname{sgn}(x)|}{3a^4}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))*x^2,x, algorithm="giac")

[Out] 1/3*(a^2*x^2 + 1)^(3/2)*abs(a)*sgn(x)/a^4 + 1/2*(a^2*x^2 + 1)/a^3 - 1/3*abs(a)*sgn(x)/a^4

Mupad [B] (verification not implemented)

Time = 5.05 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int e^{\operatorname{csch}^{-1}(ax)} x^2 dx = \left(\frac{x}{3a^2} + \frac{x^3}{3} \right) \sqrt{\frac{1}{a^2 x^2} + 1} + \frac{x^2}{2a}$$

[In] `int(x^2*((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x)),x)`

[Out] `(x/(3*a^2) + x^3/3)*(1/(a^2*x^2) + 1)^(1/2) + x^2/(2*a)`

3.30 $\int e^{\operatorname{csch}^{-1}(ax)} x dx$

Optimal result	223
Rubi [A] (verified)	223
Mathematica [A] (verified)	225
Maple [B] (verified)	225
Fricas [A] (verification not implemented)	225
Sympy [A] (verification not implemented)	226
Maxima [A] (verification not implemented)	226
Giac [A] (verification not implemented)	226
Mupad [B] (verification not implemented)	227

Optimal result

Integrand size = 8, antiderivative size = 47

$$\int e^{\operatorname{csch}^{-1}(ax)} x dx = \frac{x}{a} + \frac{1}{2} \sqrt{1 + \frac{1}{a^2 x^2}} x^2 + \frac{\operatorname{arctanh}\left(\sqrt{1 + \frac{1}{a^2 x^2}}\right)}{2a^2}$$

[Out] $x/a + 1/2 * \operatorname{arctanh}\left(\left(1 + 1/a^2/x^2\right)^{1/2}\right) / a^2 + 1/2 * x^2 * \left(1 + 1/a^2/x^2\right)^{1/2}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6471, 8, 272, 43, 65, 214}

$$\int e^{\operatorname{csch}^{-1}(ax)} x dx = \frac{\operatorname{arctanh}\left(\sqrt{\frac{1}{a^2 x^2} + 1}\right)}{2a^2} + \frac{1}{2} x^2 \sqrt{\frac{1}{a^2 x^2} + 1} + \frac{x}{a}$$

[In] `Int[E^ArcCsch[a*x]*x,x]`

[Out] $x/a + (\operatorname{Sqrt}[1 + 1/(a^2*x^2)]*x^2)/2 + \operatorname{ArcTanh}[\operatorname{Sqrt}[1 + 1/(a^2*x^2)]]/(2*a^2)$

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 43

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Dist[d*(n/(b*(m + 1))), I`

nt[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x]
 && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !IntegerQ[n] && GtQ[n, 0]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[
 {p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
 [b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
 ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
 /Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Dist[1/n, Subst[
 Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
 , m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 6471

Int[E^ArcCsch[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] :> Dist[1/a, Int[x^(m
 - p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x
]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{1}{a} dx + \int \sqrt{1 + \frac{1}{a^2 x^2}} x dx \\
 &= \frac{x}{a} - \frac{1}{2} \text{Subst} \left(\int \frac{\sqrt{1 + \frac{x}{a^2}}}{x^2} dx, x, \frac{1}{x^2} \right) \\
 &= \frac{x}{a} + \frac{1}{2} \sqrt{1 + \frac{1}{a^2 x^2}} x^2 - \frac{\text{Subst} \left(\int \frac{1}{x \sqrt{1 + \frac{x}{a^2}}} dx, x, \frac{1}{x^2} \right)}{4a^2} \\
 &= \frac{x}{a} + \frac{1}{2} \sqrt{1 + \frac{1}{a^2 x^2}} x^2 - \frac{1}{2} \text{Subst} \left(\int \frac{1}{-a^2 + a^2 x^2} dx, x, \sqrt{1 + \frac{1}{a^2 x^2}} \right) \\
 &= \frac{x}{a} + \frac{1}{2} \sqrt{1 + \frac{1}{a^2 x^2}} x^2 + \frac{\text{arctanh} \left(\sqrt{1 + \frac{1}{a^2 x^2}} \right)}{2a^2}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00

$$\int e^{\operatorname{csch}^{-1}(ax)} x dx = \frac{ax \left(2 + a \sqrt{1 + \frac{1}{a^2 x^2}} \right) + \log \left(\left(1 + \sqrt{1 + \frac{1}{a^2 x^2}} \right) x \right)}{2a^2}$$

[In] Integrate[E^ArcCsch[a*x]*x,x]

[Out] (a*x*(2 + a*Sqrt[1 + 1/(a^2*x^2)]*x) + Log[(1 + Sqrt[1 + 1/(a^2*x^2)])*x])/ (2*a^2)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 84 vs. 2(39) = 78.

Time = 0.05 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.81

method	result	size
default	$\frac{\sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} x \left(x \sqrt{\frac{a^2 x^2 + 1}{a^2}} a^2 + \ln \left(x + \sqrt{\frac{a^2 x^2 + 1}{a^2}} \right) \right)}{2 \sqrt{\frac{a^2 x^2 + 1}{a^2}} a^2} + \frac{x}{a}$	85

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))*x,x,method=_RETURNVERBOSE)

[Out] 1/2*((a^2*x^2+1)/a^2/x^2)^(1/2)*x*(x*((a^2*x^2+1)/a^2)^(1/2)*a^2+ln(x+((a^2*x^2+1)/a^2)^(1/2)))/((a^2*x^2+1)/a^2)^(1/2)/a^2+x/a

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.36

$$\int e^{\operatorname{csch}^{-1}(ax)} x dx = \frac{a^2 x^2 \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} + 2ax - \log \left(ax \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} - ax \right)}{2a^2}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))*x,x, algorithm="fricas")

[Out] 1/2*(a^2*x^2*sqrt((a^2*x^2 + 1)/(a^2*x^2)) + 2*a*x - log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x))/a^2

Sympy [A] (verification not implemented)

Time = 1.57 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.62

$$\int e^{\operatorname{csch}^{-1}(ax)} x dx = \frac{x\sqrt{a^2x^2+1}}{2a} + \frac{x}{a} + \frac{\operatorname{asinh}(ax)}{2a^2}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))*x,x)

[Out] x*sqrt(a**2*x**2 + 1)/(2*a) + x/a + asinh(a*x)/(2*a**2)

Maxima [A] (verification not implemented)

none

Time = 0.23 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.66

$$\int e^{\operatorname{csch}^{-1}(ax)} x dx = \frac{x}{a} + \frac{\sqrt{\frac{1}{a^2x^2}+1}}{2\left(a^2\left(\frac{1}{a^2x^2}+1\right)-a^2\right)} + \frac{\log\left(\sqrt{\frac{1}{a^2x^2}+1}+1\right)}{4a^2} - \frac{\log\left(\sqrt{\frac{1}{a^2x^2}+1}-1\right)}{4a^2}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))*x,x, algorithm="maxima")

[Out] x/a + 1/2*sqrt(1/(a^2*x^2) + 1)/(a^2*(1/(a^2*x^2) + 1) - a^2) + 1/4*log(sqrt(1/(a^2*x^2) + 1) + 1)/a^2 - 1/4*log(sqrt(1/(a^2*x^2) + 1) - 1)/a^2

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.11

$$\int e^{\operatorname{csch}^{-1}(ax)} x dx = \frac{\sqrt{a^2x^2+1}x|a|\operatorname{sgn}(x)}{2a^2} + \frac{x}{a} - \frac{\log(-x|a| + \sqrt{a^2x^2+1})\operatorname{sgn}(x)}{2a^2}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))*x,x, algorithm="giac")

[Out] 1/2*sqrt(a^2*x^2 + 1)*x*abs(a)*sgn(x)/a^2 + x/a - 1/2*log(-x*abs(a) + sqrt(a^2*x^2 + 1))*sgn(x)/a^2

Mupad [B] (verification not implemented)

Time = 4.88 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.83

$$\int e^{\operatorname{csch}^{-1}(ax)} x \, dx = \frac{\operatorname{atanh}\left(\sqrt{\frac{1}{a^2 x^2} + 1}\right)}{2 a^2} + \frac{x}{a} + \frac{x^2 \sqrt{\frac{1}{a^2 x^2} + 1}}{2}$$

[In] int(x*((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x)),x)

[Out] atanh((1/(a^2*x^2) + 1)^(1/2))/(2*a^2) + x/a + (x^2*(1/(a^2*x^2) + 1)^(1/2))/2

3.31 $\int e^{\operatorname{csch}^{-1}(ax)} dx$

Optimal result	228
Rubi [A] (verified)	228
Mathematica [A] (verified)	229
Maple [B] (verified)	230
Fricas [B] (verification not implemented)	230
Sympy [A] (verification not implemented)	230
Maxima [A] (verification not implemented)	231
Giac [A] (verification not implemented)	231
Mupad [B] (verification not implemented)	231

Optimal result

Integrand size = 6, antiderivative size = 24

$$\int e^{\operatorname{csch}^{-1}(ax)} dx = e^{\operatorname{csch}^{-1}(ax)} x - \frac{\operatorname{csch}^{-1}(ax)}{a} + \frac{\log(x)}{a}$$

[Out] $(1/a/x+(1+1/a^2/x^2)^{(1/2)}) * x - \operatorname{arccsch}(a*x)/a + \ln(x)/a$

Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 31, normalized size of antiderivative = 1.29, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {6466, 29, 248, 283, 221}

$$\int e^{\operatorname{csch}^{-1}(ax)} dx = x \sqrt{\frac{1}{a^2 x^2} + 1} + \frac{\log(x)}{a} - \frac{\operatorname{csch}^{-1}(ax)}{a}$$

[In] `Int[E^ArcCsch[a*x], x]`

[Out] `Sqrt[1 + 1/(a^2*x^2)]*x - ArcCsch[a*x]/a + Log[x]/a`

Rule 29

`Int[(x_)^(-1), x_Symbol] := Simp[Log[x], x]`

Rule 221

`Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

Rule 248

```
Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^2, x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0]
```

Rule 283

```
Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m+1)*((a + b*x^n)^p/(c*(m+1))), x] - Dist[b*n*(p/(c^n*(m+1))), Int[(c*x)^(m+n)*(a + b*x^n)^(p-1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && GtQ[p, 0] && LtQ[m, -1] && !ILtQ[(m+n*p+n+1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 6466

```
Int[E^ArcCsch[(a_)*(x_)^(p_)], x_Symbol] := Dist[1/a, Int[1/x^p, x], x] + Int[Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, p}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int \frac{1}{x} dx}{a} + \int \sqrt{1 + \frac{1}{a^2 x^2}} dx \\
 &= \frac{\log(x)}{a} - \text{Subst}\left(\int \frac{\sqrt{1 + \frac{x^2}{a^2}}}{x^2} dx, x, \frac{1}{x}\right) \\
 &= \sqrt{1 + \frac{1}{a^2 x^2}} x + \frac{\log(x)}{a} - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1 + \frac{x^2}{a^2}}} dx, x, \frac{1}{x}\right)}{a^2} \\
 &= \sqrt{1 + \frac{1}{a^2 x^2}} x - \frac{\text{csch}^{-1}(ax)}{a} + \frac{\log(x)}{a}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.46

$$\int e^{\text{csch}^{-1}(ax)} dx = \frac{a\sqrt{1 + \frac{1}{a^2 x^2}} x - \text{arcsinh}\left(\frac{1}{ax}\right) + \log(ax)}{a}$$

```
[In] Integrate[E^ArcCsch[a*x], x]
```

```
[Out] (a*Sqrt[1 + 1/(a^2*x^2)]*x - ArcSinh[1/(a*x)] + Log[a*x])/a
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 112 vs. $2(37) = 74$.

Time = 0.10 (sec) , antiderivative size = 113, normalized size of antiderivative = 4.71

method	result	size
default	$-\frac{\sqrt{\frac{a^2x^2+1}{a^2x^2}} x \left(-\sqrt{\frac{1}{a^2}} \sqrt{\frac{a^2x^2+1}{a^2}} a^2 + \ln \left(\frac{2\sqrt{\frac{1}{a^2}} \sqrt{\frac{a^2x^2+1}{a^2}} a^2 + 2}{a^2x} \right) \right)}{\sqrt{\frac{1}{a^2}} \sqrt{\frac{a^2x^2+1}{a^2}} a^2} + \frac{\ln(x)}{a}$	113

[In] `int(1/a/x+(1+1/a^2/x^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $-\left(\frac{a^2x^2+1}{a^2x^2}\right)^{1/2} * x * \left(-\frac{1}{a^2}\right)^{1/2} * \left(\frac{a^2x^2+1}{a^2}\right)^{1/2} * a^2 + \ln\left(2 * \left(\frac{1}{a^2}\right)^{1/2} * \left(\frac{a^2x^2+1}{a^2}\right)^{1/2} * a^2 + 1\right) / \left(\frac{1}{a^2}\right)^{1/2} / \left(\frac{a^2x^2+1}{a^2}\right)^{1/2} / a^2 + \ln(x) / a$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 86 vs. $2(37) = 74$.

Time = 0.24 (sec) , antiderivative size = 86, normalized size of antiderivative = 3.58

$$\int e^{\operatorname{csch}^{-1}(ax)} dx = \frac{ax \sqrt{\frac{a^2x^2+1}{a^2x^2}} - \log\left(ax \sqrt{\frac{a^2x^2+1}{a^2x^2}} - ax + 1\right) + \log\left(ax \sqrt{\frac{a^2x^2+1}{a^2x^2}} - ax - 1\right) + \log(x)}{a}$$

[In] `integrate(1/a/x+(1+1/a^2/x^2)^(1/2),x, algorithm="fricas")`

[Out] $(ax * \sqrt{(a^2x^2 + 1)/(a^2x^2)}) - \log(ax * \sqrt{(a^2x^2 + 1)/(a^2x^2)} - ax + 1) + \log(ax * \sqrt{(a^2x^2 + 1)/(a^2x^2)} - ax - 1) + \log(x) / a$

Sympy [A] (verification not implemented)

Time = 0.76 (sec) , antiderivative size = 48, normalized size of antiderivative = 2.00

$$\int e^{\operatorname{csch}^{-1}(ax)} dx = \frac{x}{\sqrt{1 + \frac{1}{a^2x^2}}} + \frac{\log(x)}{a} - \frac{\operatorname{asinh}\left(\frac{1}{ax}\right)}{a} + \frac{1}{a^2x \sqrt{1 + \frac{1}{a^2x^2}}}$$

[In] `integrate(1/a/x+(1+1/a**2/x**2)**(1/2),x)`

[Out] $x/\sqrt{1 + 1/(a**2*x**2)} + \log(x)/a - \operatorname{asinh}(1/(a*x))/a + 1/(a**2*x*\sqrt{1 + 1/(a**2*x**2)})$

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 64, normalized size of antiderivative = 2.67

$$\int e^{\operatorname{csch}^{-1}(ax)} dx = x \sqrt{\frac{1}{a^2 x^2} + 1} - \frac{\log\left(ax \sqrt{\frac{1}{a^2 x^2} + 1} + 1\right)}{2a} + \frac{\log\left(ax \sqrt{\frac{1}{a^2 x^2} + 1} - 1\right)}{2a} + \frac{\log(x)}{a}$$

[In] integrate(1/a/x+(1+1/a^2/x^2)^(1/2),x, algorithm="maxima")

[Out] x*sqrt(1/(a^2*x^2) + 1) - 1/2*log(a*x*sqrt(1/(a^2*x^2) + 1) + 1)/a + 1/2*log(a*x*sqrt(1/(a^2*x^2) + 1) - 1)/a + log(x)/a

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.75

$$\begin{aligned} \int e^{\operatorname{csch}^{-1}(ax)} dx \\ = -\frac{(\log(\sqrt{a^2 x^2 + 1} + 1) \operatorname{sgn}(x) - \log(\sqrt{a^2 x^2 + 1} - 1) \operatorname{sgn}(x) - 2\sqrt{a^2 x^2 + 1} \operatorname{sgn}(x))|a|}{2a^2} \\ + \frac{\log(|x|)}{a} \end{aligned}$$

[In] integrate(1/a/x+(1+1/a^2/x^2)^(1/2),x, algorithm="giac")

[Out] -1/2*(log(sqrt(a^2*x^2 + 1) + 1)*sgn(x) - log(sqrt(a^2*x^2 + 1) - 1)*sgn(x) - 2*sqrt(a^2*x^2 + 1)*sgn(x))*abs(a)/a^2 + log(abs(x))/a

Mupad [B] (verification not implemented)

Time = 4.89 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.50

$$\int e^{\operatorname{csch}^{-1}(ax)} dx = \frac{\ln(x)}{a} + x \sqrt{\frac{1}{a^2 x^2} + 1} + \frac{\operatorname{asin}\left(\frac{1i}{ax}\right) 1i}{a}$$

[In] int((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x),x)

[Out] log(x)/a + (asin(1i/(a*x))*1i)/a + x*(1/(a^2*x^2) + 1)^(1/2)

3.32 $\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x} dx$

Optimal result	232
Rubi [A] (verified)	232
Mathematica [A] (verified)	234
Maple [B] (verified)	234
Fricas [A] (verification not implemented)	234
Sympy [A] (verification not implemented)	235
Maxima [A] (verification not implemented)	235
Giac [F(-2)]	235
Mupad [B] (verification not implemented)	236

Optimal result

Integrand size = 10, antiderivative size = 38

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x} dx = -\sqrt{1 + \frac{1}{a^2 x^2}} - \frac{1}{ax} + \operatorname{arctanh}\left(\sqrt{1 + \frac{1}{a^2 x^2}}\right)$$

[Out] $-1/a/x + \operatorname{arctanh}\left(\left(1 + 1/a^2/x^2\right)^{1/2}\right) - \left(1 + 1/a^2/x^2\right)^{1/2}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6471, 30, 272, 52, 65, 214}

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x} dx = \operatorname{arctanh}\left(\sqrt{\frac{1}{a^2 x^2} + 1}\right) - \sqrt{\frac{1}{a^2 x^2} + 1} - \frac{1}{ax}$$

[In] `Int[E^ArcCsch[a*x]/x,x]`

[Out] `-Sqrt[1 + 1/(a^2*x^2)] - 1/(a*x) + ArcTanh[Sqrt[1 + 1/(a^2*x^2)]]`

Rule 30

`Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]`

Rule 52

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(`

$b*(m + n + 1))$, Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 6471

Int[E^ArcCsch[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Dist[1/a, Int[x^(m - p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int \frac{1}{x^2} dx}{a} + \int \frac{\sqrt{1 + \frac{1}{a^2 x^2}}}{x} dx \\
 &= -\frac{1}{ax} - \frac{1}{2} \text{Subst} \left(\int \frac{\sqrt{1 + \frac{x}{a^2}}}{x} dx, x, \frac{1}{x^2} \right) \\
 &= -\sqrt{1 + \frac{1}{a^2 x^2}} - \frac{1}{ax} - \frac{1}{2} \text{Subst} \left(\int \frac{1}{x \sqrt{1 + \frac{x}{a^2}}} dx, x, \frac{1}{x^2} \right) \\
 &= -\sqrt{1 + \frac{1}{a^2 x^2}} - \frac{1}{ax} - a^2 \text{Subst} \left(\int \frac{1}{-a^2 + a^2 x^2} dx, x, \sqrt{1 + \frac{1}{a^2 x^2}} \right) \\
 &= -\sqrt{1 + \frac{1}{a^2 x^2}} - \frac{1}{ax} + \text{arctanh} \left(\sqrt{1 + \frac{1}{a^2 x^2}} \right)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.11

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x} dx = -\sqrt{1 + \frac{1}{a^2 x^2}} - \frac{1}{ax} + \log \left(\left(1 + \sqrt{1 + \frac{1}{a^2 x^2}} \right) x \right)$$

[In] Integrate[E^ArcCsch[a*x]/x,x]

[Out] -Sqrt[1 + 1/(a^2*x^2)] - 1/(a*x) + Log[(1 + Sqrt[1 + 1/(a^2*x^2)])*x]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 108 vs. 2(34) = 68.

Time = 0.05 (sec) , antiderivative size = 109, normalized size of antiderivative = 2.87

method	result	size
default	$-\frac{\sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} \left(a^2 \left(\frac{a^2 x^2 + 1}{a^2} \right)^{\frac{3}{2}} - \sqrt{\frac{a^2 x^2 + 1}{a^2}} a^2 x^2 - \ln \left(x + \sqrt{\frac{a^2 x^2 + 1}{a^2}} \right) x \right)}{\sqrt{\frac{a^2 x^2 + 1}{a^2}}} - \frac{1}{ax}$	109

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))/x,x,method=_RETURNVERBOSE)

[Out] -((a^2*x^2+1)/a^2/x^2)^(1/2)*(a^2*((a^2*x^2+1)/a^2)^(3/2)-((a^2*x^2+1)/a^2)^(1/2)*a^2*x^2-ln(x+((a^2*x^2+1)/a^2)^(1/2))*x)/((a^2*x^2+1)/a^2)^(1/2)-1/a/x

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.68

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x} dx = -\frac{ax \log \left(ax \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} - ax \right) + ax \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} + ax + 1}{ax}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))/x,x, algorithm="fricas")

[Out] -(a*x*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x) + a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) + a*x + 1)/(a*x)

Sympy [A] (verification not implemented)

Time = 2.27 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.08

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x} dx = -\frac{ax}{\sqrt{a^2x^2+1}} + \operatorname{asinh}(ax) - \frac{1}{ax} - \frac{1}{ax\sqrt{a^2x^2+1}}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))/x,x)

[Out] -a*x/sqrt(a**2*x**2 + 1) + asinh(a*x) - 1/(a*x) - 1/(a*x*sqrt(a**2*x**2 + 1))

Maxima [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.42

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x} dx = -\sqrt{\frac{1}{a^2x^2}+1} - \frac{1}{ax} + \frac{1}{2} \log\left(\sqrt{\frac{1}{a^2x^2}+1}+1\right) - \frac{1}{2} \log\left(\sqrt{\frac{1}{a^2x^2}+1}-1\right)$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))/x,x, algorithm="maxima")

[Out] -sqrt(1/(a^2*x^2) + 1) - 1/(a*x) + 1/2*log(sqrt(1/(a^2*x^2) + 1) + 1) - 1/2*log(sqrt(1/(a^2*x^2) + 1) - 1)

Giac [F(-2)]

Exception generated.

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x} dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))/x,x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type

Mupad [B] (verification not implemented)

Time = 5.16 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.89

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x} dx = \operatorname{atanh}\left(\sqrt{\frac{1}{a^2 x^2} + 1}\right) - \sqrt{\frac{1}{a^2 x^2} + 1} - \frac{1}{ax}$$

[In] int(((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))/x,x)

[Out] atanh((1/(a^2*x^2) + 1)^(1/2)) - (1/(a^2*x^2) + 1)^(1/2) - 1/(a*x)

3.33 $\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^2} dx$

Optimal result	237
Rubi [A] (verified)	237
Mathematica [A] (verified)	239
Maple [B] (verified)	239
Fricas [B] (verification not implemented)	239
Sympy [A] (verification not implemented)	240
Maxima [B] (verification not implemented)	240
Giac [B] (verification not implemented)	240
Mupad [B] (verification not implemented)	241

Optimal result

Integrand size = 10, antiderivative size = 40

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^2} dx = -\frac{1}{2ax^2} - \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{2x} - \frac{1}{2} \operatorname{acsch}^{-1}(ax)$$

[Out] $-1/2/a/x^2 - 1/2*a*\operatorname{arccsch}(a*x) - 1/2*(1+1/a^2/x^2)^{(1/2)}/x$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6471, 30, 342, 201, 221}

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^2} dx = -\frac{\sqrt{\frac{1}{a^2x^2} + 1}}{2x} - \frac{1}{2ax^2} - \frac{1}{2} \operatorname{acsch}^{-1}(ax)$$

[In] $\operatorname{Int}[E^{\operatorname{ArcCsch}[a*x]}/x^2, x]$

[Out] $-1/2*1/(a*x^2) - \operatorname{Sqrt}[1 + 1/(a^2*x^2)]/(2*x) - (a*\operatorname{ArcCsch}[a*x])/2$

Rule 30

$\operatorname{Int}[(x_)^{(m_.)}, x_Symbol] \rightarrow \operatorname{Simp}[x^{(m+1)}/(m+1), x] /; \operatorname{FreeQ}[m, x] \ \&\& \operatorname{NeQ}[m, -1]$

Rule 201

$\operatorname{Int}[((a_) + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \operatorname{Simp}[x*((a + b*x^n)^p/(n*p + 1)), x] + \operatorname{Dist}[a*n*(p/(n*p + 1)), \operatorname{Int}[(a + b*x^n)^{(p-1)}, x], x] /; \operatorname{FreeQ}[a, b, n, p]$

$Q[\{a, b\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{GtQ}[p, 0] \&\& (\text{IntegerQ}[2*p] \parallel (\text{EqQ}[n, 2] \&\& \text{IntegerQ}[4*p]) \parallel (\text{EqQ}[n, 2] \&\& \text{IntegerQ}[3*p]) \parallel \text{LtQ}[\text{Denominator}[p + 1/n], \text{Denominator}[p]])$

Rule 221

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2], x_Symbol] \text{ :> } \text{Simp}[\text{ArcSinh}[\text{Rt}[b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[b, 2], x] \text{ /; } \text{FreeQ}[\{a, b\}, x] \&\& \text{GtQ}[a, 0] \&\& \text{PosQ}[b]$

Rule 342

$\text{Int}[(x_)^{(m_.)*((a_) + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \text{ :> } -\text{Subst}[\text{Int}[(a + b/x^n)^p/x^{(m+2)}, x], x, 1/x] \text{ /; } \text{FreeQ}[\{a, b, p\}, x] \&\& \text{ILtQ}[n, 0] \&\& \text{IntegerQ}[m]$

Rule 6471

$\text{Int}[\text{E}^{\text{ArcCsch}[(a_)*(x_)^{(p_.)}]}*(x_)^{(m_.)}, x_Symbol] \text{ :> } \text{Dist}[1/a, \text{Int}[x^{(m-p)}, x], x] + \text{Int}[x^m*\text{Sqrt}[1 + 1/(a^2*x^{(2*p)})], x] \text{ /; } \text{FreeQ}[\{a, m, p\}, x]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int \frac{1}{x^3} dx}{a} + \int \frac{\sqrt{1 + \frac{1}{a^2 x^2}}}{x^2} dx \\
 &= -\frac{1}{2ax^2} - \text{Subst}\left(\int \sqrt{1 + \frac{x^2}{a^2}} dx, x, \frac{1}{x}\right) \\
 &= -\frac{1}{2ax^2} - \frac{\sqrt{1 + \frac{1}{a^2 x^2}}}{2x} - \frac{1}{2} \text{Subst}\left(\int \frac{1}{\sqrt{1 + \frac{x^2}{a^2}}} dx, x, \frac{1}{x}\right) \\
 &= -\frac{1}{2ax^2} - \frac{\sqrt{1 + \frac{1}{a^2 x^2}}}{2x} - \frac{1}{2} \text{acsch}^{-1}(ax)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.08

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^2} dx = -\frac{1 + a\sqrt{1 + \frac{1}{a^2x^2}}x + a^2x^2 \operatorname{arcsinh}\left(\frac{1}{ax}\right)}{2ax^2}$$

[In] Integrate[E^ArcCsch[a*x]/x^2,x]

[Out] -1/2*(1 + a*Sqrt[1 + 1/(a^2*x^2)]*x + a^2*x^2*ArcSinh[1/(a*x)])/(a*x^2)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 144 vs. 2(32) = 64.

Time = 0.05 (sec) , antiderivative size = 145, normalized size of antiderivative = 3.62

method	result	size
default	$-\frac{\sqrt{\frac{a^2x^2+1}{a^2x^2}} \left(a^2 \left(\frac{a^2x^2+1}{a^2} \right)^{\frac{3}{2}} \sqrt{\frac{1}{a^2}} - \sqrt{\frac{a^2x^2+1}{a^2}} \sqrt{\frac{1}{a^2}} a^2x^2 + \ln \left(\frac{2\sqrt{\frac{1}{a^2}} \sqrt{\frac{a^2x^2+1}{a^2}} a^2+2}{a^2x} \right) x^2 \right)}{2x\sqrt{\frac{a^2x^2+1}{a^2}} \sqrt{\frac{1}{a^2}}} - \frac{1}{2ax^2}$	145

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))/x^2,x,method=_RETURNVERBOSE)

[Out] -1/2*((a^2*x^2+1)/a^2/x^2)^(1/2)/x*(a^2*((a^2*x^2+1)/a^2)^(3/2)*(1/a^2)^(1/2)-((a^2*x^2+1)/a^2)^(1/2)*(1/a^2)^(1/2)*a^2*x^2+ln(2*((1/a^2)^(1/2))*((a^2*x^2+1)/a^2)^(1/2)*a^2+1)/x/a^2)*x^2)/((a^2*x^2+1)/a^2)^(1/2)/(1/a^2)^(1/2)-1/2/a/x^2

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 102 vs. 2(32) = 64.

Time = 0.25 (sec) , antiderivative size = 102, normalized size of antiderivative = 2.55

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^2} dx = -\frac{a^2x^2 \log\left(ax\sqrt{\frac{a^2x^2+1}{a^2x^2}} - ax + 1\right) - a^2x^2 \log\left(ax\sqrt{\frac{a^2x^2+1}{a^2x^2}} - ax - 1\right) + ax\sqrt{\frac{a^2x^2+1}{a^2x^2}} + 1}{2ax^2}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))/x^2,x, algorithm="fricas")

[Out] -1/2*(a^2*x^2*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x + 1) - a^2*x^2*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x - 1) + a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) + 1)/(a*x^2)

Sympy [A] (verification not implemented)

Time = 0.89 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.92

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^2} dx = -a \left(\frac{\operatorname{asinh}\left(\frac{1}{ax}\right)}{2} + \frac{\sqrt{1 + \frac{1}{a^2 x^2}}}{2ax} \right) - \frac{1}{2ax^2}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))/x**2,x)

[Out] -a*(asinh(1/(a*x))/2 + sqrt(1 + 1/(a**2*x**2)))/(2*a*x) - 1/(2*a*x**2)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 86 vs. 2(32) = 64.

Time = 0.23 (sec) , antiderivative size = 86, normalized size of antiderivative = 2.15

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^2} dx = -\frac{a^2 x \sqrt{\frac{1}{a^2 x^2} + 1}}{2(a^2 x^2(\frac{1}{a^2 x^2} + 1) - 1)} - \frac{1}{4} a \log \left(ax \sqrt{\frac{1}{a^2 x^2} + 1} + 1 \right) + \frac{1}{4} a \log \left(ax \sqrt{\frac{1}{a^2 x^2} + 1} - 1 \right) - \frac{1}{2ax^2}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))/x^2,x, algorithm="maxima")

[Out] -1/2*a^2*x*sqrt(1/(a^2*x^2) + 1)/(a^2*x^2*(1/(a^2*x^2) + 1) - 1) - 1/4*a*log(a*x*sqrt(1/(a^2*x^2) + 1) + 1) + 1/4*a*log(a*x*sqrt(1/(a^2*x^2) + 1) - 1) - 1/2/(a*x^2)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 82 vs. 2(32) = 64.

Time = 0.27 (sec) , antiderivative size = 82, normalized size of antiderivative = 2.05

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^2} dx = \frac{a^4 |a| \log(\sqrt{a^2 x^2 + 1} + 1) \operatorname{sgn}(x) - a^4 |a| \log(\sqrt{a^2 x^2 + 1} - 1) \operatorname{sgn}(x) + \frac{2(\sqrt{a^2 x^2 + 1} a^4 |a| \operatorname{sgn}(x) + a^5)}{a^2 x^2}}{4 a^4}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))/x^2,x, algorithm="giac")

[Out] -1/4*(a^4*abs(a)*log(sqrt(a^2*x^2 + 1) + 1)*sgn(x) - a^4*abs(a)*log(sqrt(a^2*x^2 + 1) - 1)*sgn(x) + 2*(sqrt(a^2*x^2 + 1)*a^4*abs(a)*sgn(x) + a^5)/(a^2*x^2))/a^4

Mupad [B] (verification not implemented)

Time = 5.63 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.05

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^2} dx = -\frac{\operatorname{asinh}\left(\frac{\sqrt{\frac{1}{a^2}}}{x}\right)}{2\sqrt{\frac{1}{a^2}}} - \frac{\sqrt{\frac{1}{a^2}x^2 + 1}}{2x} - \frac{1}{2ax^2}$$

[In] int(((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))/x^2,x)

[Out] - asinh((1/a^2)^(1/2)/x)/(2*(1/a^2)^(1/2)) - (1/(a^2*x^2) + 1)^(1/2)/(2*x) - 1/(2*a*x^2)

3.34 $\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^3} dx$

Optimal result	242
Rubi [A] (verified)	242
Mathematica [A] (verified)	243
Maple [A] (verified)	243
Fricas [A] (verification not implemented)	244
Sympy [A] (verification not implemented)	244
Maxima [A] (verification not implemented)	244
Giac [B] (verification not implemented)	245
Mupad [B] (verification not implemented)	245

Optimal result

Integrand size = 10, antiderivative size = 31

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^3} dx = -\frac{1}{3}a^2 \left(1 + \frac{1}{a^2x^2}\right)^{3/2} - \frac{1}{3ax^3}$$

[Out] $-1/3*a^2*(1+1/a^2/x^2)^{(3/2)}-1/3/a/x^3$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {6471, 30, 267}

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^3} dx = -\frac{1}{3}a^2 \left(\frac{1}{a^2x^2} + 1\right)^{3/2} - \frac{1}{3ax^3}$$

[In] $\text{Int}[E^{\text{ArcCsch}[a*x]}/x^3, x]$

[Out] $-1/3*(a^2*(1 + 1/(a^2*x^2)))^{(3/2)} - 1/(3*a*x^3)$

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}/(m+1), x] /; \text{FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 267

$\text{Int}[(x_)^{(m_.)}*((a_) + (b_.)*(x_)^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x^n)^{(p+1)}/(b*n*(p+1)), x] /; \text{FreeQ}\{a, b, m, n, p\}, x \ \&\& \ \text{EqQ}[m, n-1] \ \&\& \ \text{NeQ}[p, -1]$

Rule 6471

```
Int[E^ArcCsch[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Dist[1/a, Int[x^(m
- p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x
]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \frac{1}{x^4} dx}{a} + \int \frac{\sqrt{1 + \frac{1}{a^2 x^2}}}{x^3} dx \\ &= -\frac{1}{3} a^2 \left(1 + \frac{1}{a^2 x^2}\right)^{3/2} - \frac{1}{3ax^3} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.19

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^3} dx = -\frac{1 + a\sqrt{1 + \frac{1}{a^2 x^2}} x(1 + a^2 x^2)}{3ax^3}$$

[In] Integrate[E^ArcCsch[a*x]/x^3,x]

[Out] -1/3*(1 + a*Sqrt[1 + 1/(a^2*x^2)]*x*(1 + a^2*x^2))/(a*x^3)

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.35

method	result	size
default	$-\frac{\sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} (a^2 x^2 + 1)}{3x^2} - \frac{1}{3ax^3}$	42
trager	$-\frac{\frac{1}{3x^3} - \frac{a(a^2 x^2 + 1)\sqrt{-\frac{a^2 x^2 - 1}{a^2 x^2}}}{3x^2}}{a}$	46

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))/x^3,x,method=_RETURNVERBOSE)

[Out] -1/3*((a^2*x^2+1)/a^2/x^2)^(1/2)/x^2*(a^2*x^2+1)-1/3/a/x^3

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.52

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^3} dx = -\frac{a^3 x^3 + (a^3 x^3 + ax) \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} + 1}{3 a x^3}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))/x^3,x, algorithm="fricas")

[Out] -1/3*(a^3*x^3 + (a^3*x^3 + a*x)*sqrt((a^2*x^2 + 1)/(a^2*x^2)) + 1)/(a*x^3)

Sympy [A] (verification not implemented)

Time = 1.00 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.55

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^3} dx = \begin{cases} -a \left(\begin{cases} \sqrt{1 + \frac{1}{a^2 x^2}} \left(\frac{a^2}{3} + \frac{1}{3x^2} \right) & \text{for } \frac{1}{a^2} \neq 0 \\ \frac{1}{2x^2} & \text{otherwise} \end{cases} \right)^{-\frac{1}{3x^3}} \\ \frac{\quad}{a} & \text{for } a \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))/x**3,x)

[Out] Piecewise((-a*Piecewise((sqrt(1 + 1/(a**2*x**2)))*(a**2/3 + 1/(3*x**2)), Ne(a**(-2), 0)), (1/(2*x**2), True)) - 1/(3*x**3))/a, Ne(a, 0)), (0, True))

Maxima [A] (verification not implemented)

none

Time = 0.23 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.81

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^3} dx = -\frac{1}{3} a^2 \left(\frac{1}{a^2 x^2} + 1 \right)^{\frac{3}{2}} - \frac{1}{3 a x^3}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))/x^3,x, algorithm="maxima")

[Out] -1/3*a^2*(1/(a^2*x^2) + 1)^(3/2) - 1/3/(a*x^3)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 69 vs. $2(25) = 50$.

Time = 0.28 (sec) , antiderivative size = 69, normalized size of antiderivative = 2.23

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^3} dx = \frac{2 \left(3 (x|a| - \sqrt{a^2x^2 + 1})^4 a^2 \operatorname{sgn}(x) + a^2 \operatorname{sgn}(x) \right)}{3 \left((x|a| - \sqrt{a^2x^2 + 1})^2 - 1 \right)^3} - \frac{1}{3ax^3}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))/x^3,x, algorithm="giac")

[Out] 2/3*(3*(x*abs(a) - sqrt(a^2*x^2 + 1))^4*a^2*sgn(x) + a^2*sgn(x))/((x*abs(a) - sqrt(a^2*x^2 + 1))^2 - 1)^3 - 1/3/(a*x^3)

Mupad [B] (verification not implemented)

Time = 4.80 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.35

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^3} dx = -\frac{x\sqrt{\frac{1}{a^2x^2}+1}}{3} + \frac{1}{3a} - \frac{a^2\sqrt{\frac{1}{a^2x^2}+1}}{3}$$

[In] int(((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))/x^3,x)

[Out] - ((x*(1/(a^2*x^2) + 1)^(1/2))/3 + 1/(3*a))/x^3 - (a^2*(1/(a^2*x^2) + 1)^(1/2))/3

3.35 $\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^4} dx$

Optimal result	246
Rubi [A] (verified)	246
Mathematica [A] (verified)	248
Maple [B] (verified)	248
Fricas [B] (verification not implemented)	248
Sympy [A] (verification not implemented)	249
Maxima [B] (verification not implemented)	249
Giac [A] (verification not implemented)	250
Mupad [B] (verification not implemented)	250

Optimal result

Integrand size = 10, antiderivative size = 65

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^4} dx = -\frac{1}{4ax^4} - \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{4x^3} - \frac{a^2\sqrt{1 + \frac{1}{a^2x^2}}}{8x} + \frac{1}{8}a^3\operatorname{csch}^{-1}(ax)$$

[Out] $-1/4/a/x^4 + 1/8*a^3*\operatorname{arccsch}(a*x) - 1/4*(1 + 1/a^2/x^2)^{(1/2)}/x^3 - 1/8*a^2*(1 + 1/a^2/x^2)^{(1/2)}/x$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6471, 30, 342, 285, 327, 221}

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^4} dx = \frac{1}{8}a^3\operatorname{csch}^{-1}(ax) - \frac{a^2\sqrt{\frac{1}{a^2x^2} + 1}}{8x} - \frac{\sqrt{\frac{1}{a^2x^2} + 1}}{4x^3} - \frac{1}{4ax^4}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcCsch}[a*x]}/x^4, x]$

[Out] $-1/4*1/(a*x^4) - \operatorname{Sqrt}[1 + 1/(a^2*x^2)]/(4*x^3) - (a^2*\operatorname{Sqrt}[1 + 1/(a^2*x^2)])/(8*x) + (a^3*\operatorname{ArcCsch}[a*x])/8$

Rule 30

$\operatorname{Int}[(x_)^m, x_Symbol] \rightarrow \operatorname{Simp}[x^{(m+1)}/(m+1), x] /; \operatorname{FreeQ}[m, x] \ \&\& \operatorname{N} \operatorname{eQ}[m, -1]$

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 285

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^p/(c*(m + n*p + 1))), x] + Dist[a*n*(p/(m + n*p + 1)), Int[(c*x)^m*(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[n, 0] && GtQ[p, 0] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 327

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 342

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]

Rule 6471

Int[E^ArcCsch[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Dist[1/a, Int[x^(m - p), x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{1}{x^5} dx + \int \frac{\sqrt{1 + \frac{1}{a^2 x^2}}}{x^4} dx \\
 &= -\frac{1}{4ax^4} - \text{Subst}\left(\int x^2 \sqrt{1 + \frac{x^2}{a^2}} dx, x, \frac{1}{x}\right) \\
 &= -\frac{1}{4ax^4} - \frac{\sqrt{1 + \frac{1}{a^2 x^2}}}{4x^3} - \frac{1}{4} \text{Subst}\left(\int \frac{x^2}{\sqrt{1 + \frac{x^2}{a^2}}} dx, x, \frac{1}{x}\right) \\
 &= -\frac{1}{4ax^4} - \frac{\sqrt{1 + \frac{1}{a^2 x^2}}}{4x^3} - \frac{a^2 \sqrt{1 + \frac{1}{a^2 x^2}}}{8x} + \frac{1}{8} a^2 \text{Subst}\left(\int \frac{1}{\sqrt{1 + \frac{x^2}{a^2}}} dx, x, \frac{1}{x}\right)
 \end{aligned}$$

$$= -\frac{1}{4ax^4} - \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{4x^3} - \frac{a^2\sqrt{1 + \frac{1}{a^2x^2}}}{8x} + \frac{1}{8}a^3\operatorname{csch}^{-1}(ax)$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.82

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^4} dx = \frac{-2 - a\sqrt{1 + \frac{1}{a^2x^2}}x(2 + a^2x^2) + a^4x^4\operatorname{arcsinh}\left(\frac{1}{ax}\right)}{8ax^4}$$

[In] Integrate[E^ArcCsch[a*x]/x^4,x]

[Out] (-2 - a*Sqrt[1 + 1/(a^2*x^2)]*x*(2 + a^2*x^2) + a^4*x^4*ArcSinh[1/(a*x)])/(8*a*x^4)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 172 vs. 2(53) = 106.

Time = 0.06 (sec) , antiderivative size = 173, normalized size of antiderivative = 2.66

method	result	size
default	$\frac{\sqrt{\frac{a^2x^2+1}{a^2x^2}} a^2 \left(\left(\frac{a^2x^2+1}{a^2} \right)^{\frac{3}{2}} \sqrt{\frac{1}{a^2}} a^2 x^2 - \sqrt{\frac{a^2x^2+1}{a^2}} \sqrt{\frac{1}{a^2}} a^2 x^4 + \ln \left(\frac{2\sqrt{\frac{1}{a^2}} \sqrt{\frac{a^2x^2+1}{a^2}} a^2 + 2}{a^2 x} \right) x^4 - 2 \left(\frac{a^2x^2+1}{a^2} \right)^{\frac{3}{2}} \sqrt{\frac{1}{a^2}} \right)}{8x^3 \sqrt{\frac{a^2x^2+1}{a^2}} \sqrt{\frac{1}{a^2}}} - \frac{1}{4ax^4}$	173

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))/x^4,x,method=_RETURNVERBOSE)

[Out] 1/8*((a^2*x^2+1)/a^2/x^2)^(1/2)/x^3*a^2*((a^2*x^2+1)/a^2)^(3/2)*(1/a^2)^(1/2)*a^2*x^2-((a^2*x^2+1)/a^2)^(1/2)*(1/a^2)^(1/2)*a^2*x^4+ln(2*((1/a^2)^(1/2))*((a^2*x^2+1)/a^2)^(1/2)*a^2+1)/x/a^2)*x^4-2*((a^2*x^2+1)/a^2)^(3/2)*(1/a^2)^(1/2))/((a^2*x^2+1)/a^2)^(1/2)/(1/a^2)^(1/2)-1/4/a/x^4

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 113 vs. 2(53) = 106.

Time = 0.26 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.74

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^4} dx = \frac{a^4x^4 \log \left(ax\sqrt{\frac{a^2x^2+1}{a^2x^2}} - ax + 1 \right) - a^4x^4 \log \left(ax\sqrt{\frac{a^2x^2+1}{a^2x^2}} - ax - 1 \right) - (a^3x^3 + 2ax)\sqrt{\frac{a^2x^2+1}{a^2x^2}} - 2}{8ax^4}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))/x^4,x, algorithm="fricas")

[Out] 1/8*(a^4*x^4*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x + 1) - a^4*x^4*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x - 1) - (a^3*x^3 + 2*a*x)*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - 2)/(a*x^4)

Sympy [A] (verification not implemented)

Time = 1.03 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.46

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^4} dx = \begin{cases} -a \left(\begin{cases} -\frac{a^2 \log\left(2\sqrt{1+\frac{1}{a^2x^2}}\sqrt{\frac{1}{a^2}+\frac{2}{a^2x}}\right) + \sqrt{1+\frac{1}{a^2x^2}}\left(\frac{a^2}{8x} + \frac{1}{4x^3}\right)}{8\sqrt{\frac{1}{a^2}}} & \text{for } \frac{1}{a^2} \neq 0 \\ \frac{1}{3x^3} & \text{otherwise} \end{cases} \right)_{-\frac{1}{4x^4}} & \text{for } a \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))/x**4,x)

[Out] Piecewise(((-a*Piecewise((-a**2*log(2*sqrt(1 + 1/(a**2*x**2)))*sqrt(a**(-2)) + 2/(a**2*x))/(8*sqrt(a**(-2))) + sqrt(1 + 1/(a**2*x**2))*(a**2/(8*x) + 1/(4*x**3)), Ne(a**(-2), 0)), (1/(3*x**3), True)) - 1/(4*x**4))/a, Ne(a, 0)), (0, True))

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 129 vs. 2(53) = 106.

Time = 0.23 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.98

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^4} dx = \frac{1}{16} a^3 \log\left(ax\sqrt{\frac{1}{a^2x^2} + 1} + 1\right) - \frac{1}{16} a^3 \log\left(ax\sqrt{\frac{1}{a^2x^2} + 1} - 1\right) - \frac{a^6x^3\left(\frac{1}{a^2x^2} + 1\right)^{\frac{3}{2}} + a^4x\sqrt{\frac{1}{a^2x^2} + 1}}{8\left(a^4x^4\left(\frac{1}{a^2x^2} + 1\right)^2 - 2a^2x^2\left(\frac{1}{a^2x^2} + 1\right) + 1\right)} - \frac{1}{4ax^4}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))/x^4,x, algorithm="maxima")

[Out] 1/16*a^3*log(a*x*sqrt(1/(a^2*x^2) + 1) + 1) - 1/16*a^3*log(a*x*sqrt(1/(a^2*x^2) + 1) - 1) - 1/8*(a^6*x^3*(1/(a^2*x^2) + 1)^(3/2) + a^4*x*sqrt(1/(a^2*x^2) + 1))/(a^4*x^4*(1/(a^2*x^2) + 1)^2 - 2*a^2*x^2*(1/(a^2*x^2) + 1) + 1) - 1/4/(a*x^4)

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.58

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^4} dx$$

$$= \frac{a^6 |a| \log(\sqrt{a^2 x^2 + 1} + 1) \operatorname{sgn}(x) - a^6 |a| \log(\sqrt{a^2 x^2 + 1} - 1) \operatorname{sgn}(x) - \frac{2 \left((a^2 x^2 + 1)^{\frac{3}{2}} a^6 |a| \operatorname{sgn}(x) + \sqrt{a^2 x^2 + 1} a^6 |a| \operatorname{sgn}(x) \right)}{a^4 x^4}}{16 a^4}$$

```
[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))/x^4,x, algorithm="giac")
```

```
[Out] 1/16*(a^6*abs(a)*log(sqrt(a^2*x^2 + 1) + 1)*sgn(x) - a^6*abs(a)*log(sqrt(a^2*x^2 + 1) - 1)*sgn(x) - 2*((a^2*x^2 + 1)^(3/2)*a^6*abs(a)*sgn(x) + sqrt(a^2*x^2 + 1)*a^6*abs(a)*sgn(x) + 2*a^7)/(a^4*x^4)/a^4
```

Mupad [B] (verification not implemented)

Time = 5.62 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.94

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^4} dx = \frac{\operatorname{asinh}\left(\frac{\sqrt{\frac{1}{a^2}}}{x}\right)}{8 \left(\frac{1}{a^2}\right)^{3/2}} - \frac{\sqrt{\frac{1}{a^2 x^2} + 1}}{4 x^3} - \frac{1}{4 a x^4} - \frac{a^2 \sqrt{\frac{1}{a^2 x^2} + 1}}{8 x}$$

```
[In] int(((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))/x^4,x)
```

```
[Out] asinh((1/a^2)^(1/2)/x)/(8*(1/a^2)^(3/2)) - (1/(a^2*x^2) + 1)^(1/2)/(4*x^3) - 1/(4*a*x^4) - (a^2*(1/(a^2*x^2) + 1)^(1/2))/(8*x)
```

3.36 $\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^5} dx$

Optimal result	251
Rubi [A] (verified)	251
Mathematica [A] (verified)	252
Maple [A] (verified)	253
Fricas [A] (verification not implemented)	253
Sympy [A] (verification not implemented)	253
Maxima [A] (verification not implemented)	254
Giac [B] (verification not implemented)	254
Mupad [B] (verification not implemented)	254

Optimal result

Integrand size = 10, antiderivative size = 51

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^5} dx = \frac{1}{3}a^4 \left(1 + \frac{1}{a^2x^2}\right)^{3/2} - \frac{1}{5}a^4 \left(1 + \frac{1}{a^2x^2}\right)^{5/2} - \frac{1}{5ax^5}$$

[Out] $1/3*a^4*(1+1/a^2/x^2)^{(3/2)}-1/5*a^4*(1+1/a^2/x^2)^{(5/2)}-1/5/a/x^5$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {6471, 30, 272, 45}

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^5} dx = -\frac{1}{5}a^4 \left(\frac{1}{a^2x^2} + 1\right)^{5/2} + \frac{1}{3}a^4 \left(\frac{1}{a^2x^2} + 1\right)^{3/2} - \frac{1}{5ax^5}$$

[In] Int[E^ArcCsch[a*x]/x^5,x]

[Out] $(a^4*(1 + 1/(a^2*x^2))^{(3/2)})/3 - (a^4*(1 + 1/(a^2*x^2))^{(5/2)})/5 - 1/(5*a*x^5)$

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 45

Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},

$x]$ && NeQ[$b*c - a*d, 0]$ && IGtQ[$m, 0]$ && (!IntegerQ[n] || (EqQ[$c, 0]$ && LeQ[$7*m + 4*n + 4, 0]$) || LtQ[$9*m + 5*(n + 1), 0]$ || GtQ[$m + n + 2, 0]$)

Rule 272

Int[($x_$)^($m_$)*($(a_ + (b_)*(x_)$)^($n_$))^($p_$), $x_Symbol]$:= Dist[$1/n$, Subst[
Int[$x^((Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x]$, $x, x^n]$, $x]$ /; FreeQ[{ a, b ,
 m, n, p }, $x]$ && IntegerQ[Simplify[($m + 1$)/ n]]

Rule 6471

Int[E^ArcCsch[($a_$)*($x_$)^($p_$)]*($x_$)^($m_$), $x_Symbol]$:= Dist[$1/a$, Int[$x^(m$
 $- p)$, $x]$, $x]$ + Int[$x^m*sqrt[1 + 1/(a^2*x^(2*p))]$, $x]$ /; FreeQ[{ a, m, p }, x
]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \frac{1}{x^6} dx}{a} + \int \frac{\sqrt{1 + \frac{1}{a^2 x^2}}}{x^5} dx \\ &= -\frac{1}{5ax^5} - \frac{1}{2} \text{Subst}\left(\int x \sqrt{1 + \frac{x}{a^2}} dx, x, \frac{1}{x^2}\right) \\ &= -\frac{1}{5ax^5} - \frac{1}{2} \text{Subst}\left(\int \left(-a^2 \sqrt{1 + \frac{x}{a^2}} + a^2 \left(1 + \frac{x}{a^2}\right)^{3/2}\right) dx, x, \frac{1}{x^2}\right) \\ &= \frac{1}{3} a^4 \left(1 + \frac{1}{a^2 x^2}\right)^{3/2} - \frac{1}{5} a^4 \left(1 + \frac{1}{a^2 x^2}\right)^{5/2} - \frac{1}{5ax^5} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.90

$$\int \frac{e^{\text{csch}^{-1}(ax)}}{x^5} dx = \frac{-3 + a \sqrt{1 + \frac{1}{a^2 x^2}} x (-3 - a^2 x^2 + 2a^4 x^4)}{15ax^5}$$

[In] Integrate[E^ArcCsch[a*x]/x^5,x]

[Out] (-3 + a*Sqrt[1 + 1/(a^2*x^2)]*x*(-3 - a^2*x^2 + 2*a^4*x^4))/(15*a*x^5)

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.02

method	result	size
default	$\frac{\sqrt{\frac{a^2x^2+1}{a^2x^2}}(a^2x^2+1)(2a^2x^2-3)}{15x^4} - \frac{1}{5ax^5}$	52
trager	$-\frac{1}{5x^5} + \frac{a(2a^4x^4 - a^2x^2 - 3)\sqrt{-\frac{a^2x^2-1}{a^2x^2}}}{15x^4 a}$	55

[In] `int((1/a/x+(1+1/a^2/x^2)^(1/2))/x^5,x,method=_RETURNVERBOSE)`

[Out] $1/15*((a^2*x^2+1)/a^2/x^2)^(1/2)/x^4*(a^2*x^2+1)*(2*a^2*x^2-3)-1/5/a/x^5$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.14

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^5} dx = \frac{2a^5x^5 + (2a^5x^5 - a^3x^3 - 3ax)\sqrt{\frac{a^2x^2+1}{a^2x^2}} - 3}{15ax^5}$$

[In] `integrate((1/a/x+(1+1/a^2/x^2)^(1/2))/x^5,x, algorithm="fricas")`

[Out] $1/15*(2*a^5*x^5 + (2*a^5*x^5 - a^3*x^3 - 3*a*x)*\operatorname{sqrt}((a^2*x^2 + 1)/(a^2*x^2)) - 3)/(a*x^5)$

Sympy [A] (verification not implemented)

Time = 1.11 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.14

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^5} dx = \begin{cases} -a \left(\begin{cases} \sqrt{1 + \frac{1}{a^2x^2}} \left(-\frac{2a^4}{15} + \frac{a^2}{15x^2} + \frac{1}{5x^4} \right) & \text{for } \frac{1}{a^2} \neq 0 \\ \frac{1}{4x^4} & \text{otherwise} \end{cases} \right) - \frac{1}{5x^5} & \text{for } a \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

[In] `integrate((1/a/x+(1+1/a**2/x**2)**(1/2))/x**5,x)`

[Out] `Piecewise(((-a*Piecewise((sqrt(1 + 1/(a**2*x**2)))*(-2*a**4/15 + a**2/(15*x**2) + 1/(5*x**4)), Ne(a**(-2), 0)), (1/(4*x**4), True)) - 1/(5*x**5))/a, Ne(a, 0)), (0, True))`

Maxima [A] (verification not implemented)

none

Time = 0.23 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.80

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^5} dx = -\frac{1}{5} a^4 \left(\frac{1}{a^2 x^2} + 1 \right)^{\frac{5}{2}} + \frac{1}{3} a^4 \left(\frac{1}{a^2 x^2} + 1 \right)^{\frac{3}{2}} - \frac{1}{5 a x^5}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))/x^5,x, algorithm="maxima")

[Out] -1/5*a^4*(1/(a^2*x^2) + 1)^(5/2) + 1/3*a^4*(1/(a^2*x^2) + 1)^(3/2) - 1/5/(a*x^5)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 124 vs. 2(41) = 82.

Time = 0.29 (sec) , antiderivative size = 124, normalized size of antiderivative = 2.43

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^5} dx = \frac{4 \left(15 (x|a| - \sqrt{a^2 x^2 + 1})^6 a^4 \operatorname{sgn}(x) + 5 (x|a| - \sqrt{a^2 x^2 + 1})^4 a^4 \operatorname{sgn}(x) + 5 (x|a| - \sqrt{a^2 x^2 + 1})^2 a^4 \operatorname{sgn}(x) - a^4 \operatorname{sgn}(x) \right)}{15 \left((x|a| - \sqrt{a^2 x^2 + 1})^2 - 1 \right)^5} - \frac{1}{5 a x^5}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))/x^5,x, algorithm="giac")

[Out] 4/15*(15*(x*abs(a) - sqrt(a^2*x^2 + 1))^6*a^4*sgn(x) + 5*(x*abs(a) - sqrt(a^2*x^2 + 1))^4*a^4*sgn(x) + 5*(x*abs(a) - sqrt(a^2*x^2 + 1))^2*a^4*sgn(x) - a^4*sgn(x))/((x*abs(a) - sqrt(a^2*x^2 + 1))^2 - 1)^5 - 1/5/(a*x^5)

Mupad [B] (verification not implemented)

Time = 5.17 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.20

$$\int \frac{e^{\operatorname{csch}^{-1}(ax)}}{x^5} dx = \frac{2 a^4 \sqrt{\frac{1}{a^2 x^2} + 1}}{15} - \frac{x \sqrt{\frac{1}{a^2 x^2} + 1}}{5 x^5} + \frac{1}{5 a} - \frac{a^2 \sqrt{\frac{1}{a^2 x^2} + 1}}{15 x^2}$$

[In] int(((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))/x^5,x)

[Out] (2*a^4*(1/(a^2*x^2) + 1)^(1/2))/15 - ((x*(1/(a^2*x^2) + 1)^(1/2))/5 + 1/(5*a))/x^5 - (a^2*(1/(a^2*x^2) + 1)^(1/2))/(15*x^2)

3.37 $\int e^{\operatorname{csch}^{-1}(ax^2)} x^m dx$

Optimal result	255
Rubi [A] (verified)	255
Mathematica [A] (verified)	256
Maple [F]	257
Fricas [F]	257
Sympy [A] (verification not implemented)	257
Maxima [F(-2)]	257
Giac [F(-2)]	258
Mupad [F(-1)]	258

Optimal result

Integrand size = 12, antiderivative size = 59

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^m dx = -\frac{x^{-1+m}}{a(1-m)} + \frac{x^{1+m} \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, \frac{1}{4}(-1-m), \frac{3-m}{4}, -\frac{1}{a^2x^4}\right)}{1+m}$$

[Out] $-x^{(-1+m)}/a/(1-m)+x^{(1+m)}*\operatorname{hypergeom}([-1/2, -1/4-1/4*m], [3/4-1/4*m], -1/a^2/x^4)/(1+m)$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6471, 30, 346, 371}

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^m dx = \frac{x^{m+1} \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, \frac{1}{4}(-m-1), \frac{3-m}{4}, -\frac{1}{a^2x^4}\right)}{m+1} - \frac{x^{m-1}}{a(1-m)}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcCsch}[a*x^2]}*x^m, x]$

[Out] $-(x^{(-1+m)}/(a*(1-m))) + (x^{(1+m)}*\operatorname{Hypergeometric2F1}[-1/2, (-1-m)/4, (3-m)/4, -(1/(a^2*x^4))])/(1+m)$

Rule 30

$\operatorname{Int}[(x_)^{(m_.)}, x_Symbol] := \operatorname{Simp}[x^{(m+1)}/(m+1), x] /; \operatorname{FreeQ}[m, x] \ \&\& \ \operatorname{NeQ}[m, -1]$

Rule 346

$\operatorname{Int}[((c_.)*(x_))^{(m_.)}*((a_.) + (b_.)*(x_)^{(n_.)})^{(p_.)}, x_Symbol] := \operatorname{Dist}[(-c^{(-1)})*(c*x)^{(m+1)}*(1/x)^{(m+1)}, \operatorname{Subst}[\operatorname{Int}[(a + b/x^n)^p/x^{(m+2)}, x], x]$

, 1/x], x] /; FreeQ[{a, b, c, m, p}, x] && ILtQ[n, 0] && !RationalQ[m]

Rule 371

Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 6471

Int[E^ArcCsch[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Dist[1/a, Int[x^(m - p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int x^{-2+m} dx}{a} + \int \sqrt{1 + \frac{1}{a^2 x^4}} x^m dx \\ &= -\frac{x^{-1+m}}{a(1-m)} - \left(\left(\frac{1}{x} \right)^m x^m \right) \text{Subst} \left(\int x^{-2-m} \sqrt{1 + \frac{x^4}{a^2}} dx, x, \frac{1}{x} \right) \\ &= -\frac{x^{-1+m}}{a(1-m)} + \frac{x^{1+m} \text{Hypergeometric2F1} \left(-\frac{1}{2}, \frac{1}{4}(-1-m), \frac{3-m}{4}, -\frac{1}{a^2 x^4} \right)}{1+m} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.93

$$\int e^{\text{csch}^{-1}(ax^2)} x^m dx = x^{-1+m} \left(\frac{1}{a(-1+m)} + \frac{x^2 \text{Hypergeometric2F1} \left(-\frac{1}{2}, -\frac{1}{4} - \frac{m}{4}, \frac{3}{4} - \frac{m}{4}, -\frac{1}{a^2 x^4} \right)}{1+m} \right)$$

[In] Integrate[E^ArcCsch[a*x^2]*x^m,x]

[Out] x^(-1 + m)*(1/(a*(-1 + m)) + (x^2*Hypergeometric2F1[-1/2, -1/4 - m/4, 3/4 - m/4, -(1/(a^2*x^4))]))/(1 + m)

Maple [F]

$$\int \left(\frac{1}{ax^2} + \sqrt{1 + \frac{1}{a^2x^4}} \right) x^m dx$$

[In] int((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^m,x)

[Out] int((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^m,x)

Fricas [F]

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^m dx = \int x^m \left(\sqrt{\frac{1}{a^2x^4} + 1} + \frac{1}{ax^2} \right) dx$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^m,x, algorithm="fricas")

[Out] integral((a*x^2*x^m*sqrt((a^2*x^4 + 1)/(a^2*x^4)) + x^m)/(a*x^2), x)

Sympy [A] (verification not implemented)

Time = 2.73 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.20

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^m dx = -\frac{x^{m+1} \Gamma\left(-\frac{m}{4} - \frac{1}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{1}{2}, -\frac{m}{4} - \frac{1}{4} \\ \frac{3}{4} - \frac{m}{4} \end{matrix} \middle| \frac{e^{i\pi}}{a^2x^4}\right)}{4\Gamma\left(\frac{3}{4} - \frac{m}{4}\right)} + \frac{\begin{cases} \frac{x^m}{mx-x} & \text{for } m \neq 1 \\ \frac{x^m \log(x)}{x} & \text{otherwise} \end{cases}}{a}$$

[In] integrate((1/a/x**2+(1+1/a**2/x**4)**(1/2))*x**m,x)

[Out] -x**(m + 1)*gamma(-m/4 - 1/4)*hyper((-1/2, -m/4 - 1/4), (3/4 - m/4,), exp_polar(I*pi)/(a**2*x**4))/(4*gamma(3/4 - m/4)) + Piecewise((x**m/(m*x - x), Ne(m, 1)), (x**m*log(x)/x, True))/a

Maxima [F(-2)]

Exception generated.

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^m dx = \text{Exception raised: ValueError}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^m,x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(m-2>0)', see 'assume?' for more details)Is

Giac [F(-2)]

Exception generated.

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^m dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^m,x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
 UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
 dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^m dx = \int x^m \left(\sqrt{\frac{1}{a^2 x^4} + 1} + \frac{1}{a x^2} \right) dx$$

[In] int(x^m*((1/(a^2*x^4) + 1)^(1/2) + 1/(a*x^2)),x)

[Out] int(x^m*((1/(a^2*x^4) + 1)^(1/2) + 1/(a*x^2)), x)

3.38 $\int e^{\operatorname{csch}^{-1}(ax^2)} x^4 dx$

Optimal result	259
Rubi [A] (verified)	260
Mathematica [C] (verified)	262
Maple [C] (verified)	263
Fricas [A] (verification not implemented)	263
Sympy [C] (verification not implemented)	263
Maxima [F]	264
Giac [F]	264
Mupad [F(-1)]	264

Optimal result

Integrand size = 12, antiderivative size = 202

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^4 dx = -\frac{2\sqrt{1+\frac{1}{a^2x^4}}}{5a^2\left(a+\frac{1}{x^2}\right)x} + \frac{2\sqrt{1+\frac{1}{a^2x^4}}x}{5a^2} + \frac{x^3}{3a} + \frac{1}{5}\sqrt{1+\frac{1}{a^2x^4}}x^5$$

$$+ \frac{2\sqrt{\frac{a^2+\frac{1}{x^4}}{\left(a+\frac{1}{x^2}\right)^2}}\left(a+\frac{1}{x^2}\right)E\left(2\cot^{-1}\left(\sqrt{ax}\right)\left|\frac{1}{2}\right.\right)}{5a^{7/2}\sqrt{1+\frac{1}{a^2x^4}}}$$

$$- \frac{\sqrt{\frac{a^2+\frac{1}{x^4}}{\left(a+\frac{1}{x^2}\right)^2}}\left(a+\frac{1}{x^2}\right)\operatorname{EllipticF}\left(2\cot^{-1}\left(\sqrt{ax}\right),\frac{1}{2}\right)}{5a^{7/2}\sqrt{1+\frac{1}{a^2x^4}}}$$

```
[Out] 1/3*x^3/a-2/5*(1+1/a^2/x^4)^(1/2)/a^2/(a+1/x^2)/x+2/5*x*(1+1/a^2/x^4)^(1/2)
/a^2+1/5*x^5*(1+1/a^2/x^4)^(1/2)+2/5*(a+1/x^2)*(cos(2*arccot(x*a^(1/2))))^2
^(1/2)/cos(2*arccot(x*a^(1/2)))*EllipticE(sin(2*arccot(x*a^(1/2))),1/2*2^(1
/2))*((a^2+1/x^4)/(a+1/x^2)^2)^(1/2)/a^(7/2)/(1+1/a^2/x^4)^(1/2)-1/5*(a+1/x
^2)*(cos(2*arccot(x*a^(1/2))))^2^(1/2)/cos(2*arccot(x*a^(1/2)))*EllipticF(s
in(2*arccot(x*a^(1/2))),1/2*2^(1/2))*((a^2+1/x^4)/(a+1/x^2)^2)^(1/2)/a^(7/2
)/(1+1/a^2/x^4)^(1/2)
```

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {6471, 30, 342, 283, 331, 311, 226, 1210}

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^4 dx = \frac{2x\sqrt{\frac{1}{a^2x^4} + 1}}{5a^2} + \frac{1}{5}x^5\sqrt{\frac{1}{a^2x^4} + 1} - \frac{2\sqrt{\frac{1}{a^2x^4} + 1}}{5a^2x\left(a + \frac{1}{x^2}\right)} - \frac{\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}}\left(a + \frac{1}{x^2}\right)\operatorname{EllipticF}\left(2\cot^{-1}\left(\sqrt{ax}\right), \frac{1}{2}\right)}{5a^{7/2}\sqrt{\frac{1}{a^2x^4} + 1}} + \frac{2\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}}\left(a + \frac{1}{x^2}\right)E\left(2\cot^{-1}\left(\sqrt{ax}\right)\left|\frac{1}{2}\right.\right)}{5a^{7/2}\sqrt{\frac{1}{a^2x^4} + 1}} + \frac{x^3}{3a}$$

[In] Int[E^ArcCsch[a*x^2]*x^4,x]

[Out] (-2*Sqrt[1 + 1/(a^2*x^4)])/(5*a^2*(a + x^(-2))*x) + (2*Sqrt[1 + 1/(a^2*x^4)]*x)/(5*a^2) + x^3/(3*a) + (Sqrt[1 + 1/(a^2*x^4)]*x^5)/5 + (2*Sqrt[(a^2 + x^(-4))/(a + x^(-2))^2]*(a + x^(-2))*EllipticE[2*ArcCot[Sqrt[a]*x], 1/2])/(5*a^(7/2)*Sqrt[1 + 1/(a^2*x^4)]) - (Sqrt[(a^2 + x^(-4))/(a + x^(-2))^2]*(a + x^(-2))*EllipticF[2*ArcCot[Sqrt[a]*x], 1/2])/(5*a^(7/2)*Sqrt[1 + 1/(a^2*x^4)])

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 226

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 283

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^p/(c*(m + 1))), x] - Dist[b*n*(p/(c^n*(m + 1))), Int[(c*x)^(m + n)*(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && GtQ[p, 0] && LtQ[m, -1] && !ILtQ[(m + n*p + n + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 311

Int[(x_)^2/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, Dist[1/q, Int[1/Sqrt[a + b*x^4], x], x] - Dist[1/q, Int[(1 - q*x^2)/Sqrt[a + b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 331

Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Dist[b*(m + n*(p + 1) + 1)/(a*c^n*(m + 1)), Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 342

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]

Rule 1210

Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + c*x^4])*EllipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]

Rule 6471

Int[E^ArcCsch[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Dist[1/a, Int[x^(m - p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int x^2 dx}{a} + \int \sqrt{1 + \frac{1}{a^2 x^4}} x^4 dx \\
 &= \frac{x^3}{3a} - \text{Subst} \left(\int \frac{\sqrt{1 + \frac{x^4}{a^2}}}{x^6} dx, x, \frac{1}{x} \right) \\
 &= \frac{x^3}{3a} + \frac{1}{5} \sqrt{1 + \frac{1}{a^2 x^4}} x^5 - \frac{2 \text{Subst} \left(\int \frac{1}{x^2 \sqrt{1 + \frac{x^4}{a^2}}} dx, x, \frac{1}{x} \right)}{5a^2}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{2\sqrt{1 + \frac{1}{a^2x^4}}}{5a^2} + \frac{x^3}{3a} + \frac{1}{5}\sqrt{1 + \frac{1}{a^2x^4}}x^5 - \frac{2\text{Subst}\left(\int \frac{x^2}{\sqrt{1+\frac{x^4}{a^2}}} dx, x, \frac{1}{x}\right)}{5a^4} \\
&= \frac{2\sqrt{1 + \frac{1}{a^2x^4}}}{5a^2} + \frac{x^3}{3a} + \frac{1}{5}\sqrt{1 + \frac{1}{a^2x^4}}x^5 \\
&\quad - \frac{2\text{Subst}\left(\int \frac{1}{\sqrt{1+\frac{x^4}{a^2}}} dx, x, \frac{1}{x}\right)}{5a^3} + \frac{2\text{Subst}\left(\int \frac{1-\frac{x^2}{a}}{\sqrt{1+\frac{x^4}{a^2}}} dx, x, \frac{1}{x}\right)}{5a^3} \\
&= -\frac{2\sqrt{1 + \frac{1}{a^2x^4}}}{5a^2\left(a + \frac{1}{x^2}\right)x} + \frac{2\sqrt{1 + \frac{1}{a^2x^4}}}{5a^2} + \frac{x^3}{3a} + \frac{1}{5}\sqrt{1 + \frac{1}{a^2x^4}}x^5 \\
&\quad + \frac{2\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}}\left(a + \frac{1}{x^2}\right)E\left(2\cot^{-1}\left(\sqrt{ax}\right)\left|\frac{1}{2}\right.\right)}{5a^{7/2}\sqrt{1 + \frac{1}{a^2x^4}}} \\
&\quad - \frac{\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}}\left(a + \frac{1}{x^2}\right)\text{EllipticF}\left(2\cot^{-1}\left(\sqrt{ax}\right), \frac{1}{2}\right)}{5a^{7/2}\sqrt{1 + \frac{1}{a^2x^4}}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.17 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.55

$$\begin{aligned}
&\int e^{\text{csch}^{-1}(ax^2)} x^4 dx \\
&= \frac{4\sqrt{2}e^{-\text{csch}^{-1}(ax^2)}\left(\frac{e^{\text{csch}^{-1}(ax^2)}}{-1+e^{2\text{csch}^{-1}(ax^2)}}\right)^{5/2}x^5\left(-4+7e^{2\text{csch}^{-1}(ax^2)}+4\left(1-e^{2\text{csch}^{-1}(ax^2)}\right)^{5/2}\text{Hypergeometric2F1}\right)}{21(ax^2)^{5/2}}
\end{aligned}$$

[In] Integrate[E^ArcCsch[a*x^2]*x^4,x]

[Out] (4*Sqrt[2]*(E^ArcCsch[a*x^2]/(-1 + E^(2*ArcCsch[a*x^2])))^(5/2)*x^5*(-4 + 7*E^(2*ArcCsch[a*x^2]) + 4*(1 - E^(2*ArcCsch[a*x^2]))^(5/2)*Hypergeometric2F1[3/4, 7/2, 7/4, E^(2*ArcCsch[a*x^2])])/(21*E^ArcCsch[a*x^2]*(a*x^2)^(5/2))

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.96 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.74

method	result
default	$\frac{\sqrt{\frac{x^4 a^2 + 1}{x^4 a^2}} x^2 \left(\sqrt{ia} a^3 x^7 + x^3 a \sqrt{ia} + 2i \sqrt{-ia x^2 + 1} \sqrt{ia x^2 + 1} \operatorname{EllipticF}(x \sqrt{ia}, i) - 2i \sqrt{-ia x^2 + 1} \sqrt{ia x^2 + 1} \operatorname{EllipticE}(x \sqrt{ia}, i) \right)}{5a(x^4 a^2 + 1) \sqrt{ia}} + \frac{x^3}{3a}$

[In] int((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^4,x,method=_RETURNVERBOSE)

[Out] 1/5*((a^2*x^4+1)/x^4/a^2)^(1/2)*x^2*((I*a)^(1/2)*a^3*x^7+x^3*a*(I*a)^(1/2)+2*I*(1-I*a*x^2)^(1/2)*(1+I*a*x^2)^(1/2)*EllipticF(x*(I*a)^(1/2),I)-2*I*(1-I*a*x^2)^(1/2)*(1+I*a*x^2)^(1/2)*EllipticE(x*(I*a)^(1/2),I))/a/(a^2*x^4+1)/(I*a)^(1/2)+1/3*x^3/a

Fricas [A] (verification not implemented)

none

Time = 0.09 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.44

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^4 dx$$

$$= \frac{5ax^3 + 6\left(-\frac{1}{a^2}\right)^{\frac{3}{4}} E\left(\arcsin\left(\frac{\left(-\frac{1}{a^2}\right)^{\frac{1}{4}}}{x}\right) \mid -1\right) - 6\left(-\frac{1}{a^2}\right)^{\frac{3}{4}} F\left(\arcsin\left(\frac{\left(-\frac{1}{a^2}\right)^{\frac{1}{4}}}{x}\right) \mid -1\right) + 3(a^2x^5 + 2x)\sqrt{\frac{a^2x^5}{a^2}}}{15a^2}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^4,x, algorithm="fricas")

[Out] 1/15*(5*a*x^3 + 6*(-1/a^2)^(3/4)*elliptic_e(arcsin((-1/a^2)^(1/4)/x), -1) - 6*(-1/a^2)^(3/4)*elliptic_f(arcsin((-1/a^2)^(1/4)/x), -1) + 3*(a^2*x^5 + 2*x)*sqrt((a^2*x^4 + 1)/(a^2*x^4)))/a^2

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.17 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.24

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^4 dx = -\frac{x^5 \Gamma\left(-\frac{5}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{5}{4}, -\frac{1}{2} \\ -\frac{1}{4} \end{matrix} \middle| \frac{e^{i\pi}}{a^2 x^4}\right)}{4\Gamma\left(-\frac{1}{4}\right)} + \frac{x^3}{3a}$$

[In] integrate((1/a/x**2+(1+1/a**2/x**4)**(1/2))*x**4,x)

[Out] $-x^{5}\gamma(-5/4)\text{hyper}((-5/4, -1/2), (-1/4,), \exp_{\text{polar}}(I\pi)/(a^{2}x^{4}))/ (4\gamma(-1/4)) + x^{3}/(3a)$

Maxima [F]

$$\int e^{\text{csch}^{-1}(ax^2)} x^4 dx = \int x^4 \left(\sqrt{\frac{1}{a^2 x^4} + 1} + \frac{1}{ax^2} \right) dx$$

[In] `integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^4,x, algorithm="maxima")`

[Out] $1/3x^3/a + \text{integrate}(\text{sqrt}(a^2x^4 + 1)*x^2, x)/a$

Giac [F]

$$\int e^{\text{csch}^{-1}(ax^2)} x^4 dx = \int x^4 \left(\sqrt{\frac{1}{a^2 x^4} + 1} + \frac{1}{ax^2} \right) dx$$

[In] `integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^4,x, algorithm="giac")`

[Out] `integrate(x^4*(sqrt(1/(a^2*x^4) + 1) + 1/(a*x^2)), x)`

Mupad [F(-1)]

Timed out.

$$\int e^{\text{csch}^{-1}(ax^2)} x^4 dx = \int x^4 \left(\sqrt{\frac{1}{a^2 x^4} + 1} + \frac{1}{ax^2} \right) dx$$

[In] `int(x^4*((1/(a^2*x^4) + 1)^(1/2) + 1/(a*x^2)),x)`

[Out] `int(x^4*((1/(a^2*x^4) + 1)^(1/2) + 1/(a*x^2)), x)`

3.39 $\int e^{\operatorname{csch}^{-1}(ax^2)} x^3 dx$

Optimal result	265
Rubi [A] (verified)	265
Mathematica [A] (verified)	267
Maple [B] (verified)	267
Fricas [A] (verification not implemented)	267
Sympy [A] (verification not implemented)	268
Maxima [A] (verification not implemented)	268
Giac [A] (verification not implemented)	268
Mupad [B] (verification not implemented)	269

Optimal result

Integrand size = 12, antiderivative size = 52

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^3 dx = \frac{x^2}{2a} + \frac{1}{4} \sqrt{1 + \frac{1}{a^2 x^4}} x^4 + \frac{\operatorname{arctanh}\left(\sqrt{1 + \frac{1}{a^2 x^4}}\right)}{4a^2}$$

[Out] $1/2*x^2/a+1/4*\operatorname{arctanh}((1+1/a^2/x^4)^{(1/2)})/a^2+1/4*x^4*(1+1/a^2/x^4)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6471, 30, 272, 43, 65, 214}

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^3 dx = \frac{\operatorname{arctanh}\left(\sqrt{\frac{1}{a^2 x^4} + 1}\right)}{4a^2} + \frac{1}{4} x^4 \sqrt{\frac{1}{a^2 x^4} + 1} + \frac{x^2}{2a}$$

[In] `Int[E^ArcCsch[a*x^2]*x^3,x]`

[Out] $x^2/(2*a) + (\operatorname{Sqrt}[1 + 1/(a^2*x^4)]*x^4)/4 + \operatorname{ArcTanh}[\operatorname{Sqrt}[1 + 1/(a^2*x^4)]]/(4*a^2)$

Rule 30

`Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]`

Rule 43

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Dist[d*(n/(b*(m + 1))), I`

```
nt[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !IntegerQ[n] && GtQ[n, 0]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 6471

```
Int[E^ArcCsch[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Dist[1/a, Int[x^(m
- p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \frac{x \, dx}{a} + \int \sqrt{1 + \frac{1}{a^2 x^4}} x^3 \, dx \\
&= \frac{x^2}{2a} - \frac{1}{4} \text{Subst} \left(\int \frac{\sqrt{1 + \frac{x}{a^2}}}{x^2} \, dx, x, \frac{1}{x^4} \right) \\
&= \frac{x^2}{2a} + \frac{1}{4} \sqrt{1 + \frac{1}{a^2 x^4}} x^4 - \frac{\text{Subst} \left(\int \frac{1}{x \sqrt{1 + \frac{x}{a^2}}} \, dx, x, \frac{1}{x^4} \right)}{8a^2} \\
&= \frac{x^2}{2a} + \frac{1}{4} \sqrt{1 + \frac{1}{a^2 x^4}} x^4 - \frac{1}{4} \text{Subst} \left(\int \frac{1}{-a^2 + a^2 x^2} \, dx, x, \sqrt{1 + \frac{1}{a^2 x^4}} \right) \\
&= \frac{x^2}{2a} + \frac{1}{4} \sqrt{1 + \frac{1}{a^2 x^4}} x^4 + \frac{\text{arctanh} \left(\sqrt{1 + \frac{1}{a^2 x^4}} \right)}{4a^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.02

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^3 dx = \frac{ax^2 \left(2 + a\sqrt{1 + \frac{1}{a^2x^4}} \right) + \log \left(\left(1 + \sqrt{1 + \frac{1}{a^2x^4}} \right) x^2 \right)}{4a^2}$$

[In] Integrate[E^ArcCsch[a*x^2]*x^3,x]

[Out] (a*x^2*(2 + a*Sqrt[1 + 1/(a^2*x^4)]*x^2) + Log[(1 + Sqrt[1 + 1/(a^2*x^4)])*x^2])/(4*a^2)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 93 vs. 2(42) = 84.

Time = 0.11 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.81

method	result	size
default	$\frac{\sqrt{\frac{x^4 a^2 + 1}{x^4 a^2}} x^2 \left(x^2 \sqrt{\frac{x^4 a^2 + 1}{a^2}} a^2 + \ln \left(x^2 + \sqrt{\frac{x^4 a^2 + 1}{a^2}} \right) \right)}{4 \sqrt{\frac{x^4 a^2 + 1}{a^2}} a^2} + \frac{x^2}{2a}$	94

[In] int((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^3,x,method=_RETURNVERBOSE)

[Out] 1/4*((a^2*x^4+1)/x^4/a^2)^(1/2)*x^2*(x^2*(1/a^2*(a^2*x^4+1))^(1/2)*a^2+ln(x^2+(1/a^2*(a^2*x^4+1))^(1/2)))/(1/a^2*(a^2*x^4+1))^(1/2)/a^2+1/2*x^2/a

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.35

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^3 dx = \frac{a^2 x^4 \sqrt{\frac{a^2 x^4 + 1}{a^2 x^4}} + 2 a x^2 - \log \left(a x^2 \sqrt{\frac{a^2 x^4 + 1}{a^2 x^4}} - a x^2 \right)}{4 a^2}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^3,x, algorithm="fricas")

[Out] 1/4*(a^2*x^4*sqrt((a^2*x^4 + 1)/(a^2*x^4)) + 2*a*x^2 - log(a*x^2*sqrt((a^2*x^4 + 1)/(a^2*x^4)) - a*x^2))/a^2

Sympy [A] (verification not implemented)

Time = 1.76 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.69

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^3 dx = \frac{x^2 \sqrt{a^2 x^4 + 1}}{4a} + \frac{x^2}{2a} + \frac{\operatorname{asinh}(ax^2)}{4a^2}$$

[In] integrate((1/a/x**2+(1+1/a**2/x**4)**(1/2))*x**3,x)

[Out] x**2*sqrt(a**2*x**4 + 1)/(4*a) + x**2/(2*a) + asinh(a*x**2)/(4*a**2)

Maxima [A] (verification not implemented)

none

Time = 0.23 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.56

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^3 dx = \frac{x^2}{2a} + \frac{\sqrt{\frac{1}{a^2 x^4} + 1}}{4(a^2(\frac{1}{a^2 x^4} + 1) - a^2)} + \frac{\log\left(\sqrt{\frac{1}{a^2 x^4} + 1} + 1\right)}{8a^2} - \frac{\log\left(\sqrt{\frac{1}{a^2 x^4} + 1} - 1\right)}{8a^2}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^3,x, algorithm="maxima")

[Out] 1/2*x^2/a + 1/4*sqrt(1/(a^2*x^4) + 1)/(a^2*(1/(a^2*x^4) + 1) - a^2) + 1/8*log(sqrt(1/(a^2*x^4) + 1) + 1)/a^2 - 1/8*log(sqrt(1/(a^2*x^4) + 1) - 1)/a^2

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.10

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^3 dx = \frac{2ax^2 + \left(\sqrt{a^2 x^4 + 1} x^2 - \frac{\log(-x^2|a| + \sqrt{a^2 x^4 + 1})}{|a|}\right) |a|}{4a^2}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^3,x, algorithm="giac")

[Out] 1/4*(2*a*x^2 + (sqrt(a^2*x^4 + 1)*x^2 - log(-x^2*abs(a) + sqrt(a^2*x^4 + 1))/abs(a))*abs(a))/a^2

Mupad [B] (verification not implemented)

Time = 5.32 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.81

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^3 dx = \frac{\operatorname{atanh}\left(\sqrt{\frac{1}{a^2 x^4} + 1}\right)}{4a^2} + \frac{x^4 \sqrt{\frac{1}{a^2 x^4} + 1}}{4} + \frac{x^2}{2a}$$

[In] int(x^3*((1/(a^2*x^4) + 1)^(1/2) + 1/(a*x^2)),x)

[Out] atanh((1/(a^2*x^4) + 1)^(1/2))/(4*a^2) + (x^4*(1/(a^2*x^4) + 1)^(1/2))/4 + x^2/(2*a)

3.40 $\int e^{\operatorname{csch}^{-1}(ax^2)} x^2 dx$

Optimal result	270
Rubi [A] (verified)	270
Mathematica [C] (verified)	272
Maple [C] (verified)	272
Fricas [A] (verification not implemented)	272
Sympy [C] (verification not implemented)	273
Maxima [F]	273
Giac [F]	273
Mupad [F(-1)]	274

Optimal result

Integrand size = 12, antiderivative size = 86

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^2 dx = \frac{x}{a} + \frac{1}{3} \sqrt{1 + \frac{1}{a^2 x^4}} x^3 - \frac{\sqrt{\frac{a^2 + \frac{1}{x^4}}{(a + \frac{1}{x^2})^2}} (a + \frac{1}{x^2}) \operatorname{EllipticF}(2 \cot^{-1}(\sqrt{ax}), \frac{1}{2})}{3a^{5/2} \sqrt{1 + \frac{1}{a^2 x^4}}}$$

[Out] $x/a + 1/3 * x^3 * (1 + 1/a^2/x^4)^{(1/2)} - 1/3 * (a + 1/x^2) * (\cos(2 * \operatorname{arccot}(x * a^{(1/2)})))^2 * (1/2) / \cos(2 * \operatorname{arccot}(x * a^{(1/2)})) * \operatorname{EllipticF}(\sin(2 * \operatorname{arccot}(x * a^{(1/2)})), 1/2 * 2^{(1/2)}) * ((a^2 + 1/x^4) / (a + 1/x^2)^2)^{(1/2)} / a^{(5/2)} / (1 + 1/a^2/x^4)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {6471, 8, 342, 283, 226}

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^2 dx = \frac{1}{3} x^3 \sqrt{\frac{1}{a^2 x^4} + 1} - \frac{\sqrt{\frac{a^2 + \frac{1}{x^4}}{(a + \frac{1}{x^2})^2}} (a + \frac{1}{x^2}) \operatorname{EllipticF}(2 \cot^{-1}(\sqrt{ax}), \frac{1}{2})}{3a^{5/2} \sqrt{\frac{1}{a^2 x^4} + 1}} + \frac{x}{a}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcCsch}[a * x^2]} * x^2, x]$

[Out] $x/a + (\operatorname{Sqrt}[1 + 1/(a^2 * x^4)] * x^3) / 3 - (\operatorname{Sqrt}[(a^2 + x^{(-4)}) / (a + x^{(-2)})^2] * (a + x^{(-2)}) * \operatorname{EllipticF}[2 * \operatorname{ArcCot}[\operatorname{Sqrt}[a] * x], 1/2]) / (3 * a^{(5/2)} * \operatorname{Sqrt}[1 + 1/(a^2 * x^4)])$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 226

Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 283

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^p/(c*(m + 1))), x] - Dist[b*n*(p/(c^n*(m + 1))), Int[(c*x)^(m + n)*(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && GtQ[p, 0] && LtQ[m, -1] && !ILtQ[(m + n*p + n + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 342

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]

Rule 6471

Int[E^ArcCsch[(a_)*(x_)^(p_)]*(x_)^(m_), x_Symbol] := Dist[1/a, Int[x^(m - p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{1}{a} dx + \int \sqrt{1 + \frac{1}{a^2 x^4}} x^2 dx \\
 &= \frac{x}{a} - \text{Subst} \left(\int \frac{\sqrt{1 + \frac{x^4}{a^2}}}{x^4} dx, x, \frac{1}{x} \right) \\
 &= \frac{x}{a} + \frac{1}{3} \sqrt{1 + \frac{1}{a^2 x^4}} x^3 - \frac{2 \text{Subst} \left(\int \frac{1}{\sqrt{1 + \frac{x^4}{a^2}}} dx, x, \frac{1}{x} \right)}{3a^2} \\
 &= \frac{x}{a} + \frac{1}{3} \sqrt{1 + \frac{1}{a^2 x^4}} x^3 - \frac{\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}} \left(a + \frac{1}{x^2}\right) \text{EllipticF} \left(2 \cot^{-1}(\sqrt{ax}), \frac{1}{2}\right)}{3a^{5/2} \sqrt{1 + \frac{1}{a^2 x^4}}}
 \end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.16 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.31

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^2 dx = \frac{2\sqrt{2}e^{-\operatorname{csch}^{-1}(ax^2)} \left(\frac{e^{\operatorname{csch}^{-1}(ax^2)}}{-1+e^{2\operatorname{csch}^{-1}(ax^2)}} \right)^{3/2} x \left(1 - 2e^{2\operatorname{csch}^{-1}(ax^2)} - \left(1 - e^{2\operatorname{csch}^{-1}(ax^2)} \right)^{3/2} \operatorname{Hypergeometric2F1} \left(\right)}{3a\sqrt{ax^2}}$$

[In] Integrate[E^ArcCsch[a*x^2]*x^2,x]

[Out] (-2*Sqrt[2]*(E^ArcCsch[a*x^2]/(-1 + E^(2*ArcCsch[a*x^2])))^(3/2)*x*(1 - 2*E^(2*ArcCsch[a*x^2]) - (1 - E^(2*ArcCsch[a*x^2]))^(3/2)*Hypergeometric2F1[1/4, 1/2, 5/4, E^(2*ArcCsch[a*x^2])])/(3*a*E^ArcCsch[a*x^2]*Sqrt[a*x^2])

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.54 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.21

method	result	size
default	$\frac{\sqrt{\frac{x^4 a^2 + 1}{x^4 a^2}} x^2 (\sqrt{ia} a^2 x^5 + 2\sqrt{-ia} x^2 + 1} \sqrt{ia} x^2 + 1} {3(x^4 a^2 + 1)\sqrt{ia}} \operatorname{EllipticF}(x\sqrt{ia}, i) + x\sqrt{ia}} + \frac{x}{a}$	104

[In] int((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^2,x,method=_RETURNVERBOSE)

[Out] 1/3*((a^2*x^4+1)/x^4/a^2)^(1/2)*x^2*((I*a)^(1/2)*a^2*x^5+2*(1-I*a*x^2)^(1/2))*(1+I*a*x^2)^(1/2)*EllipticF(x*(I*a)^(1/2),I)+x*(I*a)^(1/2)/(a^2*x^4+1)/(I*a)^(1/2)+x/a

Fricas [A] (verification not implemented)

none

Time = 0.08 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.65

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^2 dx = \frac{ax^3 \sqrt{\frac{a^2 x^4 + 1}{a^2 x^4}} + 2a \left(-\frac{1}{a^2}\right)^{\frac{3}{4}} F\left(\arcsin\left(\frac{\left(-\frac{1}{a^2}\right)^{\frac{1}{4}}}{x}\right) \mid -1\right) + 3x}{3a}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^2,x, algorithm="fricas")

[Out] $\frac{1}{3}*(a*x^3*\sqrt{(a^2*x^4 + 1)/(a^2*x^4)}) + 2*a*(-1/a^2)^{(3/4)}*\text{elliptic_f}(\arcsin((-1/a^2)^{(1/4)}/x), -1) + 3*x)/a$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.48

$$\int e^{\text{csch}^{-1}(ax^2)} x^2 dx = -\frac{x^3 \Gamma(-\frac{3}{4}) {}_2F_1\left(-\frac{3}{4}, -\frac{1}{2} \middle| \frac{e^{i\pi}}{a^2 x^4}\right)}{4 \Gamma(\frac{1}{4})} + \frac{x}{a}$$

[In] `integrate((1/a/x**2+(1+1/a**2/x**4)**(1/2))*x**2,x)`

[Out] $-x**3*\text{gamma}(-3/4)*\text{hyper}((-3/4, -1/2), (1/4,), \text{exp_polar}(I*\text{pi})/(a**2*x**4))/ (4*\text{gamma}(1/4)) + x/a$

Maxima [F]

$$\int e^{\text{csch}^{-1}(ax^2)} x^2 dx = \int x^2 \left(\sqrt{\frac{1}{a^2 x^4} + 1} + \frac{1}{ax^2} \right) dx$$

[In] `integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^2,x, algorithm="maxima")`

[Out] $x/a + \text{integrate}(\text{sqrt}(a^2*x^4 + 1), x)/a$

Giac [F]

$$\int e^{\text{csch}^{-1}(ax^2)} x^2 dx = \int x^2 \left(\sqrt{\frac{1}{a^2 x^4} + 1} + \frac{1}{ax^2} \right) dx$$

[In] `integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x^2,x, algorithm="giac")`

[Out] `integrate(x^2*(sqrt(1/(a^2*x^4) + 1) + 1/(a*x^2)), x)`

Mupad [F(-1)]

Timed out.

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x^2 dx = \int x^2 \left(\sqrt{\frac{1}{a^2 x^4} + 1} + \frac{1}{a x^2} \right) dx$$

```
[In] int(x^2*((1/(a^2*x^4) + 1)^(1/2) + 1/(a*x^2)),x)
```

```
[Out] int(x^2*((1/(a^2*x^4) + 1)^(1/2) + 1/(a*x^2)), x)
```

3.41 $\int e^{\operatorname{csch}^{-1}(ax^2)} x dx$

Optimal result	275
Rubi [A] (verified)	275
Mathematica [A] (verified)	277
Maple [B] (verified)	277
Fricas [B] (verification not implemented)	277
Sympy [A] (verification not implemented)	278
Maxima [B] (verification not implemented)	278
Giac [A] (verification not implemented)	278
Mupad [B] (verification not implemented)	279

Optimal result

Integrand size = 10, antiderivative size = 40

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x dx = \frac{1}{2} \sqrt{1 + \frac{1}{a^2 x^4}} x^2 - \frac{\operatorname{csch}^{-1}(ax^2)}{2a} + \frac{\log(x)}{a}$$

[Out] $-1/2*\operatorname{arccsch}(a*x^2)/a+\ln(x)/a+1/2*x^2*(1+1/a^2/x^4)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6471, 29, 342, 281, 283, 221}

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x dx = \frac{1}{2} x^2 \sqrt{\frac{1}{a^2 x^4} + 1} - \frac{\operatorname{csch}^{-1}(ax^2)}{2a} + \frac{\log(x)}{a}$$

[In] `Int[E^ArcCsch[a*x^2]*x,x]`

[Out] `(Sqrt[1 + 1/(a^2*x^4)]*x^2)/2 - ArcCsch[a*x^2]/(2*a) + Log[x]/a`

Rule 29

`Int[(x_)^(-1), x_Symbol] := Simp[Log[x], x]`

Rule 221

`Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

Rule 281

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m
+ 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x
^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

Rule 283

```
Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*
x)^(m + 1)*((a + b*x^n)^p/(c*(m + 1))), x] - Dist[b*n*(p/(c^n*(m + 1))), In
t[(c*x)^(m + n)*(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[
n, 0] && GtQ[p, 0] && LtQ[m, -1] && !ILtQ[(m + n*p + n + 1)/n, 0] && IntBi
nomialQ[a, b, c, n, m, p, x]
```

Rule 342

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a +
b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && Int
egerQ[m]
```

Rule 6471

```
Int[E^ArcCsch[(a_)*(x_)^(p_)]*(x_)^(m_), x_Symbol] := Dist[1/a, Int[x^(m
- p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\int \frac{1}{x} dx}{a} + \int \sqrt{1 + \frac{1}{a^2 x^4}} x dx \\
&= \frac{\log(x)}{a} - \text{Subst} \left(\int \frac{\sqrt{1 + \frac{x^4}{a^2}}}{x^3} dx, x, \frac{1}{x} \right) \\
&= \frac{\log(x)}{a} - \frac{1}{2} \text{Subst} \left(\int \frac{\sqrt{1 + \frac{x^2}{a^2}}}{x^2} dx, x, \frac{1}{x^2} \right) \\
&= \frac{1}{2} \sqrt{1 + \frac{1}{a^2 x^4}} x^2 + \frac{\log(x)}{a} - \frac{\text{Subst} \left(\int \frac{1}{\sqrt{1 + \frac{x^2}{a^2}}} dx, x, \frac{1}{x^2} \right)}{2a^2} \\
&= \frac{1}{2} \sqrt{1 + \frac{1}{a^2 x^4}} x^2 - \frac{\text{csch}^{-1}(ax^2)}{2a} + \frac{\log(x)}{a}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.05

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x dx = \frac{a\sqrt{1 + \frac{1}{a^2x^4}}x^2 - \operatorname{arcsinh}\left(\frac{1}{ax^2}\right) + \log(ax^2)}{2a}$$

[In] Integrate[E^ArcCsch[a*x^2]*x,x]

[Out] (a*Sqrt[1 + 1/(a^2*x^4)]*x^2 - ArcSinh[1/(a*x^2)] + Log[a*x^2])/(2*a)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 115 vs. 2(34) = 68.

Time = 0.12 (sec) , antiderivative size = 116, normalized size of antiderivative = 2.90

method	result	size
default	$\frac{\sqrt{\frac{x^4 a^2 + 1}{x^4 a^2}} x^2 \left(\sqrt{\frac{1}{a^2}} \sqrt{\frac{x^4 a^2 + 1}{a^2}} a^2 - \ln \left(\frac{2\sqrt{\frac{1}{a^2}} \sqrt{\frac{x^4 a^2 + 1}{a^2}} a^2 + 2}{a^2 x^2} \right) \right)}{2\sqrt{\frac{1}{a^2}} \sqrt{\frac{x^4 a^2 + 1}{a^2}} a^2} + \frac{\ln(x)}{a}$	116

[In] int((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x,x,method=_RETURNVERBOSE)

[Out] 1/2*((a^2*x^4+1)/x^4/a^2)^(1/2)*x^2*((1/a^2)^(1/2)*(1/a^2*(a^2*x^4+1))^(1/2)*a^2-ln(2*((1/a^2)^(1/2)*(1/a^2*(a^2*x^4+1))^(1/2)*a^2+1)/a^2/x^2))/(1/a^2)^(1/2)/(1/a^2*(a^2*x^4+1))^(1/2)/a^2+ln(x)/a

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 88 vs. 2(34) = 68.

Time = 0.25 (sec) , antiderivative size = 88, normalized size of antiderivative = 2.20

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x dx = \frac{2ax^2\sqrt{\frac{a^2x^4+1}{a^2x^4}} - \log\left(ax^2\sqrt{\frac{a^2x^4+1}{a^2x^4}} + 1\right) + \log\left(ax^2\sqrt{\frac{a^2x^4+1}{a^2x^4}} - 1\right) + 4\log(x)}{4a}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x,x, algorithm="fricas")

[Out] 1/4*(2*a*x^2*sqrt((a^2*x^4 + 1)/(a^2*x^4)) - log(a*x^2*sqrt((a^2*x^4 + 1)/(a^2*x^4)) + 1) + log(a*x^2*sqrt((a^2*x^4 + 1)/(a^2*x^4)) - 1) + 4*log(x))/a

Sympy [A] (verification not implemented)

Time = 2.98 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.45

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x dx = \frac{x^2}{2\sqrt{1 + \frac{1}{a^2x^4}}} + \frac{\log(x)}{a} - \frac{\operatorname{asinh}\left(\frac{1}{ax^2}\right)}{2a} + \frac{1}{2a^2x^2\sqrt{1 + \frac{1}{a^2x^4}}}$$

[In] integrate((1/a/x**2+(1+1/a**2/x**4)**(1/2))*x,x)

[Out] x**2/(2*sqrt(1 + 1/(a**2*x**4))) + log(x)/a - asinh(1/(a*x**2))/(2*a) + 1/(2*a**2*x**2*sqrt(1 + 1/(a**2*x**4)))

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 71 vs. 2(34) = 68.

Time = 0.23 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.78

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x dx = \frac{1}{2} x^2 \sqrt{\frac{1}{a^2x^4} + 1} - \frac{\log\left(ax^2 \sqrt{\frac{1}{a^2x^4} + 1} + 1\right)}{4a} + \frac{\log\left(ax^2 \sqrt{\frac{1}{a^2x^4} + 1} - 1\right)}{4a} + \frac{\log(x)}{a}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x,x, algorithm="maxima")

[Out] 1/2*x^2*sqrt(1/(a^2*x^4) + 1) - 1/4*log(a*x^2*sqrt(1/(a^2*x^4) + 1) + 1)/a + 1/4*log(a*x^2*sqrt(1/(a^2*x^4) + 1) - 1)/a + log(x)/a

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.52

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x dx = \frac{(a - |a|) \log(\sqrt{a^2x^4 + 1} + 1) + (a + |a|) \log(\sqrt{a^2x^4 + 1} - 1) + 2\sqrt{a^2x^4 + 1}|a|}{4a^2}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))*x,x, algorithm="giac")

[Out] 1/4*((a - abs(a))*log(sqrt(a^2*x^4 + 1) + 1) + (a + abs(a))*log(sqrt(a^2*x^4 + 1) - 1) + 2*sqrt(a^2*x^4 + 1)*abs(a))/a^2

Mupad [B] (verification not implemented)

Time = 5.71 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.08

$$\int e^{\operatorname{csch}^{-1}(ax^2)} x dx = \frac{x^2 \sqrt{\frac{1}{a^2 x^4} + 1}}{2} - \frac{\ln\left(\frac{1}{x^2}\right)}{2a} - \frac{\operatorname{asinh}\left(\frac{\sqrt{\frac{1}{a^2}}}{x^2}\right) \sqrt{\frac{1}{a^2}}}{2}$$

[In] `int(x*((1/(a^2*x^4) + 1)^(1/2) + 1/(a*x^2)),x)`

[Out] `(x^2*(1/(a^2*x^4) + 1)^(1/2))/2 - log(1/x^2)/(2*a) - (asinh((1/a^2)^(1/2)/x^2)*(1/a^2)^(1/2))/2`

3.42 $\int e^{\operatorname{csch}^{-1}(ax^2)} dx$

Optimal result	280
Rubi [A] (verified)	281
Mathematica [C] (verified)	283
Maple [C] (verified)	283
Fricas [F]	284
Sympy [C] (verification not implemented)	284
Maxima [F]	284
Giac [F]	284
Mupad [B] (verification not implemented)	285

Optimal result

Integrand size = 8, antiderivative size = 165

$$\int e^{\operatorname{csch}^{-1}(ax^2)} dx = -\frac{1}{ax} - \frac{2\sqrt{1 + \frac{1}{a^2x^4}}}{\left(a + \frac{1}{x^2}\right)x} + \sqrt{1 + \frac{1}{a^2x^4}}x$$

$$+ \frac{2\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}} \left(a + \frac{1}{x^2}\right) E\left(2 \cot^{-1}(\sqrt{ax}) \mid \frac{1}{2}\right)}{a^{3/2}\sqrt{1 + \frac{1}{a^2x^4}}}$$

$$- \frac{\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}} \left(a + \frac{1}{x^2}\right) \operatorname{EllipticF}\left(2 \cot^{-1}(\sqrt{ax}), \frac{1}{2}\right)}{a^{3/2}\sqrt{1 + \frac{1}{a^2x^4}}}$$

```
[Out] -1/a/x-2*(1+1/a^2/x^4)^(1/2)/(a+1/x^2)/x+x*(1+1/a^2/x^4)^(1/2)+2*(a+1/x^2)*
(cos(2*arccot(x*a^(1/2)))^2)^(1/2)/cos(2*arccot(x*a^(1/2)))*EllipticE(sin(2
*arccot(x*a^(1/2))),1/2*2^(1/2))*((a^2+1/x^4)/(a+1/x^2)^2)^(1/2)/a^(3/2)/(1
+1/a^2/x^4)^(1/2)-(a+1/x^2)*(cos(2*arccot(x*a^(1/2)))^2)^(1/2)/cos(2*arccot
(x*a^(1/2)))*EllipticF(sin(2*arccot(x*a^(1/2))),1/2*2^(1/2))*((a^2+1/x^4)/(
a+1/x^2)^2)^(1/2)/a^(3/2)/(1+1/a^2/x^4)^(1/2)
```


Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {6466, 30, 248, 283, 311, 226, 1210}

$$\int e^{\operatorname{csch}^{-1}(ax^2)} dx = x \sqrt{\frac{1}{a^2 x^4} + 1} - \frac{2 \sqrt{\frac{1}{a^2 x^4} + 1}}{x \left(a + \frac{1}{x^2}\right)} - \frac{\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}} \left(a + \frac{1}{x^2}\right) \operatorname{EllipticF}\left(2 \cot^{-1}(\sqrt{ax}), \frac{1}{2}\right)}{a^{3/2} \sqrt{\frac{1}{a^2 x^4} + 1}} + \frac{2 \sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}} \left(a + \frac{1}{x^2}\right) E\left(2 \cot^{-1}(\sqrt{ax}) \mid \frac{1}{2}\right)}{a^{3/2} \sqrt{\frac{1}{a^2 x^4} + 1}} - \frac{1}{ax}$$

[In] Int[E^ArcCsch[a*x^2], x]

[Out] $-(1/(a*x)) - (2*\sqrt{1 + 1/(a^2*x^4)})/((a + x^{(-2)})*x) + \sqrt{1 + 1/(a^2*x^4)}*x + (2*\sqrt{(a^2 + x^{(-4)})/(a + x^{(-2)})^2}*(a + x^{(-2)})*\operatorname{EllipticE}[2*\operatorname{ArcCot}[\sqrt{a}*x], 1/2])/ (a^{(3/2)}*\sqrt{1 + 1/(a^2*x^4)}) - (\sqrt{(a^2 + x^{(-4)})/(a + x^{(-2)})^2}*(a + x^{(-2)})*\operatorname{EllipticF}[2*\operatorname{ArcCot}[\sqrt{a}*x], 1/2])/ (a^{(3/2)}*\sqrt{1 + 1/(a^2*x^4)})$

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 226

Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 248

Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^2, x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0]

Rule 283

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^p/(c*(m + 1))), x] - Dist[b*n*(p/(c^n*(m + 1))), In

```
t[(c*x)^(m+n)*(a+b*x^n)^(p-1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[
n, 0] && GtQ[p, 0] && LtQ[m, -1] && !ILtQ[(m+n*p+n+1)/n, 0] && IntBi
nomialQ[a, b, c, n, m, p, x]
```

Rule 311

```
Int[(x_)^2/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, D
ist[1/q, Int[1/Sqrt[a + b*x^4], x], x] - Dist[1/q, Int[(1 - q*x^2)/Sqrt[a +
b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

Rule 1210

```
Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*
(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + c*x^4]))*E
llipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e
}, x] && PosQ[c/a]
```

Rule 6466

```
Int[E^ArcCsch[(a_)*(x_)^(p_)], x_Symbol] := Dist[1/a, Int[1/x^p, x], x] +
Int[Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, p}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\int \frac{1}{x^2} dx}{a} + \int \sqrt{1 + \frac{1}{a^2 x^4}} dx \\
&= -\frac{1}{ax} - \text{Subst} \left(\int \frac{\sqrt{1 + \frac{x^4}{a^2}}}{x^2} dx, x, \frac{1}{x} \right) \\
&= -\frac{1}{ax} + \sqrt{1 + \frac{1}{a^2 x^4}} x - \frac{2 \text{Subst} \left(\int \frac{x^2}{\sqrt{1 + \frac{x^4}{a^2}}} dx, x, \frac{1}{x} \right)}{a^2} \\
&= -\frac{1}{ax} + \sqrt{1 + \frac{1}{a^2 x^4}} x - \frac{2 \text{Subst} \left(\int \frac{1}{\sqrt{1 + \frac{x^4}{a^2}}} dx, x, \frac{1}{x} \right)}{a} + \frac{2 \text{Subst} \left(\int \frac{1 - \frac{x^2}{a}}{\sqrt{1 + \frac{x^4}{a^2}}} dx, x, \frac{1}{x} \right)}{a}
\end{aligned}$$

$$= -\frac{1}{ax} - \frac{2\sqrt{1 + \frac{1}{a^2x^4}}}{\left(a + \frac{1}{x^2}\right)x} + \sqrt{1 + \frac{1}{a^2x^4}}x + \frac{2\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}} \left(a + \frac{1}{x^2}\right) E\left(2 \cot^{-1}\left(\sqrt{ax}\right) \mid \frac{1}{2}\right)}{a^{3/2}\sqrt{1 + \frac{1}{a^2x^4}}}$$

$$- \frac{\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}} \left(a + \frac{1}{x^2}\right) \text{EllipticF}\left(2 \cot^{-1}\left(\sqrt{ax}\right), \frac{1}{2}\right)}{a^{3/2}\sqrt{1 + \frac{1}{a^2x^4}}}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.11 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.58

$$\int e^{\text{csch}^{-1}(ax^2)} dx$$

$$= \frac{\sqrt{2}e^{\text{csch}^{-1}(ax^2)} \sqrt{\frac{e^{\text{csch}^{-1}(ax^2)}}{-1+e^{2\text{csch}^{-1}(ax^2)}}} x \left(-3 + 4\sqrt{1 - e^{2\text{csch}^{-1}(ax^2)}} \text{Hypergeometric2F1}\left(\frac{3}{4}, \frac{3}{2}, \frac{7}{4}, e^{2\text{csch}^{-1}(ax^2)}\right)\right)}{3\sqrt{ax^2}}$$

[In] Integrate[E^ArcCsch[a*x^2],x]

[Out] (Sqrt[2]*E^ArcCsch[a*x^2]*Sqrt[E^ArcCsch[a*x^2]/(-1 + E^(2*ArcCsch[a*x^2]))]*x*(-3 + 4*Sqrt[1 - E^(2*ArcCsch[a*x^2])]*Hypergeometric2F1[3/4, 3/2, 7/4, E^(2*ArcCsch[a*x^2])]))/(3*Sqrt[a*x^2])

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.71 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.88

method	result
default	$\frac{\sqrt{\frac{x^4 a^2 + 1}{x^4 a^2}} x \left(-\sqrt{ia} a^2 x^4 + 2i\sqrt{-ia x^2 + 1} \sqrt{ia x^2 + 1} x \text{EllipticF}\left(x\sqrt{ia}, i\right) a - 2i\sqrt{-ia x^2 + 1} \sqrt{ia x^2 + 1} x \text{EllipticE}\left(x\sqrt{ia}, i\right) a - \sqrt{ia}\right)}{(x^4 a^2 + 1)\sqrt{ia}}$

[In] int(1/a/x^2+(1+1/a^2/x^4)^(1/2),x,method=_RETURNVERBOSE)

[Out] ((a^2*x^4+1)/x^4/a^2)^(1/2)*x*(-(I*a)^(1/2)*a^2*x^4+2*I*(1-I*a*x^2)^(1/2)*(1+I*a*x^2)^(1/2)*x*EllipticF(x*(I*a)^(1/2),I)*a-2*I*(1-I*a*x^2)^(1/2)*(1+I*a*x^2)^(1/2)*x*EllipticE(x*(I*a)^(1/2),I)*a-(I*a)^(1/2))/(a^2*x^4+1)/(I*a)^(1/2)-1/a/x

Fricas [F]

$$\int e^{\operatorname{csch}^{-1}(ax^2)} dx = \int \sqrt{\frac{1}{a^2x^4} + 1} + \frac{1}{ax^2} dx$$

[In] integrate(1/a/x^2+(1+1/a^2/x^4)^(1/2),x, algorithm="fricas")

[Out] integral((a*x^2*sqrt((a^2*x^4 + 1)/(a^2*x^4)) + 1)/(a*x^2), x)

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.50 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.25

$$\int e^{\operatorname{csch}^{-1}(ax^2)} dx = -\frac{x\Gamma(-\frac{1}{4}) {}_2F_1\left(-\frac{1}{2}, -\frac{1}{4} \middle| \frac{e^{i\pi}}{a^2x^4}\right)}{4\Gamma(\frac{3}{4})} - \frac{1}{ax}$$

[In] integrate(1/a/x**2+(1+1/a**2/x**4)**(1/2),x)

[Out] -x*gamma(-1/4)*hyper((-1/2, -1/4), (3/4,), exp_polar(I*pi)/(a**2*x**4))/(4*gamma(3/4)) - 1/(a*x)

Maxima [F]

$$\int e^{\operatorname{csch}^{-1}(ax^2)} dx = \int \sqrt{\frac{1}{a^2x^4} + 1} + \frac{1}{ax^2} dx$$

[In] integrate(1/a/x^2+(1+1/a^2/x^4)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(a^2*x^4 + 1)/x^2, x)/a - 1/(a*x)

Giac [F]

$$\int e^{\operatorname{csch}^{-1}(ax^2)} dx = \int \sqrt{\frac{1}{a^2x^4} + 1} + \frac{1}{ax^2} dx$$

[In] integrate(1/a/x^2+(1+1/a^2/x^4)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(1/(a^2*x^4) + 1) + 1/(a*x^2), x)

Mupad [B] (verification not implemented)

Time = 5.16 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.15

$$\int e^{\operatorname{csch}^{-1}(ax^2)} dx = x {}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; -\frac{1}{a^2 x^4}\right) - \frac{1}{ax}$$

[In] int((1/(a^2*x^4) + 1)^(1/2) + 1/(a*x^2),x)

[Out] x*hypergeom([-1/2, -1/4], 3/4, -1/(a^2*x^4)) - 1/(a*x)

3.43 $\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x} dx$

Optimal result	286
Rubi [A] (verified)	286
Mathematica [A] (verified)	288
Maple [B] (verified)	288
Fricas [B] (verification not implemented)	288
Sympy [A] (verification not implemented)	289
Maxima [A] (verification not implemented)	289
Giac [A] (verification not implemented)	289
Mupad [B] (verification not implemented)	290

Optimal result

Integrand size = 12, antiderivative size = 46

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x} dx = -\frac{1}{2} \sqrt{1 + \frac{1}{a^2x^4}} - \frac{1}{2ax^2} + \frac{1}{2} \operatorname{arctanh} \left(\sqrt{1 + \frac{1}{a^2x^4}} \right)$$

[Out] $-1/2/a/x^2+1/2*\operatorname{arctanh}((1+1/a^2/x^4)^{(1/2)})-1/2*(1+1/a^2/x^4)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6471, 30, 272, 52, 65, 214}

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x} dx = \frac{1}{2} \operatorname{arctanh} \left(\sqrt{\frac{1}{a^2x^4} + 1} \right) - \frac{1}{2} \sqrt{\frac{1}{a^2x^4} + 1} - \frac{1}{2ax^2}$$

[In] `Int[E^ArcCsch[a*x^2]/x,x]`

[Out] $-1/2*\operatorname{Sqrt}[1 + 1/(a^2*x^4)] - 1/(2*a*x^2) + \operatorname{ArcTanh}[\operatorname{Sqrt}[1 + 1/(a^2*x^4)]]/2$

Rule 30

`Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]`

Rule 52

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(`

$b*(m + n + 1))$, Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 6471

Int[E^ArcCsch[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Dist[1/a, Int[x^(m - p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int \frac{1}{x^3} dx}{a} + \int \frac{\sqrt{1 + \frac{1}{a^2 x^4}}}{x} dx \\
 &= -\frac{1}{2ax^2} - \frac{1}{4} \text{Subst} \left(\int \frac{\sqrt{1 + \frac{x}{a^2}}}{x} dx, x, \frac{1}{x^4} \right) \\
 &= -\frac{1}{2} \sqrt{1 + \frac{1}{a^2 x^4}} - \frac{1}{2ax^2} - \frac{1}{4} \text{Subst} \left(\int \frac{1}{x \sqrt{1 + \frac{x}{a^2}}} dx, x, \frac{1}{x^4} \right) \\
 &= -\frac{1}{2} \sqrt{1 + \frac{1}{a^2 x^4}} - \frac{1}{2ax^2} - \frac{1}{2} a^2 \text{Subst} \left(\int \frac{1}{-a^2 + a^2 x^2} dx, x, \sqrt{1 + \frac{1}{a^2 x^4}} \right) \\
 &= -\frac{1}{2} \sqrt{1 + \frac{1}{a^2 x^4}} - \frac{1}{2ax^2} + \frac{1}{2} \operatorname{arctanh} \left(\sqrt{1 + \frac{1}{a^2 x^4}} \right)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.48

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x} dx = -\frac{1}{2}e^{\operatorname{csch}^{-1}(ax^2)} + \operatorname{arctanh}\left(e^{\operatorname{csch}^{-1}(ax^2)}\right)$$

[In] Integrate[E^ArcCsch[a*x^2]/x,x]

[Out] -1/2*E^ArcCsch[a*x^2] + ArcTanh[E^ArcCsch[a*x^2]]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 85 vs. 2(36) = 72.

Time = 0.10 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.87

method	result	size
default	$-\frac{\sqrt{\frac{x^4 a^2 + 1}{x^4 a^2}} \left(-\ln \left(x^2 + \sqrt{\frac{x^4 a^2 + 1}{a^2}} \right) x^2 + \sqrt{\frac{x^4 a^2 + 1}{a^2}} \right)}{2\sqrt{\frac{x^4 a^2 + 1}{a^2}}} - \frac{1}{2a x^2}$	86

[In] int((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x,x,method=_RETURNVERBOSE)

[Out] -1/2*((a^2*x^4+1)/x^4/a^2)^(1/2)*(-ln(x^2+(1/a^2*(a^2*x^4+1))^(1/2))*x^2+(1/a^2*(a^2*x^4+1))^(1/2))/(1/a^2*(a^2*x^4+1))^(1/2)-1/2/a/x^2

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 74 vs. 2(36) = 72.

Time = 0.24 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.61

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x} dx = -\frac{ax^2 \log \left(ax^2 \sqrt{\frac{a^2 x^4 + 1}{a^2 x^4}} - ax^2 \right) + ax^2 \sqrt{\frac{a^2 x^4 + 1}{a^2 x^4}} + ax^2 + 1}{2ax^2}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x,x, algorithm="fricas")

[Out] -1/2*(a*x^2*log(a*x^2*sqrt((a^2*x^4 + 1)/(a^2*x^4)) - a*x^2) + a*x^2*sqrt((a^2*x^4 + 1)/(a^2*x^4)) + a*x^2 + 1)/(a*x^2)

Sympy [A] (verification not implemented)

Time = 4.95 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.17

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x} dx = -\frac{ax^2}{2\sqrt{a^2x^4+1}} + \frac{\operatorname{asinh}(ax^2)}{2} - \frac{1}{2ax^2} - \frac{1}{2ax^2\sqrt{a^2x^4+1}}$$

[In] integrate((1/a/x**2+(1+1/a**2/x**4)**(1/2))/x,x)

[Out] -a*x**2/(2*sqrt(a**2*x**4 + 1)) + asinh(a*x**2)/2 - 1/(2*a*x**2) - 1/(2*a*x**2*sqrt(a**2*x**4 + 1))

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.17

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x} dx = -\frac{1}{2} \sqrt{\frac{1}{a^2x^4} + 1} - \frac{1}{2ax^2} + \frac{1}{4} \log \left(\sqrt{\frac{1}{a^2x^4} + 1} + 1 \right) - \frac{1}{4} \log \left(\sqrt{\frac{1}{a^2x^4} + 1} - 1 \right)$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x,x, algorithm="maxima")

[Out] -1/2*sqrt(1/(a^2*x^4) + 1) - 1/2/(a*x^2) + 1/4*log(sqrt(1/(a^2*x^4) + 1) + 1) - 1/4*log(sqrt(1/(a^2*x^4) + 1) - 1)

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.43

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x} dx = -\frac{a^2 \log(-x^2|a| + \sqrt{a^2x^4+1}) - \frac{2a^2}{(x^2|a| - \sqrt{a^2x^4+1})^2 - 1} + \frac{a}{x^2}}{2a^2}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x,x, algorithm="giac")

[Out] -1/2*(a^2*log(-x^2*abs(a) + sqrt(a^2*x^4 + 1)) - 2*a^2/((x^2*abs(a) - sqrt(a^2*x^4 + 1))^2 - 1) + a/x^2)/a^2

Mupad [B] (verification not implemented)

Time = 5.11 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.78

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x} dx = \frac{\operatorname{atanh}\left(\sqrt{\frac{1}{a^2x^4} + 1}\right)}{2} - \frac{\sqrt{\frac{1}{a^2x^4} + 1}}{2} - \frac{1}{2ax^2}$$

[In] int(((1/(a^2*x^4) + 1)^(1/2) + 1/(a*x^2))/x,x)

[Out] atanh((1/(a^2*x^4) + 1)^(1/2))/2 - (1/(a^2*x^4) + 1)^(1/2)/2 - 1/(2*a*x^2)

3.44 $\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^2} dx$

Optimal result	291
Rubi [A] (verified)	291
Mathematica [C] (verified)	293
Maple [C] (verified)	293
Fricas [A] (verification not implemented)	293
Sympy [C] (verification not implemented)	294
Maxima [F]	294
Giac [F]	294
Mupad [B] (verification not implemented)	295

Optimal result

Integrand size = 12, antiderivative size = 91

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^2} dx = -\frac{1}{3ax^3} - \frac{\sqrt{1 + \frac{1}{a^2x^4}}}{3x} - \frac{\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}} \operatorname{EllipticF}\left(2 \cot^{-1}(\sqrt{ax}), \frac{1}{2}\right)}{3\sqrt{a}\sqrt{1 + \frac{1}{a^2x^4}}}$$

[Out] $-1/3/a/x^3 - 1/3*(1+1/a^2/x^4)^{(1/2)}/x - 1/3*(a+1/x^2)*(\cos(2*\operatorname{arccot}(x*a^{(1/2)}))^2)^{(1/2)}/\cos(2*\operatorname{arccot}(x*a^{(1/2)}))*\operatorname{EllipticF}(\sin(2*\operatorname{arccot}(x*a^{(1/2)})), 1/2*2^{(1/2)})*((a^2+1/x^4)/(a+1/x^2)^2)^{(1/2)}/a^{(1/2)}/(1+1/a^2/x^4)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {6471, 30, 342, 201, 226}

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^2} dx = -\frac{\sqrt{\frac{1}{a^2x^4} + 1}}{3x} - \frac{\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}} \operatorname{EllipticF}\left(2 \cot^{-1}(\sqrt{ax}), \frac{1}{2}\right)}{3\sqrt{a}\sqrt{\frac{1}{a^2x^4} + 1}} - \frac{1}{3ax^3}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcCsch}[a*x^2]}/x^2, x]$

[Out] $-1/3*1/(a*x^3) - \operatorname{Sqrt}[1 + 1/(a^2*x^4)]/(3*x) - (\operatorname{Sqrt}[(a^2 + x^{(-4)})/(a + x^{(-2)})^2]*(a + x^{(-2)})*\operatorname{EllipticF}[2*\operatorname{ArcCot}[\operatorname{Sqrt}[a]*x], 1/2])/ (3*\operatorname{Sqrt}[a]*\operatorname{Sqrt}[1 + 1/(a^2*x^4)])$

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 201

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p + 1)), x] + Dist[a*n*(p/(n*p + 1)), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 226

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 342

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]

Rule 6471

Int[E^ArcCsch[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Dist[1/a, Int[x^(m - p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int \frac{1}{x^4} dx}{a} + \int \frac{\sqrt{1 + \frac{1}{a^2 x^4}}}{x^2} dx \\
 &= -\frac{1}{3ax^3} - \text{Subst}\left(\int \sqrt{1 + \frac{x^4}{a^2}} dx, x, \frac{1}{x}\right) \\
 &= -\frac{1}{3ax^3} - \frac{\sqrt{1 + \frac{1}{a^2 x^4}}}{3x} - \frac{2}{3} \text{Subst}\left(\int \frac{1}{\sqrt{1 + \frac{x^4}{a^2}}} dx, x, \frac{1}{x}\right) \\
 &= -\frac{1}{3ax^3} - \frac{\sqrt{1 + \frac{1}{a^2 x^4}}}{3x} - \frac{\sqrt{\frac{a^2 + \frac{1}{x^4}}{(a + \frac{1}{x^2})^2}} (a + \frac{1}{x^2}) \text{EllipticF}\left(2 \cot^{-1}(\sqrt{ax}), \frac{1}{2}\right)}{3\sqrt{a}\sqrt{1 + \frac{1}{a^2 x^4}}}
 \end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.11 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.05

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^2} dx = \frac{a \sqrt{\frac{e^{\operatorname{csch}^{-1}(ax^2)}}{-2+2e^{2\operatorname{csch}^{-1}(ax^2)}}} x \left(-1 + e^{2\operatorname{csch}^{-1}(ax^2)} + 4\sqrt{1 - e^{2\operatorname{csch}^{-1}(ax^2)}} \operatorname{Hypergeometric2F1} \left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, e^{2\operatorname{csch}^{-1}(ax^2)} \right) \right)}{3\sqrt{ax^2}}$$

[In] Integrate[E^ArcCsch[a*x^2]/x^2,x]

[Out] $-1/3*(a*\sqrt{E^{\operatorname{ArcCsch}[a*x^2]} / (-2 + 2*E^{(2*\operatorname{ArcCsch}[a*x^2])})})*x*(-1 + E^{(2*\operatorname{ArcCsch}[a*x^2])} + 4*\sqrt{1 - E^{(2*\operatorname{ArcCsch}[a*x^2])}})*\operatorname{Hypergeometric2F1}[1/4, 1/2, 5/4, E^{(2*\operatorname{ArcCsch}[a*x^2])}]) / \sqrt{a*x^2}$

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.80 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.22

method	result	size
default	$-\frac{\sqrt{\frac{x^4 a^2 + 1}{x^4 a^2}} \left(-2\sqrt{-ia x^2 + 1} \sqrt{ia x^2 + 1} \operatorname{EllipticF}\left(x\sqrt{ia}, i\right) x^3 a^2 + \sqrt{ia} a^2 x^4 + \sqrt{ia} \right)}{3x(x^4 a^2 + 1)\sqrt{ia}} - \frac{1}{3a x^3}$	111

[In] int((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^2,x,method=_RETURNVERBOSE)

[Out] $-1/3*((a^2*x^4+1)/x^4/a^2)^(1/2)*(-2*(1-I*a*x^2)^(1/2)*(1+I*a*x^2)^(1/2)*\operatorname{EllipticF}(x*(I*a)^(1/2), I)*x^3*a^2+(I*a)^(1/2)*a^2*x^4+(I*a)^(1/2))/x/(a^2*x^4+1)/(I*a)^(1/2)-1/3/a/x^3$

Fricas [A] (verification not implemented)

none

Time = 0.07 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.63

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^2} dx = -\frac{2(-a^2)^{\frac{3}{4}} x^3 F(\arcsin\left(\frac{(-a^2)^{\frac{1}{4}} x}{-1}\right) | -1) + ax^2 \sqrt{\frac{a^2 x^4 + 1}{a^2 x^4}} + 1}{3ax^3}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^2,x, algorithm="fricas")

[Out] $-1/3*(2*(-a^2)^(3/4)*x^3*\operatorname{elliptic}_f(\arcsin((-a^2)^(1/4)*x), -1) + a*x^2*\operatorname{sqrt}((a^2*x^4 + 1)/(a^2*x^4)) + 1)/(a*x^3)$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.05 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.46

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^2} dx = -\frac{\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(-\frac{1}{2}, \frac{1}{4} \middle| \frac{5}{4}, \frac{e^{i\pi}}{a^2 x^4}\right)}{4x\Gamma\left(\frac{5}{4}\right)} - \frac{1}{3ax^3}$$

[In] integrate((1/a/x**2+(1+1/a**2/x**4)**(1/2))/x**2,x)

[Out] -gamma(1/4)*hyper((-1/2, 1/4), (5/4,), exp_polar(I*pi)/(a**2*x**4))/(4*x*gamma(5/4)) - 1/(3*a*x**3)

Maxima [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^2} dx = \int \frac{\sqrt{\frac{1}{a^2 x^4} + 1} + \frac{1}{ax^2}}{x^2} dx$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^2,x, algorithm="maxima")

[Out] integrate(sqrt(a^2*x^4 + 1)/x^4, x)/a - 1/3/(a*x^3)

Giac [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^2} dx = \int \frac{\sqrt{\frac{1}{a^2 x^4} + 1} + \frac{1}{ax^2}}{x^2} dx$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^2,x, algorithm="giac")

[Out] integrate((sqrt(1/(a^2*x^4) + 1) + 1/(a*x^2))/x^2, x)

Mupad [B] (verification not implemented)

Time = 5.26 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.30

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^2} dx = -\frac{{}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{5}{4}; -\frac{1}{a^2 x^4}\right)}{x} - \frac{1}{3 a x^3}$$

[In] int(((1/(a^2*x^4) + 1)^(1/2) + 1/(a*x^2))/x^2,x)

[Out] - hypergeom([-1/2, 1/4], 5/4, -1/(a^2*x^4))/x - 1/(3*a*x^3)

3.45 $\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^3} dx$

Optimal result	296
Rubi [A] (verified)	296
Mathematica [A] (verified)	298
Maple [B] (verified)	298
Fricas [B] (verification not implemented)	298
Sympy [A] (verification not implemented)	299
Maxima [B] (verification not implemented)	299
Giac [B] (verification not implemented)	299
Mupad [B] (verification not implemented)	300

Optimal result

Integrand size = 12, antiderivative size = 42

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^3} dx = -\frac{1}{4ax^4} - \frac{\sqrt{1 + \frac{1}{a^2x^4}}}{4x^2} - \frac{1}{4} \operatorname{acsch}^{-1}(ax^2)$$

[Out] $-1/4/a/x^4 - 1/4*a*\operatorname{arccsch}(a*x^2) - 1/4*(1+1/a^2/x^4)^{(1/2)}/x^2$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6471, 30, 342, 281, 201, 221}

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^3} dx = -\frac{\sqrt{\frac{1}{a^2x^4} + 1}}{4x^2} - \frac{1}{4ax^4} - \frac{1}{4} \operatorname{acsch}^{-1}(ax^2)$$

[In] $\operatorname{Int}[E^{\operatorname{ArcCsch}[a*x^2]}/x^3, x]$

[Out] $-1/4*1/(a*x^4) - \operatorname{Sqrt}[1 + 1/(a^2*x^4)]/(4*x^2) - (a*\operatorname{ArcCsch}[a*x^2])/4$

Rule 30

$\operatorname{Int}[(x_)^{(m_.)}, x_Symbol] := \operatorname{Simp}[x^{(m+1)}/(m+1), x] /;$ FreeQ[m, x] && NeQ[m, -1]

Rule 201

$\operatorname{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] := \operatorname{Simp}[x*((a + b*x^n)^p/(n*p + 1)), x] + \operatorname{Dist}[a*n*(p/(n*p + 1)), \operatorname{Int}[(a + b*x^n)^{(p-1)}, x], x] /;$ Free

$Q[\{a, b\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{GtQ}[p, 0] \&\& (\text{IntegerQ}[2*p] \mid\mid (\text{EqQ}[n, 2] \&\& \text{IntegerQ}[4*p]) \mid\mid (\text{EqQ}[n, 2] \&\& \text{IntegerQ}[3*p]) \mid\mid \text{LtQ}[\text{Denominator}[p + 1/n], \text{Denominator}[p]])$

Rule 221

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2], x_Symbol] \text{ :> } \text{Simp}[\text{ArcSinh}[\text{Rt}[b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[b, 2], x] \text{ /; } \text{FreeQ}[\{a, b\}, x] \&\& \text{GtQ}[a, 0] \&\& \text{PosQ}[b]$

Rule 281

$\text{Int}[(x_)^{(m_.)*((a_) + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \text{ :> } \text{With}[\{k = \text{GCD}[m + 1, n]\}, \text{Dist}[1/k, \text{Subst}[\text{Int}[x^{(m + 1)/k - 1}*(a + b*x^{(n/k)})^p, x], x, x^k], x] \text{ /; } k \neq 1] \text{ /; } \text{FreeQ}[\{a, b, p\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{IntegerQ}[m]$

Rule 342

$\text{Int}[(x_)^{(m_.)*((a_) + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \text{ :> } -\text{Subst}[\text{Int}[(a + b/x^n)^p/x^{(m + 2)}, x], x, 1/x] \text{ /; } \text{FreeQ}[\{a, b, p\}, x] \&\& \text{ILtQ}[n, 0] \&\& \text{IntegerQ}[m]$

Rule 6471

$\text{Int}[\text{E}^{\text{ArcCsch}[(a_)*(x_)^{(p_.)}]}*(x_)^{(m_.)}, x_Symbol] \text{ :> } \text{Dist}[1/a, \text{Int}[x^{(m - p)}, x], x] + \text{Int}[x^m*\text{Sqrt}[1 + 1/(a^2*x^{(2*p)})], x] \text{ /; } \text{FreeQ}[\{a, m, p\}, x]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{1}{x^5} \frac{dx}{a} + \int \frac{\sqrt{1 + \frac{1}{a^2 x^4}}}{x^3} dx \\
 &= -\frac{1}{4ax^4} - \text{Subst}\left(\int x \sqrt{1 + \frac{x^4}{a^2}} dx, x, \frac{1}{x}\right) \\
 &= -\frac{1}{4ax^4} - \frac{1}{2} \text{Subst}\left(\int \sqrt{1 + \frac{x^2}{a^2}} dx, x, \frac{1}{x^2}\right) \\
 &= -\frac{1}{4ax^4} - \frac{\sqrt{1 + \frac{1}{a^2 x^4}}}{4x^2} - \frac{1}{4} \text{Subst}\left(\int \frac{1}{\sqrt{1 + \frac{x^2}{a^2}}} dx, x, \frac{1}{x^2}\right) \\
 &= -\frac{1}{4ax^4} - \frac{\sqrt{1 + \frac{1}{a^2 x^4}}}{4x^2} - \frac{1}{4} \text{acsch}^{-1}(ax^2)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.57

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^3} dx = -\frac{1}{8}a \left(e^{2\operatorname{csch}^{-1}(ax^2)} + 2\operatorname{csch}^{-1}(ax^2) \right)$$

[In] Integrate[E^ArcCsch[a*x^2]/x^3,x]

[Out] -1/8*(a*(E^(2*ArcCsch[a*x^2]) + 2*ArcCsch[a*x^2]))

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 113 vs. 2(34) = 68.

Time = 0.11 (sec) , antiderivative size = 114, normalized size of antiderivative = 2.71

method	result	size
default	$-\frac{\sqrt{\frac{x^4 a^2 + 1}{x^4 a^2}} \left(\ln \left(\frac{2\sqrt{\frac{1}{a^2}} \sqrt{\frac{x^4 a^2 + 1}{a^2 x^2}} a^2 + 2}{a^2 x^2} \right) x^4 + \sqrt{\frac{x^4 a^2 + 1}{a^2}} \sqrt{\frac{1}{a^2}} \right)}{4x^2 \sqrt{\frac{x^4 a^2 + 1}{a^2}} \sqrt{\frac{1}{a^2}}} - \frac{1}{4a x^4}$	114

[In] int((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^3,x,method=_RETURNVERBOSE)

[Out] -1/4*((a^2*x^4+1)/x^4/a^2)^(1/2)/x^2*(ln(2*((1/a^2)^(1/2)*(1/a^2*(a^2*x^4+1)))^(1/2)*a^2+1)/a^2/x^2)*x^4+(1/a^2*(a^2*x^4+1))^(1/2)*(1/a^2)^(1/2))/(1/a^2*(a^2*x^4+1))^(1/2)/(1/a^2)^(1/2)-1/4/a/x^4

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 101 vs. 2(34) = 68.

Time = 0.24 (sec) , antiderivative size = 101, normalized size of antiderivative = 2.40

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^3} dx = -\frac{a^2 x^4 \log \left(ax^2 \sqrt{\frac{a^2 x^4 + 1}{a^2 x^4}} + 1 \right) - a^2 x^4 \log \left(ax^2 \sqrt{\frac{a^2 x^4 + 1}{a^2 x^4}} - 1 \right) + 2 a x^2 \sqrt{\frac{a^2 x^4 + 1}{a^2 x^4}} + 2}{8 a x^4}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^3,x, algorithm="fricas")

[Out] -1/8*(a^2*x^4*log(a*x^2*sqrt((a^2*x^4 + 1)/(a^2*x^4)) + 1) - a^2*x^4*log(a*x^2*sqrt((a^2*x^4 + 1)/(a^2*x^4)) - 1) + 2*a*x^2*sqrt((a^2*x^4 + 1)/(a^2*x^4)) + 2)/(a*x^4)

Sympy [A] (verification not implemented)

Time = 1.34 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^3} dx = -\frac{a\left(\frac{\operatorname{asinh}\left(\frac{1}{ax^2}\right)}{2} + \frac{\sqrt{1+\frac{1}{a^2x^4}}}{2ax^2}\right)}{2} - \frac{1}{4ax^4}$$

[In] integrate((1/a/x**2+(1+1/a**2/x**4)**(1/2))/x**3,x)

[Out] -a*(asinh(1/(a*x**2))/2 + sqrt(1 + 1/(a**2*x**4))/(2*a*x**2))/2 - 1/(4*a*x**4)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 92 vs. 2(34) = 68.

Time = 0.23 (sec) , antiderivative size = 92, normalized size of antiderivative = 2.19

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^3} dx = -\frac{a^2x^2\sqrt{\frac{1}{a^2x^4} + 1}}{4(a^2x^4(\frac{1}{a^2x^4} + 1) - 1)} - \frac{1}{8}a \log\left(ax^2\sqrt{\frac{1}{a^2x^4} + 1} + 1\right) + \frac{1}{8}a \log\left(ax^2\sqrt{\frac{1}{a^2x^4} + 1} - 1\right) - \frac{1}{4ax^4}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^3,x, algorithm="maxima")

[Out] -1/4*a^2*x^2*sqrt(1/(a^2*x^4) + 1)/(a^2*x^4*(1/(a^2*x^4) + 1) - 1) - 1/8*a*log(a*x^2*sqrt(1/(a^2*x^4) + 1) + 1) + 1/8*a*log(a*x^2*sqrt(1/(a^2*x^4) + 1) - 1) - 1/4/(a*x^4)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 76 vs. 2(34) = 68.

Time = 0.27 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.81

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^3} dx = -\frac{a^4|a| \log(\sqrt{a^2x^4 + 1} + 1) - a^4|a| \log(\sqrt{a^2x^4 + 1} - 1) + \frac{2(\sqrt{a^2x^4 + 1}a^4|a| + a^5)}{a^2x^4}}{8a^4}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^3,x, algorithm="giac")

[Out] -1/8*(a^4*abs(a)*log(sqrt(a^2*x^4 + 1) + 1) - a^4*abs(a)*log(sqrt(a^2*x^4 + 1) - 1) + 2*(sqrt(a^2*x^4 + 1)*a^4*abs(a) + a^5)/(a^2*x^4))/a^4

Mupad [B] (verification not implemented)

Time = 6.32 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^3} dx = -\frac{\operatorname{asinh}\left(\frac{\sqrt{\frac{1}{a^2}}}{x^2}\right)}{4\sqrt{\frac{1}{a^2}}} - \frac{\sqrt{\frac{1}{a^2x^4} + 1}}{4x^2} - \frac{1}{4ax^4}$$

[In] int(((1/(a^2*x^4) + 1)^(1/2) + 1/(a*x^2))/x^3,x)

[Out] - asinh((1/a^2)^(1/2)/x^2)/(4*(1/a^2)^(1/2)) - (1/(a^2*x^4) + 1)^(1/2)/(4*x^2) - 1/(4*a*x^4)

$$3.46 \quad \int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^4} dx$$

Optimal result	301
Rubi [A] (verified)	302
Mathematica [C] (verified)	304
Maple [C] (verified)	304
Fricas [A] (verification not implemented)	305
Sympy [C] (verification not implemented)	305
Maxima [F]	305
Giac [F]	306
Mupad [F(-1)]	306

Optimal result

Integrand size = 12, antiderivative size = 181

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^4} dx = -\frac{1}{5ax^5} - \frac{\sqrt{1 + \frac{1}{a^2x^4}}}{5x^3} - \frac{2a^2\sqrt{1 + \frac{1}{a^2x^4}}}{5\left(a + \frac{1}{x^2}\right)x}$$

$$+ \frac{2\sqrt{a}\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}}\left(a + \frac{1}{x^2}\right)E\left(2\cot^{-1}\left(\sqrt{ax}\right)\left|\frac{1}{2}\right.\right)}{5\sqrt{1 + \frac{1}{a^2x^4}}}$$

$$- \frac{\sqrt{a}\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}}\left(a + \frac{1}{x^2}\right)\operatorname{EllipticF}\left(2\cot^{-1}\left(\sqrt{ax}\right), \frac{1}{2}\right)}{5\sqrt{1 + \frac{1}{a^2x^4}}}$$

```
[Out] -1/5/a/x^5-1/5*(1+1/a^2/x^4)^(1/2)/x^3-2/5*a^2*(1+1/a^2/x^4)^(1/2)/(a+1/x^2)
)/x+2/5*(a+1/x^2)*(cos(2*arccot(x*a^(1/2))))^(1/2)/cos(2*arccot(x*a^(1/2)
))*EllipticE(sin(2*arccot(x*a^(1/2))),1/2*2^(1/2))*a^(1/2)*((a^2+1/x^4)/(a+
1/x^2)^2)^(1/2)/(1+1/a^2/x^4)^(1/2)-1/5*(a+1/x^2)*(cos(2*arccot(x*a^(1/2)))
)^(1/2)/cos(2*arccot(x*a^(1/2)))*EllipticF(sin(2*arccot(x*a^(1/2))),1/2*2
^(1/2))*a^(1/2)*((a^2+1/x^4)/(a+1/x^2)^2)^(1/2)/(1+1/a^2/x^4)^(1/2)
```

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {6471, 30, 342, 285, 311, 226, 1210}

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^4} dx = -\frac{\sqrt{\frac{1}{a^2x^4} + 1}}{5x^3} - \frac{2a^2\sqrt{\frac{1}{a^2x^4} + 1}}{5x\left(a + \frac{1}{x^2}\right)} - \frac{\sqrt{a}\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}}\left(a + \frac{1}{x^2}\right) \operatorname{EllipticF}\left(2 \cot^{-1}\left(\sqrt{ax}\right), \frac{1}{2}\right)}{5\sqrt{\frac{1}{a^2x^4} + 1}} + \frac{2\sqrt{a}\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}}\left(a + \frac{1}{x^2}\right) E\left(2 \cot^{-1}\left(\sqrt{ax}\right) \middle| \frac{1}{2}\right)}{5\sqrt{\frac{1}{a^2x^4} + 1}} - \frac{1}{5ax^5}$$

[In] Int[E^ArcCsch[a*x^2]/x^4,x]

[Out] -1/5*1/(a*x^5) - Sqrt[1 + 1/(a^2*x^4)]/(5*x^3) - (2*a^2*Sqrt[1 + 1/(a^2*x^4)])/(5*(a + x^(-2))*x) + (2*Sqrt[a]*Sqrt[(a^2 + x^(-4))/(a + x^(-2))^2]*(a + x^(-2))*EllipticE[2*ArcCot[Sqrt[a]*x], 1/2])/(5*Sqrt[1 + 1/(a^2*x^4)]) - (Sqrt[a]*Sqrt[(a^2 + x^(-4))/(a + x^(-2))^2]*(a + x^(-2))*EllipticF[2*ArcCot[Sqrt[a]*x], 1/2])/(5*Sqrt[1 + 1/(a^2*x^4)])

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 226

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 285

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^p/(c*(m + n*p + 1))), x] + Dist[a*n*(p/(m + n*p + 1)), Int[(c*x)^m*(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[n, 0] && GtQ[p, 0] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 311

Int[(x_)^2/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, Dist[1/q, Int[1/Sqrt[a + b*x^4], x], x] - Dist[1/q, Int[(1 - q*x^2)/Sqrt[a + b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 342

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^p, x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]

Rule 1210

Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + c*x^4])*EllipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]

Rule 6471

Int[E^ArcCsch[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Dist[1/a, Int[x^(m - p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int \frac{1}{x^6} dx}{a} + \int \frac{\sqrt{1 + \frac{1}{a^2 x^4}}}{x^4} dx \\
 &= -\frac{1}{5ax^5} - \text{Subst}\left(\int x^2 \sqrt{1 + \frac{x^4}{a^2}} dx, x, \frac{1}{x}\right) \\
 &= -\frac{1}{5ax^5} - \frac{\sqrt{1 + \frac{1}{a^2 x^4}}}{5x^3} - \frac{2}{5} \text{Subst}\left(\int \frac{x^2}{\sqrt{1 + \frac{x^4}{a^2}}} dx, x, \frac{1}{x}\right) \\
 &= -\frac{1}{5ax^5} - \frac{\sqrt{1 + \frac{1}{a^2 x^4}}}{5x^3} - \frac{1}{5}(2a) \text{Subst}\left(\int \frac{1}{\sqrt{1 + \frac{x^4}{a^2}}} dx, x, \frac{1}{x}\right) \\
 &\quad + \frac{1}{5}(2a) \text{Subst}\left(\int \frac{1 - \frac{x^2}{a}}{\sqrt{1 + \frac{x^4}{a^2}}} dx, x, \frac{1}{x}\right)
 \end{aligned}$$

$$= -\frac{1}{5ax^5} - \frac{\sqrt{1 + \frac{1}{a^2x^4}}}{5x^3} - \frac{2a^2\sqrt{1 + \frac{1}{a^2x^4}}}{5\left(a + \frac{1}{x^2}\right)x} + \frac{2\sqrt{a}\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}}\left(a + \frac{1}{x^2}\right)E\left(2\cot^{-1}\left(\sqrt{ax}\right)\left|\frac{1}{2}\right.\right)}{5\sqrt{1 + \frac{1}{a^2x^4}}}$$

$$- \frac{\sqrt{a}\sqrt{\frac{a^2 + \frac{1}{x^4}}{\left(a + \frac{1}{x^2}\right)^2}}\left(a + \frac{1}{x^2}\right)\text{EllipticF}\left(2\cot^{-1}\left(\sqrt{ax}\right), \frac{1}{2}\right)}{5\sqrt{1 + \frac{1}{a^2x^4}}}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.15 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.63

$$\int \frac{e^{\text{csch}^{-1}(ax^2)}}{x^4} dx$$

$$= \frac{(ax^2)^{3/2} \left(3 \left(1 - e^{2\text{csch}^{-1}(ax^2)} \right)^{3/2} + 4e^{2\text{csch}^{-1}(ax^2)} \text{Hypergeometric2F1} \left(-\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, e^{2\text{csch}^{-1}(ax^2)} \right) \right)}{6\sqrt{2 - 2e^{2\text{csch}^{-1}(ax^2)}} \sqrt{\frac{e^{\text{csch}^{-1}(ax^2)}}{-1 + e^{2\text{csch}^{-1}(ax^2)}}} x^3}$$

[In] Integrate[E^ArcCsch[a*x^2]/x^4,x]

[Out] ((a*x^2)^(3/2)*(3*(1 - E^(2*ArcCsch[a*x^2]))^(3/2) + 4*E^(2*ArcCsch[a*x^2])*Hypergeometric2F1[-1/2, 3/4, 7/4, E^(2*ArcCsch[a*x^2])]))/(6*Sqrt[2 - 2*E^(2*ArcCsch[a*x^2])]*Sqrt[E^ArcCsch[a*x^2]/(-1 + E^(2*ArcCsch[a*x^2]))]*x^3)

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.93 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.94

method	result
default	$\frac{\sqrt{\frac{x^4 a^2 + 1}{x^4 a^2}} \left(-2\sqrt{ia} a^4 x^8 + 2ia^3 \sqrt{-ia x^2 + 1} \sqrt{ia x^2 + 1} x^5 \text{EllipticF}\left(x\sqrt{ia}, i\right) - 2ia^3 \sqrt{-ia x^2 + 1} \sqrt{ia x^2 + 1} x^5 \text{EllipticE}\left(x\sqrt{ia}, i\right) - 3\sqrt{ia} a^2 x^4 - (I*a)^{(1/2)} \right)}{5x^3(x^4 a^2 + 1)\sqrt{ia}}$

[In] int((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^4,x,method=_RETURNVERBOSE)

[Out] 1/5*((a^2*x^4+1)/x^4/a^2)^(1/2)*(-2*(I*a)^(1/2)*a^4*x^8+2*I*a^3*(1-I*a*x^2)^(1/2)*(1+I*a*x^2)^(1/2)*x^5*EllipticF(x*(I*a)^(1/2),I)-2*I*a^3*(1-I*a*x^2)^(1/2)*(1+I*a*x^2)^(1/2)*x^5*EllipticE(x*(I*a)^(1/2),I)-3*(I*a)^(1/2)*a^2*x^4-(I*a)^(1/2))/x^3/(a^2*x^4+1)/(I*a)^(1/2)-1/5/a/x^5

Fricas [A] (verification not implemented)

none

Time = 0.08 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.54

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^4} dx = \frac{2(-a^2)^{\frac{3}{4}} a^2 x^5 E(\arcsin((-a^2)^{\frac{1}{4}} x) | -1) - 2(-a^2)^{\frac{3}{4}} a^2 x^5 F(\arcsin((-a^2)^{\frac{1}{4}} x) | -1) + (2a^3 x^6 + ax^2) \sqrt{a^2 x^4 + 1}}{5ax^5}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^4,x, algorithm="fricas")

[Out] $-1/5*(2*(-a^2)^{(3/4)}*a^2*x^5*\operatorname{elliptic}_e(\arcsin((-a^2)^{(1/4)}*x), -1) - 2*(-a^2)^{(3/4)}*a^2*x^5*\operatorname{elliptic}_f(\arcsin((-a^2)^{(1/4)}*x), -1) + (2*a^3*x^6 + a*x^2)*\operatorname{sqrt}((a^2*x^4 + 1)/(a^2*x^4)) + 1)/(a*x^5)$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.19 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.24

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^4} dx = -\frac{\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{1}{2}, \frac{3}{4} \\ \frac{7}{4} \end{matrix} \middle| \frac{e^{i\pi}}{a^2 x^4}\right)}{4x^3 \Gamma\left(\frac{7}{4}\right)} - \frac{1}{5ax^5}$$

[In] integrate((1/a/x**2+(1+1/a**2/x**4)**(1/2))/x**4,x)

[Out] $-\operatorname{gamma}(3/4)*\operatorname{hyper}((-1/2, 3/4), (7/4,), \exp_polar(I*\pi)/(a**2*x**4))/(4*x**3*\operatorname{gamma}(7/4)) - 1/(5*a*x**5)$

Maxima [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^4} dx = \int \frac{\sqrt{\frac{1}{a^2 x^4} + 1} + \frac{1}{ax^2}}{x^4} dx$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^4,x, algorithm="maxima")

[Out] integrate(sqrt(a^2*x^4 + 1)/x^6, x)/a - 1/5/(a*x^5)

Giac [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^4} dx = \int \frac{\sqrt{\frac{1}{a^2x^4} + 1} + \frac{1}{ax^2}}{x^4} dx$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^4,x, algorithm="giac")

[Out] integrate((sqrt(1/(a^2*x^4) + 1) + 1/(a*x^2))/x^4, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^4} dx = \int \frac{\sqrt{\frac{1}{a^2x^4} + 1} + \frac{1}{ax^2}}{x^4} dx$$

[In] int(((1/(a^2*x^4) + 1)^(1/2) + 1/(a*x^2))/x^4,x)

[Out] int(((1/(a^2*x^4) + 1)^(1/2) + 1/(a*x^2))/x^4, x)

$$3.47 \quad \int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^5} dx$$

Optimal result	307
Rubi [A] (verified)	307
Mathematica [A] (verified)	308
Maple [A] (verified)	308
Fricas [A] (verification not implemented)	309
Sympy [A] (verification not implemented)	309
Maxima [A] (verification not implemented)	309
Giac [B] (verification not implemented)	310
Mupad [B] (verification not implemented)	310

Optimal result

Integrand size = 12, antiderivative size = 31

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^5} dx = -\frac{1}{6}a^2 \left(1 + \frac{1}{a^2x^4}\right)^{3/2} - \frac{1}{6ax^6}$$

[Out] $-1/6*a^2*(1+1/a^2/x^4)^{(3/2)}-1/6/a/x^6$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6471, 30, 267}

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^5} dx = -\frac{1}{6}a^2 \left(\frac{1}{a^2x^4} + 1\right)^{3/2} - \frac{1}{6ax^6}$$

[In] $\text{Int}[E^{\text{ArcCsch}[a*x^2]}/x^5, x]$

[Out] $-1/6*(a^2*(1 + 1/(a^2*x^4))^{(3/2)}) - 1/(6*a*x^6)$

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}/(m+1), x] /; \text{FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 267

$\text{Int}[(x_)^{(m_.)}*((a_) + (b_.)*(x_)^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x^n)^{(p+1)}/(b*n*(p+1)), x] /; \text{FreeQ}[\{a, b, m, n, p\}, x] \ \&\& \ \text{EqQ}[m, n-1] \ \&\& \ \text{NeQ}[p, -1]$

Rule 6471

```
Int[E^ArcCsch[(a_.)*(x_)^(p_.)]*(x_)^(m_.), x_Symbol] := Dist[1/a, Int[x^(m
- p), x], x] + Int[x^m*Sqrt[1 + 1/(a^2*x^(2*p))], x] /; FreeQ[{a, m, p}, x
]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \frac{1}{x^7} dx}{a} + \int \frac{\sqrt{1 + \frac{1}{a^2 x^4}}}{x^5} dx \\ &= -\frac{1}{6} a^2 \left(1 + \frac{1}{a^2 x^4}\right)^{3/2} - \frac{1}{6 a x^6} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.26

$$\int \frac{e^{\text{csch}^{-1}(ax^2)}}{x^5} dx = -\frac{1 + a\sqrt{1 + \frac{1}{a^2 x^4}} x^2 (1 + a^2 x^4)}{6 a x^6}$$

[In] Integrate[E^ArcCsch[a*x^2]/x^5,x]

[Out] -1/6*(1 + a*Sqrt[1 + 1/(a^2*x^4)]*x^2*(1 + a^2*x^4))/(a*x^6)

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.35

method	result	size
default	$-\frac{\sqrt{\frac{x^4 a^2 + 1}{x^4 a^2}} (x^4 a^2 + 1)}{6 x^4} - \frac{1}{6 a x^6}$	42
trager	$-\frac{1}{6 x^6} - \frac{a (x^4 a^2 + 1) \sqrt{-\frac{x^4 a^2 - 1}{x^4 a^2}}}{6 x^4 a}$	46

[In] int((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^5,x,method=_RETURNVERBOSE)

[Out] -1/6*((a^2*x^4+1)/x^4/a^2)^(1/2)/x^4*(a^2*x^4+1)-1/6/a/x^6

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.58

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^5} dx = -\frac{a^3 x^6 + (a^3 x^6 + ax^2) \sqrt{\frac{a^2 x^4 + 1}{a^2 x^4}} + 1}{6 a x^6}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^5,x, algorithm="fricas")

[Out] -1/6*(a^3*x^6 + (a^3*x^6 + a*x^2)*sqrt((a^2*x^4 + 1)/(a^2*x^4)) + 1)/(a*x^6)

Sympy [A] (verification not implemented)

Time = 1.46 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.58

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^5} dx = \begin{cases} -a \left(\begin{cases} \sqrt{1 + \frac{1}{a^2 x^4}} \left(\frac{a^2}{3} + \frac{1}{3x^4} \right) & \text{for } \frac{1}{a^2} \neq 0 \\ \frac{1}{2x^4} & \text{otherwise} \end{cases} \right) - \frac{1}{3a^6} & \text{for } a \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

[In] integrate((1/a/x**2+(1+1/a**2/x**4)**(1/2))/x**5,x)

[Out] Piecewise((-a*Piecewise((sqrt(1 + 1/(a**2*x**4))*(a**2/3 + 1/(3*x**4))), Ne(a**(-2), 0)), (1/(2*x**4), True)) - 1/(3*x**6))/(2*a), Ne(a, 0)), (0, True))

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.81

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^5} dx = -\frac{1}{6} a^2 \left(\frac{1}{a^2 x^4} + 1 \right)^{\frac{3}{2}} - \frac{1}{6 a x^6}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^5,x, algorithm="maxima")

[Out] -1/6*a^2*(1/(a^2*x^4) + 1)^(3/2) - 1/6/(a*x^6)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 71 vs. 2(25) = 50.

Time = 0.27 (sec) , antiderivative size = 71, normalized size of antiderivative = 2.29

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^5} dx = \frac{2 \left(3 \left(x^2 |a| - \sqrt{a^2 x^4 + 1} \right)^4 a^4 + a^4 \right)}{\left(\left(x^2 |a| - \sqrt{a^2 x^4 + 1} \right)^2 - 1 \right)^3} - \frac{a}{x^6}$$

[In] integrate((1/a/x^2+(1+1/a^2/x^4)^(1/2))/x^5,x, algorithm="giac")

[Out] 1/6*(2*(3*(x^2*abs(a) - sqrt(a^2*x^4 + 1))^4*a^4 + a^4)/((x^2*abs(a) - sqrt(a^2*x^4 + 1))^2 - 1)^3 - a/x^6)/a^2

Mupad [B] (verification not implemented)

Time = 4.80 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.42

$$\int \frac{e^{\operatorname{csch}^{-1}(ax^2)}}{x^5} dx = -\frac{\frac{1}{6a} + \frac{x^2 \sqrt{\frac{1}{a^2 x^4} + 1}}{6}}{x^6} - \frac{a^2 \sqrt{\frac{1}{a^2 x^4} + 1}}{6}$$

[In] int(((1/(a^2*x^4) + 1)^(1/2) + 1/(a*x^2))/x^5,x)

[Out] - (1/(6*a) + (x^2*(1/(a^2*x^4) + 1)^(1/2))/6)/x^6 - (a^2*(1/(a^2*x^4) + 1)^(1/2))/6

3.48 $\int e^{2\operatorname{csch}^{-1}(ax)} x^m dx$

Optimal result	311
Rubi [A] (verified)	311
Mathematica [A] (verified)	312
Maple [F]	313
Fricas [F]	313
Sympy [A] (verification not implemented)	313
Maxima [F(-2)]	314
Giac [F(-2)]	314
Mupad [F(-1)]	314

Optimal result

Integrand size = 12, antiderivative size = 64

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^m dx = -\frac{2x^{-1+m}}{a^2(1-m)} + \frac{x^{1+m}}{1+m} + \frac{2x^m \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, -\frac{m}{2}, 1 - \frac{m}{2}, -\frac{1}{a^2x^2}\right)}{am}$$

[Out] $-2*x^{(-1+m)}/a^2/(1-m)+x^{(1+m)}/(1+m)+2*x^m*\operatorname{hypergeom}([-1/2, -1/2*m], [1-1/2*m], -1/a^2/x^2)/a/m$

Rubi [A] (verified)

Time = 0.23 (sec), antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6473, 6874, 346, 371}

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^m dx = \frac{2x^m \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, -\frac{m}{2}, 1 - \frac{m}{2}, -\frac{1}{a^2x^2}\right)}{am} - \frac{2x^{m-1}}{a^2(1-m)} + \frac{x^{m+1}}{m+1}$$

[In] $\operatorname{Int}[E^{(2*\operatorname{ArcCsch}[a*x])}*x^m, x]$

[Out] $(-2*x^{(-1+m)})/(a^2*(1-m)) + x^{(1+m)}/(1+m) + (2*x^m*\operatorname{Hypergeometric2F1}[-1/2, -1/2*m, 1-m/2, -(1/(a^2*x^2))])/(a*m)$

Rule 346

$\operatorname{Int}[(c_*)*(x_*)^{(m_*)}*((a_*) + (b_*)*(x_*)^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \operatorname{Dist}[(-c^{(-1)})*(c*x)^{(m+1)}*(1/x)^{(m+1)}, \operatorname{Subst}[\operatorname{Int}[(a + b/x^n)^p/x^{(m+2)}, x], x, 1/x], x] /; \operatorname{FreeQ}\{a, b, c, m, p\}, x \ \&\amp; \operatorname{ILtQ}[n, 0] \ \&\amp; \operatorname{!RationalQ}[m]$

Rule 371

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p
*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1
, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILt
Q[p, 0] || GtQ[a, 0])
```

Rule 6473

```
Int[E^(ArcCsch[u_]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[1 +
1/u^2])^n, x] /; FreeQ[m, x] && IntegerQ[n]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\sqrt{1 + \frac{1}{a^2 x^2}} + \frac{1}{ax} \right)^2 x^m dx \\
&= \int \left(\frac{2x^{-2+m}}{a^2} + \frac{2\sqrt{1 + \frac{1}{a^2 x^2}} x^{-1+m}}{a} + x^m \right) dx \\
&= -\frac{2x^{-1+m}}{a^2(1-m)} + \frac{x^{1+m}}{1+m} + \frac{2 \int \sqrt{1 + \frac{1}{a^2 x^2}} x^{-1+m} dx}{a} \\
&= -\frac{2x^{-1+m}}{a^2(1-m)} + \frac{x^{1+m}}{1+m} - \frac{(2(\frac{1}{x})^m x^m) \text{Subst}\left(\int x^{-1-m} \sqrt{1 + \frac{x^2}{a^2}} dx, x, \frac{1}{x}\right)}{a} \\
&= -\frac{2x^{-1+m}}{a^2(1-m)} + \frac{x^{1+m}}{1+m} + \frac{2x^m \text{Hypergeometric2F1}\left(-\frac{1}{2}, -\frac{m}{2}, 1 - \frac{m}{2}, -\frac{1}{a^2 x^2}\right)}{am}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.89

$$\int e^{2\text{csch}^{-1}(ax)} x^m dx = x^m \left(\frac{2}{a^2(-1+m)x} + \frac{x}{1+m} + \frac{2 \text{Hypergeometric2F1}\left(-\frac{1}{2}, -\frac{m}{2}, 1 - \frac{m}{2}, -\frac{1}{a^2 x^2}\right)}{am} \right)$$

[In] Integrate[E^(2*ArcCsch[a*x])*x^m,x]

[Out] $x^m \left(\frac{2}{a^2(-1+m)x} + \frac{x}{(1+m)} + \frac{(2 \text{Hypergeometric2F1}[-1/2, -1/2m, 1-m/2, -(1/(a^2x^2))])}{a^m} \right)$

Maple [F]

$$\int \left(\frac{1}{ax} + \sqrt{1 + \frac{1}{a^2x^2}} \right)^2 x^m dx$$

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^m,x)

[Out] int((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^m,x)

Fricas [F]

$$\int e^{2\text{csch}^{-1}(ax)} x^m dx = \int x^m \left(\sqrt{\frac{1}{a^2x^2} + 1} + \frac{1}{ax} \right)^2 dx$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^m,x, algorithm="fricas")

[Out] integral((2*a*x*x^m*sqrt((a^2*x^2 + 1)/(a^2*x^2)) + (a^2*x^2 + 2)*x^m)/(a^2*x^2), x)

Sympy [A] (verification not implemented)

Time = 3.10 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.19

$$\int e^{2\text{csch}^{-1}(ax)} x^m dx = \begin{cases} \frac{x^{m+1}}{m+1} & \text{for } m \neq -1 \\ \log(x) & \text{otherwise} \end{cases} - \frac{x^m \Gamma(-\frac{m}{2}) {}_2F_1\left(-\frac{1}{2}, -\frac{m}{2} \middle| 1 - \frac{m}{2} \middle| \frac{e^{i\pi}}{a^2x^2}\right)}{a \Gamma(1 - \frac{m}{2})} + \frac{2 \left(\begin{cases} \frac{x^m}{mx-x} & \text{for } m \neq 1 \\ \frac{x^m \log(x)}{x} & \text{otherwise} \end{cases} \right)}{a^2}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))**2*x**m,x)

[Out] Piecewise((x**(m + 1)/(m + 1), Ne(m, -1)), (log(x), True)) - x**m*gamma(-m/2)*hyper((-1/2, -m/2), (1 - m/2,), exp_polar(I*pi)/(a**2*x**2))/(a*gamma(1 - m/2)) + 2*Piecewise((x**m/(m*x - x), Ne(m, 1)), (x**m*log(x)/x, True))/a**2

Maxima [F(-2)]

Exception generated.

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^m dx = \text{Exception raised: ValueError}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^m,x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(m-2>0)', see 'assume?' for more details)Is

Giac [F(-2)]

Exception generated.

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^m dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^m,x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const in dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^m dx = \int x^m \left(\sqrt{\frac{1}{a^2 x^2} + 1} + \frac{1}{ax} \right)^2 dx$$

[In] int(x^m*((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))^2,x)

[Out] int(x^m*((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))^2, x)

3.49 $\int e^{2\operatorname{csch}^{-1}(ax)} x^4 dx$

Optimal result	315
Rubi [A] (verified)	315
Mathematica [A] (verified)	317
Maple [A] (verified)	317
Fricas [A] (verification not implemented)	318
Sympy [A] (verification not implemented)	318
Maxima [A] (verification not implemented)	318
Giac [A] (verification not implemented)	319
Mupad [B] (verification not implemented)	319

Optimal result

Integrand size = 12, antiderivative size = 85

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^4 dx = \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x^2}{4a^3} + \frac{2x^3}{3a^2} + \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x^4}{2a} + \frac{x^5}{5} - \frac{\operatorname{arctanh}\left(\sqrt{1 + \frac{1}{a^2 x^2}}\right)}{4a^5}$$

[Out] $2/3*x^3/a^2+1/5*x^5-1/4*\operatorname{arctanh}((1+1/a^2/x^2)^{(1/2)})/a^5+1/4*x^2*(1+1/a^2/x^2)^{(1/2)}/a^3+1/2*x^4*(1+1/a^2/x^2)^{(1/2)}/a$

Rubi [A] (verified)

Time = 0.19 (sec), antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {6473, 6874, 272, 43, 44, 65, 214}

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^4 dx = \frac{2x^3}{3a^2} + \frac{x^4 \sqrt{\frac{1}{a^2 x^2} + 1}}{2a} - \frac{\operatorname{arctanh}\left(\sqrt{\frac{1}{a^2 x^2} + 1}\right)}{4a^5} + \frac{x^2 \sqrt{\frac{1}{a^2 x^2} + 1}}{4a^3} + \frac{x^5}{5}$$

[In] $\operatorname{Int}[E^{(2*\operatorname{ArcCsch}[a*x])}*x^4, x]$

[Out] $(\operatorname{Sqrt}[1 + 1/(a^2*x^2)]*x^2)/(4*a^3) + (2*x^3)/(3*a^2) + (\operatorname{Sqrt}[1 + 1/(a^2*x^2)]*x^4)/(2*a) + x^5/5 - \operatorname{ArcTanh}[\operatorname{Sqrt}[1 + 1/(a^2*x^2)]]/(4*a^5)$

Rule 43

$\operatorname{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] := \operatorname{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + 1))), x] - \operatorname{Dist}[d*(n/(b*(m + 1))), \operatorname{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^{(n - 1)}, x], x] /;$
 $\operatorname{FreeQ}\{a, b, c, d, n\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{ILtQ}[m, -1] \&\& \operatorname{IntegerQ}[n] \&\& \operatorname{GtQ}[n, 0]$

Rule 44

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((
m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !Int
egerQ[n] && LtQ[n, 0]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 6473

```
Int[E^(ArcCsch[u_]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[1 +
1/u^2])^n, x] /; FreeQ[m, x] && IntegerQ[n]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\sqrt{1 + \frac{1}{a^2 x^2}} + \frac{1}{ax} \right)^2 x^4 dx \\ &= \int \left(\frac{2x^2}{a^2} + \frac{2\sqrt{1 + \frac{1}{a^2 x^2}} x^3}{a} + x^4 \right) dx \end{aligned}$$

$$\begin{aligned}
&= \frac{2x^3}{3a^2} + \frac{x^5}{5} + \frac{2 \int \sqrt{1 + \frac{1}{a^2 x^2}} x^3 dx}{a} \\
&= \frac{2x^3}{3a^2} + \frac{x^5}{5} - \frac{\text{Subst}\left(\int \frac{\sqrt{1 + \frac{x}{a^2}}}{x^3} dx, x, \frac{1}{x^2}\right)}{a} \\
&= \frac{2x^3}{3a^2} + \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x^4}{2a} + \frac{x^5}{5} - \frac{\text{Subst}\left(\int \frac{1}{x^2 \sqrt{1 + \frac{x}{a^2}}} dx, x, \frac{1}{x^2}\right)}{4a^3} \\
&= \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x^2}{4a^3} + \frac{2x^3}{3a^2} + \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x^4}{2a} + \frac{x^5}{5} + \frac{\text{Subst}\left(\int \frac{1}{x \sqrt{1 + \frac{x}{a^2}}} dx, x, \frac{1}{x^2}\right)}{8a^5} \\
&= \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x^2}{4a^3} + \frac{2x^3}{3a^2} + \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x^4}{2a} + \frac{x^5}{5} + \frac{\text{Subst}\left(\int \frac{1}{-a^2 + a^2 x^2} dx, x, \sqrt{1 + \frac{1}{a^2 x^2}}\right)}{4a^3} \\
&= \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x^2}{4a^3} + \frac{2x^3}{3a^2} + \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x^4}{2a} + \frac{x^5}{5} - \frac{\text{arctanh}\left(\sqrt{1 + \frac{1}{a^2 x^2}}\right)}{4a^5}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.99

$$\begin{aligned}
&\int e^{2\text{csch}^{-1}(ax)} x^4 dx \\
&= \frac{a^2 x^2 \left(15 \sqrt{1 + \frac{1}{a^2 x^2}} + 40ax + 30a^2 \sqrt{1 + \frac{1}{a^2 x^2}} x^2 + 12a^3 x^3 \right) - 15 \log\left(\left(1 + \sqrt{1 + \frac{1}{a^2 x^2}}\right) x\right)}{60a^5}
\end{aligned}$$

[In] Integrate[E^(2*ArcCsch[a*x])*x^4,x]

[Out] (a^2*x^2*(15*Sqrt[1 + 1/(a^2*x^2)] + 40*a*x + 30*a^2*Sqrt[1 + 1/(a^2*x^2)])*x^2 + 12*a^3*x^3) - 15*Log[(1 + Sqrt[1 + 1/(a^2*x^2)])*x]/(60*a^5)

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.49

method	result	size
default	$\frac{\frac{1}{5}a^2x^5 + \frac{1}{3}x^3}{a^2} - \frac{\sqrt{\frac{a^2x^2+1}{a^2}} x \left(-2x \left(\frac{a^2x^2+1}{a^2} \right)^{\frac{3}{2}} a^4 + x \sqrt{\frac{a^2x^2+1}{a^2}} a^2 + \ln \left(x + \sqrt{\frac{a^2x^2+1}{a^2}} \right) \right)}{4a^5 \sqrt{\frac{a^2x^2+1}{a^2}}} + \frac{x^3}{3a^2}$	127

[In] `int((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^4,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{a^2} \left(\frac{1}{5} a^2 x^5 + \frac{1}{3} x^3 \right) - \frac{1}{4} a^5 \left(\frac{(a^2 x^2 + 1)}{a^2 x^2} \right)^{1/2} x \left(-2 x \left(\frac{(a^2 x^2 + 1)}{a^2} \right)^{3/2} a^4 + x \left(\frac{(a^2 x^2 + 1)}{a^2} \right)^{1/2} a^2 + \ln \left(x \left(\frac{(a^2 x^2 + 1)}{a^2} \right)^{1/2} \right) \right) / \left(\frac{(a^2 x^2 + 1)}{a^2} \right)^{1/2} + \frac{1}{3} x^3 / a^2$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.02

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^4 dx = \frac{12 a^5 x^5 + 40 a^3 x^3 + 15 (2 a^4 x^4 + a^2 x^2) \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} + 15 \log \left(ax \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} - ax \right)}{60 a^5}$$

[In] `integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^4,x, algorithm="fricas")`

[Out] $\frac{1}{60} \left(12 a^5 x^5 + 40 a^3 x^3 + 15 (2 a^4 x^4 + a^2 x^2) \sqrt{(a^2 x^2 + 1) / (a^2 x^2)} + 15 \log(a x \sqrt{(a^2 x^2 + 1) / (a^2 x^2)} - a x) \right) / a^5$

Sympy [A] (verification not implemented)

Time = 2.61 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.96

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^4 dx = \frac{x^5}{5} + \frac{x^5}{2\sqrt{a^2 x^2 + 1}} + \frac{2x^3}{3a^2} + \frac{3x^3}{4a^2\sqrt{a^2 x^2 + 1}} + \frac{x}{4a^4\sqrt{a^2 x^2 + 1}} - \frac{\operatorname{asinh}(ax)}{4a^5}$$

[In] `integrate((1/a/x+(1+1/a**2/x**2)**(1/2))**2*x**4,x)`

[Out] $x^{**5}/5 + x^{**5}/(2*\sqrt{a^{**2}*x^{**2} + 1}) + 2*x^{**3}/(3*a^{**2}) + 3*x^{**3}/(4*a^{**2}*\sqrt{a^{**2}*x^{**2} + 1}) + x/(4*a^{**4}*\sqrt{a^{**2}*x^{**2} + 1}) - \operatorname{asinh}(a*x)/(4*a^{**5})$

Maxima [A] (verification not implemented)

none

Time = 0.21 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.38

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^4 dx = \frac{1}{5} x^5 + \frac{2x^3}{3a^2} + \frac{2 \left(\left(\frac{1}{a^2 x^2} + 1 \right)^{3/2} + \sqrt{\frac{1}{a^2 x^2} + 1} \right)}{a^4 \left(\frac{1}{a^2 x^2} + 1 \right)^2 - 2a^4 \left(\frac{1}{a^2 x^2} + 1 \right) + a^4} - \frac{\log \left(\sqrt{\frac{1}{a^2 x^2} + 1} + 1 \right)}{a^4} + \frac{\log \left(\sqrt{\frac{1}{a^2 x^2} + 1} - 1 \right)}{a^4}$$

$8a$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^4,x, algorithm="maxima")

[Out] 1/5*x^5 + 2/3*x^3/a^2 + 1/8*(2*((1/(a^2*x^2) + 1)^(3/2) + sqrt(1/(a^2*x^2) + 1)))/(a^4*(1/(a^2*x^2) + 1)^2 - 2*a^4*(1/(a^2*x^2) + 1) + a^4) - log(sqrt(1/(a^2*x^2) + 1) + 1)/a^4 + log(sqrt(1/(a^2*x^2) + 1) - 1)/a^4/a

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.94

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^4 dx = \frac{1}{4} \sqrt{a^2 x^2 + 1} x \left(\frac{2x^2 |a| \operatorname{sgn}(x)}{a^3} + \frac{|a| \operatorname{sgn}(x)}{a^5} \right) + \frac{3a^2 x^5 + 10x^3}{15a^2} + \frac{\log(-x|a| + \sqrt{a^2 x^2 + 1}) \operatorname{sgn}(x)}{4a^5}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^4,x, algorithm="giac")

[Out] 1/4*sqrt(a^2*x^2 + 1)*x*(2*x^2*abs(a)*sgn(x)/a^3 + abs(a)*sgn(x)/a^5) + 1/15*(3*a^2*x^5 + 10*x^3)/a^2 + 1/4*log(-x*abs(a) + sqrt(a^2*x^2 + 1))*sgn(x)/a^5

Mupad [B] (verification not implemented)

Time = 4.84 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.86

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^4 dx = \frac{x^5}{5} + \frac{2x^3}{3a^2} + \frac{x^4 \sqrt{\frac{1}{a^2 x^2} + 1}}{2a} + \frac{x^2 \sqrt{\frac{1}{a^2 x^2} + 1}}{4a^3} + \frac{\operatorname{atan}\left(\sqrt{\frac{1}{a^2 x^2} + 1}\right) \operatorname{li}}{4a^5}$$

[In] int(x^4*((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))^2,x)

[Out] (atan((1/(a^2*x^2) + 1)^(1/2)*1i)*1i)/(4*a^5) + x^5/5 + (2*x^3)/(3*a^2) + (x^4*(1/(a^2*x^2) + 1)^(1/2))/(2*a) + (x^2*(1/(a^2*x^2) + 1)^(1/2))/(4*a^3)

3.50 $\int e^{2\operatorname{csch}^{-1}(ax)} x^3 dx$

Optimal result	320
Rubi [A] (verified)	320
Mathematica [A] (verified)	321
Maple [A] (verified)	321
Fricas [A] (verification not implemented)	322
Sympy [A] (verification not implemented)	322
Maxima [A] (verification not implemented)	322
Giac [B] (verification not implemented)	323
Mupad [B] (verification not implemented)	323

Optimal result

Integrand size = 12, antiderivative size = 38

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^3 dx = \frac{x^2}{a^2} + \frac{2\left(1 + \frac{1}{a^2 x^2}\right)^{3/2} x^3}{3a} + \frac{x^4}{4}$$

[Out] $x^2/a^2 + 2/3*(1+1/a^2/x^2)^{(3/2)}*x^3/a + 1/4*x^4$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6473, 6874, 270}

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^3 dx = \frac{x^2}{a^2} + \frac{2x^3\left(\frac{1}{a^2 x^2} + 1\right)^{3/2}}{3a} + \frac{x^4}{4}$$

[In] `Int[E^(2*ArcCsch[a*x])*x^3,x]`

[Out] $x^2/a^2 + (2*(1 + 1/(a^2*x^2))^{(3/2)}*x^3)/(3*a) + x^4/4$

Rule 270

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m+1)*((a+b*x^n)^(p+1)/(a*c*(m+1))), x] /; FreeQ[{a, b, c, m, n, p}, x] && EqQ[(m+1)/n + p + 1, 0] && NeQ[m, -1]
```

Rule 6473

```
Int[E^(ArcCsch[u_]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[1 + 1/u^2])^n, x] /; FreeQ[m, x] && IntegerQ[n]
```


Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \left(\sqrt{1 + \frac{1}{a^2 x^2}} + \frac{1}{ax} \right)^2 x^3 dx \\
 &= \int \left(\frac{2x}{a^2} + \frac{2\sqrt{1 + \frac{1}{a^2 x^2}} x^2}{a} + x^3 \right) dx \\
 &= \frac{x^2}{a^2} + \frac{x^4}{4} + \frac{2 \int \sqrt{1 + \frac{1}{a^2 x^2}} x^2 dx}{a} \\
 &= \frac{x^2}{a^2} + \frac{2\left(1 + \frac{1}{a^2 x^2}\right)^{3/2} x^3}{3a} + \frac{x^4}{4}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.16

$$\int e^{2\text{csch}^{-1}(ax)} x^3 dx = \frac{x^2}{a^2} + \frac{x^4}{4} + \frac{2\sqrt{1 + \frac{1}{a^2 x^2}}(x + a^2 x^3)}{3a^3}$$

```
[In] Integrate[E^(2*ArcCsch[a*x])*x^3,x]
```

```
[Out] x^2/a^2 + x^4/4 + (2*Sqrt[1 + 1/(a^2*x^2)]*(x + a^2*x^3))/(3*a^3)
```

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.55

method	result	size
default	$\frac{(a^2 x^2 + 1)^2}{4a^4} + \frac{2\sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} x(a^2 x^2 + 1)}{3a^3} + \frac{x^2}{2a^2}$	59
trager	$\frac{(a^2 x^3 + a^2 x^2 + a^2 x + a^2 + 4x + 4)(x - 1)}{4} + \frac{2(a^2 x^2 + 1)x\sqrt{-\frac{a^2 x^2 - 1}{a^2 x^2}}}{3a}$	73

```
[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^3,x,method=_RETURNVERBOSE)
```

```
[Out] 1/4/a^4*(a^2*x^2+1)^2+2/3/a^3*((a^2*x^2+1)/a^2/x^2)^(1/2)*x*(a^2*x^2+1)+1/2*x^2/a^2
```

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.29

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^3 dx = \frac{3a^3x^4 + 12ax^2 + 8(a^2x^3 + x)\sqrt{\frac{a^2x^2+1}{a^2x^2}}}{12a^3}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^3,x, algorithm="fricas")

[Out] 1/12*(3*a^3*x^4 + 12*a*x^2 + 8*(a^2*x^3 + x)*sqrt((a^2*x^2 + 1)/(a^2*x^2)))/a^3

Sympy [A] (verification not implemented)

Time = 1.12 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.34

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^3 dx = \frac{x^4}{4} + \frac{2x^2\sqrt{a^2x^2+1}}{3a^2} + \frac{x^2}{a^2} + \frac{2\sqrt{a^2x^2+1}}{3a^4}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))**2*x**3,x)

[Out] x**4/4 + 2*x**2*sqrt(a**2*x**2 + 1)/(3*a**2) + x**2/a**2 + 2*sqrt(a**2*x**2 + 1)/(3*a**4)

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^3 dx = \frac{1}{4}x^4 + \frac{2x^3\left(\frac{1}{a^2x^2} + 1\right)^{\frac{3}{2}}}{3a} + \frac{x^2}{a^2}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^3,x, algorithm="maxima")

[Out] 1/4*x^4 + 2/3*x^3*(1/(a^2*x^2) + 1)^(3/2)/a + x^2/a^2

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 66 vs. $2(32) = 64$.

Time = 0.29 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.74

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^3 dx = \frac{a^2 x^2 + 1}{2a^4} - \frac{2|a|\operatorname{sgn}(x)}{3a^5} + \frac{8(a^2 x^2 + 1)^{\frac{3}{2}} a^2 |a|\operatorname{sgn}(x) + 3(a^2 x^2 + 1)^2 a^3}{12a^7}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^3,x, algorithm="giac")

[Out] 1/2*(a^2*x^2 + 1)/a^4 - 2/3*abs(a)*sgn(x)/a^5 + 1/12*(8*(a^2*x^2 + 1)^(3/2)*a^2*abs(a)*sgn(x) + 3*(a^2*x^2 + 1)^2*a^3)/a^7

Mupad [B] (verification not implemented)

Time = 5.23 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.05

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^3 dx = \sqrt{\frac{1}{a^2 x^2} + 1} \left(\frac{2x}{3a^3} + \frac{2x^3}{3a} \right) + \frac{x^4}{4} + \frac{x^2}{a^2}$$

[In] int(x^3*((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))^2,x)

[Out] (1/(a^2*x^2) + 1)^(1/2)*((2*x)/(3*a^3) + (2*x^3)/(3*a)) + x^4/4 + x^2/a^2

3.51 $\int e^{2\operatorname{csch}^{-1}(ax)} x^2 dx$

Optimal result	324
Rubi [A] (verified)	324
Mathematica [A] (verified)	326
Maple [B] (verified)	326
Fricas [A] (verification not implemented)	327
Sympy [A] (verification not implemented)	327
Maxima [A] (verification not implemented)	327
Giac [A] (verification not implemented)	328
Mupad [B] (verification not implemented)	328

Optimal result

Integrand size = 12, antiderivative size = 52

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^2 dx = \frac{2x}{a^2} + \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x^2}{a} + \frac{x^3}{3} + \frac{\operatorname{arctanh}\left(\sqrt{1 + \frac{1}{a^2 x^2}}\right)}{a^3}$$

[Out] $2*x/a^2+1/3*x^3+\operatorname{arctanh}((1+1/a^2/x^2)^{(1/2)})/a^3+x^2*(1+1/a^2/x^2)^{(1/2)}/a$

Rubi [A] (verified)

Time = 0.17 (sec), antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6473, 6874, 272, 43, 65, 214}

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^2 dx = \frac{x^2 \sqrt{\frac{1}{a^2 x^2} + 1}}{a} + \frac{2x}{a^2} + \frac{\operatorname{arctanh}\left(\sqrt{\frac{1}{a^2 x^2} + 1}\right)}{a^3} + \frac{x^3}{3}$$

[In] `Int[E^(2*ArcCsch[a*x])*x^2,x]`

[Out] $(2*x)/a^2 + (\operatorname{Sqrt}[1 + 1/(a^2*x^2)]*x^2)/a + x^3/3 + \operatorname{ArcTanh}[\operatorname{Sqrt}[1 + 1/(a^2*x^2)]]/a^3$

Rule 43

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Dist[d*(n/(b*(m + 1))), I
nt[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !IntegerQ[n] && GtQ[n, 0]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 6473

```
Int[E^(ArcCsch[u_]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[1 +
1/u^2])^n, x] /; FreeQ[m, x] && IntegerQ[n]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\sqrt{1 + \frac{1}{a^2 x^2}} + \frac{1}{ax} \right)^2 x^2 dx \\
&= \int \left(\frac{2}{a^2} + \frac{2\sqrt{1 + \frac{1}{a^2 x^2}} x}{a} + x^2 \right) dx \\
&= \frac{2x}{a^2} + \frac{x^3}{3} + \frac{2 \int \sqrt{1 + \frac{1}{a^2 x^2}} dx}{a} \\
&= \frac{2x}{a^2} + \frac{x^3}{3} - \frac{\text{Subst} \left(\int \frac{\sqrt{1 + \frac{x}{a^2}}}{x^2} dx, x, \frac{1}{x^2} \right)}{a} \\
&= \frac{2x}{a^2} + \frac{\sqrt{1 + \frac{1}{a^2 x^2}} x^2}{a} + \frac{x^3}{3} - \frac{\text{Subst} \left(\int \frac{1}{x \sqrt{1 + \frac{x}{a^2}}} dx, x, \frac{1}{x^2} \right)}{2a^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2x}{a^2} + \frac{\sqrt{1 + \frac{1}{a^2x^2}}x^2}{a} + \frac{x^3}{3} - \frac{\text{Subst}\left(\int \frac{1}{-a^2+a^2x^2} dx, x, \sqrt{1 + \frac{1}{a^2x^2}}\right)}{a} \\
&= \frac{2x}{a^2} + \frac{\sqrt{1 + \frac{1}{a^2x^2}}x^2}{a} + \frac{x^3}{3} + \frac{\text{arctanh}\left(\sqrt{1 + \frac{1}{a^2x^2}}\right)}{a^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.10

$$\int e^{2\text{csch}^{-1}(ax)} x^2 dx = \frac{ax\left(6 + 3a\sqrt{1 + \frac{1}{a^2x^2}}x + a^2x^2\right) + 3\log\left(\left(1 + \sqrt{1 + \frac{1}{a^2x^2}}\right)x\right)}{3a^3}$$

[In] Integrate[E^(2*ArcCsch[a*x])*x^2,x]

[Out] (a*x*(6 + 3*a*Sqrt[1 + 1/(a^2*x^2)]*x + a^2*x^2) + 3*Log[(1 + Sqrt[1 + 1/(a^2*x^2)])*x])/(3*a^3)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 97 vs. 2(46) = 92.

Time = 0.05 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.88

method	result	size
default	$\frac{\frac{1}{3}a^2x^3+x}{a^2} + \frac{\sqrt{\frac{a^2x^2+1}{a^2x^2}}x\left(x\sqrt{\frac{a^2x^2+1}{a^2}}a^2+\ln\left(x+\sqrt{\frac{a^2x^2+1}{a^2}}\right)\right)}{a^3\sqrt{\frac{a^2x^2+1}{a^2}}} + \frac{x}{a^2}$	98

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^2,x,method=_RETURNVERBOSE)

[Out] 1/a^2*(1/3*a^2*x^3+x)+1/a^3*((a^2*x^2+1)/a^2/x^2)^(1/2)*x*(x*((a^2*x^2+1)/a^2)^(1/2)*a^2+ln(x+((a^2*x^2+1)/a^2)^(1/2)))/((a^2*x^2+1)/a^2)^(1/2)+x/a^2

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.38

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^2 dx = \frac{a^3 x^3 + 3 a^2 x^2 \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} + 6 a x - 3 \log \left(a x \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} - a x \right)}{3 a^3}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^2,x, algorithm="fricas")

[Out] 1/3*(a^3*x^3 + 3*a^2*x^2*sqrt((a^2*x^2 + 1)/(a^2*x^2)) + 6*a*x - 3*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x))/a^3

Sympy [A] (verification not implemented)

Time = 1.61 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.69

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^2 dx = \frac{x^3}{3} + \frac{x\sqrt{a^2 x^2 + 1}}{a^2} + \frac{2x}{a^2} + \frac{\operatorname{asinh}(ax)}{a^3}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))**2*x**2,x)

[Out] x**3/3 + x*sqrt(a**2*x**2 + 1)/a**2 + 2*x/a**2 + asinh(a*x)/a**3

Maxima [A] (verification not implemented)

none

Time = 0.21 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.71

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^2 dx = \frac{1}{3} x^3 + \frac{\frac{2\sqrt{\frac{1}{a^2 x^2} + 1}}{\left(\frac{1}{a^2 x^2} + 1\right)^{-a^2}} + \frac{\log\left(\sqrt{\frac{1}{a^2 x^2} + 1} + 1\right)}{a^2} - \frac{\log\left(\sqrt{\frac{1}{a^2 x^2} + 1} - 1\right)}{a^2}}{2 a} + \frac{2 x}{a^2}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^2,x, algorithm="maxima")

[Out] 1/3*x^3 + 1/2*(2*sqrt(1/(a^2*x^2) + 1)/(a^2*(1/(a^2*x^2) + 1) - a^2) + log(sqrt(1/(a^2*x^2) + 1) + 1)/a^2 - log(sqrt(1/(a^2*x^2) + 1) - 1)/a^2)/a + 2*x/a^2

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.19

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^2 dx = \frac{\sqrt{a^2 x^2 + 1} x |a| \operatorname{sgn}(x)}{a^3} + \frac{a^2 x^3 + 6x}{3a^2} - \frac{\log(-x|a| + \sqrt{a^2 x^2 + 1}) \operatorname{sgn}(x)}{a^3}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x^2,x, algorithm="giac")

[Out] sqrt(a^2*x^2 + 1)*x*abs(a)*sgn(x)/a^3 + 1/3*(a^2*x^3 + 6*x)/a^2 - log(-x*abs(a) + sqrt(a^2*x^2 + 1))*sgn(x)/a^3

Mupad [B] (verification not implemented)

Time = 5.06 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.98

$$\int e^{2\operatorname{csch}^{-1}(ax)} x^2 dx = \frac{2x}{a^2} + \frac{x^3}{3} + \frac{x^2 \sqrt{\frac{1}{a^2 x^2} + 1}}{a} - \frac{\operatorname{atan}\left(\sqrt{\frac{1}{a^2 x^2} + 1} \operatorname{li}\right) \operatorname{li}}{a^3}$$

[In] int(x^2*((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))^2,x)

[Out] (2*x)/a^2 - (atan((1/(a^2*x^2) + 1)^(1/2)*1i)*1i)/a^3 + x^3/3 + (x^2*(1/(a^2*x^2) + 1)^(1/2))/a

3.52 $\int e^{2\operatorname{csch}^{-1}(ax)} x dx$

Optimal result	329
Rubi [A] (verified)	329
Mathematica [A] (verified)	331
Maple [B] (verified)	331
Fricas [B] (verification not implemented)	331
Sympy [A] (verification not implemented)	332
Maxima [A] (verification not implemented)	332
Giac [B] (verification not implemented)	332
Mupad [B] (verification not implemented)	333

Optimal result

Integrand size = 10, antiderivative size = 43

$$\int e^{2\operatorname{csch}^{-1}(ax)} x dx = \frac{2\sqrt{1 + \frac{1}{a^2x^2}}x}{a} + \frac{x^2}{2} - \frac{2\operatorname{csch}^{-1}(ax)}{a^2} + \frac{2\log(x)}{a^2}$$

[Out] $1/2*x^2-2*\operatorname{arccsch}(a*x)/a^2+2*\ln(x)/a^2+2*x*(1+1/a^2/x^2)^{(1/2)}/a$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6473, 6874, 248, 283, 221}

$$\int e^{2\operatorname{csch}^{-1}(ax)} x dx = \frac{2x\sqrt{\frac{1}{a^2x^2} + 1}}{a} + \frac{2\log(x)}{a^2} - \frac{2\operatorname{csch}^{-1}(ax)}{a^2} + \frac{x^2}{2}$$

[In] $\operatorname{Int}[E^{(2*\operatorname{ArcCsch}[a*x])}*x, x]$

[Out] $(2*\operatorname{Sqrt}[1 + 1/(a^2*x^2)]*x)/a + x^2/2 - (2*\operatorname{ArcCsch}[a*x])/a^2 + (2*\operatorname{Log}[x])/a^2$

Rule 221

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_) + (b_)*(x_)^2], x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{ArcSinh}[\operatorname{Rt}[b, 2]*(x/\operatorname{Sqrt}[a])]/\operatorname{Rt}[b, 2], x] /; \operatorname{FreeQ}[\{a, b\}, x] \ \&\& \operatorname{GtQ}[a, 0] \ \&\& \operatorname{PosQ}[b]$

Rule 248

$\operatorname{Int}[(a_) + (b_)*(x_)^{(n_)}]^{(p_)}, x_Symbol] \rightarrow -\operatorname{Subst}[\operatorname{Int}[(a + b/x^n)^p/x^2, x], x, 1/x] /; \operatorname{FreeQ}[\{a, b, p\}, x] \ \&\& \operatorname{ILtQ}[n, 0]$

Rule 283

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^p/(c^(n*(m + 1))), x] - Dist[b*n*(p/(c^n*(m + 1))), Int[(c*x)^(m + n)*(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && GtQ[p, 0] && LtQ[m, -1] && !ILtQ[(m + n*p + n + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 6473

```
Int[E^(ArcSch[u_]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[1 + 1/u^2])^n, x] /; FreeQ[m, x] && IntegerQ[n]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\sqrt{1 + \frac{1}{a^2 x^2}} + \frac{1}{ax} \right)^2 x \, dx \\
&= \int \left(\frac{2\sqrt{1 + \frac{1}{a^2 x^2}}}{a} + \frac{2}{a^2 x} + x \right) dx \\
&= \frac{x^2}{2} + \frac{2 \log(x)}{a^2} + \frac{2 \int \sqrt{1 + \frac{1}{a^2 x^2}} dx}{a} \\
&= \frac{x^2}{2} + \frac{2 \log(x)}{a^2} - \frac{2 \text{Subst} \left(\int \frac{\sqrt{1 + \frac{x^2}{a^2}}}{x^2} dx, x, \frac{1}{x} \right)}{a} \\
&= \frac{2\sqrt{1 + \frac{1}{a^2 x^2}} x}{a} + \frac{x^2}{2} + \frac{2 \log(x)}{a^2} - \frac{2 \text{Subst} \left(\int \frac{1}{\sqrt{1 + \frac{x^2}{a^2}}} dx, x, \frac{1}{x} \right)}{a^3} \\
&= \frac{2\sqrt{1 + \frac{1}{a^2 x^2}} x}{a} + \frac{x^2}{2} - \frac{2 \text{csch}^{-1}(ax)}{a^2} + \frac{2 \log(x)}{a^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.02

$$\int e^{2\operatorname{csch}^{-1}(ax)} x dx = \frac{ax \left(4\sqrt{1 + \frac{1}{a^2 x^2}} + ax \right) - 4\operatorname{arcsinh}\left(\frac{1}{ax}\right) + 4\log(x)}{2a^2}$$

[In] Integrate[E^(2*ArcCsch[a*x])*x,x]

[Out] (a*x*(4*sqrt[1 + 1/(a^2*x^2)] + a*x) - 4*ArcSinh[1/(a*x)] + 4*Log[x])/(2*a^2)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 128 vs. 2(39) = 78.

Time = 0.06 (sec) , antiderivative size = 129, normalized size of antiderivative = 3.00

method	result	size
default	$\frac{\frac{a^2 x^2}{2} + \ln(x)}{a^2} + \frac{2\sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} x \left(\sqrt{\frac{1}{a^2}} \sqrt{\frac{a^2 x^2 + 1}{a^2}} a^2 - \ln\left(\frac{2\sqrt{\frac{1}{a^2}} \sqrt{\frac{a^2 x^2 + 1}{a^2}} a^2 + 2\right)}{\frac{a^2 x}{a^2}}\right)}{a^3 \sqrt{\frac{a^2 x^2 + 1}{a^2}} \sqrt{\frac{1}{a^2}}} + \frac{\ln(x)}{a^2}$	129

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x,x,method=_RETURNVERBOSE)

[Out] 1/a^2*(1/2*a^2*x^2+ln(x))+2/a^3*((a^2*x^2+1)/a^2/x^2)^(1/2)*x*((1/a^2)^(1/2))*((a^2*x^2+1)/a^2)^(1/2)*a^2-ln(2*((1/a^2)^(1/2))*((a^2*x^2+1)/a^2)^(1/2)*a^2+1)/x/a^2)/((a^2*x^2+1)/a^2)^(1/2)/(1/a^2)^(1/2)+ln(x)/a^2

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 99 vs. 2(39) = 78.

Time = 0.26 (sec) , antiderivative size = 99, normalized size of antiderivative = 2.30

$$\int e^{2\operatorname{csch}^{-1}(ax)} x dx = \frac{a^2 x^2 + 4ax\sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} - 4\log\left(ax\sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} - ax + 1\right) + 4\log\left(ax\sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} - ax - 1\right) + 4\log(x)}{2a^2}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x,x, algorithm="fricas")

[Out] 1/2*(a^2*x^2 + 4*a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - 4*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x + 1) + 4*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x - 1) + 4*log(x))/a^2

Sympy [A] (verification not implemented)

Time = 1.73 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.47

$$\int e^{2\operatorname{csch}^{-1}(ax)} x dx = \frac{x^2}{2} + \frac{2x}{a\sqrt{1 + \frac{1}{a^2x^2}}} + \frac{2\log(x)}{a^2} - \frac{2\operatorname{asinh}\left(\frac{1}{ax}\right)}{a^2} + \frac{2}{a^3x\sqrt{1 + \frac{1}{a^2x^2}}}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))**2*x,x)

[Out] x**2/2 + 2*x/(a*sqrt(1 + 1/(a**2*x**2))) + 2*log(x)/a**2 - 2*asinh(1/(a*x))/a**2 + 2/(a**3*x*sqrt(1 + 1/(a**2*x**2)))

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.74

$$\int e^{2\operatorname{csch}^{-1}(ax)} x dx = \frac{1}{2}x^2 + \frac{2x\sqrt{\frac{1}{a^2x^2} + 1} - \frac{\log(ax\sqrt{\frac{1}{a^2x^2} + 1})}{a} + \frac{\log(ax\sqrt{\frac{1}{a^2x^2} + 1})}{a}}{a} + \frac{2\log(x)}{a^2}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x,x, algorithm="maxima")

[Out] 1/2*x^2 + (2*x*sqrt(1/(a^2*x^2) + 1) - log(a*x*sqrt(1/(a^2*x^2) + 1) + 1)/a + log(a*x*sqrt(1/(a^2*x^2) + 1) - 1)/a)/a + 2*log(x)/a^2

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 82 vs. 2(39) = 78.

Time = 0.28 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.91

$$\int e^{2\operatorname{csch}^{-1}(ax)} x dx = \frac{4\sqrt{a^2x^2 + 1}|a|\operatorname{sgn}(x) + (a^2x^2 + 1)a - 2(|a|\operatorname{sgn}(x) - a)\log(\sqrt{a^2x^2 + 1} + 1) + 2(|a|\operatorname{sgn}(x) + a)\log(\sqrt{a^2x^2 + 1} - 1)}{2a^3}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2*x,x, algorithm="giac")

[Out] 1/2*(4*sqrt(a^2*x^2 + 1)*abs(a)*sgn(x) + (a^2*x^2 + 1)*a - 2*(abs(a)*sgn(x) - a)*log(sqrt(a^2*x^2 + 1) + 1) + 2*(abs(a)*sgn(x) + a)*log(sqrt(a^2*x^2 + 1) - 1))/a^3

Mupad [B] (verification not implemented)

Time = 5.09 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.21

$$\int e^{2\operatorname{csch}^{-1}(ax)} x \, dx = \frac{x^2}{2} - \frac{2 \ln\left(\frac{1}{x}\right)}{a^2} + \frac{2x \sqrt{\frac{1}{a^2 x^2} + 1}}{a} - \frac{2 \operatorname{asinh}\left(\frac{\sqrt{\frac{1}{a^2}}}{x}\right)}{a^3 \sqrt{\frac{1}{a^2}}}$$

[In] int(x*((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))^2,x)

[Out] x^2/2 - (2*log(1/x))/a^2 + (2*x*(1/(a^2*x^2) + 1)^(1/2))/a - (2*asinh((1/a^2)^(1/2)/x))/(a^3*(1/a^2)^(1/2))

3.53 $\int e^{2\operatorname{csch}^{-1}(ax)} dx$

Optimal result	334
Rubi [A] (verified)	334
Mathematica [A] (verified)	336
Maple [B] (verified)	336
Fricas [A] (verification not implemented)	337
Sympy [A] (verification not implemented)	337
Maxima [A] (verification not implemented)	337
Giac [F(-2)]	338
Mupad [B] (verification not implemented)	338

Optimal result

Integrand size = 8, antiderivative size = 47

$$\int e^{2\operatorname{csch}^{-1}(ax)} dx = -\frac{2\sqrt{1 + \frac{1}{a^2x^2}}}{a} - \frac{2}{a^2x} + x + \frac{2\operatorname{arctanh}\left(\sqrt{1 + \frac{1}{a^2x^2}}\right)}{a}$$

[Out] $-2/a^2/x+x+2*\operatorname{arctanh}((1+1/a^2/x^2)^{(1/2)})/a-2*(1+1/a^2/x^2)^{(1/2)}/a$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6468, 6874, 272, 52, 65, 214}

$$\int e^{2\operatorname{csch}^{-1}(ax)} dx = \frac{2\operatorname{arctanh}\left(\sqrt{\frac{1}{a^2x^2} + 1}\right)}{a} - \frac{2\sqrt{\frac{1}{a^2x^2} + 1}}{a} - \frac{2}{a^2x} + x$$

[In] $\operatorname{Int}[E^{(2*\operatorname{ArcCsch}[a*x])}, x]$

[Out] $(-2*\operatorname{Sqrt}[1 + 1/(a^2*x^2)])/a - 2/(a^2*x) + x + (2*\operatorname{ArcTanh}[\operatorname{Sqrt}[1 + 1/(a^2*x^2)]])/a$

Rule 52

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(
b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 6468

```
Int[E^(ArcCsch[u_]*(n_.)), x_Symbol] := Int[(1/u + Sqrt[1 + 1/u^2])^n, x] /
; IntegerQ[n]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\sqrt{1 + \frac{1}{a^2 x^2}} + \frac{1}{ax} \right)^2 dx \\
&= \int \left(1 + \frac{2}{a^2 x^2} + \frac{2\sqrt{1 + \frac{1}{a^2 x^2}}}{ax} \right) dx \\
&= -\frac{2}{a^2 x} + x + \frac{2 \int \frac{\sqrt{1 + \frac{1}{a^2 x^2}}}{x} dx}{a} \\
&= -\frac{2}{a^2 x} + x - \frac{\text{Subst} \left(\int \frac{\sqrt{1 + \frac{x}{a^2}}}{x} dx, x, \frac{1}{x^2} \right)}{a}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2\sqrt{1+\frac{1}{a^2x^2}}}{a} - \frac{2}{a^2x} + x - \frac{\text{Subst}\left(\int \frac{1}{x\sqrt{1+\frac{x}{a^2}}} dx, x, \frac{1}{x^2}\right)}{a} \\
&= -\frac{2\sqrt{1+\frac{1}{a^2x^2}}}{a} - \frac{2}{a^2x} + x - (2a)\text{Subst}\left(\int \frac{1}{-a^2+a^2x^2} dx, x, \sqrt{1+\frac{1}{a^2x^2}}\right) \\
&= -\frac{2\sqrt{1+\frac{1}{a^2x^2}}}{a} - \frac{2}{a^2x} + x + \frac{2\text{arctanh}\left(\sqrt{1+\frac{1}{a^2x^2}}\right)}{a}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.11

$$\int e^{2\text{csch}^{-1}(ax)} dx = -\frac{2\sqrt{1+\frac{1}{a^2x^2}}}{a} - \frac{2}{a^2x} + x + \frac{2\log\left(a\left(1+\sqrt{1+\frac{1}{a^2x^2}}\right)x\right)}{a}$$

[In] Integrate[E^(2*ArcCsch[a*x]),x]

[Out] (-2*Sqrt[1 + 1/(a^2*x^2)]/a - 2/(a^2*x) + x + (2*Log[a*(1 + Sqrt[1 + 1/(a^2*x^2)])*x])/a

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 112 vs. 2(43) = 86.

Time = 0.06 (sec) , antiderivative size = 113, normalized size of antiderivative = 2.40

method	result	size
default	$x - \frac{2}{x a^2} - \frac{2\sqrt{\frac{a^2x^2+1}{a^2x^2}} \left(a^2 \left(\frac{a^2x^2+1}{a^2} \right)^{\frac{3}{2}} - \sqrt{\frac{a^2x^2+1}{a^2}} a^2x^2 - \ln \left(x + \sqrt{\frac{a^2x^2+1}{a^2}} x \right) \right)}{a\sqrt{\frac{a^2x^2+1}{a^2}}}$	113

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))^2,x,method=_RETURNVERBOSE)

[Out] x-2/x/a^2-2/a*((a^2*x^2+1)/a^2/x^2)^(1/2)*(a^2*((a^2*x^2+1)/a^2)^(3/2)-((a^2*x^2+1)/a^2)^(1/2)*a^2*x^2-ln(x+((a^2*x^2+1)/a^2)^(1/2))*x)/((a^2*x^2+1)/a^2)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.55

$$\int e^{2\operatorname{csch}^{-1}(ax)} dx = \frac{a^2 x^2 - 2ax \log\left(ax \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} - ax\right) - 2ax \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} - 2ax - 2}{a^2 x}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2,x, algorithm="fricas")

[Out] (a^2*x^2 - 2*a*x*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x) - 2*a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - 2*a*x - 2)/(a^2*x)

Sympy [A] (verification not implemented)

Time = 1.71 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.04

$$\int e^{2\operatorname{csch}^{-1}(ax)} dx = x - \frac{2x}{\sqrt{a^2 x^2 + 1}} + \frac{2 \operatorname{asinh}(ax)}{a} - \frac{2}{a^2 x} - \frac{2}{a^2 x \sqrt{a^2 x^2 + 1}}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))**2,x)

[Out] x - 2*x/sqrt(a**2*x**2 + 1) + 2*asinh(a*x)/a - 2/(a**2*x) - 2/(a**2*x*sqrt(a**2*x**2 + 1))

Maxima [A] (verification not implemented)

none

Time = 0.21 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.26

$$\int e^{2\operatorname{csch}^{-1}(ax)} dx = x - \frac{2 \sqrt{\frac{1}{a^2 x^2} + 1} - \log\left(\sqrt{\frac{1}{a^2 x^2} + 1} + 1\right) + \log\left(\sqrt{\frac{1}{a^2 x^2} + 1} - 1\right)}{a} - \frac{2}{a^2 x}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2,x, algorithm="maxima")

[Out] x - (2*sqrt(1/(a^2*x^2) + 1) - log(sqrt(1/(a^2*x^2) + 1) + 1) + log(sqrt(1/(a^2*x^2) + 1) - 1))/a - 2/(a^2*x)

Giac [F(-2)]

Exception generated.

$$\int e^{2\operatorname{csch}^{-1}(ax)} dx = \text{Exception raised: TypeError}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT>Error: Bad Argument Type

Mupad [B] (verification not implemented)

Time = 4.92 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00

$$\int e^{2\operatorname{csch}^{-1}(ax)} dx = x - \frac{2\sqrt{\frac{1}{a^2x^2} + 1}}{a} - \frac{2}{a^2x} - \frac{\operatorname{atan}\left(\sqrt{\frac{1}{a^2x^2} + 1} \operatorname{li}\right) 2i}{a}$$

[In] int(((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))^2,x)

[Out] x - (atan((1/(a^2*x^2) + 1)^(1/2)*1i)*2i)/a - (2*(1/(a^2*x^2) + 1)^(1/2))/a
- 2/(a^2*x)

3.54 $\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x} dx$

Optimal result	339
Rubi [A] (verified)	339
Mathematica [A] (verified)	341
Maple [B] (verified)	341
Fricas [B] (verification not implemented)	341
Sympy [B] (verification not implemented)	342
Maxima [B] (verification not implemented)	342
Giac [B] (verification not implemented)	343
Mupad [B] (verification not implemented)	343

Optimal result

Integrand size = 12, antiderivative size = 38

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x} dx = -\frac{1}{a^2x^2} - \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{ax} - \operatorname{csch}^{-1}(ax) + \log(x)$$

[Out] $-1/a^2/x^2 - \operatorname{arccsch}(a*x) + \ln(x) - (1 + 1/a^2/x^2)^{(1/2)}/a/x$

Rubi [A] (verified)

Time = 0.16 (sec), antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {6473, 6874, 342, 201, 221}

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x} dx = -\frac{\sqrt{\frac{1}{a^2x^2} + 1}}{ax} - \frac{1}{a^2x^2} - \operatorname{csch}^{-1}(ax) + \log(x)$$

[In] $\operatorname{Int}[E^{(2*\operatorname{ArcCsch}[a*x])}/x, x]$

[Out] $-(1/(a^2*x^2)) - \operatorname{Sqrt}[1 + 1/(a^2*x^2)]/(a*x) - \operatorname{ArcCsch}[a*x] + \operatorname{Log}[x]$

Rule 201

$\operatorname{Int}[(a + b*x^n)^p, x_Symbol] \rightarrow \operatorname{Simp}[x*(a + b*x^n)^p/(n*p + 1), x] + \operatorname{Dist}[a*n*(p/(n*p + 1)), \operatorname{Int}[(a + b*x^n)^{p-1}, x], x] /;$ Free Q[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p])) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]]

Rule 221

`Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

Rule 342

`Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]`

Rule 6473

`Int[E^(ArcCsch[u_]*(n_))*(x_)^(m_), x_Symbol] := Int[x^m*(1/u + Sqrt[1 + 1/u^2])^n, x] /; FreeQ[m, x] && IntegerQ[n]`

Rule 6874

`Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{\left(\sqrt{1 + \frac{1}{a^2x^2}} + \frac{1}{ax}\right)^2}{x} dx \\
 &= \int \left(\frac{2}{a^2x^3} + \frac{2\sqrt{1 + \frac{1}{a^2x^2}}}{ax^2} + \frac{1}{x}\right) dx \\
 &= -\frac{1}{a^2x^2} + \log(x) + \frac{2 \int \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{x^2} dx}{a} \\
 &= -\frac{1}{a^2x^2} + \log(x) - \frac{2 \text{Subst}\left(\int \sqrt{1 + \frac{x^2}{a^2}} dx, x, \frac{1}{x}\right)}{a} \\
 &= -\frac{1}{a^2x^2} - \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{ax} + \log(x) - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1 + \frac{x^2}{a^2}}} dx, x, \frac{1}{x}\right)}{a} \\
 &= -\frac{1}{a^2x^2} - \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{ax} - \text{csch}^{-1}(ax) + \log(x)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.03

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x} dx = -\frac{1 + a\sqrt{1 + \frac{1}{a^2x^2}}}{a^2x^2} - \operatorname{arcsinh}\left(\frac{1}{ax}\right) + \log(x)$$

[In] Integrate[E^(2*ArcCsch[a*x])/x,x]

[Out] -((1 + a*sqrt[1 + 1/(a^2*x^2)]*x)/(a^2*x^2)) - ArcSinh[1/(a*x)] + Log[x]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 163 vs. 2(36) = 72.

Time = 0.06 (sec) , antiderivative size = 164, normalized size of antiderivative = 4.32

method	result	size
default	$\frac{a^2 \ln(x) - \frac{1}{2x^2}}{a^2} - \frac{\sqrt{\frac{a^2x^2+1}{a^2x^2}} \left(a^2 \left(\frac{a^2x^2+1}{a^2} \right)^{\frac{3}{2}} \sqrt{\frac{1}{a^2} - \sqrt{\frac{a^2x^2+1}{a^2}}} \sqrt{\frac{1}{a^2}} a^2x^2 + \ln \left(\frac{2\sqrt{\frac{1}{a^2}} \sqrt{\frac{a^2x^2+1}{a^2}} a^2+2}{a^2x} \right) x^2 \right)}{ax\sqrt{\frac{1}{a^2}} \sqrt{\frac{a^2x^2+1}{a^2}}} - \frac{1}{2a^2x^2}$	164

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x,x,method=_RETURNVERBOSE)

[Out] 1/a^2*(a^2*ln(x)-1/2/x^2)-1/a*((a^2*x^2+1)/a^2/x^2)^(1/2)/x*(a^2*((a^2*x^2+1)/a^2)^(3/2)*(1/a^2)^(1/2)-((a^2*x^2+1)/a^2)^(1/2)*(1/a^2)^(1/2)*a^2*x^2+1
n(2*((1/a^2)^(1/2)*((a^2*x^2+1)/a^2)^(1/2)*a^2+1)/x/a^2*x^2)/(1/a^2)^(1/2)
/((a^2*x^2+1)/a^2)^(1/2)-1/2/a^2/x^2

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 112 vs. 2(36) = 72.

Time = 0.24 (sec) , antiderivative size = 112, normalized size of antiderivative = 2.95

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x} dx = \frac{a^2x^2 \log\left(ax\sqrt{\frac{a^2x^2+1}{a^2x^2}} - ax + 1\right) - a^2x^2 \log\left(ax\sqrt{\frac{a^2x^2+1}{a^2x^2}} - ax - 1\right) - a^2x^2 \log(x) + ax\sqrt{\frac{a^2x^2+1}{a^2x^2}} + 1}{a^2x^2}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x,x, algorithm="fricas")

[Out] -(a^2*x^2*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x + 1) - a^2*x^2*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x - 1) - a^2*x^2*log(x) + a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) + 1)/(a^2*x^2)

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 87 vs. 2(32) = 64.

Time = 2.00 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.29

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x} dx = \begin{cases} a^2 \log(x) - 2a \left(\begin{cases} \frac{\log\left(2\sqrt{1+\frac{1}{a^2x^2}}\sqrt{\frac{1}{a^2}+\frac{2}{a^2x}}\right) + \frac{\sqrt{1+\frac{1}{a^2x^2}}}{2x}}{2\sqrt{\frac{1}{a^2}}} & \text{for } \frac{1}{a^2} \neq 0 \\ \frac{1}{x} & \text{otherwise} \end{cases} \right)^{-\frac{1}{x^2}} & \text{for } a^2 \neq 0 \\ \text{NaN} & \text{otherwise} \end{cases}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))**2/x,x)

[Out] Piecewise(((a**2*log(x) - 2*a*Piecewise((log(2*sqrt(1 + 1/(a**2*x**2)))*sqrt(a**(-2)) + 2/(a**2*x))/(2*sqrt(a**(-2))) + sqrt(1 + 1/(a**2*x**2))/(2*x), Ne(a**(-2), 0)), (1/x, True)) - 1/x**2)/a**2, Ne(a**2, 0)), (nan, True))

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 93 vs. 2(36) = 72.

Time = 0.24 (sec) , antiderivative size = 93, normalized size of antiderivative = 2.45

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x} dx = -\frac{\frac{2a^2x\sqrt{\frac{1}{a^2x^2}+1}}{a^2x^2\left(\frac{1}{a^2x^2}+1\right)^{-1}} + a\log\left(ax\sqrt{\frac{1}{a^2x^2}+1}+1\right) - a\log\left(ax\sqrt{\frac{1}{a^2x^2}+1}-1\right)}{2a} - \frac{1}{a^2x^2} + \log(x)$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x,x, algorithm="maxima")

[Out] -1/2*(2*a^2*x*sqrt(1/(a^2*x^2) + 1)/(a^2*x^2*(1/(a^2*x^2) + 1) - 1) + a*log(a*x*sqrt(1/(a^2*x^2) + 1) + 1) - a*log(a*x*sqrt(1/(a^2*x^2) + 1) - 1))/a - 1/(a^2*x^2) + log(x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 118 vs. 2(36) = 72.

Time = 0.28 (sec) , antiderivative size = 118, normalized size of antiderivative = 3.11

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x} dx = \frac{(a^4|a|\operatorname{sgn}(x) - a^5) \log(\sqrt{a^2x^2 + 1} + 1) - (a^4|a|\operatorname{sgn}(x) + a^5) \log(\sqrt{a^2x^2 + 1} - 1) + \frac{2(\sqrt{a^2x^2 + 1}a^4|a|\operatorname{sgn}(x))}{(\sqrt{a^2x^2 + 1} + 1)(\sqrt{a^2x^2 + 1} - 1)}}{2a^5}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x,x, algorithm="giac")

[Out] -1/2*((a^4*abs(a)*sgn(x) - a^5)*log(sqrt(a^2*x^2 + 1) + 1) - (a^4*abs(a)*sgn(x) + a^5)*log(sqrt(a^2*x^2 + 1) - 1) + 2*(sqrt(a^2*x^2 + 1)*a^4*abs(a)*sgn(x) + a^5)/((sqrt(a^2*x^2 + 1) + 1)*(sqrt(a^2*x^2 + 1) - 1)))/a^5

Mupad [B] (verification not implemented)

Time = 4.87 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.16

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x} dx = -\ln\left(\frac{1}{x}\right) - \operatorname{asinh}\left(\frac{1}{ax}\right) - \frac{1}{a^2x^2} - \frac{\sqrt{\frac{1}{a^2x^2} + 1}}{ax}$$

[In] int(((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))^2/x,x)

[Out] -log(1/x) - asinh(1/(a*x)) - 1/(a^2*x^2) - (1/(a^2*x^2) + 1)^(1/2)/(a*x)

3.55 $\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^2} dx$

Optimal result	344
Rubi [A] (verified)	344
Mathematica [A] (verified)	345
Maple [A] (verified)	346
Fricas [B] (verification not implemented)	346
Sympy [A] (verification not implemented)	346
Maxima [A] (verification not implemented)	347
Giac [B] (verification not implemented)	347
Mupad [B] (verification not implemented)	347

Optimal result

Integrand size = 12, antiderivative size = 34

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^2} dx = -\frac{2}{3}a \left(1 + \frac{1}{a^2x^2}\right)^{3/2} - \frac{2}{3a^2x^3} - \frac{1}{x}$$

[Out] $-2/3*a*(1+1/a^2/x^2)^{(3/2)}-2/3/a^2/x^3-1/x$

Rubi [A] (verified)

Time = 0.15 (sec), antiderivative size = 54, normalized size of antiderivative = 1.59, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6473, 6847, 2142}

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^2} dx = -\frac{1}{6}a \left(\sqrt{\frac{1}{a^2x^2} + 1} + \frac{1}{ax}\right)^3 - \frac{1}{2}a\sqrt{\frac{1}{a^2x^2} + 1} - \frac{1}{2x}$$

[In] $\text{Int}[E^{(2*\text{ArcCsch}[a*x])}/x^2, x]$

[Out] $-1/2*(a*\text{Sqrt}[1 + 1/(a^2*x^2)]) - (a*(\text{Sqrt}[1 + 1/(a^2*x^2)] + 1/(a*x))^3)/6 - 1/(2*x)$

Rule 2142

$\text{Int}[(g_.) + (h_.)*((d_.) + (e_.)*(x_.) + (f_.)*\text{Sqrt}[(a_.) + (c_.)*(x_.)^2])^n)^p, x_Symbol] \rightarrow \text{Dist}[1/(2*e), \text{Subst}[\text{Int}[(g + h*x^n)^p*((d^2 + a*f^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*\text{Sqrt}[a + c*x^2]], x] /;$ FreeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]

Rule 6473

`Int[E^(ArcCsch[u_]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[1 + 1/u^2])^n, x] /; FreeQ[m, x] && IntegerQ[n]`

Rule 6847

`Int[(u_)*(x_)^(m_.), x_Symbol] := Dist[1/(m + 1), Subst[Int[SubstFor[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^(m + 1), u, x]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{\left(\sqrt{1 + \frac{1}{a^2 x^2}} + \frac{1}{ax}\right)^2}{x^2} dx \\
 &= -\text{Subst}\left(\int \left(\frac{x}{a} + \sqrt{1 + \frac{x^2}{a^2}}\right)^2 dx, x, \frac{1}{x}\right) \\
 &= -\left(\frac{1}{2}a \text{Subst}\left(\int (1 + x^2) dx, x, \sqrt{1 + \frac{1}{a^2 x^2}} + \frac{1}{ax}\right)\right) \\
 &= -\frac{1}{2}a \sqrt{1 + \frac{1}{a^2 x^2}} - \frac{1}{6}a \left(\sqrt{1 + \frac{1}{a^2 x^2}} + \frac{1}{ax}\right)^3 - \frac{1}{2x}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.35

$$\int \frac{e^{2\text{csch}^{-1}(ax)}}{x^2} dx = -\frac{2 + 3a^2 x^2 + 2a \sqrt{1 + \frac{1}{a^2 x^2}} x(1 + a^2 x^2)}{3a^2 x^3}$$

`[In] Integrate[E^(2*ArcCsch[a*x])/x^2,x]`

`[Out] -1/3*(2 + 3*a^2*x^2 + 2*a*Sqrt[1 + 1/(a^2*x^2)]*x*(1 + a^2*x^2))/(a^2*x^3)`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.65

method	result	size
trager	$\frac{-\frac{3a^2x^2+2}{3x^3} - \frac{2a(a^2x^2+1)\sqrt{-\frac{-a^2x^2-1}{a^2x^2}}}{a^2}}{3x^2}$	56
default	$\frac{-\frac{1}{3x^3} - \frac{a^2}{x}}{a^2} - \frac{2\sqrt{\frac{a^2x^2+1}{a^2x^2}}(a^2x^2+1)}{3ax^2} - \frac{1}{3a^2x^3}$	63

[In] `int((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^2,x,method=_RETURNVERBOSE)`

[Out] $1/a^2*(-1/3*(3*a^2*x^2+2)/x^3-2/3/x^2*a*(a^2*x^2+1)*(-(-a^2*x^2-1)/a^2/x^2)^{(1/2)})$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 57 vs. $2(28) = 56$.

Time = 0.25 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.68

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^2} dx = -\frac{2a^3x^3 + 3a^2x^2 + 2(a^3x^3 + ax)\sqrt{\frac{a^2x^2+1}{a^2x^2}} + 2}{3a^2x^3}$$

[In] `integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^2,x, algorithm="fricas")`

[Out] $-1/3*(2*a^3*x^3 + 3*a^2*x^2 + 2*(a^3*x^3 + a*x)*\operatorname{sqrt}((a^2*x^2 + 1)/(a^2*x^2)) + 2)/(a^2*x^3)$

Sympy [A] (verification not implemented)

Time = 1.19 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.71

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^2} dx = \begin{cases} -\frac{a^2}{x} - 2a \left(\begin{cases} \sqrt{1 + \frac{1}{a^2x^2}} \left(\frac{a^2}{3} + \frac{1}{3x^2} \right) & \text{for } \frac{1}{a^2} \neq 0 \\ \frac{1}{2x^2} & \text{otherwise} \end{cases} \right) - \frac{2}{3x^3} & \text{for } a^2 \neq 0 \\ \text{NaN} & \text{otherwise} \end{cases}$$

[In] `integrate((1/a/x+(1+1/a**2/x**2)**(1/2))**2/x**2,x)`

[Out] `Piecewise((((-a**2/x - 2*a*Piecewise((sqrt(1 + 1/(a**2*x**2)))*(a**2/3 + 1/(3*x**2)), Ne(a**(-2), 0)), (1/(2*x**2), True)) - 2/(3*x**3))/a**2, Ne(a**2, 0)), (nan, True))`

Maxima [A] (verification not implemented)

none

Time = 0.23 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.82

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^2} dx = -\frac{2}{3} a \left(\frac{1}{a^2 x^2} + 1 \right)^{\frac{3}{2}} - \frac{1}{x} - \frac{2}{3 a^2 x^3}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^2,x, algorithm="maxima")

[Out] -2/3*a*(1/(a^2*x^2) + 1)^(3/2) - 1/x - 2/3/(a^2*x^3)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 75 vs. 2(28) = 56.

Time = 0.30 (sec) , antiderivative size = 75, normalized size of antiderivative = 2.21

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^2} dx = \frac{4 \left(3 (x|a| - \sqrt{a^2 x^2 + 1})^4 \operatorname{asgn}(x) + a \operatorname{sgn}(x) \right)}{3 \left((x|a| - \sqrt{a^2 x^2 + 1})^2 - 1 \right)^3} - \frac{3 a^2 x^2 + 2}{3 a^2 x^3}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^2,x, algorithm="giac")

[Out] 4/3*(3*(x*abs(a) - sqrt(a^2*x^2 + 1))^4*a*sgn(x) + a*sgn(x))/((x*abs(a) - sqrt(a^2*x^2 + 1))^2 - 1)^3 - 1/3*(3*a^2*x^2 + 2)/(a^2*x^3)

Mupad [B] (verification not implemented)

Time = 4.83 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.50

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^2} dx = -\frac{2}{3a^2} + \frac{2x\sqrt{\frac{1}{a^2x^2}+1}}{3a} - \frac{2ax\sqrt{\frac{1}{a^2x^2}+1}}{3} + \frac{1}{x}$$

[In] int(((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))^2/x^2,x)

[Out] - (2/(3*a^2) + (2*x*(1/(a^2*x^2) + 1)^(1/2))/(3*a))/x^3 - ((2*a*x*(1/(a^2*x^2) + 1)^(1/2))/3 + 1)/x

3.56 $\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^3} dx$

Optimal result	348
Rubi [A] (verified)	348
Mathematica [A] (verified)	350
Maple [B] (verified)	350
Fricas [B] (verification not implemented)	351
Sympy [A] (verification not implemented)	351
Maxima [B] (verification not implemented)	352
Giac [A] (verification not implemented)	352
Mupad [B] (verification not implemented)	353

Optimal result

Integrand size = 12, antiderivative size = 73

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^3} dx = -\frac{1}{2a^2x^4} - \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{2ax^3} - \frac{1}{2x^2} - \frac{a\sqrt{1 + \frac{1}{a^2x^2}}}{4x} + \frac{1}{4}a^2\operatorname{csch}^{-1}(ax)$$

[Out] $-1/2/a^2/x^4 - 1/2/x^2 + 1/4*a^2*\operatorname{arccsch}(a*x) - 1/2*(1+1/a^2/x^2)^{(1/2)}/a/x^3 - 1/4*a*(1+1/a^2/x^2)^{(1/2)}/x$

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6473, 6874, 342, 285, 327, 221}

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^3} dx = -\frac{1}{2a^2x^4} - \frac{a\sqrt{\frac{1}{a^2x^2} + 1}}{4x} - \frac{\sqrt{\frac{1}{a^2x^2} + 1}}{2ax^3} + \frac{1}{4}a^2\operatorname{csch}^{-1}(ax) - \frac{1}{2x^2}$$

[In] $\operatorname{Int}[E^{(2*\operatorname{ArcCsch}[a*x])}/x^3, x]$

[Out] $-1/2*1/(a^2*x^4) - \operatorname{Sqrt}[1 + 1/(a^2*x^2)]/(2*a*x^3) - 1/(2*x^2) - (a*\operatorname{Sqrt}[1 + 1/(a^2*x^2)])/(4*x) + (a^2*\operatorname{ArcCsch}[a*x])/4$

Rule 221

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_) + (b_)*(x_)^2], x_Symbol] := \operatorname{Simp}[\operatorname{ArcSinh}[\operatorname{Rt}[b, 2]*(x/\operatorname{Sqrt}[a])]/\operatorname{Rt}[b, 2], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{GtQ}[a, 0] \ \&\& \operatorname{PosQ}[b]$

Rule 285

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^p/(c*(m + n*p + 1))), x] + Dist[a*n*(p/(m + n*p + 1)), Int[(c*x)^m*(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[n, 0] && GtQ[p, 0] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 327

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 342

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]
```

Rule 6473

```
Int[E^(ArcCsch[u_]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[1 + 1/u^2])^n, x] /; FreeQ[m, x] && IntegerQ[n]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{\left(\sqrt{1 + \frac{1}{a^2 x^2}} + \frac{1}{ax}\right)^2}{x^3} dx \\
 &= \int \left(\frac{2}{a^2 x^5} + \frac{2\sqrt{1 + \frac{1}{a^2 x^2}}}{ax^4} + \frac{1}{x^3}\right) dx \\
 &= -\frac{1}{2a^2 x^4} - \frac{1}{2x^2} + \frac{2 \int \frac{\sqrt{1 + \frac{1}{a^2 x^2}}}{x^4} dx}{a} \\
 &= -\frac{1}{2a^2 x^4} - \frac{1}{2x^2} - \frac{2 \text{Subst}\left(\int x^2 \sqrt{1 + \frac{x^2}{a^2}} dx, x, \frac{1}{x}\right)}{a}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{2a^2x^4} - \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{2ax^3} - \frac{1}{2x^2} - \frac{\text{Subst}\left(\int \frac{x^2}{\sqrt{1 + \frac{x^2}{a^2}}} dx, x, \frac{1}{x}\right)}{2a} \\
&= -\frac{1}{2a^2x^4} - \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{2ax^3} - \frac{1}{2x^2} - \frac{a\sqrt{1 + \frac{1}{a^2x^2}}}{4x} + \frac{1}{4}a\text{Subst}\left(\int \frac{1}{\sqrt{1 + \frac{x^2}{a^2}}} dx, x, \frac{1}{x}\right) \\
&= -\frac{1}{2a^2x^4} - \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{2ax^3} - \frac{1}{2x^2} - \frac{a\sqrt{1 + \frac{1}{a^2x^2}}}{4x} + \frac{1}{4}a^2\text{csch}^{-1}(ax)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00

$$\int \frac{e^{2\text{csch}^{-1}(ax)}}{x^3} dx = -\frac{1}{2a^2x^4} - \frac{1}{2x^2} + \left(-\frac{1}{2ax^3} - \frac{a}{4x}\right) \sqrt{\frac{1+a^2x^2}{a^2x^2}} + \frac{1}{4}a^2\text{arcsinh}\left(\frac{1}{ax}\right)$$

[In] Integrate[E^(2*ArcCsch[a*x])/x^3,x]

[Out] -1/2*1/(a^2*x^4) - 1/(2*x^2) + (-1/2*1/(a*x^3) - a/(4*x))*Sqrt[(1 + a^2*x^2)/(a^2*x^2)] + (a^2*ArcSinh[1/(a*x)])/4

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 188 vs. 2(59) = 118.

Time = 0.06 (sec) , antiderivative size = 189, normalized size of antiderivative = 2.59

method	result
default	$-\frac{1}{4x^4} - \frac{a^2}{2x^2} + \frac{a\sqrt{\frac{a^2x^2+1}{a^2x^2}} \left(\left(\frac{a^2x^2+1}{a^2}\right)^{\frac{3}{2}} \sqrt{\frac{1}{a^2}} a^2x^2 - \sqrt{\frac{a^2x^2+1}{a^2}} \sqrt{\frac{1}{a^2}} a^2x^4 + \ln\left(\frac{2\sqrt{\frac{1}{a^2}} \sqrt{\frac{a^2x^2+1}{a^2}} a^2+2}{a^2x}\right) x^4 - 2\left(\frac{a^2x^2+1}{a^2}\right)^{\frac{3}{2}} \sqrt{\frac{1}{a^2}} \right)}{4x^3 \sqrt{\frac{a^2x^2+1}{a^2}} \sqrt{\frac{1}{a^2}}}$

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^3,x,method=_RETURNVERBOSE)

[Out] 1/a^2*(-1/4/x^4-1/2*a^2/x^2)+1/4*a*((a^2*x^2+1)/a^2/x^2)^(1/2)/x^3*(((a^2*x^2+1)/a^2)^(3/2)*(1/a^2)^(1/2)*a^2*x^2-((a^2*x^2+1)/a^2)^(1/2)*(1/a^2)^(1/2))*a^2*x^4+ln(2*((1/a^2)^(1/2)*((a^2*x^2+1)/a^2)^(1/2)*a^2+1)/x/a^2)*x^4-2*((a^2*x^2+1)/a^2)^(3/2)*(1/a^2)^(1/2)/((a^2*x^2+1)/a^2)^(1/2)/(1/a^2)^(1/2)-1/4/a^2/x^4

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. 2(59) = 118.

Time = 0.25 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.66

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^3} dx$$

$$= \frac{a^4 x^4 \log\left(ax \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} - ax + 1\right) - a^4 x^4 \log\left(ax \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} - ax - 1\right) - 2a^2 x^2 - (a^3 x^3 + 2ax) \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}}}{4a^2 x^4}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^3,x, algorithm="fricas")

[Out] 1/4*(a^4*x^4*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x + 1) - a^4*x^4*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x - 1) - 2*a^2*x^2 - (a^3*x^3 + 2*a*x)*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - 2)/(a^2*x^4)

Sympy [A] (verification not implemented)

Time = 1.25 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.49

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^3} dx$$

$$= \begin{cases} \frac{-\frac{a^2}{2x^2} - 2a \left(\begin{cases} -\frac{a^2 \log\left(2\sqrt{1+\frac{1}{a^2 x^2}} \sqrt{\frac{1}{a^2} + \frac{2}{a^2 x}}\right) + \sqrt{1+\frac{1}{a^2 x^2}} \left(\frac{a^2}{8x} + \frac{1}{4x^3}\right)}{8\sqrt{\frac{1}{a^2}}} & \text{for } \frac{1}{a^2} \neq 0 \\ \frac{1}{3x^3} & \text{otherwise} \end{cases} \right)}{a^2} - \frac{1}{2x^4} & \text{for } a^2 \neq 0 \\ \text{NaN} & \text{otherwise} \end{cases}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))**2/x**3,x)

[Out] Piecewise(((-a**2/(2*x**2) - 2*a*Piecewise((-a**2*log(2*sqrt(1 + 1/(a**2*x**2))*sqrt(a**(-2)) + 2/(a**2*x))/(8*sqrt(a**(-2))) + sqrt(1 + 1/(a**2*x**2)))*(a**2/(8*x) + 1/(4*x**3)), Ne(a**(-2), 0)), (1/(3*x**3), True)) - 1/(2*x**4))/a**2, Ne(a**2, 0)), (nan, True))

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 139 vs. $2(59) = 118$.

Time = 0.26 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.90

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^3} dx$$

$$= \frac{a^3 \log\left(ax \sqrt{\frac{1}{a^2 x^2} + 1} + 1\right) - a^3 \log\left(ax \sqrt{\frac{1}{a^2 x^2} + 1} - 1\right) - \frac{2\left(a^6 x^3 \left(\frac{1}{a^2 x^2} + 1\right)^{\frac{3}{2}} + a^4 x \sqrt{\frac{1}{a^2 x^2} + 1}\right)}{a^4 x^4 \left(\frac{1}{a^2 x^2} + 1\right)^2 - 2 a^2 x^2 \left(\frac{1}{a^2 x^2} + 1\right) + 1}}{8 a} - \frac{1}{2 x^2} - \frac{1}{2 a^2 x^4}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^3,x, algorithm="maxima")

[Out] $\frac{1}{8} * (a^3 * \log(a * x * \sqrt{1/(a^2 * x^2) + 1} + 1) - a^3 * \log(a * x * \sqrt{1/(a^2 * x^2) + 1} - 1) - 2 * (a^6 * x^3 * (1/(a^2 * x^2) + 1)^{(3/2)} + a^4 * x * \sqrt{1/(a^2 * x^2) + 1})) / (a^4 * x^4 * (1/(a^2 * x^2) + 1)^2 - 2 * a^2 * x^2 * (1/(a^2 * x^2) + 1) + 1) / a - 1/2 / x^2 - 1/2 / (a^2 * x^4)$

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.53

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^3} dx$$

$$= \frac{a^6 |a| \log\left(\sqrt{a^2 x^2 + 1} + 1\right) \operatorname{sgn}(x) - a^6 |a| \log\left(\sqrt{a^2 x^2 + 1} - 1\right) \operatorname{sgn}(x) - \frac{2\left((a^2 x^2 + 1)^{\frac{3}{2}} a^6 |a| \operatorname{sgn}(x) + \sqrt{a^2 x^2 + 1} a^6 |a| \operatorname{sgn}(x)\right)}{a^4 x^4}}{8 a^5}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^3,x, algorithm="giac")

[Out] $\frac{1}{8} * (a^6 * \operatorname{abs}(a) * \log(\sqrt{a^2 * x^2 + 1} + 1) * \operatorname{sgn}(x) - a^6 * \operatorname{abs}(a) * \log(\sqrt{a^2 * x^2 + 1} - 1) * \operatorname{sgn}(x) - 2 * ((a^2 * x^2 + 1)^{(3/2)} * a^6 * \operatorname{abs}(a) * \operatorname{sgn}(x) + \sqrt{a^2 * x^2 + 1} * a^6 * \operatorname{abs}(a) * \operatorname{sgn}(x)) / (a^4 * x^4)) / a^5$

Mupad [B] (verification not implemented)

Time = 5.22 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.93

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^3} dx = \frac{a \operatorname{asinh}\left(\frac{\sqrt{\frac{1}{a^2}}}{x}\right)}{4\sqrt{\frac{1}{a^2}}} - \frac{1}{2a^2 x^4} - \frac{a\sqrt{\frac{1}{a^2 x^2} + 1}}{4x} - \frac{1}{2x^2} - \frac{\sqrt{\frac{1}{a^2 x^2} + 1}}{2ax^3}$$

[In] int(((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))^2/x^3,x)

[Out] (a*asinh((1/a^2)^(1/2)/x))/(4*(1/a^2)^(1/2)) - 1/(2*a^2*x^4) - (a*(1/(a^2*x^2) + 1)^(1/2))/(4*x) - 1/(2*x^2) - (1/(a^2*x^2) + 1)^(1/2)/(2*a*x^3)

3.57 $\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^4} dx$

Optimal result	354
Rubi [A] (verified)	354
Mathematica [A] (verified)	355
Maple [A] (verified)	356
Fricas [A] (verification not implemented)	356
Sympy [A] (verification not implemented)	356
Maxima [A] (verification not implemented)	357
Giac [B] (verification not implemented)	357
Mupad [B] (verification not implemented)	357

Optimal result

Integrand size = 12, antiderivative size = 58

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^4} dx = \frac{2}{3}a^3 \left(1 + \frac{1}{a^2x^2}\right)^{3/2} - \frac{2}{5}a^3 \left(1 + \frac{1}{a^2x^2}\right)^{5/2} - \frac{2}{5a^2x^5} - \frac{1}{3x^3}$$

[Out] $2/3*a^3*(1+1/a^2/x^2)^(3/2)-2/5*a^3*(1+1/a^2/x^2)^(5/2)-2/5/a^2/x^5-1/3/x^3$

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6473, 6874, 272, 45}

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^4} dx = -\frac{2}{5a^2x^5} - \frac{2}{5}a^3 \left(\frac{1}{a^2x^2} + 1\right)^{5/2} + \frac{2}{3}a^3 \left(\frac{1}{a^2x^2} + 1\right)^{3/2} - \frac{1}{3x^3}$$

[In] $\text{Int}[E^{(2*\text{ArcCsch}[a*x])}/x^4, x]$

[Out] $(2*a^3*(1 + 1/(a^2*x^2))^(3/2))/3 - (2*a^3*(1 + 1/(a^2*x^2))^(5/2))/5 - 2/(5*a^2*x^5) - 1/(3*x^3)$

Rule 45

$\text{Int}[(a_. + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /;$ $\text{FreeQ}\{a, b, c, d, n, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ (!\text{IntegerQ}[n] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{LeQ}[7*m + 4*n + 4, 0])) \ || \ \text{LtQ}[9*m + 5*(n + 1), 0] \ || \ \text{GtQ}[m + n + 2, 0]$

Rule 272

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 6473

```
Int[E^(ArcCsch[u_]*(n_))*(x_)^(m_), x_Symbol] := Int[x^m*(1/u + Sqrt[1 +
1/u^2])^n, x] /; FreeQ[m, x] && IntegerQ[n]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \frac{\left(\sqrt{1 + \frac{1}{a^2 x^2}} + \frac{1}{ax}\right)^2}{x^4} dx \\
&= \int \left(\frac{2}{a^2 x^6} + \frac{2\sqrt{1 + \frac{1}{a^2 x^2}}}{ax^5} + \frac{1}{x^4}\right) dx \\
&= -\frac{2}{5a^2 x^5} - \frac{1}{3x^3} + \frac{2 \int \frac{\sqrt{1 + \frac{1}{a^2 x^2}}}{x^5} dx}{a} \\
&= -\frac{2}{5a^2 x^5} - \frac{1}{3x^3} - \frac{\text{Subst}\left(\int x \sqrt{1 + \frac{x}{a^2}} dx, x, \frac{1}{x^2}\right)}{a} \\
&= -\frac{2}{5a^2 x^5} - \frac{1}{3x^3} - \frac{\text{Subst}\left(\int \left(-a^2 \sqrt{1 + \frac{x}{a^2}} + a^2 \left(1 + \frac{x}{a^2}\right)^{3/2}\right) dx, x, \frac{1}{x^2}\right)}{a} \\
&= \frac{2}{3} a^3 \left(1 + \frac{1}{a^2 x^2}\right)^{3/2} - \frac{2}{5} a^3 \left(1 + \frac{1}{a^2 x^2}\right)^{5/2} - \frac{2}{5a^2 x^5} - \frac{1}{3x^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.93

$$\int \frac{e^{2\text{csch}^{-1}(ax)}}{x^4} dx = -\frac{6 + 5a^2 x^2 + 2a\sqrt{1 + \frac{1}{a^2 x^2}}x(3 + a^2 x^2 - 2a^4 x^4)}{15a^2 x^5}$$

```
[In] Integrate[E^(2*ArcCsch[a*x])/x^4, x]
```

```
[Out] -1/15*(6 + 5*a^2*x^2 + 2*a*Sqrt[1 + 1/(a^2*x^2)]*x*(3 + a^2*x^2 - 2*a^4*x^4
))/(a^2*x^5)
```

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.12

method	result	size
trager	$\frac{-\frac{5a^2x^2+6}{15x^5} + \frac{2a(2a^4x^4-a^2x^2-3)\sqrt{-\frac{a^2x^2-1}{a^2x^2}}}{a^2}}{15x^4}$	65
default	$\frac{-\frac{a^2}{3x^3} - \frac{1}{5x^5}}{a^2} + \frac{2\sqrt{\frac{a^2x^2+1}{a^2x^2}}(a^2x^2+1)(2a^2x^2-3)}{15ax^4} - \frac{1}{5a^2x^5}$	73

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^4,x,method=_RETURNVERBOSE)

[Out] 1/a^2*(-1/15*(5*a^2*x^2+6)/x^5+2/15/x^4*a*(2*a^4*x^4-a^2*x^2-3)*(-(-a^2*x^2-1)/a^2/x^2)^(1/2))

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.16

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^4} dx = \frac{4a^5x^5 - 5a^2x^2 + 2(2a^5x^5 - a^3x^3 - 3ax)\sqrt{\frac{a^2x^2+1}{a^2x^2}} - 6}{15a^2x^5}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^4,x, algorithm="fricas")

[Out] 1/15*(4*a^5*x^5 - 5*a^2*x^2 + 2*(2*a^5*x^5 - a^3*x^3 - 3*a*x)*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - 6)/(a^2*x^5)

Sympy [A] (verification not implemented)

Time = 1.26 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.22

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^4} dx = \begin{cases} \frac{-\frac{a^2}{3x^3} - 2a \left(\begin{cases} \sqrt{1 + \frac{1}{a^2x^2}} \left(-\frac{2a^4}{15} + \frac{a^2}{15x^2} + \frac{1}{5x^4} \right) & \text{for } \frac{1}{a^2} \neq 0 \\ \frac{1}{4x^4} & \text{otherwise} \end{cases} \right) - \frac{2}{5x^5}}{a^2} & \text{for } a^2 \neq 0 \\ \text{NaN} & \text{otherwise} \end{cases}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))**2/x**4,x)

[Out] Piecewise(((-a**2/(3*x**3) - 2*a*Piecewise((sqrt(1 + 1/(a**2*x**2)))*(-2*a**4/15 + a**2/(15*x**2) + 1/(5*x**4)), Ne(a**(-2), 0)), (1/(4*x**4), True)) - 2/(5*x**5))/a**2, Ne(a**2, 0)), (nan, True))

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.90

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^4} dx = -\frac{2\left(3a^4\left(\frac{1}{a^2x^2} + 1\right)^{\frac{5}{2}} - 5a^4\left(\frac{1}{a^2x^2} + 1\right)^{\frac{3}{2}}\right)}{15a} - \frac{1}{3x^3} - \frac{2}{5a^2x^5}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^4,x, algorithm="maxima")

[Out] -2/15*(3*a^4*(1/(a^2*x^2) + 1)^(5/2) - 5*a^4*(1/(a^2*x^2) + 1)^(3/2))/a - 1/3/x^3 - 2/5/(a^2*x^5)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 134 vs. 2(46) = 92.

Time = 0.32 (sec) , antiderivative size = 134, normalized size of antiderivative = 2.31

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^4} dx = \frac{8\left(15\left(x|a| - \sqrt{a^2x^2 + 1}\right)^6 a^3 \operatorname{sgn}(x) + 5\left(x|a| - \sqrt{a^2x^2 + 1}\right)^4 a^3 \operatorname{sgn}(x) + 5\left(x|a| - \sqrt{a^2x^2 + 1}\right)^2 a^3 \operatorname{sgn}(x) - \frac{5a^2x^2 + 6}{15a^2x^5}\right)}{15\left(\left(x|a| - \sqrt{a^2x^2 + 1}\right)^2 - 1\right)^5}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^4,x, algorithm="giac")

[Out] 8/15*(15*(x*abs(a) - sqrt(a^2*x^2 + 1))^6*a^3*sgn(x) + 5*(x*abs(a) - sqrt(a^2*x^2 + 1))^4*a^3*sgn(x) + 5*(x*abs(a) - sqrt(a^2*x^2 + 1))^2*a^3*sgn(x) - a^3*sgn(x))/((x*abs(a) - sqrt(a^2*x^2 + 1))^2 - 1)^5 - 1/15*(5*a^2*x^2 + 6)/(a^2*x^5)

Mupad [B] (verification not implemented)

Time = 5.91 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.16

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^4} dx = \frac{4a^3\sqrt{\frac{1}{a^2x^2} + 1}}{15} - \frac{2ax\sqrt{\frac{1}{a^2x^2} + 1}}{15x^3} + \frac{1}{3} - \frac{2}{5a^2} + \frac{2x\sqrt{\frac{1}{a^2x^2} + 1}}{5a}$$

[In] int(((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))^2/x^4,x)

[Out] (4*a^3*(1/(a^2*x^2) + 1)^(1/2))/15 - ((2*a*x*(1/(a^2*x^2) + 1)^(1/2))/15 + 1/3)/x^3 - (2/(5*a^2) + (2*x*(1/(a^2*x^2) + 1)^(1/2))/(5*a))/x^5

3.58 $\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^5} dx$

Optimal result	358
Rubi [A] (verified)	358
Mathematica [A] (verified)	360
Maple [B] (verified)	360
Fricas [A] (verification not implemented)	361
Sympy [A] (verification not implemented)	361
Maxima [B] (verification not implemented)	362
Giac [A] (verification not implemented)	362
Mupad [B] (verification not implemented)	363

Optimal result

Integrand size = 12, antiderivative size = 96

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^5} dx = -\frac{1}{3a^2x^6} - \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{3ax^5} - \frac{1}{4x^4} - \frac{a\sqrt{1 + \frac{1}{a^2x^2}}}{12x^3} + \frac{a^3\sqrt{1 + \frac{1}{a^2x^2}}}{8x} - \frac{1}{8}a^4\operatorname{csch}^{-1}(ax)$$

[Out] $-1/3/a^2/x^6 - 1/4/x^4 - 1/8*a^4*\operatorname{arccsch}(a*x) - 1/3*(1+1/a^2/x^2)^{(1/2)}/a/x^5 - 1/12*a*(1+1/a^2/x^2)^{(1/2)}/x^3 + 1/8*a^3*(1+1/a^2/x^2)^{(1/2)}/x$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6473, 6874, 342, 285, 327, 221}

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^5} dx = -\frac{1}{8}a^4\operatorname{csch}^{-1}(ax) - \frac{1}{3a^2x^6} - \frac{\sqrt{\frac{1}{a^2x^2} + 1}}{3ax^5} - \frac{a\sqrt{\frac{1}{a^2x^2} + 1}}{12x^3} + \frac{a^3\sqrt{\frac{1}{a^2x^2} + 1}}{8x} - \frac{1}{4x^4}$$

[In] $\operatorname{Int}[E^{(2*\operatorname{ArcCsch}[a*x])}/x^5, x]$

[Out] $-1/3*1/(a^2*x^6) - \operatorname{Sqrt}[1 + 1/(a^2*x^2)]/(3*a*x^5) - 1/(4*x^4) - (a*\operatorname{Sqrt}[1 + 1/(a^2*x^2)])/(12*x^3) + (a^3*\operatorname{Sqrt}[1 + 1/(a^2*x^2)])/(8*x) - (a^4*\operatorname{ArcCsch}[a*x])/8$

Rule 221

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_) + (b_.)*(x_)^2], x_Symbol] := \operatorname{Simp}[\operatorname{ArcSinh}[\operatorname{Rt}[b, 2]*(x/\operatorname{Sqrt}[a])]/\operatorname{Rt}[b, 2], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{GtQ}[a, 0] \ \&\& \operatorname{PosQ}[b]$

Rule 285

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^p/(c*(m + n*p + 1))), x] + Dist[a*n*(p/(m + n*p + 1)), Int[(c*x)^m*(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[n, 0] && GtQ[p, 0] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 327

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 342

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]
```

Rule 6473

```
Int[E^(ArcCsch[u_]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*(1/u + Sqrt[1 + 1/u^2])^n, x] /; FreeQ[m, x] && IntegerQ[n]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{\left(\sqrt{1 + \frac{1}{a^2 x^2}} + \frac{1}{ax}\right)^2}{x^5} dx \\
 &= \int \left(\frac{2}{a^2 x^7} + \frac{2\sqrt{1 + \frac{1}{a^2 x^2}}}{ax^6} + \frac{1}{x^5}\right) dx \\
 &= -\frac{1}{3a^2 x^6} - \frac{1}{4x^4} + \frac{2 \int \frac{\sqrt{1 + \frac{1}{a^2 x^2}}}{x^6} dx}{a} \\
 &= -\frac{1}{3a^2 x^6} - \frac{1}{4x^4} - \frac{2 \text{Subst}\left(\int x^4 \sqrt{1 + \frac{x^2}{a^2}} dx, x, \frac{1}{x}\right)}{a}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{3a^2x^6} - \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{3ax^5} - \frac{1}{4x^4} - \frac{\text{Subst}\left(\int \frac{x^4}{\sqrt{1 + \frac{x^2}{a^2}}} dx, x, \frac{1}{x}\right)}{3a} \\
&= -\frac{1}{3a^2x^6} - \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{3ax^5} - \frac{1}{4x^4} - \frac{a\sqrt{1 + \frac{1}{a^2x^2}}}{12x^3} + \frac{1}{4}a\text{Subst}\left(\int \frac{x^2}{\sqrt{1 + \frac{x^2}{a^2}}} dx, x, \frac{1}{x}\right) \\
&= -\frac{1}{3a^2x^6} - \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{3ax^5} - \frac{1}{4x^4} - \frac{a\sqrt{1 + \frac{1}{a^2x^2}}}{12x^3} \\
&\quad + \frac{a^3\sqrt{1 + \frac{1}{a^2x^2}}}{8x} - \frac{1}{8}a^3\text{Subst}\left(\int \frac{1}{\sqrt{1 + \frac{x^2}{a^2}}} dx, x, \frac{1}{x}\right) \\
&= -\frac{1}{3a^2x^6} - \frac{\sqrt{1 + \frac{1}{a^2x^2}}}{3ax^5} - \frac{1}{4x^4} - \frac{a\sqrt{1 + \frac{1}{a^2x^2}}}{12x^3} + \frac{a^3\sqrt{1 + \frac{1}{a^2x^2}}}{8x} - \frac{1}{8}a^4\text{csch}^{-1}(ax)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.77

$$\int \frac{e^{2\text{csch}^{-1}(ax)}}{x^5} dx = \frac{(4+3a^2x^2)(-2-2a\sqrt{1+\frac{1}{a^2x^2}}x+a^3\sqrt{1+\frac{1}{a^2x^2}}x^3)}{x^6} - 3a^6\text{arcsinh}\left(\frac{1}{ax}\right)$$

[In] Integrate[E^(2*ArcCsch[a*x])/x^5,x]

[Out] (((4 + 3*a^2*x^2)*(-2 - 2*a*Sqrt[1 + 1/(a^2*x^2)]*x + a^3*Sqrt[1 + 1/(a^2*x^2)]*x^3))/x^6 - 3*a^6*ArcSinh[1/(a*x)])/(24*a^2)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 221 vs. 2(78) = 156.

Time = 0.06 (sec) , antiderivative size = 222, normalized size of antiderivative = 2.31

method	result
default	$ -\frac{a^2}{4x^4} - \frac{1}{6x^6} - \frac{a\sqrt{\frac{a^2x^2+1}{a^2x^2}} \left(3\sqrt{\frac{1}{a^2}} \left(\frac{a^2x^2+1}{a^2}\right)^{\frac{3}{2}} a^4x^4 - 3\sqrt{\frac{1}{a^2}} \sqrt{\frac{a^2x^2+1}{a^2}} a^4x^6 + 3\ln\left(\frac{2\sqrt{\frac{1}{a^2}} \sqrt{\frac{a^2x^2+1}{a^2}} a^2+2}{a^2x}\right) a^2x^6 - 6\left(\frac{a^2x^2+1}{a^2}\right)^{\frac{3}{2}} \sqrt{\frac{1}{a^2}} \right)}{24x^5 \sqrt{\frac{a^2x^2+1}{a^2}} \sqrt{\frac{1}{a^2}}} $

[In] int((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^5,x,method=_RETURNVERBOSE)

[Out] 1/a^2*(-1/4*a^2/x^4-1/6/x^6)-1/24*a*((a^2*x^2+1)/a^2/x^2)^(1/2)/x^5*(3*(1/a^2)^(1/2)*((a^2*x^2+1)/a^2)^(3/2)*a^4*x^4-3*(1/a^2)^(1/2)*((a^2*x^2+1)/a^2)

$$\frac{a^{1/2} a^4 x^6 + 3 \ln(2) \left(\frac{1}{a^2} \right)^{1/2} \left(\frac{a^2 x^2 + 1}{a^2} \right)^{1/2} a^2 + 1}{x a^2} a^2 x^6 - 6 \left(\frac{a^2 x^2 + 1}{a^2} \right)^{3/2} \left(\frac{1}{a^2} \right)^{1/2} a^2 x^2 + 8 \left(\frac{a^2 x^2 + 1}{a^2} \right)^{3/2} \left(\frac{1}{a^2} \right)^{1/2} \right) / \left(\frac{a^2 x^2 + 1}{a^2} \right)^{1/2} / \left(\frac{1}{a^2} \right)^{1/2} - 1/6 a^2 / x^6$$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.36

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^5} dx = \frac{3 a^6 x^6 \log \left(ax \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} - ax + 1 \right) - 3 a^6 x^6 \log \left(ax \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} - ax - 1 \right) + 6 a^2 x^2 - (3 a^5 x^5 - 2 a^3 x^3 - 5 - 2 a^3 x^3 - 8 a^2 x) \sqrt{\frac{a^2 x^2 + 1}{a^2 x^2}} + 8}{24 a^2 x^6}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^5,x, algorithm="fricas")

[Out] -1/24*(3*a^6*x^6*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x + 1) - 3*a^6*x^6*log(a*x*sqrt((a^2*x^2 + 1)/(a^2*x^2)) - a*x - 1) + 6*a^2*x^2 - (3*a^5*x^5 - 2*a^3*x^3 - 8*a*x)*sqrt((a^2*x^2 + 1)/(a^2*x^2)) + 8)/(a^2*x^6)

Sympy [A] (verification not implemented)

Time = 1.31 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.22

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^5} dx = \begin{cases} -\frac{a^2}{4x^4} - 2a \left(\begin{cases} \frac{a^4 \log \left(2\sqrt{1 + \frac{1}{a^2 x^2}} \sqrt{\frac{1}{a^2} + \frac{2}{a^2 x}} \right) + \sqrt{1 + \frac{1}{a^2 x^2}} \left(-\frac{a^4}{16x} + \frac{a^2}{24x^3} + \frac{1}{6x^5} \right)}{16\sqrt{\frac{1}{a^2}}} & \text{for } \frac{1}{a^2} \neq 0 \\ \frac{1}{5x^5} & \text{otherwise} \end{cases} \right) - \frac{1}{3x^6} & \text{for } a^2 \neq 0 \\ \text{NaN} & \text{otherwise} \end{cases}$$

[In] integrate((1/a/x+(1+1/a**2/x**2)**(1/2))**2/x**5,x)

[Out] Piecewise(((-a**2/(4*x**4) - 2*a*Piecewise((a**4*log(2*sqrt(1 + 1/(a**2*x**2)))*sqrt(a**(-2)) + 2/(a**2*x))/(16*sqrt(a**(-2))) + sqrt(1 + 1/(a**2*x**2)))*(-a**4/(16*x) + a**2/(24*x**3) + 1/(6*x**5)), Ne(a**(-2), 0)), (1/(5*x**5), True)) - 1/(3*x**6))/a**2, Ne(a**2, 0)), (nan, True))

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 180 vs. $2(78) = 156$.

Time = 0.23 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.88

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^5} dx =$$

$$\frac{3a^5 \log\left(ax\sqrt{\frac{1}{a^2x^2} + 1} + 1\right) - 3a^5 \log\left(ax\sqrt{\frac{1}{a^2x^2} + 1} - 1\right) - \frac{2\left(3a^{10}x^5\left(\frac{1}{a^2x^2} + 1\right)^{\frac{5}{2}} - 8a^8x^3\left(\frac{1}{a^2x^2} + 1\right)^{\frac{3}{2}} - 3a^6x\sqrt{\frac{1}{a^2x^2} + 1}\right)}{a^6x^6\left(\frac{1}{a^2x^2} + 1\right)^3 - 3a^4x^4\left(\frac{1}{a^2x^2} + 1\right)^2 + 3a^2x^2\left(\frac{1}{a^2x^2} + 1\right)} - \frac{1}{4x^4} - \frac{1}{3a^2x^6}}{48a}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^5,x, algorithm="maxima")

[Out] $-1/48*(3*a^5*\log(a*x*\sqrt{1/(a^2*x^2) + 1} + 1) - 3*a^5*\log(a*x*\sqrt{1/(a^2*x^2) + 1} - 1) - 2*(3*a^{10}*x^5*(1/(a^2*x^2) + 1)^{(5/2)} - 8*a^8*x^3*(1/(a^2*x^2) + 1)^{(3/2)} - 3*a^6*x*\sqrt{1/(a^2*x^2) + 1}))/a^6*x^6*(1/(a^2*x^2) + 1)^3 - 3*a^4*x^4*(1/(a^2*x^2) + 1)^2 + 3*a^2*x^2*(1/(a^2*x^2) + 1) - 1)/a - 1/4/x^4 - 1/3/(a^2*x^6)$

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.46

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^5} dx =$$

$$\frac{3a^8|a|\log\left(\sqrt{a^2x^2 + 1} + 1\right)\operatorname{sgn}(x) - 3a^8|a|\log\left(\sqrt{a^2x^2 + 1} - 1\right)\operatorname{sgn}(x) - \frac{2\left(3(a^2x^2 + 1)^{\frac{5}{2}}a^8|a|\operatorname{sgn}(x) - 8(a^2x^2 + 1)^{\frac{3}{2}}a^8|a|\operatorname{sgn}(x) - 3a^6x\sqrt{a^2x^2 + 1}\operatorname{sgn}(x)\right)}{a^6x^6(a^2x^2 + 1)^3 - 3a^4x^4(a^2x^2 + 1)^2 + 3a^2x^2(a^2x^2 + 1)}}{48a^5}$$

[In] integrate((1/a/x+(1+1/a^2/x^2)^(1/2))^2/x^5,x, algorithm="giac")

[Out] $-1/48*(3*a^8*abs(a)*\log(\sqrt{a^2*x^2 + 1} + 1)*\operatorname{sgn}(x) - 3*a^8*abs(a)*\log(\sqrt{a^2*x^2 + 1} - 1)*\operatorname{sgn}(x) - 2*(3*(a^2*x^2 + 1)^{(5/2)}*a^8*abs(a)*\operatorname{sgn}(x) - 8*(a^2*x^2 + 1)^{(3/2)}*a^8*abs(a)*\operatorname{sgn}(x) - 3*\sqrt{a^2*x^2 + 1}*a^8*abs(a)*\operatorname{sgn}(x) - 6*(a^2*x^2 + 1)*a^9 - 2*a^9)/(a^6*x^6))/a^5$

Mupad [B] (verification not implemented)

Time = 5.50 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.93

$$\int \frac{e^{2\operatorname{csch}^{-1}(ax)}}{x^5} dx = \frac{a^3 \sqrt{\frac{1}{a^2 x^2} + 1}}{8x} - \frac{1}{3a^2 x^6} - \frac{a \sqrt{\frac{1}{a^2 x^2} + 1}}{12x^3} - \frac{1}{4x^4} - \frac{\sqrt{\frac{1}{a^2 x^2} + 1}}{3ax^5} - \frac{a^3 \operatorname{asinh}\left(\frac{\sqrt{\frac{1}{a^2}}}{x}\right)}{8\sqrt{\frac{1}{a^2}}}$$

[In] int(((1/(a^2*x^2) + 1)^(1/2) + 1/(a*x))^2/x^5,x)

[Out] (a^3*(1/(a^2*x^2) + 1)^(1/2))/(8*x) - 1/(3*a^2*x^6) - (a*(1/(a^2*x^2) + 1)^(1/2))/(12*x^3) - 1/(4*x^4) - (1/(a^2*x^2) + 1)^(1/2)/(3*a*x^5) - (a^3*asinh((1/a^2)^(1/2)/x))/(8*(1/a^2)^(1/2))

$$3.59 \quad \int \frac{e^{\operatorname{csch}^{-1}(cx)}(dx)^m}{1+c^2x^2} dx$$

Optimal result	364
Rubi [A] (verified)	364
Mathematica [A] (verified)	365
Maple [F]	366
Fricas [F]	366
Sympy [F]	366
Maxima [F]	367
Giac [F(-2)]	367
Mupad [F(-1)]	367

Optimal result

Integrand size = 23, antiderivative size = 85

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}(dx)^m}{1+c^2x^2} dx = -\frac{d(dx)^{-1+m} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-m}{2}, \frac{3-m}{2}, -\frac{1}{c^2x^2}\right)}{c^2(1-m)} + \frac{(dx)^m \operatorname{Hypergeometric2F1}\left(1, \frac{m}{2}, \frac{2+m}{2}, -c^2x^2\right)}{cm}$$

[Out] -d*(d*x)^(-1+m)*hypergeom([1/2, 1/2-1/2*m], [3/2-1/2*m], -1/c^2/x^2)/c^2/(1-m)+(d*x)^m*hypergeom([1, 1/2*m], [1+1/2*m], -c^2*x^2)/c/m

Rubi [A] (verified)

Time = 0.08 (sec), antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {6477, 346, 371}

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}(dx)^m}{1+c^2x^2} dx = \frac{(dx)^m \operatorname{Hypergeometric2F1}\left(1, \frac{m}{2}, \frac{m+2}{2}, -c^2x^2\right)}{cm} - \frac{d(dx)^{m-1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-m}{2}, \frac{3-m}{2}, -\frac{1}{c^2x^2}\right)}{c^2(1-m)}$$

[In] Int[(E^ArcCsch[c*x]*(d*x)^m)/(1+c^2*x^2),x]

[Out] -(((d*(d*x)^(-1+m)*Hypergeometric2F1[1/2, (1-m)/2, (3-m)/2, -(1/(c^2*x^2))]))/(c^2*(1-m))) + ((d*x)^m*Hypergeometric2F1[1, m/2, (2+m)/2, -(c^2*x^2)])/(c*m)

Rule 346

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[(-c^
(-1))*(c*x)^(m + 1)*(1/x)^(m + 1), Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x
, 1/x], x] /; FreeQ[{a, b, c, m, p}, x] && ILtQ[n, 0] && !RationalQ[m]
```

Rule 371

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p
*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1
, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILt
Q[p, 0] || GtQ[a, 0])
```

Rule 6477

```
Int[(E^ArcCsch[(c_.)*(x_)]*((d_.)*(x_)^(m_.)))/((a_) + (b_.)*(x_)^2), x_Sym
bol] := Dist[d^2/(a*c^2), Int[(d*x)^(m - 2)/Sqrt[1 + 1/(c^2*x^2)], x], x] +
Dist[d/c, Int[(d*x)^(m - 1)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, m},
x] && EqQ[b - a*c^2, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{d \int \frac{(dx)^{-1+m}}{1+c^2x^2} dx}{c} + \frac{d^2 \int \frac{(dx)^{-2+m}}{\sqrt{1+\frac{1}{c^2x^2}}} dx}{c^2} \\
&= \frac{(dx)^m \text{Hypergeometric2F1}\left(1, \frac{m}{2}, \frac{2+m}{2}, -c^2x^2\right)}{cm} \\
&\quad - \frac{\left(d\left(\frac{1}{x}\right)^{-1+m} (dx)^{-1+m}\right) \text{Subst}\left(\int \frac{x^{-m}}{\sqrt{1+\frac{x^2}{c^2}}} dx, x, \frac{1}{x}\right)}{c^2} \\
&= -\frac{d(dx)^{-1+m} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-m}{2}, \frac{3-m}{2}, -\frac{1}{c^2x^2}\right)}{c^2(1-m)} \\
&\quad + \frac{(dx)^m \text{Hypergeometric2F1}\left(1, \frac{m}{2}, \frac{2+m}{2}, -c^2x^2\right)}{cm}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.45 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.04

$$\begin{aligned}
&\int \frac{e^{\text{csch}^{-1}(cx)} (dx)^m}{1+c^2x^2} dx \\
&= \frac{(dx)^m \left(\frac{\sqrt{1+\frac{1}{c^2x^2}} x \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m}{2}, 1+\frac{m}{2}, -c^2x^2\right)}{\sqrt{1+c^2x^2}} + \frac{\text{Hypergeometric2F1}\left(1, \frac{m}{2}, 1+\frac{m}{2}, -c^2x^2\right)}{c} \right)}{m}
\end{aligned}$$

```
[In] Integrate[(E^ArcCsch[c*x]*(d*x)^m)/(1 + c^2*x^2), x]
```

[Out] $((dx)^m (\sqrt{1 + 1/(c^2 x^2)} x \operatorname{Hypergeometric2F1}[1/2, m/2, 1 + m/2, -(c^2 x^2)])) / \sqrt{1 + c^2 x^2} + \operatorname{Hypergeometric2F1}[1, m/2, 1 + m/2, -(c^2 x^2)] / c) / m$

Maple [F]

$$\int \frac{\left(\frac{1}{cx} + \sqrt{1 + \frac{1}{c^2 x^2}}\right) (dx)^m}{c^2 x^2 + 1} dx$$

[In] `int((1/c/x+(1+1/c^2/x^2)^(1/2))*(d*x)^m/(c^2*x^2+1),x)`

[Out] `int((1/c/x+(1+1/c^2/x^2)^(1/2))*(d*x)^m/(c^2*x^2+1),x)`

Fricas [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} (dx)^m}{1 + c^2 x^2} dx = \int \frac{(dx)^m \left(\sqrt{\frac{1}{c^2 x^2} + 1} + \frac{1}{cx}\right)}{c^2 x^2 + 1} dx$$

[In] `integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*(d*x)^m/(c^2*x^2+1),x, algorithm="fricas")`

[Out] `integral(((d*x)^m*c*x*sqrt((c^2*x^2 + 1)/(c^2*x^2)) + (d*x)^m)/(c^3*x^3 + c*x), x)`

Sympy [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} (dx)^m}{1 + c^2 x^2} dx = \frac{\int \frac{(dx)^m}{c^2 x^3 + x} dx + \int \frac{cx(dx)^m \sqrt{1 + \frac{1}{c^2 x^2}}}{c^2 x^3 + x} dx}{c}$$

[In] `integrate((1/c/x+(1+1/c**2/x**2)**(1/2))*(d*x)**m/(c**2*x**2+1),x)`

[Out] `(Integral((d*x)**m/(c**2*x**3 + x), x) + Integral(c*x*(d*x)**m*sqrt(1 + 1/(c**2*x**2))/(c**2*x**3 + x), x))/c`

Maxima [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}(dx)^m}{1+c^2x^2} dx = \int \frac{(dx)^m \left(\sqrt{\frac{1}{c^2x^2} + 1} + \frac{1}{cx} \right)}{c^2x^2 + 1} dx$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*(d*x)^m/(c^2*x^2+1),x, algorithm="maxima")

[Out] integrate((d*x)^m*(sqrt(1/(c^2*x^2) + 1) + 1/(c*x))/(c^2*x^2 + 1), x)

Giac [F(-2)]

Exception generated.

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}(dx)^m}{1+c^2x^2} dx = \text{Exception raised: TypeError}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*(d*x)^m/(c^2*x^2+1),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}(dx)^m}{1+c^2x^2} dx = \int \frac{\left(\sqrt{\frac{1}{c^2x^2} + 1} + \frac{1}{cx} \right) (dx)^m}{c^2x^2 + 1} dx$$

[In] int((((1/(c^2*x^2) + 1)^(1/2) + 1/(c*x))*(d*x)^m)/(c^2*x^2 + 1),x)

[Out] int((((1/(c^2*x^2) + 1)^(1/2) + 1/(c*x))*(d*x)^m)/(c^2*x^2 + 1), x)

3.60 $\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^5}{1+c^2x^2} dx$

Optimal result	368
Rubi [A] (verified)	368
Mathematica [A] (verified)	370
Maple [B] (verified)	371
Fricas [A] (verification not implemented)	371
Sympy [F]	371
Maxima [B] (verification not implemented)	372
Giac [A] (verification not implemented)	372
Mupad [B] (verification not implemented)	373

Optimal result

Integrand size = 21, antiderivative size = 92

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^5}{1+c^2x^2} dx = -\frac{x}{c^5} - \frac{3\sqrt{1+\frac{1}{c^2x^2}}x^2}{8c^4} + \frac{x^3}{3c^3} + \frac{\sqrt{1+\frac{1}{c^2x^2}}x^4}{4c^2} + \frac{\arctan(cx)}{c^6} + \frac{3\operatorname{arctanh}\left(\sqrt{1+\frac{1}{c^2x^2}}\right)}{8c^6}$$

[Out] $-x/c^5+1/3*x^3/c^3+\arctan(c*x)/c^6+3/8*\operatorname{arctanh}((1+1/c^2/x^2)^{(1/2)})/c^6-3/8*x^2*(1+1/c^2/x^2)^{(1/2)}/c^4+1/4*x^4*(1+1/c^2/x^2)^{(1/2)}/c^2$

Rubi [A] (verified)

Time = 0.08 (sec), antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6477, 272, 44, 65, 214, 308, 209}

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^5}{1+c^2x^2} dx = \frac{\arctan(cx)}{c^6} + \frac{3\operatorname{arctanh}\left(\sqrt{\frac{1}{c^2x^2}+1}\right)}{8c^6} - \frac{x}{c^5} + \frac{x^3}{3c^3} + \frac{x^4\sqrt{\frac{1}{c^2x^2}+1}}{4c^2} - \frac{3x^2\sqrt{\frac{1}{c^2x^2}+1}}{8c^4}$$

[In] $\operatorname{Int}\left[\left(E^{\operatorname{ArcCsch}[c*x]}*x^5\right)/\left(1+c^2*x^2\right),x\right]$

[Out] $-(x/c^5) - (3*\operatorname{Sqrt}[1+1/(c^2*x^2)]*x^2)/(8*c^4) + x^3/(3*c^3) + (\operatorname{Sqrt}[1+1/(c^2*x^2)]*x^4)/(4*c^2) + \operatorname{ArcTan}[c*x]/c^6 + (3*\operatorname{ArcTanh}[\operatorname{Sqrt}[1+1/(c^2*x^2)]])/(8*c^6)$

Rule 44

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((
m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !Int
egerQ[n] && LtQ[n, 0]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 209

```
Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 214

```
Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 308

```
Int[(x_)^(m_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Int[PolynomialDivide[x
^m, a + b*x^n, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 0] && IGtQ[n, 0] && Gt
Q[m, 2*n - 1]
```

Rule 6477

```
Int[(E^ArcCsch[(c_.)*(x_)]*((d_.)*(x_))^(m_.))/((a_) + (b_.)*(x_)^2), x_Sym
bol] := Dist[d^2/(a*c^2), Int[(d*x)^(m - 2)/Sqrt[1 + 1/(c^2*x^2)], x], x] +
Dist[d/c, Int[(d*x)^(m - 1)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, m},
x] && EqQ[b - a*c^2, 0]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int \frac{x^3}{\sqrt{1+\frac{1}{c^2x^2}}} dx}{c^2} + \frac{\int \frac{x^4}{1+c^2x^2} dx}{c} \\
 &= -\frac{\text{Subst}\left(\int \frac{1}{x^3\sqrt{1+\frac{x}{c^2}}} dx, x, \frac{1}{x^2}\right)}{2c^2} + \frac{\int \left(-\frac{1}{c^4} + \frac{x^2}{c^2} + \frac{1}{c^4(1+c^2x^2)}\right) dx}{c} \\
 &= -\frac{x}{c^5} + \frac{x^3}{3c^3} + \frac{\sqrt{1+\frac{1}{c^2x^2}}x^4}{4c^2} + \frac{\int \frac{1}{1+c^2x^2} dx}{c^5} + \frac{3\text{Subst}\left(\int \frac{1}{x^2\sqrt{1+\frac{x}{c^2}}} dx, x, \frac{1}{x^2}\right)}{8c^4} \\
 &= -\frac{x}{c^5} - \frac{3\sqrt{1+\frac{1}{c^2x^2}}x^2}{8c^4} + \frac{x^3}{3c^3} + \frac{\sqrt{1+\frac{1}{c^2x^2}}x^4}{4c^2} + \frac{\arctan(cx)}{c^6} - \frac{3\text{Subst}\left(\int \frac{1}{x\sqrt{1+\frac{x}{c^2}}} dx, x, \frac{1}{x^2}\right)}{16c^6} \\
 &= -\frac{x}{c^5} - \frac{3\sqrt{1+\frac{1}{c^2x^2}}x^2}{8c^4} + \frac{x^3}{3c^3} + \frac{\sqrt{1+\frac{1}{c^2x^2}}x^4}{4c^2} \\
 &\quad + \frac{\arctan(cx)}{c^6} - \frac{3\text{Subst}\left(\int \frac{1}{-c^2+c^2x^2} dx, x, \sqrt{1+\frac{1}{c^2x^2}}\right)}{8c^4} \\
 &= -\frac{x}{c^5} - \frac{3\sqrt{1+\frac{1}{c^2x^2}}x^2}{8c^4} + \frac{x^3}{3c^3} + \frac{\sqrt{1+\frac{1}{c^2x^2}}x^4}{4c^2} + \frac{\arctan(cx)}{c^6} + \frac{3\text{arctanh}\left(\sqrt{1+\frac{1}{c^2x^2}}\right)}{8c^6}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.92

$$\begin{aligned}
 &\int \frac{e^{\text{csch}^{-1}(cx)} x^5}{1+c^2x^2} dx \\
 &= \frac{cx\left(-24-9c\sqrt{1+\frac{1}{c^2x^2}}x+8c^2x^2+6c^3\sqrt{1+\frac{1}{c^2x^2}}x^3\right)+24\arctan(cx)+9\log\left(\left(1+\sqrt{1+\frac{1}{c^2x^2}}\right)x\right)}{24c^6}
 \end{aligned}$$

[In] Integrate[(E^ArcCsch[c*x]*x^5)/(1+c^2*x^2),x]

[Out] (c*x*(-24-9*c*Sqrt[1+1/(c^2*x^2)]*x+8*c^2*x^2+6*c^3*Sqrt[1+1/(c^2*x^2)]*x^3)+24*ArcTan[c*x]+9*Log[(1+Sqrt[1+1/(c^2*x^2)])*x])/(24*c^6)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 171 vs. 2(78) = 156.

Time = 0.96 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.87

method	result
default	$\frac{\sqrt{\frac{c^2x^2+1}{c^2x^2}} x \left(2x \left(\frac{c^2x^2+1}{c^2} \right)^{\frac{3}{2}} c^4 - 5x \sqrt{\frac{c^2x^2+1}{c^2}} c^2 - 5 \ln \left(x + \sqrt{\frac{c^2x^2+1}{c^2}} \right) + 8 \ln \left(x + \sqrt{-\frac{(-c^2x + \sqrt{-c^2})(c^2x + \sqrt{-c^2})}{c^4}} \right) \right)}{8 \sqrt{\frac{c^2x^2+1}{c^2}} c^6} + \frac{\frac{1}{3} c^2 x^3 - x}{c^4} + \frac{1}{c}$

[In] int((1/c/x+(1+1/c^2/x^2)^(1/2))*x^5/(c^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] 1/8*((c^2*x^2+1)/c^2/x^2)^(1/2)*x*(2*x*(1/c^2*(c^2*x^2+1))^(3/2)*c^4-5*x*(1/c^2*(c^2*x^2+1))^(1/2)*c^2-5*ln(x+(1/c^2*(c^2*x^2+1))^(1/2))+8*ln(x+(-(-c^2*x+(-c^2)^(1/2))*(c^2*x+(-c^2)^(1/2)))/c^4)^(1/2)))/(1/c^2*(c^2*x^2+1))^(1/2)/c^6+1/c*(1/c^4*(1/3*c^2*x^3-x)+1/c^5*arctan(c*x))

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.98

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^5}{1 + c^2 x^2} dx = \frac{8c^3x^3 - 24cx + 3(2c^4x^4 - 3c^2x^2)\sqrt{\frac{c^2x^2+1}{c^2x^2}} + 24 \arctan(cx) - 9 \log\left(cx\sqrt{\frac{c^2x^2+1}{c^2x^2}} - cx\right)}{24c^6}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*x^5/(c^2*x^2+1),x, algorithm="fricas")

[Out] 1/24*(8*c^3*x^3 - 24*c*x + 3*(2*c^4*x^4 - 3*c^2*x^2)*sqrt((c^2*x^2 + 1)/(c^2*x^2)) + 24*arctan(c*x) - 9*log(c*x*sqrt((c^2*x^2 + 1)/(c^2*x^2)) - c*x))/c^6

Sympy [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^5}{1 + c^2 x^2} dx = \int \frac{x^4}{c^2 x^2 + 1} dx + \int \frac{cx^5 \sqrt{1 + \frac{1}{c^2 x^2}}}{c^2 x^2 + 1} dx$$

[In] integrate((1/c/x+(1+1/c**2/x**2)**(1/2))*x**5/(c**2*x**2+1),x)

[Out] (Integral(x**4/(c**2*x**2 + 1), x) + Integral(c*x**5*sqrt(1 + 1/(c**2*x**2)))/(c**2*x**2 + 1), x))/c

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 162 vs. $2(78) = 156$.

Time = 0.31 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.76

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^5}{1 + c^2 x^2} dx$$

$$= \frac{c^2 x^3 - 3x}{3c^5} - \frac{2 \left(\frac{5 \sqrt{\frac{c^2 x^2 + 1}{x^2}}}{c} - \frac{3 \left(\frac{c^2 x^2 + 1}{x^2} \right)^{\frac{3}{2}}}{c^3} \right)}{\frac{2(c^2 x^2 + 1)}{c^2 x^2} - \frac{(c^2 x^2 + 1)^2}{c^4 x^4} - 1} - 3 \log \left(\sqrt{\frac{c^2 x^2 + 1}{x^2}} + 1 \right) + 3 \log \left(\sqrt{\frac{c^2 x^2 + 1}{x^2}} - 1 \right)$$

$$+ \frac{\arctan(cx)}{c^6}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*x^5/(c^2*x^2+1),x, algorithm="maxima")

[Out] 1/3*(c^2*x^3 - 3*x)/c^5 - 1/16*(2*(5*sqrt((c^2*x^2 + 1)/x^2)/c - 3*((c^2*x^2 + 1)/x^2)^(3/2)/c^3)/(2*(c^2*x^2 + 1)/(c^2*x^2) - (c^2*x^2 + 1)^2/(c^4*x^4) - 1) - 3*log(sqrt((c^2*x^2 + 1)/x^2)/c + 1) + 3*log(sqrt((c^2*x^2 + 1)/x^2)/c - 1))/c^6 + arctan(c*x)/c^6

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.97

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^5}{1 + c^2 x^2} dx = \frac{1}{8} \sqrt{c^2 x^2 + 1} x \left(\frac{2x^2 |c| \operatorname{sgn}(x)}{c^4} - \frac{3 |c| \operatorname{sgn}(x)}{c^6} \right)$$

$$- \frac{3 \log(-x|c| + \sqrt{c^2 x^2 + 1}) \operatorname{sgn}(x)}{8c^6} + \frac{\arctan(cx)}{c^6} + \frac{c^6 x^3 - 3c^4 x}{3c^9}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*x^5/(c^2*x^2+1),x, algorithm="giac")

[Out] 1/8*sqrt(c^2*x^2 + 1)*x*(2*x^2*abs(c)*sgn(x)/c^4 - 3*abs(c)*sgn(x)/c^6) - 3/8*log(-x*abs(c) + sqrt(c^2*x^2 + 1))*sgn(x)/c^6 + arctan(c*x)/c^6 + 1/3*(c^6*x^3 - 3*c^4*x)/c^9

Mupad [B] (verification not implemented)

Time = 5.34 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.86

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^5}{1 + c^2 x^2} dx = \frac{3 \operatorname{atanh}\left(\sqrt{\frac{1}{c^2 x^2} + 1}\right)}{8 c^6} + \frac{3 \operatorname{atan}(cx) - 3 c x + c^3 x^3}{3 c^6} \\ + \frac{x^4 \sqrt{\frac{1}{c^2 x^2} + 1}}{4 c^2} - \frac{3 x^2 \sqrt{\frac{1}{c^2 x^2} + 1}}{8 c^4}$$

[In] int((x^5*((1/(c^2*x^2) + 1)^(1/2) + 1/(c*x)))/(c^2*x^2 + 1),x)

[Out] (3*atanh((1/(c^2*x^2) + 1)^(1/2)))/(8*c^6) + (3*atan(c*x) - 3*c*x + c^3*x^3)/(3*c^6) + (x^4*(1/(c^2*x^2) + 1)^(1/2))/(4*c^2) - (3*x^2*(1/(c^2*x^2) + 1)^(1/2))/(8*c^4)

3.61 $\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^4}{1+c^2x^2} dx$

Optimal result	374
Rubi [A] (verified)	374
Mathematica [A] (verified)	376
Maple [B] (verified)	376
Fricas [A] (verification not implemented)	376
Sympy [F]	377
Maxima [A] (verification not implemented)	377
Giac [A] (verification not implemented)	377
Mupad [B] (verification not implemented)	378

Optimal result

Integrand size = 21, antiderivative size = 72

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^4}{1+c^2x^2} dx = -\frac{2\sqrt{1+\frac{1}{c^2x^2}}x}{3c^4} + \frac{x^2}{2c^3} + \frac{\sqrt{1+\frac{1}{c^2x^2}}x^3}{3c^2} - \frac{\log(1+c^2x^2)}{2c^5}$$

[Out] $1/2*x^2/c^3-1/2*\ln(c^2*x^2+1)/c^5-2/3*x*(1+1/c^2/x^2)^{(1/2)}/c^4+1/3*x^3*(1+1/c^2/x^2)^{(1/2)}/c^2$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {6477, 277, 197, 272, 45}

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^4}{1+c^2x^2} dx = \frac{x^2}{2c^3} + \frac{x^3\sqrt{\frac{1}{c^2x^2}+1}}{3c^2} - \frac{\log(c^2x^2+1)}{2c^5} - \frac{2x\sqrt{\frac{1}{c^2x^2}+1}}{3c^4}$$

[In] `Int[(E^ArcCsch[c*x])*x^4]/(1+c^2*x^2),x]`

[Out] `(-2*Sqrt[1+1/(c^2*x^2)]*x)/(3*c^4)+x^2/(2*c^3)+(Sqrt[1+1/(c^2*x^2)]*x^3)/(3*c^2)-Log[1+c^2*x^2]/(2*c^5)`

Rule 45

`Int[((a_.)+(b_.)*(x_))^(m_.)*((c_.)+(d_.)*(x_))^(n_.),x_Symbol]>Int[ExpandIntegrand[(a+b*x)^m*(c+d*x)^n,x],x];FreeQ[{a,b,c,d,n},x]&&NeQ[b*c-a*d,0]&&IGtQ[m,0]&&(!IntegerQ[n]|| (EqQ[c,0]&&LeQ[7*m+4*n+4,0])||LtQ[9*m+5*(n+1),0]||GtQ[m+n+2,0])`

Rule 197

Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 272

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 277

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[x^(m + 1)*((a + b*x^n)^(p + 1)/(a*(m + 1))), x] - Dist[b*((m + n*(p + 1) + 1)/(a*(m + 1))), Int[x^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n + p + 1], 0] && NeQ[m, -1]

Rule 6477

Int[(E^ArcCsch[(c_)*(x_)])*((d_)*(x_)^(m_)))/((a_) + (b_)*(x_)^2), x_Symbol] := Dist[d^2/(a*c^2), Int[(d*x)^(m - 2)/Sqrt[1 + 1/(c^2*x^2)], x], x] + Dist[d/c, Int[(d*x)^(m - 1)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b - a*c^2, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int \frac{x^2}{\sqrt{1+\frac{1}{c^2x^2}}} dx}{c^2} + \frac{\int \frac{x^3}{1+c^2x^2} dx}{c} \\
 &= \frac{\sqrt{1+\frac{1}{c^2x^2}}x^3}{3c^2} - \frac{2\int \frac{1}{\sqrt{1+\frac{1}{c^2x^2}}} dx}{3c^4} + \frac{\text{Subst}\left(\int \frac{x}{1+c^2x} dx, x, x^2\right)}{2c} \\
 &= -\frac{2\sqrt{1+\frac{1}{c^2x^2}}x}{3c^4} + \frac{\sqrt{1+\frac{1}{c^2x^2}}x^3}{3c^2} + \frac{\text{Subst}\left(\int \left(\frac{1}{c^2} - \frac{1}{c^2(1+c^2x)}\right) dx, x, x^2\right)}{2c} \\
 &= -\frac{2\sqrt{1+\frac{1}{c^2x^2}}x}{3c^4} + \frac{x^2}{2c^3} + \frac{\sqrt{1+\frac{1}{c^2x^2}}x^3}{3c^2} - \frac{\log(1+c^2x^2)}{2c^5}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.89

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^4}{1 + c^2 x^2} dx = \frac{cx \left(-4\sqrt{1 + \frac{1}{c^2 x^2}} + 3cx + 2c^2 \sqrt{1 + \frac{1}{c^2 x^2}} x^2 \right) - 3 \log(1 + c^2 x^2)}{6c^5}$$

[In] Integrate[(E^ArcCsch[c*x]*x^4)/(1 + c^2*x^2),x]

[Out] (c*x*(-4*Sqrt[1 + 1/(c^2*x^2)] + 3*c*x + 2*c^2*Sqrt[1 + 1/(c^2*x^2)]*x^2) - 3*Log[1 + c^2*x^2])/(6*c^5)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 124 vs. 2(60) = 120.

Time = 0.86 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.74

method	result	size
default	$\frac{\sqrt{\frac{c^2 x^2 + 1}{c^2 x^2}} x \left(\left(\frac{c^2 x^2 + 1}{c^2} \right)^{\frac{3}{2}} c^2 - 3 \sqrt{-\frac{(-c^2 x + \sqrt{-c^2})(c^2 x + \sqrt{-c^2})}{c^4}} \right)}{3c^4 \sqrt{\frac{c^2 x^2 + 1}{c^2}}} + \frac{\frac{x^2}{2c^2} - \frac{\ln(c^2 x^2 + 1)}{2c^4}}{c}$	125

[In] int((1/c/x+(1+1/c^2/x^2)^(1/2))*x^4/(c^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] 1/3*((c^2*x^2+1)/c^2/x^2)^(1/2)*x/c^4*((1/c^2*(c^2*x^2+1))^(3/2)*c^2-3*(-c^2*x+(-c^2)^(1/2))*(c^2*x+(-c^2)^(1/2))/c^4)^(1/2))/(1/c^2*(c^2*x^2+1))^(1/2)+1/c*(1/2*x^2/c^2-1/2/c^4*ln(c^2*x^2+1))

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.81

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^4}{1 + c^2 x^2} dx = \frac{3c^2 x^2 + 2(c^3 x^3 - 2cx) \sqrt{\frac{c^2 x^2 + 1}{c^2 x^2}} - 3 \log(c^2 x^2 + 1)}{6c^5}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*x^4/(c^2*x^2+1),x, algorithm="fricas")

[Out] 1/6*(3*c^2*x^2 + 2*(c^3*x^3 - 2*c*x)*sqrt((c^2*x^2 + 1)/(c^2*x^2)) - 3*log(c^2*x^2 + 1))/c^5

Sympy [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^4}{1 + c^2 x^2} dx = \frac{\int \frac{x^3}{c^2 x^2 + 1} dx + \int \frac{cx^4 \sqrt{1 + \frac{1}{c^2 x^2}}}{c^2 x^2 + 1} dx}{c}$$

[In] integrate((1/c/x+(1+1/c**2/x**2)**(1/2))*x**4/(c**2*x**2+1),x)

[Out] (Integral(x**3/(c**2*x**2 + 1), x) + Integral(c*x**4*sqrt(1 + 1/(c**2*x**2))/(c**2*x**2 + 1), x))/c

Maxima [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.68

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^4}{1 + c^2 x^2} dx = \frac{x^2}{2c^3} + \frac{\sqrt{c^2 x^2 + 1}(c^2 x^2 - 2)}{3c^5} - \frac{\log(c^2 x^2 + 1)}{2c^5}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*x^4/(c^2*x^2+1),x, algorithm="maxima")

[Out] 1/2*x^2/c^3 + 1/3*sqrt(c^2*x^2 + 1)*(c^2*x^2 - 2)/c^5 - 1/2*log(c^2*x^2 + 1)/c^5

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.18

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^4}{1 + c^2 x^2} dx = -\frac{\log(c^2 x^2 + 1)}{2c^5} + \frac{2|c|\operatorname{sgn}(x)}{3c^6} + \frac{2(c^2 x^2 + 1)^{\frac{3}{2}} c^{12} |c|\operatorname{sgn}(x) - 6\sqrt{c^2 x^2 + 1} c^{12} |c|\operatorname{sgn}(x) + 3(c^2 x^2 + 1) c^{13}}{6c^{18}}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*x^4/(c^2*x^2+1),x, algorithm="giac")

[Out] -1/2*log(c^2*x^2 + 1)/c^5 + 2/3*abs(c)*sgn(x)/c^6 + 1/6*(2*(c^2*x^2 + 1)^(3/2)*c^12*abs(c)*sgn(x) - 6*sqrt(c^2*x^2 + 1)*c^12*abs(c)*sgn(x) + 3*(c^2*x^2 + 1)*c^13)/c^18

Mupad [B] (verification not implemented)

Time = 5.22 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.85

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^4}{1 + c^2 x^2} dx = \frac{x^3 \sqrt{\frac{1}{c^2 x^2} + 1}}{3 c^2} - \frac{2 x \sqrt{\frac{1}{c^2 x^2} + 1}}{3 c^4} - \frac{\ln(c^2 x^2 + 1) - c^2 x^2}{2 c^5}$$

[In] int((x^4*((1/(c^2*x^2) + 1)^(1/2) + 1/(c*x)))/(c^2*x^2 + 1),x)

[Out] (x^3*(1/(c^2*x^2) + 1)^(1/2))/(3*c^2) - (2*x*(1/(c^2*x^2) + 1)^(1/2))/(3*c^4) - (log(c^2*x^2 + 1) - c^2*x^2)/(2*c^5)

3.62 $\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^3}{1+c^2x^2} dx$

Optimal result	379
Rubi [A] (verified)	379
Mathematica [A] (verified)	381
Maple [B] (verified)	381
Fricas [A] (verification not implemented)	382
Sympy [F]	382
Maxima [B] (verification not implemented)	382
Giac [A] (verification not implemented)	383
Mupad [B] (verification not implemented)	383

Optimal result

Integrand size = 21, antiderivative size = 59

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^3}{1+c^2x^2} dx = \frac{x}{c^3} + \frac{\sqrt{1 + \frac{1}{c^2x^2}} x^2}{2c^2} - \frac{\arctan(cx)}{c^4} - \frac{\operatorname{arctanh}\left(\sqrt{1 + \frac{1}{c^2x^2}}\right)}{2c^4}$$

[Out] $x/c^3 - \arctan(cx)/c^4 - 1/2 * \operatorname{arctanh}((1+1/c^2/x^2)^{(1/2)})/c^4 + 1/2 * x^2 * (1+1/c^2/x^2)^{(1/2)}/c^2$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6477, 272, 44, 65, 214, 327, 209}

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^3}{1+c^2x^2} dx = -\frac{\arctan(cx)}{c^4} - \frac{\operatorname{arctanh}\left(\sqrt{\frac{1}{c^2x^2} + 1}\right)}{2c^4} + \frac{x}{c^3} + \frac{x^2 \sqrt{\frac{1}{c^2x^2} + 1}}{2c^2}$$

[In] $\text{Int}[(E^{\operatorname{ArcCsch}[c*x]} * x^3)/(1 + c^2*x^2), x]$

[Out] $x/c^3 + (\operatorname{Sqrt}[1 + 1/(c^2*x^2)] * x^2)/(2*c^2) - \operatorname{ArcTan}[c*x]/c^4 - \operatorname{ArcTanh}[\operatorname{Sqrt}[1 + 1/(c^2*x^2)]]/(2*c^4)$

Rule 44

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)} * ((c + d*x)^{(n + 1)} / ((b*c - a*d)*(m + 1))), x] - \text{Dist}[d * ((m + n + 2) / ((b*c - a*d)*(m + 1))), \text{Int}[(a + b*x)^{(m + 1)} * (c + d*x)^n, x], x] /;$ $\text{FreeQ}\{a, b, c, d, n\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{ILtQ}[m, -1] \ \&\& \ !\text{Int}$

egerQ[n] && LtQ[n, 0]

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 327

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[
a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 6477

```
Int[(E^ArcCsch[(c_.)*(x_)])*((d_.)*(x_))^(m_.))/((a_) + (b_.)*(x_)^2), x_Sym
bol] := Dist[d^2/(a*c^2), Int[(d*x)^(m - 2)/Sqrt[1 + 1/(c^2*x^2)], x], x] +
Dist[d/c, Int[(d*x)^(m - 1)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, m},
x] && EqQ[b - a*c^2, 0]
```

Rubi steps

$$\text{integral} = \frac{\int \frac{x}{\sqrt{1 + \frac{1}{c^2 x^2}}} dx}{c^2} + \frac{\int \frac{x^2}{1 + c^2 x^2} dx}{c}$$

$$\begin{aligned}
&= \frac{x}{c^3} - \frac{\int \frac{1}{1+c^2x^2} dx}{c^3} - \frac{\text{Subst}\left(\int \frac{1}{x^2\sqrt{1+\frac{x}{c^2}}} dx, x, \frac{1}{x^2}\right)}{2c^2} \\
&= \frac{x}{c^3} + \frac{\sqrt{1+\frac{1}{c^2x^2}}x^2}{2c^2} - \frac{\arctan(cx)}{c^4} + \frac{\text{Subst}\left(\int \frac{1}{x\sqrt{1+\frac{x}{c^2}}} dx, x, \frac{1}{x^2}\right)}{4c^4} \\
&= \frac{x}{c^3} + \frac{\sqrt{1+\frac{1}{c^2x^2}}x^2}{2c^2} - \frac{\arctan(cx)}{c^4} + \frac{\text{Subst}\left(\int \frac{1}{-c^2+c^2x^2} dx, x, \sqrt{1+\frac{1}{c^2x^2}}\right)}{2c^2} \\
&= \frac{x}{c^3} + \frac{\sqrt{1+\frac{1}{c^2x^2}}x^2}{2c^2} - \frac{\arctan(cx)}{c^4} - \frac{\text{arctanh}\left(\sqrt{1+\frac{1}{c^2x^2}}\right)}{2c^4}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.92

$$\int \frac{e^{\text{csch}^{-1}(cx)} x^3}{1+c^2x^2} dx = -\frac{-cx\left(2+c\sqrt{1+\frac{1}{c^2x^2}}x\right)+2\arctan(cx)+\log\left(\left(1+\sqrt{1+\frac{1}{c^2x^2}}\right)x\right)}{2c^4}$$

[In] Integrate[(E^ArcCsch[c*x]*x^3)/(1+c^2*x^2),x]

[Out] -1/2*(-(c*x*(2+c*Sqrt[1+1/(c^2*x^2)]*x))+2*ArcTan[c*x]+Log[(1+Sqrt[1+1/(c^2*x^2)])*x])/c^4

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 137 vs. 2(51) = 102.

Time = 0.86 (sec) , antiderivative size = 138, normalized size of antiderivative = 2.34

method	result	size
default	$ \frac{\sqrt{\frac{c^2x^2+1}{c^2x^2}}x\left(x\sqrt{\frac{c^2x^2+1}{c^2}}c^2+\ln\left(x+\sqrt{\frac{c^2x^2+1}{c^2}}\right)-2\ln\left(x+\sqrt{-\frac{(-c^2x+\sqrt{-c^2})(c^2x+\sqrt{-c^2})}{c^4}}\right)\right)}{2\sqrt{\frac{c^2x^2+1}{c^2}}c^4} + \frac{\frac{x}{c^2}-\frac{\arctan(cx)}{c^3}}{c} $	138

[In] int((1/c/x+(1+1/c^2/x^2)^(1/2))*x^3/(c^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] 1/2*((c^2*x^2+1)/c^2/x^2)^(1/2)*x*(x*(1/c^2*(c^2*x^2+1))^(1/2)*c^2+ln(x+(1/c^2*(c^2*x^2+1))^(1/2))-2*ln(x+(-(-c^2*x+(-c^2)^(1/2))*(c^2*x+(-c^2)^(1/2)))/c^4)^(1/2)))/(1/c^2*(c^2*x^2+1))^(1/2)/c^4+1/c*(x/c^2-1/c^3*arctan(c*x))

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.15

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^3}{1 + c^2 x^2} dx = \frac{c^2 x^2 \sqrt{\frac{c^2 x^2 + 1}{c^2 x^2}} + 2cx - 2 \arctan(cx) + \log\left(cx \sqrt{\frac{c^2 x^2 + 1}{c^2 x^2}} - cx\right)}{2c^4}$$

```
[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*x^3/(c^2*x^2+1),x, algorithm="fricas")
```

```
[Out] 1/2*(c^2*x^2*sqrt((c^2*x^2 + 1)/(c^2*x^2)) + 2*c*x - 2*arctan(c*x) + log(c*x*sqrt((c^2*x^2 + 1)/(c^2*x^2)) - c*x))/c^4
```

Sympy [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^3}{1 + c^2 x^2} dx = \int \frac{x^2}{c^2 x^2 + 1} dx + \int \frac{cx^3 \sqrt{1 + \frac{1}{c^2 x^2}}}{c^2 x^2 + 1} dx$$

```
[In] integrate((1/c/x+(1+1/c**2/x**2)**(1/2))*x**3/(c**2*x**2+1),x)
```

```
[Out] (Integral(x**2/(c**2*x**2 + 1), x) + Integral(c*x**3*sqrt(1 + 1/(c**2*x**2))/(c**2*x**2 + 1), x))/c
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 107 vs. 2(51) = 102.

Time = 0.33 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.81

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^3}{1 + c^2 x^2} dx = \frac{x}{c^3} + \frac{2 \sqrt{\frac{c^2 x^2 + 1}{x^2}} - \log\left(\frac{\sqrt{\frac{c^2 x^2 + 1}{x^2}}}{c} + 1\right) + \log\left(\frac{\sqrt{\frac{c^2 x^2 + 1}{x^2}}}{c} - 1\right)}{4c^4} - \frac{\arctan(cx)}{c^4}$$

```
[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*x^3/(c^2*x^2+1),x, algorithm="maxima")
```

```
[Out] x/c^3 + 1/4*(2*sqrt((c^2*x^2 + 1)/x^2)/(c*((c^2*x^2 + 1)/(c^2*x^2) - 1)) - log(sqrt((c^2*x^2 + 1)/x^2)/c + 1) + log(sqrt((c^2*x^2 + 1)/x^2)/c - 1))/c^4 - arctan(c*x)/c^4
```

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.03

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^3}{1 + c^2 x^2} dx = \frac{\sqrt{c^2 x^2 + 1} x |c| \operatorname{sgn}(x)}{2 c^4} + \frac{x}{c^3} + \frac{\log(-x|c| + \sqrt{c^2 x^2 + 1}) \operatorname{sgn}(x)}{2 c^4} - \frac{\arctan(cx)}{c^4}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*x^3/(c^2*x^2+1),x, algorithm="giac")

[Out] 1/2*sqrt(c^2*x^2 + 1)*x*abs(c)*sgn(x)/c^4 + x/c^3 + 1/2*log(-x*abs(c) + sqrt(c^2*x^2 + 1))*sgn(x)/c^4 - arctan(c*x)/c^4

Mupad [B] (verification not implemented)

Time = 5.25 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.86

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^3}{1 + c^2 x^2} dx = \frac{x^2 \sqrt{\frac{1}{c^2 x^2} + 1}}{2 c^2} - \frac{\operatorname{atan}(cx) - cx}{c^4} - \frac{\operatorname{atanh}\left(\sqrt{\frac{1}{c^2 x^2} + 1}\right)}{2 c^4}$$

[In] int((x^3*((1/(c^2*x^2) + 1)^(1/2) + 1/(c*x)))/(c^2*x^2 + 1),x)

[Out] (x^2*(1/(c^2*x^2) + 1)^(1/2))/(2*c^2) - (atan(c*x) - c*x)/c^4 - atanh((1/(c^2*x^2) + 1)^(1/2))/(2*c^4)

3.63 $\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^2}{1+c^2x^2} dx$

Optimal result	384
Rubi [A] (verified)	384
Mathematica [A] (verified)	385
Maple [B] (verified)	385
Fricas [A] (verification not implemented)	386
Sympy [F]	386
Maxima [A] (verification not implemented)	386
Giac [A] (verification not implemented)	387
Mupad [B] (verification not implemented)	387

Optimal result

Integrand size = 21, antiderivative size = 36

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^2}{1+c^2x^2} dx = \frac{\sqrt{1+\frac{1}{c^2x^2}} x}{c^2} + \frac{\log(1+c^2x^2)}{2c^3}$$

[Out] $1/2*\ln(c^2*x^2+1)/c^3+x*(1+1/c^2/x^2)^(1/2)/c^2$

Rubi [A] (verified)

Time = 0.05 (sec), antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6477, 197, 266}

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^2}{1+c^2x^2} dx = \frac{x\sqrt{\frac{1}{c^2x^2}+1}}{c^2} + \frac{\log(c^2x^2+1)}{2c^3}$$

[In] `Int[(E^ArcCsch[c*x]*x^2)/(1+c^2*x^2),x]`

[Out] `(Sqrt[1+1/(c^2*x^2)]*x)/c^2+Log[1+c^2*x^2]/(2*c^3)`

Rule 197

`Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

Rule 266

`Int[(x_)^(m_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]`

Rule 6477

Int[(E^ArcCsch[(c_.)*(x_)]*((d_.)*(x_))^(m_.))/((a_) + (b_.)*(x_)^2), x_Symbol] := Dist[d^2/(a*c^2), Int[(d*x)^(m - 2)/Sqrt[1 + 1/(c^2*x^2)], x], x] + Dist[d/c, Int[(d*x)^(m - 1)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b - a*c^2, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \frac{1}{\sqrt{1 + \frac{1}{c^2 x^2}}} dx}{c^2} + \frac{\int \frac{x}{1 + c^2 x^2} dx}{c} \\ &= \frac{\sqrt{1 + \frac{1}{c^2 x^2}}}{c^2} + \frac{\log(1 + c^2 x^2)}{2c^3} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.97

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^2}{1 + c^2 x^2} dx = \frac{2c \sqrt{1 + \frac{1}{c^2 x^2}} x + \log(1 + c^2 x^2)}{2c^3}$$

[In] Integrate[(E^ArcCsch[c*x]*x^2)/(1 + c^2*x^2), x]

[Out] (2*c*Sqrt[1 + 1/(c^2*x^2)]*x + Log[1 + c^2*x^2])/(2*c^3)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 88 vs. 2(32) = 64.

Time = 0.85 (sec) , antiderivative size = 89, normalized size of antiderivative = 2.47

method	result	size
default	$\frac{\sqrt{\frac{c^2 x^2 + 1}{c^2 x^2}} x \sqrt{-\frac{(-c^2 x + \sqrt{-c^2})(c^2 x + \sqrt{-c^2})}{c^4}}}{\sqrt{\frac{c^2 x^2 + 1}{c^2}} c^2} + \frac{\ln(c^2 x^2 + 1)}{2c^3}$	89

[In] int((1/c/x+(1+1/c^2/x^2)^(1/2))*x^2/(c^2*x^2+1), x, method=_RETURNVERBOSE)

[Out] ((c^2*x^2+1)/c^2/x^2)^(1/2)*x*(-(-c^2*x+(-c^2)^(1/2))*(c^2*x+(-c^2)^(1/2))/c^4)^(1/2)/(1/c^2*(c^2*x^2+1))^(1/2)/c^2+1/2*ln(c^2*x^2+1)/c^3

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.06

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^2}{1 + c^2 x^2} dx = \frac{2 cx \sqrt{\frac{c^2 x^2 + 1}{c^2 x^2}} + \log(c^2 x^2 + 1)}{2 c^3}$$

```
[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*x^2/(c^2*x^2+1),x, algorithm="fricas")
```

```
[Out] 1/2*(2*c*x*sqrt((c^2*x^2 + 1)/(c^2*x^2)) + log(c^2*x^2 + 1))/c^3
```

Sympy [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^2}{1 + c^2 x^2} dx = \int \frac{x}{c^2 x^2 + 1} dx + \int \frac{cx^2 \sqrt{1 + \frac{1}{c^2 x^2}}}{c^2 x^2 + 1} dx$$

```
[In] integrate((1/c/x+(1+1/c**2/x**2)**(1/2))*x**2/(c**2*x**2+1),x)
```

```
[Out] (Integral(x/(c**2*x**2 + 1), x) + Integral(c*x**2*sqrt(1 + 1/(c**2*x**2))/(c**2*x**2 + 1), x))/c
```

Maxima [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^2}{1 + c^2 x^2} dx = \frac{\log(c^3 x^2 + c)}{2 c^3} + \frac{\sqrt{c^2 x^2 + 1}}{c^3}$$

```
[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*x^2/(c^2*x^2+1),x, algorithm="maxima")
```

```
[Out] 1/2*log(c^3*x^2 + c)/c^3 + sqrt(c^2*x^2 + 1)/c^3
```

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.22

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^2}{1 + c^2 x^2} dx = \frac{\sqrt{c^2 x^2 + 1} |c| \operatorname{sgn}(x)}{c^4} + \frac{\log(c^2 x^2 + 1)}{2 c^3} - \frac{|c| \operatorname{sgn}(x)}{c^4}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*x^2/(c^2*x^2+1),x, algorithm="giac")

[Out] sqrt(c^2*x^2 + 1)*abs(c)*sgn(x)/c^4 + 1/2*log(c^2*x^2 + 1)/c^3 - abs(c)*sgn(x)/c^4

Mupad [B] (verification not implemented)

Time = 5.38 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x^2}{1 + c^2 x^2} dx = \frac{\ln(c^2 x^2 + 1) + 2 c x \sqrt{\frac{1}{c^2 x^2} + 1}}{2 c^3}$$

[In] int((x^2*((1/(c^2*x^2) + 1)^(1/2) + 1/(c*x)))/(c^2*x^2 + 1),x)

[Out] (log(c^2*x^2 + 1) + 2*c*x*(1/(c^2*x^2) + 1)^(1/2))/(2*c^3)

3.64 $\int \frac{e^{\operatorname{csch}^{-1}(cx)x}}{1+c^2x^2} dx$

Optimal result	388
Rubi [A] (verified)	388
Mathematica [A] (verified)	390
Maple [B] (verified)	390
Fricas [A] (verification not implemented)	390
Sympy [F]	391
Maxima [B] (verification not implemented)	391
Giac [A] (verification not implemented)	391
Mupad [B] (verification not implemented)	392

Optimal result

Integrand size = 19, antiderivative size = 27

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)x}}{1+c^2x^2} dx = \frac{\arctan(cx)}{c^2} + \frac{\operatorname{arctanh}\left(\sqrt{1+\frac{1}{c^2x^2}}\right)}{c^2}$$

[Out] $\arctan(c*x)/c^2 + \operatorname{arctanh}((1+1/c^2/x^2)^{(1/2)})/c^2$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {6477, 272, 65, 214, 209}

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)x}}{1+c^2x^2} dx = \frac{\arctan(cx)}{c^2} + \frac{\operatorname{arctanh}\left(\sqrt{\frac{1}{c^2x^2}+1}\right)}{c^2}$$

[In] $\operatorname{Int}[(E^{\operatorname{ArcCsch}[c*x]*x})/(1+c^2*x^2),x]$

[Out] $\operatorname{ArcTan}[c*x]/c^2 + \operatorname{ArcTanh}[\operatorname{Sqrt}[1+1/(c^2*x^2)]]/c^2$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] :> \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}], x]] /;$ $\operatorname{FreeQ}[\{a, b, c, d\}, x] \ \&\& \operatorname{NeQ}[b*c - a*d, 0] \ \&\& \operatorname{LtQ}[-1, m, 0] \ \&\& \operatorname{LeQ}[-1, n, 0] \ \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]] \ \&\& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 272

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 6477

Int[(E^ArcCsch[(c_)*(x_)])*((d_)*(x_)^(m_)))/((a_) + (b_)*(x_)^2), x_Symbol] := Dist[d^2/(a*c^2), Int[(d*x)^(m - 2)/Sqrt[1 + 1/(c^2*x^2)], x], x] + Dist[d/c, Int[(d*x)^(m - 1)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b - a*c^2, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int \frac{1}{\sqrt{1 + \frac{1}{c^2 x^2}}} dx}{c^2} + \frac{\int \frac{1}{1 + c^2 x^2} dx}{c} \\
 &= \frac{\arctan(cx)}{c^2} - \frac{\text{Subst}\left(\int \frac{1}{x\sqrt{1 + \frac{x}{c^2}}} dx, x, \frac{1}{x^2}\right)}{2c^2} \\
 &= \frac{\arctan(cx)}{c^2} - \text{Subst}\left(\int \frac{1}{-c^2 + c^2 x^2} dx, x, \sqrt{1 + \frac{1}{c^2 x^2}}\right) \\
 &= \frac{\arctan(cx)}{c^2} + \frac{\text{arctanh}\left(\sqrt{1 + \frac{1}{c^2 x^2}}\right)}{c^2}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.41

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x}{1 + c^2 x^2} dx = \frac{\arctan(cx)}{c^2} + \frac{\log\left(x\left(1 + \sqrt{\frac{1+c^2x^2}{c^2x^2}}\right)\right)}{c^2}$$

[In] Integrate[(E^ArcCsch[c*x]*x)/(1 + c^2*x^2),x]

[Out] ArcTan[c*x]/c^2 + Log[x*(1 + Sqrt[(1 + c^2*x^2)/(c^2*x^2)])]/c^2

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 84 vs. 2(25) = 50.

Time = 0.86 (sec) , antiderivative size = 85, normalized size of antiderivative = 3.15

method	result	size
default	$\frac{\sqrt{\frac{c^2x^2+1}{c^2x^2}} x \ln\left(x + \sqrt{-\frac{(-c^2x + \sqrt{-c^2})(c^2x + \sqrt{-c^2})}{c^4}}\right)}{\sqrt{\frac{c^2x^2+1}{c^2}} c^2} + \frac{\arctan(cx)}{c^2}$	85

[In] int((1/c/x+(1+1/c^2/x^2)^(1/2))*x/(c^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] ((c^2*x^2+1)/c^2/x^2)^(1/2)*x*ln(x+(-(-c^2*x+(-c^2)^(1/2))*(c^2*x+(-c^2)^(1/2)))/c^4)^(1/2))/(1/c^2*(c^2*x^2+1))^(1/2)/c^2+arctan(c*x)/c^2

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.41

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x}{1 + c^2 x^2} dx = \frac{\arctan(cx) - \log\left(cx\sqrt{\frac{c^2x^2+1}{c^2x^2}} - cx\right)}{c^2}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*x/(c^2*x^2+1),x, algorithm="fricas")

[Out] (arctan(c*x) - log(c*x*sqrt((c^2*x^2 + 1)/(c^2*x^2)) - c*x))/c^2

Sympy [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x}{1 + c^2 x^2} dx = \int \frac{cx \sqrt{1 + \frac{1}{c^2 x^2}}}{c^2 x^2 + 1} dx + \int \frac{1}{c^2 x^2 + 1} dx$$

[In] integrate((1/c/x+(1+1/c**2/x**2)**(1/2))*x/(c**2*x**2+1),x)

[Out] (Integral(c*x*sqrt(1 + 1/(c**2*x**2)))/(c**2*x**2 + 1), x) + Integral(1/(c**2*x**2 + 1), x))/c

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs. 2(25) = 50.

Time = 0.31 (sec) , antiderivative size = 61, normalized size of antiderivative = 2.26

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x}{1 + c^2 x^2} dx = \frac{\log\left(\frac{\sqrt{\frac{c^2 x^2 + 1}{x^2}} + 1}{c}\right) - \log\left(\frac{\sqrt{\frac{c^2 x^2 + 1}{x^2}} - 1}{c}\right)}{2c^2} + \frac{\arctan(cx)}{c^2}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*x/(c^2*x^2+1),x, algorithm="maxima")

[Out] 1/2*(log(sqrt((c^2*x^2 + 1)/x^2)/c + 1) - log(sqrt((c^2*x^2 + 1)/x^2)/c - 1))/c^2 + arctan(c*x)/c^2

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.26

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x}{1 + c^2 x^2} dx = -\frac{\log(-x|c| + \sqrt{c^2 x^2 + 1}) \operatorname{sgn}(x)}{c^2} + \frac{\arctan(cx)}{c^2}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))*x/(c^2*x^2+1),x, algorithm="giac")

[Out] -log(-x*abs(c) + sqrt(c^2*x^2 + 1))*sgn(x)/c^2 + arctan(c*x)/c^2

Mupad [B] (verification not implemented)

Time = 5.82 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)} x}{1 + c^2 x^2} dx = \frac{\operatorname{atanh}\left(\sqrt{\frac{1}{c^2 x^2} + 1}\right) + \operatorname{atan}(cx)}{c^2}$$

[In] `int((x*((1/(c^2*x^2) + 1)^(1/2) + 1/(c*x)))/(c^2*x^2 + 1),x)`

[Out] `(atanh((1/(c^2*x^2) + 1)^(1/2)) + atan(c*x))/c^2`

3.65 $\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{1+c^2x^2} dx$

Optimal result	393
Rubi [A] (verified)	393
Mathematica [A] (verified)	395
Maple [B] (verified)	395
Fricas [B] (verification not implemented)	395
Sympy [F]	396
Maxima [F]	396
Giac [B] (verification not implemented)	396
Mupad [B] (verification not implemented)	397

Optimal result

Integrand size = 18, antiderivative size = 33

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{1+c^2x^2} dx = -\frac{\operatorname{csch}^{-1}(cx)}{c} + \frac{\log(x)}{c} - \frac{\log(1+c^2x^2)}{2c}$$

[Out] $-\operatorname{arccsch}(c*x)/c+\ln(x)/c-1/2*\ln(c^2*x^2+1)/c$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {6475, 342, 221, 272, 36, 29, 31}

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{1+c^2x^2} dx = -\frac{\log(c^2x^2+1)}{2c} + \frac{\log(x)}{c} - \frac{\operatorname{csch}^{-1}(cx)}{c}$$

[In] $\operatorname{Int}[E^{\operatorname{ArcCsch}[c*x]}/(1+c^2*x^2),x]$

[Out] $-(\operatorname{ArcCsch}[c*x]/c) + \operatorname{Log}[x]/c - \operatorname{Log}[1+c^2*x^2]/(2*c)$

Rule 29

$\operatorname{Int}[(x_)^{(-1)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Log}[x], x]$

Rule 31

$\operatorname{Int}[((a_) + (b_)*(x_))^{(-1)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Log}[\operatorname{RemoveContent}[a + b*x, x]]/b, x] /; \operatorname{FreeQ}\{a, b\}, x]$

Rule 36

`Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Dist[b/(b*c - a*d), Int[1/(a + b*x), x], x] - Dist[d/(b*c - a*d), Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

Rule 221

`Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

Rule 272

`Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

Rule 342

`Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]`

Rule 6475

`Int[E^ArcCsch[(c_.)*(x_)]/((a_) + (b_.)*(x_)^2), x_Symbol] := Dist[1/(a*c^2), Int[1/(x^2*sqrt[1 + 1/(c^2*x^2)]), x], x] + Dist[1/c, Int[1/(x*(a + b*x^2)), x], x] /; FreeQ[{a, b, c}, x] && EqQ[b - a*c^2, 0]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int \frac{1}{\sqrt{1 + \frac{1}{c^2 x^2} x^2}} dx}{c^2} + \frac{\int \frac{1}{x(1 + c^2 x^2)} dx}{c} \\
 &= -\frac{\text{Subst}\left(\int \frac{1}{\sqrt{1 + \frac{x^2}{c^2}}} dx, x, \frac{1}{x}\right)}{c^2} + \frac{\text{Subst}\left(\int \frac{1}{x(1 + c^2 x)} dx, x, x^2\right)}{2c} \\
 &= -\frac{\text{csch}^{-1}(cx)}{c} + \frac{\text{Subst}\left(\int \frac{1}{x} dx, x, x^2\right)}{2c} - \frac{1}{2}c \text{Subst}\left(\int \frac{1}{1 + c^2 x} dx, x, x^2\right) \\
 &= -\frac{\text{csch}^{-1}(cx)}{c} + \frac{\log(x)}{c} - \frac{\log(1 + c^2 x^2)}{2c}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.12

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{1+c^2x^2} dx = -\frac{\operatorname{arcsinh}\left(\frac{1}{cx}\right)}{c} + \frac{\log(x)}{c} - \frac{\log(1+c^2x^2)}{2c}$$

[In] Integrate[E^ArcCsch[c*x]/(1 + c^2*x^2), x]

[Out] -(ArcSinh[1/(c*x)]/c) + Log[x]/c - Log[1 + c^2*x^2]/(2*c)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 169 vs. 2(31) = 62.

Time = 0.88 (sec) , antiderivative size = 170, normalized size of antiderivative = 5.15

method	result
default	$\frac{\sqrt{\frac{c^2x^2+1}{c^2x^2}} x \left(\sqrt{\frac{1}{c^2}} \sqrt{\frac{c^2x^2+1}{c^2}} c^2 - \sqrt{-\frac{(-c^2x+\sqrt{-c^2})(c^2x+\sqrt{-c^2})}{c^4}} c^2 \sqrt{\frac{1}{c^2}} - \ln\left(\frac{2\sqrt{\frac{1}{c^2}} \sqrt{\frac{c^2x^2+1}{c^2}} c^{2+2}}{c^2x}\right) \right)}{\sqrt{\frac{1}{c^2}} \sqrt{\frac{c^2x^2+1}{c^2}} c^2} + \frac{\ln(x) - \frac{\ln(c^2x^2+1)}{2}}{c}$

[In] int((1/c/x+(1+1/c^2/x^2)^(1/2))/(c^2*x^2+1), x, method=_RETURNVERBOSE)

[Out] ((c^2*x^2+1)/c^2/x^2)^(1/2)*x*((1/c^2)^(1/2)*(1/c^2*(c^2*x^2+1))^(1/2)*c^2-(-c^2*x+(-c^2)^(1/2))*(c^2*x+(-c^2)^(1/2))/c^4)^(1/2)*c^2*(1/c^2)^(1/2)-1/n(2*((1/c^2)^(1/2)*(1/c^2*(c^2*x^2+1))^(1/2)*c^2+1)/c^2/x)/(1/c^2)^(1/2)/(1/c^2*(c^2*x^2+1))^(1/2)/c^2+1/c*(ln(x)-1/2*ln(c^2*x^2+1))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 80 vs. 2(31) = 62.

Time = 0.25 (sec) , antiderivative size = 80, normalized size of antiderivative = 2.42

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{1+c^2x^2} dx = \frac{\log(c^2x^2+1) + 2 \log\left(cx\sqrt{\frac{c^2x^2+1}{c^2x^2}} - cx + 1\right) - 2 \log\left(cx\sqrt{\frac{c^2x^2+1}{c^2x^2}} - cx - 1\right) - 2 \log(x)}{2c}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))/(c^2*x^2+1), x, algorithm="fricas")

[Out] -1/2*(log(c^2*x^2 + 1) + 2*log(c*x*sqrt((c^2*x^2 + 1)/(c^2*x^2)) - c*x + 1) - 2*log(c*x*sqrt((c^2*x^2 + 1)/(c^2*x^2)) - c*x - 1) - 2*log(x))/c

SymPy [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{1+c^2x^2} dx = \frac{\int \frac{cx\sqrt{1+\frac{1}{c^2x^2}}}{c^2x^3+x} dx + \int \frac{1}{c^2x^3+x} dx}{c}$$

[In] integrate((1/c/x+(1+1/c**2/x**2)**(1/2))/(c**2*x**2+1),x)

[Out] (Integral(c*x*sqrt(1 + 1/(c**2*x**2)))/(c**2*x**3 + x), x) + Integral(1/(c**2*x**3 + x), x))/c

Maxima [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{1+c^2x^2} dx = \int \frac{\sqrt{\frac{1}{c^2x^2} + 1} + \frac{1}{cx}}{c^2x^2 + 1} dx$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))/(c^2*x^2+1),x, algorithm="maxima")

[Out] -1/2*log(c^2*x^2 + 1)/c + log(x)/c + integrate(sqrt(c^2*x^2 + 1)/(c^3*x^3 + c*x), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 70 vs. 2(31) = 62.

Time = 0.29 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.12

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{1+c^2x^2} dx = -\frac{\log(c^2x^2 + 1)}{2c} - \frac{(|c|\operatorname{sgn}(x) - c) \log(\sqrt{c^2x^2 + 1} + 1)}{2c^2} + \frac{(|c|\operatorname{sgn}(x) + c) \log(\sqrt{c^2x^2 + 1} - 1)}{2c^2}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))/(c^2*x^2+1),x, algorithm="giac")

[Out] -1/2*log(c^2*x^2 + 1)/c - 1/2*(abs(c)*sgn(x) - c)*log(sqrt(c^2*x^2 + 1) + 1)/c^2 + 1/2*(abs(c)*sgn(x) + c)*log(sqrt(c^2*x^2 + 1) - 1)/c^2

Mupad [B] (verification not implemented)

Time = 5.41 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.15

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{1+c^2x^2} dx = -\operatorname{asinh}\left(\frac{\sqrt{\frac{1}{c^2}}}{x}\right) \sqrt{\frac{1}{c^2}} - \frac{\ln(c^2x^2+1) - 2\ln(x)}{2c}$$

[In] int(((1/(c^2*x^2) + 1)^(1/2) + 1/(c*x))/(c^2*x^2 + 1),x)

[Out] - asinh((1/c^2)^(1/2)/x)*(1/c^2)^(1/2) - (log(c^2*x^2 + 1) - 2*log(x))/(2*c
)

3.66 $\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x(1+c^2x^2)} dx$

Optimal result	398
Rubi [A] (verified)	398
Mathematica [A] (verified)	399
Maple [B] (verified)	400
Fricas [A] (verification not implemented)	400
Sympy [A] (verification not implemented)	400
Maxima [A] (verification not implemented)	401
Giac [A] (verification not implemented)	401
Mupad [B] (verification not implemented)	401

Optimal result

Integrand size = 21, antiderivative size = 30

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x(1+c^2x^2)} dx = -\sqrt{1+\frac{1}{c^2x^2}} - \frac{1}{cx} - \arctan(cx)$$

[Out] $-1/c/x - \arctan(c*x) - (1+1/c^2/x^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec), antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {6477, 267, 331, 209}

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x(1+c^2x^2)} dx = -\arctan(cx) - \sqrt{\frac{1}{c^2x^2} + 1} - \frac{1}{cx}$$

[In] $\text{Int}[E^{\text{ArcCsch}[c*x]}/(x*(1+c^2*x^2)),x]$

[Out] $-\text{Sqrt}[1+1/(c^2*x^2)] - 1/(c*x) - \text{ArcTan}[c*x]$

Rule 209

$\text{Int}[(a_+ + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rule 267

$\text{Int}[(x_+)^{(m_+)}*((a_+ + (b_+)(x_+)^n)^{(p_+)}), x_Symbol] \rightarrow \text{Simp}[(a + b*x^n)^{(p+1)}/(b*n*(p+1)), x] /; \text{FreeQ}\{a, b, m, n, p, x\} \ \&\& \ \text{EqQ}[m, n-1] \ \&\&$

NeQ[p, -1]

Rule 331

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Dist[b*((m + n*(p + 1) + 1)/(a*c^n*(m + 1))], Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 6477

```
Int[(E^ArcCsch[(c_.)*(x_)]*((d_.)*(x_)^(m_.)))/((a_) + (b_.)*(x_)^2), x_Symbol] := Dist[d^2/(a*c^2), Int[(d*x)^(m - 2)/Sqrt[1 + 1/(c^2*x^2)], x], x] + Dist[d/c, Int[(d*x)^(m - 1)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b - a*c^2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \frac{1}{\sqrt{1 + \frac{1}{c^2 x^2}} x^3} dx}{c^2} + \frac{\int \frac{1}{x^2(1 + c^2 x^2)} dx}{c} \\ &= -\sqrt{1 + \frac{1}{c^2 x^2}} - \frac{1}{cx} - c \int \frac{1}{1 + c^2 x^2} dx \\ &= -\sqrt{1 + \frac{1}{c^2 x^2}} - \frac{1}{cx} - \arctan(cx) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{e^{\text{csch}^{-1}(cx)}}{x(1 + c^2 x^2)} dx = -\sqrt{1 + \frac{1}{c^2 x^2}} - \frac{1}{cx} - \arctan(cx)$$

```
[In] Integrate[E^ArcCsch[c*x]/(x*(1 + c^2*x^2)),x]
```

```
[Out] -Sqrt[1 + 1/(c^2*x^2)] - 1/(c*x) - ArcTan[c*x]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 156 vs. 2(28) = 56.

Time = 0.85 (sec) , antiderivative size = 157, normalized size of antiderivative = 5.23

method	result
default	$-\frac{\sqrt{\frac{c^2x^2+1}{c^2x^2}} \left(\left(\frac{c^2x^2+1}{c^2} \right)^{\frac{3}{2}} c^2 - \sqrt{\frac{c^2x^2+1}{c^2}} c^2 x^2 - \ln \left(x + \sqrt{\frac{c^2x^2+1}{c^2}} \right) x + \ln \left(x + \sqrt{-\frac{(-c^2x + \sqrt{-c^2})(c^2x + \sqrt{-c^2})}{c^4}} \right) x \right)}{\sqrt{\frac{c^2x^2+1}{c^2}}} + \frac{-\frac{1}{x} - c \arctan(cx)}{c}$

[In] int((1/c/x+(1+1/c^2/x^2)^(1/2))/x/(c^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] -((c^2*x^2+1)/c^2/x^2)^(1/2)*((1/c^2*(c^2*x^2+1))^(3/2)*c^2-(1/c^2*(c^2*x^2+1))^(1/2)*c^2*x^2-ln(x+(1/c^2*(c^2*x^2+1))^(1/2))*x+ln(x+(-(-c^2*x+(-c^2)^(1/2))*(c^2*x+(-c^2)^(1/2))/c^4)^(1/2))*x)/(1/c^2*(c^2*x^2+1))^(1/2)+1/c*(-1/x-c*arctan(c*x))

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.37

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x(1+c^2x^2)} dx = -\frac{cx \arctan(cx) + cx \sqrt{\frac{c^2x^2+1}{c^2x^2}} + cx + 1}{cx}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))/x/(c^2*x^2+1),x, algorithm="fricas")

[Out] -(c*x*arctan(c*x) + c*x*sqrt((c^2*x^2 + 1)/(c^2*x^2)) + c*x + 1)/(c*x)

Sympy [A] (verification not implemented)

Time = 2.26 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.40

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x(1+c^2x^2)} dx = -c \left(\begin{cases} \text{NaN} & \text{for } c = 0 \\ \frac{\sqrt{1+\frac{1}{c^2x^2}}}{c} & \text{otherwise} \end{cases} \right) + \frac{c \operatorname{atan}\left(\frac{1}{x\sqrt{c^2}}\right)}{\sqrt{c^2}} - \frac{1}{cx}$$

[In] integrate((1/c/x+(1+1/c**2/x**2)**(1/2))/x/(c**2*x**2+1),x)

[Out] -c*Piecewise((nan, Eq(c, 0)), (sqrt(1 + 1/(c**2*x**2))/c, True)) + c*atan(1/(x*sqrt(c**2)))/sqrt(c**2) - 1/(c*x)

Maxima [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.13

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x(1+c^2x^2)} dx = -\frac{\sqrt{c^2x^2+1}}{cx} - \frac{1}{cx} - \arctan(cx)$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))/x/(c^2*x^2+1),x, algorithm="maxima")

[Out] -sqrt(c^2*x^2 + 1)/(c*x) - 1/(c*x) - arctan(c*x)

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.43

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x(1+c^2x^2)} dx = \frac{2 \operatorname{sgn}(x)}{(x|c| - \sqrt{c^2x^2+1})^2 - 1} - \frac{1}{cx} - \arctan(cx)$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))/x/(c^2*x^2+1),x, algorithm="giac")

[Out] 2*sgn(x)/((x*abs(c) - sqrt(c^2*x^2 + 1))^2 - 1) - 1/(c*x) - arctan(c*x)

Mupad [B] (verification not implemented)

Time = 5.51 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.97

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x(1+c^2x^2)} dx = -\operatorname{atan}(cx) - \frac{x \sqrt{\frac{1}{c^2x^2} + 1} + \frac{1}{c}}{x}$$

[In] int(((1/(c^2*x^2) + 1)^(1/2) + 1/(c*x))/(x*(c^2*x^2 + 1)),x)

[Out] - atan(c*x) - (x*(1/(c^2*x^2) + 1)^(1/2) + 1/c)/x

3.67 $\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^2(1+c^2x^2)} dx$

Optimal result	402
Rubi [A] (verified)	402
Mathematica [A] (verified)	404
Maple [B] (verified)	404
Fricas [B] (verification not implemented)	404
Sympy [F]	405
Maxima [F]	405
Giac [B] (verification not implemented)	405
Mupad [B] (verification not implemented)	406

Optimal result

Integrand size = 21, antiderivative size = 60

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^2(1+c^2x^2)} dx = -\frac{1}{2cx^2} - \frac{\sqrt{1 + \frac{1}{c^2x^2}}}{2x} + \frac{1}{2}c\operatorname{csch}^{-1}(cx) - c\log(x) + \frac{1}{2}c\log(1+c^2x^2)$$

[Out] $-1/2/c/x^2+1/2*c*\operatorname{arccsch}(c*x)-c*\ln(x)+1/2*c*\ln(c^2*x^2+1)-1/2*(1+1/c^2/x^2)^{(1/2)}/x$

Rubi [A] (verified)

Time = 0.07 (sec), antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6477, 342, 327, 221, 272, 46}

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^2(1+c^2x^2)} dx = -\frac{\sqrt{\frac{1}{c^2x^2} + 1}}{2x} + \frac{1}{2}c\log(c^2x^2 + 1) - \frac{1}{2cx^2} - c\log(x) + \frac{1}{2}c\operatorname{csch}^{-1}(cx)$$

[In] $\operatorname{Int}[E^{\operatorname{ArcCsch}[c*x]}/(x^2*(1+c^2*x^2)),x]$

[Out] $-1/2*1/(c*x^2) - \operatorname{Sqrt}[1 + 1/(c^2*x^2)]/(2*x) + (c*\operatorname{ArcCsch}[c*x])/2 - c*\operatorname{Log}[x] + (c*\operatorname{Log}[1 + c^2*x^2])/2$

Rule 46

$\operatorname{Int}[(a_.) + (b_.)*(x_.)^{(m_.)}*((c_.) + (d_.)*(x_.)^{(n_.)}), x_Symbol] \rightarrow \operatorname{Int}[E^{\operatorname{xpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x]}, x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])

Rule 221

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 272

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 327

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 342

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && IntegerQ[m]

Rule 6477

Int[(E^ArcCsch[(c_)*(x_)])*((d_)*(x_))^(m_)/((a_) + (b_)*(x_)^2), x_Symbol] := Dist[d^2/(a*c^2), Int[(d*x)^(m - 2)/Sqrt[1 + 1/(c^2*x^2)], x], x] + Dist[d/c, Int[(d*x)^(m - 1)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b - a*c^2, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int \frac{1}{\sqrt{1 + \frac{1}{c^2 x^2}} x^4} dx}{c^2} + \frac{\int \frac{1}{x^3(1 + c^2 x^2)} dx}{c} \\
 &= -\frac{\text{Subst}\left(\int \frac{x^2}{\sqrt{1 + \frac{x^2}{c^2}}} dx, x, \frac{1}{x}\right)}{c^2} + \frac{\text{Subst}\left(\int \frac{1}{x^2(1 + c^2 x)} dx, x, x^2\right)}{2c} \\
 &= -\frac{\sqrt{1 + \frac{1}{c^2 x^2}}}{2x} + \frac{1}{2} \text{Subst}\left(\int \frac{1}{\sqrt{1 + \frac{x^2}{c^2}}} dx, x, \frac{1}{x}\right) + \frac{\text{Subst}\left(\int \left(\frac{1}{x^2} - \frac{c^2}{x} + \frac{c^4}{1 + c^2 x}\right) dx, x, x^2\right)}{2c} \\
 &= -\frac{1}{2cx^2} - \frac{\sqrt{1 + \frac{1}{c^2 x^2}}}{2x} + \frac{1}{2} \text{ccsch}^{-1}(cx) - c \log(x) + \frac{1}{2} c \log(1 + c^2 x^2)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.97

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^2(1+c^2x^2)} dx = \frac{1}{2} \left(-\frac{1}{cx^2} - \frac{\sqrt{1+\frac{1}{c^2x^2}}}{x} + \operatorname{carcsinh}\left(\frac{1}{cx}\right) - 2c \log(x) + c \log(1+c^2x^2) \right)$$

[In] Integrate[E^ArcCsch[c*x]/(x^2*(1+c^2*x^2)),x]

[Out] (-1/(c*x^2)) - Sqrt[1 + 1/(c^2*x^2)]/x + c*ArcSinh[1/(c*x)] - 2*c*Log[x] + c*Log[1 + c^2*x^2])/2

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 215 vs. 2(50) = 100.

Time = 0.86 (sec) , antiderivative size = 216, normalized size of antiderivative = 3.60

method	result
default	$-\frac{\sqrt{\frac{c^2x^2+1}{c^2x^2}} \left(c^2 \left(\frac{c^2x^2+1}{c^2} \right)^{\frac{3}{2}} \sqrt{\frac{1}{c^2} + \sqrt{\frac{c^2x^2+1}{c^2}}} \sqrt{\frac{1}{c^2}} c^2x^2 - 2\sqrt{\frac{1}{c^2}} \sqrt{-\frac{(-c^2x+\sqrt{-c^2})(c^2x+\sqrt{-c^2})}{c^4}} c^2x^2 - \ln\left(\frac{2\sqrt{\frac{1}{c^2}} \sqrt{\frac{c^2x^2+1}{c^2}} c^2+2}{c^2x}\right) \right)}{2x\sqrt{\frac{c^2x^2+1}{c^2}} \sqrt{\frac{1}{c^2}}} x^2$

[In] int((1/c/x+(1+1/c^2/x^2)^(1/2))/x^2/(c^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] -1/2*((c^2*x^2+1)/c^2/x^2)^(1/2)/x*(c^2*(1/c^2*(c^2*x^2+1))^(3/2)*(1/c^2)^(1/2)+(1/c^2*(c^2*x^2+1))^(1/2)*(1/c^2)^(1/2)*c^2*x^2-2*(1/c^2)^(1/2)*(-c^2*x+(-c^2)^(1/2))*(c^2*x+(-c^2)^(1/2))/c^4)^(1/2)*c^2*x^2-ln(2*((1/c^2)^(1/2)*(1/c^2*(c^2*x^2+1))^(1/2)*c^2+1)/c^2/x)*x^2)/(1/c^2*(c^2*x^2+1))^(1/2)/(1/c^2)^(1/2)+1/c*(-1/2/x^2-c^2*ln(x)+1/2*c^2*ln(c^2*x^2+1))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 130 vs. 2(50) = 100.

Time = 0.25 (sec) , antiderivative size = 130, normalized size of antiderivative = 2.17

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^2(1+c^2x^2)} dx = \frac{c^2x^2 \log(c^2x^2+1) + c^2x^2 \log\left(cx\sqrt{\frac{c^2x^2+1}{c^2x^2}} - cx + 1\right) - c^2x^2 \log\left(cx\sqrt{\frac{c^2x^2+1}{c^2x^2}} - cx - 1\right) - 2c^2x^2 \log(x)}{2cx^2}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))/x^2/(c^2*x^2+1),x, algorithm="fricas")

[Out] $\frac{1}{2}(c^2x^2 \log(c^2x^2 + 1) + c^2x^2 \log(cx \sqrt{(c^2x^2 + 1)/(c^2x^2)})) - cx + 1 - c^2x^2 \log(cx \sqrt{(c^2x^2 + 1)/(c^2x^2)}) - cx - 1 - 2c^2x^2 \log(x) - cx \sqrt{(c^2x^2 + 1)/(c^2x^2)} - 1)/(c^2x^2)$

Sympy [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^2(1+c^2x^2)} dx = \frac{\int \frac{cx \sqrt{1+\frac{1}{c^2x^2}}}{c^2x^5+x^3} dx + \int \frac{1}{c^2x^5+x^3} dx}{c}$$

[In] `integrate((1/c/x+(1+1/c**2/x**2)**(1/2))/x**2/(c**2*x**2+1),x)`

[Out] `(Integral(cx*sqrt(1 + 1/(c**2*x**2)))/(c**2*x**5 + x**3), x) + Integral(1/(c**2*x**5 + x**3), x))/c`

Maxima [F]

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^2(1+c^2x^2)} dx = \int \frac{\sqrt{\frac{1}{c^2x^2} + 1 + \frac{1}{cx}}}{(c^2x^2 + 1)x^2} dx$$

[In] `integrate((1/c/x+(1+1/c^2/x^2)^(1/2))/x^2/(c^2*x^2+1),x, algorithm="maxima")`

[Out] $\frac{1}{2}c \log(c^2x^2 + 1) - c \log(x) - \frac{1}{2}(c^2x^2) + \operatorname{integrate}(\sqrt{c^2x^2 + 1}/(c^3x^5 + cx^3), x)$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 114 vs. $2(50) = 100$.

Time = 0.28 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.90

$$\begin{aligned} \int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^2(1+c^2x^2)} dx &= \frac{1}{2}c \log(c^2x^2 + 1) + \frac{1}{4}(|c \operatorname{sgn}(x) - 2c) \log(\sqrt{c^2x^2 + 1} + 1) \\ &\quad - \frac{1}{4}(|c \operatorname{sgn}(x) + 2c) \log(\sqrt{c^2x^2 + 1} - 1) \\ &\quad - \frac{\sqrt{c^2x^2 + 1}|c \operatorname{sgn}(x) + c}{2(\sqrt{c^2x^2 + 1} + 1)(\sqrt{c^2x^2 + 1} - 1)} \end{aligned}$$

[In] `integrate((1/c/x+(1+1/c^2/x^2)^(1/2))/x^2/(c^2*x^2+1),x, algorithm="giac")`

[Out] $\frac{1}{2}c \log(c^2x^2 + 1) + \frac{1}{4}(\operatorname{abs}(c) \operatorname{sgn}(x) - 2c) \log(\sqrt{c^2x^2 + 1} + 1) - \frac{1}{4}(\operatorname{abs}(c) \operatorname{sgn}(x) + 2c) \log(\sqrt{c^2x^2 + 1} - 1) - \frac{1}{2}(\sqrt{c^2x^2 + 1} \operatorname{abs}(c) \operatorname{sgn}(x) + c)/((\sqrt{c^2x^2 + 1} + 1)(\sqrt{c^2x^2 + 1} - 1))$

Mupad [B] (verification not implemented)

Time = 5.40 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.02

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^2(1+c^2x^2)} dx = \frac{\operatorname{asinh}\left(\frac{\sqrt{\frac{1}{c^2}}}{x}\right)}{2\sqrt{\frac{1}{c^2}}} + \frac{c \ln(-c^2x^2-1)}{2} - c \ln(x) - \frac{\sqrt{\frac{1}{c^2x^2}+1}}{2x} - \frac{1}{2cx^2}$$

```
[In] int(((1/(c^2*x^2) + 1)^(1/2) + 1/(c*x))/(x^2*(c^2*x^2 + 1)),x)
```

```
[Out] asinh((1/c^2)^(1/2)/x)/(2*(1/c^2)^(1/2)) + (c*log(- c^2*x^2 - 1))/2 - c*log(x) - (1/(c^2*x^2) + 1)^(1/2)/(2*x) - 1/(2*c*x^2)
```

3.68 $\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^3(1+c^2x^2)} dx$

Optimal result	407
Rubi [A] (verified)	407
Mathematica [A] (verified)	409
Maple [B] (verified)	409
Fricas [A] (verification not implemented)	409
Sympy [A] (verification not implemented)	410
Maxima [A] (verification not implemented)	410
Giac [A] (verification not implemented)	410
Mupad [B] (verification not implemented)	411

Optimal result

Integrand size = 21, antiderivative size = 61

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^3(1+c^2x^2)} dx = c^2 \sqrt{1 + \frac{1}{c^2x^2}} - \frac{1}{3}c^2 \left(1 + \frac{1}{c^2x^2}\right)^{3/2} - \frac{1}{3cx^3} + \frac{c}{x} + c^2 \arctan(cx)$$

[Out] $-1/3*c^2*(1+1/c^2/x^2)^{(3/2)}-1/3/c/x^3+c/x+c^2*\arctan(c*x)+c^2*(1+1/c^2/x^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {6477, 272, 45, 331, 209}

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^3(1+c^2x^2)} dx = c^2 \arctan(cx) - \frac{1}{3}c^2 \left(\frac{1}{c^2x^2} + 1\right)^{3/2} + c^2 \sqrt{\frac{1}{c^2x^2} + 1} - \frac{1}{3cx^3} + \frac{c}{x}$$

[In] $\text{Int}[E^{\text{ArcCsch}[c*x]}/(x^3*(1 + c^2*x^2)),x]$

[Out] $c^2*\text{Sqrt}[1 + 1/(c^2*x^2)] - (c^2*(1 + 1/(c^2*x^2))^{(3/2)})/3 - 1/(3*c*x^3) + c/x + c^2*\text{ArcTan}[c*x]$

Rule 45

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] := \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, n, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ (!\text{IntegerQ}[n] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{LeQ}[7*m + 4*n + 4, 0]) \ || \ \text{LtQ}[9*m + 5*(n + 1), 0]) \ || \ \text{GtQ}[m + n + 2, 0])$

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 331

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x
)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Dist[b*((m + n*(p + 1)
+ 1)/(a*c^n*(m + 1)), Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a,
b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p,
x]
```

Rule 6477

```
Int[(E^ArcSch[(c_.)*(x_)])*((d_.)*(x_)^(m_.))/((a_) + (b_.)*(x_)^2), x_Sym
bol] := Dist[d^2/(a*c^2), Int[(d*x)^(m - 2)/Sqrt[1 + 1/(c^2*x^2)], x], x] +
Dist[d/c, Int[(d*x)^(m - 1)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, m},
x] && EqQ[b - a*c^2, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\int \frac{1}{\sqrt{1 + \frac{1}{c^2 x^2}} x^5} dx}{c^2} + \frac{\int \frac{1}{x^4(1 + c^2 x^2)} dx}{c} \\
&= -\frac{1}{3cx^3} - \frac{\text{Subst}\left(\int \frac{x}{\sqrt{1 + \frac{x}{c^2}}} dx, x, \frac{1}{x^2}\right)}{2c^2} - c \int \frac{1}{x^2(1 + c^2 x^2)} dx \\
&= -\frac{1}{3cx^3} + \frac{c}{x} - \frac{\text{Subst}\left(\int \left(-\frac{c^2}{\sqrt{1 + \frac{x}{c^2}}} + c^2 \sqrt{1 + \frac{x}{c^2}}\right) dx, x, \frac{1}{x^2}\right)}{2c^2} + c^3 \int \frac{1}{1 + c^2 x^2} dx \\
&= c^2 \sqrt{1 + \frac{1}{c^2 x^2}} - \frac{1}{3} c^2 \left(1 + \frac{1}{c^2 x^2}\right)^{3/2} - \frac{1}{3cx^3} + \frac{c}{x} + c^2 \arctan(cx)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.89

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^3(1+c^2x^2)} dx = -\frac{1}{3cx^3} + \frac{c}{x} + \frac{\sqrt{1+\frac{1}{c^2x^2}}(-1+2c^2x^2)}{3x^2} + c^2 \arctan(cx)$$

[In] Integrate[E^ArcCsch[c*x]/(x^3*(1+c^2*x^2)),x]

[Out] -1/3*1/(c*x^3) + c/x + (Sqrt[1+1/(c^2*x^2)]*(-1+2*c^2*x^2))/(3*x^2) + c^2*ArcTan[c*x]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 196 vs. 2(53) = 106.

Time = 0.87 (sec) , antiderivative size = 197, normalized size of antiderivative = 3.23

method	result
default	$\frac{\sqrt{\frac{c^2x^2+1}{c^2x^2}} c^2 \left(3 \left(\frac{c^2x^2+1}{c^2} \right)^{\frac{3}{2}} c^2 x^2 - 3 \sqrt{\frac{c^2x^2+1}{c^2}} c^2 x^4 - 3 \ln \left(x + \sqrt{\frac{c^2x^2+1}{c^2}} \right) x^3 + 3 \ln \left(x + \sqrt{-\frac{(-c^2x+\sqrt{-c^2})(c^2x+\sqrt{-c^2})}{c^4}} \right) x^3 - \left(\frac{c^2x^2+1}{c^2} \right)^{\frac{3}{2}} \right)}{3x^2 \sqrt{\frac{c^2x^2+1}{c^2}}}$

[In] int((1/c/x+(1+1/c^2/x^2)^(1/2))/x^3/(c^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] 1/3*((c^2*x^2+1)/c^2/x^2)^(1/2)/x^2*c^2*(3*(1/c^2*(c^2*x^2+1))^(3/2)*c^2*x^2-3*(1/c^2*(c^2*x^2+1))^(1/2)*c^2*x^4-3*ln(x+(1/c^2*(c^2*x^2+1))^(1/2))*x^3+3*ln(x+((-c^2*x+(-c^2)^(1/2))*(c^2*x+(-c^2)^(1/2))/c^4)^(1/2))*x^3-(1/c^2*(c^2*x^2+1))^(3/2))/(1/c^2*(c^2*x^2+1))^(1/2)+1/c*(-1/3/x^3+c^2/x+c^3*arctan(c*x))

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.15

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^3(1+c^2x^2)} dx = \frac{3c^3x^3 \arctan(cx) + 2c^3x^3 + 3c^2x^2 + (2c^3x^3 - cx)\sqrt{\frac{c^2x^2+1}{c^2x^2}} - 1}{3cx^3}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))/x^3/(c^2*x^2+1),x, algorithm="fricas")

[Out] 1/3*(3*c^3*x^3*arctan(c*x) + 2*c^3*x^3 + 3*c^2*x^2 + (2*c^3*x^3 - c*x)*sqrt((c^2*x^2 + 1)/(c^2*x^2)) - 1)/(c*x^3)

Sympy [A] (verification not implemented)

Time = 2.26 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.67

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^3(1+c^2x^2)} dx = -\frac{c^3 \operatorname{atan}\left(\frac{1}{x\sqrt{c^2}}\right)}{\sqrt{c^2}} - c \left(\begin{cases} 2c^4 \left(\frac{\left(1+\frac{1}{c^2x^2}\right)^{\frac{3}{2}}}{6c^3} - \frac{\sqrt{1+\frac{1}{c^2x^2}}}{2c^3} \right) & \text{for } \frac{1}{c^2} \neq 0 \\ -\frac{c \log\left(c^2+\frac{1}{x^2}\right)}{2} + \frac{1}{2cx^2} & \text{otherwise} \end{cases} \right) + \frac{c}{x} - \frac{1}{3cx^3}$$

[In] integrate((1/c/x+(1+1/c**2/x**2)**(1/2))/x**3/(c**2*x**2+1),x)

[Out] -c**3*atan(1/(x*sqrt(c**2)))/sqrt(c**2) - c*Piecewise((2*c**4*((1 + 1/(c**2*x**2))**(3/2))/(6*c**3) - sqrt(1 + 1/(c**2*x**2))/(2*c**3)), Ne(c**(-2), 0)), (-c*log(c**2 + x**(-2))/2 + 1/(2*c*x**2), True)) + c/x - 1/(3*c*x**3)

Maxima [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.92

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^3(1+c^2x^2)} dx = c^2 \arctan(cx) + \frac{(2c^2x^2-1)\sqrt{c^2x^2+1}}{3cx^3} + \frac{3c^2x^2-1}{3cx^3}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))/x^3/(c^2*x^2+1),x, algorithm="maxima")

[Out] c^2*arctan(c*x) + 1/3*(2*c^2*x^2 - 1)*sqrt(c^2*x^2 + 1)/(c*x^3) + 1/3*(3*c^2*x^2 - 1)/(c*x^3)

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.34

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^3(1+c^2x^2)} dx = c^2 \arctan(cx) + \frac{4 \left(3 \left(x|c| - \sqrt{c^2x^2+1} \right)^2 - 1 \right) c^2 \operatorname{sgn}(x)}{3 \left(\left(x|c| - \sqrt{c^2x^2+1} \right)^2 - 1 \right)^3} + \frac{3c^2x^2-1}{3cx^3}$$

[In] integrate((1/c/x+(1+1/c^2/x^2)^(1/2))/x^3/(c^2*x^2+1),x, algorithm="giac")

[Out] c^2*arctan(c*x) + 4/3*(3*(x*abs(c) - sqrt(c^2*x^2 + 1))^2 - 1)*c^2*sgn(x)/((x*abs(c) - sqrt(c^2*x^2 + 1))^2 - 1)^3 + 1/3*(3*c^2*x^2 - 1)/(c*x^3)

Mupad [B] (verification not implemented)

Time = 5.20 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.93

$$\int \frac{e^{\operatorname{csch}^{-1}(cx)}}{x^3(1+c^2x^2)} dx = \frac{c + \frac{2c^2x\sqrt{\frac{1}{c^2x^2}+1}}{3}}{x} - \frac{x\sqrt{\frac{1}{c^2x^2}+1}}{3} + \frac{1}{3c} + c^2 \operatorname{atan}(cx)$$

```
[In] int(((1/(c^2*x^2) + 1)^(1/2) + 1/(c*x))/(x^3*(c^2*x^2 + 1)),x)
```

```
[Out] (c + (2*c^2*x*(1/(c^2*x^2) + 1)^(1/2))/3)/x - ((x*(1/(c^2*x^2) + 1)^(1/2))/3 + 1/(3*c))/x^3 + c^2*atan(c*x)
```

$$3.69 \quad \int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx$$

Optimal result	412
Rubi [A] (verified)	412
Mathematica [A] (verified)	415
Maple [F]	415
Fricas [F]	415
Sympy [F]	415
Maxima [F]	416
Giac [F]	416
Mupad [F(-1)]	416

Optimal result

Integrand size = 19, antiderivative size = 61

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \frac{\operatorname{csch}^{-1}(a+bx)^2}{2d} - \frac{\operatorname{csch}^{-1}(a+bx) \log\left(1 - e^{2\operatorname{csch}^{-1}(a+bx)}\right)}{d} - \frac{\operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(a+bx)}\right)}{2d}$$

[Out] 1/2*arccsch(b*x+a)^2/d-arccsch(b*x+a)*ln(1-(1/(b*x+a)+(1+1/(b*x+a)^2)^(1/2))^2)/d-1/2*polylog(2, (1/(b*x+a)+(1+1/(b*x+a)^2)^(1/2))^2)/d

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {6455, 12, 6417, 5775, 3797, 2221, 2317, 2438}

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = -\frac{\operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(a+bx)}\right)}{2d} + \frac{\operatorname{csch}^{-1}(a+bx)^2}{2d} - \frac{\operatorname{csch}^{-1}(a+bx) \log\left(1 - e^{2\operatorname{csch}^{-1}(a+bx)}\right)}{d}$$

[In] Int[ArcCsch[a + b*x]/((a*d)/b + d*x), x]

[Out] ArcCsch[a + b*x]^2/(2*d) - (ArcCsch[a + b*x]*Log[1 - E^(2*ArcCsch[a + b*x])])/d - PolyLog[2, E^(2*ArcCsch[a + b*x])]/(2*d)

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 2221

Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)] / ((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2317

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3797

Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x)))/E^(2*I*k*Pi)], x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]

Rule 5775

Int[((a_) + ArcSinh[(c_)*(x_)])*(b_)^(n_)/(x_), x_Symbol] := Dist[1/b, Subst[Int[x^n*Coth[-a/b + x/b], x], x, a + b*ArcSinh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]

Rule 6417

Int[((a_) + ArcCsch[(c_)*(x_)])*(b_)/(x_), x_Symbol] := -Subst[Int[(a + b*ArcSinh[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]

Rule 6455

Int[((a_) + ArcCsch[(c_) + (d_)*(x_)])*(b_)^(p_)*((e_) + (f_)*(x_))^(m_), x_Symbol] := Dist[1/d, Subst[Int[(f*(x/d))^m*(a + b*ArcCsch[x])^p, x]

, x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] &
& IGtQ[p, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int \frac{b \operatorname{csch}^{-1}(x)}{dx} dx, x, a + bx\right)}{b} \\
 &= \frac{\text{Subst}\left(\int \frac{\operatorname{csch}^{-1}(x)}{x} dx, x, a + bx\right)}{d} \\
 &= -\frac{\text{Subst}\left(\int \frac{\operatorname{arcsinh}(x)}{x} dx, x, \frac{1}{a+bx}\right)}{d} \\
 &= -\frac{\text{Subst}\left(\int x \coth(x) dx, x, \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)\right)}{d} \\
 &= \frac{\operatorname{arcsinh}\left(\frac{1}{a+bx}\right)^2}{2d} + \frac{2\text{Subst}\left(\int \frac{e^{2x} x}{1-e^{2x}} dx, x, \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)\right)}{d} \\
 &= \frac{\operatorname{arcsinh}\left(\frac{1}{a+bx}\right)^2}{2d} - \frac{\operatorname{arcsinh}\left(\frac{1}{a+bx}\right) \log\left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}\right)}{d} \\
 &\quad + \frac{\text{Subst}\left(\int \log(1 - e^{2x}) dx, x, \operatorname{arcsinh}\left(\frac{1}{a+bx}\right)\right)}{d} \\
 &= \frac{\operatorname{arcsinh}\left(\frac{1}{a+bx}\right)^2}{2d} - \frac{\operatorname{arcsinh}\left(\frac{1}{a+bx}\right) \log\left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}\right)}{d} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2\operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}\right)}{2d} \\
 &= \frac{\operatorname{arcsinh}\left(\frac{1}{a+bx}\right)^2}{2d} - \frac{\operatorname{arcsinh}\left(\frac{1}{a+bx}\right) \log\left(1 - e^{2\operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}\right)}{d} - \frac{\operatorname{PolyLog}\left(2, e^{2\operatorname{arcsinh}\left(\frac{1}{a+bx}\right)}\right)}{2d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.87

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \frac{\operatorname{csch}^{-1}(a+bx)^2 - 2\operatorname{csch}^{-1}(a+bx) \log\left(1 - e^{2\operatorname{csch}^{-1}(a+bx)}\right) - \operatorname{PolyLog}\left(2, e^{2\operatorname{csch}^{-1}(a+bx)}\right)}{2d}$$

[In] Integrate[ArcCsch[a + b*x]/((a*d)/b + d*x), x]

[Out] (ArcCsch[a + b*x]^2 - 2*ArcCsch[a + b*x]*Log[1 - E^(2*ArcCsch[a + b*x])] - PolyLog[2, E^(2*ArcCsch[a + b*x])])/(2*d)

Maple [F]

$$\int \frac{\operatorname{arccsch}(bx+a)}{\frac{ad}{b}+dx} dx$$

[In] int(arccsch(b*x+a)/(a*d/b+d*x), x)

[Out] int(arccsch(b*x+a)/(a*d/b+d*x), x)

Fricas [F]

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \int \frac{\operatorname{arcsch}(bx+a)}{dx + \frac{ad}{b}} dx$$

[In] integrate(arccsch(b*x+a)/(a*d/b+d*x), x, algorithm="fricas")

[Out] integral(b*arccsch(b*x + a)/(b*d*x + a*d), x)

Sympy [F]

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \frac{b \int \frac{\operatorname{acsch}(a+bx)}{a+bx} dx}{d}$$

[In] integrate(acsch(b*x+a)/(a*d/b+d*x), x)

[Out] b*Integral(acsch(a + b*x)/(a + b*x), x)/d

Maxima [F]

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \int \frac{\operatorname{arcsch}(bx+a)}{dx+\frac{ad}{b}} dx$$

[In] integrate(arccsch(b*x+a)/(a*d/b+d*x),x, algorithm="maxima")

[Out] -1/4*(2*log(b^2*x^2 + 2*a*b*x + a^2 + 1)*log(b*x + a) + dilog(-b^2*x^2 - 2*a*b*x - a^2))/d - 1/2*(log(b*x + a)^2 - 2*log(b*x + a)*log(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) + 1))/d + integrate((b^2*x + a*b)*log(b*x + a)/(b^2*d*x^2 + 2*a*b*d*x + a^2*d + (b^2*d*x^2 + 2*a*b*d*x + a^2*d + d)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) + d), x)

Giac [F]

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \int \frac{\operatorname{arcsch}(bx+a)}{dx+\frac{ad}{b}} dx$$

[In] integrate(arccsch(b*x+a)/(a*d/b+d*x),x, algorithm="giac")

[Out] integrate(arccsch(b*x + a)/(d*x + a*d/b), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \int \frac{\operatorname{asinh}\left(\frac{1}{a+bx}\right)}{dx+\frac{ad}{b}} dx$$

[In] int(asinh(1/(a + b*x))/(d*x + (a*d)/b),x)

[Out] int(asinh(1/(a + b*x))/(d*x + (a*d)/b), x)

3.70 $\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx$

Optimal result	417
Rubi [A] (verified)	417
Mathematica [B] (verified)	419
Maple [A] (verified)	419
Fricas [B] (verification not implemented)	420
Sympy [F(-1)]	420
Maxima [A] (verification not implemented)	420
Giac [F]	421
Mupad [B] (verification not implemented)	421

Optimal result

Integrand size = 12, antiderivative size = 46

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx = \frac{(a + bx^4) \operatorname{csch}^{-1}(a + bx^4)}{4b} + \frac{\operatorname{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx^4)^2}}\right)}{4b}$$

[Out] $1/4*(b*x^4+a)*\operatorname{arccsch}(b*x^4+a)/b+1/4*\operatorname{arctanh}((1+1/(b*x^4+a)^2)^{(1/2)})/b$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6847, 6449, 379, 272, 65, 213}

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx = \frac{\operatorname{arctanh}\left(\sqrt{\frac{1}{(a+bx^4)^2} + 1}\right)}{4b} + \frac{(a + bx^4) \operatorname{csch}^{-1}(a + bx^4)}{4b}$$

[In] $\operatorname{Int}[x^3*\operatorname{ArcCsch}[a + b*x^4],x]$

[Out] $((a + b*x^4)*\operatorname{ArcCsch}[a + b*x^4])/(4*b) + \operatorname{ArcTanh}[\operatorname{Sqrt}[1 + (a + b*x^4)^{-2}]]/(4*b)$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}], x]] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{LeQ}[-1, n, 0] \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]] \&\& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 213

$\text{Int}[(a_.) + (b_.)*(x_.)^2)^{-1}, x_Symbol] := \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[b, 2])^{-1})*\text{ArcTanh}[\text{Rt}[b, 2]*(x/\text{Rt}[-a, 2])], x] /;$ FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 272

$\text{Int}[(x_.)^{(m_.)}*((a_.) + (b_.)*(x_.)^{(n_.)})^{(p_.)}, x_Symbol] := \text{Dist}[1/n, \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n] - 1)*(a + b*x)^p}, x], x, x^n], x] /;$ FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m+1)/n]]

Rule 379

$\text{Int}[(u_.)^{(m_.)}*((a_.) + (b_.)*(v_.)^{(n_.)})^{(p_.)}, x_Symbol] := \text{Dist}[u^m/(\text{Coefficient}[v, x, 1]*v^m), \text{Subst}[\text{Int}[x^m*(a + b*x^n)^p, x], x, v], x] /;$ FreeQ[{a, b, m, n, p}, x] && LinearPairQ[u, v, x]

Rule 6449

$\text{Int}[\text{ArcCsch}[(c_.) + (d_.)*(x_.)], x_Symbol] := \text{Simp}[(c + d*x)*(\text{ArcCsch}[c + d*x]/d), x] + \text{Int}[1/((c + d*x)*\text{Sqrt}[1 + 1/(c + d*x)^2]), x] /;$ FreeQ[{c, d}, x]

Rule 6847

$\text{Int}[(u_.)*(x_.)^{(m_.)}, x_Symbol] := \text{Dist}[1/(m+1), \text{Subst}[\text{Int}[\text{SubstFor}[x^{(m+1)}, u, x], x, x^{(m+1)}], x] /;$ FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^{(m+1)}, u, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{4} \text{Subst} \left(\int \text{csch}^{-1}(a + bx) dx, x, x^4 \right) \\
 &= \frac{(a + bx^4) \text{csch}^{-1}(a + bx^4)}{4b} + \frac{1}{4} \text{Subst} \left(\int \frac{1}{(a + bx) \sqrt{1 + \frac{1}{(a+bx)^2}}} dx, x, x^4 \right) \\
 &= \frac{(a + bx^4) \text{csch}^{-1}(a + bx^4)}{4b} + \frac{\text{Subst} \left(\int \frac{1}{\sqrt{1 + \frac{1}{x^2}}} dx, x, a + bx^4 \right)}{4b} \\
 &= \frac{(a + bx^4) \text{csch}^{-1}(a + bx^4)}{4b} - \frac{\text{Subst} \left(\int \frac{1}{x\sqrt{1+x}} dx, x, \frac{1}{(a+bx^4)^2} \right)}{8b}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{(a + bx^4) \operatorname{csch}^{-1}(a + bx^4)}{4b} - \frac{\operatorname{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \sqrt{1 + \frac{1}{(a+bx^4)^2}}\right)}{4b} \\
&= \frac{(a + bx^4) \operatorname{csch}^{-1}(a + bx^4)}{4b} + \frac{\operatorname{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx^4)^2}}\right)}{4b}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 95 vs. $2(46) = 92$.

Time = 0.10 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.07

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx = \frac{(a + bx^4) \operatorname{csch}^{-1}(a + bx^4)}{4b} - \frac{\sqrt{1 + (a + bx^4)^2} \log\left(-a - bx^4 + \sqrt{1 + (a + bx^4)^2}\right)}{4b(a + bx^4) \sqrt{1 + \frac{1}{(a+bx^4)^2}}}$$

[In] Integrate[x^3*ArcCsch[a + b*x^4],x]

[Out] ((a + b*x^4)*ArcCsch[a + b*x^4])/(4*b) - (Sqrt[1 + (a + b*x^4)^2]*Log[-a - b*x^4 + Sqrt[1 + (a + b*x^4)^2]])/(4*b*(a + b*x^4)*Sqrt[1 + (a + b*x^4)^(-2)])

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.13

method	result	size
derivativedivides	$\frac{(bx^4+a) \operatorname{arccsch}(bx^4+a) + \ln\left(bx^4+a + (bx^4+a) \sqrt{1 + \frac{1}{(bx^4+a)^2}}\right)}{4b}$	52
default	$\frac{(bx^4+a) \operatorname{arccsch}(bx^4+a) + \ln\left(bx^4+a + (bx^4+a) \sqrt{1 + \frac{1}{(bx^4+a)^2}}\right)}{4b}$	52

[In] int(x^3*arccsch(b*x^4+a),x,method=_RETURNVERBOSE)

[Out] 1/4/b*((b*x^4+a)*arccsch(b*x^4+a)+ln(b*x^4+a+(b*x^4+a)*(1+1/(b*x^4+a)^2)^(1/2)))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 266 vs. 2(40) = 80.

Time = 0.28 (sec) , antiderivative size = 266, normalized size of antiderivative = 5.78

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx$$

$$= \frac{bx^4 \log\left(\frac{(bx^4+a)\sqrt{\frac{b^2x^8+2abx^4+a^2+1}{b^2x^8+2abx^4+a^2}}+1}{bx^4+a}\right) + a \log\left(-bx^4 + (bx^4 + a)\sqrt{\frac{b^2x^8+2abx^4+a^2+1}{b^2x^8+2abx^4+a^2}} - a + 1\right) - a \log\left(-bx^4 + (bx^4 + a)\sqrt{\frac{b^2x^8+2abx^4+a^2+1}{b^2x^8+2abx^4+a^2}} + a - 1\right)}{4b}$$

[In] integrate(x^3*arccsch(b*x^4+a),x, algorithm="fricas")

[Out] 1/4*(b*x^4*log(((b*x^4 + a)*sqrt((b^2*x^8 + 2*a*b*x^4 + a^2 + 1)/(b^2*x^8 + 2*a*b*x^4 + a^2)) + 1)/(b*x^4 + a)) + a*log(-b*x^4 + (b*x^4 + a)*sqrt((b^2*x^8 + 2*a*b*x^4 + a^2 + 1)/(b^2*x^8 + 2*a*b*x^4 + a^2)) - a + 1) - a*log(-b*x^4 + (b*x^4 + a)*sqrt((b^2*x^8 + 2*a*b*x^4 + a^2 + 1)/(b^2*x^8 + 2*a*b*x^4 + a^2)) - a - 1) - log(-b*x^4 + (b*x^4 + a)*sqrt((b^2*x^8 + 2*a*b*x^4 + a^2 + 1)/(b^2*x^8 + 2*a*b*x^4 + a^2)) - a))/b

Sympy [F(-1)]

Timed out.

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx = \text{Timed out}$$

[In] integrate(x**3*acsch(b*x**4+a),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.24

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx$$

$$= \frac{2(bx^4 + a) \operatorname{arcsch}(bx^4 + a) + \log\left(\sqrt{\frac{1}{(bx^4+a)^2} + 1} + 1\right) - \log\left(\sqrt{\frac{1}{(bx^4+a)^2} + 1} - 1\right)}{8b}$$

[In] integrate(x^3*arccsch(b*x^4+a),x, algorithm="maxima")

[Out] 1/8*(2*(b*x^4 + a)*arccsch(b*x^4 + a) + log(sqrt(1/(b*x^4 + a)^2 + 1) + 1) - log(sqrt(1/(b*x^4 + a)^2 + 1) - 1))/b

Giac [F]

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx = \int x^3 \operatorname{arcsch}(bx^4 + a) dx$$

[In] integrate(x^3*arccsch(b*x^4+a),x, algorithm="giac")

[Out] integrate(x^3*arccsch(b*x^4 + a), x)

Mupad [B] (verification not implemented)

Time = 5.75 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.91

$$\int x^3 \operatorname{csch}^{-1}(a + bx^4) dx = \frac{\operatorname{atanh}\left(\sqrt{\frac{1}{(bx^4+a)^2} + 1}\right)}{4b} + \frac{\operatorname{asinh}\left(\frac{1}{bx^4+a}\right) (bx^4 + a)}{4b}$$

[In] int(x^3*asinh(1/(a + b*x^4)),x)

[Out] atanh((1/(a + b*x^4)^2 + 1)^(1/2))/(4*b) + (asinh(1/(a + b*x^4))*(a + b*x^4))/(4*b)

3.71 $\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx$

Optimal result	422
Rubi [A] (verified)	422
Mathematica [B] (verified)	424
Maple [F]	424
Fricas [B] (verification not implemented)	424
Sympy [F(-1)]	425
Maxima [A] (verification not implemented)	425
Giac [F]	425
Mupad [B] (verification not implemented)	426

Optimal result

Integrand size = 14, antiderivative size = 46

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx = \frac{(a + bx^n) \operatorname{csch}^{-1}(a + bx^n)}{bn} + \frac{\operatorname{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx^n)^2}}\right)}{bn}$$

[Out] (a+b*x^n)*arccsch(a+b*x^n)/b/n+arctanh((1+1/(a+b*x^n)^2)^(1/2))/b/n

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {6847, 6449, 379, 272, 65, 213}

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx = \frac{\operatorname{arctanh}\left(\sqrt{\frac{1}{(a+bx^n)^2} + 1}\right)}{bn} + \frac{(a + bx^n) \operatorname{csch}^{-1}(a + bx^n)}{bn}$$

[In] Int[x^(-1 + n)*ArcCsch[a + b*x^n],x]

[Out] ((a + b*x^n)*ArcCsch[a + b*x^n])/(b*n) + ArcTanh[Sqrt[1 + (a + b*x^n)^(-2)]]/(b*n)

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 213

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 272

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 379

Int[(u_)^(m_)*((a_) + (b_)*(v_)^(n_))^(p_), x_Symbol] := Dist[u^m/(Coefficient[v, x, 1]*v^m), Subst[Int[x^m*(a + b*x^n)^p, x], x, v], x] /; FreeQ[{a, b, m, n, p}, x] && LinearPairQ[u, v, x]

Rule 6449

Int[ArcCsch[(c_) + (d_)*(x_)], x_Symbol] := Simp[(c + d*x)*(ArcCsch[c + d*x]/d), x] + Int[1/((c + d*x)*Sqrt[1 + 1/(c + d*x)^2]), x] /; FreeQ[{c, d}, x]

Rule 6847

Int[(u_)*(x_)^(m_), x_Symbol] := Dist[1/(m + 1), Subst[Int[SubstFor[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^(m + 1), u, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int \text{csch}^{-1}(a + bx) dx, x, x^n\right)}{n} \\
 &= \frac{(a + bx^n) \text{csch}^{-1}(a + bx^n)}{bn} + \frac{\text{Subst}\left(\int \frac{1}{(a+bx)\sqrt{1+\frac{1}{(a+bx)^2}}} dx, x, x^n\right)}{n} \\
 &= \frac{(a + bx^n) \text{csch}^{-1}(a + bx^n)}{bn} + \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1+\frac{1}{x^2}}} dx, x, a + bx^n\right)}{bn} \\
 &= \frac{(a + bx^n) \text{csch}^{-1}(a + bx^n)}{bn} - \frac{\text{Subst}\left(\int \frac{1}{x\sqrt{1+x}} dx, x, \frac{1}{(a+bx^n)^2}\right)}{2bn} \\
 &= \frac{(a + bx^n) \text{csch}^{-1}(a + bx^n)}{bn} - \frac{\text{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \sqrt{1+\frac{1}{(a+bx^n)^2}}\right)}{bn}
 \end{aligned}$$

$$= \frac{(a + bx^n) \operatorname{csch}^{-1}(a + bx^n)}{bn} + \frac{\operatorname{arctanh}\left(\sqrt{1 + \frac{1}{(a+bx^n)^2}}\right)}{bn}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 93 vs. $2(46) = 92$.

Time = 0.13 (sec) , antiderivative size = 93, normalized size of antiderivative = 2.02

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx$$

$$= \frac{(a + bx^n)^2 \operatorname{csch}^{-1}(a + bx^n) - \frac{\sqrt{1+(a+bx^n)^2} \log\left(-a-bx^n + \sqrt{1+(a+bx^n)^2}\right)}{\sqrt{1+\frac{1}{(a+bx^n)^2}}}}{bn(a + bx^n)}$$

[In] Integrate[x[^](-1 + n)*ArcCsch[a + b*x[^]n],x]

[Out] ((a + b*x[^]n)[^]2*ArcCsch[a + b*x[^]n] - (Sqrt[1 + (a + b*x[^]n)[^]2]*Log[-a - b*x[^]n + Sqrt[1 + (a + b*x[^]n)[^]2]])/Sqrt[1 + (a + b*x[^]n)[^](-2)]/(b*n*(a + b*x[^]n))

Maple [F]

$$\int x^{-1+n} \operatorname{arccsch}(a + b x^n) dx$$

[In] int(x[^](-1+n)*arccsch(a+b*x[^]n),x)

[Out] int(x[^](-1+n)*arccsch(a+b*x[^]n),x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 334 vs. $2(44) = 88$.

Time = 0.29 (sec) , antiderivative size = 334, normalized size of antiderivative = 7.26

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx$$

$$= \frac{a \log\left(-b \cosh(n \log(x)) - b \sinh(n \log(x)) - a + \sqrt{\frac{2ab+(a^2+b^2+1) \cosh(n \log(x)) - (a^2-b^2+1) \sinh(n \log(x))}{\cosh(n \log(x)) - \sinh(n \log(x))} + 1}\right) - a}{bn(a + bx^n)}$$

[In] integrate(x[^](-1+n)*arccsch(a+b*x[^]n),x, algorithm="fricas")

[Out] (a*log(-b*cosh(n*log(x)) - b*sinh(n*log(x)) - a + sqrt((2*a*b + (a[^]2 + b[^]2 + 1)*cosh(n*log(x)) - (a[^]2 - b[^]2 + 1)*sinh(n*log(x))))/(cosh(n*log(x)) - sin

$$\frac{\operatorname{h}(n \cdot \log(x)) + 1 - a \cdot \log(-b \cdot \cosh(n \cdot \log(x)) - b \cdot \sinh(n \cdot \log(x)) - a + \sqrt{(2ab + (a^2 + b^2 + 1) \cosh(n \cdot \log(x)) - (a^2 - b^2 + 1) \sinh(n \cdot \log(x))) / (\cosh(n \cdot \log(x)) - \sinh(n \cdot \log(x)))}) - 1) + (b \cdot \cosh(n \cdot \log(x)) + b \cdot \sinh(n \cdot \log(x))) \cdot \log\left(\frac{\sqrt{(2ab + (a^2 + b^2 + 1) \cosh(n \cdot \log(x)) - (a^2 - b^2 + 1) \sinh(n \cdot \log(x))) / (\cosh(n \cdot \log(x)) - \sinh(n \cdot \log(x)))} + 1)}{(b \cdot \cosh(n \cdot \log(x)) + b \cdot \sinh(n \cdot \log(x)) + a)}\right) - \log(-b \cdot \cosh(n \cdot \log(x)) - b \cdot \sinh(n \cdot \log(x)) - a + \sqrt{(2ab + (a^2 + b^2 + 1) \cosh(n \cdot \log(x)) - (a^2 - b^2 + 1) \sinh(n \cdot \log(x))) / (\cosh(n \cdot \log(x)) - \sinh(n \cdot \log(x)))})}{(b \cdot n)}}{b \cdot n}$$

Sympy [F(-1)]

Timed out.

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx = \text{Timed out}$$

[In] integrate(x**(-1+n)*acsch(a+b*x**n),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.30

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx = \frac{2(bx^n + a) \operatorname{arcsch}(bx^n + a) + \log\left(\sqrt{\frac{1}{(bx^n + a)^2} + 1} + 1\right) - \log\left(\sqrt{\frac{1}{(bx^n + a)^2} + 1} - 1\right)}{2bn}$$

[In] integrate(x^(-1+n)*arccsch(a+b*x^n),x, algorithm="maxima")

[Out] 1/2*(2*(b*x^n + a)*arccsch(b*x^n + a) + log(sqrt(1/(b*x^n + a)^2 + 1) + 1) - log(sqrt(1/(b*x^n + a)^2 + 1) - 1))/(b*n)

Giac [F]

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx = \int x^{n-1} \operatorname{arcsch}(bx^n + a) dx$$

[In] integrate(x^(-1+n)*arccsch(a+b*x^n),x, algorithm="giac")

[Out] integrate(x^(n - 1)*arccsch(b*x^n + a), x)

Mupad [B] (verification not implemented)

Time = 5.63 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.87

$$\int x^{-1+n} \operatorname{csch}^{-1}(a + bx^n) dx = \frac{\operatorname{atanh}\left(\sqrt{\frac{1}{(a+bx^n)^2} + 1}\right) + \operatorname{asinh}\left(\frac{1}{a+bx^n}\right) (a + bx^n)}{bn}$$

[In] `int(x^(n - 1)*asinh(1/(a + b*x^n)),x)`

[Out] `(atanh((1/(a + b*x^n)^2 + 1)^(1/2)) + asinh(1/(a + b*x^n))*(a + b*x^n))/(b*n)`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 427

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*      Small rewrite of logic in main function to make it*)
(*      match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
```

```

(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCo
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count is
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"}
    ]
  ]
  ,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<>
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

  finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)

```

```

(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```

```

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
          If[ElementaryFunctionQ[Head[expn]],
            Max[3, ExpnType[expn[[1]]],
          If[SpecialFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
          If[HypergeometricFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
          If[AppellFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
          If[Head[expn]===RootSum,
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
          If[Head[expn]===Integrate || Head[expn]===Int,
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
          9]]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,

```

```

    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result, optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);

```

```

#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 ("
```

```

                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_c
    end if
else #result contains complex but optimal is not
    if debug then
        print("result contains complex but optimal is not");
    fi;
    return "C","Result contains complex when optimal does not.";
fi;
else # result do not contain complex
    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well")
    fi;
    if leaf_count_result<=2*leaf_count_optimal then
        if debug then
            print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
    else
        if debug then
            print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of opt
                                convert(leaf_count_result,string)," $ vs. $2(",
                                convert(leaf_count_optimal,string)," )=",convert(2*leaf_coun
    fi;
fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

```



```

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`) or type(expn,'*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else

```

```

9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u), u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#           Port of original Maple grading function by
#           Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#           added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnTy
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+' or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

#main function

```
def grade_antiderivative(result,optimal):
```

```

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

```

```

leaf_count_result = leaf_count(result)
leaf_count_optimal = leaf_count(optimal)

```

```

#print("leaf_count_result=",leaf_count_result)
#print("leaf_count_optimal=",leaf_count_optimal)

```

```

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

```

```
if str(result).find("Integral") != -1:
```

```

    grade = "F"
    grade_annotation = ""

```

```
else:
```

```
    if expnType_result <= expnType_optimal:
```

```
        if result.has(I):
```

```
            if optimal.has(I): #both result and optimal complex
```

```
                if leaf_count_result <= 2*leaf_count_optimal:
```

```

                    grade = "A"
                    grade_annotation = ""

```

```
                else:
```

```
                    grade = "B"
```

```
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
```

```
            else: #result contains complex but optimal is not
```

```
                grade = "C"
```

```
                grade_annotation = "Result contains complex when optimal does not."
```

```
        else: # result do not contain complex, this assumes optimal do not as well
```

```
            if leaf_count_result <= 2*leaf_count_optimal:
```

```

                grade = "A"
                grade_annotation = ""

```

```
            else:
```

```
                grade = "B"
```

```
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
```

```
        else:
```

```
            grade = "C"
```

```
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)
```

```

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#    Albert Rich to use with Sagemath. This is used to
#    grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#    'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#    issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr, Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception,AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```



```

    return 1
  else:
    return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
  else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. " + str(leaf_c

else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_result)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```