

Computer Algebra Independent Integration Tests

Summer 2023 edition

8-Special-functions/207-8.5-Hyperbolic-integral-functions

Nasser M. Abbasi

September 5, 2023

Compiled on September 5, 2023 at 11:57am

Contents

1	Introduction	3
2	detailed summary tables of results	21
3	Listing of integrals	59
4	Appendix	669

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	8
1.4	Performance based on number of rules Rubi used	10
1.5	Performance based on number of steps Rubi used	11
1.6	Solved integrals histogram based on leaf size of result	12
1.7	Solved integrals histogram based on CPU time used	13
1.8	Leaf size vs. CPU time used	14
1.9	list of integrals with no known antiderivative	15
1.10	List of integrals solved by CAS but has no known antiderivative	15
1.11	list of integrals solved by CAS but failed verification	15
1.12	Timing	16
1.13	Verification	16
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [136]. This is test number [207].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (136)	0.00 (0)
Mathematica	100.00 (136)	0.00 (0)
Maple	76.47 (104)	23.53 (32)
Sympy	38.24 (52)	61.76 (84)
Fricas	25.00 (34)	75.00 (102)
Mupad	25.00 (34)	75.00 (102)
Giac	25.00 (34)	75.00 (102)
Maxima	25.00 (34)	75.00 (102)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	75.000	0.000	0.000	25.000
Mathematica	75.000	0.000	0.000	25.000
Maple	50.735	0.000	0.735	48.529
Sympy	10.294	2.941	0.000	86.765
Fricas	0.000	0.000	0.000	100.000
Giac	0.000	0.000	0.000	100.000
Mupad	0.000	0.000	0.000	100.000
Maxima	0.000	0.000	0.000	100.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of

error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Maple	32	100.00	0.00	0.00
Sympy	84	100.00	0.00	0.00
Fricas	102	100.00	0.00	0.00
Mupad	102	0.00	100.00	0.00
Giac	102	100.00	0.00	0.00
Maxima	102	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Rubi	0.15
Fricas	0.24
Maxima	0.25
Giac	0.28
Mathematica	0.54
Maple	0.59
Sympy	1.05
Mupad	4.91

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Fricas	14.24	1.17	14.00	1.17
Mupad	14.24	1.17	14.00	1.17
Giac	14.24	1.17	14.00	1.17
Maxima	14.24	1.17	14.00	1.17
Sympy	34.67	1.17	14.00	1.00
Maple	47.62	0.90	30.00	0.90
Mathematica	63.36	0.93	45.00	0.95
Rubi	80.36	1.00	49.00	1.00

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{9, 14, 15, 16, 17, 22, 25, 29, 30, 31, 40, 46, 48, 58, 62, 65, 68, 77, 82, 83, 84, 85, 90, 93, 97, 98, 99, 108, 114, 116, 126, 130, 133, 136}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
```

```
x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in *Rubi Table file*

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2023
Design v1.0a

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	22
2.2	Detailed conclusion table per each integral for all CAS systems	25
2.3	Detailed conclusion table specific for Rubi results	53

2.1 List of integrals sorted by grade for each CAS

Rubi	22
Mma	22
Maple	23
Fricas	23
Maxima	23
Giac	24
Mupad	24
Sympy	24

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 13, 18, 19, 20, 21, 23, 24, 26, 27, 28, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 47, 49, 50, 51, 52, 53, 54, 55, 56, 57, 59, 60, 61, 63, 64, 66, 67, 69, 70, 71, 72, 73, 74, 75, 76, 78, 79, 80, 81, 86, 87, 88, 89, 91, 92, 94, 95, 96, 100, 101, 102, 103, 104, 105, 106, 107, 109, 110, 111, 112, 113, 115, 117, 118, 119, 120, 121, 122, 123, 124, 125, 127, 128, 129, 131, 132, 134, 135 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 13, 18, 19, 20, 21, 23, 24, 26, 27, 28, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 47, 49, 50, 51, 52, 53, 54, 55, 56, 57, 59, 60, 61, 63, 64, 66, 67, 69, 70, 71, 72, 73, 74, 75, 76, 78, 79, 80, 81, 86, 87, 88, 89, 91, 92, 94, 95, 96, 100, 101, 102, 103, 104, 105, 106, 107, 109, 110, 111, 112, 113, 115, 117, 118, 119, 120, 121, 122, 123, 124, 125, 127, 128, 129, 131, 132, 134, 135 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 13, 18, 19, 20, 21, 27, 28, 35, 41, 42, 43, 44, 45, 49, 50, 51, 52, 53, 54, 55, 56, 57, 59, 60, 61, 70, 71, 72, 73, 75, 76, 78, 79, 80, 81, 86, 87, 88, 89, 95, 96, 103, 109, 110, 111, 112, 113, 117, 118, 119, 120, 121, 122, 123, 124, 125, 127, 128, 129 }

B grade { }

C grade { 1 }

F normal fail { 23, 24, 26, 32, 33, 34, 36, 37, 38, 39, 47, 63, 64, 66, 67, 69, 74, 91, 92, 94, 100, 101, 102, 104, 105, 106, 107, 115, 131, 132, 134, 135 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 13, 18, 19, 20, 21, 23, 24, 26, 27, 28, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 47, 49, 50, 51, 52, 53, 54, 55, 56, 57, 59, 60, 61, 63, 64, 66, 67, 69, 70, 71, 72, 73, 74, 75, 76, 78, 79, 80, 81, 86, 87, 88, 89, 91, 92, 94, 95, 96, 100, 101, 102, 103, 104, 105, 106, 107, 109, 110, 111, 112, 113, 115, 117, 118, 119, 120, 121, 122, 123, 124, 125, 127, 128, 129, 131, 132, 134, 135 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maxima

A grade { }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 13, 18, 19, 20, 21, 23, 24, 26, 27, 28, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 47, 49, 50, 51, 52, 53, 54, 55, 56, 57, 59, 60, 61, 63, 64, 66, 67, 69, 70, 71, 72, 73, 74, 75, 76, 78, 79, 80, 81, 86, 87, 88, 89, 91, 92, 94, 95, 96, 100, 101, 102, 103, 104, 105, 106, 107, 109, 110, 111, 112, 113, 115, 117, 118, 119, 120, 121, 122, 123, 124, 125, 127, 128, 129, 131, 132, 134, 135 }

F(-1) timedout fail { }

F(-2) exception fail { }

Giac**A grade** { }**B grade** { }**C grade** { }**F normal fail** { 1, 2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 13, 18, 19, 20, 21, 23, 24, 26, 27, 28, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 47, 49, 50, 51, 52, 53, 54, 55, 56, 57, 59, 60, 61, 63, 64, 66, 67, 69, 70, 71, 72, 73, 74, 75, 76, 78, 79, 80, 81, 86, 87, 88, 89, 91, 92, 94, 95, 96, 100, 101, 102, 103, 104, 105, 106, 107, 109, 110, 111, 112, 113, 115, 117, 118, 119, 120, 121, 122, 123, 124, 125, 127, 128, 129, 131, 132, 134, 135 }**F(-1) timedout fail** { }**F(-2) exception fail** { }**Mupad****A grade** { }**B grade** { }**C grade** { }**F normal fail** { }**F(-1) timedout fail** { 1, 2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 13, 18, 19, 20, 21, 23, 24, 26, 27, 28, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 47, 49, 50, 51, 52, 53, 54, 55, 56, 57, 59, 60, 61, 63, 64, 66, 67, 69, 70, 71, 72, 73, 74, 75, 76, 78, 79, 80, 81, 86, 87, 88, 89, 91, 92, 94, 95, 96, 100, 101, 102, 103, 104, 105, 106, 107, 109, 110, 111, 112, 113, 115, 117, 118, 119, 120, 121, 122, 123, 124, 125, 127, 128, 129, 131, 132, 134, 135 }**F(-2) exception fail** { }**Sympy****A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 41, 70, 71, 72, 74, 109 }**B grade** { 69, 73, 75, 76 }**C grade** { }**F normal fail** { 10, 11, 12, 13, 18, 19, 20, 21, 23, 24, 26, 27, 28, 32, 33, 34, 35, 36, 37, 38, 39, 42, 43, 44, 45, 47, 49, 50, 51, 52, 53, 54, 55, 56, 57, 59, 60, 61, 63, 64, 66, 67, 78, 79, 80, 81, 86, 87, 88, 89, 91, 92, 94, 95, 96, 100, 101, 102, 103, 104, 105, 106, 107, 110, 111, 112, 113, 115, 117, 118, 119, 120, 121, 122, 123, 124, 125, 127, 128, 129, 131, 132, 134, 135 }**F(-1) timedout fail** { }**F(-2) exception fail** { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	56	37	0	0	42	0	0
N.S.	1	1.00	0.74	0.49	0.00	0.00	0.55	0.00	0.00
time (sec)	N/A	0.052	0.087	0.595	0.000	0.000	0.566	0.000	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	63	53	54	0	0	61	0	0
N.S.	1	1.00	0.84	0.86	0.00	0.00	0.97	0.00	0.00
time (sec)	N/A	0.061	0.026	0.322	0.000	0.000	0.653	0.000	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	49	49	44	42	0	0	46	0	0
N.S.	1	1.00	0.90	0.86	0.00	0.00	0.94	0.00	0.00
time (sec)	N/A	0.038	0.020	0.322	0.000	0.000	0.836	0.000	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	35	30	0	0	29	0	0
N.S.	1	1.00	1.00	0.86	0.00	0.00	0.83	0.00	0.00
time (sec)	N/A	0.019	0.007	0.304	0.000	0.000	0.495	0.000	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	17	0	0	12	0	0
N.S.	1	1.00	1.00	1.06	0.00	0.00	0.75	0.00	0.00
time (sec)	N/A	0.004	0.004	0.210	0.000	0.000	0.508	0.000	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	38	20	0	0	20	0	0
N.S.	1	1.00	1.00	0.53	0.00	0.00	0.53	0.00	0.00
time (sec)	N/A	0.016	0.005	0.381	0.000	0.000	0.340	0.000	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	30	0	0	34	0	0
N.S.	1	1.00	1.00	1.20	0.00	0.00	1.36	0.00	0.00
time (sec)	N/A	0.034	0.009	0.492	0.000	0.000	0.549	0.000	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	46	47	0	0	39	0	0
N.S.	1	1.00	1.00	1.02	0.00	0.00	0.85	0.00	0.00
time (sec)	N/A	0.050	0.015	0.395	0.000	0.000	0.517	0.000	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	10	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	1.00	1.20	1.20
time (sec)	N/A	0.022	0.447	0.221	0.211	0.242	1.289	0.270	4.882

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	149	149	107	120	0	0	0	0	0
N.S.	1	1.00	0.72	0.81	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.188	0.082	0.624	0.000	0.000	0.000	0.000	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	112	112	78	84	0	0	0	0	0
N.S.	1	1.00	0.70	0.75	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.099	0.052	0.688	0.000	0.000	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	74	58	62	0	0	0	0	0
N.S.	1	1.00	0.78	0.84	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.066	0.035	0.573	0.000	0.000	0.000	0.000	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	31	30	0	0	0	0	0
N.S.	1	1.00	1.00	0.97	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.036	0.010	0.477	0.000	0.000	0.000	0.000	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	8	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	0.80	1.20	1.20
time (sec)	N/A	0.017	0.115	0.072	0.214	0.234	1.071	0.271	4.922

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	10	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	1.00	1.20	1.20
time (sec)	N/A	0.018	0.126	0.155	0.218	0.232	0.983	0.276	4.835

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	10	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	1.00	1.20	1.20
time (sec)	N/A	0.018	0.154	0.161	0.216	0.235	1.138	0.277	4.882

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	10	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	1.00	1.20	1.20
time (sec)	N/A	0.217	6.191	0.285	0.210	0.248	0.619	0.266	4.854

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	184	184	94	155	0	0	0	0	0
N.S.	1	1.00	0.51	0.84	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.310	0.144	0.451	0.000	0.000	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	111	111	86	0	0	0	0	0	0
N.S.	1	1.00	0.77	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.273	0.192	0.000	0.000	0.000	0.000	0.000	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	14	14	12	14	14
N.S.	1	1.00	1.17	1.00	1.17	1.17	1.00	1.17	1.17
time (sec)	N/A	0.034	4.578	0.220	0.228	0.271	1.139	0.276	4.901

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	328	328	158	0	0	0	0	0	0
N.S.	1	1.00	0.48	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.166	1.024	0.000	0.000	0.000	0.000	0.000	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	154	95	113	0	0	0	0	0
N.S.	1	1.00	0.62	0.73	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.257	0.217	0.968	0.000	0.000	0.000	0.000	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	48	48	41	43	0	0	0	0	0
N.S.	1	1.00	0.85	0.90	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.054	0.012	0.454	0.000	0.000	0.000	0.000	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	96	96	96	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.154	0.014	0.000	0.000	0.000	0.000	0.000	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	14	14	14	14	14
N.S.	1	1.00	1.17	1.00	1.17	1.17	1.17	1.17	1.17
time (sec)	N/A	0.106	0.194	0.201	0.265	0.242	2.377	0.272	4.846

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	10	9	0	0	7	0	0
N.S.	1	1.00	1.00	0.90	0.00	0.00	0.70	0.00	0.00
time (sec)	N/A	0.014	0.004	0.231	0.000	0.000	0.242	0.000	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	22	0	0	0	0	0
N.S.	1	1.00	1.00	0.88	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.037	0.013	0.370	0.000	0.000	0.000	0.000	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	61	44	46	0	0	0	0	0
N.S.	1	1.00	0.72	0.75	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.067	0.063	0.635	0.000	0.000	0.000	0.000	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	90	90	64	68	0	0	0	0	0
N.S.	1	1.00	0.71	0.76	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.088	0.051	0.768	0.000	0.000	0.000	0.000	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	125	93	104	0	0	0	0	0
N.S.	1	1.00	0.74	0.83	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.136	0.120	0.956	0.000	0.000	0.000	0.000	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	14	14	14	14	14
N.S.	1	1.00	1.17	1.00	1.17	1.17	1.17	1.17	1.17
time (sec)	N/A	0.135	0.309	0.199	0.243	0.261	2.728	0.272	5.158

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	44	44	44	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.070	0.007	0.000	0.000	0.000	0.000	0.000	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	14	14	12	14	14
N.S.	1	1.00	1.17	1.00	1.17	1.17	1.00	1.17	1.17
time (sec)	N/A	0.018	0.184	0.226	0.246	0.243	2.417	0.279	4.992

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	34	36	28	0	0	0	0	0
N.S.	1	1.00	1.06	0.82	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.039	0.016	0.611	0.000	0.000	0.000	0.000	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	62	46	46	0	0	0	0	0
N.S.	1	1.00	0.74	0.74	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.050	0.036	0.803	0.000	0.000	0.000	0.000	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	72	76	0	0	0	0	0
N.S.	1	1.00	0.73	0.78	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.086	0.064	1.032	0.000	0.000	0.000	0.000	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	128	128	94	104	0	0	0	0	0
N.S.	1	1.00	0.73	0.81	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.124	0.069	1.340	0.000	0.000	0.000	0.000	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	25	24	0	0	0	0	0
N.S.	1	1.00	0.86	0.83	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.042	0.022	1.259	0.000	0.000	0.000	0.000	0.000

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	25	24	0	0	0	0	0
N.S.	1	1.00	0.86	0.83	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.042	0.022	1.346	0.000	0.000	0.000	0.000	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	186	186	123	174	0	0	0	0	0
N.S.	1	1.00	0.66	0.94	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.481	0.223	1.618	0.000	0.000	0.000	0.000	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	97	73	84	0	0	0	0	0
N.S.	1	1.00	0.75	0.87	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.237	0.127	1.053	0.000	0.000	0.000	0.000	0.000

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	32	30	0	0	0	0	0
N.S.	1	1.00	0.97	0.91	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.046	0.014	0.485	0.000	0.000	0.000	0.000	0.000

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	18	18	15	18	18
N.S.	1	1.00	1.12	1.00	1.12	1.12	0.94	1.12	1.12
time (sec)	N/A	0.093	0.412	0.269	0.320	0.247	1.107	0.287	5.039

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	153	153	137	0	0	0	0	0	0
N.S.	1	1.00	0.90	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.210	0.684	0.000	0.000	0.000	0.000	0.000	0.000

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	18	18	15	18	18
N.S.	1	1.00	1.12	1.00	1.12	1.12	0.94	1.12	1.12
time (sec)	N/A	0.100	2.058	0.404	0.320	0.243	0.900	0.290	5.065

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	371	371	313	0	0	0	0	0	0
N.S.	1	1.00	0.84	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.963	3.135	0.000	0.000	0.000	0.000	0.000	0.000

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	153	153	137	0	0	0	0	0	0
N.S.	1	1.00	0.90	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.192	0.728	0.000	0.000	0.000	0.000	0.000	0.000

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	18	18	15	18	18
N.S.	1	1.00	1.12	1.00	1.12	1.12	0.94	1.12	1.12
time (sec)	N/A	0.077	1.825	0.426	0.322	0.243	0.884	0.290	4.871

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	B	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	76	76	66	0	0	0	695	0	0
N.S.	1	1.00	0.87	0.00	0.00	0.00	9.14	0.00	0.00
time (sec)	N/A	0.057	0.058	0.000	0.000	0.000	1.269	0.000	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	63	53	54	0	0	85	0	0
N.S.	1	1.00	0.84	0.86	0.00	0.00	1.35	0.00	0.00
time (sec)	N/A	0.059	0.026	0.382	0.000	0.000	1.663	0.000	0.000

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	49	49	44	42	0	0	70	0	0
N.S.	1	1.00	0.90	0.86	0.00	0.00	1.43	0.00	0.00
time (sec)	N/A	0.040	0.019	0.415	0.000	0.000	1.499	0.000	0.000

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	35	30	0	0	53	0	0
N.S.	1	1.00	1.00	0.86	0.00	0.00	1.51	0.00	0.00
time (sec)	N/A	0.019	0.007	0.335	0.000	0.000	0.996	0.000	0.000

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	17	0	0	31	0	0
N.S.	1	1.00	1.00	1.06	0.00	0.00	1.94	0.00	0.00
time (sec)	N/A	0.004	0.004	0.243	0.000	0.000	0.992	0.000	0.000

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	A	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	52	52	52	0	0	0	42	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.81	0.00	0.00
time (sec)	N/A	0.017	0.005	0.000	0.000	0.000	0.917	0.000	0.000

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	30	0	0	39	0	0
N.S.	1	1.00	1.00	1.20	0.00	0.00	1.56	0.00	0.00
time (sec)	N/A	0.034	0.009	0.423	0.000	0.000	0.602	0.000	0.000

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	46	47	0	0	87	0	0
N.S.	1	1.00	1.00	1.02	0.00	0.00	1.89	0.00	0.00
time (sec)	N/A	0.076	0.010	0.382	0.000	0.000	2.414	0.000	0.000

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	10	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	1.00	1.20	1.20
time (sec)	N/A	0.026	0.441	0.228	0.205	0.268	0.789	0.282	4.838

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	164	164	107	120	0	0	0	0	0
N.S.	1	1.00	0.65	0.73	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.168	0.071	0.567	0.000	0.000	0.000	0.000	0.000

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	112	112	78	84	0	0	0	0	0
N.S.	1	1.00	0.70	0.75	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.106	0.054	0.615	0.000	0.000	0.000	0.000	0.000

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	74	57	62	0	0	0	0	0
N.S.	1	1.00	0.77	0.84	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.074	0.033	0.539	0.000	0.000	0.000	0.000	0.000

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	31	30	0	0	0	0	0
N.S.	1	1.00	1.00	0.97	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.038	0.010	0.160	0.000	0.000	0.000	0.000	0.000

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	8	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	0.80	1.20	1.20
time (sec)	N/A	0.015	0.109	0.072	0.215	0.255	0.503	0.291	4.874

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	10	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	1.00	1.20	1.20
time (sec)	N/A	0.017	0.129	0.161	0.218	0.234	0.418	0.292	4.808

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	10	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	1.00	1.20	1.20
time (sec)	N/A	0.017	0.142	0.194	0.216	0.244	0.456	0.287	4.900

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	10	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	1.00	1.20	1.20
time (sec)	N/A	0.199	6.300	0.280	0.212	0.243	0.598	0.282	4.746

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	184	184	94	155	0	0	0	0	0
N.S.	1	1.00	0.51	0.84	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.290	0.134	0.487	0.000	0.000	0.000	0.000	0.000

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	118	118	64	100	0	0	0	0	0
N.S.	1	1.00	0.54	0.85	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.212	0.128	0.477	0.000	0.000	0.000	0.000	0.000

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	71	71	47	58	0	0	0	0	0
N.S.	1	1.00	0.66	0.82	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.163	0.070	0.474	0.000	0.000	0.000	0.000	0.000

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	42	26	0	0	0	0	0
N.S.	1	1.00	1.56	0.96	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.005	0.025	0.418	0.000	0.000	0.000	0.000	0.000

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	8	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	0.80	1.20	1.20
time (sec)	N/A	0.021	0.196	0.280	0.200	0.225	0.377	0.288	4.738

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	46	46	39	0	0	0	0	0	0
N.S.	1	1.00	0.85	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.164	0.073	0.000	0.000	0.000	0.000	0.000	0.000

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	111	111	80	0	0	0	0	0	0
N.S.	1	1.00	0.72	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.250	0.216	0.000	0.000	0.000	0.000	0.000	0.000

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	14	14	12	14	14
N.S.	1	1.00	1.17	1.00	1.17	1.17	1.00	1.17	1.17
time (sec)	N/A	0.040	2.415	0.224	0.213	0.245	1.253	0.296	4.788

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	327	327	158	0	0	0	0	0	0
N.S.	1	1.00	0.48	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.986	0.793	0.000	0.000	0.000	0.000	0.000	0.000

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	154	95	113	0	0	0	0	0
N.S.	1	1.00	0.62	0.73	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.249	0.182	0.924	0.000	0.000	0.000	0.000	0.000

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	48	48	41	43	0	0	0	0	0
N.S.	1	1.00	0.85	0.90	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.051	0.008	0.241	0.000	0.000	0.000	0.000	0.000

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	14	14	10	14	14
N.S.	1	1.00	1.17	1.00	1.17	1.17	0.83	1.17	1.17
time (sec)	N/A	0.026	0.358	0.113	0.208	0.244	0.493	0.280	4.802

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	14	14	12	14	14
N.S.	1	1.00	1.17	1.00	1.17	1.17	1.00	1.17	1.17
time (sec)	N/A	0.028	0.770	0.212	0.213	0.230	0.318	0.286	4.898

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	14	14	12	14	14
N.S.	1	1.00	1.17	1.00	1.17	1.17	1.00	1.17	1.17
time (sec)	N/A	0.029	0.996	0.236	0.223	0.230	0.351	0.274	4.871

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	128	128	97	0	0	0	0	0	0
N.S.	1	1.00	0.76	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.197	1.125	0.000	0.000	0.000	0.000	0.000	0.000

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	128	128	97	0	0	0	0	0	0
N.S.	1	1.00	0.76	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.182	1.094	0.000	0.000	0.000	0.000	0.000	0.000

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	119	119	93	0	0	0	0	0	0
N.S.	1	1.00	0.78	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.170	1.048	0.000	0.000	0.000	0.000	0.000	0.000

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	55	55	96	56	0	0	0	0	0
N.S.	1	1.00	1.75	1.02	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.031	0.057	1.509	0.000	0.000	0.000	0.000	0.000

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	122	122	144	0	0	0	0	0	0
N.S.	1	1.00	1.18	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.182	1.256	0.000	0.000	0.000	0.000	0.000	0.000

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	130	130	146	0	0	0	0	0	0
N.S.	1	1.00	1.12	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.185	1.261	0.000	0.000	0.000	0.000	0.000	0.000

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	167	167	119	0	0	0	0	0	0
N.S.	1	1.00	0.71	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.222	1.370	0.000	0.000	0.000	0.000	0.000	0.000

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	96	96	96	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.155	0.012	0.000	0.000	0.000	0.000	0.000	0.000

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	14	14	14	14	14
N.S.	1	1.00	1.17	1.00	1.17	1.17	1.17	1.17	1.17
time (sec)	N/A	0.113	0.169	0.211	0.261	0.235	3.114	0.279	4.639

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	10	9	0	0	7	0	0
N.S.	1	1.00	1.00	0.90	0.00	0.00	0.70	0.00	0.00
time (sec)	N/A	0.013	0.004	0.251	0.000	0.000	0.235	0.000	0.000

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	22	0	0	0	0	0
N.S.	1	1.00	1.00	0.88	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.032	0.013	0.349	0.000	0.000	0.000	0.000	0.000

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	61	46	46	0	0	0	0	0
N.S.	1	1.00	0.75	0.75	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.089	0.035	0.714	0.000	0.000	0.000	0.000	0.000

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	90	90	64	68	0	0	0	0	0
N.S.	1	1.00	0.71	0.76	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.086	0.053	0.924	0.000	0.000	0.000	0.000	0.000

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	142	142	94	103	0	0	0	0	0
N.S.	1	1.00	0.66	0.73	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.135	0.074	1.246	0.000	0.000	0.000	0.000	0.000

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	14	14	14	14	14
N.S.	1	1.00	1.17	1.00	1.17	1.17	1.17	1.17	1.17
time (sec)	N/A	0.148	0.335	0.207	0.265	0.234	3.645	0.335	4.874

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	44	44	44	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.070	0.008	0.000	0.000	0.000	0.000	0.000	0.000

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	14	14	12	14	14
N.S.	1	1.00	1.17	1.00	1.17	1.17	1.00	1.17	1.17
time (sec)	N/A	0.021	0.191	0.230	0.267	0.234	3.866	0.338	4.881

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	34	36	28	0	0	0	0	0
N.S.	1	1.00	1.06	0.82	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.040	0.014	0.614	0.000	0.000	0.000	0.000	0.000

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	62	44	46	0	0	0	0	0
N.S.	1	1.00	0.71	0.74	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.046	0.053	0.714	0.000	0.000	0.000	0.000	0.000

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	109	109	72	78	0	0	0	0	0
N.S.	1	1.00	0.66	0.72	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.087	0.069	0.802	0.000	0.000	0.000	0.000	0.000

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	146	146	94	104	0	0	0	0	0
N.S.	1	1.00	0.64	0.71	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.134	0.070	1.031	0.000	0.000	0.000	0.000	0.000

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	27	24	0	0	0	0	0
N.S.	1	1.00	0.93	0.83	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.046	0.021	1.117	0.000	0.000	0.000	0.000	0.000

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	27	24	0	0	0	0	0
N.S.	1	1.00	0.93	0.83	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.044	0.018	1.078	0.000	0.000	0.000	0.000	0.000

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	220	220	134	198	0	0	0	0	0
N.S.	1	1.00	0.61	0.90	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.587	0.271	1.743	0.000	0.000	0.000	0.000	0.000

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	109	109	78	97	0	0	0	0	0
N.S.	1	1.00	0.72	0.89	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.227	0.128	1.103	0.000	0.000	0.000	0.000	0.000

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	45	38	0	0	0	0	0
N.S.	1	1.00	0.98	0.83	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.056	0.019	0.814	0.000	0.000	0.000	0.000	0.000

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	18	18	15	18	18
N.S.	1	1.00	1.12	1.00	1.12	1.12	0.94	1.12	1.12
time (sec)	N/A	0.104	0.495	0.263	0.314	0.241	1.059	0.298	5.090

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	186	186	123	175	0	0	0	0	0
N.S.	1	1.00	0.66	0.94	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.431	0.191	2.546	0.000	0.000	0.000	0.000	0.000

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	97	73	84	0	0	0	0	0
N.S.	1	1.00	0.75	0.87	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.199	0.120	1.471	0.000	0.000	0.000	0.000	0.000

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	32	30	0	0	0	0	0
N.S.	1	1.00	0.97	0.91	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.046	0.013	0.691	0.000	0.000	0.000	0.000	0.000

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	18	18	15	18	18
N.S.	1	1.00	1.12	1.00	1.12	1.12	0.94	1.12	1.12
time (sec)	N/A	0.063	0.384	0.264	0.322	0.239	1.125	0.291	5.145

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	371	371	322	0	0	0	0	0	0
N.S.	1	1.00	0.87	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.832	2.398	0.000	0.000	0.000	0.000	0.000	0.000

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	153	153	107	0	0	0	0	0	0
N.S.	1	1.00	0.70	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.192	0.663	0.000	0.000	0.000	0.000	0.000	0.000

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	18	18	15	18	18
N.S.	1	1.00	1.12	1.00	1.12	1.12	0.94	1.12	1.12
time (sec)	N/A	0.112	2.945	0.406	0.313	0.233	0.910	0.302	5.248

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	371	371	270	0	0	0	0	0	0
N.S.	1	1.00	0.73	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.642	1.969	0.000	0.000	0.000	0.000	0.000	0.000

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	153	153	137	0	0	0	0	0	0
N.S.	1	1.00	0.90	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.209	0.375	0.000	0.000	0.000	0.000	0.000	0.000

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	18	18	15	18	18
N.S.	1	1.00	1.12	1.00	1.12	1.12	0.94	1.12	1.12
time (sec)	N/A	0.071	2.725	0.440	0.312	0.239	0.899	0.296	5.137

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [26] had the largest ratio of [1.5829999999999996]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	5	4	1.00	8	0.500
2	A	6	4	1.00	8	0.500
3	A	5	4	1.00	8	0.500
4	A	4	4	1.00	6	0.667
5	A	1	1	1.00	4	0.250
6	A	1	1	1.00	8	0.125
7	A	4	4	1.00	8	0.500
8	A	5	4	1.00	8	0.500
9	N/A	0	0	1.00	10	0.000
10	A	19	11	1.00	10	1.100
11	A	15	10	1.00	10	1.000
12	A	10	8	1.00	8	1.000
13	A	6	5	1.00	6	0.833
14	N/A	0	0	1.00	10	0.000
15	N/A	0	0	1.00	10	0.000
16	N/A	0	0	1.00	10	0.000
17	N/A	0	0	1.00	10	0.000
18	A	14	6	1.00	10	0.600
19	A	10	6	1.00	10	0.600
20	A	7	6	1.00	8	0.750
21	A	1	1	1.00	6	0.167
22	N/A	0	0	1.00	10	0.000
23	A	7	5	1.00	10	0.500

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
24	A	11	6	1.00	10	0.600
25	N/A	0	0	1.00	12	0.000
26	A	39	19	1.00	12	1.583
27	A	17	14	1.00	10	1.400
28	A	5	5	1.00	8	0.625
29	N/A	0	0	1.00	12	0.000
30	N/A	0	0	1.00	12	0.000
31	N/A	0	0	1.00	12	0.000
32	A	7	5	1.00	17	0.294
33	A	7	5	1.00	15	0.333
34	A	7	5	1.00	13	0.385
35	A	3	1	1.00	17	0.059
36	A	7	5	1.00	17	0.294
37	A	7	5	1.00	17	0.294
38	A	7	5	1.00	19	0.263
39	A	14	10	1.00	12	0.833
40	N/A	0	0	1.00	12	0.000
41	A	1	1	1.00	12	0.083
42	A	5	4	1.00	9	0.444
43	A	9	7	1.00	10	0.700
44	A	14	9	1.00	12	0.750
45	A	18	10	1.00	12	0.833
46	N/A	0	0	1.00	12	0.000
47	A	7	6	1.00	12	0.500
48	N/A	0	0	1.00	12	0.000
49	A	5	4	1.00	9	0.444
50	A	9	7	1.00	10	0.700
51	A	13	9	1.00	12	0.750
52	A	20	11	1.00	12	0.917
53	A	6	4	1.00	9	0.444
54	A	6	4	1.00	9	0.444
55	A	21	16	1.00	16	1.000
56	A	11	9	1.00	14	0.643
57	A	4	4	1.00	13	0.308

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
58	N/A	0	0	1.00	16	0.000
59	A	21	14	1.00	16	0.875
60	A	12	10	1.00	14	0.714
61	A	4	3	1.00	13	0.231
62	N/A	0	0	1.00	16	0.000
63	A	24	10	1.00	14	0.714
64	A	9	5	1.00	13	0.385
65	N/A	0	0	1.00	16	0.000
66	A	24	9	1.00	14	0.643
67	A	9	5	1.00	13	0.385
68	N/A	0	0	1.00	16	0.000
69	A	5	4	1.00	8	0.500
70	A	6	4	1.00	8	0.500
71	A	5	4	1.00	8	0.500
72	A	4	4	1.00	6	0.667
73	A	1	1	1.00	4	0.250
74	A	1	1	1.00	8	0.125
75	A	4	4	1.00	8	0.500
76	A	5	4	1.00	8	0.500
77	N/A	0	0	1.00	10	0.000
78	A	19	11	1.00	10	1.100
79	A	15	10	1.00	10	1.000
80	A	10	8	1.00	8	1.000
81	A	6	5	1.00	6	0.833
82	N/A	0	0	1.00	10	0.000
83	N/A	0	0	1.00	10	0.000
84	N/A	0	0	1.00	10	0.000
85	N/A	0	0	1.00	10	0.000
86	A	14	6	1.00	10	0.600
87	A	10	6	1.00	10	0.600
88	A	7	6	1.00	8	0.750
89	A	1	1	1.00	6	0.167
90	N/A	0	0	1.00	10	0.000
91	A	7	5	1.00	10	0.500

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
92	A	11	6	1.00	10	0.600
93	N/A	0	0	1.00	12	0.000
94	A	39	19	1.00	12	1.583
95	A	17	14	1.00	10	1.400
96	A	5	5	1.00	8	0.625
97	N/A	0	0	1.00	12	0.000
98	N/A	0	0	1.00	12	0.000
99	N/A	0	0	1.00	12	0.000
100	A	7	5	1.00	17	0.294
101	A	7	5	1.00	15	0.333
102	A	7	5	1.00	13	0.385
103	A	3	1	1.00	17	0.059
104	A	7	5	1.00	17	0.294
105	A	7	5	1.00	17	0.294
106	A	7	5	1.00	19	0.263
107	A	14	10	1.00	12	0.833
108	N/A	0	0	1.00	12	0.000
109	A	1	1	1.00	12	0.083
110	A	5	4	1.00	9	0.444
111	A	9	7	1.00	10	0.700
112	A	14	9	1.00	12	0.750
113	A	18	10	1.00	12	0.833
114	N/A	0	0	1.00	12	0.000
115	A	7	6	1.00	12	0.500
116	N/A	0	0	1.00	12	0.000
117	A	5	4	1.00	9	0.444
118	A	9	7	1.00	10	0.700
119	A	13	9	1.00	12	0.750
120	A	20	11	1.00	12	0.917
121	A	6	4	1.00	9	0.444
122	A	6	4	1.00	9	0.444
123	A	21	14	1.00	16	0.875
124	A	12	10	1.00	14	0.714
125	A	4	3	1.00	13	0.231

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
126	N/A	0	0	1.00	16	0.000
127	A	21	16	1.00	16	1.000
128	A	11	9	1.00	14	0.643
129	A	4	4	1.00	13	0.308
130	N/A	0	0	1.00	16	0.000
131	A	24	9	1.00	14	0.643
132	A	9	5	1.00	13	0.385
133	N/A	0	0	1.00	16	0.000
134	A	24	10	1.00	14	0.714
135	A	9	5	1.00	13	0.385
136	N/A	0	0	1.00	16	0.000

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^m \text{Shi}(bx) dx$	63
3.2	$\int x^3 \text{Shi}(bx) dx$	67
3.3	$\int x^2 \text{Shi}(bx) dx$	71
3.4	$\int x \text{Shi}(bx) dx$	75
3.5	$\int \text{Shi}(bx) dx$	79
3.6	$\int \frac{\text{Shi}(bx)}{x} dx$	82
3.7	$\int \frac{\text{Shi}(bx)}{x^2} dx$	85
3.8	$\int \frac{\text{Shi}(bx)}{x^3} dx$	89
3.9	$\int x^m \text{Shi}(bx)^2 dx$	93
3.10	$\int x^3 \text{Shi}(bx)^2 dx$	96
3.11	$\int x^2 \text{Shi}(bx)^2 dx$	102
3.12	$\int x \text{Shi}(bx)^2 dx$	108
3.13	$\int \text{Shi}(bx)^2 dx$	113
3.14	$\int \frac{\text{Shi}(bx)^2}{x} dx$	117
3.15	$\int \frac{\text{Shi}(bx)^2}{x^2} dx$	120
3.16	$\int \frac{\text{Shi}(bx)^2}{x^3} dx$	123
3.17	$\int x^m \text{Shi}(a + bx) dx$	126
3.18	$\int x^3 \text{Shi}(a + bx) dx$	129
3.19	$\int x^2 \text{Shi}(a + bx) dx$	134
3.20	$\int x \text{Shi}(a + bx) dx$	139
3.21	$\int \text{Shi}(a + bx) dx$	143
3.22	$\int \frac{\text{Shi}(a+bx)}{x} dx$	146
3.23	$\int \frac{\text{Shi}(a+bx)}{x^2} dx$	149
3.24	$\int \frac{\text{Shi}(a+bx)}{x^3} dx$	153
3.25	$\int x^m \text{Shi}(a + bx)^2 dx$	158

3.26	$\int x^2 \text{Shi}(a + bx)^2 dx$	161
3.27	$\int x \text{Shi}(a + bx)^2 dx$	170
3.28	$\int \text{Shi}(a + bx)^2 dx$	177
3.29	$\int \frac{\text{Shi}(a+bx)^2}{x} dx$	181
3.30	$\int \frac{\text{Shi}(a+bx)^2}{x^2} dx$	184
3.31	$\int \frac{\text{Shi}(a+bx)^2}{x^3} dx$	187
3.32	$\int x^2 \text{Shi}(d(a + b \log(cx^n))) dx$	190
3.33	$\int x \text{Shi}(d(a + b \log(cx^n))) dx$	195
3.34	$\int \text{Shi}(d(a + b \log(cx^n))) dx$	200
3.35	$\int \frac{\text{Shi}(d(a+b \log(cx^n)))}{x} dx$	205
3.36	$\int \frac{\text{Shi}(d(a+b \log(cx^n)))}{x^2} dx$	209
3.37	$\int \frac{\text{Shi}(d(a+b \log(cx^n)))}{x^3} dx$	214
3.38	$\int (ex)^m \text{Shi}(d(a + b \log(cx^n))) dx$	219
3.39	$\int \frac{\sinh(bx) \text{Shi}(bx)}{x^3} dx$	224
3.40	$\int \frac{\sinh(bx) \text{Shi}(bx)}{x^2} dx$	229
3.41	$\int \frac{\sinh(bx) \text{Shi}(bx)}{x} dx$	233
3.42	$\int \sinh(bx) \text{Shi}(bx) dx$	236
3.43	$\int x \sinh(bx) \text{Shi}(bx) dx$	240
3.44	$\int x^2 \sinh(bx) \text{Shi}(bx) dx$	245
3.45	$\int x^3 \sinh(bx) \text{Shi}(bx) dx$	250
3.46	$\int \frac{\cosh(bx) \text{Shi}(bx)}{x^3} dx$	256
3.47	$\int \frac{\cosh(bx) \text{Shi}(bx)}{x^2} dx$	260
3.48	$\int \frac{\cosh(bx) \text{Shi}(bx)}{x} dx$	264
3.49	$\int \cosh(bx) \text{Shi}(bx) dx$	267
3.50	$\int x \cosh(bx) \text{Shi}(bx) dx$	271
3.51	$\int x^2 \cosh(bx) \text{Shi}(bx) dx$	276
3.52	$\int x^3 \cosh(bx) \text{Shi}(bx) dx$	281
3.53	$\int \sinh(5x) \text{Shi}(2x) dx$	287
3.54	$\int \cosh(5x) \text{Shi}(2x) dx$	291
3.55	$\int x^2 \sinh(a + bx) \text{Shi}(a + bx) dx$	295
3.56	$\int x \sinh(a + bx) \text{Shi}(a + bx) dx$	302
3.57	$\int \sinh(a + bx) \text{Shi}(a + bx) dx$	307
3.58	$\int \frac{\sinh(a+bx) \text{Shi}(a+bx)}{x} dx$	311
3.59	$\int x^2 \cosh(a + bx) \text{Shi}(a + bx) dx$	314
3.60	$\int x \cosh(a + bx) \text{Shi}(a + bx) dx$	321
3.61	$\int \cosh(a + bx) \text{Shi}(a + bx) dx$	326
3.62	$\int \frac{\cosh(a+bx) \text{Shi}(a+bx)}{x} dx$	330
3.63	$\int x \sinh(a + bx) \text{Shi}(c + dx) dx$	333
3.64	$\int \sinh(a + bx) \text{Shi}(c + dx) dx$	341
3.65	$\int \frac{\sinh(a+bx) \text{Shi}(c+dx)}{x} dx$	346

3.66	$\int x \cosh(a + bx) \operatorname{Shi}(c + dx) dx$	349
3.67	$\int \cosh(a + bx) \operatorname{Shi}(c + dx) dx$	356
3.68	$\int \frac{\cosh(a+bx) \operatorname{Shi}(c+dx)}{x} dx$	361
3.69	$\int x^m \operatorname{Chi}(bx) dx$	364
3.70	$\int x^3 \operatorname{Chi}(bx) dx$	369
3.71	$\int x^2 \operatorname{Chi}(bx) dx$	373
3.72	$\int x \operatorname{Chi}(bx) dx$	377
3.73	$\int \operatorname{Chi}(bx) dx$	381
3.74	$\int \frac{\operatorname{Chi}(bx)}{x} dx$	384
3.75	$\int \frac{\operatorname{Chi}(bx)}{x^2} dx$	387
3.76	$\int \frac{\operatorname{Chi}(bx)}{x^3} dx$	391
3.77	$\int x^m \operatorname{Chi}(bx)^2 dx$	395
3.78	$\int x^3 \operatorname{Chi}(bx)^2 dx$	398
3.79	$\int x^2 \operatorname{Chi}(bx)^2 dx$	404
3.80	$\int x \operatorname{Chi}(bx)^2 dx$	410
3.81	$\int \operatorname{Chi}(bx)^2 dx$	415
3.82	$\int \frac{\operatorname{Chi}(bx)^2}{x} dx$	419
3.83	$\int \frac{\operatorname{Chi}(bx)^2}{x^2} dx$	422
3.84	$\int \frac{\operatorname{Chi}(bx)^2}{x^3} dx$	425
3.85	$\int x^m \operatorname{Chi}(a + bx) dx$	428
3.86	$\int x^3 \operatorname{Chi}(a + bx) dx$	431
3.87	$\int x^2 \operatorname{Chi}(a + bx) dx$	436
3.88	$\int x \operatorname{Chi}(a + bx) dx$	441
3.89	$\int \operatorname{Chi}(a + bx) dx$	445
3.90	$\int \frac{\operatorname{Chi}(a+bx)}{x} dx$	448
3.91	$\int \frac{\operatorname{Chi}(a+bx)}{x^2} dx$	451
3.92	$\int \frac{\operatorname{Chi}(a+bx)}{x^3} dx$	455
3.93	$\int x^m \operatorname{Chi}(a + bx)^2 dx$	460
3.94	$\int x^2 \operatorname{Chi}(a + bx)^2 dx$	463
3.95	$\int x \operatorname{Chi}(a + bx)^2 dx$	473
3.96	$\int \operatorname{Chi}(a + bx)^2 dx$	480
3.97	$\int \frac{\operatorname{Chi}(a+bx)^2}{x} dx$	484
3.98	$\int \frac{\operatorname{Chi}(a+bx)^2}{x^2} dx$	487
3.99	$\int \frac{\operatorname{Chi}(a+bx)^2}{x^3} dx$	490
3.100	$\int x^2 \operatorname{Chi}(d(a + b \log(cx^n))) dx$	493
3.101	$\int x \operatorname{Chi}(d(a + b \log(cx^n))) dx$	498
3.102	$\int \operatorname{Chi}(d(a + b \log(cx^n))) dx$	503
3.103	$\int \frac{\operatorname{Chi}(d(a+b \log(cx^n)))}{x} dx$	508
3.104	$\int \frac{\operatorname{Chi}(d(a+b \log(cx^n)))}{x^2} dx$	512
3.105	$\int \frac{\operatorname{Chi}(d(a+b \log(cx^n)))}{x^3} dx$	517

3.106	$\int (ex)^m \text{Chi}(d(a + b \log(cx^n))) dx$	522
3.107	$\int \frac{\cosh(bx) \text{Chi}(bx)}{x^3} dx$	527
3.108	$\int \frac{\cosh(bx) \text{Chi}(bx)}{x^2} dx$	532
3.109	$\int \frac{\cosh(bx) \text{Chi}(bx)}{x} dx$	536
3.110	$\int \cosh(bx) \text{Chi}(bx) dx$	539
3.111	$\int x \cosh(bx) \text{Chi}(bx) dx$	543
3.112	$\int x^2 \cosh(bx) \text{Chi}(bx) dx$	548
3.113	$\int x^3 \cosh(bx) \text{Chi}(bx) dx$	553
3.114	$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^3} dx$	559
3.115	$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^2} dx$	563
3.116	$\int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx$	567
3.117	$\int \text{Chi}(bx) \sinh(bx) dx$	570
3.118	$\int x \text{Chi}(bx) \sinh(bx) dx$	574
3.119	$\int x^2 \text{Chi}(bx) \sinh(bx) dx$	579
3.120	$\int x^3 \text{Chi}(bx) \sinh(bx) dx$	584
3.121	$\int \text{Chi}(2x) \sinh(5x) dx$	590
3.122	$\int \cosh(5x) \text{Chi}(2x) dx$	594
3.123	$\int x^2 \text{Chi}(a + bx) \sinh(a + bx) dx$	598
3.124	$\int x \text{Chi}(a + bx) \sinh(a + bx) dx$	605
3.125	$\int \text{Chi}(a + bx) \sinh(a + bx) dx$	610
3.126	$\int \frac{\text{Chi}(a+bx) \sinh(a+bx)}{x} dx$	614
3.127	$\int x^2 \cosh(a + bx) \text{Chi}(a + bx) dx$	617
3.128	$\int x \cosh(a + bx) \text{Chi}(a + bx) dx$	624
3.129	$\int \cosh(a + bx) \text{Chi}(a + bx) dx$	629
3.130	$\int \frac{\cosh(a+bx) \text{Chi}(a+bx)}{x} dx$	633
3.131	$\int x \text{Chi}(c + dx) \sinh(a + bx) dx$	636
3.132	$\int \text{Chi}(c + dx) \sinh(a + bx) dx$	644
3.133	$\int \frac{\text{Chi}(c+dx) \sinh(a+bx)}{x} dx$	649
3.134	$\int x \cosh(a + bx) \text{Chi}(c + dx) dx$	652
3.135	$\int \cosh(a + bx) \text{Chi}(c + dx) dx$	660
3.136	$\int \frac{\cosh(a+bx) \text{Chi}(c+dx)}{x} dx$	665

3.1 $\int x^m \text{Shi}(bx) dx$

Optimal result	63
Rubi [A] (verified)	63
Mathematica [A] (verified)	64
Maple [C] (verified)	65
Fricas [F]	65
Sympy [A] (verification not implemented)	65
Maxima [F]	66
Giac [F]	66
Mupad [F(-1)]	66

Optimal result

Integrand size = 8, antiderivative size = 76

$$\int x^m \text{Shi}(bx) dx = -\frac{x^m(-bx)^{-m}\Gamma(1+m, -bx)}{2b(1+m)} - \frac{x^m(bx)^{-m}\Gamma(1+m, bx)}{2b(1+m)} + \frac{x^{1+m}\text{Shi}(bx)}{1+m}$$

[Out] $-1/2*x^m*\text{GAMMA}(1+m, -b*x)/b/(1+m)/((-b*x)^m) - 1/2*x^m*\text{GAMMA}(1+m, b*x)/b/(1+m)/((b*x)^m) + x^{(1+m)}*\text{Shi}(b*x)/(1+m)$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6667, 12, 3389, 2212}

$$\int x^m \text{Shi}(bx) dx = \frac{x^{m+1}\text{Shi}(bx)}{m+1} - \frac{x^m(-bx)^{-m}\Gamma(m+1, -bx)}{2b(m+1)} - \frac{x^m(bx)^{-m}\Gamma(m+1, bx)}{2b(m+1)}$$

[In] $\text{Int}[x^m*\text{SinhIntegral}[b*x], x]$

[Out] $-1/2*(x^m*\text{Gamma}[1+m, -(b*x)])/(b*(1+m)*(-(b*x))^m) - (x^m*\text{Gamma}[1+m, b*x])/(2*b*(1+m)*(b*x)^m) + (x^{(1+m)}*\text{SinhIntegral}[b*x])/(1+m)$

Rule 12

$\text{Int}[(a_*)*(u_), x_Symbol] := \text{Dist}[a, \text{Int}[u, x], x] /;$ FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 2212

$\text{Int}[(F_)^((g_)*((e_.) + (f_)*(x_)))*((c_.) + (d_)*(x_))^{(m_)}, x_Symbol] := \text{Simp}[(-F^{(g*(e - c*(f/d)))})*((c + d*x)^{\text{FracPart}[m]}/(d*((-f)*g*(\text{Log}[F]/d))$

```
)^(IntPart[m] + 1)*((-f)*g*Log[F]*((c + d*x)/d))^FracPart[m]]*Gamma[m + 1,
((-f)*g*(Log[F]/d))*(c + d*x)], x] /; FreeQ[{F, c, d, e, f, g, m}, x] &&
!IntegerQ[m]
```

Rule 3389

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Dist[I
/2, Int[(c + d*x)^m/E^(I*(e + f*x)), x], x] - Dist[I/2, Int[(c + d*x)^m*E^(
I*(e + f*x)), x], x] /; FreeQ[{c, d, e, f, m}, x]
```

Rule 6667

```
Int[((c_.) + (d_.)*(x_))^(m_.)*SinhIntegral[(a_.) + (b_.)*(x_)], x_Symbol]
:= Simp[(c + d*x)^(m + 1)*(SinhIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/
(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Sinh[a + b*x]/(a + b*x)), x], x] /; Fre
eQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x^{1+m}\text{Shi}(bx)}{1+m} - \frac{b \int \frac{x^m \sinh(bx)}{b} dx}{1+m} \\
 &= \frac{x^{1+m}\text{Shi}(bx)}{1+m} - \frac{\int x^m \sinh(bx) dx}{1+m} \\
 &= \frac{x^{1+m}\text{Shi}(bx)}{1+m} + \frac{\int e^{-bx} x^m dx}{2(1+m)} - \frac{\int e^{bx} x^m dx}{2(1+m)} \\
 &= -\frac{x^m (-bx)^{-m} \Gamma(1+m, -bx)}{2b(1+m)} - \frac{x^m (bx)^{-m} \Gamma(1+m, bx)}{2b(1+m)} + \frac{x^{1+m}\text{Shi}(bx)}{1+m}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.74

$$\int x^m \text{Shi}(bx) dx = -\frac{x^m((-bx)^{-m}\Gamma(1+m, -bx) + (bx)^{-m}\Gamma(1+m, bx) - 2bx\text{Shi}(bx))}{2b(1+m)}$$

```
[In] Integrate[x^m*SinhIntegral[b*x], x]
```

```
[Out] -1/2*(x^m*(Gamma[1 + m, -(b*x)]/(-(b*x))^m + Gamma[1 + m, b*x]/(b*x)^m - 2*
b*x*SinhIntegral[b*x]))/(b*(1 + m))
```


Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.60 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.49

method	result	size
meijerg	$\frac{bx^{2+m} \operatorname{hypergeom}\left(\left[\frac{1}{2}, 1+\frac{m}{2}\right], \left[\frac{3}{2}, \frac{3}{2}, 2+\frac{m}{2}\right], \frac{b^2x^2}{4}\right)}{2+m}$	37

[In] `int(x^m*Shi(b*x),x,method=_RETURNVERBOSE)`

[Out] `b/(2+m)*x^(2+m)*hypergeom([1/2,1+1/2*m],[3/2,3/2,2+1/2*m],1/4*b^2*x^2)`

Fricas [F]

$$\int x^m \operatorname{Shi}(bx) dx = \int x^m \operatorname{Shi}(bx) dx$$

[In] `integrate(x^m*Shi(b*x),x, algorithm="fricas")`

[Out] `integral(x^m*sinh_integral(b*x), x)`

Sympy [A] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.55

$$\int x^m \operatorname{Shi}(bx) dx = \frac{bx^{m+2} \Gamma\left(\frac{m}{2} + 1\right) {}_2F_3\left(\frac{1}{2}, \frac{m}{2} + 1 \mid \frac{b^2x^2}{4}\right)}{2\Gamma\left(\frac{m}{2} + 2\right)}$$

[In] `integrate(x**m*Shi(b*x),x)`

[Out] `b*x**(m + 2)*gamma(m/2 + 1)*hyper((1/2, m/2 + 1), (3/2, 3/2, m/2 + 2), b**2*x**2/4)/(2*gamma(m/2 + 2))`

Maxima [F]

$$\int x^m \operatorname{Shi}(bx) dx = \int x^m \operatorname{Shi}(bx) dx$$

[In] integrate(x^m*Shi(b*x),x, algorithm="maxima")

[Out] integrate(x^m*Shi(b*x), x)

Giac [F]

$$\int x^m \operatorname{Shi}(bx) dx = \int x^m \operatorname{Shi}(bx) dx$$

[In] integrate(x^m*Shi(b*x),x, algorithm="giac")

[Out] integrate(x^m*Shi(b*x), x)

Mupad [F(-1)]

Timed out.

$$\int x^m \operatorname{Shi}(bx) dx = \int x^m \operatorname{sinhint}(bx) dx$$

[In] int(x^m*sinhint(b*x),x)

[Out] int(x^m*sinhint(b*x), x)

3.2 $\int x^3 \text{Shi}(bx) dx$

Optimal result	67
Rubi [A] (verified)	67
Mathematica [A] (verified)	68
Maple [A] (verified)	69
Fricas [F]	69
Sympy [A] (verification not implemented)	69
Maxima [F]	70
Giac [F]	70
Mupad [F(-1)]	70

Optimal result

Integrand size = 8, antiderivative size = 63

$$\int x^3 \text{Shi}(bx) dx = -\frac{3x \cosh(bx)}{2b^3} - \frac{x^3 \cosh(bx)}{4b} + \frac{3 \sinh(bx)}{2b^4} + \frac{3x^2 \sinh(bx)}{4b^2} + \frac{1}{4}x^4 \text{Shi}(bx)$$

[Out] $-3/2*x*cosh(b*x)/b^3-1/4*x^3*cosh(b*x)/b+1/4*x^4*Shi(b*x)+3/2*sinh(b*x)/b^4+3/4*x^2*sinh(b*x)/b^2$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6667, 12, 3377, 2717}

$$\int x^3 \text{Shi}(bx) dx = \frac{3 \sinh(bx)}{2b^4} - \frac{3x \cosh(bx)}{2b^3} + \frac{3x^2 \sinh(bx)}{4b^2} + \frac{1}{4}x^4 \text{Shi}(bx) - \frac{x^3 \cosh(bx)}{4b}$$

[In] `Int[x^3*SinhIntegral[b*x],x]`

[Out] $(-3*x*Cosh[b*x])/(2*b^3) - (x^3*Cosh[b*x])/(4*b) + (3*Sinh[b*x])/(2*b^4) + (3*x^2*Sinh[b*x])/(4*b^2) + (x^4*SinhIntegral[b*x])/4$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 2717

`Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Rule 3377

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-
(c + d*x)^m)*(Cos[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Co
s[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 6667

```
Int[((c_.) + (d_.)*(x_))^(m_.)*SinhIntegral[(a_.) + (b_.)*(x_)], x_Symbol]
:= Simp[(c + d*x)^(m + 1)*(SinhIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/
(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Sinh[a + b*x]/(a + b*x)), x], x] /; Fre
eQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{4}x^4\text{Shi}(bx) - \frac{1}{4}b \int \frac{x^3 \sinh(bx)}{b} dx \\
&= \frac{1}{4}x^4\text{Shi}(bx) - \frac{1}{4} \int x^3 \sinh(bx) dx \\
&= -\frac{x^3 \cosh(bx)}{4b} + \frac{1}{4}x^4\text{Shi}(bx) + \frac{3 \int x^2 \cosh(bx) dx}{4b} \\
&= -\frac{x^3 \cosh(bx)}{4b} + \frac{3x^2 \sinh(bx)}{4b^2} + \frac{1}{4}x^4\text{Shi}(bx) - \frac{3 \int x \sinh(bx) dx}{2b^2} \\
&= -\frac{3x \cosh(bx)}{2b^3} - \frac{x^3 \cosh(bx)}{4b} + \frac{3x^2 \sinh(bx)}{4b^2} + \frac{1}{4}x^4\text{Shi}(bx) + \frac{3 \int \cosh(bx) dx}{2b^3} \\
&= -\frac{3x \cosh(bx)}{2b^3} - \frac{x^3 \cosh(bx)}{4b} + \frac{3 \sinh(bx)}{2b^4} + \frac{3x^2 \sinh(bx)}{4b^2} + \frac{1}{4}x^4\text{Shi}(bx)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.84

$$\int x^3\text{Shi}(bx) dx = -\frac{x(6 + b^2x^2) \cosh(bx)}{4b^3} + \frac{3(2 + b^2x^2) \sinh(bx)}{4b^4} + \frac{1}{4}x^4\text{Shi}(bx)$$

```
[In] Integrate[x^3*SinhIntegral[b*x],x]
```

```
[Out] -1/4*(x*(6 + b^2*x^2)*Cosh[b*x])/b^3 + (3*(2 + b^2*x^2)*Sinh[b*x])/(4*b^4)
+ (x^4*SinhIntegral[b*x])/4
```

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.86

method	result	size
parts	$\frac{x^4 \operatorname{Shi}(bx)}{4} - \frac{b^3 x^3 \cosh(bx) - 3b^2 x^2 \sinh(bx) + 6bx \cosh(bx) - 6 \sinh(bx)}{4b^4}$	54
derivativedivides	$\frac{\frac{b^4 x^4 \operatorname{Shi}(bx)}{4} - \frac{b^3 x^3 \cosh(bx)}{4} + \frac{3b^2 x^2 \sinh(bx)}{4} - \frac{3bx \cosh(bx)}{2} + \frac{3 \sinh(bx)}{2}}{b^4}$	56
default	$\frac{\frac{b^4 x^4 \operatorname{Shi}(bx)}{4} - \frac{b^3 x^3 \cosh(bx)}{4} + \frac{3b^2 x^2 \sinh(bx)}{4} - \frac{3bx \cosh(bx)}{2} + \frac{3 \sinh(bx)}{2}}{b^4}$	56
meijerg	$-\frac{4i\sqrt{\pi} \left(-\frac{ixb \left(\frac{5b^2 x^2}{2} + 15 \right) \cosh(bx)}{40\sqrt{\pi}} + \frac{i \left(\frac{15b^2 x^2}{2} + 15 \right) \sinh(bx)}{40\sqrt{\pi}} + \frac{ix^4 b^4 \operatorname{Shi}(bx)}{16\sqrt{\pi}} \right)}{b^4}$	69

```
[In] int(x^3*Shi(b*x),x,method=_RETURNVERBOSE)
```

```
[Out] 1/4*x^4*Shi(b*x)-1/4/b^4*(b^3*x^3*cosh(b*x)-3*b^2*x^2*sinh(b*x)+6*b*x*cosh(b*x)-6*sinh(b*x))
```

Fricas [F]

$$\int x^3 \operatorname{Shi}(bx) dx = \int x^3 \operatorname{Shi}(bx) dx$$

```
[In] integrate(x^3*Shi(b*x),x, algorithm="fricas")
```

```
[Out] integral(x^3*sinh_integral(b*x), x)
```

Sympy [A] (verification not implemented)

Time = 0.65 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.97

$$\int x^3 \operatorname{Shi}(bx) dx = \frac{x^4 \operatorname{Shi}(bx)}{4} - \frac{x^3 \cosh(bx)}{4b} + \frac{3x^2 \sinh(bx)}{4b^2} - \frac{3x \cosh(bx)}{2b^3} + \frac{3 \sinh(bx)}{2b^4}$$

```
[In] integrate(x**3*Shi(b*x),x)
```

```
[Out] x**4*Shi(b*x)/4 - x**3*cosh(b*x)/(4*b) + 3*x**2*sinh(b*x)/(4*b**2) - 3*x*cosh(b*x)/(2*b**3) + 3*sinh(b*x)/(2*b**4)
```

Maxima [F]

$$\int x^3 \operatorname{Shi}(bx) dx = \int x^3 \operatorname{Shi}(bx) dx$$

[In] integrate(x^3*Shi(b*x),x, algorithm="maxima")

[Out] integrate(x^3*Shi(b*x), x)

Giac [F]

$$\int x^3 \operatorname{Shi}(bx) dx = \int x^3 \operatorname{Shi}(bx) dx$$

[In] integrate(x^3*Shi(b*x),x, algorithm="giac")

[Out] integrate(x^3*Shi(b*x), x)

Mupad [F(-1)]

Timed out.

$$\int x^3 \operatorname{Shi}(bx) dx = \int x^3 \operatorname{sinhint}(bx) dx$$

[In] int(x^3*sinhint(b*x),x)

[Out] int(x^3*sinhint(b*x), x)

3.3 $\int x^2 \mathbf{Shi}(bx) dx$

Optimal result	71
Rubi [A] (verified)	71
Mathematica [A] (verified)	72
Maple [A] (verified)	73
Fricas [F]	73
Sympy [A] (verification not implemented)	73
Maxima [F]	74
Giac [F]	74
Mupad [F(-1)]	74

Optimal result

Integrand size = 8, antiderivative size = 49

$$\int x^2 \mathbf{Shi}(bx) dx = -\frac{2 \cosh(bx)}{3b^3} - \frac{x^2 \cosh(bx)}{3b} + \frac{2x \sinh(bx)}{3b^2} + \frac{1}{3} x^3 \mathbf{Shi}(bx)$$

[Out] $-2/3*\cosh(b*x)/b^3-1/3*x^2*\cosh(b*x)/b+1/3*x^3*\mathbf{Shi}(b*x)+2/3*x*\sinh(b*x)/b^2$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6667, 12, 3377, 2718}

$$\int x^2 \mathbf{Shi}(bx) dx = -\frac{2 \cosh(bx)}{3b^3} + \frac{2x \sinh(bx)}{3b^2} + \frac{1}{3} x^3 \mathbf{Shi}(bx) - \frac{x^2 \cosh(bx)}{3b}$$

[In] `Int[x^2*SinhIntegral[b*x],x]`

[Out] $(-2*\cosh[b*x])/(3*b^3) - (x^2*\cosh[b*x])/(3*b) + (2*x*\sinh[b*x])/(3*b^2) + (x^3*\mathbf{SinhIntegral}[b*x])/3$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 2718

`Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Rule 3377

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-
(c + d*x)^m)*(Cos[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Co
s[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 6667

```
Int[((c_.) + (d_.)*(x_))^(m_.)*SinhIntegral[(a_.) + (b_.)*(x_)], x_Symbol]
:= Simp[(c + d*x)^(m + 1)*(SinhIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/
(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Sinh[a + b*x]/(a + b*x)), x], x] /; Fre
eQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{3}x^3\text{Shi}(bx) - \frac{1}{3}b \int \frac{x^2 \sinh(bx)}{b} dx \\
&= \frac{1}{3}x^3\text{Shi}(bx) - \frac{1}{3} \int x^2 \sinh(bx) dx \\
&= -\frac{x^2 \cosh(bx)}{3b} + \frac{1}{3}x^3\text{Shi}(bx) + \frac{2 \int x \cosh(bx) dx}{3b} \\
&= -\frac{x^2 \cosh(bx)}{3b} + \frac{2x \sinh(bx)}{3b^2} + \frac{1}{3}x^3\text{Shi}(bx) - \frac{2 \int \sinh(bx) dx}{3b^2} \\
&= -\frac{2 \cosh(bx)}{3b^3} - \frac{x^2 \cosh(bx)}{3b} + \frac{2x \sinh(bx)}{3b^2} + \frac{1}{3}x^3\text{Shi}(bx)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.90

$$\int x^2\text{Shi}(bx) dx = -\frac{(2 + b^2x^2) \cosh(bx)}{3b^3} + \frac{2x \sinh(bx)}{3b^2} + \frac{1}{3}x^3\text{Shi}(bx)$$

```
[In] Integrate[x^2*SinhIntegral[b*x], x]
```

```
[Out] -1/3*((2 + b^2*x^2)*Cosh[b*x])/b^3 + (2*x*Sinh[b*x])/(3*b^2) + (x^3*SinhInt
egral[b*x])/3
```


Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.86

method	result	size
parts	$\frac{x^3 \operatorname{Shi}(bx)}{3} - \frac{b^2 x^2 \cosh(bx) - 2bx \sinh(bx) + 2 \cosh(bx)}{3b^3}$	42
derivativedivides	$\frac{\frac{b^3 x^3 \operatorname{Shi}(bx)}{3} - \frac{b^2 x^2 \cosh(bx)}{3} + \frac{2bx \sinh(bx)}{3} - \frac{2 \cosh(bx)}{3}}{b^3}$	44
default	$\frac{\frac{b^3 x^3 \operatorname{Shi}(bx)}{3} - \frac{b^2 x^2 \cosh(bx)}{3} + \frac{2bx \sinh(bx)}{3} - \frac{2 \cosh(bx)}{3}}{b^3}$	44
meijerg	$\frac{2\sqrt{\pi} \left(\frac{1}{3\sqrt{\pi}} - \frac{\left(\frac{b^2 x^2}{2} + 1\right) \cosh(bx)}{3\sqrt{\pi}} + \frac{bx \sinh(bx)}{3\sqrt{\pi}} + \frac{b^3 x^3 \operatorname{Shi}(bx)}{6\sqrt{\pi}} \right)}{b^3}$	60

[In] `int(x^2*Shi(b*x),x,method=_RETURNVERBOSE)`

[Out] `1/3*x^3*Shi(b*x)-1/3/b^3*(b^2*x^2*cosh(b*x)-2*b*x*sinh(b*x)+2*cosh(b*x))`

Fricas [F]

$$\int x^2 \operatorname{Shi}(bx) dx = \int x^2 \operatorname{Shi}(bx) dx$$

[In] `integrate(x^2*Shi(b*x),x, algorithm="fricas")`

[Out] `integral(x^2*sinh_integral(b*x), x)`

Sympy [A] (verification not implemented)

Time = 0.84 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.94

$$\int x^2 \operatorname{Shi}(bx) dx = \frac{x^3 \operatorname{Shi}(bx)}{3} - \frac{x^2 \cosh(bx)}{3b} + \frac{2x \sinh(bx)}{3b^2} - \frac{2 \cosh(bx)}{3b^3}$$

[In] `integrate(x**2*Shi(b*x),x)`

[Out] `x**3*Shi(b*x)/3 - x**2*cosh(b*x)/(3*b) + 2*x*sinh(b*x)/(3*b**2) - 2*cosh(b*x)/(3*b**3)`

Maxima [F]

$$\int x^2 \operatorname{Shi}(bx) dx = \int x^2 \operatorname{Shi}(bx) dx$$

[In] integrate(x^2*Shi(b*x),x, algorithm="maxima")

[Out] integrate(x^2*Shi(b*x), x)

Giac [F]

$$\int x^2 \operatorname{Shi}(bx) dx = \int x^2 \operatorname{Shi}(bx) dx$$

[In] integrate(x^2*Shi(b*x),x, algorithm="giac")

[Out] integrate(x^2*Shi(b*x), x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{Shi}(bx) dx = \frac{x^3 \operatorname{sinhint}(bx)}{3} - \frac{\frac{2 \cosh(bx)}{3} + \frac{b^2 x^2 \cosh(bx)}{3} - \frac{2 b x \sinh(bx)}{3}}{b^3}$$

[In] int(x^2*sinhint(b*x),x)

[Out] (x^3*sinhint(b*x))/3 - ((2*cosh(b*x))/3 + (b^2*x^2*cosh(b*x))/3 - (2*b*x*sinh(b*x))/3)/b^3

3.4 $\int x \operatorname{Shi}(bx) dx$

Optimal result	75
Rubi [A] (verified)	75
Mathematica [A] (verified)	76
Maple [A] (verified)	76
Fricas [F]	77
Sympy [A] (verification not implemented)	77
Maxima [F]	77
Giac [F]	78
Mupad [F(-1)]	78

Optimal result

Integrand size = 6, antiderivative size = 35

$$\int x \operatorname{Shi}(bx) dx = -\frac{x \cosh(bx)}{2b} + \frac{\sinh(bx)}{2b^2} + \frac{1}{2}x^2 \operatorname{Shi}(bx)$$

[Out] $-1/2*x*\cosh(b*x)/b+1/2*x^2*\operatorname{Shi}(b*x)+1/2*\sinh(b*x)/b^2$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {6667, 12, 3377, 2717}

$$\int x \operatorname{Shi}(bx) dx = \frac{\sinh(bx)}{2b^2} + \frac{1}{2}x^2 \operatorname{Shi}(bx) - \frac{x \cosh(bx)}{2b}$$

[In] `Int[x*SinhIntegral[b*x],x]`

[Out] $-1/2*(x*\cosh[b*x])/b + \operatorname{Sinh}[b*x]/(2*b^2) + (x^2*\operatorname{SinhIntegral}[b*x])/2$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 2717

`Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Rule 3377

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(
-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Co
s[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 6667

```
Int[((c_.) + (d_.)*(x_))^(m_.)*SinhIntegral[(a_.) + (b_.)*(x_)], x_Symbol]
:= Simp[(c + d*x)^(m + 1)*(SinhIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/
(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Sinh[a + b*x]/(a + b*x)), x], x] /; Fre
eQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{2}x^2\text{Shi}(bx) - \frac{1}{2}b \int \frac{x \sinh(bx)}{b} dx \\
&= \frac{1}{2}x^2\text{Shi}(bx) - \frac{1}{2} \int x \sinh(bx) dx \\
&= -\frac{x \cosh(bx)}{2b} + \frac{1}{2}x^2\text{Shi}(bx) + \frac{\int \cosh(bx) dx}{2b} \\
&= -\frac{x \cosh(bx)}{2b} + \frac{\sinh(bx)}{2b^2} + \frac{1}{2}x^2\text{Shi}(bx)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int x\text{Shi}(bx) dx = -\frac{x \cosh(bx)}{2b} + \frac{\sinh(bx)}{2b^2} + \frac{1}{2}x^2\text{Shi}(bx)$$

```
[In] Integrate[x*SinhIntegral[b*x],x]
```

```
[Out] -1/2*(x*Cosh[b*x])/b + Sinh[b*x]/(2*b^2) + (x^2*SinhIntegral[b*x])/2
```

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

method	result	size
parts	$\frac{x^2 \operatorname{Shi}(bx)}{2} - \frac{bx \cosh(bx) - \sinh(bx)}{2b^2}$	30
derivativedivides	$\frac{\frac{b^2 x^2 \operatorname{Shi}(bx)}{2} - \frac{bx \cosh(bx)}{2} + \frac{\sinh(bx)}{2}}{b^2}$	32
default	$\frac{\frac{b^2 x^2 \operatorname{Shi}(bx)}{2} - \frac{bx \cosh(bx)}{2} + \frac{\sinh(bx)}{2}}{b^2}$	32
meijerg	$\frac{i\sqrt{\pi} \left(\frac{ibx \cosh(bx)}{2\sqrt{\pi}} - \frac{i \sinh(bx)}{2\sqrt{\pi}} - \frac{ib^2 x^2 \operatorname{Shi}(bx)}{2\sqrt{\pi}} \right)}{b^2}$	49

[In] `int(x*Shi(b*x),x,method=_RETURNVERBOSE)`

[Out] `1/2*x^2*Shi(b*x)-1/2/b^2*(b*x*cosh(b*x)-sinh(b*x))`

Fricas [F]

$$\int x \operatorname{Shi}(bx) dx = \int x \operatorname{Shi}(bx) dx$$

[In] `integrate(x*Shi(b*x),x, algorithm="fricas")`

[Out] `integral(x*sinh_integral(b*x), x)`

Sympy [A] (verification not implemented)

Time = 0.50 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

$$\int x \operatorname{Shi}(bx) dx = \frac{x^2 \operatorname{Shi}(bx)}{2} - \frac{x \cosh(bx)}{2b} + \frac{\sinh(bx)}{2b^2}$$

[In] `integrate(x*Shi(b*x),x)`

[Out] `x**2*Shi(b*x)/2 - x*cosh(b*x)/(2*b) + sinh(b*x)/(2*b**2)`

Maxima [F]

$$\int x \operatorname{Shi}(bx) dx = \int x \operatorname{Shi}(bx) dx$$

[In] `integrate(x*Shi(b*x),x, algorithm="maxima")`

[Out] `integrate(x*Shi(b*x), x)`

Giac [F]

$$\int x\text{Shi}(bx) dx = \int x\text{Shi}(bx) dx$$

[In] integrate(x*Shi(b*x),x, algorithm="giac")

[Out] integrate(x*Shi(b*x), x)

Mupad [F(-1)]

Timed out.

$$\int x\text{Shi}(bx) dx = \frac{\frac{\sinh(bx)}{2} - \frac{bx \cosh(bx)}{2}}{b^2} + \frac{x^2 \text{sinhint}(bx)}{2}$$

[In] int(x*sinhint(b*x),x)

[Out] (sinh(b*x)/2 - (b*x*cosh(b*x))/2)/b^2 + (x^2*sinhint(b*x))/2

3.5 $\int \text{Shi}(bx) dx$

Optimal result	79
Rubi [A] (verified)	79
Mathematica [A] (verified)	80
Maple [A] (verified)	80
Fricas [F]	80
Sympy [A] (verification not implemented)	81
Maxima [F]	81
Giac [F]	81
Mupad [F(-1)]	81

Optimal result

Integrand size = 4, antiderivative size = 16

$$\int \text{Shi}(bx) dx = -\frac{\cosh(bx)}{b} + x\text{Shi}(bx)$$

[Out] $-\cosh(b*x)/b+x*\text{Shi}(b*x)$

Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6663}

$$\int \text{Shi}(bx) dx = x\text{Shi}(bx) - \frac{\cosh(bx)}{b}$$

[In] $\text{Int}[\text{SinhIntegral}[b*x], x]$

[Out] $-(\text{Cosh}[b*x]/b) + x*\text{SinhIntegral}[b*x]$

Rule 6663

$\text{Int}[\text{SinhIntegral}[(a_.) + (b_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[(a + b*x)*(\text{SinhIntegral}[a + b*x]/b), x] - \text{Simp}[\text{Cosh}[a + b*x]/b, x] /; \text{FreeQ}\{a, b\}, x]$

Rubi steps

$$\text{integral} = -\frac{\cosh(bx)}{b} + x\text{Shi}(bx)$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \operatorname{Shi}(bx) dx = -\frac{\cosh(bx)}{b} + x\operatorname{Shi}(bx)$$

[In] Integrate[SinhIntegral[b*x],x]

[Out] -(Cosh[b*x]/b) + x*SinhIntegral[b*x]

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

method	result	size
parts	$-\frac{\cosh(bx)}{b} + x \operatorname{Shi}(bx)$	17
derivativedivides	$\frac{\operatorname{Shi}(bx)bx - \cosh(bx)}{b}$	19
default	$\frac{\operatorname{Shi}(bx)bx - \cosh(bx)}{b}$	19
meijerg	$-\frac{\sqrt{\pi} \left(-\frac{2}{\sqrt{\pi}} + \frac{2 \cosh(bx)}{\sqrt{\pi}} - \frac{2bx \operatorname{Shi}(bx)}{\sqrt{\pi}} \right)}{2b}$	35

[In] int(Shi(b*x),x,method=_RETURNVERBOSE)

[Out] -cosh(b*x)/b+x*Shi(b*x)

Fricas [F]

$$\int \operatorname{Shi}(bx) dx = \int \operatorname{Shi}(bx) dx$$

[In] integrate(Shi(b*x),x, algorithm="fricas")

[Out] integral(sinh_integral(b*x), x)

Sympy [A] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \operatorname{Shi}(bx) dx = x \operatorname{Shi}(bx) - \frac{\cosh(bx)}{b}$$

[In] integrate(Shi(b*x),x)

[Out] x*Shi(b*x) - cosh(b*x)/b

Maxima [F]

$$\int \operatorname{Shi}(bx) dx = \int \operatorname{Shi}(bx) dx$$

[In] integrate(Shi(b*x),x, algorithm="maxima")

[Out] integrate(Shi(b*x), x)

Giac [F]

$$\int \operatorname{Shi}(bx) dx = \int \operatorname{Shi}(bx) dx$$

[In] integrate(Shi(b*x),x, algorithm="giac")

[Out] integrate(Shi(b*x), x)

Mupad [F(-1)]

Timed out.

$$\int \operatorname{Shi}(bx) dx = x \operatorname{sinhint}(bx) - \frac{\cosh(bx)}{b}$$

[In] int(sinhint(b*x),x)

[Out] x*sinhint(b*x) - cosh(b*x)/b

3.6 $\int \frac{\text{Shi}(bx)}{x} dx$

Optimal result	82
Rubi [A] (verified)	82
Mathematica [A] (verified)	83
Maple [A] (verified)	83
Fricas [F]	83
Sympy [A] (verification not implemented)	83
Maxima [F]	84
Giac [F]	84
Mupad [F(-1)]	84

Optimal result

Integrand size = 8, antiderivative size = 38

$$\int \frac{\text{Shi}(bx)}{x} dx = \frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; -bx) + \frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; bx)$$

[Out] 1/2*b*x*hypergeom([1, 1, 1],[2, 2, 2],-b*x)+1/2*b*x*hypergeom([1, 1, 1],[2, 2, 2],b*x)

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {6665}

$$\int \frac{\text{Shi}(bx)}{x} dx = \frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; -bx) + \frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; bx)$$

[In] Int[SinhIntegral[b*x]/x,x]

[Out] (b*x*HypergeometricPFQ[{1, 1, 1}, {2, 2, 2}, -(b*x)]/2 + (b*x*HypergeometricPFQ[{1, 1, 1}, {2, 2, 2}, b*x])/2

Rule 6665

Int[SinhIntegral[(b_.)*(x_)]/(x_), x_Symbol] := Simp[(1/2)*b*x*HypergeometricPFQ[{1, 1, 1}, {2, 2, 2}, (-b)*x], x] + Simp[(1/2)*b*x*HypergeometricPFQ[{1, 1, 1}, {2, 2, 2}, b*x], x] /; FreeQ[b, x]

Rubi steps

$$\text{integral} = \frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; -bx) + \frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; bx)$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{\text{Shi}(bx)}{x} dx = \frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; -bx) + \frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; bx)$$

[In] Integrate[SinhIntegral[b*x]/x,x]

[Out] (b*x*HypergeometricPFQ[{1, 1, 1}, {2, 2, 2}, -(b*x)])/2 + (b*x*HypergeometricPFQ[{1, 1, 1}, {2, 2, 2}, b*x])/2

Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.53

method	result	size
meijerg	$bx \text{ hypergeom} \left(\left[\frac{1}{2}, \frac{1}{2} \right], \left[\frac{3}{2}, \frac{3}{2}, \frac{3}{2} \right], \frac{b^2 x^2}{4} \right)$	20

[In] int(Shi(b*x)/x,x,method=_RETURNVERBOSE)

[Out] b*x*hypergeom([1/2,1/2],[3/2,3/2,3/2],1/4*b^2*x^2)

Fricas [F]

$$\int \frac{\text{Shi}(bx)}{x} dx = \int \frac{\text{Shi}(bx)}{x} dx$$

[In] integrate(Shi(b*x)/x,x, algorithm="fricas")

[Out] integral(sinh_integral(b*x)/x, x)

Sympy [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.53

$$\int \frac{\text{Shi}(bx)}{x} dx = bx {}_2F_3 \left(\frac{1}{2}, \frac{1}{2} \middle| \frac{b^2 x^2}{4} \right)$$

[In] integrate(Shi(b*x)/x,x)

[Out] b*x*hyper((1/2, 1/2), (3/2, 3/2, 3/2), b**2*x**2/4)

Maxima [F]

$$\int \frac{\text{Shi}(bx)}{x} dx = \int \frac{\text{Shi}(bx)}{x} dx$$

[In] integrate(Shi(b*x)/x,x, algorithm="maxima")

[Out] integrate(Shi(b*x)/x, x)

Giac [F]

$$\int \frac{\text{Shi}(bx)}{x} dx = \int \frac{\text{Shi}(bx)}{x} dx$$

[In] integrate(Shi(b*x)/x,x, algorithm="giac")

[Out] integrate(Shi(b*x)/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\text{Shi}(bx)}{x} dx = \int \frac{\text{sinhint}(bx)}{x} dx$$

[In] int(sinhint(b*x)/x,x)

[Out] int(sinhint(b*x)/x, x)

3.7 $\int \frac{\text{Shi}(bx)}{x^2} dx$

Optimal result	85
Rubi [A] (verified)	85
Mathematica [A] (verified)	86
Maple [A] (verified)	86
Fricas [F]	87
Sympy [A] (verification not implemented)	87
Maxima [F]	88
Giac [F]	88
Mupad [F(-1)]	88

Optimal result

Integrand size = 8, antiderivative size = 25

$$\int \frac{\text{Shi}(bx)}{x^2} dx = b\text{Chi}(bx) - \frac{\sinh(bx)}{x} - \frac{\text{Shi}(bx)}{x}$$

[Out] b*Chi(b*x)-Shi(b*x)/x-sinh(b*x)/x

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6667, 12, 3378, 3382}

$$\int \frac{\text{Shi}(bx)}{x^2} dx = b\text{Chi}(bx) - \frac{\text{Shi}(bx)}{x} - \frac{\sinh(bx)}{x}$$

[In] Int[SinhIntegral[b*x]/x^2,x]

[Out] b*CoshIntegral[b*x] - Sinh[b*x]/x - SinhIntegral[b*x]/x

Rule 12

Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 3378

Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] :> Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Dist[f/(d*(m + 1)), Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol]
:> Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x]
&& EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 6667

```
Int[((c_.) + (d_.)*(x_))^(m_.)*SinhIntegral[(a_.) + (b_.)*(x_)], x_Symbol]
:> Simp[(c + d*x)^(m + 1)*(SinhIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Sinh[a + b*x]/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Shi}(bx)}{x} + b \int \frac{\sinh(bx)}{bx^2} dx \\
&= -\frac{\text{Shi}(bx)}{x} + \int \frac{\sinh(bx)}{x^2} dx \\
&= -\frac{\sinh(bx)}{x} - \frac{\text{Shi}(bx)}{x} + b \int \frac{\cosh(bx)}{x} dx \\
&= b\text{Chi}(bx) - \frac{\sinh(bx)}{x} - \frac{\text{Shi}(bx)}{x}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{\text{Shi}(bx)}{x^2} dx = b\text{Chi}(bx) - \frac{\sinh(bx)}{x} - \frac{\text{Shi}(bx)}{x}$$

```
[In] Integrate[SinhIntegral[b*x]/x^2,x]
```

```
[Out] b*CoshIntegral[b*x] - Sinh[b*x]/x - SinhIntegral[b*x]/x
```

Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.20

method	result
parts	$-\frac{\text{Shi}(bx)}{x} + b\left(-\frac{\sinh(bx)}{bx} + \text{Chi}(bx)\right)$
derivativedivides	$b\left(-\frac{\text{Shi}(bx)}{bx} - \frac{\sinh(bx)}{bx} + \text{Chi}(bx)\right)$
default	$b\left(-\frac{\text{Shi}(bx)}{bx} - \frac{\sinh(bx)}{bx} + \text{Chi}(bx)\right)$
meijerg	$\frac{\sqrt{\pi} b \left(\frac{16}{\sqrt{\pi}} - \frac{4 e^{bx}}{\sqrt{\pi} bx} + \frac{4 e^{-bx}}{\sqrt{\pi} bx} - \frac{4(-9bx+9)(-\gamma-\ln(-bx)-\text{Ei}_1(-bx))}{9\sqrt{\pi} bx} + \frac{4(9bx+9)(-\gamma-\ln(bx)-\text{Ei}_1(bx))}{9\sqrt{\pi} bx} + \frac{8\gamma-16+8\ln(x)+8\ln(ib)}{\sqrt{\pi}} \right)}{8}$

```
[In] int(Shi(b*x)/x^2,x,method=_RETURNVERBOSE)
```

```
[Out] -Shi(b*x)/x+b*(-sinh(b*x)/b/x+Chi(b*x))
```

Fricas [F]

$$\int \frac{\text{Shi}(bx)}{x^2} dx = \int \frac{\text{Shi}(bx)}{x^2} dx$$

```
[In] integrate(Shi(b*x)/x^2,x, algorithm="fricas")
```

```
[Out] integral(sinh_integral(b*x)/x^2, x)
```

Sympy [A] (verification not implemented)

Time = 0.55 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.36

$$\int \frac{\text{Shi}(bx)}{x^2} dx = \frac{b^3 x^2 {}_3F_4\left(1, 1, \frac{3}{2} \mid \frac{b^2 x^2}{4}\right)}{36} + \frac{b \log(b^2 x^2)}{2}$$

```
[In] integrate(Shi(b*x)/x**2,x)
```

```
[Out] b**3*x**2*hyper((1, 1, 3/2), (2, 2, 5/2, 5/2), b**2*x**2/4)/36 + b*log(b**2*x**2)/2
```

Maxima [F]

$$\int \frac{\text{Shi}(bx)}{x^2} dx = \int \frac{\text{Shi}(bx)}{x^2} dx$$

[In] integrate(Shi(b*x)/x^2,x, algorithm="maxima")

[Out] integrate(Shi(b*x)/x^2, x)

Giac [F]

$$\int \frac{\text{Shi}(bx)}{x^2} dx = \int \frac{\text{Shi}(bx)}{x^2} dx$$

[In] integrate(Shi(b*x)/x^2,x, algorithm="giac")

[Out] integrate(Shi(b*x)/x^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\text{Shi}(bx)}{x^2} dx = \int \frac{\text{sinhint}(bx)}{x^2} dx$$

[In] int(sinhint(b*x)/x^2,x)

[Out] int(sinhint(b*x)/x^2, x)

3.8 $\int \frac{\text{Shi}(bx)}{x^3} dx$

Optimal result	89
Rubi [A] (verified)	89
Mathematica [A] (verified)	90
Maple [A] (verified)	91
Fricas [F]	91
Sympy [A] (verification not implemented)	91
Maxima [F]	92
Giac [F]	92
Mupad [F(-1)]	92

Optimal result

Integrand size = 8, antiderivative size = 46

$$\int \frac{\text{Shi}(bx)}{x^3} dx = -\frac{b \cosh(bx)}{4x} - \frac{\sinh(bx)}{4x^2} + \frac{1}{4}b^2\text{Shi}(bx) - \frac{\text{Shi}(bx)}{2x^2}$$

[Out] $-1/4*b*\cosh(b*x)/x+1/4*b^2*\text{Shi}(b*x)-1/2*\text{Shi}(b*x)/x^2-1/4*\sinh(b*x)/x^2$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6667, 12, 3378, 3379}

$$\int \frac{\text{Shi}(bx)}{x^3} dx = \frac{1}{4}b^2\text{Shi}(bx) - \frac{\text{Shi}(bx)}{2x^2} - \frac{\sinh(bx)}{4x^2} - \frac{b \cosh(bx)}{4x}$$

[In] `Int[SinhIntegral[b*x]/x^3,x]`

[Out] $-1/4*(b*\text{Cosh}[b*x])/x - \text{Sinh}[b*x]/(4*x^2) + (b^2*\text{SinhIntegral}[b*x])/4 - \text{SinhIntegral}[b*x]/(2*x^2)$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]`

Rule 3378

`Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Dist[f/(d*(m + 1)), Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]`

]

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol]
:> Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 6667

```
Int[((c_.) + (d_.)*(x_))^(m_.)*SinhIntegral[(a_.) + (b_.)*(x_)], x_Symbol]
:> Simp[(c + d*x)^(m + 1)*(SinhIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Sinh[a + b*x]/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Shi}(bx)}{2x^2} + \frac{1}{2}b \int \frac{\sinh(bx)}{bx^3} dx \\
&= -\frac{\text{Shi}(bx)}{2x^2} + \frac{1}{2} \int \frac{\sinh(bx)}{x^3} dx \\
&= -\frac{\sinh(bx)}{4x^2} - \frac{\text{Shi}(bx)}{2x^2} + \frac{1}{4}b \int \frac{\cosh(bx)}{x^2} dx \\
&= -\frac{b \cosh(bx)}{4x} - \frac{\sinh(bx)}{4x^2} - \frac{\text{Shi}(bx)}{2x^2} + \frac{1}{4}b^2 \int \frac{\sinh(bx)}{x} dx \\
&= -\frac{b \cosh(bx)}{4x} - \frac{\sinh(bx)}{4x^2} + \frac{1}{4}b^2 \text{Shi}(bx) - \frac{\text{Shi}(bx)}{2x^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{\text{Shi}(bx)}{x^3} dx = -\frac{b \cosh(bx)}{4x} - \frac{\sinh(bx)}{4x^2} + \frac{1}{4}b^2 \text{Shi}(bx) - \frac{\text{Shi}(bx)}{2x^2}$$

```
[In] Integrate[SinhIntegral[b*x]/x^3,x]
```

```
[Out] -1/4*(b*Cosh[b*x])/x - Sinh[b*x]/(4*x^2) + (b^2*SinhIntegral[b*x])/4 - SinhIntegral[b*x]/(2*x^2)
```

Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.02

method	result	size
parts	$-\frac{\text{Shi}(bx)}{2x^2} + \frac{b^2 \left(-\frac{\sinh(bx)}{2b^2x^2} - \frac{\cosh(bx)}{2bx} + \frac{\text{Shi}(bx)}{2} \right)}{2}$	47
derivativedivides	$b^2 \left(-\frac{\text{Shi}(bx)}{2b^2x^2} - \frac{\sinh(bx)}{4b^2x^2} - \frac{\cosh(bx)}{4bx} + \frac{\text{Shi}(bx)}{4} \right)$	48
default	$b^2 \left(-\frac{\text{Shi}(bx)}{2b^2x^2} - \frac{\sinh(bx)}{4b^2x^2} - \frac{\cosh(bx)}{4bx} + \frac{\text{Shi}(bx)}{4} \right)$	48
meijerg	$\frac{i\sqrt{\pi} b^2 \left(\frac{4i \cosh(bx)}{bx\sqrt{\pi}} + \frac{4i \sinh(bx)}{b^2x^2\sqrt{\pi}} + \frac{4i(-b^2x^2+2)\text{Shi}(bx)}{b^2x^2\sqrt{\pi}} \right)}{16}$	69

[In] int(Shi(b*x)/x^3,x,method=_RETURNVERBOSE)

[Out] -1/2*Shi(b*x)/x^2+1/2*b^2*(-1/2/b^2/x^2*sinh(b*x)-1/2/b/x*cosh(b*x)+1/2*Shi(b*x))

Fricas [F]

$$\int \frac{\text{Shi}(bx)}{x^3} dx = \int \frac{\text{Shi}(bx)}{x^3} dx$$

[In] integrate(Shi(b*x)/x^3,x, algorithm="fricas")

[Out] integral(sinh_integral(b*x)/x^3, x)

Sympy [A] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.85

$$\int \frac{\text{Shi}(bx)}{x^3} dx = \frac{b^2 \text{Shi}(bx)}{4} - \frac{b \cosh(bx)}{4x} - \frac{\sinh(bx)}{4x^2} - \frac{\text{Shi}(bx)}{2x^2}$$

[In] integrate(Shi(b*x)/x**3,x)

[Out] b**2*Shi(b*x)/4 - b*cosh(b*x)/(4*x) - sinh(b*x)/(4*x**2) - Shi(b*x)/(2*x**2)

Maxima [F]

$$\int \frac{\operatorname{Shi}(bx)}{x^3} dx = \int \frac{\operatorname{Shi}(bx)}{x^3} dx$$

[In] integrate(Shi(b*x)/x^3,x, algorithm="maxima")

[Out] integrate(Shi(b*x)/x^3, x)

Giac [F]

$$\int \frac{\operatorname{Shi}(bx)}{x^3} dx = \int \frac{\operatorname{Shi}(bx)}{x^3} dx$$

[In] integrate(Shi(b*x)/x^3,x, algorithm="giac")

[Out] integrate(Shi(b*x)/x^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{Shi}(bx)}{x^3} dx = \frac{b^2 \operatorname{sinhint}(bx)}{4} - \frac{\frac{\operatorname{sinhint}(bx)}{2} + \frac{\sinh(bx)}{4} + \frac{bx \cosh(bx)}{4}}{x^2}$$

[In] int(sinhint(b*x)/x^3,x)

[Out] (b^2*sinhint(b*x))/4 - (sinhint(b*x)/2 + sinh(b*x)/4 + (b*x*cosh(b*x))/4)/x^2

3.9 $\int x^m \text{Shi}(bx)^2 dx$

Optimal result	93
Rubi [N/A]	93
Mathematica [N/A]	94
Maple [N/A] (verified)	94
Fricas [N/A]	94
Sympy [N/A]	94
Maxima [N/A]	95
Giac [N/A]	95
Mupad [N/A]	95

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int x^m \text{Shi}(bx)^2 dx = \text{Int}(x^m \text{Shi}(bx)^2, x)$$

[Out] CannotIntegrate(x^m*Shi(b*x)^2,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int x^m \text{Shi}(bx)^2 dx = \int x^m \text{Shi}(bx)^2 dx$$

[In] Int[x^m*SinhIntegral[b*x]^2,x]

[Out] Defer[Int][x^m*SinhIntegral[b*x]^2, x]

Rubi steps

$$\text{integral} = \int x^m \text{Shi}(bx)^2 dx$$

Mathematica [N/A]

Not integrable

Time = 0.45 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Shi}(bx)^2 dx = \int x^m \operatorname{Shi}(bx)^2 dx$$

[In] Integrate[x^m*SinhIntegral[b*x]^2,x]

[Out] Integrate[x^m*SinhIntegral[b*x]^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.22 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int x^m \operatorname{Shi}(bx)^2 dx$$

[In] int(x^m*Shi(b*x)^2,x)

[Out] int(x^m*Shi(b*x)^2,x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Shi}(bx)^2 dx = \int x^m \operatorname{Shi}(bx)^2 dx$$

[In] integrate(x^m*Shi(b*x)^2,x, algorithm="fricas")

[Out] integral(x^m*sinh_integral(b*x)^2, x)

Sympy [N/A]

Not integrable

Time = 1.29 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int x^m \operatorname{Shi}(bx)^2 dx = \int x^m \operatorname{Shi}^2(bx) dx$$

[In] integrate(x**m*Shi(b*x)**2,x)

[Out] Integral(x**m*Shi(b*x)**2, x)

Maxima [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Shi}(bx)^2 dx = \int x^m \operatorname{Shi}(bx)^2 dx$$

[In] integrate(x^m*Shi(b*x)^2,x, algorithm="maxima")

[Out] integrate(x^m*Shi(b*x)^2, x)

Giac [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Shi}(bx)^2 dx = \int x^m \operatorname{Shi}(bx)^2 dx$$

[In] integrate(x^m*Shi(b*x)^2,x, algorithm="giac")

[Out] integrate(x^m*Shi(b*x)^2, x)

Mupad [N/A]

Not integrable

Time = 4.88 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Shi}(bx)^2 dx = \int x^m \operatorname{sinhint}(bx)^2 dx$$

[In] int(x^m*sinhint(b*x)^2,x)

[Out] int(x^m*sinhint(b*x)^2, x)

3.10 $\int x^3 \text{Shi}(bx)^2 dx$

Optimal result	96
Rubi [A] (verified)	96
Mathematica [A] (verified)	100
Maple [A] (verified)	100
Fricas [F]	100
Sympy [F]	101
Maxima [F]	101
Giac [F]	101
Mupad [F(-1)]	101

Optimal result

Integrand size = 10, antiderivative size = 149

$$\int x^3 \text{Shi}(bx)^2 dx = \frac{x^2}{2b^2} - \frac{3\text{Chi}(2bx)}{2b^4} + \frac{3\log(x)}{2b^4} - \frac{x \cosh(bx) \sinh(bx)}{b^3} + \frac{2 \sinh^2(bx)}{b^4}$$

$$+ \frac{x^2 \sinh^2(bx)}{4b^2} - \frac{3x \cosh(bx) \text{Shi}(bx)}{b^3} - \frac{x^3 \cosh(bx) \text{Shi}(bx)}{2b}$$

$$+ \frac{3 \sinh(bx) \text{Shi}(bx)}{b^4} + \frac{3x^2 \sinh(bx) \text{Shi}(bx)}{2b^2} + \frac{1}{4} x^4 \text{Shi}(bx)^2$$

[Out] $\frac{1}{2}x^2/b^2 - 3/2*\text{Chi}(2*b*x)/b^4 + 3/2*\ln(x)/b^4 - 3*x*\cosh(b*x)*\text{Shi}(b*x)/b^3 - 1/2*x^3*\cosh(b*x)*\text{Shi}(b*x)/b + 1/4*x^4*\text{Shi}(b*x)^2 - x*\cosh(b*x)*\sinh(b*x)/b^3 + 3*\text{Shi}(b*x)*\sinh(b*x)/b^4 + 3/2*x^2*\text{Shi}(b*x)*\sinh(b*x)/b^2 + 2*\sinh(b*x)^2/b^4 + 1/4*x^4*\sinh(b*x)^2/b^2$

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.00, number of steps used = 19, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 1.100$, Rules used = {6671, 6677, 12, 5480, 3391, 30, 6683, 2644, 6681, 3393, 3382}

$$\int x^3 \text{Shi}(bx)^2 dx = -\frac{3\text{Chi}(2bx)}{2b^4} + \frac{3\text{Shi}(bx) \sinh(bx)}{b^4} + \frac{3\log(x)}{2b^4} + \frac{2 \sinh^2(bx)}{b^4}$$

$$- \frac{3x \text{Shi}(bx) \cosh(bx)}{b^3} - \frac{x \sinh(bx) \cosh(bx)}{b^3} + \frac{3x^2 \text{Shi}(bx) \sinh(bx)}{2b^2}$$

$$+ \frac{x^2}{2b^2} + \frac{x^2 \sinh^2(bx)}{4b^2} + \frac{1}{4} x^4 \text{Shi}(bx)^2 - \frac{x^3 \text{Shi}(bx) \cosh(bx)}{2b}$$

[In] $\text{Int}[x^3*\text{SinhIntegral}[b*x]^2,x]$


```
[Out] x^2/(2*b^2) - (3*CoshIntegral[2*b*x])/(2*b^4) + (3*Log[x])/(2*b^4) - (x*Cosh[b*x]*Sinh[b*x])/b^3 + (2*Sinh[b*x]^2)/b^4 + (x^2*Sinh[b*x]^2)/(4*b^2) - (3*x*Cosh[b*x]*SinhIntegral[b*x])/b^3 - (x^3*Cosh[b*x]*SinhIntegral[b*x])/(2*b) + (3*Sinh[b*x]*SinhIntegral[b*x])/b^4 + (3*x^2*Sinh[b*x]*SinhIntegral[b*x])/(2*b^2) + (x^4*SinhIntegral[b*x]^2)/4
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] :=> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 30

```
Int[(x_)^(m_), x_Symbol] :=> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]
```

Rule 2644

```
Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :=> Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])
```

Rule 3382

```
Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] :=> Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3391

```
Int[((c_) + (d_)*(x_))*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :=> Simp[d*((b*Sin[e + f*x])^n/(f^2*n^2)), x] + (Dist[b^2*((n - 1)/n), Int[(c + d*x)*(b*Sin[e + f*x])^(n - 2), x], x] - Simp[b*(c + d*x)*Cos[e + f*x]*((b*Sin[e + f*x])^(n - 1)/(f*n)), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1]
```

Rule 3393

```
Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] :=> Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 5480

```
Int[Cosh[(a_) + (b_)*(x_)^(n_)]*(x_)^(m_)*Sinh[(a_) + (b_)*(x_)^(n_)]^(p_), x_Symbol] :=> Simp[x^(m - n + 1)*(Sinh[a + b*x^n]^(p + 1)/(b*n*(p +
```

1))), x] - Dist[(m - n + 1)/(b*n*(p + 1)), Int[x^(m - n)*Sinh[a + b*x^n]^(p + 1), x], x] /; FreeQ[{a, b, p}, x] && LtQ[0, n, m + 1] && NeQ[p, -1]

Rule 6671

Int[(x_)^(m_)*SinhIntegral[(b_)*(x_)]^2, x_Symbol] := Simp[x^(m + 1)*(SinhIntegral[b*x]^2/(m + 1)), x] - Dist[2/(m + 1), Int[x^m*Sinh[b*x]*SinhIntegral[b*x], x], x] /; FreeQ[b, x] && IGtQ[m, 0]

Rule 6677

Int[((e_) + (f_)*(x_))^(m_)*Sinh[(a_) + (b_)*(x_)]*SinhIntegral[(c_) + (d_)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*SinhIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rule 6681

Int[Cosh[(a_) + (b_)*(x_)]*SinhIntegral[(c_) + (d_)*(x_)], x_Symbol] := Simp[Sinh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]

Rule 6683

Int[Cosh[(a_) + (b_)*(x_)]*((e_) + (f_)*(x_))^(m_)*SinhIntegral[(c_) + (d_)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Sinh[a + b*x]*(SinhIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*SinhIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{4}x^4\text{Shi}(bx)^2 - \frac{1}{2}\int x^3\sinh(bx)\text{Shi}(bx)dx \\
 &= -\frac{x^3\cosh(bx)\text{Shi}(bx)}{2b} + \frac{1}{4}x^4\text{Shi}(bx)^2 \\
 &\quad + \frac{1}{2}\int\frac{x^2\cosh(bx)\sinh(bx)}{b}dx + \frac{3\int x^2\cosh(bx)\text{Shi}(bx)dx}{2b} \\
 &= -\frac{x^3\cosh(bx)\text{Shi}(bx)}{2b} + \frac{3x^2\sinh(bx)\text{Shi}(bx)}{2b^2} + \frac{1}{4}x^4\text{Shi}(bx)^2 \\
 &\quad - \frac{3\int x\sinh(bx)\text{Shi}(bx)dx}{b^2} + \frac{\int x^2\cosh(bx)\sinh(bx)dx}{2b} - \frac{3\int\frac{x\sinh^2(bx)}{b}dx}{2b}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{x^2 \sinh^2(bx)}{4b^2} - \frac{3x \cosh(bx) \text{Shi}(bx)}{b^3} - \frac{x^3 \cosh(bx) \text{Shi}(bx)}{2b} \\
&\quad + \frac{3x^2 \sinh(bx) \text{Shi}(bx)}{2b^2} + \frac{1}{4}x^4 \text{Shi}(bx)^2 + \frac{3 \int \cosh(bx) \text{Shi}(bx) dx}{b^3} \\
&\quad - \frac{\int x \sinh^2(bx) dx}{2b^2} - \frac{3 \int x \sinh^2(bx) dx}{2b^2} + \frac{3 \int \frac{\cosh(bx) \sinh(bx)}{b} dx}{b^2} \\
&= -\frac{x \cosh(bx) \sinh(bx)}{b^3} + \frac{\sinh^2(bx)}{2b^4} + \frac{x^2 \sinh^2(bx)}{4b^2} - \frac{3x \cosh(bx) \text{Shi}(bx)}{b^3} \\
&\quad - \frac{x^3 \cosh(bx) \text{Shi}(bx)}{2b} + \frac{3 \sinh(bx) \text{Shi}(bx)}{b^4} + \frac{3x^2 \sinh(bx) \text{Shi}(bx)}{2b^2} \\
&\quad + \frac{1}{4}x^4 \text{Shi}(bx)^2 + \frac{3 \int \cosh(bx) \sinh(bx) dx}{b^3} - \frac{3 \int \frac{\sinh^2(bx)}{bx} dx}{b^3} + \frac{\int x dx}{4b^2} + \frac{3 \int x dx}{4b^2} \\
&= \frac{x^2}{2b^2} - \frac{x \cosh(bx) \sinh(bx)}{b^3} + \frac{\sinh^2(bx)}{2b^4} + \frac{x^2 \sinh^2(bx)}{4b^2} - \frac{3x \cosh(bx) \text{Shi}(bx)}{b^3} \\
&\quad - \frac{x^3 \cosh(bx) \text{Shi}(bx)}{2b} + \frac{3 \sinh(bx) \text{Shi}(bx)}{b^4} + \frac{3x^2 \sinh(bx) \text{Shi}(bx)}{2b^2} \\
&\quad + \frac{1}{4}x^4 \text{Shi}(bx)^2 - \frac{3 \int \frac{\sinh^2(bx)}{x} dx}{b^4} - \frac{3 \text{Subst}(\int x dx, x, i \sinh(bx))}{b^4} \\
&= \frac{x^2}{2b^2} - \frac{x \cosh(bx) \sinh(bx)}{b^3} + \frac{2 \sinh^2(bx)}{b^4} + \frac{x^2 \sinh^2(bx)}{4b^2} \\
&\quad - \frac{3x \cosh(bx) \text{Shi}(bx)}{b^3} - \frac{x^3 \cosh(bx) \text{Shi}(bx)}{2b} + \frac{3 \sinh(bx) \text{Shi}(bx)}{b^4} \\
&\quad + \frac{3x^2 \sinh(bx) \text{Shi}(bx)}{2b^2} + \frac{1}{4}x^4 \text{Shi}(bx)^2 + \frac{3 \int \left(\frac{1}{2x} - \frac{\cosh(2bx)}{2x} \right) dx}{b^4} \\
&= \frac{x^2}{2b^2} + \frac{3 \log(x)}{2b^4} - \frac{x \cosh(bx) \sinh(bx)}{b^3} + \frac{2 \sinh^2(bx)}{b^4} + \frac{x^2 \sinh^2(bx)}{4b^2} \\
&\quad - \frac{3x \cosh(bx) \text{Shi}(bx)}{b^3} - \frac{x^3 \cosh(bx) \text{Shi}(bx)}{2b} + \frac{3 \sinh(bx) \text{Shi}(bx)}{b^4} \\
&\quad + \frac{3x^2 \sinh(bx) \text{Shi}(bx)}{2b^2} + \frac{1}{4}x^4 \text{Shi}(bx)^2 - \frac{3 \int \frac{\cosh(2bx)}{x} dx}{2b^4} \\
&= \frac{x^2}{2b^2} - \frac{3 \text{Chi}(2bx)}{2b^4} + \frac{3 \log(x)}{2b^4} - \frac{x \cosh(bx) \sinh(bx)}{b^3} + \frac{2 \sinh^2(bx)}{b^4} \\
&\quad + \frac{x^2 \sinh^2(bx)}{4b^2} - \frac{3x \cosh(bx) \text{Shi}(bx)}{b^3} - \frac{x^3 \cosh(bx) \text{Shi}(bx)}{2b} \\
&\quad + \frac{3 \sinh(bx) \text{Shi}(bx)}{b^4} + \frac{3x^2 \sinh(bx) \text{Shi}(bx)}{2b^2} + \frac{1}{4}x^4 \text{Shi}(bx)^2
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.72

$$\int x^3 \text{Shi}(bx)^2 dx = \frac{3b^2 x^2 + 8 \cosh(2bx) + b^2 x^2 \cosh(2bx) - 12 \text{Chi}(2bx) + 12 \log(x) - 4bx \sinh(2bx) - 4(bx(6 + b^2 x^2) \cosh(bx))}{8b^4}$$

[In] Integrate[x^3*SinhIntegral[b*x]^2,x]

[Out] (3*b^2*x^2 + 8*Cosh[2*b*x] + b^2*x^2*Cosh[2*b*x] - 12*CoshIntegral[2*b*x] + 12*Log[x] - 4*b*x*Sinh[2*b*x] - 4*(b*x*(6 + b^2*x^2)*Cosh[b*x] - 3*(2 + b^2*x^2)*Sinh[b*x])*SinhIntegral[b*x] + 2*b^4*x^4*SinhIntegral[b*x]^2)/(8*b^4)

Maple [A] (verified)

Time = 0.62 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.81

method	result
derivativedivides	$\frac{\frac{b^4 x^4 \text{Shi}(bx)^2}{4} - 2 \text{Shi}(bx) \left(\frac{b^3 x^3 \cosh(bx)}{4} - \frac{3b^2 x^2 \sinh(bx)}{4} + \frac{3bx \cosh(bx)}{2} - \frac{3 \sinh(bx)}{2} \right) + \frac{b^2 x^2 \cosh(bx)^2}{4} - bx \cosh(bx) \sinh(bx) + \dots}{b^4}$
default	$\frac{\frac{b^4 x^4 \text{Shi}(bx)^2}{4} - 2 \text{Shi}(bx) \left(\frac{b^3 x^3 \cosh(bx)}{4} - \frac{3b^2 x^2 \sinh(bx)}{4} + \frac{3bx \cosh(bx)}{2} - \frac{3 \sinh(bx)}{2} \right) + \frac{b^2 x^2 \cosh(bx)^2}{4} - bx \cosh(bx) \sinh(bx) + \dots}{b^4}$

[In] int(x^3*Shi(b*x)^2,x,method=_RETURNVERBOSE)

[Out] 1/b^4*(1/4*b^4*x^4*Shi(b*x)^2-2*Shi(b*x)*(1/4*b^3*x^3*cosh(b*x)-3/4*b^2*x^2*sinh(b*x)+3/2*b*x*cosh(b*x)-3/2*sinh(b*x))+1/4*b^2*x^2*cosh(b*x)^2-b*x*cosh(b*x)*sinh(b*x)+1/4*b^2*x^2+2*cosh(b*x)^2+3/2*ln(b*x)-3/2*Chi(2*b*x))

Fricas [F]

$$\int x^3 \text{Shi}(bx)^2 dx = \int x^3 \text{Shi}(bx)^2 dx$$

[In] integrate(x^3*Shi(b*x)^2,x, algorithm="fricas")

[Out] integral(x^3*sinh_integral(b*x)^2, x)

Sympy [F]

$$\int x^3 \operatorname{Shi}(bx)^2 dx = \int x^3 \operatorname{Shi}^2(bx) dx$$

[In] `integrate(x**3*Shi(b*x)**2,x)`

[Out] `Integral(x**3*Shi(b*x)**2, x)`

Maxima [F]

$$\int x^3 \operatorname{Shi}(bx)^2 dx = \int x^3 \operatorname{Shi}(bx)^2 dx$$

[In] `integrate(x^3*Shi(b*x)^2,x, algorithm="maxima")`

[Out] `integrate(x^3*Shi(b*x)^2, x)`

Giac [F]

$$\int x^3 \operatorname{Shi}(bx)^2 dx = \int x^3 \operatorname{Shi}(bx)^2 dx$$

[In] `integrate(x^3*Shi(b*x)^2,x, algorithm="giac")`

[Out] `integrate(x^3*Shi(b*x)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^3 \operatorname{Shi}(bx)^2 dx = \int x^3 \operatorname{sinhint}(bx)^2 dx$$

[In] `int(x^3*sinhint(b*x)^2,x)`

[Out] `int(x^3*sinhint(b*x)^2, x)`

3.11 $\int x^2 \text{Shi}(bx)^2 dx$

Optimal result	102
Rubi [A] (verified)	102
Mathematica [A] (verified)	105
Maple [A] (verified)	105
Fricas [F]	106
Sympy [F]	106
Maxima [F]	106
Giac [F]	106
Mupad [F(-1)]	107

Optimal result

Integrand size = 10, antiderivative size = 112

$$\int x^2 \text{Shi}(bx)^2 dx = \frac{5x}{6b^2} - \frac{5 \cosh(bx) \sinh(bx)}{6b^3} + \frac{x \sinh^2(bx)}{3b^2} - \frac{4 \cosh(bx) \text{Shi}(bx)}{3b^3} \\ - \frac{2x^2 \cosh(bx) \text{Shi}(bx)}{3b} + \frac{4x \sinh(bx) \text{Shi}(bx)}{3b^2} + \frac{1}{3} x^3 \text{Shi}(bx)^2 + \frac{2 \text{Shi}(2bx)}{3b^3}$$

[Out] $5/6*x/b^2 - 4/3*\cosh(b*x)*\text{Shi}(b*x)/b^3 - 2/3*x^2*\cosh(b*x)*\text{Shi}(b*x)/b + 1/3*x^3*\text{Shi}(b*x)^2 + 2/3*\text{Shi}(2*b*x)/b^3 - 5/6*\cosh(b*x)*\sinh(b*x)/b^3 + 4/3*x*\text{Shi}(b*x)*\sinh(b*x)/b^2 + 1/3*x*\sinh(b*x)^2/b^2$

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {6671, 6677, 12, 5480, 2715, 8, 6683, 6675, 5556, 3379}

$$\int x^2 \text{Shi}(bx)^2 dx = \frac{2 \text{Shi}(2bx)}{3b^3} - \frac{4 \text{Shi}(bx) \cosh(bx)}{3b^3} - \frac{5 \sinh(bx) \cosh(bx)}{6b^3} + \frac{4x \text{Shi}(bx) \sinh(bx)}{3b^2} \\ + \frac{5x}{6b^2} + \frac{x \sinh^2(bx)}{3b^2} + \frac{1}{3} x^3 \text{Shi}(bx)^2 - \frac{2x^2 \text{Shi}(bx) \cosh(bx)}{3b}$$

[In] $\text{Int}[x^2*\text{SinhIntegral}[b*x]^2, x]$

[Out] $(5*x)/(6*b^2) - (5*\text{Cosh}[b*x]*\text{Sinh}[b*x])/(6*b^3) + (x*\text{Sinh}[b*x]^2)/(3*b^2) - (4*\text{Cosh}[b*x]*\text{SinhIntegral}[b*x])/(3*b^3) - (2*x^2*\text{Cosh}[b*x]*\text{SinhIntegral}[b*x])/(3*b) + (4*x*\text{Sinh}[b*x]*\text{SinhIntegral}[b*x])/(3*b^2) + (x^3*\text{SinhIntegral}[b*x]^2)/3 + (2*\text{SinhIntegral}[2*b*x])/(3*b^3)$

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 2715

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3379

`Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`

Rule 5480

`Int[Cosh[(a_) + (b_)*(x_)^(n_)]*(x_)^(m_)*Sinh[(a_) + (b_)*(x_)^(n_)]^(p_), x_Symbol] := Simp[x^(m - n + 1)*(Sinh[a + b*x^n]^(p + 1)/(b*n*(p + 1))), x] - Dist[(m - n + 1)/(b*n*(p + 1)), Int[x^(m - n)*Sinh[a + b*x^n]^(p + 1), x], x] /; FreeQ[{a, b, p}, x] && LtQ[0, n, m + 1] && NeQ[p, -1]`

Rule 5556

`Int[Cosh[(a_) + (b_)*(x_)]^(p_)*((c_) + (d_)*(x_))^(m_)*Sinh[(a_) + (b_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

Rule 6671

`Int[(x_)^(m_)*SinhIntegral[(b_)*(x_)]^2, x_Symbol] := Simp[x^(m + 1)*(SinhIntegral[b*x]^2/(m + 1)), x] - Dist[2/(m + 1), Int[x^m*Sinh[b*x]*SinhIntegral[b*x], x], x] /; FreeQ[b, x] && IGtQ[m, 0]`

Rule 6675

`Int[Sinh[(a_) + (b_)*(x_)]*SinhIntegral[(c_) + (d_)*(x_)], x_Symbol] := Simp[Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]`

Rule 6677

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] :> Simp[(e + f*x)^m*Cosh[a + b*x]*(SinhIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Sinh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*SinhIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6683

```
Int[Cosh[(a_.) + (b_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.)*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] :> Simp[(e + f*x)^m*Sinh[a + b*x]*(SinhIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Sinh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*SinhIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{3}x^3\text{Shi}(bx)^2 - \frac{2}{3}\int x^2 \sinh(bx)\text{Shi}(bx) dx \\
&= -\frac{2x^2 \cosh(bx)\text{Shi}(bx)}{3b} + \frac{1}{3}x^3\text{Shi}(bx)^2 \\
&\quad + \frac{2}{3}\int \frac{x \cosh(bx) \sinh(bx)}{b} dx + \frac{4}{3b}\int x \cosh(bx)\text{Shi}(bx) dx \\
&= -\frac{2x^2 \cosh(bx)\text{Shi}(bx)}{3b} + \frac{4x \sinh(bx)\text{Shi}(bx)}{3b^2} + \frac{1}{3}x^3\text{Shi}(bx)^2 \\
&\quad - \frac{4}{3b^2}\int \sinh(bx)\text{Shi}(bx) dx + \frac{2}{3b}\int x \cosh(bx) \sinh(bx) dx - \frac{4}{3b}\int \frac{\sinh^2(bx)}{b} dx \\
&= \frac{x \sinh^2(bx)}{3b^2} - \frac{4 \cosh(bx)\text{Shi}(bx)}{3b^3} - \frac{2x^2 \cosh(bx)\text{Shi}(bx)}{3b} + \frac{4x \sinh(bx)\text{Shi}(bx)}{3b^2} \\
&\quad + \frac{1}{3}x^3\text{Shi}(bx)^2 - \frac{\int \sinh^2(bx) dx}{3b^2} + \frac{4 \int \frac{\cosh(bx) \sinh(bx)}{bx} dx}{3b^2} - \frac{4 \int \sinh^2(bx) dx}{3b^2} \\
&= -\frac{5 \cosh(bx) \sinh(bx)}{6b^3} + \frac{x \sinh^2(bx)}{3b^2} - \frac{4 \cosh(bx)\text{Shi}(bx)}{3b^3} - \frac{2x^2 \cosh(bx)\text{Shi}(bx)}{3b} \\
&\quad + \frac{4x \sinh(bx)\text{Shi}(bx)}{3b^2} + \frac{1}{3}x^3\text{Shi}(bx)^2 + \frac{4 \int \frac{\cosh(bx) \sinh(bx)}{x} dx}{3b^3} + \frac{\int 1 dx}{6b^2} + \frac{2 \int 1 dx}{3b^2} \\
&= \frac{5x}{6b^2} - \frac{5 \cosh(bx) \sinh(bx)}{6b^3} + \frac{x \sinh^2(bx)}{3b^2} - \frac{4 \cosh(bx)\text{Shi}(bx)}{3b^3} \\
&\quad - \frac{2x^2 \cosh(bx)\text{Shi}(bx)}{3b} + \frac{4x \sinh(bx)\text{Shi}(bx)}{3b^2} + \frac{1}{3}x^3\text{Shi}(bx)^2 + \frac{4 \int \frac{\sinh(2bx)}{2x} dx}{3b^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{5x}{6b^2} - \frac{5 \cosh(bx) \sinh(bx)}{6b^3} + \frac{x \sinh^2(bx)}{3b^2} - \frac{4 \cosh(bx) \text{Shi}(bx)}{3b^3} \\
&\quad - \frac{2x^2 \cosh(bx) \text{Shi}(bx)}{3b} + \frac{4x \sinh(bx) \text{Shi}(bx)}{3b^2} + \frac{1}{3} x^3 \text{Shi}(bx)^2 + \frac{2 \int \frac{\sinh(2bx)}{x} dx}{3b^3} \\
&= \frac{5x}{6b^2} - \frac{5 \cosh(bx) \sinh(bx)}{6b^3} + \frac{x \sinh^2(bx)}{3b^2} - \frac{4 \cosh(bx) \text{Shi}(bx)}{3b^3} \\
&\quad - \frac{2x^2 \cosh(bx) \text{Shi}(bx)}{3b} + \frac{4x \sinh(bx) \text{Shi}(bx)}{3b^2} + \frac{1}{3} x^3 \text{Shi}(bx)^2 + \frac{2 \text{Shi}(2bx)}{3b^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.70

$$\int x^2 \text{Shi}(bx)^2 dx = \frac{8bx + 2bx \cosh(2bx) - 5 \sinh(2bx) - 8((2 + b^2x^2) \cosh(bx) - 2bx \sinh(bx)) \text{Shi}(bx) + 4b^3x^3 \text{Shi}(bx)^2 + 8 \text{Shi}(2bx)}{12b^3}$$

[In] Integrate[x^2*SinhIntegral[b*x]^2,x]

[Out] (8*b*x + 2*b*x*Cosh[2*b*x] - 5*Sinh[2*b*x] - 8*((2 + b^2*x^2)*Cosh[b*x] - 2*b*x*Sinh[b*x]))*SinhIntegral[b*x] + 4*b^3*x^3*SinhIntegral[b*x]^2 + 8*SinhIntegral[2*b*x]/(12*b^3)

Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.75

method	result	S
derivativedivides	$\frac{\frac{b^3x^3 \text{Shi}(bx)^2}{3} - 2 \text{Shi}(bx) \left(\frac{b^2x^2 \cosh(bx)}{3} - \frac{2bx \sinh(bx)}{3} + \frac{2 \cosh(bx)}{3} \right) + \frac{bx \cosh(bx)^2}{3} - \frac{5 \cosh(bx) \sinh(bx)}{6} + \frac{bx}{2} + \frac{2 \text{Shi}(2bx)}{3}}{b^3}$	8
default	$\frac{\frac{b^3x^3 \text{Shi}(bx)^2}{3} - 2 \text{Shi}(bx) \left(\frac{b^2x^2 \cosh(bx)}{3} - \frac{2bx \sinh(bx)}{3} + \frac{2 \cosh(bx)}{3} \right) + \frac{bx \cosh(bx)^2}{3} - \frac{5 \cosh(bx) \sinh(bx)}{6} + \frac{bx}{2} + \frac{2 \text{Shi}(2bx)}{3}}{b^3}$	8

[In] int(x^2*Shi(b*x)^2,x,method=_RETURNVERBOSE)

[Out] 1/b^3*(1/3*b^3*x^3*Shi(b*x)^2-2*Shi(b*x)*(1/3*b^2*x^2*cosh(b*x)-2/3*b*x*sinh(b*x)+2/3*cosh(b*x))+1/3*b*x*cosh(b*x)^2-5/6*cosh(b*x)*sinh(b*x)+1/2*b*x+2/3*Shi(2*b*x))

Fricas [F]

$$\int x^2 \operatorname{Shi}(bx)^2 dx = \int x^2 \operatorname{Shi}(bx)^2 dx$$

```
[In] integrate(x^2*Shi(b*x)^2,x, algorithm="fricas")
```

```
[Out] integral(x^2*sinh_integral(b*x)^2, x)
```

Sympy [F]

$$\int x^2 \operatorname{Shi}(bx)^2 dx = \int x^2 \operatorname{Shi}^2(bx) dx$$

```
[In] integrate(x**2*Shi(b*x)**2,x)
```

```
[Out] Integral(x**2*Shi(b*x)**2, x)
```

Maxima [F]

$$\int x^2 \operatorname{Shi}(bx)^2 dx = \int x^2 \operatorname{Shi}(bx)^2 dx$$

```
[In] integrate(x^2*Shi(b*x)^2,x, algorithm="maxima")
```

```
[Out] integrate(x^2*Shi(b*x)^2, x)
```

Giac [F]

$$\int x^2 \operatorname{Shi}(bx)^2 dx = \int x^2 \operatorname{Shi}(bx)^2 dx$$

```
[In] integrate(x^2*Shi(b*x)^2,x, algorithm="giac")
```

```
[Out] integrate(x^2*Shi(b*x)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{Shi}(bx)^2 dx = \int x^2 \operatorname{sinhint}(bx)^2 dx$$

[In] `int(x^2*sinhint(b*x)^2,x)`

[Out] `int(x^2*sinhint(b*x)^2, x)`

3.12 $\int x\text{Shi}(bx)^2 dx$

Optimal result	108
Rubi [A] (verified)	108
Mathematica [A] (verified)	110
Maple [A] (verified)	111
Fricas [F]	111
Sympy [F]	111
Maxima [F]	111
Giac [F]	112
Mupad [F(-1)]	112

Optimal result

Integrand size = 8, antiderivative size = 74

$$\int x\text{Shi}(bx)^2 dx = -\frac{\text{Chi}(2bx)}{2b^2} + \frac{\log(x)}{2b^2} + \frac{\sinh^2(bx)}{2b^2} - \frac{x \cosh(bx)\text{Shi}(bx)}{b} + \frac{\sinh(bx)\text{Shi}(bx)}{b^2} + \frac{1}{2}x^2\text{Shi}(bx)^2$$

[Out] $-1/2*\text{Chi}(2*b*x)/b^2+1/2*\ln(x)/b^2-x*\cosh(b*x)*\text{Shi}(b*x)/b+1/2*x^2*\text{Shi}(b*x)^2+\text{Shi}(b*x)*\sinh(b*x)/b^2+1/2*\sinh(b*x)^2/b^2$

Rubi [A] (verified)

Time = 0.07 (sec), antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {6671, 6677, 12, 2644, 30, 6681, 3393, 3382}

$$\int x\text{Shi}(bx)^2 dx = -\frac{\text{Chi}(2bx)}{2b^2} + \frac{\text{Shi}(bx) \sinh(bx)}{b^2} + \frac{\log(x)}{2b^2} + \frac{\sinh^2(bx)}{2b^2} + \frac{1}{2}x^2\text{Shi}(bx)^2 - \frac{x\text{Shi}(bx) \cosh(bx)}{b}$$

[In] `Int[x*SinhIntegral[b*x]^2,x]`

[Out] $-1/2*\text{CoshIntegral}[2*b*x]/b^2 + \text{Log}[x]/(2*b^2) + \text{Sinh}[b*x]^2/(2*b^2) - (x*\text{Cosh}[b*x]*\text{SinhIntegral}[b*x])/b + (\text{Sinh}[b*x]*\text{SinhIntegral}[b*x])/b^2 + (x^2*\text{SinhIntegral}[b*x]^2)/2$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]`

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}/(m+1), x] \text{ ; FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 2644

$\text{Int}[\cos[(e_.) + (f_.)(x_)]^{(n_.)}((a_.)\sin[(e_.) + (f_.)(x_)]^{(m_.)}, x_Symbol] \rightarrow \text{Dist}[1/(a*f), \text{Subst}[\text{Int}[x^{m*(1-x^2/a^2)^{((n-1)/2)}, x], x, a*\sin[e+f*x]], x] \text{ ; FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[(m-1)/2] \ \&\& \ \text{LtQ}[0, m, n])$

Rule 3382

$\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])(f_.)(x_)]/((c_.) + (d_.)(x_)), x_Symbol] \rightarrow \text{Simp}[\text{CoshIntegral}[c*f*(fz/d) + f*fz*x]/d, x] \text{ ; FreeQ}[\{c, d, e, f, fz\}, x] \ \&\& \ \text{EqQ}[d*(e - \text{Pi}/2) - c*f*fz*I, 0]$

Rule 3393

$\text{Int}[(c_.) + (d_.)(x_)]^{(m)}\sin[(e_.) + (f_.)(x_)]^{(n)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(c + d*x)^m, \sin[e + f*x]^n, x], x] \text{ ; FreeQ}[\{c, d, e, f, m\}, x] \ \&\& \ \text{IGtQ}[n, 1] \ \&\& \ (!\text{RationalQ}[m] \ || \ (\text{GeQ}[m, -1] \ \&\& \ \text{LtQ}[m, 1]))$

Rule 6671

$\text{Int}[(x_)^{(m)}*\text{SinhIntegral}[(b_.)(x_)]^2, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*(\text{SinhIntegral}[b*x]^2/(m+1)), x] - \text{Dist}[2/(m+1), \text{Int}[x^m*\text{Sinh}[b*x]*\text{SinhIntegral}[b*x], x], x] \text{ ; FreeQ}[b, x] \ \&\& \ \text{IGtQ}[m, 0]$

Rule 6677

$\text{Int}[(e_.) + (f_.)(x_)]^{(m)}*\text{Sinh}[(a_.) + (b_.)(x_)]*\text{SinhIntegral}[(c_.) + (d_.)(x_)], x_Symbol] \rightarrow \text{Simp}[(e + f*x)^m*\text{Cosh}[a + b*x]*(\text{SinhIntegral}[c + d*x]/b), x] + (-\text{Dist}[d/b, \text{Int}[(e + f*x)^m*\text{Cosh}[a + b*x]*(\text{Sinh}[c + d*x]/(c + d*x)), x], x] - \text{Dist}[f*(m/b), \text{Int}[(e + f*x)^{(m-1)}*\text{Cosh}[a + b*x]*\text{SinhIntegral}[c + d*x], x], x]) \text{ ; FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{IGtQ}[m, 0]$

Rule 6681

$\text{Int}[\text{Cosh}[(a_.) + (b_.)(x_)]*\text{SinhIntegral}[(c_.) + (d_.)(x_)], x_Symbol] \rightarrow \text{Simp}[\text{Sinh}[a + b*x]*(\text{SinhIntegral}[c + d*x]/b), x] - \text{Dist}[d/b, \text{Int}[\text{Sinh}[a + b*x]*(\text{Sinh}[c + d*x]/(c + d*x)), x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x]$

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{2}x^2\text{Shi}(bx)^2 - \int x \sinh(bx)\text{Shi}(bx) dx \\
&= -\frac{x \cosh(bx)\text{Shi}(bx)}{b} + \frac{1}{2}x^2\text{Shi}(bx)^2 + \frac{\int \cosh(bx)\text{Shi}(bx) dx}{b} + \int \frac{\cosh(bx) \sinh(bx)}{b} dx \\
&= -\frac{x \cosh(bx)\text{Shi}(bx)}{b} + \frac{\sinh(bx)\text{Shi}(bx)}{b^2} + \frac{1}{2}x^2\text{Shi}(bx)^2 \\
&\quad + \frac{\int \cosh(bx) \sinh(bx) dx}{b} - \frac{\int \frac{\sinh^2(bx)}{bx} dx}{b} \\
&= -\frac{x \cosh(bx)\text{Shi}(bx)}{b} + \frac{\sinh(bx)\text{Shi}(bx)}{b^2} + \frac{1}{2}x^2\text{Shi}(bx)^2 \\
&\quad - \frac{\int \frac{\sinh^2(bx)}{x} dx}{b^2} - \frac{\text{Subst}(\int x dx, x, i \sinh(bx))}{b^2} \\
&= \frac{\sinh^2(bx)}{2b^2} - \frac{x \cosh(bx)\text{Shi}(bx)}{b} + \frac{\sinh(bx)\text{Shi}(bx)}{b^2} + \frac{1}{2}x^2\text{Shi}(bx)^2 + \frac{\int \left(\frac{1}{2x} - \frac{\cosh(2bx)}{2x}\right) dx}{b^2} \\
&= \frac{\log(x)}{2b^2} + \frac{\sinh^2(bx)}{2b^2} - \frac{x \cosh(bx)\text{Shi}(bx)}{b} + \frac{\sinh(bx)\text{Shi}(bx)}{b^2} + \frac{1}{2}x^2\text{Shi}(bx)^2 - \frac{\int \frac{\cosh(2bx)}{x} dx}{2b^2} \\
&= -\frac{\text{Chi}(2bx)}{2b^2} + \frac{\log(x)}{2b^2} + \frac{\sinh^2(bx)}{2b^2} - \frac{x \cosh(bx)\text{Shi}(bx)}{b} + \frac{\sinh(bx)\text{Shi}(bx)}{b^2} + \frac{1}{2}x^2\text{Shi}(bx)^2
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.78

$$\begin{aligned}
&\int x\text{Shi}(bx)^2 dx \\
&= \frac{\cosh(2bx) - 2\text{Chi}(2bx) + 2\log(x) + (-4bx \cosh(bx) + 4\sinh(bx))\text{Shi}(bx) + 2b^2x^2\text{Shi}(bx)^2}{4b^2}
\end{aligned}$$

[In] Integrate[x*SinhIntegral[b*x]^2,x]

[Out] (Cosh[2*b*x] - 2*CoshIntegral[2*b*x] + 2*Log[x] + (-4*b*x*Cosh[b*x] + 4*Sinh[b*x])*SinhIntegral[b*x] + 2*b^2*x^2*SinhIntegral[b*x]^2)/(4*b^2)

Maple [A] (verified)

Time = 0.57 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$\frac{\frac{b^2 x^2 \operatorname{Shi}(bx)^2}{2} - 2 \operatorname{Shi}(bx) \left(\frac{bx \cosh(bx)}{2} - \frac{\sinh(bx)}{2} \right) + \frac{\cosh(bx)^2}{2} + \frac{\ln(bx)}{2} - \frac{\operatorname{Chi}(2bx)}{2}}{b^2}$	62
default	$\frac{\frac{b^2 x^2 \operatorname{Shi}(bx)^2}{2} - 2 \operatorname{Shi}(bx) \left(\frac{bx \cosh(bx)}{2} - \frac{\sinh(bx)}{2} \right) + \frac{\cosh(bx)^2}{2} + \frac{\ln(bx)}{2} - \frac{\operatorname{Chi}(2bx)}{2}}{b^2}$	62

[In] int(x*Shi(b*x)^2,x,method=_RETURNVERBOSE)

[Out] $\frac{1}{b^2} * \left(\frac{1}{2} * b^2 * x^2 * \operatorname{Shi}(bx)^2 - 2 * \operatorname{Shi}(bx) * \left(\frac{1}{2} * b * x * \cosh(bx) - \frac{1}{2} * \sinh(bx) \right) + \frac{1}{2} * \cosh(bx)^2 + \frac{1}{2} * \ln(bx) - \frac{1}{2} * \operatorname{Chi}(2bx) \right)$

Fricas [F]

$$\int x \operatorname{Shi}(bx)^2 dx = \int x \operatorname{Shi}(bx)^2 dx$$

[In] integrate(x*Shi(b*x)^2,x, algorithm="fricas")

[Out] integral(x*sinh_integral(b*x)^2, x)

Sympy [F]

$$\int x \operatorname{Shi}(bx)^2 dx = \int x \operatorname{Shi}^2(bx) dx$$

[In] integrate(x*Shi(b*x)**2,x)

[Out] Integral(x*Shi(b*x)**2, x)

Maxima [F]

$$\int x \operatorname{Shi}(bx)^2 dx = \int x \operatorname{Shi}(bx)^2 dx$$

[In] integrate(x*Shi(b*x)^2,x, algorithm="maxima")

[Out] integrate(x*Shi(b*x)^2, x)

Giac [F]

$$\int x \operatorname{Shi}(bx)^2 dx = \int x \operatorname{Shi}(bx)^2 dx$$

[In] integrate(x*Shi(b*x)^2,x, algorithm="giac")

[Out] integrate(x*Shi(b*x)^2, x)

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{Shi}(bx)^2 dx = \int x \operatorname{sinhint}(bx)^2 dx$$

[In] int(x*sinhint(b*x)^2,x)

[Out] int(x*sinhint(b*x)^2, x)

3.13 $\int \text{Shi}(bx)^2 dx$

Optimal result	113
Rubi [A] (verified)	113
Mathematica [A] (verified)	115
Maple [A] (verified)	115
Fricas [F]	115
Sympy [F]	115
Maxima [F]	116
Giac [F]	116
Mupad [F(-1)]	116

Optimal result

Integrand size = 6, antiderivative size = 31

$$\int \text{Shi}(bx)^2 dx = -\frac{2 \cosh(bx)\text{Shi}(bx)}{b} + x\text{Shi}(bx)^2 + \frac{\text{Shi}(2bx)}{b}$$

[Out] $-2*\cosh(b*x)*\text{Shi}(b*x)/b+x*\text{Shi}(b*x)^2+\text{Shi}(2*b*x)/b$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {6669, 6675, 12, 5556, 3379}

$$\int \text{Shi}(bx)^2 dx = x\text{Shi}(bx)^2 + \frac{\text{Shi}(2bx)}{b} - \frac{2\text{Shi}(bx) \cosh(bx)}{b}$$

[In] $\text{Int}[\text{SinhIntegral}[b*x]^2, x]$

[Out] $(-2*\text{Cosh}[b*x]*\text{SinhIntegral}[b*x])/b + x*\text{SinhIntegral}[b*x]^2 + \text{SinhIntegral}[2*b*x]/b$

Rule 12

$\text{Int}[(a_*)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[u, (b_*)*(v_)] /; \text{FreeQ}[b, x]$

Rule 3379

$\text{Int}[\sin[(e_*) + (\text{Complex}[0, fz_])*(f_*)*(x_)]/((c_*) + (d_*)*(x_)), x_Symbol] \rightarrow \text{Simp}[I*(\text{SinhIntegral}[c*f*(fz/d) + f*fz*x]/d), x] /; \text{FreeQ}[\{c, d, e, f$

, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

Rule 6669

```
Int[SinhIntegral[(a_.) + (b_.)*(x_)]^2, x_Symbol] := Simp[(a + b*x)*(SinhIn
tegral[a + b*x]^2/b), x] - Dist[2, Int[Sinh[a + b*x]*SinhIntegral[a + b*x],
x], x] /; FreeQ[{a, b}, x]
```

Rule 6675

```
Int[Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
Simp[Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= x\text{Shi}(bx)^2 - 2 \int \sinh(bx)\text{Shi}(bx) dx \\
&= -\frac{2 \cosh(bx)\text{Shi}(bx)}{b} + x\text{Shi}(bx)^2 + 2 \int \frac{\cosh(bx) \sinh(bx)}{bx} dx \\
&= -\frac{2 \cosh(bx)\text{Shi}(bx)}{b} + x\text{Shi}(bx)^2 + \frac{2 \int \frac{\cosh(bx) \sinh(bx)}{x} dx}{b} \\
&= -\frac{2 \cosh(bx)\text{Shi}(bx)}{b} + x\text{Shi}(bx)^2 + \frac{2 \int \frac{\sinh(2bx)}{2x} dx}{b} \\
&= -\frac{2 \cosh(bx)\text{Shi}(bx)}{b} + x\text{Shi}(bx)^2 + \frac{\int \frac{\sinh(2bx)}{x} dx}{b} \\
&= -\frac{2 \cosh(bx)\text{Shi}(bx)}{b} + x\text{Shi}(bx)^2 + \frac{\text{Shi}(2bx)}{b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \operatorname{Shi}(bx)^2 dx = -\frac{2 \cosh(bx) \operatorname{Shi}(bx)}{b} + x \operatorname{Shi}(bx)^2 + \frac{\operatorname{Shi}(2bx)}{b}$$

[In] Integrate[SinhIntegral[b*x]^2,x]

[Out] (-2*Cosh[b*x]*SinhIntegral[b*x])/b + x*SinhIntegral[b*x]^2 + SinhIntegral[2*b*x]/b

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.97

method	result	size
derivativedivides	$\frac{\operatorname{Shi}(bx)^2 bx - 2 \cosh(bx) \operatorname{Shi}(bx) + \operatorname{Shi}(2bx)}{b}$	30
default	$\frac{\operatorname{Shi}(bx)^2 bx - 2 \cosh(bx) \operatorname{Shi}(bx) + \operatorname{Shi}(2bx)}{b}$	30

[In] int(Shi(b*x)^2,x,method=_RETURNVERBOSE)

[Out] 1/b*(Shi(b*x)^2*b*x-2*cosh(b*x)*Shi(b*x)+Shi(2*b*x))

Fricas [F]

$$\int \operatorname{Shi}(bx)^2 dx = \int \operatorname{Shi}(bx)^2 dx$$

[In] integrate(Shi(b*x)^2,x, algorithm="fricas")

[Out] integral(sinh_integral(b*x)^2, x)

Sympy [F]

$$\int \operatorname{Shi}(bx)^2 dx = \int \operatorname{Shi}^2(bx) dx$$

[In] integrate(Shi(b*x)**2,x)

[Out] Integral(Shi(b*x)**2, x)

Maxima [F]

$$\int \operatorname{Shi}(bx)^2 dx = \int \operatorname{Shi}(bx)^2 dx$$

[In] integrate(Shi(b*x)^2,x, algorithm="maxima")

[Out] integrate(Shi(b*x)^2, x)

Giac [F]

$$\int \operatorname{Shi}(bx)^2 dx = \int \operatorname{Shi}(bx)^2 dx$$

[In] integrate(Shi(b*x)^2,x, algorithm="giac")

[Out] integrate(Shi(b*x)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \operatorname{Shi}(bx)^2 dx = \int \operatorname{sinhint}(bx)^2 dx$$

[In] int(sinhint(b*x)^2,x)

[Out] int(sinhint(b*x)^2, x)

3.14 $\int \frac{\text{Shi}(bx)^2}{x} dx$

Optimal result	117
Rubi [N/A]	117
Mathematica [N/A]	118
Maple [N/A] (verified)	118
Fricas [N/A]	118
Sympy [N/A]	118
Maxima [N/A]	119
Giac [N/A]	119
Mupad [N/A]	119

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\text{Shi}(bx)^2}{x} dx = \text{Int}\left(\frac{\text{Shi}(bx)^2}{x}, x\right)$$

[Out] CannotIntegrate(Shi(b*x)^2/x,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\text{Shi}(bx)^2}{x} dx = \int \frac{\text{Shi}(bx)^2}{x} dx$$

[In] Int[SinhIntegral[b*x]^2/x,x]

[Out] Defer[Int][SinhIntegral[b*x]^2/x, x]

Rubi steps

$$\text{integral} = \int \frac{\text{Shi}(bx)^2}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Shi}(bx)^2}{x} dx = \int \frac{\text{Shi}(bx)^2}{x} dx$$

`[In] Integrate[SinhIntegral[b*x]^2/x,x]``[Out] Integrate[SinhIntegral[b*x]^2/x, x]`**Maple [N/A] (verified)**

Not integrable

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\text{Shi}(bx)^2}{x} dx$$

`[In] int(Shi(b*x)^2/x,x)``[Out] int(Shi(b*x)^2/x,x)`**Fricas [N/A]**

Not integrable

Time = 0.23 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Shi}(bx)^2}{x} dx = \int \frac{\text{Shi}(bx)^2}{x} dx$$

`[In] integrate(Shi(b*x)^2/x,x, algorithm="fricas")``[Out] integral(sinh_integral(b*x)^2/x, x)`**Sympy [N/A]**

Not integrable

Time = 1.07 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\text{Shi}(bx)^2}{x} dx = \int \frac{\text{Shi}^2(bx)}{x} dx$$

`[In] integrate(Shi(b*x)**2/x,x)``[Out] Integral(Shi(b*x)**2/x, x)`

Maxima [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Shi}(bx)^2}{x} dx = \int \frac{\text{Shi}(bx)^2}{x} dx$$

[In] integrate(Shi(b*x)^2/x,x, algorithm="maxima")

[Out] integrate(Shi(b*x)^2/x, x)

Giac [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Shi}(bx)^2}{x} dx = \int \frac{\text{Shi}(bx)^2}{x} dx$$

[In] integrate(Shi(b*x)^2/x,x, algorithm="giac")

[Out] integrate(Shi(b*x)^2/x, x)

Mupad [N/A]

Not integrable

Time = 4.92 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Shi}(bx)^2}{x} dx = \int \frac{\text{sinhint}(bx)^2}{x} dx$$

[In] int(sinhint(b*x)^2/x,x)

[Out] int(sinhint(b*x)^2/x, x)

3.15 $\int \frac{\mathbf{Shi}(bx)^2}{x^2} dx$

Optimal result	120
Rubi [N/A]	120
Mathematica [N/A]	121
Maple [N/A] (verified)	121
Fricas [N/A]	121
Sympy [N/A]	121
Maxima [N/A]	122
Giac [N/A]	122
Mupad [N/A]	122

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\mathbf{Shi}(bx)^2}{x^2} dx = \text{Int}\left(\frac{\mathbf{Shi}(bx)^2}{x^2}, x\right)$$

[Out] CannotIntegrate(Shi(b*x)^2/x^2,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\mathbf{Shi}(bx)^2}{x^2} dx = \int \frac{\mathbf{Shi}(bx)^2}{x^2} dx$$

[In] Int[SinhIntegral[b*x]^2/x^2,x]

[Out] Defer[Int][SinhIntegral[b*x]^2/x^2, x]

Rubi steps

$$\text{integral} = \int \frac{\mathbf{Shi}(bx)^2}{x^2} dx$$

Mathematica [N/A]

Not integrable

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Shi}(bx)^2}{x^2} dx = \int \frac{\text{Shi}(bx)^2}{x^2} dx$$

`[In] Integrate[SinhIntegral[b*x]^2/x^2,x]``[Out] Integrate[SinhIntegral[b*x]^2/x^2, x]`**Maple [N/A] (verified)**

Not integrable

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\text{Shi}(bx)^2}{x^2} dx$$

`[In] int(Shi(b*x)^2/x^2,x)``[Out] int(Shi(b*x)^2/x^2,x)`**Fricas [N/A]**

Not integrable

Time = 0.23 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Shi}(bx)^2}{x^2} dx = \int \frac{\text{Shi}(bx)^2}{x^2} dx$$

`[In] integrate(Shi(b*x)^2/x^2,x, algorithm="fricas")``[Out] integral(sinh_integral(b*x)^2/x^2, x)`**Sympy [N/A]**

Not integrable

Time = 0.98 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\text{Shi}(bx)^2}{x^2} dx = \int \frac{\text{Shi}^2(bx)}{x^2} dx$$

`[In] integrate(Shi(b*x)**2/x**2,x)``[Out] Integral(Shi(b*x)**2/x**2, x)`

Maxima [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\operatorname{Shi}(bx)^2}{x^2} dx = \int \frac{\operatorname{Shi}(bx)^2}{x^2} dx$$

[In] integrate(Shi(b*x)^2/x^2,x, algorithm="maxima")

[Out] integrate(Shi(b*x)^2/x^2, x)

Giac [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\operatorname{Shi}(bx)^2}{x^2} dx = \int \frac{\operatorname{Shi}(bx)^2}{x^2} dx$$

[In] integrate(Shi(b*x)^2/x^2,x, algorithm="giac")

[Out] integrate(Shi(b*x)^2/x^2, x)

Mupad [N/A]

Not integrable

Time = 4.84 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\operatorname{Shi}(bx)^2}{x^2} dx = \int \frac{\operatorname{sinhint}(bx)^2}{x^2} dx$$

[In] int(sinhint(b*x)^2/x^2,x)

[Out] int(sinhint(b*x)^2/x^2, x)

3.16 $\int \frac{\text{Shi}(bx)^2}{x^3} dx$

Optimal result	123
Rubi [N/A]	123
Mathematica [N/A]	124
Maple [N/A] (verified)	124
Fricas [N/A]	124
Sympy [N/A]	124
Maxima [N/A]	125
Giac [N/A]	125
Mupad [N/A]	125

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\text{Shi}(bx)^2}{x^3} dx = \text{Int}\left(\frac{\text{Shi}(bx)^2}{x^3}, x\right)$$

[Out] CannotIntegrate(Shi(b*x)^2/x^3,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\text{Shi}(bx)^2}{x^3} dx = \int \frac{\text{Shi}(bx)^2}{x^3} dx$$

[In] Int[SinhIntegral[b*x]^2/x^3,x]

[Out] Defer[Int][SinhIntegral[b*x]^2/x^3, x]

Rubi steps

$$\text{integral} = \int \frac{\text{Shi}(bx)^2}{x^3} dx$$

Mathematica [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Shi}(bx)^2}{x^3} dx = \int \frac{\text{Shi}(bx)^2}{x^3} dx$$

`[In] Integrate[SinhIntegral[b*x]^2/x^3,x]``[Out] Integrate[SinhIntegral[b*x]^2/x^3, x]`**Maple [N/A] (verified)**

Not integrable

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\text{Shi}(bx)^2}{x^3} dx$$

`[In] int(Shi(b*x)^2/x^3,x)``[Out] int(Shi(b*x)^2/x^3,x)`**Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Shi}(bx)^2}{x^3} dx = \int \frac{\text{Shi}(bx)^2}{x^3} dx$$

`[In] integrate(Shi(b*x)^2/x^3,x, algorithm="fricas")``[Out] integral(sinh_integral(b*x)^2/x^3, x)`**Sympy [N/A]**

Not integrable

Time = 1.14 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\text{Shi}(bx)^2}{x^3} dx = \int \frac{\text{Shi}^2(bx)}{x^3} dx$$

`[In] integrate(Shi(b*x)**2/x**3,x)``[Out] Integral(Shi(b*x)**2/x**3, x)`

Maxima [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\operatorname{Shi}(bx)^2}{x^3} dx = \int \frac{\operatorname{Shi}(bx)^2}{x^3} dx$$

[In] integrate(Shi(b*x)^2/x^3,x, algorithm="maxima")

[Out] integrate(Shi(b*x)^2/x^3, x)

Giac [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\operatorname{Shi}(bx)^2}{x^3} dx = \int \frac{\operatorname{Shi}(bx)^2}{x^3} dx$$

[In] integrate(Shi(b*x)^2/x^3,x, algorithm="giac")

[Out] integrate(Shi(b*x)^2/x^3, x)

Mupad [N/A]

Not integrable

Time = 4.88 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\operatorname{Shi}(bx)^2}{x^3} dx = \int \frac{\operatorname{sinhint}(bx)^2}{x^3} dx$$

[In] int(sinhint(b*x)^2/x^3,x)

[Out] int(sinhint(b*x)^2/x^3, x)

3.17 $\int x^m \mathbf{Shi}(a + bx) dx$

Optimal result	126
Rubi [N/A]	126
Mathematica [N/A]	127
Maple [N/A] (verified)	127
Fricas [N/A]	127
Sympy [N/A]	127
Maxima [N/A]	128
Giac [N/A]	128
Mupad [N/A]	128

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int x^m \mathbf{Shi}(a + bx) dx = \frac{x^{1+m} \mathbf{Shi}(a + bx)}{1 + m} - \frac{b \operatorname{Int}\left(\frac{x^{1+m} \sinh(a+bx)}{a+bx}, x\right)}{1 + m}$$

[Out] $-b \cdot \text{CannotIntegrate}(x^{(1+m)} \cdot \sinh(b \cdot x + a) / (b \cdot x + a), x) / (1+m) + x^{(1+m)} \cdot \mathbf{Shi}(b \cdot x + a) / (1+m)$

Rubi [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int x^m \mathbf{Shi}(a + bx) dx = \int x^m \mathbf{Shi}(a + bx) dx$$

[In] $\operatorname{Int}[x^m \cdot \operatorname{SinhIntegral}[a + b \cdot x], x]$

[Out] $(x^{(1 + m)} \cdot \operatorname{SinhIntegral}[a + b \cdot x]) / (1 + m) - (b \cdot \operatorname{Defer}[\operatorname{Int}][x^{(1 + m)} \cdot \operatorname{Sinh}[a + b \cdot x]) / (a + b \cdot x), x]) / (1 + m)$

Rubi steps

$$\text{integral} = \frac{x^{1+m} \mathbf{Shi}(a + bx)}{1 + m} - \frac{b \int \frac{x^{1+m} \sinh(a+bx)}{a+bx} dx}{1 + m}$$

Mathematica [N/A]

Not integrable

Time = 6.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Shi}(a + bx) dx = \int x^m \operatorname{Shi}(a + bx) dx$$

[In] Integrate[x^m*SinhIntegral[a + b*x],x][Out] Integrate[x^m*SinhIntegral[a + b*x], x]**Maple [N/A] (verified)**

Not integrable

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int x^m \operatorname{Shi}(bx + a) dx$$

[In] int(x^m*Shi(b*x+a),x)[Out] int(x^m*Shi(b*x+a),x)**Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Shi}(a + bx) dx = \int x^m \operatorname{Shi}(bx + a) dx$$

[In] integrate(x^m*Shi(b*x+a),x, algorithm="fricas")[Out] integral(x^m*sinh_integral(b*x + a), x)**Sympy [N/A]**

Not integrable

Time = 0.62 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int x^m \operatorname{Shi}(a + bx) dx = \int x^m \operatorname{Shi}(a + bx) dx$$

[In] integrate(x**m*Shi(b*x+a),x)

[Out] Integral(x**m*Shi(a + b*x), x)

Maxima [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Shi}(a + bx) dx = \int x^m \operatorname{Shi}(bx + a) dx$$

[In] integrate(x^m*Shi(b*x+a),x, algorithm="maxima")

[Out] integrate(x^m*Shi(b*x + a), x)

Giac [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Shi}(a + bx) dx = \int x^m \operatorname{Shi}(bx + a) dx$$

[In] integrate(x^m*Shi(b*x+a),x, algorithm="giac")

[Out] integrate(x^m*Shi(b*x + a), x)

Mupad [N/A]

Not integrable

Time = 4.85 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Shi}(a + bx) dx = \int x^m \operatorname{sinhint}(a + bx) dx$$

[In] int(x^m*sinhint(a + b*x),x)

[Out] int(x^m*sinhint(a + b*x), x)

3.18 $\int x^3 \text{Shi}(a + bx) dx$

Optimal result	129
Rubi [A] (verified)	129
Mathematica [A] (verified)	131
Maple [A] (verified)	132
Fricas [F]	132
Sympy [F]	132
Maxima [F]	133
Giac [F]	133
Mupad [F(-1)]	133

Optimal result

Integrand size = 10, antiderivative size = 184

$$\int x^3 \text{Shi}(a + bx) dx = \frac{a \cosh(a + bx)}{2b^4} + \frac{a^3 \cosh(a + bx)}{4b^4} - \frac{3x \cosh(a + bx)}{2b^3} - \frac{a^2 x \cosh(a + bx)}{4b^3} + \frac{ax^2 \cosh(a + bx)}{4b^2} - \frac{x^3 \cosh(a + bx)}{4b} + \frac{3 \sinh(a + bx)}{2b^4} + \frac{a^2 \sinh(a + bx)}{4b^4} - \frac{ax \sinh(a + bx)}{2b^3} + \frac{3x^2 \sinh(a + bx)}{4b^2} - \frac{a^4 \text{Shi}(a + bx)}{4b^4} + \frac{1}{4} x^4 \text{Shi}(a + bx)$$

[Out] $\frac{1}{2} a \cosh(bx+a)/b^4 + \frac{1}{4} a^3 \cosh(bx+a)/b^4 - \frac{3}{2} x \cosh(bx+a)/b^3 - \frac{1}{4} a^2 x^2 \cosh(bx+a)/b^3 + \frac{1}{4} a x^2 \cosh(bx+a)/b^2 - \frac{1}{4} x^3 \cosh(bx+a)/b - \frac{1}{4} a^4 \text{Shi}(bx+a)/b^4 + \frac{1}{4} x^4 \text{Shi}(bx+a) + \frac{3}{2} \sinh(bx+a)/b^4 + \frac{1}{4} a^2 \sinh(bx+a)/b^4 - \frac{1}{2} a x \sinh(bx+a)/b^3 + \frac{3}{4} x^2 \sinh(bx+a)/b^2$

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6667, 6874, 2718, 3377, 2717, 3379}

$$\int x^3 \text{Shi}(a + bx) dx = -\frac{a^4 \text{Shi}(a + bx)}{4b^4} + \frac{a^3 \cosh(a + bx)}{4b^4} + \frac{a^2 \sinh(a + bx)}{4b^4} - \frac{a^2 x \cosh(a + bx)}{4b^3} + \frac{3 \sinh(a + bx)}{2b^4} + \frac{a \cosh(a + bx)}{2b^4} - \frac{ax \sinh(a + bx)}{2b^3} - \frac{3x \cosh(a + bx)}{2b^3} + \frac{3x^2 \sinh(a + bx)}{4b^2} + \frac{ax^2 \cosh(a + bx)}{4b^2} + \frac{1}{4} x^4 \text{Shi}(a + bx) - \frac{x^3 \cosh(a + bx)}{4b}$$

[In] Int[x^3*SinhIntegral[a + b*x],x]

[Out] (a*Cosh[a + b*x])/(2*b^4) + (a^3*Cosh[a + b*x])/(4*b^4) - (3*x*Cosh[a + b*x])/(2*b^3) - (a^2*x*Cosh[a + b*x])/(4*b^3) + (a*x^2*Cosh[a + b*x])/(4*b^2) - (x^3*Cosh[a + b*x])/(4*b) + (3*Sinh[a + b*x])/(2*b^4) + (a^2*Sinh[a + b*x])/(4*b^4) - (a*x*Sinh[a + b*x])/(2*b^3) + (3*x^2*Sinh[a + b*x])/(4*b^2) - (a^4*SinhIntegral[a + b*x])/(4*b^4) + (x^4*SinhIntegral[a + b*x])/4

Rule 2717

Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rule 2718

Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rule 3377

Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 6667

Int[((c_.) + (d_.)*(x_))^(m_.)*SinhIntegral[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[(c + d*x)^(m + 1)*(SinhIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Sinh[a + b*x]/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]

Rule 6874

Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]

Rubi steps

$$\text{integral} = \frac{1}{4}x^4\text{Shi}(a + bx) - \frac{1}{4}b \int \frac{x^4 \sinh(a + bx)}{a + bx} dx$$

$$\begin{aligned}
&= \frac{1}{4}x^4\text{Shi}(a+bx) - \frac{1}{4}b \int \left(-\frac{a^3 \sinh(a+bx)}{b^4} + \frac{a^2x \sinh(a+bx)}{b^3} - \frac{ax^2 \sinh(a+bx)}{b^2} \right. \\
&\quad \left. + \frac{x^3 \sinh(a+bx)}{b} + \frac{a^4 \sinh(a+bx)}{b^4(a+bx)} \right) dx \\
&= \frac{1}{4}x^4\text{Shi}(a+bx) - \frac{1}{4} \int x^3 \sinh(a+bx) dx + \frac{a^3 \int \sinh(a+bx) dx}{4b^3} \\
&\quad - \frac{a^4 \int \frac{\sinh(a+bx)}{a+bx} dx}{4b^3} - \frac{a^2 \int x \sinh(a+bx) dx}{4b^2} + \frac{a \int x^2 \sinh(a+bx) dx}{4b} \\
&= \frac{a^3 \cosh(a+bx)}{4b^4} - \frac{a^2x \cosh(a+bx)}{4b^3} + \frac{ax^2 \cosh(a+bx)}{4b^2} - \frac{x^3 \cosh(a+bx)}{4b} - \frac{a^4\text{Shi}(a+bx)}{4b^4} \\
&\quad + \frac{1}{4}x^4\text{Shi}(a+bx) + \frac{a^2 \int \cosh(a+bx) dx}{4b^3} - \frac{a \int x \cosh(a+bx) dx}{2b^2} + \frac{3 \int x^2 \cosh(a+bx) dx}{4b} \\
&= \frac{a^3 \cosh(a+bx)}{4b^4} - \frac{a^2x \cosh(a+bx)}{4b^3} + \frac{ax^2 \cosh(a+bx)}{4b^2} - \frac{x^3 \cosh(a+bx)}{4b} \\
&\quad + \frac{a^2 \sinh(a+bx)}{4b^4} - \frac{ax \sinh(a+bx)}{2b^3} + \frac{3x^2 \sinh(a+bx)}{4b^2} - \frac{a^4\text{Shi}(a+bx)}{4b^4} \\
&\quad + \frac{1}{4}x^4\text{Shi}(a+bx) + \frac{a \int \sinh(a+bx) dx}{2b^3} - \frac{3 \int x \sinh(a+bx) dx}{2b^2} \\
&= \frac{a \cosh(a+bx)}{2b^4} + \frac{a^3 \cosh(a+bx)}{4b^4} - \frac{3x \cosh(a+bx)}{2b^3} - \frac{a^2x \cosh(a+bx)}{4b^3} \\
&\quad + \frac{ax^2 \cosh(a+bx)}{4b^2} - \frac{x^3 \cosh(a+bx)}{4b} + \frac{a^2 \sinh(a+bx)}{4b^4} - \frac{ax \sinh(a+bx)}{2b^3} \\
&\quad + \frac{3x^2 \sinh(a+bx)}{4b^2} - \frac{a^4\text{Shi}(a+bx)}{4b^4} + \frac{1}{4}x^4\text{Shi}(a+bx) + \frac{3 \int \cosh(a+bx) dx}{2b^3} \\
&= \frac{a \cosh(a+bx)}{2b^4} + \frac{a^3 \cosh(a+bx)}{4b^4} - \frac{3x \cosh(a+bx)}{2b^3} - \frac{a^2x \cosh(a+bx)}{4b^3} \\
&\quad + \frac{ax^2 \cosh(a+bx)}{4b^2} - \frac{x^3 \cosh(a+bx)}{4b} + \frac{3 \sinh(a+bx)}{2b^4} + \frac{a^2 \sinh(a+bx)}{4b^4} \\
&\quad - \frac{ax \sinh(a+bx)}{2b^3} + \frac{3x^2 \sinh(a+bx)}{4b^2} - \frac{a^4\text{Shi}(a+bx)}{4b^4} + \frac{1}{4}x^4\text{Shi}(a+bx)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.51

$$\begin{aligned}
&\int x^3\text{Shi}(a+bx) dx \\
&= \frac{(2a + a^3 - 6bx - a^2bx + ab^2x^2 - b^3x^3) \cosh(a+bx) + (6 + a^2 - 2abx + 3b^2x^2) \sinh(a+bx) + (-a^4 + b^4)}{4b^4}
\end{aligned}$$

[In] Integrate[x^3*SinhIntegral[a + b*x],x]

[Out] ((2*a + a^3 - 6*b*x - a^2*b*x + a*b^2*x^2 - b^3*x^3)*Cosh[a + b*x] + (6 + a^2 - 2*a*b*x + 3*b^2*x^2)*Sinh[a + b*x] + (-a^4 + b^4*x^4)*SinhIntegral[a + b*x])/(4*b^4)

Maple [A] (verified)

Time = 0.45 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.84

method	result
parts	$\frac{x^4 \operatorname{Shi}(bx+a)}{4} - \frac{a^4 \operatorname{Shi}(bx+a) - 4a^3 \cosh(bx+a) + 6a^2((bx+a) \cosh(bx+a) - \sinh(bx+a)) - 4a((bx+a)^2 \cosh(bx+a) - 2(bx+a) \sinh(bx+a))}{4}$
derivativedivides	$\frac{\operatorname{Shi}(bx+a)b^4 x^4 - \frac{a^4 \operatorname{Shi}(bx+a)}{4} + a^3 \cosh(bx+a) - \frac{3a^2((bx+a) \cosh(bx+a) - \sinh(bx+a))}{2} + a((bx+a)^2 \cosh(bx+a) - 2(bx+a) \sinh(bx+a))}{b^4}$
default	$\frac{\operatorname{Shi}(bx+a)b^4 x^4 - \frac{a^4 \operatorname{Shi}(bx+a)}{4} + a^3 \cosh(bx+a) - \frac{3a^2((bx+a) \cosh(bx+a) - \sinh(bx+a))}{2} + a((bx+a)^2 \cosh(bx+a) - 2(bx+a) \sinh(bx+a))}{b^4}$

[In] int(x^3*Shi(b*x+a),x,method=_RETURNVERBOSE)

```
[Out] 1/4*x^4*Shi(b*x+a)-1/4/b^4*(a^4*Shi(b*x+a)-4*a^3*cosh(b*x+a)+6*a^2*((b*x+a)
*cosh(b*x+a)-sinh(b*x+a))-4*a*((b*x+a)^2*cosh(b*x+a)-2*(b*x+a)*sinh(b*x+a)+
2*cosh(b*x+a))+b^4*x^4-frac(a^4*Shi(b*x+a)+a^3*cosh(b*x+a)-frac(3*a^2*((b*x+a)
*cosh(b*x+a)-sinh(b*x+a)))+a*((b*x+a)^2*cosh(b*x+a)-2*(b*x+a)*sinh(b*x+a))}{b^4}
```

Fricas [F]

$$\int x^3 \operatorname{Shi}(a + bx) dx = \int x^3 \operatorname{Shi}(bx + a) dx$$

[In] integrate(x^3*Shi(b*x+a),x, algorithm="fricas")

[Out] integral(x^3*sinh_integral(b*x + a), x)

Sympy [F]

$$\int x^3 \operatorname{Shi}(a + bx) dx = \int x^3 \operatorname{Shi}(a + bx) dx$$

[In] integrate(x**3*Shi(b*x+a),x)

[Out] Integral(x**3*Shi(a + b*x), x)

Maxima [F]

$$\int x^3 \operatorname{Shi}(a + bx) dx = \int x^3 \operatorname{Shi}(bx + a) dx$$

[In] integrate(x^3*Shi(b*x+a),x, algorithm="maxima")

[Out] integrate(x^3*Shi(b*x + a), x)

Giac [F]

$$\int x^3 \operatorname{Shi}(a + bx) dx = \int x^3 \operatorname{Shi}(bx + a) dx$$

[In] integrate(x^3*Shi(b*x+a),x, algorithm="giac")

[Out] integrate(x^3*Shi(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int x^3 \operatorname{Shi}(a + bx) dx = \int x^3 \operatorname{sinhint}(a + bx) dx$$

[In] int(x^3*sinhint(a + b*x),x)

[Out] int(x^3*sinhint(a + b*x), x)

3.19 $\int x^2 \text{Shi}(a + bx) dx$

Optimal result	134
Rubi [A] (verified)	134
Mathematica [A] (verified)	136
Maple [A] (verified)	136
Fricas [F]	137
Sympy [F]	137
Maxima [F]	137
Giac [F]	137
Mupad [F(-1)]	138

Optimal result

Integrand size = 10, antiderivative size = 118

$$\int x^2 \text{Shi}(a + bx) dx = -\frac{2 \cosh(a + bx)}{3b^3} - \frac{a^2 \cosh(a + bx)}{3b^3} + \frac{ax \cosh(a + bx)}{3b^2} - \frac{x^2 \cosh(a + bx)}{3b} \\ - \frac{a \sinh(a + bx)}{3b^3} + \frac{2x \sinh(a + bx)}{3b^2} + \frac{a^3 \text{Shi}(a + bx)}{3b^3} + \frac{1}{3} x^3 \text{Shi}(a + bx)$$

[Out] $-2/3*\cosh(b*x+a)/b^3-1/3*a^2*\cosh(b*x+a)/b^3+1/3*a*x*\cosh(b*x+a)/b^2-1/3*x^2*\cosh(b*x+a)/b+1/3*a^3*\text{Shi}(b*x+a)/b^3+1/3*x^3*\text{Shi}(b*x+a)-1/3*a*\sinh(b*x+a)/b^3+2/3*x*\sinh(b*x+a)/b^2$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6667, 6874, 2718, 3377, 2717, 3379}

$$\int x^2 \text{Shi}(a + bx) dx = \frac{a^3 \text{Shi}(a + bx)}{3b^3} - \frac{a^2 \cosh(a + bx)}{3b^3} - \frac{a \sinh(a + bx)}{3b^3} \\ - \frac{2 \cosh(a + bx)}{3b^3} + \frac{2x \sinh(a + bx)}{3b^2} + \frac{ax \cosh(a + bx)}{3b^2} \\ + \frac{1}{3} x^3 \text{Shi}(a + bx) - \frac{x^2 \cosh(a + bx)}{3b}$$

[In] $\text{Int}[x^2*\text{SinhIntegral}[a + b*x], x]$

[Out] $(-2*\text{Cosh}[a + b*x])/(3*b^3) - (a^2*\text{Cosh}[a + b*x])/(3*b^3) + (a*x*\text{Cosh}[a + b*x])/(3*b^2) - (x^2*\text{Cosh}[a + b*x])/(3*b) - (a*\text{Sinh}[a + b*x])/(3*b^3) + (2*x*\text{Sinh}[a + b*x])/(3*b^2) + (a^3*\text{SinhIntegral}[a + b*x])/(3*b^3) + (x^3*\text{SinhIntegral}[a + b*x])/3$

Rule 2717

```
Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]
```

Rule 2718

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ
[{c, d}, x]
```

Rule 3377

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-
(c + d*x)^m)*(Cos[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Co
s[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f
, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 6667

```
Int[((c_.) + (d_.)*(x_))^(m_.)*SinhIntegral[(a_.) + (b_.)*(x_)], x_Symbol]
:= Simp[(c + d*x)^(m + 1)*(SinhIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/
(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Sinh[a + b*x]/(a + b*x)), x], x] /; Fre
eQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{3}x^3\text{Shi}(a + bx) - \frac{1}{3}b \int \frac{x^3 \sinh(a + bx)}{a + bx} dx \\
&= \frac{1}{3}x^3\text{Shi}(a + bx) - \frac{1}{3}b \int \left(\frac{a^2 \sinh(a + bx)}{b^3} - \frac{ax \sinh(a + bx)}{b^2} + \frac{x^2 \sinh(a + bx)}{b} \right. \\
&\quad \left. - \frac{a^3 \sinh(a + bx)}{b^3(a + bx)} \right) dx \\
&= \frac{1}{3}x^3\text{Shi}(a + bx) - \frac{1}{3} \int x^2 \sinh(a + bx) dx - \frac{a^2 \int \sinh(a + bx) dx}{3b^2} \\
&\quad + \frac{a^3 \int \frac{\sinh(a + bx)}{a + bx} dx}{3b^2} + \frac{a \int x \sinh(a + bx) dx}{3b}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{a^2 \cosh(a+bx)}{3b^3} + \frac{ax \cosh(a+bx)}{3b^2} - \frac{x^2 \cosh(a+bx)}{3b} + \frac{a^3 \text{Shi}(a+bx)}{3b^3} \\
&\quad + \frac{1}{3}x^3 \text{Shi}(a+bx) - \frac{a \int \cosh(a+bx) dx}{3b^2} + \frac{2 \int x \cosh(a+bx) dx}{3b} \\
&= -\frac{a^2 \cosh(a+bx)}{3b^3} + \frac{ax \cosh(a+bx)}{3b^2} - \frac{x^2 \cosh(a+bx)}{3b} - \frac{a \sinh(a+bx)}{3b^3} \\
&\quad + \frac{2x \sinh(a+bx)}{3b^2} + \frac{a^3 \text{Shi}(a+bx)}{3b^3} + \frac{1}{3}x^3 \text{Shi}(a+bx) - \frac{2 \int \sinh(a+bx) dx}{3b^2} \\
&= -\frac{2 \cosh(a+bx)}{3b^3} - \frac{a^2 \cosh(a+bx)}{3b^3} + \frac{ax \cosh(a+bx)}{3b^2} - \frac{x^2 \cosh(a+bx)}{3b} \\
&\quad - \frac{a \sinh(a+bx)}{3b^3} + \frac{2x \sinh(a+bx)}{3b^2} + \frac{a^3 \text{Shi}(a+bx)}{3b^3} + \frac{1}{3}x^3 \text{Shi}(a+bx)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.54

$$\begin{aligned}
&\int x^2 \text{Shi}(a+bx) dx \\
&= -\frac{(2+a^2-abx+b^2x^2) \cosh(a+bx) + (a-2bx) \sinh(a+bx) - (a^3+b^3x^3) \text{Shi}(a+bx)}{3b^3}
\end{aligned}$$

[In] Integrate[x^2*SinhIntegral[a + b*x],x]

[Out] -1/3*((2 + a^2 - a*b*x + b^2*x^2)*Cosh[a + b*x] + (a - 2*b*x)*Sinh[a + b*x] - (a^3 + b^3*x^3)*SinhIntegral[a + b*x])/b^3

Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.85

method	result
parts	$\frac{x^3 \text{Shi}(bx+a)}{3} - \frac{-a^3 \text{Shi}(bx+a) + 3a^2 \cosh(bx+a) - 3a((bx+a) \cosh(bx+a) - \sinh(bx+a)) + (bx+a)^2 \cosh(bx+a) - 2(bx+a) \sinh(bx+a)}{3b^3}$
derivativedivides	$\frac{\text{Shi}(bx+a)b^3x^3}{3} + \frac{a^3 \text{Shi}(bx+a)}{3} - a^2 \cosh(bx+a) + a((bx+a) \cosh(bx+a) - \sinh(bx+a)) - \frac{(bx+a)^2 \cosh(bx+a)}{3} + \frac{2(bx+a) \sinh(bx+a)}{3}$
default	$\frac{\text{Shi}(bx+a)b^3x^3}{3} + \frac{a^3 \text{Shi}(bx+a)}{3} - a^2 \cosh(bx+a) + a((bx+a) \cosh(bx+a) - \sinh(bx+a)) - \frac{(bx+a)^2 \cosh(bx+a)}{3} + \frac{2(bx+a) \sinh(bx+a)}{3}$

[In] int(x^2*Shi(b*x+a),x,method=_RETURNVERBOSE)

[Out] 1/3*x^3*Shi(b*x+a)-1/3/b^3*(-a^3*Shi(b*x+a)+3*a^2*cosh(b*x+a)-3*a*((b*x+a)*cosh(b*x+a)-sinh(b*x+a))+(b*x+a)^2*cosh(b*x+a)-2*(b*x+a)*sinh(b*x+a)+2*cosh(b*x+a))

Fricas [F]

$$\int x^2 \operatorname{Shi}(a + bx) dx = \int x^2 \operatorname{Shi}(bx + a) dx$$

[In] `integrate(x^2*Shi(b*x+a),x, algorithm="fricas")`

[Out] `integral(x^2*sinh_integral(b*x + a), x)`

Sympy [F]

$$\int x^2 \operatorname{Shi}(a + bx) dx = \int x^2 \operatorname{Shi}(a + bx) dx$$

[In] `integrate(x**2*Shi(b*x+a),x)`

[Out] `Integral(x**2*Shi(a + b*x), x)`

Maxima [F]

$$\int x^2 \operatorname{Shi}(a + bx) dx = \int x^2 \operatorname{Shi}(bx + a) dx$$

[In] `integrate(x^2*Shi(b*x+a),x, algorithm="maxima")`

[Out] `integrate(x^2*Shi(b*x + a), x)`

Giac [F]

$$\int x^2 \operatorname{Shi}(a + bx) dx = \int x^2 \operatorname{Shi}(bx + a) dx$$

[In] `integrate(x^2*Shi(b*x+a),x, algorithm="giac")`

[Out] `integrate(x^2*Shi(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \text{Shi}(a + bx) dx = \int x^2 \text{sinhint}(a + bx) dx$$

```
[In] int(x^2*sinhint(a + b*x),x)
```

```
[Out] int(x^2*sinhint(a + b*x), x)
```

3.20 $\int x\text{Shi}(a + bx) dx$

Optimal result	139
Rubi [A] (verified)	139
Mathematica [A] (verified)	141
Maple [A] (verified)	141
Fricas [F]	141
Sympy [F]	142
Maxima [F]	142
Giac [F]	142
Mupad [F(-1)]	142

Optimal result

Integrand size = 8, antiderivative size = 71

$$\int x\text{Shi}(a + bx) dx = \frac{a \cosh(a + bx)}{2b^2} - \frac{x \cosh(a + bx)}{2b} + \frac{\sinh(a + bx)}{2b^2} - \frac{a^2 \text{Shi}(a + bx)}{2b^2} + \frac{1}{2} x^2 \text{Shi}(a + bx)$$

[Out] $1/2*a*\cosh(b*x+a)/b^2-1/2*x*\cosh(b*x+a)/b-1/2*a^2*\text{Shi}(b*x+a)/b^2+1/2*x^2*\text{Shi}(b*x+a)+1/2*\sinh(b*x+a)/b^2$

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6667, 6874, 2718, 3377, 2717, 3379}

$$\int x\text{Shi}(a + bx) dx = -\frac{a^2 \text{Shi}(a + bx)}{2b^2} + \frac{\sinh(a + bx)}{2b^2} + \frac{a \cosh(a + bx)}{2b^2} + \frac{1}{2} x^2 \text{Shi}(a + bx) - \frac{x \cosh(a + bx)}{2b}$$

[In] `Int[x*SinhIntegral[a + b*x],x]`

[Out] $(a*\text{Cosh}[a + b*x])/(2*b^2) - (x*\text{Cosh}[a + b*x])/(2*b) + \text{Sinh}[a + b*x]/(2*b^2) - (a^2*\text{SinhIntegral}[a + b*x])/(2*b^2) + (x^2*\text{SinhIntegral}[a + b*x])/2$

Rule 2717

`Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;`
`FreeQ[{c, d}, x]`

Rule 2718

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ
[{c, d}, x]
```

Rule 3377

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(
-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Co
s[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f
, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 6667

```
Int[((c_.) + (d_.)*(x_))^(m_.)*SinhIntegral[(a_.) + (b_.)*(x_)], x_Symbol]
:= Simp[(c + d*x)^(m + 1)*(SinhIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/
(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Sinh[a + b*x]/(a + b*x)), x], x] /; Fre
eQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{2}x^2\text{Shi}(a + bx) - \frac{1}{2}b \int \frac{x^2 \sinh(a + bx)}{a + bx} dx \\
&= \frac{1}{2}x^2\text{Shi}(a + bx) - \frac{1}{2}b \int \left(-\frac{a \sinh(a + bx)}{b^2} + \frac{x \sinh(a + bx)}{b} + \frac{a^2 \sinh(a + bx)}{b^2(a + bx)} \right) dx \\
&= \frac{1}{2}x^2\text{Shi}(a + bx) - \frac{1}{2} \int x \sinh(a + bx) dx + \frac{a \int \sinh(a + bx) dx}{2b} - \frac{a^2 \int \frac{\sinh(a + bx)}{a + bx} dx}{2b} \\
&= \frac{a \cosh(a + bx)}{2b^2} - \frac{x \cosh(a + bx)}{2b} - \frac{a^2\text{Shi}(a + bx)}{2b^2} + \frac{1}{2}x^2\text{Shi}(a + bx) + \frac{\int \cosh(a + bx) dx}{2b} \\
&= \frac{a \cosh(a + bx)}{2b^2} - \frac{x \cosh(a + bx)}{2b} + \frac{\sinh(a + bx)}{2b^2} - \frac{a^2\text{Shi}(a + bx)}{2b^2} + \frac{1}{2}x^2\text{Shi}(a + bx)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.66

$$\int x \operatorname{Shi}(a + bx) dx = \frac{(a - bx) \cosh(a + bx) + \sinh(a + bx) + (-a^2 + b^2 x^2) \operatorname{Shi}(a + bx)}{2b^2}$$

[In] Integrate[x*SinhIntegral[a + b*x],x]

[Out] ((a - b*x)*Cosh[a + b*x] + Sinh[a + b*x] + (-a^2 + b^2*x^2)*SinhIntegral[a + b*x])/(2*b^2)

Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.82

method	result	size
parts	$\frac{x^2 \operatorname{Shi}(bx+a)}{2} - \frac{a^2 \operatorname{Shi}(bx+a) - 2a \cosh(bx+a) + (bx+a) \cosh(bx+a) - \sinh(bx+a)}{2b^2}$	58
derivativedivides	$\frac{\operatorname{Shi}(bx+a) \left(-(bx+a)a + \frac{(bx+a)^2}{2} \right) + a \cosh(bx+a) - \frac{(bx+a) \cosh(bx+a)}{2} + \frac{\sinh(bx+a)}{2}}{b^2}$	60
default	$\frac{\operatorname{Shi}(bx+a) \left(-(bx+a)a + \frac{(bx+a)^2}{2} \right) + a \cosh(bx+a) - \frac{(bx+a) \cosh(bx+a)}{2} + \frac{\sinh(bx+a)}{2}}{b^2}$	60

[In] int(x*Shi(b*x+a),x,method=_RETURNVERBOSE)

[Out] 1/2*x^2*Shi(b*x+a)-1/2/b^2*(a^2*Shi(b*x+a)-2*a*cosh(b*x+a)+(b*x+a)*cosh(b*x+a)-sinh(b*x+a))

Fricas [F]

$$\int x \operatorname{Shi}(a + bx) dx = \int x \operatorname{Shi}(bx + a) dx$$

[In] integrate(x*Shi(b*x+a),x, algorithm="fricas")

[Out] integral(x*sinh_integral(b*x + a), x)

Sympy [F]

$$\int x \operatorname{Shi}(a + bx) dx = \int x \operatorname{Shi}(a + bx) dx$$

```
[In] integrate(x*Shi(b*x+a),x)
```

```
[Out] Integral(x*Shi(a + b*x), x)
```

Maxima [F]

$$\int x \operatorname{Shi}(a + bx) dx = \int x \operatorname{Shi}(bx + a) dx$$

```
[In] integrate(x*Shi(b*x+a),x, algorithm="maxima")
```

```
[Out] integrate(x*Shi(b*x + a), x)
```

Giac [F]

$$\int x \operatorname{Shi}(a + bx) dx = \int x \operatorname{Shi}(bx + a) dx$$

```
[In] integrate(x*Shi(b*x+a),x, algorithm="giac")
```

```
[Out] integrate(x*Shi(b*x + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{Shi}(a + bx) dx = \frac{e^{-a-bx} (a + e^{2a+2bx} + a e^{2a+2bx} - 2a^2 \operatorname{sinhint}(a+bx) e^{a+bx} - 1)}{4} - \frac{b e^{-a-bx} (x + x e^{2a+2bx})}{4} + \frac{x^2 \operatorname{sinhint}(a + bx)}{2}$$

```
[In] int(x*sinhint(a + b*x),x)
```

```
[Out] ((exp(- a - b*x)*(a + exp(2*a + 2*b*x) + a*exp(2*a + 2*b*x) - 2*a^2*sinhint(a + b*x)*exp(a + b*x) - 1))/4 - (b*exp(- a - b*x)*(x + x*exp(2*a + 2*b*x)))/4)/b^2 + (x^2*sinhint(a + b*x))/2
```

3.21 $\int \text{Shi}(a + bx) dx$

Optimal result	143
Rubi [A] (verified)	143
Mathematica [A] (verified)	144
Maple [A] (verified)	144
Fricas [F]	144
Sympy [F]	145
Maxima [F]	145
Giac [F]	145
Mupad [F(-1)]	145

Optimal result

Integrand size = 6, antiderivative size = 27

$$\int \text{Shi}(a + bx) dx = -\frac{\cosh(a + bx)}{b} + \frac{(a + bx)\text{Shi}(a + bx)}{b}$$

[Out] $-\cosh(b*x+a)/b+(b*x+a)*\text{Shi}(b*x+a)/b$

Rubi [A] (verified)

Time = 0.00 (sec), antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6663}

$$\int \text{Shi}(a + bx) dx = \frac{(a + bx)\text{Shi}(a + bx)}{b} - \frac{\cosh(a + bx)}{b}$$

[In] `Int[SinhIntegral[a + b*x], x]`

[Out] $-(\text{Cosh}[a + b*x]/b) + ((a + b*x)*\text{SinhIntegral}[a + b*x])/b$

Rule 6663

`Int[SinhIntegral[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[(a + b*x)*(SinhIntegral[a + b*x]/b), x] - Simp[Cosh[a + b*x]/b, x] /; FreeQ[{a, b}, x]`

Rubi steps

$$\text{integral} = -\frac{\cosh(a + bx)}{b} + \frac{(a + bx)\text{Shi}(a + bx)}{b}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.56

$$\int \text{Shi}(a + bx) dx = -\frac{\cosh(a) \cosh(bx)}{b} - \frac{\sinh(a) \sinh(bx)}{b} + \frac{a \text{Shi}(a + bx)}{b} + x \text{Shi}(a + bx)$$

[In] Integrate[SinhIntegral[a + b*x],x]

[Out] -((Cosh[a]*Cosh[b*x])/b) - (Sinh[a]*Sinh[b*x])/b + (a*SinhIntegral[a + b*x])/b + x*SinhIntegral[a + b*x]

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

method	result	size
derivativedivides	$\frac{\text{Shi}(bx+a)(bx+a) - \cosh(bx+a)}{b}$	26
default	$\frac{\text{Shi}(bx+a)(bx+a) - \cosh(bx+a)}{b}$	26
parts	$x \text{Shi}(bx+a) - \frac{-a \text{Shi}(bx+a) + \cosh(bx+a)}{b}$	31

[In] int(Shi(b*x+a),x,method=_RETURNVERBOSE)

[Out] 1/b*(Shi(b*x+a)*(b*x+a)-cosh(b*x+a))

Fricas [F]

$$\int \text{Shi}(a + bx) dx = \int \text{Shi}(bx + a) dx$$

[In] integrate(Shi(b*x+a),x, algorithm="fricas")

[Out] integral(sinh_integral(b*x + a), x)

Sympy [F]

$$\int \operatorname{Shi}(a + bx) dx = \int \operatorname{Shi}(a + bx) dx$$

[In] `integrate(Shi(b*x+a),x)`

[Out] `Integral(Shi(a + b*x), x)`

Maxima [F]

$$\int \operatorname{Shi}(a + bx) dx = \int \operatorname{Shi}(bx + a) dx$$

[In] `integrate(Shi(b*x+a),x, algorithm="maxima")`

[Out] `integrate(Shi(b*x + a), x)`

Giac [F]

$$\int \operatorname{Shi}(a + bx) dx = \int \operatorname{Shi}(bx + a) dx$$

[In] `integrate(Shi(b*x+a),x, algorithm="giac")`

[Out] `integrate(Shi(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \operatorname{Shi}(a + bx) dx = x \operatorname{sinhint}(a + bx) - \frac{e^{a+bx}}{2b} - \frac{e^{-a-bx}}{2b} + \frac{a \operatorname{sinhint}(a + bx)}{b}$$

[In] `int(sinhint(a + b*x),x)`

[Out] `x*sinhint(a + b*x) - exp(a + b*x)/(2*b) - exp(- a - b*x)/(2*b) + (a*sinhint(a + b*x))/b`

3.22 $\int \frac{\mathbf{Shi}(a+bx)}{x} dx$

Optimal result	146
Rubi [N/A]	146
Mathematica [N/A]	147
Maple [N/A] (verified)	147
Fricas [N/A]	147
Sympy [N/A]	147
Maxima [N/A]	148
Giac [N/A]	148
Mupad [N/A]	148

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\mathbf{Shi}(a+bx)}{x} dx = \text{Int}\left(\frac{\mathbf{Shi}(a+bx)}{x}, x\right)$$

[Out] CannotIntegrate(Shi(b*x+a)/x,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\mathbf{Shi}(a+bx)}{x} dx = \int \frac{\mathbf{Shi}(a+bx)}{x} dx$$

[In] Int[SinhIntegral[a + b*x]/x,x]

[Out] Defer[Int][SinhIntegral[a + b*x]/x, x]

Rubi steps

$$\text{integral} = \int \frac{\mathbf{Shi}(a+bx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Shi}(a + bx)}{x} dx = \int \frac{\text{Shi}(a + bx)}{x} dx$$

[In] Integrate[SinhIntegral[a + b*x]/x,x]

[Out] Integrate[SinhIntegral[a + b*x]/x, x]

Maple [N/A] (verified)

Not integrable

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\text{Shi}(bx + a)}{x} dx$$

[In] int(Shi(b*x+a)/x,x)

[Out] int(Shi(b*x+a)/x,x)

Fricas [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Shi}(a + bx)}{x} dx = \int \frac{\text{Shi}(bx + a)}{x} dx$$

[In] integrate(Shi(b*x+a)/x,x, algorithm="fricas")

[Out] integral(sinh_integral(b*x + a)/x, x)

Sympy [N/A]

Not integrable

Time = 0.39 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\text{Shi}(a + bx)}{x} dx = \int \frac{\text{Shi}(a + bx)}{x} dx$$

[In] integrate(Shi(b*x+a)/x,x)

[Out] Integral(Shi(a + b*x)/x, x)

Maxima [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\operatorname{Shi}(a + bx)}{x} dx = \int \frac{\operatorname{Shi}(bx + a)}{x} dx$$

[In] integrate(Shi(b*x+a)/x,x, algorithm="maxima")

[Out] integrate(Shi(b*x + a)/x, x)

Giac [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\operatorname{Shi}(a + bx)}{x} dx = \int \frac{\operatorname{Shi}(bx + a)}{x} dx$$

[In] integrate(Shi(b*x+a)/x,x, algorithm="giac")

[Out] integrate(Shi(b*x + a)/x, x)

Mupad [N/A]

Not integrable

Time = 4.88 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\operatorname{Shi}(a + bx)}{x} dx = \int \frac{\operatorname{sinhint}(a + bx)}{x} dx$$

[In] int(sinhint(a + b*x)/x,x)

[Out] int(sinhint(a + b*x)/x, x)

3.23 $\int \frac{\text{Shi}(a+bx)}{x^2} dx$

Optimal result	149
Rubi [A] (verified)	149
Mathematica [A] (verified)	151
Maple [F]	151
Fricas [F]	151
Sympy [F]	151
Maxima [F]	152
Giac [F]	152
Mupad [F(-1)]	152

Optimal result

Integrand size = 10, antiderivative size = 46

$$\int \frac{\text{Shi}(a+bx)}{x^2} dx = \frac{b\text{Chi}(bx)\sinh(a)}{a} + \frac{b\cosh(a)\text{Shi}(bx)}{a} - \frac{b\text{Shi}(a+bx)}{a} - \frac{\text{Shi}(a+bx)}{x}$$

[Out] $b*\cosh(a)*\text{Shi}(b*x)/a - b*\text{Shi}(b*x+a)/a - \text{Shi}(b*x+a)/x + b*\text{Chi}(b*x)*\sinh(a)/a$

Rubi [A] (verified)

Time = 0.21 (sec), antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6667, 6874, 3384, 3379, 3382}

$$\int \frac{\text{Shi}(a+bx)}{x^2} dx = \frac{b\sinh(a)\text{Chi}(bx)}{a} - \frac{b\text{Shi}(a+bx)}{a} - \frac{\text{Shi}(a+bx)}{x} + \frac{b\cosh(a)\text{Shi}(bx)}{a}$$

[In] `Int[SinhIntegral[a + b*x]/x^2, x]`

[Out] $(b*\text{CoshIntegral}[b*x]*\text{Sinh}[a])/a + (b*\text{Cosh}[a]*\text{SinhIntegral}[b*x])/a - (b*\text{SinhIntegral}[a + b*x])/a - \text{SinhIntegral}[a + b*x]/x$

Rule 3379

`Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`

Rule 3382

`Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}`

}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3384

Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]

Rule 6667

Int[((c_.) + (d_.)*(x_.))^(m_.)*SinhIntegral[(a_.) + (b_.)*(x_.)], x_Symbol] := Simp[(c + d*x)^(m + 1)*(SinhIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Sinh[a + b*x]/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]

Rule 6874

Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Shi}(a + bx)}{x} + b \int \frac{\sinh(a + bx)}{x(a + bx)} dx \\
 &= -\frac{\text{Shi}(a + bx)}{x} + b \int \left(\frac{\sinh(a + bx)}{ax} - \frac{b \sinh(a + bx)}{a(a + bx)} \right) dx \\
 &= -\frac{\text{Shi}(a + bx)}{x} + \frac{b \int \frac{\sinh(a+bx)}{x} dx}{a} - \frac{b^2 \int \frac{\sinh(a+bx)}{a+bx} dx}{a} \\
 &= -\frac{b \text{Shi}(a + bx)}{a} - \frac{\text{Shi}(a + bx)}{x} + \frac{(b \cosh(a)) \int \frac{\sinh(bx)}{x} dx}{a} + \frac{(b \sinh(a)) \int \frac{\cosh(bx)}{x} dx}{a} \\
 &= \frac{b \text{Chi}(bx) \sinh(a)}{a} + \frac{b \cosh(a) \text{Shi}(bx)}{a} - \frac{b \text{Shi}(a + bx)}{a} - \frac{\text{Shi}(a + bx)}{x}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.85

$$\int \frac{\text{Shi}(a + bx)}{x^2} dx = \frac{bx\text{Chi}(bx) \sinh(a) + bx \cosh(a)\text{Shi}(bx) - (a + bx)\text{Shi}(a + bx)}{ax}$$

[In] Integrate[SinhIntegral[a + b*x]/x^2,x]

[Out] (b*x*CoshIntegral[b*x]*Sinh[a] + b*x*Cosh[a]*SinhIntegral[b*x] - (a + b*x)*SinhIntegral[a + b*x])/(a*x)

Maple [F]

$$\int \frac{\text{Shi}(bx + a)}{x^2} dx$$

[In] int(Shi(b*x+a)/x^2,x)

[Out] int(Shi(b*x+a)/x^2,x)

Fricas [F]

$$\int \frac{\text{Shi}(a + bx)}{x^2} dx = \int \frac{\text{Shi}(bx + a)}{x^2} dx$$

[In] integrate(Shi(b*x+a)/x^2,x, algorithm="fricas")

[Out] integral(sinh_integral(b*x + a)/x^2, x)

Sympy [F]

$$\int \frac{\text{Shi}(a + bx)}{x^2} dx = \int \frac{\text{Shi}(a + bx)}{x^2} dx$$

[In] integrate(Shi(b*x+a)/x**2,x)

[Out] Integral(Shi(a + b*x)/x**2, x)

Maxima [F]

$$\int \frac{\text{Shi}(a + bx)}{x^2} dx = \int \frac{\text{Shi}(bx + a)}{x^2} dx$$

[In] integrate(Shi(b*x+a)/x^2,x, algorithm="maxima")

[Out] integrate(Shi(b*x + a)/x^2, x)

Giac [F]

$$\int \frac{\text{Shi}(a + bx)}{x^2} dx = \int \frac{\text{Shi}(bx + a)}{x^2} dx$$

[In] integrate(Shi(b*x+a)/x^2,x, algorithm="giac")

[Out] integrate(Shi(b*x + a)/x^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\text{Shi}(a + bx)}{x^2} dx = \int \frac{\text{sinhint}(a + bx)}{x^2} dx$$

[In] int(sinhint(a + b*x)/x^2,x)

[Out] int(sinhint(a + b*x)/x^2, x)

3.24 $\int \frac{\text{Shi}(a+bx)}{x^3} dx$

Optimal result	153
Rubi [A] (verified)	153
Mathematica [A] (verified)	155
Maple [F]	155
Fricas [F]	156
Sympy [F]	156
Maxima [F]	156
Giac [F]	156
Mupad [F(-1)]	157

Optimal result

Integrand size = 10, antiderivative size = 111

$$\int \frac{\text{Shi}(a+bx)}{x^3} dx = \frac{b^2 \cosh(a) \text{Chi}(bx)}{2a} - \frac{b^2 \text{Chi}(bx) \sinh(a)}{2a^2} - \frac{b \sinh(a+bx)}{2ax} - \frac{b^2 \cosh(a) \text{Shi}(bx)}{2a^2} + \frac{b^2 \sinh(a) \text{Shi}(bx)}{2a} + \frac{b^2 \text{Shi}(a+bx)}{2a^2} - \frac{\text{Shi}(a+bx)}{2x^2}$$

[Out] $1/2*b^2*\text{Chi}(b*x)*\cosh(a)/a-1/2*b^2*\cosh(a)*\text{Shi}(b*x)/a^2+1/2*b^2*\text{Shi}(b*x+a)/a^2-1/2*\text{Shi}(b*x+a)/x^2-1/2*b^2*\text{Chi}(b*x)*\sinh(a)/a^2+1/2*b^2*\text{Shi}(b*x)*\sinh(a)/a-1/2*b*\sinh(b*x+a)/a/x$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6667, 6874, 3378, 3384, 3379, 3382}

$$\int \frac{\text{Shi}(a+bx)}{x^3} dx = -\frac{b^2 \sinh(a) \text{Chi}(bx)}{2a^2} + \frac{b^2 \text{Shi}(a+bx)}{2a^2} - \frac{b^2 \cosh(a) \text{Shi}(bx)}{2a^2} + \frac{b^2 \cosh(a) \text{Chi}(bx)}{2a} + \frac{b^2 \sinh(a) \text{Shi}(bx)}{2a} - \frac{\text{Shi}(a+bx)}{2x^2} - \frac{b \sinh(a+bx)}{2ax}$$

[In] Int[SinhIntegral[a + b*x]/x^3,x]

[Out] $(b^2*\text{Cosh}[a]*\text{CoshIntegral}[b*x])/(2*a) - (b^2*\text{CoshIntegral}[b*x]*\text{Sinh}[a])/(2*a^2) - (b*\text{Sinh}[a + b*x])/(2*a*x) - (b^2*\text{Cosh}[a]*\text{SinhIntegral}[b*x])/(2*a^2) + (b^2*\text{Sinh}[a]*\text{SinhIntegral}[b*x])/(2*a) + (b^2*\text{SinhIntegral}[a + b*x])/(2*a^2) - \text{SinhIntegral}[a + b*x]/(2*x^2)$

Rule 3378

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c
+ d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Dist[f/(d*(m + 1)), Int[(c
+ d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1
]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f
, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz
}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3384

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*
e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f
)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] &&
NeQ[d*e - c*f, 0]
```

Rule 6667

```
Int[((c_.) + (d_.)*(x_))^(m_)*SinhIntegral[(a_.) + (b_.)*(x_)], x_Symbol]
:= Simp[(c + d*x)^(m + 1)*(SinhIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/
(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Sinh[a + b*x]/(a + b*x)), x], x] /; Fre
eQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Shi}(a + bx)}{2x^2} + \frac{1}{2}b \int \frac{\sinh(a + bx)}{x^2(a + bx)} dx \\
&= -\frac{\text{Shi}(a + bx)}{2x^2} + \frac{1}{2}b \int \left(\frac{\sinh(a + bx)}{ax^2} - \frac{b \sinh(a + bx)}{a^2x} + \frac{b^2 \sinh(a + bx)}{a^2(a + bx)} \right) dx \\
&= -\frac{\text{Shi}(a + bx)}{2x^2} + \frac{b \int \frac{\sinh(a + bx)}{x^2} dx}{2a} - \frac{b^2 \int \frac{\sinh(a + bx)}{x} dx}{2a^2} + \frac{b^3 \int \frac{\sinh(a + bx)}{a + bx} dx}{2a^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{b \sinh(a+bx)}{2ax} + \frac{b^2 \text{Shi}(a+bx)}{2a^2} - \frac{\text{Shi}(a+bx)}{2x^2} + \frac{b^2 \int \frac{\cosh(a+bx)}{x} dx}{2a} \\
&\quad - \frac{(b^2 \cosh(a)) \int \frac{\sinh(bx)}{x} dx}{2a^2} - \frac{(b^2 \sinh(a)) \int \frac{\cosh(bx)}{x} dx}{2a^2} \\
&= -\frac{b^2 \text{Chi}(bx) \sinh(a)}{2a^2} - \frac{b \sinh(a+bx)}{2ax} - \frac{b^2 \cosh(a) \text{Shi}(bx)}{2a^2} + \frac{b^2 \text{Shi}(a+bx)}{2a^2} \\
&\quad - \frac{\text{Shi}(a+bx)}{2x^2} + \frac{(b^2 \cosh(a)) \int \frac{\cosh(bx)}{x} dx}{2a} + \frac{(b^2 \sinh(a)) \int \frac{\sinh(bx)}{x} dx}{2a} \\
&= \frac{b^2 \cosh(a) \text{Chi}(bx)}{2a} - \frac{b^2 \text{Chi}(bx) \sinh(a)}{2a^2} - \frac{b \sinh(a+bx)}{2ax} \\
&\quad - \frac{b^2 \cosh(a) \text{Shi}(bx)}{2a^2} + \frac{b^2 \sinh(a) \text{Shi}(bx)}{2a} + \frac{b^2 \text{Shi}(a+bx)}{2a^2} - \frac{\text{Shi}(a+bx)}{2x^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.77

$$\int \frac{\text{Shi}(a+bx)}{x^3} dx = \frac{b^2 x^2 \text{Chi}(bx)(a \cosh(a) - \sinh(a)) - abx \sinh(a+bx) + b^2 x^2 (-\cosh(a) + a \sinh(a)) \text{Shi}(bx) - a^2 \text{Shi}(a+bx)}{2a^2 x^2}$$

[In] Integrate[SinhIntegral[a + b*x]/x^3,x]

[Out] (b^2*x^2*CoshIntegral[b*x]*(a*Cosh[a] - Sinh[a]) - a*b*x*Sinh[a + b*x] + b^2*x^2*(-Cosh[a] + a*Sinh[a])*SinhIntegral[b*x] - a^2*SinhIntegral[a + b*x] + b^2*x^2*SinhIntegral[a + b*x])/(2*a^2*x^2)

Maple [F]

$$\int \frac{\text{Shi}(bx+a)}{x^3} dx$$

[In] int(Shi(b*x+a)/x^3,x)

[Out] int(Shi(b*x+a)/x^3,x)

Fricas [F]

$$\int \frac{\operatorname{Shi}(a + bx)}{x^3} dx = \int \frac{\operatorname{Shi}(bx + a)}{x^3} dx$$

[In] integrate(Shi(b*x+a)/x^3,x, algorithm="fricas")

[Out] integral(sinh_integral(b*x + a)/x^3, x)

Sympy [F]

$$\int \frac{\operatorname{Shi}(a + bx)}{x^3} dx = \int \frac{\operatorname{Shi}(a + bx)}{x^3} dx$$

[In] integrate(Shi(b*x+a)/x**3,x)

[Out] Integral(Shi(a + b*x)/x**3, x)

Maxima [F]

$$\int \frac{\operatorname{Shi}(a + bx)}{x^3} dx = \int \frac{\operatorname{Shi}(bx + a)}{x^3} dx$$

[In] integrate(Shi(b*x+a)/x^3,x, algorithm="maxima")

[Out] integrate(Shi(b*x + a)/x^3, x)

Giac [F]

$$\int \frac{\operatorname{Shi}(a + bx)}{x^3} dx = \int \frac{\operatorname{Shi}(bx + a)}{x^3} dx$$

[In] integrate(Shi(b*x+a)/x^3,x, algorithm="giac")

[Out] integrate(Shi(b*x + a)/x^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\text{Shi}(a + bx)}{x^3} dx = \int \frac{\text{sinhint}(a + bx)}{x^3} dx$$

```
[In] int(sinhint(a + b*x)/x^3,x)
```

```
[Out] int(sinhint(a + b*x)/x^3, x)
```

3.25 $\int x^m \mathbf{Shi}(a + bx)^2 dx$

Optimal result	158
Rubi [N/A]	158
Mathematica [N/A]	159
Maple [N/A] (verified)	159
Fricas [N/A]	159
Sympy [N/A]	159
Maxima [N/A]	160
Giac [N/A]	160
Mupad [N/A]	160

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int x^m \mathbf{Shi}(a + bx)^2 dx = \text{Int}(x^m \mathbf{Shi}(a + bx)^2, x)$$

[Out] CannotIntegrate(x^m*Shi(b*x+a)²,x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int x^m \mathbf{Shi}(a + bx)^2 dx = \int x^m \mathbf{Shi}(a + bx)^2 dx$$

[In] Int[x^m*SinhIntegral[a + b*x]²,x]

[Out] Defer[Int][x^m*SinhIntegral[a + b*x]², x]

Rubi steps

$$\text{integral} = \int x^m \mathbf{Shi}(a + bx)^2 dx$$

Mathematica [N/A]

Not integrable

Time = 4.58 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \operatorname{Shi}(a + bx)^2 dx = \int x^m \operatorname{Shi}(a + bx)^2 dx$$

[In] Integrate[x^m*SinhIntegral[a + b*x]^2,x]

[Out] Integrate[x^m*SinhIntegral[a + b*x]^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.22 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int x^m \operatorname{Shi}(bx + a)^2 dx$$

[In] int(x^m*Shi(b*x+a)^2,x)

[Out] int(x^m*Shi(b*x+a)^2,x)

Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \operatorname{Shi}(a + bx)^2 dx = \int x^m \operatorname{Shi}(bx + a)^2 dx$$

[In] integrate(x^m*Shi(b*x+a)^2,x, algorithm="fricas")

[Out] integral(x^m*sinh_integral(b*x + a)^2, x)

Sympy [N/A]

Not integrable

Time = 1.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int x^m \operatorname{Shi}(a + bx)^2 dx = \int x^m \operatorname{Shi}^2(a + bx) dx$$

[In] integrate(x**m*Shi(b*x+a)**2,x)

[Out] Integral(x**m*Shi(a + b*x)**2, x)

Maxima [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \operatorname{Shi}(a + bx)^2 dx = \int x^m \operatorname{Shi}(bx + a)^2 dx$$

[In] integrate(x^m*Shi(b*x+a)^2,x, algorithm="maxima")

[Out] integrate(x^m*Shi(b*x + a)^2, x)

Giac [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \operatorname{Shi}(a + bx)^2 dx = \int x^m \operatorname{Shi}(bx + a)^2 dx$$

[In] integrate(x^m*Shi(b*x+a)^2,x, algorithm="giac")

[Out] integrate(x^m*Shi(b*x + a)^2, x)

Mupad [N/A]

Not integrable

Time = 4.90 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \operatorname{Shi}(a + bx)^2 dx = \int x^m \operatorname{sinhint}(a + bx)^2 dx$$

[In] int(x^m*sinhint(a + b*x)^2,x)

[Out] int(x^m*sinhint(a + b*x)^2, x)

3.26 $\int x^2 \text{Shi}(a + bx)^2 dx$

Optimal result	161
Rubi [A] (verified)	162
Mathematica [A] (verified)	167
Maple [F]	168
Fricas [F]	168
Sympy [F]	168
Maxima [F]	168
Giac [F]	169
Mupad [F(-1)]	169

Optimal result

Integrand size = 12, antiderivative size = 328

$$\int x^2 \text{Shi}(a + bx)^2 dx = \frac{2x}{3b^2} - \frac{a \cosh(2a + 2bx)}{3b^3} + \frac{x \cosh(2a + 2bx)}{6b^2} + \frac{a \text{Chi}(2a + 2bx)}{b^3} - \frac{a \log(a + bx)}{b^3} - \frac{2 \cosh(a + bx) \sinh(a + bx)}{3b^3} - \frac{\sinh(2a + 2bx)}{12b^3} - \frac{4 \cosh(a + bx) \text{Shi}(a + bx)}{3b^3} - \frac{2a^2 \cosh(a + bx) \text{Shi}(a + bx)}{3b^3} + \frac{2ax \cosh(a + bx) \text{Shi}(a + bx)}{3b^2} - \frac{2x^2 \cosh(a + bx) \text{Shi}(a + bx)}{3b} - \frac{2a \sinh(a + bx) \text{Shi}(a + bx)}{3b^3} + \frac{4x \sinh(a + bx) \text{Shi}(a + bx)}{3b^2} + \frac{a^2(a + bx) \text{Shi}(a + bx)^2}{3b^3} - \frac{ax(a + bx) \text{Shi}(a + bx)^2}{3b^2} + \frac{x^2(a + bx) \text{Shi}(a + bx)^2}{3b} + \frac{2 \text{Shi}(2a + 2bx)}{3b^3} + \frac{a^2 \text{Shi}(2a + 2bx)}{b^3}$$

[Out] $\frac{2}{3} \frac{x}{b^2} + \frac{a \text{Chi}(2bx + 2a)}{b^3} - \frac{1}{3} \frac{a \cosh(2bx + 2a)}{b^3} + \frac{1}{6} \frac{x \cosh(2bx + 2a)}{b^2} + \frac{2bx + 2a}{b^2} \ln(bx + a) - \frac{4}{3} \frac{\cosh(bx + a) \text{Shi}(bx + a)}{b^3} - \frac{2}{3} \frac{a^2 \cosh(bx + a) \text{Shi}(bx + a)}{b^3} + \frac{2}{3} \frac{ax \cosh(bx + a) \text{Shi}(bx + a)}{b^2} - \frac{2}{3} \frac{x^2 \cosh(bx + a) \text{Shi}(bx + a)}{b^2} + \frac{1}{3} \frac{a^2 (bx + a) \text{Shi}(bx + a)^2}{b^3} - \frac{1}{3} \frac{ax (bx + a) \text{Shi}(bx + a)^2}{b^2} + \frac{1}{3} \frac{x^2 (bx + a) \text{Shi}(bx + a)^2}{b^2} + \frac{2 \text{Shi}(2bx + 2a)}{3b^3} + \frac{a^2 \text{Shi}(2bx + 2a)}{b^3} - \frac{2}{3} \frac{\cosh(bx + a) \sinh(bx + a)}{b^3} - \frac{2}{3} \frac{a \text{Shi}(bx + a) \sinh(bx + a)}{b^3} + \frac{4}{3} \frac{x \text{Shi}(bx + a) \sinh(bx + a)}{b^2} - \frac{1}{12} \frac{\sinh(2bx + 2a)}{b^3}$

Rubi [A] (verified)

Time = 1.17 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.00, number of steps used = 39, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 1.583$, Rules used = {6673, 6677, 5736, 6873, 6874, 2718, 3377, 2717, 3379, 6683, 2715, 8, 3393, 3382, 6675, 5556, 12, 6681, 6669}

$$\int x^2 \text{Shi}(a + bx)^2 dx = \frac{a^2(a + bx)\text{Shi}(a + bx)^2}{3b^3} + \frac{a^2\text{Shi}(2a + 2bx)}{b^3} - \frac{2a^2\text{Shi}(a + bx)\cosh(a + bx)}{3b^3} + \frac{a\text{Chi}(2a + 2bx)}{b^3} + \frac{2\text{Shi}(2a + 2bx)}{3b^3} - \frac{2a\text{Shi}(a + bx)\sinh(a + bx)}{3b^3} - \frac{4\text{Shi}(a + bx)\cosh(a + bx)}{3b^3} - \frac{a\log(a + bx)}{b^3} - \frac{\sinh(2a + 2bx)}{12b^3} - \frac{a\cosh(2a + 2bx)}{3b^3} - \frac{2\sinh(a + bx)\cosh(a + bx)}{3b^3} - \frac{ax(a + bx)\text{Shi}(a + bx)^2}{3b^2} + \frac{4x\text{Shi}(a + bx)\sinh(a + bx)}{3b^2} + \frac{2ax\text{Shi}(a + bx)\cosh(a + bx)}{3b^2} + \frac{x\cosh(2a + 2bx)}{6b^2} + \frac{x^2(a + bx)\text{Shi}(a + bx)^2}{3b} - \frac{2x^2\text{Shi}(a + bx)\cosh(a + bx)}{3b} + \frac{2x}{3b^2}$$

[In] Int[x^2*SinhIntegral[a + b*x]^2,x]

[Out] (2*x)/(3*b^2) - (a*Cosh[2*a + 2*b*x])/(3*b^3) + (x*Cosh[2*a + 2*b*x])/(6*b^2) + (a*CoshIntegral[2*a + 2*b*x])/b^3 - (a*Log[a + b*x])/b^3 - (2*Cosh[a + b*x]*Sinh[a + b*x])/(3*b^3) - Sinh[2*a + 2*b*x]/(12*b^3) - (4*Cosh[a + b*x]*SinhIntegral[a + b*x])/(3*b^3) - (2*a^2*Cosh[a + b*x]*SinhIntegral[a + b*x])/(3*b^3) + (2*a*x*Cosh[a + b*x]*SinhIntegral[a + b*x])/(3*b^2) - (2*x^2*Cosh[a + b*x]*SinhIntegral[a + b*x])/(3*b) - (2*a*Sinh[a + b*x]*SinhIntegral[a + b*x])/(3*b^3) + (4*x*Sinh[a + b*x]*SinhIntegral[a + b*x])/(3*b^2) + (a^2*(a + b*x)*SinhIntegral[a + b*x]^2)/(3*b^3) - (a*x*(a + b*x)*SinhIntegral[a + b*x]^2)/(3*b^2) + (x^2*(a + b*x)*SinhIntegral[a + b*x]^2)/(3*b) + (2*SinhIntegral[2*a + 2*b*x])/(3*b^3) + (a^2*SinhIntegral[2*a + 2*b*x])/b^3

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 2717

```
Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]
```

Rule 2718

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]
```

Rule 3377

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3393

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]
```

Rule 5736

$\text{Int}[\text{Cosh}[w_]^{(p_)} \cdot (u_) \cdot \text{Sinh}[v_]^{(p_)}, x_ \text{Symbol}] \text{ :> } \text{Dist}[1/2^p, \text{Int}[u \cdot \text{Sinh}[2 \cdot v]^p, x], x] \text{ /; } \text{EqQ}[w, v] \ \&\& \ \text{IntegerQ}[p]$

Rule 6669

$\text{Int}[\text{SinhIntegral}[(a_) + (b_) \cdot (x_)]^2, x_ \text{Symbol}] \text{ :> } \text{Simp}[(a + b \cdot x) \cdot (\text{SinhIntegral}[a + b \cdot x]^2/b), x] - \text{Dist}[2, \text{Int}[\text{Sinh}[a + b \cdot x] \cdot \text{SinhIntegral}[a + b \cdot x], x], x] \text{ /; } \text{FreeQ}[\{a, b\}, x]$

Rule 6673

$\text{Int}[((c_) + (d_) \cdot (x_))^{(m_)} \cdot \text{SinhIntegral}[(a_) + (b_) \cdot (x_)]^2, x_ \text{Symbol}] \text{ :> } \text{Simp}[(a + b \cdot x) \cdot (c + d \cdot x)^m \cdot (\text{SinhIntegral}[a + b \cdot x]^2/(b \cdot (m + 1))), x] + (-\text{Dist}[2/(m + 1), \text{Int}[(c + d \cdot x)^m \cdot \text{Sinh}[a + b \cdot x] \cdot \text{SinhIntegral}[a + b \cdot x], x], x] + \text{Dist}[(b \cdot c - a \cdot d) \cdot (m/(b \cdot (m + 1))), \text{Int}[(c + d \cdot x)^{(m - 1)} \cdot \text{SinhIntegral}[a + b \cdot x]^2, x], x]) \text{ /; } \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{IGtQ}[m, 0]$

Rule 6675

$\text{Int}[\text{Sinh}[(a_) + (b_) \cdot (x_)] \cdot \text{SinhIntegral}[(c_) + (d_) \cdot (x_)], x_ \text{Symbol}] \text{ :> } \text{Simp}[\text{Cosh}[a + b \cdot x] \cdot (\text{SinhIntegral}[c + d \cdot x]/b), x] - \text{Dist}[d/b, \text{Int}[\text{Cosh}[a + b \cdot x] \cdot (\text{Sinh}[c + d \cdot x]/(c + d \cdot x)), x], x] \text{ /; } \text{FreeQ}[\{a, b, c, d\}, x]$

Rule 6677

$\text{Int}[((e_) + (f_) \cdot (x_))^{(m_)} \cdot \text{Sinh}[(a_) + (b_) \cdot (x_)] \cdot \text{SinhIntegral}[(c_) + (d_) \cdot (x_)], x_ \text{Symbol}] \text{ :> } \text{Simp}[(e + f \cdot x)^m \cdot \text{Cosh}[a + b \cdot x] \cdot (\text{SinhIntegral}[c + d \cdot x]/b), x] + (-\text{Dist}[d/b, \text{Int}[(e + f \cdot x)^m \cdot \text{Cosh}[a + b \cdot x] \cdot (\text{Sinh}[c + d \cdot x]/(c + d \cdot x)), x], x] - \text{Dist}[f \cdot (m/b), \text{Int}[(e + f \cdot x)^{(m - 1)} \cdot \text{Cosh}[a + b \cdot x] \cdot \text{SinhIntegral}[c + d \cdot x], x], x]) \text{ /; } \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{IGtQ}[m, 0]$

Rule 6681

$\text{Int}[\text{Cosh}[(a_) + (b_) \cdot (x_)] \cdot \text{SinhIntegral}[(c_) + (d_) \cdot (x_)], x_ \text{Symbol}] \text{ :> } \text{Simp}[\text{Sinh}[a + b \cdot x] \cdot (\text{SinhIntegral}[c + d \cdot x]/b), x] - \text{Dist}[d/b, \text{Int}[\text{Sinh}[a + b \cdot x] \cdot (\text{Sinh}[c + d \cdot x]/(c + d \cdot x)), x], x] \text{ /; } \text{FreeQ}[\{a, b, c, d\}, x]$

Rule 6683

$\text{Int}[\text{Cosh}[(a_) + (b_) \cdot (x_)] \cdot ((e_) + (f_) \cdot (x_))^{(m_)} \cdot \text{SinhIntegral}[(c_) + (d_) \cdot (x_)], x_ \text{Symbol}] \text{ :> } \text{Simp}[(e + f \cdot x)^m \cdot \text{Sinh}[a + b \cdot x] \cdot (\text{SinhIntegral}[c + d \cdot x]/b), x] + (-\text{Dist}[d/b, \text{Int}[(e + f \cdot x)^m \cdot \text{Sinh}[a + b \cdot x] \cdot (\text{Sinh}[c + d \cdot x]/(c + d \cdot x)), x], x] - \text{Dist}[f \cdot (m/b), \text{Int}[(e + f \cdot x)^{(m - 1)} \cdot \text{Sinh}[a + b \cdot x] \cdot \text{SinhIntegral}[c + d \cdot x], x], x]) \text{ /; } \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{IGtQ}[m, 0]$

Rule 6873

Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v != u]

Rule 6874

Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x^2(a+bx)\text{Shi}(a+bx)^2}{3b} - \frac{2}{3} \int x^2 \sinh(a+bx)\text{Shi}(a+bx) dx \\
 &\quad - \frac{(2a) \int x\text{Shi}(a+bx)^2 dx}{3b} \\
 &= -\frac{2x^2 \cosh(a+bx)\text{Shi}(a+bx)}{3b} - \frac{ax(a+bx)\text{Shi}(a+bx)^2}{3b^2} + \frac{x^2(a+bx)\text{Shi}(a+bx)^2}{3b} \\
 &\quad + \frac{2}{3} \int \frac{x^2 \cosh(a+bx) \sinh(a+bx)}{a+bx} dx + \frac{a^2 \int \text{Shi}(a+bx)^2 dx}{3b^2} \\
 &\quad + \frac{4 \int x \cosh(a+bx)\text{Shi}(a+bx) dx}{3b} + \frac{(2a) \int x \sinh(a+bx)\text{Shi}(a+bx) dx}{3b} \\
 &= \frac{2ax \cosh(a+bx)\text{Shi}(a+bx)}{3b^2} - \frac{2x^2 \cosh(a+bx)\text{Shi}(a+bx)}{3b} \\
 &\quad + \frac{4x \sinh(a+bx)\text{Shi}(a+bx)}{3b^2} + \frac{a^2(a+bx)\text{Shi}(a+bx)^2}{3b^3} \\
 &\quad - \frac{ax(a+bx)\text{Shi}(a+bx)^2}{3b^2} + \frac{x^2(a+bx)\text{Shi}(a+bx)^2}{3b} \\
 &\quad + \frac{1}{3} \int \frac{x^2 \sinh(2(a+bx))}{a+bx} dx - \frac{4 \int \sinh(a+bx)\text{Shi}(a+bx) dx}{3b^2} \\
 &\quad - \frac{(2a) \int \cosh(a+bx)\text{Shi}(a+bx) dx}{3b^2} - \frac{(2a^2) \int \sinh(a+bx)\text{Shi}(a+bx) dx}{3b^2} \\
 &\quad - \frac{4 \int \frac{x \sinh^2(a+bx)}{a+bx} dx}{3b} - \frac{(2a) \int \frac{x \cosh(a+bx) \sinh(a+bx)}{a+bx} dx}{3b}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{4 \cosh(a+bx)\text{Shi}(a+bx)}{3b^3} - \frac{2a^2 \cosh(a+bx)\text{Shi}(a+bx)}{3b^3} + \frac{2ax \cosh(a+bx)\text{Shi}(a+bx)}{3b^2} \\
&\quad - \frac{2x^2 \cosh(a+bx)\text{Shi}(a+bx)}{3b} - \frac{2a \sinh(a+bx)\text{Shi}(a+bx)}{3b^3} + \frac{4x \sinh(a+bx)\text{Shi}(a+bx)}{3b^2} \\
&\quad + \frac{a^2(a+bx)\text{Shi}(a+bx)^2}{3b^3} - \frac{ax(a+bx)\text{Shi}(a+bx)^2}{3b^2} + \frac{x^2(a+bx)\text{Shi}(a+bx)^2}{3b} \\
&\quad + \frac{1}{3} \int \frac{x^2 \sinh(2a+2bx)}{a+bx} dx + \frac{4 \int \frac{\cosh(a+bx) \sinh(a+bx)}{a+bx} dx}{3b^2} + \frac{(2a) \int \frac{\sinh^2(a+bx)}{a+bx} dx}{3b^2} \\
&\quad + \frac{(2a^2) \int \frac{\cosh(a+bx) \sinh(a+bx)}{a+bx} dx}{3b^2} - \frac{4 \int \left(\frac{\sinh^2(a+bx)}{b} - \frac{a \sinh^2(a+bx)}{b(a+bx)} \right) dx}{3b} \\
&\quad - \frac{a \int \frac{x \sinh(2(a+bx))}{a+bx} dx}{3b} \\
&= -\frac{4 \cosh(a+bx)\text{Shi}(a+bx)}{3b^3} - \frac{2a^2 \cosh(a+bx)\text{Shi}(a+bx)}{3b^3} \\
&\quad + \frac{2ax \cosh(a+bx)\text{Shi}(a+bx)}{3b^2} - \frac{2x^2 \cosh(a+bx)\text{Shi}(a+bx)}{3b} \\
&\quad - \frac{2a \sinh(a+bx)\text{Shi}(a+bx)}{3b^3} + \frac{4x \sinh(a+bx)\text{Shi}(a+bx)}{3b^2} \\
&\quad + \frac{a^2(a+bx)\text{Shi}(a+bx)^2}{3b^3} - \frac{ax(a+bx)\text{Shi}(a+bx)^2}{3b^2} + \frac{x^2(a+bx)\text{Shi}(a+bx)^2}{3b} \\
&\quad + \frac{1}{3} \int \left(-\frac{a \sinh(2a+2bx)}{b^2} + \frac{x \sinh(2a+2bx)}{b} + \frac{a^2 \sinh(2a+2bx)}{b^2(a+bx)} \right) dx \\
&\quad - \frac{4 \int \sinh^2(a+bx) dx}{3b^2} + \frac{4 \int \frac{\sinh(2a+2bx)}{2(a+bx)} dx}{3b^2} - \frac{(2a) \int \left(\frac{1}{2(a+bx)} - \frac{\cosh(2a+2bx)}{2(a+bx)} \right) dx}{3b^2} \\
&\quad + \frac{(4a) \int \frac{\sinh^2(a+bx)}{a+bx} dx}{3b^2} + \frac{(2a^2) \int \frac{\sinh(2a+2bx)}{2(a+bx)} dx}{3b^2} - \frac{a \int \frac{x \sinh(2a+2bx)}{a+bx} dx}{3b} \\
&= -\frac{a \log(a+bx)}{3b^3} - \frac{2 \cosh(a+bx) \sinh(a+bx)}{3b^3} - \frac{4 \cosh(a+bx)\text{Shi}(a+bx)}{3b^3} \\
&\quad - \frac{2a^2 \cosh(a+bx)\text{Shi}(a+bx)}{3b^3} + \frac{2ax \cosh(a+bx)\text{Shi}(a+bx)}{3b^2} \\
&\quad - \frac{2x^2 \cosh(a+bx)\text{Shi}(a+bx)}{3b} - \frac{2a \sinh(a+bx)\text{Shi}(a+bx)}{3b^3} \\
&\quad + \frac{4x \sinh(a+bx)\text{Shi}(a+bx)}{3b^2} + \frac{a^2(a+bx)\text{Shi}(a+bx)^2}{3b^3} - \frac{ax(a+bx)\text{Shi}(a+bx)^2}{3b^2} \\
&\quad + \frac{x^2(a+bx)\text{Shi}(a+bx)^2}{3b} + \frac{2 \int 1 dx}{3b^2} + \frac{2 \int \frac{\sinh(2a+2bx)}{a+bx} dx}{3b^2} + \frac{a \int \frac{\cosh(2a+2bx)}{a+bx} dx}{3b^2} \\
&\quad - \frac{a \int \sinh(2a+2bx) dx}{3b^2} - \frac{(4a) \int \left(\frac{1}{2(a+bx)} - \frac{\cosh(2a+2bx)}{2(a+bx)} \right) dx}{3b^2} + 2 \frac{a^2 \int \frac{\sinh(2a+2bx)}{a+bx} dx}{3b^2} \\
&\quad + \frac{\int x \sinh(2a+2bx) dx}{3b} - \frac{a \int \left(\frac{\sinh(2a+2bx)}{b} + \frac{a \sinh(2a+2bx)}{b(-a-bx)} \right) dx}{3b}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2x}{3b^2} - \frac{a \cosh(2a + 2bx)}{6b^3} + \frac{x \cosh(2a + 2bx)}{6b^2} + \frac{a \operatorname{Chi}(2a + 2bx)}{3b^3} \\
&\quad - \frac{a \log(a + bx)}{b^3} - \frac{2 \cosh(a + bx) \sinh(a + bx)}{3b^3} - \frac{4 \cosh(a + bx) \operatorname{Shi}(a + bx)}{3b^3} \\
&\quad - \frac{2a^2 \cosh(a + bx) \operatorname{Shi}(a + bx)}{3b^3} + \frac{2ax \cosh(a + bx) \operatorname{Shi}(a + bx)}{3b^2} \\
&\quad - \frac{2x^2 \cosh(a + bx) \operatorname{Shi}(a + bx)}{3b} - \frac{2a \sinh(a + bx) \operatorname{Shi}(a + bx)}{3b^3} \\
&\quad + \frac{4x \sinh(a + bx) \operatorname{Shi}(a + bx)}{3b^2} + \frac{a^2(a + bx) \operatorname{Shi}(a + bx)^2}{3b^3} \\
&\quad - \frac{ax(a + bx) \operatorname{Shi}(a + bx)^2}{3b^2} + \frac{x^2(a + bx) \operatorname{Shi}(a + bx)^2}{3b} \\
&\quad + \frac{2 \operatorname{Shi}(2a + 2bx)}{3b^3} + \frac{2a^2 \operatorname{Shi}(2a + 2bx)}{3b^3} - \frac{\int \cosh(2a + 2bx) dx}{6b^2} \\
&\quad - \frac{a \int \sinh(2a + 2bx) dx}{3b^2} + \frac{(2a) \int \frac{\cosh(2a+2bx)}{a+bx} dx}{3b^2} - \frac{a^2 \int \frac{\sinh(2a+2bx)}{-a-bx} dx}{3b^2} \\
&= \frac{2x}{3b^2} - \frac{a \cosh(2a + 2bx)}{3b^3} + \frac{x \cosh(2a + 2bx)}{6b^2} + \frac{a \operatorname{Chi}(2a + 2bx)}{b^3} - \frac{a \log(a + bx)}{b^3} \\
&\quad - \frac{2 \cosh(a + bx) \sinh(a + bx)}{3b^3} - \frac{\sinh(2a + 2bx)}{12b^3} - \frac{4 \cosh(a + bx) \operatorname{Shi}(a + bx)}{3b^3} \\
&\quad - \frac{2a^2 \cosh(a + bx) \operatorname{Shi}(a + bx)}{3b^3} + \frac{2ax \cosh(a + bx) \operatorname{Shi}(a + bx)}{3b^2} \\
&\quad - \frac{2x^2 \cosh(a + bx) \operatorname{Shi}(a + bx)}{3b} - \frac{2a \sinh(a + bx) \operatorname{Shi}(a + bx)}{3b^3} \\
&\quad + \frac{4x \sinh(a + bx) \operatorname{Shi}(a + bx)}{3b^2} + \frac{a^2(a + bx) \operatorname{Shi}(a + bx)^2}{3b^3} - \frac{ax(a + bx) \operatorname{Shi}(a + bx)^2}{3b^2} \\
&\quad + \frac{x^2(a + bx) \operatorname{Shi}(a + bx)^2}{3b} + \frac{2 \operatorname{Shi}(2a + 2bx)}{3b^3} + \frac{a^2 \operatorname{Shi}(2a + 2bx)}{b^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.02 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.48

$$\int x^2 \operatorname{Shi}(a + bx)^2 dx = \frac{8a + 8bx - 4a \cosh(2(a + bx)) + 2bx \cosh(2(a + bx)) + 12a \operatorname{Chi}(2(a + bx)) - 12a \log(a + bx) - 5 \sinh(2(a + bx))}{12b^3}$$

[In] Integrate[x^2*SinhIntegral[a + b*x]^2,x]

[Out] (8*a + 8*b*x - 4*a*Cosh[2*(a + b*x)] + 2*b*x*Cosh[2*(a + b*x)] + 12*a*CoshIntegral[2*(a + b*x)] - 12*a*Log[a + b*x] - 5*Sinh[2*(a + b*x)] - 8*((2 + a^2 - a*b*x + b^2*x^2)*Cosh[a + b*x] + (a - 2*b*x)*Sinh[a + b*x])*SinhIntegral[a + b*x] + 4*(a^3 + b^3*x^3)*SinhIntegral[a + b*x]^2 + 8*SinhIntegral[2*(a + b*x)] + 12*a^2*SinhIntegral[2*(a + b*x)])/(12*b^3)

Maple [F]

$$\int x^2 \operatorname{Shi}(bx + a)^2 dx$$

```
[In] int(x^2*Shi(b*x+a)^2,x)
```

```
[Out] int(x^2*Shi(b*x+a)^2,x)
```

Fricas [F]

$$\int x^2 \operatorname{Shi}(a + bx)^2 dx = \int x^2 \operatorname{Shi}(bx + a)^2 dx$$

```
[In] integrate(x^2*Shi(b*x+a)^2,x, algorithm="fricas")
```

```
[Out] integral(x^2*sinh_integral(b*x + a)^2, x)
```

Sympy [F]

$$\int x^2 \operatorname{Shi}(a + bx)^2 dx = \int x^2 \operatorname{Shi}^2(a + bx) dx$$

```
[In] integrate(x**2*Shi(b*x+a)**2,x)
```

```
[Out] Integral(x**2*Shi(a + b*x)**2, x)
```

Maxima [F]

$$\int x^2 \operatorname{Shi}(a + bx)^2 dx = \int x^2 \operatorname{Shi}(bx + a)^2 dx$$

```
[In] integrate(x^2*Shi(b*x+a)^2,x, algorithm="maxima")
```

```
[Out] integrate(x^2*Shi(b*x + a)^2, x)
```


Giac [F]

$$\int x^2 \operatorname{Shi}(a + bx)^2 dx = \int x^2 \operatorname{Shi}(bx + a)^2 dx$$

[In] integrate(x^2*Shi(b*x+a)^2,x, algorithm="giac")

[Out] integrate(x^2*Shi(b*x + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{Shi}(a + bx)^2 dx = \int x^2 \operatorname{sinhint}(a + bx)^2 dx$$

[In] int(x^2*sinhint(a + b*x)^2,x)

[Out] int(x^2*sinhint(a + b*x)^2, x)

3.27 $\int x\text{Shi}(a + bx)^2 dx$

Optimal result	170
Rubi [A] (verified)	170
Mathematica [A] (verified)	174
Maple [A] (verified)	174
Fricas [F]	175
Sympy [F]	175
Maxima [F]	175
Giac [F]	175
Mupad [F(-1)]	176

Optimal result

Integrand size = 10, antiderivative size = 154

$$\begin{aligned} \int x\text{Shi}(a + bx)^2 dx = & \frac{\cosh(2a + 2bx)}{4b^2} - \frac{\text{Chi}(2a + 2bx)}{2b^2} + \frac{\log(a + bx)}{2b^2} \\ & + \frac{a \cosh(a + bx)\text{Shi}(a + bx)}{b^2} - \frac{x \cosh(a + bx)\text{Shi}(a + bx)}{b} \\ & + \frac{\sinh(a + bx)\text{Shi}(a + bx)}{b^2} - \frac{a(a + bx)\text{Shi}(a + bx)^2}{2b^2} \\ & + \frac{x(a + bx)\text{Shi}(a + bx)^2}{2b} - \frac{a\text{Shi}(2a + 2bx)}{b^2} \end{aligned}$$

[Out] $-1/2*\text{Chi}(2*b*x+2*a)/b^2+1/4*\cosh(2*b*x+2*a)/b^2+1/2*\ln(b*x+a)/b^2+a*\cosh(b*x+a)*\text{Shi}(b*x+a)/b^2-x*\cosh(b*x+a)*\text{Shi}(b*x+a)/b-1/2*a*(b*x+a)*\text{Shi}(b*x+a)^2/b^2+1/2*x*(b*x+a)*\text{Shi}(b*x+a)^2/b-a*\text{Shi}(2*b*x+2*a)/b^2+\text{Shi}(b*x+a)*\sinh(b*x+a)/b^2$

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 1.400$, Rules used = {6673, 6677, 5736, 6873, 6874, 2718, 3379, 6681, 3393, 3382, 6669, 6675, 5556, 12}

$$\begin{aligned} \int x\text{Shi}(a + bx)^2 dx = & -\frac{\text{Chi}(2a + 2bx)}{2b^2} - \frac{a(a + bx)\text{Shi}(a + bx)^2}{2b^2} \\ & - \frac{a\text{Shi}(2a + 2bx)}{b^2} + \frac{\text{Shi}(a + bx)\sinh(a + bx)}{b^2} \\ & + \frac{a\text{Shi}(a + bx)\cosh(a + bx)}{b^2} + \frac{\log(a + bx)}{2b^2} + \frac{\cosh(2a + 2bx)}{4b^2} \\ & + \frac{x(a + bx)\text{Shi}(a + bx)^2}{2b} - \frac{x\text{Shi}(a + bx)\cosh(a + bx)}{b} \end{aligned}$$

[In] Int[x*SinhIntegral[a + b*x]^2,x]

[Out] Cosh[2*a + 2*b*x]/(4*b^2) - CoshIntegral[2*a + 2*b*x]/(2*b^2) + Log[a + b*x]/(2*b^2) + (a*Cosh[a + b*x]*SinhIntegral[a + b*x])/b^2 - (x*Cosh[a + b*x]*SinhIntegral[a + b*x])/b + (Sinh[a + b*x]*SinhIntegral[a + b*x])/b^2 - (a*(a + b*x)*SinhIntegral[a + b*x]^2)/(2*b^2) + (x*(a + b*x)*SinhIntegral[a + b*x]^2)/(2*b) - (a*SinhIntegral[2*a + 2*b*x])/b^2

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 2718

Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3393

Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))

Rule 5556

Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]

Rule 5736

Int[Cosh[w_]^(p_.)*(u_.)*Sinh[v_]^(p_.), x_Symbol] := Dist[1/2^p, Int[u*Sinh[2*v]^p, x], x] /; EqQ[w, v] && IntegerQ[p]

Rule 6669

```
Int[SinhIntegral[(a_.) + (b_.)*(x_)^2, x_Symbol] := Simp[(a + b*x)*(SinhIntegral[a + b*x]^2/b), x] - Dist[2, Int[Sinh[a + b*x]*SinhIntegral[a + b*x], x], x] /; FreeQ[{a, b}, x]
```

Rule 6673

```
Int[((c_.) + (d_.)*(x_)^(m_.))*SinhIntegral[(a_.) + (b_.)*(x_)^2, x_Symbol] := Simp[(a + b*x)*(c + d*x)^m*(SinhIntegral[a + b*x]^2/(b*(m + 1))), x] + (-Dist[2/(m + 1), Int[(c + d*x)^m*Sinh[a + b*x]*SinhIntegral[a + b*x], x], x] + Dist[(b*c - a*d)*(m/(b*(m + 1))), Int[(c + d*x)^(m - 1)*SinhIntegral[a + b*x]^2, x], x]) /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0]
```

Rule 6675

```
Int[Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6677

```
Int[((e_.) + (f_.)*(x_)^(m_.))*Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*SinhIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6681

```
Int[Cosh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sinh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6873

```
Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v != u]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x(a+bx)\text{Shi}(a+bx)^2}{2b} - \frac{a \int \text{Shi}(a+bx)^2 dx}{2b} - \int x \sinh(a+bx)\text{Shi}(a+bx) dx \\
&= -\frac{x \cosh(a+bx)\text{Shi}(a+bx)}{b} - \frac{a(a+bx)\text{Shi}(a+bx)^2}{2b^2} \\
&\quad + \frac{x(a+bx)\text{Shi}(a+bx)^2}{2b} + \frac{\int \cosh(a+bx)\text{Shi}(a+bx) dx}{b} \\
&\quad + \frac{a \int \sinh(a+bx)\text{Shi}(a+bx) dx}{b} + \int \frac{x \cosh(a+bx) \sinh(a+bx)}{a+bx} dx \\
&= \frac{a \cosh(a+bx)\text{Shi}(a+bx)}{b^2} - \frac{x \cosh(a+bx)\text{Shi}(a+bx)}{b} \\
&\quad + \frac{\sinh(a+bx)\text{Shi}(a+bx)}{b^2} - \frac{a(a+bx)\text{Shi}(a+bx)^2}{2b^2} + \frac{x(a+bx)\text{Shi}(a+bx)^2}{2b} \\
&\quad + \frac{1}{2} \int \frac{x \sinh(2(a+bx))}{a+bx} dx - \frac{\int \frac{\sinh^2(a+bx)}{a+bx} dx}{b} - \frac{a \int \frac{\cosh(a+bx) \sinh(a+bx)}{a+bx} dx}{b} \\
&= \frac{a \cosh(a+bx)\text{Shi}(a+bx)}{b^2} - \frac{x \cosh(a+bx)\text{Shi}(a+bx)}{b} \\
&\quad + \frac{\sinh(a+bx)\text{Shi}(a+bx)}{b^2} - \frac{a(a+bx)\text{Shi}(a+bx)^2}{2b^2} + \frac{x(a+bx)\text{Shi}(a+bx)^2}{2b} \\
&\quad + \frac{1}{2} \int \frac{x \sinh(2a+2bx)}{a+bx} dx + \frac{\int \left(\frac{1}{2(a+bx)} - \frac{\cosh(2a+2bx)}{2(a+bx)} \right) dx}{b} - \frac{a \int \frac{\sinh(2a+2bx)}{2(a+bx)} dx}{b} \\
&= \frac{\log(a+bx)}{2b^2} + \frac{a \cosh(a+bx)\text{Shi}(a+bx)}{b^2} - \frac{x \cosh(a+bx)\text{Shi}(a+bx)}{b} \\
&\quad + \frac{\sinh(a+bx)\text{Shi}(a+bx)}{b^2} - \frac{a(a+bx)\text{Shi}(a+bx)^2}{2b^2} \\
&\quad + \frac{x(a+bx)\text{Shi}(a+bx)^2}{2b} + \frac{1}{2} \int \left(\frac{\sinh(2a+2bx)}{b} + \frac{a \sinh(2a+2bx)}{b(-a-bx)} \right) dx \\
&\quad - \frac{\int \frac{\cosh(2a+2bx)}{a+bx} dx}{2b} - \frac{a \int \frac{\sinh(2a+2bx)}{a+bx} dx}{2b} \\
&= -\frac{\text{Chi}(2a+2bx)}{2b^2} + \frac{\log(a+bx)}{2b^2} + \frac{a \cosh(a+bx)\text{Shi}(a+bx)}{b^2} \\
&\quad - \frac{x \cosh(a+bx)\text{Shi}(a+bx)}{b} + \frac{\sinh(a+bx)\text{Shi}(a+bx)}{b^2} \\
&\quad - \frac{a(a+bx)\text{Shi}(a+bx)^2}{2b^2} + \frac{x(a+bx)\text{Shi}(a+bx)^2}{2b} \\
&\quad - \frac{a\text{Shi}(2a+2bx)}{2b^2} + \frac{\int \sinh(2a+2bx) dx}{2b} + \frac{a \int \frac{\sinh(2a+2bx)}{-a-bx} dx}{2b}
\end{aligned}$$

$$= \frac{\cosh(2a + 2bx)}{4b^2} - \frac{\text{Chi}(2a + 2bx)}{2b^2} + \frac{\log(a + bx)}{2b^2} + \frac{a \cosh(a + bx)\text{Shi}(a + bx)}{b^2}$$

$$- \frac{x \cosh(a + bx)\text{Shi}(a + bx)}{b} + \frac{\sinh(a + bx)\text{Shi}(a + bx)}{b^2}$$

$$- \frac{a(a + bx)\text{Shi}(a + bx)^2}{2b^2} + \frac{x(a + bx)\text{Shi}(a + bx)^2}{2b} - \frac{a\text{Shi}(2a + 2bx)}{b^2}$$

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.62

$$\int x\text{Shi}(a + bx)^2 dx$$

$$= \frac{\cosh(2(a + bx)) - 2\text{Chi}(2(a + bx)) + 2\log(a + bx) + 4((a - bx)\cosh(a + bx) + \sinh(a + bx))\text{Shi}(a + bx)}{4b^2}$$

[In] Integrate[x*SinhIntegral[a + b*x]^2,x]

[Out] (Cosh[2*(a + b*x)] - 2*CoshIntegral[2*(a + b*x)] + 2*Log[a + b*x] + 4*((a - b*x)*Cosh[a + b*x] + Sinh[a + b*x])*SinhIntegral[a + b*x] - 2*(a^2 - b^2*x^2)*SinhIntegral[a + b*x]^2 - 4*a*SinhIntegral[2*(a + b*x)])/(4*b^2)

Maple [A] (verified)

Time = 0.97 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.73

method	result
derivativedivides	$\frac{\text{Shi}(bx+a)^2 \left(-(bx+a)a + \frac{(bx+a)^2}{2} \right) - 2 \text{Shi}(bx+a) \left(-a \cosh(bx+a) + \frac{(bx+a) \cosh(bx+a)}{2} - \frac{\sinh(bx+a)}{2} \right) - a \text{Shi}(2bx+2a) + \cosh(2bx+2a)}{b^2}$
default	$\frac{\text{Shi}(bx+a)^2 \left(-(bx+a)a + \frac{(bx+a)^2}{2} \right) - 2 \text{Shi}(bx+a) \left(-a \cosh(bx+a) + \frac{(bx+a) \cosh(bx+a)}{2} - \frac{\sinh(bx+a)}{2} \right) - a \text{Shi}(2bx+2a) + \cosh(2bx+2a)}{b^2}$

[In] int(x*Shi(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/b^2*(Shi(b*x+a)^2*(-(b*x+a)*a+1/2*(b*x+a)^2)-2*Shi(b*x+a)*(-a*cosh(b*x+a)+1/2*(b*x+a)*cosh(b*x+a)-1/2*sinh(b*x+a))-a*Shi(2*b*x+2*a)+1/2*cosh(b*x+a)^2+1/2*ln(b*x+a)-1/2*Chi(2*b*x+2*a))

Fricas [F]

$$\int x\text{Shi}(a + bx)^2 dx = \int x\text{Shi}(bx + a)^2 dx$$

[In] integrate(x*Shi(b*x+a)^2,x, algorithm="fricas")

[Out] integral(x*sinh_integral(b*x + a)^2, x)

Sympy [F]

$$\int x\text{Shi}(a + bx)^2 dx = \int x\text{Shi}^2(a + bx) dx$$

[In] integrate(x*Shi(b*x+a)**2,x)

[Out] Integral(x*Shi(a + b*x)**2, x)

Maxima [F]

$$\int x\text{Shi}(a + bx)^2 dx = \int x\text{Shi}(bx + a)^2 dx$$

[In] integrate(x*Shi(b*x+a)^2,x, algorithm="maxima")

[Out] integrate(x*Shi(b*x + a)^2, x)

Giac [F]

$$\int x\text{Shi}(a + bx)^2 dx = \int x\text{Shi}(bx + a)^2 dx$$

[In] integrate(x*Shi(b*x+a)^2,x, algorithm="giac")

[Out] integrate(x*Shi(b*x + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{Shi}(a + bx)^2 dx = \int x \operatorname{sinhint}(a + bx)^2 dx$$

```
[In] int(x*sinhint(a + b*x)^2,x)
```

```
[Out] int(x*sinhint(a + b*x)^2, x)
```


3.28 $\int \text{Shi}(a + bx)^2 dx$

Optimal result	177
Rubi [A] (verified)	177
Mathematica [A] (verified)	179
Maple [A] (verified)	179
Fricas [F]	179
Sympy [F]	179
Maxima [F]	180
Giac [F]	180
Mupad [F(-1)]	180

Optimal result

Integrand size = 8, antiderivative size = 48

$$\int \text{Shi}(a + bx)^2 dx = -\frac{2 \cosh(a + bx)\text{Shi}(a + bx)}{b} + \frac{(a + bx)\text{Shi}(a + bx)^2}{b} + \frac{\text{Shi}(2a + 2bx)}{b}$$

[Out] $-2*\cosh(b*x+a)*\text{Shi}(b*x+a)/b+(b*x+a)*\text{Shi}(b*x+a)^2/b+\text{Shi}(2*b*x+2*a)/b$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {6669, 6675, 5556, 12, 3379}

$$\int \text{Shi}(a + bx)^2 dx = \frac{(a + bx)\text{Shi}(a + bx)^2}{b} + \frac{\text{Shi}(2a + 2bx)}{b} - \frac{2\text{Shi}(a + bx) \cosh(a + bx)}{b}$$

[In] $\text{Int}[\text{SinhIntegral}[a + b*x]^2, x]$

[Out] $(-2*\text{Cosh}[a + b*x]*\text{SinhIntegral}[a + b*x])/b + ((a + b*x)*\text{SinhIntegral}[a + b*x]^2)/b + \text{SinhIntegral}[2*a + 2*b*x]/b$

Rule 12

$\text{Int}[(a_*)(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{Match} Q[u, (b_*)(v_)] /; \text{FreeQ}[b, x]$

Rule 3379

$\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] \rightarrow \text{Simp}[I*(\text{SinhIntegral}[c*f*(fz/d) + f*fz*x]/d), x] /; \text{FreeQ}\{c, d, e, f$

, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

Rule 6669

```
Int[SinhIntegral[(a_.) + (b_.)*(x_)]^2, x_Symbol] := Simp[(a + b*x)*(SinhIn
tegral[a + b*x]^2/b), x] - Dist[2, Int[Sinh[a + b*x]*SinhIntegral[a + b*x],
x], x] /; FreeQ[{a, b}, x]
```

Rule 6675

```
Int[Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
Simp[Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(a + bx)\text{Shi}(a + bx)^2}{b} - 2 \int \sinh(a + bx)\text{Shi}(a + bx) dx \\
&= -\frac{2 \cosh(a + bx)\text{Shi}(a + bx)}{b} + \frac{(a + bx)\text{Shi}(a + bx)^2}{b} + 2 \int \frac{\cosh(a + bx) \sinh(a + bx)}{a + bx} dx \\
&= -\frac{2 \cosh(a + bx)\text{Shi}(a + bx)}{b} + \frac{(a + bx)\text{Shi}(a + bx)^2}{b} + 2 \int \frac{\sinh(2a + 2bx)}{2(a + bx)} dx \\
&= -\frac{2 \cosh(a + bx)\text{Shi}(a + bx)}{b} + \frac{(a + bx)\text{Shi}(a + bx)^2}{b} + \int \frac{\sinh(2a + 2bx)}{a + bx} dx \\
&= -\frac{2 \cosh(a + bx)\text{Shi}(a + bx)}{b} + \frac{(a + bx)\text{Shi}(a + bx)^2}{b} + \frac{\text{Shi}(2a + 2bx)}{b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.85

$$\int \operatorname{Shi}(a + bx)^2 dx = \frac{-2 \cosh(a + bx) \operatorname{Shi}(a + bx) + (a + bx) \operatorname{Shi}(a + bx)^2 + \operatorname{Shi}(2(a + bx))}{b}$$

[In] Integrate[SinhIntegral[a + b*x]^2,x]

[Out] (-2*Cosh[a + b*x]*SinhIntegral[a + b*x] + (a + b*x)*SinhIntegral[a + b*x]^2 + SinhIntegral[2*(a + b*x)])/b

Maple [A] (verified)

Time = 0.45 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.90

method	result	size
derivativedivides	$\frac{\operatorname{Shi}(bx+a)^2 (bx+a) - 2 \cosh(bx+a) \operatorname{Shi}(bx+a) + \operatorname{Shi}(2bx+2a)}{b}$	43
default	$\frac{\operatorname{Shi}(bx+a)^2 (bx+a) - 2 \cosh(bx+a) \operatorname{Shi}(bx+a) + \operatorname{Shi}(2bx+2a)}{b}$	43

[In] int(Shi(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/b*(Shi(b*x+a)^2*(b*x+a)-2*cosh(b*x+a)*Shi(b*x+a)+Shi(2*b*x+2*a))

Fricas [F]

$$\int \operatorname{Shi}(a + bx)^2 dx = \int \operatorname{Shi}(bx + a)^2 dx$$

[In] integrate(Shi(b*x+a)^2,x, algorithm="fricas")

[Out] integral(sinh_integral(b*x + a)^2, x)

Sympy [F]

$$\int \operatorname{Shi}(a + bx)^2 dx = \int \operatorname{Shi}^2(a + bx) dx$$

[In] integrate(Shi(b*x+a)**2,x)

[Out] Integral(Shi(a + b*x)**2, x)

Maxima [F]

$$\int \operatorname{Shi}(a + bx)^2 dx = \int \operatorname{Shi}(bx + a)^2 dx$$

[In] integrate(Shi(b*x+a)^2,x, algorithm="maxima")

[Out] integrate(Shi(b*x + a)^2, x)

Giac [F]

$$\int \operatorname{Shi}(a + bx)^2 dx = \int \operatorname{Shi}(bx + a)^2 dx$$

[In] integrate(Shi(b*x+a)^2,x, algorithm="giac")

[Out] integrate(Shi(b*x + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \operatorname{Shi}(a + bx)^2 dx = \int \operatorname{sinhint}(a + bx)^2 dx$$

[In] int(sinhint(a + b*x)^2,x)

[Out] int(sinhint(a + b*x)^2, x)

3.29 $\int \frac{\text{Shi}(a+bx)^2}{x} dx$

Optimal result	181
Rubi [N/A]	181
Mathematica [N/A]	182
Maple [N/A] (verified)	182
Fricas [N/A]	182
Sympy [N/A]	182
Maxima [N/A]	183
Giac [N/A]	183
Mupad [N/A]	183

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\text{Shi}(a+bx)^2}{x} dx = \text{Int}\left(\frac{\text{Shi}(a+bx)^2}{x}, x\right)$$

[Out] CannotIntegrate(Shi(b*x+a)^2/x,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\text{Shi}(a+bx)^2}{x} dx = \int \frac{\text{Shi}(a+bx)^2}{x} dx$$

[In] Int[SinhIntegral[a + b*x]^2/x,x]

[Out] Defer[Int][SinhIntegral[a + b*x]^2/x, x]

Rubi steps

$$\text{integral} = \int \frac{\text{Shi}(a+bx)^2}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 0.46 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Shi}(a + bx)^2}{x} dx = \int \frac{\operatorname{Shi}(a + bx)^2}{x} dx$$

`[In] Integrate[SinhIntegral[a + b*x]^2/x, x]``[Out] Integrate[SinhIntegral[a + b*x]^2/x, x]`**Maple [N/A] (verified)**

Not integrable

Time = 0.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{Shi}(bx + a)^2}{x} dx$$

`[In] int(Shi(b*x+a)^2/x, x)``[Out] int(Shi(b*x+a)^2/x, x)`**Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Shi}(a + bx)^2}{x} dx = \int \frac{\operatorname{Shi}(bx + a)^2}{x} dx$$

`[In] integrate(Shi(b*x+a)^2/x, x, algorithm="fricas")``[Out] integral(sinh_integral(b*x + a)^2/x, x)`**Sympy [N/A]**

Not integrable

Time = 0.41 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{Shi}(a + bx)^2}{x} dx = \int \frac{\operatorname{Shi}^2(a + bx)}{x} dx$$

`[In] integrate(Shi(b*x+a)**2/x, x)``[Out] Integral(Shi(a + b*x)**2/x, x)`

Maxima [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Shi}(a + bx)^2}{x} dx = \int \frac{\operatorname{Shi}(bx + a)^2}{x} dx$$

[In] integrate(Shi(b*x+a)^2/x,x, algorithm="maxima")

[Out] integrate(Shi(b*x + a)^2/x, x)

Giac [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Shi}(a + bx)^2}{x} dx = \int \frac{\operatorname{Shi}(bx + a)^2}{x} dx$$

[In] integrate(Shi(b*x+a)^2/x,x, algorithm="giac")

[Out] integrate(Shi(b*x + a)^2/x, x)

Mupad [N/A]

Not integrable

Time = 4.88 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Shi}(a + bx)^2}{x} dx = \int \frac{\operatorname{sinhint}(a + bx)^2}{x} dx$$

[In] int(sinhint(a + b*x)^2/x,x)

[Out] int(sinhint(a + b*x)^2/x, x)

3.30 $\int \frac{\mathbf{Shi}(a+bx)^2}{x^2} dx$

Optimal result	184
Rubi [N/A]	184
Mathematica [N/A]	185
Maple [N/A] (verified)	185
Fricas [N/A]	185
Sympy [N/A]	185
Maxima [N/A]	186
Giac [N/A]	186
Mupad [N/A]	186

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\mathbf{Shi}(a+bx)^2}{x^2} dx = \text{Int}\left(\frac{\mathbf{Shi}(a+bx)^2}{x^2}, x\right)$$

[Out] CannotIntegrate(Shi(b*x+a)^2/x^2,x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\mathbf{Shi}(a+bx)^2}{x^2} dx = \int \frac{\mathbf{Shi}(a+bx)^2}{x^2} dx$$

[In] Int[SinhIntegral[a + b*x]^2/x^2,x]

[Out] Defer[Int][SinhIntegral[a + b*x]^2/x^2, x]

Rubi steps

$$\text{integral} = \int \frac{\mathbf{Shi}(a+bx)^2}{x^2} dx$$

Mathematica [N/A]

Not integrable

Time = 0.93 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Shi}(a + bx)^2}{x^2} dx = \int \frac{\text{Shi}(a + bx)^2}{x^2} dx$$

[In] Integrate[SinhIntegral[a + b*x]^2/x^2,x]

[Out] Integrate[SinhIntegral[a + b*x]^2/x^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.21 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\text{Shi}(bx + a)^2}{x^2} dx$$

[In] int(Shi(b*x+a)^2/x^2,x)

[Out] int(Shi(b*x+a)^2/x^2,x)

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Shi}(a + bx)^2}{x^2} dx = \int \frac{\text{Shi}(bx + a)^2}{x^2} dx$$

[In] integrate(Shi(b*x+a)^2/x^2,x, algorithm="fricas")

[Out] integral(sinh_integral(b*x + a)^2/x^2, x)

Sympy [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\text{Shi}(a + bx)^2}{x^2} dx = \int \frac{\text{Shi}^2(a + bx)}{x^2} dx$$

[In] integrate(Shi(b*x+a)**2/x**2,x)

[Out] Integral(Shi(a + b*x)**2/x**2, x)

Maxima [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Shi}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{Shi}(bx + a)^2}{x^2} dx$$

[In] integrate(Shi(b*x+a)^2/x^2,x, algorithm="maxima")

[Out] integrate(Shi(b*x + a)^2/x^2, x)

Giac [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Shi}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{Shi}(bx + a)^2}{x^2} dx$$

[In] integrate(Shi(b*x+a)^2/x^2,x, algorithm="giac")

[Out] integrate(Shi(b*x + a)^2/x^2, x)

Mupad [N/A]

Not integrable

Time = 4.93 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Shi}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{sinhint}(a + bx)^2}{x^2} dx$$

[In] int(sinhint(a + b*x)^2/x^2,x)

[Out] int(sinhint(a + b*x)^2/x^2, x)

3.31 $\int \frac{\text{Shi}(a+bx)^2}{x^3} dx$

Optimal result	187
Rubi [N/A]	187
Mathematica [N/A]	188
Maple [N/A] (verified)	188
Fricas [N/A]	188
Sympy [N/A]	188
Maxima [N/A]	189
Giac [N/A]	189
Mupad [N/A]	189

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\text{Shi}(a+bx)^2}{x^3} dx = \text{Int}\left(\frac{\text{Shi}(a+bx)^2}{x^3}, x\right)$$

[Out] CannotIntegrate(Shi(b*x+a)^2/x^3,x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\text{Shi}(a+bx)^2}{x^3} dx = \int \frac{\text{Shi}(a+bx)^2}{x^3} dx$$

[In] Int[SinhIntegral[a + b*x]^2/x^3,x]

[Out] Defer[Int][SinhIntegral[a + b*x]^2/x^3, x]

Rubi steps

$$\text{integral} = \int \frac{\text{Shi}(a+bx)^2}{x^3} dx$$

Mathematica [N/A]

Not integrable

Time = 1.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Shi}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{Shi}(a + bx)^2}{x^3} dx$$

`[In] Integrate[SinhIntegral[a + b*x]^2/x^3,x]``[Out] Integrate[SinhIntegral[a + b*x]^2/x^3, x]`**Maple [N/A] (verified)**

Not integrable

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{Shi}(bx + a)^2}{x^3} dx$$

`[In] int(Shi(b*x+a)^2/x^3,x)``[Out] int(Shi(b*x+a)^2/x^3,x)`**Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Shi}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{Shi}(bx + a)^2}{x^3} dx$$

`[In] integrate(Shi(b*x+a)^2/x^3,x, algorithm="fricas")``[Out] integral(sinh_integral(b*x + a)^2/x^3, x)`**Sympy [N/A]**

Not integrable

Time = 0.41 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{Shi}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{Shi}^2(a + bx)}{x^3} dx$$

`[In] integrate(Shi(b*x+a)**2/x**3,x)``[Out] Integral(Shi(a + b*x)**2/x**3, x)`

Maxima [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Shi}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{Shi}(bx + a)^2}{x^3} dx$$

[In] integrate(Shi(b*x+a)^2/x^3,x, algorithm="maxima")

[Out] integrate(Shi(b*x + a)^2/x^3, x)

Giac [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Shi}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{Shi}(bx + a)^2}{x^3} dx$$

[In] integrate(Shi(b*x+a)^2/x^3,x, algorithm="giac")

[Out] integrate(Shi(b*x + a)^2/x^3, x)

Mupad [N/A]

Not integrable

Time = 4.92 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Shi}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{sinhint}(a + bx)^2}{x^3} dx$$

[In] int(sinhint(a + b*x)^2/x^3,x)

[Out] int(sinhint(a + b*x)^2/x^3, x)

3.32 $\int x^2 \text{Shi}(d(a + b \log(cx^n))) dx$

Optimal result	190
Rubi [A] (verified)	190
Mathematica [A] (verified)	192
Maple [F]	193
Fricas [F]	193
Sympy [F]	193
Maxima [F]	193
Giac [F]	194
Mupad [F(-1)]	194

Optimal result

Integrand size = 17, antiderivative size = 128

$$\begin{aligned} & \int x^2 \text{Shi}(d(a + b \log(cx^n))) dx \\ &= \frac{1}{6} e^{-\frac{3a}{bn}} x^3 (cx^n)^{-3/n} \text{ExpIntegralEi} \left(\frac{(3 - bdn)(a + b \log(cx^n))}{bn} \right) \\ & \quad - \frac{1}{6} e^{-\frac{3a}{bn}} x^3 (cx^n)^{-3/n} \text{ExpIntegralEi} \left(\frac{(3 + bdn)(a + b \log(cx^n))}{bn} \right) \\ & \quad + \frac{1}{3} x^3 \text{Shi}(d(a + b \log(cx^n))) \end{aligned}$$

[Out] 1/6*x^3*Ei((-b*d*n+3)*(a+b*ln(c*x^n))/b/n)/exp(3*a/b/n)/((c*x^n)^(3/n))-1/6*x^3*Ei((b*d*n+3)*(a+b*ln(c*x^n))/b/n)/exp(3*a/b/n)/((c*x^n)^(3/n))+1/3*x^3*Shi(d*(a+b*ln(c*x^n)))

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {6690, 12, 5650, 2347, 2209}

$$\begin{aligned} & \int x^2 \text{Shi}(d(a + b \log(cx^n))) dx \\ &= \frac{1}{6} x^3 e^{-\frac{3a}{bn}} (cx^n)^{-3/n} \text{ExpIntegralEi} \left(\frac{(3 - bdn)(a + b \log(cx^n))}{bn} \right) \\ & \quad - \frac{1}{6} x^3 e^{-\frac{3a}{bn}} (cx^n)^{-3/n} \text{ExpIntegralEi} \left(\frac{(bdn + 3)(a + b \log(cx^n))}{bn} \right) \\ & \quad + \frac{1}{3} x^3 \text{Shi}(d(a + b \log(cx^n))) \end{aligned}$$

[In] Int[x^2*SinhIntegral[d*(a + b*Log[c*x^n]),x]

[Out] (x^3*ExpIntegralEi[((3 - b*d*n)*(a + b*Log[c*x^n]))/(b*n)])/(6*E^((3*a)/(b*n))*(c*x^n)^(3/n)) - (x^3*ExpIntegralEi[((3 + b*d*n)*(a + b*Log[c*x^n]))/(b*n)])/(6*E^((3*a)/(b*n))*(c*x^n)^(3/n)) + (x^3*SinhIntegral[d*(a + b*Log[c*x^n])])/3

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 2209

Int[(F_)^((g_)*((e_) + (f_)*(x_)))/((c_) + (d_)*(x_)), x_Symbol] := Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[\$UseGamma]

Rule 2347

Int[((a_) + Log[(c_)*(x_)^(n_)])*(b_)^(p_)*((d_)*(x_)^(m_)), x_Symbol] := Dist[(d*x)^(m + 1)/(d*n*(c*x^n)^((m + 1)/n)), Subst[Int[E^((m + 1)/n)*x]*(a + b*x)^p, x], x, Log[c*x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x]

Rule 5650

Int[((e_) + Log[(g_)*(x_)^(m_)])*(f_)*(h_)^(q_)*((i_)*(x_)^(r_))*Sinh[((a_) + Log[(c_)*(x_)^(n_)])*(b_)]*(d_), x_Symbol] := Dist[(-E^((-a)*d))*(i*x)^r*(1/((c*x^n)^(b*d)*(2*x^(r - b*d*n))))], Int[x^(r - b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] + Dist[E^(a*d)*(i*x)^r*((c*x^n)^(b*d)/(2*x^(r + b*d*n))), Int[x^(r + b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, m, n, q, r}, x]

Rule 6690

Int[((e_)*(x_)^(m_))*SinhIntegral[((a_) + Log[(c_)*(x_)^(n_)])*(b_)]*(d_), x_Symbol] := Simp[(e*x)^(m + 1)*(SinhIntegral[d*(a + b*Log[c*x^n])]/(e*(m + 1))), x] - Dist[b*d*(n/(m + 1)), Int[(e*x)^m*(Sinh[d*(a + b*Log[c*x^n])]/(d*(a + b*Log[c*x^n]))], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{3}x^3\text{Shi}(d(a + b \log(cx^n))) - \frac{1}{3}(bdn) \int \frac{x^2 \sinh(d(a + b \log(cx^n)))}{d(a + b \log(cx^n))} dx \\ &= \frac{1}{3}x^3\text{Shi}(d(a + b \log(cx^n))) - \frac{1}{3}(bn) \int \frac{x^2 \sinh(d(a + b \log(cx^n)))}{a + b \log(cx^n)} dx \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{3}x^3\text{Shi}(d(a + b \log(cx^n))) + \frac{1}{6}\left(be^{-ad}nx^{bdn}(cx^n)^{-bd} \right) \int \frac{x^{2-bdn}}{a + b \log(cx^n)} dx \\
&\quad - \frac{1}{6}\left(be^{ad}nx^{-bdn}(cx^n)^{bd} \right) \int \frac{x^{2+bdn}}{a + b \log(cx^n)} dx \\
&= \frac{1}{3}x^3\text{Shi}(d(a + b \log(cx^n))) \\
&\quad + \frac{1}{6}\left(be^{-ad}x^3(cx^n)^{-bd-\frac{3-bdn}{n}} \right) \text{Subst}\left(\int \frac{e^{\frac{(3-bdn)x}{n}}}{a + bx} dx, x, \log(cx^n) \right) \\
&\quad - \frac{1}{6}\left(be^{ad}x^3(cx^n)^{bd-\frac{3+bdn}{n}} \right) \text{Subst}\left(\int \frac{e^{\frac{(3+bdn)x}{n}}}{a + bx} dx, x, \log(cx^n) \right) \\
&= \frac{1}{6}e^{-\frac{3a}{bn}}x^3(cx^n)^{-3/n} \text{ExpIntegralEi}\left(\frac{(3 - bdn)(a + b \log(cx^n))}{bn} \right) \\
&\quad - \frac{1}{6}e^{-\frac{3a}{bn}}x^3(cx^n)^{-3/n} \text{ExpIntegralEi}\left(\frac{(3 + bdn)(a + b \log(cx^n))}{bn} \right) \\
&\quad + \frac{1}{3}x^3\text{Shi}(d(a + b \log(cx^n)))
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.09 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.77

$$\begin{aligned}
&\int x^2\text{Shi}(d(a + b \log(cx^n))) dx \\
&= \frac{1}{6}x^3\left(e^{-\frac{3a}{bn}}(cx^n)^{-3/n} \left(\text{ExpIntegralEi}\left(-\frac{(-3 + bdn)(a + b \log(cx^n))}{bn} \right) - \text{ExpIntegralEi}\left(\frac{(3 + bdn)(a + b \log(cx^n))}{bn} \right) \right) \right. \\
&\quad \left. + 2\text{Shi}(d(a + b \log(cx^n))) \right)
\end{aligned}$$

[In] Integrate[x^2*SinhIntegral[d*(a + b*Log[c*x^n]),x]

[Out] (x^3*((ExpIntegralEi[-(((3 + b*d*n)*(a + b*Log[c*x^n]))/(b*n))] - ExpIntegralEi[(((3 + b*d*n)*(a + b*Log[c*x^n]))/(b*n))]/(E^((3*a)/(b*n))*(c*x^n)^(3/n)) + 2*SinhIntegral[d*(a + b*Log[c*x^n]))])/6

Maple [F]

$$\int x^2 \operatorname{Shi}(d(a + b \ln(cx^n))) dx$$

[In] `int(x^2*Shi(d*(a+b*ln(c*x^n))),x)`

[Out] `int(x^2*Shi(d*(a+b*ln(c*x^n))),x)`

Fricas [F]

$$\int x^2 \operatorname{Shi}(d(a + b \log(cx^n))) dx = \int x^2 \operatorname{Shi}((b \log(cx^n) + a)d) dx$$

[In] `integrate(x^2*Shi(d*(a+b*log(c*x^n))),x, algorithm="fricas")`

[Out] `integral(x^2*sinh_integral(b*d*log(c*x^n) + a*d), x)`

Sympy [F]

$$\int x^2 \operatorname{Shi}(d(a + b \log(cx^n))) dx = \int x^2 \operatorname{Shi}(ad + bd \log(cx^n)) dx$$

[In] `integrate(x**2*Shi(d*(a+b*ln(c*x**n))),x)`

[Out] `Integral(x**2*Shi(a*d + b*d*log(c*x**n)), x)`

Maxima [F]

$$\int x^2 \operatorname{Shi}(d(a + b \log(cx^n))) dx = \int x^2 \operatorname{Shi}((b \log(cx^n) + a)d) dx$$

[In] `integrate(x^2*Shi(d*(a+b*log(c*x^n))),x, algorithm="maxima")`

[Out] `integrate(x^2*Shi((b*log(c*x^n) + a)*d), x)`

Giac [F]

$$\int x^2 \operatorname{Shi}(d(a + b \log(cx^n))) dx = \int x^2 \operatorname{Shi}((b \log(cx^n) + a)d) dx$$

[In] integrate(x^2*Shi(d*(a+b*log(c*x^n))),x, algorithm="giac")

[Out] integrate(x^2*Shi((b*log(c*x^n) + a)*d), x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{Shi}(d(a + b \log(cx^n))) dx = \int x^2 \operatorname{sinhint}(d(a + b \ln(cx^n))) dx$$

[In] int(x^2*sinhint(d*(a + b*log(c*x^n))),x)

[Out] int(x^2*sinhint(d*(a + b*log(c*x^n))), x)

3.33 $\int x \text{Shi}(d(a + b \log(cx^n))) dx$

Optimal result	195
Rubi [A] (verified)	195
Mathematica [A] (verified)	197
Maple [F]	198
Fricas [F]	198
Sympy [F]	198
Maxima [F]	198
Giac [F]	199
Mupad [F(-1)]	199

Optimal result

Integrand size = 15, antiderivative size = 128

$$\begin{aligned} & \int x \text{Shi}(d(a + b \log(cx^n))) dx \\ &= \frac{1}{4} e^{-\frac{2a}{bn}} x^2 (cx^n)^{-2/n} \text{ExpIntegralEi} \left(\frac{(2 - bdn)(a + b \log(cx^n))}{bn} \right) \\ & \quad - \frac{1}{4} e^{-\frac{2a}{bn}} x^2 (cx^n)^{-2/n} \text{ExpIntegralEi} \left(\frac{(2 + bdn)(a + b \log(cx^n))}{bn} \right) \\ & \quad + \frac{1}{2} x^2 \text{Shi}(d(a + b \log(cx^n))) \end{aligned}$$

[Out] 1/4*x^2*Ei((-b*d*n+2)*(a+b*ln(c*x^n))/b/n)/exp(2*a/b/n)/((c*x^n)^(2/n))-1/4*x^2*Ei((b*d*n+2)*(a+b*ln(c*x^n))/b/n)/exp(2*a/b/n)/((c*x^n)^(2/n))+1/2*x^2*Shi(d*(a+b*ln(c*x^n)))

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6690, 12, 5650, 2347, 2209}

$$\begin{aligned} & \int x \text{Shi}(d(a + b \log(cx^n))) dx \\ &= \frac{1}{4} x^2 e^{-\frac{2a}{bn}} (cx^n)^{-2/n} \text{ExpIntegralEi} \left(\frac{(2 - bdn)(a + b \log(cx^n))}{bn} \right) \\ & \quad - \frac{1}{4} x^2 e^{-\frac{2a}{bn}} (cx^n)^{-2/n} \text{ExpIntegralEi} \left(\frac{(bdn + 2)(a + b \log(cx^n))}{bn} \right) \\ & \quad + \frac{1}{2} x^2 \text{Shi}(d(a + b \log(cx^n))) \end{aligned}$$

```
[In] Int[x*SinhIntegral[d*(a + b*Log[c*x^n])],x]
[Out] (x^2*ExpIntegralEi[((2 - b*d*n)*(a + b*Log[c*x^n]))/(b*n)]/(4*E^((2*a)/(b*n)))*(c*x^n)^(2/n)) - (x^2*ExpIntegralEi[((2 + b*d*n)*(a + b*Log[c*x^n]))/(b*n)]/(4*E^((2*a)/(b*n)))*(c*x^n)^(2/n)) + (x^2*SinhIntegral[d*(a + b*Log[c*x^n])])/2
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 2209

```
Int[(F_)^((g_)*((e_) + (f_)*(x_)))/((c_) + (d_)*(x_)), x_Symbol] := Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]
```

Rule 2347

```
Int[((a_) + Log[(c_)*(x_)^(n_)])*(b_)^(p_)*((d_)*(x_)^(m_)), x_Symbol] := Dist[(d*x)^(m + 1)/(d*n*(c*x^n)^((m + 1)/n)), Subst[Int[E^(((m + 1)/n)*x)*(a + b*x)^p, x], x, Log[c*x^n]], x] /; FreeQ[{a, b, c, d, m, n, p}, x]
```

Rule 5650

```
Int[(((e_) + Log[(g_)*(x_)^(m_)])*(f_)*(h_)^(q_)*((i_)*(x_)^(r_))*Sinh[((a_) + Log[(c_)*(x_)^(n_)])*(b_)]*(d_)), x_Symbol] := Dist[(-E^((-a)*d))*(i*x)^r*(1/((c*x^n)^(b*d)*(2*x^(r - b*d*n))))], Int[x^(r - b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] + Dist[E^(a*d)*(i*x)^r*((c*x^n)^(b*d)/(2*x^(r + b*d*n))), Int[x^(r + b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, m, n, q, r}, x]
```

Rule 6690

```
Int[((e_)*(x_)^(m_))*SinhIntegral[((a_) + Log[(c_)*(x_)^(n_)])*(b_)]*(d_), x_Symbol] := Simp[(e*x)^(m + 1)*(SinhIntegral[d*(a + b*Log[c*x^n])]/(e*(m + 1))), x] - Dist[b*d*(n/(m + 1)), Int[(e*x)^m*(Sinh[d*(a + b*Log[c*x^n])]/(d*(a + b*Log[c*x^n]))], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{2}x^2\text{Shi}(d(a + b \log(cx^n))) - \frac{1}{2}(bdn) \int \frac{x \sinh(d(a + b \log(cx^n)))}{d(a + b \log(cx^n))} dx \\ &= \frac{1}{2}x^2\text{Shi}(d(a + b \log(cx^n))) - \frac{1}{2}(bn) \int \frac{x \sinh(d(a + b \log(cx^n)))}{a + b \log(cx^n)} dx \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2}x^2 \text{Shi}(d(a + b \log(cx^n))) + \frac{1}{4} \left(b e^{-ad} n x^{bdn} (cx^n)^{-bd} \right) \int \frac{x^{1-bdn}}{a + b \log(cx^n)} dx \\
&\quad - \frac{1}{4} \left(b e^{ad} n x^{-bdn} (cx^n)^{bd} \right) \int \frac{x^{1+bdn}}{a + b \log(cx^n)} dx \\
&= \frac{1}{2}x^2 \text{Shi}(d(a + b \log(cx^n))) \\
&\quad + \frac{1}{4} \left(b e^{-ad} x^2 (cx^n)^{-bd - \frac{2-bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(2-bdn)x}{n}}}{a + bx} dx, x, \log(cx^n) \right) \\
&\quad - \frac{1}{4} \left(b e^{ad} x^2 (cx^n)^{bd - \frac{2+bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(2+bdn)x}{n}}}{a + bx} dx, x, \log(cx^n) \right) \\
&= \frac{1}{4} e^{-\frac{2a}{bn}} x^2 (cx^n)^{-2/n} \text{ExpIntegralEi} \left(\frac{(2 - bdn)(a + b \log(cx^n))}{bn} \right) \\
&\quad - \frac{1}{4} e^{-\frac{2a}{bn}} x^2 (cx^n)^{-2/n} \text{ExpIntegralEi} \left(\frac{(2 + bdn)(a + b \log(cx^n))}{bn} \right) \\
&\quad + \frac{1}{2} x^2 \text{Shi}(d(a + b \log(cx^n)))
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.08 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.77

$$\begin{aligned}
&\int x \text{Shi}(d(a + b \log(cx^n))) dx \\
&= \frac{1}{4} x^2 \left(e^{-\frac{2a}{bn}} (cx^n)^{-2/n} \left(\text{ExpIntegralEi} \left(-\frac{(-2 + bdn)(a + b \log(cx^n))}{bn} \right) - \text{ExpIntegralEi} \left(\frac{(2 + bdn)(a + b \log(cx^n))}{bn} \right) \right. \right. \\
&\quad \left. \left. + 2 \text{Shi}(d(a + b \log(cx^n))) \right) \right)
\end{aligned}$$

[In] Integrate[x*SinhIntegral[d*(a + b*Log[c*x^n])],x]

[Out] (x^2*((ExpIntegralEi[-(((-2 + b*d*n)*(a + b*Log[c*x^n]))/(b*n))] - ExpIntegralEi[((2 + b*d*n)*(a + b*Log[c*x^n]))/(b*n)])/(E^((2*a)/(b*n))*(c*x^n)^(2/n)) + 2*SinhIntegral[d*(a + b*Log[c*x^n])))/4

Maple [F]

$$\int x \operatorname{Shi}(d(a + b \ln(cx^n))) dx$$

```
[In] int(x*Shi(d*(a+b*ln(c*x^n))),x)
```

```
[Out] int(x*Shi(d*(a+b*ln(c*x^n))),x)
```

Fricas [F]

$$\int x \operatorname{Shi}(d(a + b \log(cx^n))) dx = \int x \operatorname{Shi}((b \log(cx^n) + a)d) dx$$

```
[In] integrate(x*Shi(d*(a+b*log(c*x^n))),x, algorithm="fricas")
```

```
[Out] integral(x*sinh_integral(b*d*log(c*x^n) + a*d), x)
```

Sympy [F]

$$\int x \operatorname{Shi}(d(a + b \log(cx^n))) dx = \int x \operatorname{Shi}(ad + bd \log(cx^n)) dx$$

```
[In] integrate(x*Shi(d*(a+b*ln(c*x**n))),x)
```

```
[Out] Integral(x*Shi(a*d + b*d*log(c*x**n)), x)
```

Maxima [F]

$$\int x \operatorname{Shi}(d(a + b \log(cx^n))) dx = \int x \operatorname{Shi}((b \log(cx^n) + a)d) dx$$

```
[In] integrate(x*Shi(d*(a+b*log(c*x^n))),x, algorithm="maxima")
```

```
[Out] integrate(x*Shi((b*log(c*x^n) + a)*d), x)
```

Giac [F]

$$\int x \operatorname{Shi}(d(a + b \log(cx^n))) dx = \int x \operatorname{Shi}((b \log(cx^n) + a)d) dx$$

[In] integrate(x*Shi(d*(a+b*log(c*x^n))),x, algorithm="giac")

[Out] integrate(x*Shi((b*log(c*x^n) + a)*d), x)

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{Shi}(d(a + b \log(cx^n))) dx = \int x \operatorname{sinhint}(d(a + b \ln(cx^n))) dx$$

[In] int(x*sinhint(d*(a + b*log(c*x^n))),x)

[Out] int(x*sinhint(d*(a + b*log(c*x^n))), x)

3.34 $\int \text{Shi}(d(a + b \log(cx^n))) dx$

Optimal result	200
Rubi [A] (verified)	200
Mathematica [A] (verified)	202
Maple [F]	202
Fricas [F]	203
Sympy [F]	203
Maxima [F]	203
Giac [F]	203
Mupad [F(-1)]	204

Optimal result

Integrand size = 13, antiderivative size = 119

$$\int \text{Shi}(d(a + b \log(cx^n))) dx$$

$$= \frac{1}{2} e^{-\frac{a}{bn}} x (cx^n)^{-1/n} \text{ExpIntegralEi} \left(\frac{(1 - bdn)(a + b \log(cx^n))}{bn} \right)$$

$$- \frac{1}{2} e^{-\frac{a}{bn}} x (cx^n)^{-1/n} \text{ExpIntegralEi} \left(\frac{(1 + bdn)(a + b \log(cx^n))}{bn} \right)$$

$$+ x \text{Shi}(d(a + b \log(cx^n)))$$

[Out] 1/2*x*Ei((-b*d*n+1)*(a+b*ln(c*x^n))/b/n)/exp(a/b/n)/((c*x^n)^(1/n))-1/2*x*Ei((b*d*n+1)*(a+b*ln(c*x^n))/b/n)/exp(a/b/n)/((c*x^n)^(1/n))+x*Shi(d*(a+b*ln(c*x^n)))

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {6687, 12, 5648, 2347, 2209}

$$\int \text{Shi}(d(a + b \log(cx^n))) dx$$

$$= \frac{1}{2} x e^{-\frac{a}{bn}} (cx^n)^{-1/n} \text{ExpIntegralEi} \left(\frac{(1 - bdn)(a + b \log(cx^n))}{bn} \right)$$

$$- \frac{1}{2} x e^{-\frac{a}{bn}} (cx^n)^{-1/n} \text{ExpIntegralEi} \left(\frac{(bdn + 1)(a + b \log(cx^n))}{bn} \right)$$

$$+ x \text{Shi}(d(a + b \log(cx^n)))$$

[In] Int[SinhIntegral[d*(a + b*Log[c*x^n]),x]


```
[Out] (x*ExpIntegralEi[((1 - b*d*n)*(a + b*Log[c*x^n])/(b*n))]/(2*E^(a/(b*n)))*(c*x^n)^n^(-1)) - (x*ExpIntegralEi[((1 + b*d*n)*(a + b*Log[c*x^n])/(b*n))]/(2*E^(a/(b*n)))*(c*x^n)^n^(-1)) + x*SinhIntegral[d*(a + b*Log[c*x^n])]
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] :=> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 2209

```
Int[(F_)^((g_)*((e_) + (f_)*(x_)))/((c_) + (d_)*(x_)), x_Symbol] :=> Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]
```

Rule 2347

```
Int[((a_) + Log[(c_)*(x_)^(n_)])*(b_)^(p_)*((d_)*(x_)^(m_)), x_Symbol] :=> Dist[(d*x)^(m + 1)/(d*n*(c*x^n)^((m + 1)/n)), Subst[Int[E^((m + 1)/n)*x*(a + b*x)^p, x], x, Log[c*x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x]
```

Rule 5648

```
Int[((e_) + Log[(g_)*(x_)^(m_)])*(f_)*(h_)^(q_)*Sinh[((a_) + Log[(c_)*(x_)^(n_)])*(b_)]*(d_), x_Symbol] :=> Dist[(-E^((-a)*d))*(1/((c*x^n)^(b*d)*(2/x^(b*d*n))))], Int[(h*(e + f*Log[g*x^m]))^q/x^(b*d*n), x], x] + Dist[E^(a*d)*((c*x^n)^(b*d)/(2*x^(b*d*n))), Int[x^(b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n, q}, x]
```

Rule 6687

```
Int[SinhIntegral[((a_) + Log[(c_)*(x_)^(n_)])*(b_)]*(d_), x_Symbol] :=> Simp[x*SinhIntegral[d*(a + b*Log[c*x^n])], x] - Dist[b*d*n, Int[Sinh[d*(a + b*Log[c*x^n])]/(d*(a + b*Log[c*x^n])), x], x] /; FreeQ[{a, b, c, d, n}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= x\text{Shi}(d(a + b \log(cx^n))) - (bdn) \int \frac{\sinh(d(a + b \log(cx^n)))}{d(a + b \log(cx^n))} dx \\
 &= x\text{Shi}(d(a + b \log(cx^n))) - (bn) \int \frac{\sinh(d(a + b \log(cx^n)))}{a + b \log(cx^n)} dx \\
 &= x\text{Shi}(d(a + b \log(cx^n))) + \frac{1}{2} \left(be^{-ad} n x^{bdn} (cx^n)^{-bd} \right) \int \frac{x^{-bdn}}{a + b \log(cx^n)} dx \\
 &\quad - \frac{1}{2} \left(be^{ad} n x^{-bdn} (cx^n)^{bd} \right) \int \frac{x^{bdn}}{a + b \log(cx^n)} dx
 \end{aligned}$$

$$\begin{aligned}
&= x\operatorname{Shi}(d(a + b \log(cx^n))) \\
&\quad + \frac{1}{2} \left(b e^{-ad} x (cx^n)^{-bd - \frac{1-bdn}{n}} \right) \operatorname{Subst} \left(\int \frac{e^{\frac{(1-bdn)x}{n}}}{a + bx} dx, x, \log(cx^n) \right) \\
&\quad - \frac{1}{2} \left(b e^{ad} x (cx^n)^{bd - \frac{1+bdn}{n}} \right) \operatorname{Subst} \left(\int \frac{e^{\frac{(1+bdn)x}{n}}}{a + bx} dx, x, \log(cx^n) \right) \\
&= \frac{1}{2} e^{-\frac{a}{bn}} x (cx^n)^{-1/n} \operatorname{ExpIntegralEi} \left(\frac{(1 - bdn)(a + b \log(cx^n))}{bn} \right) \\
&\quad - \frac{1}{2} e^{-\frac{a}{bn}} x (cx^n)^{-1/n} \operatorname{ExpIntegralEi} \left(\frac{(1 + bdn)(a + b \log(cx^n))}{bn} \right) \\
&\quad + x\operatorname{Shi}(d(a + b \log(cx^n)))
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.05 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.80

$$\begin{aligned}
&\int \operatorname{Shi}(d(a + b \log(cx^n))) dx \\
&= \frac{1}{2} e^{-\frac{a}{bn}} x (cx^n)^{-1/n} \left(\operatorname{ExpIntegralEi} \left(-\frac{(-1 + bdn)(a + b \log(cx^n))}{bn} \right) \right. \\
&\quad \left. - \operatorname{ExpIntegralEi} \left(\frac{(1 + bdn)(a + b \log(cx^n))}{bn} \right) \right) + x\operatorname{Shi}(d(a + b \log(cx^n)))
\end{aligned}$$

[In] Integrate[SinhIntegral[d*(a + b*Log[c*x^n])],x]

[Out] (x*(ExpIntegralEi[-(((1 + b*d*n)*(a + b*Log[c*x^n]))/(b*n))]) - ExpIntegralEi[(((1 + b*d*n)*(a + b*Log[c*x^n]))/(b*n))])/(2*E^(a/(b*n))*(c*x^n)^n^(-1)) + x*SinhIntegral[d*(a + b*Log[c*x^n])]

Maple [F]

$$\int \operatorname{Shi}(d(a + b \ln(cx^n))) dx$$

[In] int(Shi(d*(a+b*ln(c*x^n))),x)

[Out] int(Shi(d*(a+b*ln(c*x^n))),x)

Fricas [F]

$$\int \operatorname{Shi}(d(a + b \log(cx^n))) dx = \int \operatorname{Shi}((b \log(cx^n) + a)d) dx$$

```
[In] integrate(Shi(d*(a+b*log(c*x^n))),x, algorithm="fricas")
```

```
[Out] integral(sinh_integral(b*d*log(c*x^n) + a*d), x)
```

Sympy [F]

$$\int \operatorname{Shi}(d(a + b \log(cx^n))) dx = \int \operatorname{Shi}(d(a + b \log(cx^n))) dx$$

```
[In] integrate(Shi(d*(a+b*ln(c*x**n))),x)
```

```
[Out] Integral(Shi(d*(a + b*log(c*x**n))), x)
```

Maxima [F]

$$\int \operatorname{Shi}(d(a + b \log(cx^n))) dx = \int \operatorname{Shi}((b \log(cx^n) + a)d) dx$$

```
[In] integrate(Shi(d*(a+b*log(c*x^n))),x, algorithm="maxima")
```

```
[Out] integrate(Shi((b*log(c*x^n) + a)*d), x)
```

Giac [F]

$$\int \operatorname{Shi}(d(a + b \log(cx^n))) dx = \int \operatorname{Shi}((b \log(cx^n) + a)d) dx$$

```
[In] integrate(Shi(d*(a+b*log(c*x^n))),x, algorithm="giac")
```

```
[Out] integrate(Shi((b*log(c*x^n) + a)*d), x)
```

Mupad [F(-1)]

Timed out.

$$\int \operatorname{Shi}(d(a + b \log(cx^n))) dx = \int \operatorname{sinhint}(d(a + b \ln(cx^n))) dx$$

```
[In] int(sinhint(d*(a + b*log(c*x^n))),x)
```

```
[Out] int(sinhint(d*(a + b*log(c*x^n))), x)
```

3.35 $\int \frac{\text{Shi}(d(a+b \log(cx^n)))}{x} dx$

Optimal result	205
Rubi [A] (verified)	205
Mathematica [A] (verified)	206
Maple [A] (verified)	206
Fricas [F]	207
Sympy [F]	207
Maxima [F]	207
Giac [F]	207
Mupad [F(-1)]	208

Optimal result

Integrand size = 17, antiderivative size = 55

$$\int \frac{\text{Shi}(d(a + b \log(cx^n)))}{x} dx = -\frac{\cosh(d(a + b \log(cx^n)))}{bdn} + \frac{(a + b \log(cx^n)) \text{Shi}(d(a + b \log(cx^n)))}{bn}$$

[Out] $-\cosh(d*(a+b*\ln(c*x^n)))/b/d/n+(a+b*\ln(c*x^n))*\text{Shi}(d*(a+b*\ln(c*x^n)))/b/n$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {6663}

$$\int \frac{\text{Shi}(d(a + b \log(cx^n)))}{x} dx = \frac{(a + b \log(cx^n)) \text{Shi}(d(a + b \log(cx^n)))}{bn} - \frac{\cosh(d(a + b \log(cx^n)))}{bdn}$$

[In] $\text{Int}[\text{SinhIntegral}[d*(a + b*\text{Log}[c*x^n])]/x, x]$

[Out] $-(\text{Cosh}[d*(a + b*\text{Log}[c*x^n])]/(b*d*n)) + ((a + b*\text{Log}[c*x^n])*\text{SinhIntegral}[d*(a + b*\text{Log}[c*x^n])])/b/n$

Rule 6663

$\text{Int}[\text{SinhIntegral}[(a_.) + (b_.)*(x_.)], x_Symbol] \text{ :> } \text{Simp}[(a + b*x)*(\text{SinhIntegral}[a + b*x]/b), x] - \text{Simp}[\text{Cosh}[a + b*x]/b, x] \text{ ;/; } \text{FreeQ}\{a, b\}, x]$

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Subst}\left(\int \text{Shi}(d(a + bx)) dx, x, \log(cx^n)\right)}{n} \\
&= \frac{\text{Subst}\left(\int \text{Shi}(x) dx, x, ad + bd \log(cx^n)\right)}{bdn} \\
&= -\frac{\cosh(ad + bd \log(cx^n))}{bdn} + \frac{(a + b \log(cx^n)) \text{Shi}(ad + bd \log(cx^n))}{bn}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.75

$$\int \frac{\text{Shi}(d(a + b \log(cx^n)))}{x} dx = -\frac{\cosh(ad) \cosh(bd \log(cx^n))}{bdn} - \frac{\sinh(ad) \sinh(bd \log(cx^n))}{bdn} + \frac{\log(cx^n) \text{Shi}(d(a + b \log(cx^n)))}{n} + \frac{a \text{Shi}(ad + bd \log(cx^n))}{bn}$$

[In] Integrate[SinhIntegral[d*(a + b*Log[c*x^n])]/x,x]

[Out] -((Cosh[a*d]*Cosh[b*d*Log[c*x^n]])/(b*d*n)) - (Sinh[a*d]*Sinh[b*d*Log[c*x^n]])/(b*d*n) + (Log[c*x^n]*SinhIntegral[d*(a + b*Log[c*x^n])])/n + (a*SinhIntegral[a*d + b*d*Log[c*x^n]])/(b*n)

Maple [A] (verified)

Time = 1.46 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.02

method	result
derivativedivides	$\frac{\text{Shi}(ad+bd \ln(cx^n))(ad+bd \ln(cx^n))-\cosh(ad+bd \ln(cx^n))}{ndb}$
default	$\frac{\text{Shi}(ad+bd \ln(cx^n))(ad+bd \ln(cx^n))-\cosh(ad+bd \ln(cx^n))}{ndb}$
parts	$\ln(x) \text{Shi}(d(a + b \ln(cx^n))) - nb \left(-\frac{(\ln(cx^n) - n \ln(x)) \text{Shi}(\ln(x) bdn + d(b(\ln(cx^n) - n \ln(x)) + a))}{bn^2} - \frac{a \text{Shi}(ad + bd \ln(cx^n))}{bn} \right)$

[In] int(Shi(d*(a+b*ln(c*x^n)))/x,x,method=_RETURNVERBOSE)

[Out] 1/n/d/b*(Shi(a*d+b*d*ln(c*x^n))*(a*d+b*d*ln(c*x^n))-cosh(a*d+b*d*ln(c*x^n)))

Fricas [F]

$$\int \frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{x} dx = \int \frac{\operatorname{Shi}((b \log(cx^n) + a)d)}{x} dx$$

[In] integrate(Shi(d*(a+b*log(c*x^n)))/x,x, algorithm="fricas")

[Out] integral(sinh_integral(b*d*log(c*x^n) + a*d)/x, x)

Sympy [F]

$$\int \frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{x} dx = \int \frac{\operatorname{Shi}(ad + bd \log(cx^n))}{x} dx$$

[In] integrate(Shi(d*(a+b*ln(c*x**n)))/x,x)

[Out] Integral(Shi(a*d + b*d*log(c*x**n))/x, x)

Maxima [F]

$$\int \frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{x} dx = \int \frac{\operatorname{Shi}((b \log(cx^n) + a)d)}{x} dx$$

[In] integrate(Shi(d*(a+b*log(c*x^n)))/x,x, algorithm="maxima")

[Out] integrate(Shi((b*log(c*x^n) + a)*d)/x, x)

Giac [F]

$$\int \frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{x} dx = \int \frac{\operatorname{Shi}((b \log(cx^n) + a)d)}{x} dx$$

[In] integrate(Shi(d*(a+b*log(c*x^n)))/x,x, algorithm="giac")

[Out] integrate(Shi((b*log(c*x^n) + a)*d)/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\text{Shi}(d(a + b \log(cx^n)))}{x} dx = \frac{\text{sinhint}(d(a + b \ln(cx^n))) \ln(cx^n)}{n} + \frac{a \text{sinhint}(d(a + b \ln(cx^n)))}{bn} - \frac{e^{ad}(cx^n)^{bd}}{2bdn} - \frac{e^{-ad}}{2bdn(cx^n)^{bd}}$$

```
[In] int(sinhint(d*(a + b*log(c*x^n)))/x,x)
```

```
[Out] (sinhint(d*(a + b*log(c*x^n))*log(c*x^n))/n + (a*sinhint(d*(a + b*log(c*x^n))))/(b*n) - (exp(a*d)*(c*x^n)^(b*d))/(2*b*d*n) - exp(-a*d)/(2*b*d*n*(c*x^n)^(b*d)))
```


3.36 $\int \frac{\text{Shi}(d(a+b \log(cx^n)))}{x^2} dx$

Optimal result	209
Rubi [A] (verified)	209
Mathematica [A] (verified)	211
Maple [F]	212
Fricas [F]	212
Sympy [F]	212
Maxima [F]	212
Giac [F]	213
Mupad [F(-1)]	213

Optimal result

Integrand size = 17, antiderivative size = 122

$$\int \frac{\text{Shi}(d(a+b \log(cx^n)))}{x^2} dx = \frac{e^{\frac{a}{bn}} (cx^n)^{\frac{1}{n}} \text{ExpIntegralEi}\left(-\frac{(1-bdn)(a+b \log(cx^n))}{bn}\right)}{2x} - \frac{e^{\frac{a}{bn}} (cx^n)^{\frac{1}{n}} \text{ExpIntegralEi}\left(-\frac{(1+bdn)(a+b \log(cx^n))}{bn}\right)}{2x} - \frac{\text{Shi}(d(a+b \log(cx^n)))}{x}$$

[Out] 1/2*exp(a/b/n)*(c*x^n)^(1/n)*Ei(-(-b*d*n+1)*(a+b*ln(c*x^n))/b/n)/x-1/2*exp(a/b/n)*(c*x^n)^(1/n)*Ei(-(b*d*n+1)*(a+b*ln(c*x^n))/b/n)/x-Shi(d*(a+b*ln(c*x^n)))/x

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {6690, 12, 5650, 2347, 2209}

$$\int \frac{\text{Shi}(d(a+b \log(cx^n)))}{x^2} dx = \frac{e^{\frac{a}{bn}} (cx^n)^{\frac{1}{n}} \text{ExpIntegralEi}\left(-\frac{(1-bdn)(a+b \log(cx^n))}{bn}\right)}{2x} - \frac{e^{\frac{a}{bn}} (cx^n)^{\frac{1}{n}} \text{ExpIntegralEi}\left(-\frac{(bdn+1)(a+b \log(cx^n))}{bn}\right)}{2x} - \frac{\text{Shi}(d(a+b \log(cx^n)))}{x}$$

[In] Int[SinhIntegral[d*(a + b*Log[c*x^n])]/x^2,x]

```
[Out] (E^(a/(b*n))*(c*x^n)^n^(-1)*ExpIntegralEi[-(((1 - b*d*n)*(a + b*Log[c*x^n])
)/(b*n))])/(2*x) - (E^(a/(b*n))*(c*x^n)^n^(-1)*ExpIntegralEi[-(((1 + b*d*n)
*(a + b*Log[c*x^n]))/(b*n))])/(2*x) - SinhIntegral[d*(a + b*Log[c*x^n])/x
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 2209

```
Int[(F_)^((g_)*((e_) + (f_)*(x_)))/((c_) + (d_)*(x_)), x_Symbol] := Si
mp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; F
reeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]
```

Rule 2347

```
Int[((a_) + Log[(c_)*(x_)^(n_)])*(b_)^(p_)*((d_)*(x_)^(m_)), x_Symbol
] := Dist[(d*x)^(m + 1)/(d*n*(c*x^n)^((m + 1)/n)), Subst[Int[E^(((m + 1)/n)
*x)*(a + b*x)^p, x], x, Log[c*x^n]], x] /; FreeQ[{a, b, c, d, m, n, p}, x]
```

Rule 5650

```
Int[(((e_) + Log[(g_)*(x_)^(m_)])*(f_)*(h_)^(q_)*((i_)*(x_)^(r_))*
Sinh[((a_) + Log[(c_)*(x_)^(n_)])*(b_)]*(d_)), x_Symbol] := Dist[(-E^((
-a)*d))*(i*x)^r*(1/((c*x^n)^(b*d)*(2*x^(r - b*d*n))))], Int[x^(r - b*d*n)*(h
*(e + f*Log[g*x^m]))^q, x], x] + Dist[E^(a*d)*(i*x)^r*((c*x^n)^(b*d)/(2*x^(
r + b*d*n))), Int[x^(r + b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] /; FreeQ[{
a, b, c, d, e, f, g, h, i, m, n, q, r}, x]
```

Rule 6690

```
Int[((e_)*(x_)^(m_))*SinhIntegral[((a_) + Log[(c_)*(x_)^(n_)])*(b_)]*(
d_)], x_Symbol] := Simp[(e*x)^(m + 1)*(SinhIntegral[d*(a + b*Log[c*x^n])]/
(e*(m + 1))), x] - Dist[b*d*(n/(m + 1)), Int[(e*x)^m*(Sinh[d*(a + b*Log[c*x
^n])]/(d*(a + b*Log[c*x^n]))], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] &&
NeQ[m, -1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Shi}(d(a + b \log(cx^n)))}{x} + (bdn) \int \frac{\sinh(d(a + b \log(cx^n)))}{dx^2(a + b \log(cx^n))} dx \\ &= -\frac{\text{Shi}(d(a + b \log(cx^n)))}{x} + (bn) \int \frac{\sinh(d(a + b \log(cx^n)))}{x^2(a + b \log(cx^n))} dx \end{aligned}$$

$$\begin{aligned}
&= -\frac{\text{Shi}(d(a + b \log(cx^n)))}{x} - \frac{1}{2} \left(b e^{-ad} n x^{bdn} (cx^n)^{-bd} \right) \int \frac{x^{-2-bdn}}{a + b \log(cx^n)} dx \\
&\quad + \frac{1}{2} \left(b e^{ad} n x^{-bdn} (cx^n)^{bd} \right) \int \frac{x^{-2+bdn}}{a + b \log(cx^n)} dx \\
&= -\frac{\text{Shi}(d(a + b \log(cx^n)))}{x} \\
&\quad - \frac{\left(b e^{-ad} (cx^n)^{-bd - \frac{-1-bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(-1-bdn)x}{a+bx}}}{a+bx} dx, x, \log(cx^n) \right)}{2x} \\
&\quad + \frac{\left(b e^{ad} (cx^n)^{bd - \frac{-1+bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(-1+bdn)x}{a+bx}}}{a+bx} dx, x, \log(cx^n) \right)}{2x} \\
&= \frac{e^{\frac{a}{bn}} (cx^n)^{\frac{1}{n}} \text{ExpIntegralEi} \left(-\frac{(1-bdn)(a+b \log(cx^n))}{bn} \right)}{2x} \\
&\quad - \frac{e^{\frac{a}{bn}} (cx^n)^{\frac{1}{n}} \text{ExpIntegralEi} \left(-\frac{(1+bdn)(a+b \log(cx^n))}{bn} \right)}{2x} - \frac{\text{Shi}(d(a + b \log(cx^n)))}{x}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.25 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.20

$$\begin{aligned}
&\int \frac{\text{Shi}(d(a + b \log(cx^n)))}{x^2} dx \\
&= \frac{1}{2} e^{-\frac{(-1+bdn)(a+b(-n \log(x)+\log(cx^n)))}{bn}} \left(\text{ExpIntegralEi} \left(\frac{(-1+bdn)(a+b \log(cx^n))}{bn} \right) \right. \\
&\quad \left. - \text{ExpIntegralEi} \left(-\frac{(1+bdn)(a+b \log(cx^n))}{bn} \right) \right) (\cosh(d(a+b(-n \log(x)+\log(cx^n)))) \\
&\quad + \sinh(d(a+b(-n \log(x)+\log(cx^n)))))) - \frac{\text{Shi}(d(a + b \log(cx^n)))}{x}
\end{aligned}$$

[In] Integrate[SinhIntegral[d*(a + b*Log[c*x^n])/x^2,x]

[Out] ((ExpIntegralEi[((-1 + b*d*n)*(a + b*Log[c*x^n]))/(b*n)] - ExpIntegralEi[-((1 + b*d*n)*(a + b*Log[c*x^n]))/(b*n)])*(Cosh[d*(a + b*(-n*Log[x]) + Log[c*x^n]))] + Sinh[d*(a + b*(-n*Log[x]) + Log[c*x^n]))])/(2*E^(((-1 + b*d*n)*(a + b*(-n*Log[x]) + Log[c*x^n]))/(b*n))) - SinhIntegral[d*(a + b*Log[c*x^n])/x]

Maple [F]

$$\int \frac{\operatorname{Shi}(d(a + b \ln(cx^n)))}{x^2} dx$$

[In] `int(Shi(d*(a+b*ln(c*x^n)))/x^2,x)`

[Out] `int(Shi(d*(a+b*ln(c*x^n)))/x^2,x)`

Fricas [F]

$$\int \frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\operatorname{Shi}((b \log(cx^n) + a)d)}{x^2} dx$$

[In] `integrate(Shi(d*(a+b*log(c*x^n)))/x^2,x, algorithm="fricas")`

[Out] `integral(sinh_integral(b*d*log(c*x^n) + a*d)/x^2, x)`

Sympy [F]

$$\int \frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\operatorname{Shi}(ad + bd \log(cx^n))}{x^2} dx$$

[In] `integrate(Shi(d*(a+b*ln(c*x**n)))/x**2,x)`

[Out] `Integral(Shi(a*d + b*d*log(c*x**n))/x**2, x)`

Maxima [F]

$$\int \frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\operatorname{Shi}((b \log(cx^n) + a)d)}{x^2} dx$$

[In] `integrate(Shi(d*(a+b*log(c*x^n)))/x^2,x, algorithm="maxima")`

[Out] `integrate(Shi((b*log(c*x^n) + a)*d)/x^2, x)`

Giac [F]

$$\int \frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\operatorname{Shi}((b \log(cx^n) + a)d)}{x^2} dx$$

[In] integrate(Shi(d*(a+b*log(c*x^n)))/x^2,x, algorithm="giac")

[Out] integrate(Shi((b*log(c*x^n) + a)*d)/x^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\operatorname{sinhint}(d(a + b \ln(cx^n)))}{x^2} dx$$

[In] int(sinhint(d*(a + b*log(c*x^n)))/x^2,x)

[Out] int(sinhint(d*(a + b*log(c*x^n)))/x^2, x)

3.37 $\int \frac{\text{Shi}(d(a+b \log(cx^n)))}{x^3} dx$

Optimal result	214
Rubi [A] (verified)	214
Mathematica [A] (verified)	216
Maple [F]	217
Fricas [F]	217
Sympy [F]	217
Maxima [F]	217
Giac [F]	218
Mupad [F(-1)]	218

Optimal result

Integrand size = 17, antiderivative size = 130

$$\int \frac{\text{Shi}(d(a+b \log(cx^n)))}{x^3} dx = \frac{e^{\frac{2a}{bn}} (cx^n)^{2/n} \text{ExpIntegralEi}\left(-\frac{(2-bdn)(a+b \log(cx^n))}{bn}\right)}{4x^2} - \frac{e^{\frac{2a}{bn}} (cx^n)^{2/n} \text{ExpIntegralEi}\left(-\frac{(2+bdn)(a+b \log(cx^n))}{bn}\right)}{4x^2} - \frac{\text{Shi}(d(a+b \log(cx^n)))}{2x^2}$$

[Out] $\frac{1}{4} \exp(2a/b/n) * (c*x^n)^{(2/n)} * \text{Ei}(-(-b*d*n+2)*(a+b*\ln(c*x^n))/b/n) / x^{2-1/4} * \exp(2a/b/n) * (c*x^n)^{(2/n)} * \text{Ei}(-(b*d*n+2)*(a+b*\ln(c*x^n))/b/n) / x^{2-1/2} * \text{Shi}(d*(a+b*\ln(c*x^n))) / x^2$

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {6690, 12, 5650, 2347, 2209}

$$\int \frac{\text{Shi}(d(a+b \log(cx^n)))}{x^3} dx = \frac{e^{\frac{2a}{bn}} (cx^n)^{2/n} \text{ExpIntegralEi}\left(-\frac{(2-bdn)(a+b \log(cx^n))}{bn}\right)}{4x^2} - \frac{e^{\frac{2a}{bn}} (cx^n)^{2/n} \text{ExpIntegralEi}\left(-\frac{(bdn+2)(a+b \log(cx^n))}{bn}\right)}{4x^2} - \frac{\text{Shi}(d(a+b \log(cx^n)))}{2x^2}$$

[In] $\text{Int}[\text{SinhIntegral}[d*(a + b*\text{Log}[c*x^n])]/x^3, x]$

```
[Out] (E^((2*a)/(b*n))*(c*x^n)^(2/n)*ExpIntegralEi[-(((2 - b*d*n)*(a + b*Log[c*x^n]))/(b*n))])/(4*x^2) - (E^((2*a)/(b*n))*(c*x^n)^(2/n)*ExpIntegralEi[-(((2 + b*d*n)*(a + b*Log[c*x^n]))/(b*n))])/(4*x^2) - SinhIntegral[d*(a + b*Log[c*x^n])]/(2*x^2)
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 2209

```
Int[(F_)^((g_)*((e_) + (f_)*(x_)))/((c_) + (d_)*(x_)), x_Symbol] := Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]
```

Rule 2347

```
Int[((a_) + Log[(c_)*(x_)^(n_)])*(b_)^(p_)*((d_)*(x_))^(m_), x_Symbol] := Dist[(d*x)^(m + 1)/(d*n*(c*x^n)^(m + 1/n)), Subst[Int[E^((m + 1)/n)*x]*(a + b*x)^p, x], x, Log[c*x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x]
```

Rule 5650

```
Int[(((e_) + Log[(g_)*(x_)^(m_)])*(f_))*(h_)^(q_)*((i_)*(x_))^(r_)*Sinh[((a_) + Log[(c_)*(x_)^(n_)])*(b_)]*(d_), x_Symbol] := Dist[(-E^((-a)*d))*(i*x)^r*(1/((c*x^n)^(b*d)*(2*x^(r - b*d*n))))], Int[x^(r - b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] + Dist[E^(a*d)*(i*x)^r*((c*x^n)^(b*d)/(2*x^(r + b*d*n))), Int[x^(r + b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, m, n, q, r}, x]
```

Rule 6690

```
Int[((e_)*(x_))^(m_)*SinhIntegral[((a_) + Log[(c_)*(x_)^(n_)])*(b_)]*(d_), x_Symbol] := Simp[(e*x)^(m + 1)*(SinhIntegral[d*(a + b*Log[c*x^n])]/(e*(m + 1))), x] - Dist[b*d*(n/(m + 1)), Int[(e*x)^m*(Sinh[d*(a + b*Log[c*x^n])]/(d*(a + b*Log[c*x^n]))], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Shi}(d(a + b \log(cx^n)))}{2x^2} + \frac{1}{2}(bdn) \int \frac{\sinh(d(a + b \log(cx^n)))}{dx^3 (a + b \log(cx^n))} dx \\ &= -\frac{\text{Shi}(d(a + b \log(cx^n)))}{2x^2} + \frac{1}{2}(bn) \int \frac{\sinh(d(a + b \log(cx^n)))}{x^3 (a + b \log(cx^n))} dx \end{aligned}$$

$$\begin{aligned}
&= -\frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{2x^2} - \frac{1}{4} \left(be^{-ad} n x^{bdn} (cx^n)^{-bd} \right) \int \frac{x^{-3-bdn}}{a + b \log(cx^n)} dx \\
&\quad + \frac{1}{4} \left(be^{ad} n x^{-bdn} (cx^n)^{bd} \right) \int \frac{x^{-3+bdn}}{a + b \log(cx^n)} dx \\
&= -\frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{2x^2} \\
&\quad - \frac{\left(be^{-ad} (cx^n)^{-bd - \frac{-2-bdn}{n}} \right) \operatorname{Subst} \left(\int \frac{e^{\frac{(-2-bdn)x}{a+bx}}}{a+bx} dx, x, \log(cx^n) \right)}{4x^2} \\
&\quad + \frac{\left(be^{ad} (cx^n)^{bd - \frac{-2+bdn}{n}} \right) \operatorname{Subst} \left(\int \frac{e^{\frac{(-2+bdn)x}{a+bx}}}{a+bx} dx, x, \log(cx^n) \right)}{4x^2} \\
&= \frac{e^{\frac{2a}{bn}} (cx^n)^{2/n} \operatorname{ExpIntegralEi} \left(-\frac{(2-bdn)(a+b \log(cx^n))}{bn} \right)}{4x^2} \\
&\quad - \frac{e^{\frac{2a}{bn}} (cx^n)^{2/n} \operatorname{ExpIntegralEi} \left(-\frac{(2+bdn)(a+b \log(cx^n))}{bn} \right)}{4x^2} - \frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{2x^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.25 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.14

$$\begin{aligned}
&\int \frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{x^3} dx \\
&= \frac{1}{4} e^{-\frac{(-2+bdn)(a+b(-n \log(x)+\log(cx^n)))}{bn}} \left(\operatorname{ExpIntegralEi} \left(\frac{(-2+bdn)(a+b \log(cx^n))}{bn} \right) \right. \\
&\quad \left. - \operatorname{ExpIntegralEi} \left(-\frac{(2+bdn)(a+b \log(cx^n))}{bn} \right) \right) (\cosh(d(a+b(-n \log(x)+\log(cx^n)))) \\
&\quad \quad + \sinh(d(a+b(-n \log(x)+\log(cx^n)))))) - \frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{2x^2}
\end{aligned}$$

[In] Integrate[SinhIntegral[d*(a + b*Log[c*x^n])/x^3,x]

[Out] ((ExpIntegralEi[(-2 + b*d*n)*(a + b*Log[c*x^n])]/(b*n)] - ExpIntegralEi[-((2 + b*d*n)*(a + b*Log[c*x^n])]/(b*n))])*(Cosh[d*(a + b*(-n*Log[x]) + Log[c*x^n]))] + Sinh[d*(a + b*(-n*Log[x]) + Log[c*x^n]))])/(4*E^(((-2 + b*d*n)*(a + b*(-n*Log[x]) + Log[c*x^n]))/(b*n))) - SinhIntegral[d*(a + b*Log[c*x^n])/x^2]

Maple [F]

$$\int \frac{\text{Shi}(d(a + b \ln(cx^n)))}{x^3} dx$$

```
[In] int(Shi(d*(a+b*ln(c*x^n)))/x^3,x)
```

```
[Out] int(Shi(d*(a+b*ln(c*x^n)))/x^3,x)
```

Fricas [F]

$$\int \frac{\text{Shi}(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\text{Shi}((b \log(cx^n) + a)d)}{x^3} dx$$

```
[In] integrate(Shi(d*(a+b*log(c*x^n)))/x^3,x, algorithm="fricas")
```

```
[Out] integral(sinh_integral(b*d*log(c*x^n) + a*d)/x^3, x)
```

Sympy [F]

$$\int \frac{\text{Shi}(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\text{Shi}(ad + bd \log(cx^n))}{x^3} dx$$

```
[In] integrate(Shi(d*(a+b*ln(c*x**n)))/x**3,x)
```

```
[Out] Integral(Shi(a*d + b*d*log(c*x**n))/x**3, x)
```

Maxima [F]

$$\int \frac{\text{Shi}(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\text{Shi}((b \log(cx^n) + a)d)}{x^3} dx$$

```
[In] integrate(Shi(d*(a+b*log(c*x^n)))/x^3,x, algorithm="maxima")
```

```
[Out] integrate(Shi((b*log(c*x^n) + a)*d)/x^3, x)
```

Giac [F]

$$\int \frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\operatorname{Shi}((b \log(cx^n) + a)d)}{x^3} dx$$

[In] integrate(Shi(d*(a+b*log(c*x^n)))/x^3,x, algorithm="giac")

[Out] integrate(Shi((b*log(c*x^n) + a)*d)/x^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{Shi}(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\operatorname{sinhint}(d(a + b \ln(cx^n)))}{x^3} dx$$

[In] int(sinhint(d*(a + b*log(c*x^n)))/x^3,x)

[Out] int(sinhint(d*(a + b*log(c*x^n)))/x^3, x)

3.38 $\int (ex)^m \text{Shi}(d(a + b \log(cx^n))) dx$

Optimal result	219
Rubi [A] (verified)	219
Mathematica [A] (verified)	222
Maple [F]	222
Fricas [F]	222
Sympy [F]	222
Maxima [F]	223
Giac [F]	223
Mupad [F(-1)]	223

Optimal result

Integrand size = 19, antiderivative size = 167

$$\int (ex)^m \text{Shi}(d(a + b \log(cx^n))) dx$$

$$= \frac{e^{-\frac{a(1+m)}{bn}} x (ex)^m (cx^n)^{-\frac{1+m}{n}} \text{ExpIntegralEi}\left(\frac{(1+m-bdn)(a+b \log(cx^n))}{bn}\right)}{2(1+m)} - \frac{e^{-\frac{a(1+m)}{bn}} x (ex)^m (cx^n)^{-\frac{1+m}{n}} \text{ExpIntegralEi}\left(\frac{(1+m+bdn)(a+b \log(cx^n))}{bn}\right)}{2(1+m)} + \frac{(ex)^{1+m} \text{Shi}(d(a + b \log(cx^n)))}{e(1+m)}$$

```
[Out] 1/2*x*(e*x)^(m+1)*Ei((-b*d*n+m+1)*(a+b*ln(c*x^n))/b/n)/exp(a*(1+m)/b/n)/(1+m)/((c*x^n)^((1+m)/n))-1/2*x*(e*x)^(m+1)*Ei((b*d*n+m+1)*(a+b*ln(c*x^n))/b/n)/exp(a*(1+m)/b/n)/(1+m)/((c*x^n)^((1+m)/n))+(e*x)^(1+m)*Shi(d*(a+b*ln(c*x^n)))/e/(1+m)
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used

= {6690, 12, 5650, 2347, 2209}

$$\int (ex)^m \text{Shi}(d(a + b \log(cx^n))) dx$$

$$= \frac{x(ex)^m e^{-\frac{a(m+1)}{bn}} (cx^n)^{-\frac{m+1}{n}} \text{ExpIntegralEi}\left(\frac{(m-bdn+1)(a+b \log(cx^n))}{bn}\right)}{2(m+1)} - \frac{x(ex)^m e^{-\frac{a(m+1)}{bn}} (cx^n)^{-\frac{m+1}{n}} \text{ExpIntegralEi}\left(\frac{(m+bdn+1)(a+b \log(cx^n))}{bn}\right)}{2(m+1)} + \frac{(ex)^{m+1} \text{Shi}(d(a + b \log(cx^n)))}{e(m+1)}$$

[In] Int[(e*x)^m*SinhIntegral[d*(a + b*Log[c*x^n]),x]

[Out] (x*(e*x)^m*ExpIntegralEi[((1 + m - b*d*n)*(a + b*Log[c*x^n])/(b*n))]/(2*E^((a*(1 + m))/(b*n))*(1 + m)*(c*x^n)^((1 + m)/n)) - (x*(e*x)^m*ExpIntegralEi[((1 + m + b*d*n)*(a + b*Log[c*x^n])/(b*n))]/(2*E^((a*(1 + m))/(b*n))*(1 + m)*(c*x^n)^((1 + m)/n)) + ((e*x)^(1 + m)*SinhIntegral[d*(a + b*Log[c*x^n])])/(e*(1 + m))

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 2209

Int[(F_)^((g_)*((e_) + (f_)*(x_)))/((c_) + (d_)*(x_)), x_Symbol] := Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[\$UseGamma]

Rule 2347

Int[((a_) + Log[(c_)*(x_)^(n_)])*(b_)^(p_)*((d_)*(x_)^(m_)), x_Symbol] := Dist[(d*x)^(m + 1)/(d*n*(c*x^n)^((m + 1)/n)), Subst[Int[E^(((m + 1)/n)*x)*(a + b*x)^p, x], x, Log[c*x^n]], x] /; FreeQ[{a, b, c, d, m, n, p}, x]

Rule 5650

Int[(((e_) + Log[(g_)*(x_)^(m_)])*(f_))*(h_)^(q_)*((i_)*(x_)^(r_))*Sinh[((a_) + Log[(c_)*(x_)^(n_)])*(b_)]*(d_), x_Symbol] := Dist[(-E^((-a*d))*(i*x)^r*(1/((c*x^n)^(b*d)*(2*x^(r - b*d*n))))), Int[x^(r - b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] + Dist[E^(a*d)*(i*x)^r*((c*x^n)^(b*d)/(2*x^(r + b*d*n))), Int[x^(r + b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, m, n, q, r}, x]

Rule 6690

Int[((e_.)*(x_))^(m_.)*SinhIntegral[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)], x_Symbol] :> Simp[(e*x)^(m + 1)*(SinhIntegral[d*(a + b*Log[c*x^n])]/(e*(m + 1))), x] - Dist[b*d*(n/(m + 1)), Int[(e*x)^m*(Sinh[d*(a + b*Log[c*x^n])])]/(d*(a + b*Log[c*x^n]))], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(ex)^{1+m} \text{Shi}(d(a + b \log(cx^n)))}{e(1+m)} - \frac{(bdn) \int \frac{(ex)^m \sinh(d(a+b \log(cx^n)))}{d(a+b \log(cx^n))} dx}{1+m} \\
&= \frac{(ex)^{1+m} \text{Shi}(d(a + b \log(cx^n)))}{e(1+m)} - \frac{(bn) \int \frac{(ex)^m \sinh(d(a+b \log(cx^n)))}{a+b \log(cx^n)} dx}{1+m} \\
&= \frac{(ex)^{1+m} \text{Shi}(d(a + b \log(cx^n)))}{e(1+m)} + \frac{\left(be^{-ad} n x^{-m+bdn} (ex)^m (cx^n)^{-bd} \right) \int \frac{x^{m-bdn}}{a+b \log(cx^n)} dx}{2(1+m)} \\
&\quad - \frac{\left(be^{ad} n x^{-m-bdn} (ex)^m (cx^n)^{bd} \right) \int \frac{x^{m+bdn}}{a+b \log(cx^n)} dx}{2(1+m)} \\
&= \frac{(ex)^{1+m} \text{Shi}(d(a + b \log(cx^n)))}{e(1+m)} \\
&\quad + \frac{\left(be^{-ad} x (ex)^m (cx^n)^{-bd - \frac{1+m-bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(1+m-bdn)x}{a+bx}}}{a+bx} dx, x, \log(cx^n) \right)}{2(1+m)} \\
&\quad - \frac{\left(be^{ad} x (ex)^m (cx^n)^{bd - \frac{1+m+bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(1+m+bdn)x}{a+bx}}}{a+bx} dx, x, \log(cx^n) \right)}{2(1+m)} \\
&= \frac{e^{-\frac{a(1+m)}{bn}} x (ex)^m (cx^n)^{-\frac{1+m}{n}} \text{ExpIntegralEi} \left(\frac{(1+m-bdn)(a+b \log(cx^n))}{bn} \right)}{2(1+m)} \\
&\quad - \frac{e^{-\frac{a(1+m)}{bn}} x (ex)^m (cx^n)^{-\frac{1+m}{n}} \text{ExpIntegralEi} \left(\frac{(1+m+bdn)(a+b \log(cx^n))}{bn} \right)}{2(1+m)} \\
&\quad + \frac{(ex)^{1+m} \text{Shi}(d(a + b \log(cx^n)))}{e(1+m)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.98 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.72

$$\int (ex)^m \text{Shi}(d(a + b \log(cx^n))) dx$$

$$= \frac{(ex)^m \left(e^{-\frac{(1+m)(a-bn \log(x)+b \log(cx^n))}{bn}} x^{-m} \left(\text{ExpIntegralEi} \left(\frac{(1+m-bdn)(a+b \log(cx^n))}{bn} \right) - \text{ExpIntegralEi} \left(\frac{(1+m+bdn)(a+b \log(cx^n))}{bn} \right) \right) \right)}{2(1+m)}$$

[In] Integrate[(e*x)^m*SinhIntegral[d*(a + b*Log[c*x^n]),x]

[Out] ((e*x)^m*((ExpIntegralEi[((1 + m - b*d*n)*(a + b*Log[c*x^n]))/(b*n)] - ExpIntegralEi[((1 + m + b*d*n)*(a + b*Log[c*x^n]))/(b*n)]))/(E^(((1 + m)*(a - b*n*Log[x] + b*Log[c*x^n]))/(b*n))*x^m) + 2*x*SinhIntegral[d*(a + b*Log[c*x^n])]))/(2*(1 + m))

Maple [F]

$$\int (ex)^m \text{Shi}(d(a + b \ln(cx^n))) dx$$

[In] int((e*x)^m*Shi(d*(a+b*ln(c*x^n))),x)

[Out] int((e*x)^m*Shi(d*(a+b*ln(c*x^n))),x)

Fricas [F]

$$\int (ex)^m \text{Shi}(d(a + b \log(cx^n))) dx = \int (ex)^m \text{Shi}((b \log(cx^n) + a)d) dx$$

[In] integrate((e*x)^m*Shi(d*(a+b*log(c*x^n))),x, algorithm="fricas")

[Out] integral((e*x)^m*sinh_integral(b*d*log(c*x^n) + a*d), x)

Sympy [F]

$$\int (ex)^m \text{Shi}(d(a + b \log(cx^n))) dx = \int (ex)^m \text{Shi}(ad + bd \log(cx^n)) dx$$

[In] integrate((e*x)**m*Shi(d*(a+b*ln(c*x**n))),x)

[Out] Integral((e*x)**m*Shi(a*d + b*d*log(c*x**n)), x)

Maxima [F]

$$\int (ex)^m \text{Shi}(d(a + b \log(cx^n))) dx = \int (ex)^m \text{Shi}((b \log(cx^n) + a)d) dx$$

[In] integrate((e*x)^m*Shi(d*(a+b*log(c*x^n))),x, algorithm="maxima")

[Out] integrate((e*x)^m*Shi((b*log(c*x^n) + a)*d), x)

Giac [F]

$$\int (ex)^m \text{Shi}(d(a + b \log(cx^n))) dx = \int (ex)^m \text{Shi}((b \log(cx^n) + a)d) dx$$

[In] integrate((e*x)^m*Shi(d*(a+b*log(c*x^n))),x, algorithm="giac")

[Out] integrate((e*x)^m*Shi((b*log(c*x^n) + a)*d), x)

Mupad [F(-1)]

Timed out.

$$\int (ex)^m \text{Shi}(d(a + b \log(cx^n))) dx = \int \text{sinhint}(d(a + b \ln(cx^n))) (ex)^m dx$$

[In] int(sinhint(d*(a + b*log(c*x^n)))*(e*x)^m,x)

[Out] int(sinhint(d*(a + b*log(c*x^n)))*(e*x)^m, x)

3.39 $\int \frac{\sinh(bx)\mathbf{Shi}(bx)}{x^3} dx$

Optimal result	224
Rubi [A] (verified)	224
Mathematica [A] (verified)	227
Maple [F]	227
Fricas [F]	227
Sympy [F]	228
Maxima [F]	228
Giac [F]	228
Mupad [F(-1)]	228

Optimal result

Integrand size = 12, antiderivative size = 96

$$\int \frac{\sinh(bx)\mathbf{Shi}(bx)}{x^3} dx = b^2\mathbf{Chi}(2bx) - \frac{b \cosh(bx) \sinh(bx)}{2x} - \frac{\sinh^2(bx)}{4x^2} - \frac{b \sinh(2bx)}{4x} \\ - \frac{b \cosh(bx)\mathbf{Shi}(bx)}{2x} - \frac{\sinh(bx)\mathbf{Shi}(bx)}{2x^2} + \frac{1}{4}b^2\mathbf{Shi}(bx)^2$$

[Out] $b^2\mathbf{Chi}(2bx) - 1/2b\cosh(bx)*\mathbf{Shi}(bx)/x + 1/4b^2\mathbf{Shi}(bx)^2 - 1/2b\cosh(bx)*\sinh(bx)/x - 1/2\mathbf{Shi}(bx)*\sinh(bx)/x^2 - 1/4\sinh(bx)^2/x^2 - 1/4b*\sinh(2bx)/x$

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {6679, 6685, 6818, 12, 5556, 3378, 3382, 3395, 29, 3393}

$$\int \frac{\sinh(bx)\mathbf{Shi}(bx)}{x^3} dx = b^2\mathbf{Chi}(2bx) + \frac{1}{4}b^2\mathbf{Shi}(bx)^2 - \frac{\mathbf{Shi}(bx) \sinh(bx)}{2x^2} - \frac{b\mathbf{Shi}(bx) \cosh(bx)}{2x} \\ - \frac{\sinh^2(bx)}{4x^2} - \frac{b \sinh(2bx)}{4x} - \frac{b \sinh(bx) \cosh(bx)}{2x}$$

[In] $\text{Int}[(\text{Sinh}[bx]*\text{SinhIntegral}[bx])/x^3, x]$

[Out] $b^2\text{CoshIntegral}[2bx] - (b\text{Cosh}[bx]*\text{Sinh}[bx])/(2x) - \text{Sinh}[bx]^2/(4x^2) - (b\text{Sinh}[2bx])/(4x) - (b\text{Cosh}[bx]*\text{SinhIntegral}[bx])/(2x) - (\text{Sinh}[bx]*\text{SinhIntegral}[bx])/(2x^2) + (b^2\text{SinhIntegral}[bx]^2)/4$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 29

Int[(x_)^(-1), x_Symbol] := Simp[Log[x], x]

Rule 3378

Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Dist[f/(d*(m + 1)), Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3393

Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))

Rule 3395

Int[((c_.) + (d_.)*(x_))^(m_)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(c + d*x)^(m + 1)*((b*Sine[e + f*x])^n/(d*(m + 1))), x] + (Dist[b^2*f^2*n*((n - 1)/(d^2*(m + 1)*(m + 2))), Int[(c + d*x)^(m + 2)*(b*Sine[e + f*x])^(n - 2), x], x] - Dist[f^2*(n^2/(d^2*(m + 1)*(m + 2))), Int[(c + d*x)^(m + 2)*(b*Sine[e + f*x])^n, x], x] - Simp[b*f*n*(c + d*x)^(m + 2)*Cos[e + f*x]*((b*Sine[e + f*x])^(n - 1)/(d^2*(m + 1)*(m + 2))), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && LtQ[m, -2]

Rule 5556

Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]

Rule 6679

Int[((e_.) + (f_.)*(x_))^(m_)*Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^(m + 1)*Sinh[a + b*x]*(SinhIntegr

al[c + d*x]/(f*(m + 1)), x] + (-Dist[b/(f*(m + 1)), Int[(e + f*x)^(m + 1)*Cosh[a + b*x]*SinhIntegral[c + d*x], x], x] - Dist[d/(f*(m + 1)), Int[(e + f*x)^(m + 1)*Sinh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[m, -1]

Rule 6685

Int[Cosh[(a_.) + (b_.)*(x_.)]*((e_.) + (f_.)*(x_.))^(m_.)*SinhIntegral[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[(e + f*x)^(m + 1)*Cosh[a + b*x]*(SinhIntegral[c + d*x]/(f*(m + 1))), x] + (-Dist[b/(f*(m + 1)), Int[(e + f*x)^(m + 1)*Sinh[a + b*x]*SinhIntegral[c + d*x], x], x] - Dist[d/(f*(m + 1)), Int[(e + f*x)^(m + 1)*Cosh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[m, -1]

Rule 6818

Int[(u_)*(y_)^(m_.), x_Symbol] := With[{q = DerivativeDivides[y, u, x]}, Simp[q*(y^(m + 1)/(m + 1)), x] /; !FalseQ[q]] /; FreeQ[m, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\sinh(bx)\text{Shi}(bx)}{2x^2} + \frac{1}{2}b \int \frac{\sinh^2(bx)}{bx^3} dx + \frac{1}{2}b \int \frac{\cosh(bx)\text{Shi}(bx)}{x^2} dx \\
 &= -\frac{b \cosh(bx)\text{Shi}(bx)}{2x} - \frac{\sinh(bx)\text{Shi}(bx)}{2x^2} + \frac{1}{2} \int \frac{\sinh^2(bx)}{x^3} dx \\
 &\quad + \frac{1}{2}b^2 \int \frac{\cosh(bx) \sinh(bx)}{bx^2} dx + \frac{1}{2}b^2 \int \frac{\sinh(bx)\text{Shi}(bx)}{x} dx \\
 &= -\frac{b \cosh(bx) \sinh(bx)}{2x} - \frac{\sinh^2(bx)}{4x^2} - \frac{b \cosh(bx)\text{Shi}(bx)}{2x} - \frac{\sinh(bx)\text{Shi}(bx)}{2x^2} \\
 &\quad + \frac{1}{4}b^2\text{Shi}(bx)^2 + \frac{1}{2}b \int \frac{\cosh(bx) \sinh(bx)}{x^2} dx + \frac{1}{2}b^2 \int \frac{1}{x} dx + b^2 \int \frac{\sinh^2(bx)}{x} dx \\
 &= \frac{1}{2}b^2 \log(x) - \frac{b \cosh(bx) \sinh(bx)}{2x} - \frac{\sinh^2(bx)}{4x^2} - \frac{b \cosh(bx)\text{Shi}(bx)}{2x} - \frac{\sinh(bx)\text{Shi}(bx)}{2x^2} \\
 &\quad + \frac{1}{4}b^2\text{Shi}(bx)^2 + \frac{1}{2}b \int \frac{\sinh(2bx)}{2x^2} dx - b^2 \int \left(\frac{1}{2x} - \frac{\cosh(2bx)}{2x} \right) dx \\
 &= -\frac{b \cosh(bx) \sinh(bx)}{2x} - \frac{\sinh^2(bx)}{4x^2} - \frac{b \cosh(bx)\text{Shi}(bx)}{2x} - \frac{\sinh(bx)\text{Shi}(bx)}{2x^2} \\
 &\quad + \frac{1}{4}b^2\text{Shi}(bx)^2 + \frac{1}{4}b \int \frac{\sinh(2bx)}{x^2} dx + \frac{1}{2}b^2 \int \frac{\cosh(2bx)}{x} dx \\
 &= \frac{1}{2}b^2\text{Chi}(2bx) - \frac{b \cosh(bx) \sinh(bx)}{2x} - \frac{\sinh^2(bx)}{4x^2} - \frac{b \sinh(2bx)}{4x} \\
 &\quad - \frac{b \cosh(bx)\text{Shi}(bx)}{2x} - \frac{\sinh(bx)\text{Shi}(bx)}{2x^2} + \frac{1}{4}b^2\text{Shi}(bx)^2 + \frac{1}{2}b^2 \int \frac{\cosh(2bx)}{x} dx
 \end{aligned}$$

$$= b^2 \text{Chi}(2bx) - \frac{b \cosh(bx) \sinh(bx)}{2x} - \frac{\sinh^2(bx)}{4x^2} - \frac{b \sinh(2bx)}{4x} \\ - \frac{b \cosh(bx) \text{Shi}(bx)}{2x} - \frac{\sinh(bx) \text{Shi}(bx)}{2x^2} + \frac{1}{4} b^2 \text{Shi}(bx)^2$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.00

$$\int \frac{\sinh(bx) \text{Shi}(bx)}{x^3} dx = b^2 \text{Chi}(2bx) - \frac{b \cosh(bx) \sinh(bx)}{2x} - \frac{\sinh^2(bx)}{4x^2} - \frac{b \sinh(2bx)}{4x} \\ - \frac{b \cosh(bx) \text{Shi}(bx)}{2x} - \frac{\sinh(bx) \text{Shi}(bx)}{2x^2} + \frac{1}{4} b^2 \text{Shi}(bx)^2$$

[In] Integrate[(Sinh[b*x]*SinhIntegral[b*x])/x^3,x]

[Out] b^2*CoshIntegral[2*b*x] - (b*Cosh[b*x]*Sinh[b*x])/(2*x) - Sinh[b*x]^2/(4*x^2) - (b*Sinh[2*b*x])/(4*x) - (b*Cosh[b*x]*SinhIntegral[b*x])/(2*x) - (Sinh[b*x]*SinhIntegral[b*x])/(2*x^2) + (b^2*SinhIntegral[b*x]^2)/4

Maple [F]

$$\int \frac{\text{Shi}(bx) \sinh(bx)}{x^3} dx$$

[In] int(Shi(b*x)*sinh(b*x)/x^3,x)

[Out] int(Shi(b*x)*sinh(b*x)/x^3,x)

Fricas [F]

$$\int \frac{\sinh(bx) \text{Shi}(bx)}{x^3} dx = \int \frac{\text{Shi}(bx) \sinh(bx)}{x^3} dx$$

[In] integrate(Shi(b*x)*sinh(b*x)/x^3,x, algorithm="fricas")

[Out] integral(sinh(b*x)*sinh_integral(b*x)/x^3, x)

Sympy [F]

$$\int \frac{\sinh(bx)\operatorname{Shi}(bx)}{x^3} dx = \int \frac{\sinh(bx)\operatorname{Shi}(bx)}{x^3} dx$$

```
[In] integrate(Shi(b*x)*sinh(b*x)/x**3,x)
```

```
[Out] Integral(sinh(b*x)*Shi(b*x)/x**3, x)
```

Maxima [F]

$$\int \frac{\sinh(bx)\operatorname{Shi}(bx)}{x^3} dx = \int \frac{\operatorname{Shi}(bx)\sinh(bx)}{x^3} dx$$

```
[In] integrate(Shi(b*x)*sinh(b*x)/x^3,x, algorithm="maxima")
```

```
[Out] integrate(Shi(b*x)*sinh(b*x)/x^3, x)
```

Giac [F]

$$\int \frac{\sinh(bx)\operatorname{Shi}(bx)}{x^3} dx = \int \frac{\operatorname{Shi}(bx)\sinh(bx)}{x^3} dx$$

```
[In] integrate(Shi(b*x)*sinh(b*x)/x^3,x, algorithm="giac")
```

```
[Out] integrate(Shi(b*x)*sinh(b*x)/x^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\sinh(bx)\operatorname{Shi}(bx)}{x^3} dx = \int \frac{\sinhint(bx)\sinh(bx)}{x^3} dx$$

```
[In] int((sinhint(b*x)*sinh(b*x))/x^3,x)
```

```
[Out] int((sinhint(b*x)*sinh(b*x))/x^3, x)
```

3.40 $\int \frac{\sinh(bx)\mathbf{Shi}(bx)}{x^2} dx$

Optimal result	229
Rubi [N/A]	229
Mathematica [N/A]	230
Maple [N/A] (verified)	230
Fricas [N/A]	231
Sympy [N/A]	231
Maxima [N/A]	231
Giac [N/A]	232
Mupad [N/A]	232

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\sinh(bx)\mathbf{Shi}(bx)}{x^2} dx = -\frac{\sinh^2(bx)}{x} - \frac{\sinh(bx)\mathbf{Shi}(bx)}{x} + b\mathbf{Shi}(2bx) + b\mathbf{Int}\left(\frac{\cosh(bx)\mathbf{Shi}(bx)}{x}, x\right)$$

[Out] b*CannotIntegrate(cosh(b*x)*Shi(b*x)/x,x)+b*Shi(2*b*x)-Shi(b*x)*sinh(b*x)/x-sinh(b*x)^2/x

Rubi [N/A]

Not integrable

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\sinh(bx)\mathbf{Shi}(bx)}{x^2} dx = \int \frac{\sinh(bx)\mathbf{Shi}(bx)}{x^2} dx$$

[In] Int[(Sinh[b*x]*SinhIntegral[b*x])/x^2,x]

[Out] -(Sinh[b*x]^2/x) - (Sinh[b*x]*SinhIntegral[b*x])/x + b*SinhIntegral[2*b*x] + b*Defer[Int] [(Cosh[b*x]*SinhIntegral[b*x])/x, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\sinh(bx)\text{Shi}(bx)}{x} + b \int \frac{\sinh^2(bx)}{bx^2} dx + b \int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx \\
 &= -\frac{\sinh(bx)\text{Shi}(bx)}{x} + b \int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx + \int \frac{\sinh^2(bx)}{x^2} dx \\
 &= -\frac{\sinh^2(bx)}{x} - \frac{\sinh(bx)\text{Shi}(bx)}{x} - (2ib) \int \frac{i \sinh(2bx)}{2x} dx + b \int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx \\
 &= -\frac{\sinh^2(bx)}{x} - \frac{\sinh(bx)\text{Shi}(bx)}{x} + b \int \frac{\sinh(2bx)}{x} dx + b \int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx \\
 &= -\frac{\sinh^2(bx)}{x} - \frac{\sinh(bx)\text{Shi}(bx)}{x} + b\text{Shi}(2bx) + b \int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx
 \end{aligned}$$

Mathematica [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\sinh(bx)\text{Shi}(bx)}{x^2} dx = \int \frac{\sinh(bx)\text{Shi}(bx)}{x^2} dx$$

[In] Integrate[(Sinh[b*x]*SinhIntegral[b*x])/x^2,x]

[Out] Integrate[(Sinh[b*x]*SinhIntegral[b*x])/x^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\text{Shi}(bx) \sinh(bx)}{x^2} dx$$

[In] int(Shi(b*x)*sinh(b*x)/x^2,x)

[Out] int(Shi(b*x)*sinh(b*x)/x^2,x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\sinh(bx)\text{Shi}(bx)}{x^2} dx = \int \frac{\text{Shi}(bx) \sinh(bx)}{x^2} dx$$

[In] integrate(Shi(b*x)*sinh(b*x)/x^2,x, algorithm="fricas")

[Out] integral(sinh(b*x)*sinh_integral(b*x)/x^2, x)

Sympy [N/A]

Not integrable

Time = 2.38 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\sinh(bx)\text{Shi}(bx)}{x^2} dx = \int \frac{\sinh(bx) \text{Shi}(bx)}{x^2} dx$$

[In] integrate(Shi(b*x)*sinh(b*x)/x**2,x)

[Out] Integral(sinh(b*x)*Shi(b*x)/x**2, x)

Maxima [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\sinh(bx)\text{Shi}(bx)}{x^2} dx = \int \frac{\text{Shi}(bx) \sinh(bx)}{x^2} dx$$

[In] integrate(Shi(b*x)*sinh(b*x)/x^2,x, algorithm="maxima")

[Out] integrate(Shi(b*x)*sinh(b*x)/x^2, x)

Giac [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\sinh(bx)\text{Shi}(bx)}{x^2} dx = \int \frac{\text{Shi}(bx) \sinh(bx)}{x^2} dx$$

[In] integrate(Shi(b*x)*sinh(b*x)/x^2,x, algorithm="giac")

[Out] integrate(Shi(b*x)*sinh(b*x)/x^2, x)

Mupad [N/A]

Not integrable

Time = 4.85 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\sinh(bx)\text{Shi}(bx)}{x^2} dx = \int \frac{\sinhint(bx) \sinh(bx)}{x^2} dx$$

[In] int((sinhint(b*x)*sinh(b*x))/x^2,x)

[Out] int((sinhint(b*x)*sinh(b*x))/x^2, x)

3.41 $\int \frac{\sinh(bx)\mathbf{Shi}(bx)}{x} dx$

Optimal result	233
Rubi [A] (verified)	233
Mathematica [A] (verified)	234
Maple [A] (verified)	234
Fricas [F]	234
Sympy [A] (verification not implemented)	234
Maxima [F]	235
Giac [F]	235
Mupad [F(-1)]	235

Optimal result

Integrand size = 12, antiderivative size = 10

$$\int \frac{\sinh(bx)\mathbf{Shi}(bx)}{x} dx = \frac{\mathbf{Shi}(bx)^2}{2}$$

[Out] 1/2*Shi(b*x)^2

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {6818}

$$\int \frac{\sinh(bx)\mathbf{Shi}(bx)}{x} dx = \frac{\mathbf{Shi}(bx)^2}{2}$$

[In] Int[(Sinh[b*x]*SinhIntegral[b*x])/x,x]

[Out] SinhIntegral[b*x]^2/2

Rule 6818

Int[(u)*(y_)^(m_.), x_Symbol] :> With[{q = DerivativeDivides[y, u, x]}, Si
mp[q*(y^(m + 1)/(m + 1)), x] /; !FalseQ[q] /; FreeQ[m, x] && NeQ[m, -1]

Rubi steps

$$\text{integral} = \frac{\mathbf{Shi}(bx)^2}{2}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\sinh(bx)\text{Shi}(bx)}{x} dx = \frac{\text{Shi}(bx)^2}{2}$$

[In] Integrate[(Sinh[b*x]*SinhIntegral[b*x])/x,x]

[Out] SinhIntegral[b*x]^2/2

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

method	result	size
derivativedivides	$\frac{\text{Shi}(bx)^2}{2}$	9
default	$\frac{\text{Shi}(bx)^2}{2}$	9

[In] int(Shi(b*x)*sinh(b*x)/x,x,method=_RETURNVERBOSE)

[Out] 1/2*Shi(b*x)^2

Fricas [F]

$$\int \frac{\sinh(bx)\text{Shi}(bx)}{x} dx = \int \frac{\text{Shi}(bx)\sinh(bx)}{x} dx$$

[In] integrate(Shi(b*x)*sinh(b*x)/x,x, algorithm="fricas")

[Out] integral(sinh(b*x)*sinh_integral(b*x)/x, x)

Sympy [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \frac{\sinh(bx)\text{Shi}(bx)}{x} dx = \frac{\text{Shi}^2(bx)}{2}$$

[In] integrate(Shi(b*x)*sinh(b*x)/x,x)

[Out] Shi(b*x)**2/2

Maxima [F]

$$\int \frac{\sinh(bx)\text{Shi}(bx)}{x} dx = \int \frac{\text{Shi}(bx) \sinh(bx)}{x} dx$$

[In] integrate(Shi(b*x)*sinh(b*x)/x,x, algorithm="maxima")

[Out] integrate(Shi(b*x)*sinh(b*x)/x, x)

Giac [F]

$$\int \frac{\sinh(bx)\text{Shi}(bx)}{x} dx = \int \frac{\text{Shi}(bx) \sinh(bx)}{x} dx$$

[In] integrate(Shi(b*x)*sinh(b*x)/x,x, algorithm="giac")

[Out] integrate(Shi(b*x)*sinh(b*x)/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sinh(bx)\text{Shi}(bx)}{x} dx = \frac{\sinhint(bx)^2}{2}$$

[In] int((sinhint(b*x)*sinh(b*x))/x,x)

[Out] sinhint(b*x)^2/2

3.42 $\int \sinh(bx) \mathbf{Shi}(bx) dx$

Optimal result	236
Rubi [A] (verified)	236
Mathematica [A] (verified)	237
Maple [A] (verified)	238
Fricas [F]	238
Sympy [F]	238
Maxima [F]	238
Giac [F]	239
Mupad [F(-1)]	239

Optimal result

Integrand size = 9, antiderivative size = 25

$$\int \sinh(bx) \mathbf{Shi}(bx) dx = \frac{\cosh(bx) \mathbf{Shi}(bx)}{b} - \frac{\mathbf{Shi}(2bx)}{2b}$$

[Out] $\cosh(b*x)*\mathbf{Shi}(b*x)/b - 1/2*\mathbf{Shi}(2*b*x)/b$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {6675, 12, 5556, 3379}

$$\int \sinh(bx) \mathbf{Shi}(bx) dx = \frac{\mathbf{Shi}(bx) \cosh(bx)}{b} - \frac{\mathbf{Shi}(2bx)}{2b}$$

[In] $\text{Int}[\text{Sinh}[b*x]*\text{SinhIntegral}[b*x], x]$

[Out] $(\text{Cosh}[b*x]*\text{SinhIntegral}[b*x])/b - \text{SinhIntegral}[2*b*x]/(2*b)$

Rule 12

$\text{Int}[(a_*)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[u, (b_*)*(v_)] /; \text{FreeQ}[b, x]$

Rule 3379

$\text{Int}[\sin[(e_*) + (\text{Complex}[0, fz_*])*(f_*)*(x_*)]/((c_*) + (d_*)*(x_*)), x_Symbol] \rightarrow \text{Simp}[I*(\text{SinhIntegral}[c*f*(fz/d) + f*fz*x]/d), x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \ \&\& \ \text{EqQ}[d*e - c*f*fz*I, 0]$

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

Rule 6675

```
Int[Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
Simp[Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\cosh(bx)\text{Shi}(bx)}{b} - \int \frac{\cosh(bx)\sinh(bx)}{bx} dx \\
&= \frac{\cosh(bx)\text{Shi}(bx)}{b} - \int \frac{\cosh(bx)\sinh(bx)}{x} dx \\
&= \frac{\cosh(bx)\text{Shi}(bx)}{b} - \int \frac{\sinh(2bx)}{2x} dx \\
&= \frac{\cosh(bx)\text{Shi}(bx)}{b} - \frac{\int \frac{\sinh(2bx)}{x} dx}{2b} \\
&= \frac{\cosh(bx)\text{Shi}(bx)}{b} - \frac{\text{Shi}(2bx)}{2b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \sinh(bx)\text{Shi}(bx) dx = \frac{\cosh(bx)\text{Shi}(bx)}{b} - \frac{\text{Shi}(2bx)}{2b}$$

```
[In] Integrate[Sinh[b*x]*SinhIntegral[b*x], x]
```

```
[Out] (Cosh[b*x]*SinhIntegral[b*x])/b - SinhIntegral[2*b*x]/(2*b)
```

Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{\cosh(bx) \operatorname{Shi}(bx) - \frac{\operatorname{Shi}(2bx)}{2}}{b}$	22
default	$\frac{\cosh(bx) \operatorname{Shi}(bx) - \frac{\operatorname{Shi}(2bx)}{2}}{b}$	22

[In] `int(Shi(b*x)*sinh(b*x),x,method=_RETURNVERBOSE)`

[Out] `1/b*(cosh(b*x)*Shi(b*x)-1/2*Shi(2*b*x))`

Fricas [F]

$$\int \sinh(bx) \operatorname{Shi}(bx) dx = \int \operatorname{Shi}(bx) \sinh(bx) dx$$

[In] `integrate(Shi(b*x)*sinh(b*x),x, algorithm="fricas")`

[Out] `integral(sinh(b*x)*sinh_integral(b*x), x)`

Sympy [F]

$$\int \sinh(bx) \operatorname{Shi}(bx) dx = \int \sinh(bx) \operatorname{Shi}(bx) dx$$

[In] `integrate(Shi(b*x)*sinh(b*x),x)`

[Out] `Integral(sinh(b*x)*Shi(b*x), x)`

Maxima [F]

$$\int \sinh(bx) \operatorname{Shi}(bx) dx = \int \operatorname{Shi}(bx) \sinh(bx) dx$$

[In] `integrate(Shi(b*x)*sinh(b*x),x, algorithm="maxima")`

[Out] `integrate(Shi(b*x)*sinh(b*x), x)`

Giac [**F**]

$$\int \sinh(bx) \operatorname{Shi}(bx) dx = \int \operatorname{Shi}(bx) \sinh(bx) dx$$

[In] integrate(Shi(b*x)*sinh(b*x),x, algorithm="giac")

[Out] integrate(Shi(b*x)*sinh(b*x), x)

Mupad [**F(-1)**]

Timed out.

$$\int \sinh(bx) \operatorname{Shi}(bx) dx = \int \operatorname{sinhint}(bx) \sinh(bx) dx$$

[In] int(sinhint(b*x)*sinh(b*x),x)

[Out] int(sinhint(b*x)*sinh(b*x), x)

3.43 $\int x \sinh(bx) \mathbf{Shi}(bx) dx$

Optimal result	240
Rubi [A] (verified)	240
Mathematica [A] (verified)	242
Maple [A] (verified)	242
Fricas [F]	243
Sympy [F]	243
Maxima [F]	243
Giac [F]	243
Mupad [F(-1)]	244

Optimal result

Integrand size = 10, antiderivative size = 61

$$\int x \sinh(bx) \mathbf{Shi}(bx) dx = \frac{\mathbf{Chi}(2bx)}{2b^2} - \frac{\log(x)}{2b^2} - \frac{\sinh^2(bx)}{2b^2} + \frac{x \cosh(bx) \mathbf{Shi}(bx)}{b} - \frac{\sinh(bx) \mathbf{Shi}(bx)}{b^2}$$

[Out] $1/2*\mathbf{Chi}(2*b*x)/b^2-1/2*\ln(x)/b^2+x*\cosh(b*x)*\mathbf{Shi}(b*x)/b-\mathbf{Shi}(b*x)*\sinh(b*x)/b^2-1/2*\sinh(b*x)^2/b^2$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {6677, 12, 2644, 30, 6681, 3393, 3382}

$$\int x \sinh(bx) \mathbf{Shi}(bx) dx = \frac{\mathbf{Chi}(2bx)}{2b^2} - \frac{\mathbf{Shi}(bx) \sinh(bx)}{b^2} - \frac{\log(x)}{2b^2} - \frac{\sinh^2(bx)}{2b^2} + \frac{x \mathbf{Shi}(bx) \cosh(bx)}{b}$$

[In] `Int[x*Sinh[b*x]*SinhIntegral[b*x],x]`

[Out] `CoshIntegral[2*b*x]/(2*b^2) - Log[x]/(2*b^2) - Sinh[b*x]^2/(2*b^2) + (x*Cosh[b*x]*SinhIntegral[b*x])/b - (Sinh[b*x]*SinhIntegral[b*x])/b^2`

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2644

Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)]^(m_), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rule 3382

Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3393

Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))

Rule 6677

Int[((e_) + (f_)*(x_))^(m_)*Sinh[(a_) + (b_)*(x_)]*SinhIntegral[(c_) + (d_)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*SinhIntegral[c + d*x], x], x) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rule 6681

Int[Cosh[(a_) + (b_)*(x_)]*SinhIntegral[(c_) + (d_)*(x_)], x_Symbol] := Simp[Sinh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x \cosh(bx) \text{Shi}(bx)}{b} - \frac{\int \cosh(bx) \text{Shi}(bx) dx}{b} - \int \frac{\cosh(bx) \sinh(bx)}{b} dx \\
 &= \frac{x \cosh(bx) \text{Shi}(bx)}{b} - \frac{\sinh(bx) \text{Shi}(bx)}{b^2} - \frac{\int \cosh(bx) \sinh(bx) dx}{b} + \frac{\int \frac{\sinh^2(bx)}{bx} dx}{b} \\
 &= \frac{x \cosh(bx) \text{Shi}(bx)}{b} - \frac{\sinh(bx) \text{Shi}(bx)}{b^2} + \frac{\int \frac{\sinh^2(bx)}{x} dx}{b^2} + \frac{\text{Subst}(\int x dx, x, i \sinh(bx))}{b^2}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{\sinh^2(bx)}{2b^2} + \frac{x \cosh(bx)\text{Shi}(bx)}{b} - \frac{\sinh(bx)\text{Shi}(bx)}{b^2} - \frac{\int \left(\frac{1}{2x} - \frac{\cosh(2bx)}{2x} \right) dx}{b^2} \\
&= -\frac{\log(x)}{2b^2} - \frac{\sinh^2(bx)}{2b^2} + \frac{x \cosh(bx)\text{Shi}(bx)}{b} - \frac{\sinh(bx)\text{Shi}(bx)}{b^2} + \frac{\int \frac{\cosh(2bx)}{x} dx}{2b^2} \\
&= \frac{\text{Chi}(2bx)}{2b^2} - \frac{\log(x)}{2b^2} - \frac{\sinh^2(bx)}{2b^2} + \frac{x \cosh(bx)\text{Shi}(bx)}{b} - \frac{\sinh(bx)\text{Shi}(bx)}{b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.72

$$\begin{aligned}
&\int x \sinh(bx)\text{Shi}(bx) dx \\
&= -\frac{\cosh(2bx) - 2\text{Chi}(2bx) + 2\log(x) + (-4bx \cosh(bx) + 4 \sinh(bx))\text{Shi}(bx)}{4b^2}
\end{aligned}$$

[In] Integrate[x*Sinh[b*x]*SinhIntegral[b*x],x]

[Out] -1/4*(Cosh[2*b*x] - 2*CoshIntegral[2*b*x] + 2*Log[x] + (-4*b*x*Cosh[b*x] + 4*Sinh[b*x])*SinhIntegral[b*x])/b^2

Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.75

method	result	size
derivativedivides	$\frac{\text{Shi}(bx)(bx \cosh(bx) - \sinh(bx)) - \frac{\cosh(bx)^2}{2} - \frac{\ln(bx)}{2} + \frac{\text{Chi}(2bx)}{2}}{b^2}$	46
default	$\frac{\text{Shi}(bx)(bx \cosh(bx) - \sinh(bx)) - \frac{\cosh(bx)^2}{2} - \frac{\ln(bx)}{2} + \frac{\text{Chi}(2bx)}{2}}{b^2}$	46

[In] int(x*Shi(b*x)*sinh(b*x),x,method=_RETURNVERBOSE)

[Out] 1/b^2*(Shi(b*x)*(b*x*cosh(b*x)-sinh(b*x))-1/2*cosh(b*x)^2-1/2*ln(b*x)+1/2*Chi(2*b*x))

Fricas [F]

$$\int x \sinh(bx) \operatorname{Shi}(bx) dx = \int x \operatorname{Shi}(bx) \sinh(bx) dx$$

[In] `integrate(x*Shi(b*x)*sinh(b*x),x, algorithm="fricas")`

[Out] `integral(x*sinh(b*x)*sinh_integral(b*x), x)`

Sympy [F]

$$\int x \sinh(bx) \operatorname{Shi}(bx) dx = \int x \sinh(bx) \operatorname{Shi}(bx) dx$$

[In] `integrate(x*Shi(b*x)*sinh(b*x),x)`

[Out] `Integral(x*sinh(b*x)*Shi(b*x), x)`

Maxima [F]

$$\int x \sinh(bx) \operatorname{Shi}(bx) dx = \int x \operatorname{Shi}(bx) \sinh(bx) dx$$

[In] `integrate(x*Shi(b*x)*sinh(b*x),x, algorithm="maxima")`

[Out] `integrate(x*Shi(b*x)*sinh(b*x), x)`

Giac [F]

$$\int x \sinh(bx) \operatorname{Shi}(bx) dx = \int x \operatorname{Shi}(bx) \sinh(bx) dx$$

[In] `integrate(x*Shi(b*x)*sinh(b*x),x, algorithm="giac")`

[Out] `integrate(x*Shi(b*x)*sinh(b*x), x)`

Mupad [F(-1)]

Timed out.

$$\int x \sinh(bx) \operatorname{Shi}(bx) dx = \int x \operatorname{sinhint}(bx) \sinh(bx) dx$$

```
[In] int(x*sinhint(b*x)*sinh(b*x),x)
```

```
[Out] int(x*sinhint(b*x)*sinh(b*x), x)
```

3.44 $\int x^2 \sinh(bx) \mathbf{Shi}(bx) dx$

Optimal result	245
Rubi [A] (verified)	245
Mathematica [A] (verified)	248
Maple [A] (verified)	248
Fricas [F]	248
Sympy [F]	249
Maxima [F]	249
Giac [F]	249
Mupad [F(-1)]	249

Optimal result

Integrand size = 12, antiderivative size = 90

$$\int x^2 \sinh(bx) \mathbf{Shi}(bx) dx = -\frac{5x}{4b^2} + \frac{5 \cosh(bx) \sinh(bx)}{4b^3} - \frac{x \sinh^2(bx)}{2b^2} + \frac{2 \cosh(bx) \mathbf{Shi}(bx)}{b^3} \\ + \frac{x^2 \cosh(bx) \mathbf{Shi}(bx)}{b} - \frac{2x \sinh(bx) \mathbf{Shi}(bx)}{b^2} - \frac{\mathbf{Shi}(2bx)}{b^3}$$

[Out] $-5/4*x/b^2+2*\cosh(b*x)*\mathbf{Shi}(b*x)/b^3+x^2*\cosh(b*x)*\mathbf{Shi}(b*x)/b-\mathbf{Shi}(2*b*x)/b^3$
 $+5/4*\cosh(b*x)*\sinh(b*x)/b^3-2*x*\mathbf{Shi}(b*x)*\sinh(b*x)/b^2-1/2*x*\sinh(b*x)^2/b$
 2

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6677, 12, 5480, 2715, 8, 6683, 6675, 5556, 3379}

$$\int x^2 \sinh(bx) \mathbf{Shi}(bx) dx = -\frac{\mathbf{Shi}(2bx)}{b^3} + \frac{2\mathbf{Shi}(bx) \cosh(bx)}{b^3} + \frac{5 \sinh(bx) \cosh(bx)}{4b^3} \\ - \frac{2x\mathbf{Shi}(bx) \sinh(bx)}{b^2} - \frac{5x}{4b^2} - \frac{x \sinh^2(bx)}{2b^2} + \frac{x^2\mathbf{Shi}(bx) \cosh(bx)}{b}$$

[In] $\text{Int}[x^2*\text{Sinh}[b*x]*\text{SinhIntegral}[b*x], x]$

[Out] $(-5*x)/(4*b^2) + (5*\text{Cosh}[b*x]*\text{Sinh}[b*x])/(4*b^3) - (x*\text{Sinh}[b*x]^2)/(2*b^2)$
 $+ (2*\text{Cosh}[b*x]*\text{SinhIntegral}[b*x])/b^3 + (x^2*\text{Cosh}[b*x]*\text{SinhIntegral}[b*x])/b$
 $- (2*x*\text{Sinh}[b*x]*\text{SinhIntegral}[b*x])/b^2 - \text{SinhIntegral}[2*b*x]/b^3$

Rule 8

```
Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 5480

```
Int[Cosh[(a_.) + (b_.)*(x_)^(n_.)]*(x_)^(m_.)*Sinh[(a_.) + (b_.)*(x_)^(n_.)]^(p_.), x_Symbol] := Simp[x^(m - n + 1)*(Sinh[a + b*x^n]^(p + 1)/(b*n*(p + 1))), x] - Dist[(m - n + 1)/(b*n*(p + 1)), Int[x^(m - n)*Sinh[a + b*x^n]^(p + 1), x], x] /; FreeQ[{a, b, p}, x] && LtQ[0, n, m + 1] && NeQ[p, -1]
```

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_)^(p_.)]*((c_.) + (d_.)*(x_)^(m_.))*Sinh[(a_.) + (b_.)*(x_)^(n_.)], x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]
```

Rule 6675

```
Int[Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6677

```
Int[((e_.) + (f_.)*(x_)^(m_.))*Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*SinhIn
```

tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rule 6683

Int[Cosh[(a_.) + (b_.)*(x_.)]*((e_.) + (f_.)*(x_.))^(m_.)*SinhIntegral[(c_.) + (d_.)*(x_.)], x_Symbol] :> Simp[(e + f*x)^m*Sinh[a + b*x]*(SinhIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*SinhIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x^2 \cosh(bx) \text{Shi}(bx)}{b} - \frac{2 \int x \cosh(bx) \text{Shi}(bx) dx}{b} - \int \frac{x \cosh(bx) \sinh(bx)}{b} dx \\
 &= \frac{x^2 \cosh(bx) \text{Shi}(bx)}{b} - \frac{2x \sinh(bx) \text{Shi}(bx)}{b^2} + \frac{2 \int \sinh(bx) \text{Shi}(bx) dx}{b^2} \\
 &\quad - \frac{\int x \cosh(bx) \sinh(bx) dx}{b} + \frac{2 \int \frac{\sinh^2(bx)}{b} dx}{b} \\
 &= -\frac{x \sinh^2(bx)}{2b^2} + \frac{2 \cosh(bx) \text{Shi}(bx)}{b^3} + \frac{x^2 \cosh(bx) \text{Shi}(bx)}{b} - \frac{2x \sinh(bx) \text{Shi}(bx)}{b^2} \\
 &\quad + \frac{\int \sinh^2(bx) dx}{2b^2} - \frac{2 \int \frac{\cosh(bx) \sinh(bx)}{bx} dx}{b^2} + \frac{2 \int \sinh^2(bx) dx}{b^2} \\
 &= \frac{5 \cosh(bx) \sinh(bx)}{4b^3} - \frac{x \sinh^2(bx)}{2b^2} + \frac{2 \cosh(bx) \text{Shi}(bx)}{b^3} + \frac{x^2 \cosh(bx) \text{Shi}(bx)}{b} \\
 &\quad - \frac{2x \sinh(bx) \text{Shi}(bx)}{b^2} - \frac{2 \int \frac{\cosh(bx) \sinh(bx)}{x} dx}{b^3} - \frac{\int 1 dx}{4b^2} - \frac{\int 1 dx}{b^2} \\
 &= -\frac{5x}{4b^2} + \frac{5 \cosh(bx) \sinh(bx)}{4b^3} - \frac{x \sinh^2(bx)}{2b^2} + \frac{2 \cosh(bx) \text{Shi}(bx)}{b^3} \\
 &\quad + \frac{x^2 \cosh(bx) \text{Shi}(bx)}{b} - \frac{2x \sinh(bx) \text{Shi}(bx)}{b^2} - \frac{2 \int \frac{\sinh(2bx)}{2x} dx}{b^3} \\
 &= -\frac{5x}{4b^2} + \frac{5 \cosh(bx) \sinh(bx)}{4b^3} - \frac{x \sinh^2(bx)}{2b^2} + \frac{2 \cosh(bx) \text{Shi}(bx)}{b^3} \\
 &\quad + \frac{x^2 \cosh(bx) \text{Shi}(bx)}{b} - \frac{2x \sinh(bx) \text{Shi}(bx)}{b^2} - \frac{\int \frac{\sinh(2bx)}{x} dx}{b^3} \\
 &= -\frac{5x}{4b^2} + \frac{5 \cosh(bx) \sinh(bx)}{4b^3} - \frac{x \sinh^2(bx)}{2b^2} + \frac{2 \cosh(bx) \text{Shi}(bx)}{b^3} \\
 &\quad + \frac{x^2 \cosh(bx) \text{Shi}(bx)}{b} - \frac{2x \sinh(bx) \text{Shi}(bx)}{b^2} - \frac{\text{Shi}(2bx)}{b^3}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.71

$$\int x^2 \sinh(bx) \operatorname{Shi}(bx) dx$$

$$= \frac{-8bx - 2bx \cosh(2bx) + 5 \sinh(2bx) + 8((2 + b^2x^2) \cosh(bx) - 2bx \sinh(bx)) \operatorname{Shi}(bx) - 8 \operatorname{Shi}(2bx)}{8b^3}$$

[In] Integrate[x^2*Sinh[b*x]*SinhIntegral[b*x],x]

[Out] (-8*b*x - 2*b*x*Cosh[2*b*x] + 5*Sinh[2*b*x] + 8*((2 + b^2*x^2)*Cosh[b*x] - 2*b*x*Sinh[b*x])*SinhIntegral[b*x] - 8*SinhIntegral[2*b*x])/(8*b^3)

Maple [A] (verified)

Time = 0.77 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.76

method	result	size
derivativedivides	$\frac{\operatorname{Shi}(bx)(b^2x^2 \cosh(bx) - 2bx \sinh(bx) + 2 \cosh(bx)) - \frac{bx \cosh(bx)^2}{2} + \frac{5 \cosh(bx) \sinh(bx)}{4} - \frac{3bx}{4} - \operatorname{Shi}(2bx)}{b^3}$	68
default	$\frac{\operatorname{Shi}(bx)(b^2x^2 \cosh(bx) - 2bx \sinh(bx) + 2 \cosh(bx)) - \frac{bx \cosh(bx)^2}{2} + \frac{5 \cosh(bx) \sinh(bx)}{4} - \frac{3bx}{4} - \operatorname{Shi}(2bx)}{b^3}$	68

[In] int(x^2*Shi(b*x)*sinh(b*x),x,method=_RETURNVERBOSE)

[Out] 1/b^3*(Shi(b*x)*(b^2*x^2*cosh(b*x)-2*b*x*sinh(b*x)+2*cosh(b*x))-1/2*b*x*cosh(b*x)^2+5/4*cosh(b*x)*sinh(b*x)-3/4*b*x-Shi(2*b*x))

Fricas [F]

$$\int x^2 \sinh(bx) \operatorname{Shi}(bx) dx = \int x^2 \operatorname{Shi}(bx) \sinh(bx) dx$$

[In] integrate(x^2*Shi(b*x)*sinh(b*x),x, algorithm="fricas")

[Out] integral(x^2*sinh(b*x)*sinh_integral(b*x), x)

Sympy [F]

$$\int x^2 \sinh(bx) \operatorname{Shi}(bx) dx = \int x^2 \sinh(bx) \operatorname{Shi}(bx) dx$$

```
[In] integrate(x**2*Shi(b*x)*sinh(b*x),x)
```

```
[Out] Integral(x**2*sinh(b*x)*Shi(b*x), x)
```

Maxima [F]

$$\int x^2 \sinh(bx) \operatorname{Shi}(bx) dx = \int x^2 \operatorname{Shi}(bx) \sinh(bx) dx$$

```
[In] integrate(x^2*Shi(b*x)*sinh(b*x),x, algorithm="maxima")
```

```
[Out] integrate(x^2*Shi(b*x)*sinh(b*x), x)
```

Giac [F]

$$\int x^2 \sinh(bx) \operatorname{Shi}(bx) dx = \int x^2 \operatorname{Shi}(bx) \sinh(bx) dx$$

```
[In] integrate(x^2*Shi(b*x)*sinh(b*x),x, algorithm="giac")
```

```
[Out] integrate(x^2*Shi(b*x)*sinh(b*x), x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \sinh(bx) \operatorname{Shi}(bx) dx = \int x^2 \operatorname{sinhint}(bx) \sinh(bx) dx$$

```
[In] int(x^2*sinhint(b*x)*sinh(b*x),x)
```

```
[Out] int(x^2*sinhint(b*x)*sinh(b*x), x)
```

3.45 $\int x^3 \sinh(bx) \text{Shi}(bx) dx$

Optimal result	250
Rubi [A] (verified)	250
Mathematica [A] (verified)	253
Maple [A] (verified)	254
Fricas [F]	254
Sympy [F]	254
Maxima [F]	254
Giac [F]	255
Mupad [F(-1)]	255

Optimal result

Integrand size = 12, antiderivative size = 125

$$\int x^3 \sinh(bx) \text{Shi}(bx) dx = -\frac{x^2}{b^2} + \frac{3\text{Chi}(2bx)}{b^4} - \frac{3 \log(x)}{b^4} + \frac{2x \cosh(bx) \sinh(bx)}{b^3} - \frac{4 \sinh^2(bx)}{b^4} - \frac{x^2 \sinh^2(bx)}{2b^2} + \frac{6x \cosh(bx) \text{Shi}(bx)}{b^3} + \frac{x^3 \cosh(bx) \text{Shi}(bx)}{b} - \frac{6 \sinh(bx) \text{Shi}(bx)}{b^4} - \frac{3x^2 \sinh(bx) \text{Shi}(bx)}{b^2}$$

[Out] $-x^2/b^2+3*\text{Chi}(2*b*x)/b^4-3*\ln(x)/b^4+6*x*\cosh(b*x)*\text{Shi}(b*x)/b^3+x^3*\cosh(b*x)*\text{Shi}(b*x)/b+2*x*\cosh(b*x)*\sinh(b*x)/b^3-6*\text{Shi}(b*x)*\sinh(b*x)/b^4-3*x^2*\text{Shi}(b*x)*\sinh(b*x)/b^2-4*\sinh(b*x)^2/b^4-1/2*x^2*\sinh(b*x)^2/b^2$

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.00, number of steps used = 18, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {6677, 12, 5480, 3391, 30, 6683, 2644, 6681, 3393, 3382}

$$\int x^3 \sinh(bx) \text{Shi}(bx) dx = \frac{3\text{Chi}(2bx)}{b^4} - \frac{6\text{Shi}(bx) \sinh(bx)}{b^4} - \frac{3 \log(x)}{b^4} - \frac{4 \sinh^2(bx)}{b^4} + \frac{6x \text{Shi}(bx) \cosh(bx)}{b^3} + \frac{2x \sinh(bx) \cosh(bx)}{b^3} - \frac{3x^2 \text{Shi}(bx) \sinh(bx)}{b^2} - \frac{x^2}{b^2} - \frac{x^2 \sinh^2(bx)}{2b^2} + \frac{x^3 \text{Shi}(bx) \cosh(bx)}{b}$$

[In] $\text{Int}[x^3*\text{Sinh}[b*x]*\text{SinhIntegral}[b*x],x]$

[Out] $-(x^2/b^2) + (3*\text{CoshIntegral}[2*b*x])/b^4 - (3*\text{Log}[x])/b^4 + (2*x*\text{Cosh}[b*x]*\text{Sinh}[b*x])/b^3 - (4*\text{Sinh}[b*x]^2)/b^4 - (x^2*\text{Sinh}[b*x]^2)/(2*b^2) + (6*x*\text{Cos}$

$$\frac{h[b*x]*\text{SinhIntegral}[b*x]}{b^3} + \frac{(x^3*\text{Cosh}[b*x]*\text{SinhIntegral}[b*x])}{b} - \frac{(6*\text{Sinh}[b*x]*\text{SinhIntegral}[b*x])}{b^4} - \frac{(3*x^2*\text{Sinh}[b*x]*\text{SinhIntegral}[b*x])}{b^2}$$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 30

`Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]`

Rule 2644

`Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])`

Rule 3382

`Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`

Rule 3391

`Int[((c_) + (d_)*(x_))*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[d*((b*Sine[e + f*x])^n/(f^2*n^2)), x] + (Dist[b^2*((n - 1)/n), Int[(c + d*x)*(b*Sine[e + f*x])^(n - 2), x], x] - Simp[b*(c + d*x)*Cos[e + f*x]*((b*Sine[e + f*x])^(n - 1)/(f*n)), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1]`

Rule 3393

`Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

Rule 5480

`Int[Cosh[(a_) + (b_)*(x_)]^(n_)*Sinh[(a_) + (b_)*(x_)]^(p_), x_Symbol] := Simp[x^(m - n + 1)*(Sinh[a + b*x]^n)^(p + 1)/(b*n*(p + 1)), x] - Dist[(m - n + 1)/(b*n*(p + 1)), Int[x^(m - n)*Sinh[a + b*x]^n)^(p + 1), x], x] /; FreeQ[{a, b, p}, x] && LtQ[0, n, m + 1] && NeQ[p, -1]`

Rule 6677

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(SinhIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Sinh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*SinhIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6681

```
Int[Cosh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
Simp[Sinh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6683

```
Int[Cosh[(a_.) + (b_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.)*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Sinh[a + b*x]*(SinhIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Sinh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*SinhIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x^3 \cosh(bx) \text{Shi}(bx)}{b} - \frac{3 \int x^2 \cosh(bx) \text{Shi}(bx) dx}{b} - \int \frac{x^2 \cosh(bx) \sinh(bx)}{b} dx \\
&= \frac{x^3 \cosh(bx) \text{Shi}(bx)}{b} - \frac{3x^2 \sinh(bx) \text{Shi}(bx)}{b^2} + \frac{6 \int x \sinh(bx) \text{Shi}(bx) dx}{b^2} \\
&\quad - \frac{\int x^2 \cosh(bx) \sinh(bx) dx}{b} + \frac{3 \int \frac{x \sinh^2(bx)}{b} dx}{b} \\
&= -\frac{x^2 \sinh^2(bx)}{2b^2} + \frac{6x \cosh(bx) \text{Shi}(bx)}{b^3} + \frac{x^3 \cosh(bx) \text{Shi}(bx)}{b} - \frac{3x^2 \sinh(bx) \text{Shi}(bx)}{b^2} \\
&\quad - \frac{6 \int \cosh(bx) \text{Shi}(bx) dx}{b^3} + \frac{\int x \sinh^2(bx) dx}{b^2} + \frac{3 \int x \sinh^2(bx) dx}{b^2} - \frac{6 \int \frac{\cosh(bx) \sinh(bx)}{b} dx}{b^2} \\
&= \frac{2x \cosh(bx) \sinh(bx)}{b^3} - \frac{\sinh^2(bx)}{b^4} - \frac{x^2 \sinh^2(bx)}{2b^2} + \frac{6x \cosh(bx) \text{Shi}(bx)}{b^3} \\
&\quad + \frac{x^3 \cosh(bx) \text{Shi}(bx)}{b} - \frac{6 \sinh(bx) \text{Shi}(bx)}{b^4} - \frac{3x^2 \sinh(bx) \text{Shi}(bx)}{b^2} \\
&\quad - \frac{6 \int \cosh(bx) \sinh(bx) dx}{b^3} + \frac{6 \int \frac{\sinh^2(bx)}{bx} dx}{b^3} - \frac{\int x dx}{2b^2} - \frac{3 \int x dx}{2b^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{x^2}{b^2} + \frac{2x \cosh(bx) \sinh(bx)}{b^3} - \frac{\sinh^2(bx)}{b^4} - \frac{x^2 \sinh^2(bx)}{2b^2} \\
&\quad + \frac{6x \cosh(bx) \text{Shi}(bx)}{b^3} + \frac{x^3 \cosh(bx) \text{Shi}(bx)}{b} - \frac{6 \sinh(bx) \text{Shi}(bx)}{b^4} \\
&\quad - \frac{3x^2 \sinh(bx) \text{Shi}(bx)}{b^2} + \frac{6 \int \frac{\sinh^2(bx)}{x} dx}{b^4} + \frac{6 \text{Subst}(\int x dx, x, i \sinh(bx))}{b^4} \\
&= -\frac{x^2}{b^2} + \frac{2x \cosh(bx) \sinh(bx)}{b^3} - \frac{4 \sinh^2(bx)}{b^4} - \frac{x^2 \sinh^2(bx)}{2b^2} + \frac{6x \cosh(bx) \text{Shi}(bx)}{b^3} \\
&\quad + \frac{x^3 \cosh(bx) \text{Shi}(bx)}{b} - \frac{6 \sinh(bx) \text{Shi}(bx)}{b^4} - \frac{3x^2 \sinh(bx) \text{Shi}(bx)}{b^2} - \frac{6 \int \left(\frac{1}{2x} - \frac{\cosh(2bx)}{2x} \right) dx}{b^4} \\
&= -\frac{x^2}{b^2} - \frac{3 \log(x)}{b^4} + \frac{2x \cosh(bx) \sinh(bx)}{b^3} - \frac{4 \sinh^2(bx)}{b^4} - \frac{x^2 \sinh^2(bx)}{2b^2} + \frac{6x \cosh(bx) \text{Shi}(bx)}{b^3} \\
&\quad + \frac{x^3 \cosh(bx) \text{Shi}(bx)}{b} - \frac{6 \sinh(bx) \text{Shi}(bx)}{b^4} - \frac{3x^2 \sinh(bx) \text{Shi}(bx)}{b^2} + \frac{3 \int \frac{\cosh(2bx)}{x} dx}{b^4} \\
&= -\frac{x^2}{b^2} + \frac{3 \text{Chi}(2bx)}{b^4} - \frac{3 \log(x)}{b^4} + \frac{2x \cosh(bx) \sinh(bx)}{b^3} - \frac{4 \sinh^2(bx)}{b^4} - \frac{x^2 \sinh^2(bx)}{2b^2} \\
&\quad + \frac{6x \cosh(bx) \text{Shi}(bx)}{b^3} + \frac{x^3 \cosh(bx) \text{Shi}(bx)}{b} - \frac{6 \sinh(bx) \text{Shi}(bx)}{b^4} - \frac{3x^2 \sinh(bx) \text{Shi}(bx)}{b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.74

$$\int x^3 \sinh(bx) \text{Shi}(bx) dx = \frac{3b^2 x^2 + 8 \cosh(2bx) + b^2 x^2 \cosh(2bx) - 12 \text{Chi}(2bx) + 12 \log(x) - 4bx \sinh(2bx) - 4(bx(6 + b^2 x^2) \cosh(2bx))}{4b^4}$$

[In] Integrate[x^3*Sinh[b*x]*SinhIntegral[b*x],x]

[Out] -1/4*(3*b^2*x^2 + 8*Cosh[2*b*x] + b^2*x^2*Cosh[2*b*x] - 12*CoshIntegral[2*b*x] + 12*Log[x] - 4*b*x*Sinh[2*b*x] - 4*(b*x*(6 + b^2*x^2)*Cosh[b*x] - 3*(2 + b^2*x^2)*Sinh[b*x])*SinhIntegral[b*x])/b^4

Maple [A] (verified)

Time = 0.96 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.83

method	result
derivativedivides	$\frac{\text{Shi}(bx)(b^3x^3 \cosh(bx) - 3b^2x^2 \sinh(bx) + 6bx \cosh(bx) - 6 \sinh(bx)) - \frac{b^2x^2 \cosh(bx)^2}{b^4} + 2bx \cosh(bx) \sinh(bx) - \frac{b^2x^2}{2} - 4 \cosh(bx)}{b^4}$
default	$\frac{\text{Shi}(bx)(b^3x^3 \cosh(bx) - 3b^2x^2 \sinh(bx) + 6bx \cosh(bx) - 6 \sinh(bx)) - \frac{b^2x^2 \cosh(bx)^2}{b^4} + 2bx \cosh(bx) \sinh(bx) - \frac{b^2x^2}{2} - 4 \cosh(bx)}{b^4}$

```
[In] int(x^3*Shi(b*x)*sinh(b*x),x,method=_RETURNVERBOSE)
```

```
[Out] 1/b^4*(Shi(b*x)*(b^3*x^3*cosh(b*x)-3*b^2*x^2*sinh(b*x)+6*b*x*cosh(b*x)-6*sinh(b*x))-1/2*b^2*x^2*cosh(b*x)^2+2*b*x*cosh(b*x)*sinh(b*x)-1/2*b^2*x^2-4*cosh(b*x)^2-3*ln(b*x)+3*Chi(2*b*x))
```

Fricas [F]

$$\int x^3 \sinh(bx) \text{Shi}(bx) dx = \int x^3 \text{Shi}(bx) \sinh(bx) dx$$

```
[In] integrate(x^3*Shi(b*x)*sinh(b*x),x, algorithm="fricas")
```

```
[Out] integral(x^3*sinh(b*x)*sinh_integral(b*x), x)
```

Sympy [F]

$$\int x^3 \sinh(bx) \text{Shi}(bx) dx = \int x^3 \sinh(bx) \text{Shi}(bx) dx$$

```
[In] integrate(x**3*Shi(b*x)*sinh(b*x),x)
```

```
[Out] Integral(x**3*sinh(b*x)*Shi(b*x), x)
```

Maxima [F]

$$\int x^3 \sinh(bx) \text{Shi}(bx) dx = \int x^3 \text{Shi}(bx) \sinh(bx) dx$$

```
[In] integrate(x^3*Shi(b*x)*sinh(b*x),x, algorithm="maxima")
```

```
[Out] integrate(x^3*Shi(b*x)*sinh(b*x), x)
```

Giac [**F**]

$$\int x^3 \sinh(bx) \operatorname{Shi}(bx) dx = \int x^3 \operatorname{Shi}(bx) \sinh(bx) dx$$

[In] integrate(x^3*Shi(b*x)*sinh(b*x),x, algorithm="giac")

[Out] integrate(x^3*Shi(b*x)*sinh(b*x), x)

Mupad [**F(-1)**]

Timed out.

$$\int x^3 \sinh(bx) \operatorname{Shi}(bx) dx = \int x^3 \operatorname{sinhint}(bx) \sinh(bx) dx$$

[In] int(x^3*sinhint(b*x)*sinh(b*x),x)

[Out] int(x^3*sinhint(b*x)*sinh(b*x), x)

3.46 $\int \frac{\cosh(bx)\mathbf{Shi}(bx)}{x^3} dx$

Optimal result	256
Rubi [N/A]	256
Mathematica [N/A]	257
Maple [N/A] (verified)	258
Fricas [N/A]	258
Sympy [N/A]	258
Maxima [N/A]	258
Giac [N/A]	259
Mupad [N/A]	259

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\cosh(bx)\mathbf{Shi}(bx)}{x^3} dx = -\frac{b \cosh(2bx)}{4x} - \frac{b \sinh^2(bx)}{2x} - \frac{\sinh(2bx)}{8x^2} - \frac{\cosh(bx)\mathbf{Shi}(bx)}{2x^2} - \frac{b \sinh(bx)\mathbf{Shi}(bx)}{2x} + b^2 \mathbf{Shi}(2bx) + \frac{1}{2} b^2 \text{Int}\left(\frac{\cosh(bx)\mathbf{Shi}(bx)}{x}, x\right)$$

[Out] 1/2*b^2*CannotIntegrate(cosh(b*x)*Shi(b*x)/x,x)-1/4*b*cosh(2*b*x)/x-1/2*cosh(b*x)*Shi(b*x)/x^2+b^2*Shi(2*b*x)-1/2*b*Shi(b*x)*sinh(b*x)/x-1/2*b*sinh(b*x)^2/x-1/8*sinh(2*b*x)/x^2

Rubi [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\cosh(bx)\mathbf{Shi}(bx)}{x^3} dx = \int \frac{\cosh(bx)\mathbf{Shi}(bx)}{x^3} dx$$

[In] Int[(Cosh[b*x]*SinhIntegral[b*x])/x^3,x]

[Out] -1/4*(b*Cosh[2*b*x])/x - (b*Sinh[b*x]^2)/(2*x) - Sinh[2*b*x]/(8*x^2) - (Cosh[b*x]*SinhIntegral[b*x])/(2*x^2) - (b*Sinh[b*x]*SinhIntegral[b*x])/(2*x) + b^2*SinhIntegral[2*b*x] + (b^2*Defer[Int]((Cosh[b*x]*SinhIntegral[b*x])/x,x))/2

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\cosh(bx)\text{Shi}(bx)}{2x^2} + \frac{1}{2}b \int \frac{\cosh(bx)\sinh(bx)}{bx^3} dx + \frac{1}{2}b \int \frac{\sinh(bx)\text{Shi}(bx)}{x^2} dx \\
&= -\frac{\cosh(bx)\text{Shi}(bx)}{2x^2} - \frac{b\sinh(bx)\text{Shi}(bx)}{2x} + \frac{1}{2} \int \frac{\cosh(bx)\sinh(bx)}{x^3} dx \\
&\quad + \frac{1}{2}b^2 \int \frac{\sinh^2(bx)}{bx^2} dx + \frac{1}{2}b^2 \int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx \\
&= -\frac{\cosh(bx)\text{Shi}(bx)}{2x^2} - \frac{b\sinh(bx)\text{Shi}(bx)}{2x} + \frac{1}{2} \int \frac{\sinh(2bx)}{2x^3} dx \\
&\quad + \frac{1}{2}b \int \frac{\sinh^2(bx)}{x^2} dx + \frac{1}{2}b^2 \int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx \\
&= -\frac{b\sinh^2(bx)}{2x} - \frac{\cosh(bx)\text{Shi}(bx)}{2x^2} - \frac{b\sinh(bx)\text{Shi}(bx)}{2x} + \frac{1}{4} \int \frac{\sinh(2bx)}{x^3} dx \\
&\quad - (ib^2) \int \frac{i\sinh(2bx)}{2x} dx + \frac{1}{2}b^2 \int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx \\
&= -\frac{b\sinh^2(bx)}{2x} - \frac{\sinh(2bx)}{8x^2} - \frac{\cosh(bx)\text{Shi}(bx)}{2x^2} - \frac{b\sinh(bx)\text{Shi}(bx)}{2x} \\
&\quad + \frac{1}{4}b \int \frac{\cosh(2bx)}{x^2} dx + \frac{1}{2}b^2 \int \frac{\sinh(2bx)}{x} dx + \frac{1}{2}b^2 \int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx \\
&= -\frac{b\cosh(2bx)}{4x} - \frac{b\sinh^2(bx)}{2x} - \frac{\sinh(2bx)}{8x^2} - \frac{\cosh(bx)\text{Shi}(bx)}{2x^2} - \frac{b\sinh(bx)\text{Shi}(bx)}{2x} \\
&\quad + \frac{1}{2}b^2\text{Shi}(2bx) + \frac{1}{2}b^2 \int \frac{\sinh(2bx)}{x} dx + \frac{1}{2}b^2 \int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx \\
&= -\frac{b\cosh(2bx)}{4x} - \frac{b\sinh^2(bx)}{2x} - \frac{\sinh(2bx)}{8x^2} - \frac{\cosh(bx)\text{Shi}(bx)}{2x^2} \\
&\quad - \frac{b\sinh(bx)\text{Shi}(bx)}{2x} + b^2\text{Shi}(2bx) + \frac{1}{2}b^2 \int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx
\end{aligned}$$

Mathematica [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx)\text{Shi}(bx)}{x^3} dx = \int \frac{\cosh(bx)\text{Shi}(bx)}{x^3} dx$$

[In] Integrate[(Cosh[b*x]*SinhIntegral[b*x])/x^3,x]

[Out] Integrate[(Cosh[b*x]*SinhIntegral[b*x])/x^3, x]

Maple [N/A] (verified)

Not integrable

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\cosh(bx) \operatorname{Shi}(bx)}{x^3} dx$$

[In] int(cosh(b*x)*Shi(b*x)/x^3,x)

[Out] int(cosh(b*x)*Shi(b*x)/x^3,x)

Fricas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx) \operatorname{Shi}(bx)}{x^3} dx = \int \frac{\operatorname{Shi}(bx) \cosh(bx)}{x^3} dx$$

[In] integrate(cosh(b*x)*Shi(b*x)/x^3,x, algorithm="fricas")

[Out] integral(cosh(b*x)*sinh_integral(b*x)/x^3, x)

Sympy [N/A]

Not integrable

Time = 2.73 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx) \operatorname{Shi}(bx)}{x^3} dx = \int \frac{\cosh(bx) \operatorname{Shi}(bx)}{x^3} dx$$

[In] integrate(cosh(b*x)*Shi(b*x)/x**3,x)

[Out] Integral(cosh(b*x)*Shi(b*x)/x**3, x)

Maxima [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx) \operatorname{Shi}(bx)}{x^3} dx = \int \frac{\operatorname{Shi}(bx) \cosh(bx)}{x^3} dx$$

[In] integrate(cosh(b*x)*Shi(b*x)/x^3,x, algorithm="maxima")

[Out] integrate(Shi(b*x)*cosh(b*x)/x^3, x)

Giac [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx)\text{Shi}(bx)}{x^3} dx = \int \frac{\text{Shi}(bx) \cosh(bx)}{x^3} dx$$

[In] integrate(cosh(b*x)*Shi(b*x)/x^3,x, algorithm="giac")

[Out] integrate(Shi(b*x)*cosh(b*x)/x^3, x)

Mupad [N/A]

Not integrable

Time = 5.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx)\text{Shi}(bx)}{x^3} dx = \int \frac{\sinhint(bx) \cosh(bx)}{x^3} dx$$

[In] int((sinhint(b*x)*cosh(b*x))/x^3,x)

[Out] int((sinhint(b*x)*cosh(b*x))/x^3, x)

3.47 $\int \frac{\cosh(bx)\mathbf{Shi}(bx)}{x^2} dx$

Optimal result	260
Rubi [A] (verified)	260
Mathematica [A] (verified)	262
Maple [F]	262
Fricas [F]	262
Sympy [F]	262
Maxima [F]	263
Giac [F]	263
Mupad [F(-1)]	263

Optimal result

Integrand size = 12, antiderivative size = 44

$$\int \frac{\cosh(bx)\mathbf{Shi}(bx)}{x^2} dx = b\mathbf{Chi}(2bx) - \frac{\sinh(2bx)}{2x} - \frac{\cosh(bx)\mathbf{Shi}(bx)}{x} + \frac{1}{2}b\mathbf{Shi}(bx)^2$$

[Out] $b*\mathbf{Chi}(2*b*x) - \cosh(b*x)*\mathbf{Shi}(b*x)/x + 1/2*b*\mathbf{Shi}(b*x)^2 - 1/2*\sinh(2*b*x)/x$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6685, 6818, 12, 5556, 3378, 3382}

$$\int \frac{\cosh(bx)\mathbf{Shi}(bx)}{x^2} dx = b\mathbf{Chi}(2bx) + \frac{1}{2}b\mathbf{Shi}(bx)^2 - \frac{\mathbf{Shi}(bx)\cosh(bx)}{x} - \frac{\sinh(2bx)}{2x}$$

[In] $\text{Int}[(\text{Cosh}[b*x]*\text{SinhIntegral}[b*x])/x^2, x]$

[Out] $b*\text{CoshIntegral}[2*b*x] - \text{Sinh}[2*b*x]/(2*x) - (\text{Cosh}[b*x]*\text{SinhIntegral}[b*x])/x + (b*\text{SinhIntegral}[b*x]^2)/2$

Rule 12

$\text{Int}[(a_*)(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /;$ FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 3378

$\text{Int}[(c_*) + (d_*)(x_)^{(m_*)}\sin[(e_*) + (f_*)(x_)], x_Symbol] \rightarrow \text{Simp}[(c + d*x)^{(m+1)}*(\text{Sin}[e + f*x]/(d*(m+1))), x] - \text{Dist}[f/(d*(m+1)), \text{Int}[(c + d*x)^{(m+1)}*\text{Cos}[e + f*x], x], x] /;$ FreeQ[{c, d, e, f}, x] && LtQ[m, -1

]

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol]
:> Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x]
&& EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] :> Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

Rule 6685

```
Int[Cosh[(a_.) + (b_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.)*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] :> Simp[(e + f*x)^(m + 1)*Cosh[a + b*x]*(SinhIntegral[c + d*x]/(f*(m + 1))), x]
+ (-Dist[b/(f*(m + 1)), Int[(e + f*x)^(m + 1)*Sinh[a + b*x]*SinhIntegral[c + d*x], x], x]
- Dist[d/(f*(m + 1)), Int[(e + f*x)^(m + 1)*Cosh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x]) /; FreeQ[{a,
b, c, d, e, f}, x] && ILtQ[m, -1]
```

Rule 6818

```
Int[(u_)*(y_)^(m_.), x_Symbol] :> With[{q = DerivativeDivides[y, u, x]}, Simp[q*(y^(m + 1)/(m + 1)), x] /;
!FalseQ[q]] /; FreeQ[m, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\cosh(bx)\text{Shi}(bx)}{x} + b \int \frac{\cosh(bx)\sinh(bx)}{bx^2} dx + b \int \frac{\sinh(bx)\text{Shi}(bx)}{x} dx \\
&= -\frac{\cosh(bx)\text{Shi}(bx)}{x} + \frac{1}{2}b\text{Shi}(bx)^2 + \int \frac{\cosh(bx)\sinh(bx)}{x^2} dx \\
&= -\frac{\cosh(bx)\text{Shi}(bx)}{x} + \frac{1}{2}b\text{Shi}(bx)^2 + \int \frac{\sinh(2bx)}{2x^2} dx \\
&= -\frac{\cosh(bx)\text{Shi}(bx)}{x} + \frac{1}{2}b\text{Shi}(bx)^2 + \frac{1}{2} \int \frac{\sinh(2bx)}{x^2} dx \\
&= -\frac{\sinh(2bx)}{2x} - \frac{\cosh(bx)\text{Shi}(bx)}{x} + \frac{1}{2}b\text{Shi}(bx)^2 + b \int \frac{\cosh(2bx)}{x} dx \\
&= b\text{Chi}(2bx) - \frac{\sinh(2bx)}{2x} - \frac{\cosh(bx)\text{Shi}(bx)}{x} + \frac{1}{2}b\text{Shi}(bx)^2
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00

$$\int \frac{\cosh(bx)\text{Shi}(bx)}{x^2} dx = b\text{Chi}(2bx) - \frac{\sinh(2bx)}{2x} - \frac{\cosh(bx)\text{Shi}(bx)}{x} + \frac{1}{2}b\text{Shi}(bx)^2$$

[In] Integrate[(Cosh[b*x]*SinhIntegral[b*x])/x^2,x]

[Out] b*CoshIntegral[2*b*x] - Sinh[2*b*x]/(2*x) - (Cosh[b*x]*SinhIntegral[b*x])/x + (b*SinhIntegral[b*x]^2)/2

Maple [F]

$$\int \frac{\cosh(bx)\text{Shi}(bx)}{x^2} dx$$

[In] int(cosh(b*x)*Shi(b*x)/x^2,x)

[Out] int(cosh(b*x)*Shi(b*x)/x^2,x)

Fricas [F]

$$\int \frac{\cosh(bx)\text{Shi}(bx)}{x^2} dx = \int \frac{\text{Shi}(bx)\cosh(bx)}{x^2} dx$$

[In] integrate(cosh(b*x)*Shi(b*x)/x^2,x, algorithm="fricas")

[Out] integral(cosh(b*x)*sinh_integral(b*x)/x^2, x)

Sympy [F]

$$\int \frac{\cosh(bx)\text{Shi}(bx)}{x^2} dx = \int \frac{\cosh(bx)\text{Shi}(bx)}{x^2} dx$$

[In] integrate(cosh(b*x)*Shi(b*x)/x**2,x)

[Out] Integral(cosh(b*x)*Shi(b*x)/x**2, x)

Maxima [F]

$$\int \frac{\cosh(bx)\operatorname{Shi}(bx)}{x^2} dx = \int \frac{\operatorname{Shi}(bx)\cosh(bx)}{x^2} dx$$

[In] integrate(cosh(b*x)*Shi(b*x)/x^2,x, algorithm="maxima")

[Out] integrate(Shi(b*x)*cosh(b*x)/x^2, x)

Giac [F]

$$\int \frac{\cosh(bx)\operatorname{Shi}(bx)}{x^2} dx = \int \frac{\operatorname{Shi}(bx)\cosh(bx)}{x^2} dx$$

[In] integrate(cosh(b*x)*Shi(b*x)/x^2,x, algorithm="giac")

[Out] integrate(Shi(b*x)*cosh(b*x)/x^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cosh(bx)\operatorname{Shi}(bx)}{x^2} dx = \int \frac{\operatorname{sinhint}(bx)\cosh(bx)}{x^2} dx$$

[In] int((sinhint(b*x)*cosh(b*x))/x^2,x)

[Out] int((sinhint(b*x)*cosh(b*x))/x^2, x)

3.48 $\int \frac{\cosh(bx)\mathbf{Shi}(bx)}{x} dx$

Optimal result	264
Rubi [N/A]	264
Mathematica [N/A]	265
Maple [N/A] (verified)	265
Fricas [N/A]	265
Sympy [N/A]	265
Maxima [N/A]	266
Giac [N/A]	266
Mupad [N/A]	266

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\cosh(bx)\mathbf{Shi}(bx)}{x} dx = \text{Int}\left(\frac{\cosh(bx)\mathbf{Shi}(bx)}{x}, x\right)$$

[Out] CannotIntegrate(cosh(b*x)*Shi(b*x)/x,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\cosh(bx)\mathbf{Shi}(bx)}{x} dx = \int \frac{\cosh(bx)\mathbf{Shi}(bx)}{x} dx$$

[In] Int[(Cosh[b*x]*SinhIntegral[b*x])/x,x]

[Out] Defer[Int] [(Cosh[b*x]*SinhIntegral[b*x])/x, x]

Rubi steps

$$\text{integral} = \int \frac{\cosh(bx)\mathbf{Shi}(bx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx = \int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx$$

[In] Integrate[(Cosh[b*x]*SinhIntegral[b*x])/x,x]

[Out] Integrate[(Cosh[b*x]*SinhIntegral[b*x])/x, x]

Maple [N/A] (verified)

Not integrable

Time = 0.23 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx$$

[In] int(cosh(b*x)*Shi(b*x)/x,x)

[Out] int(cosh(b*x)*Shi(b*x)/x,x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx = \int \frac{\text{Shi}(bx)\cosh(bx)}{x} dx$$

[In] integrate(cosh(b*x)*Shi(b*x)/x,x, algorithm="fricas")

[Out] integral(cosh(b*x)*sinh_integral(b*x)/x, x)

Sympy [N/A]

Not integrable

Time = 2.42 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx = \int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx$$

[In] integrate(cosh(b*x)*Shi(b*x)/x,x)

[Out] Integral(cosh(b*x)*Shi(b*x)/x, x)

Maxima [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx = \int \frac{\text{Shi}(bx) \cosh(bx)}{x} dx$$

[In] integrate(cosh(b*x)*Shi(b*x)/x,x, algorithm="maxima")

[Out] integrate(Shi(b*x)*cosh(b*x)/x, x)

Giac [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx = \int \frac{\text{Shi}(bx) \cosh(bx)}{x} dx$$

[In] integrate(cosh(b*x)*Shi(b*x)/x,x, algorithm="giac")

[Out] integrate(Shi(b*x)*cosh(b*x)/x, x)

Mupad [N/A]

Not integrable

Time = 4.99 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx)\text{Shi}(bx)}{x} dx = \int \frac{\sinhint(bx) \cosh(bx)}{x} dx$$

[In] int((sinhint(b*x)*cosh(b*x))/x,x)

[Out] int((sinhint(b*x)*cosh(b*x))/x, x)

3.49 $\int \cosh(bx)\mathbf{Shi}(bx) dx$

Optimal result	267
Rubi [A] (verified)	267
Mathematica [A] (verified)	268
Maple [A] (verified)	269
Fricas [F]	269
Sympy [F]	269
Maxima [F]	269
Giac [F]	270
Mupad [F(-1)]	270

Optimal result

Integrand size = 9, antiderivative size = 34

$$\int \cosh(bx)\mathbf{Shi}(bx) dx = -\frac{\text{Chi}(2bx)}{2b} + \frac{\log(x)}{2b} + \frac{\sinh(bx)\mathbf{Shi}(bx)}{b}$$

[Out] $-1/2*\text{Chi}(2*b*x)/b+1/2*\ln(x)/b+\text{Shi}(b*x)*\sinh(b*x)/b$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {6681, 12, 3393, 3382}

$$\int \cosh(bx)\mathbf{Shi}(bx) dx = -\frac{\text{Chi}(2bx)}{2b} + \frac{\mathbf{Shi}(bx) \sinh(bx)}{b} + \frac{\log(x)}{2b}$$

[In] $\text{Int}[\text{Cosh}[b*x]*\text{SinhIntegral}[b*x], x]$

[Out] $-1/2*\text{CoshIntegral}[2*b*x]/b + \text{Log}[x]/(2*b) + (\text{Sinh}[b*x]*\text{SinhIntegral}[b*x])/b$

Rule 12

$\text{Int}[(a_*)(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \&\& \text{!MatchQ}[u, (b_*)(v_)] /; \text{FreeQ}[b, x]$

Rule 3382

$\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] \rightarrow \text{Simp}[\text{CoshIntegral}[c*f*(fz/d) + f*fz*x]/d, x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \&\& \text{EqQ}[d*(e - \text{Pi}/2) - c*f*fz*I, 0]$

Rule 3393

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] :> Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 6681

```
Int[Cosh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[Sinh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\sinh(bx)\text{Shi}(bx)}{b} - \int \frac{\sinh^2(bx)}{bx} dx \\
 &= \frac{\sinh(bx)\text{Shi}(bx)}{b} - \frac{\int \frac{\sinh^2(bx)}{x} dx}{b} \\
 &= \frac{\sinh(bx)\text{Shi}(bx)}{b} + \frac{\int \left(\frac{1}{2x} - \frac{\cosh(2bx)}{2x} \right) dx}{b} \\
 &= \frac{\log(x)}{2b} + \frac{\sinh(bx)\text{Shi}(bx)}{b} - \frac{\int \frac{\cosh(2bx)}{x} dx}{2b} \\
 &= -\frac{\text{Chi}(2bx)}{2b} + \frac{\log(x)}{2b} + \frac{\sinh(bx)\text{Shi}(bx)}{b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.06

$$\int \cosh(bx)\text{Shi}(bx) dx = -\frac{\text{Chi}(2bx)}{2b} + \frac{\log(bx)}{2b} + \frac{\sinh(bx)\text{Shi}(bx)}{b}$$

```
[In] Integrate[Cosh[b*x]*SinhIntegral[b*x],x]
```

```
[Out] -1/2*CoshIntegral[2*b*x]/b + Log[b*x]/(2*b) + (Sinh[b*x]*SinhIntegral[b*x])/b
```

Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$\frac{\text{Shi}(bx) \sinh(bx) + \frac{\ln(bx)}{2} - \frac{\text{Chi}(2bx)}{2}}{b}$	28
default	$\frac{\text{Shi}(bx) \sinh(bx) + \frac{\ln(bx)}{2} - \frac{\text{Chi}(2bx)}{2}}{b}$	28

[In] `int(cosh(b*x)*Shi(b*x),x,method=_RETURNVERBOSE)`

[Out] `1/b*(Shi(b*x)*sinh(b*x)+1/2*ln(b*x)-1/2*Chi(2*b*x))`

Fricas [F]

$$\int \cosh(bx) \text{Shi}(bx) dx = \int \text{Shi}(bx) \cosh(bx) dx$$

[In] `integrate(cosh(b*x)*Shi(b*x),x, algorithm="fricas")`

[Out] `integral(cosh(b*x)*sinh_integral(b*x), x)`

Sympy [F]

$$\int \cosh(bx) \text{Shi}(bx) dx = \int \cosh(bx) \text{Shi}(bx) dx$$

[In] `integrate(cosh(b*x)*Shi(b*x),x)`

[Out] `Integral(cosh(b*x)*Shi(b*x), x)`

Maxima [F]

$$\int \cosh(bx) \text{Shi}(bx) dx = \int \text{Shi}(bx) \cosh(bx) dx$$

[In] `integrate(cosh(b*x)*Shi(b*x),x, algorithm="maxima")`

[Out] `integrate(Shi(b*x)*cosh(b*x), x)`

Giac [F]

$$\int \cosh(bx)\operatorname{Shi}(bx) dx = \int \operatorname{Shi}(bx) \cosh(bx) dx$$

[In] `integrate(cosh(b*x)*Shi(b*x),x, algorithm="giac")`

[Out] `integrate(Shi(b*x)*cosh(b*x), x)`

Mupad [F(-1)]

Timed out.

$$\int \cosh(bx)\operatorname{Shi}(bx) dx = \int \operatorname{sinhint}(bx) \cosh(bx) dx$$

[In] `int(sinhint(b*x)*cosh(b*x),x)`

[Out] `int(sinhint(b*x)*cosh(b*x), x)`

3.50 $\int x \cosh(bx) \text{Shi}(bx) dx$

Optimal result	271
Rubi [A] (verified)	271
Mathematica [A] (verified)	273
Maple [A] (verified)	273
Fricas [F]	274
Sympy [F]	274
Maxima [F]	274
Giac [F]	274
Mupad [F(-1)]	275

Optimal result

Integrand size = 10, antiderivative size = 62

$$\int x \cosh(bx) \text{Shi}(bx) dx = \frac{x}{2b} - \frac{\cosh(bx) \sinh(bx)}{2b^2} - \frac{\cosh(bx) \text{Shi}(bx)}{b^2} + \frac{x \sinh(bx) \text{Shi}(bx)}{b} + \frac{\text{Shi}(2bx)}{2b^2}$$

[Out] $1/2*x/b - \cosh(b*x)*\text{Shi}(b*x)/b^2 + 1/2*\text{Shi}(2*b*x)/b^2 - 1/2*\cosh(b*x)*\sinh(b*x)/b^2 + x*\text{Shi}(b*x)*\sinh(b*x)/b$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {6683, 12, 2715, 8, 6675, 5556, 3379}

$$\int x \cosh(bx) \text{Shi}(bx) dx = \frac{\text{Shi}(2bx)}{2b^2} - \frac{\text{Shi}(bx) \cosh(bx)}{b^2} - \frac{\sinh(bx) \cosh(bx)}{2b^2} + \frac{x \text{Shi}(bx) \sinh(bx)}{b} + \frac{x}{2b}$$

[In] $\text{Int}[x*\text{Cosh}[b*x]*\text{SinhIntegral}[b*x], x]$

[Out] $x/(2*b) - (\text{Cosh}[b*x]*\text{Sinh}[b*x])/(2*b^2) - (\text{Cosh}[b*x]*\text{SinhIntegral}[b*x])/b^2 + (x*\text{Sinh}[b*x]*\text{SinhIntegral}[b*x])/b + \text{SinhIntegral}[2*b*x]/(2*b^2)$

Rule 8

$\text{Int}[a_, x_Symbol] \text{ :> } \text{Simp}[a*x, x] \text{ /; } \text{FreeQ}[a, x]$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 2715

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_)*(x_)]/((c_.) + (d_)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 5556

Int[Cosh[(a_.) + (b_)*(x_)]^(p_)*((c_.) + (d_)*(x_))^(m_)*Sinh[(a_.) + (b_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]

Rule 6675

Int[Sinh[(a_.) + (b_)*(x_)]*SinhIntegral[(c_.) + (d_)*(x_)], x_Symbol] := Simp[Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]

Rule 6683

Int[Cosh[(a_.) + (b_)*(x_)]*((e_.) + (f_)*(x_))^(m_)*SinhIntegral[(c_.) + (d_)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Sinh[a + b*x]*(SinhIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*SinhIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x \sinh(bx) \text{Shi}(bx)}{b} - \frac{\int \sinh(bx) \text{Shi}(bx) dx}{b} - \int \frac{\sinh^2(bx)}{b} dx \\
 &= -\frac{\cosh(bx) \text{Shi}(bx)}{b^2} + \frac{x \sinh(bx) \text{Shi}(bx)}{b} + \frac{\int \frac{\cosh(bx) \sinh(bx)}{bx} dx}{b} - \frac{\int \sinh^2(bx) dx}{b} \\
 &= -\frac{\cosh(bx) \sinh(bx)}{2b^2} - \frac{\cosh(bx) \text{Shi}(bx)}{b^2} + \frac{x \sinh(bx) \text{Shi}(bx)}{b} + \frac{\int \frac{\cosh(bx) \sinh(bx)}{x} dx}{b^2} + \frac{\int 1 dx}{2b}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{x}{2b} - \frac{\cosh(bx) \sinh(bx)}{2b^2} - \frac{\cosh(bx) \operatorname{Shi}(bx)}{b^2} + \frac{x \sinh(bx) \operatorname{Shi}(bx)}{b} + \frac{\int \frac{\sinh(2bx)}{2x} dx}{b^2} \\
&= \frac{x}{2b} - \frac{\cosh(bx) \sinh(bx)}{2b^2} - \frac{\cosh(bx) \operatorname{Shi}(bx)}{b^2} + \frac{x \sinh(bx) \operatorname{Shi}(bx)}{b} + \frac{\int \frac{\sinh(2bx)}{x} dx}{2b^2} \\
&= \frac{x}{2b} - \frac{\cosh(bx) \sinh(bx)}{2b^2} - \frac{\cosh(bx) \operatorname{Shi}(bx)}{b^2} + \frac{x \sinh(bx) \operatorname{Shi}(bx)}{b} + \frac{\operatorname{Shi}(2bx)}{2b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.74

$$\int x \cosh(bx) \operatorname{Shi}(bx) dx = \frac{2bx - \sinh(2bx) + 4(-\cosh(bx) + bx \sinh(bx)) \operatorname{Shi}(bx) + 2\operatorname{Shi}(2bx)}{4b^2}$$

[In] Integrate[x*Cosh[b*x]*SinhIntegral[b*x],x]

[Out] (2*b*x - Sinh[2*b*x] + 4*(-Cosh[b*x] + b*x*Sinh[b*x])*SinhIntegral[b*x] + 2*SinhIntegral[2*b*x])/(4*b^2)

Maple [A] (verified)

Time = 0.80 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.74

method	result	size
derivativedivides	$\frac{\operatorname{Shi}(bx)(bx \sinh(bx) - \cosh(bx)) - \frac{\cosh(bx) \sinh(bx)}{2} + \frac{bx}{2} + \frac{\operatorname{Shi}(2bx)}{2}}{b^2}$	46
default	$\frac{\operatorname{Shi}(bx)(bx \sinh(bx) - \cosh(bx)) - \frac{\cosh(bx) \sinh(bx)}{2} + \frac{bx}{2} + \frac{\operatorname{Shi}(2bx)}{2}}{b^2}$	46

[In] int(x*cosh(b*x)*Shi(b*x),x,method=_RETURNVERBOSE)

[Out] 1/b^2*(Shi(b*x)*(b*x*sinh(b*x)-cosh(b*x))-1/2*cosh(b*x)*sinh(b*x)+1/2*b*x+1/2*Shi(2*b*x))

Fricas [F]

$$\int x \cosh(bx) \operatorname{Shi}(bx) dx = \int x \operatorname{Shi}(bx) \cosh(bx) dx$$

```
[In] integrate(x*cosh(b*x)*Shi(b*x),x, algorithm="fricas")
```

```
[Out] integral(x*cosh(b*x)*sinh_integral(b*x), x)
```

Sympy [F]

$$\int x \cosh(bx) \operatorname{Shi}(bx) dx = \int x \cosh(bx) \operatorname{Shi}(bx) dx$$

```
[In] integrate(x*cosh(b*x)*Shi(b*x),x)
```

```
[Out] Integral(x*cosh(b*x)*Shi(b*x), x)
```

Maxima [F]

$$\int x \cosh(bx) \operatorname{Shi}(bx) dx = \int x \operatorname{Shi}(bx) \cosh(bx) dx$$

```
[In] integrate(x*cosh(b*x)*Shi(b*x),x, algorithm="maxima")
```

```
[Out] integrate(x*Shi(b*x)*cosh(b*x), x)
```

Giac [F]

$$\int x \cosh(bx) \operatorname{Shi}(bx) dx = \int x \operatorname{Shi}(bx) \cosh(bx) dx$$

```
[In] integrate(x*cosh(b*x)*Shi(b*x),x, algorithm="giac")
```

```
[Out] integrate(x*Shi(b*x)*cosh(b*x), x)
```

Mupad [F(-1)]

Timed out.

$$\int x \cosh(bx) \operatorname{Shi}(bx) dx = \int x \operatorname{sinhint}(bx) \cosh(bx) dx$$

```
[In] int(x*sinhint(b*x)*cosh(b*x),x)
```

```
[Out] int(x*sinhint(b*x)*cosh(b*x), x)
```

3.51 $\int x^2 \cosh(bx) \operatorname{Shi}(bx) dx$

Optimal result	276
Rubi [A] (verified)	276
Mathematica [A] (verified)	279
Maple [A] (verified)	279
Fricas [F]	279
Sympy [F]	280
Maxima [F]	280
Giac [F]	280
Mupad [F(-1)]	280

Optimal result

Integrand size = 12, antiderivative size = 98

$$\int x^2 \cosh(bx) \operatorname{Shi}(bx) dx = \frac{x^2}{4b} - \frac{\operatorname{Chi}(2bx)}{b^3} + \frac{\log(x)}{b^3} - \frac{x \cosh(bx) \sinh(bx)}{2b^2} + \frac{5 \sinh^2(bx)}{4b^3} - \frac{2x \cosh(bx) \operatorname{Shi}(bx)}{b^2} + \frac{2 \sinh(bx) \operatorname{Shi}(bx)}{b^3} + \frac{x^2 \sinh(bx) \operatorname{Shi}(bx)}{b}$$

[Out] $1/4*x^2/b - \operatorname{Chi}(2*b*x)/b^3 + \ln(x)/b^3 - 2*x*\cosh(b*x)*\operatorname{Shi}(b*x)/b^2 - 1/2*x*\cosh(b*x)*\sinh(b*x)/b^2 + 2*\operatorname{Shi}(b*x)*\sinh(b*x)/b^3 + x^2*\operatorname{Shi}(b*x)*\sinh(b*x)/b + 5/4*\sinh(b*x)^2/b^3$

Rubi [A] (verified)

Time = 0.09 (sec), antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6683, 12, 3391, 30, 6677, 2644, 6681, 3393, 3382}

$$\int x^2 \cosh(bx) \operatorname{Shi}(bx) dx = -\frac{\operatorname{Chi}(2bx)}{b^3} + \frac{2\operatorname{Shi}(bx) \sinh(bx)}{b^3} + \frac{\log(x)}{b^3} + \frac{5 \sinh^2(bx)}{4b^3} - \frac{2x \operatorname{Shi}(bx) \cosh(bx)}{b^2} - \frac{x \sinh(bx) \cosh(bx)}{2b^2} + \frac{x^2 \operatorname{Shi}(bx) \sinh(bx)}{b} + \frac{x^2}{4b}$$

[In] $\operatorname{Int}[x^2*\operatorname{Cosh}[b*x]*\operatorname{SinhIntegral}[b*x],x]$

[Out] $x^2/(4*b) - \operatorname{CoshIntegral}[2*b*x]/b^3 + \operatorname{Log}[x]/b^3 - (x*\operatorname{Cosh}[b*x]*\operatorname{Sinh}[b*x])/(2*b^2) + (5*\operatorname{Sinh}[b*x]^2)/(4*b^3) - (2*x*\operatorname{Cosh}[b*x]*\operatorname{SinhIntegral}[b*x])/b^2 + (2*\operatorname{Sinh}[b*x]*\operatorname{SinhIntegral}[b*x])/b^3 + (x^2*\operatorname{Sinh}[b*x]*\operatorname{SinhIntegral}[b*x])/b$

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 30

```
Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && N
eQ[m, -1]
```

Rule 2644

```
Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)]^(m_.), x_
Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*
Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(In
tegerQ[(m - 1)/2] && LtQ[0, m, n])
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz
}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3391

```
Int[((c_.) + (d_.)*(x_))*((b_.)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] :=
Simp[d*((b*SIN[e + f*x])^n/(f^2*n^2)), x] + (Dist[b^2*((n - 1)/n), Int[(c
+ d*x)*(b*SIN[e + f*x])^(n - 2), x], x] - Simp[b*(c + d*x)*Cos[e + f*x]*((b
*SIN[e + f*x])^(n - 1)/(f*n)), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1
]
```

Rule 3393

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := In
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 6677

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(SinhIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Sinh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*SinhIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6681

```
Int[Cosh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :>
Simp[Sinh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6683

```
Int[Cosh[(a_.) + (b_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.)*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] :> Simp[(e + f*x)^m*Sinh[a + b*x]*(SinhIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Sinh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*SinhIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x^2 \sinh(bx) \text{Shi}(bx)}{b} - \frac{2 \int x \sinh(bx) \text{Shi}(bx) dx}{b} - \int \frac{x \sinh^2(bx)}{b} dx \\
&= -\frac{2x \cosh(bx) \text{Shi}(bx)}{b^2} + \frac{x^2 \sinh(bx) \text{Shi}(bx)}{b} \\
&\quad + \frac{2 \int \cosh(bx) \text{Shi}(bx) dx}{b^2} - \frac{\int x \sinh^2(bx) dx}{b} + \frac{2 \int \frac{\cosh(bx) \sinh(bx)}{b} dx}{b} \\
&= -\frac{x \cosh(bx) \sinh(bx)}{2b^2} + \frac{\sinh^2(bx)}{4b^3} - \frac{2x \cosh(bx) \text{Shi}(bx)}{b^2} + \frac{2 \sinh(bx) \text{Shi}(bx)}{b^3} \\
&\quad + \frac{x^2 \sinh(bx) \text{Shi}(bx)}{b} + \frac{2 \int \cosh(bx) \sinh(bx) dx}{b^2} - \frac{2 \int \frac{\sinh^2(bx)}{bx} dx}{b^2} + \frac{\int x dx}{2b} \\
&= \frac{x^2}{4b} - \frac{x \cosh(bx) \sinh(bx)}{2b^2} + \frac{\sinh^2(bx)}{4b^3} - \frac{2x \cosh(bx) \text{Shi}(bx)}{b^2} + \frac{2 \sinh(bx) \text{Shi}(bx)}{b^3} \\
&\quad + \frac{x^2 \sinh(bx) \text{Shi}(bx)}{b} - \frac{2 \int \frac{\sinh^2(bx)}{x} dx}{b^3} - \frac{2 \text{Subst}(\int x dx, x, i \sinh(bx))}{b^3} \\
&= \frac{x^2}{4b} - \frac{x \cosh(bx) \sinh(bx)}{2b^2} + \frac{5 \sinh^2(bx)}{4b^3} - \frac{2x \cosh(bx) \text{Shi}(bx)}{b^2} \\
&\quad + \frac{2 \sinh(bx) \text{Shi}(bx)}{b^3} + \frac{x^2 \sinh(bx) \text{Shi}(bx)}{b} + \frac{2 \int \left(\frac{1}{2x} - \frac{\cosh(2bx)}{2x} \right) dx}{b^3} \\
&= \frac{x^2}{4b} + \frac{\log(x)}{b^3} - \frac{x \cosh(bx) \sinh(bx)}{2b^2} + \frac{5 \sinh^2(bx)}{4b^3} - \frac{2x \cosh(bx) \text{Shi}(bx)}{b^2} \\
&\quad + \frac{2 \sinh(bx) \text{Shi}(bx)}{b^3} + \frac{x^2 \sinh(bx) \text{Shi}(bx)}{b} - \frac{\int \frac{\cosh(2bx)}{x} dx}{b^3} \\
&= \frac{x^2}{4b} - \frac{\text{Chi}(2bx)}{b^3} + \frac{\log(x)}{b^3} - \frac{x \cosh(bx) \sinh(bx)}{2b^2} + \frac{5 \sinh^2(bx)}{4b^3} \\
&\quad - \frac{2x \cosh(bx) \text{Shi}(bx)}{b^2} + \frac{2 \sinh(bx) \text{Shi}(bx)}{b^3} + \frac{x^2 \sinh(bx) \text{Shi}(bx)}{b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.73

$$\int x^2 \cosh(bx) \operatorname{Shi}(bx) dx = \frac{2b^2x^2 + 5 \cosh(2bx) - 8\operatorname{Chi}(2bx) + 8 \log(x) - 2bx \sinh(2bx) + 8(-2bx \cosh(bx) + (2 + b^2x^2) \sinh(bx)) \operatorname{Shi}(bx)}{8b^3}$$

[In] Integrate[x^2*Cosh[b*x]*SinhIntegral[b*x],x]

[Out] (2*b^2*x^2 + 5*Cosh[2*b*x] - 8*CoshIntegral[2*b*x] + 8*Log[x] - 2*b*x*Sinh[2*b*x] + 8*(-2*b*x*Cosh[b*x] + (2 + b^2*x^2)*Sinh[b*x])*SinhIntegral[b*x])/(8*b^3)

Maple [A] (verified)

Time = 1.03 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.78

method	result	size
derivativedivides	$\frac{\operatorname{Shi}(bx)(b^2x^2 \sinh(bx) - 2bx \cosh(bx) + 2 \sinh(bx)) - \frac{bx \cosh(bx) \sinh(bx)}{2} + \frac{b^2x^2}{4} + \frac{5 \cosh(bx)^2}{4} + \ln(bx) - \operatorname{Chi}(2bx)}{b^3}$	76
default	$\frac{\operatorname{Shi}(bx)(b^2x^2 \sinh(bx) - 2bx \cosh(bx) + 2 \sinh(bx)) - \frac{bx \cosh(bx) \sinh(bx)}{2} + \frac{b^2x^2}{4} + \frac{5 \cosh(bx)^2}{4} + \ln(bx) - \operatorname{Chi}(2bx)}{b^3}$	76

[In] int(x^2*cosh(b*x)*Shi(b*x),x,method=_RETURNVERBOSE)

[Out] 1/b^3*(Shi(b*x)*(b^2*x^2*sinh(b*x)-2*b*x*cosh(b*x)+2*sinh(b*x))-1/2*b*x*cosh(b*x)*sinh(b*x)+1/4*b^2*x^2+5/4*cosh(b*x)^2+ln(b*x)-Chi(2*b*x))

Fricas [F]

$$\int x^2 \cosh(bx) \operatorname{Shi}(bx) dx = \int x^2 \operatorname{Shi}(bx) \cosh(bx) dx$$

[In] integrate(x^2*cosh(b*x)*Shi(b*x),x, algorithm="fricas")

[Out] integral(x^2*cosh(b*x)*sinh_integral(b*x), x)

Sympy [F]

$$\int x^2 \cosh(bx) \operatorname{Shi}(bx) dx = \int x^2 \cosh(bx) \operatorname{Shi}(bx) dx$$

```
[In] integrate(x**2*cosh(b*x)*Shi(b*x),x)
```

```
[Out] Integral(x**2*cosh(b*x)*Shi(b*x), x)
```

Maxima [F]

$$\int x^2 \cosh(bx) \operatorname{Shi}(bx) dx = \int x^2 \operatorname{Shi}(bx) \cosh(bx) dx$$

```
[In] integrate(x^2*cosh(b*x)*Shi(b*x),x, algorithm="maxima")
```

```
[Out] integrate(x^2*Shi(b*x)*cosh(b*x), x)
```

Giac [F]

$$\int x^2 \cosh(bx) \operatorname{Shi}(bx) dx = \int x^2 \operatorname{Shi}(bx) \cosh(bx) dx$$

```
[In] integrate(x^2*cosh(b*x)*Shi(b*x),x, algorithm="giac")
```

```
[Out] integrate(x^2*Shi(b*x)*cosh(b*x), x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \cosh(bx) \operatorname{Shi}(bx) dx = \int x^2 \operatorname{sinhint}(bx) \cosh(bx) dx$$

```
[In] int(x^2*sinhint(b*x)*cosh(b*x),x)
```

```
[Out] int(x^2*sinhint(b*x)*cosh(b*x), x)
```


3.52 $\int x^3 \cosh(bx) \text{Shi}(bx) dx$

Optimal result	281
Rubi [A] (verified)	281
Mathematica [A] (verified)	284
Maple [A] (verified)	285
Fricas [F]	285
Sympy [F]	285
Maxima [F]	285
Giac [F]	286
Mupad [F(-1)]	286

Optimal result

Integrand size = 12, antiderivative size = 128

$$\int x^3 \cosh(bx) \text{Shi}(bx) dx = \frac{4x}{b^3} + \frac{x^3}{6b} - \frac{4 \cosh(bx) \sinh(bx)}{b^4} - \frac{x^2 \cosh(bx) \sinh(bx)}{2b^2} + \frac{2x \sinh^2(bx)}{b^3} - \frac{6 \cosh(bx) \text{Shi}(bx)}{b^4} - \frac{3x^2 \cosh(bx) \text{Shi}(bx)}{b^2} + \frac{6x \sinh(bx) \text{Shi}(bx)}{b^3} + \frac{x^3 \sinh(bx) \text{Shi}(bx)}{b} + \frac{3 \text{Shi}(2bx)}{b^4}$$

[Out] $4*x/b^3 + 1/6*x^3/b - 6*\cosh(b*x)*\text{Shi}(b*x)/b^4 - 3*x^2*\cosh(b*x)*\text{Shi}(b*x)/b^2 + 3*\text{Shi}(2*b*x)/b^4 - 4*\cosh(b*x)*\sinh(b*x)/b^4 - 1/2*x^2*\cosh(b*x)*\sinh(b*x)/b^2 + 6*x*\text{Shi}(b*x)*\sinh(b*x)/b^3 + x^3*\text{Shi}(b*x)*\sinh(b*x)/b + 2*x*\sinh(b*x)^2/b^3$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 20, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$, Rules used = {6683, 12, 3392, 30, 2715, 8, 6677, 5480, 6675, 5556, 3379}

$$\int x^3 \cosh(bx) \text{Shi}(bx) dx = \frac{3 \text{Shi}(2bx)}{b^4} - \frac{6 \text{Shi}(bx) \cosh(bx)}{b^4} - \frac{4 \sinh(bx) \cosh(bx)}{b^4} + \frac{6x \text{Shi}(bx) \sinh(bx)}{b^3} + \frac{4x}{b^3} + \frac{2x \sinh^2(bx)}{b^3} - \frac{3x^2 \text{Shi}(bx) \cosh(bx)}{b^2} - \frac{x^2 \sinh(bx) \cosh(bx)}{2b^2} + \frac{x^3 \text{Shi}(bx) \sinh(bx)}{b} + \frac{x^3}{6b}$$

[In] $\text{Int}[x^3 * \text{Cosh}[b*x] * \text{SinhIntegral}[b*x], x]$

[Out] $(4*x)/b^3 + x^3/(6*b) - (4*\text{Cosh}[b*x]*\text{Sinh}[b*x])/b^4 - (x^2*\text{Cosh}[b*x]*\text{Sinh}[b*x])/(2*b^2) + (2*x*\text{Sinh}[b*x]^2)/b^3 - (6*\text{Cosh}[b*x]*\text{SinhIntegral}[b*x])/b^4$

$$- (3*x^2*\text{Cosh}[b*x]*\text{SinhIntegral}[b*x])/b^2 + (6*x*\text{Sinh}[b*x]*\text{SinhIntegral}[b*x])/b^3 + (x^3*\text{Sinh}[b*x]*\text{SinhIntegral}[b*x])/b + (3*\text{SinhIntegral}[2*b*x])/b^4$$

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 30

`Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]`

Rule 2715

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3379

`Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`

Rule 3392

`Int[((c_) + (d_)*(x_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[d*m*(c + d*x)^(m - 1)*((b*Sin[e + f*x])^n/(f^2*n^2)), x] + (Dist[b^2*((n - 1)/n), Int[(c + d*x)^m*(b*Sin[e + f*x])^(n - 2), x], x] - Dist[d^2*m*((m - 1)/(f^2*n^2)), Int[(c + d*x)^(m - 2)*(b*Sin[e + f*x])^n, x], x] - Simp[b*(c + d*x)^m*Cos[e + f*x]*((b*Sin[e + f*x])^(n - 1)/(f*n)), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 1]`

Rule 5480

`Int[Cosh[(a_) + (b_)*(x_)^(n_)]*(x_)^(m_)*Sinh[(a_) + (b_)*(x_)^(n_)]^(p_), x_Symbol] := Simp[x^(m - n + 1)*(Sinh[a + b*x^n]^(p + 1)/(b*n*(p + 1))), x] - Dist[(m - n + 1)/(b*n*(p + 1)), Int[x^(m - n)*Sinh[a + b*x^n]^(p + 1), x], x] /; FreeQ[{a, b, p}, x] && LtQ[0, n, m + 1] && NeQ[p, -1]`

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

Rule 6675

```
Int[Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
Simp[Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6677

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(SinhIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Sinh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*SinhIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6683

```
Int[Cosh[(a_.) + (b_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.)*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Sinh[a + b*x]*(SinhIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Sinh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*SinhIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x^3 \sinh(bx) \text{Shi}(bx)}{b} - \frac{3 \int x^2 \sinh(bx) \text{Shi}(bx) dx}{b} - \int \frac{x^2 \sinh^2(bx)}{b} dx \\
&= -\frac{3x^2 \cosh(bx) \text{Shi}(bx)}{b^2} + \frac{x^3 \sinh(bx) \text{Shi}(bx)}{b} + \frac{6 \int x \cosh(bx) \text{Shi}(bx) dx}{b^2} \\
&\quad - \frac{\int x^2 \sinh^2(bx) dx}{b} + \frac{3 \int \frac{x \cosh(bx) \sinh(bx)}{b} dx}{b} \\
&= -\frac{x^2 \cosh(bx) \sinh(bx)}{2b^2} + \frac{x \sinh^2(bx)}{2b^3} - \frac{3x^2 \cosh(bx) \text{Shi}(bx)}{b^2} \\
&\quad + \frac{6x \sinh(bx) \text{Shi}(bx)}{b^3} + \frac{x^3 \sinh(bx) \text{Shi}(bx)}{b} - \frac{\int \sinh^2(bx) dx}{2b^3} \\
&\quad - \frac{6 \int \sinh(bx) \text{Shi}(bx) dx}{b^3} + \frac{3 \int x \cosh(bx) \sinh(bx) dx}{b^2} - \frac{6 \int \frac{\sinh^2(bx)}{b} dx}{b^2} + \frac{\int x^2 dx}{2b}
\end{aligned}$$

$$\begin{aligned}
&= \frac{x^3}{6b} - \frac{\cosh(bx) \sinh(bx)}{4b^4} - \frac{x^2 \cosh(bx) \sinh(bx)}{2b^2} + \frac{2x \sinh^2(bx)}{b^3} - \frac{6 \cosh(bx) \text{Shi}(bx)}{b^4} \\
&\quad - \frac{3x^2 \cosh(bx) \text{Shi}(bx)}{b^2} + \frac{6x \sinh(bx) \text{Shi}(bx)}{b^3} + \frac{x^3 \sinh(bx) \text{Shi}(bx)}{b} \\
&\quad + \frac{\int 1 dx}{4b^3} - \frac{3 \int \sinh^2(bx) dx}{2b^3} + \frac{6 \int \frac{\cosh(bx) \sinh(bx)}{bx} dx}{b^3} - \frac{6 \int \sinh^2(bx) dx}{b^3} \\
&= \frac{x}{4b^3} + \frac{x^3}{6b} - \frac{4 \cosh(bx) \sinh(bx)}{b^4} - \frac{x^2 \cosh(bx) \sinh(bx)}{2b^2} + \frac{2x \sinh^2(bx)}{b^3} \\
&\quad - \frac{6 \cosh(bx) \text{Shi}(bx)}{b^4} - \frac{3x^2 \cosh(bx) \text{Shi}(bx)}{b^2} + \frac{6x \sinh(bx) \text{Shi}(bx)}{b^3} \\
&\quad + \frac{x^3 \sinh(bx) \text{Shi}(bx)}{b} + \frac{6 \int \frac{\cosh(bx) \sinh(bx)}{x} dx}{b^4} + \frac{3 \int 1 dx}{4b^3} + \frac{3 \int 1 dx}{b^3} \\
&= \frac{4x}{b^3} + \frac{x^3}{6b} - \frac{4 \cosh(bx) \sinh(bx)}{b^4} - \frac{x^2 \cosh(bx) \sinh(bx)}{2b^2} + \frac{2x \sinh^2(bx)}{b^3} - \frac{6 \cosh(bx) \text{Shi}(bx)}{b^4} \\
&\quad - \frac{3x^2 \cosh(bx) \text{Shi}(bx)}{b^2} + \frac{6x \sinh(bx) \text{Shi}(bx)}{b^3} + \frac{x^3 \sinh(bx) \text{Shi}(bx)}{b} + \frac{6 \int \frac{\sinh(2bx)}{2x} dx}{b^4} \\
&= \frac{4x}{b^3} + \frac{x^3}{6b} - \frac{4 \cosh(bx) \sinh(bx)}{b^4} - \frac{x^2 \cosh(bx) \sinh(bx)}{2b^2} + \frac{2x \sinh^2(bx)}{b^3} - \frac{6 \cosh(bx) \text{Shi}(bx)}{b^4} \\
&\quad - \frac{3x^2 \cosh(bx) \text{Shi}(bx)}{b^2} + \frac{6x \sinh(bx) \text{Shi}(bx)}{b^3} + \frac{x^3 \sinh(bx) \text{Shi}(bx)}{b} + \frac{3 \int \frac{\sinh(2bx)}{x} dx}{b^4} \\
&= \frac{4x}{b^3} + \frac{x^3}{6b} - \frac{4 \cosh(bx) \sinh(bx)}{b^4} - \frac{x^2 \cosh(bx) \sinh(bx)}{2b^2} + \frac{2x \sinh^2(bx)}{b^3} - \frac{6 \cosh(bx) \text{Shi}(bx)}{b^4} \\
&\quad - \frac{3x^2 \cosh(bx) \text{Shi}(bx)}{b^2} + \frac{6x \sinh(bx) \text{Shi}(bx)}{b^3} + \frac{x^3 \sinh(bx) \text{Shi}(bx)}{b} + \frac{3 \text{Shi}(2bx)}{b^4}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.73

$$\begin{aligned}
&\int x^3 \cosh(bx) \text{Shi}(bx) dx \\
&= \frac{36bx + 2b^3x^3 + 12bx \cosh(2bx) - 24 \sinh(2bx) - 3b^2x^2 \sinh(2bx) + 12(-3(2 + b^2x^2) \cosh(bx) + bx(6 + b^2x^2) \text{Shi}(bx))}{12b^4}
\end{aligned}$$

[In] Integrate[x^3*Cosh[b*x]*SinhIntegral[b*x],x]

[Out] (36*b*x + 2*b^3*x^3 + 12*b*x*Cosh[2*b*x] - 24*Sinh[2*b*x] - 3*b^2*x^2*Sinh[2*b*x] + 12*(-3*(2 + b^2*x^2)*Cosh[b*x] + b*x*(6 + b^2*x^2)*Sinh[b*x])*SinhIntegral[b*x] + 36*SinhIntegral[2*b*x])/(12*b^4)

Maple [A] (verified)

Time = 1.34 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.81

method	result
derivativedivides	$\frac{\text{Shi}(bx)(b^3x^3 \sinh(bx) - 3b^2x^2 \cosh(bx) + 6bx \sinh(bx) - 6 \cosh(bx)) - \frac{b^2x^2 \cosh(bx) \sinh(bx)}{2} + \frac{b^3x^3}{6} + 2bx \cosh(bx)^2 - 4 \cosh(bx)}{b^4}$
default	$\frac{\text{Shi}(bx)(b^3x^3 \sinh(bx) - 3b^2x^2 \cosh(bx) + 6bx \sinh(bx) - 6 \cosh(bx)) - \frac{b^2x^2 \cosh(bx) \sinh(bx)}{2} + \frac{b^3x^3}{6} + 2bx \cosh(bx)^2 - 4 \cosh(bx)}{b^4}$

[In] int(x^3*cosh(b*x)*Shi(b*x),x,method=_RETURNVERBOSE)

[Out] 1/b^4*(Shi(b*x)*(b^3*x^3*sinh(b*x)-3*b^2*x^2*cosh(b*x)+6*b*x*sinh(b*x)-6*cosh(b*x))-1/2*b^2*x^2*cosh(b*x)*sinh(b*x)+1/6*b^3*x^3+2*b*x*cosh(b*x)^2-4*cosh(b*x)*sinh(b*x)+2*b*x+3*Shi(2*b*x))

Fricas [F]

$$\int x^3 \cosh(bx) \text{Shi}(bx) dx = \int x^3 \text{Shi}(bx) \cosh(bx) dx$$

[In] integrate(x^3*cosh(b*x)*Shi(b*x),x, algorithm="fricas")

[Out] integral(x^3*cosh(b*x)*sinh_integral(b*x), x)

Sympy [F]

$$\int x^3 \cosh(bx) \text{Shi}(bx) dx = \int x^3 \cosh(bx) \text{Shi}(bx) dx$$

[In] integrate(x**3*cosh(b*x)*Shi(b*x),x)

[Out] Integral(x**3*cosh(b*x)*Shi(b*x), x)

Maxima [F]

$$\int x^3 \cosh(bx) \text{Shi}(bx) dx = \int x^3 \text{Shi}(bx) \cosh(bx) dx$$

[In] integrate(x^3*cosh(b*x)*Shi(b*x),x, algorithm="maxima")

[Out] integrate(x^3*Shi(b*x)*cosh(b*x), x)

Giac [F]

$$\int x^3 \cosh(bx) \operatorname{Shi}(bx) dx = \int x^3 \operatorname{Shi}(bx) \cosh(bx) dx$$

[In] integrate(x^3*cosh(b*x)*Shi(b*x),x, algorithm="giac")

[Out] integrate(x^3*Shi(b*x)*cosh(b*x), x)

Mupad [F(-1)]

Timed out.

$$\int x^3 \cosh(bx) \operatorname{Shi}(bx) dx = \int x^3 \operatorname{sinhint}(bx) \cosh(bx) dx$$

[In] int(x^3*sinhint(b*x)*cosh(b*x),x)

[Out] int(x^3*sinhint(b*x)*cosh(b*x), x)

3.53 $\int \sinh(5x)\mathbf{Shi}(2x) dx$

Optimal result	287
Rubi [A] (verified)	287
Mathematica [A] (verified)	288
Maple [A] (verified)	289
Fricas [F]	289
Sympy [F]	289
Maxima [F]	289
Giac [F]	290
Mupad [F(-1)]	290

Optimal result

Integrand size = 9, antiderivative size = 29

$$\int \sinh(5x)\mathbf{Shi}(2x) dx = \frac{1}{5} \cosh(5x)\mathbf{Shi}(2x) + \frac{\mathbf{Shi}(3x)}{10} - \frac{\mathbf{Shi}(7x)}{10}$$

[Out] 1/5*cosh(5*x)*Shi(2*x)+1/10*Shi(3*x)-1/10*Shi(7*x)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {6675, 12, 5580, 3379}

$$\int \sinh(5x)\mathbf{Shi}(2x) dx = \frac{\mathbf{Shi}(3x)}{10} - \frac{\mathbf{Shi}(7x)}{10} + \frac{1}{5}\mathbf{Shi}(2x) \cosh(5x)$$

[In] Int[Sinh[5*x]*SinhIntegral[2*x],x]

[Out] (Cosh[5*x]*SinhIntegral[2*x])/5 + SinhIntegral[3*x]/10 - SinhIntegral[7*x]/10

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f

, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 5580

```
Int[Cosh[(c_.) + (d_.)*(x_)]^(q_.)*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(p_.), x_Symbol] := Int[ExpandTrigReduce[(e + f*x)^m, Sinh[a +
b*x]^p*Cosh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[p
, 0] && IGtQ[q, 0]
```

Rule 6675

```
Int[Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
Simp[Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{5} \cosh(5x) \text{Shi}(2x) - \frac{2}{5} \int \frac{\cosh(5x) \sinh(2x)}{2x} dx \\
 &= \frac{1}{5} \cosh(5x) \text{Shi}(2x) - \frac{1}{5} \int \frac{\cosh(5x) \sinh(2x)}{x} dx \\
 &= \frac{1}{5} \cosh(5x) \text{Shi}(2x) - \frac{1}{5} \int \left(-\frac{\sinh(3x)}{2x} + \frac{\sinh(7x)}{2x} \right) dx \\
 &= \frac{1}{5} \cosh(5x) \text{Shi}(2x) + \frac{1}{10} \int \frac{\sinh(3x)}{x} dx - \frac{1}{10} \int \frac{\sinh(7x)}{x} dx \\
 &= \frac{1}{5} \cosh(5x) \text{Shi}(2x) + \frac{\text{Shi}(3x)}{10} - \frac{\text{Shi}(7x)}{10}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.86

$$\int \sinh(5x) \text{Shi}(2x) dx = \frac{1}{10} (2 \cosh(5x) \text{Shi}(2x) + \text{Shi}(3x) - \text{Shi}(7x))$$

[In] Integrate[Sinh[5*x]*SinhIntegral[2*x], x]

[Out] (2*Cosh[5*x]*SinhIntegral[2*x] + SinhIntegral[3*x] - SinhIntegral[7*x])/10

Maple [A] (verified)

Time = 1.26 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{\cosh(5x) \operatorname{Shi}(2x)}{5} + \frac{\operatorname{Shi}(3x)}{10} - \frac{\operatorname{Shi}(7x)}{10}$	24

[In] `int(Shi(2*x)*sinh(5*x),x,method=_RETURNVERBOSE)`

[Out] `1/5*cosh(5*x)*Shi(2*x)+1/10*Shi(3*x)-1/10*Shi(7*x)`

Fricas [F]

$$\int \sinh(5x) \operatorname{Shi}(2x) dx = \int \operatorname{Shi}(2x) \sinh(5x) dx$$

[In] `integrate(Shi(2*x)*sinh(5*x),x, algorithm="fricas")`

[Out] `integral(sinh(5*x)*sinh_integral(2*x), x)`

Sympy [F]

$$\int \sinh(5x) \operatorname{Shi}(2x) dx = \int \sinh(5x) \operatorname{Shi}(2x) dx$$

[In] `integrate(Shi(2*x)*sinh(5*x),x)`

[Out] `Integral(sinh(5*x)*Shi(2*x), x)`

Maxima [F]

$$\int \sinh(5x) \operatorname{Shi}(2x) dx = \int \operatorname{Shi}(2x) \sinh(5x) dx$$

[In] `integrate(Shi(2*x)*sinh(5*x),x, algorithm="maxima")`

[Out] `integrate(Shi(2*x)*sinh(5*x), x)`

Giac [F]

$$\int \sinh(5x)\operatorname{Shi}(2x) dx = \int \operatorname{Shi}(2x) \sinh(5x) dx$$

[In] integrate(Shi(2*x)*sinh(5*x),x, algorithm="giac")

[Out] integrate(Shi(2*x)*sinh(5*x), x)

Mupad [F(-1)]

Timed out.

$$\int \sinh(5x)\operatorname{Shi}(2x) dx = \int \operatorname{sinhint}(2x) \sinh(5x) dx$$

[In] int(sinhint(2*x)*sinh(5*x),x)

[Out] int(sinhint(2*x)*sinh(5*x), x)

3.54 $\int \cosh(5x)\text{Shi}(2x) dx$

Optimal result	291
Rubi [A] (verified)	291
Mathematica [A] (verified)	292
Maple [A] (verified)	293
Fricas [F]	293
Sympy [F]	293
Maxima [F]	293
Giac [F]	294
Mupad [F(-1)]	294

Optimal result

Integrand size = 9, antiderivative size = 29

$$\int \cosh(5x)\text{Shi}(2x) dx = \frac{\text{Chi}(3x)}{10} - \frac{\text{Chi}(7x)}{10} + \frac{1}{5} \sinh(5x)\text{Shi}(2x)$$

[Out] 1/10*Chi(3*x)-1/10*Chi(7*x)+1/5*Shi(2*x)*sinh(5*x)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {6681, 12, 5578, 3382}

$$\int \cosh(5x)\text{Shi}(2x) dx = \frac{\text{Chi}(3x)}{10} - \frac{\text{Chi}(7x)}{10} + \frac{1}{5} \text{Shi}(2x) \sinh(5x)$$

[In] Int[Cosh[5*x]*SinhIntegral[2*x],x]

[Out] CoshIntegral[3*x]/10 - CoshIntegral[7*x]/10 + (Sinh[5*x]*SinhIntegral[2*x])/5

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz

```
}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 5578

```
Int[((e_.) + (f_.)*(x_.))^(m_.)*Sinh[(a_.) + (b_.)*(x_.)]^(p_.)*Sinh[(c_.) +
(d_.)*(x_.)]^(q_.), x_Symbol] := Int[ExpandTrigReduce[(e + f*x)^m, Sinh[a +
b*x]^p*Sinh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0]
] && IGtQ[q, 0] && IntegerQ[m]
```

Rule 6681

```
Int[Cosh[(a_.) + (b_.)*(x_.)]*SinhIntegral[(c_.) + (d_.)*(x_.)], x_Symbol] :=
Simp[Sinh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{5} \sinh(5x) \text{Shi}(2x) - \frac{2}{5} \int \frac{\sinh(2x) \sinh(5x)}{2x} dx \\
&= \frac{1}{5} \sinh(5x) \text{Shi}(2x) - \frac{1}{5} \int \frac{\sinh(2x) \sinh(5x)}{x} dx \\
&= \frac{1}{5} \sinh(5x) \text{Shi}(2x) - \frac{1}{5} \int \left(-\frac{\cosh(3x)}{2x} + \frac{\cosh(7x)}{2x} \right) dx \\
&= \frac{1}{5} \sinh(5x) \text{Shi}(2x) + \frac{1}{10} \int \frac{\cosh(3x)}{x} dx - \frac{1}{10} \int \frac{\cosh(7x)}{x} dx \\
&= \frac{\text{Chi}(3x)}{10} - \frac{\text{Chi}(7x)}{10} + \frac{1}{5} \sinh(5x) \text{Shi}(2x)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.86

$$\int \cosh(5x) \text{Shi}(2x) dx = \frac{1}{10} (\text{Chi}(3x) - \text{Chi}(7x) + 2 \sinh(5x) \text{Shi}(2x))$$

```
[In] Integrate[Cosh[5*x]*SinhIntegral[2*x], x]
```

```
[Out] (CoshIntegral[3*x] - CoshIntegral[7*x] + 2*Sinh[5*x]*SinhIntegral[2*x])/10
```

Maple [A] (verified)

Time = 1.35 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{\text{Chi}(3x)}{10} - \frac{\text{Chi}(7x)}{10} + \frac{\text{Shi}(2x)\sinh(5x)}{5}$	24

[In] `int(cosh(5*x)*Shi(2*x),x,method=_RETURNVERBOSE)`

[Out] `1/10*Chi(3*x)-1/10*Chi(7*x)+1/5*Shi(2*x)*sinh(5*x)`

Fricas [F]

$$\int \cosh(5x)\text{Shi}(2x) dx = \int \text{Shi}(2x) \cosh(5x) dx$$

[In] `integrate(cosh(5*x)*Shi(2*x),x, algorithm="fricas")`

[Out] `integral(cosh(5*x)*sinh_integral(2*x), x)`

Sympy [F]

$$\int \cosh(5x)\text{Shi}(2x) dx = \int \cosh(5x) \text{Shi}(2x) dx$$

[In] `integrate(cosh(5*x)*Shi(2*x),x)`

[Out] `Integral(cosh(5*x)*Shi(2*x), x)`

Maxima [F]

$$\int \cosh(5x)\text{Shi}(2x) dx = \int \text{Shi}(2x) \cosh(5x) dx$$

[In] `integrate(cosh(5*x)*Shi(2*x),x, algorithm="maxima")`

[Out] `integrate(Shi(2*x)*cosh(5*x), x)`

Giac [F]

$$\int \cosh(5x)\operatorname{Shi}(2x) dx = \int \operatorname{Shi}(2x) \cosh(5x) dx$$

[In] integrate(cosh(5*x)*Shi(2*x),x, algorithm="giac")

[Out] integrate(Shi(2*x)*cosh(5*x), x)

Mupad [F(-1)]

Timed out.

$$\int \cosh(5x)\operatorname{Shi}(2x) dx = \int \operatorname{sinhint}(2x) \cosh(5x) dx$$

[In] int(sinhint(2*x)*cosh(5*x),x)

[Out] int(sinhint(2*x)*cosh(5*x), x)

3.55 $\int x^2 \sinh(a + bx) \mathbf{Shi}(a + bx) dx$

Optimal result	295
Rubi [A] (verified)	296
Mathematica [A] (verified)	299
Maple [A] (verified)	300
Fricas [F]	300
Sympy [F]	300
Maxima [F]	301
Giac [F]	301
Mupad [F(-1)]	301

Optimal result

Integrand size = 16, antiderivative size = 186

$$\int x^2 \sinh(a + bx) \mathbf{Shi}(a + bx) dx = -\frac{x}{b^2} + \frac{a \cosh(2a + 2bx)}{4b^3} - \frac{x \cosh(2a + 2bx)}{4b^2}$$

$$- \frac{a \mathbf{Chi}(2a + 2bx)}{b^3} + \frac{a \log(a + bx)}{b^3}$$

$$+ \frac{\cosh(a + bx) \sinh(a + bx)}{b^3} + \frac{\sinh(2a + 2bx)}{8b^3}$$

$$+ \frac{2 \cosh(a + bx) \mathbf{Shi}(a + bx)}{b^3}$$

$$+ \frac{x^2 \cosh(a + bx) \mathbf{Shi}(a + bx)}{b}$$

$$- \frac{2x \sinh(a + bx) \mathbf{Shi}(a + bx)}{b^2}$$

$$- \frac{\mathbf{Shi}(2a + 2bx)}{b^3} - \frac{a^2 \mathbf{Shi}(2a + 2bx)}{2b^3}$$

```
[Out] -x/b^2-a*Chi(2*b*x+2*a)/b^3+1/4*a*cosh(2*b*x+2*a)/b^3-1/4*x*cosh(2*b*x+2*a)
/b^2+a*ln(b*x+a)/b^3+2*cosh(b*x+a)*Shi(b*x+a)/b^3+x^2*cosh(b*x+a)*Shi(b*x+a)
)/b-Shi(2*b*x+2*a)/b^3-1/2*a^2*Shi(2*b*x+2*a)/b^3+cosh(b*x+a)*sinh(b*x+a)/b
^3-2*x*Shi(b*x+a)*sinh(b*x+a)/b^2+1/8*sinh(2*b*x+2*a)/b^3
```

Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.00, number of steps used = 21, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {6677, 5736, 6873, 6874, 2718, 3377, 2717, 3379, 6683, 2715, 8, 3393, 3382, 6675, 5556, 12}

$$\int x^2 \sinh(a + bx) \text{Shi}(a + bx) dx = -\frac{a^2 \text{Shi}(2a + 2bx)}{2b^3} - \frac{a \text{Chi}(2a + 2bx)}{b^3} - \frac{\text{Shi}(2a + 2bx)}{b^3} + \frac{2 \text{Shi}(a + bx) \cosh(a + bx)}{b^3} + \frac{a \log(a + bx)}{b^3} + \frac{\sinh(2a + 2bx)}{8b^3} + \frac{a \cosh(2a + 2bx)}{4b^3} + \frac{\sinh(a + bx) \cosh(a + bx)}{b^3} - \frac{2x \text{Shi}(a + bx) \sinh(a + bx)}{b^2} - \frac{x \cosh(2a + 2bx)}{4b^2} + \frac{x^2 \text{Shi}(a + bx) \cosh(a + bx)}{b} - \frac{x}{b^2}$$

[In] Int[x^2*Sinh[a + b*x]*SinhIntegral[a + b*x],x]

[Out] -(x/b^2) + (a*Cosh[2*a + 2*b*x])/(4*b^3) - (x*Cosh[2*a + 2*b*x])/(4*b^2) - (a*CoshIntegral[2*a + 2*b*x])/b^3 + (a*Log[a + b*x])/b^3 + (Cosh[a + b*x]*SinhIntegral[a + b*x])/b^3 + Sinh[2*a + 2*b*x]/(8*b^3) + (2*Cosh[a + b*x]*SinhIntegral[a + b*x])/b^3 + (x^2*Cosh[a + b*x]*SinhIntegral[a + b*x])/b - (2*x*Sinh[a + b*x]*SinhIntegral[a + b*x])/b^2 - SinhIntegral[2*a + 2*b*x]/b^3 - (a^2*SinhIntegral[2*a + 2*b*x])/(2*b^3)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n-1)/(d*n)), x] + Dist[b^2*((n-1)/n), Int[(b*Sin[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2717

Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]

Rule 2718

Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ
[{c, d}, x]

Rule 3377

Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(
-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Co
s[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f
, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz
}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3393

Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := In
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))

Rule 5556

Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]

Rule 5736

Int[Cosh[w_]^(p_.)*(u_.)*Sinh[v_]^(p_.), x_Symbol] := Dist[1/2^p, Int[u*Sin
h[2*v]^p, x], x] /; EqQ[w, v] && IntegerQ[p]

Rule 6675

```
Int[Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
  Simp[Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
  b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6677

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(SinhIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Sinh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*SinhIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6683

```
Int[Cosh[(a_.) + (b_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.)*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Sinh[a + b*x]*(SinhIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Sinh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*SinhIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6873

```
Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v !=
= u]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x^2 \cosh(a + bx) \text{Shi}(a + bx)}{b} - \frac{2 \int x \cosh(a + bx) \text{Shi}(a + bx) dx}{b} \\
&\quad - \int \frac{x^2 \cosh(a + bx) \sinh(a + bx)}{a + bx} dx \\
&= \frac{x^2 \cosh(a + bx) \text{Shi}(a + bx)}{b} - \frac{2x \sinh(a + bx) \text{Shi}(a + bx)}{b^2} \\
&\quad - \frac{1}{2} \int \frac{x^2 \sinh(2(a + bx))}{a + bx} dx + \frac{2 \int \sinh(a + bx) \text{Shi}(a + bx) dx}{b^2} + \frac{2 \int \frac{x \sinh^2(a + bx)}{a + bx} dx}{b} \\
&= \frac{2 \cosh(a + bx) \text{Shi}(a + bx)}{b^3} + \frac{x^2 \cosh(a + bx) \text{Shi}(a + bx)}{b} - \frac{2x \sinh(a + bx) \text{Shi}(a + bx)}{b^2} \\
&\quad - \frac{1}{2} \int \frac{x^2 \sinh(2a + 2bx)}{a + bx} dx - \frac{2 \int \frac{\cosh(a + bx) \sinh(a + bx)}{a + bx} dx}{b^2} + \frac{2 \int \left(\frac{\sinh^2(a + bx)}{b} - \frac{a \sinh^2(a + bx)}{b(a + bx)} \right) dx}{b}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2 \cosh(a + bx) \operatorname{Shi}(a + bx)}{b^3} + \frac{x^2 \cosh(a + bx) \operatorname{Shi}(a + bx)}{b} \\
&\quad - \frac{2x \sinh(a + bx) \operatorname{Shi}(a + bx)}{b^2} \\
&\quad - \frac{1}{2} \int \left(-\frac{a \sinh(2a + 2bx)}{b^2} + \frac{x \sinh(2a + 2bx)}{b} + \frac{a^2 \sinh(2a + 2bx)}{b^2(a + bx)} \right) dx \\
&\quad + \frac{2 \int \sinh^2(a + bx) dx}{b^2} - \frac{2 \int \frac{\sinh(2a+2bx)}{2(a+bx)} dx}{b^2} - \frac{(2a) \int \frac{\sinh^2(a+bx)}{a+bx} dx}{b^2} \\
&= \frac{\cosh(a + bx) \sinh(a + bx)}{b^3} + \frac{2 \cosh(a + bx) \operatorname{Shi}(a + bx)}{b^3} + \frac{x^2 \cosh(a + bx) \operatorname{Shi}(a + bx)}{b} \\
&\quad - \frac{2x \sinh(a + bx) \operatorname{Shi}(a + bx)}{b^2} - \frac{\int 1 dx}{b^2} - \frac{\int \frac{\sinh(2a+2bx)}{a+bx} dx}{b^2} + \frac{a \int \sinh(2a + 2bx) dx}{2b^2} \\
&\quad + \frac{(2a) \int \left(\frac{1}{2(a+bx)} - \frac{\cosh(2a+2bx)}{2(a+bx)} \right) dx}{b^2} - \frac{a^2 \int \frac{\sinh(2a+2bx)}{a+bx} dx}{2b^2} - \frac{\int x \sinh(2a + 2bx) dx}{2b} \\
&= -\frac{x}{b^2} + \frac{a \cosh(2a + 2bx)}{4b^3} - \frac{x \cosh(2a + 2bx)}{4b^2} + \frac{a \log(a + bx)}{b^3} + \frac{\cosh(a + bx) \sinh(a + bx)}{b^3} \\
&\quad + \frac{2 \cosh(a + bx) \operatorname{Shi}(a + bx)}{b^3} + \frac{x^2 \cosh(a + bx) \operatorname{Shi}(a + bx)}{b} - \frac{2x \sinh(a + bx) \operatorname{Shi}(a + bx)}{b^2} \\
&\quad - \frac{\operatorname{Shi}(2a + 2bx)}{b^3} - \frac{a^2 \operatorname{Shi}(2a + 2bx)}{2b^3} + \frac{\int \cosh(2a + 2bx) dx}{4b^2} - \frac{a \int \frac{\cosh(2a+2bx)}{a+bx} dx}{b^2} \\
&= -\frac{x}{b^2} + \frac{a \cosh(2a + 2bx)}{4b^3} - \frac{x \cosh(2a + 2bx)}{4b^2} - \frac{a \operatorname{Chi}(2a + 2bx)}{b^3} \\
&\quad + \frac{a \log(a + bx)}{b^3} + \frac{\cosh(a + bx) \sinh(a + bx)}{b^3} + \frac{\sinh(2a + 2bx)}{8b^3} \\
&\quad + \frac{2 \cosh(a + bx) \operatorname{Shi}(a + bx)}{b^3} + \frac{x^2 \cosh(a + bx) \operatorname{Shi}(a + bx)}{b} \\
&\quad - \frac{2x \sinh(a + bx) \operatorname{Shi}(a + bx)}{b^2} - \frac{\operatorname{Shi}(2a + 2bx)}{b^3} - \frac{a^2 \operatorname{Shi}(2a + 2bx)}{2b^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.66

$$\int x^2 \sinh(a + bx) \operatorname{Shi}(a + bx) dx$$

$$= \frac{-8bx + 2a \cosh(2(a + bx)) - 2bx \cosh(2(a + bx)) - 8a \operatorname{Chi}(2(a + bx)) + 8a \log(a + bx) + 5 \sinh(2(a + bx))}{8b^3}$$

[In] Integrate[x^2*Sinh[a + b*x]*SinhIntegral[a + b*x],x]

[Out] (-8*b*x + 2*a*Cosh[2*(a + b*x)] - 2*b*x*Cosh[2*(a + b*x)] - 8*a*CoshIntegral[2*(a + b*x)] + 8*a*Log[a + b*x] + 5*Sinh[2*(a + b*x)] + 8*((2 + b^2*x^2)*Cosh[a + b*x] - 2*b*x*Sinh[a + b*x])*SinhIntegral[a + b*x] - 8*SinhIntegral[2*(a + b*x)] - 4*a^2*SinhIntegral[2*(a + b*x)])/(8*b^3)

Maple [A] (verified)

Time = 1.62 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.94

method	result
derivativedivides	$\frac{\text{Shi}(bx+a) \left(a^2 \cosh(bx+a) - 2a((bx+a) \cosh(bx+a) - \sinh(bx+a)) + (bx+a)^2 \cosh(bx+a) - 2(bx+a) \sinh(bx+a) + 2 \cosh(bx+a) \right)}{\dots}$
default	$\text{Shi}(bx+a) \left(a^2 \cosh(bx+a) - 2a((bx+a) \cosh(bx+a) - \sinh(bx+a)) + (bx+a)^2 \cosh(bx+a) - 2(bx+a) \sinh(bx+a) + 2 \cosh(bx+a) \right)$

```
[In] int(x^2*Shi(b*x+a)*sinh(b*x+a),x,method=_RETURNVERBOSE)
```

```
[Out] 1/b^3*(Shi(b*x+a)*(a^2*cosh(b*x+a)-2*a*((b*x+a)*cosh(b*x+a)-sinh(b*x+a)))+(b*x+a)^2*cosh(b*x+a)-2*(b*x+a)*sinh(b*x+a)+2*cosh(b*x+a))-1/2*a^2*Shi(2*b*x+2*a)+a*cosh(b*x+a)^2+a*ln(b*x+a)-a*Chi(2*b*x+2*a)-1/2*(b*x+a)*cosh(b*x+a)^2+5/4*cosh(b*x+a)*sinh(b*x+a)-3/4*b*x-3/4*a-Shi(2*b*x+2*a))
```

Fricas [F]

$$\int x^2 \sinh(a + bx) \text{Shi}(a + bx) dx = \int x^2 \text{Shi}(bx + a) \sinh(bx + a) dx$$

```
[In] integrate(x^2*Shi(b*x+a)*sinh(b*x+a),x, algorithm="fricas")
```

```
[Out] integral(x^2*sinh(b*x + a)*sinh_integral(b*x + a), x)
```

Sympy [F]

$$\int x^2 \sinh(a + bx) \text{Shi}(a + bx) dx = \int x^2 \sinh(a + bx) \text{Shi}(a + bx) dx$$

```
[In] integrate(x**2*Shi(b*x+a)*sinh(b*x+a),x)
```

```
[Out] Integral(x**2*sinh(a + b*x)*Shi(a + b*x), x)
```

Maxima [F]

$$\int x^2 \sinh(a + bx) \operatorname{Shi}(a + bx) dx = \int x^2 \operatorname{Shi}(bx + a) \sinh(bx + a) dx$$

[In] integrate(x^2*Shi(b*x+a)*sinh(b*x+a),x, algorithm="maxima")

[Out] integrate(x^2*Shi(b*x + a)*sinh(b*x + a), x)

Giac [F]

$$\int x^2 \sinh(a + bx) \operatorname{Shi}(a + bx) dx = \int x^2 \operatorname{Shi}(bx + a) \sinh(bx + a) dx$$

[In] integrate(x^2*Shi(b*x+a)*sinh(b*x+a),x, algorithm="giac")

[Out] integrate(x^2*Shi(b*x + a)*sinh(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \sinh(a + bx) \operatorname{Shi}(a + bx) dx = \int x^2 \operatorname{sinhint}(a + bx) \sinh(a + bx) dx$$

[In] int(x^2*sinhint(a + b*x)*sinh(a + b*x),x)

[Out] int(x^2*sinhint(a + b*x)*sinh(a + b*x), x)

3.56 $\int x \sinh(a + bx) \text{Shi}(a + bx) dx$

Optimal result	302
Rubi [A] (verified)	302
Mathematica [A] (verified)	305
Maple [A] (verified)	305
Fricas [F]	305
Sympy [F]	306
Maxima [F]	306
Giac [F]	306
Mupad [F(-1)]	306

Optimal result

Integrand size = 14, antiderivative size = 97

$$\int x \sinh(a + bx) \text{Shi}(a + bx) dx = -\frac{\cosh(2a + 2bx)}{4b^2} + \frac{\text{Chi}(2a + 2bx)}{2b^2} - \frac{\log(a + bx)}{2b^2} + \frac{x \cosh(a + bx) \text{Shi}(a + bx)}{b} - \frac{\sinh(a + bx) \text{Shi}(a + bx)}{b^2} + \frac{a \text{Shi}(2a + 2bx)}{2b^2}$$

[Out] $1/2*\text{Chi}(2*b*x+2*a)/b^2-1/4*\cosh(2*b*x+2*a)/b^2-1/2*\ln(b*x+a)/b^2+x*\cosh(b*x+a)*\text{Shi}(b*x+a)/b+1/2*a*\text{Shi}(2*b*x+2*a)/b^2-\text{Shi}(b*x+a)*\sinh(b*x+a)/b^2$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {6677, 5736, 6873, 6874, 2718, 3379, 6681, 3393, 3382}

$$\int x \sinh(a + bx) \text{Shi}(a + bx) dx = \frac{\text{Chi}(2a + 2bx)}{2b^2} + \frac{a \text{Shi}(2a + 2bx)}{2b^2} - \frac{\text{Shi}(a + bx) \sinh(a + bx)}{b^2} - \frac{\log(a + bx)}{2b^2} - \frac{\cosh(2a + 2bx)}{4b^2} + \frac{x \text{Shi}(a + bx) \cosh(a + bx)}{b}$$

[In] $\text{Int}[x*\text{Sinh}[a + b*x]*\text{SinhIntegral}[a + b*x], x]$

[Out] $-1/4*\text{Cosh}[2*a + 2*b*x]/b^2 + \text{CoshIntegral}[2*a + 2*b*x]/(2*b^2) - \text{Log}[a + b*x]/(2*b^2) + (x*\text{Cosh}[a + b*x]*\text{SinhIntegral}[a + b*x])/b - (\text{Sinh}[a + b*x]*\text{SinhIntegral}[a + b*x])/b^2 + (a*\text{SinhIntegral}[2*a + 2*b*x])/(2*b^2)$

Rule 2718

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ
[{c, d}, x]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3393

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 5736

```
Int[Cosh[w_]^(p_.)*(u_.)*Sinh[v_]^(p_.), x_Symbol] := Dist[1/2^p, Int[u*Sinh
h[2*v]^p, x], x] /; EqQ[w, v] && IntegerQ[p]
```

Rule 6677

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(SinhIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Sinh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*SinhIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6681

```
Int[Cosh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
Simp[Sinh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6873

```
Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v !=
= u]
```

Rule 6874

`Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]`
`]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x \cosh(a + bx) \text{Shi}(a + bx)}{b} - \frac{\int \cosh(a + bx) \text{Shi}(a + bx) dx}{b} \\
 &\quad - \int \frac{x \cosh(a + bx) \sinh(a + bx)}{a + bx} dx \\
 &= \frac{x \cosh(a + bx) \text{Shi}(a + bx)}{b} - \frac{\sinh(a + bx) \text{Shi}(a + bx)}{b^2} \\
 &\quad - \frac{1}{2} \int \frac{x \sinh(2(a + bx))}{a + bx} dx + \frac{\int \frac{\sinh^2(a + bx)}{a + bx} dx}{b} \\
 &= \frac{x \cosh(a + bx) \text{Shi}(a + bx)}{b} - \frac{\sinh(a + bx) \text{Shi}(a + bx)}{b^2} \\
 &\quad - \frac{1}{2} \int \frac{x \sinh(2a + 2bx)}{a + bx} dx - \frac{\int \left(\frac{1}{2(a + bx)} - \frac{\cosh(2a + 2bx)}{2(a + bx)} \right) dx}{b} \\
 &= -\frac{\log(a + bx)}{2b^2} + \frac{x \cosh(a + bx) \text{Shi}(a + bx)}{b} - \frac{\sinh(a + bx) \text{Shi}(a + bx)}{b^2} \\
 &\quad - \frac{1}{2} \int \left(\frac{\sinh(2a + 2bx)}{b} + \frac{a \sinh(2a + 2bx)}{b(-a - bx)} \right) dx + \frac{\int \frac{\cosh(2a + 2bx)}{a + bx} dx}{2b} \\
 &= \frac{\text{Chi}(2a + 2bx)}{2b^2} - \frac{\log(a + bx)}{2b^2} + \frac{x \cosh(a + bx) \text{Shi}(a + bx)}{b} \\
 &\quad - \frac{\sinh(a + bx) \text{Shi}(a + bx)}{b^2} - \frac{\int \sinh(2a + 2bx) dx}{2b} - \frac{a \int \frac{\sinh(2a + 2bx)}{-a - bx} dx}{2b} \\
 &= -\frac{\cosh(2a + 2bx)}{4b^2} + \frac{\text{Chi}(2a + 2bx)}{2b^2} - \frac{\log(a + bx)}{2b^2} \\
 &\quad + \frac{x \cosh(a + bx) \text{Shi}(a + bx)}{b} - \frac{\sinh(a + bx) \text{Shi}(a + bx)}{b^2} + \frac{a \text{Shi}(2a + 2bx)}{2b^2}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.75

$$\int x \sinh(a + bx) \operatorname{Shi}(a + bx) dx = \frac{-\cosh(2(a + bx)) + 2\operatorname{Chi}(2(a + bx)) - 2\log(a + bx) + 4(bx \cosh(a + bx) - \sinh(a + bx))\operatorname{Shi}(a + bx) + 2a \operatorname{Shi}(a + bx)}{4b^2}$$

[In] Integrate[x*Sinh[a + b*x]*SinhIntegral[a + b*x],x]

[Out] (-Cosh[2*(a + b*x)] + 2*CoshIntegral[2*(a + b*x)] - 2*Log[a + b*x] + 4*(b*x *Cosh[a + b*x] - Sinh[a + b*x])*SinhIntegral[a + b*x] + 2*a*SinhIntegral[2*(a + b*x)])/(4*b^2)

Maple [A] (verified)

Time = 1.05 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.87

method	result
derivativedivides	$\frac{\operatorname{Shi}(bx+a)(-a \cosh(bx+a) + (bx+a) \cosh(bx+a) - \sinh(bx+a)) + \frac{a \operatorname{Shi}(2bx+2a)}{2} - \frac{\cosh(bx+a)^2}{2} - \frac{\ln(bx+a)}{2} + \frac{\operatorname{Chi}(2bx+2a)}{2}}{b^2}$
default	$\frac{\operatorname{Shi}(bx+a)(-a \cosh(bx+a) + (bx+a) \cosh(bx+a) - \sinh(bx+a)) + \frac{a \operatorname{Shi}(2bx+2a)}{2} - \frac{\cosh(bx+a)^2}{2} - \frac{\ln(bx+a)}{2} + \frac{\operatorname{Chi}(2bx+2a)}{2}}{b^2}$

[In] int(x*Shi(b*x+a)*sinh(b*x+a),x,method=_RETURNVERBOSE)

[Out] 1/b^2*(Shi(b*x+a)*(-a*cosh(b*x+a)+(b*x+a)*cosh(b*x+a)-sinh(b*x+a))+1/2*a*Shi(2*b*x+2*a)-1/2*cosh(b*x+a)^2-1/2*ln(b*x+a)+1/2*Chi(2*b*x+2*a))

Fricas [F]

$$\int x \sinh(a + bx) \operatorname{Shi}(a + bx) dx = \int x \operatorname{Shi}(bx + a) \sinh(bx + a) dx$$

[In] integrate(x*Shi(b*x+a)*sinh(b*x+a),x, algorithm="fricas")

[Out] integral(x*sinh(b*x + a)*sinh_integral(b*x + a), x)

Sympy [F]

$$\int x \sinh(a + bx) \operatorname{Shi}(a + bx) dx = \int x \sinh(a + bx) \operatorname{Shi}(a + bx) dx$$

[In] `integrate(x*Shi(b*x+a)*sinh(b*x+a),x)`

[Out] `Integral(x*sinh(a + b*x)*Shi(a + b*x), x)`

Maxima [F]

$$\int x \sinh(a + bx) \operatorname{Shi}(a + bx) dx = \int x \operatorname{Shi}(bx + a) \sinh(bx + a) dx$$

[In] `integrate(x*Shi(b*x+a)*sinh(b*x+a),x, algorithm="maxima")`

[Out] `integrate(x*Shi(b*x + a)*sinh(b*x + a), x)`

Giac [F]

$$\int x \sinh(a + bx) \operatorname{Shi}(a + bx) dx = \int x \operatorname{Shi}(bx + a) \sinh(bx + a) dx$$

[In] `integrate(x*Shi(b*x+a)*sinh(b*x+a),x, algorithm="giac")`

[Out] `integrate(x*Shi(b*x + a)*sinh(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x \sinh(a + bx) \operatorname{Shi}(a + bx) dx = \int x \operatorname{sinhint}(a + bx) \sinh(a + bx) dx$$

[In] `int(x*sinhint(a + b*x)*sinh(a + b*x),x)`

[Out] `int(x*sinhint(a + b*x)*sinh(a + b*x), x)`

3.57 $\int \sinh(a + bx)\mathbf{Shi}(a + bx) dx$

Optimal result	307
Rubi [A] (verified)	307
Mathematica [A] (verified)	308
Maple [A] (verified)	308
Fricas [F]	309
Sympy [F]	309
Maxima [F]	309
Giac [F]	310
Mupad [F(-1)]	310

Optimal result

Integrand size = 13, antiderivative size = 33

$$\int \sinh(a + bx)\mathbf{Shi}(a + bx) dx = \frac{\cosh(a + bx)\mathbf{Shi}(a + bx)}{b} - \frac{\mathbf{Shi}(2a + 2bx)}{2b}$$

[Out] $\cosh(b*x+a)*\mathbf{Shi}(b*x+a)/b-1/2*\mathbf{Shi}(2*b*x+2*a)/b$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {6675, 5556, 12, 3379}

$$\int \sinh(a + bx)\mathbf{Shi}(a + bx) dx = \frac{\mathbf{Shi}(a + bx) \cosh(a + bx)}{b} - \frac{\mathbf{Shi}(2a + 2bx)}{2b}$$

[In] $\text{Int}[\text{Sinh}[a + b*x]*\text{SinhIntegral}[a + b*x], x]$

[Out] $(\text{Cosh}[a + b*x]*\text{SinhIntegral}[a + b*x])/b - \text{SinhIntegral}[2*a + 2*b*x]/(2*b)$

Rule 12

$\text{Int}[(a_*)(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \&\& \text{!MatchQ}[u, (b_*)(v_)] /; \text{FreeQ}[b, x]$

Rule 3379

$\text{Int}[\sin[(e_*) + (\text{Complex}[0, fz_])*(f_*)(x_)]/((c_*) + (d_*)(x_)), x_Symbol] \rightarrow \text{Simp}[I*(\text{SinhIntegral}[c*f*(fz/d) + f*fz*x]/d), x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \&\& \text{EqQ}[d*e - c*f*fz*I, 0]$

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

Rule 6675

```
Int[Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
Simp[Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\cosh(a + bx)\text{Shi}(a + bx)}{b} - \int \frac{\cosh(a + bx)\sinh(a + bx)}{a + bx} dx \\
&= \frac{\cosh(a + bx)\text{Shi}(a + bx)}{b} - \int \frac{\sinh(2a + 2bx)}{2(a + bx)} dx \\
&= \frac{\cosh(a + bx)\text{Shi}(a + bx)}{b} - \frac{1}{2} \int \frac{\sinh(2a + 2bx)}{a + bx} dx \\
&= \frac{\cosh(a + bx)\text{Shi}(a + bx)}{b} - \frac{\text{Shi}(2a + 2bx)}{2b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.97

$$\int \sinh(a + bx)\text{Shi}(a + bx) dx = \frac{\cosh(a + bx)\text{Shi}(a + bx)}{b} - \frac{\text{Shi}(2(a + bx))}{2b}$$

```
[In] Integrate[Sinh[a + b*x]*SinhIntegral[a + b*x],x]
```

```
[Out] (Cosh[a + b*x]*SinhIntegral[a + b*x])/b - SinhIntegral[2*(a + b*x)]/(2*b)
```

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.91

method	result	size
derivativedivides	$\frac{\cosh(bx+a) \operatorname{Shi}(bx+a) - \frac{\operatorname{Shi}(2bx+2a)}{2}}{b}$	30
default	$\frac{\cosh(bx+a) \operatorname{Shi}(bx+a) - \frac{\operatorname{Shi}(2bx+2a)}{2}}{b}$	30

```
[In] int(Shi(b*x+a)*sinh(b*x+a),x,method=_RETURNVERBOSE)
```

```
[Out] 1/b*(cosh(b*x+a)*Shi(b*x+a)-1/2*Shi(2*b*x+2*a))
```

Fricas [F]

$$\int \sinh(a + bx) \operatorname{Shi}(a + bx) dx = \int \operatorname{Shi}(bx + a) \sinh(bx + a) dx$$

```
[In] integrate(Shi(b*x+a)*sinh(b*x+a),x, algorithm="fricas")
```

```
[Out] integral(sinh(b*x + a)*sinh_integral(b*x + a), x)
```

Sympy [F]

$$\int \sinh(a + bx) \operatorname{Shi}(a + bx) dx = \int \sinh(a + bx) \operatorname{Shi}(a + bx) dx$$

```
[In] integrate(Shi(b*x+a)*sinh(b*x+a),x)
```

```
[Out] Integral(sinh(a + b*x)*Shi(a + b*x), x)
```

Maxima [F]

$$\int \sinh(a + bx) \operatorname{Shi}(a + bx) dx = \int \operatorname{Shi}(bx + a) \sinh(bx + a) dx$$

```
[In] integrate(Shi(b*x+a)*sinh(b*x+a),x, algorithm="maxima")
```

```
[Out] integrate(Shi(b*x + a)*sinh(b*x + a), x)
```

Giac [F]

$$\int \sinh(a + bx)\text{Shi}(a + bx) dx = \int \text{Shi}(bx + a) \sinh(bx + a) dx$$

[In] integrate(Shi(b*x+a)*sinh(b*x+a),x, algorithm="giac")

[Out] integrate(Shi(b*x + a)*sinh(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int \sinh(a + bx)\text{Shi}(a + bx) dx = \int \sinhint(a + bx) \sinh(a + bx) dx$$

[In] int(sinhint(a + b*x)*sinh(a + b*x),x)

[Out] int(sinhint(a + b*x)*sinh(a + b*x), x)

3.58 $\int \frac{\sinh(a+bx)\mathbf{Shi}(a+bx)}{x} dx$

Optimal result	311
Rubi [N/A]	311
Mathematica [N/A]	312
Maple [N/A] (verified)	312
Fricas [N/A]	312
Sympy [N/A]	312
Maxima [N/A]	313
Giac [N/A]	313
Mupad [N/A]	313

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\sinh(a+bx)\mathbf{Shi}(a+bx)}{x} dx = \text{Int}\left(\frac{\sinh(a+bx)\mathbf{Shi}(a+bx)}{x}, x\right)$$

[Out] CannotIntegrate(Shi(b*x+a)*sinh(b*x+a)/x,x)

Rubi [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\sinh(a+bx)\mathbf{Shi}(a+bx)}{x} dx = \int \frac{\sinh(a+bx)\mathbf{Shi}(a+bx)}{x} dx$$

[In] Int[(Sinh[a + b*x]*SinhIntegral[a + b*x])/x,x]

[Out] Defer[Int] [(Sinh[a + b*x]*SinhIntegral[a + b*x])/x, x]

Rubi steps

$$\text{integral} = \int \frac{\sinh(a+bx)\mathbf{Shi}(a+bx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 0.41 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sinh(a + bx)\text{Shi}(a + bx)}{x} dx = \int \frac{\sinh(a + bx)\text{Shi}(a + bx)}{x} dx$$

[In] Integrate[(Sinh[a + b*x]*SinhIntegral[a + b*x])/x,x]

[Out] Integrate[(Sinh[a + b*x]*SinhIntegral[a + b*x])/x, x]

Maple [N/A] (verified)

Not integrable

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\text{Shi}(bx + a)\sinh(bx + a)}{x} dx$$

[In] int(Shi(b*x+a)*sinh(b*x+a)/x,x)

[Out] int(Shi(b*x+a)*sinh(b*x+a)/x,x)

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sinh(a + bx)\text{Shi}(a + bx)}{x} dx = \int \frac{\text{Shi}(bx + a)\sinh(bx + a)}{x} dx$$

[In] integrate(Shi(b*x+a)*sinh(b*x+a)/x,x, algorithm="fricas")

[Out] integral(sinh(b*x + a)*sinh_integral(b*x + a)/x, x)

Sympy [N/A]

Not integrable

Time = 1.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{\sinh(a + bx)\text{Shi}(a + bx)}{x} dx = \int \frac{\sinh(a + bx)\text{Shi}(a + bx)}{x} dx$$

[In] integrate(Shi(b*x+a)*sinh(b*x+a)/x,x)

[Out] Integral(sinh(a + b*x)*Shi(a + b*x)/x, x)

Maxima [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sinh(a + bx)\text{Shi}(a + bx)}{x} dx = \int \frac{\text{Shi}(bx + a) \sinh(bx + a)}{x} dx$$

[In] integrate(Shi(b*x+a)*sinh(b*x+a)/x,x, algorithm="maxima")

[Out] integrate(Shi(b*x + a)*sinh(b*x + a)/x, x)

Giac [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sinh(a + bx)\text{Shi}(a + bx)}{x} dx = \int \frac{\text{Shi}(bx + a) \sinh(bx + a)}{x} dx$$

[In] integrate(Shi(b*x+a)*sinh(b*x+a)/x,x, algorithm="giac")

[Out] integrate(Shi(b*x + a)*sinh(b*x + a)/x, x)

Mupad [N/A]

Not integrable

Time = 5.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sinh(a + bx)\text{Shi}(a + bx)}{x} dx = \int \frac{\sinhint(a + bx) \sinh(a + bx)}{x} dx$$

[In] int((sinhint(a + b*x)*sinh(a + b*x))/x,x)

[Out] int((sinhint(a + b*x)*sinh(a + b*x))/x, x)

3.59 $\int x^2 \cosh(a + bx) \mathbf{Shi}(a + bx) dx$

Optimal result	314
Rubi [A] (verified)	315
Mathematica [A] (verified)	319
Maple [A] (verified)	319
Fricas [F]	319
Sympy [F]	320
Maxima [F]	320
Giac [F]	320
Mupad [F(-1)]	320

Optimal result

Integrand size = 16, antiderivative size = 219

$$\begin{aligned}
 \int x^2 \cosh(a + bx) \mathbf{Shi}(a + bx) dx = & -\frac{ax}{2b^2} + \frac{x^2}{4b} + \frac{\cosh(2a + 2bx)}{2b^3} - \frac{\mathbf{Chi}(2a + 2bx)}{b^3} \\
 & - \frac{a^2 \mathbf{Chi}(2a + 2bx)}{2b^3} + \frac{\log(a + bx)}{b^3} \\
 & + \frac{a^2 \log(a + bx)}{2b^3} + \frac{a \cosh(a + bx) \sinh(a + bx)}{2b^3} \\
 & - \frac{x \cosh(a + bx) \sinh(a + bx)}{2b^2} \\
 & + \frac{\sinh^2(a + bx)}{4b^3} - \frac{2x \cosh(a + bx) \mathbf{Shi}(a + bx)}{b^2} \\
 & + \frac{2 \sinh(a + bx) \mathbf{Shi}(a + bx)}{b^3} \\
 & + \frac{x^2 \sinh(a + bx) \mathbf{Shi}(a + bx)}{b} - \frac{a \mathbf{Shi}(2a + 2bx)}{b^3}
 \end{aligned}$$

```
[Out] -1/2*a*x/b^2+1/4*x^2/b-Chi(2*b*x+2*a)/b^3-1/2*a^2*Chi(2*b*x+2*a)/b^3+1/2*cosh(2*b*x+2*a)/b^3+ln(b*x+a)/b^3+1/2*a^2*ln(b*x+a)/b^3-2*x*cosh(b*x+a)*Shi(b*x+a)/b^2-a*Shi(2*b*x+2*a)/b^3+1/2*a*cosh(b*x+a)*sinh(b*x+a)/b^3-1/2*x*cosh(b*x+a)*sinh(b*x+a)/b^2+2*Shi(b*x+a)*sinh(b*x+a)/b^3+x^2*Shi(b*x+a)*sinh(b*x+a)/b+1/4*sinh(b*x+a)^2/b^3
```

Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.00, number of steps used = 21, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {6683, 6874, 2715, 8, 3391, 30, 3393, 3382, 6677, 5736, 6873, 2718, 3379, 6681}

$$\int x^2 \cosh(a + bx) \operatorname{Shi}(a + bx) dx = -\frac{a^2 \operatorname{Chi}(2a + 2bx)}{2b^3} + \frac{a^2 \log(a + bx)}{2b^3} - \frac{\operatorname{Chi}(2a + 2bx)}{b^3} - \frac{a \operatorname{Shi}(2a + 2bx)}{b^3} + \frac{2 \operatorname{Shi}(a + bx) \sinh(a + bx)}{b^3} + \frac{\log(a + bx)}{b^3} + \frac{\sinh^2(a + bx)}{4b^3} + \frac{\cosh(2a + 2bx)}{2b^3} + \frac{a \sinh(a + bx) \cosh(a + bx)}{2b^3} - \frac{2x \operatorname{Shi}(a + bx) \cosh(a + bx)}{b^2} - \frac{ax}{2b^2} - \frac{x \sinh(a + bx) \cosh(a + bx)}{2b^2} + \frac{x^2 \operatorname{Shi}(a + bx) \sinh(a + bx)}{b} + \frac{x^2}{4b}$$

[In] Int[x^2*Cosh[a + b*x]*SinhIntegral[a + b*x],x]

[Out] -1/2*(a*x)/b^2 + x^2/(4*b) + Cosh[2*a + 2*b*x]/(2*b^3) - CoshIntegral[2*a + 2*b*x]/b^3 - (a^2*CoshIntegral[2*a + 2*b*x])/(2*b^3) + Log[a + b*x]/b^3 + (a^2*Log[a + b*x])/(2*b^3) + (a*Cosh[a + b*x]*Sinh[a + b*x])/(2*b^3) - (x*Cosh[a + b*x]*Sinh[a + b*x])/(2*b^2) + Sinh[a + b*x]^2/(4*b^3) - (2*x*Cosh[a + b*x]*SinhIntegral[a + b*x])/b^2 + (2*Sinh[a + b*x]*SinhIntegral[a + b*x])/b^3 + (x^2*Sinh[a + b*x]*SinhIntegral[a + b*x])/b - (a*SinhIntegral[2*a + 2*b*x])/b^3

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Sine[c + d*x])^(n - 1)/(d*n), x] + Dist[b^2*((n - 1)/n), Int[(b*Sine[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2718

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ
[{c, d}, x]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f
, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz
}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3391

```
Int[((c_.) + (d_.)*(x_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :=>
Simp[d*((b*Sin[e + f*x])^n/(f^2*n^2)), x] + (Dist[b^2*((n - 1)/n), Int[(c
+ d*x)*(b*Sin[e + f*x])^(n - 2), x], x] - Simp[b*(c + d*x)*Cos[e + f*x]*((b
*Sin[e + f*x])^(n - 1)/(f*n)), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1
]
```

Rule 3393

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := In
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 5736

```
Int[Cosh[w_]^(p_.)*(u_.)*Sinh[v_]^(p_.), x_Symbol] := Dist[1/2^p, Int[u*Sin
h[2*v]^p, x], x] /; EqQ[w, v] && IntegerQ[p]
```

Rule 6677

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(SinhIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Sinh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*SinhIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6681

```
Int[Cosh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=>
Simp[Sinh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
```

$b*x]*(\text{Sinh}[c + d*x]/(c + d*x)), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x]$

Rule 6683

$\text{Int}[\text{Cosh}[(a_.) + (b_.)*(x_.)]*((e_.) + (f_.)*(x_.))^{(m_.)*\text{SinhIntegral}[(c_.) + (d_.)*(x_.)]}, x_Symbol] \rightarrow \text{Simp}[(e + f*x)^m*\text{Sinh}[a + b*x]*(\text{SinhIntegral}[c + d*x]/b), x] + (-\text{Dist}[d/b, \text{Int}[(e + f*x)^m*\text{Sinh}[a + b*x]*(\text{Sinh}[c + d*x]/(c + d*x)), x], x] - \text{Dist}[f*(m/b), \text{Int}[(e + f*x)^{(m-1)}*\text{Sinh}[a + b*x]*\text{SinhIntegral}[c + d*x], x], x]) /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{IGtQ}[m, 0]$

Rule 6873

$\text{Int}[u_, x_Symbol] \rightarrow \text{With}[\{v = \text{NormalizeIntegrand}[u, x]\}, \text{Int}[v, x] /; v \neq u]$

Rule 6874

$\text{Int}[u_, x_Symbol] \rightarrow \text{With}[\{v = \text{ExpandIntegrand}[u, x]\}, \text{Int}[v, x] /; \text{SumQ}[v]]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{x^2 \sinh(a + bx) \text{Shi}(a + bx)}{b} \\ &\quad - \frac{2 \int x \sinh(a + bx) \text{Shi}(a + bx) dx}{b} - \int \frac{x^2 \sinh^2(a + bx)}{a + bx} dx \\ &= -\frac{2x \cosh(a + bx) \text{Shi}(a + bx)}{b^2} + \frac{x^2 \sinh(a + bx) \text{Shi}(a + bx)}{b} \\ &\quad + \frac{2 \int \cosh(a + bx) \text{Shi}(a + bx) dx}{b^2} + \frac{2 \int \frac{x \cosh(a + bx) \sinh(a + bx)}{a + bx} dx}{b} \\ &\quad - \int \left(-\frac{a \sinh^2(a + bx)}{b^2} + \frac{x \sinh^2(a + bx)}{b} + \frac{a^2 \sinh^2(a + bx)}{b^2(a + bx)} \right) dx \\ &= -\frac{2x \cosh(a + bx) \text{Shi}(a + bx)}{b^2} + \frac{2 \sinh(a + bx) \text{Shi}(a + bx)}{b^3} \\ &\quad + \frac{x^2 \sinh(a + bx) \text{Shi}(a + bx)}{b} - \frac{2 \int \frac{\sinh^2(a + bx)}{a + bx} dx}{b^2} + \frac{a \int \sinh^2(a + bx) dx}{b^2} \\ &\quad - \frac{a^2 \int \frac{\sinh^2(a + bx)}{a + bx} dx}{b^2} - \frac{\int x \sinh^2(a + bx) dx}{b} + \frac{\int \frac{x \sinh(2(a + bx))}{a + bx} dx}{b} \end{aligned}$$

$$\begin{aligned}
&= \frac{a \cosh(a+bx) \sinh(a+bx)}{2b^3} - \frac{x \cosh(a+bx) \sinh(a+bx)}{2b^2} + \frac{\sinh^2(a+bx)}{4b^3} \\
&\quad - \frac{2x \cosh(a+bx) \text{Shi}(a+bx)}{b^2} + \frac{2 \sinh(a+bx) \text{Shi}(a+bx)}{b^3} \\
&\quad + \frac{x^2 \sinh(a+bx) \text{Shi}(a+bx)}{b} + \frac{2 \int \left(\frac{1}{2(a+bx)} - \frac{\cosh(2a+2bx)}{2(a+bx)} \right) dx}{b^2} - \frac{a \int 1 dx}{2b^2} \\
&\quad + \frac{a^2 \int \left(\frac{1}{2(a+bx)} - \frac{\cosh(2a+2bx)}{2(a+bx)} \right) dx}{b^2} + \frac{\int x dx}{2b} + \frac{\int \frac{x \sinh(2a+2bx)}{a+bx} dx}{b} \\
&= -\frac{ax}{2b^2} + \frac{x^2}{4b} + \frac{\log(a+bx)}{b^3} + \frac{a^2 \log(a+bx)}{2b^3} + \frac{a \cosh(a+bx) \sinh(a+bx)}{2b^3} \\
&\quad - \frac{x \cosh(a+bx) \sinh(a+bx)}{2b^2} + \frac{\sinh^2(a+bx)}{4b^3} - \frac{2x \cosh(a+bx) \text{Shi}(a+bx)}{b^2} \\
&\quad + \frac{2 \sinh(a+bx) \text{Shi}(a+bx)}{b^3} + \frac{x^2 \sinh(a+bx) \text{Shi}(a+bx)}{b} \\
&\quad - \frac{\int \frac{\cosh(2a+2bx)}{a+bx} dx}{b^2} - \frac{a^2 \int \frac{\cosh(2a+2bx)}{a+bx} dx}{2b^2} + \frac{\int \left(\frac{\sinh(2a+2bx)}{b} + \frac{a \sinh(2a+2bx)}{b(-a-bx)} \right) dx}{b} \\
&= -\frac{ax}{2b^2} + \frac{x^2}{4b} - \frac{\text{Chi}(2a+2bx)}{b^3} - \frac{a^2 \text{Chi}(2a+2bx)}{2b^3} + \frac{\log(a+bx)}{b^3} \\
&\quad + \frac{a^2 \log(a+bx)}{2b^3} + \frac{a \cosh(a+bx) \sinh(a+bx)}{2b^3} - \frac{x \cosh(a+bx) \sinh(a+bx)}{2b^2} \\
&\quad + \frac{\sinh^2(a+bx)}{4b^3} - \frac{2x \cosh(a+bx) \text{Shi}(a+bx)}{b^2} + \frac{2 \sinh(a+bx) \text{Shi}(a+bx)}{b^3} \\
&\quad + \frac{x^2 \sinh(a+bx) \text{Shi}(a+bx)}{b} + \frac{\int \sinh(2a+2bx) dx}{b^2} + \frac{a \int \frac{\sinh(2a+2bx)}{-a-bx} dx}{b^2} \\
&= -\frac{ax}{2b^2} + \frac{x^2}{4b} + \frac{\cosh(2a+2bx)}{2b^3} - \frac{\text{Chi}(2a+2bx)}{b^3} - \frac{a^2 \text{Chi}(2a+2bx)}{2b^3} \\
&\quad + \frac{\log(a+bx)}{b^3} + \frac{a^2 \log(a+bx)}{2b^3} + \frac{a \cosh(a+bx) \sinh(a+bx)}{2b^3} \\
&\quad - \frac{x \cosh(a+bx) \sinh(a+bx)}{2b^2} + \frac{\sinh^2(a+bx)}{4b^3} - \frac{2x \cosh(a+bx) \text{Shi}(a+bx)}{b^2} \\
&\quad + \frac{2 \sinh(a+bx) \text{Shi}(a+bx)}{b^3} + \frac{x^2 \sinh(a+bx) \text{Shi}(a+bx)}{b} - \frac{a \text{Shi}(2a+2bx)}{b^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.61

$$\int x^2 \cosh(a + bx) \operatorname{Shi}(a + bx) dx$$

$$= \frac{-4abx + 2b^2x^2 + 5 \cosh(2(a + bx)) - 4(2 + a^2) \operatorname{Chi}(2(a + bx)) + 8 \log(a + bx) + 4a^2 \log(a + bx) + 2a \operatorname{Shi}(2(a + bx))}{8b^3}$$

[In] Integrate[x^2*Cosh[a + b*x]*SinhIntegral[a + b*x],x]

[Out] $(-4*a*b*x + 2*b^2*x^2 + 5*\operatorname{Cosh}[2*(a + b*x)] - 4*(2 + a^2)*\operatorname{CoshIntegral}[2*(a + b*x)] + 8*\operatorname{Log}[a + b*x] + 4*a^2*\operatorname{Log}[a + b*x] + 2*a*\operatorname{Sinh}[2*(a + b*x)] - 2*b*x*\operatorname{Sinh}[2*(a + b*x)] + 8*(-2*b*x*\operatorname{Cosh}[a + b*x] + (2 + b^2*x^2)*\operatorname{Sinh}[a + b*x])*\operatorname{SinhIntegral}[a + b*x] - 8*a*\operatorname{SinhIntegral}[2*(a + b*x)])/(8*b^3)$

Maple [A] (verified)

Time = 2.45 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.90

method	result
derivativedivides	$\frac{\operatorname{Shi}(bx+a)(a^2 \sinh(bx+a) - 2a((bx+a) \sinh(bx+a) - \cosh(bx+a)) + (bx+a)^2 \sinh(bx+a) - 2(bx+a) \cosh(bx+a) + 2 \sinh(bx+a))}{8b^3}$
default	$\frac{\operatorname{Shi}(bx+a)(a^2 \sinh(bx+a) - 2a((bx+a) \sinh(bx+a) - \cosh(bx+a)) + (bx+a)^2 \sinh(bx+a) - 2(bx+a) \cosh(bx+a) + 2 \sinh(bx+a))}{8b^3}$

[In] int(x^2*cosh(b*x+a)*Shi(b*x+a),x,method=_RETURNVERBOSE)

[Out] $1/b^3*(\operatorname{Shi}(b*x+a)*(a^2*\sinh(b*x+a)-2*a*((b*x+a)*\sinh(b*x+a)-\cosh(b*x+a))+(b*x+a)^2*\sinh(b*x+a)-2*(b*x+a)*\cosh(b*x+a)+2*\sinh(b*x+a))+1/2*a^2*\ln(b*x+a)-1/2*a^2*\operatorname{Chi}(2*b*x+2*a)+\cosh(b*x+a)*\sinh(b*x+a)*a-(b*x+a)*a-a*\operatorname{Shi}(2*b*x+2*a)-1/2*(b*x+a)*\cosh(b*x+a)*\sinh(b*x+a)+1/4*(b*x+a)^2+5/4*\cosh(b*x+a)^2+\ln(b*x+a)-\operatorname{Chi}(2*b*x+2*a))$

Fricas [F]

$$\int x^2 \cosh(a + bx) \operatorname{Shi}(a + bx) dx = \int x^2 \operatorname{Shi}(bx + a) \cosh(bx + a) dx$$

[In] integrate(x^2*cosh(b*x+a)*Shi(b*x+a),x, algorithm="fricas")

[Out] integral(x^2*cosh(b*x + a)*sinh_integral(b*x + a), x)

Sympy [F]

$$\int x^2 \cosh(a + bx) \operatorname{Shi}(a + bx) dx = \int x^2 \cosh(a + bx) \operatorname{Shi}(a + bx) dx$$

[In] integrate(x**2*cosh(b*x+a)*Shi(b*x+a),x)

[Out] Integral(x**2*cosh(a + b*x)*Shi(a + b*x), x)

Maxima [F]

$$\int x^2 \cosh(a + bx) \operatorname{Shi}(a + bx) dx = \int x^2 \operatorname{Shi}(bx + a) \cosh(bx + a) dx$$

[In] integrate(x^2*cosh(b*x+a)*Shi(b*x+a),x, algorithm="maxima")

[Out] integrate(x^2*Shi(b*x + a)*cosh(b*x + a), x)

Giac [F]

$$\int x^2 \cosh(a + bx) \operatorname{Shi}(a + bx) dx = \int x^2 \operatorname{Shi}(bx + a) \cosh(bx + a) dx$$

[In] integrate(x^2*cosh(b*x+a)*Shi(b*x+a),x, algorithm="giac")

[Out] integrate(x^2*Shi(b*x + a)*cosh(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \cosh(a + bx) \operatorname{Shi}(a + bx) dx = \int x^2 \operatorname{sinhint}(a + bx) \cosh(a + bx) dx$$

[In] int(x^2*sinhint(a + b*x)*cosh(a + b*x),x)

[Out] int(x^2*sinhint(a + b*x)*cosh(a + b*x), x)

3.60 $\int x \cosh(a + bx) \text{Shi}(a + bx) dx$

Optimal result	321
Rubi [A] (verified)	321
Mathematica [A] (verified)	324
Maple [A] (verified)	324
Fricas [F]	324
Sympy [F]	325
Maxima [F]	325
Giac [F]	325
Mupad [F(-1)]	325

Optimal result

Integrand size = 14, antiderivative size = 109

$$\int x \cosh(a + bx) \text{Shi}(a + bx) dx = \frac{x}{2b} + \frac{a \text{Chi}(2a + 2bx)}{2b^2} - \frac{a \log(a + bx)}{2b^2} - \frac{\cosh(a + bx) \sinh(a + bx)}{2b^2} - \frac{\cosh(a + bx) \text{Shi}(a + bx)}{b^2} + \frac{x \sinh(a + bx) \text{Shi}(a + bx)}{b} + \frac{\text{Shi}(2a + 2bx)}{2b^2}$$

[Out] $1/2*x/b + 1/2*a*\text{Chi}(2*b*x + 2*a)/b^2 - 1/2*a*\ln(b*x + a)/b^2 - \cosh(b*x + a)*\text{Shi}(b*x + a)/b^2 + 1/2*\text{Shi}(2*b*x + 2*a)/b^2 - 1/2*\cosh(b*x + a)*\sinh(b*x + a)/b^2 + x*\text{Shi}(b*x + a)*\sinh(b*x + a)/b$

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {6683, 6874, 2715, 8, 3393, 3382, 6675, 5556, 12, 3379}

$$\int x \cosh(a + bx) \text{Shi}(a + bx) dx = \frac{a \text{Chi}(2a + 2bx)}{2b^2} + \frac{\text{Shi}(2a + 2bx)}{2b^2} - \frac{\text{Shi}(a + bx) \cosh(a + bx)}{b^2} - \frac{a \log(a + bx)}{2b^2} - \frac{\sinh(a + bx) \cosh(a + bx)}{2b^2} + \frac{x \text{Shi}(a + bx) \sinh(a + bx)}{b} + \frac{x}{2b}$$

[In] $\text{Int}[x*\text{Cosh}[a + b*x]*\text{SinhIntegral}[a + b*x], x]$

```
[Out] x/(2*b) + (a*CoshIntegral[2*a + 2*b*x])/(2*b^2) - (a*Log[a + b*x])/(2*b^2)
- (Cosh[a + b*x]*Sinh[a + b*x])/(2*b^2) - (Cosh[a + b*x]*SinhIntegral[a + b
*x])/b^2 + (x*Sinh[a + b*x]*SinhIntegral[a + b*x])/b + SinhIntegral[2*a + 2
*b*x]/(2*b^2)
```

Rule 8

```
Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[
c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2
*n]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f
, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz
}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3393

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)^(n_)], x_Symbol] := In
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_)^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)^(n_.)], x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

Rule 6675

```
Int[Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
  Simp[Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
  b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6683

```
Int[Cosh[(a_.) + (b_.)*(x_)]*((e_.) + (f_.)*(x_)^(m_.)*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Sinh[a + b*x]*(SinhIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Sinh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*SinhIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x \sinh(a + bx) \text{Shi}(a + bx)}{b} - \frac{\int \sinh(a + bx) \text{Shi}(a + bx) dx}{b} - \int \frac{x \sinh^2(a + bx)}{a + bx} dx \\
&= -\frac{\cosh(a + bx) \text{Shi}(a + bx)}{b^2} + \frac{x \sinh(a + bx) \text{Shi}(a + bx)}{b} \\
&\quad + \frac{\int \frac{\cosh(a + bx) \sinh(a + bx)}{a + bx} dx}{b} - \int \left(\frac{\sinh^2(a + bx)}{b} - \frac{a \sinh^2(a + bx)}{b(a + bx)} \right) dx \\
&= -\frac{\cosh(a + bx) \text{Shi}(a + bx)}{b^2} + \frac{x \sinh(a + bx) \text{Shi}(a + bx)}{b} \\
&\quad - \frac{\int \sinh^2(a + bx) dx}{b} + \frac{\int \frac{\sinh(2a + 2bx)}{2(a + bx)} dx}{b} + \frac{a \int \frac{\sinh^2(a + bx)}{a + bx} dx}{b} \\
&= -\frac{\cosh(a + bx) \sinh(a + bx)}{2b^2} - \frac{\cosh(a + bx) \text{Shi}(a + bx)}{b^2} + \frac{x \sinh(a + bx) \text{Shi}(a + bx)}{b} \\
&\quad + \frac{\int 1 dx}{2b} + \frac{\int \frac{\sinh(2a + 2bx)}{a + bx} dx}{2b} - \frac{a \int \left(\frac{1}{2(a + bx)} - \frac{\cosh(2a + 2bx)}{2(a + bx)} \right) dx}{b} \\
&= \frac{x}{2b} - \frac{a \log(a + bx)}{2b^2} - \frac{\cosh(a + bx) \sinh(a + bx)}{2b^2} - \frac{\cosh(a + bx) \text{Shi}(a + bx)}{b^2} \\
&\quad + \frac{x \sinh(a + bx) \text{Shi}(a + bx)}{b} + \frac{\text{Shi}(2a + 2bx)}{2b^2} + \frac{a \int \frac{\cosh(2a + 2bx)}{a + bx} dx}{2b} \\
&= \frac{x}{2b} + \frac{a \text{Chi}(2a + 2bx)}{2b^2} - \frac{a \log(a + bx)}{2b^2} - \frac{\cosh(a + bx) \sinh(a + bx)}{2b^2} \\
&\quad - \frac{\cosh(a + bx) \text{Shi}(a + bx)}{b^2} + \frac{x \sinh(a + bx) \text{Shi}(a + bx)}{b} + \frac{\text{Shi}(2a + 2bx)}{2b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.72

$$\int x \cosh(a + bx) \operatorname{Shi}(a + bx) dx$$

$$= \frac{2bx + 2a \operatorname{Chi}(2(a + bx)) - 2a \log(a + bx) - \sinh(2(a + bx)) + 4(-\cosh(a + bx) + bx \sinh(a + bx)) \operatorname{Shi}(a + bx)}{4b^2}$$

[In] Integrate[x*Cosh[a + b*x]*SinhIntegral[a + b*x],x]

[Out] (2*b*x + 2*a*CoshIntegral[2*(a + b*x)] - 2*a*Log[a + b*x] - Sinh[2*(a + b*x)]) + 4*(-Cosh[a + b*x] + b*x*Sinh[a + b*x])*SinhIntegral[a + b*x] + 2*SinhIntegral[2*(a + b*x)]/(4*b^2)

Maple [A] (verified)

Time = 1.54 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.89

method	result
derivativedivides	$\frac{\operatorname{Shi}(bx+a)(-a \sinh(bx+a) + (bx+a) \sinh(bx+a) - \cosh(bx+a)) + a \left(-\frac{\ln(bx+a)}{2} + \frac{\operatorname{Chi}(2bx+2a)}{2} \right) - \frac{\cosh(bx+a) \sinh(bx+a)}{2} + \frac{bx}{2}}{b^2}$
default	$\frac{\operatorname{Shi}(bx+a)(-a \sinh(bx+a) + (bx+a) \sinh(bx+a) - \cosh(bx+a)) + a \left(-\frac{\ln(bx+a)}{2} + \frac{\operatorname{Chi}(2bx+2a)}{2} \right) - \frac{\cosh(bx+a) \sinh(bx+a)}{2} + \frac{bx}{2}}{b^2}$

[In] int(x*cosh(b*x+a)*Shi(b*x+a),x,method=_RETURNVERBOSE)

[Out] 1/b^2*(Shi(b*x+a)*(-a*sinh(b*x+a)+(b*x+a)*sinh(b*x+a)-cosh(b*x+a))+a*(-1/2*ln(b*x+a)+1/2*Chi(2*b*x+2*a))-1/2*cosh(b*x+a)*sinh(b*x+a)+1/2*b*x+1/2*a+1/2*Shi(2*b*x+2*a))

Fricas [F]

$$\int x \cosh(a + bx) \operatorname{Shi}(a + bx) dx = \int x \operatorname{Shi}(bx + a) \cosh(bx + a) dx$$

[In] integrate(x*cosh(b*x+a)*Shi(b*x+a),x, algorithm="fricas")

[Out] integral(x*cosh(b*x + a)*sinh_integral(b*x + a), x)

Sympy [F]

$$\int x \cosh(a + bx) \operatorname{Shi}(a + bx) dx = \int x \cosh(a + bx) \operatorname{Shi}(a + bx) dx$$

[In] `integrate(x*cosh(b*x+a)*Shi(b*x+a),x)`

[Out] `Integral(x*cosh(a + b*x)*Shi(a + b*x), x)`

Maxima [F]

$$\int x \cosh(a + bx) \operatorname{Shi}(a + bx) dx = \int x \operatorname{Shi}(bx + a) \cosh(bx + a) dx$$

[In] `integrate(x*cosh(b*x+a)*Shi(b*x+a),x, algorithm="maxima")`

[Out] `integrate(x*Shi(b*x + a)*cosh(b*x + a), x)`

Giac [F]

$$\int x \cosh(a + bx) \operatorname{Shi}(a + bx) dx = \int x \operatorname{Shi}(bx + a) \cosh(bx + a) dx$$

[In] `integrate(x*cosh(b*x+a)*Shi(b*x+a),x, algorithm="giac")`

[Out] `integrate(x*Shi(b*x + a)*cosh(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x \cosh(a + bx) \operatorname{Shi}(a + bx) dx = \int x \operatorname{sinhint}(a + bx) \cosh(a + bx) dx$$

[In] `int(x*sinhint(a + b*x)*cosh(a + b*x),x)`

[Out] `int(x*sinhint(a + b*x)*cosh(a + b*x), x)`

3.61 $\int \cosh(a + bx)\text{Shi}(a + bx) dx$

Optimal result	326
Rubi [A] (verified)	326
Mathematica [A] (verified)	327
Maple [A] (verified)	327
Fricas [F]	328
Sympy [F]	328
Maxima [F]	328
Giac [F]	328
Mupad [F(-1)]	329

Optimal result

Integrand size = 13, antiderivative size = 46

$$\int \cosh(a + bx)\text{Shi}(a + bx) dx = -\frac{\text{Chi}(2a + 2bx)}{2b} + \frac{\log(a + bx)}{2b} + \frac{\sinh(a + bx)\text{Shi}(a + bx)}{b}$$

[Out] $-1/2*\text{Chi}(2*b*x+2*a)/b+1/2*\ln(b*x+a)/b+\text{Shi}(b*x+a)*\sinh(b*x+a)/b$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {6681, 3393, 3382}

$$\int \cosh(a + bx)\text{Shi}(a + bx) dx = -\frac{\text{Chi}(2a + 2bx)}{2b} + \frac{\text{Shi}(a + bx) \sinh(a + bx)}{b} + \frac{\log(a + bx)}{2b}$$

[In] `Int[Cosh[a + b*x]*SinhIntegral[a + b*x],x]`

[Out] $-1/2*\text{CoshIntegral}[2*a + 2*b*x]/b + \text{Log}[a + b*x]/(2*b) + (\text{Sinh}[a + b*x]*\text{SinhIntegral}[a + b*x])/b$

Rule 3382

`Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] :> Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`

Rule 3393

`Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] :> Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f`

, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))

Rule 6681

```
Int[Cosh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :>
Simp[Sinh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\sinh(a + bx)\text{Shi}(a + bx)}{b} - \int \frac{\sinh^2(a + bx)}{a + bx} dx \\
 &= \frac{\sinh(a + bx)\text{Shi}(a + bx)}{b} + \int \left(\frac{1}{2(a + bx)} - \frac{\cosh(2a + 2bx)}{2(a + bx)} \right) dx \\
 &= \frac{\log(a + bx)}{2b} + \frac{\sinh(a + bx)\text{Shi}(a + bx)}{b} - \frac{1}{2} \int \frac{\cosh(2a + 2bx)}{a + bx} dx \\
 &= -\frac{\text{Chi}(2a + 2bx)}{2b} + \frac{\log(a + bx)}{2b} + \frac{\sinh(a + bx)\text{Shi}(a + bx)}{b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.98

$$\int \cosh(a + bx)\text{Shi}(a + bx) dx = -\frac{\text{Chi}(2(a + bx))}{2b} + \frac{\log(a + bx)}{2b} + \frac{\sinh(a + bx)\text{Shi}(a + bx)}{b}$$

[In] Integrate[Cosh[a + b*x]*SinhIntegral[a + b*x],x]

[Out] -1/2*CoshIntegral[2*(a + b*x)]/b + Log[a + b*x]/(2*b) + (Sinh[a + b*x]*SinhIntegral[a + b*x])/b

Maple [A] (verified)

Time = 0.88 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$\frac{\text{Shi}(bx+a) \sinh(bx+a) + \frac{\ln(bx+a)}{2} - \frac{\text{Chi}(2bx+2a)}{2}}{b}$	38
default	$\frac{\text{Shi}(bx+a) \sinh(bx+a) + \frac{\ln(bx+a)}{2} - \frac{\text{Chi}(2bx+2a)}{2}}{b}$	38

[In] int(cosh(b*x+a)*Shi(b*x+a),x,method=_RETURNVERBOSE)

[Out] 1/b*(Shi(b*x+a)*sinh(b*x+a)+1/2*ln(b*x+a)-1/2*Chi(2*b*x+2*a))

Fricas [F]

$$\int \cosh(a + bx)\operatorname{Shi}(a + bx) dx = \int \operatorname{Shi}(bx + a) \cosh(bx + a) dx$$

[In] `integrate(cosh(b*x+a)*Shi(b*x+a),x, algorithm="fricas")`

[Out] `integral(cosh(b*x + a)*sinh_integral(b*x + a), x)`

Sympy [F]

$$\int \cosh(a + bx)\operatorname{Shi}(a + bx) dx = \int \cosh(a + bx) \operatorname{Shi}(a + bx) dx$$

[In] `integrate(cosh(b*x+a)*Shi(b*x+a),x)`

[Out] `Integral(cosh(a + b*x)*Shi(a + b*x), x)`

Maxima [F]

$$\int \cosh(a + bx)\operatorname{Shi}(a + bx) dx = \int \operatorname{Shi}(bx + a) \cosh(bx + a) dx$$

[In] `integrate(cosh(b*x+a)*Shi(b*x+a),x, algorithm="maxima")`

[Out] `integrate(Shi(b*x + a)*cosh(b*x + a), x)`

Giac [F]

$$\int \cosh(a + bx)\operatorname{Shi}(a + bx) dx = \int \operatorname{Shi}(bx + a) \cosh(bx + a) dx$$

[In] `integrate(cosh(b*x+a)*Shi(b*x+a),x, algorithm="giac")`

[Out] `integrate(Shi(b*x + a)*cosh(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \cosh(a + bx)\text{Shi}(a + bx) dx = \int \sinhint(a + bx) \cosh(a + bx) dx$$

```
[In] int(sinhint(a + b*x)*cosh(a + b*x),x)
```

```
[Out] int(sinhint(a + b*x)*cosh(a + b*x), x)
```

3.62 $\int \frac{\cosh(a+bx)\mathbf{Shi}(a+bx)}{x} dx$

Optimal result	330
Rubi [N/A]	330
Mathematica [N/A]	331
Maple [N/A] (verified)	331
Fricas [N/A]	331
Sympy [N/A]	331
Maxima [N/A]	332
Giac [N/A]	332
Mupad [N/A]	332

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\cosh(a+bx)\mathbf{Shi}(a+bx)}{x} dx = \text{Int}\left(\frac{\cosh(a+bx)\mathbf{Shi}(a+bx)}{x}, x\right)$$

[Out] CannotIntegrate(cosh(b*x+a)*Shi(b*x+a)/x,x)

Rubi [N/A]

Not integrable

Time = 0.08 (sec), antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\cosh(a+bx)\mathbf{Shi}(a+bx)}{x} dx = \int \frac{\cosh(a+bx)\mathbf{Shi}(a+bx)}{x} dx$$

[In] Int[(Cosh[a + b*x]*SinhIntegral[a + b*x])/x,x]

[Out] Defer[Int] [(Cosh[a + b*x]*SinhIntegral[a + b*x])/x, x]

Rubi steps

$$\text{integral} = \int \frac{\cosh(a+bx)\mathbf{Shi}(a+bx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 0.53 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Shi}(a + bx)}{x} dx = \int \frac{\cosh(a + bx)\text{Shi}(a + bx)}{x} dx$$

[In] Integrate[(Cosh[a + b*x]*SinhIntegral[a + b*x])/x,x]

[Out] Integrate[(Cosh[a + b*x]*SinhIntegral[a + b*x])/x, x]

Maple [N/A] (verified)

Not integrable

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\cosh(bx + a)\text{Shi}(bx + a)}{x} dx$$

[In] int(cosh(b*x+a)*Shi(b*x+a)/x,x)

[Out] int(cosh(b*x+a)*Shi(b*x+a)/x,x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Shi}(a + bx)}{x} dx = \int \frac{\text{Shi}(bx + a)\cosh(bx + a)}{x} dx$$

[In] integrate(cosh(b*x+a)*Shi(b*x+a)/x,x, algorithm="fricas")

[Out] integral(cosh(b*x + a)*sinh_integral(b*x + a)/x, x)

Sympy [N/A]

Not integrable

Time = 1.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{\cosh(a + bx)\text{Shi}(a + bx)}{x} dx = \int \frac{\cosh(a + bx)\text{Shi}(a + bx)}{x} dx$$

[In] integrate(cosh(b*x+a)*Shi(b*x+a)/x,x)

[Out] Integral(cosh(a + b*x)*Shi(a + b*x)/x, x)

Maxima [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Shi}(a + bx)}{x} dx = \int \frac{\text{Shi}(bx + a) \cosh(bx + a)}{x} dx$$

[In] integrate(cosh(b*x+a)*Shi(b*x+a)/x,x, algorithm="maxima")

[Out] integrate(Shi(b*x + a)*cosh(b*x + a)/x, x)

Giac [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Shi}(a + bx)}{x} dx = \int \frac{\text{Shi}(bx + a) \cosh(bx + a)}{x} dx$$

[In] integrate(cosh(b*x+a)*Shi(b*x+a)/x,x, algorithm="giac")

[Out] integrate(Shi(b*x + a)*cosh(b*x + a)/x, x)

Mupad [N/A]

Not integrable

Time = 4.86 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Shi}(a + bx)}{x} dx = \int \frac{\sinhint(a + bx) \cosh(a + bx)}{x} dx$$

[In] int((sinhint(a + b*x)*cosh(a + b*x))/x,x)

[Out] int((sinhint(a + b*x)*cosh(a + b*x))/x, x)

3.63 $\int x \sinh(a + bx) \mathbf{Shi}(c + dx) dx$

Optimal result	333
Rubi [A] (verified)	334
Mathematica [A] (verified)	338
Maple [F]	338
Fricas [F]	339
Sympy [F]	339
Maxima [F]	339
Giac [F]	339
Mupad [F(-1)]	340

Optimal result

Integrand size = 14, antiderivative size = 371

$$\begin{aligned}
 \int x \sinh(a + bx) \mathbf{Shi}(c + dx) dx = & \frac{\cosh(a - c + (b - d)x)}{2b(b - d)} - \frac{\cosh(a + c + (b + d)x)}{2b(b + d)} \\
 & - \frac{\cosh\left(a - \frac{bc}{d}\right) \mathbf{Chi}\left(\frac{c(b-d)}{d} + (b - d)x\right)}{2b^2} \\
 & + \frac{\cosh\left(a - \frac{bc}{d}\right) \mathbf{Chi}\left(\frac{c(b+d)}{d} + (b + d)x\right)}{2b^2} \\
 & - \frac{c \mathbf{Chi}\left(\frac{c(b-d)}{d} + (b - d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2bd} \\
 & + \frac{c \mathbf{Chi}\left(\frac{c(b+d)}{d} + (b + d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2bd} \\
 & - \frac{c \cosh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b-d)}{d} + (b - d)x\right)}{2bd} \\
 & - \frac{\sinh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b-d)}{d} + (b - d)x\right)}{2b^2} \\
 & + \frac{x \cosh(a + bx) \mathbf{Shi}(c + dx)}{b} - \frac{\sinh(a + bx) \mathbf{Shi}(c + dx)}{b^2} \\
 & + \frac{c \cosh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b+d)}{d} + (b + d)x\right)}{2bd} \\
 & + \frac{\sinh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b+d)}{d} + (b + d)x\right)}{2b^2}
 \end{aligned}$$

[Out] $-1/2*\mathbf{Chi}(c*(b-d)/d+(b-d)*x)*\cosh(a-b*c/d)/b^2+1/2*\mathbf{Chi}(c*(b+d)/d+(b+d)*x)*\cosh(a-b*c/d)/b^2+1/2*\cosh(a-c+(b-d)*x)/b/(b-d)-1/2*\cosh(a+c+(b+d)*x)/b/(b+d)$

$$\begin{aligned}
& -1/2*c*cosh(a-b*c/d)*Shi(c*(b-d)/d+(b-d)*x)/b/d+x*cosh(b*x+a)*Shi(d*x+c)/b+ \\
& 1/2*c*cosh(a-b*c/d)*Shi(c*(b+d)/d+(b+d)*x)/b/d-1/2*c*Chi(c*(b-d)/d+(b-d)*x) \\
& *sinh(a-b*c/d)/b/d+1/2*c*Chi(c*(b+d)/d+(b+d)*x)*sinh(a-b*c/d)/b/d-1/2*Shi(c \\
& *(b-d)/d+(b-d)*x)*sinh(a-b*c/d)/b^2+1/2*Shi(c*(b+d)/d+(b+d)*x)*sinh(a-b*c/d \\
&)/b^2-Shi(d*x+c)*sinh(b*x+a)/b^2
\end{aligned}$$

Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 371, normalized size of antiderivative = 1.00, number of steps used = 24, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {6677, 6874, 5737, 2718, 5580, 3384, 3379, 3382, 6681, 5578}

$$\begin{aligned}
\int x \sinh(a + bx) \text{Shi}(c + dx) dx = & -\frac{\cosh\left(a - \frac{bc}{d}\right) \text{Chi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2b^2} \\
& + \frac{\cosh\left(a - \frac{bc}{d}\right) \text{Chi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2b^2} \\
& - \frac{\sinh\left(a - \frac{bc}{d}\right) \text{Shi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2b^2} \\
& - \frac{\sinh(a + bx) \text{Shi}(c + dx)}{b^2} \\
& + \frac{\sinh\left(a - \frac{bc}{d}\right) \text{Shi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2b^2} \\
& - \frac{c \sinh\left(a - \frac{bc}{d}\right) \text{Chi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2bd} \\
& + \frac{c \sinh\left(a - \frac{bc}{d}\right) \text{Chi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2bd} \\
& - \frac{c \cosh\left(a - \frac{bc}{d}\right) \text{Shi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2bd} \\
& + \frac{x \cosh(a + bx) \text{Shi}(c + dx)}{b} \\
& + \frac{c \cosh\left(a - \frac{bc}{d}\right) \text{Shi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2bd} \\
& + \frac{\cosh(a + x(b-d) - c)}{2b(b-d)} - \frac{\cosh(a + x(b+d) + c)}{2b(b+d)}
\end{aligned}$$

[In] Int[x*Sinh[a + b*x]*SinhIntegral[c + d*x],x]

[Out] Cosh[a - c + (b - d)*x]/(2*b*(b - d)) - Cosh[a + c + (b + d)*x]/(2*b*(b + d)) - (Cosh[a - (b*c)/d]*CoshIntegral[(c*(b - d))/d + (b - d)*x])/(2*b^2) + (Cosh[a - (b*c)/d]*CoshIntegral[(c*(b + d))/d + (b + d)*x])/(2*b^2) - (c*Co

$$\begin{aligned} & \text{shIntegral}[(c*(b-d))/d + (b-d)*x]*\text{Sinh}[a - (b*c)/d]/(2*b*d) + (c*\text{CoshIntegral}[(c*(b+d))/d + (b+d)*x]*\text{Sinh}[a - (b*c)/d]/(2*b*d) - (c*\text{Cosh}[a - (b*c)/d]*\text{SinhIntegral}[(c*(b-d))/d + (b-d)*x])/(2*b*d) - (\text{Sinh}[a - (b*c)/d]*\text{SinhIntegral}[(c*(b-d))/d + (b-d)*x])/(2*b^2) + (x*\text{Cosh}[a + b*x]*\text{SinhIntegral}[c + d*x])/b - (\text{Sinh}[a + b*x]*\text{SinhIntegral}[c + d*x])/b^2 + (c*\text{Cosh}[a - (b*c)/d]*\text{SinhIntegral}[(c*(b+d))/d + (b+d)*x])/(2*b*d) + (\text{Sinh}[a - (b*c)/d]*\text{SinhIntegral}[(c*(b+d))/d + (b+d)*x])/(2*b^2) \end{aligned}$$
Rule 2718

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3384

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]
```

Rule 5578

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(p_.)*Sinh[(c_.) + (d_.)*(x_)]^(q_.), x_Symbol] := Int[ExpandTrigReduce[(e + f*x)^m, Sinh[a + b*x]^p*Sinh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IntegerQ[m]
```

Rule 5580

```
Int[Cosh[(c_.) + (d_.)*(x_)]^(q_.)*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(p_.), x_Symbol] := Int[ExpandTrigReduce[(e + f*x)^m, Sinh[a + b*x]^p*Cosh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[p, 0] && IGtQ[q, 0]
```

Rule 5737

```
Int[Cosh[w_]^(q_.)*Sinh[v_]^(p_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]
]~p*Cosh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))
```

Rule 6677

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(SinhIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Sinh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*SinhIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6681

```
Int[Cosh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
Simp[Sinh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x \cosh(a + bx) \text{Shi}(c + dx)}{b} - \frac{\int \cosh(a + bx) \text{Shi}(c + dx) dx}{b} \\
&\quad - \frac{d \int \frac{x \cosh(a+bx) \sinh(c+dx)}{c+dx} dx}{b} \\
&= \frac{x \cosh(a + bx) \text{Shi}(c + dx)}{b} - \frac{\sinh(a + bx) \text{Shi}(c + dx)}{b^2} \\
&\quad + \frac{d \int \frac{\sinh(a+bx) \sinh(c+dx)}{c+dx} dx}{b^2} - \frac{d \int \left(\frac{\cosh(a+bx) \sinh(c+dx)}{d} - \frac{c \cosh(a+bx) \sinh(c+dx)}{d(c+dx)} \right) dx}{b} \\
&= \frac{x \cosh(a + bx) \text{Shi}(c + dx)}{b} - \frac{\sinh(a + bx) \text{Shi}(c + dx)}{b^2} \\
&\quad - \frac{\int \cosh(a + bx) \sinh(c + dx) dx}{b} + \frac{c \int \frac{\cosh(a+bx) \sinh(c+dx)}{c+dx} dx}{b} \\
&\quad + \frac{d \int \left(-\frac{\cosh(a-c+(b-d)x)}{2(c+dx)} + \frac{\cosh(a+c+(b+d)x)}{2(c+dx)} \right) dx}{b^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{x \cosh(a + bx) \operatorname{Shi}(c + dx)}{b} - \frac{\sinh(a + bx) \operatorname{Shi}(c + dx)}{b^2} \\
&\quad - \frac{\int \left(-\frac{1}{2} \sinh(a - c + (b - d)x) + \frac{1}{2} \sinh(a + c + (b + d)x) \right) dx}{b} \\
&\quad + \frac{c \int \left(-\frac{\sinh(a - c + (b - d)x)}{2(c + dx)} + \frac{\sinh(a + c + (b + d)x)}{2(c + dx)} \right) dx}{b} \\
&\quad - \frac{d \int \frac{\cosh(a - c + (b - d)x)}{c + dx} dx}{2b^2} + \frac{d \int \frac{\cosh(a + c + (b + d)x)}{c + dx} dx}{2b^2} \\
&= \frac{x \cosh(a + bx) \operatorname{Shi}(c + dx)}{b} - \frac{\sinh(a + bx) \operatorname{Shi}(c + dx)}{b^2} + \frac{\int \sinh(a - c + (b - d)x) dx}{2b} \\
&\quad - \frac{\int \sinh(a + c + (b + d)x) dx}{2b} - \frac{c \int \frac{\sinh(a - c + (b - d)x)}{c + dx} dx}{2b} + \frac{c \int \frac{\sinh(a + c + (b + d)x)}{c + dx} dx}{2b} \\
&\quad - \frac{(d \cosh(a - \frac{bc}{d})) \int \frac{\cosh(\frac{c(b-d)}{d} + (b-d)x)}{c + dx} dx}{2b^2} + \frac{(d \cosh(a - \frac{bc}{d})) \int \frac{\cosh(\frac{c(b+d)}{d} + (b+d)x)}{c + dx} dx}{2b^2} \\
&\quad - \frac{(d \sinh(a - \frac{bc}{d})) \int \frac{\sinh(\frac{c(b-d)}{d} + (b-d)x)}{c + dx} dx}{2b^2} + \frac{(d \sinh(a - \frac{bc}{d})) \int \frac{\sinh(\frac{c(b+d)}{d} + (b+d)x)}{c + dx} dx}{2b^2} \\
&= \frac{\cosh(a - c + (b - d)x)}{2b(b - d)} - \frac{\cosh(a + c + (b + d)x)}{2b(b + d)} \\
&\quad - \frac{\cosh(a - \frac{bc}{d}) \operatorname{Chi}\left(\frac{c(b-d)}{d} + (b - d)x\right)}{2b^2} + \frac{\cosh(a - \frac{bc}{d}) \operatorname{Chi}\left(\frac{c(b+d)}{d} + (b + d)x\right)}{2b^2} \\
&\quad - \frac{\sinh(a - \frac{bc}{d}) \operatorname{Shi}\left(\frac{c(b-d)}{d} + (b - d)x\right)}{2b^2} + \frac{x \cosh(a + bx) \operatorname{Shi}(c + dx)}{b} \\
&\quad - \frac{\sinh(a + bx) \operatorname{Shi}(c + dx)}{b^2} + \frac{\sinh(a - \frac{bc}{d}) \operatorname{Shi}\left(\frac{c(b+d)}{d} + (b + d)x\right)}{2b^2} \\
&\quad - \frac{(c \cosh(a - \frac{bc}{d})) \int \frac{\sinh(\frac{c(b-d)}{d} + (b-d)x)}{c + dx} dx}{2b} \\
&\quad + \frac{(c \cosh(a - \frac{bc}{d})) \int \frac{\sinh(\frac{c(b+d)}{d} + (b+d)x)}{c + dx} dx}{2b} \\
&\quad - \frac{(c \sinh(a - \frac{bc}{d})) \int \frac{\cosh(\frac{c(b-d)}{d} + (b-d)x)}{c + dx} dx}{2b} \\
&\quad + \frac{(c \sinh(a - \frac{bc}{d})) \int \frac{\cosh(\frac{c(b+d)}{d} + (b+d)x)}{c + dx} dx}{2b}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\cosh(a - c + (b - d)x)}{2b(b - d)} - \frac{\cosh(a + c + (b + d)x)}{2b(b + d)} \\
&\quad - \frac{\cosh\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(\frac{c(b-d)}{d} + (b - d)x\right)}{2b^2} + \frac{\cosh\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(\frac{c(b+d)}{d} + (b + d)x\right)}{2b^2} \\
&\quad - \frac{c \operatorname{Chi}\left(\frac{c(b-d)}{d} + (b - d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2bd} + \frac{c \operatorname{Chi}\left(\frac{c(b+d)}{d} + (b + d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2bd} \\
&\quad - \frac{c \cosh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(\frac{c(b-d)}{d} + (b - d)x\right)}{2bd} - \frac{\sinh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(\frac{c(b-d)}{d} + (b - d)x\right)}{2b^2} \\
&\quad + \frac{x \cosh(a + bx) \operatorname{Shi}(c + dx)}{b} - \frac{\sinh(a + bx) \operatorname{Shi}(c + dx)}{b^2} \\
&\quad + \frac{c \cosh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(\frac{c(b+d)}{d} + (b + d)x\right)}{2bd} + \frac{\sinh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(\frac{c(b+d)}{d} + (b + d)x\right)}{2b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 2.61 (sec) , antiderivative size = 271, normalized size of antiderivative = 0.73

$$\int x \sinh(a + bx) \operatorname{Shi}(c + dx) dx$$

$$= \frac{e^{-a} \left(bde^{-c} \left(-\frac{e^{-((b+d)x}}{b+d} + \frac{e^{2a+bx-dx}}{b-d} \right) - (bc+d)e^{2a-\frac{bc}{d}} \operatorname{ExpIntegralEi}\left(\frac{(b-d)(c+dx)}{d}\right) - (bc-d)e^{\frac{bc}{d}} \operatorname{ExpIntegralEi}\left(-\frac{(b+d)(c+dx)}{d}\right) \right)}{d} - \frac{e^{-a} (bde^{-c} \dots)}{d}$$

[In] Integrate[x*Sinh[a + b*x]*SinhIntegral[c + d*x],x]

[Out] (((b*d*(-1/((b + d)*E^((b + d)*x))) + E^(2*a + b*x - d*x)/(b - d)))/E^c - (b*c + d)*E^(2*a - (b*c)/d)*ExpIntegralEi[((b - d)*(c + d*x))/d] - (b*c - d)*E^((b*c)/d)*ExpIntegralEi[-(((b + d)*(c + d*x))/d)])/(d*E^a) - (b*d*E^c*(E^((-b + d)*x)/(-b + d) + E^(2*a + (b + d)*x)/(b + d)) + (-b*c) + d)*E^((b*c)/d)*ExpIntegralEi[-(((b - d)*(c + d*x))/d)] - (b*c + d)*E^(2*a - (b*c)/d)*ExpIntegralEi[((b + d)*(c + d*x))/d])/(d*E^a) + 4*(b*x*Cosh[a + b*x] - Sinh[a + b*x])*SinhIntegral[c + d*x])/(4*b^2)

Maple [F]

$$\int x \operatorname{Shi}(dx + c) \sinh(bx + a) dx$$

[In] int(x*Shi(d*x+c)*sinh(b*x+a),x)

[Out] int(x*Shi(d*x+c)*sinh(b*x+a),x)

Fricas [F]

$$\int x \sinh(a + bx) \operatorname{Shi}(c + dx) dx = \int x \operatorname{Shi}(dx + c) \sinh(bx + a) dx$$

[In] `integrate(x*Shi(d*x+c)*sinh(b*x+a),x, algorithm="fricas")`

[Out] `integral(x*sinh(b*x + a)*sinh_integral(d*x + c), x)`

Sympy [F]

$$\int x \sinh(a + bx) \operatorname{Shi}(c + dx) dx = \int x \sinh(a + bx) \operatorname{Shi}(c + dx) dx$$

[In] `integrate(x*Shi(d*x+c)*sinh(b*x+a),x)`

[Out] `Integral(x*sinh(a + b*x)*Shi(c + d*x), x)`

Maxima [F]

$$\int x \sinh(a + bx) \operatorname{Shi}(c + dx) dx = \int x \operatorname{Shi}(dx + c) \sinh(bx + a) dx$$

[In] `integrate(x*Shi(d*x+c)*sinh(b*x+a),x, algorithm="maxima")`

[Out] `integrate(x*Shi(d*x + c)*sinh(b*x + a), x)`

Giac [F]

$$\int x \sinh(a + bx) \operatorname{Shi}(c + dx) dx = \int x \operatorname{Shi}(dx + c) \sinh(bx + a) dx$$

[In] `integrate(x*Shi(d*x+c)*sinh(b*x+a),x, algorithm="giac")`

[Out] `integrate(x*Shi(d*x + c)*sinh(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x \sinh(a + bx) \operatorname{Shi}(c + dx) dx = \int x \operatorname{sinhint}(c + dx) \sinh(a + bx) dx$$

```
[In] int(x*sinhint(c + d*x)*sinh(a + b*x),x)
```

```
[Out] int(x*sinhint(c + d*x)*sinh(a + b*x), x)
```

3.64 $\int \sinh(a + bx)\mathbf{Shi}(c + dx) dx$

Optimal result	341
Rubi [A] (verified)	341
Mathematica [A] (verified)	344
Maple [F]	344
Fricas [F]	344
Sympy [F]	344
Maxima [F]	345
Giac [F]	345
Mupad [F(-1)]	345

Optimal result

Integrand size = 13, antiderivative size = 153

$$\int \sinh(a + bx)\mathbf{Shi}(c + dx) dx = \frac{\mathbf{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b} - \frac{\mathbf{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b} + \frac{\cosh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b} + \frac{\cosh(a + bx)\mathbf{Shi}(c + dx)}{b} - \frac{\cosh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b}$$

```
[Out] 1/2*cosh(a-b*c/d)*Shi(c*(b-d)/d+(b-d)*x)/b+cosh(b*x+a)*Shi(d*x+c)/b-1/2*cosh(a-b*c/d)*Shi(c*(b+d)/d+(b+d)*x)/b+1/2*Chi(c*(b-d)/d+(b-d)*x)*sinh(a-b*c/d)/b-1/2*Chi(c*(b+d)/d+(b+d)*x)*sinh(a-b*c/d)/b
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used

= {6675, 5580, 3384, 3379, 3382}

$$\int \sinh(a + bx) \operatorname{Shi}(c + dx) dx = \frac{\sinh\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2b} - \frac{\sinh\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2b} + \frac{\cosh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2b} + \frac{\cosh(a + bx) \operatorname{Shi}(c + dx)}{b} - \frac{\cosh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2b}$$

[In] Int[Sinh[a + b*x]*SinhIntegral[c + d*x],x]

[Out] (CoshIntegral[(c*(b - d))/d + (b - d)*x]*Sinh[a - (b*c)/d])/(2*b) - (CoshIntegral[(c*(b + d))/d + (b + d)*x]*Sinh[a - (b*c)/d])/(2*b) + (Cosh[a - (b*c)/d]*SinhIntegral[(c*(b - d))/d + (b - d)*x])/(2*b) + (Cosh[a + b*x]*SinhIntegral[c + d*x])/b - (Cosh[a - (b*c)/d]*SinhIntegral[(c*(b + d))/d + (b + d)*x])/(2*b)

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3384

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]

Rule 5580

Int[Cosh[(c_.) + (d_.)*(x_)]^(q_.)*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(p_.), x_Symbol] := Int[ExpandTrigReduce[(e + f*x)^m, Sinh[a + b*x]^p*Cosh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[p

, 0] && IGtQ[q, 0]

Rule 6675

```
Int[Sinh[(a_.) + (b_.)*(x_.)]*SinhIntegral[(c_.) + (d_.)*(x_.)], x_Symbol] :>
  Simp[Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
  b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\cosh(a + bx)\text{Shi}(c + dx)}{b} - \frac{d \int \frac{\cosh(a+bx) \sinh(c+dx)}{c+dx} dx}{b} \\
 &= \frac{\cosh(a + bx)\text{Shi}(c + dx)}{b} - \frac{d \int \left(-\frac{\sinh(a-c+(b-d)x)}{2(c+dx)} + \frac{\sinh(a+c+(b+d)x)}{2(c+dx)} \right) dx}{b} \\
 &= \frac{\cosh(a + bx)\text{Shi}(c + dx)}{b} + \frac{d \int \frac{\sinh(a-c+(b-d)x)}{c+dx} dx}{2b} - \frac{d \int \frac{\sinh(a+c+(b+d)x)}{c+dx} dx}{2b} \\
 &= \frac{\cosh(a + bx)\text{Shi}(c + dx)}{b} + \frac{(d \cosh(a - \frac{bc}{d})) \int \frac{\sinh(\frac{c(b-d)}{d} + (b-d)x)}{c+dx} dx}{2b} \\
 &\quad - \frac{(d \cosh(a - \frac{bc}{d})) \int \frac{\sinh(\frac{c(b+d)}{d} + (b+d)x)}{c+dx} dx}{2b} \\
 &\quad + \frac{(d \sinh(a - \frac{bc}{d})) \int \frac{\cosh(\frac{c(b-d)}{d} + (b-d)x)}{c+dx} dx}{2b} \\
 &\quad - \frac{(d \sinh(a - \frac{bc}{d})) \int \frac{\cosh(\frac{c(b+d)}{d} + (b+d)x)}{c+dx} dx}{2b} \\
 &= \frac{\text{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b} - \frac{\text{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b} \\
 &\quad + \frac{\cosh\left(a - \frac{bc}{d}\right) \text{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b} \\
 &\quad + \frac{\cosh(a + bx)\text{Shi}(c + dx)}{b} - \frac{\cosh\left(a - \frac{bc}{d}\right) \text{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.68 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.90

$$\int \sinh(a + bx) \operatorname{Shi}(c + dx) dx$$

$$= \frac{e^{-a - \frac{bc}{d}} \left(-e^{\frac{2bc}{d}} \operatorname{ExpIntegralEi} \left(-\frac{(b-d)(c+dx)}{d} \right) + e^{2a} \operatorname{ExpIntegralEi} \left(\frac{(b-d)(c+dx)}{d} \right) + e^{\frac{2bc}{d}} \operatorname{ExpIntegralEi} \left(-\frac{(b+d)(c+dx)}{d} \right) \right)}{4b}$$

[In] Integrate[Sinh[a + b*x]*SinhIntegral[c + d*x],x]

[Out] (E^(-a - (b*c)/d)*(-E^((2*b*c)/d)*ExpIntegralEi[-(((b - d)*(c + d*x))/d)]) + E^(2*a)*ExpIntegralEi[((b - d)*(c + d*x))/d] + E^((2*b*c)/d)*ExpIntegralEi[-(((b + d)*(c + d*x))/d)] - E^(2*a)*ExpIntegralEi[((b + d)*(c + d*x))/d] + 4*E^(a + (b*c)/d)*Cosh[a + b*x]*SinhIntegral[c + d*x])/(4*b)

Maple [F]

$$\int \operatorname{Shi}(dx + c) \sinh(bx + a) dx$$

[In] int(Shi(d*x+c)*sinh(b*x+a),x)

[Out] int(Shi(d*x+c)*sinh(b*x+a),x)

Fricas [F]

$$\int \sinh(a + bx) \operatorname{Shi}(c + dx) dx = \int \operatorname{Shi}(dx + c) \sinh(bx + a) dx$$

[In] integrate(Shi(d*x+c)*sinh(b*x+a),x, algorithm="fricas")

[Out] integral(sinh(b*x + a)*sinh_integral(d*x + c), x)

Sympy [F]

$$\int \sinh(a + bx) \operatorname{Shi}(c + dx) dx = \int \sinh(a + bx) \operatorname{Shi}(c + dx) dx$$

[In] integrate(Shi(d*x+c)*sinh(b*x+a),x)

[Out] Integral(sinh(a + b*x)*Shi(c + d*x), x)

Maxima [F]

$$\int \sinh(a + bx)\text{Shi}(c + dx) dx = \int \text{Shi}(dx + c) \sinh(bx + a) dx$$

[In] integrate(Shi(d*x+c)*sinh(b*x+a),x, algorithm="maxima")

[Out] integrate(Shi(d*x + c)*sinh(b*x + a), x)

Giac [F]

$$\int \sinh(a + bx)\text{Shi}(c + dx) dx = \int \text{Shi}(dx + c) \sinh(bx + a) dx$$

[In] integrate(Shi(d*x+c)*sinh(b*x+a),x, algorithm="giac")

[Out] integrate(Shi(d*x + c)*sinh(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int \sinh(a + bx)\text{Shi}(c + dx) dx = \int \sinhint(c + dx) \sinh(a + bx) dx$$

[In] int(sinhint(c + d*x)*sinh(a + b*x),x)

[Out] int(sinhint(c + d*x)*sinh(a + b*x), x)

3.65 $\int \frac{\sinh(a+bx)\mathbf{Shi}(c+dx)}{x} dx$

Optimal result	346
Rubi [N/A]	346
Mathematica [N/A]	347
Maple [N/A] (verified)	347
Fricas [N/A]	347
Sympy [N/A]	347
Maxima [N/A]	348
Giac [N/A]	348
Mupad [N/A]	348

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\sinh(a+bx)\mathbf{Shi}(c+dx)}{x} dx = \text{Int}\left(\frac{\sinh(a+bx)\mathbf{Shi}(c+dx)}{x}, x\right)$$

[Out] CannotIntegrate(Shi(d*x+c)*sinh(b*x+a)/x,x)

Rubi [N/A]

Not integrable

Time = 0.10 (sec), antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\sinh(a+bx)\mathbf{Shi}(c+dx)}{x} dx = \int \frac{\sinh(a+bx)\mathbf{Shi}(c+dx)}{x} dx$$

[In] Int[(Sinh[a + b*x]*SinhIntegral[c + d*x])/x,x]

[Out] Defer[Int] [(Sinh[a + b*x]*SinhIntegral[c + d*x])/x, x]

Rubi steps

$$\text{integral} = \int \frac{\sinh(a+bx)\mathbf{Shi}(c+dx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 2.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sinh(a + bx)\text{Shi}(c + dx)}{x} dx = \int \frac{\sinh(a + bx)\text{Shi}(c + dx)}{x} dx$$

[In] Integrate[(Sinh[a + b*x]*SinhIntegral[c + d*x])/x,x]

[Out] Integrate[(Sinh[a + b*x]*SinhIntegral[c + d*x])/x, x]

Maple [N/A] (verified)

Not integrable

Time = 0.40 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\text{Shi}(dx + c) \sinh(bx + a)}{x} dx$$

[In] int(Shi(d*x+c)*sinh(b*x+a)/x,x)

[Out] int(Shi(d*x+c)*sinh(b*x+a)/x,x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sinh(a + bx)\text{Shi}(c + dx)}{x} dx = \int \frac{\text{Shi}(dx + c) \sinh(bx + a)}{x} dx$$

[In] integrate(Shi(d*x+c)*sinh(b*x+a)/x,x, algorithm="fricas")

[Out] integral(sinh(b*x + a)*sinh_integral(d*x + c)/x, x)

Sympy [N/A]

Not integrable

Time = 0.90 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{\sinh(a + bx)\text{Shi}(c + dx)}{x} dx = \int \frac{\sinh(a + bx)\text{Shi}(c + dx)}{x} dx$$

[In] integrate(Shi(d*x+c)*sinh(b*x+a)/x,x)

[Out] Integral(sinh(a + b*x)*Shi(c + d*x)/x, x)

Maxima [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sinh(a + bx)\text{Shi}(c + dx)}{x} dx = \int \frac{\text{Shi}(dx + c) \sinh(bx + a)}{x} dx$$

[In] integrate(Shi(d*x+c)*sinh(b*x+a)/x,x, algorithm="maxima")

[Out] integrate(Shi(d*x + c)*sinh(b*x + a)/x, x)

Giac [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sinh(a + bx)\text{Shi}(c + dx)}{x} dx = \int \frac{\text{Shi}(dx + c) \sinh(bx + a)}{x} dx$$

[In] integrate(Shi(d*x+c)*sinh(b*x+a)/x,x, algorithm="giac")

[Out] integrate(Shi(d*x + c)*sinh(b*x + a)/x, x)

Mupad [N/A]

Not integrable

Time = 5.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sinh(a + bx)\text{Shi}(c + dx)}{x} dx = \int \frac{\sinhint(c + dx) \sinh(a + bx)}{x} dx$$

[In] int((sinhint(c + d*x)*sinh(a + b*x))/x,x)

[Out] int((sinhint(c + d*x)*sinh(a + b*x))/x, x)

3.66 $\int x \cosh(a + bx) \mathbf{Shi}(c + dx) dx$

Optimal result	349
Rubi [A] (verified)	350
Mathematica [A] (verified)	354
Maple [F]	354
Fricas [F]	354
Sympy [F]	355
Maxima [F]	355
Giac [F]	355
Mupad [F(-1)]	355

Optimal result

Integrand size = 14, antiderivative size = 371

$$\begin{aligned}
 \int x \cosh(a + bx) \mathbf{Shi}(c + dx) dx = & -\frac{c \cosh\left(a - \frac{bc}{d}\right) \mathbf{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2bd} \\
 & + \frac{c \cosh\left(a - \frac{bc}{d}\right) \mathbf{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2bd} \\
 & - \frac{\mathbf{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b^2} \\
 & + \frac{\mathbf{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b^2} \\
 & + \frac{\sinh(a - c + (b-d)x)}{2b(b-d)} - \frac{\sinh(a + c + (b+d)x)}{2b(b+d)} \\
 & - \frac{\cosh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b^2} \\
 & - \frac{c \sinh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2bd} \\
 & - \frac{\cosh(a + bx) \mathbf{Shi}(c + dx)}{b^2} + \frac{x \sinh(a + bx) \mathbf{Shi}(c + dx)}{b} \\
 & + \frac{\cosh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b^2} \\
 & + \frac{c \sinh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2bd}
 \end{aligned}$$

[Out] $-1/2*c*\mathbf{Chi}(c*(b-d)/d+(b-d)*x)*\cosh(a-b*c/d)/b/d+1/2*c*\mathbf{Chi}(c*(b+d)/d+(b+d)*x)*\cosh(a-b*c/d)/b/d-1/2*\cosh(a-b*c/d)*\mathbf{Shi}(c*(b-d)/d+(b-d)*x)/b^2-\cosh(b*x+a$

) * Shi(d*x+c)/b^2+1/2*cosh(a-b*c/d)*Shi(c*(b+d)/d+(b+d)*x)/b^2-1/2*Chi(c*(b-d)/d+(b-d)*x)*sinh(a-b*c/d)/b^2+1/2*Chi(c*(b+d)/d+(b+d)*x)*sinh(a-b*c/d)/b^2-1/2*c*Shi(c*(b-d)/d+(b-d)*x)*sinh(a-b*c/d)/b/d+1/2*c*Shi(c*(b+d)/d+(b+d)*x)*sinh(a-b*c/d)/b/d+x*Shi(d*x+c)*sinh(b*x+a)/b+1/2*sinh(a-c+(b-d)*x)/b/(b-d)-1/2*sinh(a+c+(b+d)*x)/b/(b+d)

Rubi [A] (verified)

Time = 0.96 (sec) , antiderivative size = 371, normalized size of antiderivative = 1.00, number of steps used = 24, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {6683, 5761, 6874, 2717, 3384, 3379, 3382, 6675, 5580}

$$\int x \cosh(a + bx) \text{Shi}(c + dx) dx = -\frac{\sinh\left(a - \frac{bc}{d}\right) \text{Chi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2b^2} + \frac{\sinh\left(a - \frac{bc}{d}\right) \text{Chi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2b^2} - \frac{\cosh\left(a - \frac{bc}{d}\right) \text{Shi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2b^2} - \frac{\cosh(a + bx) \text{Shi}(c + dx)}{b^2} + \frac{\cosh\left(a - \frac{bc}{d}\right) \text{Shi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2b^2} - \frac{c \cosh\left(a - \frac{bc}{d}\right) \text{Chi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2bd} + \frac{c \cosh\left(a - \frac{bc}{d}\right) \text{Chi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2bd} - \frac{c \sinh\left(a - \frac{bc}{d}\right) \text{Shi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2bd} + \frac{x \sinh(a + bx) \text{Shi}(c + dx)}{b} + \frac{c \sinh\left(a - \frac{bc}{d}\right) \text{Shi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2bd} + \frac{\sinh(a + x(b-d) - c)}{2b(b-d)} - \frac{\sinh(a + x(b+d) + c)}{2b(b+d)}$$

[In] Int[x*Cosh[a + b*x]*SinhIntegral[c + d*x],x]

[Out] -1/2*(c*Cosh[a - (b*c)/d]*CoshIntegral[(c*(b - d))/d + (b - d)*x])/(b*d) + (c*Cosh[a - (b*c)/d]*CoshIntegral[(c*(b + d))/d + (b + d)*x])/(2*b*d) - (CoshIntegral[(c*(b - d))/d + (b - d)*x]*Sinh[a - (b*c)/d])/(2*b^2) + (CoshInt

```

egral[(c*(b + d))/d + (b + d)*x]*Sinh[a - (b*c)/d]/(2*b^2) + Sinh[a - c +
(b - d)*x]/(2*b*(b - d)) - Sinh[a + c + (b + d)*x]/(2*b*(b + d)) - (Cosh[a
- (b*c)/d]*SinhIntegral[(c*(b - d))/d + (b - d)*x]/(2*b^2) - (c*Sinh[a - (
b*c)/d]*SinhIntegral[(c*(b - d))/d + (b - d)*x])/(2*b*d) - (Cosh[a + b*x]*S
inhIntegral[c + d*x])/b^2 + (x*Sinh[a + b*x]*SinhIntegral[c + d*x])/b + (Co
sh[a - (b*c)/d]*SinhIntegral[(c*(b + d))/d + (b + d)*x]/(2*b^2) + (c*Sinh[
a - (b*c)/d]*SinhIntegral[(c*(b + d))/d + (b + d)*x])/(2*b*d)

```

Rule 2717

```

Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]

```

Rule 3379

```

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol]
:= Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f
, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

```

Rule 3382

```

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol]
:= Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz
}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

```

Rule 3384

```

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*
e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f
)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] &&
NeQ[d*e - c*f, 0]

```

Rule 5580

```

Int[Cosh[(c_.) + (d_.)*(x_)]^(q_.)*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(p_.), x_Symbol] := Int[ExpandTrigReduce[(e + f*x)^m, Sinh[a +
b*x]^p*Cosh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[p
, 0] && IGtQ[q, 0]

```

Rule 5761

```

Int[(u_.)*Sinh[(a_.) + (b_.)*(x_)]^(m_.)*Sinh[(c_.) + (d_.)*(x_)]^(n_.), x_
Symbol] := Int[ExpandTrigReduce[u, Sinh[a + b*x]^m*Sinh[c + d*x]^n, x], x]
/; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[n, 0]

```

Rule 6675

```
Int[Sinh[(a_.) + (b_.)*(x_)]*SinhIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
  Simp[Cosh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
  b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6683

```
Int[Cosh[(a_.) + (b_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.)*SinhIntegral[(c_.)
+ (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Sinh[a + b*x]*(SinhIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Sinh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*SinhIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x \sinh(a + bx) \text{Shi}(c + dx)}{b} - \frac{\int \sinh(a + bx) \text{Shi}(c + dx) dx}{b} \\
 &\quad - \frac{d \int \frac{x \sinh(a+bx) \sinh(c+dx)}{c+dx} dx}{b} \\
 &= -\frac{\cosh(a + bx) \text{Shi}(c + dx)}{b^2} + \frac{x \sinh(a + bx) \text{Shi}(c + dx)}{b} \\
 &\quad + \frac{d \int \frac{\cosh(a+bx) \sinh(c+dx)}{c+dx} dx}{b^2} - \frac{d \int \left(-\frac{x \cosh(a-c+(b-d)x)}{2(c+dx)} + \frac{x \cosh(a+c+(b+d)x)}{2(c+dx)} \right) dx}{b} \\
 &= -\frac{\cosh(a + bx) \text{Shi}(c + dx)}{b^2} + \frac{x \sinh(a + bx) \text{Shi}(c + dx)}{b} \\
 &\quad + \frac{d \int \left(-\frac{\sinh(a-c+(b-d)x)}{2(c+dx)} + \frac{\sinh(a+c+(b+d)x)}{2(c+dx)} \right) dx}{b^2} \\
 &\quad + \frac{d \int \frac{x \cosh(a-c+(b-d)x)}{c+dx} dx}{2b} - \frac{d \int \frac{x \cosh(a+c+(b+d)x)}{c+dx} dx}{2b} \\
 &= -\frac{\cosh(a + bx) \text{Shi}(c + dx)}{b^2} + \frac{x \sinh(a + bx) \text{Shi}(c + dx)}{b} - \frac{d \int \frac{\sinh(a-c+(b-d)x)}{c+dx} dx}{2b^2} \\
 &\quad + \frac{d \int \frac{\sinh(a+c+(b+d)x)}{c+dx} dx}{2b^2} + \frac{d \int \left(\frac{\cosh(a-c+(b-d)x)}{d} - \frac{c \cosh(a-c+(b-d)x)}{d(c+dx)} \right) dx}{2b} \\
 &\quad - \frac{d \int \left(\frac{\cosh(a+c+(b+d)x)}{d} - \frac{c \cosh(a+c+(b+d)x)}{d(c+dx)} \right) dx}{2b}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{\cosh(a+bx)\text{Shi}(c+dx)}{b^2} + \frac{x \sinh(a+bx)\text{Shi}(c+dx)}{b} + \frac{\int \cosh(a-c+(b-d)x) dx}{2b} \\
&\quad - \frac{\int \cosh(a+c+(b+d)x) dx}{2b} - \frac{c \int \frac{\cosh(a-c+(b-d)x)}{c+dx} dx}{2b} + \frac{c \int \frac{\cosh(a+c+(b+d)x)}{c+dx} dx}{2b} \\
&\quad - \frac{(d \cosh(a-\frac{bc}{d})) \int \frac{\sinh(\frac{c(b-d)}{d}+(b-d)x)}{c+dx} dx}{2b^2} + \frac{(d \cosh(a-\frac{bc}{d})) \int \frac{\sinh(\frac{c(b+d)}{d}+(b+d)x)}{c+dx} dx}{2b^2} \\
&\quad - \frac{(d \sinh(a-\frac{bc}{d})) \int \frac{\cosh(\frac{c(b-d)}{d}+(b-d)x)}{c+dx} dx}{2b^2} + \frac{(d \sinh(a-\frac{bc}{d})) \int \frac{\cosh(\frac{c(b+d)}{d}+(b+d)x)}{c+dx} dx}{2b^2} \\
&= -\frac{\text{Chi}\left(\frac{c(b-d)}{d}+(b-d)x\right) \sinh(a-\frac{bc}{d})}{2b^2} + \frac{\text{Chi}\left(\frac{c(b+d)}{d}+(b+d)x\right) \sinh(a-\frac{bc}{d})}{2b^2} \\
&\quad + \frac{\sinh(a-c+(b-d)x)}{2b(b-d)} - \frac{\sinh(a+c+(b+d)x)}{2b(b+d)} \\
&\quad - \frac{\cosh(a-\frac{bc}{d}) \text{Shi}\left(\frac{c(b-d)}{d}+(b-d)x\right)}{2b^2} - \frac{\cosh(a+bx)\text{Shi}(c+dx)}{b^2} \\
&\quad + \frac{x \sinh(a+bx)\text{Shi}(c+dx)}{b} + \frac{\cosh(a-\frac{bc}{d}) \text{Shi}\left(\frac{c(b+d)}{d}+(b+d)x\right)}{2b^2} \\
&\quad - \frac{(c \cosh(a-\frac{bc}{d})) \int \frac{\cosh(\frac{c(b-d)}{d}+(b-d)x)}{c+dx} dx}{2b} \\
&\quad + \frac{(c \cosh(a-\frac{bc}{d})) \int \frac{\cosh(\frac{c(b+d)}{d}+(b+d)x)}{c+dx} dx}{2b} \\
&\quad - \frac{(c \sinh(a-\frac{bc}{d})) \int \frac{\sinh(\frac{c(b-d)}{d}+(b-d)x)}{c+dx} dx}{2b} \\
&\quad + \frac{(c \sinh(a-\frac{bc}{d})) \int \frac{\sinh(\frac{c(b+d)}{d}+(b+d)x)}{c+dx} dx}{2b} \\
&= -\frac{c \cosh(a-\frac{bc}{d}) \text{Chi}\left(\frac{c(b-d)}{d}+(b-d)x\right)}{2bd} + \frac{c \cosh(a-\frac{bc}{d}) \text{Chi}\left(\frac{c(b+d)}{d}+(b+d)x\right)}{2bd} \\
&\quad - \frac{\text{Chi}\left(\frac{c(b-d)}{d}+(b-d)x\right) \sinh(a-\frac{bc}{d})}{2b^2} + \frac{\text{Chi}\left(\frac{c(b+d)}{d}+(b+d)x\right) \sinh(a-\frac{bc}{d})}{2b^2} \\
&\quad + \frac{\sinh(a-c+(b-d)x)}{2b(b-d)} - \frac{\sinh(a+c+(b+d)x)}{2b(b+d)} \\
&\quad - \frac{\cosh(a-\frac{bc}{d}) \text{Shi}\left(\frac{c(b-d)}{d}+(b-d)x\right)}{2b^2} - \frac{c \sinh(a-\frac{bc}{d}) \text{Shi}\left(\frac{c(b-d)}{d}+(b-d)x\right)}{2bd} \\
&\quad - \frac{\cosh(a+bx)\text{Shi}(c+dx)}{b^2} + \frac{x \sinh(a+bx)\text{Shi}(c+dx)}{b} \\
&\quad + \frac{\cosh(a-\frac{bc}{d}) \text{Shi}\left(\frac{c(b+d)}{d}+(b+d)x\right)}{2b^2} + \frac{c \sinh(a-\frac{bc}{d}) \text{Shi}\left(\frac{c(b+d)}{d}+(b+d)x\right)}{2bd}
\end{aligned}$$

Mathematica [A] (verified)

Time = 3.14 (sec) , antiderivative size = 313, normalized size of antiderivative = 0.84

$$\int x \cosh(a + bx) \operatorname{Shi}(c + dx) dx$$

$$= \frac{e^{-a - \frac{(b+d)(c+dx)}{d}} \left((bc+d)(b^2-d^2)e^{2a+c+(b+d)x} \operatorname{ExpIntegralEi}\left(\frac{(b-d)(c+dx)}{d}\right) - e^{\frac{bc}{d}} \left(bd(d(-1+e^{2(a+bx)})+b(1+e^{2(a+bx)})) + (bc-d)(b^2-d^2)e^{\frac{(b+d)(c+dx)}{d}} \right) \right)}{d(-b+d)(b+d)}$$

[In] Integrate[x*Cosh[a + b*x]*SinhIntegral[c + d*x], x]

[Out] ((E^(-a - ((b + d)*(c + d*x))/d))*((b*c + d)*(b^2 - d^2)*E^(2*a + c + (b + d)*x))*ExpIntegralEi[((b - d)*(c + d*x))/d] - E^((b*c)/d)*(b*d*(d*(-1 + E^(2*(a + b*x)))) + b*(1 + E^(2*(a + b*x)))) + (b*c - d)*(b^2 - d^2)*E^(((b + d)*(c + d*x))/d)*ExpIntegralEi[-(((b + d)*(c + d*x))/d)])))/(d*(-b + d)*(b + d)) + (b*d*E^c*(E^((-b + d)*x)/(-b + d) - E^(2*a + (b + d)*x)/(b + d)) + (-b*c + d)*E^((b*c)/d)*ExpIntegralEi[-(((b - d)*(c + d*x))/d)] + (b*c + d)*E^(2*a - (b*c)/d)*ExpIntegralEi[((b + d)*(c + d*x))/d])/(d*E^a) + 4*(-Cosh[a + b*x] + b*x*Sinh[a + b*x])*SinhIntegral[c + d*x])/(4*b^2)

Maple [F]

$$\int x \cosh(bx + a) \operatorname{Shi}(dx + c) dx$$

[In] int(x*cosh(b*x+a)*Shi(d*x+c), x)

[Out] int(x*cosh(b*x+a)*Shi(d*x+c), x)

Fricas [F]

$$\int x \cosh(a + bx) \operatorname{Shi}(c + dx) dx = \int x \operatorname{Shi}(dx + c) \cosh(bx + a) dx$$

[In] integrate(x*cosh(b*x+a)*Shi(d*x+c), x, algorithm="fricas")

[Out] integral(x*cosh(b*x + a)*sinh_integral(d*x + c), x)

Sympy [F]

$$\int x \cosh(a + bx) \operatorname{Shi}(c + dx) dx = \int x \cosh(a + bx) \operatorname{Shi}(c + dx) dx$$

[In] `integrate(x*cosh(b*x+a)*Shi(d*x+c),x)`

[Out] `Integral(x*cosh(a + b*x)*Shi(c + d*x), x)`

Maxima [F]

$$\int x \cosh(a + bx) \operatorname{Shi}(c + dx) dx = \int x \operatorname{Shi}(dx + c) \cosh(bx + a) dx$$

[In] `integrate(x*cosh(b*x+a)*Shi(d*x+c),x, algorithm="maxima")`

[Out] `integrate(x*Shi(d*x + c)*cosh(b*x + a), x)`

Giac [F]

$$\int x \cosh(a + bx) \operatorname{Shi}(c + dx) dx = \int x \operatorname{Shi}(dx + c) \cosh(bx + a) dx$$

[In] `integrate(x*cosh(b*x+a)*Shi(d*x+c),x, algorithm="giac")`

[Out] `integrate(x*Shi(d*x + c)*cosh(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x \cosh(a + bx) \operatorname{Shi}(c + dx) dx = \int x \operatorname{sinhint}(c + dx) \cosh(a + bx) dx$$

[In] `int(x*sinhint(c + d*x)*cosh(a + b*x),x)`

[Out] `int(x*sinhint(c + d*x)*cosh(a + b*x), x)`

3.67 $\int \cosh(a + bx)\text{Shi}(c + dx) dx$

Optimal result	356
Rubi [A] (verified)	356
Mathematica [A] (verified)	359
Maple [F]	359
Fricas [F]	359
Sympy [F]	359
Maxima [F]	360
Giac [F]	360
Mupad [F(-1)]	360

Optimal result

Integrand size = 13, antiderivative size = 153

$$\int \cosh(a + bx)\text{Shi}(c + dx) dx = \frac{\cosh\left(a - \frac{bc}{d}\right) \text{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b} - \frac{\cosh\left(a - \frac{bc}{d}\right) \text{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b} + \frac{\sinh\left(a - \frac{bc}{d}\right) \text{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b} + \frac{\sinh(a + bx)\text{Shi}(c + dx)}{b} - \frac{\sinh\left(a - \frac{bc}{d}\right) \text{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b}$$

[Out] 1/2*Chi(c*(b-d)/d+(b-d)*x)*cosh(a-b*c/d)/b-1/2*Chi(c*(b+d)/d+(b+d)*x)*cosh(a-b*c/d)/b+1/2*Shi(c*(b-d)/d+(b-d)*x)*sinh(a-b*c/d)/b-1/2*Shi(c*(b+d)/d+(b+d)*x)*sinh(a-b*c/d)/b+Shi(d*x+c)*sinh(b*x+a)/b

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used

= {6681, 5578, 3384, 3379, 3382}

$$\int \cosh(a + bx) \operatorname{Shi}(c + dx) dx = \frac{\cosh\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2b} - \frac{\cosh\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2b} + \frac{\sinh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2b} + \frac{\sinh(a + bx) \operatorname{Shi}(c + dx)}{b} - \frac{\sinh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2b}$$

[In] Int[Cosh[a + b*x]*SinhIntegral[c + d*x], x]

[Out] (Cosh[a - (b*c)/d]*CoshIntegral[(c*(b - d))/d + (b - d)*x])/(2*b) - (Cosh[a - (b*c)/d]*CoshIntegral[(c*(b + d))/d + (b + d)*x])/(2*b) + (Sinh[a - (b*c)/d]*SinhIntegral[(c*(b - d))/d + (b - d)*x])/(2*b) + (Sinh[a + b*x]*SinhIntegral[c + d*x])/b - (Sinh[a - (b*c)/d]*SinhIntegral[(c*(b + d))/d + (b + d)*x])/(2*b)

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3384

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]

Rule 5578

Int[((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(p_.)*Sinh[(c_.) + (d_.)*(x_)]^(q_.), x_Symbol] := Int[ExpandTrigReduce[(e + f*x)^m, Sinh[a + b*x]^p*Sinh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0]

] && IGtQ[q, 0] && IntegerQ[m]

Rule 6681

```
Int[Cosh[(a_.) + (b_.)*(x_.)]*SinhIntegral[(c_.) + (d_.)*(x_.)], x_Symbol] :>
  Simp[Sinh[a + b*x]*(SinhIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
  b*x]*(Sinh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\sinh(a + bx)\text{Shi}(c + dx)}{b} - \frac{d \int \frac{\sinh(a+bx)\sinh(c+dx)}{c+dx} dx}{b} \\
 &= \frac{\sinh(a + bx)\text{Shi}(c + dx)}{b} - \frac{d \int \left(-\frac{\cosh(a-c+(b-d)x)}{2(c+dx)} + \frac{\cosh(a+c+(b+d)x)}{2(c+dx)} \right) dx}{b} \\
 &= \frac{\sinh(a + bx)\text{Shi}(c + dx)}{b} + \frac{d \int \frac{\cosh(a-c+(b-d)x)}{c+dx} dx}{2b} - \frac{d \int \frac{\cosh(a+c+(b+d)x)}{c+dx} dx}{2b} \\
 &= \frac{\sinh(a + bx)\text{Shi}(c + dx)}{b} + \frac{(d \cosh(a - \frac{bc}{d})) \int \frac{\cosh(\frac{c(b-d)}{d} + (b-d)x)}{c+dx} dx}{2b} \\
 &\quad - \frac{(d \cosh(a - \frac{bc}{d})) \int \frac{\cosh(\frac{c(b+d)}{d} + (b+d)x)}{c+dx} dx}{2b} \\
 &\quad + \frac{(d \sinh(a - \frac{bc}{d})) \int \frac{\sinh(\frac{c(b-d)}{d} + (b-d)x)}{c+dx} dx}{2b} \\
 &\quad - \frac{(d \sinh(a - \frac{bc}{d})) \int \frac{\sinh(\frac{c(b+d)}{d} + (b+d)x)}{c+dx} dx}{2b} \\
 &= \frac{\cosh(a - \frac{bc}{d}) \text{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b} - \frac{\cosh(a - \frac{bc}{d}) \text{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b} \\
 &\quad + \frac{\sinh(a - \frac{bc}{d}) \text{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b} \\
 &\quad + \frac{\sinh(a + bx)\text{Shi}(c + dx)}{b} - \frac{\sinh(a - \frac{bc}{d}) \text{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.73 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.90

$$\int \cosh(a + bx) \operatorname{Shi}(c + dx) dx$$

$$= \frac{e^{-a - \frac{bc}{d}} \left(e^{\frac{2bc}{d}} \operatorname{ExpIntegralEi} \left(-\frac{(b-d)(c+dx)}{d} \right) + e^{2a} \operatorname{ExpIntegralEi} \left(\frac{(b-d)(c+dx)}{d} \right) - e^{\frac{2bc}{d}} \operatorname{ExpIntegralEi} \left(-\frac{(b+d)(c+dx)}{d} \right) \right)}{4b}$$

[In] Integrate[Cosh[a + b*x]*SinhIntegral[c + d*x],x]

[Out] (E^(-a - (b*c)/d)*(E^((2*b*c)/d)*ExpIntegralEi[-((b - d)*(c + d*x))/d]) + E^(2*a)*ExpIntegralEi[((b - d)*(c + d*x))/d] - E^((2*b*c)/d)*ExpIntegralEi[-((b + d)*(c + d*x))/d]) - E^(2*a)*ExpIntegralEi[((b + d)*(c + d*x))/d] + 4*E^(a + (b*c)/d)*Sinh[a + b*x]*SinhIntegral[c + d*x])/(4*b)

Maple [F]

$$\int \cosh(bx + a) \operatorname{Shi}(dx + c) dx$$

[In] int(cosh(b*x+a)*Shi(d*x+c),x)

[Out] int(cosh(b*x+a)*Shi(d*x+c),x)

Fricas [F]

$$\int \cosh(a + bx) \operatorname{Shi}(c + dx) dx = \int \operatorname{Shi}(dx + c) \cosh(bx + a) dx$$

[In] integrate(cosh(b*x+a)*Shi(d*x+c),x, algorithm="fricas")

[Out] integral(cosh(b*x + a)*sinh_integral(d*x + c), x)

Sympy [F]

$$\int \cosh(a + bx) \operatorname{Shi}(c + dx) dx = \int \cosh(a + bx) \operatorname{Shi}(c + dx) dx$$

[In] integrate(cosh(b*x+a)*Shi(d*x+c),x)

[Out] Integral(cosh(a + b*x)*Shi(c + d*x), x)

Maxima [F]

$$\int \cosh(a + bx)\text{Shi}(c + dx) dx = \int \text{Shi}(dx + c) \cosh(bx + a) dx$$

[In] integrate(cosh(b*x+a)*Shi(d*x+c),x, algorithm="maxima")

[Out] integrate(Shi(d*x + c)*cosh(b*x + a), x)

Giac [F]

$$\int \cosh(a + bx)\text{Shi}(c + dx) dx = \int \text{Shi}(dx + c) \cosh(bx + a) dx$$

[In] integrate(cosh(b*x+a)*Shi(d*x+c),x, algorithm="giac")

[Out] integrate(Shi(d*x + c)*cosh(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int \cosh(a + bx)\text{Shi}(c + dx) dx = \int \sinhint(c + dx) \cosh(a + bx) dx$$

[In] int(sinhint(c + d*x)*cosh(a + b*x),x)

[Out] int(sinhint(c + d*x)*cosh(a + b*x), x)

3.68 $\int \frac{\cosh(ax+bx)\mathbf{Shi}(c+dx)}{x} dx$

Optimal result	361
Rubi [N/A]	361
Mathematica [N/A]	362
Maple [N/A] (verified)	362
Fricas [N/A]	362
Sympy [N/A]	362
Maxima [N/A]	363
Giac [N/A]	363
Mupad [N/A]	363

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\cosh(a+bx)\mathbf{Shi}(c+dx)}{x} dx = \text{Int}\left(\frac{\cosh(a+bx)\mathbf{Shi}(c+dx)}{x}, x\right)$$

[Out] `CannotIntegrate(cosh(b*x+a)*Shi(d*x+c)/x,x)`

Rubi [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\cosh(a+bx)\mathbf{Shi}(c+dx)}{x} dx = \int \frac{\cosh(a+bx)\mathbf{Shi}(c+dx)}{x} dx$$

[In] `Int[(Cosh[a + b*x]*SinhIntegral[c + d*x])/x,x]`

[Out] `Defer[Int] [(Cosh[a + b*x]*SinhIntegral[c + d*x])/x, x]`

Rubi steps

$$\text{integral} = \int \frac{\cosh(a+bx)\mathbf{Shi}(c+dx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 1.83 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\operatorname{Shi}(c + dx)}{x} dx = \int \frac{\cosh(a + bx)\operatorname{Shi}(c + dx)}{x} dx$$

[In] Integrate[(Cosh[a + b*x]*SinhIntegral[c + d*x])/x,x]

[Out] Integrate[(Cosh[a + b*x]*SinhIntegral[c + d*x])/x, x]

Maple [N/A] (verified)

Not integrable

Time = 0.43 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\cosh(bx + a)\operatorname{Shi}(dx + c)}{x} dx$$

[In] int(cosh(b*x+a)*Shi(d*x+c)/x,x)

[Out] int(cosh(b*x+a)*Shi(d*x+c)/x,x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\operatorname{Shi}(c + dx)}{x} dx = \int \frac{\operatorname{Shi}(dx + c)\cosh(bx + a)}{x} dx$$

[In] integrate(cosh(b*x+a)*Shi(d*x+c)/x,x, algorithm="fricas")

[Out] integral(cosh(b*x + a)*sinh_integral(d*x + c)/x, x)

Sympy [N/A]

Not integrable

Time = 0.88 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{\cosh(a + bx)\operatorname{Shi}(c + dx)}{x} dx = \int \frac{\cosh(a + bx)\operatorname{Shi}(c + dx)}{x} dx$$

[In] integrate(cosh(b*x+a)*Shi(d*x+c)/x,x)

[Out] Integral(cosh(a + b*x)*Shi(c + d*x)/x, x)

Maxima [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Shi}(c + dx)}{x} dx = \int \frac{\text{Shi}(dx + c) \cosh(bx + a)}{x} dx$$

[In] integrate(cosh(b*x+a)*Shi(d*x+c)/x,x, algorithm="maxima")

[Out] integrate(Shi(d*x + c)*cosh(b*x + a)/x, x)

Giac [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Shi}(c + dx)}{x} dx = \int \frac{\text{Shi}(dx + c) \cosh(bx + a)}{x} dx$$

[In] integrate(cosh(b*x+a)*Shi(d*x+c)/x,x, algorithm="giac")

[Out] integrate(Shi(d*x + c)*cosh(b*x + a)/x, x)

Mupad [N/A]

Not integrable

Time = 4.87 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Shi}(c + dx)}{x} dx = \int \frac{\sinhint(c + dx) \cosh(a + bx)}{x} dx$$

[In] int((sinhint(c + d*x)*cosh(a + b*x))/x,x)

[Out] int((sinhint(c + d*x)*cosh(a + b*x))/x, x)

3.69 $\int x^m \mathbf{Chi}(bx) dx$

Optimal result	364
Rubi [A] (verified)	364
Mathematica [A] (verified)	365
Maple [F]	366
Fricas [F]	366
Sympy [B] (verification not implemented)	366
Maxima [F]	368
Giac [F]	368
Mupad [F(-1)]	368

Optimal result

Integrand size = 8, antiderivative size = 76

$$\int x^m \mathbf{Chi}(bx) dx = \frac{x^{1+m} \mathbf{Chi}(bx)}{1+m} - \frac{x^m (-bx)^{-m} \Gamma(1+m, -bx)}{2b(1+m)} + \frac{x^m (bx)^{-m} \Gamma(1+m, bx)}{2b(1+m)}$$

[Out] $x^{(1+m)} * \mathbf{Chi}(b*x) / (1+m) - 1/2 * x^m * \mathbf{GAMMA}(1+m, -b*x) / b / (1+m) / ((-b*x)^m) + 1/2 * x^m * \mathbf{GAMMA}(1+m, b*x) / b / (1+m) / ((b*x)^m)$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6668, 12, 3388, 2212}

$$\int x^m \mathbf{Chi}(bx) dx = \frac{x^{m+1} \mathbf{Chi}(bx)}{m+1} - \frac{x^m (-bx)^{-m} \Gamma(m+1, -bx)}{2b(m+1)} + \frac{x^m (bx)^{-m} \Gamma(m+1, bx)}{2b(m+1)}$$

[In] `Int[x^m*CoshIntegral[b*x],x]`

[Out] $(x^{(1+m)} * \mathbf{CoshIntegral}[b*x]) / (1+m) - (x^m * \mathbf{Gamma}[1+m, -(b*x)]) / (2*b*(1+m)*(-b*x)^m) + (x^m * \mathbf{Gamma}[1+m, b*x]) / (2*b*(1+m)*(b*x)^m)$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 2212

`Int[(F_)^((g_.)*((e_.) + (f_.)*(x_))) * ((c_.) + (d_.)*(x_))^(m_), x_Symbol] := Simp[(-F^(g*(e - c*(f/d)))) * ((c + d*x)^FracPart[m] / (d * ((-f)*g*(Log[F]/d)))]`

```
)^(IntPart[m] + 1)*((-f)*g*Log[F]*((c + d*x)/d))^FracPart[m])*Gamma[m + 1,
((-f)*g*(Log[F]/d))*(c + d*x)], x] /; FreeQ[{F, c, d, e, f, g, m}, x] &&
!IntegerQ[m]
```

Rule 3388

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + Pi*(k_.) + (f_.)*(x_)], x_Symbol]
:> Dist[I/2, Int[(c + d*x)^m/(E^(I*k*Pi)*E^(I*(e + f*x))), x], x] - Dist[
I/2, Int[(c + d*x)^m*E^(I*k*Pi)*E^(I*(e + f*x)), x], x] /; FreeQ[{c, d, e,
f, m}, x] && IntegerQ[2*k]
```

Rule 6668

```
Int[CoshIntegral[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol]
:> Simp[(c + d*x)^(m + 1)*(CoshIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/
(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Cosh[a + b*x]/(a + b*x)), x], x] /; Fre
eQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x^{1+m}\text{Chi}(bx)}{1+m} - \frac{b \int \frac{x^m \cosh(bx)}{b} dx}{1+m} \\
 &= \frac{x^{1+m}\text{Chi}(bx)}{1+m} - \frac{\int x^m \cosh(bx) dx}{1+m} \\
 &= \frac{x^{1+m}\text{Chi}(bx)}{1+m} - \frac{\int e^{-bx} x^m dx}{2(1+m)} - \frac{\int e^{bx} x^m dx}{2(1+m)} \\
 &= \frac{x^{1+m}\text{Chi}(bx)}{1+m} - \frac{x^m(-bx)^{-m}\Gamma(1+m, -bx)}{2b(1+m)} + \frac{x^m(bx)^{-m}\Gamma(1+m, bx)}{2b(1+m)}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.87

$$\int x^m \text{Chi}(bx) dx = \frac{x^m \left(2x \text{Chi}(bx) + \frac{(-b^2 x^2)^{-m} (-bx)^m \Gamma(1+m, -bx) + (-bx)^m \Gamma(1+m, bx)}{b} \right)}{2(1+m)}$$

```
[In] Integrate[x^m*CoshIntegral[b*x], x]
```

```
[Out] (x^m*(2*x*CoshIntegral[b*x] + (-((b*x)^m*Gamma[1 + m, -(b*x)])) + (-b*x))^m
*Gamma[1 + m, b*x])/(b*(-b^2*x^2)^m)/(2*(1 + m))
```

Maple [F]

$$\int x^m \operatorname{Chi}(bx) dx$$

[In] `int(x^m*Chi(b*x),x)`

[Out] `int(x^m*Chi(b*x),x)`

Fricas [F]

$$\int x^m \operatorname{Chi}(bx) dx = \int x^m \operatorname{Chi}(bx) dx$$

[In] `integrate(x^m*Chi(b*x),x, algorithm="fricas")`

[Out] `integral(x^m*cosh_integral(b*x), x)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 695 vs. $2(60) = 120$.

Time = 1.27 (sec) , antiderivative size = 695, normalized size of antiderivative = 9.14

$$\begin{aligned}
 \int x^m \text{Chi}(bx) dx = & \frac{4 \cdot 2^m b b^{-m-1} m x \sqrt{e^{-2m \log(2)} e^{m \log(b^2 x^2)}} \log(b^2 x^2) \Gamma\left(\frac{m}{2} + \frac{5}{2}\right)}{8m^2 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 16m \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 8 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right)} \\
 & + \frac{8 \cdot 2^m \gamma b b^{-m-1} m x \sqrt{e^{-2m \log(2)} e^{m \log(b^2 x^2)}} \Gamma\left(\frac{m}{2} + \frac{5}{2}\right)}{8m^2 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 16m \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 8 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right)} \\
 & + \frac{4 \cdot 2^m b b^{-m-1} x \sqrt{e^{-2m \log(2)} e^{m \log(b^2 x^2)}} \log(b^2 x^2) \Gamma\left(\frac{m}{2} + \frac{5}{2}\right)}{8m^2 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 16m \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 8 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right)} \\
 & - \frac{8 \cdot 2^m b b^{-m-1} x \sqrt{e^{-2m \log(2)} e^{m \log(b^2 x^2)}} \Gamma\left(\frac{m}{2} + \frac{5}{2}\right)}{8m^2 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 16m \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 8 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right)} \\
 & + \frac{8 \cdot 2^m \gamma b b^{-m-1} x \sqrt{e^{-2m \log(2)} e^{m \log(b^2 x^2)}} \Gamma\left(\frac{m}{2} + \frac{5}{2}\right)}{8m^2 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 16m \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 8 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right)} \\
 & + \frac{b^{-m-1} b^{m+3} m^2 x^{m+3} \Gamma\left(\frac{m}{2} + \frac{3}{2}\right) {}_3F_4\left(\begin{matrix} 1, 1, \frac{m}{2} + \frac{3}{2} \\ \frac{3}{2}, 2, 2, \frac{m}{2} + \frac{5}{2} \end{matrix} \middle| \frac{b^2 x^2}{4}\right)}{8m^2 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 16m \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 8 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right)} \\
 & + \frac{2b^{-m-1} b^{m+3} m x^{m+3} \Gamma\left(\frac{m}{2} + \frac{3}{2}\right) {}_3F_4\left(\begin{matrix} 1, 1, \frac{m}{2} + \frac{3}{2} \\ \frac{3}{2}, 2, 2, \frac{m}{2} + \frac{5}{2} \end{matrix} \middle| \frac{b^2 x^2}{4}\right)}{8m^2 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 16m \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 8 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right)} \\
 & + \frac{b^{-m-1} b^{m+3} x^{m+3} \Gamma\left(\frac{m}{2} + \frac{3}{2}\right) {}_3F_4\left(\begin{matrix} 1, 1, \frac{m}{2} + \frac{3}{2} \\ \frac{3}{2}, 2, 2, \frac{m}{2} + \frac{5}{2} \end{matrix} \middle| \frac{b^2 x^2}{4}\right)}{8m^2 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 16m \Gamma\left(\frac{m}{2} + \frac{5}{2}\right) + 8 \Gamma\left(\frac{m}{2} + \frac{5}{2}\right)}
 \end{aligned}$$

[In] integrate(x**m*Chi(b*x),x)

[Out] 4*2**m*b*b**(-m - 1)*m*x*sqrt(exp(-2*m*log(2))*exp(m*log(b**2*x**2)))*log(b**2*x**2)*gamma(m/2 + 5/2)/(8*m**2*gamma(m/2 + 5/2) + 16*m*gamma(m/2 + 5/2) + 8*gamma(m/2 + 5/2)) + 8*2**m*EulerGamma*b*b**(-m - 1)*m*x*sqrt(exp(-2*m*log(2))*exp(m*log(b**2*x**2)))*gamma(m/2 + 5/2)/(8*m**2*gamma(m/2 + 5/2) + 16*m*gamma(m/2 + 5/2) + 8*gamma(m/2 + 5/2)) + 4*2**m*b*b**(-m - 1)*x*sqrt(exp(-2*m*log(2))*exp(m*log(b**2*x**2)))*log(b**2*x**2)*gamma(m/2 + 5/2)/(8*m**2*gamma(m/2 + 5/2) + 16*m*gamma(m/2 + 5/2) + 8*gamma(m/2 + 5/2)) - 8*2**m*b*b**(-m - 1)*x*sqrt(exp(-2*m*log(2))*exp(m*log(b**2*x**2)))*gamma(m/2 + 5/2)/(8*m**2*gamma(m/2 + 5/2) + 16*m*gamma(m/2 + 5/2) + 8*gamma(m/2 + 5/2)) + 8*2**m*EulerGamma*b*b**(-m - 1)*x*sqrt(exp(-2*m*log(2))*exp(m*log(b**2*x**2)))*gamma(m/2 + 5/2)/(8*m**2*gamma(m/2 + 5/2) + 16*m*gamma(m/2 + 5/2) + 8*gamma(m/2 + 5/2)) + b**(-m - 1)*b**(m + 3)*m**2*x**(m + 3)*gamma(m/2 + 3/2)*hyper((1, 1, m/2 + 3/2), (3/2, 2, 2, m/2 + 5/2), b**2*x**2/4)/(8*m**2*gamma(m/2 + 5/2) + 16*m*gamma(m/2 + 5/2) + 8*gamma(m/2 + 5/2)) + 2*b**(-m - 1)

```
*b**(m + 3)*m*x**(m + 3)*gamma(m/2 + 3/2)*hyper((1, 1, m/2 + 3/2), (3/2, 2,
2, m/2 + 5/2), b**2*x**2/4)/(8*m**2*gamma(m/2 + 5/2) + 16*m*gamma(m/2 + 5/
2) + 8*gamma(m/2 + 5/2)) + b**(-m - 1)*b**(m + 3)*x**(m + 3)*gamma(m/2 + 3/
2)*hyper((1, 1, m/2 + 3/2), (3/2, 2, 2, m/2 + 5/2), b**2*x**2/4)/(8*m**2*ga
mma(m/2 + 5/2) + 16*m*gamma(m/2 + 5/2) + 8*gamma(m/2 + 5/2))
```

Maxima [F]

$$\int x^m \operatorname{Chi}(bx) dx = \int x^m \operatorname{Chi}(bx) dx$$

```
[In] integrate(x^m*Chi(b*x),x, algorithm="maxima")
```

```
[Out] integrate(x^m*Chi(b*x), x)
```

Giac [F]

$$\int x^m \operatorname{Chi}(bx) dx = \int x^m \operatorname{Chi}(bx) dx$$

```
[In] integrate(x^m*Chi(b*x),x, algorithm="giac")
```

```
[Out] integrate(x^m*Chi(b*x), x)
```

Mupad [F(-1)]

Timed out.

$$\int x^m \operatorname{Chi}(bx) dx = \int x^m \operatorname{coshint}(bx) dx$$

```
[In] int(x^m*coshint(b*x),x)
```

```
[Out] int(x^m*coshint(b*x), x)
```


3.70 $\int x^3 \text{Chi}(bx) dx$

Optimal result	369
Rubi [A] (verified)	369
Mathematica [A] (verified)	370
Maple [A] (verified)	371
Fricas [F]	371
Sympy [A] (verification not implemented)	371
Maxima [F]	372
Giac [F]	372
Mupad [F(-1)]	372

Optimal result

Integrand size = 8, antiderivative size = 63

$$\int x^3 \text{Chi}(bx) dx = \frac{3 \cosh(bx)}{2b^4} + \frac{3x^2 \cosh(bx)}{4b^2} + \frac{1}{4}x^4 \text{Chi}(bx) - \frac{3x \sinh(bx)}{2b^3} - \frac{x^3 \sinh(bx)}{4b}$$

[Out] $\frac{1}{4}x^4 \text{Chi}(bx) + \frac{3}{2} \frac{\cosh(bx)}{b^4} + \frac{3}{4}x^2 \frac{\cosh(bx)}{b^2} - \frac{3}{2}x \frac{\sinh(bx)}{b^3} - \frac{x^3 \sinh(bx)}{4b}$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6668, 12, 3377, 2718}

$$\int x^3 \text{Chi}(bx) dx = \frac{3 \cosh(bx)}{2b^4} - \frac{3x \sinh(bx)}{2b^3} + \frac{3x^2 \cosh(bx)}{4b^2} + \frac{1}{4}x^4 \text{Chi}(bx) - \frac{x^3 \sinh(bx)}{4b}$$

[In] $\text{Int}[x^3 \text{CoshIntegral}[bx], x]$

[Out] $\frac{(3 \text{Cosh}[bx])}{(2b^4)} + \frac{(3x^2 \text{Cosh}[bx])}{(4b^2)} + \frac{(x^4 \text{CoshIntegral}[bx])}{4} - \frac{(3x \text{Sinh}[bx])}{(2b^3)} - \frac{(x^3 \text{Sinh}[bx])}{(4b)}$

Rule 12

$\text{Int}[(a_*)(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /;$ FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 2718

$\text{Int}[\sin[(c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Cos}[c + d*x]/d, x] /;$ FreeQ[{c, d}, x]

Rule 3377

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-
(c + d*x)^m)*(Cos[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Co
s[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 6668

```
Int[CoshIntegral[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol]
:= Simp[(c + d*x)^(m + 1)*(CoshIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/
(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Cosh[a + b*x]/(a + b*x)), x], x] /; Fre
eQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{4}x^4\text{Chi}(bx) - \frac{1}{4}b \int \frac{x^3 \cosh(bx)}{b} dx \\
&= \frac{1}{4}x^4\text{Chi}(bx) - \frac{1}{4} \int x^3 \cosh(bx) dx \\
&= \frac{1}{4}x^4\text{Chi}(bx) - \frac{x^3 \sinh(bx)}{4b} + \frac{3 \int x^2 \sinh(bx) dx}{4b} \\
&= \frac{3x^2 \cosh(bx)}{4b^2} + \frac{1}{4}x^4\text{Chi}(bx) - \frac{x^3 \sinh(bx)}{4b} - \frac{3 \int x \cosh(bx) dx}{2b^2} \\
&= \frac{3x^2 \cosh(bx)}{4b^2} + \frac{1}{4}x^4\text{Chi}(bx) - \frac{3x \sinh(bx)}{2b^3} - \frac{x^3 \sinh(bx)}{4b} + \frac{3 \int \sinh(bx) dx}{2b^3} \\
&= \frac{3 \cosh(bx)}{2b^4} + \frac{3x^2 \cosh(bx)}{4b^2} + \frac{1}{4}x^4\text{Chi}(bx) - \frac{3x \sinh(bx)}{2b^3} - \frac{x^3 \sinh(bx)}{4b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.84

$$\int x^3 \text{Chi}(bx) dx = \frac{3(2 + b^2 x^2) \cosh(bx)}{4b^4} + \frac{1}{4}x^4 \text{Chi}(bx) - \frac{x(6 + b^2 x^2) \sinh(bx)}{4b^3}$$

[In] Integrate[x^3*CoshIntegral[b*x],x]

[Out] (3*(2 + b^2*x^2)*Cosh[b*x])/(4*b^4) + (x^4*CoshIntegral[b*x])/4 - (x*(6 + b^2*x^2)*Sinh[b*x])/(4*b^3)

Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.86

method	result	size
parts	$\frac{x^4 \operatorname{Chi}(bx)}{4} - \frac{b^3 x^3 \sinh(bx) - 3b^2 x^2 \cosh(bx) + 6bx \sinh(bx) - 6 \cosh(bx)}{4b^4}$	54
derivativedivides	$\frac{\frac{b^4 x^4 \operatorname{Chi}(bx)}{4} - \frac{b^3 x^3 \sinh(bx)}{4} + \frac{3b^2 x^2 \cosh(bx)}{4} - \frac{3bx \sinh(bx)}{2} + \frac{3 \cosh(bx)}{2}}{b^4}$	56
default	$\frac{\frac{b^4 x^4 \operatorname{Chi}(bx)}{4} - \frac{b^3 x^3 \sinh(bx)}{4} + \frac{3b^2 x^2 \cosh(bx)}{4} - \frac{3bx \sinh(bx)}{2} + \frac{3 \cosh(bx)}{2}}{b^4}$	56

[In] `int(x^3*Chi(b*x),x,method=_RETURNVERBOSE)`[Out] `1/4*x^4*Chi(b*x)-1/4/b^4*(b^3*x^3*sinh(b*x)-3*b^2*x^2*cosh(b*x)+6*b*x*sinh(b*x)-6*cosh(b*x))`**Fricas [F]**

$$\int x^3 \operatorname{Chi}(bx) dx = \int x^3 \operatorname{Chi}(bx) dx$$

[In] `integrate(x^3*Chi(b*x),x, algorithm="fricas")`[Out] `integral(x^3*cosh_integral(b*x), x)`**Sympy [A] (verification not implemented)**

Time = 1.66 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.35

$$\int x^3 \operatorname{Chi}(bx) dx = -\frac{x^4 \log(bx)}{4} + \frac{x^4 \log(b^2 x^2)}{8} + \frac{x^4 \operatorname{Chi}(bx)}{4} - \frac{x^3 \sinh(bx)}{4b} + \frac{3x^2 \cosh(bx)}{4b^2} - \frac{3x \sinh(bx)}{2b^3} + \frac{3 \cosh(bx)}{2b^4}$$

[In] `integrate(x**3*Chi(b*x),x)`[Out] `-x**4*log(b*x)/4 + x**4*log(b**2*x**2)/8 + x**4*Chi(b*x)/4 - x**3*sinh(b*x)/(4*b) + 3*x**2*cosh(b*x)/(4*b**2) - 3*x*sinh(b*x)/(2*b**3) + 3*cosh(b*x)/(2*b**4)`

Maxima [F]

$$\int x^3 \operatorname{Chi}(bx) dx = \int x^3 \operatorname{Chi}(bx) dx$$

[In] integrate(x^3*Chi(b*x),x, algorithm="maxima")

[Out] integrate(x^3*Chi(b*x), x)

Giac [F]

$$\int x^3 \operatorname{Chi}(bx) dx = \int x^3 \operatorname{Chi}(bx) dx$$

[In] integrate(x^3*Chi(b*x),x, algorithm="giac")

[Out] integrate(x^3*Chi(b*x), x)

Mupad [F(-1)]

Timed out.

$$\int x^3 \operatorname{Chi}(bx) dx = \int x^3 \operatorname{coshint}(bx) dx$$

[In] int(x^3*coshint(b*x),x)

[Out] int(x^3*coshint(b*x), x)

3.71 $\int x^2 \text{Chi}(bx) dx$

Optimal result	373
Rubi [A] (verified)	373
Mathematica [A] (verified)	374
Maple [A] (verified)	375
Fricas [F]	375
Sympy [A] (verification not implemented)	375
Maxima [F]	376
Giac [F]	376
Mupad [F(-1)]	376

Optimal result

Integrand size = 8, antiderivative size = 49

$$\int x^2 \text{Chi}(bx) dx = \frac{2x \cosh(bx)}{3b^2} + \frac{1}{3}x^3 \text{Chi}(bx) - \frac{2 \sinh(bx)}{3b^3} - \frac{x^2 \sinh(bx)}{3b}$$

[Out] $1/3*x^3*\text{Chi}(b*x)+2/3*x*\cosh(b*x)/b^2-2/3*\sinh(b*x)/b^3-1/3*x^2*\sinh(b*x)/b$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6668, 12, 3377, 2717}

$$\int x^2 \text{Chi}(bx) dx = -\frac{2 \sinh(bx)}{3b^3} + \frac{2x \cosh(bx)}{3b^2} + \frac{1}{3}x^3 \text{Chi}(bx) - \frac{x^2 \sinh(bx)}{3b}$$

[In] `Int[x^2*CoshIntegral[b*x],x]`

[Out] $(2*x*\text{Cosh}[b*x])/(3*b^2) + (x^3*\text{CoshIntegral}[b*x])/3 - (2*\text{Sinh}[b*x])/(3*b^3) - (x^2*\text{Sinh}[b*x])/(3*b)$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 2717

`Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Rule 3377

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-
(c + d*x)^m)*(Cos[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Co
s[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 6668

```
Int[CoshIntegral[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol]
:= Simp[(c + d*x)^(m + 1)*(CoshIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/
(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Cosh[a + b*x]/(a + b*x)), x], x] /; Fre
eQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{3}x^3\text{Chi}(bx) - \frac{1}{3}b \int \frac{x^2 \cosh(bx)}{b} dx \\
&= \frac{1}{3}x^3\text{Chi}(bx) - \frac{1}{3} \int x^2 \cosh(bx) dx \\
&= \frac{1}{3}x^3\text{Chi}(bx) - \frac{x^2 \sinh(bx)}{3b} + \frac{2 \int x \sinh(bx) dx}{3b} \\
&= \frac{2x \cosh(bx)}{3b^2} + \frac{1}{3}x^3\text{Chi}(bx) - \frac{x^2 \sinh(bx)}{3b} - \frac{2 \int \cosh(bx) dx}{3b^2} \\
&= \frac{2x \cosh(bx)}{3b^2} + \frac{1}{3}x^3\text{Chi}(bx) - \frac{2 \sinh(bx)}{3b^3} - \frac{x^2 \sinh(bx)}{3b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.90

$$\int x^2\text{Chi}(bx) dx = \frac{2x \cosh(bx)}{3b^2} + \frac{1}{3}x^3\text{Chi}(bx) - \frac{(2 + b^2x^2) \sinh(bx)}{3b^3}$$

```
[In] Integrate[x^2*CoshIntegral[b*x], x]
```

```
[Out] (2*x*Cosh[b*x])/(3*b^2) + (x^3*CoshIntegral[b*x])/3 - ((2 + b^2*x^2)*Sinh[b
*x])/(3*b^3)
```

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.86

method	result	size
parts	$\frac{x^3 \operatorname{Chi}(bx)}{3} - \frac{b^2 x^2 \sinh(bx) - 2bx \cosh(bx) + 2 \sinh(bx)}{3b^3}$	42
derivativedivides	$\frac{\frac{b^3 x^3 \operatorname{Chi}(bx)}{3} - \frac{b^2 x^2 \sinh(bx)}{3} + \frac{2bx \cosh(bx)}{3} - \frac{2 \sinh(bx)}{3}}{b^3}$	44
default	$\frac{\frac{b^3 x^3 \operatorname{Chi}(bx)}{3} - \frac{b^2 x^2 \sinh(bx)}{3} + \frac{2bx \cosh(bx)}{3} - \frac{2 \sinh(bx)}{3}}{b^3}$	44

```
[In] int(x^2*Chi(b*x),x,method=_RETURNVERBOSE)
```

```
[Out] 1/3*x^3*Chi(b*x)-1/3/b^3*(b^2*x^2*sinh(b*x)-2*b*x*cosh(b*x)+2*sinh(b*x))
```

Fricas [F]

$$\int x^2 \operatorname{Chi}(bx) dx = \int x^2 \operatorname{Chi}(bx) dx$$

```
[In] integrate(x^2*Chi(b*x),x, algorithm="fricas")
```

```
[Out] integral(x^2*cosh_integral(b*x), x)
```

Sympy [A] (verification not implemented)

Time = 1.50 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.43

$$\int x^2 \operatorname{Chi}(bx) dx = -\frac{x^3 \log(bx)}{3} + \frac{x^3 \log(b^2 x^2)}{6} + \frac{x^3 \operatorname{Chi}(bx)}{3} - \frac{x^2 \sinh(bx)}{3b} + \frac{2x \cosh(bx)}{3b^2} - \frac{2 \sinh(bx)}{3b^3}$$

```
[In] integrate(x**2*Chi(b*x),x)
```

```
[Out] -x**3*log(b*x)/3 + x**3*log(b**2*x**2)/6 + x**3*Chi(b*x)/3 - x**2*sinh(b*x)/(3*b) + 2*x*cosh(b*x)/(3*b**2) - 2*sinh(b*x)/(3*b**3)
```

Maxima [F]

$$\int x^2 \operatorname{Chi}(bx) dx = \int x^2 \operatorname{Chi}(bx) dx$$

[In] integrate(x^2*Chi(b*x),x, algorithm="maxima")

[Out] integrate(x^2*Chi(b*x), x)

Giac [F]

$$\int x^2 \operatorname{Chi}(bx) dx = \int x^2 \operatorname{Chi}(bx) dx$$

[In] integrate(x^2*Chi(b*x),x, algorithm="giac")

[Out] integrate(x^2*Chi(b*x), x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{Chi}(bx) dx = \frac{x^3 \operatorname{coshint}(bx)}{3} - \frac{\frac{2 \sinh(bx)}{3} + \frac{b^2 x^2 \sinh(bx)}{3} - \frac{2 b x \cosh(bx)}{3}}{b^3}$$

[In] int(x^2*coshint(b*x),x)

[Out] (x^3*coshint(b*x))/3 - ((2*sinh(b*x))/3 + (b^2*x^2*sinh(b*x))/3 - (2*b*x*cosh(b*x))/3)/b^3

3.72 $\int x \operatorname{Chi}(bx) dx$

Optimal result	377
Rubi [A] (verified)	377
Mathematica [A] (verified)	378
Maple [A] (verified)	378
Fricas [F]	379
Sympy [A] (verification not implemented)	379
Maxima [F]	379
Giac [F]	379
Mupad [F(-1)]	380

Optimal result

Integrand size = 6, antiderivative size = 35

$$\int x \operatorname{Chi}(bx) dx = \frac{\cosh(bx)}{2b^2} + \frac{1}{2}x^2 \operatorname{Chi}(bx) - \frac{x \sinh(bx)}{2b}$$

[Out] $1/2*x^2*\operatorname{Chi}(b*x)+1/2*\cosh(b*x)/b^2-1/2*x*\sinh(b*x)/b$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {6668, 12, 3377, 2718}

$$\int x \operatorname{Chi}(bx) dx = \frac{\cosh(bx)}{2b^2} + \frac{1}{2}x^2 \operatorname{Chi}(bx) - \frac{x \sinh(bx)}{2b}$$

[In] `Int[x*CoshIntegral[b*x],x]`

[Out] `Cosh[b*x]/(2*b^2) + (x^2*CoshIntegral[b*x])/2 - (x*Sinh[b*x])/(2*b)`

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 2718

`Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Rule 3377

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(
-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Co
s[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 6668

```
Int[CoshIntegral[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol]
:= Simp[(c + d*x)^(m + 1)*(CoshIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/
(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Cosh[a + b*x]/(a + b*x)), x], x] /; Fre
eQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{2}x^2\text{Chi}(bx) - \frac{1}{2}b \int \frac{x \cosh(bx)}{b} dx \\
&= \frac{1}{2}x^2\text{Chi}(bx) - \frac{1}{2} \int x \cosh(bx) dx \\
&= \frac{1}{2}x^2\text{Chi}(bx) - \frac{x \sinh(bx)}{2b} + \frac{\int \sinh(bx) dx}{2b} \\
&= \frac{\cosh(bx)}{2b^2} + \frac{1}{2}x^2\text{Chi}(bx) - \frac{x \sinh(bx)}{2b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int x\text{Chi}(bx) dx = \frac{\cosh(bx)}{2b^2} + \frac{1}{2}x^2\text{Chi}(bx) - \frac{x \sinh(bx)}{2b}$$

```
[In] Integrate[x*CoshIntegral[b*x],x]
```

```
[Out] Cosh[b*x]/(2*b^2) + (x^2*CoshIntegral[b*x])/2 - (x*Sinh[b*x])/(2*b)
```

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

method	result	size
parts	$\frac{x^2 \text{Chi}(bx)}{2} - \frac{bx \sinh(bx) - \cosh(bx)}{2b^2}$	30
derivativedivides	$\frac{b^2 x^2 \text{Chi}(bx) - bx \sinh(bx) + \cosh(bx)}{2b^2}$	32
default	$\frac{b^2 x^2 \text{Chi}(bx) - bx \sinh(bx) + \cosh(bx)}{2b^2}$	32

[In] `int(x*Chi(b*x),x,method=_RETURNVERBOSE)`

[Out] `1/2*x^2*Chi(b*x)-1/2/b^2*(b*x*sinh(b*x)-cosh(b*x))`

Fricas [F]

$$\int x\text{Chi}(bx) dx = \int x\text{Chi}(bx) dx$$

[In] `integrate(x*Chi(b*x),x, algorithm="fricas")`

[Out] `integral(x*cosh_integral(b*x), x)`

Sympy [A] (verification not implemented)

Time = 1.00 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.51

$$\int x\text{Chi}(bx) dx = -\frac{x^2 \log(bx)}{2} + \frac{x^2 \log(b^2x^2)}{4} + \frac{x^2 \text{Chi}(bx)}{2} - \frac{x \sinh(bx)}{2b} + \frac{\cosh(bx)}{2b^2}$$

[In] `integrate(x*Chi(b*x),x)`

[Out] `-x**2*log(b*x)/2 + x**2*log(b**2*x**2)/4 + x**2*Chi(b*x)/2 - x*sinh(b*x)/(2*b) + cosh(b*x)/(2*b**2)`

Maxima [F]

$$\int x\text{Chi}(bx) dx = \int x\text{Chi}(bx) dx$$

[In] `integrate(x*Chi(b*x),x, algorithm="maxima")`

[Out] `integrate(x*Chi(b*x), x)`

Giac [F]

$$\int x\text{Chi}(bx) dx = \int x\text{Chi}(bx) dx$$

[In] `integrate(x*Chi(b*x),x, algorithm="giac")`

[Out] `integrate(x*Chi(b*x), x)`

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{Chi}(bx) dx = \frac{\frac{\cosh(bx)}{2} - \frac{bx \sinh(bx)}{2}}{b^2} + \frac{x^2 \operatorname{coshint}(bx)}{2}$$

```
[In] int(x*coshint(b*x),x)
```

```
[Out] (cosh(b*x)/2 - (b*x*sinh(b*x))/2)/b^2 + (x^2*coshint(b*x))/2
```

3.73 $\int \text{Chi}(bx) dx$

Optimal result	381
Rubi [A] (verified)	381
Mathematica [A] (verified)	382
Maple [A] (verified)	382
Fricas [F]	382
Sympy [B] (verification not implemented)	383
Maxima [F]	383
Giac [F]	383
Mupad [F(-1)]	383

Optimal result

Integrand size = 4, antiderivative size = 16

$$\int \text{Chi}(bx) dx = x\text{Chi}(bx) - \frac{\sinh(bx)}{b}$$

[Out] $x*\text{Chi}(b*x) - \sinh(b*x)/b$

Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {6664}

$$\int \text{Chi}(bx) dx = x\text{Chi}(bx) - \frac{\sinh(bx)}{b}$$

[In] `Int[CoshIntegral[b*x], x]`

[Out] `x*CoshIntegral[b*x] - Sinh[b*x]/b`

Rule 6664

`Int[CoshIntegral[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[(a + b*x)*(CoshIntegral[a + b*x]/b), x] - Simp[Sinh[a + b*x]/b, x] /; FreeQ[{a, b}, x]`

Rubi steps

$$\text{integral} = x\text{Chi}(bx) - \frac{\sinh(bx)}{b}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \text{Chi}(bx) dx = x\text{Chi}(bx) - \frac{\sinh(bx)}{b}$$

[In] Integrate[CoshIntegral[b*x],x]

[Out] x*CoshIntegral[b*x] - Sinh[b*x]/b

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

method	result	size
parts	$x \text{Chi}(bx) - \frac{\sinh(bx)}{b}$	17
derivativedivides	$\frac{\text{Chi}(bx)bx - \sinh(bx)}{b}$	19
default	$\frac{\text{Chi}(bx)bx - \sinh(bx)}{b}$	19

[In] int(Chi(b*x),x,method=_RETURNVERBOSE)

[Out] x*Chi(b*x)-sinh(b*x)/b

Fricas [F]

$$\int \text{Chi}(bx) dx = \int \text{Chi}(bx) dx$$

[In] integrate(Chi(b*x),x, algorithm="fricas")

[Out] integral(cosh_integral(b*x), x)

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. $2(12) = 24$.

Time = 0.99 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.94

$$\int \text{Chi}(bx) dx = -x \log(bx) + \frac{x \log(b^2 x^2)}{2} + x \text{Chi}(bx) - \frac{\sinh(bx)}{b}$$

[In] integrate(Chi(b*x),x)

[Out] -x*log(b*x) + x*log(b**2*x**2)/2 + x*Chi(b*x) - sinh(b*x)/b

Maxima [F]

$$\int \text{Chi}(bx) dx = \int \text{Chi}(bx) dx$$

[In] integrate(Chi(b*x),x, algorithm="maxima")

[Out] integrate(Chi(b*x), x)

Giac [F]

$$\int \text{Chi}(bx) dx = \int \text{Chi}(bx) dx$$

[In] integrate(Chi(b*x),x, algorithm="giac")

[Out] integrate(Chi(b*x), x)

Mupad [F(-1)]

Timed out.

$$\int \text{Chi}(bx) dx = x \text{coshint}(bx) - \frac{\sinh(bx)}{b}$$

[In] int(coshint(b*x),x)

[Out] x*coshint(b*x) - sinh(b*x)/b

3.74 $\int \frac{\text{Chi}(bx)}{x} dx$

Optimal result	384
Rubi [A] (verified)	384
Mathematica [A] (verified)	385
Maple [F]	385
Fricas [F]	385
Sympy [A] (verification not implemented)	386
Maxima [F]	386
Giac [F]	386
Mupad [F(-1)]	386

Optimal result

Integrand size = 8, antiderivative size = 52

$$\int \frac{\text{Chi}(bx)}{x} dx = -\frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; -bx) + \frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; bx) + \gamma \log(x) + \frac{1}{2} \log^2(bx)$$

[Out] $-1/2*b*x*\text{hypergeom}([1, 1, 1], [2, 2, 2], -b*x) + 1/2*b*x*\text{hypergeom}([1, 1, 1], [2, 2, 2], b*x) + \text{EulerGamma}*\ln(x) + 1/2*\ln(b*x)^2$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {6666}

$$\int \frac{\text{Chi}(bx)}{x} dx = -\frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; -bx) + \frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; bx) + \frac{1}{2} \log^2(bx) + \gamma \log(x)$$

[In] `Int[CoshIntegral[b*x]/x,x]`

[Out] $-1/2*(b*x*\text{HypergeometricPFQ}[\{1, 1, 1\}, \{2, 2, 2\}, -(b*x)]) + (b*x*\text{HypergeometricPFQ}[\{1, 1, 1\}, \{2, 2, 2\}, b*x])/2 + \text{EulerGamma}*\text{Log}[x] + \text{Log}[b*x]^2/2$

Rule 6666

`Int[CoshIntegral[(b_.)*(x_)]/(x_), x_Symbol] := Simp[(-2^(-1))*b*x*HypergeometricPFQ[{1, 1, 1}, {2, 2, 2}, (-b)*x], x] + (Simp[(1/2)*b*x*HypergeometricPFQ[{1, 1, 1}, {2, 2, 2}, b*x], x] + Simp[EulerGamma*Log[x], x] + Simp[(1/`

2)*Log[b*x]^2, x]) /; FreeQ[b, x]

Rubi steps

$$\text{integral} = -\frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; -bx) + \frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; bx) + \gamma \log(x) + \frac{1}{2} \log^2(bx)$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx)}{x} dx = -\frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; -bx) + \frac{1}{2}bx {}_3F_3(1, 1, 1; 2, 2, 2; bx) + \gamma \log(x) + \frac{1}{2} \log^2(bx)$$

[In] Integrate[CoshIntegral[b*x]/x,x]

[Out] -1/2*(b*x*HypergeometricPFQ[{1, 1, 1}, {2, 2, 2}, -(b*x)]) + (b*x*HypergeometricPFQ[{1, 1, 1}, {2, 2, 2}, b*x])/2 + EulerGamma*Log[x] + Log[b*x]^2/2

Maple [F]

$$\int \frac{\text{Chi}(bx)}{x} dx$$

[In] int(Chi(b*x)/x,x)

[Out] int(Chi(b*x)/x,x)

Fricas [F]

$$\int \frac{\text{Chi}(bx)}{x} dx = \int \frac{\text{Chi}(bx)}{x} dx$$

[In] integrate(Chi(b*x)/x,x, algorithm="fricas")

[Out] integral(cosh_integral(b*x)/x, x)

Sympy [A] (verification not implemented)

Time = 0.92 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.81

$$\int \frac{\text{Chi}(bx)}{x} dx = \frac{b^2 x^2 {}_3F_4\left(\begin{matrix} 1, 1, 1 \\ \frac{3}{2}, 2, 2, 2 \end{matrix} \middle| \frac{b^2 x^2}{4}\right)}{8} + \frac{\log(b^2 x^2)^2}{8} + \frac{\gamma \log(b^2 x^2)}{2}$$

[In] integrate(Chi(b*x)/x,x)

[Out] b**2*x**2*hyper((1, 1, 1), (3/2, 2, 2, 2), b**2*x**2/4)/8 + log(b**2*x**2)*
*2/8 + EulerGamma*log(b**2*x**2)/2

Maxima [F]

$$\int \frac{\text{Chi}(bx)}{x} dx = \int \frac{\text{Chi}(bx)}{x} dx$$

[In] integrate(Chi(b*x)/x,x, algorithm="maxima")

[Out] integrate(Chi(b*x)/x, x)

Giac [F]

$$\int \frac{\text{Chi}(bx)}{x} dx = \int \frac{\text{Chi}(bx)}{x} dx$$

[In] integrate(Chi(b*x)/x,x, algorithm="giac")

[Out] integrate(Chi(b*x)/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\text{Chi}(bx)}{x} dx = \int \frac{\text{coshint}(bx)}{x} dx$$

[In] int(coshint(b*x)/x,x)

[Out] int(coshint(b*x)/x, x)

3.75 $\int \frac{\text{Chi}(bx)}{x^2} dx$

Optimal result	387
Rubi [A] (verified)	387
Mathematica [A] (verified)	388
Maple [A] (verified)	388
Fricas [F]	389
Sympy [B] (verification not implemented)	389
Maxima [F]	390
Giac [F]	390
Mupad [F(-1)]	390

Optimal result

Integrand size = 8, antiderivative size = 25

$$\int \frac{\text{Chi}(bx)}{x^2} dx = -\frac{\cosh(bx)}{x} - \frac{\text{Chi}(bx)}{x} + b\text{Shi}(bx)$$

[Out] -Chi(b*x)/x-cosh(b*x)/x+b*Shi(b*x)

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6668, 12, 3378, 3379}

$$\int \frac{\text{Chi}(bx)}{x^2} dx = -\frac{\text{Chi}(bx)}{x} + b\text{Shi}(bx) - \frac{\cosh(bx)}{x}$$

[In] Int[CoshIntegral[b*x]/x^2,x]

[Out] -(Cosh[b*x]/x) - CoshIntegral[b*x]/x + b*SinhIntegral[b*x]

Rule 12

Int[(a_)*(u_), x_Symbol] :=> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 3378

Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] :=> Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Dist[f/(d*(m + 1)), Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol]
:> Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 6668

```
Int[CoshIntegral[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol]
:> Simp[(c + d*x)^(m + 1)*(CoshIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Cosh[a + b*x]/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Chi}(bx)}{x} + b \int \frac{\cosh(bx)}{bx^2} dx \\
&= -\frac{\text{Chi}(bx)}{x} + \int \frac{\cosh(bx)}{x^2} dx \\
&= -\frac{\cosh(bx)}{x} - \frac{\text{Chi}(bx)}{x} + b \int \frac{\sinh(bx)}{x} dx \\
&= -\frac{\cosh(bx)}{x} - \frac{\text{Chi}(bx)}{x} + b\text{Shi}(bx)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx)}{x^2} dx = -\frac{\cosh(bx)}{x} - \frac{\text{Chi}(bx)}{x} + b\text{Shi}(bx)$$

```
[In] Integrate[CoshIntegral[b*x]/x^2,x]
```

```
[Out] -(Cosh[b*x]/x) - CoshIntegral[b*x]/x + b*SinhIntegral[b*x]
```

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.20

method	result	size
parts	$-\frac{\text{Chi}(bx)}{x} + b\left(-\frac{\cosh(bx)}{bx} + \text{Shi}(bx)\right)$	30
derivativedivides	$b\left(-\frac{\text{Chi}(bx)}{bx} - \frac{\cosh(bx)}{bx} + \text{Shi}(bx)\right)$	32
default	$b\left(-\frac{\text{Chi}(bx)}{bx} - \frac{\cosh(bx)}{bx} + \text{Shi}(bx)\right)$	32

[In] `int(Chi(b*x)/x^2,x,method=_RETURNVERBOSE)`

[Out] `-Chi(b*x)/x+b*(-1/b/x*cosh(b*x)+Shi(b*x))`

Fricas [F]

$$\int \frac{\text{Chi}(bx)}{x^2} dx = \int \frac{\text{Chi}(bx)}{x^2} dx$$

[In] `integrate(Chi(b*x)/x^2,x, algorithm="fricas")`

[Out] `integral(cosh_integral(b*x)/x^2, x)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. $2(19) = 38$.

Time = 0.60 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.56

$$\int \frac{\text{Chi}(bx)}{x^2} dx = \frac{b^2 x {}_3F_4\left(\frac{1}{2}, 1, 1 \mid \frac{b^2 x^2}{4}\right)}{4} - \frac{\log(b^2 x^2)}{2x} - \frac{1}{x} - \frac{\gamma}{x}$$

[In] `integrate(Chi(b*x)/x**2,x)`

[Out] `b**2*x*hyper((1/2, 1, 1), (3/2, 3/2, 2, 2), b**2*x**2/4)/4 - log(b**2*x**2)/(2*x) - 1/x - EulerGamma/x`

Maxima [F]

$$\int \frac{\text{Chi}(bx)}{x^2} dx = \int \frac{\text{Chi}(bx)}{x^2} dx$$

[In] integrate(Chi(b*x)/x^2,x, algorithm="maxima")

[Out] integrate(Chi(b*x)/x^2, x)

Giac [F]

$$\int \frac{\text{Chi}(bx)}{x^2} dx = \int \frac{\text{Chi}(bx)}{x^2} dx$$

[In] integrate(Chi(b*x)/x^2,x, algorithm="giac")

[Out] integrate(Chi(b*x)/x^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\text{Chi}(bx)}{x^2} dx = \int \frac{\text{coshint}(b x)}{x^2} dx$$

[In] int(coshint(b*x)/x^2,x)

[Out] int(coshint(b*x)/x^2, x)

3.76 $\int \frac{\text{Chi}(bx)}{x^3} dx$

Optimal result	391
Rubi [A] (verified)	391
Mathematica [A] (verified)	392
Maple [A] (verified)	393
Fricas [F]	393
Sympy [B] (verification not implemented)	393
Maxima [F]	394
Giac [F]	394
Mupad [F(-1)]	394

Optimal result

Integrand size = 8, antiderivative size = 46

$$\int \frac{\text{Chi}(bx)}{x^3} dx = -\frac{\cosh(bx)}{4x^2} + \frac{1}{4}b^2\text{Chi}(bx) - \frac{\text{Chi}(bx)}{2x^2} - \frac{b \sinh(bx)}{4x}$$

[Out] $1/4*b^2*\text{Chi}(b*x)-1/2*\text{Chi}(b*x)/x^2-1/4*\cosh(b*x)/x^2-1/4*b*\sinh(b*x)/x$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6668, 12, 3378, 3382}

$$\int \frac{\text{Chi}(bx)}{x^3} dx = \frac{1}{4}b^2\text{Chi}(bx) - \frac{\text{Chi}(bx)}{2x^2} - \frac{\cosh(bx)}{4x^2} - \frac{b \sinh(bx)}{4x}$$

[In] Int[CoshIntegral[b*x]/x^3,x]

[Out] $-1/4*\text{Cosh}[b*x]/x^2 + (b^2*\text{CoshIntegral}[b*x])/4 - \text{CoshIntegral}[b*x]/(2*x^2) - (b*\text{Sinh}[b*x])/(4*x)$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 3378

Int[((c_.) + (d_)*(x_))^(m_)*sin[(e_.) + (f_)*(x_)], x_Symbol] := Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Dist[f/(d*(m + 1)), Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]

]

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol]
:> Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x]
&& EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 6668

```
Int[CoshIntegral[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol]
:> Simp[(c + d*x)^(m + 1)*(CoshIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Cosh[a + b*x]/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Chi}(bx)}{2x^2} + \frac{1}{2}b \int \frac{\cosh(bx)}{bx^3} dx \\
&= -\frac{\text{Chi}(bx)}{2x^2} + \frac{1}{2} \int \frac{\cosh(bx)}{x^3} dx \\
&= -\frac{\cosh(bx)}{4x^2} - \frac{\text{Chi}(bx)}{2x^2} + \frac{1}{4}b \int \frac{\sinh(bx)}{x^2} dx \\
&= -\frac{\cosh(bx)}{4x^2} - \frac{\text{Chi}(bx)}{2x^2} - \frac{b \sinh(bx)}{4x} + \frac{1}{4}b^2 \int \frac{\cosh(bx)}{x} dx \\
&= -\frac{\cosh(bx)}{4x^2} + \frac{1}{4}b^2 \text{Chi}(bx) - \frac{\text{Chi}(bx)}{2x^2} - \frac{b \sinh(bx)}{4x}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx)}{x^3} dx = -\frac{\cosh(bx)}{4x^2} + \frac{1}{4}b^2 \text{Chi}(bx) - \frac{\text{Chi}(bx)}{2x^2} - \frac{b \sinh(bx)}{4x}$$

```
[In] Integrate[CoshIntegral[b*x]/x^3,x]
```

```
[Out] -1/4*Cosh[b*x]/x^2 + (b^2*CoshIntegral[b*x])/4 - CoshIntegral[b*x]/(2*x^2)
- (b*Sinh[b*x])/(4*x)
```


Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.02

method	result	size
parts	$-\frac{\text{Chi}(bx)}{2x^2} + \frac{b^2 \left(-\frac{\cosh(bx)}{2b^2x^2} - \frac{\sinh(bx)}{2bx} + \frac{\text{Chi}(bx)}{2} \right)}{2}$	47
derivativedivides	$b^2 \left(-\frac{\text{Chi}(bx)}{2b^2x^2} - \frac{\cosh(bx)}{4b^2x^2} - \frac{\sinh(bx)}{4bx} + \frac{\text{Chi}(bx)}{4} \right)$	48
default	$b^2 \left(-\frac{\text{Chi}(bx)}{2b^2x^2} - \frac{\cosh(bx)}{4b^2x^2} - \frac{\sinh(bx)}{4bx} + \frac{\text{Chi}(bx)}{4} \right)$	48

[In] `int(Chi(b*x)/x^3,x,method=_RETURNVERBOSE)`

[Out] `-1/2*Chi(b*x)/x^2+1/2*b^2*(-1/2/b^2/x^2*cosh(b*x)-1/2*sinh(b*x)/b/x+1/2*Chi(b*x))`

Fricas [F]

$$\int \frac{\text{Chi}(bx)}{x^3} dx = \int \frac{\text{Chi}(bx)}{x^3} dx$$

[In] `integrate(Chi(b*x)/x^3,x, algorithm="fricas")`

[Out] `integral(cosh_integral(b*x)/x^3, x)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 87 vs. 2(39) = 78.

Time = 2.41 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.89

$$\int \frac{\text{Chi}(bx)}{x^3} dx = -\frac{b^2 \log(bx)}{4} + \frac{b^2 \log(b^2x^2)}{8} + \frac{b^2 \text{Chi}(bx)}{4} - \frac{b \sinh(bx)}{4x} + \frac{\log(bx)}{2x^2} - \frac{\log(b^2x^2)}{4x^2} - \frac{\cosh(bx)}{4x^2} - \frac{\text{Chi}(bx)}{2x^2}$$

[In] `integrate(Chi(b*x)/x**3,x)`

[Out] `-b**2*log(b*x)/4 + b**2*log(b**2*x**2)/8 + b**2*Chi(b*x)/4 - b*sinh(b*x)/(4*x) + log(b*x)/(2*x**2) - log(b**2*x**2)/(4*x**2) - cosh(b*x)/(4*x**2) - Chi(b*x)/(2*x**2)`

Maxima [F]

$$\int \frac{\text{Chi}(bx)}{x^3} dx = \int \frac{\text{Chi}(bx)}{x^3} dx$$

[In] integrate(Chi(b*x)/x^3,x, algorithm="maxima")

[Out] integrate(Chi(b*x)/x^3, x)

Giac [F]

$$\int \frac{\text{Chi}(bx)}{x^3} dx = \int \frac{\text{Chi}(bx)}{x^3} dx$$

[In] integrate(Chi(b*x)/x^3,x, algorithm="giac")

[Out] integrate(Chi(b*x)/x^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\text{Chi}(bx)}{x^3} dx = \frac{b^2 \text{coshint}(bx)}{4} - \frac{\frac{\text{coshint}(bx)}{2} + \frac{\cosh(bx)}{4} + \frac{bx \sinh(bx)}{4}}{x^2}$$

[In] int(coshint(b*x)/x^3,x)

[Out] (b^2*coshint(b*x))/4 - (coshint(b*x)/2 + cosh(b*x)/4 + (b*x*sinh(b*x))/4)/x^2

3.77 $\int x^m \mathbf{Chi}(bx)^2 dx$

Optimal result	395
Rubi [N/A]	395
Mathematica [N/A]	396
Maple [N/A] (verified)	396
Fricas [N/A]	396
Sympy [N/A]	396
Maxima [N/A]	397
Giac [N/A]	397
Mupad [N/A]	397

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int x^m \mathbf{Chi}(bx)^2 dx = \text{Int}(x^m \mathbf{Chi}(bx)^2, x)$$

[Out] CannotIntegrate(x^m*Chi(b*x)^2,x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int x^m \mathbf{Chi}(bx)^2 dx = \int x^m \mathbf{Chi}(bx)^2 dx$$

[In] Int[x^m*CoshIntegral[b*x]^2,x]

[Out] Defer[Int][x^m*CoshIntegral[b*x]^2, x]

Rubi steps

$$\text{integral} = \int x^m \mathbf{Chi}(bx)^2 dx$$

Mathematica [N/A]

Not integrable

Time = 0.44 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Chi}(bx)^2 dx = \int x^m \operatorname{Chi}(bx)^2 dx$$

[In] Integrate[x^m*CoshIntegral[b*x]^2,x]

[Out] Integrate[x^m*CoshIntegral[b*x]^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.23 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int x^m \operatorname{Chi}(bx)^2 dx$$

[In] int(x^m*Chi(b*x)^2,x)

[Out] int(x^m*Chi(b*x)^2,x)

Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Chi}(bx)^2 dx = \int x^m \operatorname{Chi}(bx)^2 dx$$

[In] integrate(x^m*Chi(b*x)^2,x, algorithm="fricas")

[Out] integral(x^m*cosh_integral(b*x)^2, x)

Sympy [N/A]

Not integrable

Time = 0.79 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int x^m \operatorname{Chi}(bx)^2 dx = \int x^m \operatorname{Chi}^2(bx) dx$$

[In] integrate(x**m*Chi(b*x)**2,x)

[Out] Integral(x**m*Chi(b*x)**2, x)

Maxima [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Chi}(bx)^2 dx = \int x^m \operatorname{Chi}(bx)^2 dx$$

[In] integrate(x^m*Chi(b*x)^2,x, algorithm="maxima")

[Out] integrate(x^m*Chi(b*x)^2, x)

Giac [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Chi}(bx)^2 dx = \int x^m \operatorname{Chi}(bx)^2 dx$$

[In] integrate(x^m*Chi(b*x)^2,x, algorithm="giac")

[Out] integrate(x^m*Chi(b*x)^2, x)

Mupad [N/A]

Not integrable

Time = 4.84 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Chi}(bx)^2 dx = \int x^m \operatorname{coshint}(bx)^2 dx$$

[In] int(x^m*coshint(b*x)^2,x)

[Out] int(x^m*coshint(b*x)^2, x)

3.78 $\int x^3 \text{Chi}(bx)^2 dx$

Optimal result	398
Rubi [A] (verified)	398
Mathematica [A] (verified)	402
Maple [A] (verified)	402
Fricas [F]	402
Sympy [F]	403
Maxima [F]	403
Giac [F]	403
Mupad [F(-1)]	403

Optimal result

Integrand size = 10, antiderivative size = 164

$$\int x^3 \text{Chi}(bx)^2 dx = -\frac{x^2}{4b^2} + \frac{3 \cosh^2(bx)}{8b^4} + \frac{3 \cosh(bx) \text{Chi}(bx)}{b^4} + \frac{3x^2 \cosh(bx) \text{Chi}(bx)}{2b^2}$$

$$+ \frac{1}{4} x^4 \text{Chi}(bx)^2 - \frac{3 \text{Chi}(2bx)}{2b^4} - \frac{3 \log(x)}{2b^4} - \frac{x \cosh(bx) \sinh(bx)}{b^3}$$

$$- \frac{3x \text{Chi}(bx) \sinh(bx)}{b^3} - \frac{x^3 \text{Chi}(bx) \sinh(bx)}{2b} + \frac{13 \sinh^2(bx)}{8b^4} + \frac{x^2 \sinh^2(bx)}{4b^2}$$

[Out] $-1/4*x^2/b^2+1/4*x^4*\text{Chi}(b*x)^2-3/2*\text{Chi}(2*b*x)/b^4+3*\text{Chi}(b*x)*\cosh(b*x)/b^4$
 $+3/2*x^2*\text{Chi}(b*x)*\cosh(b*x)/b^2+3/8*\cosh(b*x)^2/b^4-3/2*\ln(x)/b^4-3*x*\text{Chi}(b$
 $*x)*\sinh(b*x)/b^3-1/2*x^3*\text{Chi}(b*x)*\sinh(b*x)/b-x*\cosh(b*x)*\sinh(b*x)/b^3+13$
 $/8*\sinh(b*x)^2/b^4+1/4*x^2*\sinh(b*x)^2/b^2$

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.00,
 number of steps used = 19, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 1.100$, Rules
 used = {6672, 6678, 12, 5480, 3391, 30, 6684, 2644, 6682, 3393, 3382}

$$\int x^3 \text{Chi}(bx)^2 dx = -\frac{3 \text{Chi}(2bx)}{2b^4} + \frac{3 \text{Chi}(bx) \cosh(bx)}{b^4} - \frac{3 \log(x)}{2b^4} + \frac{13 \sinh^2(bx)}{8b^4} + \frac{3 \cosh^2(bx)}{8b^4}$$

$$- \frac{3x \text{Chi}(bx) \sinh(bx)}{b^3} - \frac{x \sinh(bx) \cosh(bx)}{b^3} + \frac{3x^2 \text{Chi}(bx) \cosh(bx)}{2b^2}$$

$$- \frac{x^2}{4b^2} + \frac{x^2 \sinh^2(bx)}{4b^2} + \frac{1}{4} x^4 \text{Chi}(bx)^2 - \frac{x^3 \text{Chi}(bx) \sinh(bx)}{2b}$$

[In] $\text{Int}[x^3*\text{CoshIntegral}[b*x]^2,x]$

```
[Out] -1/4*x^2/b^2 + (3*Cosh[b*x]^2)/(8*b^4) + (3*Cosh[b*x]*CoshIntegral[b*x])/b^
4 + (3*x^2*Cosh[b*x]*CoshIntegral[b*x])/(2*b^2) + (x^4*CoshIntegral[b*x]^2)
/4 - (3*CoshIntegral[2*b*x])/(2*b^4) - (3*Log[x])/(2*b^4) - (x*Cosh[b*x]*Si
nh[b*x])/b^3 - (3*x*CoshIntegral[b*x]*Sinh[b*x])/b^3 - (x^3*CoshIntegral[b*
x]*Sinh[b*x])/(2*b) + (13*Sinh[b*x]^2)/(8*b^4) + (x^2*Sinh[b*x]^2)/(4*b^2)
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 30

```
Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && N
eQ[m, -1]
```

Rule 2644

```
Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_
Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*
Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(In
tegerQ[(m - 1)/2] && LtQ[0, m, n])
```

Rule 3382

```
Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbo
l] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz
}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3391

```
Int[((c_) + (d_)*(x_))*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :=
Simp[d*((b*SIN[e + f*x])^n/(f^2*n^2)), x] + (Dist[b^2*((n - 1)/n), Int[(c
+ d*x)*(b*SIN[e + f*x])^(n - 2), x], x] - Simp[b*(c + d*x)*Cos[e + f*x]*((b
*SIN[e + f*x])^(n - 1)/(f*n)), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1
]
```

Rule 3393

```
Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] := In
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 5480

```
Int[Cosh[(a_) + (b_)*(x_)^(n_)]*(x_)^(m_)*Sinh[(a_) + (b_)*(x_)^(n_)]
]^(p_), x_Symbol] := Simp[x^(m - n + 1)*(Sinh[a + b*x^n]^(p + 1)/(b*n*(p +
```

1))), x] - Dist[(m - n + 1)/(b*n*(p + 1)), Int[x^(m - n)*Sinh[a + b*x^n]^(p + 1), x], x] /; FreeQ[{a, b, p}, x] && LtQ[0, n, m + 1] && NeQ[p, -1]

Rule 6672

Int[CoshIntegral[(b_.)*(x_)]^2*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*(CoshIntegral[b*x]^2/(m + 1)), x] - Dist[2/(m + 1), Int[x^m*Cosh[b*x]*CoshIntegral[b*x], x], x] /; FreeQ[b, x] && IGtQ[m, 0]

Rule 6678

Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^m*Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*CoshIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rule 6682

Int[CoshIntegral[(c_.) + (d_.)*(x_)]*Sinh[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]

Rule 6684

Int[CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*CoshIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{4}x^4\text{Chi}(bx)^2 - \frac{1}{2}\int x^3 \cosh(bx)\text{Chi}(bx) dx \\
 &= \frac{1}{4}x^4\text{Chi}(bx)^2 - \frac{x^3\text{Chi}(bx) \sinh(bx)}{2b} \\
 &\quad + \frac{1}{2}\int \frac{x^2 \cosh(bx) \sinh(bx)}{b} dx + \frac{3\int x^2\text{Chi}(bx) \sinh(bx) dx}{2b} \\
 &= \frac{3x^2 \cosh(bx)\text{Chi}(bx)}{2b^2} + \frac{1}{4}x^4\text{Chi}(bx)^2 - \frac{x^3\text{Chi}(bx) \sinh(bx)}{2b} \\
 &\quad - \frac{3\int x \cosh(bx)\text{Chi}(bx) dx}{b^2} + \frac{\int x^2 \cosh(bx) \sinh(bx) dx}{2b} - \frac{3\int \frac{x \cosh^2(bx)}{b} dx}{2b}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{3x^2 \cosh(bx) \operatorname{Chi}(bx)}{2b^2} + \frac{1}{4} x^4 \operatorname{Chi}(bx)^2 - \frac{3x \operatorname{Chi}(bx) \sinh(bx)}{b^3} \\
&\quad - \frac{x^3 \operatorname{Chi}(bx) \sinh(bx)}{2b} + \frac{x^2 \sinh^2(bx)}{4b^2} + \frac{3 \int \operatorname{Chi}(bx) \sinh(bx) dx}{b^3} \\
&\quad - \frac{\int x \sinh^2(bx) dx}{2b^2} - \frac{3 \int x \cosh^2(bx) dx}{2b^2} + \frac{3 \int \frac{\cosh(bx) \sinh(bx)}{b} dx}{b^2} \\
&= \frac{3 \cosh^2(bx)}{8b^4} + \frac{3 \cosh(bx) \operatorname{Chi}(bx)}{b^4} + \frac{3x^2 \cosh(bx) \operatorname{Chi}(bx)}{2b^2} + \frac{1}{4} x^4 \operatorname{Chi}(bx)^2 \\
&\quad - \frac{x \cosh(bx) \sinh(bx)}{b^3} - \frac{3x \operatorname{Chi}(bx) \sinh(bx)}{b^3} - \frac{x^3 \operatorname{Chi}(bx) \sinh(bx)}{2b} + \frac{\sinh^2(bx)}{8b^4} \\
&\quad + \frac{x^2 \sinh^2(bx)}{4b^2} - \frac{3 \int \frac{\cosh^2(bx)}{bx} dx}{b^3} + \frac{3 \int \cosh(bx) \sinh(bx) dx}{b^3} + \frac{\int x dx}{4b^2} - \frac{3 \int x dx}{4b^2} \\
&= -\frac{x^2}{4b^2} + \frac{3 \cosh^2(bx)}{8b^4} + \frac{3 \cosh(bx) \operatorname{Chi}(bx)}{b^4} + \frac{3x^2 \cosh(bx) \operatorname{Chi}(bx)}{2b^2} + \frac{1}{4} x^4 \operatorname{Chi}(bx)^2 \\
&\quad - \frac{x \cosh(bx) \sinh(bx)}{b^3} - \frac{3x \operatorname{Chi}(bx) \sinh(bx)}{b^3} - \frac{x^3 \operatorname{Chi}(bx) \sinh(bx)}{2b} \\
&\quad + \frac{\sinh^2(bx)}{8b^4} + \frac{x^2 \sinh^2(bx)}{4b^2} - \frac{3 \int \frac{\cosh^2(bx)}{x} dx}{b^4} - \frac{3 \operatorname{Subst}(\int x dx, x, i \sinh(bx))}{b^4} \\
&= -\frac{x^2}{4b^2} + \frac{3 \cosh^2(bx)}{8b^4} + \frac{3 \cosh(bx) \operatorname{Chi}(bx)}{b^4} + \frac{3x^2 \cosh(bx) \operatorname{Chi}(bx)}{2b^2} \\
&\quad + \frac{1}{4} x^4 \operatorname{Chi}(bx)^2 - \frac{x \cosh(bx) \sinh(bx)}{b^3} - \frac{3x \operatorname{Chi}(bx) \sinh(bx)}{b^3} \\
&\quad - \frac{x^3 \operatorname{Chi}(bx) \sinh(bx)}{2b} + \frac{13 \sinh^2(bx)}{8b^4} + \frac{x^2 \sinh^2(bx)}{4b^2} - \frac{3 \int \left(\frac{1}{2x} + \frac{\cosh(2bx)}{2x} \right) dx}{b^4} \\
&= -\frac{x^2}{4b^2} + \frac{3 \cosh^2(bx)}{8b^4} + \frac{3 \cosh(bx) \operatorname{Chi}(bx)}{b^4} + \frac{3x^2 \cosh(bx) \operatorname{Chi}(bx)}{2b^2} \\
&\quad + \frac{1}{4} x^4 \operatorname{Chi}(bx)^2 - \frac{3 \log(x)}{2b^4} - \frac{x \cosh(bx) \sinh(bx)}{b^3} - \frac{3x \operatorname{Chi}(bx) \sinh(bx)}{b^3} \\
&\quad - \frac{x^3 \operatorname{Chi}(bx) \sinh(bx)}{2b} + \frac{13 \sinh^2(bx)}{8b^4} + \frac{x^2 \sinh^2(bx)}{4b^2} - \frac{3 \int \frac{\cosh(2bx)}{x} dx}{2b^4} \\
&= -\frac{x^2}{4b^2} + \frac{3 \cosh^2(bx)}{8b^4} + \frac{3 \cosh(bx) \operatorname{Chi}(bx)}{b^4} + \frac{3x^2 \cosh(bx) \operatorname{Chi}(bx)}{2b^2} \\
&\quad + \frac{1}{4} x^4 \operatorname{Chi}(bx)^2 - \frac{3 \operatorname{Chi}(2bx)}{2b^4} - \frac{3 \log(x)}{2b^4} - \frac{x \cosh(bx) \sinh(bx)}{b^3} \\
&\quad - \frac{3x \operatorname{Chi}(bx) \sinh(bx)}{b^3} - \frac{x^3 \operatorname{Chi}(bx) \sinh(bx)}{2b} + \frac{13 \sinh^2(bx)}{8b^4} + \frac{x^2 \sinh^2(bx)}{4b^2}
\end{aligned}$$

Sympy [F]

$$\int x^3 \operatorname{Chi}(bx)^2 dx = \int x^3 \operatorname{Chi}^2(bx) dx$$

[In] `integrate(x**3*Chi(b*x)**2,x)`

[Out] `Integral(x**3*Chi(b*x)**2, x)`

Maxima [F]

$$\int x^3 \operatorname{Chi}(bx)^2 dx = \int x^3 \operatorname{Chi}(bx)^2 dx$$

[In] `integrate(x^3*Chi(b*x)^2,x, algorithm="maxima")`

[Out] `integrate(x^3*Chi(b*x)^2, x)`

Giac [F]

$$\int x^3 \operatorname{Chi}(bx)^2 dx = \int x^3 \operatorname{Chi}(bx)^2 dx$$

[In] `integrate(x^3*Chi(b*x)^2,x, algorithm="giac")`

[Out] `integrate(x^3*Chi(b*x)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^3 \operatorname{Chi}(bx)^2 dx = \int x^3 \operatorname{coshint}(bx)^2 dx$$

[In] `int(x^3*coshint(b*x)^2,x)`

[Out] `int(x^3*coshint(b*x)^2, x)`

3.79 $\int x^2 \text{Chi}(bx)^2 dx$

Optimal result	404
Rubi [A] (verified)	404
Mathematica [A] (verified)	407
Maple [A] (verified)	407
Fricas [F]	408
Sympy [F]	408
Maxima [F]	408
Giac [F]	408
Mupad [F(-1)]	409

Optimal result

Integrand size = 10, antiderivative size = 112

$$\int x^2 \text{Chi}(bx)^2 dx = -\frac{x}{2b^2} + \frac{4x \cosh(bx) \text{Chi}(bx)}{3b^2} + \frac{1}{3} x^3 \text{Chi}(bx)^2 - \frac{5 \cosh(bx) \sinh(bx)}{6b^3} \\ - \frac{4 \text{Chi}(bx) \sinh(bx)}{3b^3} - \frac{2x^2 \text{Chi}(bx) \sinh(bx)}{3b} + \frac{x \sinh^2(bx)}{3b^2} + \frac{2 \text{Shi}(2bx)}{3b^3}$$

[Out] $-1/2*x/b^2 + 1/3*x^3*\text{Chi}(b*x)^2 + 4/3*x*\text{Chi}(b*x)*\cosh(b*x)/b^2 + 2/3*\text{Shi}(2*b*x)/b^3 - 4/3*\text{Chi}(b*x)*\sinh(b*x)/b^3 - 2/3*x^2*\text{Chi}(b*x)*\sinh(b*x)/b - 5/6*\cosh(b*x)*\sinh(b*x)/b^3 + 1/3*x*\sinh(b*x)^2/b^2$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {6672, 6678, 12, 5480, 2715, 8, 6684, 6676, 5556, 3379}

$$\int x^2 \text{Chi}(bx)^2 dx = -\frac{4 \text{Chi}(bx) \sinh(bx)}{3b^3} + \frac{2 \text{Shi}(2bx)}{3b^3} - \frac{5 \sinh(bx) \cosh(bx)}{6b^3} \\ + \frac{4x \text{Chi}(bx) \cosh(bx)}{3b^2} - \frac{x}{2b^2} + \frac{x \sinh^2(bx)}{3b^2} \\ + \frac{1}{3} x^3 \text{Chi}(bx)^2 - \frac{2x^2 \text{Chi}(bx) \sinh(bx)}{3b}$$

[In] $\text{Int}[x^2*\text{CoshIntegral}[b*x]^2, x]$

[Out] $-1/2*x/b^2 + (4*x*\text{Cosh}[b*x]*\text{CoshIntegral}[b*x])/(3*b^2) + (x^3*\text{CoshIntegral}[b*x]^2)/3 - (5*\text{Cosh}[b*x]*\text{Sinh}[b*x])/(6*b^3) - (4*\text{CoshIntegral}[b*x]*\text{Sinh}[b*x])/ (3*b^3) - (2*x^2*\text{CoshIntegral}[b*x]*\text{Sinh}[b*x])/(3*b) + (x*\text{Sinh}[b*x]^2)/(3*b^2) + (2*\text{SinhIntegral}[2*b*x])/(3*b^3)$

Rule 8

```
Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 2715

```
Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[
c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2
*n]
```

Rule 3379

```
Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol]
:= Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f
, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 5480

```
Int[Cosh[(a_) + (b_)*(x_)^(n_)]*(x_)^(m_)*Sinh[(a_) + (b_)*(x_)^(n_)]
]^(p_), x_Symbol] := Simp[x^(m - n + 1)*(Sinh[a + b*x^n]^(p + 1)/(b*n*(p +
1))), x] - Dist[(m - n + 1)/(b*n*(p + 1)), Int[x^(m - n)*Sinh[a + b*x^n]^(
p + 1), x], x] /; FreeQ[{a, b, p}, x] && LtQ[0, n, m + 1] && NeQ[p, -1]
```

Rule 5556

```
Int[Cosh[(a_) + (b_)*(x_)]^(p_)*((c_) + (d_)*(x_))^(m_)*Sinh[(a_) +
(b_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

Rule 6672

```
Int[CoshIntegral[(b_)*(x_)]^2*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*(Cos
hIntegral[b*x]^2/(m + 1), x] - Dist[2/(m + 1), Int[x^m*Cosh[b*x]*CoshInteg
ral[b*x], x], x] /; FreeQ[b, x] && IGtQ[m, 0]
```

Rule 6676

```
Int[Cosh[(a_) + (b_)*(x_)]*CoshIntegral[(c_) + (d_)*(x_)], x_Symbol] :=
Simp[Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
```

$b*x]*(\text{Cosh}[c + d*x]/(c + d*x)), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x]$

Rule 6678

$\text{Int}[\text{Cosh}[(a_.) + (b_.)*(x_.)]*\text{CoshIntegral}[(c_.) + (d_.)*(x_.)]*((e_.) + (f_.)*(x_.))^{(m_.)}, x_Symbol] :> \text{Simp}[(e + f*x)^m*\text{Sinh}[a + b*x]*(\text{CoshIntegral}[c + d*x]/b), x] + (-\text{Dist}[d/b, \text{Int}[(e + f*x)^m*\text{Sinh}[a + b*x]*(\text{Cosh}[c + d*x]/(c + d*x)), x], x] - \text{Dist}[f*(m/b), \text{Int}[(e + f*x)^{(m-1)}*\text{Sinh}[a + b*x]*\text{CoshIntegral}[c + d*x], x], x]) /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{IGtQ}[m, 0]$

Rule 6684

$\text{Int}[\text{CoshIntegral}[(c_.) + (d_.)*(x_.)]*((e_.) + (f_.)*(x_.))^{(m_.)}*\text{Sinh}[(a_.) + (b_.)*(x_.)], x_Symbol] :> \text{Simp}[(e + f*x)^m*\text{Cosh}[a + b*x]*(\text{CoshIntegral}[c + d*x]/b), x] + (-\text{Dist}[d/b, \text{Int}[(e + f*x)^m*\text{Cosh}[a + b*x]*(\text{Cosh}[c + d*x]/(c + d*x)), x], x] - \text{Dist}[f*(m/b), \text{Int}[(e + f*x)^{(m-1)}*\text{Cosh}[a + b*x]*\text{CoshIntegral}[c + d*x], x], x]) /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{IGtQ}[m, 0]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{3}x^3\text{Chi}(bx)^2 - \frac{2}{3}\int x^2 \cosh(bx)\text{Chi}(bx) dx \\
 &= \frac{1}{3}x^3\text{Chi}(bx)^2 - \frac{2x^2\text{Chi}(bx)\sinh(bx)}{3b} + \frac{2}{3}\int \frac{x \cosh(bx)\sinh(bx)}{b} dx + \frac{4\int x\text{Chi}(bx)\sinh(bx) dx}{3b} \\
 &= \frac{4x \cosh(bx)\text{Chi}(bx)}{3b^2} + \frac{1}{3}x^3\text{Chi}(bx)^2 - \frac{2x^2\text{Chi}(bx)\sinh(bx)}{3b} \\
 &\quad - \frac{4\int \cosh(bx)\text{Chi}(bx) dx}{3b^2} + \frac{2\int x \cosh(bx)\sinh(bx) dx}{3b} - \frac{4\int \frac{\cosh^2(bx)}{b} dx}{3b} \\
 &= \frac{4x \cosh(bx)\text{Chi}(bx)}{3b^2} + \frac{1}{3}x^3\text{Chi}(bx)^2 - \frac{4\text{Chi}(bx)\sinh(bx)}{3b^3} - \frac{2x^2\text{Chi}(bx)\sinh(bx)}{3b} \\
 &\quad + \frac{x \sinh^2(bx)}{3b^2} - \frac{\int \sinh^2(bx) dx}{3b^2} - \frac{4\int \cosh^2(bx) dx}{3b^2} + \frac{4\int \frac{\cosh(bx)\sinh(bx)}{bx} dx}{3b^2} \\
 &= \frac{4x \cosh(bx)\text{Chi}(bx)}{3b^2} + \frac{1}{3}x^3\text{Chi}(bx)^2 - \frac{5 \cosh(bx)\sinh(bx)}{6b^3} - \frac{4\text{Chi}(bx)\sinh(bx)}{3b^3} \\
 &\quad - \frac{2x^2\text{Chi}(bx)\sinh(bx)}{3b} + \frac{x \sinh^2(bx)}{3b^2} + \frac{4\int \frac{\cosh(bx)\sinh(bx)}{x} dx}{3b^3} + \frac{\int 1 dx}{6b^2} - \frac{2\int 1 dx}{3b^2} \\
 &= -\frac{x}{2b^2} + \frac{4x \cosh(bx)\text{Chi}(bx)}{3b^2} + \frac{1}{3}x^3\text{Chi}(bx)^2 - \frac{5 \cosh(bx)\sinh(bx)}{6b^3} \\
 &\quad - \frac{4\text{Chi}(bx)\sinh(bx)}{3b^3} - \frac{2x^2\text{Chi}(bx)\sinh(bx)}{3b} + \frac{x \sinh^2(bx)}{3b^2} + \frac{4\int \frac{\sinh(2bx)}{2x} dx}{3b^3}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{x}{2b^2} + \frac{4x \cosh(bx) \operatorname{Chi}(bx)}{3b^2} + \frac{1}{3}x^3 \operatorname{Chi}(bx)^2 - \frac{5 \cosh(bx) \sinh(bx)}{6b^3} \\
&\quad - \frac{4 \operatorname{Chi}(bx) \sinh(bx)}{3b^3} - \frac{2x^2 \operatorname{Chi}(bx) \sinh(bx)}{3b} + \frac{x \sinh^2(bx)}{3b^2} + \frac{2 \int \frac{\sinh(2bx)}{x} dx}{3b^3} \\
&= -\frac{x}{2b^2} + \frac{4x \cosh(bx) \operatorname{Chi}(bx)}{3b^2} + \frac{1}{3}x^3 \operatorname{Chi}(bx)^2 - \frac{5 \cosh(bx) \sinh(bx)}{6b^3} \\
&\quad - \frac{4 \operatorname{Chi}(bx) \sinh(bx)}{3b^3} - \frac{2x^2 \operatorname{Chi}(bx) \sinh(bx)}{3b} + \frac{x \sinh^2(bx)}{3b^2} + \frac{2 \operatorname{Shi}(2bx)}{3b^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.70

$$\int x^2 \operatorname{Chi}(bx)^2 dx = \frac{-8bx + 2bx \cosh(2bx) + 4b^3 x^3 \operatorname{Chi}(bx)^2 - 8 \operatorname{Chi}(bx) (-2bx \cosh(bx) + (2 + b^2 x^2) \sinh(bx)) - 5 \sinh(2bx)}{12b^3}$$

[In] Integrate[x^2*CoshIntegral[b*x]^2,x]

[Out] (-8*b*x + 2*b*x*Cosh[2*b*x] + 4*b^3*x^3*CoshIntegral[b*x]^2 - 8*CoshIntegral[b*x]*(-2*b*x*Cosh[b*x] + (2 + b^2*x^2)*Sinh[b*x]) - 5*Sinh[2*b*x] + 8*ShiIntegral[2*b*x])/(12*b^3)

Maple [A] (verified)

Time = 0.62 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.75

method	result
derivativedivides	$\frac{b^3 x^3 \operatorname{Chi}(bx)^2 - 2 \operatorname{Chi}(bx) \left(\frac{b^2 x^2 \sinh(bx)}{3} - \frac{2bx \cosh(bx)}{3} + \frac{2 \sinh(bx)}{3} \right) + \frac{bx \cosh(bx)^2}{3} - \frac{5 \cosh(bx) \sinh(bx)}{6} - \frac{5bx}{6} + \frac{2 \operatorname{Shi}(2bx)}{3}}{b^3}$
default	$\frac{b^3 x^3 \operatorname{Chi}(bx)^2 - 2 \operatorname{Chi}(bx) \left(\frac{b^2 x^2 \sinh(bx)}{3} - \frac{2bx \cosh(bx)}{3} + \frac{2 \sinh(bx)}{3} \right) + \frac{bx \cosh(bx)^2}{3} - \frac{5 \cosh(bx) \sinh(bx)}{6} - \frac{5bx}{6} + \frac{2 \operatorname{Shi}(2bx)}{3}}{b^3}$

[In] int(x^2*Chi(b*x)^2,x,method=_RETURNVERBOSE)

[Out] 1/b^3*(1/3*b^3*x^3*Chi(b*x)^2-2*Chi(b*x)*(1/3*b^2*x^2*sinh(b*x)-2/3*b*x*cosh(b*x)+2/3*sinh(b*x))+1/3*b*x*cosh(b*x)^2-5/6*cosh(b*x)*sinh(b*x)-5/6*b*x+2/3*Shi(2*b*x))

Fricas [F]

$$\int x^2 \operatorname{Chi}(bx)^2 dx = \int x^2 \operatorname{Chi}(bx)^2 dx$$

[In] integrate(x^2*Chi(b*x)^2,x, algorithm="fricas")

[Out] integral(x^2*cosh_integral(b*x)^2, x)

Sympy [F]

$$\int x^2 \operatorname{Chi}(bx)^2 dx = \int x^2 \operatorname{Chi}^2(bx) dx$$

[In] integrate(x**2*Chi(b*x)**2,x)

[Out] Integral(x**2*Chi(b*x)**2, x)

Maxima [F]

$$\int x^2 \operatorname{Chi}(bx)^2 dx = \int x^2 \operatorname{Chi}(bx)^2 dx$$

[In] integrate(x^2*Chi(b*x)^2,x, algorithm="maxima")

[Out] integrate(x^2*Chi(b*x)^2, x)

Giac [F]

$$\int x^2 \operatorname{Chi}(bx)^2 dx = \int x^2 \operatorname{Chi}(bx)^2 dx$$

[In] integrate(x^2*Chi(b*x)^2,x, algorithm="giac")

[Out] integrate(x^2*Chi(b*x)^2, x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \text{Chi}(bx)^2 dx = \int x^2 \text{coshint}(bx)^2 dx$$

```
[In] int(x^2*coshint(b*x)^2,x)
```

```
[Out] int(x^2*coshint(b*x)^2, x)
```

3.80 $\int x\text{Chi}(bx)^2 dx$

Optimal result	410
Rubi [A] (verified)	410
Mathematica [A] (verified)	412
Maple [A] (verified)	413
Fricas [F]	413
Sympy [F]	413
Maxima [F]	413
Giac [F]	414
Mupad [F(-1)]	414

Optimal result

Integrand size = 8, antiderivative size = 74

$$\int x\text{Chi}(bx)^2 dx = \frac{\cosh(bx)\text{Chi}(bx)}{b^2} + \frac{1}{2}x^2\text{Chi}(bx)^2 - \frac{\text{Chi}(2bx)}{2b^2} - \frac{\log(x)}{2b^2} - \frac{x\text{Chi}(bx)\sinh(bx)}{b} + \frac{\sinh^2(bx)}{2b^2}$$

[Out] $1/2*x^2*\text{Chi}(b*x)^2 - 1/2*\text{Chi}(2*b*x)/b^2 + \text{Chi}(b*x)*\cosh(b*x)/b^2 - 1/2*\ln(x)/b^2 - x*\text{Chi}(b*x)*\sinh(b*x)/b + 1/2*\sinh(b*x)^2/b^2$

Rubi [A] (verified)

Time = 0.07 (sec), antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {6672, 6678, 12, 2644, 30, 6682, 3393, 3382}

$$\int x\text{Chi}(bx)^2 dx = -\frac{\text{Chi}(2bx)}{2b^2} + \frac{\text{Chi}(bx)\cosh(bx)}{b^2} - \frac{\log(x)}{2b^2} + \frac{\sinh^2(bx)}{2b^2} + \frac{1}{2}x^2\text{Chi}(bx)^2 - \frac{x\text{Chi}(bx)\sinh(bx)}{b}$$

[In] `Int[x*CoshIntegral[b*x]^2,x]`

[Out] $(\text{Cosh}[b*x]*\text{CoshIntegral}[b*x])/b^2 + (x^2*\text{CoshIntegral}[b*x]^2)/2 - \text{CoshIntegral}[2*b*x]/(2*b^2) - \text{Log}[x]/(2*b^2) - (x*\text{CoshIntegral}[b*x]*\text{Sinh}[b*x])/b + \text{Sinh}[b*x]^2/(2*b^2)$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]`

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}/(m+1), x] /; \text{FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 2644

$\text{Int}[\cos[(e_.) + (f_.)(x_)]^{(n_.)} * ((a_.) * \sin[(e_.) + (f_.)(x_)]^{(m_.)}), x_Symbol] \rightarrow \text{Dist}[1/(a*f), \text{Subst}[\text{Int}[x^{m*(1-x^2/a^2)^{((n-1)/2)}], x], x, a*\sin[e+f*x]], x] /; \text{FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[(m-1)/2] \ \&\& \ \text{LtQ}[0, m, n])$

Rule 3382

$\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])*(f_.)(x_)]/((c_.) + (d_.)(x_)), x_Symbol] \rightarrow \text{Simp}[\text{CoshIntegral}[c*f*(fz/d) + f*fz*x]/d, x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \ \&\& \ \text{EqQ}[d*(e - \text{Pi}/2) - c*f*fz*I, 0]$

Rule 3393

$\text{Int}[(c_.) + (d_.)(x_)]^{(m_.)} * \sin[(e_.) + (f_.)(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(c + d*x)^m, \sin[e + f*x]^n, x], x] /; \text{FreeQ}[\{c, d, e, f, m\}, x] \ \&\& \ \text{IGtQ}[n, 1] \ \&\& \ (!\text{RationalQ}[m] \ || \ (\text{GeQ}[m, -1] \ \&\& \ \text{LtQ}[m, 1]))$

Rule 6672

$\text{Int}[\text{CoshIntegral}[(b_.)(x_)]^2 * (x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)} * (\text{CoshIntegral}[b*x]^2/(m+1)), x] - \text{Dist}[2/(m+1), \text{Int}[x^m * \text{Cosh}[b*x] * \text{CoshIntegral}[b*x], x], x] /; \text{FreeQ}[b, x] \ \&\& \ \text{IGtQ}[m, 0]$

Rule 6678

$\text{Int}[\text{Cosh}[(a_.) + (b_.)(x_)] * \text{CoshIntegral}[(c_.) + (d_.)(x_)] * ((e_.) + (f_.)(x_))^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(e + f*x)^m * \text{Sinh}[a + b*x] * (\text{CoshIntegral}[c + d*x]/b), x] + (-\text{Dist}[d/b, \text{Int}[(e + f*x)^m * \text{Sinh}[a + b*x] * (\text{Cosh}[c + d*x]/(c + d*x)), x], x] - \text{Dist}[f*(m/b), \text{Int}[(e + f*x)^{(m-1)} * \text{Sinh}[a + b*x] * \text{CoshIntegral}[c + d*x], x], x]) /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{IGtQ}[m, 0]$

Rule 6682

$\text{Int}[\text{CoshIntegral}[(c_.) + (d_.)(x_)] * \text{Sinh}[(a_.) + (b_.)(x_)], x_Symbol] \rightarrow \text{Simp}[\text{Cosh}[a + b*x] * (\text{CoshIntegral}[c + d*x]/b), x] - \text{Dist}[d/b, \text{Int}[\text{Cosh}[a + b*x] * (\text{Cosh}[c + d*x]/(c + d*x)), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x]$

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{2}x^2\text{Chi}(bx)^2 - \int x \cosh(bx)\text{Chi}(bx) dx \\
&= \frac{1}{2}x^2\text{Chi}(bx)^2 - \frac{x\text{Chi}(bx) \sinh(bx)}{b} + \frac{\int \text{Chi}(bx) \sinh(bx) dx}{b} + \int \frac{\cosh(bx) \sinh(bx)}{b} dx \\
&= \frac{\cosh(bx)\text{Chi}(bx)}{b^2} + \frac{1}{2}x^2\text{Chi}(bx)^2 - \frac{x\text{Chi}(bx) \sinh(bx)}{b} \\
&\quad - \frac{\int \frac{\cosh^2(bx)}{bx} dx}{b} + \frac{\int \cosh(bx) \sinh(bx) dx}{b} \\
&= \frac{\cosh(bx)\text{Chi}(bx)}{b^2} + \frac{1}{2}x^2\text{Chi}(bx)^2 - \frac{x\text{Chi}(bx) \sinh(bx)}{b} \\
&\quad - \frac{\int \frac{\cosh^2(bx)}{x} dx}{b^2} - \frac{\text{Subst}(\int x dx, x, i \sinh(bx))}{b^2} \\
&= \frac{\cosh(bx)\text{Chi}(bx)}{b^2} + \frac{1}{2}x^2\text{Chi}(bx)^2 - \frac{x\text{Chi}(bx) \sinh(bx)}{b} + \frac{\sinh^2(bx)}{2b^2} - \frac{\int \left(\frac{1}{2x} + \frac{\cosh(2bx)}{2x}\right) dx}{b^2} \\
&= \frac{\cosh(bx)\text{Chi}(bx)}{b^2} + \frac{1}{2}x^2\text{Chi}(bx)^2 - \frac{\log(x)}{2b^2} - \frac{x\text{Chi}(bx) \sinh(bx)}{b} + \frac{\sinh^2(bx)}{2b^2} - \frac{\int \frac{\cosh(2bx)}{x} dx}{2b^2} \\
&= \frac{\cosh(bx)\text{Chi}(bx)}{b^2} + \frac{1}{2}x^2\text{Chi}(bx)^2 - \frac{\text{Chi}(2bx)}{2b^2} - \frac{\log(x)}{2b^2} - \frac{x\text{Chi}(bx) \sinh(bx)}{b} + \frac{\sinh^2(bx)}{2b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.77

$$\begin{aligned}
&\int x\text{Chi}(bx)^2 dx \\
&= \frac{\cosh(2bx) + 2b^2x^2\text{Chi}(bx)^2 - 2\text{Chi}(2bx) - 2\log(x) + 4\text{Chi}(bx)(\cosh(bx) - bx \sinh(bx))}{4b^2}
\end{aligned}$$

[In] Integrate[x*CoshIntegral[b*x]^2,x]

[Out] (Cosh[2*b*x] + 2*b^2*x^2*CoshIntegral[b*x]^2 - 2*CoshIntegral[2*b*x] - 2*Log[x] + 4*CoshIntegral[b*x]*(Cosh[b*x] - b*x*Sinh[b*x]))/(4*b^2)

Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$\frac{b^2 x^2 \operatorname{Chi}(bx)^2 - 2 \operatorname{Chi}(bx) \left(\frac{bx \sinh(bx)}{2} - \frac{\cosh(bx)}{2} \right) + \frac{\cosh(bx)^2}{2} - \frac{\ln(bx)}{2} - \frac{\operatorname{Chi}(2bx)}{2}}{b^2}$	62
default	$\frac{b^2 x^2 \operatorname{Chi}(bx)^2 - 2 \operatorname{Chi}(bx) \left(\frac{bx \sinh(bx)}{2} - \frac{\cosh(bx)}{2} \right) + \frac{\cosh(bx)^2}{2} - \frac{\ln(bx)}{2} - \frac{\operatorname{Chi}(2bx)}{2}}{b^2}$	62

[In] int(x*Chi(b*x)^2,x,method=_RETURNVERBOSE)

[Out] $1/b^2*(1/2*b^2*x^2*Chi(b*x)^2-2*Chi(b*x)*(1/2*b*x*sinh(b*x)-1/2*cosh(b*x))+1/2*cosh(b*x)^2-1/2*ln(b*x)-1/2*Chi(2*b*x))$

Fricas [F]

$$\int x \operatorname{Chi}(bx)^2 dx = \int x \operatorname{Chi}(bx)^2 dx$$

[In] integrate(x*Chi(b*x)^2,x, algorithm="fricas")

[Out] integral(x*cosh_integral(b*x)^2, x)

Sympy [F]

$$\int x \operatorname{Chi}(bx)^2 dx = \int x \operatorname{Chi}^2(bx) dx$$

[In] integrate(x*Chi(b*x)**2,x)

[Out] Integral(x*Chi(b*x)**2, x)

Maxima [F]

$$\int x \operatorname{Chi}(bx)^2 dx = \int x \operatorname{Chi}(bx)^2 dx$$

[In] integrate(x*Chi(b*x)^2,x, algorithm="maxima")

[Out] integrate(x*Chi(b*x)^2, x)

Giac [F]

$$\int x\text{Chi}(bx)^2 dx = \int x\text{Chi}(bx)^2 dx$$

[In] integrate(x*Chi(b*x)^2,x, algorithm="giac")

[Out] integrate(x*Chi(b*x)^2, x)

Mupad [F(-1)]

Timed out.

$$\int x\text{Chi}(bx)^2 dx = \int x \coshint(bx)^2 dx$$

[In] int(x*coshint(b*x)^2,x)

[Out] int(x*coshint(b*x)^2, x)

3.81 $\int \text{Chi}(bx)^2 dx$

Optimal result	415
Rubi [A] (verified)	415
Mathematica [A] (verified)	417
Maple [A] (verified)	417
Fricas [F]	417
Sympy [F]	417
Maxima [F]	418
Giac [F]	418
Mupad [F(-1)]	418

Optimal result

Integrand size = 6, antiderivative size = 31

$$\int \text{Chi}(bx)^2 dx = x\text{Chi}(bx)^2 - \frac{2\text{Chi}(bx) \sinh(bx)}{b} + \frac{\text{Shi}(2bx)}{b}$$

[Out] $x*\text{Chi}(b*x)^2+\text{Shi}(2*b*x)/b-2*\text{Chi}(b*x)*\sinh(b*x)/b$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {6670, 6676, 12, 5556, 3379}

$$\int \text{Chi}(bx)^2 dx = x\text{Chi}(bx)^2 - \frac{2\text{Chi}(bx) \sinh(bx)}{b} + \frac{\text{Shi}(2bx)}{b}$$

[In] `Int[CoshIntegral[b*x]^2,x]`

[Out] $x*\text{CoshIntegral}[b*x]^2 - (2*\text{CoshIntegral}[b*x]*\text{Sinh}[b*x])/b + \text{SinhIntegral}[2*b*x]/b$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]`

Rule 3379

`Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f`

, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

Rule 6670

```
Int[CoshIntegral[(a_.) + (b_.)*(x_)]^2, x_Symbol] := Simp[(a + b*x)*(CoshIn
tegral[a + b*x]^2/b), x] - Dist[2, Int[Cosh[a + b*x]*CoshIntegral[a + b*x],
x], x] /; FreeQ[{a, b}, x]
```

Rule 6676

```
Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
Simp[Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= x\text{Chi}(bx)^2 - 2 \int \cosh(bx)\text{Chi}(bx) dx \\
&= x\text{Chi}(bx)^2 - \frac{2\text{Chi}(bx) \sinh(bx)}{b} + 2 \int \frac{\cosh(bx) \sinh(bx)}{bx} dx \\
&= x\text{Chi}(bx)^2 - \frac{2\text{Chi}(bx) \sinh(bx)}{b} + \frac{2 \int \frac{\cosh(bx) \sinh(bx)}{x} dx}{b} \\
&= x\text{Chi}(bx)^2 - \frac{2\text{Chi}(bx) \sinh(bx)}{b} + \frac{2 \int \frac{\sinh(2bx)}{2x} dx}{b} \\
&= x\text{Chi}(bx)^2 - \frac{2\text{Chi}(bx) \sinh(bx)}{b} + \frac{\int \frac{\sinh(2bx)}{x} dx}{b} \\
&= x\text{Chi}(bx)^2 - \frac{2\text{Chi}(bx) \sinh(bx)}{b} + \frac{\text{Shi}(2bx)}{b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \text{Chi}(bx)^2 dx = x\text{Chi}(bx)^2 - \frac{2\text{Chi}(bx) \sinh(bx)}{b} + \frac{\text{Shi}(2bx)}{b}$$

[In] Integrate[CoshIntegral[b*x]^2,x]

[Out] x*CoshIntegral[b*x]^2 - (2*CoshIntegral[b*x]*Sinh[b*x])/b + SinhIntegral[2*b*x]/b

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.97

method	result	size
derivativedivides	$\frac{\text{Chi}(bx)^2 bx - 2 \text{Chi}(bx) \sinh(bx) + \text{Shi}(2bx)}{b}$	30
default	$\frac{\text{Chi}(bx)^2 bx - 2 \text{Chi}(bx) \sinh(bx) + \text{Shi}(2bx)}{b}$	30

[In] int(Chi(b*x)^2,x,method=_RETURNVERBOSE)

[Out] 1/b*(Chi(b*x)^2*b*x-2*Chi(b*x)*sinh(b*x)+Shi(2*b*x))

Fricas [F]

$$\int \text{Chi}(bx)^2 dx = \int \text{Chi}(bx)^2 dx$$

[In] integrate(Chi(b*x)^2,x, algorithm="fricas")

[Out] integral(cosh_integral(b*x)^2, x)

Sympy [F]

$$\int \text{Chi}(bx)^2 dx = \int \text{Chi}^2(bx) dx$$

[In] integrate(Chi(b*x)**2,x)

[Out] Integral(Chi(b*x)**2, x)

Maxima [F]

$$\int \operatorname{Chi}(bx)^2 dx = \int \operatorname{Chi}(bx)^2 dx$$

[In] integrate(Chi(b*x)^2,x, algorithm="maxima")

[Out] integrate(Chi(b*x)^2, x)

Giac [F]

$$\int \operatorname{Chi}(bx)^2 dx = \int \operatorname{Chi}(bx)^2 dx$$

[In] integrate(Chi(b*x)^2,x, algorithm="giac")

[Out] integrate(Chi(b*x)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \operatorname{Chi}(bx)^2 dx = \int \operatorname{coshint}(bx)^2 dx$$

[In] int(coshint(b*x)^2,x)

[Out] int(coshint(b*x)^2, x)

3.82 $\int \frac{\text{Chi}(bx)^2}{x} dx$

Optimal result	419
Rubi [N/A]	419
Mathematica [N/A]	420
Maple [N/A] (verified)	420
Fricas [N/A]	420
Sympy [N/A]	420
Maxima [N/A]	421
Giac [N/A]	421
Mupad [N/A]	421

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\text{Chi}(bx)^2}{x} dx = \text{Int}\left(\frac{\text{Chi}(bx)^2}{x}, x\right)$$

[Out] CannotIntegrate(Chi(b*x)^2/x,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\text{Chi}(bx)^2}{x} dx = \int \frac{\text{Chi}(bx)^2}{x} dx$$

[In] Int[CoshIntegral[b*x]^2/x,x]

[Out] Defer[Int][CoshIntegral[b*x]^2/x, x]

Rubi steps

$$\text{integral} = \int \frac{\text{Chi}(bx)^2}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(bx)^2}{x} dx = \int \frac{\text{Chi}(bx)^2}{x} dx$$

`[In] Integrate[CoshIntegral[b*x]^2/x,x]``[Out] Integrate[CoshIntegral[b*x]^2/x, x]`**Maple [N/A] (verified)**

Not integrable

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx)^2}{x} dx$$

`[In] int(Chi(b*x)^2/x,x)``[Out] int(Chi(b*x)^2/x,x)`**Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(bx)^2}{x} dx = \int \frac{\text{Chi}(bx)^2}{x} dx$$

`[In] integrate(Chi(b*x)^2/x,x, algorithm="fricas")``[Out] integral(cosh_integral(b*x)^2/x, x)`**Sympy [N/A]**

Not integrable

Time = 0.50 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\text{Chi}(bx)^2}{x} dx = \int \frac{\text{Chi}^2(bx)}{x} dx$$

`[In] integrate(Chi(b*x)**2/x,x)``[Out] Integral(Chi(b*x)**2/x, x)`

Maxima [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(bx)^2}{x} dx = \int \frac{\text{Chi}(bx)^2}{x} dx$$

[In] integrate(Chi(b*x)^2/x,x, algorithm="maxima")

[Out] integrate(Chi(b*x)^2/x, x)

Giac [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(bx)^2}{x} dx = \int \frac{\text{Chi}(bx)^2}{x} dx$$

[In] integrate(Chi(b*x)^2/x,x, algorithm="giac")

[Out] integrate(Chi(b*x)^2/x, x)

Mupad [N/A]

Not integrable

Time = 4.87 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(bx)^2}{x} dx = \int \frac{\text{coshint}(bx)^2}{x} dx$$

[In] int(coshint(b*x)^2/x,x)

[Out] int(coshint(b*x)^2/x, x)

3.83 $\int \frac{\text{Chi}(bx)^2}{x^2} dx$

Optimal result	422
Rubi [N/A]	422
Mathematica [N/A]	423
Maple [N/A] (verified)	423
Fricas [N/A]	423
Sympy [N/A]	423
Maxima [N/A]	424
Giac [N/A]	424
Mupad [N/A]	424

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\text{Chi}(bx)^2}{x^2} dx = \text{Int}\left(\frac{\text{Chi}(bx)^2}{x^2}, x\right)$$

[Out] CannotIntegrate(Chi(b*x)^2/x^2,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\text{Chi}(bx)^2}{x^2} dx = \int \frac{\text{Chi}(bx)^2}{x^2} dx$$

[In] Int[CoshIntegral[b*x]^2/x^2,x]

[Out] Defer[Int][CoshIntegral[b*x]^2/x^2, x]

Rubi steps

$$\text{integral} = \int \frac{\text{Chi}(bx)^2}{x^2} dx$$

Mathematica [N/A]

Not integrable

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(bx)^2}{x^2} dx = \int \frac{\text{Chi}(bx)^2}{x^2} dx$$

[In] Integrate[CoshIntegral[b*x]^2/x^2,x]

[Out] Integrate[CoshIntegral[b*x]^2/x^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx)^2}{x^2} dx$$

[In] int(Chi(b*x)^2/x^2,x)

[Out] int(Chi(b*x)^2/x^2,x)

Fricas [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(bx)^2}{x^2} dx = \int \frac{\text{Chi}(bx)^2}{x^2} dx$$

[In] integrate(Chi(b*x)^2/x^2,x, algorithm="fricas")

[Out] integral(cosh_integral(b*x)^2/x^2, x)

Sympy [N/A]

Not integrable

Time = 0.42 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx)^2}{x^2} dx = \int \frac{\text{Chi}^2(bx)}{x^2} dx$$

[In] integrate(Chi(b*x)**2/x**2,x)

[Out] Integral(Chi(b*x)**2/x**2, x)

Maxima [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(bx)^2}{x^2} dx = \int \frac{\text{Chi}(bx)^2}{x^2} dx$$

[In] integrate(Chi(b*x)^2/x^2,x, algorithm="maxima")

[Out] integrate(Chi(b*x)^2/x^2, x)

Giac [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(bx)^2}{x^2} dx = \int \frac{\text{Chi}(bx)^2}{x^2} dx$$

[In] integrate(Chi(b*x)^2/x^2,x, algorithm="giac")

[Out] integrate(Chi(b*x)^2/x^2, x)

Mupad [N/A]

Not integrable

Time = 4.81 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(bx)^2}{x^2} dx = \int \frac{\text{coshint}(bx)^2}{x^2} dx$$

[In] int(coshint(b*x)^2/x^2,x)

[Out] int(coshint(b*x)^2/x^2, x)

3.84 $\int \frac{\text{Chi}(bx)^2}{x^3} dx$

Optimal result	425
Rubi [N/A]	425
Mathematica [N/A]	426
Maple [N/A] (verified)	426
Fricas [N/A]	426
Sympy [N/A]	426
Maxima [N/A]	427
Giac [N/A]	427
Mupad [N/A]	427

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\text{Chi}(bx)^2}{x^3} dx = \text{Int}\left(\frac{\text{Chi}(bx)^2}{x^3}, x\right)$$

[Out] CannotIntegrate(Chi(b*x)^2/x^3,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\text{Chi}(bx)^2}{x^3} dx = \int \frac{\text{Chi}(bx)^2}{x^3} dx$$

[In] Int[CoshIntegral[b*x]^2/x^3,x]

[Out] Defer[Int][CoshIntegral[b*x]^2/x^3, x]

Rubi steps

$$\text{integral} = \int \frac{\text{Chi}(bx)^2}{x^3} dx$$

Mathematica [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(bx)^2}{x^3} dx = \int \frac{\text{Chi}(bx)^2}{x^3} dx$$

`[In] Integrate[CoshIntegral[b*x]^2/x^3,x]``[Out] Integrate[CoshIntegral[b*x]^2/x^3, x]`**Maple [N/A] (verified)**

Not integrable

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx)^2}{x^3} dx$$

`[In] int(Chi(b*x)^2/x^3,x)``[Out] int(Chi(b*x)^2/x^3,x)`**Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(bx)^2}{x^3} dx = \int \frac{\text{Chi}(bx)^2}{x^3} dx$$

`[In] integrate(Chi(b*x)^2/x^3,x, algorithm="fricas")``[Out] integral(cosh_integral(b*x)^2/x^3, x)`**Sympy [N/A]**

Not integrable

Time = 0.46 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx)^2}{x^3} dx = \int \frac{\text{Chi}^2(bx)}{x^3} dx$$

`[In] integrate(Chi(b*x)**2/x**3,x)``[Out] Integral(Chi(b*x)**2/x**3, x)`

Maxima [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(bx)^2}{x^3} dx = \int \frac{\text{Chi}(bx)^2}{x^3} dx$$

[In] integrate(Chi(b*x)^2/x^3,x, algorithm="maxima")

[Out] integrate(Chi(b*x)^2/x^3, x)

Giac [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(bx)^2}{x^3} dx = \int \frac{\text{Chi}(bx)^2}{x^3} dx$$

[In] integrate(Chi(b*x)^2/x^3,x, algorithm="giac")

[Out] integrate(Chi(b*x)^2/x^3, x)

Mupad [N/A]

Not integrable

Time = 4.90 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(bx)^2}{x^3} dx = \int \frac{\text{coshint}(bx)^2}{x^3} dx$$

[In] int(coshint(b*x)^2/x^3,x)

[Out] int(coshint(b*x)^2/x^3, x)

3.85 $\int x^m \mathbf{Chi}(a + bx) dx$

Optimal result	428
Rubi [N/A]	428
Mathematica [N/A]	429
Maple [N/A] (verified)	429
Fricas [N/A]	429
Sympy [N/A]	429
Maxima [N/A]	430
Giac [N/A]	430
Mupad [N/A]	430

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int x^m \mathbf{Chi}(a + bx) dx = \frac{x^{1+m} \mathbf{Chi}(a + bx)}{1 + m} - \frac{b \operatorname{Int}\left(\frac{x^{1+m} \cosh(a+bx)}{a+bx}, x\right)}{1 + m}$$

[Out] $-b \cdot \text{CannotIntegrate}(x^{(1+m)} \cdot \cosh(b \cdot x + a) / (b \cdot x + a), x) / (1+m) + x^{(1+m)} \cdot \mathbf{Chi}(b \cdot x + a) / (1+m)$

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int x^m \mathbf{Chi}(a + bx) dx = \int x^m \mathbf{Chi}(a + bx) dx$$

[In] $\operatorname{Int}[x^m \cdot \text{CoshIntegral}[a + b \cdot x], x]$

[Out] $(x^{(1 + m)} \cdot \text{CoshIntegral}[a + b \cdot x]) / (1 + m) - (b \cdot \text{Defer}[\operatorname{Int}][x^{(1 + m)} \cdot \text{Cosh}[a + b \cdot x]) / (a + b \cdot x), x] / (1 + m)$

Rubi steps

$$\text{integral} = \frac{x^{1+m} \mathbf{Chi}(a + bx)}{1 + m} - \frac{b \int \frac{x^{1+m} \cosh(a+bx)}{a+bx} dx}{1 + m}$$

Mathematica [N/A]

Not integrable

Time = 6.30 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \text{Chi}(a + bx) dx = \int x^m \text{Chi}(a + bx) dx$$

[In] Integrate[x^m*CoshIntegral[a + b*x],x][Out] Integrate[x^m*CoshIntegral[a + b*x], x]**Maple [N/A] (verified)**

Not integrable

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int x^m \text{Chi}(bx + a) dx$$

[In] int(x^m*Chi(b*x+a),x)[Out] int(x^m*Chi(b*x+a),x)**Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \text{Chi}(a + bx) dx = \int x^m \text{Chi}(bx + a) dx$$

[In] integrate(x^m*Chi(b*x+a),x, algorithm="fricas")[Out] integral(x^m*cosh_integral(b*x + a), x)**Sympy [N/A]**

Not integrable

Time = 0.60 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int x^m \text{Chi}(a + bx) dx = \int x^m \text{Chi}(a + bx) dx$$

[In] integrate(x**m*Chi(b*x+a),x)

[Out] Integral(x**m*Chi(a + b*x), x)

Maxima [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Chi}(a + bx) dx = \int x^m \operatorname{Chi}(bx + a) dx$$

[In] integrate(x^m*Chi(b*x+a),x, algorithm="maxima")

[Out] integrate(x^m*Chi(b*x + a), x)

Giac [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Chi}(a + bx) dx = \int x^m \operatorname{Chi}(bx + a) dx$$

[In] integrate(x^m*Chi(b*x+a),x, algorithm="giac")

[Out] integrate(x^m*Chi(b*x + a), x)

Mupad [N/A]

Not integrable

Time = 4.75 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \operatorname{Chi}(a + bx) dx = \int x^m \operatorname{coshint}(a + bx) dx$$

[In] int(x^m*coshint(a + b*x),x)

[Out] int(x^m*coshint(a + b*x), x)

3.86 $\int x^3 \text{Chi}(a + bx) dx$

Optimal result	431
Rubi [A] (verified)	431
Mathematica [A] (verified)	433
Maple [A] (verified)	434
Fricas [F]	434
Sympy [F]	434
Maxima [F]	435
Giac [F]	435
Mupad [F(-1)]	435

Optimal result

Integrand size = 10, antiderivative size = 184

$$\int x^3 \text{Chi}(a + bx) dx = \frac{3 \cosh(a + bx)}{2b^4} + \frac{a^2 \cosh(a + bx)}{4b^4} - \frac{ax \cosh(a + bx)}{2b^3} + \frac{3x^2 \cosh(a + bx)}{4b^2} - \frac{a^4 \text{Chi}(a + bx)}{4b^4} + \frac{1}{4} x^4 \text{Chi}(a + bx) + \frac{a \sinh(a + bx)}{2b^4} + \frac{a^3 \sinh(a + bx)}{4b^4} - \frac{3x \sinh(a + bx)}{2b^3} - \frac{a^2 x \sinh(a + bx)}{4b^3} + \frac{ax^2 \sinh(a + bx)}{4b^2} - \frac{x^3 \sinh(a + bx)}{4b}$$

[Out] $-1/4*a^4*\text{Chi}(b*x+a)/b^4+1/4*x^4*\text{Chi}(b*x+a)+3/2*\cosh(b*x+a)/b^4+1/4*a^2*\cosh(b*x+a)/b^4-1/2*a*x*\cosh(b*x+a)/b^3+3/4*x^2*\cosh(b*x+a)/b^2+1/2*a*\sinh(b*x+a)/b^4+1/4*a^3*\sinh(b*x+a)/b^4-3/2*x*\sinh(b*x+a)/b^3-1/4*a^2*x*\sinh(b*x+a)/b^3+1/4*a*x^2*\sinh(b*x+a)/b^2-1/4*x^3*\sinh(b*x+a)/b$

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6668, 6874, 2717, 3377, 2718, 3382}

$$\int x^3 \text{Chi}(a + bx) dx = -\frac{a^4 \text{Chi}(a + bx)}{4b^4} + \frac{a^3 \sinh(a + bx)}{4b^4} + \frac{a^2 \cosh(a + bx)}{4b^4} - \frac{a^2 x \sinh(a + bx)}{4b^3} + \frac{a \sinh(a + bx)}{2b^4} + \frac{3 \cosh(a + bx)}{2b^4} - \frac{3x \sinh(a + bx)}{2b^3} - \frac{ax \cosh(a + bx)}{2b^3} + \frac{ax^2 \sinh(a + bx)}{4b^2} + \frac{3x^2 \cosh(a + bx)}{4b^2} + \frac{1}{4} x^4 \text{Chi}(a + bx) - \frac{x^3 \sinh(a + bx)}{4b}$$

[In] Int[x^3*CoshIntegral[a + b*x], x]

[Out] (3*Cosh[a + b*x])/(2*b^4) + (a^2*Cosh[a + b*x])/(4*b^4) - (a*x*Cosh[a + b*x])/(2*b^3) + (3*x^2*Cosh[a + b*x])/(4*b^2) - (a^4*CoshIntegral[a + b*x])/(4*b^4) + (x^4*CoshIntegral[a + b*x])/4 + (a*Sinh[a + b*x])/(2*b^4) + (a^3*Sinh[a + b*x])/(4*b^4) - (3*x*Sinh[a + b*x])/(2*b^3) - (a^2*x*Sinh[a + b*x])/(4*b^3) + (a*x^2*Sinh[a + b*x])/(4*b^2) - (x^3*Sinh[a + b*x])/(4*b)

Rule 2717

Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rule 2718

Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rule 3377

Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 6668

Int[CoshIntegral[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*(CoshIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Cosh[a + b*x]/(a + b*x)), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1]

Rule 6874

Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]

Rubi steps

$$\text{integral} = \frac{1}{4}x^4\text{Chi}(a + bx) - \frac{1}{4}b \int \frac{x^4 \cosh(a + bx)}{a + bx} dx$$

$$\begin{aligned}
&= \frac{1}{4}x^4\text{Chi}(a+bx) - \frac{1}{4}b \int \left(-\frac{a^3 \cosh(a+bx)}{b^4} + \frac{a^2x \cosh(a+bx)}{b^3} - \frac{ax^2 \cosh(a+bx)}{b^2} \right. \\
&\quad \left. + \frac{x^3 \cosh(a+bx)}{b} + \frac{a^4 \cosh(a+bx)}{b^4(a+bx)} \right) dx \\
&= \frac{1}{4}x^4\text{Chi}(a+bx) - \frac{1}{4} \int x^3 \cosh(a+bx) dx + \frac{a^3 \int \cosh(a+bx) dx}{4b^3} \\
&\quad - \frac{a^4 \int \frac{\cosh(a+bx)}{a+bx} dx}{4b^3} - \frac{a^2 \int x \cosh(a+bx) dx}{4b^2} + \frac{a \int x^2 \cosh(a+bx) dx}{4b} \\
&= -\frac{a^4\text{Chi}(a+bx)}{4b^4} + \frac{1}{4}x^4\text{Chi}(a+bx) + \frac{a^3 \sinh(a+bx)}{4b^4} - \frac{a^2x \sinh(a+bx)}{4b^3} + \frac{ax^2 \sinh(a+bx)}{4b^2} \\
&\quad - \frac{x^3 \sinh(a+bx)}{4b} + \frac{a^2 \int \sinh(a+bx) dx}{4b^3} - \frac{a \int x \sinh(a+bx) dx}{2b^2} + \frac{3 \int x^2 \sinh(a+bx) dx}{4b} \\
&= \frac{a^2 \cosh(a+bx)}{4b^4} - \frac{ax \cosh(a+bx)}{2b^3} + \frac{3x^2 \cosh(a+bx)}{4b^2} - \frac{a^4\text{Chi}(a+bx)}{4b^4} \\
&\quad + \frac{1}{4}x^4\text{Chi}(a+bx) + \frac{a^3 \sinh(a+bx)}{4b^4} - \frac{a^2x \sinh(a+bx)}{4b^3} + \frac{ax^2 \sinh(a+bx)}{4b^2} \\
&\quad - \frac{x^3 \sinh(a+bx)}{4b} + \frac{a \int \cosh(a+bx) dx}{2b^3} - \frac{3 \int x \cosh(a+bx) dx}{2b^2} \\
&= \frac{a^2 \cosh(a+bx)}{4b^4} - \frac{ax \cosh(a+bx)}{2b^3} + \frac{3x^2 \cosh(a+bx)}{4b^2} - \frac{a^4\text{Chi}(a+bx)}{4b^4} \\
&\quad + \frac{1}{4}x^4\text{Chi}(a+bx) + \frac{a \sinh(a+bx)}{2b^4} + \frac{a^3 \sinh(a+bx)}{4b^4} - \frac{3x \sinh(a+bx)}{2b^3} \\
&\quad - \frac{a^2x \sinh(a+bx)}{4b^3} + \frac{ax^2 \sinh(a+bx)}{4b^2} - \frac{x^3 \sinh(a+bx)}{4b} + \frac{3 \int \sinh(a+bx) dx}{2b^3} \\
&= \frac{3 \cosh(a+bx)}{2b^4} + \frac{a^2 \cosh(a+bx)}{4b^4} - \frac{ax \cosh(a+bx)}{2b^3} + \frac{3x^2 \cosh(a+bx)}{4b^2} \\
&\quad - \frac{a^4\text{Chi}(a+bx)}{4b^4} + \frac{1}{4}x^4\text{Chi}(a+bx) + \frac{a \sinh(a+bx)}{2b^4} + \frac{a^3 \sinh(a+bx)}{4b^4} \\
&\quad - \frac{3x \sinh(a+bx)}{2b^3} - \frac{a^2x \sinh(a+bx)}{4b^3} + \frac{ax^2 \sinh(a+bx)}{4b^2} - \frac{x^3 \sinh(a+bx)}{4b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.51

$$\int x^3\text{Chi}(a+bx) dx = \frac{(6+a^2-2abx+3b^2x^2)\cosh(a+bx) + (-a^4+b^4x^4)\text{Chi}(a+bx) + (2a+a^3-6bx-a^2bx+ab^2x^2-b^3x^3)\sinh(a+bx)}{4b^4}$$

[In] Integrate[x^3*CoshIntegral[a + b*x],x]

[Out] ((6 + a^2 - 2*a*b*x + 3*b^2*x^2)*Cosh[a + b*x] + (-a^4 + b^4*x^4)*CoshIntegral[a + b*x] + (2*a + a^3 - 6*b*x - a^2*b*x + a*b^2*x^2 - b^3*x^3)*Sinh[a + b*x])/(4*b^4)

Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.84

method	result
parts	$\frac{x^4 \operatorname{Chi}(bx+a)}{4} - \frac{a^4 \operatorname{Chi}(bx+a) - 4a^3 \sinh(bx+a) + 6a^2((bx+a) \sinh(bx+a) - \cosh(bx+a)) - 4a((bx+a)^2 \sinh(bx+a) - 2(bx+a) \cosh(bx+a))}{4}$
derivativedivides	$\frac{\operatorname{Chi}(bx+a)b^4x^4}{4} - \frac{a^4 \operatorname{Chi}(bx+a)}{4} + a^3 \sinh(bx+a) - \frac{3a^2((bx+a) \sinh(bx+a) - \cosh(bx+a))}{2} + a \frac{((bx+a)^2 \sinh(bx+a) - 2(bx+a) \cosh(bx+a))}{b^4}$
default	$\frac{\operatorname{Chi}(bx+a)b^4x^4}{4} - \frac{a^4 \operatorname{Chi}(bx+a)}{4} + a^3 \sinh(bx+a) - \frac{3a^2((bx+a) \sinh(bx+a) - \cosh(bx+a))}{2} + a \frac{((bx+a)^2 \sinh(bx+a) - 2(bx+a) \cosh(bx+a))}{b^4}$

[In] int(x^3*Chi(b*x+a),x,method=_RETURNVERBOSE)

```
[Out] 1/4*x^4*Chi(b*x+a)-1/4/b^4*(a^4*Chi(b*x+a)-4*a^3*sinh(b*x+a)+6*a^2*((b*x+a)*sinh(b*x+a)-cosh(b*x+a))-4*a*((b*x+a)^2*sinh(b*x+a)-2*(b*x+a)*cosh(b*x+a)+2*sinh(b*x+a))+(b*x+a)^3*sinh(b*x+a)-3*(b*x+a)^2*cosh(b*x+a)+6*(b*x+a)*sinh(b*x+a)-6*cosh(b*x+a))
```

Fricas [F]

$$\int x^3 \operatorname{Chi}(a + bx) dx = \int x^3 \operatorname{Chi}(bx + a) dx$$

[In] integrate(x^3*Chi(b*x+a),x, algorithm="fricas")

[Out] integral(x^3*cosh_integral(b*x + a), x)

Sympy [F]

$$\int x^3 \operatorname{Chi}(a + bx) dx = \int x^3 \operatorname{Chi}(a + bx) dx$$

[In] integrate(x**3*Chi(b*x+a),x)

[Out] Integral(x**3*Chi(a + b*x), x)

Maxima [F]

$$\int x^3 \operatorname{Chi}(a + bx) dx = \int x^3 \operatorname{Chi}(bx + a) dx$$

[In] integrate(x^3*Chi(b*x+a),x, algorithm="maxima")

[Out] integrate(x^3*Chi(b*x + a), x)

Giac [F]

$$\int x^3 \operatorname{Chi}(a + bx) dx = \int x^3 \operatorname{Chi}(bx + a) dx$$

[In] integrate(x^3*Chi(b*x+a),x, algorithm="giac")

[Out] integrate(x^3*Chi(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int x^3 \operatorname{Chi}(a + bx) dx = \int x^3 \operatorname{coshint}(a + bx) dx$$

[In] int(x^3*coshint(a + b*x),x)

[Out] int(x^3*coshint(a + b*x), x)

3.87 $\int x^2 \text{Chi}(a + bx) dx$

Optimal result	436
Rubi [A] (verified)	436
Mathematica [A] (verified)	438
Maple [A] (verified)	438
Fricas [F]	439
Sympy [F]	439
Maxima [F]	439
Giac [F]	439
Mupad [F(-1)]	440

Optimal result

Integrand size = 10, antiderivative size = 118

$$\int x^2 \text{Chi}(a + bx) dx = -\frac{a \cosh(a + bx)}{3b^3} + \frac{2x \cosh(a + bx)}{3b^2} + \frac{a^3 \text{Chi}(a + bx)}{3b^3} + \frac{1}{3} x^3 \text{Chi}(a + bx) - \frac{2 \sinh(a + bx)}{3b^3} - \frac{a^2 \sinh(a + bx)}{3b^3} + \frac{ax \sinh(a + bx)}{3b^2} - \frac{x^2 \sinh(a + bx)}{3b}$$

[Out] $\frac{1}{3} a^3 \text{Chi}(b*x+a)/b^3 + \frac{1}{3} x^3 \text{Chi}(b*x+a) - \frac{1}{3} a \cosh(b*x+a)/b^3 + \frac{2}{3} x \cosh(b*x+a)/b^2 - \frac{2}{3} \sinh(b*x+a)/b^3 - \frac{1}{3} a^2 \sinh(b*x+a)/b^3 + \frac{1}{3} a x \sinh(b*x+a)/b^2 - \frac{1}{3} x^2 \sinh(b*x+a)/b$

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6668, 6874, 2717, 3377, 2718, 3382}

$$\int x^2 \text{Chi}(a + bx) dx = \frac{a^3 \text{Chi}(a + bx)}{3b^3} - \frac{a^2 \sinh(a + bx)}{3b^3} - \frac{2 \sinh(a + bx)}{3b^3} - \frac{a \cosh(a + bx)}{3b^3} + \frac{ax \sinh(a + bx)}{3b^2} + \frac{2x \cosh(a + bx)}{3b^2} + \frac{1}{3} x^3 \text{Chi}(a + bx) - \frac{x^2 \sinh(a + bx)}{3b}$$

[In] `Int[x^2*CoshIntegral[a + b*x],x]`

[Out] $-\frac{1}{3} (a \text{Cosh}[a + b*x])/b^3 + (2*x*\text{Cosh}[a + b*x])/(3*b^2) + (a^3*\text{CoshIntegral}[a + b*x])/(3*b^3) + (x^3*\text{CoshIntegral}[a + b*x])/3 - (2*\text{Sinh}[a + b*x])/(3*b^3) - (a^2*\text{Sinh}[a + b*x])/(3*b^3) + (a*x*\text{Sinh}[a + b*x])/(3*b^2) - (x^2*\text{Sinh}[a + b*x])/(3*b)$

Rule 2717

```
Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]
```

Rule 2718

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ
[{c, d}, x]
```

Rule 3377

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-
(c + d*x)^m)*(Cos[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Co
s[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz
}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 6668

```
Int[CoshIntegral[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol]
:= Simp[(c + d*x)^(m + 1)*(CoshIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/
(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Cosh[a + b*x]/(a + b*x)), x], x] /; Fre
eQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{3}x^3\text{Chi}(a + bx) - \frac{1}{3}b \int \frac{x^3 \cosh(a + bx)}{a + bx} dx \\
&= \frac{1}{3}x^3\text{Chi}(a + bx) - \frac{1}{3}b \int \left(\frac{a^2 \cosh(a + bx)}{b^3} - \frac{ax \cosh(a + bx)}{b^2} + \frac{x^2 \cosh(a + bx)}{b} \right. \\
&\quad \left. - \frac{a^3 \cosh(a + bx)}{b^3(a + bx)} \right) dx \\
&= \frac{1}{3}x^3\text{Chi}(a + bx) - \frac{1}{3} \int x^2 \cosh(a + bx) dx - \frac{a^2 \int \cosh(a + bx) dx}{3b^2} \\
&\quad + \frac{a^3 \int \frac{\cosh(a + bx)}{a + bx} dx}{3b^2} + \frac{a \int x \cosh(a + bx) dx}{3b}
\end{aligned}$$

$$\begin{aligned}
&= \frac{a^3 \text{Chi}(a+bx)}{3b^3} + \frac{1}{3} x^3 \text{Chi}(a+bx) - \frac{a^2 \sinh(a+bx)}{3b^3} + \frac{ax \sinh(a+bx)}{3b^2} \\
&\quad - \frac{x^2 \sinh(a+bx)}{3b} - \frac{a \int \sinh(a+bx) dx}{3b^2} + \frac{2 \int x \sinh(a+bx) dx}{3b} \\
&= -\frac{a \cosh(a+bx)}{3b^3} + \frac{2x \cosh(a+bx)}{3b^2} + \frac{a^3 \text{Chi}(a+bx)}{3b^3} + \frac{1}{3} x^3 \text{Chi}(a+bx) \\
&\quad - \frac{a^2 \sinh(a+bx)}{3b^3} + \frac{ax \sinh(a+bx)}{3b^2} - \frac{x^2 \sinh(a+bx)}{3b} - \frac{2 \int \cosh(a+bx) dx}{3b^2} \\
&= -\frac{a \cosh(a+bx)}{3b^3} + \frac{2x \cosh(a+bx)}{3b^2} + \frac{a^3 \text{Chi}(a+bx)}{3b^3} + \frac{1}{3} x^3 \text{Chi}(a+bx) \\
&\quad - \frac{2 \sinh(a+bx)}{3b^3} - \frac{a^2 \sinh(a+bx)}{3b^3} + \frac{ax \sinh(a+bx)}{3b^2} - \frac{x^2 \sinh(a+bx)}{3b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.54

$$\begin{aligned}
&\int x^2 \text{Chi}(a+bx) dx \\
&= \\
&\quad - \frac{(a-2bx) \cosh(a+bx) - (a^3 + b^3 x^3) \text{Chi}(a+bx) + (2 + a^2 - abx + b^2 x^2) \sinh(a+bx)}{3b^3}
\end{aligned}$$

[In] Integrate[x^2*CoshIntegral[a + b*x],x]

[Out] -1/3*((a - 2*b*x)*Cosh[a + b*x] - (a^3 + b^3*x^3)*CoshIntegral[a + b*x] + (2 + a^2 - a*b*x + b^2*x^2)*Sinh[a + b*x])/b^3

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.85

method	result
parts	$\frac{x^3 \text{Chi}(bx+a)}{3} - \frac{-a^3 \text{Chi}(bx+a) + 3a^2 \sinh(bx+a) - 3a((bx+a) \sinh(bx+a) - \cosh(bx+a)) + (bx+a)^2 \sinh(bx+a) - 2(bx+a) \cosh(bx+a)}{3b^3}$
derivativedivides	$\frac{\text{Chi}(bx+a)b^3x^3 + a^3 \text{Chi}(bx+a) - a^2 \sinh(bx+a) + a((bx+a) \sinh(bx+a) - \cosh(bx+a)) - \frac{(bx+a)^2 \sinh(bx+a)}{3} + \frac{2(bx+a) \cosh(bx+a)}{3}}{b^3}$
default	$\frac{\text{Chi}(bx+a)b^3x^3 + a^3 \text{Chi}(bx+a) - a^2 \sinh(bx+a) + a((bx+a) \sinh(bx+a) - \cosh(bx+a)) - \frac{(bx+a)^2 \sinh(bx+a)}{3} + \frac{2(bx+a) \cosh(bx+a)}{3}}{b^3}$

[In] int(x^2*Chi(b*x+a),x,method=_RETURNVERBOSE)

[Out] 1/3*x^3*Chi(b*x+a)-1/3/b^3*(-a^3*Chi(b*x+a)+3*a^2*sinh(b*x+a)-3*a*((b*x+a)*sinh(b*x+a)-cosh(b*x+a))+(b*x+a)^2*sinh(b*x+a)-2*(b*x+a)*cosh(b*x+a)+2*sinh(b*x+a))

Fricas [F]

$$\int x^2 \operatorname{Chi}(a + bx) dx = \int x^2 \operatorname{Chi}(bx + a) dx$$

[In] `integrate(x^2*Chi(b*x+a),x, algorithm="fricas")`

[Out] `integral(x^2*cosh_integral(b*x + a), x)`

Sympy [F]

$$\int x^2 \operatorname{Chi}(a + bx) dx = \int x^2 \operatorname{Chi}(a + bx) dx$$

[In] `integrate(x**2*Chi(b*x+a),x)`

[Out] `Integral(x**2*Chi(a + b*x), x)`

Maxima [F]

$$\int x^2 \operatorname{Chi}(a + bx) dx = \int x^2 \operatorname{Chi}(bx + a) dx$$

[In] `integrate(x^2*Chi(b*x+a),x, algorithm="maxima")`

[Out] `integrate(x^2*Chi(b*x + a), x)`

Giac [F]

$$\int x^2 \operatorname{Chi}(a + bx) dx = \int x^2 \operatorname{Chi}(bx + a) dx$$

[In] `integrate(x^2*Chi(b*x+a),x, algorithm="giac")`

[Out] `integrate(x^2*Chi(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{Chi}(a + bx) dx = \int x^2 \operatorname{coshint}(a + bx) dx$$

```
[In] int(x^2*coshint(a + b*x),x)
```

```
[Out] int(x^2*coshint(a + b*x), x)
```


3.88 $\int x \operatorname{Chi}(a + bx) dx$

Optimal result	441
Rubi [A] (verified)	441
Mathematica [A] (verified)	443
Maple [A] (verified)	443
Fricas [F]	443
Sympy [F]	444
Maxima [F]	444
Giac [F]	444
Mupad [F(-1)]	444

Optimal result

Integrand size = 8, antiderivative size = 71

$$\int x \operatorname{Chi}(a + bx) dx = \frac{\cosh(a + bx)}{2b^2} - \frac{a^2 \operatorname{Chi}(a + bx)}{2b^2} + \frac{1}{2} x^2 \operatorname{Chi}(a + bx) + \frac{a \sinh(a + bx)}{2b^2} - \frac{x \sinh(a + bx)}{2b}$$

[Out] $-1/2*a^2*\operatorname{Chi}(b*x+a)/b^2+1/2*x^2*\operatorname{Chi}(b*x+a)+1/2*\cosh(b*x+a)/b^2+1/2*a*\sinh(b*x+a)/b^2-1/2*x*\sinh(b*x+a)/b$

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6668, 6874, 2717, 3377, 2718, 3382}

$$\int x \operatorname{Chi}(a + bx) dx = -\frac{a^2 \operatorname{Chi}(a + bx)}{2b^2} + \frac{a \sinh(a + bx)}{2b^2} + \frac{\cosh(a + bx)}{2b^2} + \frac{1}{2} x^2 \operatorname{Chi}(a + bx) - \frac{x \sinh(a + bx)}{2b}$$

[In] `Int[x*CoshIntegral[a + b*x],x]`

[Out] `Cosh[a + b*x]/(2*b^2) - (a^2*CoshIntegral[a + b*x])/(2*b^2) + (x^2*CoshIntegral[a + b*x])/2 + (a*Sinh[a + b*x])/(2*b^2) - (x*Sinh[a + b*x])/(2*b)`

Rule 2717

`Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;`
`FreeQ[{c, d}, x]`

Rule 2718

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ
[{c, d}, x]
```

Rule 3377

```
Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(
-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Co
s[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz
}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 6668

```
Int[CoshIntegral[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol]
:= Simp[(c + d*x)^(m + 1)*(CoshIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/
(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Cosh[a + b*x]/(a + b*x)), x], x] /; Fre
eQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{2}x^2\text{Chi}(a + bx) - \frac{1}{2}b \int \frac{x^2 \cosh(a + bx)}{a + bx} dx \\
&= \frac{1}{2}x^2\text{Chi}(a + bx) - \frac{1}{2}b \int \left(-\frac{a \cosh(a + bx)}{b^2} + \frac{x \cosh(a + bx)}{b} + \frac{a^2 \cosh(a + bx)}{b^2(a + bx)} \right) dx \\
&= \frac{1}{2}x^2\text{Chi}(a + bx) - \frac{1}{2} \int x \cosh(a + bx) dx + \frac{a \int \cosh(a + bx) dx}{2b} - \frac{a^2 \int \frac{\cosh(a + bx)}{a + bx} dx}{2b} \\
&= -\frac{a^2\text{Chi}(a + bx)}{2b^2} + \frac{1}{2}x^2\text{Chi}(a + bx) + \frac{a \sinh(a + bx)}{2b^2} - \frac{x \sinh(a + bx)}{2b} + \frac{\int \sinh(a + bx) dx}{2b} \\
&= \frac{\cosh(a + bx)}{2b^2} - \frac{a^2\text{Chi}(a + bx)}{2b^2} + \frac{1}{2}x^2\text{Chi}(a + bx) + \frac{a \sinh(a + bx)}{2b^2} - \frac{x \sinh(a + bx)}{2b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.66

$$\int x \operatorname{Chi}(a + bx) dx = \frac{\cosh(a + bx) + (-a^2 + b^2 x^2) \operatorname{Chi}(a + bx) + (a - bx) \sinh(a + bx)}{2b^2}$$

[In] Integrate[x*CoshIntegral[a + b*x],x]

[Out] (Cosh[a + b*x] + (-a^2 + b^2*x^2)*CoshIntegral[a + b*x] + (a - b*x)*Sinh[a + b*x])/(2*b^2)

Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.82

method	result	size
parts	$\frac{x^2 \operatorname{Chi}(bx+a)}{2} - \frac{a^2 \operatorname{Chi}(bx+a) - 2a \sinh(bx+a) + (bx+a) \sinh(bx+a) - \cosh(bx+a)}{2b^2}$	58
derivativedivides	$\frac{\operatorname{Chi}(bx+a) \left(-(bx+a)a + \frac{(bx+a)^2}{2} \right) + a \sinh(bx+a) - \frac{(bx+a) \sinh(bx+a)}{2} + \frac{\cosh(bx+a)}{2}}{b^2}$	60
default	$\frac{\operatorname{Chi}(bx+a) \left(-(bx+a)a + \frac{(bx+a)^2}{2} \right) + a \sinh(bx+a) - \frac{(bx+a) \sinh(bx+a)}{2} + \frac{\cosh(bx+a)}{2}}{b^2}$	60

[In] int(x*Chi(b*x+a),x,method=_RETURNVERBOSE)

[Out] 1/2*x^2*Chi(b*x+a)-1/2/b^2*(a^2*Chi(b*x+a)-2*a*sinh(b*x+a)+(b*x+a)*sinh(b*x+a)-cosh(b*x+a))

Fricas [F]

$$\int x \operatorname{Chi}(a + bx) dx = \int x \operatorname{Chi}(bx + a) dx$$

[In] integrate(x*Chi(b*x+a),x, algorithm="fricas")

[Out] integral(x*cosh_integral(b*x + a), x)

Sympy [F]

$$\int x \operatorname{Chi}(a + bx) dx = \int x \operatorname{Chi}(a + bx) dx$$

```
[In] integrate(x*Chi(b*x+a),x)
```

```
[Out] Integral(x*Chi(a + b*x), x)
```

Maxima [F]

$$\int x \operatorname{Chi}(a + bx) dx = \int x \operatorname{Chi}(bx + a) dx$$

```
[In] integrate(x*Chi(b*x+a),x, algorithm="maxima")
```

```
[Out] integrate(x*Chi(b*x + a), x)
```

Giac [F]

$$\int x \operatorname{Chi}(a + bx) dx = \int x \operatorname{Chi}(bx + a) dx$$

```
[In] integrate(x*Chi(b*x+a),x, algorithm="giac")
```

```
[Out] integrate(x*Chi(b*x + a), x)
```

Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int x \operatorname{Chi}(a + bx) dx \\ &= \frac{x^2 \operatorname{coshint}(a + bx)}{2} \\ &+ \frac{e^{-a-bx} (e^{2a+2bx} - a + a e^{2a+2bx} - 2a^2 \operatorname{coshint}(a+bx) e^{a+bx} + 1)}{4} + \frac{b e^{-a-bx} (x - x e^{2a+2bx})}{4} \\ & \qquad \qquad \qquad b^2 \end{aligned}$$

```
[In] int(x*coshint(a + b*x),x)
```

```
[Out] (x^2*coshint(a + b*x))/2 + ((exp(- a - b*x)*(exp(2*a + 2*b*x) - a + a*exp(2
*a + 2*b*x) - 2*a^2*coshint(a + b*x)*exp(a + b*x) + 1))/4 + (b*exp(- a - b*
x)*(x - x*exp(2*a + 2*b*x)))/4)/b^2
```

3.89 $\int \text{Chi}(a + bx) dx$

Optimal result	445
Rubi [A] (verified)	445
Mathematica [A] (verified)	446
Maple [A] (verified)	446
Fricas [F]	446
Sympy [F]	447
Maxima [F]	447
Giac [F]	447
Mupad [F(-1)]	447

Optimal result

Integrand size = 6, antiderivative size = 27

$$\int \text{Chi}(a + bx) dx = \frac{(a + bx)\text{Chi}(a + bx)}{b} - \frac{\sinh(a + bx)}{b}$$

[Out] (b*x+a)*Chi(b*x+a)/b-sinh(b*x+a)/b

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {6664}

$$\int \text{Chi}(a + bx) dx = \frac{(a + bx)\text{Chi}(a + bx)}{b} - \frac{\sinh(a + bx)}{b}$$

[In] Int[CoshIntegral[a + b*x],x]

[Out] ((a + b*x)*CoshIntegral[a + b*x])/b - Sinh[a + b*x]/b

Rule 6664

Int[CoshIntegral[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[(a + b*x)*(CoshIntegral[a + b*x]/b), x] - Simp[Sinh[a + b*x]/b, x] /; FreeQ[{a, b}, x]

Rubi steps

$$\text{integral} = \frac{(a + bx)\text{Chi}(a + bx)}{b} - \frac{\sinh(a + bx)}{b}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.56

$$\int \text{Chi}(a + bx) dx = \frac{a \text{Chi}(a + bx)}{b} + x \text{Chi}(a + bx) - \frac{\cosh(bx) \sinh(a)}{b} - \frac{\cosh(a) \sinh(bx)}{b}$$

[In] Integrate[CoshIntegral[a + b*x], x]

[Out] (a*CoshIntegral[a + b*x])/b + x*CoshIntegral[a + b*x] - (Cosh[b*x]*Sinh[a])/b - (Cosh[a]*Sinh[b*x])/b

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

method	result	size
derivativedivides	$\frac{\text{Chi}(bx+a)(bx+a) - \sinh(bx+a)}{b}$	26
default	$\frac{\text{Chi}(bx+a)(bx+a) - \sinh(bx+a)}{b}$	26
parts	$x \text{Chi}(bx+a) - \frac{-a \text{Chi}(bx+a) + \sinh(bx+a)}{b}$	31

[In] int(Chi(b*x+a), x, method=_RETURNVERBOSE)

[Out] 1/b*(Chi(b*x+a)*(b*x+a)-sinh(b*x+a))

Fricas [F]

$$\int \text{Chi}(a + bx) dx = \int \text{Chi}(bx + a) dx$$

[In] integrate(Chi(b*x+a), x, algorithm="fricas")

[Out] integral(cosh_integral(b*x + a), x)

Sympy [F]

$$\int \operatorname{Chi}(a + bx) dx = \int \operatorname{Chi}(a + bx) dx$$

[In] integrate(Chi(b*x+a),x)

[Out] Integral(Chi(a + b*x), x)

Maxima [F]

$$\int \operatorname{Chi}(a + bx) dx = \int \operatorname{Chi}(bx + a) dx$$

[In] integrate(Chi(b*x+a),x, algorithm="maxima")

[Out] integrate(Chi(b*x + a), x)

Giac [F]

$$\int \operatorname{Chi}(a + bx) dx = \int \operatorname{Chi}(bx + a) dx$$

[In] integrate(Chi(b*x+a),x, algorithm="giac")

[Out] integrate(Chi(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int \operatorname{Chi}(a + bx) dx = x \operatorname{coshint}(a + bx) - \frac{e^{a+bx}}{2b} + \frac{e^{-a-bx}}{2b} + \frac{a \operatorname{coshint}(a + bx)}{b}$$

[In] int(coshint(a + b*x),x)

[Out] x*coshint(a + b*x) - exp(a + b*x)/(2*b) + exp(- a - b*x)/(2*b) + (a*coshint(a + b*x))/b

3.90 $\int \frac{\mathbf{Chi}(a+bx)}{x} dx$

Optimal result	448
Rubi [N/A]	448
Mathematica [N/A]	449
Maple [N/A] (verified)	449
Fricas [N/A]	449
Sympy [N/A]	449
Maxima [N/A]	450
Giac [N/A]	450
Mupad [N/A]	450

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\mathbf{Chi}(a+bx)}{x} dx = \text{Int}\left(\frac{\mathbf{Chi}(a+bx)}{x}, x\right)$$

[Out] CannotIntegrate(Chi(b*x+a)/x,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\mathbf{Chi}(a+bx)}{x} dx = \int \frac{\mathbf{Chi}(a+bx)}{x} dx$$

[In] Int[CoshIntegral[a + b*x]/x,x]

[Out] Defer[Int][CoshIntegral[a + b*x]/x, x]

Rubi steps

$$\text{integral} = \int \frac{\mathbf{Chi}(a+bx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(a + bx)}{x} dx = \int \frac{\text{Chi}(a + bx)}{x} dx$$

[In] Integrate[CoshIntegral[a + b*x]/x,x]

[Out] Integrate[CoshIntegral[a + b*x]/x, x]

Maple [N/A] (verified)

Not integrable

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx + a)}{x} dx$$

[In] int(Chi(b*x+a)/x,x)

[Out] int(Chi(b*x+a)/x,x)

Fricas [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(a + bx)}{x} dx = \int \frac{\text{Chi}(bx + a)}{x} dx$$

[In] integrate(Chi(b*x+a)/x,x, algorithm="fricas")

[Out] integral(cosh_integral(b*x + a)/x, x)

Sympy [N/A]

Not integrable

Time = 0.38 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\text{Chi}(a + bx)}{x} dx = \int \frac{\text{Chi}(a + bx)}{x} dx$$

[In] integrate(Chi(b*x+a)/x,x)

[Out] Integral(Chi(a + b*x)/x, x)

Maxima [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(a + bx)}{x} dx = \int \frac{\text{Chi}(bx + a)}{x} dx$$

[In] integrate(Chi(b*x+a)/x,x, algorithm="maxima")

[Out] integrate(Chi(b*x + a)/x, x)

Giac [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(a + bx)}{x} dx = \int \frac{\text{Chi}(bx + a)}{x} dx$$

[In] integrate(Chi(b*x+a)/x,x, algorithm="giac")

[Out] integrate(Chi(b*x + a)/x, x)

Mupad [N/A]

Not integrable

Time = 4.74 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\text{Chi}(a + bx)}{x} dx = \int \frac{\text{coshint}(a + bx)}{x} dx$$

[In] int(coshint(a + b*x)/x,x)

[Out] int(coshint(a + b*x)/x, x)

3.91 $\int \frac{\text{Chi}(a+bx)}{x^2} dx$

Optimal result	451
Rubi [A] (verified)	451
Mathematica [A] (verified)	453
Maple [F]	453
Fricas [F]	453
Sympy [F]	453
Maxima [F]	454
Giac [F]	454
Mupad [F(-1)]	454

Optimal result

Integrand size = 10, antiderivative size = 46

$$\int \frac{\text{Chi}(a+bx)}{x^2} dx = \frac{b \cosh(a) \text{Chi}(bx)}{a} - \frac{b \text{Chi}(a+bx)}{a} - \frac{\text{Chi}(a+bx)}{x} + \frac{b \sinh(a) \text{Shi}(bx)}{a}$$

[Out] $-b \cdot \text{Chi}(b \cdot x + a) / a - \text{Chi}(b \cdot x + a) / x + b \cdot \text{Chi}(b \cdot x) \cdot \cosh(a) / a + b \cdot \text{Shi}(b \cdot x) \cdot \sinh(a) / a$

Rubi [A] (verified)

Time = 0.16 (sec), antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6668, 6874, 3384, 3379, 3382}

$$\int \frac{\text{Chi}(a+bx)}{x^2} dx = -\frac{b \text{Chi}(a+bx)}{a} - \frac{\text{Chi}(a+bx)}{x} + \frac{b \cosh(a) \text{Chi}(bx)}{a} + \frac{b \sinh(a) \text{Shi}(bx)}{a}$$

[In] `Int[CoshIntegral[a + b*x]/x^2,x]`

[Out] $(b \cdot \text{Cosh}[a] \cdot \text{CoshIntegral}[b \cdot x]) / a - (b \cdot \text{CoshIntegral}[a + b \cdot x]) / a - \text{CoshIntegral}[a + b \cdot x] / x + (b \cdot \text{Sinh}[a] \cdot \text{SinhIntegral}[b \cdot x]) / a$

Rule 3379

`Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`

Rule 3382

`Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}`

}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3384

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*
e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f
)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] &&
NeQ[d*e - c*f, 0]
```

Rule 6668

```
Int[CoshIntegral[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol]
:= Simp[(c + d*x)^(m + 1)*(CoshIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/
(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Cosh[a + b*x]/(a + b*x)), x], x] /; Fre
eQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Chi}(a + bx)}{x} + b \int \frac{\cosh(a + bx)}{x(a + bx)} dx \\
 &= -\frac{\text{Chi}(a + bx)}{x} + b \int \left(\frac{\cosh(a + bx)}{ax} - \frac{b \cosh(a + bx)}{a(a + bx)} \right) dx \\
 &= -\frac{\text{Chi}(a + bx)}{x} + \frac{b \int \frac{\cosh(a + bx)}{x} dx}{a} - \frac{b^2 \int \frac{\cosh(a + bx)}{a + bx} dx}{a} \\
 &= -\frac{b \text{Chi}(a + bx)}{a} - \frac{\text{Chi}(a + bx)}{x} + \frac{(b \cosh(a)) \int \frac{\cosh(bx)}{x} dx}{a} + \frac{(b \sinh(a)) \int \frac{\sinh(bx)}{x} dx}{a} \\
 &= \frac{b \cosh(a) \text{Chi}(bx)}{a} - \frac{b \text{Chi}(a + bx)}{a} - \frac{\text{Chi}(a + bx)}{x} + \frac{b \sinh(a) \text{Shi}(bx)}{a}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.85

$$\int \frac{\text{Chi}(a + bx)}{x^2} dx = \frac{bx \cosh(a)\text{Chi}(bx) - (a + bx)\text{Chi}(a + bx) + bx \sinh(a)\text{Shi}(bx)}{ax}$$

[In] Integrate[CoshIntegral[a + b*x]/x^2,x]

[Out] (b*x*Cosh[a]*CoshIntegral[b*x] - (a + b*x)*CoshIntegral[a + b*x] + b*x*Sinh[a]*SinhIntegral[b*x])/(a*x)

Maple [F]

$$\int \frac{\text{Chi}(bx + a)}{x^2} dx$$

[In] int(Chi(b*x+a)/x^2,x)

[Out] int(Chi(b*x+a)/x^2,x)

Fricas [F]

$$\int \frac{\text{Chi}(a + bx)}{x^2} dx = \int \frac{\text{Chi}(bx + a)}{x^2} dx$$

[In] integrate(Chi(b*x+a)/x^2,x, algorithm="fricas")

[Out] integral(cosh_integral(b*x + a)/x^2, x)

Sympy [F]

$$\int \frac{\text{Chi}(a + bx)}{x^2} dx = \int \frac{\text{Chi}(a + bx)}{x^2} dx$$

[In] integrate(Chi(b*x+a)/x**2,x)

[Out] Integral(Chi(a + b*x)/x**2, x)

Maxima [F]

$$\int \frac{\text{Chi}(a + bx)}{x^2} dx = \int \frac{\text{Chi}(bx + a)}{x^2} dx$$

[In] integrate(Chi(b*x+a)/x^2,x, algorithm="maxima")

[Out] integrate(Chi(b*x + a)/x^2, x)

Giac [F]

$$\int \frac{\text{Chi}(a + bx)}{x^2} dx = \int \frac{\text{Chi}(bx + a)}{x^2} dx$$

[In] integrate(Chi(b*x+a)/x^2,x, algorithm="giac")

[Out] integrate(Chi(b*x + a)/x^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\text{Chi}(a + bx)}{x^2} dx = \int \frac{\text{coshint}(a + bx)}{x^2} dx$$

[In] int(coshint(a + b*x)/x^2,x)

[Out] int(coshint(a + b*x)/x^2, x)

3.92 $\int \frac{\text{Chi}(a+bx)}{x^3} dx$

Optimal result	455
Rubi [A] (verified)	455
Mathematica [A] (verified)	457
Maple [F]	457
Fricas [F]	458
Sympy [F]	458
Maxima [F]	458
Giac [F]	458
Mupad [F(-1)]	459

Optimal result

Integrand size = 10, antiderivative size = 111

$$\int \frac{\text{Chi}(a+bx)}{x^3} dx = -\frac{b \cosh(a+bx)}{2ax} - \frac{b^2 \cosh(a)\text{Chi}(bx)}{2a^2} + \frac{b^2 \text{Chi}(a+bx)}{2a^2} - \frac{\text{Chi}(a+bx)}{2x^2} \\ + \frac{b^2 \text{Chi}(bx) \sinh(a)}{2a} + \frac{b^2 \cosh(a)\text{Shi}(bx)}{2a} - \frac{b^2 \sinh(a)\text{Shi}(bx)}{2a^2}$$

[Out] $1/2*b^2*\text{Chi}(b*x+a)/a^2-1/2*\text{Chi}(b*x+a)/x^2-1/2*b^2*\text{Chi}(b*x)*\cosh(a)/a^2-1/2*b*\cosh(b*x+a)/a/x+1/2*b^2*\cosh(a)*\text{Shi}(b*x)/a+1/2*b^2*\text{Chi}(b*x)*\sinh(a)/a-1/2*b^2*\text{Shi}(b*x)*\sinh(a)/a^2$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6668, 6874, 3378, 3384, 3379, 3382}

$$\int \frac{\text{Chi}(a+bx)}{x^3} dx = \frac{b^2 \text{Chi}(a+bx)}{2a^2} - \frac{b^2 \cosh(a)\text{Chi}(bx)}{2a^2} - \frac{b^2 \sinh(a)\text{Shi}(bx)}{2a^2} \\ + \frac{b^2 \sinh(a)\text{Chi}(bx)}{2a} + \frac{b^2 \cosh(a)\text{Shi}(bx)}{2a} - \frac{\text{Chi}(a+bx)}{2x^2} - \frac{b \cosh(a+bx)}{2ax}$$

[In] Int[CoshIntegral[a + b*x]/x^3,x]

[Out] $-1/2*(b*\text{Cosh}[a + b*x])/(a*x) - (b^2*\text{Cosh}[a]*\text{CoshIntegral}[b*x])/(2*a^2) + (b^2*\text{CoshIntegral}[a + b*x])/(2*a^2) - \text{CoshIntegral}[a + b*x]/(2*x^2) + (b^2*\text{CoshIntegral}[b*x]*\text{Sinh}[a])/(2*a) + (b^2*\text{Cosh}[a]*\text{SinhIntegral}[b*x])/(2*a) - (b^2*\text{Sinh}[a]*\text{SinhIntegral}[b*x])/(2*a^2)$

Rule 3378

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c
+ d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Dist[f/(d*(m + 1)), Int[(c
+ d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1
]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f
, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz
}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3384

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*
e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f
)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] &&
NeQ[d*e - c*f, 0]
```

Rule 6668

```
Int[CoshIntegral[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol]
:= Simp[(c + d*x)^(m + 1)*(CoshIntegral[a + b*x]/(d*(m + 1))), x] - Dist[b/
(d*(m + 1)), Int[(c + d*x)^(m + 1)*(Cosh[a + b*x]/(a + b*x)), x], x] /; Fre
eQ[{a, b, c, d, m}, x] && NeQ[m, -1]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Chi}(a + bx)}{2x^2} + \frac{1}{2}b \int \frac{\cosh(a + bx)}{x^2(a + bx)} dx \\
&= -\frac{\text{Chi}(a + bx)}{2x^2} + \frac{1}{2}b \int \left(\frac{\cosh(a + bx)}{ax^2} - \frac{b \cosh(a + bx)}{a^2x} + \frac{b^2 \cosh(a + bx)}{a^2(a + bx)} \right) dx \\
&= -\frac{\text{Chi}(a + bx)}{2x^2} + \frac{b \int \frac{\cosh(a + bx)}{x^2} dx}{2a} - \frac{b^2 \int \frac{\cosh(a + bx)}{x} dx}{2a^2} + \frac{b^3 \int \frac{\cosh(a + bx)}{a + bx} dx}{2a^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{b \cosh(a+bx)}{2ax} + \frac{b^2 \text{Chi}(a+bx)}{2a^2} - \frac{\text{Chi}(a+bx)}{2x^2} + \frac{b^2 \int \frac{\sinh(a+bx)}{x} dx}{2a} \\
&\quad - \frac{(b^2 \cosh(a)) \int \frac{\cosh(bx)}{x} dx}{2a^2} - \frac{(b^2 \sinh(a)) \int \frac{\sinh(bx)}{x} dx}{2a^2} \\
&= -\frac{b \cosh(a+bx)}{2ax} - \frac{b^2 \cosh(a) \text{Chi}(bx)}{2a^2} + \frac{b^2 \text{Chi}(a+bx)}{2a^2} - \frac{\text{Chi}(a+bx)}{2x^2} \\
&\quad - \frac{b^2 \sinh(a) \text{Shi}(bx)}{2a^2} + \frac{(b^2 \cosh(a)) \int \frac{\sinh(bx)}{x} dx}{2a} + \frac{(b^2 \sinh(a)) \int \frac{\cosh(bx)}{x} dx}{2a} \\
&= -\frac{b \cosh(a+bx)}{2ax} - \frac{b^2 \cosh(a) \text{Chi}(bx)}{2a^2} + \frac{b^2 \text{Chi}(a+bx)}{2a^2} - \frac{\text{Chi}(a+bx)}{2x^2} \\
&\quad + \frac{b^2 \text{Chi}(bx) \sinh(a)}{2a} + \frac{b^2 \cosh(a) \text{Shi}(bx)}{2a} - \frac{b^2 \sinh(a) \text{Shi}(bx)}{2a^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.72

$$\begin{aligned}
&\int \frac{\text{Chi}(a+bx)}{x^3} dx \\
&= \frac{(-a^2 + b^2 x^2) \text{Chi}(a+bx) + b^2 x^2 \text{Chi}(bx) (-\cosh(a) + a \sinh(a)) + bx(-a \cosh(a+bx) + bx(a \cosh(a) - \text{Chi}(a+bx)))}{2a^2 x^2}
\end{aligned}$$

[In] Integrate[CoshIntegral[a + b*x]/x^3,x]

[Out] ((-a^2 + b^2*x^2)*CoshIntegral[a + b*x] + b^2*x^2*CoshIntegral[b*x]*(-Cosh[a] + a*Sinh[a]) + b*x*(-(a*Cosh[a + b*x]) + b*x*(a*Cosh[a] - Sinh[a])*SinhIntegral[b*x]))/(2*a^2*x^2)

Maple [F]

$$\int \frac{\text{Chi}(bx+a)}{x^3} dx$$

[In] int(Chi(b*x+a)/x^3,x)

[Out] int(Chi(b*x+a)/x^3,x)

Fricas [F]

$$\int \frac{\text{Chi}(a + bx)}{x^3} dx = \int \frac{\text{Chi}(bx + a)}{x^3} dx$$

[In] integrate(Chi(b*x+a)/x^3,x, algorithm="fricas")

[Out] integral(cosh_integral(b*x + a)/x^3, x)

Sympy [F]

$$\int \frac{\text{Chi}(a + bx)}{x^3} dx = \int \frac{\text{Chi}(a + bx)}{x^3} dx$$

[In] integrate(Chi(b*x+a)/x**3,x)

[Out] Integral(Chi(a + b*x)/x**3, x)

Maxima [F]

$$\int \frac{\text{Chi}(a + bx)}{x^3} dx = \int \frac{\text{Chi}(bx + a)}{x^3} dx$$

[In] integrate(Chi(b*x+a)/x^3,x, algorithm="maxima")

[Out] integrate(Chi(b*x + a)/x^3, x)

Giac [F]

$$\int \frac{\text{Chi}(a + bx)}{x^3} dx = \int \frac{\text{Chi}(bx + a)}{x^3} dx$$

[In] integrate(Chi(b*x+a)/x^3,x, algorithm="giac")

[Out] integrate(Chi(b*x + a)/x^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\text{Chi}(a + bx)}{x^3} dx = \int \frac{\text{coshint}(a + bx)}{x^3} dx$$

```
[In] int(coshint(a + b*x)/x^3,x)
```

```
[Out] int(coshint(a + b*x)/x^3, x)
```

3.93 $\int x^m \mathbf{Chi}(a + bx)^2 dx$

Optimal result	460
Rubi [N/A]	460
Mathematica [N/A]	461
Maple [N/A] (verified)	461
Fricas [N/A]	461
Sympy [N/A]	461
Maxima [N/A]	462
Giac [N/A]	462
Mupad [N/A]	462

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int x^m \mathbf{Chi}(a + bx)^2 dx = \text{Int}(x^m \mathbf{Chi}(a + bx)^2, x)$$

[Out] CannotIntegrate(x^m*Chi(b*x+a)²,x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int x^m \mathbf{Chi}(a + bx)^2 dx = \int x^m \mathbf{Chi}(a + bx)^2 dx$$

[In] Int[x^m*CoshIntegral[a + b*x]²,x]

[Out] Defer[Int][x^m*CoshIntegral[a + b*x]², x]

Rubi steps

$$\text{integral} = \int x^m \mathbf{Chi}(a + bx)^2 dx$$

Mathematica [N/A]

Not integrable

Time = 2.41 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \operatorname{Chi}(a + bx)^2 dx = \int x^m \operatorname{Chi}(a + bx)^2 dx$$

[In] Integrate[x^m*CoshIntegral[a + b*x]^2,x]

[Out] Integrate[x^m*CoshIntegral[a + b*x]^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.22 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int x^m \operatorname{Chi}(bx + a)^2 dx$$

[In] int(x^m*Chi(b*x+a)^2,x)

[Out] int(x^m*Chi(b*x+a)^2,x)

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \operatorname{Chi}(a + bx)^2 dx = \int x^m \operatorname{Chi}(bx + a)^2 dx$$

[In] integrate(x^m*Chi(b*x+a)^2,x, algorithm="fricas")

[Out] integral(x^m*cosh_integral(b*x + a)^2, x)

Sympy [N/A]

Not integrable

Time = 1.25 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int x^m \operatorname{Chi}(a + bx)^2 dx = \int x^m \operatorname{Chi}^2(a + bx) dx$$

[In] integrate(x**m*Chi(b*x+a)**2,x)

[Out] Integral(x**m*Chi(a + b*x)**2, x)

Maxima [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \operatorname{Chi}(a + bx)^2 dx = \int x^m \operatorname{Chi}(bx + a)^2 dx$$

[In] integrate(x^m*Chi(b*x+a)^2,x, algorithm="maxima")

[Out] integrate(x^m*Chi(b*x + a)^2, x)

Giac [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \operatorname{Chi}(a + bx)^2 dx = \int x^m \operatorname{Chi}(bx + a)^2 dx$$

[In] integrate(x^m*Chi(b*x+a)^2,x, algorithm="giac")

[Out] integrate(x^m*Chi(b*x + a)^2, x)

Mupad [N/A]

Not integrable

Time = 4.79 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \operatorname{Chi}(a + bx)^2 dx = \int x^m \operatorname{coshint}(a + bx)^2 dx$$

[In] int(x^m*coshint(a + b*x)^2,x)

[Out] int(x^m*coshint(a + b*x)^2, x)

3.94 $\int x^2 \text{Chi}(a + bx)^2 dx$

Optimal result	463
Rubi [A] (verified)	464
Mathematica [A] (verified)	470
Maple [F]	470
Fricas [F]	471
Sympy [F]	471
Maxima [F]	471
Giac [F]	471
Mupad [F(-1)]	472

Optimal result

Integrand size = 12, antiderivative size = 327

$$\begin{aligned} \int x^2 \text{Chi}(a + bx)^2 dx = & -\frac{2x}{3b^2} - \frac{a \cosh(2a + 2bx)}{3b^3} + \frac{x \cosh(2a + 2bx)}{6b^2} \\ & - \frac{2a \cosh(a + bx) \text{Chi}(a + bx)}{3b^3} + \frac{4x \cosh(a + bx) \text{Chi}(a + bx)}{3b^2} \\ & + \frac{a^2(a + bx) \text{Chi}(a + bx)^2}{3b^3} - \frac{ax(a + bx) \text{Chi}(a + bx)^2}{3b^2} \\ & + \frac{x^2(a + bx) \text{Chi}(a + bx)^2}{3b} + \frac{a \text{Chi}(2a + 2bx)}{b^3} \\ & + \frac{a \log(a + bx)}{b^3} - \frac{2 \cosh(a + bx) \sinh(a + bx)}{3b^3} \\ & - \frac{4 \text{Chi}(a + bx) \sinh(a + bx)}{3b^3} - \frac{2a^2 \text{Chi}(a + bx) \sinh(a + bx)}{3b^3} \\ & + \frac{2ax \text{Chi}(a + bx) \sinh(a + bx)}{3b^2} - \frac{2x^2 \text{Chi}(a + bx) \sinh(a + bx)}{3b} \\ & - \frac{\sinh(2a + 2bx)}{12b^3} + \frac{2 \text{Shi}(2a + 2bx)}{3b^3} + \frac{a^2 \text{Shi}(2a + 2bx)}{b^3} \end{aligned}$$

```
[Out] -2/3*x/b^2+1/3*a^2*(b*x+a)*Chi(b*x+a)^2/b^3-1/3*a*x*(b*x+a)*Chi(b*x+a)^2/b^
2+1/3*x^2*(b*x+a)*Chi(b*x+a)^2/b+a*Chi(2*b*x+2*a)/b^3-2/3*a*Chi(b*x+a)*cosh
(b*x+a)/b^3+4/3*x*Chi(b*x+a)*cosh(b*x+a)/b^2-1/3*a*cosh(2*b*x+2*a)/b^3+1/6*
x*cosh(2*b*x+2*a)/b^2+a*ln(b*x+a)/b^3+2/3*Shi(2*b*x+2*a)/b^3+a^2*Shi(2*b*x+
2*a)/b^3-4/3*Chi(b*x+a)*sinh(b*x+a)/b^3-2/3*a^2*Chi(b*x+a)*sinh(b*x+a)/b^3+
2/3*a*x*Chi(b*x+a)*sinh(b*x+a)/b^2-2/3*x^2*Chi(b*x+a)*sinh(b*x+a)/b-2/3*cos
h(b*x+a)*sinh(b*x+a)/b^3-1/12*sinh(2*b*x+2*a)/b^3
```

Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.00, number of steps used = 39, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 1.583$, Rules used = {6674, 6678, 5736, 6873, 6874, 2718, 3377, 2717, 3379, 6684, 2715, 8, 3393, 3382, 6676, 5556, 12, 6682, 6670}

$$\int x^2 \text{Chi}(a + bx)^2 dx = \frac{a^2(a + bx)\text{Chi}(a + bx)^2}{3b^3} - \frac{2a^2\text{Chi}(a + bx)\sinh(a + bx)}{3b^3} + \frac{a^2\text{Shi}(2a + 2bx)}{b^3} + \frac{a\text{Chi}(2a + 2bx)}{b^3} - \frac{4\text{Chi}(a + bx)\sinh(a + bx)}{3b^3} - \frac{2a\text{Chi}(a + bx)\cosh(a + bx)}{3b^3} + \frac{2\text{Shi}(2a + 2bx)}{3b^3} + \frac{a\log(a + bx)}{b^3} - \frac{\sinh(2a + 2bx)}{12b^3} - \frac{a\cosh(2a + 2bx)}{3b^3} - \frac{2\sinh(a + bx)\cosh(a + bx)}{3b^3} - \frac{ax(a + bx)\text{Chi}(a + bx)^2}{3b^2} + \frac{2ax\text{Chi}(a + bx)\sinh(a + bx)}{3b^2} + \frac{4x\text{Chi}(a + bx)\cosh(a + bx)}{3b^2} + \frac{x\cosh(2a + 2bx)}{6b^2} + \frac{x^2(a + bx)\text{Chi}(a + bx)^2}{3b} - \frac{2x^2\text{Chi}(a + bx)\sinh(a + bx)}{3b} - \frac{2x}{3b^2}$$

[In] Int[x^2*CoshIntegral[a + b*x]^2,x]

[Out] (-2*x)/(3*b^2) - (a*Cosh[2*a + 2*b*x])/(3*b^3) + (x*Cosh[2*a + 2*b*x])/(6*b^2) - (2*a*Cosh[a + b*x]*CoshIntegral[a + b*x])/(3*b^3) + (4*x*Cosh[a + b*x]*CoshIntegral[a + b*x])/(3*b^2) + (a^2*(a + b*x)*CoshIntegral[a + b*x]^2)/(3*b^3) - (a*x*(a + b*x)*CoshIntegral[a + b*x]^2)/(3*b^2) + (x^2*(a + b*x)*CoshIntegral[a + b*x]^2)/(3*b) + (a*CoshIntegral[2*a + 2*b*x])/b^3 + (a*Log[a + b*x])/b^3 - (2*Cosh[a + b*x]*Sinh[a + b*x])/(3*b^3) - (4*CoshIntegral[a + b*x]*Sinh[a + b*x])/(3*b^3) - (2*a^2*CoshIntegral[a + b*x]*Sinh[a + b*x])/(3*b^3) + (2*a*x*CoshIntegral[a + b*x]*Sinh[a + b*x])/(3*b^2) - (2*x^2*CoshIntegral[a + b*x]*Sinh[a + b*x])/(3*b) - Sinh[2*a + 2*b*x]/(12*b^3) + (2*SinhIntegral[2*a + 2*b*x])/(3*b^3) + (a^2*SinhIntegral[2*a + 2*b*x])/b^3

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[

$(c + d*x)^{(n - 2)}, x], x] /;$ FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2717

Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]

Rule 2718

Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ
[{c, d}, x]

Rule 3377

Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-
(c + d*x)^m)*(Cos[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Co
s[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f
, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz
}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3393

Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := In
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))

Rule 5556

Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]

Rule 5736

Int[Cosh[w_]^(p_)*(u_)*Sinh[v_]^(p_), x_Symbol] := Dist[1/2^p, Int[u*Sinh[2*v]^p, x], x] /; EqQ[w, v] && IntegerQ[p]

Rule 6670

Int[CoshIntegral[(a_)+(b_)*(x_)]², x_Symbol] := Simp[(a+b*x)*(CoshIntegral[a+b*x]²/b), x] - Dist[2, Int[Cosh[a+b*x]*CoshIntegral[a+b*x], x], x] /; FreeQ[{a, b}, x]

Rule 6674

Int[CoshIntegral[(a_)+(b_)*(x_)]²*((c_)+(d_)*(x_))^(m_), x_Symbol] := Simp[(a+b*x)*(c+d*x)^m(CoshIntegral[a+b*x]²/(b*(m+1))), x] + (-Dist[2/(m+1), Int[(c+d*x)^m*Cosh[a+b*x]*CoshIntegral[a+b*x], x], x] + Dist[(b*c-a*d)*(m/(b*(m+1))), Int[(c+d*x)^(m-1)*CoshIntegral[a+b*x]², x], x]) /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0]

Rule 6676

Int[Cosh[(a_)+(b_)*(x_)]*CoshIntegral[(c_)+(d_)*(x_)], x_Symbol] := Simp[Sinh[a+b*x]*(CoshIntegral[c+d*x]/b), x] - Dist[d/b, Int[Sinh[a+b*x]*(Cosh[c+d*x]/(c+d*x)), x], x] /; FreeQ[{a, b, c, d}, x]

Rule 6678

Int[Cosh[(a_)+(b_)*(x_)]*CoshIntegral[(c_)+(d_)*(x_)]*((e_)+(f_)*(x_))^(m_), x_Symbol] := Simp[(e+f*x)^m*Sinh[a+b*x]*(CoshIntegral[c+d*x]/b), x] + (-Dist[d/b, Int[(e+f*x)^m*Sinh[a+b*x]*(Cosh[c+d*x]/(c+d*x)), x], x] - Dist[f*(m/b), Int[(e+f*x)^(m-1)*Sinh[a+b*x]*CoshIntegral[c+d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rule 6682

Int[CoshIntegral[(c_)+(d_)*(x_)]*Sinh[(a_)+(b_)*(x_)], x_Symbol] := Simp[Cosh[a+b*x]*(CoshIntegral[c+d*x]/b), x] - Dist[d/b, Int[Cosh[a+b*x]*(Cosh[c+d*x]/(c+d*x)), x], x] /; FreeQ[{a, b, c, d}, x]

Rule 6684

Int[CoshIntegral[(c_)+(d_)*(x_)]*((e_)+(f_)*(x_))^(m_)*Sinh[(a_)+(b_)*(x_)], x_Symbol] := Simp[(e+f*x)^m*Cosh[a+b*x]*(CoshIntegral[c+d*x]/b), x] + (-Dist[d/b, Int[(e+f*x)^m*Cosh[a+b*x]*(Cosh[c+d*x]/(c+d*x)), x], x] - Dist[f*(m/b), Int[(e+f*x)^(m-1)*Cosh[a+b*x]*CoshIntegral[c+d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rule 6873

Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v != u]

Rule 6874

Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x^2(a+bx)\text{Chi}(a+bx)^2}{3b} - \frac{2}{3} \int x^2 \cosh(a+bx)\text{Chi}(a+bx) dx \\
 &\quad - \frac{(2a) \int x\text{Chi}(a+bx)^2 dx}{3b} \\
 &= -\frac{ax(a+bx)\text{Chi}(a+bx)^2}{3b^2} + \frac{x^2(a+bx)\text{Chi}(a+bx)^2}{3b} - \frac{2x^2\text{Chi}(a+bx)\sinh(a+bx)}{3b} \\
 &\quad + \frac{2}{3} \int \frac{x^2 \cosh(a+bx)\sinh(a+bx)}{a+bx} dx + \frac{a^2 \int \text{Chi}(a+bx)^2 dx}{3b^2} \\
 &\quad + \frac{4 \int x\text{Chi}(a+bx)\sinh(a+bx) dx}{3b} + \frac{(2a) \int x \cosh(a+bx)\text{Chi}(a+bx) dx}{3b} \\
 &= \frac{4x \cosh(a+bx)\text{Chi}(a+bx)}{3b^2} + \frac{a^2(a+bx)\text{Chi}(a+bx)^2}{3b^3} \\
 &\quad - \frac{ax(a+bx)\text{Chi}(a+bx)^2}{3b^2} + \frac{x^2(a+bx)\text{Chi}(a+bx)^2}{3b} \\
 &\quad + \frac{2ax\text{Chi}(a+bx)\sinh(a+bx)}{3b^2} - \frac{2x^2\text{Chi}(a+bx)\sinh(a+bx)}{3b} \\
 &\quad + \frac{1}{3} \int \frac{x^2 \sinh(2(a+bx))}{a+bx} dx - \frac{4 \int \cosh(a+bx)\text{Chi}(a+bx) dx}{3b^2} \\
 &\quad - \frac{(2a) \int \text{Chi}(a+bx)\sinh(a+bx) dx}{3b^2} - \frac{(2a^2) \int \cosh(a+bx)\text{Chi}(a+bx) dx}{3b^2} \\
 &\quad - \frac{4 \int \frac{x \cosh^2(a+bx)}{a+bx} dx}{3b} - \frac{(2a) \int \frac{x \cosh(a+bx)\sinh(a+bx)}{a+bx} dx}{3b}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{2a \cosh(a+bx)\text{Chi}(a+bx)}{3b^3} + \frac{4x \cosh(a+bx)\text{Chi}(a+bx)}{3b^2} \\
&+ \frac{a^2(a+bx)\text{Chi}(a+bx)^2}{3b^3} - \frac{ax(a+bx)\text{Chi}(a+bx)^2}{3b^2} + \frac{x^2(a+bx)\text{Chi}(a+bx)^2}{3b} \\
&- \frac{4\text{Chi}(a+bx)\sinh(a+bx)}{3b^3} - \frac{2a^2\text{Chi}(a+bx)\sinh(a+bx)}{3b^3} \\
&+ \frac{2ax\text{Chi}(a+bx)\sinh(a+bx)}{3b^2} - \frac{2x^2\text{Chi}(a+bx)\sinh(a+bx)}{3b} \\
&+ \frac{1}{3} \int \frac{x^2 \sinh(2a+2bx)}{a+bx} dx + \frac{4}{3b^2} \int \frac{\cosh(a+bx)\sinh(a+bx)}{a+bx} dx \\
&+ \frac{(2a) \int \frac{\cosh^2(a+bx)}{a+bx} dx}{3b^2} + \frac{(2a^2) \int \frac{\cosh(a+bx)\sinh(a+bx)}{a+bx} dx}{3b^2} \\
&- \frac{4 \int \left(\frac{\cosh^2(a+bx)}{b} - \frac{a \cosh^2(a+bx)}{b(a+bx)} \right) dx}{3b} - \frac{a \int \frac{x \sinh(2(a+bx))}{a+bx} dx}{3b} \\
&= -\frac{2a \cosh(a+bx)\text{Chi}(a+bx)}{3b^3} + \frac{4x \cosh(a+bx)\text{Chi}(a+bx)}{3b^2} \\
&+ \frac{a^2(a+bx)\text{Chi}(a+bx)^2}{3b^3} - \frac{ax(a+bx)\text{Chi}(a+bx)^2}{3b^2} + \frac{x^2(a+bx)\text{Chi}(a+bx)^2}{3b} \\
&- \frac{4\text{Chi}(a+bx)\sinh(a+bx)}{3b^3} - \frac{2a^2\text{Chi}(a+bx)\sinh(a+bx)}{3b^3} \\
&+ \frac{2ax\text{Chi}(a+bx)\sinh(a+bx)}{3b^2} - \frac{2x^2\text{Chi}(a+bx)\sinh(a+bx)}{3b} \\
&+ \frac{1}{3} \int \left(-\frac{a \sinh(2a+2bx)}{b^2} + \frac{x \sinh(2a+2bx)}{b} + \frac{a^2 \sinh(2a+2bx)}{b^2(a+bx)} \right) dx \\
&- \frac{4 \int \cosh^2(a+bx) dx}{3b^2} + \frac{4 \int \frac{\sinh(2a+2bx)}{2(a+bx)} dx}{3b^2} + \frac{(2a) \int \left(\frac{1}{2(a+bx)} + \frac{\cosh(2a+2bx)}{2(a+bx)} \right) dx}{3b^2} \\
&+ \frac{(4a) \int \frac{\cosh^2(a+bx)}{a+bx} dx}{3b^2} + \frac{(2a^2) \int \frac{\sinh(2a+2bx)}{2(a+bx)} dx}{3b^2} - \frac{a \int \frac{x \sinh(2a+2bx)}{a+bx} dx}{3b}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2a \cosh(a+bx)\text{Chi}(a+bx)}{3b^3} + \frac{4x \cosh(a+bx)\text{Chi}(a+bx)}{3b^2} \\
&+ \frac{a^2(a+bx)\text{Chi}(a+bx)^2}{3b^3} - \frac{ax(a+bx)\text{Chi}(a+bx)^2}{3b^2} + \frac{x^2(a+bx)\text{Chi}(a+bx)^2}{3b} \\
&+ \frac{a \log(a+bx)}{3b^3} - \frac{2 \cosh(a+bx) \sinh(a+bx)}{3b^3} - \frac{4\text{Chi}(a+bx) \sinh(a+bx)}{3b^3} \\
&- \frac{2a^2\text{Chi}(a+bx) \sinh(a+bx)}{3b^3} + \frac{2ax\text{Chi}(a+bx) \sinh(a+bx)}{3b^2} \\
&- \frac{2x^2\text{Chi}(a+bx) \sinh(a+bx)}{3b} - \frac{2 \int 1 dx}{3b^2} + \frac{2 \int \frac{\sinh(2a+2bx)}{a+bx} dx}{3b^2} + \frac{a \int \frac{\cosh(2a+2bx)}{a+bx} dx}{3b^2} \\
&- \frac{a \int \sinh(2a+2bx) dx}{3b^2} + \frac{(4a) \int \left(\frac{1}{2(a+bx)} + \frac{\cosh(2a+2bx)}{2(a+bx)} \right) dx}{3b^2} + 2 \frac{a^2 \int \frac{\sinh(2a+2bx)}{a+bx} dx}{3b^2} \\
&+ \frac{\int x \sinh(2a+2bx) dx}{3b} - \frac{a \int \left(\frac{\sinh(2a+2bx)}{b} + \frac{a \sinh(2a+2bx)}{b(-a-bx)} \right) dx}{3b} \\
&= -\frac{2x}{3b^2} - \frac{a \cosh(2a+2bx)}{6b^3} + \frac{x \cosh(2a+2bx)}{6b^2} - \frac{2a \cosh(a+bx)\text{Chi}(a+bx)}{3b^3} \\
&+ \frac{4x \cosh(a+bx)\text{Chi}(a+bx)}{3b^2} + \frac{a^2(a+bx)\text{Chi}(a+bx)^2}{3b^3} \\
&- \frac{ax(a+bx)\text{Chi}(a+bx)^2}{3b^2} + \frac{x^2(a+bx)\text{Chi}(a+bx)^2}{3b} \\
&+ \frac{a\text{Chi}(2a+2bx)}{3b^3} + \frac{a \log(a+bx)}{b^3} - \frac{2 \cosh(a+bx) \sinh(a+bx)}{3b^3} \\
&- \frac{4\text{Chi}(a+bx) \sinh(a+bx)}{3b^3} - \frac{2a^2\text{Chi}(a+bx) \sinh(a+bx)}{3b^3} \\
&+ \frac{2ax\text{Chi}(a+bx) \sinh(a+bx)}{3b^2} - \frac{2x^2\text{Chi}(a+bx) \sinh(a+bx)}{3b} \\
&+ \frac{2\text{Shi}(2a+2bx)}{3b^3} + \frac{2a^2\text{Shi}(2a+2bx)}{3b^3} - \frac{\int \cosh(2a+2bx) dx}{6b^2} \\
&- \frac{a \int \sinh(2a+2bx) dx}{3b^2} + \frac{(2a) \int \frac{\cosh(2a+2bx)}{a+bx} dx}{3b^2} - \frac{a^2 \int \frac{\sinh(2a+2bx)}{-a-bx} dx}{3b^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2x}{3b^2} - \frac{a \cosh(2a + 2bx)}{3b^3} + \frac{x \cosh(2a + 2bx)}{6b^2} - \frac{2a \cosh(a + bx) \text{Chi}(a + bx)}{3b^3} \\
&+ \frac{4x \cosh(a + bx) \text{Chi}(a + bx)}{3b^2} + \frac{a^2(a + bx) \text{Chi}(a + bx)^2}{3b^3} \\
&- \frac{ax(a + bx) \text{Chi}(a + bx)^2}{3b^2} + \frac{x^2(a + bx) \text{Chi}(a + bx)^2}{3b} \\
&+ \frac{a \text{Chi}(2a + 2bx)}{b^3} + \frac{a \log(a + bx)}{b^3} - \frac{2 \cosh(a + bx) \sinh(a + bx)}{3b^3} \\
&- \frac{4 \text{Chi}(a + bx) \sinh(a + bx)}{3b^3} - \frac{2a^2 \text{Chi}(a + bx) \sinh(a + bx)}{3b^3} \\
&+ \frac{2ax \text{Chi}(a + bx) \sinh(a + bx)}{3b^2} - \frac{2x^2 \text{Chi}(a + bx) \sinh(a + bx)}{3b} \\
&- \frac{\sinh(2a + 2bx)}{12b^3} + \frac{2 \text{Shi}(2a + 2bx)}{3b^3} + \frac{a^2 \text{Shi}(2a + 2bx)}{b^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.79 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.48

$$\int x^2 \text{Chi}(a + bx)^2 dx = \frac{-8a - 8bx - 4a \cosh(2(a + bx)) + 2bx \cosh(2(a + bx)) + 4(a^3 + b^3x^3) \text{Chi}(a + bx)^2 + 12a \text{Chi}(2(a + bx)) -}{12b^3}$$

[In] Integrate[x^2*CoshIntegral[a + b*x]^2,x]

[Out] (-8*a - 8*b*x - 4*a*Cosh[2*(a + b*x)] + 2*b*x*Cosh[2*(a + b*x)] + 4*(a^3 + b^3*x^3)*CoshIntegral[a + b*x]^2 + 12*a*CoshIntegral[2*(a + b*x)] + 12*a*Log[a + b*x] - 8*CoshIntegral[a + b*x]*((a - 2*b*x)*Cosh[a + b*x] + (2 + a^2 - a*b*x + b^2*x^2)*Sinh[a + b*x]) - 5*Sinh[2*(a + b*x)] + 8*SinhIntegral[2*(a + b*x)] + 12*a^2*SinhIntegral[2*(a + b*x)])/(12*b^3)

Maple [F]

$$\int x^2 \text{Chi}(bx + a)^2 dx$$

[In] int(x^2*Chi(b*x+a)^2,x)

[Out] int(x^2*Chi(b*x+a)^2,x)

Fricas [F]

$$\int x^2 \operatorname{Chi}(a + bx)^2 dx = \int x^2 \operatorname{Chi}(bx + a)^2 dx$$

[In] `integrate(x^2*Chi(b*x+a)^2,x, algorithm="fricas")`

[Out] `integral(x^2*cosh_integral(b*x + a)^2, x)`

Sympy [F]

$$\int x^2 \operatorname{Chi}(a + bx)^2 dx = \int x^2 \operatorname{Chi}^2(a + bx) dx$$

[In] `integrate(x**2*Chi(b*x+a)**2,x)`

[Out] `Integral(x**2*Chi(a + b*x)**2, x)`

Maxima [F]

$$\int x^2 \operatorname{Chi}(a + bx)^2 dx = \int x^2 \operatorname{Chi}(bx + a)^2 dx$$

[In] `integrate(x^2*Chi(b*x+a)^2,x, algorithm="maxima")`

[Out] `integrate(x^2*Chi(b*x + a)^2, x)`

Giac [F]

$$\int x^2 \operatorname{Chi}(a + bx)^2 dx = \int x^2 \operatorname{Chi}(bx + a)^2 dx$$

[In] `integrate(x^2*Chi(b*x+a)^2,x, algorithm="giac")`

[Out] `integrate(x^2*Chi(b*x + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{Chi}(a + bx)^2 dx = \int x^2 \operatorname{coshint}(a + bx)^2 dx$$

```
[In] int(x^2*coshint(a + b*x)^2,x)
```

```
[Out] int(x^2*coshint(a + b*x)^2, x)
```


3.95 $\int x \operatorname{Chi}(a + bx)^2 dx$

Optimal result	473
Rubi [A] (verified)	473
Mathematica [A] (verified)	477
Maple [A] (verified)	477
Fricas [F]	478
Sympy [F]	478
Maxima [F]	478
Giac [F]	478
Mupad [F(-1)]	479

Optimal result

Integrand size = 10, antiderivative size = 154

$$\int x \operatorname{Chi}(a + bx)^2 dx = \frac{\cosh(2a + 2bx)}{4b^2} + \frac{\cosh(a + bx)\operatorname{Chi}(a + bx)}{b^2} - \frac{a(a + bx)\operatorname{Chi}(a + bx)^2}{2b^2} + \frac{x(a + bx)\operatorname{Chi}(a + bx)^2}{2b} - \frac{\operatorname{Chi}(2a + 2bx)}{2b^2} - \frac{\log(a + bx)}{2b^2} + \frac{a\operatorname{Chi}(a + bx)\sinh(a + bx)}{b^2} - \frac{x\operatorname{Chi}(a + bx)\sinh(a + bx)}{b} - \frac{a\operatorname{Shi}(2a + 2bx)}{b^2}$$

[Out] $-1/2*a*(b*x+a)*\operatorname{Chi}(b*x+a)^2/b^2+1/2*x*(b*x+a)*\operatorname{Chi}(b*x+a)^2/b-1/2*\operatorname{Chi}(2*b*x+2*a)/b^2+\operatorname{Chi}(b*x+a)*\cosh(b*x+a)/b^2+1/4*\cosh(2*b*x+2*a)/b^2-1/2*\ln(b*x+a)/b^2-a*\operatorname{Shi}(2*b*x+2*a)/b^2+a*\operatorname{Chi}(b*x+a)*\sinh(b*x+a)/b^2-x*\operatorname{Chi}(b*x+a)*\sinh(b*x+a)/b$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 1.400$, Rules used = {6674, 6678, 5736, 6873, 6874, 2718, 3379, 6682, 3393, 3382, 6670, 6676, 5556, 12}

$$\int x \operatorname{Chi}(a + bx)^2 dx = -\frac{a(a + bx)\operatorname{Chi}(a + bx)^2}{2b^2} - \frac{\operatorname{Chi}(2a + 2bx)}{2b^2} + \frac{a\operatorname{Chi}(a + bx)\sinh(a + bx)}{b^2} + \frac{\operatorname{Chi}(a + bx)\cosh(a + bx)}{b^2} - \frac{a\operatorname{Shi}(2a + 2bx)}{b^2} - \frac{\log(a + bx)}{2b^2} + \frac{\cosh(2a + 2bx)}{4b^2} + \frac{x(a + bx)\operatorname{Chi}(a + bx)^2}{2b} - \frac{x\operatorname{Chi}(a + bx)\sinh(a + bx)}{b}$$

[In] Int[x*CoshIntegral[a + b*x]^2,x]

[Out] Cosh[2*a + 2*b*x]/(4*b^2) + (Cosh[a + b*x]*CoshIntegral[a + b*x])/b^2 - (a*(a + b*x)*CoshIntegral[a + b*x]^2)/(2*b^2) + (x*(a + b*x)*CoshIntegral[a + b*x]^2)/(2*b) - CoshIntegral[2*a + 2*b*x]/(2*b^2) - Log[a + b*x]/(2*b^2) + (a*CoshIntegral[a + b*x]*Sinh[a + b*x])/b^2 - (x*CoshIntegral[a + b*x]*Sinh[a + b*x])/b - (a*SinhIntegral[2*a + 2*b*x])/b^2

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 2718

Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3393

Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))

Rule 5556

Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]

Rule 5736

Int[Cosh[w_]^(p_.)*(u_.)*Sinh[v_]^(p_.), x_Symbol] := Dist[1/2^p, Int[u*Sinh[2*v]^p, x], x] /; EqQ[w, v] && IntegerQ[p]

Rule 6670

```
Int[CoshIntegral[(a_.) + (b_.)*(x_)]^2, x_Symbol] := Simp[(a + b*x)*(CoshIntegral[a + b*x]^2/b), x] - Dist[2, Int[Cosh[a + b*x]*CoshIntegral[a + b*x], x], x] /; FreeQ[{a, b}, x]
```

Rule 6674

```
Int[CoshIntegral[(a_.) + (b_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(a + b*x)*(c + d*x)^m*(CoshIntegral[a + b*x]^2/(b*(m + 1))), x] + (-Dist[2/(m + 1), Int[(c + d*x)^m*Cosh[a + b*x]*CoshIntegral[a + b*x], x], x] + Dist[(b*c - a*d)*(m/(b*(m + 1))), Int[(c + d*x)^(m - 1)*CoshIntegral[a + b*x]^2, x], x]) /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0]
```

Rule 6676

```
Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6678

```
Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^m*Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*CoshIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6682

```
Int[CoshIntegral[(c_.) + (d_.)*(x_)]*Sinh[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6873

```
Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v != u]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x(a+bx)\text{Chi}(a+bx)^2}{2b} - \frac{a \int \text{Chi}(a+bx)^2 dx}{2b} - \int x \cosh(a+bx)\text{Chi}(a+bx) dx \\
&= -\frac{a(a+bx)\text{Chi}(a+bx)^2}{2b^2} + \frac{x(a+bx)\text{Chi}(a+bx)^2}{2b} \\
&\quad - \frac{x\text{Chi}(a+bx) \sinh(a+bx)}{b} + \frac{\int \text{Chi}(a+bx) \sinh(a+bx) dx}{b} \\
&\quad + \frac{a \int \cosh(a+bx)\text{Chi}(a+bx) dx}{b} + \int \frac{x \cosh(a+bx) \sinh(a+bx)}{a+bx} dx \\
&= \frac{\cosh(a+bx)\text{Chi}(a+bx)}{b^2} - \frac{a(a+bx)\text{Chi}(a+bx)^2}{2b^2} + \frac{x(a+bx)\text{Chi}(a+bx)^2}{2b} \\
&\quad + \frac{a\text{Chi}(a+bx) \sinh(a+bx)}{b^2} - \frac{x\text{Chi}(a+bx) \sinh(a+bx)}{b} \\
&\quad + \frac{1}{2} \int \frac{x \sinh(2(a+bx))}{a+bx} dx - \frac{\int \frac{\cosh^2(a+bx)}{a+bx} dx}{b} - \frac{a \int \frac{\cosh(a+bx) \sinh(a+bx)}{a+bx} dx}{b} \\
&= \frac{\cosh(a+bx)\text{Chi}(a+bx)}{b^2} - \frac{a(a+bx)\text{Chi}(a+bx)^2}{2b^2} + \frac{x(a+bx)\text{Chi}(a+bx)^2}{2b} \\
&\quad + \frac{a\text{Chi}(a+bx) \sinh(a+bx)}{b^2} - \frac{x\text{Chi}(a+bx) \sinh(a+bx)}{b} \\
&\quad + \frac{1}{2} \int \frac{x \sinh(2a+2bx)}{a+bx} dx - \frac{\int \left(\frac{1}{2(a+bx)} + \frac{\cosh(2a+2bx)}{2(a+bx)} \right) dx}{b} - \frac{a \int \frac{\sinh(2a+2bx)}{2(a+bx)} dx}{b} \\
&= \frac{\cosh(a+bx)\text{Chi}(a+bx)}{b^2} - \frac{a(a+bx)\text{Chi}(a+bx)^2}{2b^2} \\
&\quad + \frac{x(a+bx)\text{Chi}(a+bx)^2}{2b} - \frac{\log(a+bx)}{2b^2} + \frac{a\text{Chi}(a+bx) \sinh(a+bx)}{b^2} \\
&\quad - \frac{x\text{Chi}(a+bx) \sinh(a+bx)}{b} + \frac{1}{2} \int \left(\frac{\sinh(2a+2bx)}{b} \right. \\
&\quad \quad \left. + \frac{a \sinh(2a+2bx)}{b(-a-bx)} \right) dx - \frac{\int \frac{\cosh(2a+2bx)}{a+bx} dx}{2b} - \frac{a \int \frac{\sinh(2a+2bx)}{a+bx} dx}{2b} \\
&= \frac{\cosh(a+bx)\text{Chi}(a+bx)}{b^2} - \frac{a(a+bx)\text{Chi}(a+bx)^2}{2b^2} \\
&\quad + \frac{x(a+bx)\text{Chi}(a+bx)^2}{2b} - \frac{\text{Chi}(2a+2bx)}{2b^2} - \frac{\log(a+bx)}{2b^2} \\
&\quad + \frac{a\text{Chi}(a+bx) \sinh(a+bx)}{b^2} - \frac{x\text{Chi}(a+bx) \sinh(a+bx)}{b} \\
&\quad - \frac{a\text{Shi}(2a+2bx)}{2b^2} + \frac{\int \sinh(2a+2bx) dx}{2b} + \frac{a \int \frac{\sinh(2a+2bx)}{-a-bx} dx}{2b}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\cosh(2a + 2bx)}{4b^2} + \frac{\cosh(a + bx)\text{Chi}(a + bx)}{b^2} - \frac{a(a + bx)\text{Chi}(a + bx)^2}{2b^2} \\
&+ \frac{x(a + bx)\text{Chi}(a + bx)^2}{2b} - \frac{\text{Chi}(2a + 2bx)}{2b^2} - \frac{\log(a + bx)}{2b^2} \\
&+ \frac{a\text{Chi}(a + bx)\sinh(a + bx)}{b^2} - \frac{x\text{Chi}(a + bx)\sinh(a + bx)}{b} - \frac{a\text{Shi}(2a + 2bx)}{b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.62

$$\int x\text{Chi}(a + bx)^2 dx = \frac{\cosh(2(a + bx)) - 2(a^2 - b^2x^2)\text{Chi}(a + bx)^2 - 2\text{Chi}(2(a + bx)) - 2\log(a + bx) + 4\text{Chi}(a + bx)(\cosh(a + bx) + (a - bx)\text{Sinh}[a + bx]) - 4a*\text{SinhIntegral}[2*(a + b*x)]}{4b^2}$$

[In] Integrate[x*CoshIntegral[a + b*x]^2,x]

[Out] (Cosh[2*(a + b*x)] - 2*(a^2 - b^2*x^2)*CoshIntegral[a + b*x]^2 - 2*CoshIntegral[2*(a + b*x)] - 2*Log[a + b*x] + 4*CoshIntegral[a + b*x]*(Cosh[a + b*x] + (a - b*x)*Sinh[a + b*x]) - 4*a*SinhIntegral[2*(a + b*x)])/(4*b^2)

Maple [A] (verified)

Time = 0.92 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.73

method	result
derivativedivides	$\frac{\text{Chi}(bx+a)^2 \left(-(bx+a)a + \frac{(bx+a)^2}{2} \right) - 2 \text{Chi}(bx+a) \left(-a \sinh(bx+a) + \frac{(bx+a)\sinh(bx+a)}{2} - \frac{\cosh(bx+a)}{2} \right) - a \text{Shi}(2bx+2a) + \dots}{b^2}$
default	$\frac{\text{Chi}(bx+a)^2 \left(-(bx+a)a + \frac{(bx+a)^2}{2} \right) - 2 \text{Chi}(bx+a) \left(-a \sinh(bx+a) + \frac{(bx+a)\sinh(bx+a)}{2} - \frac{\cosh(bx+a)}{2} \right) - a \text{Shi}(2bx+2a) + \dots}{b^2}$

[In] int(x*Chi(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/b^2*(Chi(b*x+a)^2*(-(b*x+a)*a+1/2*(b*x+a)^2)-2*Chi(b*x+a)*(-a*sinh(b*x+a)+1/2*(b*x+a)*sinh(b*x+a)-1/2*cosh(b*x+a))-a*Shi(2*b*x+2*a)+1/2*cosh(b*x+a)^2-1/2*ln(b*x+a)-1/2*Chi(2*b*x+2*a))

Fricas [F]

$$\int x\text{Chi}(a + bx)^2 dx = \int x\text{Chi}(bx + a)^2 dx$$

```
[In] integrate(x*Chi(b*x+a)^2,x, algorithm="fricas")
```

```
[Out] integral(x*cosh_integral(b*x + a)^2, x)
```

Sympy [F]

$$\int x\text{Chi}(a + bx)^2 dx = \int x\text{Chi}^2(a + bx) dx$$

```
[In] integrate(x*Chi(b*x+a)**2,x)
```

```
[Out] Integral(x*Chi(a + b*x)**2, x)
```

Maxima [F]

$$\int x\text{Chi}(a + bx)^2 dx = \int x\text{Chi}(bx + a)^2 dx$$

```
[In] integrate(x*Chi(b*x+a)^2,x, algorithm="maxima")
```

```
[Out] integrate(x*Chi(b*x + a)^2, x)
```

Giac [F]

$$\int x\text{Chi}(a + bx)^2 dx = \int x\text{Chi}(bx + a)^2 dx$$

```
[In] integrate(x*Chi(b*x+a)^2,x, algorithm="giac")
```

```
[Out] integrate(x*Chi(b*x + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int x\text{Chi}(a + bx)^2 dx = \int x \text{coshint}(a + bx)^2 dx$$

```
[In] int(x*coshint(a + b*x)^2,x)
```

```
[Out] int(x*coshint(a + b*x)^2, x)
```

3.96 $\int \text{Chi}(a + bx)^2 dx$

Optimal result	480
Rubi [A] (verified)	480
Mathematica [A] (verified)	482
Maple [A] (verified)	482
Fricas [F]	482
Sympy [F]	482
Maxima [F]	483
Giac [F]	483
Mupad [F(-1)]	483

Optimal result

Integrand size = 8, antiderivative size = 48

$$\int \text{Chi}(a + bx)^2 dx = \frac{(a + bx)\text{Chi}(a + bx)^2}{b} - \frac{2\text{Chi}(a + bx)\sinh(a + bx)}{b} + \frac{\text{Shi}(2a + 2bx)}{b}$$

[Out] (b*x+a)*Chi(b*x+a)^2/b+Shi(2*b*x+2*a)/b-2*Chi(b*x+a)*sinh(b*x+a)/b

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {6670, 6676, 5556, 12, 3379}

$$\int \text{Chi}(a + bx)^2 dx = \frac{(a + bx)\text{Chi}(a + bx)^2}{b} - \frac{2\text{Chi}(a + bx)\sinh(a + bx)}{b} + \frac{\text{Shi}(2a + 2bx)}{b}$$

[In] Int[CoshIntegral[a + b*x]^2,x]

[Out] ((a + b*x)*CoshIntegral[a + b*x]^2)/b - (2*CoshIntegral[a + b*x]*Sinh[a + b*x])/b + SinhIntegral[2*a + 2*b*x]/b

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f

, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 5556

Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]

Rule 6670

Int[CoshIntegral[(a_.) + (b_.)*(x_)]^2, x_Symbol] := Simp[(a + b*x)*(CoshIntegral[a + b*x]^2/b), x] - Dist[2, Int[Cosh[a + b*x]*CoshIntegral[a + b*x], x], x] /; FreeQ[{a, b}, x]

Rule 6676

Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{(a + bx)\text{Chi}(a + bx)^2}{b} - 2 \int \cosh(a + bx)\text{Chi}(a + bx) dx \\
 &= \frac{(a + bx)\text{Chi}(a + bx)^2}{b} - \frac{2\text{Chi}(a + bx)\sinh(a + bx)}{b} + 2 \int \frac{\cosh(a + bx)\sinh(a + bx)}{a + bx} dx \\
 &= \frac{(a + bx)\text{Chi}(a + bx)^2}{b} - \frac{2\text{Chi}(a + bx)\sinh(a + bx)}{b} + 2 \int \frac{\sinh(2a + 2bx)}{2(a + bx)} dx \\
 &= \frac{(a + bx)\text{Chi}(a + bx)^2}{b} - \frac{2\text{Chi}(a + bx)\sinh(a + bx)}{b} + \int \frac{\sinh(2a + 2bx)}{a + bx} dx \\
 &= \frac{(a + bx)\text{Chi}(a + bx)^2}{b} - \frac{2\text{Chi}(a + bx)\sinh(a + bx)}{b} + \frac{\text{Shi}(2a + 2bx)}{b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.85

$$\int \operatorname{Chi}(a + bx)^2 dx = \frac{(a + bx)\operatorname{Chi}(a + bx)^2 - 2\operatorname{Chi}(a + bx) \sinh(a + bx) + \operatorname{Shi}(2(a + bx))}{b}$$

[In] Integrate[CoshIntegral[a + b*x]^2,x]

[Out] ((a + b*x)*CoshIntegral[a + b*x]^2 - 2*CoshIntegral[a + b*x]*Sinh[a + b*x] + SinhIntegral[2*(a + b*x)])/b

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.90

method	result	size
derivativedivides	$\frac{\operatorname{Chi}(bx+a)^2(bx+a) - 2 \operatorname{Chi}(bx+a) \sinh(bx+a) + \operatorname{Shi}(2bx+2a)}{b}$	43
default	$\frac{\operatorname{Chi}(bx+a)^2(bx+a) - 2 \operatorname{Chi}(bx+a) \sinh(bx+a) + \operatorname{Shi}(2bx+2a)}{b}$	43

[In] int(Chi(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/b*(Chi(b*x+a)^2*(b*x+a)-2*Chi(b*x+a)*sinh(b*x+a)+Shi(2*b*x+2*a))

Fricas [F]

$$\int \operatorname{Chi}(a + bx)^2 dx = \int \operatorname{Chi}(bx + a)^2 dx$$

[In] integrate(Chi(b*x+a)^2,x, algorithm="fricas")

[Out] integral(cosh_integral(b*x + a)^2, x)

Sympy [F]

$$\int \operatorname{Chi}(a + bx)^2 dx = \int \operatorname{Chi}^2(a + bx) dx$$

[In] integrate(Chi(b*x+a)**2,x)

[Out] Integral(Chi(a + b*x)**2, x)

Maxima [F]

$$\int \operatorname{Chi}(a + bx)^2 dx = \int \operatorname{Chi}(bx + a)^2 dx$$

[In] integrate(Chi(b*x+a)^2,x, algorithm="maxima")

[Out] integrate(Chi(b*x + a)^2, x)

Giac [F]

$$\int \operatorname{Chi}(a + bx)^2 dx = \int \operatorname{Chi}(bx + a)^2 dx$$

[In] integrate(Chi(b*x+a)^2,x, algorithm="giac")

[Out] integrate(Chi(b*x + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \operatorname{Chi}(a + bx)^2 dx = \int \operatorname{coshint}(a + bx)^2 dx$$

[In] int(coshint(a + b*x)^2,x)

[Out] int(coshint(a + b*x)^2, x)

3.97 $\int \frac{\text{Chi}(a+bx)^2}{x} dx$

Optimal result	484
Rubi [N/A]	484
Mathematica [N/A]	485
Maple [N/A] (verified)	485
Fricas [N/A]	485
Sympy [N/A]	485
Maxima [N/A]	486
Giac [N/A]	486
Mupad [N/A]	486

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\text{Chi}(a+bx)^2}{x} dx = \text{Int}\left(\frac{\text{Chi}(a+bx)^2}{x}, x\right)$$

[Out] CannotIntegrate(Chi(b*x+a)^2/x,x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\text{Chi}(a+bx)^2}{x} dx = \int \frac{\text{Chi}(a+bx)^2}{x} dx$$

[In] Int[CoshIntegral[a + b*x]^2/x,x]

[Out] Defer[Int][CoshIntegral[a + b*x]^2/x, x]

Rubi steps

$$\text{integral} = \int \frac{\text{Chi}(a+bx)^2}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(a + bx)^2}{x} dx = \int \frac{\text{Chi}(a + bx)^2}{x} dx$$

[In] Integrate[CoshIntegral[a + b*x]^2/x, x]

[Out] Integrate[CoshIntegral[a + b*x]^2/x, x]

Maple [N/A] (verified)

Not integrable

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx + a)^2}{x} dx$$

[In] int(Chi(b*x+a)^2/x, x)

[Out] int(Chi(b*x+a)^2/x, x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(a + bx)^2}{x} dx = \int \frac{\text{Chi}(bx + a)^2}{x} dx$$

[In] integrate(Chi(b*x+a)^2/x, x, algorithm="fricas")

[Out] integral(cosh_integral(b*x + a)^2/x, x)

Sympy [N/A]

Not integrable

Time = 0.49 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\text{Chi}(a + bx)^2}{x} dx = \int \frac{\text{Chi}^2(a + bx)}{x} dx$$

[In] integrate(Chi(b*x+a)**2/x, x)

[Out] Integral(Chi(a + b*x)**2/x, x)

Maxima [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(a + bx)^2}{x} dx = \int \frac{\text{Chi}(bx + a)^2}{x} dx$$

[In] integrate(Chi(b*x+a)^2/x,x, algorithm="maxima")

[Out] integrate(Chi(b*x + a)^2/x, x)

Giac [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(a + bx)^2}{x} dx = \int \frac{\text{Chi}(bx + a)^2}{x} dx$$

[In] integrate(Chi(b*x+a)^2/x,x, algorithm="giac")

[Out] integrate(Chi(b*x + a)^2/x, x)

Mupad [N/A]

Not integrable

Time = 4.80 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(a + bx)^2}{x} dx = \int \frac{\text{coshint}(a + bx)^2}{x} dx$$

[In] int(coshint(a + b*x)^2/x,x)

[Out] int(coshint(a + b*x)^2/x, x)

3.98 $\int \frac{\text{Chi}(a+bx)^2}{x^2} dx$

Optimal result	487
Rubi [N/A]	487
Mathematica [N/A]	488
Maple [N/A] (verified)	488
Fricas [N/A]	488
Sympy [N/A]	488
Maxima [N/A]	489
Giac [N/A]	489
Mupad [N/A]	489

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\text{Chi}(a+bx)^2}{x^2} dx = \text{Int}\left(\frac{\text{Chi}(a+bx)^2}{x^2}, x\right)$$

[Out] CannotIntegrate(Chi(b*x+a)^2/x^2,x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\text{Chi}(a+bx)^2}{x^2} dx = \int \frac{\text{Chi}(a+bx)^2}{x^2} dx$$

[In] Int[CoshIntegral[a + b*x]^2/x^2,x]

[Out] Defer[Int][CoshIntegral[a + b*x]^2/x^2, x]

Rubi steps

$$\text{integral} = \int \frac{\text{Chi}(a+bx)^2}{x^2} dx$$

Mathematica [N/A]

Not integrable

Time = 0.77 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(a + bx)^2}{x^2} dx = \int \frac{\text{Chi}(a + bx)^2}{x^2} dx$$

[In] Integrate[CoshIntegral[a + b*x]^2/x^2,x]

[Out] Integrate[CoshIntegral[a + b*x]^2/x^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.21 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx + a)^2}{x^2} dx$$

[In] int(Chi(b*x+a)^2/x^2,x)

[Out] int(Chi(b*x+a)^2/x^2,x)

Fricas [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(a + bx)^2}{x^2} dx = \int \frac{\text{Chi}(bx + a)^2}{x^2} dx$$

[In] integrate(Chi(b*x+a)^2/x^2,x, algorithm="fricas")

[Out] integral(cosh_integral(b*x + a)^2/x^2, x)

Sympy [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(a + bx)^2}{x^2} dx = \int \frac{\text{Chi}^2(a + bx)}{x^2} dx$$

[In] integrate(Chi(b*x+a)**2/x**2,x)

[Out] Integral(Chi(a + b*x)**2/x**2, x)

Maxima [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(a + bx)^2}{x^2} dx = \int \frac{\text{Chi}(bx + a)^2}{x^2} dx$$

[In] integrate(Chi(b*x+a)^2/x^2,x, algorithm="maxima")

[Out] integrate(Chi(b*x + a)^2/x^2, x)

Giac [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(a + bx)^2}{x^2} dx = \int \frac{\text{Chi}(bx + a)^2}{x^2} dx$$

[In] integrate(Chi(b*x+a)^2/x^2,x, algorithm="giac")

[Out] integrate(Chi(b*x + a)^2/x^2, x)

Mupad [N/A]

Not integrable

Time = 4.90 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(a + bx)^2}{x^2} dx = \int \frac{\text{coshint}(a + bx)^2}{x^2} dx$$

[In] int(coshint(a + b*x)^2/x^2,x)

[Out] int(coshint(a + b*x)^2/x^2, x)

3.99 $\int \frac{\text{Chi}(a+bx)^2}{x^3} dx$

Optimal result	490
Rubi [N/A]	490
Mathematica [N/A]	491
Maple [N/A] (verified)	491
Fricas [N/A]	491
Sympy [N/A]	491
Maxima [N/A]	492
Giac [N/A]	492
Mupad [N/A]	492

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\text{Chi}(a+bx)^2}{x^3} dx = \text{Int}\left(\frac{\text{Chi}(a+bx)^2}{x^3}, x\right)$$

[Out] CannotIntegrate(Chi(b*x+a)^2/x^3,x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\text{Chi}(a+bx)^2}{x^3} dx = \int \frac{\text{Chi}(a+bx)^2}{x^3} dx$$

[In] Int[CoshIntegral[a + b*x]^2/x^3,x]

[Out] Defer[Int][CoshIntegral[a + b*x]^2/x^3, x]

Rubi steps

$$\text{integral} = \int \frac{\text{Chi}(a+bx)^2}{x^3} dx$$

Mathematica [N/A]

Not integrable

Time = 1.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(a + bx)^2}{x^3} dx = \int \frac{\text{Chi}(a + bx)^2}{x^3} dx$$

[In] Integrate[CoshIntegral[a + b*x]^2/x^3,x]

[Out] Integrate[CoshIntegral[a + b*x]^2/x^3, x]

Maple [N/A] (verified)

Not integrable

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx + a)^2}{x^3} dx$$

[In] int(Chi(b*x+a)^2/x^3,x)

[Out] int(Chi(b*x+a)^2/x^3,x)

Fricas [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(a + bx)^2}{x^3} dx = \int \frac{\text{Chi}(bx + a)^2}{x^3} dx$$

[In] integrate(Chi(b*x+a)^2/x^3,x, algorithm="fricas")

[Out] integral(cosh_integral(b*x + a)^2/x^3, x)

Sympy [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(a + bx)^2}{x^3} dx = \int \frac{\text{Chi}^2(a + bx)}{x^3} dx$$

[In] integrate(Chi(b*x+a)**2/x**3,x)

[Out] Integral(Chi(a + b*x)**2/x**3, x)

Maxima [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Chi}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{Chi}(bx + a)^2}{x^3} dx$$

[In] integrate(Chi(b*x+a)^2/x^3,x, algorithm="maxima")

[Out] integrate(Chi(b*x + a)^2/x^3, x)

Giac [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Chi}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{Chi}(bx + a)^2}{x^3} dx$$

[In] integrate(Chi(b*x+a)^2/x^3,x, algorithm="giac")

[Out] integrate(Chi(b*x + a)^2/x^3, x)

Mupad [N/A]

Not integrable

Time = 4.87 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{Chi}(a + bx)^2}{x^3} dx = \int \frac{\operatorname{coshint}(a + bx)^2}{x^3} dx$$

[In] int(coshint(a + b*x)^2/x^3,x)

[Out] int(coshint(a + b*x)^2/x^3, x)

3.100 $\int x^2 \text{Chi}(d(a + b \log(cx^n))) dx$

Optimal result	493
Rubi [A] (verified)	493
Mathematica [A] (verified)	495
Maple [F]	495
Fricas [F]	496
Sympy [F]	496
Maxima [F]	496
Giac [F]	496
Mupad [F(-1)]	497

Optimal result

Integrand size = 17, antiderivative size = 128

$$\begin{aligned} & \int x^2 \text{Chi}(d(a + b \log(cx^n))) dx \\ &= \frac{1}{3} x^3 \text{Chi}(d(a + b \log(cx^n))) \\ & \quad - \frac{1}{6} e^{-\frac{3a}{bn}} x^3 (cx^n)^{-3/n} \text{ExpIntegralEi} \left(\frac{(3 - bdn)(a + b \log(cx^n))}{bn} \right) \\ & \quad - \frac{1}{6} e^{-\frac{3a}{bn}} x^3 (cx^n)^{-3/n} \text{ExpIntegralEi} \left(\frac{(3 + bdn)(a + b \log(cx^n))}{bn} \right) \end{aligned}$$

[Out] $\frac{1}{3} x^3 \text{Chi}(d(a + b \ln(cx^n))) - \frac{1}{6} x^3 \text{Ei} \left(\frac{(-b d n + 3)(a + b \ln(cx^n))}{b n} \right) / e^{3 a / b n} / ((c x^n)^{3 / n}) - \frac{1}{6} x^3 \text{Ei} \left(\frac{(b d n + 3)(a + b \ln(cx^n))}{b n} \right) / \exp(3 a / b n) / ((c x^n)^{3 / n})$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {6691, 12, 5651, 2347, 2209}

$$\begin{aligned} & \int x^2 \text{Chi}(d(a + b \log(cx^n))) dx \\ &= \frac{1}{3} x^3 \text{Chi}(d(a + b \log(cx^n))) \\ & \quad - \frac{1}{6} x^3 e^{-\frac{3a}{bn}} (cx^n)^{-3/n} \text{ExpIntegralEi} \left(\frac{(3 - bdn)(a + b \log(cx^n))}{bn} \right) \\ & \quad - \frac{1}{6} x^3 e^{-\frac{3a}{bn}} (cx^n)^{-3/n} \text{ExpIntegralEi} \left(\frac{(bdn + 3)(a + b \log(cx^n))}{bn} \right) \end{aligned}$$

[In] Int[x^2*CoshIntegral[d*(a + b*Log[c*x^n]),x]

[Out] (x^3*CoshIntegral[d*(a + b*Log[c*x^n]))/3 - (x^3*ExpIntegralEi[((3 - b*d*n)*(a + b*Log[c*x^n]))/(b*n)])/(6*E^((3*a)/(b*n))*(c*x^n)^(3/n)) - (x^3*ExpIntegralEi[((3 + b*d*n)*(a + b*Log[c*x^n]))/(b*n)])/(6*E^((3*a)/(b*n))*(c*x^n)^(3/n))

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 2209

Int[(F_)^((g_)*((e_) + (f_)*(x_)))/((c_) + (d_)*(x_)), x_Symbol] := Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[\$UseGamma]

Rule 2347

Int[((a_) + Log[(c_)*(x_)^(n_)])*(b_)^(p_)*((d_)*(x_)^(m_)), x_Symbol] := Dist[(d*x)^(m + 1)/(d*n*(c*x^n)^((m + 1)/n)), Subst[Int[E^(((m + 1)/n)*x)*(a + b*x)^p, x], x, Log[c*x^n]], x] /; FreeQ[{a, b, c, d, m, n, p}, x]

Rule 5651

Int[Cosh[((a_) + Log[(c_)*(x_)^(n_)])*(b_)]*(d_)*((e_) + Log[(g_)*(x_)^(m_)]*(f_))*(h_)^(q_)*((i_)*(x_)^(r_)), x_Symbol] := Dist[((i*x)^r*(1/((c*x^n)^(b*d)*(2*x^(r - b*d*n)))))/E^(a*d), Int[x^(r - b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] + Dist[E^(a*d)*(i*x)^r*((c*x^n)^(b*d)/(2*x^(r + b*d*n))), Int[x^(r + b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, m, n, q, r}, x]

Rule 6691

Int[CoshIntegral[((a_) + Log[(c_)*(x_)^(n_)])*(b_)]*(d_)*((e_)*(x_)^(m_)), x_Symbol] := Simp[(e*x)^(m + 1)*(CoshIntegral[d*(a + b*Log[c*x^n])]/(e*(m + 1))), x] - Dist[b*d*(n/(m + 1)), Int[(e*x)^m*(Cosh[d*(a + b*Log[c*x^n])]/(d*(a + b*Log[c*x^n]))], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{3}x^3\text{Chi}(d(a + b \log(cx^n))) - \frac{1}{3}(bdn) \int \frac{x^2 \cosh(d(a + b \log(cx^n)))}{d(a + b \log(cx^n))} dx \\ &= \frac{1}{3}x^3\text{Chi}(d(a + b \log(cx^n))) - \frac{1}{3}(bn) \int \frac{x^2 \cosh(d(a + b \log(cx^n)))}{a + b \log(cx^n)} dx \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{3}x^3\text{Chi}(d(a + b \log(cx^n))) - \frac{1}{6} \left(be^{-ad} n x^{bdn} (cx^n)^{-bd} \right) \int \frac{x^{2-bdn}}{a + b \log(cx^n)} dx \\
&\quad - \frac{1}{6} \left(be^{ad} n x^{-bdn} (cx^n)^{bd} \right) \int \frac{x^{2+bdn}}{a + b \log(cx^n)} dx \\
&= \frac{1}{3}x^3\text{Chi}(d(a + b \log(cx^n))) \\
&\quad - \frac{1}{6} \left(be^{-ad} x^3 (cx^n)^{-bd - \frac{3-bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(3-bdn)x}{n}}}{a + bx} dx, x, \log(cx^n) \right) \\
&\quad - \frac{1}{6} \left(be^{ad} x^3 (cx^n)^{bd - \frac{3+bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(3+bdn)x}{n}}}{a + bx} dx, x, \log(cx^n) \right) \\
&= \frac{1}{3}x^3\text{Chi}(d(a + b \log(cx^n))) \\
&\quad - \frac{1}{6} e^{-\frac{3a}{bn}} x^3 (cx^n)^{-3/n} \text{ExpIntegralEi} \left(\frac{(3 - bdn)(a + b \log(cx^n))}{bn} \right) \\
&\quad - \frac{1}{6} e^{-\frac{3a}{bn}} x^3 (cx^n)^{-3/n} \text{ExpIntegralEi} \left(\frac{(3 + bdn)(a + b \log(cx^n))}{bn} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.13 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.76

$$\int x^2 \text{Chi}(d(a + b \log(cx^n))) dx = \frac{1}{6} x^3 \left(2 \text{Chi}(d(a + b \log(cx^n))) \right. \\
\left. - e^{-\frac{3a}{bn}} (cx^n)^{-3/n} \left(\text{ExpIntegralEi} \left(-\frac{(-3 + bdn)(a + b \log(cx^n))}{bn} \right) + \text{ExpIntegralEi} \left(\frac{(3 + bdn)(a + b \log(cx^n))}{bn} \right) \right) \right)$$

[In] Integrate[x^2*CoshIntegral[d*(a + b*Log[c*x^n])],x]

[Out] (x^3*(2*CoshIntegral[d*(a + b*Log[c*x^n])] - (ExpIntegralEi[-(((-3 + b*d*n)*(a + b*Log[c*x^n]))/(b*n))] + ExpIntegralEi[((3 + b*d*n)*(a + b*Log[c*x^n]))/(b*n)]))/(E^((3*a)/(b*n))*(c*x^n)^(3/n)))/6

Maple [F]

$$\int x^2 \text{Chi}(d(a + b \ln(cx^n))) dx$$

[In] int(x^2*Chi(d*(a+b*ln(c*x^n))),x)

[Out] int(x^2*Chi(d*(a+b*ln(c*x^n))),x)

Fricas [F]

$$\int x^2 \operatorname{Chi}(d(a + b \log(cx^n))) dx = \int x^2 \operatorname{Chi}((b \log(cx^n) + a)d) dx$$

[In] integrate(x^2*Chi(d*(a+b*log(c*x^n))),x, algorithm="fricas")

[Out] integral(x^2*cosh_integral(b*d*log(c*x^n) + a*d), x)

Sympy [F]

$$\int x^2 \operatorname{Chi}(d(a + b \log(cx^n))) dx = \int x^2 \operatorname{Chi}(ad + bd \log(cx^n)) dx$$

[In] integrate(x**2*Chi(d*(a+b*ln(c*x**n))),x)

[Out] Integral(x**2*Chi(a*d + b*d*log(c*x**n)), x)

Maxima [F]

$$\int x^2 \operatorname{Chi}(d(a + b \log(cx^n))) dx = \int x^2 \operatorname{Chi}((b \log(cx^n) + a)d) dx$$

[In] integrate(x^2*Chi(d*(a+b*log(c*x^n))),x, algorithm="maxima")

[Out] integrate(x^2*Chi((b*log(c*x^n) + a)*d), x)

Giac [F]

$$\int x^2 \operatorname{Chi}(d(a + b \log(cx^n))) dx = \int x^2 \operatorname{Chi}((b \log(cx^n) + a)d) dx$$

[In] integrate(x^2*Chi(d*(a+b*log(c*x^n))),x, algorithm="giac")

[Out] integrate(x^2*Chi((b*log(c*x^n) + a)*d), x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{Chi}(d(a + b \log(cx^n))) dx = \int x^2 \operatorname{coshint}(d(a + b \ln(cx^n))) dx$$

```
[In] int(x^2*coshint(d*(a + b*log(c*x^n))),x)
```

```
[Out] int(x^2*coshint(d*(a + b*log(c*x^n))), x)
```

3.101 $\int x \text{Chi}(d(a + b \log(cx^n))) dx$

Optimal result	498
Rubi [A] (verified)	498
Mathematica [A] (verified)	500
Maple [F]	500
Fricas [F]	501
Sympy [F]	501
Maxima [F]	501
Giac [F]	501
Mupad [F(-1)]	502

Optimal result

Integrand size = 15, antiderivative size = 128

$$\begin{aligned} & \int x \text{Chi}(d(a + b \log(cx^n))) dx \\ &= \frac{1}{2} x^2 \text{Chi}(d(a + b \log(cx^n))) \\ & \quad - \frac{1}{4} e^{-\frac{2a}{bn}} x^2 (cx^n)^{-2/n} \text{ExpIntegralEi}\left(\frac{(2 - bdn)(a + b \log(cx^n))}{bn}\right) \\ & \quad - \frac{1}{4} e^{-\frac{2a}{bn}} x^2 (cx^n)^{-2/n} \text{ExpIntegralEi}\left(\frac{(2 + bdn)(a + b \log(cx^n))}{bn}\right) \end{aligned}$$

[Out] $\frac{1}{2} x^2 \text{Chi}(d(a + b \ln(cx^n))) - \frac{1}{4} x^2 \text{Ei}\left(\frac{(-b*d*n+2)*(a+b*\ln(cx^n))}{b/n}\right) / e^{2*a/b/n} / ((cx^n)^{(2/n)}) - \frac{1}{4} x^2 \text{Ei}\left(\frac{(b*d*n+2)*(a+b*\ln(cx^n))}{b/n}\right) / \exp(2*a/b/n) / ((cx^n)^{(2/n)})$

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {6691, 12, 5651, 2347, 2209}

$$\begin{aligned} & \int x \text{Chi}(d(a + b \log(cx^n))) dx \\ &= \frac{1}{2} x^2 \text{Chi}(d(a + b \log(cx^n))) \\ & \quad - \frac{1}{4} x^2 e^{-\frac{2a}{bn}} (cx^n)^{-2/n} \text{ExpIntegralEi}\left(\frac{(2 - bdn)(a + b \log(cx^n))}{bn}\right) \\ & \quad - \frac{1}{4} x^2 e^{-\frac{2a}{bn}} (cx^n)^{-2/n} \text{ExpIntegralEi}\left(\frac{(bdn + 2)(a + b \log(cx^n))}{bn}\right) \end{aligned}$$

[In] Int[x*CoshIntegral[d*(a + b*Log[c*x^n]), x]

[Out] (x^2*CoshIntegral[d*(a + b*Log[c*x^n]))/2 - (x^2*ExpIntegralEi[((2 - b*d*n)*(a + b*Log[c*x^n]))/(b*n)])/(4*E^((2*a)/(b*n))*(c*x^n)^(2/n)) - (x^2*ExpIntegralEi[((2 + b*d*n)*(a + b*Log[c*x^n]))/(b*n)])/(4*E^((2*a)/(b*n))*(c*x^n)^(2/n))

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 2209

Int[(F_)^((g_)*((e_) + (f_)*(x_)))/((c_) + (d_)*(x_)), x_Symbol] := Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[\$UseGamma]

Rule 2347

Int[((a_) + Log[(c_)*(x_)^(n_)])*(b_)^(p_)*((d_)*(x_))^(m_), x_Symbol] := Dist[(d*x)^(m + 1)/(d*n*(c*x^n)^((m + 1)/n)), Subst[Int[E^((m + 1)/n)*x*(a + b*x)^p, x], x, Log[c*x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x]

Rule 5651

Int[Cosh[((a_) + Log[(c_)*(x_)^(n_)])*(b_)*(d_)]*(((e_) + Log[(g_)*(x_)^(m_)])*(f_)*(h_))^(q_)*((i_)*(x_))^(r_), x_Symbol] := Dist[((i*x)^r*(1/((c*x^n)^(b*d)*(2*x^(r - b*d*n)))))/E^(a*d), Int[x^(r - b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] + Dist[E^(a*d)*(i*x)^r*((c*x^n)^(b*d)/(2*x^(r + b*d*n))), Int[x^(r + b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, m, n, q, r}, x]

Rule 6691

Int[CoshIntegral[((a_) + Log[(c_)*(x_)^(n_)])*(b_)*(d_)]*((e_)*(x_))^(m_), x_Symbol] := Simp[(e*x)^(m + 1)*(CoshIntegral[d*(a + b*Log[c*x^n])]/(e*(m + 1))), x] - Dist[b*d*(n/(m + 1)), Int[(e*x)^m*(Cosh[d*(a + b*Log[c*x^n])]/(d*(a + b*Log[c*x^n]))], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{2}x^2\text{Chi}(d(a + b\log(cx^n))) - \frac{1}{2}(bdn) \int \frac{x \cosh(d(a + b\log(cx^n)))}{d(a + b\log(cx^n))} dx \\ &= \frac{1}{2}x^2\text{Chi}(d(a + b\log(cx^n))) - \frac{1}{2}(bn) \int \frac{x \cosh(d(a + b\log(cx^n)))}{a + b\log(cx^n)} dx \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2}x^2\text{Chi}(d(a + b \log(cx^n))) - \frac{1}{4}\left(be^{-ad}n x^{bdn}(cx^n)^{-bd} \right) \int \frac{x^{1-bdn}}{a + b \log(cx^n)} dx \\
&\quad - \frac{1}{4}\left(be^{ad}n x^{-bdn}(cx^n)^{bd} \right) \int \frac{x^{1+bdn}}{a + b \log(cx^n)} dx \\
&= \frac{1}{2}x^2\text{Chi}(d(a + b \log(cx^n))) \\
&\quad - \frac{1}{4}\left(be^{-ad}x^2(cx^n)^{-bd-\frac{2-bdn}{n}} \right) \text{Subst}\left(\int \frac{e^{\frac{(2-bdn)x}{n}}}{a + bx} dx, x, \log(cx^n) \right) \\
&\quad - \frac{1}{4}\left(be^{ad}x^2(cx^n)^{bd-\frac{2+bdn}{n}} \right) \text{Subst}\left(\int \frac{e^{\frac{(2+bdn)x}{n}}}{a + bx} dx, x, \log(cx^n) \right) \\
&= \frac{1}{2}x^2\text{Chi}(d(a + b \log(cx^n))) \\
&\quad - \frac{1}{4}e^{-\frac{2a}{bn}}x^2(cx^n)^{-2/n} \text{ExpIntegralEi}\left(\frac{(2 - bdn)(a + b \log(cx^n))}{bn} \right) \\
&\quad - \frac{1}{4}e^{-\frac{2a}{bn}}x^2(cx^n)^{-2/n} \text{ExpIntegralEi}\left(\frac{(2 + bdn)(a + b \log(cx^n))}{bn} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.09 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.76

$$\int x\text{Chi}(d(a + b \log(cx^n))) dx = \frac{1}{4}x^2\left(2\text{Chi}(d(a + b \log(cx^n))) \right. \\
\left. - e^{-\frac{2a}{bn}}(cx^n)^{-2/n} \left(\text{ExpIntegralEi}\left(-\frac{(-2 + bdn)(a + b \log(cx^n))}{bn} \right) + \text{ExpIntegralEi}\left(\frac{(2 + bdn)(a + b \log(cx^n))}{bn} \right) \right) \right)$$

[In] Integrate[x*CoshIntegral[d*(a + b*Log[c*x^n])],x]

[Out] (x^2*(2*CoshIntegral[d*(a + b*Log[c*x^n])] - (ExpIntegralEi[-((-2 + b*d*n)*(a + b*Log[c*x^n])]/(b*n))] + ExpIntegralEi[((2 + b*d*n)*(a + b*Log[c*x^n])/]/(b*n)))/(E^((2*a)/(b*n))*(c*x^n)^(2/n)))/4

Maple [F]

$$\int x \text{Chi}(d(a + b \ln(cx^n))) dx$$

[In] int(x*Chi(d*(a+b*ln(c*x^n))),x)

[Out] int(x*Chi(d*(a+b*ln(c*x^n))),x)

Fricas [F]

$$\int x \operatorname{Chi}(d(a + b \log(cx^n))) dx = \int x \operatorname{Chi}((b \log(cx^n) + a)d) dx$$

[In] integrate(x*Chi(d*(a+b*log(c*x^n))),x, algorithm="fricas")

[Out] integral(x*cosh_integral(b*d*log(c*x^n) + a*d), x)

Sympy [F]

$$\int x \operatorname{Chi}(d(a + b \log(cx^n))) dx = \int x \operatorname{Chi}(ad + bd \log(cx^n)) dx$$

[In] integrate(x*Chi(d*(a+b*ln(c*x**n))),x)

[Out] Integral(x*Chi(a*d + b*d*log(c*x**n)), x)

Maxima [F]

$$\int x \operatorname{Chi}(d(a + b \log(cx^n))) dx = \int x \operatorname{Chi}((b \log(cx^n) + a)d) dx$$

[In] integrate(x*Chi(d*(a+b*log(c*x^n))),x, algorithm="maxima")

[Out] integrate(x*Chi((b*log(c*x^n) + a)*d), x)

Giac [F]

$$\int x \operatorname{Chi}(d(a + b \log(cx^n))) dx = \int x \operatorname{Chi}((b \log(cx^n) + a)d) dx$$

[In] integrate(x*Chi(d*(a+b*log(c*x^n))),x, algorithm="giac")

[Out] integrate(x*Chi((b*log(c*x^n) + a)*d), x)

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{Chi}(d(a + b \log(cx^n))) dx = \int x \operatorname{coshint}(d(a + b \ln(cx^n))) dx$$

```
[In] int(x*coshint(d*(a + b*log(c*x^n))),x)
```

```
[Out] int(x*coshint(d*(a + b*log(c*x^n))), x)
```

3.102 $\int \text{Chi}(d(a + b \log(cx^n))) dx$

Optimal result	503
Rubi [A] (verified)	503
Mathematica [A] (verified)	505
Maple [F]	505
Fricas [F]	506
Sympy [F]	506
Maxima [F]	506
Giac [F]	506
Mupad [F(-1)]	507

Optimal result

Integrand size = 13, antiderivative size = 119

$$\int \text{Chi}(d(a + b \log(cx^n))) dx$$

$$= x \text{Chi}(d(a + b \log(cx^n))) - \frac{1}{2} e^{-\frac{a}{bn}} x (cx^n)^{-1/n} \text{ExpIntegralEi}\left(\frac{(1 - bdn)(a + b \log(cx^n))}{bn}\right)$$

$$- \frac{1}{2} e^{-\frac{a}{bn}} x (cx^n)^{-1/n} \text{ExpIntegralEi}\left(\frac{(1 + bdn)(a + b \log(cx^n))}{bn}\right)$$

[Out] $x \text{Chi}(d(a + b \ln(cx^n))) - 1/2 * x * \text{Ei}((-b*d*n+1)*(a+b*\ln(cx^n))/b/n) / \exp(a/b/n) / ((cx^n)^{(1/n)}) - 1/2 * x * \text{Ei}((b*d*n+1)*(a+b*\ln(cx^n))/b/n) / \exp(a/b/n) / ((cx^n)^{(1/n)})$

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {6688, 12, 5649, 2347, 2209}

$$\int \text{Chi}(d(a + b \log(cx^n))) dx$$

$$= x \text{Chi}(d(a + b \log(cx^n))) - \frac{1}{2} x e^{-\frac{a}{bn}} (cx^n)^{-1/n} \text{ExpIntegralEi}\left(\frac{(1 - bdn)(a + b \log(cx^n))}{bn}\right)$$

$$- \frac{1}{2} x e^{-\frac{a}{bn}} (cx^n)^{-1/n} \text{ExpIntegralEi}\left(\frac{(bdn + 1)(a + b \log(cx^n))}{bn}\right)$$

[In] $\text{Int}[\text{CoshIntegral}[d(a + b*\text{Log}[c*x^n])], x]$

```
[Out] x*CoshIntegral[d*(a + b*Log[c*x^n])] - (x*ExpIntegralEi[((1 - b*d*n)*(a + b
*Log[c*x^n]))/(b*n)])/(2*E^(a/(b*n))*(c*x^n)^n^(-1)) - (x*ExpIntegralEi[((1
+ b*d*n)*(a + b*Log[c*x^n]))/(b*n)])/(2*E^(a/(b*n))*(c*x^n)^n^(-1))
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 2209

```
Int[(F_)^((g_.)*((e_.) + (f_.)*(x_)))/((c_.) + (d_.)*(x_)), x_Symbol] := Si
mp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; F
reeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]
```

Rule 2347

```
Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^((p_.)*((d_.)*(x_)^(m_.)), x_Symbol
] := Dist[(d*x)^(m + 1)/(d*n*(c*x^n)^((m + 1)/n)), Subst[Int[E^(((m + 1)/n)
*x)*(a + b*x)^p, x], x, Log[c*x^n]], x] /; FreeQ[{a, b, c, d, m, n, p}, x]
```

Rule 5649

```
Int[Cosh[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.))*((e_.) + Log[(g_.)*(
x_)^(m_.)]*(f_.))*(h_.)^((q_.)), x_Symbol] := Dist[1/((c*x^n)^(b*d)*(2/x^(b*
d*n)))/E^(a*d), Int[(h*(e + f*Log[g*x^m]))^q/x^(b*d*n), x], x] + Dist[E^(a*
d)*((c*x^n)^(b*d)/(2*x^(b*d*n))), Int[x^(b*d*n)*(h*(e + f*Log[g*x^m]))^q, x
], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n, q}, x]
```

Rule 6688

```
Int[CoshIntegral[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)], x_Symbol] :=
Simp[x*CoshIntegral[d*(a + b*Log[c*x^n])], x] - Dist[b*d*n, Int[Cosh[d*(a
+ b*Log[c*x^n])]/(d*(a + b*Log[c*x^n])), x], x] /; FreeQ[{a, b, c, d, n}, x
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= x\text{Chi}(d(a + b \log(cx^n))) - (bdn) \int \frac{\cosh(d(a + b \log(cx^n)))}{d(a + b \log(cx^n))} dx \\
&= x\text{Chi}(d(a + b \log(cx^n))) - (bn) \int \frac{\cosh(d(a + b \log(cx^n)))}{a + b \log(cx^n)} dx \\
&= x\text{Chi}(d(a + b \log(cx^n))) - \frac{1}{2} \left(be^{-ad} n x^{bdn} (cx^n)^{-bd} \right) \int \frac{x^{-bdn}}{a + b \log(cx^n)} dx \\
&\quad - \frac{1}{2} \left(be^{ad} n x^{-bdn} (cx^n)^{bd} \right) \int \frac{x^{bdn}}{a + b \log(cx^n)} dx
\end{aligned}$$

$$\begin{aligned}
&= x\text{Chi}(d(a + b \log(cx^n))) \\
&\quad - \frac{1}{2} \left(b e^{-ad} x (cx^n)^{-bd - \frac{1-bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(1-bdn)x}{n}}}{a + bx} dx, x, \log(cx^n) \right) \\
&\quad - \frac{1}{2} \left(b e^{ad} x (cx^n)^{bd - \frac{1+bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(1+bdn)x}{n}}}{a + bx} dx, x, \log(cx^n) \right) \\
&= x\text{Chi}(d(a + b \log(cx^n))) \\
&\quad - \frac{1}{2} e^{-\frac{a}{bn}} x (cx^n)^{-1/n} \text{ExpIntegralEi} \left(\frac{(1 - bdn)(a + b \log(cx^n))}{bn} \right) \\
&\quad - \frac{1}{2} e^{-\frac{a}{bn}} x (cx^n)^{-1/n} \text{ExpIntegralEi} \left(\frac{(1 + bdn)(a + b \log(cx^n))}{bn} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.05 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.78

$$\begin{aligned}
&\int \text{Chi}(d(a + b \log(cx^n))) dx \\
&= x\text{Chi}(d(a + b \log(cx^n))) \\
&\quad - \frac{1}{2} e^{-\frac{a}{bn}} x (cx^n)^{-1/n} \left(\text{ExpIntegralEi} \left(-\frac{(-1 + bdn)(a + b \log(cx^n))}{bn} \right) \right. \\
&\quad \quad \quad \left. + \text{ExpIntegralEi} \left(\frac{(1 + bdn)(a + b \log(cx^n))}{bn} \right) \right)
\end{aligned}$$

[In] Integrate[CoshIntegral[d*(a + b*Log[c*x^n])],x]

[Out] x*CoshIntegral[d*(a + b*Log[c*x^n])] - (x*(ExpIntegralEi[-(((-1 + b*d*n)*(a + b*Log[c*x^n]))/(b*n))]) + ExpIntegralEi[(((1 + b*d*n)*(a + b*Log[c*x^n]))/(b*n))]))/(2*E^(a/(b*n))*(c*x^n)^n^(-1))

Maple [F]

$$\int \text{Chi}(d(a + b \ln(cx^n))) dx$$

[In] int(Chi(d*(a+b*ln(c*x^n))),x)

[Out] int(Chi(d*(a+b*ln(c*x^n))),x)

Fricas [F]

$$\int \operatorname{Chi}(d(a + b \log(cx^n))) dx = \int \operatorname{Chi}((b \log(cx^n) + a)d) dx$$

```
[In] integrate(Chi(d*(a+b*log(c*x^n))),x, algorithm="fricas")
```

```
[Out] integral(cosh_integral(b*d*log(c*x^n) + a*d), x)
```

Sympy [F]

$$\int \operatorname{Chi}(d(a + b \log(cx^n))) dx = \int \operatorname{Chi}(d(a + b \log(cx^n))) dx$$

```
[In] integrate(Chi(d*(a+b*ln(c*x**n))),x)
```

```
[Out] Integral(Chi(d*(a + b*log(c*x**n))), x)
```

Maxima [F]

$$\int \operatorname{Chi}(d(a + b \log(cx^n))) dx = \int \operatorname{Chi}((b \log(cx^n) + a)d) dx$$

```
[In] integrate(Chi(d*(a+b*log(c*x^n))),x, algorithm="maxima")
```

```
[Out] integrate(Chi((b*log(c*x^n) + a)*d), x)
```

Giac [F]

$$\int \operatorname{Chi}(d(a + b \log(cx^n))) dx = \int \operatorname{Chi}((b \log(cx^n) + a)d) dx$$

```
[In] integrate(Chi(d*(a+b*log(c*x^n))),x, algorithm="giac")
```

```
[Out] integrate(Chi((b*log(c*x^n) + a)*d), x)
```

Mupad [F(-1)]

Timed out.

$$\int \text{Chi}(d(a + b \log(cx^n))) dx = \int \text{coshint}(d(a + b \ln(cx^n))) dx$$

```
[In] int(coshint(d*(a + b*log(c*x^n))),x)
```

```
[Out] int(coshint(d*(a + b*log(c*x^n))), x)
```

3.103 $\int \frac{\text{Chi}(d(a+b \log(cx^n)))}{x} dx$

Optimal result	508
Rubi [A] (verified)	508
Mathematica [A] (verified)	509
Maple [A] (verified)	509
Fricas [F]	510
Sympy [F]	510
Maxima [F]	510
Giac [F]	510
Mupad [F(-1)]	511

Optimal result

Integrand size = 17, antiderivative size = 55

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x} dx = \frac{\text{Chi}(d(a + b \log(cx^n))) (a + b \log(cx^n))}{bn} - \frac{\sinh(d(a + b \log(cx^n)))}{bdn}$$

[Out] Chi(d*(a+b*ln(c*x^n))*(a+b*ln(c*x^n))/b/n-sinh(d*(a+b*ln(c*x^n)))/b/d/n

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {6664}

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x} dx = \frac{(a + b \log(cx^n)) \text{Chi}(d(a + b \log(cx^n)))}{bn} - \frac{\sinh(d(a + b \log(cx^n)))}{bdn}$$

[In] Int[CoshIntegral[d*(a + b*Log[c*x^n])]/x,x]

[Out] (CoshIntegral[d*(a + b*Log[c*x^n])*(a + b*Log[c*x^n])]/(b*n) - Sinh[d*(a + b*Log[c*x^n])]/(b*d*n)

Rule 6664

Int[CoshIntegral[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[(a + b*x)*(CoshIntegral[a + b*x]/b), x] - Simp[Sinh[a + b*x]/b, x] /; FreeQ[{a, b}, x]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\text{Subst}\left(\int \text{Chi}(d(a+bx)) dx, x, \log(cx^n)\right)}{n} \\ &= \frac{\text{Subst}\left(\int \text{Chi}(x) dx, x, ad+bd \log(cx^n)\right)}{bdn} \\ &= \frac{\text{Chi}(ad+bd \log(cx^n))(a+b \log(cx^n))}{bn} - \frac{\sinh(ad+bd \log(cx^n))}{bdn} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.75

$$\int \frac{\text{Chi}(d(a+b \log(cx^n)))}{x} dx = \frac{a \text{Chi}(ad+bd \log(cx^n))}{bn} + \frac{\text{Chi}(d(a+b \log(cx^n))) \log(cx^n)}{n} - \frac{\cosh(bd \log(cx^n)) \sinh(ad)}{bdn} - \frac{\cosh(ad) \sinh(bd \log(cx^n))}{bdn}$$

[In] Integrate[CoshIntegral[d*(a + b*Log[c*x^n])]/x,x]

[Out] (a*CoshIntegral[a*d + b*d*Log[c*x^n])/(b*n) + (CoshIntegral[d*(a + b*Log[c*x^n])]*Log[c*x^n])/n - (Cosh[b*d*Log[c*x^n])*Sinh[a*d]/(b*d*n) - (Cosh[a*d]*Sinh[b*d*Log[c*x^n])/(b*d*n)

Maple [A] (verified)

Time = 1.51 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.02

method	result
derivativedivides	$\frac{\text{Chi}(ad+bd \ln(cx^n))(ad+bd \ln(cx^n))-\sinh(ad+bd \ln(cx^n))}{ndb}$
default	$\frac{\text{Chi}(ad+bd \ln(cx^n))(ad+bd \ln(cx^n))-\sinh(ad+bd \ln(cx^n))}{ndb}$
parts	$\ln(x) \text{Chi}(d(a+b \ln(cx^n))) - nb \left(-\frac{(\ln(cx^n)-n \ln(x)) \text{Chi}(\ln(x)bdn+d(b(\ln(cx^n)-n \ln(x))+a))}{bn^2} - \frac{a}{bn} \right)$

[In] int(Chi(d*(a+b*ln(c*x^n)))/x,x,method=_RETURNVERBOSE)

[Out] 1/n/d/b*(Chi(a*d+b*d*ln(c*x^n))*(a*d+b*d*ln(c*x^n))-sinh(a*d+b*d*ln(c*x^n)))

Fricas [F]

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x} dx = \int \frac{\text{Chi}((b \log(cx^n) + a)d)}{x} dx$$

[In] integrate(Chi(d*(a+b*log(c*x^n)))/x,x, algorithm="fricas")

[Out] integral(cosh_integral(b*d*log(c*x^n) + a*d)/x, x)

Sympy [F]

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x} dx = \int \frac{\text{Chi}(ad + bd \log(cx^n))}{x} dx$$

[In] integrate(Chi(d*(a+b*ln(c*x**n)))/x,x)

[Out] Integral(Chi(a*d + b*d*log(c*x**n))/x, x)

Maxima [F]

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x} dx = \int \frac{\text{Chi}((b \log(cx^n) + a)d)}{x} dx$$

[In] integrate(Chi(d*(a+b*log(c*x^n)))/x,x, algorithm="maxima")

[Out] integrate(Chi((b*log(c*x^n) + a)*d)/x, x)

Giac [F]

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x} dx = \int \frac{\text{Chi}((b \log(cx^n) + a)d)}{x} dx$$

[In] integrate(Chi(d*(a+b*log(c*x^n)))/x,x, algorithm="giac")

[Out] integrate(Chi((b*log(c*x^n) + a)*d)/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x} dx = \frac{\ln(cx^n) \text{coshint}(d(a + b \ln(cx^n)))}{n} + \frac{a \text{coshint}(d(a + b \ln(cx^n)))}{bn} - \frac{e^{ad}(cx^n)^{bd}}{2bdn} + \frac{e^{-ad}}{2bdn(cx^n)^{bd}}$$

```
[In] int(coshint(d*(a + b*log(c*x^n)))/x,x)
```

```
[Out] (log(c*x^n)*coshint(d*(a + b*log(c*x^n))))/n + (a*coshint(d*(a + b*log(c*x^n))))/(b*n) - (exp(a*d)*(c*x^n)^(b*d))/(2*b*d*n) + exp(-a*d)/(2*b*d*n*(c*x^n)^(b*d))
```

3.104 $\int \frac{\mathbf{Chi}(d(a+b \log(cx^n)))}{x^2} dx$

Optimal result	512
Rubi [A] (verified)	512
Mathematica [A] (verified)	514
Maple [F]	515
Fricas [F]	515
Sympy [F]	515
Maxima [F]	515
Giac [F]	516
Mupad [F(-1)]	516

Optimal result

Integrand size = 17, antiderivative size = 122

$$\int \frac{\mathbf{Chi}(d(a + b \log(cx^n)))}{x^2} dx = -\frac{\mathbf{Chi}(d(a + b \log(cx^n)))}{x} + \frac{e^{\frac{a}{bn}}(cx^n)^{\frac{1}{n}} \text{ExpIntegralEi}\left(-\frac{(1-bdn)(a+b \log(cx^n))}{bn}\right)}{2x} + \frac{e^{\frac{a}{bn}}(cx^n)^{\frac{1}{n}} \text{ExpIntegralEi}\left(-\frac{(1+bdn)(a+b \log(cx^n))}{bn}\right)}{2x}$$

[Out] $-\mathbf{Chi}(d*(a+b*\ln(c*x^n)))/x+1/2*\exp(a/b/n)*(c*x^n)^{(1/n)}*Ei(-(-b*d*n+1)*(a+b*\ln(c*x^n))/b/n)/x+1/2*\exp(a/b/n)*(c*x^n)^{(1/n)}*Ei(-(b*d*n+1)*(a+b*\ln(c*x^n))/b/n)/x$

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {6691, 12, 5651, 2347, 2209}

$$\int \frac{\mathbf{Chi}(d(a + b \log(cx^n)))}{x^2} dx = -\frac{\mathbf{Chi}(d(a + b \log(cx^n)))}{x} + \frac{e^{\frac{a}{bn}}(cx^n)^{\frac{1}{n}} \text{ExpIntegralEi}\left(-\frac{(1-bdn)(a+b \log(cx^n))}{bn}\right)}{2x} + \frac{e^{\frac{a}{bn}}(cx^n)^{\frac{1}{n}} \text{ExpIntegralEi}\left(-\frac{(bdn+1)(a+b \log(cx^n))}{bn}\right)}{2x}$$

[In] $\text{Int}[\text{CoshIntegral}[d*(a + b*\text{Log}[c*x^n])]/x^2, x]$


```
[Out] -(CoshIntegral[d*(a + b*Log[c*x^n])/x] + (E^(a/(b*n))*(c*x^n)^n^(-1)*ExpIntegralEi[-(((1 - b*d*n)*(a + b*Log[c*x^n]))/(b*n))])/(2*x) + (E^(a/(b*n))*(c*x^n)^n^(-1)*ExpIntegralEi[-(((1 + b*d*n)*(a + b*Log[c*x^n]))/(b*n))])/(2*x)
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] :=> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 2209

```
Int[(F_)^((g_)*((e_) + (f_)*(x_)))/((c_) + (d_)*(x_)), x_Symbol] :=> Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]
```

Rule 2347

```
Int[((a_) + Log[(c_)*(x_)^(n_)])*(b_)^(p_)*((d_)*(x_))^(m_), x_Symbol] :=> Dist[(d*x)^(m + 1)/(d*n*(c*x^n)^(m + 1/n)), Subst[Int[E^((m + 1)/n)*x]*(a + b*x)^p, x], x, Log[c*x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x]
```

Rule 5651

```
Int[Cosh[((a_) + Log[(c_)*(x_)^(n_)])*(b_)*(d_)]*(((e_) + Log[(g_)*(x_)^(m_)])*(f_)*(h_)^(q_)*((i_)*(x_))^(r_)), x_Symbol] :=> Dist[((i*x)^r*(1/((c*x^n)^(b*d)*(2*x^(r - b*d*n)))))/E^(a*d), Int[x^(r - b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] + Dist[E^(a*d)*(i*x)^r*((c*x^n)^(b*d)/(2*x^(r + b*d*n))), Int[x^(r + b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, m, n, q, r}, x]
```

Rule 6691

```
Int[CoshIntegral[((a_) + Log[(c_)*(x_)^(n_)])*(b_)*(d_)]*((e_)*(x_))^(m_), x_Symbol] :=> Simp[(e*x)^(m + 1)*(CoshIntegral[d*(a + b*Log[c*x^n])]/(e*(m + 1))), x] - Dist[b*d*(n/(m + 1)), Int[(e*x)^m*(Cosh[d*(a + b*Log[c*x^n])]/(d*(a + b*Log[c*x^n]))], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Chi}(d(a + b \log(cx^n)))}{x} + (bdn) \int \frac{\cosh(d(a + b \log(cx^n)))}{dx^2 (a + b \log(cx^n))} dx \\ &= -\frac{\text{Chi}(d(a + b \log(cx^n)))}{x} + (bn) \int \frac{\cosh(d(a + b \log(cx^n)))}{x^2 (a + b \log(cx^n))} dx \end{aligned}$$

$$\begin{aligned}
&= -\frac{\text{Chi}(d(a + b \log(cx^n)))}{x} + \frac{1}{2} \left(be^{-ad} n x^{bdn} (cx^n)^{-bd} \right) \int \frac{x^{-2-bdn}}{a + b \log(cx^n)} dx \\
&\quad + \frac{1}{2} \left(be^{ad} n x^{-bdn} (cx^n)^{bd} \right) \int \frac{x^{-2+bdn}}{a + b \log(cx^n)} dx \\
&= -\frac{\text{Chi}(d(a + b \log(cx^n)))}{x} \\
&\quad + \frac{\left(be^{-ad} (cx^n)^{-bd - \frac{-1-bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(-1-bdn)x}{a+bx}}}{a+bx} dx, x, \log(cx^n) \right)}{2x} \\
&\quad + \frac{\left(be^{ad} (cx^n)^{bd - \frac{-1+bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(-1+bdn)x}{a+bx}}}{a+bx} dx, x, \log(cx^n) \right)}{2x} \\
&= -\frac{\text{Chi}(d(a + b \log(cx^n)))}{x} + \frac{e^{\frac{a}{bn}} (cx^n)^{\frac{1}{n}} \text{ExpIntegralEi} \left(-\frac{(1-bdn)(a+b \log(cx^n))}{bn} \right)}{2x} \\
&\quad + \frac{e^{\frac{a}{bn}} (cx^n)^{\frac{1}{n}} \text{ExpIntegralEi} \left(-\frac{(1+bdn)(a+b \log(cx^n))}{bn} \right)}{2x}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.26 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.18

$$\begin{aligned}
&\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x^2} dx \\
&= -\frac{\text{Chi}(d(a + b \log(cx^n)))}{x} \\
&\quad + \frac{1}{2} e^{-\frac{(-1+bdn)(a+b(-n \log(x)+\log(cx^n)))}{bn}} \left(\text{ExpIntegralEi} \left(\frac{(-1+bdn)(a+b \log(cx^n))}{bn} \right) \right) \\
&\quad + \text{ExpIntegralEi} \left(-\frac{(1+bdn)(a+b \log(cx^n))}{bn} \right) \left(\cosh(d(a+b(-n \log(x)+\log(cx^n)))) \right. \\
&\quad \quad \left. + \sinh(d(a+b(-n \log(x)+\log(cx^n)))) \right)
\end{aligned}$$

[In] Integrate[CoshIntegral[d*(a + b*Log[c*x^n])]/x^2,x]

[Out] -(CoshIntegral[d*(a + b*Log[c*x^n])]/x) + ((ExpIntegralEi[((-1 + b*d*n)*(a + b*Log[c*x^n])/(b*n)] + ExpIntegralEi[-(((1 + b*d*n)*(a + b*Log[c*x^n])/(b*n))])*(Cosh[d*(a + b*(-n*Log[x]) + Log[c*x^n])]) + Sinh[d*(a + b*(-n*Log[x]) + Log[c*x^n])])]/(2*E^(((1 + b*d*n)*(a + b*(-n*Log[x]) + Log[c*x^n])/(b*n))))))

Maple [F]

$$\int \frac{\text{Chi}(d(a + b \ln(cx^n)))}{x^2} dx$$

[In] int(Chi(d*(a+b*ln(c*x^n)))/x^2,x)

[Out] int(Chi(d*(a+b*ln(c*x^n)))/x^2,x)

Fricas [F]

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\text{Chi}((b \log(cx^n) + a)d)}{x^2} dx$$

[In] integrate(Chi(d*(a+b*log(c*x^n)))/x^2,x, algorithm="fricas")

[Out] integral(cosh_integral(b*d*log(c*x^n) + a*d)/x^2, x)

Sympy [F]

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\text{Chi}(ad + bd \log(cx^n))}{x^2} dx$$

[In] integrate(Chi(d*(a+b*ln(c*x**n)))/x**2,x)

[Out] Integral(Chi(a*d + b*d*log(c*x**n))/x**2, x)

Maxima [F]

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\text{Chi}((b \log(cx^n) + a)d)}{x^2} dx$$

[In] integrate(Chi(d*(a+b*log(c*x^n)))/x^2,x, algorithm="maxima")

[Out] integrate(Chi((b*log(c*x^n) + a)*d)/x^2, x)

Giac [F]

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\text{Chi}((b \log(cx^n) + a)d)}{x^2} dx$$

[In] integrate(Chi(d*(a+b*log(c*x^n)))/x^2,x, algorithm="giac")

[Out] integrate(Chi((b*log(c*x^n) + a)*d)/x^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\text{coshint}(d(a + b \ln(cx^n)))}{x^2} dx$$

[In] int(coshint(d*(a + b*log(c*x^n)))/x^2,x)

[Out] int(coshint(d*(a + b*log(c*x^n)))/x^2, x)

3.105 $\int \frac{\text{Chi}(d(a+b \log(cx^n)))}{x^3} dx$

Optimal result	517
Rubi [A] (verified)	517
Mathematica [A] (verified)	519
Maple [F]	520
Fricas [F]	520
Sympy [F]	520
Maxima [F]	520
Giac [F]	521
Mupad [F(-1)]	521

Optimal result

Integrand size = 17, antiderivative size = 130

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x^3} dx = -\frac{\text{Chi}(d(a + b \log(cx^n)))}{2x^2} + \frac{e^{\frac{2a}{bn}}(cx^n)^{2/n} \text{ExpIntegralEi}\left(-\frac{(2-bdn)(a+b \log(cx^n))}{bn}\right)}{4x^2} + \frac{e^{\frac{2a}{bn}}(cx^n)^{2/n} \text{ExpIntegralEi}\left(-\frac{(2+bdn)(a+b \log(cx^n))}{bn}\right)}{4x^2}$$

[Out] $-1/2*\text{Chi}(d*(a+b*\ln(c*x^n)))/x^2+1/4*\exp(2*a/b/n)*(c*x^n)^{(2/n)}*Ei(-(-b*d*n+2)*(a+b*\ln(c*x^n))/b/n)/x^2+1/4*\exp(2*a/b/n)*(c*x^n)^{(2/n)}*Ei(-(b*d*n+2)*(a+b*\ln(c*x^n))/b/n)/x^2$

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {6691, 12, 5651, 2347, 2209}

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x^3} dx = -\frac{\text{Chi}(d(a + b \log(cx^n)))}{2x^2} + \frac{e^{\frac{2a}{bn}}(cx^n)^{2/n} \text{ExpIntegralEi}\left(-\frac{(2-bdn)(a+b \log(cx^n))}{bn}\right)}{4x^2} + \frac{e^{\frac{2a}{bn}}(cx^n)^{2/n} \text{ExpIntegralEi}\left(-\frac{(bdn+2)(a+b \log(cx^n))}{bn}\right)}{4x^2}$$

[In] $\text{Int}[\text{CoshIntegral}[d*(a + b*\text{Log}[c*x^n])]/x^3, x]$

```
[Out] -1/2*CoshIntegral[d*(a + b*Log[c*x^n])/x^2 + (E^((2*a)/(b*n))*(c*x^n)^(2/n)
)*ExpIntegralEi[-(((2 - b*d*n)*(a + b*Log[c*x^n]))/(b*n)))]/(4*x^2) + (E^((
2*a)/(b*n))*(c*x^n)^(2/n)*ExpIntegralEi[-(((2 + b*d*n)*(a + b*Log[c*x^n]))/
(b*n)))]/(4*x^2)
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 2209

```
Int[(F_)^((g_)*(e_) + (f_)*(x_)))/((c_) + (d_)*(x_)), x_Symbol] := Si
mp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; F
reeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]
```

Rule 2347

```
Int[((a_) + Log[(c_)*(x_)^(n_)])*(b_)^(p_)*((d_)*(x_)^(m_)), x_Symbol
] := Dist[(d*x)^(m + 1)/(d*n*(c*x^n)^(m + 1)/n), Subst[Int[E^(((m + 1)/n)
*x)*(a + b*x)^p, x], x, Log[c*x^n]], x] /; FreeQ[{a, b, c, d, m, n, p}, x]
```

Rule 5651

```
Int[Cosh[((a_) + Log[(c_)*(x_)^(n_)])*(b_)*(d_)]*(((e_) + Log[(g_)*(
x_)^(m_)]*(f_)*(h_))^(q_)*((i_)*(x_)^(r_)), x_Symbol] := Dist[((i*x)
^r*(1/((c*x^n)^(b*d)*(2*x^(r - b*d*n)))))/E^(a*d), Int[x^(r - b*d*n)*(h*(e
+ f*Log[g*x^m]))^q, x], x] + Dist[E^(a*d)*(i*x)^r*((c*x^n)^(b*d)/(2*x^(r +
b*d*n))), Int[x^(r + b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] /; FreeQ[{a, b
, c, d, e, f, g, h, i, m, n, q, r}, x]
```

Rule 6691

```
Int[CoshIntegral[((a_) + Log[(c_)*(x_)^(n_)])*(b_)*(d_)]*((e_)*(x_)^(
m_)), x_Symbol] := Simp[(e*x)^(m + 1)*(CoshIntegral[d*(a + b*Log[c*x^n])]/
(e*(m + 1))), x] - Dist[b*d*(n/(m + 1)), Int[(e*x)^m*(Cosh[d*(a + b*Log[c*x
^n])]/(d*(a + b*Log[c*x^n]))), x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] &&
NeQ[m, -1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Chi}(d(a + b \log(cx^n)))}{2x^2} + \frac{1}{2}(bdn) \int \frac{\cosh(d(a + b \log(cx^n)))}{dx^3(a + b \log(cx^n))} dx \\ &= -\frac{\text{Chi}(d(a + b \log(cx^n)))}{2x^2} + \frac{1}{2}(bn) \int \frac{\cosh(d(a + b \log(cx^n)))}{x^3(a + b \log(cx^n))} dx \end{aligned}$$

$$\begin{aligned}
&= -\frac{\text{Chi}(d(a + b \log(cx^n)))}{2x^2} + \frac{1}{4} \left(be^{-ad} n x^{bdn} (cx^n)^{-bd} \right) \int \frac{x^{-3-bdn}}{a + b \log(cx^n)} dx \\
&\quad + \frac{1}{4} \left(be^{ad} n x^{-bdn} (cx^n)^{bd} \right) \int \frac{x^{-3+bdn}}{a + b \log(cx^n)} dx \\
&= -\frac{\text{Chi}(d(a + b \log(cx^n)))}{2x^2} \\
&\quad + \frac{\left(be^{-ad} (cx^n)^{-bd - \frac{-2-bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(-2-bdn)x}{a+bx}}}{a+bx} dx, x, \log(cx^n) \right)}{4x^2} \\
&\quad + \frac{\left(be^{ad} (cx^n)^{bd - \frac{-2+bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(-2+bdn)x}{a+bx}}}{a+bx} dx, x, \log(cx^n) \right)}{4x^2} \\
&= -\frac{\text{Chi}(d(a + b \log(cx^n)))}{2x^2} + \frac{e^{\frac{2a}{bn}} (cx^n)^{2/n} \text{ExpIntegralEi} \left(-\frac{(2-bdn)(a+b \log(cx^n))}{bn} \right)}{4x^2} \\
&\quad + \frac{e^{\frac{2a}{bn}} (cx^n)^{2/n} \text{ExpIntegralEi} \left(-\frac{(2+bdn)(a+b \log(cx^n))}{bn} \right)}{4x^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.26 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.12

$$\begin{aligned}
&\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x^3} dx \\
&= -\frac{\text{Chi}(d(a + b \log(cx^n)))}{2x^2} \\
&\quad + \frac{1}{4} e^{-\frac{(-2+bdn)(a+b(-n \log(x)+\log(cx^n)))}{bn}} \left(\text{ExpIntegralEi} \left(\frac{(-2 + bdn)(a + b \log(cx^n))}{bn} \right) \right. \\
&\quad \left. + \text{ExpIntegralEi} \left(-\frac{(2 + bdn)(a + b \log(cx^n))}{bn} \right) \right) (\cosh(d(a + b(-n \log(x) + \log(cx^n)))) \\
&\quad \quad \quad + \sinh(d(a + b(-n \log(x) + \log(cx^n))))))
\end{aligned}$$

[In] Integrate[CoshIntegral[d*(a + b*Log[c*x^n])/x^3,x]

[Out] -1/2*CoshIntegral[d*(a + b*Log[c*x^n])/x^2 + ((ExpIntegralEi[((-2 + b*d*n)*(a + b*Log[c*x^n])/(b*n)] + ExpIntegralEi[-(((2 + b*d*n)*(a + b*Log[c*x^n]))/(b*n))])*(Cosh[d*(a + b*(-n*Log[x]) + Log[c*x^n]))] + Sinh[d*(a + b*(-n*Log[x]) + Log[c*x^n]))])/(4*E^(((2 + b*d*n)*(a + b*(-n*Log[x]) + Log[c*x^n]))/(b*n)))

Maple [F]

$$\int \frac{\text{Chi}(d(a + b \ln(cx^n)))}{x^3} dx$$

[In] int(Chi(d*(a+b*ln(c*x^n)))/x^3,x)

[Out] int(Chi(d*(a+b*ln(c*x^n)))/x^3,x)

Fricas [F]

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\text{Chi}((b \log(cx^n) + a)d)}{x^3} dx$$

[In] integrate(Chi(d*(a+b*log(c*x^n)))/x^3,x, algorithm="fricas")

[Out] integral(cosh_integral(b*d*log(c*x^n) + a*d)/x^3, x)

Sympy [F]

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\text{Chi}(ad + bd \log(cx^n))}{x^3} dx$$

[In] integrate(Chi(d*(a+b*ln(c*x**n)))/x**3,x)

[Out] Integral(Chi(a*d + b*d*log(c*x**n))/x**3, x)

Maxima [F]

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\text{Chi}((b \log(cx^n) + a)d)}{x^3} dx$$

[In] integrate(Chi(d*(a+b*log(c*x^n)))/x^3,x, algorithm="maxima")

[Out] integrate(Chi((b*log(c*x^n) + a)*d)/x^3, x)

Giac [F]

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\text{Chi}((b \log(cx^n) + a)d)}{x^3} dx$$

[In] integrate(Chi(d*(a+b*log(c*x^n)))/x^3,x, algorithm="giac")

[Out] integrate(Chi((b*log(c*x^n) + a)*d)/x^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\text{Chi}(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\text{coshint}(d(a + b \ln(cx^n)))}{x^3} dx$$

[In] int(coshint(d*(a + b*log(c*x^n)))/x^3,x)

[Out] int(coshint(d*(a + b*log(c*x^n)))/x^3, x)

3.106 $\int (ex)^m \mathbf{Chi}(d(a + b \log(cx^n))) dx$

Optimal result	522
Rubi [A] (verified)	522
Mathematica [A] (verified)	525
Maple [F]	525
Fricas [F]	525
Sympy [F]	525
Maxima [F]	526
Giac [F]	526
Mupad [F(-1)]	526

Optimal result

Integrand size = 19, antiderivative size = 167

$$\begin{aligned} & \int (ex)^m \mathbf{Chi}(d(a + b \log(cx^n))) dx \\ &= \frac{(ex)^{1+m} \mathbf{Chi}(d(a + b \log(cx^n)))}{e(1+m)} \\ & \quad - \frac{e^{-\frac{a(1+m)}{bn}} x (ex)^m (cx^n)^{-\frac{1+m}{n}} \text{ExpIntegralEi}\left(\frac{(1+m-bdn)(a+b \log(cx^n))}{bn}\right)}{2(1+m)} \\ & \quad - \frac{e^{-\frac{a(1+m)}{bn}} x (ex)^m (cx^n)^{-\frac{1+m}{n}} \text{ExpIntegralEi}\left(\frac{(1+m+bdn)(a+b \log(cx^n))}{bn}\right)}{2(1+m)} \end{aligned}$$

[Out] (e*x)^(1+m)*Chi(d*(a+b*ln(c*x^n)))/e/(1+m)-1/2*x*(e*x)^m*Ei((-b*d*n+m+1)*(a+b*ln(c*x^n))/b/n)/exp(a*(1+m)/b/n)/(1+m)/((c*x^n)^((1+m)/n))-1/2*x*(e*x)^m*Ei((b*d*n+m+1)*(a+b*ln(c*x^n))/b/n)/exp(a*(1+m)/b/n)/(1+m)/((c*x^n)^((1+m)/n))

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used

= {6691, 12, 5651, 2347, 2209}

$$\int (ex)^m \text{Chi}(d(a + b \log(cx^n))) dx$$

$$= \frac{(ex)^{m+1} \text{Chi}(d(a + b \log(cx^n)))}{e(m+1)}$$

$$- \frac{x(ex)^m e^{-\frac{a(m+1)}{bn}} (cx^n)^{-\frac{m+1}{n}} \text{ExpIntegralEi}\left(\frac{(m-bdn+1)(a+b \log(cx^n))}{bn}\right)}{2(m+1)}$$

$$- \frac{x(ex)^m e^{-\frac{a(m+1)}{bn}} (cx^n)^{-\frac{m+1}{n}} \text{ExpIntegralEi}\left(\frac{(m+bdn+1)(a+b \log(cx^n))}{bn}\right)}{2(m+1)}$$

[In] Int[(e*x)^m*CoshIntegral[d*(a + b*Log[c*x^n]),x]

[Out] ((e*x)^(1 + m)*CoshIntegral[d*(a + b*Log[c*x^n])]/(e*(1 + m)) - (x*(e*x)^m*ExpIntegralEi[((1 + m - b*d*n)*(a + b*Log[c*x^n])/(b*n))]/(2*E^((a*(1 + m))/(b*n)))*(1 + m)*(c*x^n)^((1 + m)/n)) - (x*(e*x)^m*ExpIntegralEi[((1 + m + b*d*n)*(a + b*Log[c*x^n])/(b*n))]/(2*E^((a*(1 + m))/(b*n)))*(1 + m)*(c*x^n)^((1 + m)/n))

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 2209

Int[(F_)^((g_)*((e_) + (f_)*(x_)))/((c_) + (d_)*(x_)), x_Symbol] := Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[\$UseGamma]

Rule 2347

Int[((a_) + Log[(c_)*(x_)^(n_)])*(b_)^(p_)*((d_)*(x_)^(m_)), x_Symbol] := Dist[(d*x)^(m + 1)/(d*n*(c*x^n)^((m + 1)/n)), Subst[Int[E^((m + 1)/n)*x*(a + b*x)^p, x], x, Log[c*x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x]

Rule 5651

Int[Cosh[((a_) + Log[(c_)*(x_)^(n_)])*(b_)]*(d_)]*(((e_) + Log[(g_)*(x_)^(m_)])*(f_))*(h_)^(q_)*((i_)*(x_)^(r_)), x_Symbol] := Dist[((i*x)^r*(1/((c*x^n)^(b*d)*(2*x^(r - b*d*n))))/E^(a*d), Int[x^(r - b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] + Dist[E^(a*d)*(i*x)^r*((c*x^n)^(b*d)/(2*x^(r + b*d*n))), Int[x^(r + b*d*n)*(h*(e + f*Log[g*x^m]))^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, m, n, q, r}, x]

Rule 6691

Int[CoshIntegral[(a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.)]*(d_.)]*((e_.)*(x_)^(m_.), x_Symbol] :> Simp[(e*x)^(m + 1)*(CoshIntegral[d*(a + b*Log[c*x^n])]/(e*(m + 1))), x] - Dist[b*d*(n/(m + 1)), Int[(e*x)^m*(Cosh[d*(a + b*Log[c*x^n])])]/(d*(a + b*Log[c*x^n]))], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(ex)^{1+m} \text{Chi}(d(a + b \log(cx^n)))}{e(1+m)} - \frac{(bdn) \int \frac{(ex)^m \cosh(d(a+b \log(cx^n)))}{d(a+b \log(cx^n))} dx}{1+m} \\
&= \frac{(ex)^{1+m} \text{Chi}(d(a + b \log(cx^n)))}{e(1+m)} - \frac{(bn) \int \frac{(ex)^m \cosh(d(a+b \log(cx^n)))}{a+b \log(cx^n)} dx}{1+m} \\
&= \frac{(ex)^{1+m} \text{Chi}(d(a + b \log(cx^n)))}{e(1+m)} - \frac{\left(be^{-ad} n x^{-m+bdn} (ex)^m (cx^n)^{-bd} \right) \int \frac{x^{m-bdn}}{a+b \log(cx^n)} dx}{2(1+m)} \\
&\quad - \frac{\left(be^{ad} n x^{-m-bdn} (ex)^m (cx^n)^{bd} \right) \int \frac{x^{m+bdn}}{a+b \log(cx^n)} dx}{2(1+m)} \\
&= \frac{(ex)^{1+m} \text{Chi}(d(a + b \log(cx^n)))}{e(1+m)} \\
&\quad - \frac{\left(be^{-ad} x (ex)^m (cx^n)^{-bd - \frac{1+m-bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(1+m-bdn)x}{a+bx}}}{a+bx} dx, x, \log(cx^n) \right)}{2(1+m)} \\
&\quad - \frac{\left(be^{ad} x (ex)^m (cx^n)^{bd - \frac{1+m+bdn}{n}} \right) \text{Subst} \left(\int \frac{e^{\frac{(1+m+bdn)x}{a+bx}}}{a+bx} dx, x, \log(cx^n) \right)}{2(1+m)} \\
&= \frac{(ex)^{1+m} \text{Chi}(d(a + b \log(cx^n)))}{e(1+m)} \\
&\quad - \frac{e^{-\frac{a(1+m)}{bn}} x (ex)^m (cx^n)^{-\frac{1+m}{n}} \text{ExpIntegralEi} \left(\frac{(1+m-bdn)(a+b \log(cx^n))}{bn} \right)}{2(1+m)} \\
&\quad - \frac{e^{-\frac{a(1+m)}{bn}} x (ex)^m (cx^n)^{-\frac{1+m}{n}} \text{ExpIntegralEi} \left(\frac{(1+m+bdn)(a+b \log(cx^n))}{bn} \right)}{2(1+m)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.37 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.71

$$\int (ex)^m \text{Chi}(d(a + b \log(cx^n))) dx$$

$$= \frac{(ex)^m \left(2x \text{Chi}(d(a + b \log(cx^n))) - e^{-\frac{(1+m)(a - bn \log(x) + b \log(cx^n))}{bn}} x^{-m} \left(\text{ExpIntegralEi} \left(\frac{(1+m - bdn)(a + b \log(cx^n))}{bn} \right) \right) \right)}{2(1+m)}$$

[In] Integrate[(e*x)^m*CoshIntegral[d*(a + b*Log[c*x^n])],x]

[Out] ((e*x)^m*(2*x*CoshIntegral[d*(a + b*Log[c*x^n])] - (ExpIntegralEi[((1 + m - b*d*n)*(a + b*Log[c*x^n])]/(b*n)] + ExpIntegralEi[((1 + m + b*d*n)*(a + b*Log[c*x^n])]/(b*n)]))/(E^(((1 + m)*(a - b*n*Log[x] + b*Log[c*x^n])/(b*n))*x^m)))/(2*(1 + m))

Maple [F]

$$\int (ex)^m \text{Chi}(d(a + b \ln(cx^n))) dx$$

[In] int((e*x)^m*Chi(d*(a+b*ln(c*x^n))),x)

[Out] int((e*x)^m*Chi(d*(a+b*ln(c*x^n))),x)

Fricas [F]

$$\int (ex)^m \text{Chi}(d(a + b \log(cx^n))) dx = \int (ex)^m \text{Chi}((b \log(cx^n) + a)d) dx$$

[In] integrate((e*x)^m*Chi(d*(a+b*log(c*x^n))),x, algorithm="fricas")

[Out] integral((e*x)^m*cosh_integral(b*d*log(c*x^n) + a*d), x)

Sympy [F]

$$\int (ex)^m \text{Chi}(d(a + b \log(cx^n))) dx = \int (ex)^m \text{Chi}(ad + bd \log(cx^n)) dx$$

[In] integrate((e*x)**m*Chi(d*(a+b*ln(c*x**n))),x)

[Out] Integral((e*x)**m*Chi(a*d + b*d*log(c*x**n)), x)

Maxima [F]

$$\int (ex)^m \text{Chi}(d(a + b \log(cx^n))) dx = \int (ex)^m \text{Chi}((b \log(cx^n) + a)d) dx$$

[In] integrate((e*x)^m*Chi(d*(a+b*log(c*x^n))),x, algorithm="maxima")

[Out] integrate((e*x)^m*Chi((b*log(c*x^n) + a)*d), x)

Giac [F]

$$\int (ex)^m \text{Chi}(d(a + b \log(cx^n))) dx = \int (ex)^m \text{Chi}((b \log(cx^n) + a)d) dx$$

[In] integrate((e*x)^m*Chi(d*(a+b*log(c*x^n))),x, algorithm="giac")

[Out] integrate((e*x)^m*Chi((b*log(c*x^n) + a)*d), x)

Mupad [F(-1)]

Timed out.

$$\int (ex)^m \text{Chi}(d(a + b \log(cx^n))) dx = \int \text{coshint}(d(a + b \ln(cx^n))) (ex)^m dx$$

[In] int(coshint(d*(a + b*log(c*x^n)))*(e*x)^m,x)

[Out] int(coshint(d*(a + b*log(c*x^n)))*(e*x)^m, x)

3.107 $\int \frac{\cosh(bx)\mathbf{Chi}(bx)}{x^3} dx$

Optimal result	527
Rubi [A] (verified)	527
Mathematica [A] (verified)	530
Maple [F]	530
Fricas [F]	530
Sympy [F]	531
Maxima [F]	531
Giac [F]	531
Mupad [F(-1)]	531

Optimal result

Integrand size = 12, antiderivative size = 96

$$\int \frac{\cosh(bx)\mathbf{Chi}(bx)}{x^3} dx = -\frac{\cosh^2(bx)}{4x^2} - \frac{\cosh(bx)\mathbf{Chi}(bx)}{2x^2} + \frac{1}{4}b^2\mathbf{Chi}(bx)^2 + b^2\mathbf{Chi}(2bx) - \frac{b\cosh(bx)\sinh(bx)}{2x} - \frac{b\mathbf{Chi}(bx)\sinh(bx)}{2x} - \frac{b\sinh(2bx)}{4x}$$

[Out] 1/4*b^2*Chi(b*x)^2+b^2*Chi(2*b*x)-1/2*Chi(b*x)*cosh(b*x)/x^2-1/4*cosh(b*x)^2/x^2-1/2*b*Chi(b*x)*sinh(b*x)/x-1/2*b*cosh(b*x)*sinh(b*x)/x-1/4*b*sinh(2*b*x)/x

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {6680, 6686, 6818, 12, 5556, 3378, 3382, 3395, 29, 3393}

$$\int \frac{\cosh(bx)\mathbf{Chi}(bx)}{x^3} dx = \frac{1}{4}b^2\mathbf{Chi}(bx)^2 + b^2\mathbf{Chi}(2bx) - \frac{\mathbf{Chi}(bx)\cosh(bx)}{2x^2} - \frac{b\mathbf{Chi}(bx)\sinh(bx)}{2x} - \frac{\cosh^2(bx)}{4x^2} - \frac{b\sinh(2bx)}{4x} - \frac{b\sinh(bx)\cosh(bx)}{2x}$$

[In] Int[(Cosh[b*x]*CoshIntegral[b*x])/x^3,x]

[Out] -1/4*Cosh[b*x]^2/x^2 - (Cosh[b*x]*CoshIntegral[b*x])/(2*x^2) + (b^2*CoshIntegral[b*x]^2)/4 + b^2*CoshIntegral[2*b*x] - (b*Cosh[b*x]*Sinh[b*x])/(2*x) - (b*CoshIntegral[b*x]*Sinh[b*x])/(2*x) - (b*Sinh[2*b*x])/(4*x)

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 29

`Int[(x_)^(-1), x_Symbol] := Simp[Log[x], x]`

Rule 3378

`Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Dist[f/(d*(m + 1)), Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]`

Rule 3382

`Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`

Rule 3393

`Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

Rule 3395

`Int[((c_.) + (d_.)*(x_))^(m_)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(c + d*x)^(m + 1)*((b*Sine[e + f*x])^n/(d*(m + 1))), x] + (Dist[b^2*f^2*n*((n - 1)/(d^2*(m + 1)*(m + 2))), Int[(c + d*x)^(m + 2)*(b*Sine[e + f*x])^(n - 2), x], x] - Dist[f^2*(n^2/(d^2*(m + 1)*(m + 2))), Int[(c + d*x)^(m + 2)*(b*Sine[e + f*x])^n, x], x] - Simp[b*f*n*(c + d*x)^(m + 2)*Cos[e + f*x]*((b*Sine[e + f*x])^(n - 1)/(d^2*(m + 1)*(m + 2))), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && LtQ[m, -2]`

Rule 5556

`Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

Rule 6680

`Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*Cosh[a + b*x]*(CoshIntegral`


```

ral[c + d*x]/(f*(m + 1)), x] + (-Dist[b/(f*(m + 1)), Int[(e + f*x)^(m + 1)
*Sinh[a + b*x]*CoshIntegral[c + d*x], x], x] - Dist[d/(f*(m + 1)), Int[(e +
f*x)^(m + 1)*Cosh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x]) /; FreeQ[{a,
b, c, d, e, f}, x] && ILtQ[m, -1]

```

Rule 6686

```

Int[CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_)*Sinh[(a_.) +
(b_.)*(x_)], x_Symbol] := Simp[(e + f*x)^(m + 1)*Sinh[a + b*x]*(CoshIntegr
al[c + d*x]/(f*(m + 1))), x] + (-Dist[b/(f*(m + 1)), Int[(e + f*x)^(m + 1)*
Cosh[a + b*x]*CoshIntegral[c + d*x], x], x] - Dist[d/(f*(m + 1)), Int[(e +
f*x)^(m + 1)*Sinh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x]) /; FreeQ[{a,
b, c, d, e, f}, x] && ILtQ[m, -1]

```

Rule 6818

```

Int[(u_)*(y_)^(m_), x_Symbol] := With[{q = DerivativeDivides[y, u, x]}, Si
mp[q*(y^(m + 1)/(m + 1)), x] /; !FalseQ[q]] /; FreeQ[m, x] && NeQ[m, -1]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\cosh(bx)\text{Chi}(bx)}{2x^2} + \frac{1}{2}b \int \frac{\cosh^2(bx)}{bx^3} dx + \frac{1}{2}b \int \frac{\text{Chi}(bx) \sinh(bx)}{x^2} dx \\
&= -\frac{\cosh(bx)\text{Chi}(bx)}{2x^2} - \frac{b\text{Chi}(bx) \sinh(bx)}{2x} + \frac{1}{2} \int \frac{\cosh^2(bx)}{x^3} dx \\
&\quad + \frac{1}{2}b^2 \int \frac{\cosh(bx)\text{Chi}(bx)}{x} dx + \frac{1}{2}b^2 \int \frac{\cosh(bx) \sinh(bx)}{bx^2} dx \\
&= -\frac{\cosh^2(bx)}{4x^2} - \frac{\cosh(bx)\text{Chi}(bx)}{2x^2} + \frac{1}{4}b^2\text{Chi}(bx)^2 \\
&\quad - \frac{b \cosh(bx) \sinh(bx)}{2x} - \frac{b\text{Chi}(bx) \sinh(bx)}{2x} \\
&\quad + \frac{1}{2}b \int \frac{\cosh(bx) \sinh(bx)}{x^2} dx - \frac{1}{2}b^2 \int \frac{1}{x} dx + b^2 \int \frac{\cosh^2(bx)}{x} dx \\
&= -\frac{\cosh^2(bx)}{4x^2} - \frac{\cosh(bx)\text{Chi}(bx)}{2x^2} + \frac{1}{4}b^2\text{Chi}(bx)^2 - \frac{1}{2}b^2 \log(x) - \frac{b \cosh(bx) \sinh(bx)}{2x} \\
&\quad - \frac{b\text{Chi}(bx) \sinh(bx)}{2x} + \frac{1}{2}b \int \frac{\sinh(2bx)}{2x^2} dx + b^2 \int \left(\frac{1}{2x} + \frac{\cosh(2bx)}{2x} \right) dx \\
&= -\frac{\cosh^2(bx)}{4x^2} - \frac{\cosh(bx)\text{Chi}(bx)}{2x^2} + \frac{1}{4}b^2\text{Chi}(bx)^2 - \frac{b \cosh(bx) \sinh(bx)}{2x} \\
&\quad - \frac{b\text{Chi}(bx) \sinh(bx)}{2x} + \frac{1}{4}b \int \frac{\sinh(2bx)}{x^2} dx + \frac{1}{2}b^2 \int \frac{\cosh(2bx)}{x} dx
\end{aligned}$$

$$\begin{aligned}
&= -\frac{\cosh^2(bx)}{4x^2} - \frac{\cosh(bx)\text{Chi}(bx)}{2x^2} + \frac{1}{4}b^2\text{Chi}(bx)^2 + \frac{1}{2}b^2\text{Chi}(2bx) \\
&\quad - \frac{b \cosh(bx) \sinh(bx)}{2x} - \frac{b\text{Chi}(bx) \sinh(bx)}{2x} - \frac{b \sinh(2bx)}{4x} + \frac{1}{2}b^2 \int \frac{\cosh(2bx)}{x} dx \\
&= -\frac{\cosh^2(bx)}{4x^2} - \frac{\cosh(bx)\text{Chi}(bx)}{2x^2} + \frac{1}{4}b^2\text{Chi}(bx)^2 + b^2\text{Chi}(2bx) \\
&\quad - \frac{b \cosh(bx) \sinh(bx)}{2x} - \frac{b\text{Chi}(bx) \sinh(bx)}{2x} - \frac{b \sinh(2bx)}{4x}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.00

$$\int \frac{\cosh(bx)\text{Chi}(bx)}{x^3} dx = -\frac{\cosh^2(bx)}{4x^2} - \frac{\cosh(bx)\text{Chi}(bx)}{2x^2} + \frac{1}{4}b^2\text{Chi}(bx)^2 + b^2\text{Chi}(2bx) \\
- \frac{b \cosh(bx) \sinh(bx)}{2x} - \frac{b\text{Chi}(bx) \sinh(bx)}{2x} - \frac{b \sinh(2bx)}{4x}$$

[In] Integrate[(Cosh[b*x]*CoshIntegral[b*x])/x^3,x]

[Out] -1/4*Cosh[b*x]^2/x^2 - (Cosh[b*x]*CoshIntegral[b*x])/(2*x^2) + (b^2*CoshIntegral[b*x]^2)/4 + b^2*CoshIntegral[2*b*x] - (b*Cosh[b*x]*Sinh[b*x])/(2*x) - (b*CoshIntegral[b*x]*Sinh[b*x])/(2*x) - (b*Sinh[2*b*x])/(4*x)

Maple [F]

$$\int \frac{\text{Chi}(bx) \cosh(bx)}{x^3} dx$$

[In] int(Chi(b*x)*cosh(b*x)/x^3,x)

[Out] int(Chi(b*x)*cosh(b*x)/x^3,x)

Fricas [F]

$$\int \frac{\cosh(bx)\text{Chi}(bx)}{x^3} dx = \int \frac{\text{Chi}(bx) \cosh(bx)}{x^3} dx$$

[In] integrate(Chi(b*x)*cosh(b*x)/x^3,x, algorithm="fricas")

[Out] integral(cosh(b*x)*cosh_integral(b*x)/x^3, x)

Sympy [F]

$$\int \frac{\cosh(bx)\operatorname{Chi}(bx)}{x^3} dx = \int \frac{\cosh(bx)\operatorname{Chi}(bx)}{x^3} dx$$

[In] integrate(Chi(b*x)*cosh(b*x)/x**3,x)

[Out] Integral(cosh(b*x)*Chi(b*x)/x**3, x)

Maxima [F]

$$\int \frac{\cosh(bx)\operatorname{Chi}(bx)}{x^3} dx = \int \frac{\operatorname{Chi}(bx)\cosh(bx)}{x^3} dx$$

[In] integrate(Chi(b*x)*cosh(b*x)/x^3,x, algorithm="maxima")

[Out] integrate(Chi(b*x)*cosh(b*x)/x^3, x)

Giac [F]

$$\int \frac{\cosh(bx)\operatorname{Chi}(bx)}{x^3} dx = \int \frac{\operatorname{Chi}(bx)\cosh(bx)}{x^3} dx$$

[In] integrate(Chi(b*x)*cosh(b*x)/x^3,x, algorithm="giac")

[Out] integrate(Chi(b*x)*cosh(b*x)/x^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cosh(bx)\operatorname{Chi}(bx)}{x^3} dx = \int \frac{\operatorname{coshint}(bx)\cosh(bx)}{x^3} dx$$

[In] int((coshint(b*x)*cosh(b*x))/x^3,x)

[Out] int((coshint(b*x)*cosh(b*x))/x^3, x)

3.108 $\int \frac{\cosh(bx)\mathbf{Chi}(bx)}{x^2} dx$

Optimal result	532
Rubi [N/A]	532
Mathematica [N/A]	533
Maple [N/A] (verified)	533
Fricas [N/A]	534
Sympy [N/A]	534
Maxima [N/A]	534
Giac [N/A]	535
Mupad [N/A]	535

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\cosh(bx)\mathbf{Chi}(bx)}{x^2} dx = -\frac{\cosh^2(bx)}{x} - \frac{\cosh(bx)\mathbf{Chi}(bx)}{x} + b\mathbf{Shi}(2bx) + b\mathbf{Int}\left(\frac{\mathbf{Chi}(bx)\sinh(bx)}{x}, x\right)$$

[Out] b*CannotIntegrate(Chi(b*x)*sinh(b*x)/x,x)-Chi(b*x)*cosh(b*x)/x-cosh(b*x)^2/x+b*Shi(2*b*x)

Rubi [N/A]

Not integrable

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\cosh(bx)\mathbf{Chi}(bx)}{x^2} dx = \int \frac{\cosh(bx)\mathbf{Chi}(bx)}{x^2} dx$$

[In] Int[(Cosh[b*x]*CoshIntegral[b*x])/x^2,x]

[Out] -(Cosh[b*x]^2/x) - (Cosh[b*x]*CoshIntegral[b*x])/x + b*SinhIntegral[2*b*x] + b*Defer[Int] [(CoshIntegral[b*x]*Sinh[b*x])/x, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\cosh(bx)\text{Chi}(bx)}{x} + b \int \frac{\cosh^2(bx)}{bx^2} dx + b \int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx \\
 &= -\frac{\cosh(bx)\text{Chi}(bx)}{x} + b \int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx + \int \frac{\cosh^2(bx)}{x^2} dx \\
 &= -\frac{\cosh^2(bx)}{x} - \frac{\cosh(bx)\text{Chi}(bx)}{x} + (2ib) \int -\frac{i \sinh(2bx)}{2x} dx + b \int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx \\
 &= -\frac{\cosh^2(bx)}{x} - \frac{\cosh(bx)\text{Chi}(bx)}{x} + b \int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx + b \int \frac{\sinh(2bx)}{x} dx \\
 &= -\frac{\cosh^2(bx)}{x} - \frac{\cosh(bx)\text{Chi}(bx)}{x} + b\text{Shi}(2bx) + b \int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx
 \end{aligned}$$

Mathematica [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx)\text{Chi}(bx)}{x^2} dx = \int \frac{\cosh(bx)\text{Chi}(bx)}{x^2} dx$$

[In] Integrate[(Cosh[b*x]*CoshIntegral[b*x])/x^2,x]

[Out] Integrate[(Cosh[b*x]*CoshIntegral[b*x])/x^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.21 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx) \cosh(bx)}{x^2} dx$$

[In] int(Chi(b*x)*cosh(b*x)/x^2,x)

[Out] int(Chi(b*x)*cosh(b*x)/x^2,x)

Fricas [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx)\text{Chi}(bx)}{x^2} dx = \int \frac{\text{Chi}(bx)\cosh(bx)}{x^2} dx$$

[In] integrate(Chi(b*x)*cosh(b*x)/x^2,x, algorithm="fricas")

[Out] integral(cosh(b*x)*cosh_integral(b*x)/x^2, x)

Sympy [N/A]

Not integrable

Time = 3.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx)\text{Chi}(bx)}{x^2} dx = \int \frac{\cosh(bx)\text{Chi}(bx)}{x^2} dx$$

[In] integrate(Chi(b*x)*cosh(b*x)/x**2,x)

[Out] Integral(cosh(b*x)*Chi(b*x)/x**2, x)

Maxima [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx)\text{Chi}(bx)}{x^2} dx = \int \frac{\text{Chi}(bx)\cosh(bx)}{x^2} dx$$

[In] integrate(Chi(b*x)*cosh(b*x)/x^2,x, algorithm="maxima")

[Out] integrate(Chi(b*x)*cosh(b*x)/x^2, x)

Giac [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx)\text{Chi}(bx)}{x^2} dx = \int \frac{\text{Chi}(bx)\cosh(bx)}{x^2} dx$$

[In] integrate(Chi(b*x)*cosh(b*x)/x^2,x, algorithm="giac")

[Out] integrate(Chi(b*x)*cosh(b*x)/x^2, x)

Mupad [N/A]

Not integrable

Time = 4.64 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cosh(bx)\text{Chi}(bx)}{x^2} dx = \int \frac{\text{coshint}(bx)\cosh(bx)}{x^2} dx$$

[In] int((coshint(b*x)*cosh(b*x))/x^2,x)

[Out] int((coshint(b*x)*cosh(b*x))/x^2, x)

3.109 $\int \frac{\cosh(bx)\mathbf{Chi}(bx)}{x} dx$

Optimal result	536
Rubi [A] (verified)	536
Mathematica [A] (verified)	537
Maple [A] (verified)	537
Fricas [F]	537
Sympy [A] (verification not implemented)	537
Maxima [F]	538
Giac [F]	538
Mupad [F(-1)]	538

Optimal result

Integrand size = 12, antiderivative size = 10

$$\int \frac{\cosh(bx)\mathbf{Chi}(bx)}{x} dx = \frac{\mathbf{Chi}(bx)^2}{2}$$

[Out] 1/2*Chi(b*x)^2

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {6818}

$$\int \frac{\cosh(bx)\mathbf{Chi}(bx)}{x} dx = \frac{\mathbf{Chi}(bx)^2}{2}$$

[In] Int[(Cosh[b*x]*CoshIntegral[b*x])/x,x]

[Out] CoshIntegral[b*x]^2/2

Rule 6818

Int[(u_)*(y_)^(m_.), x_Symbol] := With[{q = DerivativeDivides[y, u, x]}, Si
mp[q*(y^(m + 1)/(m + 1)), x] /; !FalseQ[q]] /; FreeQ[m, x] && NeQ[m, -1]

Rubi steps

$$\text{integral} = \frac{\mathbf{Chi}(bx)^2}{2}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\cosh(bx)\text{Chi}(bx)}{x} dx = \frac{\text{Chi}(bx)^2}{2}$$

[In] Integrate[(Cosh[b*x]*CoshIntegral[b*x])/x,x]

[Out] CoshIntegral[b*x]^2/2

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

method	result	size
derivativedivides	$\frac{\text{Chi}(bx)^2}{2}$	9
default	$\frac{\text{Chi}(bx)^2}{2}$	9

[In] int(Chi(b*x)*cosh(b*x)/x,x,method=_RETURNVERBOSE)

[Out] 1/2*Chi(b*x)^2

Fricas [F]

$$\int \frac{\cosh(bx)\text{Chi}(bx)}{x} dx = \int \frac{\text{Chi}(bx)\cosh(bx)}{x} dx$$

[In] integrate(Chi(b*x)*cosh(b*x)/x,x, algorithm="fricas")

[Out] integral(cosh(b*x)*cosh_integral(b*x)/x, x)

Sympy [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \frac{\cosh(bx)\text{Chi}(bx)}{x} dx = \frac{\text{Chi}^2(bx)}{2}$$

[In] integrate(Chi(b*x)*cosh(b*x)/x,x)

[Out] Chi(b*x)**2/2

Maxima [F]

$$\int \frac{\cosh(bx)\text{Chi}(bx)}{x} dx = \int \frac{\text{Chi}(bx) \cosh(bx)}{x} dx$$

[In] integrate(Chi(b*x)*cosh(b*x)/x,x, algorithm="maxima")

[Out] integrate(Chi(b*x)*cosh(b*x)/x, x)

Giac [F]

$$\int \frac{\cosh(bx)\text{Chi}(bx)}{x} dx = \int \frac{\text{Chi}(bx) \cosh(bx)}{x} dx$$

[In] integrate(Chi(b*x)*cosh(b*x)/x,x, algorithm="giac")

[Out] integrate(Chi(b*x)*cosh(b*x)/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cosh(bx)\text{Chi}(bx)}{x} dx = \frac{\text{coshint}(bx)^2}{2}$$

[In] int((coshint(b*x)*cosh(b*x))/x,x)

[Out] coshint(b*x)^2/2

3.110 $\int \cosh(bx)\mathbf{Chi}(bx) dx$

Optimal result	539
Rubi [A] (verified)	539
Mathematica [A] (verified)	540
Maple [A] (verified)	541
Fricas [F]	541
Sympy [F]	541
Maxima [F]	541
Giac [F]	542
Mupad [F(-1)]	542

Optimal result

Integrand size = 9, antiderivative size = 25

$$\int \cosh(bx)\mathbf{Chi}(bx) dx = \frac{\mathbf{Chi}(bx) \sinh(bx)}{b} - \frac{\mathbf{Shi}(2bx)}{2b}$$

[Out] $-1/2*\mathbf{Shi}(2*b*x)/b+\mathbf{Chi}(b*x)*\sinh(b*x)/b$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {6676, 12, 5556, 3379}

$$\int \cosh(bx)\mathbf{Chi}(bx) dx = \frac{\mathbf{Chi}(bx) \sinh(bx)}{b} - \frac{\mathbf{Shi}(2bx)}{2b}$$

[In] $\text{Int}[\text{Cosh}[b*x]*\text{CoshIntegral}[b*x], x]$

[Out] $(\text{CoshIntegral}[b*x]*\text{Sinh}[b*x])/b - \text{SinhIntegral}[2*b*x]/(2*b)$

Rule 12

$\text{Int}[(a_*)(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[u, (b_*)(v_)] /; \text{FreeQ}[b, x]$

Rule 3379

$\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] \rightarrow \text{Simp}[I*(\text{SinhIntegral}[c*f*(fz/d) + f*fz*x]/d), x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \ \&\& \ \text{EqQ}[d*e - c*f*fz*I, 0]$

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

Rule 6676

```
Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
Simp[Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Chi}(bx) \sinh(bx)}{b} - \int \frac{\cosh(bx) \sinh(bx)}{bx} dx \\
&= \frac{\text{Chi}(bx) \sinh(bx)}{b} - \int \frac{\cosh(bx) \sinh(bx)}{x} dx \\
&= \frac{\text{Chi}(bx) \sinh(bx)}{b} - \int \frac{\sinh(2bx)}{2x} dx \\
&= \frac{\text{Chi}(bx) \sinh(bx)}{b} - \frac{\int \frac{\sinh(2bx)}{x} dx}{2b} \\
&= \frac{\text{Chi}(bx) \sinh(bx)}{b} - \frac{\text{Shi}(2bx)}{2b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \cosh(bx) \text{Chi}(bx) dx = \frac{\text{Chi}(bx) \sinh(bx)}{b} - \frac{\text{Shi}(2bx)}{2b}$$

```
[In] Integrate[Cosh[b*x]*CoshIntegral[b*x], x]
```

```
[Out] (CoshIntegral[b*x]*Sinh[b*x])/b - SinhIntegral[2*b*x]/(2*b)
```

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{\text{Chi}(bx) \sinh(bx) - \frac{\text{Shi}(2bx)}{2}}{b}$	22
default	$\frac{\text{Chi}(bx) \sinh(bx) - \frac{\text{Shi}(2bx)}{2}}{b}$	22

[In] `int(Chi(b*x)*cosh(b*x),x,method=_RETURNVERBOSE)`

[Out] `1/b*(Chi(b*x)*sinh(b*x)-1/2*Shi(2*b*x))`

Fricas [F]

$$\int \cosh(bx)\text{Chi}(bx) dx = \int \text{Chi}(bx) \cosh(bx) dx$$

[In] `integrate(Chi(b*x)*cosh(b*x),x, algorithm="fricas")`

[Out] `integral(cosh(b*x)*cosh_integral(b*x), x)`

Sympy [F]

$$\int \cosh(bx)\text{Chi}(bx) dx = \int \cosh(bx) \text{Chi}(bx) dx$$

[In] `integrate(Chi(b*x)*cosh(b*x),x)`

[Out] `Integral(cosh(b*x)*Chi(b*x), x)`

Maxima [F]

$$\int \cosh(bx)\text{Chi}(bx) dx = \int \text{Chi}(bx) \cosh(bx) dx$$

[In] `integrate(Chi(b*x)*cosh(b*x),x, algorithm="maxima")`

[Out] `integrate(Chi(b*x)*cosh(b*x), x)`

Giac [F]

$$\int \cosh(bx)\operatorname{Chi}(bx) dx = \int \operatorname{Chi}(bx) \cosh(bx) dx$$

[In] integrate(Chi(b*x)*cosh(b*x),x, algorithm="giac")

[Out] integrate(Chi(b*x)*cosh(b*x), x)

Mupad [F(-1)]

Timed out.

$$\int \cosh(bx)\operatorname{Chi}(bx) dx = \int \operatorname{coshint}(bx) \cosh(bx) dx$$

[In] int(coshint(b*x)*cosh(b*x),x)

[Out] int(coshint(b*x)*cosh(b*x), x)

3.111 $\int x \cosh(bx) \mathbf{Chi}(bx) dx$

Optimal result	543
Rubi [A] (verified)	543
Mathematica [A] (verified)	545
Maple [A] (verified)	545
Fricas [F]	546
Sympy [F]	546
Maxima [F]	546
Giac [F]	546
Mupad [F(-1)]	547

Optimal result

Integrand size = 10, antiderivative size = 61

$$\int x \cosh(bx) \mathbf{Chi}(bx) dx = -\frac{\cosh(bx) \mathbf{Chi}(bx)}{b^2} + \frac{\mathbf{Chi}(2bx)}{2b^2} + \frac{\log(x)}{2b^2} + \frac{x \mathbf{Chi}(bx) \sinh(bx)}{b} - \frac{\sinh^2(bx)}{2b^2}$$

[Out] $1/2*\mathbf{Chi}(2*b*x)/b^2 - \mathbf{Chi}(b*x)*\cosh(b*x)/b^2 + 1/2*\ln(x)/b^2 + x*\mathbf{Chi}(b*x)*\sinh(b*x)/b - 1/2*\sinh(b*x)^2/b^2$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {6678, 12, 2644, 30, 6682, 3393, 3382}

$$\int x \cosh(bx) \mathbf{Chi}(bx) dx = \frac{\mathbf{Chi}(2bx)}{2b^2} - \frac{\mathbf{Chi}(bx) \cosh(bx)}{b^2} + \frac{\log(x)}{2b^2} - \frac{\sinh^2(bx)}{2b^2} + \frac{x \mathbf{Chi}(bx) \sinh(bx)}{b}$$

[In] `Int[x*Cosh[b*x]*CoshIntegral[b*x],x]`

[Out] $-((\text{Cosh}[b*x]*\text{CoshIntegral}[b*x])/b^2) + \text{CoshIntegral}[2*b*x]/(2*b^2) + \text{Log}[x]/(2*b^2) + (x*\text{CoshIntegral}[b*x]*\text{Sinh}[b*x])/b - \text{Sinh}[b*x]^2/(2*b^2)$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 30

```
Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]
```

Rule 2644

```
Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)]^(m_), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])
```

Rule 3382

```
Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3393

```
Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 6678

```
Int[Cosh[(a_) + (b_)*(x_)]*CoshIntegral[(c_) + (d_)*(x_)]*((e_) + (f_)*(x_))^(m_), x_Symbol] := Simp[(e + f*x)^m*Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*CoshIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6682

```
Int[CoshIntegral[(c_) + (d_)*(x_)]*Sinh[(a_) + (b_)*(x_)], x_Symbol] := Simp[Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{x\text{Chi}(bx) \sinh(bx)}{b} - \frac{\int \text{Chi}(bx) \sinh(bx) dx}{b} - \int \frac{\cosh(bx) \sinh(bx)}{b} dx \\ &= -\frac{\cosh(bx)\text{Chi}(bx)}{b^2} + \frac{x\text{Chi}(bx) \sinh(bx)}{b} + \frac{\int \frac{\cosh^2(bx)}{bx} dx}{b} - \frac{\int \cosh(bx) \sinh(bx) dx}{b} \\ &= -\frac{\cosh(bx)\text{Chi}(bx)}{b^2} + \frac{x\text{Chi}(bx) \sinh(bx)}{b} + \frac{\int \frac{\cosh^2(bx)}{x} dx}{b^2} + \frac{\text{Subst}(\int x dx, x, i \sinh(bx))}{b^2} \end{aligned}$$

$$\begin{aligned}
&= -\frac{\cosh(bx)\text{Chi}(bx)}{b^2} + \frac{x\text{Chi}(bx)\sinh(bx)}{b} - \frac{\sinh^2(bx)}{2b^2} + \frac{\int\left(\frac{1}{2x} + \frac{\cosh(2bx)}{2x}\right)dx}{b^2} \\
&= -\frac{\cosh(bx)\text{Chi}(bx)}{b^2} + \frac{\log(x)}{2b^2} + \frac{x\text{Chi}(bx)\sinh(bx)}{b} - \frac{\sinh^2(bx)}{2b^2} + \frac{\int\frac{\cosh(2bx)}{x}dx}{2b^2} \\
&= -\frac{\cosh(bx)\text{Chi}(bx)}{b^2} + \frac{\text{Chi}(2bx)}{2b^2} + \frac{\log(x)}{2b^2} + \frac{x\text{Chi}(bx)\sinh(bx)}{b} - \frac{\sinh^2(bx)}{2b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.75

$$\begin{aligned}
&\int x \cosh(bx)\text{Chi}(bx) dx \\
&= \frac{-\cosh(2bx) + 2\text{Chi}(2bx) + 2\log(x) + 4\text{Chi}(bx)(-\cosh(bx) + bx\sinh(bx))}{4b^2}
\end{aligned}$$

[In] Integrate[x*Cosh[b*x]*CoshIntegral[b*x],x]

[Out] (-Cosh[2*b*x] + 2*CoshIntegral[2*b*x] + 2*Log[x] + 4*CoshIntegral[b*x]*(-Cosh[b*x] + b*x*Sinh[b*x]))/(4*b^2)

Maple [A] (verified)

Time = 0.71 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.75

method	result	size
derivativedivides	$\frac{\text{Chi}(bx)(bx\sinh(bx) - \cosh(bx)) - \frac{\cosh(bx)^2}{2} + \frac{\ln(bx)}{2} + \frac{\text{Chi}(2bx)}{2}}{b^2}$	46
default	$\frac{\text{Chi}(bx)(bx\sinh(bx) - \cosh(bx)) - \frac{\cosh(bx)^2}{2} + \frac{\ln(bx)}{2} + \frac{\text{Chi}(2bx)}{2}}{b^2}$	46

[In] int(x*Chi(b*x)*cosh(b*x),x,method=_RETURNVERBOSE)

[Out] 1/b^2*(Chi(b*x)*(b*x*sinh(b*x)-cosh(b*x))-1/2*cosh(b*x)^2+1/2*ln(b*x)+1/2*Chi(2*b*x))

Fricas [F]

$$\int x \cosh(bx) \operatorname{Chi}(bx) dx = \int x \operatorname{Chi}(bx) \cosh(bx) dx$$

[In] `integrate(x*Chi(b*x)*cosh(b*x),x, algorithm="fricas")`

[Out] `integral(x*cosh(b*x)*cosh_integral(b*x), x)`

Sympy [F]

$$\int x \cosh(bx) \operatorname{Chi}(bx) dx = \int x \cosh(bx) \operatorname{Chi}(bx) dx$$

[In] `integrate(x*Chi(b*x)*cosh(b*x),x)`

[Out] `Integral(x*cosh(b*x)*Chi(b*x), x)`

Maxima [F]

$$\int x \cosh(bx) \operatorname{Chi}(bx) dx = \int x \operatorname{Chi}(bx) \cosh(bx) dx$$

[In] `integrate(x*Chi(b*x)*cosh(b*x),x, algorithm="maxima")`

[Out] `integrate(x*Chi(b*x)*cosh(b*x), x)`

Giac [F]

$$\int x \cosh(bx) \operatorname{Chi}(bx) dx = \int x \operatorname{Chi}(bx) \cosh(bx) dx$$

[In] `integrate(x*Chi(b*x)*cosh(b*x),x, algorithm="giac")`

[Out] `integrate(x*Chi(b*x)*cosh(b*x), x)`

Mupad [F(-1)]

Timed out.

$$\int x \cosh(bx) \operatorname{Chi}(bx) dx = \int x \operatorname{coshint}(bx) \cosh(bx) dx$$

```
[In] int(x*coshint(b*x)*cosh(b*x),x)
```

```
[Out] int(x*coshint(b*x)*cosh(b*x), x)
```

3.112 $\int x^2 \cosh(bx) \mathbf{Chi}(bx) dx$

Optimal result	548
Rubi [A] (verified)	548
Mathematica [A] (verified)	551
Maple [A] (verified)	551
Fricas [F]	551
Sympy [F]	552
Maxima [F]	552
Giac [F]	552
Mupad [F(-1)]	552

Optimal result

Integrand size = 12, antiderivative size = 90

$$\int x^2 \cosh(bx) \mathbf{Chi}(bx) dx = \frac{3x}{4b^2} - \frac{2x \cosh(bx) \mathbf{Chi}(bx)}{b^2} + \frac{5 \cosh(bx) \sinh(bx)}{4b^3} + \frac{2 \mathbf{Chi}(bx) \sinh(bx)}{b^3} + \frac{x^2 \mathbf{Chi}(bx) \sinh(bx)}{b} - \frac{x \sinh^2(bx)}{2b^2} - \frac{\mathbf{Shi}(2bx)}{b^3}$$

[Out] $\frac{3}{4}x/b^2 - 2x*\mathbf{Chi}(b*x)*\cosh(b*x)/b^2 - \mathbf{Shi}(2*b*x)/b^3 + 2*\mathbf{Chi}(b*x)*\sinh(b*x)/b^3 + x^2*\mathbf{Chi}(b*x)*\sinh(b*x)/b + 5/4*\cosh(b*x)*\sinh(b*x)/b^3 - 1/2*x*\sinh(b*x)^2/b^2$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6678, 12, 5480, 2715, 8, 6684, 6676, 5556, 3379}

$$\int x^2 \cosh(bx) \mathbf{Chi}(bx) dx = \frac{2 \mathbf{Chi}(bx) \sinh(bx)}{b^3} - \frac{\mathbf{Shi}(2bx)}{b^3} + \frac{5 \sinh(bx) \cosh(bx)}{4b^3} - \frac{2x \mathbf{Chi}(bx) \cosh(bx)}{b^2} + \frac{3x}{4b^2} - \frac{x \sinh^2(bx)}{2b^2} + \frac{x^2 \mathbf{Chi}(bx) \sinh(bx)}{b}$$

[In] $\text{Int}[x^2*\text{Cosh}[b*x]*\text{CoshIntegral}[b*x],x]$

[Out] $(3*x)/(4*b^2) - (2*x*\text{Cosh}[b*x]*\text{CoshIntegral}[b*x])/b^2 + (5*\text{Cosh}[b*x]*\text{Sinh}[b*x])/(4*b^3) + (2*\text{CoshIntegral}[b*x]*\text{Sinh}[b*x])/b^3 + (x^2*\text{CoshIntegral}[b*x]*\text{Sinh}[b*x])/b - (x*\text{Sinh}[b*x]^2)/(2*b^2) - \text{SinhIntegral}[2*b*x]/b^3$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] \text{ /; FreeQ}[a, x]$

Rule 12

$\text{Int}[(a_)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] \text{ /; FreeQ}[a, x] \ \&\& \ !\text{Match}$
 $\text{Q}[u, (b_)*(v_)] \text{ /; FreeQ}[b, x]$

Rule 2715

$\text{Int}[(b_)*\sin[(c_.) + (d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]$
 $*((b*\text{Sin}[c + d*x])^{(n-1)})/(d*n), x] + \text{Dist}[b^2*((n-1)/n), \text{Int}[(b*\text{Sin}[$
 $c + d*x])^{(n-2)}, x], x] \text{ /; FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2$
 $*n]$

Rule 3379

$\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])*(f_)*(x_)]/((c_.) + (d_)*(x_)), x_Symbol]$
 $\rightarrow \text{Simp}[I*(\text{SinhIntegral}[c*f*(fz/d) + f*fz*x]/d), x] \text{ /; FreeQ}[\{c, d, e, f,$
 $fz\}, x] \ \&\& \ \text{EqQ}[d*e - c*f*fz*I, 0]$

Rule 5480

$\text{Int}[\text{Cosh}[(a_.) + (b_)*(x_)]^{(n_)}*(x_)]^{(m_)}*\text{Sinh}[(a_.) + (b_)*(x_)]^{(n_)}$
 $]^{(p_)}, x_Symbol] \rightarrow \text{Simp}[x^{(m-n+1)}*(\text{Sinh}[a + b*x^n]^{(p+1)})/(b*n*(p+1))$
 $], x] - \text{Dist}[(m-n+1)/(b*n*(p+1)), \text{Int}[x^{(m-n)}*\text{Sinh}[a + b*x^n]^{(p+1)}$
 $], x], x] \text{ /; FreeQ}[\{a, b, p\}, x] \ \&\& \ \text{LtQ}[0, n, m+1] \ \&\& \ \text{NeQ}[p, -1]$

Rule 5556

$\text{Int}[\text{Cosh}[(a_.) + (b_)*(x_)]^{(p_)}*((c_.) + (d_)*(x_)]^{(m_)}*\text{Sinh}[(a_.) +$
 $(b_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(c + d*x)^m, \text{Sinh}[a +$
 $b*x]^{n*}\text{Cosh}[a + b*x]^{p}, x], x] \text{ /; FreeQ}[\{a, b, c, d, m\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&$
 $\ \& \ \text{IGtQ}[p, 0]$

Rule 6676

$\text{Int}[\text{Cosh}[(a_.) + (b_)*(x_)]*\text{CoshIntegral}[(c_.) + (d_)*(x_)], x_Symbol] \rightarrow$
 $\text{Simp}[\text{Sinh}[a + b*x]*(\text{CoshIntegral}[c + d*x]/b), x] - \text{Dist}[d/b, \text{Int}[\text{Sinh}[a +$
 $b*x]*(\text{Cosh}[c + d*x]/(c + d*x)), x], x] \text{ /; FreeQ}[\{a, b, c, d\}, x]$

Rule 6678

$\text{Int}[\text{Cosh}[(a_.) + (b_)*(x_)]*\text{CoshIntegral}[(c_.) + (d_)*(x_)]*((e_.) + (f_.$
 $)*(x_)]^{(m_)}, x_Symbol] \rightarrow \text{Simp}[(e + f*x)^m*\text{Sinh}[a + b*x]*(\text{CoshIntegral}[c$
 $+ d*x]/b), x] + (-\text{Dist}[d/b, \text{Int}[(e + f*x)^m*\text{Sinh}[a + b*x]*(\text{Cosh}[c + d*x]/(c$

+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*CoshIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rule 6684

Int[CoshIntegral[(c_.) + (d_.)*(x_.)]*((e_.) + (f_.)*(x_.))^(m_.)*Sinh[(a_.) + (b_.)*(x_.)], x_Symbol] :> Simp[(e + f*x)^m*Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*CoshIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x^2 \text{Chi}(bx) \sinh(bx)}{b} - \frac{2 \int x \text{Chi}(bx) \sinh(bx) dx}{b} - \int \frac{x \cosh(bx) \sinh(bx)}{b} dx \\
 &= -\frac{2x \cosh(bx) \text{Chi}(bx)}{b^2} + \frac{x^2 \text{Chi}(bx) \sinh(bx)}{b} + \frac{2 \int \cosh(bx) \text{Chi}(bx) dx}{b^2} \\
 &\quad - \frac{\int x \cosh(bx) \sinh(bx) dx}{b} + \frac{2 \int \frac{\cosh^2(bx)}{b} dx}{b} \\
 &= -\frac{2x \cosh(bx) \text{Chi}(bx)}{b^2} + \frac{2 \text{Chi}(bx) \sinh(bx)}{b^3} + \frac{x^2 \text{Chi}(bx) \sinh(bx)}{b} \\
 &\quad - \frac{x \sinh^2(bx)}{2b^2} + \frac{\int \sinh^2(bx) dx}{2b^2} + \frac{2 \int \cosh^2(bx) dx}{b^2} - \frac{2 \int \frac{\cosh(bx) \sinh(bx)}{bx} dx}{b^2} \\
 &= -\frac{2x \cosh(bx) \text{Chi}(bx)}{b^2} + \frac{5 \cosh(bx) \sinh(bx)}{4b^3} + \frac{2 \text{Chi}(bx) \sinh(bx)}{b^3} \\
 &\quad + \frac{x^2 \text{Chi}(bx) \sinh(bx)}{b} - \frac{x \sinh^2(bx)}{2b^2} - \frac{2 \int \frac{\cosh(bx) \sinh(bx)}{x} dx}{b^3} - \frac{\int 1 dx}{4b^2} + \frac{\int 1 dx}{b^2} \\
 &= \frac{3x}{4b^2} - \frac{2x \cosh(bx) \text{Chi}(bx)}{b^2} + \frac{5 \cosh(bx) \sinh(bx)}{4b^3} + \frac{2 \text{Chi}(bx) \sinh(bx)}{b^3} \\
 &\quad + \frac{x^2 \text{Chi}(bx) \sinh(bx)}{b} - \frac{x \sinh^2(bx)}{2b^2} - \frac{2 \int \frac{\sinh(2bx)}{2x} dx}{b^3} \\
 &= \frac{3x}{4b^2} - \frac{2x \cosh(bx) \text{Chi}(bx)}{b^2} + \frac{5 \cosh(bx) \sinh(bx)}{4b^3} + \frac{2 \text{Chi}(bx) \sinh(bx)}{b^3} \\
 &\quad + \frac{x^2 \text{Chi}(bx) \sinh(bx)}{b} - \frac{x \sinh^2(bx)}{2b^2} - \frac{\int \frac{\sinh(2bx)}{x} dx}{b^3} \\
 &= \frac{3x}{4b^2} - \frac{2x \cosh(bx) \text{Chi}(bx)}{b^2} + \frac{5 \cosh(bx) \sinh(bx)}{4b^3} + \frac{2 \text{Chi}(bx) \sinh(bx)}{b^3} \\
 &\quad + \frac{x^2 \text{Chi}(bx) \sinh(bx)}{b} - \frac{x \sinh^2(bx)}{2b^2} - \frac{\text{Shi}(2bx)}{b^3}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.71

$$\int x^2 \cosh(bx) \operatorname{Chi}(bx) dx = \frac{8bx - 2bx \cosh(2bx) + 8\operatorname{Chi}(bx) (-2bx \cosh(bx) + (2 + b^2x^2) \sinh(bx)) + 5 \sinh(2bx) - 8\operatorname{Shi}(2bx)}{8b^3}$$

[In] Integrate[x^2*Cosh[b*x]*CoshIntegral[b*x],x]

[Out] (8*b*x - 2*b*x*Cosh[2*b*x] + 8*CoshIntegral[b*x]*(-2*b*x*Cosh[b*x] + (2 + b^2*x^2)*Sinh[b*x]) + 5*Sinh[2*b*x] - 8*SinhIntegral[2*b*x])/(8*b^3)

Maple [A] (verified)

Time = 0.92 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.76

method	result	size
derivativedivides	$\frac{\operatorname{Chi}(bx)(b^2x^2 \sinh(bx) - 2bx \cosh(bx) + 2 \sinh(bx)) - \frac{bx \cosh(bx)^2}{2} + \frac{5 \cosh(bx) \sinh(bx)}{4} + \frac{5bx}{4} - \operatorname{Shi}(2bx)}{b^3}$	68
default	$\frac{\operatorname{Chi}(bx)(b^2x^2 \sinh(bx) - 2bx \cosh(bx) + 2 \sinh(bx)) - \frac{bx \cosh(bx)^2}{2} + \frac{5 \cosh(bx) \sinh(bx)}{4} + \frac{5bx}{4} - \operatorname{Shi}(2bx)}{b^3}$	68

[In] int(x^2*Chi(b*x)*cosh(b*x),x,method=_RETURNVERBOSE)

[Out] 1/b^3*(Chi(b*x)*(b^2*x^2*sinh(b*x)-2*b*x*cosh(b*x)+2*sinh(b*x))-1/2*b*x*cosh(b*x)^2+5/4*cosh(b*x)*sinh(b*x)+5/4*b*x-Shi(2*b*x))

Fricas [F]

$$\int x^2 \cosh(bx) \operatorname{Chi}(bx) dx = \int x^2 \operatorname{Chi}(bx) \cosh(bx) dx$$

[In] integrate(x^2*Chi(b*x)*cosh(b*x),x, algorithm="fricas")

[Out] integral(x^2*cosh(b*x)*cosh_integral(b*x), x)

Sympy [F]

$$\int x^2 \cosh(bx) \operatorname{Chi}(bx) dx = \int x^2 \cosh(bx) \operatorname{Chi}(bx) dx$$

[In] integrate(x**2*Chi(b*x)*cosh(b*x), x)

[Out] Integral(x**2*cosh(b*x)*Chi(b*x), x)

Maxima [F]

$$\int x^2 \cosh(bx) \operatorname{Chi}(bx) dx = \int x^2 \operatorname{Chi}(bx) \cosh(bx) dx$$

[In] integrate(x^2*Chi(b*x)*cosh(b*x), x, algorithm="maxima")

[Out] integrate(x^2*Chi(b*x)*cosh(b*x), x)

Giac [F]

$$\int x^2 \cosh(bx) \operatorname{Chi}(bx) dx = \int x^2 \operatorname{Chi}(bx) \cosh(bx) dx$$

[In] integrate(x^2*Chi(b*x)*cosh(b*x), x, algorithm="giac")

[Out] integrate(x^2*Chi(b*x)*cosh(b*x), x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \cosh(bx) \operatorname{Chi}(bx) dx = \int x^2 \operatorname{coshint}(bx) \cosh(bx) dx$$

[In] int(x^2*coshint(b*x)*cosh(b*x), x)

[Out] int(x^2*coshint(b*x)*cosh(b*x), x)

3.113 $\int x^3 \cosh(bx) \mathbf{Chi}(bx) dx$

Optimal result	553
Rubi [A] (verified)	553
Mathematica [A] (verified)	556
Maple [A] (verified)	557
Fricas [F]	557
Sympy [F]	557
Maxima [F]	557
Giac [F]	558
Mupad [F(-1)]	558

Optimal result

Integrand size = 12, antiderivative size = 142

$$\int x^3 \cosh(bx) \mathbf{Chi}(bx) dx = \frac{x^2}{2b^2} - \frac{3 \cosh^2(bx)}{4b^4} - \frac{6 \cosh(bx) \mathbf{Chi}(bx)}{b^4} - \frac{3x^2 \cosh(bx) \mathbf{Chi}(bx)}{b^2} + \frac{3 \mathbf{Chi}(2bx)}{b^4} + \frac{3 \log(x)}{b^4} + \frac{2x \cosh(bx) \sinh(bx)}{b^3} + \frac{6x \mathbf{Chi}(bx) \sinh(bx)}{b^3} + \frac{x^3 \mathbf{Chi}(bx) \sinh(bx)}{b} - \frac{13 \sinh^2(bx)}{4b^4} - \frac{x^2 \sinh^2(bx)}{2b^2}$$

[Out] $\frac{1}{2}x^2/b^2 + 3\mathbf{Chi}(2bx)/b^4 - 6\mathbf{Chi}(bx)\cosh(bx)/b^4 - 3x^2\mathbf{Chi}(bx)\cosh(bx)/b^2 - 3/4\cosh(bx)^2/b^4 + 3\ln(x)/b^4 + 6xx\mathbf{Chi}(bx)\sinh(bx)/b^3 + x^3\mathbf{Chi}(bx)\sinh(bx)/b + 2xx\cosh(bx)\sinh(bx)/b^3 - 13/4\sinh(bx)^2/b^4 - 1/2x^2\sinh(bx)^2/b^2$

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.00, number of steps used = 18, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {6678, 12, 5480, 3391, 30, 6684, 2644, 6682, 3393, 3382}

$$\int x^3 \cosh(bx) \mathbf{Chi}(bx) dx = \frac{3\mathbf{Chi}(2bx)}{b^4} - \frac{6\mathbf{Chi}(bx) \cosh(bx)}{b^4} + \frac{3 \log(x)}{b^4} - \frac{13 \sinh^2(bx)}{4b^4} - \frac{3 \cosh^2(bx)}{4b^4} + \frac{6x \mathbf{Chi}(bx) \sinh(bx)}{b^3} + \frac{2x \sinh(bx) \cosh(bx)}{b^3} - \frac{3x^2 \mathbf{Chi}(bx) \cosh(bx)}{b^2} + \frac{x^2}{2b^2} - \frac{x^2 \sinh^2(bx)}{2b^2} + \frac{x^3 \mathbf{Chi}(bx) \sinh(bx)}{b}$$

[In] Int[x^3*Cosh[b*x]*CoshIntegral[b*x],x]

[Out] $x^2/(2*b^2) - (3*Cosh[b*x]^2)/(4*b^4) - (6*Cosh[b*x]*CoshIntegral[b*x])/b^4 - (3*x^2*Cosh[b*x]*CoshIntegral[b*x])/b^2 + (3*CoshIntegral[2*b*x])/b^4 + (3*Log[x])/b^4 + (2*x*Cosh[b*x]*Sinh[b*x])/b^3 + (6*x*CoshIntegral[b*x]*Sinh[b*x])/b^3 + (x^3*CoshIntegral[b*x]*Sinh[b*x])/b - (13*Sinh[b*x]^2)/(4*b^4) - (x^2*Sinh[b*x]^2)/(2*b^2)$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2644

Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)]^(m_), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rule 3382

Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3391

Int[((c_) + (d_)*(x_))*((b_)*sin[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[d*((b*Sin[e + f*x])^n/(f^2*n^2), x] + (Dist[b^2*((n - 1)/n), Int[(c + d*x)*(b*Sin[e + f*x])^(n - 2), x], x] - Simp[b*(c + d*x)*Cos[e + f*x]*(b*Sin[e + f*x])^(n - 1)/(f*n), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1]

Rule 3393

Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))

Rule 5480

Int[Cosh[(a_) + (b_)*(x_)]^(n_)*(x_)^(m_)*Sinh[(a_) + (b_)*(x_)]^(p_), x_Symbol] := Simp[x^(m - n + 1)*(Sinh[a + b*x^n]^(p + 1)/(b*n*(p + 1))

1))), x] - Dist[(m - n + 1)/(b*n*(p + 1)), Int[x^(m - n)*Sinh[a + b*x^n]^(p + 1), x], x] /; FreeQ[{a, b, p}, x] && LtQ[0, n, m + 1] && NeQ[p, -1]

Rule 6678

Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[(e + f*x)^m*Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*CoshIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rule 6682

Int[CoshIntegral[(c_.) + (d_.)*(x_)]*Sinh[(a_.) + (b_.)*(x_)], x_Symbol] :> Simp[Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]

Rule 6684

Int[CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)], x_Symbol] :> Simp[(e + f*x)^m*Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*CoshIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x^3 \text{Chi}(bx) \sinh(bx)}{b} - \frac{3 \int x^2 \text{Chi}(bx) \sinh(bx) dx}{b} - \int \frac{x^2 \cosh(bx) \sinh(bx)}{b} dx \\
 &= -\frac{3x^2 \cosh(bx) \text{Chi}(bx)}{b^2} + \frac{x^3 \text{Chi}(bx) \sinh(bx)}{b} + \frac{6 \int x \cosh(bx) \text{Chi}(bx) dx}{b^2} \\
 &\quad - \frac{\int x^2 \cosh(bx) \sinh(bx) dx}{b} + \frac{3 \int \frac{x \cosh^2(bx)}{b} dx}{b} \\
 &= -\frac{3x^2 \cosh(bx) \text{Chi}(bx)}{b^2} + \frac{6x \text{Chi}(bx) \sinh(bx)}{b^3} + \frac{x^3 \text{Chi}(bx) \sinh(bx)}{b} - \frac{x^2 \sinh^2(bx)}{2b^2} \\
 &\quad - \frac{6 \int \text{Chi}(bx) \sinh(bx) dx}{b^3} + \frac{\int x \sinh^2(bx) dx}{b^2} + \frac{3 \int x \cosh^2(bx) dx}{b^2} - \frac{6 \int \frac{\cosh(bx) \sinh(bx)}{b} dx}{b^2} \\
 &= -\frac{3 \cosh^2(bx)}{4b^4} - \frac{6 \cosh(bx) \text{Chi}(bx)}{b^4} - \frac{3x^2 \cosh(bx) \text{Chi}(bx)}{b^2} + \frac{2x \cosh(bx) \sinh(bx)}{b^3} \\
 &\quad + \frac{6x \text{Chi}(bx) \sinh(bx)}{b^3} + \frac{x^3 \text{Chi}(bx) \sinh(bx)}{b} - \frac{\sinh^2(bx)}{4b^4} - \frac{x^2 \sinh^2(bx)}{2b^2} \\
 &\quad + \frac{6 \int \frac{\cosh^2(bx)}{bx} dx}{b^3} - \frac{6 \int \cosh(bx) \sinh(bx) dx}{b^3} - \frac{\int x dx}{2b^2} + \frac{3 \int x dx}{2b^2}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{x^2}{2b^2} - \frac{3 \cosh^2(bx)}{4b^4} - \frac{6 \cosh(bx)\text{Chi}(bx)}{b^4} - \frac{3x^2 \cosh(bx)\text{Chi}(bx)}{b^2} \\
&\quad + \frac{2x \cosh(bx) \sinh(bx)}{b^3} + \frac{6x\text{Chi}(bx) \sinh(bx)}{b^3} + \frac{x^3\text{Chi}(bx) \sinh(bx)}{b} \\
&\quad - \frac{\sinh^2(bx)}{4b^4} - \frac{x^2 \sinh^2(bx)}{2b^2} + \frac{6 \int \frac{\cosh^2(bx)}{x} dx}{b^4} + \frac{6\text{Subst}(\int x dx, x, i \sinh(bx))}{b^4} \\
&= \frac{x^2}{2b^2} - \frac{3 \cosh^2(bx)}{4b^4} - \frac{6 \cosh(bx)\text{Chi}(bx)}{b^4} - \frac{3x^2 \cosh(bx)\text{Chi}(bx)}{b^2} \\
&\quad + \frac{2x \cosh(bx) \sinh(bx)}{b^3} + \frac{6x\text{Chi}(bx) \sinh(bx)}{b^3} + \frac{x^3\text{Chi}(bx) \sinh(bx)}{b} \\
&\quad - \frac{13 \sinh^2(bx)}{4b^4} - \frac{x^2 \sinh^2(bx)}{2b^2} + \frac{6 \int \left(\frac{1}{2x} + \frac{\cosh(2bx)}{2x} \right) dx}{b^4} \\
&= \frac{x^2}{2b^2} - \frac{3 \cosh^2(bx)}{4b^4} - \frac{6 \cosh(bx)\text{Chi}(bx)}{b^4} - \frac{3x^2 \cosh(bx)\text{Chi}(bx)}{b^2} \\
&\quad + \frac{3 \log(x)}{b^4} + \frac{2x \cosh(bx) \sinh(bx)}{b^3} + \frac{6x\text{Chi}(bx) \sinh(bx)}{b^3} \\
&\quad + \frac{x^3\text{Chi}(bx) \sinh(bx)}{b} - \frac{13 \sinh^2(bx)}{4b^4} - \frac{x^2 \sinh^2(bx)}{2b^2} + \frac{3 \int \frac{\cosh(2bx)}{x} dx}{b^4} \\
&= \frac{x^2}{2b^2} - \frac{3 \cosh^2(bx)}{4b^4} - \frac{6 \cosh(bx)\text{Chi}(bx)}{b^4} - \frac{3x^2 \cosh(bx)\text{Chi}(bx)}{b^2} \\
&\quad + \frac{3\text{Chi}(2bx)}{b^4} + \frac{3 \log(x)}{b^4} + \frac{2x \cosh(bx) \sinh(bx)}{b^3} + \frac{6x\text{Chi}(bx) \sinh(bx)}{b^3} \\
&\quad + \frac{x^3\text{Chi}(bx) \sinh(bx)}{b} - \frac{13 \sinh^2(bx)}{4b^4} - \frac{x^2 \sinh^2(bx)}{2b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.66

$$\int x^3 \cosh(bx)\text{Chi}(bx) dx = \frac{3b^2x^2 - 8 \cosh(2bx) - b^2x^2 \cosh(2bx) + 12\text{Chi}(2bx) + 12 \log(x) + 4\text{Chi}(bx) (-3(2 + b^2x^2) \cosh(bx) + bx(6 + b^2x^2)) + 4b^2x \sinh(bx)}{4b^4}$$

[In] Integrate[x^3*Cosh[b*x]*CoshIntegral[b*x],x]

[Out] (3*b^2*x^2 - 8*Cosh[2*b*x] - b^2*x^2*Cosh[2*b*x] + 12*CoshIntegral[2*b*x] + 12*Log[x] + 4*CoshIntegral[b*x]*(-3*(2 + b^2*x^2)*Cosh[b*x] + b*x*(6 + b^2*x^2)*Sinh[b*x]) + 4*b*x*Sinh[2*b*x])/(4*b^4)

Maple [A] (verified)

Time = 1.25 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.73

method	result
derivativedivides	$\frac{\text{Chi}(bx)(b^3x^3 \sinh(bx) - 3b^2x^2 \cosh(bx) + 6bx \sinh(bx) - 6 \cosh(bx)) - \frac{b^2x^2 \cosh(bx)^2}{b^4} + 2bx \cosh(bx) \sinh(bx) + b^2x^2 - 4 \cosh(bx)}{b^4}$
default	$\frac{\text{Chi}(bx)(b^3x^3 \sinh(bx) - 3b^2x^2 \cosh(bx) + 6bx \sinh(bx) - 6 \cosh(bx)) - \frac{b^2x^2 \cosh(bx)^2}{b^4} + 2bx \cosh(bx) \sinh(bx) + b^2x^2 - 4 \cosh(bx)}{b^4}$

[In] `int(x^3*Chi(b*x)*cosh(b*x),x,method=_RETURNVERBOSE)`[Out] `1/b^4*(Chi(b*x)*(b^3*x^3*sinh(b*x)-3*b^2*x^2*cosh(b*x)+6*b*x*sinh(b*x)-6*cosh(b*x))-1/2*b^2*x^2*cosh(b*x)^2+2*b*x*cosh(b*x)*sinh(b*x)+b^2*x^2-4*cosh(b*x)^2+3*ln(b*x)+3*Chi(2*b*x))`**Fricas [F]**

$$\int x^3 \cosh(bx) \text{Chi}(bx) dx = \int x^3 \text{Chi}(bx) \cosh(bx) dx$$

[In] `integrate(x^3*Chi(b*x)*cosh(b*x),x, algorithm="fricas")`[Out] `integral(x^3*cosh(b*x)*cosh_integral(b*x), x)`**Sympy [F]**

$$\int x^3 \cosh(bx) \text{Chi}(bx) dx = \int x^3 \cosh(bx) \text{Chi}(bx) dx$$

[In] `integrate(x**3*Chi(b*x)*cosh(b*x),x)`[Out] `Integral(x**3*cosh(b*x)*Chi(b*x), x)`**Maxima [F]**

$$\int x^3 \cosh(bx) \text{Chi}(bx) dx = \int x^3 \text{Chi}(bx) \cosh(bx) dx$$

[In] `integrate(x^3*Chi(b*x)*cosh(b*x),x, algorithm="maxima")`[Out] `integrate(x^3*Chi(b*x)*cosh(b*x), x)`

Giac [F]

$$\int x^3 \cosh(bx) \operatorname{Chi}(bx) dx = \int x^3 \operatorname{Chi}(bx) \cosh(bx) dx$$

[In] integrate(x^3*Chi(b*x)*cosh(b*x),x, algorithm="giac")

[Out] integrate(x^3*Chi(b*x)*cosh(b*x), x)

Mupad [F(-1)]

Timed out.

$$\int x^3 \cosh(bx) \operatorname{Chi}(bx) dx = \int x^3 \operatorname{coshint}(bx) \cosh(bx) dx$$

[In] int(x^3*coshint(b*x)*cosh(b*x),x)

[Out] int(x^3*coshint(b*x)*cosh(b*x), x)

3.114 $\int \frac{\mathbf{Chi}(bx) \sinh(bx)}{x^3} dx$

Optimal result	559
Rubi [N/A]	559
Mathematica [N/A]	560
Maple [N/A] (verified)	561
Fricas [N/A]	561
Sympy [N/A]	561
Maxima [N/A]	561
Giac [N/A]	562
Mupad [N/A]	562

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\mathbf{Chi}(bx) \sinh(bx)}{x^3} dx = -\frac{b \cosh^2(bx)}{2x} - \frac{b \cosh(2bx)}{4x} - \frac{b \cosh(bx) \mathbf{Chi}(bx)}{2x} - \frac{\mathbf{Chi}(bx) \sinh(bx)}{2x^2} - \frac{\sinh(2bx)}{8x^2} + b^2 \text{Shi}(2bx) + \frac{1}{2} b^2 \text{Int}\left(\frac{\mathbf{Chi}(bx) \sinh(bx)}{x}, x\right)$$

[Out] 1/2*b^2*CannotIntegrate(Chi(b*x)*sinh(b*x)/x,x)-1/2*b*Chi(b*x)*cosh(b*x)/x-1/2*b*cosh(b*x)^2/x-1/4*b*cosh(2*b*x)/x+b^2*Shi(2*b*x)-1/2*Chi(b*x)*sinh(b*x)/x^2-1/8*sinh(2*b*x)/x^2

Rubi [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\mathbf{Chi}(bx) \sinh(bx)}{x^3} dx = \int \frac{\mathbf{Chi}(bx) \sinh(bx)}{x^3} dx$$

[In] Int[(CoshIntegral[b*x]*Sinh[b*x])/x^3,x]

[Out] -1/2*(b*Cosh[b*x]^2)/x - (b*Cosh[2*b*x])/(4*x) - (b*Cosh[b*x]*CoshIntegral[b*x])/(2*x) - (CoshIntegral[b*x]*Sinh[b*x])/(2*x^2) - Sinh[2*b*x]/(8*x^2) + b^2*SinhIntegral[2*b*x] + (b^2*Defer[Int]((CoshIntegral[b*x]*Sinh[b*x])/x,x))/2

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Chi}(bx) \sinh(bx)}{2x^2} + \frac{1}{2}b \int \frac{\cosh(bx)\text{Chi}(bx)}{x^2} dx + \frac{1}{2}b \int \frac{\cosh(bx) \sinh(bx)}{bx^3} dx \\
&= -\frac{b \cosh(bx)\text{Chi}(bx)}{2x} - \frac{\text{Chi}(bx) \sinh(bx)}{2x^2} + \frac{1}{2} \int \frac{\cosh(bx) \sinh(bx)}{x^3} dx \\
&\quad + \frac{1}{2}b^2 \int \frac{\cosh^2(bx)}{bx^2} dx + \frac{1}{2}b^2 \int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx \\
&= -\frac{b \cosh(bx)\text{Chi}(bx)}{2x} - \frac{\text{Chi}(bx) \sinh(bx)}{2x^2} + \frac{1}{2} \int \frac{\sinh(2bx)}{2x^3} dx \\
&\quad + \frac{1}{2}b \int \frac{\cosh^2(bx)}{x^2} dx + \frac{1}{2}b^2 \int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx \\
&= -\frac{b \cosh^2(bx)}{2x} - \frac{b \cosh(bx)\text{Chi}(bx)}{2x} - \frac{\text{Chi}(bx) \sinh(bx)}{2x^2} + \frac{1}{4} \int \frac{\sinh(2bx)}{x^3} dx \\
&\quad + (ib^2) \int -\frac{i \sinh(2bx)}{2x} dx + \frac{1}{2}b^2 \int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx \\
&= -\frac{b \cosh^2(bx)}{2x} - \frac{b \cosh(bx)\text{Chi}(bx)}{2x} - \frac{\text{Chi}(bx) \sinh(bx)}{2x^2} - \frac{\sinh(2bx)}{8x^2} \\
&\quad + \frac{1}{4}b \int \frac{\cosh(2bx)}{x^2} dx + \frac{1}{2}b^2 \int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx + \frac{1}{2}b^2 \int \frac{\sinh(2bx)}{x} dx \\
&= -\frac{b \cosh^2(bx)}{2x} - \frac{b \cosh(2bx)}{4x} - \frac{b \cosh(bx)\text{Chi}(bx)}{2x} - \frac{\text{Chi}(bx) \sinh(bx)}{2x^2} \\
&\quad - \frac{\sinh(2bx)}{8x^2} + \frac{1}{2}b^2 \text{Shi}(2bx) + \frac{1}{2}b^2 \int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx + \frac{1}{2}b^2 \int \frac{\sinh(2bx)}{x} dx \\
&= -\frac{b \cosh^2(bx)}{2x} - \frac{b \cosh(2bx)}{4x} - \frac{b \cosh(bx)\text{Chi}(bx)}{2x} - \frac{\text{Chi}(bx) \sinh(bx)}{2x^2} \\
&\quad - \frac{\sinh(2bx)}{8x^2} + b^2 \text{Shi}(2bx) + \frac{1}{2}b^2 \int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx
\end{aligned}$$

Mathematica [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^3} dx = \int \frac{\text{Chi}(bx) \sinh(bx)}{x^3} dx$$

[In] Integrate[(CoshIntegral[b*x]*Sinh[b*x])/x^3,x]

[Out] Integrate[(CoshIntegral[b*x]*Sinh[b*x])/x^3, x]

Maple [N/A] (verified)

Not integrable

Time = 0.21 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^3} dx$$

[In] int(Chi(b*x)*sinh(b*x)/x^3,x)

[Out] int(Chi(b*x)*sinh(b*x)/x^3,x)

Fricas [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^3} dx = \int \frac{\text{Chi}(bx) \sinh(bx)}{x^3} dx$$

[In] integrate(Chi(b*x)*sinh(b*x)/x^3,x, algorithm="fricas")

[Out] integral(cosh_integral(b*x)*sinh(b*x)/x^3, x)

Sympy [N/A]

Not integrable

Time = 3.64 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^3} dx = \int \frac{\sinh(bx) \text{Chi}(bx)}{x^3} dx$$

[In] integrate(Chi(b*x)*sinh(b*x)/x**3,x)

[Out] Integral(sinh(b*x)*Chi(b*x)/x**3, x)

Maxima [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^3} dx = \int \frac{\text{Chi}(bx) \sinh(bx)}{x^3} dx$$

[In] integrate(Chi(b*x)*sinh(b*x)/x^3,x, algorithm="maxima")

[Out] integrate(Chi(b*x)*sinh(b*x)/x^3, x)

Giac [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^3} dx = \int \frac{\text{Chi}(bx) \sinh(bx)}{x^3} dx$$

`[In] integrate(Chi(b*x)*sinh(b*x)/x^3,x, algorithm="giac")``[Out] integrate(Chi(b*x)*sinh(b*x)/x^3, x)`**Mupad [N/A]**

Not integrable

Time = 4.87 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^3} dx = \int \frac{\text{coshint}(bx) \sinh(bx)}{x^3} dx$$

`[In] int((coshint(b*x)*sinh(b*x))/x^3,x)``[Out] int((coshint(b*x)*sinh(b*x))/x^3, x)`

3.115 $\int \frac{\text{Chi}(bx) \sinh(bx)}{x^2} dx$

Optimal result	563
Rubi [A] (verified)	563
Mathematica [A] (verified)	565
Maple [F]	565
Fricas [F]	565
Sympy [F]	565
Maxima [F]	566
Giac [F]	566
Mupad [F(-1)]	566

Optimal result

Integrand size = 12, antiderivative size = 44

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^2} dx = \frac{1}{2}b\text{Chi}(bx)^2 + b\text{Chi}(2bx) - \frac{\text{Chi}(bx) \sinh(bx)}{x} - \frac{\sinh(2bx)}{2x}$$

[Out] $1/2*b*\text{Chi}(b*x)^2+b*\text{Chi}(2*b*x)-\text{Chi}(b*x)*\sinh(b*x)/x-1/2*\sinh(2*b*x)/x$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {6686, 6818, 12, 5556, 3378, 3382}

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^2} dx = \frac{1}{2}b\text{Chi}(bx)^2 + b\text{Chi}(2bx) - \frac{\text{Chi}(bx) \sinh(bx)}{x} - \frac{\sinh(2bx)}{2x}$$

[In] $\text{Int}[(\text{CoshIntegral}[b*x]*\text{Sinh}[b*x])/x^2,x]$

[Out] $(b*\text{CoshIntegral}[b*x]^2)/2 + b*\text{CoshIntegral}[2*b*x] - (\text{CoshIntegral}[b*x]*\text{Sinh}[b*x])/x - \text{Sinh}[2*b*x]/(2*x)$

Rule 12

$\text{Int}[(a_*)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /;$ FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 3378

$\text{Int}[(c_*) + (d_*)*(x_)^m*\sin[(e_*) + (f_*)*(x_)], x_Symbol] \rightarrow \text{Simp}[(c + d*x)^{m+1}*(\text{Sin}[e + f*x]/(d*(m+1))), x] - \text{Dist}[f/(d*(m+1)), \text{Int}[(c + d*x)^{m+1}*\text{Cos}[e + f*x], x], x] /;$ FreeQ[{c, d, e, f}, x] && LtQ[m, -1

]

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol]
:> Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x]
&& EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] :> Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

Rule 6686

```
Int[CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)], x_Symbol] :> Simp[(e + f*x)^(m + 1)*Sinh[a + b*x]*(CoshIntegral[c + d*x]/(f*(m + 1))), x] + (-Dist[b/(f*(m + 1)), Int[(e + f*x)^(m + 1)*Cosh[a + b*x]*CoshIntegral[c + d*x], x], x] - Dist[d/(f*(m + 1)), Int[(e + f*x)^(m + 1)*Sinh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[m, -1]
```

Rule 6818

```
Int[(u_.)*(y_)^(m_.), x_Symbol] :> With[{q = DerivativeDivides[y, u, x]}, Simp[q*(y^(m + 1)/(m + 1)), x] /; !FalseQ[q]] /; FreeQ[m, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Chi}(bx) \sinh(bx)}{x} + b \int \frac{\cosh(bx) \text{Chi}(bx)}{x} dx + b \int \frac{\cosh(bx) \sinh(bx)}{bx^2} dx \\
&= \frac{1}{2} b \text{Chi}(bx)^2 - \frac{\text{Chi}(bx) \sinh(bx)}{x} + \int \frac{\cosh(bx) \sinh(bx)}{x^2} dx \\
&= \frac{1}{2} b \text{Chi}(bx)^2 - \frac{\text{Chi}(bx) \sinh(bx)}{x} + \int \frac{\sinh(2bx)}{2x^2} dx \\
&= \frac{1}{2} b \text{Chi}(bx)^2 - \frac{\text{Chi}(bx) \sinh(bx)}{x} + \frac{1}{2} \int \frac{\sinh(2bx)}{x^2} dx \\
&= \frac{1}{2} b \text{Chi}(bx)^2 - \frac{\text{Chi}(bx) \sinh(bx)}{x} - \frac{\sinh(2bx)}{2x} + b \int \frac{\cosh(2bx)}{x} dx \\
&= \frac{1}{2} b \text{Chi}(bx)^2 + b \text{Chi}(2bx) - \frac{\text{Chi}(bx) \sinh(bx)}{x} - \frac{\sinh(2bx)}{2x}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^2} dx = \frac{1}{2} b \text{Chi}(bx)^2 + b \text{Chi}(2bx) - \frac{\text{Chi}(bx) \sinh(bx)}{x} - \frac{\sinh(2bx)}{2x}$$

[In] Integrate[(CoshIntegral[b*x]*Sinh[b*x])/x^2,x]

[Out] (b*CoshIntegral[b*x]^2)/2 + b*CoshIntegral[2*b*x] - (CoshIntegral[b*x]*Sinh[b*x])/x - Sinh[2*b*x]/(2*x)

Maple [F]

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^2} dx$$

[In] int(Chi(b*x)*sinh(b*x)/x^2,x)

[Out] int(Chi(b*x)*sinh(b*x)/x^2,x)

Fricas [F]

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^2} dx = \int \frac{\text{Chi}(bx) \sinh(bx)}{x^2} dx$$

[In] integrate(Chi(b*x)*sinh(b*x)/x^2,x, algorithm="fricas")

[Out] integral(cosh_integral(b*x)*sinh(b*x)/x^2, x)

Sympy [F]

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^2} dx = \int \frac{\sinh(bx) \text{Chi}(bx)}{x^2} dx$$

[In] integrate(Chi(b*x)*sinh(b*x)/x**2,x)

[Out] Integral(sinh(b*x)*Chi(b*x)/x**2, x)

Maxima [F]

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^2} dx = \int \frac{\text{Chi}(bx) \sinh(bx)}{x^2} dx$$

[In] integrate(Chi(b*x)*sinh(b*x)/x^2,x, algorithm="maxima")

[Out] integrate(Chi(b*x)*sinh(b*x)/x^2, x)

Giac [F]

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^2} dx = \int \frac{\text{Chi}(bx) \sinh(bx)}{x^2} dx$$

[In] integrate(Chi(b*x)*sinh(b*x)/x^2,x, algorithm="giac")

[Out] integrate(Chi(b*x)*sinh(b*x)/x^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x^2} dx = \int \frac{\text{coshint}(bx) \sinh(bx)}{x^2} dx$$

[In] int((coshint(b*x)*sinh(b*x))/x^2,x)

[Out] int((coshint(b*x)*sinh(b*x))/x^2, x)

3.116 $\int \frac{\mathbf{Chi}(bx) \sinh(bx)}{x} dx$

Optimal result	567
Rubi [N/A]	567
Mathematica [N/A]	568
Maple [N/A] (verified)	568
Fricas [N/A]	568
Sympy [N/A]	568
Maxima [N/A]	569
Giac [N/A]	569
Mupad [N/A]	569

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\mathbf{Chi}(bx) \sinh(bx)}{x} dx = \text{Int}\left(\frac{\mathbf{Chi}(bx) \sinh(bx)}{x}, x\right)$$

[Out] `CannotIntegrate(Chi(b*x)*sinh(b*x)/x,x)`

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\mathbf{Chi}(bx) \sinh(bx)}{x} dx = \int \frac{\mathbf{Chi}(bx) \sinh(bx)}{x} dx$$

[In] `Int[(CoshIntegral[b*x]*Sinh[b*x])/x,x]`

[Out] `Defer[Int] [(CoshIntegral[b*x]*Sinh[b*x])/x, x]`

Rubi steps

$$\text{integral} = \int \frac{\mathbf{Chi}(bx) \sinh(bx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx = \int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx$$

[In] Integrate[(CoshIntegral[b*x]*Sinh[b*x])/x,x]

[Out] Integrate[(CoshIntegral[b*x]*Sinh[b*x])/x, x]

Maple [N/A] (verified)

Not integrable

Time = 0.23 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx$$

[In] int(Chi(b*x)*sinh(b*x)/x,x)

[Out] int(Chi(b*x)*sinh(b*x)/x,x)

Fricas [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx = \int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx$$

[In] integrate(Chi(b*x)*sinh(b*x)/x,x, algorithm="fricas")

[Out] integral(cosh_integral(b*x)*sinh(b*x)/x, x)

Sympy [N/A]

Not integrable

Time = 3.87 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx = \int \frac{\sinh(bx) \text{Chi}(bx)}{x} dx$$

[In] integrate(Chi(b*x)*sinh(b*x)/x,x)

[Out] Integral(sinh(b*x)*Chi(b*x)/x, x)

Maxima [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx = \int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx$$

[In] integrate(Chi(b*x)*sinh(b*x)/x,x, algorithm="maxima")

[Out] integrate(Chi(b*x)*sinh(b*x)/x, x)

Giac [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx = \int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx$$

[In] integrate(Chi(b*x)*sinh(b*x)/x,x, algorithm="giac")

[Out] integrate(Chi(b*x)*sinh(b*x)/x, x)

Mupad [N/A]

Not integrable

Time = 4.88 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\text{Chi}(bx) \sinh(bx)}{x} dx = \int \frac{\text{coshint}(bx) \sinh(bx)}{x} dx$$

[In] int((coshint(b*x)*sinh(b*x))/x,x)

[Out] int((coshint(b*x)*sinh(b*x))/x, x)

3.117 $\int \text{Chi}(bx) \sinh(bx) dx$

Optimal result	570
Rubi [A] (verified)	570
Mathematica [A] (verified)	571
Maple [A] (verified)	572
Fricas [F]	572
Sympy [F]	572
Maxima [F]	572
Giac [F]	573
Mupad [F(-1)]	573

Optimal result

Integrand size = 9, antiderivative size = 34

$$\int \text{Chi}(bx) \sinh(bx) dx = \frac{\cosh(bx)\text{Chi}(bx)}{b} - \frac{\text{Chi}(2bx)}{2b} - \frac{\log(x)}{2b}$$

[Out] $-1/2*\text{Chi}(2*b*x)/b+\text{Chi}(b*x)*\cosh(b*x)/b-1/2*\ln(x)/b$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {6682, 12, 3393, 3382}

$$\int \text{Chi}(bx) \sinh(bx) dx = -\frac{\text{Chi}(2bx)}{2b} + \frac{\text{Chi}(bx) \cosh(bx)}{b} - \frac{\log(x)}{2b}$$

[In] `Int[CoshIntegral[b*x]*Sinh[b*x],x]`

[Out] `(Cosh[b*x]*CoshIntegral[b*x])/b - CoshIntegral[2*b*x]/(2*b) - Log[x]/(2*b)`

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 3382

`Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`

Rule 3393

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] :> Int
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 6682

```
Int[CoshIntegral[(c_.) + (d_.)*(x_)]*Sinh[(a_.) + (b_.)*(x_)], x_Symbol] :>
Simp[Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\cosh(bx)\text{Chi}(bx)}{b} - \int \frac{\cosh^2(bx)}{bx} dx \\
&= \frac{\cosh(bx)\text{Chi}(bx)}{b} - \frac{\int \frac{\cosh^2(bx)}{x} dx}{b} \\
&= \frac{\cosh(bx)\text{Chi}(bx)}{b} - \frac{\int \left(\frac{1}{2x} + \frac{\cosh(2bx)}{2x} \right) dx}{b} \\
&= \frac{\cosh(bx)\text{Chi}(bx)}{b} - \frac{\log(x)}{2b} - \frac{\int \frac{\cosh(2bx)}{x} dx}{2b} \\
&= \frac{\cosh(bx)\text{Chi}(bx)}{b} - \frac{\text{Chi}(2bx)}{2b} - \frac{\log(x)}{2b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.06

$$\int \text{Chi}(bx) \sinh(bx) dx = \frac{\cosh(bx)\text{Chi}(bx)}{b} - \frac{\text{Chi}(2bx)}{2b} - \frac{\log(bx)}{2b}$$

```
[In] Integrate[CoshIntegral[b*x]*Sinh[b*x],x]
```

```
[Out] (Cosh[b*x]*CoshIntegral[b*x])/b - CoshIntegral[2*b*x]/(2*b) - Log[b*x]/(2*b
)
```

Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$\frac{\text{Chi}(bx) \cosh(bx) - \frac{\ln(bx)}{2} - \frac{\text{Chi}(2bx)}{2}}{b}$	28
default	$\frac{\text{Chi}(bx) \cosh(bx) - \frac{\ln(bx)}{2} - \frac{\text{Chi}(2bx)}{2}}{b}$	28

[In] `int(Chi(b*x)*sinh(b*x),x,method=_RETURNVERBOSE)`

[Out] `1/b*(Chi(b*x)*cosh(b*x)-1/2*ln(b*x)-1/2*Chi(2*b*x))`

Fricas [F]

$$\int \text{Chi}(bx) \sinh(bx) dx = \int \text{Chi}(bx) \sinh(bx) dx$$

[In] `integrate(Chi(b*x)*sinh(b*x),x, algorithm="fricas")`

[Out] `integral(cosh_integral(b*x)*sinh(b*x), x)`

Sympy [F]

$$\int \text{Chi}(bx) \sinh(bx) dx = \int \sinh(bx) \text{Chi}(bx) dx$$

[In] `integrate(Chi(b*x)*sinh(b*x),x)`

[Out] `Integral(sinh(b*x)*Chi(b*x), x)`

Maxima [F]

$$\int \text{Chi}(bx) \sinh(bx) dx = \int \text{Chi}(bx) \sinh(bx) dx$$

[In] `integrate(Chi(b*x)*sinh(b*x),x, algorithm="maxima")`

[Out] `integrate(Chi(b*x)*sinh(b*x), x)`

Giac [**F**]

$$\int \operatorname{Chi}(bx) \sinh(bx) dx = \int \operatorname{Chi}(bx) \sinh(bx) dx$$

[In] integrate(Chi(b*x)*sinh(b*x),x, algorithm="giac")

[Out] integrate(Chi(b*x)*sinh(b*x), x)

Mupad [**F(-1)**]

Timed out.

$$\int \operatorname{Chi}(bx) \sinh(bx) dx = \int \operatorname{coshint}(bx) \sinh(bx) dx$$

[In] int(coshint(b*x)*sinh(b*x),x)

[Out] int(coshint(b*x)*sinh(b*x), x)

3.118 $\int x \operatorname{Chi}(bx) \sinh(bx) dx$

Optimal result	574
Rubi [A] (verified)	574
Mathematica [A] (verified)	576
Maple [A] (verified)	576
Fricas [F]	577
Sympy [F]	577
Maxima [F]	577
Giac [F]	577
Mupad [F(-1)]	578

Optimal result

Integrand size = 10, antiderivative size = 62

$$\int x \operatorname{Chi}(bx) \sinh(bx) dx = -\frac{x}{2b} + \frac{x \cosh(bx) \operatorname{Chi}(bx)}{b} - \frac{\cosh(bx) \sinh(bx)}{2b^2} - \frac{\operatorname{Chi}(bx) \sinh(bx)}{b^2} + \frac{\operatorname{Shi}(2bx)}{2b^2}$$

[Out] $-1/2*x/b+x*\operatorname{Chi}(b*x)*\cosh(b*x)/b+1/2*\operatorname{Shi}(2*b*x)/b^2-\operatorname{Chi}(b*x)*\sinh(b*x)/b^2-1/2*\cosh(b*x)*\sinh(b*x)/b^2$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {6684, 12, 2715, 8, 6676, 5556, 3379}

$$\int x \operatorname{Chi}(bx) \sinh(bx) dx = -\frac{\operatorname{Chi}(bx) \sinh(bx)}{b^2} + \frac{\operatorname{Shi}(2bx)}{2b^2} - \frac{\sinh(bx) \cosh(bx)}{2b^2} + \frac{x \operatorname{Chi}(bx) \cosh(bx)}{b} - \frac{x}{2b}$$

[In] `Int[x*CoshIntegral[b*x]*Sinh[b*x],x]`

[Out] $-1/2*x/b + (x*\operatorname{Cosh}[b*x]*\operatorname{CoshIntegral}[b*x])/b - (\operatorname{Cosh}[b*x]*\operatorname{Sinh}[b*x])/(2*b^2) - (\operatorname{CoshIntegral}[b*x]*\operatorname{Sinh}[b*x])/b^2 + \operatorname{SinhIntegral}[2*b*x]/(2*b^2)$

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 2715

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3379

Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 5556

Int[Cosh[(a_) + (b_)*(x_)]^(p_)*((c_) + (d_)*(x_))^(m_)*Sinh[(a_) + (b_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]

Rule 6676

Int[Cosh[(a_) + (b_)*(x_)]*CoshIntegral[(c_) + (d_)*(x_)], x_Symbol] := Simp[Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]

Rule 6684

Int[CoshIntegral[(c_) + (d_)*(x_)]*((e_) + (f_)*(x_))^(m_)*Sinh[(a_) + (b_)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*CoshIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x \cosh(bx) \text{Chi}(bx)}{b} - \frac{\int \cosh(bx) \text{Chi}(bx) dx}{b} - \int \frac{\cosh^2(bx)}{b} dx \\
 &= \frac{x \cosh(bx) \text{Chi}(bx)}{b} - \frac{\text{Chi}(bx) \sinh(bx)}{b^2} - \frac{\int \cosh^2(bx) dx}{b} + \frac{\int \frac{\cosh(bx) \sinh(bx)}{bx} dx}{b} \\
 &= \frac{x \cosh(bx) \text{Chi}(bx)}{b} - \frac{\cosh(bx) \sinh(bx)}{2b^2} - \frac{\text{Chi}(bx) \sinh(bx)}{b^2} + \frac{\int \frac{\cosh(bx) \sinh(bx)}{x} dx}{b^2} - \frac{\int 1 dx}{2b}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{x}{2b} + \frac{x \cosh(bx) \operatorname{Chi}(bx)}{b} - \frac{\cosh(bx) \sinh(bx)}{2b^2} - \frac{\operatorname{Chi}(bx) \sinh(bx)}{b^2} + \frac{\int \frac{\sinh(2bx)}{2x} dx}{b^2} \\
&= -\frac{x}{2b} + \frac{x \cosh(bx) \operatorname{Chi}(bx)}{b} - \frac{\cosh(bx) \sinh(bx)}{2b^2} - \frac{\operatorname{Chi}(bx) \sinh(bx)}{b^2} + \frac{\int \frac{\sinh(2bx)}{x} dx}{2b^2} \\
&= -\frac{x}{2b} + \frac{x \cosh(bx) \operatorname{Chi}(bx)}{b} - \frac{\cosh(bx) \sinh(bx)}{2b^2} - \frac{\operatorname{Chi}(bx) \sinh(bx)}{b^2} + \frac{\operatorname{Shi}(2bx)}{2b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.71

$$\begin{aligned}
&\int x \operatorname{Chi}(bx) \sinh(bx) dx \\
&= -\frac{2bx + \operatorname{Chi}(bx)(-4bx \cosh(bx) + 4 \sinh(bx)) + \sinh(2bx) - 2\operatorname{Shi}(2bx)}{4b^2}
\end{aligned}$$

[In] Integrate[x*CoshIntegral[b*x]*Sinh[b*x],x]

[Out] -1/4*(2*b*x + CoshIntegral[b*x]*(-4*b*x*Cosh[b*x] + 4*Sinh[b*x]) + Sinh[2*b*x] - 2*SinhIntegral[2*b*x])/b^2

Maple [A] (verified)

Time = 0.71 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.74

method	result	size
derivativedivides	$\frac{\operatorname{Chi}(bx)(bx \cosh(bx) - \sinh(bx)) - \frac{\cosh(bx) \sinh(bx)}{2} - \frac{bx}{2} + \frac{\operatorname{Shi}(2bx)}{2}}{b^2}$	46
default	$\frac{\operatorname{Chi}(bx)(bx \cosh(bx) - \sinh(bx)) - \frac{\cosh(bx) \sinh(bx)}{2} - \frac{bx}{2} + \frac{\operatorname{Shi}(2bx)}{2}}{b^2}$	46

[In] int(x*Chi(b*x)*sinh(b*x),x,method=_RETURNVERBOSE)

[Out] 1/b^2*(Chi(b*x)*(b*x*cosh(b*x)-sinh(b*x))-1/2*cosh(b*x)*sinh(b*x)-1/2*b*x+1/2*Shi(2*b*x))

Fricas [F]

$$\int x\text{Chi}(bx) \sinh(bx) dx = \int x\text{Chi}(bx) \sinh (bx) dx$$

[In] integrate(x*Chi(b*x)*sinh(b*x),x, algorithm="fricas")

[Out] integral(x*cosh_integral(b*x)*sinh(b*x), x)

Sympy [F]

$$\int x\text{Chi}(bx) \sinh(bx) dx = \int x \sinh (bx) \text{Chi} (bx) dx$$

[In] integrate(x*Chi(b*x)*sinh(b*x),x)

[Out] Integral(x*sinh(b*x)*Chi(b*x), x)

Maxima [F]

$$\int x\text{Chi}(bx) \sinh(bx) dx = \int x\text{Chi}(bx) \sinh (bx) dx$$

[In] integrate(x*Chi(b*x)*sinh(b*x),x, algorithm="maxima")

[Out] integrate(x*Chi(b*x)*sinh(b*x), x)

Giac [F]

$$\int x\text{Chi}(bx) \sinh(bx) dx = \int x\text{Chi}(bx) \sinh (bx) dx$$

[In] integrate(x*Chi(b*x)*sinh(b*x),x, algorithm="giac")

[Out] integrate(x*Chi(b*x)*sinh(b*x), x)

Mupad [F(-1)]

Timed out.

$$\int x\text{Chi}(bx) \sinh(bx) dx = \int x \text{coshint}(bx) \sinh(bx) dx$$

```
[In] int(x*coshint(b*x)*sinh(b*x),x)
```

```
[Out] int(x*coshint(b*x)*sinh(b*x), x)
```

3.119 $\int x^2 \mathbf{Chi}(bx) \sinh(bx) dx$

Optimal result	579
Rubi [A] (verified)	579
Mathematica [A] (verified)	582
Maple [A] (verified)	582
Fricas [F]	582
Sympy [F]	583
Maxima [F]	583
Giac [F]	583
Mupad [F(-1)]	583

Optimal result

Integrand size = 12, antiderivative size = 109

$$\int x^2 \mathbf{Chi}(bx) \sinh(bx) dx = -\frac{x^2}{4b} + \frac{\cosh^2(bx)}{4b^3} + \frac{2 \cosh(bx) \mathbf{Chi}(bx)}{b^3} \\ + \frac{x^2 \cosh(bx) \mathbf{Chi}(bx)}{b} - \frac{\mathbf{Chi}(2bx)}{b^3} - \frac{\log(x)}{b^3} \\ - \frac{x \cosh(bx) \sinh(bx)}{2b^2} - \frac{2x \mathbf{Chi}(bx) \sinh(bx)}{b^2} + \frac{\sinh^2(bx)}{b^3}$$

[Out] $-1/4*x^2/b - \mathbf{Chi}(2*b*x)/b^3 + 2*\mathbf{Chi}(b*x)*\cosh(b*x)/b^3 + x^2*\mathbf{Chi}(b*x)*\cosh(b*x)/b$
 $+ 1/4*\cosh(b*x)^2/b^3 - \ln(x)/b^3 - 2*x*\mathbf{Chi}(b*x)*\sinh(b*x)/b^2 - 1/2*x*\cosh(b*x)*\sinh(b*x)/b^2 + \sinh(b*x)^2/b^3$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {6684, 12, 3391, 30, 6678, 2644, 6682, 3393, 3382}

$$\int x^2 \mathbf{Chi}(bx) \sinh(bx) dx = -\frac{\mathbf{Chi}(2bx)}{b^3} + \frac{2\mathbf{Chi}(bx) \cosh(bx)}{b^3} - \frac{\log(x)}{b^3} \\ + \frac{\sinh^2(bx)}{b^3} + \frac{\cosh^2(bx)}{4b^3} - \frac{2x \mathbf{Chi}(bx) \sinh(bx)}{b^2} \\ - \frac{x \sinh(bx) \cosh(bx)}{2b^2} + \frac{x^2 \mathbf{Chi}(bx) \cosh(bx)}{b} - \frac{x^2}{4b}$$

[In] $\text{Int}[x^2*\text{CoshIntegral}[b*x]*\text{Sinh}[b*x], x]$

[Out] $-1/4*x^2/b + \text{Cosh}[b*x]^2/(4*b^3) + (2*\text{Cosh}[b*x]*\text{CoshIntegral}[b*x])/b^3 + (x^2*\text{Cosh}[b*x]*\text{CoshIntegral}[b*x])/b - \text{CoshIntegral}[2*b*x]/b^3 - \text{Log}[x]/b^3 -$

$(x \cosh[bx] \sinh[bx]) / (2b^2) - (2x \operatorname{CoshIntegral}[bx] \sinh[bx]) / b^2 + \operatorname{Sinh}[bx]^2 / b^3$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 30

`Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]`

Rule 2644

`Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)]^(m_), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])`

Rule 3382

`Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`

Rule 3391

`Int[((c_) + (d_)*(x_))*((b_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Simp[d*((b*Sin[e + f*x])^n/(f^2*n^2), x] + (Dist[b^2*((n - 1)/n), Int[(c + d*x)*(b*Sin[e + f*x])^(n - 2), x], x] - Simp[b*(c + d*x)*Cos[e + f*x]*((b*Sin[e + f*x])^(n - 1)/(f*n)), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1]`

Rule 3393

`Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

Rule 6678

`Int[Cosh[(a_) + (b_)*(x_)]*CoshIntegral[(c_) + (d_)*(x_)]*((e_) + (f_)*(x_))^(m_), x_Symbol] := Simp[(e + f*x)^m*Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*CoshIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]`

Rule 6682

```
Int[CoshIntegral[(c_.) + (d_.)*(x_)]*Sinh[(a_.) + (b_.)*(x_)], x_Symbol] :>
  Simp[Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
  b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6684

```
Int[CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.)
+ (b_.)*(x_)], x_Symbol] :> Simp[(e + f*x)^m*Cosh[a + b*x]*(CoshIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Cosh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*CoshIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x^2 \cosh(bx) \text{Chi}(bx)}{b} - \frac{2 \int x \cosh(bx) \text{Chi}(bx) dx}{b} - \int \frac{x \cosh^2(bx)}{b} dx \\
&= \frac{x^2 \cosh(bx) \text{Chi}(bx)}{b} - \frac{2x \text{Chi}(bx) \sinh(bx)}{b^2} + \frac{2 \int \text{Chi}(bx) \sinh(bx) dx}{b^2} \\
&\quad - \frac{\int x \cosh^2(bx) dx}{b} + \frac{2 \int \frac{\cosh(bx) \sinh(bx)}{b} dx}{b} \\
&= \frac{\cosh^2(bx)}{4b^3} + \frac{2 \cosh(bx) \text{Chi}(bx)}{b^3} + \frac{x^2 \cosh(bx) \text{Chi}(bx)}{b} - \frac{x \cosh(bx) \sinh(bx)}{2b^2} \\
&\quad - \frac{2x \text{Chi}(bx) \sinh(bx)}{b^2} - \frac{2 \int \frac{\cosh^2(bx)}{bx} dx}{b^2} + \frac{2 \int \cosh(bx) \sinh(bx) dx}{b^2} - \frac{\int x dx}{2b} \\
&= -\frac{x^2}{4b} + \frac{\cosh^2(bx)}{4b^3} + \frac{2 \cosh(bx) \text{Chi}(bx)}{b^3} + \frac{x^2 \cosh(bx) \text{Chi}(bx)}{b} - \frac{x \cosh(bx) \sinh(bx)}{2b^2} \\
&\quad - \frac{2x \text{Chi}(bx) \sinh(bx)}{b^2} - \frac{2 \int \frac{\cosh^2(bx)}{x} dx}{b^3} - \frac{2 \text{Subst}(\int x dx, x, i \sinh(bx))}{b^3} \\
&= -\frac{x^2}{4b} + \frac{\cosh^2(bx)}{4b^3} + \frac{2 \cosh(bx) \text{Chi}(bx)}{b^3} + \frac{x^2 \cosh(bx) \text{Chi}(bx)}{b} \\
&\quad - \frac{x \cosh(bx) \sinh(bx)}{2b^2} - \frac{2x \text{Chi}(bx) \sinh(bx)}{b^2} + \frac{\sinh^2(bx)}{b^3} - \frac{2 \int \left(\frac{1}{2x} + \frac{\cosh(2bx)}{2x} \right) dx}{b^3} \\
&= -\frac{x^2}{4b} + \frac{\cosh^2(bx)}{4b^3} + \frac{2 \cosh(bx) \text{Chi}(bx)}{b^3} + \frac{x^2 \cosh(bx) \text{Chi}(bx)}{b} - \frac{\log(x)}{b^3} \\
&\quad - \frac{x \cosh(bx) \sinh(bx)}{2b^2} - \frac{2x \text{Chi}(bx) \sinh(bx)}{b^2} + \frac{\sinh^2(bx)}{b^3} - \frac{\int \frac{\cosh(2bx)}{x} dx}{b^3} \\
&= -\frac{x^2}{4b} + \frac{\cosh^2(bx)}{4b^3} + \frac{2 \cosh(bx) \text{Chi}(bx)}{b^3} + \frac{x^2 \cosh(bx) \text{Chi}(bx)}{b} - \frac{\text{Chi}(2bx)}{b^3} \\
&\quad - \frac{\log(x)}{b^3} - \frac{x \cosh(bx) \sinh(bx)}{2b^2} - \frac{2x \text{Chi}(bx) \sinh(bx)}{b^2} + \frac{\sinh^2(bx)}{b^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.66

$$\int x^2 \text{Chi}(bx) \sinh(bx) dx = \frac{2b^2 x^2 - 5 \cosh(2bx) + 8 \text{Chi}(2bx) + 8 \log(x) - 8 \text{Chi}(bx) ((2 + b^2 x^2) \cosh(bx) - 2bx \sinh(bx)) + 2bx \sinh(bx)}{8b^3}$$

[In] Integrate[x^2*CoshIntegral[b*x]*Sinh[b*x],x]

[Out] -1/8*(2*b^2*x^2 - 5*Cosh[2*b*x] + 8*CoshIntegral[2*b*x] + 8*Log[x] - 8*CoshIntegral[b*x]*((2 + b^2*x^2)*Cosh[b*x] - 2*b*x*Sinh[b*x]) + 2*b*x*Sinh[2*b*x])/b^3

Maple [A] (verified)

Time = 0.80 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.72

method	result	size
derivativedivides	$\frac{\text{Chi}(bx)(b^2 x^2 \cosh(bx) - 2bx \sinh(bx) + 2 \cosh(bx)) - \frac{bx \cosh(bx) \sinh(bx)}{2} - \frac{b^2 x^2}{4} + \frac{5 \cosh(bx)^2}{4} - \ln(bx) - \text{Chi}(2bx)}{b^3}$	78
default	$\frac{\text{Chi}(bx)(b^2 x^2 \cosh(bx) - 2bx \sinh(bx) + 2 \cosh(bx)) - \frac{bx \cosh(bx) \sinh(bx)}{2} - \frac{b^2 x^2}{4} + \frac{5 \cosh(bx)^2}{4} - \ln(bx) - \text{Chi}(2bx)}{b^3}$	78

[In] int(x^2*Chi(b*x)*sinh(b*x),x,method=_RETURNVERBOSE)

[Out] 1/b^3*(Chi(b*x)*(b^2*x^2*cosh(b*x)-2*b*x*sinh(b*x)+2*cosh(b*x))-1/2*b*x*cosh(b*x)*sinh(b*x)-1/4*b^2*x^2+5/4*cosh(b*x)^2-ln(b*x)-Chi(2*b*x))

Fricas [F]

$$\int x^2 \text{Chi}(bx) \sinh(bx) dx = \int x^2 \text{Chi}(bx) \sinh(bx) dx$$

[In] integrate(x^2*Chi(b*x)*sinh(b*x),x, algorithm="fricas")

[Out] integral(x^2*cosh_integral(b*x)*sinh(b*x), x)

Sympy [F]

$$\int x^2 \operatorname{Chi}(bx) \sinh(bx) dx = \int x^2 \sinh(bx) \operatorname{Chi}(bx) dx$$

[In] `integrate(x**2*Chi(b*x)*sinh(b*x),x)`

[Out] `Integral(x**2*sinh(b*x)*Chi(b*x), x)`

Maxima [F]

$$\int x^2 \operatorname{Chi}(bx) \sinh(bx) dx = \int x^2 \operatorname{Chi}(bx) \sinh(bx) dx$$

[In] `integrate(x^2*Chi(b*x)*sinh(b*x),x, algorithm="maxima")`

[Out] `integrate(x^2*Chi(b*x)*sinh(b*x), x)`

Giac [F]

$$\int x^2 \operatorname{Chi}(bx) \sinh(bx) dx = \int x^2 \operatorname{Chi}(bx) \sinh(bx) dx$$

[In] `integrate(x^2*Chi(b*x)*sinh(b*x),x, algorithm="giac")`

[Out] `integrate(x^2*Chi(b*x)*sinh(b*x), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{Chi}(bx) \sinh(bx) dx = \int x^2 \operatorname{coshint}(bx) \sinh(bx) dx$$

[In] `int(x^2*coshint(b*x)*sinh(b*x),x)`

[Out] `int(x^2*coshint(b*x)*sinh(b*x), x)`

3.120 $\int x^3 \text{Chi}(bx) \sinh(bx) dx$

Optimal result	584
Rubi [A] (verified)	584
Mathematica [A] (verified)	588
Maple [A] (verified)	588
Fricas [F]	588
Sympy [F]	589
Maxima [F]	589
Giac [F]	589
Mupad [F(-1)]	589

Optimal result

Integrand size = 12, antiderivative size = 146

$$\int x^3 \text{Chi}(bx) \sinh(bx) dx = -\frac{5x}{2b^3} - \frac{x^3}{6b} + \frac{x \cosh^2(bx)}{2b^3} + \frac{6x \cosh(bx) \text{Chi}(bx)}{b^3} \\ + \frac{x^3 \cosh(bx) \text{Chi}(bx)}{b} - \frac{4 \cosh(bx) \sinh(bx)}{b^4} \\ - \frac{x^2 \cosh(bx) \sinh(bx)}{2b^2} - \frac{6 \text{Chi}(bx) \sinh(bx)}{b^4} \\ - \frac{3x^2 \text{Chi}(bx) \sinh(bx)}{b^2} + \frac{3x \sinh^2(bx)}{2b^3} + \frac{3 \text{Shi}(2bx)}{b^4}$$

[Out] $-5/2*x/b^3-1/6*x^3/b+6*x*\text{Chi}(b*x)*\cosh(b*x)/b^3+x^3*\text{Chi}(b*x)*\cosh(b*x)/b+1/2*x*\cosh(b*x)^2/b^3+3*\text{Shi}(2*b*x)/b^4-6*\text{Chi}(b*x)*\sinh(b*x)/b^4-3*x^2*\text{Chi}(b*x)*\sinh(b*x)/b^2-4*\cosh(b*x)*\sinh(b*x)/b^4-1/2*x^2*\cosh(b*x)*\sinh(b*x)/b^2+3/2*x*\sinh(b*x)^2/b^3$

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.00, number of steps used = 20, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$, Rules used = {6684, 12, 3392, 30, 2715, 8, 6678, 5480, 6676, 5556, 3379}

$$\int x^3 \text{Chi}(bx) \sinh(bx) dx = -\frac{6 \text{Chi}(bx) \sinh(bx)}{b^4} + \frac{3 \text{Shi}(2bx)}{b^4} - \frac{4 \sinh(bx) \cosh(bx)}{b^4} \\ + \frac{6x \text{Chi}(bx) \cosh(bx)}{b^3} - \frac{5x}{2b^3} + \frac{3x \sinh^2(bx)}{2b^3} \\ + \frac{x \cosh^2(bx)}{2b^3} - \frac{3x^2 \text{Chi}(bx) \sinh(bx)}{b^2} \\ - \frac{x^2 \sinh(bx) \cosh(bx)}{2b^2} + \frac{x^3 \text{Chi}(bx) \cosh(bx)}{b} - \frac{x^3}{6b}$$

[In] Int[x^3*CoshIntegral[b*x]*Sinh[b*x],x]

[Out] (-5*x)/(2*b^3) - x^3/(6*b) + (x*Cosh[b*x]^2)/(2*b^3) + (6*x*Cosh[b*x]*CoshIntegral[b*x])/b^3 + (x^3*Cosh[b*x]*CoshIntegral[b*x])/b - (4*Cosh[b*x]*Sinh[b*x])/b^4 - (x^2*Cosh[b*x]*Sinh[b*x])/(2*b^2) - (6*CoshIntegral[b*x]*Sinh[b*x])/b^4 - (3*x^2*CoshIntegral[b*x]*Sinh[b*x])/b^2 + (3*x*Sinh[b*x]^2)/(2*b^3) + (3*SinhIntegral[2*b*x])/b^4

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2715

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3379

Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 3392

Int[((c_) + (d_)*(x_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[d*m*(c + d*x)^(m - 1)*((b*Sin[e + f*x])^n/(f^2*n^2)), x] + (Dist[b^2*((n - 1)/n), Int[(c + d*x)^m*(b*Sin[e + f*x])^(n - 2), x], x] - Dist[d^2*m*((m - 1)/(f^2*n^2)), Int[(c + d*x)^(m - 2)*(b*Sin[e + f*x])^n, x], x] - Simp[b*(c + d*x)^m*Cos[e + f*x]*((b*Sin[e + f*x])^(n - 1)/(f*n)), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 1]

Rule 5480

Int[Cosh[(a_) + (b_)*(x_)^(n_)]*(x_)^(m_)*Sinh[(a_) + (b_)*(x_)^(n_)]^(p_), x_Symbol] := Simp[x^(m - n + 1)*(Sinh[a + b*x^n]^(p + 1)/(b*n*(p +

1))), x] - Dist[(m - n + 1)/(b*n*(p + 1)), Int[x^(m - n)*Sinh[a + b*x^n]^(p + 1), x], x] /; FreeQ[{a, b, p}, x] && LtQ[0, n, m + 1] && NeQ[p, -1]

Rule 5556

Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^(n)*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]

Rule 6676

Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]

Rule 6678

Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^m*Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*CoshIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rule 6684

Int[CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*CoshIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{x^3 \cosh(bx) \text{Chi}(bx)}{b} - \frac{3 \int x^2 \cosh(bx) \text{Chi}(bx) dx}{b} - \int \frac{x^2 \cosh^2(bx)}{b} dx \\ &= \frac{x^3 \cosh(bx) \text{Chi}(bx)}{b} - \frac{3x^2 \text{Chi}(bx) \sinh(bx)}{b^2} + \frac{6 \int x \text{Chi}(bx) \sinh(bx) dx}{b^2} \\ &\quad - \frac{\int x^2 \cosh^2(bx) dx}{b} + \frac{3 \int \frac{x \cosh(bx) \sinh(bx)}{b} dx}{b} \end{aligned}$$

$$\begin{aligned}
&= \frac{x \cosh^2(bx)}{2b^3} + \frac{6x \cosh(bx)\text{Chi}(bx)}{b^3} + \frac{x^3 \cosh(bx)\text{Chi}(bx)}{b} - \frac{x^2 \cosh(bx) \sinh(bx)}{2b^2} \\
&\quad - \frac{3x^2 \text{Chi}(bx) \sinh(bx)}{b^2} - \frac{\int \cosh^2(bx) dx}{2b^3} - \frac{6 \int \cosh(bx)\text{Chi}(bx) dx}{b^3} \\
&\quad + \frac{3 \int x \cosh(bx) \sinh(bx) dx}{b^2} - \frac{6 \int \frac{\cosh^2(bx)}{b} dx}{b^2} - \frac{\int x^2 dx}{2b} \\
&= -\frac{x^3}{6b} + \frac{x \cosh^2(bx)}{2b^3} + \frac{6x \cosh(bx)\text{Chi}(bx)}{b^3} + \frac{x^3 \cosh(bx)\text{Chi}(bx)}{b} - \frac{\cosh(bx) \sinh(bx)}{4b^4} \\
&\quad - \frac{x^2 \cosh(bx) \sinh(bx)}{2b^2} - \frac{6\text{Chi}(bx) \sinh(bx)}{b^4} - \frac{3x^2 \text{Chi}(bx) \sinh(bx)}{b^2} + \frac{3x \sinh^2(bx)}{2b^3} \\
&\quad - \frac{\int 1 dx}{4b^3} - \frac{3 \int \sinh^2(bx) dx}{2b^3} - \frac{6 \int \cosh^2(bx) dx}{b^3} + \frac{6 \int \frac{\cosh(bx) \sinh(bx)}{bx} dx}{b^3} \\
&= -\frac{x}{4b^3} - \frac{x^3}{6b} + \frac{x \cosh^2(bx)}{2b^3} + \frac{6x \cosh(bx)\text{Chi}(bx)}{b^3} + \frac{x^3 \cosh(bx)\text{Chi}(bx)}{b} \\
&\quad - \frac{4 \cosh(bx) \sinh(bx)}{b^4} - \frac{x^2 \cosh(bx) \sinh(bx)}{2b^2} - \frac{6\text{Chi}(bx) \sinh(bx)}{b^4} \\
&\quad - \frac{3x^2 \text{Chi}(bx) \sinh(bx)}{b^2} + \frac{3x \sinh^2(bx)}{2b^3} + \frac{6 \int \frac{\cosh(bx) \sinh(bx)}{x} dx}{b^4} + \frac{3 \int 1 dx}{4b^3} - \frac{3 \int 1 dx}{b^3} \\
&= -\frac{5x}{2b^3} - \frac{x^3}{6b} + \frac{x \cosh^2(bx)}{2b^3} + \frac{6x \cosh(bx)\text{Chi}(bx)}{b^3} + \frac{x^3 \cosh(bx)\text{Chi}(bx)}{b} \\
&\quad - \frac{4 \cosh(bx) \sinh(bx)}{b^4} - \frac{x^2 \cosh(bx) \sinh(bx)}{2b^2} - \frac{6\text{Chi}(bx) \sinh(bx)}{b^4} \\
&\quad - \frac{3x^2 \text{Chi}(bx) \sinh(bx)}{b^2} + \frac{3x \sinh^2(bx)}{2b^3} + \frac{6 \int \frac{\sinh(2bx)}{2x} dx}{b^4} \\
&= -\frac{5x}{2b^3} - \frac{x^3}{6b} + \frac{x \cosh^2(bx)}{2b^3} + \frac{6x \cosh(bx)\text{Chi}(bx)}{b^3} + \frac{x^3 \cosh(bx)\text{Chi}(bx)}{b} \\
&\quad - \frac{4 \cosh(bx) \sinh(bx)}{b^4} - \frac{x^2 \cosh(bx) \sinh(bx)}{2b^2} - \frac{6\text{Chi}(bx) \sinh(bx)}{b^4} \\
&\quad - \frac{3x^2 \text{Chi}(bx) \sinh(bx)}{b^2} + \frac{3x \sinh^2(bx)}{2b^3} + \frac{3 \int \frac{\sinh(2bx)}{x} dx}{b^4} \\
&= -\frac{5x}{2b^3} - \frac{x^3}{6b} + \frac{x \cosh^2(bx)}{2b^3} + \frac{6x \cosh(bx)\text{Chi}(bx)}{b^3} + \frac{x^3 \cosh(bx)\text{Chi}(bx)}{b} \\
&\quad - \frac{4 \cosh(bx) \sinh(bx)}{b^4} - \frac{x^2 \cosh(bx) \sinh(bx)}{2b^2} - \frac{6\text{Chi}(bx) \sinh(bx)}{b^4} \\
&\quad - \frac{3x^2 \text{Chi}(bx) \sinh(bx)}{b^2} + \frac{3x \sinh^2(bx)}{2b^3} + \frac{3\text{Shi}(2bx)}{b^4}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.64

$$\int x^3 \text{Chi}(bx) \sinh(bx) dx = \frac{-36bx - 2b^3x^3 + 12bx \cosh(2bx) + 12\text{Chi}(bx) (bx(6 + b^2x^2) \cosh(bx) - 3(2 + b^2x^2) \sinh(bx)) - 24 \sinh(2bx)}{12b^4}$$

[In] Integrate[x^3*CoshIntegral[b*x]*Sinh[b*x],x]

[Out] (-36*b*x - 2*b^3*x^3 + 12*b*x*Cosh[2*b*x] + 12*CoshIntegral[b*x]*(b*x*(6 + b^2*x^2)*Cosh[b*x] - 3*(2 + b^2*x^2)*Sinh[b*x]) - 24*Sinh[2*b*x] - 3*b^2*x^2*Sinh[2*b*x] + 36*SinhIntegral[2*b*x])/(12*b^4)

Maple [A] (verified)

Time = 1.03 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.71

method	result
derivativedivides	$\frac{\text{Chi}(bx)(b^3x^3 \cosh(bx) - 3b^2x^2 \sinh(bx) + 6bx \cosh(bx) - 6 \sinh(bx)) - \frac{b^2x^2 \cosh(bx) \sinh(bx)}{2} - \frac{b^3x^3}{6} + 2bx \cosh(bx)^2 - 4 \cosh(bx)}{b^4}$
default	$\frac{\text{Chi}(bx)(b^3x^3 \cosh(bx) - 3b^2x^2 \sinh(bx) + 6bx \cosh(bx) - 6 \sinh(bx)) - \frac{b^2x^2 \cosh(bx) \sinh(bx)}{2} - \frac{b^3x^3}{6} + 2bx \cosh(bx)^2 - 4 \cosh(bx)}{b^4}$

[In] int(x^3*Chi(b*x)*sinh(b*x),x,method=_RETURNVERBOSE)

[Out] 1/b^4*(Chi(b*x)*(b^3*x^3*cosh(b*x)-3*b^2*x^2*sinh(b*x)+6*b*x*cosh(b*x)-6*sinh(b*x))-1/2*b^2*x^2*cosh(b*x)*sinh(b*x)-1/6*b^3*x^3+2*b*x*cosh(b*x)^2-4*cosh(b*x)*sinh(b*x)-4*b*x+3*Shi(2*b*x))

Fricas [F]

$$\int x^3 \text{Chi}(bx) \sinh(bx) dx = \int x^3 \text{Chi}(bx) \sinh(bx) dx$$

[In] integrate(x^3*Chi(b*x)*sinh(b*x),x, algorithm="fricas")

[Out] integral(x^3*cosh_integral(b*x)*sinh(b*x), x)

Sympy [F]

$$\int x^3 \operatorname{Chi}(bx) \sinh(bx) dx = \int x^3 \sinh(bx) \operatorname{Chi}(bx) dx$$

```
[In] integrate(x**3*Chi(b*x)*sinh(b*x),x)
```

```
[Out] Integral(x**3*sinh(b*x)*Chi(b*x), x)
```

Maxima [F]

$$\int x^3 \operatorname{Chi}(bx) \sinh(bx) dx = \int x^3 \operatorname{Chi}(bx) \sinh(bx) dx$$

```
[In] integrate(x^3*Chi(b*x)*sinh(b*x),x, algorithm="maxima")
```

```
[Out] integrate(x^3*Chi(b*x)*sinh(b*x), x)
```

Giac [F]

$$\int x^3 \operatorname{Chi}(bx) \sinh(bx) dx = \int x^3 \operatorname{Chi}(bx) \sinh(bx) dx$$

```
[In] integrate(x^3*Chi(b*x)*sinh(b*x),x, algorithm="giac")
```

```
[Out] integrate(x^3*Chi(b*x)*sinh(b*x), x)
```

Mupad [F(-1)]

Timed out.

$$\int x^3 \operatorname{Chi}(bx) \sinh(bx) dx = \int x^3 \operatorname{coshint}(bx) \sinh(bx) dx$$

```
[In] int(x^3*coshint(b*x)*sinh(b*x),x)
```

```
[Out] int(x^3*coshint(b*x)*sinh(b*x), x)
```

3.121 $\int \operatorname{Chi}(2x) \sinh(5x) dx$

Optimal result	590
Rubi [A] (verified)	590
Mathematica [A] (verified)	591
Maple [A] (verified)	592
Fricas [F]	592
Sympy [F]	592
Maxima [F]	592
Giac [F]	593
Mupad [F(-1)]	593

Optimal result

Integrand size = 9, antiderivative size = 29

$$\int \operatorname{Chi}(2x) \sinh(5x) dx = \frac{1}{5} \cosh(5x) \operatorname{Chi}(2x) - \frac{\operatorname{Chi}(3x)}{10} - \frac{\operatorname{Chi}(7x)}{10}$$

[Out] $-1/10*\operatorname{Chi}(3*x)-1/10*\operatorname{Chi}(7*x)+1/5*\operatorname{Chi}(2*x)*\cosh(5*x)$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {6682, 12, 5579, 3382}

$$\int \operatorname{Chi}(2x) \sinh(5x) dx = -\frac{\operatorname{Chi}(3x)}{10} - \frac{\operatorname{Chi}(7x)}{10} + \frac{1}{5} \operatorname{Chi}(2x) \cosh(5x)$$

[In] `Int[CoshIntegral[2*x]*Sinh[5*x],x]`

[Out] $(\operatorname{Cosh}[5*x]*\operatorname{CoshIntegral}[2*x])/5 - \operatorname{CoshIntegral}[3*x]/10 - \operatorname{CoshIntegral}[7*x]/10$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 3382

`Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz`

}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 5579

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*Cosh[(c_.) + (d_.)*(x_)]^(q_.)*((e_.) +
(f_.)*(x_))^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e + f*x)^m, Cosh[a +
b*x]^p*Cosh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0]
&& IGtQ[q, 0] && IntegerQ[m]
```

Rule 6682

```
Int[CoshIntegral[(c_.) + (d_.)*(x_)]*Sinh[(a_.) + (b_.)*(x_)], x_Symbol] :=
Simp[Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{5} \cosh(5x) \text{Chi}(2x) - \frac{2}{5} \int \frac{\cosh(2x) \cosh(5x)}{2x} dx \\
&= \frac{1}{5} \cosh(5x) \text{Chi}(2x) - \frac{1}{5} \int \frac{\cosh(2x) \cosh(5x)}{x} dx \\
&= \frac{1}{5} \cosh(5x) \text{Chi}(2x) - \frac{1}{5} \int \left(\frac{\cosh(3x)}{2x} + \frac{\cosh(7x)}{2x} \right) dx \\
&= \frac{1}{5} \cosh(5x) \text{Chi}(2x) - \frac{1}{10} \int \frac{\cosh(3x)}{x} dx - \frac{1}{10} \int \frac{\cosh(7x)}{x} dx \\
&= \frac{1}{5} \cosh(5x) \text{Chi}(2x) - \frac{\text{Chi}(3x)}{10} - \frac{\text{Chi}(7x)}{10}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \text{Chi}(2x) \sinh(5x) dx = \frac{1}{10} (2 \cosh(5x) \text{Chi}(2x) - \text{Chi}(3x) - \text{Chi}(7x))$$

[In] Integrate[CoshIntegral[2*x]*Sinh[5*x],x]

[Out] (2*Cosh[5*x]*CoshIntegral[2*x] - CoshIntegral[3*x] - CoshIntegral[7*x])/10

Maple [A] (verified)

Time = 1.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

method	result	size
default	$-\frac{\text{Chi}(3x)}{10} - \frac{\text{Chi}(7x)}{10} + \frac{\text{Chi}(2x)\cosh(5x)}{5}$	24

[In] `int(Chi(2*x)*sinh(5*x),x,method=_RETURNVERBOSE)`

[Out] `-1/10*Chi(3*x)-1/10*Chi(7*x)+1/5*Chi(2*x)*cosh(5*x)`

Fricas [F]

$$\int \text{Chi}(2x) \sinh(5x) dx = \int \text{Chi}(2x) \sinh(5x) dx$$

[In] `integrate(Chi(2*x)*sinh(5*x),x, algorithm="fricas")`

[Out] `integral(cosh_integral(2*x)*sinh(5*x), x)`

Sympy [F]

$$\int \text{Chi}(2x) \sinh(5x) dx = \int \sinh(5x) \text{Chi}(2x) dx$$

[In] `integrate(Chi(2*x)*sinh(5*x),x)`

[Out] `Integral(sinh(5*x)*Chi(2*x), x)`

Maxima [F]

$$\int \text{Chi}(2x) \sinh(5x) dx = \int \text{Chi}(2x) \sinh(5x) dx$$

[In] `integrate(Chi(2*x)*sinh(5*x),x, algorithm="maxima")`

[Out] `integrate(Chi(2*x)*sinh(5*x), x)`

Giac [**F**]

$$\int \operatorname{Chi}(2x) \sinh(5x) dx = \int \operatorname{Chi}(2x) \sinh(5x) dx$$

[In] integrate(Chi(2*x)*sinh(5*x),x, algorithm="giac")

[Out] integrate(Chi(2*x)*sinh(5*x), x)

Mupad [**F(-1)**]

Timed out.

$$\int \operatorname{Chi}(2x) \sinh(5x) dx = \int \operatorname{coshint}(2x) \sinh(5x) dx$$

[In] int(coshint(2*x)*sinh(5*x),x)

[Out] int(coshint(2*x)*sinh(5*x), x)

3.122 $\int \cosh(5x)\mathbf{Chi}(2x) dx$

Optimal result	594
Rubi [A] (verified)	594
Mathematica [A] (verified)	595
Maple [A] (verified)	596
Fricas [F]	596
Sympy [F]	596
Maxima [F]	596
Giac [F]	597
Mupad [F(-1)]	597

Optimal result

Integrand size = 9, antiderivative size = 29

$$\int \cosh(5x)\mathbf{Chi}(2x) dx = \frac{1}{5}\mathbf{Chi}(2x) \sinh(5x) - \frac{\mathbf{Shi}(3x)}{10} - \frac{\mathbf{Shi}(7x)}{10}$$

[Out] -1/10*Shi(3*x)-1/10*Shi(7*x)+1/5*Chi(2*x)*sinh(5*x)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {6676, 12, 5580, 3379}

$$\int \cosh(5x)\mathbf{Chi}(2x) dx = \frac{1}{5}\mathbf{Chi}(2x) \sinh(5x) - \frac{\mathbf{Shi}(3x)}{10} - \frac{\mathbf{Shi}(7x)}{10}$$

[In] Int[Cosh[5*x]*CoshIntegral[2*x],x]

[Out] (CoshIntegral[2*x]*Sinh[5*x])/5 - SinhIntegral[3*x]/10 - SinhIntegral[7*x]/10

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x/d), x] /; FreeQ[{c, d, e, f

, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 5580

```
Int[Cosh[(c_.) + (d_.)*(x_)]^(q_.)*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(p_.), x_Symbol] := Int[ExpandTrigReduce[(e + f*x)^m, Sinh[a +
b*x]^p*Cosh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[p
, 0] && IGtQ[q, 0]
```

Rule 6676

```
Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
Simp[Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{5} \text{Chi}(2x) \sinh(5x) - \frac{2}{5} \int \frac{\cosh(2x) \sinh(5x)}{2x} dx \\
 &= \frac{1}{5} \text{Chi}(2x) \sinh(5x) - \frac{1}{5} \int \frac{\cosh(2x) \sinh(5x)}{x} dx \\
 &= \frac{1}{5} \text{Chi}(2x) \sinh(5x) - \frac{1}{5} \int \left(\frac{\sinh(3x)}{2x} + \frac{\sinh(7x)}{2x} \right) dx \\
 &= \frac{1}{5} \text{Chi}(2x) \sinh(5x) - \frac{1}{10} \int \frac{\sinh(3x)}{x} dx - \frac{1}{10} \int \frac{\sinh(7x)}{x} dx \\
 &= \frac{1}{5} \text{Chi}(2x) \sinh(5x) - \frac{\text{Shi}(3x)}{10} - \frac{\text{Shi}(7x)}{10}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \cosh(5x) \text{Chi}(2x) dx = \frac{1}{10} (2 \text{Chi}(2x) \sinh(5x) - \text{Shi}(3x) - \text{Shi}(7x))$$

[In] Integrate[Cosh[5*x]*CoshIntegral[2*x], x]

[Out] (2*CoshIntegral[2*x]*Sinh[5*x] - SinhIntegral[3*x] - SinhIntegral[7*x])/10

Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

method	result	size
default	$-\frac{\text{Shi}(3x)}{10} - \frac{\text{Shi}(7x)}{10} + \frac{\text{Chi}(2x)\sinh(5x)}{5}$	24

[In] `int(Chi(2*x)*cosh(5*x),x,method=_RETURNVERBOSE)`

[Out] `-1/10*Shi(3*x)-1/10*Shi(7*x)+1/5*Chi(2*x)*sinh(5*x)`

Fricas [F]

$$\int \cosh(5x)\text{Chi}(2x) dx = \int \text{Chi}(2x) \cosh(5x) dx$$

[In] `integrate(Chi(2*x)*cosh(5*x),x, algorithm="fricas")`

[Out] `integral(cosh(5*x)*cosh_integral(2*x), x)`

Sympy [F]

$$\int \cosh(5x)\text{Chi}(2x) dx = \int \cosh(5x) \text{Chi}(2x) dx$$

[In] `integrate(Chi(2*x)*cosh(5*x),x)`

[Out] `Integral(cosh(5*x)*Chi(2*x), x)`

Maxima [F]

$$\int \cosh(5x)\text{Chi}(2x) dx = \int \text{Chi}(2x) \cosh(5x) dx$$

[In] `integrate(Chi(2*x)*cosh(5*x),x, algorithm="maxima")`

[Out] `integrate(Chi(2*x)*cosh(5*x), x)`

Giac [F]

$$\int \cosh(5x)\operatorname{Chi}(2x) dx = \int \operatorname{Chi}(2x) \cosh(5x) dx$$

[In] integrate(Chi(2*x)*cosh(5*x),x, algorithm="giac")

[Out] integrate(Chi(2*x)*cosh(5*x), x)

Mupad [F(-1)]

Timed out.

$$\int \cosh(5x)\operatorname{Chi}(2x) dx = \int \operatorname{coshint}(2x) \cosh(5x) dx$$

[In] int(coshint(2*x)*cosh(5*x),x)

[Out] int(coshint(2*x)*cosh(5*x), x)

3.123 $\int x^2 \text{Chi}(a + bx) \sinh(a + bx) dx$

Optimal result	598
Rubi [A] (verified)	599
Mathematica [A] (verified)	603
Maple [A] (verified)	603
Fricas [F]	603
Sympy [F]	604
Maxima [F]	604
Giac [F]	604
Mupad [F(-1)]	604

Optimal result

Integrand size = 16, antiderivative size = 220

$$\begin{aligned}
 \int x^2 \text{Chi}(a + bx) \sinh(a + bx) dx = & \frac{ax}{2b^2} - \frac{x^2}{4b} + \frac{\cosh^2(a + bx)}{4b^3} + \frac{\cosh(2a + 2bx)}{2b^3} \\
 & + \frac{2 \cosh(a + bx) \text{Chi}(a + bx)}{b^3} \\
 & + \frac{x^2 \cosh(a + bx) \text{Chi}(a + bx)}{b} - \frac{\text{Chi}(2a + 2bx)}{b^3} \\
 & - \frac{a^2 \text{Chi}(2a + 2bx)}{2b^3} - \frac{\log(a + bx)}{b^3} \\
 & - \frac{a^2 \log(a + bx)}{2b^3} + \frac{a \cosh(a + bx) \sinh(a + bx)}{2b^3} \\
 & - \frac{x \cosh(a + bx) \sinh(a + bx)}{2b^2} \\
 & - \frac{2x \text{Chi}(a + bx) \sinh(a + bx)}{b^2} - \frac{a \text{Shi}(2a + 2bx)}{b^3}
 \end{aligned}$$

[Out] 1/2*a*x/b^2-1/4*x^2/b-Chi(2*b*x+2*a)/b^3-1/2*a^2*Chi(2*b*x+2*a)/b^3+2*Chi(b*x+a)*cosh(b*x+a)/b^3+x^2*Chi(b*x+a)*cosh(b*x+a)/b+1/4*cosh(b*x+a)^2/b^3+1/2*cosh(2*b*x+2*a)/b^3-ln(b*x+a)/b^3-1/2*a^2*ln(b*x+a)/b^3-a*Shi(2*b*x+2*a)/b^3-2*x*Chi(b*x+a)*sinh(b*x+a)/b^2+1/2*a*cosh(b*x+a)*sinh(b*x+a)/b^3-1/2*x*cosh(b*x+a)*sinh(b*x+a)/b^2

Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.00, number of steps used = 21, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {6684, 6874, 2715, 8, 3391, 30, 3393, 3382, 6678, 5736, 6873, 2718, 3379, 6682}

$$\int x^2 \text{Chi}(a + bx) \sinh(a + bx) dx = -\frac{a^2 \text{Chi}(2a + 2bx)}{2b^3} - \frac{a^2 \log(a + bx)}{2b^3} - \frac{\text{Chi}(2a + 2bx)}{b^3} + \frac{2\text{Chi}(a + bx) \cosh(a + bx)}{b^3} - \frac{a \text{Shi}(2a + 2bx)}{b^3} - \frac{\log(a + bx)}{b^3} + \frac{\cosh^2(a + bx)}{4b^3} + \frac{\cosh(2a + 2bx)}{2b^3} + \frac{a \sinh(a + bx) \cosh(a + bx)}{2b^3} - \frac{2x \text{Chi}(a + bx) \sinh(a + bx)}{b^2} + \frac{ax}{2b^2} - \frac{x \sinh(a + bx) \cosh(a + bx)}{2b^2} + \frac{x^2 \text{Chi}(a + bx) \cosh(a + bx)}{b} - \frac{x^2}{4b}$$

[In] Int[x^2*CoshIntegral[a + b*x]*Sinh[a + b*x],x]

[Out] (a*x)/(2*b^2) - x^2/(4*b) + Cosh[a + b*x]^2/(4*b^3) + Cosh[2*a + 2*b*x]/(2*b^3) + (2*Cosh[a + b*x]*CoshIntegral[a + b*x])/b^3 + (x^2*Cosh[a + b*x]*CoshIntegral[a + b*x])/b - CoshIntegral[2*a + 2*b*x]/b^3 - (a^2*CoshIntegral[2*a + 2*b*x])/(2*b^3) - Log[a + b*x]/b^3 - (a^2*Log[a + b*x])/(2*b^3) + (a*Cosh[a + b*x]*Sinh[a + b*x])/(2*b^3) - (x*Cosh[a + b*x]*Sinh[a + b*x])/(2*b^2) - (2*x*CoshIntegral[a + b*x]*Sinh[a + b*x])/b^2 - (a*SinhIntegral[2*a + 2*b*x])/b^3

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Sine[c + d*x])^(n - 1)/(d*n), x] + Dist[b^2*((n - 1)/n), Int[(b*Sine[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2718

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ
[{c, d}, x]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f
, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz
}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3391

```
Int[((c_.) + (d_.)*(x_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :=>
Simp[d*((b*Sine[e + f*x])^n/(f^2*n^2)), x] + (Dist[b^2*((n - 1)/n), Int[(c
+ d*x)*(b*Sine[e + f*x])^(n - 2), x], x] - Simp[b*(c + d*x)*Cos[e + f*x]*((b
*Sine[e + f*x])^(n - 1)/(f*n)), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1
]
```

Rule 3393

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] :=> In
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 5736

```
Int[Cosh[w_]^(p_.)*(u_.)*Sinh[v_]^(p_.), x_Symbol] :=> Dist[1/2^p, Int[u*Sin
h[2*v]^p, x], x] /; EqQ[w, v] && IntegerQ[p]
```

Rule 6678

```
Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.
)*(x_))^(m_.), x_Symbol] :=> Simp[(e + f*x)^m*Sinh[a + b*x]*(CoshIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Cosh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*CoshIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6682

```
Int[CoshIntegral[(c_.) + (d_.)*(x_)]*Sinh[(a_.) + (b_.)*(x_)], x_Symbol] :=>
Simp[Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
```


$b*x]*(\text{Cosh}[c + d*x]/(c + d*x)), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x]$

Rule 6684

$\text{Int}[\text{CoshIntegral}[(c_.) + (d_.)*(x_.)]*((e_.) + (f_.)*(x_.))^{(m_.)*\text{Sinh}[(a_.) + (b_.)*(x_.)]}, x_Symbol] \text{ :> } \text{Simp}[(e + f*x)^m*\text{Cosh}[a + b*x]*(\text{CoshIntegral}[c + d*x]/b), x] + (-\text{Dist}[d/b, \text{Int}[(e + f*x)^m*\text{Cosh}[a + b*x]*(\text{Cosh}[c + d*x]/(c + d*x)), x], x] - \text{Dist}[f*(m/b), \text{Int}[(e + f*x)^{(m-1)}*\text{Cosh}[a + b*x]*\text{CoshIntegral}[c + d*x], x], x]) /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{IGtQ}[m, 0]$

Rule 6873

$\text{Int}[u_, x_Symbol] \text{ :> } \text{With}[\{v = \text{NormalizeIntegrand}[u, x]\}, \text{Int}[v, x] /; v \neq u]$

Rule 6874

$\text{Int}[u_, x_Symbol] \text{ :> } \text{With}[\{v = \text{ExpandIntegrand}[u, x]\}, \text{Int}[v, x] /; \text{SumQ}[v]]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x^2 \cosh(a + bx) \text{Chi}(a + bx)}{b} \\
 &\quad - \frac{2 \int x \cosh(a + bx) \text{Chi}(a + bx) dx}{b} - \int \frac{x^2 \cosh^2(a + bx)}{a + bx} dx \\
 &= \frac{x^2 \cosh(a + bx) \text{Chi}(a + bx)}{b} - \frac{2x \text{Chi}(a + bx) \sinh(a + bx)}{b^2} \\
 &\quad + \frac{2 \int \text{Chi}(a + bx) \sinh(a + bx) dx}{b^2} + \frac{2 \int \frac{x \cosh(a + bx) \sinh(a + bx)}{a + bx} dx}{b} \\
 &\quad - \int \left(-\frac{a \cosh^2(a + bx)}{b^2} + \frac{x \cosh^2(a + bx)}{b} + \frac{a^2 \cosh^2(a + bx)}{b^2(a + bx)} \right) dx \\
 &= \frac{2 \cosh(a + bx) \text{Chi}(a + bx)}{b^3} + \frac{x^2 \cosh(a + bx) \text{Chi}(a + bx)}{b} \\
 &\quad - \frac{2x \text{Chi}(a + bx) \sinh(a + bx)}{b^2} - \frac{2 \int \frac{\cosh^2(a + bx)}{a + bx} dx}{b^2} + \frac{a \int \cosh^2(a + bx) dx}{b^2} \\
 &\quad - \frac{a^2 \int \frac{\cosh^2(a + bx)}{a + bx} dx}{b^2} - \frac{\int x \cosh^2(a + bx) dx}{b} + \frac{\int \frac{x \sinh(2(a + bx))}{a + bx} dx}{b}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{\cosh^2(a+bx)}{4b^3} + \frac{2\cosh(a+bx)\text{Chi}(a+bx)}{b^3} + \frac{x^2\cosh(a+bx)\text{Chi}(a+bx)}{b} \\
&\quad + \frac{a\cosh(a+bx)\sinh(a+bx)}{2b^3} - \frac{x\cosh(a+bx)\sinh(a+bx)}{2b^2} \\
&\quad - \frac{2x\text{Chi}(a+bx)\sinh(a+bx)}{b^2} - \frac{2\int\left(\frac{1}{2(a+bx)} + \frac{\cosh(2a+2bx)}{2(a+bx)}\right)dx}{b^2} + \frac{a\int 1 dx}{2b^2} \\
&\quad - \frac{a^2\int\left(\frac{1}{2(a+bx)} + \frac{\cosh(2a+2bx)}{2(a+bx)}\right)dx}{b^2} - \frac{\int x dx}{2b} + \frac{\int\frac{x\sinh(2a+2bx)}{a+bx}dx}{b} \\
&= \frac{ax}{2b^2} - \frac{x^2}{4b} + \frac{\cosh^2(a+bx)}{4b^3} + \frac{2\cosh(a+bx)\text{Chi}(a+bx)}{b^3} \\
&\quad + \frac{x^2\cosh(a+bx)\text{Chi}(a+bx)}{b} - \frac{\log(a+bx)}{b^3} - \frac{a^2\log(a+bx)}{2b^3} \\
&\quad + \frac{a\cosh(a+bx)\sinh(a+bx)}{2b^3} - \frac{x\cosh(a+bx)\sinh(a+bx)}{2b^2} \\
&\quad - \frac{2x\text{Chi}(a+bx)\sinh(a+bx)}{b^2} - \frac{\int\frac{\cosh(2a+2bx)}{a+bx}dx}{b^2} \\
&\quad - \frac{a^2\int\frac{\cosh(2a+2bx)}{a+bx}dx}{2b^2} + \frac{\int\left(\frac{\sinh(2a+2bx)}{b} + \frac{a\sinh(2a+2bx)}{b(-a-bx)}\right)dx}{b} \\
&= \frac{ax}{2b^2} - \frac{x^2}{4b} + \frac{\cosh^2(a+bx)}{4b^3} + \frac{2\cosh(a+bx)\text{Chi}(a+bx)}{b^3} \\
&\quad + \frac{x^2\cosh(a+bx)\text{Chi}(a+bx)}{b} - \frac{\text{Chi}(2a+2bx)}{b^3} - \frac{a^2\text{Chi}(2a+2bx)}{2b^3} - \frac{\log(a+bx)}{b^3} \\
&\quad - \frac{a^2\log(a+bx)}{2b^3} + \frac{a\cosh(a+bx)\sinh(a+bx)}{2b^3} - \frac{x\cosh(a+bx)\sinh(a+bx)}{2b^2} \\
&\quad - \frac{2x\text{Chi}(a+bx)\sinh(a+bx)}{b^2} + \frac{\int\sinh(2a+2bx)dx}{b^2} + \frac{a\int\frac{\sinh(2a+2bx)}{-a-bx}dx}{b^2} \\
&= \frac{ax}{2b^2} - \frac{x^2}{4b} + \frac{\cosh^2(a+bx)}{4b^3} + \frac{\cosh(2a+2bx)}{2b^3} + \frac{2\cosh(a+bx)\text{Chi}(a+bx)}{b^3} \\
&\quad + \frac{x^2\cosh(a+bx)\text{Chi}(a+bx)}{b} - \frac{\text{Chi}(2a+2bx)}{b^3} - \frac{a^2\text{Chi}(2a+2bx)}{2b^3} \\
&\quad - \frac{\log(a+bx)}{b^3} - \frac{a^2\log(a+bx)}{2b^3} + \frac{a\cosh(a+bx)\sinh(a+bx)}{2b^3} \\
&\quad - \frac{x\cosh(a+bx)\sinh(a+bx)}{2b^2} - \frac{2x\text{Chi}(a+bx)\sinh(a+bx)}{b^2} - \frac{a\text{Shi}(2a+2bx)}{b^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.61

$$\int x^2 \text{Chi}(a + bx) \sinh(a + bx) dx = \frac{-4abx + 2b^2x^2 - 5 \cosh(2(a + bx)) + 4(2 + a^2) \text{Chi}(2(a + bx)) + 8 \log(a + bx) + 4a^2 \log(a + bx) - 8C}{-}$$

[In] Integrate[x^2*CoshIntegral[a + b*x]*Sinh[a + b*x],x]

[Out]
$$\frac{-1/8*(-4*a*b*x + 2*b^2*x^2 - 5*\text{Cosh}[2*(a + b*x)] + 4*(2 + a^2)*\text{CoshIntegral}[2*(a + b*x)] + 8*\text{Log}[a + b*x] + 4*a^2*\text{Log}[a + b*x] - 8*\text{CoshIntegral}[a + b*x]*((2 + b^2*x^2)*\text{Cosh}[a + b*x] - 2*b*x*\text{Sinh}[a + b*x]) - 2*a*\text{Sinh}[2*(a + b*x)] + 2*b*x*\text{Sinh}[2*(a + b*x)] + 8*a*\text{SinhIntegral}[2*(a + b*x)])}{b^3}$$

Maple [A] (verified)

Time = 1.74 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.90

method	result
derivativedivides	$\text{Chi}(bx+a) \left(\frac{a^2 \cosh(bx+a) - 2a((bx+a) \cosh(bx+a) - \sinh(bx+a)) + (bx+a)^2 \cosh(bx+a) - 2(bx+a) \sinh(bx+a) + 2 \cosh(bx+a)}{b^3} \right)$
default	$\text{Chi}(bx+a) \left(\frac{a^2 \cosh(bx+a) - 2a((bx+a) \cosh(bx+a) - \sinh(bx+a)) + (bx+a)^2 \cosh(bx+a) - 2(bx+a) \sinh(bx+a) + 2 \cosh(bx+a)}{b^3} \right)$

[In] int(x^2*Chi(b*x+a)*sinh(b*x+a),x,method=_RETURNVERBOSE)

[Out]
$$\frac{1}{b^3} * (\text{Chi}(b*x+a) * (a^2 * \cosh(b*x+a) - 2*a * ((b*x+a) * \cosh(b*x+a) - \sinh(b*x+a))) + (b*x+a)^2 * \cosh(b*x+a) - 2*(b*x+a) * \sinh(b*x+a) + 2 * \cosh(b*x+a)) - \frac{1}{2} * a^2 * \ln(b*x+a) - \frac{1}{2} * a^2 * \text{Chi}(2*b*x+2*a) + \cosh(b*x+a) * \sinh(b*x+a) * a + (b*x+a) * a - a * \text{Shi}(2*b*x+2*a) - \frac{1}{2} * (b*x+a) * \cosh(b*x+a) * \sinh(b*x+a) - \frac{1}{4} * (b*x+a)^2 + \frac{5}{4} * \cosh(b*x+a)^2 - \ln(b*x+a) - \text{Chi}(2*b*x+2*a))$$

Fricas [F]

$$\int x^2 \text{Chi}(a + bx) \sinh(a + bx) dx = \int x^2 \text{Chi}(bx + a) \sinh(bx + a) dx$$

[In] integrate(x^2*Chi(b*x+a)*sinh(b*x+a),x, algorithm="fricas")

[Out] integral(x^2*cosh_integral(b*x + a)*sinh(b*x + a), x)

Sympy [F]

$$\int x^2 \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \int x^2 \sinh(a + bx) \operatorname{Chi}(a + bx) dx$$

[In] `integrate(x**2*Chi(b*x+a)*sinh(b*x+a),x)`

[Out] `Integral(x**2*sinh(a + b*x)*Chi(a + b*x), x)`

Maxima [F]

$$\int x^2 \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \int x^2 \operatorname{Chi}(bx + a) \sinh(bx + a) dx$$

[In] `integrate(x^2*Chi(b*x+a)*sinh(b*x+a),x, algorithm="maxima")`

[Out] `integrate(x^2*Chi(b*x + a)*sinh(b*x + a), x)`

Giac [F]

$$\int x^2 \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \int x^2 \operatorname{Chi}(bx + a) \sinh(bx + a) dx$$

[In] `integrate(x^2*Chi(b*x+a)*sinh(b*x+a),x, algorithm="giac")`

[Out] `integrate(x^2*Chi(b*x + a)*sinh(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \int x^2 \operatorname{coshint}(a + bx) \sinh(a + bx) dx$$

[In] `int(x^2*coshint(a + b*x)*sinh(a + b*x),x)`

[Out] `int(x^2*coshint(a + b*x)*sinh(a + b*x), x)`

3.124 $\int x \operatorname{Chi}(a + bx) \sinh(a + bx) dx$

Optimal result	605
Rubi [A] (verified)	605
Mathematica [A] (verified)	608
Maple [A] (verified)	608
Fricas [F]	608
Sympy [F]	609
Maxima [F]	609
Giac [F]	609
Mupad [F(-1)]	609

Optimal result

Integrand size = 14, antiderivative size = 109

$$\int x \operatorname{Chi}(a + bx) \sinh(a + bx) dx = -\frac{x}{2b} + \frac{x \cosh(a + bx) \operatorname{Chi}(a + bx)}{b} + \frac{a \operatorname{Chi}(2a + 2bx)}{2b^2} + \frac{a \log(a + bx)}{2b^2} - \frac{\cosh(a + bx) \sinh(a + bx)}{2b^2} - \frac{\operatorname{Chi}(a + bx) \sinh(a + bx)}{b^2} + \frac{\operatorname{Shi}(2a + 2bx)}{2b^2}$$

[Out] $-1/2*x/b+1/2*a*\operatorname{Chi}(2*b*x+2*a)/b^2+x*\operatorname{Chi}(b*x+a)*\cosh(b*x+a)/b+1/2*a*\ln(b*x+a)/b^2+1/2*\operatorname{Shi}(2*b*x+2*a)/b^2-\operatorname{Chi}(b*x+a)*\sinh(b*x+a)/b^2-1/2*\cosh(b*x+a)*\sinh(b*x+a)/b^2$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {6684, 6874, 2715, 8, 3393, 3382, 6676, 5556, 12, 3379}

$$\int x \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \frac{a \operatorname{Chi}(2a + 2bx)}{2b^2} - \frac{\operatorname{Chi}(a + bx) \sinh(a + bx)}{b^2} + \frac{\operatorname{Shi}(2a + 2bx)}{2b^2} + \frac{a \log(a + bx)}{2b^2} - \frac{\sinh(a + bx) \cosh(a + bx)}{2b^2} + \frac{x \operatorname{Chi}(a + bx) \cosh(a + bx)}{b} - \frac{x}{2b}$$

[In] $\operatorname{Int}[x*\operatorname{CoshIntegral}[a + b*x]*\operatorname{Sinh}[a + b*x], x]$

```
[Out] -1/2*x/b + (x*Cosh[a + b*x]*CoshIntegral[a + b*x])/b + (a*CoshIntegral[2*a
+ 2*b*x])/(2*b^2) + (a*Log[a + b*x])/(2*b^2) - (Cosh[a + b*x]*Sinh[a + b*x]
)/(2*b^2) - (CoshIntegral[a + b*x]*Sinh[a + b*x])/b^2 + SinhIntegral[2*a +
2*b*x]/(2*b^2)
```

Rule 8

```
Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 2715

```
Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[
c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2
*n]
```

Rule 3379

```
Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbo
l] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f
, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbo
l] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz
}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3393

```
Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] := In
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 5556

```
Int[Cosh[(a_) + (b_)*(x_)]^(p_)*((c_) + (d_)*(x_))^(m_)*Sinh[(a_) +
(b_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

Rule 6676

```
Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
  Simp[Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
  b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6684

```
Int[CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.)
+ (b_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(CoshIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Cosh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*CoshIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x \cosh(a + bx) \text{Chi}(a + bx)}{b} - \frac{\int \cosh(a + bx) \text{Chi}(a + bx) dx}{b} - \int \frac{x \cosh^2(a + bx)}{a + bx} dx \\
&= \frac{x \cosh(a + bx) \text{Chi}(a + bx)}{b} - \frac{\text{Chi}(a + bx) \sinh(a + bx)}{b^2} \\
&\quad + \frac{\int \frac{\cosh(a + bx) \sinh(a + bx)}{a + bx} dx}{b} - \int \left(\frac{\cosh^2(a + bx)}{b} - \frac{a \cosh^2(a + bx)}{b(a + bx)} \right) dx \\
&= \frac{x \cosh(a + bx) \text{Chi}(a + bx)}{b} - \frac{\text{Chi}(a + bx) \sinh(a + bx)}{b^2} \\
&\quad - \frac{\int \cosh^2(a + bx) dx}{b} + \frac{\int \frac{\sinh(2a + 2bx)}{2(a + bx)} dx}{b} + \frac{a \int \frac{\cosh^2(a + bx)}{a + bx} dx}{b} \\
&= \frac{x \cosh(a + bx) \text{Chi}(a + bx)}{b} - \frac{\cosh(a + bx) \sinh(a + bx)}{2b^2} - \frac{\text{Chi}(a + bx) \sinh(a + bx)}{b^2} \\
&\quad - \frac{\int 1 dx}{2b} + \frac{\int \frac{\sinh(2a + 2bx)}{a + bx} dx}{2b} + \frac{a \int \left(\frac{1}{2(a + bx)} + \frac{\cosh(2a + 2bx)}{2(a + bx)} \right) dx}{b} \\
&= -\frac{x}{2b} + \frac{x \cosh(a + bx) \text{Chi}(a + bx)}{b} + \frac{a \log(a + bx)}{2b^2} - \frac{\cosh(a + bx) \sinh(a + bx)}{2b^2} \\
&\quad - \frac{\text{Chi}(a + bx) \sinh(a + bx)}{b^2} + \frac{\text{Shi}(2a + 2bx)}{2b^2} + \frac{a \int \frac{\cosh(2a + 2bx)}{a + bx} dx}{2b} \\
&= -\frac{x}{2b} + \frac{x \cosh(a + bx) \text{Chi}(a + bx)}{b} + \frac{a \text{Chi}(2a + 2bx)}{2b^2} + \frac{a \log(a + bx)}{2b^2} \\
&\quad - \frac{\cosh(a + bx) \sinh(a + bx)}{2b^2} - \frac{\text{Chi}(a + bx) \sinh(a + bx)}{b^2} + \frac{\text{Shi}(2a + 2bx)}{2b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.72

$$\int x \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \frac{-2bx + 2a \operatorname{Chi}(2(a + bx)) + 2a \log(a + bx) + 4 \operatorname{Chi}(a + bx)(bx \cosh(a + bx) - \sinh(a + bx)) - \sinh(2(a + bx))}{4b^2}$$

[In] Integrate[x*CoshIntegral[a + b*x]*Sinh[a + b*x],x]

[Out] (-2*b*x + 2*a*CoshIntegral[2*(a + b*x)] + 2*a*Log[a + b*x] + 4*CoshIntegral[a + b*x]*(b*x*Cosh[a + b*x] - Sinh[a + b*x]) - Sinh[2*(a + b*x)] + 2*SinhIntegral[2*(a + b*x)])/(4*b^2)

Maple [A] (verified)

Time = 1.10 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.89

method	result
derivativedivides	$\frac{\operatorname{Chi}(bx+a)(-a \cosh(bx+a) + (bx+a) \cosh(bx+a) - \sinh(bx+a)) + a \left(\frac{\ln(bx+a)}{2} + \frac{\operatorname{Chi}(2bx+2a)}{2} \right) - \frac{\cosh(bx+a) \sinh(bx+a)}{2} - \frac{bx}{2}}{b^2}$
default	$\frac{\operatorname{Chi}(bx+a)(-a \cosh(bx+a) + (bx+a) \cosh(bx+a) - \sinh(bx+a)) + a \left(\frac{\ln(bx+a)}{2} + \frac{\operatorname{Chi}(2bx+2a)}{2} \right) - \frac{\cosh(bx+a) \sinh(bx+a)}{2} - \frac{bx}{2}}{b^2}$

[In] int(x*Chi(b*x+a)*sinh(b*x+a),x,method=_RETURNVERBOSE)

[Out] 1/b^2*(Chi(b*x+a)*(-a*cosh(b*x+a)+(b*x+a)*cosh(b*x+a)-sinh(b*x+a))+a*(1/2*ln(b*x+a)+1/2*Chi(2*b*x+2*a))-1/2*cosh(b*x+a)*sinh(b*x+a)-1/2*b*x-1/2*a+1/2*Shi(2*b*x+2*a))

Fricas [F]

$$\int x \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \int x \operatorname{Chi}(bx + a) \sinh(bx + a) dx$$

[In] integrate(x*Chi(b*x+a)*sinh(b*x+a),x, algorithm="fricas")

[Out] integral(x*cosh_integral(b*x + a)*sinh(b*x + a), x)

Sympy [F]

$$\int x \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \int x \sinh(a + bx) \operatorname{Chi}(a + bx) dx$$

[In] `integrate(x*Chi(b*x+a)*sinh(b*x+a),x)`

[Out] `Integral(x*sinh(a + b*x)*Chi(a + b*x), x)`

Maxima [F]

$$\int x \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \int x \operatorname{Chi}(bx + a) \sinh(bx + a) dx$$

[In] `integrate(x*Chi(b*x+a)*sinh(b*x+a),x, algorithm="maxima")`

[Out] `integrate(x*Chi(b*x + a)*sinh(b*x + a), x)`

Giac [F]

$$\int x \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \int x \operatorname{Chi}(bx + a) \sinh(bx + a) dx$$

[In] `integrate(x*Chi(b*x+a)*sinh(b*x+a),x, algorithm="giac")`

[Out] `integrate(x*Chi(b*x + a)*sinh(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \int x \operatorname{coshint}(a + bx) \sinh(a + bx) dx$$

[In] `int(x*coshint(a + b*x)*sinh(a + b*x),x)`

[Out] `int(x*coshint(a + b*x)*sinh(a + b*x), x)`

3.125 $\int \text{Chi}(a + bx) \sinh(a + bx) dx$

Optimal result	610
Rubi [A] (verified)	610
Mathematica [A] (verified)	611
Maple [A] (verified)	611
Fricas [F]	612
Sympy [F]	612
Maxima [F]	612
Giac [F]	612
Mupad [F(-1)]	613

Optimal result

Integrand size = 13, antiderivative size = 46

$$\int \text{Chi}(a + bx) \sinh(a + bx) dx = \frac{\cosh(a + bx)\text{Chi}(a + bx)}{b} - \frac{\text{Chi}(2a + 2bx)}{2b} - \frac{\log(a + bx)}{2b}$$

[Out] $-1/2*\text{Chi}(2*b*x+2*a)/b+\text{Chi}(b*x+a)*\cosh(b*x+a)/b-1/2*\ln(b*x+a)/b$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {6682, 3393, 3382}

$$\int \text{Chi}(a + bx) \sinh(a + bx) dx = -\frac{\text{Chi}(2a + 2bx)}{2b} + \frac{\text{Chi}(a + bx) \cosh(a + bx)}{b} - \frac{\log(a + bx)}{2b}$$

[In] `Int[CoshIntegral[a + b*x]*Sinh[a + b*x],x]`

[Out] $(\text{Cosh}[a + b*x]*\text{CoshIntegral}[a + b*x])/b - \text{CoshIntegral}[2*a + 2*b*x]/(2*b) - \text{Log}[a + b*x]/(2*b)$

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol]
:> Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3393

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol]
:> Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f}
```

, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))

Rule 6682

```
Int[CoshIntegral[(c_.) + (d_.)*(x_)]*Sinh[(a_.) + (b_.)*(x_)], x_Symbol] :>
Simp[Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\cosh(a + bx)\text{Chi}(a + bx)}{b} - \int \frac{\cosh^2(a + bx)}{a + bx} dx \\
 &= \frac{\cosh(a + bx)\text{Chi}(a + bx)}{b} - \int \left(\frac{1}{2(a + bx)} + \frac{\cosh(2a + 2bx)}{2(a + bx)} \right) dx \\
 &= \frac{\cosh(a + bx)\text{Chi}(a + bx)}{b} - \frac{\log(a + bx)}{2b} - \frac{1}{2} \int \frac{\cosh(2a + 2bx)}{a + bx} dx \\
 &= \frac{\cosh(a + bx)\text{Chi}(a + bx)}{b} - \frac{\text{Chi}(2a + 2bx)}{2b} - \frac{\log(a + bx)}{2b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.98

$$\int \text{Chi}(a + bx) \sinh(a + bx) dx = \frac{\cosh(a + bx)\text{Chi}(a + bx)}{b} - \frac{\text{Chi}(2(a + bx))}{2b} - \frac{\log(a + bx)}{2b}$$

```
[In] Integrate[CoshIntegral[a + b*x]*Sinh[a + b*x],x]
```

```
[Out] (Cosh[a + b*x]*CoshIntegral[a + b*x])/b - CoshIntegral[2*(a + b*x)]/(2*b) -
Log[a + b*x]/(2*b)
```

Maple [A] (verified)

Time = 0.81 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$\frac{\text{Chi}(bx+a) \cosh(bx+a) - \frac{\ln(bx+a)}{2} - \frac{\text{Chi}(2bx+2a)}{2}}{b}$	38
default	$\frac{\text{Chi}(bx+a) \cosh(bx+a) - \frac{\ln(bx+a)}{2} - \frac{\text{Chi}(2bx+2a)}{2}}{b}$	38

```
[In] int(Chi(b*x+a)*sinh(b*x+a),x,method=_RETURNVERBOSE)
```

```
[Out] 1/b*(Chi(b*x+a)*cosh(b*x+a)-1/2*ln(b*x+a)-1/2*Chi(2*b*x+2*a))
```

Fricas [F]

$$\int \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \int \operatorname{Chi}(bx + a) \sinh(bx + a) dx$$

[In] integrate(Chi(b*x+a)*sinh(b*x+a),x, algorithm="fricas")

[Out] integral(cosh_integral(b*x + a)*sinh(b*x + a), x)

Sympy [F]

$$\int \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \int \sinh(a + bx) \operatorname{Chi}(a + bx) dx$$

[In] integrate(Chi(b*x+a)*sinh(b*x+a),x)

[Out] Integral(sinh(a + b*x)*Chi(a + b*x), x)

Maxima [F]

$$\int \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \int \operatorname{Chi}(bx + a) \sinh(bx + a) dx$$

[In] integrate(Chi(b*x+a)*sinh(b*x+a),x, algorithm="maxima")

[Out] integrate(Chi(b*x + a)*sinh(b*x + a), x)

Giac [F]

$$\int \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \int \operatorname{Chi}(bx + a) \sinh(bx + a) dx$$

[In] integrate(Chi(b*x+a)*sinh(b*x+a),x, algorithm="giac")

[Out] integrate(Chi(b*x + a)*sinh(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int \operatorname{Chi}(a + bx) \sinh(a + bx) dx = \int \operatorname{coshint}(a + bx) \sinh(a + bx) dx$$

```
[In] int(coshint(a + b*x)*sinh(a + b*x),x)
```

```
[Out] int(coshint(a + b*x)*sinh(a + b*x), x)
```

3.126 $\int \frac{\mathbf{Chi}(a+bx) \sinh(a+bx)}{x} dx$

Optimal result	614
Rubi [N/A]	614
Mathematica [N/A]	615
Maple [N/A] (verified)	615
Fricas [N/A]	615
Sympy [N/A]	615
Maxima [N/A]	616
Giac [N/A]	616
Mupad [N/A]	616

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\mathbf{Chi}(a+bx) \sinh(a+bx)}{x} dx = \text{Int}\left(\frac{\mathbf{Chi}(a+bx) \sinh(a+bx)}{x}, x\right)$$

[Out] CannotIntegrate(Chi(b*x+a)*sinh(b*x+a)/x,x)

Rubi [N/A]

Not integrable

Time = 0.10 (sec), antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\mathbf{Chi}(a+bx) \sinh(a+bx)}{x} dx = \int \frac{\mathbf{Chi}(a+bx) \sinh(a+bx)}{x} dx$$

[In] Int[(CoshIntegral[a + b*x]*Sinh[a + b*x])/x,x]

[Out] Defer[Int] [(CoshIntegral[a + b*x]*Sinh[a + b*x])/x, x]

Rubi steps

$$\text{integral} = \int \frac{\mathbf{Chi}(a+bx) \sinh(a+bx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 0.50 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\text{Chi}(a + bx) \sinh(a + bx)}{x} dx = \int \frac{\text{Chi}(a + bx) \sinh(a + bx)}{x} dx$$

[In] Integrate[(CoshIntegral[a + b*x]*Sinh[a + b*x])/x,x]

[Out] Integrate[(CoshIntegral[a + b*x]*Sinh[a + b*x])/x, x]

Maple [N/A] (verified)

Not integrable

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx + a) \sinh(bx + a)}{x} dx$$

[In] int(Chi(b*x+a)*sinh(b*x+a)/x,x)

[Out] int(Chi(b*x+a)*sinh(b*x+a)/x,x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\text{Chi}(a + bx) \sinh(a + bx)}{x} dx = \int \frac{\text{Chi}(bx + a) \sinh(bx + a)}{x} dx$$

[In] integrate(Chi(b*x+a)*sinh(b*x+a)/x,x, algorithm="fricas")

[Out] integral(cosh_integral(b*x + a)*sinh(b*x + a)/x, x)

Sympy [N/A]

Not integrable

Time = 1.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{\text{Chi}(a + bx) \sinh(a + bx)}{x} dx = \int \frac{\sinh(a + bx) \text{Chi}(a + bx)}{x} dx$$

[In] integrate(Chi(b*x+a)*sinh(b*x+a)/x,x)

[Out] Integral(sinh(a + b*x)*Chi(a + b*x)/x, x)

Maxima [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\text{Chi}(a + bx) \sinh(a + bx)}{x} dx = \int \frac{\text{Chi}(bx + a) \sinh(bx + a)}{x} dx$$

[In] integrate(Chi(b*x+a)*sinh(b*x+a)/x,x, algorithm="maxima")

[Out] integrate(Chi(b*x + a)*sinh(b*x + a)/x, x)

Giac [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\text{Chi}(a + bx) \sinh(a + bx)}{x} dx = \int \frac{\text{Chi}(bx + a) \sinh(bx + a)}{x} dx$$

[In] integrate(Chi(b*x+a)*sinh(b*x+a)/x,x, algorithm="giac")

[Out] integrate(Chi(b*x + a)*sinh(b*x + a)/x, x)

Mupad [N/A]

Not integrable

Time = 5.09 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\text{Chi}(a + bx) \sinh(a + bx)}{x} dx = \int \frac{\text{coshint}(a + bx) \sinh(a + bx)}{x} dx$$

[In] int((coshint(a + b*x)*sinh(a + b*x))/x,x)

[Out] int((coshint(a + b*x)*sinh(a + b*x))/x, x)

3.127 $\int x^2 \cosh(a + bx) \mathbf{Chi}(a + bx) dx$

Optimal result	617
Rubi [A] (verified)	617
Mathematica [A] (verified)	621
Maple [A] (verified)	622
Fricas [F]	622
Sympy [F]	622
Maxima [F]	623
Giac [F]	623
Mupad [F(-1)]	623

Optimal result

Integrand size = 16, antiderivative size = 186

$$\int x^2 \cosh(a + bx) \mathbf{Chi}(a + bx) dx = \frac{x}{b^2} + \frac{a \cosh(2a + 2bx)}{4b^3} - \frac{x \cosh(2a + 2bx)}{4b^2} - \frac{2x \cosh(a + bx) \mathbf{Chi}(a + bx)}{b^2} - \frac{a \mathbf{Chi}(2a + 2bx)}{b^3} - \frac{a \log(a + bx)}{b^3} + \frac{\cosh(a + bx) \sinh(a + bx)}{b^3} + \frac{2 \mathbf{Chi}(a + bx) \sinh(a + bx)}{b^3} + \frac{x^2 \mathbf{Chi}(a + bx) \sinh(a + bx)}{b} + \frac{\sinh(2a + 2bx)}{8b^3} - \frac{\mathbf{Shi}(2a + 2bx)}{b^3} - \frac{a^2 \mathbf{Shi}(2a + 2bx)}{2b^3}$$

[Out] $x/b^2 - a \mathbf{Chi}(2bx + 2a)/b^3 - 2x \mathbf{Chi}(bx + a) \cosh(bx + a)/b^2 + 1/4 a \cosh(2bx + 2a)/b^3 - 1/4 x \cosh(2bx + 2a)/b^2 - a \ln(bx + a)/b^3 - \mathbf{Shi}(2bx + 2a)/b^3 - 1/2 a^2 \mathbf{Shi}(2bx + 2a)/b^3 + 2 \mathbf{Chi}(bx + a) \sinh(bx + a)/b^3 + x^2 \mathbf{Chi}(bx + a) \sinh(bx + a)/b + \cosh(bx + a) \sinh(bx + a)/b^3 + 1/8 \sinh(2bx + 2a)/b^3$

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.00, number of steps used = 21, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {6678, 5736, 6873, 6874, 2718, 3377, 2717, 3379, 6684, 2715, 8, 3393, 3382, 6676,

5556, 12}

$$\begin{aligned}
\int x^2 \cosh(a + bx) \operatorname{Chi}(a + bx) dx = & -\frac{a^2 \operatorname{Shi}(2a + 2bx)}{2b^3} - \frac{a \operatorname{Chi}(2a + 2bx)}{b^3} \\
& + \frac{2 \operatorname{Chi}(a + bx) \sinh(a + bx)}{b^3} - \frac{\operatorname{Shi}(2a + 2bx)}{b^3} \\
& - \frac{a \log(a + bx)}{b^3} + \frac{\sinh(2a + 2bx)}{8b^3} \\
& + \frac{a \cosh(2a + 2bx)}{4b^3} + \frac{\sinh(a + bx) \cosh(a + bx)}{b^3} \\
& - \frac{2x \operatorname{Chi}(a + bx) \cosh(a + bx)}{b^2} - \frac{x \cosh(2a + 2bx)}{4b^2} \\
& + \frac{x^2 \operatorname{Chi}(a + bx) \sinh(a + bx)}{b} + \frac{x}{b^2}
\end{aligned}$$

```
[In] Int[x^2*Cosh[a + b*x]*CoshIntegral[a + b*x],x]
```

```
[Out] x/b^2 + (a*Cosh[2*a + 2*b*x])/(4*b^3) - (x*Cosh[2*a + 2*b*x])/(4*b^2) - (2*
x*Cosh[a + b*x]*CoshIntegral[a + b*x])/b^2 - (a*CoshIntegral[2*a + 2*b*x])/
b^3 - (a*Log[a + b*x])/b^3 + (Cosh[a + b*x]*Sinh[a + b*x])/b^3 + (2*CoshInt
egral[a + b*x]*Sinh[a + b*x])/b^3 + (x^2*CoshIntegral[a + b*x]*Sinh[a + b*x
])/b + Sinh[2*a + 2*b*x]/(8*b^3) - SinhIntegral[2*a + 2*b*x]/b^3 - (a^2*Sinh
Integral[2*a + 2*b*x])/(2*b^3)
```

Rule 8

```
Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Sine[c + d*x])^(n - 1)/(d*n), x] + Dist[b^2*((n - 1)/n), Int[(b*Sine[
c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2
*n]
```

Rule 2717

```
Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]
```

Rule 2718

$\text{Int}[\sin[(c_.) + (d_.)(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Cos}[c + d*x]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 3377

$\text{Int}[\left((c_.) + (d_.)(x_.)\right)^{(m_.)} \sin[(e_.) + (f_.)(x_.)], x_Symbol] \rightarrow \text{Simp}[\left(-\left(c + d*x\right)^m \text{Cos}[e + f*x]/f\right), x] + \text{Dist}[d*(m/f), \text{Int}[\left(c + d*x\right)^{(m-1)} \text{Cos}[e + f*x], x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \ \&\& \ \text{GtQ}[m, 0]$

Rule 3379

$\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])*(f_.)(x_.)]/((c_.) + (d_.)(x_.)), x_Symbol] \rightarrow \text{Simp}[I*(\text{SinhIntegral}[c*f*(fz/d) + f*fz*x]/d), x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \ \&\& \ \text{EqQ}[d*e - c*f*fz*I, 0]$

Rule 3382

$\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])*(f_.)(x_.)]/((c_.) + (d_.)(x_.)), x_Symbol] \rightarrow \text{Simp}[\text{CoshIntegral}[c*f*(fz/d) + f*fz*x]/d, x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \ \&\& \ \text{EqQ}[d*(e - \text{Pi}/2) - c*f*fz*I, 0]$

Rule 3393

$\text{Int}[\left((c_.) + (d_.)(x_.)\right)^{(m_.)} \sin[(e_.) + (f_.)(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandTrigReduce}[\left(c + d*x\right)^m, \text{Sin}[e + f*x]^n, x], x] /; \text{FreeQ}[\{c, d, e, f, m\}, x] \ \&\& \ \text{IGtQ}[n, 1] \ \&\& \ (\text{!RationalQ}[m] \ || \ (\text{GeQ}[m, -1] \ \&\& \ \text{LtQ}[m, 1]))$

Rule 5556

$\text{Int}[\text{Cosh}[(a_.) + (b_.)(x_.)]^{(p_.)} \left((c_.) + (d_.)(x_.)\right)^{(m_.)} \text{Sinh}[(a_.) + (b_.)(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandTrigReduce}[\left(c + d*x\right)^m, \text{Sinh}[a + b*x]^n * \text{Cosh}[a + b*x]^p, x], x] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IGtQ}[p, 0]$

Rule 5736

$\text{Int}[\text{Cosh}[w_]^{(p_.)} (u_.) \text{Sinh}[v_]^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[1/2^p, \text{Int}[u * \text{Sinh}[2*v]^p, x], x] /; \text{EqQ}[w, v] \ \&\& \ \text{IntegerQ}[p]$

Rule 6676

$\text{Int}[\text{Cosh}[(a_.) + (b_.)(x_.)] * \text{CoshIntegral}[(c_.) + (d_.)(x_.)], x_Symbol] \rightarrow \text{Simp}[\text{Sinh}[a + b*x] * (\text{CoshIntegral}[c + d*x]/b), x] - \text{Dist}[d/b, \text{Int}[\text{Sinh}[a + b*x] * (\text{Cosh}[c + d*x]/(c + d*x)), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x]$

Rule 6678

```
Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)
)*(x_)^(m_.), x_Symbol] := Simp[(e + f*x)^m*Sinh[a + b*x]*(CoshIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Cosh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*CoshIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6684

```
Int[CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_)^(m_.)*Sinh[(a_.)
+ (b_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(CoshIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Cosh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*CoshIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6873

```
Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v !=
= u]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x^2 \text{Chi}(a + bx) \sinh(a + bx)}{b} - \frac{2 \int x \text{Chi}(a + bx) \sinh(a + bx) dx}{b} \\
&\quad - \int \frac{x^2 \cosh(a + bx) \sinh(a + bx)}{a + bx} dx \\
&= -\frac{2x \cosh(a + bx) \text{Chi}(a + bx)}{b^2} + \frac{x^2 \text{Chi}(a + bx) \sinh(a + bx)}{b} \\
&\quad - \frac{1}{2} \int \frac{x^2 \sinh(2(a + bx))}{a + bx} dx + \frac{2 \int \cosh(a + bx) \text{Chi}(a + bx) dx}{b^2} + \frac{2 \int \frac{x \cosh^2(a + bx)}{a + bx} dx}{b} \\
&= -\frac{2x \cosh(a + bx) \text{Chi}(a + bx)}{b^2} + \frac{2 \text{Chi}(a + bx) \sinh(a + bx)}{b^3} \\
&\quad + \frac{x^2 \text{Chi}(a + bx) \sinh(a + bx)}{b} - \frac{1}{2} \int \frac{x^2 \sinh(2a + 2bx)}{a + bx} dx \\
&\quad - \frac{2 \int \frac{\cosh(a + bx) \sinh(a + bx)}{a + bx} dx}{b^2} + \frac{2 \int \left(\frac{\cosh^2(a + bx)}{b} - \frac{a \cosh^2(a + bx)}{b(a + bx)} \right) dx}{b}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2x \cosh(a+bx)\text{Chi}(a+bx)}{b^2} + \frac{2\text{Chi}(a+bx)\sinh(a+bx)}{b^3} \\
&\quad + \frac{x^2\text{Chi}(a+bx)\sinh(a+bx)}{b} \\
&\quad - \frac{1}{2} \int \left(-\frac{a \sinh(2a+2bx)}{b^2} + \frac{x \sinh(2a+2bx)}{b} + \frac{a^2 \sinh(2a+2bx)}{b^2(a+bx)} \right) dx \\
&\quad + \frac{2 \int \cosh^2(a+bx) dx}{b^2} - \frac{2 \int \frac{\sinh(2a+2bx)}{2(a+bx)} dx}{b^2} - \frac{(2a) \int \frac{\cosh^2(a+bx)}{a+bx} dx}{b^2} \\
&= -\frac{2x \cosh(a+bx)\text{Chi}(a+bx)}{b^2} + \frac{\cosh(a+bx)\sinh(a+bx)}{b^3} + \frac{2\text{Chi}(a+bx)\sinh(a+bx)}{b^3} \\
&\quad + \frac{x^2\text{Chi}(a+bx)\sinh(a+bx)}{b} + \frac{\int 1 dx}{b^2} - \frac{\int \frac{\sinh(2a+2bx)}{a+bx} dx}{b^2} + \frac{a \int \sinh(2a+2bx) dx}{2b^2} \\
&\quad - \frac{(2a) \int \left(\frac{1}{2(a+bx)} + \frac{\cosh(2a+2bx)}{2(a+bx)} \right) dx}{b^2} - \frac{a^2 \int \frac{\sinh(2a+2bx)}{a+bx} dx}{2b^2} - \frac{\int x \sinh(2a+2bx) dx}{2b} \\
&= \frac{x}{b^2} + \frac{a \cosh(2a+2bx)}{4b^3} - \frac{x \cosh(2a+2bx)}{4b^2} - \frac{2x \cosh(a+bx)\text{Chi}(a+bx)}{b^2} - \frac{a \log(a+bx)}{b^3} \\
&\quad + \frac{\cosh(a+bx)\sinh(a+bx)}{b^3} + \frac{2\text{Chi}(a+bx)\sinh(a+bx)}{b^3} + \frac{x^2\text{Chi}(a+bx)\sinh(a+bx)}{b} \\
&\quad - \frac{\text{Shi}(2a+2bx)}{b^3} - \frac{a^2\text{Shi}(2a+2bx)}{2b^3} + \frac{\int \cosh(2a+2bx) dx}{4b^2} - \frac{a \int \frac{\cosh(2a+2bx)}{a+bx} dx}{b^2} \\
&= \frac{x}{b^2} + \frac{a \cosh(2a+2bx)}{4b^3} - \frac{x \cosh(2a+2bx)}{4b^2} - \frac{2x \cosh(a+bx)\text{Chi}(a+bx)}{b^2} \\
&\quad - \frac{a\text{Chi}(2a+2bx)}{b^3} - \frac{a \log(a+bx)}{b^3} + \frac{\cosh(a+bx)\sinh(a+bx)}{b^3} \\
&\quad + \frac{2\text{Chi}(a+bx)\sinh(a+bx)}{b^3} + \frac{x^2\text{Chi}(a+bx)\sinh(a+bx)}{b} \\
&\quad + \frac{\sinh(2a+2bx)}{8b^3} - \frac{\text{Shi}(2a+2bx)}{b^3} - \frac{a^2\text{Shi}(2a+2bx)}{2b^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.66

$$\int x^2 \cosh(a+bx)\text{Chi}(a+bx) dx$$

$$= \frac{8bx + 2a \cosh(2(a+bx)) - 2bx \cosh(2(a+bx)) - 8a\text{Chi}(2(a+bx)) - 8a \log(a+bx) + 8\text{Chi}(a+bx) (-2a^2 \text{Shi}(2(a+bx)) + 2a \cosh(2(a+bx)) - 2bx \cosh(2(a+bx)) - 8a\text{Chi}(2(a+bx)) - 8a \log(a+bx) + 8\text{Chi}(a+bx))}{8b^3}$$

[In] Integrate[x^2*Cosh[a + b*x]*CoshIntegral[a + b*x],x]

[Out] (8*b*x + 2*a*Cosh[2*(a + b*x)] - 2*b*x*Cosh[2*(a + b*x)] - 8*a*CoshIntegral[2*(a + b*x)] - 8*a*Log[a + b*x] + 8*CoshIntegral[a + b*x]*(-2*b*x*Cosh[a + b*x] + (2 + b^2*x^2)*Sinh[a + b*x]) + 5*Sinh[2*(a + b*x)] - 8*SinhIntegral[2*(a + b*x)] - 4*a^2*SinhIntegral[2*(a + b*x)])/(8*b^3)

Maple [A] (verified)

Time = 2.55 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.94

method	result
derivativedivides	$\frac{\text{Chi}(bx+a) \left(a^2 \sinh(bx+a) - 2a((bx+a) \sinh(bx+a) - \cosh(bx+a)) + (bx+a)^2 \sinh(bx+a) - 2(bx+a) \cosh(bx+a) + 2 \sinh(bx+a) \right)}{\dots}$
default	$\text{Chi}(bx+a) \left(a^2 \sinh(bx+a) - 2a((bx+a) \sinh(bx+a) - \cosh(bx+a)) + (bx+a)^2 \sinh(bx+a) - 2(bx+a) \cosh(bx+a) + 2 \sinh(bx+a) \right)$

[In] `int(x^2*Chi(b*x+a)*cosh(b*x+a),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{b^3} \left(\text{Chi}(bx+a) \left(a^2 \sinh(bx+a) - 2a((bx+a) \sinh(bx+a) - \cosh(bx+a)) + (bx+a)^2 \sinh(bx+a) - 2(bx+a) \cosh(bx+a) + 2 \sinh(bx+a) \right) - \frac{1}{2} a^2 \text{Shi}(2bx+2a) + a \cosh(bx+a)^2 - a \ln(bx+a) - a \text{Chi}(2bx+2a) - \frac{1}{2} (bx+a) \cosh(bx+a)^2 + \frac{5}{4} \cosh(bx+a) \sinh(bx+a) + \frac{5}{4} bx + \frac{5}{4} a - \text{Shi}(2bx+2a) \right)$$

Fricas [F]

$$\int x^2 \cosh(a + bx) \text{Chi}(a + bx) dx = \int x^2 \text{Chi}(bx + a) \cosh(bx + a) dx$$

[In] `integrate(x^2*Chi(b*x+a)*cosh(b*x+a),x, algorithm="fricas")`

[Out] `integral(x^2*cosh(b*x + a)*cosh_integral(b*x + a), x)`

Sympy [F]

$$\int x^2 \cosh(a + bx) \text{Chi}(a + bx) dx = \int x^2 \cosh(a + bx) \text{Chi}(a + bx) dx$$

[In] `integrate(x**2*Chi(b*x+a)*cosh(b*x+a),x)`

[Out] `Integral(x**2*cosh(a + b*x)*Chi(a + b*x), x)`

Maxima [F]

$$\int x^2 \cosh(a + bx) \operatorname{Chi}(a + bx) dx = \int x^2 \operatorname{Chi}(bx + a) \cosh(bx + a) dx$$

[In] integrate(x^2*Chi(b*x+a)*cosh(b*x+a),x, algorithm="maxima")

[Out] integrate(x^2*Chi(b*x + a)*cosh(b*x + a), x)

Giac [F]

$$\int x^2 \cosh(a + bx) \operatorname{Chi}(a + bx) dx = \int x^2 \operatorname{Chi}(bx + a) \cosh(bx + a) dx$$

[In] integrate(x^2*Chi(b*x+a)*cosh(b*x+a),x, algorithm="giac")

[Out] integrate(x^2*Chi(b*x + a)*cosh(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \cosh(a + bx) \operatorname{Chi}(a + bx) dx = \int x^2 \operatorname{coshint}(a + bx) \cosh(a + bx) dx$$

[In] int(x^2*coshint(a + b*x)*cosh(a + b*x),x)

[Out] int(x^2*coshint(a + b*x)*cosh(a + b*x), x)

3.128 $\int x \cosh(a + bx) \mathbf{Chi}(a + bx) dx$

Optimal result	624
Rubi [A] (verified)	624
Mathematica [A] (verified)	627
Maple [A] (verified)	627
Fricas [F]	627
Sympy [F]	628
Maxima [F]	628
Giac [F]	628
Mupad [F(-1)]	628

Optimal result

Integrand size = 14, antiderivative size = 97

$$\int x \cosh(a + bx) \mathbf{Chi}(a + bx) dx = -\frac{\cosh(2a + 2bx)}{4b^2} - \frac{\cosh(a + bx) \mathbf{Chi}(a + bx)}{b^2} + \frac{\mathbf{Chi}(2a + 2bx)}{2b^2} + \frac{\log(a + bx)}{2b^2} + \frac{x \mathbf{Chi}(a + bx) \sinh(a + bx)}{b} + \frac{a \mathbf{Shi}(2a + 2bx)}{2b^2}$$

[Out] $1/2*\mathbf{Chi}(2*b*x+2*a)/b^2 - \mathbf{Chi}(b*x+a)*\cosh(b*x+a)/b^2 - 1/4*\cosh(2*b*x+2*a)/b^2 + 1/2*\ln(b*x+a)/b^2 + 1/2*a*\mathbf{Shi}(2*b*x+2*a)/b^2 + x*\mathbf{Chi}(b*x+a)*\sinh(b*x+a)/b$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {6678, 5736, 6873, 6874, 2718, 3379, 6682, 3393, 3382}

$$\int x \cosh(a + bx) \mathbf{Chi}(a + bx) dx = \frac{\mathbf{Chi}(2a + 2bx)}{2b^2} - \frac{\mathbf{Chi}(a + bx) \cosh(a + bx)}{b^2} + \frac{a \mathbf{Shi}(2a + 2bx)}{2b^2} + \frac{\log(a + bx)}{2b^2} - \frac{\cosh(2a + 2bx)}{4b^2} + \frac{x \mathbf{Chi}(a + bx) \sinh(a + bx)}{b}$$

[In] `Int[x*Cosh[a + b*x]*CoshIntegral[a + b*x],x]`

[Out] $-1/4*\mathbf{Cosh}[2*a + 2*b*x]/b^2 - (\mathbf{Cosh}[a + b*x]*\mathbf{CoshIntegral}[a + b*x])/b^2 + \mathbf{CoshIntegral}[2*a + 2*b*x]/(2*b^2) + \mathbf{Log}[a + b*x]/(2*b^2) + (x*\mathbf{CoshIntegral}[a + b*x]*\mathbf{Sinh}[a + b*x])/b + (a*\mathbf{SinhIntegral}[2*a + 2*b*x])/(2*b^2)$

Rule 2718

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ
[{c, d}, x]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3393

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 5736

```
Int[Cosh[w_]^(p_.)*(u_.)*Sinh[v_]^(p_.), x_Symbol] := Dist[1/2^p, Int[u*Sinh[2*v]^p, x], x] /; EqQ[w, v] && IntegerQ[p]
```

Rule 6678

```
Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^m*Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*CoshIntegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6682

```
Int[CoshIntegral[(c_.) + (d_.)*(x_)]*Sinh[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a + b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6873

```
Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v != u]
```

Rule 6874

`Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]`
`]`

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x\text{Chi}(a+bx)\sinh(a+bx)}{b} - \frac{\int \text{Chi}(a+bx)\sinh(a+bx) dx}{b} \\
&\quad - \int \frac{x \cosh(a+bx)\sinh(a+bx)}{a+bx} dx \\
&= -\frac{\cosh(a+bx)\text{Chi}(a+bx)}{b^2} + \frac{x\text{Chi}(a+bx)\sinh(a+bx)}{b} \\
&\quad - \frac{1}{2} \int \frac{x \sinh(2(a+bx))}{a+bx} dx + \frac{\int \frac{\cosh^2(a+bx)}{a+bx} dx}{b} \\
&= -\frac{\cosh(a+bx)\text{Chi}(a+bx)}{b^2} + \frac{x\text{Chi}(a+bx)\sinh(a+bx)}{b} \\
&\quad - \frac{1}{2} \int \frac{x \sinh(2a+2bx)}{a+bx} dx + \frac{\int \left(\frac{1}{2(a+bx)} + \frac{\cosh(2a+2bx)}{2(a+bx)} \right) dx}{b} \\
&= -\frac{\cosh(a+bx)\text{Chi}(a+bx)}{b^2} + \frac{\log(a+bx)}{2b^2} + \frac{x\text{Chi}(a+bx)\sinh(a+bx)}{b} \\
&\quad - \frac{1}{2} \int \left(\frac{\sinh(2a+2bx)}{b} + \frac{a \sinh(2a+2bx)}{b(-a-bx)} \right) dx + \frac{\int \frac{\cosh(2a+2bx)}{a+bx} dx}{2b} \\
&= -\frac{\cosh(a+bx)\text{Chi}(a+bx)}{b^2} + \frac{\text{Chi}(2a+2bx)}{2b^2} + \frac{\log(a+bx)}{2b^2} \\
&\quad + \frac{x\text{Chi}(a+bx)\sinh(a+bx)}{b} - \frac{\int \sinh(2a+2bx) dx}{2b} - \frac{a \int \frac{\sinh(2a+2bx)}{-a-bx} dx}{2b} \\
&= -\frac{\cosh(2a+2bx)}{4b^2} - \frac{\cosh(a+bx)\text{Chi}(a+bx)}{b^2} + \frac{\text{Chi}(2a+2bx)}{2b^2} \\
&\quad + \frac{\log(a+bx)}{2b^2} + \frac{x\text{Chi}(a+bx)\sinh(a+bx)}{b} + \frac{a\text{Shi}(2a+2bx)}{2b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.75

$$\int x \cosh(a + bx) \operatorname{Chi}(a + bx) dx = \frac{-\cosh(2(a + bx)) + 2\operatorname{Chi}(2(a + bx)) + 2\log(a + bx) + 4\operatorname{Chi}(a + bx)(-\cosh(a + bx) + bx \sinh(a + bx))}{4b^2}$$

[In] Integrate[x*Cosh[a + b*x]*CoshIntegral[a + b*x],x]

[Out] (-Cosh[2*(a + b*x)] + 2*CoshIntegral[2*(a + b*x)] + 2*Log[a + b*x] + 4*CoshIntegral[a + b*x]*(-Cosh[a + b*x] + b*x*Sinh[a + b*x]) + 2*a*SinhIntegral[2*(a + b*x)])/(4*b^2)

Maple [A] (verified)

Time = 1.47 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.87

method	result
derivativedivides	$\frac{\operatorname{Chi}(bx+a)(-a \sinh(bx+a) + (bx+a) \sinh(bx+a) - \cosh(bx+a)) + \frac{a \operatorname{Shi}(2bx+2a)}{2} - \frac{\cosh(bx+a)^2}{2} + \frac{\ln(bx+a)}{2} + \frac{\operatorname{Chi}(2bx+2a)}{2}}{b^2}$
default	$\frac{\operatorname{Chi}(bx+a)(-a \sinh(bx+a) + (bx+a) \sinh(bx+a) - \cosh(bx+a)) + \frac{a \operatorname{Shi}(2bx+2a)}{2} - \frac{\cosh(bx+a)^2}{2} + \frac{\ln(bx+a)}{2} + \frac{\operatorname{Chi}(2bx+2a)}{2}}{b^2}$

[In] int(x*Chi(b*x+a)*cosh(b*x+a),x,method=_RETURNVERBOSE)

[Out] 1/b^2*(Chi(b*x+a)*(-a*sinh(b*x+a)+(b*x+a)*sinh(b*x+a)-cosh(b*x+a))+1/2*a*Shi(2*b*x+2*a)-1/2*cosh(b*x+a)^2+1/2*ln(b*x+a)+1/2*Chi(2*b*x+2*a))

Fricas [F]

$$\int x \cosh(a + bx) \operatorname{Chi}(a + bx) dx = \int x \operatorname{Chi}(bx + a) \cosh(bx + a) dx$$

[In] integrate(x*Chi(b*x+a)*cosh(b*x+a),x, algorithm="fricas")

[Out] integral(x*cosh(b*x + a)*cosh_integral(b*x + a), x)

Sympy [F]

$$\int x \cosh(a + bx) \operatorname{Chi}(a + bx) dx = \int x \cosh(a + bx) \operatorname{Chi}(a + bx) dx$$

[In] `integrate(x*Chi(b*x+a)*cosh(b*x+a),x)`

[Out] `Integral(x*cosh(a + b*x)*Chi(a + b*x), x)`

Maxima [F]

$$\int x \cosh(a + bx) \operatorname{Chi}(a + bx) dx = \int x \operatorname{Chi}(bx + a) \cosh(bx + a) dx$$

[In] `integrate(x*Chi(b*x+a)*cosh(b*x+a),x, algorithm="maxima")`

[Out] `integrate(x*Chi(b*x + a)*cosh(b*x + a), x)`

Giac [F]

$$\int x \cosh(a + bx) \operatorname{Chi}(a + bx) dx = \int x \operatorname{Chi}(bx + a) \cosh(bx + a) dx$$

[In] `integrate(x*Chi(b*x+a)*cosh(b*x+a),x, algorithm="giac")`

[Out] `integrate(x*Chi(b*x + a)*cosh(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x \cosh(a + bx) \operatorname{Chi}(a + bx) dx = \int x \operatorname{coshint}(a + bx) \cosh(a + bx) dx$$

[In] `int(x*coshint(a + b*x)*cosh(a + b*x),x)`

[Out] `int(x*coshint(a + b*x)*cosh(a + b*x), x)`

3.129 $\int \cosh(a + bx) \mathbf{Chi}(a + bx) dx$

Optimal result	629
Rubi [A] (verified)	629
Mathematica [A] (verified)	630
Maple [A] (verified)	630
Fricas [F]	631
Sympy [F]	631
Maxima [F]	631
Giac [F]	632
Mupad [F(-1)]	632

Optimal result

Integrand size = 13, antiderivative size = 33

$$\int \cosh(a + bx) \mathbf{Chi}(a + bx) dx = \frac{\mathbf{Chi}(a + bx) \sinh(a + bx)}{b} - \frac{\mathbf{Shi}(2a + 2bx)}{2b}$$

[Out] $-1/2*\mathbf{Shi}(2*b*x+2*a)/b+\mathbf{Chi}(b*x+a)*\sinh(b*x+a)/b$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {6676, 5556, 12, 3379}

$$\int \cosh(a + bx) \mathbf{Chi}(a + bx) dx = \frac{\mathbf{Chi}(a + bx) \sinh(a + bx)}{b} - \frac{\mathbf{Shi}(2a + 2bx)}{2b}$$

[In] `Int[Cosh[a + b*x]*CoshIntegral[a + b*x],x]`

[Out] `(CoshIntegral[a + b*x]*Sinh[a + b*x])/b - SinhIntegral[2*a + 2*b*x]/(2*b)`

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 3379

`Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

Rule 6676

```
Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
Simp[Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Chi}(a + bx) \sinh(a + bx)}{b} - \int \frac{\cosh(a + bx) \sinh(a + bx)}{a + bx} dx \\
&= \frac{\text{Chi}(a + bx) \sinh(a + bx)}{b} - \int \frac{\sinh(2a + 2bx)}{2(a + bx)} dx \\
&= \frac{\text{Chi}(a + bx) \sinh(a + bx)}{b} - \frac{1}{2} \int \frac{\sinh(2a + 2bx)}{a + bx} dx \\
&= \frac{\text{Chi}(a + bx) \sinh(a + bx)}{b} - \frac{\text{Shi}(2a + 2bx)}{2b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.97

$$\int \cosh(a + bx) \text{Chi}(a + bx) dx = \frac{\text{Chi}(a + bx) \sinh(a + bx)}{b} - \frac{\text{Shi}(2(a + bx))}{2b}$$

```
[In] Integrate[Cosh[a + b*x]*CoshIntegral[a + b*x], x]
```

```
[Out] (CoshIntegral[a + b*x]*Sinh[a + b*x])/b - SinhIntegral[2*(a + b*x)]/(2*b)
```

Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.91

method	result	size
derivativedivides	$\frac{\text{Chi}(bx+a) \sinh(bx+a) - \frac{\text{Shi}(2bx+2a)}{2}}{b}$	30
default	$\frac{\text{Chi}(bx+a) \sinh(bx+a) - \frac{\text{Shi}(2bx+2a)}{2}}{b}$	30

[In] `int(Chi(b*x+a)*cosh(b*x+a),x,method=_RETURNVERBOSE)`

[Out] `1/b*(Chi(b*x+a)*sinh(b*x+a)-1/2*Shi(2*b*x+2*a))`

Fricas [F]

$$\int \cosh(a + bx) \text{Chi}(a + bx) dx = \int \text{Chi}(bx + a) \cosh(bx + a) dx$$

[In] `integrate(Chi(b*x+a)*cosh(b*x+a),x, algorithm="fricas")`

[Out] `integral(cosh(b*x + a)*cosh_integral(b*x + a), x)`

Sympy [F]

$$\int \cosh(a + bx) \text{Chi}(a + bx) dx = \int \cosh(a + bx) \text{Chi}(a + bx) dx$$

[In] `integrate(Chi(b*x+a)*cosh(b*x+a),x)`

[Out] `Integral(cosh(a + b*x)*Chi(a + b*x), x)`

Maxima [F]

$$\int \cosh(a + bx) \text{Chi}(a + bx) dx = \int \text{Chi}(bx + a) \cosh(bx + a) dx$$

[In] `integrate(Chi(b*x+a)*cosh(b*x+a),x, algorithm="maxima")`

[Out] `integrate(Chi(b*x + a)*cosh(b*x + a), x)`

Giac [F]

$$\int \cosh(a + bx)\text{Chi}(a + bx) dx = \int \text{Chi}(bx + a) \cosh(bx + a) dx$$

[In] integrate(Chi(b*x+a)*cosh(b*x+a),x, algorithm="giac")

[Out] integrate(Chi(b*x + a)*cosh(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int \cosh(a + bx)\text{Chi}(a + bx) dx = \int \text{coshint}(a + bx) \cosh(a + bx) dx$$

[In] int(coshint(a + b*x)*cosh(a + b*x),x)

[Out] int(coshint(a + b*x)*cosh(a + b*x), x)

3.130 $\int \frac{\cosh(a+bx)\mathbf{Chi}(a+bx)}{x} dx$

Optimal result	633
Rubi [N/A]	633
Mathematica [N/A]	634
Maple [N/A] (verified)	634
Fricas [N/A]	634
Sympy [N/A]	634
Maxima [N/A]	635
Giac [N/A]	635
Mupad [N/A]	635

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\cosh(a+bx)\mathbf{Chi}(a+bx)}{x} dx = \text{Int}\left(\frac{\cosh(a+bx)\mathbf{Chi}(a+bx)}{x}, x\right)$$

[Out] CannotIntegrate(Chi(b*x+a)*cosh(b*x+a)/x,x)

Rubi [N/A]

Not integrable

Time = 0.06 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\cosh(a+bx)\mathbf{Chi}(a+bx)}{x} dx = \int \frac{\cosh(a+bx)\mathbf{Chi}(a+bx)}{x} dx$$

[In] Int[(Cosh[a + b*x]*CoshIntegral[a + b*x])/x,x]

[Out] Defer[Int] [(Cosh[a + b*x]*CoshIntegral[a + b*x])/x, x]

Rubi steps

$$\text{integral} = \int \frac{\cosh(a+bx)\mathbf{Chi}(a+bx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 0.38 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Chi}(a + bx)}{x} dx = \int \frac{\cosh(a + bx)\text{Chi}(a + bx)}{x} dx$$

[In] Integrate[(Cosh[a + b*x]*CoshIntegral[a + b*x])/x,x]

[Out] Integrate[(Cosh[a + b*x]*CoshIntegral[a + b*x])/x, x]

Maple [N/A] (verified)

Not integrable

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(bx + a) \cosh(bx + a)}{x} dx$$

[In] int(Chi(b*x+a)*cosh(b*x+a)/x,x)

[Out] int(Chi(b*x+a)*cosh(b*x+a)/x,x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Chi}(a + bx)}{x} dx = \int \frac{\text{Chi}(bx + a) \cosh(bx + a)}{x} dx$$

[In] integrate(Chi(b*x+a)*cosh(b*x+a)/x,x, algorithm="fricas")

[Out] integral(cosh(b*x + a)*cosh_integral(b*x + a)/x, x)

Sympy [N/A]

Not integrable

Time = 1.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{\cosh(a + bx)\text{Chi}(a + bx)}{x} dx = \int \frac{\cosh(a + bx)\text{Chi}(a + bx)}{x} dx$$

[In] integrate(Chi(b*x+a)*cosh(b*x+a)/x,x)

[Out] Integral(cosh(a + b*x)*Chi(a + b*x)/x, x)

Maxima [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Chi}(a + bx)}{x} dx = \int \frac{\text{Chi}(bx + a) \cosh(bx + a)}{x} dx$$

[In] integrate(Chi(b*x+a)*cosh(b*x+a)/x,x, algorithm="maxima")

[Out] integrate(Chi(b*x + a)*cosh(b*x + a)/x, x)

Giac [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Chi}(a + bx)}{x} dx = \int \frac{\text{Chi}(bx + a) \cosh(bx + a)}{x} dx$$

[In] integrate(Chi(b*x+a)*cosh(b*x+a)/x,x, algorithm="giac")

[Out] integrate(Chi(b*x + a)*cosh(b*x + a)/x, x)

Mupad [N/A]

Not integrable

Time = 5.15 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Chi}(a + bx)}{x} dx = \int \frac{\coshint(a + bx) \cosh(a + bx)}{x} dx$$

[In] int((coshint(a + b*x)*cosh(a + b*x))/x,x)

[Out] int((coshint(a + b*x)*cosh(a + b*x))/x, x)

3.131 $\int x \mathbf{Chi}(c + dx) \sinh(a + bx) dx$

Optimal result	636
Rubi [A] (verified)	637
Mathematica [A] (verified)	641
Maple [F]	642
Fricas [F]	642
Sympy [F]	642
Maxima [F]	642
Giac [F]	643
Mupad [F(-1)]	643

Optimal result

Integrand size = 14, antiderivative size = 371

$$\begin{aligned}
 \int x \mathbf{Chi}(c + dx) \sinh(a + bx) dx = & \frac{c \cosh\left(a - \frac{bc}{d}\right) \mathbf{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2bd} \\
 & + \frac{x \cosh(a + bx) \mathbf{Chi}(c + dx)}{b} \\
 & + \frac{c \cosh\left(a - \frac{bc}{d}\right) \mathbf{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2bd} \\
 & + \frac{\mathbf{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b^2} \\
 & + \frac{\mathbf{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b^2} \\
 & - \frac{\mathbf{Chi}(c + dx) \sinh(a + bx)}{b^2} \\
 & - \frac{\sinh(a - c + (b-d)x)}{2b(b-d)} - \frac{\sinh(a + c + (b+d)x)}{2b(b+d)} \\
 & + \frac{\cosh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b^2} \\
 & + \frac{c \sinh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2bd} \\
 & + \frac{\cosh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b^2} \\
 & + \frac{c \sinh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2bd}
 \end{aligned}$$

[Out] $\frac{1}{2}c \operatorname{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right) \operatorname{cosh}\left(\frac{a-bc}{d}\right) / \frac{b}{d} + \frac{1}{2}c \operatorname{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right) \operatorname{cosh}\left(\frac{a-bc}{d}\right) / \frac{b}{d} + x \operatorname{Chi}(dx+c) \operatorname{cosh}(bx+a) / b + \frac{1}{2} \operatorname{cosh}\left(\frac{a-bc}{d}\right) \operatorname{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right) / b^2 + \frac{1}{2} \operatorname{cosh}\left(\frac{a-bc}{d}\right) \operatorname{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right) / b^2 + \frac{1}{2} \operatorname{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right) \operatorname{sinh}\left(\frac{a-bc}{d}\right) / b^2 + \frac{1}{2} \operatorname{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right) \operatorname{sinh}\left(\frac{a-bc}{d}\right) / b^2 + \frac{1}{2}c \operatorname{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right) \operatorname{sinh}\left(\frac{a-bc}{d}\right) / \frac{b}{d} + \frac{1}{2}c \operatorname{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right) \operatorname{sinh}\left(\frac{a-bc}{d}\right) / \frac{b}{d} - \operatorname{Chi}(dx+c) \operatorname{sinh}(bx+a) / b^2 - \frac{1}{2} \operatorname{sinh}\left(\frac{a-c+(b-d)x}{b}\right) / (b-d) - \frac{1}{2} \operatorname{sinh}\left(\frac{a+c+(b+d)x}{b}\right) / (b+d)$

Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 371, normalized size of antiderivative = 1.00, number of steps used = 24, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {6684, 5762, 6874, 2717, 3384, 3379, 3382, 6676, 5580}

$$\int x \operatorname{Chi}(c+dx) \operatorname{sinh}(a+bx) dx = \frac{\operatorname{sinh}\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2b^2} + \frac{\operatorname{sinh}\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2b^2} - \frac{\operatorname{sinh}(a+bx) \operatorname{Chi}(c+dx)}{b^2} + \frac{\operatorname{cosh}\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2b^2} + \frac{\operatorname{cosh}\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2b^2} + \frac{c \operatorname{cosh}\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2bd} + \frac{x \operatorname{cosh}(a+bx) \operatorname{Chi}(c+dx)}{b} + \frac{c \operatorname{cosh}\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2bd} + \frac{c \operatorname{sinh}\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2bd} + \frac{c \operatorname{sinh}\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2bd} - \frac{\operatorname{sinh}(a+x(b-d)-c)}{2b(b-d)} - \frac{\operatorname{sinh}(a+x(b+d)+c)}{2b(b+d)}$$

[In] Int[x*CoshIntegral[c+d*x]*Sinh[a+b*x],x]

[Out] $(c \operatorname{Cosh}[a - (b*c)/d] \operatorname{CoshIntegral}[(c*(b-d))/d + (b-d)*x]) / (2*b*d) + (x \operatorname{Cosh}[a + b*x] \operatorname{CoshIntegral}[c + d*x]) / b + (c \operatorname{Cosh}[a - (b*c)/d] \operatorname{CoshIntegral}[$

$$\begin{aligned} & (c*(b + d))/d + (b + d)*x]/(2*b*d) + (\text{CoshIntegral}[(c*(b - d))/d + (b - d) \\ & *x]*\text{Sinh}[a - (b*c)/d])/(2*b^2) + (\text{CoshIntegral}[(c*(b + d))/d + (b + d)*x]*\text{S} \\ & \text{inh}[a - (b*c)/d])/(2*b^2) - (\text{CoshIntegral}[c + d*x]*\text{Sinh}[a + b*x])/b^2 - \text{Sinh}[a - c + (b - d)*x]/(2*b*(b - d)) - \text{Sinh}[a + c + (b + d)*x]/(2*b*(b + d)) \\ & + (\text{Cosh}[a - (b*c)/d]*\text{SinhIntegral}[(c*(b - d))/d + (b - d)*x])/(2*b^2) + (c* \\ & \text{Sinh}[a - (b*c)/d]*\text{SinhIntegral}[(c*(b - d))/d + (b - d)*x])/(2*b*d) + (\text{Cosh}[\\ & a - (b*c)/d]*\text{SinhIntegral}[(c*(b + d))/d + (b + d)*x])/(2*b^2) + (c*\text{Sinh}[a - \\ & (b*c)/d]*\text{SinhIntegral}[(c*(b + d))/d + (b + d)*x])/(2*b*d) \end{aligned}$$
Rule 2717

```
Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3384

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]
```

Rule 5580

```
Int[Cosh[(c_.) + (d_.)*(x_)]^(q_.)*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(p_.), x_Symbol] := Int[ExpandTrigReduce[(e + f*x)^m, Sinh[a + b*x]^p*Cosh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[p, 0] && IGtQ[q, 0]
```

Rule 5762

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(m_.)*Cosh[(c_.) + (d_.)*(x_)]^(n_.)*(u_.), x_Symbol] := Int[ExpandTrigReduce[u, Cosh[a + b*x]^m*Cosh[c + d*x]^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 6676

```
Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)], x_Symbol] :=
  Simp[Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
  b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6684

```
Int[CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.)
+ (b_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*Cosh[a + b*x]*(CoshIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Cosh[a + b*x]*(Cosh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Cosh[a + b*x]*CoshIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x \cosh(a + bx) \text{Chi}(c + dx)}{b} - \frac{\int \cosh(a + bx) \text{Chi}(c + dx) dx}{b} \\
&\quad - \frac{d \int \frac{x \cosh(a + bx) \cosh(c + dx)}{c + dx} dx}{b} \\
&= \frac{x \cosh(a + bx) \text{Chi}(c + dx)}{b} - \frac{\text{Chi}(c + dx) \sinh(a + bx)}{b^2} \\
&\quad + \frac{d \int \frac{\cosh(c + dx) \sinh(a + bx)}{c + dx} dx}{b^2} - \frac{d \int \left(\frac{x \cosh(a - c + (b - d)x)}{2(c + dx)} + \frac{x \cosh(a + c + (b + d)x)}{2(c + dx)} \right) dx}{b} \\
&= \frac{x \cosh(a + bx) \text{Chi}(c + dx)}{b} - \frac{\text{Chi}(c + dx) \sinh(a + bx)}{b^2} \\
&\quad + \frac{d \int \left(\frac{\sinh(a - c + (b - d)x)}{2(c + dx)} + \frac{\sinh(a + c + (b + d)x)}{2(c + dx)} \right) dx}{b^2} \\
&\quad - \frac{d \int \frac{x \cosh(a - c + (b - d)x)}{c + dx} dx}{2b} - \frac{d \int \frac{x \cosh(a + c + (b + d)x)}{c + dx} dx}{2b} \\
&= \frac{x \cosh(a + bx) \text{Chi}(c + dx)}{b} - \frac{\text{Chi}(c + dx) \sinh(a + bx)}{b^2} + \frac{d \int \frac{\sinh(a - c + (b - d)x)}{c + dx} dx}{2b^2} \\
&\quad + \frac{d \int \frac{\sinh(a + c + (b + d)x)}{c + dx} dx}{2b^2} - \frac{d \int \left(\frac{\cosh(a - c + (b - d)x)}{d} - \frac{c \cosh(a - c + (b - d)x)}{d(c + dx)} \right) dx}{2b} \\
&\quad - \frac{d \int \left(\frac{\cosh(a + c + (b + d)x)}{d} - \frac{c \cosh(a + c + (b + d)x)}{d(c + dx)} \right) dx}{2b}
\end{aligned}$$

$$\begin{aligned}
&= \frac{x \cosh(a + bx) \operatorname{Chi}(c + dx)}{b} - \frac{\operatorname{Chi}(c + dx) \sinh(a + bx)}{b^2} - \frac{\int \cosh(a - c + (b - d)x) dx}{2b} \\
&\quad - \frac{\int \cosh(a + c + (b + d)x) dx}{2b} + \frac{c \int \frac{\cosh(a - c + (b - d)x)}{c + dx} dx}{2b} + \frac{c \int \frac{\cosh(a + c + (b + d)x)}{c + dx} dx}{2b} \\
&\quad + \frac{(d \cosh(a - \frac{bc}{d})) \int \frac{\sinh(\frac{c(b-d)}{d} + (b-d)x)}{c + dx} dx}{2b^2} + \frac{(d \cosh(a - \frac{bc}{d})) \int \frac{\sinh(\frac{c(b+d)}{d} + (b+d)x)}{c + dx} dx}{2b^2} \\
&\quad + \frac{(d \sinh(a - \frac{bc}{d})) \int \frac{\cosh(\frac{c(b-d)}{d} + (b-d)x)}{c + dx} dx}{2b^2} + \frac{(d \sinh(a - \frac{bc}{d})) \int \frac{\cosh(\frac{c(b+d)}{d} + (b+d)x)}{c + dx} dx}{2b^2} \\
&= \frac{x \cosh(a + bx) \operatorname{Chi}(c + dx)}{b} + \frac{\operatorname{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b^2} \\
&\quad + \frac{\operatorname{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b^2} - \frac{\operatorname{Chi}(c + dx) \sinh(a + bx)}{b^2} \\
&\quad - \frac{\sinh(a - c + (b - d)x)}{2b(b - d)} - \frac{\sinh(a + c + (b + d)x)}{2b(b + d)} \\
&\quad + \frac{\cosh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b^2} + \frac{\cosh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b^2} \\
&\quad + \frac{(c \cosh\left(a - \frac{bc}{d}\right)) \int \frac{\cosh\left(\frac{c(b-d)}{d} + (b-d)x\right)}{c + dx} dx}{2b} \\
&\quad + \frac{(c \cosh\left(a - \frac{bc}{d}\right)) \int \frac{\cosh\left(\frac{c(b+d)}{d} + (b+d)x\right)}{c + dx} dx}{2b} \\
&\quad + \frac{(c \sinh\left(a - \frac{bc}{d}\right)) \int \frac{\sinh\left(\frac{c(b-d)}{d} + (b-d)x\right)}{c + dx} dx}{2b} \\
&\quad + \frac{(c \sinh\left(a - \frac{bc}{d}\right)) \int \frac{\sinh\left(\frac{c(b+d)}{d} + (b+d)x\right)}{c + dx} dx}{2b}
\end{aligned}$$

$$\begin{aligned}
&= \frac{c \cosh\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2bd} + \frac{x \cosh(a+bx) \operatorname{Chi}(c+dx)}{b} \\
&+ \frac{c \cosh\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2bd} + \frac{\operatorname{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b^2} \\
&+ \frac{\operatorname{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b^2} - \frac{\operatorname{Chi}(c+dx) \sinh(a+bx)}{b^2} \\
&- \frac{\sinh(a-c+(b-d)x)}{2b(b-d)} - \frac{\sinh(a+c+(b+d)x)}{2b(b+d)} \\
&+ \frac{\cosh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b^2} + \frac{c \sinh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2bd} \\
&+ \frac{\cosh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b^2} + \frac{c \sinh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2bd}
\end{aligned}$$

Mathematica [A] (verified)

Time = 2.40 (sec) , antiderivative size = 322, normalized size of antiderivative = 0.87

$$\int x \operatorname{Chi}(c+dx) \sinh(a+bx) dx$$

$$e^{-a-c-(b+d)x} \left(bd(d(-1+e^{2(c+dx)}) + b(1+e^{2(c+dx)})) + (bc-d)(b^2-d^2) e^{\frac{(b+d)(c+dx)}{d}} \operatorname{ExpIntegralEi}\left(-\frac{(b-d)(c+dx)}{d}\right) \right)$$

[In] Integrate[x*CoshIntegral[c + d*x]*Sinh[a + b*x],x]

[Out] (E^(-a - c - (b + d)*x)*(b*d*(d*(-1 + E^(2*(c + d*x))) + b*(1 + E^(2*(c + d*x)))) + (b*c - d)*(b^2 - d^2)*E^(((b + d)*(c + d*x))/d)*ExpIntegralEi[-(((b - d)*(c + d*x))/d)] + (b*c - d)*(b^2 - d^2)*E^(((b + d)*(c + d*x))/d)*ExpIntegralEi[-(((b + d)*(c + d*x))/d)]) + E^(a - (c*(b + d))/d)*(-(b*d*E^((b*c)/d + b*x - d*x)*(b + d + b*E^(2*(c + d*x)) - d*E^(2*(c + d*x)))) + (b*c + d)*(b^2 - d^2)*E^c*ExpIntegralEi[((b - d)*(c + d*x))/d] + (b*c + d)*(b^2 - d^2)*E^c*ExpIntegralEi[((b + d)*(c + d*x))/d]) + 4*(b - d)*d*(b + d)*CoshIntegral[c + d*x]*(b*x*Cosh[a + b*x] - Sinh[a + b*x]))/(4*b^2*(b - d)*d*(b + d))

Maple [F]

$$\int x \operatorname{Chi}(dx + c) \sinh(bx + a) dx$$

```
[In] int(x*Chi(d*x+c)*sinh(b*x+a),x)
```

```
[Out] int(x*Chi(d*x+c)*sinh(b*x+a),x)
```

Fricas [F]

$$\int x \operatorname{Chi}(c + dx) \sinh(a + bx) dx = \int x \operatorname{Chi}(dx + c) \sinh(bx + a) dx$$

```
[In] integrate(x*Chi(d*x+c)*sinh(b*x+a),x, algorithm="fricas")
```

```
[Out] integral(x*cosh_integral(d*x + c)*sinh(b*x + a), x)
```

Sympy [F]

$$\int x \operatorname{Chi}(c + dx) \sinh(a + bx) dx = \int x \sinh(a + bx) \operatorname{Chi}(c + dx) dx$$

```
[In] integrate(x*Chi(d*x+c)*sinh(b*x+a),x)
```

```
[Out] Integral(x*sinh(a + b*x)*Chi(c + d*x), x)
```

Maxima [F]

$$\int x \operatorname{Chi}(c + dx) \sinh(a + bx) dx = \int x \operatorname{Chi}(dx + c) \sinh(bx + a) dx$$

```
[In] integrate(x*Chi(d*x+c)*sinh(b*x+a),x, algorithm="maxima")
```

```
[Out] integrate(x*Chi(d*x + c)*sinh(b*x + a), x)
```

Giac [F]

$$\int x\text{Chi}(c + dx) \sinh(a + bx) dx = \int x\text{Chi}(dx + c) \sinh(bx + a) dx$$

[In] integrate(x*Chi(d*x+c)*sinh(b*x+a),x, algorithm="giac")

[Out] integrate(x*Chi(d*x + c)*sinh(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int x\text{Chi}(c + dx) \sinh(a + bx) dx = \int x \text{coshint}(c + dx) \sinh(a + bx) dx$$

[In] int(x*coshint(c + d*x)*sinh(a + b*x),x)

[Out] int(x*coshint(c + d*x)*sinh(a + b*x), x)

3.132 $\int \mathbf{Chi}(c + dx) \sinh(a + bx) dx$

Optimal result	644
Rubi [A] (verified)	644
Mathematica [A] (verified)	647
Maple [F]	647
Fricas [F]	647
Sympy [F]	647
Maxima [F]	648
Giac [F]	648
Mupad [F(-1)]	648

Optimal result

Integrand size = 13, antiderivative size = 153

$$\int \mathbf{Chi}(c + dx) \sinh(a + bx) dx = -\frac{\cosh\left(a - \frac{bc}{d}\right) \mathbf{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b} + \frac{\cosh(a + bx) \mathbf{Chi}(c + dx)}{b} - \frac{\cosh\left(a - \frac{bc}{d}\right) \mathbf{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b} - \frac{\sinh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b} - \frac{\sinh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b}$$

[Out] -1/2*Chi(c*(b-d)/d+(b-d)*x)*cosh(a-b*c/d)/b-1/2*Chi(c*(b+d)/d+(b+d)*x)*cosh(a-b*c/d)/b+Chi(d*x+c)*cosh(b*x+a)/b-1/2*Shi(c*(b-d)/d+(b-d)*x)*sinh(a-b*c/d)/b-1/2*Shi(c*(b+d)/d+(b+d)*x)*sinh(a-b*c/d)/b

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used

= {6682, 5579, 3384, 3379, 3382}

$$\int \text{Chi}(c + dx) \sinh(a + bx) dx = -\frac{\cosh\left(a - \frac{bc}{d}\right) \text{Chi}\left(x(b - d) + \frac{c(b-d)}{d}\right)}{2b} + \frac{\cosh(a + bx) \text{Chi}(c + dx)}{b} - \frac{\cosh\left(a - \frac{bc}{d}\right) \text{Chi}\left(x(b + d) + \frac{c(b+d)}{d}\right)}{2b} - \frac{\sinh\left(a - \frac{bc}{d}\right) \text{Shi}\left(x(b - d) + \frac{c(b-d)}{d}\right)}{2b} - \frac{\sinh\left(a - \frac{bc}{d}\right) \text{Shi}\left(x(b + d) + \frac{c(b+d)}{d}\right)}{2b}$$

[In] Int[CoshIntegral[c + d*x]*Sinh[a + b*x], x]

[Out] -1/2*(Cosh[a - (b*c)/d]*CoshIntegral[(c*(b - d))/d + (b - d)*x])/b + (Cosh[a + b*x]*CoshIntegral[c + d*x])/b - (Cosh[a - (b*c)/d]*CoshIntegral[(c*(b + d))/d + (b + d)*x])/(2*b) - (Sinh[a - (b*c)/d]*SinhIntegral[(c*(b - d))/d + (b - d)*x])/(2*b) - (Sinh[a - (b*c)/d]*SinhIntegral[(c*(b + d))/d + (b + d)*x])/(2*b)

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3384

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]

Rule 5579

Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*Cosh[(c_.) + (d_.)*(x_)]^(q_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e + f*x)^m, Cosh[a + b*x]^p*Cosh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0]

] && IGtQ[q, 0] && IntegerQ[m]

Rule 6682

```
Int[CoshIntegral[(c_.) + (d_.)*(x_)]*Sinh[(a_.) + (b_.)*(x_)], x_Symbol] :>
Simp[Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\cosh(a + bx)\text{Chi}(c + dx)}{b} - \frac{d \int \frac{\cosh(a+bx)\cosh(c+dx)}{c+dx} dx}{b} \\
&= \frac{\cosh(a + bx)\text{Chi}(c + dx)}{b} - \frac{d \int \left(\frac{\cosh(a-c+(b-d)x)}{2(c+dx)} + \frac{\cosh(a+c+(b+d)x)}{2(c+dx)} \right) dx}{b} \\
&= \frac{\cosh(a + bx)\text{Chi}(c + dx)}{b} - \frac{d \int \frac{\cosh(a-c+(b-d)x)}{c+dx} dx}{2b} - \frac{d \int \frac{\cosh(a+c+(b+d)x)}{c+dx} dx}{2b} \\
&= \frac{\cosh(a + bx)\text{Chi}(c + dx)}{b} - \frac{(d \cosh(a - \frac{bc}{d})) \int \frac{\cosh(\frac{c(b-d)}{d} + (b-d)x)}{c+dx} dx}{2b} \\
&\quad - \frac{(d \cosh(a - \frac{bc}{d})) \int \frac{\cosh(\frac{c(b+d)}{d} + (b+d)x)}{c+dx} dx}{2b} \\
&\quad - \frac{(d \sinh(a - \frac{bc}{d})) \int \frac{\sinh(\frac{c(b-d)}{d} + (b-d)x)}{c+dx} dx}{2b} \\
&\quad - \frac{(d \sinh(a - \frac{bc}{d})) \int \frac{\sinh(\frac{c(b+d)}{d} + (b+d)x)}{c+dx} dx}{2b} \\
&= -\frac{\cosh(a - \frac{bc}{d}) \text{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b} + \frac{\cosh(a + bx)\text{Chi}(c + dx)}{b} \\
&\quad - \frac{\cosh(a - \frac{bc}{d}) \text{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b} - \frac{\sinh(a - \frac{bc}{d}) \text{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b} \\
&\quad - \frac{\sinh(a - \frac{bc}{d}) \text{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.66 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.70

$$\int \text{Chi}(c + dx) \sinh(a + bx) dx = \frac{-4 \cosh(a + bx) \text{Chi}(c + dx) + e^{-a + \frac{bc}{d}} \left(\text{ExpIntegralEi} \left(-\frac{(b-d)(c+dx)}{d} \right) + \text{ExpIntegralEi} \left(-\frac{(b+d)(c+dx)}{d} \right) \right)}{4b}$$

[In] Integrate[CoshIntegral[c + d*x]*Sinh[a + b*x],x]

[Out] -1/4*(-4*Cosh[a + b*x]*CoshIntegral[c + d*x] + E^(-a + (b*c)/d)*(ExpIntegralEi[-(((b - d)*(c + d*x))/d)] + ExpIntegralEi[-(((b + d)*(c + d*x))/d])) + E^(a - (b*c)/d)*(ExpIntegralEi[((b - d)*(c + d*x))/d] + ExpIntegralEi[((b + d)*(c + d*x))/d]))/b

Maple [F]

$$\int \text{Chi}(dx + c) \sinh(bx + a) dx$$

[In] int(Chi(d*x+c)*sinh(b*x+a),x)

[Out] int(Chi(d*x+c)*sinh(b*x+a),x)

Fricas [F]

$$\int \text{Chi}(c + dx) \sinh(a + bx) dx = \int \text{Chi}(dx + c) \sinh(bx + a) dx$$

[In] integrate(Chi(d*x+c)*sinh(b*x+a),x, algorithm="fricas")

[Out] integral(cosh_integral(d*x + c)*sinh(b*x + a), x)

Sympy [F]

$$\int \text{Chi}(c + dx) \sinh(a + bx) dx = \int \sinh(a + bx) \text{Chi}(c + dx) dx$$

[In] integrate(Chi(d*x+c)*sinh(b*x+a),x)

[Out] Integral(sinh(a + b*x)*Chi(c + d*x), x)

Maxima [F]

$$\int \text{Chi}(c + dx) \sinh(a + bx) dx = \int \text{Chi}(dx + c) \sinh(bx + a) dx$$

[In] integrate(Chi(d*x+c)*sinh(b*x+a),x, algorithm="maxima")

[Out] integrate(Chi(d*x + c)*sinh(b*x + a), x)

Giac [F]

$$\int \text{Chi}(c + dx) \sinh(a + bx) dx = \int \text{Chi}(dx + c) \sinh(bx + a) dx$$

[In] integrate(Chi(d*x+c)*sinh(b*x+a),x, algorithm="giac")

[Out] integrate(Chi(d*x + c)*sinh(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int \text{Chi}(c + dx) \sinh(a + bx) dx = \int \text{coshint}(c + dx) \sinh(a + bx) dx$$

[In] int(coshint(c + d*x)*sinh(a + b*x),x)

[Out] int(coshint(c + d*x)*sinh(a + b*x), x)

3.133 $\int \frac{\text{Chi}(c+dx) \sinh(a+bx)}{x} dx$

Optimal result	649
Rubi [N/A]	649
Mathematica [N/A]	650
Maple [N/A] (verified)	650
Fricas [N/A]	650
Sympy [N/A]	650
Maxima [N/A]	651
Giac [N/A]	651
Mupad [N/A]	651

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\text{Chi}(c+dx) \sinh(a+bx)}{x} dx = \text{Int}\left(\frac{\text{Chi}(c+dx) \sinh(a+bx)}{x}, x\right)$$

[Out] CannotIntegrate(Chi(d*x+c)*sinh(b*x+a)/x,x)

Rubi [N/A]

Not integrable

Time = 0.11 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\text{Chi}(c+dx) \sinh(a+bx)}{x} dx = \int \frac{\text{Chi}(c+dx) \sinh(a+bx)}{x} dx$$

[In] Int[(CoshIntegral[c + d*x]*Sinh[a + b*x])/x,x]

[Out] Defer[Int] [(CoshIntegral[c + d*x]*Sinh[a + b*x])/x, x]

Rubi steps

$$\text{integral} = \int \frac{\text{Chi}(c+dx) \sinh(a+bx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 2.94 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\text{Chi}(c + dx) \sinh(a + bx)}{x} dx = \int \frac{\text{Chi}(c + dx) \sinh(a + bx)}{x} dx$$

[In] Integrate[(CoshIntegral[c + d*x]*Sinh[a + b*x])/x,x]

[Out] Integrate[(CoshIntegral[c + d*x]*Sinh[a + b*x])/x, x]

Maple [N/A] (verified)

Not integrable

Time = 0.41 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(dx + c) \sinh(bx + a)}{x} dx$$

[In] int(Chi(d*x+c)*sinh(b*x+a)/x,x)

[Out] int(Chi(d*x+c)*sinh(b*x+a)/x,x)

Fricas [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\text{Chi}(c + dx) \sinh(a + bx)}{x} dx = \int \frac{\text{Chi}(dx + c) \sinh(bx + a)}{x} dx$$

[In] integrate(Chi(d*x+c)*sinh(b*x+a)/x,x, algorithm="fricas")

[Out] integral(cosh_integral(d*x + c)*sinh(b*x + a)/x, x)

Sympy [N/A]

Not integrable

Time = 0.91 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{\text{Chi}(c + dx) \sinh(a + bx)}{x} dx = \int \frac{\sinh(a + bx) \text{Chi}(c + dx)}{x} dx$$

[In] integrate(Chi(d*x+c)*sinh(b*x+a)/x,x)

[Out] Integral(sinh(a + b*x)*Chi(c + d*x)/x, x)

Maxima [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\text{Chi}(c + dx) \sinh(a + bx)}{x} dx = \int \frac{\text{Chi}(dx + c) \sinh(bx + a)}{x} dx$$

[In] integrate(Chi(d*x+c)*sinh(b*x+a)/x,x, algorithm="maxima")

[Out] integrate(Chi(d*x + c)*sinh(b*x + a)/x, x)

Giac [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\text{Chi}(c + dx) \sinh(a + bx)}{x} dx = \int \frac{\text{Chi}(dx + c) \sinh(bx + a)}{x} dx$$

[In] integrate(Chi(d*x+c)*sinh(b*x+a)/x,x, algorithm="giac")

[Out] integrate(Chi(d*x + c)*sinh(b*x + a)/x, x)

Mupad [N/A]

Not integrable

Time = 5.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\text{Chi}(c + dx) \sinh(a + bx)}{x} dx = \int \frac{\text{coshint}(c + dx) \sinh(a + bx)}{x} dx$$

[In] int((coshint(c + d*x)*sinh(a + b*x))/x,x)

[Out] int((coshint(c + d*x)*sinh(a + b*x))/x, x)

3.134 $\int x \cosh(a + bx) \mathbf{Chi}(c + dx) dx$

Optimal result	652
Rubi [A] (verified)	653
Mathematica [A] (verified)	657
Maple [F]	657
Fricas [F]	658
Sympy [F]	658
Maxima [F]	658
Giac [F]	658
Mupad [F(-1)]	659

Optimal result

Integrand size = 14, antiderivative size = 371

$$\begin{aligned}
 \int x \cosh(a + bx) \mathbf{Chi}(c + dx) dx = & -\frac{\cosh(a - c + (b - d)x)}{2b(b - d)} - \frac{\cosh(a + c + (b + d)x)}{2b(b + d)} \\
 & + \frac{\cosh\left(a - \frac{bc}{d}\right) \mathbf{Chi}\left(\frac{c(b-d)}{d} + (b - d)x\right)}{2b^2} \\
 & - \frac{\cosh(a + bx) \mathbf{Chi}(c + dx)}{b^2} \\
 & + \frac{\cosh\left(a - \frac{bc}{d}\right) \mathbf{Chi}\left(\frac{c(b+d)}{d} + (b + d)x\right)}{2b^2} \\
 & + \frac{c \mathbf{Chi}\left(\frac{c(b-d)}{d} + (b - d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2bd} \\
 & + \frac{c \mathbf{Chi}\left(\frac{c(b+d)}{d} + (b + d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2bd} \\
 & + \frac{x \mathbf{Chi}(c + dx) \sinh(a + bx)}{b} \\
 & + \frac{c \cosh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b-d)}{d} + (b - d)x\right)}{2bd} \\
 & + \frac{\sinh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b-d)}{d} + (b - d)x\right)}{2b^2} \\
 & + \frac{c \cosh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b+d)}{d} + (b + d)x\right)}{2bd} \\
 & + \frac{\sinh\left(a - \frac{bc}{d}\right) \mathbf{Shi}\left(\frac{c(b+d)}{d} + (b + d)x\right)}{2b^2}
 \end{aligned}$$

[Out] $\frac{1}{2} \operatorname{Chi}\left(\frac{c(b-d)}{d+(b-d)x}\right) \operatorname{cosh}\left(\frac{a-bc}{d}\right) / b^2 + \frac{1}{2} \operatorname{Chi}\left(\frac{c(b+d)}{d+(b+d)x}\right) \operatorname{cosh}\left(\frac{a-bc}{d}\right) / b^2 - \operatorname{Chi}(dx+c) \operatorname{cosh}(bx+a) / b^2 - \frac{1}{2} \operatorname{cosh}\left(\frac{a-c+(b-d)x}{b}\right) / (b-d) - \frac{1}{2} \operatorname{cosh}\left(\frac{a+c+(b+d)x}{b}\right) / (b+d) + \frac{1}{2} c \operatorname{cosh}\left(\frac{a-bc}{d}\right) \operatorname{Shi}\left(\frac{c(b-d)}{d+(b-d)x}\right) / b/d + \frac{1}{2} c \operatorname{cosh}\left(\frac{a-bc}{d}\right) \operatorname{Shi}\left(\frac{c(b+d)}{d+(b+d)x}\right) / b/d + \frac{1}{2} c \operatorname{Chi}\left(\frac{c(b-d)}{d+(b-d)x}\right) \operatorname{sinh}\left(\frac{a-bc}{d}\right) / b/d + \frac{1}{2} c \operatorname{Chi}\left(\frac{c(b+d)}{d+(b+d)x}\right) \operatorname{sinh}\left(\frac{a-bc}{d}\right) / b/d + \frac{1}{2} \operatorname{Shi}\left(\frac{c(b-d)}{d+(b-d)x}\right) \operatorname{sinh}\left(\frac{a-bc}{d}\right) / b^2 + \frac{1}{2} \operatorname{Shi}\left(\frac{c(b+d)}{d+(b+d)x}\right) \operatorname{sinh}\left(\frac{a-bc}{d}\right) / b^2 + x \operatorname{Chi}(dx+c) \operatorname{sinh}(bx+a) / b$

Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 371, normalized size of antiderivative = 1.00, number of steps used = 24, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {6678, 6874, 5737, 2718, 5580, 3384, 3379, 3382, 6682, 5579}

$$\int x \operatorname{cosh}(a+bx) \operatorname{Chi}(c+dx) dx = \frac{\operatorname{cosh}\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2b^2} - \frac{\operatorname{cosh}(a+bx) \operatorname{Chi}(c+dx)}{b^2} + \frac{\operatorname{cosh}\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2b^2} + \frac{\operatorname{sinh}\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2b^2} + \frac{\operatorname{sinh}\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2b^2} + \frac{c \operatorname{sinh}\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2bd} + \frac{c \operatorname{sinh}\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2bd} + \frac{x \operatorname{sinh}(a+bx) \operatorname{Chi}(c+dx)}{b} + \frac{c \operatorname{cosh}\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(x(b-d) + \frac{c(b-d)}{d}\right)}{2bd} + \frac{c \operatorname{cosh}\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(x(b+d) + \frac{c(b+d)}{d}\right)}{2bd} - \frac{\operatorname{cosh}(a+x(b-d)-c)}{2b(b-d)} - \frac{\operatorname{cosh}(a+x(b+d)+c)}{2b(b+d)}$$

[In] $\operatorname{Int}[x \operatorname{Cosh}[a+bx] \operatorname{CoshIntegral}[c+dx], x]$

[Out] $-1/2 \operatorname{Cosh}[a-c+(b-d)x] / (b(b-d)) - \operatorname{Cosh}[a+c+(b+d)x] / (2b(b+d)) + (\operatorname{Cosh}[a-(bc)/d] \operatorname{CoshIntegral}[(c(b-d))/d+(b-d)x]) / (2b^2)$

$$\begin{aligned}
& - (\text{Cosh}[a + b*x]*\text{CoshIntegral}[c + d*x])/b^2 + (\text{Cosh}[a - (b*c)/d]*\text{CoshIntegral}[(c*(b + d))/d + (b + d)*x])/(2*b^2) + (c*\text{CoshIntegral}[(c*(b - d))/d + (b - d)*x]*\text{Sinh}[a - (b*c)/d])/(2*b*d) + (c*\text{CoshIntegral}[(c*(b + d))/d + (b + d)*x]*\text{Sinh}[a - (b*c)/d])/(2*b*d) + (x*\text{CoshIntegral}[c + d*x]*\text{Sinh}[a + b*x])/b + (c*\text{Cosh}[a - (b*c)/d]*\text{SinhIntegral}[(c*(b - d))/d + (b - d)*x])/(2*b*d) + (\text{Sinh}[a - (b*c)/d]*\text{SinhIntegral}[(c*(b - d))/d + (b - d)*x])/(2*b^2) + (c*\text{Cosh}[a - (b*c)/d]*\text{SinhIntegral}[(c*(b + d))/d + (b + d)*x])/(2*b*d) + (\text{Sinh}[a - (b*c)/d]*\text{SinhIntegral}[(c*(b + d))/d + (b + d)*x])/(2*b^2)
\end{aligned}$$
Rule 2718

```
Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3384

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]
```

Rule 5579

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*Cosh[(c_.) + (d_.)*(x_)]^(q_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e + f*x)^m, Cosh[a + b*x]^p*Cosh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IntegerQ[m]
```

Rule 5580

```
Int[Cosh[(c_.) + (d_.)*(x_)]^(q_.)*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(p_.), x_Symbol] := Int[ExpandTrigReduce[(e + f*x)^m, Sinh[a + b*x]^p*Cosh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[p, 0] && IGtQ[q, 0]
```

Rule 5737

```
Int[Cosh[w_]^(q_.)*Sinh[v_]^(p_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]
]^(p)*Cosh[w]^q, x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))
```

Rule 6678

```
Int[Cosh[(a_.) + (b_.)*(x_)]*CoshIntegral[(c_.) + (d_.)*(x_)]*((e_.) + (f_.
)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^m*Sinh[a + b*x]*(CoshIntegral[c
+ d*x]/b), x] + (-Dist[d/b, Int[(e + f*x)^m*Sinh[a + b*x]*(Cosh[c + d*x]/(c
+ d*x)), x], x] - Dist[f*(m/b), Int[(e + f*x)^(m - 1)*Sinh[a + b*x]*CoshIn
tegral[c + d*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6682

```
Int[CoshIntegral[(c_.) + (d_.)*(x_)]*Sinh[(a_.) + (b_.)*(x_)], x_Symbol] :=
Simp[Cosh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Cosh[a +
b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x\text{Chi}(c + dx) \sinh(a + bx)}{b} - \frac{\int \text{Chi}(c + dx) \sinh(a + bx) dx}{b} \\
&\quad - \frac{d \int \frac{x \cosh(c + dx) \sinh(a + bx)}{c + dx} dx}{b} \\
&= -\frac{\cosh(a + bx)\text{Chi}(c + dx)}{b^2} + \frac{x\text{Chi}(c + dx) \sinh(a + bx)}{b} \\
&\quad + \frac{d \int \frac{\cosh(a + bx) \cosh(c + dx)}{c + dx} dx}{b^2} - \frac{d \int \left(\frac{\cosh(c + dx) \sinh(a + bx)}{d} - \frac{c \cosh(c + dx) \sinh(a + bx)}{d(c + dx)} \right) dx}{b} \\
&= -\frac{\cosh(a + bx)\text{Chi}(c + dx)}{b^2} + \frac{x\text{Chi}(c + dx) \sinh(a + bx)}{b} \\
&\quad - \frac{\int \cosh(c + dx) \sinh(a + bx) dx}{b} + \frac{c \int \frac{\cosh(c + dx) \sinh(a + bx)}{c + dx} dx}{b} \\
&\quad + \frac{d \int \left(\frac{\cosh(a - c + (b - d)x)}{2(c + dx)} + \frac{\cosh(a + c + (b + d)x)}{2(c + dx)} \right) dx}{b^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{\cosh(a+bx)\text{Chi}(c+dx)}{b^2} + \frac{x\text{Chi}(c+dx)\sinh(a+bx)}{b} \\
&\quad - \frac{\int \left(\frac{1}{2}\sinh(a-c+(b-d)x) + \frac{1}{2}\sinh(a+c+(b+d)x)\right) dx}{b} \\
&\quad + \frac{c \int \left(\frac{\sinh(a-c+(b-d)x)}{2(c+dx)} + \frac{\sinh(a+c+(b+d)x)}{2(c+dx)}\right) dx}{b} \\
&\quad + \frac{d \int \frac{\cosh(a-c+(b-d)x)}{c+dx} dx}{2b^2} + \frac{d \int \frac{\cosh(a+c+(b+d)x)}{c+dx} dx}{2b^2} \\
&= -\frac{\cosh(a+bx)\text{Chi}(c+dx)}{b^2} + \frac{x\text{Chi}(c+dx)\sinh(a+bx)}{b} - \frac{\int \sinh(a-c+(b-d)x) dx}{2b} \\
&\quad - \frac{\int \sinh(a+c+(b+d)x) dx}{2b} + \frac{c \int \frac{\sinh(a-c+(b-d)x)}{c+dx} dx}{2b} + \frac{c \int \frac{\sinh(a+c+(b+d)x)}{c+dx} dx}{2b} \\
&\quad + \frac{(d \cosh(a-\frac{bc}{d})) \int \frac{\cosh(\frac{c(b-d)}{d}+(b-d)x)}{c+dx} dx}{2b^2} + \frac{(d \cosh(a-\frac{bc}{d})) \int \frac{\cosh(\frac{c(b+d)}{d}+(b+d)x)}{c+dx} dx}{2b^2} \\
&\quad + \frac{(d \sinh(a-\frac{bc}{d})) \int \frac{\sinh(\frac{c(b-d)}{d}+(b-d)x)}{c+dx} dx}{2b^2} + \frac{(d \sinh(a-\frac{bc}{d})) \int \frac{\sinh(\frac{c(b+d)}{d}+(b+d)x)}{c+dx} dx}{2b^2} \\
&= -\frac{\cosh(a-c+(b-d)x)}{2b(b-d)} - \frac{\cosh(a+c+(b+d)x)}{2b(b+d)} \\
&\quad + \frac{\cosh(a-\frac{bc}{d})\text{Chi}\left(\frac{c(b-d)}{d}+(b-d)x\right)}{2b^2} - \frac{\cosh(a+bx)\text{Chi}(c+dx)}{b^2} \\
&\quad + \frac{\cosh(a-\frac{bc}{d})\text{Chi}\left(\frac{c(b+d)}{d}+(b+d)x\right)}{2b^2} + \frac{x\text{Chi}(c+dx)\sinh(a+bx)}{b} \\
&\quad + \frac{\sinh(a-\frac{bc}{d})\text{Shi}\left(\frac{c(b-d)}{d}+(b-d)x\right)}{2b^2} + \frac{\sinh(a-\frac{bc}{d})\text{Shi}\left(\frac{c(b+d)}{d}+(b+d)x\right)}{2b^2} \\
&\quad + \frac{(c \cosh(a-\frac{bc}{d})) \int \frac{\sinh(\frac{c(b-d)}{d}+(b-d)x)}{c+dx} dx}{2b} \\
&\quad + \frac{(c \cosh(a-\frac{bc}{d})) \int \frac{\sinh(\frac{c(b+d)}{d}+(b+d)x)}{c+dx} dx}{2b} \\
&\quad + \frac{(c \sinh(a-\frac{bc}{d})) \int \frac{\cosh(\frac{c(b-d)}{d}+(b-d)x)}{c+dx} dx}{2b} \\
&\quad + \frac{(c \sinh(a-\frac{bc}{d})) \int \frac{\cosh(\frac{c(b+d)}{d}+(b+d)x)}{c+dx} dx}{2b}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{\cosh(a - c + (b - d)x)}{2b(b - d)} - \frac{\cosh(a + c + (b + d)x)}{2b(b + d)} \\
&+ \frac{\cosh\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(\frac{c(b-d)}{d} + (b - d)x\right)}{2b^2} - \frac{\cosh(a + bx) \operatorname{Chi}(c + dx)}{b^2} \\
&+ \frac{\cosh\left(a - \frac{bc}{d}\right) \operatorname{Chi}\left(\frac{c(b+d)}{d} + (b + d)x\right)}{2b^2} + \frac{c \operatorname{Chi}\left(\frac{c(b-d)}{d} + (b - d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2bd} \\
&+ \frac{c \operatorname{Chi}\left(\frac{c(b+d)}{d} + (b + d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2bd} + \frac{x \operatorname{Chi}(c + dx) \sinh(a + bx)}{b} \\
&+ \frac{c \cosh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(\frac{c(b-d)}{d} + (b - d)x\right)}{2bd} + \frac{\sinh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(\frac{c(b-d)}{d} + (b - d)x\right)}{2b^2} \\
&+ \frac{c \cosh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(\frac{c(b+d)}{d} + (b + d)x\right)}{2bd} + \frac{\sinh\left(a - \frac{bc}{d}\right) \operatorname{Shi}\left(\frac{c(b+d)}{d} + (b + d)x\right)}{2b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.97 (sec) , antiderivative size = 270, normalized size of antiderivative = 0.73

$$\int x \cosh(a + bx) \operatorname{Chi}(c + dx) dx$$

$$= \frac{e^{-a} \left(bde^{-c} \left(-\frac{e^{-((b+d)x)}}{b+d} - \frac{e^{2a+bx-dx}}{b-d} \right) + (bc+d)e^{2a-\frac{bc}{d}} \operatorname{ExpIntegralEi}\left(\frac{(b-d)(c+dx)}{d}\right) - (bc-d)e^{\frac{bc}{d}} \operatorname{ExpIntegralEi}\left(-\frac{(b+d)(c+dx)}{d}\right) \right)}{d} + \frac{e^{-a} (b \dots)}{d}$$

[In] Integrate[x*Cosh[a + b*x]*CoshIntegral[c + d*x],x]

[Out] (((b*d*(-1/((b + d)*E^((b + d)*x))) - E^(2*a + b*x - d*x)/(b - d)))/E^c + (b*c + d)*E^(2*a - (b*c)/d)*ExpIntegralEi[((b - d)*(c + d*x))/d] - (b*c - d)*E^((b*c)/d)*ExpIntegralEi[-(((b + d)*(c + d*x))/d)])/ (d*E^a) + (b*d*E^c*(E^((-b + d)*x)/(-b + d) - E^(2*a + (b + d)*x)/(b + d)) + (-b*c) + d)*E^((b*c)/d)*ExpIntegralEi[-(((b - d)*(c + d*x))/d)] + (b*c + d)*E^(2*a - (b*c)/d)*ExpIntegralEi[((b + d)*(c + d*x))/d])/ (d*E^a) + 4*CoshIntegral[c + d*x]*(-Cosh[a + b*x] + b*x*Sinh[a + b*x]))/(4*b^2)

Maple [F]

$$\int x \operatorname{Chi}(dx + c) \cosh(bx + a) dx$$

[In] int(x*Chi(d*x+c)*cosh(b*x+a),x)

[Out] int(x*Chi(d*x+c)*cosh(b*x+a),x)

Fricas [F]

$$\int x \cosh(a + bx) \operatorname{Chi}(c + dx) dx = \int x \operatorname{Chi}(dx + c) \cosh(bx + a) dx$$

[In] `integrate(x*Chi(d*x+c)*cosh(b*x+a),x, algorithm="fricas")`

[Out] `integral(x*cosh(b*x + a)*cosh_integral(d*x + c), x)`

Sympy [F]

$$\int x \cosh(a + bx) \operatorname{Chi}(c + dx) dx = \int x \cosh(a + bx) \operatorname{Chi}(c + dx) dx$$

[In] `integrate(x*Chi(d*x+c)*cosh(b*x+a),x)`

[Out] `Integral(x*cosh(a + b*x)*Chi(c + d*x), x)`

Maxima [F]

$$\int x \cosh(a + bx) \operatorname{Chi}(c + dx) dx = \int x \operatorname{Chi}(dx + c) \cosh(bx + a) dx$$

[In] `integrate(x*Chi(d*x+c)*cosh(b*x+a),x, algorithm="maxima")`

[Out] `integrate(x*Chi(d*x + c)*cosh(b*x + a), x)`

Giac [F]

$$\int x \cosh(a + bx) \operatorname{Chi}(c + dx) dx = \int x \operatorname{Chi}(dx + c) \cosh(bx + a) dx$$

[In] `integrate(x*Chi(d*x+c)*cosh(b*x+a),x, algorithm="giac")`

[Out] `integrate(x*Chi(d*x + c)*cosh(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x \cosh(a + bx) \operatorname{Chi}(c + dx) dx = \int x \operatorname{coshint}(c + dx) \cosh(a + bx) dx$$

```
[In] int(x*coshint(c + d*x)*cosh(a + b*x),x)
```

```
[Out] int(x*coshint(c + d*x)*cosh(a + b*x), x)
```

3.135 $\int \cosh(a + bx)\text{Chi}(c + dx) dx$

Optimal result	660
Rubi [A] (verified)	660
Mathematica [A] (verified)	663
Maple [F]	663
Fricas [F]	663
Sympy [F]	663
Maxima [F]	664
Giac [F]	664
Mupad [F(-1)]	664

Optimal result

Integrand size = 13, antiderivative size = 153

$$\int \cosh(a + bx)\text{Chi}(c + dx) dx = -\frac{\text{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b}$$

$$- \frac{\text{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b}$$

$$+ \frac{\text{Chi}(c + dx) \sinh(a + bx)}{b}$$

$$- \frac{\cosh\left(a - \frac{bc}{d}\right) \text{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b}$$

$$- \frac{\cosh\left(a - \frac{bc}{d}\right) \text{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b}$$

```
[Out] -1/2*cosh(a-b*c/d)*Shi(c*(b-d)/d+(b-d)*x)/b-1/2*cosh(a-b*c/d)*Shi(c*(b+d)/d
+(b+d)*x)/b-1/2*Chi(c*(b-d)/d+(b-d)*x)*sinh(a-b*c/d)/b-1/2*Chi(c*(b+d)/d+(b
+d)*x)*sinh(a-b*c/d)/b+Chi(d*x+c)*sinh(b*x+a)/b
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used

= {6676, 5580, 3384, 3379, 3382}

$$\int \cosh(a + bx)\text{Chi}(c + dx) dx = -\frac{\sinh\left(a - \frac{bc}{d}\right)\text{Chi}\left(x(b - d) + \frac{c(b-d)}{d}\right)}{2b} - \frac{\sinh\left(a - \frac{bc}{d}\right)\text{Chi}\left(x(b + d) + \frac{c(b+d)}{d}\right)}{2b} + \frac{\sinh(a + bx)\text{Chi}(c + dx)}{b} - \frac{\cosh\left(a - \frac{bc}{d}\right)\text{Shi}\left(x(b - d) + \frac{c(b-d)}{d}\right)}{2b} - \frac{\cosh\left(a - \frac{bc}{d}\right)\text{Shi}\left(x(b + d) + \frac{c(b+d)}{d}\right)}{2b}$$

[In] Int[Cosh[a + b*x]*CoshIntegral[c + d*x], x]

[Out] -1/2*(CoshIntegral[(c*(b - d))/d + (b - d)*x]*Sinh[a - (b*c)/d])/b - (CoshIntegral[(c*(b + d))/d + (b + d)*x]*Sinh[a - (b*c)/d])/(2*b) + (CoshIntegral[c + d*x]*Sinh[a + b*x])/b - (Cosh[a - (b*c)/d]*SinhIntegral[(c*(b - d))/d + (b - d)*x])/(2*b) - (Cosh[a - (b*c)/d]*SinhIntegral[(c*(b + d))/d + (b + d)*x])/(2*b)

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3384

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]

Rule 5580

Int[Cosh[(c_.) + (d_.)*(x_)]^(q_.)*((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(p_.), x_Symbol] := Int[ExpandTrigReduce[(e + f*x)^m, Sinh[a + b*x]^p*Cosh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[p

, 0] && IGtQ[q, 0]

Rule 6676

```
Int[Cosh[(a_.) + (b_.)*(x_.)]*CoshIntegral[(c_.) + (d_.)*(x_.)], x_Symbol] :>
Simp[Sinh[a + b*x]*(CoshIntegral[c + d*x]/b), x] - Dist[d/b, Int[Sinh[a +
b*x]*(Cosh[c + d*x]/(c + d*x)), x], x] /; FreeQ[{a, b, c, d}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Chi}(c + dx) \sinh(a + bx)}{b} - \frac{d \int \frac{\cosh(c+dx) \sinh(a+bx)}{c+dx} dx}{b} \\
&= \frac{\text{Chi}(c + dx) \sinh(a + bx)}{b} - \frac{d \int \left(\frac{\sinh(a-c+(b-d)x)}{2(c+dx)} + \frac{\sinh(a+c+(b+d)x)}{2(c+dx)} \right) dx}{b} \\
&= \frac{\text{Chi}(c + dx) \sinh(a + bx)}{b} - \frac{d \int \frac{\sinh(a-c+(b-d)x)}{c+dx} dx}{2b} - \frac{d \int \frac{\sinh(a+c+(b+d)x)}{c+dx} dx}{2b} \\
&= \frac{\text{Chi}(c + dx) \sinh(a + bx)}{b} - \frac{(d \cosh(a - \frac{bc}{d})) \int \frac{\sinh(\frac{c(b-d)}{d} + (b-d)x)}{c+dx} dx}{2b} \\
&\quad - \frac{(d \cosh(a - \frac{bc}{d})) \int \frac{\sinh(\frac{c(b+d)}{d} + (b+d)x)}{c+dx} dx}{2b} \\
&\quad - \frac{(d \sinh(a - \frac{bc}{d})) \int \frac{\cosh(\frac{c(b-d)}{d} + (b-d)x)}{c+dx} dx}{2b} \\
&\quad - \frac{(d \sinh(a - \frac{bc}{d})) \int \frac{\cosh(\frac{c(b+d)}{d} + (b+d)x)}{c+dx} dx}{2b} \\
&= -\frac{\text{Chi}\left(\frac{c(b-d)}{d} + (b-d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b} - \frac{\text{Chi}\left(\frac{c(b+d)}{d} + (b+d)x\right) \sinh\left(a - \frac{bc}{d}\right)}{2b} \\
&\quad + \frac{\text{Chi}(c + dx) \sinh(a + bx)}{b} - \frac{\cosh\left(a - \frac{bc}{d}\right) \text{Shi}\left(\frac{c(b-d)}{d} + (b-d)x\right)}{2b} \\
&\quad - \frac{\cosh\left(a - \frac{bc}{d}\right) \text{Shi}\left(\frac{c(b+d)}{d} + (b+d)x\right)}{2b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.90

$$\int \cosh(a + bx) \operatorname{Chi}(c + dx) dx$$

$$= \frac{e^{-a - \frac{bc}{d}} \left(e^{\frac{2bc}{d}} \operatorname{ExpIntegralEi} \left(-\frac{(b-d)(c+dx)}{d} \right) - e^{2a} \operatorname{ExpIntegralEi} \left(\frac{(b-d)(c+dx)}{d} \right) + e^{\frac{2bc}{d}} \operatorname{ExpIntegralEi} \left(-\frac{(b+d)(c+dx)}{d} \right) \right)}{4b}$$

[In] Integrate[Cosh[a + b*x]*CoshIntegral[c + d*x],x]

[Out] (E^(-a - (b*c)/d)*(E^((2*b*c)/d)*ExpIntegralEi[-((b - d)*(c + d*x))/d]) - E^(2*a)*ExpIntegralEi[((b - d)*(c + d*x))/d] + E^((2*b*c)/d)*ExpIntegralEi[-((b + d)*(c + d*x))/d]) - E^(2*a)*ExpIntegralEi[((b + d)*(c + d*x))/d] + 4*E^(a + (b*c)/d)*CoshIntegral[c + d*x]*Sinh[a + b*x])/(4*b)

Maple [F]

$$\int \operatorname{Chi}(dx + c) \cosh(bx + a) dx$$

[In] int(Chi(d*x+c)*cosh(b*x+a),x)

[Out] int(Chi(d*x+c)*cosh(b*x+a),x)

Fricas [F]

$$\int \cosh(a + bx) \operatorname{Chi}(c + dx) dx = \int \operatorname{Chi}(dx + c) \cosh(bx + a) dx$$

[In] integrate(Chi(d*x+c)*cosh(b*x+a),x, algorithm="fricas")

[Out] integral(cosh(b*x + a)*cosh_integral(d*x + c), x)

Sympy [F]

$$\int \cosh(a + bx) \operatorname{Chi}(c + dx) dx = \int \cosh(a + bx) \operatorname{Chi}(c + dx) dx$$

[In] integrate(Chi(d*x+c)*cosh(b*x+a),x)

[Out] Integral(cosh(a + b*x)*Chi(c + d*x), x)

Maxima [F]

$$\int \cosh(a + bx)\text{Chi}(c + dx) dx = \int \text{Chi}(dx + c) \cosh(bx + a) dx$$

[In] integrate(Chi(d*x+c)*cosh(b*x+a),x, algorithm="maxima")

[Out] integrate(Chi(d*x + c)*cosh(b*x + a), x)

Giac [F]

$$\int \cosh(a + bx)\text{Chi}(c + dx) dx = \int \text{Chi}(dx + c) \cosh(bx + a) dx$$

[In] integrate(Chi(d*x+c)*cosh(b*x+a),x, algorithm="giac")

[Out] integrate(Chi(d*x + c)*cosh(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int \cosh(a + bx)\text{Chi}(c + dx) dx = \int \text{coshint}(c + dx) \cosh(a + bx) dx$$

[In] int(coshint(c + d*x)*cosh(a + b*x),x)

[Out] int(coshint(c + d*x)*cosh(a + b*x), x)

3.136 $\int \frac{\cosh(a+bx)\mathbf{Chi}(c+dx)}{x} dx$

Optimal result	665
Rubi [N/A]	665
Mathematica [N/A]	666
Maple [N/A] (verified)	666
Fricas [N/A]	666
Sympy [N/A]	666
Maxima [N/A]	667
Giac [N/A]	667
Mupad [N/A]	667

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\cosh(a+bx)\mathbf{Chi}(c+dx)}{x} dx = \text{Int}\left(\frac{\cosh(a+bx)\mathbf{Chi}(c+dx)}{x}, x\right)$$

[Out] CannotIntegrate(Chi(d*x+c)*cosh(b*x+a)/x,x)

Rubi [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\cosh(a+bx)\mathbf{Chi}(c+dx)}{x} dx = \int \frac{\cosh(a+bx)\mathbf{Chi}(c+dx)}{x} dx$$

[In] Int[(Cosh[a + b*x]*CoshIntegral[c + d*x])/x,x]

[Out] Defer[Int] [(Cosh[a + b*x]*CoshIntegral[c + d*x])/x, x]

Rubi steps

$$\text{integral} = \int \frac{\cosh(a+bx)\mathbf{Chi}(c+dx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 2.73 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Chi}(c + dx)}{x} dx = \int \frac{\cosh(a + bx)\text{Chi}(c + dx)}{x} dx$$

[In] Integrate[(Cosh[a + b*x]*CoshIntegral[c + d*x])/x,x]

[Out] Integrate[(Cosh[a + b*x]*CoshIntegral[c + d*x])/x, x]

Maple [N/A] (verified)

Not integrable

Time = 0.44 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\text{Chi}(dx + c) \cosh(bx + a)}{x} dx$$

[In] int(Chi(d*x+c)*cosh(b*x+a)/x,x)

[Out] int(Chi(d*x+c)*cosh(b*x+a)/x,x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Chi}(c + dx)}{x} dx = \int \frac{\text{Chi}(dx + c) \cosh(bx + a)}{x} dx$$

[In] integrate(Chi(d*x+c)*cosh(b*x+a)/x,x, algorithm="fricas")

[Out] integral(cosh(b*x + a)*cosh_integral(d*x + c)/x, x)

Sympy [N/A]

Not integrable

Time = 0.90 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{\cosh(a + bx)\text{Chi}(c + dx)}{x} dx = \int \frac{\cosh(a + bx)\text{Chi}(c + dx)}{x} dx$$

[In] integrate(Chi(d*x+c)*cosh(b*x+a)/x,x)

[Out] Integral(cosh(a + b*x)*Chi(c + d*x)/x, x)

Maxima [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Chi}(c + dx)}{x} dx = \int \frac{\text{Chi}(dx + c) \cosh(bx + a)}{x} dx$$

[In] integrate(Chi(d*x+c)*cosh(b*x+a)/x,x, algorithm="maxima")

[Out] integrate(Chi(d*x + c)*cosh(b*x + a)/x, x)

Giac [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Chi}(c + dx)}{x} dx = \int \frac{\text{Chi}(dx + c) \cosh(bx + a)}{x} dx$$

[In] integrate(Chi(d*x+c)*cosh(b*x+a)/x,x, algorithm="giac")

[Out] integrate(Chi(d*x + c)*cosh(b*x + a)/x, x)

Mupad [N/A]

Not integrable

Time = 5.14 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cosh(a + bx)\text{Chi}(c + dx)}{x} dx = \int \frac{\coshint(c + dx) \cosh(a + bx)}{x} dx$$

[In] int((coshint(c + d*x)*cosh(a + b*x))/x,x)

[Out] int((coshint(c + d*x)*cosh(a + b*x))/x, x)

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 669

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*      Small rewrite of logic in main function to make it*)
(*      match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
```

```

(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCo
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count is
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"}
    ]
  ]
  ,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<>
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

  finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)

```



```

    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result, optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);

```



```

#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 ("
```

```

                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_c
    end if
  else #result contains complex but optimal is not
    if debug then
      print("result contains complex but optimal is not");
    fi;
    return "C","Result contains complex when optimal does not.";
  fi;
else # result do not contain complex
  # this assumes optimal do not as well. No check is needed here.
  if debug then
    print("result do not contain complex, this assumes optimal do not as well");
  fi;
  if leaf_count_result<=2*leaf_count_optimal then
    if debug then
      print("leaf_count_result<=2*leaf_count_optimal");
    fi;
    return "A"," ";
  else
    if debug then
      print("leaf_count_result>2*leaf_count_optimal");
    fi;
    return "B",cat("Leaf count of result is larger than twice the leaf count of opt
                                convert(leaf_count_result,string)," $ vs. $2(",
                                convert(leaf_count_optimal,string)," )=",convert(2*leaf_coun
    fi;
  fi;
else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
                convert(ExpnType_result,string)," vs. order ",
                convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

```

```

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`) or type(expn,'*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else

```

```

9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u), u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#           Port of original Maple grading function by
#           Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#           added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
                    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
                    asinh,acosh,atanh,acoth,asech,acsch
                    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
                    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
                    gamma,loggamma,digamma,zeta,polylog,LambertW,
                    elliptic_f,elliptic_e,elliptic_pi,exp_polar
                    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnTy
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+' or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal."
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)+"/"+str(leaf_count_optimal)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)+"/"+str(ExpnType_optimal)

```

```

# print("Before returning. grade=", grade, " grade_annotation=", grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

# Dec 24, 2019. Nasser: Ported original Maple grading function by
# Albert Rich to use with Sagemath. This is used to
# grade Fricas, Giac and Maxima results.
# Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
# 'arctan2', 'floor', 'abs', 'log_integral'
# June 4, 2022 Made default grade_annotation "none" instead of "" due
# issue later when reading the file.
# July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    # print("Enter tree_size, expr is ", expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: # isinstance(expr, Pow):
        if expr.operands()[1] == 1/2: # expr.args[1] == Rational(1,2):
            if debug: print("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```



```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception,AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

    return 1
  else:
    return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
  else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. " + str(leaf_c
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_result)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```