

Computer Algebra Independent Integration Tests

Summer 2023 edition with Rubi V 4.17.3

0-Independent-test-suites/9-Stewart-Problems

Nasser M. Abbasi

December 4, 2023

Compiled on December 4, 2023 at 7:05pm

Contents

1	Introduction	2
2	detailed summary tables of results	20
3	Listing of integrals	134
4	Appendix	1891

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	3
1.2	Results	4
1.3	Time and leaf size Performance	7
1.4	Performance based on number of rules Rubi used	9
1.5	Performance based on number of steps Rubi used	10
1.6	Solved integrals histogram based on leaf size of result	11
1.7	Solved integrals histogram based on CPU time used	12
1.8	Leaf size vs. CPU time used	13
1.9	list of integrals with no known antiderivative	14
1.10	List of integrals solved by CAS but has no known antiderivative	14
1.11	list of integrals solved by CAS but failed verification	14
1.12	Timing	15
1.13	Verification	15
1.14	Important notes about some of the results	15
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [376]. This is test number [9].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 13.3.1 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (376)	0.00 (0)
Mathematica	100.00 (376)	0.00 (0)
Maple	100.00 (376)	0.00 (0)
Fricas	100.00 (376)	0.00 (0)
Giac	99.73 (375)	0.27 (1)
Maxima	99.47 (374)	0.53 (2)
Mupad	98.94 (372)	1.06 (4)
Sympy	96.54 (363)	3.46 (13)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	98.936	0.266	0.798	0.000
Maple	98.936	0.532	0.532	0.000
Mathematica	93.085	5.319	1.596	0.000
Maxima	91.489	7.979	0.000	0.532
Giac	90.426	9.309	0.000	0.266
Fricas	85.904	14.096	0.000	0.000
Sympy	79.787	11.702	5.053	3.457
Mupad	0.000	98.936	0.000	1.064

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates

an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Maple	0	0.00	0.00	0.00
Giac	1	0.00	100.00	0.00
Maxima	2	100.00	0.00	0.00
Mupad	4	0.00	100.00	0.00
Sympy	13	76.92	23.08	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Mathematica	0.03
Mupad	0.14
Rubi	0.17
Sympy	0.22
Maxima	0.23
Fricas	0.25
Giac	0.29
Maple	0.30

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Maple	18.76	0.89	16.00	0.83
Mupad	19.37	0.90	16.00	0.80
Giac	21.24	1.10	18.00	0.82
Maxima	21.78	1.04	16.00	0.80
Mathematica	23.20	1.12	20.00	1.00
Rubi	24.17	1.05	19.50	1.00
Fricas	24.64	1.10	18.00	0.86
Sympy	223.04	14.37	19.00	0.88

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following gives the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {77, 79, 115}

Mathematica {}

Maple {235, 323}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

1.14.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.14.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

1.14.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

1.14.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2013
Design v0.6

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	21
2.2	Detailed conclusion table per each integral for all CAS systems	27
2.3	Detailed conclusion table specific for Rubi results	122

2.1 List of integrals sorted by grade for each CAS

2.1.1	Rubi	21
2.1.2	Mma	22
2.1.3	Maple	22
2.1.4	Fricas	23
2.1.5	Maxima	24
2.1.6	Giac	24
2.1.7	Mupad	25
2.1.8	Sympy	26

2.1.1 Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376 }

B grade { 112 }

C grade { 34, 276, 300 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.2 Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 99, 101, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 122, 123, 124, 125, 126, 127, 128, 129, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 146, 147, 148, 149, 150, 151, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 213, 214, 215, 216, 217, 218, 219, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 236, 237, 238, 239, 240, 241, 242, 243, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 313, 314, 315, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 329, 330, 331, 332, 333, 334, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 371, 372, 373, 374, 375, 376 }

B grade { 14, 81, 100, 102, 103, 104, 121, 130, 145, 152, 195, 212, 221, 245, 246, 270, 297, 312, 328, 370 }

C grade { 98, 220, 235, 244, 316, 335 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.3 Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245,

246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376 }

B grade { 270, 359 }

C grade { 195, 323 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.4 Fricas

A grade { 1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 80, 81, 82, 83, 84, 85, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 106, 107, 109, 110, 111, 112, 114, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 131, 132, 133, 134, 135, 136, 137, 138, 140, 141, 142, 144, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 222, 223, 224, 226, 227, 228, 229, 230, 231, 232, 233, 234, 236, 237, 238, 239, 240, 242, 243, 245, 247, 252, 253, 254, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 292, 293, 294, 296, 297, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 329, 330, 331, 332, 333, 335, 336, 338, 339, 340, 342, 343, 345, 346, 347, 348, 349, 350, 352, 353, 354, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 371, 372, 373, 374, 375, 376 }

B grade { 7, 13, 14, 29, 41, 67, 79, 86, 97, 98, 99, 100, 101, 102, 103, 104, 105, 108, 113, 115, 130, 139, 143, 145, 195, 220, 221, 225, 235, 241, 244, 246, 248, 249, 250, 251, 255, 270, 291, 295, 298, 311, 312, 313, 314, 328, 334, 337, 341, 344, 351, 355, 370 }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.5 Maxima

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 30, 31, 32, 33, 36, 37, 38, 39, 40, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 96, 98, 99, 100, 101, 102, 105, 106, 107, 108, 109, 110, 111, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 222, 223, 224, 226, 227, 228, 229, 230, 231, 232, 233, 234, 236, 237, 238, 239, 240, 242, 243, 247, 249, 250, 251, 252, 254, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 272, 273, 274, 275, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 292, 293, 294, 295, 296, 297, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 331, 332, 333, 334, 335, 336, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 356, 357, 358, 359, 360, 361, 362, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376 }

B grade { 29, 34, 35, 41, 95, 97, 103, 104, 112, 113, 220, 221, 225, 235, 241, 244, 245, 246, 248, 253, 255, 271, 276, 291, 298, 313, 328, 329, 355, 363 }

C grade { }

F normal fail { 330, 337 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.6 Giac

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 99, 100, 101, 102, 105, 106, 107, 108, 109, 110, 111, 112, 114, 115, 116, 117, 118, 119, 120, 122, 123, 125, 126, 127, 128, 129, 131, 132, 134, 135, 136, 137, 139, 140, 141, 142, 143, 144, 146, 147, 148, 149, 150, 151, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 196, 197, 198, 199, 200, 201, 202, 203, 204, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220,

221, 222, 223, 224, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 242, 243, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 256, 257, 258, 259, 260, 261, 262, 264, 265, 266, 267, 268, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 292, 294, 296, 297, 299, 300, 301, 302, 303, 304, 305, 307, 308, 309, 310, 311, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 345, 346, 347, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376 }

B grade { 11, 12, 29, 41, 97, 98, 103, 104, 113, 121, 124, 130, 133, 138, 145, 152, 195, 205, 225, 241, 244, 255, 263, 270, 291, 293, 295, 298, 306, 312, 328, 329, 344, 348, 363 }

C grade { }

F normal fail { }

F(-1) timedout fail { 269 }

F(-2) exception fail { }

2.1.7 Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 360, 361, 362, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376 }

C grade { }

F normal fail { }

F(-1) timeout fail { 147, 323, 359, 363 }

F(-2) exception fail { }

2.1.8 Sympy

A grade { 1, 2, 3, 4, 5, 6, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 38, 39, 40, 41, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 58, 59, 60, 61, 62, 63, 64, 65, 66, 68, 69, 70, 71, 72, 73, 77, 78, 79, 81, 83, 85, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 101, 103, 105, 107, 109, 110, 112, 114, 115, 116, 117, 118, 119, 120, 122, 123, 125, 126, 128, 129, 130, 131, 137, 138, 140, 144, 145, 146, 147, 148, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 221, 222, 223, 224, 227, 231, 232, 233, 234, 236, 237, 239, 240, 241, 242, 243, 244, 245, 246, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 267, 268, 269, 271, 272, 273, 275, 276, 277, 278, 279, 280, 281, 282, 284, 285, 286, 287, 288, 289, 290, 292, 293, 294, 295, 299, 300, 302, 303, 304, 305, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 321, 323, 325, 326, 327, 328, 329, 331, 332, 333, 335, 338, 339, 340, 342, 343, 345, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 360, 361, 362, 364, 366, 367, 368, 371, 373, 374, 375, 376 }

B grade { 7, 8, 37, 42, 57, 67, 75, 76, 80, 82, 84, 86, 87, 100, 102, 104, 106, 108, 111, 113, 127, 139, 142, 181, 195, 219, 225, 226, 230, 251, 270, 283, 291, 296, 297, 298, 306, 320, 330, 334, 341, 344, 370, 372 }

C grade { 121, 124, 132, 133, 134, 135, 136, 141, 143, 228, 229, 250, 266, 274, 324, 336, 346, 363, 369 }

F normal fail { 149, 220, 235, 238, 247, 248, 249, 301, 322, 359 }

F(-1) timeout fail { 74, 337, 365 }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	11	11	11	10	12	11	20
N.S.	1	1.00	1.00	1.00	1.00	0.91	1.09	1.00	1.82
time (sec)	N/A	0.114	0.001	0.020	0.193	0.242	0.014	0.279	0.371

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	3	3	2	2	2	2	2
N.S.	1	1.00	1.00	1.00	0.67	0.67	0.67	0.67	0.67
time (sec)	N/A	0.115	0.000	0.024	0.187	0.236	0.029	0.264	0.006

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	2	2	2	3	2
N.S.	1	1.00	1.00	1.50	1.00	1.00	1.00	1.50	1.00
time (sec)	N/A	0.109	0.000	0.012	0.189	0.238	0.033	0.270	0.002

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	9	8	8	8	8	8
N.S.	1	1.00	1.00	1.12	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.122	0.001	0.030	0.195	0.235	0.033	0.294	0.209

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	4	4	4	5	4	4	3	4	4
N.S.	1	1.00	1.00	1.25	1.00	1.00	0.75	1.00	1.00
time (sec)	N/A	0.129	0.001	0.045	0.205	0.243	0.033	0.270	0.021

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	2	2	2	2	2
N.S.	1	1.00	1.00	1.50	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.128	0.001	0.027	0.193	0.246	0.033	0.269	0.003

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	2	7	5	2	2
N.S.	1	1.00	1.00	1.50	1.00	3.50	2.50	1.00	1.00
time (sec)	N/A	0.146	0.000	0.174	0.180	0.239	0.038	0.261	0.033

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	4	4	4	5	6	8	7	6	4
N.S.	1	1.00	1.00	1.25	1.50	2.00	1.75	1.50	1.00
time (sec)	N/A	0.139	0.009	0.175	0.180	0.248	0.035	0.300	0.011

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	4	4	3	4	12
N.S.	1	1.00	1.00	1.50	2.00	2.00	1.50	2.00	6.00
time (sec)	N/A	0.131	0.006	0.078	0.176	0.238	0.021	0.301	0.280

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	4	4	4	5	6	6	5	6	6
N.S.	1	1.00	1.00	1.25	1.50	1.50	1.25	1.50	1.50
time (sec)	N/A	0.142	0.006	0.053	0.185	0.234	0.038	0.287	0.252

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	2	2	2	11	2
N.S.	1	1.00	1.00	1.50	1.00	1.00	1.00	5.50	1.00
time (sec)	N/A	0.131	0.011	0.067	0.185	0.225	0.063	0.290	0.019

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	2	2	2	11	2
N.S.	1	1.00	1.00	1.50	1.00	1.00	1.00	5.50	1.00
time (sec)	N/A	0.131	0.008	0.056	0.178	0.225	0.064	0.300	0.016

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	5	5	6	3	11	5	6	5
N.S.	1	1.00	1.00	1.20	0.60	2.20	1.00	1.20	1.00
time (sec)	N/A	0.133	0.001	0.019	0.181	0.240	0.041	0.269	0.029

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	7	4	3	11	3	4	3
N.S.	1	1.00	2.33	1.33	1.00	3.67	1.00	1.33	1.00
time (sec)	N/A	0.137	0.001	0.029	0.181	0.245	0.036	0.278	0.002

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	9	8	8	7	8	8
N.S.	1	1.00	1.00	1.12	1.00	1.00	0.88	1.00	1.00
time (sec)	N/A	0.170	0.003	0.047	0.194	0.248	0.059	0.258	0.023

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	9	8	8	5	8	6
N.S.	1	1.00	1.00	1.12	1.00	1.00	0.62	1.00	0.75
time (sec)	N/A	0.123	0.000	0.013	0.192	0.229	0.030	0.271	0.019

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	21	12	12	11	11	10	11	11
N.S.	1	1.11	0.63	0.63	0.58	0.58	0.53	0.58	0.58
time (sec)	N/A	0.174	0.002	0.026	0.203	0.235	0.028	0.267	0.028

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	14	12	11	13	15	11	11
N.S.	1	1.00	0.74	0.63	0.58	0.68	0.79	0.58	0.58
time (sec)	N/A	0.144	0.008	0.076	0.187	0.243	0.086	0.275	0.003

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	12	13	13
N.S.	1	1.00	1.00	0.93	0.87	0.87	0.80	0.87	0.87
time (sec)	N/A	0.142	0.002	0.073	0.186	0.251	0.068	0.288	0.214

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	15	11	11	11	10	11	11
N.S.	1	1.00	0.75	0.55	0.55	0.55	0.50	0.55	0.55
time (sec)	N/A	0.147	0.013	0.037	0.184	0.225	0.032	0.269	0.023

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	7	7	7	8	7	7	7	7	7
N.S.	1	1.00	1.00	1.14	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.176	0.003	0.064	0.186	0.248	0.063	0.266	0.002

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	15	14	14	14	14	14
N.S.	1	1.00	1.00	0.83	0.78	0.78	0.78	0.78	0.78
time (sec)	N/A	0.179	0.033	0.136	0.184	0.248	0.066	0.269	0.028

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	13	13	12	13	9
N.S.	1	1.00	1.00	0.82	0.76	0.76	0.71	0.76	0.53
time (sec)	N/A	0.128	0.000	0.017	0.178	0.243	0.030	0.273	0.002

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	34	25	22	21	21	27	21	23
N.S.	1	1.17	0.86	0.76	0.72	0.72	0.93	0.72	0.79
time (sec)	N/A	0.252	0.032	0.191	0.179	0.258	0.090	0.268	0.073

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	25	21	21	21	24	21	24
N.S.	1	1.00	0.86	0.72	0.72	0.72	0.83	0.72	0.83
time (sec)	N/A	0.240	0.031	0.171	0.194	0.247	0.090	0.267	0.032

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	17	15	16	12	15	15	15	12
N.S.	1	1.13	1.00	1.07	0.80	1.00	1.00	1.00	0.80
time (sec)	N/A	0.142	0.000	0.017	0.179	0.236	0.040	0.263	0.032

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	14	14	12	14	14
N.S.	1	1.00	1.00	0.94	0.88	0.88	0.75	0.88	0.88
time (sec)	N/A	0.138	0.001	0.009	0.256	0.243	0.052	0.273	0.002

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	28	18	15	14	17	24	14	18
N.S.	1	1.22	0.78	0.65	0.61	0.74	1.04	0.61	0.78
time (sec)	N/A	0.179	0.004	0.102	0.173	0.239	0.086	0.263	0.048

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	9	74	18	8	103	8
N.S.	1	1.00	1.00	1.12	9.25	2.25	1.00	12.88	1.00
time (sec)	N/A	0.194	0.007	0.230	0.263	0.271	0.304	0.286	0.023

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	13	13	12	13	9
N.S.	1	1.00	1.00	0.82	0.76	0.76	0.71	0.76	0.53
time (sec)	N/A	0.136	0.003	0.026	0.176	0.238	0.032	0.278	0.031

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	31	17	17	16	16	15	16	16
N.S.	1	1.15	0.63	0.63	0.59	0.59	0.56	0.59	0.59
time (sec)	N/A	0.204	0.017	0.032	0.186	0.243	0.032	0.278	0.019

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	22	20	19	21	26	19	19
N.S.	1	1.00	0.81	0.74	0.70	0.78	0.96	0.70	0.70
time (sec)	N/A	0.158	0.056	0.124	0.179	0.249	0.090	0.322	0.028

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	20	18	17	21	20	17	17
N.S.	1	1.00	0.74	0.67	0.63	0.78	0.74	0.63	0.63
time (sec)	N/A	0.153	0.048	0.137	0.186	0.238	0.167	0.270	0.028

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	9	18	9	10	34	9	7	17	9
N.S.	1	2.00	1.00	1.11	3.78	1.00	0.78	1.89	1.00
time (sec)	N/A	0.185	0.013	0.083	0.196	0.237	0.069	0.289	0.020

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	19	57	18	20	30	18
N.S.	1	1.00	1.00	1.00	3.00	0.95	1.05	1.58	0.95
time (sec)	N/A	0.201	0.017	0.131	0.183	0.235	0.136	0.280	0.062

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	11	10	9	9	7	9	9
N.S.	1	1.00	0.69	0.62	0.56	0.56	0.44	0.56	0.56
time (sec)	N/A	0.155	0.002	0.026	0.177	0.234	0.028	0.283	0.021

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	15	14	13	14	105	13	9
N.S.	1	1.00	0.71	0.67	0.62	0.67	5.00	0.62	0.43
time (sec)	N/A	0.139	0.005	0.099	0.176	0.247	0.917	0.302	0.035

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	15	14	14	14	14	14
N.S.	1	1.00	1.00	0.83	0.78	0.78	0.78	0.78	0.78
time (sec)	N/A	0.193	0.031	0.130	0.187	0.248	0.060	0.294	0.023

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	29	16	15	14	14	12	14	14
N.S.	1	1.12	0.62	0.58	0.54	0.54	0.46	0.54	0.54
time (sec)	N/A	0.184	0.018	0.031	0.195	0.230	0.031	0.287	0.029

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	17	16	16	12	16	16
N.S.	1	1.00	1.00	0.94	0.89	0.89	0.67	0.89	0.89
time (sec)	N/A	0.144	0.004	0.015	0.277	0.253	0.056	0.279	0.183

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	9	9	9	10	104	20	8	52	9
N.S.	1	1.00	1.00	1.11	11.56	2.22	0.89	5.78	1.00
time (sec)	N/A	0.196	0.020	0.158	0.194	0.263	0.276	0.313	0.159

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	13	25	26	13	13
N.S.	1	1.00	1.00	0.82	0.76	1.47	1.53	0.76	0.76
time (sec)	N/A	0.160	0.032	0.257	0.176	0.243	0.129	0.274	0.056

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	9	13	14	22	8	13
N.S.	1	1.00	1.00	0.53	0.76	0.82	1.29	0.47	0.76
time (sec)	N/A	0.161	0.008	0.159	0.183	0.242	0.131	0.294	0.206

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	11	9	11	11	10	11	8
N.S.	1	1.00	1.00	0.82	1.00	1.00	0.91	1.00	0.73
time (sec)	N/A	0.169	0.005	0.288	0.183	0.246	0.199	0.280	0.222

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	14	12	11	11	10	11	11
N.S.	1	1.00	0.64	0.55	0.50	0.50	0.45	0.50	0.50
time (sec)	N/A	0.165	0.023	0.042	0.189	0.238	0.030	0.290	0.041

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	9	9	12	8	7	8	8
N.S.	1	1.00	0.60	0.60	0.80	0.53	0.47	0.53	0.53
time (sec)	N/A	0.151	0.027	0.030	0.191	0.224	0.033	0.294	0.035

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	14	15	14	14	14	14	14
N.S.	1	1.00	0.74	0.79	0.74	0.74	0.74	0.74	0.74
time (sec)	N/A	0.152	0.016	0.046	0.349	0.236	0.039	0.290	0.025

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	11	10	13	15	13	13
N.S.	1	1.00	1.00	0.65	0.59	0.76	0.88	0.76	0.76
time (sec)	N/A	0.142	0.003	0.073	0.208	0.241	0.118	0.277	0.206

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	25	16	16	11	11	20	11	11
N.S.	1	1.04	0.67	0.67	0.46	0.46	0.83	0.46	0.46
time (sec)	N/A	0.168	0.003	0.026	0.223	0.235	0.068	0.294	0.022

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	12	10	9	9	8	9	7
N.S.	1	1.00	0.86	0.71	0.64	0.64	0.57	0.64	0.50
time (sec)	N/A	0.129	0.000	0.017	0.222	0.245	0.029	0.278	0.030

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	13	12	13	15	13	13
N.S.	1	1.00	1.00	0.76	0.71	0.76	0.88	0.76	0.76
time (sec)	N/A	0.139	0.003	0.066	0.195	0.260	0.117	0.299	0.023

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	22	17	16	16	20	16	16
N.S.	1	1.00	1.00	0.77	0.73	0.73	0.91	0.73	0.73
time (sec)	N/A	0.205	0.007	0.066	0.207	0.244	0.099	0.286	0.264

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	17	20	17	16	16	15	16	16
N.S.	1	0.85	1.00	0.85	0.80	0.80	0.75	0.80	0.80
time (sec)	N/A	0.211	0.014	0.141	0.203	0.261	0.244	0.276	0.217

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	37	19	17	16	16	15	16	16
N.S.	1	1.32	0.68	0.61	0.57	0.57	0.54	0.57	0.57
time (sec)	N/A	0.196	0.025	0.040	0.278	0.232	0.036	0.295	0.031

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	20	16	16	15	13	15	15	14
N.S.	1	0.95	0.76	0.76	0.71	0.62	0.71	0.71	0.67
time (sec)	N/A	0.154	0.002	0.029	0.277	0.239	0.084	0.291	0.020

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	17	16	16	15	18	16
N.S.	1	1.00	1.00	0.94	0.89	0.89	0.83	1.00	0.89
time (sec)	N/A	0.195	0.019	0.276	0.191	0.248	0.128	0.280	0.026

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	15	14	13	14	105	13	9
N.S.	1	1.00	0.71	0.67	0.62	0.67	5.00	0.62	0.43
time (sec)	N/A	0.132	0.005	0.100	0.180	0.240	0.930	0.276	0.018

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	14	11	10	14	14	10	10
N.S.	1	1.00	0.78	0.61	0.56	0.78	0.78	0.56	0.56
time (sec)	N/A	0.151	0.038	0.125	0.199	0.237	0.025	0.283	0.052

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	14	11	10	10	10	10	10
N.S.	1	1.00	1.00	0.79	0.71	0.71	0.71	0.71	0.71
time (sec)	N/A	0.148	0.002	0.056	0.189	0.240	0.017	0.275	0.002

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	29	22	17	16	19	24	16	16
N.S.	1	1.21	0.92	0.71	0.67	0.79	1.00	0.67	0.67
time (sec)	N/A	0.192	0.002	0.233	0.187	0.242	0.017	0.282	0.034

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	15	11	11	11	8	11	10
N.S.	1	1.00	1.15	0.85	0.85	0.85	0.62	0.85	0.77
time (sec)	N/A	0.159	0.002	0.211	0.183	0.245	0.019	0.282	0.046

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	31	14	13	13	12	13	14
N.S.	1	1.00	1.82	0.82	0.76	0.76	0.71	0.76	0.82
time (sec)	N/A	0.194	0.058	0.202	0.186	0.238	0.020	0.284	0.040

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	31	14	13	22	12	13	14
N.S.	1	1.00	1.82	0.82	0.76	1.29	0.71	0.76	0.82
time (sec)	N/A	0.172	0.057	0.187	0.182	0.244	0.020	0.278	0.202

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	46	30	23	18	25	31	22	24
N.S.	1	1.28	0.83	0.64	0.50	0.69	0.86	0.61	0.67
time (sec)	N/A	0.264	0.009	0.172	0.179	0.253	0.019	0.281	0.043

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	29	14	11	10	19	14	10	18
N.S.	1	1.21	0.58	0.46	0.42	0.79	0.58	0.42	0.75
time (sec)	N/A	0.207	0.001	0.095	0.182	0.249	0.022	0.294	0.045

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	18	15	14	18	37	14	14
N.S.	1	1.00	0.82	0.68	0.64	0.82	1.68	0.64	0.64
time (sec)	N/A	0.159	0.061	0.204	0.181	0.240	0.068	0.295	0.394

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	20	15	14	31	37	14	18
N.S.	1	1.00	1.00	0.75	0.70	1.55	1.85	0.70	0.90
time (sec)	N/A	0.163	0.017	0.426	0.184	0.248	0.129	0.305	0.171

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	29	25	20	19	19	19	19	19
N.S.	1	1.16	1.00	0.80	0.76	0.76	0.76	0.76	0.76
time (sec)	N/A	0.187	0.034	0.228	0.186	0.264	0.022	0.294	0.054

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	44	30	23	24	25	36	22	22
N.S.	1	1.29	0.88	0.68	0.71	0.74	1.06	0.65	0.65
time (sec)	N/A	0.236	0.002	0.254	0.201	0.245	0.017	0.288	0.046

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	44	30	23	24	25	36	22	22
N.S.	1	1.29	0.88	0.68	0.71	0.74	1.06	0.65	0.65
time (sec)	N/A	0.241	0.009	0.264	0.186	0.249	0.018	0.281	0.039

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	56	30	23	18	33	41	22	37
N.S.	1	1.22	0.65	0.50	0.39	0.72	0.89	0.48	0.80
time (sec)	N/A	0.265	0.057	0.312	0.183	0.245	0.020	0.272	0.081

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	23	17	17	17	17	17	17
N.S.	1	1.00	1.10	0.81	0.81	0.81	0.81	0.81	0.81
time (sec)	N/A	0.159	0.002	0.230	0.188	0.239	0.021	0.273	0.038

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	61	22	17	16	31	31	16	32
N.S.	1	1.33	0.48	0.37	0.35	0.67	0.67	0.35	0.70
time (sec)	N/A	0.331	0.007	0.299	0.207	0.258	0.022	0.287	0.040

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	34	14	13	17	0	13	13
N.S.	1	1.00	1.62	0.67	0.62	0.81	0.00	0.62	0.62
time (sec)	N/A	0.175	0.071	0.201	0.209	0.252	0.000	0.272	0.114

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	18	14	13	14	170	13	25
N.S.	1	1.00	0.86	0.67	0.62	0.67	8.10	0.62	1.19
time (sec)	N/A	0.175	0.015	0.186	0.189	0.240	4.176	0.292	0.249

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	28	18	14	12	13	39	12	12
N.S.	1	1.47	0.95	0.74	0.63	0.68	2.05	0.63	0.63
time (sec)	N/A	0.187	0.034	0.082	0.217	0.249	0.108	0.281	0.303

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	18	19	15	15	15	22	15	14
N.S.	1	0.95	1.00	0.79	0.79	0.79	1.16	0.79	0.74
time (sec)	N/A	0.194	0.022	0.223	0.193	0.243	0.116	0.289	0.203

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	14	13	16	14	10	13	16
N.S.	1	1.00	1.00	0.93	1.14	1.00	0.71	0.93	1.14
time (sec)	N/A	0.174	0.008	1.193	0.197	0.259	0.038	0.276	0.233

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	18	20	29	20	37	20	28	32
N.S.	1	0.82	0.91	1.32	0.91	1.68	0.91	1.27	1.45
time (sec)	N/A	0.198	0.027	3.375	0.185	0.262	0.044	0.276	0.259

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	5	7	6	5	5	19	5	5
N.S.	1	1.00	1.40	1.20	1.00	1.00	3.80	1.00	1.00
time (sec)	N/A	0.170	0.008	0.184	0.195	0.253	0.139	0.277	0.203

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	25	11	15	17	8	10	10
N.S.	1	1.00	2.27	1.00	1.36	1.55	0.73	0.91	0.91
time (sec)	N/A	0.157	0.006	0.095	0.183	0.236	0.183	0.286	0.024

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	8	7	6	6	7	6	6
N.S.	1	1.00	1.33	1.17	1.00	1.00	1.17	1.00	1.00
time (sec)	N/A	0.146	0.001	0.026	0.285	0.257	0.021	0.282	0.027

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	16	13	12	12	19	12	12
N.S.	1	1.00	1.14	0.93	0.86	0.86	1.36	0.86	0.86
time (sec)	N/A	0.191	0.001	0.030	0.273	0.244	0.028	0.277	0.028

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	11	11	9	16	19	9	17
N.S.	1	1.00	1.00	1.00	0.82	1.45	1.73	0.82	1.55
time (sec)	N/A	0.167	0.005	0.260	0.187	0.246	0.020	0.280	0.029

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	19	15	22	31	15	27
N.S.	1	1.00	1.00	1.00	0.79	1.16	1.63	0.79	1.42
time (sec)	N/A	0.175	0.006	0.341	0.195	0.243	0.023	0.298	0.037

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	20	29	6	6
N.S.	1	1.00	1.00	0.88	0.75	2.50	3.62	0.75	0.75
time (sec)	N/A	0.163	0.002	0.547	0.189	0.245	0.021	0.315	0.023

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	27	14	13	20	29	13	13
N.S.	1	1.00	1.59	0.82	0.76	1.18	1.71	0.76	0.76
time (sec)	N/A	0.181	0.036	1.950	0.191	0.244	0.021	0.283	0.212

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	7	6	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.88	0.75	0.75
time (sec)	N/A	0.160	0.007	0.468	0.183	0.260	0.022	0.301	0.304

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	14	14	14	14	13
N.S.	1	1.00	1.00	0.82	0.82	0.82	0.82	0.82	0.76
time (sec)	N/A	0.183	0.039	1.060	0.205	0.236	0.044	0.276	0.446

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	22	23	34	24	20	22	18
N.S.	1	1.00	1.00	1.05	1.55	1.09	0.91	1.00	0.82
time (sec)	N/A	0.231	0.010	0.063	0.190	0.251	0.052	0.277	0.036

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	24	19	18	18	31	18	18
N.S.	1	1.00	1.09	0.86	0.82	0.82	1.41	0.82	0.82
time (sec)	N/A	0.230	0.006	0.062	0.271	0.252	0.027	0.285	0.036

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	16	20	20	22	20	17
N.S.	1	1.00	1.00	0.84	1.05	1.05	1.16	1.05	0.89
time (sec)	N/A	0.174	0.021	0.352	0.190	0.254	0.048	0.295	0.276

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	20	20	20	22	20	19
N.S.	1	1.00	1.00	0.80	0.80	0.80	0.88	0.80	0.76
time (sec)	N/A	0.187	0.024	2.250	0.188	0.244	0.052	0.313	0.540

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	10	6	7	6	18
N.S.	1	1.00	1.00	0.88	1.25	0.75	0.88	0.75	2.25
time (sec)	N/A	0.149	0.007	9.019	0.207	0.256	0.023	0.296	0.185

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	36	14	14	14	20
N.S.	1	1.00	1.00	0.82	2.12	0.82	0.82	0.82	1.18
time (sec)	N/A	0.182	0.016	18.177	0.189	0.267	0.045	0.280	0.188

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	10	6	7	6	6
N.S.	1	1.00	1.00	0.88	1.25	0.75	0.88	0.75	0.75
time (sec)	N/A	0.152	0.001	0.171	0.187	0.248	0.022	0.277	0.047

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	24	27	34	27	29	30
N.S.	1	1.00	1.00	1.50	1.69	2.12	1.69	1.81	1.88
time (sec)	N/A	0.203	0.010	0.131	0.192	0.257	0.051	0.295	0.310

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	18	13	10	20	8	18	8
N.S.	1	1.00	2.25	1.62	1.25	2.50	1.00	2.25	1.00
time (sec)	N/A	0.150	0.001	0.039	0.270	0.247	0.021	0.292	0.018

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	19	17	14	28	14	22	18
N.S.	1	1.00	1.36	1.21	1.00	2.00	1.00	1.57	1.29
time (sec)	N/A	0.207	0.001	0.122	0.196	0.254	0.044	0.285	0.025

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	37	14	14	39	41	14	14
N.S.	1	1.00	2.18	0.82	0.82	2.29	2.41	0.82	0.82
time (sec)	N/A	0.182	0.046	0.355	0.188	0.251	0.026	0.285	0.232

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	14	30	15	14	14
N.S.	1	1.00	1.00	0.82	0.82	1.76	0.88	0.82	0.82
time (sec)	N/A	0.189	0.010	0.241	0.192	0.253	0.048	0.300	0.222

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	5	17	6	8	19	15	6	5
N.S.	1	1.00	3.40	1.20	1.60	3.80	3.00	1.20	1.00
time (sec)	N/A	0.139	0.003	0.049	0.178	0.251	0.067	0.285	0.047

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	47	18	27	44	27	54	16
N.S.	1	1.00	2.94	1.12	1.69	2.75	1.69	3.38	1.00
time (sec)	N/A	0.184	0.012	0.230	0.191	0.252	0.055	0.280	0.211

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	19	12	17	21	19	19	8
N.S.	1	1.00	2.38	1.50	2.12	2.62	2.38	2.38	1.00
time (sec)	N/A	0.161	0.016	0.108	0.181	0.241	0.044	0.282	0.176

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	17	12	14	25	20	14	17
N.S.	1	1.00	1.31	0.92	1.08	1.92	1.54	1.08	1.31
time (sec)	N/A	0.162	0.010	0.241	0.187	0.244	0.019	0.277	0.034

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	13	24	26	13	13
N.S.	1	1.00	1.00	0.82	0.76	1.41	1.53	0.76	0.76
time (sec)	N/A	0.162	0.008	0.273	0.185	0.249	0.127	0.276	0.077

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	13	13	22	13	13
N.S.	1	1.00	1.00	0.82	0.76	0.76	1.29	0.76	0.76
time (sec)	N/A	0.160	0.006	0.220	0.185	0.252	0.143	0.288	0.030

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	12	11	24	26	11	11
N.S.	1	1.00	1.00	0.80	0.73	1.60	1.73	0.73	0.73
time (sec)	N/A	0.162	0.007	0.232	0.184	0.257	0.136	0.278	0.225

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	9	13	14	24	8	13
N.S.	1	1.00	1.00	0.53	0.76	0.82	1.41	0.47	0.76
time (sec)	N/A	0.162	0.030	0.181	0.176	0.259	0.132	0.271	0.067

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	7	6	19
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.88	0.75	2.38
time (sec)	N/A	0.155	0.002	0.151	0.185	0.264	0.021	0.290	0.032

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	30	23	22	25	114	22	22
N.S.	1	1.00	1.00	0.77	0.73	0.83	3.80	0.73	0.73
time (sec)	N/A	0.195	0.002	1.353	0.189	0.260	0.995	0.270	0.343

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	11	8	6	11	5	7	9	6
N.S.	1	2.20	1.60	1.20	2.20	1.00	1.40	1.80	1.20
time (sec)	N/A	0.171	0.002	0.769	0.185	0.265	0.186	0.282	0.202

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	30	18	69	35	32	29	24
N.S.	1	1.00	2.00	1.20	4.60	2.33	2.13	1.93	1.60
time (sec)	N/A	0.205	0.029	0.569	0.276	0.258	0.525	0.317	0.490

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	14	13	16	14	10	18	16
N.S.	1	1.00	1.00	0.93	1.14	1.00	0.71	1.29	1.14
time (sec)	N/A	0.174	0.001	0.314	0.194	0.263	0.029	0.290	0.161

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	18	20	29	20	37	20	28	32
N.S.	1	0.82	0.91	1.32	0.91	1.68	0.91	1.27	1.45
time (sec)	N/A	0.197	0.001	1.089	0.185	0.252	0.042	0.296	0.186

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	7	6	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.88	0.75	0.75
time (sec)	N/A	0.155	0.002	0.419	0.190	0.236	0.024	0.281	0.002

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	14	14	14	14	13
N.S.	1	1.00	1.00	0.82	0.82	0.82	0.82	0.82	0.76
time (sec)	N/A	0.187	0.007	1.004	0.175	0.240	0.043	0.281	0.002

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	37	26	21	35	15	39	21
N.S.	1	1.00	1.48	1.04	0.84	1.40	0.60	1.56	0.84
time (sec)	N/A	0.133	0.053	0.429	0.262	0.238	0.096	0.290	0.039

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	13	12	14	12	19	12
N.S.	1	1.00	1.00	0.81	0.75	0.88	0.75	1.19	0.75
time (sec)	N/A	0.123	0.030	0.141	0.268	0.233	0.392	0.277	0.030

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	9	9	9	8	7	7	7	7	7
N.S.	1	1.00	1.00	0.89	0.78	0.78	0.78	0.78	0.78
time (sec)	N/A	0.117	0.001	0.125	0.191	0.244	0.061	0.274	0.035

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	46	15	18	18	19	37	14
N.S.	1	1.00	2.88	0.94	1.12	1.12	1.19	2.31	0.88
time (sec)	N/A	0.126	0.004	0.166	0.174	0.247	0.522	0.276	0.085

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	35	22	19	26	18	27	23	24
N.S.	1	1.13	0.71	0.61	0.84	0.58	0.87	0.74	0.77
time (sec)	N/A	0.148	0.025	0.141	0.276	0.248	0.209	0.276	0.050

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	40	24	21	42	20	23	38
N.S.	1	1.00	1.48	0.89	0.78	1.56	0.74	0.85	1.41
time (sec)	N/A	0.149	0.086	0.234	0.259	0.268	0.280	0.303	0.238

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	14	14	27	33	14
N.S.	1	1.00	1.00	0.94	0.88	0.88	1.69	2.06	0.88
time (sec)	N/A	0.125	0.035	0.142	0.266	0.240	0.412	0.294	0.272

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	35	27	19	26	23	39	30	23
N.S.	1	1.13	0.87	0.61	0.84	0.74	1.26	0.97	0.74
time (sec)	N/A	0.149	0.021	0.149	0.265	0.230	0.143	0.291	0.037

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	11	11	8	11	11
N.S.	1	1.00	1.00	0.92	0.85	0.85	0.62	0.85	0.85
time (sec)	N/A	0.119	0.001	0.144	0.196	0.239	0.062	0.282	0.047

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	12	11	16	24	11	11
N.S.	1	1.00	1.00	0.80	0.73	1.07	1.60	0.73	0.73
time (sec)	N/A	0.125	0.003	0.138	0.209	0.249	0.088	0.278	0.029

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	41	20	19	32	19	19	18
N.S.	1	1.00	1.64	0.80	0.76	1.28	0.76	0.76	0.72
time (sec)	N/A	0.133	0.057	0.444	0.290	0.233	0.083	0.266	0.034

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	29	18	15	22	14	24	19	14
N.S.	1	1.16	0.72	0.60	0.88	0.56	0.96	0.76	0.56
time (sec)	N/A	0.145	0.017	0.136	0.302	0.228	0.111	0.277	0.024

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	16	5	4	14	3	25	4
N.S.	1	1.00	2.67	0.83	0.67	2.33	0.50	4.17	0.67
time (sec)	N/A	0.115	0.017	0.141	0.285	0.244	0.060	0.283	0.032

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	33	16	15	25	15	25	15
N.S.	1	1.00	1.57	0.76	0.71	1.19	0.71	1.19	0.71
time (sec)	N/A	0.125	0.028	0.197	0.289	0.241	0.084	0.282	0.033

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	39	35	26	22	33	66	25	25
N.S.	1	1.11	1.00	0.74	0.63	0.94	1.89	0.71	0.71
time (sec)	N/A	0.143	0.036	0.332	0.317	0.245	1.144	0.275	0.378

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	23	20	19	23	76	48	19
N.S.	1	1.00	1.00	0.87	0.83	1.00	3.30	2.09	0.83
time (sec)	N/A	0.132	0.050	0.167	0.294	0.242	0.427	0.300	0.388

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	36	30	25	19	28	92	24	24
N.S.	1	1.20	1.00	0.83	0.63	0.93	3.07	0.80	0.80
time (sec)	N/A	0.147	0.026	0.315	0.278	0.242	0.813	0.279	0.379

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	15	14	18	37	23	14
N.S.	1	1.00	1.00	0.83	0.78	1.00	2.06	1.28	0.78
time (sec)	N/A	0.136	0.031	0.174	0.296	0.240	0.448	0.284	0.305

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	34	34	31	22	58	49	24	34
N.S.	1	1.00	1.00	0.91	0.65	1.71	1.44	0.71	1.00
time (sec)	N/A	0.149	0.118	0.218	0.288	0.242	0.842	0.313	0.272

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	43	23	22	29	24	22	22
N.S.	1	1.00	1.48	0.79	0.76	1.00	0.83	0.76	0.76
time (sec)	N/A	0.132	0.074	0.428	0.277	0.246	0.086	0.398	0.046

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	23	18	14	24	15	37	18
N.S.	1	1.00	1.00	0.78	0.61	1.04	0.65	1.61	0.78
time (sec)	N/A	0.138	0.047	0.176	0.274	0.249	0.537	0.272	0.059

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	10	9	21	26	9	9
N.S.	1	1.00	1.00	0.77	0.69	1.62	2.00	0.69	0.69
time (sec)	N/A	0.124	0.003	0.134	0.188	0.232	0.398	0.285	0.222

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	35	22	19	26	23	44	32	23
N.S.	1	1.13	0.71	0.61	0.84	0.74	1.42	1.03	0.74
time (sec)	N/A	0.151	0.023	0.153	0.284	0.246	0.149	0.284	0.187

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	45	50	46	32	31	39	110	26	27
N.S.	1	1.11	1.02	0.71	0.69	0.87	2.44	0.58	0.60
time (sec)	N/A	0.144	0.104	0.241	0.335	0.237	1.746	0.276	0.047

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	10	9	9	26	9	9
N.S.	1	1.00	1.00	0.77	0.69	0.69	2.00	0.69	0.69
time (sec)	N/A	0.122	0.002	0.131	0.198	0.235	0.075	0.290	0.035

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	13	12	30	34	12	12
N.S.	1	1.00	1.00	0.81	0.75	1.88	2.12	0.75	0.75
time (sec)	N/A	0.126	0.035	0.164	0.440	0.232	0.458	0.394	0.297

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	37	47	25	36	35	22	23	24
N.S.	1	1.12	1.42	0.76	1.09	1.06	0.67	0.70	0.73
time (sec)	N/A	0.151	0.066	0.208	0.303	0.229	0.217	0.282	0.232

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	10	20	7	6	18	5	34	14
N.S.	1	1.25	2.50	0.88	0.75	2.25	0.62	4.25	1.75
time (sec)	N/A	0.129	0.066	0.237	0.373	0.240	0.281	0.295	0.249

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	24	21	22	20	22	41	20
N.S.	1	1.00	0.96	0.84	0.88	0.80	0.88	1.64	0.80
time (sec)	N/A	0.139	0.069	0.197	0.351	0.238	0.278	0.311	0.324

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	49	55	27	36	35	24	25	0
N.S.	1	1.11	1.25	0.61	0.82	0.80	0.55	0.57	0.00
time (sec)	N/A	0.172	0.072	0.427	0.332	0.262	0.262	0.273	0.000

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	23	24	22	28	19	22	23
N.S.	1	1.00	0.88	0.92	0.85	1.08	0.73	0.85	0.88
time (sec)	N/A	0.143	0.009	0.238	0.331	0.244	0.053	0.286	0.184

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	43	36	59	49	0	36	29
N.S.	1	1.00	1.00	0.84	1.37	1.14	0.00	0.84	0.67
time (sec)	N/A	0.148	0.208	0.192	0.234	0.246	0.000	0.320	0.056

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	45	23	22	35	24	22	22
N.S.	1	1.00	1.36	0.70	0.67	1.06	0.73	0.67	0.67
time (sec)	N/A	0.160	0.089	0.059	0.372	0.251	0.353	0.283	0.225

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	36	30	23	22	22	24	22	34
N.S.	1	1.20	1.00	0.77	0.73	0.73	0.80	0.73	1.13
time (sec)	N/A	0.156	0.033	0.088	0.393	0.238	0.291	0.276	0.252

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	42	13	6	16	3	32	12
N.S.	1	1.00	3.00	0.93	0.43	1.14	0.21	2.29	0.86
time (sec)	N/A	0.124	0.003	0.167	0.234	0.242	0.523	0.285	0.203

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	10	15	13
N.S.	1	1.00	1.00	0.93	0.87	0.87	0.67	1.00	0.87
time (sec)	N/A	0.146	0.005	0.168	0.205	0.233	0.040	0.277	0.192

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	25	21	20	20	19	21	20
N.S.	1	1.00	0.96	0.81	0.77	0.77	0.73	0.81	0.77
time (sec)	N/A	0.161	0.007	0.135	0.222	0.241	0.025	0.295	0.036

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	18	19	19	19	22	19
N.S.	1	1.00	1.00	0.72	0.76	0.76	0.76	0.88	0.76
time (sec)	N/A	0.203	0.008	0.037	0.214	0.243	0.076	0.287	0.254

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	29	25	24	36	20	26	22
N.S.	1	1.00	0.97	0.83	0.80	1.20	0.67	0.87	0.73
time (sec)	N/A	0.185	0.018	0.042	0.224	0.228	0.041	0.272	0.053

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	23	18	17	17	17	18	21
N.S.	1	1.00	1.00	0.78	0.74	0.74	0.74	0.78	0.91
time (sec)	N/A	0.202	0.007	0.165	0.329	0.252	0.062	0.270	0.056

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	38	32	31	31	34	31	30
N.S.	1	1.00	1.00	0.84	0.82	0.82	0.89	0.82	0.79
time (sec)	N/A	0.185	0.012	0.424	0.327	0.242	0.053	0.321	0.275

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	103	103	93	72	77	136	88	74	96
N.S.	1	1.00	0.90	0.70	0.75	1.32	0.85	0.72	0.93
time (sec)	N/A	0.557	0.048	0.454	0.329	0.247	0.302	0.295	0.347

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	33	28	29	44	27	30	33
N.S.	1	1.00	1.00	0.85	0.88	1.33	0.82	0.91	1.00
time (sec)	N/A	0.206	0.022	0.168	0.303	0.237	0.060	0.282	0.238

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	16	16	15	19	12	15	16
N.S.	1	1.00	0.84	0.84	0.79	1.00	0.63	0.79	0.84
time (sec)	N/A	0.128	0.006	0.144	0.303	0.249	0.042	0.277	0.032

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	14	13	13	12	15	12
N.S.	1	1.00	1.00	0.74	0.68	0.68	0.63	0.79	0.63
time (sec)	N/A	0.128	0.004	0.152	0.222	0.238	0.038	0.272	0.100

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	21	21	14	15	15	17	17	8
N.S.	1	1.11	1.11	0.74	0.79	0.79	0.89	0.89	0.42
time (sec)	N/A	0.154	0.004	0.170	0.224	0.238	0.050	0.261	0.136

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	32	25	26	37	26	43	22
N.S.	1	1.00	1.00	0.78	0.81	1.16	0.81	1.34	0.69
time (sec)	N/A	0.175	0.019	0.166	0.226	0.253	0.079	0.274	0.110

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	33	30	34	53	32	31	29
N.S.	1	1.00	0.77	0.70	0.79	1.23	0.74	0.72	0.67
time (sec)	N/A	0.193	0.026	0.152	0.208	0.234	0.086	0.282	0.250

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	21	17	19	26	17	21	16
N.S.	1	1.00	1.00	0.81	0.90	1.24	0.81	1.00	0.76
time (sec)	N/A	0.175	0.003	0.148	0.214	0.234	0.042	0.270	0.044

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	20	23	19	17	26	19
N.S.	1	1.00	1.00	0.80	0.92	0.76	0.68	1.04	0.76
time (sec)	N/A	0.214	0.008	0.160	0.221	0.225	0.037	0.276	0.197

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	19	11	12	13	11	8	13	11
N.S.	1	1.73	1.00	1.09	1.18	1.00	0.73	1.18	1.00
time (sec)	N/A	0.149	0.005	0.142	0.217	0.236	0.037	0.274	0.218

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	36	32	31	31	36	31	56
N.S.	1	1.00	1.00	0.89	0.86	0.86	1.00	0.86	1.56
time (sec)	N/A	0.276	0.017	0.235	0.301	0.240	0.095	0.271	0.120

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	29	22	21	33	22	21	23
N.S.	1	1.00	1.00	0.76	0.72	1.14	0.76	0.72	0.79
time (sec)	N/A	0.266	0.018	0.200	0.291	0.239	0.071	0.265	0.192

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	60	60	54	48	47	60	63	60	61
N.S.	1	1.00	0.90	0.80	0.78	1.00	1.05	1.00	1.02
time (sec)	N/A	0.245	0.055	0.377	0.365	0.249	0.131	0.276	0.128

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	42	28	25	30	39	27	25	30
N.S.	1	1.14	0.76	0.68	0.81	1.05	0.73	0.68	0.81
time (sec)	N/A	0.146	0.015	0.170	0.307	0.224	0.051	0.288	0.191

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	104	78	68	75	134	88	71	84
N.S.	1	1.07	0.80	0.70	0.77	1.38	0.91	0.73	0.87
time (sec)	N/A	0.252	0.043	0.721	0.292	0.236	0.116	0.277	0.260

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	46	34	35	39	46	36	49
N.S.	1	1.00	1.00	0.74	0.76	0.85	1.00	0.78	1.07
time (sec)	N/A	0.231	0.028	0.051	0.295	0.233	0.072	0.265	0.266

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	48	53	48	38	37	46	48	38	51
N.S.	1	1.10	1.00	0.79	0.77	0.96	1.00	0.79	1.06
time (sec)	N/A	0.204	0.014	0.164	0.307	0.239	0.069	0.284	0.093

Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	19	14	13	13	10	14	13
N.S.	1	1.00	1.27	0.93	0.87	0.87	0.67	0.93	0.87
time (sec)	N/A	0.133	0.004	0.128	0.209	0.231	0.023	0.289	0.027

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	8	9	8	8	7	9	8
N.S.	1	1.00	0.80	0.90	0.80	0.80	0.70	0.90	0.80
time (sec)	N/A	0.135	0.002	0.144	0.204	0.231	0.026	0.267	0.030

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	11	11	10	13	11
N.S.	1	1.00	1.00	0.92	0.85	0.85	0.77	1.00	0.85
time (sec)	N/A	0.135	0.005	0.145	0.204	0.231	0.046	0.268	0.188

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	11	12	11	11	8	13	10
N.S.	1	1.00	1.00	1.09	1.00	1.00	0.73	1.18	0.91
time (sec)	N/A	0.122	0.005	0.135	0.198	0.238	0.053	0.279	0.081

Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	15	11	12	12	10	13	10
N.S.	1	1.00	1.25	0.92	1.00	1.00	0.83	1.08	0.83
time (sec)	N/A	0.135	0.004	0.135	0.195	0.234	0.030	0.271	0.187

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	19	21	26	20	80	28	18
N.S.	1	1.00	0.73	0.81	1.00	0.77	3.08	1.08	0.69
time (sec)	N/A	0.135	0.009	0.193	0.207	0.245	0.116	0.264	0.249

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	14	13	12	12	10	14	12
N.S.	1	1.00	1.00	0.93	0.86	0.86	0.71	1.00	0.86
time (sec)	N/A	0.156	0.004	0.148	0.229	0.242	0.041	0.273	0.050

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	26	19	18	18	17	20	14
N.S.	1	1.00	1.00	0.73	0.69	0.69	0.65	0.77	0.54
time (sec)	N/A	0.166	0.006	0.190	0.193	0.241	0.039	0.268	0.041

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	14	15	14	17	10	15	14
N.S.	1	1.00	1.00	1.07	1.00	1.21	0.71	1.07	1.00
time (sec)	N/A	0.138	0.005	0.128	0.191	0.238	0.032	0.266	0.031

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	23	18	19	19	19	22	17
N.S.	1	1.00	1.00	0.78	0.83	0.83	0.83	0.96	0.74
time (sec)	N/A	0.144	0.007	0.157	0.198	0.237	0.064	0.266	0.086

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	16	15	15	15	18	15
N.S.	1	1.00	1.00	0.94	0.88	0.88	0.88	1.06	0.88
time (sec)	N/A	0.200	0.007	0.042	0.191	0.252	0.058	0.267	0.067

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	13	12	16	8	13	12
N.S.	1	1.00	1.00	1.08	1.00	1.33	0.67	1.08	1.00
time (sec)	N/A	0.141	0.005	0.167	0.191	0.233	0.028	0.274	0.034

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	22	21	20	26	19	21	22
N.S.	1	1.00	0.73	0.70	0.67	0.87	0.63	0.70	0.73
time (sec)	N/A	0.149	0.010	0.154	0.198	0.235	0.057	0.277	0.063

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	28	26	21	20	27	22	26	22
N.S.	1	1.00	0.93	0.75	0.71	0.96	0.79	0.93	0.79
time (sec)	N/A	0.147	0.019	0.155	0.224	0.237	0.048	0.268	0.238

Problem 190	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	14	15	14	18	14	16	14
N.S.	1	1.00	1.00	1.07	1.00	1.29	1.00	1.14	1.00
time (sec)	N/A	0.171	0.005	0.153	0.195	0.241	0.059	0.267	0.043

Problem 191	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	25	18	17	17	17	20	17
N.S.	1	1.00	1.32	0.95	0.89	0.89	0.89	1.05	0.89
time (sec)	N/A	0.176	0.008	0.037	0.196	0.240	0.056	0.294	0.195

Problem 192	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	12	14	13
N.S.	1	1.00	1.00	0.93	0.87	0.87	0.80	0.93	0.87
time (sec)	N/A	0.143	0.006	0.024	0.194	0.235	0.033	0.289	0.052

Problem 193	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	24	27	40	20	30	27
N.S.	1	1.00	1.00	0.96	1.08	1.60	0.80	1.20	1.08
time (sec)	N/A	0.147	0.013	0.184	0.200	0.236	0.045	0.280	0.051

Problem 194	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	21	17	22	31	19	18	21
N.S.	1	1.00	1.00	0.81	1.05	1.48	0.90	0.86	1.00
time (sec)	N/A	0.141	0.011	0.129	0.184	0.244	0.035	0.343	0.026

Problem 195	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	C	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	22	16	16	20	15	18	8
N.S.	1	1.00	2.75	2.00	2.00	2.50	1.88	2.25	1.00
time (sec)	N/A	0.128	0.004	0.148	0.206	0.249	0.041	0.273	0.180

Problem 196	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	10	13	13
N.S.	1	1.00	1.00	0.93	0.87	0.87	0.67	0.87	0.87
time (sec)	N/A	0.140	0.006	0.030	0.189	0.244	0.036	0.270	0.046

Problem 197	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	16	18	15	14	14	12	14	14
N.S.	1	0.89	1.00	0.83	0.78	0.78	0.67	0.78	0.78
time (sec)	N/A	0.139	0.004	0.130	0.207	0.238	0.026	0.278	0.027

Problem 198	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	20	19	18	18	17	18	18
N.S.	1	1.00	1.00	0.95	0.90	0.90	0.85	0.90	0.90
time (sec)	N/A	0.156	0.005	0.235	0.274	0.238	0.047	0.290	0.221

Problem 199	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	31	27	26	26	34	26	28
N.S.	1	1.00	1.00	0.87	0.84	0.84	1.10	0.84	0.90
time (sec)	N/A	0.161	0.008	0.331	0.279	0.230	0.045	0.286	0.188

Problem 200	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	31	22	21	21	22	21	17
N.S.	1	1.00	1.15	0.81	0.78	0.78	0.81	0.78	0.63
time (sec)	N/A	0.175	0.006	0.253	0.294	0.237	0.050	0.281	0.044

Problem 201	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	28	20	19	19	19	20	25
N.S.	1	1.00	1.22	0.87	0.83	0.83	0.83	0.87	1.09
time (sec)	N/A	0.188	0.009	0.181	0.282	0.242	0.066	0.295	0.055

Problem 202	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	28	28	24	23	23	29	24	55
N.S.	1	1.00	1.00	0.86	0.82	0.82	1.04	0.86	1.96
time (sec)	N/A	0.172	0.010	0.160	0.282	0.236	0.064	0.281	0.310

Problem 203	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	46	41	31	32	32	41	33	46
N.S.	1	1.12	1.00	0.76	0.78	0.78	1.00	0.80	1.12
time (sec)	N/A	0.185	0.006	0.145	0.275	0.256	0.060	0.271	0.065

Problem 204	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	47	42	34	35	35	42	36	47
N.S.	1	1.15	1.02	0.83	0.85	0.85	1.02	0.88	1.15
time (sec)	N/A	0.188	0.009	0.151	0.273	0.268	0.058	0.290	0.243

Problem 205	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	22	21	20	36	20	47	28
N.S.	1	1.00	0.92	0.88	0.83	1.50	0.83	1.96	1.17
time (sec)	N/A	0.186	0.016	0.184	0.280	0.238	0.068	0.290	0.045

Problem 206	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	26	19	18	18	19	20	10
N.S.	1	1.00	1.86	1.36	1.29	1.29	1.36	1.43	0.71
time (sec)	N/A	0.138	0.005	0.180	0.287	0.235	0.054	0.292	0.057

Problem 207	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	29	25	24	24	29	24	51
N.S.	1	1.00	1.00	0.86	0.83	0.83	1.00	0.83	1.76
time (sec)	N/A	0.293	0.015	0.174	0.295	0.248	0.087	0.295	0.079

Problem 208	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	23	18	17	17	19	17	33
N.S.	1	1.00	1.00	0.78	0.74	0.74	0.83	0.74	1.43
time (sec)	N/A	0.189	0.010	0.070	0.283	0.254	0.089	0.277	0.201

Problem 209	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	44	39	32	32	39	41	32	36
N.S.	1	1.13	1.00	0.82	0.82	1.00	1.05	0.82	0.92
time (sec)	N/A	0.156	0.025	0.634	0.302	0.239	0.055	0.285	0.044

Problem 210	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	18	10	11	14	18	8	14	10
N.S.	1	1.80	1.00	1.10	1.40	1.80	0.80	1.40	1.00
time (sec)	N/A	0.158	0.006	0.136	0.193	0.237	0.033	0.293	0.173

Problem 211	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	17	12	11	15	12	15	11
N.S.	1	1.00	1.55	1.09	1.00	1.36	1.09	1.36	1.00
time (sec)	N/A	0.212	0.199	0.296	0.188	0.261	0.093	0.319	0.088

Problem 212	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	82	18	17	17	19	17	17
N.S.	1	1.00	4.10	0.90	0.85	0.85	0.95	0.85	0.85
time (sec)	N/A	0.212	0.214	0.309	0.272	0.263	0.160	0.314	0.061

Problem 213	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	14	13	13	12	15	8
N.S.	1	1.00	1.00	0.74	0.68	0.68	0.63	0.79	0.42
time (sec)	N/A	0.146	0.003	0.175	0.202	0.238	0.045	0.331	0.078

Problem 214	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	12	11	11	10	13	6
N.S.	1	1.00	1.00	0.71	0.65	0.65	0.59	0.76	0.35
time (sec)	N/A	0.145	0.004	0.147	0.187	0.247	0.038	0.271	0.130

Problem 215	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	23	21	14	17	17	17	19	13
N.S.	1	1.10	1.00	0.67	0.81	0.81	0.81	0.90	0.62
time (sec)	N/A	0.152	0.006	0.181	0.196	0.238	0.045	0.267	0.239

Problem 216	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	49	49	44	27	37	44	46	40	36
N.S.	1	1.00	0.90	0.55	0.76	0.90	0.94	0.82	0.73
time (sec)	N/A	0.191	0.018	0.227	0.288	0.248	0.042	0.291	0.153

Problem 217	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	63	57	51	50	50	68	53	58
N.S.	1	1.00	0.90	0.81	0.79	0.79	1.08	0.84	0.92
time (sec)	N/A	0.247	0.028	0.074	0.291	0.260	0.199	0.277	0.337

Problem 218	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	86	86	67	54	59	103	65	59	71
N.S.	1	1.00	0.78	0.63	0.69	1.20	0.76	0.69	0.83
time (sec)	N/A	0.263	0.044	0.084	0.277	0.251	0.106	0.290	0.156

Problem 219	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	24	28	28	28	42	29	18
N.S.	1	1.00	1.00	1.17	1.17	1.17	1.75	1.21	0.75
time (sec)	N/A	0.137	0.020	0.142	0.227	0.253	0.506	0.282	0.041

Problem 220	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	200	205	126	124	272	638	0	139	223
N.S.	1	1.02	0.63	0.62	1.36	3.19	0.00	0.70	1.12
time (sec)	N/A	0.401	0.070	0.148	0.286	0.907	0.000	0.352	0.277

Problem 221	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	41	22	30	27	20	23	11
N.S.	1	1.00	2.28	1.22	1.67	1.50	1.11	1.28	0.61
time (sec)	N/A	0.178	0.030	0.221	0.200	0.249	0.126	0.290	0.493

Problem 222	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	15	15	14	15	14	14
N.S.	1	1.00	1.00	0.83	0.83	0.78	0.83	0.78	0.78
time (sec)	N/A	0.144	0.013	0.135	0.207	0.236	0.050	0.285	0.060

Problem 223	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	28	21	28	20	26	20	20
N.S.	1	1.00	0.88	0.66	0.88	0.62	0.81	0.62	0.62
time (sec)	N/A	0.159	0.021	0.143	0.202	0.231	0.052	0.269	0.028

Problem 224	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	13	12	12	14	12	12
N.S.	1	1.00	1.00	0.81	0.75	0.75	0.88	0.75	0.75
time (sec)	N/A	0.132	0.016	0.183	0.276	0.237	0.070	0.270	0.032

Problem 225	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	10	9	19	19	26	20	8
N.S.	1	1.00	1.00	0.90	1.90	1.90	2.60	2.00	0.80
time (sec)	N/A	0.125	0.015	0.147	0.210	0.246	0.346	0.276	0.171

Problem 226	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	25	11	17	8	22	18	8
N.S.	1	1.00	1.79	0.79	1.21	0.57	1.57	1.29	0.57
time (sec)	N/A	0.141	0.015	0.203	0.224	0.226	0.070	0.276	0.155

Problem 227	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	33	29	22	21	21	36	22	25
N.S.	1	1.06	0.94	0.71	0.68	0.68	1.16	0.71	0.81
time (sec)	N/A	0.207	0.019	0.119	0.204	0.242	0.657	0.278	0.202

Problem 228	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	21	17	22	17	76	22	19
N.S.	1	1.00	0.66	0.53	0.69	0.53	2.38	0.69	0.59
time (sec)	N/A	0.144	0.018	0.151	0.192	0.231	0.731	0.263	0.036

Problem 229	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	31	25	24	24	75	24	24
N.S.	1	1.00	1.00	0.81	0.77	0.77	2.42	0.77	0.77
time (sec)	N/A	0.135	0.033	0.199	0.314	0.240	0.758	0.269	0.164

Problem 230	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	31	22	20	19	14	117	19	12
N.S.	1	1.07	0.76	0.69	0.66	0.48	4.03	0.66	0.41
time (sec)	N/A	0.144	0.017	0.124	0.190	0.235	0.504	0.273	0.277

Problem 231	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	7	6	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.88	0.75	0.75
time (sec)	N/A	0.120	0.013	0.167	0.283	0.245	0.106	0.274	0.285

Problem 232	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	27	19	16	15	15	17	16	15
N.S.	1	1.29	0.90	0.76	0.71	0.71	0.81	0.76	0.71
time (sec)	N/A	0.152	0.016	0.142	0.200	0.261	0.055	0.274	0.216

Problem 233	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	34	28	23	22	22	26	23	22
N.S.	1	1.13	0.93	0.77	0.73	0.73	0.87	0.77	0.73
time (sec)	N/A	0.162	0.022	0.154	0.199	0.238	0.063	0.266	0.042

Problem 234	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	31	20	16	19	16	26	19	16
N.S.	1	1.15	0.74	0.59	0.70	0.59	0.96	0.70	0.59
time (sec)	N/A	0.142	0.020	0.147	0.210	0.237	0.539	0.258	0.294

Problem 235	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	B	F	A	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	201	208	127	132	293	547	0	140	208
N.S.	1	1.03	0.63	0.66	1.46	2.72	0.00	0.70	1.03
time (sec)	N/A	0.404	0.070	0.151	0.317	0.913	0.000	0.342	0.063

Problem 236	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	71	62	46	45	47	68	45	73
N.S.	1	1.15	1.00	0.74	0.73	0.76	1.10	0.73	1.18
time (sec)	N/A	0.232	0.063	0.135	0.280	0.251	0.180	0.263	0.194

Problem 237	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	130	140	117	83	82	76	121	82	82
N.S.	1	1.08	0.90	0.64	0.63	0.58	0.93	0.63	0.63
time (sec)	N/A	0.222	0.063	0.166	0.205	0.240	1.106	0.281	0.135

Problem 238	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	24	40	37	26	0	28	20
N.S.	1	1.00	1.00	1.67	1.54	1.08	0.00	1.17	0.83
time (sec)	N/A	0.155	0.032	0.190	0.278	0.253	0.000	0.277	0.188

Problem 239	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	11	12	11	13	10	12	9
N.S.	1	1.00	1.00	1.09	1.00	1.18	0.91	1.09	0.82
time (sec)	N/A	0.197	0.010	0.171	0.201	0.255	0.078	0.274	0.145

Problem 240	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	16	15	15	14	15	15
N.S.	1	1.00	1.00	0.94	0.88	0.88	0.82	0.88	0.88
time (sec)	N/A	0.187	0.043	0.067	0.195	0.259	0.064	0.273	0.215

Problem 241	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	10	21	21	22	21	9
N.S.	1	1.00	1.00	0.83	1.75	1.75	1.83	1.75	0.75
time (sec)	N/A	0.149	0.025	0.069	0.202	0.249	0.231	0.267	0.029

Problem 242	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	28	28	27	35	35	32	37	40
N.S.	1	1.00	1.00	0.96	1.25	1.25	1.14	1.32	1.43
time (sec)	N/A	0.158	0.034	0.086	0.202	0.250	0.396	0.272	0.205

Problem 243	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	33	43	22	30	27	20	23	11
N.S.	1	0.77	1.00	0.51	0.70	0.63	0.47	0.53	0.26
time (sec)	N/A	0.197	0.022	0.137	0.199	0.279	0.107	0.290	0.462

Problem 244	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	24	19	39	38	39	37	21
N.S.	1	1.00	1.14	0.90	1.86	1.81	1.86	1.76	1.00
time (sec)	N/A	0.170	0.022	0.143	0.375	0.256	0.226	0.307	0.392

Problem 245	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	24	16	25	17	14	17	11
N.S.	1	1.00	2.18	1.45	2.27	1.55	1.27	1.55	1.00
time (sec)	N/A	0.163	0.028	0.231	0.206	0.249	0.114	0.287	0.068

Problem 246	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	43	22	30	27	20	23	11
N.S.	1	1.00	2.39	1.22	1.67	1.50	1.11	1.28	0.61
time (sec)	N/A	0.169	0.028	0.224	0.323	0.271	0.150	0.291	0.499

Problem 247	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	35	24	25	35	0	28	16
N.S.	1	1.00	1.46	1.00	1.04	1.46	0.00	1.17	0.67
time (sec)	N/A	0.351	0.034	0.217	0.189	0.263	0.000	0.274	0.316

Problem 248	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	39	24	220	35	0	28	16
N.S.	1	1.00	1.62	1.00	9.17	1.46	0.00	1.17	0.67
time (sec)	N/A	0.193	0.035	0.358	0.212	0.274	0.000	0.281	0.353

Problem 249	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	24	23	33	0	25	22
N.S.	1	1.00	1.00	1.33	1.28	1.83	0.00	1.39	1.22
time (sec)	N/A	0.192	0.021	0.267	0.194	0.265	0.000	0.274	0.136

Problem 250	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	38	35	61	98	121	61	31
N.S.	1	1.00	1.06	0.97	1.69	2.72	3.36	1.69	0.86
time (sec)	N/A	0.192	0.061	0.282	0.297	0.273	1.557	0.298	0.792

Problem 251	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	16	15	43	71839	26	15
N.S.	1	1.00	1.00	1.07	1.00	2.87	4789.27	1.73	1.00
time (sec)	N/A	0.194	0.034	0.492	0.275	0.256	16.133	0.292	0.512

Problem 252	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	10	9	8	8	7	9	8
N.S.	1	1.00	0.83	0.75	0.67	0.67	0.58	0.75	0.67
time (sec)	N/A	0.121	0.002	0.144	0.203	0.231	0.031	0.291	0.046

Problem 253	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	12	26	11	12	11	11
N.S.	1	1.00	1.00	0.71	1.53	0.65	0.71	0.65	0.65
time (sec)	N/A	0.137	0.003	0.029	0.204	0.233	0.050	0.285	0.027

Problem 254	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	8	9	10	10	7	8	6
N.S.	1	1.00	0.67	0.75	0.83	0.83	0.58	0.67	0.50
time (sec)	N/A	0.154	0.005	0.065	0.211	0.281	0.162	0.281	0.211

Problem 255	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	24	27	34	27	29	30
N.S.	1	1.00	1.00	1.50	1.69	2.12	1.69	1.81	1.88
time (sec)	N/A	0.197	0.002	0.104	0.191	0.251	0.051	0.266	0.002

Problem 256	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	14	14	14	14	13
N.S.	1	1.00	1.00	0.82	0.82	0.82	0.82	0.82	0.76
time (sec)	N/A	0.182	0.008	1.012	0.195	0.238	0.051	0.257	0.002

Problem 257	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	25	16	16	11	11	20	11	11
N.S.	1	1.04	0.67	0.67	0.46	0.46	0.83	0.46	0.46
time (sec)	N/A	0.168	0.003	0.023	0.209	0.233	0.067	0.283	0.002

Problem 258	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	42	31	30	30	36	33	30
N.S.	1	1.00	1.00	0.74	0.71	0.71	0.86	0.79	0.71
time (sec)	N/A	0.212	0.008	0.046	0.196	0.256	0.072	0.274	0.225

Problem 259	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	7	6	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.88	0.75	0.75
time (sec)	N/A	0.142	0.004	0.042	0.192	0.237	0.121	0.260	0.070

Problem 260	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	11	10	10	8	11	10
N.S.	1	1.00	1.00	0.92	0.83	0.83	0.67	0.92	0.83
time (sec)	N/A	0.137	0.004	0.147	0.188	0.246	0.031	0.294	0.032

Problem 261	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	5	5	4	3	3	3	3	3
N.S.	1	1.00	1.00	0.80	0.60	0.60	0.60	0.60	0.60
time (sec)	N/A	0.145	0.007	0.036	0.218	0.263	0.253	0.271	0.032

Problem 262	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	29	14	11	10	19	14	10	18
N.S.	1	1.21	0.58	0.46	0.42	0.79	0.58	0.42	0.75
time (sec)	N/A	0.211	0.002	0.096	0.185	0.272	0.022	0.281	0.002

Problem 263	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	9	8	11	8	18	32
N.S.	1	1.00	1.00	1.12	1.00	1.38	1.00	2.25	4.00
time (sec)	N/A	0.170	0.050	0.304	0.190	0.274	0.064	0.295	0.500

Problem 264	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	11	11	8	11	11
N.S.	1	1.00	1.00	0.92	0.85	0.85	0.62	0.85	0.85
time (sec)	N/A	0.123	0.001	0.144	0.189	0.262	0.064	0.288	0.002

Problem 265	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	13	13	12	13	9
N.S.	1	1.00	1.00	0.82	0.76	0.76	0.71	0.76	0.53
time (sec)	N/A	0.141	0.003	0.029	0.192	0.258	0.039	0.287	0.228

Problem 266	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	24	19	18	18	109	18	18
N.S.	1	1.00	1.00	0.79	0.75	0.75	4.54	0.75	0.75
time (sec)	N/A	0.135	0.023	0.201	0.271	0.246	0.769	0.284	0.047

Problem 267	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	13	12	16	8	13	12
N.S.	1	1.00	1.00	1.08	1.00	1.33	0.67	1.08	1.00
time (sec)	N/A	0.134	0.004	0.141	0.189	0.245	0.025	0.286	0.037

Problem 268	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	17	16	17	16	16	15	16	16
N.S.	1	1.06	1.00	1.06	1.00	1.00	0.94	1.00	1.00
time (sec)	N/A	0.142	0.003	0.108	0.264	0.252	0.048	0.260	0.232

Problem 269	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	22	30	29	29	32	0	18
N.S.	1	1.00	1.00	1.36	1.32	1.32	1.45	0.00	0.82
time (sec)	N/A	0.193	0.042	0.063	0.192	0.232	0.630	0.000	0.287

Problem 270	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	29	56	43	19	46	54	42	42
N.S.	1	1.07	2.07	1.59	0.70	1.70	2.00	1.56	1.56
time (sec)	N/A	0.146	0.019	0.122	0.186	0.241	0.132	0.466	0.033

Problem 271	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	30	14	14	14	14
N.S.	1	1.00	1.00	0.82	1.76	0.82	0.82	0.82	0.82
time (sec)	N/A	0.186	0.014	2.787	0.200	0.246	0.050	0.381	0.241

Problem 272	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	22	17	16	16	15	16	16
N.S.	1	1.00	1.00	0.77	0.73	0.73	0.68	0.73	0.73
time (sec)	N/A	0.158	0.005	0.376	0.261	0.239	0.040	0.385	0.236

Problem 273	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	37	28	25	24	24	24	26	24
N.S.	1	1.16	0.88	0.78	0.75	0.75	0.75	0.81	0.75
time (sec)	N/A	0.178	0.010	0.017	0.274	0.269	0.087	0.327	0.029

Problem 274	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	36	30	25	35	28	66	40	30
N.S.	1	1.20	1.00	0.83	1.17	0.93	2.20	1.33	1.00
time (sec)	N/A	0.143	0.029	0.189	0.267	0.241	0.780	0.345	0.066

Problem 275	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	14	13	13	10	15	13
N.S.	1	1.00	1.00	1.08	1.00	1.00	0.77	1.15	1.00
time (sec)	N/A	0.138	0.004	0.182	0.195	0.245	0.035	0.345	0.053

Problem 276	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	25	14	17	44	14	17	27	16
N.S.	1	1.56	0.88	1.06	2.75	0.88	1.06	1.69	1.00
time (sec)	N/A	0.244	0.017	0.108	0.200	0.252	0.096	0.351	0.209

Problem 277	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	20	14	15	15	14	18	13
N.S.	1	1.00	1.18	0.82	0.88	0.88	0.82	1.06	0.76
time (sec)	N/A	0.138	0.007	0.032	0.192	0.236	0.037	0.348	0.074

Problem 278	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	3	3	3	3	3
N.S.	1	1.00	1.00	1.33	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.161	0.007	0.440	0.296	0.253	0.084	0.320	0.076

Problem 279	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	21	22	17	16	16	20	16	16
N.S.	1	0.95	1.00	0.77	0.73	0.73	0.91	0.73	0.73
time (sec)	N/A	0.209	0.007	0.073	0.196	0.250	0.105	0.330	0.297

Problem 280	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	9	9	9	10	9	9	7	9	9
N.S.	1	1.00	1.00	1.11	1.00	1.00	0.78	1.00	1.00
time (sec)	N/A	0.140	0.011	0.180	0.191	0.247	0.041	0.322	0.019

Problem 281	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	11	10	10	8	10	10
N.S.	1	1.00	1.00	0.92	0.83	0.83	0.67	0.83	0.83
time (sec)	N/A	0.159	0.003	0.036	0.187	0.237	0.038	0.328	0.049

Problem 282	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	22	20	19	21	26	19	19
N.S.	1	1.00	0.81	0.74	0.70	0.78	0.96	0.70	0.70
time (sec)	N/A	0.159	0.049	0.138	0.192	0.239	0.104	0.330	0.031

Problem 283	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	13	22	26	13	13
N.S.	1	1.00	1.00	0.82	0.76	1.29	1.53	0.76	0.76
time (sec)	N/A	0.164	0.008	0.272	0.199	0.248	0.140	0.321	0.068

Problem 284	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	20	19	19	19	20	25
N.S.	1	1.00	1.00	0.80	0.76	0.76	0.76	0.80	1.00
time (sec)	N/A	0.158	0.006	0.065	0.273	0.268	0.055	0.353	0.228

Problem 285	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	38	28	28	29	25	29	49	25
N.S.	1	0.97	0.72	0.72	0.74	0.64	0.74	1.26	0.64
time (sec)	N/A	0.167	0.012	0.066	0.202	0.245	0.044	0.325	0.043

Problem 286	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	16	14	13	13	12	13	13
N.S.	1	1.00	0.62	0.54	0.50	0.50	0.46	0.50	0.50
time (sec)	N/A	0.172	0.028	0.052	0.188	0.261	0.041	0.327	0.066

Problem 287	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	18	11	10	10	12	10	10
N.S.	1	1.00	1.50	0.92	0.83	0.83	1.00	0.83	0.83
time (sec)	N/A	0.152	0.011	0.037	0.292	0.274	0.034	0.405	0.230

Problem 288	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	24	21	22	20	22	41	20
N.S.	1	1.00	0.96	0.84	0.88	0.80	0.88	1.64	0.80
time (sec)	N/A	0.135	0.069	0.223	0.273	0.254	0.310	0.324	0.336

Problem 289	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	26	23	22	21	21	20	21	21
N.S.	1	0.96	0.85	0.81	0.78	0.78	0.74	0.78	0.78
time (sec)	N/A	0.168	0.008	0.076	0.212	0.253	0.107	0.345	0.223

Problem 290	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	12	11	11	15	11	12
N.S.	1	1.00	1.00	0.63	0.58	0.58	0.79	0.58	0.63
time (sec)	N/A	0.134	0.004	0.048	0.215	0.257	0.384	0.346	0.026

Problem 291	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	6	6	19	15	19	16	15
N.S.	1	1.00	1.00	1.00	3.17	2.50	3.17	2.67	2.50
time (sec)	N/A	0.135	0.035	0.036	0.209	0.253	0.047	0.338	0.111

Problem 292	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	16	14	11	10	10	10	10	10
N.S.	1	1.14	1.00	0.79	0.71	0.71	0.71	0.71	0.71
time (sec)	N/A	0.147	0.006	0.063	0.278	0.232	0.046	0.558	0.247

Problem 293	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	9	8	8	10	32	8
N.S.	1	1.00	1.00	0.75	0.67	0.67	0.83	2.67	0.67
time (sec)	N/A	0.131	0.017	0.250	0.268	0.250	0.078	0.320	0.278

Problem 294	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	44	30	23	18	25	31	22	26
N.S.	1	1.29	0.88	0.68	0.53	0.74	0.91	0.65	0.76
time (sec)	N/A	0.253	0.010	0.315	0.194	0.257	0.027	0.354	0.045

Problem 295	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	23	7	8	29	7	26	6
N.S.	1	1.00	1.92	0.58	0.67	2.42	0.58	2.17	0.50
time (sec)	N/A	0.128	0.078	0.237	0.272	0.247	0.284	0.312	0.190

Problem 296	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	14	21	48	14	21
N.S.	1	1.00	1.00	0.94	0.88	1.31	3.00	0.88	1.31
time (sec)	N/A	0.213	0.019	0.088	0.206	0.260	1.218	0.301	0.131

Problem 297	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	7	7	23	13	5	7	12	7	5
N.S.	1	1.00	3.29	1.86	0.71	1.00	1.71	1.00	0.71
time (sec)	N/A	0.165	0.005	0.159	0.195	0.253	0.798	0.302	0.050

Problem 298	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	6	6	15	15	15	16	15
N.S.	1	1.00	1.00	1.00	2.50	2.50	2.50	2.67	2.50
time (sec)	N/A	0.147	0.001	0.044	0.236	0.249	0.053	0.285	0.027

Problem 299	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	51	43	33	32	32	41	33	46
N.S.	1	1.19	1.00	0.77	0.74	0.74	0.95	0.77	1.07
time (sec)	N/A	0.185	0.008	0.166	0.281	0.253	0.056	0.285	0.091

Problem 300	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	59	29	33	74	30	42	57	37
N.S.	1	1.59	0.78	0.89	2.00	0.81	1.14	1.54	1.00
time (sec)	N/A	0.453	0.019	0.136	0.200	0.241	0.279	0.284	0.238

Problem 301	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	9	9	9	8	7	12	0	7	27
N.S.	1	1.00	1.00	0.89	0.78	1.33	0.00	0.78	3.00
time (sec)	N/A	0.169	0.008	1.143	0.183	0.259	0.000	0.298	2.664

Problem 302	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	20	16	16	16	12	16	15
N.S.	1	1.00	1.00	0.80	0.80	0.80	0.60	0.80	0.75
time (sec)	N/A	0.140	0.000	0.020	0.204	0.228	0.019	0.319	0.026

Problem 303	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	15	13	10	9	9	8	10	9
N.S.	1	1.25	1.08	0.83	0.75	0.75	0.67	0.83	0.75
time (sec)	N/A	0.171	0.023	0.037	0.205	0.259	0.034	0.282	0.184

Problem 304	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	25	21	18	17	17	15	17	17
N.S.	1	1.19	1.00	0.86	0.81	0.81	0.71	0.81	0.81
time (sec)	N/A	0.135	0.006	0.162	0.203	0.244	0.044	0.296	0.053

Problem 305	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	33	43	22	30	27	20	23	11
N.S.	1	0.77	1.00	0.51	0.70	0.63	0.47	0.53	0.26
time (sec)	N/A	0.189	0.021	0.134	0.197	0.263	0.120	0.318	0.426

Problem 306	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	18	15	16	22	144	43	14
N.S.	1	1.00	0.75	0.62	0.67	0.92	6.00	1.79	0.58
time (sec)	N/A	0.142	0.020	0.167	0.181	0.257	0.628	0.319	0.212

Problem 307	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	41	24	20	16	16	34	16	21
N.S.	1	1.08	0.63	0.53	0.42	0.42	0.89	0.42	0.55
time (sec)	N/A	0.202	0.011	0.032	0.202	0.242	0.100	0.291	0.035

Problem 308	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	43	37	31	30	32	39	30	32
N.S.	1	1.16	1.00	0.84	0.81	0.86	1.05	0.81	0.86
time (sec)	N/A	0.218	0.050	0.234	0.278	0.251	0.601	0.300	0.208

Problem 309	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	16	15	21	14	17	15
N.S.	1	1.00	1.00	0.94	0.88	1.24	0.82	1.00	0.88
time (sec)	N/A	0.194	0.005	0.181	0.190	0.247	0.045	0.286	0.042

Problem 310	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	40	40	29	26	38	26	39	26	34
N.S.	1	1.00	0.72	0.65	0.95	0.65	0.98	0.65	0.85
time (sec)	N/A	0.240	0.089	0.220	0.201	0.252	0.093	0.272	0.198

Problem 311	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	5	5	6	5	20	5	5	5
N.S.	1	1.00	1.00	1.20	1.00	4.00	1.00	1.00	1.00
time (sec)	N/A	0.196	4.174	0.201	0.194	0.256	0.154	0.303	0.247

Problem 312	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	20	5	4	18	3	19	4
N.S.	1	1.00	3.33	0.83	0.67	3.00	0.50	3.17	0.67
time (sec)	N/A	0.118	0.035	0.422	0.300	0.240	0.068	0.279	0.008

Problem 313	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	37	24	22	62	62	61	22	29
N.S.	1	1.00	0.65	0.59	1.68	1.68	1.65	0.59	0.78
time (sec)	N/A	0.157	0.008	0.156	0.192	0.247	0.061	0.291	0.100

Problem 314	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	25	21	18	18	36	19	18	18
N.S.	1	1.19	1.00	0.86	0.86	1.71	0.90	0.86	0.86
time (sec)	N/A	0.199	0.028	0.177	0.215	0.248	0.049	0.299	0.395

Problem 315	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	30	25	24	22	41	24	24
N.S.	1	1.00	1.00	0.83	0.80	0.73	1.37	0.80	0.80
time (sec)	N/A	0.197	0.058	0.172	0.199	0.256	0.074	0.266	0.255

Problem 316	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	4	4	27	4	3	3	3	3	3
N.S.	1	1.00	6.75	1.00	0.75	0.75	0.75	0.75	0.75
time (sec)	N/A	0.160	0.007	0.188	0.207	0.260	0.219	0.277	0.236

Problem 317	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	17	13	12	15	11	10	15	11
N.S.	1	1.31	1.00	0.92	1.15	0.85	0.77	1.15	0.85
time (sec)	N/A	0.134	0.005	0.151	0.200	0.234	0.040	0.317	0.046

Problem 318	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	49	24	21	21	21	22	21	21
N.S.	1	1.11	0.55	0.48	0.48	0.48	0.50	0.48	0.48
time (sec)	N/A	0.230	0.022	0.046	0.210	0.233	0.042	0.296	0.027

Problem 319	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	45	42	28	27	25	37	27	27
N.S.	1	1.10	1.02	0.68	0.66	0.61	0.90	0.66	0.66
time (sec)	N/A	0.159	0.039	0.149	0.276	0.230	1.704	0.283	0.031

Problem 320	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	20	19	17	114	13	14
N.S.	1	1.00	1.00	0.80	0.76	0.68	4.56	0.52	0.56
time (sec)	N/A	0.197	0.002	1.335	0.185	0.263	0.962	0.316	0.225

Problem 321	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	11	10	10	7	10	8
N.S.	1	1.00	1.00	0.92	0.83	0.83	0.58	0.83	0.67
time (sec)	N/A	0.131	0.003	0.023	0.192	0.234	0.029	0.287	0.186

Problem 322	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	57	65	41	43	32	0	30	43
N.S.	1	1.39	1.59	1.00	1.05	0.78	0.00	0.73	1.05
time (sec)	N/A	0.154	0.101	0.190	0.275	0.251	0.000	0.322	0.050

Problem 323	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	A	A	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	34	41	27	119	27	27	29	28	0
N.S.	1	1.21	0.79	3.50	0.79	0.79	0.85	0.82	0.00
time (sec)	N/A	0.185	0.015	0.157	0.272	0.268	1.221	0.303	0.000

Problem 324	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	18	17	17	42	17	17
N.S.	1	1.00	1.00	0.95	0.89	0.89	2.21	0.89	0.89
time (sec)	N/A	0.138	0.006	0.184	0.275	0.261	0.044	0.294	0.042

Problem 325	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	45	30	39	37	34	31	28
N.S.	1	1.00	1.18	0.79	1.03	0.97	0.89	0.82	0.74
time (sec)	N/A	0.157	0.110	0.311	0.278	0.250	0.224	0.268	0.047

Problem 326	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	9	8	8	7	8	8
N.S.	1	1.00	1.00	0.75	0.67	0.67	0.58	0.67	0.67
time (sec)	N/A	0.131	0.006	0.210	0.278	0.246	0.048	0.295	0.199

Problem 327	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	31	27	26	26	36	26	28
N.S.	1	1.00	1.00	0.87	0.84	0.84	1.16	0.84	0.90
time (sec)	N/A	0.163	0.013	0.708	0.282	0.237	0.044	0.289	0.040

Problem 328	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	37	16	121	29	12	150	19
N.S.	1	1.00	3.70	1.60	12.10	2.90	1.20	15.00	1.90
time (sec)	N/A	0.180	0.014	0.124	0.280	0.257	0.442	0.361	0.112

Problem 329	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	27	29	26	22	30	13
N.S.	1	1.00	1.00	1.80	1.93	1.73	1.47	2.00	0.87
time (sec)	N/A	0.133	0.006	0.202	0.193	0.256	0.067	0.288	0.061

Problem 330	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	21	14	0	13	63	13	21
N.S.	1	1.00	1.00	0.67	0.00	0.62	3.00	0.62	1.00
time (sec)	N/A	0.138	0.103	0.029	0.000	0.232	0.160	0.279	0.237

Problem 331	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	25	18	16	19	17	19	18	17
N.S.	1	1.09	0.78	0.70	0.83	0.74	0.83	0.78	0.74
time (sec)	N/A	0.176	0.050	0.040	0.203	0.241	0.060	0.267	0.302

Problem 332	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	20	17	16	16	17	16	16
N.S.	1	1.00	1.00	0.85	0.80	0.80	0.85	0.80	0.80
time (sec)	N/A	0.157	0.009	0.037	0.206	0.261	0.126	0.274	0.313

Problem 333	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	16	18	19	14	20	18
N.S.	1	1.00	1.00	0.89	1.00	1.06	0.78	1.11	1.00
time (sec)	N/A	0.147	0.006	0.065	0.202	0.256	0.054	0.281	0.216

Problem 334	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	20	19	25	20	20	19
N.S.	1	1.00	1.00	1.67	1.58	2.08	1.67	1.67	1.58
time (sec)	N/A	0.157	0.046	0.052	0.192	0.243	0.051	0.285	0.249

Problem 335	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	23	9	10	15	12	16	8
N.S.	1	1.00	2.88	1.12	1.25	1.88	1.50	2.00	1.00
time (sec)	N/A	0.260	0.007	0.145	0.279	0.262	0.353	0.303	0.228

Problem 336	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	13	12	12	44	12	12
N.S.	1	1.00	1.00	0.72	0.67	0.67	2.44	0.67	0.67
time (sec)	N/A	0.125	0.018	0.308	0.274	0.242	0.486	0.295	0.146

Problem 337	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	15	11	10	0	24	0	9	18
N.S.	1	1.36	1.00	0.91	0.00	2.18	0.00	0.82	1.64
time (sec)	N/A	0.222	0.021	0.479	0.000	0.316	0.000	0.289	0.438

Problem 338	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	46	23	22	30	27	22	22
N.S.	1	1.00	1.53	0.77	0.73	1.00	0.90	0.73	0.73
time (sec)	N/A	0.135	0.077	0.465	0.310	0.256	0.107	0.328	0.177

Problem 339	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	27	20	23	21	21	26	21	23
N.S.	1	1.12	0.83	0.96	0.88	0.88	1.08	0.88	0.96
time (sec)	N/A	0.300	0.009	0.111	0.206	0.261	0.127	0.285	0.030

Problem 340	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	50	55	47	33	49	39	37	40	39
N.S.	1	1.10	0.94	0.66	0.98	0.78	0.74	0.80	0.78
time (sec)	N/A	0.167	0.084	0.373	0.283	0.249	0.250	0.286	0.228

Problem 341	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	11	10	9	46	51	9	9
N.S.	1	1.00	1.00	0.91	0.82	4.18	4.64	0.82	0.82
time (sec)	N/A	0.118	0.002	0.155	0.198	0.244	0.028	0.266	0.224

Problem 342	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	31	20	19	19	20	19	19
N.S.	1	1.00	1.24	0.80	0.76	0.76	0.80	0.76	0.76
time (sec)	N/A	0.187	0.011	0.175	0.201	0.242	0.022	0.279	0.190

Problem 343	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	22	20	19	21	26	19	19
N.S.	1	1.00	0.81	0.74	0.70	0.78	0.96	0.70	0.70
time (sec)	N/A	0.154	0.050	0.151	0.191	0.242	0.167	0.272	0.030

Problem 344	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	41	23	34	56	36	70	18
N.S.	1	1.00	1.71	0.96	1.42	2.33	1.50	2.92	0.75
time (sec)	N/A	0.200	0.043	0.295	0.192	0.248	0.057	0.282	0.070

Problem 345	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	34	35	32	33	35	27	44	32
N.S.	1	1.00	1.03	0.94	0.97	1.03	0.79	1.29	0.94
time (sec)	N/A	0.144	0.042	0.280	0.316	0.232	0.098	0.289	0.495

Problem 346	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	36	30	25	35	28	73	38	37
N.S.	1	1.20	1.00	0.83	1.17	0.93	2.43	1.27	1.23
time (sec)	N/A	0.144	0.028	0.218	0.275	0.232	0.841	0.288	0.111

Problem 347	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	37	19	16	16	16	15	16	16
N.S.	1	1.16	0.59	0.50	0.50	0.50	0.47	0.50	0.50
time (sec)	N/A	0.189	0.018	0.047	0.208	0.238	0.036	0.295	0.033

Problem 348	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	31	18	17	12	26	42	12
N.S.	1	1.00	1.35	0.78	0.74	0.52	1.13	1.83	0.52
time (sec)	N/A	0.201	0.025	0.158	0.194	0.254	0.119	0.282	0.113

Problem 349	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	24	24	22	21	21	19	21	21
N.S.	1	0.89	0.89	0.81	0.78	0.78	0.70	0.78	0.78
time (sec)	N/A	0.201	0.008	0.023	0.285	0.254	0.072	0.267	0.290

Problem 350	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	43	32	31	53	28	29	32	28
N.S.	1	1.13	0.84	0.82	1.39	0.74	0.76	0.84	0.74
time (sec)	N/A	0.185	0.014	0.037	0.279	0.248	0.144	0.275	0.234

Problem 351	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	5	5	5	4	16	8	6	6
N.S.	1	1.00	1.00	1.00	0.80	3.20	1.60	1.20	1.20
time (sec)	N/A	0.189	0.008	0.319	0.202	0.244	0.375	0.270	0.048

Problem 352	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	34	25	22	21	21	27	21	23
N.S.	1	1.17	0.86	0.76	0.72	0.72	0.93	0.72	0.79
time (sec)	N/A	0.258	0.018	0.194	0.206	0.274	0.088	0.285	0.202

Problem 353	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	45	29	36	47	27	26	27
N.S.	1	1.00	1.25	0.81	1.00	1.31	0.75	0.72	0.75
time (sec)	N/A	0.148	0.090	0.240	0.294	0.251	0.283	0.287	0.187

Problem 354	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	34	31	23	22	22	24	22	22
N.S.	1	1.21	1.11	0.82	0.79	0.79	0.86	0.79	0.79
time (sec)	N/A	0.150	0.012	0.210	0.292	0.254	0.056	0.301	0.056

Problem 355	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	31	26	25	42	43	46	38	29
N.S.	1	1.19	1.00	0.96	1.62	1.65	1.77	1.46	1.12
time (sec)	N/A	0.255	0.006	0.362	0.205	0.253	0.065	0.272	0.059

Problem 356	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	56	30	23	24	33	46	22	22
N.S.	1	1.22	0.65	0.50	0.52	0.72	1.00	0.48	0.48
time (sec)	N/A	0.249	0.039	0.316	0.210	0.256	0.025	0.291	0.221

Problem 357	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	15	17	16	24	17	16	11
N.S.	1	1.00	0.75	0.85	0.80	1.20	0.85	0.80	0.55
time (sec)	N/A	0.206	0.017	0.419	0.204	0.274	0.641	0.260	0.286

Problem 358	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	24	24	18	16	24	14	19	19
N.S.	1	1.14	1.14	0.86	0.76	1.14	0.67	0.90	0.90
time (sec)	N/A	0.169	0.032	0.051	0.211	0.253	0.048	0.282	0.068

Problem 359	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	37	33	77	47	58	0	50	0
N.S.	1	1.00	0.89	2.08	1.27	1.57	0.00	1.35	0.00
time (sec)	N/A	0.204	0.024	0.678	0.291	0.307	0.000	0.287	0.000

Problem 360	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	28	46	26	29	27	29	28	23
N.S.	1	1.00	1.64	0.93	1.04	0.96	1.04	1.00	0.82
time (sec)	N/A	0.147	0.058	0.254	0.192	0.247	0.227	0.292	0.145

Problem 361	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	17	15	18	17	15	21
N.S.	1	1.00	1.00	0.89	0.79	0.95	0.89	0.79	1.11
time (sec)	N/A	0.169	0.008	0.292	0.213	0.250	0.020	0.298	0.035

Problem 362	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	55	26	25	24	24	22	24	24
N.S.	1	1.20	0.57	0.54	0.52	0.52	0.48	0.52	0.52
time (sec)	N/A	0.253	0.021	0.038	0.195	0.245	0.034	0.278	0.027

Problem 363	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	C	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	15	33	16	32	30	0
N.S.	1	1.00	1.00	0.83	1.83	0.89	1.78	1.67	0.00
time (sec)	N/A	0.135	0.284	0.736	0.214	0.242	0.528	0.298	0.000

Problem 364	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	22	20	19	21	24	19	19
N.S.	1	1.00	0.81	0.74	0.70	0.78	0.89	0.70	0.70
time (sec)	N/A	0.156	0.074	0.152	0.217	0.254	0.088	0.305	0.220

Problem 365	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	22	18	17	16	15	0	16	18
N.S.	1	1.22	1.00	0.94	0.89	0.83	0.00	0.89	1.00
time (sec)	N/A	0.200	0.009	0.039	0.202	0.266	0.000	0.283	0.248

Problem 366	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	26	23	22	21	21	20	21	21
N.S.	1	0.96	0.85	0.81	0.78	0.78	0.74	0.78	0.78
time (sec)	N/A	0.172	0.001	0.030	0.198	0.260	0.096	0.276	0.002

Problem 367	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	32	26	21	20	20	20	20	31
N.S.	1	1.23	1.00	0.81	0.77	0.77	0.77	0.77	1.19
time (sec)	N/A	0.153	0.031	0.083	0.306	0.251	0.282	0.298	0.245

Problem 368	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	16	11	14	9	9	15	9	9
N.S.	1	1.07	0.73	0.93	0.60	0.60	1.00	0.60	0.60
time (sec)	N/A	0.183	0.033	1.704	0.205	0.250	0.779	0.275	0.287

Problem 369	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	52	50	35	34	37	121	29	30
N.S.	1	1.11	1.06	0.74	0.72	0.79	2.57	0.62	0.64
time (sec)	N/A	0.147	0.129	0.250	0.289	0.241	1.713	0.299	0.038

Problem 370	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	36	10	9	26	27	9	26
N.S.	1	1.00	3.27	0.91	0.82	2.36	2.45	0.82	2.36
time (sec)	N/A	0.124	0.003	0.177	0.200	0.238	0.014	0.303	0.024

Problem 371	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	13	13	12	13	14
N.S.	1	1.00	1.00	0.82	0.76	0.76	0.71	0.76	0.82
time (sec)	N/A	0.179	0.007	0.167	0.207	0.257	0.021	0.293	0.052

Problem 372	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	27	14	13	20	29	13	13
N.S.	1	1.00	1.59	0.82	0.76	1.18	1.71	0.76	0.76
time (sec)	N/A	0.181	0.012	1.822	0.198	0.274	0.020	0.265	0.199

Problem 373	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	18	15	19	17	36	19	14
N.S.	1	1.00	0.67	0.56	0.70	0.63	1.33	0.70	0.52
time (sec)	N/A	0.140	0.012	0.194	0.207	0.242	0.532	0.279	0.036

Problem 374	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	29	22	17	16	19	24	16	16
N.S.	1	1.21	0.92	0.71	0.67	0.79	1.00	0.67	0.67
time (sec)	N/A	0.191	0.002	0.274	0.210	0.266	0.017	0.271	0.033

Problem 375	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	17	20	18	12	16	16
N.S.	1	1.00	1.00	1.42	1.67	1.50	1.00	1.33	1.33
time (sec)	N/A	0.176	0.001	0.027	0.207	0.256	0.037	0.308	0.020

Problem 376	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	40	44	30	22	34	26	53	28	25
N.S.	1	1.10	0.75	0.55	0.85	0.65	1.32	0.70	0.62
time (sec)	N/A	0.146	0.021	0.177	0.292	0.251	0.280	0.282	0.024

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [300] had the largest ratio of [3]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	1	1	1.00	3	0.333
2	A	1	1	1.00	3	0.333
3	A	1	1	1.00	3	0.333
4	A	1	1	1.00	3	0.333
5	A	2	2	1.00	2	1.000
6	A	2	2	1.00	2	1.000
7	A	4	3	1.00	4	0.750
8	A	4	3	1.00	4	0.750
9	A	4	3	1.00	5	0.600
10	A	5	4	1.00	5	0.800
11	A	3	3	1.00	2	1.500
12	A	2	2	1.00	2	1.000
13	A	2	2	1.00	2	1.000
14	A	3	3	1.00	2	1.500
15	A	4	4	1.00	4	1.000
16	A	1	1	1.00	2	0.500
17	A	3	3	1.11	7	0.429
18	A	1	1	1.00	6	0.167
19	A	2	2	1.00	2	1.000
20	A	2	2	1.00	7	0.286
21	A	5	5	1.00	4	1.250
22	A	4	4	1.00	6	0.667

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
23	A	1	1	1.00	4	0.250
24	A	7	7	1.17	8	0.875
25	A	7	7	1.00	8	0.875
26	A	2	2	1.13	4	0.500
27	A	2	2	1.00	2	1.000
28	A	4	4	1.22	6	0.667
29	A	5	5	1.00	6	0.833
30	A	1	1	1.00	6	0.167
31	A	4	4	1.15	7	0.571
32	A	1	1	1.00	10	0.100
33	A	1	1	1.00	10	0.100
34	C	5	5	2.00	4	1.250
35	A	6	6	1.00	6	1.000
36	A	2	2	1.00	7	0.286
37	A	1	1	1.00	8	0.125
38	A	5	5	1.00	6	0.833
39	A	3	3	1.12	9	0.333
40	A	2	2	1.00	2	1.000
41	A	5	5	1.00	6	0.833
42	A	2	2	1.00	9	0.222
43	A	2	2	1.00	9	0.222
44	A	3	3	1.00	6	0.500
45	A	2	2	1.00	9	0.222
46	A	2	2	1.00	9	0.222
47	A	2	2	1.00	5	0.400
48	A	1	1	1.00	3	0.333
49	A	4	3	1.04	7	0.429
50	A	1	1	1.00	6	0.167
51	A	1	1	1.00	3	0.333
52	A	6	5	1.00	6	0.833
53	A	7	6	0.85	8	0.750
54	A	3	3	1.32	9	0.333
55	A	3	3	0.95	4	0.750
56	A	5	5	1.00	6	0.833

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
57	A	1	1	1.00	8	0.125
58	A	3	3	1.00	6	0.500
59	A	3	3	1.00	4	0.750
60	A	5	5	1.21	4	1.250
61	A	4	3	1.00	4	0.750
62	A	5	4	1.00	9	0.444
63	A	5	4	1.00	9	0.444
64	A	7	7	1.28	9	0.778
65	A	5	5	1.21	9	0.556
66	A	2	2	1.00	10	0.200
67	A	2	2	1.00	11	0.182
68	A	6	5	1.16	9	0.556
69	A	7	7	1.29	4	1.750
70	A	7	7	1.29	4	1.750
71	A	7	7	1.22	13	0.538
72	A	4	3	1.00	4	0.750
73	A	9	9	1.33	9	1.000
74	A	5	4	1.00	11	0.364
75	A	5	4	1.00	11	0.364
76	A	5	4	1.47	14	0.286
77	A	5	4	0.95	8	0.500
78	A	5	4	1.00	7	0.571
79	A	7	6	0.82	9	0.667
80	A	4	3	1.00	9	0.333
81	A	2	2	1.00	8	0.250
82	A	3	3	1.00	4	0.750
83	A	5	5	1.00	4	1.250
84	A	4	3	1.00	4	0.750
85	A	4	3	1.00	4	0.750
86	A	4	3	1.00	9	0.333
87	A	5	4	1.00	9	0.444
88	A	4	3	1.00	7	0.429
89	A	6	5	1.00	9	0.556
90	A	6	6	1.00	4	1.500

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
91	A	7	7	1.00	4	1.750
92	A	5	4	1.00	7	0.571
93	A	5	4	1.00	9	0.444
94	A	4	3	1.00	7	0.429
95	A	6	5	1.00	9	0.556
96	A	4	3	1.00	7	0.429
97	A	4	4	1.00	7	0.571
98	A	3	3	1.00	4	0.750
99	A	7	7	1.00	4	1.750
100	A	5	4	1.00	9	0.444
101	A	7	6	1.00	9	0.667
102	A	2	2	1.00	2	1.000
103	A	4	4	1.00	4	1.000
104	A	6	5	1.00	5	1.000
105	A	4	3	1.00	4	0.750
106	A	2	2	1.00	9	0.222
107	A	2	2	1.00	7	0.286
108	A	2	2	1.00	9	0.222
109	A	2	2	1.00	9	0.222
110	A	4	3	1.00	7	0.429
111	A	3	3	1.00	11	0.273
112	B	4	3	2.20	13	0.231
113	A	3	3	1.00	10	0.300
114	A	5	4	1.00	7	0.571
115	A	7	6	0.82	9	0.667
116	A	4	3	1.00	7	0.429
117	A	6	5	1.00	9	0.556
118	A	2	2	1.00	15	0.133
119	A	1	1	1.00	13	0.077
120	A	1	1	1.00	11	0.091
121	A	3	2	1.00	13	0.154
122	A	4	3	1.13	15	0.200
123	A	4	3	1.00	16	0.188
124	A	1	1	1.00	15	0.067

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
125	A	4	3	1.13	15	0.200
126	A	1	1	1.00	13	0.077
127	A	1	1	1.00	13	0.077
128	A	2	2	1.00	11	0.182
129	A	4	3	1.16	13	0.231
130	A	1	1	1.00	9	0.111
131	A	2	2	1.00	9	0.222
132	A	5	4	1.11	13	0.308
133	A	1	1	1.00	17	0.059
134	A	5	4	1.20	15	0.267
135	A	1	1	1.00	15	0.067
136	A	4	3	1.00	17	0.176
137	A	2	2	1.00	15	0.133
138	A	4	3	1.00	13	0.231
139	A	1	1	1.00	11	0.091
140	A	4	3	1.13	15	0.200
141	A	3	3	1.11	15	0.200
142	A	2	2	1.00	12	0.167
143	A	1	1	1.00	11	0.091
144	A	4	3	1.12	13	0.231
145	A	3	2	1.25	12	0.167
146	A	3	2	1.00	14	0.143
147	A	5	4	1.11	17	0.235
148	A	4	3	1.00	10	0.300
149	A	2	2	1.00	14	0.143
150	A	4	3	1.00	17	0.176
151	A	5	4	1.20	11	0.364
152	A	3	2	1.00	11	0.182
153	A	2	2	1.00	12	0.167
154	A	3	3	1.00	11	0.273
155	A	3	3	1.00	25	0.120
156	A	2	2	1.00	29	0.069
157	A	3	3	1.00	20	0.150
158	A	2	2	1.00	23	0.087

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
159	A	2	2	1.00	32	0.062
160	A	4	4	1.00	26	0.154
161	A	2	2	1.00	7	0.286
162	A	2	2	1.00	11	0.182
163	A	3	3	1.11	14	0.214
164	A	2	2	1.00	23	0.087
165	A	3	3	1.00	22	0.136
166	A	3	3	1.00	11	0.273
167	A	3	3	1.00	26	0.115
168	A	5	4	1.73	16	0.250
169	A	2	2	1.00	25	0.080
170	A	2	2	1.00	23	0.087
171	A	2	2	1.00	26	0.077
172	A	3	3	1.14	11	0.273
173	A	6	6	1.07	20	0.300
174	A	3	3	1.00	23	0.130
175	A	10	9	1.10	11	0.818
176	A	2	2	1.00	9	0.222
177	A	2	2	1.00	7	0.286
178	A	2	2	1.00	16	0.125
179	A	2	2	1.00	11	0.182
180	A	2	2	1.00	13	0.154
181	A	2	2	1.00	11	0.182
182	A	3	3	1.00	15	0.200
183	A	2	2	1.00	20	0.100
184	A	2	2	1.00	11	0.182
185	A	2	2	1.00	16	0.125
186	A	3	3	1.00	25	0.120
187	A	3	3	1.00	12	0.250
188	A	2	2	1.00	11	0.182
189	A	2	2	1.00	14	0.143
190	A	3	3	1.00	22	0.136
191	A	2	2	1.00	24	0.083
192	A	1	1	1.00	20	0.050

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
193	A	2	2	1.00	9	0.222
194	A	2	2	1.00	9	0.222
195	A	3	3	1.00	11	0.273
196	A	1	1	1.00	22	0.045
197	A	4	3	0.89	11	0.273
198	A	6	5	1.00	14	0.357
199	A	5	4	1.00	10	0.400
200	A	2	2	1.00	23	0.087
201	A	2	2	1.00	23	0.087
202	A	3	3	1.00	15	0.200
203	A	8	7	1.12	7	1.000
204	A	9	8	1.15	11	0.727
205	A	2	2	1.00	21	0.095
206	A	4	4	1.00	11	0.364
207	A	2	2	1.00	30	0.067
208	A	5	5	1.00	24	0.208
209	A	4	3	1.13	14	0.214
210	A	4	3	1.80	16	0.188
211	A	5	4	1.00	21	0.190
212	A	5	4	1.00	15	0.267
213	A	2	2	1.00	10	0.200
214	A	2	2	1.00	9	0.222
215	A	2	2	1.10	18	0.111
216	A	2	2	1.00	10	0.200
217	A	2	2	1.00	43	0.047
218	A	2	2	1.00	50	0.040
219	A	4	3	1.00	11	0.273
220	A	12	11	1.02	15	0.733
221	A	4	3	1.00	11	0.273
222	A	4	3	1.00	9	0.333
223	A	5	4	1.00	9	0.444
224	A	4	3	1.00	11	0.273
225	A	3	2	1.00	11	0.182
226	A	2	2	1.00	11	0.182

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
227	A	5	4	1.06	13	0.308
228	A	2	2	1.00	11	0.182
229	A	4	3	1.00	13	0.231
230	A	4	3	1.07	11	0.273
231	A	4	3	1.00	13	0.231
232	A	5	4	1.29	17	0.235
233	A	5	4	1.13	17	0.235
234	A	4	3	1.15	13	0.231
235	A	6	5	1.03	21	0.238
236	A	11	10	1.15	13	0.769
237	A	5	4	1.08	13	0.308
238	A	6	5	1.00	13	0.385
239	A	5	4	1.00	12	0.333
240	A	4	3	1.00	20	0.150
241	A	4	3	1.00	9	0.333
242	A	5	4	1.00	11	0.364
243	A	5	4	0.77	8	0.500
244	A	4	3	1.00	7	0.429
245	A	4	3	1.00	10	0.300
246	A	4	3	1.00	11	0.273
247	A	14	13	1.00	7	1.857
248	A	6	5	1.00	11	0.455
249	A	5	4	1.00	9	0.444
250	A	4	3	1.00	11	0.273
251	A	4	3	1.00	19	0.158
252	A	1	1	1.00	9	0.111
253	A	2	2	1.00	13	0.154
254	A	2	2	1.00	8	0.250
255	A	4	4	1.00	7	0.571
256	A	6	5	1.00	9	0.556
257	A	4	3	1.04	7	0.429
258	A	3	3	1.00	20	0.150
259	A	3	2	1.00	10	0.200
260	A	2	2	1.00	11	0.182

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
261	A	3	2	1.00	7	0.286
262	A	5	5	1.21	9	0.556
263	A	2	2	1.00	15	0.133
264	A	1	1	1.00	13	0.077
265	A	1	1	1.00	6	0.167
266	A	4	3	1.00	13	0.231
267	A	2	2	1.00	7	0.286
268	A	3	3	1.06	6	0.500
269	A	5	4	1.00	16	0.250
270	A	4	3	1.07	9	0.333
271	A	6	5	1.00	9	0.556
272	A	6	5	1.00	12	0.417
273	A	3	3	1.16	4	0.750
274	A	5	4	1.20	15	0.267
275	A	2	2	1.00	12	0.167
276	C	8	8	1.56	6	1.333
277	A	1	1	1.00	21	0.048
278	A	4	3	1.00	11	0.273
279	A	7	6	0.95	6	1.000
280	A	2	2	1.00	4	0.500
281	A	4	3	1.00	13	0.231
282	A	1	1	1.00	10	0.100
283	A	2	2	1.00	9	0.222
284	A	2	2	1.00	11	0.182
285	A	3	3	0.97	8	0.375
286	A	2	2	1.00	11	0.182
287	A	3	3	1.00	6	0.500
288	A	3	2	1.00	14	0.143
289	A	5	4	0.96	6	0.667
290	A	2	2	1.00	15	0.133
291	A	3	2	1.00	13	0.154
292	A	4	3	1.14	14	0.214
293	A	2	2	1.00	9	0.222
294	A	7	7	1.29	9	0.778

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
295	A	3	2	1.00	14	0.143
296	A	4	3	1.00	22	0.136
297	A	4	3	1.00	7	0.429
298	A	3	2	1.00	13	0.154
299	A	9	8	1.19	7	1.143
300	C	18	18	1.59	6	3.000
301	A	1	1	1.00	8	0.125
302	A	1	1	1.00	10	0.100
303	A	4	3	1.25	15	0.200
304	A	4	3	1.19	16	0.188
305	A	5	4	0.77	8	0.500
306	A	2	2	1.00	9	0.222
307	A	5	4	1.08	7	0.571
308	A	6	5	1.16	12	0.417
309	A	3	3	1.00	17	0.176
310	A	2	2	1.00	13	0.154
311	A	4	3	1.00	6	0.500
312	A	1	1	1.00	11	0.091
313	A	2	2	1.00	9	0.222
314	A	7	6	1.19	13	0.462
315	A	2	2	1.00	6	0.333
316	A	1	1	1.00	12	0.083
317	A	5	4	1.31	11	0.364
318	A	4	4	1.11	9	0.444
319	A	8	7	1.10	15	0.467
320	A	3	3	1.00	11	0.273
321	A	1	1	1.00	6	0.167
322	A	4	3	1.39	15	0.200
323	A	6	5	1.21	13	0.385
324	A	3	3	1.00	13	0.231
325	A	4	3	1.00	12	0.250
326	A	3	2	1.00	11	0.182
327	A	5	4	1.00	12	0.333
328	A	3	3	1.00	6	0.500

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
329	A	3	2	1.00	13	0.154
330	A	3	3	1.00	15	0.200
331	A	4	3	1.09	16	0.188
332	A	2	2	1.00	12	0.167
333	A	4	4	1.00	8	0.500
334	A	5	4	1.00	13	0.308
335	A	8	7	1.00	17	0.412
336	A	3	2	1.00	13	0.154
337	A	7	6	1.36	17	0.353
338	A	2	2	1.00	15	0.133
339	A	9	9	1.12	6	1.500
340	A	5	4	1.10	14	0.286
341	A	1	1	1.00	9	0.111
342	A	5	4	1.00	9	0.444
343	A	1	1	1.00	10	0.100
344	A	4	4	1.00	8	0.500
345	A	4	3	1.00	15	0.200
346	A	5	4	1.20	15	0.267
347	A	3	3	1.16	9	0.333
348	A	5	4	1.00	13	0.308
349	A	4	3	0.89	6	0.500
350	A	6	5	1.13	8	0.625
351	A	4	3	1.00	8	0.375
352	A	7	7	1.17	8	0.875
353	A	4	3	1.00	14	0.214
354	A	4	3	1.21	15	0.200
355	A	6	6	1.19	4	1.500
356	A	7	7	1.22	6	1.167
357	A	6	5	1.00	10	0.500
358	A	4	3	1.14	15	0.200
359	A	7	6	1.00	13	0.462
360	A	4	3	1.00	13	0.231
361	A	4	3	1.00	4	0.750
362	A	5	5	1.20	9	0.556

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
363	A	4	3	1.00	13	0.231
364	A	1	1	1.00	10	0.100
365	A	5	4	1.22	10	0.400
366	A	5	4	0.96	6	0.667
367	A	5	4	1.23	11	0.364
368	A	5	4	1.07	9	0.444
369	A	3	3	1.11	15	0.200
370	A	1	1	1.00	11	0.091
371	A	5	4	1.00	9	0.444
372	A	5	4	1.00	9	0.444
373	A	2	2	1.00	11	0.182
374	A	5	5	1.21	4	1.250
375	A	4	4	1.00	4	1.000
376	A	4	3	1.10	13	0.231

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^n dx$	145
3.2	$\int e^x dx$	149
3.3	$\int \frac{1}{x} dx$	153
3.4	$\int a^x dx$	157
3.5	$\int \sin(x) dx$	161
3.6	$\int \cos(x) dx$	165
3.7	$\int \sec^2(x) dx$	169
3.8	$\int \csc^2(x) dx$	173
3.9	$\int \sec(x) \tan(x) dx$	177
3.10	$\int \cot(x) \csc(x) dx$	181
3.11	$\int \sinh(x) dx$	186
3.12	$\int \cosh(x) dx$	190
3.13	$\int \tan(x) dx$	194
3.14	$\int \cot(x) dx$	199
3.15	$\int x \sin(x) dx$	204
3.16	$\int \log(x) dx$	209
3.17	$\int e^x x^2 dx$	213
3.18	$\int e^x \sin(x) dx$	217
3.19	$\int \arctan(x) dx$	221
3.20	$\int e^{2x} x dx$	226
3.21	$\int x \cos(x) dx$	231
3.22	$\int x \sin(4x) dx$	236
3.23	$\int x \log(x) dx$	241
3.24	$\int x^2 \cos(3x) dx$	245
3.25	$\int x^2 \sin(2x) dx$	250
3.26	$\int \log^2(x) dx$	255
3.27	$\int \arcsin(x) dx$	259
3.28	$\int t \cos(t) \sin(t) dt$	263

3.29	$\int t \sec^2(t) dt$	268
3.30	$\int t^2 \log(t) dt$	273
3.31	$\int e^{t^3} dt$	277
3.32	$\int e^{2t} \sin(3t) dt$	282
3.33	$\int e^{-t} \cos(3t) dt$	286
3.34	$\int y \sinh(y) dy$	290
3.35	$\int y \cosh(ay) dy$	295
3.36	$\int e^{-t} t dt$	300
3.37	$\int \sqrt{t} \log(t) dt$	304
3.38	$\int x \cos(2x) dx$	308
3.39	$\int e^{-x} x^2 dx$	313
3.40	$\int \arccos(x) dx$	317
3.41	$\int x \csc^2(x) dx$	321
3.42	$\int \cos(5x) \sin(3x) dx$	326
3.43	$\int \sin(2x) \sin(4x) dx$	330
3.44	$\int \cos(x) \log(\sin(x)) dx$	334
3.45	$\int e^{x^2} x^3 dx$	338
3.46	$\int e^x (3 + 2x) dx$	343
3.47	$\int 5^x dx$	347
3.48	$\int \cos(\log(x)) dx$	351
3.49	$\int e^{\sqrt{x}} dx$	355
3.50	$\int \log(\sqrt{x}) dx$	360
3.51	$\int \sin(\log(x)) dx$	364
3.52	$\int \sin(\sqrt{x}) dx$	368
3.53	$\int x^5 \cos(x^3) dx$	373
3.54	$\int e^{x^2} x^5 dx$	378
3.55	$\int x \arctan(x) dx$	383
3.56	$\int x \cos(\pi x) dx$	388
3.57	$\int \sqrt{x} \log(x) dx$	393
3.58	$\int \sin^2(3x) dx$	397
3.59	$\int \cos^2(x) dx$	402
3.60	$\int \cos^4(x) dx$	406
3.61	$\int \sin^3(x) dx$	411
3.62	$\int \cos^4(x) \sin^3(x) dx$	415
3.63	$\int \cos^3(x) \sin^4(x) dx$	420
3.64	$\int \cos^2(x) \sin^4(x) dx$	425
3.65	$\int \cos^2(x) \sin^2(x) dx$	430
3.66	$\int (1 - \sin(2x))^2 dx$	435
3.67	$\int \cos(x) \sin\left(\frac{\pi}{6} + x\right) dx$	439
3.68	$\int \cos^5(x) \sin^5(x) dx$	443

3.69	$\int \sin^6(x) dx$	448
3.70	$\int \cos^6(x) dx$	453
3.71	$\int \cos^4(2x) \sin^2(2x) dx$	458
3.72	$\int \sin^5(x) dx$	463
3.73	$\int \cos^4(x) \sin^4(x) dx$	467
3.74	$\int \sqrt{\cos(x)} \sin^3(x) dx$	472
3.75	$\int \cos^3(x) \sqrt{\sin(x)} dx$	477
3.76	$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx$	482
3.77	$\int x \sin^3(x^2) dx$	487
3.78	$\int \sin^2(x) \tan(x) dx$	492
3.79	$\int \cos^2(x) \cot^3(x) dx$	497
3.80	$\int \sec(x)(1 - \sin(x)) dx$	502
3.81	$\int \frac{1}{1 - \sin(x)} dx$	507
3.82	$\int \tan^2(x) dx$	511
3.83	$\int \tan^4(x) dx$	515
3.84	$\int \sec^4(x) dx$	520
3.85	$\int \sec^6(x) dx$	524
3.86	$\int \sec^2(x) \tan^4(x) dx$	528
3.87	$\int \sec^4(x) \tan^2(x) dx$	532
3.88	$\int \sec^3(x) \tan(x) dx$	537
3.89	$\int \sec^3(x) \tan^3(x) dx$	541
3.90	$\int \tan^5(x) dx$	546
3.91	$\int \tan^6(x) dx$	551
3.92	$\int \sec(x) \tan^5(x) dx$	556
3.93	$\int \sec^3(x) \tan^5(x) dx$	561
3.94	$\int \sec^6(x) \tan(x) dx$	566
3.95	$\int \sec^6(x) \tan^3(x) dx$	570
3.96	$\int \sec^2(x) \tan(x) dx$	575
3.97	$\int \sec(x) \tan^2(x) dx$	579
3.98	$\int \cot^2(x) dx$	584
3.99	$\int \cot^3(x) dx$	588
3.100	$\int \cot^4(x) \csc^4(x) dx$	593
3.101	$\int \cot^3(x) \csc^4(x) dx$	598
3.102	$\int \csc(x) dx$	603
3.103	$\int \csc^3(x) dx$	607
3.104	$\int \cos(x) \cot(x) dx$	612
3.105	$\int \csc^4(x) dx$	617
3.106	$\int \sin(2x) \sin(5x) dx$	621
3.107	$\int \cos(x) \sin(3x) dx$	625
3.108	$\int \cos(3x) \cos(4x) dx$	629

3.109	$\int \sin(3x) \sin(6x) dx$	633
3.110	$\int \cos^5(x) \sin(x) dx$	637
3.111	$\int \cos(x) \cos(2x) \cos(3x) dx$	641
3.112	$\int \cos^2(x) (1 - \tan^2(x)) dx$	645
3.113	$\int \csc(2x)(\cos(x) + \sin(x)) dx$	650
3.114	$\int \sin^2(x) \tan(x) dx$	655
3.115	$\int \cos^2(x) \cot^3(x) dx$	660
3.116	$\int \sec^3(x) \tan(x) dx$	665
3.117	$\int \sec^3(x) \tan^3(x) dx$	669
3.118	$\int \frac{\sqrt{9-x^2}}{x^2} dx$	674
3.119	$\int \frac{1}{x^2\sqrt{4+x^2}} dx$	679
3.120	$\int \frac{x}{\sqrt{4+x^2}} dx$	683
3.121	$\int \frac{1}{\sqrt{-a^2+x^2}} dx$	687
3.122	$\int \frac{x^3}{(9+4x^2)^{3/2}} dx$	691
3.123	$\int \frac{x}{\sqrt{3-2x-x^2}} dx$	696
3.124	$\int \frac{1}{x^2\sqrt{1-x^2}} dx$	701
3.125	$\int x^3\sqrt{4-x^2} dx$	705
3.126	$\int \frac{x}{\sqrt{1-x^2}} dx$	710
3.127	$\int x\sqrt{4-x^2} dx$	714
3.128	$\int \sqrt{1-4x^2} dx$	719
3.129	$\int \frac{x^3}{\sqrt{4+x^2}} dx$	724
3.130	$\int \frac{1}{\sqrt{9+x^2}} dx$	729
3.131	$\int \sqrt{1+x^2} dx$	733
3.132	$\int \frac{1}{x^3\sqrt{-16+x^2}} dx$	737
3.133	$\int \frac{\sqrt{-a^2+x^2}}{x^4} dx$	743
3.134	$\int \frac{\sqrt{-4+9x^2}}{x} dx$	747
3.135	$\int \frac{1}{x^2\sqrt{-9+16x^2}} dx$	752
3.136	$\int \frac{x^2}{(a^2-x^2)^{3/2}} dx$	757
3.137	$\int \frac{x^2}{\sqrt{5-x^2}} dx$	762
3.138	$\int \frac{1}{x\sqrt{3+x^2}} dx$	767
3.139	$\int \frac{x}{(4+x^2)^{5/2}} dx$	772
3.140	$\int x^3\sqrt{4-9x^2} dx$	777
3.141	$\int x^2\sqrt{9-x^2} dx$	782
3.142	$\int 5x\sqrt{1+x^2} dx$	787
3.143	$\int \frac{1}{(-25+4x^2)^{3/2}} dx$	792
3.144	$\int \sqrt{2x-x^2} dx$	796
3.145	$\int \frac{1}{\sqrt{8+4x+x^2}} dx$	801
3.146	$\int \frac{1}{\sqrt{-8+6x+9x^2}} dx$	805

3.147	$\int \frac{x^2}{\sqrt{4x-x^2}} dx$	809
3.148	$\int \frac{1}{(2+2x+x^2)^2} dx$	814
3.149	$\int \frac{1}{(5-4x-x^2)^{5/2}} dx$	819
3.150	$\int e^t \sqrt{9-e^{2t}} dt$	823
3.151	$\int \sqrt{-9+e^{2t}} dt$	828
3.152	$\int \frac{1}{\sqrt{a^2+x^2}} dx$	833
3.153	$\int \frac{5+x}{-2+x+x^2} dx$	837
3.154	$\int \frac{x+x^3}{-1+x} dx$	841
3.155	$\int \frac{-1+2x+x^2}{-2x+3x^2+2x^3} dx$	845
3.156	$\int \frac{1+4x-2x^2+x^4}{1-x-x^2+x^3} dx$	849
3.157	$\int \frac{4-x+2x^2}{4x+x^3} dx$	853
3.158	$\int \frac{2-3x+4x^2}{3-4x+4x^2} dx$	857
3.159	$\int \frac{1+x^2+x^3}{(-1+x)x(1+x^2)^3(1+x+x^2)} dx$	861
3.160	$\int \frac{1-3x+2x^2-x^3}{x(1+x^2)^2} dx$	867
3.161	$\int \frac{1}{(1+x^2)^2} dx$	872
3.162	$\int \frac{1}{(-1+x)(2+x)} dx$	876
3.163	$\int \frac{7}{-12+5x+2x^2} dx$	880
3.164	$\int \frac{-4+3x+x^2}{(-1+2x)^2(3+2x)} dx$	884
3.165	$\int \frac{-x^2+x^3}{(-6+x)(3+5x)^3} dx$	888
3.166	$\int \frac{1}{-x^3+x^4} dx$	893
3.167	$\int \frac{1-x-x^2+x^3+x^4}{-x+x^3} dx$	897
3.168	$\int \frac{-2+x^2}{x(2+x^2)} dx$	901
3.169	$\int \frac{2-4x^2+x^3}{(1+x^2)(2+x^2)} dx$	906
3.170	$\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx$	910
3.171	$\int \frac{1+16x}{(5+x)^2(-3+2x)(1+x+x^2)} dx$	914
3.172	$\int \frac{x^4}{(9+x^2)^3} dx$	919
3.173	$\int \frac{19x}{(-1+x)^3(3+5x+4x^2)^2} dx$	923
3.174	$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx$	929
3.175	$\int \frac{1}{-x^3+x^6} dx$	934
3.176	$\int \frac{x^2}{1+x} dx$	940
3.177	$\int \frac{x}{-5+x} dx$	944
3.178	$\int \frac{-1+4x}{(-1+x)(2+x)} dx$	948
3.179	$\int \frac{1}{(1+x)(2+x)} dx$	952
3.180	$\int \frac{-5+6x}{3+2x} dx$	956
3.181	$\int \frac{1}{(a+x)(b+x)} dx$	960
3.182	$\int \frac{1+x^2}{-x+x^2} dx$	965

3.183	$\int \frac{1-12x+x^2+x^3}{-12+x+x^2} dx$	969
3.184	$\int \frac{3+2x}{(1+x)^2} dx$	973
3.185	$\int \frac{1}{x(1+x)(3+2x)} dx$	977
3.186	$\int \frac{-3+5x+6x^2}{-3x+2x^2+x^3} dx$	981
3.187	$\int \frac{x}{4+4x+x^2} dx$	985
3.188	$\int \frac{1}{(-1+x)^2(4+x)} dx$	989
3.189	$\int \frac{x^2}{(-3+x)(2+x)^2} dx$	993
3.190	$\int \frac{-2+3x+5x^2}{2x^2+x^3} dx$	997
3.191	$\int \frac{18-2x-4x^2}{-6+x+4x^2+x^3} dx$	1002
3.192	$\int \frac{2x+x^2}{4+3x^2+x^3} dx$	1006
3.193	$\int \frac{1}{(-1+x)^2x^2} dx$	1010
3.194	$\int \frac{x^2}{(1+x)^3} dx$	1014
3.195	$\int \frac{1}{-x^2+x^4} dx$	1018
3.196	$\int \frac{-x+2x^3}{1-x^2+x^4} dx$	1023
3.197	$\int \frac{x^3}{1+x^2} dx$	1027
3.198	$\int \frac{-1+x}{2+2x+x^2} dx$	1031
3.199	$\int \frac{x}{1+x+x^2} dx$	1036
3.200	$\int \frac{7+5x+4x^2}{5+4x+4x^2} dx$	1041
3.201	$\int \frac{5-4x+3x^2}{(-1+x)(1+x^2)} dx$	1045
3.202	$\int \frac{3+2x}{3x+x^3} dx$	1049
3.203	$\int \frac{1}{-1+x^3} dx$	1053
3.204	$\int \frac{x^3}{1+x^3} dx$	1059
3.205	$\int \frac{-1-2x+x^2}{(-1+x)^2(1+x^2)} dx$	1065
3.206	$\int \frac{x^4}{-1+x^4} dx$	1069
3.207	$\int \frac{-4+6x-x^2+3x^3}{(1+x^2)(2+x^2)} dx$	1074
3.208	$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx$	1078
3.209	$\int \frac{-3+x}{(4+2x+x^2)^2} dx$	1083
3.210	$\int \frac{1+x^4}{x(1+x^2)^2} dx$	1088
3.211	$\int \frac{\cos(x)(-3+2\sin(x))}{2-3\sin(x)+\sin^2(x)} dx$	1092
3.212	$\int \frac{\cos^2(x)\sin(x)}{5+\cos^2(x)} dx$	1097
3.213	$\int \frac{1}{-3+2x+x^2} dx$	1102
3.214	$\int \frac{1}{-2x+x^2} dx$	1106
3.215	$\int \frac{1+2x}{-7+12x+4x^2} dx$	1110
3.216	$\int \frac{x}{-1+x+x^2} dx$	1114
3.217	$\int \frac{-32+5x-27x^2+4x^3}{-70-299x-286x^2+50x^3-13x^4+30x^5} dx$	1118
3.218	$\int \frac{8-13x^2-7x^3+12x^5}{4-20x+41x^2-80x^3+116x^4-80x^5+100x^6} dx$	1123
3.219	$\int \frac{\sqrt{4+x}}{x} dx$	1129

3.220	$\int \frac{1}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx$	1134
3.221	$\int \frac{1}{-4 \cos(x) + 3 \sin(x)} dx$	1144
3.222	$\int \frac{1}{1 + \sqrt{x}} dx$	1149
3.223	$\int \frac{1}{1 + \frac{1}{\sqrt[3]{x}}} dx$	1153
3.224	$\int \frac{\sqrt{x}}{1+x} dx$	1158
3.225	$\int \frac{1}{x\sqrt{1+x}} dx$	1163
3.226	$\int \frac{1}{-\sqrt[3]{x+x}} dx$	1168
3.227	$\int \frac{1}{x - \sqrt{2+x}} dx$	1172
3.228	$\int \frac{x^2}{\sqrt{-1+x}} dx$	1177
3.229	$\int \frac{\sqrt{-1+x}}{1+x} dx$	1182
3.230	$\int \frac{1}{\sqrt{1+\sqrt{x}}} dx$	1187
3.231	$\int \frac{\sqrt{x}}{x+x^2} dx$	1192
3.232	$\int \frac{1+\sqrt{x}}{-1+\sqrt{x}} dx$	1197
3.233	$\int \frac{1+\frac{1}{\sqrt[3]{x}}}{-1+\frac{1}{\sqrt[3]{x}}} dx$	1202
3.234	$\int \frac{x^3}{\sqrt[3]{1+x^2}} dx$	1207
3.235	$\int \frac{\sqrt{x}}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx$	1212
3.236	$\int \frac{1}{\frac{1}{\sqrt[4]{x}} + \sqrt{x}} dx$	1221
3.237	$\int \frac{1}{\frac{1}{\sqrt[3]{x}} + \frac{1}{\sqrt[4]{x}}} dx$	1228
3.238	$\int \sqrt{\frac{1-x}{x}} dx$	1234
3.239	$\int \frac{\cos(x)}{\sin(x) + \sin^2(x)} dx$	1239
3.240	$\int \frac{e^{2x}}{2+3e^x+e^{2x}} dx$	1244
3.241	$\int \frac{1}{\sqrt{1+e^x}} dx$	1249
3.242	$\int \sqrt{1-e^x} dx$	1254
3.243	$\int \frac{1}{3-5 \sin(x)} dx$	1259
3.244	$\int \frac{1}{\cos(x) + \sin(x)} dx$	1264
3.245	$\int \frac{1}{1 - \cos(x) + \sin(x)} dx$	1269
3.246	$\int \frac{1}{4 \cos(x) + 3 \sin(x)} dx$	1273
3.247	$\int \frac{1}{\sin(x) + \tan(x)} dx$	1278
3.248	$\int \frac{1}{2 \sin(x) + \sin(2x)} dx$	1284
3.249	$\int \frac{\sec(x)}{1 + \sin(x)} dx$	1289
3.250	$\int \frac{1}{b \cos(x) + a \sin(x)} dx$	1294
3.251	$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx$	1299

3.252	$\int \frac{x}{-1+x^2} dx$	1304
3.253	$\int (1 + \sqrt{x}) \sqrt{x} dx$	1308
3.254	$\int \frac{1}{1-\cos(x)} dx$	1312
3.255	$\int \sec(x) \tan^2(x) dx$	1316
3.256	$\int \sec^3(x) \tan^3(x) dx$	1321
3.257	$\int e^{\sqrt{x}} dx$	1326
3.258	$\int \frac{1+x^5}{-10x-3x^2+x^3} dx$	1331
3.259	$\int \frac{1}{x\sqrt{\log(x)}} dx$	1336
3.260	$\int \frac{5+2x}{-3+x} dx$	1340
3.261	$\int e^{e^x+x} dx$	1344
3.262	$\int \cos^2(x) \sin^2(x) dx$	1348
3.263	$\int \frac{-\cos(x)+\sin(x)}{\cos(x)+\sin(x)} dx$	1353
3.264	$\int \frac{x}{\sqrt{1-x^2}} dx$	1357
3.265	$\int x^3 \log(x) dx$	1361
3.266	$\int \frac{\sqrt{-2+x}}{2+x} dx$	1365
3.267	$\int \frac{x}{(2+x)^2} dx$	1370
3.268	$\int \log(1+x^2) dx$	1374
3.269	$\int \frac{\sqrt{1+\log(x)}}{x \log(x)} dx$	1379
3.270	$\int (1 + \sqrt{x})^8 dx$	1384
3.271	$\int \sec^4(x) \tan^3(x) dx$	1389
3.272	$\int \frac{x}{2-2x+x^2} dx$	1394
3.273	$\int x \arcsin(x) dx$	1399
3.274	$\int \frac{\sqrt{9-x^2}}{x} dx$	1404
3.275	$\int \frac{x}{2+3x+x^2} dx$	1409
3.276	$\int x^2 \cosh(x) dx$	1413
3.277	$\int \frac{1+x+x^3}{4x+2x^2+x^4} dx$	1418
3.278	$\int \frac{\cos(x)}{1+\sin^2(x)} dx$	1422
3.279	$\int \cos(\sqrt{x}) dx$	1427
3.280	$\int \sin(\pi x) dx$	1432
3.281	$\int \frac{e^{2x}}{1+e^x} dx$	1436
3.282	$\int e^{3x} \cos(5x) dx$	1441
3.283	$\int \cos(3x) \cos(5x) dx$	1445
3.284	$\int \frac{1}{1+x+x^2+x^3} dx$	1449
3.285	$\int x^2 \log(1+x) dx$	1453
3.286	$\int e^{-x^3} x^5 dx$	1458
3.287	$\int \tan^2(4x) dx$	1463
3.288	$\int \frac{1}{\sqrt{-5+12x+9x^2}} dx$	1467
3.289	$\int x^2 \arctan(x) dx$	1471
3.290	$\int \frac{1-\sqrt{x}}{\sqrt[3]{x}} dx$	1476

3.291	$\int \frac{1}{-e^{-x}+e^x} dx$	1480
3.292	$\int \frac{x}{10+2x^2+x^4} dx$	1485
3.293	$\int \frac{1}{\frac{1}{\sqrt[3]{x}}+x} dx$	1489
3.294	$\int \cos^4(x) \sin^2(x) dx$	1494
3.295	$\int \frac{1}{\sqrt{5-4x-x^2}} dx$	1499
3.296	$\int \frac{x}{1-x^2+\sqrt{1-x^2}} dx$	1503
3.297	$\int (1 + \cos(x)) \csc(x) dx$	1508
3.298	$\int \frac{e^x}{-1+e^{2x}} dx$	1513
3.299	$\int \frac{1}{-8+x^3} dx$	1517
3.300	$\int x^5 \cosh(x) dx$	1523
3.301	$\int \csc(x) \log(\tan(x)) \sec(x) dx$	1529
3.302	$\int (-2x + x^2 + x^3) dx$	1533
3.303	$\int \frac{1+e^x}{1-e^x} dx$	1537
3.304	$\int \frac{x}{(1+x^2)(4+x^2)} dx$	1542
3.305	$\int \frac{1}{4-5\sin(x)} dx$	1547
3.306	$\int x \sqrt[3]{c+x} dx$	1552
3.307	$\int e^{\sqrt[3]{x}} dx$	1557
3.308	$\int \frac{1}{4+x+\sqrt{1+x}} dx$	1562
3.309	$\int \frac{1+x^3}{-x^2+x^3} dx$	1567
3.310	$\int (-3 + 4x + x^2) \sin(2x) dx$	1571
3.311	$\int \cos(\cos(x)) \sin(x) dx$	1576
3.312	$\int \frac{1}{\sqrt{16-x^2}} dx$	1581
3.313	$\int \frac{x^3}{(1+x)^{10}} dx$	1585
3.314	$\int \cot^3(2x) \csc^3(2x) dx$	1590
3.315	$\int (x + \sin(x))^2 dx$	1595
3.316	$\int \frac{e^{\arctan(x)}}{1+x^2} dx$	1599
3.317	$\int \frac{1}{x(1+x^4)} dx$	1603
3.318	$\int e^{-2t} t^3 dt$	1608
3.319	$\int \frac{\sqrt{t}}{1+\sqrt[3]{t}} dt$	1613
3.320	$\int \sin(x) \sin(2x) \sin(3x) dx$	1618
3.321	$\int \log\left(\frac{x}{2}\right) dx$	1622
3.322	$\int \sqrt{\frac{1+x}{1-x}} dx$	1626
3.323	$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx$	1631
3.324	$\int \frac{a+x}{a^2+x^2} dx$	1636
3.325	$\int \sqrt{1+x-x^2} dx$	1640
3.326	$\int \frac{x^4}{16+x^{10}} dx$	1645
3.327	$\int \frac{2+x}{2+x+x^2} dx$	1649
3.328	$\int x \sec(x) \tan(x) dx$	1654

3.329	$\int \frac{x}{-a^4+x^4} dx$	1659
3.330	$\int \frac{1}{\sqrt{x+\sqrt{1+x}}} dx$	1663
3.331	$\int \frac{1}{1-e^{-x}+2e^x} dx$	1668
3.332	$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}} dx$	1673
3.333	$\int \frac{\log(1+x)}{x^2} dx$	1677
3.334	$\int \frac{1}{-e^x+e^{3x}} dx$	1682
3.335	$\int \frac{1+\cos^2(x)}{1-\cos^2(x)} dx$	1687
3.336	$\int \frac{1}{x\sqrt{-25+2x}} dx$	1692
3.337	$\int \frac{\sin(2x)}{\sqrt{9-\cos^4(x)}} dx$	1697
3.338	$\int \frac{x^2}{\sqrt{5-4x^2}} dx$	1702
3.339	$\int x^3 \sin(x) dx$	1707
3.340	$\int x\sqrt{4+2x+x^2} dx$	1712
3.341	$\int x(5+x^2)^8 dx$	1717
3.342	$\int \cos^2(x) \sin^5(x) dx$	1721
3.343	$\int e^{-3x} \cos(4x) dx$	1726
3.344	$\int \csc^3\left(\frac{x}{2}\right) dx$	1730
3.345	$\int \frac{\sqrt{-1+9x^2}}{x^2} dx$	1735
3.346	$\int \frac{\sqrt{4-3x^2}}{x} dx$	1740
3.347	$\int e^{3x} x^2 dx$	1745
3.348	$\int \frac{\cos(x) \sin(x)}{\sqrt{1+\sin(x)}} dx$	1750
3.349	$\int x \arcsin(x^2) dx$	1755
3.350	$\int x^3 \arcsin(x^2) dx$	1759
3.351	$\int e^x \operatorname{sech}(e^x) dx$	1764
3.352	$\int x^2 \cos(3x) dx$	1769
3.353	$\int \sqrt{5-4x-x^2} dx$	1774
3.354	$\int \frac{x^5}{\sqrt{2+x^2}} dx$	1779
3.355	$\int \sec^5(x) dx$	1783
3.356	$\int \sin^6(2x) dx$	1788
3.357	$\int \cos(x) \log(\sin(x)) \sin^2(x) dx$	1793
3.358	$\int \frac{e^{-x}}{1+2e^x} dx$	1798
3.359	$\int \sqrt{2+3\cos(x)} \tan(x) dx$	1803
3.360	$\int \frac{x}{\sqrt{-4x+x^2}} dx$	1809
3.361	$\int \cos^5(x) dx$	1814
3.362	$\int e^{-x} x^4 dx$	1818
3.363	$\int \frac{x^4}{\sqrt{-2+x^{10}}} dx$	1823
3.364	$\int e^x \cos(4+3x) dx$	1828
3.365	$\int e^x \log(1+e^x) dx$	1832
3.366	$\int x^2 \arctan(x) dx$	1837
3.367	$\int \sqrt{-1+e^{2x}} dx$	1842

3.368	$\int e^{\sin(x)} \sin(2x) dx$	1847
3.369	$\int x^2 \sqrt{5-x^2} dx$	1852
3.370	$\int x^2(1+x^3)^4 dx$	1857
3.371	$\int \cos^3(x) \sin^3(x) dx$	1861
3.372	$\int \sec^4(x) \tan^2(x) dx$	1866
3.373	$\int x\sqrt{1+2x} dx$	1871
3.374	$\int \sin^4(x) dx$	1876
3.375	$\int \tan^3(x) dx$	1881
3.376	$\int x^5 \sqrt{1+x^2} dx$	1886

3.1 $\int x^n dx$

3.1.1	Optimal result	145
3.1.2	Mathematica [A] (verified)	145
3.1.3	Rubi [A] (verified)	146
3.1.4	Maple [A] (verified)	146
3.1.5	Fricas [A] (verification not implemented)	147
3.1.6	Sympy [A] (verification not implemented)	147
3.1.7	Maxima [A] (verification not implemented)	147
3.1.8	Giac [A] (verification not implemented)	148
3.1.9	Mupad [B] (verification not implemented)	148

3.1.1 Optimal result

Integrand size = 3, antiderivative size = 11

$$\int x^n dx = \frac{x^{1+n}}{1+n}$$

output `x^(1+n)/(1+n)`

3.1.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int x^n dx = \frac{x^{1+n}}{1+n}$$

input `Integrate[x^n,x]`

output `x^(1 + n)/(1 + n)`

3.1.3 Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^n dx$$

$$\downarrow 15$$

$$\frac{x^{n+1}}{n+1}$$

input `Int[x^n,x]`

output `x^(1+n)/(1+n)`

3.1.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m+1)/(m+1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

3.1.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

method	result	size
risch	$\frac{x x^n}{1+n}$	11
parallelrisk	$\frac{x x^n}{1+n}$	11
gosp	$\frac{x^{1+n}}{1+n}$	12
default	$\frac{x^{1+n}}{1+n}$	12
norman	$\frac{x e^{n \ln(x)}}{1+n}$	13

input `int(x^n,x,method=_RETURNVERBOSE)`

output `x/(1+n)*x^n`

3.1.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int x^n dx = \frac{xx^n}{n+1}$$

input `integrate(x^n,x, algorithm="fricas")`

output `x*x^n/(n + 1)`

3.1.6 Sympy [A] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

$$\int x^n dx = \begin{cases} \frac{x^{n+1}}{n+1} & \text{for } n \neq -1 \\ \log(x) & \text{otherwise} \end{cases}$$

input `integrate(x**n,x)`

output `Piecewise((x**(n + 1)/(n + 1), Ne(n, -1)), (log(x), True))`

3.1.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int x^n dx = \frac{x^{n+1}}{n+1}$$

input `integrate(x^n,x, algorithm="maxima")`

output `x^(n + 1)/(n + 1)`

3.1.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int x^n dx = \frac{x^{n+1}}{n+1}$$

input `integrate(x^n,x, algorithm="giac")`

output `x^(n + 1)/(n + 1)`

3.1.9 Mupad [B] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.82

$$\int x^n dx = \begin{cases} \ln(x) & \text{if } n = -1 \\ \frac{x^{n+1}}{n+1} & \text{if } n \neq -1 \end{cases}$$

input `int(x^n,x)`

output `piecewise(n == -1, log(x), n ~= -1, x^(n + 1)/(n + 1))`

3.2 $\int e^x dx$

3.2.1	Optimal result	149
3.2.2	Mathematica [A] (verified)	149
3.2.3	Rubi [A] (verified)	150
3.2.4	Maple [A] (verified)	150
3.2.5	Fricas [A] (verification not implemented)	151
3.2.6	Sympy [A] (verification not implemented)	151
3.2.7	Maxima [A] (verification not implemented)	151
3.2.8	Giac [A] (verification not implemented)	152
3.2.9	Mupad [B] (verification not implemented)	152

3.2.1 Optimal result

Integrand size = 3, antiderivative size = 3

$$\int e^x dx = e^x$$

output `exp(x)`

3.2.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int e^x dx = e^x$$

input `Integrate[E^x,x]`

output `E^x`

3.2.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x dx$$

↓ 2624

$$e^x$$

input `Int[E^x,x]`

output `E^x`

3.2.3.1 Defintions of rubi rules used

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;`
`FreeQ[{F, n}, x] && LinearQ[v, x]`

3.2.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

method	result	size
gospers	e^x	3
lookup	e^x	3
derivativedivides	e^x	3
default	e^x	3
norman	e^x	3
risch	e^x	3
parallelrisc	e^x	3
meijerg	$-1 + e^x$	5

input `int(exp(x),x,method=_RETURNVERBOSE)`

output `exp(x)`

3.2.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int e^x dx = e^x$$

input `integrate(exp(x),x, algorithm="fricas")`

output `e^x`

3.2.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int e^x dx = e^x$$

input `integrate(exp(x),x)`

output `exp(x)`

3.2.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int e^x dx = e^x$$

input `integrate(exp(x),x, algorithm="maxima")`

output `e^x`

3.2.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int e^x dx = e^x$$

input `integrate(exp(x),x, algorithm="giac")`

output `e^x`

3.2.9 Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int e^x dx = e^x$$

input `int(exp(x),x)`

output `exp(x)`

3.3 $\int \frac{1}{x} dx$

3.3.1	Optimal result	153
3.3.2	Mathematica [A] (verified)	153
3.3.3	Rubi [A] (verified)	154
3.3.4	Maple [A] (verified)	154
3.3.5	Fricas [A] (verification not implemented)	155
3.3.6	Sympy [A] (verification not implemented)	155
3.3.7	Maxima [A] (verification not implemented)	155
3.3.8	Giac [A] (verification not implemented)	156
3.3.9	Mupad [B] (verification not implemented)	156

3.3.1 Optimal result

Integrand size = 3, antiderivative size = 2

$$\int \frac{1}{x} dx = \log(x)$$

output

`ln(x)`

3.3.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \log(x)$$

input

`Integrate[x^(-1), x]`

output

`Log[x]`

3.3.3 Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x} dx$$

↓ 14

$$\log(x)$$

input `Int [x(-1), x]`

output `Log [x]`

3.3.3.1 Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] :> Simp[a*Log[x], x] /; FreeQ[a, x]`

3.3.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\ln(x)$	3
norman	$\ln(x)$	3
risch	$\ln(x)$	3
parallelrisch	$\ln(x)$	3

input `int(1/x,x,method=_RETURNVERBOSE)`

output `ln(x)`

3.3.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \log(x)$$

input `integrate(1/x,x, algorithm="fricas")`

output `log(x)`

3.3.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \log(x)$$

input `integrate(1/x,x)`

output `log(x)`

3.3.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \log(x)$$

input `integrate(1/x,x, algorithm="maxima")`

output `log(x)`

3.3.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

$$\int \frac{1}{x} dx = \log(|x|)$$

input `integrate(1/x,x, algorithm="giac")`

output `log(abs(x))`

3.3.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \ln(x)$$

input `int(1/x,x)`

output `log(x)`

3.4 $\int a^x dx$

3.4.1	Optimal result	157
3.4.2	Mathematica [A] (verified)	157
3.4.3	Rubi [A] (verified)	158
3.4.4	Maple [A] (verified)	158
3.4.5	Fricas [A] (verification not implemented)	159
3.4.6	Sympy [A] (verification not implemented)	159
3.4.7	Maxima [A] (verification not implemented)	159
3.4.8	Giac [A] (verification not implemented)	160
3.4.9	Mupad [B] (verification not implemented)	160

3.4.1 Optimal result

Integrand size = 3, antiderivative size = 8

$$\int a^x dx = \frac{a^x}{\log(a)}$$

output `ax/ln(a)`

3.4.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int a^x dx = \frac{a^x}{\log(a)}$$

input `Integrate[ax, x]`

output `ax/Log[a]`

3.4.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int a^x dx$$

$$\downarrow \text{2624}$$

$$\frac{a^x}{\log(a)}$$

input `Int[a^x,x]`

output `a^x/Log[a]`

3.4.3.1 Defintions of rubi rules used

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;`
`FreeQ[{F, n}, x] && LinearQ[v, x]`

3.4.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
gospers	$\frac{a^x}{\ln(a)}$	9
derivativedivides	$\frac{a^x}{\ln(a)}$	9
default	$\frac{a^x}{\ln(a)}$	9
risch	$\frac{a^x}{\ln(a)}$	9
parallelrisch	$\frac{a^x}{\ln(a)}$	9
norman	$\frac{e^{x \ln(a)}}{\ln(a)}$	11
meijerg	$-\frac{1-e^{x \ln(a)}}{\ln(a)}$	16

input `int(a^x,x,method=_RETURNVERBOSE)`

output `a^x/ln(a)`

3.4.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int a^x dx = \frac{a^x}{\log(a)}$$

input `integrate(a^x,x, algorithm="fricas")`

output `a^x/log(a)`

3.4.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int a^x dx = \begin{cases} \frac{a^x}{\log(a)} & \text{for } \log(a) \neq 0 \\ x & \text{otherwise} \end{cases}$$

input `integrate(a**x,x)`

output `Piecewise((a**x/log(a), Ne(log(a), 0)), (x, True))`

3.4.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int a^x dx = \frac{a^x}{\log(a)}$$

input `integrate(a^x,x, algorithm="maxima")`

output `a^x/log(a)`

3.4.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int a^x dx = \frac{a^x}{\log(a)}$$

input `integrate(a^x,x, algorithm="giac")`

output `a^x/log(a)`

3.4.9 Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int a^x dx = \frac{a^x}{\ln(a)}$$

input `int(a^x,x)`

output `a^x/log(a)`

3.5 $\int \sin(x) dx$

3.5.1	Optimal result	161
3.5.2	Mathematica [A] (verified)	161
3.5.3	Rubi [A] (verified)	162
3.5.4	Maple [A] (verified)	163
3.5.5	Fricas [A] (verification not implemented)	163
3.5.6	Sympy [A] (verification not implemented)	163
3.5.7	Maxima [A] (verification not implemented)	164
3.5.8	Giac [A] (verification not implemented)	164
3.5.9	Mupad [B] (verification not implemented)	164

3.5.1 Optimal result

Integrand size = 2, antiderivative size = 4

$$\int \sin(x) dx = -\cos(x)$$

output `-cos(x)`

3.5.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \sin(x) dx = -\cos(x)$$

input `Integrate[Sin[x],x]`

output `-Cos[x]`

3.5.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sin(x) dx \\ \downarrow 3042 \\ \int \sin(x) dx \\ \downarrow 3118 \\ -\cos(x) \end{array}$$

input `Int [Sin [x] , x]`

output `-Cos [x]`

3.5.3.1 Defintions of rubi rules used

rule 3042 `Int [u_ , x_Symbol] :> Int [DeactivateTrig [u , x] , x] /; FunctionOfTrigOfLinear Q [u , x]`

rule 3118 `Int [sin [(c_.) + (d_.)*(x_)] , x_Symbol] :> Simp [-Cos [c + d*x] / d , x] /; FreeQ [{c , d} , x]`

3.5.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

method	result	size
lookup	$-\cos(x)$	5
default	$-\cos(x)$	5
risch	$-\cos(x)$	5
parallelrisc	$-\cos(x) - 1$	7
norman	$-\frac{2}{1+\tan^2(\frac{x}{2})}$	13
meijerg	$\sqrt{\pi} \left(\frac{1}{\sqrt{\pi}} - \frac{\cos(x)}{\sqrt{\pi}} \right)$	16

input `int(sin(x),x,method=_RETURNVERBOSE)`

output `-cos(x)`

3.5.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \sin(x) dx = -\cos(x)$$

input `integrate(sin(x),x, algorithm="fricas")`

output `-cos(x)`

3.5.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \sin(x) dx = -\cos(x)$$

input `integrate(sin(x),x)`

output `-cos(x)`

3.5.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \sin(x) dx = -\cos(x)$$

input `integrate(sin(x),x, algorithm="maxima")`

output `-cos(x)`

3.5.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \sin(x) dx = -\cos(x)$$

input `integrate(sin(x),x, algorithm="giac")`

output `-cos(x)`

3.5.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \sin(x) dx = -\cos(x)$$

input `int(sin(x),x)`

output `-cos(x)`

3.6 $\int \cos(x) dx$

3.6.1	Optimal result	165
3.6.2	Mathematica [A] (verified)	165
3.6.3	Rubi [A] (verified)	166
3.6.4	Maple [A] (verified)	167
3.6.5	Fricas [A] (verification not implemented)	167
3.6.6	Sympy [A] (verification not implemented)	167
3.6.7	Maxima [A] (verification not implemented)	168
3.6.8	Giac [A] (verification not implemented)	168
3.6.9	Mupad [B] (verification not implemented)	168

3.6.1 Optimal result

Integrand size = 2, antiderivative size = 2

$$\int \cos(x) dx = \sin(x)$$

output `sin(x)`

3.6.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `Integrate[Cos[x],x]`

output `Sin[x]`

3.6.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \cos(x) dx \\ \downarrow \text{3042} \\ \int \sin\left(x + \frac{\pi}{2}\right) dx \\ \downarrow \text{3117} \\ \sin(x) \end{array}$$

input `Int[Cos[x], x]`

output `Sin[x]`

3.6.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

3.6.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
lookup	$\sin(x)$	3
default	$\sin(x)$	3
meijerg	$\sin(x)$	3
risch	$\sin(x)$	3
parallelrisch	$\sin(x)$	3
norman	$\frac{2 \tan(\frac{x}{2})}{1 + \tan^2(\frac{x}{2})}$	17

input `int(cos(x),x,method=_RETURNVERBOSE)`

output `sin(x)`

3.6.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `integrate(cos(x),x, algorithm="fricas")`

output `sin(x)`

3.6.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `integrate(cos(x),x)`

output `sin(x)`

3.6.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `integrate(cos(x),x, algorithm="maxima")`

output `sin(x)`

3.6.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `integrate(cos(x),x, algorithm="giac")`

output `sin(x)`

3.6.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `int(cos(x),x)`

output `sin(x)`

3.7 $\int \sec^2(x) dx$

3.7.1	Optimal result	169
3.7.2	Mathematica [A] (verified)	169
3.7.3	Rubi [A] (verified)	170
3.7.4	Maple [A] (verified)	171
3.7.5	Fricas [B] (verification not implemented)	171
3.7.6	Sympy [B] (verification not implemented)	171
3.7.7	Maxima [A] (verification not implemented)	172
3.7.8	Giac [A] (verification not implemented)	172
3.7.9	Mupad [B] (verification not implemented)	172

3.7.1 Optimal result

Integrand size = 4, antiderivative size = 2

$$\int \sec^2(x) dx = \tan(x)$$

output `tan(x)`

3.7.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sec^2(x) dx = \tan(x)$$

input `Integrate[Sec[x]^2,x]`

output `Tan[x]`

3.7.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \sec^2(x) dx \\
 \downarrow 3042 \\
 \int \csc\left(x + \frac{\pi}{2}\right)^2 dx \\
 \downarrow 4254 \\
 - \int 1d(-\tan(x)) \\
 \downarrow 24 \\
 \tan(x)
 \end{array}$$

input `Int[Sec[x]^2,x]`

output `Tan[x]`

3.7.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

3.7.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\tan(x)$	3
parallelrisch	$\tan(x)$	3
risch	$\frac{2i}{e^{2ix}+1}$	13
norman	$-\frac{2 \tan(\frac{x}{2})}{\tan^2(\frac{x}{2})-1}$	17

input `int(sec(x)^2,x,method=_RETURNVERBOSE)`

output `tan(x)`

3.7.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7 vs. $2(2) = 4$.

Time = 0.24 (sec) , antiderivative size = 7, normalized size of antiderivative = 3.50

$$\int \sec^2(x) dx = \frac{\sin(x)}{\cos(x)}$$

input `integrate(sec(x)^2,x, algorithm="fricas")`

output `sin(x)/cos(x)`

3.7.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5 vs. $2(2) = 4$.

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 2.50

$$\int \sec^2(x) dx = \frac{\sin(x)}{\cos(x)}$$

input `integrate(sec(x)**2,x)`

output `sin(x)/cos(x)`

3.7.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sec^2(x) dx = \tan(x)$$

input `integrate(sec(x)^2,x, algorithm="maxima")`

output `tan(x)`

3.7.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sec^2(x) dx = \tan(x)$$

input `integrate(sec(x)^2,x, algorithm="giac")`

output `tan(x)`

3.7.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sec^2(x) dx = \tan(x)$$

input `int(1/cos(x)^2,x)`

output `tan(x)`

3.8 $\int \csc^2(x) dx$

3.8.1	Optimal result	173
3.8.2	Mathematica [A] (verified)	173
3.8.3	Rubi [A] (verified)	174
3.8.4	Maple [A] (verified)	175
3.8.5	Fricas [A] (verification not implemented)	175
3.8.6	Sympy [B] (verification not implemented)	175
3.8.7	Maxima [A] (verification not implemented)	176
3.8.8	Giac [A] (verification not implemented)	176
3.8.9	Mupad [B] (verification not implemented)	176

3.8.1 Optimal result

Integrand size = 4, antiderivative size = 4

$$\int \csc^2(x) dx = -\cot(x)$$

output `-cot(x)`

3.8.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \csc^2(x) dx = -\cot(x)$$

input `Integrate[Csc[x]^2,x]`

output `-Cot[x]`

3.8.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \csc^2(x) dx \\
 \downarrow 3042 \\
 \int \csc(x)^2 dx \\
 \downarrow 4254 \\
 - \int 1 d \cot(x) \\
 \downarrow 24 \\
 - \cot(x)
 \end{array}$$

input `Int[Csc[x]^2,x]`

output `-Cot[x]`

3.8.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

3.8.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

method	result	size
default	$-\cot(x)$	5
parallelrisch	$-\cot(x)$	5
risch	$-\frac{2i}{e^{2ix}-1}$	13
norman	$\frac{-\frac{1}{2} + \frac{\tan^2(\frac{x}{2})}{2}}{\tan(\frac{x}{2})}$	18

input `int(csc(x)^2,x,method=_RETURNVERBOSE)`

output `-cot(x)`

3.8.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 8, normalized size of antiderivative = 2.00

$$\int \csc^2(x) dx = -\frac{\cos(x)}{\sin(x)}$$

input `integrate(csc(x)^2,x, algorithm="fricas")`

output `-cos(x)/sin(x)`

3.8.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7 vs. $2(3) = 6$.

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.75

$$\int \csc^2(x) dx = -\frac{\cos(x)}{\sin(x)}$$

input `integrate(csc(x)**2,x)`

output `-cos(x)/sin(x)`

3.8.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.50

$$\int \csc^2(x) dx = -\frac{1}{\tan(x)}$$

input `integrate(csc(x)^2,x, algorithm="maxima")`

output `-1/tan(x)`

3.8.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.50

$$\int \csc^2(x) dx = -\frac{1}{\tan(x)}$$

input `integrate(csc(x)^2,x, algorithm="giac")`

output `-1/tan(x)`

3.8.9 Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \csc^2(x) dx = -\cot(x)$$

input `int(1/sin(x)^2,x)`

output `-\cot(x)`

3.9 $\int \sec(x) \tan(x) dx$

3.9.1	Optimal result	177
3.9.2	Mathematica [A] (verified)	177
3.9.3	Rubi [A] (verified)	178
3.9.4	Maple [A] (verified)	179
3.9.5	Fricas [A] (verification not implemented)	179
3.9.6	Sympy [A] (verification not implemented)	179
3.9.7	Maxima [A] (verification not implemented)	180
3.9.8	Giac [A] (verification not implemented)	180
3.9.9	Mupad [B] (verification not implemented)	180

3.9.1 Optimal result

Integrand size = 5, antiderivative size = 2

$$\int \sec(x) \tan(x) dx = \sec(x)$$

output

```
sec(x)
```

3.9.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sec(x) \tan(x) dx = \sec(x)$$

input `Integrate[Sec[x]*Tan[x],x]`

output

```
Sec[x]
```

3.9.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 3086, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \tan(x) \sec(x) dx \\
 \downarrow \text{3042} \\
 \int \tan(x) \sec(x) dx \\
 \downarrow \text{3086} \\
 \int 1 d \sec(x) \\
 \downarrow \text{24} \\
 \sec(x)
 \end{array}$$

input `Int[Sec[x]*Tan[x],x]`

output `Sec[x]`

3.9.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

3.9.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
derivativedivides	$\sec(x)$	3
default	$\sec(x)$	3
risch	$\frac{2e^{ix}}{e^{2ix}+1}$	17

input `int(sec(x)*tan(x),x,method=_RETURNVERBOSE)`

output `sec(x)`

3.9.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 4, normalized size of antiderivative = 2.00

$$\int \sec(x) \tan(x) dx = \frac{1}{\cos(x)}$$

input `integrate(sec(x)*tan(x),x, algorithm="fricas")`

output `1/cos(x)`

3.9.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

$$\int \sec(x) \tan(x) dx = \frac{1}{\cos(x)}$$

input `integrate(sec(x)*tan(x),x)`

output `1/cos(x)`

3.9.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 4, normalized size of antiderivative = 2.00

$$\int \sec(x) \tan(x) dx = \frac{1}{\cos(x)}$$

input `integrate(sec(x)*tan(x),x, algorithm="maxima")`output `1/cos(x)`**3.9.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 4, normalized size of antiderivative = 2.00

$$\int \sec(x) \tan(x) dx = \frac{1}{\cos(x)}$$

input `integrate(sec(x)*tan(x),x, algorithm="giac")`output `1/cos(x)`**3.9.9 Mupad [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 6.00

$$\int \sec(x) \tan(x) dx = -\frac{2}{\tan\left(\frac{x}{2}\right)^2 - 1}$$

input `int(tan(x)/cos(x),x)`output `-2/(tan(x/2)^2 - 1)`

3.10 $\int \cot(x) \csc(x) dx$

3.10.1	Optimal result	181
3.10.2	Mathematica [A] (verified)	181
3.10.3	Rubi [A] (verified)	182
3.10.4	Maple [A] (verified)	183
3.10.5	Fricas [A] (verification not implemented)	183
3.10.6	Sympy [A] (verification not implemented)	184
3.10.7	Maxima [A] (verification not implemented)	184
3.10.8	Giac [A] (verification not implemented)	184
3.10.9	Mupad [B] (verification not implemented)	185

3.10.1 Optimal result

Integrand size = 5, antiderivative size = 4

$$\int \cot(x) \csc(x) dx = -\csc(x)$$

output `-csc(x)`

3.10.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \cot(x) \csc(x) dx = -\csc(x)$$

input `Integrate[Cot[x]*Csc[x],x]`

output `-Csc[x]`

3.10.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {3042, 25, 3086, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot(x) \csc(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan\left(x - \frac{\pi}{2}\right) \left(-\sec\left(x - \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{25} \\
 & - \int \sec\left(x - \frac{\pi}{2}\right) \tan\left(x - \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3086} \\
 & - \int 1 d \csc(x) \\
 & \quad \downarrow \text{24} \\
 & - \csc(x)
 \end{aligned}$$

input `Int[Cot[x]*Csc[x],x]`

output `-Csc[x]`

3.10.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3086 Int[((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] :> Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])
```

3.10.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

method	result	size
derivativedivides	$-\csc(x)$	5
default	$-\csc(x)$	5
risch	$-\frac{2ie^{ix}}{e^{2ix}-1}$	18

```
input int(csc(x)*cot(x),x,method=_RETURNVERBOSE)
```

```
output -csc(x)
```

3.10.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.50

$$\int \cot(x) \csc(x) dx = -\frac{1}{\sin(x)}$$

```
input integrate(cot(x)*csc(x),x, algorithm="fricas")
```

```
output -1/sin(x)
```


3.10.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

$$\int \cot(x) \csc(x) dx = -\frac{1}{\sin(x)}$$

input `integrate(cot(x)*csc(x),x)`

output `-1/sin(x)`

3.10.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.50

$$\int \cot(x) \csc(x) dx = -\frac{1}{\sin(x)}$$

input `integrate(cot(x)*csc(x),x, algorithm="maxima")`

output `-1/sin(x)`

3.10.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.50

$$\int \cot(x) \csc(x) dx = -\frac{1}{\sin(x)}$$

input `integrate(cot(x)*csc(x),x, algorithm="giac")`

output `-1/sin(x)`

3.10.9 Mupad [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.50

$$\int \cot(x) \csc(x) dx = -\frac{1}{\sin(x)}$$

input `int(cot(x)/sin(x),x)`

output `-1/sin(x)`

3.11 $\int \sinh(x) dx$

3.11.1	Optimal result	186
3.11.2	Mathematica [A] (verified)	186
3.11.3	Rubi [A] (verified)	187
3.11.4	Maple [A] (verified)	188
3.11.5	Fricas [A] (verification not implemented)	188
3.11.6	Sympy [A] (verification not implemented)	188
3.11.7	Maxima [A] (verification not implemented)	189
3.11.8	Giac [B] (verification not implemented)	189
3.11.9	Mupad [B] (verification not implemented)	189

3.11.1 Optimal result

Integrand size = 2, antiderivative size = 2

$$\int \sinh(x) dx = \cosh(x)$$

output `cosh(x)`

3.11.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sinh(x) dx = \cosh(x)$$

input `Integrate[Sinh[x],x]`

output `Cosh[x]`

3.11.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {3042, 26, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sinh(x) dx \\ \downarrow 3042 \\ \int -i \sin(ix) dx \\ \downarrow 26 \\ -i \int \sin(ix) dx \\ \downarrow 3118 \\ \cosh(x) \end{array}$$

input `Int[Sinh[x],x]`

output `Cosh[x]`

3.11.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

3.11.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
lookup	$\cosh(x)$	3
default	$\cosh(x)$	3
risch	$\frac{e^x}{2} + \frac{e^{-x}}{2}$	12
parallelrisc	$-\frac{2}{\tanh^2(\frac{x}{2})-1}$	13
meijerg	$-\sqrt{\pi} \left(\frac{1}{\sqrt{\pi}} - \frac{\cosh(x)}{\sqrt{\pi}} \right)$	17

input `int(sinh(x),x,method=_RETURNVERBOSE)`

output `cosh(x)`

3.11.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sinh(x) dx = \cosh(x)$$

input `integrate(sinh(x),x, algorithm="fricas")`

output `cosh(x)`

3.11.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sinh(x) dx = \cosh(x)$$

input `integrate(sinh(x),x)`

output `cosh(x)`

3.11.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sinh(x) dx = \cosh(x)$$

input `integrate(sinh(x),x, algorithm="maxima")`

output `cosh(x)`

3.11.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. 2(2) = 4.

Time = 0.29 (sec) , antiderivative size = 11, normalized size of antiderivative = 5.50

$$\int \sinh(x) dx = \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x$$

input `integrate(sinh(x),x, algorithm="giac")`

output `1/2*e^(-x) + 1/2*e^x`

3.11.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sinh(x) dx = \cosh(x)$$

input `int(sinh(x),x)`

output `cosh(x)`

3.12 $\int \cosh(x) dx$

3.12.1	Optimal result	190
3.12.2	Mathematica [A] (verified)	190
3.12.3	Rubi [A] (verified)	191
3.12.4	Maple [A] (verified)	192
3.12.5	Fricas [A] (verification not implemented)	192
3.12.6	Sympy [A] (verification not implemented)	192
3.12.7	Maxima [A] (verification not implemented)	193
3.12.8	Giac [B] (verification not implemented)	193
3.12.9	Mupad [B] (verification not implemented)	193

3.12.1 Optimal result

Integrand size = 2, antiderivative size = 2

$$\int \cosh(x) dx = \sinh(x)$$

output `sinh(x)`

3.12.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cosh(x) dx = \sinh(x)$$

input `Integrate[Cosh[x],x]`

output `Sinh[x]`

3.12.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \cosh(x) dx \\ \downarrow \text{3042} \\ \int \sin\left(\frac{\pi}{2} + ix\right) dx \\ \downarrow \text{3117} \\ \sinh(x) \end{array}$$

input `Int[Cosh[x], x]`

output `Sinh[x]`

3.12.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

3.12.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
lookup	$\sinh(x)$	3
default	$\sinh(x)$	3
meijerg	$\sinh(x)$	3
parallelsch	$\sinh(x)$	3
risch	$-\frac{e^{-x}}{2} + \frac{e^x}{2}$	12

input `int(cosh(x),x,method=_RETURNVERBOSE)`

output `sinh(x)`

3.12.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cosh(x) dx = \sinh(x)$$

input `integrate(cosh(x),x, algorithm="fricas")`

output `sinh(x)`

3.12.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cosh(x) dx = \sinh(x)$$

input `integrate(cosh(x),x)`

output `sinh(x)`

3.12.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cosh(x) dx = \sinh(x)$$

input `integrate(cosh(x),x, algorithm="maxima")`

output `sinh(x)`

3.12.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. 2(2) = 4.

Time = 0.30 (sec) , antiderivative size = 11, normalized size of antiderivative = 5.50

$$\int \cosh(x) dx = -\frac{1}{2} e^{(-x)} + \frac{1}{2} e^x$$

input `integrate(cosh(x),x, algorithm="giac")`

output `-1/2*e^(-x) + 1/2*e^x`

3.12.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cosh(x) dx = \sinh(x)$$

input `int(cosh(x),x)`

output `sinh(x)`

3.13 $\int \tan(x) dx$

3.13.1	Optimal result	194
3.13.2	Mathematica [A] (verified)	194
3.13.3	Rubi [A] (verified)	195
3.13.4	Maple [A] (verified)	196
3.13.5	Fricas [B] (verification not implemented)	196
3.13.6	Sympy [A] (verification not implemented)	197
3.13.7	Maxima [A] (verification not implemented)	197
3.13.8	Giac [A] (verification not implemented)	197
3.13.9	Mupad [B] (verification not implemented)	198

3.13.1 Optimal result

Integrand size = 2, antiderivative size = 5

$$\int \tan(x) dx = -\log(\cos(x))$$

output `-ln(cos(x))`

3.13.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \tan(x) dx = -\log(\cos(x))$$

input `Integrate[Tan[x],x]`

output `-Log[Cos[x]]`

3.13.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \tan(x) dx \\ \downarrow 3042 \\ \int \tan(x) dx \\ \downarrow 3956 \\ -\log(\cos(x)) \end{array}$$

input `Int [Tan [x] , x]`

output `-Log [Cos [x]]`

3.13.3.1 Defintions of rubi rules used

rule 3042 `Int [u_ , x_Symbol] := Int [DeactivateTrig [u , x] , x] /; FunctionOfTrigOfLinear Q [u , x]`

rule 3956 `Int [tan [(c_) + (d_)*(x_)] , x_Symbol] := Simp [-Log [RemoveContent [Cos [c + d *x] , x]]/d , x] /; FreeQ [{c , d} , x]`

3.13.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
lookup	$-\ln(\cos(x))$	6
default	$-\ln(\cos(x))$	6
derivativedivides	$\frac{\ln(1+\tan^2(x))}{2}$	10
norman	$\frac{\ln(1+\tan^2(x))}{2}$	10
parallelrisc	$\frac{\ln(1+\tan^2(x))}{2}$	10
risc	$ix - \ln(e^{2ix} + 1)$	16

input `int(tan(x),x,method=_RETURNVERBOSE)`

output `-ln(cos(x))`

3.13.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. $2(5) = 10$.

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 2.20

$$\int \tan(x) dx = -\frac{1}{2} \log\left(\frac{1}{\tan(x)^2 + 1}\right)$$

input `integrate(tan(x),x, algorithm="fracas")`

output `-1/2*log(1/(tan(x)^2 + 1))`

3.13.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \tan(x) dx = -\log(\cos(x))$$

input `integrate(tan(x),x)`

output `-log(cos(x))`

3.13.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.60

$$\int \tan(x) dx = \log(\sec(x))$$

input `integrate(tan(x),x, algorithm="maxima")`

output `log(sec(x))`

3.13.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

$$\int \tan(x) dx = -\log(|\cos(x)|)$$

input `integrate(tan(x),x, algorithm="giac")`

output `-log(abs(cos(x)))`

3.13.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \tan(x) dx = -\ln(\cos(x))$$

input `int(tan(x),x)`

output `-log(cos(x))`

3.14 $\int \cot(x) dx$

3.14.1	Optimal result	199
3.14.2	Mathematica [B] (verified)	199
3.14.3	Rubi [A] (verified)	200
3.14.4	Maple [A] (verified)	201
3.14.5	Fricas [B] (verification not implemented)	201
3.14.6	Sympy [A] (verification not implemented)	202
3.14.7	Maxima [A] (verification not implemented)	202
3.14.8	Giac [A] (verification not implemented)	202
3.14.9	Mupad [B] (verification not implemented)	203

3.14.1 Optimal result

Integrand size = 2, antiderivative size = 3

$$\int \cot(x) dx = \log(\sin(x))$$

output `ln(sin(x))`

3.14.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 7 vs. $2(3) = 6$.

Time = 0.00 (sec) , antiderivative size = 7, normalized size of antiderivative = 2.33

$$\int \cot(x) dx = \log(\cos(x)) + \log(\tan(x))$$

input `Integrate[Cot[x], x]`

output `Log[Cos[x]] + Log[Tan[x]]`

3.14.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \cot(x) dx \\
 \downarrow \text{3042} \\
 \int -\tan\left(x + \frac{\pi}{2}\right) dx \\
 \downarrow \text{25} \\
 -\int \tan\left(x + \frac{\pi}{2}\right) dx \\
 \downarrow \text{3956} \\
 \log(\sin(x))
 \end{array}$$

input `Int[Cot[x], x]`

output `Log[Sin[x]]`

3.14.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

3.14.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
lookup	$\ln(\sin(x))$	4
default	$\ln(\sin(x))$	4
derivativedivides	$-\frac{\ln(\cot^2(x)+1)}{2}$	10
parallelrisc	$\ln\left(\frac{1}{\sqrt{\sec^2(x)}}\right) + \ln(\tan(x))$	12
norman	$-\frac{\ln(1+\tan^2(x))}{2} + \ln(\tan(x))$	14
risc	$-ix + \ln(e^{2ix} - 1)$	14

input `int(cot(x),x,method=_RETURNVERBOSE)`

output `ln(sin(x))`

3.14.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. $2(3) = 6$.

Time = 0.25 (sec) , antiderivative size = 11, normalized size of antiderivative = 3.67

$$\int \cot(x) dx = \frac{1}{2} \log\left(-\frac{1}{2} \cos(2x) + \frac{1}{2}\right)$$

input `integrate(cot(x),x, algorithm="fricas")`

output `1/2*log(-1/2*cos(2*x) + 1/2)`

3.14.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \cot(x) dx = \log(\sin(x))$$

input `integrate(cot(x),x)`

output `log(sin(x))`

3.14.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \cot(x) dx = \log(\sin(x))$$

input `integrate(cot(x),x, algorithm="maxima")`

output `log(sin(x))`

3.14.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

$$\int \cot(x) dx = \log(|\sin(x)|)$$

input `integrate(cot(x),x, algorithm="giac")`

output `log(abs(sin(x)))`

3.14.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \cot(x) dx = \ln(\sin(x))$$

input `int(cot(x),x)`

output `log(sin(x))`

3.15 $\int x \sin(x) dx$

3.15.1	Optimal result	204
3.15.2	Mathematica [A] (verified)	204
3.15.3	Rubi [A] (verified)	205
3.15.4	Maple [A] (verified)	206
3.15.5	Fricas [A] (verification not implemented)	206
3.15.6	Sympy [A] (verification not implemented)	207
3.15.7	Maxima [A] (verification not implemented)	207
3.15.8	Giac [A] (verification not implemented)	207
3.15.9	Mupad [B] (verification not implemented)	208

3.15.1 Optimal result

Integrand size = 4, antiderivative size = 8

$$\int x \sin(x) dx = -x \cos(x) + \sin(x)$$

output `-x*cos(x)+sin(x)`

3.15.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int x \sin(x) dx = -x \cos(x) + \sin(x)$$

input `Integrate[x*Sin[x],x]`

output `-(x*Cos[x]) + Sin[x]`

3.15.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int x \sin(x) dx \\
 \downarrow \text{3042} \\
 \int x \sin(x) dx \\
 \downarrow \text{3777} \\
 \int \cos(x) dx - x \cos(x) \\
 \downarrow \text{3042} \\
 \int \sin\left(x + \frac{\pi}{2}\right) dx - x \cos(x) \\
 \downarrow \text{3117} \\
 \sin(x) - x \cos(x)
 \end{array}$$

input `Int[x*Sin[x],x]`

output `-(x*Cos[x]) + Sin[x]`

3.15.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

```
rule 3777 Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-
(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*C
os[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

3.15.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
default	$-x \cos(x) + \sin(x)$	9
risch	$-x \cos(x) + \sin(x)$	9
parallelrisch	$-x \cos(x) + \sin(x)$	9
parts	$-x \cos(x) + \sin(x)$	9
meijerg	$2\sqrt{\pi} \left(-\frac{x \cos(x)}{2\sqrt{\pi}} + \frac{\sin(x)}{2\sqrt{\pi}} \right)$	22
norman	$\frac{x(\tan^2(\frac{x}{2}) - x + 2 \tan(\frac{x}{2}))}{1 + \tan^2(\frac{x}{2})}$	30

```
input int(x*sin(x),x,method=_RETURNVERBOSE)
```

```
output -x*cos(x)+sin(x)
```

3.15.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int x \sin(x) dx = -x \cos(x) + \sin(x)$$

```
input integrate(x*sin(x),x, algorithm="fricas")
```

```
output -x*cos(x) + sin(x)
```

3.15.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int x \sin(x) dx = -x \cos(x) + \sin(x)$$

input `integrate(x*sin(x),x)`

output `-x*cos(x) + sin(x)`

3.15.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int x \sin(x) dx = -x \cos(x) + \sin(x)$$

input `integrate(x*sin(x),x, algorithm="maxima")`

output `-x*cos(x) + sin(x)`

3.15.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int x \sin(x) dx = -x \cos(x) + \sin(x)$$

input `integrate(x*sin(x),x, algorithm="giac")`

output `-x*cos(x) + sin(x)`

3.15.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int x \sin(x) dx = \sin(x) - x \cos(x)$$

input `int(x*sin(x),x)`

output `sin(x) - x*cos(x)`

3.16 $\int \log(x) dx$

3.16.1	Optimal result	209
3.16.2	Mathematica [A] (verified)	209
3.16.3	Rubi [A] (verified)	210
3.16.4	Maple [A] (verified)	210
3.16.5	Fricas [A] (verification not implemented)	211
3.16.6	Sympy [A] (verification not implemented)	211
3.16.7	Maxima [A] (verification not implemented)	211
3.16.8	Giac [A] (verification not implemented)	212
3.16.9	Mupad [B] (verification not implemented)	212

3.16.1 Optimal result

Integrand size = 2, antiderivative size = 8

$$\int \log(x) dx = -x + x \log(x)$$

output `-x+x*ln(x)`

3.16.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \log(x) dx = -x + x \log(x)$$

input `Integrate[Log[x],x]`

output `-x + x*Log[x]`

3.16.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {2732}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \log(x) dx$$

$$\downarrow \text{2732}$$

$$x \log(x) - x$$

input `Int [Log[x] , x]`

output `-x + x*Log[x]`

3.16.3.1 Defintions of rubi rules used

rule 2732 `Int [Log[(c_.)*(x_)^(n_.)], x_Symbol] :> Simp[x*Log[c*x^n], x] - Simp[n*x, x] /; FreeQ[{c, n}, x]`

3.16.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
lookup	$-x + x \ln(x)$	9
default	$-x + x \ln(x)$	9
norman	$-x + x \ln(x)$	9
risch	$-x + x \ln(x)$	9
parallelrisch	$-x + x \ln(x)$	9
parts	$-x + x \ln(x)$	9

input `int(ln(x), x, method=_RETURNVERBOSE)`

output `-x+x*ln(x)`

3.16.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \log(x) dx = x \log(x) - x$$

input `integrate(log(x),x, algorithm="fricas")`

output `x*log(x) - x`

3.16.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \log(x) dx = x \log(x) - x$$

input `integrate(ln(x),x)`

output `x*log(x) - x`

3.16.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \log(x) dx = x \log(x) - x$$

input `integrate(log(x),x, algorithm="maxima")`

output `x*log(x) - x`

3.16.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \log(x) dx = x \log(x) - x$$

input `integrate(log(x),x, algorithm="giac")`

output `x*log(x) - x`

3.16.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \log(x) dx = x (\ln(x) - 1)$$

input `int(log(x),x)`

output `x*(log(x) - 1)`

3.17 $\int e^x x^2 dx$

3.17.1	Optimal result	213
3.17.2	Mathematica [A] (verified)	213
3.17.3	Rubi [A] (verified)	214
3.17.4	Maple [A] (verified)	215
3.17.5	Fricas [A] (verification not implemented)	215
3.17.6	Sympy [A] (verification not implemented)	215
3.17.7	Maxima [A] (verification not implemented)	216
3.17.8	Giac [A] (verification not implemented)	216
3.17.9	Mupad [B] (verification not implemented)	216

3.17.1 Optimal result

Integrand size = 7, antiderivative size = 19

$$\int e^x x^2 dx = 2e^x - 2e^x x + e^x x^2$$

output `2*exp(x)-2*exp(x)*x+exp(x)*x^2`

3.17.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int e^x x^2 dx = e^x (2 - 2x + x^2)$$

input `Integrate[E^x*x^2,x]`

output `E^x*(2 - 2*x + x^2)`

3.17.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {2607, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int e^x x^2 dx \\
 \downarrow \text{2607} \\
 e^x x^2 - 2 \int e^x x dx \\
 \downarrow \text{2607} \\
 e^x x^2 - 2 \left(e^x x - \int e^x dx \right) \\
 \downarrow \text{2624} \\
 e^x x^2 - 2(e^x x - e^x)
 \end{array}$$

input `Int[E^x*x^2,x]`

output `E^x*x^2 - 2*(-E^x + E^x*x)`

3.17.3.1 Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2624 `Int[((F_)^(v_))^(n_.), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`

3.17.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

method	result	size
gospers	$(x^2 - 2x + 2)e^x$	12
risch	$(x^2 - 2x + 2)e^x$	12
default	$2e^x - 2e^x x + e^x x^2$	17
norman	$2e^x - 2e^x x + e^x x^2$	17
meijerg	$-2 + \frac{(3x^2 - 6x + 6)e^x}{3}$	17
parallelrisc	$2e^x - 2e^x x + e^x x^2$	17
parts	$2e^x - 2e^x x + e^x x^2$	17

input `int(exp(x)*x^2,x,method=_RETURNVERBOSE)`

output `(x^2-2*x+2)*exp(x)`

3.17.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x x^2 dx = (x^2 - 2x + 2)e^x$$

input `integrate(exp(x)*x^2,x, algorithm="fricas")`

output `(x^2 - 2*x + 2)*e^x`

3.17.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.53

$$\int e^x x^2 dx = (x^2 - 2x + 2)e^x$$

input `integrate(exp(x)*x**2,x)`

output `(x**2 - 2*x + 2)*exp(x)`

3.17.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x x^2 dx = (x^2 - 2x + 2)e^x$$

input `integrate(exp(x)*x^2,x, algorithm="maxima")`output `(x^2 - 2*x + 2)*e^x`**3.17.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x x^2 dx = (x^2 - 2x + 2)e^x$$

input `integrate(exp(x)*x^2,x, algorithm="giac")`output `(x^2 - 2*x + 2)*e^x`**3.17.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x x^2 dx = e^x (x^2 - 2x + 2)$$

input `int(x^2*exp(x),x)`output `exp(x)*(x^2 - 2*x + 2)`

3.18 $\int e^x \sin(x) dx$

3.18.1	Optimal result	217
3.18.2	Mathematica [A] (verified)	217
3.18.3	Rubi [A] (verified)	218
3.18.4	Maple [A] (verified)	218
3.18.5	Fricas [A] (verification not implemented)	219
3.18.6	Sympy [A] (verification not implemented)	219
3.18.7	Maxima [A] (verification not implemented)	219
3.18.8	Giac [A] (verification not implemented)	220
3.18.9	Mupad [B] (verification not implemented)	220

3.18.1 Optimal result

Integrand size = 6, antiderivative size = 19

$$\int e^x \sin(x) dx = -\frac{1}{2}e^x \cos(x) + \frac{1}{2}e^x \sin(x)$$

output `-1/2*exp(x)*cos(x)+1/2*exp(x)*sin(x)`

3.18.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int e^x \sin(x) dx = \frac{1}{2}e^x(-\cos(x) + \sin(x))$$

input `Integrate[E^x*Sin[x],x]`

output `(E^x*(-Cos[x] + Sin[x]))/2`

3.18.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {4932}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x \sin(x) dx$$

$$\downarrow 4932$$

$$\frac{1}{2}e^x \sin(x) - \frac{1}{2}e^x \cos(x)$$

input `Int[E^x*Sin[x],x]`

output `-1/2*(E^x*Cos[x]) + (E^x*Sin[x])/2`

3.18.3.1 Defintions of rubi rules used

rule 4932 `Int[(F_)^((c_.)*((a_.) + (b_.)*(x_)))*Sin[(d_.) + (e_.)*(x_)], x_Symbol] :=
Simp[b*c*Log[F]*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x
] - Simp[e*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; F
reeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]`

3.18.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

method	result	size
parallelrisch	$-\frac{e^x(\cos(x)-\sin(x))}{2}$	12
default	$-\frac{e^x \cos(x)}{2} + \frac{e^x \sin(x)}{2}$	14
norman	$\frac{e^x \tan(\frac{x}{2}) + \frac{e^x (\tan^2(\frac{x}{2}))}{2} - \frac{e^x}{2}}{1 + \tan^2(\frac{x}{2})}$	34
risch	$-\frac{e^{(1+i)x}}{4} - \frac{ie^{(1+i)x}}{4} - \frac{e^{(1-i)x}}{4} + \frac{ie^{(1-i)x}}{4}$	36

input `int(exp(x)*sin(x),x,method=_RETURNVERBOSE)`

output `-1/2*exp(x)*(cos(x)-sin(x))`

3.18.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int e^x \sin(x) dx = -\frac{1}{2} \cos(x) e^x + \frac{1}{2} e^x \sin(x)$$

input `integrate(exp(x)*sin(x),x, algorithm="fricas")`

output `-1/2*cos(x)*e^x + 1/2*e^x*sin(x)`

3.18.6 Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int e^x \sin(x) dx = \frac{e^x \sin(x)}{2} - \frac{e^x \cos(x)}{2}$$

input `integrate(exp(x)*sin(x),x)`

output `exp(x)*sin(x)/2 - exp(x)*cos(x)/2`

3.18.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x \sin(x) dx = -\frac{1}{2} (\cos(x) - \sin(x)) e^x$$

input `integrate(exp(x)*sin(x),x, algorithm="maxima")`

output `-1/2*(cos(x) - sin(x))*e^x`

3.18.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x \sin(x) dx = -\frac{1}{2} (\cos(x) - \sin(x))e^x$$

input `integrate(exp(x)*sin(x),x, algorithm="giac")`

output `-1/2*(cos(x) - sin(x))*e^x`

3.18.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int e^x \sin(x) dx = -\frac{e^x (\cos(x) - \sin(x))}{2}$$

input `int(exp(x)*sin(x),x)`

output `-(exp(x)*(cos(x) - sin(x)))/2`

3.19 $\int \arctan(x) dx$

3.19.1	Optimal result	221
3.19.2	Mathematica [A] (verified)	221
3.19.3	Rubi [A] (verified)	222
3.19.4	Maple [A] (verified)	223
3.19.5	Fricas [A] (verification not implemented)	223
3.19.6	Sympy [A] (verification not implemented)	224
3.19.7	Maxima [A] (verification not implemented)	224
3.19.8	Giac [A] (verification not implemented)	224
3.19.9	Mupad [B] (verification not implemented)	225

3.19.1 Optimal result

Integrand size = 2, antiderivative size = 15

$$\int \arctan(x) dx = x \arctan(x) - \frac{1}{2} \log(1 + x^2)$$

output `x*arctan(x)-1/2*ln(x^2+1)`

3.19.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \arctan(x) dx = x \arctan(x) - \frac{1}{2} \log(1 + x^2)$$

input `Integrate[ArcTan[x],x]`

output `x*ArcTan[x] - Log[1 + x^2]/2`

3.19.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {5345, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan(x) dx$$

$$\downarrow \text{5345}$$

$$x \arctan(x) - \int \frac{x}{x^2 + 1} dx$$

$$\downarrow \text{240}$$

$$x \arctan(x) - \frac{1}{2} \log(x^2 + 1)$$

input `Int[ArcTan[x], x]`

output `x*ArcTan[x] - Log[1 + x^2]/2`

3.19.3.1 Defintions of rubi rules used

rule 240 `Int[(x_)/((a_) + (b_)*(x_)^2), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`

rule 5345 `Int[((a_) + ArcTan[(c_)*(x_)^(n_)])*(b_)^(p_), x_Symbol] :> Simp[x*(a + b*ArcTan[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`

3.19.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
lookup	$x \arctan(x) - \frac{\ln(x^2+1)}{2}$	14
default	$x \arctan(x) - \frac{\ln(x^2+1)}{2}$	14
parallelsch	$x \arctan(x) - \frac{\ln(x^2+1)}{2}$	14
parts	$x \arctan(x) - \frac{\ln(x^2+1)}{2}$	14
meijerg	$\frac{x^2 \arctan(\sqrt{x^2})}{\sqrt{x^2}} - \frac{\ln(x^2+1)}{2}$	25
risch	$-\frac{ix \ln(ix+1)}{2} + \frac{ix \ln(-ix+1)}{2} - \frac{\ln(x^2+1)}{2}$	32

input `int(arctan(x),x,method=_RETURNVERBOSE)`output `x*arctan(x)-1/2*ln(x^2+1)`**3.19.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \arctan(x) dx = x \arctan(x) - \frac{1}{2} \log(x^2 + 1)$$

input `integrate(arctan(x),x, algorithm="fricas")`output `x*arctan(x) - 1/2*log(x^2 + 1)`

3.19.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \arctan(x) dx = x \operatorname{atan}(x) - \frac{\log(x^2 + 1)}{2}$$

input `integrate(atan(x),x)`output `x*atan(x) - log(x**2 + 1)/2`**3.19.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \arctan(x) dx = x \arctan(x) - \frac{1}{2} \log(x^2 + 1)$$

input `integrate(arctan(x),x, algorithm="maxima")`output `x*arctan(x) - 1/2*log(x^2 + 1)`**3.19.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \arctan(x) dx = x \arctan(x) - \frac{1}{2} \log(x^2 + 1)$$

input `integrate(arctan(x),x, algorithm="giac")`output `x*arctan(x) - 1/2*log(x^2 + 1)`

3.19.9 Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \arctan(x) dx = x \operatorname{atan}(x) - \frac{\ln(x^2 + 1)}{2}$$

input `int(atan(x),x)`

output `x*atan(x) - log(x^2 + 1)/2`

3.20 $\int e^{2x} x dx$

3.20.1	Optimal result	226
3.20.2	Mathematica [A] (verified)	226
3.20.3	Rubi [A] (verified)	227
3.20.4	Maple [A] (verified)	228
3.20.5	Fricas [A] (verification not implemented)	228
3.20.6	Sympy [A] (verification not implemented)	229
3.20.7	Maxima [A] (verification not implemented)	229
3.20.8	Giac [A] (verification not implemented)	229
3.20.9	Mupad [B] (verification not implemented)	230

3.20.1 Optimal result

Integrand size = 7, antiderivative size = 20

$$\int e^{2x} x dx = -\frac{e^{2x}}{4} + \frac{1}{2}e^{2x}x$$

output `-1/4*exp(2*x)+1/2*exp(2*x)*x`

3.20.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int e^{2x} x dx = e^{2x} \left(-\frac{1}{4} + \frac{x}{2} \right)$$

input `Integrate[E^(2*x)*x,x]`

output `E^(2*x)*(-1/4 + x/2)`

3.20.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{2x} x dx$$

$$\downarrow \text{2607}$$

$$\frac{1}{2}e^{2x}x - \frac{\int e^{2x} dx}{2}$$

$$\downarrow \text{2624}$$

$$\frac{1}{2}e^{2x}x - \frac{e^{2x}}{4}$$

input `Int[E^(2*x)*x,x]`

output `-1/4*E^(2*x) + (E^(2*x)*x)/2`

3.20.3.1 Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*(b*F^(g*(e + f*x)))^n/(f*g*n*Log[F]), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2624 `Int[((F_)^(v_))^(n_.), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`

3.20.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.55

method	result	size
risch	$\left(-\frac{1}{4} + \frac{x}{2}\right) e^{2x}$	11
gospers	$\frac{(2x-1)e^{2x}}{4}$	12
meijerg	$\frac{1}{4} - \frac{(2-4x)e^{2x}}{8}$	14
derivativdivides	$-\frac{e^{2x}}{4} + \frac{e^{2x}x}{2}$	15
default	$-\frac{e^{2x}}{4} + \frac{e^{2x}x}{2}$	15
norman	$-\frac{e^{2x}}{4} + \frac{e^{2x}x}{2}$	15
parallelrisch	$-\frac{e^{2x}}{4} + \frac{e^{2x}x}{2}$	15
parts	$-\frac{e^{2x}}{4} + \frac{e^{2x}x}{2}$	15

input `int(exp(2*x)*x,x,method=_RETURNVERBOSE)`output `(-1/4+1/2*x)*exp(2*x)`**3.20.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.55

$$\int e^{2x} x dx = \frac{1}{4} (2x - 1) e^{(2x)}$$

input `integrate(exp(2*x)*x,x, algorithm="fricas")`output `1/4*(2*x - 1)*e^(2*x)`

3.20.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.50

$$\int e^{2x} x dx = \frac{(2x - 1) e^{2x}}{4}$$

input `integrate(exp(2*x)*x,x)`output `(2*x - 1)*exp(2*x)/4`**3.20.7 Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.55

$$\int e^{2x} x dx = \frac{1}{4} (2x - 1) e^{(2x)}$$

input `integrate(exp(2*x)*x,x, algorithm="maxima")`output `1/4*(2*x - 1)*e^(2*x)`**3.20.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.55

$$\int e^{2x} x dx = \frac{1}{4} (2x - 1) e^{(2x)}$$

input `integrate(exp(2*x)*x,x, algorithm="giac")`output `1/4*(2*x - 1)*e^(2*x)`

3.20.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.55

$$\int e^{2x} x dx = \frac{e^{2x} (2x - 1)}{4}$$

input `int(x*exp(2*x),x)`

output `(exp(2*x)*(2*x - 1))/4`

3.21 $\int x \cos(x) dx$

3.21.1	Optimal result	231
3.21.2	Mathematica [A] (verified)	231
3.21.3	Rubi [A] (verified)	232
3.21.4	Maple [A] (verified)	233
3.21.5	Fricas [A] (verification not implemented)	234
3.21.6	Sympy [A] (verification not implemented)	234
3.21.7	Maxima [A] (verification not implemented)	234
3.21.8	Giac [A] (verification not implemented)	235
3.21.9	Mupad [B] (verification not implemented)	235

3.21.1 Optimal result

Integrand size = 4, antiderivative size = 7

$$\int x \cos(x) dx = \cos(x) + x \sin(x)$$

output `cos(x)+x*sin(x)`

3.21.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = \cos(x) + x \sin(x)$$

input `Integrate[x*Cos[x],x]`

output `Cos[x] + x*Sin[x]`

3.21.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \cos(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x \sin\left(x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3777} \\
 & \int -\sin(x) dx + x \sin(x) \\
 & \quad \downarrow \text{25} \\
 & x \sin(x) - \int \sin(x) dx \\
 & \quad \downarrow \text{3042} \\
 & x \sin(x) - \int \sin(x) dx \\
 & \quad \downarrow \text{3118} \\
 & x \sin(x) + \cos(x)
 \end{aligned}$$

input `Int [x*Cos [x] , x]`

output `Cos [x] + x*Sin [x]`

3.21.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

3.21.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

method	result	size
default	$\cos(x) + x \sin(x)$	8
risch	$\cos(x) + x \sin(x)$	8
parts	$\cos(x) + x \sin(x)$	8
parallelrisc	$x \sin(x) + \cos(x) + 1$	9
norman	$\frac{2x \tan(\frac{x}{2}) + 2}{1 + \tan^2(\frac{x}{2})}$	21
meijerg	$2\sqrt{\pi} \left(-\frac{1}{2\sqrt{\pi}} + \frac{\cos(x)}{2\sqrt{\pi}} + \frac{x \sin(x)}{2\sqrt{\pi}} \right)$	27

input `int(x*cos(x),x,method=_RETURNVERBOSE)`

output `cos(x)+x*sin(x)`

3.21.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = x \sin(x) + \cos(x)$$

input `integrate(x*cos(x),x, algorithm="fricas")`

output `x*sin(x) + cos(x)`

3.21.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = x \sin(x) + \cos(x)$$

input `integrate(x*cos(x),x)`

output `x*sin(x) + cos(x)`

3.21.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = x \sin(x) + \cos(x)$$

input `integrate(x*cos(x),x, algorithm="maxima")`

output `x*sin(x) + cos(x)`

3.21.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = x \sin(x) + \cos(x)$$

input `integrate(x*cos(x),x, algorithm="giac")`

output `x*sin(x) + cos(x)`

3.21.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int x \cos(x) dx = \cos(x) + x \sin(x)$$

input `int(x*cos(x),x)`

output `cos(x) + x*sin(x)`

3.22 $\int x \sin(4x) dx$

3.22.1	Optimal result	236
3.22.2	Mathematica [A] (verified)	236
3.22.3	Rubi [A] (verified)	237
3.22.4	Maple [A] (verified)	238
3.22.5	Fricas [A] (verification not implemented)	238
3.22.6	Sympy [A] (verification not implemented)	239
3.22.7	Maxima [A] (verification not implemented)	239
3.22.8	Giac [A] (verification not implemented)	239
3.22.9	Mupad [B] (verification not implemented)	240

3.22.1 Optimal result

Integrand size = 6, antiderivative size = 18

$$\int x \sin(4x) dx = -\frac{1}{4}x \cos(4x) + \frac{1}{16} \sin(4x)$$

output `-1/4*x*cos(4*x)+1/16*sin(4*x)`

3.22.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int x \sin(4x) dx = -\frac{1}{4}x \cos(4x) + \frac{1}{16} \sin(4x)$$

input `Integrate[x*Sin[4*x],x]`

output `-1/4*(x*Cos[4*x]) + Sin[4*x]/16`

3.22.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \sin(4x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x \sin(4x) dx \\
 & \quad \downarrow \text{3777} \\
 & \frac{1}{4} \int \cos(4x) dx - \frac{1}{4} x \cos(4x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4} \int \sin\left(4x + \frac{\pi}{2}\right) dx - \frac{1}{4} x \cos(4x) \\
 & \quad \downarrow \text{3117} \\
 & \frac{1}{16} \sin(4x) - \frac{1}{4} x \cos(4x)
 \end{aligned}$$

input `Int[x*Sin[4*x],x]`

output `-1/4*(x*Cos[4*x]) + Sin[4*x]/16`

3.22.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

```
rule 3777 Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-
(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*C
os[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

3.22.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$-\frac{x \cos(4x)}{4} + \frac{\sin(4x)}{16}$	15
default	$-\frac{x \cos(4x)}{4} + \frac{\sin(4x)}{16}$	15
risch	$-\frac{x \cos(4x)}{4} + \frac{\sin(4x)}{16}$	15
parallelrisc	$-\frac{x \cos(4x)}{4} + \frac{\sin(4x)}{16}$	15
parts	$-\frac{x \cos(4x)}{4} + \frac{\sin(4x)}{16}$	15
meijerg	$\frac{\sqrt{\pi} \left(-\frac{2x \cos(4x)}{\sqrt{\pi}} + \frac{\sin(4x)}{2\sqrt{\pi}} \right)}{8}$	26
norman	$-\frac{x}{4} + \frac{x(\tan^2(2x))}{4} + \frac{\tan(2x)}{8}$ $1 + \tan^2(2x)$	31

```
input int(x*sin(4*x),x,method=_RETURNVERBOSE)
```

```
output -1/4*x*cos(4*x)+1/16*sin(4*x)
```

3.22.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int x \sin(4x) dx = -\frac{1}{4} x \cos(4x) + \frac{1}{16} \sin(4x)$$

```
input integrate(x*sin(4*x),x, algorithm="fracas")
```

```
output -1/4*x*cos(4*x) + 1/16*sin(4*x)
```

3.22.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int x \sin(4x) dx = -\frac{x \cos(4x)}{4} + \frac{\sin(4x)}{16}$$

input `integrate(x*sin(4*x),x)`output `-x*cos(4*x)/4 + sin(4*x)/16`**3.22.7 Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int x \sin(4x) dx = -\frac{1}{4} x \cos(4x) + \frac{1}{16} \sin(4x)$$

input `integrate(x*sin(4*x),x, algorithm="maxima")`output `-1/4*x*cos(4*x) + 1/16*sin(4*x)`**3.22.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int x \sin(4x) dx = -\frac{1}{4} x \cos(4x) + \frac{1}{16} \sin(4x)$$

input `integrate(x*sin(4*x),x, algorithm="giac")`output `-1/4*x*cos(4*x) + 1/16*sin(4*x)`

3.22.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int x \sin(4x) dx = \frac{\sin(4x)}{16} - \frac{x \cos(4x)}{4}$$

input `int(x*sin(4*x),x)`

output `sin(4*x)/16 - (x*cos(4*x))/4`

3.23 $\int x \log(x) dx$

3.23.1	Optimal result	241
3.23.2	Mathematica [A] (verified)	241
3.23.3	Rubi [A] (verified)	242
3.23.4	Maple [A] (verified)	242
3.23.5	Fricas [A] (verification not implemented)	243
3.23.6	Sympy [A] (verification not implemented)	243
3.23.7	Maxima [A] (verification not implemented)	243
3.23.8	Giac [A] (verification not implemented)	244
3.23.9	Mupad [B] (verification not implemented)	244

3.23.1 Optimal result

Integrand size = 4, antiderivative size = 17

$$\int x \log(x) dx = -\frac{x^2}{4} + \frac{1}{2}x^2 \log(x)$$

output `-1/4*x^2+1/2*x^2*ln(x)`

3.23.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int x \log(x) dx = -\frac{x^2}{4} + \frac{1}{2}x^2 \log(x)$$

input `Integrate[x*Log[x],x]`

output `-1/4*x^2 + (x^2*Log[x])/2`

3.23.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \log(x) dx$$

$$\downarrow 2741$$

$$\frac{1}{2}x^2 \log(x) - \frac{x^2}{4}$$

input `Int[x*Log[x],x]`

output `-1/4*x^2 + (x^2*Log[x])/2`

3.23.3.1 Defintions of rubi rules used

rule 2741 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_.), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(m + 1))/(d*(m + 1)^2), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]`

3.23.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
norman	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
risch	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
parallelrisch	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14
parts	$-\frac{x^2}{4} + \frac{x^2 \ln(x)}{2}$	14

input `int(x*ln(x),x,method=_RETURNVERBOSE)`

output `-1/4*x^2+1/2*x^2*ln(x)`

3.23.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int x \log(x) dx = \frac{1}{2} x^2 \log(x) - \frac{1}{4} x^2$$

input `integrate(x*log(x),x, algorithm="fricas")`

output `1/2*x^2*log(x) - 1/4*x^2`

3.23.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int x \log(x) dx = \frac{x^2 \log(x)}{2} - \frac{x^2}{4}$$

input `integrate(x*ln(x),x)`

output `x**2*log(x)/2 - x**2/4`

3.23.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int x \log(x) dx = \frac{1}{2} x^2 \log(x) - \frac{1}{4} x^2$$

input `integrate(x*log(x),x, algorithm="maxima")`

output `1/2*x^2*log(x) - 1/4*x^2`

3.23.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int x \log(x) dx = \frac{1}{2} x^2 \log(x) - \frac{1}{4} x^2$$

input `integrate(x*log(x),x, algorithm="giac")`

output `1/2*x^2*log(x) - 1/4*x^2`

3.23.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.53

$$\int x \log(x) dx = \frac{x^2 (\ln(x) - \frac{1}{2})}{2}$$

input `int(x*log(x),x)`

output `(x^2*(log(x) - 1/2))/2`

3.24 $\int x^2 \cos(3x) dx$

3.24.1	Optimal result	245
3.24.2	Mathematica [A] (verified)	245
3.24.3	Rubi [A] (verified)	246
3.24.4	Maple [A] (verified)	247
3.24.5	Fricas [A] (verification not implemented)	248
3.24.6	Sympy [A] (verification not implemented)	248
3.24.7	Maxima [A] (verification not implemented)	248
3.24.8	Giac [A] (verification not implemented)	249
3.24.9	Mupad [B] (verification not implemented)	249

3.24.1 Optimal result

Integrand size = 8, antiderivative size = 29

$$\int x^2 \cos(3x) dx = \frac{2}{9}x \cos(3x) - \frac{2}{27} \sin(3x) + \frac{1}{3}x^2 \sin(3x)$$

output `2/9*x*cos(3*x)-2/27*sin(3*x)+1/3*x^2*sin(3*x)`

3.24.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.86

$$\int x^2 \cos(3x) dx = \frac{2}{9}x \cos(3x) + \frac{1}{27}(-2 + 9x^2) \sin(3x)$$

input `Integrate[x^2*Cos[3*x],x]`

output `(2*x*Cos[3*x])/9 + ((-2 + 9*x^2)*Sin[3*x])/27`

3.24.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.17, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {3042, 3777, 25, 3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \cos(3x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x^2 \sin\left(3x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3777} \\
 & \frac{2}{3} \int -x \sin(3x) dx + \frac{1}{3} x^2 \sin(3x) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{3} x^2 \sin(3x) - \frac{2}{3} \int x \sin(3x) dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} x^2 \sin(3x) - \frac{2}{3} \int x \sin(3x) dx \\
 & \quad \downarrow \text{3777} \\
 & \frac{1}{3} x^2 \sin(3x) - \frac{2}{3} \left(\frac{1}{3} \int \cos(3x) dx - \frac{1}{3} x \cos(3x) \right) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} x^2 \sin(3x) - \frac{2}{3} \left(\frac{1}{3} \int \sin\left(3x + \frac{\pi}{2}\right) dx - \frac{1}{3} x \cos(3x) \right) \\
 & \quad \downarrow \text{3117} \\
 & \frac{1}{3} x^2 \sin(3x) - \frac{2}{3} \left(\frac{1}{9} \sin(3x) - \frac{1}{3} x \cos(3x) \right)
 \end{aligned}$$

input `Int [x^2*Cos [3*x] , x]`

output `(-2*(-1/3*(x*Cos [3*x]) + Sin [3*x]/9))/3 + (x^2*Sin [3*x])/3`

3.24.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

3.24.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

method	result	size
risch	$\frac{2x \cos(3x)}{9} + \frac{(9x^2 - 2) \sin(3x)}{27}$	22
derivativedivides	$\frac{2x \cos(3x)}{9} - \frac{2 \sin(3x)}{27} + \frac{x^2 \sin(3x)}{3}$	24
default	$\frac{2x \cos(3x)}{9} - \frac{2 \sin(3x)}{27} + \frac{x^2 \sin(3x)}{3}$	24
parts	$\frac{2x \cos(3x)}{9} - \frac{2 \sin(3x)}{27} + \frac{x^2 \sin(3x)}{3}$	24
meijerg	$\frac{4\sqrt{\pi} \left(\frac{3x \cos(3x)}{2\sqrt{\pi}} - \frac{(-27x^2 + 3) \sin(3x)}{6\sqrt{\pi}} \right)}{27}$	33
norman	$\frac{\frac{2x}{9} - \frac{2x \tan^2\left(\frac{3x}{2}\right)}{9} + \frac{2x^2 \tan\left(\frac{3x}{2}\right)}{3} - \frac{4 \tan\left(\frac{3x}{2}\right)}{27}}{1 + \tan^2\left(\frac{3x}{2}\right)}$	40
parallelrisc	$\frac{18x^2 \tan\left(\frac{3x}{2}\right) - 6x \tan^2\left(\frac{3x}{2}\right) + 6x - 4 \tan\left(\frac{3x}{2}\right)}{27 \tan^2\left(\frac{3x}{2}\right) + 27}$	42

input `int(x^2*cos(3*x), x, method=_RETURNVERBOSE)`

output `2/9*x*cos(3*x)+1/27*(9*x^2-2)*sin(3*x)`

3.24.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int x^2 \cos(3x) dx = \frac{2}{9} x \cos(3x) + \frac{1}{27} (9x^2 - 2) \sin(3x)$$

input `integrate(x^2*cos(3*x),x, algorithm="fricas")`

output `2/9*x*cos(3*x) + 1/27*(9*x^2 - 2)*sin(3*x)`

3.24.6 Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int x^2 \cos(3x) dx = \frac{x^2 \sin(3x)}{3} + \frac{2x \cos(3x)}{9} - \frac{2 \sin(3x)}{27}$$

input `integrate(x**2*cos(3*x),x)`

output `x**2*sin(3*x)/3 + 2*x*cos(3*x)/9 - 2*sin(3*x)/27`

3.24.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int x^2 \cos(3x) dx = \frac{2}{9} x \cos(3x) + \frac{1}{27} (9x^2 - 2) \sin(3x)$$

input `integrate(x^2*cos(3*x),x, algorithm="maxima")`

output `2/9*x*cos(3*x) + 1/27*(9*x^2 - 2)*sin(3*x)`

3.24.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int x^2 \cos(3x) dx = \frac{2}{9} x \cos(3x) + \frac{1}{27} (9x^2 - 2) \sin(3x)$$

input `integrate(x^2*cos(3*x),x, algorithm="giac")`

output `2/9*x*cos(3*x) + 1/27*(9*x^2 - 2)*sin(3*x)`

3.24.9 Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

$$\int x^2 \cos(3x) dx = \frac{2x \cos(3x)}{9} - \frac{2 \sin(3x)}{27} + \frac{x^2 \sin(3x)}{3}$$

input `int(x^2*cos(3*x),x)`

output `(2*x*cos(3*x))/9 - (2*sin(3*x))/27 + (x^2*sin(3*x))/3`

3.25 $\int x^2 \sin(2x) dx$

3.25.1	Optimal result	250
3.25.2	Mathematica [A] (verified)	250
3.25.3	Rubi [A] (verified)	251
3.25.4	Maple [A] (verified)	252
3.25.5	Fricas [A] (verification not implemented)	253
3.25.6	Sympy [A] (verification not implemented)	253
3.25.7	Maxima [A] (verification not implemented)	253
3.25.8	Giac [A] (verification not implemented)	254
3.25.9	Mupad [B] (verification not implemented)	254

3.25.1 Optimal result

Integrand size = 8, antiderivative size = 29

$$\int x^2 \sin(2x) dx = \frac{1}{4} \cos(2x) - \frac{1}{2} x^2 \cos(2x) + \frac{1}{2} x \sin(2x)$$

output `1/4*cos(2*x)-1/2*x^2*cos(2*x)+1/2*x*sin(2*x)`

3.25.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.86

$$\int x^2 \sin(2x) dx = -\frac{1}{4}(-1 + 2x^2) \cos(2x) + \frac{1}{2} x \sin(2x)$$

input `Integrate[x^2*Sin[2*x],x]`

output `-1/4*((-1 + 2*x^2)*Cos[2*x]) + (x*Sin[2*x])/2`

3.25.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {3042, 3777, 3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \sin(2x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x^2 \sin(2x) dx \\
 & \quad \downarrow \text{3777} \\
 & \int x \cos(2x) dx - \frac{1}{2} x^2 \cos(2x) \\
 & \quad \downarrow \text{3042} \\
 & \int x \sin\left(2x + \frac{\pi}{2}\right) dx - \frac{1}{2} x^2 \cos(2x) \\
 & \quad \downarrow \text{3777} \\
 & \frac{1}{2} \int -\sin(2x) dx - \frac{1}{2} x^2 \cos(2x) + \frac{1}{2} x \sin(2x) \\
 & \quad \downarrow \text{25} \\
 & -\frac{1}{2} \int \sin(2x) dx - \frac{1}{2} x^2 \cos(2x) + \frac{1}{2} x \sin(2x) \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{2} \int \sin(2x) dx - \frac{1}{2} x^2 \cos(2x) + \frac{1}{2} x \sin(2x) \\
 & \quad \downarrow \text{3118} \\
 & -\frac{1}{2} x^2 \cos(2x) + \frac{1}{2} x \sin(2x) + \frac{1}{4} \cos(2x)
 \end{aligned}$$

input `Int[x^2*Sin[2*x],x]`

output `Cos[2*x]/4 - (x^2*Cos[2*x])/2 + (x*Sin[2*x])/2`

3.25.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

3.25.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

method	result	size
risch	$\left(-\frac{x^2}{2} + \frac{1}{4}\right) \cos(2x) + \frac{x \sin(2x)}{2}$	21
derivativedivides	$\frac{\cos(2x)}{4} - \frac{x^2 \cos(2x)}{2} + \frac{x \sin(2x)}{2}$	24
default	$\frac{\cos(2x)}{4} - \frac{x^2 \cos(2x)}{2} + \frac{x \sin(2x)}{2}$	24
parts	$\frac{\cos(2x)}{4} - \frac{x^2 \cos(2x)}{2} + \frac{x \sin(2x)}{2}$	24
parallelrisch	$\frac{1}{4} - \frac{x^2 \cos(2x)}{2} + \frac{x \sin(2x)}{2} + \frac{\cos(2x)}{4}$	25
norman	$\frac{x \tan(x) - \frac{x^2}{2} + \frac{x^2 \tan^2(x)}{2} + \frac{1}{2}}{1 + \tan^2(x)}$	30
meijerg	$\frac{\sqrt{\pi} \left(-\frac{1}{2\sqrt{\pi}} + \frac{(-2x^2+1) \cos(2x)}{2\sqrt{\pi}} + \frac{x \sin(2x)}{\sqrt{\pi}} \right)}{2}$	37

input `int(x^2*sin(2*x), x, method=_RETURNVERBOSE)`

output `(-1/2*x^2+1/4)*cos(2*x)+1/2*x*sin(2*x)`

3.25.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int x^2 \sin(2x) dx = -\frac{1}{4} (2x^2 - 1) \cos(2x) + \frac{1}{2} x \sin(2x)$$

input `integrate(x^2*sin(2*x),x, algorithm="fricas")`output `-1/4*(2*x^2 - 1)*cos(2*x) + 1/2*x*sin(2*x)`**3.25.6 Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

$$\int x^2 \sin(2x) dx = -\frac{x^2 \cos(2x)}{2} + \frac{x \sin(2x)}{2} + \frac{\cos(2x)}{4}$$

input `integrate(x**2*sin(2*x),x)`output `-x**2*cos(2*x)/2 + x*sin(2*x)/2 + cos(2*x)/4`**3.25.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int x^2 \sin(2x) dx = -\frac{1}{4} (2x^2 - 1) \cos(2x) + \frac{1}{2} x \sin(2x)$$

input `integrate(x^2*sin(2*x),x, algorithm="maxima")`output `-1/4*(2*x^2 - 1)*cos(2*x) + 1/2*x*sin(2*x)`

3.25.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int x^2 \sin(2x) dx = -\frac{1}{4} (2x^2 - 1) \cos(2x) + \frac{1}{2} x \sin(2x)$$

input `integrate(x^2*sin(2*x),x, algorithm="giac")`output `-1/4*(2*x^2 - 1)*cos(2*x) + 1/2*x*sin(2*x)`**3.25.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

$$\int x^2 \sin(2x) dx = \frac{x \sin(2x)}{2} + (2 \sin(x)^2 - 1) \left(\frac{x^2}{2} - \frac{1}{4} \right)$$

input `int(x^2*sin(2*x),x)`output `(x*sin(2*x))/2 + (2*sin(x)^2 - 1)*(x^2/2 - 1/4)`

3.26 $\int \log^2(x) dx$

3.26.1	Optimal result	255
3.26.2	Mathematica [A] (verified)	255
3.26.3	Rubi [A] (verified)	256
3.26.4	Maple [A] (verified)	257
3.26.5	Fricas [A] (verification not implemented)	257
3.26.6	Sympy [A] (verification not implemented)	257
3.26.7	Maxima [A] (verification not implemented)	258
3.26.8	Giac [A] (verification not implemented)	258
3.26.9	Mupad [B] (verification not implemented)	258

3.26.1 Optimal result

Integrand size = 4, antiderivative size = 15

$$\int \log^2(x) dx = 2x - 2x \log(x) + x \log^2(x)$$

output `2*x-2*x*ln(x)+x*ln(x)^2`

3.26.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \log^2(x) dx = 2x - 2x \log(x) + x \log^2(x)$$

input `Integrate[Log[x]^2,x]`

output `2*x - 2*x*Log[x] + x*Log[x]^2`

3.26.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {2733, 2732}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \log^2(x) dx \\ \downarrow \text{2733} \\ x \log^2(x) - 2 \int \log(x) dx \\ \downarrow \text{2732} \\ x \log^2(x) - 2(x \log(x) - x) \end{array}$$

input `Int [Log[x]^2, x]`

output `x*Log[x]^2 - 2*(-x + x*Log[x])`

3.26.3.1 Defintions of rubi rules used

rule 2732 `Int [Log[(c_.)*(x_)^(n_.)], x_Symbol] := Simp[x*Log[c*x^n], x] - Simp[n*x, x] /; FreeQ[{c, n}, x]`

rule 2733 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.), x_Symbol] := Simp[x*(a + b *Log[c*x^n])^p, x] - Simp[b*n*p Int[(a + b*Log[c*x^n])^(p - 1), x], x] /; FreeQ[{a, b, c, n}, x] && GtQ[p, 0] && IntegerQ[2*p]`

3.26.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

method	result	size
default	$2x - 2x \ln(x) + x \ln(x)^2$	16
norman	$2x - 2x \ln(x) + x \ln(x)^2$	16
risch	$2x - 2x \ln(x) + x \ln(x)^2$	16
parallelrisc	$2x - 2x \ln(x) + x \ln(x)^2$	16

input `int(ln(x)^2,x,method=_RETURNVERBOSE)`output `2*x-2*x*ln(x)+x*ln(x)^2`**3.26.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \log^2(x) dx = x \log(x)^2 - 2x \log(x) + 2x$$

input `integrate(log(x)^2,x, algorithm="fricas")`output `x*log(x)^2 - 2*x*log(x) + 2*x`**3.26.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \log^2(x) dx = x \log(x)^2 - 2x \log(x) + 2x$$

input `integrate(ln(x)**2,x)`output `x*log(x)**2 - 2*x*log(x) + 2*x`

3.26.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \log^2(x) dx = (\log(x)^2 - 2 \log(x) + 2)x$$

input `integrate(log(x)^2,x, algorithm="maxima")`output `(log(x)^2 - 2*log(x) + 2)*x`**3.26.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \log^2(x) dx = x \log(x)^2 - 2x \log(x) + 2x$$

input `integrate(log(x)^2,x, algorithm="giac")`output `x*log(x)^2 - 2*x*log(x) + 2*x`**3.26.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \log^2(x) dx = x (\ln(x)^2 - 2 \ln(x) + 2)$$

input `int(log(x)^2,x)`output `x*(log(x)^2 - 2*log(x) + 2)`

3.27 $\int \arcsin(x) dx$

3.27.1	Optimal result	259
3.27.2	Mathematica [A] (verified)	259
3.27.3	Rubi [A] (verified)	260
3.27.4	Maple [A] (verified)	261
3.27.5	Fricas [A] (verification not implemented)	261
3.27.6	Sympy [A] (verification not implemented)	261
3.27.7	Maxima [A] (verification not implemented)	262
3.27.8	Giac [A] (verification not implemented)	262
3.27.9	Mupad [B] (verification not implemented)	262

3.27.1 Optimal result

Integrand size = 2, antiderivative size = 16

$$\int \arcsin(x) dx = \sqrt{1-x^2} + x \arcsin(x)$$

output `x*arcsin(x)+(-x^2+1)^(1/2)`

3.27.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \arcsin(x) dx = \sqrt{1-x^2} + x \arcsin(x)$$

input `Integrate[ArcSin[x],x]`

output `Sqrt[1 - x^2] + x*ArcSin[x]`

3.27.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {5130, 241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arcsin(x) dx$$

$$\downarrow \text{5130}$$

$$x \arcsin(x) - \int \frac{x}{\sqrt{1-x^2}} dx$$

$$\downarrow \text{241}$$

$$x \arcsin(x) + \sqrt{1-x^2}$$

input `Int[ArcSin[x], x]`

output `Sqrt[1 - x^2] + x*ArcSin[x]`

3.27.3.1 Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

rule 5130 `Int[((a_.) + ArcSin[(c_.)*(x_)])*(b_.))^(n_.), x_Symbol] := Simp[x*(a + b*ArcSin[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcSin[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

3.27.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result	size
lookup	$\arcsin(x) x + \sqrt{-x^2 + 1}$	15
default	$\arcsin(x) x + \sqrt{-x^2 + 1}$	15
parts	$\arcsin(x) x + \sqrt{-x^2 + 1}$	15

input `int(arcsin(x),x,method=_RETURNVERBOSE)`

output `arcsin(x)*x+(-x^2+1)^(1/2)`

3.27.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \arcsin(x) dx = x \arcsin(x) + \sqrt{-x^2 + 1}$$

input `integrate(arcsin(x),x, algorithm="fricas")`

output `x*arcsin(x) + sqrt(-x^2 + 1)`

3.27.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \arcsin(x) dx = x \operatorname{asin}(x) + \sqrt{1 - x^2}$$

input `integrate(asin(x),x)`

output `x*asin(x) + sqrt(1 - x**2)`

3.27.7 Maxima [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \arcsin(x) dx = x \arcsin(x) + \sqrt{-x^2 + 1}$$

input `integrate(arcsin(x),x, algorithm="maxima")`output `x*arcsin(x) + sqrt(-x^2 + 1)`**3.27.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \arcsin(x) dx = x \arcsin(x) + \sqrt{-x^2 + 1}$$

input `integrate(arcsin(x),x, algorithm="giac")`output `x*arcsin(x) + sqrt(-x^2 + 1)`**3.27.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \arcsin(x) dx = x \arcsin(x) + \sqrt{1 - x^2}$$

input `int(asin(x),x)`output `x*asin(x) + (1 - x^2)^(1/2)`

3.28 $\int t \cos(t) \sin(t) dt$

3.28.1	Optimal result	263
3.28.2	Mathematica [A] (verified)	263
3.28.3	Rubi [A] (verified)	264
3.28.4	Maple [A] (verified)	265
3.28.5	Fricas [A] (verification not implemented)	265
3.28.6	Sympy [A] (verification not implemented)	266
3.28.7	Maxima [A] (verification not implemented)	266
3.28.8	Giac [A] (verification not implemented)	266
3.28.9	Mupad [B] (verification not implemented)	267

3.28.1 Optimal result

Integrand size = 6, antiderivative size = 23

$$\int t \cos(t) \sin(t) dt = -\frac{t}{4} + \frac{1}{4} \cos(t) \sin(t) + \frac{1}{2} t \sin^2(t)$$

output `-1/4*t+1/4*cos(t)*sin(t)+1/2*t*sin(t)^2`

3.28.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int t \cos(t) \sin(t) dt = -\frac{1}{4} t \cos(2t) + \frac{1}{8} \sin(2t)$$

input `Integrate[t*Cos[t]*Sin[t],t]`

output `-1/4*(t*Cos[2*t]) + Sin[2*t]/8`

3.28.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.22, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3924, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int t \sin(t) \cos(t) dt \\
 & \quad \downarrow \text{3924} \\
 & \frac{1}{2} t \sin^2(t) - \frac{1}{2} \int \sin^2(t) dt \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} t \sin^2(t) - \frac{1}{2} \int \sin(t)^2 dt \\
 & \quad \downarrow \text{3115} \\
 & \frac{1}{2} \left(\frac{1}{2} \sin(t) \cos(t) - \frac{\int 1 dt}{2} \right) + \frac{1}{2} t \sin^2(t) \\
 & \quad \downarrow \text{24} \\
 & \frac{1}{2} t \sin^2(t) + \frac{1}{2} \left(\frac{1}{2} \sin(t) \cos(t) - \frac{t}{2} \right)
 \end{aligned}$$

input `Int[t*Cos[t]*Sin[t],t]`

output `(t*Sin[t]^2)/2 + (-1/2*t + (Cos[t]*Sin[t])/2)/2`

3.28.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3924 `Int[Cos[(a_.) + (b_.)*(x_)^(n_.)]*(x_)^(m_.)*Sin[(a_.) + (b_.)*(x_)^(n_.)]^(p_.), x_Symbol] := Simp[x^(m - n + 1)*(Sin[a + b*x^n]^(p + 1)/(b*n*(p + 1))), x] - Simp[(m - n + 1)/(b*n*(p + 1)) Int[x^(m - n)*Sin[a + b*x^n]^(p + 1), x], x] /; FreeQ[{a, b, p}, x] && LtQ[0, n, m + 1] && NeQ[p, -1]`

3.28.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

method	result	size
risch	$-\frac{t \cos(2t)}{4} + \frac{\sin(2t)}{8}$	15
parallelrisc	$-\frac{t \cos(2t)}{4} + \frac{\sin(2t)}{8}$	15
default	$-\frac{t(\cos^2(t))}{2} + \frac{\cos(t)\sin(t)}{4} + \frac{t}{4}$	18
meijerg	$\frac{\sqrt{\pi} \left(-\frac{t \cos(2t)}{\sqrt{\pi}} + \frac{\sin(2t)}{2\sqrt{\pi}} \right)}{4}$	26
norman	$\frac{-\frac{t}{4} - \frac{(\tan^3(\frac{t}{2}))}{2} + \frac{3t(\tan^2(\frac{t}{2}))}{2} - \frac{t(\tan^4(\frac{t}{2}))}{4} + \frac{\tan(\frac{t}{2})}{2}}{(1+\tan^2(\frac{t}{2}))^2}$	48

input `int(t*cos(t)*sin(t),t,method=_RETURNVERBOSE)`

output `-1/4*t*cos(2*t)+1/8*sin(2*t)`

3.28.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int t \cos(t) \sin(t) dt = -\frac{1}{2} t \cos(t)^2 + \frac{1}{4} \cos(t) \sin(t) + \frac{1}{4} t$$

input `integrate(t*cos(t)*sin(t),t, algorithm="fricas")`

output `-1/2*t*cos(t)^2 + 1/4*cos(t)*sin(t) + 1/4*t`

3.28.6 Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.04

$$\int t \cos(t) \sin(t) dt = \frac{t \sin^2(t)}{4} - \frac{t \cos^2(t)}{4} + \frac{\sin(t) \cos(t)}{4}$$

input `integrate(t*cos(t)*sin(t),t)`

output `t*sin(t)**2/4 - t*cos(t)**2/4 + sin(t)*cos(t)/4`

3.28.7 Maxima [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.61

$$\int t \cos(t) \sin(t) dt = -\frac{1}{4} t \cos(2t) + \frac{1}{8} \sin(2t)$$

input `integrate(t*cos(t)*sin(t),t, algorithm="maxima")`

output `-1/4*t*cos(2*t) + 1/8*sin(2*t)`

3.28.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.61

$$\int t \cos(t) \sin(t) dt = -\frac{1}{4} t \cos(2t) + \frac{1}{8} \sin(2t)$$

input `integrate(t*cos(t)*sin(t),t, algorithm="giac")`

output `-1/4*t*cos(2*t) + 1/8*sin(2*t)`

3.28.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int t \cos(t) \sin(t) dt = \frac{\sin(2t)}{8} + \frac{t(2\sin(t)^2 - 1)}{4}$$

input `int(t*cos(t)*sin(t),t)`

output `sin(2*t)/8 + (t*(2*sin(t)^2 - 1))/4`

3.29 $\int t \sec^2(t) dt$

3.29.1	Optimal result	268
3.29.2	Mathematica [A] (verified)	268
3.29.3	Rubi [A] (verified)	269
3.29.4	Maple [A] (verified)	270
3.29.5	Fricas [B] (verification not implemented)	271
3.29.6	Sympy [A] (verification not implemented)	271
3.29.7	Maxima [B] (verification not implemented)	271
3.29.8	Giac [B] (verification not implemented)	272
3.29.9	Mupad [B] (verification not implemented)	272

3.29.1 Optimal result

Integrand size = 6, antiderivative size = 8

$$\int t \sec^2(t) dt = \log(\cos(t)) + t \tan(t)$$

output `ln(cos(t))+t*tan(t)`

3.29.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int t \sec^2(t) dt = \log(\cos(t)) + t \tan(t)$$

input `Integrate[t*Sec[t]^2,t]`

output `Log[Cos[t]] + t*Tan[t]`

3.29.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {3042, 4672, 25, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int t \sec^2(t) dt \\
 & \quad \downarrow \text{3042} \\
 & \int t \csc\left(t + \frac{\pi}{2}\right)^2 dt \\
 & \quad \downarrow \text{4672} \\
 & \int -\tan(t) dt + t \tan(t) \\
 & \quad \downarrow \text{25} \\
 & t \tan(t) - \int \tan(t) dt \\
 & \quad \downarrow \text{3042} \\
 & t \tan(t) - \int \tan(t) dt \\
 & \quad \downarrow \text{3956} \\
 & t \tan(t) + \log(\cos(t))
 \end{aligned}$$

input `Int [t*Sec[t]^2, t]`

output `Log[Cos[t]] + t*Tan[t]`

3.29.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4672 `Int[csc[(e_.) + (f_.)*(x_)^2*((c_.) + (d_.)*(x_)^(m_.)), x_Symbol] := Simp[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

3.29.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
default	$\ln(\cos(t)) + t \tan(t)$	9
risch	$-2it + \frac{2it}{e^{2it}+1} + \ln(e^{2it} + 1)$	27
parallelrisc	$-\ln\left(\frac{2}{\cos(t)+1}\right) + \ln(\csc(t) - \cot(t) - 1) + \ln(\csc(t) - \cot(t) + 1) + t \tan(t)$	35
norman	$-\frac{2 \tan(\frac{t}{2})t}{\tan^2(\frac{t}{2})-1} - \ln(1 + \tan^2(\frac{t}{2})) + \ln(\tan(\frac{t}{2}) - 1) + \ln(\tan(\frac{t}{2}) + 1)$	44

input `int(t*sec(t)^2,t,method=_RETURNVERBOSE)`

output `ln(cos(t))+t*tan(t)`

3.29.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18 vs. $2(8) = 16$.

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 2.25

$$\int t \sec^2(t) dt = \frac{\cos(t) \log(-\cos(t)) + t \sin(t)}{\cos(t)}$$

input `integrate(t*sec(t)^2,t, algorithm="fricas")`

output `(cos(t)*log(-cos(t)) + t*sin(t))/cos(t)`

3.29.6 Sympy [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int t \sec^2(t) dt = t \tan(t) + \log(\cos(t))$$

input `integrate(t*sec(t)**2,t)`

output `t*tan(t) + log(cos(t))`

3.29.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 74 vs. $2(8) = 16$.

Time = 0.26 (sec) , antiderivative size = 74, normalized size of antiderivative = 9.25

$$\int t \sec^2(t) dt = \frac{(\cos(2t)^2 + \sin(2t)^2 + 2 \cos(2t) + 1) \log(\cos(2t)^2 + \sin(2t)^2 + 2 \cos(2t) + 1) + 4t \sin(2t)}{2(\cos(2t)^2 + \sin(2t)^2 + 2 \cos(2t) + 1)}$$

input `integrate(t*sec(t)^2,t, algorithm="maxima")`

output `1/2*((cos(2*t)^2 + sin(2*t)^2 + 2*cos(2*t) + 1)*log(cos(2*t)^2 + sin(2*t)^2 + 2*cos(2*t) + 1) + 4*t*sin(2*t))/(cos(2*t)^2 + sin(2*t)^2 + 2*cos(2*t) + 1)`

3.29.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 103 vs. $2(8) = 16$.

Time = 0.29 (sec) , antiderivative size = 103, normalized size of antiderivative = 12.88

$$\int t \sec^2(t) dt$$

$$= \frac{\log\left(\frac{4\left(\tan\left(\frac{1}{2}t\right)^4 - 2\tan\left(\frac{1}{2}t\right)^2 + 1\right)}{\tan\left(\frac{1}{2}t\right)^4 + 2\tan\left(\frac{1}{2}t\right)^2 + 1}\right) \tan\left(\frac{1}{2}t\right)^2 - 4t \tan\left(\frac{1}{2}t\right) - \log\left(\frac{4\left(\tan\left(\frac{1}{2}t\right)^4 - 2\tan\left(\frac{1}{2}t\right)^2 + 1\right)}{\tan\left(\frac{1}{2}t\right)^4 + 2\tan\left(\frac{1}{2}t\right)^2 + 1}\right)}{2\left(\tan\left(\frac{1}{2}t\right)^2 - 1\right)}$$

input `integrate(t*sec(t)^2,t, algorithm="giac")`

output `1/2*(log(4*(tan(1/2*t)^4 - 2*tan(1/2*t)^2 + 1)/(tan(1/2*t)^4 + 2*tan(1/2*t)^2 + 1))*tan(1/2*t)^2 - 4*t*tan(1/2*t) - log(4*(tan(1/2*t)^4 - 2*tan(1/2*t)^2 + 1)/(tan(1/2*t)^4 + 2*tan(1/2*t)^2 + 1)))/(tan(1/2*t)^2 - 1)`

3.29.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int t \sec^2(t) dt = \ln(\cos(t)) + t \tan(t)$$

input `int(t/cos(t)^2,t)`

output `log(cos(t)) + t*tan(t)`

3.30 $\int t^2 \log(t) dt$

3.30.1	Optimal result	273
3.30.2	Mathematica [A] (verified)	273
3.30.3	Rubi [A] (verified)	274
3.30.4	Maple [A] (verified)	274
3.30.5	Fricas [A] (verification not implemented)	275
3.30.6	Sympy [A] (verification not implemented)	275
3.30.7	Maxima [A] (verification not implemented)	275
3.30.8	Giac [A] (verification not implemented)	276
3.30.9	Mupad [B] (verification not implemented)	276

3.30.1 Optimal result

Integrand size = 6, antiderivative size = 17

$$\int t^2 \log(t) dt = -\frac{t^3}{9} + \frac{1}{3}t^3 \log(t)$$

output `-1/9*t^3+1/3*t^3*ln(t)`

3.30.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int t^2 \log(t) dt = -\frac{t^3}{9} + \frac{1}{3}t^3 \log(t)$$

input `Integrate[t^2*Log[t],t]`

output `-1/9*t^3 + (t^3*Log[t])/3`

3.30.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int t^2 \log(t) dt$$

↓ 2741

$$\frac{1}{3}t^3 \log(t) - \frac{t^3}{9}$$

input `Int[t^2*Log[t],t]`

output `-1/9*t^3 + (t^3*Log[t])/3`

3.30.3.1 Defintions of rubi rules used

rule 2741 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_.), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(m + 1))/(d*(m + 1)^2), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]`

3.30.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{t^3}{9} + \frac{t^3 \ln(t)}{3}$	14
norman	$-\frac{t^3}{9} + \frac{t^3 \ln(t)}{3}$	14
risch	$-\frac{t^3}{9} + \frac{t^3 \ln(t)}{3}$	14
parallelrisch	$-\frac{t^3}{9} + \frac{t^3 \ln(t)}{3}$	14
parts	$-\frac{t^3}{9} + \frac{t^3 \ln(t)}{3}$	14

input `int(t^2*ln(t),t,method=_RETURNVERBOSE)`

output `-1/9*t^3+1/3*t^3*ln(t)`

3.30.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int t^2 \log(t) dt = \frac{1}{3} t^3 \log(t) - \frac{1}{9} t^3$$

input `integrate(t^2*log(t),t, algorithm="fricas")`

output `1/3*t^3*log(t) - 1/9*t^3`

3.30.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int t^2 \log(t) dt = \frac{t^3 \log(t)}{3} - \frac{t^3}{9}$$

input `integrate(t**2*ln(t),t)`

output `t**3*log(t)/3 - t**3/9`

3.30.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int t^2 \log(t) dt = \frac{1}{3} t^3 \log(t) - \frac{1}{9} t^3$$

input `integrate(t^2*log(t),t, algorithm="maxima")`

output `1/3*t^3*log(t) - 1/9*t^3`

3.30.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int t^2 \log(t) dt = \frac{1}{3} t^3 \log(t) - \frac{1}{9} t^3$$

input `integrate(t^2*log(t),t, algorithm="giac")`

output `1/3*t^3*log(t) - 1/9*t^3`

3.30.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.53

$$\int t^2 \log(t) dt = \frac{t^3 (\ln(t) - \frac{1}{3})}{3}$$

input `int(t^2*log(t),t)`

output `(t^3*(log(t) - 1/3))/3`

3.31 $\int e^{tt^3} dt$

3.31.1	Optimal result	277
3.31.2	Mathematica [A] (verified)	277
3.31.3	Rubi [A] (verified)	278
3.31.4	Maple [A] (verified)	279
3.31.5	Fricas [A] (verification not implemented)	279
3.31.6	Sympy [A] (verification not implemented)	280
3.31.7	Maxima [A] (verification not implemented)	280
3.31.8	Giac [A] (verification not implemented)	280
3.31.9	Mupad [B] (verification not implemented)	281

3.31.1 Optimal result

Integrand size = 7, antiderivative size = 27

$$\int e^{tt^3} dt = -6e^t + 6e^{tt} - 3e^{tt^2} + e^{tt^3}$$

output `-6*exp(t)+6*exp(t)*t-3*exp(t)*t^2+exp(t)*t^3`

3.31.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.63

$$\int e^{tt^3} dt = e^t(-6 + 6t - 3t^2 + t^3)$$

input `Integrate[E^t*t^3,t]`

output `E^t*(-6 + 6*t - 3*t^2 + t^3)`

3.31.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.15, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {2607, 2607, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{t^3} dt \\
 & \quad \downarrow \text{2607} \\
 & e^{t^3} - 3 \int e^{t^2} dt \\
 & \quad \downarrow \text{2607} \\
 & e^{t^3} - 3 \left(e^{t^2} - 2 \int e^t dt \right) \\
 & \quad \downarrow \text{2607} \\
 & e^{t^3} - 3 \left(e^{t^2} - 2 \left(e^t - \int e^t dt \right) \right) \\
 & \quad \downarrow \text{2624} \\
 & e^{t^3} - 3(e^{t^2} - 2(e^t - e^t))
 \end{aligned}$$

input `Int[E^t*t^3,t]`

output `E^t*t^3 - 3*(E^t*t^2 - 2*(-E^t + E^t*t))`

3.31.3.1 Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

```
rule 2624 Int[((F_)^(v_))^(n_), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;
FreeQ[{F, n}, x] && LinearQ[v, x]
```

3.31.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.63

method	result	size
gospers	$(t^3 - 3t^2 + 6t - 6)e^t$	17
risch	$(t^3 - 3t^2 + 6t - 6)e^t$	17
meijerg	$6 - \frac{(-4t^3 + 12t^2 - 24t + 24)e^t}{4}$	22
default	$-6e^t + 6e^t t - 3e^t t^2 + e^t t^3$	24
norman	$-6e^t + 6e^t t - 3e^t t^2 + e^t t^3$	24
parallelrisch	$-6e^t + 6e^t t - 3e^t t^2 + e^t t^3$	24
parts	$-6e^t + 6e^t t - 3e^t t^2 + e^t t^3$	24

```
input int(exp(t)*t^3,t,method=_RETURNVERBOSE)
```

```
output (t^3-3*t^2+6*t-6)*exp(t)
```

3.31.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.59

$$\int e^t t^3 dt = (t^3 - 3t^2 + 6t - 6)e^t$$

```
input integrate(exp(t)*t^3,t, algorithm="fricas")
```

```
output (t^3 - 3*t^2 + 6*t - 6)*e^t
```


3.31.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.56

$$\int e^t t^3 dt = (t^3 - 3t^2 + 6t - 6) e^t$$

input `integrate(exp(t)*t**3,t)`output `(t**3 - 3*t**2 + 6*t - 6)*exp(t)`**3.31.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.59

$$\int e^t t^3 dt = (t^3 - 3t^2 + 6t - 6) e^t$$

input `integrate(exp(t)*t^3,t, algorithm="maxima")`output `(t^3 - 3*t^2 + 6*t - 6)*e^t`**3.31.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.59

$$\int e^t t^3 dt = (t^3 - 3t^2 + 6t - 6) e^t$$

input `integrate(exp(t)*t^3,t, algorithm="giac")`output `(t^3 - 3*t^2 + 6*t - 6)*e^t`

3.31.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.59

$$\int e^t t^3 dt = e^t (t^3 - 3t^2 + 6t - 6)$$

input `int(t^3*exp(t),t)`

output `exp(t)*(6*t - 3*t^2 + t^3 - 6)`

3.32 $\int e^{2t} \sin(3t) dt$

3.32.1	Optimal result	282
3.32.2	Mathematica [A] (verified)	282
3.32.3	Rubi [A] (verified)	283
3.32.4	Maple [A] (verified)	283
3.32.5	Fricas [A] (verification not implemented)	284
3.32.6	Sympy [A] (verification not implemented)	284
3.32.7	Maxima [A] (verification not implemented)	284
3.32.8	Giac [A] (verification not implemented)	285
3.32.9	Mupad [B] (verification not implemented)	285

3.32.1 Optimal result

Integrand size = 10, antiderivative size = 27

$$\int e^{2t} \sin(3t) dt = -\frac{3}{13}e^{2t} \cos(3t) + \frac{2}{13}e^{2t} \sin(3t)$$

output `-3/13*exp(2*t)*cos(3*t)+2/13*exp(2*t)*sin(3*t)`

3.32.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int e^{2t} \sin(3t) dt = \frac{1}{13}e^{2t}(-3 \cos(3t) + 2 \sin(3t))$$

input `Integrate[E^(2*t)*Sin[3*t],t]`

output `(E^(2*t))*(-3*Cos[3*t] + 2*Sin[3*t])/13`

3.32.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {4932}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{2t} \sin(3t) dt$$

$$\downarrow 4932$$

$$\frac{2}{13} e^{2t} \sin(3t) - \frac{3}{13} e^{2t} \cos(3t)$$

input `Int[E^(2*t)*Sin[3*t],t]`

output `(-3*E^(2*t)*Cos[3*t])/13 + (2*E^(2*t)*Sin[3*t])/13`

3.32.3.1 Defintions of rubi rules used

rule 4932 `Int[(F_)^((c_.)*(a_.) + (b_.)*(x_))*Sin[(d_.) + (e_.)*(x_)], x_Symbol] :=
Simp[b*c*Log[F]*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x
] - Simp[e*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; F
reeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]`

3.32.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result	size
parallelrisch	$\frac{e^{2t}(-3 \cos(3t) + 2 \sin(3t))}{13}$	20
default	$-\frac{3 e^{2t} \cos(3t)}{13} + \frac{2 e^{2t} \sin(3t)}{13}$	22
risch	$-\frac{3 e^{(2+3i)t}}{26} - \frac{i e^{(2+3i)t}}{13} - \frac{3 e^{(2-3i)t}}{26} + \frac{i e^{(2-3i)t}}{13}$	36
norman	$\frac{4 e^{2t} \tan\left(\frac{3t}{2}\right)}{13} + \frac{3 e^{2t} \left(\tan^2\left(\frac{3t}{2}\right)\right)}{13} - \frac{3 e^{2t}}{13}$ $\frac{1}{1 + \tan^2\left(\frac{3t}{2}\right)}$	41

input `int(exp(2*t)*sin(3*t),t,method=_RETURNVERBOSE)`

output `1/13*exp(2*t)*(-3*cos(3*t)+2*sin(3*t))`

3.32.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int e^{2t} \sin(3t) dt = -\frac{3}{13} \cos(3t) e^{(2t)} + \frac{2}{13} e^{(2t)} \sin(3t)$$

input `integrate(exp(2*t)*sin(3*t),t, algorithm="fricas")`

output `-3/13*cos(3*t)*e^(2*t) + 2/13*e^(2*t)*sin(3*t)`

3.32.6 Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int e^{2t} \sin(3t) dt = \frac{2e^{2t} \sin(3t)}{13} - \frac{3e^{2t} \cos(3t)}{13}$$

input `integrate(exp(2*t)*sin(3*t),t)`

output `2*exp(2*t)*sin(3*t)/13 - 3*exp(2*t)*cos(3*t)/13`

3.32.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^{2t} \sin(3t) dt = -\frac{1}{13} (3 \cos(3t) - 2 \sin(3t)) e^{(2t)}$$

input `integrate(exp(2*t)*sin(3*t),t, algorithm="maxima")`

output `-1/13*(3*cos(3*t) - 2*sin(3*t))*e^(2*t)`

3.32.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^{2t} \sin(3t) dt = -\frac{1}{13} (3 \cos(3t) - 2 \sin(3t))e^{(2t)}$$

input `integrate(exp(2*t)*sin(3*t),t, algorithm="giac")`

output `-1/13*(3*cos(3*t) - 2*sin(3*t))*e^(2*t)`

3.32.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^{2t} \sin(3t) dt = -\frac{e^{2t} (3 \cos(3t) - 2 \sin(3t))}{13}$$

input `int(sin(3*t)*exp(2*t),t)`

output `-(exp(2*t)*(3*cos(3*t) - 2*sin(3*t)))/13`

3.33 $\int e^{-t} \cos(3t) dt$

3.33.1	Optimal result	286
3.33.2	Mathematica [A] (verified)	286
3.33.3	Rubi [A] (verified)	287
3.33.4	Maple [A] (verified)	287
3.33.5	Fricas [A] (verification not implemented)	288
3.33.6	Sympy [A] (verification not implemented)	288
3.33.7	Maxima [A] (verification not implemented)	288
3.33.8	Giac [A] (verification not implemented)	289
3.33.9	Mupad [B] (verification not implemented)	289

3.33.1 Optimal result

Integrand size = 10, antiderivative size = 27

$$\int e^{-t} \cos(3t) dt = -\frac{1}{10}e^{-t} \cos(3t) + \frac{3}{10}e^{-t} \sin(3t)$$

output `-1/10*cos(3*t)/exp(t)+3/10*sin(3*t)/exp(t)`

3.33.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

$$\int e^{-t} \cos(3t) dt = -\frac{1}{10}e^{-t}(\cos(3t) - 3 \sin(3t))$$

input `Integrate[Cos[3*t]/E^t,t]`

output `-1/10*(Cos[3*t] - 3*Sin[3*t])/E^t`

3.33.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {4933}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{-t} \cos(3t) dt$$

↓ 4933

$$\frac{3}{10}e^{-t} \sin(3t) - \frac{1}{10}e^{-t} \cos(3t)$$

input `Int[Cos[3*t]/E^t,t]`

output `-1/10*Cos[3*t]/E^t + (3*Sin[3*t])/(10*E^t)`

3.33.3.1 Defintions of rubi rules used

rule 4933 `Int[Cos[(d_.) + (e_.)*(x_)]*(F_)^(c_.*((a_.) + (b_.)*(x_))), x_Symbol] :=
Simp[b*c*Log[F]*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x
] + Simp[e*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; F
reeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]`

3.33.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.67

method	result	size
parallelrisc	$-\frac{e^{-t}(\cos(3t)-3\sin(3t))}{10}$	18
default	$-\frac{e^{-t} \cos(3t)}{10} + \frac{3 e^{-t} \sin(3t)}{10}$	22
norman	$\frac{\left(-\frac{1}{10} + \frac{\tan^2\left(\frac{3t}{2}\right)}{10} + \frac{3 \tan\left(\frac{3t}{2}\right)}{5}\right) e^{-t}}{1 + \tan^2\left(\frac{3t}{2}\right)}$	32
risc	$-\frac{e^{(-1+3i)t}}{20} - \frac{3ie^{(-1+3i)t}}{20} - \frac{e^{(-1-3i)t}}{20} + \frac{3ie^{(-1-3i)t}}{20}$	36

input `int(cos(3*t)/exp(t),t,method=_RETURNVERBOSE)`

output `-1/10*exp(-t)*(cos(3*t)-3*sin(3*t))`

3.33.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int e^{-t} \cos(3t) dt = -\frac{1}{10} \cos(3t) e^{(-t)} + \frac{3}{10} e^{(-t)} \sin(3t)$$

input `integrate(cos(3*t)/exp(t),t, algorithm="fricas")`

output `-1/10*cos(3*t)*e^(-t) + 3/10*e^(-t)*sin(3*t)`

3.33.6 Sympy [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

$$\int e^{-t} \cos(3t) dt = \frac{3e^{-t} \sin(3t)}{10} - \frac{e^{-t} \cos(3t)}{10}$$

input `integrate(cos(3*t)/exp(t),t)`

output `3*exp(-t)*sin(3*t)/10 - exp(-t)*cos(3*t)/10`

3.33.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.63

$$\int e^{-t} \cos(3t) dt = -\frac{1}{10} (\cos(3t) - 3 \sin(3t)) e^{(-t)}$$

input `integrate(cos(3*t)/exp(t),t, algorithm="maxima")`

output `-1/10*(cos(3*t) - 3*sin(3*t))*e^(-t)`

3.33.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.63

$$\int e^{-t} \cos(3t) dt = -\frac{1}{10} (\cos(3t) - 3 \sin(3t))e^{(-t)}$$

input `integrate(cos(3*t)/exp(t),t, algorithm="giac")`

output `-1/10*(cos(3*t) - 3*sin(3*t))*e^(-t)`

3.33.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.63

$$\int e^{-t} \cos(3t) dt = -\frac{e^{-t} (\cos(3t) - 3 \sin(3t))}{10}$$

input `int(cos(3*t)*exp(-t),t)`

output `-(exp(-t)*(cos(3*t) - 3*sin(3*t)))/10`

3.34 $\int y \sinh(y) dy$

3.34.1	Optimal result	290
3.34.2	Mathematica [A] (verified)	290
3.34.3	Rubi [C] (verified)	291
3.34.4	Maple [A] (verified)	292
3.34.5	Fricas [A] (verification not implemented)	293
3.34.6	Sympy [A] (verification not implemented)	293
3.34.7	Maxima [B] (verification not implemented)	293
3.34.8	Giac [A] (verification not implemented)	294
3.34.9	Mupad [B] (verification not implemented)	294

3.34.1 Optimal result

Integrand size = 4, antiderivative size = 9

$$\int y \sinh(y) dy = y \cosh(y) - \sinh(y)$$

output `y*cosh(y)-sinh(y)`

3.34.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int y \sinh(y) dy = y \cosh(y) - \sinh(y)$$

input `Integrate[y*Sinh[y],y]`

output `y*Cosh[y] - Sinh[y]`

3.34.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.18 (sec) , antiderivative size = 18, normalized size of antiderivative = 2.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3042, 26, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int y \sinh(y) dy \\
 & \quad \downarrow \text{3042} \\
 & \int -iy \sin(iy) dy \\
 & \quad \downarrow \text{26} \\
 & -i \int y \sin(iy) dy \\
 & \quad \downarrow \text{3777} \\
 & -i(iy \cosh(y) - i \int \cosh(y) dy) \\
 & \quad \downarrow \text{3042} \\
 & -i \left(iy \cosh(y) - i \int \sin \left(iy + \frac{\pi}{2} \right) dy \right) \\
 & \quad \downarrow \text{3117} \\
 & -i(iy \cosh(y) - i \sinh(y))
 \end{aligned}$$

input `Int[y*Sinh[y],y]`

output `(-I)*(I*y*Cosh[y] - I*Sinh[y])`

3.34.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

3.34.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.11

method	result	size
default	$y \cosh(y) - \sinh(y)$	10
meijerg	$y \cosh(y) - \sinh(y)$	10
parallelrisch	$y \cosh(y) - \sinh(y)$	10
parts	$y \cosh(y) - \sinh(y)$	10
risch	$(-\frac{1}{2} + \frac{y}{2}) e^y + (\frac{1}{2} + \frac{y}{2}) e^{-y}$	20

input `int(y*sinh(y),y,method=_RETURNVERBOSE)`

output `y*cosh(y)-sinh(y)`

3.34.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int y \sinh(y) dy = y \cosh(y) - \sinh(y)$$

input `integrate(y*sinh(y),y, algorithm="fricas")`

output `y*cosh(y) - sinh(y)`

3.34.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int y \sinh(y) dy = y \cosh(y) - \sinh(y)$$

input `integrate(y*sinh(y),y)`

output `y*cosh(y) - sinh(y)`

3.34.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. 2(9) = 18.

Time = 0.20 (sec) , antiderivative size = 34, normalized size of antiderivative = 3.78

$$\int y \sinh(y) dy = \frac{1}{2} y^2 \sinh(y) + \frac{1}{4} (y^2 + 2y + 2)e^{-y} - \frac{1}{4} (y^2 - 2y + 2)e^y$$

input `integrate(y*sinh(y),y, algorithm="maxima")`

output `1/2*y^2*sinh(y) + 1/4*(y^2 + 2*y + 2)*e^(-y) - 1/4*(y^2 - 2*y + 2)*e^y`

3.34.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.89

$$\int y \sinh(y) dy = \frac{1}{2} (y + 1)e^{(-y)} + \frac{1}{2} (y - 1)e^y$$

input `integrate(y*sinh(y),y, algorithm="giac")`

output `1/2*(y + 1)*e^(-y) + 1/2*(y - 1)*e^y`

3.34.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int y \sinh(y) dy = y \cosh(y) - \sinh(y)$$

input `int(y*sinh(y),y)`

output `y*cosh(y) - sinh(y)`

3.35 $\int y \cosh(ay) dy$

3.35.1	Optimal result	295
3.35.2	Mathematica [A] (verified)	295
3.35.3	Rubi [A] (verified)	296
3.35.4	Maple [A] (verified)	297
3.35.5	Fricas [A] (verification not implemented)	298
3.35.6	Sympy [A] (verification not implemented)	298
3.35.7	Maxima [B] (verification not implemented)	298
3.35.8	Giac [A] (verification not implemented)	299
3.35.9	Mupad [B] (verification not implemented)	299

3.35.1 Optimal result

Integrand size = 6, antiderivative size = 19

$$\int y \cosh(ay) dy = -\frac{\cosh(ay)}{a^2} + \frac{y \sinh(ay)}{a}$$

output `-cosh(a*y)/a^2+y*sinh(a*y)/a`

3.35.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int y \cosh(ay) dy = -\frac{\cosh(ay)}{a^2} + \frac{y \sinh(ay)}{a}$$

input `Integrate[y*Cosh[a*y],y]`

output `-(Cosh[a*y]/a^2) + (y*Sinh[a*y])/a`

3.35.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 3777, 26, 3042, 26, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int y \cosh(ay) dy \\
 & \quad \downarrow \text{3042} \\
 & \int y \sin\left(\frac{\pi}{2} + iay\right) dy \\
 & \quad \downarrow \text{3777} \\
 & \frac{y \sinh(ay)}{a} - \frac{i \int -i \sinh(ay) dy}{a} \\
 & \quad \downarrow \text{26} \\
 & \frac{y \sinh(ay)}{a} - \frac{\int \sinh(ay) dy}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{y \sinh(ay)}{a} - \frac{\int -i \sin(iay) dy}{a} \\
 & \quad \downarrow \text{26} \\
 & \frac{y \sinh(ay)}{a} + \frac{i \int \sin(iay) dy}{a} \\
 & \quad \downarrow \text{3118} \\
 & \frac{y \sinh(ay)}{a} - \frac{\cosh(ay)}{a^2}
 \end{aligned}$$

input `Int [y*Cosh[a*y] ,y]`

output `-(Cosh[a*y]/a^2) + (y*Sinh[a*y])/a`

3.35.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

3.35.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

method	result	size
derivativedivides	$\frac{ya \sinh(ay) - \cosh(ay)}{a^2}$	19
default	$\frac{ya \sinh(ay) - \cosh(ay)}{a^2}$	19
parts	$-\frac{\cosh(ay)}{a^2} + \frac{y \sinh(ay)}{a}$	20
parallelrisc	$\frac{2 - 2y \tanh(\frac{ay}{2}) a}{a^2 (\tanh^2(\frac{ay}{2}) - 1)}$	27
risch	$\frac{(ay-1)e^{ay}}{2a^2} - \frac{(ay+1)e^{-ay}}{2a^2}$	32
meijerg	$-\frac{2\sqrt{\pi} \left(-\frac{1}{2\sqrt{\pi}} + \frac{\cosh(ay)}{2\sqrt{\pi}} - \frac{ya \sinh(ay)}{2\sqrt{\pi}} \right)}{a^2}$	35

input `int(y*cosh(a*y), y, method=_RETURNVERBOSE)`

output `1/a^2*(y*a*sinh(a*y)-cosh(a*y))`

3.35.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

$$\int y \cosh(ay) dy = \frac{ay \sinh(ay) - \cosh(ay)}{a^2}$$

input `integrate(y*cosh(a*y),y, algorithm="fricas")`

output `(a*y*sinh(a*y) - cosh(a*y))/a^2`

3.35.6 Sympy [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.05

$$\int y \cosh(ay) dy = \begin{cases} \frac{y \sinh(ay)}{a} - \frac{\cosh(ay)}{a^2} & \text{for } a \neq 0 \\ \frac{y^2}{2} & \text{otherwise} \end{cases}$$

input `integrate(y*cosh(a*y),y)`

output `Piecewise((y*sinh(a*y)/a - cosh(a*y)/a**2, Ne(a, 0)), (y**2/2, True))`

3.35.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 57 vs. 2(19) = 38.

Time = 0.18 (sec) , antiderivative size = 57, normalized size of antiderivative = 3.00

$$\int y \cosh(ay) dy = \frac{1}{2} y^2 \cosh(ay) - \frac{1}{4} a \left(\frac{(a^2 y^2 - 2ay + 2)e^{ay}}{a^3} + \frac{(a^2 y^2 + 2ay + 2)e^{-ay}}{a^3} \right)$$

input `integrate(y*cosh(a*y),y, algorithm="maxima")`

output `1/2*y^2*cosh(a*y) - 1/4*a*((a^2*y^2 - 2*a*y + 2)*e^(a*y)/a^3 + (a^2*y^2 + 2*a*y + 2)*e^(-a*y)/a^3)`

3.35.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.58

$$\int y \cosh(ay) dy = \frac{(ay - 1)e^{(ay)}}{2a^2} - \frac{(ay + 1)e^{(-ay)}}{2a^2}$$

input `integrate(y*cosh(a*y),y, algorithm="giac")`output `1/2*(a*y - 1)*e^(a*y)/a^2 - 1/2*(a*y + 1)*e^(-a*y)/a^2`**3.35.9 Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

$$\int y \cosh(ay) dy = -\frac{\cosh(ay) - ay \sinh(ay)}{a^2}$$

input `int(y*cosh(a*y),y)`output `-(cosh(a*y) - a*y*sinh(a*y))/a^2`

3.36 $\int e^{-t}t dt$

3.36.1	Optimal result	300
3.36.2	Mathematica [A] (verified)	300
3.36.3	Rubi [A] (verified)	301
3.36.4	Maple [A] (verified)	302
3.36.5	Fricas [A] (verification not implemented)	302
3.36.6	Sympy [A] (verification not implemented)	302
3.36.7	Maxima [A] (verification not implemented)	303
3.36.8	Giac [A] (verification not implemented)	303
3.36.9	Mupad [B] (verification not implemented)	303

3.36.1 Optimal result

Integrand size = 7, antiderivative size = 16

$$\int e^{-t}t dt = -e^{-t} - e^{-t}t$$

output `-1/exp(t)-t/exp(t)`

3.36.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.69

$$\int e^{-t}t dt = e^{-t}(-1 - t)$$

input `Integrate[t/E^t,t]`

output `(-1 - t)/E^t`

3.36.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int e^{-t} dt \\ \downarrow \text{2607} \\ \int e^{-t} dt - e^{-t} \\ \downarrow \text{2624} \\ -e^{-t} - e^{-t} \end{array}$$

input `Int[t/E^t,t]`

output `-E^(-t) - t/E^t`

3.36.3.1 Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`

3.36.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

method	result	size
gospers	$-(1+t)e^{-t}$	10
norman	$(-1-t)e^{-t}$	11
risch	$(-1-t)e^{-t}$	11
parallelrisch	$(-1-t)e^{-t}$	11
meijerg	$1 - \frac{(2+2t)e^{-t}}{2}$	14
default	$-e^{-t} - te^{-t}$	15

input `int(t/exp(t),t,method=_RETURNVERBOSE)`

output `-(1+t)/exp(t)`

3.36.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.56

$$\int e^{-t}t dt = -(t+1)e^{(-t)}$$

input `integrate(t/exp(t),t, algorithm="fracas")`

output `-(t + 1)*e^(-t)`

3.36.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.44

$$\int e^{-t}t dt = (-t-1)e^{-t}$$

input `integrate(t/exp(t),t)`

output `(-t - 1)*exp(-t)`

3.36.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.56

$$\int e^{-t}t dt = -(t+1)e^{(-t)}$$

input `integrate(t/exp(t),t, algorithm="maxima")`

output `-(t + 1)*e^(-t)`

3.36.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.56

$$\int e^{-t}t dt = -(t+1)e^{(-t)}$$

input `integrate(t/exp(t),t, algorithm="giac")`

output `-(t + 1)*e^(-t)`

3.36.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.56

$$\int e^{-t}t dt = -e^{-t}(t+1)$$

input `int(t*exp(-t),t)`

output `-exp(-t)*(t + 1)`

3.37 $\int \sqrt{t} \log(t) dt$

3.37.1	Optimal result	304
3.37.2	Mathematica [A] (verified)	304
3.37.3	Rubi [A] (verified)	305
3.37.4	Maple [A] (verified)	305
3.37.5	Fricas [A] (verification not implemented)	306
3.37.6	Sympy [B] (verification not implemented)	306
3.37.7	Maxima [A] (verification not implemented)	307
3.37.8	Giac [A] (verification not implemented)	307
3.37.9	Mupad [B] (verification not implemented)	307

3.37.1 Optimal result

Integrand size = 8, antiderivative size = 21

$$\int \sqrt{t} \log(t) dt = -\frac{4t^{3/2}}{9} + \frac{2}{3}t^{3/2} \log(t)$$

output `-4/9*t^(3/2)+2/3*t^(3/2)*ln(t)`

3.37.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \sqrt{t} \log(t) dt = \frac{2}{9}t^{3/2}(-2 + 3 \log(t))$$

input `Integrate[Sqrt[t]*Log[t],t]`

output `(2*t^(3/2)*(-2 + 3*Log[t]))/9`

3.37.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{t} \log(t) dt$$

↓ 2741

$$\frac{2}{3}t^{3/2} \log(t) - \frac{4t^{3/2}}{9}$$

input `Int[Sqrt[t]*Log[t],t]`

output `(-4*t^(3/2))/9 + (2*t^(3/2)*Log[t])/3`

3.37.3.1 Defintions of rubi rules used

rule 2741 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_.), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(m + 1))/(d*(m + 1)^2), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]`

3.37.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

method	result	size
derivativedivides	$-\frac{4t^{\frac{3}{2}}}{9} + \frac{2t^{\frac{3}{2}} \ln(t)}{3}$	14
default	$-\frac{4t^{\frac{3}{2}}}{9} + \frac{2t^{\frac{3}{2}} \ln(t)}{3}$	14
risch	$-\frac{4t^{\frac{3}{2}}}{9} + \frac{2t^{\frac{3}{2}} \ln(t)}{3}$	14

input `int(ln(t)*t^(1/2),t,method=_RETURNVERBOSE)`

output `-4/9*t^(3/2)+2/3*t^(3/2)*ln(t)`

3.37.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

$$\int \sqrt{t} \log(t) dt = \frac{2}{9} (3t \log(t) - 2t) \sqrt{t}$$

input `integrate(log(t)*t^(1/2),t, algorithm="fricas")`

output `2/9*(3*t*log(t) - 2*t)*sqrt(t)`

3.37.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. 2(19) = 38.

Time = 0.92 (sec) , antiderivative size = 105, normalized size of antiderivative = 5.00

$$\int \sqrt{t} \log(t) dt = \begin{cases} -\frac{2t^{\frac{3}{2}} \log(\frac{1}{t})}{3} + \frac{2t^{\frac{3}{2}} \log(t)}{3} - \frac{8t^{\frac{3}{2}}}{9} & \text{for } \frac{1}{|t|} < 1 \wedge |t| < 1 \\ \frac{2t^{\frac{3}{2}} \log(t)}{3} - \frac{4t^{\frac{3}{2}}}{9} & \text{for } |t| < 1 \\ -\frac{2t^{\frac{3}{2}} \log(\frac{1}{t})}{3} - \frac{4t^{\frac{3}{2}}}{9} & \text{for } \frac{1}{|t|} < 1 \\ -G_{3,3}^{2,1} \left(\begin{matrix} 1 & \frac{5}{2}, \frac{5}{2} \\ \frac{3}{2}, \frac{3}{2} \end{matrix} \middle| t \right) + G_{3,3}^{0,3} \left(\begin{matrix} \frac{5}{2}, \frac{5}{2}, 1 \\ \frac{3}{2}, \frac{3}{2}, 0 \end{matrix} \middle| t \right) & \text{otherwise} \end{cases}$$

input `integrate(ln(t)*t**(1/2),t)`

output `Piecewise((-2*t**(3/2)*log(1/t)/3 + 2*t**(3/2)*log(t)/3 - 8*t**(3/2)/9, (Abs(t) < 1) & (1/Abs(t) < 1)), (2*t**(3/2)*log(t)/3 - 4*t**(3/2)/9, Abs(t) < 1), (-2*t**(3/2)*log(1/t)/3 - 4*t**(3/2)/9, 1/Abs(t) < 1), (-meijerg(((1, (5/2, 5/2)), ((3/2, 3/2), (0,)), t) + meijerg(((5/2, 5/2, 1), ()), ((3/2, 3/2, 0)), t), True))`

3.37.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \sqrt{t} \log(t) dt = \frac{2}{3} t^{\frac{3}{2}} \log(t) - \frac{4}{9} t^{\frac{3}{2}}$$

input `integrate(log(t)*t^(1/2),t, algorithm="maxima")`output `2/3*t^(3/2)*log(t) - 4/9*t^(3/2)`**3.37.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \sqrt{t} \log(t) dt = \frac{2}{3} t^{\frac{3}{2}} \log(t) - \frac{4}{9} t^{\frac{3}{2}}$$

input `integrate(log(t)*t^(1/2),t, algorithm="giac")`output `2/3*t^(3/2)*log(t) - 4/9*t^(3/2)`**3.37.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.43

$$\int \sqrt{t} \log(t) dt = \frac{2t^{3/2} (\ln(t) - \frac{2}{3})}{3}$$

input `int(t^(1/2)*log(t),t)`output `(2*t^(3/2)*(log(t) - 2/3))/3`

3.38 $\int x \cos(2x) dx$

3.38.1	Optimal result	308
3.38.2	Mathematica [A] (verified)	308
3.38.3	Rubi [A] (verified)	309
3.38.4	Maple [A] (verified)	310
3.38.5	Fricas [A] (verification not implemented)	311
3.38.6	Sympy [A] (verification not implemented)	311
3.38.7	Maxima [A] (verification not implemented)	311
3.38.8	Giac [A] (verification not implemented)	312
3.38.9	Mupad [B] (verification not implemented)	312

3.38.1 Optimal result

Integrand size = 6, antiderivative size = 18

$$\int x \cos(2x) dx = \frac{1}{4} \cos(2x) + \frac{1}{2} x \sin(2x)$$

output `1/4*cos(2*x)+1/2*x*sin(2*x)`

3.38.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int x \cos(2x) dx = \frac{1}{4} \cos(2x) + \frac{1}{2} x \sin(2x)$$

input `Integrate[x*Cos[2*x],x]`

output `Cos[2*x]/4 + (x*Sin[2*x])/2`

3.38.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \cos(2x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x \sin\left(2x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3777} \\
 & \frac{1}{2} \int -\sin(2x) dx + \frac{1}{2} x \sin(2x) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{2} x \sin(2x) - \frac{1}{2} \int \sin(2x) dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} x \sin(2x) - \frac{1}{2} \int \sin(2x) dx \\
 & \quad \downarrow \text{3118} \\
 & \frac{1}{2} x \sin(2x) + \frac{1}{4} \cos(2x)
 \end{aligned}$$

input `Int[x*Cos[2*x],x]`

output `Cos[2*x]/4 + (x*Sin[2*x])/2`

3.38.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

3.38.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$\frac{\cos(2x)}{4} + \frac{x \sin(2x)}{2}$	15
default	$\frac{\cos(2x)}{4} + \frac{x \sin(2x)}{2}$	15
risch	$\frac{\cos(2x)}{4} + \frac{x \sin(2x)}{2}$	15
parts	$\frac{\cos(2x)}{4} + \frac{x \sin(2x)}{2}$	15
norman	$\frac{x \tan(x) + \frac{1}{2}}{1 + \tan^2(x)}$	16
parallelrisc	$\frac{x \sin(2x)}{2} + \frac{\cos(2x)}{4} + \frac{1}{4}$	16
meijerg	$\frac{\sqrt{\pi} \left(-\frac{1}{2\sqrt{\pi}} + \frac{\cos(2x)}{2\sqrt{\pi}} + \frac{x \sin(2x)}{\sqrt{\pi}} \right)}{2}$	30

input `int(x*cos(2*x),x,method=_RETURNVERBOSE)`

output `1/4*cos(2*x)+1/2*x*sin(2*x)`

3.38.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int x \cos(2x) dx = \frac{1}{2} x \sin(2x) + \frac{1}{4} \cos(2x)$$

input `integrate(x*cos(2*x),x, algorithm="fricas")`output `1/2*x*sin(2*x) + 1/4*cos(2*x)`**3.38.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int x \cos(2x) dx = \frac{x \sin(2x)}{2} + \frac{\cos(2x)}{4}$$

input `integrate(x*cos(2*x),x)`output `x*sin(2*x)/2 + cos(2*x)/4`**3.38.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int x \cos(2x) dx = \frac{1}{2} x \sin(2x) + \frac{1}{4} \cos(2x)$$

input `integrate(x*cos(2*x),x, algorithm="maxima")`output `1/2*x*sin(2*x) + 1/4*cos(2*x)`

3.38.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int x \cos(2x) dx = \frac{1}{2} x \sin(2x) + \frac{1}{4} \cos(2x)$$

input `integrate(x*cos(2*x),x, algorithm="giac")`

output `1/2*x*sin(2*x) + 1/4*cos(2*x)`

3.38.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int x \cos(2x) dx = \frac{\cos(2x)}{4} + \frac{x \sin(2x)}{2}$$

input `int(x*cos(2*x),x)`

output `cos(2*x)/4 + (x*sin(2*x))/2`

3.39 $\int e^{-x} x^2 dx$

3.39.1	Optimal result	313
3.39.2	Mathematica [A] (verified)	313
3.39.3	Rubi [A] (verified)	314
3.39.4	Maple [A] (verified)	315
3.39.5	Fricas [A] (verification not implemented)	315
3.39.6	Sympy [A] (verification not implemented)	315
3.39.7	Maxima [A] (verification not implemented)	316
3.39.8	Giac [A] (verification not implemented)	316
3.39.9	Mupad [B] (verification not implemented)	316

3.39.1 Optimal result

Integrand size = 9, antiderivative size = 26

$$\int e^{-x} x^2 dx = -2e^{-x} - 2e^{-x}x - e^{-x}x^2$$

output `-2/exp(x)-2*x/exp(x)-x^2/exp(x)`

3.39.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.62

$$\int e^{-x} x^2 dx = e^{-x}(-2 - 2x - x^2)$$

input `Integrate[x^2/E^x,x]`

output `(-2 - 2*x - x^2)/E^x`

3.39.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.12, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2607, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^{-x} x^2 dx \\ & \quad \downarrow \text{2607} \\ & 2 \int e^{-x} x dx - e^{-x} x^2 \\ & \quad \downarrow \text{2607} \\ & 2 \left(\int e^{-x} dx - e^{-x} x \right) - e^{-x} x^2 \\ & \quad \downarrow \text{2624} \\ & 2(-e^{-x} x - e^{-x}) - e^{-x} x^2 \end{aligned}$$

input `Int[x^2/E^x,x]`

output `-(x^2/E^x) + 2*(-E^(-x) - x/E^x)`

3.39.3.1 Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2624 `Int[((F_)^(v_))^(n_.), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`

3.39.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.58

method	result	size
gospers	$-(x^2 + 2x + 2)e^{-x}$	15
norman	$(-x^2 - 2x - 2)e^{-x}$	16
risch	$(-x^2 - 2x - 2)e^{-x}$	16
parallelrisc	$(-x^2 - 2x - 2)e^{-x}$	16
meijerg	$2 - \frac{(3x^2+6x+6)e^{-x}}{3}$	19
default	$-2e^{-x} - 2xe^{-x} - x^2e^{-x}$	24

input `int(x^2/exp(x),x,method=_RETURNVERBOSE)`output `-(x^2+2*x+2)/exp(x)`**3.39.5 Fricas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.54

$$\int e^{-x}x^2 dx = -(x^2 + 2x + 2)e^{(-x)}$$

input `integrate(x^2/exp(x),x, algorithm="fricas")`output `-(x^2 + 2*x + 2)*e^(-x)`**3.39.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.46

$$\int e^{-x}x^2 dx = (-x^2 - 2x - 2)e^{-x}$$

input `integrate(x**2/exp(x),x)`output `(-x**2 - 2*x - 2)*exp(-x)`

3.39.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.54

$$\int e^{-x}x^2 dx = -(x^2 + 2x + 2)e^{(-x)}$$

input `integrate(x^2/exp(x),x, algorithm="maxima")`output `-(x^2 + 2*x + 2)*e^(-x)`**3.39.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.54

$$\int e^{-x}x^2 dx = -(x^2 + 2x + 2)e^{(-x)}$$

input `integrate(x^2/exp(x),x, algorithm="giac")`output `-(x^2 + 2*x + 2)*e^(-x)`**3.39.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.54

$$\int e^{-x}x^2 dx = -e^{-x}(x^2 + 2x + 2)$$

input `int(x^2*exp(-x),x)`output `-exp(-x)*(2*x + x^2 + 2)`

3.40 $\int \arccos(x) dx$

3.40.1	Optimal result	317
3.40.2	Mathematica [A] (verified)	317
3.40.3	Rubi [A] (verified)	318
3.40.4	Maple [A] (verified)	319
3.40.5	Fricas [A] (verification not implemented)	319
3.40.6	Sympy [A] (verification not implemented)	319
3.40.7	Maxima [A] (verification not implemented)	320
3.40.8	Giac [A] (verification not implemented)	320
3.40.9	Mupad [B] (verification not implemented)	320

3.40.1 Optimal result

Integrand size = 2, antiderivative size = 18

$$\int \arccos(x) dx = -\sqrt{1-x^2} + x \arccos(x)$$

output `x*arccos(x)-(-x^2+1)^(1/2)`

3.40.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \arccos(x) dx = -\sqrt{1-x^2} + x \arccos(x)$$

input `Integrate[ArcCos[x],x]`

output `-Sqrt[1 - x^2] + x*ArcCos[x]`

3.40.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {5131, 241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arccos(x) dx$$

$$\downarrow \text{5131}$$

$$\int \frac{x}{\sqrt{1-x^2}} dx + x \arccos(x)$$

$$\downarrow \text{241}$$

$$x \arccos(x) - \sqrt{1-x^2}$$

input `Int[ArcCos[x], x]`

output `-Sqrt[1 - x^2] + x*ArcCos[x]`

3.40.3.1 Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

rule 5131 `Int[((a_) + ArcCos[(c_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[x*(a + b*ArcCos[c*x])^n, x] + Simp[b*c*n Int[x*((a + b*ArcCos[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

3.40.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

method	result	size
lookup	$x \arccos(x) - \sqrt{-x^2 + 1}$	17
default	$x \arccos(x) - \sqrt{-x^2 + 1}$	17
parts	$x \arccos(x) - \sqrt{-x^2 + 1}$	17

input `int(arccos(x), x, method=_RETURNVERBOSE)`

output `x*arccos(x)-(-x^2+1)^(1/2)`

3.40.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \arccos(x) dx = x \arccos(x) - \sqrt{-x^2 + 1}$$

input `integrate(arccos(x), x, algorithm="fricas")`

output `x*arccos(x) - sqrt(-x^2 + 1)`

3.40.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \arccos(x) dx = x \arccos(x) - \sqrt{1 - x^2}$$

input `integrate(acos(x), x)`

output `x*acos(x) - sqrt(1 - x**2)`

3.40.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \arccos(x) dx = x \arccos(x) - \sqrt{-x^2 + 1}$$

input `integrate(arccos(x),x, algorithm="maxima")`output `x*arccos(x) - sqrt(-x^2 + 1)`**3.40.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \arccos(x) dx = x \arccos(x) - \sqrt{-x^2 + 1}$$

input `integrate(arccos(x),x, algorithm="giac")`output `x*arccos(x) - sqrt(-x^2 + 1)`**3.40.9 Mupad [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \arccos(x) dx = x \arccos(x) - \sqrt{1 - x^2}$$

input `int(acos(x),x)`output `x*acos(x) - (1 - x^2)^(1/2)`

3.41 $\int x \csc^2(x) dx$

3.41.1	Optimal result	321
3.41.2	Mathematica [A] (verified)	321
3.41.3	Rubi [A] (verified)	322
3.41.4	Maple [A] (verified)	323
3.41.5	Fricas [B] (verification not implemented)	324
3.41.6	Sympy [A] (verification not implemented)	324
3.41.7	Maxima [B] (verification not implemented)	324
3.41.8	Giac [B] (verification not implemented)	325
3.41.9	Mupad [B] (verification not implemented)	325

3.41.1 Optimal result

Integrand size = 6, antiderivative size = 9

$$\int x \csc^2(x) dx = -x \cot(x) + \log(\sin(x))$$

output `-x*cot(x)+ln(sin(x))`

3.41.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int x \csc^2(x) dx = -x \cot(x) + \log(\sin(x))$$

input `Integrate[x*Csc[x]^2,x]`

output `-(x*Cot[x]) + Log[Sin[x]]`

3.41.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {3042, 4672, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \csc^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x \csc(x)^2 dx \\
 & \quad \downarrow \text{4672} \\
 & \int \cot(x) dx - x \cot(x) \\
 & \quad \downarrow \text{3042} \\
 & \int -\tan\left(x + \frac{\pi}{2}\right) dx - x \cot(x) \\
 & \quad \downarrow \text{25} \\
 & -\int \tan\left(x + \frac{\pi}{2}\right) dx - x \cot(x) \\
 & \quad \downarrow \text{3956} \\
 & \log(\sin(x)) - x \cot(x)
 \end{aligned}$$

input `Int [x*Csc [x] ^2, x]`

output `-(x*Cot [x]) + Log [Sin [x]]`

3.41.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4672 `Int[csc[(e_.) + (f_.)*(x_.)]^2*((c_.) + (d_.)*(x_.))^(m_.), x_Symbol] := Simp[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

3.41.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.11

method	result	size
default	$-x \cot(x) + \ln(\sin(x))$	10
parallelrisch	$-\ln\left(\frac{2}{\cos(x)+1}\right) + \ln(\csc(x) - \cot(x)) - x \cot(x)$	26
risch	$-2ix - \frac{2ix}{e^{2ix}-1} + \ln(e^{2ix} - 1)$	27
norman	$\frac{-\frac{x}{2} + \frac{x \tan^2(\frac{x}{2})}{2}}{\tan(\frac{x}{2})} - \ln(1 + \tan^2(\frac{x}{2})) + \ln(\tan(\frac{x}{2}))$	38

input `int(x*csc(x)^2,x,method=_RETURNVERBOSE)`

output `-x*cot(x)+ln(sin(x))`

3.41.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. $2(9) = 18$.

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 2.22

$$\int x \csc^2(x) dx = -\frac{x \cos(x) - \log\left(\frac{1}{2} \sin(x)\right) \sin(x)}{\sin(x)}$$

input `integrate(x*csc(x)^2,x, algorithm="fricas")`

output `-(x*cos(x) - log(1/2*sin(x))*sin(x))/sin(x)`

3.41.6 Sympy [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int x \csc^2(x) dx = -x \cot(x) + \log(\sin(x))$$

input `integrate(x*csc(x)**2,x)`

output `-x*cot(x) + log(sin(x))`

3.41.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 104 vs. $2(9) = 18$.

Time = 0.19 (sec) , antiderivative size = 104, normalized size of antiderivative = 11.56

$$\int x \csc^2(x) dx = \frac{(\cos(2x)^2 + \sin(2x)^2 - 2 \cos(2x) + 1) \log(\cos(x)^2 + \sin(x)^2 + 2 \cos(x) + 1) + (\cos(2x)^2 + \sin(2x)^2 - 2 \cos(2x) + 1) \log(\cos(x)^2 + \sin(x)^2 - 2 \cos(x) + 1) - 4x \sin(2x)}{2(\cos(2x)^2 + \sin(2x)^2 - 2 \cos(2x) + 1)}$$

input `integrate(x*csc(x)^2,x, algorithm="maxima")`

output `1/2*((cos(2*x)^2 + sin(2*x)^2 - 2*cos(2*x) + 1)*log(cos(x)^2 + sin(x)^2 + 2*cos(x) + 1) + (cos(2*x)^2 + sin(2*x)^2 - 2*cos(2*x) + 1)*log(cos(x)^2 + sin(x)^2 - 2*cos(x) + 1) - 4*x*sin(2*x))/(cos(2*x)^2 + sin(2*x)^2 - 2*cos(2*x) + 1)`

3.41.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 52 vs. 2(9) = 18.

Time = 0.31 (sec) , antiderivative size = 52, normalized size of antiderivative = 5.78

$$\int x \csc^2(x) dx = \frac{x \tan\left(\frac{1}{2}x\right)^2 + \log\left(\frac{16 \tan\left(\frac{1}{2}x\right)^2}{\tan\left(\frac{1}{2}x\right)^4 + 2 \tan\left(\frac{1}{2}x\right)^2 + 1}\right) \tan\left(\frac{1}{2}x\right) - x}{2 \tan\left(\frac{1}{2}x\right)}$$

input `integrate(x*csc(x)^2,x, algorithm="giac")`

output `1/2*(x*tan(1/2*x)^2 + log(16*tan(1/2*x)^2/(tan(1/2*x)^4 + 2*tan(1/2*x)^2 + 1))*tan(1/2*x) - x)/tan(1/2*x)`

3.41.9 Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int x \csc^2(x) dx = \ln(\sin(x)) - x \cot(x)$$

input `int(x/sin(x)^2,x)`

output `log(sin(x)) - x*cot(x)`

3.42 $\int \cos(5x) \sin(3x) dx$

3.42.1	Optimal result	326
3.42.2	Mathematica [A] (verified)	326
3.42.3	Rubi [A] (verified)	327
3.42.4	Maple [A] (verified)	328
3.42.5	Fricas [A] (verification not implemented)	328
3.42.6	Sympy [B] (verification not implemented)	328
3.42.7	Maxima [A] (verification not implemented)	329
3.42.8	Giac [A] (verification not implemented)	329
3.42.9	Mupad [B] (verification not implemented)	329

3.42.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \cos(5x) \sin(3x) dx = \frac{1}{4} \cos(2x) - \frac{1}{16} \cos(8x)$$

output `1/4*cos(2*x)-1/16*cos(8*x)`

3.42.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cos(5x) \sin(3x) dx = \frac{\cos^2(x)}{2} - \frac{1}{16} \cos(8x)$$

input `Integrate[Cos[5*x]*Sin[3*x],x]`

output `Cos[x]^2/2 - Cos[8*x]/16`

3.42.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4772}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sin(3x) \cos(5x) dx \\ \downarrow 3042 \\ \int \sin(3x) \cos(5x) dx \\ \downarrow 4772 \\ \frac{1}{4} \cos(2x) - \frac{1}{16} \cos(8x) \end{array}$$

input `Int[Cos[5*x]*Sin[3*x],x]`

output `Cos[2*x]/4 - Cos[8*x]/16`

3.42.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4772 `Int[cos[(c_.) + (d_.)*(x_)]*sin[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[-Cos[a - c + (b - d)*x]/(2*(b - d)), x] - Simp[Cos[a + c + (b + d)*x]/(2*(b + d)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

3.42.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{\cos(2x)}{4} - \frac{\cos(8x)}{16}$	14
risch	$\frac{\cos(2x)}{4} - \frac{\cos(8x)}{16}$	14
parallelrisc	$-\frac{3}{16} + \frac{(2-\cos(6x))\cos(2x)}{8} + \frac{\cos(4x)}{16}$	23
norman	$\frac{-\frac{3(\tan^2(\frac{3x}{2}))}{8} - \frac{3(\tan^2(\frac{5x}{2}))}{8} + \frac{5 \tan(\frac{3x}{2}) \tan(\frac{5x}{2})}{4}}{(1+\tan^2(\frac{5x}{2}))(1+\tan^2(\frac{3x}{2}))}$	49

input `int(cos(5*x)*sin(3*x),x,method=_RETURNVERBOSE)`output `1/4*cos(2*x)-1/16*cos(8*x)`**3.42.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.47

$$\int \cos(5x) \sin(3x) dx = -8 \cos(x)^8 + 16 \cos(x)^6 - 10 \cos(x)^4 + \frac{5}{2} \cos(x)^2$$

input `integrate(cos(5*x)*sin(3*x),x, algorithm="fricas")`output `-8*cos(x)^8 + 16*cos(x)^6 - 10*cos(x)^4 + 5/2*cos(x)^2`**3.42.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. 2(12) = 24.

Time = 0.13 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.53

$$\int \cos(5x) \sin(3x) dx = \frac{5 \sin(3x) \sin(5x)}{16} + \frac{3 \cos(3x) \cos(5x)}{16}$$

input `integrate(cos(5*x)*sin(3*x),x)`output `5*sin(3*x)*sin(5*x)/16 + 3*cos(3*x)*cos(5*x)/16`

3.42.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(5x) \sin(3x) dx = -\frac{1}{16} \cos(8x) + \frac{1}{4} \cos(2x)$$

input `integrate(cos(5*x)*sin(3*x),x, algorithm="maxima")`output `-1/16*cos(8*x) + 1/4*cos(2*x)`**3.42.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(5x) \sin(3x) dx = -\frac{1}{16} \cos(8x) + \frac{1}{4} \cos(2x)$$

input `integrate(cos(5*x)*sin(3*x),x, algorithm="giac")`output `-1/16*cos(8*x) + 1/4*cos(2*x)`**3.42.9 Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(5x) \sin(3x) dx = \frac{\cos(2x)}{4} - \frac{\cos(8x)}{16}$$

input `int(cos(5*x)*sin(3*x),x)`output `cos(2*x)/4 - cos(8*x)/16`

3.43 $\int \sin(2x) \sin(4x) dx$

3.43.1	Optimal result	330
3.43.2	Mathematica [A] (verified)	330
3.43.3	Rubi [A] (verified)	331
3.43.4	Maple [A] (verified)	332
3.43.5	Fricas [A] (verification not implemented)	332
3.43.6	Sympy [A] (verification not implemented)	332
3.43.7	Maxima [A] (verification not implemented)	333
3.43.8	Giac [A] (verification not implemented)	333
3.43.9	Mupad [B] (verification not implemented)	333

3.43.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \sin(2x) \sin(4x) dx = \frac{1}{4} \sin(2x) - \frac{1}{12} \sin(6x)$$

output `1/4*sin(2*x)-1/12*sin(6*x)`

3.43.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sin(2x) \sin(4x) dx = \frac{1}{4} \sin(2x) - \frac{1}{12} \sin(6x)$$

input `Integrate[Sin[2*x]*Sin[4*x],x]`

output `Sin[2*x]/4 - Sin[6*x]/12`

3.43.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4770}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sin(2x) \sin(4x) dx \\ \downarrow 3042 \\ \int \sin(2x) \sin(4x) dx \\ \downarrow 4770 \\ \frac{1}{4} \sin(2x) - \frac{1}{12} \sin(6x) \end{array}$$

input `Int[Sin[2*x]*Sin[4*x],x]`

output `Sin[2*x]/4 - Sin[6*x]/12`

3.43.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4770 `Int[sin[(a_.) + (b_.)*(x_)]*sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[a - c + (b - d)*x]/(2*(b - d)), x] - Simp[Sin[a + c + (b + d)*x]/(2*(b + d)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

3.43.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.53

method	result	size
derivativedivides	$\frac{(\sin^3(2x))}{3}$	9
default	$\frac{(\sin^3(2x))}{3}$	9
risch	$\frac{\sin(2x)}{4} - \frac{\sin(6x)}{12}$	14
parallelrisch	$\frac{\sin(2x)}{4} - \frac{\sin(6x)}{12}$	14
norman	$\frac{2 \tan(x) (\tan^2(2x))}{3} - \frac{(\tan^2(x)) \tan(2x)}{3} - \frac{2 \tan(x)}{3} + \frac{\tan(2x)}{3}$ $(1+\tan^2(x))(1+\tan^2(2x))$	51

input `int(sin(2*x)*sin(4*x),x,method=_RETURNVERBOSE)`output `1/3*sin(2*x)^3`**3.43.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sin(2x) \sin(4x) dx = -\frac{1}{3} (\cos(2x))^2 - 1) \sin(2x)$$

input `integrate(sin(2*x)*sin(4*x),x, algorithm="fricas")`output `-1/3*(cos(2*x)^2 - 1)*sin(2*x)`**3.43.6 Sympy [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.29

$$\int \sin(2x) \sin(4x) dx = -\frac{\sin(2x) \cos(4x)}{3} + \frac{\sin(4x) \cos(2x)}{6}$$

input `integrate(sin(2*x)*sin(4*x),x)`output `-sin(2*x)*cos(4*x)/3 + sin(4*x)*cos(2*x)/6`

3.43.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sin(2x) \sin(4x) dx = -\frac{1}{12} \sin(6x) + \frac{1}{4} \sin(2x)$$

input `integrate(sin(2*x)*sin(4*x),x, algorithm="maxima")`

output `-1/12*sin(6*x) + 1/4*sin(2*x)`

3.43.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.47

$$\int \sin(2x) \sin(4x) dx = \frac{1}{3} \sin(2x)^3$$

input `integrate(sin(2*x)*sin(4*x),x, algorithm="giac")`

output `1/3*sin(2*x)^3`

3.43.9 Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sin(2x) \sin(4x) dx = \frac{\sin(2x)}{4} - \frac{\sin(6x)}{12}$$

input `int(sin(2*x)*sin(4*x),x)`

output `sin(2*x)/4 - sin(6*x)/12`

3.44 $\int \cos(x) \log(\sin(x)) dx$

3.44.1	Optimal result	334
3.44.2	Mathematica [A] (verified)	334
3.44.3	Rubi [A] (verified)	335
3.44.4	Maple [A] (verified)	336
3.44.5	Fricas [A] (verification not implemented)	336
3.44.6	Sympy [A] (verification not implemented)	336
3.44.7	Maxima [A] (verification not implemented)	337
3.44.8	Giac [A] (verification not implemented)	337
3.44.9	Mupad [B] (verification not implemented)	337

3.44.1 Optimal result

Integrand size = 6, antiderivative size = 11

$$\int \cos(x) \log(\sin(x)) dx = -\sin(x) + \log(\sin(x)) \sin(x)$$

output `-sin(x)+ln(sin(x))*sin(x)`

3.44.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \cos(x) \log(\sin(x)) dx = -\sin(x) + \log(\sin(x)) \sin(x)$$

input `Integrate[Cos[x]*Log[Sin[x]],x]`

output `-Sin[x] + Log[Sin[x]]*Sin[x]`

3.44.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3034, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(x) \log(\sin(x)) dx \\
 & \quad \downarrow \text{3034} \\
 & \sin(x) \log(\sin(x)) - \int \cos(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \sin(x) \log(\sin(x)) - \int \sin\left(x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3117} \\
 & \sin(x) \log(\sin(x)) - \sin(x)
 \end{aligned}$$

input `Int[Cos[x]*Log[Sin[x]],x]`

output `-Sin[x] + Log[Sin[x]]*Sin[x]`

3.44.3.1 Defintions of rubi rules used

rule 3034 `Int[Log[u_]*(v_), x_Symbol] :=> With[{w = IntHide[v, x]}, Simp[Log[u] w, x] - Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w, x]] /; InverseFunctionFreeQ[u, x]`

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] :=> Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

3.44.4 Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

method	result
parallelrisch	$(\ln(\sin(x)) - 1) \sin(x)$
derivativdivides	$-\sin(x) + \ln(\sin(x)) \sin(x)$
default	$-\sin(x) + \ln(\sin(x)) \sin(x)$
norman	$\frac{2 \tan\left(\frac{x}{2}\right) \ln\left(\frac{2 \tan\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)}\right) - 2 \tan\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)}$
risch	$-\ln(e^{ix}) \sin(x) - \frac{e^{ix}\pi}{4} + \frac{e^{-ix}\pi}{4} - \frac{e^{-ix} \operatorname{csgn}(\sin(x))^2 \operatorname{csgn}(ie^{-ix})\pi}{4} - \frac{e^{-ix} \operatorname{csgn}(i \sin(x)) \operatorname{csgn}(\sin(x))\pi}{4}$

input `int(cos(x)*ln(sin(x)),x,method=_RETURNVERBOSE)`output `(ln(sin(x))-1)*sin(x)`**3.44.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \cos(x) \log(\sin(x)) dx = \log(\sin(x)) \sin(x) - \sin(x)$$

input `integrate(cos(x)*log(sin(x)),x, algorithm="fricas")`output `log(sin(x))*sin(x) - sin(x)`**3.44.6 Sympy [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int \cos(x) \log(\sin(x)) dx = \log(\sin(x)) \sin(x) - \sin(x)$$

input `integrate(cos(x)*ln(sin(x)),x)`output `log(sin(x))*sin(x) - sin(x)`

3.44.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \cos(x) \log(\sin(x)) dx = \log(\sin(x)) \sin(x) - \sin(x)$$

input `integrate(cos(x)*log(sin(x)),x, algorithm="maxima")`output `log(sin(x))*sin(x) - sin(x)`**3.44.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \cos(x) \log(\sin(x)) dx = \log(\sin(x)) \sin(x) - \sin(x)$$

input `integrate(cos(x)*log(sin(x)),x, algorithm="giac")`output `log(sin(x))*sin(x) - sin(x)`**3.44.9 Mupad [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \cos(x) \log(\sin(x)) dx = \sin(x) (\ln(\sin(x)) - 1)$$

input `int(log(sin(x))*cos(x),x)`output `sin(x)*(log(sin(x)) - 1)`

3.45 $\int e^{x^2} x^3 dx$

3.45.1	Optimal result	338
3.45.2	Mathematica [A] (verified)	338
3.45.3	Rubi [A] (verified)	339
3.45.4	Maple [A] (verified)	340
3.45.5	Fricas [A] (verification not implemented)	340
3.45.6	Sympy [A] (verification not implemented)	341
3.45.7	Maxima [A] (verification not implemented)	341
3.45.8	Giac [A] (verification not implemented)	341
3.45.9	Mupad [B] (verification not implemented)	342

3.45.1 Optimal result

Integrand size = 9, antiderivative size = 22

$$\int e^{x^2} x^3 dx = -\frac{e^{x^2}}{2} + \frac{1}{2}e^{x^2} x^2$$

output `-1/2*exp(x^2)+1/2*exp(x^2)*x^2`

3.45.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.64

$$\int e^{x^2} x^3 dx = \frac{1}{2}e^{x^2}(-1 + x^2)$$

input `Integrate[E^x^2*x^3,x]`

output `(E^x^2*(-1 + x^2))/2`

3.45.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {2641, 2638}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{x^2} x^3 dx$$

$$\downarrow \text{2641}$$

$$\frac{1}{2} e^{x^2} x^2 - \int e^{x^2} x dx$$

$$\downarrow \text{2638}$$

$$\frac{1}{2} e^{x^2} x^2 - \frac{e^{x^2}}{2}$$

input `Int[E^x^2*x^3,x]`

output `-1/2*E^x^2 + (E^x^2*x^2)/2`

3.45.3.1 Defintions of rubi rules used

rule 2638 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[(e + f*x)^n*(F^(a + b*(c + d*x)^n)/(b*f*n*(c + d*x)^n *Log[F])), x] /; FreeQ[{F, a, b, c, d, e, f, n}, x] && EqQ[m, n - 1] && EqQ[d*e - c*f, 0]`

rule 2641 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^(m - n + 1)*(F^(a + b*(c + d*x)^n)/(b*d*n*Log[F])), x] - Simp[(m - n + 1)/(b*n*Log[F]) Int[(c + d*x)^(m - n)*F^(a + b*(c + d*x)^n), x], x] /; FreeQ[{F, a, b, c, d}, x] && IntegerQ[2*((m + 1)/n)] && LtQ[0, (m + 1)/n, 5] && IntegerQ[n] && (LtQ[0, n, m + 1] || LtQ[m, n, 0])`

3.45.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.55

method	result	size
gospers	$\frac{(x^2-1)e^{x^2}}{2}$	12
risch	$\left(\frac{x^2}{2} - \frac{1}{2}\right) e^{x^2}$	13
meijerg	$\frac{1}{2} - \frac{(-2x^2+2)e^{x^2}}{4}$	16
derivativedivides	$-\frac{e^{x^2}}{2} + \frac{e^{x^2}x^2}{2}$	17
default	$-\frac{e^{x^2}}{2} + \frac{e^{x^2}x^2}{2}$	17
norman	$-\frac{e^{x^2}}{2} + \frac{e^{x^2}x^2}{2}$	17
parallelrisch	$-\frac{e^{x^2}}{2} + \frac{e^{x^2}x^2}{2}$	17
parts	$\frac{\operatorname{erfi}(x)\sqrt{\pi}x^3}{2} - \frac{3\sqrt{\pi}\left(\frac{x^3\operatorname{erfi}(x)}{3} - \frac{2\left(-\frac{e^{x^2}}{2} + \frac{e^{x^2}x^2}{2}\right)}{3\sqrt{\pi}}\right)}{2}$	46

input `int(x^3*exp(x^2),x,method=_RETURNVERBOSE)`output `1/2*(x^2-1)*exp(x^2)`**3.45.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.50

$$\int e^{x^2} x^3 dx = \frac{1}{2} (x^2 - 1) e^{(x^2)}$$

input `integrate(exp(x^2)*x^3,x, algorithm="fricas")`output `1/2*(x^2 - 1)*e^(x^2)`

3.45.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.45

$$\int e^{x^2} x^3 dx = \frac{(x^2 - 1) e^{x^2}}{2}$$

input `integrate(exp(x**2)*x**3,x)`output `(x**2 - 1)*exp(x**2)/2`**3.45.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.50

$$\int e^{x^2} x^3 dx = \frac{1}{2} (x^2 - 1) e^{(x^2)}$$

input `integrate(exp(x^2)*x^3,x, algorithm="maxima")`output `1/2*(x^2 - 1)*e^(x^2)`**3.45.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.50

$$\int e^{x^2} x^3 dx = \frac{1}{2} (x^2 - 1) e^{(x^2)}$$

input `integrate(exp(x^2)*x^3,x, algorithm="giac")`output `1/2*(x^2 - 1)*e^(x^2)`

3.45.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.50

$$\int e^{x^2} x^3 dx = \frac{e^{x^2} (x^2 - 1)}{2}$$

input `int(x^3*exp(x^2),x)`

output `(exp(x^2)*(x^2 - 1))/2`

3.46 $\int e^x(3 + 2x) dx$

3.46.1	Optimal result	343
3.46.2	Mathematica [A] (verified)	343
3.46.3	Rubi [A] (verified)	344
3.46.4	Maple [A] (verified)	345
3.46.5	Fricas [A] (verification not implemented)	345
3.46.6	Sympy [A] (verification not implemented)	345
3.46.7	Maxima [A] (verification not implemented)	346
3.46.8	Giac [A] (verification not implemented)	346
3.46.9	Mupad [B] (verification not implemented)	346

3.46.1 Optimal result

Integrand size = 9, antiderivative size = 15

$$\int e^x(3 + 2x) dx = -2e^x + e^x(3 + 2x)$$

output `-2*exp(x)+exp(x)*(3+2*x)`

3.46.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.60

$$\int e^x(3 + 2x) dx = e^x(1 + 2x)$$

input `Integrate[E^x*(3 + 2*x),x]`

output `E^x*(1 + 2*x)`

3.46.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int e^x(2x + 3) dx \\ \downarrow \text{2607} \\ e^x(2x + 3) - 2 \int e^x dx \\ \downarrow \text{2624} \\ e^x(2x + 3) - 2e^x \end{array}$$

input `Int[E^x*(3 + 2*x),x]`

output `-2*E^x + E^x*(3 + 2*x)`

3.46.3.1 Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2624 `Int[((F_)^(v_))^(n_.), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`

3.46.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.60

method	result	size
gospers	$(1 + 2x) e^x$	9
default	$e^x + 2 e^x x$	9
norman	$e^x + 2 e^x x$	9
risch	$(1 + 2x) e^x$	9
parallelrisch	$e^x + 2 e^x x$	9
parts	$e^x + 2 e^x x$	9
meijerg	$-1 + 3 e^x - (-2x + 2) e^x$	16

input `int(exp(x)*(3+2*x),x,method=_RETURNVERBOSE)`output `(1+2*x)*exp(x)`**3.46.5 Fracas [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.53

$$\int e^x(3 + 2x) dx = (2x + 1)e^x$$

input `integrate(exp(x)*(3+2*x),x, algorithm="fricas")`output `(2*x + 1)*e^x`**3.46.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.47

$$\int e^x(3 + 2x) dx = (2x + 1)e^x$$

input `integrate(exp(x)*(3+2*x),x)`output `(2*x + 1)*exp(x)`

3.46.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int e^x(3 + 2x) dx = 2(x - 1)e^x + 3e^x$$

input `integrate(exp(x)*(3+2*x),x, algorithm="maxima")`

output `2*(x - 1)*e^x + 3*e^x`

3.46.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.53

$$\int e^x(3 + 2x) dx = (2x + 1)e^x$$

input `integrate(exp(x)*(3+2*x),x, algorithm="giac")`

output `(2*x + 1)*e^x`

3.46.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.53

$$\int e^x(3 + 2x) dx = e^x(2x + 1)$$

input `int(exp(x)*(2*x + 3),x)`

output `exp(x)*(2*x + 1)`

3.47 $\int 5^x x dx$

3.47.1	Optimal result	347
3.47.2	Mathematica [A] (verified)	347
3.47.3	Rubi [A] (verified)	348
3.47.4	Maple [A] (verified)	349
3.47.5	Fricas [A] (verification not implemented)	349
3.47.6	Sympy [A] (verification not implemented)	349
3.47.7	Maxima [A] (verification not implemented)	350
3.47.8	Giac [A] (verification not implemented)	350
3.47.9	Mupad [B] (verification not implemented)	350

3.47.1 Optimal result

Integrand size = 5, antiderivative size = 19

$$\int 5^x x dx = -\frac{5^x}{\log^2(5)} + \frac{5^x x}{\log(5)}$$

output `-5^x/ln(5)^2+5^x*x/ln(5)`

3.47.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int 5^x x dx = \frac{5^x(-1 + x \log(5))}{\log^2(5)}$$

input `Integrate[5^x*x,x]`

output `(5^x*(-1 + x*Log[5]))/Log[5]^2`

3.47.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int 5^x x dx$$

$$\downarrow \text{2607}$$

$$\frac{5^x x}{\log(5)} - \frac{\int 5^x dx}{\log(5)}$$

$$\downarrow \text{2624}$$

$$\frac{5^x x}{\log(5)} - \frac{5^x}{\log^2(5)}$$

input `Int[5^x*x,x]`

output `-(5^x/Log[5]^2) + (5^x*x)/Log[5]`

3.47.3.1 Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`

3.47.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

method	result	size
gospers	$\frac{(x \ln(5) - 1)5^x}{\ln(5)^2}$	15
risch	$\frac{(x \ln(5) - 1)5^x}{\ln(5)^2}$	15
parallelrisch	$\frac{5^x \ln(5)x - 5^x}{\ln(5)^2}$	19
meijerg	$\frac{1 - \frac{(2 - 2x \ln(5))e^{x \ln(5)}}{2}}{\ln(5)^2}$	22
norman	$\frac{x e^{x \ln(5)}}{\ln(5)} - \frac{e^{x \ln(5)}}{\ln(5)^2}$	24

input `int(5^x*x,x,method=_RETURNVERBOSE)`output `(x*ln(5)-1)*5^x/ln(5)^2`**3.47.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int 5^x x dx = \frac{(x \log(5) - 1)5^x}{\log(5)^2}$$

input `integrate(5^x*x,x, algorithm="fricas")`output `(x*log(5) - 1)*5^x/log(5)^2`**3.47.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int 5^x x dx = \frac{5^x(x \log(5) - 1)}{\log(5)^2}$$

input `integrate(5**x*x,x)`

output `5**x*(x*log(5) - 1)/log(5)**2`

3.47.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int 5^x x dx = \frac{(x \log(5) - 1)5^x}{\log(5)^2}$$

input `integrate(5^x*x,x, algorithm="maxima")`

output `(x*log(5) - 1)*5^x/log(5)^2`

3.47.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int 5^x x dx = \frac{(x \log(5) - 1)5^x}{\log(5)^2}$$

input `integrate(5^x*x,x, algorithm="giac")`

output `(x*log(5) - 1)*5^x/log(5)^2`

3.47.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int 5^x x dx = \frac{5^x (x \ln(5) - 1)}{\ln(5)^2}$$

input `int(5^x*x,x)`

output `(5^x*(x*log(5) - 1))/log(5)^2`

3.48 $\int \cos(\log(x)) dx$

3.48.1	Optimal result	351
3.48.2	Mathematica [A] (verified)	351
3.48.3	Rubi [A] (verified)	352
3.48.4	Maple [A] (verified)	352
3.48.5	Fricas [A] (verification not implemented)	353
3.48.6	Sympy [A] (verification not implemented)	353
3.48.7	Maxima [A] (verification not implemented)	353
3.48.8	Giac [A] (verification not implemented)	354
3.48.9	Mupad [B] (verification not implemented)	354

3.48.1 Optimal result

Integrand size = 3, antiderivative size = 17

$$\int \cos(\log(x)) dx = \frac{1}{2}x \cos(\log(x)) + \frac{1}{2}x \sin(\log(x))$$

output `1/2*x*cos(ln(x))+1/2*x*sin(ln(x))`

3.48.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cos(\log(x)) dx = \frac{1}{2}x \cos(\log(x)) + \frac{1}{2}x \sin(\log(x))$$

input `Integrate[Cos[Log[x]],x]`

output `(x*Cos[Log[x]])/2 + (x*Sin[Log[x]])/2`

3.48.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {4979}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(\log(x)) dx$$

$$\downarrow 4979$$

$$\frac{1}{2}x \sin(\log(x)) + \frac{1}{2}x \cos(\log(x))$$

input `Int[Cos[Log[x]],x]`

output `(x*Cos[Log[x]])/2 + (x*Sin[Log[x]])/2`

3.48.3.1 Defintions of rubi rules used

rule 4979 `Int[Cos[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)], x_Symbol] := Simp[x*(Cos[d*(a + b*Log[c*x^n])]/(b^2*d^2*n^2 + 1)), x] + Simp[b*d*n*x*(Sin[d*(a + b*Log[c*x^n])]/(b^2*d^2*n^2 + 1)), x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b^2*d^2*n^2 + 1, 0]`

3.48.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

method	result	size
parallelrisch	$\frac{x(\cos(\ln(x)) + \sin(\ln(x)))}{2}$	11
lookup	$\frac{x \cos(\ln(x))}{2} + \frac{x \sin(\ln(x))}{2}$	14
default	$\frac{x \cos(\ln(x))}{2} + \frac{x \sin(\ln(x))}{2}$	14
risch	$(\frac{1}{4} - \frac{i}{4}) x x^i + (\frac{1}{4} + \frac{i}{4}) x x^{-i}$	22

input `int(cos(ln(x)),x,method=_RETURNVERBOSE)`

output `1/2*x*(cos(ln(x))+sin(ln(x)))`

3.48.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(\log(x)) dx = \frac{1}{2} x \cos(\log(x)) + \frac{1}{2} x \sin(\log(x))$$

input `integrate(cos(log(x)),x, algorithm="fricas")`

output `1/2*x*cos(log(x)) + 1/2*x*sin(log(x))`

3.48.6 Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \cos(\log(x)) dx = \frac{x \sin(\log(x))}{2} + \frac{x \cos(\log(x))}{2}$$

input `integrate(cos(ln(x)),x)`

output `x*sin(log(x))/2 + x*cos(log(x))/2`

3.48.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.59

$$\int \cos(\log(x)) dx = \frac{1}{2} x(\cos(\log(x)) + \sin(\log(x)))$$

input `integrate(cos(log(x)),x, algorithm="maxima")`

output `1/2*x*(cos(log(x)) + sin(log(x)))`

3.48.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(\log(x)) dx = \frac{1}{2} x \cos(\log(x)) + \frac{1}{2} x \sin(\log(x))$$

input `integrate(cos(log(x)),x, algorithm="giac")`

output `1/2*x*cos(log(x)) + 1/2*x*sin(log(x))`

3.48.9 Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(\log(x)) dx = \frac{\sqrt{2} x \sin\left(\frac{\pi}{4} + \ln(x)\right)}{2}$$

input `int(cos(log(x)),x)`

output `(2^(1/2)*x*sin(pi/4 + log(x)))/2`

3.49 $\int e^{\sqrt{x}} dx$

3.49.1	Optimal result	355
3.49.2	Mathematica [A] (verified)	355
3.49.3	Rubi [A] (verified)	356
3.49.4	Maple [A] (verified)	357
3.49.5	Fricas [A] (verification not implemented)	357
3.49.6	Sympy [A] (verification not implemented)	358
3.49.7	Maxima [A] (verification not implemented)	358
3.49.8	Giac [A] (verification not implemented)	358
3.49.9	Mupad [B] (verification not implemented)	359

3.49.1 Optimal result

Integrand size = 7, antiderivative size = 24

$$\int e^{\sqrt{x}} dx = -2e^{\sqrt{x}} + 2e^{\sqrt{x}}\sqrt{x}$$

output `-2*exp(x^(1/2))+2*exp(x^(1/2))*x^(1/2)`

3.49.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int e^{\sqrt{x}} dx = 2e^{\sqrt{x}}(-1 + \sqrt{x})$$

input `Integrate[E^Sqrt[x],x]`

output `2*E^Sqrt[x]*(-1 + Sqrt[x])`

3.49.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.04, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {2636, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int e^{\sqrt{x}} dx \\
 \downarrow \text{2636} \\
 2 \int e^{\sqrt{x}} \sqrt{x} d\sqrt{x} \\
 \downarrow \text{2607} \\
 2 \left(e^{\sqrt{x}} \sqrt{x} - \int e^{\sqrt{x}} d\sqrt{x} \right) \\
 \downarrow \text{2624} \\
 2 \left(e^{\sqrt{x}} \sqrt{x} - e^{\sqrt{x}} \right)
 \end{array}$$

input `Int[E^Sqrt[x], x]`

output `2*(-E^Sqrt[x] + E^Sqrt[x]*Sqrt[x])`

3.49.3.1 Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`

```
rule 2636 Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] :> With[{k =
Denominator[n]}, Simp[k/d Subst[Int[x^(k - 1)*F^(a + b*x^(k*n)), x], x, (
c + d*x)^(1/k)], x]] /; FreeQ[{F, a, b, c, d}, x] && IntegerQ[2/n] && !Int
egerQ[n]
```

3.49.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

method	result	size
meijerg	$2 - (-2\sqrt{x} + 2)e^{\sqrt{x}}$	16
derivativedivides	$-2e^{\sqrt{x}} + 2e^{\sqrt{x}}\sqrt{x}$	17
default	$-2e^{\sqrt{x}} + 2e^{\sqrt{x}}\sqrt{x}$	17

```
input int(exp(x^(1/2)),x,method=_RETURNVERBOSE)
```

```
output 2-(-2*x^(1/2)+2)*exp(x^(1/2))
```

3.49.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46

$$\int e^{\sqrt{x}} dx = 2(\sqrt{x} - 1)e^{\sqrt{x}}$$

```
input integrate(exp(x^(1/2)),x, algorithm="fracas")
```

```
output 2*(sqrt(x) - 1)*e^sqrt(x)
```

3.49.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int e^{\sqrt{x}} dx = 2\sqrt{x}e^{\sqrt{x}} - 2e^{\sqrt{x}}$$

input `integrate(exp(x**(1/2)),x)`output `2*sqrt(x)*exp(sqrt(x)) - 2*exp(sqrt(x))`**3.49.7 Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46

$$\int e^{\sqrt{x}} dx = 2(\sqrt{x} - 1)e^{\sqrt{x}}$$

input `integrate(exp(x^(1/2)),x, algorithm="maxima")`output `2*(sqrt(x) - 1)*e^sqrt(x)`**3.49.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46

$$\int e^{\sqrt{x}} dx = 2(\sqrt{x} - 1)e^{\sqrt{x}}$$

input `integrate(exp(x^(1/2)),x, algorithm="giac")`output `2*(sqrt(x) - 1)*e^sqrt(x)`

3.49.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46

$$\int e^{\sqrt{x}} dx = 2e^{\sqrt{x}}(\sqrt{x} - 1)$$

input `int(exp(x^(1/2)),x)`

output `2*exp(x^(1/2))*(x^(1/2) - 1)`

3.50 $\int \log(\sqrt{x}) dx$

3.50.1	Optimal result	360
3.50.2	Mathematica [A] (verified)	360
3.50.3	Rubi [A] (verified)	361
3.50.4	Maple [A] (verified)	361
3.50.5	Fricas [A] (verification not implemented)	362
3.50.6	Sympy [A] (verification not implemented)	362
3.50.7	Maxima [A] (verification not implemented)	362
3.50.8	Giac [A] (verification not implemented)	363
3.50.9	Mupad [B] (verification not implemented)	363

3.50.1 Optimal result

Integrand size = 6, antiderivative size = 14

$$\int \log(\sqrt{x}) dx = -\frac{x}{2} + x \log(\sqrt{x})$$

output `-1/2*x+1/2*x*ln(x)`

3.50.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \log(\sqrt{x}) dx = \frac{1}{2}(-x + x \log(x))$$

input `Integrate[Log[Sqrt[x]],x]`

output `(-x + x*Log[x])/2`

3.50.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2732}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \log(\sqrt{x}) dx$$

$$\downarrow 2732$$

$$x \log(\sqrt{x}) - \frac{x}{2}$$

input `Int[Log[Sqrt[x]], x]`

output `-1/2*x + x*Log[Sqrt[x]]`

3.50.3.1 Defintions of rubi rules used

rule 2732 `Int[Log[(c_.)*(x_)^(n_.)], x_Symbol] :> Simp[x*Log[c*x^n], x] - Simp[n*x, x] /; FreeQ[{c, n}, x]`

3.50.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

method	result	size
lookup	$-\frac{x}{2} + \frac{x \ln(x)}{2}$	10
default	$-\frac{x}{2} + \frac{x \ln(x)}{2}$	10
norman	$-\frac{x}{2} + \frac{x \ln(x)}{2}$	10
risch	$-\frac{x}{2} + \frac{x \ln(x)}{2}$	10
parallelrisch	$-\frac{x}{2} + \frac{x \ln(x)}{2}$	10
parts	$-\frac{x}{2} + \frac{x \ln(x)}{2}$	10

input `int(1/2*ln(x), x, method=_RETURNVERBOSE)`

output `-1/2*x+1/2*x*ln(x)`

3.50.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.64

$$\int \log(\sqrt{x}) dx = \frac{1}{2} x \log(x) - \frac{1}{2} x$$

input `integrate(1/2*log(x),x, algorithm="fricas")`

output `1/2*x*log(x) - 1/2*x`

3.50.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.57

$$\int \log(\sqrt{x}) dx = \frac{x \log(x)}{2} - \frac{x}{2}$$

input `integrate(1/2*ln(x),x)`

output `x*log(x)/2 - x/2`

3.50.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.64

$$\int \log(\sqrt{x}) dx = \frac{1}{2} x \log(x) - \frac{1}{2} x$$

input `integrate(1/2*log(x),x, algorithm="maxima")`

output `1/2*x*log(x) - 1/2*x`

3.50.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.64

$$\int \log(\sqrt{x}) dx = \frac{1}{2} x \log(x) - \frac{1}{2} x$$

input `integrate(1/2*log(x),x, algorithm="giac")`

output `1/2*x*log(x) - 1/2*x`

3.50.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.50

$$\int \log(\sqrt{x}) dx = \frac{x(\ln(x) - 1)}{2}$$

input `int(log(x)/2,x)`

output `(x*(log(x) - 1))/2`

3.51 $\int \sin(\log(x)) dx$

3.51.1	Optimal result	364
3.51.2	Mathematica [A] (verified)	364
3.51.3	Rubi [A] (verified)	365
3.51.4	Maple [A] (verified)	365
3.51.5	Fricas [A] (verification not implemented)	366
3.51.6	Sympy [A] (verification not implemented)	366
3.51.7	Maxima [A] (verification not implemented)	366
3.51.8	Giac [A] (verification not implemented)	367
3.51.9	Mupad [B] (verification not implemented)	367

3.51.1 Optimal result

Integrand size = 3, antiderivative size = 17

$$\int \sin(\log(x)) dx = -\frac{1}{2}x \cos(\log(x)) + \frac{1}{2}x \sin(\log(x))$$

output `-1/2*x*cos(ln(x))+1/2*x*sin(ln(x))`

3.51.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sin(\log(x)) dx = -\frac{1}{2}x \cos(\log(x)) + \frac{1}{2}x \sin(\log(x))$$

input `Integrate[Sin[Log[x]],x]`

output `-1/2*(x*Cos[Log[x]]) + (x*Sin[Log[x]])/2`

3.51.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {4978}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sin(\log(x)) dx$$

$$\downarrow 4978$$

$$\frac{1}{2}x \sin(\log(x)) - \frac{1}{2}x \cos(\log(x))$$

input `Int[Sin[Log[x]],x]`

output `-1/2*(x*Cos[Log[x]]) + (x*Sin[Log[x]])/2`

3.51.3.1 Defintions of rubi rules used

rule 4978 `Int[Sin[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)], x_Symbol] := Simp[x*(Sin[d*(a + b*Log[c*x^n])]/(b^2*d^2*n^2 + 1)), x] - Simp[b*d*n*x*(Cos[d*(a + b*Log[c*x^n])]/(b^2*d^2*n^2 + 1)), x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b^2*d^2*n^2 + 1, 0]`

3.51.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

method	result	size
parallelrisch	$-\frac{x(\cos(\ln(x)) - \sin(\ln(x)))}{2}$	13
lookup	$-\frac{x \cos(\ln(x))}{2} + \frac{x \sin(\ln(x))}{2}$	14
default	$-\frac{x \cos(\ln(x))}{2} + \frac{x \sin(\ln(x))}{2}$	14
risch	$\left(-\frac{1}{4} - \frac{i}{4}\right) x x^i + \left(-\frac{1}{4} + \frac{i}{4}\right) x x^{-i}$	22
norman	$\frac{x \tan\left(\frac{\ln(x)}{2}\right) - \frac{x}{2} + \frac{x \left(\tan^2\left(\frac{\ln(x)}{2}\right)\right)}{2}}{1 + \tan^2\left(\frac{\ln(x)}{2}\right)}$	34

input `int(sin(ln(x)),x,method=_RETURNVERBOSE)`

output `-1/2*x*(cos(ln(x))-sin(ln(x)))`

3.51.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sin(\log(x)) dx = -\frac{1}{2} x \cos(\log(x)) + \frac{1}{2} x \sin(\log(x))$$

input `integrate(sin(log(x)),x, algorithm="fricas")`

output `-1/2*x*cos(log(x)) + 1/2*x*sin(log(x))`

3.51.6 Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \sin(\log(x)) dx = \frac{x \sin(\log(x))}{2} - \frac{x \cos(\log(x))}{2}$$

input `integrate(sin(ln(x)),x)`

output `x*sin(log(x))/2 - x*cos(log(x))/2`

3.51.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int \sin(\log(x)) dx = -\frac{1}{2} x (\cos(\log(x)) - \sin(\log(x)))$$

input `integrate(sin(log(x)),x, algorithm="maxima")`

output `-1/2*x*(cos(log(x)) - sin(log(x)))`

3.51.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sin(\log(x)) dx = -\frac{1}{2} x \cos(\log(x)) + \frac{1}{2} x \sin(\log(x))$$

input `integrate(sin(log(x)),x, algorithm="giac")`output `-1/2*x*cos(log(x)) + 1/2*x*sin(log(x))`**3.51.9 Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sin(\log(x)) dx = -\frac{\sqrt{2} x \cos\left(\frac{\pi}{4} + \ln(x)\right)}{2}$$

input `int(sin(log(x)),x)`output `-(2^(1/2)*x*cos(pi/4 + log(x)))/2`

3.52 $\int \sin(\sqrt{x}) dx$

3.52.1	Optimal result	368
3.52.2	Mathematica [A] (verified)	368
3.52.3	Rubi [A] (verified)	369
3.52.4	Maple [A] (verified)	370
3.52.5	Fricas [A] (verification not implemented)	371
3.52.6	Sympy [A] (verification not implemented)	371
3.52.7	Maxima [A] (verification not implemented)	371
3.52.8	Giac [A] (verification not implemented)	372
3.52.9	Mupad [B] (verification not implemented)	372

3.52.1 Optimal result

Integrand size = 6, antiderivative size = 22

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

output `2*sin(x^(1/2))-2*cos(x^(1/2))*x^(1/2)`

3.52.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

input `Integrate[Sin[Sqrt[x]],x]`

output `-2*Sqrt[x]*Cos[Sqrt[x]] + 2*Sin[Sqrt[x]]`

3.52.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {3842, 3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(\sqrt{x}) \, dx \\
 & \quad \downarrow \text{3842} \\
 & 2 \int \sqrt{x} \sin(\sqrt{x}) \, d\sqrt{x} \\
 & \quad \downarrow \text{3042} \\
 & 2 \int \sqrt{x} \sin(\sqrt{x}) \, d\sqrt{x} \\
 & \quad \downarrow \text{3777} \\
 & 2 \left(\int \cos(\sqrt{x}) \, d\sqrt{x} - \sqrt{x} \cos(\sqrt{x}) \right) \\
 & \quad \downarrow \text{3042} \\
 & 2 \left(\int \sin\left(\sqrt{x} + \frac{\pi}{2}\right) \, d\sqrt{x} - \sqrt{x} \cos(\sqrt{x}) \right) \\
 & \quad \downarrow \text{3117} \\
 & 2(\sin(\sqrt{x}) - \sqrt{x} \cos(\sqrt{x}))
 \end{aligned}$$

input `Int[Sin[Sqrt[x]], x]`

output `2*(-(Sqrt[x]*Cos[Sqrt[x]]) + Sin[Sqrt[x]])`

3.52.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3842 `Int[((a_.) + (b_.)*Sin[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)])^(p_.), x_Symbol] := Simp[1/(n*f) Subst[Int[x^(1/n - 1)*(a + b*SIN[c + d*x])^p, x], (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]`

3.52.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

method	result	size
derivativedivides	$2 \sin(\sqrt{x}) - 2 \cos(\sqrt{x}) \sqrt{x}$	17
default	$2 \sin(\sqrt{x}) - 2 \cos(\sqrt{x}) \sqrt{x}$	17
meijerg	$4\sqrt{\pi} \left(-\frac{\sqrt{x} \cos(\sqrt{x})}{2\sqrt{\pi}} + \frac{\sin(\sqrt{x})}{2\sqrt{\pi}} \right)$	28

input `int(sin(x^(1/2)),x,method=_RETURNVERBOSE)`

output `2*sin(x^(1/2))-2*cos(x^(1/2))*x^(1/2)`

3.52.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

input `integrate(sin(x^(1/2)),x, algorithm="fricas")`

output `-2*sqrt(x)*cos(sqrt(x)) + 2*sin(sqrt(x))`

3.52.6 Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

input `integrate(sin(x**(1/2)),x)`

output `-2*sqrt(x)*cos(sqrt(x)) + 2*sin(sqrt(x))`

3.52.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

input `integrate(sin(x^(1/2)),x, algorithm="maxima")`

output `-2*sqrt(x)*cos(sqrt(x)) + 2*sin(sqrt(x))`

3.52.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

input `integrate(sin(x^(1/2)),x, algorithm="giac")`

output `-2*sqrt(x)*cos(sqrt(x)) + 2*sin(sqrt(x))`

3.52.9 Mupad [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \sin(\sqrt{x}) dx = 2 \sin(\sqrt{x}) - 2\sqrt{x} \cos(\sqrt{x})$$

input `int(sin(x^(1/2)),x)`

output `2*sin(x^(1/2)) - 2*x^(1/2)*cos(x^(1/2))`

3.53 $\int x^5 \cos(x^3) dx$

3.53.1	Optimal result	373
3.53.2	Mathematica [A] (verified)	373
3.53.3	Rubi [A] (verified)	374
3.53.4	Maple [A] (verified)	375
3.53.5	Fricas [A] (verification not implemented)	376
3.53.6	Sympy [A] (verification not implemented)	376
3.53.7	Maxima [A] (verification not implemented)	376
3.53.8	Giac [A] (verification not implemented)	377
3.53.9	Mupad [B] (verification not implemented)	377

3.53.1 Optimal result

Integrand size = 8, antiderivative size = 20

$$\int x^5 \cos(x^3) dx = \frac{\cos(x^3)}{3} + \frac{1}{3}x^3 \sin(x^3)$$

output `1/3*cos(x^3)+1/3*x^3*sin(x^3)`

3.53.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int x^5 \cos(x^3) dx = \frac{\cos(x^3)}{3} + \frac{1}{3}x^3 \sin(x^3)$$

input `Integrate[x^5*Cos[x^3],x]`

output `Cos[x^3]/3 + (x^3*Sin[x^3])/3`

3.53.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3861, 3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^5 \cos(x^3) dx \\
 & \quad \downarrow \text{3861} \\
 & \frac{1}{3} \int x^3 \cos(x^3) dx^3 \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int x^3 \sin\left(x^3 + \frac{\pi}{2}\right) dx^3 \\
 & \quad \downarrow \text{3777} \\
 & \frac{1}{3} \left(\int -\sin(x^3) dx^3 + x^3 \sin(x^3) \right) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{3} \left(x^3 \sin(x^3) - \int \sin(x^3) dx^3 \right) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \left(x^3 \sin(x^3) - \int \sin(x^3) dx^3 \right) \\
 & \quad \downarrow \text{3118} \\
 & \frac{1}{3} (x^3 \sin(x^3) + \cos(x^3))
 \end{aligned}$$

input `Int[x^5*Cos[x^3],x]`

output `(Cos[x^3] + x^3*Sin[x^3])/3`

3.53.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`
- rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`
- rule 3861 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))`

3.53.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

method	result	size
derivativedivides	$\frac{\cos(x^3)}{3} + \frac{x^3 \sin(x^3)}{3}$	17
default	$\frac{\cos(x^3)}{3} + \frac{x^3 \sin(x^3)}{3}$	17
risch	$\frac{\cos(x^3)}{3} + \frac{x^3 \sin(x^3)}{3}$	17
norman	$\frac{2x^3 \tan\left(\frac{x^3}{2}\right)}{3} + \frac{2}{3} \frac{1}{1 + \tan^2\left(\frac{x^3}{2}\right)}$	27
parallelrisch	$\frac{2x^3 \tan\left(\frac{x^3}{2}\right) + 2}{3\left(\tan^2\left(\frac{x^3}{2}\right) + 3\right)}$	29
meijerg	$\frac{2\sqrt{\pi} \left(-\frac{1}{2\sqrt{\pi}} + \frac{\cos(x^3)}{2\sqrt{\pi}} + \frac{x^3 \sin(x^3)}{2\sqrt{\pi}} \right)}{3}$	33

input `int(x^5*cos(x^3),x,method=_RETURNVERBOSE)`

output `1/3*cos(x^3)+1/3*x^3*sin(x^3)`

3.53.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int x^5 \cos(x^3) dx = \frac{1}{3} x^3 \sin(x^3) + \frac{1}{3} \cos(x^3)$$

input `integrate(x^5*cos(x^3),x, algorithm="fricas")`

output `1/3*x^3*sin(x^3) + 1/3*cos(x^3)`

3.53.6 Sympy [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int x^5 \cos(x^3) dx = \frac{x^3 \sin(x^3)}{3} + \frac{\cos(x^3)}{3}$$

input `integrate(x**5*cos(x**3),x)`

output `x**3*sin(x**3)/3 + cos(x**3)/3`

3.53.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int x^5 \cos(x^3) dx = \frac{1}{3} x^3 \sin(x^3) + \frac{1}{3} \cos(x^3)$$

input `integrate(x^5*cos(x^3),x, algorithm="maxima")`

output `1/3*x^3*sin(x^3) + 1/3*cos(x^3)`

3.53.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int x^5 \cos(x^3) dx = \frac{1}{3} x^3 \sin(x^3) + \frac{1}{3} \cos(x^3)$$

input `integrate(x^5*cos(x^3),x, algorithm="giac")`

output `1/3*x^3*sin(x^3) + 1/3*cos(x^3)`

3.53.9 Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int x^5 \cos(x^3) dx = \frac{\cos(x^3)}{3} + \frac{x^3 \sin(x^3)}{3}$$

input `int(x^5*cos(x^3),x)`

output `cos(x^3)/3 + (x^3*sin(x^3))/3`

3.54 $\int e^{x^2} x^5 dx$

3.54.1	Optimal result	378
3.54.2	Mathematica [A] (verified)	378
3.54.3	Rubi [A] (verified)	379
3.54.4	Maple [A] (verified)	380
3.54.5	Fricas [A] (verification not implemented)	381
3.54.6	Sympy [A] (verification not implemented)	381
3.54.7	Maxima [A] (verification not implemented)	381
3.54.8	Giac [A] (verification not implemented)	382
3.54.9	Mupad [B] (verification not implemented)	382

3.54.1 Optimal result

Integrand size = 9, antiderivative size = 28

$$\int e^{x^2} x^5 dx = e^{x^2} - e^{x^2} x^2 + \frac{1}{2} e^{x^2} x^4$$

output `exp(x^2)-exp(x^2)*x^2+1/2*exp(x^2)*x^4`

3.54.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.68

$$\int e^{x^2} x^5 dx = \frac{1}{2} e^{x^2} (2 - 2x^2 + x^4)$$

input `Integrate[E^x^2*x^5,x]`

output `(E^x^2*(2 - 2*x^2 + x^4))/2`

3.54.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.32, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2641, 2641, 2638}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^{x^2} x^5 dx \\ & \quad \downarrow \text{2641} \\ & \frac{1}{2} e^{x^2} x^4 - 2 \int e^{x^2} x^3 dx \\ & \quad \downarrow \text{2641} \\ & \frac{1}{2} e^{x^2} x^4 - 2 \left(\frac{1}{2} e^{x^2} x^2 - \int e^{x^2} x dx \right) \\ & \quad \downarrow \text{2638} \\ & \frac{1}{2} e^{x^2} x^4 - 2 \left(\frac{1}{2} e^{x^2} x^2 - \frac{e^{x^2}}{2} \right) \end{aligned}$$

input `Int [E^x^2*x^5, x]`

output `(E^x^2*x^4)/2 - 2*(-1/2*E^x^2 + (E^x^2*x^2)/2)`

3.54.3.1 Defintions of rubi rules used

rule 2638 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^n*(F^(a + b*(c + d*x)^n)/(b*f*n*(c + d*x)^n *Log[F])), x] /; FreeQ[{F, a, b, c, d, e, f, n}, x] && EqQ[m, n - 1] && EqQ[d*e - c*f, 0]`

rule 2641 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m - n + 1)*(F^(a + b*(c + d*x)^n)/(b*d*n*Log[F])), x] - Simp[(m - n + 1)/(b*n*Log[F]) Int[(c + d*x)^(m - n)*F^(a + b*(c + d*x)^n), x], x] /; FreeQ[{F, a, b, c, d}, x] && IntegerQ[2*((m + 1)/n)] && LtQ[0, (m + 1)/n, 5] && IntegerQ[n] && (LtQ[0, n, m + 1] || LtQ[m, n, 0])`

3.54.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.61

method	result	size
gospers	$\frac{(x^4 - 2x^2 + 2)e^{x^2}}{2}$	17
risch	$\left(\frac{1}{2}x^4 - x^2 + 1\right)e^{x^2}$	18
meijerg	$-1 + \frac{(3x^4 - 6x^2 + 6)e^{x^2}}{6}$	21
default	$e^{x^2} - e^{x^2}x^2 + \frac{e^{x^2}x^4}{2}$	24
norman	$e^{x^2} - e^{x^2}x^2 + \frac{e^{x^2}x^4}{2}$	24
parallelrisch	$e^{x^2} - e^{x^2}x^2 + \frac{e^{x^2}x^4}{2}$	24
parts	$\frac{\operatorname{erfi}(x)\sqrt{\pi}x^5}{2} - \frac{5\sqrt{\pi} \left(\frac{x^5 \operatorname{erfi}(x)}{5} - \frac{2(e^{x^2} - e^{x^2}x^2 + \frac{e^{x^2}x^4}{2})}{5\sqrt{\pi}} \right)}{2}$	53

input `int(exp(x^2)*x^5,x,method=_RETURNVERBOSE)`

output `1/2*(x^4-2*x^2+2)*exp(x^2)`

3.54.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.57

$$\int e^{x^2} x^5 dx = \frac{1}{2} (x^4 - 2x^2 + 2) e^{(x^2)}$$

input `integrate(exp(x^2)*x^5,x, algorithm="fricas")`output `1/2*(x^4 - 2*x^2 + 2)*e^(x^2)`**3.54.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.54

$$\int e^{x^2} x^5 dx = \frac{(x^4 - 2x^2 + 2) e^{x^2}}{2}$$

input `integrate(exp(x**2)*x**5,x)`output `(x**4 - 2*x**2 + 2)*exp(x**2)/2`**3.54.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.57

$$\int e^{x^2} x^5 dx = \frac{1}{2} (x^4 - 2x^2 + 2) e^{(x^2)}$$

input `integrate(exp(x^2)*x^5,x, algorithm="maxima")`output `1/2*(x^4 - 2*x^2 + 2)*e^(x^2)`

3.54.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.57

$$\int e^{x^2} x^5 dx = \frac{1}{2} (x^4 - 2x^2 + 2)e^{(x^2)}$$

input `integrate(exp(x^2)*x^5,x, algorithm="giac")`

output `1/2*(x^4 - 2*x^2 + 2)*e^(x^2)`

3.54.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.57

$$\int e^{x^2} x^5 dx = \frac{e^{x^2} (x^4 - 2x^2 + 2)}{2}$$

input `int(x^5*exp(x^2),x)`

output `(exp(x^2)*(x^4 - 2*x^2 + 2))/2`

3.55 $\int x \arctan(x) dx$

3.55.1	Optimal result	383
3.55.2	Mathematica [A] (verified)	383
3.55.3	Rubi [A] (verified)	384
3.55.4	Maple [A] (verified)	385
3.55.5	Fricas [A] (verification not implemented)	385
3.55.6	Sympy [A] (verification not implemented)	386
3.55.7	Maxima [A] (verification not implemented)	386
3.55.8	Giac [A] (verification not implemented)	386
3.55.9	Mupad [B] (verification not implemented)	387

3.55.1 Optimal result

Integrand size = 4, antiderivative size = 21

$$\int x \arctan(x) dx = -\frac{x}{2} + \frac{\arctan(x)}{2} + \frac{1}{2}x^2 \arctan(x)$$

output `-1/2*x+1/2*arctan(x)+1/2*x^2*arctan(x)`

3.55.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.76

$$\int x \arctan(x) dx = \frac{1}{2}(-x + (1 + x^2) \arctan(x))$$

input `Integrate[x*ArcTan[x],x]`

output `(-x + (1 + x^2)*ArcTan[x])/2`

3.55.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.95, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5361, 262, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \arctan(x) dx$$

$$\downarrow \text{5361}$$

$$\frac{1}{2}x^2 \arctan(x) - \frac{1}{2} \int \frac{x^2}{x^2 + 1} dx$$

$$\downarrow \text{262}$$

$$\frac{1}{2} \left(\int \frac{1}{x^2 + 1} dx - x \right) + \frac{1}{2}x^2 \arctan(x)$$

$$\downarrow \text{216}$$

$$\frac{1}{2}x^2 \arctan(x) + \frac{1}{2}(\arctan(x) - x)$$

input `Int[x*ArcTan[x],x]`

output `(x^2*ArcTan[x])/2 + (-x + ArcTan[x])/2`

3.55.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 262 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m-1)*((a + b*x^2)^(p+1)/(b*(m+2*p+1))), x] - Simp[a*c^2*((m-1)/(b*(m+2*p+1)) Int[(c*x)^(m-2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2-1] && NeQ[m+2*p+1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

```
rule 5361 Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :>
  Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m +
  1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x],
  x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] &
  & IntegerQ[m])) && NeQ[m, -1]
```

3.55.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.76

method	result	size
default	$-\frac{x}{2} + \frac{\arctan(x)}{2} + \frac{x^2 \arctan(x)}{2}$	16
meijerg	$-\frac{x}{2} + \frac{(3x^2+3) \arctan(x)}{6}$	16
parallelrisch	$-\frac{x}{2} + \frac{\arctan(x)}{2} + \frac{x^2 \arctan(x)}{2}$	16
parts	$-\frac{x}{2} + \frac{\arctan(x)}{2} + \frac{x^2 \arctan(x)}{2}$	16
risch	$-\frac{ix^2 \ln(ix+1)}{4} + \frac{ix^2 \ln(-ix+1)}{4} - \frac{x}{2} + \frac{\arctan(x)}{2}$	35

```
input int(x*arctan(x), x, method=_RETURNVERBOSE)
```

```
output -1/2*x+1/2*arctan(x)+1/2*x^2*arctan(x)
```

3.55.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int x \arctan(x) dx = \frac{1}{2} (x^2 + 1) \arctan(x) - \frac{1}{2} x$$

```
input integrate(x*arctan(x), x, algorithm="fricas")
```

```
output 1/2*(x^2 + 1)*arctan(x) - 1/2*x
```

3.55.6 Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int x \arctan(x) dx = \frac{x^2 \operatorname{atan}(x)}{2} - \frac{x}{2} + \frac{\operatorname{atan}(x)}{2}$$

input `integrate(x*atan(x),x)`output `x**2*atan(x)/2 - x/2 + atan(x)/2`**3.55.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int x \arctan(x) dx = \frac{1}{2} x^2 \arctan(x) - \frac{1}{2} x + \frac{1}{2} \arctan(x)$$

input `integrate(x*arctan(x),x, algorithm="maxima")`output `1/2*x^2*arctan(x) - 1/2*x + 1/2*arctan(x)`**3.55.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int x \arctan(x) dx = \frac{1}{2} x^2 \arctan(x) - \frac{1}{2} x + \frac{1}{2} \arctan(x)$$

input `integrate(x*arctan(x),x, algorithm="giac")`output `1/2*x^2*arctan(x) - 1/2*x + 1/2*arctan(x)`

3.55.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

$$\int x \arctan(x) dx = \operatorname{atan}(x) \left(\frac{x^2}{2} + \frac{1}{2} \right) - \frac{x}{2}$$

input `int(x*atan(x),x)`

output `atan(x)*(x^2/2 + 1/2) - x/2`

3.56 $\int x \cos(\pi x) dx$

3.56.1	Optimal result	388
3.56.2	Mathematica [A] (verified)	388
3.56.3	Rubi [A] (verified)	389
3.56.4	Maple [A] (verified)	390
3.56.5	Fricas [A] (verification not implemented)	391
3.56.6	Sympy [A] (verification not implemented)	391
3.56.7	Maxima [A] (verification not implemented)	391
3.56.8	Giac [A] (verification not implemented)	392
3.56.9	Mupad [B] (verification not implemented)	392

3.56.1 Optimal result

Integrand size = 6, antiderivative size = 18

$$\int x \cos(\pi x) dx = \frac{\cos(\pi x)}{\pi^2} + \frac{x \sin(\pi x)}{\pi}$$

output `cos(Pi*x)/Pi^2+x*sin(Pi*x)/Pi`

3.56.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int x \cos(\pi x) dx = \frac{\cos(\pi x)}{\pi^2} + \frac{x \sin(\pi x)}{\pi}$$

input `Integrate[x*Cos[Pi*x],x]`

output `Cos[Pi*x]/Pi^2 + (x*Sin[Pi*x])/Pi`

3.56.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \cos(\pi x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x \sin\left(\pi x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3777} \\
 & \frac{\int -\sin(\pi x) dx}{\pi} + \frac{x \sin(\pi x)}{\pi} \\
 & \quad \downarrow \text{25} \\
 & \frac{x \sin(\pi x)}{\pi} - \frac{\int \sin(\pi x) dx}{\pi} \\
 & \quad \downarrow \text{3042} \\
 & \frac{x \sin(\pi x)}{\pi} - \frac{\int \sin(\pi x) dx}{\pi} \\
 & \quad \downarrow \text{3118} \\
 & \frac{x \sin(\pi x)}{\pi} + \frac{\cos(\pi x)}{\pi^2}
 \end{aligned}$$

input `Int[x*Cos[Pi*x],x]`

output `Cos[Pi*x]/Pi^2 + (x*Sin[Pi*x])/Pi`

3.56.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_.))^(m_.)*sin[(e_.) + (f_.)*(x_.)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

3.56.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

method	result	size
derivativedivides	$\frac{\cos(\pi x) + x\pi \sin(\pi x)}{\pi^2}$	17
default	$\frac{\cos(\pi x) + x\pi \sin(\pi x)}{\pi^2}$	17
risch	$\frac{\cos(\pi x)}{\pi^2} + \frac{x \sin(\pi x)}{\pi}$	19
parts	$\frac{\cos(\pi x)}{\pi^2} + \frac{x \sin(\pi x)}{\pi}$	19
parallelrisch	$\frac{2 + 2\pi x \tan(\frac{\pi x}{2})}{\pi^2(1 + \tan^2(\frac{\pi x}{2}))}$	27
norman	$\frac{2x \tan(\frac{\pi x}{2}) + \frac{2}{\pi}}{1 + \tan^2(\frac{\pi x}{2})}$	30
meijerg	$\frac{-\frac{1}{\sqrt{\pi}} + \frac{\cos(\pi x)}{\sqrt{\pi}} + \sqrt{\pi} x \sin(\pi x)}{\pi^{\frac{3}{2}}}$	31

input `int(x*cos(Pi*x), x, method=_RETURNVERBOSE)`

output `1/Pi^2*(cos(Pi*x)+x*Pi*sin(Pi*x))`

3.56.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int x \cos(\pi x) dx = \frac{\pi x \sin(\pi x) + \cos(\pi x)}{\pi^2}$$

input `integrate(x*cos(pi*x),x, algorithm="fricas")`output `(pi*x*sin(pi*x) + cos(pi*x))/pi^2`**3.56.6 Sympy [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int x \cos(\pi x) dx = \frac{x \sin(\pi x)}{\pi} + \frac{\cos(\pi x)}{\pi^2}$$

input `integrate(x*cos(pi*x),x)`output `x*sin(pi*x)/pi + cos(pi*x)/pi**2`**3.56.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int x \cos(\pi x) dx = \frac{\pi x \sin(\pi x) + \cos(\pi x)}{\pi^2}$$

input `integrate(x*cos(pi*x),x, algorithm="maxima")`output `(pi*x*sin(pi*x) + cos(pi*x))/pi^2`

3.56.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int x \cos(\pi x) dx = \frac{x \sin(\pi x)}{\pi} + \frac{\cos(\pi x)}{\pi^2}$$

input `integrate(x*cos(pi*x),x, algorithm="giac")`output `x*sin(pi*x)/pi + cos(pi*x)/pi^2`**3.56.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int x \cos(\pi x) dx = \frac{\cos(\Pi x) + \Pi x \sin(\Pi x)}{\Pi^2}$$

input `int(x*cos(Pi*x),x)`output `(cos(Pi*x) + Pi*x*sin(Pi*x))/Pi^2`

3.57 $\int \sqrt{x} \log(x) dx$

3.57.1	Optimal result	393
3.57.2	Mathematica [A] (verified)	393
3.57.3	Rubi [A] (verified)	394
3.57.4	Maple [A] (verified)	394
3.57.5	Fricas [A] (verification not implemented)	395
3.57.6	Sympy [B] (verification not implemented)	395
3.57.7	Maxima [A] (verification not implemented)	396
3.57.8	Giac [A] (verification not implemented)	396
3.57.9	Mupad [B] (verification not implemented)	396

3.57.1 Optimal result

Integrand size = 8, antiderivative size = 21

$$\int \sqrt{x} \log(x) dx = -\frac{4x^{3/2}}{9} + \frac{2}{3}x^{3/2} \log(x)$$

output `-4/9*x^(3/2)+2/3*x^(3/2)*ln(x)`

3.57.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \sqrt{x} \log(x) dx = \frac{2}{9}x^{3/2}(-2 + 3 \log(x))$$

input `Integrate[Sqrt[x]*Log[x],x]`

output `(2*x^(3/2)*(-2 + 3*Log[x]))/9`

3.57.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x} \log(x) dx$$

$$\downarrow 2741$$

$$\frac{2}{3}x^{3/2} \log(x) - \frac{4x^{3/2}}{9}$$

input `Int[Sqrt[x]*Log[x],x]`

output `(-4*x^(3/2))/9 + (2*x^(3/2)*Log[x])/3`

3.57.3.1 Defintions of rubi rules used

rule 2741 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_))^(m_.), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(m + 1))/(d*(m + 1)^2), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]`

3.57.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

method	result	size
derivativedivides	$-\frac{4x^{\frac{3}{2}}}{9} + \frac{2x^{\frac{3}{2}} \ln(x)}{3}$	14
default	$-\frac{4x^{\frac{3}{2}}}{9} + \frac{2x^{\frac{3}{2}} \ln(x)}{3}$	14
risch	$-\frac{4x^{\frac{3}{2}}}{9} + \frac{2x^{\frac{3}{2}} \ln(x)}{3}$	14

input `int(ln(x)*x^(1/2),x,method=_RETURNVERBOSE)`

output `-4/9*x^(3/2)+2/3*x^(3/2)*ln(x)`

3.57.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

$$\int \sqrt{x} \log(x) dx = \frac{2}{9} (3x \log(x) - 2x) \sqrt{x}$$

input `integrate(log(x)*x^(1/2),x, algorithm="fricas")`

output `2/9*(3*x*log(x) - 2*x)*sqrt(x)`

3.57.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. 2(19) = 38.

Time = 0.93 (sec) , antiderivative size = 105, normalized size of antiderivative = 5.00

$$\int \sqrt{x} \log(x) dx = \begin{cases} -\frac{2x^{\frac{3}{2}} \log\left(\frac{1}{x}\right)}{3} + \frac{2x^{\frac{3}{2}} \log(x)}{3} - \frac{8x^{\frac{3}{2}}}{9} & \text{for } \frac{1}{|x|} < 1 \wedge |x| < 1 \\ \frac{2x^{\frac{3}{2}} \log(x)}{3} - \frac{4x^{\frac{3}{2}}}{9} & \text{for } |x| < 1 \\ -\frac{2x^{\frac{3}{2}} \log\left(\frac{1}{x}\right)}{3} - \frac{4x^{\frac{3}{2}}}{9} & \text{for } \frac{1}{|x|} < 1 \\ -G_{3,3}^{2,1} \left(\begin{matrix} 1 & \frac{5}{2}, \frac{5}{2} \\ \frac{3}{2}, \frac{3}{2} \end{matrix} \middle| x \right) + G_{3,3}^{0,3} \left(\begin{matrix} \frac{5}{2}, \frac{5}{2}, 1 \\ \frac{3}{2}, \frac{3}{2}, 0 \end{matrix} \middle| x \right) & \text{otherwise} \end{cases}$$

input `integrate(ln(x)*x**(1/2),x)`

output `Piecewise((-2*x**(3/2)*log(1/x)/3 + 2*x**(3/2)*log(x)/3 - 8*x**(3/2)/9, (Abs(x) < 1) & (1/Abs(x) < 1)), (2*x**(3/2)*log(x)/3 - 4*x**(3/2)/9, Abs(x) < 1), (-2*x**(3/2)*log(1/x)/3 - 4*x**(3/2)/9, 1/Abs(x) < 1), (-meijerg(((1, (5/2, 5/2)), ((3/2, 3/2), (0,)), x) + meijerg(((5/2, 5/2, 1), ()), ((3/2, 3/2, 0)), x), True))`

3.57.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \sqrt{x} \log(x) dx = \frac{2}{3} x^{\frac{3}{2}} \log(x) - \frac{4}{9} x^{\frac{3}{2}}$$

input `integrate(log(x)*x^(1/2),x, algorithm="maxima")`output `2/3*x^(3/2)*log(x) - 4/9*x^(3/2)`**3.57.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \sqrt{x} \log(x) dx = \frac{2}{3} x^{\frac{3}{2}} \log(x) - \frac{4}{9} x^{\frac{3}{2}}$$

input `integrate(log(x)*x^(1/2),x, algorithm="giac")`output `2/3*x^(3/2)*log(x) - 4/9*x^(3/2)`**3.57.9 Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.43

$$\int \sqrt{x} \log(x) dx = \frac{2x^{3/2} (\ln(x) - \frac{2}{3})}{3}$$

input `int(x^(1/2)*log(x),x)`output `(2*x^(3/2)*(log(x) - 2/3))/3`

3.58 $\int \sin^2(3x) dx$

3.58.1	Optimal result	397
3.58.2	Mathematica [A] (verified)	397
3.58.3	Rubi [A] (verified)	398
3.58.4	Maple [A] (verified)	399
3.58.5	Fricas [A] (verification not implemented)	399
3.58.6	Sympy [A] (verification not implemented)	400
3.58.7	Maxima [A] (verification not implemented)	400
3.58.8	Giac [A] (verification not implemented)	400
3.58.9	Mupad [B] (verification not implemented)	401

3.58.1 Optimal result

Integrand size = 6, antiderivative size = 18

$$\int \sin^2(3x) dx = \frac{x}{2} - \frac{1}{6} \cos(3x) \sin(3x)$$

output `1/2*x-1/6*cos(3*x)*sin(3*x)`

3.58.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \sin^2(3x) dx = \frac{x}{2} - \frac{1}{12} \sin(6x)$$

input `Integrate[Sin[3*x]^2,x]`

output `x/2 - Sin[6*x]/12`

3.58.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin^2(3x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin(3x)^2 dx \\ & \quad \downarrow \text{3115} \\ & \frac{\int 1 dx}{2} - \frac{1}{6} \sin(3x) \cos(3x) \\ & \quad \downarrow \text{24} \\ & \frac{x}{2} - \frac{1}{6} \sin(3x) \cos(3x) \end{aligned}$$

input `Int[Sin[3*x]^2,x]`

output `x/2 - (Cos[3*x]*Sin[3*x])/6`

3.58.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sine[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sine[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

3.58.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.61

method	result	size
risch	$\frac{x}{2} - \frac{\sin(6x)}{12}$	11
parallelrisch	$\frac{x}{2} - \frac{\sin(6x)}{12}$	11
derivativedivides	$\frac{x}{2} - \frac{\sin(3x)\cos(3x)}{6}$	15
default	$\frac{x}{2} - \frac{\sin(3x)\cos(3x)}{6}$	15
meijerg	$\frac{\sqrt{\pi} \left(\frac{6x}{\sqrt{\pi}} - \frac{\sin(6x)}{\sqrt{\pi}} \right)}{12}$	22
norman	$\frac{x \tan^2\left(\frac{3x}{2}\right) + \frac{x}{2} + \frac{\tan^3\left(\frac{3x}{2}\right)}{3} + \frac{x \tan^4\left(\frac{3x}{2}\right)}{2} - \frac{\tan\left(\frac{3x}{2}\right)}{3}}{\left(1 + \tan^2\left(\frac{3x}{2}\right)\right)^2}$	47

input `int(sin(3*x)^2,x,method=_RETURNVERBOSE)`output `1/2*x-1/12*sin(6*x)`**3.58.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \sin^2(3x) dx = -\frac{1}{6} \cos(3x) \sin(3x) + \frac{1}{2} x$$

input `integrate(sin(3*x)^2,x, algorithm="fracas")`output `-1/6*cos(3*x)*sin(3*x) + 1/2*x`

3.58.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \sin^2(3x) dx = \frac{x}{2} - \frac{\sin(3x) \cos(3x)}{6}$$

input `integrate(sin(3*x)**2,x)`

output `x/2 - sin(3*x)*cos(3*x)/6`

3.58.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.56

$$\int \sin^2(3x) dx = \frac{1}{2} x - \frac{1}{12} \sin(6x)$$

input `integrate(sin(3*x)^2,x, algorithm="maxima")`

output `1/2*x - 1/12*sin(6*x)`

3.58.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.56

$$\int \sin^2(3x) dx = \frac{1}{2} x - \frac{1}{12} \sin(6x)$$

input `integrate(sin(3*x)^2,x, algorithm="giac")`

output `1/2*x - 1/12*sin(6*x)`

3.58.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.56

$$\int \sin^2(3x) dx = \frac{x}{2} - \frac{\sin(6x)}{12}$$

input `int(sin(3*x)^2,x)`

output `x/2 - sin(6*x)/12`

3.59 $\int \cos^2(x) dx$

3.59.1	Optimal result	402
3.59.2	Mathematica [A] (verified)	402
3.59.3	Rubi [A] (verified)	403
3.59.4	Maple [A] (verified)	404
3.59.5	Fricas [A] (verification not implemented)	404
3.59.6	Sympy [A] (verification not implemented)	404
3.59.7	Maxima [A] (verification not implemented)	405
3.59.8	Giac [A] (verification not implemented)	405
3.59.9	Mupad [B] (verification not implemented)	405

3.59.1 Optimal result

Integrand size = 4, antiderivative size = 14

$$\int \cos^2(x) dx = \frac{x}{2} + \frac{1}{2} \cos(x) \sin(x)$$

output `1/2*x+1/2*cos(x)*sin(x)`

3.59.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \cos^2(x) dx = \frac{x}{2} + \frac{1}{4} \sin(2x)$$

input `Integrate[Cos[x]^2,x]`

output `x/2 + Sin[2*x]/4`

3.59.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^2(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin\left(x + \frac{\pi}{2}\right)^2 dx \\ & \quad \downarrow \text{3115} \\ & \frac{\int 1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \\ & \quad \downarrow \text{24} \\ & \frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \end{aligned}$$

input `Int[Cos[x]^2,x]`

output `x/2 + (Cos[x]*Sin[x])/2`

3.59.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sine[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sine[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

3.59.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{x}{2} + \frac{\cos(x) \sin(x)}{2}$	11
risch	$\frac{x}{2} + \frac{\sin(2x)}{4}$	11
parallelrisch	$\frac{x}{2} + \frac{\sin(2x)}{4}$	11
norman	$\frac{x(\tan^2(\frac{x}{2})) + \frac{x}{2} - (\tan^3(\frac{x}{2})) + \frac{x(\tan^4(\frac{x}{2}))}{2} + \tan(\frac{x}{2})}{(1+\tan^2(\frac{x}{2}))^2}$	45

input `int(cos(x)^2,x,method=_RETURNVERBOSE)`output `1/2*x+1/2*cos(x)*sin(x)`**3.59.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{1}{2} \cos(x) \sin(x) + \frac{1}{2} x$$

input `integrate(cos(x)^2,x, algorithm="fricas")`output `1/2*cos(x)*sin(x) + 1/2*x`**3.59.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{x}{2} + \frac{\sin(x) \cos(x)}{2}$$

input `integrate(cos(x)**2,x)`output `x/2 + sin(x)*cos(x)/2`

3.59.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{1}{2}x + \frac{1}{4}\sin(2x)$$

input `integrate(cos(x)^2,x, algorithm="maxima")`

output `1/2*x + 1/4*sin(2*x)`

3.59.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{1}{2}x + \frac{1}{4}\sin(2x)$$

input `integrate(cos(x)^2,x, algorithm="giac")`

output `1/2*x + 1/4*sin(2*x)`

3.59.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) dx = \frac{x}{2} + \frac{\sin(2x)}{4}$$

input `int(cos(x)^2,x)`

output `x/2 + sin(2*x)/4`

3.60 $\int \cos^4(x) dx$

3.60.1	Optimal result	406
3.60.2	Mathematica [A] (verified)	406
3.60.3	Rubi [A] (verified)	407
3.60.4	Maple [A] (verified)	408
3.60.5	Fricas [A] (verification not implemented)	409
3.60.6	Sympy [A] (verification not implemented)	409
3.60.7	Maxima [A] (verification not implemented)	409
3.60.8	Giac [A] (verification not implemented)	410
3.60.9	Mupad [B] (verification not implemented)	410

3.60.1 Optimal result

Integrand size = 4, antiderivative size = 24

$$\int \cos^4(x) dx = \frac{3x}{8} + \frac{3}{8} \cos(x) \sin(x) + \frac{1}{4} \cos^3(x) \sin(x)$$

output `3/8*x+3/8*cos(x)*sin(x)+1/4*cos(x)^3*sin(x)`

3.60.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \cos^4(x) dx = \frac{3x}{8} + \frac{1}{4} \sin(2x) + \frac{1}{32} \sin(4x)$$

input `Integrate[Cos[x]^4,x]`

output `(3*x)/8 + Sin[2*x]/4 + Sin[4*x]/32`

3.60.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(x + \frac{\pi}{2}\right)^4 dx \\
 & \quad \downarrow \text{3115} \\
 & \frac{3}{4} \int \cos^2(x) dx + \frac{1}{4} \sin(x) \cos^3(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{4} \int \sin\left(x + \frac{\pi}{2}\right)^2 dx + \frac{1}{4} \sin(x) \cos^3(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{3}{4} \left(\int \frac{1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \right) + \frac{1}{4} \sin(x) \cos^3(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{1}{4} \sin(x) \cos^3(x) + \frac{3}{4} \left(\frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \right)
 \end{aligned}$$

input `Int[Cos[x]^4,x]`

output `(Cos[x]^3*Sin[x])/4 + (3*(x/2 + (Cos[x]*Sin[x])/2))/4`

3.60.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

3.60.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.71

method	result	size
risch	$\frac{3x}{8} + \frac{\sin(4x)}{32} + \frac{\sin(2x)}{4}$	17
parallelrisch	$\frac{3x}{8} + \frac{\sin(4x)}{32} + \frac{\sin(2x)}{4}$	17
default	$\frac{(\cos^3(x) + \frac{3\cos(x)}{2})\sin(x)}{4} + \frac{3x}{8}$	18
norman	$\frac{\frac{3x}{8} - \frac{3(\tan^3(\frac{x}{2}))}{4} + \frac{3(\tan^5(\frac{x}{2}))}{4} - \frac{5(\tan^7(\frac{x}{2}))}{4} + \frac{3x(\tan^2(\frac{x}{2}))}{2} + \frac{9x(\tan^4(\frac{x}{2}))}{4} + \frac{3x(\tan^6(\frac{x}{2}))}{2} + \frac{3x(\tan^8(\frac{x}{2}))}{8} + \frac{5\tan(\frac{x}{2})}{4}}{(1+\tan^2(\frac{x}{2}))^4}$	82

input `int(cos(x)^4,x,method=_RETURNVERBOSE)`

output `3/8*x+1/32*sin(4*x)+1/4*sin(2*x)`

3.60.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \cos^4(x) dx = \frac{1}{8} (2 \cos(x)^3 + 3 \cos(x)) \sin(x) + \frac{3}{8} x$$

input `integrate(cos(x)^4,x, algorithm="fricas")`output `1/8*(2*cos(x)^3 + 3*cos(x))*sin(x) + 3/8*x`**3.60.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \cos^4(x) dx = \frac{3x}{8} + \frac{\sin(x) \cos^3(x)}{4} + \frac{3 \sin(x) \cos(x)}{8}$$

input `integrate(cos(x)**4,x)`output `3*x/8 + sin(x)*cos(x)**3/4 + 3*sin(x)*cos(x)/8`**3.60.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \cos^4(x) dx = \frac{3}{8} x + \frac{1}{32} \sin(4x) + \frac{1}{4} \sin(2x)$$

input `integrate(cos(x)^4,x, algorithm="maxima")`output `3/8*x + 1/32*sin(4*x) + 1/4*sin(2*x)`

3.60.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \cos^4(x) dx = \frac{3}{8}x + \frac{1}{32}\sin(4x) + \frac{1}{4}\sin(2x)$$

input `integrate(cos(x)^4,x, algorithm="giac")`

output `3/8*x + 1/32*sin(4*x) + 1/4*sin(2*x)`

3.60.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \cos^4(x) dx = \frac{3x}{8} + \frac{\sin(2x)}{4} + \frac{\sin(4x)}{32}$$

input `int(cos(x)^4,x)`

output `(3*x)/8 + sin(2*x)/4 + sin(4*x)/32`

3.61 $\int \sin^3(x) dx$

3.61.1	Optimal result	411
3.61.2	Mathematica [A] (verified)	411
3.61.3	Rubi [A] (verified)	412
3.61.4	Maple [A] (verified)	413
3.61.5	Fricas [A] (verification not implemented)	413
3.61.6	Sympy [A] (verification not implemented)	413
3.61.7	Maxima [A] (verification not implemented)	414
3.61.8	Giac [A] (verification not implemented)	414
3.61.9	Mupad [B] (verification not implemented)	414

3.61.1 Optimal result

Integrand size = 4, antiderivative size = 13

$$\int \sin^3(x) dx = -\cos(x) + \frac{\cos^3(x)}{3}$$

output `-cos(x)+1/3*cos(x)^3`

3.61.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \sin^3(x) dx = -\frac{3 \cos(x)}{4} + \frac{1}{12} \cos(3x)$$

input `Integrate[Sin[x]^3,x]`

output `(-3*Cos[x])/4 + Cos[3*x]/12`

3.61.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \sin^3(x) dx \\
 \downarrow \text{3042} \\
 \int \sin(x)^3 dx \\
 \downarrow \text{3113} \\
 - \int (1 - \cos^2(x)) d \cos(x) \\
 \downarrow \text{2009} \\
 \frac{\cos^3(x)}{3} - \cos(x)
 \end{array}$$

input `Int[Sin[x]^3,x]`

output `-Cos[x] + Cos[x]^3/3`

3.61.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^(n - 1)/2], x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

3.61.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

method	result	size
default	$-\frac{(2+\sin^2(x))\cos(x)}{3}$	11
risch	$-\frac{3\cos(x)}{4} + \frac{\cos(3x)}{12}$	12
parallelrisch	$-\frac{2}{3} - \frac{3\cos(x)}{4} + \frac{\cos(3x)}{12}$	13
norman	$\frac{-4(\tan^2(\frac{x}{2})) - \frac{4}{3}}{(1+\tan^2(\frac{x}{2}))^3}$	22

input `int(sin(x)^3,x,method=_RETURNVERBOSE)`output `-1/3*(2+sin(x)^2)*cos(x)`**3.61.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \sin^3(x) dx = \frac{1}{3} \cos^3(x) - \cos(x)$$

input `integrate(sin(x)^3,x, algorithm="fricas")`output `1/3*cos(x)^3 - cos(x)`**3.61.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.62

$$\int \sin^3(x) dx = \frac{\cos^3(x)}{3} - \cos(x)$$

input `integrate(sin(x)**3,x)`output `cos(x)**3/3 - cos(x)`

3.61.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \sin^3(x) dx = \frac{1}{3} \cos(x)^3 - \cos(x)$$

input `integrate(sin(x)^3,x, algorithm="maxima")`

output `1/3*cos(x)^3 - cos(x)`

3.61.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \sin^3(x) dx = \frac{1}{3} \cos(x)^3 - \cos(x)$$

input `integrate(sin(x)^3,x, algorithm="giac")`

output `1/3*cos(x)^3 - cos(x)`

3.61.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \sin^3(x) dx = \frac{\cos(x) (\cos(x)^2 - 3)}{3}$$

input `int(sin(x)^3,x)`

output `(cos(x)*(cos(x)^2 - 3))/3`

3.62 $\int \cos^4(x) \sin^3(x) dx$

3.62.1	Optimal result	415
3.62.2	Mathematica [A] (verified)	415
3.62.3	Rubi [A] (verified)	416
3.62.4	Maple [A] (verified)	417
3.62.5	Fricas [A] (verification not implemented)	417
3.62.6	Sympy [A] (verification not implemented)	418
3.62.7	Maxima [A] (verification not implemented)	418
3.62.8	Giac [A] (verification not implemented)	418
3.62.9	Mupad [B] (verification not implemented)	419

3.62.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \cos^4(x) \sin^3(x) dx = -\frac{1}{5} \cos^5(x) + \frac{\cos^7(x)}{7}$$

output `-1/5*cos(x)^5+1/7*cos(x)^7`

3.62.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.82

$$\int \cos^4(x) \sin^3(x) dx = -\frac{3 \cos(x)}{64} - \frac{1}{64} \cos(3x) + \frac{1}{320} \cos(5x) + \frac{1}{448} \cos(7x)$$

input `Integrate[Cos[x]^4*Sin[x]^3,x]`

output `(-3*Cos[x])/64 - Cos[3*x]/64 + Cos[5*x]/320 + Cos[7*x]/448`

3.62.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 3045, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^3(x) \cos^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^3 \cos(x)^4 dx \\
 & \quad \downarrow \text{3045} \\
 & - \int \cos^4(x) (1 - \cos^2(x)) d \cos(x) \\
 & \quad \downarrow \text{244} \\
 & - \int (\cos^4(x) - \cos^6(x)) d \cos(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\cos^7(x)}{7} - \frac{\cos^5(x)}{5}
 \end{aligned}$$

input `Int[Cos[x]^4*Sin[x]^3,x]`

output `-1/5*Cos[x]^5 + Cos[x]^7/7`

3.62.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] :> Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3045 Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_.)]^(n_.), x_
Symbol] := Simp[-(a*f)^(-1) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x],
x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] &&
!(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

3.62.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativdivides	$-\frac{\cos^5(x)}{5} + \frac{\cos^7(x)}{7}$	14
default	$-\frac{\cos^5(x)}{5} + \frac{\cos^7(x)}{7}$	14
risch	$-\frac{3 \cos(x)}{64} + \frac{\cos(7x)}{448} + \frac{\cos(5x)}{320} - \frac{\cos(3x)}{64}$	24
parallelrisch	$\frac{6}{35} - \frac{3 \cos(x)}{64} + \frac{\cos(7x)}{448} + \frac{\cos(5x)}{320} - \frac{\cos(3x)}{64}$	25
norman	$\frac{-8(\tan^6(\frac{x}{2})) - 4(\tan^{10}(\frac{x}{2})) + 4(\tan^8(\frac{x}{2})) - \frac{4(\tan^2(\frac{x}{2}))}{5} + \frac{8(\tan^4(\frac{x}{2}))}{5} - \frac{4}{35}}{(1 + \tan^2(\frac{x}{2}))^7}$	54

```
input int(cos(x)^4*sin(x)^3,x,method=_RETURNVERBOSE)
```

```
output -1/5*cos(x)^5+1/7*cos(x)^7
```

3.62.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos^4(x) \sin^3(x) dx = \frac{1}{7} \cos(x)^7 - \frac{1}{5} \cos(x)^5$$

```
input integrate(cos(x)^4*sin(x)^3,x, algorithm="fracas")
```

```
output 1/7*cos(x)^7 - 1/5*cos(x)^5
```

3.62.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int \cos^4(x) \sin^3(x) dx = \frac{\cos^7(x)}{7} - \frac{\cos^5(x)}{5}$$

input `integrate(cos(x)**4*sin(x)**3,x)`output `cos(x)**7/7 - cos(x)**5/5`**3.62.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos^4(x) \sin^3(x) dx = \frac{1}{7} \cos(x)^7 - \frac{1}{5} \cos(x)^5$$

input `integrate(cos(x)^4*sin(x)^3,x, algorithm="maxima")`output `1/7*cos(x)^7 - 1/5*cos(x)^5`**3.62.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos^4(x) \sin^3(x) dx = \frac{1}{7} \cos(x)^7 - \frac{1}{5} \cos(x)^5$$

input `integrate(cos(x)^4*sin(x)^3,x, algorithm="giac")`output `1/7*cos(x)^7 - 1/5*cos(x)^5`

3.62.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \cos^4(x) \sin^3(x) dx = \frac{\cos(x)^5 (5 \cos(x)^2 - 7)}{35}$$

input `int(cos(x)^4*sin(x)^3,x)`

output `(cos(x)^5*(5*cos(x)^2 - 7))/35`

3.63 $\int \cos^3(x) \sin^4(x) dx$

3.63.1	Optimal result	420
3.63.2	Mathematica [A] (verified)	420
3.63.3	Rubi [A] (verified)	421
3.63.4	Maple [A] (verified)	422
3.63.5	Fricas [A] (verification not implemented)	422
3.63.6	Sympy [A] (verification not implemented)	423
3.63.7	Maxima [A] (verification not implemented)	423
3.63.8	Giac [A] (verification not implemented)	423
3.63.9	Mupad [B] (verification not implemented)	424

3.63.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \cos^3(x) \sin^4(x) dx = \frac{\sin^5(x)}{5} - \frac{\sin^7(x)}{7}$$

output `1/5*sin(x)^5-1/7*sin(x)^7`

3.63.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.82

$$\int \cos^3(x) \sin^4(x) dx = \frac{3 \sin(x)}{64} - \frac{1}{64} \sin(3x) - \frac{1}{320} \sin(5x) + \frac{1}{448} \sin(7x)$$

input `Integrate[Cos[x]^3*Sin[x]^4,x]`

output `(3*Sin[x])/64 - Sin[3*x]/64 - Sin[5*x]/320 + Sin[7*x]/448`

3.63.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 3044, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^4(x) \cos^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^4 \cos(x)^3 dx \\
 & \quad \downarrow \text{3044} \\
 & \int \sin^4(x) (1 - \sin^2(x)) d \sin(x) \\
 & \quad \downarrow \text{244} \\
 & \int (\sin^4(x) - \sin^6(x)) d \sin(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sin^5(x)}{5} - \frac{\sin^7(x)}{7}
 \end{aligned}$$

input `Int[Cos[x]^3*Sin[x]^4,x]`

output `Sin[x]^5/5 - Sin[x]^7/7`

3.63.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] :> Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3044 Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_
Symbol] := Simp[1/(a*f) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a
*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(I
ntegerQ[(m - 1)/2] && LtQ[0, m, n])
```

3.63.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativdivides	$\frac{\sin^5(x)}{5} - \frac{\sin^7(x)}{7}$	14
default	$\frac{\sin^5(x)}{5} - \frac{\sin^7(x)}{7}$	14
risch	$\frac{3 \sin(x)}{64} + \frac{\sin(7x)}{448} - \frac{\sin(5x)}{320} - \frac{\sin(3x)}{64}$	24
paralelrisch	$\frac{3 \sin(x)}{64} + \frac{\sin(7x)}{448} - \frac{\sin(5x)}{320} - \frac{\sin(3x)}{64}$	24
norman	$\frac{32(\tan^5(\frac{x}{2}))}{5} - \frac{192(\tan^7(\frac{x}{2}))}{35} + \frac{32(\tan^9(\frac{x}{2}))}{5}$ $(1+\tan^2(\frac{x}{2}))^7$	37

```
input int(cos(x)^3*sin(x)^4,x,method=_RETURNVERBOSE)
```

```
output 1/5*sin(x)^5-1/7*sin(x)^7
```

3.63.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.29

$$\int \cos^3(x) \sin^4(x) dx = \frac{1}{35} (5 \cos(x)^6 - 8 \cos(x)^4 + \cos(x)^2 + 2) \sin(x)$$

```
input integrate(cos(x)^3*sin(x)^4,x, algorithm="fracas")
```

```
output 1/35*(5*cos(x)^6 - 8*cos(x)^4 + cos(x)^2 + 2)*sin(x)
```

3.63.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int \cos^3(x) \sin^4(x) dx = -\frac{\sin^7(x)}{7} + \frac{\sin^5(x)}{5}$$

input `integrate(cos(x)**3*sin(x)**4,x)`output `-sin(x)**7/7 + sin(x)**5/5`**3.63.7 Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos^3(x) \sin^4(x) dx = -\frac{1}{7} \sin(x)^7 + \frac{1}{5} \sin(x)^5$$

input `integrate(cos(x)^3*sin(x)^4,x, algorithm="maxima")`output `-1/7*sin(x)^7 + 1/5*sin(x)^5`**3.63.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos^3(x) \sin^4(x) dx = -\frac{1}{7} \sin(x)^7 + \frac{1}{5} \sin(x)^5$$

input `integrate(cos(x)^3*sin(x)^4,x, algorithm="giac")`output `-1/7*sin(x)^7 + 1/5*sin(x)^5`

3.63.9 Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \cos^3(x) \sin^4(x) dx = -\frac{\sin(x)^5 (5 \sin(x)^2 - 7)}{35}$$

input `int(cos(x)^3*sin(x)^4,x)`

output `-(sin(x)^5*(5*sin(x)^2 - 7))/35`

3.64 $\int \cos^2(x) \sin^4(x) dx$

3.64.1	Optimal result	425
3.64.2	Mathematica [A] (verified)	425
3.64.3	Rubi [A] (verified)	426
3.64.4	Maple [A] (verified)	427
3.64.5	Fricas [A] (verification not implemented)	428
3.64.6	Sympy [A] (verification not implemented)	428
3.64.7	Maxima [A] (verification not implemented)	428
3.64.8	Giac [A] (verification not implemented)	429
3.64.9	Mupad [B] (verification not implemented)	429

3.64.1 Optimal result

Integrand size = 9, antiderivative size = 36

$$\int \cos^2(x) \sin^4(x) dx = \frac{x}{16} + \frac{1}{16} \cos(x) \sin(x) - \frac{1}{8} \cos^3(x) \sin(x) - \frac{1}{6} \cos^3(x) \sin^3(x)$$

output `1/16*x+1/16*cos(x)*sin(x)-1/8*cos(x)^3*sin(x)-1/6*cos(x)^3*sin(x)^3`

3.64.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.83

$$\int \cos^2(x) \sin^4(x) dx = \frac{x}{16} - \frac{1}{64} \sin(2x) - \frac{1}{64} \sin(4x) + \frac{1}{192} \sin(6x)$$

input `Integrate[Cos[x]^2*Sin[x]^4,x]`

output `x/16 - Sin[2*x]/64 - Sin[4*x]/64 + Sin[6*x]/192`

3.64.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.28, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.778$, Rules used = {3042, 3048, 3042, 3048, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^4(x) \cos^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^4 \cos(x)^2 dx \\
 & \quad \downarrow \text{3048} \\
 & \frac{1}{2} \int \cos^2(x) \sin^2(x) dx - \frac{1}{6} \sin^3(x) \cos^3(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \cos(x)^2 \sin(x)^2 dx - \frac{1}{6} \sin^3(x) \cos^3(x) \\
 & \quad \downarrow \text{3048} \\
 & \frac{1}{2} \left(\frac{1}{4} \int \cos^2(x) dx - \frac{1}{4} \sin(x) \cos^3(x) \right) - \frac{1}{6} \sin^3(x) \cos^3(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \left(\frac{1}{4} \int \sin \left(x + \frac{\pi}{2} \right)^2 dx - \frac{1}{4} \sin(x) \cos^3(x) \right) - \frac{1}{6} \sin^3(x) \cos^3(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{1}{2} \left(\frac{1}{4} \left(\int \frac{1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin(x) \cos^3(x) \right) - \frac{1}{6} \sin^3(x) \cos^3(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{1}{2} \left(\frac{1}{4} \left(\frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin(x) \cos^3(x) \right) - \frac{1}{6} \sin^3(x) \cos^3(x)
 \end{aligned}$$

input `Int[Cos[x]^2*Sin[x]^4,x]`

```
output -1/6*(Cos[x]^3*Sin[x]^3) + (-1/4*(Cos[x]^3*Sin[x]) + (x/2 + (Cos[x]*Sin[x])
)/2)/4)/2
```

3.64.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3048 Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^m
_, x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m -
1)/(b*f*(m + n))), x] + Simp[a^2*((m - 1)/(m + n)) Int[(b*Cos[e + f*x])^n
*(a*Sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1]
&& NeQ[m + n, 0] && IntegersQ[2*m, 2*n]
```

```
rule 3115 Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^n_, x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin
[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[
2*n]
```

3.64.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.64

method	result
risch	$\frac{x}{16} + \frac{\sin(6x)}{192} - \frac{\sin(4x)}{64} - \frac{\sin(2x)}{64}$
parallelrisch	$\frac{x}{16} + \frac{\sin(6x)}{192} - \frac{\sin(4x)}{64} - \frac{\sin(2x)}{64}$
default	$\frac{x}{16} + \frac{\cos(x)\sin(x)}{16} - \frac{(\cos^3(x))\sin(x)}{8} - \frac{(\sin^3(x))(\cos^3(x))}{6}$
norman	$\frac{x}{16} - \frac{17(\tan^3(\frac{x}{2}))}{24} + \frac{19(\tan^5(\frac{x}{2}))}{4} - \frac{19(\tan^7(\frac{x}{2}))}{4} + \frac{17(\tan^9(\frac{x}{2}))}{24} + \frac{(\tan^{11}(\frac{x}{2}))}{8} + \frac{3x(\tan^2(\frac{x}{2}))}{8} + \frac{15x(\tan^4(\frac{x}{2}))}{16} + \frac{5x(\tan^6(\frac{x}{2}))}{4} + \frac{1}{(1+\tan^2(\frac{x}{2}))^6}$

```
input int(sin(x)^4*cos(x)^2,x,method=_RETURNVERBOSE)
```

3.64. $\int \cos^2(x) \sin^4(x) dx$

output $1/16*x+1/192*\sin(6*x)-1/64*\sin(4*x)-1/64*\sin(2*x)$

3.64.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.69

$$\int \cos^2(x) \sin^4(x) dx = \frac{1}{48} (8 \cos(x)^5 - 14 \cos(x)^3 + 3 \cos(x)) \sin(x) + \frac{1}{16} x$$

input `integrate(cos(x)^2*sin(x)^4,x, algorithm="fricas")`

output $1/48*(8*\cos(x)^5 - 14*\cos(x)^3 + 3*\cos(x))*\sin(x) + 1/16*x$

3.64.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

$$\int \cos^2(x) \sin^4(x) dx = \frac{x}{16} + \frac{\sin^5(x) \cos(x)}{6} - \frac{\sin^3(x) \cos(x)}{24} - \frac{\sin(x) \cos(x)}{16}$$

input `integrate(cos(x)**2*sin(x)**4,x)`

output $x/16 + \sin(x)**5*\cos(x)/6 - \sin(x)**3*\cos(x)/24 - \sin(x)*\cos(x)/16$

3.64.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.50

$$\int \cos^2(x) \sin^4(x) dx = -\frac{1}{48} \sin(2x)^3 + \frac{1}{16} x - \frac{1}{64} \sin(4x)$$

input `integrate(cos(x)^2*sin(x)^4,x, algorithm="maxima")`

output $-1/48*\sin(2*x)^3 + 1/16*x - 1/64*\sin(4*x)$

3.64.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.61

$$\int \cos^2(x) \sin^4(x) dx = \frac{1}{16} x + \frac{1}{192} \sin(6x) - \frac{1}{64} \sin(4x) - \frac{1}{64} \sin(2x)$$

input `integrate(cos(x)^2*sin(x)^4,x, algorithm="giac")`output `1/16*x + 1/192*sin(6*x) - 1/64*sin(4*x) - 1/64*sin(2*x)`**3.64.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.67

$$\int \cos^2(x) \sin^4(x) dx = \frac{\cos(x) \sin(x)^5}{6} + \frac{x}{16} - \frac{\sin(2x)}{24} + \frac{\sin(4x)}{192}$$

input `int(cos(x)^2*sin(x)^4,x)`output `x/16 - sin(2*x)/24 + sin(4*x)/192 + (cos(x)*sin(x)^5)/6`

3.65 $\int \cos^2(x) \sin^2(x) dx$

3.65.1	Optimal result	430
3.65.2	Mathematica [A] (verified)	430
3.65.3	Rubi [A] (verified)	431
3.65.4	Maple [A] (verified)	432
3.65.5	Fricas [A] (verification not implemented)	433
3.65.6	Sympy [A] (verification not implemented)	433
3.65.7	Maxima [A] (verification not implemented)	433
3.65.8	Giac [A] (verification not implemented)	434
3.65.9	Mupad [B] (verification not implemented)	434

3.65.1 Optimal result

Integrand size = 9, antiderivative size = 24

$$\int \cos^2(x) \sin^2(x) dx = \frac{x}{8} + \frac{1}{8} \cos(x) \sin(x) - \frac{1}{4} \cos^3(x) \sin(x)$$

output `1/8*x+1/8*cos(x)*sin(x)-1/4*cos(x)^3*sin(x)`

3.65.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.58

$$\int \cos^2(x) \sin^2(x) dx = \frac{x}{8} - \frac{1}{32} \sin(4x)$$

input `Integrate[Cos[x]^2*Sin[x]^2,x]`

output `x/8 - Sin[4*x]/32`

3.65.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 3048, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^2(x) \cos^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^2 \cos(x)^2 dx \\
 & \quad \downarrow \text{3048} \\
 & \frac{1}{4} \int \cos^2(x) dx - \frac{1}{4} \sin(x) \cos^3(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4} \int \sin\left(x + \frac{\pi}{2}\right)^2 dx - \frac{1}{4} \sin(x) \cos^3(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{1}{4} \left(\int \frac{1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin(x) \cos^3(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{1}{4} \left(\frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin(x) \cos^3(x)
 \end{aligned}$$

input `Int[Cos[x]^2*Sin[x]^2,x]`

output `-1/4*(Cos[x]^3*Sin[x]) + (x/2 + (Cos[x]*Sin[x])/2)/4`

3.65.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3048 Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Simp[a^2*((m - 1)/(m + n)) Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]
```

```
rule 3115 Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^n_, x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

3.65.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46

method	result	size
risch	$\frac{x}{8} - \frac{\sin(4x)}{32}$	11
parallelrisch	$\frac{x}{8} - \frac{\sin(4x)}{32}$	11
default	$\frac{x}{8} + \frac{\cos(x)\sin(x)}{8} - \frac{(\cos^3(x))\sin(x)}{4}$	19
norman	$\frac{x}{8} + \frac{7(\tan^3(\frac{x}{2}))}{4} - \frac{7(\tan^5(\frac{x}{2}))}{4} + \frac{(\tan^7(\frac{x}{2}))}{4} + \frac{x(\tan^2(\frac{x}{2}))}{2} + \frac{3x(\tan^4(\frac{x}{2}))}{4} + \frac{x(\tan^6(\frac{x}{2}))}{2} + \frac{x(\tan^8(\frac{x}{2}))}{8} - \frac{\tan(\frac{x}{2})}{4}$ $(1 + \tan^2(\frac{x}{2}))^4$	82

```
input int(cos(x)^2*sin(x)^2,x,method=_RETURNVERBOSE)
```

```
output 1/8*x-1/32*sin(4*x)
```

3.65. $\int \cos^2(x) \sin^2(x) dx$

3.65.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \cos^2(x) \sin^2(x) dx = -\frac{1}{8} (2 \cos(x)^3 - \cos(x)) \sin(x) + \frac{1}{8} x$$

input `integrate(cos(x)^2*sin(x)^2,x, algorithm="fracas")`output `-1/8*(2*cos(x)^3 - cos(x))*sin(x) + 1/8*x`**3.65.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.58

$$\int \cos^2(x) \sin^2(x) dx = \frac{x}{8} - \frac{\sin(2x) \cos(2x)}{16}$$

input `integrate(cos(x)**2*sin(x)**2,x)`output `x/8 - sin(2*x)*cos(2*x)/16`**3.65.7 Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.42

$$\int \cos^2(x) \sin^2(x) dx = \frac{1}{8} x - \frac{1}{32} \sin(4x)$$

input `integrate(cos(x)^2*sin(x)^2,x, algorithm="maxima")`output `1/8*x - 1/32*sin(4*x)`

3.65.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.42

$$\int \cos^2(x) \sin^2(x) dx = \frac{1}{8} x - \frac{1}{32} \sin(4x)$$

input `integrate(cos(x)^2*sin(x)^2,x, algorithm="giac")`

output `1/8*x - 1/32*sin(4*x)`

3.65.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.75

$$\int \cos^2(x) \sin^2(x) dx = \frac{\cos(x) \sin(x)^3}{4} - \frac{\cos(x) \sin(x)}{8} + \frac{x}{8}$$

input `int(cos(x)^2*sin(x)^2,x)`

output `x/8 - (cos(x)*sin(x))/8 + (cos(x)*sin(x)^3)/4`

3.66 $\int (1 - \sin(2x))^2 dx$

3.66.1	Optimal result	435
3.66.2	Mathematica [A] (verified)	435
3.66.3	Rubi [A] (verified)	436
3.66.4	Maple [A] (verified)	437
3.66.5	Fricas [A] (verification not implemented)	437
3.66.6	Sympy [A] (verification not implemented)	437
3.66.7	Maxima [A] (verification not implemented)	438
3.66.8	Giac [A] (verification not implemented)	438
3.66.9	Mupad [B] (verification not implemented)	438

3.66.1 Optimal result

Integrand size = 10, antiderivative size = 22

$$\int (1 - \sin(2x))^2 dx = \frac{3x}{2} + \cos(2x) - \frac{1}{4} \cos(2x) \sin(2x)$$

output `3/2*x+cos(2*x)-1/4*cos(2*x)*sin(2*x)`

3.66.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int (1 - \sin(2x))^2 dx = \frac{3x}{2} + \cos(2x) - \frac{1}{8} \sin(4x)$$

input `Integrate[(1 - Sin[2*x])^2,x]`

output `(3*x)/2 + Cos[2*x] - Sin[4*x]/8`

3.66.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3123}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int (1 - \sin(2x))^2 dx \\ \downarrow \text{3042} \\ \int (1 - \sin(2x))^2 dx \\ \downarrow \text{3123} \\ \frac{3x}{2} + \cos(2x) - \frac{1}{4} \sin(2x) \cos(2x) \end{array}$$

input `Int[(1 - Sin[2*x])^2,x]`

output `(3*x)/2 + Cos[2*x] - (Cos[2*x]*Sin[2*x])/4`

3.66.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3123 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^2, x_Symbol] := Simp[(2*a^2 + b^2)*(x/2), x] + (-Simp[2*a*b*(Cos[c + d*x]/d), x] - Simp[b^2*Cos[c + d*x]*(Sin[c + d*x]/(2*d)), x]) /; FreeQ[{a, b, c, d}, x]`

3.66.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.68

method	result	size
risch	$\frac{3x}{2} - \frac{\sin(4x)}{8} + \cos(2x)$	15
parallelrisc	$\frac{3x}{2} + 1 - \frac{\sin(4x)}{8} + \cos(2x)$	16
derivativedivides	$\frac{3x}{2} + \cos(2x) - \frac{\sin(2x)\cos(2x)}{4}$	19
default	$\frac{3x}{2} + \cos(2x) - \frac{\sin(2x)\cos(2x)}{4}$	19
parts	$\frac{3x}{2} + \cos(2x) - \frac{\sin(2x)\cos(2x)}{4}$	19
norman	$\frac{2(\tan^2(x) + \frac{3x}{2} + \frac{(\tan^3(x))}{2} + 3x(\tan^2(x)) + \frac{3x(\tan^4(x))}{2} - \frac{\tan(x)}{2} + 2}{(1+\tan^2(x))^2}$	45

input `int((1-sin(2*x))^2,x,method=_RETURNVERBOSE)`output `3/2*x-1/8*sin(4*x)+cos(2*x)`**3.66.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int (1 - \sin(2x))^2 dx = -\frac{1}{4} \cos(2x) \sin(2x) + \frac{3}{2} x + \cos(2x)$$

input `integrate((1-sin(2*x))^2,x, algorithm="fricas")`output `-1/4*cos(2*x)*sin(2*x) + 3/2*x + cos(2*x)`**3.66.6 Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.68

$$\int (1 - \sin(2x))^2 dx = \frac{x \sin^2(2x)}{2} + \frac{x \cos^2(2x)}{2} + x - \frac{\sin(2x) \cos(2x)}{4} + \cos(2x)$$

input `integrate((1-sin(2*x))**2,x)`

output `x*sin(2*x)**2/2 + x*cos(2*x)**2/2 + x - sin(2*x)*cos(2*x)/4 + cos(2*x)`

3.66.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.64

$$\int (1 - \sin(2x))^2 dx = \frac{3}{2}x + \cos(2x) - \frac{1}{8}\sin(4x)$$

input `integrate((1-sin(2*x))^2,x, algorithm="maxima")`

output `3/2*x + cos(2*x) - 1/8*sin(4*x)`

3.66.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.64

$$\int (1 - \sin(2x))^2 dx = \frac{3}{2}x + \cos(2x) - \frac{1}{8}\sin(4x)$$

input `integrate((1-sin(2*x))^2,x, algorithm="giac")`

output `3/2*x + cos(2*x) - 1/8*sin(4*x)`

3.66.9 Mupad [B] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.64

$$\int (1 - \sin(2x))^2 dx = \frac{3x}{2} + \cos(2x) - \frac{\sin(4x)}{8}$$

input `int((sin(2*x) - 1)^2,x)`

output `(3*x)/2 + cos(2*x) - sin(4*x)/8`

3.67 $\int \cos(x) \sin\left(\frac{\pi}{6} + x\right) dx$

3.67.1	Optimal result	439
3.67.2	Mathematica [A] (verified)	439
3.67.3	Rubi [A] (verified)	440
3.67.4	Maple [A] (verified)	441
3.67.5	Fricas [B] (verification not implemented)	441
3.67.6	Sympy [B] (verification not implemented)	441
3.67.7	Maxima [A] (verification not implemented)	442
3.67.8	Giac [A] (verification not implemented)	442
3.67.9	Mupad [B] (verification not implemented)	442

3.67.1 Optimal result

Integrand size = 11, antiderivative size = 20

$$\int \cos(x) \sin\left(\frac{\pi}{6} + x\right) dx = \frac{x}{4} - \frac{1}{4} \cos\left(\frac{\pi}{6} + 2x\right)$$

output `1/4*x-1/4*cos(1/6*Pi+2*x)`

3.67.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \cos(x) \sin\left(\frac{\pi}{6} + x\right) dx = \frac{x}{4} - \frac{1}{4} \cos\left(\frac{\pi}{6} + 2x\right)$$

input `Integrate[Cos[x]*Sin[Pi/6 + x],x]`

output `x/4 - Cos[Pi/6 + 2*x]/4`

3.67.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {5085, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sin\left(x + \frac{\pi}{6}\right) \cos(x) dx$$

$$\downarrow \text{5085}$$

$$\int \left(\frac{1}{2} \sin\left(2x + \frac{\pi}{6}\right) + \frac{1}{4}\right) dx$$

$$\downarrow \text{2009}$$

$$\frac{x}{4} - \frac{1}{4} \cos\left(2x + \frac{\pi}{6}\right)$$

input `Int[Cos[x]*Sin[Pi/6 + x],x]`

output `x/4 - Cos[Pi/6 + 2*x]/4`

3.67.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5085 `Int[Cos[w_]^(q_.)*Sin[v_]^(p_.), x_Symbol] := Int[ExpandTrigReduce[Sin[v]^p * Cos[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))`

3.67.4 Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

method	result	size
default	$\frac{x}{4} - \frac{\cos(\frac{\pi}{6}+2x)}{4}$	15
risch	$\frac{x}{4} - \frac{\sqrt{3} \cos(2x)}{8} + \frac{\sin(2x)}{8}$	20
parallelrisch	$\frac{\sin(\frac{\pi}{3}+2x)}{8} - \frac{\cos(\frac{\pi}{6}+2x)}{8} - \frac{\sqrt{3} \cos(2x)}{8} + \frac{\sqrt{3}}{8} + \frac{x}{4}$	39
norman	$\frac{x \tan(\frac{\pi}{12}+\frac{x}{2})+x \tan(\frac{x}{2})(\tan^2(\frac{\pi}{12}+\frac{x}{2}))+2 \tan(\frac{x}{2}) \tan(\frac{\pi}{12}+\frac{x}{2})-x \tan(\frac{x}{2})-x(\tan^2(\frac{x}{2})) \tan(\frac{\pi}{12}+\frac{x}{2})}{(1+\tan^2(\frac{x}{2}))(1+\tan^2(\frac{\pi}{12}+\frac{x}{2}))}$	91

input `int(cos(x)*sin(1/6*Pi+x),x,method=_RETURNVERBOSE)`output `1/4*x-1/4*cos(1/6*Pi+2*x)`**3.67.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(14) = 28.

Time = 0.25 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.55

$$\int \cos(x) \sin\left(\frac{\pi}{6} + x\right) dx = -\frac{1}{4} \sqrt{3} \cos\left(\frac{1}{6} \pi + x\right)^2 - \frac{1}{4} \cos\left(\frac{1}{6} \pi + x\right) \sin\left(\frac{1}{6} \pi + x\right) + \frac{1}{4} x$$

input `integrate(cos(x)*sin(1/6*pi+x),x, algorithm="fracas")`output `-1/4*sqrt(3)*cos(1/6*pi + x)^2 - 1/4*cos(1/6*pi + x)*sin(1/6*pi + x) + 1/4*x`**3.67.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. 2(12) = 24.

Time = 0.13 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.85

$$\int \cos(x) \sin\left(\frac{\pi}{6} + x\right) dx = -\frac{x \sin(x) \cos\left(x + \frac{\pi}{6}\right)}{2} + \frac{x \sin\left(x + \frac{\pi}{6}\right) \cos(x)}{2} + \frac{\sin(x) \sin\left(x + \frac{\pi}{6}\right)}{2}$$

input `integrate(cos(x)*sin(1/6*pi+x),x)`

output `-x*sin(x)*cos(x + pi/6)/2 + x*sin(x + pi/6)*cos(x)/2 + sin(x)*sin(x + pi/6)/2`

3.67.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \cos(x) \sin\left(\frac{\pi}{6} + x\right) dx = \frac{1}{4}x - \frac{1}{4}\cos\left(\frac{1}{6}\pi + 2x\right)$$

input `integrate(cos(x)*sin(1/6*pi+x),x, algorithm="maxima")`

output `1/4*x - 1/4*cos(1/6*pi + 2*x)`

3.67.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \cos(x) \sin\left(\frac{\pi}{6} + x\right) dx = \frac{1}{4}x - \frac{1}{4}\cos\left(\frac{1}{6}\pi + 2x\right)$$

input `integrate(cos(x)*sin(1/6*pi+x),x, algorithm="giac")`

output `1/4*x - 1/4*cos(1/6*pi + 2*x)`

3.67.9 Mupad [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \cos(x) \sin\left(\frac{\pi}{6} + x\right) dx = \frac{x \sin\left(\frac{\pi}{6}\right)}{2} - \frac{\cos\left(\frac{\pi}{6} + 2x\right)}{4}$$

input `int(cos(x)*sin(Pi/6 + x),x)`

output `(x*sin(Pi/6))/2 - cos(Pi/6 + 2*x)/4`

3.68 $\int \cos^5(x) \sin^5(x) dx$

3.68.1	Optimal result	443
3.68.2	Mathematica [A] (verified)	443
3.68.3	Rubi [A] (verified)	444
3.68.4	Maple [A] (verified)	445
3.68.5	Fricas [A] (verification not implemented)	446
3.68.6	Sympy [A] (verification not implemented)	446
3.68.7	Maxima [A] (verification not implemented)	446
3.68.8	Giac [A] (verification not implemented)	447
3.68.9	Mupad [B] (verification not implemented)	447

3.68.1 Optimal result

Integrand size = 9, antiderivative size = 25

$$\int \cos^5(x) \sin^5(x) dx = \frac{\sin^6(x)}{6} - \frac{\sin^8(x)}{4} + \frac{\sin^{10}(x)}{10}$$

output `1/6*sin(x)^6-1/4*sin(x)^8+1/10*sin(x)^10`

3.68.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \cos^5(x) \sin^5(x) dx = -\frac{5}{512} \cos(2x) + \frac{5 \cos(6x)}{3072} - \frac{\cos(10x)}{5120}$$

input `Integrate[Cos[x]^5*Sin[x]^5,x]`

output `(-5*Cos[2*x])/512 + (5*Cos[6*x])/3072 - Cos[10*x]/5120`

3.68.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.16, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 3044, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^5(x) \cos^5(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^5 \cos(x)^5 dx \\
 & \quad \downarrow \text{3044} \\
 & \int \sin^5(x) (1 - \sin^2(x))^2 d \sin(x) \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \int \sin^4(x) (1 - \sin^2(x))^2 d \sin^2(x) \\
 & \quad \downarrow \text{49} \\
 & \frac{1}{2} \int (\sin^8(x) - 2 \sin^6(x) + \sin^4(x)) d \sin^2(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} \left(\frac{\sin^{10}(x)}{5} - \frac{\sin^8(x)}{2} + \frac{\sin^6(x)}{3} \right)
 \end{aligned}$$

input `Int[Cos[x]^5*Sin[x]^5,x]`

output `(Sin[x]^6/3 - Sin[x]^8/2 + Sin[x]^10/5)/2`

3.68.3.1 Defintions of rubi rules used

- rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int [ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`
- rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3044 `Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Simp[1/(a*f) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a *Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !IntegerQ[(m - 1)/2] && LtQ[0, m, n]`

3.68.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

method	result	size
derivativedivides	$\frac{\sin^6(x)}{6} - \frac{\sin^8(x)}{4} + \frac{\sin^{10}(x)}{10}$	20
default	$\frac{\sin^6(x)}{6} - \frac{\sin^8(x)}{4} + \frac{\sin^{10}(x)}{10}$	20
risch	$-\frac{\cos(10x)}{5120} + \frac{5 \cos(6x)}{3072} - \frac{5 \cos(2x)}{512}$	20
parallelrisch	$-\frac{121}{840} - \frac{\cos(10x)}{5120} + \frac{5 \cos(6x)}{3072} - \frac{5 \cos(2x)}{512}$	21

input `int(cos(x)^5*sin(x)^5,x,method=_RETURNVERBOSE)`

output `1/6*sin(x)^6-1/4*sin(x)^8+1/10*sin(x)^10`

3.68.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \cos^5(x) \sin^5(x) dx = -\frac{1}{10} \cos(x)^{10} + \frac{1}{4} \cos(x)^8 - \frac{1}{6} \cos(x)^6$$

input `integrate(cos(x)^5*sin(x)^5,x, algorithm="fracas")`output `-1/10*cos(x)^10 + 1/4*cos(x)^8 - 1/6*cos(x)^6`**3.68.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \cos^5(x) \sin^5(x) dx = \frac{\sin^{10}(x)}{10} - \frac{\sin^8(x)}{4} + \frac{\sin^6(x)}{6}$$

input `integrate(cos(x)**5*sin(x)**5,x)`output `sin(x)**10/10 - sin(x)**8/4 + sin(x)**6/6`**3.68.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \cos^5(x) \sin^5(x) dx = \frac{1}{10} \sin(x)^{10} - \frac{1}{4} \sin(x)^8 + \frac{1}{6} \sin(x)^6$$

input `integrate(cos(x)^5*sin(x)^5,x, algorithm="maxima")`output `1/10*sin(x)^10 - 1/4*sin(x)^8 + 1/6*sin(x)^6`

3.68.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \cos^5(x) \sin^5(x) dx = -\frac{1}{10} \cos(x)^{10} + \frac{1}{4} \cos(x)^8 - \frac{1}{6} \cos(x)^6$$

input `integrate(cos(x)^5*sin(x)^5,x, algorithm="giac")`output `-1/10*cos(x)^10 + 1/4*cos(x)^8 - 1/6*cos(x)^6`**3.68.9 Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \cos^5(x) \sin^5(x) dx = \frac{\sin(x)^{10}}{10} - \frac{\sin(x)^8}{4} + \frac{\sin(x)^6}{6}$$

input `int(cos(x)^5*sin(x)^5,x)`output `sin(x)^6/6 - sin(x)^8/4 + sin(x)^10/10`

3.69 $\int \sin^6(x) dx$

3.69.1	Optimal result	448
3.69.2	Mathematica [A] (verified)	448
3.69.3	Rubi [A] (verified)	449
3.69.4	Maple [A] (verified)	450
3.69.5	Fricas [A] (verification not implemented)	451
3.69.6	Sympy [A] (verification not implemented)	451
3.69.7	Maxima [A] (verification not implemented)	451
3.69.8	Giac [A] (verification not implemented)	452
3.69.9	Mupad [B] (verification not implemented)	452

3.69.1 Optimal result

Integrand size = 4, antiderivative size = 34

$$\int \sin^6(x) dx = \frac{5x}{16} - \frac{5}{16} \cos(x) \sin(x) - \frac{5}{24} \cos(x) \sin^3(x) - \frac{1}{6} \cos(x) \sin^5(x)$$

output `5/16*x-5/16*cos(x)*sin(x)-5/24*cos(x)*sin(x)^3-1/6*cos(x)*sin(x)^5`

3.69.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.88

$$\int \sin^6(x) dx = \frac{5x}{16} - \frac{15}{64} \sin(2x) + \frac{3}{64} \sin(4x) - \frac{1}{192} \sin(6x)$$

input `Integrate[Sin[x]^6,x]`

output `(5*x)/16 - (15*Sin[2*x])/64 + (3*Sin[4*x])/64 - Sin[6*x]/192`

3.69.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.29, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.750$, Rules used = {3042, 3115, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^6(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^6 dx \\
 & \quad \downarrow \text{3115} \\
 & \frac{5}{6} \int \sin^4(x) dx - \frac{1}{6} \sin^5(x) \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{6} \int \sin(x)^4 dx - \frac{1}{6} \sin^5(x) \cos(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{5}{6} \left(\frac{3}{4} \int \sin^2(x) dx - \frac{1}{4} \sin^3(x) \cos(x) \right) - \frac{1}{6} \sin^5(x) \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{6} \left(\frac{3}{4} \int \sin(x)^2 dx - \frac{1}{4} \sin^3(x) \cos(x) \right) - \frac{1}{6} \sin^5(x) \cos(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{5}{6} \left(\frac{3}{4} \left(\frac{\int 1 dx}{2} - \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin^3(x) \cos(x) \right) - \frac{1}{6} \sin^5(x) \cos(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{5}{6} \left(\frac{3}{4} \left(\frac{x}{2} - \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin^3(x) \cos(x) \right) - \frac{1}{6} \sin^5(x) \cos(x)
 \end{aligned}$$

input `Int [Sin [x] ^6, x]`

output
$$-1/6*(\text{Cos}[x]*\text{Sin}[x]^5) + (5*(-1/4*(\text{Cos}[x]*\text{Sin}[x]^3) + (3*(x/2 - (\text{Cos}[x]*\text{Sin}[x])/2))/4))/6$$

3.69.3.1 Defintions of rubi rules used

rule 24 $\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3115 $\text{Int}[(b_)*\text{sin}[(c_)+(d_)*(x_)]^(n_), x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^(n-1)/(d*n), x] + \text{Simp}[b^2*((n-1)/n) \text{Int}[(b*\text{Sin}[c + d*x])^(n-2), x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

3.69.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.68

method	result
risch	$\frac{5x}{16} - \frac{\sin(6x)}{192} + \frac{3\sin(4x)}{64} - \frac{15\sin(2x)}{64}$
parallelrisc	$\frac{5x}{16} - \frac{\sin(6x)}{192} + \frac{3\sin(4x)}{64} - \frac{15\sin(2x)}{64}$
default	$-\frac{\left(\sin^5(x) + \frac{5(\sin^3(x))}{4} + \frac{15\sin(x)}{8}\right)\cos(x)}{6} + \frac{5x}{16}$
norman	$\frac{5x}{16} - \frac{85(\tan^3(\frac{x}{2}))}{24} - \frac{33(\tan^5(\frac{x}{2}))}{4} + \frac{33(\tan^7(\frac{x}{2}))}{4} + \frac{85(\tan^9(\frac{x}{2}))}{24} + \frac{5(\tan^{11}(\frac{x}{2}))}{8} + \frac{15x(\tan^2(\frac{x}{2}))}{8} + \frac{75x(\tan^4(\frac{x}{2}))}{16} + \frac{25x(\tan^6(\frac{x}{2}))}{4} + \frac{1}{(1+\tan^2(\frac{x}{2}))^6}$

input $\text{int}(\sin(x)^6, x, \text{method}=_RETURNVERBOSE)$

output $5/16*x - 1/192*\sin(6*x) + 3/64*\sin(4*x) - 15/64*\sin(2*x)$

3.69.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.74

$$\int \sin^6(x) dx = -\frac{1}{48} (8 \cos(x)^5 - 26 \cos(x)^3 + 33 \cos(x)) \sin(x) + \frac{5}{16} x$$

input `integrate(sin(x)^6,x, algorithm="fricas")`output `-1/48*(8*cos(x)^5 - 26*cos(x)^3 + 33*cos(x))*sin(x) + 5/16*x`**3.69.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.06

$$\int \sin^6(x) dx = \frac{5x}{16} - \frac{\sin^5(x) \cos(x)}{6} - \frac{5 \sin^3(x) \cos(x)}{24} - \frac{5 \sin(x) \cos(x)}{16}$$

input `integrate(sin(x)**6,x)`output `5*x/16 - sin(x)**5*cos(x)/6 - 5*sin(x)**3*cos(x)/24 - 5*sin(x)*cos(x)/16`**3.69.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.71

$$\int \sin^6(x) dx = \frac{1}{48} \sin(2x)^3 + \frac{5}{16} x + \frac{3}{64} \sin(4x) - \frac{1}{4} \sin(2x)$$

input `integrate(sin(x)^6,x, algorithm="maxima")`output `1/48*sin(2*x)^3 + 5/16*x + 3/64*sin(4*x) - 1/4*sin(2*x)`

3.69.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.65

$$\int \sin^6(x) dx = \frac{5}{16} x - \frac{1}{192} \sin(6x) + \frac{3}{64} \sin(4x) - \frac{15}{64} \sin(2x)$$

input `integrate(sin(x)^6,x, algorithm="giac")`

output `5/16*x - 1/192*sin(6*x) + 3/64*sin(4*x) - 15/64*sin(2*x)`

3.69.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.65

$$\int \sin^6(x) dx = \frac{5x}{16} - \frac{15 \sin(2x)}{64} + \frac{3 \sin(4x)}{64} - \frac{\sin(6x)}{192}$$

input `int(sin(x)^6,x)`

output `(5*x)/16 - (15*sin(2*x))/64 + (3*sin(4*x))/64 - sin(6*x)/192`

3.70 $\int \cos^6(x) dx$

3.70.1	Optimal result	453
3.70.2	Mathematica [A] (verified)	453
3.70.3	Rubi [A] (verified)	454
3.70.4	Maple [A] (verified)	455
3.70.5	Fricas [A] (verification not implemented)	456
3.70.6	Sympy [A] (verification not implemented)	456
3.70.7	Maxima [A] (verification not implemented)	456
3.70.8	Giac [A] (verification not implemented)	457
3.70.9	Mupad [B] (verification not implemented)	457

3.70.1 Optimal result

Integrand size = 4, antiderivative size = 34

$$\int \cos^6(x) dx = \frac{5x}{16} + \frac{5}{16} \cos(x) \sin(x) + \frac{5}{24} \cos^3(x) \sin(x) + \frac{1}{6} \cos^5(x) \sin(x)$$

output `5/16*x+5/16*cos(x)*sin(x)+5/24*cos(x)^3*sin(x)+1/6*cos(x)^5*sin(x)`

3.70.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.88

$$\int \cos^6(x) dx = \frac{5x}{16} + \frac{15}{64} \sin(2x) + \frac{3}{64} \sin(4x) + \frac{1}{192} \sin(6x)$$

input `Integrate[Cos[x]^6,x]`

output `(5*x)/16 + (15*Sin[2*x])/64 + (3*Sin[4*x])/64 + Sin[6*x]/192`

3.70.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.29, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.750$, Rules used = {3042, 3115, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^6(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(x + \frac{\pi}{2}\right)^6 dx \\
 & \quad \downarrow \text{3115} \\
 & \frac{5}{6} \int \cos^4(x) dx + \frac{1}{6} \sin(x) \cos^5(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{6} \int \sin\left(x + \frac{\pi}{2}\right)^4 dx + \frac{1}{6} \sin(x) \cos^5(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{5}{6} \left(\frac{3}{4} \int \cos^2(x) dx + \frac{1}{4} \sin(x) \cos^3(x) \right) + \frac{1}{6} \sin(x) \cos^5(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{6} \left(\frac{3}{4} \int \sin\left(x + \frac{\pi}{2}\right)^2 dx + \frac{1}{4} \sin(x) \cos^3(x) \right) + \frac{1}{6} \sin(x) \cos^5(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{5}{6} \left(\frac{3}{4} \left(\frac{\int 1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \right) + \frac{1}{4} \sin(x) \cos^3(x) \right) + \frac{1}{6} \sin(x) \cos^5(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{1}{6} \sin(x) \cos^5(x) + \frac{5}{6} \left(\frac{1}{4} \sin(x) \cos^3(x) + \frac{3}{4} \left(\frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \right) \right)
 \end{aligned}$$

input `Int[Cos[x]^6,x]`

output $(\cos[x]^5 \sin[x])/6 + (5*((\cos[x]^3 \sin[x])/4 + (3*(x/2 + (\cos[x] \sin[x])/2))/4))/6$

3.70.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sine[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sine[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

3.70.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.68

method	result
risch	$\frac{5x}{16} + \frac{\sin(6x)}{192} + \frac{3 \sin(4x)}{64} + \frac{15 \sin(2x)}{64}$
parallelrisc	$\frac{5x}{16} + \frac{\sin(6x)}{192} + \frac{3 \sin(4x)}{64} + \frac{15 \sin(2x)}{64}$
default	$\frac{\left(\cos^5(x) + \frac{5 \cos^3(x)}{4} + \frac{15 \cos(x)}{8}\right) \sin(x)}{6} + \frac{5x}{16}$
norman	$\frac{5x}{16} - \frac{5 \tan^3\left(\frac{x}{2}\right)}{24} + \frac{15 \tan^5\left(\frac{x}{2}\right)}{4} - \frac{15 \tan^7\left(\frac{x}{2}\right)}{4} + \frac{5 \tan^9\left(\frac{x}{2}\right)}{24} - \frac{11 \tan^{11}\left(\frac{x}{2}\right)}{8} + \frac{15x \tan^2\left(\frac{x}{2}\right)}{8} + \frac{75x \tan^4\left(\frac{x}{2}\right)}{16} + \frac{25x \tan^6\left(\frac{x}{2}\right)}{4} \frac{1}{(1 + \tan^2\left(\frac{x}{2}\right))^6}$

input `int(cos(x)^6,x,method=_RETURNVERBOSE)`

output $5/16*x+1/192*\sin(6*x)+3/64*\sin(4*x)+15/64*\sin(2*x)$

3.70.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.74

$$\int \cos^6(x) dx = \frac{1}{48} (8 \cos(x)^5 + 10 \cos(x)^3 + 15 \cos(x)) \sin(x) + \frac{5}{16} x$$

input `integrate(cos(x)^6,x, algorithm="fricas")`output `1/48*(8*cos(x)^5 + 10*cos(x)^3 + 15*cos(x))*sin(x) + 5/16*x`**3.70.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.06

$$\int \cos^6(x) dx = \frac{5x}{16} + \frac{\sin(x) \cos^5(x)}{6} + \frac{5 \sin(x) \cos^3(x)}{24} + \frac{5 \sin(x) \cos(x)}{16}$$

input `integrate(cos(x)**6,x)`output `5*x/16 + sin(x)*cos(x)**5/6 + 5*sin(x)*cos(x)**3/24 + 5*sin(x)*cos(x)/16`**3.70.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.71

$$\int \cos^6(x) dx = -\frac{1}{48} \sin(2x)^3 + \frac{5}{16} x + \frac{3}{64} \sin(4x) + \frac{1}{4} \sin(2x)$$

input `integrate(cos(x)^6,x, algorithm="maxima")`output `-1/48*sin(2*x)^3 + 5/16*x + 3/64*sin(4*x) + 1/4*sin(2*x)`

3.70.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.65

$$\int \cos^6(x) dx = \frac{5}{16} x + \frac{1}{192} \sin(6x) + \frac{3}{64} \sin(4x) + \frac{15}{64} \sin(2x)$$

input `integrate(cos(x)^6,x, algorithm="giac")`

output `5/16*x + 1/192*sin(6*x) + 3/64*sin(4*x) + 15/64*sin(2*x)`

3.70.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.65

$$\int \cos^6(x) dx = \frac{5x}{16} + \frac{15 \sin(2x)}{64} + \frac{3 \sin(4x)}{64} + \frac{\sin(6x)}{192}$$

input `int(cos(x)^6,x)`

output `(5*x)/16 + (15*sin(2*x))/64 + (3*sin(4*x))/64 + sin(6*x)/192`

3.71 $\int \cos^4(2x) \sin^2(2x) dx$

3.71.1	Optimal result	458
3.71.2	Mathematica [A] (verified)	458
3.71.3	Rubi [A] (verified)	459
3.71.4	Maple [A] (verified)	460
3.71.5	Fricas [A] (verification not implemented)	461
3.71.6	Sympy [A] (verification not implemented)	461
3.71.7	Maxima [A] (verification not implemented)	461
3.71.8	Giac [A] (verification not implemented)	462
3.71.9	Mupad [B] (verification not implemented)	462

3.71.1 Optimal result

Integrand size = 13, antiderivative size = 46

$$\int \cos^4(2x) \sin^2(2x) dx = \frac{x}{16} + \frac{1}{32} \cos(2x) \sin(2x) + \frac{1}{48} \cos^3(2x) \sin(2x) - \frac{1}{12} \cos^5(2x) \sin(2x)$$

output `1/16*x+1/32*cos(2*x)*sin(2*x)+1/48*cos(2*x)^3*sin(2*x)-1/12*cos(2*x)^5*sin(2*x)`

3.71.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.65

$$\int \cos^4(2x) \sin^2(2x) dx = \frac{x}{16} + \frac{1}{128} \sin(4x) - \frac{1}{128} \sin(8x) - \frac{1}{384} \sin(12x)$$

input `Integrate[Cos[2*x]^4*Sin[2*x]^2,x]`

output `x/16 + Sin[4*x]/128 - Sin[8*x]/128 - Sin[12*x]/384`

3.71.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.22, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$, Rules used = {3042, 3048, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^2(2x) \cos^4(2x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(2x)^2 \cos(2x)^4 dx \\
 & \quad \downarrow \text{3048} \\
 & \frac{1}{6} \int \cos^4(2x) dx - \frac{1}{12} \sin(2x) \cos^5(2x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{6} \int \sin\left(2x + \frac{\pi}{2}\right)^4 dx - \frac{1}{12} \sin(2x) \cos^5(2x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{1}{6} \left(\frac{3}{4} \int \cos^2(2x) dx + \frac{1}{8} \sin(2x) \cos^3(2x) \right) - \frac{1}{12} \sin(2x) \cos^5(2x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{6} \left(\frac{3}{4} \int \sin\left(2x + \frac{\pi}{2}\right)^2 dx + \frac{1}{8} \sin(2x) \cos^3(2x) \right) - \frac{1}{12} \sin(2x) \cos^5(2x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{1}{6} \left(\frac{3}{4} \left(\frac{\int 1 dx}{2} + \frac{1}{4} \sin(2x) \cos(2x) \right) + \frac{1}{8} \sin(2x) \cos^3(2x) \right) - \frac{1}{12} \sin(2x) \cos^5(2x) \\
 & \quad \downarrow \text{24} \\
 & \frac{1}{6} \left(\frac{1}{8} \sin(2x) \cos^3(2x) + \frac{3}{4} \left(\frac{x}{2} + \frac{1}{4} \sin(2x) \cos(2x) \right) \right) - \frac{1}{12} \sin(2x) \cos^5(2x)
 \end{aligned}$$

input `Int[Cos[2*x]^4*Sin[2*x]^2,x]`

output $-1/12*(\text{Cos}[2*x]^5*\text{Sin}[2*x]) + ((\text{Cos}[2*x]^3*\text{Sin}[2*x])/8 + (3*(x/2 + (\text{Cos}[2*x]*\text{Sin}[2*x])/4))/4)/6$

3.71.3.1 Defintions of rubi rules used

rule 24 $\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3048 $\text{Int}[(\text{cos}[(e_.) + (f_.)*(x_.)]*(b_.))^n*((a_.)*\text{sin}[(e_.) + (f_.)*(x_.)])^m, x_Symbol] \rightarrow \text{Simp}[(-a)*(b*\text{Cos}[e + f*x])^{n+1}*((a*\text{Sin}[e + f*x])^{m-1})/(b*f*(m+n)), x] + \text{Simp}[a^2*((m-1)/(m+n)) \text{Int}[(b*\text{Cos}[e + f*x])^n*(a*\text{Sin}[e + f*x])^{m-2}, x], x] /; \text{FreeQ}[\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{NeQ}[m+n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

rule 3115 $\text{Int}[(b_.)*\text{sin}[(c_.) + (d_.)*(x_.)]^n, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{n-1}/(d*n), x] + \text{Simp}[b^2*((n-1)/n) \text{Int}[(b*\text{Sin}[c + d*x])^{n-2}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

3.71.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.50

method	result
risch	$\frac{x}{16} - \frac{\sin(12x)}{384} - \frac{\sin(8x)}{128} + \frac{\sin(4x)}{128}$
parallelrisch	$\frac{x}{16} - \frac{\sin(12x)}{384} - \frac{\sin(8x)}{128} + \frac{\sin(4x)}{128}$
derivativedivides	$-\frac{(\cos^5(2x)) \sin(2x)}{12} + \frac{(\cos^3(2x) + \frac{3 \cos(2x)}{2}) \sin(2x)}{48} + \frac{x}{16}$
default	$-\frac{(\cos^5(2x)) \sin(2x)}{12} + \frac{(\cos^3(2x) + \frac{3 \cos(2x)}{2}) \sin(2x)}{48} + \frac{x}{16}$
norman	$\frac{x}{16} + \frac{47(\tan^3(x))}{48} - \frac{13(\tan^5(x))}{8} + \frac{13(\tan^7(x))}{8} - \frac{47(\tan^9(x))}{48} + \frac{(\tan^{11}(x))}{16} + \frac{3x(\tan^2(x))}{8(1+\tan^2(x))^6} + \frac{15x(\tan^4(x))}{16(1+\tan^2(x))^6} + \frac{5x(\tan^6(x))}{4(1+\tan^2(x))^6} + \frac{15x(\tan^8(x))}{16(1+\tan^2(x))^6}$

input `int(cos(2*x)^4*sin(2*x)^2,x,method=_RETURNVERBOSE)`

output `1/16*x-1/384*sin(12*x)-1/128*sin(8*x)+1/128*sin(4*x)`

3.71.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.72

$$\int \cos^4(2x) \sin^2(2x) dx = -\frac{1}{96} (8 \cos(2x)^5 - 2 \cos(2x)^3 - 3 \cos(2x)) \sin(2x) + \frac{1}{16} x$$

input `integrate(cos(2*x)^4*sin(2*x)^2,x, algorithm="fricas")`

output `-1/96*(8*cos(2*x)^5 - 2*cos(2*x)^3 - 3*cos(2*x))*sin(2*x) + 1/16*x`

3.71.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.89

$$\int \cos^4(2x) \sin^2(2x) dx = \frac{x}{16} - \frac{\sin(2x) \cos^5(2x)}{12} + \frac{\sin(2x) \cos^3(2x)}{48} + \frac{\sin(2x) \cos(2x)}{32}$$

input `integrate(cos(2*x)**4*sin(2*x)**2,x)`

output `x/16 - sin(2*x)*cos(2*x)**5/12 + sin(2*x)*cos(2*x)**3/48 + sin(2*x)*cos(2*x)/32`

3.71.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.39

$$\int \cos^4(2x) \sin^2(2x) dx = \frac{1}{96} \sin(4x)^3 + \frac{1}{16} x - \frac{1}{128} \sin(8x)$$

input `integrate(cos(2*x)^4*sin(2*x)^2,x, algorithm="maxima")`

output `1/96*sin(4*x)^3 + 1/16*x - 1/128*sin(8*x)`

3.71.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.48

$$\int \cos^4(2x) \sin^2(2x) dx = \frac{1}{16} x - \frac{1}{384} \sin(12x) - \frac{1}{128} \sin(8x) + \frac{1}{128} \sin(4x)$$

input `integrate(cos(2*x)^4*sin(2*x)^2,x, algorithm="giac")`

output `1/16*x - 1/384*sin(12*x) - 1/128*sin(8*x) + 1/128*sin(4*x)`

3.71.9 Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\int \cos^4(2x) \sin^2(2x) dx = \frac{x}{16} - \frac{\cos(2x) \sin(2x)}{32} + \frac{\sin(2x)^3 \left(\frac{\cos(2x)^3}{6} + \frac{\cos(2x)}{8} \right)}{2}$$

input `int(cos(2*x)^4*sin(2*x)^2,x)`

output `x/16 - (cos(2*x)*sin(2*x))/32 + (sin(2*x)^3*(cos(2*x)/8 + cos(2*x)^3/6))/2`

3.72 $\int \sin^5(x) dx$

3.72.1	Optimal result	463
3.72.2	Mathematica [A] (verified)	463
3.72.3	Rubi [A] (verified)	464
3.72.4	Maple [A] (verified)	465
3.72.5	Fricas [A] (verification not implemented)	465
3.72.6	Sympy [A] (verification not implemented)	465
3.72.7	Maxima [A] (verification not implemented)	466
3.72.8	Giac [A] (verification not implemented)	466
3.72.9	Mupad [B] (verification not implemented)	466

3.72.1 Optimal result

Integrand size = 4, antiderivative size = 21

$$\int \sin^5(x) dx = -\cos(x) + \frac{2 \cos^3(x)}{3} - \frac{\cos^5(x)}{5}$$

output `-cos(x)+2/3*cos(x)^3-1/5*cos(x)^5`

3.72.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \sin^5(x) dx = -\frac{5 \cos(x)}{8} + \frac{5}{48} \cos(3x) - \frac{1}{80} \cos(5x)$$

input `Integrate[Sin[x]^5,x]`

output `(-5*Cos[x])/8 + (5*Cos[3*x])/48 - Cos[5*x]/80`

3.72.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^5(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^5 dx \\
 & \quad \downarrow \text{3113} \\
 & - \int (\cos^4(x) - 2 \cos^2(x) + 1) d \cos(x) \\
 & \quad \downarrow \text{2009} \\
 & -\frac{1}{5} \cos^5(x) + \frac{2 \cos^3(x)}{3} - \cos(x)
 \end{aligned}$$

input `Int[Sin[x]^5,x]`

output `-Cos[x] + (2*Cos[x]^3)/3 - Cos[x]^5/5`

3.72.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^(n - 1)/2], x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

3.72.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

method	result	size
default	$-\frac{\left(\frac{8}{3} + \sin^4(x) + \frac{4(\sin^2(x))}{3}\right) \cos(x)}{5}$	17
risch	$-\frac{5 \cos(x)}{8} - \frac{\cos(5x)}{80} + \frac{5 \cos(3x)}{48}$	18
parallelrisch	$\frac{8}{15} - \frac{5 \cos(x)}{8} + \frac{5 \cos(3x)}{48} - \frac{\cos(5x)}{80}$	19
norman	$\frac{-\frac{32(\tan^4(\frac{x}{2}))}{3} - \frac{16(\tan^2(\frac{x}{2}))}{3} - \frac{16}{15}}{(1 + \tan^2(\frac{x}{2}))^5}$	30

input `int(sin(x)^5,x,method=_RETURNVERBOSE)`output `-1/5*(8/3+sin(x)^4+4/3*sin(x)^2)*cos(x)`**3.72.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \sin^5(x) dx = -\frac{1}{5} \cos(x)^5 + \frac{2}{3} \cos(x)^3 - \cos(x)$$

input `integrate(sin(x)^5,x, algorithm="fricas")`output `-1/5*cos(x)^5 + 2/3*cos(x)^3 - cos(x)`**3.72.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \sin^5(x) dx = -\frac{\cos^5(x)}{5} + \frac{2 \cos^3(x)}{3} - \cos(x)$$

input `integrate(sin(x)**5,x)`output `-cos(x)**5/5 + 2*cos(x)**3/3 - cos(x)`

3.72.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \sin^5(x) dx = -\frac{1}{5} \cos(x)^5 + \frac{2}{3} \cos(x)^3 - \cos(x)$$

input `integrate(sin(x)^5,x, algorithm="maxima")`output `-1/5*cos(x)^5 + 2/3*cos(x)^3 - cos(x)`**3.72.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \sin^5(x) dx = -\frac{1}{5} \cos(x)^5 + \frac{2}{3} \cos(x)^3 - \cos(x)$$

input `integrate(sin(x)^5,x, algorithm="giac")`output `-1/5*cos(x)^5 + 2/3*cos(x)^3 - cos(x)`**3.72.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \sin^5(x) dx = -\frac{\cos(x)^5}{5} + \frac{2 \cos(x)^3}{3} - \cos(x)$$

input `int(sin(x)^5,x)`output `(2*cos(x)^3)/3 - cos(x) - cos(x)^5/5`

3.73 $\int \cos^4(x) \sin^4(x) dx$

3.73.1	Optimal result	467
3.73.2	Mathematica [A] (verified)	467
3.73.3	Rubi [A] (verified)	468
3.73.4	Maple [A] (verified)	470
3.73.5	Fricas [A] (verification not implemented)	470
3.73.6	Sympy [A] (verification not implemented)	470
3.73.7	Maxima [A] (verification not implemented)	471
3.73.8	Giac [A] (verification not implemented)	471
3.73.9	Mupad [B] (verification not implemented)	471

3.73.1 Optimal result

Integrand size = 9, antiderivative size = 46

$$\int \cos^4(x) \sin^4(x) dx = \frac{3x}{128} + \frac{3}{128} \cos(x) \sin(x) + \frac{1}{64} \cos^3(x) \sin(x) - \frac{1}{16} \cos^5(x) \sin(x) - \frac{1}{8} \cos^5(x) \sin^3(x)$$

output `3/128*x+3/128*cos(x)*sin(x)+1/64*cos(x)^3*sin(x)-1/16*cos(x)^5*sin(x)-1/8*cos(x)^5*sin(x)^3`

3.73.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.48

$$\int \cos^4(x) \sin^4(x) dx = \frac{3x}{128} - \frac{1}{128} \sin(4x) + \frac{\sin(8x)}{1024}$$

input `Integrate[Cos[x]^4*Sin[x]^4,x]`

output `(3*x)/128 - Sin[4*x]/128 + Sin[8*x]/1024`

3.73.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.33, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 3048, 3042, 3048, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^4(x) \cos^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^4 \cos(x)^4 dx \\
 & \quad \downarrow \text{3048} \\
 & \frac{3}{8} \int \cos^4(x) \sin^2(x) dx - \frac{1}{8} \sin^3(x) \cos^5(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{8} \int \cos(x)^4 \sin(x)^2 dx - \frac{1}{8} \sin^3(x) \cos^5(x) \\
 & \quad \downarrow \text{3048} \\
 & \frac{3}{8} \left(\frac{1}{6} \int \cos^4(x) dx - \frac{1}{6} \sin(x) \cos^5(x) \right) - \frac{1}{8} \sin^3(x) \cos^5(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{8} \left(\frac{1}{6} \int \sin \left(x + \frac{\pi}{2} \right)^4 dx - \frac{1}{6} \sin(x) \cos^5(x) \right) - \frac{1}{8} \sin^3(x) \cos^5(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{3}{8} \left(\frac{1}{6} \left(\frac{3}{4} \int \cos^2(x) dx + \frac{1}{4} \sin(x) \cos^3(x) \right) - \frac{1}{6} \sin(x) \cos^5(x) \right) - \frac{1}{8} \sin^3(x) \cos^5(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{8} \left(\frac{1}{6} \left(\frac{3}{4} \int \sin \left(x + \frac{\pi}{2} \right)^2 dx + \frac{1}{4} \sin(x) \cos^3(x) \right) - \frac{1}{6} \sin(x) \cos^5(x) \right) - \frac{1}{8} \sin^3(x) \cos^5(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{3}{8} \left(\frac{1}{6} \left(\frac{3}{4} \left(\frac{\int 1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \right) + \frac{1}{4} \sin(x) \cos^3(x) \right) - \frac{1}{6} \sin(x) \cos^5(x) \right) - \frac{1}{8} \sin^3(x) \cos^5(x) \\
 & \quad \downarrow \text{24}
 \end{aligned}$$

$$\frac{3}{8} \left(\frac{1}{6} \left(\frac{1}{4} \sin(x) \cos^3(x) + \frac{3}{4} \left(\frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \right) \right) - \frac{1}{6} \sin(x) \cos^5(x) \right) - \frac{1}{8} \sin^3(x) \cos^5(x)$$

input `Int[Cos[x]^4*Sin[x]^4,x]`

output `-1/8*(Cos[x]^5*Sin[x]^3) + (3*(-1/6*(Cos[x]^5*Sin[x]) + ((Cos[x]^3*Sin[x])
/4 + (3*(x/2 + (Cos[x]*Sin[x])/2))/4)/6))/8`

3.73.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3048 `Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m -
1)/(b*f*(m + n))), x] + Simp[a^2*((m - 1)/(m + n)) Int[(b*Cos[e + f*x])^n
*(a*Sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1]
&& NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin
[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[
2*n]`

3.73.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.37

method	result
risch	$\frac{3x}{128} + \frac{\sin(8x)}{1024} - \frac{\sin(4x)}{128}$
parallelrisch	$\frac{3x}{128} + \frac{\sin(8x)}{1024} - \frac{\sin(4x)}{128}$
default	$-\frac{(\cos^5(x))(\sin^3(x))}{8} - \frac{(\cos^5(x))\sin(x)}{16} + \frac{(\cos^3(x) + \frac{3\cos(x)}{2})\sin(x)}{64} + \frac{3x}{128}$
norman	$\frac{3x}{128} - \frac{23(\tan^3(\frac{x}{2}))}{64} + \frac{333(\tan^5(\frac{x}{2}))}{64} - \frac{671(\tan^7(\frac{x}{2}))}{64} + \frac{671(\tan^9(\frac{x}{2}))}{64} - \frac{333(\tan^{11}(\frac{x}{2}))}{64} + \frac{23(\tan^{13}(\frac{x}{2}))}{64} + \frac{3(\tan^{15}(\frac{x}{2}))}{64} + \frac{3x(\tan^{17}(\frac{x}{2}))}{16}$

input `int(cos(x)^4*sin(x)^4,x,method=_RETURNVERBOSE)`output `3/128*x+1/1024*sin(8*x)-1/128*sin(4*x)`**3.73.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.67

$$\int \cos^4(x) \sin^4(x) dx = \frac{1}{128} (16 \cos(x)^7 - 24 \cos(x)^5 + 2 \cos(x)^3 + 3 \cos(x)) \sin(x) + \frac{3}{128} x$$

input `integrate(cos(x)^4*sin(x)^4,x, algorithm="fricas")`output `1/128*(16*cos(x)^7 - 24*cos(x)^5 + 2*cos(x)^3 + 3*cos(x))*sin(x) + 3/128*x`**3.73.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.67

$$\int \cos^4(x) \sin^4(x) dx = \frac{3x}{128} - \frac{\sin^3(2x) \cos(2x)}{128} - \frac{3 \sin(2x) \cos(2x)}{256}$$

input `integrate(cos(x)**4*sin(x)**4,x)`output `3*x/128 - sin(2*x)**3*cos(2*x)/128 - 3*sin(2*x)*cos(2*x)/256`

3.73.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.35

$$\int \cos^4(x) \sin^4(x) dx = \frac{3}{128} x + \frac{1}{1024} \sin(8x) - \frac{1}{128} \sin(4x)$$

input `integrate(cos(x)^4*sin(x)^4,x, algorithm="maxima")`output `3/128*x + 1/1024*sin(8*x) - 1/128*sin(4*x)`**3.73.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.35

$$\int \cos^4(x) \sin^4(x) dx = \frac{3}{128} x + \frac{1}{1024} \sin(8x) - \frac{1}{128} \sin(4x)$$

input `integrate(cos(x)^4*sin(x)^4,x, algorithm="giac")`output `3/128*x + 1/1024*sin(8*x) - 1/128*sin(4*x)`**3.73.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.70

$$\int \cos^4(x) \sin^4(x) dx = \left(\frac{\cos(x)^3}{8} + \frac{\cos(x)}{16} \right) \sin(x)^5 + \frac{3x}{128} - \frac{\sin(2x)}{64} + \frac{\sin(4x)}{512}$$

input `int(cos(x)^4*sin(x)^4,x)`output `(3*x)/128 - sin(2*x)/64 + sin(4*x)/512 + sin(x)^5*(cos(x)/16 + cos(x)^3/8)`

3.74 $\int \sqrt{\cos(x)} \sin^3(x) dx$

3.74.1	Optimal result	472
3.74.2	Mathematica [A] (verified)	472
3.74.3	Rubi [A] (verified)	473
3.74.4	Maple [A] (verified)	474
3.74.5	Fricas [A] (verification not implemented)	474
3.74.6	Sympy [F(-1)]	475
3.74.7	Maxima [A] (verification not implemented)	475
3.74.8	Giac [A] (verification not implemented)	475
3.74.9	Mupad [B] (verification not implemented)	476

3.74.1 Optimal result

Integrand size = 11, antiderivative size = 21

$$\int \sqrt{\cos(x)} \sin^3(x) dx = -\frac{2}{3} \cos^{\frac{3}{2}}(x) + \frac{2}{7} \cos^{\frac{7}{2}}(x)$$

output `-2/3*cos(x)^(3/2)+2/7*cos(x)^(7/2)`

3.74.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.62

$$\int \sqrt{\cos(x)} \sin^3(x) dx = \frac{8\sqrt[4]{\cos^2(x)} + \cos^2(x)(-11 + 3 \cos(2x))}{21\sqrt{\cos(x)}}$$

input `Integrate[Sqrt[Cos[x]]*Sin[x]^3,x]`

output `(8*(Cos[x]^2)^(1/4) + Cos[x]^2*(-11 + 3*Cos[2*x]))/(21*Sqrt[Cos[x]])`

3.74.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3045, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^3(x) \sqrt{\cos(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^3 \sqrt{\cos(x)} dx \\
 & \quad \downarrow \text{3045} \\
 & - \int \sqrt{\cos(x)} (1 - \cos^2(x)) d \cos(x) \\
 & \quad \downarrow \text{244} \\
 & - \int \left(\sqrt{\cos(x)} - \cos^{\frac{5}{2}}(x) \right) d \cos(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{2}{7} \cos^{\frac{7}{2}}(x) - \frac{2}{3} \cos^{\frac{3}{2}}(x)
 \end{aligned}$$

input `Int[Sqrt[Cos[x]]*Sin[x]^3,x]`

output `(-2*Cos[x]^(3/2))/3 + (2*Cos[x]^(7/2))/7`

3.74.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3045 `Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[-(a*f)^(-1) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])`

3.74.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

method	result	size
derivativedivides	$-\frac{2(\cos^{\frac{3}{2}}(x))}{3} + \frac{2(\cos^{\frac{7}{2}}(x))}{7}$	14
default	$-\frac{2(\cos^{\frac{3}{2}}(x))}{3} + \frac{2(\cos^{\frac{7}{2}}(x))}{7}$	14

input `int(sin(x)^3*cos(x)^(1/2),x,method=_RETURNVERBOSE)`

output `-2/3*cos(x)^(3/2)+2/7*cos(x)^(7/2)`

3.74.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \sqrt{\cos(x)} \sin^3(x) dx = \frac{2}{21} (3 \cos(x)^3 - 7 \cos(x)) \sqrt{\cos(x)}$$

input `integrate(sin(x)^3*cos(x)^(1/2),x, algorithm="fricas")`

output `2/21*(3*cos(x)^3 - 7*cos(x))*sqrt(cos(x))`

3.74.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{\cos(x)} \sin^3(x) dx = \text{Timed out}$$

input `integrate(sin(x)**3*cos(x)**(1/2),x)`output `Timed out`**3.74.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \sqrt{\cos(x)} \sin^3(x) dx = \frac{2}{7} \cos(x)^{\frac{7}{2}} - \frac{2}{3} \cos(x)^{\frac{3}{2}}$$

input `integrate(sin(x)^3*cos(x)^(1/2),x, algorithm="maxima")`output `2/7*cos(x)^(7/2) - 2/3*cos(x)^(3/2)`**3.74.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \sqrt{\cos(x)} \sin^3(x) dx = \frac{2}{7} \cos(x)^{\frac{7}{2}} - \frac{2}{3} \cos(x)^{\frac{3}{2}}$$

input `integrate(sin(x)^3*cos(x)^(1/2),x, algorithm="giac")`output `2/7*cos(x)^(7/2) - 2/3*cos(x)^(3/2)`

3.74.9 Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \sqrt{\cos(x)} \sin^3(x) dx = \cos(x)^{3/2} \left(\frac{2 \cos(x)^2}{7} - \frac{2}{3} \right)$$

input `int(cos(x)^(1/2)*sin(x)^3,x)`

output `cos(x)^(3/2)*((2*cos(x)^2)/7 - 2/3)`

3.75 $\int \cos^3(x) \sqrt{\sin(x)} dx$

3.75.1	Optimal result	477
3.75.2	Mathematica [A] (verified)	477
3.75.3	Rubi [A] (verified)	478
3.75.4	Maple [A] (verified)	479
3.75.5	Fricas [A] (verification not implemented)	479
3.75.6	Sympy [B] (verification not implemented)	480
3.75.7	Maxima [A] (verification not implemented)	480
3.75.8	Giac [A] (verification not implemented)	481
3.75.9	Mupad [B] (verification not implemented)	481

3.75.1 Optimal result

Integrand size = 11, antiderivative size = 21

$$\int \cos^3(x) \sqrt{\sin(x)} dx = \frac{2}{3} \sin^{\frac{3}{2}}(x) - \frac{2}{7} \sin^{\frac{7}{2}}(x)$$

output `2/3*sin(x)^(3/2)-2/7*sin(x)^(7/2)`

3.75.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

$$\int \cos^3(x) \sqrt{\sin(x)} dx = \frac{1}{21} (11 + 3 \cos(2x)) \sin^{\frac{3}{2}}(x)$$

input `Integrate[Cos[x]^3*Sqrt[Sin[x]],x]`

output `((11 + 3*Cos[2*x])*Sin[x]^(3/2))/21`

3.75.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3044, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{\sin(x)} \cos^3(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{\sin(x)} \cos(x)^3 dx \\ & \quad \downarrow \text{3044} \\ & \int \sqrt{\sin(x)} (1 - \sin^2(x)) d \sin(x) \\ & \quad \downarrow \text{244} \\ & \int \left(\sqrt{\sin(x)} - \sin^{\frac{5}{2}}(x) \right) d \sin(x) \\ & \quad \downarrow \text{2009} \\ & \frac{2}{3} \sin^{\frac{3}{2}}(x) - \frac{2}{7} \sin^{\frac{7}{2}}(x) \end{aligned}$$

input `Int[Cos[x]^3*Sqrt[Sin[x]],x]`

output `(2*Sin[x]^(3/2))/3 - (2*Sin[x]^(7/2))/7`

3.75.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3044 Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)]^(m_.), x_
Symbol] := Simp[1/(a*f) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a
*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(I
ntegerQ[(m - 1)/2] && LtQ[0, m, n])
```

3.75.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

method	result	size
derivativedivides	$\frac{2(\sin^{\frac{3}{2}}(x))}{3} - \frac{2(\sin^{\frac{7}{2}}(x))}{7}$	14
default	$\frac{2(\sin^{\frac{3}{2}}(x))}{3} - \frac{2(\sin^{\frac{7}{2}}(x))}{7}$	14

```
input int(cos(x)^3*sin(x)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 2/3*sin(x)^(3/2)-2/7*sin(x)^(7/2)
```

3.75.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

$$\int \cos^3(x) \sqrt{\sin(x)} dx = \frac{2}{21} (3 \cos^2(x) + 4) \sin(x)^{\frac{3}{2}}$$

```
input integrate(cos(x)^3*sin(x)^(1/2),x, algorithm="fracas")
```

```
output 2/21*(3*cos(x)^2 + 4)*sin(x)^(3/2)
```


3.75.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 170 vs. $2(19) = 38$.

Time = 4.18 (sec) , antiderivative size = 170, normalized size of antiderivative = 8.10

$$\int \cos^3(x) \sqrt{\sin(x)} dx = \frac{28\sqrt{2} \sqrt{\frac{\tan(\frac{x}{2})}{\tan^2(\frac{x}{2})+1}} \tan^5(\frac{x}{2})}{21 \tan^6(\frac{x}{2}) + 63 \tan^4(\frac{x}{2}) + 63 \tan^2(\frac{x}{2}) + 21} + \frac{8\sqrt{2} \sqrt{\frac{\tan(\frac{x}{2})}{\tan^2(\frac{x}{2})+1}} \tan^3(\frac{x}{2})}{21 \tan^6(\frac{x}{2}) + 63 \tan^4(\frac{x}{2}) + 63 \tan^2(\frac{x}{2}) + 21} + \frac{28\sqrt{2} \sqrt{\frac{\tan(\frac{x}{2})}{\tan^2(\frac{x}{2})+1}} \tan(\frac{x}{2})}{21 \tan^6(\frac{x}{2}) + 63 \tan^4(\frac{x}{2}) + 63 \tan^2(\frac{x}{2}) + 21}$$

input `integrate(cos(x)**3*sin(x)**(1/2),x)`

output `28*sqrt(2)*sqrt(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)**5/(21*tan(x/2)**6 + 63*tan(x/2)**4 + 63*tan(x/2)**2 + 21) + 8*sqrt(2)*sqrt(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)**3/(21*tan(x/2)**6 + 63*tan(x/2)**4 + 63*tan(x/2)**2 + 21) + 28*sqrt(2)*sqrt(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)/(21*tan(x/2)**6 + 63*tan(x/2)**4 + 63*tan(x/2)**2 + 21)`

3.75.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \cos^3(x) \sqrt{\sin(x)} dx = -\frac{2}{7} \sin(x)^{\frac{7}{2}} + \frac{2}{3} \sin(x)^{\frac{3}{2}}$$

input `integrate(cos(x)^3*sin(x)^(1/2),x, algorithm="maxima")`

output `-2/7*sin(x)^(7/2) + 2/3*sin(x)^(3/2)`

3.75.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \cos^3(x) \sqrt{\sin(x)} dx = -\frac{2}{7} \sin(x)^{\frac{7}{2}} + \frac{2}{3} \sin(x)^{\frac{3}{2}}$$

input `integrate(cos(x)^3*sin(x)^(1/2),x, algorithm="giac")`output `-2/7*sin(x)^(7/2) + 2/3*sin(x)^(3/2)`**3.75.9 Mupad [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.19

$$\int \cos^3(x) \sqrt{\sin(x)} dx = -\frac{\cos(x)^4 \sin(x)^{3/2} {}_2F_1\left(\frac{1}{4}, 2; 3; \cos(x)^2\right)}{4 (\sin(x)^2)^{3/4}}$$

input `int(cos(x)^3*sin(x)^(1/2),x)`output `-(cos(x)^4*sin(x)^(3/2)*hypergeom([1/4, 2], 3, cos(x)^2))/(4*(sin(x)^2)^(3/4))`

3.76 $\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx$

3.76.1	Optimal result	482
3.76.2	Mathematica [A] (verified)	482
3.76.3	Rubi [A] (verified)	483
3.76.4	Maple [A] (verified)	484
3.76.5	Fricas [A] (verification not implemented)	484
3.76.6	Sympy [B] (verification not implemented)	485
3.76.7	Maxima [A] (verification not implemented)	485
3.76.8	Giac [A] (verification not implemented)	485
3.76.9	Mupad [B] (verification not implemented)	486

3.76.1 Optimal result

Integrand size = 14, antiderivative size = 19

$$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx = \sqrt{x} + \cos(\sqrt{x}) \sin(\sqrt{x})$$

output `cos(x^(1/2))*sin(x^(1/2))+x^(1/2)`

3.76.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

$$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx = \sqrt{x} + \frac{1}{2} \sin(2\sqrt{x})$$

input `Integrate[Cos[Sqrt[x]]^2/Sqrt[x],x]`

output `Sqrt[x] + Sin[2*Sqrt[x]]/2`

3.76.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.47, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3861, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx \\
 & \quad \downarrow \text{3861} \\
 & 2 \int \cos^2(\sqrt{x}) d\sqrt{x} \\
 & \quad \downarrow \text{3042} \\
 & 2 \int \sin\left(\sqrt{x} + \frac{\pi}{2}\right)^2 d\sqrt{x} \\
 & \quad \downarrow \text{3115} \\
 & 2 \left(\frac{\int 1 d\sqrt{x}}{2} + \frac{1}{2} \sin(\sqrt{x}) \cos(\sqrt{x}) \right) \\
 & \quad \downarrow \text{24} \\
 & 2 \left(\frac{\sqrt{x}}{2} + \frac{1}{2} \sin(\sqrt{x}) \cos(\sqrt{x}) \right)
 \end{aligned}$$

input `Int[Cos[Sqrt[x]]^2/Sqrt[x],x]`

output `2*(Sqrt[x]/2 + (Cos[Sqrt[x]]*Sin[Sqrt[x]])/2)`

3.76.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3115 Int[((b_.)*sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

```
rule 3861 Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*cos[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))
```

3.76.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

method	result	size
derivativedivides	$\cos(\sqrt{x}) \sin(\sqrt{x}) + \sqrt{x}$	14
default	$\cos(\sqrt{x}) \sin(\sqrt{x}) + \sqrt{x}$	14

```
input int(cos(x^(1/2))^2/x^(1/2),x,method=_RETURNVERBOSE)
```

```
output cos(x^(1/2))*sin(x^(1/2))+x^(1/2)
```

3.76.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx = \cos(\sqrt{x}) \sin(\sqrt{x}) + \sqrt{x}$$

```
input integrate(cos(x^(1/2))^2/x^(1/2),x, algorithm="fricas")
```

```
output cos(sqrt(x))*sin(sqrt(x)) + sqrt(x)
```

3.76.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. $2(17) = 34$.

Time = 0.11 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.05

$$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx = \sqrt{x} \sin^2(\sqrt{x}) + \sqrt{x} \cos^2(\sqrt{x}) + \sin(\sqrt{x}) \cos(\sqrt{x})$$

input `integrate(cos(x**(1/2))**2/x**(1/2),x)`

output `sqrt(x)*sin(sqrt(x))**2 + sqrt(x)*cos(sqrt(x))**2 + sin(sqrt(x))*cos(sqrt(x))`

3.76.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx = \sqrt{x} + \frac{1}{2} \sin(2\sqrt{x})$$

input `integrate(cos(x^(1/2))**2/x^(1/2),x, algorithm="maxima")`

output `sqrt(x) + 1/2*sin(2*sqrt(x))`

3.76.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx = \sqrt{x} + \frac{1}{2} \sin(2\sqrt{x})$$

input `integrate(cos(x^(1/2))**2/x^(1/2),x, algorithm="giac")`

output `sqrt(x) + 1/2*sin(2*sqrt(x))`

3.76.9 Mupad [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx = \frac{\sin(2\sqrt{x})}{2} + \sqrt{x}$$

input `int(cos(x^(1/2))^2/x^(1/2),x)`

output `sin(2*x^(1/2))/2 + x^(1/2)`

3.77 $\int x \sin^3(x^2) dx$

3.77.1	Optimal result	487
3.77.2	Mathematica [A] (verified)	487
3.77.3	Rubi [A] (warning: unable to verify)	488
3.77.4	Maple [A] (verified)	489
3.77.5	Fricas [A] (verification not implemented)	489
3.77.6	Sympy [A] (verification not implemented)	490
3.77.7	Maxima [A] (verification not implemented)	490
3.77.8	Giac [A] (verification not implemented)	490
3.77.9	Mupad [B] (verification not implemented)	491

3.77.1 Optimal result

Integrand size = 8, antiderivative size = 19

$$\int x \sin^3(x^2) dx = -\frac{1}{2} \cos(x^2) + \frac{1}{6} \cos^3(x^2)$$

output `-1/2*cos(x^2)+1/6*cos(x^2)^3`

3.77.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int x \sin^3(x^2) dx = -\frac{3}{8} \cos(x^2) + \frac{1}{24} \cos(3x^2)$$

input `Integrate[x*Sin[x^2]^3,x]`

output `(-3*Cos[x^2])/8 + Cos[3*x^2]/24`

3.77.3 Rubi [A] (warning: unable to verify)

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3860, 3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \sin^3(x^2) dx \\
 & \quad \downarrow \text{3860} \\
 & \frac{1}{2} \int \sin^3(x^2) dx^2 \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \sin(x^2)^3 dx^2 \\
 & \quad \downarrow \text{3113} \\
 & -\frac{1}{2} \int (1 - x^4) d \cos(x^2) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} \left(\frac{x^6}{3} - \cos(x^2) \right)
 \end{aligned}$$

input `Int[x*Sin[x^2]^3,x]`

output `(x^6/3 - Cos[x^2])/2`

3.77.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3113 Int[sin[(c_.) + (d_.)*(x_)^(n_), x_Symbol] :> Simp[-d^(-1) Subst[Int[Exp
and[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x]
&& IGtQ[(n - 1)/2, 0]
```

```
rule 3860 Int[(x_)^(m_.)*((a_.) + (b_.)*Sin[(c_.) + (d_.)*(x_)^(n_)])^(p_.), x_Symbol
] :> Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Sin[c + d*x])^
p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[
(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[
(m + 1)/n], 0]))
```

3.77.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

method	result	size
derivativedivides	$-\frac{(2+\sin^2(x^2))\cos(x^2)}{6}$	15
default	$-\frac{(2+\sin^2(x^2))\cos(x^2)}{6}$	15
risch	$-\frac{3\cos(x^2)}{8} + \frac{\cos(3x^2)}{24}$	16
parallelrisch	$-\frac{1}{3} - \frac{3\cos(x^2)}{8} + \frac{\cos(3x^2)}{24}$	17
norman	$\frac{-2\left(\tan^2\left(\frac{x^2}{2}\right)\right) - \frac{2}{3}}{\left(1 + \tan^2\left(\frac{x^2}{2}\right)\right)^3}$	26

```
input int(x*sin(x^2)^3,x,method=_RETURNVERBOSE)
```

```
output -1/6*(2+sin(x^2)^2)*cos(x^2)
```

3.77.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int x \sin^3(x^2) dx = \frac{1}{6} \cos(x^2)^3 - \frac{1}{2} \cos(x^2)$$

```
input integrate(x*sin(x^2)^3,x, algorithm="fracas")
```

output $1/6*\cos(x^2)^3 - 1/2*\cos(x^2)$

3.77.6 Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.16

$$\int x \sin^3(x^2) dx = -\frac{\sin^2(x^2) \cos(x^2)}{2} - \frac{\cos^3(x^2)}{3}$$

input `integrate(x*sin(x**2)**3,x)`

output $-\sin(x^2)**2*\cos(x^2)/2 - \cos(x^2)**3/3$

3.77.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int x \sin^3(x^2) dx = \frac{1}{24} \cos(3x^2) - \frac{3}{8} \cos(x^2)$$

input `integrate(x*sin(x^2)^3,x, algorithm="maxima")`

output $1/24*\cos(3*x^2) - 3/8*\cos(x^2)$

3.77.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int x \sin^3(x^2) dx = \frac{1}{6} \cos(x^2)^3 - \frac{1}{2} \cos(x^2)$$

input `integrate(x*sin(x^2)^3,x, algorithm="giac")`

output $1/6*\cos(x^2)^3 - 1/2*\cos(x^2)$

3.77.9 Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int x \sin^3(x^2) dx = \frac{\cos(x^2) (\cos(x^2)^2 - 3)}{6}$$

input `int(x*sin(x^2)^3,x)`

output `(cos(x^2)*(cos(x^2)^2 - 3))/6`

3.78 $\int \sin^2(x) \tan(x) dx$

3.78.1	Optimal result	492
3.78.2	Mathematica [A] (verified)	492
3.78.3	Rubi [A] (verified)	493
3.78.4	Maple [A] (verified)	494
3.78.5	Fricas [A] (verification not implemented)	494
3.78.6	Sympy [A] (verification not implemented)	495
3.78.7	Maxima [A] (verification not implemented)	495
3.78.8	Giac [A] (verification not implemented)	495
3.78.9	Mupad [B] (verification not implemented)	496

3.78.1 Optimal result

Integrand size = 7, antiderivative size = 14

$$\int \sin^2(x) \tan(x) dx = \frac{\cos^2(x)}{2} - \log(\cos(x))$$

output `1/2*cos(x)^2-ln(cos(x))`

3.78.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \sin^2(x) \tan(x) dx = \frac{\cos^2(x)}{2} - \log(\cos(x))$$

input `Integrate[Sin[x]^2*Tan[x],x]`

output `Cos[x]^2/2 - Log[Cos[x]]`

3.78.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 3070, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^2(x) \tan(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^2 \tan(x) dx \\
 & \quad \downarrow \text{3070} \\
 & - \int (1 - \cos^2(x)) \sec(x) d \cos(x) \\
 & \quad \downarrow \text{244} \\
 & - \int (\sec(x) - \cos(x)) d \cos(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\cos^2(x)}{2} - \log(\cos(x))
 \end{aligned}$$

input `Int[Sin[x]^2*Tan[x],x]`

output `Cos[x]^2/2 - Log[Cos[x]]`

3.78.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3070 Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol]
:= Simp[-f^(-1) Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f
*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]
```

3.78.4 Maple [A] (verified)

Time = 1.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
default	$-\frac{\sin^2(x)}{2} - \ln(\cos(x))$	13
risch	$ix + \frac{e^{2ix}}{8} + \frac{e^{-2ix}}{8} - \ln(e^{2ix} + 1)$	30

```
input int(cos(x)^2*tan(x)^3,x,method=_RETURNVERBOSE)
```

```
output -1/2*sin(x)^2-ln(cos(x))
```

3.78.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \sin^2(x) \tan(x) dx = \frac{1}{2} \cos(x)^2 - \log(-\cos(x))$$

```
input integrate(cos(x)^2*tan(x)^3,x, algorithm="fracas")
```

```
output 1/2*cos(x)^2 - log(-cos(x))
```

3.78.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \sin^2(x) \tan(x) dx = -\log(\cos(x)) + \frac{\cos^2(x)}{2}$$

input `integrate(cos(x)**2*tan(x)**3,x)`output `-log(cos(x)) + cos(x)**2/2`**3.78.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \sin^2(x) \tan(x) dx = -\frac{1}{2} \sin(x)^2 - \frac{1}{2} \log(\sin(x)^2 - 1)$$

input `integrate(cos(x)^2*tan(x)^3,x, algorithm="maxima")`output `-1/2*sin(x)^2 - 1/2*log(sin(x)^2 - 1)`**3.78.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

$$\int \sin^2(x) \tan(x) dx = \frac{1}{2} \cos(x)^2 - \log(|\cos(x)|)$$

input `integrate(cos(x)^2*tan(x)^3,x, algorithm="giac")`output `1/2*cos(x)^2 - log(abs(cos(x)))`

3.78.9 Mupad [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \sin^2(x) \tan(x) dx = \frac{\cos(x)^2}{2} + \frac{\ln(\tan(x)^2 + 1)}{2}$$

input `int(cos(x)^2*tan(x)^3,x)`

output `log(tan(x)^2 + 1)/2 + cos(x)^2/2`

3.79 $\int \cos^2(x) \cot^3(x) dx$

3.79.1	Optimal result	497
3.79.2	Mathematica [A] (verified)	497
3.79.3	Rubi [A] (warning: unable to verify)	498
3.79.4	Maple [A] (verified)	499
3.79.5	Fricas [B] (verification not implemented)	500
3.79.6	Sympy [A] (verification not implemented)	500
3.79.7	Maxima [A] (verification not implemented)	500
3.79.8	Giac [A] (verification not implemented)	501
3.79.9	Mupad [B] (verification not implemented)	501

3.79.1 Optimal result

Integrand size = 9, antiderivative size = 22

$$\int \cos^2(x) \cot^3(x) dx = -\frac{1}{2} \csc^2(x) - 2 \log(\sin(x)) + \frac{\sin^2(x)}{2}$$

output `-1/2*csc(x)^2-2*ln(sin(x))+1/2*sin(x)^2`

3.79.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \cos^2(x) \cot^3(x) dx = \frac{1}{2} (-\csc^2(x) - 4 \log(\sin(x)) + \sin^2(x))$$

input `Integrate[Cos[x]^2*Cot[x]^3,x]`

output `(-Csc[x]^2 - 4*Log[Sin[x]] + Sin[x]^2)/2`

3.79.3 Rubi [A] (warning: unable to verify)

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 25, 3070, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^2(x) \cot^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\sin\left(x + \frac{\pi}{2}\right)^2 \tan\left(x + \frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{25} \\
 & -\int \sin\left(x + \frac{\pi}{2}\right)^2 \tan\left(x + \frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{3070} \\
 & \int (1 - \sin^2(x))^2 (-\csc^3(x)) d(-\sin(x)) \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \int \csc^2(x) (\sin(x) + 1)^2 d\sin^2(x) \\
 & \quad \downarrow \text{49} \\
 & \frac{1}{2} \int (\csc^2(x) + 2 \csc(x) + 1) d\sin^2(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} (\sin^2(x) + \csc(x) - 2 \log(\sin^2(x)))
 \end{aligned}$$

input `Int[Cos[x]^2*Cot[x]^3,x]`

output `(Csc[x] - 2*Log[Sin[x]^2] + Sin[x]^2)/2`

3.79.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3070 `Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[-f^(-1) Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]`

3.79.4 Maple [A] (verified)

Time = 3.38 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.32

method	result	size
default	$-\frac{\cos^6(x)}{2\sin(x)^2} - \frac{\cos^4(x)}{2} - \cos^2(x) - 2\ln(\sin(x))$	29
risch	$2ix - \frac{e^{2ix}}{8} - \frac{e^{-2ix}}{8} + \frac{2e^{2ix}}{(e^{2ix}-1)^2} - 2\ln(e^{2ix}-1)$	46

input `int(cot(x)^5*sin(x)^2,x,method=_RETURNVERBOSE)`

output `-1/2/sin(x)^2*cos(x)^6-1/2*cos(x)^4-cos(x)^2-2*ln(sin(x))`

3.79.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. $2(18) = 36$.

Time = 0.26 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.68

$$\int \cos^2(x) \cot^3(x) dx = -\frac{2 \cos(x)^4 - 3 \cos(x)^2 + 8 (\cos(x)^2 - 1) \log\left(\frac{1}{2} \sin(x)\right) - 1}{4 (\cos(x)^2 - 1)}$$

input `integrate(cot(x)^5*sin(x)^2,x, algorithm="fracas")`

output `-1/4*(2*cos(x)^4 - 3*cos(x)^2 + 8*(cos(x)^2 - 1)*log(1/2*sin(x)) - 1)/(cos(x)^2 - 1)`

3.79.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \cos^2(x) \cot^3(x) dx = -2 \log(\sin(x)) + \frac{\sin^2(x)}{2} - \frac{1}{2 \sin^2(x)}$$

input `integrate(cot(x)**5*sin(x)**2,x)`

output `-2*log(sin(x)) + sin(x)**2/2 - 1/(2*sin(x)**2)`

3.79.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \cos^2(x) \cot^3(x) dx = \frac{1}{2} \sin(x)^2 - \frac{1}{2 \sin(x)^2} - \log(\sin(x)^2)$$

input `integrate(cot(x)^5*sin(x)^2,x, algorithm="maxima")`

output `1/2*sin(x)^2 - 1/2/sin(x)^2 - log(sin(x)^2)`

3.79.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.27

$$\int \cos^2(x) \cot^3(x) dx = -\frac{1}{2} \cos(x)^2 + \frac{1}{2(\cos(x)^2 - 1)} - \log(-\cos(x)^2 + 1)$$

input `integrate(cot(x)^5*sin(x)^2,x, algorithm="giac")`output `-1/2*cos(x)^2 + 1/2/(cos(x)^2 - 1) - log(-cos(x)^2 + 1)`**3.79.9 Mupad [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.45

$$\int \cos^2(x) \cot^3(x) dx = \ln(\tan(x)^2 + 1) - 2 \ln(\tan(x)) - \frac{\tan(x)^2 + \frac{1}{2}}{\tan(x)^4 + \tan(x)^2}$$

input `int(cot(x)^5*sin(x)^2,x)`output `log(tan(x)^2 + 1) - 2*log(tan(x)) - (tan(x)^2 + 1/2)/(tan(x)^2 + tan(x)^4)`

3.80 $\int \sec(x)(1 - \sin(x)) dx$

3.80.1	Optimal result	502
3.80.2	Mathematica [A] (verified)	502
3.80.3	Rubi [A] (verified)	503
3.80.4	Maple [A] (verified)	504
3.80.5	Fricas [A] (verification not implemented)	504
3.80.6	Sympy [B] (verification not implemented)	505
3.80.7	Maxima [A] (verification not implemented)	505
3.80.8	Giac [A] (verification not implemented)	505
3.80.9	Mupad [B] (verification not implemented)	506

3.80.1 Optimal result

Integrand size = 9, antiderivative size = 5

$$\int \sec(x)(1 - \sin(x)) dx = \log(1 + \sin(x))$$

output `ln(1+sin(x))`

3.80.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.40

$$\int \sec(x)(1 - \sin(x)) dx = \operatorname{arctanh}(\sin(x)) + \log(\cos(x))$$

input `Integrate[Sec[x]*(1 - Sin[x]),x]`

output `ArcTanh[Sin[x]] + Log[Cos[x]]`

3.80.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3146, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (1 - \sin(x)) \sec(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1 - \sin(x)}{\cos(x)} dx \\
 & \quad \downarrow \text{3146} \\
 & - \int \frac{1}{\sin(x) + 1} d(-\sin(x)) \\
 & \quad \downarrow \text{16} \\
 & \log(\sin(x) + 1)
 \end{aligned}$$

input `Int[Sec[x]*(1 - Sin[x]),x]`

output `Log[1 + Sin[x]]`

3.80.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`


```
rule 3146 Int[cos[(e_.) + (f_.)*(x_.)]^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)])^(m
_.), x_Symbol] :> Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x
)^(p - 1)/2], x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && I
ntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/
2])
```

3.80.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
derivativedivides	$\ln(\sin(x) + 1)$	6
default	$\ln(\sin(x) + 1)$	6
risch	$-ix + 2 \ln(i + e^{ix})$	17
norman	$2 \ln(1 + \tan(\frac{x}{2})) - \ln(1 + \tan^2(\frac{x}{2}))$	22
parallelrisch	$2 \ln(-\cot(x) + 1 + \csc(x)) - \ln\left(\frac{2}{\cos(x)+1}\right)$	24

```
input int((-sin(x)+1)/cos(x),x,method=_RETURNVERBOSE)
```

```
output ln(sin(x)+1)
```

3.80.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \sec(x)(1 - \sin(x)) dx = \log(\sin(x) + 1)$$

```
input integrate((1-sin(x))/cos(x),x, algorithm="fricas")
```

```
output log(sin(x) + 1)
```

3.80.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(5) = 10$.

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 3.80

$$\int \sec(x)(1 - \sin(x)) dx = 2 \log \left(\tan \left(\frac{x}{2} \right) + 1 \right) - \log \left(\tan^2 \left(\frac{x}{2} \right) + 1 \right)$$

input `integrate((1-sin(x))/cos(x),x)`

output `2*log(tan(x/2) + 1) - log(tan(x/2)**2 + 1)`

3.80.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \sec(x)(1 - \sin(x)) dx = \log(\sin(x) + 1)$$

input `integrate((1-sin(x))/cos(x),x, algorithm="maxima")`

output `log(sin(x) + 1)`

3.80.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \sec(x)(1 - \sin(x)) dx = \log(\sin(x) + 1)$$

input `integrate((1-sin(x))/cos(x),x, algorithm="giac")`

output `log(sin(x) + 1)`

3.80.9 Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \sec(x)(1 - \sin(x)) dx = \ln(\sin(x) + 1)$$

input `int(-(sin(x) - 1)/cos(x),x)`

output `log(sin(x) + 1)`

3.81 $\int \frac{1}{1-\sin(x)} dx$

3.81.1	Optimal result	507
3.81.2	Mathematica [B] (verified)	507
3.81.3	Rubi [A] (verified)	508
3.81.4	Maple [A] (verified)	509
3.81.5	Fricas [A] (verification not implemented)	509
3.81.6	Sympy [A] (verification not implemented)	509
3.81.7	Maxima [A] (verification not implemented)	510
3.81.8	Giac [A] (verification not implemented)	510
3.81.9	Mupad [B] (verification not implemented)	510

3.81.1 Optimal result

Integrand size = 8, antiderivative size = 11

$$\int \frac{1}{1-\sin(x)} dx = \frac{\cos(x)}{1-\sin(x)}$$

output `cos(x)/(1-sin(x))`

3.81.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 25 vs. 2(11) = 22.

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 2.27

$$\int \frac{1}{1-\sin(x)} dx = \frac{2 \sin\left(\frac{x}{2}\right)}{\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)}$$

input `Integrate[(1 - Sin[x])^(-1),x]`

output `(2*Sin[x/2])/(Cos[x/2] - Sin[x/2])`

3.81.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{1 - \sin(x)} dx$$

↓ 3042

$$\int \frac{1}{1 - \sin(x)} dx$$

↓ 3127

$$\frac{\cos(x)}{1 - \sin(x)}$$

input `Int[(1 - Sin[x])^(-1),x]`

output `Cos[x]/(1 - Sin[x])`

3.81.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

3.81.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

method	result	size
default	$-\frac{2}{\tan(\frac{x}{2})-1}$	11
norman	$-\frac{2}{\tan(\frac{x}{2})-1}$	11
parallelrisch	$-\frac{2}{\tan(\frac{x}{2})-1}$	11
risch	$\frac{2}{e^{ix}-i}$	13

input `int(1/(-sin(x)+1),x,method=_RETURNVERBOSE)`output `-2/(tan(1/2*x)-1)`**3.81.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.55

$$\int \frac{1}{1-\sin(x)} dx = \frac{\cos(x) + \sin(x) + 1}{\cos(x) - \sin(x) + 1}$$

input `integrate(1/(1-sin(x)),x, algorithm="fracas")`output `(cos(x) + sin(x) + 1)/(cos(x) - sin(x) + 1)`**3.81.6 Sympy [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \frac{1}{1-\sin(x)} dx = -\frac{2}{\tan(\frac{x}{2})-1}$$

input `integrate(1/(1-sin(x)),x)`output `-2/(tan(x/2) - 1)`

3.81.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.36

$$\int \frac{1}{1 - \sin(x)} dx = -\frac{2}{\frac{\sin(x)}{\cos(x)+1} - 1}$$

input `integrate(1/(1-sin(x)),x, algorithm="maxima")`output `-2/(sin(x)/(cos(x) + 1) - 1)`**3.81.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int \frac{1}{1 - \sin(x)} dx = -\frac{2}{\tan\left(\frac{1}{2}x\right) - 1}$$

input `integrate(1/(1-sin(x)),x, algorithm="giac")`output `-2/(tan(1/2*x) - 1)`**3.81.9 Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int \frac{1}{1 - \sin(x)} dx = -\frac{2}{\tan\left(\frac{x}{2}\right) - 1}$$

input `int(-1/(sin(x) - 1),x)`output `-2/(tan(x/2) - 1)`

3.82 $\int \tan^2(x) dx$

3.82.1	Optimal result	511
3.82.2	Mathematica [A] (verified)	511
3.82.3	Rubi [A] (verified)	512
3.82.4	Maple [A] (verified)	513
3.82.5	Fricas [A] (verification not implemented)	513
3.82.6	Sympy [B] (verification not implemented)	513
3.82.7	Maxima [A] (verification not implemented)	514
3.82.8	Giac [A] (verification not implemented)	514
3.82.9	Mupad [B] (verification not implemented)	514

3.82.1 Optimal result

Integrand size = 4, antiderivative size = 6

$$\int \tan^2(x) dx = -x + \tan(x)$$

output `-x+tan(x)`

3.82.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.33

$$\int \tan^2(x) dx = -\arctan(\tan(x)) + \tan(x)$$

input `Integrate[Tan[x]^2,x]`

output `-ArcTan[Tan[x]] + Tan[x]`

3.82.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \tan^2(x) dx \\
 \downarrow 3042 \\
 \int \tan(x)^2 dx \\
 \downarrow 3954 \\
 \tan(x) - \int 1 dx \\
 \downarrow 24 \\
 \tan(x) - x
 \end{array}$$

input `Int[Tan[x]^2,x]`

output `-x + Tan[x]`

3.82.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

3.82.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

method	result	size
norman	$-x + \tan(x)$	7
parallelrisch	$-x + \tan(x)$	7
derivativedivides	$\tan(x) - \arctan(\tan(x))$	9
default	$\tan(x) - \arctan(\tan(x))$	9
risch	$-x + \frac{2i}{e^{2ix}+1}$	17

input `int(tan(x)^2,x,method=_RETURNVERBOSE)`

output `-x+tan(x)`

3.82.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \tan^2(x) dx = -x + \tan(x)$$

input `integrate(tan(x)^2,x, algorithm="fricas")`

output `-x + tan(x)`

3.82.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7 vs. $2(3) = 6$.

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

$$\int \tan^2(x) dx = -x + \frac{\sin(x)}{\cos(x)}$$

input `integrate(tan(x)**2,x)`

output `-x + sin(x)/cos(x)`

3.82.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \tan^2(x) dx = -x + \tan(x)$$

input `integrate(tan(x)^2,x, algorithm="maxima")`

output `-x + tan(x)`

3.82.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \tan^2(x) dx = -x + \tan(x)$$

input `integrate(tan(x)^2,x, algorithm="giac")`

output `-x + tan(x)`

3.82.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \tan^2(x) dx = \tan(x) - x$$

input `int(tan(x)^2,x)`

output `tan(x) - x`

3.83 $\int \tan^4(x) dx$

3.83.1	Optimal result	515
3.83.2	Mathematica [A] (verified)	515
3.83.3	Rubi [A] (verified)	516
3.83.4	Maple [A] (verified)	517
3.83.5	Fricas [A] (verification not implemented)	518
3.83.6	Sympy [A] (verification not implemented)	518
3.83.7	Maxima [A] (verification not implemented)	518
3.83.8	Giac [A] (verification not implemented)	519
3.83.9	Mupad [B] (verification not implemented)	519

3.83.1 Optimal result

Integrand size = 4, antiderivative size = 14

$$\int \tan^4(x) dx = x - \tan(x) + \frac{\tan^3(x)}{3}$$

output `x-tan(x)+1/3*tan(x)^3`

3.83.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \tan^4(x) dx = \arctan(\tan(x)) - \tan(x) + \frac{\tan^3(x)}{3}$$

input `Integrate[Tan[x]^4,x]`

output `ArcTan[Tan[x]] - Tan[x] + Tan[x]^3/3`

3.83.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3042, 3954, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \tan^4(x) dx \\
 \downarrow 3042 \\
 \int \tan(x)^4 dx \\
 \downarrow 3954 \\
 \frac{\tan^3(x)}{3} - \int \tan^2(x) dx \\
 \downarrow 3042 \\
 \frac{\tan^3(x)}{3} - \int \tan(x)^2 dx \\
 \downarrow 3954 \\
 \int 1 dx + \frac{\tan^3(x)}{3} - \tan(x) \\
 \downarrow 24 \\
 x + \frac{\tan^3(x)}{3} - \tan(x)
 \end{array}$$

input `Int [Tan [x] ^4, x]`

output `x - Tan [x] + Tan [x] ^3/3`

3.83.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

3.83.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
norman	$x - \tan(x) + \frac{\tan^3(x)}{3}$	13
parallelrisch	$x - \tan(x) + \frac{\tan^3(x)}{3}$	13
derivativedivides	$\frac{\tan^3(x)}{3} - \tan(x) + \arctan(\tan(x))$	15
default	$\frac{\tan^3(x)}{3} - \tan(x) + \arctan(\tan(x))$	15
risch	$x - \frac{4i(3e^{4ix} + 3e^{2ix} + 2)}{3(e^{2ix} + 1)^3}$	31

input `int(tan(x)^4,x,method=_RETURNVERBOSE)`

output `x-tan(x)+1/3*tan(x)^3`

3.83.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(x) dx = \frac{1}{3} \tan(x)^3 + x - \tan(x)$$

input `integrate(tan(x)^4,x, algorithm="fricas")`

output `1/3*tan(x)^3 + x - tan(x)`

3.83.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.36

$$\int \tan^4(x) dx = x + \frac{\sin^3(x)}{3 \cos^3(x)} - \frac{\sin(x)}{\cos(x)}$$

input `integrate(tan(x)**4,x)`

output `x + sin(x)**3/(3*cos(x)**3) - sin(x)/cos(x)`

3.83.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(x) dx = \frac{1}{3} \tan(x)^3 + x - \tan(x)$$

input `integrate(tan(x)^4,x, algorithm="maxima")`

output `1/3*tan(x)^3 + x - tan(x)`

3.83.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(x) dx = \frac{1}{3} \tan(x)^3 + x - \tan(x)$$

input `integrate(tan(x)^4,x, algorithm="giac")`

output `1/3*tan(x)^3 + x - tan(x)`

3.83.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(x) dx = \frac{\tan(x)^3}{3} - \tan(x) + x$$

input `int(tan(x)^4,x)`

output `x - tan(x) + tan(x)^3/3`

3.84 $\int \sec^4(x) dx$

3.84.1	Optimal result	520
3.84.2	Mathematica [A] (verified)	520
3.84.3	Rubi [A] (verified)	521
3.84.4	Maple [A] (verified)	522
3.84.5	Fricas [A] (verification not implemented)	522
3.84.6	Sympy [B] (verification not implemented)	522
3.84.7	Maxima [A] (verification not implemented)	523
3.84.8	Giac [A] (verification not implemented)	523
3.84.9	Mupad [B] (verification not implemented)	523

3.84.1 Optimal result

Integrand size = 4, antiderivative size = 11

$$\int \sec^4(x) dx = \tan(x) + \frac{\tan^3(x)}{3}$$

output `tan(x)+1/3*tan(x)^3`

3.84.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \sec^4(x) dx = \tan(x) + \frac{\tan^3(x)}{3}$$

input `Integrate[Sec[x]^4,x]`

output `Tan[x] + Tan[x]^3/3`

3.84.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \sec^4(x) dx \\
 \downarrow 3042 \\
 \int \csc\left(x + \frac{\pi}{2}\right)^4 dx \\
 \downarrow 4254 \\
 - \int (\tan^2(x) + 1) d(-\tan(x)) \\
 \downarrow 2009 \\
 \frac{\tan^3(x)}{3} + \tan(x)
 \end{array}$$

input `Int[Sec[x]^4,x]`

output `Tan[x] + Tan[x]^3/3`

3.84.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

3.84.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

method	result	size
parallelrisch	$\frac{\tan(x)(2+\sec^2(x))}{3}$	11
default	$-\left(-\frac{2}{3} - \frac{(\sec^2(x))}{3}\right) \tan(x)$	13
risch	$\frac{4i(3e^{2ix}+1)}{3(e^{2ix}+1)^3}$	22
norman	$\frac{4\left(\tan^3\left(\frac{x}{2}\right)\right) - 2\left(\tan^5\left(\frac{x}{2}\right)\right) - 2\tan\left(\frac{x}{2}\right)}{\left(\tan^2\left(\frac{x}{2}\right) - 1\right)^3}$	35

input `int(sec(x)^4,x,method=_RETURNVERBOSE)`

output `1/3*tan(x)*(2+sec(x)^2)`

3.84.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.45

$$\int \sec^4(x) dx = \frac{(2 \cos(x)^2 + 1) \sin(x)}{3 \cos(x)^3}$$

input `integrate(sec(x)^4,x, algorithm="fricas")`

output `1/3*(2*cos(x)^2 + 1)*sin(x)/cos(x)^3`

3.84.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. 2(8) = 16.

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.73

$$\int \sec^4(x) dx = \frac{2 \sin(x)}{3 \cos(x)} + \frac{\sin(x)}{3 \cos^3(x)}$$

input `integrate(sec(x)**4,x)`

output `2*sin(x)/(3*cos(x)) + sin(x)/(3*cos(x)**3)`

3.84.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \sec^4(x) dx = \frac{1}{3} \tan(x)^3 + \tan(x)$$

input `integrate(sec(x)^4,x, algorithm="maxima")`

output `1/3*tan(x)^3 + tan(x)`

3.84.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \sec^4(x) dx = \frac{1}{3} \tan(x)^3 + \tan(x)$$

input `integrate(sec(x)^4,x, algorithm="giac")`

output `1/3*tan(x)^3 + tan(x)`

3.84.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.55

$$\int \sec^4(x) dx = \frac{2 \sin(x) \cos(x)^2 + \sin(x)}{3 \cos(x)^3}$$

input `int(1/cos(x)^4,x)`

output `(sin(x) + 2*cos(x)^2*sin(x))/(3*cos(x)^3)`

3.85 $\int \sec^6(x) dx$

3.85.1	Optimal result	524
3.85.2	Mathematica [A] (verified)	524
3.85.3	Rubi [A] (verified)	525
3.85.4	Maple [A] (verified)	526
3.85.5	Fricas [A] (verification not implemented)	526
3.85.6	Sympy [A] (verification not implemented)	526
3.85.7	Maxima [A] (verification not implemented)	527
3.85.8	Giac [A] (verification not implemented)	527
3.85.9	Mupad [B] (verification not implemented)	527

3.85.1 Optimal result

Integrand size = 4, antiderivative size = 19

$$\int \sec^6(x) dx = \tan(x) + \frac{2 \tan^3(x)}{3} + \frac{\tan^5(x)}{5}$$

output `tan(x)+2/3*tan(x)^3+1/5*tan(x)^5`

3.85.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \sec^6(x) dx = \tan(x) + \frac{2 \tan^3(x)}{3} + \frac{\tan^5(x)}{5}$$

input `Integrate[Sec[x]^6,x]`

output `Tan[x] + (2*Tan[x]^3)/3 + Tan[x]^5/5`

3.85.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^6(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(x + \frac{\pi}{2}\right)^6 dx \\
 & \quad \downarrow \text{4254} \\
 & - \int (\tan^4(x) + 2\tan^2(x) + 1) d(-\tan(x)) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\tan^5(x)}{5} + \frac{2\tan^3(x)}{3} + \tan(x)
 \end{aligned}$$

input `Int[Sec[x]^6,x]`

output `Tan[x] + (2*Tan[x]^3)/3 + Tan[x]^5/5`

3.85.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

3.85.4 Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

method	result	size
default	$-\left(-\frac{8}{15} - \frac{\sec^4(x)}{5} - \frac{4(\sec^2(x))}{15}\right) \tan(x)$	19
parallelrisc	$\frac{\tan(x)(3(\sec^4(x))+4(\sec^2(x))+8)}{15}$	19
risc	$\frac{16i(10e^{4ix}+5e^{2ix}+1)}{15(e^{2ix}+1)^5}$	29
norman	$\frac{\frac{8(\tan^3(\frac{x}{2}))}{3} - \frac{116(\tan^5(\frac{x}{2}))}{15} + \frac{8(\tan^7(\frac{x}{2}))}{3} - 2(\tan^9(\frac{x}{2})) - 2\tan(\frac{x}{2})}{(\tan^2(\frac{x}{2})-1)^5}$	51

input `int(sec(x)^6,x,method=_RETURNVERBOSE)`output `-(-8/15-1/5*sec(x)^4-4/15*sec(x)^2)*tan(x)`**3.85.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.16

$$\int \sec^6(x) dx = \frac{(8 \cos(x)^4 + 4 \cos(x)^2 + 3) \sin(x)}{15 \cos(x)^5}$$

input `integrate(sec(x)^6,x, algorithm="fricas")`output `1/15*(8*cos(x)^4 + 4*cos(x)^2 + 3)*sin(x)/cos(x)^5`**3.85.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.63

$$\int \sec^6(x) dx = \frac{8 \sin(x)}{15 \cos(x)} + \frac{4 \sin(x)}{15 \cos^3(x)} + \frac{\sin(x)}{5 \cos^5(x)}$$

input `integrate(sec(x)**6,x)`output `8*sin(x)/(15*cos(x)) + 4*sin(x)/(15*cos(x)**3) + sin(x)/(5*cos(x)**5)`

3.85.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \sec^6(x) dx = \frac{1}{5} \tan(x)^5 + \frac{2}{3} \tan(x)^3 + \tan(x)$$

input `integrate(sec(x)^6,x, algorithm="maxima")`output `1/5*tan(x)^5 + 2/3*tan(x)^3 + tan(x)`**3.85.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \sec^6(x) dx = \frac{1}{5} \tan(x)^5 + \frac{2}{3} \tan(x)^3 + \tan(x)$$

input `integrate(sec(x)^6,x, algorithm="giac")`output `1/5*tan(x)^5 + 2/3*tan(x)^3 + tan(x)`**3.85.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.42

$$\int \sec^6(x) dx = \frac{8 \sin(x) \cos(x)^4 + 4 \sin(x) \cos(x)^2 + 3 \sin(x)}{15 \cos(x)^5}$$

input `int(1/cos(x)^6,x)`output `(3*sin(x) + 4*cos(x)^2*sin(x) + 8*cos(x)^4*sin(x))/(15*cos(x)^5)`

3.86 $\int \sec^2(x) \tan^4(x) dx$

3.86.1	Optimal result	528
3.86.2	Mathematica [A] (verified)	528
3.86.3	Rubi [A] (verified)	529
3.86.4	Maple [A] (verified)	530
3.86.5	Fricas [B] (verification not implemented)	530
3.86.6	Sympy [B] (verification not implemented)	530
3.86.7	Maxima [A] (verification not implemented)	531
3.86.8	Giac [A] (verification not implemented)	531
3.86.9	Mupad [B] (verification not implemented)	531

3.86.1 Optimal result

Integrand size = 9, antiderivative size = 8

$$\int \sec^2(x) \tan^4(x) dx = \frac{\tan^5(x)}{5}$$

output `1/5*tan(x)^5`

3.86.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \sec^2(x) \tan^4(x) dx = \frac{\tan^5(x)}{5}$$

input `Integrate[Sec[x]^2*Tan[x]^4,x]`

output `Tan[x]^5/5`

3.86.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3087, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan^4(x) \sec^2(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(x)^4 \sec(x)^2 dx \\ & \quad \downarrow \text{3087} \\ & \int \tan^4(x) d \tan(x) \\ & \quad \downarrow \text{15} \\ & \frac{\tan^5(x)}{5} \end{aligned}$$

input `Int[Sec[x]^2*Tan[x]^4,x]`

output `Tan[x]^5/5`

3.86.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3087 `Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[1/f Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])`

3.86.4 Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{(\tan^5(x))}{5}$	7
default	$\frac{(\tan^5(x))}{5}$	7
risch	$\frac{2i(5e^{8ix}+10e^{4ix}+1)}{5(e^{2ix}+1)^5}$	29

input `int(sec(x)^2*tan(x)^4,x,method=_RETURNVERBOSE)`

output `1/5*tan(x)^5`

3.86.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. 2(6) = 12.

Time = 0.25 (sec) , antiderivative size = 20, normalized size of antiderivative = 2.50

$$\int \sec^2(x) \tan^4(x) dx = \frac{(\cos(x)^4 - 2 \cos(x)^2 + 1) \sin(x)}{5 \cos(x)^5}$$

input `integrate(sec(x)^2*tan(x)^4,x, algorithm="fricas")`

output `1/5*(cos(x)^4 - 2*cos(x)^2 + 1)*sin(x)/cos(x)^5`

3.86.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. 2(5) = 10.

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 3.62

$$\int \sec^2(x) \tan^4(x) dx = \frac{\sin(x)}{5 \cos(x)} - \frac{2 \sin(x)}{5 \cos^3(x)} + \frac{\sin(x)}{5 \cos^5(x)}$$

input `integrate(sec(x)**2*tan(x)**4,x)`

output `sin(x)/(5*cos(x)) - 2*sin(x)/(5*cos(x)**3) + sin(x)/(5*cos(x)**5)`

3.86.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan^4(x) dx = \frac{1}{5} \tan(x)^5$$

input `integrate(sec(x)^2*tan(x)^4,x, algorithm="maxima")`output `1/5*tan(x)^5`**3.86.8 Giac [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan^4(x) dx = \frac{1}{5} \tan(x)^5$$

input `integrate(sec(x)^2*tan(x)^4,x, algorithm="giac")`output `1/5*tan(x)^5`**3.86.9 Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan^4(x) dx = \frac{\tan(x)^5}{5}$$

input `int(tan(x)^4/cos(x)^2,x)`output `tan(x)^5/5`

3.87 $\int \sec^4(x) \tan^2(x) dx$

3.87.1	Optimal result	532
3.87.2	Mathematica [A] (verified)	532
3.87.3	Rubi [A] (verified)	533
3.87.4	Maple [A] (verified)	534
3.87.5	Fricas [A] (verification not implemented)	534
3.87.6	Sympy [B] (verification not implemented)	535
3.87.7	Maxima [A] (verification not implemented)	535
3.87.8	Giac [A] (verification not implemented)	535
3.87.9	Mupad [B] (verification not implemented)	536

3.87.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \sec^4(x) \tan^2(x) dx = \frac{\tan^3(x)}{3} + \frac{\tan^5(x)}{5}$$

output `1/3*tan(x)^3+1/5*tan(x)^5`

3.87.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \sec^4(x) \tan^2(x) dx = -\frac{2 \tan(x)}{15} - \frac{1}{15} \sec^2(x) \tan(x) + \frac{1}{5} \sec^4(x) \tan(x)$$

input `Integrate[Sec[x]^4*Tan[x]^2,x]`

output `(-2*Tan[x])/15 - (Sec[x]^2*Tan[x])/15 + (Sec[x]^4*Tan[x])/5`

3.87.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 3087, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^2(x) \sec^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^2 \sec(x)^4 dx \\
 & \quad \downarrow \text{3087} \\
 & \int \tan^2(x) (\tan^2(x) + 1) d \tan(x) \\
 & \quad \downarrow \text{244} \\
 & \int (\tan^4(x) + \tan^2(x)) d \tan(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\tan^5(x)}{5} + \frac{\tan^3(x)}{3}
 \end{aligned}$$

input `Int[Sec[x]^4*Tan[x]^2,x]`

output `Tan[x]^3/3 + Tan[x]^5/5`

3.87.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] :> Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3087 Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_S
ymbol] := Simp[1/f Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e +
f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)
/2] && LtQ[0, n, m - 1])
```

3.87.4 Maple [A] (verified)

Time = 1.95 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$\frac{\tan^3(x)}{3} + \frac{\tan^5(x)}{5}$	14
default	$\frac{\tan^3(x)}{3} + \frac{\tan^5(x)}{5}$	14
risch	$-\frac{4i(15e^{6ix} - 5e^{4ix} + 5e^{2ix} + 1)}{15(e^{2ix} + 1)^5}$	36

```
input int(sec(x)^4*tan(x)^2,x,method=_RETURNVERBOSE)
```

```
output 1/3*tan(x)^3+1/5*tan(x)^5
```

3.87.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \sec^4(x) \tan^2(x) dx = -\frac{(2 \cos(x)^4 + \cos(x)^2 - 3) \sin(x)}{15 \cos(x)^5}$$

```
input integrate(sec(x)^4*tan(x)^2,x, algorithm="fracas")
```

```
output -1/15*(2*cos(x)^4 + cos(x)^2 - 3)*sin(x)/cos(x)^5
```

3.87.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(12) = 24$.

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.71

$$\int \sec^4(x) \tan^2(x) dx = -\frac{2 \sin(x)}{15 \cos(x)} - \frac{\sin(x)}{15 \cos^3(x)} + \frac{\sin(x)}{5 \cos^5(x)}$$

input `integrate(sec(x)**4*tan(x)**2,x)`

output `-2*sin(x)/(15*cos(x)) - sin(x)/(15*cos(x)**3) + sin(x)/(5*cos(x)**5)`

3.87.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sec^4(x) \tan^2(x) dx = \frac{1}{5} \tan(x)^5 + \frac{1}{3} \tan(x)^3$$

input `integrate(sec(x)^4*tan(x)^2,x, algorithm="maxima")`

output `1/5*tan(x)^5 + 1/3*tan(x)^3`

3.87.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sec^4(x) \tan^2(x) dx = \frac{1}{5} \tan(x)^5 + \frac{1}{3} \tan(x)^3$$

input `integrate(sec(x)^4*tan(x)^2,x, algorithm="giac")`

output `1/5*tan(x)^5 + 1/3*tan(x)^3`

3.87.9 Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sec^4(x) \tan^2(x) dx = \frac{\tan(x)^5}{5} + \frac{\tan(x)^3}{3}$$

input `int(tan(x)^2/cos(x)^4,x)`

output `tan(x)^3/3 + tan(x)^5/5`

3.88 $\int \sec^3(x) \tan(x) dx$

3.88.1	Optimal result	537
3.88.2	Mathematica [A] (verified)	537
3.88.3	Rubi [A] (verified)	538
3.88.4	Maple [A] (verified)	539
3.88.5	Fricas [A] (verification not implemented)	539
3.88.6	Sympy [A] (verification not implemented)	539
3.88.7	Maxima [A] (verification not implemented)	540
3.88.8	Giac [A] (verification not implemented)	540
3.88.9	Mupad [B] (verification not implemented)	540

3.88.1 Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \sec^3(x) \tan(x) dx = \frac{\sec^3(x)}{3}$$

output `1/3*sec(x)^3`

3.88.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \sec^3(x) \tan(x) dx = \frac{\sec^3(x)}{3}$$

input `Integrate[Sec[x]^3*Tan[x],x]`

output `Sec[x]^3/3`

3.88.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3086, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan(x) \sec^3(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(x) \sec(x)^3 dx \\ & \quad \downarrow \text{3086} \\ & \int \sec^2(x) d \sec(x) \\ & \quad \downarrow \text{15} \\ & \frac{\sec^3(x)}{3} \end{aligned}$$

input `Int[Sec[x]^3*Tan[x],x]`

output `Sec[x]^3/3`

3.88.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

3.88.4 Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{(\sec^3(x))}{3}$	7
default	$\frac{(\sec^3(x))}{3}$	7
risch	$\frac{8 e^{3ix}}{3(e^{2ix}+1)^3}$	17

input `int(sec(x)^3*tan(x),x,method=_RETURNVERBOSE)`

output `1/3*sec(x)^3`

3.88.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^3(x) \tan(x) dx = \frac{1}{3 \cos^3(x)}$$

input `integrate(sec(x)^3*tan(x),x, algorithm="fricas")`

output `1/3/cos(x)^3`

3.88.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \sec^3(x) \tan(x) dx = \frac{1}{3 \cos^3(x)}$$

input `integrate(sec(x)**3*tan(x),x)`

output `1/(3*cos(x)**3)`

3.88.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^3(x) \tan(x) dx = \frac{1}{3 \cos(x)^3}$$

input `integrate(sec(x)^3*tan(x),x, algorithm="maxima")`output `1/3/cos(x)^3`**3.88.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^3(x) \tan(x) dx = \frac{1}{3 \cos(x)^3}$$

input `integrate(sec(x)^3*tan(x),x, algorithm="giac")`output `1/3/cos(x)^3`**3.88.9 Mupad [B] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^3(x) \tan(x) dx = \frac{1}{3 \cos(x)^3}$$

input `int(tan(x)/cos(x)^3,x)`output `1/(3*cos(x)^3)`

3.89 $\int \sec^3(x) \tan^3(x) dx$

3.89.1	Optimal result	541
3.89.2	Mathematica [A] (verified)	541
3.89.3	Rubi [A] (verified)	542
3.89.4	Maple [A] (verified)	543
3.89.5	Fricas [A] (verification not implemented)	544
3.89.6	Sympy [A] (verification not implemented)	544
3.89.7	Maxima [A] (verification not implemented)	544
3.89.8	Giac [A] (verification not implemented)	545
3.89.9	Mupad [B] (verification not implemented)	545

3.89.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \sec^3(x) \tan^3(x) dx = -\frac{1}{3} \sec^3(x) + \frac{\sec^5(x)}{5}$$

output `-1/3*sec(x)^3+1/5*sec(x)^5`

3.89.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sec^3(x) \tan^3(x) dx = -\frac{1}{3} \sec^3(x) + \frac{\sec^5(x)}{5}$$

input `Integrate[Sec[x]^3*Tan[x]^3,x]`

output `-1/3*Sec[x]^3 + Sec[x]^5/5`

3.89.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 3086, 25, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^3(x) \sec^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^3 \sec(x)^3 dx \\
 & \quad \downarrow \text{3086} \\
 & \int -\sec^2(x) (1 - \sec^2(x)) d \sec(x) \\
 & \quad \downarrow \text{25} \\
 & - \int \sec^2(x) (1 - \sec^2(x)) d \sec(x) \\
 & \quad \downarrow \text{244} \\
 & - \int (\sec^2(x) - \sec^4(x)) d \sec(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sec^5(x)}{5} - \frac{\sec^3(x)}{3}
 \end{aligned}$$

input `Int [Sec [x] ^3*Tan [x] ^3,x]`

output `-1/3*Sec [x] ^3 + Sec [x] ^5/5`

3.89.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 244 `Int[((c_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

3.89.4 Maple [A] (verified)

Time = 1.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$-\frac{(\sec^3(x))}{3} + \frac{(\sec^5(x))}{5}$	14
default	$-\frac{(\sec^3(x))}{3} + \frac{(\sec^5(x))}{5}$	14
risch	$-\frac{8(5e^{7ix} - 2e^{5ix} + 5e^{3ix})}{15(e^{2ix} + 1)^5}$	34

input `int(sec(x)^3*tan(x)^3,x,method=_RETURNVERBOSE)`

output `-1/3*sec(x)^3+1/5*sec(x)^5`

3.89.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^3(x) \tan^3(x) dx = -\frac{5 \cos(x)^2 - 3}{15 \cos(x)^5}$$

input `integrate(sec(x)^3*tan(x)^3,x, algorithm="fracas")`output `-1/15*(5*cos(x)^2 - 3)/cos(x)^5`**3.89.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^3(x) \tan^3(x) dx = \frac{3 - 5 \cos^2(x)}{15 \cos^5(x)}$$

input `integrate(sec(x)**3*tan(x)**3,x)`output `(3 - 5*cos(x)**2)/(15*cos(x)**5)`**3.89.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^3(x) \tan^3(x) dx = -\frac{5 \cos(x)^2 - 3}{15 \cos(x)^5}$$

input `integrate(sec(x)^3*tan(x)^3,x, algorithm="maxima")`output `-1/15*(5*cos(x)^2 - 3)/cos(x)^5`

3.89.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^3(x) \tan^3(x) dx = -\frac{5 \cos(x)^2 - 3}{15 \cos(x)^5}$$

input `integrate(sec(x)^3*tan(x)^3,x, algorithm="giac")`output `-1/15*(5*cos(x)^2 - 3)/cos(x)^5`**3.89.9 Mupad [B] (verification not implemented)**

Time = 0.45 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sec^3(x) \tan^3(x) dx = \frac{1}{5 \cos(x)^5} - \frac{1}{3 \cos(x)^3}$$

input `int(tan(x)^3/cos(x)^3,x)`output `1/(5*cos(x)^5) - 1/(3*cos(x)^3)`

3.90 $\int \tan^5(x) dx$

3.90.1	Optimal result	546
3.90.2	Mathematica [A] (verified)	546
3.90.3	Rubi [A] (verified)	547
3.90.4	Maple [A] (verified)	548
3.90.5	Fricas [A] (verification not implemented)	549
3.90.6	Sympy [A] (verification not implemented)	549
3.90.7	Maxima [A] (verification not implemented)	549
3.90.8	Giac [A] (verification not implemented)	550
3.90.9	Mupad [B] (verification not implemented)	550

3.90.1 Optimal result

Integrand size = 4, antiderivative size = 22

$$\int \tan^5(x) dx = -\log(\cos(x)) - \frac{\tan^2(x)}{2} + \frac{\tan^4(x)}{4}$$

output `-ln(cos(x))-1/2*tan(x)^2+1/4*tan(x)^4`

3.90.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \tan^5(x) dx = -\log(\cos(x)) - \frac{\tan^2(x)}{2} + \frac{\tan^4(x)}{4}$$

input `Integrate[Tan[x]^5,x]`

output `-Log[Cos[x]] - Tan[x]^2/2 + Tan[x]^4/4`

3.90.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {3042, 3954, 3042, 3954, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^5(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^5 dx \\
 & \quad \downarrow \text{3954} \\
 & \frac{\tan^4(x)}{4} - \int \tan^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{\tan^4(x)}{4} - \int \tan(x)^3 dx \\
 & \quad \downarrow \text{3954} \\
 & \int \tan(x) dx + \frac{\tan^4(x)}{4} - \frac{\tan^2(x)}{2} \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x) dx + \frac{\tan^4(x)}{4} - \frac{\tan^2(x)}{2} \\
 & \quad \downarrow \text{3956} \\
 & \frac{\tan^4(x)}{4} - \frac{\tan^2(x)}{2} - \log(\cos(x))
 \end{aligned}$$

input `Int [Tan [x] ^5, x]`

output `-Log [Cos [x]] - Tan [x] ^2/2 + Tan [x] ^4/4`

3.90.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

3.90.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.05

method	result	size
derivativedivides	$\frac{\tan^4(x)}{4} - \frac{\tan^2(x)}{2} + \frac{\ln(1+\tan^2(x))}{2}$	23
default	$\frac{\tan^4(x)}{4} - \frac{\tan^2(x)}{2} + \frac{\ln(1+\tan^2(x))}{2}$	23
norman	$\frac{\tan^4(x)}{4} - \frac{\tan^2(x)}{2} + \frac{\ln(1+\tan^2(x))}{2}$	23
parallelrisc	$\frac{\tan^4(x)}{4} - \frac{\tan^2(x)}{2} + \frac{\ln(1+\tan^2(x))}{2}$	23
risc	$ix - \frac{4(e^{6ix} + e^{4ix} + e^{2ix})}{(e^{2ix} + 1)^4} - \ln(e^{2ix} + 1)$	43

input `int(tan(x)^5,x,method=_RETURNVERBOSE)`

output `1/4*tan(x)^4-1/2*tan(x)^2+1/2*ln(1+tan(x)^2)`

3.90.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \tan^5(x) dx = \frac{1}{4} \tan(x)^4 - \frac{1}{2} \tan(x)^2 - \frac{1}{2} \log\left(\frac{1}{\tan(x)^2 + 1}\right)$$

input `integrate(tan(x)^5,x, algorithm="fracas")`output `1/4*tan(x)^4 - 1/2*tan(x)^2 - 1/2*log(1/(tan(x)^2 + 1))`**3.90.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \tan^5(x) dx = -\frac{4 \cos^2(x) - 1}{4 \cos^4(x)} - \log(\cos(x))$$

input `integrate(tan(x)**5,x)`output `-(4*cos(x)**2 - 1)/(4*cos(x)**4) - log(cos(x))`**3.90.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.55

$$\int \tan^5(x) dx = \frac{4 \sin(x)^2 - 3}{4 (\sin(x)^4 - 2 \sin(x)^2 + 1)} - \frac{1}{2} \log(\sin(x)^2 - 1)$$

input `integrate(tan(x)^5,x, algorithm="maxima")`output `1/4*(4*sin(x)^2 - 3)/(sin(x)^4 - 2*sin(x)^2 + 1) - 1/2*log(sin(x)^2 - 1)`

3.90.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \tan^5(x) dx = \frac{1}{4} \tan(x)^4 - \frac{1}{2} \tan(x)^2 + \frac{1}{2} \log(\tan(x)^2 + 1)$$

input `integrate(tan(x)^5,x, algorithm="giac")`

output `1/4*tan(x)^4 - 1/2*tan(x)^2 + 1/2*log(tan(x)^2 + 1)`

3.90.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \tan^5(x) dx = \frac{\tan(x)^4}{4} - \frac{\tan(x)^2}{2} - \ln(\cos(x))$$

input `int(tan(x)^5,x)`

output `tan(x)^4/4 - tan(x)^2/2 - log(cos(x))`

3.91 $\int \tan^6(x) dx$

3.91.1	Optimal result	551
3.91.2	Mathematica [A] (verified)	551
3.91.3	Rubi [A] (verified)	552
3.91.4	Maple [A] (verified)	553
3.91.5	Fricas [A] (verification not implemented)	554
3.91.6	Sympy [A] (verification not implemented)	554
3.91.7	Maxima [A] (verification not implemented)	554
3.91.8	Giac [A] (verification not implemented)	555
3.91.9	Mupad [B] (verification not implemented)	555

3.91.1 Optimal result

Integrand size = 4, antiderivative size = 22

$$\int \tan^6(x) dx = -x + \tan(x) - \frac{\tan^3(x)}{3} + \frac{\tan^5(x)}{5}$$

output `-x+tan(x)-1/3*tan(x)^3+1/5*tan(x)^5`

3.91.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \tan^6(x) dx = -\arctan(\tan(x)) + \tan(x) - \frac{\tan^3(x)}{3} + \frac{\tan^5(x)}{5}$$

input `Integrate[Tan[x]^6,x]`

output `-ArcTan[Tan[x]] + Tan[x] - Tan[x]^3/3 + Tan[x]^5/5`

3.91.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.750$, Rules used = {3042, 3954, 3042, 3954, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^6(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^6 dx \\
 & \quad \downarrow \text{3954} \\
 & \frac{\tan^5(x)}{5} - \int \tan^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{\tan^5(x)}{5} - \int \tan(x)^4 dx \\
 & \quad \downarrow \text{3954} \\
 & \int \tan^2(x) dx + \frac{\tan^5(x)}{5} - \frac{\tan^3(x)}{3} \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^2 dx + \frac{\tan^5(x)}{5} - \frac{\tan^3(x)}{3} \\
 & \quad \downarrow \text{3954} \\
 & - \int 1 dx + \frac{\tan^5(x)}{5} - \frac{\tan^3(x)}{3} + \tan(x) \\
 & \quad \downarrow \text{24} \\
 & -x + \frac{\tan^5(x)}{5} - \frac{\tan^3(x)}{3} + \tan(x)
 \end{aligned}$$

input `Int [Tan[x]^6, x]`

output `-x + Tan[x] - Tan[x]^3/3 + Tan[x]^5/5`

3.91.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

3.91.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.86

method	result	size
norman	$-x + \tan(x) - \frac{\tan^3(x)}{3} + \frac{\tan^5(x)}{5}$	19
parallelrisch	$-x + \tan(x) - \frac{\tan^3(x)}{3} + \frac{\tan^5(x)}{5}$	19
derivativedivides	$\frac{\tan^5(x)}{5} - \frac{\tan^3(x)}{3} + \tan(x) - \arctan(\tan(x))$	21
default	$\frac{\tan^5(x)}{5} - \frac{\tan^3(x)}{3} + \tan(x) - \arctan(\tan(x))$	21
risch	$-x + \frac{2i(45 e^{8ix} + 90 e^{6ix} + 140 e^{4ix} + 70 e^{2ix} + 23)}{15(e^{2ix} + 1)^5}$	47

input `int(tan(x)^6,x,method=_RETURNVERBOSE)`

output `-x+tan(x)-1/3*tan(x)^3+1/5*tan(x)^5`

3.91.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \tan^6(x) dx = \frac{1}{5} \tan(x)^5 - \frac{1}{3} \tan(x)^3 - x + \tan(x)$$

input `integrate(tan(x)^6,x, algorithm="fricas")`output `1/5*tan(x)^5 - 1/3*tan(x)^3 - x + tan(x)`**3.91.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.41

$$\int \tan^6(x) dx = -x + \frac{\sin^5(x)}{5 \cos^5(x)} - \frac{\sin^3(x)}{3 \cos^3(x)} + \frac{\sin(x)}{\cos(x)}$$

input `integrate(tan(x)**6,x)`output `-x + sin(x)**5/(5*cos(x)**5) - sin(x)**3/(3*cos(x)**3) + sin(x)/cos(x)`**3.91.7 Maxima [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \tan^6(x) dx = \frac{1}{5} \tan(x)^5 - \frac{1}{3} \tan(x)^3 - x + \tan(x)$$

input `integrate(tan(x)^6,x, algorithm="maxima")`output `1/5*tan(x)^5 - 1/3*tan(x)^3 - x + tan(x)`

3.91.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \tan^6(x) dx = \frac{1}{5} \tan(x)^5 - \frac{1}{3} \tan(x)^3 - x + \tan(x)$$

input `integrate(tan(x)^6,x, algorithm="giac")`

output `1/5*tan(x)^5 - 1/3*tan(x)^3 - x + tan(x)`

3.91.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \tan^6(x) dx = \frac{\tan(x)^5}{5} - \frac{\tan(x)^3}{3} + \tan(x) - x$$

input `int(tan(x)^6,x)`

output `tan(x) - x - tan(x)^3/3 + tan(x)^5/5`

3.92 $\int \sec(x) \tan^5(x) dx$

3.92.1	Optimal result	556
3.92.2	Mathematica [A] (verified)	556
3.92.3	Rubi [A] (verified)	557
3.92.4	Maple [A] (verified)	558
3.92.5	Fricas [A] (verification not implemented)	558
3.92.6	Sympy [A] (verification not implemented)	559
3.92.7	Maxima [A] (verification not implemented)	559
3.92.8	Giac [A] (verification not implemented)	559
3.92.9	Mupad [B] (verification not implemented)	560

3.92.1 Optimal result

Integrand size = 7, antiderivative size = 19

$$\int \sec(x) \tan^5(x) dx = \sec(x) - \frac{2 \sec^3(x)}{3} + \frac{\sec^5(x)}{5}$$

output `sec(x)-2/3*sec(x)^3+1/5*sec(x)^5`

3.92.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \sec(x) \tan^5(x) dx = \sec(x) - \frac{2 \sec^3(x)}{3} + \frac{\sec^5(x)}{5}$$

input `Integrate[Sec[x]*Tan[x]^5,x]`

output `Sec[x] - (2*Sec[x]^3)/3 + Sec[x]^5/5`

3.92.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 3086, 210, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^5(x) \sec(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^5 \sec(x) dx \\
 & \quad \downarrow \text{3086} \\
 & \int (\sec^2(x) - 1)^2 d\sec(x) \\
 & \quad \downarrow \text{210} \\
 & \int (\sec^4(x) - 2\sec^2(x) + 1) d\sec(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sec^5(x)}{5} - \frac{2\sec^3(x)}{3} + \sec(x)
 \end{aligned}$$

input `Int[Sec[x]*Tan[x]^5,x]`

output `Sec[x] - (2*Sec[x]^3)/3 + Sec[x]^5/5`

3.92.3.1 Defintions of rubi rules used

rule 210 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Int[ExpandIntegrand[(a + b*x^2)^(p, x), x] /; FreeQ[{a, b}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3086 Int[((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_.))*((b_.)*tan[(e_.) + (f_.)*(x_)]^(
n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2
), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2
] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])
```

3.92.4 Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$\sec(x) - \frac{2(\sec^3(x))}{3} + \frac{(\sec^5(x))}{5}$	16
default	$\sec(x) - \frac{2(\sec^3(x))}{3} + \frac{(\sec^5(x))}{5}$	16
risch	$\frac{2e^{9ix} + \frac{8e^{7ix}}{3} + \frac{116e^{5ix}}{15} + \frac{8e^{3ix}}{3} + 2e^{ix}}{(e^{2ix} + 1)^5}$	48

```
input int(sec(x)*tan(x)^5,x,method=_RETURNVERBOSE)
```

```
output sec(x)-2/3*sec(x)^3+1/5*sec(x)^5
```

3.92.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.05

$$\int \sec(x) \tan^5(x) dx = \frac{15 \cos(x)^4 - 10 \cos(x)^2 + 3}{15 \cos(x)^5}$$

```
input integrate(sec(x)*tan(x)^5,x, algorithm="fracas")
```

```
output 1/15*(15*cos(x)^4 - 10*cos(x)^2 + 3)/cos(x)^5
```

3.92.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.16

$$\int \sec(x) \tan^5(x) dx = -\frac{-15 \cos^4(x) + 10 \cos^2(x) - 3}{15 \cos^5(x)}$$

input `integrate(sec(x)*tan(x)**5,x)`output `-(-15*cos(x)**4 + 10*cos(x)**2 - 3)/(15*cos(x)**5)`**3.92.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.05

$$\int \sec(x) \tan^5(x) dx = \frac{15 \cos(x)^4 - 10 \cos(x)^2 + 3}{15 \cos(x)^5}$$

input `integrate(sec(x)*tan(x)^5,x, algorithm="maxima")`output `1/15*(15*cos(x)^4 - 10*cos(x)^2 + 3)/cos(x)^5`**3.92.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.05

$$\int \sec(x) \tan^5(x) dx = \frac{15 \cos(x)^4 - 10 \cos(x)^2 + 3}{15 \cos(x)^5}$$

input `integrate(sec(x)*tan(x)^5,x, algorithm="giac")`output `1/15*(15*cos(x)^4 - 10*cos(x)^2 + 3)/cos(x)^5`

3.92.9 Mupad [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \sec(x) \tan^5(x) dx = \frac{\cos(x)^4 - \frac{2\cos(x)^2}{3} + \frac{1}{5}}{\cos(x)^5}$$

input `int(tan(x)^5/cos(x),x)`

output `(cos(x)^4 - (2*cos(x)^2)/3 + 1/5)/cos(x)^5`

3.93 $\int \sec^3(x) \tan^5(x) dx$

3.93.1	Optimal result	561
3.93.2	Mathematica [A] (verified)	561
3.93.3	Rubi [A] (verified)	562
3.93.4	Maple [A] (verified)	563
3.93.5	Fricas [A] (verification not implemented)	563
3.93.6	Sympy [A] (verification not implemented)	564
3.93.7	Maxima [A] (verification not implemented)	564
3.93.8	Giac [A] (verification not implemented)	564
3.93.9	Mupad [B] (verification not implemented)	565

3.93.1 Optimal result

Integrand size = 9, antiderivative size = 25

$$\int \sec^3(x) \tan^5(x) dx = \frac{\sec^3(x)}{3} - \frac{2 \sec^5(x)}{5} + \frac{\sec^7(x)}{7}$$

output `1/3*sec(x)^3-2/5*sec(x)^5+1/7*sec(x)^7`

3.93.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \sec^3(x) \tan^5(x) dx = \frac{\sec^3(x)}{3} - \frac{2 \sec^5(x)}{5} + \frac{\sec^7(x)}{7}$$

input `Integrate[Sec[x]^3*Tan[x]^5,x]`

output `Sec[x]^3/3 - (2*Sec[x]^5)/5 + Sec[x]^7/7`

3.93.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 3086, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^5(x) \sec^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^5 \sec(x)^3 dx \\
 & \quad \downarrow \text{3086} \\
 & \int \sec^2(x) (1 - \sec^2(x))^2 d \sec(x) \\
 & \quad \downarrow \text{244} \\
 & \int (\sec^6(x) - 2 \sec^4(x) + \sec^2(x)) d \sec(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sec^7(x)}{7} - \frac{2 \sec^5(x)}{5} + \frac{\sec^3(x)}{3}
 \end{aligned}$$

input `Int[Sec[x]^3*Tan[x]^5,x]`

output `Sec[x]^3/3 - (2*Sec[x]^5)/5 + Sec[x]^7/7`

3.93.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3086 Int[((a_)*sec[(e_) + (f_)*(x_)]^(m_))*((b_)*tan[(e_) + (f_)*(x_)]^(
n_), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2
), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2
] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])
```

3.93.4 Maple [A] (verified)

Time = 2.25 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

method	result	size
derivativedivides	$\frac{\sec^3(x)}{3} - \frac{2(\sec^5(x))}{5} + \frac{\sec^7(x)}{7}$	20
default	$\frac{\sec^3(x)}{3} - \frac{2(\sec^5(x))}{5} + \frac{\sec^7(x)}{7}$	20
risch	$\frac{8e^{11ix} - 32e^{9ix} + 304e^{7ix} - 32e^{5ix} + 8e^{3ix}}{3(e^{2ix} + 1)^7} - \frac{32e^{9ix} - 304e^{7ix} + 32e^{5ix} - 8e^{3ix}}{15(e^{2ix} + 1)^7} + \frac{304e^{7ix} - 32e^{5ix} + 8e^{3ix}}{35(e^{2ix} + 1)^7} - \frac{32e^{5ix} - 8e^{3ix}}{15(e^{2ix} + 1)^7} + \frac{8e^{3ix}}{3(e^{2ix} + 1)^7}$	48

```
input int(sec(x)^3*tan(x)^5,x,method=_RETURNVERBOSE)
```

```
output 1/3*sec(x)^3-2/5*sec(x)^5+1/7*sec(x)^7
```

3.93.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \sec^3(x) \tan^5(x) dx = \frac{35 \cos(x)^4 - 42 \cos(x)^2 + 15}{105 \cos(x)^7}$$

```
input integrate(sec(x)^3*tan(x)^5,x, algorithm="fricas")
```

```
output 1/105*(35*cos(x)^4 - 42*cos(x)^2 + 15)/cos(x)^7
```

3.93.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \sec^3(x) \tan^5(x) dx = -\frac{-35 \cos^4(x) + 42 \cos^2(x) - 15}{105 \cos^7(x)}$$

input `integrate(sec(x)**3*tan(x)**5,x)`output `-(-35*cos(x)**4 + 42*cos(x)**2 - 15)/(105*cos(x)**7)`**3.93.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \sec^3(x) \tan^5(x) dx = \frac{35 \cos(x)^4 - 42 \cos(x)^2 + 15}{105 \cos(x)^7}$$

input `integrate(sec(x)^3*tan(x)^5,x, algorithm="maxima")`output `1/105*(35*cos(x)^4 - 42*cos(x)^2 + 15)/cos(x)^7`**3.93.8 Giac [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \sec^3(x) \tan^5(x) dx = \frac{35 \cos(x)^4 - 42 \cos(x)^2 + 15}{105 \cos(x)^7}$$

input `integrate(sec(x)^3*tan(x)^5,x, algorithm="giac")`output `1/105*(35*cos(x)^4 - 42*cos(x)^2 + 15)/cos(x)^7`

3.93.9 Mupad [B] (verification not implemented)

Time = 0.54 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \sec^3(x) \tan^5(x) dx = \frac{\frac{\cos(x)^4}{3} - \frac{2\cos(x)^2}{5} + \frac{1}{7}}{\cos(x)^7}$$

input `int(tan(x)^5/cos(x)^3,x)`

output `(cos(x)^4/3 - (2*cos(x)^2)/5 + 1/7)/cos(x)^7`

3.94 $\int \sec^6(x) \tan(x) dx$

3.94.1	Optimal result	566
3.94.2	Mathematica [A] (verified)	566
3.94.3	Rubi [A] (verified)	567
3.94.4	Maple [A] (verified)	568
3.94.5	Fricas [A] (verification not implemented)	568
3.94.6	Sympy [A] (verification not implemented)	568
3.94.7	Maxima [A] (verification not implemented)	569
3.94.8	Giac [A] (verification not implemented)	569
3.94.9	Mupad [B] (verification not implemented)	569

3.94.1 Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \sec^6(x) \tan(x) dx = \frac{\sec^6(x)}{6}$$

output `1/6*sec(x)^6`

3.94.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \sec^6(x) \tan(x) dx = \frac{\sec^6(x)}{6}$$

input `Integrate[Sec[x]^6*Tan[x],x]`

output `Sec[x]^6/6`

3.94.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3086, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan(x) \sec^6(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(x) \sec(x)^6 dx \\ & \quad \downarrow \text{3086} \\ & \int \sec^5(x) d \sec(x) \\ & \quad \downarrow \text{15} \\ & \frac{\sec^6(x)}{6} \end{aligned}$$

input `Int[Sec[x]^6*Tan[x],x]`

output `Sec[x]^6/6`

3.94.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

3.94.4 Maple [A] (verified)

Time = 9.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{(\sec^6(x))}{6}$	7
default	$\frac{(\sec^6(x))}{6}$	7
risch	$\frac{32 e^{6ix}}{3(e^{2ix}+1)^6}$	17

input `int(sec(x)^6*tan(x),x,method=_RETURNVERBOSE)`

output `1/6*sec(x)^6`

3.94.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^6(x) \tan(x) dx = \frac{1}{6 \cos^6(x)}$$

input `integrate(sec(x)^6*tan(x),x, algorithm="fracas")`

output `1/6/cos(x)^6`

3.94.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \sec^6(x) \tan(x) dx = \frac{1}{6 \cos^6(x)}$$

input `integrate(sec(x)**6*tan(x),x)`

output `1/(6*cos(x)**6)`

3.94.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.25

$$\int \sec^6(x) \tan(x) dx = -\frac{1}{6 (\sin(x)^2 - 1)^3}$$

input `integrate(sec(x)^6*tan(x),x, algorithm="maxima")`output `-1/6/(sin(x)^2 - 1)^3`**3.94.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^6(x) \tan(x) dx = \frac{1}{6 \cos(x)^6}$$

input `integrate(sec(x)^6*tan(x),x, algorithm="giac")`output `1/6/cos(x)^6`**3.94.9 Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 2.25

$$\int \sec^6(x) \tan(x) dx = \frac{\tan(x)^2 (\tan(x)^4 + 3 \tan(x)^2 + 3)}{6}$$

input `int(tan(x)/cos(x)^6,x)`output `(tan(x)^2*(3*tan(x)^2 + tan(x)^4 + 3))/6`

3.95 $\int \sec^6(x) \tan^3(x) dx$

3.95.1	Optimal result	570
3.95.2	Mathematica [A] (verified)	570
3.95.3	Rubi [A] (verified)	571
3.95.4	Maple [A] (verified)	572
3.95.5	Fricas [A] (verification not implemented)	573
3.95.6	Sympy [A] (verification not implemented)	573
3.95.7	Maxima [B] (verification not implemented)	573
3.95.8	Giac [A] (verification not implemented)	574
3.95.9	Mupad [B] (verification not implemented)	574

3.95.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \sec^6(x) \tan^3(x) dx = -\frac{1}{6} \sec^6(x) + \frac{\sec^8(x)}{8}$$

output `-1/6*sec(x)^6+1/8*sec(x)^8`

3.95.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sec^6(x) \tan^3(x) dx = -\frac{1}{6} \sec^6(x) + \frac{\sec^8(x)}{8}$$

input `Integrate[Sec[x]^6*Tan[x]^3,x]`

output `-1/6*Sec[x]^6 + Sec[x]^8/8`

3.95.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 3086, 25, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^3(x) \sec^6(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^3 \sec(x)^6 dx \\
 & \quad \downarrow \text{3086} \\
 & \int -\sec^5(x) (1 - \sec^2(x)) d \sec(x) \\
 & \quad \downarrow \text{25} \\
 & - \int \sec^5(x) (1 - \sec^2(x)) d \sec(x) \\
 & \quad \downarrow \text{244} \\
 & - \int (\sec^5(x) - \sec^7(x)) d \sec(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sec^8(x)}{8} - \frac{\sec^6(x)}{6}
 \end{aligned}$$

input `Int[Sec[x]^6*Tan[x]^3,x]`

output `-1/6*Sec[x]^6 + Sec[x]^8/8`

3.95.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

3.95.4 Maple [A] (verified)

Time = 18.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$-\frac{(\sec^6(x))}{6} + \frac{(\sec^8(x))}{8}$	14
default	$-\frac{(\sec^6(x))}{6} + \frac{(\sec^8(x))}{8}$	14
risch	$-\frac{32(e^{10ix} - e^{8ix} + e^{6ix})}{3(e^{2ix} + 1)^8}$	30

input `int(sec(x)^6*tan(x)^3,x,method=_RETURNVERBOSE)`

output `-1/6*sec(x)^6+1/8*sec(x)^8`

3.95.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^6(x) \tan^3(x) dx = -\frac{4 \cos(x)^2 - 3}{24 \cos(x)^8}$$

input `integrate(sec(x)^6*tan(x)^3,x, algorithm="fracas")`

output `-1/24*(4*cos(x)^2 - 3)/cos(x)^8`

3.95.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^6(x) \tan^3(x) dx = \frac{3 - 4 \cos^2(x)}{24 \cos^8(x)}$$

input `integrate(sec(x)**6*tan(x)**3,x)`

output `(3 - 4*cos(x)**2)/(24*cos(x)**8)`

3.95.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 36 vs. $2(13) = 26$.

Time = 0.19 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.12

$$\int \sec^6(x) \tan^3(x) dx = \frac{4 \sin(x)^2 - 1}{24 (\sin(x)^8 - 4 \sin(x)^6 + 6 \sin(x)^4 - 4 \sin(x)^2 + 1)}$$

input `integrate(sec(x)^6*tan(x)^3,x, algorithm="maxima")`

output `1/24*(4*sin(x)^2 - 1)/(sin(x)^8 - 4*sin(x)^6 + 6*sin(x)^4 - 4*sin(x)^2 + 1)`

3.95.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^6(x) \tan^3(x) dx = -\frac{4 \cos(x)^2 - 3}{24 \cos(x)^8}$$

input `integrate(sec(x)^6*tan(x)^3,x, algorithm="giac")`

output `-1/24*(4*cos(x)^2 - 3)/cos(x)^8`

3.95.9 Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \sec^6(x) \tan^3(x) dx = \frac{\tan(x)^4 (3 \tan(x)^4 + 8 \tan(x)^2 + 6)}{24}$$

input `int(tan(x)^3/cos(x)^6,x)`

output `(tan(x)^4*(8*tan(x)^2 + 3*tan(x)^4 + 6))/24`

3.96 $\int \sec^2(x) \tan(x) dx$

3.96.1	Optimal result	575
3.96.2	Mathematica [A] (verified)	575
3.96.3	Rubi [A] (verified)	576
3.96.4	Maple [A] (verified)	577
3.96.5	Fricas [A] (verification not implemented)	577
3.96.6	Sympy [A] (verification not implemented)	577
3.96.7	Maxima [A] (verification not implemented)	578
3.96.8	Giac [A] (verification not implemented)	578
3.96.9	Mupad [B] (verification not implemented)	578

3.96.1 Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \sec^2(x) \tan(x) dx = \frac{\sec^2(x)}{2}$$

output `1/2*sec(x)^2`

3.96.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \sec^2(x) \tan(x) dx = \frac{\sec^2(x)}{2}$$

input `Integrate[Sec[x]^2*Tan[x],x]`

output `Sec[x]^2/2`

3.96.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3086, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan(x) \sec^2(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(x) \sec(x)^2 dx \\ & \quad \downarrow \text{3086} \\ & \int \sec(x) d \sec(x) \\ & \quad \downarrow \text{15} \\ & \frac{\sec^2(x)}{2} \end{aligned}$$

input `Int[Sec[x]^2*Tan[x],x]`

output `Sec[x]^2/2`

3.96.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

3.96.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{(\sec^2(x))}{2}$	7
default	$\frac{(\sec^2(x))}{2}$	7
risch	$\frac{2 e^{2ix}}{(e^{2ix}+1)^2}$	17

input `int(sec(x)^2/cot(x),x,method=_RETURNVERBOSE)`

output `1/2*sec(x)^2`

3.96.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan(x) dx = \frac{1}{2 \cos(x)^2}$$

input `integrate(sec(x)^2/cot(x),x, algorithm="fricas")`

output `1/2/cos(x)^2`

3.96.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \sec^2(x) \tan(x) dx = \frac{1}{2 \cos^2(x)}$$

input `integrate(sec(x)**2/cot(x),x)`

output `1/(2*cos(x)**2)`

3.96.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.25

$$\int \sec^2(x) \tan(x) dx = -\frac{1}{2(\sin(x)^2 - 1)}$$

input `integrate(sec(x)^2/cot(x),x, algorithm="maxima")`output `-1/2/(sin(x)^2 - 1)`**3.96.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan(x) dx = \frac{1}{2 \cos(x)^2}$$

input `integrate(sec(x)^2/cot(x),x, algorithm="giac")`output `1/2/cos(x)^2`**3.96.9 Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^2(x) \tan(x) dx = \frac{\tan(x)^2}{2}$$

input `int(1/(cos(x)^2*cot(x)),x)`output `tan(x)^2/2`

3.97 $\int \sec(x) \tan^2(x) dx$

3.97.1	Optimal result	579
3.97.2	Mathematica [A] (verified)	579
3.97.3	Rubi [A] (verified)	580
3.97.4	Maple [A] (verified)	581
3.97.5	Fricas [B] (verification not implemented)	581
3.97.6	Sympy [A] (verification not implemented)	582
3.97.7	Maxima [B] (verification not implemented)	582
3.97.8	Giac [B] (verification not implemented)	582
3.97.9	Mupad [B] (verification not implemented)	583

3.97.1 Optimal result

Integrand size = 7, antiderivative size = 16

$$\int \sec(x) \tan^2(x) dx = -\frac{1}{2} \operatorname{arctanh}(\sin(x)) + \frac{1}{2} \sec(x) \tan(x)$$

output `-1/2*arctanh(sin(x))+1/2*sec(x)*tan(x)`

3.97.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \sec(x) \tan^2(x) dx = -\frac{1}{2} \operatorname{arctanh}(\sin(x)) + \frac{1}{2} \sec(x) \tan(x)$$

input `Integrate[Sec[x]*Tan[x]^2,x]`

output `-1/2*ArcTanh[Sin[x]] + (Sec[x]*Tan[x])/2`

3.97.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 3091, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^2(x) \sec(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^2 \sec(x) dx \\
 & \quad \downarrow \text{3091} \\
 & \frac{1}{2} \tan(x) \sec(x) - \frac{\int \sec(x) dx}{2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \tan(x) \sec(x) - \frac{1}{2} \int \csc\left(x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{4257} \\
 & \frac{1}{2} \tan(x) \sec(x) - \frac{1}{2} \operatorname{arctanh}(\sin(x))
 \end{aligned}$$

input `Int[Sec[x]*Tan[x]^2,x]`

output `-1/2*ArcTanh[Sin[x]] + (Sec[x]*Tan[x])/2`

3.97.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

```
rule 3091 Int[((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] :> Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Simp[b^2*((n - 1)/(m + n - 1)) Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

3.97.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.50

method	result	size
default	$\frac{\sin^3(x)}{2 \cos(x)^2} + \frac{\sin(x)}{2} - \frac{\ln(\sec(x)+\tan(x))}{2}$	24
risch	$-\frac{i(e^{3ix}-e^{ix})}{(e^{2ix}+1)^2} - \frac{\ln(i+e^{ix})}{2} + \frac{\ln(e^{ix}-i)}{2}$	49

```
input int(sec(x)*tan(x)^2,x,method=_RETURNVERBOSE)
```

```
output 1/2*sin(x)^3/cos(x)^2+1/2*sin(x)-1/2*ln(sec(x)+tan(x))
```

3.97.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. $2(12) = 24$.

Time = 0.26 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.12

$$\int \sec(x) \tan^2(x) dx = -\frac{\cos(x)^2 \log(\sin(x) + 1) - \cos(x)^2 \log(-\sin(x) + 1) - 2 \sin(x)}{4 \cos(x)^2}$$

```
input integrate(sec(x)*tan(x)^2,x, algorithm="fricas")
```

```
output -1/4*(cos(x)^2*log(sin(x) + 1) - cos(x)^2*log(-sin(x) + 1) - 2*sin(x))/cos(x)^2
```

3.97.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.69

$$\int \sec(x) \tan^2(x) dx = \frac{\log(\sin(x) - 1)}{4} - \frac{\log(\sin(x) + 1)}{4} - \frac{\sin(x)}{2 \sin^2(x) - 2}$$

input `integrate(sec(x)*tan(x)**2,x)`

output `log(sin(x) - 1)/4 - log(sin(x) + 1)/4 - sin(x)/(2*sin(x)**2 - 2)`

3.97.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(12) = 24.

Time = 0.19 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.69

$$\int \sec(x) \tan^2(x) dx = -\frac{\sin(x)}{2(\sin(x)^2 - 1)} - \frac{1}{4} \log(\sin(x) + 1) + \frac{1}{4} \log(\sin(x) - 1)$$

input `integrate(sec(x)*tan(x)^2,x, algorithm="maxima")`

output `-1/2*sin(x)/(sin(x)^2 - 1) - 1/4*log(sin(x) + 1) + 1/4*log(sin(x) - 1)`

3.97.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. 2(12) = 24.

Time = 0.29 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.81

$$\int \sec(x) \tan^2(x) dx = -\frac{\sin(x)}{2(\sin(x)^2 - 1)} - \frac{1}{4} \log(\sin(x) + 1) + \frac{1}{4} \log(-\sin(x) + 1)$$

input `integrate(sec(x)*tan(x)^2,x, algorithm="giac")`

output `-1/2*sin(x)/(sin(x)^2 - 1) - 1/4*log(sin(x) + 1) + 1/4*log(-sin(x) + 1)`

3.97.9 Mupad [B] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.88

$$\int \sec(x) \tan^2(x) dx = \frac{\tan\left(\frac{x}{2}\right)^3 + \tan\left(\frac{x}{2}\right)}{\left(\tan\left(\frac{x}{2}\right)^2 - 1\right)^2} - \operatorname{atanh}\left(\tan\left(\frac{x}{2}\right)\right)$$

input `int(tan(x)^2/cos(x),x)`

output `(tan(x/2) + tan(x/2)^3)/(tan(x/2)^2 - 1)^2 - atanh(tan(x/2))`

3.98 $\int \cot^2(x) dx$

3.98.1	Optimal result	584
3.98.2	Mathematica [C] (verified)	584
3.98.3	Rubi [A] (verified)	585
3.98.4	Maple [A] (verified)	586
3.98.5	Fricas [B] (verification not implemented)	586
3.98.6	Sympy [A] (verification not implemented)	586
3.98.7	Maxima [A] (verification not implemented)	587
3.98.8	Giac [B] (verification not implemented)	587
3.98.9	Mupad [B] (verification not implemented)	587

3.98.1 Optimal result

Integrand size = 4, antiderivative size = 8

$$\int \cot^2(x) dx = -x - \cot(x)$$

output `-x-cot(x)`

3.98.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 2.25

$$\int \cot^2(x) dx = -\cot(x) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(x)\right)$$

input `Integrate[Cot[x]^2,x]`

output `-(Cot[x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[x]^2])`

3.98.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \cot^2(x) dx \\
 \downarrow 3042 \\
 \int \tan\left(x + \frac{\pi}{2}\right)^2 dx \\
 \downarrow 3954 \\
 - \int 1 dx - \cot(x) \\
 \downarrow 24 \\
 -x - \cot(x)
 \end{array}$$

input `Int[Cot[x]^2,x]`

output `-x - Cot[x]`

3.98.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

3.98.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.62

method	result	size
norman	$\frac{-1-x \tan(x)}{\tan(x)}$	13
parallelrisc	$\frac{-1-x \tan(x)}{\tan(x)}$	13
derivativedivides	$-\cot(x) + \frac{\pi}{2} - \operatorname{arccot}(\cot(x))$	14
default	$-\cot(x) + \frac{\pi}{2} - \operatorname{arccot}(\cot(x))$	14
risc	$-x - \frac{2i}{e^{2ix}-1}$	17

input `int(cot(x)^2,x,method=_RETURNVERBOSE)`

output `(-1-x*tan(x))/tan(x)`

3.98.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. 2(8) = 16.

Time = 0.25 (sec) , antiderivative size = 20, normalized size of antiderivative = 2.50

$$\int \cot^2(x) dx = -\frac{x \sin(2x) + \cos(2x) + 1}{\sin(2x)}$$

input `integrate(cot(x)^2,x, algorithm="fricas")`

output `-(x*sin(2*x) + cos(2*x) + 1)/sin(2*x)`

3.98.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \cot^2(x) dx = -x - \frac{\cos(x)}{\sin(x)}$$

input `integrate(cot(x)**2,x)`

output `-x - cos(x)/sin(x)`

3.98.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.25

$$\int \cot^2(x) dx = -x - \frac{1}{\tan(x)}$$

input `integrate(cot(x)^2,x, algorithm="maxima")`

output `-x - 1/tan(x)`

3.98.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18 vs. 2(8) = 16.

Time = 0.29 (sec) , antiderivative size = 18, normalized size of antiderivative = 2.25

$$\int \cot^2(x) dx = -x - \frac{1}{2 \tan\left(\frac{1}{2}x\right)} + \frac{1}{2} \tan\left(\frac{1}{2}x\right)$$

input `integrate(cot(x)^2,x, algorithm="giac")`

output `-x - 1/2/tan(1/2*x) + 1/2*tan(1/2*x)`

3.98.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \cot^2(x) dx = -x - \cot(x)$$

input `int(cot(x)^2,x)`

output `- x - cot(x)`

3.99 $\int \cot^3(x) dx$

3.99.1	Optimal result	588
3.99.2	Mathematica [A] (verified)	588
3.99.3	Rubi [A] (verified)	589
3.99.4	Maple [A] (verified)	590
3.99.5	Fricas [B] (verification not implemented)	591
3.99.6	Sympy [A] (verification not implemented)	591
3.99.7	Maxima [A] (verification not implemented)	591
3.99.8	Giac [A] (verification not implemented)	592
3.99.9	Mupad [B] (verification not implemented)	592

3.99.1 Optimal result

Integrand size = 4, antiderivative size = 14

$$\int \cot^3(x) dx = -\frac{1}{2} \cot^2(x) - \log(\sin(x))$$

output `-1/2*cot(x)^2-ln(sin(x))`

3.99.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.36

$$\int \cot^3(x) dx = -\frac{1}{2} \cot^2(x) - \log(\cos(x)) - \log(\tan(x))$$

input `Integrate[Cot[x]^3,x]`

output `-1/2*Cot[x]^2 - Log[Cos[x]] - Log[Tan[x]]`

3.99.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.750$, Rules used = {3042, 25, 3954, 25, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\tan\left(x + \frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{25} \\
 & -\int \tan\left(x + \frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{3954} \\
 & \int -\cot(x) dx - \frac{\cot^2(x)}{2} \\
 & \quad \downarrow \text{25} \\
 & -\int \cot(x) dx - \frac{1}{2} \cot^2(x) \\
 & \quad \downarrow \text{3042} \\
 & -\int -\tan\left(x + \frac{\pi}{2}\right) dx - \frac{1}{2} \cot^2(x) \\
 & \quad \downarrow \text{25} \\
 & \int \tan\left(x + \frac{\pi}{2}\right) dx - \frac{\cot^2(x)}{2} \\
 & \quad \downarrow \text{3956} \\
 & -\frac{1}{2} \cot^2(x) - \log(\sin(x))
 \end{aligned}$$

input `Int[Cot[x]^3,x]`

output `-1/2*Cot[x]^2 - Log[Sin[x]]`

3.99.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

3.99.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.21

method	result	size
derivativedivides	$-\frac{(\cot^2(x))}{2} + \frac{\ln(\cot^2(x)+1)}{2}$	17
default	$-\frac{(\cot^2(x))}{2} + \frac{\ln(\cot^2(x)+1)}{2}$	17
parallelrisc	$-\ln(\tan(x)) + \ln(\sqrt{\sec^2(x)}) - \frac{(\cot^2(x))}{2}$	20
norman	$-\frac{1}{2 \tan(x)^2} - \ln(\tan(x)) + \frac{\ln(1+\tan^2(x))}{2}$	22
risc	$ix + \frac{2e^{2ix}}{(e^{2ix}-1)^2} - \ln(e^{2ix} - 1)$	32

input `int(cot(x)^3,x,method=_RETURNVERBOSE)`

output `-1/2*cot(x)^2+1/2*ln(cot(x)^2+1)`

3.99.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 28 vs. $2(12) = 24$.

Time = 0.25 (sec) , antiderivative size = 28, normalized size of antiderivative = 2.00

$$\int \cot^3(x) dx = -\frac{(\cos(2x) - 1) \log\left(-\frac{1}{2} \cos(2x) + \frac{1}{2}\right) - 2}{2(\cos(2x) - 1)}$$

input `integrate(cot(x)^3,x, algorithm="fricas")`

output `-1/2*((cos(2*x) - 1)*log(-1/2*cos(2*x) + 1/2) - 2)/(cos(2*x) - 1)`

3.99.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \cot^3(x) dx = -\log(\sin(x)) - \frac{1}{2 \sin^2(x)}$$

input `integrate(cot(x)**3,x)`

output `-log(sin(x)) - 1/(2*sin(x)**2)`

3.99.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \cot^3(x) dx = -\frac{1}{2 \sin(x)^2} - \frac{1}{2} \log(\sin(x)^2)$$

input `integrate(cot(x)^3,x, algorithm="maxima")`

output `-1/2/sin(x)^2 - 1/2*log(sin(x)^2)`

3.99.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.57

$$\int \cot^3(x) dx = \frac{1}{2(\cos(x)^2 - 1)} - \frac{1}{2} \log(-\cos(x)^2 + 1)$$

input `integrate(cot(x)^3,x, algorithm="giac")`

output `1/2/(cos(x)^2 - 1) - 1/2*log(-cos(x)^2 + 1)`

3.99.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.29

$$\int \cot^3(x) dx = \frac{\sin(x)^2 - 1}{2 \sin(x)^2} - \ln(\sin(x))$$

input `int(cot(x)^3,x)`

output `(sin(x)^2 - 1)/(2*sin(x)^2) - log(sin(x))`

3.100 $\int \cot^4(x) \csc^4(x) dx$

3.100.1 Optimal result	593
3.100.2 Mathematica [B] (verified)	593
3.100.3 Rubi [A] (verified)	594
3.100.4 Maple [A] (verified)	595
3.100.5 Fricas [B] (verification not implemented)	595
3.100.6 Sympy [B] (verification not implemented)	596
3.100.7 Maxima [A] (verification not implemented)	596
3.100.8 Giac [A] (verification not implemented)	596
3.100.9 Mupad [B] (verification not implemented)	597

3.100.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \cot^4(x) \csc^4(x) dx = -\frac{1}{5} \cot^5(x) - \frac{\cot^7(x)}{7}$$

output `-1/5*cot(x)^5-1/7*cot(x)^7`

3.100.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 37 vs. 2(17) = 34.

Time = 0.05 (sec) , antiderivative size = 37, normalized size of antiderivative = 2.18

$$\int \cot^4(x) \csc^4(x) dx = -\frac{2 \cot(x)}{35} - \frac{1}{35} \cot(x) \csc^2(x) + \frac{8}{35} \cot(x) \csc^4(x) - \frac{1}{7} \cot(x) \csc^6(x)$$

input `Integrate[Cot[x]^4*Csc[x]^4,x]`

output `(-2*Cot[x])/35 - (Cot[x]*Csc[x]^2)/35 + (8*Cot[x]*Csc[x]^4)/35 - (Cot[x]*Csc[x]^6)/7`

3.100.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 3087, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^4(x) \csc^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan\left(x - \frac{\pi}{2}\right)^4 \sec\left(x - \frac{\pi}{2}\right)^4 dx \\
 & \quad \downarrow \text{3087} \\
 & \int \cot^4(x) (\cot^2(x) + 1) d(-\cot(x)) \\
 & \quad \downarrow \text{244} \\
 & \int (\cot^6(x) + \cot^4(x)) d(-\cot(x)) \\
 & \quad \downarrow \text{2009} \\
 & -\frac{1}{7} \cot^7(x) - \frac{\cot^5(x)}{5}
 \end{aligned}$$

input `Int[Cot[x]^4*Csc[x]^4,x]`

output `-1/5*Cot[x]^5 - Cot[x]^7/7`

3.100.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] :> Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3087 `Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[1/f Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])`

3.100.4 Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$-\frac{(\cot^5(x))}{5} - \frac{(\cot^7(x))}{7}$	14
default	$-\frac{(\cot^5(x))}{5} - \frac{(\cot^7(x))}{7}$	14
risch	$\frac{4i(35e^{10ix} + 35e^{8ix} + 70e^{6ix} + 14e^{4ix} + 7e^{2ix} - 1)}{35(e^{2ix} - 1)^7}$	50

input `int(cot(x)^4*csc(x)^4,x,method=_RETURNVERBOSE)`

output `-1/5*cot(x)^5-1/7*cot(x)^7`

3.100.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. $2(13) = 26$.

Time = 0.25 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.29

$$\int \cot^4(x) \csc^4(x) dx = -\frac{2 \cos(x)^7 - 7 \cos(x)^5}{35 (\cos(x)^6 - 3 \cos(x)^4 + 3 \cos(x)^2 - 1) \sin(x)}$$

input `integrate(cot(x)^4*csc(x)^4,x, algorithm="fracas")`

output `-1/35*(2*cos(x)^7 - 7*cos(x)^5)/((cos(x)^6 - 3*cos(x)^4 + 3*cos(x)^2 - 1)*sin(x))`

3.100.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 41 vs. $2(14) = 28$.

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 2.41

$$\int \cot^4(x) \csc^4(x) dx = -\frac{2 \cos(x)}{35 \sin(x)} - \frac{\cos(x)}{35 \sin^3(x)} + \frac{8 \cos(x)}{35 \sin^5(x)} - \frac{\cos(x)}{7 \sin^7(x)}$$

input `integrate(cot(x)**4*csc(x)**4,x)`

output `-2*cos(x)/(35*sin(x)) - cos(x)/(35*sin(x)**3) + 8*cos(x)/(35*sin(x)**5) - cos(x)/(7*sin(x)**7)`

3.100.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \cot^4(x) \csc^4(x) dx = -\frac{7 \tan(x)^2 + 5}{35 \tan(x)^7}$$

input `integrate(cot(x)^4*csc(x)^4,x, algorithm="maxima")`

output `-1/35*(7*tan(x)^2 + 5)/tan(x)^7`

3.100.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \cot^4(x) \csc^4(x) dx = -\frac{7 \tan(x)^2 + 5}{35 \tan(x)^7}$$

input `integrate(cot(x)^4*csc(x)^4,x, algorithm="giac")`

output `-1/35*(7*tan(x)^2 + 5)/tan(x)^7`

3.100.9 Mupad [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \cot^4(x) \csc^4(x) dx = -\frac{\cot(x)^5 (5 \cot(x)^2 + 7)}{35}$$

input `int(cot(x)^4/sin(x)^4,x)`

output `-(cot(x)^5*(5*cot(x)^2 + 7))/35`

3.101 $\int \cot^3(x) \csc^4(x) dx$

3.101.1 Optimal result	598
3.101.2 Mathematica [A] (verified)	598
3.101.3 Rubi [A] (verified)	599
3.101.4 Maple [A] (verified)	600
3.101.5 Fricas [B] (verification not implemented)	601
3.101.6 Sympy [A] (verification not implemented)	601
3.101.7 Maxima [A] (verification not implemented)	601
3.101.8 Giac [A] (verification not implemented)	602
3.101.9 Mupad [B] (verification not implemented)	602

3.101.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \cot^3(x) \csc^4(x) dx = \frac{\csc^4(x)}{4} - \frac{\csc^6(x)}{6}$$

output `1/4*csc(x)^4-1/6*csc(x)^6`

3.101.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cot^3(x) \csc^4(x) dx = \frac{\csc^4(x)}{4} - \frac{\csc^6(x)}{6}$$

input `Integrate[Cot[x]^3*Csc[x]^4,x]`

output `Csc[x]^4/4 - Csc[x]^6/6`

3.101.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 25, 3086, 25, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^3(x) \csc^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan\left(x - \frac{\pi}{2}\right)^3 \left(-\sec\left(x - \frac{\pi}{2}\right)\right)^4 dx \\
 & \quad \downarrow \text{25} \\
 & - \int \sec\left(x - \frac{\pi}{2}\right)^4 \tan\left(x - \frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{3086} \\
 & - \int -\csc^3(x) (1 - \csc^2(x)) d \csc(x) \\
 & \quad \downarrow \text{25} \\
 & \int \csc^3(x) (1 - \csc^2(x)) d \csc(x) \\
 & \quad \downarrow \text{244} \\
 & \int (\csc^3(x) - \csc^5(x)) d \csc(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\csc^4(x)}{4} - \frac{\csc^6(x)}{6}
 \end{aligned}$$

input `Int[Cot[x]^3*Csc[x]^4,x]`

output `Csc[x]^4/4 - Csc[x]^6/6`

3.101.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 244 `Int[((c_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

3.101.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$-\frac{(\cot^6(x))}{6} - \frac{(\cot^4(x))}{4}$	14
default	$-\frac{(\cot^6(x))}{6} - \frac{(\cot^4(x))}{4}$	14
risch	$\frac{4e^{8ix} + \frac{8e^{6ix}}{3} + 4e^{4ix}}{(e^{2ix} - 1)^6}$	34

input `int(cot(x)^3*csc(x)^4,x,method=_RETURNVERBOSE)`

output `-1/6*cot(x)^6-1/4*cot(x)^4`

3.101.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 30 vs. $2(13) = 26$.

Time = 0.25 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.76

$$\int \cot^3(x) \csc^4(x) dx = \frac{3 \cos(x)^2 - 1}{12 (\cos(x)^6 - 3 \cos(x)^4 + 3 \cos(x)^2 - 1)}$$

input `integrate(cot(x)^3*csc(x)^4,x, algorithm="fricas")`

output `1/12*(3*cos(x)^2 - 1)/(cos(x)^6 - 3*cos(x)^4 + 3*cos(x)^2 - 1)`

3.101.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \cot^3(x) \csc^4(x) dx = -\frac{2 - 3 \sin^2(x)}{12 \sin^6(x)}$$

input `integrate(cot(x)**3*csc(x)**4,x)`

output `-(2 - 3*sin(x)**2)/(12*sin(x)**6)`

3.101.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \cot^3(x) \csc^4(x) dx = \frac{3 \sin(x)^2 - 2}{12 \sin(x)^6}$$

input `integrate(cot(x)^3*csc(x)^4,x, algorithm="maxima")`

output `1/12*(3*sin(x)^2 - 2)/sin(x)^6`

3.101.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \cot^3(x) \csc^4(x) dx = \frac{3 \sin(x)^2 - 2}{12 \sin(x)^6}$$

input `integrate(cot(x)^3*csc(x)^4,x, algorithm="giac")`output `1/12*(3*sin(x)^2 - 2)/sin(x)^6`**3.101.9 Mupad [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \cot^3(x) \csc^4(x) dx = -\frac{\cot(x)^4 (2 \cot(x)^2 + 3)}{12}$$

input `int(cot(x)^3/sin(x)^4,x)`output `-(cot(x)^4*(2*cot(x)^2 + 3))/12`

3.102 $\int \csc(x) dx$

3.102.1 Optimal result	603
3.102.2 Mathematica [B] (verified)	603
3.102.3 Rubi [A] (verified)	604
3.102.4 Maple [A] (verified)	605
3.102.5 Fricas [B] (verification not implemented)	605
3.102.6 Sympy [B] (verification not implemented)	605
3.102.7 Maxima [A] (verification not implemented)	606
3.102.8 Giac [A] (verification not implemented)	606
3.102.9 Mupad [B] (verification not implemented)	606

3.102.1 Optimal result

Integrand size = 2, antiderivative size = 5

$$\int \csc(x) dx = -\operatorname{arctanh}(\cos(x))$$

output `-arctanh(cos(x))`

3.102.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 17 vs. $2(5) = 10$.

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 3.40

$$\int \csc(x) dx = -\log\left(\cos\left(\frac{x}{2}\right)\right) + \log\left(\sin\left(\frac{x}{2}\right)\right)$$

input `Integrate[Csc[x], x]`

output `-Log[Cos[x/2]] + Log[Sin[x/2]]`

3.102.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \csc(x) dx \\ \downarrow 3042 \\ \int \csc(x) dx \\ \downarrow 4257 \\ -\operatorname{arctanh}(\cos(x)) \end{array}$$

input `Int [Csc [x] , x]`

output `-ArcTanh [Cos [x]]`

3.102.3.1 Defintions of rubi rules used

rule 3042 `Int [u_ , x_Symbol] :> Int [DeactivateTrig [u , x] , x] /; FunctionOfTrigOfLinear Q [u , x]`

rule 4257 `Int [csc [(c_.) + (d_.)*(x_)], x_Symbol] :> Simp [-ArcTanh [Cos [c + d*x]]/d , x] /; FreeQ [{c , d} , x]`

3.102.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
norman	$\ln\left(\tan\left(\frac{x}{2}\right)\right)$	6
parallelrisc	$\ln\left(\tan\left(\frac{x}{2}\right)\right)$	6
lookup	$-\ln(\csc(x) + \cot(x))$	9
default	$-\ln(\csc(x) + \cot(x))$	9
risc	$\ln(e^{ix} - 1) - \ln(e^{ix} + 1)$	20

input `int(csc(x),x,method=_RETURNVERBOSE)`

output `ln(tan(1/2*x))`

3.102.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(5) = 10$.

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 3.80

$$\int \csc(x) dx = -\frac{1}{2} \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) + \frac{1}{2} \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right)$$

input `integrate(csc(x),x, algorithm="fricas")`

output `-1/2*log(1/2*cos(x) + 1/2) + 1/2*log(-1/2*cos(x) + 1/2)`

3.102.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(5) = 10$.

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 3.00

$$\int \csc(x) dx = \frac{\log(\cos(x) - 1)}{2} - \frac{\log(\cos(x) + 1)}{2}$$

input `integrate(csc(x),x)`

output `log(cos(x) - 1)/2 - log(cos(x) + 1)/2`

3.102.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.60

$$\int \csc(x) dx = -\log(\cot(x) + \csc(x))$$

input `integrate(csc(x),x, algorithm="maxima")`

output `-log(cot(x) + csc(x))`

3.102.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

$$\int \csc(x) dx = \log\left(\left|\tan\left(\frac{1}{2}x\right)\right|\right)$$

input `integrate(csc(x),x, algorithm="giac")`

output `log(abs(tan(1/2*x)))`

3.102.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \csc(x) dx = \ln\left(\tan\left(\frac{x}{2}\right)\right)$$

input `int(1/sin(x),x)`

output `log(tan(x/2))`

3.103 $\int \csc^3(x) dx$

3.103.1 Optimal result	607
3.103.2 Mathematica [B] (verified)	607
3.103.3 Rubi [A] (verified)	608
3.103.4 Maple [A] (verified)	609
3.103.5 Fricas [B] (verification not implemented)	609
3.103.6 Sympy [A] (verification not implemented)	610
3.103.7 Maxima [B] (verification not implemented)	610
3.103.8 Giac [B] (verification not implemented)	610
3.103.9 Mupad [B] (verification not implemented)	611

3.103.1 Optimal result

Integrand size = 4, antiderivative size = 16

$$\int \csc^3(x) dx = -\frac{1}{2} \operatorname{arctanh}(\cos(x)) - \frac{1}{2} \cot(x) \csc(x)$$

output `-1/2*arctanh(cos(x))-1/2*cot(x)*csc(x)`

3.103.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 47 vs. $2(16) = 32$.

Time = 0.01 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.94

$$\int \csc^3(x) dx = -\frac{1}{8} \csc^2\left(\frac{x}{2}\right) - \frac{1}{2} \log\left(\cos\left(\frac{x}{2}\right)\right) + \frac{1}{2} \log\left(\sin\left(\frac{x}{2}\right)\right) + \frac{1}{8} \sec^2\left(\frac{x}{2}\right)$$

input `Integrate[Csc[x]^3,x]`

output `-1/8*Csc[x/2]^2 - Log[Cos[x/2]]/2 + Log[Sin[x/2]]/2 + Sec[x/2]^2/8`

3.103.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \csc^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc(x)^3 dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{\int \csc(x) dx}{2} - \frac{1}{2} \cot(x) \csc(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \csc(x) dx}{2} - \frac{1}{2} \cot(x) \csc(x) \\
 & \quad \downarrow \text{4257} \\
 & -\frac{1}{2} \operatorname{arctanh}(\cos(x)) - \frac{1}{2} \cot(x) \csc(x)
 \end{aligned}$$

input `Int[Csc[x]^3,x]`

output `-1/2*ArcTanh[Cos[x]] - (Cot[x]*Csc[x])/2`

3.103.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
;/; FreeQ[{c, d}, x]
```

3.103.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

method	result	size
default	$-\frac{\csc(x)\cot(x)}{2} + \frac{\ln(\csc(x)-\cot(x))}{2}$	18
parallelrisc	$-\frac{\csc(x)\cot(x)}{2} + \ln\left(\sqrt{\csc(x)-\cot(x)}\right)$	18
norman	$-\frac{\frac{1}{8} + \frac{(\tan^4(\frac{x}{2}))}{8}}{\tan(\frac{x}{2})^2} + \frac{\ln(\tan(\frac{x}{2}))}{2}$	26
risc	$\frac{e^{3ix}+e^{ix}}{(e^{2ix}-1)^2} + \frac{\ln(e^{ix}-1)}{2} - \frac{\ln(e^{ix}+1)}{2}$	43

```
input int(csc(x)^3,x,method=_RETURNVERBOSE)
```

```
output -1/2*csc(x)*cot(x)+1/2*ln(csc(x)-cot(x))
```

3.103.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 44 vs. $2(12) = 24$.

Time = 0.25 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.75

$$\int \csc^3(x) dx$$

$$= -\frac{(\cos(x)^2 - 1) \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - (\cos(x)^2 - 1) \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right) - 2 \cos(x)}{4(\cos(x)^2 - 1)}$$

```
input integrate(csc(x)^3,x, algorithm="fricas")
```

```
output -1/4*((cos(x)^2 - 1)*log(1/2*cos(x) + 1/2) - (cos(x)^2 - 1)*log(-1/2*cos(x)
) + 1/2) - 2*cos(x))/(cos(x)^2 - 1)
```

3.103.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.69

$$\int \csc^3(x) dx = \frac{\log(\cos(x) - 1)}{4} - \frac{\log(\cos(x) + 1)}{4} + \frac{\cos(x)}{2\cos^2(x) - 2}$$

input `integrate(csc(x)**3,x)`

output `log(cos(x) - 1)/4 - log(cos(x) + 1)/4 + cos(x)/(2*cos(x)**2 - 2)`

3.103.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(12) = 24.

Time = 0.19 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.69

$$\int \csc^3(x) dx = \frac{\cos(x)}{2(\cos(x)^2 - 1)} - \frac{1}{4} \log(\cos(x) + 1) + \frac{1}{4} \log(\cos(x) - 1)$$

input `integrate(csc(x)^3,x, algorithm="maxima")`

output `1/2*cos(x)/(cos(x)^2 - 1) - 1/4*log(cos(x) + 1) + 1/4*log(cos(x) - 1)`

3.103.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 54 vs. 2(12) = 24.

Time = 0.28 (sec) , antiderivative size = 54, normalized size of antiderivative = 3.38

$$\int \csc^3(x) dx = -\frac{\left(\frac{2(\cos(x)-1)}{\cos(x)+1} - 1\right)(\cos(x) + 1)}{8(\cos(x) - 1)} - \frac{\cos(x) - 1}{8(\cos(x) + 1)} + \frac{1}{4} \log\left(-\frac{\cos(x) - 1}{\cos(x) + 1}\right)$$

input `integrate(csc(x)^3,x, algorithm="giac")`

output `-1/8*(2*(cos(x) - 1)/(cos(x) + 1) - 1)*(cos(x) + 1)/(cos(x) - 1) - 1/8*(cos(x) - 1)/(cos(x) + 1) + 1/4*log(-(cos(x) - 1)/(cos(x) + 1))`

3.103.9 Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \csc^3(x) dx = \frac{\ln\left(\tan\left(\frac{x}{2}\right)\right)}{2} - \frac{\cos(x)}{2\sin(x)^2}$$

input `int(1/sin(x)^3,x)`

output `log(tan(x/2))/2 - cos(x)/(2*sin(x)^2)`

3.104 $\int \cos(x) \cot(x) dx$

3.104.1 Optimal result	612
3.104.2 Mathematica [B] (verified)	612
3.104.3 Rubi [A] (verified)	613
3.104.4 Maple [A] (verified)	614
3.104.5 Fricas [B] (verification not implemented)	615
3.104.6 Sympy [B] (verification not implemented)	615
3.104.7 Maxima [B] (verification not implemented)	615
3.104.8 Giac [B] (verification not implemented)	616
3.104.9 Mupad [B] (verification not implemented)	616

3.104.1 Optimal result

Integrand size = 5, antiderivative size = 8

$$\int \cos(x) \cot(x) dx = -\operatorname{arctanh}(\cos(x)) + \cos(x)$$

output `-arctanh(cos(x))+cos(x)`

3.104.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 19 vs. $2(8) = 16$.

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.38

$$\int \cos(x) \cot(x) dx = \cos(x) - \log\left(\cos\left(\frac{x}{2}\right)\right) + \log\left(\sin\left(\frac{x}{2}\right)\right)$$

input `Integrate[Cos[x]*Cot[x],x]`

output `Cos[x] - Log[Cos[x/2]] + Log[Sin[x/2]]`

3.104.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 25, 3072, 262, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(x) \cot(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\sin\left(x + \frac{\pi}{2}\right) \tan\left(x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{25} \\
 & -\int \sin\left(x + \frac{\pi}{2}\right) \tan\left(x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3072} \\
 & -\int \frac{\cos^2(x)}{1 - \cos^2(x)} d\cos(x) \\
 & \quad \downarrow \text{262} \\
 & \cos(x) - \int \frac{1}{1 - \cos^2(x)} d\cos(x) \\
 & \quad \downarrow \text{219} \\
 & \cos(x) - \operatorname{arctanh}(\cos(x))
 \end{aligned}$$

input `Int[Cos[x]*Cot[x],x]`

output `-ArcTanh[Cos[x]] + Cos[x]`

3.104.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 262 `Int[((c_)*(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*(m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3072 `Int[((a_)*sin[(e_) + (f_)*(x_)])^(m_)*tan[(e_) + (f_)*(x_)^(n_)], x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Simp[ff/f Subst[Int[(ff*x)^(m + n)/(a^2 - ff^2*x^2)^((n + 1)/2), x], x, a*(Sin[e + f*x]/ff)], x]] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2]`

3.104.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.50

method	result	size
default	$\cos(x) + \ln(\csc(x) - \cot(x))$	12
parallelrisc	$\cos(x) + \ln(\csc(x) - \cot(x)) + 1$	13
norman	$\frac{2}{1 + \tan^2(\frac{x}{2})} + \ln\left(\tan\left(\frac{x}{2}\right)\right)$	19
risc	$\frac{e^{ix}}{2} + \frac{e^{-ix}}{2} + \ln(e^{ix} - 1) - \ln(e^{ix} + 1)$	34

input `int(cos(x)^2/sin(x), x, method=_RETURNVERBOSE)`

output `cos(x)+ln(csc(x)-cot(x))`

3.104.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. $2(8) = 16$.

Time = 0.24 (sec) , antiderivative size = 21, normalized size of antiderivative = 2.62

$$\int \cos(x) \cot(x) dx = \cos(x) - \frac{1}{2} \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) + \frac{1}{2} \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right)$$

input `integrate(cos(x)^2/sin(x),x, algorithm="fracas")`

output `cos(x) - 1/2*log(1/2*cos(x) + 1/2) + 1/2*log(-1/2*cos(x) + 1/2)`

3.104.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(7) = 14$.

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.38

$$\int \cos(x) \cot(x) dx = \frac{\log(\cos(x) - 1)}{2} - \frac{\log(\cos(x) + 1)}{2} + \cos(x)$$

input `integrate(cos(x)**2/sin(x),x)`

output `log(cos(x) - 1)/2 - log(cos(x) + 1)/2 + cos(x)`

3.104.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(8) = 16$.

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.12

$$\int \cos(x) \cot(x) dx = \cos(x) - \frac{1}{2} \log(\cos(x) + 1) + \frac{1}{2} \log(\cos(x) - 1)$$

input `integrate(cos(x)^2/sin(x),x, algorithm="maxima")`

output `cos(x) - 1/2*log(cos(x) + 1) + 1/2*log(cos(x) - 1)`

3.104.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(8) = 16$.

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.38

$$\int \cos(x) \cot(x) dx = \cos(x) - \frac{1}{2} \log(\cos(x) + 1) + \frac{1}{2} \log(-\cos(x) + 1)$$

input `integrate(cos(x)^2/sin(x),x, algorithm="giac")`

output `cos(x) - 1/2*log(cos(x) + 1) + 1/2*log(-cos(x) + 1)`

3.104.9 Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \cos(x) \cot(x) dx = \ln\left(\tan\left(\frac{x}{2}\right)\right) + \cos(x)$$

input `int(cos(x)^2/sin(x),x)`

output `log(tan(x/2)) + cos(x)`

3.105 $\int \csc^4(x) dx$

3.105.1 Optimal result	617
3.105.2 Mathematica [A] (verified)	617
3.105.3 Rubi [A] (verified)	618
3.105.4 Maple [A] (verified)	619
3.105.5 Fricas [B] (verification not implemented)	619
3.105.6 Sympy [A] (verification not implemented)	619
3.105.7 Maxima [A] (verification not implemented)	620
3.105.8 Giac [A] (verification not implemented)	620
3.105.9 Mupad [B] (verification not implemented)	620

3.105.1 Optimal result

Integrand size = 4, antiderivative size = 13

$$\int \csc^4(x) dx = -\cot(x) - \frac{\cot^3(x)}{3}$$

output `-cot(x)-1/3*cot(x)^3`

3.105.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31

$$\int \csc^4(x) dx = -\frac{2 \cot(x)}{3} - \frac{1}{3} \cot(x) \csc^2(x)$$

input `Integrate[Csc[x]^4,x]`

output `(-2*Cot[x])/3 - (Cot[x]*Csc[x]^2)/3`

3.105.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \csc^4(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \csc(x)^4 dx \\ & \quad \downarrow \text{4254} \\ & - \int (\cot^2(x) + 1) d \cot(x) \\ & \quad \downarrow \text{2009} \\ & -\frac{1}{3} \cot^3(x) - \cot(x) \end{aligned}$$

input `Int[Csc[x]^4,x]`

output `-Cot[x] - Cot[x]^3/3`

3.105.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

3.105.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
default	$\left(-\frac{2}{3} - \frac{\csc^2(x)}{3}\right) \cot(x)$	12
parallelrisc	$\frac{2(\cot^3(x))}{3} - \cot(x) (\csc^2(x))$	16
risc	$\frac{4i(3e^{2ix}-1)}{3(e^{2ix}-1)^3}$	22
norman	$\frac{-\frac{1}{24} - \frac{3(\tan^2(\frac{x}{2}))}{8} + \frac{3(\tan^4(\frac{x}{2}))}{8} + \frac{(\tan^6(\frac{x}{2}))}{24}}{\tan(\frac{x}{2})^3}$	34

input `int(1/sin(x)^4,x,method=_RETURNVERBOSE)`

output `(-2/3-1/3*csc(x)^2)*cot(x)`

3.105.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. $2(11) = 22$.

Time = 0.24 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.92

$$\int \csc^4(x) dx = -\frac{2 \cos(x)^3 - 3 \cos(x)}{3 (\cos(x)^2 - 1) \sin(x)}$$

input `integrate(1/sin(x)^4,x, algorithm="fricas")`

output `-1/3*(2*cos(x)^3 - 3*cos(x))/((cos(x)^2 - 1)*sin(x))`

3.105.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.54

$$\int \csc^4(x) dx = -\frac{2 \cos(x)}{3 \sin(x)} - \frac{\cos(x)}{3 \sin^3(x)}$$

input `integrate(1/sin(x)**4,x)`

output `-2*cos(x)/(3*sin(x)) - cos(x)/(3*sin(x)**3)`

3.105.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.08

$$\int \csc^4(x) dx = -\frac{3 \tan(x)^2 + 1}{3 \tan(x)^3}$$

input `integrate(1/sin(x)^4,x, algorithm="maxima")`

output `-1/3*(3*tan(x)^2 + 1)/tan(x)^3`

3.105.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.08

$$\int \csc^4(x) dx = -\frac{3 \tan(x)^2 + 1}{3 \tan(x)^3}$$

input `integrate(1/sin(x)^4,x, algorithm="giac")`

output `-1/3*(3*tan(x)^2 + 1)/tan(x)^3`

3.105.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31

$$\int \csc^4(x) dx = -\frac{2 \cos(x) \sin(x)^2 + \cos(x)}{3 \sin(x)^3}$$

input `int(1/sin(x)^4,x)`

output `-(cos(x) + 2*cos(x)*sin(x)^2)/(3*sin(x)^3)`

3.106 $\int \sin(2x) \sin(5x) dx$

3.106.1 Optimal result	621
3.106.2 Mathematica [A] (verified)	621
3.106.3 Rubi [A] (verified)	622
3.106.4 Maple [A] (verified)	623
3.106.5 Fricas [A] (verification not implemented)	623
3.106.6 Sympy [B] (verification not implemented)	623
3.106.7 Maxima [A] (verification not implemented)	624
3.106.8 Giac [A] (verification not implemented)	624
3.106.9 Mupad [B] (verification not implemented)	624

3.106.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \sin(2x) \sin(5x) dx = \frac{1}{6} \sin(3x) - \frac{1}{14} \sin(7x)$$

output `1/6*sin(3*x)-1/14*sin(7*x)`

3.106.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sin(2x) \sin(5x) dx = \frac{1}{6} \sin(3x) - \frac{1}{14} \sin(7x)$$

input `Integrate[Sin[2*x]*Sin[5*x],x]`

output `Sin[3*x]/6 - Sin[7*x]/14`

3.106.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4770}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sin(2x) \sin(5x) dx$$

$$\downarrow \text{3042}$$

$$\int \sin(2x) \sin(5x) dx$$

$$\downarrow \text{4770}$$

$$\frac{1}{6} \sin(3x) - \frac{1}{14} \sin(7x)$$

input `Int[Sin[2*x]*Sin[5*x],x]`

output `Sin[3*x]/6 - Sin[7*x]/14`

3.106.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4770 `Int[sin[(a_.) + (b_.)*(x_)]*sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[a - c + (b - d)*x]/(2*(b - d)), x] - Simp[Sin[a + c + (b + d)*x]/(2*(b + d)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

3.106.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{\sin(3x)}{6} - \frac{\sin(7x)}{14}$	14
risch	$\frac{\sin(3x)}{6} - \frac{\sin(7x)}{14}$	14
parallelrisch	$\frac{(-\sin(\frac{3x}{2}) + 3\sin(\frac{x}{2}))(\cos(\frac{3x}{2}) + 3\cos(\frac{x}{2}))(11 + 6\cos(4x) + 18\cos(2x))}{21}$	41
norman	$\frac{10 \tan(x) \tan^2(\frac{5x}{2})}{21} - \frac{4(\tan^2(x) \tan(\frac{5x}{2}))}{21} - \frac{10 \tan(x)}{21} + \frac{4 \tan(\frac{5x}{2})}{21}$ $\frac{1}{(1 + \tan^2(x))(1 + \tan^2(\frac{5x}{2}))}$	51

input `int(sin(2*x)*sin(5*x),x,method=_RETURNVERBOSE)`output `1/6*sin(3*x)-1/14*sin(7*x)`**3.106.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.41

$$\int \sin(2x) \sin(5x) dx = -\frac{2}{21} (48 \cos(x)^6 - 60 \cos(x)^4 + 11 \cos(x)^2 + 1) \sin(x)$$

input `integrate(sin(2*x)*sin(5*x),x, algorithm="fracas")`output `-2/21*(48*cos(x)^6 - 60*cos(x)^4 + 11*cos(x)^2 + 1)*sin(x)`**3.106.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. 2(12) = 24.

Time = 0.13 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.53

$$\int \sin(2x) \sin(5x) dx = -\frac{5 \sin(2x) \cos(5x)}{21} + \frac{2 \sin(5x) \cos(2x)}{21}$$

input `integrate(sin(2*x)*sin(5*x),x)`output `-5*sin(2*x)*cos(5*x)/21 + 2*sin(5*x)*cos(2*x)/21`

3.106.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sin(2x) \sin(5x) dx = -\frac{1}{14} \sin(7x) + \frac{1}{6} \sin(3x)$$

input `integrate(sin(2*x)*sin(5*x),x, algorithm="maxima")`output `-1/14*sin(7*x) + 1/6*sin(3*x)`**3.106.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sin(2x) \sin(5x) dx = -\frac{1}{14} \sin(7x) + \frac{1}{6} \sin(3x)$$

input `integrate(sin(2*x)*sin(5*x),x, algorithm="giac")`output `-1/14*sin(7*x) + 1/6*sin(3*x)`**3.106.9 Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sin(2x) \sin(5x) dx = \frac{\sin(3x)}{6} - \frac{\sin(7x)}{14}$$

input `int(sin(2*x)*sin(5*x),x)`output `sin(3*x)/6 - sin(7*x)/14`

3.107 $\int \cos(x) \sin(3x) dx$

3.107.1 Optimal result	625
3.107.2 Mathematica [A] (verified)	625
3.107.3 Rubi [A] (verified)	626
3.107.4 Maple [A] (verified)	627
3.107.5 Fricas [A] (verification not implemented)	627
3.107.6 Sympy [A] (verification not implemented)	627
3.107.7 Maxima [A] (verification not implemented)	628
3.107.8 Giac [A] (verification not implemented)	628
3.107.9 Mupad [B] (verification not implemented)	628

3.107.1 Optimal result

Integrand size = 7, antiderivative size = 17

$$\int \cos(x) \sin(3x) dx = -\frac{1}{4} \cos(2x) - \frac{1}{8} \cos(4x)$$

output `-1/4*cos(2*x)-1/8*cos(4*x)`

3.107.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cos(x) \sin(3x) dx = -\frac{1}{2} \cos^2(x) - \frac{1}{8} \cos(4x)$$

input `Integrate[Cos[x]*Sin[3*x],x]`

output `-1/2*Cos[x]^2 - Cos[4*x]/8`

3.107.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4772}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sin(3x) \cos(x) dx \\ \downarrow 3042 \\ \int \sin(3x) \cos(x) dx \\ \downarrow 4772 \\ -\frac{1}{4} \cos(2x) - \frac{1}{8} \cos(4x) \end{array}$$

input `Int[Cos[x]*Sin[3*x],x]`

output `-1/4*Cos[2*x] - Cos[4*x]/8`

3.107.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4772 `Int[cos[(c_.) + (d_.)*(x_)]*sin[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[-Cos[a - c + (b - d)*x]/(2*(b - d)), x] - Simp[Cos[a + c + (b + d)*x]/(2*(b + d)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

3.107.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{\cos(2x)}{4} - \frac{\cos(4x)}{8}$	14
risch	$-\frac{\cos(2x)}{4} - \frac{\cos(4x)}{8}$	14
parallelrisch	$-\frac{\cos(4x)}{8} + \frac{3}{8} - \frac{\cos(2x)}{4}$	15
norman	$\frac{3(\tan^2(\frac{x}{2}))}{4} + \frac{3(\tan^2(\frac{3x}{2}))}{4} - \frac{\tan(\frac{x}{2})\tan(\frac{3x}{2})}{2}$ $\frac{1}{(1+\tan^2(\frac{x}{2}))(1+\tan^2(\frac{3x}{2}))}$	49

input `int(cos(x)*sin(3*x),x,method=_RETURNVERBOSE)`output `-1/4*cos(2*x)-1/8*cos(4*x)`**3.107.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(x) \sin(3x) dx = -\cos(x)^4 + \frac{1}{2} \cos(x)^2$$

input `integrate(cos(x)*sin(3*x),x, algorithm="fricas")`output `-cos(x)^4 + 1/2*cos(x)^2`**3.107.6 Sympy [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.29

$$\int \cos(x) \sin(3x) dx = -\frac{\sin(x) \sin(3x)}{8} - \frac{3 \cos(x) \cos(3x)}{8}$$

input `integrate(cos(x)*sin(3*x),x)`output `-sin(x)*sin(3*x)/8 - 3*cos(x)*cos(3*x)/8`

3.107.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(x) \sin(3x) dx = -\frac{1}{8} \cos(4x) - \frac{1}{4} \cos(2x)$$

input `integrate(cos(x)*sin(3*x),x, algorithm="maxima")`output `-1/8*cos(4*x) - 1/4*cos(2*x)`**3.107.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(x) \sin(3x) dx = -\cos(x)^4 + \frac{1}{2} \cos(x)^2$$

input `integrate(cos(x)*sin(3*x),x, algorithm="giac")`output `cos(x)^2/2 - cos(x)^4`**3.107.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(x) \sin(3x) dx = \frac{\cos(x)^2}{2} - \cos(x)^4$$

input `int(sin(3*x)*cos(x),x)`output `cos(x)^2/2 - cos(x)^4`

3.108 $\int \cos(3x) \cos(4x) dx$

3.108.1 Optimal result	629
3.108.2 Mathematica [A] (verified)	629
3.108.3 Rubi [A] (verified)	630
3.108.4 Maple [A] (verified)	631
3.108.5 Fricas [B] (verification not implemented)	631
3.108.6 Sympy [B] (verification not implemented)	631
3.108.7 Maxima [A] (verification not implemented)	632
3.108.8 Giac [A] (verification not implemented)	632
3.108.9 Mupad [B] (verification not implemented)	632

3.108.1 Optimal result

Integrand size = 9, antiderivative size = 15

$$\int \cos(3x) \cos(4x) dx = \frac{\sin(x)}{2} + \frac{1}{14} \sin(7x)$$

output `1/2*sin(x)+1/14*sin(7*x)`

3.108.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \cos(3x) \cos(4x) dx = \frac{\sin(x)}{2} + \frac{1}{14} \sin(7x)$$

input `Integrate[Cos[3*x]*Cos[4*x],x]`

output `Sin[x]/2 + Sin[7*x]/14`

3.108.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4771}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(3x) \cos(4x) dx$$

$$\downarrow \text{3042}$$

$$\int \cos(3x) \cos(4x) dx$$

$$\downarrow \text{4771}$$

$$\frac{\sin(x)}{2} + \frac{1}{14} \sin(7x)$$

input `Int[Cos[3*x]*Cos[4*x],x]`

output `Sin[x]/2 + Sin[7*x]/14`

3.108.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4771 `Int[cos[(a_.) + (b_.)*(x_)]*cos[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[a - c + (b - d)*x]/(2*(b - d)), x] + Simp[Sin[a + c + (b + d)*x]/(2*(b + d)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

3.108.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

method	result	size
default	$\frac{\sin(x)}{2} + \frac{\sin(7x)}{14}$	12
risch	$\frac{\sin(x)}{2} + \frac{\sin(7x)}{14}$	12
parallelrisch	$\frac{\sin(x)}{2} + \frac{\sin(7x)}{14}$	12
norman	$-\frac{8 \tan(2x) \left(\tan^2\left(\frac{3x}{2}\right) \right) + 6 \left(\tan^2(2x) \right) \tan\left(\frac{3x}{2}\right) + 8 \tan(2x) - 6 \tan\left(\frac{3x}{2}\right)}{(1+\tan^2\left(\frac{3x}{2}\right))(1+\tan^2(2x))}$	59

input `int(cos(3*x)*cos(4*x),x,method=_RETURNVERBOSE)`

output `1/2*sin(x)+1/14*sin(7*x)`

3.108.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 24 vs. $2(11) = 22$.

Time = 0.26 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.60

$$\int \cos(3x) \cos(4x) dx = \frac{1}{7} (32 \cos(x)^6 - 40 \cos(x)^4 + 12 \cos(x)^2 + 3) \sin(x)$$

input `integrate(cos(3*x)*cos(4*x),x, algorithm="fricas")`

output `1/7*(32*cos(x)^6 - 40*cos(x)^4 + 12*cos(x)^2 + 3)*sin(x)`

3.108.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. $2(10) = 20$.

Time = 0.14 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.73

$$\int \cos(3x) \cos(4x) dx = -\frac{3 \sin(3x) \cos(4x)}{7} + \frac{4 \sin(4x) \cos(3x)}{7}$$

input `integrate(cos(3*x)*cos(4*x),x)`

output `-3*sin(3*x)*cos(4*x)/7 + 4*sin(4*x)*cos(3*x)/7`

3.108.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \cos(3x) \cos(4x) dx = \frac{1}{14} \sin(7x) + \frac{1}{2} \sin(x)$$

input `integrate(cos(3*x)*cos(4*x),x, algorithm="maxima")`

output `1/14*sin(7*x) + 1/2*sin(x)`

3.108.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \cos(3x) \cos(4x) dx = \frac{1}{14} \sin(7x) + \frac{1}{2} \sin(x)$$

input `integrate(cos(3*x)*cos(4*x),x, algorithm="giac")`

output `1/14*sin(7*x) + 1/2*sin(x)`

3.108.9 Mupad [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \cos(3x) \cos(4x) dx = \frac{\sin(7x)}{14} + \frac{\sin(x)}{2}$$

input `int(cos(3*x)*cos(4*x),x)`

output `sin(7*x)/14 + sin(x)/2`

3.109 $\int \sin(3x) \sin(6x) dx$

3.109.1 Optimal result	633
3.109.2 Mathematica [A] (verified)	633
3.109.3 Rubi [A] (verified)	634
3.109.4 Maple [A] (verified)	635
3.109.5 Fricas [A] (verification not implemented)	635
3.109.6 Sympy [A] (verification not implemented)	635
3.109.7 Maxima [A] (verification not implemented)	636
3.109.8 Giac [A] (verification not implemented)	636
3.109.9 Mupad [B] (verification not implemented)	636

3.109.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \sin(3x) \sin(6x) dx = \frac{1}{6} \sin(3x) - \frac{1}{18} \sin(9x)$$

output `1/6*sin(3*x)-1/18*sin(9*x)`

3.109.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sin(3x) \sin(6x) dx = \frac{1}{6} \sin(3x) - \frac{1}{18} \sin(9x)$$

input `Integrate[Sin[3*x]*Sin[6*x],x]`

output `Sin[3*x]/6 - Sin[9*x]/18`

3.109.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4770}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sin(3x) \sin(6x) dx \\ \downarrow \text{3042} \\ \int \sin(3x) \sin(6x) dx \\ \downarrow \text{4770} \\ \frac{1}{6} \sin(3x) - \frac{1}{18} \sin(9x) \end{array}$$

input `Int[Sin[3*x]*Sin[6*x],x]`

output `Sin[3*x]/6 - Sin[9*x]/18`

3.109.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4770 `Int[sin[(a_.) + (b_.)*(x_)]*sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[a - c + (b - d)*x]/(2*(b - d)), x] - Simp[Sin[a + c + (b + d)*x]/(2*(b + d)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

3.109.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.53

method	result	size
derivativdivides	$\frac{2(\sin^3(3x))}{9}$	9
default	$\frac{2(\sin^3(3x))}{9}$	9
risch	$\frac{\sin(3x)}{6} - \frac{\sin(9x)}{18}$	14
parallelrisch	$\frac{\sin(3x)}{6} - \frac{\sin(9x)}{18}$	14
norman	$-\frac{2 \tan(3x) \left(\tan^2\left(\frac{3x}{2}\right) \right)}{9} + \frac{4 \left(\tan^2(3x) \right) \tan\left(\frac{3x}{2}\right)}{9} + \frac{2 \tan(3x)}{9} - \frac{4 \tan\left(\frac{3x}{2}\right)}{9}$ $\frac{\hspace{10em}}{(1+\tan^2\left(\frac{3x}{2}\right))(1+\tan^2(3x))}$	59

input `int(sin(3*x)*sin(6*x),x,method=_RETURNVERBOSE)`output `2/9*sin(3*x)^3`**3.109.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sin(3x) \sin(6x) dx = -\frac{2}{9} (\cos(3x))^2 - 1) \sin(3x)$$

input `integrate(sin(3*x)*sin(6*x),x, algorithm="fracas")`output `-2/9*(cos(3*x)^2 - 1)*sin(3*x)`**3.109.6 Sympy [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.41

$$\int \sin(3x) \sin(6x) dx = -\frac{2 \sin(3x) \cos(6x)}{9} + \frac{\sin(6x) \cos(3x)}{9}$$

input `integrate(sin(3*x)*sin(6*x),x)`output `-2*sin(3*x)*cos(6*x)/9 + sin(6*x)*cos(3*x)/9`

3.109.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sin(3x) \sin(6x) dx = -\frac{1}{18} \sin(9x) + \frac{1}{6} \sin(3x)$$

input `integrate(sin(3*x)*sin(6*x),x, algorithm="maxima")`output `-1/18*sin(9*x) + 1/6*sin(3*x)`**3.109.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.47

$$\int \sin(3x) \sin(6x) dx = \frac{2}{9} \sin(3x)^3$$

input `integrate(sin(3*x)*sin(6*x),x, algorithm="giac")`output `2/9*sin(3*x)^3`**3.109.9 Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sin(3x) \sin(6x) dx = \frac{\sin(3x)}{6} - \frac{\sin(9x)}{18}$$

input `int(sin(3*x)*sin(6*x),x)`output `sin(3*x)/6 - sin(9*x)/18`

3.110 $\int \cos^5(x) \sin(x) dx$

3.110.1 Optimal result	637
3.110.2 Mathematica [A] (verified)	637
3.110.3 Rubi [A] (verified)	638
3.110.4 Maple [A] (verified)	639
3.110.5 Fricas [A] (verification not implemented)	639
3.110.6 Sympy [A] (verification not implemented)	639
3.110.7 Maxima [A] (verification not implemented)	640
3.110.8 Giac [A] (verification not implemented)	640
3.110.9 Mupad [B] (verification not implemented)	640

3.110.1 Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \cos^5(x) \sin(x) dx = -\frac{1}{6} \cos^6(x)$$

output `-1/6*cos(x)^6`

3.110.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \cos^5(x) \sin(x) dx = -\frac{1}{6} \cos^6(x)$$

input `Integrate[Cos[x]^5*Sin[x],x]`

output `-1/6*Cos[x]^6`

3.110.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3045, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin(x) \cos^5(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin(x) \cos(x)^5 dx \\ & \quad \downarrow \text{3045} \\ & - \int \cos^5(x) d \cos(x) \\ & \quad \downarrow \text{15} \\ & -\frac{1}{6} \cos^6(x) \end{aligned}$$

input `Int[Cos[x]^5*Sin[x],x]`

output `-1/6*Cos[x]^6`

3.110.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3045 `Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[-(a*f)^(-1) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])`

3.110.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$-\frac{(\cos^6(x))}{6}$	7
default	$-\frac{(\cos^6(x))}{6}$	7
risch	$-\frac{\cos(6x)}{192} - \frac{\cos(4x)}{32} - \frac{5 \cos(2x)}{64}$	20
parallelrisch	$-\frac{7}{32} - \frac{\cos(6x)}{192} - \frac{\cos(4x)}{32} - \frac{5 \cos(2x)}{64}$	21
norman	$\frac{2(\tan^2(\frac{x}{2})) + 2(\tan^{10}(\frac{x}{2})) + \frac{20(\tan^6(\frac{x}{2}))}{3}}{(1 + \tan^2(\frac{x}{2}))^6}$	37

input `int(cos(x)^5*sin(x),x,method=_RETURNVERBOSE)`output `-1/6*cos(x)^6`**3.110.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos^5(x) \sin(x) dx = -\frac{1}{6} \cos(x)^6$$

input `integrate(cos(x)^5*sin(x),x, algorithm="fracas")`output `-1/6*cos(x)^6`**3.110.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \cos^5(x) \sin(x) dx = -\frac{\cos^6(x)}{6}$$

input `integrate(cos(x)**5*sin(x),x)`output `-cos(x)**6/6`

3.110.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos^5(x) \sin(x) dx = -\frac{1}{6} \cos(x)^6$$

input `integrate(cos(x)^5*sin(x),x, algorithm="maxima")`output `-1/6*cos(x)^6`**3.110.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cos^5(x) \sin(x) dx = -\frac{1}{6} \cos(x)^6$$

input `integrate(cos(x)^5*sin(x),x, algorithm="giac")`output `-1/6*cos(x)^6`**3.110.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.38

$$\int \cos^5(x) \sin(x) dx = \frac{\sin(x)^6}{6} - \frac{\sin(x)^4}{2} + \frac{\sin(x)^2}{2}$$

input `int(cos(x)^5*sin(x),x)`output `sin(x)^2/2 - sin(x)^4/2 + sin(x)^6/6`

3.111 $\int \cos(x) \cos(2x) \cos(3x) dx$

3.111.1 Optimal result	641
3.111.2 Mathematica [A] (verified)	641
3.111.3 Rubi [A] (verified)	642
3.111.4 Maple [A] (verified)	643
3.111.5 Fricas [A] (verification not implemented)	643
3.111.6 Sympy [B] (verification not implemented)	643
3.111.7 Maxima [A] (verification not implemented)	644
3.111.8 Giac [A] (verification not implemented)	644
3.111.9 Mupad [B] (verification not implemented)	644

3.111.1 Optimal result

Integrand size = 11, antiderivative size = 30

$$\int \cos(x) \cos(2x) \cos(3x) dx = \frac{x}{4} + \frac{1}{8} \sin(2x) + \frac{1}{16} \sin(4x) + \frac{1}{24} \sin(6x)$$

output `1/4*x+1/8*sin(2*x)+1/16*sin(4*x)+1/24*sin(6*x)`

3.111.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \cos(x) \cos(2x) \cos(3x) dx = \frac{x}{4} + \frac{1}{8} \sin(2x) + \frac{1}{16} \sin(4x) + \frac{1}{24} \sin(6x)$$

input `Integrate[Cos[x]*Cos[2*x]*Cos[3*x],x]`

output `x/4 + Sin[2*x]/8 + Sin[4*x]/16 + Sin[6*x]/24`

3.111.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 4855, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(x) \cos(2x) \cos(3x) dx$$

$$\downarrow \text{3042}$$

$$\int \cos(x) \cos(2x) \cos(3x) dx$$

$$\downarrow \text{4855}$$

$$\int \left(\frac{1}{4} \cos(2x) + \frac{1}{4} \cos(4x) + \frac{1}{4} \cos(6x) + \frac{1}{4} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{x}{4} + \frac{1}{8} \sin(2x) + \frac{1}{16} \sin(4x) + \frac{1}{24} \sin(6x)$$

input `Int[Cos[x]*Cos[2*x]*Cos[3*x],x]`

output `x/4 + Sin[2*x]/8 + Sin[4*x]/16 + Sin[6*x]/24`

3.111.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4855 `Int[(F_)[(a_.) + (b_.)*(x_)]^(p_.)*(G_)[(c_.) + (d_.)*(x_)]^(q_.)*(H_)[(e_.) + (f_.)*(x_)]^(r_.), x_Symbol] := Int[ExpandTrigReduce[ActivateTrig[F[a + b*x]^p*G[c + d*x]^q*H[e + f*x]^r], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && (EqQ[F, sin] || EqQ[F, cos]) && (EqQ[G, sin] || EqQ[G, cos]) && (EqQ[H, sin] || EqQ[H, cos]) && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]`

3.111.4 Maple [A] (verified)

Time = 1.35 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.77

method	result	size
default	$\frac{x}{4} + \frac{\sin(2x)}{8} + \frac{\sin(4x)}{16} + \frac{\sin(6x)}{24}$	23
risch	$\frac{x}{4} + \frac{\sin(2x)}{8} + \frac{\sin(4x)}{16} + \frac{\sin(6x)}{24}$	23
parallelrisch	$\frac{x}{4} + \frac{\sin(2x)}{8} + \frac{\sin(4x)}{16} + \frac{\sin(6x)}{24}$	23

input `int(cos(x)*cos(2*x)*cos(3*x),x,method=_RETURNVERBOSE)`output `1/4*x+1/8*sin(2*x)+1/16*sin(4*x)+1/24*sin(6*x)`**3.111.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.83

$$\int \cos(x) \cos(2x) \cos(3x) dx = \frac{1}{12} (16 \cos(x)^5 - 10 \cos(x)^3 + 3 \cos(x)) \sin(x) + \frac{1}{4} x$$

input `integrate(cos(x)*cos(2*x)*cos(3*x),x, algorithm="fricas")`output `1/12*(16*cos(x)^5 - 10*cos(x)^3 + 3*cos(x))*sin(x) + 1/4*x`**3.111.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 114 vs. 2(22) = 44.

Time = 0.99 (sec) , antiderivative size = 114, normalized size of antiderivative = 3.80

$$\begin{aligned} \int \cos(x) \cos(2x) \cos(3x) dx = & -\frac{x \sin(x) \sin(2x) \cos(3x)}{4} + \frac{x \sin(x) \sin(3x) \cos(2x)}{4} \\ & + \frac{x \sin(2x) \sin(3x) \cos(x)}{4} + \frac{x \cos(x) \cos(2x) \cos(3x)}{4} \\ & - \frac{\sin(x) \cos(2x) \cos(3x)}{24} - \frac{\sin(2x) \cos(x) \cos(3x)}{6} \\ & + \frac{3 \sin(3x) \cos(x) \cos(2x)}{8} \end{aligned}$$

input `integrate(cos(x)*cos(2*x)*cos(3*x),x)`

output `-x*sin(x)*sin(2*x)*cos(3*x)/4 + x*sin(x)*sin(3*x)*cos(2*x)/4 + x*sin(2*x)*sin(3*x)*cos(x)/4 + x*cos(x)*cos(2*x)*cos(3*x)/4 - sin(x)*cos(2*x)*cos(3*x)/24 - sin(2*x)*cos(x)*cos(3*x)/6 + 3*sin(3*x)*cos(x)*cos(2*x)/8`

3.111.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \cos(x) \cos(2x) \cos(3x) dx = \frac{1}{4}x + \frac{1}{24} \sin(6x) + \frac{1}{16} \sin(4x) + \frac{1}{8} \sin(2x)$$

input `integrate(cos(x)*cos(2*x)*cos(3*x),x, algorithm="maxima")`

output `1/4*x + 1/24*sin(6*x) + 1/16*sin(4*x) + 1/8*sin(2*x)`

3.111.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \cos(x) \cos(2x) \cos(3x) dx = \frac{1}{4}x + \frac{1}{24} \sin(6x) + \frac{1}{16} \sin(4x) + \frac{1}{8} \sin(2x)$$

input `integrate(cos(x)*cos(2*x)*cos(3*x),x, algorithm="giac")`

output `1/4*x + 1/24*sin(6*x) + 1/16*sin(4*x) + 1/8*sin(2*x)`

3.111.9 Mupad [B] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \cos(x) \cos(2x) \cos(3x) dx = \frac{x}{4} + \frac{\sin(2x)}{8} + \frac{\sin(4x)}{16} + \frac{\sin(6x)}{24}$$

input `int(cos(2*x)*cos(3*x)*cos(x),x)`

output `x/4 + sin(2*x)/8 + sin(4*x)/16 + sin(6*x)/24`

3.112 $\int \cos^2(x) (1 - \tan^2(x)) dx$

3.112.1 Optimal result	645
3.112.2 Mathematica [A] (verified)	645
3.112.3 Rubi [B] (verified)	646
3.112.4 Maple [A] (verified)	647
3.112.5 Fricas [A] (verification not implemented)	647
3.112.6 Sympy [A] (verification not implemented)	648
3.112.7 Maxima [B] (verification not implemented)	648
3.112.8 Giac [A] (verification not implemented)	648
3.112.9 Mupad [B] (verification not implemented)	649

3.112.1 Optimal result

Integrand size = 13, antiderivative size = 5

$$\int \cos^2(x) (1 - \tan^2(x)) dx = \cos(x) \sin(x)$$

output `cos(x)*sin(x)`

3.112.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.60

$$\int \cos^2(x) (1 - \tan^2(x)) dx = \frac{1}{2} \sin(2x)$$

input `Integrate[Cos[x]^2*(1 - Tan[x]^2),x]`

output `Sin[2*x]/2`

3.112.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 11 vs. $2(5) = 10$.

Time = 0.17 (sec) , antiderivative size = 11, normalized size of antiderivative = 2.20, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3042, 4158, 297}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^2(x) (1 - \tan^2(x)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1 - \tan(x)^2}{\sec(x)^2} dx \\ & \quad \downarrow \text{4158} \\ & \int \frac{1 - \tan^2(x)}{(\tan^2(x) + 1)^2} d \tan(x) \\ & \quad \downarrow \text{297} \\ & \frac{\tan(x)}{\tan^2(x) + 1} \end{aligned}$$

input `Int[Cos[x]^2*(1 - Tan[x]^2), x]`

output `Tan[x]/(1 + Tan[x]^2)`

3.112.3.1 Defintions of rubi rules used

rule 297 `Int[((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2), x_Symbol] :> Simp[c*x*((a + b*x^2)^(p + 1)/a), x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a*d - b*c*(2*p + 3), 0]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4158 Int[sec[(e_.) + (f_.)*(x_.)]^(m_)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_.)
)])^(n_)]^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp
p[ff/(c^(m - 1)*f) Subst[Int[(c^2 + ff^2*x^2)^(m/2 - 1)*(a + b*(ff*x)^n)^
p, x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && I
negerQ[m/2] && (IntegersQ[n, p] || IGtQ[m, 0] || IGtQ[p, 0] || EqQ[n^2, 4]
|| EqQ[n^2, 16])
```

3.112.4 Maple [A] (verified)

Time = 0.77 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
default	$\cos(x) \sin(x)$	6
risch	$\frac{\sin(2x)}{2}$	7

```
input int((1-tan(x)^2)/sec(x)^2,x,method=_RETURNVERBOSE)
```

```
output cos(x)*sin(x)
```

3.112.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \cos^2(x) (1 - \tan^2(x)) dx = \cos(x) \sin(x)$$

```
input integrate((1-tan(x)^2)/sec(x)^2,x, algorithm="fracas")
```

```
output cos(x)*sin(x)
```


3.112.6 Sympy [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.40

$$\int \cos^2(x) (1 - \tan^2(x)) dx = \frac{\tan(x)}{\sec^2(x)}$$

input `integrate((1-tan(x)**2)/sec(x)**2,x)`

output `tan(x)/sec(x)**2`

3.112.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. 2(5) = 10.

Time = 0.18 (sec) , antiderivative size = 11, normalized size of antiderivative = 2.20

$$\int \cos^2(x) (1 - \tan^2(x)) dx = \frac{\tan(x)}{\tan(x)^2 + 1}$$

input `integrate((1-tan(x)^2)/sec(x)^2,x, algorithm="maxima")`

output `tan(x)/(tan(x)^2 + 1)`

3.112.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.80

$$\int \cos^2(x) (1 - \tan^2(x)) dx = \frac{1}{\frac{1}{\tan(x)} + \tan(x)}$$

input `integrate((1-tan(x)^2)/sec(x)^2,x, algorithm="giac")`

output `1/(1/tan(x) + tan(x))`

3.112.9 Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

$$\int \cos^2(x) (1 - \tan^2(x)) dx = \frac{\sin(2x)}{2}$$

input `int(-cos(x)^2*(tan(x)^2 - 1),x)`

output `sin(2*x)/2`

3.113 $\int \csc(2x)(\cos(x) + \sin(x)) dx$

3.113.1 Optimal result	650
3.113.2 Mathematica [A] (verified)	650
3.113.3 Rubi [A] (verified)	651
3.113.4 Maple [A] (verified)	652
3.113.5 Fricas [B] (verification not implemented)	652
3.113.6 Sympy [B] (verification not implemented)	652
3.113.7 Maxima [B] (verification not implemented)	653
3.113.8 Giac [B] (verification not implemented)	653
3.113.9 Mupad [B] (verification not implemented)	654

3.113.1 Optimal result

Integrand size = 10, antiderivative size = 15

$$\int \csc(2x)(\cos(x) + \sin(x)) dx = -\frac{1}{2}\operatorname{arctanh}(\cos(x)) + \frac{1}{2}\operatorname{arctanh}(\sin(x))$$

output `-1/2*arctanh(cos(x))+1/2*arctanh(sin(x))`

3.113.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 30, normalized size of antiderivative = 2.00

$$\int \csc(2x)(\cos(x) + \sin(x)) dx = \frac{1}{2}\operatorname{arctanh}(\sin(x)) - \frac{1}{2}\log\left(\cos\left(\frac{x}{2}\right)\right) + \frac{1}{2}\log\left(\sin\left(\frac{x}{2}\right)\right)$$

input `Integrate[Csc[2*x]*(Cos[x] + Sin[x]),x]`

output `ArcTanh[Sin[x]]/2 - Log[Cos[x/2]]/2 + Log[Sin[x/2]]/2`

3.113.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3042, 4901, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \csc(2x)(\sin(x) + \cos(x)) dx \\ & \quad \downarrow \text{3042} \\ & \int \csc(2x)(\sin(x) + \cos(x)) dx \\ & \quad \downarrow \text{4901} \\ & \int (\cos(x) \csc(2x) + \sin(x) \csc(2x)) dx \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \operatorname{arctanh}(\sin(x)) - \frac{1}{2} \operatorname{arctanh}(\cos(x)) \end{aligned}$$

input `Int[Csc[2*x]*(Cos[x] + Sin[x]),x]`

output `-1/2*ArcTanh[Cos[x]] + ArcTanh[Sin[x]]/2`

3.113.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4901 `Int[u_, x_Symbol] := With[{v = ExpandTrig[u, x]}, Int[v, x] /; SumQ[v]] /; !InertTrigFreeQ[u]`

3.113.4 Maple [A] (verified)

Time = 0.57 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.20

method	result	size
parts	$-\frac{\ln(\csc(x)+\cot(x))}{2} + \frac{\ln(\sec(x)+\tan(x))}{2}$	18
default	$\frac{\ln(\sec(x)+\tan(x))}{2} + \frac{\ln(\csc(x)-\cot(x))}{2}$	20
risch	$\frac{\ln(e^{2ix}+(-1+i)e^{ix}-i)}{2} - \frac{\ln(e^{2ix}+(1-i)e^{ix}-i)}{2}$	42

input `int((cos(x)+sin(x))/sin(2*x),x,method=_RETURNVERBOSE)`

output `-1/2*ln(csc(x)+cot(x))+1/2*ln(sec(x)+tan(x))`

3.113.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 35 vs. $2(11) = 22$.

Time = 0.26 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.33

$$\int \csc(2x)(\cos(x) + \sin(x)) dx = -\frac{1}{4} \log \left(-\frac{1}{2} (\cos(x) + 1) \sin(x) + \frac{1}{2} \cos(x) + \frac{1}{2} \right) + \frac{1}{4} \log \left(-\frac{1}{2} (\cos(x) - 1) \sin(x) - \frac{1}{2} \cos(x) + \frac{1}{2} \right)$$

input `integrate((cos(x)+sin(x))/sin(2*x),x, algorithm="fracas")`

output `-1/4*log(-1/2*(cos(x) + 1)*sin(x) + 1/2*cos(x) + 1/2) + 1/4*log(-1/2*(cos(x) - 1)*sin(x) - 1/2*cos(x) + 1/2)`

3.113.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 32 vs. $2(12) = 24$.

Time = 0.52 (sec) , antiderivative size = 32, normalized size of antiderivative = 2.13

$$\int \csc(2x)(\cos(x) + \sin(x)) dx = -\frac{\log(\sin(x) - 1)}{4} + \frac{\log(\sin(x) + 1)}{4} + \frac{\log(\cos(x) - 1)}{4} - \frac{\log(\cos(x) + 1)}{4}$$

input `integrate((cos(x)+sin(x))/sin(2*x),x)`

output `-log(sin(x) - 1)/4 + log(sin(x) + 1)/4 + log(cos(x) - 1)/4 - log(cos(x) + 1)/4`

3.113.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 69 vs. $2(11) = 22$.

Time = 0.28 (sec) , antiderivative size = 69, normalized size of antiderivative = 4.60

$$\begin{aligned} \int \csc(2x)(\cos(x) + \sin(x)) dx = & -\frac{1}{4} \log(\cos(x)^2 + \sin(x)^2 + 2\cos(x) + 1) \\ & + \frac{1}{4} \log(\cos(x)^2 + \sin(x)^2 - 2\cos(x) + 1) \\ & + \frac{1}{4} \log(\cos(x)^2 + \sin(x)^2 + 2\sin(x) + 1) \\ & - \frac{1}{4} \log(\cos(x)^2 + \sin(x)^2 - 2\sin(x) + 1) \end{aligned}$$

input `integrate((cos(x)+sin(x))/sin(2*x),x, algorithm="maxima")`

output `-1/4*log(cos(x)^2 + sin(x)^2 + 2*cos(x) + 1) + 1/4*log(cos(x)^2 + sin(x)^2 - 2*cos(x) + 1) + 1/4*log(cos(x)^2 + sin(x)^2 + 2*sin(x) + 1) - 1/4*log(cos(x)^2 + sin(x)^2 - 2*sin(x) + 1)`

3.113.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(11) = 22$.

Time = 0.32 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.93

$$\begin{aligned} \int \csc(2x)(\cos(x) + \sin(x)) dx = & \frac{1}{2} \log\left(\left|\tan\left(\frac{1}{2}x\right) + 1\right|\right) \\ & - \frac{1}{2} \log\left(\left|\tan\left(\frac{1}{2}x\right) - 1\right|\right) + \frac{1}{2} \log\left(\left|\tan\left(\frac{1}{2}x\right)\right|\right) \end{aligned}$$

input `integrate((cos(x)+sin(x))/sin(2*x),x, algorithm="giac")`

output `1/2*log(abs(tan(1/2*x) + 1)) - 1/2*log(abs(tan(1/2*x) - 1)) + 1/2*log(abs(tan(1/2*x)))`

3.113.9 Mupad [B] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.60

$$\int \csc(2x)(\cos(x) + \sin(x)) dx = \frac{\ln\left(\tan\left(\frac{x}{2}\right)^2 + \tan\left(\frac{x}{2}\right)\right)}{2} - \frac{\ln\left(\tan\left(\frac{x}{2}\right) - 1\right)}{2}$$

input `int((cos(x) + sin(x))/sin(2*x),x)`

output `log(tan(x/2) + tan(x/2)^2)/2 - log(tan(x/2) - 1)/2`

3.114 $\int \sin^2(x) \tan(x) dx$

3.114.1 Optimal result	655
3.114.2 Mathematica [A] (verified)	655
3.114.3 Rubi [A] (verified)	656
3.114.4 Maple [A] (verified)	657
3.114.5 Fricas [A] (verification not implemented)	657
3.114.6 Sympy [A] (verification not implemented)	658
3.114.7 Maxima [A] (verification not implemented)	658
3.114.8 Giac [A] (verification not implemented)	658
3.114.9 Mupad [B] (verification not implemented)	659

3.114.1 Optimal result

Integrand size = 7, antiderivative size = 14

$$\int \sin^2(x) \tan(x) dx = \frac{\cos^2(x)}{2} - \log(\cos(x))$$

output `1/2*cos(x)^2-ln(cos(x))`

3.114.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \sin^2(x) \tan(x) dx = \frac{\cos^2(x)}{2} - \log(\cos(x))$$

input `Integrate[Sin[x]^2*Tan[x],x]`

output `Cos[x]^2/2 - Log[Cos[x]]`

3.114.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 3070, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^2(x) \tan(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^2 \tan(x) dx \\
 & \quad \downarrow \text{3070} \\
 & - \int (1 - \cos^2(x)) \sec(x) d \cos(x) \\
 & \quad \downarrow \text{244} \\
 & - \int (\sec(x) - \cos(x)) d \cos(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\cos^2(x)}{2} - \log(\cos(x))
 \end{aligned}$$

input `Int[Sin[x]^2*Tan[x],x]`

output `Cos[x]^2/2 - Log[Cos[x]]`

3.114.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] :> Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3070 Int[sin[(e_.) + (f_.)*(x_.)]^(m_.)*tan[(e_.) + (f_.)*(x_.)]^(n_.), x_Symbol]
:= Simp[-f^(-1) Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f
*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]
```

3.114.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
default	$-\frac{\sin^2(x)}{2} - \ln(\cos(x))$	13
risch	$ix + \frac{e^{2ix}}{8} + \frac{e^{-2ix}}{8} - \ln(e^{2ix} + 1)$	30

```
input int(sin(x)^2*tan(x),x,method=_RETURNVERBOSE)
```

```
output -1/2*sin(x)^2-ln(cos(x))
```

3.114.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \sin^2(x) \tan(x) dx = \frac{1}{2} \cos(x)^2 - \log(-\cos(x))$$

```
input integrate(sin(x)^2*tan(x),x, algorithm="fricas")
```

```
output 1/2*cos(x)^2 - log(-cos(x))
```

3.114.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \sin^2(x) \tan(x) dx = -\log(\cos(x)) + \frac{\cos^2(x)}{2}$$

input `integrate(sin(x)**2*tan(x),x)`output `-log(cos(x)) + cos(x)**2/2`**3.114.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \sin^2(x) \tan(x) dx = -\frac{1}{2} \sin(x)^2 - \frac{1}{2} \log(\sin(x)^2 - 1)$$

input `integrate(sin(x)^2*tan(x),x, algorithm="maxima")`output `-1/2*sin(x)^2 - 1/2*log(sin(x)^2 - 1)`**3.114.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.29

$$\int \sin^2(x) \tan(x) dx = -\frac{1}{2} \sin(x)^2 - \frac{1}{2} \log(-\sin(x)^2 + 1)$$

input `integrate(sin(x)^2*tan(x),x, algorithm="giac")`output `-1/2*sin(x)^2 - 1/2*log(-sin(x)^2 + 1)`

3.114.9 Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \sin^2(x) \tan(x) dx = \frac{\cos(x)^2}{2} + \frac{\ln(\tan(x)^2 + 1)}{2}$$

input `int(sin(x)^2*tan(x),x)`

output `log(tan(x)^2 + 1)/2 + cos(x)^2/2`

3.115 $\int \cos^2(x) \cot^3(x) dx$

3.115.1 Optimal result	660
3.115.2 Mathematica [A] (verified)	660
3.115.3 Rubi [A] (warning: unable to verify)	661
3.115.4 Maple [A] (verified)	662
3.115.5 Fricas [B] (verification not implemented)	663
3.115.6 Sympy [A] (verification not implemented)	663
3.115.7 Maxima [A] (verification not implemented)	663
3.115.8 Giac [A] (verification not implemented)	664
3.115.9 Mupad [B] (verification not implemented)	664

3.115.1 Optimal result

Integrand size = 9, antiderivative size = 22

$$\int \cos^2(x) \cot^3(x) dx = -\frac{1}{2} \csc^2(x) - 2 \log(\sin(x)) + \frac{\sin^2(x)}{2}$$

output `-1/2*csc(x)^2-2*ln(sin(x))+1/2*sin(x)^2`

3.115.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \cos^2(x) \cot^3(x) dx = \frac{1}{2} (-\csc^2(x) - 4 \log(\sin(x)) + \sin^2(x))$$

input `Integrate[Cos[x]^2*Cot[x]^3,x]`

output `(-Csc[x]^2 - 4*Log[Sin[x]] + Sin[x]^2)/2`

3.115.3 Rubi [A] (warning: unable to verify)

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 25, 3070, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^2(x) \cot^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\sin\left(x + \frac{\pi}{2}\right)^2 \tan\left(x + \frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{25} \\
 & -\int \sin\left(x + \frac{\pi}{2}\right)^2 \tan\left(x + \frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{3070} \\
 & \int (1 - \sin^2(x))^2 (-\csc^3(x)) d(-\sin(x)) \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \int \csc^2(x) (\sin(x) + 1)^2 d\sin^2(x) \\
 & \quad \downarrow \text{49} \\
 & \frac{1}{2} \int (\csc^2(x) + 2 \csc(x) + 1) d\sin^2(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} (\sin^2(x) + \csc(x) - 2 \log(\sin^2(x)))
 \end{aligned}$$

input `Int[Cos[x]^2*Cot[x]^3,x]`

output `(Csc[x] - 2*Log[Sin[x]^2] + Sin[x]^2)/2`

3.115.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3070 `Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[-f^(-1) Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]`

3.115.4 Maple [A] (verified)

Time = 1.09 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.32

method	result	size
default	$-\frac{\cos^6(x)}{2\sin(x)^2} - \frac{(\cos^4(x))}{2} - (\cos^2(x)) - 2\ln(\sin(x))$	29
risch	$2ix - \frac{e^{2ix}}{8} - \frac{e^{-2ix}}{8} + \frac{2e^{2ix}}{(e^{2ix}-1)^2} - 2\ln(e^{2ix} - 1)$	46

input `int(cos(x)^2*cot(x)^3,x,method=_RETURNVERBOSE)`

output `-1/2/sin(x)^2*cos(x)^6-1/2*cos(x)^4-cos(x)^2-2*ln(sin(x))`

3.115.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. $2(18) = 36$.

Time = 0.25 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.68

$$\int \cos^2(x) \cot^3(x) dx = -\frac{2 \cos(x)^4 - 3 \cos(x)^2 + 8 (\cos(x)^2 - 1) \log\left(\frac{1}{2} \sin(x)\right) - 1}{4 (\cos(x)^2 - 1)}$$

input `integrate(cos(x)^2*cot(x)^3,x, algorithm="fracas")`

output `-1/4*(2*cos(x)^4 - 3*cos(x)^2 + 8*(cos(x)^2 - 1)*log(1/2*sin(x)) - 1)/(cos(x)^2 - 1)`

3.115.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \cos^2(x) \cot^3(x) dx = -2 \log(\sin(x)) + \frac{\sin^2(x)}{2} - \frac{1}{2 \sin^2(x)}$$

input `integrate(cos(x)**2*cot(x)**3,x)`

output `-2*log(sin(x)) + sin(x)**2/2 - 1/(2*sin(x)**2)`

3.115.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \cos^2(x) \cot^3(x) dx = \frac{1}{2} \sin(x)^2 - \frac{1}{2 \sin(x)^2} - \log(\sin(x)^2)$$

input `integrate(cos(x)^2*cot(x)^3,x, algorithm="maxima")`

output `1/2*sin(x)^2 - 1/2/sin(x)^2 - log(sin(x)^2)`

3.115.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.27

$$\int \cos^2(x) \cot^3(x) dx = -\frac{1}{2} \cos(x)^2 + \frac{1}{2(\cos(x)^2 - 1)} - \log(-\cos(x)^2 + 1)$$

input `integrate(cos(x)^2*cot(x)^3,x, algorithm="giac")`output `-1/2*cos(x)^2 + 1/2/(cos(x)^2 - 1) - log(-cos(x)^2 + 1)`**3.115.9 Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.45

$$\int \cos^2(x) \cot^3(x) dx = \ln(\tan(x)^2 + 1) - 2 \ln(\tan(x)) - \frac{\tan(x)^2 + \frac{1}{2}}{\tan(x)^4 + \tan(x)^2}$$

input `int(cos(x)^2*cot(x)^3,x)`output `log(tan(x)^2 + 1) - 2*log(tan(x)) - (tan(x)^2 + 1/2)/(tan(x)^2 + tan(x)^4)`

3.116 $\int \sec^3(x) \tan(x) dx$

3.116.1 Optimal result	665
3.116.2 Mathematica [A] (verified)	665
3.116.3 Rubi [A] (verified)	666
3.116.4 Maple [A] (verified)	667
3.116.5 Fricas [A] (verification not implemented)	667
3.116.6 Sympy [A] (verification not implemented)	667
3.116.7 Maxima [A] (verification not implemented)	668
3.116.8 Giac [A] (verification not implemented)	668
3.116.9 Mupad [B] (verification not implemented)	668

3.116.1 Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \sec^3(x) \tan(x) dx = \frac{\sec^3(x)}{3}$$

output `1/3*sec(x)^3`

3.116.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \sec^3(x) \tan(x) dx = \frac{\sec^3(x)}{3}$$

input `Integrate[Sec[x]^3*Tan[x],x]`

output `Sec[x]^3/3`

3.116.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3086, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan(x) \sec^3(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(x) \sec(x)^3 dx \\ & \quad \downarrow \text{3086} \\ & \int \sec^2(x) d \sec(x) \\ & \quad \downarrow \text{15} \\ & \frac{\sec^3(x)}{3} \end{aligned}$$

input `Int[Sec[x]^3*Tan[x],x]`

output `Sec[x]^3/3`

3.116.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

3.116.4 Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{(\sec^3(x))}{3}$	7
default	$\frac{(\sec^3(x))}{3}$	7
risch	$\frac{8 e^{3ix}}{3(e^{2ix}+1)^3}$	17

input `int(sec(x)^3*tan(x),x,method=_RETURNVERBOSE)`

output `1/3*sec(x)^3`

3.116.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^3(x) \tan(x) dx = \frac{1}{3 \cos(x)^3}$$

input `integrate(sec(x)^3*tan(x),x, algorithm="fricas")`

output `1/3/cos(x)^3`

3.116.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \sec^3(x) \tan(x) dx = \frac{1}{3 \cos^3(x)}$$

input `integrate(sec(x)**3*tan(x),x)`

output `1/(3*cos(x)**3)`

3.116.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^3(x) \tan(x) dx = \frac{1}{3 \cos(x)^3}$$

input `integrate(sec(x)^3*tan(x),x, algorithm="maxima")`output `1/3/cos(x)^3`**3.116.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^3(x) \tan(x) dx = \frac{1}{3 \cos(x)^3}$$

input `integrate(sec(x)^3*tan(x),x, algorithm="giac")`output `1/3/cos(x)^3`**3.116.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sec^3(x) \tan(x) dx = \frac{1}{3 \cos(x)^3}$$

input `int(tan(x)/cos(x)^3,x)`output `1/(3*cos(x)^3)`

3.117 $\int \sec^3(x) \tan^3(x) dx$

3.117.1 Optimal result	669
3.117.2 Mathematica [A] (verified)	669
3.117.3 Rubi [A] (verified)	670
3.117.4 Maple [A] (verified)	671
3.117.5 Fricas [A] (verification not implemented)	672
3.117.6 Sympy [A] (verification not implemented)	672
3.117.7 Maxima [A] (verification not implemented)	672
3.117.8 Giac [A] (verification not implemented)	673
3.117.9 Mupad [B] (verification not implemented)	673

3.117.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \sec^3(x) \tan^3(x) dx = -\frac{1}{3} \sec^3(x) + \frac{\sec^5(x)}{5}$$

output `-1/3*sec(x)^3+1/5*sec(x)^5`

3.117.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sec^3(x) \tan^3(x) dx = -\frac{1}{3} \sec^3(x) + \frac{\sec^5(x)}{5}$$

input `Integrate[Sec[x]^3*Tan[x]^3,x]`

output `-1/3*Sec[x]^3 + Sec[x]^5/5`

3.117.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 3086, 25, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^3(x) \sec^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^3 \sec(x)^3 dx \\
 & \quad \downarrow \text{3086} \\
 & \int -\sec^2(x) (1 - \sec^2(x)) d\sec(x) \\
 & \quad \downarrow \text{25} \\
 & - \int \sec^2(x) (1 - \sec^2(x)) d\sec(x) \\
 & \quad \downarrow \text{244} \\
 & - \int (\sec^2(x) - \sec^4(x)) d\sec(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sec^5(x)}{5} - \frac{\sec^3(x)}{3}
 \end{aligned}$$

input `Int[Sec[x]^3*Tan[x]^3,x]`

output `-1/3*Sec[x]^3 + Sec[x]^5/5`

3.117.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 244 `Int[((c_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

3.117.4 Maple [A] (verified)

Time = 1.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$-\frac{(\sec^3(x))}{3} + \frac{(\sec^5(x))}{5}$	14
default	$-\frac{(\sec^3(x))}{3} + \frac{(\sec^5(x))}{5}$	14
risch	$-\frac{8(5e^{7ix} - 2e^{5ix} + 5e^{3ix})}{15(e^{2ix} + 1)^5}$	34

input `int(sec(x)^3*tan(x)^3,x,method=_RETURNVERBOSE)`

output `-1/3*sec(x)^3+1/5*sec(x)^5`

3.117.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^3(x) \tan^3(x) dx = -\frac{5 \cos(x)^2 - 3}{15 \cos(x)^5}$$

input `integrate(sec(x)^3*tan(x)^3,x, algorithm="fricas")`output `-1/15*(5*cos(x)^2 - 3)/cos(x)^5`**3.117.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^3(x) \tan^3(x) dx = \frac{3 - 5 \cos^2(x)}{15 \cos^5(x)}$$

input `integrate(sec(x)**3*tan(x)**3,x)`output `(3 - 5*cos(x)**2)/(15*cos(x)**5)`**3.117.7 Maxima [A] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^3(x) \tan^3(x) dx = -\frac{5 \cos(x)^2 - 3}{15 \cos(x)^5}$$

input `integrate(sec(x)^3*tan(x)^3,x, algorithm="maxima")`output `-1/15*(5*cos(x)^2 - 3)/cos(x)^5`

3.117.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^3(x) \tan^3(x) dx = -\frac{5 \cos(x)^2 - 3}{15 \cos(x)^5}$$

input `integrate(sec(x)^3*tan(x)^3,x, algorithm="giac")`output `-1/15*(5*cos(x)^2 - 3)/cos(x)^5`**3.117.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sec^3(x) \tan^3(x) dx = \frac{1}{5 \cos(x)^5} - \frac{1}{3 \cos(x)^3}$$

input `int(tan(x)^3/cos(x)^3,x)`output `1/(5*cos(x)^5) - 1/(3*cos(x)^3)`

3.118 $\int \frac{\sqrt{9-x^2}}{x^2} dx$

3.118.1 Optimal result	674
3.118.2 Mathematica [A] (verified)	674
3.118.3 Rubi [A] (verified)	675
3.118.4 Maple [A] (verified)	676
3.118.5 Fricas [A] (verification not implemented)	676
3.118.6 Sympy [A] (verification not implemented)	677
3.118.7 Maxima [A] (verification not implemented)	677
3.118.8 Giac [A] (verification not implemented)	677
3.118.9 Mupad [B] (verification not implemented)	678

3.118.1 Optimal result

Integrand size = 15, antiderivative size = 25

$$\int \frac{\sqrt{9-x^2}}{x^2} dx = -\frac{\sqrt{9-x^2}}{x} - \arcsin\left(\frac{x}{3}\right)$$

output `-arcsin(1/3*x)-(-x^2+9)^(1/2)/x`

3.118.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.48

$$\int \frac{\sqrt{9-x^2}}{x^2} dx = -\frac{\sqrt{9-x^2}}{x} + 2 \arctan\left(\frac{\sqrt{9-x^2}}{3+x}\right)$$

input `Integrate[Sqrt[9 - x^2]/x^2,x]`

output `-(Sqrt[9 - x^2]/x) + 2*ArcTan[Sqrt[9 - x^2]/(3 + x)]`

3.118.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {247, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{9-x^2}}{x^2} dx$$

↓ 247

$$-\int \frac{1}{\sqrt{9-x^2}} dx - \frac{\sqrt{9-x^2}}{x}$$

↓ 223

$$-\arcsin\left(\frac{x}{3}\right) - \frac{\sqrt{9-x^2}}{x}$$

input `Int[Sqrt[9 - x^2]/x^2,x]`

output `-(Sqrt[9 - x^2]/x) - ArcSin[x/3]`

3.118.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 247 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^p/(c*(m + 1))), x] - Simp[2*b*(p/(c^2*(m + 1))) Int[(c*x)^(m + 2)*(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && GtQ[p, 0] && LtQ[m, -1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

3.118.4 Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

method	result	size
risch	$\frac{x^2-9}{x\sqrt{-x^2+9}} - \arcsin\left(\frac{x}{3}\right)$	26
pseudoelliptic	$\frac{\arctan\left(\frac{\sqrt{-x^2+9}}{x}\right)x - \sqrt{-x^2+9}}{x}$	33
default	$-\frac{(-x^2+9)^{\frac{3}{2}}}{9x} - \frac{x\sqrt{-x^2+9}}{9} - \arcsin\left(\frac{x}{3}\right)$	34
meijerg	$i \frac{\left(-\frac{12i\sqrt{\pi}\sqrt{-\frac{x^2}{9}+1}}{x} - 4i\sqrt{\pi} \arcsin\left(\frac{x}{3}\right)\right)}{4\sqrt{\pi}}$	36
trager	$-\frac{\sqrt{-x^2+9}}{x} + \text{RootOf}(-Z^2+1) \ln(\text{RootOf}(-Z^2+1)x + \sqrt{-x^2+9})$	42

input `int((-x^2+9)^(1/2)/x^2,x,method=_RETURNVERBOSE)`output `(x^2-9)/x/(-x^2+9)^(1/2)-arcsin(1/3*x)`**3.118.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.40

$$\int \frac{\sqrt{9-x^2}}{x^2} dx = \frac{2x \arctan\left(\frac{\sqrt{-x^2+9}-3}{x}\right) - \sqrt{-x^2+9}}{x}$$

input `integrate((-x^2+9)^(1/2)/x^2,x, algorithm="fracas")`output `(2*x*arctan((sqrt(-x^2 + 9) - 3)/x) - sqrt(-x^2 + 9))/x`

3.118.6 Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.60

$$\int \frac{\sqrt{9-x^2}}{x^2} dx = -\operatorname{asin}\left(\frac{x}{3}\right) - \frac{\sqrt{9-x^2}}{x}$$

input `integrate((-x**2+9)**(1/2)/x**2,x)`output `-asin(x/3) - sqrt(9 - x**2)/x`**3.118.7 Maxima [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int \frac{\sqrt{9-x^2}}{x^2} dx = -\frac{\sqrt{-x^2+9}}{x} - \arcsin\left(\frac{1}{3}x\right)$$

input `integrate((-x^2+9)^(1/2)/x^2,x, algorithm="maxima")`output `-sqrt(-x^2 + 9)/x - arcsin(1/3*x)`**3.118.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.56

$$\int \frac{\sqrt{9-x^2}}{x^2} dx = \frac{x}{2(\sqrt{-x^2+9}-3)} - \frac{\sqrt{-x^2+9}-3}{2x} - \arcsin\left(\frac{1}{3}x\right)$$

input `integrate((-x^2+9)^(1/2)/x^2,x, algorithm="giac")`output `1/2*x/(sqrt(-x^2 + 9) - 3) - 1/2*(sqrt(-x^2 + 9) - 3)/x - arcsin(1/3*x)`

3.118.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int \frac{\sqrt{9-x^2}}{x^2} dx = -\operatorname{asin}\left(\frac{x}{3}\right) - \frac{\sqrt{9-x^2}}{x}$$

input `int((9 - x^2)^(1/2)/x^2,x)`

output `- asin(x/3) - (9 - x^2)^(1/2)/x`

3.119 $\int \frac{1}{x^2\sqrt{4+x^2}} dx$

3.119.1 Optimal result	679
3.119.2 Mathematica [A] (verified)	679
3.119.3 Rubi [A] (verified)	680
3.119.4 Maple [A] (verified)	680
3.119.5 Fricas [A] (verification not implemented)	681
3.119.6 Sympy [A] (verification not implemented)	681
3.119.7 Maxima [A] (verification not implemented)	681
3.119.8 Giac [A] (verification not implemented)	682
3.119.9 Mupad [B] (verification not implemented)	682

3.119.1 Optimal result

Integrand size = 13, antiderivative size = 16

$$\int \frac{1}{x^2\sqrt{4+x^2}} dx = -\frac{\sqrt{4+x^2}}{4x}$$

output `-1/4*(x^2+4)^(1/2)/x`

3.119.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2\sqrt{4+x^2}} dx = -\frac{\sqrt{4+x^2}}{4x}$$

input `Integrate[1/(x^2*Sqrt[4 + x^2]),x]`

output `-1/4*Sqrt[4 + x^2]/x`

3.119.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 \sqrt{x^2 + 4}} dx$$

$$\downarrow \text{242}$$

$$-\frac{\sqrt{x^2 + 4}}{4x}$$

input `Int[1/(x^2*Sqrt[4 + x^2]),x]`

output `-1/4*Sqrt[4 + x^2]/x`

3.119.3.1 Defintions of rubi rules used

rule 242 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

3.119.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

method	result	size
gospers	$-\frac{\sqrt{x^2+4}}{4x}$	13
default	$-\frac{\sqrt{x^2+4}}{4x}$	13
trager	$-\frac{\sqrt{x^2+4}}{4x}$	13
risch	$-\frac{\sqrt{x^2+4}}{4x}$	13
pseudoelliptic	$-\frac{\sqrt{x^2+4}}{4x}$	13
meijerg	$-\frac{\sqrt{1+\frac{x^2}{4}}}{2x}$	15

input `int(1/x^2/(x^2+4)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/4*(x^2+4)^(1/2)/x`

3.119.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{x^2\sqrt{4+x^2}} dx = -\frac{x + \sqrt{x^2+4}}{4x}$$

input `integrate(1/x^2/(x^2+4)^(1/2),x, algorithm="fricas")`

output `-1/4*(x + sqrt(x^2 + 4))/x`

3.119.6 Sympy [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{x^2\sqrt{4+x^2}} dx = -\frac{\sqrt{1+\frac{4}{x^2}}}{4}$$

input `integrate(1/x**2/(x**2+4)**(1/2),x)`

output `-sqrt(1 + 4/x**2)/4`

3.119.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{x^2\sqrt{4+x^2}} dx = -\frac{\sqrt{x^2+4}}{4x}$$

input `integrate(1/x^2/(x^2+4)^(1/2),x, algorithm="maxima")`

output `-1/4*sqrt(x^2 + 4)/x`

3.119.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.19

$$\int \frac{1}{x^2\sqrt{4+x^2}} dx = \frac{2}{(x - \sqrt{x^2+4})^2 - 4}$$

input `integrate(1/x^2/(x^2+4)^(1/2),x, algorithm="giac")`output `2/((x - sqrt(x^2 + 4))^2 - 4)`**3.119.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{x^2\sqrt{4+x^2}} dx = -\frac{\sqrt{x^2+4}}{4x}$$

input `int(1/(x^2*(x^2 + 4)^(1/2)),x)`output `-(x^2 + 4)^(1/2)/(4*x)`

3.120 $\int \frac{x}{\sqrt{4+x^2}} dx$

3.120.1 Optimal result	683
3.120.2 Mathematica [A] (verified)	683
3.120.3 Rubi [A] (verified)	684
3.120.4 Maple [A] (verified)	684
3.120.5 Fricas [A] (verification not implemented)	685
3.120.6 Sympy [A] (verification not implemented)	685
3.120.7 Maxima [A] (verification not implemented)	685
3.120.8 Giac [A] (verification not implemented)	686
3.120.9 Mupad [B] (verification not implemented)	686

3.120.1 Optimal result

Integrand size = 11, antiderivative size = 9

$$\int \frac{x}{\sqrt{4+x^2}} dx = \sqrt{4+x^2}$$

output $(x^2+4)^{(1/2)}$

3.120.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{x}{\sqrt{4+x^2}} dx = \sqrt{4+x^2}$$

input `Integrate[x/Sqrt[4 + x^2], x]`

output `Sqrt[4 + x^2]`

3.120.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt{x^2 + 4}} dx$$

↓ 241

$$\sqrt{x^2 + 4}$$

input `Int[x/Sqrt[4 + x^2],x]`

output `Sqrt[4 + x^2]`

3.120.3.1 Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

3.120.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

method	result	size
gospers	$\sqrt{x^2 + 4}$	8
derivativedivides	$\sqrt{x^2 + 4}$	8
default	$\sqrt{x^2 + 4}$	8
trager	$\sqrt{x^2 + 4}$	8
risch	$\sqrt{x^2 + 4}$	8
pseudoelliptic	$\sqrt{x^2 + 4}$	8
meijerg	$\frac{-2\sqrt{\pi}+2\sqrt{\pi}\sqrt{1+\frac{x^2}{4}}}{\sqrt{\pi}}$	25

input `int(x/(x^2+4)^(1/2),x,method=_RETURNVERBOSE)`

output `(x^2+4)^(1/2)`

3.120.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{x}{\sqrt{4+x^2}} dx = \sqrt{x^2+4}$$

input `integrate(x/(x^2+4)^(1/2),x, algorithm="fricas")`

output `sqrt(x^2 + 4)`

3.120.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{x}{\sqrt{4+x^2}} dx = \sqrt{x^2+4}$$

input `integrate(x/(x**2+4)**(1/2),x)`

output `sqrt(x**2 + 4)`

3.120.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{x}{\sqrt{4+x^2}} dx = \sqrt{x^2+4}$$

input `integrate(x/(x^2+4)^(1/2),x, algorithm="maxima")`

output `sqrt(x^2 + 4)`

3.120.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{x}{\sqrt{4+x^2}} dx = \sqrt{x^2+4}$$

input `integrate(x/(x^2+4)^(1/2),x, algorithm="giac")`

output `sqrt(x^2 + 4)`

3.120.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{x}{\sqrt{4+x^2}} dx = \sqrt{x^2+4}$$

input `int(x/(x^2 + 4)^(1/2),x)`

output `(x^2 + 4)^(1/2)`

3.121 $\int \frac{1}{\sqrt{-a^2+x^2}} dx$

3.121.1 Optimal result	687
3.121.2 Mathematica [B] (verified)	687
3.121.3 Rubi [A] (verified)	688
3.121.4 Maple [A] (verified)	689
3.121.5 Fricas [A] (verification not implemented)	689
3.121.6 Sympy [C] (verification not implemented)	689
3.121.7 Maxima [A] (verification not implemented)	690
3.121.8 Giac [B] (verification not implemented)	690
3.121.9 Mupad [B] (verification not implemented)	690

3.121.1 Optimal result

Integrand size = 13, antiderivative size = 16

$$\int \frac{1}{\sqrt{-a^2+x^2}} dx = \operatorname{arctanh}\left(\frac{x}{\sqrt{-a^2+x^2}}\right)$$

output `arctanh(x/(-a^2+x^2)^(1/2))`

3.121.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 46 vs. 2(16) = 32.

Time = 0.00 (sec) , antiderivative size = 46, normalized size of antiderivative = 2.88

$$\int \frac{1}{\sqrt{-a^2+x^2}} dx = -\frac{1}{2} \log\left(1 - \frac{x}{\sqrt{-a^2+x^2}}\right) + \frac{1}{2} \log\left(1 + \frac{x}{\sqrt{-a^2+x^2}}\right)$$

input `Integrate[1/Sqrt[-a^2 + x^2],x]`

output `-1/2*Log[1 - x/Sqrt[-a^2 + x^2]] + Log[1 + x/Sqrt[-a^2 + x^2]]/2`

3.121.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx$$

↓ 224

$$\int \frac{1}{1 - \frac{x^2}{x^2 - a^2}} d \frac{x}{\sqrt{x^2 - a^2}}$$

↓ 219

$$\operatorname{arctanh}\left(\frac{x}{\sqrt{x^2 - a^2}}\right)$$

input `Int[1/Sqrt[-a^2 + x^2],x]`

output `ArcTanh[x/Sqrt[-a^2 + x^2]]`

3.121.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

3.121.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result	size
default	$\ln(x + \sqrt{-a^2 + x^2})$	15
pseudoelliptic	$\operatorname{arctanh}\left(\frac{\sqrt{-a^2+x^2}}{x}\right)$	17

input `int(1/(-a^2+x^2)^(1/2),x,method=_RETURNVERBOSE)`

output `ln(x+(-a^2+x^2)^(1/2))`

3.121.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{1}{\sqrt{-a^2 + x^2}} dx = -\log(-x + \sqrt{-a^2 + x^2})$$

input `integrate(1/(-a^2+x^2)^(1/2),x, algorithm="fricas")`

output `-log(-x + sqrt(-a^2 + x^2))`

3.121.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.19

$$\int \frac{1}{\sqrt{-a^2 + x^2}} dx = \begin{cases} \operatorname{acosh}\left(\frac{x}{a}\right) & \text{for } \left|\frac{x^2}{a^2}\right| > 1 \\ -i \operatorname{asin}\left(\frac{x}{a}\right) & \text{otherwise} \end{cases}$$

input `integrate(1/(-a**2+x**2)**(1/2),x)`

output `Piecewise((acosh(x/a), Abs(x**2/a**2) > 1), (-I*asin(x/a), True))`

3.121.7 Maxima [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{1}{\sqrt{-a^2 + x^2}} dx = \log \left(2x + 2\sqrt{-a^2 + x^2} \right)$$

input `integrate(1/(-a^2+x^2)^(1/2),x, algorithm="maxima")`

output `log(2*x + 2*sqrt(-a^2 + x^2))`

3.121.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. 2(14) = 28.

Time = 0.28 (sec) , antiderivative size = 37, normalized size of antiderivative = 2.31

$$\int \frac{1}{\sqrt{-a^2 + x^2}} dx = \frac{1}{2} a^2 \log \left(\left| -x + \sqrt{-a^2 + x^2} \right| \right) + \frac{1}{2} \sqrt{-a^2 + x^2} x$$

input `integrate(1/(-a^2+x^2)^(1/2),x, algorithm="giac")`

output `1/2*a^2*log(abs(-x + sqrt(-a^2 + x^2))) + 1/2*sqrt(-a^2 + x^2)*x`

3.121.9 Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{\sqrt{-a^2 + x^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right)$$

input `int(1/(x^2 - a^2)^(1/2),x)`

output `log(x + (x^2 - a^2)^(1/2))`

3.122 $\int \frac{x^3}{(9+4x^2)^{3/2}} dx$

3.122.1 Optimal result	691
3.122.2 Mathematica [A] (verified)	691
3.122.3 Rubi [A] (verified)	692
3.122.4 Maple [A] (verified)	693
3.122.5 Fricas [A] (verification not implemented)	693
3.122.6 Sympy [A] (verification not implemented)	694
3.122.7 Maxima [A] (verification not implemented)	694
3.122.8 Giac [A] (verification not implemented)	694
3.122.9 Mupad [B] (verification not implemented)	695

3.122.1 Optimal result

Integrand size = 15, antiderivative size = 31

$$\int \frac{x^3}{(9 + 4x^2)^{3/2}} dx = \frac{9}{16\sqrt{9 + 4x^2}} + \frac{1}{16}\sqrt{9 + 4x^2}$$

output `9/16/(4*x^2+9)^(1/2)+1/16*(4*x^2+9)^(1/2)`

3.122.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.71

$$\int \frac{x^3}{(9 + 4x^2)^{3/2}} dx = \frac{9 + 2x^2}{8\sqrt{9 + 4x^2}}$$

input `Integrate[x^3/(9 + 4*x^2)^(3/2),x]`

output `(9 + 2*x^2)/(8*Sqrt[9 + 4*x^2])`

3.122.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.13, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^3}{(4x^2 + 9)^{3/2}} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int \frac{x^2}{(4x^2 + 9)^{3/2}} dx^2 \\ & \quad \downarrow \text{53} \\ & \frac{1}{2} \int \left(\frac{1}{4\sqrt{4x^2 + 9}} - \frac{9}{4(4x^2 + 9)^{3/2}} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \left(\frac{1}{8} \sqrt{4x^2 + 9} + \frac{9}{8\sqrt{4x^2 + 9}} \right) \end{aligned}$$

input `Int[x^3/(9 + 4*x^2)^(3/2),x]`

output `(9/(8*Sqrt[9 + 4*x^2]) + Sqrt[9 + 4*x^2]/8)/2`

3.122.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.122.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.61

method	result	size
gospers	$\frac{2x^2+9}{8\sqrt{4x^2+9}}$	19
trager	$\frac{2x^2+9}{8\sqrt{4x^2+9}}$	19
risch	$\frac{2x^2+9}{8\sqrt{4x^2+9}}$	19
pseudoelliptic	$\frac{2x^2+9}{8\sqrt{4x^2+9}}$	19
default	$\frac{x^2}{4\sqrt{4x^2+9}} + \frac{9}{8\sqrt{4x^2+9}}$	27
meijerg	$-\frac{3\sqrt{\pi}}{8} + \frac{3\sqrt{\pi}\left(\frac{16x^2}{9}+8\right)}{64\sqrt{1+\frac{4x^2}{9}}}$	33

input `int(x^3/(4*x^2+9)^(3/2),x,method=_RETURNVERBOSE)`

output `1/8*(2*x^2+9)/(4*x^2+9)^(1/2)`

3.122.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.58

$$\int \frac{x^3}{(9+4x^2)^{3/2}} dx = \frac{2x^2+9}{8\sqrt{4x^2+9}}$$

input `integrate(x^3/(4*x^2+9)^(3/2),x, algorithm="fricas")`

output `1/8*(2*x^2 + 9)/sqrt(4*x^2 + 9)`

3.122.6 Sympy [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

$$\int \frac{x^3}{(9+4x^2)^{3/2}} dx = \frac{x^2}{4\sqrt{4x^2+9}} + \frac{9}{8\sqrt{4x^2+9}}$$

input `integrate(x**3/(4*x**2+9)**(3/2),x)`output `x**2/(4*sqrt(4*x**2 + 9)) + 9/(8*sqrt(4*x**2 + 9))`**3.122.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{x^3}{(9+4x^2)^{3/2}} dx = \frac{x^2}{4\sqrt{4x^2+9}} + \frac{9}{8\sqrt{4x^2+9}}$$

input `integrate(x^3/(4*x^2+9)^(3/2),x, algorithm="maxima")`output `1/4*x^2/sqrt(4*x^2 + 9) + 9/8/sqrt(4*x^2 + 9)`**3.122.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int \frac{x^3}{(9+4x^2)^{3/2}} dx = \frac{1}{16}\sqrt{4x^2+9} + \frac{9}{16\sqrt{4x^2+9}}$$

input `integrate(x^3/(4*x^2+9)^(3/2),x, algorithm="giac")`output `1/16*sqrt(4*x^2 + 9) + 9/16/sqrt(4*x^2 + 9)`

3.122.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.77

$$\int \frac{x^3}{(9 + 4x^2)^{3/2}} dx = \frac{\sqrt{x^2 + \frac{9}{4}} (2x^2 + 9)}{4(4x^2 + 9)}$$

input `int(x^3/(4*x^2 + 9)^(3/2),x)`output `((x^2 + 9/4)^(1/2)*(2*x^2 + 9))/(4*(4*x^2 + 9))`

3.123 $\int \frac{x}{\sqrt{3-2x-x^2}} dx$

3.123.1 Optimal result	696
3.123.2 Mathematica [A] (verified)	696
3.123.3 Rubi [A] (verified)	697
3.123.4 Maple [A] (verified)	698
3.123.5 Fricas [A] (verification not implemented)	698
3.123.6 Sympy [A] (verification not implemented)	699
3.123.7 Maxima [A] (verification not implemented)	699
3.123.8 Giac [A] (verification not implemented)	699
3.123.9 Mupad [B] (verification not implemented)	700

3.123.1 Optimal result

Integrand size = 16, antiderivative size = 27

$$\int \frac{x}{\sqrt{3-2x-x^2}} dx = -\sqrt{3-2x-x^2} + \arcsin\left(\frac{1}{2}(-1-x)\right)$$

output `-arcsin(1/2+1/2*x)-(-x^2-2*x+3)^(1/2)`

3.123.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.48

$$\int \frac{x}{\sqrt{3-2x-x^2}} dx = -\sqrt{3-2x-x^2} + 2 \arctan\left(\frac{\sqrt{3-2x-x^2}}{3+x}\right)$$

input `Integrate[x/Sqrt[3 - 2*x - x^2], x]`

output `-Sqrt[3 - 2*x - x^2] + 2*ArcTan[Sqrt[3 - 2*x - x^2]/(3 + x)]`

3.123.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {1160, 1090, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x}{\sqrt{-x^2 - 2x + 3}} dx \\ & \quad \downarrow \text{1160} \\ & - \int \frac{1}{\sqrt{-x^2 - 2x + 3}} dx - \sqrt{-x^2 - 2x + 3} \\ & \quad \downarrow \text{1090} \\ & \frac{1}{4} \int \frac{1}{\sqrt{1 - \frac{1}{16}(-2x - 2)^2}} d(-2x - 2) - \sqrt{-x^2 - 2x + 3} \\ & \quad \downarrow \text{223} \\ & \arcsin\left(\frac{1}{4}(-2x - 2)\right) - \sqrt{-x^2 - 2x + 3} \end{aligned}$$

input `Int[x/Sqrt[3 - 2*x - x^2],x]`

output `-Sqrt[3 - 2*x - x^2] + ArcSin[(-2 - 2*x)/4]`

3.123.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 1090 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

```
rule 1160 Int[((d_.) + (e_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol
] := Simp[e*((a + b*x + c*x^2)^(p + 1)/(2*c*(p + 1))), x] + Simp[(2*c*d - b
*e)/(2*c) Int[(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x]
&& NeQ[p, -1]
```

3.123.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

method	result
default	$-\arcsin\left(\frac{1}{2} + \frac{x}{2}\right) - \sqrt{-x^2 - 2x + 3}$
risch	$\frac{x^2+2x-3}{\sqrt{-x^2-2x+3}} - \arcsin\left(\frac{1}{2} + \frac{x}{2}\right)$
trager	$-\sqrt{-x^2 - 2x + 3} + \text{RootOf}(_Z^2 + 1) \ln(\text{RootOf}(_Z^2 + 1)x + \text{RootOf}(_Z^2 + 1)) + \sqrt{-x^2 - 2x + 3}$

```
input int(x/(-x^2-2*x+3)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -arcsin(1/2+1/2*x)-(-x^2-2*x+3)^(1/2)
```

3.123.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.56

$$\int \frac{x}{\sqrt{3-2x-x^2}} dx = -\sqrt{-x^2 - 2x + 3} + \arctan\left(\frac{\sqrt{-x^2 - 2x + 3}(x + 1)}{x^2 + 2x - 3}\right)$$

```
input integrate(x/(-x^2-2*x+3)^(1/2),x, algorithm="fricas")
```

```
output -sqrt(-x^2 - 2*x + 3) + arctan(sqrt(-x^2 - 2*x + 3)*(x + 1)/(x^2 + 2*x - 3
))
```

3.123.6 Sympy [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

$$\int \frac{x}{\sqrt{3-2x-x^2}} dx = -\sqrt{-x^2-2x+3} - \operatorname{asin}\left(\frac{x}{2} + \frac{1}{2}\right)$$

input `integrate(x/(-x**2-2*x+3)**(1/2),x)`output `-sqrt(-x**2 - 2*x + 3) - asin(x/2 + 1/2)`**3.123.7 Maxima [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int \frac{x}{\sqrt{3-2x-x^2}} dx = -\sqrt{-x^2-2x+3} + \arcsin\left(-\frac{1}{2}x - \frac{1}{2}\right)$$

input `integrate(x/(-x^2-2*x+3)^(1/2),x, algorithm="maxima")`output `-sqrt(-x^2 - 2*x + 3) + arcsin(-1/2*x - 1/2)`**3.123.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{3-2x-x^2}} dx = -\sqrt{-x^2-2x+3} - \arcsin\left(\frac{1}{2}x + \frac{1}{2}\right)$$

input `integrate(x/(-x^2-2*x+3)^(1/2),x, algorithm="giac")`output `-sqrt(-x^2 - 2*x + 3) - arcsin(1/2*x + 1/2)`

3.123.9 Mupad [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.41

$$\int \frac{x}{\sqrt{3-2x-x^2}} dx = -\sqrt{-x^2-2x+3} + \ln\left(x + \sqrt{-x^2-2x+3}\right)$$

input `int(x/(3 - x^2 - 2*x)^(1/2),x)`

output `log(x*1i + (3 - x^2 - 2*x)^(1/2) + 1i)*1i - (3 - x^2 - 2*x)^(1/2)`

3.124 $\int \frac{1}{x^2\sqrt{1-x^2}} dx$

3.124.1 Optimal result	701
3.124.2 Mathematica [A] (verified)	701
3.124.3 Rubi [A] (verified)	702
3.124.4 Maple [A] (verified)	702
3.124.5 Fricas [A] (verification not implemented)	703
3.124.6 Sympy [C] (verification not implemented)	703
3.124.7 Maxima [A] (verification not implemented)	704
3.124.8 Giac [B] (verification not implemented)	704
3.124.9 Mupad [B] (verification not implemented)	704

3.124.1 Optimal result

Integrand size = 15, antiderivative size = 16

$$\int \frac{1}{x^2\sqrt{1-x^2}} dx = -\frac{\sqrt{1-x^2}}{x}$$

output `-(-x^2+1)^(1/2)/x`

3.124.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2\sqrt{1-x^2}} dx = -\frac{\sqrt{1-x^2}}{x}$$

input `Integrate[1/(x^2*Sqrt[1 - x^2]),x]`

output `-(Sqrt[1 - x^2]/x)`

3.124.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$, Rules used = {242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 \sqrt{1-x^2}} dx$$

↓ 242

$$-\frac{\sqrt{1-x^2}}{x}$$

input `Int[1/(x^2*Sqrt[1 - x^2]),x]`

output `-(Sqrt[1 - x^2]/x)`

3.124.3.1 Defintions of rubi rules used

rule 242 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

3.124.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result	size
default	$-\frac{\sqrt{-x^2+1}}{x}$	15
trager	$-\frac{\sqrt{-x^2+1}}{x}$	15
meijerg	$-\frac{\sqrt{-x^2+1}}{x}$	15
pseudoelliptic	$-\frac{\sqrt{-x^2+1}}{x}$	15
risch	$\frac{x^2-1}{x\sqrt{-x^2+1}}$	19
gosper	$\frac{(-1+x)(1+x)}{x\sqrt{-x^2+1}}$	20

input `int(1/x^2/(-x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `-(-x^2+1)^(1/2)/x`

3.124.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{x^2\sqrt{1-x^2}} dx = -\frac{\sqrt{-x^2+1}}{x}$$

input `integrate(1/x^2/(-x^2+1)^(1/2),x, algorithm="fracas")`

output `-sqrt(-x^2 + 1)/x`

3.124.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.41 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.69

$$\int \frac{1}{x^2\sqrt{1-x^2}} dx = \begin{cases} -\frac{i\sqrt{x^2-1}}{x} & \text{for } |x^2| > 1 \\ -\frac{\sqrt{1-x^2}}{x} & \text{otherwise} \end{cases}$$

input `integrate(1/x**2/(-x**2+1)**(1/2),x)`

output `Piecewise((-I*sqrt(x**2 - 1)/x, Abs(x**2) > 1), (-sqrt(1 - x**2)/x, True))`

3.124.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{x^2\sqrt{1-x^2}} dx = -\frac{\sqrt{-x^2+1}}{x}$$

input `integrate(1/x^2/(-x^2+1)^(1/2),x, algorithm="maxima")`

output `-sqrt(-x^2 + 1)/x`

3.124.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 33 vs. 2(14) = 28.

Time = 0.29 (sec) , antiderivative size = 33, normalized size of antiderivative = 2.06

$$\int \frac{1}{x^2\sqrt{1-x^2}} dx = \frac{x}{2(\sqrt{-x^2+1}-1)} - \frac{\sqrt{-x^2+1}-1}{2x}$$

input `integrate(1/x^2/(-x^2+1)^(1/2),x, algorithm="giac")`

output `1/2*x/(sqrt(-x^2 + 1) - 1) - 1/2*(sqrt(-x^2 + 1) - 1)/x`

3.124.9 Mupad [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{x^2\sqrt{1-x^2}} dx = -\frac{\sqrt{1-x^2}}{x}$$

input `int(1/(x^2*(1 - x^2)^(1/2)),x)`

output `-(1 - x^2)^(1/2)/x`

3.125 $\int x^3 \sqrt{4 - x^2} dx$

3.125.1 Optimal result	705
3.125.2 Mathematica [A] (verified)	705
3.125.3 Rubi [A] (verified)	706
3.125.4 Maple [A] (verified)	707
3.125.5 Fricas [A] (verification not implemented)	707
3.125.6 Sympy [A] (verification not implemented)	708
3.125.7 Maxima [A] (verification not implemented)	708
3.125.8 Giac [A] (verification not implemented)	708
3.125.9 Mupad [B] (verification not implemented)	709

3.125.1 Optimal result

Integrand size = 15, antiderivative size = 31

$$\int x^3 \sqrt{4 - x^2} dx = -\frac{4}{3}(4 - x^2)^{3/2} + \frac{1}{5}(4 - x^2)^{5/2}$$

output `-4/3*(-x^2+4)^(3/2)+1/5*(-x^2+4)^(5/2)`

3.125.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

$$\int x^3 \sqrt{4 - x^2} dx = \frac{1}{15} \sqrt{4 - x^2} (-32 - 4x^2 + 3x^4)$$

input `Integrate[x^3*Sqrt[4 - x^2],x]`

output `(Sqrt[4 - x^2]*(-32 - 4*x^2 + 3*x^4))/15`

3.125.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.13, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^3 \sqrt{4-x^2} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int x^2 \sqrt{4-x^2} dx^2 \\ & \quad \downarrow \text{53} \\ & \frac{1}{2} \int \left(4\sqrt{4-x^2} - (4-x^2)^{3/2} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \left(\frac{2}{5} (4-x^2)^{5/2} - \frac{8}{3} (4-x^2)^{3/2} \right) \end{aligned}$$

input `Int[x^3*Sqrt[4 - x^2],x]`

output `((-8*(4 - x^2)^(3/2))/3 + (2*(4 - x^2)^(5/2))/5)/2`

3.125.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.125.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.61

method	result	size
pseudoelliptic	$-\frac{(3x^2+8)(-x^2+4)^{\frac{3}{2}}}{15}$	19
trager	$\left(\frac{1}{5}x^4 - \frac{4}{15}x^2 - \frac{32}{15}\right)\sqrt{-x^2+4}$	23
gosper	$\frac{(-2+x)(2+x)(3x^2+8)\sqrt{-x^2+4}}{15}$	25
default	$-\frac{x^2(-x^2+4)^{\frac{3}{2}}}{5} - \frac{8(-x^2+4)^{\frac{3}{2}}}{15}$	27
risch	$-\frac{(3x^4-4x^2-32)(x^2-4)}{15\sqrt{-x^2+4}}$	29
meijerg	$-\frac{8\left(-\frac{8\sqrt{\pi}}{15} + \frac{4\sqrt{\pi}\left(1-\frac{x^2}{4}\right)^{\frac{3}{2}}\left(\frac{3x^2}{4}+2\right)}{15}\right)}{\sqrt{\pi}}$	33

input `int(x^3*(-x^2+4)^(1/2),x,method=_RETURNVERBOSE)`output `-1/15*(3*x^2+8)*(-x^2+4)^(3/2)`**3.125.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int x^3\sqrt{4-x^2} dx = \frac{1}{15}(3x^4 - 4x^2 - 32)\sqrt{-x^2+4}$$

input `integrate(x^3*(-x^2+4)^(1/2),x, algorithm="fracas")`output `1/15*(3*x^4 - 4*x^2 - 32)*sqrt(-x^2 + 4)`

3.125.6 Sympy [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.26

$$\int x^3 \sqrt{4-x^2} dx = \frac{x^4 \sqrt{4-x^2}}{5} - \frac{4x^2 \sqrt{4-x^2}}{15} - \frac{32 \sqrt{4-x^2}}{15}$$

input `integrate(x**3*(-x**2+4)**(1/2),x)`output `x**4*sqrt(4 - x**2)/5 - 4*x**2*sqrt(4 - x**2)/15 - 32*sqrt(4 - x**2)/15`**3.125.7 Maxima [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int x^3 \sqrt{4-x^2} dx = -\frac{1}{5} (-x^2 + 4)^{\frac{3}{2}} x^2 - \frac{8}{15} (-x^2 + 4)^{\frac{3}{2}}$$

input `integrate(x^3*(-x^2+4)^(1/2),x, algorithm="maxima")`output `-1/5*(-x^2 + 4)^(3/2)*x^2 - 8/15*(-x^2 + 4)^(3/2)`**3.125.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.97

$$\int x^3 \sqrt{4-x^2} dx = \frac{1}{5} (x^2 - 4)^2 \sqrt{-x^2 + 4} - \frac{4}{3} (-x^2 + 4)^{\frac{3}{2}}$$

input `integrate(x^3*(-x^2+4)^(1/2),x, algorithm="giac")`output `1/5*(x^2 - 4)^2*sqrt(-x^2 + 4) - 4/3*(-x^2 + 4)^(3/2)`

3.125.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int x^3 \sqrt{4 - x^2} dx = -\sqrt{4 - x^2} \left(-\frac{x^4}{5} + \frac{4x^2}{15} + \frac{32}{15} \right)$$

input `int(x^3*(4 - x^2)^(1/2),x)`

output `-(4 - x^2)^(1/2)*((4*x^2)/15 - x^4/5 + 32/15)`

3.126 $\int \frac{x}{\sqrt{1-x^2}} dx$

3.126.1 Optimal result	710
3.126.2 Mathematica [A] (verified)	710
3.126.3 Rubi [A] (verified)	711
3.126.4 Maple [A] (verified)	711
3.126.5 Fricas [A] (verification not implemented)	712
3.126.6 Sympy [A] (verification not implemented)	712
3.126.7 Maxima [A] (verification not implemented)	712
3.126.8 Giac [A] (verification not implemented)	713
3.126.9 Mupad [B] (verification not implemented)	713

3.126.1 Optimal result

Integrand size = 13, antiderivative size = 13

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{1-x^2}$$

output `-(-x^2+1)^(1/2)`

3.126.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{1-x^2}$$

input `Integrate[x/Sqrt[1 - x^2],x]`

output `-Sqrt[1 - x^2]`

3.126.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt{1-x^2}} dx$$

↓ 241

$$-\sqrt{1-x^2}$$

input `Int[x/Sqrt[1 - x^2],x]`

output `-Sqrt[1 - x^2]`

3.126.3.1 Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

3.126.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
derivativedivides	$-\sqrt{-x^2+1}$	12
default	$-\sqrt{-x^2+1}$	12
trager	$-\sqrt{-x^2+1}$	12
pseudoelliptic	$-\sqrt{-x^2+1}$	12
risch	$\frac{x^2-1}{\sqrt{-x^2+1}}$	16
gospers	$\frac{(-1+x)(1+x)}{\sqrt{-x^2+1}}$	17
meijerg	$-\frac{-2\sqrt{\pi}+2\sqrt{\pi}\sqrt{-x^2+1}}{2\sqrt{\pi}}$	26

input `int(1/(-x^2+1)^(1/2)*x,x,method=_RETURNVERBOSE)`

output `-(-x^2+1)^(1/2)`

3.126.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{-x^2+1}$$

input `integrate(x/(-x^2+1)^(1/2),x, algorithm="fricas")`

output `-sqrt(-x^2 + 1)`

3.126.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.62

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{1-x^2}$$

input `integrate(x/(-x**2+1)**(1/2),x)`

output `-sqrt(1 - x**2)`

3.126.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{-x^2+1}$$

input `integrate(x/(-x^2+1)^(1/2),x, algorithm="maxima")`

output `-sqrt(-x^2 + 1)`

3.126.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{-x^2+1}$$

input `integrate(x/(-x^2+1)^(1/2),x, algorithm="giac")`

output `-sqrt(-x^2 + 1)`

3.126.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{1-x^2}$$

input `int(x/(1 - x^2)^(1/2),x)`

output `-(1 - x^2)^(1/2)`

3.127 $\int x\sqrt{4-x^2} dx$

3.127.1 Optimal result	714
3.127.2 Mathematica [A] (verified)	714
3.127.3 Rubi [A] (verified)	715
3.127.4 Maple [A] (verified)	715
3.127.5 Fricas [A] (verification not implemented)	716
3.127.6 Sympy [B] (verification not implemented)	717
3.127.7 Maxima [A] (verification not implemented)	717
3.127.8 Giac [A] (verification not implemented)	717
3.127.9 Mupad [B] (verification not implemented)	718

3.127.1 Optimal result

Integrand size = 13, antiderivative size = 15

$$\int x\sqrt{4-x^2} dx = -\frac{1}{3}(4-x^2)^{3/2}$$

output `-1/3*(-x^2+4)^(3/2)`

3.127.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int x\sqrt{4-x^2} dx = -\frac{1}{3}(4-x^2)^{3/2}$$

input `Integrate[x*Sqrt[4 - x^2],x]`

output `-1/3*(4 - x^2)^(3/2)`

3.127.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x\sqrt{4-x^2} dx$$

$$\downarrow \text{241}$$

$$-\frac{1}{3}(4-x^2)^{3/2}$$

input `Int[x*Sqrt[4 - x^2],x]`

output `-1/3*(4 - x^2)^(3/2)`

3.127.3.1 Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

3.127.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

method	result	size
derivativedivides	$-\frac{(-x^2+4)^{\frac{3}{2}}}{3}$	12
default	$-\frac{(-x^2+4)^{\frac{3}{2}}}{3}$	12
pseudoelliptic	$-\frac{(-x^2+4)^{\frac{3}{2}}}{3}$	12
gosper	$\frac{(-2+x)(2+x)\sqrt{-x^2+4}}{3}$	18
trager	$\left(\frac{x^2}{3} - \frac{4}{3}\right)\sqrt{-x^2+4}$	18
risch	$-\frac{(x^2-4)^2}{3\sqrt{-x^2+4}}$	19
meijerg	$\frac{\frac{8\sqrt{\pi}}{3} - \frac{4\sqrt{\pi}\left(-\frac{x^2}{2}+2\right)\sqrt{1-\frac{x^2}{4}}}{3}}{\sqrt{\pi}}$	33

input `int(x*(-x^2+4)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/3*(-x^2+4)^(3/2)`

3.127.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

$$\int x\sqrt{4-x^2} dx = \frac{1}{3}(x^2-4)\sqrt{-x^2+4}$$

input `integrate(x*(-x^2+4)^(1/2),x, algorithm="fracas")`

output `1/3*(x^2 - 4)*sqrt(-x^2 + 4)`

3.127.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 24 vs. $2(10) = 20$.

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.60

$$\int x\sqrt{4-x^2} dx = \frac{x^2\sqrt{4-x^2}}{3} - \frac{4\sqrt{4-x^2}}{3}$$

input `integrate(x*(-x**2+4)**(1/2),x)`

output `x**2*sqrt(4 - x**2)/3 - 4*sqrt(4 - x**2)/3`

3.127.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int x\sqrt{4-x^2} dx = -\frac{1}{3}(-x^2+4)^{\frac{3}{2}}$$

input `integrate(x*(-x^2+4)^(1/2),x, algorithm="maxima")`

output `-1/3*(-x^2 + 4)^(3/2)`

3.127.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int x\sqrt{4-x^2} dx = -\frac{1}{3}(-x^2+4)^{\frac{3}{2}}$$

input `integrate(x*(-x^2+4)^(1/2),x, algorithm="giac")`

output `-1/3*(-x^2 + 4)^(3/2)`

3.127.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int x\sqrt{4-x^2} dx = -\frac{(4-x^2)^{3/2}}{3}$$

input `int(x*(4 - x^2)^(1/2),x)`

output `-(4 - x^2)^(3/2)/3`

3.128 $\int \sqrt{1 - 4x^2} dx$

3.128.1 Optimal result	719
3.128.2 Mathematica [A] (verified)	719
3.128.3 Rubi [A] (verified)	720
3.128.4 Maple [A] (verified)	721
3.128.5 Fricas [A] (verification not implemented)	721
3.128.6 Sympy [A] (verification not implemented)	722
3.128.7 Maxima [A] (verification not implemented)	722
3.128.8 Giac [A] (verification not implemented)	722
3.128.9 Mupad [B] (verification not implemented)	723

3.128.1 Optimal result

Integrand size = 11, antiderivative size = 25

$$\int \sqrt{1 - 4x^2} dx = \frac{1}{2}x\sqrt{1 - 4x^2} + \frac{1}{4} \arcsin(2x)$$

output `1/4*arcsin(2*x)+1/2*x*(-4*x^2+1)^(1/2)`

3.128.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.64

$$\int \sqrt{1 - 4x^2} dx = \frac{1}{2}x\sqrt{1 - 4x^2} - \frac{1}{2} \arctan\left(\frac{\sqrt{1 - 4x^2}}{1 + 2x}\right)$$

input `Integrate[Sqrt[1 - 4*x^2],x]`

output `(x*Sqrt[1 - 4*x^2])/2 - ArcTan[Sqrt[1 - 4*x^2]/(1 + 2*x)]/2`

3.128.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {211, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{1-4x^2} dx$$

$$\downarrow \text{211}$$

$$\frac{1}{2} \int \frac{1}{\sqrt{1-4x^2}} dx + \frac{1}{2} \sqrt{1-4x^2} x$$

$$\downarrow \text{223}$$

$$\frac{1}{4} \arcsin(2x) + \frac{1}{2} \sqrt{1-4x^2} x$$

input `Int[Sqrt[1 - 4*x^2], x]`

output `(x*Sqrt[1 - 4*x^2])/2 + ArcSin[2*x]/4`

3.128.3.1 Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

3.128.4 Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

method	result	size
default	$\frac{\arcsin(2x)}{4} + \frac{x\sqrt{-4x^2+1}}{2}$	20
risch	$-\frac{(4x^2-1)x}{2\sqrt{-4x^2+1}} + \frac{\arcsin(2x)}{4}$	27
pseudoelliptic	$\frac{x\sqrt{-4x^2+1}}{2} - \frac{\arctan\left(\frac{\sqrt{-4x^2+1}}{2x}\right)}{4}$	31
meijerg	$\frac{i(-4i\sqrt{\pi}x\sqrt{-4x^2+1}-2i\sqrt{\pi}\arcsin(2x))}{8\sqrt{\pi}}$	34
trager	$\frac{x\sqrt{-4x^2+1}}{2} - \frac{\text{RootOf}(-Z^2+1)\ln(-\text{RootOf}(-Z^2+1)\sqrt{-4x^2+1}+2x)}{4}$	44

input `int((-4*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`output `1/4*arcsin(2*x)+1/2*x*(-4*x^2+1)^(1/2)`**3.128.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.28

$$\int \sqrt{1-4x^2} dx = \frac{1}{2} \sqrt{-4x^2+1}x - \frac{1}{2} \arctan\left(\frac{\sqrt{-4x^2+1}-1}{2x}\right)$$

input `integrate((-4*x^2+1)^(1/2),x, algorithm="fracas")`output `1/2*sqrt(-4*x^2 + 1)*x - 1/2*arctan(1/2*(sqrt(-4*x^2 + 1) - 1)/x)`

3.128.6 Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \sqrt{1-4x^2} dx = \frac{x\sqrt{1-4x^2}}{2} + \frac{\operatorname{asin}(2x)}{4}$$

input `integrate((-4*x**2+1)**(1/2),x)`output `x*sqrt(1 - 4*x**2)/2 + asin(2*x)/4`**3.128.7 Maxima [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \sqrt{1-4x^2} dx = \frac{1}{2} \sqrt{-4x^2+1}x + \frac{1}{4} \arcsin(2x)$$

input `integrate((-4*x^2+1)^(1/2),x, algorithm="maxima")`output `1/2*sqrt(-4*x^2 + 1)*x + 1/4*arcsin(2*x)`**3.128.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \sqrt{1-4x^2} dx = \frac{1}{2} \sqrt{-4x^2+1}x + \frac{1}{4} \arcsin(2x)$$

input `integrate((-4*x^2+1)^(1/2),x, algorithm="giac")`output `1/2*sqrt(-4*x^2 + 1)*x + 1/4*arcsin(2*x)`

3.128.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.72

$$\int \sqrt{1 - 4x^2} dx = \frac{\arcsin(2x)}{4} + x \sqrt{\frac{1}{4} - x^2}$$

input `int((1 - 4*x^2)^(1/2),x)`

output `asin(2*x)/4 + x*(1/4 - x^2)^(1/2)`

3.129 $\int \frac{x^3}{\sqrt{4+x^2}} dx$

3.129.1 Optimal result	724
3.129.2 Mathematica [A] (verified)	724
3.129.3 Rubi [A] (verified)	725
3.129.4 Maple [A] (verified)	726
3.129.5 Fricas [A] (verification not implemented)	726
3.129.6 Sympy [A] (verification not implemented)	727
3.129.7 Maxima [A] (verification not implemented)	727
3.129.8 Giac [A] (verification not implemented)	727
3.129.9 Mupad [B] (verification not implemented)	728

3.129.1 Optimal result

Integrand size = 13, antiderivative size = 25

$$\int \frac{x^3}{\sqrt{4+x^2}} dx = -4\sqrt{4+x^2} + \frac{1}{3}(4+x^2)^{3/2}$$

output `1/3*(x^2+4)^(3/2)-4*(x^2+4)^(1/2)`

3.129.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.72

$$\int \frac{x^3}{\sqrt{4+x^2}} dx = \frac{1}{3}(-8+x^2)\sqrt{4+x^2}$$

input `Integrate[x^3/Sqrt[4 + x^2],x]`

output `((-8 + x^2)*Sqrt[4 + x^2])/3`

3.129.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.16, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^3}{\sqrt{x^2+4}} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int \frac{x^2}{\sqrt{x^2+4}} dx^2 \\ & \quad \downarrow \text{53} \\ & \frac{1}{2} \int \left(\sqrt{x^2+4} - \frac{4}{\sqrt{x^2+4}} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \left(\frac{2}{3} (x^2+4)^{3/2} - 8\sqrt{x^2+4} \right) \end{aligned}$$

input `Int[x^3/Sqrt[4 + x^2],x]`

output `(-8*Sqrt[4 + x^2] + (2*(4 + x^2)^(3/2))/3)/2`

3.129.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.129.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.60

method	result	size
gosper	$\frac{\sqrt{x^2+4}(x^2-8)}{3}$	15
risch	$\frac{\sqrt{x^2+4}(x^2-8)}{3}$	15
pseudoelliptic	$\frac{\sqrt{x^2+4}(x^2-8)}{3}$	15
trager	$\sqrt{x^2+4} \left(\frac{x^2}{3} - \frac{8}{3} \right)$	16
default	$\frac{x^2\sqrt{x^2+4}}{3} - \frac{8\sqrt{x^2+4}}{3}$	23
meijerg	$\frac{\frac{16\sqrt{\pi}}{3} - \frac{2\sqrt{\pi}(-x^2+8)\sqrt{1+\frac{x^2}{4}}}{3}}{\sqrt{\pi}}$	33

input `int(x^3/(x^2+4)^(1/2),x,method=_RETURNVERBOSE)`

output `1/3*(x^2+4)^(1/2)*(x^2-8)`

3.129.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.56

$$\int \frac{x^3}{\sqrt{4+x^2}} dx = \frac{1}{3} \sqrt{x^2+4}(x^2-8)$$

input `integrate(x^3/(x^2+4)^(1/2),x, algorithm="fricas")`

output `1/3*sqrt(x^2 + 4)*(x^2 - 8)`

3.129.6 Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{x^3}{\sqrt{4+x^2}} dx = \frac{x^2\sqrt{x^2+4}}{3} - \frac{8\sqrt{x^2+4}}{3}$$

input `integrate(x**3/(x**2+4)**(1/2),x)`output `x**2*sqrt(x**2 + 4)/3 - 8*sqrt(x**2 + 4)/3`**3.129.7 Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{x^3}{\sqrt{4+x^2}} dx = \frac{1}{3} \sqrt{x^2+4} x^2 - \frac{8}{3} \sqrt{x^2+4}$$

input `integrate(x^3/(x^2+4)^(1/2),x, algorithm="maxima")`output `1/3*sqrt(x^2 + 4)*x^2 - 8/3*sqrt(x^2 + 4)`**3.129.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{x^3}{\sqrt{4+x^2}} dx = \frac{1}{3} (x^2+4)^{\frac{3}{2}} - 4\sqrt{x^2+4}$$

input `integrate(x^3/(x^2+4)^(1/2),x, algorithm="giac")`output `1/3*(x^2 + 4)^(3/2) - 4*sqrt(x^2 + 4)`

3.129.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.56

$$\int \frac{x^3}{\sqrt{4+x^2}} dx = \frac{\sqrt{x^2+4}(x^2-8)}{3}$$

input `int(x^3/(x^2 + 4)^(1/2),x)`

output `((x^2 + 4)^(1/2)*(x^2 - 8))/3`

3.130 $\int \frac{1}{\sqrt{9+x^2}} dx$

3.130.1 Optimal result	729
3.130.2 Mathematica [B] (verified)	729
3.130.3 Rubi [A] (verified)	730
3.130.4 Maple [A] (verified)	730
3.130.5 Fricas [B] (verification not implemented)	731
3.130.6 Sympy [A] (verification not implemented)	731
3.130.7 Maxima [A] (verification not implemented)	731
3.130.8 Giac [B] (verification not implemented)	732
3.130.9 Mupad [B] (verification not implemented)	732

3.130.1 Optimal result

Integrand size = 9, antiderivative size = 6

$$\int \frac{1}{\sqrt{9+x^2}} dx = \operatorname{arcsinh}\left(\frac{x}{3}\right)$$

output `arcsinh(1/3*x)`

3.130.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 16 vs. $2(6) = 12$.

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 2.67

$$\int \frac{1}{\sqrt{9+x^2}} dx = -\log\left(-x + \sqrt{9+x^2}\right)$$

input `Integrate[1/Sqrt[9 + x^2],x]`

output `-Log[-x + Sqrt[9 + x^2]]`

3.130.3 Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x^2 + 9}} dx$$

↓ 222

$$\operatorname{arcsinh}\left(\frac{x}{3}\right)$$

input `Int[1/Sqrt[9 + x^2], x]`

output `ArcSinh[x/3]`

3.130.3.1 Defintions of rubi rules used

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

3.130.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

method	result	size
default	$\operatorname{arcsinh}\left(\frac{x}{3}\right)$	5
meijerg	$\operatorname{arcsinh}\left(\frac{x}{3}\right)$	5
pseudoelliptic	$\operatorname{arctanh}\left(\frac{\sqrt{x^2+9}}{x}\right)$	13
trager	$-\ln(x - \sqrt{x^2 + 9})$	15

input `int(1/(x^2+9)^(1/2), x, method=_RETURNVERBOSE)`

output `arcsinh(1/3*x)`

3.130.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 14 vs. $2(4) = 8$.

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 2.33

$$\int \frac{1}{\sqrt{9+x^2}} dx = -\log\left(-x + \sqrt{x^2+9}\right)$$

input `integrate(1/(x^2+9)^(1/2),x, algorithm="fricas")`

output `-log(-x + sqrt(x^2 + 9))`

3.130.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.50

$$\int \frac{1}{\sqrt{9+x^2}} dx = \operatorname{asinh}\left(\frac{x}{3}\right)$$

input `integrate(1/(x**2+9)**(1/2),x)`

output `asinh(x/3)`

3.130.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt{9+x^2}} dx = \operatorname{arsinh}\left(\frac{1}{3}x\right)$$

input `integrate(1/(x^2+9)^(1/2),x, algorithm="maxima")`

output `arcsinh(1/3*x)`

3.130.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. $2(4) = 8$.

Time = 0.28 (sec) , antiderivative size = 25, normalized size of antiderivative = 4.17

$$\int \frac{1}{\sqrt{9+x^2}} dx = \frac{1}{2} \sqrt{x^2+9}x - \frac{9}{2} \log(-x + \sqrt{x^2+9})$$

input `integrate(1/(x^2+9)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(x^2 + 9)*x - 9/2*log(-x + sqrt(x^2 + 9))`

3.130.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt{9+x^2}} dx = \operatorname{asinh}\left(\frac{x}{3}\right)$$

input `int(1/(x^2 + 9)^(1/2),x)`

output `asinh(x/3)`

3.131 $\int \sqrt{1+x^2} dx$

3.131.1 Optimal result	733
3.131.2 Mathematica [A] (verified)	733
3.131.3 Rubi [A] (verified)	734
3.131.4 Maple [A] (verified)	735
3.131.5 Fricas [A] (verification not implemented)	735
3.131.6 Sympy [A] (verification not implemented)	735
3.131.7 Maxima [A] (verification not implemented)	736
3.131.8 Giac [A] (verification not implemented)	736
3.131.9 Mupad [B] (verification not implemented)	736

3.131.1 Optimal result

Integrand size = 9, antiderivative size = 21

$$\int \sqrt{1+x^2} dx = \frac{1}{2}x\sqrt{1+x^2} + \frac{\operatorname{arcsinh}(x)}{2}$$

output `1/2*arcsinh(x)+1/2*x*(x^2+1)^(1/2)`

3.131.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.57

$$\int \sqrt{1+x^2} dx = \frac{1}{2}x\sqrt{1+x^2} - \frac{1}{2} \log(-x + \sqrt{1+x^2})$$

input `Integrate[Sqrt[1 + x^2],x]`

output `(x*Sqrt[1 + x^2])/2 - Log[-x + Sqrt[1 + x^2]]/2`

3.131.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {211, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x^2 + 1} dx$$

$$\downarrow \text{211}$$

$$\frac{1}{2} \int \frac{1}{\sqrt{x^2 + 1}} dx + \frac{1}{2} \sqrt{x^2 + 1} x$$

$$\downarrow \text{222}$$

$$\frac{\operatorname{arcsinh}(x)}{2} + \frac{1}{2} \sqrt{x^2 + 1} x$$

input `Int[Sqrt[1 + x^2], x]`

output `(x*Sqrt[1 + x^2])/2 + ArcSinh[x]/2`

3.131.3.1 Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

3.131.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.76

method	result	size
default	$\frac{\operatorname{arcsinh}(x)}{2} + \frac{x\sqrt{x^2+1}}{2}$	16
risch	$\frac{\operatorname{arcsinh}(x)}{2} + \frac{x\sqrt{x^2+1}}{2}$	16
trager	$\frac{x\sqrt{x^2+1}}{2} + \frac{\ln(x+\sqrt{x^2+1})}{2}$	24
meijerg	$-\frac{-2\sqrt{\pi}x\sqrt{x^2+1}-2\sqrt{\pi}\operatorname{arcsinh}(x)}{4\sqrt{\pi}}$	27
pseudoelliptic	$\frac{x\sqrt{x^2+1}}{2} + \frac{\ln\left(\frac{x+\sqrt{x^2+1}}{x}\right)}{4} - \frac{\ln\left(\frac{\sqrt{x^2+1}-x}{x}\right)}{4}$	46

input `int((x^2+1)^(1/2),x,method=_RETURNVERBOSE)`output `1/2*arcsinh(x)+1/2*x*(x^2+1)^(1/2)`**3.131.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.19

$$\int \sqrt{1+x^2} dx = \frac{1}{2} \sqrt{x^2+1}x - \frac{1}{2} \log(-x + \sqrt{x^2+1})$$

input `integrate((x^2+1)^(1/2),x, algorithm="fricas")`output `1/2*sqrt(x^2 + 1)*x - 1/2*log(-x + sqrt(x^2 + 1))`**3.131.6 Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \sqrt{1+x^2} dx = \frac{x\sqrt{x^2+1}}{2} + \frac{\operatorname{asinh}(x)}{2}$$

input `integrate((x**2+1)**(1/2),x)`

output `x*sqrt(x**2 + 1)/2 + asinh(x)/2`

3.131.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \sqrt{1+x^2} dx = \frac{1}{2} \sqrt{x^2+1}x + \frac{1}{2} \operatorname{arsinh}(x)$$

input `integrate((x^2+1)^(1/2),x, algorithm="maxima")`

output `1/2*sqrt(x^2 + 1)*x + 1/2*arcsinh(x)`

3.131.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.19

$$\int \sqrt{1+x^2} dx = \frac{1}{2} \sqrt{x^2+1}x - \frac{1}{2} \log(-x + \sqrt{x^2+1})$$

input `integrate((x^2+1)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(x^2 + 1)*x - 1/2*log(-x + sqrt(x^2 + 1))`

3.131.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \sqrt{1+x^2} dx = \frac{\operatorname{asinh}(x)}{2} + \frac{x\sqrt{x^2+1}}{2}$$

input `int((x^2 + 1)^(1/2),x)`

output `asinh(x)/2 + (x*(x^2 + 1)^(1/2))/2`

3.132 $\int \frac{1}{x^3\sqrt{-16+x^2}} dx$

3.132.1 Optimal result	737
3.132.2 Mathematica [A] (verified)	737
3.132.3 Rubi [A] (verified)	738
3.132.4 Maple [A] (verified)	739
3.132.5 Fricas [A] (verification not implemented)	740
3.132.6 Sympy [C] (verification not implemented)	741
3.132.7 Maxima [A] (verification not implemented)	741
3.132.8 Giac [A] (verification not implemented)	741
3.132.9 Mupad [B] (verification not implemented)	742

3.132.1 Optimal result

Integrand size = 13, antiderivative size = 35

$$\int \frac{1}{x^3\sqrt{-16+x^2}} dx = \frac{\sqrt{-16+x^2}}{32x^2} + \frac{1}{128} \arctan\left(\frac{1}{4}\sqrt{-16+x^2}\right)$$

output `1/128*arctan(1/4*(x^2-16)^(1/2))+1/32*(x^2-16)^(1/2)/x^2`

3.132.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^3\sqrt{-16+x^2}} dx = \frac{\sqrt{-16+x^2}}{32x^2} + \frac{1}{128} \arctan\left(\frac{1}{4}\sqrt{-16+x^2}\right)$$

input `Integrate[1/(x^3*Sqrt[-16 + x^2]),x]`

output `Sqrt[-16 + x^2]/(32*x^2) + ArcTan[Sqrt[-16 + x^2]/4]/128`

3.132.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.11, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {243, 52, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^3 \sqrt{x^2 - 16}} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \int \frac{1}{x^4 \sqrt{x^2 - 16}} dx^2 \\
 & \quad \downarrow \text{52} \\
 & \frac{1}{2} \left(\frac{1}{32} \int \frac{1}{x^2 \sqrt{x^2 - 16}} dx^2 + \frac{\sqrt{x^2 - 16}}{16x^2} \right) \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{2} \left(\frac{1}{16} \int \frac{1}{x^4 + 16} d\sqrt{x^2 - 16} + \frac{\sqrt{x^2 - 16}}{16x^2} \right) \\
 & \quad \downarrow \text{216} \\
 & \frac{1}{2} \left(\frac{1}{64} \arctan \left(\frac{\sqrt{x^2 - 16}}{4} \right) + \frac{\sqrt{x^2 - 16}}{16x^2} \right)
 \end{aligned}$$

input `Int[1/(x^3*sqrt[-16 + x^2]),x]`

output `(sqrt[-16 + x^2]/(16*x^2) + ArcTan[sqrt[-16 + x^2]/4]/64)/2`

3.132.3.1 Defintions of rubi rules used

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

3.132.4 Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

method	result	si
default	$\frac{\sqrt{x^2-16}}{32x^2} - \frac{\arctan\left(\frac{4}{\sqrt{x^2-16}}\right)}{128}$	20
risch	$\frac{\sqrt{x^2-16}}{32x^2} - \frac{\arctan\left(\frac{4}{\sqrt{x^2-16}}\right)}{128}$	20
pseudoelliptic	$\frac{\arctan\left(\frac{\sqrt{x^2-16}}{4}\right)x^2 + 4\sqrt{x^2-16}}{128x^2}$	30
trager	$\frac{\sqrt{x^2-16}}{32x^2} - \frac{\text{RootOf}(_Z^2+1) \ln\left(-\frac{4 \text{RootOf}(_Z^2+1) - \sqrt{x^2-16}}{x}\right)}{128}$	40
meijerg	$-\frac{\sqrt{-\text{signum}\left(-1+\frac{x^2}{16}\right)} \left(\frac{16\sqrt{\pi}}{x^2} - \frac{(1-6\ln(2)+2\ln(x)+i\pi)\sqrt{\pi}}{2} - \frac{2\sqrt{\pi}\left(-\frac{x^2}{4}+8\right)}{x^2} + \frac{16\sqrt{\pi}\sqrt{1-\frac{x^2}{16}}}{x^2} + \sqrt{\pi} \ln\left(\frac{1}{2} + \frac{\sqrt{1-\frac{x^2}{16}}}{2}\right)\right)}{128\sqrt{\pi}\sqrt{\text{signum}\left(-1+\frac{x^2}{16}\right)}}$	10

input `int(1/x^3/(x^2-16)^(1/2),x,method=_RETURNVERBOSE)`

output `1/32*(x^2-16)^(1/2)/x^2-1/128*arctan(4/(x^2-16)^(1/2))`

3.132.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{1}{x^3\sqrt{-16+x^2}} dx = \frac{x^2 \arctan\left(-\frac{1}{4}x + \frac{1}{4}\sqrt{x^2-16}\right) + 2\sqrt{x^2-16}}{64x^2}$$

input `integrate(1/x^3/(x^2-16)^(1/2),x, algorithm="fracas")`

output `1/64*(x^2*arctan(-1/4*x + 1/4*sqrt(x^2 - 16)) + 2*sqrt(x^2 - 16))/x^2`

3.132.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.14 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.89

$$\int \frac{1}{x^3 \sqrt{-16 + x^2}} dx = \begin{cases} \frac{i \operatorname{acosh}\left(\frac{4}{x}\right)}{128} - \frac{i}{32x \sqrt{-1 + \frac{16}{x^2}}} + \frac{i}{2x^3 \sqrt{-1 + \frac{16}{x^2}}} & \text{for } \frac{1}{|x^2|} > \frac{1}{16} \\ -\frac{\operatorname{asin}\left(\frac{4}{x}\right)}{128} + \frac{\sqrt{1 - \frac{16}{x^2}}}{32x} & \text{otherwise} \end{cases}$$

input `integrate(1/x**3/(x**2-16)**(1/2),x)`

output `Piecewise((I*acosh(4/x)/128 - I/(32*x*sqrt(-1 + 16/x**2)) + I/(2*x**3*sqrt(-1 + 16/x**2)), 1/Abs(x**2) > 1/16), (-asin(4/x)/128 + sqrt(1 - 16/x**2)/(32*x), True))`

3.132.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.63

$$\int \frac{1}{x^3 \sqrt{-16 + x^2}} dx = \frac{\sqrt{x^2 - 16}}{32 x^2} - \frac{1}{128} \arcsin\left(\frac{4}{|x|}\right)$$

input `integrate(1/x^3/(x^2-16)^(1/2),x, algorithm="maxima")`

output `1/32*sqrt(x^2 - 16)/x^2 - 1/128*arcsin(4/abs(x))`

3.132.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int \frac{1}{x^3 \sqrt{-16 + x^2}} dx = \frac{\sqrt{x^2 - 16}}{32 x^2} + \frac{1}{128} \arctan\left(\frac{1}{4} \sqrt{x^2 - 16}\right)$$

input `integrate(1/x^3/(x^2-16)^(1/2),x, algorithm="giac")`

output `1/32*sqrt(x^2 - 16)/x^2 + 1/128*arctan(1/4*sqrt(x^2 - 16))`

3.132.9 Mupad [B] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int \frac{1}{x^3 \sqrt{-16 + x^2}} dx = \frac{\operatorname{atan}\left(\frac{\sqrt{x^2-16}}{4}\right)}{128} + \frac{\sqrt{x^2-16}}{32x^2}$$

input `int(1/(x^3*(x^2 - 16)^(1/2)),x)`

output `atan((x^2 - 16)^(1/2)/4)/128 + (x^2 - 16)^(1/2)/(32*x^2)`

3.133 $\int \frac{\sqrt{-a^2+x^2}}{x^4} dx$

3.133.1 Optimal result	743
3.133.2 Mathematica [A] (verified)	743
3.133.3 Rubi [A] (verified)	744
3.133.4 Maple [A] (verified)	744
3.133.5 Fricas [A] (verification not implemented)	745
3.133.6 Sympy [C] (verification not implemented)	745
3.133.7 Maxima [A] (verification not implemented)	746
3.133.8 Giac [B] (verification not implemented)	746
3.133.9 Mupad [B] (verification not implemented)	746

3.133.1 Optimal result

Integrand size = 17, antiderivative size = 23

$$\int \frac{\sqrt{-a^2+x^2}}{x^4} dx = \frac{(-a^2+x^2)^{3/2}}{3a^2x^3}$$

output $1/3*(-a^2+x^2)^{(3/2)}/a^2/x^3$

3.133.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{-a^2+x^2}}{x^4} dx = \frac{(-a^2+x^2)^{3/2}}{3a^2x^3}$$

input `Integrate[Sqrt[-a^2 + x^2]/x^4,x]`

output $(-a^2+x^2)^{(3/2)}/(3*a^2*x^3)$

3.133.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{x^2 - a^2}}{x^4} dx$$

↓ 242

$$\frac{(x^2 - a^2)^{3/2}}{3a^2x^3}$$

input `Int[Sqrt[-a^2 + x^2]/x^4,x]`

output `(-a^2 + x^2)^(3/2)/(3*a^2*x^3)`

3.133.3.1 Defintions of rubi rules used

rule 242 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

3.133.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

method	result	size
default	$\frac{(-a^2+x^2)^{\frac{3}{2}}}{3a^2x^3}$	20
pseudoelliptic	$\frac{(-a^2+x^2)^{\frac{3}{2}}}{3a^2x^3}$	20
gospers	$-\frac{(a-x)(a+x)\sqrt{-a^2+x^2}}{3x^3a^2}$	28
trager	$-\frac{(a^2-x^2)\sqrt{-a^2+x^2}}{3a^2x^3}$	29
risch	$\frac{(a^2-x^2)^2}{3x^3\sqrt{-a^2+x^2}a^2}$	31

input `int((-a^2+x^2)^(1/2)/x^4,x,method=_RETURNVERBOSE)`

output `1/3*(-a^2+x^2)^(3/2)/a^2/x^3`

3.133.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{-a^2 + x^2}}{x^4} dx = \frac{x^3 + (-a^2 + x^2)^{\frac{3}{2}}}{3a^2x^3}$$

input `integrate((-a^2+x^2)^(1/2)/x^4,x, algorithm="fricas")`

output `1/3*(x^3 + (-a^2 + x^2)^(3/2))/(a^2*x^3)`

3.133.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.43 (sec) , antiderivative size = 76, normalized size of antiderivative = 3.30

$$\int \frac{\sqrt{-a^2 + x^2}}{x^4} dx = \begin{cases} -\frac{i\sqrt{\frac{a^2}{x^2}-1}}{3x^2} + \frac{i\sqrt{\frac{a^2}{x^2}-1}}{3a^2} & \text{for } \left|\frac{a^2}{x^2}\right| > 1 \\ -\frac{\sqrt{-\frac{a^2}{x^2}+1}}{3x^2} + \frac{\sqrt{-\frac{a^2}{x^2}+1}}{3a^2} & \text{otherwise} \end{cases}$$

input `integrate((-a**2+x**2)**(1/2)/x**4,x)`

output `Piecewise((-I*sqrt(a**2/x**2 - 1)/(3*x**2) + I*sqrt(a**2/x**2 - 1)/(3*a**2), Abs(a**2/x**2) > 1), (-sqrt(-a**2/x**2 + 1)/(3*x**2) + sqrt(-a**2/x**2 + 1)/(3*a**2), True))`

3.133.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{\sqrt{-a^2 + x^2}}{x^4} dx = \frac{(-a^2 + x^2)^{\frac{3}{2}}}{3a^2x^3}$$

input `integrate((-a^2+x^2)^(1/2)/x^4,x, algorithm="maxima")`

output `1/3*(-a^2 + x^2)^(3/2)/(a^2*x^3)`

3.133.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 48 vs. 2(19) = 38.

Time = 0.30 (sec) , antiderivative size = 48, normalized size of antiderivative = 2.09

$$\int \frac{\sqrt{-a^2 + x^2}}{x^4} dx = \frac{2 \left(a^4 + 3 (x - \sqrt{-a^2 + x^2})^4 \right)}{3 \left(a^2 + (x - \sqrt{-a^2 + x^2})^2 \right)^3}$$

input `integrate((-a^2+x^2)^(1/2)/x^4,x, algorithm="giac")`

output `2/3*(a^4 + 3*(x - sqrt(-a^2 + x^2))^4)/(a^2 + (x - sqrt(-a^2 + x^2))^2)^3`

3.133.9 Mupad [B] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{\sqrt{-a^2 + x^2}}{x^4} dx = \frac{(x^2 - a^2)^{3/2}}{3a^2x^3}$$

input `int((x^2 - a^2)^(1/2)/x^4,x)`

output `(x^2 - a^2)^(3/2)/(3*a^2*x^3)`

3.134 $\int \frac{\sqrt{-4+9x^2}}{x} dx$

3.134.1 Optimal result	747
3.134.2 Mathematica [A] (verified)	747
3.134.3 Rubi [A] (verified)	748
3.134.4 Maple [A] (verified)	749
3.134.5 Fricas [A] (verification not implemented)	750
3.134.6 Sympy [C] (verification not implemented)	750
3.134.7 Maxima [A] (verification not implemented)	750
3.134.8 Giac [A] (verification not implemented)	751
3.134.9 Mupad [B] (verification not implemented)	751

3.134.1 Optimal result

Integrand size = 15, antiderivative size = 30

$$\int \frac{\sqrt{-4+9x^2}}{x} dx = \sqrt{-4+9x^2} - 2 \arctan\left(\frac{1}{2}\sqrt{-4+9x^2}\right)$$

output `-2*arctan(1/2*(9*x^2-4)^(1/2))+ (9*x^2-4)^(1/2)`

3.134.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{-4+9x^2}}{x} dx = \sqrt{-4+9x^2} - 2 \arctan\left(\frac{1}{2}\sqrt{-4+9x^2}\right)$$

input `Integrate[Sqrt[-4 + 9*x^2]/x,x]`

output `Sqrt[-4 + 9*x^2] - 2*ArcTan[Sqrt[-4 + 9*x^2]/2]`

3.134.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.20, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {243, 60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{9x^2 - 4}}{x} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \int \frac{\sqrt{9x^2 - 4}}{x^2} dx^2 \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2} \left(2\sqrt{9x^2 - 4} - 4 \int \frac{1}{x^2 \sqrt{9x^2 - 4}} dx^2 \right) \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{2} \left(2\sqrt{9x^2 - 4} - \frac{8}{9} \int \frac{1}{\frac{x^4}{9} + \frac{4}{9}} d\sqrt{9x^2 - 4} \right) \\
 & \quad \downarrow \text{216} \\
 & \frac{1}{2} \left(2\sqrt{9x^2 - 4} - 4 \arctan \left(\frac{1}{2} \sqrt{9x^2 - 4} \right) \right)
 \end{aligned}$$

input `Int[Sqrt[-4 + 9*x^2]/x,x]`

output `(2*Sqrt[-4 + 9*x^2] - 4*ArcTan[Sqrt[-4 + 9*x^2]/2])/2`

3.134.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

```
rule 243 Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[In
t[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I
ntegerQ[(m - 1)/2]
```

3.134.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.83

method	result	size
default	$\sqrt{9x^2 - 4} + 2 \arctan\left(\frac{2}{\sqrt{9x^2 - 4}}\right)$	25
pseudoelliptic	$-2 \arctan\left(\frac{\sqrt{9x^2 - 4}}{2}\right) + \sqrt{9x^2 - 4}$	25
trager	$\sqrt{9x^2 - 4} - 2 \operatorname{RootOf}(_Z^2 + 1) \ln\left(\frac{2 \operatorname{RootOf}(_Z^2 + 1) + \sqrt{9x^2 - 4}}{x}\right)$	42
meijerg	$-\frac{\sqrt{\operatorname{signum}\left(-1 + \frac{9x^2}{4}\right)} \left(-2(2 - 4 \ln(2) + 2 \ln(x) + 2 \ln(3) + i\pi)\sqrt{\pi} + 4\sqrt{\pi} - 4\sqrt{\pi} \sqrt{1 - \frac{9x^2}{4}} + 4\sqrt{\pi} \ln\left(\frac{1}{2} + \frac{\sqrt{1 - \frac{9x^2}{4}}}{2}\right)\right)}{2\sqrt{\pi} \sqrt{-\operatorname{signum}\left(-1 + \frac{9x^2}{4}\right)}}$	90

```
input int((9*x^2-4)^(1/2)/x,x,method=_RETURNVERBOSE)
```

```
output (9*x^2-4)^(1/2)+2*arctan(2/(9*x^2-4)^(1/2))
```

3.134.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{-4+9x^2}}{x} dx = \sqrt{9x^2-4} - 4 \arctan\left(-\frac{3}{2}x + \frac{1}{2}\sqrt{9x^2-4}\right)$$

input `integrate((9*x^2-4)^(1/2)/x,x, algorithm="fricas")`output `sqrt(9*x^2 - 4) - 4*arctan(-3/2*x + 1/2*sqrt(9*x^2 - 4))`**3.134.6 Sympy [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.81 (sec) , antiderivative size = 92, normalized size of antiderivative = 3.07

$$\int \frac{\sqrt{-4+9x^2}}{x} dx = \begin{cases} -\frac{3ix}{\sqrt{-1+\frac{4}{9x^2}}} - 2i \operatorname{acosh}\left(\frac{2}{3x}\right) + \frac{4i}{3x\sqrt{-1+\frac{4}{9x^2}}} & \text{for } \frac{1}{|x^2|} > \frac{9}{4} \\ \frac{3x}{\sqrt{1-\frac{4}{9x^2}}} + 2 \operatorname{asin}\left(\frac{2}{3x}\right) - \frac{4}{3x\sqrt{1-\frac{4}{9x^2}}} & \text{otherwise} \end{cases}$$

input `integrate((9*x**2-4)**(1/2)/x,x)`output `Piecewise((-3*I*x/sqrt(-1 + 4/(9*x**2)) - 2*I*acosh(2/(3*x)) + 4*I/(3*x*sqrt(-1 + 4/(9*x**2))), 1/Abs(x**2) > 9/4), (3*x/sqrt(1 - 4/(9*x**2)) + 2*asin(2/(3*x)) - 4/(3*x*sqrt(1 - 4/(9*x**2))), True))`**3.134.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.63

$$\int \frac{\sqrt{-4+9x^2}}{x} dx = \sqrt{9x^2-4} + 2 \arcsin\left(\frac{2}{3|x|}\right)$$

input `integrate((9*x^2-4)^(1/2)/x,x, algorithm="maxima")`output `sqrt(9*x^2 - 4) + 2*arcsin(2/3/abs(x))`

3.134.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int \frac{\sqrt{-4+9x^2}}{x} dx = \sqrt{9x^2-4} - 2 \arctan\left(\frac{1}{2}\sqrt{9x^2-4}\right)$$

input `integrate((9*x^2-4)^(1/2)/x,x, algorithm="giac")`output `sqrt(9*x^2 - 4) - 2*arctan(1/2*sqrt(9*x^2 - 4))`**3.134.9 Mupad [B] (verification not implemented)**

Time = 0.38 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int \frac{\sqrt{-4+9x^2}}{x} dx = \sqrt{9x^2-4} - 2 \operatorname{atan}\left(\frac{\sqrt{9x^2-4}}{2}\right)$$

input `int((9*x^2 - 4)^(1/2)/x,x)`output `(9*x^2 - 4)^(1/2) - 2*atan((9*x^2 - 4)^(1/2)/2)`

$$\mathbf{3.135} \quad \int \frac{1}{x^2 \sqrt{-9+16x^2}} dx$$

3.135.1 Optimal result	752
3.135.2 Mathematica [A] (verified)	752
3.135.3 Rubi [A] (verified)	753
3.135.4 Maple [A] (verified)	754
3.135.5 Fricas [A] (verification not implemented)	754
3.135.6 Sympy [C] (verification not implemented)	755
3.135.7 Maxima [A] (verification not implemented)	755
3.135.8 Giac [A] (verification not implemented)	755
3.135.9 Mupad [B] (verification not implemented)	756

3.135.1 Optimal result

Integrand size = 15, antiderivative size = 18

$$\int \frac{1}{x^2 \sqrt{-9+16x^2}} dx = \frac{\sqrt{-9+16x^2}}{9x}$$

output `1/9*(16*x^2-9)^(1/2)/x`

3.135.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2 \sqrt{-9+16x^2}} dx = \frac{\sqrt{-9+16x^2}}{9x}$$

input `Integrate[1/(x^2*Sqrt[-9 + 16*x^2]),x]`

output `Sqrt[-9 + 16*x^2]/(9*x)`

3.135.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$, Rules used = {242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 \sqrt{16x^2 - 9}} dx$$

↓ 242

$$\frac{\sqrt{16x^2 - 9}}{9x}$$

input `Int[1/(x^2*Sqrt[-9 + 16*x^2]),x]`

output `Sqrt[-9 + 16*x^2]/(9*x)`

3.135.3.1 Defintions of rubi rules used

rule 242 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

3.135.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{\sqrt{16x^2-9}}{9x}$	15
trager	$\frac{\sqrt{16x^2-9}}{9x}$	15
risch	$\frac{\sqrt{16x^2-9}}{9x}$	15
pseudoelliptic	$\frac{\sqrt{16x^2-9}}{9x}$	15
gospers	$\frac{(4x-3)(4x+3)}{9x\sqrt{16x^2-9}}$	25
meijerg	$-\frac{\sqrt{-\operatorname{signum}\left(-1+\frac{16x^2}{9}\right)}\sqrt{1-\frac{16x^2}{9}}}{3\sqrt{\operatorname{signum}\left(-1+\frac{16x^2}{9}\right)}x}$	37

input `int(1/x^2/(16*x^2-9)^(1/2),x,method=_RETURNVERBOSE)`output `1/9*(16*x^2-9)^(1/2)/x`**3.135.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2\sqrt{-9+16x^2}} dx = \frac{4x + \sqrt{16x^2 - 9}}{9x}$$

input `integrate(1/x^2/(16*x^2-9)^(1/2),x, algorithm="fracas")`output `1/9*(4*x + sqrt(16*x^2 - 9))/x`

3.135.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.45 (sec) , antiderivative size = 37, normalized size of antiderivative = 2.06

$$\int \frac{1}{x^2 \sqrt{-9 + 16x^2}} dx = \begin{cases} \frac{4i\sqrt{-1 + \frac{9}{16x^2}}}{9} & \text{for } \frac{1}{|x^2|} > \frac{16}{9} \\ \frac{4\sqrt{1 - \frac{9}{16x^2}}}{9} & \text{otherwise} \end{cases}$$

input `integrate(1/x**2/(16*x**2-9)**(1/2),x)`

output `Piecewise((4*I*sqrt(-1 + 9/(16*x**2)))/9, 1/Abs(x**2) > 16/9), (4*sqrt(1 - 9/(16*x**2)))/9, True))`

3.135.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{1}{x^2 \sqrt{-9 + 16x^2}} dx = \frac{\sqrt{16x^2 - 9}}{9x}$$

input `integrate(1/x^2/(16*x^2-9)^(1/2),x, algorithm="maxima")`

output `1/9*sqrt(16*x^2 - 9)/x`

3.135.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.28

$$\int \frac{1}{x^2 \sqrt{-9 + 16x^2}} dx = \frac{8}{(4x - \sqrt{16x^2 - 9})^2 + 9}$$

input `integrate(1/x^2/(16*x^2-9)^(1/2),x, algorithm="giac")`

output `8/((4*x - sqrt(16*x^2 - 9))^2 + 9)`

3.135.9 Mupad [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{1}{x^2\sqrt{-9+16x^2}} dx = \frac{\sqrt{16x^2-9}}{9x}$$

input `int(1/(x^2*(16*x^2 - 9)^(1/2)),x)`

output `(16*x^2 - 9)^(1/2)/(9*x)`

3.136 $\int \frac{x^2}{(a^2-x^2)^{3/2}} dx$

3.136.1 Optimal result	757
3.136.2 Mathematica [A] (verified)	757
3.136.3 Rubi [A] (verified)	758
3.136.4 Maple [A] (verified)	759
3.136.5 Fricas [A] (verification not implemented)	759
3.136.6 Sympy [C] (verification not implemented)	760
3.136.7 Maxima [A] (verification not implemented)	760
3.136.8 Giac [A] (verification not implemented)	760
3.136.9 Mupad [B] (verification not implemented)	761

3.136.1 Optimal result

Integrand size = 17, antiderivative size = 34

$$\int \frac{x^2}{(a^2-x^2)^{3/2}} dx = \frac{x}{\sqrt{a^2-x^2}} - \arctan\left(\frac{x}{\sqrt{a^2-x^2}}\right)$$

output `-arctan(x/(a^2-x^2)^(1/2))+x/(a^2-x^2)^(1/2)`

3.136.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{(a^2-x^2)^{3/2}} dx = \frac{x}{\sqrt{a^2-x^2}} - \arctan\left(\frac{x}{\sqrt{a^2-x^2}}\right)$$

input `Integrate[x^2/(a^2 - x^2)^(3/2),x]`

output `x/Sqrt[a^2 - x^2] - ArcTan[x/Sqrt[a^2 - x^2]]`

3.136.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {252, 224, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{(a^2 - x^2)^{3/2}} dx$$

$$\downarrow 252$$

$$\frac{x}{\sqrt{a^2 - x^2}} - \int \frac{1}{\sqrt{a^2 - x^2}} dx$$

$$\downarrow 224$$

$$\frac{x}{\sqrt{a^2 - x^2}} - \int \frac{1}{\frac{x^2}{a^2 - x^2} + 1} d\frac{x}{\sqrt{a^2 - x^2}}$$

$$\downarrow 216$$

$$\frac{x}{\sqrt{a^2 - x^2}} - \arctan\left(\frac{x}{\sqrt{a^2 - x^2}}\right)$$

input `Int[x^2/(a^2 - x^2)^(3/2),x]`

output `x/Sqrt[a^2 - x^2] - ArcTan[x/Sqrt[a^2 - x^2]]`

3.136.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

```
rule 252 Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*((m - 1)/(2*b*(p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

3.136.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.91

method	result	size
default	$-\arctan\left(\frac{x}{\sqrt{a^2-x^2}}\right) + \frac{x}{\sqrt{a^2-x^2}}$	31
pseudoelliptic	$\frac{\arctan\left(\frac{\sqrt{a^2-x^2}}{x}\right)\sqrt{a^2-x^2}+x}{\sqrt{a^2-x^2}}$	43

```
input int(x^2/(a^2-x^2)^(3/2),x,method=_RETURNVERBOSE)
```

```
output -arctan(x/(a^2-x^2)^(1/2))+x/(a^2-x^2)^(1/2)
```

3.136.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.71

$$\int \frac{x^2}{(a^2 - x^2)^{3/2}} dx = \frac{2(a^2 - x^2) \arctan\left(-\frac{a - \sqrt{a^2 - x^2}}{x}\right) + \sqrt{a^2 - x^2}x}{a^2 - x^2}$$

```
input integrate(x^2/(a^2-x^2)^(3/2),x, algorithm="fracas")
```

```
output (2*(a^2 - x^2)*arctan(-(a - sqrt(a^2 - x^2))/x) + sqrt(a^2 - x^2)*x)/(a^2 - x^2)
```


3.136.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.84 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.44

$$\int \frac{x^2}{(a^2 - x^2)^{3/2}} dx = \begin{cases} i \operatorname{acosh}\left(\frac{x}{a}\right) - \frac{ix}{a\sqrt{-1 + \frac{x^2}{a^2}}} & \text{for } \left|\frac{x^2}{a^2}\right| > 1 \\ -\operatorname{asin}\left(\frac{x}{a}\right) + \frac{x}{a\sqrt{1 - \frac{x^2}{a^2}}} & \text{otherwise} \end{cases}$$

input `integrate(x**2/(a**2-x**2)**(3/2),x)`

output `Piecewise((I*acosh(x/a) - I*x/(a*sqrt(-1 + x**2/a**2)), Abs(x**2/a**2) > 1), (-asin(x/a) + x/(a*sqrt(1 - x**2/a**2)), True))`

3.136.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.65

$$\int \frac{x^2}{(a^2 - x^2)^{3/2}} dx = \frac{x}{\sqrt{a^2 - x^2}} - \operatorname{arcsin}\left(\frac{x}{a}\right)$$

input `integrate(x^2/(a^2-x^2)^(3/2),x, algorithm="maxima")`

output `x/sqrt(a^2 - x^2) - arcsin(x/a)`

3.136.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.71

$$\int \frac{x^2}{(a^2 - x^2)^{3/2}} dx = -\operatorname{arcsin}\left(\frac{x}{a}\right) \operatorname{sgn}(a) + \frac{x}{\sqrt{a^2 - x^2}}$$

input `integrate(x^2/(a^2-x^2)^(3/2),x, algorithm="giac")`

output `-arcsin(x/a)*sgn(a) + x/sqrt(a^2 - x^2)`

3.136.9 Mupad [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{(a^2 - x^2)^{3/2}} dx = \frac{x}{\sqrt{a^2 - x^2}} + \ln \left(\sqrt{a^2 - x^2} + x \operatorname{li} \right) \operatorname{li}$$

input `int(x^2/(a^2 - x^2)^(3/2),x)`

output `log(x*1i + (a^2 - x^2)^(1/2))*1i + x/(a^2 - x^2)^(1/2)`

3.137 $\int \frac{x^2}{\sqrt{5-x^2}} dx$

3.137.1 Optimal result	762
3.137.2 Mathematica [A] (verified)	762
3.137.3 Rubi [A] (verified)	763
3.137.4 Maple [A] (verified)	764
3.137.5 Fricas [A] (verification not implemented)	764
3.137.6 Sympy [A] (verification not implemented)	765
3.137.7 Maxima [A] (verification not implemented)	765
3.137.8 Giac [A] (verification not implemented)	765
3.137.9 Mupad [B] (verification not implemented)	766

3.137.1 Optimal result

Integrand size = 15, antiderivative size = 29

$$\int \frac{x^2}{\sqrt{5-x^2}} dx = -\frac{1}{2}x\sqrt{5-x^2} + \frac{5}{2} \arcsin\left(\frac{x}{\sqrt{5}}\right)$$

output `5/2*arcsin(1/5*x*5^(1/2))-1/2*x*(-x^2+5)^(1/2)`

3.137.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.48

$$\int \frac{x^2}{\sqrt{5-x^2}} dx = -\frac{1}{2}x\sqrt{5-x^2} - 5 \arctan\left(\frac{x}{\sqrt{5}-\sqrt{5-x^2}}\right)$$

input `Integrate[x^2/Sqrt[5 - x^2],x]`

output `-1/2*(x*Sqrt[5 - x^2]) - 5*ArcTan[x/(Sqrt[5] - Sqrt[5 - x^2])]`

3.137.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {262, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{\sqrt{5-x^2}} dx$$

↓ 262

$$\frac{5}{2} \int \frac{1}{\sqrt{5-x^2}} dx - \frac{1}{2} x \sqrt{5-x^2}$$

↓ 223

$$\frac{5}{2} \arcsin\left(\frac{x}{\sqrt{5}}\right) - \frac{1}{2} x \sqrt{5-x^2}$$

input `Int[x^2/Sqrt[5 - x^2],x]`

output `-1/2*(x*Sqrt[5 - x^2]) + (5*ArcSin[x/Sqrt[5]])/2`

3.137.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 262 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m-1)*((a + b*x^2)^(p+1)/(b*(m+2*p+1))), x] - Simp[a*c^2*((m-1)/(b*(m+2*p+1)) Int[(c*x)^(m-2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2-1] && NeQ[m+2*p+1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

3.137.4 Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{5 \arcsin\left(\frac{x\sqrt{5}}{5}\right)}{2} - \frac{x\sqrt{-x^2+5}}{2}$	23
risch	$\frac{x(x^2-5)}{2\sqrt{-x^2+5}} + \frac{5 \arcsin\left(\frac{x\sqrt{5}}{5}\right)}{2}$	28
pseudoelliptic	$-\frac{x\sqrt{-x^2+5}}{2} - \frac{5 \arctan\left(\frac{\sqrt{-x^2+5}}{x}\right)}{2}$	30
meijerg	$\frac{5i \left(\frac{i\sqrt{\pi} x\sqrt{5} \sqrt{-\frac{x^2}{5}+1}}{5} - i\sqrt{\pi} \arcsin\left(\frac{x\sqrt{5}}{5}\right) \right)}{2\sqrt{\pi}}$	40
trager	$-\frac{x\sqrt{-x^2+5}}{2} - \frac{5 \operatorname{RootOf}\left(-Z^2+1\right) \ln\left(-\operatorname{RootOf}\left(-Z^2+1\right)\sqrt{-x^2+5}+x\right)}{2}$	42

input `int(x^2/(-x^2+5)^(1/2),x,method=_RETURNVERBOSE)`output `5/2*arcsin(1/5*x*5^(1/2))-1/2*x*(-x^2+5)^(1/2)`**3.137.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{\sqrt{5-x^2}} dx = -\frac{1}{2} \sqrt{-x^2+5}x - \frac{5}{2} \arctan\left(\frac{\sqrt{-x^2+5}}{x}\right)$$

input `integrate(x^2/(-x^2+5)^(1/2),x, algorithm="fricas")`output `-1/2*sqrt(-x^2 + 5)*x - 5/2*arctan(sqrt(-x^2 + 5)/x)`

3.137.6 Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

$$\int \frac{x^2}{\sqrt{5-x^2}} dx = -\frac{x\sqrt{5-x^2}}{2} + \frac{5 \operatorname{asin}\left(\frac{\sqrt{5}x}{5}\right)}{2}$$

input `integrate(x**2/(-x**2+5)**(1/2),x)`output `-x*sqrt(5 - x**2)/2 + 5*asin(sqrt(5)*x/5)/2`**3.137.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

$$\int \frac{x^2}{\sqrt{5-x^2}} dx = -\frac{1}{2} \sqrt{-x^2+5}x + \frac{5}{2} \arcsin\left(\frac{1}{5} \sqrt{5}x\right)$$

input `integrate(x^2/(-x^2+5)^(1/2),x, algorithm="maxima")`output `-1/2*sqrt(-x^2 + 5)*x + 5/2*arcsin(1/5*sqrt(5)*x)`**3.137.8 Giac [A] (verification not implemented)**

Time = 0.40 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

$$\int \frac{x^2}{\sqrt{5-x^2}} dx = -\frac{1}{2} \sqrt{-x^2+5}x + \frac{5}{2} \arcsin\left(\frac{1}{5} \sqrt{5}x\right)$$

input `integrate(x^2/(-x^2+5)^(1/2),x, algorithm="giac")`output `-1/2*sqrt(-x^2 + 5)*x + 5/2*arcsin(1/5*sqrt(5)*x)`

3.137.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

$$\int \frac{x^2}{\sqrt{5-x^2}} dx = \frac{5 \operatorname{asin}\left(\frac{\sqrt{5}x}{5}\right)}{2} - \frac{x\sqrt{5-x^2}}{2}$$

input `int(x^2/(5 - x^2)^(1/2),x)`

output `(5*asin((5^(1/2)*x)/5))/2 - (x*(5 - x^2)^(1/2))/2`

3.138 $\int \frac{1}{x\sqrt{3+x^2}} dx$

3.138.1 Optimal result	767
3.138.2 Mathematica [A] (verified)	767
3.138.3 Rubi [A] (verified)	768
3.138.4 Maple [A] (verified)	769
3.138.5 Fricas [A] (verification not implemented)	769
3.138.6 Sympy [A] (verification not implemented)	770
3.138.7 Maxima [A] (verification not implemented)	770
3.138.8 Giac [B] (verification not implemented)	770
3.138.9 Mupad [B] (verification not implemented)	771

3.138.1 Optimal result

Integrand size = 13, antiderivative size = 23

$$\int \frac{1}{x\sqrt{3+x^2}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{3+x^2}}{\sqrt{3}}\right)}{\sqrt{3}}$$

output `-1/3*arctanh(1/3*(x^2+3)^(1/2)*3^(1/2))*3^(1/2)`

3.138.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{3+x^2}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{3+x^2}}{\sqrt{3}}\right)}{\sqrt{3}}$$

input `Integrate[1/(x*Sqrt[3 + x^2]),x]`

output `-(ArcTanh[Sqrt[3 + x^2]/Sqrt[3]]/Sqrt[3])`

3.138.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {243, 73, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x\sqrt{x^2+3}} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int \frac{1}{x^2\sqrt{x^2+3}} dx^2 \\ & \quad \downarrow \text{73} \\ & \int \frac{1}{x^4-3} d\sqrt{x^2+3} \\ & \quad \downarrow \text{220} \\ & -\frac{\operatorname{arctanh}\left(\frac{\sqrt{x^2+3}}{\sqrt{3}}\right)}{\sqrt{3}} \end{aligned}$$

input `Int[1/(x*Sqrt[3 + x^2]),x]`

output `-(ArcTanh[Sqrt[3 + x^2]/Sqrt[3]]/Sqrt[3])`

3.138.3.1 Defintions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

```
rule 243 Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int
t[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I
ntegerQ[(m - 1)/2]
```

3.138.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

method	result	size
default	$-\frac{\sqrt{3} \operatorname{arctanh}\left(\frac{\sqrt{3}}{\sqrt{x^2+3}}\right)}{3}$	18
pseudoelliptic	$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{x^2+3}\sqrt{3}}{3}\right)\sqrt{3}}{3}$	19
trager	$\frac{\operatorname{RootOf}\left(-Z^2-3\right) \ln\left(\frac{-\operatorname{RootOf}\left(-Z^2-3\right)+\sqrt{x^2+3}}{x}\right)}{3}$	30
meijerg	$\frac{\sqrt{3}\left(-2\ln(2)+2\ln(x)-\ln(3)\right)\sqrt{\pi}-2\sqrt{\pi}\ln\left(\frac{1}{2}+\frac{\sqrt{\frac{x^2}{3}+1}}{2}\right)}{6\sqrt{\pi}}$	46

```
input int(1/x/(x^2+3)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/3*3^(1/2)*arctanh(3^(1/2)/(x^2+3)^(1/2))
```

3.138.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.04

$$\int \frac{1}{x\sqrt{3+x^2}} dx = \frac{1}{3} \sqrt{3} \log\left(-\frac{\sqrt{3}-\sqrt{x^2+3}}{x}\right)$$

```
input integrate(1/x/(x^2+3)^(1/2),x, algorithm="fricas")
```

```
output 1/3*sqrt(3)*log(-(sqrt(3) - sqrt(x^2 + 3))/x)
```

3.138.6 Sympy [A] (verification not implemented)

Time = 0.54 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int \frac{1}{x\sqrt{3+x^2}} dx = -\frac{\sqrt{3} \operatorname{asinh}\left(\frac{\sqrt{3}}{x}\right)}{3}$$

input `integrate(1/x/(x**2+3)**(1/2),x)`

output `-sqrt(3)*asinh(sqrt(3)/x)/3`

3.138.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.61

$$\int \frac{1}{x\sqrt{3+x^2}} dx = -\frac{1}{3} \sqrt{3} \operatorname{arsinh}\left(\frac{\sqrt{3}}{|x|}\right)$$

input `integrate(1/x/(x^2+3)^(1/2),x, algorithm="maxima")`

output `-1/3*sqrt(3)*arcsinh(sqrt(3)/abs(x))`

3.138.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. 2(18) = 36.

Time = 0.27 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.61

$$\int \frac{1}{x\sqrt{3+x^2}} dx = -\frac{1}{6} \sqrt{3} \log\left(\sqrt{3} + \sqrt{x^2+3}\right) + \frac{1}{6} \sqrt{3} \log\left(-\sqrt{3} + \sqrt{x^2+3}\right)$$

input `integrate(1/x/(x^2+3)^(1/2),x, algorithm="giac")`

output `-1/6*sqrt(3)*log(sqrt(3) + sqrt(x^2 + 3)) + 1/6*sqrt(3)*log(-sqrt(3) + sqrt(x^2 + 3))`

3.138.9 Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int \frac{1}{x\sqrt{3+x^2}} dx = -\frac{\sqrt{3} \operatorname{atanh}\left(\frac{\sqrt{3}\sqrt{x^2+3}}{3}\right)}{3}$$

input `int(1/(x*(x^2 + 3)^(1/2)),x)`output `-(3^(1/2)*atanh((3^(1/2)*(x^2 + 3)^(1/2))/3))/3`

$$\mathbf{3.139} \quad \int \frac{x}{(4+x^2)^{5/2}} dx$$

3.139.1 Optimal result	772
3.139.2 Mathematica [A] (verified)	772
3.139.3 Rubi [A] (verified)	773
3.139.4 Maple [A] (verified)	774
3.139.5 Fricas [B] (verification not implemented)	774
3.139.6 Sympy [B] (verification not implemented)	775
3.139.7 Maxima [A] (verification not implemented)	775
3.139.8 Giac [A] (verification not implemented)	775
3.139.9 Mupad [B] (verification not implemented)	776

3.139.1 Optimal result

Integrand size = 11, antiderivative size = 13

$$\int \frac{x}{(4+x^2)^{5/2}} dx = -\frac{1}{3(4+x^2)^{3/2}}$$

output `-1/3/(x^2+4)^(3/2)`

3.139.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{x}{(4+x^2)^{5/2}} dx = -\frac{1}{3(4+x^2)^{3/2}}$$

input `Integrate[x/(4 + x^2)^(5/2), x]`

output `-1/3*1/(4 + x^2)^(3/2)`

3.139.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(x^2 + 4)^{5/2}} dx$$

↓ 241

$$-\frac{1}{3(x^2 + 4)^{3/2}}$$

input `Int[x/(4 + x^2)^(5/2),x]`

output `-1/3*1/(4 + x^2)^(3/2)`

3.139.3.1 Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

3.139.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

method	result	size
gospers	$-\frac{1}{3(x^2+4)^{\frac{3}{2}}}$	10
derivativedivides	$-\frac{1}{3(x^2+4)^{\frac{3}{2}}}$	10
default	$-\frac{1}{3(x^2+4)^{\frac{3}{2}}}$	10
trager	$-\frac{1}{3(x^2+4)^{\frac{3}{2}}}$	10
risch	$-\frac{1}{3(x^2+4)^{\frac{3}{2}}}$	10
pseudoelliptic	$-\frac{1}{3(x^2+4)^{\frac{3}{2}}}$	10
meijerg	$\frac{\frac{\sqrt{\pi}}{2} - \frac{\sqrt{\pi}}{2\left(1+\frac{x^2}{4}\right)^{\frac{3}{2}}}}{12\sqrt{\pi}}$	26

input `int(x/(x^2+4)^(5/2),x,method=_RETURNVERBOSE)`

output `-1/3/(x^2+4)^(3/2)`

3.139.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. 2(9) = 18.

Time = 0.23 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.62

$$\int \frac{x}{(4+x^2)^{5/2}} dx = -\frac{\sqrt{x^2+4}}{3(x^4+8x^2+16)}$$

input `integrate(x/(x^2+4)^(5/2),x, algorithm="fracas")`

output `-1/3*sqrt(x^2 + 4)/(x^4 + 8*x^2 + 16)`

3.139.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. $2(12) = 24$.

Time = 0.40 (sec) , antiderivative size = 26, normalized size of antiderivative = 2.00

$$\int \frac{x}{(4+x^2)^{5/2}} dx = -\frac{1}{3x^2\sqrt{x^2+4} + 12\sqrt{x^2+4}}$$

input `integrate(x/(x**2+4)**(5/2),x)`

output `-1/(3*x**2*sqrt(x**2 + 4) + 12*sqrt(x**2 + 4))`

3.139.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \frac{x}{(4+x^2)^{5/2}} dx = -\frac{1}{3(x^2+4)^{3/2}}$$

input `integrate(x/(x^2+4)^(5/2),x, algorithm="maxima")`

output `-1/3/(x^2 + 4)^(3/2)`

3.139.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \frac{x}{(4+x^2)^{5/2}} dx = -\frac{1}{3(x^2+4)^{3/2}}$$

input `integrate(x/(x^2+4)^(5/2),x, algorithm="giac")`

output `-1/3/(x^2 + 4)^(3/2)`

3.139.9 Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \frac{x}{(4+x^2)^{5/2}} dx = -\frac{1}{3(x^2+4)^{3/2}}$$

input `int(x/(x^2 + 4)^(5/2),x)`

output `-1/(3*(x^2 + 4)^(3/2))`

3.140 $\int x^3 \sqrt{4 - 9x^2} dx$

3.140.1 Optimal result	777
3.140.2 Mathematica [A] (verified)	777
3.140.3 Rubi [A] (verified)	778
3.140.4 Maple [A] (verified)	779
3.140.5 Fricas [A] (verification not implemented)	779
3.140.6 Sympy [A] (verification not implemented)	780
3.140.7 Maxima [A] (verification not implemented)	780
3.140.8 Giac [A] (verification not implemented)	780
3.140.9 Mupad [B] (verification not implemented)	781

3.140.1 Optimal result

Integrand size = 15, antiderivative size = 31

$$\int x^3 \sqrt{4 - 9x^2} dx = -\frac{4}{243} (4 - 9x^2)^{3/2} + \frac{1}{405} (4 - 9x^2)^{5/2}$$

output `-4/243*(-9*x^2+4)^(3/2)+1/405*(-9*x^2+4)^(5/2)`

3.140.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.71

$$\int x^3 \sqrt{4 - 9x^2} dx = \frac{(-8 - 27x^2)(4 - 9x^2)^{3/2}}{1215}$$

input `Integrate[x^3*Sqrt[4 - 9*x^2],x]`

output `((-8 - 27*x^2)*(4 - 9*x^2)^(3/2))/1215`

3.140.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.13, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^3 \sqrt{4 - 9x^2} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int x^2 \sqrt{4 - 9x^2} dx^2 \\ & \quad \downarrow \text{53} \\ & \frac{1}{2} \int \left(\frac{4}{9} \sqrt{4 - 9x^2} - \frac{1}{9} (4 - 9x^2)^{3/2} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \left(\frac{2}{405} (4 - 9x^2)^{5/2} - \frac{8}{243} (4 - 9x^2)^{3/2} \right) \end{aligned}$$

input `Int[x^3*Sqrt[4 - 9*x^2],x]`

output `((-8*(4 - 9*x^2)^(3/2))/243 + (2*(4 - 9*x^2)^(5/2))/405)/2`

3.140.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.140.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.61

method	result	size
pseudoelliptic	$-\frac{(27x^2+8)(-9x^2+4)^{\frac{3}{2}}}{1215}$	19
trager	$\left(\frac{1}{5}x^4 - \frac{4}{135}x^2 - \frac{32}{1215}\right)\sqrt{-9x^2+4}$	23
default	$-\frac{x^2(-9x^2+4)^{\frac{3}{2}}}{45} - \frac{8(-9x^2+4)^{\frac{3}{2}}}{1215}$	27
gospers	$\frac{(-2+3x)(2+3x)(27x^2+8)\sqrt{-9x^2+4}}{1215}$	29
risch	$-\frac{(243x^4-36x^2-32)(9x^2-4)}{1215\sqrt{-9x^2+4}}$	31
meijerg	$-\frac{8\left(-\frac{8\sqrt{\pi}}{15} + \frac{4\sqrt{\pi}\left(1-\frac{9x^2}{4}\right)^{\frac{3}{2}}\left(\frac{27x^2}{4}+2\right)}{15}\right)}{81\sqrt{\pi}}$	33

input `int(x^3*(-9*x^2+4)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/1215*(27*x^2+8)*(-9*x^2+4)^(3/2)`

3.140.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int x^3\sqrt{4-9x^2} dx = \frac{1}{1215} (243x^4 - 36x^2 - 32)\sqrt{-9x^2+4}$$

input `integrate(x^3*(-9*x^2+4)^(1/2),x, algorithm="fracas")`

output `1/1215*(243*x^4 - 36*x^2 - 32)*sqrt(-9*x^2 + 4)`

3.140.6 Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.42

$$\int x^3 \sqrt{4-9x^2} dx = \frac{x^4 \sqrt{4-9x^2}}{5} - \frac{4x^2 \sqrt{4-9x^2}}{135} - \frac{32 \sqrt{4-9x^2}}{1215}$$

input `integrate(x**3*(-9*x**2+4)**(1/2),x)`output `x**4*sqrt(4 - 9*x**2)/5 - 4*x**2*sqrt(4 - 9*x**2)/135 - 32*sqrt(4 - 9*x**2)/1215`**3.140.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int x^3 \sqrt{4-9x^2} dx = -\frac{1}{45} (-9x^2 + 4)^{\frac{3}{2}} x^2 - \frac{8}{1215} (-9x^2 + 4)^{\frac{3}{2}}$$

input `integrate(x^3*(-9*x^2+4)^(1/2),x, algorithm="maxima")`output `-1/45*(-9*x^2 + 4)^(3/2)*x^2 - 8/1215*(-9*x^2 + 4)^(3/2)`**3.140.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.03

$$\int x^3 \sqrt{4-9x^2} dx = \frac{1}{405} (9x^2 - 4)^2 \sqrt{-9x^2 + 4} - \frac{4}{243} (-9x^2 + 4)^{\frac{3}{2}}$$

input `integrate(x^3*(-9*x^2+4)^(1/2),x, algorithm="giac")`output `1/405*(9*x^2 - 4)^2*sqrt(-9*x^2 + 4) - 4/243*(-9*x^2 + 4)^(3/2)`

3.140.9 Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int x^3 \sqrt{4 - 9x^2} dx = -\frac{\sqrt{\frac{4}{9} - x^2} \left(-\frac{9x^4}{5} + \frac{4x^2}{15} + \frac{32}{135} \right)}{3}$$

input `int(x^3*(4 - 9*x^2)^(1/2),x)`

output `-((4/9 - x^2)^(1/2)*((4*x^2)/15 - (9*x^4)/5 + 32/135))/3`

3.141 $\int x^2\sqrt{9-x^2} dx$

3.141.1 Optimal result	782
3.141.2 Mathematica [A] (verified)	782
3.141.3 Rubi [A] (verified)	783
3.141.4 Maple [A] (verified)	784
3.141.5 Fricas [A] (verification not implemented)	784
3.141.6 Sympy [C] (verification not implemented)	785
3.141.7 Maxima [A] (verification not implemented)	785
3.141.8 Giac [A] (verification not implemented)	785
3.141.9 Mupad [B] (verification not implemented)	786

3.141.1 Optimal result

Integrand size = 15, antiderivative size = 45

$$\int x^2\sqrt{9-x^2} dx = -\frac{9}{8}x\sqrt{9-x^2} + \frac{1}{4}x^3\sqrt{9-x^2} + \frac{81}{8} \arcsin\left(\frac{x}{3}\right)$$

output `81/8*arcsin(1/3*x)-9/8*x*(-x^2+9)^(1/2)+1/4*x^3*(-x^2+9)^(1/2)`

3.141.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.02

$$\int x^2\sqrt{9-x^2} dx = \frac{1}{8}x\sqrt{9-x^2}(-9+2x^2) - \frac{81}{4} \arctan\left(\frac{\sqrt{9-x^2}}{3+x}\right)$$

input `Integrate[x^2*Sqrt[9-x^2],x]`

output `(x*Sqrt[9-x^2]*(-9+2*x^2))/8 - (81*ArcTan[Sqrt[9-x^2]/(3+x)]/4`

3.141.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.11, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {248, 262, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \sqrt{9 - x^2} dx$$

$$\downarrow \text{248}$$

$$\frac{9}{4} \int \frac{x^2}{\sqrt{9 - x^2}} dx + \frac{1}{4} \sqrt{9 - x^2} x^3$$

$$\downarrow \text{262}$$

$$\frac{9}{4} \left(\frac{9}{2} \int \frac{1}{\sqrt{9 - x^2}} dx - \frac{1}{2} x \sqrt{9 - x^2} \right) + \frac{1}{4} \sqrt{9 - x^2} x^3$$

$$\downarrow \text{223}$$

$$\frac{9}{4} \left(\frac{9}{2} \arcsin \left(\frac{x}{3} \right) - \frac{1}{2} x \sqrt{9 - x^2} \right) + \frac{1}{4} \sqrt{9 - x^2} x^3$$

input `Int[x^2*Sqrt[9 - x^2],x]`

output `(x^3*Sqrt[9 - x^2])/4 + (9*(-1/2*(x*Sqrt[9 - x^2]) + (9*ArcSin[x/3])/2))/4`

3.141.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 248 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^p/(c*(m + 2*p + 1))), x] + Simp[2*a*(p/(m + 2*p + 1)) Int[(c*x)^m*(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c, m}, x] && GtQ[p, 0] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 262 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

3.141.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.71

method	result	size
default	$-\frac{x(-x^2+9)^{\frac{3}{2}}}{4} + \frac{9x\sqrt{-x^2+9}}{8} + \frac{81 \arcsin\left(\frac{x}{3}\right)}{8}$	32
risch	$-\frac{x(2x^2-9)(x^2-9)}{8\sqrt{-x^2+9}} + \frac{81 \arcsin\left(\frac{x}{3}\right)}{8}$	32
pseudoelliptic	$-\frac{81 \arctan\left(\frac{\sqrt{-x^2+9}}{x}\right)}{8} + \frac{(2x^3-9x)\sqrt{-x^2+9}}{8}$	38
meijerg	$81i \left(-\frac{i\sqrt{\pi}x\left(-\frac{2x^2}{3}+3\right)\sqrt{-\frac{x^2}{9}+1}}{18} + \frac{i\sqrt{\pi} \arcsin\left(\frac{x}{3}\right)}{2} \right)$	41
trager	$\frac{x(2x^2-9)\sqrt{-x^2+9}}{8} + \frac{81 \operatorname{RootOf}(_Z^2+1) \ln(\operatorname{RootOf}(_Z^2+1)\sqrt{-x^2+9}+x)}{8}$	48

input `int(x^2*(-x^2+9)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/4*x*(-x^2+9)^(3/2)+9/8*x*(-x^2+9)^(1/2)+81/8*arcsin(1/3*x)`

3.141.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.87

$$\int x^2 \sqrt{9 - x^2} dx = \frac{1}{8} (2x^3 - 9x) \sqrt{-x^2 + 9} - \frac{81}{4} \arctan\left(\frac{\sqrt{-x^2 + 9} - 3}{x}\right)$$

input `integrate(x^2*(-x^2+9)^(1/2),x, algorithm="fricas")`

output `1/8*(2*x^3 - 9*x)*sqrt(-x^2 + 9) - 81/4*arctan((sqrt(-x^2 + 9) - 3)/x)`

3.141.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.75 (sec) , antiderivative size = 110, normalized size of antiderivative = 2.44

$$\int x^2 \sqrt{9 - x^2} dx = \begin{cases} \frac{ix^5}{4\sqrt{x^2-9}} - \frac{27ix^3}{8\sqrt{x^2-9}} + \frac{81ix}{8\sqrt{x^2-9}} - \frac{81i \operatorname{acosh}\left(\frac{x}{3}\right)}{8} & \text{for } |x^2| > 9 \\ -\frac{x^5}{4\sqrt{9-x^2}} + \frac{27x^3}{8\sqrt{9-x^2}} - \frac{81x}{8\sqrt{9-x^2}} + \frac{81 \operatorname{asin}\left(\frac{x}{3}\right)}{8} & \text{otherwise} \end{cases}$$

input `integrate(x**2*(-x**2+9)**(1/2),x)`

output `Piecewise((I*x**5/(4*sqrt(x**2 - 9)) - 27*I*x**3/(8*sqrt(x**2 - 9)) + 81*I*x/(8*sqrt(x**2 - 9)) - 81*I*acosh(x/3)/8, Abs(x**2) > 9), (-x**5/(4*sqrt(9 - x**2)) + 27*x**3/(8*sqrt(9 - x**2)) - 81*x/(8*sqrt(9 - x**2)) + 81*asin(x/3)/8, True))`

3.141.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.69

$$\int x^2 \sqrt{9 - x^2} dx = -\frac{1}{4} (-x^2 + 9)^{\frac{3}{2}} x + \frac{9}{8} \sqrt{-x^2 + 9} x + \frac{81}{8} \arcsin\left(\frac{1}{3} x\right)$$

input `integrate(x^2*(-x^2+9)^(1/2),x, algorithm="maxima")`

output `-1/4*(-x^2 + 9)^(3/2)*x + 9/8*sqrt(-x^2 + 9)*x + 81/8*arcsin(1/3*x)`

3.141.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.58

$$\int x^2 \sqrt{9 - x^2} dx = \frac{1}{8} (2x^2 - 9) \sqrt{-x^2 + 9} x + \frac{81}{8} \arcsin\left(\frac{1}{3} x\right)$$

input `integrate(x^2*(-x^2+9)^(1/2),x, algorithm="giac")`

output `1/8*(2*x^2 - 9)*sqrt(-x^2 + 9)*x + 81/8*arcsin(1/3*x)`

3.141.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.60

$$\int x^2 \sqrt{9 - x^2} dx = \frac{81 \operatorname{asin}\left(\frac{x}{3}\right)}{8} - \sqrt{9 - x^2} \left(\frac{9x}{8} - \frac{x^3}{4} \right)$$

input `int(x^2*(9 - x^2)^(1/2),x)`

output `(81*asin(x/3))/8 - (9 - x^2)^(1/2)*((9*x)/8 - x^3/4)`

3.142 $\int 5x\sqrt{1+x^2} dx$

3.142.1 Optimal result	787
3.142.2 Mathematica [A] (verified)	787
3.142.3 Rubi [A] (verified)	788
3.142.4 Maple [A] (verified)	789
3.142.5 Fricas [A] (verification not implemented)	789
3.142.6 Sympy [B] (verification not implemented)	790
3.142.7 Maxima [A] (verification not implemented)	790
3.142.8 Giac [A] (verification not implemented)	790
3.142.9 Mupad [B] (verification not implemented)	791

3.142.1 Optimal result

Integrand size = 12, antiderivative size = 13

$$\int 5x\sqrt{1+x^2} dx = \frac{5}{3}(1+x^2)^{3/2}$$

output `5/3*(x^2+1)^(3/2)`

3.142.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int 5x\sqrt{1+x^2} dx = \frac{5}{3}(1+x^2)^{3/2}$$

input `Integrate[5*x*Sqrt[1 + x^2],x]`

output `(5*(1 + x^2)^(3/2))/3`

3.142.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {27, 241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int 5x\sqrt{x^2+1} dx \\ \downarrow 27 \\ 5 \int x\sqrt{x^2+1} dx \\ \downarrow 241 \\ \frac{5}{3}(x^2+1)^{3/2} \end{array}$$

input `Int[5*x*Sqrt[1 + x^2],x]`

output `(5*(1 + x^2)^(3/2))/3`

3.142.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

3.142.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

method	result	size
gospers	$\frac{5(x^2+1)^{\frac{3}{2}}}{3}$	10
derivativdivides	$\frac{5(x^2+1)^{\frac{3}{2}}}{3}$	10
default	$\frac{5(x^2+1)^{\frac{3}{2}}}{3}$	10
risch	$\frac{5(x^2+1)^{\frac{3}{2}}}{3}$	10
pseudoelliptic	$\frac{5(x^2+1)^{\frac{3}{2}}}{3}$	10
trager	$5\left(\frac{x^2}{3} + \frac{1}{3}\right)\sqrt{x^2+1}$	17
meijerg	$-\frac{5\left(\frac{4\sqrt{\pi}}{3} - \frac{2\sqrt{\pi}(2x^2+2)\sqrt{x^2+1}}{3}\right)}{4\sqrt{\pi}}$	31

input `int(5*x*(x^2+1)^(1/2),x,method=_RETURNVERBOSE)`output `5/3*(x^2+1)^(3/2)`**3.142.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int 5x\sqrt{1+x^2} dx = \frac{5}{3}(x^2+1)^{\frac{3}{2}}$$

input `integrate(5*x*(x^2+1)^(1/2),x, algorithm="fracas")`output `5/3*(x^2 + 1)^(3/2)`

3.142.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. $2(10) = 20$.

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 2.00

$$\int 5x\sqrt{1+x^2} dx = \frac{5x^2\sqrt{x^2+1}}{3} + \frac{5\sqrt{x^2+1}}{3}$$

input `integrate(5*x*(x**2+1)**(1/2),x)`

output `5*x**2*sqrt(x**2 + 1)/3 + 5*sqrt(x**2 + 1)/3`

3.142.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int 5x\sqrt{1+x^2} dx = \frac{5}{3} (x^2 + 1)^{\frac{3}{2}}$$

input `integrate(5*x*(x^2+1)^(1/2),x, algorithm="maxima")`

output `5/3*(x^2 + 1)^(3/2)`

3.142.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int 5x\sqrt{1+x^2} dx = \frac{5}{3} (x^2 + 1)^{\frac{3}{2}}$$

input `integrate(5*x*(x^2+1)^(1/2),x, algorithm="giac")`

output `5/3*(x^2 + 1)^(3/2)`

3.142.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int 5x\sqrt{1+x^2} dx = \frac{5(x^2+1)^{3/2}}{3}$$

input `int(5*x*(x^2 + 1)^(1/2),x)`

output `(5*(x^2 + 1)^(3/2))/3`

3.143 $\int \frac{1}{(-25+4x^2)^{3/2}} dx$

3.143.1 Optimal result	792
3.143.2 Mathematica [A] (verified)	792
3.143.3 Rubi [A] (verified)	793
3.143.4 Maple [A] (verified)	793
3.143.5 Fricas [B] (verification not implemented)	794
3.143.6 Sympy [C] (verification not implemented)	794
3.143.7 Maxima [A] (verification not implemented)	795
3.143.8 Giac [A] (verification not implemented)	795
3.143.9 Mupad [B] (verification not implemented)	795

3.143.1 Optimal result

Integrand size = 11, antiderivative size = 16

$$\int \frac{1}{(-25 + 4x^2)^{3/2}} dx = -\frac{x}{25\sqrt{-25 + 4x^2}}$$

output -1/25*x/(4*x^2-25)^(1/2)

3.143.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{1}{(-25 + 4x^2)^{3/2}} dx = -\frac{x}{25\sqrt{-25 + 4x^2}}$$

input Integrate[(-25 + 4*x^2)^(-3/2),x]

output -1/25*x/Sqrt[-25 + 4*x^2]

3.143.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(4x^2 - 25)^{3/2}} dx$$

↓ 208

$$-\frac{x}{25\sqrt{4x^2 - 25}}$$

input `Int[(-25 + 4*x^2)^(-3/2),x]`

output `-1/25*x/Sqrt[-25 + 4*x^2]`

3.143.3.1 Defintions of rubi rules used

rule 208 `Int[((a_) + (b_.)*(x_)^2)^(-3/2), x_Symbol] :> Simp[x/(a*Sqrt[a + b*x^2]), x] /; FreeQ[{a, b}, x]`

3.143.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

method	result	size
default	$-\frac{x}{25\sqrt{4x^2-25}}$	13
trager	$-\frac{x}{25\sqrt{4x^2-25}}$	13
risch	$-\frac{x}{25\sqrt{4x^2-25}}$	13
pseudoelliptic	$-\frac{x}{25\sqrt{4x^2-25}}$	13
gosper	$-\frac{(2x-5)(5+2x)x}{25(4x^2-25)^{3/2}}$	23
meijerg	$\frac{\left(-\text{signum}\left(-1+\frac{4x^2}{25}\right)\right)^{3/2} x}{125 \text{signum}\left(-1+\frac{4x^2}{25}\right)^{3/2} \sqrt{1-\frac{4x^2}{25}}}$	35

input `int(1/(4*x^2-25)^(3/2),x,method=_RETURNVERBOSE)`

output `-1/25*x/(4*x^2-25)^(1/2)`

3.143.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 30 vs. $2(12) = 24$.

Time = 0.23 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.88

$$\int \frac{1}{(-25 + 4x^2)^{3/2}} dx = -\frac{4x^2 + 2\sqrt{4x^2 - 25}x - 25}{50(4x^2 - 25)}$$

input `integrate(1/(4*x^2-25)^(3/2),x, algorithm="fricas")`

output `-1/50*(4*x^2 + 2*sqrt(4*x^2 - 25)*x - 25)/(4*x^2 - 25)`

3.143.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.46 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.12

$$\int \frac{1}{(-25 + 4x^2)^{3/2}} dx = \begin{cases} -\frac{x}{25\sqrt{4x^2-25}} & \text{for } |x^2| > \frac{25}{4} \\ \frac{ix}{25\sqrt{25-4x^2}} & \text{otherwise} \end{cases}$$

input `integrate(1/(4*x**2-25)**(3/2),x)`

output `Piecewise((-x/(25*sqrt(4*x**2 - 25))), Abs(x**2) > 25/4, (I*x/(25*sqrt(25 - 4*x**2))), True))`

3.143.7 Maxima [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{(-25 + 4x^2)^{3/2}} dx = -\frac{x}{25\sqrt{4x^2 - 25}}$$

input `integrate(1/(4*x^2-25)^(3/2),x, algorithm="maxima")`output `-1/25*x/sqrt(4*x^2 - 25)`**3.143.8 Giac [A] (verification not implemented)**

Time = 0.39 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{(-25 + 4x^2)^{3/2}} dx = -\frac{x}{25\sqrt{4x^2 - 25}}$$

input `integrate(1/(4*x^2-25)^(3/2),x, algorithm="giac")`output `-1/25*x/sqrt(4*x^2 - 25)`**3.143.9 Mupad [B] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{(-25 + 4x^2)^{3/2}} dx = -\frac{x}{25\sqrt{4x^2 - 25}}$$

input `int(1/(4*x^2 - 25)^(3/2),x)`output `-x/(25*(4*x^2 - 25)^(1/2))`

3.144 $\int \sqrt{2x - x^2} dx$

3.144.1 Optimal result	796
3.144.2 Mathematica [A] (verified)	796
3.144.3 Rubi [A] (verified)	797
3.144.4 Maple [A] (verified)	798
3.144.5 Fricas [A] (verification not implemented)	798
3.144.6 Sympy [A] (verification not implemented)	799
3.144.7 Maxima [A] (verification not implemented)	799
3.144.8 Giac [A] (verification not implemented)	799
3.144.9 Mupad [B] (verification not implemented)	800

3.144.1 Optimal result

Integrand size = 13, antiderivative size = 33

$$\int \sqrt{2x - x^2} dx = -\frac{1}{2}(1 - x)\sqrt{2x - x^2} - \frac{1}{2} \arcsin(1 - x)$$

output `1/2*arcsin(-1+x)-1/2*(1-x)*(-x^2+2*x)^(1/2)`

3.144.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.42

$$\int \sqrt{2x - x^2} dx = \frac{1}{2} \sqrt{-((-2 + x)x)} \left(-1 + x + \frac{2 \log(\sqrt{-2 + x} - \sqrt{x})}{\sqrt{-2 + x}\sqrt{x}} \right)$$

input `Integrate[Sqrt[2*x - x^2],x]`

output `(Sqrt[-((-2 + x)*x)]*(-1 + x + (2*Log[Sqrt[-2 + x] - Sqrt[x]])/(Sqrt[-2 + x]*Sqrt[x])))/2`

3.144.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.12, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {1087, 1090, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{2x - x^2} dx$$

$$\downarrow 1087$$

$$\frac{1}{2} \int \frac{1}{\sqrt{2x - x^2}} dx - \frac{1}{2}(1 - x)\sqrt{2x - x^2}$$

$$\downarrow 1090$$

$$-\frac{1}{4} \int \frac{1}{\sqrt{1 - \frac{1}{4}(2 - 2x)^2}} d(2 - 2x) - \frac{1}{2}\sqrt{2x - x^2}(1 - x)$$

$$\downarrow 223$$

$$-\frac{1}{2} \arcsin\left(\frac{1}{2}(2 - 2x)\right) - \frac{1}{2}\sqrt{2x - x^2}(1 - x)$$

input `Int[Sqrt[2*x - x^2], x]`

output `-1/2*((1 - x)*Sqrt[2*x - x^2]) - ArcSin[(2 - 2*x)/2]/2`

3.144.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 1087 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] - Simp[p*(b^2 - 4*a*c)/(2*c*(2*p + 1)) Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[3*p])`

rule 1090 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

3.144.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.76

method	result	size
risch	$-\frac{(-1+x)x(-2+x)}{2\sqrt{-x(-2+x)}} + \frac{\arcsin(-1+x)}{2}$	25
default	$-\frac{(-2x+2)\sqrt{-x^2+2x}}{4} + \frac{\arcsin(-1+x)}{2}$	26
pseudoelliptic	$-\arctan\left(\frac{\sqrt{-x(-2+x)}}{x}\right) + \frac{(-1+x)\sqrt{-x(-2+x)}}{2}$	30
meijerg	$-\frac{2i\left(-\frac{i\sqrt{\pi}\sqrt{x}\sqrt{2}(-3x+3)\sqrt{1-\frac{x}{2}}}{12} + \frac{i\sqrt{\pi}\arcsin\left(\frac{\sqrt{2}\sqrt{x}}{2}\right)}{2}\right)}{\sqrt{\pi}}$	47
trager	$\left(-\frac{1}{2} + \frac{x}{2}\right)\sqrt{-x^2+2x} + \frac{\text{RootOf}(_Z^2+1)\ln(\text{RootOf}(_Z^2+1)\sqrt{-x^2+2x+x-1})}{2}$	49

input `int((-x^2+2*x)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2*(-1+x)*x*(-2+x)/(-x*(-2+x))^(1/2)+1/2*arcsin(-1+x)`

3.144.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.06

$$\int \sqrt{2x - x^2} dx = \frac{1}{2} \sqrt{-x^2 + 2x}(x - 1) - \arctan\left(\frac{\sqrt{-x^2 + 2x}}{x}\right)$$

input `integrate((-x^2+2*x)^(1/2),x, algorithm="fricas")`

output `1/2*sqrt(-x^2 + 2*x)*(x - 1) - arctan(sqrt(-x^2 + 2*x)/x)`

3.144.6 Sympy [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.67

$$\int \sqrt{2x - x^2} dx = \left(\frac{x}{2} - \frac{1}{2}\right) \sqrt{-x^2 + 2x} + \frac{\operatorname{asin}(x - 1)}{2}$$

input `integrate((-x**2+2*x)**(1/2),x)`output `(x/2 - 1/2)*sqrt(-x**2 + 2*x) + asin(x - 1)/2`**3.144.7 Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.09

$$\int \sqrt{2x - x^2} dx = \frac{1}{2} \sqrt{-x^2 + 2x} x - \frac{1}{2} \sqrt{-x^2 + 2x} - \frac{1}{2} \arcsin(-x + 1)$$

input `integrate((-x^2+2*x)^(1/2),x, algorithm="maxima")`output `1/2*sqrt(-x^2 + 2*x)*x - 1/2*sqrt(-x^2 + 2*x) - 1/2*arcsin(-x + 1)`**3.144.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.70

$$\int \sqrt{2x - x^2} dx = \frac{1}{2} \sqrt{-x^2 + 2x}(x - 1) + \frac{1}{2} \arcsin(x - 1)$$

input `integrate((-x^2+2*x)^(1/2),x, algorithm="giac")`output `1/2*sqrt(-x^2 + 2*x)*(x - 1) + 1/2*arcsin(x - 1)`

3.144.9 Mupad [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.73

$$\int \sqrt{2x - x^2} dx = \frac{\operatorname{asin}(x - 1)}{2} + \left(\frac{x}{2} - \frac{1}{2}\right) \sqrt{2x - x^2}$$

input `int((2*x - x^2)^(1/2),x)`

output `asin(x - 1)/2 + (x/2 - 1/2)*(2*x - x^2)^(1/2)`

3.145 $\int \frac{1}{\sqrt{8+4x+x^2}} dx$

3.145.1 Optimal result	801
3.145.2 Mathematica [B] (verified)	801
3.145.3 Rubi [A] (verified)	802
3.145.4 Maple [A] (verified)	803
3.145.5 Fricas [B] (verification not implemented)	803
3.145.6 Sympy [A] (verification not implemented)	803
3.145.7 Maxima [A] (verification not implemented)	804
3.145.8 Giac [B] (verification not implemented)	804
3.145.9 Mupad [B] (verification not implemented)	804

3.145.1 Optimal result

Integrand size = 12, antiderivative size = 8

$$\int \frac{1}{\sqrt{8+4x+x^2}} dx = \operatorname{arcsinh}\left(\frac{2+x}{2}\right)$$

output `arcsinh(1+1/2*x)`

3.145.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 20 vs. $2(8) = 16$.

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 2.50

$$\int \frac{1}{\sqrt{8+4x+x^2}} dx = -\log\left(-2-x+\sqrt{8+4x+x^2}\right)$$

input `Integrate[1/Sqrt[8 + 4*x + x^2],x]`

output `-Log[-2 - x + Sqrt[8 + 4*x + x^2]]`

3.145.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.25, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {1090, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x^2 + 4x + 8}} dx$$

↓ 1090

$$\frac{1}{4} \int \frac{1}{\sqrt{\frac{1}{16}(2x + 4)^2 + 1}} d(2x + 4)$$

↓ 222

$$\operatorname{arcsinh}\left(\frac{1}{4}(2x + 4)\right)$$

input `Int[1/Sqrt[8 + 4*x + x^2],x]`

output `ArcSinh[(4 + 2*x)/4]`

3.145.3.1 Defintions of rubi rules used

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

rule 1090 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

3.145.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
default	$\operatorname{arcsinh}\left(1 + \frac{x}{2}\right)$	7
trager	$\ln\left(x + 2 + \sqrt{x^2 + 4x + 8}\right)$	15

input `int(1/(x^2+4*x+8)^(1/2),x,method=_RETURNVERBOSE)`

output `arcsinh(1+1/2*x)`

3.145.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18 vs. 2(6) = 12.

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 2.25

$$\int \frac{1}{\sqrt{8 + 4x + x^2}} dx = -\log\left(-x + \sqrt{x^2 + 4x + 8} - 2\right)$$

input `integrate(1/(x^2+4*x+8)^(1/2),x, algorithm="fricas")`

output `-log(-x + sqrt(x^2 + 4*x + 8) - 2)`

3.145.6 Sympy [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \frac{1}{\sqrt{8 + 4x + x^2}} dx = \operatorname{asinh}\left(\frac{x}{2} + 1\right)$$

input `integrate(1/(x**2+4*x+8)**(1/2),x)`

output `asinh(x/2 + 1)`

3.145.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{8+4x+x^2}} dx = \operatorname{arsinh}\left(\frac{1}{2}x+1\right)$$

input `integrate(1/(x^2+4*x+8)^(1/2),x, algorithm="maxima")`

output `arcsinh(1/2*x + 1)`

3.145.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. 2(6) = 12.

Time = 0.29 (sec) , antiderivative size = 34, normalized size of antiderivative = 4.25

$$\int \frac{1}{\sqrt{8+4x+x^2}} dx = \frac{1}{2} \sqrt{x^2+4x+8}(x+2) - 2 \log(-x + \sqrt{x^2+4x+8} - 2)$$

input `integrate(1/(x^2+4*x+8)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(x^2 + 4*x + 8)*(x + 2) - 2*log(-x + sqrt(x^2 + 4*x + 8) - 2)`

3.145.9 Mupad [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.75

$$\int \frac{1}{\sqrt{8+4x+x^2}} dx = \ln\left(x + \sqrt{x^2+4x+8} + 2\right)$$

input `int(1/(4*x + x^2 + 8)^(1/2),x)`

output `log(x + (4*x + x^2 + 8)^(1/2) + 2)`

3.146 $\int \frac{1}{\sqrt{-8+6x+9x^2}} dx$

3.146.1 Optimal result	805
3.146.2 Mathematica [A] (verified)	805
3.146.3 Rubi [A] (verified)	806
3.146.4 Maple [A] (verified)	807
3.146.5 Fricas [A] (verification not implemented)	807
3.146.6 Sympy [A] (verification not implemented)	807
3.146.7 Maxima [A] (verification not implemented)	808
3.146.8 Giac [A] (verification not implemented)	808
3.146.9 Mupad [B] (verification not implemented)	808

3.146.1 Optimal result

Integrand size = 14, antiderivative size = 25

$$\int \frac{1}{\sqrt{-8+6x+9x^2}} dx = \frac{1}{3} \operatorname{arctanh} \left(\frac{1+3x}{\sqrt{-8+6x+9x^2}} \right)$$

output `1/3*arctanh((1+3*x)/(9*x^2+6*x-8)^(1/2))`

3.146.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{1}{\sqrt{-8+6x+9x^2}} dx = -\frac{1}{3} \log \left(-1 - 3x + \sqrt{-8+6x+9x^2} \right)$$

input `Integrate[1/Sqrt[-8 + 6*x + 9*x^2],x]`

output `-1/3*Log[-1 - 3*x + Sqrt[-8 + 6*x + 9*x^2]]`

3.146.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {1092, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{9x^2 + 6x - 8}} dx$$

↓ 1092

$$2 \int \frac{1}{36 - \frac{36(3x+1)^2}{9x^2+6x-8}} d \frac{6(3x+1)}{\sqrt{9x^2 + 6x - 8}}$$

↓ 219

$$\frac{1}{3} \operatorname{arctanh} \left(\frac{3x+1}{\sqrt{9x^2 + 6x - 8}} \right)$$

input `Int[1/Sqrt[-8 + 6*x + 9*x^2],x]`

output `ArcTanh[(1 + 3*x)/Sqrt[-8 + 6*x + 9*x^2]]/3`

3.146.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1092 `Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Simp[2 Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x]`

3.146.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

method	result	size
trager	$\frac{\ln(\sqrt{9x^2+6x-8}+1+3x)}{3}$	21
default	$\frac{\ln\left(\frac{(3+9x)\sqrt{9}+\sqrt{9x^2+6x-8}}{9}\right)\sqrt{9}}{9}$	30

input `int(1/(9*x^2+6*x-8)^(1/2),x,method=_RETURNVERBOSE)`output `1/3*ln((9*x^2+6*x-8)^(1/2)+1+3*x)`**3.146.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{1}{\sqrt{-8+6x+9x^2}} dx = -\frac{1}{3} \log\left(-3x + \sqrt{9x^2+6x-8} - 1\right)$$

input `integrate(1/(9*x^2+6*x-8)^(1/2),x, algorithm="fricas")`output `-1/3*log(-3*x + sqrt(9*x^2 + 6*x - 8) - 1)`**3.146.6 Sympy [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{1}{\sqrt{-8+6x+9x^2}} dx = \frac{\log(18x + 6\sqrt{9x^2+6x-8} + 6)}{3}$$

input `integrate(1/(9*x**2+6*x-8)**(1/2),x)`output `log(18*x + 6*sqrt(9*x**2 + 6*x - 8) + 6)/3`

3.146.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{1}{\sqrt{-8+6x+9x^2}} dx = \frac{1}{3} \log \left(18x + 6\sqrt{9x^2+6x-8} + 6 \right)$$

input `integrate(1/(9*x^2+6*x-8)^(1/2),x, algorithm="maxima")`output `1/3*log(18*x + 6*sqrt(9*x^2 + 6*x - 8) + 6)`**3.146.8 Giac [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.64

$$\int \frac{1}{\sqrt{-8+6x+9x^2}} dx = \frac{1}{6} \sqrt{9x^2+6x-8}(3x+1) + \frac{3}{2} \log \left(\left| -3x + \sqrt{9x^2+6x-8} - 1 \right| \right)$$

input `integrate(1/(9*x^2+6*x-8)^(1/2),x, algorithm="giac")`output `1/6*sqrt(9*x^2 + 6*x - 8)*(3*x + 1) + 3/2*log(abs(-3*x + sqrt(9*x^2 + 6*x - 8) - 1))`**3.146.9 Mupad [B] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{1}{\sqrt{-8+6x+9x^2}} dx = \frac{\ln(3x + \sqrt{9x^2+6x-8} + 1)}{3}$$

input `int(1/(6*x + 9*x^2 - 8)^(1/2),x)`output `log(3*x + (6*x + 9*x^2 - 8)^(1/2) + 1)/3`

3.147 $\int \frac{x^2}{\sqrt{4x-x^2}} dx$

3.147.1 Optimal result	809
3.147.2 Mathematica [A] (verified)	809
3.147.3 Rubi [A] (verified)	810
3.147.4 Maple [A] (verified)	811
3.147.5 Fricas [A] (verification not implemented)	812
3.147.6 Sympy [A] (verification not implemented)	812
3.147.7 Maxima [A] (verification not implemented)	812
3.147.8 Giac [A] (verification not implemented)	813
3.147.9 Mupad [F(-1)]	813

3.147.1 Optimal result

Integrand size = 17, antiderivative size = 44

$$\int \frac{x^2}{\sqrt{4x-x^2}} dx = -3\sqrt{4x-x^2} - \frac{1}{2}x\sqrt{4x-x^2} - 6 \arcsin\left(1 - \frac{x}{2}\right)$$

output `6*arcsin(-1+1/2*x)-3*(-x^2+4*x)^(1/2)-1/2*x*(-x^2+4*x)^(1/2)`

3.147.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.25

$$\int \frac{x^2}{\sqrt{4x-x^2}} dx = \frac{x(-24+2x+x^2) - 24\sqrt{-4+x}\sqrt{x} \log(\sqrt{-4+x} - \sqrt{x})}{2\sqrt{-((-4+x)x)}}$$

input `Integrate[x^2/Sqrt[4*x - x^2],x]`

output `(x*(-24 + 2*x + x^2) - 24*Sqrt[-4 + x]*Sqrt[x]*Log[Sqrt[-4 + x] - Sqrt[x]])/(2*Sqrt[-((-4 + x)*x]))`

3.147.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.11, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {1134, 1160, 1090, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2}{\sqrt{4x-x^2}} dx \\
 & \quad \downarrow \text{1134} \\
 & 3 \int \frac{x}{\sqrt{4x-x^2}} dx - \frac{1}{2} x \sqrt{4x-x^2} \\
 & \quad \downarrow \text{1160} \\
 & 3 \left(2 \int \frac{1}{\sqrt{4x-x^2}} dx - \sqrt{4x-x^2} \right) - \frac{1}{2} x \sqrt{4x-x^2} \\
 & \quad \downarrow \text{1090} \\
 & 3 \left(-\frac{1}{2} \int \frac{1}{\sqrt{1-\frac{1}{16}(4-2x)^2}} d(4-2x) - \sqrt{4x-x^2} \right) - \frac{1}{2} x \sqrt{4x-x^2} \\
 & \quad \downarrow \text{223} \\
 & 3 \left(-2 \arcsin \left(\frac{1}{4}(4-2x) \right) - \sqrt{4x-x^2} \right) - \frac{1}{2} x \sqrt{4x-x^2}
 \end{aligned}$$

input `Int[x^2/Sqrt[4*x - x^2],x]`

output `-1/2*(x*Sqrt[4*x - x^2]) + 3*(-Sqrt[4*x - x^2] - 2*ArcSin[(4 - 2*x)/4])`

3.147.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 1090 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

rule 1134 `Int[((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[e*(d + e*x)^(m - 1)*((a + b*x + c*x^2)^(p + 1)/(c*(m + 2*p + 1))), x] + Simp[(m + p)*((2*c*d - b*e)/(c*(m + 2*p + 1))) Int[(d + e*x)^(m - 1)*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c*d^2 - b*d*e + a*e^2, 0] && GtQ[m, 1] && NeQ[m + 2*p + 1, 0] && IntegerQ[2*p]`

rule 1160 `Int[((d_.) + (e_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[e*((a + b*x + c*x^2)^(p + 1)/(2*c*(p + 1))), x] + Simp[(2*c*d - b*e)/(2*c) Int[(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[p, -1]`

3.147.4 Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.61

method	result
risch	$\frac{(6+x)x(x-4)}{2\sqrt{-x(x-4)}} + 6 \arcsin\left(-1 + \frac{x}{2}\right)$
default	$6 \arcsin\left(-1 + \frac{x}{2}\right) - 3\sqrt{-x^2 + 4x} - \frac{x\sqrt{-x^2 + 4x}}{2}$
pseudoelliptic	$-\frac{x\sqrt{-x(x-4)}}{2} - 3\sqrt{-x(x-4)} - 12 \arctan\left(\frac{\sqrt{-x(x-4)}}{x}\right)$
meijerg	$-\frac{16i\left(-\frac{i\sqrt{\pi}\sqrt{x}\left(\frac{5x}{2}+15\right)\sqrt{-\frac{x}{4}+1}}{40} + \frac{3i\sqrt{\pi}\arcsin\left(\frac{\sqrt{x}}{2}\right)}{4}\right)}{\sqrt{\pi}}$
trager	$\left(-3 - \frac{x}{2}\right)\sqrt{-x^2 + 4x} - 6 \operatorname{RootOf}\left(_Z^2 + 1\right) \ln\left(\operatorname{RootOf}\left(_Z^2 + 1\right) x - 2 \operatorname{RootOf}\left(_Z^2 + 1\right)\right)$

input `int(x^2/(-x^2+4*x)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*(6+x)*x*(x-4)/(-x*(x-4))^(1/2)+6*arcsin(-1+1/2*x)`

3.147. $\int \frac{x^2}{\sqrt{4x-x^2}} dx$

3.147.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.80

$$\int \frac{x^2}{\sqrt{4x-x^2}} dx = -\frac{1}{2} \sqrt{-x^2+4x}(x+6) - 12 \arctan\left(\frac{\sqrt{-x^2+4x}}{x}\right)$$

input `integrate(x^2/(-x^2+4*x)^(1/2),x, algorithm="fricas")`output `-1/2*sqrt(-x^2 + 4*x)*(x + 6) - 12*arctan(sqrt(-x^2 + 4*x)/x)`**3.147.6 Sympy [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.55

$$\int \frac{x^2}{\sqrt{4x-x^2}} dx = \left(-\frac{x}{2} - 3\right) \sqrt{-x^2+4x} + 6 \arcsin\left(\frac{x}{2} - 1\right)$$

input `integrate(x**2/(-x**2+4*x)**(1/2),x)`output `(-x/2 - 3)*sqrt(-x**2 + 4*x) + 6*asin(x/2 - 1)`**3.147.7 Maxima [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.82

$$\int \frac{x^2}{\sqrt{4x-x^2}} dx = -\frac{1}{2} \sqrt{-x^2+4x}x - 3 \sqrt{-x^2+4x} - 6 \arcsin\left(-\frac{1}{2}x + 1\right)$$

input `integrate(x^2/(-x^2+4*x)^(1/2),x, algorithm="maxima")`output `-1/2*sqrt(-x^2 + 4*x)*x - 3*sqrt(-x^2 + 4*x) - 6*arcsin(-1/2*x + 1)`

3.147.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.57

$$\int \frac{x^2}{\sqrt{4x-x^2}} dx = -\frac{1}{2} \sqrt{-x^2+4x}(x+6) + 6 \arcsin\left(\frac{1}{2}x-1\right)$$

input `integrate(x^2/(-x^2+4*x)^(1/2),x, algorithm="giac")`output `-1/2*sqrt(-x^2 + 4*x)*(x + 6) + 6*arcsin(1/2*x - 1)`**3.147.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2}{\sqrt{4x-x^2}} dx = \int \frac{x^2}{\sqrt{4x-x^2}} dx$$

input `int(x^2/(4*x - x^2)^(1/2),x)`output `int(x^2/(4*x - x^2)^(1/2), x)`

$$3.148 \quad \int \frac{1}{(2+2x+x^2)^2} dx$$

3.148.1 Optimal result	814
3.148.2 Mathematica [A] (verified)	814
3.148.3 Rubi [A] (verified)	815
3.148.4 Maple [A] (verified)	816
3.148.5 Fracas [A] (verification not implemented)	816
3.148.6 Sympy [A] (verification not implemented)	817
3.148.7 Maxima [A] (verification not implemented)	817
3.148.8 Giac [A] (verification not implemented)	817
3.148.9 Mupad [B] (verification not implemented)	818

3.148.1 Optimal result

Integrand size = 10, antiderivative size = 26

$$\int \frac{1}{(2+2x+x^2)^2} dx = \frac{1+x}{2(2+2x+x^2)} + \frac{1}{2} \arctan(1+x)$$

output `1/2*(1+x)/(x^2+2*x+2)+1/2*arctan(1+x)`

3.148.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.88

$$\int \frac{1}{(2+2x+x^2)^2} dx = \frac{1}{2} \left(\frac{1+x}{2+2x+x^2} + \arctan(1+x) \right)$$

input `Integrate[(2 + 2*x + x^2)^(-2), x]`

output `((1 + x)/(2 + 2*x + x^2) + ArcTan[1 + x])/2`

3.148.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {1086, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(x^2 + 2x + 2)^2} dx \\
 & \quad \downarrow \text{1086} \\
 & \frac{1}{2} \int \frac{1}{x^2 + 2x + 2} dx + \frac{x + 1}{2(x^2 + 2x + 2)} \\
 & \quad \downarrow \text{1082} \\
 & \frac{x + 1}{2(x^2 + 2x + 2)} - \frac{1}{2} \int \frac{1}{-(x + 1)^2 - 1} d(x + 1) \\
 & \quad \downarrow \text{217} \\
 & \frac{1}{2} \arctan(x + 1) + \frac{x + 1}{2(x^2 + 2x + 2)}
 \end{aligned}$$

input `Int[(2 + 2*x + x^2)^(-2),x]`

output `(1 + x)/(2*(2 + 2*x + x^2)) + ArcTan[1 + x]/2`

3.148.3.1 Defintions of rubi rules used

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1086 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x) * ((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] - Simp[2*c*((2*p + 3)/((p + 1)*(b^2 - 4*a*c))) Int[(a + b*x + c*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && ILtQ[p, -1]`

3.148.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

method	result	size
risch	$\frac{\frac{1}{2} + \frac{x}{2}}{x^2 + 2x + 2} + \frac{\arctan(1+x)}{2}$	24
default	$\frac{2x+2}{4x^2+8x+8} + \frac{\arctan(1+x)}{2}$	25
parallelrisch	$-\frac{i \ln(x+1-i)x^2 - i \ln(x+1+i)x^2 + 2i \ln(x+1-i)x - 2i \ln(x+1+i)x + 2i \ln(x+1-i) - 2i \ln(x+1+i) + x^2}{4(x^2 + 2x + 2)}$	79

input `int(1/(x^2+2*x+2)^2,x,method=_RETURNVERBOSE)`

output `(1/2+1/2*x)/(x^2+2*x+2)+1/2*arctan(1+x)`

3.148.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

$$\int \frac{1}{(2 + 2x + x^2)^2} dx = \frac{(x^2 + 2x + 2) \arctan(x + 1) + x + 1}{2(x^2 + 2x + 2)}$$

input `integrate(1/(x^2+2*x+2)^2,x, algorithm="fricas")`

output `1/2*((x^2 + 2*x + 2)*arctan(x + 1) + x + 1)/(x^2 + 2*x + 2)`

3.148.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.73

$$\int \frac{1}{(2+2x+x^2)^2} dx = \frac{x+1}{2x^2+4x+4} + \frac{\operatorname{atan}(x+1)}{2}$$

input `integrate(1/(x**2+2*x+2)**2,x)`output `(x + 1)/(2*x**2 + 4*x + 4) + atan(x + 1)/2`**3.148.7 Maxima [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{1}{(2+2x+x^2)^2} dx = \frac{x+1}{2(x^2+2x+2)} + \frac{1}{2} \arctan(x+1)$$

input `integrate(1/(x^2+2*x+2)^2,x, algorithm="maxima")`output `1/2*(x + 1)/(x^2 + 2*x + 2) + 1/2*arctan(x + 1)`**3.148.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{1}{(2+2x+x^2)^2} dx = \frac{x+1}{2(x^2+2x+2)} + \frac{1}{2} \arctan(x+1)$$

input `integrate(1/(x^2+2*x+2)^2,x, algorithm="giac")`output `1/2*(x + 1)/(x^2 + 2*x + 2) + 1/2*arctan(x + 1)`

3.148.9 Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.88

$$\int \frac{1}{(2 + 2x + x^2)^2} dx = \frac{\operatorname{atan}(x + 1)}{2} + \frac{\frac{x}{2} + \frac{1}{2}}{x^2 + 2x + 2}$$

input `int(1/(2*x + x^2 + 2)^2,x)`

output `atan(x + 1)/2 + (x/2 + 1/2)/(2*x + x^2 + 2)`

3.149 $\int \frac{1}{(5-4x-x^2)^{5/2}} dx$

3.149.1 Optimal result	819
3.149.2 Mathematica [A] (verified)	819
3.149.3 Rubi [A] (verified)	820
3.149.4 Maple [A] (verified)	821
3.149.5 Fricas [A] (verification not implemented)	821
3.149.6 Sympy [F]	821
3.149.7 Maxima [A] (verification not implemented)	822
3.149.8 Giac [A] (verification not implemented)	822
3.149.9 Mupad [B] (verification not implemented)	822

3.149.1 Optimal result

Integrand size = 14, antiderivative size = 43

$$\int \frac{1}{(5-4x-x^2)^{5/2}} dx = \frac{2+x}{27(5-4x-x^2)^{3/2}} + \frac{2(2+x)}{243\sqrt{5-4x-x^2}}$$

output `1/27*(2+x)/(-x^2-4*x+5)^(3/2)+2/243*(2+x)/(-x^2-4*x+5)^(1/2)`

3.149.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int \frac{1}{(5-4x-x^2)^{5/2}} dx = \frac{\sqrt{5-4x-x^2}(38+3x-12x^2-2x^3)}{243(-1+x)^2(5+x)^2}$$

input `Integrate[(5 - 4*x - x^2)^(-5/2), x]`

output `(Sqrt[5 - 4*x - x^2]*(38 + 3*x - 12*x^2 - 2*x^3))/(243*(-1 + x)^2*(5 + x)^2)`

3.149.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {1089, 1088}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(-x^2 - 4x + 5)^{5/2}} dx$$

↓ 1089

$$\frac{2}{27} \int \frac{1}{(-x^2 - 4x + 5)^{3/2}} dx + \frac{x + 2}{27(-x^2 - 4x + 5)^{3/2}}$$

↓ 1088

$$\frac{2(x + 2)}{243\sqrt{-x^2 - 4x + 5}} + \frac{x + 2}{27(-x^2 - 4x + 5)^{3/2}}$$

input `Int[(5 - 4*x - x^2)^(-5/2), x]`

output `(2 + x)/(27*(5 - 4*x - x^2)^(3/2)) + (2*(2 + x))/(243*Sqrt[5 - 4*x - x^2])`

3.149.3.1 Defintions of rubi rules used

rule 1088 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-3/2), x_Symbol] :> Simp[-2*((b + 2*c*x)/((b^2 - 4*a*c)*Sqrt[a + b*x + c*x^2])), x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]`

rule 1089 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(b + 2*c*x)*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] - Simp[2*c*((2*p + 3)/((p + 1)*(b^2 - 4*a*c)) Int[(a + b*x + c*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[3*p])`

3.149.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.84

method	result	size
gospers	$\frac{(5+x)(-1+x)(2x^3+12x^2-3x-38)}{243(-x^2-4x+5)^{\frac{5}{2}}}$	36
default	$-\frac{-2x-4}{54(-x^2-4x+5)^{\frac{3}{2}}} - \frac{-2x-4}{243\sqrt{-x^2-4x+5}}$	40
trager	$-\frac{(2x^3+12x^2-3x-38)\sqrt{-x^2-4x+5}}{243(x^2+4x-5)^2}$	40
risch	$\frac{2x^3+12x^2-3x-38}{243(x^2+4x-5)\sqrt{-x^2-4x+5}}$	40

input `int(1/(-x^2-4*x+5)^(5/2),x,method=_RETURNVERBOSE)`output `1/243*(5+x)*(-1+x)*(2*x^3+12*x^2-3*x-38)/(-x^2-4*x+5)^(5/2)`**3.149.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.14

$$\int \frac{1}{(5-4x-x^2)^{5/2}} dx = -\frac{(2x^3+12x^2-3x-38)\sqrt{-x^2-4x+5}}{243(x^4+8x^3+6x^2-40x+25)}$$

input `integrate(1/(-x^2-4*x+5)^(5/2),x, algorithm="fricas")`output `-1/243*(2*x^3 + 12*x^2 - 3*x - 38)*sqrt(-x^2 - 4*x + 5)/(x^4 + 8*x^3 + 6*x^2 - 40*x + 25)`**3.149.6 Sympy [F]**

$$\int \frac{1}{(5-4x-x^2)^{5/2}} dx = \int \frac{1}{(-x^2-4x+5)^{\frac{5}{2}}} dx$$

input `integrate(1/(-x**2-4*x+5)**(5/2),x)`output `Integral((-x**2 - 4*x + 5)**(-5/2), x)`

3.149.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.37

$$\int \frac{1}{(5-4x-x^2)^{5/2}} dx = \frac{2x}{243\sqrt{-x^2-4x+5}} + \frac{4}{243\sqrt{-x^2-4x+5}} + \frac{x}{27(-x^2-4x+5)^{3/2}} + \frac{2}{27(-x^2-4x+5)^{3/2}}$$

input `integrate(1/(-x^2-4*x+5)^(5/2),x, algorithm="maxima")`output `2/243*x/sqrt(-x^2 - 4*x + 5) + 4/243/sqrt(-x^2 - 4*x + 5) + 1/27*x/(-x^2 - 4*x + 5)^(3/2) + 2/27/(-x^2 - 4*x + 5)^(3/2)`**3.149.8 Giac [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.84

$$\int \frac{1}{(5-4x-x^2)^{5/2}} dx = -\frac{((2(x+6)x-3)x-38)\sqrt{-x^2-4x+5}}{243(x^2+4x-5)^2}$$

input `integrate(1/(-x^2-4*x+5)^(5/2),x, algorithm="giac")`output `-1/243*((2*(x+6)*x-3)*x-38)*sqrt(-x^2-4*x+5)/(x^2+4*x-5)^2`**3.149.9 Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.67

$$\int \frac{1}{(5-4x-x^2)^{5/2}} dx = -\frac{(4x+8)(8x^2+32x-76)}{3888(-x^2-4x+5)^{3/2}}$$

input `int(1/(5 - x^2 - 4*x)^(5/2),x)`output `-((4*x + 8)*(32*x + 8*x^2 - 76))/(3888*(5 - x^2 - 4*x)^(3/2))`

3.150 $\int e^t \sqrt{9 - e^{2t}} dt$

3.150.1 Optimal result	823
3.150.2 Mathematica [A] (verified)	823
3.150.3 Rubi [A] (verified)	824
3.150.4 Maple [A] (verified)	825
3.150.5 Fricas [A] (verification not implemented)	825
3.150.6 Sympy [A] (verification not implemented)	826
3.150.7 Maxima [A] (verification not implemented)	826
3.150.8 Giac [A] (verification not implemented)	826
3.150.9 Mupad [B] (verification not implemented)	827

3.150.1 Optimal result

Integrand size = 17, antiderivative size = 33

$$\int e^t \sqrt{9 - e^{2t}} dt = \frac{1}{2} e^t \sqrt{9 - e^{2t}} + \frac{9}{2} \arcsin\left(\frac{e^t}{3}\right)$$

output `9/2*arcsin(1/3*exp(t))+1/2*exp(t)*(9-exp(2*t))^(1/2)`

3.150.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.36

$$\int e^t \sqrt{9 - e^{2t}} dt = \frac{1}{2} e^t \sqrt{9 - e^{2t}} - 9 \arctan\left(\frac{\sqrt{9 - e^{2t}}}{3 + e^t}\right)$$

input `Integrate[E^t*Sqrt[9 - E^(2*t)],t]`

output `(E^t*Sqrt[9 - E^(2*t)])/2 - 9*ArcTan[Sqrt[9 - E^(2*t)]/(3 + E^t)]`

3.150.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2679, 211, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^t \sqrt{9 - e^{2t}} dt \\ & \quad \downarrow \text{2679} \\ & \int \sqrt{9 - e^{2t}} de^t \\ & \quad \downarrow \text{211} \\ & \frac{9}{2} \int \frac{1}{\sqrt{9 - e^{2t}}} de^t + \frac{1}{2} e^t \sqrt{9 - e^{2t}} \\ & \quad \downarrow \text{223} \\ & \frac{9}{2} \arcsin\left(\frac{e^t}{3}\right) + \frac{1}{2} e^t \sqrt{9 - e^{2t}} \end{aligned}$$

input `Int[E^t*Sqrt[9 - E^(2*t)],t]`

output `(E^t*Sqrt[9 - E^(2*t)])/2 + (9*ArcSin[E^t/3])/2`

3.150.3.1 Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

```
rule 2679 Int[((a_) + (b_)*(F_)^((e_)*((c_) + (d_)*(x_)))^(p_)*(G_)^((h_)*((f_
.) + (g_)*(x_))), x_Symbol] := With[{m = FullSimplify[d*e*(Log[F]/(g*h*Log
[G]))]}, Simp[Denominator[m]/(g*h*Log[G]) Subst[Int[x^(Denominator[m] - 1
)*(a + b*F^(c*e - d*e*(f/g))*x^Numerator[m])^p, x], x, G^(h*((f + g*x)/Deno
minator[m]))], x] /; LtQ[m, -1] || GtQ[m, 1]] /; FreeQ[{F, G, a, b, c, d, e
, f, g, h, p}, x]
```

3.150.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.70

method	result	size
default	$\frac{9 \arcsin\left(\frac{e^t}{3}\right)}{2} + \frac{e^t \sqrt{9-e^{2t}}}{2}$	23
risch	$-\frac{e^t(-9+e^{2t})}{2\sqrt{9-e^{2t}}} + \frac{9 \arcsin\left(\frac{e^t}{3}\right)}{2}$	29

```
input int(exp(t)*(9-exp(2*t))^(1/2),t,method=_RETURNVERBOSE)
```

```
output 1/2*exp(t)*(9-exp(t)^2)^(1/2)+9/2*arcsin(1/3*exp(t))
```

3.150.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.06

$$\int e^t \sqrt{9 - e^{2t}} dt = \frac{1}{2} \sqrt{-e^{(2t)} + 9} e^t - 9 \arctan\left(\left(\sqrt{-e^{(2t)} + 9} - 3\right) e^{(-t)}\right)$$

```
input integrate(exp(t)*(9-exp(2*t))^(1/2),t, algorithm="fricas")
```

```
output 1/2*sqrt(-e^(2*t) + 9)*e^t - 9*arctan((sqrt(-e^(2*t) + 9) - 3)*e^(-t))
```

3.150.6 Sympy [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.73

$$\int e^t \sqrt{9 - e^{2t}} dt = \frac{\sqrt{9 - e^{2t}} e^t}{2} + \frac{9 \operatorname{asin}\left(\frac{e^t}{3}\right)}{2}$$

input `integrate(exp(t)*(9-exp(2*t))**(1/2),t)`output `sqrt(9 - exp(2*t))*exp(t)/2 + 9*asin(exp(t)/3)/2`**3.150.7 Maxima [A] (verification not implemented)**

Time = 0.37 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.67

$$\int e^t \sqrt{9 - e^{2t}} dt = \frac{1}{2} \sqrt{-e^{(2t)} + 9e^t} + \frac{9}{2} \arcsin\left(\frac{1}{3} e^t\right)$$

input `integrate(exp(t)*(9-exp(2*t))^(1/2),t, algorithm="maxima")`output `1/2*sqrt(-e^(2*t) + 9)*e^t + 9/2*arcsin(1/3*e^t)`**3.150.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.67

$$\int e^t \sqrt{9 - e^{2t}} dt = \frac{1}{2} \sqrt{-e^{(2t)} + 9e^t} + \frac{9}{2} \arcsin\left(\frac{1}{3} e^t\right)$$

input `integrate(exp(t)*(9-exp(2*t))^(1/2),t, algorithm="giac")`output `1/2*sqrt(-e^(2*t) + 9)*e^t + 9/2*arcsin(1/3*e^t)`

3.150.9 Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.67

$$\int e^t \sqrt{9 - e^{2t}} dt = \frac{9 \operatorname{asin}\left(\frac{e^t}{3}\right)}{2} + \frac{e^t \sqrt{9 - e^{2t}}}{2}$$

input `int(exp(t)*(9 - exp(2*t))^(1/2),t)`

output `(9*asin(exp(t)/3))/2 + (exp(t)*(9 - exp(2*t))^(1/2))/2`

3.151 $\int \sqrt{-9 + e^{2t}} dt$

3.151.1 Optimal result	828
3.151.2 Mathematica [A] (verified)	828
3.151.3 Rubi [A] (verified)	829
3.151.4 Maple [A] (verified)	830
3.151.5 Fricas [A] (verification not implemented)	831
3.151.6 Sympy [A] (verification not implemented)	831
3.151.7 Maxima [A] (verification not implemented)	831
3.151.8 Giac [A] (verification not implemented)	832
3.151.9 Mupad [B] (verification not implemented)	832

3.151.1 Optimal result

Integrand size = 11, antiderivative size = 30

$$\int \sqrt{-9 + e^{2t}} dt = \sqrt{-9 + e^{2t}} - 3 \arctan \left(\frac{1}{3} \sqrt{-9 + e^{2t}} \right)$$

output `-3*arctan(1/3*(-9+exp(2*t))^(1/2))+(-9+exp(2*t))^(1/2)`

3.151.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \sqrt{-9 + e^{2t}} dt = \sqrt{-9 + e^{2t}} - 3 \arctan \left(\frac{1}{3} \sqrt{-9 + e^{2t}} \right)$$

input `Integrate[Sqrt[-9 + E^(2*t)],t]`

output `Sqrt[-9 + E^(2*t)] - 3*ArcTan[Sqrt[-9 + E^(2*t)]]/3`

3.151.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.20, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {2720, 60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{e^{2t} - 9} dt \\
 & \quad \downarrow \text{2720} \\
 & \frac{1}{2} \int e^{-2t} \sqrt{-9 + e^{2t}} de^{2t} \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2} \left(2\sqrt{e^{2t} - 9} - 9 \int \frac{e^{-2t}}{\sqrt{-9 + e^{2t}}} de^{2t} \right) \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{2} \left(2\sqrt{e^{2t} - 9} - 18 \int \frac{1}{9 + e^{4t}} d\sqrt{-9 + e^{2t}} \right) \\
 & \quad \downarrow \text{216} \\
 & \frac{1}{2} \left(2\sqrt{e^{2t} - 9} - 6 \arctan \left(\frac{1}{3} \sqrt{e^{2t} - 9} \right) \right)
 \end{aligned}$$

input `Int[Sqrt[-9 + E^(2*t)],t]`

output `(2*Sqrt[-9 + E^(2*t)] - 6*ArcTan[Sqrt[-9 + E^(2*t)]/3])/2`

3.151.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
  {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
  d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
  Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
  inearQ[a, b, c, d, m, n, x]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
  rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
  , 0] || GtQ[b, 0])
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
  ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
  [{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
  *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]
```

3.151.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.77

method	result	size
derivativedivides	$-3 \arctan\left(\frac{\sqrt{-9+e^{2t}}}{3}\right) + \sqrt{-9+e^{2t}}$	23
default	$-3 \arctan\left(\frac{\sqrt{-9+e^{2t}}}{3}\right) + \sqrt{-9+e^{2t}}$	23
risch	$-3 \arctan\left(\frac{\sqrt{-9+e^{2t}}}{3}\right) + \sqrt{-9+e^{2t}}$	23

```
input int((-9+exp(2*t))^(1/2),t,method=_RETURNVERBOSE)
```

```
output -3*arctan(1/3*(-9+exp(2*t))^(1/2))+(-9+exp(2*t))^(1/2)
```

3.151.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \sqrt{-9 + e^{2t}} dt = \sqrt{e^{2t} - 9} - 3 \arctan\left(\frac{1}{3} \sqrt{e^{2t} - 9}\right)$$

input `integrate((-9+exp(2*t))^(1/2),t, algorithm="fricas")`output `sqrt(e^(2*t) - 9) - 3*arctan(1/3*sqrt(e^(2*t) - 9))`**3.151.6 Sympy [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int \sqrt{-9 + e^{2t}} dt = \sqrt{e^{2t} - 9} - 3 \operatorname{atan}\left(\frac{\sqrt{e^{2t} - 9}}{3}\right)$$

input `integrate((-9+exp(2*t))**(1/2),t)`output `sqrt(exp(2*t) - 9) - 3*atan(sqrt(exp(2*t) - 9)/3)`**3.151.7 Maxima [A] (verification not implemented)**

Time = 0.39 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \sqrt{-9 + e^{2t}} dt = \sqrt{e^{2t} - 9} - 3 \arctan\left(\frac{1}{3} \sqrt{e^{2t} - 9}\right)$$

input `integrate((-9+exp(2*t))^(1/2),t, algorithm="maxima")`output `sqrt(e^(2*t) - 9) - 3*arctan(1/3*sqrt(e^(2*t) - 9))`

3.151.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \sqrt{-9 + e^{2t}} dt = \sqrt{e^{2t} - 9} - 3 \arctan\left(\frac{1}{3} \sqrt{e^{2t} - 9}\right)$$

input `integrate((-9+exp(2*t))^(1/2),t, algorithm="giac")`output `sqrt(e^(2*t) - 9) - 3*arctan(1/3*sqrt(e^(2*t) - 9))`**3.151.9 Mupad [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.13

$$\int \sqrt{-9 + e^{2t}} dt = \left(\frac{3e^{-t} \operatorname{asin}(3e^{-t})}{\sqrt{1 - 9e^{-2t}}} + 1 \right) \sqrt{e^{2t} - 9}$$

input `int((exp(2*t) - 9)^(1/2),t)`output `((3*exp(-t)*asin(3*exp(-t)))/(1 - 9*exp(-2*t))^(1/2) + 1)*(exp(2*t) - 9)^(1/2)`

3.152 $\int \frac{1}{\sqrt{a^2+x^2}} dx$

3.152.1 Optimal result	833
3.152.2 Mathematica [B] (verified)	833
3.152.3 Rubi [A] (verified)	834
3.152.4 Maple [A] (verified)	835
3.152.5 Fricas [A] (verification not implemented)	835
3.152.6 Sympy [A] (verification not implemented)	835
3.152.7 Maxima [A] (verification not implemented)	836
3.152.8 Giac [B] (verification not implemented)	836
3.152.9 Mupad [B] (verification not implemented)	836

3.152.1 Optimal result

Integrand size = 11, antiderivative size = 14

$$\int \frac{1}{\sqrt{a^2+x^2}} dx = \operatorname{arctanh}\left(\frac{x}{\sqrt{a^2+x^2}}\right)$$

output `arctanh(x/(a^2+x^2)^(1/2))`

3.152.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 42 vs. $2(14) = 28$.

Time = 0.00 (sec) , antiderivative size = 42, normalized size of antiderivative = 3.00

$$\int \frac{1}{\sqrt{a^2+x^2}} dx = -\frac{1}{2} \log\left(1 - \frac{x}{\sqrt{a^2+x^2}}\right) + \frac{1}{2} \log\left(1 + \frac{x}{\sqrt{a^2+x^2}}\right)$$

input `Integrate[1/Sqrt[a^2 + x^2],x]`

output `-1/2*Log[1 - x/Sqrt[a^2 + x^2]] + Log[1 + x/Sqrt[a^2 + x^2]]/2`

3.152.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{1}{\sqrt{a^2 + x^2}} dx \\ \downarrow 224 \\ \int \frac{1}{1 - \frac{x^2}{a^2 + x^2}} d \frac{x}{\sqrt{a^2 + x^2}} \\ \downarrow 219 \\ \operatorname{arctanh} \left(\frac{x}{\sqrt{a^2 + x^2}} \right) \end{array}$$

input `Int[1/Sqrt[a^2 + x^2],x]`

output `ArcTanh[x/Sqrt[a^2 + x^2]]`

3.152.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

3.152.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
default	$\ln(x + \sqrt{a^2 + x^2})$	13
pseudoelliptic	$\operatorname{arctanh}\left(\frac{\sqrt{a^2+x^2}}{x}\right)$	15

input `int(1/(a^2+x^2)^(1/2),x,method=_RETURNVERBOSE)`

output `ln(x+(a^2+x^2)^(1/2))`

3.152.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{1}{\sqrt{a^2 + x^2}} dx = -\log(-x + \sqrt{a^2 + x^2})$$

input `integrate(1/(a^2+x^2)^(1/2),x, algorithm="fracas")`

output `-log(-x + sqrt(a^2 + x^2))`

3.152.6 Sympy [A] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.21

$$\int \frac{1}{\sqrt{a^2 + x^2}} dx = \operatorname{asinh}\left(\frac{x}{a}\right)$$

input `integrate(1/(a**2+x**2)**(1/2),x)`

output `asinh(x/a)`

3.152.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.43

$$\int \frac{1}{\sqrt{a^2 + x^2}} dx = \operatorname{arsinh}\left(\frac{x}{a}\right)$$

input `integrate(1/(a^2+x^2)^(1/2),x, algorithm="maxima")`

output `arcsinh(x/a)`

3.152.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 32 vs. 2(12) = 24.

Time = 0.28 (sec) , antiderivative size = 32, normalized size of antiderivative = 2.29

$$\int \frac{1}{\sqrt{a^2 + x^2}} dx = -\frac{1}{2} a^2 \log\left(-x + \sqrt{a^2 + x^2}\right) + \frac{1}{2} \sqrt{a^2 + x^2} x$$

input `integrate(1/(a^2+x^2)^(1/2),x, algorithm="giac")`

output `-1/2*a^2*log(-x + sqrt(a^2 + x^2)) + 1/2*sqrt(a^2 + x^2)*x`

3.152.9 Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sqrt{a^2 + x^2}} dx = \ln\left(x + \sqrt{a^2 + x^2}\right)$$

input `int(1/(a^2 + x^2)^(1/2),x)`

output `log(x + (a^2 + x^2)^(1/2))`

3.153 $\int \frac{5+x}{-2+x+x^2} dx$

3.153.1 Optimal result	837
3.153.2 Mathematica [A] (verified)	837
3.153.3 Rubi [A] (verified)	838
3.153.4 Maple [A] (verified)	839
3.153.5 Fricas [A] (verification not implemented)	839
3.153.6 Sympy [A] (verification not implemented)	839
3.153.7 Maxima [A] (verification not implemented)	840
3.153.8 Giac [A] (verification not implemented)	840
3.153.9 Mupad [B] (verification not implemented)	840

3.153.1 Optimal result

Integrand size = 12, antiderivative size = 15

$$\int \frac{5+x}{-2+x+x^2} dx = 2 \log(1-x) - \log(2+x)$$

output `2*ln(1-x)-ln(2+x)`

3.153.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{5+x}{-2+x+x^2} dx = 2 \log(1-x) - \log(2+x)$$

input `Integrate[(5 + x)/(-2 + x + x^2), x]`

output `2*Log[1 - x] - Log[2 + x]`

3.153.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x+5}{x^2+x-2} dx$$

↓ 1141

$$\int \left(\frac{1}{-x-2} - \frac{2}{1-x} \right) dx$$

↓ 2009

$$2 \log(1-x) - \log(x+2)$$

input `Int[(5 + x)/(-2 + x + x^2),x]`

output `2*Log[1 - x] - Log[2 + x]`

3.153.3.1 Defintions of rubi rules used

rule 1141 `Int[((d_.) + (e_.)*(x_)^(m_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x], x] /; EqQ[p, -1] || !FractionalPowerFactorQ[q] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p, 0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.153.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
default	$2 \ln(-1+x) - \ln(2+x)$	14
norman	$2 \ln(-1+x) - \ln(2+x)$	14
risch	$2 \ln(-1+x) - \ln(2+x)$	14
parallelrisch	$2 \ln(-1+x) - \ln(2+x)$	14

input `int((5+x)/(x^2+x-2),x,method=_RETURNVERBOSE)`output `2*ln(-1+x)-ln(2+x)`**3.153.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{5+x}{-2+x+x^2} dx = -\log(x+2) + 2 \log(x-1)$$

input `integrate((5+x)/(x^2+x-2),x, algorithm="fracas")`output `-log(x + 2) + 2*log(x - 1)`**3.153.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int \frac{5+x}{-2+x+x^2} dx = 2 \log(x-1) - \log(x+2)$$

input `integrate((5+x)/(x**2+x-2),x)`output `2*log(x - 1) - log(x + 2)`

3.153.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{5+x}{-2+x+x^2} dx = -\log(x+2) + 2 \log(x-1)$$

input `integrate((5+x)/(x^2+x-2),x, algorithm="maxima")`output `-log(x + 2) + 2*log(x - 1)`**3.153.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{5+x}{-2+x+x^2} dx = -\log(|x+2|) + 2 \log(|x-1|)$$

input `integrate((5+x)/(x^2+x-2),x, algorithm="giac")`output `-log(abs(x + 2)) + 2*log(abs(x - 1))`**3.153.9 Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{5+x}{-2+x+x^2} dx = 2 \ln(x-1) - \ln(x+2)$$

input `int((x + 5)/(x + x^2 - 2),x)`output `2*log(x - 1) - log(x + 2)`

3.154 $\int \frac{x+x^3}{-1+x} dx$

3.154.1 Optimal result	841
3.154.2 Mathematica [A] (verified)	841
3.154.3 Rubi [A] (verified)	842
3.154.4 Maple [A] (verified)	843
3.154.5 Fricas [A] (verification not implemented)	843
3.154.6 Sympy [A] (verification not implemented)	843
3.154.7 Maxima [A] (verification not implemented)	844
3.154.8 Giac [A] (verification not implemented)	844
3.154.9 Mupad [B] (verification not implemented)	844

3.154.1 Optimal result

Integrand size = 11, antiderivative size = 26

$$\int \frac{x+x^3}{-1+x} dx = 2x + \frac{x^2}{2} + \frac{x^3}{3} + 2\log(1-x)$$

output `2*x+1/2*x^2+1/3*x^3+2*ln(1-x)`

3.154.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.96

$$\int \frac{x+x^3}{-1+x} dx = \frac{1}{6}(-17 + 12x + 3x^2 + 2x^3 + 12\log(-1+x))$$

input `Integrate[(x + x^3)/(-1 + x),x]`

output `(-17 + 12*x + 3*x^2 + 2*x^3 + 12*Log[-1 + x])/6`

3.154.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {2027, 522, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^3 + x}{x - 1} dx \\ & \quad \downarrow \text{2027} \\ & \int \frac{x(x^2 + 1)}{x - 1} dx \\ & \quad \downarrow \text{522} \\ & \int \left(x^2 + x + \frac{2}{x - 1} + 2 \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{x^3}{3} + \frac{x^2}{2} + 2x + 2 \log(1 - x) \end{aligned}$$

input `Int[(x + x^3)/(-1 + x),x]`

output `2*x + x^2/2 + x^3/3 + 2*Log[1 - x]`

3.154.3.1 Defintions of rubi rules used

rule 522 `Int[((e_.)*(x_.))^(m_.)*((c_) + (d_.)*(x_.))^(n_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(e*x)^m*(c + d*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2027 `Int[(Fx_.)*((a_.)*(x_)^(r_.) + (b_.)*(x_)^(s_.))^(p_.), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r))^p*Fx, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] && PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])`

3.154.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.81

method	result	size
default	$\frac{x^3}{3} + \frac{x^2}{2} + 2x + 2 \ln(-1 + x)$	21
norman	$\frac{x^3}{3} + \frac{x^2}{2} + 2x + 2 \ln(-1 + x)$	21
risch	$\frac{x^3}{3} + \frac{x^2}{2} + 2x + 2 \ln(-1 + x)$	21
parallelrisch	$\frac{x^3}{3} + \frac{x^2}{2} + 2x + 2 \ln(-1 + x)$	21
meijerg	$\frac{x(4x^2+6x+12)}{12} + 2 \ln(1 - x) + x$	24

input `int((x^3+x)/(-1+x),x,method=_RETURNVERBOSE)`output `1/3*x^3+1/2*x^2+2*x+2*ln(-1+x)`**3.154.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \frac{x + x^3}{-1 + x} dx = \frac{1}{3} x^3 + \frac{1}{2} x^2 + 2x + 2 \log(x - 1)$$

input `integrate((x^3+x)/(-1+x),x, algorithm="fracas")`output `1/3*x^3 + 1/2*x^2 + 2*x + 2*log(x - 1)`**3.154.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.73

$$\int \frac{x + x^3}{-1 + x} dx = \frac{x^3}{3} + \frac{x^2}{2} + 2x + 2 \log(x - 1)$$

input `integrate((x**3+x)/(-1+x),x)`output `x**3/3 + x**2/2 + 2*x + 2*log(x - 1)`

3.154.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \frac{x + x^3}{-1 + x} dx = \frac{1}{3} x^3 + \frac{1}{2} x^2 + 2x + 2 \log(x - 1)$$

input `integrate((x^3+x)/(-1+x),x, algorithm="maxima")`output `1/3*x^3 + 1/2*x^2 + 2*x + 2*log(x - 1)`**3.154.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.81

$$\int \frac{x + x^3}{-1 + x} dx = \frac{1}{3} x^3 + \frac{1}{2} x^2 + 2x + 2 \log(|x - 1|)$$

input `integrate((x^3+x)/(-1+x),x, algorithm="giac")`output `1/3*x^3 + 1/2*x^2 + 2*x + 2*log(abs(x - 1))`**3.154.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \frac{x + x^3}{-1 + x} dx = 2x + 2 \ln(x - 1) + \frac{x^2}{2} + \frac{x^3}{3}$$

input `int((x + x^3)/(x - 1),x)`output `2*x + 2*log(x - 1) + x^2/2 + x^3/3`

3.155 $\int \frac{-1+2x+x^2}{-2x+3x^2+2x^3} dx$

3.155.1 Optimal result	845
3.155.2 Mathematica [A] (verified)	845
3.155.3 Rubi [A] (verified)	846
3.155.4 Maple [A] (verified)	847
3.155.5 Fricas [A] (verification not implemented)	847
3.155.6 Sympy [A] (verification not implemented)	847
3.155.7 Maxima [A] (verification not implemented)	848
3.155.8 Giac [A] (verification not implemented)	848
3.155.9 Mupad [B] (verification not implemented)	848

3.155.1 Optimal result

Integrand size = 25, antiderivative size = 25

$$\int \frac{-1+2x+x^2}{-2x+3x^2+2x^3} dx = \frac{1}{10} \log(1-2x) + \frac{\log(x)}{2} - \frac{1}{10} \log(2+x)$$

output `1/10*ln(1-2*x)+1/2*ln(x)-1/10*ln(2+x)`

3.155.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{-1+2x+x^2}{-2x+3x^2+2x^3} dx = \frac{1}{10} \log(1-2x) + \frac{\log(x)}{2} - \frac{1}{10} \log(2+x)$$

input `Integrate[(-1 + 2*x + x^2)/(-2*x + 3*x^2 + 2*x^3),x]`

output `Log[1 - 2*x]/10 + Log[x]/2 - Log[2 + x]/10`

3.155.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2026, 2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^2 + 2x - 1}{2x^3 + 3x^2 - 2x} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{x^2 + 2x - 1}{x(2x^2 + 3x - 2)} dx \\ & \quad \downarrow \text{2159} \\ & \int \left(-\frac{1}{10(x+2)} + \frac{1}{5(2x-1)} + \frac{1}{2x} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{1}{10} \log(1-2x) + \frac{\log(x)}{2} - \frac{1}{10} \log(x+2) \end{aligned}$$

input `Int[(-1 + 2*x + x^2)/(-2*x + 3*x^2 + 2*x^3), x]`

output `Log[1 - 2*x]/10 + Log[x]/2 - Log[2 + x]/10`

3.155.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2159 `Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

3.155. $\int \frac{-1+2x+x^2}{-2x+3x^2+2x^3} dx$

3.155.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.72

method	result	size
parallelrisch	$\frac{\ln(x)}{2} - \frac{\ln(2+x)}{10} + \frac{\ln(x-\frac{1}{2})}{10}$	18
default	$-\frac{\ln(2+x)}{10} + \frac{\ln(x)}{2} + \frac{\ln(2x-1)}{10}$	20
norman	$-\frac{\ln(2+x)}{10} + \frac{\ln(x)}{2} + \frac{\ln(2x-1)}{10}$	20
risch	$-\frac{\ln(2+x)}{10} + \frac{\ln(x)}{2} + \frac{\ln(2x-1)}{10}$	20

input `int((x^2+2*x-1)/(2*x^3+3*x^2-2*x),x,method=_RETURNVERBOSE)`output `1/2*ln(x)-1/10*ln(2+x)+1/10*ln(x-1/2)`**3.155.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{-1 + 2x + x^2}{-2x + 3x^2 + 2x^3} dx = \frac{1}{10} \log(2x - 1) - \frac{1}{10} \log(x + 2) + \frac{1}{2} \log(x)$$

input `integrate((x^2+2*x-1)/(2*x^3+3*x^2-2*x),x, algorithm="fricas")`output `1/10*log(2*x - 1) - 1/10*log(x + 2) + 1/2*log(x)`**3.155.6 Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{-1 + 2x + x^2}{-2x + 3x^2 + 2x^3} dx = \frac{\log(x)}{2} + \frac{\log(x - \frac{1}{2})}{10} - \frac{\log(x + 2)}{10}$$

input `integrate((x**2+2*x-1)/(2*x**3+3*x**2-2*x),x)`output `log(x)/2 + log(x - 1/2)/10 - log(x + 2)/10`

3.155. $\int \frac{-1+2x+x^2}{-2x+3x^2+2x^3} dx$

3.155.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{-1 + 2x + x^2}{-2x + 3x^2 + 2x^3} dx = \frac{1}{10} \log(2x - 1) - \frac{1}{10} \log(x + 2) + \frac{1}{2} \log(x)$$

input `integrate((x^2+2*x-1)/(2*x^3+3*x^2-2*x),x, algorithm="maxima")`output `1/10*log(2*x - 1) - 1/10*log(x + 2) + 1/2*log(x)`**3.155.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{-1 + 2x + x^2}{-2x + 3x^2 + 2x^3} dx = \frac{1}{10} \log(|2x - 1|) - \frac{1}{10} \log(|x + 2|) + \frac{1}{2} \log(|x|)$$

input `integrate((x^2+2*x-1)/(2*x^3+3*x^2-2*x),x, algorithm="giac")`output `1/10*log(abs(2*x - 1)) - 1/10*log(abs(x + 2)) + 1/2*log(abs(x))`**3.155.9 Mupad [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{-1 + 2x + x^2}{-2x + 3x^2 + 2x^3} dx = \frac{\operatorname{atanh}\left(\frac{24}{145\left(\frac{29x}{100} - \frac{11}{50}\right)} + \frac{35}{29}\right)}{5} + \frac{\ln(x)}{2}$$

input `int((2*x + x^2 - 1)/(3*x^2 - 2*x + 2*x^3),x)`output `atanh(24/(145*((29*x)/100 - 11/50)) + 35/29)/5 + log(x)/2`

3.156 $\int \frac{1+4x-2x^2+x^4}{1-x-x^2+x^3} dx$

3.156.1 Optimal result	849
3.156.2 Mathematica [A] (verified)	849
3.156.3 Rubi [A] (verified)	850
3.156.4 Maple [A] (verified)	851
3.156.5 Fricas [A] (verification not implemented)	851
3.156.6 Sympy [A] (verification not implemented)	851
3.156.7 Maxima [A] (verification not implemented)	852
3.156.8 Giac [A] (verification not implemented)	852
3.156.9 Mupad [B] (verification not implemented)	852

3.156.1 Optimal result

Integrand size = 29, antiderivative size = 30

$$\int \frac{1+4x-2x^2+x^4}{1-x-x^2+x^3} dx = \frac{2}{1-x} + x + \frac{x^2}{2} + \log(1-x) - \log(1+x)$$

output `2/(1-x)+x+1/2*x^2+ln(1-x)-ln(1+x)`

3.156.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.97

$$\int \frac{1+4x-2x^2+x^4}{1-x-x^2+x^3} dx = -\frac{2}{-1+x} + \frac{1}{2}(1+x)^2 + \log(1-x) - \log(1+x)$$

input `Integrate[(1 + 4*x - 2*x^2 + x^4)/(1 - x - x^2 + x^3),x]`

output `-2/(-1 + x) + (1 + x)^2/2 + Log[1 - x] - Log[1 + x]`

3.156.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.069$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 - 2x^2 + 4x + 1}{x^3 - x^2 - x + 1} dx$$

$$\downarrow \text{2462}$$

$$\int \left(x + \frac{1}{-x-1} + \frac{1}{x-1} + \frac{2}{(x-1)^2} + 1 \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^2}{2} + x + \frac{2}{1-x} + \log(1-x) - \log(x+1)$$

input `Int[(1 + 4*x - 2*x^2 + x^4)/(1 - x - x^2 + x^3),x]`

output `2/(1 - x) + x + x^2/2 + Log[1 - x] - Log[1 + x]`

3.156.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2462 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ [Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

3.156.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.83

method	result	size
default	$\ln(-1+x) - \frac{2}{-1+x} + x + \frac{x^2}{2} - \ln(1+x)$	25
risch	$\ln(-1+x) - \frac{2}{-1+x} + x + \frac{x^2}{2} - \ln(1+x)$	25
norman	$\frac{\frac{1}{2}x^2 + \frac{1}{2}x^3 - 3}{-1+x} - \ln(1+x) + \ln(-1+x)$	30
parallelrisch	$\frac{x^3 + 2\ln(-1+x)x - 2\ln(1+x)x + x^2 - 6 - 2\ln(-1+x) + 2\ln(1+x)}{-2+2x}$	42

input `int((x^4-2*x^2+4*x+1)/(x^3-x^2-x+1),x,method=_RETURNVERBOSE)`output `ln(-1+x)-2/(-1+x)+x+1/2*x^2-ln(1+x)`**3.156.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.20

$$\int \frac{1+4x-2x^2+x^4}{1-x-x^2+x^3} dx = \frac{x^3+x^2-2(x-1)\log(x+1)+2(x-1)\log(x-1)-2x-4}{2(x-1)}$$

input `integrate((x^4-2*x^2+4*x+1)/(x^3-x^2-x+1),x, algorithm="fracas")`output `1/2*(x^3 + x^2 - 2*(x - 1)*log(x + 1) + 2*(x - 1)*log(x - 1) - 2*x - 4)/(x - 1)`**3.156.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.67

$$\int \frac{1+4x-2x^2+x^4}{1-x-x^2+x^3} dx = \frac{x^2}{2} + x + \log(x-1) - \log(x+1) - \frac{2}{x-1}$$

input `integrate((x**4-2*x**2+4*x+1)/(x**3-x**2-x+1),x)`output `x**2/2 + x + log(x - 1) - log(x + 1) - 2/(x - 1)`

3.156. $\int \frac{1+4x-2x^2+x^4}{1-x-x^2+x^3} dx$

3.156.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int \frac{1 + 4x - 2x^2 + x^4}{1 - x - x^2 + x^3} dx = \frac{1}{2}x^2 + x - \frac{2}{x-1} - \log(x+1) + \log(x-1)$$

input `integrate((x^4-2*x^2+4*x+1)/(x^3-x^2-x+1),x, algorithm="maxima")`output `1/2*x^2 + x - 2/(x - 1) - log(x + 1) + log(x - 1)`**3.156.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.87

$$\int \frac{1 + 4x - 2x^2 + x^4}{1 - x - x^2 + x^3} dx = \frac{1}{2}x^2 + x - \frac{2}{x-1} - \log(|x+1|) + \log(|x-1|)$$

input `integrate((x^4-2*x^2+4*x+1)/(x^3-x^2-x+1),x, algorithm="giac")`output `1/2*x^2 + x - 2/(x - 1) - log(abs(x + 1)) + log(abs(x - 1))`**3.156.9 Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \frac{1 + 4x - 2x^2 + x^4}{1 - x - x^2 + x^3} dx = x - \frac{2}{x-1} + \frac{x^2}{2} + \operatorname{atan}(x \operatorname{li} 2i)$$

input `int(-(4*x - 2*x^2 + x^4 + 1)/(x + x^2 - x^3 - 1),x)`output `x + atan(x*1i)*2i - 2/(x - 1) + x^2/2`

3.157 $\int \frac{4-x+2x^2}{4x+x^3} dx$

3.157.1 Optimal result	853
3.157.2 Mathematica [A] (verified)	853
3.157.3 Rubi [A] (verified)	854
3.157.4 Maple [A] (verified)	855
3.157.5 Fricas [A] (verification not implemented)	855
3.157.6 Sympy [A] (verification not implemented)	855
3.157.7 Maxima [A] (verification not implemented)	856
3.157.8 Giac [A] (verification not implemented)	856
3.157.9 Mupad [B] (verification not implemented)	856

3.157.1 Optimal result

Integrand size = 20, antiderivative size = 23

$$\int \frac{4-x+2x^2}{4x+x^3} dx = -\frac{1}{2} \arctan\left(\frac{x}{2}\right) + \log(x) + \frac{1}{2} \log(4+x^2)$$

output `-1/2*arctan(1/2*x)+ln(x)+1/2*ln(x^2+4)`

3.157.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{4-x+2x^2}{4x+x^3} dx = -\frac{1}{2} \arctan\left(\frac{x}{2}\right) + \log(x) + \frac{1}{2} \log(4+x^2)$$

input `Integrate[(4 - x + 2*x^2)/(4*x + x^3), x]`

output `-1/2*ArcTan[x/2] + Log[x] + Log[4 + x^2]/2`

3.157.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {2026, 2333, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{2x^2 - x + 4}{x^3 + 4x} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{2x^2 - x + 4}{x(x^2 + 4)} dx \\ & \quad \downarrow \text{2333} \\ & \int \left(\frac{x-1}{x^2+4} + \frac{1}{x} \right) dx \\ & \quad \downarrow \text{2009} \\ & -\frac{1}{2} \arctan\left(\frac{x}{2}\right) + \frac{1}{2} \log(x^2 + 4) + \log(x) \end{aligned}$$

input `Int[(4 - x + 2*x^2)/(4*x + x^3), x]`

output `-1/2*ArcTan[x/2] + Log[x] + Log[4 + x^2]/2`

3.157.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2333 `Int[(Pq_)*((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

3.157.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

method	result	size
default	$-\frac{\arctan(\frac{x}{2})}{2} + \ln(x) + \frac{\ln(x^2+4)}{2}$	18
risch	$-\frac{\arctan(\frac{x}{2})}{2} + \ln(x) + \frac{\ln(x^2+4)}{2}$	18
meijerg	$\ln(x) - \ln(2) + \frac{\ln(1+\frac{x^2}{4})}{2} - \frac{\arctan(\frac{x}{2})}{2}$	24
parallelrisch	$\ln(x) + \frac{\ln(x-2i)}{2} + \frac{i \ln(x-2i)}{4} + \frac{\ln(x+2i)}{2} - \frac{i \ln(x+2i)}{4}$	34

input `int((2*x^2-x+4)/(x^3+4*x),x,method=_RETURNVERBOSE)`output `-1/2*arctan(1/2*x)+ln(x)+1/2*ln(x^2+4)`**3.157.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{4-x+2x^2}{4x+x^3} dx = -\frac{1}{2} \arctan\left(\frac{1}{2}x\right) + \frac{1}{2} \log(x^2+4) + \log(x)$$

input `integrate((2*x^2-x+4)/(x^3+4*x),x, algorithm="fracas")`output `-1/2*arctan(1/2*x) + 1/2*log(x^2 + 4) + log(x)`**3.157.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{4-x+2x^2}{4x+x^3} dx = \log(x) + \frac{\log(x^2+4)}{2} - \frac{\operatorname{atan}\left(\frac{x}{2}\right)}{2}$$

input `integrate((2*x**2-x+4)/(x**3+4*x),x)`output `log(x) + log(x**2 + 4)/2 - atan(x/2)/2`

3.157.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{4-x+2x^2}{4x+x^3} dx = -\frac{1}{2} \arctan\left(\frac{1}{2}x\right) + \frac{1}{2} \log(x^2+4) + \log(x)$$

input `integrate((2*x^2-x+4)/(x^3+4*x),x, algorithm="maxima")`output `-1/2*arctan(1/2*x) + 1/2*log(x^2 + 4) + log(x)`**3.157.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int \frac{4-x+2x^2}{4x+x^3} dx = -\frac{1}{2} \arctan\left(\frac{1}{2}x\right) + \frac{1}{2} \log(x^2+4) + \log(|x|)$$

input `integrate((2*x^2-x+4)/(x^3+4*x),x, algorithm="giac")`output `-1/2*arctan(1/2*x) + 1/2*log(x^2 + 4) + log(abs(x))`**3.157.9 Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{4-x+2x^2}{4x+x^3} dx = \ln(x) + \ln(x-2i) \left(\frac{1}{2} + \frac{1}{4}i\right) + \ln(x+2i) \left(\frac{1}{2} - \frac{1}{4}i\right)$$

input `int((2*x^2 - x + 4)/(4*x + x^3),x)`output `log(x - 2i)*(1/2 + 1i/4) + log(x + 2i)*(1/2 - 1i/4) + log(x)`

3.158 $\int \frac{2-3x+4x^2}{3-4x+4x^2} dx$

3.158.1 Optimal result	857
3.158.2 Mathematica [A] (verified)	857
3.158.3 Rubi [A] (verified)	858
3.158.4 Maple [A] (verified)	859
3.158.5 Fricas [A] (verification not implemented)	859
3.158.6 Sympy [A] (verification not implemented)	859
3.158.7 Maxima [A] (verification not implemented)	860
3.158.8 Giac [A] (verification not implemented)	860
3.158.9 Mupad [B] (verification not implemented)	860

3.158.1 Optimal result

Integrand size = 23, antiderivative size = 38

$$\int \frac{2-3x+4x^2}{3-4x+4x^2} dx = x + \frac{\arctan\left(\frac{1-2x}{\sqrt{2}}\right)}{4\sqrt{2}} + \frac{1}{8} \log(3-4x+4x^2)$$

output `x+1/8*ln(4*x^2-4*x+3)+1/8*arctan(1/2*(1-2*x)*2^(1/2))*2^(1/2)`

3.158.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{2-3x+4x^2}{3-4x+4x^2} dx = x - \frac{\arctan\left(\frac{-1+2x}{\sqrt{2}}\right)}{4\sqrt{2}} + \frac{1}{8} \log(3-4x+4x^2)$$

input `Integrate[(2 - 3*x + 4*x^2)/(3 - 4*x + 4*x^2),x]`

output `x - ArcTan[(-1 + 2*x)/Sqrt[2]]/(4*Sqrt[2]) + Log[3 - 4*x + 4*x^2]/8`

3.158.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2188, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{4x^2 - 3x + 2}{4x^2 - 4x + 3} dx$$

↓ 2188

$$\int \left(1 - \frac{1-x}{4x^2 - 4x + 3} \right) dx$$

↓ 2009

$$\frac{\arctan\left(\frac{1-2x}{\sqrt{2}}\right)}{4\sqrt{2}} + \frac{1}{8} \log(4x^2 - 4x + 3) + x$$

input `Int[(2 - 3*x + 4*x^2)/(3 - 4*x + 4*x^2),x]`

output `x + ArcTan[(1 - 2*x)/Sqrt[2]]/(4*Sqrt[2]) + Log[3 - 4*x + 4*x^2]/8`

3.158.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2188 `Int[(Pq_)*((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[Pq*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

3.158.4 Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

method	result	size
default	$x + \frac{\ln(4x^2-4x+3)}{8} - \frac{\sqrt{2} \arctan\left(\frac{(8x-4)\sqrt{2}}{8}\right)}{8}$	32
risch	$x + \frac{\ln(4x^2-4x+3)}{8} - \frac{\sqrt{2} \arctan\left(\frac{(2x-1)\sqrt{2}}{2}\right)}{8}$	32

input `int((4*x^2-3*x+2)/(4*x^2-4*x+3),x,method=_RETURNVERBOSE)`output `x+1/8*ln(4*x^2-4*x+3)-1/8*2^(1/2)*arctan(1/8*(8*x-4)*2^(1/2))`**3.158.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int \frac{2-3x+4x^2}{3-4x+4x^2} dx = -\frac{1}{8} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(2x-1)\right) + x + \frac{1}{8} \log(4x^2-4x+3)$$

input `integrate((4*x^2-3*x+2)/(4*x^2-4*x+3),x, algorithm="fricas")`output `-1/8*sqrt(2)*arctan(1/2*sqrt(2)*(2*x - 1)) + x + 1/8*log(4*x^2 - 4*x + 3)`**3.158.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.89

$$\int \frac{2-3x+4x^2}{3-4x+4x^2} dx = x + \frac{\log\left(x^2-x+\frac{3}{4}\right)}{8} - \frac{\sqrt{2} \operatorname{atan}\left(\sqrt{2}x - \frac{\sqrt{2}}{2}\right)}{8}$$

input `integrate((4*x**2-3*x+2)/(4*x**2-4*x+3),x)`output `x + log(x**2 - x + 3/4)/8 - sqrt(2)*atan(sqrt(2)*x - sqrt(2)/2)/8`

3.158.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int \frac{2-3x+4x^2}{3-4x+4x^2} dx = -\frac{1}{8}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}(2x-1)\right) + x + \frac{1}{8}\log(4x^2-4x+3)$$

input `integrate((4*x^2-3*x+2)/(4*x^2-4*x+3),x, algorithm="maxima")`output `-1/8*sqrt(2)*arctan(1/2*sqrt(2)*(2*x - 1)) + x + 1/8*log(4*x^2 - 4*x + 3)`**3.158.8 Giac [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int \frac{2-3x+4x^2}{3-4x+4x^2} dx = -\frac{1}{8}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}(2x-1)\right) + x + \frac{1}{8}\log(4x^2-4x+3)$$

input `integrate((4*x^2-3*x+2)/(4*x^2-4*x+3),x, algorithm="giac")`output `-1/8*sqrt(2)*arctan(1/2*sqrt(2)*(2*x - 1)) + x + 1/8*log(4*x^2 - 4*x + 3)`**3.158.9 Mupad [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.79

$$\int \frac{2-3x+4x^2}{3-4x+4x^2} dx = x + \frac{\ln(x^2-x+\frac{3}{4})}{8} - \frac{\sqrt{2}\operatorname{atan}\left(\sqrt{2}x-\frac{\sqrt{2}}{2}\right)}{8}$$

input `int((4*x^2 - 3*x + 2)/(4*x^2 - 4*x + 3),x)`output `x + log(x^2 - x + 3/4)/8 - (2^(1/2)*atan(2^(1/2)*x - 2^(1/2)/2))/8`

3.159 $\int \frac{1+x^2+x^3}{(-1+x)x(1+x^2)^3(1+x+x^2)} dx$

3.159.1 Optimal result 861
 3.159.2 Mathematica [A] (verified) 861
 3.159.3 Rubi [A] (verified) 862
 3.159.4 Maple [A] (verified) 863
 3.159.5 Fracas [A] (verification not implemented) 863
 3.159.6 Sympy [A] (verification not implemented) 864
 3.159.7 Maxima [A] (verification not implemented) 864
 3.159.8 Giac [A] (verification not implemented) 865
 3.159.9 Mupad [B] (verification not implemented) 865

3.159.1 Optimal result

Integrand size = 32, antiderivative size = 103

$$\int \frac{1+x^2+x^3}{(-1+x)x(1+x^2)^3(1+x+x^2)} dx = \frac{1+x}{8(1+x^2)^2} - \frac{3(1-x)}{8(1+x^2)} + \frac{3x}{16(1+x^2)} + \frac{7 \arctan(x)}{16} - \frac{\arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{8} \log(1-x) - \log(x) + \frac{15}{16} \log(1+x^2) - \frac{1}{2} \log(1+x+x^2)$$

output `1/8*(1+x)/(x^2+1)^2-3/8*(1-x)/(x^2+1)+3/16*x/(x^2+1)+7/16*arctan(x)+1/8*ln(1-x)-ln(x)+15/16*ln(x^2+1)-1/2*ln(x^2+x+1)-1/3*arctan(1/3*(1+2*x)*3^(1/2))*3^(1/2)`

3.159.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.90

$$\int \frac{1+x^2+x^3}{(-1+x)x(1+x^2)^3(1+x+x^2)} dx = \frac{1}{48} \left(\frac{6(1+x)}{(1+x^2)^2} + \frac{9(-2+3x)}{1+x^2} + 21 \arctan(x) - 16\sqrt{3} \arctan\left(\frac{1+2x}{\sqrt{3}}\right) + 20 \log(1-x) - 48 \log(x) + 45 \log(1+x^2) - 10 \log(1+x+x^2) - 14 \log(1-x^3) \right)$$

input `Integrate[(1 + x^2 + x^3)/((-1 + x)*x*(1 + x^2)^3*(1 + x + x^2)),x]`

output `((6*(1 + x))/(1 + x^2)^2 + (9*(-2 + 3*x))/(1 + x^2) + 21*ArcTan[x] - 16*sqrt[3]*ArcTan[(1 + 2*x)/sqrt[3]] + 20*Log[1 - x] - 48*Log[x] + 45*Log[1 + x^2] - 10*Log[1 + x + x^2] - 14*Log[1 - x^3])/48`

3.159.3 Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.062$, Rules used = {7279, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 + x^2 + 1}{(x-1)x(x^2+1)^3(x^2+x+1)} dx$$

↓ 7279

$$\int \left(\frac{-x-1}{x^2+x+1} + \frac{15x-1}{8(x^2+1)} + \frac{3(x+1)}{4(x^2+1)^2} + \frac{1-x}{2(x^2+1)^3} + \frac{1}{8(x-1)} - \frac{1}{x} \right) dx$$

↓ 2009

$$\frac{7 \arctan(x)}{16} - \frac{\arctan\left(\frac{2x+1}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{3(1-x)}{8(x^2+1)} + \frac{3x}{16(x^2+1)} + \frac{x+1}{8(x^2+1)^2} + \frac{15}{16} \log(x^2+1) - \frac{1}{2} \log(x^2+x+1) + \frac{1}{8} \log(1-x) - \log(x)$$

input `Int[(1 + x^2 + x^3)/((-1 + x)*x*(1 + x^2)^3*(1 + x + x^2)),x]`

output `(1 + x)/(8*(1 + x^2)^2) - (3*(1 - x))/(8*(1 + x^2)) + (3*x)/(16*(1 + x^2)) + (7*ArcTan[x])/16 - ArcTan[(1 + 2*x)/sqrt[3]]/sqrt[3] + Log[1 - x]/8 - Log[x] + (15*Log[1 + x^2])/16 - Log[1 + x + x^2]/2`

3.159.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7279 `Int[(u_)/((a_.) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.)), x_Symbol] := With[
{v = RationalFunctionExpand[u/(a + b*x^n + c*x^(2*n)), x]}, Int[v, x] /; Su
mQ[v]] /; FreeQ[{a, b, c}, x] && EqQ[n2, 2*n] && IGtQ[n, 0]`

3.159.4 Maple [A] (verified)

Time = 0.45 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.70

method	result
risch	$\frac{\frac{9}{16}x^3 - \frac{3}{8}x^2 + \frac{11}{16}x - \frac{1}{4}}{(x^2+1)^2} + \frac{\ln(-1+x)}{8} + \frac{15 \ln(49x^2+49)}{16} + \frac{7 \arctan(x)}{16} - \frac{\sqrt{3} \arctan\left(\frac{2\sqrt{3}\left(x+\frac{1}{2}\right)}{3}\right)}{3} - \frac{\ln(x^2+x+1)}{2} - \ln(x)$
default	$\frac{\ln(-1+x)}{8} + \frac{\frac{9}{2}x^3 - 3x^2 + \frac{11}{2}x - 2}{8(x^2+1)^2} + \frac{15 \ln(x^2+1)}{16} + \frac{7 \arctan(x)}{16} - \frac{\ln(x^2+x+1)}{2} - \frac{\arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3} - \ln(x)$

input `int((x^3+x^2+1)/(-1+x)/x/(x^2+1)^3/(x^2+x+1),x,method=_RETURNVERBOSE)`

output
$$\frac{(9/16*x^3-3/8*x^2+11/16*x-1/4)/(x^2+1)^2+1/8*\ln(-1+x)+15/16*\ln(49*x^2+49)+7/16*\arctan(x)-1/3*3^{(1/2)}*\arctan(2/3*3^{(1/2)}*(x+1/2))-1/2*\ln(x^2+x+1)-\ln(x)}{x}$$

3.159.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.32

$$\int \frac{1+x^2+x^3}{(-1+x)x(1+x^2)^3(1+x+x^2)} dx$$

$$= \frac{27x^3 - 16\sqrt{3}(x^4 + 2x^2 + 1) \arctan\left(\frac{1}{3}\sqrt{3}(2x+1)\right) - 18x^2 + 21(x^4 + 2x^2 + 1) \arctan(x) - 24(x^4 + 2x^2 + 1) \arctan\left(\frac{2\sqrt{3}\left(x+\frac{1}{2}\right)}{3}\right)}{(x^2+1)^3(x^2+x+1)}$$

input `integrate((x^3+x^2+1)/(-1+x)/x/(x^2+1)^3/(x^2+x+1),x, algorithm="fracas")`

3.159.
$$\int \frac{1+x^2+x^3}{(-1+x)x(1+x^2)^3(1+x+x^2)} dx$$

output $1/48*(27*x^3 - 16*\sqrt{3}*(x^4 + 2*x^2 + 1)*\arctan(1/3*\sqrt{3}*(2*x + 1)) - 18*x^2 + 21*(x^4 + 2*x^2 + 1)*\arctan(x) - 24*(x^4 + 2*x^2 + 1)*\log(x^2 + x + 1) + 45*(x^4 + 2*x^2 + 1)*\log(x^2 + 1) + 6*(x^4 + 2*x^2 + 1)*\log(x - 1) - 48*(x^4 + 2*x^2 + 1)*\log(x) + 33*x - 12)/(x^4 + 2*x^2 + 1)$

3.159.6 Sympy [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.85

$$\int \frac{1 + x^2 + x^3}{(-1 + x)x(1 + x^2)^3(1 + x + x^2)} dx = -\log(x) + \frac{\log(x - 1)}{8} + \frac{15 \log(x^2 + 1)}{16} - \frac{\log(x^2 + x + 1)}{2} + \frac{7 \operatorname{atan}(x)}{16} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x}{3} + \frac{\sqrt{3}}{3}\right)}{3} + \frac{9x^3 - 6x^2 + 11x - 4}{16x^4 + 32x^2 + 16}$$

input `integrate((x**3+x**2+1)/(-1+x)/x/(x**2+1)**3/(x**2+x+1),x)`

output $-\log(x) + \log(x - 1)/8 + 15*\log(x**2 + 1)/16 - \log(x**2 + x + 1)/2 + 7*\operatorname{atan}(x)/16 - \sqrt{3}*\operatorname{atan}(2*\sqrt{3}*x/3 + \sqrt{3}/3)/3 + (9*x**3 - 6*x**2 + 11*x - 4)/(16*x**4 + 32*x**2 + 16)$

3.159.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.75

$$\int \frac{1 + x^2 + x^3}{(-1 + x)x(1 + x^2)^3(1 + x + x^2)} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x + 1)\right) + \frac{9x^3 - 6x^2 + 11x - 4}{16(x^4 + 2x^2 + 1)} + \frac{7}{16} \arctan(x) - \frac{1}{2} \log(x^2 + x + 1) + \frac{15}{16} \log(x^2 + 1) + \frac{1}{8} \log(x - 1) - \log(x)$$

input `integrate((x^3+x^2+1)/(-1+x)/x/(x^2+1)^3/(x^2+x+1),x, algorithm="maxima")`

output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + 1/16*(9*x^3 - 6*x^2 + 11*x - 4)/(x^4 + 2*x^2 + 1) + 7/16*arctan(x) - 1/2*log(x^2 + x + 1) + 15/16*log(x^2 + 1) + 1/8*log(x - 1) - log(x)`

3.159.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.72

$$\int \frac{1 + x^2 + x^3}{(-1 + x)x(1 + x^2)^3(1 + x + x^2)} dx = -\frac{1}{3}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}(2x + 1)\right) + \frac{9x^3 - 6x^2 + 11x - 4}{16(x^2 + 1)^2} + \frac{7}{16}\arctan(x) - \frac{1}{2}\log(x^2 + x + 1) + \frac{15}{16}\log(x^2 + 1) + \frac{1}{8}\log(|x - 1|) - \log(|x|)$$

input `integrate((x^3+x^2+1)/(-1+x)/x/(x^2+1)^3/(x^2+x+1),x, algorithm="giac")`

output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + 1/16*(9*x^3 - 6*x^2 + 11*x - 4)/(x^2 + 1)^2 + 7/16*arctan(x) - 1/2*log(x^2 + x + 1) + 15/16*log(x^2 + 1) + 1/8*log(abs(x - 1)) - log(abs(x))`

3.159.9 Mupad [B] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.93

$$\int \frac{1 + x^2 + x^3}{(-1 + x)x(1 + x^2)^3(1 + x + x^2)} dx = \frac{\ln(x - 1)}{8} - \ln(x) + \ln\left(x + \frac{1}{2} - \frac{\sqrt{3}1i}{2}\right)\left(-\frac{1}{2} + \frac{\sqrt{3}1i}{6}\right) - \ln\left(x + \frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)\left(\frac{1}{2} + \frac{\sqrt{3}1i}{6}\right) + \frac{\frac{9x^3}{16} - \frac{3x^2}{8} + \frac{11x}{16} - \frac{1}{4}}{x^4 + 2x^2 + 1} + \ln(x - i)\left(\frac{15}{16} - \frac{7}{32}i\right) + \ln(x + 1i)\left(\frac{15}{16} + \frac{7}{32}i\right)$$

input `int((x^2 + x^3 + 1)/(x*(x^2 + 1)^3*(x - 1)*(x + x^2 + 1)),x)`

output `log(x - 1)/8 + log(x - 1i)*(15/16 - 7i/32) + log(x + 1i)*(15/16 + 7i/32) -
log(x) + log(x - (3^(1/2)*1i)/2 + 1/2)*((3^(1/2)*1i)/6 - 1/2) - log(x + (
3^(1/2)*1i)/2 + 1/2)*((3^(1/2)*1i)/6 + 1/2) + ((11*x)/16 - (3*x^2)/8 + (9*
x^3)/16 - 1/4)/(2*x^2 + x^4 + 1)`

3.159. $\int \frac{1+x^2+x^3}{(-1+x)x(1+x^2)^3(1+x+x^2)} dx$

$$3.160 \quad \int \frac{1-3x+2x^2-x^3}{x(1+x^2)^2} dx$$

3.160.1 Optimal result	867
3.160.2 Mathematica [A] (verified)	867
3.160.3 Rubi [A] (verified)	868
3.160.4 Maple [A] (verified)	869
3.160.5 Fricas [A] (verification not implemented)	869
3.160.6 Sympy [A] (verification not implemented)	870
3.160.7 Maxima [A] (verification not implemented)	870
3.160.8 Giac [A] (verification not implemented)	870
3.160.9 Mupad [B] (verification not implemented)	871

3.160.1 Optimal result

Integrand size = 26, antiderivative size = 33

$$\int \frac{1-3x+2x^2-x^3}{x(1+x^2)^2} dx = -\frac{1+2x}{2(1+x^2)} - 2 \arctan(x) + \log(x) - \frac{1}{2} \log(1+x^2)$$

output `1/2*(-2*x-1)/(x^2+1)-2*arctan(x)+ln(x)-1/2*ln(x^2+1)`

3.160.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \frac{1-3x+2x^2-x^3}{x(1+x^2)^2} dx = \frac{-1-2x}{2(1+x^2)} - 2 \arctan(x) + \log(x) - \frac{1}{2} \log(1+x^2)$$

input `Integrate[(1 - 3*x + 2*x^2 - x^3)/(x*(1 + x^2)^2),x]`

output `(-1 - 2*x)/(2*(1 + x^2)) - 2*ArcTan[x] + Log[x] - Log[1 + x^2]/2`

3.160.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {2336, 27, 523, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{-x^3 + 2x^2 - 3x + 1}{x(x^2 + 1)^2} dx \\ & \quad \downarrow \text{2336} \\ & -\frac{1}{2} \int \frac{2(1-2x)}{x(x^2+1)} dx - \frac{2x+1}{2(x^2+1)} \\ & \quad \downarrow \text{27} \\ & \int \frac{1-2x}{x(x^2+1)} dx - \frac{2x+1}{2(x^2+1)} \\ & \quad \downarrow \text{523} \\ & \int \left(\frac{-x-2}{x^2+1} + \frac{1}{x} \right) dx - \frac{2x+1}{2(x^2+1)} \\ & \quad \downarrow \text{2009} \\ & -2 \arctan(x) - \frac{2x+1}{2(x^2+1)} - \frac{1}{2} \log(x^2+1) + \log(x) \end{aligned}$$

input `Int[(1 - 3*x + 2*x^2 - x^3)/(x*(1 + x^2)^2),x]`

output `-1/2*(1 + 2*x)/(1 + x^2) - 2*ArcTan[x] + Log[x] - Log[1 + x^2]/2`

3.160.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 523 `Int[((x_)^(m_.)*((c_) + (d_.)*(x_)))/((a_) + (b_.)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[x^m*((c + d*x)/(a + b*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && IntegerQ[m]`

3.160. $\int \frac{1-3x+2x^2-x^3}{x(1+x^2)^2} dx$

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2336 `Int[(Pq_)*((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[
{Q = PolynomialQuotient[(c*x)^m*Pq, a + b*x^2, x], f = Coeff[PolynomialRema
inder[(c*x)^m*Pq, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[(c*x)
^m*Pq, a + b*x^2, x], x, 1]}, Simp[(a*g - b*f*x)*((a + b*x^2)^(p + 1)/(2*a*
b*(p + 1))), x] + Simp[1/(2*a*(p + 1)) Int[(c*x)^m*(a + b*x^2)^(p + 1)*Ex
pandToSum[(2*a*(p + 1)*Q)/(c*x)^m + (f*(2*p + 3))/(c*x)^m, x], x]] /; F
reeQ[{a, b, c}, x] && PolyQ[Pq, x] && LtQ[p, -1] && ILtQ[m, 0]`

3.160.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.85

method	result
default	$-\frac{x+\frac{1}{2}}{x^2+1} - \frac{\ln(x^2+1)}{2} - 2 \arctan(x) + \ln(x)$
risch	$-\frac{x-\frac{1}{2}}{x^2+1} - \frac{\ln(x^2+1)}{2} - 2 \arctan(x) + \ln(x)$
meijerg	$-\frac{2x}{2x^2+2} - 2 \arctan(x) + \frac{x^2}{x^2+1} + \frac{1}{2} + \ln(x) - \frac{x^2}{2x^2+2} - \frac{\ln(x^2+1)}{2}$
parallelrisch	$\frac{2i \ln(x-i)x^2 - 2i \ln(x+i)x^2 + 2x^2 \ln(x) - x^2 \ln(x-i) - \ln(x+i)x^2 - 1 + 2i \ln(x-i) - 2i \ln(x+i) + 2 \ln(x) - \ln(x-i) - \ln(x+i) - 2x}{2x^2+2}$

input `int((-x^3+2*x^2-3*x+1)/x/(x^2+1)^2,x,method=_RETURNVERBOSE)`

output `-(x+1/2)/(x^2+1)-1/2*ln(x^2+1)-2*arctan(x)+ln(x)`

3.160.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.33

$$\int \frac{1 - 3x + 2x^2 - x^3}{x(1+x^2)^2} dx$$

$$= -\frac{4(x^2+1) \arctan(x) + (x^2+1) \log(x^2+1) - 2(x^2+1) \log(x) + 2x+1}{2(x^2+1)}$$

input `integrate((-x^3+2*x^2-3*x+1)/x/(x^2+1)^2,x, algorithm="fricas")`

output $-1/2*(4*(x^2 + 1)*\arctan(x) + (x^2 + 1)*\log(x^2 + 1) - 2*(x^2 + 1)*\log(x) + 2*x + 1)/(x^2 + 1)$

3.160.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.82

$$\int \frac{1 - 3x + 2x^2 - x^3}{x(1 + x^2)^2} dx = -\frac{2x + 1}{2x^2 + 2} + \log(x) - \frac{\log(x^2 + 1)}{2} - 2 \operatorname{atan}(x)$$

input `integrate((-x**3+2*x**2-3*x+1)/x/(x**2+1)**2,x)`

output $-(2*x + 1)/(2*x**2 + 2) + \log(x) - \log(x**2 + 1)/2 - 2*\operatorname{atan}(x)$

3.160.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.88

$$\int \frac{1 - 3x + 2x^2 - x^3}{x(1 + x^2)^2} dx = -\frac{2x + 1}{2(x^2 + 1)} - 2 \arctan(x) - \frac{1}{2} \log(x^2 + 1) + \log(x)$$

input `integrate((-x^3+2*x^2-3*x+1)/x/(x^2+1)^2,x, algorithm="maxima")`

output $-1/2*(2*x + 1)/(x^2 + 1) - 2*\arctan(x) - 1/2*\log(x^2 + 1) + \log(x)$

3.160.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.91

$$\int \frac{1 - 3x + 2x^2 - x^3}{x(1 + x^2)^2} dx = -\frac{2x + 1}{2(x^2 + 1)} - 2 \arctan(x) - \frac{1}{2} \log(x^2 + 1) + \log(|x|)$$

input `integrate((-x^3+2*x^2-3*x+1)/x/(x^2+1)^2,x, algorithm="giac")`

output $-1/2*(2*x + 1)/(x^2 + 1) - 2*\arctan(x) - 1/2*\log(x^2 + 1) + \log(\operatorname{abs}(x))$

3.160.9 Mupad [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \frac{1 - 3x + 2x^2 - x^3}{x(1 + x^2)^2} dx = \ln(x) - \frac{x + \frac{1}{2}}{x^2 + 1} + \ln(x - i) \left(-\frac{1}{2} + i\right) + \ln(x + i) \left(-\frac{1}{2} - i\right)$$

input `int(-(3*x - 2*x^2 + x^3 - 1)/(x*(x^2 + 1)^2),x)`output `log(x) - log(x + 1i)*(1/2 + 1i) - log(x - 1i)*(1/2 - 1i) - (x + 1/2)/(x^2 + 1)`

3.161 $\int \frac{1}{(1+x^2)^2} dx$

3.161.1 Optimal result	872
3.161.2 Mathematica [A] (verified)	872
3.161.3 Rubi [A] (verified)	873
3.161.4 Maple [A] (verified)	874
3.161.5 Fricas [A] (verification not implemented)	874
3.161.6 Sympy [A] (verification not implemented)	874
3.161.7 Maxima [A] (verification not implemented)	875
3.161.8 Giac [A] (verification not implemented)	875
3.161.9 Mupad [B] (verification not implemented)	875

3.161.1 Optimal result

Integrand size = 7, antiderivative size = 19

$$\int \frac{1}{(1+x^2)^2} dx = \frac{x}{2(1+x^2)} + \frac{\arctan(x)}{2}$$

output `1/2*x/(x^2+1)+1/2*arctan(x)`

3.161.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{1}{(1+x^2)^2} dx = \frac{1}{2} \left(\frac{x}{1+x^2} + \arctan(x) \right)$$

input `Integrate[(1 + x^2)^(-2),x]`

output `(x/(1 + x^2) + ArcTan[x])/2`

3.161.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {215, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x^2 + 1)^2} dx$$

↓ 215

$$\frac{1}{2} \int \frac{1}{x^2 + 1} dx + \frac{x}{2(x^2 + 1)}$$

↓ 216

$$\frac{\arctan(x)}{2} + \frac{x}{2(x^2 + 1)}$$

input `Int[(1 + x^2)^(-2), x]`

output `x/(2*(1 + x^2)) + ArcTan[x]/2`

3.161.3.1 Defintions of rubi rules used

rule 215 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

3.161.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

method	result	size
default	$\frac{x}{2x^2+2} + \frac{\arctan(x)}{2}$	16
risch	$\frac{x}{2x^2+2} + \frac{\arctan(x)}{2}$	16
meijerg	$\frac{x}{2x^2+2} + \frac{\arctan(x)}{2}$	17
parallelrisch	$-\frac{i \ln(x-i)x^2 - i \ln(x+i)x^2 + i \ln(x-i) - i \ln(x+i) - 2x}{4(x^2+1)}$	52

input `int(1/(x^2+1)^2,x,method=_RETURNVERBOSE)`output `1/2*x/(x^2+1)+1/2*arctan(x)`**3.161.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x^2)^2} dx = \frac{(x^2+1)\arctan(x)+x}{2(x^2+1)}$$

input `integrate(1/(x^2+1)^2,x, algorithm="fricas")`output `1/2*((x^2 + 1)*arctan(x) + x)/(x^2 + 1)`**3.161.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{1}{(1+x^2)^2} dx = \frac{x}{2x^2+2} + \frac{\operatorname{atan}(x)}{2}$$

input `integrate(1/(x**2+1)**2,x)`output `x/(2*x**2 + 2) + atan(x)/2`

3.161.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{(1+x^2)^2} dx = \frac{x}{2(x^2+1)} + \frac{1}{2} \arctan(x)$$

input `integrate(1/(x^2+1)^2,x, algorithm="maxima")`output `1/2*x/(x^2 + 1) + 1/2*arctan(x)`**3.161.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{(1+x^2)^2} dx = \frac{x}{2(x^2+1)} + \frac{1}{2} \arctan(x)$$

input `integrate(1/(x^2+1)^2,x, algorithm="giac")`output `1/2*x/(x^2 + 1) + 1/2*arctan(x)`**3.161.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{1}{(1+x^2)^2} dx = \frac{\operatorname{atan}(x)}{2} + \frac{x}{2(x^2+1)}$$

input `int(1/(x^2 + 1)^2,x)`output `atan(x)/2 + x/(2*(x^2 + 1))`

3.162 $\int \frac{1}{(-1+x)(2+x)} dx$

3.162.1 Optimal result	876
3.162.2 Mathematica [A] (verified)	876
3.162.3 Rubi [A] (verified)	877
3.162.4 Maple [A] (verified)	878
3.162.5 Fricas [A] (verification not implemented)	878
3.162.6 Sympy [A] (verification not implemented)	878
3.162.7 Maxima [A] (verification not implemented)	879
3.162.8 Giac [A] (verification not implemented)	879
3.162.9 Mupad [B] (verification not implemented)	879

3.162.1 Optimal result

Integrand size = 11, antiderivative size = 19

$$\int \frac{1}{(-1+x)(2+x)} dx = \frac{1}{3} \log(1-x) - \frac{1}{3} \log(2+x)$$

output `1/3*ln(1-x)-1/3*ln(2+x)`

3.162.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{1}{(-1+x)(2+x)} dx = \frac{1}{3} \log(1-x) - \frac{1}{3} \log(2+x)$$

input `Integrate[1/((-1 + x)*(2 + x)),x]`

output `Log[1 - x]/3 - Log[2 + x]/3`

3.162.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {47, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x-1)(x+2)} dx$$

$$\downarrow 47$$

$$\frac{1}{3} \int \frac{1}{x-1} dx - \frac{1}{3} \int \frac{1}{x+2} dx$$

$$\downarrow 16$$

$$\frac{1}{3} \log(1-x) - \frac{1}{3} \log(x+2)$$

input `Int[1/((-1 + x)*(2 + x)),x]`

output `Log[1 - x]/3 - Log[2 + x]/3`

3.162.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

3.162.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

method	result	size
default	$\frac{\ln(-1+x)}{3} - \frac{\ln(2+x)}{3}$	14
norman	$\frac{\ln(-1+x)}{3} - \frac{\ln(2+x)}{3}$	14
risch	$\frac{\ln(-1+x)}{3} - \frac{\ln(2+x)}{3}$	14
parallelrisch	$\frac{\ln(-1+x)}{3} - \frac{\ln(2+x)}{3}$	14

input `int(1/(-1+x)/(2+x),x,method=_RETURNVERBOSE)`

output `1/3*ln(-1+x)-1/3*ln(2+x)`

3.162.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int \frac{1}{(-1+x)(2+x)} dx = -\frac{1}{3} \log(x+2) + \frac{1}{3} \log(x-1)$$

input `integrate(1/(-1+x)/(2+x),x, algorithm="fricas")`

output `-1/3*log(x + 2) + 1/3*log(x - 1)`

3.162.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{1}{(-1+x)(2+x)} dx = \frac{\log(x-1)}{3} - \frac{\log(x+2)}{3}$$

input `integrate(1/(-1+x)/(2+x),x)`

output `log(x - 1)/3 - log(x + 2)/3`

3.162.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int \frac{1}{(-1+x)(2+x)} dx = -\frac{1}{3} \log(x+2) + \frac{1}{3} \log(x-1)$$

input `integrate(1/(-1+x)/(2+x),x, algorithm="maxima")`output `-1/3*log(x + 2) + 1/3*log(x - 1)`**3.162.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{(-1+x)(2+x)} dx = -\frac{1}{3} \log(|x+2|) + \frac{1}{3} \log(|x-1|)$$

input `integrate(1/(-1+x)/(2+x),x, algorithm="giac")`output `-1/3*log(abs(x + 2)) + 1/3*log(abs(x - 1))`**3.162.9 Mupad [B] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{1}{(-1+x)(2+x)} dx = \frac{\ln\left(\frac{x-1}{x+2}\right)}{3}$$

input `int(1/((x - 1)*(x + 2)),x)`output `log((x - 1)/(x + 2))/3`

3.163 $\int \frac{7}{-12+5x+2x^2} dx$

3.163.1 Optimal result	880
3.163.2 Mathematica [A] (verified)	880
3.163.3 Rubi [A] (verified)	881
3.163.4 Maple [A] (verified)	882
3.163.5 Fricas [A] (verification not implemented)	882
3.163.6 Sympy [A] (verification not implemented)	882
3.163.7 Maxima [A] (verification not implemented)	883
3.163.8 Giac [A] (verification not implemented)	883
3.163.9 Mupad [B] (verification not implemented)	883

3.163.1 Optimal result

Integrand size = 14, antiderivative size = 19

$$\int \frac{7}{-12+5x+2x^2} dx = \frac{7}{11} \log(3-2x) - \frac{7}{11} \log(4+x)$$

output `7/11*ln(3-2*x)-7/11*ln(4+x)`

3.163.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{7}{-12+5x+2x^2} dx = 7 \left(\frac{1}{11} \log(3-2x) - \frac{1}{11} \log(4+x) \right)$$

input `Integrate[7/(-12 + 5*x + 2*x^2), x]`

output `7*(Log[3 - 2*x]/11 - Log[4 + x]/11)`

3.163.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {27, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{7}{2x^2 + 5x - 12} dx \\ & \quad \downarrow \text{27} \\ & 7 \int \frac{1}{2x^2 + 5x - 12} dx \\ & \quad \downarrow \text{1081} \\ & 14 \int \left(-\frac{1}{22(x+4)} - \frac{1}{11(3-2x)} \right) dx \\ & \quad \downarrow \text{2009} \\ & 14 \left(\frac{1}{22} \log(3-2x) - \frac{1}{22} \log(x+4) \right) \end{aligned}$$

input `Int[7/(-12 + 5*x + 2*x^2),x]`

output `14*(Log[3 - 2*x]/22 - Log[4 + x]/22)`

3.163.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1081 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.163.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

method	result	size
parallelrisch	$-\frac{7\ln(4+x)}{11} + \frac{7\ln(x-\frac{3}{2})}{11}$	14
default	$-\frac{7\ln(4+x)}{11} + \frac{7\ln(2x-3)}{11}$	16
norman	$-\frac{7\ln(4+x)}{11} + \frac{7\ln(2x-3)}{11}$	16
risch	$-\frac{7\ln(4+x)}{11} + \frac{7\ln(2x-3)}{11}$	16

input `int(7/(2*x^2+5*x-12),x,method=_RETURNVERBOSE)`output `-7/11*ln(4+x)+7/11*ln(x-3/2)`**3.163.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{7}{-12 + 5x + 2x^2} dx = \frac{7}{11} \log(2x - 3) - \frac{7}{11} \log(x + 4)$$

input `integrate(7/(2*x^2+5*x-12),x, algorithm="fracas")`output `7/11*log(2*x - 3) - 7/11*log(x + 4)`**3.163.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{7}{-12 + 5x + 2x^2} dx = \frac{7 \log(x - \frac{3}{2})}{11} - \frac{7 \log(x + 4)}{11}$$

input `integrate(7/(2*x**2+5*x-12),x)`output `7*log(x - 3/2)/11 - 7*log(x + 4)/11`

3.163.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{7}{-12 + 5x + 2x^2} dx = \frac{7}{11} \log(2x - 3) - \frac{7}{11} \log(x + 4)$$

input `integrate(7/(2*x^2+5*x-12),x, algorithm="maxima")`output `7/11*log(2*x - 3) - 7/11*log(x + 4)`**3.163.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{7}{-12 + 5x + 2x^2} dx = \frac{7}{11} \log(|2x - 3|) - \frac{7}{11} \log(|x + 4|)$$

input `integrate(7/(2*x^2+5*x-12),x, algorithm="giac")`output `7/11*log(abs(2*x - 3)) - 7/11*log(abs(x + 4))`**3.163.9 Mupad [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.42

$$\int \frac{7}{-12 + 5x + 2x^2} dx = -\frac{14 \operatorname{atanh}\left(\frac{4x}{11} + \frac{5}{11}\right)}{11}$$

input `int(7/(5*x + 2*x^2 - 12),x)`output `-(14*atanh((4*x)/11 + 5/11))/11`

$$3.164 \quad \int \frac{-4+3x+x^2}{(-1+2x)^2(3+2x)} dx$$

3.164.1 Optimal result	884
3.164.2 Mathematica [A] (verified)	884
3.164.3 Rubi [A] (verified)	885
3.164.4 Maple [A] (verified)	886
3.164.5 Fricas [A] (verification not implemented)	886
3.164.6 Sympy [A] (verification not implemented)	886
3.164.7 Maxima [A] (verification not implemented)	887
3.164.8 Giac [A] (verification not implemented)	887
3.164.9 Mupad [B] (verification not implemented)	887

3.164.1 Optimal result

Integrand size = 23, antiderivative size = 32

$$\int \frac{-4+3x+x^2}{(-1+2x)^2(3+2x)} dx = -\frac{9}{32(1-2x)} + \frac{41}{128} \log(1-2x) - \frac{25}{128} \log(3+2x)$$

output `-9/32/(1-2*x)+41/128*ln(1-2*x)-25/128*ln(3+2*x)`

3.164.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \frac{-4+3x+x^2}{(-1+2x)^2(3+2x)} dx = \frac{9}{32(-1+2x)} + \frac{41}{128} \log(1-2x) - \frac{25}{128} \log(3+2x)$$

input `Integrate[(-4 + 3*x + x^2)/((-1 + 2*x)^2*(3 + 2*x)),x]`

output `9/(32*(-1 + 2*x)) + (41*Log[1 - 2*x])/128 - (25*Log[3 + 2*x])/128`

3.164.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2 + 3x - 4}{(2x - 1)^2(2x + 3)} dx$$

↓ 1195

$$\int \left(-\frac{25}{64(2x + 3)} + \frac{41}{64(2x - 1)} - \frac{9}{16(2x - 1)^2} \right) dx$$

↓ 2009

$$-\frac{9}{32(1 - 2x)} + \frac{41}{128} \log(1 - 2x) - \frac{25}{128} \log(2x + 3)$$

input `Int[(-4 + 3*x + x^2)/((-1 + 2*x)^2*(3 + 2*x)),x]`

output `-9/(32*(1 - 2*x)) + (41*Log[1 - 2*x])/128 - (25*Log[3 + 2*x])/128`

3.164.3.1 Defintions of rubi rules used

rule 1195 `Int[((d_.) + (e_.)*(x_)^(m_.))*((f_.) + (g_.)*(x_)^(n_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.164.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.78

method	result	size
risch	$\frac{9}{64(x-\frac{1}{2})} - \frac{25 \ln(3+2x)}{128} + \frac{41 \ln(2x-1)}{128}$	25
default	$-\frac{25 \ln(3+2x)}{128} + \frac{9}{32(2x-1)} + \frac{41 \ln(2x-1)}{128}$	27
norman	$\frac{9x}{16(2x-1)} - \frac{25 \ln(3+2x)}{128} + \frac{41 \ln(2x-1)}{128}$	28
parallelrisch	$\frac{82 \ln(x-\frac{1}{2})x - 50 \ln(\frac{3}{2}+x)x - 41 \ln(x-\frac{1}{2}) + 25 \ln(\frac{3}{2}+x) + 72x}{256x-128}$	40

input `int((x^2+3*x-4)/(2*x-1)^2/(3+2*x),x,method=_RETURNVERBOSE)`output `9/64/(x-1/2)-25/128*ln(3+2*x)+41/128*ln(2*x-1)`**3.164.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.16

$$\int \frac{-4 + 3x + x^2}{(-1 + 2x)^2(3 + 2x)} dx = -\frac{25(2x-1)\log(2x+3) - 41(2x-1)\log(2x-1) - 36}{128(2x-1)}$$

input `integrate((x^2+3*x-4)/(-1+2*x)^2/(3+2*x),x, algorithm="fricas")`output `-1/128*(25*(2*x - 1)*log(2*x + 3) - 41*(2*x - 1)*log(2*x - 1) - 36)/(2*x - 1)`**3.164.6 Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

$$\int \frac{-4 + 3x + x^2}{(-1 + 2x)^2(3 + 2x)} dx = \frac{41 \log(x - \frac{1}{2})}{128} - \frac{25 \log(x + \frac{3}{2})}{128} + \frac{9}{64x - 32}$$

input `integrate((x**2+3*x-4)/(-1+2*x)**2/(3+2*x),x)`output `41*log(x - 1/2)/128 - 25*log(x + 3/2)/128 + 9/(64*x - 32)`

3.164. $\int \frac{-4+3x+x^2}{(-1+2x)^2(3+2x)} dx$

3.164.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

$$\int \frac{-4 + 3x + x^2}{(-1 + 2x)^2(3 + 2x)} dx = \frac{9}{32(2x - 1)} - \frac{25}{128} \log(2x + 3) + \frac{41}{128} \log(2x - 1)$$

input `integrate((x^2+3*x-4)/(-1+2*x)^2/(3+2*x),x, algorithm="maxima")`output `9/32/(2*x - 1) - 25/128*log(2*x + 3) + 41/128*log(2*x - 1)`**3.164.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.34

$$\int \frac{-4 + 3x + x^2}{(-1 + 2x)^2(3 + 2x)} dx = \frac{9}{32(2x - 1)} - \frac{1}{8} \log\left(\frac{|2x - 1|}{2(2x - 1)^2}\right) - \frac{25}{128} \log\left(\left|-\frac{4}{2x - 1} - 1\right|\right)$$

input `integrate((x^2+3*x-4)/(-1+2*x)^2/(3+2*x),x, algorithm="giac")`output `9/32/(2*x - 1) - 1/8*log(1/2*abs(2*x - 1)/(2*x - 1)^2) - 25/128*log(abs(-4/(2*x - 1) - 1))`**3.164.9 Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.69

$$\int \frac{-4 + 3x + x^2}{(-1 + 2x)^2(3 + 2x)} dx = \frac{41 \ln(x - \frac{1}{2})}{128} - \frac{25 \ln(x + \frac{3}{2})}{128} + \frac{9}{64(x - \frac{1}{2})}$$

input `int((3*x + x^2 - 4)/((2*x - 1)^2*(2*x + 3)),x)`output `(41*log(x - 1/2))/128 - (25*log(x + 3/2))/128 + 9/(64*(x - 1/2))`

3.165 $\int \frac{-x^2+x^3}{(-6+x)(3+5x)^3} dx$

3.165.1 Optimal result	888
3.165.2 Mathematica [A] (verified)	888
3.165.3 Rubi [A] (verified)	889
3.165.4 Maple [A] (verified)	890
3.165.5 Fricas [A] (verification not implemented)	890
3.165.6 Sympy [A] (verification not implemented)	891
3.165.7 Maxima [A] (verification not implemented)	891
3.165.8 Giac [A] (verification not implemented)	891
3.165.9 Mupad [B] (verification not implemented)	892

3.165.1 Optimal result

Integrand size = 22, antiderivative size = 43

$$\int \frac{-x^2 + x^3}{(-6 + x)(3 + 5x)^3} dx = -\frac{12}{1375(3 + 5x)^2} + \frac{201}{15125(3 + 5x)} + \frac{20 \log(6 - x)}{3993} + \frac{1493 \log(3 + 5x)}{499125}$$

output `-12/1375/(3+5*x)^2+201/15125/(3+5*x)+20/3993*ln(6-x)+1493/499125*ln(3+5*x)`

3.165.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.77

$$\int \frac{-x^2 + x^3}{(-6 + x)(3 + 5x)^3} dx = \frac{99(157+335x)}{(3+5x)^2} + \frac{2500 \log(-6 + x) + 1493 \log(3 + 5x)}{499125}$$

input `Integrate[(-x^2 + x^3)/((-6 + x)*(3 + 5*x)^3),x]`

output `((99*(157 + 335*x))/(3 + 5*x)^2 + 2500*Log[-6 + x] + 1493*Log[3 + 5*x])/499125`

3.165.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$, Rules used = {2027, 165, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^3 - x^2}{(x-6)(5x+3)^3} dx \\ & \quad \downarrow \text{2027} \\ & \int \frac{(x-1)x^2}{(x-6)(5x+3)^3} dx \\ & \quad \downarrow \text{165} \\ & \int \left(\frac{1493}{99825(5x+3)} - \frac{201}{3025(5x+3)^2} + \frac{24}{275(5x+3)^3} + \frac{20}{3993(x-6)} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{201}{15125(5x+3)} - \frac{12}{1375(5x+3)^2} + \frac{20 \log(6-x)}{3993} + \frac{1493 \log(5x+3)}{499125} \end{aligned}$$

input `Int[(-x^2 + x^3)/((-6 + x)*(3 + 5*x)^3), x]`

output `-12/(1375*(3 + 5*x)^2) + 201/(15125*(3 + 5*x)) + (20*Log[6 - x])/3993 + (1493*Log[3 + 5*x])/499125`

3.165.3.1 Defintions of rubi rules used

rule 165 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p*(g + h*x), x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m}, x] && (IntegersQ[m, n, p] || (IGtQ[n, 0] && IGtQ[p, 0]))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2027 `Int[(Fx_.)*((a_.)*(x_)^(r_.) + (b_.)*(x_)^(s_.))^(p_.), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r))^p*Fx, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] && PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])`

3.165.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.70

method	result
risch	$\frac{201x + 471}{3025 + 15125} + \frac{20 \ln(-6+x)}{3993} + \frac{1493 \ln(3+5x)}{499125}$
norman	$\frac{-\frac{113}{3025}x - \frac{157}{1815}x^2}{(3+5x)^2} + \frac{20 \ln(-6+x)}{3993} + \frac{1493 \ln(3+5x)}{499125}$
default	$\frac{20 \ln(-6+x)}{3993} - \frac{12}{1375(3+5x)^2} + \frac{201}{15125(3+5x)} + \frac{1493 \ln(3+5x)}{499125}$
parallelrisch	$\frac{187500 \ln(-6+x)x^2 + 111975 \ln(x + \frac{3}{5})x^2 + 225000 \ln(-6+x)x + 134370 \ln(x + \frac{3}{5})x - 129525x^2 + 67500 \ln(-6+x) + 40311 \ln(x + \frac{3}{5})}{1497375(3+5x)^2}$

input `int((x^3-x^2)/(-6+x)/(3+5*x)^3,x,method=_RETURNVERBOSE)`

output `25*(201/75625*x+471/378125)/(3+5*x)^2+20/3993*ln(-6+x)+1493/499125*ln(3+5*x)`

3.165.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.23

$$\int \frac{-x^2 + x^3}{(-6+x)(3+5x)^3} dx$$

$$= \frac{1493(25x^2 + 30x + 9) \log(5x + 3) + 2500(25x^2 + 30x + 9) \log(x - 6) + 33165x + 15543}{499125(25x^2 + 30x + 9)}$$

input `integrate((x^3-x^2)/(-6+x)/(3+5*x)^3,x, algorithm="fricas")`

output `1/499125*(1493*(25*x^2 + 30*x + 9)*log(5*x + 3) + 2500*(25*x^2 + 30*x + 9)*log(x - 6) + 33165*x + 15543)/(25*x^2 + 30*x + 9)`

3.165. $\int \frac{-x^2+x^3}{(-6+x)(3+5x)^3} dx$

3.165.6 Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.74

$$\int \frac{-x^2 + x^3}{(-6 + x)(3 + 5x)^3} dx = \frac{1005x + 471}{378125x^2 + 453750x + 136125} + \frac{20 \log(x - 6)}{3993} + \frac{1493 \log(x + \frac{3}{5})}{499125}$$

input `integrate((x**3-x**2)/(-6+x)/(3+5*x)**3,x)`output `(1005*x + 471)/(378125*x**2 + 453750*x + 136125) + 20*log(x - 6)/3993 + 1493*log(x + 3/5)/499125`**3.165.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.79

$$\int \frac{-x^2 + x^3}{(-6 + x)(3 + 5x)^3} dx = \frac{3(335x + 157)}{15125(25x^2 + 30x + 9)} + \frac{1493}{499125} \log(5x + 3) + \frac{20}{3993} \log(x - 6)$$

input `integrate((x^3-x^2)/(-6+x)/(3+5*x)^3,x, algorithm="maxima")`output `3/15125*(335*x + 157)/(25*x^2 + 30*x + 9) + 1493/499125*log(5*x + 3) + 20/3993*log(x - 6)`**3.165.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.72

$$\int \frac{-x^2 + x^3}{(-6 + x)(3 + 5x)^3} dx = \frac{3(335x + 157)}{15125(5x + 3)^2} + \frac{1493}{499125} \log(|5x + 3|) + \frac{20}{3993} \log(|x - 6|)$$

input `integrate((x^3-x^2)/(-6+x)/(3+5*x)^3,x, algorithm="giac")`output `3/15125*(335*x + 157)/(5*x + 3)^2 + 1493/499125*log(abs(5*x + 3)) + 20/3993*log(abs(x - 6))`

3.165.9 Mupad [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.67

$$\int \frac{-x^2 + x^3}{(-6 + x)(3 + 5x)^3} dx = \frac{20 \ln(x - 6)}{3993} + \frac{1493 \ln\left(x + \frac{3}{5}\right)}{499125} + \frac{\frac{201x}{75625} + \frac{471}{378125}}{x^2 + \frac{6x}{5} + \frac{9}{25}}$$

input `int(-(x^2 - x^3)/((5*x + 3)^3*(x - 6)),x)`output `(20*log(x - 6))/3993 + (1493*log(x + 3/5))/499125 + ((201*x)/75625 + 471/378125)/((6*x)/5 + x^2 + 9/25)`

3.166 $\int \frac{1}{-x^3+x^4} dx$

3.166.1 Optimal result	893
3.166.2 Mathematica [A] (verified)	893
3.166.3 Rubi [A] (verified)	894
3.166.4 Maple [A] (verified)	895
3.166.5 Fricas [A] (verification not implemented)	895
3.166.6 Sympy [A] (verification not implemented)	895
3.166.7 Maxima [A] (verification not implemented)	896
3.166.8 Giac [A] (verification not implemented)	896
3.166.9 Mupad [B] (verification not implemented)	896

3.166.1 Optimal result

Integrand size = 11, antiderivative size = 21

$$\int \frac{1}{-x^3+x^4} dx = \frac{1}{2x^2} + \frac{1}{x} + \log(1-x) - \log(x)$$

output `1/2/x^2+1/x+ln(1-x)-ln(x)`

3.166.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{1}{-x^3+x^4} dx = \frac{1}{2x^2} + \frac{1}{x} + \log(1-x) - \log(x)$$

input `Integrate[(-x^3 + x^4)^(-1), x]`

output `1/(2*x^2) + x^(-1) + Log[1 - x] - Log[x]`

3.166.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {2026, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x^4 - x^3} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{1}{(x-1)x^3} dx \\ & \quad \downarrow \text{54} \\ & \int \left(-\frac{1}{x^3} - \frac{1}{x^2} - \frac{1}{x} + \frac{1}{x-1} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2x^2} + \frac{1}{x} + \log(1-x) - \log(x) \end{aligned}$$

input `Int[(-x^3 + x^4)^(-1), x]`

output `1/(2*x^2) + x^(-1) + Log[1 - x] - Log[x]`

3.166.3.1 Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

3.166.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

method	result	size
norman	$\frac{x+\frac{1}{2}}{x^2} - \ln(x) + \ln(-1+x)$	17
risch	$\frac{x+\frac{1}{2}}{x^2} - \ln(x) + \ln(-1+x)$	17
default	$\ln(-1+x) + \frac{1}{2x^2} + \frac{1}{x} - \ln(x)$	18
meijerg	$\frac{1}{2x^2} + \frac{1}{x} - \ln(x) - i\pi + \ln(1-x)$	24
parallelrisch	$-\frac{2x^2 \ln(x) - 2 \ln(-1+x)x^2 - 1 - 2x}{2x^2}$	27

input `int(1/(x^4-x^3),x,method=_RETURNVERBOSE)`output `(x+1/2)/x^2-ln(x)+ln(-1+x)`**3.166.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.24

$$\int \frac{1}{-x^3 + x^4} dx = \frac{2x^2 \log(x-1) - 2x^2 \log(x) + 2x + 1}{2x^2}$$

input `integrate(1/(x^4-x^3),x, algorithm="fricas")`output `1/2*(2*x^2*log(x - 1) - 2*x^2*log(x) + 2*x + 1)/x^2`**3.166.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{1}{-x^3 + x^4} dx = -\log(x) + \log(x-1) + \frac{2x+1}{2x^2}$$

input `integrate(1/(x**4-x**3),x)`output `-log(x) + log(x - 1) + (2*x + 1)/(2*x**2)`

3.166.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{1}{-x^3 + x^4} dx = \frac{2x + 1}{2x^2} + \log(x - 1) - \log(x)$$

input `integrate(1/(x^4-x^3),x, algorithm="maxima")`output `1/2*(2*x + 1)/x^2 + log(x - 1) - log(x)`**3.166.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{1}{-x^3 + x^4} dx = \frac{2x + 1}{2x^2} + \log(|x - 1|) - \log(|x|)$$

input `integrate(1/(x^4-x^3),x, algorithm="giac")`output `1/2*(2*x + 1)/x^2 + log(abs(x - 1)) - log(abs(x))`**3.166.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.76

$$\int \frac{1}{-x^3 + x^4} dx = \frac{x + \frac{1}{2}}{x^2} - 2 \operatorname{atanh}(2x - 1)$$

input `int(-1/(x^3 - x^4),x)`output `(x + 1/2)/x^2 - 2*atanh(2*x - 1)`

3.167 $\int \frac{1-x-x^2+x^3+x^4}{-x+x^3} dx$

3.167.1 Optimal result 897
 3.167.2 Mathematica [A] (verified) 897
 3.167.3 Rubi [A] (verified) 898
 3.167.4 Maple [A] (verified) 899
 3.167.5 Fracas [A] (verification not implemented) 899
 3.167.6 Sympy [A] (verification not implemented) 899
 3.167.7 Maxima [A] (verification not implemented) 900
 3.167.8 Giac [A] (verification not implemented) 900
 3.167.9 Mupad [B] (verification not implemented) 900

3.167.1 Optimal result

Integrand size = 26, antiderivative size = 25

$$\int \frac{1-x-x^2+x^3+x^4}{-x+x^3} dx = x + \frac{x^2}{2} - \log(x) + \frac{1}{2} \log(1-x^2)$$

output `x+1/2*x^2-ln(x)+1/2*ln(-x^2+1)`

3.167.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1-x-x^2+x^3+x^4}{-x+x^3} dx = x + \frac{x^2}{2} - \log(x) + \frac{1}{2} \log(1-x^2)$$

input `Integrate[(1 - x - x^2 + x^3 + x^4)/(-x + x^3),x]`

output `x + x^2/2 - Log[x] + Log[1 - x^2]/2`

3.167.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {2026, 2333, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 + x^3 - x^2 - x + 1}{x^3 - x} dx$$

↓ 2026

$$\int \frac{x^4 + x^3 - x^2 - x + 1}{x(x^2 - 1)} dx$$

↓ 2333

$$\int \left(\frac{x}{x^2 - 1} + x - \frac{1}{x} + 1 \right) dx$$

↓ 2009

$$\frac{x^2}{2} + \frac{1}{2} \log(1 - x^2) + x - \log(x)$$

input `Int[(1 - x - x^2 + x^3 + x^4)/(-x + x^3),x]`

output `x + x^2/2 - Log[x] + Log[1 - x^2]/2`

3.167.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2333 `Int[(Pq_)*((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

3.167. $\int \frac{1-x-x^2+x^3+x^4}{-x+x^3} dx$

3.167.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

method	result	size
risch	$x + \frac{x^2}{2} - \ln(x) + \frac{\ln(x^2-1)}{2}$	20
default	$x + \frac{x^2}{2} + \frac{\ln(-1+x)}{2} - \ln(x) + \frac{\ln(1+x)}{2}$	24
norman	$x + \frac{x^2}{2} + \frac{\ln(-1+x)}{2} - \ln(x) + \frac{\ln(1+x)}{2}$	24
parallelrisc	$x + \frac{x^2}{2} + \frac{\ln(-1+x)}{2} - \ln(x) + \frac{\ln(1+x)}{2}$	24
meijerg	$-\ln(x) - \frac{i\pi}{2} + \frac{\ln(-x^2+1)}{2} + \frac{x^2}{2} - \frac{i(2ix-2i \operatorname{arctanh}(x))}{2} + \operatorname{arctanh}(x)$	40

input `int((x^4+x^3-x^2-x+1)/(x^3-x),x,method=_RETURNVERBOSE)`output `x+1/2*x^2-ln(x)+1/2*ln(x^2-1)`**3.167.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{1-x-x^2+x^3+x^4}{-x+x^3} dx = \frac{1}{2}x^2 + x + \frac{1}{2} \log(x^2-1) - \log(x)$$

input `integrate((x^4+x^3-x^2-x+1)/(x^3-x),x, algorithm="fracas")`output `1/2*x^2 + x + 1/2*log(x^2 - 1) - log(x)`**3.167.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.68

$$\int \frac{1-x-x^2+x^3+x^4}{-x+x^3} dx = \frac{x^2}{2} + x - \log(x) + \frac{\log(x^2-1)}{2}$$

input `integrate((x**4+x**3-x**2-x+1)/(x**3-x),x)`output `x**2/2 + x - log(x) + log(x**2 - 1)/2`

3.167. $\int \frac{1-x-x^2+x^3+x^4}{-x+x^3} dx$

3.167.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{1 - x - x^2 + x^3 + x^4}{-x + x^3} dx = \frac{1}{2}x^2 + x + \frac{1}{2} \log(x + 1) + \frac{1}{2} \log(x - 1) - \log(x)$$

input `integrate((x^4+x^3-x^2-x+1)/(x^3-x),x, algorithm="maxima")`output `1/2*x^2 + x + 1/2*log(x + 1) + 1/2*log(x - 1) - log(x)`**3.167.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

$$\int \frac{1 - x - x^2 + x^3 + x^4}{-x + x^3} dx = \frac{1}{2}x^2 + x + \frac{1}{2} \log(|x + 1|) + \frac{1}{2} \log(|x - 1|) - \log(|x|)$$

input `integrate((x^4+x^3-x^2-x+1)/(x^3-x),x, algorithm="giac")`output `1/2*x^2 + x + 1/2*log(abs(x + 1)) + 1/2*log(abs(x - 1)) - log(abs(x))`**3.167.9 Mupad [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{1 - x - x^2 + x^3 + x^4}{-x + x^3} dx = x + \frac{\ln(x^2 - 1)}{2} - \ln(x) + \frac{x^2}{2}$$

input `int(-(x^3 - x^2 - x + x^4 + 1)/(x - x^3),x)`output `x + log(x^2 - 1)/2 - log(x) + x^2/2`

$$3.168 \quad \int \frac{-2+x^2}{x(2+x^2)} dx$$

3.168.1 Optimal result	901
3.168.2 Mathematica [A] (verified)	901
3.168.3 Rubi [A] (verified)	902
3.168.4 Maple [A] (verified)	903
3.168.5 Fracas [A] (verification not implemented)	904
3.168.6 Sympy [A] (verification not implemented)	904
3.168.7 Maxima [A] (verification not implemented)	904
3.168.8 Giac [A] (verification not implemented)	905
3.168.9 Mupad [B] (verification not implemented)	905

3.168.1 Optimal result

Integrand size = 16, antiderivative size = 11

$$\int \frac{-2+x^2}{x(2+x^2)} dx = -\log(x) + \log(2+x^2)$$

output `-ln(x)+ln(x^2+2)`

3.168.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{-2+x^2}{x(2+x^2)} dx = -\log(x) + \log(2+x^2)$$

input `Integrate[(-2 + x^2)/(x*(2 + x^2)), x]`

output `-Log[x] + Log[2 + x^2]`

3.168.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.73, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {354, 25, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2 - 2}{x(x^2 + 2)} dx \\
 & \quad \downarrow \text{354} \\
 & \frac{1}{2} \int -\frac{2 - x^2}{x^2(x^2 + 2)} dx^2 \\
 & \quad \downarrow \text{25} \\
 & -\frac{1}{2} \int \frac{2 - x^2}{x^2(x^2 + 2)} dx^2 \\
 & \quad \downarrow \text{86} \\
 & -\frac{1}{2} \int \left(\frac{1}{x^2} - \frac{2}{x^2 + 2} \right) dx^2 \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} (2 \log(x^2 + 2) - \log(x^2))
 \end{aligned}$$

input `Int[(-2 + x^2)/(x*(2 + x^2)),x]`

output `(-Log[x^2] + 2*Log[2 + x^2])/2`

3.168.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 354 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.168.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
default	$-\ln(x) + \ln(x^2 + 2)$	12
norman	$-\ln(x) + \ln(x^2 + 2)$	12
risch	$-\ln(x) + \ln(x^2 + 2)$	12
parallelrisc	$-\ln(x) + \ln(x^2 + 2)$	12
meijerg	$-\ln(x) + \frac{\ln(2)}{2} + \ln\left(1 + \frac{x^2}{2}\right)$	18

input `int((x^2-2)/x/(x^2+2),x,method=_RETURNVERBOSE)`

output `-ln(x)+ln(x^2+2)`

3.168.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{-2 + x^2}{x(2 + x^2)} dx = \log(x^2 + 2) - \log(x)$$

input `integrate((x^2-2)/x/(x^2+2),x, algorithm="fricas")`

output `log(x^2 + 2) - log(x)`

3.168.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \frac{-2 + x^2}{x(2 + x^2)} dx = -\log(x) + \log(x^2 + 2)$$

input `integrate((x**2-2)/x/(x**2+2),x)`

output `-log(x) + log(x**2 + 2)`

3.168.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.18

$$\int \frac{-2 + x^2}{x(2 + x^2)} dx = \log(x^2 + 2) - \frac{1}{2} \log(x^2)$$

input `integrate((x^2-2)/x/(x^2+2),x, algorithm="maxima")`

output `log(x^2 + 2) - 1/2*log(x^2)`

3.168.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.18

$$\int \frac{-2 + x^2}{x(2 + x^2)} dx = \log(x^2 + 2) - \frac{1}{2} \log(x^2)$$

input `integrate((x^2-2)/x/(x^2+2),x, algorithm="giac")`

output `log(x^2 + 2) - 1/2*log(x^2)`

3.168.9 Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{-2 + x^2}{x(2 + x^2)} dx = \ln(x^2 + 2) - \ln(x)$$

input `int((x^2 - 2)/(x*(x^2 + 2)),x)`

output `log(x^2 + 2) - log(x)`

$$3.169 \quad \int \frac{2-4x^2+x^3}{(1+x^2)(2+x^2)} dx$$

3.169.1 Optimal result	906
3.169.2 Mathematica [A] (verified)	906
3.169.3 Rubi [A] (verified)	907
3.169.4 Maple [A] (verified)	908
3.169.5 Fricas [A] (verification not implemented)	908
3.169.6 Sympy [A] (verification not implemented)	908
3.169.7 Maxima [A] (verification not implemented)	909
3.169.8 Giac [A] (verification not implemented)	909
3.169.9 Mupad [B] (verification not implemented)	909

3.169.1 Optimal result

Integrand size = 25, antiderivative size = 36

$$\int \frac{2-4x^2+x^3}{(1+x^2)(2+x^2)} dx = 6 \arctan(x) - 5\sqrt{2} \arctan\left(\frac{x}{\sqrt{2}}\right) - \frac{1}{2} \log(1+x^2) + \log(2+x^2)$$

output `6*arctan(x)-1/2*ln(x^2+1)+ln(x^2+2)-5*arctan(1/2*x*2^(1/2))*2^(1/2)`

3.169.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00

$$\int \frac{2-4x^2+x^3}{(1+x^2)(2+x^2)} dx = 6 \arctan(x) - 5\sqrt{2} \arctan\left(\frac{x}{\sqrt{2}}\right) - \frac{1}{2} \log(1+x^2) + \log(2+x^2)$$

input `Integrate[(2 - 4*x^2 + x^3)/((1 + x^2)*(2 + x^2)),x]`

output `6*ArcTan[x] - 5*sqrt[2]*ArcTan[x/sqrt[2]] - Log[1 + x^2]/2 + Log[2 + x^2]`

3.169.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 - 4x^2 + 2}{(x^2 + 1)(x^2 + 2)} dx$$

↓ 7276

$$\int \left(\frac{6 - x}{x^2 + 1} + \frac{2(x - 5)}{x^2 + 2} \right) dx$$

↓ 2009

$$6 \arctan(x) - 5\sqrt{2} \arctan\left(\frac{x}{\sqrt{2}}\right) - \frac{1}{2} \log(x^2 + 1) + \log(x^2 + 2)$$

input `Int[(2 - 4*x^2 + x^3)/((1 + x^2)*(2 + x^2)),x]`

output `6*ArcTan[x] - 5*Sqrt[2]*ArcTan[x/Sqrt[2]] - Log[1 + x^2]/2 + Log[2 + x^2]`

3.169.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
x
p
a
n
x
}], Int[v, x] /; SumQ[v] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]`

3.169.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.89

method	result	size
default	$6 \arctan(x) - \frac{\ln(x^2+1)}{2} + \ln(x^2+2) - 5 \arctan\left(\frac{x\sqrt{2}}{2}\right) \sqrt{2}$	32
risch	$6 \arctan(x) - \frac{\ln(x^2+1)}{2} + \ln(x^2+2) - 5 \arctan\left(\frac{x\sqrt{2}}{2}\right) \sqrt{2}$	32

input `int((x^3-4*x^2+2)/(x^2+1)/(x^2+2),x,method=_RETURNVERBOSE)`output `6*arctan(x)-1/2*ln(x^2+1)+ln(x^2+2)-5*arctan(1/2*x*2^(1/2))*2^(1/2)`**3.169.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

$$\int \frac{2 - 4x^2 + x^3}{(1 + x^2)(2 + x^2)} dx = -5\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}x\right) + 6 \arctan(x) + \log(x^2 + 2) - \frac{1}{2} \log(x^2 + 1)$$

input `integrate((x^3-4*x^2+2)/(x^2+1)/(x^2+2),x, algorithm="fricas")`output `-5*sqrt(2)*arctan(1/2*sqrt(2)*x) + 6*arctan(x) + log(x^2 + 2) - 1/2*log(x^2 + 1)`**3.169.6 Sympy [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00

$$\int \frac{2 - 4x^2 + x^3}{(1 + x^2)(2 + x^2)} dx = -\frac{\log(x^2 + 1)}{2} + \log(x^2 + 2) + 6 \operatorname{atan}(x) - 5\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}x}{2}\right)$$

input `integrate((x**3-4*x**2+2)/(x**2+1)/(x**2+2),x)`output `-log(x**2 + 1)/2 + log(x**2 + 2) + 6*atan(x) - 5*sqrt(2)*atan(sqrt(2)*x/2)`

3.169. $\int \frac{2-4x^2+x^3}{(1+x^2)(2+x^2)} dx$

3.169.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

$$\int \frac{2 - 4x^2 + x^3}{(1 + x^2)(2 + x^2)} dx = -5\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}x\right) + 6 \arctan(x) + \log(x^2 + 2) - \frac{1}{2} \log(x^2 + 1)$$

input `integrate((x^3-4*x^2+2)/(x^2+1)/(x^2+2),x, algorithm="maxima")`output `-5*sqrt(2)*arctan(1/2*sqrt(2)*x) + 6*arctan(x) + log(x^2 + 2) - 1/2*log(x^2 + 1)`**3.169.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

$$\int \frac{2 - 4x^2 + x^3}{(1 + x^2)(2 + x^2)} dx = -5\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}x\right) + 6 \arctan(x) + \log(x^2 + 2) - \frac{1}{2} \log(x^2 + 1)$$

input `integrate((x^3-4*x^2+2)/(x^2+1)/(x^2+2),x, algorithm="giac")`output `-5*sqrt(2)*arctan(1/2*sqrt(2)*x) + 6*arctan(x) + log(x^2 + 2) - 1/2*log(x^2 + 1)`**3.169.9 Mupad [B] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.56

$$\begin{aligned} \int \frac{2 - 4x^2 + x^3}{(1 + x^2)(2 + x^2)} dx = & \ln(x - i) \left(-\frac{1}{2} - 3i\right) + \ln(x + i) \left(-\frac{1}{2} + 3i\right) \\ & + \ln\left(x - \sqrt{2}li\right) \left(1 + \frac{\sqrt{2}5i}{2}\right) - \ln\left(x + \sqrt{2}li\right) \left(-1 + \frac{\sqrt{2}5i}{2}\right) \end{aligned}$$

input `int((x^3 - 4*x^2 + 2)/((x^2 + 1)*(x^2 + 2)),x)`output `log(x - 2^(1/2)*1i)*((2^(1/2)*5i)/2 + 1) - log(x + 1i)*(1/2 - 3i) - log(x - 1i)*(1/2 + 3i) - log(x + 2^(1/2)*1i)*((2^(1/2)*5i)/2 - 1)`

3.169. $\int \frac{2-4x^2+x^3}{(1+x^2)(2+x^2)} dx$

3.170 $\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx$

3.170.1 Optimal result 910
 3.170.2 Mathematica [A] (verified) 910
 3.170.3 Rubi [A] (verified) 911
 3.170.4 Maple [A] (verified) 912
 3.170.5 Fricas [A] (verification not implemented) 912
 3.170.6 Sympy [A] (verification not implemented) 912
 3.170.7 Maxima [A] (verification not implemented) 913
 3.170.8 Giac [A] (verification not implemented) 913
 3.170.9 Mupad [B] (verification not implemented) 913

3.170.1 Optimal result

Integrand size = 23, antiderivative size = 29

$$\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx = -\frac{13x}{24(4+x^2)} + \frac{25}{144} \arctan\left(\frac{x}{2}\right) + \frac{\arctan(x)}{9}$$

output `-13/24*x/(x^2+4)+25/144*arctan(1/2*x)+1/9*arctan(x)`

3.170.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx = -\frac{13x}{24(4+x^2)} + \frac{25}{144} \arctan\left(\frac{x}{2}\right) + \frac{\arctan(x)}{9}$$

input `Integrate[(1 + x^2 + x^4)/((1 + x^2)*(4 + x^2)^2),x]`

output `(-13*x)/(24*(4 + x^2)) + (25*ArcTan[x/2])/144 + ArcTan[x]/9`

3.170.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 + x^2 + 1}{(x^2 + 1)(x^2 + 4)^2} dx$$

↓ 7276

$$\int \left(\frac{8}{9(x^2 + 4)} - \frac{13}{3(x^2 + 4)^2} + \frac{1}{9(x^2 + 1)} \right) dx$$

↓ 2009

$$\frac{25}{144} \arctan\left(\frac{x}{2}\right) + \frac{\arctan(x)}{9} - \frac{13x}{24(x^2 + 4)}$$

input `Int[(1 + x^2 + x^4)/((1 + x^2)*(4 + x^2)^2),x]`

output `(-13*x)/(24*(4 + x^2)) + (25*ArcTan[x/2])/144 + ArcTan[x]/9`

3.170.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
x
p
a
n
x
b
n
x
v
x
v
a
b
x
n
0`

3.170.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

method	result
default	$-\frac{13x}{24(x^2+4)} + \frac{25 \arctan(\frac{x}{2})}{144} + \frac{\arctan(x)}{9}$
risch	$-\frac{13x}{24(x^2+4)} + \frac{25 \arctan(\frac{x}{2})}{144} + \frac{\arctan(x)}{9}$
parallelrisch	$-\frac{25i \ln(x-2i)x^2 + 16i \ln(x-i)x^2 - 16i \ln(x+i)x^2 - 25i \ln(x+2i)x^2 + 100i \ln(x-2i) + 64i \ln(x-i) - 64i \ln(x+i) - 100i \ln(x+2i) + 100}{288(x^2+4)}$

input `int((x^4+x^2+1)/(x^2+1)/(x^2+4)^2,x,method=_RETURNVERBOSE)`output `-13/24*x/(x^2+4)+25/144*arctan(1/2*x)+1/9*arctan(x)`**3.170.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.14

$$\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx = \frac{25(x^2+4) \arctan\left(\frac{1}{2}x\right) + 16(x^2+4) \arctan(x) - 78x}{144(x^2+4)}$$

input `integrate((x^4+x^2+1)/(x^2+1)/(x^2+4)^2,x, algorithm="fracas")`output `1/144*(25*(x^2 + 4)*arctan(1/2*x) + 16*(x^2 + 4)*arctan(x) - 78*x)/(x^2 + 4)`**3.170.6 Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

$$\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx = -\frac{13x}{24x^2+96} + \frac{25 \operatorname{atan}\left(\frac{x}{2}\right)}{144} + \frac{\operatorname{atan}(x)}{9}$$

input `integrate((x**4+x**2+1)/(x**2+1)/(x**2+4)**2,x)`output `-13*x/(24*x**2 + 96) + 25*atan(x/2)/144 + atan(x)/9`

3.170. $\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx$

3.170.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx = -\frac{13x}{24(x^2+4)} + \frac{25}{144} \arctan\left(\frac{1}{2}x\right) + \frac{1}{9} \arctan(x)$$

input `integrate((x^4+x^2+1)/(x^2+1)/(x^2+4)^2,x, algorithm="maxima")`output `-13/24*x/(x^2 + 4) + 25/144*arctan(1/2*x) + 1/9*arctan(x)`**3.170.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx = -\frac{13x}{24(x^2+4)} + \frac{25}{144} \arctan\left(\frac{1}{2}x\right) + \frac{1}{9} \arctan(x)$$

input `integrate((x^4+x^2+1)/(x^2+1)/(x^2+4)^2,x, algorithm="giac")`output `-13/24*x/(x^2 + 4) + 25/144*arctan(1/2*x) + 1/9*arctan(x)`**3.170.9 Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

$$\int \frac{1+x^2+x^4}{(1+x^2)(4+x^2)^2} dx = \frac{25 \operatorname{atan}\left(\frac{x}{2}\right)}{144} + \frac{\operatorname{atan}(x)}{9} - \frac{13x}{24(x^2+4)}$$

input `int((x^2 + x^4 + 1)/((x^2 + 1)*(x^2 + 4)^2),x)`output `(25*atan(x/2))/144 + atan(x)/9 - (13*x)/(24*(x^2 + 4))`

3.171 $\int \frac{1+16x}{(5+x)^2(-3+2x)(1+x+x^2)} dx$

3.171.1 Optimal result 914
 3.171.2 Mathematica [A] (verified) 914
 3.171.3 Rubi [A] (verified) 915
 3.171.4 Maple [A] (verified) 916
 3.171.5 Fricas [A] (verification not implemented) 916
 3.171.6 Sympy [A] (verification not implemented) 917
 3.171.7 Maxima [A] (verification not implemented) 917
 3.171.8 Giac [A] (verification not implemented) 918
 3.171.9 Mupad [B] (verification not implemented) 918

3.171.1 Optimal result

Integrand size = 26, antiderivative size = 60

$$\int \frac{1 + 16x}{(5 + x)^2(-3 + 2x)(1 + x + x^2)} dx = -\frac{79}{273(5 + x)} + \frac{451 \arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{2793\sqrt{3}} + \frac{200 \log(3 - 2x)}{3211} + \frac{2731 \log(5 + x)}{24843} - \frac{481 \log(1 + x + x^2)}{5586}$$

output `-79/273/(5+x)+200/3211*ln(3-2*x)+2731/24843*ln(5+x)-481/5586*ln(x^2+x+1)+451/8379*arctan(1/3*(1+2*x)*3^(1/2))*3^(1/2)`

3.171.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.90

$$\int \frac{1 + 16x}{(5 + x)^2(-3 + 2x)(1 + x + x^2)} dx = \frac{-\frac{819546}{5+x} + 152438\sqrt{3} \arctan\left(\frac{1+2x}{\sqrt{3}}\right) + 176400 \log(3 - 2x) + 311334 \log(5 + x) - 243867 \log(1 + x + x^2)}{2832102}$$

input `Integrate[(1 + 16*x)/((5 + x)^2*(-3 + 2*x)*(1 + x + x^2)),x]`

output `(-819546/(5 + x) + 152438*sqrt[3]*ArcTan[(1 + 2*x)/sqrt[3]] + 176400*Log[3 - 2*x] + 311334*Log[5 + x] - 243867*Log[1 + x + x^2])/2832102`

3.171. $\int \frac{1+16x}{(5+x)^2(-3+2x)(1+x+x^2)} dx$

3.171.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {2153, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{16x + 1}{(x + 5)^2(2x - 3)(x^2 + x + 1)} dx$$

↓ 2153

$$\int \left(\frac{-481x - 15}{2793(x^2 + x + 1)} + \frac{2731}{24843(x + 5)} + \frac{400}{3211(2x - 3)} + \frac{79}{273(x + 5)^2} \right) dx$$

↓ 2009

$$\frac{451 \arctan\left(\frac{2x+1}{\sqrt{3}}\right)}{2793\sqrt{3}} - \frac{481 \log(x^2 + x + 1)}{5586} - \frac{79}{273(x + 5)} + \frac{200 \log(3 - 2x)}{3211} + \frac{2731 \log(x + 5)}{24843}$$

input `Int[(1 + 16*x)/((5 + x)^2*(-3 + 2*x)*(1 + x + x^2)),x]`

output `-79/(273*(5 + x)) + (451*ArcTan[(1 + 2*x)/Sqrt[3]])/(2793*Sqrt[3]) + (200*Log[3 - 2*x])/3211 + (2731*Log[5 + x])/24843 - (481*Log[1 + x + x^2])/5586`

3.171.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2153 `Int[(Px_)*((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && PolyQ[Px, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0])) && !(IGtQ[m, 0] && IGtQ[n, 0])`

3.171.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.80

method	result
default	$-\frac{79}{273(5+x)} + \frac{2731 \ln(5+x)}{24843} - \frac{481 \ln(x^2+x+1)}{5586} + \frac{451 \arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{8379} + \frac{200 \ln(2x-3)}{3211}$
risch	$-\frac{79}{273(5+x)} + \frac{2731 \ln(5+x)}{24843} - \frac{481 \ln(203401x^2+203401x+203401)}{5586} + \frac{451\sqrt{3} \arctan\left(\frac{2(451x+\frac{451}{2})\sqrt{3}}{1353}\right)}{8379} + \frac{200 \ln(2x-3)}{3211}$

input `int((1+16*x)/(5+x)^2/(2*x-3)/(x^2+x+1),x,method=_RETURNVERBOSE)`output
$$-79/273/(5+x)+2731/24843*\ln(5+x)-481/5586*\ln(x^2+x+1)+451/8379*\arctan(1/3*(1+2*x)*3^{(1/2)})*3^{(1/2)}+200/3211*\ln(2*x-3)$$
3.171.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00

$$\int \frac{1+16x}{(5+x)^2(-3+2x)(1+x+x^2)} dx$$

$$= \frac{152438 \sqrt{3}(x+5) \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) - 243867(x+5) \log(x^2+x+1) + 176400(x+5) \log(2x-3)}{2832102(x+5)}$$

input `integrate((1+16*x)/(5+x)^2/(-3+2*x)/(x^2+x+1),x, algorithm="fricas")`output
$$1/2832102*(152438*\sqrt{3}*(x+5)*\arctan(1/3*\sqrt{3}*(2*x+1)) - 243867*(x+5)*\log(x^2+x+1) + 176400*(x+5)*\log(2*x-3) + 311334*(x+5)*\log(x+5) - 819546)/(x+5)$$

3.171.6 Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.05

$$\int \frac{1 + 16x}{(5 + x)^2(-3 + 2x)(1 + x + x^2)} dx = \frac{200 \log(x - \frac{3}{2})}{3211} + \frac{2731 \log(x + 5)}{24843} - \frac{481 \log(x^2 + x + 1)}{5586} + \frac{451\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x}{3} + \frac{\sqrt{3}}{3}\right)}{8379} - \frac{79}{273x + 1365}$$

input `integrate((1+16*x)/(5+x)**2/(-3+2*x)/(x**2+x+1),x)`output `200*log(x - 3/2)/3211 + 2731*log(x + 5)/24843 - 481*log(x**2 + x + 1)/5586 + 451*sqrt(3)*atan(2*sqrt(3)*x/3 + sqrt(3)/3)/8379 - 79/(273*x + 1365)`**3.171.7 Maxima [A] (verification not implemented)**

Time = 0.37 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.78

$$\int \frac{1 + 16x}{(5 + x)^2(-3 + 2x)(1 + x + x^2)} dx = \frac{451}{8379} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x + 1)\right) - \frac{79}{273(x + 5)} - \frac{481}{5586} \log(x^2 + x + 1) + \frac{200}{3211} \log(2x - 3) + \frac{2731}{24843} \log(x + 5)$$

input `integrate((1+16*x)/(5+x)^2/(-3+2*x)/(x^2+x+1),x, algorithm="maxima")`output `451/8379*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) - 79/273/(x + 5) - 481/5586*log(x^2 + x + 1) + 200/3211*log(2*x - 3) + 2731/24843*log(x + 5)`

3.171.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00

$$\int \frac{1+16x}{(5+x)^2(-3+2x)(1+x+x^2)} dx = \frac{451}{8379} \sqrt{3} \arctan \left(-\sqrt{3} \left(\frac{14}{x+5} - 3 \right) \right) - \frac{79}{273(x+5)} - \frac{481}{5586} \log \left(-\frac{9}{x+5} + \frac{21}{(x+5)^2} + 1 \right) + \frac{200}{3211} \log \left(\left| -\frac{13}{x+5} + 2 \right| \right)$$

input `integrate((1+16*x)/(5+x)^2/(-3+2*x)/(x^2+x+1),x, algorithm="giac")`output `451/8379*sqrt(3)*arctan(-sqrt(3)*(14/(x + 5) - 3)) - 79/273/(x + 5) - 481/5586*log(-9/(x + 5) + 21/(x + 5)^2 + 1) + 200/3211*log(abs(-13/(x + 5) + 2))`**3.171.9 Mupad [B] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.02

$$\int \frac{1+16x}{(5+x)^2(-3+2x)(1+x+x^2)} dx = \frac{200 \ln \left(x - \frac{3}{2} \right)}{3211} + \frac{2731 \ln(x+5)}{24843} - \frac{79}{273(x+5)} - \ln \left(x + \frac{1}{2} - \frac{\sqrt{3} \text{li}}{2} \right) \left(\frac{481}{5586} + \frac{\sqrt{3} 451 \text{i}}{16758} \right) + \ln \left(x + \frac{1}{2} + \frac{\sqrt{3} \text{li}}{2} \right) \left(-\frac{481}{5586} + \frac{\sqrt{3} 451 \text{i}}{16758} \right)$$

input `int((16*x + 1)/((2*x - 3)*(x + 5)^2*(x + x^2 + 1)),x)`output `(200*log(x - 3/2))/3211 + (2731*log(x + 5))/24843 - 79/(273*(x + 5)) - log(x - (3^(1/2)*1i)/2 + 1/2)*((3^(1/2)*451i)/16758 + 481/5586) + log(x + (3^(1/2)*1i)/2 + 1/2)*((3^(1/2)*451i)/16758 - 481/5586)`

3.172 $\int \frac{x^4}{(9+x^2)^3} dx$

3.172.1 Optimal result	919
3.172.2 Mathematica [A] (verified)	919
3.172.3 Rubi [A] (verified)	920
3.172.4 Maple [A] (verified)	921
3.172.5 Fricas [A] (verification not implemented)	921
3.172.6 Sympy [A] (verification not implemented)	921
3.172.7 Maxima [A] (verification not implemented)	922
3.172.8 Giac [A] (verification not implemented)	922
3.172.9 Mupad [B] (verification not implemented)	922

3.172.1 Optimal result

Integrand size = 11, antiderivative size = 37

$$\int \frac{x^4}{(9+x^2)^3} dx = -\frac{x^3}{4(9+x^2)^2} - \frac{3x}{8(9+x^2)} + \frac{1}{8} \arctan\left(\frac{x}{3}\right)$$

output `-1/4*x^3/(x^2+9)^2-3/8*x/(x^2+9)+1/8*arctan(1/3*x)`

3.172.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.76

$$\int \frac{x^4}{(9+x^2)^3} dx = \frac{1}{8} \left(-\frac{x(27+5x^2)}{(9+x^2)^2} + \arctan\left(\frac{x}{3}\right) \right)$$

input `Integrate[x^4/(9 + x^2)^3,x]`

output `(-((x*(27 + 5*x^2))/(9 + x^2)^2) + ArcTan[x/3])/8`

3.172.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.14, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {252, 252, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^4}{(x^2 + 9)^3} dx \\ & \quad \downarrow \text{252} \\ & \frac{3}{4} \int \frac{x^2}{(x^2 + 9)^2} dx - \frac{x^3}{4(x^2 + 9)^2} \\ & \quad \downarrow \text{252} \\ & \frac{3}{4} \left(\frac{1}{2} \int \frac{1}{x^2 + 9} dx - \frac{x}{2(x^2 + 9)} \right) - \frac{x^3}{4(x^2 + 9)^2} \\ & \quad \downarrow \text{216} \\ & \frac{3}{4} \left(\frac{1}{6} \arctan\left(\frac{x}{3}\right) - \frac{x}{2(x^2 + 9)} \right) - \frac{x^3}{4(x^2 + 9)^2} \end{aligned}$$

input `Int[x^4/(9 + x^2)^3,x]`

output `-1/4*x^3/(9 + x^2)^2 + (3*(-1/2*x/(9 + x^2) + ArcTan[x/3]/6))/4`

3.172.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 252 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*((m - 1)/(2*b*(p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !IlTQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

3.172.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.68

method	result
default	$\frac{-\frac{5}{8}x^3 - \frac{27}{8}x}{(x^2+9)^2} + \frac{\arctan(\frac{x}{3})}{8}$
risch	$\frac{-\frac{5}{8}x^3 - \frac{27}{8}x}{(x^2+9)^2} + \frac{\arctan(\frac{x}{3})}{8}$
meijerg	$-\frac{x\left(\frac{25x^2}{9}+15\right)}{360\left(\frac{x^2}{9}+1\right)^2} + \frac{\arctan(\frac{x}{3})}{8}$
parallelrisch	$-\frac{81i \ln(x-3i)x^4 - 81i \ln(x+3i)x^4 + 1458i \ln(x-3i)x^2 - 1458i \ln(x+3i)x^2 + 810x^3 + 6561i \ln(x-3i) - 6561i \ln(x+3i) + 4374x}{1296(x^2+9)^2}$

input `int(x^4/(x^2+9)^3,x,method=_RETURNVERBOSE)`output `(-5/8*x^3-27/8*x)/(x^2+9)^2+1/8*arctan(1/3*x)`**3.172.5 Fracas [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.05

$$\int \frac{x^4}{(9+x^2)^3} dx = -\frac{5x^3 - (x^4 + 18x^2 + 81) \arctan\left(\frac{1}{3}x\right) + 27x}{8(x^4 + 18x^2 + 81)}$$

input `integrate(x^4/(x^2+9)^3,x, algorithm="fricas")`output `-1/8*(5*x^3 - (x^4 + 18*x^2 + 81)*arctan(1/3*x) + 27*x)/(x^4 + 18*x^2 + 81)`**3.172.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.73

$$\int \frac{x^4}{(9+x^2)^3} dx = \frac{-5x^3 - 27x}{8x^4 + 144x^2 + 648} + \frac{\operatorname{atan}\left(\frac{x}{3}\right)}{8}$$

input `integrate(x**4/(x**2+9)**3,x)`

output `(-5*x**3 - 27*x)/(8*x**4 + 144*x**2 + 648) + atan(x/3)/8`

3.172.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.81

$$\int \frac{x^4}{(9+x^2)^3} dx = -\frac{5x^3+27x}{8(x^4+18x^2+81)} + \frac{1}{8} \arctan\left(\frac{1}{3}x\right)$$

input `integrate(x^4/(x^2+9)^3,x, algorithm="maxima")`

output `-1/8*(5*x^3 + 27*x)/(x^4 + 18*x^2 + 81) + 1/8*arctan(1/3*x)`

3.172.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.68

$$\int \frac{x^4}{(9+x^2)^3} dx = -\frac{5x^3+27x}{8(x^2+9)^2} + \frac{1}{8} \arctan\left(\frac{1}{3}x\right)$$

input `integrate(x^4/(x^2+9)^3,x, algorithm="giac")`

output `-1/8*(5*x^3 + 27*x)/(x^2 + 9)^2 + 1/8*arctan(1/3*x)`

3.172.9 Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.81

$$\int \frac{x^4}{(9+x^2)^3} dx = \frac{\operatorname{atan}\left(\frac{x}{3}\right)}{8} - \frac{\frac{5x^3}{8} + \frac{27x}{8}}{x^4 + 18x^2 + 81}$$

input `int(x^4/(x^2 + 9)^3,x)`

output `atan(x/3)/8 - ((27*x)/8 + (5*x^3)/8)/(18*x^2 + x^4 + 81)`

3.173 $\int \frac{19x}{(-1+x)^3(3+5x+4x^2)^2} dx$

3.173.1 Optimal result	923
3.173.2 Mathematica [A] (verified)	923
3.173.3 Rubi [A] (verified)	924
3.173.4 Maple [A] (verified)	926
3.173.5 Fricas [A] (verification not implemented)	926
3.173.6 Sympy [A] (verification not implemented)	927
3.173.7 Maxima [A] (verification not implemented)	927
3.173.8 Giac [A] (verification not implemented)	928
3.173.9 Mupad [B] (verification not implemented)	928

3.173.1 Optimal result

Integrand size = 20, antiderivative size = 97

$$\int \frac{19x}{(-1+x)^3(3+5x+4x^2)^2} dx = -\frac{399}{736(1-x)^2} - \frac{1843}{4416(1-x)} + \frac{19(39+44x)}{276(1-x)^2(3+5x+4x^2)} + \frac{114437 \arctan\left(\frac{5+8x}{\sqrt{23}}\right)}{52992\sqrt{23}} + \frac{209 \log(1-x)}{2304} - \frac{209 \log(3+5x+4x^2)}{4608}$$

```
output -399/736/(1-x)^2-1843/4416/(1-x)+19/276*(39+44*x)/(1-x)^2/(4*x^2+5*x+3)+20
9/2304*ln(1-x)-209/4608*ln(4*x^2+5*x+3)+114437/1218816*arctan(1/23*(5+8*x)
*23^(1/2))*23^(1/2)
```

3.173.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.80

$$\int \frac{19x}{(-1+x)^3(3+5x+4x^2)^2} dx = \frac{19\left(-\frac{25392}{(-1+x)^2} + \frac{59248}{-1+x} + \frac{184(975+2204x)}{3+5x+4x^2} + 36138\sqrt{23} \arctan\left(\frac{5+8x}{\sqrt{23}}\right) + 34914 \log(1-x) - 17457 \log(3+5x+4x^2)\right)}{7312896}$$

input `Integrate[(19*x)/((-1 + x)^3*(3 + 5*x + 4*x^2)^2),x]`

output $(19*(-25392/(-1 + x)^2 + 59248/(-1 + x) + (184*(975 + 2204*x))/(3 + 5*x + 4*x^2) + 36138*\text{Sqrt}[23]*\text{ArcTan}[(5 + 8*x)/\text{Sqrt}[23]] + 34914*\text{Log}[1 - x] - 17457*\text{Log}[3 + 5*x + 4*x^2]))/7312896$

3.173.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.07, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {27, 25, 1235, 27, 1200, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{19x}{(x-1)^3(4x^2+5x+3)^2} dx \\
 & \quad \downarrow 27 \\
 & 19 \int -\frac{x}{(1-x)^3(4x^2+5x+3)^2} dx \\
 & \quad \downarrow 25 \\
 & -19 \int \frac{x}{(1-x)^3(4x^2+5x+3)^2} dx \\
 & \quad \downarrow 1235 \\
 & -19 \left(\frac{1}{276} \int \frac{3(44x+19)}{(1-x)^3(4x^2+5x+3)} dx - \frac{44x+39}{276(1-x)^2(4x^2+5x+3)} \right) \\
 & \quad \downarrow 27 \\
 & -19 \left(\frac{1}{92} \int \frac{44x+19}{(1-x)^3(4x^2+5x+3)} dx - \frac{44x+39}{276(1-x)^2(4x^2+5x+3)} \right) \\
 & \quad \downarrow 1200 \\
 & -19 \left(\frac{1}{92} \int \left(\frac{1012x-2379}{576(4x^2+5x+3)} - \frac{253}{576(x-1)} + \frac{97}{48(x-1)^2} - \frac{21}{4(x-1)^3} \right) dx - \frac{44x+39}{276(1-x)^2(4x^2+5x+3)} \right) \\
 & \quad \downarrow 2009
 \end{aligned}$$

$$-19 \left(\frac{1}{92} \left(-\frac{6023 \arctan\left(\frac{8x+5}{\sqrt{23}}\right)}{576\sqrt{23}} + \frac{253 \log(4x^2 + 5x + 3)}{1152} + \frac{97}{48(1-x)} + \frac{21}{8(1-x)^2} - \frac{253}{576} \log(1-x) \right) - \frac{1}{276(1-x)} \right)$$

input `Int[(19*x)/((-1 + x)^3*(3 + 5*x + 4*x^2)^2),x]`

output `-19*(-1/276*(39 + 44*x)/((1 - x)^2*(3 + 5*x + 4*x^2)) + (21/(8*(1 - x)^2) + 97/(48*(1 - x)) - (6023*ArcTan[(5 + 8*x)/Sqrt[23]])/(576*Sqrt[23]) - (253*Log[1 - x])/576 + (253*Log[3 + 5*x + 4*x^2])/1152)/92)`

3.173.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 1200 `Int[(((d_) + (e_)*(x_))^(m_))*((f_) + (g_)*(x_))^(n_)]/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)^n/(a + b*x + c*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && IntegerQ[n]`

rule 1235 `Int[(((d_) + (e_)*(x_))^(m_))*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*(f*(b*c*d - b^2*e + 2*a*c*e) - a*g*(2*c*d - b*e) + c*(f*(2*c*d - b*e) - g*(b*d - 2*a*e))*x)/((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2))), x] + Simp[1/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)) Int[(d + e*x)^m*(a + b*x + c*x^2)^(p + 1)*Simp[f*(b*c*d*e*(2*p - m + 2) + b^2*e^2*(p + m + 2) - 2*c^2*d^2*(2*p + 3) - 2*a*c*e^2*(m + 2*p + 3) - g*(a*e*(b*e - 2*c*d*m + b*e*m) - b*d*(3*c*d - b*e + 2*c*d*p - b*e*p)) + c*e*(g*(b*d - 2*a*e) - f*(2*c*d - b*e))*(m + 2*p + 4)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && LtQ[p, -1] && (IntegerQ[m] || IntegerQ[p] || IntegerQ[2*m, 2*p])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.173. $\int \frac{19x}{(-1+x)^3(3+5x+4x^2)^2} dx$

3.173.4 Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.70

method	result
default	$-\frac{19}{288(-1+x)^2} + \frac{133}{864(-1+x)} + \frac{209 \ln(-1+x)}{2304} - \frac{19(-\frac{2204x}{23} - \frac{975}{23})}{6912(x^2 + \frac{5}{4}x + \frac{3}{4})} - \frac{209 \ln(4x^2+5x+3)}{4608} + \frac{114437 \arctan\left(\frac{(5+8x)\sqrt{23}}{23}\right)\sqrt{23}}{1218816}$
risch	$\frac{\frac{1843}{1104}x^3 - \frac{7733}{4416}x^2 - \frac{95}{184}x - \frac{285}{1472}}{(-1+x)^2(4x^2+5x+3)} + \frac{209 \ln(-1+x)}{2304} - \frac{209 \ln(580424464x^2+725530580x+435318348)}{4608} + \frac{114437\sqrt{23} \arctan\left(\frac{2(2409}{1218816}\right)}{1218816}$

input `int(19*x/(-1+x)^3/(4*x^2+5*x+3)^2,x,method=_RETURNVERBOSE)`output
$$-19/288/(-1+x)^2+133/864/(-1+x)+209/2304*\ln(-1+x)-19/6912*(-2204/23*x-975/23)/(x^2+5/4*x+3/4)-209/4608*\ln(4*x^2+5*x+3)+114437/1218816*\arctan(1/23*(5+8*x)*23^(1/2))*23^(1/2)$$
3.173.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.38

$$\int \frac{19x}{(-1+x)^3(3+5x+4x^2)^2} dx$$

$$= \frac{19(214176x^3 + 12046\sqrt{23}(4x^4 - 3x^3 - 3x^2 - x + 3)) \arctan\left(\frac{1}{23}\sqrt{23}(8x+5)\right) - 224664x^2 - 5819(4x^4 - 3x^3 - 3x^2 - x + 3) \log(4x^2 + 5x + 3) + 11638(4x^4 - 3x^3 - 3x^2 - x + 3) \log(x - 1) - 66240x - 24840}{2437632(4x^4 - 3x^3 - 3x^2 - x + 3)}$$

input `integrate(19*x/(-1+x)^3/(4*x^2+5*x+3)^2,x, algorithm="fricas")`output
$$19/2437632*(214176*x^3 + 12046*\sqrt{23}*(4*x^4 - 3*x^3 - 3*x^2 - x + 3)*\arctan(1/23*\sqrt{23}*(8*x + 5)) - 224664*x^2 - 5819*(4*x^4 - 3*x^3 - 3*x^2 - x + 3)*\log(4*x^2 + 5*x + 3) + 11638*(4*x^4 - 3*x^3 - 3*x^2 - x + 3)*\log(x - 1) - 66240*x - 24840)/(4*x^4 - 3*x^3 - 3*x^2 - x + 3)$$

3.173.6 Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.91

$$\int \frac{19x}{(-1+x)^3(3+5x+4x^2)^2} dx = \frac{19 \cdot (388x^3 - 407x^2 - 120x - 45)}{17664x^4 - 13248x^3 - 13248x^2 - 4416x + 13248} + \frac{209 \log(x-1)}{2304} - \frac{209 \log\left(x^2 + \frac{5x}{4} + \frac{3}{4}\right)}{4608} + \frac{114437\sqrt{23} \operatorname{atan}\left(\frac{8\sqrt{23}x}{23} + \frac{5\sqrt{23}}{23}\right)}{1218816}$$

input `integrate(19*x/(-1+x)**3/(4*x**2+5*x+3)**2,x)`output `19*(388*x**3 - 407*x**2 - 120*x - 45)/(17664*x**4 - 13248*x**3 - 13248*x**2 - 4416*x + 13248) + 209*log(x - 1)/2304 - 209*log(x**2 + 5*x/4 + 3/4)/4608 + 114437*sqrt(23)*atan(8*sqrt(23)*x/23 + 5*sqrt(23)/23)/1218816`**3.173.7 Maxima [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.77

$$\int \frac{19x}{(-1+x)^3(3+5x+4x^2)^2} dx = \frac{114437}{1218816} \sqrt{23} \arctan\left(\frac{1}{23} \sqrt{23}(8x+5)\right) + \frac{19(388x^3 - 407x^2 - 120x - 45)}{4416(4x^4 - 3x^3 - 3x^2 - x + 3)} - \frac{209}{4608} \log(4x^2 + 5x + 3) + \frac{209}{2304} \log(x-1)$$

input `integrate(19*x/(-1+x)^3/(4*x^2+5*x+3)^2,x, algorithm="maxima")`output `114437/1218816*sqrt(23)*arctan(1/23*sqrt(23)*(8*x + 5)) + 19/4416*(388*x^3 - 407*x^2 - 120*x - 45)/(4*x^4 - 3*x^3 - 3*x^2 - x + 3) - 209/4608*log(4*x^2 + 5*x + 3) + 209/2304*log(x - 1)`

3.173.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.73

$$\int \frac{19x}{(-1+x)^3(3+5x+4x^2)^2} dx = \frac{114437}{1218816} \sqrt{23} \arctan\left(\frac{1}{23} \sqrt{23}(8x+5)\right) + \frac{19(388x^3 - 407x^2 - 120x - 45)}{4416(4x^2 + 5x + 3)(x-1)^2} - \frac{209}{4608} \log(4x^2 + 5x + 3) + \frac{209}{2304} \log(|x-1|)$$

input `integrate(19*x/(-1+x)^3/(4*x^2+5*x+3)^2,x, algorithm="giac")`output `114437/1218816*sqrt(23)*arctan(1/23*sqrt(23)*(8*x + 5)) + 19/4416*(388*x^3 - 407*x^2 - 120*x - 45)/((4*x^2 + 5*x + 3)*(x - 1)^2) - 209/4608*log(4*x^2 + 5*x + 3) + 209/2304*log(abs(x - 1))`**3.173.9 Mupad [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.87

$$\int \frac{19x}{(-1+x)^3(3+5x+4x^2)^2} dx = \frac{209 \ln(x-1)}{2304} + \frac{-\frac{1843x^3}{4416} + \frac{7733x^2}{17664} + \frac{95x}{736} + \frac{285}{5888}}{-x^4 + \frac{3x^3}{4} + \frac{3x^2}{4} + \frac{x}{4} - \frac{3}{4}} - \ln\left(x + \frac{5}{8} - \frac{\sqrt{23}i}{8}\right) \left(\frac{209}{4608} + \frac{\sqrt{23}114437i}{2437632}\right) + \ln\left(x + \frac{5}{8} + \frac{\sqrt{23}i}{8}\right) \left(-\frac{209}{4608} + \frac{\sqrt{23}114437i}{2437632}\right)$$

input `int((19*x)/((x - 1)^3*(5*x + 4*x^2 + 3)^2),x)`output `(209*log(x - 1))/2304 + ((95*x)/736 + (7733*x^2)/17664 - (1843*x^3)/4416 + 285/5888)/(x/4 + (3*x^2)/4 + (3*x^3)/4 - x^4 - 3/4) - log(x - (23^(1/2)*1i)/8 + 5/8)*((23^(1/2)*114437i)/2437632 + 209/4608) + log(x + (23^(1/2)*1i)/8 + 5/8)*((23^(1/2)*114437i)/2437632 - 209/4608)`

3.174 $\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx$

3.174.1 Optimal result	929
3.174.2 Mathematica [A] (verified)	929
3.174.3 Rubi [A] (verified)	930
3.174.4 Maple [A] (verified)	931
3.174.5 Fricas [A] (verification not implemented)	931
3.174.6 Sympy [A] (verification not implemented)	932
3.174.7 Maxima [A] (verification not implemented)	932
3.174.8 Giac [A] (verification not implemented)	932
3.174.9 Mupad [B] (verification not implemented)	933

3.174.1 Optimal result

Integrand size = 23, antiderivative size = 46

$$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx = -\frac{1}{2x} + \frac{\arctan\left(\frac{1+2x}{\sqrt{7}}\right)}{4\sqrt{7}} - \frac{\log(x)}{4} + \frac{5}{8} \log(2+x+x^2)$$

output `-1/2/x-1/4*ln(x)+5/8*ln(x^2+x+2)+1/28*arctan(1/7*(1+2*x)*7^(1/2))*7^(1/2)`

3.174.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx = -\frac{1}{2x} + \frac{\arctan\left(\frac{1+2x}{\sqrt{7}}\right)}{4\sqrt{7}} - \frac{\log(x)}{4} + \frac{5}{8} \log(2+x+x^2)$$

input `Integrate[(1 + x^2 + x^3)/(2*x^2 + x^3 + x^4),x]`

output `-1/2*1/x + ArcTan[(1 + 2*x)/Sqrt[7]]/(4*Sqrt[7]) - Log[x]/4 + (5*Log[2 + x + x^2])/8`

3.174.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {2026, 2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^3 + x^2 + 1}{x^4 + x^3 + 2x^2} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{x^3 + x^2 + 1}{x^2(x^2 + x + 2)} dx \\ & \quad \downarrow \text{2159} \\ & \int \left(\frac{5x + 3}{4(x^2 + x + 2)} + \frac{1}{2x^2} - \frac{1}{4x} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{\arctan\left(\frac{2x+1}{\sqrt{7}}\right)}{4\sqrt{7}} + \frac{5}{8} \log(x^2 + x + 2) - \frac{1}{2x} - \frac{\log(x)}{4} \end{aligned}$$

input `Int[(1 + x^2 + x^3)/(2*x^2 + x^3 + x^4), x]`

output `-1/2*1/x + ArcTan[(1 + 2*x)/Sqrt[7]]/(4*Sqrt[7]) - Log[x]/4 + (5*Log[2 + x + x^2])/8`

3.174.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2159 `Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^(m)*Pq*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

3.174.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.74

method	result	size
risch	$-\frac{1}{2x} - \frac{\ln(x)}{4} + \frac{5 \ln(x^2+x+2)}{8} + \frac{\sqrt{7} \arctan\left(\frac{2(x+\frac{1}{2})\sqrt{7}}{7}\right)}{28}$	34
default	$-\frac{1}{2x} - \frac{\ln(x)}{4} + \frac{5 \ln(x^2+x+2)}{8} + \frac{\arctan\left(\frac{(1+2x)\sqrt{7}}{7}\right)\sqrt{7}}{28}$	36

input `int((x^3+x^2+1)/(x^4+x^3+2*x^2),x,method=_RETURNVERBOSE)`

output `-1/2/x-1/4*ln(x)+5/8*ln(x^2+x+2)+1/28*7^(1/2)*arctan(2/7*(x+1/2)*7^(1/2))`

3.174.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.85

$$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx$$

$$= \frac{2\sqrt{7}x \arctan\left(\frac{1}{7}\sqrt{7}(2x+1)\right) + 35x \log(x^2+x+2) - 14x \log(x) - 28}{56x}$$

input `integrate((x^3+x^2+1)/(x^4+x^3+2*x^2),x, algorithm="fricas")`

output `1/56*(2*sqrt(7)*x*arctan(1/7*sqrt(7)*(2*x + 1)) + 35*x*log(x^2 + x + 2) - 14*x*log(x) - 28)/x`

3.174.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx = -\frac{\log(x)}{4} + \frac{5\log(x^2+x+2)}{8} + \frac{\sqrt{7}\operatorname{atan}\left(\frac{2\sqrt{7}x}{7} + \frac{\sqrt{7}}{7}\right)}{28} - \frac{1}{2x}$$

input `integrate((x**3+x**2+1)/(x**4+x**3+2*x**2),x)`output `-log(x)/4 + 5*log(x**2 + x + 2)/8 + sqrt(7)*atan(2*sqrt(7)*x/7 + sqrt(7)/7)/28 - 1/(2*x)`**3.174.7 Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.76

$$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx = \frac{1}{28} \sqrt{7} \arctan\left(\frac{1}{7} \sqrt{7}(2x+1)\right) - \frac{1}{2x} + \frac{5}{8} \log(x^2+x+2) - \frac{1}{4} \log(x)$$

input `integrate((x^3+x^2+1)/(x^4+x^3+2*x^2),x, algorithm="maxima")`output `1/28*sqrt(7)*arctan(1/7*sqrt(7)*(2*x + 1)) - 1/2/x + 5/8*log(x^2 + x + 2) - 1/4*log(x)`**3.174.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.78

$$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx = \frac{1}{28} \sqrt{7} \arctan\left(\frac{1}{7} \sqrt{7}(2x+1)\right) - \frac{1}{2x} + \frac{5}{8} \log(x^2+x+2) - \frac{1}{4} \log(|x|)$$

input `integrate((x^3+x^2+1)/(x^4+x^3+2*x^2),x, algorithm="giac")`output `1/28*sqrt(7)*arctan(1/7*sqrt(7)*(2*x + 1)) - 1/2/x + 5/8*log(x^2 + x + 2) - 1/4*log(abs(x))`

3.174.9 Mupad [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.07

$$\int \frac{1+x^2+x^3}{2x^2+x^3+x^4} dx = -\frac{\ln(x)}{4} - \ln\left(x + \frac{1}{2} - \frac{\sqrt{7}1i}{2}\right) \left(-\frac{5}{8} + \frac{\sqrt{7}1i}{56}\right) \\ + \ln\left(x + \frac{1}{2} + \frac{\sqrt{7}1i}{2}\right) \left(\frac{5}{8} + \frac{\sqrt{7}1i}{56}\right) - \frac{1}{2x}$$

input `int((x^2 + x^3 + 1)/(2*x^2 + x^3 + x^4),x)`output `log(x + (7^(1/2)*1i)/2 + 1/2)*((7^(1/2)*1i)/56 + 5/8) - log(x - (7^(1/2)*1i)/2 + 1/2)*((7^(1/2)*1i)/56 - 5/8) - log(x)/4 - 1/(2*x)`

3.175 $\int \frac{1}{-x^3+x^6} dx$

3.175.1 Optimal result	934
3.175.2 Mathematica [A] (verified)	934
3.175.3 Rubi [A] (verified)	935
3.175.4 Maple [A] (verified)	937
3.175.5 Fricas [A] (verification not implemented)	938
3.175.6 Sympy [A] (verification not implemented)	938
3.175.7 Maxima [A] (verification not implemented)	938
3.175.8 Giac [A] (verification not implemented)	939
3.175.9 Mupad [B] (verification not implemented)	939

3.175.1 Optimal result

Integrand size = 11, antiderivative size = 48

$$\int \frac{1}{-x^3+x^6} dx = \frac{1}{2x^2} - \frac{\arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{3}\log(1-x) - \frac{1}{6}\log(1+x+x^2)$$

output `1/2/x^2+1/3*ln(1-x)-1/6*ln(x^2+x+1)-1/3*arctan(1/3*(1+2*x)*3^(1/2))*3^(1/2)`
`)`

3.175.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00

$$\int \frac{1}{-x^3+x^6} dx = \frac{1}{2x^2} - \frac{\arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{3}\log(1-x) - \frac{1}{6}\log(1+x+x^2)$$

input `Integrate[(-x^3 + x^6)^(-1),x]`

output `1/(2*x^2) - ArcTan[(1 + 2*x)/Sqrt[3]]/Sqrt[3] + Log[1 - x]/3 - Log[1 + x + x^2]/6`

3.175.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.10, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.818$, Rules used = {2026, 847, 750, 16, 25, 1142, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^6 - x^3} dx \\
 & \quad \downarrow \text{2026} \\
 & \int \frac{1}{x^3(x^3 - 1)} dx \\
 & \quad \downarrow \text{847} \\
 & \int \frac{1}{x^3 - 1} dx + \frac{1}{2x^2} \\
 & \quad \downarrow \text{750} \\
 & \frac{1}{3} \int -\frac{x+2}{x^2+x+1} dx + \frac{1}{3} \int \frac{1}{x-1} dx + \frac{1}{2x^2} \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{3} \int -\frac{x+2}{x^2+x+1} dx + \frac{1}{2x^2} + \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{25} \\
 & -\frac{1}{3} \int \frac{x+2}{x^2+x+1} dx + \frac{1}{2x^2} + \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{1142} \\
 & \frac{1}{3} \left(-\frac{3}{2} \int \frac{1}{x^2+x+1} dx - \frac{1}{2} \int \frac{2x+1}{x^2+x+1} dx \right) + \frac{1}{2x^2} + \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{1083} \\
 & \frac{1}{3} \left(3 \int \frac{1}{-(2x+1)^2-3} d(2x+1) - \frac{1}{2} \int \frac{2x+1}{x^2+x+1} dx \right) + \frac{1}{2x^2} + \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{217} \\
 & \frac{1}{3} \left(-\frac{1}{2} \int \frac{2x+1}{x^2+x+1} dx - \sqrt{3} \arctan \left(\frac{2x+1}{\sqrt{3}} \right) \right) + \frac{1}{2x^2} + \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{1103}
 \end{aligned}$$

$$\frac{1}{3} \left(-\sqrt{3} \arctan \left(\frac{2x+1}{\sqrt{3}} \right) - \frac{1}{2} \log(x^2 + x + 1) \right) + \frac{1}{2x^2} + \frac{1}{3} \log(1-x)$$

input `Int[(-x^3 + x^6)^(-1),x]`

output `1/(2*x^2) + Log[1 - x]/3 + (- (Sqrt[3]*ArcTan[(1 + 2*x)/Sqrt[3]]) - Log[1 + x + x^2]/2)/3`

3.175.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(- (Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])`

rule 750 `Int[((a_) + (b_.)*(x_)^3)^(-1), x_Symbol] := Simp[1/(3*Rt[a, 3]^2) Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Simp[1/(3*Rt[a, 3]^2) Int[(2*Rt[a, 3] - Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]`

rule 847 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*((m + n*(p + 1) + 1)/(a*c^n*(m + 1)) Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c)
Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 2026 `Int[(F*_)(P*_)^*(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p
*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && Integ
erQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

3.175.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{\ln(-1+x)}{3} - \frac{\ln(x^2+x+1)}{6} - \frac{\arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3} + \frac{1}{2x^2}$	38
risch	$\frac{1}{2x^2} - \frac{\ln(4x^2+4x+4)}{6} - \frac{\arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3} + \frac{\ln(-1+x)}{3}$	42
meijerg	$- \frac{(-1)^{\frac{2}{3}} \left(\frac{3(-1)^{\frac{1}{3}}}{2x^2} + \frac{x(-1)^{\frac{1}{3}} \left(\ln\left(1 - (x^3)^{\frac{1}{3}}\right) - \frac{\ln\left(1 + (x^3)^{\frac{1}{3}} + (x^3)^{\frac{2}{3}}\right)}{2} - \sqrt{3} \arctan\left(\frac{\sqrt{3}(x^3)^{\frac{1}{3}}}{2 + (x^3)^{\frac{1}{3}}}\right) \right)}{(x^3)^{\frac{1}{3}}} \right)}{3}$	78

input `int(1/(x^6-x^3), x, method=_RETURNVERBOSE)`

output `1/3*ln(-1+x)-1/6*ln(x^2+x+1)-1/3*arctan(1/3*(1+2*x)*3^(1/2))*3^(1/2)+1/2/x
^2`

3.175.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.96

$$\int \frac{1}{-x^3 + x^6} dx = -\frac{2\sqrt{3}x^2 \arctan\left(\frac{1}{3}\sqrt{3}(2x+1)\right) + x^2 \log(x^2 + x + 1) - 2x^2 \log(x-1) - 3}{6x^2}$$

input `integrate(1/(x^6-x^3),x, algorithm="fricas")`output `-1/6*(2*sqrt(3)*x^2*arctan(1/3*sqrt(3)*(2*x + 1)) + x^2*log(x^2 + x + 1) - 2*x^2*log(x - 1) - 3)/x^2`**3.175.6 Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00

$$\int \frac{1}{-x^3 + x^6} dx = \frac{\log(x-1)}{3} - \frac{\log(x^2 + x + 1)}{6} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x}{3} + \frac{\sqrt{3}}{3}\right)}{3} + \frac{1}{2x^2}$$

input `integrate(1/(x**6-x**3),x)`output `log(x - 1)/3 - log(x**2 + x + 1)/6 - sqrt(3)*atan(2*sqrt(3)*x/3 + sqrt(3)/3)/3 + 1/(2*x**2)`**3.175.7 Maxima [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.77

$$\int \frac{1}{-x^3 + x^6} dx = -\frac{1}{3}\sqrt{3} \arctan\left(\frac{1}{3}\sqrt{3}(2x+1)\right) + \frac{1}{2x^2} - \frac{1}{6} \log(x^2 + x + 1) + \frac{1}{3} \log(x-1)$$

input `integrate(1/(x^6-x^3),x, algorithm="maxima")`output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + 1/2/x^2 - 1/6*log(x^2 + x + 1) + 1/3*log(x - 1)`

3.175.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.79

$$\int \frac{1}{-x^3 + x^6} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x + 1)\right) + \frac{1}{2x^2} - \frac{1}{6} \log(x^2 + x + 1) + \frac{1}{3} \log(|x - 1|)$$

input `integrate(1/(x^6-x^3),x, algorithm="giac")`output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + 1/2/x^2 - 1/6*log(x^2 + x + 1) + 1/3*log(abs(x - 1))`**3.175.9 Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.06

$$\int \frac{1}{-x^3 + x^6} dx = \frac{\ln(x - 1)}{3} + \ln\left(x + \frac{1}{2} - \frac{\sqrt{3} \text{li}}{2}\right) \left(-\frac{1}{6} + \frac{\sqrt{3} \text{li}}{6}\right) - \ln\left(x + \frac{1}{2} + \frac{\sqrt{3} \text{li}}{2}\right) \left(\frac{1}{6} + \frac{\sqrt{3} \text{li}}{6}\right) + \frac{1}{2x^2}$$

input `int(-1/(x^3 - x^6),x)`output `log(x - 1)/3 + log(x - (3^(1/2)*1i)/2 + 1/2)*((3^(1/2)*1i)/6 - 1/6) - log(x + (3^(1/2)*1i)/2 + 1/2)*((3^(1/2)*1i)/6 + 1/6) + 1/(2*x^2)`

3.176 $\int \frac{x^2}{1+x} dx$

3.176.1 Optimal result	940
3.176.2 Mathematica [A] (verified)	940
3.176.3 Rubi [A] (verified)	941
3.176.4 Maple [A] (verified)	942
3.176.5 Fricas [A] (verification not implemented)	942
3.176.6 Sympy [A] (verification not implemented)	942
3.176.7 Maxima [A] (verification not implemented)	943
3.176.8 Giac [A] (verification not implemented)	943
3.176.9 Mupad [B] (verification not implemented)	943

3.176.1 Optimal result

Integrand size = 9, antiderivative size = 15

$$\int \frac{x^2}{1+x} dx = -x + \frac{x^2}{2} + \log(1+x)$$

output `-x+1/2*x^2+ln(1+x)`

3.176.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.27

$$\int \frac{x^2}{1+x} dx = -2(1+x) + \frac{1}{2}(1+x)^2 + \log(1+x)$$

input `Integrate[x^2/(1 + x),x]`

output `-2*(1 + x) + (1 + x)^2/2 + Log[1 + x]`

3.176.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{x+1} dx$$

$$\downarrow 49$$

$$\int \left(x + \frac{1}{x+1} - 1 \right) dx$$

$$\downarrow 2009$$

$$\frac{x^2}{2} - x + \log(x+1)$$

input `Int[x^2/(1 + x),x]`

output `-x + x^2/2 + Log[1 + x]`

3.176.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.176.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
default	$-x + \frac{x^2}{2} + \ln(1+x)$	14
norman	$-x + \frac{x^2}{2} + \ln(1+x)$	14
meijerg	$-\frac{x(-3x+6)}{6} + \ln(1+x)$	14
risch	$-x + \frac{x^2}{2} + \ln(1+x)$	14
parallelrisc	$-x + \frac{x^2}{2} + \ln(1+x)$	14

input `int(x^2/(1+x),x,method=_RETURNVERBOSE)`output `-x+1/2*x^2+ln(1+x)`**3.176.5 Fricas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{x^2}{1+x} dx = \frac{1}{2}x^2 - x + \log(x+1)$$

input `integrate(x^2/(1+x),x, algorithm="fricas")`output `1/2*x^2 - x + log(x + 1)`**3.176.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int \frac{x^2}{1+x} dx = \frac{x^2}{2} - x + \log(x+1)$$

input `integrate(x**2/(1+x),x)`output `x**2/2 - x + log(x + 1)`

3.176.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{x^2}{1+x} dx = \frac{1}{2}x^2 - x + \log(x+1)$$

input `integrate(x^2/(1+x),x, algorithm="maxima")`output `1/2*x^2 - x + log(x + 1)`**3.176.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{x^2}{1+x} dx = \frac{1}{2}x^2 - x + \log(|x+1|)$$

input `integrate(x^2/(1+x),x, algorithm="giac")`output `1/2*x^2 - x + log(abs(x + 1))`**3.176.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{x^2}{1+x} dx = \ln(x+1) - x + \frac{x^2}{2}$$

input `int(x^2/(x + 1),x)`output `log(x + 1) - x + x^2/2`

3.177 $\int \frac{x}{-5+x} dx$

3.177.1 Optimal result	944
3.177.2 Mathematica [A] (verified)	944
3.177.3 Rubi [A] (verified)	945
3.177.4 Maple [A] (verified)	946
3.177.5 Fricas [A] (verification not implemented)	946
3.177.6 Sympy [A] (verification not implemented)	946
3.177.7 Maxima [A] (verification not implemented)	947
3.177.8 Giac [A] (verification not implemented)	947
3.177.9 Mupad [B] (verification not implemented)	947

3.177.1 Optimal result

Integrand size = 7, antiderivative size = 10

$$\int \frac{x}{-5+x} dx = x + 5 \log(5-x)$$

output `x+5*ln(5-x)`

3.177.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{x}{-5+x} dx = x + 5 \log(-5+x)$$

input `Integrate[x/(-5 + x),x]`

output `x + 5*Log[-5 + x]`

3.177.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{x}{x-5} dx \\ \downarrow 49 \\ \int \left(\frac{5}{x-5} + 1 \right) dx \\ \downarrow 2009 \\ x + 5 \log(5-x) \end{array}$$

input `Int[x/(-5 + x), x]`

output `x + 5*Log[5 - x]`

3.177.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.177.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

method	result	size
default	$x + 5 \ln(x - 5)$	9
norman	$x + 5 \ln(x - 5)$	9
risch	$x + 5 \ln(x - 5)$	9
parallelrisch	$x + 5 \ln(x - 5)$	9
meijerg	$x + 5 \ln\left(1 - \frac{x}{5}\right)$	11

input `int(x/(x-5),x,method=_RETURNVERBOSE)`output `x+5*ln(x-5)`**3.177.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{x}{-5+x} dx = x + 5 \log(x - 5)$$

input `integrate(x/(-5+x),x, algorithm="fricas")`output `x + 5*log(x - 5)`**3.177.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \frac{x}{-5+x} dx = x + 5 \log(x - 5)$$

input `integrate(x/(-5+x),x)`output `x + 5*log(x - 5)`

3.177.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{x}{-5+x} dx = x + 5 \log(x - 5)$$

input `integrate(x/(-5+x),x, algorithm="maxima")`output `x + 5*log(x - 5)`**3.177.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{x}{-5+x} dx = x + 5 \log(|x - 5|)$$

input `integrate(x/(-5+x),x, algorithm="giac")`output `x + 5*log(abs(x - 5))`**3.177.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{x}{-5+x} dx = x + 5 \ln(x - 5)$$

input `int(x/(x - 5),x)`output `x + 5*log(x - 5)`

3.178 $\int \frac{-1+4x}{(-1+x)(2+x)} dx$

3.178.1 Optimal result	948
3.178.2 Mathematica [A] (verified)	948
3.178.3 Rubi [A] (verified)	949
3.178.4 Maple [A] (verified)	950
3.178.5 Fricas [A] (verification not implemented)	950
3.178.6 Sympy [A] (verification not implemented)	950
3.178.7 Maxima [A] (verification not implemented)	951
3.178.8 Giac [A] (verification not implemented)	951
3.178.9 Mupad [B] (verification not implemented)	951

3.178.1 Optimal result

Integrand size = 16, antiderivative size = 13

$$\int \frac{-1 + 4x}{(-1 + x)(2 + x)} dx = \log(1 - x) + 3 \log(2 + x)$$

output `ln(1-x)+3*ln(2+x)`

3.178.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{-1 + 4x}{(-1 + x)(2 + x)} dx = \log(1 - x) + 3 \log(2 + x)$$

input `Integrate[(-1 + 4*x)/((-1 + x)*(2 + x)),x]`

output `Log[1 - x] + 3*Log[2 + x]`

3.178.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{4x-1}{(x-1)(x+2)} dx$$

↓ 86

$$\int \left(\frac{3}{x+2} + \frac{1}{x-1} \right) dx$$

↓ 2009

$$\log(1-x) + 3\log(x+2)$$

input `Int[(-1 + 4*x)/((-1 + x)*(2 + x)),x]`

output `Log[1 - x] + 3*Log[2 + x]`

3.178.3.1 Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.178.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
default	$\ln(-1+x) + 3\ln(2+x)$	12
norman	$\ln(-1+x) + 3\ln(2+x)$	12
risch	$\ln(-1+x) + 3\ln(2+x)$	12
parallelrisch	$\ln(-1+x) + 3\ln(2+x)$	12

input `int((-1+4*x)/(-1+x)/(2+x),x,method=_RETURNVERBOSE)`output `ln(-1+x)+3*ln(2+x)`**3.178.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{-1+4x}{(-1+x)(2+x)} dx = 3 \log(x+2) + \log(x-1)$$

input `integrate((-1+4*x)/(-1+x)/(2+x),x, algorithm="fracas")`output `3*log(x + 2) + log(x - 1)`**3.178.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \frac{-1+4x}{(-1+x)(2+x)} dx = \log(x-1) + 3\log(x+2)$$

input `integrate((-1+4*x)/(-1+x)/(2+x),x)`output `log(x - 1) + 3*log(x + 2)`

3.178.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{-1 + 4x}{(-1 + x)(2 + x)} dx = 3 \log(x + 2) + \log(x - 1)$$

input `integrate((-1+4*x)/(-1+x)/(2+x),x, algorithm="maxima")`output `3*log(x + 2) + log(x - 1)`**3.178.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{-1 + 4x}{(-1 + x)(2 + x)} dx = 3 \log(|x + 2|) + \log(|x - 1|)$$

input `integrate((-1+4*x)/(-1+x)/(2+x),x, algorithm="giac")`output `3*log(abs(x + 2)) + log(abs(x - 1))`**3.178.9 Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{-1 + 4x}{(-1 + x)(2 + x)} dx = \ln(x - 1) + 3 \ln(x + 2)$$

input `int((4*x - 1)/((x - 1)*(x + 2)),x)`output `log(x - 1) + 3*log(x + 2)`

3.179 $\int \frac{1}{(1+x)(2+x)} dx$

3.179.1 Optimal result	952
3.179.2 Mathematica [A] (verified)	952
3.179.3 Rubi [A] (verified)	953
3.179.4 Maple [A] (verified)	954
3.179.5 Fricas [A] (verification not implemented)	954
3.179.6 Sympy [A] (verification not implemented)	954
3.179.7 Maxima [A] (verification not implemented)	955
3.179.8 Giac [A] (verification not implemented)	955
3.179.9 Mupad [B] (verification not implemented)	955

3.179.1 Optimal result

Integrand size = 11, antiderivative size = 11

$$\int \frac{1}{(1+x)(2+x)} dx = \log(1+x) - \log(2+x)$$

output `ln(1+x)-ln(2+x)`

3.179.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x)(2+x)} dx = \log(1+x) - \log(2+x)$$

input `Integrate[1/((1 + x)*(2 + x)),x]`

output `Log[1 + x] - Log[2 + x]`

3.179.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {47, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x+1)(x+2)} dx$$

$$\downarrow 47$$

$$\int \frac{1}{x+1} dx - \int \frac{1}{x+2} dx$$

$$\downarrow 16$$

$$\log(x+1) - \log(x+2)$$

input `Int[1/((1 + x)*(2 + x)),x]`

output `Log[1 + x] - Log[2 + x]`

3.179.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

3.179.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
default	$\ln(1+x) - \ln(2+x)$	12
norman	$\ln(1+x) - \ln(2+x)$	12
risch	$\ln(1+x) - \ln(2+x)$	12
parallelrisc	$\ln(1+x) - \ln(2+x)$	12

input `int(1/(1+x)/(2+x),x,method=_RETURNVERBOSE)`output `ln(1+x)-ln(2+x)`**3.179.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x)(2+x)} dx = -\log(x+2) + \log(x+1)$$

input `integrate(1/(1+x)/(2+x),x, algorithm="fracas")`output `-log(x + 2) + log(x + 1)`**3.179.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \frac{1}{(1+x)(2+x)} dx = \log(x+1) - \log(x+2)$$

input `integrate(1/(1+x)/(2+x),x)`output `log(x + 1) - log(x + 2)`

3.179.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x)(2+x)} dx = -\log(x+2) + \log(x+1)$$

input `integrate(1/(1+x)/(2+x),x, algorithm="maxima")`output `-log(x + 2) + log(x + 1)`**3.179.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.18

$$\int \frac{1}{(1+x)(2+x)} dx = -\log(|x+2|) + \log(|x+1|)$$

input `integrate(1/(1+x)/(2+x),x, algorithm="giac")`output `-log(abs(x + 2)) + log(abs(x + 1))`**3.179.9 Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int \frac{1}{(1+x)(2+x)} dx = \ln\left(1 - \frac{1}{x+2}\right)$$

input `int(1/((x + 1)*(x + 2)),x)`output `log(1 - 1/(x + 2))`

3.180 $\int \frac{-5+6x}{3+2x} dx$

3.180.1 Optimal result	956
3.180.2 Mathematica [A] (verified)	956
3.180.3 Rubi [A] (verified)	957
3.180.4 Maple [A] (verified)	958
3.180.5 Fricas [A] (verification not implemented)	958
3.180.6 Sympy [A] (verification not implemented)	958
3.180.7 Maxima [A] (verification not implemented)	959
3.180.8 Giac [A] (verification not implemented)	959
3.180.9 Mupad [B] (verification not implemented)	959

3.180.1 Optimal result

Integrand size = 13, antiderivative size = 12

$$\int \frac{-5+6x}{3+2x} dx = 3x - 7\log(3+2x)$$

output `3*x-7*ln(3+2*x)`

3.180.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.25

$$\int \frac{-5+6x}{3+2x} dx = \frac{9}{2} + 3x - 7\log(3+2x)$$

input `Integrate[(-5 + 6*x)/(3 + 2*x),x]`

output `9/2 + 3*x - 7*Log[3 + 2*x]`

3.180.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{6x - 5}{2x + 3} dx$$

$$\downarrow 49$$

$$\int \left(3 - \frac{14}{2x + 3} \right) dx$$

$$\downarrow 2009$$

$$3x - 7 \log(2x + 3)$$

input `Int[(-5 + 6*x)/(3 + 2*x),x]`

output `3*x - 7*Log[3 + 2*x]`

3.180.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.180.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
parallelrisc	$3x - 7 \ln\left(\frac{3}{2} + x\right)$	11
default	$3x - 7 \ln(3 + 2x)$	13
norman	$3x - 7 \ln(3 + 2x)$	13
meijerg	$-7 \ln\left(1 + \frac{2x}{3}\right) + 3x$	13
risc	$3x - 7 \ln(3 + 2x)$	13

input `int((6*x-5)/(3+2*x),x,method=_RETURNVERBOSE)`output `3*x-7*ln(3/2+x)`**3.180.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{-5 + 6x}{3 + 2x} dx = 3x - 7 \log(2x + 3)$$

input `integrate((-5+6*x)/(3+2*x),x, algorithm="fricas")`output `3*x - 7*log(2*x + 3)`**3.180.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{-5 + 6x}{3 + 2x} dx = 3x - 7 \log(2x + 3)$$

input `integrate((-5+6*x)/(3+2*x),x)`output `3*x - 7*log(2*x + 3)`

3.180.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{-5 + 6x}{3 + 2x} dx = 3x - 7 \log(2x + 3)$$

input `integrate((-5+6*x)/(3+2*x),x, algorithm="maxima")`output `3*x - 7*log(2*x + 3)`**3.180.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

$$\int \frac{-5 + 6x}{3 + 2x} dx = 3x - 7 \log(|2x + 3|)$$

input `integrate((-5+6*x)/(3+2*x),x, algorithm="giac")`output `3*x - 7*log(abs(2*x + 3))`**3.180.9 Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{-5 + 6x}{3 + 2x} dx = 3x - 7 \ln\left(x + \frac{3}{2}\right)$$

input `int((6*x - 5)/(2*x + 3),x)`output `3*x - 7*log(x + 3/2)`

3.181 $\int \frac{1}{(a+x)(b+x)} dx$

3.181.1 Optimal result	960
3.181.2 Mathematica [A] (verified)	960
3.181.3 Rubi [A] (verified)	961
3.181.4 Maple [A] (verified)	962
3.181.5 Fricas [A] (verification not implemented)	962
3.181.6 Sympy [B] (verification not implemented)	962
3.181.7 Maxima [A] (verification not implemented)	963
3.181.8 Giac [A] (verification not implemented)	963
3.181.9 Mupad [B] (verification not implemented)	963

3.181.1 Optimal result

Integrand size = 11, antiderivative size = 26

$$\int \frac{1}{(a+x)(b+x)} dx = -\frac{\log(a+x)}{a-b} + \frac{\log(b+x)}{a-b}$$

output `-ln(a+x)/(a-b)+ln(b+x)/(a-b)`

3.181.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.73

$$\int \frac{1}{(a+x)(b+x)} dx = \frac{-\log(a+x) + \log(b+x)}{a-b}$$

input `Integrate[1/((a + x)*(b + x)),x]`

output `(-Log[a + x] + Log[b + x])/(a - b)`

3.181.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {47, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a+x)(b+x)} dx$$

$$\downarrow 47$$

$$\frac{\int \frac{1}{b+x} dx}{a-b} - \frac{\int \frac{1}{a+x} dx}{a-b}$$

$$\downarrow 16$$

$$\frac{\log(b+x)}{a-b} - \frac{\log(a+x)}{a-b}$$

input `Int[1/((a + x)*(b + x)),x]`

output `-(Log[a + x]/(a - b)) + Log[b + x]/(a - b)`

3.181.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

3.181.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.81

method	result	size
parallelrisch	$-\frac{\ln(a+x)-\ln(b+x)}{a-b}$	21
default	$-\frac{\ln(a+x)}{a-b} + \frac{\ln(b+x)}{a-b}$	27
norman	$-\frac{\ln(a+x)}{a-b} + \frac{\ln(b+x)}{a-b}$	27
risch	$-\frac{\ln(a+x)}{a-b} + \frac{\ln(b+x)}{a-b}$	27

input `int(1/(a+x)/(b+x),x,method=_RETURNVERBOSE)`

output $-(\ln(a+x)-\ln(b+x))/(a-b)$

3.181.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \frac{1}{(a+x)(b+x)} dx = -\frac{\log(a+x) - \log(b+x)}{a-b}$$

input `integrate(1/(a+x)/(b+x),x, algorithm="fricas")`

output $-(\log(a+x) - \log(b+x))/(a-b)$

3.181.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 80 vs. $2(15) = 30$.

Time = 0.12 (sec) , antiderivative size = 80, normalized size of antiderivative = 3.08

$$\int \frac{1}{(a+x)(b+x)} dx = \frac{\log\left(-\frac{a^2}{2(a-b)} + \frac{ab}{a-b} + \frac{a}{2} - \frac{b^2}{2(a-b)} + \frac{b}{2} + x\right)}{a-b} - \frac{\log\left(\frac{a^2}{2(a-b)} - \frac{ab}{a-b} + \frac{a}{2} + \frac{b^2}{2(a-b)} + \frac{b}{2} + x\right)}{a-b}$$

input `integrate(1/(a+x)/(b+x),x)`

output `log(-a**2/(2*(a - b)) + a*b/(a - b) + a/2 - b**2/(2*(a - b)) + b/2 + x)/(a - b) - log(a**2/(2*(a - b)) - a*b/(a - b) + a/2 + b**2/(2*(a - b)) + b/2 + x)/(a - b)`

3.181.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a+x)(b+x)} dx = -\frac{\log(a+x)}{a-b} + \frac{\log(b+x)}{a-b}$$

input `integrate(1/(a+x)/(b+x),x, algorithm="maxima")`

output `-log(a + x)/(a - b) + log(b + x)/(a - b)`

3.181.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

$$\int \frac{1}{(a+x)(b+x)} dx = -\frac{\log(|a+x|)}{a-b} + \frac{\log(|b+x|)}{a-b}$$

input `integrate(1/(a+x)/(b+x),x, algorithm="giac")`

output `-log(abs(a + x))/(a - b) + log(abs(b + x))/(a - b)`

3.181.9 Mupad [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.69

$$\int \frac{1}{(a+x)(b+x)} dx = \frac{\ln\left(\frac{b+x}{a+x}\right)}{a-b}$$

input `int(1/((a + x)*(b + x)),x)`

output `log((b + x)/(a + x))/(a - b)`

3.182 $\int \frac{1+x^2}{-x+x^2} dx$

3.182.1 Optimal result	965
3.182.2 Mathematica [A] (verified)	965
3.182.3 Rubi [A] (verified)	966
3.182.4 Maple [A] (verified)	967
3.182.5 Fricas [A] (verification not implemented)	967
3.182.6 Sympy [A] (verification not implemented)	967
3.182.7 Maxima [A] (verification not implemented)	968
3.182.8 Giac [A] (verification not implemented)	968
3.182.9 Mupad [B] (verification not implemented)	968

3.182.1 Optimal result

Integrand size = 15, antiderivative size = 14

$$\int \frac{1+x^2}{-x+x^2} dx = x + 2\log(1-x) - \log(x)$$

output `x+2*ln(1-x)-ln(x)`

3.182.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1+x^2}{-x+x^2} dx = x + 2\log(1-x) - \log(x)$$

input `Integrate[(1 + x^2)/(-x + x^2),x]`

output `x + 2*Log[1 - x] - Log[x]`

3.182.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2026, 522, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^2 + 1}{x^2 - x} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{x^2 + 1}{(x - 1)x} dx \\ & \quad \downarrow \text{522} \\ & \int \left(-\frac{1}{x} + \frac{2}{x - 1} + 1 \right) dx \\ & \quad \downarrow \text{2009} \\ & x + 2 \log(1 - x) - \log(x) \end{aligned}$$

input `Int[(1 + x^2)/(-x + x^2),x]`

output `x + 2*Log[1 - x] - Log[x]`

3.182.3.1 Defintions of rubi rules used

rule 522 `Int[((e_.)*(x_)^(m_.)*((c_) + (d_.)*(x_)^(n_.))*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(e*x)^m*(c + d*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

3.182.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
default	$x + 2 \ln(-1 + x) - \ln(x)$	13
norman	$x + 2 \ln(-1 + x) - \ln(x)$	13
risch	$x + 2 \ln(-1 + x) - \ln(x)$	13
parallelrisc	$x + 2 \ln(-1 + x) - \ln(x)$	13
meijerg	$-\ln(x) - i\pi + 2 \ln(1 - x) + x$	19

input `int((x^2+1)/(x^2-x),x,method=_RETURNVERBOSE)`output `x+2*ln(-1+x)-ln(x)`**3.182.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1+x^2}{-x+x^2} dx = x + 2 \log(x-1) - \log(x)$$

input `integrate((x^2+1)/(x^2-x),x, algorithm="fracas")`output `x + 2*log(x - 1) - log(x)`**3.182.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{1+x^2}{-x+x^2} dx = x - \log(x) + 2 \log(x-1)$$

input `integrate((x**2+1)/(x**2-x),x)`output `x - log(x) + 2*log(x - 1)`

3.182.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1+x^2}{-x+x^2} dx = x + 2 \log(x-1) - \log(x)$$

input `integrate((x^2+1)/(x^2-x),x, algorithm="maxima")`output `x + 2*log(x - 1) - log(x)`**3.182.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1+x^2}{-x+x^2} dx = x + 2 \log(|x-1|) - \log(|x|)$$

input `integrate((x^2+1)/(x^2-x),x, algorithm="giac")`output `x + 2*log(abs(x - 1)) - log(abs(x))`**3.182.9 Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1+x^2}{-x+x^2} dx = x + 2 \ln(x-1) - \ln(x)$$

input `int(-(x^2 + 1)/(x - x^2),x)`output `x + 2*log(x - 1) - log(x)`

$$3.183 \quad \int \frac{1-12x+x^2+x^3}{-12+x+x^2} dx$$

3.183.1 Optimal result	969
3.183.2 Mathematica [A] (verified)	969
3.183.3 Rubi [A] (verified)	970
3.183.4 Maple [A] (verified)	971
3.183.5 Fricas [A] (verification not implemented)	971
3.183.6 Sympy [A] (verification not implemented)	971
3.183.7 Maxima [A] (verification not implemented)	972
3.183.8 Giac [A] (verification not implemented)	972
3.183.9 Mupad [B] (verification not implemented)	972

3.183.1 Optimal result

Integrand size = 20, antiderivative size = 26

$$\int \frac{1-12x+x^2+x^3}{-12+x+x^2} dx = \frac{x^2}{2} + \frac{1}{7} \log(3-x) - \frac{1}{7} \log(4+x)$$

output `1/2*x^2+1/7*ln(3-x)-1/7*ln(4+x)`

3.183.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{1-12x+x^2+x^3}{-12+x+x^2} dx = \frac{x^2}{2} + \frac{1}{7} \log(3-x) - \frac{1}{7} \log(4+x)$$

input `Integrate[(1 - 12*x + x^2 + x^3)/(-12 + x + x^2),x]`

output `x^2/2 + Log[3 - x]/7 - Log[4 + x]/7`

3.183.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2188, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 + x^2 - 12x + 1}{x^2 + x - 12} dx$$

↓ 2188

$$\int \left(\frac{1}{x^2 + x - 12} + x \right) dx$$

↓ 2009

$$\frac{x^2}{2} + \frac{1}{7} \log(3 - x) - \frac{1}{7} \log(x + 4)$$

input `Int[(1 - 12*x + x^2 + x^3)/(-12 + x + x^2),x]`

output `x^2/2 + Log[3 - x]/7 - Log[4 + x]/7`

3.183.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2188 `Int[(Pq_)*((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] :> Int[Expand Integrand[Pq*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

3.183.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.73

method	result	size
default	$\frac{x^2}{2} - \frac{\ln(4+x)}{7} + \frac{\ln(-3+x)}{7}$	19
norman	$\frac{x^2}{2} - \frac{\ln(4+x)}{7} + \frac{\ln(-3+x)}{7}$	19
risch	$\frac{x^2}{2} - \frac{\ln(4+x)}{7} + \frac{\ln(-3+x)}{7}$	19
parallelrisch	$\frac{x^2}{2} - \frac{\ln(4+x)}{7} + \frac{\ln(-3+x)}{7}$	19

input `int((x^3+x^2-12*x+1)/(x^2+x-12),x,method=_RETURNVERBOSE)`output `1/2*x^2-1/7*ln(4+x)+1/7*ln(-3+x)`**3.183.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.69

$$\int \frac{1 - 12x + x^2 + x^3}{-12 + x + x^2} dx = \frac{1}{2} x^2 - \frac{1}{7} \log(x + 4) + \frac{1}{7} \log(x - 3)$$

input `integrate((x^3+x^2-12*x+1)/(x^2+x-12),x, algorithm="fricas")`output `1/2*x^2 - 1/7*log(x + 4) + 1/7*log(x - 3)`**3.183.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.65

$$\int \frac{1 - 12x + x^2 + x^3}{-12 + x + x^2} dx = \frac{x^2}{2} + \frac{\log(x - 3)}{7} - \frac{\log(x + 4)}{7}$$

input `integrate((x**3+x**2-12*x+1)/(x**2+x-12),x)`output `x**2/2 + log(x - 3)/7 - log(x + 4)/7`

3.183. $\int \frac{1-12x+x^2+x^3}{-12+x+x^2} dx$

3.183.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.69

$$\int \frac{1 - 12x + x^2 + x^3}{-12 + x + x^2} dx = \frac{1}{2} x^2 - \frac{1}{7} \log(x + 4) + \frac{1}{7} \log(x - 3)$$

input `integrate((x^3+x^2-12*x+1)/(x^2+x-12),x, algorithm="maxima")`output `1/2*x^2 - 1/7*log(x + 4) + 1/7*log(x - 3)`**3.183.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \frac{1 - 12x + x^2 + x^3}{-12 + x + x^2} dx = \frac{1}{2} x^2 - \frac{1}{7} \log(|x + 4|) + \frac{1}{7} \log(|x - 3|)$$

input `integrate((x^3+x^2-12*x+1)/(x^2+x-12),x, algorithm="giac")`output `1/2*x^2 - 1/7*log(abs(x + 4)) + 1/7*log(abs(x - 3))`**3.183.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.54

$$\int \frac{1 - 12x + x^2 + x^3}{-12 + x + x^2} dx = \frac{x^2}{2} - \frac{2 \operatorname{atanh}\left(\frac{2x}{7} + \frac{1}{7}\right)}{7}$$

input `int((x^2 - 12*x + x^3 + 1)/(x + x^2 - 12),x)`output `x^2/2 - (2*atanh((2*x)/7 + 1/7))/7`

3.184 $\int \frac{3+2x}{(1+x)^2} dx$

3.184.1 Optimal result	973
3.184.2 Mathematica [A] (verified)	973
3.184.3 Rubi [A] (verified)	974
3.184.4 Maple [A] (verified)	975
3.184.5 Fricas [A] (verification not implemented)	975
3.184.6 Sympy [A] (verification not implemented)	975
3.184.7 Maxima [A] (verification not implemented)	976
3.184.8 Giac [A] (verification not implemented)	976
3.184.9 Mupad [B] (verification not implemented)	976

3.184.1 Optimal result

Integrand size = 11, antiderivative size = 14

$$\int \frac{3 + 2x}{(1 + x)^2} dx = -\frac{1}{1 + x} + 2 \log(1 + x)$$

output `-1/(1+x)+2*ln(1+x)`

3.184.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{3 + 2x}{(1 + x)^2} dx = -\frac{1}{1 + x} + 2 \log(1 + x)$$

input `Integrate[(3 + 2*x)/(1 + x)^2,x]`

output `-(1 + x)^(-1) + 2*Log[1 + x]`

3.184.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x+3}{(x+1)^2} dx$$

↓ 49

$$\int \left(\frac{2}{x+1} + \frac{1}{(x+1)^2} \right) dx$$

↓ 2009

$$2 \log(x+1) - \frac{1}{x+1}$$

input `Int[(3 + 2*x)/(1 + x)^2,x]`

output `-(1 + x)^(-1) + 2*Log[1 + x]`

3.184.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.184.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

method	result	size
default	$-\frac{1}{1+x} + 2 \ln(1+x)$	15
norman	$-\frac{1}{1+x} + 2 \ln(1+x)$	15
meijerg	$\frac{x}{1+x} + 2 \ln(1+x)$	15
risch	$-\frac{1}{1+x} + 2 \ln(1+x)$	15
parallelrisch	$\frac{2 \ln(1+x)x - 1 + 2 \ln(1+x)}{1+x}$	22

input `int((3+2*x)/(1+x)^2,x,method=_RETURNVERBOSE)`output `-1/(1+x)+2*ln(1+x)`**3.184.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.21

$$\int \frac{3+2x}{(1+x)^2} dx = \frac{2(x+1)\log(x+1) - 1}{x+1}$$

input `integrate((3+2*x)/(1+x)^2,x, algorithm="fracas")`output `(2*(x + 1)*log(x + 1) - 1)/(x + 1)`**3.184.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{3+2x}{(1+x)^2} dx = 2 \log(x+1) - \frac{1}{x+1}$$

input `integrate((3+2*x)/(1+x)**2,x)`output `2*log(x + 1) - 1/(x + 1)`

3.184.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{3+2x}{(1+x)^2} dx = -\frac{1}{x+1} + 2 \log(x+1)$$

input `integrate((3+2*x)/(1+x)^2,x, algorithm="maxima")`output `-1/(x + 1) + 2*log(x + 1)`**3.184.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

$$\int \frac{3+2x}{(1+x)^2} dx = -\frac{1}{x+1} + 2 \log(|x+1|)$$

input `integrate((3+2*x)/(1+x)^2,x, algorithm="giac")`output `-1/(x + 1) + 2*log(abs(x + 1))`**3.184.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{3+2x}{(1+x)^2} dx = 2 \ln(x+1) - \frac{1}{x+1}$$

input `int((2*x + 3)/(x + 1)^2,x)`output `2*log(x + 1) - 1/(x + 1)`

3.185 $\int \frac{1}{x(1+x)(3+2x)} dx$

3.185.1 Optimal result	977
3.185.2 Mathematica [A] (verified)	977
3.185.3 Rubi [A] (verified)	978
3.185.4 Maple [A] (verified)	979
3.185.5 Fricas [A] (verification not implemented)	979
3.185.6 Sympy [A] (verification not implemented)	979
3.185.7 Maxima [A] (verification not implemented)	980
3.185.8 Giac [A] (verification not implemented)	980
3.185.9 Mupad [B] (verification not implemented)	980

3.185.1 Optimal result

Integrand size = 16, antiderivative size = 23

$$\int \frac{1}{x(1+x)(3+2x)} dx = \frac{\log(x)}{3} - \log(1+x) + \frac{2}{3} \log(3+2x)$$

output `1/3*ln(x)-ln(1+x)+2/3*ln(3+2*x)`

3.185.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+x)(3+2x)} dx = \frac{\log(x)}{3} - \log(1+x) + \frac{2}{3} \log(3+2x)$$

input `Integrate[1/(x*(1+x)*(3+2*x)),x]`

output `Log[x]/3 - Log[1+x] + (2*Log[3+2*x])/3`

3.185.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {93, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x(x+1)(2x+3)} dx$$

↓ 93

$$\int \left(\frac{1}{3x} + \frac{4}{3(2x+3)} + \frac{1}{-x-1} \right) dx$$

↓ 2009

$$\frac{\log(x)}{3} - \log(x+1) + \frac{2}{3} \log(2x+3)$$

input `Int[1/(x*(1 + x)*(3 + 2*x)),x]`

output `Log[x]/3 - Log[1 + x] + (2*Log[3 + 2*x])/3`

3.185.3.1 Defintions of rubi rules used

rule 93 `Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_] := Int[ExpandIntegrand[(e + f*x)^p/((a + b*x)*(c + d*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IntegerQ[p]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.185.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

method	result	size
parallelsch	$\frac{\ln(x)}{3} - \ln(1+x) + \frac{2\ln(\frac{3}{2}+x)}{3}$	18
default	$\frac{\ln(x)}{3} - \ln(1+x) + \frac{2\ln(3+2x)}{3}$	20
norman	$\frac{\ln(x)}{3} - \ln(1+x) + \frac{2\ln(3+2x)}{3}$	20
risch	$\frac{\ln(x)}{3} - \ln(1+x) + \frac{2\ln(3+2x)}{3}$	20

input `int(1/x/(1+x)/(3+2*x),x,method=_RETURNVERBOSE)`output `1/3*ln(x)-ln(1+x)+2/3*ln(3/2+x)`**3.185.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{1}{x(1+x)(3+2x)} dx = \frac{2}{3} \log(2x+3) - \log(x+1) + \frac{1}{3} \log(x)$$

input `integrate(1/x/(1+x)/(3+2*x),x, algorithm="fracas")`output `2/3*log(2*x + 3) - log(x + 1) + 1/3*log(x)`**3.185.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{1}{x(1+x)(3+2x)} dx = \frac{\log(x)}{3} - \log(x+1) + \frac{2\log(x+\frac{3}{2})}{3}$$

input `integrate(1/x/(1+x)/(3+2*x),x)`output `log(x)/3 - log(x + 1) + 2*log(x + 3/2)/3`

3.185.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{1}{x(1+x)(3+2x)} dx = \frac{2}{3} \log(2x+3) - \log(x+1) + \frac{1}{3} \log(x)$$

input `integrate(1/x/(1+x)/(3+2*x),x, algorithm="maxima")`output `2/3*log(2*x + 3) - log(x + 1) + 1/3*log(x)`**3.185.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int \frac{1}{x(1+x)(3+2x)} dx = \frac{2}{3} \log(|2x+3|) - \log(|x+1|) + \frac{1}{3} \log(|x|)$$

input `integrate(1/x/(1+x)/(3+2*x),x, algorithm="giac")`output `2/3*log(abs(2*x + 3)) - log(abs(x + 1)) + 1/3*log(abs(x))`**3.185.9 Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{1}{x(1+x)(3+2x)} dx = \frac{2 \ln\left(x + \frac{3}{2}\right)}{3} - \ln(x+1) + \frac{\ln(x)}{3}$$

input `int(1/(x*(2*x + 3)*(x + 1)),x)`output `(2*log(x + 3/2))/3 - log(x + 1) + log(x)/3`

$$3.186 \quad \int \frac{-3+5x+6x^2}{-3x+2x^2+x^3} dx$$

3.186.1 Optimal result	981
3.186.2 Mathematica [A] (verified)	981
3.186.3 Rubi [A] (verified)	982
3.186.4 Maple [A] (verified)	983
3.186.5 Fricas [A] (verification not implemented)	983
3.186.6 Sympy [A] (verification not implemented)	983
3.186.7 Maxima [A] (verification not implemented)	984
3.186.8 Giac [A] (verification not implemented)	984
3.186.9 Mupad [B] (verification not implemented)	984

3.186.1 Optimal result

Integrand size = 25, antiderivative size = 17

$$\int \frac{-3 + 5x + 6x^2}{-3x + 2x^2 + x^3} dx = 2 \log(1 - x) + \log(x) + 3 \log(3 + x)$$

output `2*ln(1-x)+ln(x)+3*ln(3+x)`

3.186.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{-3 + 5x + 6x^2}{-3x + 2x^2 + x^3} dx = 2 \log(1 - x) + \log(x) + 3 \log(3 + x)$$

input `Integrate[(-3 + 5*x + 6*x^2)/(-3*x + 2*x^2 + x^3),x]`

output `2*Log[1 - x] + Log[x] + 3*Log[3 + x]`

3.186.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2026, 2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{6x^2 + 5x - 3}{x^3 + 2x^2 - 3x} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{6x^2 + 5x - 3}{x(x^2 + 2x - 3)} dx \\ & \quad \downarrow \text{2159} \\ & \int \left(\frac{1}{x} + \frac{3}{x+3} + \frac{2}{x-1} \right) dx \\ & \quad \downarrow \text{2009} \\ & 2 \log(1-x) + \log(x) + 3 \log(x+3) \end{aligned}$$

input `Int[(-3 + 5*x + 6*x^2)/(-3*x + 2*x^2 + x^3),x]`

output `2*Log[1 - x] + Log[x] + 3*Log[3 + x]`

3.186.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])]`

rule 2159 `Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

3.186.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

method	result	size
default	$2 \ln(-1+x) + \ln(x) + 3 \ln(3+x)$	16
norman	$2 \ln(-1+x) + \ln(x) + 3 \ln(3+x)$	16
risch	$2 \ln(-1+x) + \ln(x) + 3 \ln(3+x)$	16
parallelrisch	$2 \ln(-1+x) + \ln(x) + 3 \ln(3+x)$	16

input `int((6*x^2+5*x-3)/(x^3+2*x^2-3*x),x,method=_RETURNVERBOSE)`output `2*ln(-1+x)+ln(x)+3*ln(3+x)`**3.186.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{-3 + 5x + 6x^2}{-3x + 2x^2 + x^3} dx = 3 \log(x + 3) + 2 \log(x - 1) + \log(x)$$

input `integrate((6*x^2+5*x-3)/(x^3+2*x^2-3*x),x, algorithm="fricas")`output `3*log(x + 3) + 2*log(x - 1) + log(x)`**3.186.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{-3 + 5x + 6x^2}{-3x + 2x^2 + x^3} dx = \log(x) + 2 \log(x - 1) + 3 \log(x + 3)$$

input `integrate((6*x**2+5*x-3)/(x**3+2*x**2-3*x),x)`output `log(x) + 2*log(x - 1) + 3*log(x + 3)`

3.186.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{-3 + 5x + 6x^2}{-3x + 2x^2 + x^3} dx = 3 \log(x + 3) + 2 \log(x - 1) + \log(x)$$

input `integrate((6*x^2+5*x-3)/(x^3+2*x^2-3*x),x, algorithm="maxima")`output `3*log(x + 3) + 2*log(x - 1) + log(x)`**3.186.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

$$\int \frac{-3 + 5x + 6x^2}{-3x + 2x^2 + x^3} dx = 3 \log(|x + 3|) + 2 \log(|x - 1|) + \log(|x|)$$

input `integrate((6*x^2+5*x-3)/(x^3+2*x^2-3*x),x, algorithm="giac")`output `3*log(abs(x + 3)) + 2*log(abs(x - 1)) + log(abs(x))`**3.186.9 Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{-3 + 5x + 6x^2}{-3x + 2x^2 + x^3} dx = 2 \ln(x - 1) + 3 \ln(x + 3) + \ln(x)$$

input `int((5*x + 6*x^2 - 3)/(2*x^2 - 3*x + x^3),x)`output `2*log(x - 1) + 3*log(x + 3) + log(x)`

3.187 $\int \frac{x}{4+4x+x^2} dx$

3.187.1 Optimal result	985
3.187.2 Mathematica [A] (verified)	985
3.187.3 Rubi [A] (verified)	986
3.187.4 Maple [A] (verified)	987
3.187.5 Fricas [A] (verification not implemented)	987
3.187.6 Sympy [A] (verification not implemented)	987
3.187.7 Maxima [A] (verification not implemented)	988
3.187.8 Giac [A] (verification not implemented)	988
3.187.9 Mupad [B] (verification not implemented)	988

3.187.1 Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{x}{4+4x+x^2} dx = \frac{2}{2+x} + \log(2+x)$$

output `2/(2+x)+ln(2+x)`

3.187.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{x}{4+4x+x^2} dx = \frac{2}{2+x} + \log(2+x)$$

input `Integrate[x/(4 + 4*x + x^2),x]`

output `2/(2 + x) + Log[2 + x]`

3.187.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {1098, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x}{x^2 + 4x + 4} dx \\ & \quad \downarrow \text{1098} \\ & \int \frac{x}{(x+2)^2} dx \\ & \quad \downarrow \text{49} \\ & \int \left(\frac{1}{x+2} - \frac{2}{(x+2)^2} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{2}{x+2} + \log(x+2) \end{aligned}$$

input `Int[x/(4 + 4*x + x^2),x]`

output `2/(2 + x) + Log[2 + x]`

3.187.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 1098 `Int[((d_.) + (e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[1/c^p Int[(d + e*x)^m*(b/2 + c*x)^(2*p), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.187.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

method	result	size
default	$\frac{2}{2+x} + \ln(2+x)$	13
norman	$\frac{2}{2+x} + \ln(2+x)$	13
risch	$\frac{2}{2+x} + \ln(2+x)$	13
meijerg	$-\frac{x}{2(1+\frac{x}{2})} + \ln(1+\frac{x}{2})$	18
parallelrisch	$\frac{\ln(2+x)x+2+2\ln(2+x)}{2+x}$	21

input `int(x/(x^2+4*x+4),x,method=_RETURNVERBOSE)`output `2/(2+x)+ln(2+x)`**3.187.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.33

$$\int \frac{x}{4+4x+x^2} dx = \frac{(x+2)\log(x+2)+2}{x+2}$$

input `integrate(x/(x^2+4*x+4),x, algorithm="fricas")`output `((x+2)*log(x+2)+2)/(x+2)`**3.187.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x}{4+4x+x^2} dx = \log(x+2) + \frac{2}{x+2}$$

input `integrate(x/(x**2+4*x+4),x)`output `log(x+2)+2/(x+2)`

3.187.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{x}{4 + 4x + x^2} dx = \frac{2}{x + 2} + \log(x + 2)$$

input `integrate(x/(x^2+4*x+4),x, algorithm="maxima")`output `2/(x + 2) + log(x + 2)`**3.187.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

$$\int \frac{x}{4 + 4x + x^2} dx = \frac{2}{x + 2} + \log(|x + 2|)$$

input `integrate(x/(x^2+4*x+4),x, algorithm="giac")`output `2/(x + 2) + log(abs(x + 2))`**3.187.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{x}{4 + 4x + x^2} dx = \ln(x + 2) + \frac{2}{x + 2}$$

input `int(x/(4*x + x^2 + 4),x)`output `log(x + 2) + 2/(x + 2)`

$$3.188 \quad \int \frac{1}{(-1+x)^2(4+x)} dx$$

3.188.1 Optimal result	989
3.188.2 Mathematica [A] (verified)	989
3.188.3 Rubi [A] (verified)	990
3.188.4 Maple [A] (verified)	991
3.188.5 Fricas [A] (verification not implemented)	991
3.188.6 Sympy [A] (verification not implemented)	991
3.188.7 Maxima [A] (verification not implemented)	992
3.188.8 Giac [A] (verification not implemented)	992
3.188.9 Mupad [B] (verification not implemented)	992

3.188.1 Optimal result

Integrand size = 11, antiderivative size = 30

$$\int \frac{1}{(-1+x)^2(4+x)} dx = \frac{1}{5(1-x)} - \frac{1}{25} \log(1-x) + \frac{1}{25} \log(4+x)$$

output `1/5/(1-x)-1/25*ln(1-x)+1/25*ln(4+x)`

3.188.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \frac{1}{(-1+x)^2(4+x)} dx = \frac{1}{25} \left(-\frac{5}{-1+x} - \log(-1+x) + \log(4+x) \right)$$

input `Integrate[1/((-1 + x)^2*(4 + x)),x]`

output `(-5/(-1 + x) - Log[-1 + x] + Log[4 + x])/25`

3.188.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x-1)^2(x+4)} dx$$

↓ 54

$$\int \left(\frac{1}{25(x+4)} - \frac{1}{25(x-1)} + \frac{1}{5(x-1)^2} \right) dx$$

↓ 2009

$$\frac{1}{5(1-x)} - \frac{1}{25} \log(1-x) + \frac{1}{25} \log(x+4)$$

input `Int[1/((-1 + x)^2*(4 + x)),x]`

output `1/(5*(1 - x)) - Log[1 - x]/25 + Log[4 + x]/25`

3.188.3.1 Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.188.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.70

method	result	size
default	$-\frac{1}{5(-1+x)} - \frac{\ln(-1+x)}{25} + \frac{\ln(4+x)}{25}$	21
norman	$-\frac{1}{5(-1+x)} - \frac{\ln(-1+x)}{25} + \frac{\ln(4+x)}{25}$	21
risch	$-\frac{1}{5(-1+x)} - \frac{\ln(-1+x)}{25} + \frac{\ln(4+x)}{25}$	21
parallelrisch	$-\frac{\ln(-1+x)x - \ln(4+x)x + 5 - \ln(-1+x) + \ln(4+x)}{25(-1+x)}$	33

input `int(1/(-1+x)^2/(4+x),x,method=_RETURNVERBOSE)`output `-1/5/(-1+x)-1/25*ln(-1+x)+1/25*ln(4+x)`**3.188.5 Fricas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.87

$$\int \frac{1}{(-1+x)^2(4+x)} dx = \frac{(x-1)\log(x+4) - (x-1)\log(x-1) - 5}{25(x-1)}$$

input `integrate(1/(-1+x)^2/(4+x),x, algorithm="fricas")`output `1/25*((x - 1)*log(x + 4) - (x - 1)*log(x - 1) - 5)/(x - 1)`**3.188.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.63

$$\int \frac{1}{(-1+x)^2(4+x)} dx = -\frac{\log(x-1)}{25} + \frac{\log(x+4)}{25} - \frac{1}{5x-5}$$

input `integrate(1/(-1+x)**2/(4+x),x)`output `-log(x - 1)/25 + log(x + 4)/25 - 1/(5*x - 5)`

3.188.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.67

$$\int \frac{1}{(-1+x)^2(4+x)} dx = -\frac{1}{5(x-1)} + \frac{1}{25} \log(x+4) - \frac{1}{25} \log(x-1)$$

input `integrate(1/(-1+x)^2/(4+x),x, algorithm="maxima")`output `-1/5/(x - 1) + 1/25*log(x + 4) - 1/25*log(x - 1)`**3.188.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.70

$$\int \frac{1}{(-1+x)^2(4+x)} dx = -\frac{1}{5(x-1)} + \frac{1}{25} \log\left(\left|-\frac{5}{x-1} - 1\right|\right)$$

input `integrate(1/(-1+x)^2/(4+x),x, algorithm="giac")`output `-1/5/(x - 1) + 1/25*log(abs(-5/(x - 1) - 1))`**3.188.9 Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \frac{1}{(-1+x)^2(4+x)} dx = -\frac{\ln\left(\frac{x-1}{x+4}\right)}{25} - \frac{1}{5(x-1)}$$

input `int(1/((x - 1)^2*(x + 4)),x)`output `- log((x - 1)/(x + 4))/25 - 1/(5*(x - 1))`

$$3.189 \quad \int \frac{x^2}{(-3+x)(2+x)^2} dx$$

3.189.1 Optimal result	993
3.189.2 Mathematica [A] (verified)	993
3.189.3 Rubi [A] (verified)	994
3.189.4 Maple [A] (verified)	995
3.189.5 Fricas [A] (verification not implemented)	995
3.189.6 Sympy [A] (verification not implemented)	995
3.189.7 Maxima [A] (verification not implemented)	996
3.189.8 Giac [A] (verification not implemented)	996
3.189.9 Mupad [B] (verification not implemented)	996

3.189.1 Optimal result

Integrand size = 14, antiderivative size = 28

$$\int \frac{x^2}{(-3+x)(2+x)^2} dx = \frac{4}{5(2+x)} + \frac{9}{25} \log(3-x) + \frac{16}{25} \log(2+x)$$

output `4/5/(2+x)+9/25*ln(3-x)+16/25*ln(2+x)`

3.189.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{x^2}{(-3+x)(2+x)^2} dx = \frac{4}{5(2+x)} + \frac{9}{25} \log(-3+x) + \frac{16}{25} \log(2+x)$$

input `Integrate[x^2/((-3 + x)*(2 + x)^2),x]`

output `4/(5*(2 + x)) + (9*Log[-3 + x])/25 + (16*Log[2 + x])/25`

3.189.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {99, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{(x-3)(x+2)^2} dx$$

↓ 99

$$\int \left(\frac{16}{25(x+2)} - \frac{4}{5(x+2)^2} + \frac{9}{25(x-3)} \right) dx$$

↓ 2009

$$\frac{4}{5(x+2)} + \frac{9}{25} \log(3-x) + \frac{16}{25} \log(x+2)$$

input `Int[x^2/((-3 + x)*(2 + x)^2),x]`

output `4/(5*(2 + x)) + (9*Log[3 - x])/25 + (16*Log[2 + x])/25`

3.189.3.1 Defintions of rubi rules used

rule 99 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] | (GtQ[m, 0] && GeQ[n, -1]))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.189.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.75

method	result	size
default	$\frac{4}{5(2+x)} + \frac{16 \ln(2+x)}{25} + \frac{9 \ln(-3+x)}{25}$	21
norman	$\frac{4}{5(2+x)} + \frac{16 \ln(2+x)}{25} + \frac{9 \ln(-3+x)}{25}$	21
risch	$\frac{4}{5(2+x)} + \frac{16 \ln(2+x)}{25} + \frac{9 \ln(-3+x)}{25}$	21
parallelrisc	$\frac{9 \ln(-3+x)x + 16 \ln(2+x)x + 20 + 18 \ln(-3+x) + 32 \ln(2+x)}{50 + 25x}$	36

input `int(x^2/(-3+x)/(2+x)^2,x,method=_RETURNVERBOSE)`output `4/5/(2+x)+16/25*ln(2+x)+9/25*ln(-3+x)`**3.189.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{x^2}{(-3+x)(2+x)^2} dx = \frac{16(x+2) \log(x+2) + 9(x+2) \log(x-3) + 20}{25(x+2)}$$

input `integrate(x^2/(-3+x)/(2+x)^2,x, algorithm="fricas")`output `1/25*(16*(x+2)*log(x+2)+9*(x+2)*log(x-3)+20)/(x+2)`**3.189.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int \frac{x^2}{(-3+x)(2+x)^2} dx = \frac{9 \log(x-3)}{25} + \frac{16 \log(x+2)}{25} + \frac{4}{5x+10}$$

input `integrate(x**2/(-3+x)/(2+x)**2,x)`output `9*log(x-3)/25+16*log(x+2)/25+4/(5*x+10)`

3.189. $\int \frac{x^2}{(-3+x)(2+x)^2} dx$

3.189.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.71

$$\int \frac{x^2}{(-3+x)(2+x)^2} dx = \frac{4}{5(x+2)} + \frac{16}{25} \log(x+2) + \frac{9}{25} \log(x-3)$$

input `integrate(x^2/(-3+x)/(2+x)^2,x, algorithm="maxima")`output `4/5/(x + 2) + 16/25*log(x + 2) + 9/25*log(x - 3)`**3.189.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{x^2}{(-3+x)(2+x)^2} dx = \frac{4}{5(x+2)} + \log(|x+2|) + \frac{9}{25} \log\left(\left|-\frac{5}{x+2} + 1\right|\right)$$

input `integrate(x^2/(-3+x)/(2+x)^2,x, algorithm="giac")`output `4/5/(x + 2) + log(abs(x + 2)) + 9/25*log(abs(-5/(x + 2) + 1))`**3.189.9 Mupad [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int \frac{x^2}{(-3+x)(2+x)^2} dx = \frac{16 \ln(x+2)}{25} + \frac{9 \ln(x-3)}{25} + \frac{4}{5(x+2)}$$

input `int(x^2/((x + 2)^2*(x - 3)),x)`output `(16*log(x + 2))/25 + (9*log(x - 3))/25 + 4/(5*(x + 2))`

$$3.190 \quad \int \frac{-2+3x+5x^2}{2x^2+x^3} dx$$

3.190.1 Optimal result	997
3.190.2 Mathematica [A] (verified)	997
3.190.3 Rubi [A] (verified)	998
3.190.4 Maple [A] (verified)	999
3.190.5 Fricas [A] (verification not implemented)	999
3.190.6 Sympy [A] (verification not implemented)	1000
3.190.7 Maxima [A] (verification not implemented)	1000
3.190.8 Giac [A] (verification not implemented)	1000
3.190.9 Mupad [B] (verification not implemented)	1001

3.190.1 Optimal result

Integrand size = 22, antiderivative size = 14

$$\int \frac{-2+3x+5x^2}{2x^2+x^3} dx = \frac{1}{x} + 2 \log(x) + 3 \log(2+x)$$

output `1/x+2*ln(x)+3*ln(2+x)`

3.190.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{-2+3x+5x^2}{2x^2+x^3} dx = \frac{1}{x} + 2 \log(x) + 3 \log(2+x)$$

input `Integrate[(-2 + 3*x + 5*x^2)/(2*x^2 + x^3),x]`

output `x^(-1) + 2*Log[x] + 3*Log[2 + x]`

3.190.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$, Rules used = {2026, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{5x^2 + 3x - 2}{x^3 + 2x^2} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{5x^2 + 3x - 2}{x^2(x+2)} dx \\ & \quad \downarrow \text{1195} \\ & \int \left(-\frac{1}{x^2} + \frac{3}{x+2} + \frac{2}{x} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{1}{x} + 2 \log(x) + 3 \log(x+2) \end{aligned}$$

input `Int[(-2 + 3*x + 5*x^2)/(2*x^2 + x^3), x]`

output `x^(-1) + 2*Log[x] + 3*Log[2 + x]`

3.190.3.1 Defintions of rubi rules used

rule 1195 `Int[((d_.) + (e_.)*(x_)^(m_.))*((f_.) + (g_.)*(x_)^(n_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

3.190.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

method	result	size
default	$\frac{1}{x} + 2 \ln(x) + 3 \ln(2 + x)$	15
norman	$\frac{1}{x} + 2 \ln(x) + 3 \ln(2 + x)$	15
risch	$\frac{1}{x} + 2 \ln(x) + 3 \ln(2 + x)$	15
parallelrisch	$\frac{2x \ln(x) + 3 \ln(2+x)x + 1}{x}$	19
meijerg	$\frac{1}{x} + 2 \ln(x) - 2 \ln(2) + 3 \ln\left(1 + \frac{x}{2}\right)$	21

input `int((5*x^2+3*x-2)/(x^3+2*x^2),x,method=_RETURNVERBOSE)`

output `1/x+2*ln(x)+3*ln(2+x)`

3.190.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.29

$$\int \frac{-2 + 3x + 5x^2}{2x^2 + x^3} dx = \frac{3x \log(x + 2) + 2x \log(x) + 1}{x}$$

input `integrate((5*x^2+3*x-2)/(x^3+2*x^2),x, algorithm="fricas")`

output `(3*x*log(x + 2) + 2*x*log(x) + 1)/x`

3.190.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{-2 + 3x + 5x^2}{2x^2 + x^3} dx = 2 \log(x) + 3 \log(x + 2) + \frac{1}{x}$$

input `integrate((5*x**2+3*x-2)/(x**3+2*x**2),x)`output `2*log(x) + 3*log(x + 2) + 1/x`**3.190.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{-2 + 3x + 5x^2}{2x^2 + x^3} dx = \frac{1}{x} + 3 \log(x + 2) + 2 \log(x)$$

input `integrate((5*x^2+3*x-2)/(x^3+2*x^2),x, algorithm="maxima")`output `1/x + 3*log(x + 2) + 2*log(x)`**3.190.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{-2 + 3x + 5x^2}{2x^2 + x^3} dx = \frac{1}{x} + 3 \log(|x + 2|) + 2 \log(|x|)$$

input `integrate((5*x^2+3*x-2)/(x^3+2*x^2),x, algorithm="giac")`output `1/x + 3*log(abs(x + 2)) + 2*log(abs(x))`

3.190.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{-2 + 3x + 5x^2}{2x^2 + x^3} dx = 3 \ln(x + 2) + 2 \ln(x) + \frac{1}{x}$$

input `int((3*x + 5*x^2 - 2)/(2*x^2 + x^3),x)`

output `3*log(x + 2) + 2*log(x) + 1/x`

$$3.191 \quad \int \frac{18-2x-4x^2}{-6+x+4x^2+x^3} dx$$

3.191.1 Optimal result	1002
3.191.2 Mathematica [A] (verified)	1002
3.191.3 Rubi [A] (verified)	1003
3.191.4 Maple [A] (verified)	1004
3.191.5 Fricas [A] (verification not implemented)	1004
3.191.6 Sympy [A] (verification not implemented)	1004
3.191.7 Maxima [A] (verification not implemented)	1005
3.191.8 Giac [A] (verification not implemented)	1005
3.191.9 Mupad [B] (verification not implemented)	1005

3.191.1 Optimal result

Integrand size = 24, antiderivative size = 19

$$\int \frac{18 - 2x - 4x^2}{-6 + x + 4x^2 + x^3} dx = \log(1 - x) - 2\log(2 + x) - 3\log(3 + x)$$

output `ln(1-x)-2*ln(2+x)-3*ln(3+x)`

3.191.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.32

$$\int \frac{18 - 2x - 4x^2}{-6 + x + 4x^2 + x^3} dx = -2 \left(-\frac{1}{2} \log(1 - x) + \log(2 + x) + \frac{3}{2} \log(3 + x) \right)$$

input `Integrate[(18 - 2*x - 4*x^2)/(-6 + x + 4*x^2 + x^3),x]`

output `-2*(-1/2*Log[1 - x] + Log[2 + x] + (3*Log[3 + x])/2)`

3.191.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{-4x^2 - 2x + 18}{x^3 + 4x^2 + x - 6} dx$$

↓ 2462

$$\int \left(-\frac{2}{x+2} - \frac{3}{x+3} + \frac{1}{x-1} \right) dx$$

↓ 2009

$$\log(1-x) - 2\log(x+2) - 3\log(x+3)$$

input `Int[(18 - 2*x - 4*x^2)/(-6 + x + 4*x^2 + x^3),x]`

output `Log[1 - x] - 2*Log[2 + x] - 3*Log[3 + x]`

3.191.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2462 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ [Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

3.191.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

method	result	size
default	$\ln(-1+x) - 2\ln(2+x) - 3\ln(3+x)$	18
norman	$\ln(-1+x) - 2\ln(2+x) - 3\ln(3+x)$	18
risch	$\ln(-1+x) - 2\ln(2+x) - 3\ln(3+x)$	18
parallelrisch	$\ln(-1+x) - 2\ln(2+x) - 3\ln(3+x)$	18

input `int((-4*x^2-2*x+18)/(x^3+4*x^2+x-6),x,method=_RETURNVERBOSE)`output `ln(-1+x)-2*ln(2+x)-3*ln(3+x)`**3.191.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{18 - 2x - 4x^2}{-6 + x + 4x^2 + x^3} dx = -3 \log(x + 3) - 2 \log(x + 2) + \log(x - 1)$$

input `integrate((-4*x^2-2*x+18)/(x^3+4*x^2+x-6),x, algorithm="fracas")`output `-3*log(x + 3) - 2*log(x + 2) + log(x - 1)`**3.191.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{18 - 2x - 4x^2}{-6 + x + 4x^2 + x^3} dx = \log(x - 1) - 2 \log(x + 2) - 3 \log(x + 3)$$

input `integrate((-4*x**2-2*x+18)/(x**3+4*x**2+x-6),x)`output `log(x - 1) - 2*log(x + 2) - 3*log(x + 3)`

3.191.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{18 - 2x - 4x^2}{-6 + x + 4x^2 + x^3} dx = -3 \log(x + 3) - 2 \log(x + 2) + \log(x - 1)$$

input `integrate((-4*x^2-2*x+18)/(x^3+4*x^2+x-6),x, algorithm="maxima")`output `-3*log(x + 3) - 2*log(x + 2) + log(x - 1)`**3.191.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.05

$$\int \frac{18 - 2x - 4x^2}{-6 + x + 4x^2 + x^3} dx = -3 \log(|x + 3|) - 2 \log(|x + 2|) + \log(|x - 1|)$$

input `integrate((-4*x^2-2*x+18)/(x^3+4*x^2+x-6),x, algorithm="giac")`output `-3*log(abs(x + 3)) - 2*log(abs(x + 2)) + log(abs(x - 1))`**3.191.9 Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{18 - 2x - 4x^2}{-6 + x + 4x^2 + x^3} dx = \ln(x - 1) - 2 \ln(x + 2) - 3 \ln(x + 3)$$

input `int(-(2*x + 4*x^2 - 18)/(x + 4*x^2 + x^3 - 6),x)`output `log(x - 1) - 2*log(x + 2) - 3*log(x + 3)`

$$3.192 \quad \int \frac{2x+x^2}{4+3x^2+x^3} dx$$

3.192.1 Optimal result	1006
3.192.2 Mathematica [A] (verified)	1006
3.192.3 Rubi [A] (verified)	1007
3.192.4 Maple [A] (verified)	1007
3.192.5 Fricas [A] (verification not implemented)	1008
3.192.6 Sympy [A] (verification not implemented)	1008
3.192.7 Maxima [A] (verification not implemented)	1008
3.192.8 Giac [A] (verification not implemented)	1009
3.192.9 Mupad [B] (verification not implemented)	1009

3.192.1 Optimal result

Integrand size = 20, antiderivative size = 15

$$\int \frac{2x+x^2}{4+3x^2+x^3} dx = \frac{1}{3} \log(4+3x^2+x^3)$$

output `1/3*ln(x^3+3*x^2+4)`

3.192.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{2x+x^2}{4+3x^2+x^3} dx = \frac{1}{3} \log(4+3x^2+x^3)$$

input `Integrate[(2*x + x^2)/(4 + 3*x^2 + x^3),x]`

output `Log[4 + 3*x^2 + x^3]/3`

3.192.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {2020}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2 + 2x}{x^3 + 3x^2 + 4} dx$$

↓ 2020

$$\frac{1}{3} \log(x^3 + 3x^2 + 4)$$

input `Int[(2*x + x^2)/(4 + 3*x^2 + x^3),x]`

output `Log[4 + 3*x^2 + x^3]/3`

3.192.3.1 Defintions of rubi rules used

rule 2020 `Int[(Pp_)/(Qq_), x_Symbol] := With[{p = Expon[Pp, x], q = Expon[Qq, x]}, Simp[Coeff[Pp, x, p]*(Log[RemoveContent[Qq, x]]/(q*Coeff[Qq, x, q])), x] /; EqQ[p, q - 1] && EqQ[Pp, Simplify[(Coeff[Pp, x, p]/(q*Coeff[Qq, x, q]))*D[Qq, x]]] /; PolyQ[Pp, x] && PolyQ[Qq, x]`

3.192.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
default	$\frac{\ln(x^3+3x^2+4)}{3}$	14
norman	$\frac{\ln(x^3+3x^2+4)}{3}$	14
risch	$\frac{\ln(x^3+3x^2+4)}{3}$	14
parallelrisch	$\frac{\ln(x^3+3x^2+4)}{3}$	14

input `int((x^2+2*x)/(x^3+3*x^2+4),x,method=_RETURNVERBOSE)`

output $1/3*\ln(x^3+3*x^2+4)$

3.192.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{2x + x^2}{4 + 3x^2 + x^3} dx = \frac{1}{3} \log(x^3 + 3x^2 + 4)$$

input `integrate((x^2+2*x)/(x^3+3*x^2+4),x, algorithm="fricas")`

output $1/3*\log(x^3 + 3*x^2 + 4)$

3.192.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \frac{2x + x^2}{4 + 3x^2 + x^3} dx = \frac{\log(x^3 + 3x^2 + 4)}{3}$$

input `integrate((x**2+2*x)/(x**3+3*x**2+4),x)`

output $\log(x**3 + 3*x**2 + 4)/3$

3.192.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{2x + x^2}{4 + 3x^2 + x^3} dx = \frac{1}{3} \log(x^3 + 3x^2 + 4)$$

input `integrate((x^2+2*x)/(x^3+3*x^2+4),x, algorithm="maxima")`

output $1/3*\log(x^3 + 3*x^2 + 4)$

3.192.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{2x + x^2}{4 + 3x^2 + x^3} dx = \frac{1}{3} \log(|x^3 + 3x^2 + 4|)$$

input `integrate((x^2+2*x)/(x^3+3*x^2+4),x, algorithm="giac")`output `1/3*log(abs(x^3 + 3*x^2 + 4))`**3.192.9 Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{2x + x^2}{4 + 3x^2 + x^3} dx = \frac{\ln(x^3 + 3x^2 + 4)}{3}$$

input `int((2*x + x^2)/(3*x^2 + x^3 + 4),x)`output `log(3*x^2 + x^3 + 4)/3`

3.193 $\int \frac{1}{(-1+x)^2 x^2} dx$

3.193.1 Optimal result	1010
3.193.2 Mathematica [A] (verified)	1010
3.193.3 Rubi [A] (verified)	1011
3.193.4 Maple [A] (verified)	1012
3.193.5 Fricas [A] (verification not implemented)	1012
3.193.6 Sympy [A] (verification not implemented)	1012
3.193.7 Maxima [A] (verification not implemented)	1013
3.193.8 Giac [A] (verification not implemented)	1013
3.193.9 Mupad [B] (verification not implemented)	1013

3.193.1 Optimal result

Integrand size = 9, antiderivative size = 25

$$\int \frac{1}{(-1+x)^2 x^2} dx = \frac{1}{1-x} - \frac{1}{x} - 2 \log(1-x) + 2 \log(x)$$

output `1/(1-x)-1/x-2*ln(1-x)+2*ln(x)`

3.193.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1}{(-1+x)^2 x^2} dx = -\frac{1}{-1+x} - \frac{1}{x} - 2 \log(1-x) + 2 \log(x)$$

input `Integrate[1/((-1 + x)^2*x^2),x]`

output `-(-1 + x)^(-1) - x^(-1) - 2*Log[1 - x] + 2*Log[x]`

3.193.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x-1)^2 x^2} dx$$

$$\downarrow 54$$

$$\int \left(\frac{1}{x^2} + \frac{2}{x} - \frac{2}{x-1} + \frac{1}{(x-1)^2} \right) dx$$

$$\downarrow 2009$$

$$\frac{1}{1-x} - \frac{1}{x} - 2 \log(1-x) + 2 \log(x)$$

input `Int[1/((-1 + x)^2*x^2),x]`

output `(1 - x)^(-1) - x^(-1) - 2*Log[1 - x] + 2*Log[x]`

3.193.3.1 Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.193.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

method	result	size
default	$-\frac{1}{-1+x} - 2 \ln(-1+x) - \frac{1}{x} + 2 \ln(x)$	24
norman	$\frac{1-2x}{x(-1+x)} + 2 \ln(x) - 2 \ln(-1+x)$	26
risch	$\frac{1-2x}{x(-1+x)} + 2 \ln(x) - 2 \ln(-1+x)$	26
meijerg	$-\frac{1}{x} + 1 + 2 \ln(x) + 2i\pi + \frac{3x}{-3x+3} - 2 \ln(1-x)$	34
parallelrisch	$\frac{2x^2 \ln(x) - 2 \ln(-1+x)x^2 + 1 - 2x \ln(x) + 2 \ln(-1+x)x - 2x}{x(-1+x)}$	43

input `int(1/(-1+x)^2/x^2,x,method=_RETURNVERBOSE)`output `-1/(-1+x)-2*ln(-1+x)-1/x+2*ln(x)`**3.193.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.60

$$\int \frac{1}{(-1+x)^2 x^2} dx = -\frac{2(x^2-x)\log(x-1) - 2(x^2-x)\log(x) + 2x-1}{x^2-x}$$

input `integrate(1/(-1+x)^2/x^2,x, algorithm="fracas")`output `-(2*(x^2 - x)*log(x - 1) - 2*(x^2 - x)*log(x) + 2*x - 1)/(x^2 - x)`**3.193.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{1}{(-1+x)^2 x^2} dx = \frac{1-2x}{x^2-x} + 2 \log(x) - 2 \log(x-1)$$

input `integrate(1/(-1+x)**2/x**2,x)`output `(1 - 2*x)/(x**2 - x) + 2*log(x) - 2*log(x - 1)`

3.193.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08

$$\int \frac{1}{(-1+x)^2 x^2} dx = -\frac{2x-1}{x^2-x} - 2 \log(x-1) + 2 \log(x)$$

input `integrate(1/(-1+x)^2/x^2,x, algorithm="maxima")`output `-(2*x - 1)/(x^2 - x) - 2*log(x - 1) + 2*log(x)`**3.193.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.20

$$\int \frac{1}{(-1+x)^2 x^2} dx = -\frac{1}{x-1} + \frac{1}{\frac{1}{x-1} + 1} + 2 \log\left(\left|-\frac{1}{x-1} - 1\right|\right)$$

input `integrate(1/(-1+x)^2/x^2,x, algorithm="giac")`output `-1/(x - 1) + 1/(1/(x - 1) + 1) + 2*log(abs(-1/(x - 1) - 1))`**3.193.9 Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08

$$\int \frac{1}{(-1+x)^2 x^2} dx = \frac{1}{x(x-1)} - \frac{2}{x-1} - 2 \ln\left(\frac{x-1}{x}\right)$$

input `int(1/(x^2*(x - 1)^2),x)`output `1/(x*(x - 1)) - 2/(x - 1) - 2*log((x - 1)/x)`

3.194 $\int \frac{x^2}{(1+x)^3} dx$

3.194.1 Optimal result	1014
3.194.2 Mathematica [A] (verified)	1014
3.194.3 Rubi [A] (verified)	1015
3.194.4 Maple [A] (verified)	1016
3.194.5 Fricas [A] (verification not implemented)	1016
3.194.6 Sympy [A] (verification not implemented)	1016
3.194.7 Maxima [A] (verification not implemented)	1017
3.194.8 Giac [A] (verification not implemented)	1017
3.194.9 Mupad [B] (verification not implemented)	1017

3.194.1 Optimal result

Integrand size = 9, antiderivative size = 21

$$\int \frac{x^2}{(1+x)^3} dx = -\frac{1}{2(1+x)^2} + \frac{2}{1+x} + \log(1+x)$$

output `-1/2/(1+x)^2+2/(1+x)+ln(1+x)`

3.194.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{(1+x)^3} dx = -\frac{1}{2(1+x)^2} + \frac{2}{1+x} + \log(1+x)$$

input `Integrate[x^2/(1 + x)^3,x]`

output `-1/2*1/(1 + x)^2 + 2/(1 + x) + Log[1 + x]`

3.194.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{(x+1)^3} dx$$

$$\downarrow 49$$

$$\int \left(\frac{1}{x+1} - \frac{2}{(x+1)^2} + \frac{1}{(x+1)^3} \right) dx$$

$$\downarrow 2009$$

$$\frac{2}{x+1} - \frac{1}{2(x+1)^2} + \log(x+1)$$

input `Int[x^2/(1 + x)^3,x]`

output `-1/2*1/(1 + x)^2 + 2/(1 + x) + Log[1 + x]`

3.194.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.194.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

method	result	size
norman	$\frac{2x+\frac{3}{2}}{(1+x)^2} + \ln(1+x)$	17
risch	$\frac{2x+\frac{3}{2}}{(1+x)^2} + \ln(1+x)$	17
meijerg	$-\frac{x(9x+6)}{6(1+x)^2} + \ln(1+x)$	19
default	$-\frac{1}{2(1+x)^2} + \frac{2}{1+x} + \ln(1+x)$	20
parallelrisc	$\frac{2\ln(1+x)x^2+3+4\ln(1+x)x+2\ln(1+x)+4x}{2(1+x)^2}$	35

input `int(x^2/(1+x)^3,x,method=_RETURNVERBOSE)`output `(2*x+3/2)/(1+x)^2+ln(1+x)`**3.194.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.48

$$\int \frac{x^2}{(1+x)^3} dx = \frac{2(x^2 + 2x + 1) \log(x + 1) + 4x + 3}{2(x^2 + 2x + 1)}$$

input `integrate(x^2/(1+x)^3,x, algorithm="fricas")`output `1/2*(2*(x^2 + 2*x + 1)*log(x + 1) + 4*x + 3)/(x^2 + 2*x + 1)`**3.194.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{x^2}{(1+x)^3} dx = \frac{4x + 3}{2x^2 + 4x + 2} + \log(x + 1)$$

input `integrate(x**2/(1+x)**3,x)`output `(4*x + 3)/(2*x**2 + 4*x + 2) + log(x + 1)`

3.194.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \frac{x^2}{(1+x)^3} dx = \frac{4x+3}{2(x^2+2x+1)} + \log(x+1)$$

input `integrate(x^2/(1+x)^3,x, algorithm="maxima")`output `1/2*(4*x + 3)/(x^2 + 2*x + 1) + log(x + 1)`**3.194.8 Giac [A] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

$$\int \frac{x^2}{(1+x)^3} dx = \frac{4x+3}{2(x+1)^2} + \log(|x+1|)$$

input `integrate(x^2/(1+x)^3,x, algorithm="giac")`output `1/2*(4*x + 3)/(x + 1)^2 + log(abs(x + 1))`**3.194.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{(1+x)^3} dx = \ln(x+1) + \frac{2x+\frac{3}{2}}{x^2+2x+1}$$

input `int(x^2/(x + 1)^3,x)`output `log(x + 1) + (2*x + 3/2)/(2*x + x^2 + 1)`

3.195 $\int \frac{1}{-x^2+x^4} dx$

3.195.1 Optimal result	1018
3.195.2 Mathematica [B] (verified)	1018
3.195.3 Rubi [A] (verified)	1019
3.195.4 Maple [C] (verified)	1020
3.195.5 Fricas [B] (verification not implemented)	1020
3.195.6 Sympy [B] (verification not implemented)	1021
3.195.7 Maxima [A] (verification not implemented)	1021
3.195.8 Giac [B] (verification not implemented)	1021
3.195.9 Mupad [B] (verification not implemented)	1022

3.195.1 Optimal result

Integrand size = 11, antiderivative size = 8

$$\int \frac{1}{-x^2+x^4} dx = \frac{1}{x} - \operatorname{arctanh}(x)$$

output `1/x-arctanh(x)`

3.195.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 22 vs. $2(8) = 16$.

Time = 0.00 (sec) , antiderivative size = 22, normalized size of antiderivative = 2.75

$$\int \frac{1}{-x^2+x^4} dx = \frac{1}{x} + \frac{1}{2} \log(1-x) - \frac{1}{2} \log(1+x)$$

input `Integrate[(-x^2 + x^4)^(-1),x]`

output `x^(-1) + Log[1 - x]/2 - Log[1 + x]/2`

3.195.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {1397, 264, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x^4 - x^2} dx \\ & \quad \downarrow \text{1397} \\ & \int \frac{1}{x^2(x^2 - 1)} dx \\ & \quad \downarrow \text{264} \\ & \int \frac{1}{x^2 - 1} dx + \frac{1}{x} \\ & \quad \downarrow \text{220} \\ & \frac{1}{x} - \operatorname{arctanh}(x) \end{aligned}$$

input `Int[(-x^2 + x^4)^(-1),x]`

output `x^(-1) - ArcTanh[x]`

3.195.3.1 Defintions of rubi rules used

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 264 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*((m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 1397 `Int[((b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := Int[x^(2*p)*(b + c*x^2)^p, x] /; FreeQ[{b, c}, x] && IntegerQ[p]`

3.195.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 2.00

method	result	size
meijerg	$-\frac{i\left(\frac{2i}{x}-2i\operatorname{arctanh}(x)\right)}{2}$	16
default	$\frac{\ln(-1+x)}{2} + \frac{1}{x} - \frac{\ln(1+x)}{2}$	17
norman	$\frac{\ln(-1+x)}{2} + \frac{1}{x} - \frac{\ln(1+x)}{2}$	17
risch	$\frac{\ln(-1+x)}{2} + \frac{1}{x} - \frac{\ln(1+x)}{2}$	17
parallelrisch	$\frac{\ln(-1+x)x - \ln(1+x)x + 2}{2x}$	21

input `int(1/(x^4-x^2),x,method=_RETURNVERBOSE)`

output `-1/2*I*(2*I/x-2*I*arctanh(x))`

3.195.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. 2(8) = 16.

Time = 0.25 (sec) , antiderivative size = 20, normalized size of antiderivative = 2.50

$$\int \frac{1}{-x^2 + x^4} dx = -\frac{x \log(x+1) - x \log(x-1) - 2}{2x}$$

input `integrate(1/(x^4-x^2),x, algorithm="fricas")`

output `-1/2*(x*log(x + 1) - x*log(x - 1) - 2)/x`

3.195.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(5) = 10$.

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.88

$$\int \frac{1}{-x^2 + x^4} dx = \frac{\log(x-1)}{2} - \frac{\log(x+1)}{2} + \frac{1}{x}$$

input `integrate(1/(x**4-x**2),x)`

output `log(x - 1)/2 - log(x + 1)/2 + 1/x`

3.195.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 2.00

$$\int \frac{1}{-x^2 + x^4} dx = \frac{1}{x} - \frac{1}{2} \log(x+1) + \frac{1}{2} \log(x-1)$$

input `integrate(1/(x^4-x^2),x, algorithm="maxima")`

output `1/x - 1/2*log(x + 1) + 1/2*log(x - 1)`

3.195.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18 vs. $2(8) = 16$.

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 2.25

$$\int \frac{1}{-x^2 + x^4} dx = \frac{1}{x} - \frac{1}{2} \log(|x+1|) + \frac{1}{2} \log(|x-1|)$$

input `integrate(1/(x^4-x^2),x, algorithm="giac")`

output `1/x - 1/2*log(abs(x + 1)) + 1/2*log(abs(x - 1))`

3.195.9 Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{1}{-x^2 + x^4} dx = \frac{1}{x} - \operatorname{atanh}(x)$$

input `int(-1/(x^2 - x^4),x)`

output `1/x - atanh(x)`

3.196 $\int \frac{-x+2x^3}{1-x^2+x^4} dx$

3.196.1 Optimal result	1023
3.196.2 Mathematica [A] (verified)	1023
3.196.3 Rubi [A] (verified)	1024
3.196.4 Maple [A] (verified)	1024
3.196.5 Fricas [A] (verification not implemented)	1025
3.196.6 Sympy [A] (verification not implemented)	1025
3.196.7 Maxima [A] (verification not implemented)	1025
3.196.8 Giac [A] (verification not implemented)	1026
3.196.9 Mupad [B] (verification not implemented)	1026

3.196.1 Optimal result

Integrand size = 22, antiderivative size = 15

$$\int \frac{-x + 2x^3}{1 - x^2 + x^4} dx = \frac{1}{2} \log(1 - x^2 + x^4)$$

output `1/2*ln(x^4-x^2+1)`

3.196.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{-x + 2x^3}{1 - x^2 + x^4} dx = \frac{1}{2} \log(1 - x^2 + x^4)$$

input `Integrate[(-x + 2*x^3)/(1 - x^2 + x^4), x]`

output `Log[1 - x^2 + x^4]/2`

3.196.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {2020}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x^3 - x}{x^4 - x^2 + 1} dx$$

↓ 2020

$$\frac{1}{2} \log(x^4 - x^2 + 1)$$

input `Int[(-x + 2*x^3)/(1 - x^2 + x^4),x]`

output `Log[1 - x^2 + x^4]/2`

3.196.3.1 Defintions of rubi rules used

rule 2020 `Int[(Pp_)/(Qq_), x_Symbol] := With[{p = Expon[Pp, x], q = Expon[Qq, x]}, Simp[Coeff[Pp, x, p]*(Log[RemoveContent[Qq, x]]/(q*Coeff[Qq, x, q])), x] /; EqQ[p, q - 1] && EqQ[Pp, Simplify[(Coeff[Pp, x, p]/(q*Coeff[Qq, x, q]))*D[Qq, x]]] /; PolyQ[Pp, x] && PolyQ[Qq, x]`

3.196.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
default	$\frac{\ln(x^4 - x^2 + 1)}{2}$	14
norman	$\frac{\ln(x^4 - x^2 + 1)}{2}$	14
risch	$\frac{\ln(x^4 - x^2 + 1)}{2}$	14
parallelrisch	$\frac{\ln(x^4 - x^2 + 1)}{2}$	14

input `int((2*x^3-x)/(x^4-x^2+1),x,method=_RETURNVERBOSE)`

output $1/2*\ln(x^4-x^2+1)$

3.196.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{-x + 2x^3}{1 - x^2 + x^4} dx = \frac{1}{2} \log(x^4 - x^2 + 1)$$

input `integrate((2*x^3-x)/(x^4-x^2+1),x, algorithm="fricas")`

output $1/2*\log(x^4 - x^2 + 1)$

3.196.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int \frac{-x + 2x^3}{1 - x^2 + x^4} dx = \frac{\log(x^4 - x^2 + 1)}{2}$$

input `integrate((2*x**3-x)/(x**4-x**2+1),x)`

output $\log(x**4 - x**2 + 1)/2$

3.196.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{-x + 2x^3}{1 - x^2 + x^4} dx = \frac{1}{2} \log(x^4 - x^2 + 1)$$

input `integrate((2*x^3-x)/(x^4-x^2+1),x, algorithm="maxima")`

output $1/2*\log(x^4 - x^2 + 1)$

3.196.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{-x + 2x^3}{1 - x^2 + x^4} dx = \frac{1}{2} \log(x^4 - x^2 + 1)$$

input `integrate((2*x^3-x)/(x^4-x^2+1),x, algorithm="giac")`output `1/2*log(x^4 - x^2 + 1)`**3.196.9 Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{-x + 2x^3}{1 - x^2 + x^4} dx = \frac{\ln(x^4 - x^2 + 1)}{2}$$

input `int(-(x - 2*x^3)/(x^4 - x^2 + 1),x)`output `log(x^4 - x^2 + 1)/2`

3.197 $\int \frac{x^3}{1+x^2} dx$

3.197.1 Optimal result	1027
3.197.2 Mathematica [A] (verified)	1027
3.197.3 Rubi [A] (verified)	1028
3.197.4 Maple [A] (verified)	1029
3.197.5 Fricas [A] (verification not implemented)	1029
3.197.6 Sympy [A] (verification not implemented)	1029
3.197.7 Maxima [A] (verification not implemented)	1030
3.197.8 Giac [A] (verification not implemented)	1030
3.197.9 Mupad [B] (verification not implemented)	1030

3.197.1 Optimal result

Integrand size = 11, antiderivative size = 18

$$\int \frac{x^3}{1+x^2} dx = \frac{x^2}{2} - \frac{1}{2} \log(1+x^2)$$

output `1/2*x^2-1/2*ln(x^2+1)`

3.197.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{x^3}{1+x^2} dx = \frac{x^2}{2} - \frac{1}{2} \log(1+x^2)$$

input `Integrate[x^3/(1 + x^2),x]`

output `x^2/2 - Log[1 + x^2]/2`

3.197.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^3}{x^2+1} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int \frac{x^2}{x^2+1} dx^2 \\ & \quad \downarrow \text{49} \\ & \frac{1}{2} \int \left(1 + \frac{1}{-x^2-1} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} (x^2 - \log(x^2+1)) \end{aligned}$$

input `Int[x^3/(1 + x^2), x]`

output `(x^2 - Log[1 + x^2])/2`

3.197.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.197.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{x^2}{2} - \frac{\ln(x^2+1)}{2}$	15
norman	$\frac{x^2}{2} - \frac{\ln(x^2+1)}{2}$	15
meijerg	$\frac{x^2}{2} - \frac{\ln(x^2+1)}{2}$	15
risch	$\frac{x^2}{2} - \frac{\ln(x^2+1)}{2}$	15
parallelrisch	$\frac{x^2}{2} - \frac{\ln(x^2+1)}{2}$	15

input `int(x^3/(x^2+1),x,method=_RETURNVERBOSE)`output `1/2*x^2-1/2*ln(x^2+1)`**3.197.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{x^3}{1+x^2} dx = \frac{1}{2} x^2 - \frac{1}{2} \log(x^2 + 1)$$

input `integrate(x^3/(x^2+1),x, algorithm="fricas")`output `1/2*x^2 - 1/2*log(x^2 + 1)`**3.197.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \frac{x^3}{1+x^2} dx = \frac{x^2}{2} - \frac{\log(x^2 + 1)}{2}$$

input `integrate(x**3/(x**2+1),x)`output `x**2/2 - log(x**2 + 1)/2`

3.197.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{x^3}{1+x^2} dx = \frac{1}{2}x^2 - \frac{1}{2}\log(x^2+1)$$

input `integrate(x^3/(x^2+1),x, algorithm="maxima")`output `1/2*x^2 - 1/2*log(x^2 + 1)`**3.197.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{x^3}{1+x^2} dx = \frac{1}{2}x^2 - \frac{1}{2}\log(x^2+1)$$

input `integrate(x^3/(x^2+1),x, algorithm="giac")`output `1/2*x^2 - 1/2*log(x^2 + 1)`**3.197.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{x^3}{1+x^2} dx = \frac{x^2}{2} - \frac{\ln(x^2+1)}{2}$$

input `int(x^3/(x^2 + 1),x)`output `x^2/2 - log(x^2 + 1)/2`

3.198 $\int \frac{-1+x}{2+2x+x^2} dx$

3.198.1 Optimal result	1031
3.198.2 Mathematica [A] (verified)	1031
3.198.3 Rubi [A] (verified)	1032
3.198.4 Maple [A] (verified)	1033
3.198.5 Fricas [A] (verification not implemented)	1034
3.198.6 Sympy [A] (verification not implemented)	1034
3.198.7 Maxima [A] (verification not implemented)	1034
3.198.8 Giac [A] (verification not implemented)	1035
3.198.9 Mupad [B] (verification not implemented)	1035

3.198.1 Optimal result

Integrand size = 14, antiderivative size = 20

$$\int \frac{-1+x}{2+2x+x^2} dx = -2 \arctan(1+x) + \frac{1}{2} \log(2+2x+x^2)$$

output `-2*arctan(1+x)+1/2*ln(x^2+2*x+2)`

3.198.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{-1+x}{2+2x+x^2} dx = -2 \arctan(1+x) + \frac{1}{2} \log(2+2x+x^2)$$

input `Integrate[(-1 + x)/(2 + 2*x + x^2), x]`

output `-2*ArcTan[1 + x] + Log[2 + 2*x + x^2]/2`

3.198.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {1142, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x-1}{x^2+2x+2} dx \\
 & \quad \downarrow \text{1142} \\
 & \frac{1}{2} \int \frac{2(x+1)}{x^2+2x+2} dx - 2 \int \frac{1}{x^2+2x+2} dx \\
 & \quad \downarrow \text{27} \\
 & \int \frac{x+1}{x^2+2x+2} dx - 2 \int \frac{1}{x^2+2x+2} dx \\
 & \quad \downarrow \text{1082} \\
 & \int \frac{x+1}{x^2+2x+2} dx + 2 \int \frac{1}{-(x+1)^2-1} d(x+1) \\
 & \quad \downarrow \text{217} \\
 & \int \frac{x+1}{x^2+2x+2} dx - 2 \arctan(x+1) \\
 & \quad \downarrow \text{1103} \\
 & \frac{1}{2} \log(x^2+2x+2) - 2 \arctan(x+1)
 \end{aligned}$$

input `Int[(-1 + x)/(2 + 2*x + x^2), x]`

output `-2*ArcTan[1 + x] + Log[2 + 2*x + x^2]/2`

3.198.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1142 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

3.198.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

method	result	size
default	$-2 \arctan(1+x) + \frac{\ln(x^2+2x+2)}{2}$	19
risch	$-2 \arctan(1+x) + \frac{\ln(x^2+2x+2)}{2}$	19
parallelrisch	$\frac{\ln(x+1-i)}{2} + i \ln(x+1-i) + \frac{\ln(x+1+i)}{2} - i \ln(x+1+i)$	36

input `int((-1+x)/(x^2+2*x+2),x,method=_RETURNVERBOSE)`

output `-2*arctan(1+x)+1/2*ln(x^2+2*x+2)`

3.198.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{-1+x}{2+2x+x^2} dx = -2 \arctan(x+1) + \frac{1}{2} \log(x^2+2x+2)$$

input `integrate((-1+x)/(x^2+2*x+2),x, algorithm="fricas")`output `-2*arctan(x + 1) + 1/2*log(x^2 + 2*x + 2)`**3.198.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{-1+x}{2+2x+x^2} dx = \frac{\log(x^2+2x+2)}{2} - 2 \operatorname{atan}(x+1)$$

input `integrate((-1+x)/(x**2+2*x+2),x)`output `log(x**2 + 2*x + 2)/2 - 2*atan(x + 1)`**3.198.7 Maxima [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{-1+x}{2+2x+x^2} dx = -2 \arctan(x+1) + \frac{1}{2} \log(x^2+2x+2)$$

input `integrate((-1+x)/(x^2+2*x+2),x, algorithm="maxima")`output `-2*arctan(x + 1) + 1/2*log(x^2 + 2*x + 2)`

3.198.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{-1+x}{2+2x+x^2} dx = -2 \arctan(x+1) + \frac{1}{2} \log(x^2+2x+2)$$

input `integrate((-1+x)/(x^2+2*x+2),x, algorithm="giac")`output `-2*arctan(x + 1) + 1/2*log(x^2 + 2*x + 2)`**3.198.9 Mupad [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{-1+x}{2+2x+x^2} dx = \frac{\ln(x^2+2x+2)}{2} - 2 \operatorname{atan}(x+1)$$

input `int((x - 1)/(2*x + x^2 + 2),x)`output `log(2*x + x^2 + 2)/2 - 2*atan(x + 1)`

3.199 $\int \frac{x}{1+x+x^2} dx$

3.199.1 Optimal result	1036
3.199.2 Mathematica [A] (verified)	1036
3.199.3 Rubi [A] (verified)	1037
3.199.4 Maple [A] (verified)	1038
3.199.5 Fricas [A] (verification not implemented)	1038
3.199.6 Sympy [A] (verification not implemented)	1039
3.199.7 Maxima [A] (verification not implemented)	1039
3.199.8 Giac [A] (verification not implemented)	1039
3.199.9 Mupad [B] (verification not implemented)	1040

3.199.1 Optimal result

Integrand size = 10, antiderivative size = 31

$$\int \frac{x}{1+x+x^2} dx = -\frac{\arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{2} \log(1+x+x^2)$$

output `1/2*ln(x^2+x+1)-1/3*arctan(1/3*(1+2*x)*3^(1/2))*3^(1/2)`

3.199.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{x}{1+x+x^2} dx = -\frac{\arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{2} \log(1+x+x^2)$$

input `Integrate[x/(1+x+x^2),x]`

output `-(ArcTan[(1+2*x)/Sqrt[3]]/Sqrt[3]) + Log[1+x+x^2]/2`

3.199.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {1142, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x}{x^2 + x + 1} dx \\
 & \quad \downarrow \text{1142} \\
 & \frac{1}{2} \int \frac{2x + 1}{x^2 + x + 1} dx - \frac{1}{2} \int \frac{1}{x^2 + x + 1} dx \\
 & \quad \downarrow \text{1083} \\
 & \frac{1}{2} \int \frac{2x + 1}{x^2 + x + 1} dx + \int \frac{1}{-(2x + 1)^2 - 3} d(2x + 1) \\
 & \quad \downarrow \text{217} \\
 & \frac{1}{2} \int \frac{2x + 1}{x^2 + x + 1} dx - \frac{\arctan\left(\frac{2x+1}{\sqrt{3}}\right)}{\sqrt{3}} \\
 & \quad \downarrow \text{1103} \\
 & \frac{1}{2} \log(x^2 + x + 1) - \frac{\arctan\left(\frac{2x+1}{\sqrt{3}}\right)}{\sqrt{3}}
 \end{aligned}$$

input `Int[x/(1 + x + x^2),x]`

output `-(ArcTan[(1 + 2*x)/Sqrt[3]]/Sqrt[3]) + Log[1 + x + x^2]/2`

3.199.3.1 Defintions of rubi rules used

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & (LtQ[a, 0] || LtQ[b, 0])`

rule 1083 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

3.199.4 Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

method	result	size
default	$\frac{\ln(x^2+x+1)}{2} - \frac{\arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3}$	27
risch	$-\frac{\arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3} + \frac{\ln(4x^2+4x+4)}{2}$	31

input `int(x/(x^2+x+1),x,method=_RETURNVERBOSE)`

output `1/2*ln(x^2+x+1)-1/3*arctan(1/3*(1+2*x)*3^(1/2))*3^(1/2)`

3.199.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{x}{1+x+x^2} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) + \frac{1}{2} \log(x^2+x+1)$$

input `integrate(x/(x^2+x+1),x, algorithm="fricas")`

output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + 1/2*log(x^2 + x + 1)`

3.199.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.10

$$\int \frac{x}{1+x+x^2} dx = \frac{\log(x^2+x+1)}{2} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x}{3} + \frac{\sqrt{3}}{3}\right)}{3}$$

input `integrate(x/(x**2+x+1),x)`output `log(x**2 + x + 1)/2 - sqrt(3)*atan(2*sqrt(3)*x/3 + sqrt(3)/3)/3`**3.199.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{x}{1+x+x^2} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) + \frac{1}{2} \log(x^2+x+1)$$

input `integrate(x/(x^2+x+1),x, algorithm="maxima")`output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + 1/2*log(x^2 + x + 1)`**3.199.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{x}{1+x+x^2} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) + \frac{1}{2} \log(x^2+x+1)$$

input `integrate(x/(x^2+x+1),x, algorithm="giac")`output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + 1/2*log(x^2 + x + 1)`

3.199.9 Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

$$\int \frac{x}{1+x+x^2} dx = \frac{\ln(x^2+x+1)}{2} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x}{3} + \frac{\sqrt{3}}{3}\right)}{3}$$

input `int(x/(x + x^2 + 1),x)`output `log(x + x^2 + 1)/2 - (3^(1/2)*atan((2*3^(1/2)*x)/3 + 3^(1/2)/3))/3`

3.200 $\int \frac{7+5x+4x^2}{5+4x+4x^2} dx$

3.200.1 Optimal result1041
3.200.2 Mathematica [A] (verified)1041
3.200.3 Rubi [A] (verified)1042
3.200.4 Maple [A] (verified)1043
3.200.5 Fricas [A] (verification not implemented)1043
3.200.6 Sympy [A] (verification not implemented)1043
3.200.7 Maxima [A] (verification not implemented)1044
3.200.8 Giac [A] (verification not implemented)1044
3.200.9 Mupad [B] (verification not implemented)1044

3.200.1 Optimal result

Integrand size = 23, antiderivative size = 27

$$\int \frac{7+5x+4x^2}{5+4x+4x^2} dx = x + \frac{3}{8} \arctan\left(\frac{1}{2} + x\right) + \frac{1}{8} \log(5+4x+4x^2)$$

output `x+3/8*arctan(1/2+x)+1/8*ln(4*x^2+4*x+5)`

3.200.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.15

$$\int \frac{7+5x+4x^2}{5+4x+4x^2} dx = x + \frac{3}{8} \arctan\left(\frac{1}{2}(1+2x)\right) + \frac{1}{8} \log(5+4x+4x^2)$$

input `Integrate[(7 + 5*x + 4*x^2)/(5 + 4*x + 4*x^2),x]`

output `x + (3*ArcTan[(1 + 2*x)/2])/8 + Log[5 + 4*x + 4*x^2]/8`

3.200.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2188, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{4x^2 + 5x + 7}{4x^2 + 4x + 5} dx$$

↓ 2188

$$\int \left(\frac{x+2}{4x^2 + 4x + 5} + 1 \right) dx$$

↓ 2009

$$\frac{3}{8} \arctan \left(x + \frac{1}{2} \right) + \frac{1}{8} \log(4x^2 + 4x + 5) + x$$

input `Int[(7 + 5*x + 4*x^2)/(5 + 4*x + 4*x^2),x]`

output `x + (3*ArcTan[1/2 + x])/8 + Log[5 + 4*x + 4*x^2]/8`

3.200.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2188 `Int[(Pq_)*((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[Pq*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

3.200.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

method	result	size
default	$x + \frac{3 \arctan(x + \frac{1}{2})}{8} + \frac{\ln(4x^2 + 4x + 5)}{8}$	22
risch	$x + \frac{3 \arctan(x + \frac{1}{2})}{8} + \frac{\ln(4x^2 + 4x + 5)}{8}$	22
parallelrisch	$x + \frac{\ln(x + \frac{1}{2} - i)}{8} - \frac{3i \ln(x + \frac{1}{2} - i)}{16} + \frac{\ln(x + \frac{1}{2} + i)}{8} + \frac{3i \ln(x + \frac{1}{2} + i)}{16}$	37

input `int((4*x^2+5*x+7)/(4*x^2+4*x+5),x,method=_RETURNVERBOSE)`output `x+3/8*arctan(x+1/2)+1/8*ln(4*x^2+4*x+5)`**3.200.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int \frac{7 + 5x + 4x^2}{5 + 4x + 4x^2} dx = x + \frac{3}{8} \arctan\left(x + \frac{1}{2}\right) + \frac{1}{8} \log(4x^2 + 4x + 5)$$

input `integrate((4*x^2+5*x+7)/(4*x^2+4*x+5),x, algorithm="fricas")`output `x + 3/8*arctan(x + 1/2) + 1/8*log(4*x^2 + 4*x + 5)`**3.200.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int \frac{7 + 5x + 4x^2}{5 + 4x + 4x^2} dx = x + \frac{\log(x^2 + x + \frac{5}{4})}{8} + \frac{3 \operatorname{atan}(x + \frac{1}{2})}{8}$$

input `integrate((4*x**2+5*x+7)/(4*x**2+4*x+5),x)`output `x + log(x**2 + x + 5/4)/8 + 3*atan(x + 1/2)/8`

3.200.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int \frac{7 + 5x + 4x^2}{5 + 4x + 4x^2} dx = x + \frac{3}{8} \arctan\left(x + \frac{1}{2}\right) + \frac{1}{8} \log(4x^2 + 4x + 5)$$

input `integrate((4*x^2+5*x+7)/(4*x^2+4*x+5),x, algorithm="maxima")`output `x + 3/8*arctan(x + 1/2) + 1/8*log(4*x^2 + 4*x + 5)`**3.200.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int \frac{7 + 5x + 4x^2}{5 + 4x + 4x^2} dx = x + \frac{3}{8} \arctan\left(x + \frac{1}{2}\right) + \frac{1}{8} \log(4x^2 + 4x + 5)$$

input `integrate((4*x^2+5*x+7)/(4*x^2+4*x+5),x, algorithm="giac")`output `x + 3/8*arctan(x + 1/2) + 1/8*log(4*x^2 + 4*x + 5)`**3.200.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.63

$$\int \frac{7 + 5x + 4x^2}{5 + 4x + 4x^2} dx = x + \frac{\ln\left(x^2 + x + \frac{5}{4}\right)}{8} + \frac{3 \operatorname{atan}\left(x + \frac{1}{2}\right)}{8}$$

input `int((5*x + 4*x^2 + 7)/(4*x + 4*x^2 + 5),x)`output `x + log(x + x^2 + 5/4)/8 + (3*atan(x + 1/2))/8`

$$\mathbf{3.201} \quad \int \frac{5-4x+3x^2}{(-1+x)(1+x^2)} dx$$

3.201.1 Optimal result	1045
3.201.2 Mathematica [A] (verified)	1045
3.201.3 Rubi [A] (verified)	1046
3.201.4 Maple [A] (verified)	1047
3.201.5 Fricas [A] (verification not implemented)	1047
3.201.6 Sympy [A] (verification not implemented)	1047
3.201.7 Maxima [A] (verification not implemented)	1048
3.201.8 Giac [A] (verification not implemented)	1048
3.201.9 Mupad [B] (verification not implemented)	1048

3.201.1 Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{5-4x+3x^2}{(-1+x)(1+x^2)} dx = -3 \arctan(x) + 2 \log(1-x) + \frac{1}{2} \log(1+x^2)$$

output `-3*arctan(x)+2*ln(1-x)+1/2*ln(x^2+1)`

3.201.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.22

$$\int \frac{5-4x+3x^2}{(-1+x)(1+x^2)} dx = -3 \arctan(x) + \frac{1}{2} \log(2+2(-1+x)+(-1+x)^2) + 2 \log(-1+x)$$

input `Integrate[(5 - 4*x + 3*x^2)/((-1 + x)*(1 + x^2)),x]`

output `-3*ArcTan[x] + Log[2 + 2*(-1 + x) + (-1 + x)^2]/2 + 2*Log[-1 + x]`

3.201.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2160, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{3x^2 - 4x + 5}{(x-1)(x^2+1)} dx$$

$$\downarrow \text{2160}$$

$$\int \left(\frac{x-3}{x^2+1} + \frac{2}{x-1} \right) dx$$

$$\downarrow \text{2009}$$

$$-3 \arctan(x) + \frac{1}{2} \log(x^2+1) + 2 \log(1-x)$$

input `Int[(5 - 4*x + 3*x^2)/((-1 + x)*(1 + x^2)),x]`

output `-3*ArcTan[x] + 2*Log[1 - x] + Log[1 + x^2]/2`

3.201.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2160 `Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

3.201.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

method	result	size
default	$2 \ln(-1+x) + \frac{\ln(x^2+1)}{2} - 3 \arctan(x)$	20
risch	$2 \ln(-1+x) + \frac{\ln(9x^2+9)}{2} - 3 \arctan(x)$	22
parallelrisch	$2 \ln(-1+x) + \frac{\ln(x-i)}{2} + \frac{3i \ln(x-i)}{2} + \frac{\ln(x+i)}{2} - \frac{3i \ln(x+i)}{2}$	38

input `int((3*x^2-4*x+5)/(-1+x)/(x^2+1),x,method=_RETURNVERBOSE)`output `2*ln(-1+x)+1/2*ln(x^2+1)-3*arctan(x)`**3.201.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{5-4x+3x^2}{(-1+x)(1+x^2)} dx = -3 \arctan(x) + \frac{1}{2} \log(x^2+1) + 2 \log(x-1)$$

input `integrate((3*x^2-4*x+5)/(-1+x)/(x^2+1),x, algorithm="fricas")`output `-3*arctan(x) + 1/2*log(x^2 + 1) + 2*log(x - 1)`**3.201.6 Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{5-4x+3x^2}{(-1+x)(1+x^2)} dx = 2 \log(x-1) + \frac{\log(x^2+1)}{2} - 3 \operatorname{atan}(x)$$

input `integrate((3*x**2-4*x+5)/(-1+x)/(x**2+1),x)`output `2*log(x - 1) + log(x**2 + 1)/2 - 3*atan(x)`

3.201.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{5 - 4x + 3x^2}{(-1 + x)(1 + x^2)} dx = -3 \arctan(x) + \frac{1}{2} \log(x^2 + 1) + 2 \log(x - 1)$$

input `integrate((3*x^2-4*x+5)/(-1+x)/(x^2+1),x, algorithm="maxima")`output `-3*arctan(x) + 1/2*log(x^2 + 1) + 2*log(x - 1)`**3.201.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{5 - 4x + 3x^2}{(-1 + x)(1 + x^2)} dx = -3 \arctan(x) + \frac{1}{2} \log(x^2 + 1) + 2 \log(|x - 1|)$$

input `integrate((3*x^2-4*x+5)/(-1+x)/(x^2+1),x, algorithm="giac")`output `-3*arctan(x) + 1/2*log(x^2 + 1) + 2*log(abs(x - 1))`**3.201.9 Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{5 - 4x + 3x^2}{(-1 + x)(1 + x^2)} dx = 2 \ln(x - 1) + \ln(x - i) \left(\frac{1}{2} + \frac{3}{2}i \right) + \ln(x + i) \left(\frac{1}{2} - \frac{3}{2}i \right)$$

input `int((3*x^2 - 4*x + 5)/((x^2 + 1)*(x - 1)),x)`output `2*log(x - 1) + log(x - 1i)*(1/2 + 3i/2) + log(x + 1i)*(1/2 - 3i/2)`

3.202 $\int \frac{3+2x}{3x+x^3} dx$

3.202.1 Optimal result	1049
3.202.2 Mathematica [A] (verified)	1049
3.202.3 Rubi [A] (verified)	1050
3.202.4 Maple [A] (verified)	1051
3.202.5 Fricas [A] (verification not implemented)	1051
3.202.6 Sympy [A] (verification not implemented)	1051
3.202.7 Maxima [A] (verification not implemented)	1052
3.202.8 Giac [A] (verification not implemented)	1052
3.202.9 Mupad [B] (verification not implemented)	1052

3.202.1 Optimal result

Integrand size = 15, antiderivative size = 28

$$\int \frac{3+2x}{3x+x^3} dx = \frac{2 \arctan\left(\frac{x}{\sqrt{3}}\right)}{\sqrt{3}} + \log(x) - \frac{1}{2} \log(3+x^2)$$

output `ln(x)-1/2*ln(x^2+3)+2/3*arctan(1/3*x*3^(1/2))*3^(1/2)`

3.202.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{3+2x}{3x+x^3} dx = \frac{2 \arctan\left(\frac{x}{\sqrt{3}}\right)}{\sqrt{3}} + \log(x) - \frac{1}{2} \log(3+x^2)$$

input `Integrate[(3 + 2*x)/(3*x + x^3),x]`

output `(2*ArcTan[x/Sqrt[3]])/Sqrt[3] + Log[x] - Log[3 + x^2]/2`

3.202.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2026, 523, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{2x+3}{x^3+3x} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{2x+3}{x(x^2+3)} dx \\ & \quad \downarrow \text{523} \\ & \int \left(\frac{2-x}{x^2+3} + \frac{1}{x} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{2 \arctan\left(\frac{x}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{2} \log(x^2+3) + \log(x) \end{aligned}$$

input `Int[(3 + 2*x)/(3*x + x^3),x]`

output `(2*ArcTan[x/Sqrt[3]])/Sqrt[3] + Log[x] - Log[3 + x^2]/2`

3.202.3.1 Defintions of rubi rules used

rule 523 `Int[((x_)^(m_.)*((c_) + (d_.)*(x_)))/((a_) + (b_.)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[x^m*((c + d*x)/(a + b*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && IntegerQ[m]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(F_x_.)*(P_x_)^(p_.), x_Symbol] := With[{r = Expon[P_x, x, Min]}, Int[x^(p*r)*ExpandToSum[P_x/x^r, x]^p*F_x, x] /; IGtQ[r, 0] /; PolyQ[P_x, x] && IntegerQ[p] && !MonomialQ[P_x, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

3.202.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

method	result	size
default	$\ln(x) - \frac{\ln(x^2+3)}{2} + \frac{2 \arctan\left(\frac{x\sqrt{3}}{3}\right)\sqrt{3}}{3}$	24
risch	$\ln(x) - \frac{\ln(x^2+3)}{2} + \frac{2 \arctan\left(\frac{x\sqrt{3}}{3}\right)\sqrt{3}}{3}$	24
meijerg	$\ln(x) - \frac{\ln(3)}{2} - \frac{\ln\left(\frac{x^2}{3}+1\right)}{2} + \frac{2 \arctan\left(\frac{x\sqrt{3}}{3}\right)\sqrt{3}}{3}$	30

input `int((3+2*x)/(x^3+3*x),x,method=_RETURNVERBOSE)`output `ln(x)-1/2*ln(x^2+3)+2/3*arctan(1/3*x*3^(1/2))*3^(1/2)`**3.202.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

$$\int \frac{3+2x}{3x+x^3} dx = \frac{2}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}x\right) - \frac{1}{2} \log(x^2+3) + \log(x)$$

input `integrate((3+2*x)/(x^3+3*x),x, algorithm="fracas")`output `2/3*sqrt(3)*arctan(1/3*sqrt(3)*x) - 1/2*log(x^2 + 3) + log(x)`**3.202.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{3+2x}{3x+x^3} dx = \log(x) - \frac{\log(x^2+3)}{2} + \frac{2\sqrt{3} \operatorname{atan}\left(\frac{\sqrt{3}x}{3}\right)}{3}$$

input `integrate((3+2*x)/(x**3+3*x),x)`output `log(x) - log(x**2 + 3)/2 + 2*sqrt(3)*atan(sqrt(3)*x/3)/3`

3.202.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

$$\int \frac{3+2x}{3x+x^3} dx = \frac{2}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}x\right) - \frac{1}{2} \log(x^2+3) + \log(x)$$

input `integrate((3+2*x)/(x^3+3*x),x, algorithm="maxima")`output `2/3*sqrt(3)*arctan(1/3*sqrt(3)*x) - 1/2*log(x^2 + 3) + log(x)`**3.202.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int \frac{3+2x}{3x+x^3} dx = \frac{2}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}x\right) - \frac{1}{2} \log(x^2+3) + \log(|x|)$$

input `integrate((3+2*x)/(x^3+3*x),x, algorithm="giac")`output `2/3*sqrt(3)*arctan(1/3*sqrt(3)*x) - 1/2*log(x^2 + 3) + log(abs(x))`**3.202.9 Mupad [B] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.96

$$\int \frac{3+2x}{3x+x^3} dx = \ln(x) - \frac{\ln(x + \sqrt{3} \text{li})}{2} - \frac{\ln(x - \sqrt{3} \text{li})}{2} - \frac{\sqrt{3} \ln(x - \sqrt{3} \text{li}) \text{li}}{3} + \frac{\sqrt{3} \ln(x + \sqrt{3} \text{li}) \text{li}}{3}$$

input `int((2*x + 3)/(3*x + x^3),x)`output `log(x) - log(x + 3^(1/2)*1i)/2 - log(x - 3^(1/2)*1i)/2 - (3^(1/2)*log(x - 3^(1/2)*1i)*1i)/3 + (3^(1/2)*log(x + 3^(1/2)*1i)*1i)/3`

3.203 $\int \frac{1}{-1+x^3} dx$

3.203.1 Optimal result	1053
3.203.2 Mathematica [A] (verified)	1053
3.203.3 Rubi [A] (verified)	1054
3.203.4 Maple [A] (verified)	1056
3.203.5 Fricas [A] (verification not implemented)	1056
3.203.6 Sympy [A] (verification not implemented)	1056
3.203.7 Maxima [A] (verification not implemented)	1057
3.203.8 Giac [A] (verification not implemented)	1057
3.203.9 Mupad [B] (verification not implemented)	1058

3.203.1 Optimal result

Integrand size = 7, antiderivative size = 41

$$\int \frac{1}{-1+x^3} dx = -\frac{\arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{3} \log(1-x) - \frac{1}{6} \log(1+x+x^2)$$

output `1/3*ln(1-x)-1/6*ln(x^2+x+1)-1/3*arctan(1/3*(1+2*x)*3^(1/2))*3^(1/2)`

3.203.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{1}{-1+x^3} dx = -\frac{\arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{3} \log(1-x) - \frac{1}{6} \log(1+x+x^2)$$

input `Integrate[(-1 + x^3)^(-1),x]`

output `-(ArcTan[(1 + 2*x)/Sqrt[3]]/Sqrt[3]) + Log[1 - x]/3 - Log[1 + x + x^2]/6`

3.203.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.12, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {750, 16, 25, 1142, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^3 - 1} dx \\
 & \quad \downarrow \text{750} \\
 & \frac{1}{3} \int -\frac{x+2}{x^2+x+1} dx + \frac{1}{3} \int \frac{1}{x-1} dx \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{3} \int -\frac{x+2}{x^2+x+1} dx + \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{3} \log(1-x) - \frac{1}{3} \int \frac{x+2}{x^2+x+1} dx \\
 & \quad \downarrow \text{1142} \\
 & \frac{1}{3} \left(-\frac{3}{2} \int \frac{1}{x^2+x+1} dx - \frac{1}{2} \int \frac{2x+1}{x^2+x+1} dx \right) + \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{1083} \\
 & \frac{1}{3} \left(3 \int \frac{1}{-(2x+1)^2-3} d(2x+1) - \frac{1}{2} \int \frac{2x+1}{x^2+x+1} dx \right) + \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{217} \\
 & \frac{1}{3} \left(-\frac{1}{2} \int \frac{2x+1}{x^2+x+1} dx - \sqrt{3} \arctan \left(\frac{2x+1}{\sqrt{3}} \right) \right) + \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{1103} \\
 & \frac{1}{3} \left(-\sqrt{3} \arctan \left(\frac{2x+1}{\sqrt{3}} \right) - \frac{1}{2} \log(x^2+x+1) \right) + \frac{1}{3} \log(1-x)
 \end{aligned}$$

input `Int[(-1 + x^3)^(-1), x]`

output $\text{Log}[1 - x]/3 + (-\text{Sqrt}[3]*\text{ArcTan}[(1 + 2*x)/\text{Sqrt}[3]]) - \text{Log}[1 + x + x^2]/2) / 3$

3.203.3.1 Defintions of rubi rules used

rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /;$ $\text{FreeQ}[\{a, b, c\}, x]$

rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{Int}[Fx, x], x]$

rule 217 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1} * \text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /;$ $\text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 750 $\text{Int}[(a_)+(b_)*(x_)^3)^{-1}, x_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /;$ $\text{FreeQ}[\{a, b\}, x]$

rule 1083 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[-2 \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /;$ $\text{FreeQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /;$ $\text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$

rule 1142 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /;$ $\text{FreeQ}[\{a, b, c, d, e\}, x]$

3.203.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

method	result	size
risch	$\frac{\ln(-1+x)}{3} - \frac{\ln(x^2+x+1)}{6} - \frac{\sqrt{3} \arctan\left(\frac{2\sqrt{3}\left(x+\frac{1}{2}\right)}{3}\right)}{3}$	31
default	$\frac{\ln(-1+x)}{3} - \frac{\ln(x^2+x+1)}{6} - \frac{\arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3}$	33
meijerg	$\frac{x \left(\ln\left(1-(x^3)^{\frac{1}{3}}\right) - \frac{\ln\left(1+(x^3)^{\frac{1}{3}}+(x^3)^{\frac{2}{3}}\right)}{2} - \sqrt{3} \arctan\left(\frac{\sqrt{3}(x^3)^{\frac{1}{3}}}{2+(x^3)^{\frac{1}{3}}}\right) \right)}{3(x^3)^{\frac{1}{3}}}$	62

input `int(1/(x^3-1),x,method=_RETURNVERBOSE)`output `1/3*ln(-1+x)-1/6*ln(x^2+x+1)-1/3*3^(1/2)*arctan(2/3*3^(1/2)*(x+1/2))`**3.203.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.78

$$\int \frac{1}{-1+x^3} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) - \frac{1}{6} \log(x^2+x+1) + \frac{1}{3} \log(x-1)$$

input `integrate(1/(x^3-1),x, algorithm="fracas")`output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) - 1/6*log(x^2 + x + 1) + 1/3*log(x - 1)`**3.203.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{1}{-1+x^3} dx = \frac{\log(x-1)}{3} - \frac{\log(x^2+x+1)}{6} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x}{3} + \frac{\sqrt{3}}{3}\right)}{3}$$

input `integrate(1/(x**3-1),x)`

output `log(x - 1)/3 - log(x**2 + x + 1)/6 - sqrt(3)*atan(2*sqrt(3)*x/3 + sqrt(3)/3)/3`

3.203.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.78

$$\int \frac{1}{-1+x^3} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) - \frac{1}{6} \log(x^2+x+1) + \frac{1}{3} \log(x-1)$$

input `integrate(1/(x^3-1),x, algorithm="maxima")`

output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) - 1/6*log(x^2 + x + 1) + 1/3*log(x - 1)`

3.203.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.80

$$\int \frac{1}{-1+x^3} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) - \frac{1}{6} \log(x^2+x+1) + \frac{1}{3} \log(|x-1|)$$

input `integrate(1/(x^3-1),x, algorithm="giac")`

output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) - 1/6*log(x^2 + x + 1) + 1/3*log(abs(x - 1))`

3.203.9 Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.12

$$\int \frac{1}{-1+x^3} dx = \frac{\ln(x-1)}{3} + \ln\left(x + \frac{1}{2} - \frac{\sqrt{3}1i}{2}\right) \left(-\frac{1}{6} + \frac{\sqrt{3}1i}{6}\right) - \ln\left(x + \frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) \left(\frac{1}{6} + \frac{\sqrt{3}1i}{6}\right)$$

input `int(1/(x^3 - 1),x)`output `log(x - 1)/3 + log(x - (3^(1/2)*1i)/2 + 1/2)*((3^(1/2)*1i)/6 - 1/6) - log(x + (3^(1/2)*1i)/2 + 1/2)*((3^(1/2)*1i)/6 + 1/6)`

3.204 $\int \frac{x^3}{1+x^3} dx$

3.204.1 Optimal result	1059
3.204.2 Mathematica [A] (verified)	1059
3.204.3 Rubi [A] (verified)	1060
3.204.4 Maple [A] (verified)	1062
3.204.5 Fricas [A] (verification not implemented)	1062
3.204.6 Sympy [A] (verification not implemented)	1063
3.204.7 Maxima [A] (verification not implemented)	1063
3.204.8 Giac [A] (verification not implemented)	1063
3.204.9 Mupad [B] (verification not implemented)	1064

3.204.1 Optimal result

Integrand size = 11, antiderivative size = 41

$$\int \frac{x^3}{1+x^3} dx = x + \frac{\arctan\left(\frac{1-2x}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{3} \log(1+x) + \frac{1}{6} \log(1-x+x^2)$$

output `x-1/3*ln(1+x)+1/6*ln(x^2-x+1)+1/3*arctan(1/3*(1-2*x)*3^(1/2))*3^(1/2)`

3.204.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.02

$$\int \frac{x^3}{1+x^3} dx = x - \frac{\arctan\left(\frac{-1+2x}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{3} \log(1+x) + \frac{1}{6} \log(1-x+x^2)$$

input `Integrate[x^3/(1 + x^3),x]`

output `x - ArcTan[(-1 + 2*x)/Sqrt[3]]/Sqrt[3] - Log[1 + x]/3 + Log[1 - x + x^2]/6`

3.204.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.15, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$, Rules used = {843, 750, 16, 1142, 25, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3}{x^3+1} dx \\
 & \quad \downarrow \text{843} \\
 & x - \int \frac{1}{x^3+1} dx \\
 & \quad \downarrow \text{750} \\
 & -\frac{1}{3} \int \frac{2-x}{x^2-x+1} dx - \frac{1}{3} \int \frac{1}{x+1} dx + x \\
 & \quad \downarrow \text{16} \\
 & -\frac{1}{3} \int \frac{2-x}{x^2-x+1} dx + x - \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{1142} \\
 & \frac{1}{3} \left(\frac{1}{2} \int -\frac{1-2x}{x^2-x+1} dx - \frac{3}{2} \int \frac{1}{x^2-x+1} dx \right) + x - \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{3} \left(-\frac{3}{2} \int \frac{1}{x^2-x+1} dx - \frac{1}{2} \int \frac{1-2x}{x^2-x+1} dx \right) + x - \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{1083} \\
 & \frac{1}{3} \left(3 \int \frac{1}{-(2x-1)^2-3} d(2x-1) - \frac{1}{2} \int \frac{1-2x}{x^2-x+1} dx \right) + x - \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{217} \\
 & \frac{1}{3} \left(-\frac{1}{2} \int \frac{1-2x}{x^2-x+1} dx - \sqrt{3} \arctan \left(\frac{2x-1}{\sqrt{3}} \right) \right) + x - \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{1103} \\
 & \frac{1}{3} \left(\frac{1}{2} \log(x^2-x+1) - \sqrt{3} \arctan \left(\frac{2x-1}{\sqrt{3}} \right) \right) + x - \frac{1}{3} \log(x+1)
 \end{aligned}$$

input `Int[x^3/(1 + x^3),x]`

output `x - Log[1 + x]/3 + (- (Sqrt[3]*ArcTan[(-1 + 2*x)/Sqrt[3]]) + Log[1 - x + x^2]/2)/3`

3.204.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(- (Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 750 `Int[((a_) + (b_.)*(x_)^3)^(-1), x_Symbol] := Simp[1/(3*Rt[a, 3]^2) Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Simp[1/(3*Rt[a, 3]^2) Int[(2*Rt[a, 3] - Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]`

rule 843 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Simp[a*c^n*((m - n + 1)/(b*(m + n*p + 1))) Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

```
rule 1142 Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c)
Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]
```

3.204.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

method	result	size
risch	$x - \frac{\ln(1+x)}{3} + \frac{\ln(x^2-x+1)}{6} - \frac{\sqrt{3} \arctan\left(\frac{2(x-\frac{1}{2})\sqrt{3}}{3}\right)}{3}$	34
default	$x + \frac{\ln(x^2-x+1)}{6} - \frac{\sqrt{3} \arctan\left(\frac{(2x-1)\sqrt{3}}{3}\right)}{3} - \frac{\ln(1+x)}{3}$	36
meijerg	$x - \frac{\left(\frac{\ln\left(1+(x^3)^{\frac{1}{3}}\right)}{(x^3)^{\frac{1}{3}}} - \frac{\ln\left(1-(x^3)^{\frac{1}{3}}+(x^3)^{\frac{2}{3}}\right)}{2(x^3)^{\frac{1}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}(x^3)^{\frac{1}{3}}}{2-(x^3)^{\frac{1}{3}}}\right)}{(x^3)^{\frac{1}{3}}} \right)}{3}$	74

```
input int(x^3/(x^3+1),x,method=_RETURNVERBOSE)
```

```
output x-1/3*ln(1+x)+1/6*ln(x^2-x+1)-1/3*3^(1/2)*arctan(2/3*(x-1/2)*3^(1/2))
```

3.204.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.85

$$\int \frac{x^3}{1+x^3} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x-1)\right) + x + \frac{1}{6} \log(x^2-x+1) - \frac{1}{3} \log(x+1)$$

```
input integrate(x^3/(x^3+1),x, algorithm="fricas")
```

```
output -1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) + x + 1/6*log(x^2 - x + 1) - 1/
3*log(x + 1)
```

3.204.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.02

$$\int \frac{x^3}{1+x^3} dx = x - \frac{\log(x+1)}{3} + \frac{\log(x^2-x+1)}{6} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x}{3} - \frac{\sqrt{3}}{3}\right)}{3}$$

input `integrate(x**3/(x**3+1),x)`output `x - log(x + 1)/3 + log(x**2 - x + 1)/6 - sqrt(3)*atan(2*sqrt(3)*x/3 - sqrt(3)/3)/3`**3.204.7 Maxima [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.85

$$\int \frac{x^3}{1+x^3} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x-1)\right) + x + \frac{1}{6} \log(x^2-x+1) - \frac{1}{3} \log(x+1)$$

input `integrate(x^3/(x^3+1),x, algorithm="maxima")`output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) + x + 1/6*log(x^2 - x + 1) - 1/3*log(x + 1)`**3.204.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.88

$$\int \frac{x^3}{1+x^3} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x-1)\right) + x + \frac{1}{6} \log(x^2-x+1) - \frac{1}{3} \log(|x+1|)$$

input `integrate(x^3/(x^3+1),x, algorithm="giac")`output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) + x + 1/6*log(x^2 - x + 1) - 1/3*log(abs(x + 1))`

3.204.9 Mupad [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.15

$$\int \frac{x^3}{1+x^3} dx = x - \frac{\ln(x+1)}{3} + \ln\left(x - \frac{1}{2} - \frac{\sqrt{3}1i}{2}\right) \left(\frac{1}{6} + \frac{\sqrt{3}1i}{6}\right) - \ln\left(x - \frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) \left(-\frac{1}{6} + \frac{\sqrt{3}1i}{6}\right)$$

input `int(x^3/(x^3 + 1),x)`output `x - log(x + 1)/3 + log(x - (3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/6 + 1/6) - log(x + (3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/6 - 1/6)`

3.205 $\int \frac{-1-2x+x^2}{(-1+x)^2(1+x^2)} dx$

3.205.1 Optimal result	1065
3.205.2 Mathematica [A] (verified)	1065
3.205.3 Rubi [A] (verified)	1066
3.205.4 Maple [A] (verified)	1067
3.205.5 Fricas [A] (verification not implemented)	1067
3.205.6 Sympy [A] (verification not implemented)	1067
3.205.7 Maxima [A] (verification not implemented)	1068
3.205.8 Giac [B] (verification not implemented)	1068
3.205.9 Mupad [B] (verification not implemented)	1068

3.205.1 Optimal result

Integrand size = 21, antiderivative size = 24

$$\int \frac{-1-2x+x^2}{(-1+x)^2(1+x^2)} dx = \frac{1}{-1+x} + \arctan(x) + \log(1-x) - \frac{1}{2} \log(1+x^2)$$

output `1/(-1+x)+arctan(x)+ln(1-x)-1/2*ln(x^2+1)`

3.205.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{-1-2x+x^2}{(-1+x)^2(1+x^2)} dx = \frac{1}{-1+x} + \arctan(x) + \log(-1+x) - \frac{1}{2} \log(1+x^2)$$

input `Integrate[(-1 - 2*x + x^2)/((-1 + x)^2*(1 + x^2)),x]`

output `(-1 + x)^(-1) + ArcTan[x] + Log[-1 + x] - Log[1 + x^2]/2`

3.205.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2160, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2 - 2x - 1}{(x-1)^2(x^2+1)} dx$$

↓ 2160

$$\int \left(\frac{1-x}{x^2+1} + \frac{1}{x-1} - \frac{1}{(x-1)^2} \right) dx$$

↓ 2009

$$\arctan(x) - \frac{1}{2} \log(x^2+1) + \frac{1}{x-1} + \log(1-x)$$

input `Int[(-1 - 2*x + x^2)/((-1 + x)^2*(1 + x^2)),x]`

output `(-1 + x)^(-1) + ArcTan[x] + Log[1 - x] - Log[1 + x^2]/2`

3.205.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2160 `Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

3.205.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

method	result	size
default	$\ln(-1+x) + \frac{1}{-1+x} - \frac{\ln(x^2+1)}{2} + \arctan(x)$	21
risch	$\ln(-1+x) + \frac{1}{-1+x} - \frac{\ln(x^2+1)}{2} + \arctan(x)$	21
parallelrisch	$\frac{-i \ln(x-i)x+i \ln(x+i)x+2 \ln(-1+x)x+i \ln(x-i)-\ln(x-i)x-i \ln(x+i)-\ln(x+i)x+2-2 \ln(-1+x)+\ln(x-i)+\ln(x+i)}{-2+2x}$	83

input `int((x^2-2*x-1)/(-1+x)^2/(x^2+1),x,method=_RETURNVERBOSE)`output `ln(-1+x)+1/(-1+x)-1/2*ln(x^2+1)+arctan(x)`**3.205.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.50

$$\int \frac{-1-2x+x^2}{(-1+x)^2(1+x^2)} dx$$

$$= \frac{2(x-1)\arctan(x) - (x-1)\log(x^2+1) + 2(x-1)\log(x-1) + 2}{2(x-1)}$$

input `integrate((x^2-2*x-1)/(-1+x)^2/(x^2+1),x, algorithm="fricas")`output `1/2*(2*(x - 1)*arctan(x) - (x - 1)*log(x^2 + 1) + 2*(x - 1)*log(x - 1) + 2)/(x - 1)`**3.205.6 Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{-1-2x+x^2}{(-1+x)^2(1+x^2)} dx = \log(x-1) - \frac{\log(x^2+1)}{2} + \operatorname{atan}(x) + \frac{1}{x-1}$$

input `integrate((x**2-2*x-1)/(-1+x)**2/(x**2+1),x)`output `log(x - 1) - log(x**2 + 1)/2 + atan(x) + 1/(x - 1)`

3.205. $\int \frac{-1-2x+x^2}{(-1+x)^2(1+x^2)} dx$

3.205.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{-1 - 2x + x^2}{(-1 + x)^2 (1 + x^2)} dx = \frac{1}{x - 1} + \arctan(x) - \frac{1}{2} \log(x^2 + 1) + \log(x - 1)$$

input `integrate((x^2-2*x-1)/(-1+x)^2/(x^2+1),x, algorithm="maxima")`

output `1/(x - 1) + arctan(x) - 1/2*log(x^2 + 1) + log(x - 1)`

3.205.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 47 vs. 2(22) = 44.

Time = 0.29 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.96

$$\int \frac{-1 - 2x + x^2}{(-1 + x)^2 (1 + x^2)} dx = \frac{1}{4} \pi - \pi \left[\frac{\pi + 4 \arctan(x)}{4 \pi} + \frac{1}{2} \right] + \frac{1}{x - 1} + \arctan(x) - \frac{1}{2} \log \left(\frac{2}{x - 1} + \frac{2}{(x - 1)^2} + 1 \right)$$

input `integrate((x^2-2*x-1)/(-1+x)^2/(x^2+1),x, algorithm="giac")`

output `1/4*pi - pi*floor(1/4*(pi + 4*arctan(x))/pi + 1/2) + 1/(x - 1) + arctan(x) - 1/2*log(2/(x - 1) + 2/(x - 1)^2 + 1)`

3.205.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17

$$\int \frac{-1 - 2x + x^2}{(-1 + x)^2 (1 + x^2)} dx = \ln(x - 1) + \frac{1}{x - 1} + \ln(x - i) \left(-\frac{1}{2} - \frac{1}{2}i \right) + \ln(x + i) \left(-\frac{1}{2} + \frac{1}{2}i \right)$$

input `int(-(2*x - x^2 + 1)/((x^2 + 1)*(x - 1)^2),x)`

output `log(x - 1) - log(x - 1i)*(1/2 + 1i/2) - log(x + 1i)*(1/2 - 1i/2) + 1/(x - 1)`

3.205. $\int \frac{-1-2x+x^2}{(-1+x)^2(1+x^2)} dx$

3.206 $\int \frac{x^4}{-1+x^4} dx$

3.206.1 Optimal result	1069
3.206.2 Mathematica [A] (verified)	1069
3.206.3 Rubi [A] (verified)	1070
3.206.4 Maple [A] (verified)	1071
3.206.5 Fricas [A] (verification not implemented)	1071
3.206.6 Sympy [A] (verification not implemented)	1072
3.206.7 Maxima [A] (verification not implemented)	1072
3.206.8 Giac [A] (verification not implemented)	1072
3.206.9 Mupad [B] (verification not implemented)	1073

3.206.1 Optimal result

Integrand size = 11, antiderivative size = 14

$$\int \frac{x^4}{-1+x^4} dx = x - \frac{\arctan(x)}{2} - \frac{\operatorname{arctanh}(x)}{2}$$

output `x-1/2*arctan(x)-1/2*arctanh(x)`

3.206.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.86

$$\int \frac{x^4}{-1+x^4} dx = x - \frac{\arctan(x)}{2} + \frac{1}{4} \log(1-x) - \frac{1}{4} \log(1+x)$$

input `Integrate[x^4/(-1 + x^4),x]`

output `x - ArcTan[x]/2 + Log[1 - x]/4 - Log[1 + x]/4`

3.206.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {843, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4}{x^4 - 1} dx \\
 & \quad \downarrow \text{843} \\
 & \int \frac{1}{x^4 - 1} dx + x \\
 & \quad \downarrow \text{756} \\
 & -\frac{1}{2} \int \frac{1}{1 - x^2} dx - \frac{1}{2} \int \frac{1}{x^2 + 1} dx + x \\
 & \quad \downarrow \text{216} \\
 & -\frac{1}{2} \int \frac{1}{1 - x^2} dx - \frac{\arctan(x)}{2} + x \\
 & \quad \downarrow \text{219} \\
 & -\frac{\arctan(x)}{2} - \frac{\operatorname{arctanh}(x)}{2} + x
 \end{aligned}$$

input `Int[x^4/(-1 + x^4),x]`

output `x - ArcTan[x]/2 - ArcTanh[x]/2`

3.206.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 756 `Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 843 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n-1)*(c*x)^(m-n+1)*((a+b*x^n)^(p+1)/(b*(m+n*p+1))), x] - Simp[a*c^n*(m-n+1)/(b*(m+n*p+1)) Int[(c*x)^(m-n)*(a+b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n-1] && NeQ[m+n*p+1, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

3.206.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.36

method	result	size
default	$x + \frac{\ln(-1+x)}{4} - \frac{\ln(1+x)}{4} - \frac{\arctan(x)}{2}$	19
risch	$x + \frac{\ln(-1+x)}{4} - \frac{\ln(1+x)}{4} - \frac{\arctan(x)}{2}$	19
parallelrisc	$x + \frac{i \ln(x-i)}{4} - \frac{i \ln(x+i)}{4} - \frac{\ln(1+x)}{4} + \frac{\ln(-1+x)}{4}$	31
meijerg	$- \frac{(-1)^{\frac{3}{4}} \left(4(-1)^{\frac{1}{4}} x + \frac{x(-1)^{\frac{1}{4}} \left(\ln \left(1 - (x^4)^{\frac{1}{4}} \right) - \ln \left(1 + (x^4)^{\frac{1}{4}} \right) - 2 \arctan \left((x^4)^{\frac{1}{4}} \right) \right)}{(x^4)^{\frac{1}{4}}} \right)}{4}$	52

input `int(x^4/(x^4-1),x,method=_RETURNVERBOSE)`

output `x+1/4*ln(-1+x)-1/4*ln(1+x)-1/2*arctan(x)`

3.206.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.29

$$\int \frac{x^4}{-1+x^4} dx = x - \frac{1}{2} \arctan(x) - \frac{1}{4} \log(x+1) + \frac{1}{4} \log(x-1)$$

input `integrate(x^4/(x^4-1),x, algorithm="fricas")`

output $x - 1/2*\arctan(x) - 1/4*\log(x + 1) + 1/4*\log(x - 1)$

3.206.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.36

$$\int \frac{x^4}{-1+x^4} dx = x + \frac{\log(x-1)}{4} - \frac{\log(x+1)}{4} - \frac{\operatorname{atan}(x)}{2}$$

input `integrate(x**4/(x**4-1),x)`

output $x + \log(x - 1)/4 - \log(x + 1)/4 - \operatorname{atan}(x)/2$

3.206.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.29

$$\int \frac{x^4}{-1+x^4} dx = x - \frac{1}{2} \arctan(x) - \frac{1}{4} \log(x+1) + \frac{1}{4} \log(x-1)$$

input `integrate(x^4/(x^4-1),x, algorithm="maxima")`

output $x - 1/2*\arctan(x) - 1/4*\log(x + 1) + 1/4*\log(x - 1)$

3.206.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.43

$$\int \frac{x^4}{-1+x^4} dx = x - \frac{1}{2} \arctan(x) - \frac{1}{4} \log(|x+1|) + \frac{1}{4} \log(|x-1|)$$

input `integrate(x^4/(x^4-1),x, algorithm="giac")`

output $x - 1/2*\arctan(x) - 1/4*\log(\operatorname{abs}(x + 1)) + 1/4*\log(\operatorname{abs}(x - 1))$

3.206.9 Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{x^4}{-1+x^4} dx = x - \frac{\operatorname{atan}(x)}{2} - \frac{\operatorname{atanh}(x)}{2}$$

input `int(x^4/(x^4 - 1),x)`

output `x - atan(x)/2 - atanh(x)/2`

$$3.207 \quad \int \frac{-4+6x-x^2+3x^3}{(1+x^2)(2+x^2)} dx$$

3.207.1 Optimal result	1074
3.207.2 Mathematica [A] (verified)	1074
3.207.3 Rubi [A] (verified)	1075
3.207.4 Maple [A] (verified)	1076
3.207.5 Fricas [A] (verification not implemented)	1076
3.207.6 Sympy [A] (verification not implemented)	1076
3.207.7 Maxima [A] (verification not implemented)	1077
3.207.8 Giac [A] (verification not implemented)	1077
3.207.9 Mupad [B] (verification not implemented)	1077

3.207.1 Optimal result

Integrand size = 30, antiderivative size = 29

$$\int \frac{-4 + 6x - x^2 + 3x^3}{(1 + x^2)(2 + x^2)} dx = -3 \arctan(x) + \sqrt{2} \arctan\left(\frac{x}{\sqrt{2}}\right) + \frac{3}{2} \log(1 + x^2)$$

output `-3*arctan(x)+3/2*ln(x^2+1)+arctan(1/2*x*2^(1/2))*2^(1/2)`

3.207.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{-4 + 6x - x^2 + 3x^3}{(1 + x^2)(2 + x^2)} dx = -3 \arctan(x) + \sqrt{2} \arctan\left(\frac{x}{\sqrt{2}}\right) + \frac{3}{2} \log(1 + x^2)$$

input `Integrate[(-4 + 6*x - x^2 + 3*x^3)/((1 + x^2)*(2 + x^2)),x]`

output `-3*ArcTan[x] + Sqrt[2]*ArcTan[x/Sqrt[2]] + (3*Log[1 + x^2])/2`

3.207.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{3x^3 - x^2 + 6x - 4}{(x^2 + 1)(x^2 + 2)} dx$$

↓ 7276

$$\int \left(\frac{3(x-1)}{x^2+1} + \frac{2}{x^2+2} \right) dx$$

↓ 2009

$$-3 \arctan(x) + \sqrt{2} \arctan\left(\frac{x}{\sqrt{2}}\right) + \frac{3}{2} \log(x^2 + 1)$$

input `Int[(-4 + 6*x - x^2 + 3*x^3)/((1 + x^2)*(2 + x^2)),x]`

output `-3*ArcTan[x] + Sqrt[2]*ArcTan[x/Sqrt[2]] + (3*Log[1 + x^2])/2`

3.207.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
x
pand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]`

3.207.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.86

method	result	size
default	$-3 \arctan(x) + \frac{3 \ln(x^2+1)}{2} + \arctan\left(\frac{x\sqrt{2}}{2}\right) \sqrt{2}$	25
risch	$-3 \arctan(x) + \frac{3 \ln(x^2+1)}{2} + \arctan\left(\frac{x\sqrt{2}}{2}\right) \sqrt{2}$	25

input `int((3*x^3-x^2+6*x-4)/(x^2+1)/(x^2+2),x,method=_RETURNVERBOSE)`output `-3*arctan(x)+3/2*ln(x^2+1)+arctan(1/2*x*2^(1/2))*2^(1/2)`**3.207.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

$$\int \frac{-4 + 6x - x^2 + 3x^3}{(1 + x^2)(2 + x^2)} dx = \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}x\right) - 3 \arctan(x) + \frac{3}{2} \log(x^2 + 1)$$

input `integrate((3*x^3-x^2+6*x-4)/(x^2+1)/(x^2+2),x, algorithm="fricas")`output `sqrt(2)*arctan(1/2*sqrt(2)*x) - 3*arctan(x) + 3/2*log(x^2 + 1)`**3.207.6 Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{-4 + 6x - x^2 + 3x^3}{(1 + x^2)(2 + x^2)} dx = \frac{3 \log(x^2 + 1)}{2} - 3 \operatorname{atan}(x) + \sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}x}{2}\right)$$

input `integrate((3*x**3-x**2+6*x-4)/(x**2+1)/(x**2+2),x)`output `3*log(x**2 + 1)/2 - 3*atan(x) + sqrt(2)*atan(sqrt(2)*x/2)`

3.207.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

$$\int \frac{-4 + 6x - x^2 + 3x^3}{(1 + x^2)(2 + x^2)} dx = \sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}x\right) - 3 \arctan(x) + \frac{3}{2} \log(x^2 + 1)$$

input `integrate((3*x^3-x^2+6*x-4)/(x^2+1)/(x^2+2),x, algorithm="maxima")`output `sqrt(2)*arctan(1/2*sqrt(2)*x) - 3*arctan(x) + 3/2*log(x^2 + 1)`**3.207.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

$$\int \frac{-4 + 6x - x^2 + 3x^3}{(1 + x^2)(2 + x^2)} dx = \sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}x\right) - 3 \arctan(x) + \frac{3}{2} \log(x^2 + 1)$$

input `integrate((3*x^3-x^2+6*x-4)/(x^2+1)/(x^2+2),x, algorithm="giac")`output `sqrt(2)*arctan(1/2*sqrt(2)*x) - 3*arctan(x) + 3/2*log(x^2 + 1)`**3.207.9 Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.76

$$\int \frac{-4 + 6x - x^2 + 3x^3}{(1 + x^2)(2 + x^2)} dx = -\sqrt{2} \operatorname{atan}\left(\frac{24\sqrt{2}}{24x - 64} + \frac{32\sqrt{2}x}{24x - 64}\right) + \ln(x - i) \left(\frac{3}{2} + \frac{3i}{2}\right) + \ln(x + i) \left(\frac{3}{2} - \frac{3i}{2}\right)$$

input `int((6*x - x^2 + 3*x^3 - 4)/((x^2 + 1)*(x^2 + 2)),x)`output `log(x - 1i)*(3/2 + 3i/2) + log(x + 1i)*(3/2 - 3i/2) - 2^(1/2)*atan((24*2^(1/2))/(24*x - 64) + (32*2^(1/2)*x)/(24*x - 64))`

$$3.208 \quad \int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx$$

3.208.1 Optimal result	1078
3.208.2 Mathematica [A] (verified)	1078
3.208.3 Rubi [A] (verified)	1079
3.208.4 Maple [A] (verified)	1080
3.208.5 Fricas [A] (verification not implemented)	1081
3.208.6 Sympy [A] (verification not implemented)	1081
3.208.7 Maxima [A] (verification not implemented)	1081
3.208.8 Giac [A] (verification not implemented)	1082
3.208.9 Mupad [B] (verification not implemented)	1082

3.208.1 Optimal result

Integrand size = 24, antiderivative size = 23

$$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx = -\frac{3}{2} \arctan\left(\frac{x}{2}\right) + \arctan(x) + \frac{1}{2} \log(4+x^2)$$

output `-3/2*arctan(1/2*x)+arctan(x)+1/2*ln(x^2+4)`

3.208.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx = -\frac{3}{2} \arctan\left(\frac{x}{2}\right) + \arctan(x) + \frac{1}{2} \log(4+x^2)$$

input `Integrate[(1 + x - 2*x^2 + x^3)/(4 + 5*x^2 + x^4),x]`

output `(-3*ArcTan[x/2])/2 + ArcTan[x] + Log[4 + x^2]/2`

3.208.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {2202, 1387, 240, 1480, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3 - 2x^2 + x + 1}{x^4 + 5x^2 + 4} dx \\
 & \quad \downarrow \text{2202} \\
 & \int \frac{1 - 2x^2}{x^4 + 5x^2 + 4} dx + \int \frac{x(x^2 + 1)}{x^4 + 5x^2 + 4} dx \\
 & \quad \downarrow \text{1387} \\
 & \int \frac{x}{x^2 + 4} dx + \int \frac{1 - 2x^2}{x^4 + 5x^2 + 4} dx \\
 & \quad \downarrow \text{240} \\
 & \int \frac{1 - 2x^2}{x^4 + 5x^2 + 4} dx + \frac{1}{2} \log(x^2 + 4) \\
 & \quad \downarrow \text{1480} \\
 & \int \frac{1}{x^2 + 1} dx - 3 \int \frac{1}{x^2 + 4} dx + \frac{1}{2} \log(x^2 + 4) \\
 & \quad \downarrow \text{216} \\
 & -\frac{3}{2} \arctan\left(\frac{x}{2}\right) + \arctan(x) + \frac{1}{2} \log(x^2 + 4)
 \end{aligned}$$

input `Int[(1 + x - 2*x^2 + x^3)/(4 + 5*x^2 + x^4), x]`

output `(-3*ArcTan[x/2])/2 + ArcTan[x] + Log[4 + x^2]/2`

3.208.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 240 `Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`

rule 1387 `Int[(u_.)*((a_) + (c_.)*(x_)^(n2_.) + (b_.)*(x_)^(n_.))^ (p_.)*((d_) + (e_.)*(x_)^(n_.))^ (q_.), x_Symbol] := Int[u*(d + e*x^n)^(p + q)*(a/d + (c/e)*x^n)^p, x] /; FreeQ[{a, b, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[c*d^2 - b*d*e + a*e^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[d, 0] && LtQ[c, 0]))`

rule 1480 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[(e/2 + (2*c*d - b*e)/(2*q)) Int[1/(b/2 - q/2 + c*x^2), x], x] + Simp[(e/2 - (2*c*d - b*e)/(2*q)) Int[1/(b/2 + q/2 + c*x^2), x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[b^2 - 4*a*c]`

rule 2202 `Int[(Pn_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^ (p_), x_Symbol] := Module[{n = Expon[Pn, x], k}, Int[Sum[Coeff[Pn, x, 2*k]*x^(2*k), {k, 0, n/2}]*(a + b*x^2 + c*x^4)^p, x] + Int[x*Sum[Coeff[Pn, x, 2*k + 1]*x^(2*k), {k, 0, (n - 1)/2}]*(a + b*x^2 + c*x^4)^p, x]] /; FreeQ[{a, b, c, p}, x] && PolyQ[Pn, x] && !PolyQ[Pn, x^2]`

3.208.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

method	result	size
default	$-\frac{3 \arctan\left(\frac{x}{2}\right)}{2} + \arctan(x) + \frac{\ln(x^2+4)}{2}$	18
risch	$-\frac{3 \arctan\left(\frac{x}{2}\right)}{2} + \arctan(x) + \frac{\ln(x^2+4)}{2}$	18
parallelrisch	$\frac{\ln(x-2i)}{2} + \frac{3i \ln(x-2i)}{4} - \frac{i \ln(x-i)}{2} + \frac{i \ln(x+i)}{2} + \frac{\ln(x+2i)}{2} - \frac{3i \ln(x+2i)}{4}$	48

input `int((x^3-2*x^2+x+1)/(x^4+5*x^2+4),x,method=_RETURNVERBOSE)`

output `-3/2*arctan(1/2*x)+arctan(x)+1/2*ln(x^2+4)`

3.208.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx = -\frac{3}{2} \arctan\left(\frac{1}{2}x\right) + \arctan(x) + \frac{1}{2} \log(x^2+4)$$

input `integrate((x^3-2*x^2+x+1)/(x^4+5*x^2+4),x, algorithm="fricas")`

output `-3/2*arctan(1/2*x) + arctan(x) + 1/2*log(x^2 + 4)`

3.208.6 Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx = \frac{\log(x^2+4)}{2} - \frac{3 \operatorname{atan}\left(\frac{x}{2}\right)}{2} + \operatorname{atan}(x)$$

input `integrate((x**3-2*x**2+x+1)/(x**4+5*x**2+4),x)`

output `log(x**2 + 4)/2 - 3*atan(x/2)/2 + atan(x)`

3.208.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx = -\frac{3}{2} \arctan\left(\frac{1}{2}x\right) + \arctan(x) + \frac{1}{2} \log(x^2+4)$$

input `integrate((x^3-2*x^2+x+1)/(x^4+5*x^2+4),x, algorithm="maxima")`

output `-3/2*arctan(1/2*x) + arctan(x) + 1/2*log(x^2 + 4)`

3.208.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx = -\frac{3}{2} \arctan\left(\frac{1}{2}x\right) + \arctan(x) + \frac{1}{2} \log(x^2+4)$$

input `integrate((x^3-2*x^2+x+1)/(x^4+5*x^2+4),x, algorithm="giac")`output `-3/2*arctan(1/2*x) + arctan(x) + 1/2*log(x^2 + 4)`**3.208.9 Mupad [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.43

$$\int \frac{1+x-2x^2+x^3}{4+5x^2+x^4} dx = -\operatorname{atan}\left(\frac{1305}{4(144x-162)} + \frac{9}{8}\right) + \ln(x-2i)\left(\frac{1}{2} + \frac{3i}{4}\right) + \ln(x+2i)\left(\frac{1}{2} - \frac{3i}{4}\right)$$

input `int((x - 2*x^2 + x^3 + 1)/(5*x^2 + x^4 + 4),x)`output `log(x - 2i)*(1/2 + 3i/4) + log(x + 2i)*(1/2 - 3i/4) - atan(1305/(4*(144*x - 162)) + 9/8)`

3.209 $\int \frac{-3+x}{(4+2x+x^2)^2} dx$

3.209.1 Optimal result 1083
 3.209.2 Mathematica [A] (verified) 1083
 3.209.3 Rubi [A] (verified) 1084
 3.209.4 Maple [A] (verified) 1085
 3.209.5 Fricas [A] (verification not implemented) 1085
 3.209.6 Sympy [A] (verification not implemented) 1086
 3.209.7 Maxima [A] (verification not implemented) 1086
 3.209.8 Giac [A] (verification not implemented) 1086
 3.209.9 Mupad [B] (verification not implemented) 1087

3.209.1 Optimal result

Integrand size = 14, antiderivative size = 39

$$\int \frac{-3+x}{(4+2x+x^2)^2} dx = \frac{-7-4x}{6(4+2x+x^2)} - \frac{2 \arctan\left(\frac{1+x}{\sqrt{3}}\right)}{3\sqrt{3}}$$

output `1/6*(-7-4*x)/(x^2+2*x+4)-2/9*arctan(1/3*(1+x)*3^(1/2))*3^(1/2)`

3.209.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

$$\int \frac{-3+x}{(4+2x+x^2)^2} dx = \frac{-7-4x}{6(4+2x+x^2)} - \frac{2 \arctan\left(\frac{1+x}{\sqrt{3}}\right)}{3\sqrt{3}}$$

input `Integrate[(-3 + x)/(4 + 2*x + x^2)^2,x]`

output `(-7 - 4*x)/(6*(4 + 2*x + x^2)) - (2*ArcTan[(1 + x)/Sqrt[3]])/(3*Sqrt[3])`

3.209.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.13, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {1159, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x-3}{(x^2+2x+4)^2} dx$$

$$\downarrow 1159$$

$$-\frac{2}{3} \int \frac{1}{x^2+2x+4} dx - \frac{4x+7}{6(x^2+2x+4)}$$

$$\downarrow 1083$$

$$\frac{4}{3} \int \frac{1}{-(2x+2)^2-12} d(2x+2) - \frac{4x+7}{6(x^2+2x+4)}$$

$$\downarrow 217$$

$$-\frac{2 \arctan\left(\frac{2x+2}{2\sqrt{3}}\right)}{3\sqrt{3}} - \frac{4x+7}{6(x^2+2x+4)}$$

input `Int[(-3 + x)/(4 + 2*x + x^2)^2,x]`

output `-1/6*(7 + 4*x)/(4 + 2*x + x^2) - (2*ArcTan[(2 + 2*x)/(2*sqrt[3])])/(3*sqrt[3])`

3.209.3.1 Defintions of rubi rules used

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])`

rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] :> Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

```
rule 1159 Int[((d_.) + (e_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol
] := Simp[((b*d - 2*a*e + (2*c*d - b*e)*x)/((p + 1)*(b^2 - 4*a*c)))*(a + b*
x + c*x^2)^(p + 1), x] - Simp[(2*p + 3)*((2*c*d - b*e)/((p + 1)*(b^2 - 4*a*
c))) Int[(a + b*x + c*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e}, x] &
& LtQ[p, -1] && NeQ[p, -3/2]
```

3.209.4 Maple [A] (verified)

Time = 0.63 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.82

method	result	size
risch	$\frac{-\frac{2x}{3} - \frac{7}{6}}{x^2 + 2x + 4} - \frac{2 \arctan\left(\frac{(1+x)\sqrt{3}}{3}\right)\sqrt{3}}{9}$	32
default	$\frac{-8x - 14}{12x^2 + 24x + 48} - \frac{2\sqrt{3} \arctan\left(\frac{(2x+2)\sqrt{3}}{6}\right)}{9}$	35

```
input int((-3+x)/(x^2+2*x+4)^2,x,method=_RETURNVERBOSE)
```

```
output (-2/3*x-7/6)/(x^2+2*x+4)-2/9*arctan(1/3*(1+x)*3^(1/2))*3^(1/2)
```

3.209.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

$$\int \frac{-3+x}{(4+2x+x^2)^2} dx = -\frac{4\sqrt{3}(x^2+2x+4)\arctan\left(\frac{1}{3}\sqrt{3}(x+1)\right)+12x+21}{18(x^2+2x+4)}$$

```
input integrate((-3+x)/(x^2+2*x+4)^2,x, algorithm="fracas")
```

```
output -1/18*(4*sqrt(3)*(x^2 + 2*x + 4)*arctan(1/3*sqrt(3)*(x + 1)) + 12*x + 21)/
(x^2 + 2*x + 4)
```

3.209.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.05

$$\int \frac{-3+x}{(4+2x+x^2)^2} dx = \frac{-4x-7}{6x^2+12x+24} - \frac{2\sqrt{3} \operatorname{atan}\left(\frac{\sqrt{3}x}{3} + \frac{\sqrt{3}}{3}\right)}{9}$$

input `integrate((-3+x)/(x**2+2*x+4)**2,x)`output `(-4*x - 7)/(6*x**2 + 12*x + 24) - 2*sqrt(3)*atan(sqrt(3)*x/3 + sqrt(3)/3)/9`**3.209.7 Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.82

$$\int \frac{-3+x}{(4+2x+x^2)^2} dx = -\frac{2}{9} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(x+1)\right) - \frac{4x+7}{6(x^2+2x+4)}$$

input `integrate((-3+x)/(x^2+2*x+4)^2,x, algorithm="maxima")`output `-2/9*sqrt(3)*arctan(1/3*sqrt(3)*(x + 1)) - 1/6*(4*x + 7)/(x^2 + 2*x + 4)`**3.209.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.82

$$\int \frac{-3+x}{(4+2x+x^2)^2} dx = -\frac{2}{9} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(x+1)\right) - \frac{4x+7}{6(x^2+2x+4)}$$

input `integrate((-3+x)/(x^2+2*x+4)^2,x, algorithm="giac")`output `-2/9*sqrt(3)*arctan(1/3*sqrt(3)*(x + 1)) - 1/6*(4*x + 7)/(x^2 + 2*x + 4)`

3.209.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.92

$$\int \frac{-3+x}{(4+2x+x^2)^2} dx = -\frac{\frac{2x}{3} + \frac{7}{6}}{x^2 + 2x + 4} - \frac{2\sqrt{3} \operatorname{atan}\left(\frac{\sqrt{3}x}{3} + \frac{\sqrt{3}}{3}\right)}{9}$$

input `int((x - 3)/(2*x + x^2 + 4)^2,x)`output `- ((2*x)/3 + 7/6)/(2*x + x^2 + 4) - (2*3^(1/2)*atan((3^(1/2)*x)/3 + 3^(1/2)/3))/9`

$$3.210 \quad \int \frac{1+x^4}{x(1+x^2)^2} dx$$

3.210.1 Optimal result	1088
3.210.2 Mathematica [A] (verified)	1088
3.210.3 Rubi [A] (verified)	1089
3.210.4 Maple [A] (verified)	1090
3.210.5 Fricas [A] (verification not implemented)	1090
3.210.6 Sympy [A] (verification not implemented)	1090
3.210.7 Maxima [A] (verification not implemented)	1091
3.210.8 Giac [A] (verification not implemented)	1091
3.210.9 Mupad [B] (verification not implemented)	1091

3.210.1 Optimal result

Integrand size = 16, antiderivative size = 10

$$\int \frac{1+x^4}{x(1+x^2)^2} dx = \frac{1}{1+x^2} + \log(x)$$

output `1/(x^2+1)+ln(x)`

3.210.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1+x^4}{x(1+x^2)^2} dx = \frac{1}{1+x^2} + \log(x)$$

input `Integrate[(1 + x^4)/(x*(1 + x^2)^2), x]`

output `(1 + x^2)^(-1) + Log[x]`

3.210.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.80, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {1579, 522, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^4 + 1}{x(x^2 + 1)^2} dx \\ & \quad \downarrow \text{1579} \\ & \frac{1}{2} \int \frac{x^4 + 1}{x^2(x^2 + 1)^2} dx^2 \\ & \quad \downarrow \text{522} \\ & \frac{1}{2} \int \left(\frac{1}{x^2} - \frac{2}{(x^2 + 1)^2} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \left(\frac{2}{x^2 + 1} + \log(x^2) \right) \end{aligned}$$

input `Int[(1 + x^4)/(x*(1 + x^2)^2), x]`

output `(2/(1 + x^2) + Log[x^2])/2`

3.210.3.1 Defintions of rubi rules used

rule 522 `Int[((e_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(e*x)^m*(c + d*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]`

rule 1579 `Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, c, d, e, p, q}, x] && IntegerQ[(m + 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.210. $\int \frac{1+x^4}{x(1+x^2)^2} dx$

3.210.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

method	result	size
default	$\frac{1}{x^2+1} + \ln(x)$	11
norman	$\frac{1}{x^2+1} + \ln(x)$	11
risch	$\frac{1}{x^2+1} + \ln(x)$	11
parallelrisch	$\frac{x^2 \ln(x)+1+\ln(x)}{x^2+1}$	19
meijerg	$-\frac{x^2}{2(x^2+1)} + \frac{1}{2} + \ln(x) - \frac{x^2}{2x^2+2}$	31

input `int((x^4+1)/x/(x^2+1)^2,x,method=_RETURNVERBOSE)`output `1/(x^2+1)+ln(x)`**3.210.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.80

$$\int \frac{1+x^4}{x(1+x^2)^2} dx = \frac{(x^2+1)\log(x)+1}{x^2+1}$$

input `integrate((x^4+1)/x/(x^2+1)^2,x, algorithm="fricas")`output `((x^2 + 1)*log(x) + 1)/(x^2 + 1)`**3.210.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{1+x^4}{x(1+x^2)^2} dx = \log(x) + \frac{1}{x^2+1}$$

input `integrate((x**4+1)/x/(x**2+1)**2,x)`output `log(x) + 1/(x**2 + 1)`

3.210. $\int \frac{1+x^4}{x(1+x^2)^2} dx$

3.210.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.40

$$\int \frac{1+x^4}{x(1+x^2)^2} dx = \frac{1}{x^2+1} + \frac{1}{2} \log(x^2)$$

input `integrate((x^4+1)/x/(x^2+1)^2,x, algorithm="maxima")`output `1/(x^2 + 1) + 1/2*log(x^2)`**3.210.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.40

$$\int \frac{1+x^4}{x(1+x^2)^2} dx = \frac{1}{x^2+1} + \frac{1}{2} \log(x^2)$$

input `integrate((x^4+1)/x/(x^2+1)^2,x, algorithm="giac")`output `1/(x^2 + 1) + 1/2*log(x^2)`**3.210.9 Mupad [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1+x^4}{x(1+x^2)^2} dx = \ln(x) + \frac{1}{x^2+1}$$

input `int((x^4 + 1)/(x*(x^2 + 1)^2),x)`output `log(x) + 1/(x^2 + 1)`

3.211 $\int \frac{\cos(x)(-3+2\sin(x))}{2-3\sin(x)+\sin^2(x)} dx$

3.211.1 Optimal result 1092
 3.211.2 Mathematica [A] (verified) 1092
 3.211.3 Rubi [A] (verified) 1093
 3.211.4 Maple [A] (verified) 1094
 3.211.5 Fricas [A] (verification not implemented) 1094
 3.211.6 Sympy [A] (verification not implemented) 1095
 3.211.7 Maxima [A] (verification not implemented) 1095
 3.211.8 Giac [A] (verification not implemented) 1095
 3.211.9 Mupad [B] (verification not implemented) 1096

3.211.1 Optimal result

Integrand size = 21, antiderivative size = 11

$$\int \frac{\cos(x)(-3 + 2 \sin(x))}{2 - 3 \sin(x) + \sin^2(x)} dx = \log (2 - 3 \sin(x) + \sin^2(x))$$

output `ln(2-3*sin(x)+sin(x)^2)`

3.211.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.55

$$\int \frac{\cos(x)(-3 + 2 \sin(x))}{2 - 3 \sin(x) + \sin^2(x)} dx = 2(\operatorname{arctanh}(3 - 2 \sin(x)) + \log(1 - \sin(x)))$$

input `Integrate[(Cos[x]*(-3 + 2*Sin[x]))/(2 - 3*Sin[x] + Sin[x]^2),x]`

output `2*(ArcTanh[3 - 2*Sin[x]] + Log[1 - Sin[x]])`

3.211.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {3042, 4834, 25, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(2 \sin(x) - 3) \cos(x)}{\sin^2(x) - 3 \sin(x) + 2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(2 \sin(x) - 3) \cos(x)}{\sin(x)^2 - 3 \sin(x) + 2} dx \\
 & \quad \downarrow \text{4834} \\
 & \int -\frac{3 - 2 \sin(x)}{\sin^2(x) - 3 \sin(x) + 2} d \sin(x) \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{3 - 2 \sin(x)}{\sin^2(x) - 3 \sin(x) + 2} d \sin(x) \\
 & \quad \downarrow \text{1103} \\
 & \log(\sin^2(x) - 3 \sin(x) + 2)
 \end{aligned}$$

input `Int[(Cos[x]*(-3 + 2*Sin[x]))/(2 - 3*Sin[x] + Sin[x]^2),x]`

output `Log[2 - 3*Sin[x] + Sin[x]^2]`

3.211.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4834 Int[(u_)*(F_)[(c_)*((a_) + (b_)*(x_))], x_Symbol] := With[{d = FreeFacto
rs[Sin[c*(a + b*x)], x]}, Simp[d/(b*c) Subst[Int[SubstFor[1, Sin[c*(a + b
*x)]]/d, u, x], x], x, Sin[c*(a + b*x)]/d, x] /; FunctionOfQ[Sin[c*(a + b*x
)]/d, u, x, True]] /; FreeQ[{a, b, c}, x] && (EqQ[F, Cos] || EqQ[F, cos])
```

3.211.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
derivatividivides	$\ln(2 - 3 \sin(x) + \sin^2(x))$	12
default	$\ln(2 - 3 \sin(x) + \sin^2(x))$	12
risch	$-2ix + 2 \ln(e^{ix} - i) + \ln(-4ie^{ix} + e^{2ix} - 1)$	33
norman	$2 \ln(\tan(\frac{x}{2}) - 1) - 2 \ln(1 + \tan^2(\frac{x}{2})) + \ln(\tan^2(\frac{x}{2}) - \tan(\frac{x}{2}) + 1)$	37
parallelrisch	$2 \ln(-\cot(x) + \csc(x) - 1) - 2 \ln\left(\frac{1}{\cos(x)+1}\right) + \ln\left(\frac{-\sin(x)+2}{4 \cos(x)+4}\right)$	38

```
input int(cos(x)*(-3+2*sin(x))/(2-3*sin(x)+sin(x)^2),x,method=_RETURNVERBOSE)
```

```
output ln(2-3*sin(x)+sin(x)^2)
```

3.211.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.36

$$\int \frac{\cos(x)(-3 + 2 \sin(x))}{2 - 3 \sin(x) + \sin^2(x)} dx = \log\left(-\frac{1}{2} \sin(x) + 1\right) + \log(-\sin(x) + 1)$$

```
input integrate(cos(x)*(-3+2*sin(x))/(2-3*sin(x)+sin(x)^2),x, algorithm="fracas"
)
```

```
output log(-1/2*sin(x) + 1) + log(-sin(x) + 1)
```

3.211. $\int \frac{\cos(x)(-3+2 \sin(x))}{2-3 \sin(x)+\sin^2(x)} dx$

3.211.6 Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

$$\int \frac{\cos(x)(-3 + 2 \sin(x))}{2 - 3 \sin(x) + \sin^2(x)} dx = \log(\sin(x) - 2) + \log(\sin(x) - 1)$$

input `integrate(cos(x)*(-3+2*sin(x))/(2-3*sin(x)+sin(x)**2),x)`output `log(sin(x) - 2) + log(sin(x) - 1)`**3.211.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)(-3 + 2 \sin(x))}{2 - 3 \sin(x) + \sin^2(x)} dx = \log(\sin(x)^2 - 3 \sin(x) + 2)$$

input `integrate(cos(x)*(-3+2*sin(x))/(2-3*sin(x)+sin(x)^2),x, algorithm="maxima")`output `log(sin(x)^2 - 3*sin(x) + 2)`**3.211.8 Giac [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.36

$$\int \frac{\cos(x)(-3 + 2 \sin(x))}{2 - 3 \sin(x) + \sin^2(x)} dx = \log(-\sin(x) + 2) + \log(-\sin(x) + 1)$$

input `integrate(cos(x)*(-3+2*sin(x))/(2-3*sin(x)+sin(x)^2),x, algorithm="giac")`output `log(-sin(x) + 2) + log(-sin(x) + 1)`

3.211.9 Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)(-3 + 2 \sin(x))}{2 - 3 \sin(x) + \sin^2(x)} dx = \ln(\sin(x)^2 - 3 \sin(x) + 2)$$

input `int((cos(x)*(2*sin(x) - 3))/(sin(x)^2 - 3*sin(x) + 2),x)`

output `log(sin(x)^2 - 3*sin(x) + 2)`

3.212 $\int \frac{\cos^2(x) \sin(x)}{5 + \cos^2(x)} dx$

3.212.1 Optimal result	1097
3.212.2 Mathematica [B] (verified)	1097
3.212.3 Rubi [A] (verified)	1098
3.212.4 Maple [A] (verified)	1099
3.212.5 Fricas [A] (verification not implemented)	1100
3.212.6 Sympy [A] (verification not implemented)	1100
3.212.7 Maxima [A] (verification not implemented)	1100
3.212.8 Giac [A] (verification not implemented)	1101
3.212.9 Mupad [B] (verification not implemented)	1101

3.212.1 Optimal result

Integrand size = 15, antiderivative size = 20

$$\int \frac{\cos^2(x) \sin(x)}{5 + \cos^2(x)} dx = \sqrt{5} \arctan\left(\frac{\cos(x)}{\sqrt{5}}\right) - \cos(x)$$

output `-cos(x)+arctan(1/5*cos(x)*5^(1/2))*5^(1/2)`

3.212.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 82 vs. $2(20) = 40$.

Time = 0.21 (sec) , antiderivative size = 82, normalized size of antiderivative = 4.10

$$\int \frac{\cos^2(x) \sin(x)}{5 + \cos^2(x)} dx = \frac{1}{20} \left(-\sqrt{5} \arctan\left(\frac{\cos(x)}{\sqrt{5}}\right) + 21\sqrt{5} \arctan\left(\frac{1}{\sqrt{5}} - \sqrt{\frac{6}{5}} \tan\left(\frac{x}{2}\right)\right) \right. \\ \left. + 21\sqrt{5} \arctan\left(\frac{1}{\sqrt{5}} + \sqrt{\frac{6}{5}} \tan\left(\frac{x}{2}\right)\right) - 20 \cos(x) \right)$$

input `Integrate[(Cos[x]^2*Sin[x])/(5 + Cos[x]^2),x]`

output `(-(Sqrt[5]*ArcTan[Cos[x]/Sqrt[5]]) + 21*Sqrt[5]*ArcTan[1/Sqrt[5] - Sqrt[6/5]*Tan[x/2]] + 21*Sqrt[5]*ArcTan[1/Sqrt[5] + Sqrt[6/5]*Tan[x/2]] - 20*Cos[x])/20`

3.212.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 4835, 262, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin(x) \cos^2(x)}{\cos^2(x) + 5} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(x) \cos(x)^2}{\cos(x)^2 + 5} dx \\
 & \quad \downarrow \text{4835} \\
 & - \int \frac{\cos^2(x)}{\cos^2(x) + 5} d \cos(x) \\
 & \quad \downarrow \text{262} \\
 & 5 \int \frac{1}{\cos^2(x) + 5} d \cos(x) - \cos(x) \\
 & \quad \downarrow \text{216} \\
 & \sqrt{5} \arctan\left(\frac{\cos(x)}{\sqrt{5}}\right) - \cos(x)
 \end{aligned}$$

input `Int[(Cos[x]^2*Sin[x])/(5 + Cos[x]^2),x]`

output `Sqrt[5]*ArcTan[Cos[x]/Sqrt[5]] - Cos[x]`

3.212.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

```
rule 262 Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)
^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*(m - 1)/
(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b
, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c
, 2, m, p, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4835 Int[(u_)*(F_)[(c_.)*((a_.) + (b_.)*(x_))], x_Symbol] := With[{d = FreeFacto
rs[Cos[c*(a + b*x)], x]}, Simp[-d/(b*c) Subst[Int[SubstFor[1, Cos[c*(a +
b*x)]]/d, u, x], x], x, Cos[c*(a + b*x)]/d, x] /; FunctionOfQ[Cos[c*(a + b*
x)]/d, u, x, True]] /; FreeQ[{a, b, c}, x] && (EqQ[F, Sin] || EqQ[F, sin])
```

3.212.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

method	result	size
derivativedivides	$-\cos(x) + \arctan\left(\frac{\cos(x)\sqrt{5}}{5}\right)\sqrt{5}$	18
default	$-\cos(x) + \arctan\left(\frac{\cos(x)\sqrt{5}}{5}\right)\sqrt{5}$	18
risch	$-\frac{e^{ix}}{2} - \frac{e^{-ix}}{2} - \frac{i\sqrt{5} \ln(e^{2ix} - 2i\sqrt{5}e^{ix} + 1)}{2} + \frac{i\sqrt{5} \ln(e^{2ix} + 2i\sqrt{5}e^{ix} + 1)}{2}$	66

```
input int(cos(x)^2*sin(x)/(5+cos(x)^2), x, method=_RETURNVERBOSE)
```

```
output -cos(x)+arctan(1/5*cos(x)*5^(1/2))*5^(1/2)
```

3.212.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{\cos^2(x) \sin(x)}{5 + \cos^2(x)} dx = \sqrt{5} \arctan\left(\frac{1}{5} \sqrt{5} \cos(x)\right) - \cos(x)$$

input `integrate(cos(x)^2*sin(x)/(5+cos(x)^2),x, algorithm="fricas")`output `sqrt(5)*arctan(1/5*sqrt(5)*cos(x)) - cos(x)`**3.212.6 Sympy [A] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{\cos^2(x) \sin(x)}{5 + \cos^2(x)} dx = -\cos(x) + \sqrt{5} \operatorname{atan}\left(\frac{\sqrt{5} \cos(x)}{5}\right)$$

input `integrate(cos(x)**2*sin(x)/(5+cos(x)**2),x)`output `-cos(x) + sqrt(5)*atan(sqrt(5)*cos(x)/5)`**3.212.7 Maxima [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{\cos^2(x) \sin(x)}{5 + \cos^2(x)} dx = \sqrt{5} \arctan\left(\frac{1}{5} \sqrt{5} \cos(x)\right) - \cos(x)$$

input `integrate(cos(x)^2*sin(x)/(5+cos(x)^2),x, algorithm="maxima")`output `sqrt(5)*arctan(1/5*sqrt(5)*cos(x)) - cos(x)`

3.212.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{\cos^2(x) \sin(x)}{5 + \cos^2(x)} dx = \sqrt{5} \arctan\left(\frac{1}{5} \sqrt{5} \cos(x)\right) - \cos(x)$$

input `integrate(cos(x)^2*sin(x)/(5+cos(x)^2),x, algorithm="giac")`output `sqrt(5)*arctan(1/5*sqrt(5)*cos(x)) - cos(x)`**3.212.9 Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{\cos^2(x) \sin(x)}{5 + \cos^2(x)} dx = \sqrt{5} \operatorname{atan}\left(\frac{\sqrt{5} \cos(x)}{5}\right) - \cos(x)$$

input `int((cos(x)^2*sin(x))/(cos(x)^2 + 5),x)`output `5^(1/2)*atan((5^(1/2)*cos(x))/5) - cos(x)`

3.213 $\int \frac{1}{-3+2x+x^2} dx$

3.213.1 Optimal result	1102
3.213.2 Mathematica [A] (verified)	1102
3.213.3 Rubi [A] (verified)	1103
3.213.4 Maple [A] (verified)	1104
3.213.5 Fricas [A] (verification not implemented)	1104
3.213.6 Sympy [A] (verification not implemented)	1104
3.213.7 Maxima [A] (verification not implemented)	1105
3.213.8 Giac [A] (verification not implemented)	1105
3.213.9 Mupad [B] (verification not implemented)	1105

3.213.1 Optimal result

Integrand size = 10, antiderivative size = 19

$$\int \frac{1}{-3 + 2x + x^2} dx = \frac{1}{4} \log(1 - x) - \frac{1}{4} \log(3 + x)$$

output `1/4*ln(1-x)-1/4*ln(3+x)`

3.213.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{1}{-3 + 2x + x^2} dx = \frac{1}{4} \log(1 - x) - \frac{1}{4} \log(3 + x)$$

input `Integrate[(-3 + 2*x + x^2)^(-1), x]`

output `Log[1 - x]/4 - Log[3 + x]/4`

3.213.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 + 2x - 3} dx$$

$$\downarrow \text{1081}$$

$$\int \left(-\frac{1}{4(x+3)} - \frac{1}{4(1-x)} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{1}{4} \log(1-x) - \frac{1}{4} \log(x+3)$$

input `Int[(-3 + 2*x + x^2)^(-1),x]`

output `Log[1 - x]/4 - Log[3 + x]/4`

3.213.3.1 Defintions of rubi rules used

rule 1081 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.213.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

method	result	size
default	$\frac{\ln(-1+x)}{4} - \frac{\ln(3+x)}{4}$	14
norman	$\frac{\ln(-1+x)}{4} - \frac{\ln(3+x)}{4}$	14
risch	$\frac{\ln(-1+x)}{4} - \frac{\ln(3+x)}{4}$	14
parallelrisch	$\frac{\ln(-1+x)}{4} - \frac{\ln(3+x)}{4}$	14

input `int(1/(x^2+2*x-3),x,method=_RETURNVERBOSE)`output `1/4*ln(-1+x)-1/4*ln(3+x)`**3.213.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int \frac{1}{-3 + 2x + x^2} dx = -\frac{1}{4} \log(x + 3) + \frac{1}{4} \log(x - 1)$$

input `integrate(1/(x^2+2*x-3),x, algorithm="fricas")`output `-1/4*log(x + 3) + 1/4*log(x - 1)`**3.213.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{1}{-3 + 2x + x^2} dx = \frac{\log(x - 1)}{4} - \frac{\log(x + 3)}{4}$$

input `integrate(1/(x**2+2*x-3),x)`output `log(x - 1)/4 - log(x + 3)/4`

3.213.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int \frac{1}{-3 + 2x + x^2} dx = -\frac{1}{4} \log(x + 3) + \frac{1}{4} \log(x - 1)$$

input `integrate(1/(x^2+2*x-3),x, algorithm="maxima")`output `-1/4*log(x + 3) + 1/4*log(x - 1)`**3.213.8 Giac [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{-3 + 2x + x^2} dx = -\frac{1}{4} \log(|x + 3|) + \frac{1}{4} \log(|x - 1|)$$

input `integrate(1/(x^2+2*x-3),x, algorithm="giac")`output `-1/4*log(abs(x + 3)) + 1/4*log(abs(x - 1))`**3.213.9 Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.42

$$\int \frac{1}{-3 + 2x + x^2} dx = -\frac{\operatorname{atanh}\left(\frac{x}{2} + \frac{1}{2}\right)}{2}$$

input `int(1/(2*x + x^2 - 3),x)`output `-atanh(x/2 + 1/2)/2`

3.214 $\int \frac{1}{-2x+x^2} dx$

3.214.1 Optimal result	1106
3.214.2 Mathematica [A] (verified)	1106
3.214.3 Rubi [A] (verified)	1107
3.214.4 Maple [A] (verified)	1108
3.214.5 Fricas [A] (verification not implemented)	1108
3.214.6 Sympy [A] (verification not implemented)	1108
3.214.7 Maxima [A] (verification not implemented)	1109
3.214.8 Giac [A] (verification not implemented)	1109
3.214.9 Mupad [B] (verification not implemented)	1109

3.214.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \frac{1}{-2x+x^2} dx = \frac{1}{2} \log(2-x) - \frac{\log(x)}{2}$$

output `1/2*ln(2-x)-1/2*ln(x)`

3.214.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{1}{-2x+x^2} dx = \frac{1}{2} \log(2-x) - \frac{\log(x)}{2}$$

input `Integrate[(-2*x + x^2)^(-1),x]`

output `Log[2 - x]/2 - Log[x]/2`

3.214.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {1080, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 - 2x} dx$$

$$\downarrow 1080$$

$$\int \left(\frac{1}{2(x-2)} - \frac{1}{2x} \right) dx$$

$$\downarrow 2009$$

$$\frac{1}{2} \log(2-x) - \frac{\log(x)}{2}$$

input `Int[(-2*x + x^2)^(-1), x]`

output `Log[2 - x]/2 - Log[x]/2`

3.214.3.1 Defintions of rubi rules used

rule 1080 `Int[((b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[x^p*(b + c*x)^p, x], x] /; FreeQ[{b, c}, x] && IntegerQ[p]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.214.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

method	result	size
default	$-\frac{\ln(x)}{2} + \frac{\ln(-2+x)}{2}$	12
norman	$-\frac{\ln(x)}{2} + \frac{\ln(-2+x)}{2}$	12
risch	$-\frac{\ln(x)}{2} + \frac{\ln(-2+x)}{2}$	12
parallelrisch	$-\frac{\ln(x)}{2} + \frac{\ln(-2+x)}{2}$	12
meijerg	$-\frac{\ln(x)}{2} + \frac{\ln(2)}{2} - \frac{i\pi}{2} + \frac{\ln(1-\frac{x}{2})}{2}$	22

input `int(1/(x^2-2*x),x,method=_RETURNVERBOSE)`output `-1/2*ln(x)+1/2*ln(-2+x)`**3.214.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \frac{1}{-2x + x^2} dx = \frac{1}{2} \log(x - 2) - \frac{1}{2} \log(x)$$

input `integrate(1/(x^2-2*x),x, algorithm="fricas")`output `1/2*log(x - 2) - 1/2*log(x)`**3.214.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.59

$$\int \frac{1}{-2x + x^2} dx = -\frac{\log(x)}{2} + \frac{\log(x - 2)}{2}$$

input `integrate(1/(x**2-2*x),x)`output `-log(x)/2 + log(x - 2)/2`

3.214.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \frac{1}{-2x + x^2} dx = \frac{1}{2} \log(x - 2) - \frac{1}{2} \log(x)$$

input `integrate(1/(x^2-2*x),x, algorithm="maxima")`output `1/2*log(x - 2) - 1/2*log(x)`**3.214.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{1}{-2x + x^2} dx = \frac{1}{2} \log(|x - 2|) - \frac{1}{2} \log(|x|)$$

input `integrate(1/(x^2-2*x),x, algorithm="giac")`output `1/2*log(abs(x - 2)) - 1/2*log(abs(x))`**3.214.9 Mupad [B] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.35

$$\int \frac{1}{-2x + x^2} dx = -\operatorname{atanh}(x - 1)$$

input `int(-1/(2*x - x^2),x)`output `-atanh(x - 1)`

3.215 $\int \frac{1+2x}{-7+12x+4x^2} dx$

3.215.1 Optimal result	1110
3.215.2 Mathematica [A] (verified)	1110
3.215.3 Rubi [A] (verified)	1111
3.215.4 Maple [A] (verified)	1112
3.215.5 Fricas [A] (verification not implemented)	1112
3.215.6 Sympy [A] (verification not implemented)	1112
3.215.7 Maxima [A] (verification not implemented)	1113
3.215.8 Giac [A] (verification not implemented)	1113
3.215.9 Mupad [B] (verification not implemented)	1113

3.215.1 Optimal result

Integrand size = 18, antiderivative size = 21

$$\int \frac{1 + 2x}{-7 + 12x + 4x^2} dx = \frac{1}{8} \log(1 - 2x) + \frac{3}{8} \log(7 + 2x)$$

output `1/8*ln(1-2*x)+3/8*ln(7+2*x)`

3.215.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{1 + 2x}{-7 + 12x + 4x^2} dx = \frac{1}{8} \log(1 - 2x) + \frac{3}{8} \log(7 + 2x)$$

input `Integrate[(1 + 2*x)/(-7 + 12*x + 4*x^2), x]`

output `Log[1 - 2*x]/8 + (3*Log[7 + 2*x])/8`

3.215.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x+1}{4x^2+12x-7} dx$$

↓ 1141

$$4 \int \left(\frac{3}{16(2x+7)} - \frac{1}{16(1-2x)} \right) dx$$

↓ 2009

$$4 \left(\frac{1}{32} \log(1-2x) + \frac{3}{32} \log(2x+7) \right)$$

input `Int[(1 + 2*x)/(-7 + 12*x + 4*x^2), x]`

output `4*(Log[1 - 2*x]/32 + (3*Log[7 + 2*x])/32)`

3.215.3.1 Defintions of rubi rules used

rule 1141 `Int[((d_.) + (e_.)*(x_)^(m_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x], x] /; EqQ[p, -1] || !FractionalPowerFactorQ[q] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p, 0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.215.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

method	result	size
parallelrisc	$\frac{\ln(x-\frac{1}{2})}{8} + \frac{3\ln(x+\frac{7}{2})}{8}$	14
default	$\frac{3\ln(7+2x)}{8} + \frac{\ln(2x-1)}{8}$	18
norman	$\frac{3\ln(7+2x)}{8} + \frac{\ln(2x-1)}{8}$	18
risc	$\frac{3\ln(7+2x)}{8} + \frac{\ln(2x-1)}{8}$	18

input `int((1+2*x)/(4*x^2+12*x-7),x,method=_RETURNVERBOSE)`output `1/8*ln(x-1/2)+3/8*ln(x+7/2)`**3.215.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{1+2x}{-7+12x+4x^2} dx = \frac{3}{8} \log(2x+7) + \frac{1}{8} \log(2x-1)$$

input `integrate((1+2*x)/(4*x^2+12*x-7),x, algorithm="fracas")`output `3/8*log(2*x + 7) + 1/8*log(2*x - 1)`**3.215.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{1+2x}{-7+12x+4x^2} dx = \frac{\log(x-\frac{1}{2})}{8} + \frac{3\log(x+\frac{7}{2})}{8}$$

input `integrate((1+2*x)/(4*x**2+12*x-7),x)`output `log(x - 1/2)/8 + 3*log(x + 7/2)/8`

3.215.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{1+2x}{-7+12x+4x^2} dx = \frac{3}{8} \log(2x+7) + \frac{1}{8} \log(2x-1)$$

input `integrate((1+2*x)/(4*x^2+12*x-7),x, algorithm="maxima")`output `3/8*log(2*x + 7) + 1/8*log(2*x - 1)`**3.215.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{1+2x}{-7+12x+4x^2} dx = \frac{3}{8} \log(|2x+7|) + \frac{1}{8} \log(|2x-1|)$$

input `integrate((1+2*x)/(4*x^2+12*x-7),x, algorithm="giac")`output `3/8*log(abs(2*x + 7)) + 1/8*log(abs(2*x - 1))`**3.215.9 Mupad [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \frac{1+2x}{-7+12x+4x^2} dx = \frac{\ln(x-\frac{1}{2})}{8} + \frac{3 \ln(x+\frac{7}{2})}{8}$$

input `int((2*x + 1)/(12*x + 4*x^2 - 7),x)`output `log(x - 1/2)/8 + (3*log(x + 7/2))/8`

3.216 $\int \frac{x}{-1+x+x^2} dx$

3.216.1 Optimal result	1114
3.216.2 Mathematica [A] (verified)	1114
3.216.3 Rubi [A] (verified)	1115
3.216.4 Maple [A] (verified)	1116
3.216.5 Fricas [A] (verification not implemented)	1116
3.216.6 Sympy [A] (verification not implemented)	1116
3.216.7 Maxima [A] (verification not implemented)	1117
3.216.8 Giac [A] (verification not implemented)	1117
3.216.9 Mupad [B] (verification not implemented)	1117

3.216.1 Optimal result

Integrand size = 10, antiderivative size = 49

$$\int \frac{x}{-1+x+x^2} dx = \frac{1}{10} (5 - \sqrt{5}) \log(1 - \sqrt{5} + 2x) + \frac{1}{10} (5 + \sqrt{5}) \log(1 + \sqrt{5} + 2x)$$

output `1/10*ln(1+2*x-5^(1/2))*(5-5^(1/2))+1/10*ln(1+2*x+5^(1/2))*(5+5^(1/2))`

3.216.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.90

$$\int \frac{x}{-1+x+x^2} dx = \frac{1}{10} \left(- \left((-5 + \sqrt{5}) \log(-1 + \sqrt{5} - 2x) \right) + (5 + \sqrt{5}) \log(1 + \sqrt{5} + 2x) \right)$$

input `Integrate[x/(-1 + x + x^2),x]`

output `((-((-5 + Sqrt[5])*Log[-1 + Sqrt[5] - 2*x]) + (5 + Sqrt[5])*Log[1 + Sqrt[5] + 2*x])/10`

3.216.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{x^2 + x - 1} dx$$

↓ 1141

$$\int \left(\frac{1 + \sqrt{5}}{2\sqrt{5}x + \sqrt{5} + 5} + \frac{5 - \sqrt{5}}{5(2x - \sqrt{5} + 1)} \right) dx$$

↓ 2009

$$\frac{1}{10}(5 - \sqrt{5}) \log(2x - \sqrt{5} + 1) + \frac{1}{10}(5 + \sqrt{5}) \log(2x + \sqrt{5} + 1)$$

input `Int[x/(-1 + x + x^2),x]`

output `((5 - Sqrt[5])*Log[1 - Sqrt[5] + 2*x])/10 + ((5 + Sqrt[5])*Log[1 + Sqrt[5] + 2*x])/10`

3.216.3.1 Defintions of rubi rules used

rule 1141 `Int[((d_.) + (e_.)*(x_))^(m_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x], x] /; EqQ[p, -1] || !FractionalPowerFactorQ[q] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p, 0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.216.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.55

method	result	size
default	$\frac{\ln(x^2+x-1)}{2} + \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{(1+2x)\sqrt{5}}{5}\right)}{5}$	27
risch	$\frac{\ln(2x+\sqrt{5}+1)}{2} + \frac{\ln(2x+\sqrt{5}+1)\sqrt{5}}{10} + \frac{\ln(2x+1-\sqrt{5})}{2} - \frac{\ln(2x+1-\sqrt{5})\sqrt{5}}{10}$	56

input `int(x/(x^2+x-1),x,method=_RETURNVERBOSE)`output `1/2*ln(x^2+x-1)+1/5*5^(1/2)*arctanh(1/5*(1+2*x)*5^(1/2))`**3.216.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.90

$$\int \frac{x}{-1+x+x^2} dx = \frac{1}{10} \sqrt{5} \log \left(\frac{2x^2 + \sqrt{5}(2x+1) + 2x+3}{x^2+x-1} \right) + \frac{1}{2} \log(x^2+x-1)$$

input `integrate(x/(x^2+x-1),x, algorithm="fricas")`output `1/10*sqrt(5)*log((2*x^2 + sqrt(5)*(2*x + 1) + 2*x + 3)/(x^2 + x - 1)) + 1/2*log(x^2 + x - 1)`**3.216.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.94

$$\int \frac{x}{-1+x+x^2} dx = \left(\frac{\sqrt{5}}{10} + \frac{1}{2} \right) \log \left(x + \frac{1}{2} + \frac{\sqrt{5}}{2} \right) + \left(\frac{1}{2} - \frac{\sqrt{5}}{10} \right) \log \left(x - \frac{\sqrt{5}}{2} + \frac{1}{2} \right)$$

input `integrate(x/(x**2+x-1),x)`output `(sqrt(5)/10 + 1/2)*log(x + 1/2 + sqrt(5)/2) + (1/2 - sqrt(5)/10)*log(x - sqrt(5)/2 + 1/2)`

3.216.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.76

$$\int \frac{x}{-1+x+x^2} dx = -\frac{1}{10} \sqrt{5} \log \left(\frac{2x - \sqrt{5} + 1}{2x + \sqrt{5} + 1} \right) + \frac{1}{2} \log(x^2 + x - 1)$$

input `integrate(x/(x^2+x-1),x, algorithm="maxima")`output `-1/10*sqrt(5)*log((2*x - sqrt(5) + 1)/(2*x + sqrt(5) + 1)) + 1/2*log(x^2 + x - 1)`**3.216.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.82

$$\int \frac{x}{-1+x+x^2} dx = -\frac{1}{10} \sqrt{5} \log \left(\frac{|2x - \sqrt{5} + 1|}{|2x + \sqrt{5} + 1|} \right) + \frac{1}{2} \log(|x^2 + x - 1|)$$

input `integrate(x/(x^2+x-1),x, algorithm="giac")`output `-1/10*sqrt(5)*log(abs(2*x - sqrt(5) + 1)/abs(2*x + sqrt(5) + 1)) + 1/2*log(abs(x^2 + x - 1))`**3.216.9 Mupad [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.73

$$\int \frac{x}{-1+x+x^2} dx = \ln \left(x + \frac{\sqrt{5}}{2} + \frac{1}{2} \right) \left(\frac{\sqrt{5}}{10} + \frac{1}{2} \right) - \ln \left(x - \frac{\sqrt{5}}{2} + \frac{1}{2} \right) \left(\frac{\sqrt{5}}{10} - \frac{1}{2} \right)$$

input `int(x/(x + x^2 - 1),x)`output `log(x + 5^(1/2)/2 + 1/2)*(5^(1/2)/10 + 1/2) - log(x - 5^(1/2)/2 + 1/2)*(5^(1/2)/10 - 1/2)`

3.217 $\int \frac{-32+5x-27x^2+4x^3}{-70-299x-286x^2+50x^3-13x^4+30x^5} dx$

3.217.1 Optimal result 1118
 3.217.2 Mathematica [A] (verified) 1118
 3.217.3 Rubi [A] (verified) 1119
 3.217.4 Maple [A] (verified) 1120
 3.217.5 Fricas [A] (verification not implemented) 1120
 3.217.6 Sympy [A] (verification not implemented) 1121
 3.217.7 Maxima [A] (verification not implemented) 1121
 3.217.8 Giac [A] (verification not implemented) 1122
 3.217.9 Mupad [B] (verification not implemented) 1122

3.217.1 Optimal result

Integrand size = 43, antiderivative size = 63

$$\int \frac{-32 + 5x - 27x^2 + 4x^3}{-70 - 299x - 286x^2 + 50x^3 - 13x^4 + 30x^5} dx$$

$$= \frac{3988 \arctan\left(\frac{1+2x}{\sqrt{19}}\right)}{13685\sqrt{19}} - \frac{3146 \log(7 - 3x)}{80155} - \frac{334}{323} \log(1 + 2x)$$

$$+ \frac{4822 \log(2 + 5x)}{4879} + \frac{11049 \log(5 + x + x^2)}{260015}$$

output `-3146/80155*ln(7-3*x)-334/323*ln(1+2*x)+4822/4879*ln(2+5*x)+11049/260015*ln(x^2+x+5)+3988/260015*arctan(1/19*(1+2*x)*19^(1/2))*19^(1/2)`

3.217.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.90

$$\int \frac{-32 + 5x - 27x^2 + 4x^3}{-70 - 299x - 286x^2 + 50x^3 - 13x^4 + 30x^5} dx$$

$$= \frac{163508\sqrt{19} \arctan\left(\frac{1+2x}{\sqrt{19}}\right) - 418418 \log(7 - 3x) - 11023670 \log(1 + 2x) + 10536070 \log(2 + 5x) + 45300}{10660615}$$

input `Integrate[(-32 + 5*x - 27*x^2 + 4*x^3)/(-70 - 299*x - 286*x^2 + 50*x^3 - 13*x^4 + 30*x^5), x]`

3.217. $\int \frac{-32+5x-27x^2+4x^3}{-70-299x-286x^2+50x^3-13x^4+30x^5} dx$

output (163508* $\sqrt{19}$ *ArcTan[(1 + 2*x)/ $\sqrt{19}$] - 418418*Log[7 - 3*x] - 110236
70*Log[1 + 2*x] + 10536070*Log[2 + 5*x] + 453009*Log[5 + x + x^2])/1066061
5

3.217.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.047$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{4x^3 - 27x^2 + 5x - 32}{30x^5 - 13x^4 + 50x^3 - 286x^2 - 299x - 70} dx$$

↓ 2462

$$\int \left(\frac{22098x + 48935}{260015(x^2 + x + 5)} - \frac{668}{323(2x + 1)} - \frac{9438}{80155(3x - 7)} + \frac{24110}{4879(5x + 2)} \right) dx$$

↓ 2009

$$\frac{3988 \arctan\left(\frac{2x+1}{\sqrt{19}}\right)}{13685\sqrt{19}} + \frac{11049 \log(x^2 + x + 5)}{260015} - \frac{3146 \log(7 - 3x)}{80155} - \frac{334}{323} \log(2x+1) + \frac{4822 \log(5x + 2)}{4879}$$

input Int[(-32 + 5*x - 27*x^2 + 4*x^3)/(-70 - 299*x - 286*x^2 + 50*x^3 - 13*x^4 + 30*x^5), x]

output (3988*ArcTan[(1 + 2*x)/ $\sqrt{19}$])/(13685* $\sqrt{19}$) - (3146*Log[7 - 3*x])/8
0155 - (334*Log[1 + 2*x])/323 + (4822*Log[2 + 5*x])/4879 + (11049*Log[5 +
x + x^2])/260015

3.217. $\int \frac{-32+5x-27x^2+4x^3}{-70-299x-286x^2+50x^3-13x^4+30x^5} dx$

3.217.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2462 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr
and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ
[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0
] && RationalFunctionQ[u, x]`

3.217.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.81

method	result
default	$\frac{4822 \ln(5x+2)}{4879} - \frac{3146 \ln(3x-7)}{80155} + \frac{11049 \ln(x^2+x+5)}{260015} + \frac{3988 \arctan\left(\frac{(1+2x)\sqrt{19}}{19}\right)\sqrt{19}}{260015} - \frac{334 \ln(1+2x)}{323}$
risch	$-\frac{3146 \ln(3x-7)}{80155} + \frac{4822 \ln(5x+2)}{4879} - \frac{334 \ln(1+2x)}{323} + \frac{11049 \ln(15904144x^2+15904144x+79520720)}{260015} + \frac{3988\sqrt{19} \arctan\left(\frac{3988}{260015}\right)}{260015}$

input `int((4*x^3-27*x^2+5*x-32)/(30*x^5-13*x^4+50*x^3-286*x^2-299*x-70),x,method
=_RETURNVERBOSE)`

output `4822/4879*ln(5*x+2)-3146/80155*ln(3*x-7)+11049/260015*ln(x^2+x+5)+3988/260
015*arctan(1/19*(1+2*x)*19^(1/2))*19^(1/2)-334/323*ln(1+2*x)`

3.217.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.79

$$\int \frac{-32 + 5x - 27x^2 + 4x^3}{-70 - 299x - 286x^2 + 50x^3 - 13x^4 + 30x^5} dx$$

$$= \frac{3988}{260015} \sqrt{19} \arctan\left(\frac{1}{19} \sqrt{19}(2x+1)\right) + \frac{11049}{260015} \log(x^2+x+5)$$

$$+ \frac{4822}{4879} \log(5x+2) - \frac{3146}{80155} \log(3x-7) - \frac{334}{323} \log(2x+1)$$

input `integrate((4*x^3-27*x^2+5*x-32)/(30*x^5-13*x^4+50*x^3-286*x^2-299*x-70),x,
algorithm="fracas")`

3.217. $\int \frac{-32+5x-27x^2+4x^3}{-70-299x-286x^2+50x^3-13x^4+30x^5} dx$

output $3988/260015*\sqrt{19}*\arctan(1/19*\sqrt{19}*(2*x + 1)) + 11049/260015*\log(x^2 + x + 5) + 4822/4879*\log(5*x + 2) - 3146/80155*\log(3*x - 7) - 334/323*\log(2*x + 1)$

3.217.6 Sympy [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.08

$$\int \frac{-32 + 5x - 27x^2 + 4x^3}{-70 - 299x - 286x^2 + 50x^3 - 13x^4 + 30x^5} dx$$

$$= -\frac{3146 \log\left(x - \frac{7}{3}\right)}{80155} + \frac{4822 \log\left(x + \frac{2}{5}\right)}{4879} - \frac{334 \log\left(x + \frac{1}{2}\right)}{323}$$

$$+ \frac{11049 \log(x^2 + x + 5)}{260015} + \frac{3988\sqrt{19} \operatorname{atan}\left(\frac{2\sqrt{19}x}{19} + \frac{\sqrt{19}}{19}\right)}{260015}$$

input `integrate((4*x**3-27*x**2+5*x-32)/(30*x**5-13*x**4+50*x**3-286*x**2-299*x-70),x)`

output $-3146*\log(x - 7/3)/80155 + 4822*\log(x + 2/5)/4879 - 334*\log(x + 1/2)/323 + 11049*\log(x**2 + x + 5)/260015 + 3988*\sqrt{19}*\operatorname{atan}(2*\sqrt{19}*x/19 + \sqrt{19}/19)/260015$

3.217.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.79

$$\int \frac{-32 + 5x - 27x^2 + 4x^3}{-70 - 299x - 286x^2 + 50x^3 - 13x^4 + 30x^5} dx$$

$$= \frac{3988}{260015} \sqrt{19} \arctan\left(\frac{1}{19} \sqrt{19}(2x + 1)\right) + \frac{11049}{260015} \log(x^2 + x + 5)$$

$$+ \frac{4822}{4879} \log(5x + 2) - \frac{3146}{80155} \log(3x - 7) - \frac{334}{323} \log(2x + 1)$$

input `integrate((4*x^3-27*x^2+5*x-32)/(30*x^5-13*x^4+50*x^3-286*x^2-299*x-70),x, algorithm="maxima")`

output $3988/260015*\sqrt{19}*\arctan(1/19*\sqrt{19}*(2*x + 1)) + 11049/260015*\log(x^2 + x + 5) + 4822/4879*\log(5*x + 2) - 3146/80155*\log(3*x - 7) - 334/323*\log(2*x + 1)$

3.217. $\int \frac{-32+5x-27x^2+4x^3}{-70-299x-286x^2+50x^3-13x^4+30x^5} dx$

3.217.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.84

$$\int \frac{-32 + 5x - 27x^2 + 4x^3}{-70 - 299x - 286x^2 + 50x^3 - 13x^4 + 30x^5} dx$$

$$= \frac{3988}{260015} \sqrt{19} \arctan\left(\frac{1}{19} \sqrt{19}(2x + 1)\right) + \frac{11049}{260015} \log(x^2 + x + 5)$$

$$+ \frac{4822}{4879} \log(|5x + 2|) - \frac{3146}{80155} \log(|3x - 7|) - \frac{334}{323} \log(|2x + 1|)$$

```
input integrate((4*x^3-27*x^2+5*x-32)/(30*x^5-13*x^4+50*x^3-286*x^2-299*x-70),x,
algorithm="giac")
```

```
output 3988/260015*sqrt(19)*arctan(1/19*sqrt(19)*(2*x + 1)) + 11049/260015*log(x^
2 + x + 5) + 4822/4879*log(abs(5*x + 2)) - 3146/80155*log(abs(3*x - 7)) -
334/323*log(abs(2*x + 1))
```

3.217.9 Mupad [B] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.92

$$\int \frac{-32 + 5x - 27x^2 + 4x^3}{-70 - 299x - 286x^2 + 50x^3 - 13x^4 + 30x^5} dx$$

$$= \frac{4822 \ln\left(x + \frac{2}{5}\right)}{4879} - \frac{334 \ln\left(x + \frac{1}{2}\right)}{323} - \frac{3146 \ln\left(x - \frac{7}{3}\right)}{80155}$$

$$- \ln\left(x + \frac{1}{2} - \frac{\sqrt{19}i}{2}\right) \left(-\frac{11049}{260015} + \frac{\sqrt{19}1994i}{260015}\right)$$

$$+ \ln\left(x + \frac{1}{2} + \frac{\sqrt{19}i}{2}\right) \left(\frac{11049}{260015} + \frac{\sqrt{19}1994i}{260015}\right)$$

```
input int(-(5*x - 27*x^2 + 4*x^3 - 32)/(299*x + 286*x^2 - 50*x^3 + 13*x^4 - 30*x
^5 + 70),x)
```

```
output (4822*log(x + 2/5))/4879 - (334*log(x + 1/2))/323 - (3146*log(x - 7/3))/80
155 - log(x - (19^(1/2)*1i)/2 + 1/2)*((19^(1/2)*1994i)/260015 - 11049/2600
15) + log(x + (19^(1/2)*1i)/2 + 1/2)*((19^(1/2)*1994i)/260015 + 11049/2600
15)
```

$$3.218 \quad \int \frac{8-13x^2-7x^3+12x^5}{4-20x+41x^2-80x^3+116x^4-80x^5+100x^6} dx$$

3.218.1 Optimal result	1123
3.218.2 Mathematica [A] (verified)	1123
3.218.3 Rubi [A] (verified)	1124
3.218.4 Maple [A] (verified)	1125
3.218.5 Fricas [A] (verification not implemented)	1125
3.218.6 Sympy [A] (verification not implemented)	1126
3.218.7 Maxima [A] (verification not implemented)	1126
3.218.8 Giac [A] (verification not implemented)	1127
3.218.9 Mupad [B] (verification not implemented)	1127

3.218.1 Optimal result

Integrand size = 50, antiderivative size = 86

$$\begin{aligned} & \int \frac{8-13x^2-7x^3+12x^5}{4-20x+41x^2-80x^3+116x^4-80x^5+100x^6} dx \\ &= \frac{5828}{9075(2-5x)} - \frac{313+502x}{1452(1+2x^2)} - \frac{251 \arctan(\sqrt{2}x)}{726\sqrt{2}} \\ & \quad + \frac{272\sqrt{2} \arctan(\sqrt{2}x)}{1331} - \frac{59096 \log(2-5x)}{99825} + \frac{2843 \log(1+2x^2)}{7986} \end{aligned}$$

output `5828/9075/(2-5*x)+1/1452*(-313-502*x)/(2*x^2+1)-59096/99825*ln(2-5*x)+2843/7986*ln(2*x^2+1)+503/15972*arctan(x*2^(1/2))*2^(1/2)`

3.218.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.78

$$\begin{aligned} & \int \frac{8-13x^2-7x^3+12x^5}{4-20x+41x^2-80x^3+116x^4-80x^5+100x^6} dx \\ &= \frac{-\frac{33(2554+4675x+36458x^2)}{-2+5x-4x^2+10x^3} + 12575\sqrt{2} \arctan(\sqrt{2}x) - 236384 \log(2-5x) + 142150 \log(1+2x^2)}{399300} \end{aligned}$$

input `Integrate[(8 - 13*x^2 - 7*x^3 + 12*x^5)/(4 - 20*x + 41*x^2 - 80*x^3 + 116*x^4 - 80*x^5 + 100*x^6), x]`

$$3.218. \quad \int \frac{8-13x^2-7x^3+12x^5}{4-20x+41x^2-80x^3+116x^4-80x^5+100x^6} dx$$

output $((-33*(2554 + 4675*x + 36458*x^2))/(-2 + 5*x - 4*x^2 + 10*x^3) + 12575*sqrt{2}*ArcTan[Sqrt{2}*x] - 236384*Log[2 - 5*x] + 142150*Log[1 + 2*x^2])/399300$

3.218.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{12x^5 - 7x^3 - 13x^2 + 8}{100x^6 - 80x^5 + 116x^4 - 80x^3 + 41x^2 - 20x + 4} dx$$

↓ 2462

$$\int \left(\frac{313x - 251}{363(2x^2 + 1)^2} + \frac{2(2843x + 816)}{3993(2x^2 + 1)} - \frac{59096}{19965(5x - 2)} + \frac{5828}{1815(5x - 2)^2} \right) dx$$

↓ 2009

$$\frac{272\sqrt{2} \arctan(\sqrt{2}x)}{1331} - \frac{251 \arctan(\sqrt{2}x)}{726\sqrt{2}} - \frac{502x + 313}{1452(2x^2 + 1)} + \frac{2843 \log(2x^2 + 1)}{7986} + \frac{5828}{9075(2 - 5x)} - \frac{59096 \log(2 - 5x)}{99825}$$

input $Int[(8 - 13*x^2 - 7*x^3 + 12*x^5)/(4 - 20*x + 41*x^2 - 80*x^3 + 116*x^4 - 80*x^5 + 100*x^6), x]$

output $5828/(9075*(2 - 5*x)) - (313 + 502*x)/(1452*(1 + 2*x^2)) - (251*ArcTan[Sqrt{2}*x])/(726*sqrt{2}) + (272*sqrt{2}*ArcTan[Sqrt{2}*x])/1331 - (59096*Log[2 - 5*x])/99825 + (2843*Log[1 + 2*x^2])/7986$

3.218.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2462 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr
and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ
[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0
] && RationalFunctionQ[u, x]`

3.218.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.63

method	result	size
default	$\frac{-\frac{2761x}{4} - \frac{3443}{8}}{3993x^2 + \frac{3993}{2}} + \frac{2843 \ln(2x^2+1)}{7986} + \frac{503 \arctan(x\sqrt{2})\sqrt{2}}{15972} - \frac{5828}{9075(5x-2)} - \frac{59096 \ln(5x-2)}{99825}$	54
risch	$\frac{-\frac{18229}{60500}x^2 - \frac{17}{440}x - \frac{1277}{60500}}{x^3 - \frac{2}{5}x^2 + \frac{1}{2}x - \frac{1}{5}} + \frac{2843 \ln(4x^2+2)}{7986} + \frac{503 \arctan(x\sqrt{2})\sqrt{2}}{15972} - \frac{59096 \ln(5x-2)}{99825}$	57

input `int((12*x^5-7*x^3-13*x^2+8)/(100*x^6-80*x^5+116*x^4-80*x^3+41*x^2-20*x+4),
x,method=_RETURNVERBOSE)`

output `1/3993*(-2761/4*x-3443/8)/(x^2+1/2)+2843/7986*ln(2*x^2+1)+503/15972*arctan
(x*2^(1/2))*2^(1/2)-5828/9075/(5*x-2)-59096/99825*ln(5*x-2)`

3.218.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.20

$$\int \frac{8 - 13x^2 - 7x^3 + 12x^5}{4 - 20x + 41x^2 - 80x^3 + 116x^4 - 80x^5 + 100x^6} dx$$

$$= \frac{12575 \sqrt{2}(10x^3 - 4x^2 + 5x - 2) \arctan(\sqrt{2}x) - 1203114x^2 + 142150(10x^3 - 4x^2 + 5x - 2) \log(2x^2 - 10x + 4)}{399300(10x^3 - 4x^2 + 5x - 2)}$$

input `integrate((12*x^5-7*x^3-13*x^2+8)/(100*x^6-80*x^5+116*x^4-80*x^3+41*x^2-20
*x+4),x, algorithm="fracas")`

3.218.
$$\int \frac{8 - 13x^2 - 7x^3 + 12x^5}{4 - 20x + 41x^2 - 80x^3 + 116x^4 - 80x^5 + 100x^6} dx$$

output `1/399300*(12575*sqrt(2)*(10*x^3 - 4*x^2 + 5*x - 2)*arctan(sqrt(2)*x) - 120
3114*x^2 + 142150*(10*x^3 - 4*x^2 + 5*x - 2)*log(2*x^2 + 1) - 236384*(10*x
^3 - 4*x^2 + 5*x - 2)*log(5*x - 2) - 154275*x - 84282)/(10*x^3 - 4*x^2 + 5
*x - 2)`

3.218.6 Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.76

$$\int \frac{8 - 13x^2 - 7x^3 + 12x^5}{4 - 20x + 41x^2 - 80x^3 + 116x^4 - 80x^5 + 100x^6} dx$$

$$= \frac{-36458x^2 - 4675x - 2554}{121000x^3 - 48400x^2 + 60500x - 24200} - \frac{59096 \log\left(x - \frac{2}{5}\right)}{99825}$$

$$+ \frac{2843 \log\left(x^2 + \frac{1}{2}\right)}{7986} + \frac{503\sqrt{2} \operatorname{atan}\left(\sqrt{2}x\right)}{15972}$$

input `integrate((12*x**5-7*x**3-13*x**2+8)/(100*x**6-80*x**5+116*x**4-80*x**3+41
*x**2-20*x+4),x)`

output `(-36458*x**2 - 4675*x - 2554)/(121000*x**3 - 48400*x**2 + 60500*x - 24200)
- 59096*log(x - 2/5)/99825 + 2843*log(x**2 + 1/2)/7986 + 503*sqrt(2)*atan
(sqrt(2)*x)/15972`

3.218.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.69

$$\int \frac{8 - 13x^2 - 7x^3 + 12x^5}{4 - 20x + 41x^2 - 80x^3 + 116x^4 - 80x^5 + 100x^6} dx$$

$$= \frac{503}{15972} \sqrt{2} \operatorname{arctan}\left(\sqrt{2}x\right) - \frac{36458x^2 + 4675x + 2554}{12100(10x^3 - 4x^2 + 5x - 2)}$$

$$+ \frac{2843}{7986} \log(2x^2 + 1) - \frac{59096}{99825} \log(5x - 2)$$

input `integrate((12*x^5-7*x^3-13*x^2+8)/(100*x^6-80*x^5+116*x^4-80*x^3+41*x^2-20
*x+4),x, algorithm="maxima")`

output $503/15972*\sqrt{2}*\arctan(\sqrt{2}*x) - 1/12100*(36458*x^2 + 4675*x + 2554)/(10*x^3 - 4*x^2 + 5*x - 2) + 2843/7986*\log(2*x^2 + 1) - 59096/99825*\log(5*x - 2)$

3.218.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.69

$$\int \frac{8 - 13x^2 - 7x^3 + 12x^5}{4 - 20x + 41x^2 - 80x^3 + 116x^4 - 80x^5 + 100x^6} dx$$

$$= \frac{503}{15972} \sqrt{2} \arctan(\sqrt{2}x) - \frac{36458x^2 + 4675x + 2554}{12100(2x^2 + 1)(5x - 2)}$$

$$+ \frac{2843}{7986} \log(2x^2 + 1) - \frac{59096}{99825} \log(|5x - 2|)$$

input `integrate((12*x^5-7*x^3-13*x^2+8)/(100*x^6-80*x^5+116*x^4-80*x^3+41*x^2-20*x+4),x, algorithm="giac")`

output $503/15972*\sqrt{2}*\arctan(\sqrt{2}*x) - 1/12100*(36458*x^2 + 4675*x + 2554)/((2*x^2 + 1)*(5*x - 2)) + 2843/7986*\log(2*x^2 + 1) - 59096/99825*\log(\text{abs}(5*x - 2))$

3.218.9 Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.83

$$\int \frac{8 - 13x^2 - 7x^3 + 12x^5}{4 - 20x + 41x^2 - 80x^3 + 116x^4 - 80x^5 + 100x^6} dx$$

$$= -\frac{59096 \ln\left(x - \frac{2}{5}\right)}{99825} - \frac{\frac{18229x^2}{60500} + \frac{17x}{440} + \frac{1277}{60500}}{x^3 - \frac{2x^2}{5} + \frac{x}{2} - \frac{1}{5}}$$

$$- \ln\left(x - \frac{\sqrt{2} \text{li}}{2}\right) \left(-\frac{2843}{7986} + \frac{\sqrt{2} 503i}{31944}\right) + \ln\left(x + \frac{\sqrt{2} \text{li}}{2}\right) \left(\frac{2843}{7986} + \frac{\sqrt{2} 503i}{31944}\right)$$

input `int(-(13*x^2 + 7*x^3 - 12*x^5 - 8)/(41*x^2 - 20*x - 80*x^3 + 116*x^4 - 80*x^5 + 100*x^6 + 4),x)`

output $\log(x + (2^{(1/2)}*1i)/2)*((2^{(1/2)}*503i)/31944 + 2843/7986) - ((17*x)/440 + (18229*x^2)/60500 + 1277/60500)/(x/2 - (2*x^2)/5 + x^3 - 1/5) - \log(x - (2^{(1/2)}*1i)/2)*((2^{(1/2)}*503i)/31944 - 2843/7986) - (59096*\log(x - 2/5))/99825$

3.218. $\int \frac{8-13x^2-7x^3+12x^5}{4-20x+41x^2-80x^3+116x^4-80x^5+100x^6} dx$

3.219 $\int \frac{\sqrt{4+x}}{x} dx$

3.219.1 Optimal result	1129
3.219.2 Mathematica [A] (verified)	1129
3.219.3 Rubi [A] (verified)	1130
3.219.4 Maple [A] (verified)	1131
3.219.5 Fricas [A] (verification not implemented)	1131
3.219.6 Sympy [B] (verification not implemented)	1132
3.219.7 Maxima [A] (verification not implemented)	1132
3.219.8 Giac [A] (verification not implemented)	1132
3.219.9 Mupad [B] (verification not implemented)	1133

3.219.1 Optimal result

Integrand size = 11, antiderivative size = 24

$$\int \frac{\sqrt{4+x}}{x} dx = 2\sqrt{4+x} - 4\operatorname{arctanh}\left(\frac{\sqrt{4+x}}{2}\right)$$

output `-4*arctanh(1/2*(4+x)^(1/2))+2*(4+x)^(1/2)`

3.219.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{4+x}}{x} dx = 2\sqrt{4+x} - 4\operatorname{arctanh}\left(\frac{\sqrt{4+x}}{2}\right)$$

input `Integrate[Sqrt[4 + x]/x,x]`

output `2*Sqrt[4 + x] - 4*ArcTanh[Sqrt[4 + x]/2]`

3.219.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {60, 73, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{x+4}}{x} dx \\ & \quad \downarrow \text{60} \\ & 4 \int \frac{1}{x\sqrt{x+4}} dx + 2\sqrt{x+4} \\ & \quad \downarrow \text{73} \\ & 8 \int \frac{1}{x} d\sqrt{x+4} + 2\sqrt{x+4} \\ & \quad \downarrow \text{220} \\ & 2\sqrt{x+4} - 4\operatorname{arctanh}\left(\frac{\sqrt{x+4}}{2}\right) \end{aligned}$$

input `Int[Sqrt[4 + x]/x,x]`

output `2*Sqrt[4 + x] - 4*ArcTanh[Sqrt[4 + x]/2]`

3.219.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 220 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-
1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] &&
(LtQ[a, 0] || GtQ[b, 0])
```

3.219.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17

method	result	size
trager	$2\sqrt{4+x} + 2\ln\left(\frac{-8-x+4\sqrt{4+x}}{x}\right)$	28
derivativedivides	$2\sqrt{4+x} - 2\ln(\sqrt{4+x} + 2) + 2\ln(\sqrt{4+x} - 2)$	29
default	$2\sqrt{4+x} - 2\ln(\sqrt{4+x} + 2) + 2\ln(\sqrt{4+x} - 2)$	29
meijerg	$-\frac{-2(2-4\ln(2)+\ln(x))\sqrt{\pi}+4\sqrt{\pi}-4\sqrt{\pi}\sqrt{1+\frac{x}{4}}+4\sqrt{\pi}\ln\left(\frac{1}{2}+\frac{\sqrt{1+\frac{x}{4}}}{2}\right)}{\sqrt{\pi}}$	54

```
input int((4+x)^(1/2)/x,x,method=_RETURNVERBOSE)
```

```
output 2*(4+x)^(1/2)+2*ln((-8-x+4*(4+x)^(1/2))/x)
```

3.219.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17

$$\int \frac{\sqrt{4+x}}{x} dx = 2\sqrt{x+4} - 2\log(\sqrt{x+4} + 2) + 2\log(\sqrt{x+4} - 2)$$

```
input integrate((4+x)^(1/2)/x,x, algorithm="fricas")
```

```
output 2*sqrt(x + 4) - 2*log(sqrt(x + 4) + 2) + 2*log(sqrt(x + 4) - 2)
```

3.219.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. $2(19) = 38$.

Time = 0.51 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.75

$$\int \frac{\sqrt{4+x}}{x} dx = \begin{cases} 2\sqrt{x+4} - 4 \operatorname{acoth}\left(\frac{\sqrt{x+4}}{2}\right) & \text{for } |x+4| > 4 \\ 2\sqrt{x+4} - 4 \operatorname{atanh}\left(\frac{\sqrt{x+4}}{2}\right) & \text{otherwise} \end{cases}$$

input `integrate((4+x)**(1/2)/x,x)`

output `Piecewise((2*sqrt(x + 4) - 4*acoth(sqrt(x + 4)/2), Abs(x + 4) > 4), (2*sqrt(x + 4) - 4*atanh(sqrt(x + 4)/2), True))`

3.219.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17

$$\int \frac{\sqrt{4+x}}{x} dx = 2\sqrt{x+4} - 2 \log(\sqrt{x+4} + 2) + 2 \log(\sqrt{x+4} - 2)$$

input `integrate((4+x)^(1/2)/x,x, algorithm="maxima")`

output `2*sqrt(x + 4) - 2*log(sqrt(x + 4) + 2) + 2*log(sqrt(x + 4) - 2)`

3.219.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21

$$\int \frac{\sqrt{4+x}}{x} dx = 2\sqrt{x+4} - 2 \log(\sqrt{x+4} + 2) + 2 \log\left(\left|\sqrt{x+4} - 2\right|\right)$$

input `integrate((4+x)^(1/2)/x,x, algorithm="giac")`

output `2*sqrt(x + 4) - 2*log(sqrt(x + 4) + 2) + 2*log(abs(sqrt(x + 4) - 2))`

3.219.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{4+x}}{x} dx = 2\sqrt{x+4} - 4 \operatorname{atanh}\left(\frac{\sqrt{x+4}}{2}\right)$$

input `int((x + 4)^(1/2)/x,x)`

output `2*(x + 4)^(1/2) - 4*atanh((x + 4)^(1/2)/2)`

3.220 $\int \frac{1}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx$

3.220.1 Optimal result 1134
 3.220.2 Mathematica [C] (verified) 1135
 3.220.3 Rubi [A] (verified) 1135
 3.220.4 Maple [A] (verified) 1138
 3.220.5 Fricas [B] (verification not implemented) 1139
 3.220.6 Sympy [F] 1140
 3.220.7 Maxima [B] (verification not implemented) 1141
 3.220.8 Giac [A] (verification not implemented) 1142
 3.220.9 Mupad [B] (verification not implemented) 1142

3.220.1 Optimal result

Integrand size = 15, antiderivative size = 200

$$\int \frac{1}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx = 2\sqrt{x} + \frac{3}{5}\sqrt{2(5-\sqrt{5})} \arctan\left(\frac{1-\sqrt{5}+4\sqrt[6]{x}}{\sqrt{2(5+\sqrt{5})}}\right) - \frac{3}{5}\sqrt{2(5+\sqrt{5})} \arctan\left(\frac{1}{2}\sqrt{\frac{1}{10}(5+\sqrt{5})}(1+\sqrt{5}+4\sqrt[6]{x})\right) + \frac{6}{5}\log(1-\sqrt[6]{x}) - \frac{3}{10}(1+\sqrt{5})\log(2+\sqrt[6]{x}-\sqrt{5}\sqrt[6]{x}+2\sqrt[3]{x}) - \frac{3}{10}(1-\sqrt{5})\log(2+\sqrt[6]{x}+\sqrt{5}\sqrt[6]{x}+2\sqrt[3]{x})$$

```
output 6/5*ln(1-x^(1/6))-3/10*ln(2+x^(1/6)+2*x^(1/3)+x^(1/6)*5^(1/2))*(-5^(1/2)+1)
-3/10*ln(2+x^(1/6)+2*x^(1/3)-x^(1/6)*5^(1/2))*(5^(1/2)+1)+2*x^(1/2)+3/5*a
rctan((1+4*x^(1/6)-5^(1/2))/(10+2*5^(1/2))^(1/2))*(10-2*5^(1/2))^(1/2)-3/5
*arctan(1/20*(1+4*x^(1/6)+5^(1/2))*(50+10*5^(1/2))^(1/2))*(10+2*5^(1/2))^(
1/2)
```

3.220.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.07 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.63

$$\int \frac{1}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx = 2\sqrt{x} + \frac{6}{5} \log(-1 + \sqrt[6]{x}) - \frac{6}{5} \text{RootSum} \left[1 + \#1 + \#1^2 + \#1^3 \right. \\ \left. + \#1^4 \&, \frac{-\log(\sqrt[6]{x} - \#1) - 2\log(\sqrt[6]{x} - \#1)\#1 + 2\log(\sqrt[6]{x} - \#1)\#1^2 + \log(\sqrt[6]{x} - \#1)\#1^3}{1 + 2\#1 + 3\#1^2 + 4\#1^3} \& \right]$$

input `Integrate[(-x^(-1/3) + Sqrt[x])^(-1), x]`

output `2*Sqrt[x] + (6*Log[-1 + x^(1/6)])/5 - (6*RootSum[1 + #1 + #1^2 + #1^3 + #1^4 & , (-Log[x^(1/6) - #1] - 2*Log[x^(1/6) - #1]*#1 + 2*Log[x^(1/6) - #1]*#1^2 + Log[x^(1/6) - #1]*#1^3)/(1 + 2*#1 + 3*#1^2 + 4*#1^3) &])/5`

3.220.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.733$, Rules used = {2027, 864, 25, 843, 823, 16, 27, 1142, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt{x} - \frac{1}{\sqrt[3]{x}}} dx \\ & \quad \downarrow \text{2027} \\ & \int \frac{\sqrt[3]{x}}{x^{5/6} - 1} dx \\ & \quad \downarrow \text{864} \\ & 6 \int -\frac{x^{7/6}}{1 - x^{5/6}} d\sqrt[6]{x} \\ & \quad \downarrow \text{25} \\ & -6 \int \frac{x^{7/6}}{1 - x^{5/6}} d\sqrt[6]{x} \end{aligned}$$

$$\begin{aligned}
& \downarrow 843 \\
& 6 \left(\frac{\sqrt{x}}{3} - \int \frac{\sqrt[3]{x}}{1-x^{5/6}} d\sqrt[6]{x} \right) \\
& \downarrow 823 \\
& 6 \left(-\frac{1}{5} \int \frac{1}{1-\sqrt[6]{x}} d\sqrt[6]{x} - \frac{2}{5} \int -\frac{-((1+\sqrt{5})\sqrt[6]{x}) + \sqrt{5} + 1}{2(2\sqrt[3]{x} + (1-\sqrt{5})\sqrt[6]{x} + 2)} d\sqrt[6]{x} - \frac{2}{5} \int -\frac{-((1-\sqrt{5})\sqrt[6]{x}) - \sqrt{5} + 1}{2(2\sqrt[3]{x} + (1+\sqrt{5})\sqrt[6]{x} + 2)} d\sqrt[6]{x} + \frac{\sqrt{x}}{3} \right) \\
& \downarrow 16 \\
& 6 \left(-\frac{2}{5} \int -\frac{-((1+\sqrt{5})\sqrt[6]{x}) + \sqrt{5} + 1}{2(2\sqrt[3]{x} + (1-\sqrt{5})\sqrt[6]{x} + 2)} d\sqrt[6]{x} - \frac{2}{5} \int -\frac{-((1-\sqrt{5})\sqrt[6]{x}) - \sqrt{5} + 1}{2(2\sqrt[3]{x} + (1+\sqrt{5})\sqrt[6]{x} + 2)} d\sqrt[6]{x} + \frac{\sqrt{x}}{3} + \frac{1}{5} \log(1-\sqrt[6]{x}) \right) \\
& \downarrow 27 \\
& 6 \left(\frac{1}{5} \int \frac{-((1+\sqrt{5})\sqrt[6]{x}) + \sqrt{5} + 1}{2\sqrt[3]{x} + (1-\sqrt{5})\sqrt[6]{x} + 2} d\sqrt[6]{x} + \frac{1}{5} \int \frac{-((1-\sqrt{5})\sqrt[6]{x}) - \sqrt{5} + 1}{2\sqrt[3]{x} + (1+\sqrt{5})\sqrt[6]{x} + 2} d\sqrt[6]{x} + \frac{\sqrt{x}}{3} + \frac{1}{5} \log(1-\sqrt[6]{x}) \right) \\
& \downarrow 1142 \\
& 6 \left(\frac{1}{5} \left(\sqrt{5} \int \frac{1}{2\sqrt[3]{x} + (1-\sqrt{5})\sqrt[6]{x} + 2} d\sqrt[6]{x} - \frac{1}{4} (1+\sqrt{5}) \int \frac{4\sqrt[6]{x} - \sqrt{5} + 1}{2\sqrt[3]{x} + (1-\sqrt{5})\sqrt[6]{x} + 2} d\sqrt[6]{x} \right) + \frac{1}{5} \left(-\sqrt{5} \int \frac{1}{2\sqrt[3]{x} + (1+\sqrt{5})\sqrt[6]{x} + 2} d\sqrt[6]{x} - \frac{1}{4} (1+\sqrt{5}) \int \frac{4\sqrt[6]{x} - \sqrt{5} + 1}{2\sqrt[3]{x} + (1+\sqrt{5})\sqrt[6]{x} + 2} d\sqrt[6]{x} \right) \right) \\
& \downarrow 1083 \\
& 6 \left(\frac{1}{5} \left(-2\sqrt{5} \int \frac{1}{-\sqrt[3]{x} - 2(5+\sqrt{5})} d(4\sqrt[6]{x} - \sqrt{5} + 1) - \frac{1}{4} (1+\sqrt{5}) \int \frac{4\sqrt[6]{x} - \sqrt{5} + 1}{2\sqrt[3]{x} + (1-\sqrt{5})\sqrt[6]{x} + 2} d\sqrt[6]{x} \right) + \frac{1}{5} \left(2\sqrt{5} \int \frac{1}{\sqrt[3]{x} - 2(5-\sqrt{5})} d(4\sqrt[6]{x} - \sqrt{5} + 1) - \frac{1}{4} (1+\sqrt{5}) \int \frac{4\sqrt[6]{x} - \sqrt{5} + 1}{2\sqrt[3]{x} + (1+\sqrt{5})\sqrt[6]{x} + 2} d\sqrt[6]{x} \right) \right) \\
& \downarrow 217 \\
& 6 \left(\frac{1}{5} \left(\sqrt{\frac{10}{5+\sqrt{5}}} \arctan \left(\frac{4\sqrt[6]{x} - \sqrt{5} + 1}{\sqrt{2}(5+\sqrt{5})} \right) - \frac{1}{4} (1+\sqrt{5}) \int \frac{4\sqrt[6]{x} - \sqrt{5} + 1}{2\sqrt[3]{x} + (1-\sqrt{5})\sqrt[6]{x} + 2} d\sqrt[6]{x} \right) + \frac{1}{5} \left(-\frac{1}{4} (1-\sqrt{5}) \int \frac{4\sqrt[6]{x} - \sqrt{5} + 1}{2\sqrt[3]{x} + (1+\sqrt{5})\sqrt[6]{x} + 2} d\sqrt[6]{x} \right) \right) \\
& \downarrow 1103 \\
& 6 \left(\frac{1}{5} \left(\sqrt{\frac{10}{5+\sqrt{5}}} \arctan \left(\frac{4\sqrt[6]{x} - \sqrt{5} + 1}{\sqrt{2}(5+\sqrt{5})} \right) - \frac{1}{4} (1+\sqrt{5}) \log(2\sqrt[3]{x} + (1-\sqrt{5})\sqrt[6]{x} + 2) \right) + \frac{1}{5} \left(-\sqrt{\frac{10}{5-\sqrt{5}}} \arctan \left(\frac{4\sqrt[6]{x} - \sqrt{5} + 1}{\sqrt{2}(5-\sqrt{5})} \right) - \frac{1}{4} (1+\sqrt{5}) \log(2\sqrt[3]{x} + (1+\sqrt{5})\sqrt[6]{x} + 2) \right) \right)
\end{aligned}$$

input `Int[(-x^(-1/3) + Sqrt[x])^(-1),x]`

output `6*(Sqrt[x]/3 + Log[1 - x^(1/6)]/5 + (Sqrt[10/(5 + Sqrt[5])]*ArcTan[(1 - Sqrt[5] + 4*x^(1/6))/Sqrt[2*(5 + Sqrt[5])]]) - ((1 + Sqrt[5])*Log[2 + (1 - Sqrt[5])*x^(1/6) + 2*x^(1/3)]/4)/5 + (-Sqrt[10/(5 - Sqrt[5])]*ArcTan[(1 + Sqrt[5] + 4*x^(1/6))/Sqrt[2*(5 - Sqrt[5])]]) - ((1 - Sqrt[5])*Log[2 + (1 + Sqrt[5])*x^(1/6) + 2*x^(1/3)]/4)/5)`

3.220.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 823 `Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Module[{r = Numerator[Rt[-a/b, n]], s = Denominator[Rt[-a/b, n]], k, u}, Simp[u = Int[(r*cos[(2*k - 1)*m*(Pi/n)] + s*cos[(2*k - 1)*(m + 1)*(Pi/n)]*x)/(r^2 + 2*r*s*cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x]; r^(m + 1)/(a*n*s^m) Int[1/(r - s*x), x] - 2*((-r)^(m + 1)/(a*n*s^m)) Sum[u, {k, 1, (n - 1)/2}], x] /; FreeQ[{a, b}, x] && IGtQ[(n - 1)/2, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && NegQ[a/b]`

rule 843 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Simp[a*c^n*((m - n + 1)/(b*(m + n*p + 1))) Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

```
rule 864 Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denomi
nator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a + b*x^(k*n))^(p), x], x, x
^(1/k)], x] /; FreeQ[{a, b, m, p}, x] && FractionQ[n]
```

```
rule 1083 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[I
nt[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x]
```

```
rule 1103 Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

```
rule 1142 Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c)
Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]
```

```
rule 2027 Int[(F*x_.)*((a_.)*(x_)^(r_.) + (b_.)*(x_)^(s_.))^(p_.), x_Symbol] := Int[x^
(p*r)*(a + b*x^(s - r))^p*F*x, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] &
& PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])
```

3.220.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.62

method	result
meijerg	$\frac{6(-1)^{\frac{2}{5}} \left(\frac{5\sqrt{x}(-1)^{\frac{3}{5}}}{3} + (-1)^{\frac{3}{5}} \left(\ln(1-x^{\frac{1}{6}}) - \cos\left(\frac{\pi}{5}\right) \ln\left(1-2\cos\left(\frac{2\pi}{5}\right)x^{\frac{1}{6}}+x^{\frac{1}{3}}\right) + 2\sin\left(\frac{\pi}{5}\right) \arctan\left(\frac{\sin\left(\frac{2\pi}{5}\right)x^{\frac{1}{6}}}{1-\cos\left(\frac{2\pi}{5}\right)x^{\frac{1}{6}}}\right) \right) \right)}{5} + c$
derivativedivides	$2\sqrt{x} + \frac{6\ln(x^{\frac{1}{6}}-1)}{5} + \frac{3(\sqrt{5}-1)\ln(2+x^{\frac{1}{6}}+2x^{\frac{1}{3}}+x^{\frac{1}{6}}\sqrt{5})}{10} + \frac{12\left(-\sqrt{5}+1-\frac{(\sqrt{5}-1)(\sqrt{5}+1)}{4}\right)\arctan\left(\frac{1+4x^{\frac{1}{6}}+\sqrt{5}}{\sqrt{10-2\sqrt{5}}}\right)}{5\sqrt{10-2\sqrt{5}}}$
default	$2\sqrt{x} + \frac{6\ln(x^{\frac{1}{6}}-1)}{5} + \frac{3(\sqrt{5}-1)\ln(2+x^{\frac{1}{6}}+2x^{\frac{1}{3}}+x^{\frac{1}{6}}\sqrt{5})}{10} + \frac{12\left(-\sqrt{5}+1-\frac{(\sqrt{5}-1)(\sqrt{5}+1)}{4}\right)\arctan\left(\frac{1+4x^{\frac{1}{6}}+\sqrt{5}}{\sqrt{10-2\sqrt{5}}}\right)}{5\sqrt{10-2\sqrt{5}}}$

```
input int(1/(-1/x^(1/3)+x^(1/2)),x,method=_RETURNVERBOSE)
```

3.220. $\int \frac{1}{-\frac{1}{\sqrt[3]{x}}+\sqrt{x}} dx$

output $-6/5*(-1)^{(2/5)}*(5/3*x^{(1/2)}*(-1)^{(3/5)}+(-1)^{(3/5)}*(\ln(1-x^{(1/6))}-\cos(1/5*\text{Pi})*\ln(1-2*\cos(2/5*\text{Pi})*x^{(1/6)}+x^{(1/3))}+2*\sin(1/5*\text{Pi})*\arctan(\sin(2/5*\text{Pi})*x^{(1/6)}/(1-\cos(2/5*\text{Pi})*x^{(1/6)})))+\cos(2/5*\text{Pi})*\ln(1+2*\cos(1/5*\text{Pi})*x^{(1/6)}+x^{(1/3))}-2*\sin(2/5*\text{Pi})*\arctan(\sin(1/5*\text{Pi})*x^{(1/6)}/(1+\cos(1/5*\text{Pi})*x^{(1/6)})))$

3.220.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 638 vs. $2(133) = 266$.

Time = 0.91 (sec) , antiderivative size = 638, normalized size of antiderivative = 3.19

$$\int \frac{1}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx$$

$$= \frac{1}{10} \left(3\sqrt{5} - \sqrt{-\frac{27}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} + 1 \right)^2} + \frac{9}{2} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} - 3 \right) \left(\sqrt{2}\sqrt{\sqrt{5}-5} - \sqrt{5} - 1 \right) + \frac{9}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} - \sqrt{5} - 1 \right)^2 \right.$$

$$\left. + 3\sqrt{-\frac{27}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} + 1 \right)^2} + \frac{9}{2} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} - 3 \right) \left(\sqrt{2}\sqrt{\sqrt{5}-5} - \sqrt{5} - 1 \right) - \frac{27}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} + 1 \right) \right.$$

$$\left. + 72x^{\frac{1}{6}} + 36 \right)$$

$$+ \frac{1}{10} \left(3\sqrt{5} + \sqrt{-\frac{27}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} + 1 \right)^2} + \frac{9}{2} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} - 3 \right) \left(\sqrt{2}\sqrt{\sqrt{5}-5} - \sqrt{5} - 1 \right) + \frac{9}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} - \sqrt{5} - 1 \right)^2 \right.$$

$$\left. - 3\sqrt{-\frac{27}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} + 1 \right)^2} + \frac{9}{2} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} - 3 \right) \left(\sqrt{2}\sqrt{\sqrt{5}-5} - \sqrt{5} - 1 \right) - \frac{27}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} + 1 \right) \right.$$

$$\left. + 72x^{\frac{1}{6}} + 36 \right)$$

$$- \frac{3}{10} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} + 1 \right) \log \left(-\frac{9}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} + 1 \right)^2 + 36x^{\frac{1}{6}} \right)$$

$$+ \frac{3}{10} \left(\sqrt{2}\sqrt{\sqrt{5}-5} - \sqrt{5} - 1 \right) \log \left(-\frac{9}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} - \sqrt{5} - 1 \right)^2 + 36x^{\frac{1}{6}} \right)$$

$$+ 2\sqrt{x} + \frac{6}{5} \log \left(x^{\frac{1}{6}} - 1 \right)$$

input `integrate(1/(-1/x^(1/3)+x^(1/2)),x, algorithm="fricas")`

output `1/10*(3*sqrt(5) - sqrt(-27/4*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) + 1)^2 + 9/2*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) - 3)*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1) - 27/4*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1)^2 + 18*sqrt(2)*sqrt(sqrt(5) - 5) + 18*sqrt(5) - 90) - 3*log(9/4*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) + 1)^2 + 9/4*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1)^2 + 3*sqrt(-27/4*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) + 1)^2 + 9/2*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) - 3)*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1) - 27/4*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1)^2 + 18*sqrt(2)*sqrt(sqrt(5) - 5) + 18*sqrt(5) - 90)*(sqrt(5) - 1) + 72*x^(1/6) + 36) + 1/10*(3*sqrt(5) + sqrt(-27/4*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) + 1)^2 + 9/2*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) - 3)*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1) - 27/4*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1)^2 + 18*sqrt(2)*sqrt(sqrt(5) - 5) + 18*sqrt(5) - 90) - 3*log(9/4*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) + 1)^2 + 9/4*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1)^2 - 3*sqrt(-27/4*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) + 1)^2 + 9/2*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) - 3)*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1) - 27/4*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1)^2 + 18*sqrt(2)*sqrt(sqrt(5) - 5) + 18*sqrt(5) - 90)*(sqrt(5) - 1) + 72*x^(1/6) + 36) - 3/10*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) + 1)*log(-9/4*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) + 1)^2 + 36*x^(1/6)) + 3/10*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1)*lo...`

3.220.6 Sympy [F]

$$\int \frac{1}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx = \int \frac{\sqrt[3]{x}}{(\sqrt[6]{x} - 1) (\sqrt[6]{x} + x^{2/3} + \sqrt[3]{x} + \sqrt{x} + 1)} dx$$

input `integrate(1/(-1/x**(1/3)+x**(1/2)),x)`

output `Integral(x**(1/3)/((x**(1/6) - 1)*(x**(1/6) + x**(2/3) + x**(1/3) + sqrt(x) + 1)), x)`

3.220.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 272 vs. $2(133) = 266$.

Time = 0.29 (sec) , antiderivative size = 272, normalized size of antiderivative = 1.36

$$\int \frac{1}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx = -\frac{6}{5} (-1)^{\frac{3}{5}} \log \left((-1)^{\frac{1}{5}} + x^{\frac{1}{6}} \right) \\ - \frac{6\sqrt{5}(-1)^{\frac{3}{5}} \log \left(\frac{\sqrt{5}(-1)^{\frac{1}{5}} + (-1)^{\frac{1}{5}} \sqrt{2\sqrt{5}-10} + (-1)^{\frac{1}{5}} - 4x^{\frac{1}{6}}}{\sqrt{5}(-1)^{\frac{1}{5}} - (-1)^{\frac{1}{5}} \sqrt{2\sqrt{5}-10} + (-1)^{\frac{1}{5}} - 4x^{\frac{1}{6}}} \right)}{5\sqrt{2}\sqrt{5}-10} \\ + \frac{6\sqrt{5}(-1)^{\frac{3}{5}} \log \left(\frac{\sqrt{5}(-1)^{\frac{1}{5}} - (-1)^{\frac{1}{5}} \sqrt{-2\sqrt{5}-10} - (-1)^{\frac{1}{5}} + 4x^{\frac{1}{6}}}{\sqrt{5}(-1)^{\frac{1}{5}} + (-1)^{\frac{1}{5}} \sqrt{-2\sqrt{5}-10} - (-1)^{\frac{1}{5}} + 4x^{\frac{1}{6}}} \right)}{5\sqrt{-2}\sqrt{5}-10} + 2\sqrt{x} \\ + \frac{6 \log \left(-x^{\frac{1}{6}} \left(\sqrt{5}(-1)^{\frac{1}{5}} + (-1)^{\frac{1}{5}} \right) + 2(-1)^{\frac{2}{5}} + 2x^{\frac{1}{3}} \right)}{5 \left(\sqrt{5}(-1)^{\frac{2}{5}} + (-1)^{\frac{2}{5}} \right)} \\ - \frac{6 \log \left(x^{\frac{1}{6}} \left(\sqrt{5}(-1)^{\frac{1}{5}} - (-1)^{\frac{1}{5}} \right) + 2(-1)^{\frac{2}{5}} + 2x^{\frac{1}{3}} \right)}{5 \left(\sqrt{5}(-1)^{\frac{2}{5}} - (-1)^{\frac{2}{5}} \right)}$$

input `integrate(1/(-1/x^(1/3)+x^(1/2)),x, algorithm="maxima")`

output `-6/5*(-1)^(3/5)*log((-1)^(1/5) + x^(1/6)) - 6/5*sqrt(5)*(-1)^(3/5)*log((sqrt(5)*(-1)^(1/5) + (-1)^(1/5)*sqrt(2*sqrt(5) - 10) + (-1)^(1/5) - 4*x^(1/6))/(sqrt(5)*(-1)^(1/5) - (-1)^(1/5)*sqrt(2*sqrt(5) - 10) + (-1)^(1/5) - 4*x^(1/6)))/sqrt(2*sqrt(5) - 10) + 6/5*sqrt(5)*(-1)^(3/5)*log((sqrt(5)*(-1)^(1/5) - (-1)^(1/5)*sqrt(-2*sqrt(5) - 10) - (-1)^(1/5) + 4*x^(1/6))/(sqrt(5)*(-1)^(1/5) + (-1)^(1/5)*sqrt(-2*sqrt(5) - 10) - (-1)^(1/5) + 4*x^(1/6)))/sqrt(-2*sqrt(5) - 10) + 2*sqrt(x) + 6/5*log(-x^(1/6)*(sqrt(5)*(-1)^(1/5) + (-1)^(1/5)) + 2*(-1)^(2/5) + 2*x^(1/3))/(sqrt(5)*(-1)^(2/5) + (-1)^(2/5)) - 6/5*log(x^(1/6)*(sqrt(5)*(-1)^(1/5) - (-1)^(1/5)) + 2*(-1)^(2/5) + 2*x^(1/3))/(sqrt(5)*(-1)^(2/5) - (-1)^(2/5))`

3.220.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.70

$$\int \frac{1}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx = \frac{3}{5} \sqrt{-2\sqrt{5} + 10} \arctan \left(-\frac{\sqrt{5} - 4x^{\frac{1}{6}} - 1}{\sqrt{2\sqrt{5} + 10}} \right) - \frac{3}{5} \sqrt{2\sqrt{5} + 10} \arctan \left(\frac{\sqrt{5} + 4x^{\frac{1}{6}} + 1}{\sqrt{-2\sqrt{5} + 10}} \right) + \frac{3}{10} \sqrt{5} \log \left(\frac{1}{2} x^{\frac{1}{6}} (\sqrt{5} + 1) + x^{\frac{1}{3}} + 1 \right) - \frac{3}{10} \sqrt{5} \log \left(-\frac{1}{2} x^{\frac{1}{6}} (\sqrt{5} - 1) + x^{\frac{1}{3}} + 1 \right) + 2\sqrt{x} - \frac{3}{10} \log \left(x^{\frac{2}{3}} + \sqrt{x} + x^{\frac{1}{3}} + x^{\frac{1}{6}} + 1 \right) + \frac{6}{5} \log \left(\left| x^{\frac{1}{6}} - 1 \right| \right)$$

input `integrate(1/(-1/x^(1/3)+x^(1/2)),x, algorithm="giac")`output `3/5*sqrt(-2*sqrt(5) + 10)*arctan(-(sqrt(5) - 4*x^(1/6) - 1)/sqrt(2*sqrt(5) + 10)) - 3/5*sqrt(2*sqrt(5) + 10)*arctan((sqrt(5) + 4*x^(1/6) + 1)/sqrt(-2*sqrt(5) + 10)) + 3/10*sqrt(5)*log(1/2*x^(1/6)*(sqrt(5) + 1) + x^(1/3) + 1) - 3/10*sqrt(5)*log(-1/2*x^(1/6)*(sqrt(5) - 1) + x^(1/3) + 1) + 2*sqrt(x) - 3/10*log(x^(2/3) + sqrt(x) + x^(1/3) + x^(1/6) + 1) + 6/5*log(abs(x^(1/6) - 1))`**3.220.9 Mupad [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.12

$$\int \frac{1}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx = \frac{6 \ln(1296 x^{1/6} - 1296)}{5} - \ln \left(-750 x^{1/6} \left(\frac{3\sqrt{2}\sqrt{-\sqrt{5}-5}}{10} - \frac{3\sqrt{5}}{10} + \frac{3}{10} \right)^3 - 1296 \right) \left(\frac{3\sqrt{2}\sqrt{-\sqrt{5}-5}}{10} - \frac{3\sqrt{5}}{10} + \frac{3}{10} \right) + \ln \left(750 x^{1/6} \left(\frac{3\sqrt{2}\sqrt{-\sqrt{5}-5}}{10} + \frac{3\sqrt{5}}{10} - \frac{3}{10} \right)^3 - 1296 \right) \left(\frac{3\sqrt{2}\sqrt{-\sqrt{5}-5}}{10} + \frac{3\sqrt{5}}{10} - \frac{3}{10} \right) - \ln \left(-750 x^{1/6} \left(\frac{3\sqrt{2}\sqrt{-\sqrt{5}-5}}{10} + \frac{3\sqrt{5}}{10} - \frac{3}{10} \right)^3 - 1296 \right) \left(\frac{3\sqrt{2}\sqrt{-\sqrt{5}-5}}{10} + \frac{3\sqrt{5}}{10} - \frac{3}{10} \right)$$

input `int(1/(x^(1/2) - 1/x^(1/3)),x)`

output $(6*\log(1296*x^{1/6} - 1296))/5 - \log(-750*x^{1/6}*((3*2^{1/2})*(-5^{1/2} - 5^{1/2}))/10 - (3*5^{1/2})/10 + 3/10)^3 - 1296*((3*2^{1/2})*(-5^{1/2} - 5^{1/2}))/10 - (3*5^{1/2})/10 + 3/10) + \log(750*x^{1/6}*((3*2^{1/2})*(-5^{1/2} - 5^{1/2}))/10 + (3*5^{1/2})/10 - 3/10)^3 - 1296*((3*2^{1/2})*(-5^{1/2} - 5^{1/2}))/10 + (3*5^{1/2})/10 - 3/10) - \log(-750*x^{1/6}*((3*5^{1/2})/10 - (3*2^{1/2})*(5^{1/2} - 5^{1/2}))/10 + 3/10)^3 - 1296*((3*5^{1/2})/10 - (3*2^{1/2})*(5^{1/2} - 5^{1/2}))/10 + 3/10) - \log(-750*x^{1/6}*((3*5^{1/2})/10 + (3*2^{1/2})*(5^{1/2} - 5^{1/2}))/10 + 3/10)^3 - 1296*((3*5^{1/2})/10 + (3*2^{1/2})*(5^{1/2} - 5^{1/2}))/10 + 3/10) + 2*x^{1/2}$

3.220. $\int \frac{1}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx$

$$\mathbf{3.221} \quad \int \frac{1}{-4 \cos(x) + 3 \sin(x)} dx$$

3.221.1 Optimal result	1144
3.221.2 Mathematica [B] (verified)	1144
3.221.3 Rubi [A] (verified)	1145
3.221.4 Maple [A] (verified)	1146
3.221.5 Fricas [B] (verification not implemented)	1146
3.221.6 Sympy [A] (verification not implemented)	1147
3.221.7 Maxima [B] (verification not implemented)	1147
3.221.8 Giac [A] (verification not implemented)	1147
3.221.9 Mupad [B] (verification not implemented)	1148

3.221.1 Optimal result

Integrand size = 11, antiderivative size = 18

$$\int \frac{1}{-4 \cos(x) + 3 \sin(x)} dx = -\frac{1}{5} \operatorname{arctanh} \left(\frac{1}{5} (3 \cos(x) + 4 \sin(x)) \right)$$

output `-1/5*arctanh(3/5*cos(x)+4/5*sin(x))`

3.221.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 41 vs. $2(18) = 36$.

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 2.28

$$\int \frac{1}{-4 \cos(x) + 3 \sin(x)} dx = \frac{1}{5} \log \left(\cos \left(\frac{x}{2} \right) - 2 \sin \left(\frac{x}{2} \right) \right) - \frac{1}{5} \log \left(2 \cos \left(\frac{x}{2} \right) + \sin \left(\frac{x}{2} \right) \right)$$

input `Integrate[(-4*Cos[x] + 3*Sin[x])^(-1), x]`

output `Log[Cos[x/2] - 2*Sin[x/2]]/5 - Log[2*Cos[x/2] + Sin[x/2]]/5`

3.221.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 3553, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{3 \sin(x) - 4 \cos(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{3 \sin(x) - 4 \cos(x)} dx \\ & \quad \downarrow \text{3553} \\ & - \int \frac{1}{25 - (3 \cos(x) + 4 \sin(x))^2} d(3 \cos(x) + 4 \sin(x)) \\ & \quad \downarrow \text{219} \\ & -\frac{1}{5} \operatorname{arctanh}\left(\frac{1}{5}(4 \sin(x) + 3 \cos(x))\right) \end{aligned}$$

input `Int[(-4*Cos[x] + 3*Sin[x])^(-1),x]`

output `-1/5*ArcTanh[(3*Cos[x] + 4*Sin[x])/5]`

3.221.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3553 Int[(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x
_Symbol] := Simp[-d^(-1) Subst[Int[1/(a^2 + b^2 - x^2), x], x, b*Cos[c +
d*x] - a*Sin[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]
```

3.221.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

method	result	size
default	$\frac{\ln(2 \tan(\frac{x}{2}) - 1)}{5} - \frac{\ln(\tan(\frac{x}{2}) + 2)}{5}$	22
norman	$\frac{\ln(2 \tan(\frac{x}{2}) - 1)}{5} - \frac{\ln(\tan(\frac{x}{2}) + 2)}{5}$	22
parallelrisch	$\ln\left(\frac{1}{(2 \tan(\frac{x}{2}) + 4)^{\frac{1}{5}}}\right) + \ln\left((2 \tan(\frac{x}{2}) - 1)^{\frac{1}{5}}\right)$	24
risch	$-\frac{\ln(e^{ix} + \frac{3}{5} + \frac{4i}{5})}{5} + \frac{\ln(e^{ix} - \frac{3}{5} - \frac{4i}{5})}{5}$	26

```
input int(1/(-4*cos(x)+3*sin(x)),x,method=_RETURNVERBOSE)
```

```
output 1/5*ln(2*tan(1/2*x)-1)-1/5*ln(tan(1/2*x)+2)
```

3.221.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(12) = 24.

Time = 0.25 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.50

$$\int \frac{1}{-4 \cos(x) + 3 \sin(x)} dx = -\frac{1}{10} \log\left(\frac{3}{2} \cos(x) + 2 \sin(x) + \frac{5}{2}\right) + \frac{1}{10} \log\left(-\frac{3}{2} \cos(x) - 2 \sin(x) + \frac{5}{2}\right)$$

```
input integrate(1/(-4*cos(x)+3*sin(x)),x, algorithm="fricas")
```

```
output -1/10*log(3/2*cos(x) + 2*sin(x) + 5/2) + 1/10*log(-3/2*cos(x) - 2*sin(x) +
5/2)
```

3.221.6 Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{1}{-4 \cos(x) + 3 \sin(x)} dx = -\frac{\log\left(\tan\left(\frac{x}{2}\right) + 2\right)}{5} + \frac{\log\left(2 \tan\left(\frac{x}{2}\right) - 1\right)}{5}$$

input `integrate(1/(-4*cos(x)+3*sin(x)),x)`

output `-log(tan(x/2) + 2)/5 + log(2*tan(x/2) - 1)/5`

3.221.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 30 vs. 2(12) = 24.

Time = 0.20 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.67

$$\int \frac{1}{-4 \cos(x) + 3 \sin(x)} dx = \frac{1}{5} \log\left(\frac{2 \sin(x)}{\cos(x) + 1} - 1\right) - \frac{1}{5} \log\left(\frac{\sin(x)}{\cos(x) + 1} + 2\right)$$

input `integrate(1/(-4*cos(x)+3*sin(x)),x, algorithm="maxima")`

output `1/5*log(2*sin(x)/(cos(x) + 1) - 1) - 1/5*log(sin(x)/(cos(x) + 1) + 2)`

3.221.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.28

$$\int \frac{1}{-4 \cos(x) + 3 \sin(x)} dx = \frac{1}{5} \log\left(\left|2 \tan\left(\frac{1}{2}x\right) - 1\right|\right) - \frac{1}{5} \log\left(\left|\tan\left(\frac{1}{2}x\right) + 2\right|\right)$$

input `integrate(1/(-4*cos(x)+3*sin(x)),x, algorithm="giac")`

output `1/5*log(abs(2*tan(1/2*x) - 1)) - 1/5*log(abs(tan(1/2*x) + 2))`

3.221.9 Mupad [B] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.61

$$\int \frac{1}{-4 \cos(x) + 3 \sin(x)} dx = -\frac{2 \operatorname{atanh}\left(\frac{4 \tan\left(\frac{x}{2}\right) + \frac{3}{5}}{5}\right)}{5}$$

input `int(-1/(4*cos(x) - 3*sin(x)),x)`

output `-(2*atanh((4*tan(x/2))/5 + 3/5))/5`

3.222 $\int \frac{1}{1+\sqrt{x}} dx$

3.222.1 Optimal result	1149
3.222.2 Mathematica [A] (verified)	1149
3.222.3 Rubi [A] (verified)	1150
3.222.4 Maple [A] (verified)	1151
3.222.5 Fricas [A] (verification not implemented)	1151
3.222.6 Sympy [A] (verification not implemented)	1151
3.222.7 Maxima [A] (verification not implemented)	1152
3.222.8 Giac [A] (verification not implemented)	1152
3.222.9 Mupad [B] (verification not implemented)	1152

3.222.1 Optimal result

Integrand size = 9, antiderivative size = 18

$$\int \frac{1}{1+\sqrt{x}} dx = 2\sqrt{x} - 2\log(1+\sqrt{x})$$

output `-2*ln(1+x^(1/2))+2*x^(1/2)`

3.222.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+\sqrt{x}} dx = 2\sqrt{x} - 2\log(1+\sqrt{x})$$

input `Integrate[(1 + Sqrt[x])^(-1),x]`

output `2*Sqrt[x] - 2*Log[1 + Sqrt[x]]`

3.222.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {774, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{1}{\sqrt{x}+1} dx \\
 \downarrow 774 \\
 2 \int \frac{\sqrt{x}}{\sqrt{x}+1} d\sqrt{x} \\
 \downarrow 49 \\
 2 \int \left(1 + \frac{1}{-\sqrt{x}-1}\right) d\sqrt{x} \\
 \downarrow 2009 \\
 2(\sqrt{x} - \log(\sqrt{x}+1))
 \end{array}$$

input `Int[(1 + Sqrt[x])^(-1),x]`

output `2*(Sqrt[x] - Log[1 + Sqrt[x]])`

3.222.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 774 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> With[{k = Denominator[n]}, Simp[k Subst[Int[x^(k - 1)*(a + b*x^(k*n))^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, p}, x] && FractionQ[n]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.222.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

method	result	size
derivativdivides	$-2 \ln(\sqrt{x} + 1) + 2\sqrt{x}$	15
meijerg	$-2 \ln(\sqrt{x} + 1) + 2\sqrt{x}$	15
trager	$2\sqrt{x} - \ln(2\sqrt{x} + 1 + x)$	18
default	$2\sqrt{x} + \ln(\sqrt{x} - 1) - \ln(\sqrt{x} + 1) - \ln(-1 + x)$	27

input `int(1/(x^(1/2)+1),x,method=_RETURNVERBOSE)`output `-2*ln(x^(1/2)+1)+2*x^(1/2)`**3.222.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{1}{1 + \sqrt{x}} dx = 2\sqrt{x} - 2 \log(\sqrt{x} + 1)$$

input `integrate(1/(1+x^(1/2)),x, algorithm="fricas")`output `2*sqrt(x) - 2*log(sqrt(x) + 1)`**3.222.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{1}{1 + \sqrt{x}} dx = 2\sqrt{x} - 2 \log(\sqrt{x} + 1)$$

input `integrate(1/(1+x**(1/2)),x)`output `2*sqrt(x) - 2*log(sqrt(x) + 1)`

3.222.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{1}{1+\sqrt{x}} dx = 2\sqrt{x} - 2 \log(\sqrt{x} + 1) + 2$$

input `integrate(1/(1+x^(1/2)),x, algorithm="maxima")`output `2*sqrt(x) - 2*log(sqrt(x) + 1) + 2`**3.222.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{1}{1+\sqrt{x}} dx = 2\sqrt{x} - 2 \log(\sqrt{x} + 1)$$

input `integrate(1/(1+x^(1/2)),x, algorithm="giac")`output `2*sqrt(x) - 2*log(sqrt(x) + 1)`**3.222.9 Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{1}{1+\sqrt{x}} dx = 2\sqrt{x} - 2 \ln(\sqrt{x} + 1)$$

input `int(1/(x^(1/2) + 1),x)`output `2*x^(1/2) - 2*log(x^(1/2) + 1)`

$$3.223 \quad \int \frac{1}{1 + \frac{1}{\sqrt[3]{x}}} dx$$

3.223.1 Optimal result	1153
3.223.2 Mathematica [A] (verified)	1153
3.223.3 Rubi [A] (verified)	1154
3.223.4 Maple [A] (verified)	1155
3.223.5 Fricas [A] (verification not implemented)	1156
3.223.6 Sympy [A] (verification not implemented)	1156
3.223.7 Maxima [A] (verification not implemented)	1156
3.223.8 Giac [A] (verification not implemented)	1157
3.223.9 Mupad [B] (verification not implemented)	1157

3.223.1 Optimal result

Integrand size = 9, antiderivative size = 32

$$\int \frac{1}{1 + \frac{1}{\sqrt[3]{x}}} dx = 3\sqrt[3]{x} - \frac{3x^{2/3}}{2} + x - 3 \log \left(1 + \frac{1}{\sqrt[3]{x}} \right) - \log(x)$$

output `3*x^(1/3)-3/2*x^(2/3)+x-3*ln(1+1/x^(1/3))-ln(x)`

3.223.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.88

$$\int \frac{1}{1 + \frac{1}{\sqrt[3]{x}}} dx = 3\sqrt[3]{x} - \frac{3x^{2/3}}{2} + x - 3 \log (1 + \sqrt[3]{x})$$

input `Integrate[(1 + x^(-1/3))^(-1), x]`

output `3*x^(1/3) - (3*x^(2/3))/2 + x - 3*Log[1 + x^(1/3)]`

3.223.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {774, 795, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt[3]{x} + 1} dx \\
 & \quad \downarrow 774 \\
 & 3 \int \frac{x^{2/3}}{1 + \frac{1}{\sqrt[3]{x}}} d\sqrt[3]{x} \\
 & \quad \downarrow 795 \\
 & 3 \int \frac{x}{\sqrt[3]{x} + 1} d\sqrt[3]{x} \\
 & \quad \downarrow 49 \\
 & 3 \int \left(x^{2/3} - \sqrt[3]{x} + \frac{1}{-\sqrt[3]{x} - 1} + 1 \right) d\sqrt[3]{x} \\
 & \quad \downarrow 2009 \\
 & 3 \left(-\frac{x^{2/3}}{2} + \frac{x}{3} + \sqrt[3]{x} - \log(\sqrt[3]{x} + 1) \right)
 \end{aligned}$$

input `Int[(1 + x^(-1/3))^-1, x]`

output `3*(x^(1/3) - x^(2/3)/2 + x/3 - Log[1 + x^(1/3)])`

3.223.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 774 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[n]},
Simp[k Subst[Int[x^(k - 1)*(a + b*x^(k*n))^p, x], x, x^(1/k)], x] /; Fre
eQ[{a, b, p}, x] && FractionQ[n]`

rule 795 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Int[x^(m + n*p)*
(b + a/x^n)^p, x] /; FreeQ[{a, b, m, n}, x] && IntegerQ[p] && NegQ[n]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.223.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.66

method	result	size
derivativedivides	$x - \frac{3x^{\frac{2}{3}}}{2} + 3x^{\frac{1}{3}} - 3 \ln(x^{\frac{1}{3}} + 1)$	21
default	$x - \frac{3x^{\frac{2}{3}}}{2} + 3x^{\frac{1}{3}} - 3 \ln(x^{\frac{1}{3}} + 1)$	21
meijerg	$\frac{x^{\frac{1}{3}}(4x^{\frac{2}{3}} - 6x^{\frac{1}{3}} + 12)}{4} - 3 \ln(x^{\frac{1}{3}} + 1)$	27
trager	$-1 + x + 3x^{\frac{1}{3}} - \frac{3x^{\frac{2}{3}}}{2} - \ln(-3x^{\frac{2}{3}} - 3x^{\frac{1}{3}} - x - 1)$	32

input `int(1/(1+1/x^(1/3)),x,method=_RETURNVERBOSE)`

output `x-3/2*x^(2/3)+3*x^(1/3)-3*ln(x^(1/3)+1)`

3.223.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.62

$$\int \frac{1}{1 + \frac{1}{\sqrt[3]{x}}} dx = x - \frac{3}{2} x^{\frac{2}{3}} + 3x^{\frac{1}{3}} - 3 \log(x^{\frac{1}{3}} + 1)$$

input `integrate(1/(1+1/x^(1/3)),x, algorithm="fricas")`output `x - 3/2*x^(2/3) + 3*x^(1/3) - 3*log(x^(1/3) + 1)`**3.223.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

$$\int \frac{1}{1 + \frac{1}{\sqrt[3]{x}}} dx = -\frac{3x^{\frac{2}{3}}}{2} + 3\sqrt[3]{x} + x - 3 \log(\sqrt[3]{x} + 1)$$

input `integrate(1/(1+1/x**(1/3)),x)`output `-3*x**(2/3)/2 + 3*x**(1/3) + x - 3*log(x**(1/3) + 1)`**3.223.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.88

$$\int \frac{1}{1 + \frac{1}{\sqrt[3]{x}}} dx = -\frac{1}{2} x \left(\frac{3}{x^{\frac{1}{3}}} - \frac{6}{x^{\frac{2}{3}}} - 2 \right) - \log(x) - 3 \log\left(\frac{1}{x^{\frac{1}{3}}} + 1\right)$$

input `integrate(1/(1+1/x^(1/3)),x, algorithm="maxima")`output `-1/2*x*(3/x^(1/3) - 6/x^(2/3) - 2) - log(x) - 3*log(1/x^(1/3) + 1)`

3.223.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.62

$$\int \frac{1}{1 + \frac{1}{\sqrt[3]{x}}} dx = x - \frac{3}{2} x^{\frac{2}{3}} + 3x^{\frac{1}{3}} - 3 \log(x^{\frac{1}{3}} + 1)$$

input `integrate(1/(1+1/x^(1/3)),x, algorithm="giac")`output `x - 3/2*x^(2/3) + 3*x^(1/3) - 3*log(x^(1/3) + 1)`**3.223.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.62

$$\int \frac{1}{1 + \frac{1}{\sqrt[3]{x}}} dx = x - 3 \ln(x^{1/3} + 1) + 3x^{1/3} - \frac{3x^{2/3}}{2}$$

input `int(1/(1/x^(1/3) + 1),x)`output `x - 3*log(x^(1/3) + 1) + 3*x^(1/3) - (3*x^(2/3))/2`

3.224 $\int \frac{\sqrt{x}}{1+x} dx$

3.224.1 Optimal result	1158
3.224.2 Mathematica [A] (verified)	1158
3.224.3 Rubi [A] (verified)	1159
3.224.4 Maple [A] (verified)	1160
3.224.5 Fricas [A] (verification not implemented)	1160
3.224.6 Sympy [A] (verification not implemented)	1161
3.224.7 Maxima [A] (verification not implemented)	1161
3.224.8 Giac [A] (verification not implemented)	1161
3.224.9 Mupad [B] (verification not implemented)	1162

3.224.1 Optimal result

Integrand size = 11, antiderivative size = 16

$$\int \frac{\sqrt{x}}{1+x} dx = 2\sqrt{x} - 2 \arctan(\sqrt{x})$$

output `-2*arctan(x^(1/2))+2*x^(1/2)`

3.224.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{x}}{1+x} dx = 2\sqrt{x} - 2 \arctan(\sqrt{x})$$

input `Integrate[Sqrt[x]/(1 + x),x]`

output `2*Sqrt[x] - 2*ArcTan[Sqrt[x]]`

3.224.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{x}}{x+1} dx \\ & \quad \downarrow \text{60} \\ & 2\sqrt{x} - \int \frac{1}{\sqrt{x}(x+1)} dx \\ & \quad \downarrow \text{73} \\ & 2\sqrt{x} - 2 \int \frac{1}{x+1} d\sqrt{x} \\ & \quad \downarrow \text{216} \\ & 2\sqrt{x} - 2 \arctan(\sqrt{x}) \end{aligned}$$

input `Int[Sqrt[x]/(1 + x),x]`

output `2*Sqrt[x] - 2*ArcTan[Sqrt[x]]`

3.224.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
 rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
 , 0] || GtQ[b, 0])`

3.224.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$-2 \arctan(\sqrt{x}) + 2\sqrt{x}$	13
default	$-2 \arctan(\sqrt{x}) + 2\sqrt{x}$	13
meijerg	$-2 \arctan(\sqrt{x}) + 2\sqrt{x}$	13
risch	$-2 \arctan(\sqrt{x}) + 2\sqrt{x}$	13
trager	$2\sqrt{x} + \text{RootOf}(-Z^2 + 1) \ln\left(-\frac{2\text{RootOf}(-Z^2 + 1)\sqrt{x-x+1}}{1+x}\right)$	38

input `int(x^(1/2)/(1+x),x,method=_RETURNVERBOSE)`

output `-2*arctan(x^(1/2))+2*x^(1/2)`

3.224.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{x}}{1+x} dx = 2\sqrt{x} - 2 \arctan(\sqrt{x})$$

input `integrate(x^(1/2)/(1+x),x, algorithm="fricas")`

output `2*sqrt(x) - 2*arctan(sqrt(x))`

3.224.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{\sqrt{x}}{1+x} dx = 2\sqrt{x} - 2 \operatorname{atan}(\sqrt{x})$$

input `integrate(x**(1/2)/(1+x),x)`output `2*sqrt(x) - 2*atan(sqrt(x))`**3.224.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{x}}{1+x} dx = 2\sqrt{x} - 2 \operatorname{arctan}(\sqrt{x})$$

input `integrate(x^(1/2)/(1+x),x, algorithm="maxima")`output `2*sqrt(x) - 2*arctan(sqrt(x))`**3.224.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{x}}{1+x} dx = 2\sqrt{x} - 2 \operatorname{arctan}(\sqrt{x})$$

input `integrate(x^(1/2)/(1+x),x, algorithm="giac")`output `2*sqrt(x) - 2*arctan(sqrt(x))`

3.224.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{x}}{1+x} dx = 2\sqrt{x} - 2\operatorname{atan}(\sqrt{x})$$

input `int(x^(1/2)/(x + 1),x)`

output `2*x^(1/2) - 2*atan(x^(1/2))`

3.225 $\int \frac{1}{x\sqrt{1+x}} dx$

3.225.1 Optimal result	1163
3.225.2 Mathematica [A] (verified)	1163
3.225.3 Rubi [A] (verified)	1164
3.225.4 Maple [A] (verified)	1165
3.225.5 Fricas [B] (verification not implemented)	1165
3.225.6 Sympy [B] (verification not implemented)	1165
3.225.7 Maxima [B] (verification not implemented)	1166
3.225.8 Giac [B] (verification not implemented)	1166
3.225.9 Mupad [B] (verification not implemented)	1167

3.225.1 Optimal result

Integrand size = 11, antiderivative size = 10

$$\int \frac{1}{x\sqrt{1+x}} dx = -2\operatorname{arctanh}(\sqrt{1+x})$$

output `-2*arctanh((1+x)^(1/2))`

3.225.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{1+x}} dx = -2\operatorname{arctanh}(\sqrt{1+x})$$

input `Integrate[1/(x*Sqrt[1 + x]),x]`

output `-2*ArcTanh[Sqrt[1 + x]]`

3.225.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {73, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x\sqrt{x+1}} dx$$

↓ 73

$$2 \int \frac{1}{x} d\sqrt{x+1}$$

↓ 220

$$-2\operatorname{arctanh}(\sqrt{x+1})$$

input `Int[1/(x*Sqrt[1 + x]),x]`

output `-2*ArcTanh[Sqrt[1 + x]]`

3.225.3.1 Defintions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]`

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-Rt[-a, 2]*Rt[b, 2])^(-
1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] &&
(LtQ[a, 0] || GtQ[b, 0])`

3.225.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

method	result	size
derivativedivides	$-2 \operatorname{arctanh}(\sqrt{1+x})$	9
default	$-2 \operatorname{arctanh}(\sqrt{1+x})$	9
trager	$-\ln\left(\frac{2\sqrt{1+x}+2+x}{x}\right)$	18
meijerg	$\frac{(-2\ln(2)+\ln(x))\sqrt{\pi}-2\sqrt{\pi}\ln\left(\frac{1}{2}+\frac{\sqrt{1+x}}{2}\right)}{\sqrt{\pi}}$	32

input `int(1/x/(1+x)^(1/2),x,method=_RETURNVERBOSE)`output `-2*arctanh((1+x)^(1/2))`**3.225.5 Fracas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(8) = 16$.

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.90

$$\int \frac{1}{x\sqrt{1+x}} dx = -\log(\sqrt{x+1} + 1) + \log(\sqrt{x+1} - 1)$$

input `integrate(1/x/(1+x)^(1/2),x, algorithm="fracas")`output `-log(sqrt(x + 1) + 1) + log(sqrt(x + 1) - 1)`**3.225.6 Sympy [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 26 vs. $2(10) = 20$.

Time = 0.35 (sec) , antiderivative size = 26, normalized size of antiderivative = 2.60

$$\int \frac{1}{x\sqrt{1+x}} dx = \begin{cases} -2 \operatorname{acoth}(\sqrt{x+1}) & \text{for } |x+1| > 1 \\ -2 \operatorname{atanh}(\sqrt{x+1}) & \text{otherwise} \end{cases}$$

input `integrate(1/x/(1+x)**(1/2),x)`

output `Piecewise((-2*acoth(sqrt(x + 1)), Abs(x + 1) > 1), (-2*atanh(sqrt(x + 1)), True))`

3.225.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(8) = 16$.

Time = 0.21 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.90

$$\int \frac{1}{x\sqrt{1+x}} dx = -\log(\sqrt{x+1} + 1) + \log(\sqrt{x+1} - 1)$$

input `integrate(1/x/(1+x)^(1/2),x, algorithm="maxima")`

output `-log(sqrt(x + 1) + 1) + log(sqrt(x + 1) - 1)`

3.225.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. $2(8) = 16$.

Time = 0.28 (sec) , antiderivative size = 20, normalized size of antiderivative = 2.00

$$\int \frac{1}{x\sqrt{1+x}} dx = -\log(\sqrt{x+1} + 1) + \log(|\sqrt{x+1} - 1|)$$

input `integrate(1/x/(1+x)^(1/2),x, algorithm="giac")`

output `-log(sqrt(x + 1) + 1) + log(abs(sqrt(x + 1) - 1))`

3.225.9 Mupad [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{1}{x\sqrt{1+x}} dx = -2 \operatorname{atanh}(\sqrt{x+1})$$

input `int(1/(x*(x + 1)^(1/2)),x)`

output `-2*atanh((x + 1)^(1/2))`

$$3.226 \quad \int \frac{1}{-\sqrt[3]{x+x}} dx$$

3.226.1 Optimal result	1168
3.226.2 Mathematica [A] (verified)	1168
3.226.3 Rubi [A] (verified)	1169
3.226.4 Maple [A] (verified)	1170
3.226.5 Fracas [A] (verification not implemented)	1170
3.226.6 Sympy [B] (verification not implemented)	1170
3.226.7 Maxima [A] (verification not implemented)	1171
3.226.8 Giac [A] (verification not implemented)	1171
3.226.9 Mupad [B] (verification not implemented)	1171

3.226.1 Optimal result

Integrand size = 11, antiderivative size = 14

$$\int \frac{1}{-\sqrt[3]{x+x}} dx = \frac{3}{2} \log(1 - x^{2/3})$$

output `3/2*ln(1-x^(2/3))`

3.226.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.79

$$\int \frac{1}{-\sqrt[3]{x+x}} dx = \frac{3}{2} \log(-1 + \sqrt[3]{x}) + \frac{3}{2} \log(1 + \sqrt[3]{x})$$

input `Integrate[(-x^(1/3) + x)^(-1),x]`

output `(3*Log[-1 + x^(1/3)])/2 + (3*Log[1 + x^(1/3)])/2`

3.226.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2027, 792}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x - \sqrt[3]{x}} dx$$

↓ 2027

$$\int \frac{1}{(x^{2/3} - 1)\sqrt[3]{x}} dx$$

↓ 792

$$\frac{3}{2} \log(1 - x^{2/3})$$

input `Int[(-x^(1/3) + x)^(-1),x]`

output `(3*Log[1 - x^(2/3)])/2`

3.226.3.1 Defintions of rubi rules used

rule 792 `Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]`

rule 2027 `Int[(Fx_)*((a_)*(x_)^(r_) + (b_)*(x_)^(s_))^(p_), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r))^p*Fx, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] && PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])`

3.226.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

method	result	size
meijerg	$\frac{3 \ln(1-x^{\frac{2}{3}})}{2}$	11
derivativedivides	$\frac{3 \ln(x^{\frac{1}{3}}-1)}{2} + \frac{3 \ln(x^{\frac{1}{3}}+1)}{2}$	18
trager	$\frac{\ln(3x^{\frac{2}{3}}-3x^{\frac{4}{3}}+x^2-1)}{2}$	19
default	$\frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2} - \frac{\ln(x^{\frac{2}{3}}+x^{\frac{1}{3}}+1)}{2} + \ln(x^{\frac{1}{3}}-1) + \ln(x^{\frac{1}{3}}+1) - \frac{\ln(x^{\frac{2}{3}}-x^{\frac{1}{3}}+1)}{2}$	50

input `int(1/(-x^(1/3)+x),x,method=_RETURNVERBOSE)`output `3/2*ln(1-x^(2/3))`**3.226.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.57

$$\int \frac{1}{-\sqrt[3]{x}+x} dx = \frac{3}{2} \log(x^{\frac{2}{3}}-1)$$

input `integrate(1/(-x^(1/3)+x),x, algorithm="fricas")`output `3/2*log(x^(2/3) - 1)`**3.226.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 22 vs. 2(10) = 20.

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.57

$$\int \frac{1}{-\sqrt[3]{x}+x} dx = \frac{3 \log(\sqrt[3]{x}-1)}{2} + \frac{3 \log(\sqrt[3]{x}+1)}{2}$$

input `integrate(1/(-x**(1/3)+x),x)`

output `3*log(x**(1/3) - 1)/2 + 3*log(x**(1/3) + 1)/2`

3.226.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.21

$$\int \frac{1}{-\sqrt[3]{x} + x} dx = \frac{3}{2} \log(x^{\frac{1}{3}} + 1) + \frac{3}{2} \log(x^{\frac{1}{3}} - 1)$$

input `integrate(1/(-x^(1/3)+x),x, algorithm="maxima")`

output `3/2*log(x^(1/3) + 1) + 3/2*log(x^(1/3) - 1)`

3.226.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.29

$$\int \frac{1}{-\sqrt[3]{x} + x} dx = \frac{3}{2} \log(x^{\frac{1}{3}} + 1) + \frac{3}{2} \log(|x^{\frac{1}{3}} - 1|)$$

input `integrate(1/(-x^(1/3)+x),x, algorithm="giac")`

output `3/2*log(x^(1/3) + 1) + 3/2*log(abs(x^(1/3) - 1))`

3.226.9 Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.57

$$\int \frac{1}{-\sqrt[3]{x} + x} dx = \frac{3 \ln(x^{2/3} - 1)}{2}$$

input `int(1/(x - x^(1/3)),x)`

output `(3*log(x^(2/3) - 1))/2`

$$3.227 \quad \int \frac{1}{x - \sqrt{2+x}} dx$$

3.227.1 Optimal result	1172
3.227.2 Mathematica [A] (verified)	1172
3.227.3 Rubi [A] (verified)	1173
3.227.4 Maple [A] (verified)	1174
3.227.5 Fricas [A] (verification not implemented)	1174
3.227.6 Sympy [A] (verification not implemented)	1175
3.227.7 Maxima [A] (verification not implemented)	1175
3.227.8 Giac [A] (verification not implemented)	1175
3.227.9 Mupad [B] (verification not implemented)	1176

3.227.1 Optimal result

Integrand size = 13, antiderivative size = 31

$$\int \frac{1}{x - \sqrt{2+x}} dx = \frac{4}{3} \log(2 - \sqrt{2+x}) + \frac{2}{3} \log(1 + \sqrt{2+x})$$

output `4/3*ln(2-(2+x)^(1/2))+2/3*ln(1+(2+x)^(1/2))`

3.227.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{1}{x - \sqrt{2+x}} dx = \frac{4}{3} \log(-2 + \sqrt{2+x}) + \frac{2}{3} \log(1 + \sqrt{2+x})$$

input `Integrate[(x - Sqrt[2 + x])^(-1),x]`

output `(4*Log[-2 + Sqrt[2 + x]])/3 + (2*Log[1 + Sqrt[2 + x]])/3`

3.227.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {7267, 25, 1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x - \sqrt{x+2}} dx \\
 & \quad \downarrow \text{7267} \\
 & 2 \int -\frac{\sqrt{x+2}}{\sqrt{x+2} - x} d\sqrt{x+2} \\
 & \quad \downarrow \text{25} \\
 & -2 \int \frac{\sqrt{x+2}}{\sqrt{x+2} - x} d\sqrt{x+2} \\
 & \quad \downarrow \text{1141} \\
 & 2 \int \left(\frac{1}{3(\sqrt{x+2} + 1)} - \frac{2}{3(2 - \sqrt{x+2})} \right) d\sqrt{x+2} \\
 & \quad \downarrow \text{2009} \\
 & 2 \left(\frac{2}{3} \log(2 - \sqrt{x+2}) + \frac{1}{3} \log(\sqrt{x+2} + 1) \right)
 \end{aligned}$$

input `Int[(x - Sqrt[2 + x])^(-1),x]`

output `2*((2*Log[2 - Sqrt[2 + x]])/3 + Log[1 + Sqrt[2 + x]]/3)`

3.227.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

```
rule 1141 Int[((d_.) + (e_.)*(x_)^(m_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_
Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[
(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x] /; EqQ[p, -
1] || !FractionalPowerFactorQ[q]] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p,
0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]
```

```
rule 2009 Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 7267 Int[u_, x_Symbol] :> With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]]
```

3.227.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.71

method	result	size
derivativedivides	$\frac{2 \ln(1+\sqrt{2+x})}{3} + \frac{4 \ln(\sqrt{2+x}-2)}{3}$	22
trager	$\frac{\ln(6\sqrt{2+x}x^2-x^3+16\sqrt{2+x}x-15x^2+8\sqrt{2+x}-24x-12)}{3}$	44
default	$\frac{\ln(1+x)}{3} + \frac{2 \ln(-2+x)}{3} + \frac{\ln(1+\sqrt{2+x})}{3} - \frac{\ln(\sqrt{2+x}-1)}{3} + \frac{2 \ln(\sqrt{2+x}-2)}{3} - \frac{2 \ln(\sqrt{2+x}+2)}{3}$	54

```
input int(1/(x-(2+x)^(1/2)),x,method=_RETURNVERBOSE)
```

```
output 2/3*ln(1+(2+x)^(1/2))+4/3*ln((2+x)^(1/2)-2)
```

3.227.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.68

$$\int \frac{1}{x - \sqrt{2+x}} dx = \frac{2}{3} \log(\sqrt{x+2} + 1) + \frac{4}{3} \log(\sqrt{x+2} - 2)$$

```
input integrate(1/(x-(2+x)^(1/2)),x, algorithm="fricas")
```

```
output 2/3*log(sqrt(x + 2) + 1) + 4/3*log(sqrt(x + 2) - 2)
```

3.227.6 Sympy [A] (verification not implemented)

Time = 0.66 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.16

$$\int \frac{1}{x - \sqrt{2+x}} dx = \log(x - \sqrt{x+2}) + \frac{\log(2\sqrt{x+2} - 4)}{3} - \frac{\log(2\sqrt{x+2} + 2)}{3}$$

input `integrate(1/(x-(2+x)**(1/2)),x)`output `log(x - sqrt(x + 2)) + log(2*sqrt(x + 2) - 4)/3 - log(2*sqrt(x + 2) + 2)/3`**3.227.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.68

$$\int \frac{1}{x - \sqrt{2+x}} dx = \frac{2}{3} \log(\sqrt{x+2} + 1) + \frac{4}{3} \log(\sqrt{x+2} - 2)$$

input `integrate(1/(x-(2+x)^(1/2)),x, algorithm="maxima")`output `2/3*log(sqrt(x + 2) + 1) + 4/3*log(sqrt(x + 2) - 2)`**3.227.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.71

$$\int \frac{1}{x - \sqrt{2+x}} dx = \frac{2}{3} \log(\sqrt{x+2} + 1) + \frac{4}{3} \log(|\sqrt{x+2} - 2|)$$

input `integrate(1/(x-(2+x)^(1/2)),x, algorithm="giac")`output `2/3*log(sqrt(x + 2) + 1) + 4/3*log(abs(sqrt(x + 2) - 2))`

3.227.9 Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.81

$$\int \frac{1}{x - \sqrt{2+x}} dx = \frac{2 \ln \left(\frac{2\sqrt{x+2}}{3} + \frac{2}{3} \right)}{3} + \frac{4 \ln \left(\frac{4}{3} - \frac{2\sqrt{x+2}}{3} \right)}{3}$$

input `int(1/(x - (x + 2)^(1/2)),x)`output `(2*log((2*(x + 2)^(1/2))/3 + 2/3))/3 + (4*log(4/3 - (2*(x + 2)^(1/2))/3))/3`

$$3.228 \quad \int \frac{x^2}{\sqrt{-1+x}} dx$$

3.228.1 Optimal result	1177
3.228.2 Mathematica [A] (verified)	1177
3.228.3 Rubi [A] (verified)	1178
3.228.4 Maple [A] (verified)	1179
3.228.5 Fricas [A] (verification not implemented)	1179
3.228.6 Sympy [C] (verification not implemented)	1180
3.228.7 Maxima [A] (verification not implemented)	1180
3.228.8 Giac [A] (verification not implemented)	1180
3.228.9 Mupad [B] (verification not implemented)	1181

3.228.1 Optimal result

Integrand size = 11, antiderivative size = 32

$$\int \frac{x^2}{\sqrt{-1+x}} dx = 2\sqrt{-1+x} + \frac{4}{3}(-1+x)^{3/2} + \frac{2}{5}(-1+x)^{5/2}$$

output `4/3*(-1+x)^(3/2)+2/5*(-1+x)^(5/2)+2*(-1+x)^(1/2)`

3.228.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.66

$$\int \frac{x^2}{\sqrt{-1+x}} dx = \frac{2}{15}\sqrt{-1+x}(8+4x+3x^2)$$

input `Integrate[x^2/Sqrt[-1 + x],x]`

output `(2*Sqrt[-1 + x]*(8 + 4*x + 3*x^2))/15`

3.228.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{\sqrt{x-1}} dx$$

↓ 53

$$\int \left((x-1)^{3/2} + 2\sqrt{x-1} + \frac{1}{\sqrt{x-1}} \right) dx$$

↓ 2009

$$\frac{2}{5}(x-1)^{5/2} + \frac{4}{3}(x-1)^{3/2} + 2\sqrt{x-1}$$

input `Int[x^2/Sqrt[-1 + x],x]`

output `2*Sqrt[-1 + x] + (4*(-1 + x)^(3/2))/3 + (2*(-1 + x)^(5/2))/5`

3.228.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.228.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.53

method	result	size
trager	$\left(\frac{2}{5}x^2 + \frac{8}{15}x + \frac{16}{15}\right) \sqrt{-1+x}$	17
gosper	$\frac{2\sqrt{-1+x}(3x^2+4x+8)}{15}$	18
risch	$\frac{2\sqrt{-1+x}(3x^2+4x+8)}{15}$	18
derivativedivides	$\frac{4(-1+x)^{\frac{3}{2}}}{3} + \frac{2(-1+x)^{\frac{5}{2}}}{5} + 2\sqrt{-1+x}$	23
default	$\frac{4(-1+x)^{\frac{3}{2}}}{3} + \frac{2(-1+x)^{\frac{5}{2}}}{5} + 2\sqrt{-1+x}$	23
meijerg	$-\frac{\sqrt{-\text{signum}(-1+x)} \left(-\frac{16\sqrt{\pi}}{15} + \frac{\sqrt{\pi}(6x^2+8x+16)\sqrt{1-x}}{15} \right)}{\sqrt{\pi} \sqrt{\text{signum}(-1+x)}}$	48

input `int(x^2/(-1+x)^(1/2),x,method=_RETURNVERBOSE)`output `(2/5*x^2+8/15*x+16/15)*(-1+x)^(1/2)`**3.228.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.53

$$\int \frac{x^2}{\sqrt{-1+x}} dx = \frac{2}{15} (3x^2 + 4x + 8) \sqrt{x-1}$$

input `integrate(x^2/(-1+x)^(1/2),x, algorithm="fracas")`output `2/15*(3*x^2 + 4*x + 8)*sqrt(x - 1)`

3.228.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.73 (sec) , antiderivative size = 76, normalized size of antiderivative = 2.38

$$\int \frac{x^2}{\sqrt{-1+x}} dx = \begin{cases} \frac{2x^2\sqrt{x-1}}{5} + \frac{8x\sqrt{x-1}}{15} + \frac{16\sqrt{x-1}}{15} & \text{for } |x| > 1 \\ \frac{2ix^2\sqrt{1-x}}{5} + \frac{8ix\sqrt{1-x}}{15} + \frac{16i\sqrt{1-x}}{15} & \text{otherwise} \end{cases}$$

input `integrate(x**2/(-1+x)**(1/2),x)`

output `Piecewise((2*x**2*sqrt(x - 1)/5 + 8*x*sqrt(x - 1)/15 + 16*sqrt(x - 1)/15, Abs(x) > 1), (2*I*x**2*sqrt(1 - x)/5 + 8*I*x*sqrt(1 - x)/15 + 16*I*sqrt(1 - x)/15, True))`

3.228.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.69

$$\int \frac{x^2}{\sqrt{-1+x}} dx = \frac{2}{5}(x-1)^{\frac{5}{2}} + \frac{4}{3}(x-1)^{\frac{3}{2}} + 2\sqrt{x-1}$$

input `integrate(x^2/(-1+x)^(1/2),x, algorithm="maxima")`

output `2/5*(x - 1)^(5/2) + 4/3*(x - 1)^(3/2) + 2*sqrt(x - 1)`

3.228.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.69

$$\int \frac{x^2}{\sqrt{-1+x}} dx = \frac{2}{5}(x-1)^{\frac{5}{2}} + \frac{4}{3}(x-1)^{\frac{3}{2}} + 2\sqrt{x-1}$$

input `integrate(x^2/(-1+x)^(1/2),x, algorithm="giac")`

output `2/5*(x - 1)^(5/2) + 4/3*(x - 1)^(3/2) + 2*sqrt(x - 1)`

3.228.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.59

$$\int \frac{x^2}{\sqrt{-1+x}} dx = \frac{2\sqrt{x-1}(10x + 3(x-1)^2 + 5)}{15}$$

input `int(x^2/(x - 1)^(1/2),x)`

output `(2*(x - 1)^(1/2)*(10*x + 3*(x - 1)^2 + 5))/15`

3.229 $\int \frac{\sqrt{-1+x}}{1+x} dx$

3.229.1 Optimal result	1182
3.229.2 Mathematica [A] (verified)	1182
3.229.3 Rubi [A] (verified)	1183
3.229.4 Maple [A] (verified)	1184
3.229.5 Fricas [A] (verification not implemented)	1184
3.229.6 Sympy [C] (verification not implemented)	1185
3.229.7 Maxima [A] (verification not implemented)	1185
3.229.8 Giac [A] (verification not implemented)	1185
3.229.9 Mupad [B] (verification not implemented)	1186

3.229.1 Optimal result

Integrand size = 13, antiderivative size = 31

$$\int \frac{\sqrt{-1+x}}{1+x} dx = 2\sqrt{-1+x} - 2\sqrt{2} \arctan\left(\frac{\sqrt{-1+x}}{\sqrt{2}}\right)$$

output `-2*arctan(1/2*(-1+x)^(1/2)*2^(1/2))*2^(1/2)+2*(-1+x)^(1/2)`

3.229.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{-1+x}}{1+x} dx = 2\sqrt{-1+x} - 2\sqrt{2} \arctan\left(\frac{\sqrt{-1+x}}{\sqrt{2}}\right)$$

input `Integrate[Sqrt[-1 + x]/(1 + x),x]`

output `2*Sqrt[-1 + x] - 2*Sqrt[2]*ArcTan[Sqrt[-1 + x]/Sqrt[2]]`

3.229.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{x-1}}{x+1} dx \\ & \quad \downarrow \text{60} \\ & 2\sqrt{x-1} - 2 \int \frac{1}{\sqrt{x-1}(x+1)} dx \\ & \quad \downarrow \text{73} \\ & 2\sqrt{x-1} - 4 \int \frac{1}{x+1} d\sqrt{x-1} \\ & \quad \downarrow \text{216} \\ & 2\sqrt{x-1} - 2\sqrt{2} \arctan\left(\frac{\sqrt{x-1}}{\sqrt{2}}\right) \end{aligned}$$

input `Int[Sqrt[-1 + x]/(1 + x),x]`

output `2*Sqrt[-1 + x] - 2*Sqrt[2]*ArcTan[Sqrt[-1 + x]/Sqrt[2]]`

3.229.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`


```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

3.229.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$-2 \arctan\left(\frac{\sqrt{-1+x}\sqrt{2}}{2}\right) \sqrt{2} + 2\sqrt{-1+x}$	25
default	$-2 \arctan\left(\frac{\sqrt{-1+x}\sqrt{2}}{2}\right) \sqrt{2} + 2\sqrt{-1+x}$	25
risch	$-2 \arctan\left(\frac{\sqrt{-1+x}\sqrt{2}}{2}\right) \sqrt{2} + 2\sqrt{-1+x}$	25
trager	$2\sqrt{-1+x} - \text{RootOf}(_Z^2 + 2) \ln\left(-\frac{\text{RootOf}(_Z^2 + 2)x - 3\text{RootOf}(_Z^2 + 2) - 4\sqrt{-1+x}}{1+x}\right)$	49

```
input int((-1+x)^(1/2)/(1+x),x,method=_RETURNVERBOSE)
```

```
output -2*arctan(1/2*(-1+x)^(1/2)*2^(1/2))*2^(1/2)+2*(-1+x)^(1/2)
```

3.229.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.77

$$\int \frac{\sqrt{-1+x}}{1+x} dx = -2\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}\sqrt{x-1}\right) + 2\sqrt{x-1}$$

```
input integrate((-1+x)^(1/2)/(1+x),x, algorithm="fricas")
```

```
output -2*sqrt(2)*arctan(1/2*sqrt(2)*sqrt(x - 1)) + 2*sqrt(x - 1)
```

3.229.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.76 (sec) , antiderivative size = 75, normalized size of antiderivative = 2.42

$$\int \frac{\sqrt{-1+x}}{1+x} dx = \begin{cases} 2\sqrt{x-1} + 2\sqrt{2} \operatorname{asin}\left(\frac{\sqrt{2}}{\sqrt{x+1}}\right) & \text{for } |x+1| > 2 \\ 2i\sqrt{1-x} + \sqrt{2}i \log(x+1) - 2\sqrt{2}i \log\left(\sqrt{\frac{1}{2} - \frac{x}{2}} + 1\right) & \text{otherwise} \end{cases}$$

input `integrate((-1+x)**(1/2)/(1+x),x)`

output `Piecewise((2*sqrt(x - 1) + 2*sqrt(2)*asin(sqrt(2)/sqrt(x + 1)), Abs(x + 1) > 2), (2*I*sqrt(1 - x) + sqrt(2)*I*log(x + 1) - 2*sqrt(2)*I*log(sqrt(1/2 - x/2) + 1), True))`

3.229.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.77

$$\int \frac{\sqrt{-1+x}}{1+x} dx = -2\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}\sqrt{x-1}\right) + 2\sqrt{x-1}$$

input `integrate((-1+x)^(1/2)/(1+x),x, algorithm="maxima")`

output `-2*sqrt(2)*arctan(1/2*sqrt(2)*sqrt(x - 1)) + 2*sqrt(x - 1)`

3.229.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.77

$$\int \frac{\sqrt{-1+x}}{1+x} dx = -2\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}\sqrt{x-1}\right) + 2\sqrt{x-1}$$

input `integrate((-1+x)^(1/2)/(1+x),x, algorithm="giac")`

output `-2*sqrt(2)*arctan(1/2*sqrt(2)*sqrt(x - 1)) + 2*sqrt(x - 1)`

3.229.9 Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.77

$$\int \frac{\sqrt{-1+x}}{1+x} dx = 2\sqrt{x-1} - 2\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}\sqrt{x-1}}{2}\right)$$

input `int((x - 1)^(1/2)/(x + 1),x)`

output `2*(x - 1)^(1/2) - 2*2^(1/2)*atan((2^(1/2)*(x - 1)^(1/2))/2)`

3.230 $\int \frac{1}{\sqrt{1+\sqrt{x}}} dx$

3.230.1 Optimal result	1187
3.230.2 Mathematica [A] (verified)	1187
3.230.3 Rubi [A] (verified)	1188
3.230.4 Maple [A] (verified)	1189
3.230.5 Fricas [A] (verification not implemented)	1189
3.230.6 Sympy [B] (verification not implemented)	1190
3.230.7 Maxima [A] (verification not implemented)	1190
3.230.8 Giac [A] (verification not implemented)	1190
3.230.9 Mupad [B] (verification not implemented)	1191

3.230.1 Optimal result

Integrand size = 11, antiderivative size = 29

$$\int \frac{1}{\sqrt{1+\sqrt{x}}} dx = -4\sqrt{1+\sqrt{x}} + \frac{4}{3}(1+\sqrt{x})^{3/2}$$

output `4/3*(1+x^(1/2))^(3/2)-4*(1+x^(1/2))^(1/2)`

3.230.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

$$\int \frac{1}{\sqrt{1+\sqrt{x}}} dx = \frac{4}{3}(-2+\sqrt{x})\sqrt{1+\sqrt{x}}$$

input `Integrate[1/Sqrt[1 + Sqrt[x]],x]`

output `(4*(-2 + Sqrt[x])*Sqrt[1 + Sqrt[x]])/3`

3.230.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {774, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{\sqrt{x}+1}} dx \\
 & \quad \downarrow 774 \\
 & 2 \int \frac{\sqrt{x}}{\sqrt{\sqrt{x}+1}} d\sqrt{x} \\
 & \quad \downarrow 53 \\
 & 2 \int \left(\sqrt{\sqrt{x}+1} - \frac{1}{\sqrt{\sqrt{x}+1}} \right) d\sqrt{x} \\
 & \quad \downarrow 2009 \\
 & 2 \left(\frac{2}{3} (\sqrt{x}+1)^{3/2} - 2\sqrt{\sqrt{x}+1} \right)
 \end{aligned}$$

input `Int[1/Sqrt[1 + Sqrt[x]],x]`

output `2*(-2*Sqrt[1 + Sqrt[x]] + (2*(1 + Sqrt[x])^(3/2))/3)`

3.230.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 774 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[n]}, Simp[k Subst[Int[x^(k - 1)*(a + b*x^(k*n))^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, p}, x] && FractionQ[n]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.230.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.69

method	result	size
derivativedivides	$\frac{4(\sqrt{x}+1)^{\frac{3}{2}}}{3} - 4\sqrt{\sqrt{x}+1}$	20
default	$\frac{4(\sqrt{x}+1)^{\frac{3}{2}}}{3} - 4\sqrt{\sqrt{x}+1}$	20
meijerg	$\frac{8\sqrt{\pi} - \sqrt{\pi}(-4\sqrt{x}+8)\sqrt{\sqrt{x}+1}}{3\sqrt{\pi}}$	31

input `int(1/(x^(1/2)+1)^(1/2),x,method=_RETURNVERBOSE)`

output `4/3*(x^(1/2)+1)^(3/2)-4*(x^(1/2)+1)^(1/2)`

3.230.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.48

$$\int \frac{1}{\sqrt{1+\sqrt{x}}} dx = \frac{4}{3} \sqrt{\sqrt{x}+1}(\sqrt{x}-2)$$

input `integrate(1/(1+x^(1/2))^(1/2),x, algorithm="fricas")`

output `4/3*sqrt(sqrt(x) + 1)*(sqrt(x) - 2)`

3.230.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 117 vs. $2(24) = 48$.

Time = 0.50 (sec) , antiderivative size = 117, normalized size of antiderivative = 4.03

$$\int \frac{1}{\sqrt{1+\sqrt{x}}} dx = -\frac{4x^{\frac{5}{2}}\sqrt{\sqrt{x}+1}}{3x^{\frac{5}{2}}+3x^2} + \frac{8x^{\frac{5}{2}}}{3x^{\frac{5}{2}}+3x^2} + \frac{4x^3\sqrt{\sqrt{x}+1}}{3x^{\frac{5}{2}}+3x^2} - \frac{8x^2\sqrt{\sqrt{x}+1}}{3x^{\frac{5}{2}}+3x^2} + \frac{8x^2}{3x^{\frac{5}{2}}+3x^2}$$

input `integrate(1/(1+x**(1/2))**(1/2),x)`

output `-4*x**(5/2)*sqrt(sqrt(x) + 1)/(3*x**(5/2) + 3*x**2) + 8*x**(5/2)/(3*x**(5/2) + 3*x**2) + 4*x**3*sqrt(sqrt(x) + 1)/(3*x**(5/2) + 3*x**2) - 8*x**2*sqrt(sqrt(x) + 1)/(3*x**(5/2) + 3*x**2) + 8*x**2/(3*x**(5/2) + 3*x**2)`

3.230.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.66

$$\int \frac{1}{\sqrt{1+\sqrt{x}}} dx = \frac{4}{3} (\sqrt{x} + 1)^{\frac{3}{2}} - 4 \sqrt{\sqrt{x} + 1}$$

input `integrate(1/(1+x^(1/2))^(1/2),x, algorithm="maxima")`

output `4/3*(sqrt(x) + 1)^(3/2) - 4*sqrt(sqrt(x) + 1)`

3.230.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.66

$$\int \frac{1}{\sqrt{1+\sqrt{x}}} dx = \frac{4}{3} (\sqrt{x} + 1)^{\frac{3}{2}} - 4 \sqrt{\sqrt{x} + 1}$$

input `integrate(1/(1+x^(1/2))^(1/2),x, algorithm="giac")`

output `4/3*(sqrt(x) + 1)^(3/2) - 4*sqrt(sqrt(x) + 1)`

3.230.9 Mupad [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.41

$$\int \frac{1}{\sqrt{1+\sqrt{x}}} dx = x {}_2F_1\left(\frac{1}{2}, 2; 3; -\sqrt{x}\right)$$

input `int(1/(x^(1/2) + 1)^(1/2),x)`

output `x*hypergeom([1/2, 2], 3, -x^(1/2))`

3.231 $\int \frac{\sqrt{x}}{x+x^2} dx$

3.231.1 Optimal result	1192
3.231.2 Mathematica [A] (verified)	1192
3.231.3 Rubi [A] (verified)	1193
3.231.4 Maple [A] (verified)	1194
3.231.5 Fricas [A] (verification not implemented)	1194
3.231.6 Sympy [A] (verification not implemented)	1195
3.231.7 Maxima [A] (verification not implemented)	1195
3.231.8 Giac [A] (verification not implemented)	1195
3.231.9 Mupad [B] (verification not implemented)	1196

3.231.1 Optimal result

Integrand size = 13, antiderivative size = 8

$$\int \frac{\sqrt{x}}{x+x^2} dx = 2 \arctan(\sqrt{x})$$

output `2*arctan(x^(1/2))`

3.231.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{x}}{x+x^2} dx = 2 \arctan(\sqrt{x})$$

input `Integrate[Sqrt[x]/(x + x^2),x]`

output `2*ArcTan[Sqrt[x]]`

3.231.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {9, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{x}}{x^2 + x} dx \\ & \quad \downarrow \text{9} \\ & \int \frac{1}{\sqrt{x}(x+1)} dx \\ & \quad \downarrow \text{73} \\ & 2 \int \frac{1}{x+1} d\sqrt{x} \\ & \quad \downarrow \text{216} \\ & 2 \arctan(\sqrt{x}) \end{aligned}$$

input `Int[Sqrt[x]/(x + x^2),x]`

output `2*ArcTan[Sqrt[x]]`

3.231.3.1 Defintions of rubi rules used

rule 9 `Int[(u_.)*(Px_)^(p_.)*((e_.)*(x_))^(m_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Simp[1/e^(p*r) Int[u*(e*x)^(m + p*r)*ExpandToSum[Px/x^r, x]^p, x], x] /; IGtQ[r, 0]] /; FreeQ[{e, m}, x] && PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

3.231.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$2 \arctan(\sqrt{x})$	7
default	$2 \arctan(\sqrt{x})$	7
meijerg	$2 \arctan(\sqrt{x})$	7
trager	$\text{RootOf}(_Z^2 + 1) \ln\left(\frac{2\text{RootOf}(_Z^2 + 1)\sqrt{x+x-1}}{1+x}\right)$	29

input `int(x^(1/2)/(x^2+x),x,method=_RETURNVERBOSE)`

output `2*arctan(x^(1/2))`

3.231.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{x}}{x+x^2} dx = 2 \arctan(\sqrt{x})$$

input `integrate(x^(1/2)/(x^2+x),x, algorithm="fricas")`

output `2*arctan(sqrt(x))`

3.231.6 Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \frac{\sqrt{x}}{x+x^2} dx = 2 \operatorname{atan}(\sqrt{x})$$

input `integrate(x**(1/2)/(x**2+x),x)`

output `2*atan(sqrt(x))`

3.231.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{x}}{x+x^2} dx = 2 \operatorname{arctan}(\sqrt{x})$$

input `integrate(x^(1/2)/(x^2+x),x, algorithm="maxima")`

output `2*arctan(sqrt(x))`

3.231.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{x}}{x+x^2} dx = 2 \operatorname{arctan}(\sqrt{x})$$

input `integrate(x^(1/2)/(x^2+x),x, algorithm="giac")`

output `2*arctan(sqrt(x))`

3.231.9 Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{x}}{x+x^2} dx = 2 \operatorname{atan}(\sqrt{x})$$

input `int(x^(1/2)/(x + x^2),x)`

output `2*atan(x^(1/2))`

$$\mathbf{3.232} \quad \int \frac{1+\sqrt{x}}{-1+\sqrt{x}} dx$$

3.232.1 Optimal result	1197
3.232.2 Mathematica [A] (verified)	1197
3.232.3 Rubi [A] (verified)	1198
3.232.4 Maple [A] (verified)	1199
3.232.5 Fricas [A] (verification not implemented)	1200
3.232.6 Sympy [A] (verification not implemented)	1200
3.232.7 Maxima [A] (verification not implemented)	1200
3.232.8 Giac [A] (verification not implemented)	1201
3.232.9 Mupad [B] (verification not implemented)	1201

3.232.1 Optimal result

Integrand size = 17, antiderivative size = 21

$$\int \frac{1+\sqrt{x}}{-1+\sqrt{x}} dx = 4\sqrt{x} + x + 4 \log(1 - \sqrt{x})$$

output `x+4*ln(1-x^(1/2))+4*x^(1/2)`

3.232.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{1+\sqrt{x}}{-1+\sqrt{x}} dx = 4\sqrt{x} + x + 4 \log(-1 + \sqrt{x})$$

input `Integrate[(1 + Sqrt[x])/(-1 + Sqrt[x]),x]`

output `4*Sqrt[x] + x + 4*Log[-1 + Sqrt[x]]`

3.232.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.29, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {900, 25, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{x} + 1}{\sqrt{x} - 1} dx \\
 & \quad \downarrow \text{900} \\
 & 2 \int -\frac{(\sqrt{x} + 1)\sqrt{x}}{1 - \sqrt{x}} d\sqrt{x} \\
 & \quad \downarrow \text{25} \\
 & -2 \int \frac{(\sqrt{x} + 1)\sqrt{x}}{1 - \sqrt{x}} d\sqrt{x} \\
 & \quad \downarrow \text{86} \\
 & -2 \int \left(-\sqrt{x} - \frac{2}{\sqrt{x} - 1} - 2 \right) d\sqrt{x} \\
 & \quad \downarrow \text{2009} \\
 & 2 \left(\frac{x}{2} + 2\sqrt{x} + 2 \log(1 - \sqrt{x}) \right)
 \end{aligned}$$

input `Int[(1 + Sqrt[x])/(-1 + Sqrt[x]),x]`

output `2*(2*Sqrt[x] + x/2 + 2*Log[1 - Sqrt[x]])`

3.232.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 900 `Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := With[{g = Denominator[n]}, Simp[g Subst[Int[x^(g - 1)*(a + b*x^(g*n))^p*(c + d*x^(g*n))^q, x], x, x^(1/g)], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && FractionQ[n]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.232.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.76

method	result	size
derivativedivides	$x + 4\sqrt{x} + 4 \ln(\sqrt{x} - 1)$	16
default	$x + 4\sqrt{x} + 4 \ln(\sqrt{x} - 1)$	16
trager	$-2 + x + 4\sqrt{x} + 2 \ln(2\sqrt{x} - 1 - x)$	22
meijerg	$2\sqrt{x} + 4 \ln(1 - \sqrt{x}) + \frac{\sqrt{x}(3\sqrt{x}+6)}{3}$	29

input `int((x^(1/2)+1)/(x^(1/2)-1),x,method=_RETURNVERBOSE)`

output `x+4*x^(1/2)+4*ln(x^(1/2)-1)`

3.232.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \frac{1 + \sqrt{x}}{-1 + \sqrt{x}} dx = x + 4\sqrt{x} + 4 \log(\sqrt{x} - 1)$$

input `integrate((1+x^(1/2))/(-1+x^(1/2)),x, algorithm="fracas")`output `x + 4*sqrt(x) + 4*log(sqrt(x) - 1)`**3.232.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{1 + \sqrt{x}}{-1 + \sqrt{x}} dx = 4\sqrt{x} + x + 4 \log(\sqrt{x} - 1)$$

input `integrate((1+x**(1/2))/(-1+x**(1/2)),x)`output `4*sqrt(x) + x + 4*log(sqrt(x) - 1)`**3.232.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \frac{1 + \sqrt{x}}{-1 + \sqrt{x}} dx = x + 4\sqrt{x} + 4 \log(\sqrt{x} - 1)$$

input `integrate((1+x^(1/2))/(-1+x^(1/2)),x, algorithm="maxima")`output `x + 4*sqrt(x) + 4*log(sqrt(x) - 1)`

3.232.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.76

$$\int \frac{1 + \sqrt{x}}{-1 + \sqrt{x}} dx = x + 4\sqrt{x} + 4 \log(|\sqrt{x} - 1|)$$

input `integrate((1+x^(1/2))/(-1+x^(1/2)),x, algorithm="giac")`output `x + 4*sqrt(x) + 4*log(abs(sqrt(x) - 1))`**3.232.9 Mupad [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \frac{1 + \sqrt{x}}{-1 + \sqrt{x}} dx = x + 4 \ln(\sqrt{x} - 1) + 4\sqrt{x}$$

input `int((x^(1/2) + 1)/(x^(1/2) - 1),x)`output `x + 4*log(x^(1/2) - 1) + 4*x^(1/2)`

$$3.233 \quad \int \frac{1 + \frac{1}{\sqrt[3]{x}}}{-1 + \frac{1}{\sqrt[3]{x}}} dx$$

3.233.1 Optimal result	1202
3.233.2 Mathematica [A] (verified)	1202
3.233.3 Rubi [A] (verified)	1203
3.233.4 Maple [A] (verified)	1204
3.233.5 Fricas [A] (verification not implemented)	1205
3.233.6 Sympy [A] (verification not implemented)	1205
3.233.7 Maxima [A] (verification not implemented)	1205
3.233.8 Giac [A] (verification not implemented)	1206
3.233.9 Mupad [B] (verification not implemented)	1206

3.233.1 Optimal result

Integrand size = 17, antiderivative size = 30

$$\int \frac{1 + \frac{1}{\sqrt[3]{x}}}{-1 + \frac{1}{\sqrt[3]{x}}} dx = -6\sqrt[3]{x} - 3x^{2/3} - x - 6 \log(1 - \sqrt[3]{x})$$

output `-6*x^(1/3)-3*x^(2/3)-x-6*ln(1-x^(1/3))`

3.233.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{1 + \frac{1}{\sqrt[3]{x}}}{-1 + \frac{1}{\sqrt[3]{x}}} dx = -6\sqrt[3]{x} - 3x^{2/3} - x - 6 \log(-1 + \sqrt[3]{x})$$

input `Integrate[(1 + x^(-1/3))/(-1 + x^(-1/3)), x]`

output `-6*x^(1/3) - 3*x^(2/3) - x - 6*Log[-1 + x^(1/3)]`

$$3.233. \quad \int \frac{1 + \frac{1}{\sqrt[3]{x}}}{-1 + \frac{1}{\sqrt[3]{x}}} dx$$

3.233.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.13, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {898, 900, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\frac{1}{\sqrt[3]{x}} + 1}{\frac{1}{\sqrt[3]{x}} - 1} dx \\
 & \quad \downarrow \text{898} \\
 & \int \frac{\sqrt[3]{x} + 1}{1 - \sqrt[3]{x}} dx \\
 & \quad \downarrow \text{900} \\
 & 3 \int \frac{(\sqrt[3]{x} + 1) x^{2/3}}{1 - \sqrt[3]{x}} d\sqrt[3]{x} \\
 & \quad \downarrow \text{86} \\
 & 3 \int \left(-x^{2/3} - 2\sqrt[3]{x} - \frac{2}{\sqrt[3]{x} - 1} - 2 \right) d\sqrt[3]{x} \\
 & \quad \downarrow \text{2009} \\
 & 3 \left(-x^{2/3} - \frac{x}{3} - 2\sqrt[3]{x} - 2 \log(1 - \sqrt[3]{x}) \right)
 \end{aligned}$$

input `Int[(1 + x^(-1/3))/(-1 + x^(-1/3)),x]`

output `3*(-2*x^(1/3) - x^(2/3) - x/3 - 2*Log[1 - x^(1/3)])`

3.233.3.1 Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

3.233. $\int \frac{1 + \frac{1}{\sqrt[3]{x}}}{-1 + \frac{1}{\sqrt[3]{x}}} dx$

rule 898 `Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Int[x^(n*(p + q))*(b + a/x^n)^p*(d + c/x^n)^q, x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IntegersQ[p, q] && NegQ[n]`

rule 900 `Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := With[{g = Denominator[n]}, Simp[g Subst[Int[x^(g - 1)*(a + b*x^(g*n))]^p*(c + d*x^(g*n))^q, x], x, x^(1/g)], x]] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && FractionQ[n]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.233.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.77

method	result	size
derivativedivides	$-x - 3x^{\frac{2}{3}} - 6x^{\frac{1}{3}} - 6 \ln(x^{\frac{1}{3}} - 1)$	23
default	$-x - 3x^{\frac{2}{3}} - 6x^{\frac{1}{3}} - 6 \ln(x^{\frac{1}{3}} - 1)$	23
trager	$2 - x - 6x^{\frac{1}{3}} - 3x^{\frac{2}{3}} - 2 \ln(-3x^{\frac{2}{3}} + 3x^{\frac{1}{3}} + x - 1)$	32
meijerg	$-\frac{x^{\frac{1}{3}}(4x^{\frac{2}{3}} + 6x^{\frac{1}{3}} + 12)}{4} - 6 \ln(1 - x^{\frac{1}{3}}) - \frac{x^{\frac{1}{3}}(3x^{\frac{1}{3}} + 6)}{2}$	41

input `int((1+1/x^(1/3))/(-1+1/x^(1/3)),x,method=_RETURNVERBOSE)`

output `-x-3*x^(2/3)-6*x^(1/3)-6*ln(x^(1/3)-1)`

3.233.
$$\int \frac{1 + \frac{1}{\sqrt[3]{x}}}{-1 + \frac{1}{\sqrt[3]{x}}} dx$$

3.233.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \frac{1 + \frac{1}{\sqrt[3]{x}}}{-1 + \frac{1}{\sqrt[3]{x}}} dx = -x - 3x^{\frac{2}{3}} - 6x^{\frac{1}{3}} - 6 \log(x^{\frac{1}{3}} - 1)$$

input `integrate((1+1/x^(1/3))/(-1+1/x^(1/3)),x, algorithm="fricas")`output `-x - 3*x^(2/3) - 6*x^(1/3) - 6*log(x^(1/3) - 1)`**3.233.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.87

$$\int \frac{1 + \frac{1}{\sqrt[3]{x}}}{-1 + \frac{1}{\sqrt[3]{x}}} dx = -3x^{\frac{2}{3}} - 6\sqrt[3]{x} - x - 6 \log(\sqrt[3]{x} - 1)$$

input `integrate((1+1/x**(1/3))/(-1+1/x**(1/3)),x)`output `-3*x**(2/3) - 6*x**(1/3) - x - 6*log(x**(1/3) - 1)`**3.233.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \frac{1 + \frac{1}{\sqrt[3]{x}}}{-1 + \frac{1}{\sqrt[3]{x}}} dx = -x - 3x^{\frac{2}{3}} - 6x^{\frac{1}{3}} - 6 \log(x^{\frac{1}{3}} - 1)$$

input `integrate((1+1/x^(1/3))/(-1+1/x^(1/3)),x, algorithm="maxima")`output `-x - 3*x^(2/3) - 6*x^(1/3) - 6*log(x^(1/3) - 1)`

3.233. $\int \frac{1 + \frac{1}{\sqrt[3]{x}}}{-1 + \frac{1}{\sqrt[3]{x}}} dx$

3.233.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.77

$$\int \frac{1 + \frac{1}{\sqrt[3]{x}}}{-1 + \frac{1}{\sqrt[3]{x}}} dx = -x - 3x^{\frac{2}{3}} - 6x^{\frac{1}{3}} - 6 \log \left(\left| x^{\frac{1}{3}} - 1 \right| \right)$$

input `integrate((1+1/x^(1/3))/(-1+1/x^(1/3)),x, algorithm="giac")`output `-x - 3*x^(2/3) - 6*x^(1/3) - 6*log(abs(x^(1/3) - 1))`**3.233.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \frac{1 + \frac{1}{\sqrt[3]{x}}}{-1 + \frac{1}{\sqrt[3]{x}}} dx = -x - 6 \ln (x^{1/3} - 1) - 6x^{1/3} - 3x^{2/3}$$

input `int((1/x^(1/3) + 1)/(1/x^(1/3) - 1),x)`output `- x - 6*log(x^(1/3) - 1) - 6*x^(1/3) - 3*x^(2/3)`

$$3.234 \quad \int \frac{x^3}{\sqrt[3]{1+x^2}} dx$$

3.234.1 Optimal result	1207
3.234.2 Mathematica [A] (verified)	1207
3.234.3 Rubi [A] (verified)	1208
3.234.4 Maple [A] (verified)	1209
3.234.5 Fricas [A] (verification not implemented)	1209
3.234.6 Sympy [A] (verification not implemented)	1210
3.234.7 Maxima [A] (verification not implemented)	1210
3.234.8 Giac [A] (verification not implemented)	1210
3.234.9 Mupad [B] (verification not implemented)	1211

3.234.1 Optimal result

Integrand size = 13, antiderivative size = 27

$$\int \frac{x^3}{\sqrt[3]{1+x^2}} dx = -\frac{3}{4}(1+x^2)^{2/3} + \frac{3}{10}(1+x^2)^{5/3}$$

output `-3/4*(x^2+1)^(2/3)+3/10*(x^2+1)^(5/3)`

3.234.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

$$\int \frac{x^3}{\sqrt[3]{1+x^2}} dx = \frac{3}{20}(1+x^2)^{2/3}(-3+2x^2)$$

input `Integrate[x^3/(1+x^2)^(1/3),x]`

output `(3*(1+x^2)^(2/3)*(-3+2*x^2))/20`

3.234.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.15, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^3}{\sqrt[3]{x^2+1}} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int \frac{x^2}{\sqrt[3]{x^2+1}} dx^2 \\ & \quad \downarrow \text{53} \\ & \frac{1}{2} \int \left((x^2+1)^{2/3} - \frac{1}{\sqrt[3]{x^2+1}} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \left(\frac{3}{5} (x^2+1)^{5/3} - \frac{3}{2} (x^2+1)^{2/3} \right) \end{aligned}$$

input `Int[x^3/(1 + x^2)^(1/3),x]`

output `((-3*(1 + x^2)^(2/3))/2 + (3*(1 + x^2)^(5/3))/5)/2`

3.234.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.234.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.59

method	result	size
trager	$\left(\frac{3x^2}{10} - \frac{9}{20}\right) (x^2 + 1)^{\frac{2}{3}}$	16
gosper	$\frac{3(x^2+1)^{\frac{2}{3}}(2x^2-3)}{20}$	17
meijerg	$\frac{x^4 {}_2F_1\left(\frac{1}{3}, 2; 3; -x^2\right)}{4}$	17
risch	$\frac{3(x^2+1)^{\frac{2}{3}}(2x^2-3)}{20}$	17
pseudoelliptic	$\frac{3(x^2+1)^{\frac{2}{3}}(2x^2-3)}{20}$	17

input `int(x^3/(x^2+1)^(1/3),x,method=_RETURNVERBOSE)`

output `(3/10*x^2-9/20)*(x^2+1)^(2/3)`

3.234.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.59

$$\int \frac{x^3}{\sqrt[3]{1+x^2}} dx = \frac{3}{20} (2x^2 - 3)(x^2 + 1)^{\frac{2}{3}}$$

input `integrate(x^3/(x^2+1)^(1/3),x, algorithm="fracas")`

output `3/20*(2*x^2 - 3)*(x^2 + 1)^(2/3)`

3.234.6 Sympy [A] (verification not implemented)

Time = 0.54 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \frac{x^3}{\sqrt[3]{1+x^2}} dx = \frac{3x^2(x^2+1)^{\frac{2}{3}}}{10} - \frac{9(x^2+1)^{\frac{2}{3}}}{20}$$

input `integrate(x**3/(x**2+1)**(1/3),x)`output `3*x**2*(x**2 + 1)**(2/3)/10 - 9*(x**2 + 1)**(2/3)/20`**3.234.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int \frac{x^3}{\sqrt[3]{1+x^2}} dx = \frac{3}{10} (x^2+1)^{\frac{5}{3}} - \frac{3}{4} (x^2+1)^{\frac{2}{3}}$$

input `integrate(x^3/(x^2+1)^(1/3),x, algorithm="maxima")`output `3/10*(x^2 + 1)^(5/3) - 3/4*(x^2 + 1)^(2/3)`**3.234.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int \frac{x^3}{\sqrt[3]{1+x^2}} dx = \frac{3}{10} (x^2+1)^{\frac{5}{3}} - \frac{3}{4} (x^2+1)^{\frac{2}{3}}$$

input `integrate(x^3/(x^2+1)^(1/3),x, algorithm="giac")`output `3/10*(x^2 + 1)^(5/3) - 3/4*(x^2 + 1)^(2/3)`

3.234.9 Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.59

$$\int \frac{x^3}{\sqrt[3]{1+x^2}} dx = \frac{3(x^2+1)^{2/3}(2x^2-3)}{20}$$

input `int(x^3/(x^2 + 1)^(1/3),x)`output `(3*(x^2 + 1)^(2/3)*(2*x^2 - 3))/20`

3.235 $\int \frac{\sqrt{x}}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx$

3.235.1 Optimal result 1212
 3.235.2 Mathematica [C] (verified) 1213
 3.235.3 Rubi [A] (verified) 1213
 3.235.4 Maple [A] (warning: unable to verify) 1215
 3.235.5 Fricas [B] (verification not implemented) 1215
 3.235.6 Sympy [F] 1217
 3.235.7 Maxima [B] (verification not implemented) 1218
 3.235.8 Giac [A] (verification not implemented) 1219
 3.235.9 Mupad [B] (verification not implemented) 1219

3.235.1 Optimal result

Integrand size = 21, antiderivative size = 201

$$\int \frac{\sqrt{x}}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx = 6\sqrt[6]{x} + x - \frac{3}{5}\sqrt{2(5 + \sqrt{5})} \arctan\left(\frac{1 - \sqrt{5} + 4\sqrt[6]{x}}{\sqrt{2(5 + \sqrt{5})}}\right) - \frac{3}{5}\sqrt{2(5 - \sqrt{5})} \arctan\left(\frac{1}{2}\sqrt{\frac{1}{10}(5 + \sqrt{5})}(1 + \sqrt{5} + 4\sqrt[6]{x})\right) + \frac{6}{5} \log(1 - \sqrt[6]{x}) - \frac{3}{10}(1 - \sqrt{5}) \log(2 + \sqrt[6]{x} - \sqrt{5}\sqrt[6]{x} + 2\sqrt[3]{x}) - \frac{3}{10}(1 + \sqrt{5}) \log(2 + \sqrt[6]{x} + \sqrt{5}\sqrt[6]{x} + 2\sqrt[3]{x})$$

output

```
6*x^(1/6)+x+6/5*ln(1-x^(1/6))-3/10*ln(2+x^(1/6)+2*x^(1/3)-x^(1/6)*5^(1/2))
*(-5^(1/2)+1)-3/10*ln(2+x^(1/6)+2*x^(1/3)+x^(1/6)*5^(1/2))*(5^(1/2)+1)-3/5
*arctan(1/20*(1+4*x^(1/6)+5^(1/2))*(50+10*5^(1/2))^(1/2))*(10-2*5^(1/2))^(
1/2)-3/5*arctan((1+4*x^(1/6)-5^(1/2))/(10+2*5^(1/2))^(1/2))*(10+2*5^(1/2))
^(1/2)
```

3.235.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.07 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.63

$$\int \frac{\sqrt{x}}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx = 6\sqrt[6]{x} + x + \frac{6}{5} \log(-1 + \sqrt[6]{x}) - \frac{6}{5} \text{RootSum} \left[1 + \#1 + \#1^2 + \#1^3 + \#1^4 \&, \frac{4 \log(\sqrt[6]{x} - \#1) + 3 \log(\sqrt[6]{x} - \#1) \#1 + 2 \log(\sqrt[6]{x} - \#1) \#1^2 + \log(\sqrt[6]{x} - \#1) \#1^3}{1 + 2\#1 + 3\#1^2 + 4\#1^3} \& \right]$$

input `Integrate[Sqrt[x]/(-x^(-1/3) + Sqrt[x]),x]`

output `6*x^(1/6) + x + (6*Log[-1 + x^(1/6)])/5 - (6*RootSum[1 + #1 + #1^2 + #1^3 + #1^4 & , (4*Log[x^(1/6) - #1] + 3*Log[x^(1/6) - #1]*#1 + 2*Log[x^(1/6) - #1]*#1^2 + Log[x^(1/6) - #1]*#1^3)/(1 + 2*#1 + 3*#1^2 + 4*#1^3) &])/5`

3.235.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {10, 25, 864, 831, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{x}}{\sqrt{x} - \frac{1}{\sqrt[3]{x}}} dx \\ & \quad \downarrow \text{10} \\ & \int -\frac{x^{5/6}}{1 - x^{5/6}} dx \\ & \quad \downarrow \text{25} \\ & -\int \frac{x^{5/6}}{1 - x^{5/6}} dx \\ & \quad \downarrow \text{864} \\ & -6 \int \frac{x^{5/3}}{1 - x^{5/6}} d\sqrt[6]{x} \end{aligned}$$

3.235. $\int \frac{\sqrt{x}}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx$

$$\begin{array}{c} \downarrow 831 \\ -6 \int \left(-x^{5/6} + \frac{1}{1-x^{5/6}} - 1 \right) d\sqrt[6]{x} \\ \downarrow 2009 \end{array}$$

$$-6 \left(\frac{1}{5} \sqrt{\frac{1}{2} (5 + \sqrt{5})} \arctan \left(\frac{4\sqrt[6]{x} - \sqrt{5} + 1}{\sqrt{2} (5 + \sqrt{5})} \right) + \frac{1}{5} \sqrt{\frac{1}{2} (5 - \sqrt{5})} \arctan \left(\frac{1}{2} \sqrt{\frac{1}{10} (5 + \sqrt{5})} (4\sqrt[6]{x} + \sqrt{5} + 1) \right) \right)$$

input `Int[Sqrt[x]/(-x^(-1/3) + Sqrt[x]),x]`

output `-6*(-x^(1/6) - x/6 + (Sqrt[(5 + Sqrt[5])/2]*ArcTan[(1 - Sqrt[5] + 4*x^(1/6))/Sqrt[2*(5 + Sqrt[5])]])/5 + (Sqrt[(5 - Sqrt[5])/2]*ArcTan[(Sqrt[(5 + Sqrt[5])/10]*(1 + Sqrt[5] + 4*x^(1/6)))/2])/5 - Log[1 - x^(1/6)]/5 + ((1 - Sqrt[5])*Log[1 + ((1 - Sqrt[5])*x^(1/6))/2 + x^(1/3)])/20 + ((1 + Sqrt[5])*Log[1 + ((1 + Sqrt[5])*x^(1/6))/2 + x^(1/3)])/20`

3.235.3.1 Defintions of rubi rules used

rule 10 `Int[(u_)*((e_)*(x_)^(m_))*((a_)*(x_)^(r_) + (b_)*(x_)^(s_))^(p_), x_Symbol] := Simp[1/e^(p*r) Int[u*(e*x)^(m + p*r)*(a + b*x^(s - r))^p, x], x] /; FreeQ[{a, b, e, m, r, s}, x] && IntegerQ[p] && (IntegerQ[p*r] || GtQ[e, 0]) && PosQ[s - r]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 831 `Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^n, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 0] && IGtQ[n, 0] && GtQ[m, 2*n - 1]`

rule 864 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a + b*x^(k*n))^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, m, p}, x] && FractionQ[n]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.235. $\int \frac{\sqrt{x}}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx$

3.235.4 Maple [A] (warning: unable to verify)

Time = 0.15 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.66

method	result
meijerg	$\frac{6(-1)^{\frac{4}{5}} \left(-\frac{5x^{\frac{1}{6}}(-1)^{\frac{1}{5}}(11x^{\frac{5}{6}}+66)}{66} - (-1)^{\frac{1}{5}} \left(\ln(1-x^{\frac{1}{6}}) + \cos\left(\frac{2\pi}{5}\right) \ln(1-2\cos\left(\frac{2\pi}{5}\right)x^{\frac{1}{6}}+x^{\frac{1}{3}}) - 2\sin\left(\frac{2\pi}{5}\right) \arctan\left(\frac{\sin\left(\frac{2\pi}{5}\right)x^{\frac{1}{6}}}{1-\cos\left(\frac{2\pi}{5}\right)x^{\frac{1}{6}}}\right) \right)}{5} \right)}{x + 6x^{\frac{1}{6}} - \frac{3\ln(2+x^{\frac{1}{6}}+2x^{\frac{1}{3}}-x^{\frac{1}{6}}\sqrt{5})(-\sqrt{5}+1)}{10} - \frac{12\left(4-\frac{(-\sqrt{5}+1)^2}{4}\right)\arctan\left(\frac{1+4x^{\frac{1}{6}}-\sqrt{5}}{\sqrt{10+2\sqrt{5}}}\right)}{5\sqrt{10+2\sqrt{5}}} + \frac{3(-\sqrt{5}-1)\ln(1-x^{\frac{1}{6}})}{5}}$
derivativedivides	$x + 6x^{\frac{1}{6}} - \frac{3\ln(2+x^{\frac{1}{6}}+2x^{\frac{1}{3}}-x^{\frac{1}{6}}\sqrt{5})(-\sqrt{5}+1)}{10} - \frac{12\left(4-\frac{(-\sqrt{5}+1)^2}{4}\right)\arctan\left(\frac{1+4x^{\frac{1}{6}}-\sqrt{5}}{\sqrt{10+2\sqrt{5}}}\right)}{5\sqrt{10+2\sqrt{5}}} + \frac{3(-\sqrt{5}-1)\ln(1-x^{\frac{1}{6}})}{5}$
default	$x + 6x^{\frac{1}{6}} - \frac{3\ln(2+x^{\frac{1}{6}}+2x^{\frac{1}{3}}-x^{\frac{1}{6}}\sqrt{5})(-\sqrt{5}+1)}{10} - \frac{12\left(4-\frac{(-\sqrt{5}+1)^2}{4}\right)\arctan\left(\frac{1+4x^{\frac{1}{6}}-\sqrt{5}}{\sqrt{10+2\sqrt{5}}}\right)}{5\sqrt{10+2\sqrt{5}}} + \frac{3(-\sqrt{5}-1)\ln(1-x^{\frac{1}{6}})}{5}$

input `int(x^(1/2)/(-1/x^(1/3)+x^(1/2)),x,method=_RETURNVERBOSE)`output `6/5*(-1)^(4/5)*(-5/66*x^(1/6)*(-1)^(1/5)*(11*x^(5/6)+66)-(-1)^(1/5)*(ln(1-x^(1/6))+cos(2/5*Pi)*ln(1-2*cos(2/5*Pi)*x^(1/6)+x^(1/3))-2*sin(2/5*Pi)*arctan(sin(2/5*Pi)*x^(1/6)/(1-cos(2/5*Pi)*x^(1/6)))-cos(1/5*Pi)*ln(1+2*cos(1/5*Pi)*x^(1/6)+x^(1/3))-2*sin(1/5*Pi)*arctan(sin(1/5*Pi)*x^(1/6)/(1+cos(1/5*Pi)*x^(1/6))))`**3.235.5 Fracas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 547 vs. $2(134) = 268$.

3.235. $\int \frac{\sqrt{x}}{-\frac{1}{\sqrt[3]{x}}+\sqrt{x}} dx$

Time = 0.91 (sec) , antiderivative size = 547, normalized size of antiderivative = 2.72

$$\begin{aligned}
 \int \frac{\sqrt{x}}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx = & -\frac{3}{10} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} + 1 \right) \log \left(\frac{3}{2} \sqrt{2}\sqrt{\sqrt{5}-5} + \frac{3}{2} \sqrt{5} + 6x^{\frac{1}{6}} + \frac{3}{2} \right) \\
 & + \frac{3}{10} \left(\sqrt{2}\sqrt{\sqrt{5}-5} - \sqrt{5} - 1 \right) \log \left(-\frac{3}{2} \sqrt{2}\sqrt{\sqrt{5}-5} + \frac{3}{2} \sqrt{5} + 6x^{\frac{1}{6}} + \frac{3}{2} \right) \\
 & + \frac{1}{10} \left(3\sqrt{5} - \sqrt{-\frac{27}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} + 1 \right)^2} + \frac{9}{2} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} - 3 \right) \left(\sqrt{2}\sqrt{\sqrt{5}-5} - \sqrt{5} - \right. \right. \\
 & \left. \left. + \sqrt{-\frac{27}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} + 1 \right)^2} + \frac{9}{2} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} - 3 \right) \left(\sqrt{2}\sqrt{\sqrt{5}-5} - \sqrt{5} - 1 \right) - \frac{27}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} + 1 \right) \right. \right. \\
 & \left. \left. + 12x^{\frac{1}{6}} + 3 \right) \right) \\
 & + \frac{1}{10} \left(3\sqrt{5} + \sqrt{-\frac{27}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} + 1 \right)^2} + \frac{9}{2} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} - 3 \right) \left(\sqrt{2}\sqrt{\sqrt{5}-5} - \sqrt{5} - \right. \right. \\
 & \left. \left. - \sqrt{-\frac{27}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} + 1 \right)^2} + \frac{9}{2} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} - 3 \right) \left(\sqrt{2}\sqrt{\sqrt{5}-5} - \sqrt{5} - 1 \right) - \frac{27}{4} \left(\sqrt{2}\sqrt{\sqrt{5}-5} + \sqrt{5} + 1 \right) \right. \right. \\
 & \left. \left. + 12x^{\frac{1}{6}} + 3 \right) + x + 6x^{\frac{1}{6}} + \frac{6}{5} \log \left(x^{\frac{1}{6}} - 1 \right)
 \end{aligned}$$

input `integrate(x^(1/2)/(-1/x^(1/3)+x^(1/2)),x, algorithm="fricas")`

output

```
-3/10*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) + 1)*log(3/2*sqrt(2)*sqrt(sqrt(5) - 5) + 3/2*sqrt(5) + 6*x^(1/6) + 3/2) + 3/10*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1)*log(-3/2*sqrt(2)*sqrt(sqrt(5) - 5) + 3/2*sqrt(5) + 6*x^(1/6) + 3/2) + 1/10*(3*sqrt(5) - sqrt(-27/4*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) + 1)^2 + 9/2*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) - 3)*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1) - 27/4*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1)^2 + 18*sqrt(2)*sqrt(sqrt(5) - 5) + 18*sqrt(5) - 90) - 3)*log(-3*sqrt(5) + sqrt(-27/4*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) + 1)^2 + 9/2*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) - 3)*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1) - 27/4*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1)^2 + 18*sqrt(2)*sqrt(sqrt(5) - 5) + 18*sqrt(5) - 90) + 12*x^(1/6) + 3) + 1/10*(3*sqrt(5) + sqrt(-27/4*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) + 1)^2 + 9/2*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) - 3)*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1) - 27/4*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1)^2 + 18*sqrt(2)*sqrt(sqrt(5) - 5) + 18*sqrt(5) - 90) - 3)*log(-3*sqrt(5) - sqrt(-27/4*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) + 1)^2 + 9/2*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) - 3)*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1) - 27/4*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1)^2 + 18*sqrt(2)*sqrt(sqrt(5) - 5) + 18*sqrt(5) - 90) - 3)*log(-3*sqrt(5) - sqrt(-27/4*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) + 1)^2 + 9/2*(sqrt(2)*sqrt(sqrt(5) - 5) + sqrt(5) - 3)*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1) - 27/4*(sqrt(2)*sqrt(sqrt(5) - 5) - sqrt(5) - 1)^2 + 18*sqrt(2)*sqrt(sqrt(5) - 5) + 18*sqrt(5) - 90) + 12*x^(1/6) + 3) + x + 6*x^(1/6) + 6/5*log(x^(1/6) - 1)
```

3.235.6 Sympy [F]

$$\int \frac{\sqrt{x}}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx = \int \frac{x^{\frac{5}{6}}}{(\sqrt[6]{x} - 1) (\sqrt[6]{x} + x^{\frac{2}{3}} + \sqrt[3]{x} + \sqrt{x} + 1)} dx$$

input `integrate(x**(1/2)/(-1/x**(1/3)+x**(1/2)),x)`

output `Integral(x**(5/6)/((x**(1/6) - 1)*(x**(1/6) + x**(2/3) + x**(1/3) + sqrt(x) + 1)), x)`

3.235.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 293 vs. $2(134) = 268$.

Time = 0.32 (sec) , antiderivative size = 293, normalized size of antiderivative = 1.46

$$\int \frac{\sqrt{x}}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx = -\frac{3\sqrt{5}(-1)^{\frac{1}{5}}(\sqrt{5}-1)\log\left(\frac{\sqrt{5}(-1)^{\frac{1}{5}}+(-1)^{\frac{1}{5}}\sqrt{2\sqrt{5}-10}+(-1)^{\frac{1}{5}}-4x^{\frac{1}{6}}}{\sqrt{5}(-1)^{\frac{1}{5}}-(-1)^{\frac{1}{5}}\sqrt{2\sqrt{5}-10}+(-1)^{\frac{1}{5}}-4x^{\frac{1}{6}}}\right)}{5\sqrt{2\sqrt{5}-10}}$$

$$-\frac{3\sqrt{5}(-1)^{\frac{1}{5}}(\sqrt{5}+1)\log\left(\frac{\sqrt{5}(-1)^{\frac{1}{5}}-(-1)^{\frac{1}{5}}\sqrt{-2\sqrt{5}-10}-(-1)^{\frac{1}{5}}+4x^{\frac{1}{6}}}{\sqrt{5}(-1)^{\frac{1}{5}}+(-1)^{\frac{1}{5}}\sqrt{-2\sqrt{5}-10}-(-1)^{\frac{1}{5}}+4x^{\frac{1}{6}}}\right)}{5\sqrt{-2\sqrt{5}-10}}$$

$$-\frac{6}{5}(-1)^{\frac{1}{5}}\log\left((-1)^{\frac{1}{5}}+x^{\frac{1}{6}}\right)+x$$

$$-\frac{3(\sqrt{5}+3)\log\left(-x^{\frac{1}{6}}\left(\sqrt{5}(-1)^{\frac{1}{5}}+(-1)^{\frac{1}{5}}\right)+2(-1)^{\frac{2}{5}}+2x^{\frac{1}{3}}\right)}{5\left(\sqrt{5}(-1)^{\frac{4}{5}}+(-1)^{\frac{4}{5}}\right)}$$

$$-\frac{3(\sqrt{5}-3)\log\left(x^{\frac{1}{6}}\left(\sqrt{5}(-1)^{\frac{1}{5}}-(-1)^{\frac{1}{5}}\right)+2(-1)^{\frac{2}{5}}+2x^{\frac{1}{3}}\right)}{5\left(\sqrt{5}(-1)^{\frac{4}{5}}-(-1)^{\frac{4}{5}}\right)}+6x^{\frac{1}{6}}$$

input `integrate(x^(1/2)/(-1/x^(1/3)+x^(1/2)),x, algorithm="maxima")`

output `-3/5*sqrt(5)*(-1)^(1/5)*(sqrt(5) - 1)*log((sqrt(5)*(-1)^(1/5) + (-1)^(1/5) *sqrt(2*sqrt(5) - 10) + (-1)^(1/5) - 4*x^(1/6))/(sqrt(5)*(-1)^(1/5) - (-1)^(1/5)*sqrt(2*sqrt(5) - 10) + (-1)^(1/5) - 4*x^(1/6)))/sqrt(2*sqrt(5) - 10) - 3/5*sqrt(5)*(-1)^(1/5)*(sqrt(5) + 1)*log((sqrt(5)*(-1)^(1/5) - (-1)^(1/5)*sqrt(-2*sqrt(5) - 10) - (-1)^(1/5) + 4*x^(1/6))/(sqrt(5)*(-1)^(1/5) + (-1)^(1/5)*sqrt(-2*sqrt(5) - 10) - (-1)^(1/5) + 4*x^(1/6)))/sqrt(-2*sqrt(5) - 10) - 6/5*(-1)^(1/5)*log((-1)^(1/5) + x^(1/6)) + x - 3/5*(sqrt(5) + 3) *log(-x^(1/6)*(sqrt(5)*(-1)^(1/5) + (-1)^(1/5)) + 2*(-1)^(2/5) + 2*x^(1/3))/(sqrt(5)*(-1)^(4/5) + (-1)^(4/5)) - 3/5*(sqrt(5) - 3)*log(x^(1/6)*(sqrt(5)*(-1)^(1/5) - (-1)^(1/5)) + 2*(-1)^(2/5) + 2*x^(1/3))/(sqrt(5)*(-1)^(4/5) - (-1)^(4/5)) + 6*x^(1/6)`

3.235.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.70

$$\int \frac{\sqrt{x}}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx = -\frac{3}{5} \sqrt{2\sqrt{5} + 10} \arctan\left(-\frac{\sqrt{5} - 4x^{\frac{1}{6}} - 1}{\sqrt{2\sqrt{5} + 10}}\right) - \frac{3}{5} \sqrt{-2\sqrt{5} + 10} \arctan\left(\frac{\sqrt{5} + 4x^{\frac{1}{6}} + 1}{\sqrt{-2\sqrt{5} + 10}}\right) - \frac{3}{10} \sqrt{5} \log\left(\frac{1}{2} x^{\frac{1}{6}} (\sqrt{5} + 1) + x^{\frac{1}{3}} + 1\right) + \frac{3}{10} \sqrt{5} \log\left(-\frac{1}{2} x^{\frac{1}{6}} (\sqrt{5} - 1) + x^{\frac{1}{3}} + 1\right) + x + 6x^{\frac{1}{6}} - \frac{3}{10} \log\left(x^{\frac{2}{3}} + \sqrt{x} + x^{\frac{1}{3}} + x^{\frac{1}{6}} + 1\right) + \frac{6}{5} \log\left(|x^{\frac{1}{6}} - 1|\right)$$

input `integrate(x^(1/2)/(-1/x^(1/3)+x^(1/2)),x, algorithm="giac")`output `-3/5*sqrt(2*sqrt(5) + 10)*arctan(-(sqrt(5) - 4*x^(1/6) - 1)/sqrt(2*sqrt(5) + 10)) - 3/5*sqrt(-2*sqrt(5) + 10)*arctan((sqrt(5) + 4*x^(1/6) + 1)/sqrt(-2*sqrt(5) + 10)) - 3/10*sqrt(5)*log(1/2*x^(1/6)*(sqrt(5) + 1) + x^(1/3) + 1) + 3/10*sqrt(5)*log(-1/2*x^(1/6)*(sqrt(5) - 1) + x^(1/3) + 1) + x + 6*x^(1/6) - 3/10*log(x^(2/3) + sqrt(x) + x^(1/3) + x^(1/6) + 1) + 6/5*log(abs(x^(1/6) - 1))`**3.235.9 Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.03

$$\int \frac{\sqrt{x}}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx = x + \frac{6 \ln(1296 x^{1/6} - 1296)}{5} - \ln\left(270 \sqrt{2} \sqrt{-\sqrt{5} - 5} - 270 \sqrt{5} + 1080 x^{1/6} + 270\right) \left(\frac{3 \sqrt{2} \sqrt{-\sqrt{5} - 5}}{10} - \frac{3 \sqrt{5}}{10} + \frac{3}{10}\right) + \ln\left(270 \sqrt{2} \sqrt{-\sqrt{5} - 5} + 270 \sqrt{5} - 1080 x^{1/6} - 270\right) \left(\frac{3 \sqrt{2} \sqrt{-\sqrt{5} - 5}}{10} + \frac{3 \sqrt{5}}{10} - \frac{3}{10}\right) + 6x^{1/6} - \ln\left(270\right)$$

input `int(x^(1/2)/(x^(1/2) - 1/x^(1/3)),x)`

3.235. $\int \frac{\sqrt{x}}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx$

output $x + (6*\log(1296*x^{(1/6)} - 1296))/5 - \log(270*2^{(1/2)}*(-5^{(1/2)} - 5)^{(1/2)} - 270*5^{(1/2)} + 1080*x^{(1/6)} + 270)*((3*2^{(1/2)}*(-5^{(1/2)} - 5)^{(1/2)})/10 - (3*5^{(1/2)})/10 + 3/10) + \log(270*2^{(1/2)}*(-5^{(1/2)} - 5)^{(1/2)} + 270*5^{(1/2)} - 1080*x^{(1/6)} - 270)*((3*2^{(1/2)}*(-5^{(1/2)} - 5)^{(1/2)})/10 + (3*5^{(1/2)})/10 - 3/10) + 6*x^{(1/6)} - \log(270*5^{(1/2)} + 1080*x^{(1/6)} - 270*2^{(1/2)}*(5^{(1/2)} - 5)^{(1/2)} + 270)*((3*5^{(1/2)})/10 - (3*2^{(1/2)}*(5^{(1/2)} - 5)^{(1/2)})/10 + 3/10) - \log(270*5^{(1/2)} + 1080*x^{(1/6)} + 270*2^{(1/2)}*(5^{(1/2)} - 5)^{(1/2)} + 270)*((3*5^{(1/2)})/10 + (3*2^{(1/2)}*(5^{(1/2)} - 5)^{(1/2)})/10 + 3/10)$

3.235. $\int \frac{\sqrt{x}}{-\frac{1}{\sqrt[3]{x}} + \sqrt{x}} dx$

$$3.236 \quad \int \frac{1}{\frac{1}{\sqrt[4]{x}} + \sqrt{x}} dx$$

3.236.1 Optimal result	1221
3.236.2 Mathematica [A] (verified)	1221
3.236.3 Rubi [A] (verified)	1222
3.236.4 Maple [A] (verified)	1224
3.236.5 Fricas [A] (verification not implemented)	1225
3.236.6 Sympy [A] (verification not implemented)	1225
3.236.7 Maxima [A] (verification not implemented)	1226
3.236.8 Giac [A] (verification not implemented)	1226
3.236.9 Mupad [B] (verification not implemented)	1226

3.236.1 Optimal result

Integrand size = 13, antiderivative size = 62

$$\int \frac{1}{\frac{1}{\sqrt[4]{x}} + \sqrt{x}} dx = 2\sqrt{x} + \frac{4 \arctan\left(\frac{1-2\sqrt[4]{x}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{4}{3} \log(1 + \sqrt[4]{x}) - \frac{2}{3} \log(1 - \sqrt[4]{x} + \sqrt{x})$$

output `4/3*ln(1+x^(1/4))-2/3*ln(1-x^(1/4)+x^(1/2))+4/3*arctan(1/3*(1-2*x^(1/4))*3^(1/2))*3^(1/2)+2*x^(1/2)`

3.236.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00

$$\int \frac{1}{\frac{1}{\sqrt[4]{x}} + \sqrt{x}} dx = \frac{2}{3} \left(3\sqrt{x} + 2\sqrt{3} \arctan\left(\frac{1-2\sqrt[4]{x}}{\sqrt{3}}\right) + 2 \log(1 + \sqrt[4]{x}) - \log(1 - \sqrt[4]{x} + \sqrt{x}) \right)$$

input `Integrate[(x^(-1/4) + Sqrt[x])^(-1), x]`

output `(2*(3*Sqrt[x] + 2*Sqrt[3]*ArcTan[(1 - 2*x^(1/4))/Sqrt[3]] + 2*Log[1 + x^(1/4)] - Log[1 - x^(1/4) + Sqrt[x]]))/3`

3.236. $\int \frac{1}{\frac{1}{\sqrt[4]{x}} + \sqrt{x}} dx$

3.236.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.15, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.769$, Rules used = {2027, 864, 843, 821, 16, 1142, 25, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{x} + \frac{1}{\sqrt[4]{x}}} dx \\
 & \quad \downarrow \text{2027} \\
 & \int \frac{\sqrt[4]{x}}{x^{3/4} + 1} dx \\
 & \quad \downarrow \text{864} \\
 & 4 \int \frac{x}{x^{3/4} + 1} d\sqrt[4]{x} \\
 & \quad \downarrow \text{843} \\
 & 4 \left(\frac{\sqrt{x}}{2} - \int \frac{\sqrt[4]{x}}{x^{3/4} + 1} d\sqrt[4]{x} \right) \\
 & \quad \downarrow \text{821} \\
 & 4 \left(\frac{1}{3} \int \frac{1}{\sqrt[4]{x} + 1} d\sqrt[4]{x} - \frac{1}{3} \int \frac{\sqrt[4]{x} + 1}{\sqrt{x} - \sqrt[4]{x} + 1} d\sqrt[4]{x} + \frac{\sqrt{x}}{2} \right) \\
 & \quad \downarrow \text{16} \\
 & 4 \left(-\frac{1}{3} \int \frac{\sqrt[4]{x} + 1}{\sqrt{x} - \sqrt[4]{x} + 1} d\sqrt[4]{x} + \frac{\sqrt{x}}{2} + \frac{1}{3} \log(\sqrt[4]{x} + 1) \right) \\
 & \quad \downarrow \text{1142} \\
 & 4 \left(\frac{1}{3} \left(-\frac{3}{2} \int \frac{1}{\sqrt{x} - \sqrt[4]{x} + 1} d\sqrt[4]{x} - \frac{1}{2} \int -\frac{1 - 2\sqrt[4]{x}}{\sqrt{x} - \sqrt[4]{x} + 1} d\sqrt[4]{x} \right) + \frac{\sqrt{x}}{2} + \frac{1}{3} \log(\sqrt[4]{x} + 1) \right) \\
 & \quad \downarrow \text{25} \\
 & 4 \left(\frac{1}{3} \left(\frac{1}{2} \int \frac{1 - 2\sqrt[4]{x}}{\sqrt{x} - \sqrt[4]{x} + 1} d\sqrt[4]{x} - \frac{3}{2} \int \frac{1}{\sqrt{x} - \sqrt[4]{x} + 1} d\sqrt[4]{x} \right) + \frac{\sqrt{x}}{2} + \frac{1}{3} \log(\sqrt[4]{x} + 1) \right) \\
 & \quad \downarrow \text{1083} \\
 & 4 \left(\frac{1}{3} \left(3 \int \frac{1}{-\sqrt{x} - 3} d(2\sqrt[4]{x} - 1) + \frac{1}{2} \int \frac{1 - 2\sqrt[4]{x}}{\sqrt{x} - \sqrt[4]{x} + 1} d\sqrt[4]{x} \right) + \frac{\sqrt{x}}{2} + \frac{1}{3} \log(\sqrt[4]{x} + 1) \right)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 217 \\
 & 4 \left(\frac{1}{3} \left(\frac{1}{2} \int \frac{1 - 2\sqrt[4]{x}}{\sqrt{x} - \sqrt[4]{x} + 1} d\sqrt[4]{x} - \sqrt{3} \arctan \left(\frac{2\sqrt[4]{x} - 1}{\sqrt{3}} \right) \right) + \frac{\sqrt{x}}{2} + \frac{1}{3} \log(\sqrt[4]{x} + 1) \right) \\
 & \downarrow 1103 \\
 & 4 \left(\frac{1}{3} \left(-\sqrt{3} \arctan \left(\frac{2\sqrt[4]{x} - 1}{\sqrt{3}} \right) - \frac{1}{2} \log(\sqrt{x} - \sqrt[4]{x} + 1) \right) + \frac{\sqrt{x}}{2} + \frac{1}{3} \log(\sqrt[4]{x} + 1) \right)
 \end{aligned}$$

input `Int[(x^(-1/4) + Sqrt[x])^(-1), x]`

output `4*(Sqrt[x]/2 + Log[1 + x^(1/4)]/3 + (-Sqrt[3]*ArcTan[(-1 + 2*x^(1/4))/Sqrt[3]]) - Log[1 - x^(1/4) + Sqrt[x]]/2)/3`

3.236.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 821 `Int[(x_)/((a_) + (b_.)*(x_)^3), x_Symbol] := Simp[-(3*Rt[a, 3]*Rt[b, 3])^(-1) Int[1/(Rt[a, 3] + Rt[b, 3]*x), x, x] + Simp[1/(3*Rt[a, 3]*Rt[b, 3]) Int[(Rt[a, 3] + Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]`

rule 843 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Simp[a*c^n*((m - n + 1)/(b*(m + n*p + 1))) Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 864 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a + b*x^(k*n))^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, m, p}, x] && FractionQ[n]`

rule 1083 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 2027 `Int[(F*x_)*((a_)*(x_)^(r_) + (b_)*(x_)^(s_))^(p_), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r))^p*F*x, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] && PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])`

3.236.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.74

method	result	size
derivativedivides	$2\sqrt{x} - \frac{2\ln(1-x^{\frac{1}{4}}+\sqrt{x})}{3} - \frac{4\sqrt{3} \arctan\left(\frac{(2x^{\frac{1}{4}}-1)\sqrt{3}}{3}\right)}{3} + \frac{4\ln(1+x^{\frac{1}{4}})}{3}$	46
default	$2\sqrt{x} - \frac{2\ln(1-x^{\frac{1}{4}}+\sqrt{x})}{3} - \frac{4\sqrt{3} \arctan\left(\frac{(2x^{\frac{1}{4}}-1)\sqrt{3}}{3}\right)}{3} + \frac{4\ln(1+x^{\frac{1}{4}})}{3}$	46
meijerg	$2\sqrt{x} - \frac{4\sqrt{x} \left(-\frac{\ln(1+x^{\frac{1}{4}})}{\sqrt{x}} + \frac{\ln(1-x^{\frac{1}{4}}+\sqrt{x})}{2\sqrt{x}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}x^{\frac{1}{4}}}{2-x^{\frac{1}{4}}}\right)}{\sqrt{x}} \right)}{3}$	65

input `int(1/(1/x^(1/4)+x^(1/2)),x,method=_RETURNVERBOSE)`

output `2*x^(1/2)-2/3*ln(1-x^(1/4)+x^(1/2))-4/3*3^(1/2)*arctan(1/3*(2*x^(1/4)-1)*3^(1/2))+4/3*ln(1+x^(1/4))`

3.236.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.76

$$\int \frac{1}{\frac{1}{\sqrt[4]{x}} + \sqrt{x}} dx = -\frac{4}{3} \sqrt{3} \arctan\left(\frac{2}{3} \sqrt{3} x^{\frac{1}{4}} - \frac{1}{3} \sqrt{3}\right) + 2\sqrt{x} - \frac{2}{3} \log\left(\sqrt{x} - x^{\frac{1}{4}} + 1\right) + \frac{4}{3} \log\left(x^{\frac{1}{4}} + 1\right)$$

input `integrate(1/(1/x^(1/4)+x^(1/2)),x, algorithm="fricas")`

output `-4/3*sqrt(3)*arctan(2/3*sqrt(3)*x^(1/4) - 1/3*sqrt(3)) + 2*sqrt(x) - 2/3*log(sqrt(x) - x^(1/4) + 1) + 4/3*log(x^(1/4) + 1)`

3.236.6 Sympy [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.10

$$\int \frac{1}{\frac{1}{\sqrt[4]{x}} + \sqrt{x}} dx = 2\sqrt{x} + \frac{4 \log(\sqrt[4]{x} + 1)}{3} - \frac{2 \log(-4\sqrt[4]{x} + 4\sqrt{x} + 4)}{3} - \frac{4\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}\sqrt[4]{x}}{3} - \frac{\sqrt{3}}{3}\right)}{3}$$

input `integrate(1/(1/x**(1/4)+x**(1/2)),x)`

output `2*sqrt(x) + 4*log(x**(1/4) + 1)/3 - 2*log(-4*x**(1/4) + 4*sqrt(x) + 4)/3 - 4*sqrt(3)*atan(2*sqrt(3)*x**(1/4)/3 - sqrt(3)/3)/3`

3.236.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.73

$$\int \frac{1}{\frac{1}{\sqrt[4]{x}} + \sqrt{x}} dx = -\frac{4}{3} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} (2x^{1/4} - 1) \right) + 2\sqrt{x} \\ - \frac{2}{3} \log \left(\sqrt{x} - x^{1/4} + 1 \right) + \frac{4}{3} \log \left(x^{1/4} + 1 \right)$$

input `integrate(1/(1/x^(1/4)+x^(1/2)),x, algorithm="maxima")`output `-4/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x^(1/4) - 1)) + 2*sqrt(x) - 2/3*log(sqrt(x) - x^(1/4) + 1) + 4/3*log(x^(1/4) + 1)`**3.236.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.73

$$\int \frac{1}{\frac{1}{\sqrt[4]{x}} + \sqrt{x}} dx = -\frac{4}{3} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} (2x^{1/4} - 1) \right) + 2\sqrt{x} \\ - \frac{2}{3} \log \left(\sqrt{x} - x^{1/4} + 1 \right) + \frac{4}{3} \log \left(x^{1/4} + 1 \right)$$

input `integrate(1/(1/x^(1/4)+x^(1/2)),x, algorithm="giac")`output `-4/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x^(1/4) - 1)) + 2*sqrt(x) - 2/3*log(sqrt(x) - x^(1/4) + 1) + 4/3*log(x^(1/4) + 1)`**3.236.9 Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.18

$$\int \frac{1}{\frac{1}{\sqrt[4]{x}} + \sqrt{x}} dx = \frac{4 \ln(16x^{1/4} + 16)}{3} + \ln \left(9 \left(-\frac{2}{3} + \frac{\sqrt{3}2i}{3} \right)^2 + 16x^{1/4} \right) \left(-\frac{2}{3} + \frac{\sqrt{3}2i}{3} \right) \\ - \ln \left(9 \left(\frac{2}{3} + \frac{\sqrt{3}2i}{3} \right)^2 + 16x^{1/4} \right) \left(\frac{2}{3} + \frac{\sqrt{3}2i}{3} \right) + 2\sqrt{x}$$

input `int(1/(x^(1/2) + 1/x^(1/4)),x)`

output `(4*log(16*x^(1/4) + 16))/3 + log(9*((3^(1/2)*2i)/3 - 2/3)^2 + 16*x^(1/4))*
((3^(1/2)*2i)/3 - 2/3) - log(9*((3^(1/2)*2i)/3 + 2/3)^2 + 16*x^(1/4))*((3^(1/2)*2i)/3 + 2/3) + 2*x^(1/2)`

3.237 $\int \frac{1}{\frac{1}{\sqrt[3]{x}} + \frac{1}{\sqrt[4]{x}}} dx$

3.237.1 Optimal result 1228
 3.237.2 Mathematica [A] (verified) 1228
 3.237.3 Rubi [A] (verified) 1229
 3.237.4 Maple [A] (verified) 1230
 3.237.5 Fricas [A] (verification not implemented) 1231
 3.237.6 Sympy [A] (verification not implemented) 1231
 3.237.7 Maxima [A] (verification not implemented) 1232
 3.237.8 Giac [A] (verification not implemented) 1232
 3.237.9 Mupad [B] (verification not implemented) 1233

3.237.1 Optimal result

Integrand size = 13, antiderivative size = 130

$$\int \frac{1}{\frac{1}{\sqrt[3]{x}} + \frac{1}{\sqrt[4]{x}}} dx = 12 \sqrt[12]{x} - 6\sqrt[6]{x} + 4\sqrt[4]{x} - 3\sqrt[3]{x} + \frac{12x^{5/12}}{5} - 2\sqrt{x} + \frac{12x^{7/12}}{7} - \frac{3x^{2/3}}{2} + \frac{4x^{3/4}}{3} - \frac{6x^{5/6}}{5} + \frac{12x^{11/12}}{11} - x + \frac{12x^{13/12}}{13} - \frac{6x^{7/6}}{7} + \frac{4x^{5/4}}{5} - 12 \log(1 + \sqrt[12]{x})$$

output

```
12*x^(1/12)-6*x^(1/6)+4*x^(1/4)-3*x^(1/3)+12/5*x^(5/12)+12/7*x^(7/12)-3/2*x^(2/3)+4/3*x^(3/4)-6/5*x^(5/6)+12/11*x^(11/12)-x+12/13*x^(13/12)-6/7*x^(7/6)+4/5*x^(5/4)-12*ln(1+x^(1/12))-2*x^(1/2)
```

3.237.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.90

$$\int \frac{1}{\frac{1}{\sqrt[3]{x}} + \frac{1}{\sqrt[4]{x}}} dx = \frac{360360 \sqrt[12]{x} - 180180 \sqrt[6]{x} + 120120 \sqrt[4]{x} - 90090 \sqrt[3]{x} + 72072x^{5/12} - 60060\sqrt{x} + 51480x^{7/12} - 45045x^{2/3} + 30030}{30030} - 12 \log(1 + \sqrt[12]{x})$$

input `Integrate[(x^(-1/3) + x^(-1/4))^-1, x]`

output $(360360*x^{(1/12)} - 180180*x^{(1/6)} + 120120*x^{(1/4)} - 90090*x^{(1/3)} + 72072*x^{(5/12)} - 60060*\text{Sqrt}[x] + 51480*x^{(7/12)} - 45045*x^{(2/3)} + 40040*x^{(3/4)} - 36036*x^{(5/6)} + 32760*x^{(11/12)} - 30030*x + 27720*x^{(13/12)} - 25740*x^{(7/6)} + 24024*x^{(5/4)})/30030 - 12*\text{Log}[1 + x^{(1/12)}]$

3.237.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.08, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {2027, 798, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\frac{1}{\sqrt[4]{x}} + \frac{1}{\sqrt[3]{x}}} dx$$

$$\downarrow 2027$$

$$\int \frac{\sqrt[3]{x}}{\sqrt[12]{x} + 1} dx$$

$$\downarrow 798$$

$$12 \int \frac{x^{5/4}}{\sqrt[12]{x} + 1} d\sqrt[12]{x}$$

$$\downarrow 49$$

$$12 \int \left(x^{7/6} - x^{13/12} + x - x^{11/12} + x^{5/6} - x^{3/4} + x^{2/3} - x^{7/12} + \sqrt{x} - x^{5/12} + \sqrt[3]{x} - \sqrt[4]{x} + \sqrt[6]{x} - \sqrt[12]{x} + \frac{1}{-\sqrt[12]{x}} \right) dx$$

$$\downarrow 2009$$

$$12 \left(\frac{x^{5/4}}{15} - \frac{x^{7/6}}{14} + \frac{x^{13/12}}{13} + \frac{x^{11/12}}{11} - \frac{x^{5/6}}{10} + \frac{x^{3/4}}{9} - \frac{x^{2/3}}{8} + \frac{x^{7/12}}{7} + \frac{x^{5/12}}{5} - \frac{x}{12} - \frac{\sqrt{x}}{6} - \frac{\sqrt[3]{x}}{4} + \frac{\sqrt[4]{x}}{3} - \frac{\sqrt[6]{x}}{2} + \sqrt[12]{x} \right)$$

input `Int[(x^(-1/3) + x^(-1/4))^-1, x]`

output $12*(x^{(1/12)} - x^{(1/6)}/2 + x^{(1/4)}/3 - x^{(1/3)}/4 + x^{(5/12)}/5 - \text{Sqrt}[x]/6 + x^{(7/12)}/7 - x^{(2/3)}/8 + x^{(3/4)}/9 - x^{(5/6)}/10 + x^{(11/12)}/11 - x/12 + x^{(13/12)}/13 - x^{(7/6)}/14 + x^{(5/4)}/15 - \text{Log}[1 + x^{(1/12)}])$

3.237.3.1 Defintions of rubi rules used

rule 49 $\text{Int}[(a + b*x)^m*(c + d*x)^n, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x$ && $\text{IGtQ}[m, 0]$ && $\text{IGtQ}[m + n + 2, 0]$

rule 798 $\text{Int}[x^m*(a + b*x^n)^p, x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[m + 1)/n} - 1)*(a + b*x)^p, x], x, x^n], x] /;$ $\text{FreeQ}\{a, b, m, n, p\}, x$ && $\text{IntegerQ}[\text{Simplify}[m + 1)/n]]$

rule 2009 $\text{Int}[u, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /;$ $\text{SumQ}[u]$

rule 2027 $\text{Int}[(F*x)^r*(a + b*x^s)^p, x_Symbol] \rightarrow \text{Int}[x^{(p*r)*(a + b*x^{(s - r)})^p}*F, x] /;$ $\text{FreeQ}\{a, b, r, s\}, x$ && $\text{IntegerQ}[p]$ && $\text{PosQ}[s - r]$ && $!(\text{EqQ}[p, 1] \&\& \text{EqQ}[u, 1])$

3.237.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.64

method	result
derivativedivides	$12x^{1/12} - 6x^{1/6} + 4x^{1/4} - 3x^{1/3} + \frac{12x^{5/12}}{5} + \frac{12x^{7/12}}{7} - \frac{3x^{2/3}}{2} + \frac{4x^{3/4}}{3} - \frac{6x^{5/6}}{5} + \frac{12x^{11/12}}{11} - x + \frac{12x^{13/12}}{13} - 6$
default	$12x^{1/12} - 6x^{1/6} + 4x^{1/4} - 3x^{1/3} + \frac{12x^{5/12}}{5} + \frac{12x^{7/12}}{7} - \frac{3x^{2/3}}{2} + \frac{4x^{3/4}}{3} - \frac{6x^{5/6}}{5} + \frac{12x^{11/12}}{11} - x + \frac{12x^{13/12}}{13} - 6$
meijerg	$\frac{x^{1/12} (48048x^{7/6} - 51480x^{13/12} + 55440x - 60060x^{11/12} + 65520x^{5/6} - 72072x^{3/4} + 80080x^{2/3} - 90090x^{7/12} + 102960\sqrt{x} - 120120x^{5/12} + 60060)}{60060}$

input $\text{int}(1/(1/x^{(1/3)}+1/x^{(1/4)}),x,\text{method}=_RETURNVERBOSE)$

3.237. $\int \frac{1}{\sqrt[3]{x} + \sqrt[4]{x}} dx$

output $12*x^{(1/12)}-6*x^{(1/6)}+4*x^{(1/4)}-3*x^{(1/3)}+12/5*x^{(5/12)}+12/7*x^{(7/12)}-3/2*x^{(2/3)}+4/3*x^{(3/4)}-6/5*x^{(5/6)}+12/11*x^{(11/12)}-x+12/13*x^{(13/12)}-6/7*x^{(7/6)}+4/5*x^{(5/4)}-12*\ln(1+x^{(1/12)})-2*x^{(1/2)}$

3.237.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.58

$$\int \frac{1}{\sqrt[3]{x} + \sqrt[4]{x}} dx = \frac{4}{5} (x+5)x^{\frac{1}{4}} - \frac{6}{7} (x+7)x^{\frac{1}{6}} + \frac{12}{13} (x+13)x^{\frac{1}{12}} - x + \frac{12}{11} x^{\frac{11}{12}} - \frac{6}{5} x^{\frac{5}{6}} + \frac{4}{3} x^{\frac{3}{4}} - \frac{3}{2} x^{\frac{2}{3}} + \frac{12}{7} x^{\frac{7}{12}} - 2\sqrt{x} + \frac{12}{5} x^{\frac{5}{12}} - 3x^{\frac{1}{3}} - 12 \log\left(x^{\frac{1}{12}} + 1\right)$$

input `integrate(1/(1/x^(1/3)+1/x^(1/4)),x, algorithm="fricas")`

output $4/5*(x + 5)*x^{(1/4)} - 6/7*(x + 7)*x^{(1/6)} + 12/13*(x + 13)*x^{(1/12)} - x + 12/11*x^{(11/12)} - 6/5*x^{(5/6)} + 4/3*x^{(3/4)} - 3/2*x^{(2/3)} + 12/7*x^{(7/12)} - 2*\sqrt{x} + 12/5*x^{(5/12)} - 3*x^{(1/3)} - 12*\log(x^{(1/12)} + 1)$

3.237.6 Sympy [A] (verification not implemented)

Time = 1.11 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.93

$$\int \frac{1}{\sqrt[3]{x} + \sqrt[4]{x}} dx = \frac{12x^{\frac{13}{12}}}{13} + \frac{12x^{\frac{11}{12}}}{11} + \frac{12x^{\frac{7}{12}}}{7} + \frac{12x^{\frac{5}{12}}}{5} + 12\sqrt[12]{x} - \frac{6x^{\frac{7}{6}}}{7} - \frac{6x^{\frac{5}{6}}}{5} - 6\sqrt[6]{x} + \frac{4x^{\frac{5}{4}}}{5} + \frac{4x^{\frac{3}{4}}}{3} + 4\sqrt[4]{x} - \frac{3x^{\frac{2}{3}}}{2} - 3\sqrt[3]{x} - 2\sqrt{x} - x - 12 \log\left(\sqrt[12]{x} + 1\right)$$

input `integrate(1/(1/x**(1/3)+1/x**(1/4)),x)`

output $12*x^{(13/12)}/13 + 12*x^{(11/12)}/11 + 12*x^{(7/12)}/7 + 12*x^{(5/12)}/5 + 12*x^{(1/12)} - 6*x^{(7/6)}/7 - 6*x^{(5/6)}/5 - 6*x^{(1/6)} + 4*x^{(5/4)}/5 + 4*x^{(3/4)}/3 + 4*x^{(1/4)} - 3*x^{(2/3)}/2 - 3*x^{(1/3)} - 2*\sqrt{x} - x - 12*\log(x^{(1/12)} + 1)$

3.237.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.63

$$\int \frac{1}{\frac{1}{\sqrt[3]{x}} + \frac{1}{\sqrt[4]{x}}} dx = \frac{4}{5} x^{\frac{5}{4}} - \frac{6}{7} x^{\frac{7}{6}} + \frac{12}{13} x^{\frac{13}{12}} - x + \frac{12}{11} x^{\frac{11}{12}} - \frac{6}{5} x^{\frac{5}{6}} + \frac{4}{3} x^{\frac{3}{4}} - \frac{3}{2} x^{\frac{2}{3}} + \frac{12}{7} x^{\frac{7}{12}} - 2\sqrt{x} + \frac{12}{5} x^{\frac{5}{12}} - 3x^{\frac{1}{3}} + 4x^{\frac{1}{4}} - 6x^{\frac{1}{6}} + 12x^{\frac{1}{12}} - 12 \log\left(x^{\frac{1}{12}} + 1\right)$$

input `integrate(1/(1/x^(1/3)+1/x^(1/4)),x, algorithm="maxima")`output `4/5*x^(5/4) - 6/7*x^(7/6) + 12/13*x^(13/12) - x + 12/11*x^(11/12) - 6/5*x^(5/6) + 4/3*x^(3/4) - 3/2*x^(2/3) + 12/7*x^(7/12) - 2*sqrt(x) + 12/5*x^(5/12) - 3*x^(1/3) + 4*x^(1/4) - 6*x^(1/6) + 12*x^(1/12) - 12*log(x^(1/12) + 1)`**3.237.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.63

$$\int \frac{1}{\frac{1}{\sqrt[3]{x}} + \frac{1}{\sqrt[4]{x}}} dx = \frac{4}{5} x^{\frac{5}{4}} - \frac{6}{7} x^{\frac{7}{6}} + \frac{12}{13} x^{\frac{13}{12}} - x + \frac{12}{11} x^{\frac{11}{12}} - \frac{6}{5} x^{\frac{5}{6}} + \frac{4}{3} x^{\frac{3}{4}} - \frac{3}{2} x^{\frac{2}{3}} + \frac{12}{7} x^{\frac{7}{12}} - 2\sqrt{x} + \frac{12}{5} x^{\frac{5}{12}} - 3x^{\frac{1}{3}} + 4x^{\frac{1}{4}} - 6x^{\frac{1}{6}} + 12x^{\frac{1}{12}} - 12 \log\left(x^{\frac{1}{12}} + 1\right)$$

input `integrate(1/(1/x^(1/3)+1/x^(1/4)),x, algorithm="giac")`output `4/5*x^(5/4) - 6/7*x^(7/6) + 12/13*x^(13/12) - x + 12/11*x^(11/12) - 6/5*x^(5/6) + 4/3*x^(3/4) - 3/2*x^(2/3) + 12/7*x^(7/12) - 2*sqrt(x) + 12/5*x^(5/12) - 3*x^(1/3) + 4*x^(1/4) - 6*x^(1/6) + 12*x^(1/12) - 12*log(x^(1/12) + 1)`

3.237.9 Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.63

$$\int \frac{1}{\sqrt[3]{x} + \sqrt[4]{x}} dx = 4x^{1/4}$$

$$- 12 \ln(x^{1/12} + 1) - 2\sqrt{x} - 3x^{1/3} - x - \frac{3x^{2/3}}{2} - 6x^{1/6} + \frac{4x^{3/4}}{3} + \frac{4x^{5/4}}{5}$$

$$- \frac{6x^{5/6}}{5} + 12x^{1/12} - \frac{6x^{7/6}}{7} + \frac{12x^{5/12}}{5} + \frac{12x^{7/12}}{7} + \frac{12x^{11/12}}{11} + \frac{12x^{13/12}}{13}$$

input `int(1/(1/x^(1/3) + 1/x^(1/4)),x)`output `4*x^(1/4) - 12*log(x^(1/12) + 1) - 2*x^(1/2) - 3*x^(1/3) - x - (3*x^(2/3))/2 - 6*x^(1/6) + (4*x^(3/4))/3 + (4*x^(5/4))/5 - (6*x^(5/6))/5 + 12*x^(1/12) - (6*x^(7/6))/7 + (12*x^(5/12))/5 + (12*x^(7/12))/7 + (12*x^(11/12))/11 + (12*x^(13/12))/13`

3.238 $\int \sqrt{\frac{1-x}{x}} dx$

3.238.1 Optimal result 1234
 3.238.2 Mathematica [A] (verified) 1234
 3.238.3 Rubi [A] (verified) 1235
 3.238.4 Maple [A] (verified) 1236
 3.238.5 Fricas [A] (verification not implemented) 1237
 3.238.6 Sympy [F] 1237
 3.238.7 Maxima [A] (verification not implemented) 1237
 3.238.8 Giac [A] (verification not implemented) 1238
 3.238.9 Mupad [B] (verification not implemented) 1238

3.238.1 Optimal result

Integrand size = 13, antiderivative size = 24

$$\int \sqrt{\frac{1-x}{x}} dx = \sqrt{-1 + \frac{1}{x}}x - \arctan\left(\sqrt{-1 + \frac{1}{x}}\right)$$

output `-arctan((-1+1/x)^(1/2))+x*(-1+1/x)^(1/2)`

3.238.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \sqrt{\frac{1-x}{x}} dx = \sqrt{-1 + \frac{1}{x}}x - \arctan\left(\sqrt{-1 + \frac{1}{x}}\right)$$

input `Integrate[Sqrt[(1 - x)/x],x]`

output `Sqrt[-1 + x^(-1)]*x - ArcTan[Sqrt[-1 + x^(-1)]]`

3.238.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {2072, 773, 51, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\frac{1-x}{x}} dx \\
 & \quad \downarrow \text{2072} \\
 & \int \sqrt{\frac{1}{x} - 1} dx \\
 & \quad \downarrow \text{773} \\
 & - \int \sqrt{\frac{1}{x} - 1} x^2 d\frac{1}{x} \\
 & \quad \downarrow \text{51} \\
 & \sqrt{\frac{1}{x} - 1} x - \frac{1}{2} \int \frac{x}{\sqrt{\frac{1}{x} - 1}} d\frac{1}{x} \\
 & \quad \downarrow \text{73} \\
 & \sqrt{\frac{1}{x} - 1} x - \int \frac{1}{1 + \frac{1}{x^2}} d\sqrt{\frac{1}{x} - 1} \\
 & \quad \downarrow \text{216} \\
 & \sqrt{\frac{1}{x} - 1} x - \arctan\left(\sqrt{\frac{1}{x} - 1}\right)
 \end{aligned}$$

input `Int[Sqrt[(1 - x)/x], x]`

output `Sqrt[-1 + x^(-1)]*x - ArcTan[Sqrt[-1 + x^(-1)]]`

3.238.3.1 Defintions of rubi rules used

rule 51 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))*Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 773 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^2, x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && !IntegerQ[p]`

rule 2072 `Int[(u_)^(p_), x_Symbol] := Int[ExpandToSum[u, x]^p, x] /; FreeQ[p, x] && BinomialQ[u, x] && !BinomialMatchQ[u, x]`

3.238.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.67

method	result	size
default	$\frac{\sqrt{-\frac{-1+x}{x}} x (2\sqrt{-x^2+x} + \arcsin(2x-1))}{2\sqrt{-x(-1+x)}}$	40
risch	$\sqrt{-\frac{-1+x}{x}} x - \frac{\arcsin(2x-1)\sqrt{-\frac{-1+x}{x}}\sqrt{-x(-1+x)}}{2(-1+x)}$	45
trager	$\sqrt{-\frac{-1+x}{x}} x - \frac{\text{RootOf}(_Z^2+1)\ln\left(2\sqrt{-\frac{-1+x}{x}}x+2\text{RootOf}(_Z^2+1)x-\text{RootOf}(_Z^2+1)\right)}{2}$	54

input `int(((1-x)/x)^(1/2), x, method=_RETURNVERBOSE)`

3.238. $\int \sqrt{\frac{1-x}{x}} dx$

output $1/2*(-(-1+x)/x)^{(1/2)}*x*(2*(-x^2+x)^{(1/2)}+\arcsin(2*x-1))/(-x*(-1+x))^{(1/2)}$

3.238.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \sqrt{\frac{1-x}{x}} dx = x\sqrt{-\frac{x-1}{x}} - \arctan\left(\sqrt{-\frac{x-1}{x}}\right)$$

input `integrate(((1-x)/x)^(1/2),x, algorithm="fricas")`

output `x*sqrt(-(x - 1)/x) - arctan(sqrt(-(x - 1)/x))`

3.238.6 Sympy [F]

$$\int \sqrt{\frac{1-x}{x}} dx = \int \sqrt{\frac{1-x}{x}} dx$$

input `integrate(((1-x)/x)**(1/2),x)`

output `Integral(sqrt((1 - x)/x), x)`

3.238.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.54

$$\int \sqrt{\frac{1-x}{x}} dx = -\frac{\sqrt{-\frac{x-1}{x}}}{\frac{x-1}{x} - 1} - \arctan\left(\sqrt{-\frac{x-1}{x}}\right)$$

input `integrate(((1-x)/x)^(1/2),x, algorithm="maxima")`

output `-sqrt(-(x - 1)/x)/((x - 1)/x - 1) - arctan(sqrt(-(x - 1)/x))`

3.238. $\int \sqrt{\frac{1-x}{x}} dx$

3.238.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17

$$\int \sqrt{\frac{1-x}{x}} dx = \frac{1}{4} \pi \operatorname{sgn}(x) + \frac{1}{2} \arcsin(2x-1) \operatorname{sgn}(x) + \sqrt{-x^2+x} \operatorname{sgn}(x)$$

input `integrate(((1-x)/x)^(1/2),x, algorithm="giac")`output `1/4*pi*sgn(x) + 1/2*arcsin(2*x - 1)*sgn(x) + sqrt(-x^2 + x)*sgn(x)`**3.238.9 Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \sqrt{\frac{1-x}{x}} dx = x \sqrt{\frac{1}{x} - 1} - \operatorname{atan}\left(\sqrt{\frac{1}{x} - 1}\right)$$

input `int((-x - 1)/x)^(1/2),x)`output `x*(1/x - 1)^(1/2) - atan((1/x - 1)^(1/2))`

$$3.239 \quad \int \frac{\cos(x)}{\sin(x) + \sin^2(x)} dx$$

3.239.1 Optimal result	1239
3.239.2 Mathematica [A] (verified)	1239
3.239.3 Rubi [A] (verified)	1240
3.239.4 Maple [A] (verified)	1241
3.239.5 Fricas [A] (verification not implemented)	1241
3.239.6 Sympy [A] (verification not implemented)	1242
3.239.7 Maxima [A] (verification not implemented)	1242
3.239.8 Giac [A] (verification not implemented)	1242
3.239.9 Mupad [B] (verification not implemented)	1243

3.239.1 Optimal result

Integrand size = 12, antiderivative size = 11

$$\int \frac{\cos(x)}{\sin(x) + \sin^2(x)} dx = \log(\sin(x)) - \log(1 + \sin(x))$$

output `ln(sin(x))-ln(1+sin(x))`

3.239.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{\sin(x) + \sin^2(x)} dx = \log(\sin(x)) - \log(1 + \sin(x))$$

input `Integrate[Cos[x]/(Sin[x] + Sin[x]^2),x]`

output `Log[Sin[x]] - Log[1 + Sin[x]]`

3.239.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3739, 1080, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(x)}{\sin^2(x) + \sin(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cos(x)}{\sin(x)^2 + \sin(x)} dx \\
 & \quad \downarrow \text{3739} \\
 & \int \frac{1}{\sin^2(x) + \sin(x)} d\sin(x) \\
 & \quad \downarrow \text{1080} \\
 & \int \left(\frac{1}{-\sin(x) - 1} + \csc(x) \right) d\sin(x) \\
 & \quad \downarrow \text{2009} \\
 & \log(\sin(x)) - \log(\sin(x) + 1)
 \end{aligned}$$

input `Int[Cos[x]/(Sin[x] + Sin[x]^2),x]`

output `Log[Sin[x]] - Log[1 + Sin[x]]`

3.239.3.1 Defintions of rubi rules used

rule 1080 `Int[((b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[x^p*(b + c*x)^p, x], x] /; FreeQ[{b, c}, x] && IntegerQ[p]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3739 Int[cos[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*((f_.)*sin[(d_.) + (e_.)*(
x_)])^(n_.) + (c_.)*((f_.)*sin[(d_.) + (e_.)*(x_)])^(n2_.))^(p_.), x_Symbol
] := Module[{g = FreeFactors[Sin[d + e*x], x]}, Simp[g/e Subst[Int[(1 - g
^2*x^2)^((m - 1)/2)*(a + b*(f*g*x)^n + c*(f*g*x)^(2*n))^p, x], x, Sin[d + e
*x]/g], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && EqQ[n2, 2*n] && Intege
rQ[(m - 1)/2]
```

3.239.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
derivativedivides	$\ln(\sin(x)) - \ln(\sin(x) + 1)$	12
default	$\ln(\sin(x)) - \ln(\sin(x) + 1)$	12
norman	$-2 \ln(1 + \tan(\frac{x}{2})) + \ln(\tan(\frac{x}{2}))$	16
parallelrisc	$-2 \ln(1 + \tan(\frac{x}{2})) + \ln(\tan(\frac{x}{2}))$	16
risc	$-2 \ln(i + e^{ix}) + \ln(e^{2ix} - 1)$	21

```
input int(cos(x)/(sin(x)+sin(x)^2),x,method=_RETURNVERBOSE)
```

```
output ln(sin(x))-ln(sin(x)+1)
```

3.239.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.18

$$\int \frac{\cos(x)}{\sin(x) + \sin^2(x)} dx = \log\left(\frac{1}{2} \sin(x)\right) - \log(\sin(x) + 1)$$

```
input integrate(cos(x)/(sin(x)+sin(x)^2),x, algorithm="fracas")
```

```
output log(1/2*sin(x)) - log(sin(x) + 1)
```

3.239. $\int \frac{\cos(x)}{\sin(x) + \sin^2(x)} dx$

3.239.6 Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int \frac{\cos(x)}{\sin(x) + \sin^2(x)} dx = -\log(\sin(x) + 1) + \log(\sin(x))$$

input `integrate(cos(x)/(sin(x)+sin(x)**2),x)`output `-log(sin(x) + 1) + log(sin(x))`**3.239.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{\sin(x) + \sin^2(x)} dx = -\log(\sin(x) + 1) + \log(\sin(x))$$

input `integrate(cos(x)/(sin(x)+sin(x)^2),x, algorithm="maxima")`output `-log(sin(x) + 1) + log(sin(x))`**3.239.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

$$\int \frac{\cos(x)}{\sin(x) + \sin^2(x)} dx = -\log(\sin(x) + 1) + \log(|\sin(x)|)$$

input `integrate(cos(x)/(sin(x)+sin(x)^2),x, algorithm="giac")`output `-log(sin(x) + 1) + log(abs(sin(x)))`

3.239.9 Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \frac{\cos(x)}{\sin(x) + \sin^2(x)} dx = -2 \operatorname{atanh}(2 \sin(x) + 1)$$

input `int(cos(x)/(sin(x) + sin(x)^2),x)`

output `-2*atanh(2*sin(x) + 1)`

3.240 $\int \frac{e^{2x}}{2+3e^x+e^{2x}} dx$

3.240.1 Optimal result	1244
3.240.2 Mathematica [A] (verified)	1244
3.240.3 Rubi [A] (verified)	1245
3.240.4 Maple [A] (verified)	1246
3.240.5 Fricas [A] (verification not implemented)	1246
3.240.6 Sympy [A] (verification not implemented)	1247
3.240.7 Maxima [A] (verification not implemented)	1247
3.240.8 Giac [A] (verification not implemented)	1247
3.240.9 Mupad [B] (verification not implemented)	1248

3.240.1 Optimal result

Integrand size = 20, antiderivative size = 17

$$\int \frac{e^{2x}}{2+3e^x+e^{2x}} dx = -\log(1+e^x) + 2\log(2+e^x)$$

output `-ln(1+exp(x))+2*ln(2+exp(x))`

3.240.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{e^{2x}}{2+3e^x+e^{2x}} dx = -\log(1+e^x) + 2\log(2+e^x)$$

input `Integrate[E^(2*x)/(2 + 3*E^x + E^(2*x)),x]`

output `-Log[1 + E^x] + 2*Log[2 + E^x]`

3.240.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {2720, 1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{e^{2x}}{3e^x + e^{2x} + 2} dx \\ & \quad \downarrow \text{2720} \\ & \int \frac{e^x}{3e^x + e^{2x} + 2} de^x \\ & \quad \downarrow \text{1141} \\ & \int \left(\frac{2}{e^x + 2} + \frac{1}{-e^x - 1} \right) de^x \\ & \quad \downarrow \text{2009} \\ & 2 \log(e^x + 2) - \log(e^x + 1) \end{aligned}$$

input `Int[E^(2*x)/(2 + 3*E^x + E^(2*x)),x]`

output `-Log[1 + E^x] + 2*Log[2 + E^x]`

3.240.3.1 Defintions of rubi rules used

rule 1141 `Int[((d_.) + (e_.)*(x_))^(m_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x] /; EqQ[p, -1] || !FractionalPowerFactorQ[q]] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p, 0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.240.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

method	result	size
default	$-\ln(1 + e^x) + 2 \ln(2 + e^x)$	16
norman	$-\ln(1 + e^x) + 2 \ln(2 + e^x)$	16
risch	$-\ln(1 + e^x) + 2 \ln(2 + e^x)$	16

```
input int(exp(2*x)/(2+3*exp(x)+exp(2*x)),x,method=_RETURNVERBOSE)
```

```
output -ln(1+exp(x))+2*ln(2+exp(x))
```

3.240.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{e^{2x}}{2 + 3e^x + e^{2x}} dx = 2 \log(e^x + 2) - \log(e^x + 1)$$

```
input integrate(exp(2*x)/(2+3*exp(x)+exp(2*x)),x, algorithm="fracas")
```

```
output 2*log(e^x + 2) - log(e^x + 1)
```

3.240.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \frac{e^{2x}}{2 + 3e^x + e^{2x}} dx = -\log(e^x + 1) + 2\log(e^x + 2)$$

input `integrate(exp(2*x)/(2+3*exp(x)+exp(2*x)),x)`output `-log(exp(x) + 1) + 2*log(exp(x) + 2)`**3.240.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{e^{2x}}{2 + 3e^x + e^{2x}} dx = 2\log(e^x + 2) - \log(e^x + 1)$$

input `integrate(exp(2*x)/(2+3*exp(x)+exp(2*x)),x, algorithm="maxima")`output `2*log(e^x + 2) - log(e^x + 1)`**3.240.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{e^{2x}}{2 + 3e^x + e^{2x}} dx = 2\log(e^x + 2) - \log(e^x + 1)$$

input `integrate(exp(2*x)/(2+3*exp(x)+exp(2*x)),x, algorithm="giac")`output `2*log(e^x + 2) - log(e^x + 1)`

3.240.9 Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{e^{2x}}{2 + 3e^x + e^{2x}} dx = 2 \ln(e^x + 2) - \ln(e^x + 1)$$

input `int(exp(2*x)/(exp(2*x) + 3*exp(x) + 2),x)`

output `2*log(exp(x) + 2) - log(exp(x) + 1)`

3.241 $\int \frac{1}{\sqrt{1+e^x}} dx$

3.241.1 Optimal result	1249
3.241.2 Mathematica [A] (verified)	1249
3.241.3 Rubi [A] (verified)	1250
3.241.4 Maple [A] (verified)	1251
3.241.5 Fricas [B] (verification not implemented)	1251
3.241.6 Sympy [A] (verification not implemented)	1252
3.241.7 Maxima [B] (verification not implemented)	1252
3.241.8 Giac [B] (verification not implemented)	1252
3.241.9 Mupad [B] (verification not implemented)	1253

3.241.1 Optimal result

Integrand size = 9, antiderivative size = 12

$$\int \frac{1}{\sqrt{1+e^x}} dx = -2\operatorname{arctanh}\left(\sqrt{1+e^x}\right)$$

output `-2*arctanh((1+exp(x))^(1/2))`

3.241.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1+e^x}} dx = -2\operatorname{arctanh}\left(\sqrt{1+e^x}\right)$$

input `Integrate[1/Sqrt[1 + E^x],x]`

output `-2*ArcTanh[Sqrt[1 + E^x]]`

3.241.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2720, 73, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt{e^x + 1}} dx \\ & \quad \downarrow \text{2720} \\ & \int \frac{e^{-x}}{\sqrt{e^x + 1}} de^x \\ & \quad \downarrow \text{73} \\ & 2 \int \frac{1}{-1 + e^{2x}} d\sqrt{1 + e^x} \\ & \quad \downarrow \text{220} \\ & -2\operatorname{arctanh}(\sqrt{e^x + 1}) \end{aligned}$$

input `Int[1/Sqrt[1 + E^x],x]`

output `-2*ArcTanh[Sqrt[1 + E^x]]`

3.241.3.1 Defintions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.241.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$-2 \operatorname{arctanh}(\sqrt{1+e^x})$	10
default	$-2 \operatorname{arctanh}(\sqrt{1+e^x})$	10

```
input int(1/(1+exp(x))^(1/2),x,method=_RETURNVERBOSE)
```

```
output -2*arctanh((1+exp(x))^(1/2))
```

3.241.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. 2(9) = 18.

Time = 0.25 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.75

$$\int \frac{1}{\sqrt{1+e^x}} dx = -\log(\sqrt{e^x+1}+1) + \log(\sqrt{e^x+1}-1)$$

```
input integrate(1/(1+exp(x))^(1/2),x, algorithm="fracas")
```

```
output -log(sqrt(e^x + 1) + 1) + log(sqrt(e^x + 1) - 1)
```

3.241.6 Sympy [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.83

$$\int \frac{1}{\sqrt{1+e^x}} dx = \log(\sqrt{e^x+1}-1) - \log(\sqrt{e^x+1}+1)$$

input `integrate(1/(1+exp(x))**(1/2),x)`

output `log(sqrt(exp(x) + 1) - 1) - log(sqrt(exp(x) + 1) + 1)`

3.241.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. 2(9) = 18.

Time = 0.20 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.75

$$\int \frac{1}{\sqrt{1+e^x}} dx = -\log(\sqrt{e^x+1}+1) + \log(\sqrt{e^x+1}-1)$$

input `integrate(1/(1+exp(x))^(1/2),x, algorithm="maxima")`

output `-log(sqrt(e^x + 1) + 1) + log(sqrt(e^x + 1) - 1)`

3.241.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. 2(9) = 18.

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.75

$$\int \frac{1}{\sqrt{1+e^x}} dx = -\log(\sqrt{e^x+1}+1) + \log(\sqrt{e^x+1}-1)$$

input `integrate(1/(1+exp(x))^(1/2),x, algorithm="giac")`

output `-log(sqrt(e^x + 1) + 1) + log(sqrt(e^x + 1) - 1)`

3.241.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{1+e^x}} dx = -2 \operatorname{atanh}(\sqrt{e^x+1})$$

input `int(1/(exp(x) + 1)^(1/2),x)`

output `-2*atanh((exp(x) + 1)^(1/2))`

3.242 $\int \sqrt{1 - e^x} dx$

3.242.1 Optimal result	1254
3.242.2 Mathematica [A] (verified)	1254
3.242.3 Rubi [A] (verified)	1255
3.242.4 Maple [A] (verified)	1256
3.242.5 Fricas [A] (verification not implemented)	1257
3.242.6 Sympy [A] (verification not implemented)	1257
3.242.7 Maxima [A] (verification not implemented)	1257
3.242.8 Giac [A] (verification not implemented)	1258
3.242.9 Mupad [B] (verification not implemented)	1258

3.242.1 Optimal result

Integrand size = 11, antiderivative size = 28

$$\int \sqrt{1 - e^x} dx = 2\sqrt{1 - e^x} - 2\operatorname{arctanh}(\sqrt{1 - e^x})$$

output `-2*arctanh((1-exp(x))^(1/2))+2*(1-exp(x))^(1/2)`

3.242.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \sqrt{1 - e^x} dx = 2\sqrt{1 - e^x} - 2\operatorname{arctanh}(\sqrt{1 - e^x})$$

input `Integrate[Sqrt[1 - E^x], x]`

output `2*Sqrt[1 - E^x] - 2*ArcTanh[Sqrt[1 - E^x]]`

3.242.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {2720, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{1 - e^x} dx \\
 & \quad \downarrow \text{2720} \\
 & \int e^{-x} \sqrt{1 - e^x} de^x \\
 & \quad \downarrow \text{60} \\
 & \int \frac{e^{-x}}{\sqrt{1 - e^x}} de^x + 2\sqrt{1 - e^x} \\
 & \quad \downarrow \text{73} \\
 & 2\sqrt{1 - e^x} - 2 \int \frac{1}{1 - e^{2x}} d\sqrt{1 - e^x} \\
 & \quad \downarrow \text{219} \\
 & 2\sqrt{1 - e^x} - 2\operatorname{arctanh}(\sqrt{1 - e^x})
 \end{aligned}$$

input `Int[Sqrt[1 - E^x], x]`

output `2*Sqrt[1 - E^x] - 2*ArcTanh[Sqrt[1 - E^x]]`

3.242.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`


```
rule 73 Int[((a_.) + (b_.)*(x_)^(m))*((c_.) + (d_.)*(x_)^(n)), x_Symbol] := With[
  {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
  d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
  Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
  inearQ[a, b, c, d, m, n, x]
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
  ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
  Q[a, 0] || LtQ[b, 0])
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
  ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
  [{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
  *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.242.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

method	result	size
risch	$-\frac{2(-1+e^x)}{\sqrt{1-e^x}} - 2 \operatorname{arctanh}(\sqrt{1-e^x})$	27
derivativedivides	$2\sqrt{1-e^x} + \ln(\sqrt{1-e^x}-1) - \ln(\sqrt{1-e^x}+1)$	36
default	$2\sqrt{1-e^x} + \ln(\sqrt{1-e^x}-1) - \ln(\sqrt{1-e^x}+1)$	36

```
input int((1-exp(x))^(1/2),x,method=_RETURNVERBOSE)
```

```
output -2*(-1+exp(x))/(1-exp(x))^(1/2)-2*arctanh((1-exp(x))^(1/2))
```

3.242.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.25

$$\int \sqrt{1 - e^x} dx = 2\sqrt{-e^x + 1} - \log(\sqrt{-e^x + 1} + 1) + \log(\sqrt{-e^x + 1} - 1)$$

input `integrate((1-exp(x))^(1/2),x, algorithm="fricas")`output `2*sqrt(-e^x + 1) - log(sqrt(-e^x + 1) + 1) + log(sqrt(-e^x + 1) - 1)`**3.242.6 Sympy [A] (verification not implemented)**

Time = 0.40 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.14

$$\int \sqrt{1 - e^x} dx = 2\sqrt{1 - e^x} + \log(\sqrt{1 - e^x} - 1) - \log(\sqrt{1 - e^x} + 1)$$

input `integrate((1-exp(x))**(1/2),x)`output `2*sqrt(1 - exp(x)) + log(sqrt(1 - exp(x)) - 1) - log(sqrt(1 - exp(x)) + 1)`**3.242.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.25

$$\int \sqrt{1 - e^x} dx = 2\sqrt{-e^x + 1} - \log(\sqrt{-e^x + 1} + 1) + \log(\sqrt{-e^x + 1} - 1)$$

input `integrate((1-exp(x))^(1/2),x, algorithm="maxima")`output `2*sqrt(-e^x + 1) - log(sqrt(-e^x + 1) + 1) + log(sqrt(-e^x + 1) - 1)`

3.242.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.32

$$\int \sqrt{1 - e^x} dx = 2\sqrt{-e^x + 1} - \log(\sqrt{-e^x + 1} + 1) + \log(-\sqrt{-e^x + 1} + 1)$$

input `integrate((1-exp(x))^(1/2),x, algorithm="giac")`output `2*sqrt(-e^x + 1) - log(sqrt(-e^x + 1) + 1) + log(-sqrt(-e^x + 1) + 1)`**3.242.9 Mupad [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.43

$$\int \sqrt{1 - e^x} dx = 2\sqrt{1 - e^x} + \frac{2e^{-\frac{x}{2}} \operatorname{asin}(e^{-\frac{x}{2}}) \sqrt{1 - e^x}}{\sqrt{1 - e^{-x}}}$$

input `int((1 - exp(x))^(1/2),x)`output `2*(1 - exp(x))^(1/2) + (2*exp(-x/2)*asin(exp(-x/2))*(1 - exp(x))^(1/2))/(1 - exp(-x))^(1/2)`

3.243 $\int \frac{1}{3-5\sin(x)} dx$

3.243.1 Optimal result	1259
3.243.2 Mathematica [A] (verified)	1259
3.243.3 Rubi [A] (verified)	1260
3.243.4 Maple [A] (verified)	1261
3.243.5 Fricas [A] (verification not implemented)	1261
3.243.6 Sympy [A] (verification not implemented)	1262
3.243.7 Maxima [A] (verification not implemented)	1262
3.243.8 Giac [A] (verification not implemented)	1262
3.243.9 Mupad [B] (verification not implemented)	1263

3.243.1 Optimal result

Integrand size = 8, antiderivative size = 43

$$\int \frac{1}{3-5\sin(x)} dx = -\frac{1}{4} \log \left(\cos \left(\frac{x}{2} \right) - 3 \sin \left(\frac{x}{2} \right) \right) + \frac{1}{4} \log \left(3 \cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right)$$

output `-1/4*ln(cos(1/2*x)-3*sin(1/2*x))+1/4*ln(3*cos(1/2*x)-sin(1/2*x))`

3.243.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int \frac{1}{3-5\sin(x)} dx = -\frac{1}{4} \log \left(\cos \left(\frac{x}{2} \right) - 3 \sin \left(\frac{x}{2} \right) \right) + \frac{1}{4} \log \left(3 \cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right)$$

input `Integrate[(3 - 5*Sin[x])^(-1),x]`

output `-1/4*Log[Cos[x/2] - 3*Sin[x/2]] + Log[3*Cos[x/2] - Sin[x/2]]/4`

3.243.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.77, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{3 - 5 \sin(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{3 - 5 \sin(x)} dx \\
 & \quad \downarrow \text{3139} \\
 & 2 \int \frac{1}{3 \tan^2\left(\frac{x}{2}\right) - 10 \tan\left(\frac{x}{2}\right) + 3} d \tan\left(\frac{x}{2}\right) \\
 & \quad \downarrow \text{1081} \\
 & 6 \int \left(\frac{1}{8(1 - 3 \tan\left(\frac{x}{2}\right))} - \frac{1}{24(3 - \tan\left(\frac{x}{2}\right))} \right) d \tan\left(\frac{x}{2}\right) \\
 & \quad \downarrow \text{2009} \\
 & 6 \left(\frac{1}{24} \log\left(3 - \tan\left(\frac{x}{2}\right)\right) - \frac{1}{24} \log\left(1 - 3 \tan\left(\frac{x}{2}\right)\right) \right)
 \end{aligned}$$

input `Int[(3 - 5*Sin[x])^(-1),x]`

output `6*(-1/24*Log[1 - 3*Tan[x/2]] + Log[3 - Tan[x/2]]/24)`

3.243.3.1 Defintions of rubi rules used

rule 1081 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3139 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

3.243.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.51

method	result	size
default	$-\frac{\ln(3 \tan(\frac{x}{2}) - 1)}{4} + \frac{\ln(\tan(\frac{x}{2}) - 3)}{4}$	22
norman	$-\frac{\ln(3 \tan(\frac{x}{2}) - 1)}{4} + \frac{\ln(\tan(\frac{x}{2}) - 3)}{4}$	22
parallelrisc	$\ln\left(\left(3 \tan\left(\frac{x}{2}\right) - 9\right)^{\frac{1}{4}}\right) + \ln\left(\frac{1}{\left(3 \tan\left(\frac{x}{2}\right) - 1\right)^{\frac{1}{4}}}\right)$	24
risc	$\frac{\ln\left(\frac{4}{5} - \frac{3i}{5} + e^{ix}\right)}{4} - \frac{\ln\left(e^{ix} - \frac{4}{5} - \frac{3i}{5}\right)}{4}$	26

input `int(1/(3-5*sin(x)),x,method=_RETURNVERBOSE)`

output `-1/4*ln(3*tan(1/2*x)-1)+1/4*ln(tan(1/2*x)-3)`

3.243.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.63

$$\int \frac{1}{3 - 5 \sin(x)} dx = \frac{1}{8} \log(4 \cos(x) - 3 \sin(x) + 5) - \frac{1}{8} \log(-4 \cos(x) - 3 \sin(x) + 5)$$

input `integrate(1/(3-5*sin(x)),x, algorithm="fricas")`

output `1/8*log(4*cos(x) - 3*sin(x) + 5) - 1/8*log(-4*cos(x) - 3*sin(x) + 5)`

3.243.6 Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.47

$$\int \frac{1}{3-5\sin(x)} dx = \frac{\log\left(\tan\left(\frac{x}{2}\right) - 3\right)}{4} - \frac{\log\left(3\tan\left(\frac{x}{2}\right) - 1\right)}{4}$$

input `integrate(1/(3-5*sin(x)),x)`output `log(tan(x/2) - 3)/4 - log(3*tan(x/2) - 1)/4`**3.243.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.70

$$\int \frac{1}{3-5\sin(x)} dx = -\frac{1}{4} \log\left(\frac{3\sin(x)}{\cos(x)+1} - 1\right) + \frac{1}{4} \log\left(\frac{\sin(x)}{\cos(x)+1} - 3\right)$$

input `integrate(1/(3-5*sin(x)),x, algorithm="maxima")`output `-1/4*log(3*sin(x)/(cos(x) + 1) - 1) + 1/4*log(sin(x)/(cos(x) + 1) - 3)`**3.243.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.53

$$\int \frac{1}{3-5\sin(x)} dx = -\frac{1}{4} \log\left(\left|3\tan\left(\frac{1}{2}x\right) - 1\right|\right) + \frac{1}{4} \log\left(\left|\tan\left(\frac{1}{2}x\right) - 3\right|\right)$$

input `integrate(1/(3-5*sin(x)),x, algorithm="giac")`output `-1/4*log(abs(3*tan(1/2*x) - 1)) + 1/4*log(abs(tan(1/2*x) - 3))`

3.243.9 Mupad [B] (verification not implemented)

Time = 0.46 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.26

$$\int \frac{1}{3 - 5 \sin(x)} dx = -\frac{\operatorname{atanh}\left(\frac{3 \tan\left(\frac{x}{2}\right)}{4} - \frac{5}{4}\right)}{2}$$

input `int(-1/(5*sin(x) - 3),x)`

output `-atanh((3*tan(x/2))/4 - 5/4)/2`

3.244 $\int \frac{1}{\cos(x)+\sin(x)} dx$

3.244.1 Optimal result	1264
3.244.2 Mathematica [C] (verified)	1264
3.244.3 Rubi [A] (verified)	1265
3.244.4 Maple [A] (verified)	1266
3.244.5 Fricas [B] (verification not implemented)	1266
3.244.6 Sympy [A] (verification not implemented)	1267
3.244.7 Maxima [B] (verification not implemented)	1267
3.244.8 Giac [B] (verification not implemented)	1267
3.244.9 Mupad [B] (verification not implemented)	1268

3.244.1 Optimal result

Integrand size = 7, antiderivative size = 21

$$\int \frac{1}{\cos(x) + \sin(x)} dx = -\frac{\operatorname{arctanh}\left(\frac{\cos(x)-\sin(x)}{\sqrt{2}}\right)}{\sqrt{2}}$$

output `-1/2*arctanh(1/2*(cos(x)-sin(x))*2^(1/2))*2^(1/2)`

3.244.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.14

$$\int \frac{1}{\cos(x) + \sin(x)} dx = (-1 - i)(-1)^{3/4} \operatorname{arctanh}\left(\frac{-1 + \tan\left(\frac{x}{2}\right)}{\sqrt{2}}\right)$$

input `Integrate[(Cos[x] + Sin[x])^(-1),x]`

output `(-1 - I)*(-1)^(3/4)*ArcTanh[(-1 + Tan[x/2])/Sqrt[2]]`

3.244.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3553, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sin(x) + \cos(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{\sin(x) + \cos(x)} dx \\ & \quad \downarrow \text{3553} \\ & - \int \frac{1}{2 - (\cos(x) - \sin(x))^2} d(\cos(x) - \sin(x)) \\ & \quad \downarrow \text{219} \\ & - \frac{\operatorname{arctanh}\left(\frac{\cos(x) - \sin(x)}{\sqrt{2}}\right)}{\sqrt{2}} \end{aligned}$$

input `Int[(Cos[x] + Sin[x])^(-1),x]`

output `-(ArcTanh[(Cos[x] - Sin[x])/Sqrt[2]]/Sqrt[2])`

3.244.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3553 Int[(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x
_Symbol] := Simp[-d^(-1) Subst[Int[1/(a^2 + b^2 - x^2), x], x, b*Cos[c +
d*x] - a*Sin[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]
```

3.244.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

method	result	size
default	$\sqrt{2} \operatorname{arctanh}\left(\frac{(2 \tan(\frac{x}{2}) - 2)\sqrt{2}}{4}\right)$	19
risch	$\frac{\sqrt{2} \ln\left(e^{ix} - \frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right)}{2} - \frac{\sqrt{2} \ln\left(e^{ix} + \frac{\sqrt{2}}{2} - \frac{i\sqrt{2}}{2}\right)}{2}$	48

```
input int(1/(cos(x)+sin(x)),x,method=_RETURNVERBOSE)
```

```
output 2^(1/2)*arctanh(1/4*(2*tan(1/2*x)-2)*2^(1/2))
```

3.244.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 38 vs. $2(18) = 36$.

Time = 0.26 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.81

$$\int \frac{1}{\cos(x) + \sin(x)} dx = \frac{1}{4} \sqrt{2} \log \left(\frac{2(\sqrt{2} - \cos(x)) \sin(x) - 2\sqrt{2} \cos(x) + 3}{2 \cos(x) \sin(x) + 1} \right)$$

```
input integrate(1/(cos(x)+sin(x)),x, algorithm="fricas")
```

```
output 1/4*sqrt(2)*log((2*(sqrt(2) - cos(x))*sin(x) - 2*sqrt(2)*cos(x) + 3)/(2*co
s(x)*sin(x) + 1))
```

3.244.6 Sympy [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.86

$$\int \frac{1}{\cos(x) + \sin(x)} dx = \frac{\sqrt{2} \log(\tan(\frac{x}{2}) - 1 + \sqrt{2})}{2} - \frac{\sqrt{2} \log(\tan(\frac{x}{2}) - \sqrt{2} - 1)}{2}$$

input `integrate(1/(cos(x)+sin(x)),x)`

output `sqrt(2)*log(tan(x/2) - 1 + sqrt(2))/2 - sqrt(2)*log(tan(x/2) - sqrt(2) - 1)/2`

3.244.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. 2(18) = 36.

Time = 0.38 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.86

$$\int \frac{1}{\cos(x) + \sin(x)} dx = -\frac{1}{2} \sqrt{2} \log \left(-\frac{\sqrt{2} - \frac{\sin(x)}{\cos(x)+1} + 1}{\sqrt{2} + \frac{\sin(x)}{\cos(x)+1} - 1} \right)$$

input `integrate(1/(cos(x)+sin(x)),x, algorithm="maxima")`

output `-1/2*sqrt(2)*log(-(sqrt(2) - sin(x)/(cos(x) + 1) + 1)/(sqrt(2) + sin(x)/(cos(x) + 1) - 1))`

3.244.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. 2(18) = 36.

Time = 0.31 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.76

$$\int \frac{1}{\cos(x) + \sin(x)} dx = -\frac{1}{2} \sqrt{2} \log \left(\frac{|-2\sqrt{2} + 2 \tan(\frac{1}{2}x) - 2|}{|2\sqrt{2} + 2 \tan(\frac{1}{2}x) - 2|} \right)$$

input `integrate(1/(cos(x)+sin(x)),x, algorithm="giac")`

output `-1/2*sqrt(2)*log(abs(-2*sqrt(2) + 2*tan(1/2*x) - 2)/abs(2*sqrt(2) + 2*tan(1/2*x) - 2))`

3.244.9 Mupad [B] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{1}{\cos(x) + \sin(x)} dx = -\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2} \tan\left(\frac{x}{2}\right)}{2}\right)$$

input `int(1/(cos(x) + sin(x)),x)`

output `-2^(1/2)*atanh(2^(1/2)/2 - (2^(1/2)*tan(x/2))/2)`

3.245 $\int \frac{1}{1-\cos(x)+\sin(x)} dx$

3.245.1 Optimal result	1269
3.245.2 Mathematica [B] (verified)	1269
3.245.3 Rubi [A] (verified)	1270
3.245.4 Maple [A] (verified)	1271
3.245.5 Fricas [A] (verification not implemented)	1271
3.245.6 Sympy [A] (verification not implemented)	1271
3.245.7 Maxima [B] (verification not implemented)	1272
3.245.8 Giac [A] (verification not implemented)	1272
3.245.9 Mupad [B] (verification not implemented)	1272

3.245.1 Optimal result

Integrand size = 10, antiderivative size = 11

$$\int \frac{1}{1-\cos(x)+\sin(x)} dx = -\log\left(1+\cot\left(\frac{x}{2}\right)\right)$$

output `-ln(1+cot(1/2*x))`

3.245.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 24 vs. 2(11) = 22.

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 2.18

$$\int \frac{1}{1-\cos(x)+\sin(x)} dx = \log\left(\sin\left(\frac{x}{2}\right)\right) - \log\left(\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)\right)$$

input `Integrate[(1 - Cos[x] + Sin[x])^(-1),x]`

output `Log[Sin[x/2]] - Log[Cos[x/2] + Sin[x/2]]`

3.245.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3042, 3600, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{1}{\sin(x) - \cos(x) + 1} dx \\
 \downarrow \text{3042} \\
 \int \frac{1}{\sin(x) - \cos(x) + 1} dx \\
 \downarrow \text{3600} \\
 - \int \frac{1}{\cot\left(\frac{x}{2}\right) + 1} d \cot\left(\frac{x}{2}\right) \\
 \downarrow \text{16} \\
 - \log\left(\cot\left(\frac{x}{2}\right) + 1\right)
 \end{array}$$

input `Int[(1 - Cos[x] + Sin[x])^(-1),x]`

output `-Log[1 + Cot[x/2]]`

3.245.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_.)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3600 `Int[(cos[(d_.) + (e_.)*(x_.)]*(b_.) + (a_.) + (c_.)*sin[(d_.) + (e_.)*(x_.)])^(-1), x_Symbol] := Module[{f = FreeFactors[Cot[(d + e*x)/2], x]}, Simp[-f/e Subst[Int[1/(a + c*f*x), x], x, Cot[(d + e*x)/2]/f], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[a + b, 0]`

3.245.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.45

method	result	size
default	$\ln\left(\tan\left(\frac{x}{2}\right)\right) - \ln\left(1 + \tan\left(\frac{x}{2}\right)\right)$	16
norman	$\ln\left(\tan\left(\frac{x}{2}\right)\right) - \ln\left(1 + \tan\left(\frac{x}{2}\right)\right)$	16
parallelrisch	$\ln\left(\tan\left(\frac{x}{2}\right)\right) - \ln\left(1 + \tan\left(\frac{x}{2}\right)\right)$	16
risch	$\ln\left(e^{ix} - 1\right) - \ln\left(i + e^{ix}\right)$	21

input `int(1/(1-cos(x)+sin(x)),x,method=_RETURNVERBOSE)`output `ln(tan(1/2*x))-ln(1+tan(1/2*x))`**3.245.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.55

$$\int \frac{1}{1 - \cos(x) + \sin(x)} dx = \frac{1}{2} \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right) - \frac{1}{2} \log(\sin(x) + 1)$$

input `integrate(1/(1-cos(x)+sin(x)),x, algorithm="fricas")`output `1/2*log(-1/2*cos(x) + 1/2) - 1/2*log(sin(x) + 1)`**3.245.6 Sympy [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.27

$$\int \frac{1}{1 - \cos(x) + \sin(x)} dx = -\log\left(\tan\left(\frac{x}{2}\right) + 1\right) + \log\left(\tan\left(\frac{x}{2}\right)\right)$$

input `integrate(1/(1-cos(x)+sin(x)),x)`output `-log(tan(x/2) + 1) + log(tan(x/2))`

3.245.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. 2(9) = 18.

Time = 0.21 (sec) , antiderivative size = 25, normalized size of antiderivative = 2.27

$$\int \frac{1}{1 - \cos(x) + \sin(x)} dx = -\log\left(\frac{\sin(x)}{\cos(x) + 1} + 1\right) + \log\left(\frac{\sin(x)}{\cos(x) + 1}\right)$$

input `integrate(1/(1-cos(x)+sin(x)),x, algorithm="maxima")`

output `-log(sin(x)/(cos(x) + 1) + 1) + log(sin(x)/(cos(x) + 1))`

3.245.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.55

$$\int \frac{1}{1 - \cos(x) + \sin(x)} dx = -\log\left(\left|\tan\left(\frac{1}{2}x\right) + 1\right|\right) + \log\left(\left|\tan\left(\frac{1}{2}x\right)\right|\right)$$

input `integrate(1/(1-cos(x)+sin(x)),x, algorithm="giac")`

output `-log(abs(tan(1/2*x) + 1)) + log(abs(tan(1/2*x)))`

3.245.9 Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 - \cos(x) + \sin(x)} dx = -2 \operatorname{atanh}\left(2 \tan\left(\frac{x}{2}\right) + 1\right)$$

input `int(1/(sin(x) - cos(x) + 1),x)`

output `-2*atanh(2*tan(x/2) + 1)`

$$3.246 \quad \int \frac{1}{4 \cos(x) + 3 \sin(x)} dx$$

3.246.1 Optimal result	1273
3.246.2 Mathematica [B] (verified)	1273
3.246.3 Rubi [A] (verified)	1274
3.246.4 Maple [A] (verified)	1275
3.246.5 Fricas [B] (verification not implemented)	1275
3.246.6 Sympy [A] (verification not implemented)	1276
3.246.7 Maxima [B] (verification not implemented)	1276
3.246.8 Giac [A] (verification not implemented)	1276
3.246.9 Mupad [B] (verification not implemented)	1277

3.246.1 Optimal result

Integrand size = 11, antiderivative size = 18

$$\int \frac{1}{4 \cos(x) + 3 \sin(x)} dx = -\frac{1}{5} \operatorname{arctanh} \left(\frac{1}{5} (3 \cos(x) - 4 \sin(x)) \right)$$

output `-1/5*arctanh(3/5*cos(x)-4/5*sin(x))`

3.246.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 43 vs. $2(18) = 36$.

Time = 0.03 (sec) , antiderivative size = 43, normalized size of antiderivative = 2.39

$$\int \frac{1}{4 \cos(x) + 3 \sin(x)} dx = -\frac{1}{5} \log \left(2 \cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right) + \frac{1}{5} \log \left(\cos \left(\frac{x}{2} \right) + 2 \sin \left(\frac{x}{2} \right) \right)$$

input `Integrate[(4*Cos[x] + 3*Sin[x])^(-1),x]`

output `-1/5*Log[2*Cos[x/2] - Sin[x/2]] + Log[Cos[x/2] + 2*Sin[x/2]]/5`

3.246.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 3553, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{3 \sin(x) + 4 \cos(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{3 \sin(x) + 4 \cos(x)} dx \\ & \quad \downarrow \text{3553} \\ & - \int \frac{1}{25 - (3 \cos(x) - 4 \sin(x))^2} d(3 \cos(x) - 4 \sin(x)) \\ & \quad \downarrow \text{219} \\ & -\frac{1}{5} \operatorname{arctanh}\left(\frac{1}{5}(3 \cos(x) - 4 \sin(x))\right) \end{aligned}$$

input `Int[(4*Cos[x] + 3*Sin[x])^(-1),x]`

output `-1/5*ArcTanh[(3*Cos[x] - 4*Sin[x])/5]`

3.246.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3553 Int[(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x
_Symbol] := Simp[-d^(-1) Subst[Int[1/(a^2 + b^2 - x^2), x], x, b*Cos[c +
d*x] - a*Sin[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]
```

3.246.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

method	result	size
default	$\frac{\ln(2 \tan(\frac{x}{2}) + 1)}{5} - \frac{\ln(\tan(\frac{x}{2}) - 2)}{5}$	22
norman	$\frac{\ln(2 \tan(\frac{x}{2}) + 1)}{5} - \frac{\ln(\tan(\frac{x}{2}) - 2)}{5}$	22
parallelrisch	$\ln\left(\frac{1}{(2 \tan(\frac{x}{2}) - 4)^{\frac{1}{5}}}\right) + \ln\left((2 \tan(\frac{x}{2}) + 1)^{\frac{1}{5}}\right)$	24
risch	$\frac{\ln(e^{ix} - \frac{3}{5} + \frac{4i}{5})}{5} - \frac{\ln(e^{ix} + \frac{3}{5} - \frac{4i}{5})}{5}$	26

```
input int(1/(4*cos(x)+3*sin(x)),x,method=_RETURNVERBOSE)
```

```
output 1/5*ln(2*tan(1/2*x)+1)-1/5*ln(tan(1/2*x)-2)
```

3.246.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(12) = 24.

Time = 0.27 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.50

$$\int \frac{1}{4 \cos(x) + 3 \sin(x)} dx = -\frac{1}{10} \log\left(\frac{3}{2} \cos(x) - 2 \sin(x) + \frac{5}{2}\right) + \frac{1}{10} \log\left(-\frac{3}{2} \cos(x) + 2 \sin(x) + \frac{5}{2}\right)$$

```
input integrate(1/(4*cos(x)+3*sin(x)),x, algorithm="fricas")
```

```
output -1/10*log(3/2*cos(x) - 2*sin(x) + 5/2) + 1/10*log(-3/2*cos(x) + 2*sin(x) +
5/2)
```

3.246.6 Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{1}{4 \cos(x) + 3 \sin(x)} dx = -\frac{\log\left(\tan\left(\frac{x}{2}\right) - 2\right)}{5} + \frac{\log\left(2 \tan\left(\frac{x}{2}\right) + 1\right)}{5}$$

input `integrate(1/(4*cos(x)+3*sin(x)),x)`

output `-log(tan(x/2) - 2)/5 + log(2*tan(x/2) + 1)/5`

3.246.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 30 vs. 2(12) = 24.

Time = 0.32 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.67

$$\int \frac{1}{4 \cos(x) + 3 \sin(x)} dx = \frac{1}{5} \log\left(\frac{2 \sin(x)}{\cos(x) + 1} + 1\right) - \frac{1}{5} \log\left(\frac{\sin(x)}{\cos(x) + 1} - 2\right)$$

input `integrate(1/(4*cos(x)+3*sin(x)),x, algorithm="maxima")`

output `1/5*log(2*sin(x)/(cos(x) + 1) + 1) - 1/5*log(sin(x)/(cos(x) + 1) - 2)`

3.246.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.28

$$\int \frac{1}{4 \cos(x) + 3 \sin(x)} dx = \frac{1}{5} \log\left(\left|2 \tan\left(\frac{1}{2} x\right) + 1\right|\right) - \frac{1}{5} \log\left(\left|\tan\left(\frac{1}{2} x\right) - 2\right|\right)$$

input `integrate(1/(4*cos(x)+3*sin(x)),x, algorithm="giac")`

output `1/5*log(abs(2*tan(1/2*x) + 1)) - 1/5*log(abs(tan(1/2*x) - 2))`

3.246.9 Mupad [B] (verification not implemented)

Time = 0.50 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.61

$$\int \frac{1}{4 \cos(x) + 3 \sin(x)} dx = \frac{2 \operatorname{atanh}\left(\frac{4 \tan\left(\frac{x}{2}\right) - 3}{5}\right)}{5}$$

input `int(1/(4*cos(x) + 3*sin(x)),x)`

output `(2*atanh((4*tan(x/2))/5 - 3/5))/5`

3.247 $\int \frac{1}{\sin(x)+\tan(x)} dx$

3.247.1 Optimal result	1278
3.247.2 Mathematica [A] (verified)	1278
3.247.3 Rubi [A] (verified)	1279
3.247.4 Maple [A] (verified)	1281
3.247.5 Fricas [A] (verification not implemented)	1282
3.247.6 Sympy [F]	1282
3.247.7 Maxima [A] (verification not implemented)	1282
3.247.8 Giac [A] (verification not implemented)	1283
3.247.9 Mupad [B] (verification not implemented)	1283

3.247.1 Optimal result

Integrand size = 7, antiderivative size = 24

$$\int \frac{1}{\sin(x) + \tan(x)} dx = -\frac{1}{2} \operatorname{arctanh}(\cos(x)) + \frac{1}{2} \cot(x) \csc(x) - \frac{\csc^2(x)}{2}$$

output `-1/2*arctanh(cos(x))+1/2*cot(x)*csc(x)-1/2*csc(x)^2`

3.247.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.46

$$\int \frac{1}{\sin(x) + \tan(x)} dx = -\frac{1}{2} \log\left(\cos\left(\frac{x}{2}\right)\right) + \frac{1}{2} \log\left(\sin\left(\frac{x}{2}\right)\right) - \frac{1}{4} \sec^2\left(\frac{x}{2}\right)$$

input `Integrate[(Sin[x] + Tan[x])^(-1), x]`

output `-1/2*Log[Cos[x/2]] + Log[Sin[x/2]]/2 - Sec[x/2]^2/4`

3.247.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 1.857$, Rules used = {3042, 4897, 3042, 25, 3185, 25, 3042, 25, 3086, 15, 3091, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sin(x) + \tan(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sin(x) + \tan(x)} dx \\
 & \quad \downarrow \text{4897} \\
 & \int \frac{\cot(x)}{\cos(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\tan\left(x - \frac{\pi}{2}\right)}{1 - \sin\left(x - \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{\tan\left(x - \frac{\pi}{2}\right)}{1 - \sin\left(x - \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{3185} \\
 & -\int \cot^2(x) \csc(x) dx - \int -\cot(x) \csc^2(x) dx \\
 & \quad \downarrow \text{25} \\
 & \int \cot(x) \csc^2(x) dx - \int \cot^2(x) \csc(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\sec\left(x - \frac{\pi}{2}\right)^2 \tan\left(x - \frac{\pi}{2}\right) dx - \int \sec\left(x - \frac{\pi}{2}\right) \tan\left(x - \frac{\pi}{2}\right)^2 dx \\
 & \quad \downarrow \text{25} \\
 & -\int \sec\left(x - \frac{\pi}{2}\right)^2 \tan\left(x - \frac{\pi}{2}\right) dx - \int \sec\left(x - \frac{\pi}{2}\right) \tan\left(x - \frac{\pi}{2}\right)^2 dx \\
 & \quad \downarrow \text{3086}
 \end{aligned}$$

$$\begin{aligned}
& - \int \csc(x) d \csc(x) - \int \sec\left(x - \frac{\pi}{2}\right) \tan\left(x - \frac{\pi}{2}\right)^2 dx \\
& \quad \downarrow \text{15} \\
& - \int \sec\left(x - \frac{\pi}{2}\right) \tan\left(x - \frac{\pi}{2}\right)^2 dx - \frac{1}{2} \csc^2(x) \\
& \quad \downarrow \text{3091} \\
& \frac{\int \csc(x) dx}{2} - \frac{1}{2} \csc^2(x) + \frac{1}{2} \cot(x) \csc(x) \\
& \quad \downarrow \text{3042} \\
& \frac{\int \csc(x) dx}{2} - \frac{1}{2} \csc^2(x) + \frac{1}{2} \cot(x) \csc(x) \\
& \quad \downarrow \text{4257} \\
& -\frac{1}{2} \operatorname{arctanh}(\cos(x)) - \frac{1}{2} \csc^2(x) + \frac{1}{2} \cot(x) \csc(x)
\end{aligned}$$

input `Int[(Sin[x] + Tan[x])^(-1),x]`

output `-1/2*ArcTanh[Cos[x]] + (Cot[x]*Csc[x])/2 - Csc[x]^2/2`

3.247.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)] + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

```
rule 3091 Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Simp[b^2*((n - 1)/(m + n - 1)) Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]
```

```
rule 3185 Int[((g_.)*tan[(e_.) + (f_.)*(x_)])^(p_.)/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/a Int[Sec[e + f*x]^2*(g*Tan[e + f*x])^p, x], x] - Simp[1/(b*g) Int[Sec[e + f*x]*(g*Tan[e + f*x])^(p + 1), x], x] /; FreeQ[{a, b, e, f, g, p}, x] && EqQ[a^2 - b^2, 0] && NeQ[p, -1]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4897 Int[u_, x_Symbol] := Int[TrigSimplify[u], x] /; TrigSimplifyQ[u]
```

3.247.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

method	result	size
default	$-\frac{1}{2(\cos(x)+1)} - \frac{\ln(\cos(x)+1)}{4} + \frac{\ln(-1+\cos(x))}{4}$	24
risch	$-\frac{e^{ix}}{(e^{ix}+1)^2} - \frac{\ln(e^{ix}+1)}{2} + \frac{\ln(e^{ix}-1)}{2}$	38

```
input int(1/(sin(x)+tan(x)),x,method=_RETURNVERBOSE)
```

```
output -1/2/(cos(x)+1)-1/4*ln(cos(x)+1)+1/4*ln(-1+cos(x))
```

3.247.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.46

$$\int \frac{1}{\sin(x) + \tan(x)} dx$$

$$= -\frac{(\cos(x) + 1) \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - (\cos(x) + 1) \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right) + 2}{4(\cos(x) + 1)}$$

input `integrate(1/(sin(x)+tan(x)),x, algorithm="fracas")`output `-1/4*((cos(x) + 1)*log(1/2*cos(x) + 1/2) - (cos(x) + 1)*log(-1/2*cos(x) + 1/2) + 2)/(cos(x) + 1)`**3.247.6 Sympy [F]**

$$\int \frac{1}{\sin(x) + \tan(x)} dx = \int \frac{1}{\sin(x) + \tan(x)} dx$$

input `integrate(1/(sin(x)+tan(x)),x)`output `Integral(1/(sin(x) + tan(x)), x)`**3.247.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.04

$$\int \frac{1}{\sin(x) + \tan(x)} dx = -\frac{\sin(x)^2}{4(\cos(x) + 1)^2} + \frac{1}{2} \log\left(\frac{\sin(x)}{\cos(x) + 1}\right)$$

input `integrate(1/(sin(x)+tan(x)),x, algorithm="maxima")`output `-1/4*sin(x)^2/(cos(x) + 1)^2 + 1/2*log(sin(x)/(cos(x) + 1))`

3.247.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17

$$\int \frac{1}{\sin(x) + \tan(x)} dx = \frac{\cos(x) - 1}{4(\cos(x) + 1)} + \frac{1}{4} \log\left(-\frac{\cos(x) - 1}{\cos(x) + 1}\right)$$

input `integrate(1/(sin(x)+tan(x)),x, algorithm="giac")`output `1/4*(cos(x) - 1)/(cos(x) + 1) + 1/4*log(-(cos(x) - 1)/(cos(x) + 1))`**3.247.9 Mupad [B] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sin(x) + \tan(x)} dx = \frac{\ln\left(\tan\left(\frac{x}{2}\right)\right)}{2} - \frac{\tan\left(\frac{x}{2}\right)^2}{4}$$

input `int(1/(sin(x) + tan(x)),x)`output `log(tan(x/2))/2 - tan(x/2)^2/4`

3.248 $\int \frac{1}{2 \sin(x) + \sin(2x)} dx$

3.248.1 Optimal result	1284
3.248.2 Mathematica [A] (verified)	1284
3.248.3 Rubi [A] (verified)	1285
3.248.4 Maple [A] (verified)	1286
3.248.5 Fricas [B] (verification not implemented)	1287
3.248.6 Sympy [F]	1287
3.248.7 Maxima [B] (verification not implemented)	1287
3.248.8 Giac [A] (verification not implemented)	1288
3.248.9 Mupad [B] (verification not implemented)	1288

3.248.1 Optimal result

Integrand size = 11, antiderivative size = 24

$$\int \frac{1}{2 \sin(x) + \sin(2x)} dx = \frac{1}{4} \log \left(\tan \left(\frac{x}{2} \right) \right) + \frac{1}{8} \tan^2 \left(\frac{x}{2} \right)$$

output `1/4*ln(tan(1/2*x))+1/8*tan(1/2*x)^2`

3.248.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.62

$$\int \frac{1}{2 \sin(x) + \sin(2x)} dx = \frac{1 - 2 \cos^2 \left(\frac{x}{2} \right) (\log (\cos \left(\frac{x}{2} \right)) - \log (\sin \left(\frac{x}{2} \right)))}{4(1 + \cos(x))}$$

input `Integrate[(2*Sin[x] + Sin[2*x])^(-1),x]`

output `(1 - 2*Cos[x/2]^2*(Log[Cos[x/2]] - Log[Sin[x/2]]))/(4*(1 + Cos[x]))`

3.248.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 4826, 27, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{2 \sin(x) + \sin(2x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{2 \sin(x) + \sin(2x)} dx \\
 & \quad \downarrow \text{4826} \\
 & 2 \int \frac{1}{8} \cot\left(\frac{x}{2}\right) \left(\tan^2\left(\frac{x}{2}\right) + 1\right) d \tan\left(\frac{x}{2}\right) \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{4} \int \cot\left(\frac{x}{2}\right) \left(\tan^2\left(\frac{x}{2}\right) + 1\right) d \tan\left(\frac{x}{2}\right) \\
 & \quad \downarrow \text{244} \\
 & \frac{1}{4} \int \left(\cot\left(\frac{x}{2}\right) + \tan\left(\frac{x}{2}\right)\right) d \tan\left(\frac{x}{2}\right) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{4} \left(\frac{1}{2} \tan^2\left(\frac{x}{2}\right) + \log\left(\tan\left(\frac{x}{2}\right)\right) \right)
 \end{aligned}$$

input `Int[(2*Sin[x] + Sin[2*x])^(-1),x]`

output `(Log[Tan[x/2]] + Tan[x/2]^2/2)/4`

3.248.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 244 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4826 `Int[((a_)*sin[(m_)*((c_) + (d_)*(x_))] + (b_)*sin[(n_)*((c_) + (d_)*(x_))])^(p_), x_Symbol] := Simp[2/d Subst[Int[Simplify[TrigExpand[a*Sin[2*m*ArcTan[x]] + b*Sin[2*n*ArcTan[x]]]]^p/(1 + x^2), x], x, Tan[(1/2)*(c + d*x)]], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IntegerQ[m/2] && IntegerQ[(n - 1)/2]`

3.248.4 Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{\ln(-1+\cos(x))}{8} + \frac{1}{4\cos(x)+4} - \frac{\ln(\cos(x)+1)}{8}$	24
risch	$\frac{e^{ix}}{2(e^{ix}+1)^2} - \frac{\ln(e^{ix}+1)}{4} + \frac{\ln(e^{ix}-1)}{4}$	38

input `int(1/(2*sin(x)+sin(2*x)),x,method=_RETURNVERBOSE)`

output `1/8*ln(-1+cos(x))+1/4/(cos(x)+1)-1/8*ln(cos(x)+1)`

3.248.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 35 vs. $2(16) = 32$.

Time = 0.27 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.46

$$\int \frac{1}{2 \sin(x) + \sin(2x)} dx$$

$$= -\frac{(\cos(x) + 1) \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - (\cos(x) + 1) \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right) - 2}{8(\cos(x) + 1)}$$

input `integrate(1/(2*sin(x)+sin(2*x)),x, algorithm="fricas")`

output `-1/8*((cos(x) + 1)*log(1/2*cos(x) + 1/2) - (cos(x) + 1)*log(-1/2*cos(x) + 1/2) - 2)/(cos(x) + 1)`

3.248.6 Sympy [F]

$$\int \frac{1}{2 \sin(x) + \sin(2x)} dx = \int \frac{1}{2 \sin(x) + \sin(2x)} dx$$

input `integrate(1/(2*sin(x)+sin(2*x)),x)`

output `Integral(1/(2*sin(x) + sin(2*x)), x)`

3.248.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 220 vs. $2(16) = 32$.

Time = 0.21 (sec) , antiderivative size = 220, normalized size of antiderivative = 9.17

$$\int \frac{1}{2 \sin(x) + \sin(2x)} dx$$

$$= \frac{4 \cos(2x) \cos(x) + 8 \cos(x)^2 - (2(2 \cos(x) + 1) \cos(2x) + \cos(2x)^2 + 4 \cos(x)^2 + \sin(2x)^2 + 4 \sin(x))}{8 \cos(x) (2 \cos(x) + 1)}$$

input `integrate(1/(2*sin(x)+sin(2*x)),x, algorithm="maxima")`

output `1/8*(4*cos(2*x)*cos(x) + 8*cos(x)^2 - (2*(2*cos(x) + 1)*cos(2*x) + cos(2*x)^2 + 4*cos(x)^2 + sin(2*x)^2 + 4*sin(2*x)*sin(x) + 4*sin(x)^2 + 4*cos(x) + 1)*log(cos(x)^2 + sin(x)^2 + 2*cos(x) + 1) + (2*(2*cos(x) + 1)*cos(2*x) + cos(2*x)^2 + 4*cos(x)^2 + sin(2*x)^2 + 4*sin(2*x)*sin(x) + 4*sin(x)^2 + 4*cos(x) + 1)*log(cos(x)^2 + sin(x)^2 - 2*cos(x) + 1) + 4*sin(2*x)*sin(x) + 8*sin(x)^2 + 4*cos(x))/(2*(2*cos(x) + 1)*cos(2*x) + cos(2*x)^2 + 4*cos(x)^2 + sin(2*x)^2 + 4*sin(2*x)*sin(x) + 4*sin(x)^2 + 4*cos(x) + 1)`

3.248.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17

$$\int \frac{1}{2\sin(x) + \sin(2x)} dx = -\frac{\cos(x) - 1}{8(\cos(x) + 1)} + \frac{1}{8} \log\left(-\frac{\cos(x) - 1}{\cos(x) + 1}\right)$$

input `integrate(1/(2*sin(x)+sin(2*x)),x, algorithm="giac")`

output `-1/8*(cos(x) - 1)/(cos(x) + 1) + 1/8*log(-(cos(x) - 1)/(cos(x) + 1))`

3.248.9 Mupad [B] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \frac{1}{2\sin(x) + \sin(2x)} dx = \frac{\ln\left(\tan\left(\frac{x}{2}\right)\right)}{4} + \frac{\tan\left(\frac{x}{2}\right)^2}{8}$$

input `int(1/(sin(2*x) + 2*sin(x)),x)`

output `log(tan(x/2))/4 + tan(x/2)^2/8`

3.249 $\int \frac{\sec(x)}{1+\sin(x)} dx$

3.249.1 Optimal result	1289
3.249.2 Mathematica [A] (verified)	1289
3.249.3 Rubi [A] (verified)	1290
3.249.4 Maple [A] (verified)	1291
3.249.5 Fricas [B] (verification not implemented)	1291
3.249.6 Sympy [F]	1292
3.249.7 Maxima [A] (verification not implemented)	1292
3.249.8 Giac [A] (verification not implemented)	1292
3.249.9 Mupad [B] (verification not implemented)	1293

3.249.1 Optimal result

Integrand size = 9, antiderivative size = 18

$$\int \frac{\sec(x)}{1+\sin(x)} dx = \frac{1}{2} \operatorname{arctanh}(\sin(x)) - \frac{1}{2(1+\sin(x))}$$

output `1/2*arctanh(sin(x))-1/2/(1+sin(x))`

3.249.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{\sec(x)}{1+\sin(x)} dx = \frac{1}{2} \operatorname{arctanh}(\sin(x)) - \frac{1}{2(1+\sin(x))}$$

input `Integrate[Sec[x]/(1 + Sin[x]),x]`

output `ArcTanh[Sin[x]]/2 - 1/(2*(1 + Sin[x]))`

3.249.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 3146, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(x)}{\sin(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(\sin(x) + 1) \cos(x)} dx \\
 & \quad \downarrow \text{3146} \\
 & \int \frac{1}{(1 - \sin(x))(\sin(x) + 1)^2} d\sin(x) \\
 & \quad \downarrow \text{54} \\
 & \int \left(\frac{1}{2(\sin(x) + 1)^2} - \frac{1}{2(\sin^2(x) - 1)} \right) d\sin(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} \operatorname{arctanh}(\sin(x)) - \frac{1}{2(\sin(x) + 1)}
 \end{aligned}$$

input `Int[Sec[x]/(1 + Sin[x]),x]`

output `ArcTanh[Sin[x]]/2 - 1/(2*(1 + Sin[x]))`

3.249.3.1 Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3146 Int[cos[(e_.) + (f_.)*(x_.)]^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)])^(m
_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x
)^(p - 1)/2], x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && I
ntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/
2])
```

3.249.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.33

method	result	size
default	$-\frac{\ln(\sin(x)-1)}{4} - \frac{1}{2(\sin(x)+1)} + \frac{\ln(\sin(x)+1)}{4}$	24
norman	$\frac{\tan(\frac{x}{2})}{(1+\tan(\frac{x}{2}))^2} - \frac{\ln(\tan(\frac{x}{2})-1)}{2} + \frac{\ln(1+\tan(\frac{x}{2}))}{2}$	33
parallelrisch	$\ln\left(\frac{1}{\sqrt{-\cot(x)+\csc(x)-1}}\right) + \ln\left(\sqrt{-\cot(x)+1+\csc(x)}\right) - \frac{\tan^2(x)}{2} + \frac{\sec(x)\tan(x)}{2}$	36
risch	$-\frac{ie^{ix}}{(i+e^{ix})^2} - \frac{\ln(e^{ix}-i)}{2} + \frac{\ln(i+e^{ix})}{2}$	42

```
input int(sec(x)/(sin(x)+1),x,method=_RETURNVERBOSE)
```

```
output -1/4*ln(sin(x)-1)-1/2/(sin(x)+1)+1/4*ln(sin(x)+1)
```

3.249.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 33 vs. $2(14) = 28$.

Time = 0.26 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.83

$$\int \frac{\sec(x)}{1 + \sin(x)} dx = \frac{(\sin(x) + 1) \log(\sin(x) + 1) - (\sin(x) + 1) \log(-\sin(x) + 1) - 2}{4(\sin(x) + 1)}$$

```
input integrate(sec(x)/(1+sin(x)),x, algorithm="fricas")
```

output $1/4*((\sin(x) + 1)*\log(\sin(x) + 1) - (\sin(x) + 1)*\log(-\sin(x) + 1) - 2)/(\sin(x) + 1)$

3.249.6 Sympy [F]

$$\int \frac{\sec(x)}{1 + \sin(x)} dx = \int \frac{\sec(x)}{\sin(x) + 1} dx$$

input `integrate(sec(x)/(1+sin(x)),x)`

output `Integral(sec(x)/(sin(x) + 1), x)`

3.249.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.28

$$\int \frac{\sec(x)}{1 + \sin(x)} dx = -\frac{1}{2(\sin(x) + 1)} + \frac{1}{4} \log(\sin(x) + 1) - \frac{1}{4} \log(\sin(x) - 1)$$

input `integrate(sec(x)/(1+sin(x)),x, algorithm="maxima")`

output `-1/2/(sin(x) + 1) + 1/4*log(sin(x) + 1) - 1/4*log(sin(x) - 1)`

3.249.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.39

$$\int \frac{\sec(x)}{1 + \sin(x)} dx = -\frac{1}{2(\sin(x) + 1)} + \frac{1}{4} \log(\sin(x) + 1) - \frac{1}{4} \log(-\sin(x) + 1)$$

input `integrate(sec(x)/(1+sin(x)),x, algorithm="giac")`

output `-1/2/(sin(x) + 1) + 1/4*log(sin(x) + 1) - 1/4*log(-sin(x) + 1)`

3.249.9 Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

$$\int \frac{\sec(x)}{1 + \sin(x)} dx = \frac{\ln\left(\tan\left(\frac{x}{2} + \frac{\pi}{4}\right)\right)}{2} - \frac{1}{2(\sin(x) + 1)}$$

input `int(1/(cos(x)*(sin(x) + 1)),x)`

output `log(tan(x/2 + pi/4))/2 - 1/(2*(sin(x) + 1))`

3.250 $\int \frac{1}{b \cos(x) + a \sin(x)} dx$

3.250.1 Optimal result	1294
3.250.2 Mathematica [A] (verified)	1294
3.250.3 Rubi [A] (verified)	1295
3.250.4 Maple [A] (verified)	1296
3.250.5 Fricas [B] (verification not implemented)	1296
3.250.6 Sympy [C] (verification not implemented)	1297
3.250.7 Maxima [A] (verification not implemented)	1297
3.250.8 Giac [A] (verification not implemented)	1298
3.250.9 Mupad [B] (verification not implemented)	1298

3.250.1 Optimal result

Integrand size = 11, antiderivative size = 36

$$\int \frac{1}{b \cos(x) + a \sin(x)} dx = -\frac{\operatorname{arctanh}\left(\frac{a \cos(x) - b \sin(x)}{\sqrt{a^2 + b^2}}\right)}{\sqrt{a^2 + b^2}}$$

output `-arctanh((a*cos(x)-b*sin(x))/(a^2+b^2)^(1/2))/(a^2+b^2)^(1/2)`

3.250.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.06

$$\int \frac{1}{b \cos(x) + a \sin(x)} dx = \frac{2 \operatorname{arctanh}\left(\frac{-a + b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2 + b^2}}\right)}{\sqrt{a^2 + b^2}}$$

input `Integrate[(b*Cos[x] + a*Sin[x])^(-1),x]`

output `(2*ArcTanh[(-a + b*Tan[x/2])/Sqrt[a^2 + b^2]]/Sqrt[a^2 + b^2])`

3.250.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 3553, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{a \sin(x) + b \cos(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{a \sin(x) + b \cos(x)} dx \\ & \quad \downarrow \text{3553} \\ & - \int \frac{1}{a^2 + b^2 - (a \cos(x) - b \sin(x))^2} d(a \cos(x) - b \sin(x)) \\ & \quad \downarrow \text{219} \\ & - \frac{\operatorname{arctanh}\left(\frac{a \cos(x) - b \sin(x)}{\sqrt{a^2 + b^2}}\right)}{\sqrt{a^2 + b^2}} \end{aligned}$$

input `Int[(b*cos[x] + a*sin[x])^(-1),x]`

output `-(ArcTanh[(a*cos[x] - b*sin[x])/Sqrt[a^2 + b^2]]/Sqrt[a^2 + b^2])`

3.250.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`


```
rule 3553 Int[(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x
_Symbol] := Simp[-d^(-1) Subst[Int[1/(a^2 + b^2 - x^2), x], x, b*Cos[c +
d*x] - a*Sin[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]
```

3.250.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.97

method	result	size
default	$-\frac{2 \operatorname{arctanh}\left(\frac{-2b \tan\left(\frac{x}{2}\right)+2a}{2\sqrt{a^2+b^2}}\right)}{\sqrt{a^2+b^2}}$	35
risch	$\frac{\ln\left(e^{ix} + \frac{ib-a}{\sqrt{a^2+b^2}}\right)}{\sqrt{a^2+b^2}} - \frac{\ln\left(e^{ix} - \frac{ib-a}{\sqrt{a^2+b^2}}\right)}{\sqrt{a^2+b^2}}$	74

```
input int(1/(b*cos(x)+a*sin(x)),x,method=_RETURNVERBOSE)
```

```
output -2/(a^2+b^2)^(1/2)*arctanh(1/2*(-2*b*tan(1/2*x)+2*a)/(a^2+b^2)^(1/2))
```

3.250.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 98 vs. $2(32) = 64$.

Time = 0.27 (sec) , antiderivative size = 98, normalized size of antiderivative = 2.72

$$\int \frac{1}{b \cos(x) + a \sin(x)} dx = \frac{\log\left(-\frac{2ab \cos(x) \sin(x) - (a^2 - b^2) \cos(x)^2 - a^2 - 2b^2 + 2\sqrt{a^2 + b^2}(a \cos(x) - b \sin(x))}{2ab \cos(x) \sin(x) - (a^2 - b^2) \cos(x)^2 + a^2}\right)}{2\sqrt{a^2 + b^2}}$$

```
input integrate(1/(b*cos(x)+a*sin(x)),x, algorithm="fracas")
```

```
output 1/2*log(-(2*a*b*cos(x)*sin(x) - (a^2 - b^2)*cos(x)^2 - a^2 - 2*b^2 + 2*sqrt
(a^2 + b^2)*(a*cos(x) - b*sin(x)))/(2*a*b*cos(x)*sin(x) - (a^2 - b^2)*cos
(x)^2 + a^2))/sqrt(a^2 + b^2)
```

3.250.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.56 (sec) , antiderivative size = 121, normalized size of antiderivative = 3.36

$$\int \frac{1}{b \cos(x) + a \sin(x)} dx = \begin{cases} \tilde{\infty}(-\log(\tan(\frac{x}{2}) - 1) + \log(\tan(\frac{x}{2}) + 1)) & \text{for } a = 0 \wedge b = 0 \\ \frac{\log(\tan(\frac{x}{2}))}{a} & \text{for } b = 0 \\ -\frac{i}{-ib \sin(x) + b \cos(x)} & \text{for } a = -ib \\ \frac{i}{ib \sin(x) + b \cos(x)} & \text{for } a = ib \\ -\frac{\log\left(-\frac{a}{b} + \tan\left(\frac{x}{2}\right) - \frac{\sqrt{a^2+b^2}}{b}\right)}{\sqrt{a^2+b^2}} + \frac{\log\left(-\frac{a}{b} + \tan\left(\frac{x}{2}\right) + \frac{\sqrt{a^2+b^2}}{b}\right)}{\sqrt{a^2+b^2}} & \text{otherwise} \end{cases}$$

input `integrate(1/(b*cos(x)+a*sin(x)),x)`

output `Piecewise((zoo*(-log(tan(x/2) - 1) + log(tan(x/2) + 1)), Eq(a, 0) & Eq(b, 0)), (log(tan(x/2))/a, Eq(b, 0)), (-I/(-I*b*sin(x) + b*cos(x)), Eq(a, -I*b)), (I/(I*b*sin(x) + b*cos(x)), Eq(a, I*b)), (-log(-a/b + tan(x/2) - sqrt(a**2 + b**2)/b)/sqrt(a**2 + b**2) + log(-a/b + tan(x/2) + sqrt(a**2 + b**2)/b)/sqrt(a**2 + b**2), True))`

3.250.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.69

$$\int \frac{1}{b \cos(x) + a \sin(x)} dx = -\frac{\log\left(\frac{a - \frac{b \sin(x)}{\cos(x)+1} + \sqrt{a^2+b^2}}{a - \frac{b \sin(x)}{\cos(x)+1} - \sqrt{a^2+b^2}}\right)}{\sqrt{a^2+b^2}}$$

input `integrate(1/(b*cos(x)+a*sin(x)),x, algorithm="maxima")`

output `-log((a - b*sin(x)/(cos(x) + 1) + sqrt(a^2 + b^2))/(a - b*sin(x)/(cos(x) + 1) - sqrt(a^2 + b^2)))/sqrt(a^2 + b^2)`

3.250.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.69

$$\int \frac{1}{b \cos(x) + a \sin(x)} dx = -\frac{\log\left(\frac{|2b \tan(\frac{1}{2}x) - 2a - 2\sqrt{a^2 + b^2}|}{|2b \tan(\frac{1}{2}x) - 2a + 2\sqrt{a^2 + b^2}|}\right)}{\sqrt{a^2 + b^2}}$$

input `integrate(1/(b*cos(x)+a*sin(x)),x, algorithm="giac")`output `-log(abs(2*b*tan(1/2*x) - 2*a - 2*sqrt(a^2 + b^2))/abs(2*b*tan(1/2*x) - 2*a + 2*sqrt(a^2 + b^2)))/sqrt(a^2 + b^2)`**3.250.9 Mupad [B] (verification not implemented)**

Time = 0.79 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

$$\int \frac{1}{b \cos(x) + a \sin(x)} dx = -\frac{2 \operatorname{atanh}\left(\frac{a - b \tan(\frac{x}{2})}{\sqrt{a^2 + b^2}}\right)}{\sqrt{a^2 + b^2}}$$

input `int(1/(b*cos(x) + a*sin(x)),x)`output `-(2*atanh((a - b*tan(x/2))/(a^2 + b^2)^(1/2)))/(a^2 + b^2)^(1/2)`

3.251 $\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx$

3.251.1 Optimal result 1299
 3.251.2 Mathematica [A] (verified) 1299
 3.251.3 Rubi [A] (verified) 1300
 3.251.4 Maple [A] (verified) 1301
 3.251.5 Fricas [B] (verification not implemented) 1301
 3.251.6 Sympy [B] (verification not implemented) 1302
 3.251.7 Maxima [A] (verification not implemented) 1302
 3.251.8 Giac [A] (verification not implemented) 1303
 3.251.9 Mupad [B] (verification not implemented) 1303

3.251.1 Optimal result

Integrand size = 19, antiderivative size = 15

$$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx = \frac{\arctan\left(\frac{a \tan(x)}{b}\right)}{ab}$$

output `arctan(a*tan(x)/b)/a/b`

3.251.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx = \frac{\arctan\left(\frac{a \tan(x)}{b}\right)}{ab}$$

input `Integrate[(b^2*Cos[x]^2 + a^2*Sin[x]^2)^(-1),x]`

output `ArcTan[(a*Tan[x])/b]/(a*b)`

3.251.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {3042, 4889, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{a^2 \sin^2(x) + b^2 \cos^2(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{a^2 \sin(x)^2 + b^2 \cos(x)^2} dx \\ & \quad \downarrow \text{4889} \\ & \int \frac{1}{a^2 \tan^2(x) + b^2} d \tan(x) \\ & \quad \downarrow \text{218} \\ & \frac{\arctan\left(\frac{a \tan(x)}{b}\right)}{ab} \end{aligned}$$

input `Int[(b^2*Cos[x]^2 + a^2*Sin[x]^2)^(-1),x]`

output `ArcTan[(a*Tan[x])/b]/(a*b)`

3.251.3.1 Defintions of rubi rules used

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4889 Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, With[{d = FreeFactors
[Tan[v], x]}, Simp[d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x
^2), Tan[v]/d, u, x], x], x, Tan[v]/d], x]] /; !FalseQ[v] && FunctionOfQ[N
onfreeFactors[Tan[v], x], u, x]] /; InverseFunctionFreeQ[u, x] && !MatchQ[
u, (v_.)*((c_.)*tan[w_]^(n_.)*tan[z_]^(n_.))^(p_.) /; FreeQ[{c, p}, x] && I
ntegerQ[n] && LinearQ[w, x] && EqQ[z, 2*w]]
```

3.251.4 Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

method	result	size
default	$\frac{\arctan\left(\frac{a \tan(x)}{b}\right)}{ab}$	16
paralelrisch	$\frac{i \left(\ln\left(\frac{ia \sin(x) - b \cos(x)}{\cos(x)+1}\right) - \ln\left(\frac{-b \cos(x) - ia \sin(x)}{\cos(x)+1}\right) \right)}{2ab}$	53
risch	$\frac{i \ln\left(e^{2ix} - \frac{a+b}{a-b}\right)}{2ab} - \frac{i \ln\left(e^{2ix} - \frac{a-b}{a+b}\right)}{2ab}$	58

input `int(1/(b^2*cos(x)^2+a^2*sin(x)^2),x,method=_RETURNVERBOSE)`

output `arctan(a*tan(x)/b)/a/b`

3.251.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. $2(15) = 30$.

Time = 0.26 (sec) , antiderivative size = 43, normalized size of antiderivative = 2.87

$$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx = -\frac{\arctan\left(\frac{(a^2+b^2) \cos(x)^2 - a^2}{2ab \cos(x) \sin(x)}\right)}{2ab}$$

input `integrate(1/(b^2*cos(x)^2+a^2*sin(x)^2),x, algorithm="fracas")`

output `-1/2*arctan(1/2*((a^2 + b^2)*cos(x)^2 - a^2)/(a*b*cos(x)*sin(x)))/(a*b)`

3.251.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 71839 vs. $2(10) = 20$.

Time = 16.13 (sec) , antiderivative size = 71839, normalized size of antiderivative = 4789.27

$$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx = \text{Too large to display}$$

input `integrate(1/(b**2*cos(x)**2+a**2*sin(x)**2),x)`

output `Piecewise((zoo*tan(x/2)/(tan(x/2)**2 - 1), Eq(a, 0) & Eq(b, 0)), ((tan(x/2)/2 - 1/(2*tan(x/2)))/a**2, Eq(b, 0)), (-2*tan(x/2)/(b**2*(tan(x/2)**2 - 1)), Eq(a, 0)), (x/(b**2*sin(x)**2 + b**2*cos(x)**2), Eq(a, b) | Eq(a, -b)), (8192*a**15*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*log(-sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) + tan(x/2))/(8192*a**15*b**2*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) - 8192*a**14*b**2*sqrt(a**2 - b**2)*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) - 30720*a**13*b**4*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) + 26624*a**12*b**4*sqrt(a**2 - b**2)*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) + 45568*a**11*b**6*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) - 33280*a**10*b**6*sqrt(a**2 - b**2)*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) - 33792*a**9*b**8*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) + 19968*a**8*b**8*sqrt(a**2 - b**2)*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1)*sqrt(-2*a**2/b**2 + 2*a*sqrt(a**2 - b**2)/b**2 + 1) + 12992*a**7*b**10*sqrt(-2*a**2/b**2 - 2*a*sqrt(a**2 - b**2)/b**2 + 1...`

3.251.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx = \frac{\arctan\left(\frac{a \tan(x)}{b}\right)}{ab}$$

input `integrate(1/(b^2*cos(x)^2+a^2*sin(x)^2),x, algorithm="maxima")`

output `arctan(a*tan(x)/b)/(a*b)`

3.251.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.73

$$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx = \frac{\pi \lfloor \frac{x}{\pi} + \frac{1}{2} \rfloor + \arctan\left(\frac{a \tan(x)}{b}\right)}{ab}$$

input `integrate(1/(b^2*cos(x)^2+a^2*sin(x)^2),x, algorithm="giac")`

output `(pi*floor(x/pi + 1/2) + arctan(a*tan(x)/b))/(a*b)`

3.251.9 Mupad [B] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{1}{b^2 \cos^2(x) + a^2 \sin^2(x)} dx = \frac{\operatorname{atan}\left(\frac{a \tan(x)}{b}\right)}{ab}$$

input `int(1/(b^2*cos(x)^2 + a^2*sin(x)^2),x)`

output `atan((a*tan(x))/b)/(a*b)`

3.252 $\int \frac{x}{-1+x^2} dx$

3.252.1 Optimal result	1304
3.252.2 Mathematica [A] (verified)	1304
3.252.3 Rubi [A] (verified)	1305
3.252.4 Maple [A] (verified)	1305
3.252.5 Fricas [A] (verification not implemented)	1306
3.252.6 Sympy [A] (verification not implemented)	1306
3.252.7 Maxima [A] (verification not implemented)	1306
3.252.8 Giac [A] (verification not implemented)	1307
3.252.9 Mupad [B] (verification not implemented)	1307

3.252.1 Optimal result

Integrand size = 9, antiderivative size = 12

$$\int \frac{x}{-1+x^2} dx = \frac{1}{2} \log(1-x^2)$$

output `1/2*ln(-x^2+1)`

3.252.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{x}{-1+x^2} dx = \frac{1}{2} \log(-1+x^2)$$

input `Integrate[x/(-1 + x^2),x]`

output `Log[-1 + x^2]/2`

3.252.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{x^2 - 1} dx$$

↓ 240

$$\frac{1}{2} \log(1 - x^2)$$

input `Int[x/(-1 + x^2), x]`

output `Log[1 - x^2]/2`

3.252.3.1 Defintions of rubi rules used

rule 240 `Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`

3.252.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

method	result	size
derivativedivides	$\frac{\ln(x^2-1)}{2}$	9
risch	$\frac{\ln(x^2-1)}{2}$	9
meijerg	$\frac{\ln(-x^2+1)}{2}$	11
default	$\frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$	14
norman	$\frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$	14
parallelrisc	$\frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$	14

input `int(x/(x^2-1),x,method=_RETURNVERBOSE)`

output `1/2*ln(x^2-1)`

3.252.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x}{-1+x^2} dx = \frac{1}{2} \log(x^2 - 1)$$

input `integrate(x/(x^2-1),x, algorithm="fricas")`

output `1/2*log(x^2 - 1)`

3.252.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.58

$$\int \frac{x}{-1+x^2} dx = \frac{\log(x^2 - 1)}{2}$$

input `integrate(x/(x**2-1),x)`

output `log(x**2 - 1)/2`

3.252.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x}{-1+x^2} dx = \frac{1}{2} \log(x^2 - 1)$$

input `integrate(x/(x^2-1),x, algorithm="maxima")`

output `1/2*log(x^2 - 1)`

3.252.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{x}{-1+x^2} dx = \frac{1}{2} \log(|x^2-1|)$$

input `integrate(x/(x^2-1),x, algorithm="giac")`output `1/2*log(abs(x^2 - 1))`**3.252.9 Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x}{-1+x^2} dx = \frac{\ln(x^2-1)}{2}$$

input `int(x/(x^2 - 1),x)`output `log(x^2 - 1)/2`

3.253 $\int (1 + \sqrt{x}) \sqrt{x} dx$

3.253.1 Optimal result	1308
3.253.2 Mathematica [A] (verified)	1308
3.253.3 Rubi [A] (verified)	1309
3.253.4 Maple [A] (verified)	1310
3.253.5 Fricas [A] (verification not implemented)	1310
3.253.6 Sympy [A] (verification not implemented)	1310
3.253.7 Maxima [B] (verification not implemented)	1311
3.253.8 Giac [A] (verification not implemented)	1311
3.253.9 Mupad [B] (verification not implemented)	1311

3.253.1 Optimal result

Integrand size = 13, antiderivative size = 17

$$\int (1 + \sqrt{x}) \sqrt{x} dx = \frac{2x^{3/2}}{3} + \frac{x^2}{2}$$

output `2/3*x^(3/2)+1/2*x^2`

3.253.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int (1 + \sqrt{x}) \sqrt{x} dx = \frac{2x^{3/2}}{3} + \frac{x^2}{2}$$

input `Integrate[(1 + Sqrt[x])*Sqrt[x], x]`

output `(2*x^(3/2))/3 + x^2/2`

3.253.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {802, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\sqrt{x} + 1) \sqrt{x} dx$$

$$\downarrow 802$$

$$\int (x + \sqrt{x}) dx$$

$$\downarrow 2009$$

$$\frac{2x^{3/2}}{3} + \frac{x^2}{2}$$

input `Int[(1 + Sqrt[x])*Sqrt[x],x]`

output `(2*x^(3/2))/3 + x^2/2`

3.253.3.1 Defintions of rubi rules used

rule 802 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.253.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

method	result	size
derivativedivides	$\frac{2x^{\frac{3}{2}}}{3} + \frac{x^2}{2}$	12
default	$\frac{2x^{\frac{3}{2}}}{3} + \frac{x^2}{2}$	12
trager	$\frac{(-1+x)(1+x)}{2} + \frac{2x^{\frac{3}{2}}}{3}$	15

input `int(x^(1/2)*(x^(1/2)+1),x,method=_RETURNVERBOSE)`output `2/3*x^(3/2)+1/2*x^2`**3.253.5 Fricas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int (1 + \sqrt{x}) \sqrt{x} dx = \frac{1}{2} x^2 + \frac{2}{3} x^{\frac{3}{2}}$$

input `integrate(x^(1/2)*(1+x^(1/2)),x, algorithm="fricas")`output `1/2*x^2 + 2/3*x^(3/2)`**3.253.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int (1 + \sqrt{x}) \sqrt{x} dx = \frac{2x^{\frac{3}{2}}}{3} + \frac{x^2}{2}$$

input `integrate(x**(1/2)*(1+x**(1/2)),x)`output `2*x**(3/2)/3 + x**2/2`

3.253.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. $2(11) = 22$.

Time = 0.20 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.53

$$\int (1 + \sqrt{x}) \sqrt{x} dx = \frac{1}{2} (\sqrt{x} + 1)^4 - \frac{4}{3} (\sqrt{x} + 1)^3 + (\sqrt{x} + 1)^2$$

input `integrate(x^(1/2)*(1+x^(1/2)),x, algorithm="maxima")`

output `1/2*(sqrt(x) + 1)^4 - 4/3*(sqrt(x) + 1)^3 + (sqrt(x) + 1)^2`

3.253.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int (1 + \sqrt{x}) \sqrt{x} dx = \frac{1}{2} x^2 + \frac{2}{3} x^{3/2}$$

input `integrate(x^(1/2)*(1+x^(1/2)),x, algorithm="giac")`

output `1/2*x^2 + 2/3*x^(3/2)`

3.253.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int (1 + \sqrt{x}) \sqrt{x} dx = \frac{x^2}{2} + \frac{2x^{3/2}}{3}$$

input `int(x^(1/2)*(x^(1/2) + 1),x)`

output `x^2/2 + (2*x^(3/2))/3`

3.254 $\int \frac{1}{1-\cos(x)} dx$

3.254.1 Optimal result	1312
3.254.2 Mathematica [A] (verified)	1312
3.254.3 Rubi [A] (verified)	1313
3.254.4 Maple [A] (verified)	1314
3.254.5 Fricas [A] (verification not implemented)	1314
3.254.6 Sympy [A] (verification not implemented)	1314
3.254.7 Maxima [A] (verification not implemented)	1315
3.254.8 Giac [A] (verification not implemented)	1315
3.254.9 Mupad [B] (verification not implemented)	1315

3.254.1 Optimal result

Integrand size = 8, antiderivative size = 12

$$\int \frac{1}{1-\cos(x)} dx = -\frac{\sin(x)}{1-\cos(x)}$$

output `-sin(x)/(1-cos(x))`

3.254.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{1-\cos(x)} dx = -\cot\left(\frac{x}{2}\right)$$

input `Integrate[(1 - Cos[x])^(-1), x]`

output `-Cot[x/2]`

3.254.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{1 - \cos(x)} dx$$

↓ 3042

$$\int \frac{1}{1 - \sin\left(x + \frac{\pi}{2}\right)} dx$$

↓ 3127

$$-\frac{\sin(x)}{1 - \cos(x)}$$

input `Int[(1 - Cos[x])^(-1),x]`

output `-(Sin[x]/(1 - Cos[x]))`

3.254.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

3.254.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

method	result	size
default	$-\frac{1}{\tan(\frac{x}{2})}$	9
norman	$-\frac{1}{\tan(\frac{x}{2})}$	9
parallelrisch	$-\frac{1}{\tan(\frac{x}{2})}$	9
risch	$-\frac{2i}{e^{ix}-1}$	13

input `int(1/(1-cos(x)),x,method=_RETURNVERBOSE)`output `-1/tan(1/2*x)`**3.254.5 Fracas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1}{1 - \cos(x)} dx = -\frac{\cos(x) + 1}{\sin(x)}$$

input `integrate(1/(1-cos(x)),x, algorithm="fricas")`output `-(cos(x) + 1)/sin(x)`**3.254.6 Sympy [A] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.58

$$\int \frac{1}{1 - \cos(x)} dx = -\frac{1}{\tan(\frac{x}{2})}$$

input `integrate(1/(1-cos(x)),x)`output `-1/tan(x/2)`

3.254.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1}{1 - \cos(x)} dx = -\frac{\cos(x) + 1}{\sin(x)}$$

input `integrate(1/(1-cos(x)),x, algorithm="maxima")`output `-(cos(x) + 1)/sin(x)`**3.254.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{1 - \cos(x)} dx = -\frac{1}{\tan\left(\frac{1}{2}x\right)}$$

input `integrate(1/(1-cos(x)),x, algorithm="giac")`output `-1/tan(1/2*x)`**3.254.9 Mupad [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.50

$$\int \frac{1}{1 - \cos(x)} dx = -\cot\left(\frac{x}{2}\right)$$

input `int(-1/(cos(x) - 1),x)`output `-cot(x/2)`

3.255 $\int \sec(x) \tan^2(x) dx$

3.255.1 Optimal result	1316
3.255.2 Mathematica [A] (verified)	1316
3.255.3 Rubi [A] (verified)	1317
3.255.4 Maple [A] (verified)	1318
3.255.5 Fricas [B] (verification not implemented)	1318
3.255.6 Sympy [A] (verification not implemented)	1319
3.255.7 Maxima [B] (verification not implemented)	1319
3.255.8 Giac [B] (verification not implemented)	1319
3.255.9 Mupad [B] (verification not implemented)	1320

3.255.1 Optimal result

Integrand size = 7, antiderivative size = 16

$$\int \sec(x) \tan^2(x) dx = -\frac{1}{2} \operatorname{arctanh}(\sin(x)) + \frac{1}{2} \sec(x) \tan(x)$$

output `-1/2*arctanh(sin(x))+1/2*sec(x)*tan(x)`

3.255.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \sec(x) \tan^2(x) dx = -\frac{1}{2} \operatorname{arctanh}(\sin(x)) + \frac{1}{2} \sec(x) \tan(x)$$

input `Integrate[Sec[x]*Tan[x]^2,x]`

output `-1/2*ArcTanh[Sin[x]] + (Sec[x]*Tan[x])/2`

3.255.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 3091, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^2(x) \sec(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^2 \sec(x) dx \\
 & \quad \downarrow \text{3091} \\
 & \frac{1}{2} \tan(x) \sec(x) - \frac{\int \sec(x) dx}{2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \tan(x) \sec(x) - \frac{1}{2} \int \csc\left(x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{4257} \\
 & \frac{1}{2} \tan(x) \sec(x) - \frac{1}{2} \operatorname{arctanh}(\sin(x))
 \end{aligned}$$

input `Int[Sec[x]*Tan[x]^2,x]`

output `-1/2*ArcTanh[Sin[x]] + (Sec[x]*Tan[x])/2`

3.255.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

```
rule 3091 Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(
n_), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m
+ n - 1))), x] - Simp[b^2*((n - 1)/(m + n - 1)) Int[(a*Sec[e + f*x])^m*(
b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] &
& NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.255.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.50

method	result	size
default	$\frac{\sin^3(x)}{2 \cos(x)^2} + \frac{\sin(x)}{2} - \frac{\ln(\sec(x)+\tan(x))}{2}$	24
risch	$-\frac{i(e^{3ix}-e^{ix})}{(e^{2ix}+1)^2} - \frac{\ln(i+e^{ix})}{2} + \frac{\ln(e^{ix}-i)}{2}$	49

```
input int(sec(x)*tan(x)^2,x,method=_RETURNVERBOSE)
```

```
output 1/2*sin(x)^3/cos(x)^2+1/2*sin(x)-1/2*ln(sec(x)+tan(x))
```

3.255.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. $2(12) = 24$.

Time = 0.25 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.12

$$\int \sec(x) \tan^2(x) dx = -\frac{\cos(x)^2 \log(\sin(x) + 1) - \cos(x)^2 \log(-\sin(x) + 1) - 2 \sin(x)}{4 \cos(x)^2}$$

```
input integrate(sec(x)*tan(x)^2,x, algorithm="fracas")
```

```
output -1/4*(cos(x)^2*log(sin(x) + 1) - cos(x)^2*log(-sin(x) + 1) - 2*sin(x))/cos
(x)^2
```

3.255.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.69

$$\int \sec(x) \tan^2(x) dx = \frac{\log(\sin(x) - 1)}{4} - \frac{\log(\sin(x) + 1)}{4} - \frac{\sin(x)}{2 \sin^2(x) - 2}$$

input `integrate(sec(x)*tan(x)**2,x)`

output `log(sin(x) - 1)/4 - log(sin(x) + 1)/4 - sin(x)/(2*sin(x)**2 - 2)`

3.255.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(12) = 24.

Time = 0.19 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.69

$$\int \sec(x) \tan^2(x) dx = -\frac{\sin(x)}{2(\sin(x)^2 - 1)} - \frac{1}{4} \log(\sin(x) + 1) + \frac{1}{4} \log(\sin(x) - 1)$$

input `integrate(sec(x)*tan(x)^2,x, algorithm="maxima")`

output `-1/2*sin(x)/(sin(x)^2 - 1) - 1/4*log(sin(x) + 1) + 1/4*log(sin(x) - 1)`

3.255.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. 2(12) = 24.

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.81

$$\int \sec(x) \tan^2(x) dx = -\frac{\sin(x)}{2(\sin(x)^2 - 1)} - \frac{1}{4} \log(\sin(x) + 1) + \frac{1}{4} \log(-\sin(x) + 1)$$

input `integrate(sec(x)*tan(x)^2,x, algorithm="giac")`

output `-1/2*sin(x)/(sin(x)^2 - 1) - 1/4*log(sin(x) + 1) + 1/4*log(-sin(x) + 1)`

3.255.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.88

$$\int \sec(x) \tan^2(x) dx = \frac{\tan\left(\frac{x}{2}\right)^3 + \tan\left(\frac{x}{2}\right)}{\left(\tan\left(\frac{x}{2}\right)^2 - 1\right)^2} - \operatorname{atanh}\left(\tan\left(\frac{x}{2}\right)\right)$$

input `int(tan(x)^2/cos(x),x)`

output `(tan(x/2) + tan(x/2)^3)/(tan(x/2)^2 - 1)^2 - atanh(tan(x/2))`

3.256 $\int \sec^3(x) \tan^3(x) dx$

3.256.1 Optimal result1321
3.256.2 Mathematica [A] (verified)1321
3.256.3 Rubi [A] (verified)1322
3.256.4 Maple [A] (verified)1323
3.256.5 Fricas [A] (verification not implemented)1324
3.256.6 Sympy [A] (verification not implemented)1324
3.256.7 Maxima [A] (verification not implemented)1324
3.256.8 Giac [A] (verification not implemented)1325
3.256.9 Mupad [B] (verification not implemented)1325

3.256.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \sec^3(x) \tan^3(x) dx = -\frac{1}{3} \sec^3(x) + \frac{\sec^5(x)}{5}$$

output `-1/3*sec(x)^3+1/5*sec(x)^5`

3.256.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sec^3(x) \tan^3(x) dx = -\frac{1}{3} \sec^3(x) + \frac{\sec^5(x)}{5}$$

input `Integrate[Sec[x]^3*Tan[x]^3,x]`

output `-1/3*Sec[x]^3 + Sec[x]^5/5`

3.256.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 3086, 25, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^3(x) \sec^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^3 \sec(x)^3 dx \\
 & \quad \downarrow \text{3086} \\
 & \int -\sec^2(x) (1 - \sec^2(x)) d\sec(x) \\
 & \quad \downarrow \text{25} \\
 & - \int \sec^2(x) (1 - \sec^2(x)) d\sec(x) \\
 & \quad \downarrow \text{244} \\
 & - \int (\sec^2(x) - \sec^4(x)) d\sec(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sec^5(x)}{5} - \frac{\sec^3(x)}{3}
 \end{aligned}$$

input `Int[Sec[x]^3*Tan[x]^3,x]`

output `-1/3*Sec[x]^3 + Sec[x]^5/5`

3.256.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

3.256.4 Maple [A] (verified)

Time = 1.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$-\frac{(\sec^3(x))}{3} + \frac{(\sec^5(x))}{5}$	14
default	$-\frac{(\sec^3(x))}{3} + \frac{(\sec^5(x))}{5}$	14
risch	$-\frac{8(5e^{7ix} - 2e^{5ix} + 5e^{3ix})}{15(e^{2ix} + 1)^5}$	34

input `int(sec(x)^3*tan(x)^3,x,method=_RETURNVERBOSE)`

output `-1/3*sec(x)^3+1/5*sec(x)^5`

3.256.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^3(x) \tan^3(x) dx = -\frac{5 \cos(x)^2 - 3}{15 \cos(x)^5}$$

input `integrate(sec(x)^3*tan(x)^3,x, algorithm="fricas")`output `-1/15*(5*cos(x)^2 - 3)/cos(x)^5`**3.256.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^3(x) \tan^3(x) dx = \frac{3 - 5 \cos^2(x)}{15 \cos^5(x)}$$

input `integrate(sec(x)**3*tan(x)**3,x)`output `(3 - 5*cos(x)**2)/(15*cos(x)**5)`**3.256.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^3(x) \tan^3(x) dx = -\frac{5 \cos(x)^2 - 3}{15 \cos(x)^5}$$

input `integrate(sec(x)^3*tan(x)^3,x, algorithm="maxima")`output `-1/15*(5*cos(x)^2 - 3)/cos(x)^5`

3.256.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^3(x) \tan^3(x) dx = -\frac{5 \cos(x)^2 - 3}{15 \cos(x)^5}$$

input `integrate(sec(x)^3*tan(x)^3,x, algorithm="giac")`output `-1/15*(5*cos(x)^2 - 3)/cos(x)^5`**3.256.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sec^3(x) \tan^3(x) dx = \frac{1}{5 \cos(x)^5} - \frac{1}{3 \cos(x)^3}$$

input `int(tan(x)^3/cos(x)^3,x)`output `1/(5*cos(x)^5) - 1/(3*cos(x)^3)`

3.257 $\int e^{\sqrt{x}} dx$

3.257.1 Optimal result	1326
3.257.2 Mathematica [A] (verified)	1326
3.257.3 Rubi [A] (verified)	1327
3.257.4 Maple [A] (verified)	1328
3.257.5 Fricas [A] (verification not implemented)	1328
3.257.6 Sympy [A] (verification not implemented)	1329
3.257.7 Maxima [A] (verification not implemented)	1329
3.257.8 Giac [A] (verification not implemented)	1329
3.257.9 Mupad [B] (verification not implemented)	1330

3.257.1 Optimal result

Integrand size = 7, antiderivative size = 24

$$\int e^{\sqrt{x}} dx = -2e^{\sqrt{x}} + 2e^{\sqrt{x}}\sqrt{x}$$

output `-2*exp(x^(1/2))+2*exp(x^(1/2))*x^(1/2)`

3.257.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int e^{\sqrt{x}} dx = 2e^{\sqrt{x}}(-1 + \sqrt{x})$$

input `Integrate[E^Sqrt[x],x]`

output `2*E^Sqrt[x]*(-1 + Sqrt[x])`

3.257.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.04, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {2636, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int e^{\sqrt{x}} dx \\
 \downarrow \text{2636} \\
 2 \int e^{\sqrt{x}} \sqrt{x} d\sqrt{x} \\
 \downarrow \text{2607} \\
 2 \left(e^{\sqrt{x}} \sqrt{x} - \int e^{\sqrt{x}} d\sqrt{x} \right) \\
 \downarrow \text{2624} \\
 2 \left(e^{\sqrt{x}} \sqrt{x} - e^{\sqrt{x}} \right)
 \end{array}$$

input `Int[E^Sqrt[x], x]`

output `2*(-E^Sqrt[x] + E^Sqrt[x]*Sqrt[x])`

3.257.3.1 Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`


```
rule 2636 Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{k =
Denominator[n]}, Simp[k/d Subst[Int[x^(k - 1)*F^(a + b*x^(k*n)), x], x, (
c + d*x)^(1/k)], x]] /; FreeQ[{F, a, b, c, d}, x] && IntegerQ[2/n] && !Int
egerQ[n]
```

3.257.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

method	result	size
meijerg	$2 - (-2\sqrt{x} + 2) e^{\sqrt{x}}$	16
derivativedivides	$-2e^{\sqrt{x}} + 2e^{\sqrt{x}}\sqrt{x}$	17
default	$-2e^{\sqrt{x}} + 2e^{\sqrt{x}}\sqrt{x}$	17

```
input int(exp(x^(1/2)),x,method=_RETURNVERBOSE)
```

```
output 2-(-2*x^(1/2)+2)*exp(x^(1/2))
```

3.257.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46

$$\int e^{\sqrt{x}} dx = 2(\sqrt{x} - 1)e^{\sqrt{x}}$$

```
input integrate(exp(x^(1/2)),x, algorithm="fricas")
```

```
output 2*(sqrt(x) - 1)*e^sqrt(x)
```

3.257.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int e^{\sqrt{x}} dx = 2\sqrt{x}e^{\sqrt{x}} - 2e^{\sqrt{x}}$$

input `integrate(exp(x**(1/2)),x)`output `2*sqrt(x)*exp(sqrt(x)) - 2*exp(sqrt(x))`**3.257.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46

$$\int e^{\sqrt{x}} dx = 2(\sqrt{x} - 1)e^{\sqrt{x}}$$

input `integrate(exp(x^(1/2)),x, algorithm="maxima")`output `2*(sqrt(x) - 1)*e^sqrt(x)`**3.257.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46

$$\int e^{\sqrt{x}} dx = 2(\sqrt{x} - 1)e^{\sqrt{x}}$$

input `integrate(exp(x^(1/2)),x, algorithm="giac")`output `2*(sqrt(x) - 1)*e^sqrt(x)`

3.257.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46

$$\int e^{\sqrt{x}} dx = 2e^{\sqrt{x}}(\sqrt{x} - 1)$$

input `int(exp(x^(1/2)),x)`

output `2*exp(x^(1/2))*(x^(1/2) - 1)`

3.258 $\int \frac{1+x^5}{-10x-3x^2+x^3} dx$

3.258.1 Optimal result 1331
 3.258.2 Mathematica [A] (verified) 1331
 3.258.3 Rubi [A] (verified) 1332
 3.258.4 Maple [A] (verified) 1333
 3.258.5 Fricas [A] (verification not implemented) 1333
 3.258.6 Sympy [A] (verification not implemented) 1334
 3.258.7 Maxima [A] (verification not implemented) 1334
 3.258.8 Giac [A] (verification not implemented) 1334
 3.258.9 Mupad [B] (verification not implemented) 1335

3.258.1 Optimal result

Integrand size = 20, antiderivative size = 42

$$\int \frac{1+x^5}{-10x-3x^2+x^3} dx = 19x + \frac{3x^2}{2} + \frac{x^3}{3} + \frac{3126}{35} \log(5-x) - \frac{\log(x)}{10} - \frac{31}{14} \log(2+x)$$

output `19*x+3/2*x^2+1/3*x^3+3126/35*ln(5-x)-1/10*ln(x)-31/14*ln(2+x)`

3.258.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int \frac{1+x^5}{-10x-3x^2+x^3} dx = 19x + \frac{3x^2}{2} + \frac{x^3}{3} + \frac{3126}{35} \log(5-x) - \frac{\log(x)}{10} - \frac{31}{14} \log(2+x)$$

input `Integrate[(1 + x^5)/(-10*x - 3*x^2 + x^3),x]`

output `19*x + (3*x^2)/2 + x^3/3 + (3126*Log[5 - x])/35 - Log[x]/10 - (31*Log[2 + x])/14`

3.258.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {2026, 2159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^5 + 1}{x^3 - 3x^2 - 10x} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{x^5 + 1}{x(x^2 - 3x - 10)} dx \\ & \quad \downarrow \text{2159} \\ & \int \left(x^2 + 3x + \frac{3126}{35(x-5)} - \frac{31}{14(x+2)} - \frac{1}{10x} + 19 \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{x^3}{3} + \frac{3x^2}{2} + 19x + \frac{3126}{35} \log(5-x) - \frac{\log(x)}{10} - \frac{31}{14} \log(x+2) \end{aligned}$$

input `Int[(1 + x^5)/(-10*x - 3*x^2 + x^3), x]`

output `19*x + (3*x^2)/2 + x^3/3 + (3126*Log[5 - x])/35 - Log[x]/10 - (31*Log[2 + x])/14`

3.258.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

```
rule 2159 Int[(Pq_)*((d_.) + (e_.)*(x_))^(m_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p
_.), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^(m)*Pq*(a + b*x + c*x^2)^p, x
], x] /; FreeQ[{a, b, c, d, e, m}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

3.258.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.74

method	result	size
default	$\frac{x^3}{3} + \frac{3x^2}{2} + 19x - \frac{31 \ln(2+x)}{14} + \frac{3126 \ln(x-5)}{35} - \frac{\ln(x)}{10}$	31
norman	$\frac{x^3}{3} + \frac{3x^2}{2} + 19x - \frac{31 \ln(2+x)}{14} + \frac{3126 \ln(x-5)}{35} - \frac{\ln(x)}{10}$	31
risch	$\frac{x^3}{3} + \frac{3x^2}{2} + 19x - \frac{31 \ln(2+x)}{14} + \frac{3126 \ln(x-5)}{35} - \frac{\ln(x)}{10}$	31
parallelrisch	$\frac{x^3}{3} + \frac{3x^2}{2} + 19x - \frac{31 \ln(2+x)}{14} + \frac{3126 \ln(x-5)}{35} - \frac{\ln(x)}{10}$	31

```
input int((x^5+1)/(x^3-3*x^2-10*x),x,method=_RETURNVERBOSE)
```

```
output 1/3*x^3+3/2*x^2+19*x-31/14*ln(2+x)+3126/35*ln(x-5)-1/10*ln(x)
```

3.258.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.71

$$\int \frac{1+x^5}{-10x-3x^2+x^3} dx = \frac{1}{3}x^3 + \frac{3}{2}x^2 + 19x - \frac{31}{14} \log(x+2) + \frac{3126}{35} \log(x-5) - \frac{1}{10} \log(x)$$

```
input integrate((x^5+1)/(x^3-3*x^2-10*x),x, algorithm="fracas")
```

```
output 1/3*x^3 + 3/2*x^2 + 19*x - 31/14*log(x + 2) + 3126/35*log(x - 5) - 1/10*lo
g(x)
```

3.258.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

$$\int \frac{1+x^5}{-10x-3x^2+x^3} dx = \frac{x^3}{3} + \frac{3x^2}{2} + 19x - \frac{\log(x)}{10} + \frac{3126 \log(x-5)}{35} - \frac{31 \log(x+2)}{14}$$

input `integrate((x**5+1)/(x**3-3*x**2-10*x),x)`output `x**3/3 + 3*x**2/2 + 19*x - log(x)/10 + 3126*log(x - 5)/35 - 31*log(x + 2)/14`**3.258.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.71

$$\int \frac{1+x^5}{-10x-3x^2+x^3} dx = \frac{1}{3}x^3 + \frac{3}{2}x^2 + 19x - \frac{31}{14} \log(x+2) + \frac{3126}{35} \log(x-5) - \frac{1}{10} \log(x)$$

input `integrate((x^5+1)/(x^3-3*x^2-10*x),x, algorithm="maxima")`output `1/3*x^3 + 3/2*x^2 + 19*x - 31/14*log(x + 2) + 3126/35*log(x - 5) - 1/10*log(x)`**3.258.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.79

$$\int \frac{1+x^5}{-10x-3x^2+x^3} dx = \frac{1}{3}x^3 + \frac{3}{2}x^2 + 19x - \frac{31}{14} \log(|x+2|) + \frac{3126}{35} \log(|x-5|) - \frac{1}{10} \log(|x|)$$

input `integrate((x^5+1)/(x^3-3*x^2-10*x),x, algorithm="giac")`output `1/3*x^3 + 3/2*x^2 + 19*x - 31/14*log(abs(x + 2)) + 3126/35*log(abs(x - 5)) - 1/10*log(abs(x))`

3.258.9 Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.71

$$\int \frac{1+x^5}{-10x-3x^2+x^3} dx = 19x - \frac{31 \ln(x+2)}{14} + \frac{3126 \ln(x-5)}{35} - \frac{\ln(x)}{10} + \frac{3x^2}{2} + \frac{x^3}{3}$$

input `int(-(x^5 + 1)/(10*x + 3*x^2 - x^3),x)`

output `19*x - (31*log(x + 2))/14 + (3126*log(x - 5))/35 - log(x)/10 + (3*x^2)/2 + x^3/3`

$$\mathbf{3.259} \quad \int \frac{1}{x\sqrt{\log(x)}} dx$$

3.259.1 Optimal result	1336
3.259.2 Mathematica [A] (verified)	1336
3.259.3 Rubi [A] (verified)	1337
3.259.4 Maple [A] (verified)	1338
3.259.5 Fricas [A] (verification not implemented)	1338
3.259.6 Sympy [A] (verification not implemented)	1338
3.259.7 Maxima [A] (verification not implemented)	1339
3.259.8 Giac [A] (verification not implemented)	1339
3.259.9 Mupad [B] (verification not implemented)	1339

3.259.1 Optimal result

Integrand size = 10, antiderivative size = 8

$$\int \frac{1}{x\sqrt{\log(x)}} dx = 2\sqrt{\log(x)}$$

output `2*ln(x)^(1/2)`

3.259.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{\log(x)}} dx = 2\sqrt{\log(x)}$$

input `Integrate[1/(x*Sqrt[Log[x]]),x]`

output `2*Sqrt[Log[x]]`

3.259.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2739, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{1}{x\sqrt{\log(x)}} dx \\ \downarrow 2739 \\ \int \frac{1}{\sqrt{\log(x)}} d\log(x) \\ \downarrow 15 \\ 2\sqrt{\log(x)} \end{array}$$

input `Int [1/(x*Sqrt [Log [x]]), x]`

output `2*Sqrt [Log [x]]`

3.259.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2739 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/(x_), x_Symbol] := Simp[1/(b*n) Subst[Int[x^p, x], x, a + b*Log[c*x^n]], x] /; FreeQ[{a, b, c, n, p}, x]`

3.259.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$2\sqrt{\ln(x)}$	7
default	$2\sqrt{\ln(x)}$	7

input `int(1/x/ln(x)^(1/2),x,method=_RETURNVERBOSE)`output `2*ln(x)^(1/2)`**3.259.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{x\sqrt{\log(x)}} dx = 2\sqrt{\log(x)}$$

input `integrate(1/x/log(x)^(1/2),x, algorithm="fracas")`output `2*sqrt(log(x))`**3.259.6 Sympy [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \frac{1}{x\sqrt{\log(x)}} dx = 2\sqrt{\log(x)}$$

input `integrate(1/x/ln(x)**(1/2),x)`output `2*sqrt(log(x))`

3.259.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{x\sqrt{\log(x)}} dx = 2\sqrt{\log(x)}$$

input `integrate(1/x/log(x)^(1/2),x, algorithm="maxima")`output `2*sqrt(log(x))`**3.259.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{x\sqrt{\log(x)}} dx = 2\sqrt{\log(x)}$$

input `integrate(1/x/log(x)^(1/2),x, algorithm="giac")`output `2*sqrt(log(x))`**3.259.9 Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{x\sqrt{\log(x)}} dx = 2\sqrt{\ln(x)}$$

input `int(1/(x*log(x)^(1/2)),x)`output `2*log(x)^(1/2)`

3.260 $\int \frac{5+2x}{-3+x} dx$

3.260.1 Optimal result	1340
3.260.2 Mathematica [A] (verified)	1340
3.260.3 Rubi [A] (verified)	1341
3.260.4 Maple [A] (verified)	1342
3.260.5 Fricas [A] (verification not implemented)	1342
3.260.6 Sympy [A] (verification not implemented)	1342
3.260.7 Maxima [A] (verification not implemented)	1343
3.260.8 Giac [A] (verification not implemented)	1343
3.260.9 Mupad [B] (verification not implemented)	1343

3.260.1 Optimal result

Integrand size = 11, antiderivative size = 12

$$\int \frac{5+2x}{-3+x} dx = 2x + 11 \log(3-x)$$

output `2*x+11*ln(3-x)`

3.260.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{5+2x}{-3+x} dx = 2(-3+x) + 11 \log(-3+x)$$

input `Integrate[(5 + 2*x)/(-3 + x),x]`

output `2*(-3 + x) + 11*Log[-3 + x]`

3.260.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x + 5}{x - 3} dx$$

$$\downarrow 49$$

$$\int \left(\frac{11}{x - 3} + 2 \right) dx$$

$$\downarrow 2009$$

$$2x + 11 \log(3 - x)$$

input `Int[(5 + 2*x)/(-3 + x),x]`

output `2*x + 11*Log[3 - x]`

3.260.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.260.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
default	$2x + 11 \ln(-3 + x)$	11
norman	$2x + 11 \ln(-3 + x)$	11
risch	$2x + 11 \ln(-3 + x)$	11
parallelrisc	$2x + 11 \ln(-3 + x)$	11
meijerg	$11 \ln\left(1 - \frac{x}{3}\right) + 2x$	13

input `int((5+2*x)/(-3+x),x,method=_RETURNVERBOSE)`output `2*x+11*ln(-3+x)`**3.260.5 Fricas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{5 + 2x}{-3 + x} dx = 2x + 11 \log(x - 3)$$

input `integrate((5+2*x)/(-3+x),x, algorithm="fricas")`output `2*x + 11*log(x - 3)`**3.260.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{5 + 2x}{-3 + x} dx = 2x + 11 \log(x - 3)$$

input `integrate((5+2*x)/(-3+x),x)`output `2*x + 11*log(x - 3)`

3.260.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{5 + 2x}{-3 + x} dx = 2x + 11 \log(x - 3)$$

input `integrate((5+2*x)/(-3+x),x, algorithm="maxima")`output `2*x + 11*log(x - 3)`**3.260.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

$$\int \frac{5 + 2x}{-3 + x} dx = 2x + 11 \log(|x - 3|)$$

input `integrate((5+2*x)/(-3+x),x, algorithm="giac")`output `2*x + 11*log(abs(x - 3))`**3.260.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{5 + 2x}{-3 + x} dx = 2x + 11 \ln(x - 3)$$

input `int((2*x + 5)/(x - 3),x)`output `2*x + 11*log(x - 3)`

3.261 $\int e^{e^x+x} dx$

3.261.1 Optimal result	1344
3.261.2 Mathematica [A] (verified)	1344
3.261.3 Rubi [A] (verified)	1345
3.261.4 Maple [A] (verified)	1346
3.261.5 Fricas [A] (verification not implemented)	1346
3.261.6 Sympy [A] (verification not implemented)	1346
3.261.7 Maxima [A] (verification not implemented)	1347
3.261.8 Giac [A] (verification not implemented)	1347
3.261.9 Mupad [B] (verification not implemented)	1347

3.261.1 Optimal result

Integrand size = 7, antiderivative size = 5

$$\int e^{e^x+x} dx = e^{e^x}$$

output `exp(exp(x))`

3.261.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int e^{e^x+x} dx = e^{e^x}$$

input `Integrate[E^(E^x + x), x]`

output `E^E^x`

3.261.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2720, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{x+e^x} dx$$

$$\downarrow 2720$$

$$\int e^{e^x} de^x$$

$$\downarrow 2624$$

$$e^{e^x}$$

input `Int[E^(E^x + x), x]`

output `E^E^x`

3.261.3.1 Defintions of rubi rules used

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;`
`FreeQ[{F, n}, x] && LinearQ[v, x]`

rule 2720 `Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]`
`Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /;` `FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /;` `FreeQ`
`[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))`
`*(F_)[v_] /;` `FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

3.261.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.80

method	result	size
default	e^{e^x}	4
risch	e^{e^x}	4

input `int(exp(exp(x)+x),x,method=_RETURNVERBOSE)`output `exp(exp(x))`**3.261.5 Fricas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.60

$$\int e^{e^x+x} dx = e^{(e^x)}$$

input `integrate(exp(exp(x)+x),x, algorithm="fricas")`output `e^(e^x)`**3.261.6 Sympy [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.60

$$\int e^{e^x+x} dx = e^{e^x}$$

input `integrate(exp(exp(x)+x),x)`output `exp(exp(x))`

3.261.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.60

$$\int e^{e^x+x} dx = e^{(e^x)}$$

input `integrate(exp(exp(x)+x),x, algorithm="maxima")`output `e^(e^x)`**3.261.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.60

$$\int e^{e^x+x} dx = e^{(e^x)}$$

input `integrate(exp(exp(x)+x),x, algorithm="giac")`output `e^(e^x)`**3.261.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.60

$$\int e^{e^x+x} dx = e^{e^x}$$

input `int(exp(x + exp(x)),x)`output `exp(exp(x))`

3.262 $\int \cos^2(x) \sin^2(x) dx$

3.262.1 Optimal result	1348
3.262.2 Mathematica [A] (verified)	1348
3.262.3 Rubi [A] (verified)	1349
3.262.4 Maple [A] (verified)	1350
3.262.5 Fricas [A] (verification not implemented)	1351
3.262.6 Sympy [A] (verification not implemented)	1351
3.262.7 Maxima [A] (verification not implemented)	1351
3.262.8 Giac [A] (verification not implemented)	1352
3.262.9 Mupad [B] (verification not implemented)	1352

3.262.1 Optimal result

Integrand size = 9, antiderivative size = 24

$$\int \cos^2(x) \sin^2(x) dx = \frac{x}{8} + \frac{1}{8} \cos(x) \sin(x) - \frac{1}{4} \cos^3(x) \sin(x)$$

output `1/8*x+1/8*cos(x)*sin(x)-1/4*cos(x)^3*sin(x)`

3.262.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.58

$$\int \cos^2(x) \sin^2(x) dx = \frac{x}{8} - \frac{1}{32} \sin(4x)$$

input `Integrate[Cos[x]^2*Sin[x]^2,x]`

output `x/8 - Sin[4*x]/32`

3.262.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 3048, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^2(x) \cos^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^2 \cos(x)^2 dx \\
 & \quad \downarrow \text{3048} \\
 & \frac{1}{4} \int \cos^2(x) dx - \frac{1}{4} \sin(x) \cos^3(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4} \int \sin\left(x + \frac{\pi}{2}\right)^2 dx - \frac{1}{4} \sin(x) \cos^3(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{1}{4} \left(\int \frac{1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin(x) \cos^3(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{1}{4} \left(\frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin(x) \cos^3(x)
 \end{aligned}$$

input `Int[Cos[x]^2*Sin[x]^2,x]`

output `-1/4*(Cos[x]^3*Sin[x]) + (x/2 + (Cos[x]*Sin[x])/2)/4`

3.262.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3048 `Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^n_*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m_, x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Simp[a^2*((m - 1)/(m + n)) Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

rule 3115 `Int[(b_.)*sin[(c_.) + (d_.)*(x_.)]^n_, x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

3.262.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46

method	result	size
risch	$\frac{x}{8} - \frac{\sin(4x)}{32}$	11
parallelrisch	$\frac{x}{8} - \frac{\sin(4x)}{32}$	11
default	$\frac{x}{8} + \frac{\cos(x) \sin(x)}{8} - \frac{(\cos^3(x) \sin(x))}{4}$	19
norman	$\frac{x}{8} + \frac{7(\tan^3(\frac{x}{2}))}{4} - \frac{7(\tan^5(\frac{x}{2}))}{4} + \frac{(\tan^7(\frac{x}{2}))}{4} + \frac{x(\tan^2(\frac{x}{2}))}{2} + \frac{3x(\tan^4(\frac{x}{2}))}{4} + \frac{x(\tan^6(\frac{x}{2}))}{2} + \frac{x(\tan^8(\frac{x}{2}))}{8} - \frac{\tan(\frac{x}{2})}{4}$ $(1 + \tan^2(\frac{x}{2}))^4$	82

input `int(cos(x)^2*sin(x)^2,x,method=_RETURNVERBOSE)`

output `1/8*x-1/32*sin(4*x)`

3.262.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \cos^2(x) \sin^2(x) dx = -\frac{1}{8} (2 \cos(x)^3 - \cos(x)) \sin(x) + \frac{1}{8} x$$

input `integrate(cos(x)^2*sin(x)^2,x, algorithm="fricas")`output `-1/8*(2*cos(x)^3 - cos(x))*sin(x) + 1/8*x`**3.262.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.58

$$\int \cos^2(x) \sin^2(x) dx = \frac{x}{8} - \frac{\sin(2x) \cos(2x)}{16}$$

input `integrate(cos(x)**2*sin(x)**2,x)`output `x/8 - sin(2*x)*cos(2*x)/16`**3.262.7 Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.42

$$\int \cos^2(x) \sin^2(x) dx = \frac{1}{8} x - \frac{1}{32} \sin(4x)$$

input `integrate(cos(x)^2*sin(x)^2,x, algorithm="maxima")`output `1/8*x - 1/32*sin(4*x)`

3.262.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.42

$$\int \cos^2(x) \sin^2(x) dx = \frac{1}{8} x - \frac{1}{32} \sin(4x)$$

input `integrate(cos(x)^2*sin(x)^2,x, algorithm="giac")`output `1/8*x - 1/32*sin(4*x)`**3.262.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.75

$$\int \cos^2(x) \sin^2(x) dx = \frac{\cos(x) \sin(x)^3}{4} - \frac{\cos(x) \sin(x)}{8} + \frac{x}{8}$$

input `int(cos(x)^2*sin(x)^2,x)`output `x/8 - (cos(x)*sin(x))/8 + (cos(x)*sin(x)^3)/4`

$$3.263 \quad \int \frac{-\cos(x) + \sin(x)}{\cos(x) + \sin(x)} dx$$

3.263.1 Optimal result	1353
3.263.2 Mathematica [A] (verified)	1353
3.263.3 Rubi [A] (verified)	1354
3.263.4 Maple [A] (verified)	1355
3.263.5 Fricas [A] (verification not implemented)	1355
3.263.6 Sympy [A] (verification not implemented)	1355
3.263.7 Maxima [A] (verification not implemented)	1356
3.263.8 Giac [B] (verification not implemented)	1356
3.263.9 Mupad [B] (verification not implemented)	1356

3.263.1 Optimal result

Integrand size = 15, antiderivative size = 8

$$\int \frac{-\cos(x) + \sin(x)}{\cos(x) + \sin(x)} dx = -\log(\cos(x) + \sin(x))$$

output `-ln(cos(x)+sin(x))`

3.263.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{-\cos(x) + \sin(x)}{\cos(x) + \sin(x)} dx = -\log(\cos(x) + \sin(x))$$

input `Integrate[(-Cos[x] + Sin[x])/(Cos[x] + Sin[x]),x]`

output `-Log[Cos[x] + Sin[x]]`

3.263.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 3612}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sin(x) - \cos(x)}{\sin(x) + \cos(x)} dx$$

↓ 3042

$$\int \frac{\sin(x) - \cos(x)}{\sin(x) + \cos(x)} dx$$

↓ 3612

$$-\log(\sin(x) + \cos(x))$$

input `Int[(-Cos[x] + Sin[x])/(Cos[x] + Sin[x]),x]`

output `-Log[Cos[x] + Sin[x]]`

3.263.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3612 `Int[((A_.) + cos[(d_.) + (e_.)*(x_)])*(B_.) + (C_.)*sin[(d_.) + (e_.)*(x_)]) / ((a_.) + cos[(d_.) + (e_.)*(x_)])*(b_.) + (c_.)*sin[(d_.) + (e_.)*(x_)]), x_Symbol] := Simp[(b*B + c*C)*(x/(b^2 + c^2)), x] + Simp[(c*B - b*C)*(Log[a + b*Cos[d + e*x] + c*Sin[d + e*x]]/(e*(b^2 + c^2))), x] /; FreeQ[{a, b, c, d, e, A, B, C}, x] && NeQ[b^2 + c^2, 0] && EqQ[A*(b^2 + c^2) - a*(b*B + c*C), 0]`

3.263.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
derivativedivides	$-\ln(\cos(x) + \sin(x))$	9
default	$-\ln(\cos(x) + \sin(x))$	9
risch	$ix - \ln(e^{2ix} + i)$	17
norman	$-\ln(\tan^2(\frac{x}{2}) - 2\tan(\frac{x}{2}) - 1) + \ln(1 + \tan^2(\frac{x}{2}))$	28
parallelrisc	$-\ln\left(\frac{-\sin(x)-\cos(x)}{\cos(x)+1}\right) + \ln\left(\frac{1}{\cos(x)+1}\right)$	28

input `int((-cos(x)+sin(x))/(cos(x)+sin(x)),x,method=_RETURNVERBOSE)`output `-ln(cos(x)+sin(x))`**3.263.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.38

$$\int \frac{-\cos(x) + \sin(x)}{\cos(x) + \sin(x)} dx = -\frac{1}{2} \log(2 \cos(x) \sin(x) + 1)$$

input `integrate((-cos(x)+sin(x))/(cos(x)+sin(x)),x, algorithm="fracas")`output `-1/2*log(2*cos(x)*sin(x) + 1)`**3.263.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{-\cos(x) + \sin(x)}{\cos(x) + \sin(x)} dx = -\log(\sin(x) + \cos(x))$$

input `integrate((-cos(x)+sin(x))/(cos(x)+sin(x)),x)`output `-log(sin(x) + cos(x))`

3.263.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{-\cos(x) + \sin(x)}{\cos(x) + \sin(x)} dx = -\log(\cos(x) + \sin(x))$$

input `integrate((-cos(x)+sin(x))/(cos(x)+sin(x)),x, algorithm="maxima")`

output `-log(cos(x) + sin(x))`

3.263.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18 vs. 2(8) = 16.

Time = 0.29 (sec) , antiderivative size = 18, normalized size of antiderivative = 2.25

$$\int \frac{-\cos(x) + \sin(x)}{\cos(x) + \sin(x)} dx = \frac{1}{2} \log(\tan(x)^2 + 1) - \log(|\tan(x) + 1|)$$

input `integrate((-cos(x)+sin(x))/(cos(x)+sin(x)),x, algorithm="giac")`

output `1/2*log(tan(x)^2 + 1) - log(abs(tan(x) + 1))`

3.263.9 Mupad [B] (verification not implemented)

Time = 0.50 (sec) , antiderivative size = 32, normalized size of antiderivative = 4.00

$$\int \frac{-\cos(x) + \sin(x)}{\cos(x) + \sin(x)} dx = -2 \operatorname{atanh}\left(\frac{128 \tan\left(\frac{x}{2}\right) + 128}{16 \tan\left(\frac{x}{2}\right)^2 + 32 \tan\left(\frac{x}{2}\right) + 48} - 3\right)$$

input `int(-(cos(x) - sin(x))/(cos(x) + sin(x)),x)`

output `-2*atanh((128*tan(x/2) + 128)/(32*tan(x/2) + 16*tan(x/2)^2 + 48) - 3)`

3.264 $\int \frac{x}{\sqrt{1-x^2}} dx$

3.264.1 Optimal result	1357
3.264.2 Mathematica [A] (verified)	1357
3.264.3 Rubi [A] (verified)	1358
3.264.4 Maple [A] (verified)	1358
3.264.5 Fricas [A] (verification not implemented)	1359
3.264.6 Sympy [A] (verification not implemented)	1359
3.264.7 Maxima [A] (verification not implemented)	1359
3.264.8 Giac [A] (verification not implemented)	1360
3.264.9 Mupad [B] (verification not implemented)	1360

3.264.1 Optimal result

Integrand size = 13, antiderivative size = 13

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{1-x^2}$$

output `-(-x^2+1)^(1/2)`

3.264.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{1-x^2}$$

input `Integrate[x/Sqrt[1 - x^2],x]`

output `-Sqrt[1 - x^2]`

3.264.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt{1-x^2}} dx$$

↓ 241

$$-\sqrt{1-x^2}$$

input `Int[x/Sqrt[1 - x^2],x]`

output `-Sqrt[1 - x^2]`

3.264.3.1 Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

3.264.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
derivativedivides	$-\sqrt{-x^2 + 1}$	12
default	$-\sqrt{-x^2 + 1}$	12
trager	$-\sqrt{-x^2 + 1}$	12
pseudoelliptic	$-\sqrt{-x^2 + 1}$	12
risch	$\frac{x^2-1}{\sqrt{-x^2+1}}$	16
gospers	$\frac{(-1+x)(1+x)}{\sqrt{-x^2+1}}$	17
meijerg	$-\frac{-2\sqrt{\pi}+2\sqrt{\pi}\sqrt{-x^2+1}}{2\sqrt{\pi}}$	26

input `int(1/(-x^2+1)^(1/2)*x,x,method=_RETURNVERBOSE)`

output `-(-x^2+1)^(1/2)`

3.264.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{-x^2+1}$$

input `integrate(x/(-x^2+1)^(1/2),x, algorithm="fricas")`

output `-sqrt(-x^2 + 1)`

3.264.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.62

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{1-x^2}$$

input `integrate(x/(-x**2+1)**(1/2),x)`

output `-sqrt(1 - x**2)`

3.264.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{-x^2+1}$$

input `integrate(x/(-x^2+1)^(1/2),x, algorithm="maxima")`

output `-sqrt(-x^2 + 1)`

3.264.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{-x^2+1}$$

input `integrate(x/(-x^2+1)^(1/2),x, algorithm="giac")`output `-sqrt(-x^2 + 1)`**3.264.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{1-x^2}$$

input `int(x/(1 - x^2)^(1/2),x)`output `-(1 - x^2)^(1/2)`

3.265 $\int x^3 \log(x) dx$

3.265.1 Optimal result1361
3.265.2 Mathematica [A] (verified)1361
3.265.3 Rubi [A] (verified)1362
3.265.4 Maple [A] (verified)1362
3.265.5 Fracas [A] (verification not implemented)1363
3.265.6 Sympy [A] (verification not implemented)1363
3.265.7 Maxima [A] (verification not implemented)1363
3.265.8 Giac [A] (verification not implemented)1364
3.265.9 Mupad [B] (verification not implemented)1364

3.265.1 Optimal result

Integrand size = 6, antiderivative size = 17

$$\int x^3 \log(x) dx = -\frac{x^4}{16} + \frac{1}{4}x^4 \log(x)$$

output `-1/16*x^4+1/4*x^4*ln(x)`

3.265.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int x^3 \log(x) dx = -\frac{x^4}{16} + \frac{1}{4}x^4 \log(x)$$

input `Integrate[x^3*Log[x],x]`

output `-1/16*x^4 + (x^4*Log[x])/4`

3.265.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \log(x) dx$$

$$\downarrow \text{2741}$$

$$\frac{1}{4}x^4 \log(x) - \frac{x^4}{16}$$

input `Int[x^3*Log[x],x]`

output `-1/16*x^4 + (x^4*Log[x])/4`

3.265.3.1 Defintions of rubi rules used

rule 2741 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_.), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(m + 1))/(d*(m + 1)^2), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]`

3.265.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{x^4}{16} + \frac{x^4 \ln(x)}{4}$	14
norman	$-\frac{x^4}{16} + \frac{x^4 \ln(x)}{4}$	14
risch	$-\frac{x^4}{16} + \frac{x^4 \ln(x)}{4}$	14
parallelrisch	$-\frac{x^4}{16} + \frac{x^4 \ln(x)}{4}$	14
parts	$-\frac{x^4}{16} + \frac{x^4 \ln(x)}{4}$	14

input `int(x^3*ln(x),x,method=_RETURNVERBOSE)`

output `-1/16*x^4+1/4*x^4*ln(x)`

3.265.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int x^3 \log(x) dx = \frac{1}{4} x^4 \log(x) - \frac{1}{16} x^4$$

input `integrate(x^3*log(x),x, algorithm="fricas")`

output `1/4*x^4*log(x) - 1/16*x^4`

3.265.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int x^3 \log(x) dx = \frac{x^4 \log(x)}{4} - \frac{x^4}{16}$$

input `integrate(x**3*ln(x),x)`

output `x**4*log(x)/4 - x**4/16`

3.265.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int x^3 \log(x) dx = \frac{1}{4} x^4 \log(x) - \frac{1}{16} x^4$$

input `integrate(x^3*log(x),x, algorithm="maxima")`

output `1/4*x^4*log(x) - 1/16*x^4`

3.265.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int x^3 \log(x) dx = \frac{1}{4} x^4 \log(x) - \frac{1}{16} x^4$$

input `integrate(x^3*log(x),x, algorithm="giac")`

output `1/4*x^4*log(x) - 1/16*x^4`

3.265.9 Mupad [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.53

$$\int x^3 \log(x) dx = \frac{x^4 (\ln(x) - \frac{1}{4})}{4}$$

input `int(x^3*log(x),x)`

output `(x^4*(log(x) - 1/4))/4`

3.266 $\int \frac{\sqrt{-2+x}}{2+x} dx$

3.266.1 Optimal result	1365
3.266.2 Mathematica [A] (verified)	1365
3.266.3 Rubi [A] (verified)	1366
3.266.4 Maple [A] (verified)	1367
3.266.5 Fricas [A] (verification not implemented)	1367
3.266.6 Sympy [C] (verification not implemented)	1368
3.266.7 Maxima [A] (verification not implemented)	1368
3.266.8 Giac [A] (verification not implemented)	1368
3.266.9 Mupad [B] (verification not implemented)	1369

3.266.1 Optimal result

Integrand size = 13, antiderivative size = 24

$$\int \frac{\sqrt{-2+x}}{2+x} dx = 2\sqrt{-2+x} - 4 \arctan\left(\frac{\sqrt{-2+x}}{2}\right)$$

output `-4*arctan(1/2*(-2+x)^(1/2))+2*(-2+x)^(1/2)`

3.266.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{-2+x}}{2+x} dx = 2\sqrt{-2+x} - 4 \arctan\left(\frac{\sqrt{-2+x}}{2}\right)$$

input `Integrate[Sqrt[-2 + x]/(2 + x),x]`

output `2*Sqrt[-2 + x] - 4*ArcTan[Sqrt[-2 + x]/2]`

3.266.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{x-2}}{x+2} dx$$

$$\downarrow 60$$

$$2\sqrt{x-2} - 4 \int \frac{1}{\sqrt{x-2}(x+2)} dx$$

$$\downarrow 73$$

$$2\sqrt{x-2} - 8 \int \frac{1}{x+2} d\sqrt{x-2}$$

$$\downarrow 216$$

$$2\sqrt{x-2} - 4 \arctan\left(\frac{\sqrt{x-2}}{2}\right)$$

input `Int[Sqrt[-2 + x]/(2 + x),x]`

output `2*Sqrt[-2 + x] - 4*ArcTan[Sqrt[-2 + x]/2]`

3.266.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

3.266.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

method	result	size
derivativedivides	$-4 \arctan\left(\frac{\sqrt{-2+x}}{2}\right) + 2\sqrt{-2+x}$	19
default	$-4 \arctan\left(\frac{\sqrt{-2+x}}{2}\right) + 2\sqrt{-2+x}$	19
risch	$-4 \arctan\left(\frac{\sqrt{-2+x}}{2}\right) + 2\sqrt{-2+x}$	19
trager	$2\sqrt{-2+x} + 2 \operatorname{RootOf}(_Z^2 + 1) \ln\left(\frac{\operatorname{RootOf}(_Z^2 + 1)x - 6 \operatorname{RootOf}(_Z^2 + 1) + 4\sqrt{-2+x}}{2+x}\right)$	48

```
input int((-2+x)^(1/2)/(2+x),x,method=_RETURNVERBOSE)
```

```
output -4*arctan(1/2*(-2+x)^(1/2))+2*(-2+x)^(1/2)
```

3.266.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{-2+x}}{2+x} dx = 2\sqrt{x-2} - 4 \arctan\left(\frac{1}{2}\sqrt{x-2}\right)$$

```
input integrate((-2+x)^(1/2)/(2+x),x, algorithm="fricas")
```

```
output 2*sqrt(x - 2) - 4*arctan(1/2*sqrt(x - 2))
```


3.266.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.77 (sec) , antiderivative size = 109, normalized size of antiderivative = 4.54

$$\int \frac{\sqrt{-2+x}}{2+x} dx = \begin{cases} -4i \operatorname{acosh}\left(\frac{2}{\sqrt{x+2}}\right) - \frac{2i\sqrt{x+2}}{\sqrt{-1+\frac{4}{x+2}}} + \frac{8i}{\sqrt{-1+\frac{4}{x+2}}\sqrt{x+2}} & \text{for } \frac{1}{|x+2|} > \frac{1}{4} \\ 4 \operatorname{asin}\left(\frac{2}{\sqrt{x+2}}\right) + \frac{2\sqrt{x+2}}{\sqrt{1-\frac{4}{x+2}}} - \frac{8}{\sqrt{1-\frac{4}{x+2}}\sqrt{x+2}} & \text{otherwise} \end{cases}$$

input `integrate((-2+x)**(1/2)/(2+x),x)`

output `Piecewise((-4*I*acosh(2/sqrt(x + 2)) - 2*I*sqrt(x + 2)/sqrt(-1 + 4/(x + 2)) + 8*I/(sqrt(-1 + 4/(x + 2))*sqrt(x + 2)), 1/Abs(x + 2) > 1/4), (4*asin(2/sqrt(x + 2)) + 2*sqrt(x + 2)/sqrt(1 - 4/(x + 2)) - 8/(sqrt(1 - 4/(x + 2))*sqrt(x + 2)), True))`

3.266.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{-2+x}}{2+x} dx = 2\sqrt{x-2} - 4 \arctan\left(\frac{1}{2}\sqrt{x-2}\right)$$

input `integrate((-2+x)^(1/2)/(2+x),x, algorithm="maxima")`

output `2*sqrt(x - 2) - 4*arctan(1/2*sqrt(x - 2))`

3.266.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{-2+x}}{2+x} dx = 2\sqrt{x-2} - 4 \arctan\left(\frac{1}{2}\sqrt{x-2}\right)$$

input `integrate((-2+x)^(1/2)/(2+x),x, algorithm="giac")`

output `2*sqrt(x - 2) - 4*arctan(1/2*sqrt(x - 2))`

3.266.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{-2+x}}{2+x} dx = 2\sqrt{x-2} - 4 \operatorname{atan}\left(\frac{\sqrt{x-2}}{2}\right)$$

input `int((x - 2)^(1/2)/(x + 2),x)`

output `2*(x - 2)^(1/2) - 4*atan((x - 2)^(1/2)/2)`

3.267 $\int \frac{x}{(2+x)^2} dx$

3.267.1 Optimal result	1370
3.267.2 Mathematica [A] (verified)	1370
3.267.3 Rubi [A] (verified)	1371
3.267.4 Maple [A] (verified)	1372
3.267.5 Fricas [A] (verification not implemented)	1372
3.267.6 Sympy [A] (verification not implemented)	1372
3.267.7 Maxima [A] (verification not implemented)	1373
3.267.8 Giac [A] (verification not implemented)	1373
3.267.9 Mupad [B] (verification not implemented)	1373

3.267.1 Optimal result

Integrand size = 7, antiderivative size = 12

$$\int \frac{x}{(2+x)^2} dx = \frac{2}{2+x} + \log(2+x)$$

output `2/(2+x)+ln(2+x)`

3.267.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{x}{(2+x)^2} dx = \frac{2}{2+x} + \log(2+x)$$

input `Integrate[x/(2 + x)^2,x]`

output `2/(2 + x) + Log[2 + x]`

3.267.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(x+2)^2} dx$$

$$\downarrow 49$$

$$\int \left(\frac{1}{x+2} - \frac{2}{(x+2)^2} \right) dx$$

$$\downarrow 2009$$

$$\frac{2}{x+2} + \log(x+2)$$

input `Int[x/(2 + x)^2,x]`

output `2/(2 + x) + Log[2 + x]`

3.267.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.267.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

method	result	size
default	$\frac{2}{2+x} + \ln(2+x)$	13
norman	$\frac{2}{2+x} + \ln(2+x)$	13
risch	$\frac{2}{2+x} + \ln(2+x)$	13
meijerg	$-\frac{x}{2(1+\frac{x}{2})} + \ln(1+\frac{x}{2})$	18
parallelrisch	$\frac{\ln(2+x)x+2+2\ln(2+x)}{2+x}$	21

input `int(x/(2+x)^2,x,method=_RETURNVERBOSE)`output `2/(2+x)+ln(2+x)`**3.267.5 Fricas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.33

$$\int \frac{x}{(2+x)^2} dx = \frac{(x+2) \log(x+2) + 2}{x+2}$$

input `integrate(x/(2+x)^2,x, algorithm="fricas")`output `((x + 2)*log(x + 2) + 2)/(x + 2)`**3.267.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x}{(2+x)^2} dx = \log(x+2) + \frac{2}{x+2}$$

input `integrate(x/(2+x)**2,x)`output `log(x + 2) + 2/(x + 2)`

3.267.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{x}{(2+x)^2} dx = \frac{2}{x+2} + \log(x+2)$$

input `integrate(x/(2+x)^2,x, algorithm="maxima")`output `2/(x + 2) + log(x + 2)`**3.267.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

$$\int \frac{x}{(2+x)^2} dx = \frac{2}{x+2} + \log(|x+2|)$$

input `integrate(x/(2+x)^2,x, algorithm="giac")`output `2/(x + 2) + log(abs(x + 2))`**3.267.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{x}{(2+x)^2} dx = \ln(x+2) + \frac{2}{x+2}$$

input `int(x/(x + 2)^2,x)`output `log(x + 2) + 2/(x + 2)`

3.268 $\int \log(1 + x^2) dx$

3.268.1 Optimal result	1374
3.268.2 Mathematica [A] (verified)	1374
3.268.3 Rubi [A] (verified)	1375
3.268.4 Maple [A] (verified)	1376
3.268.5 Fricas [A] (verification not implemented)	1376
3.268.6 Sympy [A] (verification not implemented)	1377
3.268.7 Maxima [A] (verification not implemented)	1377
3.268.8 Giac [A] (verification not implemented)	1377
3.268.9 Mupad [B] (verification not implemented)	1378

3.268.1 Optimal result

Integrand size = 6, antiderivative size = 16

$$\int \log(1 + x^2) dx = -2x + 2 \arctan(x) + x \log(1 + x^2)$$

output `-2*x+2*arctan(x)+x*ln(x^2+1)`

3.268.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \log(1 + x^2) dx = -2x + 2 \arctan(x) + x \log(1 + x^2)$$

input `Integrate[Log[1 + x^2],x]`

output `-2*x + 2*ArcTan[x] + x*Log[1 + x^2]`

3.268.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {2898, 262, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \log(x^2 + 1) dx \\ & \quad \downarrow \text{2898} \\ & x \log(x^2 + 1) - 2 \int \frac{x^2}{x^2 + 1} dx \\ & \quad \downarrow \text{262} \\ & x \log(x^2 + 1) - 2 \left(x - \int \frac{1}{x^2 + 1} dx \right) \\ & \quad \downarrow \text{216} \\ & x \log(x^2 + 1) - 2(x - \arctan(x)) \end{aligned}$$

input `Int[Log[1 + x^2], x]`

output `-2*(x - ArcTan[x]) + x*Log[1 + x^2]`

3.268.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 262 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*(m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2898 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_))^(p_.)], x_Symbol] := Simp[x*Log[c*(d + e*x^n)^p], x] - Simp[e*n*p Int[x^n/(d + e*x^n), x], x] /; FreeQ[{c, d, e, n, p}, x]`

3.268.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

method	result	size
default	$-2x + 2 \arctan(x) + x \ln(x^2 + 1)$	17
risch	$-2x + 2 \arctan(x) + x \ln(x^2 + 1)$	17
parts	$-2x + 2 \arctan(x) + x \ln(x^2 + 1)$	17
meijerg	$-2x + \frac{2x \arctan(\sqrt{x^2})}{\sqrt{x^2}} + x \ln(x^2 + 1)$	27
parallelrisc	$-2i \ln(x - i) + i \ln(x^2 + 1) + x \ln(x^2 + 1) - 2x$	30

input `int(ln(x^2+1),x,method=_RETURNVERBOSE)`

output `-2*x+2*arctan(x)+x*ln(x^2+1)`

3.268.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \log(1 + x^2) dx = x \log(x^2 + 1) - 2x + 2 \arctan(x)$$

input `integrate(log(x^2+1),x, algorithm="fricas")`

output `x*log(x^2 + 1) - 2*x + 2*arctan(x)`

3.268.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \log(1+x^2) dx = x \log(x^2+1) - 2x + 2 \operatorname{atan}(x)$$

input `integrate(ln(x**2+1),x)`output `x*log(x**2 + 1) - 2*x + 2*atan(x)`**3.268.7 Maxima [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \log(1+x^2) dx = x \log(x^2+1) - 2x + 2 \operatorname{arctan}(x)$$

input `integrate(log(x^2+1),x, algorithm="maxima")`output `x*log(x^2 + 1) - 2*x + 2*arctan(x)`**3.268.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \log(1+x^2) dx = x \log(x^2+1) - 2x + 2 \operatorname{arctan}(x)$$

input `integrate(log(x^2+1),x, algorithm="giac")`output `x*log(x^2 + 1) - 2*x + 2*arctan(x)`

3.268.9 Mupad [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \log(1 + x^2) dx = 2 \operatorname{atan}(x) - 2x + x \ln(x^2 + 1)$$

input `int(log(x^2 + 1),x)`

output `2*atan(x) - 2*x + x*log(x^2 + 1)`

3.269 $\int \frac{\sqrt{1+\log(x)}}{x \log(x)} dx$

3.269.1 Optimal result	1379
3.269.2 Mathematica [A] (verified)	1379
3.269.3 Rubi [A] (verified)	1380
3.269.4 Maple [A] (verified)	1381
3.269.5 Fricas [A] (verification not implemented)	1381
3.269.6 Sympy [A] (verification not implemented)	1382
3.269.7 Maxima [A] (verification not implemented)	1382
3.269.8 Giac [F(-1)]	1382
3.269.9 Mupad [B] (verification not implemented)	1383

3.269.1 Optimal result

Integrand size = 16, antiderivative size = 22

$$\int \frac{\sqrt{1+\log(x)}}{x \log(x)} dx = -2\operatorname{arctanh}\left(\sqrt{1+\log(x)}\right) + 2\sqrt{1+\log(x)}$$

output `-2*arctanh((1+ln(x))^(1/2))+2*(1+ln(x))^(1/2)`

3.269.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1+\log(x)}}{x \log(x)} dx = -2\operatorname{arctanh}\left(\sqrt{1+\log(x)}\right) + 2\sqrt{1+\log(x)}$$

input `Integrate[Sqrt[1 + Log[x]]/(x*Log[x]),x]`

output `-2*ArcTanh[Sqrt[1 + Log[x]]] + 2*Sqrt[1 + Log[x]]`

3.269.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2812, 60, 73, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{\log(x)+1}}{x \log(x)} dx \\ & \quad \downarrow \text{2812} \\ & \int \frac{\sqrt{\log(x)+1}}{\log(x)} d\log(x) \\ & \quad \downarrow \text{60} \\ & \int \frac{1}{\log(x)\sqrt{\log(x)+1}} d\log(x) + 2\sqrt{\log(x)+1} \\ & \quad \downarrow \text{73} \\ & 2 \int \frac{1}{\log(x)} d\sqrt{\log(x)+1} + 2\sqrt{\log(x)+1} \\ & \quad \downarrow \text{220} \\ & 2\sqrt{\log(x)+1} - 2\operatorname{arctanh}\left(\sqrt{\log(x)+1}\right) \end{aligned}$$

input `Int[Sqrt[1 + Log[x]]/(x*Log[x]),x]`

output `-2*ArcTanh[Sqrt[1 + Log[x]]] + 2*Sqrt[1 + Log[x]]`

3.269.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 220 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-
1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] &&
(LtQ[a, 0] || GtQ[b, 0])
```

```
rule 2812 Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((d_.) + Log[(c_.)*(x_)^(n
_.)]*(e_.))^(q_.))/(x_), x_Symbol] := Simp[1/n Subst[Int[(a + b*x)^p*(d +
e*x)^q, x], x, Log[c*x^n]], x] /; FreeQ[{a, b, c, d, e, n, p, q}, x]
```

3.269.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.36

method	result	size
derivativdivides	$2\sqrt{1 + \ln(x)} + \ln(\sqrt{1 + \ln(x)} - 1) - \ln(\sqrt{1 + \ln(x)} + 1)$	30
default	$2\sqrt{1 + \ln(x)} + \ln(\sqrt{1 + \ln(x)} - 1) - \ln(\sqrt{1 + \ln(x)} + 1)$	30

```
input int((1+ln(x))^(1/2)/x/ln(x),x,method=_RETURNVERBOSE)
```

```
output 2*(1+ln(x))^(1/2)+ln((1+ln(x))^(1/2)-1)-ln((1+ln(x))^(1/2)+1)
```

3.269.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.32

$$\int \frac{\sqrt{1 + \log(x)}}{x \log(x)} dx = 2\sqrt{\log(x) + 1} - \log(\sqrt{\log(x) + 1} + 1) + \log(\sqrt{\log(x) + 1} - 1)$$

```
input integrate((1+log(x))^(1/2)/x/log(x),x, algorithm="fracas")
```

output `2*sqrt(log(x) + 1) - log(sqrt(log(x) + 1) + 1) + log(sqrt(log(x) + 1) - 1)`

3.269.6 Sympy [A] (verification not implemented)

Time = 0.63 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.45

$$\int \frac{\sqrt{1 + \log(x)}}{x \log(x)} dx = 2\sqrt{\log(x) + 1} + \log\left(\sqrt{\log(x) + 1} - 1\right) - \log\left(\sqrt{\log(x) + 1} + 1\right)$$

input `integrate((1+ln(x))**(1/2)/x/ln(x),x)`

output `2*sqrt(log(x) + 1) + log(sqrt(log(x) + 1) - 1) - log(sqrt(log(x) + 1) + 1)`

3.269.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.32

$$\int \frac{\sqrt{1 + \log(x)}}{x \log(x)} dx = 2\sqrt{\log(x) + 1} - \log\left(\sqrt{\log(x) + 1} + 1\right) + \log\left(\sqrt{\log(x) + 1} - 1\right)$$

input `integrate((1+log(x))^(1/2)/x/log(x),x, algorithm="maxima")`

output `2*sqrt(log(x) + 1) - log(sqrt(log(x) + 1) + 1) + log(sqrt(log(x) + 1) - 1)`

3.269.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{1 + \log(x)}}{x \log(x)} dx = \text{Timed out}$$

input `integrate((1+log(x))^(1/2)/x/log(x),x, algorithm="giac")`

output `Timed out`

3.269.9 Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{\sqrt{1 + \log(x)}}{x \log(x)} dx = 2 \sqrt{\ln(x) + 1} - 2 \operatorname{atanh}\left(\sqrt{\ln(x) + 1}\right)$$

input `int((log(x) + 1)^(1/2)/(x*log(x)),x)`

output `2*(log(x) + 1)^(1/2) - 2*atanh((log(x) + 1)^(1/2))`

3.270 $\int (1 + \sqrt{x})^8 dx$

3.270.1 Optimal result	1384
3.270.2 Mathematica [B] (verified)	1384
3.270.3 Rubi [A] (verified)	1385
3.270.4 Maple [B] (verified)	1386
3.270.5 Fricas [B] (verification not implemented)	1386
3.270.6 Sympy [B] (verification not implemented)	1387
3.270.7 Maxima [A] (verification not implemented)	1387
3.270.8 Giac [B] (verification not implemented)	1387
3.270.9 Mupad [B] (verification not implemented)	1388

3.270.1 Optimal result

Integrand size = 9, antiderivative size = 27

$$\int (1 + \sqrt{x})^8 dx = -\frac{2}{9}(1 + \sqrt{x})^9 + \frac{1}{5}(1 + \sqrt{x})^{10}$$

output `-2/9*(1+x^(1/2))^9+1/5*(1+x^(1/2))^10`

3.270.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 56 vs. 2(27) = 54.

Time = 0.02 (sec) , antiderivative size = 56, normalized size of antiderivative = 2.07

$$\int (1 + \sqrt{x})^8 dx = \frac{1}{45}(45x + 240x^{3/2} + 630x^2 + 1008x^{5/2} + 1050x^3 + 720x^{7/2} + 315x^4 + 80x^{9/2} + 9x^5)$$

input `Integrate[(1 + Sqrt[x])^8,x]`

output `(45*x + 240*x^(3/2) + 630*x^2 + 1008*x^(5/2) + 1050*x^3 + 720*x^(7/2) + 315*x^4 + 80*x^(9/2) + 9*x^5)/45`

3.270.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {774, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (\sqrt{x} + 1)^8 dx \\ & \quad \downarrow 774 \\ & 2 \int (\sqrt{x} + 1)^8 \sqrt{x} d\sqrt{x} \\ & \quad \downarrow 49 \\ & 2 \int \left((\sqrt{x} + 1)^9 - (\sqrt{x} + 1)^8 \right) d\sqrt{x} \\ & \quad \downarrow 2009 \\ & 2 \left(\frac{1}{10} (\sqrt{x} + 1)^{10} - \frac{1}{9} (\sqrt{x} + 1)^9 \right) \end{aligned}$$

input `Int[(1 + Sqrt[x])^8,x]`

output `2*(-1/9*(1 + Sqrt[x])^9 + (1 + Sqrt[x])^10/10)`

3.270.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 774 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[n]}, Simp[k Subst[Int[x^(k - 1)*(a + b*x^(k*n))^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, p}, x] && FractionQ[n]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.270.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. $2(19) = 38$.

Time = 0.12 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.59

method	result	size
derivativedivides	$\frac{x^5}{5} + \frac{16x^{\frac{9}{2}}}{9} + 7x^4 + 16x^{\frac{7}{2}} + \frac{70x^3}{3} + \frac{112x^{\frac{5}{2}}}{5} + 14x^2 + \frac{16x^{\frac{3}{2}}}{3} + x$	43
default	$\frac{x^5}{5} + \frac{16x^{\frac{9}{2}}}{9} + 7x^4 + 16x^{\frac{7}{2}} + \frac{70x^3}{3} + \frac{112x^{\frac{5}{2}}}{5} + 14x^2 + \frac{16x^{\frac{3}{2}}}{3} + x$	43
trager	$\frac{(3x^4+108x^3+458x^2+668x+683)(-1+x)}{15} + \frac{16x^{\frac{3}{2}}(5x^3+45x^2+63x+15)}{45}$	47

input `int((x^(1/2)+1)^8,x,method=_RETURNVERBOSE)`

output `1/5*x^5+16/9*x^(9/2)+7*x^4+16*x^(7/2)+70/3*x^3+112/5*x^(5/2)+14*x^2+16/3*x^(3/2)+x`

3.270.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 46 vs. $2(19) = 38$.

Time = 0.24 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.70

$$\int (1 + \sqrt{x})^8 dx = \frac{1}{5} x^5 + 7x^4 + \frac{70}{3} x^3 + 14x^2 + \frac{16}{45} (5x^4 + 45x^3 + 63x^2 + 15x)\sqrt{x} + x$$

input `integrate((1+x^(1/2))^8,x, algorithm="fricas")`

output `1/5*x^5 + 7*x^4 + 70/3*x^3 + 14*x^2 + 16/45*(5*x^4 + 45*x^3 + 63*x^2 + 15*x)*sqrt(x) + x`

3.270.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 54 vs. $2(20) = 40$.

Time = 0.13 (sec) , antiderivative size = 54, normalized size of antiderivative = 2.00

$$\int (1 + \sqrt{x})^8 dx = \frac{16x^{\frac{9}{2}}}{9} + 16x^{\frac{7}{2}} + \frac{112x^{\frac{5}{2}}}{5} + \frac{16x^{\frac{3}{2}}}{3} + \frac{x^5}{5} + 7x^4 + \frac{70x^3}{3} + 14x^2 + x$$

input `integrate((1+x**(1/2))**8,x)`

output `16*x**(9/2)/9 + 16*x**(7/2) + 112*x**(5/2)/5 + 16*x**(3/2)/3 + x**5/5 + 7*x**4 + 70*x**3/3 + 14*x**2 + x`

3.270.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int (1 + \sqrt{x})^8 dx = \frac{1}{5} (\sqrt{x} + 1)^{10} - \frac{2}{9} (\sqrt{x} + 1)^9$$

input `integrate((1+x^(1/2))^8,x, algorithm="maxima")`

output `1/5*(sqrt(x) + 1)^10 - 2/9*(sqrt(x) + 1)^9`

3.270.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. $2(19) = 38$.

Time = 0.47 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.56

$$\int (1 + \sqrt{x})^8 dx = \frac{1}{5} x^5 + \frac{16}{9} x^{\frac{9}{2}} + 7x^4 + 16x^{\frac{7}{2}} + \frac{70}{3} x^3 + \frac{112}{5} x^{\frac{5}{2}} + 14x^2 + \frac{16}{3} x^{\frac{3}{2}} + x$$

input `integrate((1+x^(1/2))^8,x, algorithm="giac")`

output `1/5*x^5 + 16/9*x^(9/2) + 7*x^4 + 16*x^(7/2) + 70/3*x^3 + 112/5*x^(5/2) + 14*x^2 + 16/3*x^(3/2) + x`

3.270.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.56

$$\int (1 + \sqrt{x})^8 dx = x + 14x^2 + \frac{70x^3}{3} + 7x^4 + \frac{16x^{3/2}}{3} + \frac{x^5}{5} + \frac{112x^{5/2}}{5} + 16x^{7/2} + \frac{16x^{9/2}}{9}$$

input `int((x^(1/2) + 1)^8,x)`

output `x + 14*x^2 + (70*x^3)/3 + 7*x^4 + (16*x^(3/2))/3 + x^5/5 + (112*x^(5/2))/5 + 16*x^(7/2) + (16*x^(9/2))/9`

3.271 $\int \sec^4(x) \tan^3(x) dx$

3.271.1 Optimal result	1389
3.271.2 Mathematica [A] (verified)	1389
3.271.3 Rubi [A] (verified)	1390
3.271.4 Maple [A] (verified)	1391
3.271.5 Fricas [A] (verification not implemented)	1392
3.271.6 Sympy [A] (verification not implemented)	1392
3.271.7 Maxima [B] (verification not implemented)	1392
3.271.8 Giac [A] (verification not implemented)	1393
3.271.9 Mupad [B] (verification not implemented)	1393

3.271.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \sec^4(x) \tan^3(x) dx = -\frac{1}{4} \sec^4(x) + \frac{\sec^6(x)}{6}$$

output `-1/4*sec(x)^4+1/6*sec(x)^6`

3.271.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sec^4(x) \tan^3(x) dx = -\frac{1}{4} \sec^4(x) + \frac{\sec^6(x)}{6}$$

input `Integrate[Sec[x]^4*Tan[x]^3,x]`

output `-1/4*Sec[x]^4 + Sec[x]^6/6`

3.271.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 3086, 25, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^3(x) \sec^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^3 \sec(x)^4 dx \\
 & \quad \downarrow \text{3086} \\
 & \int -\sec^3(x) (1 - \sec^2(x)) d \sec(x) \\
 & \quad \downarrow \text{25} \\
 & - \int \sec^3(x) (1 - \sec^2(x)) d \sec(x) \\
 & \quad \downarrow \text{244} \\
 & - \int (\sec^3(x) - \sec^5(x)) d \sec(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sec^6(x)}{6} - \frac{\sec^4(x)}{4}
 \end{aligned}$$

input `Int [Sec [x]^4*Tan [x]^3,x]`

output `-1/4*Sec [x]^4 + Sec [x]^6/6`

3.271.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 244 `Int[((c_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

3.271.4 Maple [A] (verified)

Time = 2.79 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$-\frac{(\sec^4(x))}{4} + \frac{(\sec^6(x))}{6}$	14
default	$-\frac{(\sec^4(x))}{4} + \frac{(\sec^6(x))}{6}$	14
risch	$-\frac{4(3e^{8ix} - 2e^{6ix} + 3e^{4ix})}{3(e^{2ix} + 1)^6}$	34

input `int(sec(x)^4*tan(x)^3,x,method=_RETURNVERBOSE)`

output `-1/4*sec(x)^4+1/6*sec(x)^6`

3.271.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^4(x) \tan^3(x) dx = -\frac{3 \cos(x)^2 - 2}{12 \cos(x)^6}$$

input `integrate(sec(x)^4*tan(x)^3,x, algorithm="fricas")`

output `-1/12*(3*cos(x)^2 - 2)/cos(x)^6`

3.271.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^4(x) \tan^3(x) dx = \frac{2 - 3 \cos^2(x)}{12 \cos^6(x)}$$

input `integrate(sec(x)**4*tan(x)**3,x)`

output `(2 - 3*cos(x)**2)/(12*cos(x)**6)`

3.271.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 30 vs. $2(13) = 26$.

Time = 0.20 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.76

$$\int \sec^4(x) \tan^3(x) dx = -\frac{3 \sin(x)^2 - 1}{12 (\sin(x)^6 - 3 \sin(x)^4 + 3 \sin(x)^2 - 1)}$$

input `integrate(sec(x)^4*tan(x)^3,x, algorithm="maxima")`

output `-1/12*(3*sin(x)^2 - 1)/(sin(x)^6 - 3*sin(x)^4 + 3*sin(x)^2 - 1)`

3.271.8 Giac [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^4(x) \tan^3(x) dx = -\frac{3 \cos(x)^2 - 2}{12 \cos(x)^6}$$

input `integrate(sec(x)^4*tan(x)^3,x, algorithm="giac")`output `-1/12*(3*cos(x)^2 - 2)/cos(x)^6`**3.271.9 Mupad [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^4(x) \tan^3(x) dx = \frac{\tan(x)^4 (2 \tan(x)^2 + 3)}{12}$$

input `int(tan(x)^3/cos(x)^4,x)`output `(tan(x)^4*(2*tan(x)^2 + 3))/12`

3.272 $\int \frac{x}{2-2x+x^2} dx$

3.272.1 Optimal result	1394
3.272.2 Mathematica [A] (verified)	1394
3.272.3 Rubi [A] (verified)	1395
3.272.4 Maple [A] (verified)	1396
3.272.5 Fricas [A] (verification not implemented)	1397
3.272.6 Sympy [A] (verification not implemented)	1397
3.272.7 Maxima [A] (verification not implemented)	1397
3.272.8 Giac [A] (verification not implemented)	1398
3.272.9 Mupad [B] (verification not implemented)	1398

3.272.1 Optimal result

Integrand size = 12, antiderivative size = 22

$$\int \frac{x}{2-2x+x^2} dx = -\arctan(1-x) + \frac{1}{2} \log(2-2x+x^2)$$

output `arctan(-1+x)+1/2*ln(x^2-2*x+2)`

3.272.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{x}{2-2x+x^2} dx = -\arctan(1-x) + \frac{1}{2} \log(2-2x+x^2)$$

input `Integrate[x/(2 - 2*x + x^2),x]`

output `-ArcTan[1 - x] + Log[2 - 2*x + x^2]/2`

3.272.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {1142, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x}{x^2 - 2x + 2} dx \\
 & \quad \downarrow \text{1142} \\
 & \int \frac{1}{x^2 - 2x + 2} dx + \frac{1}{2} \int -\frac{2(1-x)}{x^2 - 2x + 2} dx \\
 & \quad \downarrow \text{27} \\
 & \int \frac{1}{x^2 - 2x + 2} dx - \int \frac{1-x}{x^2 - 2x + 2} dx \\
 & \quad \downarrow \text{1082} \\
 & \int \frac{1}{-(1-x)^2 - 1} d(1-x) - \int \frac{1-x}{x^2 - 2x + 2} dx \\
 & \quad \downarrow \text{217} \\
 & - \int \frac{1-x}{x^2 - 2x + 2} dx - \arctan(1-x) \\
 & \quad \downarrow \text{1103} \\
 & \frac{1}{2} \log(x^2 - 2x + 2) - \arctan(1-x)
 \end{aligned}$$

input `Int[x/(2 - 2*x + x^2), x]`

output `-ArcTan[1 - x] + Log[2 - 2*x + x^2]/2`

3.272.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1142 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

3.272.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

method	result	size
default	$\arctan(-1+x) + \frac{\ln(x^2-2x+2)}{2}$	17
risch	$\arctan(-1+x) + \frac{\ln(x^2-2x+2)}{2}$	17
parallelrisch	$\frac{\ln(x-1-i)}{2} - \frac{i \ln(x-1-i)}{2} + \frac{\ln(x-1+i)}{2} + \frac{i \ln(x-1+i)}{2}$	36

input `int(x/(x^2-2*x+2), x, method=_RETURNVERBOSE)`

output `arctan(-1+x)+1/2*ln(x^2-2*x+2)`

3.272.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \frac{x}{2 - 2x + x^2} dx = \arctan(x - 1) + \frac{1}{2} \log(x^2 - 2x + 2)$$

input `integrate(x/(x^2-2*x+2),x, algorithm="fricas")`output `arctan(x - 1) + 1/2*log(x^2 - 2*x + 2)`**3.272.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.68

$$\int \frac{x}{2 - 2x + x^2} dx = \frac{\log(x^2 - 2x + 2)}{2} + \operatorname{atan}(x - 1)$$

input `integrate(x/(x**2-2*x+2),x)`output `log(x**2 - 2*x + 2)/2 + atan(x - 1)`**3.272.7 Maxima [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \frac{x}{2 - 2x + x^2} dx = \arctan(x - 1) + \frac{1}{2} \log(x^2 - 2x + 2)$$

input `integrate(x/(x^2-2*x+2),x, algorithm="maxima")`output `arctan(x - 1) + 1/2*log(x^2 - 2*x + 2)`

3.272.8 Giac [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \frac{x}{2 - 2x + x^2} dx = \arctan(x - 1) + \frac{1}{2} \log(x^2 - 2x + 2)$$

input `integrate(x/(x^2-2*x+2),x, algorithm="giac")`output `arctan(x - 1) + 1/2*log(x^2 - 2*x + 2)`**3.272.9 Mupad [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \frac{x}{2 - 2x + x^2} dx = \operatorname{atan}(x - 1) + \frac{\ln(x^2 - 2x + 2)}{2}$$

input `int(x/(x^2 - 2*x + 2),x)`output `atan(x - 1) + log(x^2 - 2*x + 2)/2`

3.273 $\int x \arcsin(x) dx$

3.273.1 Optimal result	1399
3.273.2 Mathematica [A] (verified)	1399
3.273.3 Rubi [A] (verified)	1400
3.273.4 Maple [A] (verified)	1401
3.273.5 Fricas [A] (verification not implemented)	1401
3.273.6 Sympy [A] (verification not implemented)	1402
3.273.7 Maxima [A] (verification not implemented)	1402
3.273.8 Giac [A] (verification not implemented)	1402
3.273.9 Mupad [B] (verification not implemented)	1403

3.273.1 Optimal result

Integrand size = 4, antiderivative size = 32

$$\int x \arcsin(x) dx = \frac{1}{4}x\sqrt{1-x^2} - \frac{\arcsin(x)}{4} + \frac{1}{2}x^2 \arcsin(x)$$

output `-1/4*arcsin(x)+1/2*x^2*arcsin(x)+1/4*x*(-x^2+1)^(1/2)`

3.273.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.88

$$\int x \arcsin(x) dx = \frac{1}{4} \left(x\sqrt{1-x^2} + (-1+2x^2) \arcsin(x) \right)$$

input `Integrate[x*ArcSin[x],x]`

output `(x*Sqrt[1 - x^2] + (-1 + 2*x^2)*ArcSin[x])/4`

3.273.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.16, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5138, 262, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \arcsin(x) dx \\
 & \quad \downarrow \text{5138} \\
 & \frac{1}{2}x^2 \arcsin(x) - \frac{1}{2} \int \frac{x^2}{\sqrt{1-x^2}} dx \\
 & \quad \downarrow \text{262} \\
 & \frac{1}{2} \left(\frac{1}{2}x\sqrt{1-x^2} - \frac{1}{2} \int \frac{1}{\sqrt{1-x^2}} dx \right) + \frac{1}{2}x^2 \arcsin(x) \\
 & \quad \downarrow \text{223} \\
 & \frac{1}{2}x^2 \arcsin(x) + \frac{1}{2} \left(\frac{1}{2}x\sqrt{1-x^2} - \frac{\arcsin(x)}{2} \right)
 \end{aligned}$$

input `Int[x*ArcSin[x],x]`

output `((x*Sqrt[1 - x^2])/2 - ArcSin[x]/2)/2 + (x^2*ArcSin[x])/2`

3.273.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 262 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

```
rule 5138 Int[((a_.) + ArcSin[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol]
:> Simp[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n
/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^(n - 1)/Sqrt[1 - c^2
*x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]
```

3.273.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.78

method	result	size
default	$-\frac{\arcsin(x)}{4} + \frac{x^2 \arcsin(x)}{2} + \frac{x\sqrt{-x^2+1}}{4}$	25
parts	$-\frac{\arcsin(x)}{4} + \frac{x^2 \arcsin(x)}{2} + \frac{x\sqrt{-x^2+1}}{4}$	25

```
input int(arcsin(x)*x,x,method=_RETURNVERBOSE)
```

```
output -1/4*arcsin(x)+1/2*x^2*arcsin(x)+1/4*x*(-x^2+1)^(1/2)
```

3.273.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75

$$\int x \arcsin(x) dx = \frac{1}{4} (2x^2 - 1) \arcsin(x) + \frac{1}{4} \sqrt{-x^2 + 1}x$$

```
input integrate(x*arcsin(x),x, algorithm="fricas")
```

```
output 1/4*(2*x^2 - 1)*arcsin(x) + 1/4*sqrt(-x^2 + 1)*x
```

3.273.6 Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75

$$\int x \arcsin(x) dx = \frac{x^2 \arcsin(x)}{2} + \frac{x\sqrt{1-x^2}}{4} - \frac{\arcsin(x)}{4}$$

input `integrate(x*asin(x),x)`output `x**2*asin(x)/2 + x*sqrt(1 - x**2)/4 - asin(x)/4`**3.273.7 Maxima [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75

$$\int x \arcsin(x) dx = \frac{1}{2} x^2 \arcsin(x) + \frac{1}{4} \sqrt{-x^2 + 1} x - \frac{1}{4} \arcsin(x)$$

input `integrate(x*arcsin(x),x, algorithm="maxima")`output `1/2*x^2*arcsin(x) + 1/4*sqrt(-x^2 + 1)*x - 1/4*arcsin(x)`**3.273.8 Giac [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

$$\int x \arcsin(x) dx = \frac{1}{2} (x^2 - 1) \arcsin(x) + \frac{1}{4} \sqrt{-x^2 + 1} x + \frac{1}{4} \arcsin(x)$$

input `integrate(x*arcsin(x),x, algorithm="giac")`output `1/2*(x^2 - 1)*arcsin(x) + 1/4*sqrt(-x^2 + 1)*x + 1/4*arcsin(x)`

3.273.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75

$$\int x \arcsin(x) dx = \frac{x \sqrt{1-x^2}}{4} + \frac{\arcsin(x) (2x^2 - 1)}{4}$$

input `int(x*asin(x),x)`

output `(x*(1 - x^2)^(1/2))/4 + (asin(x)*(2*x^2 - 1))/4`

3.274 $\int \frac{\sqrt{9-x^2}}{x} dx$

3.274.1 Optimal result	1404
3.274.2 Mathematica [A] (verified)	1404
3.274.3 Rubi [A] (verified)	1405
3.274.4 Maple [A] (verified)	1406
3.274.5 Fricas [A] (verification not implemented)	1407
3.274.6 Sympy [C] (verification not implemented)	1407
3.274.7 Maxima [A] (verification not implemented)	1407
3.274.8 Giac [A] (verification not implemented)	1408
3.274.9 Mupad [B] (verification not implemented)	1408

3.274.1 Optimal result

Integrand size = 15, antiderivative size = 30

$$\int \frac{\sqrt{9-x^2}}{x} dx = \sqrt{9-x^2} - 3\operatorname{arctanh}\left(\frac{\sqrt{9-x^2}}{3}\right)$$

output `-3*arctanh(1/3*(-x^2+9)^(1/2))+(-x^2+9)^(1/2)`

3.274.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{9-x^2}}{x} dx = \sqrt{9-x^2} - 3\operatorname{arctanh}\left(\frac{\sqrt{9-x^2}}{3}\right)$$

input `Integrate[Sqrt[9 - x^2]/x,x]`

output `Sqrt[9 - x^2] - 3*ArcTanh[Sqrt[9 - x^2]/3]`

3.274.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.20, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {243, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{9-x^2}}{x} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int \frac{\sqrt{9-x^2}}{x^2} dx^2 \\ & \quad \downarrow \text{60} \\ & \frac{1}{2} \left(9 \int \frac{1}{x^2 \sqrt{9-x^2}} dx^2 + 2\sqrt{9-x^2} \right) \\ & \quad \downarrow \text{73} \\ & \frac{1}{2} \left(2\sqrt{9-x^2} - 18 \int \frac{1}{9-x^4} d\sqrt{9-x^2} \right) \\ & \quad \downarrow \text{219} \\ & \frac{1}{2} \left(2\sqrt{9-x^2} - 6 \operatorname{arctanh} \left(\frac{\sqrt{9-x^2}}{3} \right) \right) \end{aligned}$$

input `Int[Sqrt[9 - x^2]/x,x]`

output `(2*Sqrt[9 - x^2] - 6*ArcTanh[Sqrt[9 - x^2]/3])/2`

3.274.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
 ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
 Q[a, 0] || LtQ[b, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int
 t[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I
 ntegerQ[(m - 1)/2]`

3.274.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.83

method	result	size
default	$\sqrt{-x^2 + 9} - 3 \operatorname{arctanh}\left(\frac{3}{\sqrt{-x^2 + 9}}\right)$	25
trager	$\sqrt{-x^2 + 9} - 3 \ln\left(\frac{\sqrt{-x^2 + 9} + 3}{x}\right)$	29
pseudoelliptic	$\sqrt{-x^2 + 9} - \frac{3 \ln(\sqrt{-x^2 + 9} + 3)}{2} + \frac{3 \ln(\sqrt{-x^2 + 9} - 3)}{2}$	39
meijerg	$-\frac{3 \left(-2(2 - 2 \ln(2) + 2 \ln(x) - 2 \ln(3) + i\pi) \sqrt{\pi} + 4\sqrt{\pi} - 4\sqrt{\pi} \sqrt{-\frac{x^2}{9} + 1} + 4\sqrt{\pi} \ln\left(\frac{1}{2} + \frac{\sqrt{-\frac{x^2}{9} + 1}}{2}\right) \right)}{4\sqrt{\pi}}$	68

input `int((-x^2+9)^(1/2)/x,x,method=_RETURNVERBOSE)`

output `(-x^2+9)^(1/2)-3*arctanh(3/(-x^2+9)^(1/2))`

3.274.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{9-x^2}}{x} dx = \sqrt{-x^2+9} + 3 \log\left(\frac{\sqrt{-x^2+9}-3}{x}\right)$$

input `integrate((-x^2+9)^(1/2)/x,x, algorithm="fracas")`

output `sqrt(-x^2 + 9) + 3*log((sqrt(-x^2 + 9) - 3)/x)`

3.274.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.78 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.20

$$\int \frac{\sqrt{9-x^2}}{x} dx = \begin{cases} i\sqrt{x^2-9} - 3\log(x) + \frac{3\log(x^2)}{2} + 3i\operatorname{asin}\left(\frac{3}{x}\right) & \text{for } |x^2| > 9 \\ \sqrt{9-x^2} + \frac{3\log(x^2)}{2} - 3\log\left(\sqrt{1-\frac{x^2}{9}}+1\right) & \text{otherwise} \end{cases}$$

input `integrate((-x**2+9)**(1/2)/x,x)`

output `Piecewise((I*sqrt(x**2 - 9) - 3*log(x) + 3*log(x**2)/2 + 3*I*asin(3/x), Abs(x**2) > 9), (sqrt(9 - x**2) + 3*log(x**2)/2 - 3*log(sqrt(1 - x**2/9) + 1), True))`

3.274.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.17

$$\int \frac{\sqrt{9-x^2}}{x} dx = \sqrt{-x^2+9} - 3 \log\left(\frac{6\sqrt{-x^2+9}}{|x|} + \frac{18}{|x|}\right)$$

input `integrate((-x^2+9)^(1/2)/x,x, algorithm="maxima")`

output `sqrt(-x^2 + 9) - 3*log(6*sqrt(-x^2 + 9)/abs(x) + 18/abs(x))`

3.274.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.33

$$\int \frac{\sqrt{9-x^2}}{x} dx = \sqrt{-x^2+9} - \frac{3}{2} \log(\sqrt{-x^2+9}+3) + \frac{3}{2} \log(-\sqrt{-x^2+9}+3)$$

input `integrate((-x^2+9)^(1/2)/x,x, algorithm="giac")`output `sqrt(-x^2 + 9) - 3/2*log(sqrt(-x^2 + 9) + 3) + 3/2*log(-sqrt(-x^2 + 9) + 3)`**3.274.9 Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{9-x^2}}{x} dx = 3 \ln \left(\sqrt{\frac{9}{x^2} - 1} - 3 \sqrt{\frac{1}{x^2}} \right) + \sqrt{9-x^2}$$

input `int((9 - x^2)^(1/2)/x,x)`output `3*log((9/x^2 - 1)^(1/2) - 3*(1/x^2)^(1/2)) + (9 - x^2)^(1/2)`

3.275 $\int \frac{x}{2+3x+x^2} dx$

3.275.1 Optimal result	1409
3.275.2 Mathematica [A] (verified)	1409
3.275.3 Rubi [A] (verified)	1410
3.275.4 Maple [A] (verified)	1411
3.275.5 Fricas [A] (verification not implemented)	1411
3.275.6 Sympy [A] (verification not implemented)	1411
3.275.7 Maxima [A] (verification not implemented)	1412
3.275.8 Giac [A] (verification not implemented)	1412
3.275.9 Mupad [B] (verification not implemented)	1412

3.275.1 Optimal result

Integrand size = 12, antiderivative size = 13

$$\int \frac{x}{2+3x+x^2} dx = -\log(1+x) + 2\log(2+x)$$

output `-ln(1+x)+2*ln(2+x)`

3.275.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{x}{2+3x+x^2} dx = -\log(1+x) + 2\log(2+x)$$

input `Integrate[x/(2 + 3*x + x^2),x]`

output `-Log[1 + x] + 2*Log[2 + x]`

3.275.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{x^2 + 3x + 2} dx$$

↓ 1141

$$\int \left(\frac{2}{x+2} + \frac{1}{-x-1} \right) dx$$

↓ 2009

$$2 \log(x+2) - \log(x+1)$$

input `Int[x/(2 + 3*x + x^2),x]`

output `-Log[1 + x] + 2*Log[2 + x]`

3.275.3.1 Defintions of rubi rules used

rule 1141 `Int[((d_.) + (e_.)*(x_))^(m_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x], x] /; EqQ[p, -1] || !FractionalPowerFactorQ[q] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p, 0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.275.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.08

method	result	size
default	$-\ln(1+x) + 2\ln(2+x)$	14
norman	$-\ln(1+x) + 2\ln(2+x)$	14
risch	$-\ln(1+x) + 2\ln(2+x)$	14
parallelrisch	$-\ln(1+x) + 2\ln(2+x)$	14

input `int(x/(x^2+3*x+2),x,method=_RETURNVERBOSE)`output `-ln(1+x)+2*ln(2+x)`**3.275.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{x}{2+3x+x^2} dx = 2 \log(x+2) - \log(x+1)$$

input `integrate(x/(x^2+3*x+2),x, algorithm="fracas")`output `2*log(x + 2) - log(x + 1)`**3.275.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \frac{x}{2+3x+x^2} dx = -\log(x+1) + 2\log(x+2)$$

input `integrate(x/(x**2+3*x+2),x)`output `-log(x + 1) + 2*log(x + 2)`

3.275.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{x}{2 + 3x + x^2} dx = 2 \log(x + 2) - \log(x + 1)$$

input `integrate(x/(x^2+3*x+2),x, algorithm="maxima")`output `2*log(x + 2) - log(x + 1)`**3.275.8 Giac [A] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{x}{2 + 3x + x^2} dx = 2 \log(|x + 2|) - \log(|x + 1|)$$

input `integrate(x/(x^2+3*x+2),x, algorithm="giac")`output `2*log(abs(x + 2)) - log(abs(x + 1))`**3.275.9 Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{x}{2 + 3x + x^2} dx = 2 \ln(x + 2) - \ln(x + 1)$$

input `int(x/(3*x + x^2 + 2),x)`output `2*log(x + 2) - log(x + 1)`

3.276 $\int x^2 \cosh(x) dx$

3.276.1 Optimal result	1413
3.276.2 Mathematica [A] (verified)	1413
3.276.3 Rubi [C] (verified)	1414
3.276.4 Maple [A] (verified)	1415
3.276.5 Fricas [A] (verification not implemented)	1416
3.276.6 Sympy [A] (verification not implemented)	1416
3.276.7 Maxima [B] (verification not implemented)	1416
3.276.8 Giac [A] (verification not implemented)	1417
3.276.9 Mupad [B] (verification not implemented)	1417

3.276.1 Optimal result

Integrand size = 6, antiderivative size = 16

$$\int x^2 \cosh(x) dx = -2x \cosh(x) + 2 \sinh(x) + x^2 \sinh(x)$$

output `-2*x*cosh(x)+2*sinh(x)+x^2*sinh(x)`

3.276.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int x^2 \cosh(x) dx = -2x \cosh(x) + (2 + x^2) \sinh(x)$$

input `Integrate[x^2*Cosh[x],x]`

output `-2*x*Cosh[x] + (2 + x^2)*Sinh[x]`

3.276.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.24 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.56, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.333$, Rules used = {3042, 3777, 26, 3042, 26, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \cosh(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x^2 \sin\left(\frac{\pi}{2} + ix\right) dx \\
 & \quad \downarrow \text{3777} \\
 & x^2 \sinh(x) - 2i \int -ix \sinh(x) dx \\
 & \quad \downarrow \text{26} \\
 & x^2 \sinh(x) - 2 \int x \sinh(x) dx \\
 & \quad \downarrow \text{3042} \\
 & x^2 \sinh(x) - 2 \int -ix \sin(ix) dx \\
 & \quad \downarrow \text{26} \\
 & x^2 \sinh(x) + 2i \int x \sin(ix) dx \\
 & \quad \downarrow \text{3777} \\
 & x^2 \sinh(x) + 2i(ix \cosh(x) - i \int \cosh(x) dx) \\
 & \quad \downarrow \text{3042} \\
 & x^2 \sinh(x) + 2i\left(ix \cosh(x) - i \int \sin\left(ix + \frac{\pi}{2}\right) dx\right) \\
 & \quad \downarrow \text{3117} \\
 & x^2 \sinh(x) + 2i(ix \cosh(x) - i \sinh(x))
 \end{aligned}$$

input `Int[x^2*Cosh[x],x]`

output `(2*I)*(I*x*Cosh[x] - I*Sinh[x]) + x^2*Sinh[x]`

3.276.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

3.276.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

method	result	size
default	$-2x \cosh(x) + 2 \sinh(x) + x^2 \sinh(x)$	17
parallelrisch	$-2x \cosh(x) + 2 \sinh(x) + x^2 \sinh(x)$	17
parts	$-2x \cosh(x) + 2 \sinh(x) + x^2 \sinh(x)$	17
risch	$(1 - x + \frac{1}{2}x^2) e^x + (-1 - x - \frac{1}{2}x^2) e^{-x}$	30
meijerg	$4i\sqrt{\pi} \left(\frac{ix \cosh(x)}{2\sqrt{\pi}} - \frac{i\left(\frac{3x^2}{2}+3\right) \sinh(x)}{6\sqrt{\pi}} \right)$	32

input `int(x^2*cosh(x),x,method=_RETURNVERBOSE)`

output `-2*x*cosh(x)+2*sinh(x)+x^2*sinh(x)`

3.276.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int x^2 \cosh(x) dx = -2x \cosh(x) + (x^2 + 2) \sinh(x)$$

input `integrate(x^2*cosh(x),x, algorithm="fricas")`

output `-2*x*cosh(x) + (x^2 + 2)*sinh(x)`

3.276.6 Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

$$\int x^2 \cosh(x) dx = x^2 \sinh(x) - 2x \cosh(x) + 2 \sinh(x)$$

input `integrate(x**2*cosh(x),x)`

output `x**2*sinh(x) - 2*x*cosh(x) + 2*sinh(x)`

3.276.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 44 vs. $2(16) = 32$.

Time = 0.20 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.75

$$\int x^2 \cosh(x) dx = \frac{1}{3} x^3 \cosh(x) - \frac{1}{6} (x^3 + 3x^2 + 6x + 6)e^{(-x)} - \frac{1}{6} (x^3 - 3x^2 + 6x - 6)e^x$$

input `integrate(x^2*cosh(x),x, algorithm="maxima")`

output `1/3*x^3*cosh(x) - 1/6*(x^3 + 3*x^2 + 6*x + 6)*e^(-x) - 1/6*(x^3 - 3*x^2 + 6*x - 6)*e^x`

3.276.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.69

$$\int x^2 \cosh(x) dx = -\frac{1}{2} (x^2 + 2x + 2)e^{(-x)} + \frac{1}{2} (x^2 - 2x + 2)e^x$$

input `integrate(x^2*cosh(x),x, algorithm="giac")`

output `-1/2*(x^2 + 2*x + 2)*e^(-x) + 1/2*(x^2 - 2*x + 2)*e^x`

3.276.9 Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int x^2 \cosh(x) dx = 2 \sinh(x) + x^2 \sinh(x) - 2x \cosh(x)$$

input `int(x^2*cosh(x),x)`

output `2*sinh(x) + x^2*sinh(x) - 2*x*cosh(x)`

$$3.277 \quad \int \frac{1+x+x^3}{4x+2x^2+x^4} dx$$

3.277.1 Optimal result	1418
3.277.2 Mathematica [A] (verified)	1418
3.277.3 Rubi [A] (verified)	1419
3.277.4 Maple [A] (verified)	1419
3.277.5 Fricas [A] (verification not implemented)	1420
3.277.6 Sympy [A] (verification not implemented)	1420
3.277.7 Maxima [A] (verification not implemented)	1420
3.277.8 Giac [A] (verification not implemented)	1421
3.277.9 Mupad [B] (verification not implemented)	1421

3.277.1 Optimal result

Integrand size = 21, antiderivative size = 17

$$\int \frac{1+x+x^3}{4x+2x^2+x^4} dx = \frac{1}{4} \log(4x+2x^2+x^4)$$

output `1/4*ln(x^4+2*x^2+4*x)`

3.277.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \frac{1+x+x^3}{4x+2x^2+x^4} dx = \frac{\log(x)}{4} + \frac{1}{4} \log(4+2x+x^3)$$

input `Integrate[(1 + x + x^3)/(4*x + 2*x^2 + x^4),x]`

output `Log[x]/4 + Log[4 + 2*x + x^3]/4`

3.277.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2020}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 + x + 1}{x^4 + 2x^2 + 4x} dx$$

↓ 2020

$$\frac{1}{4} \log(x^4 + 2x^2 + 4x)$$

input `Int[(1 + x + x^3)/(4*x + 2*x^2 + x^4),x]`

output `Log[4*x + 2*x^2 + x^4]/4`

3.277.3.1 Defintions of rubi rules used

rule 2020 `Int[(Pp_)/(Qq_), x_Symbol] := With[{p = Expon[Pp, x], q = Expon[Qq, x]}, Simp[Coeff[Pp, x, p]*(Log[RemoveContent[Qq, x]]/(q*Coeff[Qq, x, q])), x] /; EqQ[p, q - 1] && EqQ[Pp, Simplify[(Coeff[Pp, x, p]/(q*Coeff[Qq, x, q]))*D[Qq, x]]] /; PolyQ[Pp, x] && PolyQ[Qq, x]`

3.277.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{\ln(x(x^3+2x+4))}{4}$	14
risch	$\frac{\ln(x^4+2x^2+4x)}{4}$	16
norman	$\frac{\ln(x)}{4} + \frac{\ln(x^3+2x+4)}{4}$	17
parallelrisch	$\frac{\ln(x)}{4} + \frac{\ln(x^3+2x+4)}{4}$	17

input `int((x^3+x+1)/(x^4+2*x^2+4*x),x,method=_RETURNVERBOSE)`

3.277. $\int \frac{1+x+x^3}{4x+2x^2+x^4} dx$

output `1/4*ln(x*(x^3+2*x+4))`

3.277.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{1+x+x^3}{4x+2x^2+x^4} dx = \frac{1}{4} \log(x^4+2x^2+4x)$$

input `integrate((x^3+x+1)/(x^4+2*x^2+4*x),x, algorithm="fricas")`

output `1/4*log(x^4 + 2*x^2 + 4*x)`

3.277.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \frac{1+x+x^3}{4x+2x^2+x^4} dx = \frac{\log(x^4+2x^2+4x)}{4}$$

input `integrate((x**3+x+1)/(x**4+2*x**2+4*x),x)`

output `log(x**4 + 2*x**2 + 4*x)/4`

3.277.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{1+x+x^3}{4x+2x^2+x^4} dx = \frac{1}{4} \log(x^4+2x^2+4x)$$

input `integrate((x^3+x+1)/(x^4+2*x^2+4*x),x, algorithm="maxima")`

output `1/4*log(x^4 + 2*x^2 + 4*x)`

3.277.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

$$\int \frac{1+x+x^3}{4x+2x^2+x^4} dx = \frac{1}{4} \log \left(4 \left| \frac{1}{4} x^4 + \frac{1}{2} x^2 + x \right| \right)$$

input `integrate((x^3+x+1)/(x^4+2*x^2+4*x),x, algorithm="giac")`output `1/4*log(4*abs(1/4*x^4 + 1/2*x^2 + x))`**3.277.9 Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{1+x+x^3}{4x+2x^2+x^4} dx = \frac{\ln(x(x^3+2x+4))}{4}$$

input `int((x + x^3 + 1)/(4*x + 2*x^2 + x^4),x)`output `log(x*(2*x + x^3 + 4))/4`

$$3.278 \quad \int \frac{\cos(x)}{1+\sin^2(x)} dx$$

3.278.1 Optimal result	1422
3.278.2 Mathematica [A] (verified)	1422
3.278.3 Rubi [A] (verified)	1423
3.278.4 Maple [A] (verified)	1424
3.278.5 Fricas [A] (verification not implemented)	1424
3.278.6 Sympy [A] (verification not implemented)	1425
3.278.7 Maxima [A] (verification not implemented)	1425
3.278.8 Giac [A] (verification not implemented)	1425
3.278.9 Mupad [B] (verification not implemented)	1426

3.278.1 Optimal result

Integrand size = 11, antiderivative size = 3

$$\int \frac{\cos(x)}{1+\sin^2(x)} dx = \arctan(\sin(x))$$

output `arctan(sin(x))`

3.278.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{1+\sin^2(x)} dx = \arctan(\sin(x))$$

input `Integrate[Cos[x]/(1 + Sin[x]^2),x]`

output `ArcTan[Sin[x]]`

3.278.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 3669, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(x)}{\sin^2(x) + 1} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\cos(x)}{\sin(x)^2 + 1} dx \\ & \quad \downarrow \text{3669} \\ & \int \frac{1}{\sin^2(x) + 1} d\sin(x) \\ & \quad \downarrow \text{216} \\ & \arctan(\sin(x)) \end{aligned}$$

input `Int[Cos[x]/(1 + Sin[x]^2),x]`

output `ArcTan[Sin[x]]`

3.278.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`


```
rule 3669 Int[cos[(e_.) + (f_.)*(x_.)]^(m_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)]^2)^(p_.), x_Symbol] :> With[{ff = FreeFactors[Sin[e + f*x], x]}, Simp[ff/f Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b*ff^2*x^2)^p, x], x, Sin[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

3.278.4 Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
derivativedivides	$\arctan(\sin(x))$	4
default	$\arctan(\sin(x))$	4
parallelrisc	$-\frac{i\left(-\ln\left(-\frac{2i(\sin(x)+i)}{\cos(x)+1}\right)+\ln\left(\frac{2+2i\sin(x)}{\cos(x)+1}\right)\right)}{2}$	37
risc	$\frac{i\ln(e^{2ix}-2e^{ix}-1)}{2} - \frac{i\ln(e^{2ix}+2e^{ix}-1)}{2}$	38

```
input int(cos(x)/(1+sin(x)^2),x,method=_RETURNVERBOSE)
```

```
output arctan(sin(x))
```

3.278.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{1 + \sin^2(x)} dx = \arctan(\sin(x))$$

```
input integrate(cos(x)/(1+sin(x)^2),x, algorithm="fricas")
```

```
output arctan(sin(x))
```

3.278.6 Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{1 + \sin^2(x)} dx = \operatorname{atan}(\sin(x))$$

input `integrate(cos(x)/(1+sin(x)**2),x)`output `atan(sin(x))`**3.278.7 Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{1 + \sin^2(x)} dx = \operatorname{arctan}(\sin(x))$$

input `integrate(cos(x)/(1+sin(x)^2),x, algorithm="maxima")`output `arctan(sin(x))`**3.278.8 Giac [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{1 + \sin^2(x)} dx = \operatorname{arctan}(\sin(x))$$

input `integrate(cos(x)/(1+sin(x)^2),x, algorithm="giac")`output `arctan(sin(x))`

3.278.9 Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{1 + \sin^2(x)} dx = \text{atan}(\sin(x))$$

input `int(cos(x)/(sin(x)^2 + 1),x)`

output `atan(sin(x))`

3.279 $\int \cos(\sqrt{x}) dx$

3.279.1 Optimal result	1427
3.279.2 Mathematica [A] (verified)	1427
3.279.3 Rubi [A] (verified)	1428
3.279.4 Maple [A] (verified)	1429
3.279.5 Fricas [A] (verification not implemented)	1430
3.279.6 Sympy [A] (verification not implemented)	1430
3.279.7 Maxima [A] (verification not implemented)	1430
3.279.8 Giac [A] (verification not implemented)	1431
3.279.9 Mupad [B] (verification not implemented)	1431

3.279.1 Optimal result

Integrand size = 6, antiderivative size = 22

$$\int \cos(\sqrt{x}) dx = 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$$

output `2*cos(x^(1/2))+2*sin(x^(1/2))*x^(1/2)`

3.279.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \cos(\sqrt{x}) dx = 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$$

input `Integrate[Cos[Sqrt[x]],x]`

output `2*Cos[Sqrt[x]] + 2*Sqrt[x]*Sin[Sqrt[x]]`

3.279.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.95, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3843, 3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(\sqrt{x}) \, dx \\
 & \quad \downarrow \text{3843} \\
 & 2 \int \sqrt{x} \cos(\sqrt{x}) \, d\sqrt{x} \\
 & \quad \downarrow \text{3042} \\
 & 2 \int \sqrt{x} \sin\left(\sqrt{x} + \frac{\pi}{2}\right) \, d\sqrt{x} \\
 & \quad \downarrow \text{3777} \\
 & 2 \left(\int -\sin(\sqrt{x}) \, d\sqrt{x} + \sqrt{x} \sin(\sqrt{x}) \right) \\
 & \quad \downarrow \text{25} \\
 & 2 \left(\sqrt{x} \sin(\sqrt{x}) - \int \sin(\sqrt{x}) \, d\sqrt{x} \right) \\
 & \quad \downarrow \text{3042} \\
 & 2 \left(\sqrt{x} \sin(\sqrt{x}) - \int \sin(\sqrt{x}) \, d\sqrt{x} \right) \\
 & \quad \downarrow \text{3118} \\
 & 2(\sqrt{x} \sin(\sqrt{x}) + \cos(\sqrt{x}))
 \end{aligned}$$

input `Int[Cos[Sqrt[x]], x]`

output `2*(Cos[Sqrt[x]] + Sqrt[x]*Sin[Sqrt[x]])`

3.279.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`
- rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`
- rule 3843 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)])*(b_.))^(p_.), x_Symbol] := Simp[1/(n*f) Subst[Int[x^(1/n - 1)*(a + b*Cos[c + d*x])^p, x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]`

3.279.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

method	result	size
derivativedivides	$2 \cos(\sqrt{x}) + 2 \sin(\sqrt{x}) \sqrt{x}$	17
default	$2 \cos(\sqrt{x}) + 2 \sin(\sqrt{x}) \sqrt{x}$	17
meijerg	$4\sqrt{\pi} \left(-\frac{1}{2\sqrt{\pi}} + \frac{\cos(\sqrt{x})}{2\sqrt{\pi}} + \frac{\sqrt{x} \sin(\sqrt{x})}{2\sqrt{\pi}} \right)$	33

input `int(cos(x^(1/2)), x, method=_RETURNVERBOSE)`

output `2*cos(x^(1/2))+2*sin(x^(1/2))*x^(1/2)`

3.279.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x^(1/2)),x, algorithm="fricas")`output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`**3.279.6 Sympy [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x**(1/2)),x)`output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`**3.279.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x^(1/2)),x, algorithm="maxima")`output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`

3.279.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x^(1/2)),x, algorithm="giac")`

output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`

3.279.9 Mupad [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$$

input `int(cos(x^(1/2)),x)`

output `2*cos(x^(1/2)) + 2*x^(1/2)*sin(x^(1/2))`

3.280 $\int \sin(\pi x) dx$

3.280.1 Optimal result	1432
3.280.2 Mathematica [A] (verified)	1432
3.280.3 Rubi [A] (verified)	1433
3.280.4 Maple [A] (verified)	1434
3.280.5 Fricas [A] (verification not implemented)	1434
3.280.6 Sympy [A] (verification not implemented)	1434
3.280.7 Maxima [A] (verification not implemented)	1435
3.280.8 Giac [A] (verification not implemented)	1435
3.280.9 Mupad [B] (verification not implemented)	1435

3.280.1 Optimal result

Integrand size = 4, antiderivative size = 9

$$\int \sin(\pi x) dx = -\frac{\cos(\pi x)}{\pi}$$

output `-cos(Pi*x)/Pi`

3.280.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \sin(\pi x) dx = -\frac{\cos(\pi x)}{\pi}$$

input `Integrate[Sin[Pi*x],x]`

output `-(Cos[Pi*x]/Pi)`

3.280.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sin(\pi x) dx \\ \downarrow \text{3042} \\ \int \sin(\pi x) dx \\ \downarrow \text{3118} \\ -\frac{\cos(\pi x)}{\pi} \end{array}$$

input `Int[Sin[Pi*x],x]`

output `-(Cos[Pi*x]/Pi)`

3.280.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

3.280.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.11

method	result	size
derivativedivides	$-\frac{\cos(\pi x)}{\pi}$	10
default	$-\frac{\cos(\pi x)}{\pi}$	10
risch	$-\frac{\cos(\pi x)}{\pi}$	10
parallelrisch	$\frac{-\cos(\pi x)-1}{\pi}$	13
norman	$-\frac{2}{\pi(1+\tan^2(\frac{\pi x}{2}))}$	17
meijerg	$\frac{\frac{1}{\sqrt{\pi}} - \frac{\cos(\pi x)}{\sqrt{\pi}}}{\sqrt{\pi}}$	18

input `int(sin(Pi*x),x,method=_RETURNVERBOSE)`output `-cos(Pi*x)/Pi`**3.280.5 Fricas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \sin(\pi x) dx = -\frac{\cos(\pi x)}{\pi}$$

input `integrate(sin(pi*x),x, algorithm="fricas")`output `-cos(pi*x)/pi`**3.280.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \sin(\pi x) dx = -\frac{\cos(\pi x)}{\pi}$$

input `integrate(sin(pi*x),x)`

output `-cos(pi*x)/pi`

3.280.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \sin(\pi x) dx = -\frac{\cos(\pi x)}{\pi}$$

input `integrate(sin(pi*x),x, algorithm="maxima")`

output `-cos(pi*x)/pi`

3.280.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \sin(\pi x) dx = -\frac{\cos(\pi x)}{\pi}$$

input `integrate(sin(pi*x),x, algorithm="giac")`

output `-cos(pi*x)/pi`

3.280.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \sin(\pi x) dx = -\frac{\cos(\Pi x)}{\Pi}$$

input `int(sin(Pi*x),x)`

output `-cos(Pi*x)/Pi`

3.281 $\int \frac{e^{2x}}{1+e^x} dx$

3.281.1 Optimal result	1436
3.281.2 Mathematica [A] (verified)	1436
3.281.3 Rubi [A] (verified)	1437
3.281.4 Maple [A] (verified)	1438
3.281.5 Fricas [A] (verification not implemented)	1438
3.281.6 Sympy [A] (verification not implemented)	1439
3.281.7 Maxima [A] (verification not implemented)	1439
3.281.8 Giac [A] (verification not implemented)	1439
3.281.9 Mupad [B] (verification not implemented)	1440

3.281.1 Optimal result

Integrand size = 13, antiderivative size = 12

$$\int \frac{e^{2x}}{1+e^x} dx = e^x - \log(1+e^x)$$

output `exp(x)-ln(1+exp(x))`

3.281.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{e^{2x}}{1+e^x} dx = e^x - \log(1+e^x)$$

input `Integrate[E^(2*x)/(1 + E^x), x]`

output `E^x - Log[1 + E^x]`

3.281.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {2678, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{e^{2x}}{e^x + 1} dx \\ & \quad \downarrow \text{2678} \\ & \int \frac{e^x}{e^x + 1} de^x \\ & \quad \downarrow \text{49} \\ & \int \left(\frac{1}{-e^x - 1} + 1 \right) de^x \\ & \quad \downarrow \text{2009} \\ & e^x - \log(e^x + 1) \end{aligned}$$

input `Int[E^(2*x)/(1 + E^x),x]`

output `E^x - Log[1 + E^x]`

3.281.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 2678 Int[((a_) + (b_)*(F_)^((e_)*((c_) + (d_)*(x_)))^(p_)*(G_)^((h_)*((f_
.) + (g_)*(x_))), x_Symbol] :> With[{m = FullSimplify[g*h*(Log[G]/(d*e*Log
[F]))]}, Simp[Denominator[m]*(G^(f*h - c*g*(h/d))/(d*e*Log[F])) Subst[Int
[x^(Numerator[m] - 1)*(a + b*x^Denominator[m])^p, x], x, F^(e*((c + d*x)/De
nominator[m]))], x] /; LeQ[m, -1] || GeQ[m, 1]] /; FreeQ[{F, G, a, b, c, d,
e, f, g, h, p}, x]
```

3.281.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
default	$e^x - \ln(1 + e^x)$	11
norman	$e^x - \ln(1 + e^x)$	11
risch	$e^x - \ln(1 + e^x)$	11

```
input int(exp(2*x)/(1+exp(x)),x,method=_RETURNVERBOSE)
```

```
output exp(x)-ln(1+exp(x))
```

3.281.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{e^{2x}}{1+e^x} dx = e^x - \log(e^x + 1)$$

```
input integrate(exp(2*x)/(1+exp(x)),x, algorithm="fracas")
```

```
output e^x - log(e^x + 1)
```

3.281.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{e^{2x}}{1+e^x} dx = e^x - \log(e^x + 1)$$

input `integrate(exp(2*x)/(1+exp(x)),x)`output `exp(x) - log(exp(x) + 1)`**3.281.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{e^{2x}}{1+e^x} dx = e^x - \log(e^x + 1)$$

input `integrate(exp(2*x)/(1+exp(x)),x, algorithm="maxima")`output `e^x - log(e^x + 1)`**3.281.8 Giac [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{e^{2x}}{1+e^x} dx = e^x - \log(e^x + 1)$$

input `integrate(exp(2*x)/(1+exp(x)),x, algorithm="giac")`output `e^x - log(e^x + 1)`

3.281.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{e^{2x}}{1+e^x} dx = e^x - \ln(e^x + 1)$$

input `int(exp(2*x)/(exp(x) + 1),x)`

output `exp(x) - log(exp(x) + 1)`

3.282 $\int e^{3x} \cos(5x) dx$

3.282.1 Optimal result	1441
3.282.2 Mathematica [A] (verified)	1441
3.282.3 Rubi [A] (verified)	1442
3.282.4 Maple [A] (verified)	1442
3.282.5 Fricas [A] (verification not implemented)	1443
3.282.6 Sympy [A] (verification not implemented)	1443
3.282.7 Maxima [A] (verification not implemented)	1443
3.282.8 Giac [A] (verification not implemented)	1444
3.282.9 Mupad [B] (verification not implemented)	1444

3.282.1 Optimal result

Integrand size = 10, antiderivative size = 27

$$\int e^{3x} \cos(5x) dx = \frac{3}{34} e^{3x} \cos(5x) + \frac{5}{34} e^{3x} \sin(5x)$$

output `3/34*exp(3*x)*cos(5*x)+5/34*exp(3*x)*sin(5*x)`

3.282.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int e^{3x} \cos(5x) dx = \frac{1}{34} e^{3x} (3 \cos(5x) + 5 \sin(5x))$$

input `Integrate[E^(3*x)*Cos[5*x],x]`

output `(E^(3*x)*(3*Cos[5*x] + 5*Sin[5*x]))/34`

3.282.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {4933}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{3x} \cos(5x) dx$$

↓ 4933

$$\frac{5}{34} e^{3x} \sin(5x) + \frac{3}{34} e^{3x} \cos(5x)$$

input `Int[E^(3*x)*Cos[5*x],x]`

output `(3*E^(3*x)*Cos[5*x])/34 + (5*E^(3*x)*Sin[5*x])/34`

3.282.3.1 Defintions of rubi rules used

rule 4933 `Int[Cos[(d_.) + (e_.)*(x_.)]*(F_)^((c_.)*((a_.) + (b_.)*(x_))), x_Symbol] :=
Simp[b*c*Log[F]*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x
] + Simp[e*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; F
reeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]`

3.282.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result	size
parallelrisch	$\frac{e^{3x}(3 \cos(5x) + 5 \sin(5x))}{34}$	20
default	$\frac{3 e^{3x} \cos(5x)}{34} + \frac{5 e^{3x} \sin(5x)}{34}$	22
risch	$\frac{3 e^{(3+5i)x}}{68} - \frac{5 i e^{(3+5i)x}}{68} + \frac{3 e^{(3-5i)x}}{68} + \frac{5 i e^{(3-5i)x}}{68}$	36
norman	$\frac{5 e^{3x} \tan\left(\frac{5x}{2}\right)}{17} - \frac{3 e^{3x} \left(\tan^2\left(\frac{5x}{2}\right)\right)}{34} + \frac{3 e^{3x}}{34}$ $\frac{1}{1 + \tan^2\left(\frac{5x}{2}\right)}$	41

input `int(exp(3*x)*cos(5*x),x,method=_RETURNVERBOSE)`

output `1/34*exp(3*x)*(3*cos(5*x)+5*sin(5*x))`

3.282.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int e^{3x} \cos(5x) dx = \frac{3}{34} \cos(5x) e^{(3x)} + \frac{5}{34} e^{(3x)} \sin(5x)$$

input `integrate(exp(3*x)*cos(5*x),x, algorithm="fricas")`

output `3/34*cos(5*x)*e^(3*x) + 5/34*e^(3*x)*sin(5*x)`

3.282.6 Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int e^{3x} \cos(5x) dx = \frac{5e^{3x} \sin(5x)}{34} + \frac{3e^{3x} \cos(5x)}{34}$$

input `integrate(exp(3*x)*cos(5*x),x)`

output `5*exp(3*x)*sin(5*x)/34 + 3*exp(3*x)*cos(5*x)/34`

3.282.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^{3x} \cos(5x) dx = \frac{1}{34} (3 \cos(5x) + 5 \sin(5x)) e^{(3x)}$$

input `integrate(exp(3*x)*cos(5*x),x, algorithm="maxima")`

output `1/34*(3*cos(5*x) + 5*sin(5*x))*e^(3*x)`

3.282.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^{3x} \cos(5x) dx = \frac{1}{34} (3 \cos(5x) + 5 \sin(5x))e^{(3x)}$$

input `integrate(exp(3*x)*cos(5*x),x, algorithm="giac")`

output `1/34*(3*cos(5*x) + 5*sin(5*x))*e^(3*x)`

3.282.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^{3x} \cos(5x) dx = \frac{e^{3x} (3 \cos(5x) + 5 \sin(5x))}{34}$$

input `int(cos(5*x)*exp(3*x),x)`

output `(exp(3*x)*(3*cos(5*x) + 5*sin(5*x)))/34`

3.283 $\int \cos(3x) \cos(5x) dx$

3.283.1 Optimal result	1445
3.283.2 Mathematica [A] (verified)	1445
3.283.3 Rubi [A] (verified)	1446
3.283.4 Maple [A] (verified)	1447
3.283.5 Fricas [A] (verification not implemented)	1447
3.283.6 Sympy [B] (verification not implemented)	1447
3.283.7 Maxima [A] (verification not implemented)	1448
3.283.8 Giac [A] (verification not implemented)	1448
3.283.9 Mupad [B] (verification not implemented)	1448

3.283.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \cos(3x) \cos(5x) dx = \frac{1}{4} \sin(2x) + \frac{1}{16} \sin(8x)$$

output `1/4*sin(2*x)+1/16*sin(8*x)`

3.283.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cos(3x) \cos(5x) dx = \frac{1}{4} \sin(2x) + \frac{1}{16} \sin(8x)$$

input `Integrate[Cos[3*x]*Cos[5*x],x]`

output `Sin[2*x]/4 + Sin[8*x]/16`

3.283.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4771}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \cos(3x) \cos(5x) dx \\ \downarrow \text{3042} \\ \int \cos(3x) \cos(5x) dx \\ \downarrow \text{4771} \\ \frac{1}{4} \sin(2x) + \frac{1}{16} \sin(8x) \end{array}$$

input `Int[Cos[3*x]*Cos[5*x],x]`

output `Sin[2*x]/4 + Sin[8*x]/16`

3.283.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4771 `Int[cos[(a_.) + (b_.)*(x_)]*cos[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[a - c + (b - d)*x]/(2*(b - d)), x] + Simp[Sin[a + c + (b + d)*x]/(2*(b + d)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

3.283.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{\sin(2x)}{4} + \frac{\sin(8x)}{16}$	14
risch	$\frac{\sin(2x)}{4} + \frac{\sin(8x)}{16}$	14
parallelrisc	$\frac{\sin(2x)}{4} + \frac{\sin(8x)}{16}$	14
norman	$\frac{3 \tan\left(\frac{3x}{2}\right) \left(\tan^2\left(\frac{5x}{2}\right)\right) - 5 \left(\tan^2\left(\frac{3x}{2}\right)\right) \tan\left(\frac{5x}{2}\right) - 3 \tan\left(\frac{3x}{2}\right) + 5 \tan\left(\frac{5x}{2}\right)}{(1+\tan^2\left(\frac{3x}{2}\right))(1+\tan^2\left(\frac{5x}{2}\right))}$	59

input `int(cos(3*x)*cos(5*x),x,method=_RETURNVERBOSE)`output `1/4*sin(2*x)+1/16*sin(8*x)`**3.283.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.29

$$\int \cos(3x) \cos(5x) dx = (8 \cos(x)^7 - 12 \cos(x)^5 + 5 \cos(x)^3) \sin(x)$$

input `integrate(cos(3*x)*cos(5*x),x, algorithm="fracas")`output `(8*cos(x)^7 - 12*cos(x)^5 + 5*cos(x)^3)*sin(x)`**3.283.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. 2(12) = 24.

Time = 0.14 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.53

$$\int \cos(3x) \cos(5x) dx = -\frac{3 \sin(3x) \cos(5x)}{16} + \frac{5 \sin(5x) \cos(3x)}{16}$$

input `integrate(cos(3*x)*cos(5*x),x)`output `-3*sin(3*x)*cos(5*x)/16 + 5*sin(5*x)*cos(3*x)/16`

3.283.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(3x) \cos(5x) dx = \frac{1}{16} \sin(8x) + \frac{1}{4} \sin(2x)$$

input `integrate(cos(3*x)*cos(5*x),x, algorithm="maxima")`output `1/16*sin(8*x) + 1/4*sin(2*x)`**3.283.8 Giac [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(3x) \cos(5x) dx = \frac{1}{16} \sin(8x) + \frac{1}{4} \sin(2x)$$

input `integrate(cos(3*x)*cos(5*x),x, algorithm="giac")`output `1/16*sin(8*x) + 1/4*sin(2*x)`**3.283.9 Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(3x) \cos(5x) dx = \frac{\sin(2x)}{4} + \frac{\sin(8x)}{16}$$

input `int(cos(3*x)*cos(5*x),x)`output `sin(2*x)/4 + sin(8*x)/16`

3.284 $\int \frac{1}{1+x+x^2+x^3} dx$

3.284.1 Optimal result	1449
3.284.2 Mathematica [A] (verified)	1449
3.284.3 Rubi [A] (verified)	1450
3.284.4 Maple [A] (verified)	1451
3.284.5 Fricas [A] (verification not implemented)	1451
3.284.6 Sympy [A] (verification not implemented)	1451
3.284.7 Maxima [A] (verification not implemented)	1452
3.284.8 Giac [A] (verification not implemented)	1452
3.284.9 Mupad [B] (verification not implemented)	1452

3.284.1 Optimal result

Integrand size = 11, antiderivative size = 25

$$\int \frac{1}{1+x+x^2+x^3} dx = \frac{\arctan(x)}{2} + \frac{1}{2} \log(1+x) - \frac{1}{4} \log(1+x^2)$$

output `1/2*arctan(x)+1/2*ln(1+x)-1/4*ln(x^2+1)`

3.284.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x+x^2+x^3} dx = \frac{\arctan(x)}{2} + \frac{1}{2} \log(1+x) - \frac{1}{4} \log(1+x^2)$$

input `Integrate[(1 + x + x^2 + x^3)^(-1),x]`

output `ArcTan[x]/2 + Log[1 + x]/2 - Log[1 + x^2]/4`

3.284.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3 + x^2 + x + 1} dx$$

$$\downarrow \text{2462}$$

$$\int \left(\frac{1-x}{2(x^2+1)} + \frac{1}{2(x+1)} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{\arctan(x)}{2} - \frac{1}{4} \log(x^2+1) + \frac{1}{2} \log(x+1)$$

input `Int[(1 + x + x^2 + x^3)^(-1),x]`

output `ArcTan[x]/2 + Log[1 + x]/2 - Log[1 + x^2]/4`

3.284.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2462 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegrand[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

3.284.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

method	result	size
default	$\frac{\arctan(x)}{2} + \frac{\ln(1+x)}{2} - \frac{\ln(x^2+1)}{4}$	20
risch	$\frac{\arctan(x)}{2} + \frac{\ln(1+x)}{2} - \frac{\ln(x^2+1)}{4}$	20
parallelrisc	$\frac{\ln(1+x)}{2} - \frac{\ln(x-i)}{4} - \frac{i \ln(x-i)}{4} - \frac{\ln(x+i)}{4} + \frac{i \ln(x+i)}{4}$	38

input `int(1/(x^3+x^2+x+1),x,method=_RETURNVERBOSE)`output `1/2*arctan(x)+1/2*ln(1+x)-1/4*ln(x^2+1)`**3.284.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{1}{1+x+x^2+x^3} dx = \frac{1}{2} \arctan(x) - \frac{1}{4} \log(x^2+1) + \frac{1}{2} \log(x+1)$$

input `integrate(1/(x^3+x^2+x+1),x, algorithm="fricas")`output `1/2*arctan(x) - 1/4*log(x^2 + 1) + 1/2*log(x + 1)`**3.284.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{1}{1+x+x^2+x^3} dx = \frac{\log(x+1)}{2} - \frac{\log(x^2+1)}{4} + \frac{\operatorname{atan}(x)}{2}$$

input `integrate(1/(x**3+x**2+x+1),x)`output `log(x + 1)/2 - log(x**2 + 1)/4 + atan(x)/2`

3.284.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{1}{1+x+x^2+x^3} dx = \frac{1}{2} \arctan(x) - \frac{1}{4} \log(x^2+1) + \frac{1}{2} \log(x+1)$$

input `integrate(1/(x^3+x^2+x+1),x, algorithm="maxima")`output `1/2*arctan(x) - 1/4*log(x^2 + 1) + 1/2*log(x + 1)`**3.284.8 Giac [A] (verification not implemented)**

Time = 0.35 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{1}{1+x+x^2+x^3} dx = \frac{1}{2} \arctan(x) - \frac{1}{4} \log(x^2+1) + \frac{1}{2} \log(|x+1|)$$

input `integrate(1/(x^3+x^2+x+1),x, algorithm="giac")`output `1/2*arctan(x) - 1/4*log(x^2 + 1) + 1/2*log(abs(x + 1))`**3.284.9 Mupad [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x+x^2+x^3} dx = \frac{\ln(x+1)}{2} + \ln(x-i) \left(-\frac{1}{4} - \frac{1}{4}i\right) + \ln(x+1i) \left(-\frac{1}{4} + \frac{1}{4}i\right)$$

input `int(1/(x + x^2 + x^3 + 1),x)`output `log(x + 1)/2 - log(x - 1i)*(1/4 + 1i/4) - log(x + 1i)*(1/4 - 1i/4)`

3.285 $\int x^2 \log(1 + x) dx$

3.285.1 Optimal result	1453
3.285.2 Mathematica [A] (verified)	1453
3.285.3 Rubi [A] (verified)	1454
3.285.4 Maple [A] (verified)	1455
3.285.5 Fricas [A] (verification not implemented)	1455
3.285.6 Sympy [A] (verification not implemented)	1456
3.285.7 Maxima [A] (verification not implemented)	1456
3.285.8 Giac [A] (verification not implemented)	1456
3.285.9 Mupad [B] (verification not implemented)	1457

3.285.1 Optimal result

Integrand size = 8, antiderivative size = 39

$$\int x^2 \log(1 + x) dx = -\frac{x}{3} + \frac{x^2}{6} - \frac{x^3}{9} + \frac{1}{3} \log(1 + x) + \frac{1}{3} x^3 \log(1 + x)$$

output `-1/3*x+1/6*x^2-1/9*x^3+1/3*ln(1+x)+1/3*x^3*ln(1+x)`

3.285.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.72

$$\int x^2 \log(1 + x) dx = \frac{1}{18} (x(-6 + 3x - 2x^2) + 6(1 + x^3) \log(1 + x))$$

input `Integrate[x^2*Log[1 + x],x]`

output `(x*(-6 + 3*x - 2*x^2) + 6*(1 + x^3)*Log[1 + x])/18`

3.285.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.97, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {2842, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2 \log(x+1) dx \\ & \quad \downarrow \text{2842} \\ & \frac{1}{3}x^3 \log(x+1) - \frac{1}{3} \int \frac{x^3}{x+1} dx \\ & \quad \downarrow \text{49} \\ & \frac{1}{3}x^3 \log(x+1) - \frac{1}{3} \int \left(x^2 - x + \frac{1}{-x-1} + 1 \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{1}{3}x^3 \log(x+1) + \frac{1}{3} \left(-\frac{x^3}{3} + \frac{x^2}{2} - x + \log(x+1) \right) \end{aligned}$$

input `Int[x^2*Log[1 + x],x]`

output `(x^3*Log[1 + x])/3 + (-x + x^2/2 - x^3/3 + Log[1 + x])/3`

3.285.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 2842 Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]*(b_.))*((f_.) + (g_.)*(x_)
)^(q_.), x_Symbol] := Simp[(f + g*x)^(q + 1)*((a + b*Log[c*(d + e*x)^n])/(
g*(q + 1))), x] - Simp[b*e*(n/(g*(q + 1))) Int[(f + g*x)^(q + 1)/(d + e*x
), x], x] /; FreeQ[{a, b, c, d, e, f, g, n, q}, x] && NeQ[e*f - d*g, 0] &&
NeQ[q, -1]
```

3.285.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.72

method	result	size
meijerg	$-\frac{x(4x^2-6x+12)}{36} + \frac{(4x^3+4)\ln(1+x)}{12}$	28
norman	$-\frac{x}{3} + \frac{x^2}{6} - \frac{x^3}{9} + \frac{\ln(1+x)}{3} + \frac{\ln(1+x)x^3}{3}$	30
risch	$-\frac{x}{3} + \frac{x^2}{6} - \frac{x^3}{9} + \frac{\ln(1+x)}{3} + \frac{\ln(1+x)x^3}{3}$	30
parts	$-\frac{x}{3} + \frac{x^2}{6} - \frac{x^3}{9} + \frac{\ln(1+x)}{3} + \frac{\ln(1+x)x^3}{3}$	30
parallelrisc	$\frac{\ln(1+x)x^3}{3} - \frac{x^3}{9} + \frac{x^2}{6} - \frac{x}{3} + \frac{\ln(1+x)}{3} + \frac{1}{3}$	31
derivativdivides	$\frac{(1+x)^3 \ln(1+x)}{3} - \frac{(1+x)^3}{9} - \ln(1+x)(1+x)^2 + \frac{(1+x)^2}{2} + (1+x)\ln(1+x) - 1 - x$	50
default	$\frac{(1+x)^3 \ln(1+x)}{3} - \frac{(1+x)^3}{9} - \ln(1+x)(1+x)^2 + \frac{(1+x)^2}{2} + (1+x)\ln(1+x) - 1 - x$	50

```
input int(ln(1+x)*x^2,x,method=_RETURNVERBOSE)
```

```
output -1/36*x*(4*x^2-6*x+12)+1/12*(4*x^3+4)*ln(1+x)
```

3.285.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.64

$$\int x^2 \log(1+x) dx = -\frac{1}{9}x^3 + \frac{1}{6}x^2 + \frac{1}{3}(x^3+1)\log(x+1) - \frac{1}{3}x$$

```
input integrate(x^2*log(1+x),x, algorithm="fricas")
```

```
output -1/9*x^3 + 1/6*x^2 + 1/3*(x^3 + 1)*log(x + 1) - 1/3*x
```


3.285.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.74

$$\int x^2 \log(1+x) dx = \frac{x^3 \log(x+1)}{3} - \frac{x^3}{9} + \frac{x^2}{6} - \frac{x}{3} + \frac{\log(x+1)}{3}$$

input `integrate(x**2*ln(1+x),x)`output `x**3*log(x + 1)/3 - x**3/9 + x**2/6 - x/3 + log(x + 1)/3`**3.285.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.74

$$\int x^2 \log(1+x) dx = \frac{1}{3} x^3 \log(x+1) - \frac{1}{9} x^3 + \frac{1}{6} x^2 - \frac{1}{3} x + \frac{1}{3} \log(x+1)$$

input `integrate(x^2*log(1+x),x, algorithm="maxima")`output `1/3*x^3*log(x + 1) - 1/9*x^3 + 1/6*x^2 - 1/3*x + 1/3*log(x + 1)`**3.285.8 Giac [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.26

$$\int x^2 \log(1+x) dx = \frac{1}{3} (x+1)^3 \log(x+1) - \frac{1}{9} (x+1)^3 - (x+1)^2 \log(x+1) + \frac{1}{2} (x+1)^2 + (x+1) \log(x+1) - x - 1$$

input `integrate(x^2*log(1+x),x, algorithm="giac")`output `1/3*(x + 1)^3*log(x + 1) - 1/9*(x + 1)^3 - (x + 1)^2*log(x + 1) + 1/2*(x + 1)^2 + (x + 1)*log(x + 1) - x - 1`

3.285.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.64

$$\int x^2 \log(1+x) dx = \frac{x^2}{6} - \frac{x}{3} - \frac{x^3}{9} + \frac{\ln(x+1)(x^3+1)}{3}$$

input `int(x^2*log(x + 1),x)`

output `x^2/6 - x/3 - x^3/9 + (log(x + 1)*(x^3 + 1))/3`

3.286 $\int e^{-x^3} x^5 dx$

3.286.1 Optimal result	1458
3.286.2 Mathematica [A] (verified)	1458
3.286.3 Rubi [A] (verified)	1459
3.286.4 Maple [A] (verified)	1460
3.286.5 Fricas [A] (verification not implemented)	1460
3.286.6 Sympy [A] (verification not implemented)	1461
3.286.7 Maxima [A] (verification not implemented)	1461
3.286.8 Giac [A] (verification not implemented)	1461
3.286.9 Mupad [B] (verification not implemented)	1462

3.286.1 Optimal result

Integrand size = 11, antiderivative size = 26

$$\int e^{-x^3} x^5 dx = -\frac{e^{-x^3}}{3} - \frac{1}{3}e^{-x^3} x^3$$

output `-1/3/exp(x^3)-1/3*x^3/exp(x^3)`

3.286.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.62

$$\int e^{-x^3} x^5 dx = -\frac{1}{3}e^{-x^3} (1 + x^3)$$

input `Integrate[x^5/E^x^3,x]`

output `-1/3*(1 + x^3)/E^x^3`

3.286.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2641, 2638}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{-x^3} x^5 dx$$

$$\downarrow \text{2641}$$

$$\int e^{-x^3} x^2 dx - \frac{1}{3} e^{-x^3} x^3$$

$$\downarrow \text{2638}$$

$$-\frac{1}{3} e^{-x^3} x^3 - \frac{e^{-x^3}}{3}$$

input `Int[x^5/E^x^3,x]`

output `-1/3*1/E^x^3 - x^3/(3*E^x^3)`

3.286.3.1 Defintions of rubi rules used

rule 2638 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[(e + f*x)^n*(F^(a + b*(c + d*x)^n)/(b*f*n*(c + d*x)^n *Log[F])), x] /; FreeQ[{F, a, b, c, d, e, f, n}, x] && EqQ[m, n - 1] && EqQ[d*e - c*f, 0]`

rule 2641 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^(m - n + 1)*(F^(a + b*(c + d*x)^n)/(b*d*n*Log[F])), x] - Simp[(m - n + 1)/(b*n*Log[F]) Int[(c + d*x)^(m - n)*F^(a + b*(c + d*x)^n), x], x] /; FreeQ[{F, a, b, c, d}, x] && IntegerQ[2*((m + 1)/n)] && LtQ[0, (m + 1)/n, 5] && IntegerQ[n] && (LtQ[0, n, m + 1] || LtQ[m, n, 0])`

3.286.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.54

method	result	size
gosper	$-\frac{(x^3+1)e^{-x^3}}{3}$	14
norman	$\left(-\frac{x^3}{3} - \frac{1}{3}\right)e^{-x^3}$	15
risch	$\left(-\frac{x^3}{3} - \frac{1}{3}\right)e^{-x^3}$	15
parallelrisc	$\frac{(-x^3-1)e^{-x^3}}{3}$	16
meijerg	$\frac{1}{3} - \frac{(2x^3+2)e^{-x^3}}{6}$	18
derivativedivides	$-\frac{e^{-x^3}}{3} - \frac{x^3e^{-x^3}}{3}$	21
default	$-\frac{e^{-x^3}}{3} - \frac{x^3e^{-x^3}}{3}$	21

input `int(x^5/exp(x^3),x,method=_RETURNVERBOSE)`output `-1/3*(x^3+1)/exp(x^3)`**3.286.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.50

$$\int e^{-x^3} x^5 dx = -\frac{1}{3} (x^3 + 1) e^{(-x^3)}$$

input `integrate(x^5/exp(x^3),x, algorithm="fracas")`output `-1/3*(x^3 + 1)*e^(-x^3)`

3.286.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.46

$$\int e^{-x^3} x^5 dx = \frac{(-x^3 - 1) e^{-x^3}}{3}$$

input `integrate(x**5/exp(x**3),x)`output `(-x**3 - 1)*exp(-x**3)/3`**3.286.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.50

$$\int e^{-x^3} x^5 dx = -\frac{1}{3} (x^3 + 1) e^{(-x^3)}$$

input `integrate(x^5/exp(x^3),x, algorithm="maxima")`output `-1/3*(x^3 + 1)*e^(-x^3)`**3.286.8 Giac [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.50

$$\int e^{-x^3} x^5 dx = -\frac{1}{3} (x^3 + 1) e^{(-x^3)}$$

input `integrate(x^5/exp(x^3),x, algorithm="giac")`output `-1/3*(x^3 + 1)*e^(-x^3)`

3.286.9 Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.50

$$\int e^{-x^3} x^5 dx = -\frac{e^{-x^3} (x^3 + 1)}{3}$$

input `int(x^5*exp(-x^3),x)`

output `-(exp(-x^3)*(x^3 + 1))/3`

3.287 $\int \tan^2(4x) dx$

3.287.1 Optimal result	1463
3.287.2 Mathematica [A] (verified)	1463
3.287.3 Rubi [A] (verified)	1464
3.287.4 Maple [A] (verified)	1465
3.287.5 Fricas [A] (verification not implemented)	1465
3.287.6 Sympy [A] (verification not implemented)	1465
3.287.7 Maxima [A] (verification not implemented)	1466
3.287.8 Giac [A] (verification not implemented)	1466
3.287.9 Mupad [B] (verification not implemented)	1466

3.287.1 Optimal result

Integrand size = 6, antiderivative size = 12

$$\int \tan^2(4x) dx = -x + \frac{1}{4} \tan(4x)$$

output `-x+1/4*tan(4*x)`

3.287.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.50

$$\int \tan^2(4x) dx = -\frac{1}{4} \arctan(\tan(4x)) + \frac{1}{4} \tan(4x)$$

input `Integrate[Tan[4*x]^2,x]`

output `-1/4*ArcTan[Tan[4*x]] + Tan[4*x]/4`

3.287.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan^2(4x) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(4x)^2 dx \\ & \quad \downarrow \text{3954} \\ & \frac{1}{4} \tan(4x) - \int 1 dx \\ & \quad \downarrow \text{24} \\ & \frac{1}{4} \tan(4x) - x \end{aligned}$$

input `Int[Tan[4*x]^2,x]`

output `-x + Tan[4*x]/4`

3.287.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

3.287.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
norman	$-x + \frac{\tan(4x)}{4}$	11
parallelrisch	$-x + \frac{\tan(4x)}{4}$	11
derivativedivides	$\frac{\tan(4x)}{4} - \frac{\arctan(\tan(4x))}{4}$	15
default	$\frac{\tan(4x)}{4} - \frac{\arctan(\tan(4x))}{4}$	15
risch	$-x + \frac{i}{2e^{8ix}+2}$	17

input `int(tan(4*x)^2,x,method=_RETURNVERBOSE)`output `-x+1/4*tan(4*x)`**3.287.5 Fricas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \tan^2(4x) dx = -x + \frac{1}{4} \tan(4x)$$

input `integrate(tan(4*x)^2,x, algorithm="fricas")`output `-x + 1/4*tan(4*x)`**3.287.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \tan^2(4x) dx = -x + \frac{\sin(4x)}{4 \cos(4x)}$$

input `integrate(tan(4*x)**2,x)`output `-x + sin(4*x)/(4*cos(4*x))`

3.287.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \tan^2(4x) dx = -x + \frac{1}{4} \tan(4x)$$

input `integrate(tan(4*x)^2,x, algorithm="maxima")`output `-x + 1/4*tan(4*x)`**3.287.8 Giac [A] (verification not implemented)**

Time = 0.40 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \tan^2(4x) dx = -x + \frac{1}{4} \tan(4x)$$

input `integrate(tan(4*x)^2,x, algorithm="giac")`output `-x + 1/4*tan(4*x)`**3.287.9 Mupad [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \tan^2(4x) dx = \frac{\tan(4x)}{4} - x$$

input `int(tan(4*x)^2,x)`output `tan(4*x)/4 - x`

$$3.288 \quad \int \frac{1}{\sqrt{-5+12x+9x^2}} dx$$

3.288.1 Optimal result	1467
3.288.2 Mathematica [A] (verified)	1467
3.288.3 Rubi [A] (verified)	1468
3.288.4 Maple [A] (verified)	1469
3.288.5 Fricas [A] (verification not implemented)	1469
3.288.6 Sympy [A] (verification not implemented)	1469
3.288.7 Maxima [A] (verification not implemented)	1470
3.288.8 Giac [A] (verification not implemented)	1470
3.288.9 Mupad [B] (verification not implemented)	1470

3.288.1 Optimal result

Integrand size = 14, antiderivative size = 25

$$\int \frac{1}{\sqrt{-5+12x+9x^2}} dx = \frac{1}{3} \operatorname{arctanh} \left(\frac{2+3x}{\sqrt{-5+12x+9x^2}} \right)$$

output `1/3*arctanh((2+3*x)/(9*x^2+12*x-5)^(1/2))`

3.288.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{1}{\sqrt{-5+12x+9x^2}} dx = -\frac{1}{3} \log \left(-2 - 3x + \sqrt{-5+12x+9x^2} \right)$$

input `Integrate[1/Sqrt[-5 + 12*x + 9*x^2],x]`

output `-1/3*Log[-2 - 3*x + Sqrt[-5 + 12*x + 9*x^2]]`

3.288.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {1092, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{9x^2 + 12x - 5}} dx$$

↓ 1092

$$2 \int \frac{1}{36 - \frac{36(3x+2)^2}{9x^2+12x-5}} d \frac{6(3x+2)}{\sqrt{9x^2 + 12x - 5}}$$

↓ 219

$$\frac{1}{3} \operatorname{arctanh} \left(\frac{3x+2}{\sqrt{9x^2 + 12x - 5}} \right)$$

input `Int[1/Sqrt[-5 + 12*x + 9*x^2], x]`

output `ArcTanh[(2 + 3*x)/Sqrt[-5 + 12*x + 9*x^2]]/3`

3.288.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1092 `Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Simp[2 Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x]`

3.288.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

method	result	size
trager	$-\frac{\ln(-2-3x+\sqrt{9x^2+12x-5})}{3}$	21
default	$\frac{\ln\left(\frac{(9x+6)\sqrt{9}+\sqrt{9x^2+12x-5}}{9}\right)\sqrt{9}}{9}$	30

input `int(1/(9*x^2+12*x-5)^(1/2),x,method=_RETURNVERBOSE)`output `-1/3*ln(-2-3*x+(9*x^2+12*x-5)^(1/2))`**3.288.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{1}{\sqrt{-5+12x+9x^2}} dx = -\frac{1}{3} \log(-3x + \sqrt{9x^2 + 12x - 5} - 2)$$

input `integrate(1/(9*x^2+12*x-5)^(1/2),x, algorithm="fricas")`output `-1/3*log(-3*x + sqrt(9*x^2 + 12*x - 5) - 2)`**3.288.6 Sympy [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{1}{\sqrt{-5+12x+9x^2}} dx = \frac{\log(18x + 6\sqrt{9x^2 + 12x - 5} + 12)}{3}$$

input `integrate(1/(9*x**2+12*x-5)**(1/2),x)`output `log(18*x + 6*sqrt(9*x**2 + 12*x - 5) + 12)/3`

3.288.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{1}{\sqrt{-5 + 12x + 9x^2}} dx = \frac{1}{3} \log \left(18x + 6\sqrt{9x^2 + 12x - 5} + 12 \right)$$

input `integrate(1/(9*x^2+12*x-5)^(1/2),x, algorithm="maxima")`output `1/3*log(18*x + 6*sqrt(9*x^2 + 12*x - 5) + 12)`**3.288.8 Giac [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.64

$$\int \frac{1}{\sqrt{-5 + 12x + 9x^2}} dx = \frac{1}{6} \sqrt{9x^2 + 12x - 5}(3x + 2) + \frac{3}{2} \log \left(\left| -3x + \sqrt{9x^2 + 12x - 5} - 2 \right| \right)$$

input `integrate(1/(9*x^2+12*x-5)^(1/2),x, algorithm="giac")`output `1/6*sqrt(9*x^2 + 12*x - 5)*(3*x + 2) + 3/2*log(abs(-3*x + sqrt(9*x^2 + 12*x - 5) - 2))`**3.288.9 Mupad [B] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{1}{\sqrt{-5 + 12x + 9x^2}} dx = \frac{\ln(3x + \sqrt{9x^2 + 12x - 5} + 2)}{3}$$

input `int(1/(12*x + 9*x^2 - 5)^(1/2),x)`output `log(3*x + (12*x + 9*x^2 - 5)^(1/2) + 2)/3`

3.289 $\int x^2 \arctan(x) dx$

3.289.1 Optimal result1471
3.289.2 Mathematica [A] (verified)1471
3.289.3 Rubi [A] (verified)1472
3.289.4 Maple [A] (verified)1473
3.289.5 Fricas [A] (verification not implemented)1473
3.289.6 Sympy [A] (verification not implemented)1474
3.289.7 Maxima [A] (verification not implemented)1474
3.289.8 Giac [A] (verification not implemented)1474
3.289.9 Mupad [B] (verification not implemented)1475

3.289.1 Optimal result

Integrand size = 6, antiderivative size = 27

$$\int x^2 \arctan(x) dx = -\frac{x^2}{6} + \frac{1}{3}x^3 \arctan(x) + \frac{1}{6} \log(1 + x^2)$$

output `-1/6*x^2+1/3*x^3*arctan(x)+1/6*ln(x^2+1)`

3.289.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int x^2 \arctan(x) dx = \frac{1}{6}(-x^2 + 2x^3 \arctan(x) + \log(1 + x^2))$$

input `Integrate[x^2*ArcTan[x],x]`

output `(-x^2 + 2*x^3*ArcTan[x] + Log[1 + x^2])/6`

3.289.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5361, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \arctan(x) dx \\
 & \quad \downarrow \text{5361} \\
 & \frac{1}{3}x^3 \arctan(x) - \frac{1}{3} \int \frac{x^3}{x^2+1} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{3}x^3 \arctan(x) - \frac{1}{6} \int \frac{x^2}{x^2+1} dx \\
 & \quad \downarrow \text{49} \\
 & \frac{1}{3}x^3 \arctan(x) - \frac{1}{6} \int \left(1 + \frac{1}{-x^2-1}\right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{3}x^3 \arctan(x) + \frac{1}{6}(\log(x^2+1) - x^2)
 \end{aligned}$$

input `Int[x^2*ArcTan[x],x]`

output `(x^3*ArcTan[x])/3 + (-x^2 + Log[1 + x^2])/6`

3.289.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5361 `Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m +
1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x],
x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] &
& IntegerQ[m])) && NeQ[m, -1]`

3.289.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

method	result	size
default	$-\frac{x^2}{6} + \frac{x^3 \arctan(x)}{3} + \frac{\ln(x^2+1)}{6}$	22
parts	$-\frac{x^2}{6} + \frac{x^3 \arctan(x)}{3} + \frac{\ln(x^2+1)}{6}$	22
parallelrisch	$\frac{x^3 \arctan(x)}{3} - \frac{x^2}{6} + \frac{\ln(x^2+1)}{6} + \frac{1}{6}$	23
meijerg	$-\frac{x^2}{6} + \frac{x^4 \arctan(\sqrt{x^2})}{3\sqrt{x^2}} + \frac{\ln(x^2+1)}{6}$	31
risch	$-\frac{ix^3 \ln(ix+1)}{6} + \frac{ix^3 \ln(-ix+1)}{6} - \frac{x^2}{6} + \frac{\ln(x^2+1)}{6}$	41

input `int(x^2*arctan(x),x,method=_RETURNVERBOSE)`

output `-1/6*x^2+1/3*x^3*arctan(x)+1/6*ln(x^2+1)`

3.289.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int x^2 \arctan(x) dx = \frac{1}{3} x^3 \arctan(x) - \frac{1}{6} x^2 + \frac{1}{6} \log(x^2 + 1)$$

input `integrate(x^2*arctan(x),x, algorithm="fricas")`

output `1/3*x^3*arctan(x) - 1/6*x^2 + 1/6*log(x^2 + 1)`

3.289.6 Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

$$\int x^2 \arctan(x) dx = \frac{x^3 \operatorname{atan}(x)}{3} - \frac{x^2}{6} + \frac{\log(x^2 + 1)}{6}$$

input `integrate(x**2*atan(x),x)`output `x**3*atan(x)/3 - x**2/6 + log(x**2 + 1)/6`**3.289.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int x^2 \arctan(x) dx = \frac{1}{3} x^3 \arctan(x) - \frac{1}{6} x^2 + \frac{1}{6} \log(x^2 + 1)$$

input `integrate(x^2*arctan(x),x, algorithm="maxima")`output `1/3*x^3*arctan(x) - 1/6*x^2 + 1/6*log(x^2 + 1)`**3.289.8 Giac [A] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int x^2 \arctan(x) dx = \frac{1}{3} x^3 \arctan(x) - \frac{1}{6} x^2 + \frac{1}{6} \log(x^2 + 1)$$

input `integrate(x^2*arctan(x),x, algorithm="giac")`output `1/3*x^3*arctan(x) - 1/6*x^2 + 1/6*log(x^2 + 1)`

3.289.9 Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int x^2 \arctan(x) dx = \frac{\ln(x^2 + 1)}{6} + \frac{x^3 \operatorname{atan}(x)}{3} - \frac{x^2}{6}$$

input `int(x^2*atan(x),x)`

output `log(x^2 + 1)/6 + (x^3*atan(x))/3 - x^2/6`

$$3.290 \quad \int \frac{1-\sqrt{x}}{\sqrt[3]{x}} dx$$

3.290.1 Optimal result	1476
3.290.2 Mathematica [A] (verified)	1476
3.290.3 Rubi [A] (verified)	1477
3.290.4 Maple [A] (verified)	1478
3.290.5 Fricas [A] (verification not implemented)	1478
3.290.6 Sympy [A] (verification not implemented)	1478
3.290.7 Maxima [A] (verification not implemented)	1479
3.290.8 Giac [A] (verification not implemented)	1479
3.290.9 Mupad [B] (verification not implemented)	1479

3.290.1 Optimal result

Integrand size = 15, antiderivative size = 19

$$\int \frac{1-\sqrt{x}}{\sqrt[3]{x}} dx = \frac{3x^{2/3}}{2} - \frac{6x^{7/6}}{7}$$

output `3/2*x^(2/3)-6/7*x^(7/6)`

3.290.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{1-\sqrt{x}}{\sqrt[3]{x}} dx = \frac{3x^{2/3}}{2} - \frac{6x^{7/6}}{7}$$

input `Integrate[(1 - Sqrt[x])/x^(1/3), x]`

output `(3*x^(2/3))/2 - (6*x^(7/6))/7`

3.290.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {802, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1 - \sqrt{x}}{\sqrt[3]{x}} dx$$

↓ 802

$$\int \left(\frac{1}{\sqrt[3]{x}} - \sqrt[6]{x} \right) dx$$

↓ 2009

$$\frac{3x^{2/3}}{2} - \frac{6x^{7/6}}{7}$$

input `Int[(1 - Sqrt[x])/x^(1/3),x]`

output `(3*x^(2/3))/2 - (6*x^(7/6))/7`

3.290.3.1 Defintions of rubi rules used

rule 802 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.290.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

method	result	size
derivativedivides	$\frac{3x^{\frac{2}{3}}}{2} - \frac{6x^{\frac{7}{6}}}{7}$	12
default	$\frac{3x^{\frac{2}{3}}}{2} - \frac{6x^{\frac{7}{6}}}{7}$	12

input `int((1-x^(1/2))/x^(1/3),x,method=_RETURNVERBOSE)`output `3/2*x^(2/3)-6/7*x^(7/6)`**3.290.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int \frac{1 - \sqrt{x}}{\sqrt[3]{x}} dx = -\frac{6}{7} x^{\frac{7}{6}} + \frac{3}{2} x^{\frac{2}{3}}$$

input `integrate((1-x^(1/2))/x^(1/3),x, algorithm="fricas")`output `-6/7*x^(7/6) + 3/2*x^(2/3)`**3.290.6 Sympy [A] (verification not implemented)**

Time = 0.38 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1 - \sqrt{x}}{\sqrt[3]{x}} dx = -\frac{6x^{\frac{7}{6}}}{7} + \frac{3x^{\frac{2}{3}}}{2}$$

input `integrate((1-x**(1/2))/x**(1/3),x)`output `-6*x**(7/6)/7 + 3*x**(2/3)/2`

3.290.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int \frac{1 - \sqrt{x}}{\sqrt[3]{x}} dx = -\frac{6}{7} x^{\frac{7}{6}} + \frac{3}{2} x^{\frac{2}{3}}$$

input `integrate((1-x^(1/2))/x^(1/3),x, algorithm="maxima")`output `-6/7*x^(7/6) + 3/2*x^(2/3)`**3.290.8 Giac [A] (verification not implemented)**

Time = 0.35 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int \frac{1 - \sqrt{x}}{\sqrt[3]{x}} dx = -\frac{6}{7} x^{\frac{7}{6}} + \frac{3}{2} x^{\frac{2}{3}}$$

input `integrate((1-x^(1/2))/x^(1/3),x, algorithm="giac")`output `-6/7*x^(7/6) + 3/2*x^(2/3)`**3.290.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{1 - \sqrt{x}}{\sqrt[3]{x}} dx = -\frac{3x^{2/3}(4\sqrt{x} - 7)}{14}$$

input `int(-(x^(1/2) - 1)/x^(1/3),x)`output `-(3*x^(2/3)*(4*x^(1/2) - 7))/14`

3.291 $\int \frac{1}{-e^{-x}+e^x} dx$

3.291.1 Optimal result	1480
3.291.2 Mathematica [A] (verified)	1480
3.291.3 Rubi [A] (verified)	1481
3.291.4 Maple [A] (verified)	1482
3.291.5 Fricas [B] (verification not implemented)	1482
3.291.6 Sympy [B] (verification not implemented)	1482
3.291.7 Maxima [B] (verification not implemented)	1483
3.291.8 Giac [B] (verification not implemented)	1483
3.291.9 Mupad [B] (verification not implemented)	1484

3.291.1 Optimal result

Integrand size = 13, antiderivative size = 6

$$\int \frac{1}{-e^{-x}+e^x} dx = -\operatorname{arctanh}(e^x)$$

output `-arctanh(exp(x))`

3.291.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{1}{-e^{-x}+e^x} dx = -\operatorname{arctanh}(e^x)$$

input `Integrate[(-E^(-x) + E^x)^(-1), x]`

output `-ArcTanh[E^x]`

3.291.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {2720, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{e^x - e^{-x}} dx$$

↓ 2720

$$\int \frac{1}{e^{2x} - 1} de^x$$

↓ 220

$$-\operatorname{arctanh}(e^x)$$

input `Int[(-E^(-x) + E^x)^(-1), x]`

output `-ArcTanh[E^x]`

3.291.3.1 Defintions of rubi rules used

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

3.291.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

method	result	size
derivativedivides	$-\operatorname{arctanh}(e^x)$	6
default	$-\operatorname{arctanh}(e^x)$	6
norman	$\frac{\ln(-1+e^x)}{2} - \frac{\ln(1+e^x)}{2}$	16
risch	$\frac{\ln(-1+e^x)}{2} - \frac{\ln(1+e^x)}{2}$	16
parallelrisch	$\frac{\ln(-1+e^x)}{2} - \frac{\ln(1+e^x)}{2}$	16

input `int(1/(-1/exp(x)+exp(x)),x,method=_RETURNVERBOSE)`

output `-arctanh(exp(x))`

3.291.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(5) = 10$.

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 2.50

$$\int \frac{1}{-e^{-x} + e^x} dx = -\frac{1}{2} \log(e^x + 1) + \frac{1}{2} \log(e^x - 1)$$

input `integrate(1/(-1/exp(x)+exp(x)),x, algorithm="fracas")`

output `-1/2*log(e^x + 1) + 1/2*log(e^x - 1)`

3.291.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(5) = 10$.

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 3.17

$$\int \frac{1}{-e^{-x} + e^x} dx = \frac{\log(-1 + e^{-x})}{2} - \frac{\log(1 + e^{-x})}{2}$$

input `integrate(1/(-1/exp(x)+exp(x)),x)`

output `log(-1 + exp(-x))/2 - log(1 + exp(-x))/2`

3.291.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(5) = 10$.

Time = 0.21 (sec) , antiderivative size = 19, normalized size of antiderivative = 3.17

$$\int \frac{1}{-e^{-x} + e^x} dx = -\frac{1}{2} \log(e^{-x} + 1) + \frac{1}{2} \log(e^{-x} - 1)$$

input `integrate(1/(-1/exp(x)+exp(x)),x, algorithm="maxima")`

output `-1/2*log(e^(-x) + 1) + 1/2*log(e^(-x) - 1)`

3.291.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 16 vs. $2(5) = 10$.

Time = 0.34 (sec) , antiderivative size = 16, normalized size of antiderivative = 2.67

$$\int \frac{1}{-e^{-x} + e^x} dx = -\frac{1}{2} \log(e^x + 1) + \frac{1}{2} \log(|e^x - 1|)$$

input `integrate(1/(-1/exp(x)+exp(x)),x, algorithm="giac")`

output `-1/2*log(e^x + 1) + 1/2*log(abs(e^x - 1))`

3.291.9 Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 2.50

$$\int \frac{1}{-e^{-x} + e^x} dx = \frac{\ln(e^x - 1)}{2} - \frac{\ln(e^x + 1)}{2}$$

input `int(-1/(exp(-x) - exp(x)),x)`

output `log(exp(x) - 1)/2 - log(exp(x) + 1)/2`

3.292 $\int \frac{x}{10+2x^2+x^4} dx$

3.292.1 Optimal result	1485
3.292.2 Mathematica [A] (verified)	1485
3.292.3 Rubi [A] (verified)	1486
3.292.4 Maple [A] (verified)	1487
3.292.5 Fricas [A] (verification not implemented)	1487
3.292.6 Sympy [A] (verification not implemented)	1487
3.292.7 Maxima [A] (verification not implemented)	1488
3.292.8 Giac [A] (verification not implemented)	1488
3.292.9 Mupad [B] (verification not implemented)	1488

3.292.1 Optimal result

Integrand size = 14, antiderivative size = 14

$$\int \frac{x}{10+2x^2+x^4} dx = \frac{1}{6} \arctan\left(\frac{1}{3}(1+x^2)\right)$$

output `1/6*arctan(1/3*x^2+1/3)`

3.292.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{x}{10+2x^2+x^4} dx = \frac{1}{6} \arctan\left(\frac{1}{3}(1+x^2)\right)$$

input `Integrate[x/(10 + 2*x^2 + x^4),x]`

output `ArcTan[(1 + x^2)/3]/6`

3.292.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {1432, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x}{x^4 + 2x^2 + 10} dx \\ & \quad \downarrow \text{1432} \\ & \frac{1}{2} \int \frac{1}{x^4 + 2x^2 + 10} dx^2 \\ & \quad \downarrow \text{1083} \\ & - \int \frac{1}{-x^4 - 36} d(2x^2 + 2) \\ & \quad \downarrow \text{217} \\ & \frac{1}{6} \arctan\left(\frac{1}{6}(2x^2 + 2)\right) \end{aligned}$$

input `Int[x/(10 + 2*x^2 + x^4),x]`

output `ArcTan[(2 + 2*x^2)/6]/6`

3.292.3.1 Defintions of rubi rules used

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])`

rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 1432 `Int[(x_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, p}, x]`

3.292.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{\arctan\left(\frac{x^2}{3} + \frac{1}{3}\right)}{6}$	11
risch	$\frac{\arctan\left(\frac{x^2}{3} + \frac{1}{3}\right)}{6}$	11
parallelrisc	$\frac{i \ln(x^2+3i+1)}{12} - \frac{i \ln(x^2-3i+1)}{12}$	24

input `int(x/(x^4+2*x^2+10),x,method=_RETURNVERBOSE)`output `1/6*arctan(1/3*x^2+1/3)`**3.292.5 Fricas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{x}{10 + 2x^2 + x^4} dx = \frac{1}{6} \arctan\left(\frac{1}{3}x^2 + \frac{1}{3}\right)$$

input `integrate(x/(x^4+2*x^2+10),x, algorithm="fricas")`output `1/6*arctan(1/3*x^2 + 1/3)`**3.292.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{x}{10 + 2x^2 + x^4} dx = \frac{\operatorname{atan}\left(\frac{x^2}{3} + \frac{1}{3}\right)}{6}$$

input `integrate(x/(x**4+2*x**2+10),x)`output `atan(x**2/3 + 1/3)/6`

3.292.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{x}{10 + 2x^2 + x^4} dx = \frac{1}{6} \arctan\left(\frac{1}{3}x^2 + \frac{1}{3}\right)$$

input `integrate(x/(x^4+2*x^2+10),x, algorithm="maxima")`output `1/6*arctan(1/3*x^2 + 1/3)`**3.292.8 Giac [A] (verification not implemented)**

Time = 0.56 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{x}{10 + 2x^2 + x^4} dx = \frac{1}{6} \arctan\left(\frac{1}{3}x^2 + \frac{1}{3}\right)$$

input `integrate(x/(x^4+2*x^2+10),x, algorithm="giac")`output `1/6*arctan(1/3*x^2 + 1/3)`**3.292.9 Mupad [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{x}{10 + 2x^2 + x^4} dx = \frac{\operatorname{atan}\left(\frac{x^2}{3} + \frac{1}{3}\right)}{6}$$

input `int(x/(2*x^2 + x^4 + 10),x)`output `atan(x^2/3 + 1/3)/6`

$$3.293 \quad \int \frac{1}{\frac{1}{\sqrt[3]{x}} + x} dx$$

3.293.1 Optimal result	1489
3.293.2 Mathematica [A] (verified)	1489
3.293.3 Rubi [A] (verified)	1490
3.293.4 Maple [A] (verified)	1491
3.293.5 Fricas [A] (verification not implemented)	1491
3.293.6 Sympy [A] (verification not implemented)	1491
3.293.7 Maxima [A] (verification not implemented)	1492
3.293.8 Giac [B] (verification not implemented)	1492
3.293.9 Mupad [B] (verification not implemented)	1493

3.293.1 Optimal result

Integrand size = 9, antiderivative size = 12

$$\int \frac{1}{\frac{1}{\sqrt[3]{x}} + x} dx = \frac{3}{4} \log(1 + x^{4/3})$$

output `3/4*ln(1+x^(4/3))`

3.293.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{\frac{1}{\sqrt[3]{x}} + x} dx = \frac{3}{4} \log(1 + x^{4/3})$$

input `Integrate[(x^(-1/3) + x)^(-1), x]`

output `(3*Log[1 + x^(4/3)])/4`

3.293.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {2027, 792}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x + \frac{1}{\sqrt[3]{x}}} dx$$

↓ 2027

$$\int \frac{\sqrt[3]{x}}{x^{4/3} + 1} dx$$

↓ 792

$$\frac{3}{4} \log(x^{4/3} + 1)$$

input `Int[(x^(-1/3) + x)^(-1),x]`

output `(3*Log[1 + x^(4/3)])/4`

3.293.3.1 Defintions of rubi rules used

rule 792 `Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]`

rule 2027 `Int[(F*_.)*((a_)*(x_)^(r_) + (b_)*(x_)^(s_))^(p_), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r))^p*Fx, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] && PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])`

3.293.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

method	result	size
derivativedivides	$\frac{3 \ln(1+x^{\frac{4}{3}})}{4}$	9
default	$\frac{3 \ln(1+x^{\frac{4}{3}})}{4}$	9
meijerg	$\frac{3 \ln(1+x^{\frac{4}{3}})}{4}$	9
trager	$-\frac{\ln\left(\frac{3x^{\frac{20}{3}} - x^8 - 6x^{\frac{16}{3}} - 6x^{\frac{8}{3}} + 7x^4 + 3x^{\frac{4}{3}} - 1}{(x^4+1)^3}\right)}{4}$	44

input `int(1/(1/x^(1/3)+x),x,method=_RETURNVERBOSE)`output `3/4*ln(1+x^(4/3))`**3.293.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{\frac{1}{\sqrt[3]{x}} + x} dx = \frac{3}{4} \log(x^{\frac{4}{3}} + 1)$$

input `integrate(1/(1/x^(1/3)+x),x, algorithm="fracas")`output `3/4*log(x^(4/3) + 1)`**3.293.6 Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1}{\frac{1}{\sqrt[3]{x}} + x} dx = \frac{3 \log(x^{\frac{4}{3}} + 1)}{4}$$

input `integrate(1/(1/x**(1/3)+x),x)`

output `3*log(x**(4/3) + 1)/4`

3.293.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{\frac{1}{\sqrt[3]{x}} + x} dx = \frac{3}{4} \log(x^{\frac{4}{3}} + 1)$$

input `integrate(1/(1/x^(1/3)+x),x, algorithm="maxima")`

output `3/4*log(x^(4/3) + 1)`

3.293.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 32 vs. 2(8) = 16.

Time = 0.32 (sec) , antiderivative size = 32, normalized size of antiderivative = 2.67

$$\int \frac{1}{\frac{1}{\sqrt[3]{x}} + x} dx = \frac{3}{4} \log(\sqrt{2}x^{\frac{1}{3}} + x^{\frac{2}{3}} + 1) + \frac{3}{4} \log(-\sqrt{2}x^{\frac{1}{3}} + x^{\frac{2}{3}} + 1)$$

input `integrate(1/(1/x^(1/3)+x),x, algorithm="giac")`

output `3/4*log(sqrt(2)*x^(1/3) + x^(2/3) + 1) + 3/4*log(-sqrt(2)*x^(1/3) + x^(2/3) + 1)`

3.293.9 Mupad [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{\frac{1}{\sqrt[3]{x}} + x} dx = \frac{3 \ln(x^{4/3} + 1)}{4}$$

input `int(1/(x + 1/x^(1/3)),x)`

output `(3*log(x^(4/3) + 1))/4`

3.294 $\int \cos^4(x) \sin^2(x) dx$

3.294.1 Optimal result	1494
3.294.2 Mathematica [A] (verified)	1494
3.294.3 Rubi [A] (verified)	1495
3.294.4 Maple [A] (verified)	1496
3.294.5 Fricas [A] (verification not implemented)	1497
3.294.6 Sympy [A] (verification not implemented)	1497
3.294.7 Maxima [A] (verification not implemented)	1497
3.294.8 Giac [A] (verification not implemented)	1498
3.294.9 Mupad [B] (verification not implemented)	1498

3.294.1 Optimal result

Integrand size = 9, antiderivative size = 34

$$\int \cos^4(x) \sin^2(x) dx = \frac{x}{16} + \frac{1}{16} \cos(x) \sin(x) + \frac{1}{24} \cos^3(x) \sin(x) - \frac{1}{6} \cos^5(x) \sin(x)$$

output `1/16*x+1/16*cos(x)*sin(x)+1/24*cos(x)^3*sin(x)-1/6*cos(x)^5*sin(x)`

3.294.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.88

$$\int \cos^4(x) \sin^2(x) dx = \frac{x}{16} + \frac{1}{64} \sin(2x) - \frac{1}{64} \sin(4x) - \frac{1}{192} \sin(6x)$$

input `Integrate[Cos[x]^4*Sin[x]^2,x]`

output `x/16 + Sin[2*x]/64 - Sin[4*x]/64 - Sin[6*x]/192`

3.294.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.29, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.778$, Rules used = {3042, 3048, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^2(x) \cos^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^2 \cos(x)^4 dx \\
 & \quad \downarrow \text{3048} \\
 & \frac{1}{6} \int \cos^4(x) dx - \frac{1}{6} \sin(x) \cos^5(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{6} \int \sin\left(x + \frac{\pi}{2}\right)^4 dx - \frac{1}{6} \sin(x) \cos^5(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{1}{6} \left(\frac{3}{4} \int \cos^2(x) dx + \frac{1}{4} \sin(x) \cos^3(x) \right) - \frac{1}{6} \sin(x) \cos^5(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{6} \left(\frac{3}{4} \int \sin\left(x + \frac{\pi}{2}\right)^2 dx + \frac{1}{4} \sin(x) \cos^3(x) \right) - \frac{1}{6} \sin(x) \cos^5(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{1}{6} \left(\frac{3}{4} \left(\frac{\int 1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \right) + \frac{1}{4} \sin(x) \cos^3(x) \right) - \frac{1}{6} \sin(x) \cos^5(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{1}{6} \left(\frac{1}{4} \sin(x) \cos^3(x) + \frac{3}{4} \left(\frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \right) \right) - \frac{1}{6} \sin(x) \cos^5(x)
 \end{aligned}$$

input `Int[Cos[x]^4*Sin[x]^2,x]`

output $-1/6*(\text{Cos}[x]^5*\text{Sin}[x]) + ((\text{Cos}[x]^3*\text{Sin}[x])/4 + (3*(x/2 + (\text{Cos}[x]*\text{Sin}[x])/2))/4)/6$

3.294.3.1 Defintions of rubi rules used

rule 24 $\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3048 $\text{Int}[(\text{cos}[(e_.) + (f_.)*(x_.)]*(b_.))^n*((a_.)*\text{sin}[(e_.) + (f_.)*(x_.)])^m, x_Symbol] \rightarrow \text{Simp}[(-a)*(b*\text{Cos}[e + f*x])^{n+1}*((a*\text{Sin}[e + f*x])^{m-1}/(b*f*(m+n))), x] + \text{Simp}[a^2*((m-1)/(m+n)) \text{Int}[(b*\text{Cos}[e + f*x])^n*(a*\text{Sin}[e + f*x])^{m-2}, x], x] /; \text{FreeQ}[\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{NeQ}[m+n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

rule 3115 $\text{Int}[(b_.)*\text{sin}[(c_.) + (d_.)*(x_.)]^n, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{n-1}/(d*n), x] + \text{Simp}[b^2*((n-1)/n) \text{Int}[(b*\text{Sin}[c + d*x])^{n-2}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

3.294.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.68

method	result
risch	$\frac{x}{16} - \frac{\sin(6x)}{192} - \frac{\sin(4x)}{64} + \frac{\sin(2x)}{64}$
parallelrisch	$\frac{x}{16} - \frac{\sin(6x)}{192} - \frac{\sin(4x)}{64} + \frac{\sin(2x)}{64}$
default	$-\frac{(\cos^5(x)) \sin(x)}{6} + \frac{(\cos^3(x) + \frac{3 \cos(x)}{2}) \sin(x)}{24} + \frac{x}{16}$
norman	$\frac{x}{16} + \frac{47(\tan^3(\frac{x}{2}))}{24} - \frac{13(\tan^5(\frac{x}{2}))}{4} + \frac{13(\tan^7(\frac{x}{2}))}{4} - \frac{47(\tan^9(\frac{x}{2}))}{24} + \frac{(\tan^{11}(\frac{x}{2}))}{8} + \frac{3x(\tan^2(\frac{x}{2}))}{8} + \frac{15x(\tan^4(\frac{x}{2}))}{16} + \frac{5x(\tan^6(\frac{x}{2}))}{4} + \frac{1}{(1+\tan^2(\frac{x}{2}))^6}$

input $\text{int}(\sin(x)^2*\cos(x)^4,x,\text{method}=_RETURNVERBOSE)$

output `1/16*x-1/192*sin(6*x)-1/64*sin(4*x)+1/64*sin(2*x)`

3.294.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.74

$$\int \cos^4(x) \sin^2(x) dx = -\frac{1}{48} (8 \cos(x)^5 - 2 \cos(x)^3 - 3 \cos(x)) \sin(x) + \frac{1}{16} x$$

input `integrate(cos(x)^4*sin(x)^2,x, algorithm="fricas")`

output `-1/48*(8*cos(x)^5 - 2*cos(x)^3 - 3*cos(x))*sin(x) + 1/16*x`

3.294.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.91

$$\int \cos^4(x) \sin^2(x) dx = \frac{x}{16} - \frac{\sin(x) \cos^5(x)}{6} + \frac{\sin(x) \cos^3(x)}{24} + \frac{\sin(x) \cos(x)}{16}$$

input `integrate(cos(x)**4*sin(x)**2,x)`

output `x/16 - sin(x)*cos(x)**5/6 + sin(x)*cos(x)**3/24 + sin(x)*cos(x)/16`

3.294.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.53

$$\int \cos^4(x) \sin^2(x) dx = \frac{1}{48} \sin(2x)^3 + \frac{1}{16} x - \frac{1}{64} \sin(4x)$$

input `integrate(cos(x)^4*sin(x)^2,x, algorithm="maxima")`

output `1/48*sin(2*x)^3 + 1/16*x - 1/64*sin(4*x)`

3.294.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.65

$$\int \cos^4(x) \sin^2(x) dx = \frac{1}{16} x - \frac{1}{192} \sin(6x) - \frac{1}{64} \sin(4x) + \frac{1}{64} \sin(2x)$$

input `integrate(cos(x)^4*sin(x)^2,x, algorithm="giac")`output `1/16*x - 1/192*sin(6*x) - 1/64*sin(4*x) + 1/64*sin(2*x)`**3.294.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.76

$$\int \cos^4(x) \sin^2(x) dx = \left(\frac{\cos(x)^3}{6} + \frac{\cos(x)}{8} \right) \sin(x)^3 - \frac{\cos(x) \sin(x)}{16} + \frac{x}{16}$$

input `int(cos(x)^4*sin(x)^2,x)`output `x/16 - (cos(x)*sin(x))/16 + sin(x)^3*(cos(x)/8 + cos(x)^3/6)`

3.295 $\int \frac{1}{\sqrt{5-4x-x^2}} dx$

3.295.1 Optimal result	1499
3.295.2 Mathematica [A] (verified)	1499
3.295.3 Rubi [A] (verified)	1500
3.295.4 Maple [A] (verified)	1501
3.295.5 Fricas [B] (verification not implemented)	1501
3.295.6 Sympy [A] (verification not implemented)	1501
3.295.7 Maxima [A] (verification not implemented)	1502
3.295.8 Giac [B] (verification not implemented)	1502
3.295.9 Mupad [B] (verification not implemented)	1502

3.295.1 Optimal result

Integrand size = 14, antiderivative size = 12

$$\int \frac{1}{\sqrt{5-4x-x^2}} dx = -\arcsin\left(\frac{1}{3}(-2-x)\right)$$

output `arcsin(2/3+1/3*x)`

3.295.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.92

$$\int \frac{1}{\sqrt{5-4x-x^2}} dx = -2 \arctan\left(\frac{\sqrt{5-4x-x^2}}{5+x}\right)$$

input `Integrate[1/Sqrt[5 - 4*x - x^2], x]`

output `-2*ArcTan[Sqrt[5 - 4*x - x^2]/(5 + x)]`

3.295.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {1090, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{-x^2 - 4x + 5}} dx$$

↓ 1090

$$-\frac{1}{6} \int \frac{1}{\sqrt{1 - \frac{1}{36}(-2x - 4)^2}} d(-2x - 4)$$

↓ 223

$$-\arcsin\left(\frac{1}{6}(-2x - 4)\right)$$

input `Int[1/Sqrt[5 - 4*x - x^2],x]`

output `-ArcSin[(-4 - 2*x)/6]`

3.295.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 1090 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

3.295.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.58

method	result	size
default	$\arcsin\left(\frac{2}{3} + \frac{x}{3}\right)$	7
trager	$\text{RootOf}(_Z^2 + 1) \ln\left(-\text{RootOf}(_Z^2 + 1) x + \sqrt{-x^2 - 4x + 5} - 2\text{RootOf}(_Z^2 + 1)\right)$	39

input `int(1/(-x^2-4*x+5)^(1/2),x,method=_RETURNVERBOSE)`

output `arcsin(2/3+1/3*x)`

3.295.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. 2(6) = 12.

Time = 0.25 (sec) , antiderivative size = 29, normalized size of antiderivative = 2.42

$$\int \frac{1}{\sqrt{5-4x-x^2}} dx = -\arctan\left(\frac{\sqrt{-x^2-4x+5}(x+2)}{x^2+4x-5}\right)$$

input `integrate(1/(-x^2-4*x+5)^(1/2),x, algorithm="fracas")`

output `-arctan(sqrt(-x^2 - 4*x + 5)*(x + 2)/(x^2 + 4*x - 5))`

3.295.6 Sympy [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.58

$$\int \frac{1}{\sqrt{5-4x-x^2}} dx = \text{asin}\left(\frac{x}{3} + \frac{2}{3}\right)$$

input `integrate(1/(-x**2-4*x+5)**(1/2),x)`

output `asin(x/3 + 2/3)`

3.295.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt{5-4x-x^2}} dx = -\arcsin\left(-\frac{1}{3}x - \frac{2}{3}\right)$$

input `integrate(1/(-x^2-4*x+5)^(1/2),x, algorithm="maxima")`

output `-arcsin(-1/3*x - 2/3)`

3.295.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. 2(6) = 12.

Time = 0.31 (sec) , antiderivative size = 26, normalized size of antiderivative = 2.17

$$\int \frac{1}{\sqrt{5-4x-x^2}} dx = \frac{1}{2} \sqrt{-x^2-4x+5}(x+2) + \frac{9}{2} \arcsin\left(\frac{1}{3}x + \frac{2}{3}\right)$$

input `integrate(1/(-x^2-4*x+5)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(-x^2 - 4*x + 5)*(x + 2) + 9/2*arcsin(1/3*x + 2/3)`

3.295.9 Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.50

$$\int \frac{1}{\sqrt{5-4x-x^2}} dx = \operatorname{asin}\left(\frac{x}{3} + \frac{2}{3}\right)$$

input `int(1/(5 - x^2 - 4*x)^(1/2),x)`

output `asin(x/3 + 2/3)`

$$3.296 \quad \int \frac{x}{1-x^2+\sqrt{1-x^2}} dx$$

3.296.1 Optimal result	1503
3.296.2 Mathematica [A] (verified)	1503
3.296.3 Rubi [A] (verified)	1504
3.296.4 Maple [A] (verified)	1505
3.296.5 Fricas [A] (verification not implemented)	1505
3.296.6 Sympy [B] (verification not implemented)	1506
3.296.7 Maxima [A] (verification not implemented)	1506
3.296.8 Giac [A] (verification not implemented)	1506
3.296.9 Mupad [B] (verification not implemented)	1507

3.296.1 Optimal result

Integrand size = 22, antiderivative size = 16

$$\int \frac{x}{1-x^2+\sqrt{1-x^2}} dx = -\log\left(1+\sqrt{1-x^2}\right)$$

output `-ln(1+(-x^2+1)^(1/2))`

3.296.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{x}{1-x^2+\sqrt{1-x^2}} dx = -\log\left(1+\sqrt{1-x^2}\right)$$

input `Integrate[x/(1 - x^2 + Sqrt[1 - x^2]),x]`

output `-Log[1 + Sqrt[1 - x^2]]`

3.296.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$, Rules used = {2586, 7267, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x}{-x^2 + \sqrt{1-x^2} + 1} dx \\ & \quad \downarrow \text{2586} \\ & \frac{1}{2} \int \frac{1}{-x^2 + \sqrt{1-x^2} + 1} dx^2 \\ & \quad \downarrow \text{7267} \\ & - \int \frac{1}{\sqrt{1-x^2} + 1} d\sqrt{1-x^2} \\ & \quad \downarrow \text{16} \\ & -\log(\sqrt{1-x^2} + 1) \end{aligned}$$

input `Int[x/(1 - x^2 + Sqrt[1 - x^2]),x]`

output `-Log[1 + Sqrt[1 - x^2]]`

3.296.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_.)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 2586 `Int[(x_)^(m_.)/((c_) + (d_.)*(x_)^(n_) + (e_.)*Sqrt[(a_) + (b_.)*(x_)^(n_)]), x_Symbol] :> Simp[1/n Subst[Int[x^((m + 1)/n - 1)/(c + d*x + e*Sqrt[a + b*x]], x], x, x^n], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[(m + 1)/n]`

```
rule 7267 Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Si
mp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])], x
] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]]
```

3.296.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result	size
trager	$-\ln(1 + \sqrt{-x^2 + 1})$	15
default	$-\ln(x) + \sqrt{-x^2 + 1} - \operatorname{arctanh}\left(\frac{1}{\sqrt{-x^2 + 1}}\right) - \frac{\sqrt{-(1+x)^2 + 2x + 2}}{2} - \frac{\sqrt{-(-1+x)^2 - 2x + 2}}{2}$	59

```
input int(x/(1-x^2+(-x^2+1)^(1/2)),x,method=_RETURNVERBOSE)
```

```
output -ln(1+(-x^2+1)^(1/2))
```

3.296.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.31

$$\int \frac{x}{1-x^2+\sqrt{1-x^2}} dx = -\log(x) + \log\left(\frac{\sqrt{-x^2+1}-1}{x}\right)$$

```
input integrate(x/(1-x^2+(-x^2+1)^(1/2)),x, algorithm="fricas")
```

```
output -log(x) + log((sqrt(-x^2 + 1) - 1)/x)
```

3.296.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 48 vs. $2(12) = 24$.

Time = 1.22 (sec) , antiderivative size = 48, normalized size of antiderivative = 3.00

$$\int \frac{x}{1-x^2+\sqrt{1-x^2}} dx = \frac{\log(2\sqrt{1-x^2})}{2} - \frac{\log(2\sqrt{1-x^2}+2)}{2} - \frac{\log(2x^2-2\sqrt{1-x^2}-2)}{2}$$

input `integrate(x/(1-x**2+(-x**2+1)**(1/2)),x)`

output `log(2*sqrt(1 - x**2))/2 - log(2*sqrt(1 - x**2) + 2)/2 - log(2*x**2 - 2*sqrt(1 - x**2) - 2)/2`

3.296.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{x}{1-x^2+\sqrt{1-x^2}} dx = -\log(\sqrt{-x^2+1}+1)$$

input `integrate(x/(1-x^2+(-x^2+1)^(1/2)),x, algorithm="maxima")`

output `-log(sqrt(-x^2 + 1) + 1)`

3.296.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{x}{1-x^2+\sqrt{1-x^2}} dx = -\log(\sqrt{-x^2+1}+1)$$

input `integrate(x/(1-x^2+(-x^2+1)^(1/2)),x, algorithm="giac")`

output `-log(sqrt(-x^2 + 1) + 1)`

3.296.9 Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.31

$$\int \frac{x}{1-x^2+\sqrt{1-x^2}} dx = \ln \left(\sqrt{\frac{1}{x^2}-1} - \sqrt{\frac{1}{x^2}} \right) - \ln(x)$$

input `int(x/((1 - x^2)^(1/2) - x^2 + 1),x)`

output `log((1/x^2 - 1)^(1/2) - (1/x^2)^(1/2)) - log(x)`

3.297 $\int (1 + \cos(x)) \csc(x) dx$

3.297.1 Optimal result	1508
3.297.2 Mathematica [B] (verified)	1508
3.297.3 Rubi [A] (verified)	1509
3.297.4 Maple [A] (verified)	1510
3.297.5 Fricas [A] (verification not implemented)	1510
3.297.6 Sympy [B] (verification not implemented)	1511
3.297.7 Maxima [A] (verification not implemented)	1511
3.297.8 Giac [A] (verification not implemented)	1511
3.297.9 Mupad [B] (verification not implemented)	1512

3.297.1 Optimal result

Integrand size = 7, antiderivative size = 7

$$\int (1 + \cos(x)) \csc(x) dx = \log(1 - \cos(x))$$

output `ln(1-cos(x))`

3.297.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 23 vs. $2(7) = 14$.

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 3.29

$$\int (1 + \cos(x)) \csc(x) dx = -\log\left(\cos\left(\frac{x}{2}\right)\right) + \log(\cos(x)) + \log\left(\sin\left(\frac{x}{2}\right)\right) + \log(\tan(x))$$

input `Integrate[(1 + Cos[x])*Csc[x],x]`

output `-Log[Cos[x/2]] + Log[Cos[x]] + Log[Sin[x/2]] + Log[Tan[x]]`

3.297.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3146, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (\cos(x) + 1) \csc(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1 - \sin\left(x - \frac{\pi}{2}\right)}{\cos\left(x - \frac{\pi}{2}\right)} dx \\ & \quad \downarrow \text{3146} \\ & - \int \frac{1}{1 - \cos(x)} d \cos(x) \\ & \quad \downarrow \text{16} \\ & \log(1 - \cos(x)) \end{aligned}$$

input `Int[(1 + Cos[x])*Csc[x],x]`

output `Log[1 - Cos[x]]`

3.297.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3146 Int[cos[(e_.) + (f_.)*(x_.)]^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)])^(m
_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x
)^(p - 1)/2], x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && I
ntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/
2])
```

3.297.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.86

method	result	size
default	$\ln(\sin(x)) + \ln(\csc(x) - \cot(x))$	13
parts	$-\ln(\csc(x)) - \ln(\csc(x) + \cot(x))$	15
risch	$-ix + 2 \ln(e^{ix} - 1)$	16
norman	$2 \ln\left(\tan\left(\frac{x}{2}\right)\right) - \ln\left(1 + \tan^2\left(\frac{x}{2}\right)\right)$	20
parallelrisch	$2 \ln(\csc(x) - \cot(x)) - \ln\left(\frac{2}{\cos(x)+1}\right)$	23

```
input int((cos(x)+1)*csc(x),x,method=_RETURNVERBOSE)
```

```
output ln(sin(x))+ln(csc(x)-cot(x))
```

3.297.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int (1 + \cos(x)) \csc(x) dx = \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right)$$

```
input integrate((1+cos(x))*csc(x),x, algorithm="fracas")
```

```
output log(-1/2*cos(x) + 1/2)
```

3.297.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12 vs. $2(5) = 10$.

Time = 0.80 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.71

$$\int (1 + \cos(x)) \csc(x) dx = -\log(\cot(x) + \csc(x)) + \log(\sin(x))$$

input `integrate((1+cos(x))*csc(x),x)`

output `-log(cot(x) + csc(x)) + log(sin(x))`

3.297.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71

$$\int (1 + \cos(x)) \csc(x) dx = \log(\cos(x) - 1)$$

input `integrate((1+cos(x))*csc(x),x, algorithm="maxima")`

output `log(cos(x) - 1)`

3.297.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int (1 + \cos(x)) \csc(x) dx = \log(-\cos(x) + 1)$$

input `integrate((1+cos(x))*csc(x),x, algorithm="giac")`

output `log(-cos(x) + 1)`

3.297.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71

$$\int (1 + \cos(x)) \csc(x) dx = \ln(\cos(x) - 1)$$

input `int((cos(x) + 1)/sin(x),x)`

output `log(cos(x) - 1)`

3.298 $\int \frac{e^x}{-1+e^{2x}} dx$

3.298.1 Optimal result	1513
3.298.2 Mathematica [A] (verified)	1513
3.298.3 Rubi [A] (verified)	1514
3.298.4 Maple [A] (verified)	1515
3.298.5 Fricas [B] (verification not implemented)	1515
3.298.6 Sympy [B] (verification not implemented)	1515
3.298.7 Maxima [B] (verification not implemented)	1516
3.298.8 Giac [B] (verification not implemented)	1516
3.298.9 Mupad [B] (verification not implemented)	1516

3.298.1 Optimal result

Integrand size = 13, antiderivative size = 6

$$\int \frac{e^x}{-1+e^{2x}} dx = -\operatorname{arctanh}(e^x)$$

output `-arctanh(exp(x))`

3.298.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{-1+e^{2x}} dx = -\operatorname{arctanh}(e^x)$$

input `Integrate[E^x/(-1 + E^(2*x)), x]`

output `-ArcTanh[E^x]`

3.298.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {2679, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^x}{e^{2x} - 1} dx$$

↓ 2679

$$\int \frac{1}{e^{2x} - 1} de^x$$

↓ 220

$$-\operatorname{arctanh}(e^x)$$

input `Int [E^x/(-1 + E^(2*x)), x]`

output `-ArcTanh[E^x]`

3.298.3.1 Defintions of rubi rules used

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 2679 `Int[((a_) + (b_.)*(F_)^((e_.)*((c_.) + (d_.)*(x_))))^(p_.)*(G_)^((h_.)*((f_.) + (g_.)*(x_))), x_Symbol] := With[{m = FullSimplify[d*e*(Log[F]/(g*h*Log[G]))}], Simp[Denominator[m]/(g*h*Log[G]) Subst[Int[x^(Denominator[m] - 1)*(a + b*F^(c*e - d*e*(f/g))*x^Numerator[m])^p, x], x, G^(h*((f + g*x)/Denominator[m])], x] /; LtQ[m, -1] || GtQ[m, 1]] /; FreeQ[{F, G, a, b, c, d, e, f, g, h, p}, x]`

3.298.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

method	result	size
default	$-\operatorname{arctanh}(e^x)$	6
norman	$\frac{\ln(-1+e^x)}{2} - \frac{\ln(1+e^x)}{2}$	16
risch	$\frac{\ln(-1+e^x)}{2} - \frac{\ln(1+e^x)}{2}$	16

input `int(exp(x)/(exp(2*x)-1),x,method=_RETURNVERBOSE)`

output `-arctanh(exp(x))`

3.298.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. 2(5) = 10.

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 2.50

$$\int \frac{e^x}{-1+e^{2x}} dx = -\frac{1}{2} \log(e^x + 1) + \frac{1}{2} \log(e^x - 1)$$

input `integrate(exp(x)/(-1+exp(2*x)),x, algorithm="fracas")`

output `-1/2*log(e^x + 1) + 1/2*log(e^x - 1)`

3.298.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. 2(5) = 10.

Time = 0.05 (sec) , antiderivative size = 15, normalized size of antiderivative = 2.50

$$\int \frac{e^x}{-1+e^{2x}} dx = \frac{\log(e^x - 1)}{2} - \frac{\log(e^x + 1)}{2}$$

input `integrate(exp(x)/(-1+exp(2*x)),x)`

output `log(exp(x) - 1)/2 - log(exp(x) + 1)/2`

3.298.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(5) = 10$.

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 2.50

$$\int \frac{e^x}{-1 + e^{2x}} dx = -\frac{1}{2} \log(e^x + 1) + \frac{1}{2} \log(e^x - 1)$$

input `integrate(exp(x)/(-1+exp(2*x)),x, algorithm="maxima")`

output `-1/2*log(e^x + 1) + 1/2*log(e^x - 1)`

3.298.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 16 vs. $2(5) = 10$.

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 2.67

$$\int \frac{e^x}{-1 + e^{2x}} dx = -\frac{1}{2} \log(e^x + 1) + \frac{1}{2} \log(|e^x - 1|)$$

input `integrate(exp(x)/(-1+exp(2*x)),x, algorithm="giac")`

output `-1/2*log(e^x + 1) + 1/2*log(abs(e^x - 1))`

3.298.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 2.50

$$\int \frac{e^x}{-1 + e^{2x}} dx = \frac{\ln(e^x - 1)}{2} - \frac{\ln(e^x + 1)}{2}$$

input `int(exp(x)/(exp(2*x) - 1),x)`

output `log(exp(x) - 1)/2 - log(exp(x) + 1)/2`

3.299 $\int \frac{1}{-8+x^3} dx$

3.299.1 Optimal result	1517
3.299.2 Mathematica [A] (verified)	1517
3.299.3 Rubi [A] (verified)	1518
3.299.4 Maple [A] (verified)	1520
3.299.5 Fricas [A] (verification not implemented)	1520
3.299.6 Sympy [A] (verification not implemented)	1521
3.299.7 Maxima [A] (verification not implemented)	1521
3.299.8 Giac [A] (verification not implemented)	1521
3.299.9 Mupad [B] (verification not implemented)	1522

3.299.1 Optimal result

Integrand size = 7, antiderivative size = 43

$$\int \frac{1}{-8+x^3} dx = -\frac{\arctan\left(\frac{1+x}{\sqrt{3}}\right)}{4\sqrt{3}} + \frac{1}{12} \log(2-x) - \frac{1}{24} \log(4+2x+x^2)$$

output `1/12*ln(2-x)-1/24*ln(x^2+2*x+4)-1/12*arctan(1/3*(1+x)*3^(1/2))*3^(1/2)`

3.299.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int \frac{1}{-8+x^3} dx = -\frac{\arctan\left(\frac{1+x}{\sqrt{3}}\right)}{4\sqrt{3}} + \frac{1}{12} \log(2-x) - \frac{1}{24} \log(4+2x+x^2)$$

input `Integrate[(-8 + x^3)^(-1),x]`

output `-1/4*ArcTan[(1 + x)/Sqrt[3]]/Sqrt[3] + Log[2 - x]/12 - Log[4 + 2*x + x^2]/24`

3.299.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.19, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.143$, Rules used = {750, 16, 25, 1142, 27, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^3 - 8} dx \\
 & \quad \downarrow \text{750} \\
 & \frac{1}{12} \int -\frac{x+4}{x^2+2x+4} dx + \frac{1}{12} \int \frac{1}{x-2} dx \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{12} \int -\frac{x+4}{x^2+2x+4} dx + \frac{1}{12} \log(2-x) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{12} \log(2-x) - \frac{1}{12} \int \frac{x+4}{x^2+2x+4} dx \\
 & \quad \downarrow \text{1142} \\
 & \frac{1}{12} \left(-3 \int \frac{1}{x^2+2x+4} dx - \frac{1}{2} \int \frac{2(x+1)}{x^2+2x+4} dx \right) + \frac{1}{12} \log(2-x) \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{12} \left(-3 \int \frac{1}{x^2+2x+4} dx - \int \frac{x+1}{x^2+2x+4} dx \right) + \frac{1}{12} \log(2-x) \\
 & \quad \downarrow \text{1083} \\
 & \frac{1}{12} \left(6 \int \frac{1}{-(2x+2)^2 - 12} d(2x+2) - \int \frac{x+1}{x^2+2x+4} dx \right) + \frac{1}{12} \log(2-x) \\
 & \quad \downarrow \text{217} \\
 & \frac{1}{12} \left(- \int \frac{x+1}{x^2+2x+4} dx - \sqrt{3} \arctan \left(\frac{2x+2}{2\sqrt{3}} \right) \right) + \frac{1}{12} \log(2-x) \\
 & \quad \downarrow \text{1103} \\
 & \frac{1}{12} \left(-\sqrt{3} \arctan \left(\frac{2x+2}{2\sqrt{3}} \right) - \frac{1}{2} \log(x^2+2x+4) \right) + \frac{1}{12} \log(2-x)
 \end{aligned}$$

input `Int[(-8 + x^3)^(-1), x]`

output `Log[2 - x]/12 + (-(Sqrt[3]*ArcTan[(2 + 2*x)/(2*Sqrt[3])]) - Log[4 + 2*x + x^2]/2)/12`

3.299.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 750 `Int[((a_) + (b_.)*(x_)^3)^(-1), x_Symbol] := Simp[1/(3*Rt[a, 3]^2) Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Simp[1/(3*Rt[a, 3]^2) Int[(2*Rt[a, 3] - Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]`

rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`


```
rule 1142 Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c)
Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]
```

3.299.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.77

method	result	size
risch	$-\frac{\ln(x^2+2x+4)}{24} - \frac{\arctan\left(\frac{(1+x)\sqrt{3}}{3}\right)\sqrt{3}}{12} + \frac{\ln(-2+x)}{12}$	33
default	$-\frac{\ln(x^2+2x+4)}{24} - \frac{\sqrt{3} \arctan\left(\frac{(2x+2)\sqrt{3}}{6}\right)}{12} + \frac{\ln(-2+x)}{12}$	35
meijerg	$\frac{x \left(\ln\left(1 - \frac{(x^3)^{\frac{1}{3}}}{2}\right) - \frac{\ln\left(1 + \frac{(x^3)^{\frac{1}{3}}}{2} + \frac{(x^3)^{\frac{2}{3}}}{4}\right)}{2} - \sqrt{3} \arctan\left(\frac{\sqrt{3}(x^3)^{\frac{1}{3}}}{4+(x^3)^{\frac{1}{3}}}\right) \right)}{12(x^3)^{\frac{1}{3}}}$	66

```
input int(1/(x^3-8),x,method=_RETURNVERBOSE)
```

```
output -1/24*ln(x^2+2*x+4)-1/12*arctan(1/3*(1+x)*3^(1/2))*3^(1/2)+1/12*ln(-2+x)
```

3.299.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.74

$$\int \frac{1}{-8+x^3} dx = -\frac{1}{12} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(x+1)\right) - \frac{1}{24} \log(x^2+2x+4) + \frac{1}{12} \log(x-2)$$

```
input integrate(1/(x^3-8),x, algorithm="fricas")
```

```
output -1/12*sqrt(3)*arctan(1/3*sqrt(3)*(x + 1)) - 1/24*log(x^2 + 2*x + 4) + 1/12
*log(x - 2)
```

3.299.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.95

$$\int \frac{1}{-8+x^3} dx = \frac{\log(x-2)}{12} - \frac{\log(x^2+2x+4)}{24} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{\sqrt{3}x}{3} + \frac{\sqrt{3}}{3}\right)}{12}$$

input `integrate(1/(x**3-8),x)`output `log(x - 2)/12 - log(x**2 + 2*x + 4)/24 - sqrt(3)*atan(sqrt(3)*x/3 + sqrt(3)/3)/12`**3.299.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.74

$$\int \frac{1}{-8+x^3} dx = -\frac{1}{12} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(x+1)\right) - \frac{1}{24} \log(x^2+2x+4) + \frac{1}{12} \log(x-2)$$

input `integrate(1/(x^3-8),x, algorithm="maxima")`output `-1/12*sqrt(3)*arctan(1/3*sqrt(3)*(x + 1)) - 1/24*log(x^2 + 2*x + 4) + 1/12*log(x - 2)`**3.299.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.77

$$\int \frac{1}{-8+x^3} dx = -\frac{1}{12} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(x+1)\right) - \frac{1}{24} \log(x^2+2x+4) + \frac{1}{12} \log(|x-2|)$$

input `integrate(1/(x^3-8),x, algorithm="giac")`output `-1/12*sqrt(3)*arctan(1/3*sqrt(3)*(x + 1)) - 1/24*log(x^2 + 2*x + 4) + 1/12*log(abs(x - 2))`

3.299.9 Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.07

$$\int \frac{1}{-8+x^3} dx = \frac{\ln(x-2)}{12} + \ln(x+1-\sqrt{3}i) \left(-\frac{1}{24} + \frac{\sqrt{3}i}{24} \right) - \ln(x+1+\sqrt{3}i) \left(\frac{1}{24} + \frac{\sqrt{3}i}{24} \right)$$

input `int(1/(x^3 - 8),x)`output `log(x - 2)/12 + log(x - 3^(1/2)*1i + 1)*((3^(1/2)*1i)/24 - 1/24) - log(x + 3^(1/2)*1i + 1)*((3^(1/2)*1i)/24 + 1/24)`

3.300 $\int x^5 \cosh(x) dx$

3.300.1 Optimal result	1523
3.300.2 Mathematica [A] (verified)	1523
3.300.3 Rubi [C] (verified)	1524
3.300.4 Maple [A] (verified)	1526
3.300.5 Fricas [A] (verification not implemented)	1527
3.300.6 Sympy [A] (verification not implemented)	1527
3.300.7 Maxima [A] (verification not implemented)	1527
3.300.8 Giac [A] (verification not implemented)	1528
3.300.9 Mupad [B] (verification not implemented)	1528

3.300.1 Optimal result

Integrand size = 6, antiderivative size = 37

$$\int x^5 \cosh(x) dx = -120 \cosh(x) - 60x^2 \cosh(x) - 5x^4 \cosh(x) + 120x \sinh(x) + 20x^3 \sinh(x) + x^5 \sinh(x)$$

output `-120*cosh(x)-60*x^2*cosh(x)-5*x^4*cosh(x)+120*x*sinh(x)+20*x^3*sinh(x)+x^5*sinh(x)`

3.300.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.78

$$\int x^5 \cosh(x) dx = -5(24 + 12x^2 + x^4) \cosh(x) + x(120 + 20x^2 + x^4) \sinh(x)$$

input `Integrate[x^5*Cosh[x],x]`

output `-5*(24 + 12*x^2 + x^4)*Cosh[x] + x*(120 + 20*x^2 + x^4)*Sinh[x]`

3.300.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.45 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.59, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 3.000$, Rules used = {3042, 3777, 26, 3042, 26, 3777, 3042, 3777, 26, 3042, 26, 3777, 3042, 3777, 26, 3042, 26, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^5 \cosh(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x^5 \sin\left(\frac{\pi}{2} + ix\right) dx \\
 & \quad \downarrow \text{3777} \\
 & x^5 \sinh(x) - 5i \int -ix^4 \sinh(x) dx \\
 & \quad \downarrow \text{26} \\
 & x^5 \sinh(x) - 5 \int x^4 \sinh(x) dx \\
 & \quad \downarrow \text{3042} \\
 & x^5 \sinh(x) - 5 \int -ix^4 \sin(ix) dx \\
 & \quad \downarrow \text{26} \\
 & x^5 \sinh(x) + 5i \int x^4 \sin(ix) dx \\
 & \quad \downarrow \text{3777} \\
 & x^5 \sinh(x) + 5i \left(ix^4 \cosh(x) - 4i \int x^3 \cosh(x) dx \right) \\
 & \quad \downarrow \text{3042} \\
 & x^5 \sinh(x) + 5i \left(ix^4 \cosh(x) - 4i \int x^3 \sin\left(ix + \frac{\pi}{2}\right) dx \right) \\
 & \quad \downarrow \text{3777} \\
 & x^5 \sinh(x) + 5i \left(ix^4 \cosh(x) - 4i \left(x^3 \sinh(x) - 3i \int -ix^2 \sinh(x) dx \right) \right)
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 26 \\
& x^5 \sinh(x) + 5i \left(ix^4 \cosh(x) - 4i \left(x^3 \sinh(x) - 3 \int x^2 \sinh(x) dx \right) \right) \\
& \downarrow 3042 \\
& x^5 \sinh(x) + 5i \left(ix^4 \cosh(x) - 4i \left(x^3 \sinh(x) - 3 \int -ix^2 \sin(ix) dx \right) \right) \\
& \downarrow 26 \\
& x^5 \sinh(x) + 5i \left(ix^4 \cosh(x) - 4i \left(x^3 \sinh(x) + 3i \int x^2 \sin(ix) dx \right) \right) \\
& \downarrow 3777 \\
& x^5 \sinh(x) + 5i \left(ix^4 \cosh(x) - 4i \left(x^3 \sinh(x) + 3i \left(ix^2 \cosh(x) - 2i \int x \cosh(x) dx \right) \right) \right) \\
& \downarrow 3042 \\
& x^5 \sinh(x) + 5i \left(ix^4 \cosh(x) - 4i \left(x^3 \sinh(x) + 3i \left(ix^2 \cosh(x) - 2i \int x \sin \left(ix + \frac{\pi}{2} \right) dx \right) \right) \right) \\
& \downarrow 3777 \\
& x^5 \sinh(x) + \\
& 5i \left(ix^4 \cosh(x) - 4i \left(x^3 \sinh(x) + 3i \left(ix^2 \cosh(x) - 2i(x \sinh(x) - i \int -i \sinh(x) dx) \right) \right) \right) \\
& \downarrow 26 \\
& x^5 \sinh(x) + 5i \left(ix^4 \cosh(x) - 4i \left(x^3 \sinh(x) + 3i \left(ix^2 \cosh(x) - 2i(x \sinh(x) - \int \sinh(x) dx) \right) \right) \right) \\
& \downarrow 3042 \\
& x^5 \sinh(x) + \\
& 5i \left(ix^4 \cosh(x) - 4i \left(x^3 \sinh(x) + 3i \left(ix^2 \cosh(x) - 2i(x \sinh(x) - \int -i \sin(ix) dx) \right) \right) \right) \\
& \downarrow 26 \\
& x^5 \sinh(x) + 5i \left(ix^4 \cosh(x) - 4i \left(x^3 \sinh(x) + 3i \left(ix^2 \cosh(x) - 2i(x \sinh(x) + i \int \sin(ix) dx) \right) \right) \right) \\
& \downarrow 3118 \\
& x^5 \sinh(x) + 5i \left(ix^4 \cosh(x) - 4i \left(x^3 \sinh(x) + 3i \left(ix^2 \cosh(x) - 2i(x \sinh(x) - \cosh(x)) \right) \right) \right)
\end{aligned}$$

input `Int [x^5*Cosh[x], x]`

output $x^5 \sinh(x) + (5I)(I x^4 \cosh(x) - (4I)(x^3 \sinh(x) + (3I)(I x^2 \cosh(x) - (2I)(-\cosh(x) + x \sinh(x))))))$

3.300.3.1 Defintions of rubi rules used

rule 26 $\text{Int}[(\text{Complex}[0, a_])*(F x_), x_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3118 $\text{Int}[\sin[(c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Cos}[c + d*x]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

rule 3777 $\text{Int}[(c_.) + (d_.)*(x_)]^{(m_.)} \sin[(e_.) + (f_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[(-c + d*x)^m * (\text{Cos}[e + f*x]/f), x] + \text{Simp}[d*(m/f) \text{Int}[(c + d*x)^{(m-1)} * \text{Cos}[e + f*x], x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \ \&\& \ \text{GtQ}[m, 0]$

3.300.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.89

method	result
parallelrisch	$(-5x^4 - 60x^2 - 120) \cosh(x) - 120 + (x^5 + 20x^3 + 120x) \sinh(x)$
default	$-120 \cosh(x) - 60x^2 \cosh(x) - 5x^4 \cosh(x) + 120x \sinh(x) + 20x^3 \sinh(x) + x^5 \sinh(x)$
parts	$-120 \cosh(x) - 60x^2 \cosh(x) - 5x^4 \cosh(x) + 120x \sinh(x) + 20x^3 \sinh(x) + x^5 \sinh(x)$
meijerg	$-32\sqrt{\pi} \left(-\frac{15}{4\sqrt{\pi}} + \frac{(\frac{15}{8}x^4 + \frac{45}{2}x^2 + 45) \cosh(x)}{12\sqrt{\pi}} - \frac{x(\frac{3}{8}x^4 + \frac{15}{2}x^2 + 45) \sinh(x)}{12\sqrt{\pi}} \right)$
risch	$(10x^3 - 30x^2 + 60x - 60 - \frac{5}{2}x^4 + \frac{1}{2}x^5) e^x + (-10x^3 - 30x^2 - 60x - 60 - \frac{5}{2}x^4 - \frac{1}{2}x^5) e^{-x}$

input $\text{int}(x^5 * \cosh(x), x, \text{method} = _RETURNVERBOSE)$

output $(-5*x^4 - 60*x^2 - 120) * \cosh(x) - 120 + (x^5 + 20*x^3 + 120*x) * \sinh(x)$

3.300.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.81

$$\int x^5 \cosh(x) dx = -5(x^4 + 12x^2 + 24) \cosh(x) + (x^5 + 20x^3 + 120x) \sinh(x)$$

input `integrate(x^5*cosh(x),x, algorithm="fricas")`output `-5*(x^4 + 12*x^2 + 24)*cosh(x) + (x^5 + 20*x^3 + 120*x)*sinh(x)`**3.300.6 Sympy [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.14

$$\int x^5 \cosh(x) dx = x^5 \sinh(x) - 5x^4 \cosh(x) + 20x^3 \sinh(x) - 60x^2 \cosh(x) + 120x \sinh(x) - 120 \cosh(x)$$

input `integrate(x**5*cosh(x),x)`output `x**5*sinh(x) - 5*x**4*cosh(x) + 20*x**3*sinh(x) - 60*x**2*cosh(x) + 120*x*sinh(x) - 120*cosh(x)`**3.300.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 74, normalized size of antiderivative = 2.00

$$\int x^5 \cosh(x) dx = \frac{1}{6} x^6 \cosh(x) - \frac{1}{12} (x^6 + 6x^5 + 30x^4 + 120x^3 + 360x^2 + 720x + 720) e^{-x} - \frac{1}{12} (x^6 - 6x^5 + 30x^4 - 120x^3 + 360x^2 - 720x + 720) e^x$$

input `integrate(x^5*cosh(x),x, algorithm="maxima")`output `1/6*x^6*cosh(x) - 1/12*(x^6 + 6*x^5 + 30*x^4 + 120*x^3 + 360*x^2 + 720*x + 720)*e^(-x) - 1/12*(x^6 - 6*x^5 + 30*x^4 - 120*x^3 + 360*x^2 - 720*x + 720)*e^x`

3.300.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.54

$$\int x^5 \cosh(x) dx = -\frac{1}{2} (x^5 + 5x^4 + 20x^3 + 60x^2 + 120x + 120)e^{(-x)} + \frac{1}{2} (x^5 - 5x^4 + 20x^3 - 60x^2 + 120x - 120)e^x$$

input `integrate(x^5*cosh(x),x, algorithm="giac")`output `-1/2*(x^5 + 5*x^4 + 20*x^3 + 60*x^2 + 120*x + 120)*e^(-x) + 1/2*(x^5 - 5*x^4 + 20*x^3 - 60*x^2 + 120*x - 120)*e^x`**3.300.9 Mupad [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

$$\int x^5 \cosh(x) dx = 20x^3 \sinh(x) - 60x^2 \cosh(x) - 5x^4 \cosh(x) - 120 \cosh(x) + x^5 \sinh(x) + 120x \sinh(x)$$

input `int(x^5*cosh(x),x)`output `20*x^3*sinh(x) - 60*x^2*cosh(x) - 5*x^4*cosh(x) - 120*cosh(x) + x^5*sinh(x) + 120*x*sinh(x)`

3.301 $\int \csc(x) \log(\tan(x)) \sec(x) dx$

3.301.1 Optimal result	1529
3.301.2 Mathematica [A] (verified)	1529
3.301.3 Rubi [A] (verified)	1530
3.301.4 Maple [A] (verified)	1530
3.301.5 Fricas [A] (verification not implemented)	1531
3.301.6 Sympy [F]	1531
3.301.7 Maxima [A] (verification not implemented)	1531
3.301.8 Giac [A] (verification not implemented)	1532
3.301.9 Mupad [B] (verification not implemented)	1532

3.301.1 Optimal result

Integrand size = 8, antiderivative size = 9

$$\int \csc(x) \log(\tan(x)) \sec(x) dx = \frac{1}{2} \log^2(\tan(x))$$

output `1/2*ln(tan(x))^2`

3.301.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \csc(x) \log(\tan(x)) \sec(x) dx = \frac{1}{2} \log^2(\tan(x))$$

input `Integrate[Csc[x]*Log[Tan[x]]*Sec[x],x]`

output `Log[Tan[x]]^2/2`

3.301.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {7237}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \csc(x) \sec(x) \log(\tan(x)) dx$$

$$\downarrow 7237$$

$$\frac{1}{2} \log^2(\tan(x))$$

input `Int [Csc [x] *Log [Tan [x]] *Sec [x] , x]`

output `Log [Tan [x]] ^2/2`

3.301.3.1 Defintions of rubi rules used

rule 7237 `Int[(u_)*(y_)^(m_.), x_Symbol] :> With[{q = DerivativeDivides[y, u, x]}, Si
mp[q*(y^(m + 1)/(m + 1)), x] /; !FalseQ[q]] /; FreeQ[m, x] && NeQ[m, -1]`

3.301.4 Maple [A] (verified)

Time = 1.14 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

method	result	size
derivativedivides	$\frac{\ln(\tan(x))^2}{2}$	8
default	$\frac{\ln(\tan(x))^2}{2}$	8
risch	Expression too large to display	764

input `int(ln(tan(x))/cos(x)/sin(x),x,method=_RETURNVERBOSE)`

output `1/2*ln(tan(x))^2`

3.301.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.33

$$\int \csc(x) \log(\tan(x)) \sec(x) dx = \frac{1}{2} \log\left(\frac{\sin(x)}{\cos(x)}\right)^2$$

input `integrate(log(tan(x))/cos(x)/sin(x),x, algorithm="fricas")`output `1/2*log(sin(x)/cos(x))^2`**3.301.6 Sympy [F]**

$$\int \csc(x) \log(\tan(x)) \sec(x) dx = \int \frac{\log(\tan(x))}{\sin(x) \cos(x)} dx$$

input `integrate(ln(tan(x))/cos(x)/sin(x),x)`output `Integral(log(tan(x))/(sin(x)*cos(x)), x)`**3.301.7 Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \csc(x) \log(\tan(x)) \sec(x) dx = \frac{1}{2} \log(\tan(x))^2$$

input `integrate(log(tan(x))/cos(x)/sin(x),x, algorithm="maxima")`output `1/2*log(tan(x))^2`

3.301.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \csc(x) \log(\tan(x)) \sec(x) dx = \frac{1}{2} \log(\tan(x))^2$$

input `integrate(log(tan(x))/cos(x)/sin(x),x, algorithm="giac")`output `1/2*log(tan(x))^2`**3.301.9 Mupad [B] (verification not implemented)**

Time = 2.66 (sec) , antiderivative size = 27, normalized size of antiderivative = 3.00

$$\int \csc(x) \log(\tan(x)) \sec(x) dx = \frac{\ln\left(-\frac{e^{x2i} 1i-i}{e^{x2i}+1}\right)^2}{2}$$

input `int(log(tan(x))/(cos(x)*sin(x)),x)`output `log(-(exp(x*2i)*1i - 1i)/(exp(x*2i) + 1))^2/2`

3.302 $\int (-2x + x^2 + x^3) dx$

3.302.1 Optimal result	1533
3.302.2 Mathematica [A] (verified)	1533
3.302.3 Rubi [A] (verified)	1534
3.302.4 Maple [A] (verified)	1534
3.302.5 Fricas [A] (verification not implemented)	1535
3.302.6 Sympy [A] (verification not implemented)	1535
3.302.7 Maxima [A] (verification not implemented)	1535
3.302.8 Giac [A] (verification not implemented)	1536
3.302.9 Mupad [B] (verification not implemented)	1536

3.302.1 Optimal result

Integrand size = 10, antiderivative size = 20

$$\int (-2x + x^2 + x^3) dx = -x^2 + \frac{x^3}{3} + \frac{x^4}{4}$$

output `-x^2+1/3*x^3+1/4*x^4`

3.302.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (-2x + x^2 + x^3) dx = -x^2 + \frac{x^3}{3} + \frac{x^4}{4}$$

input `Integrate[-2*x + x^2 + x^3,x]`

output `-x^2 + x^3/3 + x^4/4`

3.302.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x^3 + x^2 - 2x) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^4}{4} + \frac{x^3}{3} - x^2$$

input `Int[-2*x + x^2 + x^3,x]`

output `-x^2 + x^3/3 + x^4/4`

3.302.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.302.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

method	result	size
gospers	$\frac{x^2(3x^2+4x-12)}{12}$	16
default	$-x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4$	17
norman	$-x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4$	17
risch	$-x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4$	17
parallelrisch	$-x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4$	17
parts	$-x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4$	17

input `int(x^3+x^2-2*x,x,method=_RETURNVERBOSE)`

output `1/12*x^2*(3*x^2+4*x-12)`

3.302.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int (-2x + x^2 + x^3) dx = \frac{1}{4}x^4 + \frac{1}{3}x^3 - x^2$$

input `integrate(x^3+x^2-2*x,x, algorithm="fricas")`

output `1/4*x^4 + 1/3*x^3 - x^2`

3.302.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.60

$$\int (-2x + x^2 + x^3) dx = \frac{x^4}{4} + \frac{x^3}{3} - x^2$$

input `integrate(x**3+x**2-2*x,x)`

output `x**4/4 + x**3/3 - x**2`

3.302.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int (-2x + x^2 + x^3) dx = \frac{1}{4}x^4 + \frac{1}{3}x^3 - x^2$$

input `integrate(x^3+x^2-2*x,x, algorithm="maxima")`

output `1/4*x^4 + 1/3*x^3 - x^2`

3.302.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int (-2x + x^2 + x^3) dx = \frac{1}{4}x^4 + \frac{1}{3}x^3 - x^2$$

input `integrate(x^3+x^2-2*x,x, algorithm="giac")`

output `1/4*x^4 + 1/3*x^3 - x^2`

3.302.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int (-2x + x^2 + x^3) dx = \frac{x^2(3x^2 + 4x - 12)}{12}$$

input `int(x^2 - 2*x + x^3,x)`

output `(x^2*(4*x + 3*x^2 - 12))/12`

3.303 $\int \frac{1+e^x}{1-e^x} dx$

3.303.1 Optimal result	1537
3.303.2 Mathematica [A] (verified)	1537
3.303.3 Rubi [A] (verified)	1538
3.303.4 Maple [A] (verified)	1539
3.303.5 Fricas [A] (verification not implemented)	1539
3.303.6 Sympy [A] (verification not implemented)	1540
3.303.7 Maxima [A] (verification not implemented)	1540
3.303.8 Giac [A] (verification not implemented)	1540
3.303.9 Mupad [B] (verification not implemented)	1541

3.303.1 Optimal result

Integrand size = 15, antiderivative size = 12

$$\int \frac{1+e^x}{1-e^x} dx = x - 2 \log(1 - e^x)$$

output `x-2*ln(1-exp(x))`

3.303.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

$$\int \frac{1+e^x}{1-e^x} dx = \log(e^x) - 2 \log(-1 + e^x)$$

input `Integrate[(1 + E^x)/(1 - E^x), x]`

output `Log[E^x] - 2*Log[-1 + E^x]`

3.303.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.25, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2720, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{e^x + 1}{1 - e^x} dx \\
 & \quad \downarrow \text{2720} \\
 & \int \frac{e^{-x}(e^x + 1)}{1 - e^x} de^x \\
 & \quad \downarrow \text{86} \\
 & \int \left(e^{-x} - \frac{2}{e^x - 1} \right) de^x \\
 & \quad \downarrow \text{2009} \\
 & \log(e^x) - 2 \log(1 - e^x)
 \end{aligned}$$

input `Int[(1 + E^x)/(1 - E^x),x]`

output `Log[E^x] - 2*Log[1 - E^x]`

3.303.3.1 Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.303.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

method	result	size
norman	$x - 2 \ln(-1 + e^x)$	10
risch	$x - 2 \ln(-1 + e^x)$	10
parallelrisch	$x - 2 \ln(-1 + e^x)$	10
derivativdivides	$-2 \ln(-1 + e^x) + \ln(e^x)$	12
default	$-2 \ln(-1 + e^x) + \ln(e^x)$	12

```
input int((1+exp(x))/(1-exp(x)),x,method=_RETURNVERBOSE)
```

```
output x-2*ln(-1+exp(x))
```

3.303.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{1 + e^x}{1 - e^x} dx = x - 2 \log(e^x - 1)$$

```
input integrate((1+exp(x))/(1-exp(x)),x, algorithm="fricas")
```

```
output x - 2*log(e^x - 1)
```

3.303.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1 + e^x}{1 - e^x} dx = x - 2 \log(e^x - 1)$$

input `integrate((1+exp(x))/(1-exp(x)),x)`output `x - 2*log(exp(x) - 1)`**3.303.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{1 + e^x}{1 - e^x} dx = x - 2 \log(e^x - 1)$$

input `integrate((1+exp(x))/(1-exp(x)),x, algorithm="maxima")`output `x - 2*log(e^x - 1)`**3.303.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1 + e^x}{1 - e^x} dx = x - 2 \log(|e^x - 1|)$$

input `integrate((1+exp(x))/(1-exp(x)),x, algorithm="giac")`output `x - 2*log(abs(e^x - 1))`

3.303.9 Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{1 + e^x}{1 - e^x} dx = x - 2 \ln(e^x - 1)$$

input `int(-(exp(x) + 1)/(exp(x) - 1),x)`

output `x - 2*log(exp(x) - 1)`

3.304 $\int \frac{x}{(1+x^2)(4+x^2)} dx$

3.304.1 Optimal result	1542
3.304.2 Mathematica [A] (verified)	1542
3.304.3 Rubi [A] (verified)	1543
3.304.4 Maple [A] (verified)	1544
3.304.5 Fricas [A] (verification not implemented)	1544
3.304.6 Sympy [A] (verification not implemented)	1545
3.304.7 Maxima [A] (verification not implemented)	1545
3.304.8 Giac [A] (verification not implemented)	1545
3.304.9 Mupad [B] (verification not implemented)	1546

3.304.1 Optimal result

Integrand size = 16, antiderivative size = 21

$$\int \frac{x}{(1+x^2)(4+x^2)} dx = \frac{1}{6} \log(1+x^2) - \frac{1}{6} \log(4+x^2)$$

output `1/6*ln(x^2+1)-1/6*ln(x^2+4)`

3.304.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{x}{(1+x^2)(4+x^2)} dx = \frac{1}{6} \log(1+x^2) - \frac{1}{6} \log(4+x^2)$$

input `Integrate[x/((1 + x^2)*(4 + x^2)), x]`

output `Log[1 + x^2]/6 - Log[4 + x^2]/6`

3.304.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.19, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {353, 47, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x}{(x^2 + 1)(x^2 + 4)} dx \\ & \quad \downarrow \text{353} \\ & \frac{1}{2} \int \frac{1}{(x^2 + 1)(x^2 + 4)} dx^2 \\ & \quad \downarrow \text{47} \\ & \frac{1}{2} \left(\frac{1}{3} \int \frac{1}{x^2 + 1} dx^2 - \frac{1}{3} \int \frac{1}{x^2 + 4} dx^2 \right) \\ & \quad \downarrow \text{16} \\ & \frac{1}{2} \left(\frac{1}{3} \log(x^2 + 1) - \frac{1}{3} \log(x^2 + 4) \right) \end{aligned}$$

input `Int[x/((1 + x^2)*(4 + x^2)),x]`

output `(Log[1 + x^2]/3 - Log[4 + x^2]/3)/2`

3.304.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`


```
rule 353 Int[(x_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol]
  :> Simp[1/2 Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[
  {a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0]
```

3.304.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

method	result	size
default	$\frac{\ln(x^2+1)}{6} - \frac{\ln(x^2+4)}{6}$	18
norman	$\frac{\ln(x^2+1)}{6} - \frac{\ln(x^2+4)}{6}$	18
risch	$\frac{\ln(x^2+1)}{6} - \frac{\ln(x^2+4)}{6}$	18
parallelrisc	$\frac{\ln(x^2+1)}{6} - \frac{\ln(x^2+4)}{6}$	18

```
input int(x/(x^2+1)/(x^2+4),x,method=_RETURNVERBOSE)
```

```
output 1/6*ln(x^2+1)-1/6*ln(x^2+4)
```

3.304.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{x}{(1+x^2)(4+x^2)} dx = -\frac{1}{6} \log(x^2+4) + \frac{1}{6} \log(x^2+1)$$

```
input integrate(x/(x^2+1)/(x^2+4),x, algorithm="fracas")
```

```
output -1/6*log(x^2 + 4) + 1/6*log(x^2 + 1)
```

3.304.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \frac{x}{(1+x^2)(4+x^2)} dx = \frac{\log(x^2+1)}{6} - \frac{\log(x^2+4)}{6}$$

input `integrate(x/(x**2+1)/(x**2+4),x)`output `log(x**2 + 1)/6 - log(x**2 + 4)/6`**3.304.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{x}{(1+x^2)(4+x^2)} dx = -\frac{1}{6} \log(x^2+4) + \frac{1}{6} \log(x^2+1)$$

input `integrate(x/(x^2+1)/(x^2+4),x, algorithm="maxima")`output `-1/6*log(x^2 + 4) + 1/6*log(x^2 + 1)`**3.304.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{x}{(1+x^2)(4+x^2)} dx = -\frac{1}{6} \log(x^2+4) + \frac{1}{6} \log(x^2+1)$$

input `integrate(x/(x^2+1)/(x^2+4),x, algorithm="giac")`output `-1/6*log(x^2 + 4) + 1/6*log(x^2 + 1)`

3.304.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{x}{(1+x^2)(4+x^2)} dx = \frac{\operatorname{atanh}\left(\frac{3x^2}{5x^2+8}\right)}{3}$$

input `int(x/((x^2 + 1)*(x^2 + 4)),x)`

output `atanh((3*x^2)/(5*x^2 + 8))/3`

3.305 $\int \frac{1}{4-5\sin(x)} dx$

3.305.1 Optimal result	1547
3.305.2 Mathematica [A] (verified)	1547
3.305.3 Rubi [A] (verified)	1548
3.305.4 Maple [A] (verified)	1549
3.305.5 Fricas [A] (verification not implemented)	1549
3.305.6 Sympy [A] (verification not implemented)	1550
3.305.7 Maxima [A] (verification not implemented)	1550
3.305.8 Giac [A] (verification not implemented)	1550
3.305.9 Mupad [B] (verification not implemented)	1551

3.305.1 Optimal result

Integrand size = 8, antiderivative size = 43

$$\int \frac{1}{4-5\sin(x)} dx = -\frac{1}{3} \log \left(\cos \left(\frac{x}{2} \right) - 2 \sin \left(\frac{x}{2} \right) \right) + \frac{1}{3} \log \left(2 \cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right)$$

output `-1/3*ln(cos(1/2*x)-2*sin(1/2*x))+1/3*ln(2*cos(1/2*x)-sin(1/2*x))`

3.305.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int \frac{1}{4-5\sin(x)} dx = -\frac{1}{3} \log \left(\cos \left(\frac{x}{2} \right) - 2 \sin \left(\frac{x}{2} \right) \right) + \frac{1}{3} \log \left(2 \cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right)$$

input `Integrate[(4 - 5*Sin[x])^(-1),x]`

output `-1/3*Log[Cos[x/2] - 2*Sin[x/2]] + Log[2*Cos[x/2] - Sin[x/2]]/3`

3.305.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.77, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{4 - 5 \sin(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{4 - 5 \sin(x)} dx \\
 & \quad \downarrow \text{3139} \\
 & 2 \int \frac{1}{4 \tan^2\left(\frac{x}{2}\right) - 10 \tan\left(\frac{x}{2}\right) + 4} d \tan\left(\frac{x}{2}\right) \\
 & \quad \downarrow \text{1081} \\
 & 8 \int \left(\frac{1}{12(1 - 2 \tan\left(\frac{x}{2}\right))} - \frac{1}{24(2 - \tan\left(\frac{x}{2}\right))} \right) d \tan\left(\frac{x}{2}\right) \\
 & \quad \downarrow \text{2009} \\
 & 8 \left(\frac{1}{24} \log\left(2 - \tan\left(\frac{x}{2}\right)\right) - \frac{1}{24} \log\left(1 - 2 \tan\left(\frac{x}{2}\right)\right) \right)
 \end{aligned}$$

input `Int[(4 - 5*Sin[x])^(-1),x]`

output `8*(-1/24*Log[1 - 2*Tan[x/2]] + Log[2 - Tan[x/2]]/24)`

3.305.3.1 Defintions of rubi rules used

rule 1081 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3139 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

3.305.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.51

method	result	size
default	$-\frac{\ln(2 \tan(\frac{x}{2}) - 1)}{3} + \frac{\ln(\tan(\frac{x}{2}) - 2)}{3}$	22
norman	$-\frac{\ln(2 \tan(\frac{x}{2}) - 1)}{3} + \frac{\ln(\tan(\frac{x}{2}) - 2)}{3}$	22
parallelrisc	$\ln\left(\left(2 \tan\left(\frac{x}{2}\right) - 4\right)^{\frac{1}{3}}\right) + \ln\left(\frac{1}{\left(2 \tan\left(\frac{x}{2}\right) - 1\right)^{\frac{1}{3}}}\right)$	24
risc	$-\frac{\ln\left(e^{ix} - \frac{3}{5} - \frac{4i}{5}\right)}{3} + \frac{\ln\left(e^{ix} + \frac{3}{5} - \frac{4i}{5}\right)}{3}$	26

input `int(1/(4-5*sin(x)),x,method=_RETURNVERBOSE)`

output `-1/3*ln(2*tan(1/2*x)-1)+1/3*ln(tan(1/2*x)-2)`

3.305.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.63

$$\int \frac{1}{4 - 5 \sin(x)} dx = \frac{1}{6} \log\left(\frac{3}{2} \cos(x) - 2 \sin(x) + \frac{5}{2}\right) - \frac{1}{6} \log\left(-\frac{3}{2} \cos(x) - 2 \sin(x) + \frac{5}{2}\right)$$

input `integrate(1/(4-5*sin(x)),x, algorithm="fracas")`

output `1/6*log(3/2*cos(x) - 2*sin(x) + 5/2) - 1/6*log(-3/2*cos(x) - 2*sin(x) + 5/2)`

3.305.6 Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.47

$$\int \frac{1}{4-5\sin(x)} dx = \frac{\log\left(\tan\left(\frac{x}{2}\right) - 2\right)}{3} - \frac{\log\left(2\tan\left(\frac{x}{2}\right) - 1\right)}{3}$$

input `integrate(1/(4-5*sin(x)),x)`output `log(tan(x/2) - 2)/3 - log(2*tan(x/2) - 1)/3`**3.305.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.70

$$\int \frac{1}{4-5\sin(x)} dx = -\frac{1}{3} \log\left(\frac{2\sin(x)}{\cos(x)+1} - 1\right) + \frac{1}{3} \log\left(\frac{\sin(x)}{\cos(x)+1} - 2\right)$$

input `integrate(1/(4-5*sin(x)),x, algorithm="maxima")`output `-1/3*log(2*sin(x)/(cos(x) + 1) - 1) + 1/3*log(sin(x)/(cos(x) + 1) - 2)`**3.305.8 Giac [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.53

$$\int \frac{1}{4-5\sin(x)} dx = -\frac{1}{3} \log\left(\left|2\tan\left(\frac{1}{2}x\right) - 1\right|\right) + \frac{1}{3} \log\left(\left|\tan\left(\frac{1}{2}x\right) - 2\right|\right)$$

input `integrate(1/(4-5*sin(x)),x, algorithm="giac")`output `-1/3*log(abs(2*tan(1/2*x) - 1)) + 1/3*log(abs(tan(1/2*x) - 2))`

3.305.9 Mupad [B] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.26

$$\int \frac{1}{4 - 5 \sin(x)} dx = -\frac{2 \operatorname{atanh}\left(\frac{4 \tan\left(\frac{x}{2}\right) - 5}{3}\right)}{3}$$

input `int(-1/(5*sin(x) - 4),x)`

output `-(2*atanh((4*tan(x/2))/3 - 5/3))/3`

3.306 $\int x\sqrt[3]{c+x} dx$

3.306.1 Optimal result	1552
3.306.2 Mathematica [A] (verified)	1552
3.306.3 Rubi [A] (verified)	1553
3.306.4 Maple [A] (verified)	1554
3.306.5 Fricas [A] (verification not implemented)	1554
3.306.6 Sympy [B] (verification not implemented)	1554
3.306.7 Maxima [A] (verification not implemented)	1555
3.306.8 Giac [B] (verification not implemented)	1555
3.306.9 Mupad [B] (verification not implemented)	1556

3.306.1 Optimal result

Integrand size = 9, antiderivative size = 24

$$\int x\sqrt[3]{c+x} dx = -\frac{3}{4}c(c+x)^{4/3} + \frac{3}{7}(c+x)^{7/3}$$

output `-3/4*c*(c+x)^(4/3)+3/7*(c+x)^(7/3)`

3.306.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.75

$$\int x\sqrt[3]{c+x} dx = \frac{3}{28}(c+x)^{4/3}(-3c+4x)$$

input `Integrate[x*(c + x)^(1/3),x]`

output `(3*(c + x)^(4/3)*(-3*c + 4*x))/28`

3.306.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x\sqrt[3]{c+x} dx$$

$$\downarrow \text{53}$$

$$\int \left((c+x)^{4/3} - c\sqrt[3]{c+x} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{3}{7}(c+x)^{7/3} - \frac{3}{4}c(c+x)^{4/3}$$

input `Int[x*(c + x)^(1/3), x]`

output `(-3*c*(c + x)^(4/3))/4 + (3*(c + x)^(7/3))/7`

3.306.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.306.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.62

method	result	size
gospers	$-\frac{3(c+x)^{\frac{4}{3}}(3c-4x)}{28}$	15
derivativedivides	$-\frac{3c(c+x)^{\frac{4}{3}}}{4} + \frac{3(c+x)^{\frac{7}{3}}}{7}$	17
default	$-\frac{3c(c+x)^{\frac{4}{3}}}{4} + \frac{3(c+x)^{\frac{7}{3}}}{7}$	17
trager	$\left(-\frac{9}{28}c^2 + \frac{3}{28}cx + \frac{3}{7}x^2\right)(c+x)^{\frac{1}{3}}$	22
risch	$-\frac{3(c+x)^{\frac{1}{3}}(3c^2-cx-4x^2)}{28}$	23

input `int(x*(c+x)^(1/3),x,method=_RETURNVERBOSE)`output `-3/28*(c+x)^(4/3)*(3*c-4*x)`**3.306.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int x\sqrt[3]{c+x} dx = -\frac{3}{28}(3c^2 - cx - 4x^2)(c+x)^{\frac{1}{3}}$$

input `integrate(x*(c+x)^(1/3),x, algorithm="fricas")`output `-3/28*(3*c^2 - c*x - 4*x^2)*(c + x)^(1/3)`**3.306.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 144 vs. 2(20) = 40.

Time = 0.63 (sec) , antiderivative size = 144, normalized size of antiderivative = 6.00

$$\int x\sqrt[3]{c+x} dx = -\frac{9c^{\frac{13}{3}}\sqrt[3]{1+\frac{x}{c}}}{28c^2+28cx} + \frac{9c^{\frac{13}{3}}}{28c^2+28cx} - \frac{6c^{\frac{10}{3}}x\sqrt[3]{1+\frac{x}{c}}}{28c^2+28cx} \\ + \frac{9c^{\frac{10}{3}}x}{28c^2+28cx} + \frac{15c^{\frac{7}{3}}x^2\sqrt[3]{1+\frac{x}{c}}}{28c^2+28cx} + \frac{12c^{\frac{4}{3}}x^3\sqrt[3]{1+\frac{x}{c}}}{28c^2+28cx}$$

input `integrate(x*(c+x)**(1/3),x)`

output `-9*c**(13/3)*(1 + x/c)**(1/3)/(28*c**2 + 28*c*x) + 9*c**(13/3)/(28*c**2 + 28*c*x) - 6*c**(10/3)*x*(1 + x/c)**(1/3)/(28*c**2 + 28*c*x) + 9*c**(10/3)*x/(28*c**2 + 28*c*x) + 15*c**(7/3)*x**2*(1 + x/c)**(1/3)/(28*c**2 + 28*c*x) + 12*c**(4/3)*x**3*(1 + x/c)**(1/3)/(28*c**2 + 28*c*x)`

3.306.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int x\sqrt[3]{c+x} dx = \frac{3}{7}(c+x)^{\frac{7}{3}} - \frac{3}{4}(c+x)^{\frac{4}{3}}c$$

input `integrate(x*(c+x)^(1/3),x, algorithm="maxima")`

output `3/7*(c + x)^(7/3) - 3/4*(c + x)^(4/3)*c`

3.306.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. $2(16) = 32$.

Time = 0.32 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.79

$$\int x\sqrt[3]{c+x} dx = \frac{3}{7}(c+x)^{\frac{7}{3}} - \frac{3}{2}(c+x)^{\frac{4}{3}}c + 3(c+x)^{\frac{1}{3}}c^2 + \frac{3}{4}\left((c+x)^{\frac{4}{3}} - 4(c+x)^{\frac{1}{3}}c\right)c$$

input `integrate(x*(c+x)^(1/3),x, algorithm="giac")`

output `3/7*(c + x)^(7/3) - 3/2*(c + x)^(4/3)*c + 3*(c + x)^(1/3)*c^2 + 3/4*((c + x)^(4/3) - 4*(c + x)^(1/3)*c)*c`

3.306.9 Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.58

$$\int x \sqrt[3]{c+x} dx = -\frac{3(c+x)^{4/3}(3c-4x)}{28}$$

input `int(x*(c + x)^(1/3),x)`

output `-(3*(c + x)^(4/3)*(3*c - 4*x))/28`

3.307 $\int e^{\sqrt[3]{x}} dx$

3.307.1 Optimal result	1557
3.307.2 Mathematica [A] (verified)	1557
3.307.3 Rubi [A] (verified)	1558
3.307.4 Maple [A] (verified)	1559
3.307.5 Fricas [A] (verification not implemented)	1559
3.307.6 Sympy [A] (verification not implemented)	1560
3.307.7 Maxima [A] (verification not implemented)	1560
3.307.8 Giac [A] (verification not implemented)	1560
3.307.9 Mupad [B] (verification not implemented)	1561

3.307.1 Optimal result

Integrand size = 7, antiderivative size = 38

$$\int e^{\sqrt[3]{x}} dx = 6e^{\sqrt[3]{x}} - 6e^{\sqrt[3]{x}}\sqrt[3]{x} + 3e^{\sqrt[3]{x}}x^{2/3}$$

output `6*exp(x^(1/3))-6*exp(x^(1/3))*x^(1/3)+3*exp(x^(1/3))*x^(2/3)`

3.307.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.63

$$\int e^{\sqrt[3]{x}} dx = e^{\sqrt[3]{x}}(6 - 6\sqrt[3]{x} + 3x^{2/3})$$

input `Integrate[E^x^(1/3),x]`

output `E^x^(1/3)*(6 - 6*x^(1/3) + 3*x^(2/3))`

3.307.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.08, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {2636, 2607, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{\sqrt[3]{x}} dx \\
 & \quad \downarrow \text{2636} \\
 & 3 \int e^{\sqrt[3]{x}} x^{2/3} d\sqrt[3]{x} \\
 & \quad \downarrow \text{2607} \\
 & 3 \left(e^{\sqrt[3]{x}} x^{2/3} - 2 \int e^{\sqrt[3]{x}} \sqrt[3]{x} d\sqrt[3]{x} \right) \\
 & \quad \downarrow \text{2607} \\
 & 3 \left(e^{\sqrt[3]{x}} x^{2/3} - 2 \left(e^{\sqrt[3]{x}} \sqrt[3]{x} - \int e^{\sqrt[3]{x}} d\sqrt[3]{x} \right) \right) \\
 & \quad \downarrow \text{2624} \\
 & 3 \left(e^{\sqrt[3]{x}} x^{2/3} - 2 \left(e^{\sqrt[3]{x}} \sqrt[3]{x} - e^{\sqrt[3]{x}} \right) \right)
 \end{aligned}$$

input `Int[E^x^(1/3),x]`

output `3*(-2*(-E^x^(1/3) + E^x^(1/3)*x^(1/3)) + E^x^(1/3)*x^(2/3))`

3.307.3.1 Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

```
rule 2624 Int[((F_)^(v_))^(n_), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;
FreeQ[{F, n}, x] && LinearQ[v, x]
```

```
rule 2636 Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_))), x_Symbol] := With[{k =
Denominator[n]}, Simp[k/d Subst[Int[x^(k - 1)*F^(a + b*x^(k*n)), x], x, (
c + d*x)^(1/k)], x]] /; FreeQ[{F, a, b, c, d}, x] && IntegerQ[2/n] && !Int
egerQ[n]
```

3.307.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.53

method	result	size
meijerg	$-6 + (3x^{\frac{2}{3}} - 6x^{\frac{1}{3}} + 6)e^{x^{\frac{1}{3}}}$	20
derivativedivides	$6e^{x^{\frac{1}{3}}} - 6e^{x^{\frac{1}{3}}}x^{\frac{1}{3}} + 3e^{x^{\frac{1}{3}}}x^{\frac{2}{3}}$	26
default	$6e^{x^{\frac{1}{3}}} - 6e^{x^{\frac{1}{3}}}x^{\frac{1}{3}} + 3e^{x^{\frac{1}{3}}}x^{\frac{2}{3}}$	26

```
input int(exp(x^(1/3)),x,method=_RETURNVERBOSE)
```

```
output -6+(3*x^(2/3)-6*x^(1/3)+6)*exp(x^(1/3))
```

3.307.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.42

$$\int e^{\sqrt[3]{x}} dx = 3 \left(x^{\frac{2}{3}} - 2x^{\frac{1}{3}} + 2 \right) e^{(x^{\frac{1}{3}})}$$

```
input integrate(exp(x^(1/3)),x, algorithm="fricas")
```

```
output 3*(x^(2/3) - 2*x^(1/3) + 2)*e^(x^(1/3))
```


3.307.6 Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.89

$$\int e^{\sqrt[3]{x}} dx = 3x^{\frac{2}{3}}e^{\sqrt[3]{x}} - 6\sqrt[3]{x}e^{\sqrt[3]{x}} + 6e^{\sqrt[3]{x}}$$

input `integrate(exp(x**(1/3)),x)`output `3*x**(2/3)*exp(x**(1/3)) - 6*x**(1/3)*exp(x**(1/3)) + 6*exp(x**(1/3))`**3.307.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.42

$$\int e^{\sqrt[3]{x}} dx = 3 \left(x^{\frac{2}{3}} - 2x^{\frac{1}{3}} + 2 \right) e^{(x^{\frac{1}{3}})}$$

input `integrate(exp(x^(1/3)),x, algorithm="maxima")`output `3*(x^(2/3) - 2*x^(1/3) + 2)*e^(x^(1/3))`**3.307.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.42

$$\int e^{\sqrt[3]{x}} dx = 3 \left(x^{\frac{2}{3}} - 2x^{\frac{1}{3}} + 2 \right) e^{(x^{\frac{1}{3}})}$$

input `integrate(exp(x^(1/3)),x, algorithm="giac")`output `3*(x^(2/3) - 2*x^(1/3) + 2)*e^(x^(1/3))`

3.307.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.55

$$\int e^{\sqrt[3]{x}} dx = 3 x e^{x^{1/3}} \left(\frac{2}{x} + \frac{1}{x^{1/3}} - \frac{2}{x^{2/3}} \right)$$

input `int(exp(x^(1/3)),x)`

output `3*x*exp(x^(1/3))*(2/x + 1/x^(1/3) - 2/x^(2/3))`

3.308 $\int \frac{1}{4+x+\sqrt{1+x}} dx$

3.308.1 Optimal result	1562
3.308.2 Mathematica [A] (verified)	1562
3.308.3 Rubi [A] (verified)	1563
3.308.4 Maple [A] (verified)	1564
3.308.5 Fricas [A] (verification not implemented)	1565
3.308.6 Sympy [A] (verification not implemented)	1565
3.308.7 Maxima [A] (verification not implemented)	1565
3.308.8 Giac [A] (verification not implemented)	1566
3.308.9 Mupad [B] (verification not implemented)	1566

3.308.1 Optimal result

Integrand size = 12, antiderivative size = 37

$$\int \frac{1}{4+x+\sqrt{1+x}} dx = -\frac{2 \arctan\left(\frac{1+2\sqrt{1+x}}{\sqrt{11}}\right)}{\sqrt{11}} + \log(4+x+\sqrt{1+x})$$

output `ln(4+x+(1+x)^(1/2))-2/11*arctan(1/11*(1+2*(1+x)^(1/2))*11^(1/2))*11^(1/2)`

3.308.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

$$\int \frac{1}{4+x+\sqrt{1+x}} dx = -\frac{2 \arctan\left(\frac{1+2\sqrt{1+x}}{\sqrt{11}}\right)}{\sqrt{11}} + \log(4+x+\sqrt{1+x})$$

input `Integrate[(4 + x + Sqrt[1 + x])^(-1), x]`

output `(-2*ArcTan[(1 + 2*Sqrt[1 + x])/Sqrt[11]])/Sqrt[11] + Log[4 + x + Sqrt[1 + x]]`

3.308.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.16, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {7267, 1142, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x + \sqrt{x+1} + 4} dx \\
 & \quad \downarrow \text{7267} \\
 & 2 \int \frac{\sqrt{x+1}}{x + \sqrt{x+1} + 4} d\sqrt{x+1} \\
 & \quad \downarrow \text{1142} \\
 & 2 \left(\frac{1}{2} \int \frac{2\sqrt{x+1} + 1}{x + \sqrt{x+1} + 4} d\sqrt{x+1} - \frac{1}{2} \int \frac{1}{x + \sqrt{x+1} + 4} d\sqrt{x+1} \right) \\
 & \quad \downarrow \text{1083} \\
 & 2 \left(\int \frac{1}{-x - 12} d(2\sqrt{x+1} + 1) + \frac{1}{2} \int \frac{2\sqrt{x+1} + 1}{x + \sqrt{x+1} + 4} d\sqrt{x+1} \right) \\
 & \quad \downarrow \text{217} \\
 & 2 \left(\frac{1}{2} \int \frac{2\sqrt{x+1} + 1}{x + \sqrt{x+1} + 4} d\sqrt{x+1} - \frac{\arctan\left(\frac{2\sqrt{x+1} + 1}{\sqrt{11}}\right)}{\sqrt{11}} \right) \\
 & \quad \downarrow \text{1103} \\
 & 2 \left(\frac{1}{2} \log(x + \sqrt{x+1} + 4) - \frac{\arctan\left(\frac{2\sqrt{x+1} + 1}{\sqrt{11}}\right)}{\sqrt{11}} \right)
 \end{aligned}$$

input `Int[(4 + x + Sqrt[1 + x])^(-1),x]`

output `2*(-(ArcTan[(1 + 2*Sqrt[1 + x])/Sqrt[11]]/Sqrt[11]) + Log[4 + x + Sqrt[1 + x]]/2)`

3.308.3.1 Defintions of rubi rules used

- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & (LtQ[a, 0] || LtQ[b, 0])`
- rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1142 `Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`
- rule 7267 `Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Simp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])], x] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]]`

3.308.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.84

method	result
derivativedivides	$\ln(4 + x + \sqrt{1 + x}) - \frac{2 \arctan\left(\frac{(1+2\sqrt{1+x})\sqrt{11}}{11}\right)\sqrt{11}}{11}$
default	$-\frac{\ln(4+x-\sqrt{1+x})}{2} - \frac{\sqrt{11} \arctan\left(\frac{(2\sqrt{1+x}-1)\sqrt{11}}{11}\right)}{11} + \frac{\ln(4+x+\sqrt{1+x})}{2} - \frac{\arctan\left(\frac{(1+2\sqrt{1+x})\sqrt{11}}{11}\right)\sqrt{11}}{11} + \frac{\sqrt{11}}{11}$
trager	$-\ln(4 + x + \sqrt{1 + x}) \text{RootOf}(11_Z^2 - 22_Z + 12) + \ln(-847 \text{RootOf}(11_Z^2 - 22_Z + 12) + 11)$

```
input int(1/(4+x+(1+x)^(1/2)),x,method=_RETURNVERBOSE)
```

3.308. $\int \frac{1}{4+x+\sqrt{1+x}} dx$

output $\ln(4+x+(1+x)^{(1/2)})-2/11*\arctan(1/11*(1+2*(1+x)^{(1/2}))*11^{(1/2)})*11^{(1/2)}$

3.308.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.86

$$\int \frac{1}{4+x+\sqrt{1+x}} dx = -\frac{2}{11} \sqrt{11} \arctan \left(\frac{2}{11} \sqrt{11} \sqrt{x+1} + \frac{1}{11} \sqrt{11} \right) + \log(x + \sqrt{x+1} + 4)$$

input `integrate(1/(4+x+(1+x)^(1/2)),x, algorithm="fricas")`

output $-2/11*\sqrt{11}*\arctan(2/11*\sqrt{11}*\sqrt{x+1} + 1/11*\sqrt{11}) + \log(x + \sqrt{x+1} + 4)$

3.308.6 Sympy [A] (verification not implemented)

Time = 0.60 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.05

$$\int \frac{1}{4+x+\sqrt{1+x}} dx = \log(x + \sqrt{x+1} + 4) - \frac{2\sqrt{11} \operatorname{atan} \left(\frac{2\sqrt{11}(\sqrt{x+1} + \frac{1}{2})}{11} \right)}{11}$$

input `integrate(1/(4+x+(1+x)**(1/2)),x)`

output $\log(x + \sqrt{x+1} + 4) - 2*\sqrt{11}*\operatorname{atan}(2*\sqrt{11}*(\sqrt{x+1} + 1/2)/11)/11$

3.308.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.81

$$\int \frac{1}{4+x+\sqrt{1+x}} dx = -\frac{2}{11} \sqrt{11} \arctan \left(\frac{1}{11} \sqrt{11} (2\sqrt{x+1} + 1) \right) + \log(x + \sqrt{x+1} + 4)$$

input `integrate(1/(4+x+(1+x)^(1/2)),x, algorithm="maxima")`

output $-2/11*\sqrt{11}*\arctan(1/11*\sqrt{11}*(2*\sqrt{x+1} + 1)) + \log(x + \sqrt{x+1} + 4)$

3.308.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.81

$$\int \frac{1}{4+x+\sqrt{1+x}} dx = -\frac{2}{11} \sqrt{11} \arctan\left(\frac{1}{11} \sqrt{11} (2\sqrt{x+1} + 1)\right) + \log(x + \sqrt{x+1} + 4)$$

input `integrate(1/(4+x+(1+x)^(1/2)),x, algorithm="giac")`output `-2/11*sqrt(11)*arctan(1/11*sqrt(11)*(2*sqrt(x + 1) + 1)) + log(x + sqrt(x + 1) + 4)`**3.308.9 Mupad [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.86

$$\int \frac{1}{4+x+\sqrt{1+x}} dx = \ln(x + \sqrt{x+1} + 4) - \frac{2\sqrt{11} \operatorname{atan}\left(\frac{\sqrt{11}}{11} + \frac{2\sqrt{11}\sqrt{x+1}}{11}\right)}{11}$$

input `int(1/(x + (x + 1)^(1/2) + 4),x)`output `log(x + (x + 1)^(1/2) + 4) - (2*11^(1/2)*atan(11^(1/2)/11 + (2*11^(1/2)*(x + 1)^(1/2))/11))/11`

3.309 $\int \frac{1+x^3}{-x^2+x^3} dx$

3.309.1 Optimal result	1567
3.309.2 Mathematica [A] (verified)	1567
3.309.3 Rubi [A] (verified)	1568
3.309.4 Maple [A] (verified)	1569
3.309.5 Fricas [A] (verification not implemented)	1569
3.309.6 Sympy [A] (verification not implemented)	1569
3.309.7 Maxima [A] (verification not implemented)	1570
3.309.8 Giac [A] (verification not implemented)	1570
3.309.9 Mupad [B] (verification not implemented)	1570

3.309.1 Optimal result

Integrand size = 17, antiderivative size = 17

$$\int \frac{1+x^3}{-x^2+x^3} dx = \frac{1}{x} + x + 2\log(1-x) - \log(x)$$

output `1/x+x+2*ln(1-x)-ln(x)`

3.309.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{1+x^3}{-x^2+x^3} dx = \frac{1}{x} + x + 2\log(1-x) - \log(x)$$

input `Integrate[(1 + x^3)/(-x^2 + x^3), x]`

output `x^(-1) + x + 2*Log[1 - x] - Log[x]`

3.309.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2026, 2123, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^3 + 1}{x^3 - x^2} dx \\ & \quad \downarrow \text{2026} \\ & \int \frac{x^3 + 1}{(x - 1)x^2} dx \\ & \quad \downarrow \text{2123} \\ & \int \left(-\frac{1}{x^2} - \frac{1}{x} + \frac{2}{x - 1} + 1 \right) dx \\ & \quad \downarrow \text{2009} \\ & x + \frac{1}{x} + 2 \log(1 - x) - \log(x) \end{aligned}$$

input `Int[(1 + x^3)/(-x^2 + x^3),x]`

output `x^(-1) + x + 2*Log[1 - x] - Log[x]`

3.309.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p * r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && Integ erQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

rule 2123 `Int[(Px_)*((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && PolyQ[Px, x] && (IntegersQ[m, n] || IGtQ[m, -2])`

3.309.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

method	result	size
default	$x + 2 \ln(-1 + x) + \frac{1}{x} - \ln(x)$	16
risch	$x + 2 \ln(-1 + x) + \frac{1}{x} - \ln(x)$	16
norman	$\frac{x^2+1}{x} - \ln(x) + 2 \ln(-1 + x)$	21
meijerg	$\frac{1}{x} - \ln(x) - i\pi + 2 \ln(1 - x) + x$	22
parallelrisch	$-\frac{x \ln(x) - 2 \ln(-1+x)x - x^2 - 1}{x}$	24

input `int((x^3+1)/(x^3-x^2),x,method=_RETURNVERBOSE)`output `x+2*ln(-1+x)+1/x-ln(x)`**3.309.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.24

$$\int \frac{1+x^3}{-x^2+x^3} dx = \frac{x^2 + 2x \log(x-1) - x \log(x) + 1}{x}$$

input `integrate((x^3+1)/(x^3-x^2),x, algorithm="fricas")`output `(x^2 + 2*x*log(x - 1) - x*log(x) + 1)/x`**3.309.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \frac{1+x^3}{-x^2+x^3} dx = x - \log(x) + 2 \log(x-1) + \frac{1}{x}$$

input `integrate((x**3+1)/(x**3-x**2),x)`output `x - log(x) + 2*log(x - 1) + 1/x`

3.309.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{1+x^3}{-x^2+x^3} dx = x + \frac{1}{x} + 2 \log(x-1) - \log(x)$$

input `integrate((x^3+1)/(x^3-x^2),x, algorithm="maxima")`output `x + 1/x + 2*log(x - 1) - log(x)`**3.309.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{1+x^3}{-x^2+x^3} dx = x + \frac{1}{x} + 2 \log(|x-1|) - \log(|x|)$$

input `integrate((x^3+1)/(x^3-x^2),x, algorithm="giac")`output `x + 1/x + 2*log(abs(x - 1)) - log(abs(x))`**3.309.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{1+x^3}{-x^2+x^3} dx = x + 2 \ln(x-1) - \ln(x) + \frac{1}{x}$$

input `int(-(x^3 + 1)/(x^2 - x^3),x)`output `x + 2*log(x - 1) - log(x) + 1/x`

3.310 $\int (-3 + 4x + x^2) \sin(2x) dx$

3.310.1 Optimal result1571
3.310.2 Mathematica [A] (verified)1571
3.310.3 Rubi [A] (verified)1572
3.310.4 Maple [A] (verified)1573
3.310.5 Fricas [A] (verification not implemented)1573
3.310.6 Sympy [A] (verification not implemented)1574
3.310.7 Maxima [A] (verification not implemented)1574
3.310.8 Giac [A] (verification not implemented)1574
3.310.9 Mupad [B] (verification not implemented)1575

3.310.1 Optimal result

Integrand size = 13, antiderivative size = 40

$$\int (-3 + 4x + x^2) \sin(2x) dx = \frac{7}{4} \cos(2x) - 2x \cos(2x) - \frac{1}{2} x^2 \cos(2x) + \sin(2x) + \frac{1}{2} x \sin(2x)$$

output `7/4*cos(2*x)-2*x*cos(2*x)-1/2*x^2*cos(2*x)+sin(2*x)+1/2*x*sin(2*x)`

3.310.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.72

$$\int (-3 + 4x + x^2) \sin(2x) dx = \frac{1}{4} ((7 - 8x - 2x^2) \cos(2x) + 2(2 + x) \sin(2x))$$

input `Integrate[(-3 + 4*x + x^2)*Sin[2*x],x]`

output `((7 - 8*x - 2*x^2)*Cos[2*x] + 2*(2 + x)*Sin[2*x])/4`

3.310.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x^2 + 4x - 3) \sin(2x) dx$$

$$\downarrow \text{7293}$$

$$\int (x^2 \sin(2x) + 4x \sin(2x) - 3 \sin(2x)) dx$$

$$\downarrow \text{2009}$$

$$-\frac{1}{2}x^2 \cos(2x) + \frac{1}{2}x \sin(2x) + \sin(2x) - 2x \cos(2x) + \frac{7}{4} \cos(2x)$$

input `Int[(-3 + 4*x + x^2)*Sin[2*x],x]`

output `(7*Cos[2*x])/4 - 2*x*Cos[2*x] - (x^2*Cos[2*x])/2 + Sin[2*x] + (x*Sin[2*x])/2`

3.310.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] :> With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

3.310.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.65

method	result	size
risch	$\left(-\frac{1}{2}x^2 - 2x + \frac{7}{4}\right) \cos(2x) + \frac{(2+x)\sin(2x)}{2}$	26
derivativedivides	$\frac{7\cos(2x)}{4} - 2x\cos(2x) - \frac{x^2\cos(2x)}{2} + \sin(2x) + \frac{x\sin(2x)}{2}$	35
default	$\frac{7\cos(2x)}{4} - 2x\cos(2x) - \frac{x^2\cos(2x)}{2} + \sin(2x) + \frac{x\sin(2x)}{2}$	35
parts	$\frac{7\cos(2x)}{4} - 2x\cos(2x) - \frac{x^2\cos(2x)}{2} + \sin(2x) + \frac{x\sin(2x)}{2}$	35
parallelrisch	$\frac{(x^2+4x)(\tan^2(x))+(2x+4)\tan(x)-x^2-4x+7}{2+2(\tan^2(x))}$	42
norman	$\frac{x\tan(x)-2x-\frac{x^2}{2}+2x(\tan^2(x))+\frac{x^2(\tan^2(x))}{2}+2\tan(x)+\frac{7}{2}}{1+\tan^2(x)}$	44
meijerg	$\frac{\sqrt{\pi}\left(-\frac{1}{2\sqrt{\pi}}+\frac{(-2x^2+1)\cos(2x)}{2\sqrt{\pi}}+\frac{x\sin(2x)}{\sqrt{\pi}}\right)}{2} + 2\sqrt{\pi}\left(-\frac{x\cos(2x)}{\sqrt{\pi}}+\frac{\sin(2x)}{2\sqrt{\pi}}\right) - \frac{3\sqrt{\pi}\left(\frac{1}{\sqrt{\pi}}-\frac{\cos(2x)}{\sqrt{\pi}}\right)}{2}$	81

input `int((x^2+4*x-3)*sin(2*x),x,method=_RETURNVERBOSE)`output `(-1/2*x^2-2*x+7/4)*cos(2*x)+1/2*(2+x)*sin(2*x)`**3.310.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.65

$$\int (-3 + 4x + x^2) \sin(2x) dx = -\frac{1}{4} (2x^2 + 8x - 7) \cos(2x) + \frac{1}{2} (x + 2) \sin(2x)$$

input `integrate((x^2+4*x-3)*sin(2*x),x, algorithm="fricas")`output `-1/4*(2*x^2 + 8*x - 7)*cos(2*x) + 1/2*(x + 2)*sin(2*x)`

3.310.6 Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.98

$$\int (-3 + 4x + x^2) \sin(2x) dx = -\frac{x^2 \cos(2x)}{2} + \frac{x \sin(2x)}{2} - 2x \cos(2x) + \sin(2x) + \frac{7 \cos(2x)}{4}$$

input `integrate((x**2+4*x-3)*sin(2*x),x)`output `-x**2*cos(2*x)/2 + x*sin(2*x)/2 - 2*x*cos(2*x) + sin(2*x) + 7*cos(2*x)/4`**3.310.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.95

$$\int (-3 + 4x + x^2) \sin(2x) dx = -\frac{1}{4} (2x^2 - 1) \cos(2x) - 2x \cos(2x) + \frac{1}{2} x \sin(2x) + \frac{3}{2} \cos(2x) + \sin(2x)$$

input `integrate((x^2+4*x-3)*sin(2*x),x, algorithm="maxima")`output `-1/4*(2*x^2 - 1)*cos(2*x) - 2*x*cos(2*x) + 1/2*x*sin(2*x) + 3/2*cos(2*x) + sin(2*x)`**3.310.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.65

$$\int (-3 + 4x + x^2) \sin(2x) dx = -\frac{1}{4} (2x^2 + 8x - 7) \cos(2x) + \frac{1}{2} (x + 2) \sin(2x)$$

input `integrate((x^2+4*x-3)*sin(2*x),x, algorithm="giac")`output `-1/4*(2*x^2 + 8*x - 7)*cos(2*x) + 1/2*(x + 2)*sin(2*x)`

3.310.9 Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.85

$$\int (-3+4x+x^2) \sin(2x) dx = \frac{7 \cos(2x)}{4} + \sin(2x) - 2x \cos(2x) + \frac{x \sin(2x)}{2} - \frac{x^2 \cos(2x)}{2}$$

input `int(sin(2*x)*(4*x + x^2 - 3),x)`

output `(7*cos(2*x))/4 + sin(2*x) - 2*x*cos(2*x) + (x*sin(2*x))/2 - (x^2*cos(2*x))
/2`

3.311 $\int \cos(\cos(x)) \sin(x) dx$

3.311.1 Optimal result	1576
3.311.2 Mathematica [A] (verified)	1576
3.311.3 Rubi [A] (verified)	1577
3.311.4 Maple [A] (verified)	1578
3.311.5 Fricas [B] (verification not implemented)	1578
3.311.6 Sympy [A] (verification not implemented)	1579
3.311.7 Maxima [A] (verification not implemented)	1579
3.311.8 Giac [A] (verification not implemented)	1579
3.311.9 Mupad [B] (verification not implemented)	1580

3.311.1 Optimal result

Integrand size = 6, antiderivative size = 5

$$\int \cos(\cos(x)) \sin(x) dx = -\sin(\cos(x))$$

output `-sin(cos(x))`

3.311.2 Mathematica [A] (verified)

Time = 4.17 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \cos(\cos(x)) \sin(x) dx = -\sin(\cos(x))$$

input `Integrate[Cos[Cos[x]]*Sin[x],x]`

output `-Sin[Cos[x]]`

3.311.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {4835, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(x) \cos(\cos(x)) dx \\
 & \quad \downarrow 4835 \\
 & - \int \cos(\cos(x)) d \cos(x) \\
 & \quad \downarrow 3042 \\
 & - \int \sin\left(\cos(x) + \frac{\pi}{2}\right) d \cos(x) \\
 & \quad \downarrow 3117 \\
 & - \sin(\cos(x))
 \end{aligned}$$

input `Int[Cos[Cos[x]]*Sin[x],x]`

output `-Sin[Cos[x]]`

3.311.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 4835 `Int[(u_)*(F_)[(c_.)*((a_.) + (b_.)*(x_))], x_Symbol] := With[{d = FreeFactors[Cos[c*(a + b*x)], x]}, Simp[-d/(b*c) Subst[Int[SubstFor[1, Cos[c*(a + b*x)]]/d, u, x], x], x, Cos[c*(a + b*x)]/d, x] /; FunctionOfQ[Cos[c*(a + b*x)]/d, u, x, True]] /; FreeQ[{a, b, c}, x] && (EqQ[F, Sin] || EqQ[F, sin])`

3.311.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
derivativedivides	$-\sin(\cos(x))$	6
default	$-\sin(\cos(x))$	6
risch	$-\sin(\cos(x))$	6
parallelrisch	$-\sin(\cos(x))$	6
norman	$\frac{-2(\tan^2(\frac{x}{2})) \tan\left(\frac{1 - (\tan^2(\frac{x}{2}))}{2 + 2(\tan^2(\frac{x}{2}))}\right) - 2 \tan\left(\frac{1 - (\tan^2(\frac{x}{2}))}{2 + 2(\tan^2(\frac{x}{2}))}\right)}{\left(1 + \tan^2\left(\frac{1 - (\tan^2(\frac{x}{2}))}{2(1 + \tan^2(\frac{x}{2}))}\right)\right) (1 + \tan^2(\frac{x}{2}))}$	98

input `int(cos(cos(x))*sin(x),x,method=_RETURNVERBOSE)`output `-sin(cos(x))`**3.311.5 Fracas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 20 vs. $2(5) = 10$.

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 4.00

$$\int \cos(\cos(x)) \sin(x) dx = \sin\left(\frac{\tan\left(\frac{1}{2}x\right)^2 - 1}{\tan\left(\frac{1}{2}x\right)^2 + 1}\right)$$

input `integrate(cos(cos(x))*sin(x),x, algorithm="fracas")`output `sin((tan(1/2*x)^2 - 1)/(tan(1/2*x)^2 + 1))`

3.311.6 Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \cos(\cos(x)) \sin(x) dx = -\sin(\cos(x))$$

input `integrate(cos(cos(x))*sin(x),x)`

output `-sin(cos(x))`

3.311.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \cos(\cos(x)) \sin(x) dx = -\sin(\cos(x))$$

input `integrate(cos(cos(x))*sin(x),x, algorithm="maxima")`

output `-sin(cos(x))`

3.311.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \cos(\cos(x)) \sin(x) dx = -\sin(\cos(x))$$

input `integrate(cos(cos(x))*sin(x),x, algorithm="giac")`

output `-sin(cos(x))`

3.311.9 Mupad [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \cos(\cos(x)) \sin(x) dx = -\sin(\cos(x))$$

input `int(cos(cos(x))*sin(x),x)`

output `-sin(cos(x))`

3.312 $\int \frac{1}{\sqrt{16-x^2}} dx$

3.312.1 Optimal result	1581
3.312.2 Mathematica [B] (verified)	1581
3.312.3 Rubi [A] (verified)	1582
3.312.4 Maple [A] (verified)	1582
3.312.5 Fricas [B] (verification not implemented)	1583
3.312.6 Sympy [A] (verification not implemented)	1583
3.312.7 Maxima [A] (verification not implemented)	1583
3.312.8 Giac [B] (verification not implemented)	1584
3.312.9 Mupad [B] (verification not implemented)	1584

3.312.1 Optimal result

Integrand size = 11, antiderivative size = 6

$$\int \frac{1}{\sqrt{16-x^2}} dx = \arcsin\left(\frac{x}{4}\right)$$

output `arcsin(1/4*x)`

3.312.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 20 vs. $2(6) = 12$.

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 3.33

$$\int \frac{1}{\sqrt{16-x^2}} dx = -2 \arctan\left(\frac{\sqrt{16-x^2}}{4+x}\right)$$

input `Integrate[1/Sqrt[16 - x^2],x]`

output `-2*ArcTan[Sqrt[16 - x^2]/(4 + x)]`

3.312.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{16-x^2}} dx$$

↓ 223

$$\arcsin\left(\frac{x}{4}\right)$$

input `Int[1/Sqrt[16 - x^2],x]`

output `ArcSin[x/4]`

3.312.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

3.312.4 Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

method	result	size
default	$\arcsin\left(\frac{x}{4}\right)$	5
meijerg	$\arcsin\left(\frac{x}{4}\right)$	5
pseudoelliptic	$-\arctan\left(\frac{\sqrt{-x^2+16}}{x}\right)$	17
trager	$\text{RootOf}(-Z^2+1) \ln(\text{RootOf}(-Z^2+1)\sqrt{-x^2+16}+x)$	27

input `int(1/(-x^2+16)^(1/2),x,method=_RETURNVERBOSE)`

output `arcsin(1/4*x)`

3.312.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18 vs. $2(4) = 8$.

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 3.00

$$\int \frac{1}{\sqrt{16-x^2}} dx = -2 \arctan\left(\frac{\sqrt{-x^2+16}-4}{x}\right)$$

input `integrate(1/(-x^2+16)^(1/2),x, algorithm="fracas")`

output `-2*arctan((sqrt(-x^2 + 16) - 4)/x)`

3.312.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.50

$$\int \frac{1}{\sqrt{16-x^2}} dx = \operatorname{asin}\left(\frac{x}{4}\right)$$

input `integrate(1/(-x**2+16)**(1/2),x)`

output `asin(x/4)`

3.312.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt{16-x^2}} dx = \arcsin\left(\frac{1}{4}x\right)$$

input `integrate(1/(-x^2+16)^(1/2),x, algorithm="maxima")`

output `arcsin(1/4*x)`

3.312.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(4) = 8$.

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 3.17

$$\int \frac{1}{\sqrt{16-x^2}} dx = \frac{1}{2} \sqrt{-x^2+16}x + 8 \arcsin\left(\frac{1}{4}x\right)$$

input `integrate(1/(-x^2+16)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(-x^2 + 16)*x + 8*arcsin(1/4*x)`

3.312.9 Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt{16-x^2}} dx = \operatorname{asin}\left(\frac{x}{4}\right)$$

input `int(1/(16 - x^2)^(1/2),x)`

output `asin(x/4)`

3.313 $\int \frac{x^3}{(1+x)^{10}} dx$

3.313.1 Optimal result	1585
3.313.2 Mathematica [A] (verified)	1585
3.313.3 Rubi [A] (verified)	1586
3.313.4 Maple [A] (verified)	1587
3.313.5 Fricas [B] (verification not implemented)	1587
3.313.6 Sympy [A] (verification not implemented)	1588
3.313.7 Maxima [B] (verification not implemented)	1588
3.313.8 Giac [A] (verification not implemented)	1588
3.313.9 Mupad [B] (verification not implemented)	1589

3.313.1 Optimal result

Integrand size = 9, antiderivative size = 37

$$\int \frac{x^3}{(1+x)^{10}} dx = \frac{1}{9(1+x)^9} - \frac{3}{8(1+x)^8} + \frac{3}{7(1+x)^7} - \frac{1}{6(1+x)^6}$$

output `1/9/(1+x)^9-3/8/(1+x)^8+3/7/(1+x)^7-1/6/(1+x)^6`

3.313.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.65

$$\int \frac{x^3}{(1+x)^{10}} dx = -\frac{1+9x+36x^2+84x^3}{504(1+x)^9}$$

input `Integrate[x^3/(1+x)^10,x]`

output `-1/504*(1+9*x+36*x^2+84*x^3)/(1+x)^9`

3.313.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{(x+1)^{10}} dx$$

↓ 53

$$\int \left(\frac{1}{(x+1)^7} - \frac{3}{(x+1)^8} + \frac{3}{(x+1)^9} - \frac{1}{(x+1)^{10}} \right) dx$$

↓ 2009

$$-\frac{1}{6(x+1)^6} + \frac{3}{7(x+1)^7} - \frac{3}{8(x+1)^8} + \frac{1}{9(x+1)^9}$$

input `Int[x^3/(1 + x)^10,x]`

output `1/(9*(1 + x)^9) - 3/(8*(1 + x)^8) + 3/(7*(1 + x)^7) - 1/(6*(1 + x)^6)`

3.313.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.313.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.59

method	result	size
norman	$\frac{-\frac{1}{6}x^3 - \frac{1}{14}x^2 - \frac{1}{56}x - \frac{1}{504}}{(1+x)^9}$	22
risch	$\frac{-\frac{1}{6}x^3 - \frac{1}{14}x^2 - \frac{1}{56}x - \frac{1}{504}}{(1+x)^9}$	22
gospers	$-\frac{84x^3 + 36x^2 + 9x + 1}{504(1+x)^9}$	23
parallelrisch	$\frac{-84x^3 - 36x^2 - 9x - 1}{504(1+x)^9}$	23
default	$\frac{1}{9(1+x)^9} - \frac{3}{8(1+x)^8} + \frac{3}{7(1+x)^7} - \frac{1}{6(1+x)^6}$	30
meijerg	$\frac{x^4(x^5 + 9x^4 + 36x^3 + 84x^2 + 126x + 126)}{504(1+x)^9}$	34

input `int(x^3/(1+x)^10,x,method=_RETURNVERBOSE)`output `1/(1+x)^9*(-1/6*x^3-1/14*x^2-1/56*x-1/504)`**3.313.5 Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 62 vs. 2(29) = 58.

Time = 0.25 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.68

$$\int \frac{x^3}{(1+x)^{10}} dx$$

$$= -\frac{84x^3 + 36x^2 + 9x + 1}{504(x^9 + 9x^8 + 36x^7 + 84x^6 + 126x^5 + 126x^4 + 84x^3 + 36x^2 + 9x + 1)}$$

input `integrate(x^3/(1+x)^10,x, algorithm="fricas")`output `-1/504*(84*x^3 + 36*x^2 + 9*x + 1)/(x^9 + 9*x^8 + 36*x^7 + 84*x^6 + 126*x^5 + 126*x^4 + 84*x^3 + 36*x^2 + 9*x + 1)`

3.313.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.65

$$\int \frac{x^3}{(1+x)^{10}} dx$$

$$= \frac{-84x^3 - 36x^2 - 9x - 1}{504x^9 + 4536x^8 + 18144x^7 + 42336x^6 + 63504x^5 + 63504x^4 + 42336x^3 + 18144x^2 + 4536x + 504}$$

input `integrate(x**3/(1+x)**10,x)`

output `(-84*x**3 - 36*x**2 - 9*x - 1)/(504*x**9 + 4536*x**8 + 18144*x**7 + 42336*x**6 + 63504*x**5 + 63504*x**4 + 42336*x**3 + 18144*x**2 + 4536*x + 504)`

3.313.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 62 vs. 2(29) = 58.

Time = 0.19 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.68

$$\int \frac{x^3}{(1+x)^{10}} dx$$

$$= -\frac{84x^3 + 36x^2 + 9x + 1}{504(x^9 + 9x^8 + 36x^7 + 84x^6 + 126x^5 + 126x^4 + 84x^3 + 36x^2 + 9x + 1)}$$

input `integrate(x^3/(1+x)^10,x, algorithm="maxima")`

output `-1/504*(84*x^3 + 36*x^2 + 9*x + 1)/(x^9 + 9*x^8 + 36*x^7 + 84*x^6 + 126*x^5 + 126*x^4 + 84*x^3 + 36*x^2 + 9*x + 1)`

3.313.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.59

$$\int \frac{x^3}{(1+x)^{10}} dx = -\frac{84x^3 + 36x^2 + 9x + 1}{504(x+1)^9}$$

input `integrate(x^3/(1+x)^10,x, algorithm="giac")`

output `-1/504*(84*x^3 + 36*x^2 + 9*x + 1)/(x + 1)^9`

3.313.9 Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.78

$$\int \frac{x^3}{(1+x)^{10}} dx = \frac{3}{7(x+1)^7} - \frac{1}{6(x+1)^6} - \frac{3}{8(x+1)^8} + \frac{1}{9(x+1)^9}$$

input `int(x^3/(x + 1)^10,x)`

output `3/(7*(x + 1)^7) - 1/(6*(x + 1)^6) - 3/(8*(x + 1)^8) + 1/(9*(x + 1)^9)`

3.314 $\int \cot^3(2x) \csc^3(2x) dx$

3.314.1 Optimal result	1590
3.314.2 Mathematica [A] (verified)	1590
3.314.3 Rubi [A] (verified)	1591
3.314.4 Maple [A] (verified)	1592
3.314.5 Fricas [B] (verification not implemented)	1593
3.314.6 Sympy [A] (verification not implemented)	1593
3.314.7 Maxima [A] (verification not implemented)	1593
3.314.8 Giac [A] (verification not implemented)	1594
3.314.9 Mupad [B] (verification not implemented)	1594

3.314.1 Optimal result

Integrand size = 13, antiderivative size = 21

$$\int \cot^3(2x) \csc^3(2x) dx = \frac{1}{6} \csc^3(2x) - \frac{1}{10} \csc^5(2x)$$

output `1/6*csc(2*x)^3-1/10*csc(2*x)^5`

3.314.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \cot^3(2x) \csc^3(2x) dx = \frac{1}{6} \csc^3(2x) - \frac{1}{10} \csc^5(2x)$$

input `Integrate[Cot[2*x]^3*Csc[2*x]^3,x]`

output `Csc[2*x]^3/6 - Csc[2*x]^5/10`

3.314.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.19, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {3042, 25, 3086, 25, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^3(2x) \csc^3(2x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan\left(2x - \frac{\pi}{2}\right)^3 \left(-\sec\left(2x - \frac{\pi}{2}\right)\right)^3 dx \\
 & \quad \downarrow \text{25} \\
 & - \int \sec\left(2x - \frac{\pi}{2}\right)^3 \tan\left(2x - \frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{3086} \\
 & -\frac{1}{2} \int -\csc^2(2x) (1 - \csc^2(2x)) d \csc(2x) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{2} \int \csc^2(2x) (1 - \csc^2(2x)) d \csc(2x) \\
 & \quad \downarrow \text{244} \\
 & \frac{1}{2} \int (\csc^2(2x) - \csc^4(2x)) d \csc(2x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} \left(\frac{1}{3} \csc^3(2x) - \frac{1}{5} \csc^5(2x) \right)
 \end{aligned}$$

input `Int[Cot[2*x]^3*Csc[2*x]^3,x]`

output `(Csc[2*x]^3/3 - Csc[2*x]^5/5)/2`

3.314.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

3.314.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

method	result	size
derivativedivides	$\frac{\csc^3(2x)}{6} - \frac{\csc^5(2x)}{10}$	18
default	$\frac{\csc^3(2x)}{6} - \frac{\csc^5(2x)}{10}$	18
risch	$-\frac{4i(5e^{14ix} + 2e^{10ix} + 5e^{6ix})}{15(e^{4ix} - 1)^5}$	35

input `int(cot(2*x)^3*csc(2*x)^3,x,method=_RETURNVERBOSE)`

output `1/6*csc(2*x)^3-1/10*csc(2*x)^5`

3.314.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 36 vs. $2(17) = 34$.

Time = 0.25 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.71

$$\int \cot^3(2x) \csc^3(2x) dx = -\frac{5 \cos(2x)^2 - 2}{30 (\cos(2x)^4 - 2 \cos(2x)^2 + 1) \sin(2x)}$$

input `integrate(cot(2*x)^3*csc(2*x)^3,x, algorithm="fracas")`

output `-1/30*(5*cos(2*x)^2 - 2)/((cos(2*x)^4 - 2*cos(2*x)^2 + 1)*sin(2*x))`

3.314.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \cot^3(2x) \csc^3(2x) dx = -\frac{3 - 5 \sin^2(2x)}{30 \sin^5(2x)}$$

input `integrate(cot(2*x)**3*csc(2*x)**3,x)`

output `-(3 - 5*sin(2*x)**2)/(30*sin(2*x)**5)`

3.314.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

$$\int \cot^3(2x) \csc^3(2x) dx = \frac{5 \sin(2x)^2 - 3}{30 \sin(2x)^5}$$

input `integrate(cot(2*x)^3*csc(2*x)^3,x, algorithm="maxima")`

output `1/30*(5*sin(2*x)^2 - 3)/sin(2*x)^5`

3.314.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

$$\int \cot^3(2x) \csc^3(2x) dx = \frac{5 \sin(2x)^2 - 3}{30 \sin(2x)^5}$$

input `integrate(cot(2*x)^3*csc(2*x)^3,x, algorithm="giac")`output `1/30*(5*sin(2*x)^2 - 3)/sin(2*x)^5`**3.314.9 Mupad [B] (verification not implemented)**

Time = 0.39 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

$$\int \cot^3(2x) \csc^3(2x) dx = \frac{5 \sin(2x)^2 - 3}{30 \sin(2x)^5}$$

input `int(cot(2*x)^3/sin(2*x)^3,x)`output `(5*sin(2*x)^2 - 3)/(30*sin(2*x)^5)`

3.315 $\int (x + \sin(x))^2 dx$

3.315.1 Optimal result	1595
3.315.2 Mathematica [A] (verified)	1595
3.315.3 Rubi [A] (verified)	1596
3.315.4 Maple [A] (verified)	1597
3.315.5 Fracas [A] (verification not implemented)	1597
3.315.6 Sympy [A] (verification not implemented)	1597
3.315.7 Maxima [A] (verification not implemented)	1598
3.315.8 Giac [A] (verification not implemented)	1598
3.315.9 Mupad [B] (verification not implemented)	1598

3.315.1 Optimal result

Integrand size = 6, antiderivative size = 30

$$\int (x + \sin(x))^2 dx = \frac{x}{2} + \frac{x^3}{3} - 2x \cos(x) + 2 \sin(x) - \frac{1}{2} \cos(x) \sin(x)$$

output `1/2*x+1/3*x^3-2*x*cos(x)+2*sin(x)-1/2*cos(x)*sin(x)`

3.315.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int (x + \sin(x))^2 dx = \frac{1}{6}x(3 + 2x^2) - 2x \cos(x) + 2 \sin(x) - \frac{1}{4} \sin(2x)$$

input `Integrate[(x + Sin[x])^2,x]`

output `(x*(3 + 2*x^2))/6 - 2*x*Cos[x] + 2*Sin[x] - Sin[2*x]/4`

3.315.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x + \sin(x))^2 dx$$

$$\downarrow \text{7293}$$

$$\int (x^2 + \sin^2(x) + 2x \sin(x)) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^3}{3} + \frac{x}{2} + 2 \sin(x) - 2x \cos(x) - \frac{1}{2} \sin(x) \cos(x)$$

input `Int[(x + Sin[x])^2,x]`

output `x/2 + x^3/3 - 2*x*Cos[x] + 2*Sin[x] - (Cos[x]*Sin[x])/2`

3.315.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

3.315.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{x}{2} + \frac{x^3}{3} - 2x \cos(x) + 2 \sin(x) - \frac{\cos(x) \sin(x)}{2}$	25
risch	$\frac{x^3}{3} + \frac{x}{2} - 2x \cos(x) + 2 \sin(x) - \frac{\sin(2x)}{4}$	25
parallelrisc	$\frac{x^3}{3} + \frac{x}{2} - 2x \cos(x) + 2 \sin(x) - \frac{\sin(2x)}{4}$	25
parts	$\frac{x}{2} + \frac{x^3}{3} - 2x \cos(x) + 2 \sin(x) - \frac{\cos(x) \sin(x)}{2}$	25
norman	$\frac{x(\tan^2(\frac{x}{2}) - \frac{3x}{2} + \frac{x^3}{3} + 5(\tan^3(\frac{x}{2})) + \frac{5x(\tan^4(\frac{x}{2}))}{2} + \frac{2x^3(\tan^2(\frac{x}{2}))}{3} + \frac{x^3(\tan^4(\frac{x}{2}))}{3}) + 3 \tan(\frac{x}{2})}{(1 + \tan^2(\frac{x}{2}))^2}$	74

input `int((x+sin(x))^2,x,method=_RETURNVERBOSE)`output `1/2*x+1/3*x^3-2*x*cos(x)+2*sin(x)-1/2*cos(x)*sin(x)`**3.315.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int (x + \sin(x))^2 dx = \frac{1}{3} x^3 - 2x \cos(x) - \frac{1}{2} (\cos(x) - 4) \sin(x) + \frac{1}{2} x$$

input `integrate((x+sin(x))^2,x, algorithm="fricas")`output `1/3*x^3 - 2*x*cos(x) - 1/2*(cos(x) - 4)*sin(x) + 1/2*x`**3.315.6 Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.37

$$\int (x + \sin(x))^2 dx = \frac{x^3}{3} + \frac{x \sin^2(x)}{2} + \frac{x \cos^2(x)}{2} - 2x \cos(x) - \frac{\sin(x) \cos(x)}{2} + 2 \sin(x)$$

input `integrate((x+sin(x))**2,x)`

output `x**3/3 + x*sin(x)**2/2 + x*cos(x)**2/2 - 2*x*cos(x) - sin(x)*cos(x)/2 + 2*sin(x)`

3.315.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int (x + \sin(x))^2 dx = \frac{1}{3}x^3 - 2x \cos(x) + \frac{1}{2}x - \frac{1}{4} \sin(2x) + 2 \sin(x)$$

input `integrate((x+sin(x))^2,x, algorithm="maxima")`

output `1/3*x^3 - 2*x*cos(x) + 1/2*x - 1/4*sin(2*x) + 2*sin(x)`

3.315.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int (x + \sin(x))^2 dx = \frac{1}{3}x^3 - 2x \cos(x) + \frac{1}{2}x - \frac{1}{4} \sin(2x) + 2 \sin(x)$$

input `integrate((x+sin(x))^2,x, algorithm="giac")`

output `1/3*x^3 - 2*x*cos(x) + 1/2*x - 1/4*sin(2*x) + 2*sin(x)`

3.315.9 Mupad [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int (x + \sin(x))^2 dx = \frac{x}{2} + 2 \sin(x) - \frac{\cos(x) \sin(x)}{2} - 2x \cos(x) + \frac{x^3}{3}$$

input `int((x + sin(x))^2,x)`

output `x/2 + 2*sin(x) - (cos(x)*sin(x))/2 - 2*x*cos(x) + x^3/3`

3.316 $\int \frac{e^{\arctan(x)}}{1+x^2} dx$

3.316.1 Optimal result 1599
 3.316.2 Mathematica [C] (verified) 1599
 3.316.3 Rubi [A] (verified) 1600
 3.316.4 Maple [A] (verified) 1600
 3.316.5 Fricas [A] (verification not implemented) 1601
 3.316.6 Sympy [A] (verification not implemented) 1601
 3.316.7 Maxima [A] (verification not implemented) 1601
 3.316.8 Giac [A] (verification not implemented) 1602
 3.316.9 Mupad [B] (verification not implemented) 1602

3.316.1 Optimal result

Integrand size = 12, antiderivative size = 4

$$\int \frac{e^{\arctan(x)}}{1+x^2} dx = e^{\arctan(x)}$$

output `exp(arctan(x))`

3.316.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 6.75

$$\int \frac{e^{\arctan(x)}}{1+x^2} dx = (1-ix)^{\frac{i}{2}}(1+ix)^{-\frac{i}{2}}$$

input `Integrate[E^ArcTan[x]/(1+x^2),x]`

output `(1 - I*x)^(I/2)/(1 + I*x)^(I/2)`

3.316.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {5594}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{\arctan(x)}}{x^2 + 1} dx$$

↓ 5594

$$e^{\arctan(x)}$$

input `Int[E^ArcTan[x]/(1 + x^2),x]`

output `E^ArcTan[x]`

3.316.3.1 Defintions of rubi rules used

rule 5594 `Int[E^(ArcTan[(a_.)*(x_)])*(n_.)/((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]`

3.316.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

method	result	size
gospers	$e^{\arctan(x)}$	4
derivativedivides	$e^{\arctan(x)}$	4
default	$e^{\arctan(x)}$	4
parallelrisch	$e^{\arctan(x)}$	4
risch	$(-ix + 1)^{\frac{i}{2}} (ix + 1)^{-\frac{i}{2}}$	20

input `int(exp(arctan(x))/(x^2+1),x,method=_RETURNVERBOSE)`

output `exp(arctan(x))`

3.316.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \frac{e^{\arctan(x)}}{1+x^2} dx = e^{\arctan(x)}$$

input `integrate(exp(arctan(x))/(x^2+1),x, algorithm="fricas")`

output `e^arctan(x)`

3.316.6 Sympy [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \frac{e^{\arctan(x)}}{1+x^2} dx = e^{\arctan(x)}$$

input `integrate(exp(atan(x))/(x**2+1),x)`

output `exp(atan(x))`

3.316.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \frac{e^{\arctan(x)}}{1+x^2} dx = e^{\arctan(x)}$$

input `integrate(exp(arctan(x))/(x^2+1),x, algorithm="maxima")`

output `e^arctan(x)`

3.316.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \frac{e^{\arctan(x)}}{1+x^2} dx = e^{\arctan(x)}$$

input `integrate(exp(arctan(x))/(x^2+1),x, algorithm="giac")`

output `e^arctan(x)`

3.316.9 Mupad [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \frac{e^{\arctan(x)}}{1+x^2} dx = e^{\arctan(x)}$$

input `int(exp(atan(x))/(x^2 + 1),x)`

output `exp(atan(x))`

$$\mathbf{3.317} \quad \int \frac{1}{x(1+x^4)} dx$$

3.317.1 Optimal result	1603
3.317.2 Mathematica [A] (verified)	1603
3.317.3 Rubi [A] (verified)	1604
3.317.4 Maple [A] (verified)	1605
3.317.5 Fricas [A] (verification not implemented)	1605
3.317.6 Sympy [A] (verification not implemented)	1606
3.317.7 Maxima [A] (verification not implemented)	1606
3.317.8 Giac [A] (verification not implemented)	1606
3.317.9 Mupad [B] (verification not implemented)	1607

3.317.1 Optimal result

Integrand size = 11, antiderivative size = 13

$$\int \frac{1}{x(1+x^4)} dx = \log(x) - \frac{1}{4} \log(1+x^4)$$

output `ln(x)-1/4*ln(x^4+1)`

3.317.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+x^4)} dx = \log(x) - \frac{1}{4} \log(1+x^4)$$

input `Integrate[1/(x*(1 + x^4)),x]`

output `Log[x] - Log[1 + x^4]/4`

3.317.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {798, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x(x^4+1)} dx \\
 & \quad \downarrow 798 \\
 & \frac{1}{4} \int \frac{1}{x^4(x^4+1)} dx^4 \\
 & \quad \downarrow 47 \\
 & \frac{1}{4} \left(\int \frac{1}{x^4} dx^4 - \int \frac{1}{x^4+1} dx^4 \right) \\
 & \quad \downarrow 14 \\
 & \frac{1}{4} \left(\log(x^4) - \int \frac{1}{x^4+1} dx^4 \right) \\
 & \quad \downarrow 16 \\
 & \frac{1}{4} (\log(x^4) - \log(x^4+1))
 \end{aligned}$$

input `Int[1/(x*(1 + x^4)),x]`

output `(Log[x^4] - Log[1 + x^4])/4`

3.317.3.1 Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

```
rule 47 Int[1/((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[b/(b*c
- a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x
], x] /; FreeQ[{a, b, c, d}, x]
```

```
rule 798 Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^(p_), x_Symbol] :> Simp[1/n Subst
[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a,
b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

3.317.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
default	$\ln(x) - \frac{\ln(x^4+1)}{4}$	12
norman	$\ln(x) - \frac{\ln(x^4+1)}{4}$	12
meijerg	$\ln(x) - \frac{\ln(x^4+1)}{4}$	12
risch	$\ln(x) - \frac{\ln(x^4+1)}{4}$	12
parallelrisch	$\ln(x) - \frac{\ln(x^4+1)}{4}$	12

```
input int(1/x/(x^4+1),x,method=_RETURNVERBOSE)
```

```
output ln(x)-1/4*ln(x^4+1)
```

3.317.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1}{x(1+x^4)} dx = -\frac{1}{4} \log(x^4 + 1) + \log(x)$$

```
input integrate(1/x/(x^4+1),x, algorithm="fracas")
```

```
output -1/4*log(x^4 + 1) + log(x)
```

3.317.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \frac{1}{x(1+x^4)} dx = \log(x) - \frac{\log(x^4+1)}{4}$$

input `integrate(1/x/(x**4+1),x)`output `log(x) - log(x**4 + 1)/4`**3.317.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{1}{x(1+x^4)} dx = -\frac{1}{4} \log(x^4+1) + \frac{1}{4} \log(x^4)$$

input `integrate(1/x/(x^4+1),x, algorithm="maxima")`output `-1/4*log(x^4 + 1) + 1/4*log(x^4)`**3.317.8 Giac [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{1}{x(1+x^4)} dx = -\frac{1}{4} \log(x^4+1) + \frac{1}{4} \log(x^4)$$

input `integrate(1/x/(x^4+1),x, algorithm="giac")`output `-1/4*log(x^4 + 1) + 1/4*log(x^4)`

3.317.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1}{x(1+x^4)} dx = \ln(x) - \frac{\ln(x^4+1)}{4}$$

input `int(1/(x*(x^4 + 1)),x)`

output `log(x) - log(x^4 + 1)/4`

3.318 $\int e^{-2t}t^3 dt$

3.318.1 Optimal result	1608
3.318.2 Mathematica [A] (verified)	1608
3.318.3 Rubi [A] (verified)	1609
3.318.4 Maple [A] (verified)	1610
3.318.5 Fricas [A] (verification not implemented)	1610
3.318.6 Sympy [A] (verification not implemented)	1611
3.318.7 Maxima [A] (verification not implemented)	1611
3.318.8 Giac [A] (verification not implemented)	1611
3.318.9 Mupad [B] (verification not implemented)	1612

3.318.1 Optimal result

Integrand size = 9, antiderivative size = 44

$$\int e^{-2t}t^3 dt = -\frac{3}{8}e^{-2t} - \frac{3}{4}e^{-2t}t - \frac{3}{4}e^{-2t}t^2 - \frac{1}{2}e^{-2t}t^3$$

output `-3/8/exp(2*t)-3/4*t/exp(2*t)-3/4*t^2/exp(2*t)-1/2*t^3/exp(2*t)`

3.318.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.55

$$\int e^{-2t}t^3 dt = -\frac{1}{8}e^{-2t}(3 + 6t + 6t^2 + 4t^3)$$

input `Integrate[t^3/E^(2*t),t]`

output `-1/8*(3 + 6*t + 6*t^2 + 4*t^3)/E^(2*t)`

3.318.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.11, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {2607, 2607, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{-2t} t^3 dt \\
 & \quad \downarrow \text{2607} \\
 & \frac{3}{2} \int e^{-2t} t^2 dt - \frac{1}{2} e^{-2t} t^3 \\
 & \quad \downarrow \text{2607} \\
 & \frac{3}{2} \left(\int e^{-2t} t dt - \frac{1}{2} e^{-2t} t^2 \right) - \frac{1}{2} e^{-2t} t^3 \\
 & \quad \downarrow \text{2607} \\
 & \frac{3}{2} \left(\frac{1}{2} \int e^{-2t} dt - \frac{1}{2} e^{-2t} t^2 - \frac{1}{2} e^{-2t} t \right) - \frac{1}{2} e^{-2t} t^3 \\
 & \quad \downarrow \text{2624} \\
 & \frac{3}{2} \left(-\frac{1}{2} e^{-2t} t^2 - \frac{1}{2} e^{-2t} t - \frac{e^{-2t}}{4} \right) - \frac{1}{2} e^{-2t} t^3
 \end{aligned}$$

input `Int[t^3/E^(2*t),t]`

output `-1/2*t^3/E^(2*t) + (3*(-1/4*1/E^(2*t) - t/(2*E^(2*t)) - t^2/(2*E^(2*t))))/2`

3.318.3.1 Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

```
rule 2624 Int[((F_)^(v_))^(n_), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;
FreeQ[{F, n}, x] && LinearQ[v, x]
```

3.318.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.48

method	result	size
risch	$(-\frac{1}{2}t^3 - \frac{3}{4}t^2 - \frac{3}{4}t - \frac{3}{8})e^{-2t}$	21
norman	$(-\frac{1}{2}t^3 - \frac{3}{4}t^2 - \frac{3}{4}t - \frac{3}{8})e^{-2t}$	23
gosper	$-\frac{(4t^3+6t^2+6t+3)e^{-2t}}{8}$	24
meijerg	$\frac{3}{8} - \frac{(32t^3+48t^2+48t+24)e^{-2t}}{64}$	24
parallelrisch	$\frac{(-4t^3-6t^2-6t-3)e^{-2t}}{8}$	24
derivativedivides	$-\frac{3e^{-2t}}{8} - \frac{3te^{-2t}}{4} - \frac{3t^2e^{-2t}}{4} - \frac{t^3e^{-2t}}{2}$	41
default	$-\frac{3e^{-2t}}{8} - \frac{3te^{-2t}}{4} - \frac{3t^2e^{-2t}}{4} - \frac{t^3e^{-2t}}{2}$	41

```
input int(t^3/exp(2*t),t,method=_RETURNVERBOSE)
```

```
output (-1/2*t^3-3/4*t^2-3/4*t-3/8)*exp(-2*t)
```

3.318.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.48

$$\int e^{-2t}t^3 dt = -\frac{1}{8}(4t^3 + 6t^2 + 6t + 3)e^{(-2t)}$$

```
input integrate(t^3/exp(2*t),t, algorithm="fricas")
```

```
output -1/8*(4*t^3 + 6*t^2 + 6*t + 3)*e^(-2*t)
```

3.318.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.50

$$\int e^{-2t} t^3 dt = \frac{(-4t^3 - 6t^2 - 6t - 3) e^{-2t}}{8}$$

input `integrate(t**3/exp(2*t),t)`output `(-4*t**3 - 6*t**2 - 6*t - 3)*exp(-2*t)/8`**3.318.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.48

$$\int e^{-2t} t^3 dt = -\frac{1}{8} (4t^3 + 6t^2 + 6t + 3) e^{(-2t)}$$

input `integrate(t^3/exp(2*t),t, algorithm="maxima")`output `-1/8*(4*t^3 + 6*t^2 + 6*t + 3)*e^(-2*t)`**3.318.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.48

$$\int e^{-2t} t^3 dt = -\frac{1}{8} (4t^3 + 6t^2 + 6t + 3) e^{(-2t)}$$

input `integrate(t^3/exp(2*t),t, algorithm="giac")`output `-1/8*(4*t^3 + 6*t^2 + 6*t + 3)*e^(-2*t)`

3.318.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.48

$$\int e^{-2t} t^3 dt = -\frac{e^{-2t} (8t^3 + 12t^2 + 12t + 6)}{16}$$

input `int(t^3*exp(-2*t),t)`

output `-(exp(-2*t)*(12*t + 12*t^2 + 8*t^3 + 6))/16`

$$3.319 \quad \int \frac{\sqrt{t}}{1 + \sqrt[3]{t}} dt$$

3.319.1 Optimal result	1613
3.319.2 Mathematica [A] (verified)	1613
3.319.3 Rubi [A] (verified)	1614
3.319.4 Maple [A] (verified)	1616
3.319.5 Fracas [A] (verification not implemented)	1616
3.319.6 Sympy [A] (verification not implemented)	1616
3.319.7 Maxima [A] (verification not implemented)	1617
3.319.8 Giac [A] (verification not implemented)	1617
3.319.9 Mupad [B] (verification not implemented)	1617

3.319.1 Optimal result

Integrand size = 15, antiderivative size = 41

$$\int \frac{\sqrt{t}}{1 + \sqrt[3]{t}} dt = -6\sqrt[6]{t} + 2\sqrt{t} - \frac{6t^{5/6}}{5} + \frac{6t^{7/6}}{7} + 6 \arctan\left(\sqrt[6]{t}\right)$$

output `-6*t^(1/6)-6/5*t^(5/6)+6/7*t^(7/6)+6*arctan(t^(1/6))+2*t^(1/2)`

3.319.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.02

$$\int \frac{\sqrt{t}}{1 + \sqrt[3]{t}} dt = \frac{2}{35} \left(-105\sqrt[6]{t} + 35\sqrt{t} - 21t^{5/6} + 15t^{7/6} \right) + 6 \arctan\left(\sqrt[6]{t}\right)$$

input `Integrate[Sqrt[t]/(1 + t^(1/3)),t]`

output `(2*(-105*t^(1/6) + 35*Sqrt[t] - 21*t^(5/6) + 15*t^(7/6)))/35 + 6*ArcTan[t^(1/6)]`

3.319.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.10, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {864, 60, 60, 60, 60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{t}}{\sqrt[3]{t}+1} dt \\
 & \quad \downarrow 864 \\
 & 3 \int \frac{t^{7/6}}{\sqrt[3]{t}+1} d\sqrt[3]{t} \\
 & \quad \downarrow 60 \\
 & 3 \left(\frac{2t^{7/6}}{7} - \int \frac{t^{5/6}}{\sqrt[3]{t}+1} d\sqrt[3]{t} \right) \\
 & \quad \downarrow 60 \\
 & 3 \left(\int \frac{\sqrt{t}}{\sqrt[3]{t}+1} d\sqrt[3]{t} + \frac{2t^{7/6}}{7} - \frac{2t^{5/6}}{5} \right) \\
 & \quad \downarrow 60 \\
 & 3 \left(- \int \frac{\sqrt[6]{t}}{\sqrt[3]{t}+1} d\sqrt[3]{t} + \frac{2t^{7/6}}{7} - \frac{2t^{5/6}}{5} + \frac{2\sqrt{t}}{3} \right) \\
 & \quad \downarrow 60 \\
 & 3 \left(\int \frac{1}{(\sqrt[3]{t}+1)\sqrt[6]{t}} d\sqrt[3]{t} + \frac{2t^{7/6}}{7} - \frac{2t^{5/6}}{5} + \frac{2\sqrt{t}}{3} - 2\sqrt[6]{t} \right) \\
 & \quad \downarrow 73 \\
 & 3 \left(2 \int \frac{1}{t^{2/3}+1} d\sqrt[6]{t} + \frac{2t^{7/6}}{7} - \frac{2t^{5/6}}{5} + \frac{2\sqrt{t}}{3} - 2\sqrt[6]{t} \right) \\
 & \quad \downarrow 216 \\
 & 3 \left(2 \arctan(\sqrt[6]{t}) + \frac{2t^{7/6}}{7} - \frac{2t^{5/6}}{5} + \frac{2\sqrt{t}}{3} - 2\sqrt[6]{t} \right)
 \end{aligned}$$

input `Int[Sqrt[t]/(1 + t^(1/3)),t]`

output `3*(-2*t^(1/6) + (2*Sqrt[t])/3 - (2*t^(5/6))/5 + (2*t^(7/6))/7 + 2*ArcTan[t^(1/6)])`

3.319.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 864 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a + b*x^(k*n))^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, m, p}, x] && FractionQ[n]`

3.319.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.68

method	result	size
derivativedivides	$-6t^{\frac{1}{6}} - \frac{6t^{\frac{5}{6}}}{5} + \frac{6t^{\frac{7}{6}}}{7} + 6 \arctan\left(t^{\frac{1}{6}}\right) + 2\sqrt{t}$	28
default	$-6t^{\frac{1}{6}} - \frac{6t^{\frac{5}{6}}}{5} + \frac{6t^{\frac{7}{6}}}{7} + 6 \arctan\left(t^{\frac{1}{6}}\right) + 2\sqrt{t}$	28
meijerg	$-\frac{2t^{\frac{1}{6}}(-45t+63t^{\frac{2}{3}}-105t^{\frac{1}{3}}+315)}{105} + 6 \arctan\left(t^{\frac{1}{6}}\right)$	28

input `int(t^(1/2)/(1+t^(1/3)),t,method=_RETURNVERBOSE)`output `-6*t^(1/6)-6/5*t^(5/6)+6/7*t^(7/6)+6*arctan(t^(1/6))+2*t^(1/2)`**3.319.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.61

$$\int \frac{\sqrt{t}}{1+\sqrt[3]{t}} dt = \frac{6}{7}(t-7)t^{\frac{1}{6}} - \frac{6}{5}t^{\frac{5}{6}} + 2\sqrt{t} + 6 \arctan\left(t^{\frac{1}{6}}\right)$$

input `integrate(t^(1/2)/(1+t^(1/3)),t, algorithm="fricas")`output `6/7*(t - 7)*t^(1/6) - 6/5*t^(5/6) + 2*sqrt(t) + 6*arctan(t^(1/6))`**3.319.6 Sympy [A] (verification not implemented)**

Time = 1.70 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.90

$$\int \frac{\sqrt{t}}{1+\sqrt[3]{t}} dt = \frac{6t^{\frac{7}{6}}}{7} - \frac{6t^{\frac{5}{6}}}{5} - 6\sqrt[6]{t} + 2\sqrt{t} + 6 \operatorname{atan}\left(\sqrt[6]{t}\right)$$

input `integrate(t**(1/2)/(1+t**(1/3)),t)`output `6*t**(7/6)/7 - 6*t**(5/6)/5 - 6*t**(1/6) + 2*sqrt(t) + 6*atan(t**(1/6))`

3.319. $\int \frac{\sqrt{t}}{1+\sqrt[3]{t}} dt$

3.319.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.66

$$\int \frac{\sqrt{t}}{1 + \sqrt[3]{t}} dt = \frac{6}{7} t^{\frac{7}{6}} - \frac{6}{5} t^{\frac{5}{6}} + 2\sqrt{t} - 6t^{\frac{1}{6}} + 6 \arctan\left(t^{\frac{1}{6}}\right)$$

input `integrate(t^(1/2)/(1+t^(1/3)),t, algorithm="maxima")`output `6/7*t^(7/6) - 6/5*t^(5/6) + 2*sqrt(t) - 6*t^(1/6) + 6*arctan(t^(1/6))`**3.319.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.66

$$\int \frac{\sqrt{t}}{1 + \sqrt[3]{t}} dt = \frac{6}{7} t^{\frac{7}{6}} - \frac{6}{5} t^{\frac{5}{6}} + 2\sqrt{t} - 6t^{\frac{1}{6}} + 6 \arctan\left(t^{\frac{1}{6}}\right)$$

input `integrate(t^(1/2)/(1+t^(1/3)),t, algorithm="giac")`output `6/7*t^(7/6) - 6/5*t^(5/6) + 2*sqrt(t) - 6*t^(1/6) + 6*arctan(t^(1/6))`**3.319.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.66

$$\int \frac{\sqrt{t}}{1 + \sqrt[3]{t}} dt = 6 \operatorname{atan}\left(t^{1/6}\right) + 2\sqrt{t} - 6t^{1/6} - \frac{6t^{5/6}}{5} + \frac{6t^{7/6}}{7}$$

input `int(t^(1/2)/(t^(1/3) + 1),t)`output `6*atan(t^(1/6)) + 2*t^(1/2) - 6*t^(1/6) - (6*t^(5/6))/5 + (6*t^(7/6))/7`

3.320 $\int \sin(x) \sin(2x) \sin(3x) dx$

3.320.1 Optimal result	1618
3.320.2 Mathematica [A] (verified)	1618
3.320.3 Rubi [A] (verified)	1619
3.320.4 Maple [A] (verified)	1620
3.320.5 Fricas [A] (verification not implemented)	1620
3.320.6 Sympy [B] (verification not implemented)	1620
3.320.7 Maxima [A] (verification not implemented)	1621
3.320.8 Giac [A] (verification not implemented)	1621
3.320.9 Mupad [B] (verification not implemented)	1621

3.320.1 Optimal result

Integrand size = 11, antiderivative size = 25

$$\int \sin(x) \sin(2x) \sin(3x) dx = -\frac{1}{8} \cos(2x) - \frac{1}{16} \cos(4x) + \frac{1}{24} \cos(6x)$$

output `-1/8*cos(2*x)-1/16*cos(4*x)+1/24*cos(6*x)`

3.320.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \sin(x) \sin(2x) \sin(3x) dx = -\frac{1}{8} \cos(2x) - \frac{1}{16} \cos(4x) + \frac{1}{24} \cos(6x)$$

input `Integrate[Sin[x]*Sin[2*x]*Sin[3*x],x]`

output `-1/8*Cos[2*x] - Cos[4*x]/16 + Cos[6*x]/24`

3.320.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 4855, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin(x) \sin(2x) \sin(3x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin(x) \sin(2x) \sin(3x) dx \\ & \quad \downarrow \text{4855} \\ & \int \left(\frac{1}{4} \sin(2x) + \frac{1}{4} \sin(4x) - \frac{1}{4} \sin(6x) \right) dx \\ & \quad \downarrow \text{2009} \\ & -\frac{1}{8} \cos(2x) - \frac{1}{16} \cos(4x) + \frac{1}{24} \cos(6x) \end{aligned}$$

input `Int[Sin[x]*Sin[2*x]*Sin[3*x],x]`

output `-1/8*Cos[2*x] - Cos[4*x]/16 + Cos[6*x]/24`

3.320.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4855 `Int[(F_)[(a_.) + (b_.)*(x_)]^(p_.)*(G_)[(c_.) + (d_.)*(x_)]^(q_.)*(H_)[(e_.) + (f_.)*(x_)]^(r_.), x_Symbol] := Int[ExpandTrigReduce[ActivateTrig[F[a + b*x]^p*G[c + d*x]^q*H[e + f*x]^r], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && (EqQ[F, sin] || EqQ[F, cos]) && (EqQ[G, sin] || EqQ[G, cos]) && (EqQ[H, sin] || EqQ[H, cos]) && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]`

3.320.4 Maple [A] (verified)

Time = 1.34 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

method	result	size
default	$-\frac{\cos(2x)}{8} - \frac{\cos(4x)}{16} + \frac{\cos(6x)}{24}$	20
risch	$-\frac{\cos(2x)}{8} - \frac{\cos(4x)}{16} + \frac{\cos(6x)}{24}$	20
parallelrisch	$-\frac{29}{48} + \frac{\cos(6x)}{24} - \frac{\cos(4x)}{16} - \frac{\cos(2x)}{8}$	21

input `int(sin(x)*sin(2*x)*sin(3*x),x,method=_RETURNVERBOSE)`output `-1/8*cos(2*x)-1/16*cos(4*x)+1/24*cos(6*x)`**3.320.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.68

$$\int \sin(x) \sin(2x) \sin(3x) dx = \frac{4}{3} \cos(x)^6 - \frac{5}{2} \cos(x)^4 + \cos(x)^2$$

input `integrate(sin(x)*sin(2*x)*sin(3*x),x, algorithm="fricas")`output `4/3*cos(x)^6 - 5/2*cos(x)^4 + cos(x)^2`**3.320.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 114 vs. 2(19) = 38.

Time = 0.96 (sec) , antiderivative size = 114, normalized size of antiderivative = 4.56

$$\int \sin(x) \sin(2x) \sin(3x) dx = \frac{x \sin(x) \sin(2x) \sin(3x)}{4} + \frac{x \sin(x) \cos(2x) \cos(3x)}{4} + \frac{x \sin(2x) \cos(x) \cos(3x)}{4} - \frac{x \sin(3x) \cos(x) \cos(2x)}{4} - \frac{5 \sin(x) \sin(2x) \cos(3x)}{24} - \frac{\sin(2x) \sin(3x) \cos(x)}{8} - \frac{\cos(x) \cos(2x) \cos(3x)}{6}$$

input `integrate(sin(x)*sin(2*x)*sin(3*x),x)`

output `x*sin(x)*sin(2*x)*sin(3*x)/4 + x*sin(x)*cos(2*x)*cos(3*x)/4 + x*sin(2*x)*cos(x)*cos(3*x)/4 - x*sin(3*x)*cos(x)*cos(2*x)/4 - 5*sin(x)*sin(2*x)*cos(3*x)/24 - sin(2*x)*sin(3*x)*cos(x)/8 - cos(x)*cos(2*x)*cos(3*x)/6`

3.320.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \sin(x) \sin(2x) \sin(3x) dx = \frac{1}{24} \cos(6x) - \frac{1}{16} \cos(4x) - \frac{1}{8} \cos(2x)$$

input `integrate(sin(x)*sin(2*x)*sin(3*x),x, algorithm="maxima")`

output `1/24*cos(6*x) - 1/16*cos(4*x) - 1/8*cos(2*x)`

3.320.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.52

$$\int \sin(x) \sin(2x) \sin(3x) dx = -\frac{4}{3} \sin(x)^6 + \frac{3}{2} \sin(x)^4$$

input `integrate(sin(x)*sin(2*x)*sin(3*x),x, algorithm="giac")`

output `-4/3*sin(x)^6 + 3/2*sin(x)^4`

3.320.9 Mupad [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.56

$$\int \sin(x) \sin(2x) \sin(3x) dx = -\frac{\sin(x)^4 (8 \sin(x)^2 - 9)}{6}$$

input `int(sin(2*x)*sin(3*x)*sin(x),x)`

output `-(sin(x)^4*(8*sin(x)^2 - 9))/6`

3.321 $\int \log\left(\frac{x}{2}\right) dx$

3.321.1 Optimal result	1622
3.321.2 Mathematica [A] (verified)	1622
3.321.3 Rubi [A] (verified)	1623
3.321.4 Maple [A] (verified)	1623
3.321.5 Fricas [A] (verification not implemented)	1624
3.321.6 Sympy [A] (verification not implemented)	1624
3.321.7 Maxima [A] (verification not implemented)	1624
3.321.8 Giac [A] (verification not implemented)	1625
3.321.9 Mupad [B] (verification not implemented)	1625

3.321.1 Optimal result

Integrand size = 6, antiderivative size = 12

$$\int \log\left(\frac{x}{2}\right) dx = -x + x \log\left(\frac{x}{2}\right)$$

output `-x+x*ln(1/2*x)`

3.321.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \log\left(\frac{x}{2}\right) dx = -x + x \log\left(\frac{x}{2}\right)$$

input `Integrate[Log[x/2],x]`

output `-x + x*Log[x/2]`

3.321.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2732}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \log\left(\frac{x}{2}\right) dx$$

$$\downarrow \text{2732}$$

$$x \log\left(\frac{x}{2}\right) - x$$

input `Int[Log[x/2], x]`

output `-x + x*Log[x/2]`

3.321.3.1 Defintions of rubi rules used

rule 2732 `Int[Log[(c_.)*(x_)^(n_.)], x_Symbol] := Simp[x*Log[c*x^n], x] - Simp[n*x, x] /; FreeQ[{c, n}, x]`

3.321.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
derivativdivides	$-x + x \ln\left(\frac{x}{2}\right)$	11
default	$-x + x \ln\left(\frac{x}{2}\right)$	11
norman	$-x + x \ln\left(\frac{x}{2}\right)$	11
risch	$-x + x \ln\left(\frac{x}{2}\right)$	11
parallelrisc	$-x + x \ln\left(\frac{x}{2}\right)$	11
parts	$-x + x \ln\left(\frac{x}{2}\right)$	11

input `int(ln(1/2*x), x, method=_RETURNVERBOSE)`

output `-x+x*ln(1/2*x)`

3.321.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \log\left(\frac{x}{2}\right) dx = x \log\left(\frac{1}{2}x\right) - x$$

input `integrate(log(1/2*x),x, algorithm="fricas")`

output `x*log(1/2*x) - x`

3.321.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.58

$$\int \log\left(\frac{x}{2}\right) dx = x \log\left(\frac{x}{2}\right) - x$$

input `integrate(ln(1/2*x),x)`

output `x*log(x/2) - x`

3.321.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \log\left(\frac{x}{2}\right) dx = x \log\left(\frac{1}{2}x\right) - x$$

input `integrate(log(1/2*x),x, algorithm="maxima")`

output `x*log(1/2*x) - x`

3.321.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \log\left(\frac{x}{2}\right) dx = x \log\left(\frac{1}{2}x\right) - x$$

input `integrate(log(1/2*x),x, algorithm="giac")`

output `x*log(1/2*x) - x`

3.321.9 Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \log\left(\frac{x}{2}\right) dx = x \left(\ln\left(\frac{x}{2}\right) - 1 \right)$$

input `int(log(x/2),x)`

output `x*(log(x/2) - 1)`

$$3.322 \quad \int \sqrt{\frac{1+x}{1-x}} dx$$

3.322.1 Optimal result	1626
3.322.2 Mathematica [A] (verified)	1626
3.322.3 Rubi [A] (verified)	1627
3.322.4 Maple [A] (verified)	1628
3.322.5 Fricas [A] (verification not implemented)	1628
3.322.6 Sympy [F]	1629
3.322.7 Maxima [A] (verification not implemented)	1629
3.322.8 Giac [A] (verification not implemented)	1629
3.322.9 Mupad [B] (verification not implemented)	1630

3.322.1 Optimal result

Integrand size = 15, antiderivative size = 41

$$\int \sqrt{\frac{1+x}{1-x}} dx = -\left((1-x)\sqrt{\frac{1+x}{1-x}}\right) + 2 \arctan\left(\sqrt{\frac{1+x}{1-x}}\right)$$

output `2*arctan(((1+x)/(1-x))^(1/2))- (1-x)*((1+x)/(1-x))^(1/2)`

3.322.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.59

$$\int \sqrt{\frac{1+x}{1-x}} dx = -\frac{\sqrt{1-x}\sqrt{\frac{1+x}{1-x}}\left(\sqrt{1-x^2} + 2 \arctan\left(\frac{\sqrt{1-x^2}}{-1+x}\right)\right)}{\sqrt{1+x}}$$

input `Integrate[Sqrt[(1 + x)/(1 - x)],x]`

output `-((Sqrt[1 - x]*Sqrt[(1 + x)/(1 - x)]*(Sqrt[1 - x^2] + 2*ArcTan[Sqrt[1 - x^2]/(-1 + x)]))/Sqrt[1 + x])`

3.322. $\int \sqrt{\frac{1+x}{1-x}} dx$

3.322.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.39, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2051, 252, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\frac{x+1}{1-x}} dx \\
 & \quad \downarrow \text{2051} \\
 & 4 \int \frac{x+1}{(1-x) \left(\frac{x+1}{1-x} + 1\right)^2} d\sqrt{\frac{x+1}{1-x}} \\
 & \quad \downarrow \text{252} \\
 & 4 \left(\frac{1}{2} \int \frac{1}{\frac{x+1}{1-x} + 1} d\sqrt{\frac{x+1}{1-x}} - \frac{\sqrt{\frac{x+1}{1-x}}}{2 \left(\frac{x+1}{1-x} + 1\right)} \right) \\
 & \quad \downarrow \text{216} \\
 & 4 \left(\frac{1}{2} \arctan \left(\sqrt{\frac{x+1}{1-x}} \right) - \frac{\sqrt{\frac{x+1}{1-x}}}{2 \left(\frac{x+1}{1-x} + 1\right)} \right)
 \end{aligned}$$

input `Int[Sqrt[(1 + x)/(1 - x)],x]`

output `4*(-1/2*Sqrt[(1 + x)/(1 - x)]/(1 + (1 + x)/(1 - x)) + ArcTan[Sqrt[(1 + x)/(1 - x)]])/2`

3.322.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

```
rule 252 Int[((c_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*((m - 1)/(2*b*(p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

```
rule 2051 Int[(((e_.)*((a_.) + (b_.)*(x_)^(n_.)))/((c_) + (d_.)*(x_)^(n_.)))^(p_), x_Symbol] := With[{q = Denominator[p]}, Simp[q*e*((b*c - a*d)/n) Subst[Int[x^(q*(p + 1) - 1)*((-a)*e + c*x^q)^(1/n - 1)/(b*e - d*x^q)^(1/n + 1), x], x, (e*((a + b*x^n)/(c + d*x^n)))^(1/q)], x] /; FreeQ[{a, b, c, d, e}, x] && FractionQ[p] && IntegerQ[1/n]
```

3.322.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

method	result
default	$\frac{\sqrt{-\frac{1+x}{-1+x}}(-1+x)(\sqrt{-x^2+1}-\arcsin(x))}{\sqrt{-(-1+x)(1+x)}}$
risch	$(-1+x)\sqrt{-\frac{1+x}{-1+x}} + \frac{\arcsin(x)\sqrt{-\frac{1+x}{-1+x}}\sqrt{-(-1+x)(1+x)}}{1+x}$
trager	$(-1+x)\sqrt{-\frac{1+x}{-1+x}} + \text{RootOf}(_Z^2+1)\ln\left(-\text{RootOf}(_Z^2+1)\sqrt{-\frac{1+x}{-1+x}}x + \text{RootOf}(_Z^2+1)\right)$

```
input int(((1+x)/(1-x))^(1/2),x,method=_RETURNVERBOSE)
```

```
output ((-1+x)/(-1+x))^(1/2)*(-1+x)/(-(-1+x)*(1+x))^(1/2)*((-x^2+1)^(1/2)-arcsin(x))
```

3.322.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.78

$$\int \sqrt{\frac{1+x}{1-x}} dx = (x-1)\sqrt{-\frac{x+1}{x-1}} + 2 \arctan\left(\sqrt{-\frac{x+1}{x-1}}\right)$$

```
input integrate(((1+x)/(1-x))^(1/2),x, algorithm="fricas")
```

3.322. $\int \sqrt{\frac{1+x}{1-x}} dx$

output $(x - 1)\sqrt{-(x + 1)/(x - 1)} + 2\arctan(\sqrt{-(x + 1)/(x - 1)})$

3.322.6 Sympy [F]

$$\int \sqrt{\frac{1+x}{1-x}} dx = \int \sqrt{\frac{x+1}{1-x}} dx$$

input `integrate(((1+x)/(1-x))**(1/2),x)`

output `Integral(sqrt((x + 1)/(1 - x)), x)`

3.322.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.05

$$\int \sqrt{\frac{1+x}{1-x}} dx = \frac{2\sqrt{-\frac{x+1}{x-1}}}{\frac{x+1}{x-1} - 1} + 2 \arctan\left(\sqrt{-\frac{x+1}{x-1}}\right)$$

input `integrate(((1+x)/(1-x))^(1/2),x, algorithm="maxima")`

output $2\sqrt{-(x + 1)/(x - 1)}/((x + 1)/(x - 1) - 1) + 2\arctan(\sqrt{-(x + 1)/(x - 1)})$

3.322.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.73

$$\int \sqrt{\frac{1+x}{1-x}} dx = \frac{1}{2} \pi \operatorname{sgn}(x - 1) - \arcsin(x) \operatorname{sgn}(x - 1) + \sqrt{-x^2 + 1} \operatorname{sgn}(x - 1)$$

input `integrate(((1+x)/(1-x))^(1/2),x, algorithm="giac")`

output $1/2\pi\operatorname{sgn}(x - 1) - \arcsin(x)\operatorname{sgn}(x - 1) + \sqrt{-x^2 + 1}\operatorname{sgn}(x - 1)$

3.322.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.05

$$\int \sqrt{\frac{1+x}{1-x}} dx = 2 \operatorname{atan}\left(\sqrt{-\frac{x+1}{x-1}}\right) + \frac{2\sqrt{-\frac{x+1}{x-1}}}{\frac{x+1}{x-1} - 1}$$

input `int((-x + 1)/(x - 1)^(1/2),x)`output `2*atan((-x + 1)/(x - 1)^(1/2)) + (2*(-x + 1)/(x - 1)^(1/2))/((x + 1)/(x - 1) - 1)`

3.323 $\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx$

3.323.1 Optimal result	1631
3.323.2 Mathematica [A] (verified)	1631
3.323.3 Rubi [A] (verified)	1632
3.323.4 Maple [C] (warning: unable to verify)	1634
3.323.5 Fricas [A] (verification not implemented)	1634
3.323.6 Sympy [A] (verification not implemented)	1634
3.323.7 Maxima [A] (verification not implemented)	1635
3.323.8 Giac [A] (verification not implemented)	1635
3.323.9 Mupad [F(-1)]	1635

3.323.1 Optimal result

Integrand size = 13, antiderivative size = 34

$$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx = -\sqrt{-1+x^2} + \arctan(\sqrt{-1+x^2}) + \sqrt{-1+x^2} \log(x)$$

output `arctan((x^2-1)^(1/2))-sqrt(-1+x^2)+ln(x)*sqrt(-1+x^2)`

3.323.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.79

$$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx = -\arctan\left(\frac{1}{\sqrt{-1+x^2}}\right) + \sqrt{-1+x^2}(-1 + \log(x))$$

input `Integrate[(x*Log[x])/Sqrt[-1 + x^2],x]`

output `-ArcTan[1/Sqrt[-1 + x^2]] + Sqrt[-1 + x^2]*(-1 + Log[x])`

3.323.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.21, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {2776, 243, 60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x \log(x)}{\sqrt{x^2-1}} dx \\
 & \quad \downarrow \text{2776} \\
 & \sqrt{x^2-1} \log(x) - \int \frac{\sqrt{x^2-1}}{x} dx \\
 & \quad \downarrow \text{243} \\
 & \sqrt{x^2-1} \log(x) - \frac{1}{2} \int \frac{\sqrt{x^2-1}}{x^2} dx^2 \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2} \left(\int \frac{1}{x^2 \sqrt{x^2-1}} dx^2 - 2\sqrt{x^2-1} \right) + \sqrt{x^2-1} \log(x) \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{2} \left(2 \int \frac{1}{x^4+1} d\sqrt{x^2-1} - 2\sqrt{x^2-1} \right) + \sqrt{x^2-1} \log(x) \\
 & \quad \downarrow \text{216} \\
 & \frac{1}{2} \left(2 \arctan(\sqrt{x^2-1}) - 2\sqrt{x^2-1} \right) + \sqrt{x^2-1} \log(x)
 \end{aligned}$$

input `Int[(x*Log[x])/Sqrt[-1 + x^2],x]`

output `(-2*Sqrt[-1 + x^2] + 2*ArcTan[Sqrt[-1 + x^2]])/2 + Sqrt[-1 + x^2]*Log[x]`

3.323.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2776 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((f_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^(r_))^(q_.), x_Symbol] := Simp[f^m*(d + e*x^r)^(q + 1)*((a + b*Log[c*x^n])^p/(e*r*(q + 1))), x] - Simp[b*f^m*n*(p/(e*r*(q + 1)) Int[(d + e*x^r)^(q + 1)*((a + b*Log[c*x^n])^(p - 1)/x), x], x] /; FreeQ[{a, b, c, d, e, f, m, n, q, r}, x] && EqQ[m, r - 1] && IGtQ[p, 0] && (IntegerQ[m] || GtQ[f, 0]) && NeQ[r, n] && NeQ[q, -1]`

3.323.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.16 (sec) , antiderivative size = 119, normalized size of antiderivative = 3.50

method	result
meijerg	$-\frac{\sqrt{-\operatorname{signum}(x^2-1)}(2-2\sqrt{-x^2+1})}{4\sqrt{\operatorname{signum}(x^2-1)}} + \frac{\sqrt{-\operatorname{signum}(x^2-1)}\ln(x)(2-2\sqrt{-x^2+1})}{2\sqrt{\operatorname{signum}(x^2-1)}} + \frac{\sqrt{-\operatorname{signum}(x^2-1)}\left(-16+16\sqrt{-x^2+1}-32\sqrt{\operatorname{signum}(x^2-1)}\right)}{32\sqrt{\operatorname{signum}(x^2-1)}}$

input `int(x*ln(x)/(x^2-1)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/4/signum(x^2-1)^(1/2)*(-signum(x^2-1))^(1/2)*(2-2*(-x^2+1)^(1/2))+1/2/signum(x^2-1)^(1/2)*(-signum(x^2-1))^(1/2)*ln(x)*(2-2*(-x^2+1)^(1/2))+1/32/signum(x^2-1)^(1/2)*(-signum(x^2-1))^(1/2)*(-16+16*(-x^2+1)^(1/2)-32*ln(1/2+1/2*(-x^2+1)^(1/2)))`

3.323.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.79

$$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx = \sqrt{x^2-1}(\log(x)-1) + 2 \arctan(-x + \sqrt{x^2-1})$$

input `integrate(x*log(x)/(x^2-1)^(1/2),x, algorithm="fracas")`

output `sqrt(x^2 - 1)*(log(x) - 1) + 2*arctan(-x + sqrt(x^2 - 1))`

3.323.6 Sympy [A] (verification not implemented)

Time = 1.22 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.85

$$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx = \sqrt{x^2-1} \log(x) - \left\{ \sqrt{x^2-1} - \arccos\left(\frac{1}{x}\right) \quad \text{for } x > -1 \wedge x < 1 \right.$$

input `integrate(x*ln(x)/(x**2-1)**(1/2),x)`

output `sqrt(x**2 - 1)*log(x) - Piecewise((sqrt(x**2 - 1) - acos(1/x), (x > -1) & (x < 1)))`

3.323. $\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx$

3.323.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.79

$$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx = \sqrt{x^2-1} \log(x) - \sqrt{x^2-1} - \arcsin\left(\frac{1}{|x|}\right)$$

input `integrate(x*log(x)/(x^2-1)^(1/2),x, algorithm="maxima")`output `sqrt(x^2 - 1)*log(x) - sqrt(x^2 - 1) - arcsin(1/abs(x))`**3.323.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.82

$$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx = \sqrt{x^2-1} \log(x) - \sqrt{x^2-1} + \arctan(\sqrt{x^2-1})$$

input `integrate(x*log(x)/(x^2-1)^(1/2),x, algorithm="giac")`output `sqrt(x^2 - 1)*log(x) - sqrt(x^2 - 1) + arctan(sqrt(x^2 - 1))`**3.323.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx = \int \frac{x \ln(x)}{\sqrt{x^2-1}} dx$$

input `int((x*log(x))/(x^2 - 1)^(1/2),x)`output `int((x*log(x))/(x^2 - 1)^(1/2), x)`

3.324 $\int \frac{a+x}{a^2+x^2} dx$

3.324.1 Optimal result	1636
3.324.2 Mathematica [A] (verified)	1636
3.324.3 Rubi [A] (verified)	1637
3.324.4 Maple [A] (verified)	1638
3.324.5 Fricas [A] (verification not implemented)	1638
3.324.6 Sympy [C] (verification not implemented)	1638
3.324.7 Maxima [A] (verification not implemented)	1639
3.324.8 Giac [A] (verification not implemented)	1639
3.324.9 Mupad [B] (verification not implemented)	1639

3.324.1 Optimal result

Integrand size = 13, antiderivative size = 19

$$\int \frac{a+x}{a^2+x^2} dx = \arctan\left(\frac{x}{a}\right) + \frac{1}{2} \log(a^2+x^2)$$

output `arctan(x/a)+1/2*ln(a^2+x^2)`

3.324.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{a+x}{a^2+x^2} dx = \arctan\left(\frac{x}{a}\right) + \frac{1}{2} \log(a^2+x^2)$$

input `Integrate[(a + x)/(a^2 + x^2), x]`

output `ArcTan[x/a] + Log[a^2 + x^2]/2`

3.324.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {452, 216, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a+x}{a^2+x^2} dx \\ & \quad \downarrow \text{452} \\ & a \int \frac{1}{a^2+x^2} dx + \int \frac{x}{a^2+x^2} dx \\ & \quad \downarrow \text{216} \\ & \int \frac{x}{a^2+x^2} dx + \arctan\left(\frac{x}{a}\right) \\ & \quad \downarrow \text{240} \\ & \frac{1}{2} \log(a^2+x^2) + \arctan\left(\frac{x}{a}\right) \end{aligned}$$

input `Int[(a + x)/(a^2 + x^2),x]`

output `ArcTan[x/a] + Log[a^2 + x^2]/2`

3.324.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 240 `Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`

rule 452 `Int[((c_) + (d_.)*(x_))/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[c Int[1/(a + b*x^2), x], x] + Simp[d Int[x/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c^2 + a*d^2, 0]`

3.324.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

method	result	size
default	$\arctan\left(\frac{x}{a}\right) + \frac{\ln(a^2+x^2)}{2}$	18
risch	$\arctan\left(\frac{x}{a}\right) + \frac{\ln(a^2+x^2)}{2}$	18
parallelrisch	$\frac{\ln(-ia+x)}{2} - \frac{i \ln(-ia+x)}{2} + \frac{\ln(ia+x)}{2} + \frac{i \ln(ia+x)}{2}$	40

input `int((a+x)/(a^2+x^2),x,method=_RETURNVERBOSE)`output `arctan(x/a)+1/2*ln(a^2+x^2)`**3.324.5 Fricas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{a+x}{a^2+x^2} dx = \arctan\left(\frac{x}{a}\right) + \frac{1}{2} \log(a^2+x^2)$$

input `integrate((a+x)/(a^2+x^2),x, algorithm="fricas")`output `arctan(x/a) + 1/2*log(a^2 + x^2)`**3.324.6 Sympy [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.04 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.21

$$\int \frac{a+x}{a^2+x^2} dx = \left(\frac{1}{2} - \frac{i}{2}\right) \log\left(-a + 2a\left(\frac{1}{2} - \frac{i}{2}\right) + x\right) + \left(\frac{1}{2} + \frac{i}{2}\right) \log\left(-a + 2a\left(\frac{1}{2} + \frac{i}{2}\right) + x\right)$$

input `integrate((a+x)/(a**2+x**2),x)`

output $(1/2 - I/2)*\log(-a + 2*a*(1/2 - I/2) + x) + (1/2 + I/2)*\log(-a + 2*a*(1/2 + I/2) + x)$

3.324.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{a+x}{a^2+x^2} dx = \arctan\left(\frac{x}{a}\right) + \frac{1}{2} \log(a^2+x^2)$$

input `integrate((a+x)/(a^2+x^2),x, algorithm="maxima")`

output `arctan(x/a) + 1/2*log(a^2 + x^2)`

3.324.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{a+x}{a^2+x^2} dx = \arctan\left(\frac{x}{a}\right) + \frac{1}{2} \log(a^2+x^2)$$

input `integrate((a+x)/(a^2+x^2),x, algorithm="giac")`

output `arctan(x/a) + 1/2*log(a^2 + x^2)`

3.324.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{a+x}{a^2+x^2} dx = \frac{\ln(a^2+x^2)}{2} + \operatorname{atan}\left(\frac{x}{a}\right)$$

input `int((a + x)/(a^2 + x^2),x)`

output `log(a^2 + x^2)/2 + atan(x/a)`

3.325 $\int \sqrt{1+x-x^2} dx$

3.325.1 Optimal result	1640
3.325.2 Mathematica [A] (verified)	1640
3.325.3 Rubi [A] (verified)	1641
3.325.4 Maple [A] (verified)	1642
3.325.5 Fricas [A] (verification not implemented)	1642
3.325.6 Sympy [A] (verification not implemented)	1643
3.325.7 Maxima [A] (verification not implemented)	1643
3.325.8 Giac [A] (verification not implemented)	1643
3.325.9 Mupad [B] (verification not implemented)	1644

3.325.1 Optimal result

Integrand size = 12, antiderivative size = 38

$$\int \sqrt{1+x-x^2} dx = -\frac{1}{4}(1-2x)\sqrt{1+x-x^2} - \frac{5}{8} \arcsin\left(\frac{1-2x}{\sqrt{5}}\right)$$

output `-5/8*arcsin(1/5*(1-2*x)*5^(1/2))-1/4*(1-2*x)*(-x^2+x+1)^(1/2)`

3.325.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.18

$$\int \sqrt{1+x-x^2} dx = \frac{1}{4}(-1+2x)\sqrt{1+x-x^2} + \frac{5}{4} \arctan\left(\frac{x}{-1+\sqrt{1+x-x^2}}\right)$$

input `Integrate[Sqrt[1+x-x^2],x]`

output `((-1+2*x)*Sqrt[1+x-x^2])/4+(5*ArcTan[x/(-1+Sqrt[1+x-x^2])])/4`

3.325.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {1087, 1090, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{-x^2 + x + 1} dx$$

$$\downarrow 1087$$

$$\frac{5}{8} \int \frac{1}{\sqrt{-x^2 + x + 1}} dx - \frac{1}{4}(1 - 2x)\sqrt{-x^2 + x + 1}$$

$$\downarrow 1090$$

$$-\frac{1}{8}\sqrt{5} \int \frac{1}{\sqrt{1 - \frac{1}{5}(1 - 2x)^2}} d(1 - 2x) - \frac{1}{4}\sqrt{-x^2 + x + 1}(1 - 2x)$$

$$\downarrow 223$$

$$-\frac{5}{8} \arcsin\left(\frac{1 - 2x}{\sqrt{5}}\right) - \frac{1}{4}\sqrt{-x^2 + x + 1}(1 - 2x)$$

input `Int[Sqrt[1 + x - x^2], x]`

output `-1/4*((1 - 2*x)*Sqrt[1 + x - x^2]) - (5*ArcSin[(1 - 2*x)/Sqrt[5]])/8`

3.325.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 1087 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] - Simp[p*(b^2 - 4*a*c)/(2*c*(2*p + 1)) Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[3*p])`

rule 1090 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

3.325.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.79

method	result	size
default	$-\frac{(1-2x)\sqrt{-x^2+x+1}}{4} + \frac{5 \arcsin\left(\frac{2\sqrt{5}\left(x-\frac{1}{2}\right)}{5}\right)}{8}$	30
risch	$-\frac{(x^2-x-1)(2x-1)}{4\sqrt{-x^2+x+1}} + \frac{5 \arcsin\left(\frac{2\sqrt{5}\left(x-\frac{1}{2}\right)}{5}\right)}{8}$	38
trager	$\left(-\frac{1}{4} + \frac{x}{2}\right) \sqrt{-x^2+x+1} - \frac{5 \operatorname{RootOf}\left(_Z^2+1\right) \ln\left(2 \operatorname{RootOf}\left(_Z^2+1\right) x - \operatorname{RootOf}\left(_Z^2+1\right) + 2\sqrt{-x^2+x+1}\right)}{8}$	57

input `int((-x^2+x+1)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/4*(1-2*x)*(-x^2+x+1)^(1/2)+5/8*arcsin(2/5*5^(1/2)*(x-1/2))`

3.325.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.97

$$\int \sqrt{1+x-x^2} dx = \frac{1}{4} \sqrt{-x^2+x+1}(2x-1) - \frac{5}{4} \arctan\left(\frac{\sqrt{-x^2+x+1}-1}{x}\right)$$

input `integrate((-x^2+x+1)^(1/2),x, algorithm="fricas")`

output `1/4*sqrt(-x^2 + x + 1)*(2*x - 1) - 5/4*arctan((sqrt(-x^2 + x + 1) - 1)/x)`

3.325.6 Sympy [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.89

$$\int \sqrt{1+x-x^2} dx = \left(\frac{x}{2} - \frac{1}{4}\right) \sqrt{-x^2+x+1} + \frac{5 \operatorname{asin}\left(\frac{2\sqrt{5}(x-\frac{1}{2})}{5}\right)}{8}$$

input `integrate((-x**2+x+1)**(1/2),x)`output `(x/2 - 1/4)*sqrt(-x**2 + x + 1) + 5*asin(2*sqrt(5)*(x - 1/2)/5)/8`**3.325.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.03

$$\int \sqrt{1+x-x^2} dx = \frac{1}{2} \sqrt{-x^2+x+1} x - \frac{1}{4} \sqrt{-x^2+x+1} - \frac{5}{8} \arcsin\left(-\frac{1}{5} \sqrt{5}(2x-1)\right)$$

input `integrate((-x^2+x+1)^(1/2),x, algorithm="maxima")`output `1/2*sqrt(-x^2 + x + 1)*x - 1/4*sqrt(-x^2 + x + 1) - 5/8*arcsin(-1/5*sqrt(5)*(2*x - 1))`**3.325.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int \sqrt{1+x-x^2} dx = \frac{1}{4} \sqrt{-x^2+x+1}(2x-1) + \frac{5}{8} \arcsin\left(\frac{1}{5} \sqrt{5}(2x-1)\right)$$

input `integrate((-x^2+x+1)^(1/2),x, algorithm="giac")`output `1/4*sqrt(-x^2 + x + 1)*(2*x - 1) + 5/8*arcsin(1/5*sqrt(5)*(2*x - 1))`

3.325.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.74

$$\int \sqrt{1+x-x^2} dx = \frac{5 \operatorname{asin}\left(\frac{2\sqrt{5}\left(x-\frac{1}{2}\right)}{5}\right)}{8} + \left(\frac{x}{2} - \frac{1}{4}\right) \sqrt{-x^2+x+1}$$

input `int((x - x^2 + 1)^(1/2),x)`

output `(5*asin((2*5^(1/2)*(x - 1/2))/5))/8 + (x/2 - 1/4)*(x - x^2 + 1)^(1/2)`

3.326 $\int \frac{x^4}{16+x^{10}} dx$

3.326.1 Optimal result	1645
3.326.2 Mathematica [A] (verified)	1645
3.326.3 Rubi [A] (verified)	1646
3.326.4 Maple [A] (verified)	1647
3.326.5 Fricas [A] (verification not implemented)	1647
3.326.6 Sympy [A] (verification not implemented)	1647
3.326.7 Maxima [A] (verification not implemented)	1648
3.326.8 Giac [A] (verification not implemented)	1648
3.326.9 Mupad [B] (verification not implemented)	1648

3.326.1 Optimal result

Integrand size = 11, antiderivative size = 12

$$\int \frac{x^4}{16+x^{10}} dx = \frac{1}{20} \arctan\left(\frac{x^5}{4}\right)$$

output `1/20*arctan(1/4*x^5)`

3.326.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{x^4}{16+x^{10}} dx = \frac{1}{20} \arctan\left(\frac{x^5}{4}\right)$$

input `Integrate[x^4/(16 + x^10),x]`

output `ArcTan[x^5/4]/20`

3.326.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {807, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4}{x^{10} + 16} dx$$

↓ 807

$$\frac{1}{5} \int \frac{1}{x^{10} + 16} dx^5$$

↓ 216

$$\frac{1}{20} \arctan\left(\frac{x^5}{4}\right)$$

input `Int[x^4/(16 + x^10),x]`

output `ArcTan[x^5/4]/20`

3.326.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

3.326.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

method	result	size
default	$\frac{\arctan\left(\frac{x^5}{4}\right)}{20}$	9
meijerg	$\frac{\arctan\left(\frac{x^5}{4}\right)}{20}$	9
risch	$\frac{\arctan\left(\frac{x^5}{4}\right)}{20}$	9
parallelrisch	$\frac{i \ln(x^5+4i)}{40} - \frac{i \ln(x^5-4i)}{40}$	22

input `int(x^4/(x^10+16),x,method=_RETURNVERBOSE)`output `1/20*arctan(1/4*x^5)`**3.326.5 Fricas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x^4}{16 + x^{10}} dx = \frac{1}{20} \arctan\left(\frac{1}{4} x^5\right)$$

input `integrate(x^4/(x^10+16),x, algorithm="fricas")`output `1/20*arctan(1/4*x^5)`**3.326.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.58

$$\int \frac{x^4}{16 + x^{10}} dx = \frac{\operatorname{atan}\left(\frac{x^5}{4}\right)}{20}$$

input `integrate(x**4/(x**10+16),x)`output `atan(x**5/4)/20`

3.326.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x^4}{16 + x^{10}} dx = \frac{1}{20} \arctan\left(\frac{1}{4} x^5\right)$$

input `integrate(x^4/(x^10+16),x, algorithm="maxima")`output `1/20*arctan(1/4*x^5)`**3.326.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x^4}{16 + x^{10}} dx = \frac{1}{20} \arctan\left(\frac{1}{4} x^5\right)$$

input `integrate(x^4/(x^10+16),x, algorithm="giac")`output `1/20*arctan(1/4*x^5)`**3.326.9 Mupad [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x^4}{16 + x^{10}} dx = \frac{\operatorname{atan}\left(\frac{x^5}{4}\right)}{20}$$

input `int(x^4/(x^10 + 16),x)`output `atan(x^5/4)/20`

3.327 $\int \frac{2+x}{2+x+x^2} dx$

3.327.1 Optimal result	1649
3.327.2 Mathematica [A] (verified)	1649
3.327.3 Rubi [A] (verified)	1650
3.327.4 Maple [A] (verified)	1651
3.327.5 Fricas [A] (verification not implemented)	1651
3.327.6 Sympy [A] (verification not implemented)	1652
3.327.7 Maxima [A] (verification not implemented)	1652
3.327.8 Giac [A] (verification not implemented)	1652
3.327.9 Mupad [B] (verification not implemented)	1653

3.327.1 Optimal result

Integrand size = 12, antiderivative size = 31

$$\int \frac{2+x}{2+x+x^2} dx = \frac{3 \arctan\left(\frac{1+2x}{\sqrt{7}}\right)}{\sqrt{7}} + \frac{1}{2} \log(2+x+x^2)$$

output `1/2*ln(x^2+x+2)+3/7*arctan(1/7*(1+2*x)*7^(1/2))*7^(1/2)`

3.327.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{2+x}{2+x+x^2} dx = \frac{3 \arctan\left(\frac{1+2x}{\sqrt{7}}\right)}{\sqrt{7}} + \frac{1}{2} \log(2+x+x^2)$$

input `Integrate[(2 + x)/(2 + x + x^2), x]`

output `(3*ArcTan[(1 + 2*x)/Sqrt[7]])/Sqrt[7] + Log[2 + x + x^2]/2`

3.327.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {1142, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x+2}{x^2+x+2} dx \\
 & \quad \downarrow \text{1142} \\
 & \frac{3}{2} \int \frac{1}{x^2+x+2} dx + \frac{1}{2} \int \frac{2x+1}{x^2+x+2} dx \\
 & \quad \downarrow \text{1083} \\
 & \frac{1}{2} \int \frac{2x+1}{x^2+x+2} dx - 3 \int \frac{1}{-(2x+1)^2-7} d(2x+1) \\
 & \quad \downarrow \text{217} \\
 & \frac{1}{2} \int \frac{2x+1}{x^2+x+2} dx + \frac{3 \arctan\left(\frac{2x+1}{\sqrt{7}}\right)}{\sqrt{7}} \\
 & \quad \downarrow \text{1103} \\
 & \frac{3 \arctan\left(\frac{2x+1}{\sqrt{7}}\right)}{\sqrt{7}} + \frac{1}{2} \log(x^2+x+2)
 \end{aligned}$$

input `Int[(2 + x)/(2 + x + x^2),x]`

output `(3*ArcTan[(1 + 2*x)/Sqrt[7]])/Sqrt[7] + Log[2 + x + x^2]/2`

3.327.3.1 Defintions of rubi rules used

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & (LtQ[a, 0] || LtQ[b, 0])`

rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

3.327.4 Maple [A] (verified)

Time = 0.71 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

method	result	size
default	$\frac{\ln(x^2+x+2)}{2} + \frac{3 \arctan\left(\frac{(1+2x)\sqrt{7}}{7}\right)\sqrt{7}}{7}$	27
risch	$\frac{\ln(4x^2+4x+8)}{2} + \frac{3 \arctan\left(\frac{(1+2x)\sqrt{7}}{7}\right)\sqrt{7}}{7}$	31

input `int((2+x)/(x^2+x+2),x,method=_RETURNVERBOSE)`

output `1/2*ln(x^2+x+2)+3/7*arctan(1/7*(1+2*x)*7^(1/2))*7^(1/2)`

3.327.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{2+x}{2+x+x^2} dx = \frac{3}{7} \sqrt{7} \arctan\left(\frac{1}{7} \sqrt{7}(2x+1)\right) + \frac{1}{2} \log(x^2+x+2)$$

input `integrate((2+x)/(x^2+x+2),x, algorithm="fracas")`

output `3/7*sqrt(7)*arctan(1/7*sqrt(7)*(2*x + 1)) + 1/2*log(x^2 + x + 2)`

3.327.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.16

$$\int \frac{2+x}{2+x+x^2} dx = \frac{\log(x^2+x+2)}{2} + \frac{3\sqrt{7} \operatorname{atan}\left(\frac{2\sqrt{7}x}{7} + \frac{\sqrt{7}}{7}\right)}{7}$$

input `integrate((2+x)/(x**2+x+2),x)`output `log(x**2 + x + 2)/2 + 3*sqrt(7)*atan(2*sqrt(7)*x/7 + sqrt(7)/7)/7`**3.327.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{2+x}{2+x+x^2} dx = \frac{3}{7} \sqrt{7} \arctan\left(\frac{1}{7} \sqrt{7}(2x+1)\right) + \frac{1}{2} \log(x^2+x+2)$$

input `integrate((2+x)/(x^2+x+2),x, algorithm="maxima")`output `3/7*sqrt(7)*arctan(1/7*sqrt(7)*(2*x + 1)) + 1/2*log(x^2 + x + 2)`**3.327.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{2+x}{2+x+x^2} dx = \frac{3}{7} \sqrt{7} \arctan\left(\frac{1}{7} \sqrt{7}(2x+1)\right) + \frac{1}{2} \log(x^2+x+2)$$

input `integrate((2+x)/(x^2+x+2),x, algorithm="giac")`output `3/7*sqrt(7)*arctan(1/7*sqrt(7)*(2*x + 1)) + 1/2*log(x^2 + x + 2)`

3.327.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

$$\int \frac{2+x}{2+x+x^2} dx = \frac{\ln(x^2+x+2)}{2} + \frac{3\sqrt{7}\operatorname{atan}\left(\frac{2\sqrt{7}x}{7} + \frac{\sqrt{7}}{7}\right)}{7}$$

input `int((x + 2)/(x + x^2 + 2),x)`output `log(x + x^2 + 2)/2 + (3*7^(1/2)*atan((2*7^(1/2)*x)/7 + 7^(1/2)/7))/7`

3.328 $\int x \sec(x) \tan(x) dx$

3.328.1 Optimal result	1654
3.328.2 Mathematica [B] (verified)	1654
3.328.3 Rubi [A] (verified)	1655
3.328.4 Maple [A] (verified)	1656
3.328.5 Fricas [B] (verification not implemented)	1656
3.328.6 Sympy [A] (verification not implemented)	1656
3.328.7 Maxima [B] (verification not implemented)	1657
3.328.8 Giac [B] (verification not implemented)	1657
3.328.9 Mupad [B] (verification not implemented)	1658

3.328.1 Optimal result

Integrand size = 6, antiderivative size = 10

$$\int x \sec(x) \tan(x) dx = -\operatorname{arctanh}(\sin(x)) + x \sec(x)$$

output `-arctanh(sin(x))+x*sec(x)`

3.328.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 37 vs. $2(10) = 20$.

Time = 0.01 (sec) , antiderivative size = 37, normalized size of antiderivative = 3.70

$$\int x \sec(x) \tan(x) dx = \log\left(\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)\right) - \log\left(\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)\right) + x \sec(x)$$

input `Integrate[x*Sec[x]*Tan[x],x]`

output `Log[Cos[x/2] - Sin[x/2]] - Log[Cos[x/2] + Sin[x/2]] + x*Sec[x]`

3.328.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {4244, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \tan(x) \sec(x) dx \\ & \quad \downarrow \text{4244} \\ & x \sec(x) - \int \sec(x) dx \\ & \quad \downarrow \text{3042} \\ & x \sec(x) - \int \csc\left(x + \frac{\pi}{2}\right) dx \\ & \quad \downarrow \text{4257} \\ & x \sec(x) - \operatorname{arctanh}(\sin(x)) \end{aligned}$$

input `Int[x*Sec[x]*Tan[x],x]`

output `-ArcTanh[Sin[x]] + x*Sec[x]`

3.328.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4244 `Int[(x_)^(m_.)*Sec[(a_.) + (b_.)*(x_)^(n_.)]^(p_.)*Tan[(a_.) + (b_.)*(x_)^(n_.)]^(q_.), x_Symbol] := Simp[x^(m - n + 1)*(Sec[a + b*x^n]^p/(b*n*p)), x] - Simp[(m - n + 1)/(b*n*p) Int[x^(m - n)*Sec[a + b*x^n]^p, x], x] /; FreeQ[{a, b, p}, x] && IntegerQ[n] && GeQ[m, n] && EqQ[q, 1]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.328.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.60

method	result	size
default	$\frac{x}{\cos(x)} - \ln(\sec(x) + \tan(x))$	16
risch	$\frac{2e^{ix}x}{e^{2ix}+1} + \ln(e^{ix} - i) - \ln(i + e^{ix})$	39

input `int(x*sec(x)*tan(x),x,method=_RETURNVERBOSE)`

output `x/cos(x)-ln(sec(x)+tan(x))`

3.328.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(10) = 20$.

Time = 0.26 (sec) , antiderivative size = 29, normalized size of antiderivative = 2.90

$$\int x \sec(x) \tan(x) dx = -\frac{\cos(x) \log(\sin(x) + 1) - \cos(x) \log(-\sin(x) + 1) - 2x}{2 \cos(x)}$$

input `integrate(x*sec(x)*tan(x),x, algorithm="fracas")`

output `-1/2*(cos(x)*log(sin(x) + 1) - cos(x)*log(-sin(x) + 1) - 2*x)/cos(x)`

3.328.6 Sympy [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x \sec(x) \tan(x) dx = x \sec(x) - \log(\tan(x) + \sec(x))$$

input `integrate(x*sec(x)*tan(x),x)`

output `x*sec(x) - log(tan(x) + sec(x))`

3.328.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. $2(10) = 20$.

Time = 0.28 (sec) , antiderivative size = 121, normalized size of antiderivative = 12.10

$$\int x \sec(x) \tan(x) dx$$

$$= \frac{4x \cos(2x) \cos(x) + 4x \sin(2x) \sin(x) + 4x \cos(x) - (\cos(2x)^2 + \sin(2x)^2 + 2 \cos(2x) + 1) \log(\cos(x)^2 + \sin(x)^2 + 2 \sin(x) + 1) + (\cos(2x)^2 + \sin(2x)^2 + 2 \cos(2x) + 1) \log(\cos(x)^2 + \sin(x)^2 - 2 \sin(x) + 1)}{2(\cos(2x)^2 + \sin(2x)^2 + 2 \cos(2x) + 1)}$$

input `integrate(x*sec(x)*tan(x),x, algorithm="maxima")`

output `1/2*(4*x*cos(2*x)*cos(x) + 4*x*sin(2*x)*sin(x) + 4*x*cos(x) - (cos(2*x)^2 + sin(2*x)^2 + 2*cos(2*x) + 1)*log(cos(x)^2 + sin(x)^2 + 2*sin(x) + 1) + (cos(2*x)^2 + sin(2*x)^2 + 2*cos(2*x) + 1)*log(cos(x)^2 + sin(x)^2 - 2*sin(x) + 1))/(cos(2*x)^2 + sin(2*x)^2 + 2*cos(2*x) + 1)`

3.328.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 150 vs. $2(10) = 20$.

Time = 0.36 (sec) , antiderivative size = 150, normalized size of antiderivative = 15.00

$$\int x \sec(x) \tan(x) dx =$$

$$\frac{2x \tan\left(\frac{1}{2}x\right)^2 + \log\left(\frac{2\left(\tan\left(\frac{1}{2}x\right)^2 + 2 \tan\left(\frac{1}{2}x\right) + 1\right)}{\tan\left(\frac{1}{2}x\right)^2 + 1}\right) \tan\left(\frac{1}{2}x\right)^2 - \log\left(\frac{2\left(\tan\left(\frac{1}{2}x\right)^2 - 2 \tan\left(\frac{1}{2}x\right) + 1\right)}{\tan\left(\frac{1}{2}x\right)^2 + 1}\right) \tan\left(\frac{1}{2}x\right)^2 + 2}{2\left(\tan\left(\frac{1}{2}x\right)^2 - 1\right)}$$

input `integrate(x*sec(x)*tan(x),x, algorithm="giac")`

output `-1/2*(2*x*tan(1/2*x)^2 + log(2*(tan(1/2*x)^2 + 2*tan(1/2*x) + 1)/(tan(1/2*x)^2 + 1))*tan(1/2*x)^2 - log(2*(tan(1/2*x)^2 - 2*tan(1/2*x) + 1)/(tan(1/2*x)^2 + 1))*tan(1/2*x)^2 + 2*x - log(2*(tan(1/2*x)^2 + 2*tan(1/2*x) + 1)/(tan(1/2*x)^2 + 1)) + log(2*(tan(1/2*x)^2 - 2*tan(1/2*x) + 1)/(tan(1/2*x)^2 + 1)))/(tan(1/2*x)^2 - 1)`

3.328.9 Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.90

$$\int x \sec(x) \tan(x) dx = \frac{x}{\cos(x)} + \operatorname{atan}(\cos(x) + \sin(x) 1i) 2i$$

input `int((x*tan(x))/cos(x),x)`

output `atan(cos(x) + sin(x)*1i)*2i + x/cos(x)`

3.329 $\int \frac{x}{-a^4+x^4} dx$

3.329.1 Optimal result	1659
3.329.2 Mathematica [A] (verified)	1659
3.329.3 Rubi [A] (verified)	1660
3.329.4 Maple [A] (verified)	1661
3.329.5 Fricas [A] (verification not implemented)	1661
3.329.6 Sympy [A] (verification not implemented)	1661
3.329.7 Maxima [B] (verification not implemented)	1662
3.329.8 Giac [B] (verification not implemented)	1662
3.329.9 Mupad [B] (verification not implemented)	1662

3.329.1 Optimal result

Integrand size = 13, antiderivative size = 15

$$\int \frac{x}{-a^4+x^4} dx = -\frac{\operatorname{arctanh}\left(\frac{x^2}{a^2}\right)}{2a^2}$$

output `-1/2*arctanh(x^2/a^2)/a^2`

3.329.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{x}{-a^4+x^4} dx = -\frac{\operatorname{arctanh}\left(\frac{x^2}{a^2}\right)}{2a^2}$$

input `Integrate[x/(-a^4 + x^4),x]`

output `-1/2*ArcTanh[x^2/a^2]/a^2`

3.329.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {807, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{x^4 - a^4} dx$$

↓ 807

$$\frac{1}{2} \int \frac{1}{x^4 - a^4} dx^2$$

↓ 220

$$-\frac{\operatorname{arctanh}\left(\frac{x^2}{a^2}\right)}{2a^2}$$

input `Int[x/(-a^4 + x^4),x]`

output `-1/2*ArcTanh[x^2/a^2]/a^2`

3.329.3.1 Defintions of rubi rules used

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

3.329.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.80

method	result	size
parallelrisc	$\frac{\ln(-a+x)+\ln(a+x)-\ln(a^2+x^2)}{4a^2}$	27
default	$\frac{\ln(a^2-x^2)}{4a^2} - \frac{\ln(a^2+x^2)}{4a^2}$	30
risc	$\frac{\ln(-a^2+x^2)}{4a^2} - \frac{\ln(a^2+x^2)}{4a^2}$	30
norman	$\frac{\ln(a-x)}{4a^2} + \frac{\ln(a+x)}{4a^2} - \frac{\ln(a^2+x^2)}{4a^2}$	35

input `int(x/(-a^4+x^4),x,method=_RETURNVERBOSE)`output `1/4*(ln(-a+x)+ln(a+x)-ln(a^2+x^2))/a^2`**3.329.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.73

$$\int \frac{x}{-a^4 + x^4} dx = -\frac{\log(a^2 + x^2) - \log(-a^2 + x^2)}{4a^2}$$

input `integrate(x/(-a^4+x^4),x, algorithm="fricas")`output `-1/4*(log(a^2 + x^2) - log(-a^2 + x^2))/a^2`**3.329.6 Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.47

$$\int \frac{x}{-a^4 + x^4} dx = \frac{\frac{\log(-a^2+x^2)}{4} - \frac{\log(a^2+x^2)}{4}}{a^2}$$

input `integrate(x/(-a**4+x**4),x)`output `(log(-a**2 + x**2)/4 - log(a**2 + x**2)/4)/a**2`

3.329.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(13) = 26$.

Time = 0.19 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.93

$$\int \frac{x}{-a^4 + x^4} dx = -\frac{\log(a^2 + x^2)}{4a^2} + \frac{\log(-a^2 + x^2)}{4a^2}$$

input `integrate(x/(-a^4+x^4),x, algorithm="maxima")`

output `-1/4*log(a^2 + x^2)/a^2 + 1/4*log(-a^2 + x^2)/a^2`

3.329.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 30 vs. $2(13) = 26$.

Time = 0.29 (sec) , antiderivative size = 30, normalized size of antiderivative = 2.00

$$\int \frac{x}{-a^4 + x^4} dx = -\frac{\log(a^2 + x^2)}{4a^2} + \frac{\log(|-a^2 + x^2|)}{4a^2}$$

input `integrate(x/(-a^4+x^4),x, algorithm="giac")`

output `-1/4*log(a^2 + x^2)/a^2 + 1/4*log(abs(-a^2 + x^2))/a^2`

3.329.9 Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{x}{-a^4 + x^4} dx = -\frac{\operatorname{atanh}\left(\frac{x^2}{a^2}\right)}{2a^2}$$

input `int(-x/(a^4 - x^4),x)`

output `-atanh(x^2/a^2)/(2*a^2)`

$$\mathbf{3.330} \quad \int \frac{1}{\sqrt{x} + \sqrt{1+x}} dx$$

3.330.1 Optimal result	1663
3.330.2 Mathematica [A] (verified)	1663
3.330.3 Rubi [A] (verified)	1664
3.330.4 Maple [A] (verified)	1665
3.330.5 Fricas [A] (verification not implemented)	1665
3.330.6 Sympy [B] (verification not implemented)	1666
3.330.7 Maxima [F]	1666
3.330.8 Giac [A] (verification not implemented)	1666
3.330.9 Mupad [B] (verification not implemented)	1667

3.330.1 Optimal result

Integrand size = 15, antiderivative size = 21

$$\int \frac{1}{\sqrt{x} + \sqrt{1+x}} dx = -\frac{2x^{3/2}}{3} + \frac{2}{3}(1+x)^{3/2}$$

output `-2/3*x^(3/2)+2/3*(1+x)^(3/2)`

3.330.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{x} + \sqrt{1+x}} dx = -\frac{2x^{3/2}}{3} + \frac{2}{3}(1+x)^{3/2}$$

input `Integrate[(Sqrt[x] + Sqrt[1 + x])^(-1), x]`

output `(-2*x^(3/2))/3 + (2*(1 + x)^(3/2))/3`

3.330.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2531, 15, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt{x} + \sqrt{x+1}} dx \\ & \quad \downarrow \text{2531} \\ & \int \sqrt{x+1} dx - \int \sqrt{x} dx \\ & \quad \downarrow \text{15} \\ & \int \sqrt{x+1} dx - \frac{2x^{3/2}}{3} \\ & \quad \downarrow \text{17} \\ & \frac{2}{3}(x+1)^{3/2} - \frac{2x^{3/2}}{3} \end{aligned}$$

input `Int[(Sqrt[x] + Sqrt[1 + x])^(-1),x]`

output `(-2*x^(3/2))/3 + (2*(1 + x)^(3/2))/3`

3.330.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

```
rule 2531 Int[(u_)/((d_)*(x_)^(n_) + (c_)*Sqrt[(a_) + (b_)*(x_)^(p_)]), x_Symbol]
  := Simp[-b/(a*d) Int[u*x^n, x], x] + Simp[1/(a*c) Int[u*Sqrt[a + b*x^(2*n)], x], x]
  /; FreeQ[{a, b, c, d, n}, x] && EqQ[p, 2*n] && EqQ[b*c^2 - d^2, 0]
```

3.330.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

method	result	size
default	$-\frac{2x^{\frac{3}{2}}}{3} + \frac{2(1+x)^{\frac{3}{2}}}{3}$	14
meijerg	$-\frac{\frac{4\sqrt{\pi}x^{\frac{3}{2}}}{3} - \frac{2\sqrt{\pi}x^{\frac{3}{2}}(2+\frac{2}{x})\sqrt{1+\frac{1}{x}}}{3}}{2\sqrt{\pi}}$	37

```
input int(1/(x^(1/2)+(1+x)^(1/2)),x,method=_RETURNVERBOSE)
```

```
output -2/3*x^(3/2)+2/3*(1+x)^(3/2)
```

3.330.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \frac{1}{\sqrt{x} + \sqrt{1+x}} dx = \frac{2}{3}(x+1)^{\frac{3}{2}} - \frac{2}{3}x^{\frac{3}{2}}$$

```
input integrate(1/(x^(1/2)+(1+x)^(1/2)),x, algorithm="fracas")
```

```
output 2/3*(x + 1)^(3/2) - 2/3*x^(3/2)
```

3.330.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 63 vs. $2(17) = 34$.

Time = 0.16 (sec) , antiderivative size = 63, normalized size of antiderivative = 3.00

$$\int \frac{1}{\sqrt{x} + \sqrt{1+x}} dx = \frac{2\sqrt{x}\sqrt{x+1}}{3\sqrt{x} + 3\sqrt{x+1}} + \frac{4x}{3\sqrt{x} + 3\sqrt{x+1}} + \frac{2}{3\sqrt{x} + 3\sqrt{x+1}}$$

input `integrate(1/(x**(1/2)+(1+x)**(1/2)),x)`

output `2*sqrt(x)*sqrt(x + 1)/(3*sqrt(x) + 3*sqrt(x + 1)) + 4*x/(3*sqrt(x) + 3*sqrt(x + 1)) + 2/(3*sqrt(x) + 3*sqrt(x + 1))`

3.330.7 Maxima [F]

$$\int \frac{1}{\sqrt{x} + \sqrt{1+x}} dx = \int \frac{1}{\sqrt{x+1} + \sqrt{x}} dx$$

input `integrate(1/(x^(1/2)+(1+x)^(1/2)),x, algorithm="maxima")`

output `integrate(1/(sqrt(x + 1) + sqrt(x)), x)`

3.330.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \frac{1}{\sqrt{x} + \sqrt{1+x}} dx = \frac{2}{3}(x+1)^{\frac{3}{2}} - \frac{2}{3}x^{\frac{3}{2}}$$

input `integrate(1/(x^(1/2)+(1+x)^(1/2)),x, algorithm="giac")`

output `2/3*(x + 1)^(3/2) - 2/3*x^(3/2)`

3.330.9 Mupad [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{x} + \sqrt{1+x}} dx = \frac{2x\sqrt{x+1}}{3} + \frac{2\sqrt{x+1}}{3} - \frac{2x^{3/2}}{3}$$

input `int(1/((x + 1)^(1/2) + x^(1/2)),x)`output `(2*x*(x + 1)^(1/2))/3 + (2*(x + 1)^(1/2))/3 - (2*x^(3/2))/3`

3.331 $\int \frac{1}{1-e^{-x}+2e^x} dx$

3.331.1 Optimal result	1668
3.331.2 Mathematica [A] (verified)	1668
3.331.3 Rubi [A] (verified)	1669
3.331.4 Maple [A] (verified)	1670
3.331.5 Fricas [A] (verification not implemented)	1670
3.331.6 Sympy [A] (verification not implemented)	1671
3.331.7 Maxima [A] (verification not implemented)	1671
3.331.8 Giac [A] (verification not implemented)	1671
3.331.9 Mupad [B] (verification not implemented)	1672

3.331.1 Optimal result

Integrand size = 16, antiderivative size = 23

$$\int \frac{1}{1-e^{-x}+2e^x} dx = \frac{1}{3} \log(1-2e^x) - \frac{1}{3} \log(1+e^x)$$

output `1/3*ln(1-2*exp(x))-1/3*ln(1+exp(x))`

3.331.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int \frac{1}{1-e^{-x}+2e^x} dx = \frac{2}{3} \operatorname{arctanh}\left(\frac{1}{3} - \frac{2e^{-x}}{3}\right)$$

input `Integrate[(1 - E^(-x) + 2*E^x)^(-1), x]`

output `(2*ArcTanh[1/3 - 2/(3*E^x)]) / 3`

3.331.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {2720, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{-e^{-x} + 2e^x + 1} dx \\ & \quad \downarrow \text{2720} \\ & \int \frac{1}{e^x + 2e^{2x} - 1} de^x \\ & \quad \downarrow \text{1081} \\ & 2 \int \left(-\frac{1}{6(1+e^x)} - \frac{1}{3(1-2e^x)} \right) de^x \\ & \quad \downarrow \text{2009} \\ & 2 \left(\frac{1}{6} \log(1-2e^x) - \frac{1}{6} \log(e^x+1) \right) \end{aligned}$$

input `Int[(1 - E^(-x) + 2*E^x)^(-1), x]`

output `2*(Log[1 - 2*E^x]/6 - Log[1 + E^x]/6)`

3.331.3.1 Defintions of rubi rules used

rule 1081 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.331.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.70

method	result	size
risch	$\frac{\ln(-\frac{1}{2}+e^x)}{3} - \frac{\ln(1+e^x)}{3}$	16
parallelrisch	$\frac{\ln(-\frac{1}{2}+e^x)}{3} - \frac{\ln(1+e^x)}{3}$	16
derivativedivides	$\frac{\ln(2e^x-1)}{3} - \frac{\ln(1+e^x)}{3}$	18
default	$\frac{\ln(2e^x-1)}{3} - \frac{\ln(1+e^x)}{3}$	18
norman	$\frac{\ln(2e^x-1)}{3} - \frac{\ln(1+e^x)}{3}$	18

```
input int(1/(1-1/exp(x)+2*exp(x)),x,method=_RETURNVERBOSE)
```

```
output 1/3*ln(-1/2+exp(x))-1/3*ln(1+exp(x))
```

3.331.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{1}{1 - e^{-x} + 2e^x} dx = \frac{1}{3} \log(2e^x - 1) - \frac{1}{3} \log(e^x + 1)$$

```
input integrate(1/(1-1/exp(x)+2*exp(x)),x, algorithm="fricas")
```

```
output 1/3*log(2*e^x - 1) - 1/3*log(e^x + 1)
```

3.331.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{1}{1 - e^{-x} + 2e^x} dx = \frac{\log(-2 + e^{-x})}{3} - \frac{\log(1 + e^{-x})}{3}$$

input `integrate(1/(1-1/exp(x)+2*exp(x)),x)`output `log(-2 + exp(-x))/3 - log(1 + exp(-x))/3`**3.331.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{1}{1 - e^{-x} + 2e^x} dx = -\frac{1}{3} \log(e^{(-x)} + 1) + \frac{1}{3} \log(e^{(-x)} - 2)$$

input `integrate(1/(1-1/exp(x)+2*exp(x)),x, algorithm="maxima")`output `-1/3*log(e^(-x) + 1) + 1/3*log(e^(-x) - 2)`**3.331.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int \frac{1}{1 - e^{-x} + 2e^x} dx = -\frac{1}{3} \log(e^x + 1) + \frac{1}{3} \log(|2e^x - 1|)$$

input `integrate(1/(1-1/exp(x)+2*exp(x)),x, algorithm="giac")`output `-1/3*log(e^x + 1) + 1/3*log(abs(2*e^x - 1))`

3.331.9 Mupad [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{1}{1 - e^{-x} + 2e^x} dx = \frac{\ln(2e^x - 1)}{3} - \frac{\ln(e^x + 1)}{3}$$

input `int(1/(2*exp(x) - exp(-x) + 1),x)`

output `log(2*exp(x) - 1)/3 - log(exp(x) + 1)/3`

$$3.332 \quad \int \frac{\arctan(\sqrt{x})}{\sqrt{x}} dx$$

3.332.1 Optimal result	1673
3.332.2 Mathematica [A] (verified)	1673
3.332.3 Rubi [A] (verified)	1674
3.332.4 Maple [A] (verified)	1675
3.332.5 Fricas [A] (verification not implemented)	1675
3.332.6 Sympy [A] (verification not implemented)	1675
3.332.7 Maxima [A] (verification not implemented)	1676
3.332.8 Giac [A] (verification not implemented)	1676
3.332.9 Mupad [B] (verification not implemented)	1676

3.332.1 Optimal result

Integrand size = 12, antiderivative size = 20

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}} dx = 2\sqrt{x} \arctan(\sqrt{x}) - \log(1+x)$$

output `-ln(1+x)+2*arctan(x^(1/2))*x^(1/2)`

3.332.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}} dx = 2\sqrt{x} \arctan(\sqrt{x}) - \log(1+x)$$

input `Integrate[ArcTan[Sqrt[x]]/Sqrt[x],x]`

output `2*Sqrt[x]*ArcTan[Sqrt[x]] - Log[1 + x]`

3.332.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5361, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}} dx$$

↓ 5361

$$2\sqrt{x} \arctan(\sqrt{x}) - \int \frac{1}{x+1} dx$$

↓ 16

$$2\sqrt{x} \arctan(\sqrt{x}) - \log(x+1)$$

input `Int[ArcTan[Sqrt[x]]/Sqrt[x],x]`

output `2*Sqrt[x]*ArcTan[Sqrt[x]] - Log[1 + x]`

3.332.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 5361 `Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] & & IntegerQ[m])) && NeQ[m, -1]`

3.332.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

method	result	size
derivativedivides	$-\ln(1+x) + 2 \arctan(\sqrt{x}) \sqrt{x}$	17
default	$-\ln(1+x) + 2 \arctan(\sqrt{x}) \sqrt{x}$	17
meijerg	$-\ln(1+x) + 2 \arctan(\sqrt{x}) \sqrt{x}$	17

input `int(arctan(x^(1/2))/x^(1/2),x,method=_RETURNVERBOSE)`output `-ln(1+x)+2*arctan(x^(1/2))*x^(1/2)`**3.332.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}} dx = 2\sqrt{x} \arctan(\sqrt{x}) - \log(x+1)$$

input `integrate(arctan(x^(1/2))/x^(1/2),x, algorithm="fricas")`output `2*sqrt(x)*arctan(sqrt(x)) - log(x + 1)`**3.332.6 Sympy [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}} dx = 2\sqrt{x} \operatorname{atan}(\sqrt{x}) - \log(x+1)$$

input `integrate(atan(x**(1/2))/x**(1/2),x)`output `2*sqrt(x)*atan(sqrt(x)) - log(x + 1)`

3.332.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}} dx = 2\sqrt{x} \arctan(\sqrt{x}) - \log(x+1)$$

input `integrate(arctan(x^(1/2))/x^(1/2),x, algorithm="maxima")`output `2*sqrt(x)*arctan(sqrt(x)) - log(x + 1)`**3.332.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}} dx = 2\sqrt{x} \arctan(\sqrt{x}) - \log(x+1)$$

input `integrate(arctan(x^(1/2))/x^(1/2),x, algorithm="giac")`output `2*sqrt(x)*arctan(sqrt(x)) - log(x + 1)`**3.332.9 Mupad [B] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int \frac{\arctan(\sqrt{x})}{\sqrt{x}} dx = 2\sqrt{x} \operatorname{atan}(\sqrt{x}) - \ln(x+1)$$

input `int(atan(x^(1/2))/x^(1/2),x)`output `2*x^(1/2)*atan(x^(1/2)) - log(x + 1)`

3.333 $\int \frac{\log(1+x)}{x^2} dx$

3.333.1 Optimal result	1677
3.333.2 Mathematica [A] (verified)	1677
3.333.3 Rubi [A] (verified)	1678
3.333.4 Maple [A] (verified)	1679
3.333.5 Fricas [A] (verification not implemented)	1680
3.333.6 Sympy [A] (verification not implemented)	1680
3.333.7 Maxima [A] (verification not implemented)	1680
3.333.8 Giac [A] (verification not implemented)	1681
3.333.9 Mupad [B] (verification not implemented)	1681

3.333.1 Optimal result

Integrand size = 8, antiderivative size = 18

$$\int \frac{\log(1+x)}{x^2} dx = \log(x) - \log(1+x) - \frac{\log(1+x)}{x}$$

output `ln(x)-ln(1+x)-ln(1+x)/x`

3.333.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{\log(1+x)}{x^2} dx = \log(x) - \log(1+x) - \frac{\log(1+x)}{x}$$

input `Integrate[Log[1 + x]/x^2,x]`

output `Log[x] - Log[1 + x] - Log[1 + x]/x`

3.333.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {2842, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\log(x+1)}{x^2} dx \\
 & \quad \downarrow \text{2842} \\
 & \int \frac{1}{x(x+1)} dx - \frac{\log(x+1)}{x} \\
 & \quad \downarrow \text{47} \\
 & \int \frac{1}{x} dx - \int \frac{1}{x+1} dx - \frac{\log(x+1)}{x} \\
 & \quad \downarrow \text{14} \\
 & - \int \frac{1}{x+1} dx + \log(x) - \frac{\log(x+1)}{x} \\
 & \quad \downarrow \text{16} \\
 & \log(x) - \frac{\log(x+1)}{x} - \log(x+1)
 \end{aligned}$$

input `Int[Log[1 + x]/x^2,x]`

output `Log[x] - Log[1 + x] - Log[1 + x]/x`

3.333.3.1 Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

rule 2842 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]*(b_.))*((f_.) + (g_.)*(x_)^(q_.)), x_Symbol] := Simp[(f + g*x)^(q + 1)*((a + b*Log[c*(d + e*x)^n])/(g*(q + 1))), x] - Simp[b*e*(n/(g*(q + 1))) Int[(f + g*x)^(q + 1)/(d + e*x), x], x] /; FreeQ[{a, b, c, d, e, f, g, n, q}, x] && NeQ[e*f - d*g, 0] && NeQ[q, -1]`

3.333.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

method	result	size
derivativedivides	$\ln(x) - \frac{\ln(1+x)(1+x)}{x}$	16
default	$\ln(x) - \frac{\ln(1+x)(1+x)}{x}$	16
meijerg	$\ln(x) - \frac{(2x+2)\ln(1+x)}{2x}$	18
risch	$\ln(x) - \ln(1+x) - \frac{\ln(1+x)}{x}$	19
parts	$\ln(x) - \ln(1+x) - \frac{\ln(1+x)}{x}$	19
norman	$\frac{-\ln(1+x)x - \ln(1+x)}{x} + \ln(x)$	22
parallelrisch	$\frac{x \ln(x) - \ln(1+x)x - \ln(1+x)}{x}$	23

input `int(1/x^2*ln(1+x),x,method=_RETURNVERBOSE)`

output `ln(x)-ln(1+x)*(1+x)/x`

3.333.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.06

$$\int \frac{\log(1+x)}{x^2} dx = -\frac{(x+1)\log(x+1) - x\log(x)}{x}$$

input `integrate(log(1+x)/x^2,x, algorithm="fricas")`output `-((x + 1)*log(x + 1) - x*log(x))/x`**3.333.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{\log(1+x)}{x^2} dx = \log(x) - \log(x+1) - \frac{\log(x+1)}{x}$$

input `integrate(ln(1+x)/x**2,x)`output `log(x) - log(x + 1) - log(x + 1)/x`**3.333.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{\log(1+x)}{x^2} dx = -\frac{\log(x+1)}{x} - \log(x+1) + \log(x)$$

input `integrate(log(1+x)/x^2,x, algorithm="maxima")`output `-log(x + 1)/x - log(x + 1) + log(x)`

3.333.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{\log(1+x)}{x^2} dx = -\frac{\log(x+1)}{x} - \log(|x+1|) + \log(|x|)$$

input `integrate(log(1+x)/x^2,x, algorithm="giac")`output `-log(x + 1)/x - log(abs(x + 1)) + log(abs(x))`**3.333.9 Mupad [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{\log(1+x)}{x^2} dx = -\ln\left(\frac{1}{x} + 1\right) - \frac{\ln(x+1)}{x}$$

input `int(log(x + 1)/x^2,x)`output `- log(1/x + 1) - log(x + 1)/x`

3.334 $\int \frac{1}{-e^x + e^{3x}} dx$

3.334.1 Optimal result	1682
3.334.2 Mathematica [A] (verified)	1682
3.334.3 Rubi [A] (verified)	1683
3.334.4 Maple [A] (verified)	1684
3.334.5 Fricas [B] (verification not implemented)	1684
3.334.6 Sympy [B] (verification not implemented)	1685
3.334.7 Maxima [A] (verification not implemented)	1685
3.334.8 Giac [A] (verification not implemented)	1685
3.334.9 Mupad [B] (verification not implemented)	1686

3.334.1 Optimal result

Integrand size = 13, antiderivative size = 12

$$\int \frac{1}{-e^x + e^{3x}} dx = e^{-x} - \operatorname{arctanh}(e^x)$$

output `exp(-x)-arctanh(exp(x))`

3.334.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{-e^x + e^{3x}} dx = e^{-x} - \operatorname{arctanh}(e^x)$$

input `Integrate[(-E^x + E^(3*x))^-1, x]`

output `E^(-x) - ArcTanh[E^x]`

3.334.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {2720, 25, 264, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{1}{e^{3x} - e^x} dx \\
 \downarrow \text{2720} \\
 \int -\frac{e^{-2x}}{1 - e^{2x}} de^x \\
 \downarrow \text{25} \\
 -\int \frac{e^{-2x}}{1 - e^{2x}} de^x \\
 \downarrow \text{264} \\
 e^{-x} - \int \frac{1}{1 - e^{2x}} de^x \\
 \downarrow \text{219} \\
 e^{-x} - \operatorname{arctanh}(e^x)
 \end{array}$$

input `Int[(-E^x + E^(3*x))^-1, x]`

output `E^(-x) - ArcTanh[E^x]`

3.334.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 264 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*((m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

3.334.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.67

method	result	size
default	$e^{-x} - \frac{\ln(1+e^x)}{2} + \frac{\ln(-1+e^x)}{2}$	20
norman	$e^{-x} - \frac{\ln(1+e^x)}{2} + \frac{\ln(-1+e^x)}{2}$	20
risch	$e^{-x} - \frac{\ln(1+e^x)}{2} + \frac{\ln(-1+e^x)}{2}$	20

input `int(1/(exp(3*x)-exp(x)),x,method=_RETURNVERBOSE)`

output `1/exp(x)-1/2*ln(1+exp(x))+1/2*ln(-1+exp(x))`

3.334.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. $2(10) = 20$.

Time = 0.24 (sec) , antiderivative size = 25, normalized size of antiderivative = 2.08

$$\int \frac{1}{-e^x + e^{3x}} dx = -\frac{1}{2} (e^x \log(e^x + 1) - e^x \log(e^x - 1) - 2)e^{(-x)}$$

input `integrate(1/(-exp(x)+exp(3*x)),x, algorithm="fracas")`

output `-1/2*(e^x*log(e^x + 1) - e^x*log(e^x - 1) - 2)*e^(-x)`

3.334.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. 2(8) = 16.

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.67

$$\int \frac{1}{-e^x + e^{3x}} dx = \frac{\log(e^x - 1)}{2} - \frac{\log(e^x + 1)}{2} + e^{-x}$$

input `integrate(1/(-exp(x)+exp(3*x)),x)`

output `log(exp(x) - 1)/2 - log(exp(x) + 1)/2 + exp(-x)`

3.334.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.58

$$\int \frac{1}{-e^x + e^{3x}} dx = e^{(-x)} - \frac{1}{2} \log(e^x + 1) + \frac{1}{2} \log(e^x - 1)$$

input `integrate(1/(-exp(x)+exp(3*x)),x, algorithm="maxima")`

output `e^(-x) - 1/2*log(e^x + 1) + 1/2*log(e^x - 1)`

3.334.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.67

$$\int \frac{1}{-e^x + e^{3x}} dx = e^{(-x)} - \frac{1}{2} \log(e^x + 1) + \frac{1}{2} \log(|e^x - 1|)$$

input `integrate(1/(-exp(x)+exp(3*x)),x, algorithm="giac")`

output `e^(-x) - 1/2*log(e^x + 1) + 1/2*log(abs(e^x - 1))`

3.334.9 Mupad [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.58

$$\int \frac{1}{-e^x + e^{3x}} dx = e^{-x} + \frac{\ln(e^x - 1)}{2} - \frac{\ln(e^x + 1)}{2}$$

input `int(1/(exp(3*x) - exp(x)),x)`

output `exp(-x) + log(exp(x) - 1)/2 - log(exp(x) + 1)/2`

3.335 $\int \frac{1+\cos^2(x)}{1-\cos^2(x)} dx$

3.335.1 Optimal result	1687
3.335.2 Mathematica [C] (verified)	1687
3.335.3 Rubi [A] (verified)	1688
3.335.4 Maple [A] (verified)	1689
3.335.5 Fricas [A] (verification not implemented)	1690
3.335.6 Sympy [A] (verification not implemented)	1690
3.335.7 Maxima [A] (verification not implemented)	1690
3.335.8 Giac [A] (verification not implemented)	1691
3.335.9 Mupad [B] (verification not implemented)	1691

3.335.1 Optimal result

Integrand size = 17, antiderivative size = 8

$$\int \frac{1 + \cos^2(x)}{1 - \cos^2(x)} dx = -x - 2 \cot(x)$$

output `-x-2*cot(x)`

3.335.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 2.88

$$\int \frac{1 + \cos^2(x)}{1 - \cos^2(x)} dx = -\cot(x) - \cot(x) \text{Hypergeometric2F1} \left(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(x) \right)$$

input `Integrate[(1 + Cos[x]^2)/(1 - Cos[x]^2),x]`

output `-Cot[x] - Cot[x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[x]^2]`

3.335.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.412$, Rules used = {3042, 3650, 3042, 3654, 3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^2(x) + 1}{1 - \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(x + \frac{\pi}{2})^2 + 1}{1 - \sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3650} \\
 & 2 \int \frac{1}{1 - \cos^2(x)} dx - x \\
 & \quad \downarrow \text{3042} \\
 & 2 \int \frac{1}{1 - \sin(x + \frac{\pi}{2})^2} dx - x \\
 & \quad \downarrow \text{3654} \\
 & 2 \int \csc^2(x) dx - x \\
 & \quad \downarrow \text{3042} \\
 & 2 \int \csc(x)^2 dx - x \\
 & \quad \downarrow \text{4254} \\
 & -2 \int 1 d \cot(x) - x \\
 & \quad \downarrow \text{24} \\
 & -x - 2 \cot(x)
 \end{aligned}$$

input `Int[(1 + Cos[x]^2)/(1 - Cos[x]^2), x]`

output `-x - 2*Cot[x]`

3.335. $\int \frac{1 + \cos^2(x)}{1 - \cos^2(x)} dx$

3.335.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3650 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]^2)/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[B*(x/b), x] + Simp[(A*b - a*B)/b Int[1/(a + b*Sin[e + f*x]^2), x], x] /; FreeQ[{a, b, e, f, A, B}, x]`

rule 3654 `Int[(u_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Simp[a^p Int[ActivateTrig[u*cos[e + f*x]^(2*p)], x], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a + b, 0] && IntegerQ[p]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

3.335.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
parallelrisc	$-x - 2 \cot(x)$	9
default	$-\frac{2}{\tan(x)} - \arctan(\tan(x))$	13
risch	$-x - \frac{4i}{e^{2ix} - 1}$	17
norman	$\frac{-1 + \tan^4(\frac{x}{2}) + \tan^6(\frac{x}{2}) - (\tan^2(\frac{x}{2})) - x \tan(\frac{x}{2}) - x(\tan^5(\frac{x}{2})) - 2(\tan^3(\frac{x}{2}))x}{(1 + \tan^2(\frac{x}{2}))^2 \tan(\frac{x}{2})}$	65

input `int((1+cos(x)^2)/(1-cos(x)^2),x,method=_RETURNVERBOSE)`

output `-x-2*cot(x)`

3.335.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.88

$$\int \frac{1 + \cos^2(x)}{1 - \cos^2(x)} dx = -\frac{x \sin(x) + 2 \cos(x)}{\sin(x)}$$

input `integrate((1+cos(x)^2)/(1-cos(x)^2),x, algorithm="fricas")`output `-(x*sin(x) + 2*cos(x))/sin(x)`**3.335.6 Sympy [A] (verification not implemented)**

Time = 0.35 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.50

$$\int \frac{1 + \cos^2(x)}{1 - \cos^2(x)} dx = -x + \tan\left(\frac{x}{2}\right) - \frac{1}{\tan\left(\frac{x}{2}\right)}$$

input `integrate((1+cos(x)**2)/(1-cos(x)**2),x)`output `-x + tan(x/2) - 1/tan(x/2)`**3.335.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.25

$$\int \frac{1 + \cos^2(x)}{1 - \cos^2(x)} dx = -x - \frac{2}{\tan(x)}$$

input `integrate((1+cos(x)^2)/(1-cos(x)^2),x, algorithm="maxima")`output `-x - 2/tan(x)`

3.335.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 16, normalized size of antiderivative = 2.00

$$\int \frac{1 + \cos^2(x)}{1 - \cos^2(x)} dx = -x - \frac{1}{\tan\left(\frac{1}{2}x\right)} + \tan\left(\frac{1}{2}x\right)$$

input `integrate((1+cos(x)^2)/(1-cos(x)^2),x, algorithm="giac")`output `-x - 1/tan(1/2*x) + tan(1/2*x)`**3.335.9 Mupad [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{1 + \cos^2(x)}{1 - \cos^2(x)} dx = -x - 2 \cot(x)$$

input `int(-(cos(x)^2 + 1)/(cos(x)^2 - 1),x)`output `- x - 2*cot(x)`

$$\mathbf{3.336} \quad \int \frac{1}{x\sqrt{-25+2x}} dx$$

3.336.1 Optimal result	1692
3.336.2 Mathematica [A] (verified)	1692
3.336.3 Rubi [A] (verified)	1693
3.336.4 Maple [A] (verified)	1694
3.336.5 Fricas [A] (verification not implemented)	1694
3.336.6 Sympy [C] (verification not implemented)	1695
3.336.7 Maxima [A] (verification not implemented)	1695
3.336.8 Giac [A] (verification not implemented)	1695
3.336.9 Mupad [B] (verification not implemented)	1696

3.336.1 Optimal result

Integrand size = 13, antiderivative size = 18

$$\int \frac{1}{x\sqrt{-25+2x}} dx = \frac{2}{5} \arctan\left(\frac{1}{5}\sqrt{-25+2x}\right)$$

output `2/5*arctan(1/5*(-25+2*x)^(1/2))`

3.336.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{-25+2x}} dx = \frac{2}{5} \arctan\left(\frac{1}{5}\sqrt{-25+2x}\right)$$

input `Integrate[1/(x*Sqrt[-25 + 2*x]),x]`

output `(2*ArcTan[Sqrt[-25 + 2*x]/5])/5`

3.336.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x\sqrt{2x-25}} dx$$

↓ 73

$$\int \frac{1}{\frac{1}{2}(2x-25) + \frac{25}{2}} d\sqrt{2x-25}$$

↓ 216

$$\frac{2}{5} \arctan\left(\frac{1}{5}\sqrt{2x-25}\right)$$

input `Int[1/(x*Sqrt[-25 + 2*x]),x]`

output `(2*ArcTan[Sqrt[-25 + 2*x]/5])/5`

3.336.3.1 Defintions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

3.336.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.72

method	result	size
derivativedivides	$\frac{2 \arctan\left(\frac{\sqrt{-25+2x}}{5}\right)}{5}$	13
default	$\frac{2 \arctan\left(\frac{\sqrt{-25+2x}}{5}\right)}{5}$	13
pseudoelliptic	$\frac{2 \arctan\left(\frac{\sqrt{-25+2x}}{5}\right)}{5}$	13
trager	$\frac{\text{RootOf}\left(_Z^2+1\right) \ln\left(\frac{\text{RootOf}\left(_Z^2+1\right) x-25 \text{RootOf}\left(_Z^2+1\right)+5 \sqrt{-25+2x}}{x}\right)}{5}$	40
meijerg	$\frac{\sqrt{-\text{signum}\left(x-\frac{25}{2}\right)}\left(-\ln(2)+\ln(x)-2 \ln(5)+i \pi \sqrt{\pi}-2 \sqrt{\pi} \ln\left(\frac{1}{2}+\frac{\sqrt{1-\frac{2x}{25}}}{2}\right)\right)}{5 \sqrt{\pi} \sqrt{\text{signum}\left(x-\frac{25}{2}\right)}}$	57

input `int(1/x/(-25+2*x)^(1/2),x,method=_RETURNVERBOSE)`output `2/5*arctan(1/5*(-25+2*x)^(1/2))`**3.336.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \frac{1}{x\sqrt{-25+2x}} dx = \frac{2}{5} \arctan\left(\frac{1}{5} \sqrt{2x-25}\right)$$

input `integrate(1/x/(-25+2*x)^(1/2),x, algorithm="fricas")`output `2/5*arctan(1/5*sqrt(2*x - 25))`

3.336.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.49 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.44

$$\int \frac{1}{x\sqrt{-25+2x}} dx = \begin{cases} \frac{2i \operatorname{acosh}\left(\frac{5\sqrt{2}}{2\sqrt{x}}\right)}{5} & \text{for } \frac{1}{|x|} > \frac{2}{25} \\ -\frac{2 \operatorname{asin}\left(\frac{5\sqrt{2}}{2\sqrt{x}}\right)}{5} & \text{otherwise} \end{cases}$$

input `integrate(1/x/(-25+2*x)**(1/2),x)`

output `Piecewise((2*I*acosh(5*sqrt(2)/(2*sqrt(x)))/5, 1/Abs(x) > 2/25), (-2*asin(5*sqrt(2)/(2*sqrt(x)))/5, True))`

3.336.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \frac{1}{x\sqrt{-25+2x}} dx = \frac{2}{5} \arctan\left(\frac{1}{5}\sqrt{2x-25}\right)$$

input `integrate(1/x/(-25+2*x)^(1/2),x, algorithm="maxima")`

output `2/5*arctan(1/5*sqrt(2*x - 25))`

3.336.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \frac{1}{x\sqrt{-25+2x}} dx = \frac{2}{5} \arctan\left(\frac{1}{5}\sqrt{2x-25}\right)$$

input `integrate(1/x/(-25+2*x)^(1/2),x, algorithm="giac")`

output `2/5*arctan(1/5*sqrt(2*x - 25))`

3.336.9 Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \frac{1}{x\sqrt{-25+2x}} dx = \frac{2 \operatorname{atan}\left(\frac{\sqrt{2x-25}}{5}\right)}{5}$$

input `int(1/(x*(2*x - 25)^(1/2)),x)`

output `(2*atan((2*x - 25)^(1/2)/5))/5`

3.337 $\int \frac{\sin(2x)}{\sqrt{9-\cos^4(x)}} dx$

3.337.1 Optimal result 1697
 3.337.2 Mathematica [A] (verified) 1697
 3.337.3 Rubi [A] (verified) 1698
 3.337.4 Maple [A] (verified) 1699
 3.337.5 Fricas [B] (verification not implemented) 1700
 3.337.6 Sympy [F(-1)] 1700
 3.337.7 Maxima [F] 1700
 3.337.8 Giac [A] (verification not implemented) 1701
 3.337.9 Mupad [B] (verification not implemented) 1701

3.337.1 Optimal result

Integrand size = 17, antiderivative size = 11

$$\int \frac{\sin(2x)}{\sqrt{9-\cos^4(x)}} dx = -\arcsin\left(\frac{\cos^2(x)}{3}\right)$$

output `-arcsin(1/3*cos(x)^2)`

3.337.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{\sin(2x)}{\sqrt{9-\cos^4(x)}} dx = -\arcsin\left(\frac{\cos^2(x)}{3}\right)$$

input `Integrate[Sin[2*x]/Sqrt[9 - Cos[x]^4],x]`

output `-ArcSin[Cos[x]^2/3]`

3.337.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.36, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3042, 4878, 27, 1432, 1090, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin(2x)}{\sqrt{9 - \cos^4(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(2x)}{\sqrt{9 - \cos(x)^4}} dx \\
 & \quad \downarrow \text{4878} \\
 & \int \frac{2 \sin(x)}{\sqrt{-\sin^4(x) + 2 \sin^2(x) + 8}} d \sin(x) \\
 & \quad \downarrow \text{27} \\
 & 2 \int \frac{\sin(x)}{\sqrt{-\sin^4(x) + 2 \sin^2(x) + 8}} d \sin(x) \\
 & \quad \downarrow \text{1432} \\
 & \int \frac{1}{\sqrt{-\sin^4(x) + 2 \sin^2(x) + 8}} d \sin^2(x) \\
 & \quad \downarrow \text{1090} \\
 & -\frac{1}{6} \int \frac{1}{\sqrt{1 - \frac{\sin^4(x)}{36}}} d(2 - 2 \sin^2(x)) \\
 & \quad \downarrow \text{223} \\
 & -\arcsin\left(\frac{1}{6}(2 - 2 \sin^2(x))\right)
 \end{aligned}$$

input `Int[Sin[2*x]/Sqrt[9 - Cos[x]^4], x]`

output `-ArcSin[(2 - 2*Sin[x]^2)/6]`

3.337.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 1090 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`
- rule 1432 `Int[(x_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, p}, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4878 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Sin[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Sin[v]/d, u/Cos[v], x], x], x, Sin[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Sin[v], x], u/Cos[v], x]`

3.337.4 Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

method	result	size
derivativedivides	$-\arcsin\left(\frac{\cos^2(x)}{3}\right)$	10
default	$-\arcsin\left(\frac{\cos^2(x)}{3}\right)$	10

input `int(sin(2*x)/(9-cos(x)^4)^(1/2),x,method=_RETURNVERBOSE)`

output `-arcsin(1/3*cos(x)^2)`

3.337.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 24 vs. 2(9) = 18.

Time = 0.32 (sec) , antiderivative size = 24, normalized size of antiderivative = 2.18

$$\int \frac{\sin(2x)}{\sqrt{9 - \cos^4(x)}} dx = \arctan \left(\frac{\sqrt{-\cos(x)^4 + 9} \cos(x)^2}{\cos(x)^4 - 9} \right)$$

input `integrate(sin(2*x)/(9-cos(x)^4)^(1/2),x, algorithm="fricas")`

output `arctan(sqrt(-cos(x)^4 + 9)*cos(x)^2/(cos(x)^4 - 9))`

3.337.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sin(2x)}{\sqrt{9 - \cos^4(x)}} dx = \text{Timed out}$$

input `integrate(sin(2*x)/(9-cos(x)**4)**(1/2),x)`

output `Timed out`

3.337.7 Maxima [F]

$$\int \frac{\sin(2x)}{\sqrt{9 - \cos^4(x)}} dx = \int \frac{\sin(2x)}{\sqrt{-\cos(x)^4 + 9}} dx$$

input `integrate(sin(2*x)/(9-cos(x)^4)^(1/2),x, algorithm="maxima")`

output `integrate(sin(2*x)/sqrt(-cos(x)^4 + 9), x)`

3.337.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \frac{\sin(2x)}{\sqrt{9 - \cos^4(x)}} dx = -\arcsin\left(\frac{1}{3} \cos(x)^2\right)$$

input `integrate(sin(2*x)/(9-cos(x)^4)^(1/2),x, algorithm="giac")`output `-arcsin(1/3*cos(x)^2)`**3.337.9 Mupad [B] (verification not implemented)**

Time = 0.44 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.64

$$\int \frac{\sin(2x)}{\sqrt{9 - \cos^4(x)}} dx = -\operatorname{atan}\left(\frac{\cos(x)^2}{\sqrt{9 - \cos(x)^4}}\right)$$

input `int(sin(2*x)/(9 - cos(x)^4)^(1/2),x)`output `-atan(cos(x)^2/(9 - cos(x)^4)^(1/2))`

3.338 $\int \frac{x^2}{\sqrt{5-4x^2}} dx$

3.338.1 Optimal result	1702
3.338.2 Mathematica [A] (verified)	1702
3.338.3 Rubi [A] (verified)	1703
3.338.4 Maple [A] (verified)	1704
3.338.5 Fricas [A] (verification not implemented)	1704
3.338.6 Sympy [A] (verification not implemented)	1705
3.338.7 Maxima [A] (verification not implemented)	1705
3.338.8 Giac [A] (verification not implemented)	1705
3.338.9 Mupad [B] (verification not implemented)	1706

3.338.1 Optimal result

Integrand size = 15, antiderivative size = 30

$$\int \frac{x^2}{\sqrt{5-4x^2}} dx = -\frac{1}{8}x\sqrt{5-4x^2} + \frac{5}{16} \arcsin\left(\frac{2x}{\sqrt{5}}\right)$$

output `5/16*arcsin(2/5*x*5^(1/2))-1/8*x*(-4*x^2+5)^(1/2)`

3.338.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.53

$$\int \frac{x^2}{\sqrt{5-4x^2}} dx = -\frac{1}{8}x\sqrt{5-4x^2} - \frac{5}{8} \arctan\left(\frac{2x}{\sqrt{5}-\sqrt{5-4x^2}}\right)$$

input `Integrate[x^2/Sqrt[5 - 4*x^2],x]`

output `-1/8*(x*Sqrt[5 - 4*x^2]) - (5*ArcTan[(2*x)/(Sqrt[5] - Sqrt[5 - 4*x^2])])/8`

3.338.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {262, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{\sqrt{5-4x^2}} dx$$

↓ 262

$$\frac{5}{8} \int \frac{1}{\sqrt{5-4x^2}} dx - \frac{1}{8} x \sqrt{5-4x^2}$$

↓ 223

$$\frac{5}{16} \arcsin\left(\frac{2x}{\sqrt{5}}\right) - \frac{1}{8} x \sqrt{5-4x^2}$$

input `Int[x^2/Sqrt[5 - 4*x^2],x]`

output `-1/8*(x*Sqrt[5 - 4*x^2]) + (5*ArcSin[(2*x)/Sqrt[5]])/16`

3.338.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 262 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m-1)*((a + b*x^2)^(p+1)/(b*(m+2*p+1))), x] - Simp[a*c^2*((m-1)/(b*(m+2*p+1)) Int[(c*x)^(m-2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2-1] && NeQ[m+2*p+1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

3.338.4 Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.77

method	result	size
default	$\frac{5 \arcsin\left(\frac{2x\sqrt{5}}{5}\right)}{16} - \frac{x\sqrt{-4x^2+5}}{8}$	23
risch	$\frac{x(4x^2-5)}{8\sqrt{-4x^2+5}} + \frac{5 \arcsin\left(\frac{2x\sqrt{5}}{5}\right)}{16}$	30
pseudoelliptic	$-\frac{x\sqrt{-4x^2+5}}{8} - \frac{5 \arctan\left(\frac{\sqrt{-4x^2+5}}{2x}\right)}{16}$	31
meijerg	$\frac{5i \left(\frac{2i\sqrt{\pi} x \sqrt{5} \sqrt{-\frac{4x^2}{5}+1}}{5} - i\sqrt{\pi} \arcsin\left(\frac{2x\sqrt{5}}{5}\right) \right)}{16\sqrt{\pi}}$	40
trager	$-\frac{x\sqrt{-4x^2+5}}{8} + \frac{5 \operatorname{RootOf}\left(-Z^2+1\right) \ln\left(\operatorname{RootOf}\left(-Z^2+1\right)\sqrt{-4x^2+5}+2x\right)}{16}$	43

input `int(x^2/(-4*x^2+5)^(1/2),x,method=_RETURNVERBOSE)`output `5/16*arcsin(2/5*x*5^(1/2))-1/8*x*(-4*x^2+5)^(1/2)`**3.338.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{\sqrt{5-4x^2}} dx = -\frac{1}{8} \sqrt{-4x^2+5} - \frac{5}{16} \arctan\left(\frac{\sqrt{-4x^2+5}}{2x}\right)$$

input `integrate(x^2/(-4*x^2+5)^(1/2),x, algorithm="fricas")`output `-1/8*sqrt(-4*x^2 + 5)*x - 5/16*arctan(1/2*sqrt(-4*x^2 + 5)/x)`

3.338.6 Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.90

$$\int \frac{x^2}{\sqrt{5-4x^2}} dx = -\frac{x\sqrt{5-4x^2}}{8} + \frac{5 \operatorname{asin}\left(\frac{2\sqrt{5}x}{5}\right)}{16}$$

input `integrate(x**2/(-4*x**2+5)**(1/2),x)`output `-x*sqrt(5 - 4*x**2)/8 + 5*asin(2*sqrt(5)*x/5)/16`**3.338.7 Maxima [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \frac{x^2}{\sqrt{5-4x^2}} dx = -\frac{1}{8} \sqrt{-4x^2+5}x + \frac{5}{16} \arcsin\left(\frac{2}{5} \sqrt{5}x\right)$$

input `integrate(x^2/(-4*x^2+5)^(1/2),x, algorithm="maxima")`output `-1/8*sqrt(-4*x^2 + 5)*x + 5/16*arcsin(2/5*sqrt(5)*x)`**3.338.8 Giac [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \frac{x^2}{\sqrt{5-4x^2}} dx = -\frac{1}{8} \sqrt{-4x^2+5}x + \frac{5}{16} \arcsin\left(\frac{2}{5} \sqrt{5}x\right)$$

input `integrate(x^2/(-4*x^2+5)^(1/2),x, algorithm="giac")`output `-1/8*sqrt(-4*x^2 + 5)*x + 5/16*arcsin(2/5*sqrt(5)*x)`

3.338.9 Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \frac{x^2}{\sqrt{5-4x^2}} dx = \frac{5 \operatorname{asin}\left(\frac{2\sqrt{5}x}{5}\right)}{16} - \frac{x \sqrt{\frac{5}{4} - x^2}}{4}$$

input `int(x^2/(5 - 4*x^2)^(1/2),x)`

output `(5*asin((2*5^(1/2)*x)/5))/16 - (x*(5/4 - x^2)^(1/2))/4`

3.339 $\int x^3 \sin(x) dx$

3.339.1 Optimal result	1707
3.339.2 Mathematica [A] (verified)	1707
3.339.3 Rubi [A] (verified)	1708
3.339.4 Maple [A] (verified)	1709
3.339.5 Fracas [A] (verification not implemented)	1710
3.339.6 Sympy [A] (verification not implemented)	1710
3.339.7 Maxima [A] (verification not implemented)	1710
3.339.8 Giac [A] (verification not implemented)	1711
3.339.9 Mupad [B] (verification not implemented)	1711

3.339.1 Optimal result

Integrand size = 6, antiderivative size = 24

$$\int x^3 \sin(x) dx = 6x \cos(x) - x^3 \cos(x) - 6 \sin(x) + 3x^2 \sin(x)$$

output `6*x*cos(x)-x^3*cos(x)-6*sin(x)+3*x^2*sin(x)`

3.339.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int x^3 \sin(x) dx = -x(-6 + x^2) \cos(x) + 3(-2 + x^2) \sin(x)$$

input `Integrate[x^3*Sin[x],x]`

output `-(x*(-6 + x^2)*Cos[x]) + 3*(-2 + x^2)*Sin[x]`

3.339.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.12, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {3042, 3777, 3042, 3777, 25, 3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \sin(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x^3 \sin(x) dx \\
 & \quad \downarrow \text{3777} \\
 & 3 \int x^2 \cos(x) dx - x^3 \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & 3 \int x^2 \sin\left(x + \frac{\pi}{2}\right) dx - x^3 \cos(x) \\
 & \quad \downarrow \text{3777} \\
 & 3 \left(2 \int -x \sin(x) dx + x^2 \sin(x) \right) - x^3 \cos(x) \\
 & \quad \downarrow \text{25} \\
 & 3 \left(x^2 \sin(x) - 2 \int x \sin(x) dx \right) - x^3 \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & 3 \left(x^2 \sin(x) - 2 \int x \sin(x) dx \right) - x^3 \cos(x) \\
 & \quad \downarrow \text{3777} \\
 & 3 \left(x^2 \sin(x) - 2 \left(\int \cos(x) dx - x \cos(x) \right) \right) - x^3 \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & 3 \left(x^2 \sin(x) - 2 \left(\int \sin\left(x + \frac{\pi}{2}\right) dx - x \cos(x) \right) \right) - x^3 \cos(x) \\
 & \quad \downarrow \text{3117}
 \end{aligned}$$

$$3(x^2 \sin(x) - 2(\sin(x) - x \cos(x))) - x^3 \cos(x)$$

input `Int[x^3*Sin[x],x]`

output `-(x^3*Cos[x]) + 3*(x^2*Sin[x] - 2*(-(x*Cos[x]) + Sin[x]))`

3.339.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

3.339.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

method	result	size
risch	$(-x^3 + 6x) \cos(x) + 3(x^2 - 2) \sin(x)$	23
default	$6x \cos(x) - x^3 \cos(x) - 6 \sin(x) + 3x^2 \sin(x)$	25
parallelrisch	$6x \cos(x) - x^3 \cos(x) - 6 \sin(x) + 3x^2 \sin(x)$	25
parts	$6x \cos(x) - x^3 \cos(x) - 6 \sin(x) + 3x^2 \sin(x)$	25
meijerg	$8\sqrt{\pi} \left(\frac{x \left(-\frac{5x^2}{2} + 15\right) \cos(x)}{20\sqrt{\pi}} - \frac{\left(-\frac{15x^2}{2} + 15\right) \sin(x)}{20\sqrt{\pi}} \right)$	36
norman	$\frac{x^3 \left(\tan^2\left(\frac{x}{2}\right) + 6x - x^3 - 6x \tan^2\left(\frac{x}{2}\right) + 6x^2 \tan\left(\frac{x}{2}\right) - 12 \tan\left(\frac{x}{2}\right)\right)}{1 + \tan^2\left(\frac{x}{2}\right)}$	55

input `int(x^3*sin(x),x,method=_RETURNVERBOSE)`

output `(-x^3+6*x)*cos(x)+3*(x^2-2)*sin(x)`

3.339.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

$$\int x^3 \sin(x) dx = -(x^3 - 6x) \cos(x) + 3(x^2 - 2) \sin(x)$$

input `integrate(x^3*sin(x),x, algorithm="fricas")`

output `-(x^3 - 6*x)*cos(x) + 3*(x^2 - 2)*sin(x)`

3.339.6 Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int x^3 \sin(x) dx = -x^3 \cos(x) + 3x^2 \sin(x) + 6x \cos(x) - 6 \sin(x)$$

input `integrate(x**3*sin(x),x)`

output `-x**3*cos(x) + 3*x**2*sin(x) + 6*x*cos(x) - 6*sin(x)`

3.339.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

$$\int x^3 \sin(x) dx = -(x^3 - 6x) \cos(x) + 3(x^2 - 2) \sin(x)$$

input `integrate(x^3*sin(x),x, algorithm="maxima")`

output `(-x^3 - 6*x)*cos(x) + 3*(x^2 - 2)*sin(x)`

3.339.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

$$\int x^3 \sin(x) dx = -(x^3 - 6x) \cos(x) + 3(x^2 - 2) \sin(x)$$

input `integrate(x^3*sin(x),x, algorithm="giac")`

output `-(x^3 - 6*x)*cos(x) + 3*(x^2 - 2)*sin(x)`

3.339.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

$$\int x^3 \sin(x) dx = \cos(x) (6x - x^3) + \sin(x) (3x^2 - 6)$$

input `int(x^3*sin(x),x)`

output `cos(x)*(6*x - x^3) + sin(x)*(3*x^2 - 6)`

3.340 $\int x\sqrt{4 + 2x + x^2} dx$

3.340.1 Optimal result	1712
3.340.2 Mathematica [A] (verified)	1712
3.340.3 Rubi [A] (verified)	1713
3.340.4 Maple [A] (verified)	1714
3.340.5 Fricas [A] (verification not implemented)	1715
3.340.6 Sympy [A] (verification not implemented)	1715
3.340.7 Maxima [A] (verification not implemented)	1715
3.340.8 Giac [A] (verification not implemented)	1716
3.340.9 Mupad [B] (verification not implemented)	1716

3.340.1 Optimal result

Integrand size = 14, antiderivative size = 50

$$\int x\sqrt{4 + 2x + x^2} dx = -\frac{1}{2}(1 + x)\sqrt{4 + 2x + x^2} + \frac{1}{3}(4 + 2x + x^2)^{3/2} - \frac{3}{2}\operatorname{arcsinh}\left(\frac{1 + x}{\sqrt{3}}\right)$$

output `1/3*(x^2+2*x+4)^(3/2)-3/2*arcsinh(1/3*(1+x)*3^(1/2))-1/2*(1+x)*(x^2+2*x+4)^(1/2)`

3.340.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.94

$$\int x\sqrt{4 + 2x + x^2} dx = \frac{1}{6}\sqrt{4 + 2x + x^2}(5 + x + 2x^2) + \frac{3}{2}\log\left(-1 - x + \sqrt{4 + 2x + x^2}\right)$$

input `Integrate[x*Sqrt[4 + 2*x + x^2],x]`

output `(Sqrt[4 + 2*x + x^2]*(5 + x + 2*x^2))/6 + (3*Log[-1 - x + Sqrt[4 + 2*x + x^2]])/2`

3.340.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.10, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {1160, 1087, 1090, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x\sqrt{x^2 + 2x + 4} dx \\
 & \quad \downarrow \text{1160} \\
 & \frac{1}{3}(x^2 + 2x + 4)^{3/2} - \int \sqrt{x^2 + 2x + 4} dx \\
 & \quad \downarrow \text{1087} \\
 & -\frac{3}{2} \int \frac{1}{\sqrt{x^2 + 2x + 4}} dx + \frac{1}{3}(x^2 + 2x + 4)^{3/2} - \frac{1}{2}(x + 1)\sqrt{x^2 + 2x + 4} \\
 & \quad \downarrow \text{1090} \\
 & -\frac{1}{4}\sqrt{3} \int \frac{1}{\sqrt{\frac{1}{12}(2x + 2)^2 + 1}} d(2x + 2) + \frac{1}{3}(x^2 + 2x + 4)^{3/2} - \frac{1}{2}(x + 1)\sqrt{x^2 + 2x + 4} \\
 & \quad \downarrow \text{222} \\
 & -\frac{3}{2} \operatorname{arcsinh}\left(\frac{2x + 2}{2\sqrt{3}}\right) + \frac{1}{3}(x^2 + 2x + 4)^{3/2} - \frac{1}{2}(x + 1)\sqrt{x^2 + 2x + 4}
 \end{aligned}$$

input `Int[x*Sqrt[4 + 2*x + x^2],x]`

output `-1/2*((1 + x)*Sqrt[4 + 2*x + x^2]) + (4 + 2*x + x^2)^(3/2)/3 - (3*ArcSinh[(2 + 2*x)/(2*Sqrt[3])])/2`

3.340.3.1 Defintions of rubi rules used

- rule 222 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`
- rule 1087 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x) * ((a + b*x + c*x^2)^p / (2*c*(2*p + 1))), x] - Simp[p*(b^2 - 4*a*c) / (2*c*(2*p + 1)) Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[3*p])`
- rule 1090 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[1 / (2*c*(-4*c/(b^2 - 4*a*c)))^p] Subst[Int[Simp[1 - x^2 / (b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`
- rule 1160 `Int[((d_) + (e_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[e*((a + b*x + c*x^2)^(p + 1) / (2*c*(p + 1))), x] + Simp[(2*c*d - b*e) / (2*c) Int[(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[p, -1]`

3.340.4 Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.66

method	result	size
risch	$\frac{(2x^2+x+5)\sqrt{x^2+2x+4}}{6} - \frac{3 \operatorname{arcsinh}\left(\frac{(1+x)\sqrt{3}}{3}\right)}{2}$	33
trager	$\left(\frac{1}{3}x^2 + \frac{1}{6}x + \frac{5}{6}\right)\sqrt{x^2+2x+4} - \frac{3 \ln\left(1+x+\sqrt{x^2+2x+4}\right)}{2}$	39
default	$\frac{(x^2+2x+4)^{\frac{3}{2}}}{3} - \frac{(2x+2)\sqrt{x^2+2x+4}}{4} - \frac{3 \operatorname{arcsinh}\left(\frac{(1+x)\sqrt{3}}{3}\right)}{2}$	42

input `int(x*(x^2+2*x+4)^(1/2),x,method=_RETURNVERBOSE)`

output `1/6*(2*x^2+x+5)*(x^2+2*x+4)^(1/2)-3/2*arcsinh(1/3*(1+x)*3^(1/2))`

3.340.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.78

$$\int x\sqrt{4+2x+x^2} dx = \frac{1}{6}(2x^2+x+5)\sqrt{x^2+2x+4} + \frac{3}{2}\log(-x+\sqrt{x^2+2x+4}-1)$$

input `integrate(x*(x^2+2*x+4)^(1/2),x, algorithm="fricas")`output `1/6*(2*x^2 + x + 5)*sqrt(x^2 + 2*x + 4) + 3/2*log(-x + sqrt(x^2 + 2*x + 4) - 1)`**3.340.6 Sympy [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.74

$$\int x\sqrt{4+2x+x^2} dx = \left(\frac{x^2}{3} + \frac{x}{6} + \frac{5}{6}\right)\sqrt{x^2+2x+4} - \frac{3\operatorname{asinh}\left(\frac{\sqrt{3}(x+1)}{3}\right)}{2}$$

input `integrate(x*(x**2+2*x+4)**(1/2),x)`output `(x**2/3 + x/6 + 5/6)*sqrt(x**2 + 2*x + 4) - 3*asinh(sqrt(3)*(x + 1)/3)/2`**3.340.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.98

$$\int x\sqrt{4+2x+x^2} dx = \frac{1}{3}(x^2+2x+4)^{\frac{3}{2}} - \frac{1}{2}\sqrt{x^2+2x+4}x - \frac{1}{2}\sqrt{x^2+2x+4} - \frac{3}{2}\operatorname{arsinh}\left(\frac{1}{3}\sqrt{3}(x+1)\right)$$

input `integrate(x*(x^2+2*x+4)^(1/2),x, algorithm="maxima")`output `1/3*(x^2 + 2*x + 4)^(3/2) - 1/2*sqrt(x^2 + 2*x + 4)*x - 1/2*sqrt(x^2 + 2*x + 4) - 3/2*arcsinh(1/3*sqrt(3)*(x + 1))`

3.340.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.80

$$\int x\sqrt{4+2x+x^2} dx = \frac{1}{6}((2x+1)x+5)\sqrt{x^2+2x+4} + \frac{3}{2} \log(-x + \sqrt{x^2+2x+4} - 1)$$

input `integrate(x*(x^2+2*x+4)^(1/2),x, algorithm="giac")`output `1/6*((2*x + 1)*x + 5)*sqrt(x^2 + 2*x + 4) + 3/2*log(-x + sqrt(x^2 + 2*x + 4) - 1)`**3.340.9 Mupad [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.78

$$\int x\sqrt{4+2x+x^2} dx = \frac{\sqrt{x^2+2x+4}(8x^2+4x+20)}{24} - \frac{3 \ln(x + \sqrt{x^2+2x+4} + 1)}{2}$$

input `int(x*(2*x + x^2 + 4)^(1/2),x)`output `((2*x + x^2 + 4)^(1/2)*(4*x + 8*x^2 + 20))/24 - (3*log(x + (2*x + x^2 + 4)^(1/2) + 1))/2`

3.341 $\int x(5 + x^2)^8 dx$

3.341.1 Optimal result	1717
3.341.2 Mathematica [A] (verified)	1717
3.341.3 Rubi [A] (verified)	1718
3.341.4 Maple [A] (verified)	1718
3.341.5 Fricas [B] (verification not implemented)	1719
3.341.6 Sympy [B] (verification not implemented)	1719
3.341.7 Maxima [A] (verification not implemented)	1720
3.341.8 Giac [A] (verification not implemented)	1720
3.341.9 Mupad [B] (verification not implemented)	1720

3.341.1 Optimal result

Integrand size = 9, antiderivative size = 11

$$\int x(5 + x^2)^8 dx = \frac{1}{18}(5 + x^2)^9$$

output `1/18*(x^2+5)^9`

3.341.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int x(5 + x^2)^8 dx = \frac{1}{18}(5 + x^2)^9$$

input `Integrate[x*(5 + x^2)^8,x]`

output `(5 + x^2)^9/18`

3.341.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(x^2 + 5)^8 dx$$

↓ 241

$$\frac{1}{18}(x^2 + 5)^9$$

input `Int[x*(5 + x^2)^8,x]`

output `(5 + x^2)^9/18`

3.341.3.1 Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

3.341.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

method	result
default	$\frac{(x^2+5)^9}{18}$
gosper	$\frac{390625}{2}x^2 + 156250x^4 + \frac{218750}{3}x^6 + 21875x^8 + 4375x^{10} + \frac{1750}{3}x^{12} + 50x^{14} + \frac{5}{2}x^{16} + \frac{1}{18}x^{18}$
norman	$\frac{390625}{2}x^2 + 156250x^4 + \frac{218750}{3}x^6 + 21875x^8 + 4375x^{10} + \frac{1750}{3}x^{12} + 50x^{14} + \frac{5}{2}x^{16} + \frac{1}{18}x^{18}$
parallelrisch	$\frac{390625}{2}x^2 + 156250x^4 + \frac{218750}{3}x^6 + 21875x^8 + 4375x^{10} + \frac{1750}{3}x^{12} + 50x^{14} + \frac{5}{2}x^{16} + \frac{1}{18}x^{18}$
risch	$\frac{1}{18}x^{18} + \frac{5}{2}x^{16} + 50x^{14} + \frac{1750}{3}x^{12} + 4375x^{10} + 21875x^8 + \frac{218750}{3}x^6 + 156250x^4 + \frac{390625}{2}x^2 + \frac{1}{18}$

input `int(x*(x^2+5)^8,x,method=_RETURNVERBOSE)`

output `1/18*(x^2+5)^9`

3.341.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 46 vs. $2(9) = 18$.

Time = 0.24 (sec) , antiderivative size = 46, normalized size of antiderivative = 4.18

$$\int x(5+x^2)^8 dx = \frac{1}{18}x^{18} + \frac{5}{2}x^{16} + 50x^{14} + \frac{1750}{3}x^{12} + 4375x^{10} \\ + 21875x^8 + \frac{218750}{3}x^6 + 156250x^4 + \frac{390625}{2}x^2$$

input `integrate(x*(x^2+5)^8,x, algorithm="fricas")`

output `1/18*x^18 + 5/2*x^16 + 50*x^14 + 1750/3*x^12 + 4375*x^10 + 21875*x^8 + 218750/3*x^6 + 156250*x^4 + 390625/2*x^2`

3.341.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 51 vs. $2(7) = 14$.

Time = 0.03 (sec) , antiderivative size = 51, normalized size of antiderivative = 4.64

$$\int x(5+x^2)^8 dx = \frac{x^{18}}{18} + \frac{5x^{16}}{2} + 50x^{14} + \frac{1750x^{12}}{3} + 4375x^{10} \\ + 21875x^8 + \frac{218750x^6}{3} + 156250x^4 + \frac{390625x^2}{2}$$

input `integrate(x*(x**2+5)**8,x)`

output `x**18/18 + 5*x**16/2 + 50*x**14 + 1750*x**12/3 + 4375*x**10 + 21875*x**8 + 218750*x**6/3 + 156250*x**4 + 390625*x**2/2`

3.341.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int x(5 + x^2)^8 dx = \frac{1}{18} (x^2 + 5)^9$$

input `integrate(x*(x^2+5)^8,x, algorithm="maxima")`output `1/18*(x^2 + 5)^9`**3.341.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int x(5 + x^2)^8 dx = \frac{1}{18} (x^2 + 5)^9$$

input `integrate(x*(x^2+5)^8,x, algorithm="giac")`output `1/18*(x^2 + 5)^9`**3.341.9 Mupad [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int x(5 + x^2)^8 dx = \frac{(x^2 + 5)^9}{18}$$

input `int(x*(x^2 + 5)^8,x)`output `(x^2 + 5)^9/18`

3.342 $\int \cos^2(x) \sin^5(x) dx$

3.342.1 Optimal result1721
3.342.2 Mathematica [A] (verified)1721
3.342.3 Rubi [A] (verified)1722
3.342.4 Maple [A] (verified)1723
3.342.5 Fricas [A] (verification not implemented)1723
3.342.6 Sympy [A] (verification not implemented)1724
3.342.7 Maxima [A] (verification not implemented)1724
3.342.8 Giac [A] (verification not implemented)1724
3.342.9 Mupad [B] (verification not implemented)1725

3.342.1 Optimal result

Integrand size = 9, antiderivative size = 25

$$\int \cos^2(x) \sin^5(x) dx = -\frac{1}{3} \cos^3(x) + \frac{2 \cos^5(x)}{5} - \frac{\cos^7(x)}{7}$$

output `-1/3*cos(x)^3+2/5*cos(x)^5-1/7*cos(x)^7`

3.342.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.24

$$\int \cos^2(x) \sin^5(x) dx = -\frac{5 \cos(x)}{64} - \frac{1}{192} \cos(3x) + \frac{3}{320} \cos(5x) - \frac{1}{448} \cos(7x)$$

input `Integrate[Cos[x]^2*Sin[x]^5,x]`

output `(-5*Cos[x])/64 - Cos[3*x]/192 + (3*Cos[5*x])/320 - Cos[7*x]/448`

3.342.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 3045, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^5(x) \cos^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^5 \cos(x)^2 dx \\
 & \quad \downarrow \text{3045} \\
 & - \int \cos^2(x) (1 - \cos^2(x))^2 d \cos(x) \\
 & \quad \downarrow \text{244} \\
 & - \int (\cos^6(x) - 2 \cos^4(x) + \cos^2(x)) d \cos(x) \\
 & \quad \downarrow \text{2009} \\
 & -\frac{1}{7} \cos^7(x) + \frac{2 \cos^5(x)}{5} - \frac{\cos^3(x)}{3}
 \end{aligned}$$

input `Int[Cos[x]^2*Sin[x]^5,x]`

output `-1/3*Cos[x]^3 + (2*Cos[x]^5)/5 - Cos[x]^7/7`

3.342.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3045 `Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[-(a*f)^(-1) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])`

3.342.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

method	result	size
derivativedivides	$-\frac{\cos^3(x)}{3} + \frac{2\cos^5(x)}{5} - \frac{\cos^7(x)}{7}$	20
default	$-\frac{\cos^3(x)}{3} + \frac{2\cos^5(x)}{5} - \frac{\cos^7(x)}{7}$	20
risch	$-\frac{5\cos(x)}{64} - \frac{\cos(7x)}{448} + \frac{3\cos(5x)}{320} - \frac{\cos(3x)}{192}$	24
parallelrisch	$\frac{8}{35} - \frac{5\cos(x)}{64} - \frac{\cos(7x)}{448} + \frac{3\cos(5x)}{320} - \frac{\cos(3x)}{192}$	25
norman	$\frac{-\frac{32(\tan^8(\frac{x}{2}))}{3} - \frac{16(\tan^4(\frac{x}{2}))}{5} - \frac{16(\tan^2(\frac{x}{2}))}{15} + \frac{16(\tan^6(\frac{x}{2}))}{3} - \frac{16}{105}}{(1+\tan^2(\frac{x}{2}))^7}$	46

input `int(cos(x)^2*sin(x)^5,x,method=_RETURNVERBOSE)`

output `-1/3*cos(x)^3+2/5*cos(x)^5-1/7*cos(x)^7`

3.342.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \cos^2(x) \sin^5(x) dx = -\frac{1}{7} \cos(x)^7 + \frac{2}{5} \cos(x)^5 - \frac{1}{3} \cos(x)^3$$

input `integrate(cos(x)^2*sin(x)^5,x, algorithm="fricas")`

output `-1/7*cos(x)^7 + 2/5*cos(x)^5 - 1/3*cos(x)^3`

3.342.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \cos^2(x) \sin^5(x) dx = -\frac{\cos^7(x)}{7} + \frac{2 \cos^5(x)}{5} - \frac{\cos^3(x)}{3}$$

input `integrate(cos(x)**2*sin(x)**5,x)`output `-cos(x)**7/7 + 2*cos(x)**5/5 - cos(x)**3/3`**3.342.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \cos^2(x) \sin^5(x) dx = -\frac{1}{7} \cos(x)^7 + \frac{2}{5} \cos(x)^5 - \frac{1}{3} \cos(x)^3$$

input `integrate(cos(x)^2*sin(x)^5,x, algorithm="maxima")`output `-1/7*cos(x)^7 + 2/5*cos(x)^5 - 1/3*cos(x)^3`**3.342.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \cos^2(x) \sin^5(x) dx = -\frac{1}{7} \cos(x)^7 + \frac{2}{5} \cos(x)^5 - \frac{1}{3} \cos(x)^3$$

input `integrate(cos(x)^2*sin(x)^5,x, algorithm="giac")`output `-1/7*cos(x)^7 + 2/5*cos(x)^5 - 1/3*cos(x)^3`

3.342.9 Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \cos^2(x) \sin^5(x) dx = -\frac{\cos(x)^7}{7} + \frac{2 \cos(x)^5}{5} - \frac{\cos(x)^3}{3}$$

input `int(cos(x)^2*sin(x)^5,x)`

output `(2*cos(x)^5)/5 - cos(x)^3/3 - cos(x)^7/7`

3.343 $\int e^{-3x} \cos(4x) dx$

3.343.1 Optimal result	1726
3.343.2 Mathematica [A] (verified)	1726
3.343.3 Rubi [A] (verified)	1727
3.343.4 Maple [A] (verified)	1727
3.343.5 Fricas [A] (verification not implemented)	1728
3.343.6 Sympy [A] (verification not implemented)	1728
3.343.7 Maxima [A] (verification not implemented)	1728
3.343.8 Giac [A] (verification not implemented)	1729
3.343.9 Mupad [B] (verification not implemented)	1729

3.343.1 Optimal result

Integrand size = 10, antiderivative size = 27

$$\int e^{-3x} \cos(4x) dx = -\frac{3}{25}e^{-3x} \cos(4x) + \frac{4}{25}e^{-3x} \sin(4x)$$

output `-3/25*cos(4*x)/exp(3*x)+4/25*sin(4*x)/exp(3*x)`

3.343.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int e^{-3x} \cos(4x) dx = \frac{1}{25}e^{-3x}(-3 \cos(4x) + 4 \sin(4x))$$

input `Integrate[Cos[4*x]/E^(3*x),x]`

output `(-3*Cos[4*x] + 4*Sin[4*x])/(25*E^(3*x))`

3.343.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {4933}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{-3x} \cos(4x) dx$$

$$\downarrow 4933$$

$$\frac{4}{25}e^{-3x} \sin(4x) - \frac{3}{25}e^{-3x} \cos(4x)$$

input `Int[Cos[4*x]/E^(3*x),x]`

output `(-3*Cos[4*x])/(25*E^(3*x)) + (4*Sin[4*x])/(25*E^(3*x))`

3.343.3.1 Defintions of rubi rules used

rule 4933 `Int[Cos[(d_.) + (e_.)*(x_.)]*(F_)^((c_.)*((a_.) + (b_.)*(x_.))), x_Symbol] :=
Simp[b*c*Log[F]*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x
] + Simp[e*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; F
reeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]`

3.343.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result	size
parallelrisch	$\frac{e^{-3x}(-3 \cos(4x) + 4 \sin(4x))}{25}$	20
default	$-\frac{3e^{-3x} \cos(4x)}{25} + \frac{4e^{-3x} \sin(4x)}{25}$	22
norman	$\frac{\left(-\frac{3}{25} + \frac{3 \tan^2(2x)}{25} + \frac{8 \tan(2x)}{25}\right) e^{-3x}}{1 + \tan^2(2x)}$	34
risch	$-\frac{3e^{(-3+4i)x}}{50} - \frac{2ie^{(-3+4i)x}}{25} - \frac{3e^{(-3-4i)x}}{50} + \frac{2ie^{(-3-4i)x}}{25}$	36

input `int(cos(4*x)/exp(3*x),x,method=_RETURNVERBOSE)`

output `1/25*exp(-3*x)*(-3*cos(4*x)+4*sin(4*x))`

3.343.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int e^{-3x} \cos(4x) dx = -\frac{3}{25} \cos(4x) e^{(-3x)} + \frac{4}{25} e^{(-3x)} \sin(4x)$$

input `integrate(cos(4*x)/exp(3*x),x, algorithm="fricas")`

output `-3/25*cos(4*x)*e^(-3*x) + 4/25*e^(-3*x)*sin(4*x)`

3.343.6 Sympy [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int e^{-3x} \cos(4x) dx = \frac{4e^{-3x} \sin(4x)}{25} - \frac{3e^{-3x} \cos(4x)}{25}$$

input `integrate(cos(4*x)/exp(3*x),x)`

output `4*exp(-3*x)*sin(4*x)/25 - 3*exp(-3*x)*cos(4*x)/25`

3.343.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^{-3x} \cos(4x) dx = -\frac{1}{25} (3 \cos(4x) - 4 \sin(4x)) e^{(-3x)}$$

input `integrate(cos(4*x)/exp(3*x),x, algorithm="maxima")`

output `-1/25*(3*cos(4*x) - 4*sin(4*x))*e^(-3*x)`

3.343.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^{-3x} \cos(4x) dx = -\frac{1}{25} (3 \cos(4x) - 4 \sin(4x))e^{-3x}$$

input `integrate(cos(4*x)/exp(3*x),x, algorithm="giac")`output `-1/25*(3*cos(4*x) - 4*sin(4*x))*e^(-3*x)`**3.343.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^{-3x} \cos(4x) dx = -\frac{e^{-3x} (3 \cos(4x) - 4 \sin(4x))}{25}$$

input `int(cos(4*x)*exp(-3*x),x)`output `-(exp(-3*x)*(3*cos(4*x) - 4*sin(4*x)))/25`

3.344 $\int \csc^3\left(\frac{x}{2}\right) dx$

3.344.1 Optimal result	1730
3.344.2 Mathematica [A] (verified)	1730
3.344.3 Rubi [A] (verified)	1731
3.344.4 Maple [A] (verified)	1732
3.344.5 Fracas [B] (verification not implemented)	1732
3.344.6 Sympy [B] (verification not implemented)	1733
3.344.7 Maxima [A] (verification not implemented)	1733
3.344.8 Giac [B] (verification not implemented)	1733
3.344.9 Mupad [B] (verification not implemented)	1734

3.344.1 Optimal result

Integrand size = 8, antiderivative size = 24

$$\int \csc^3\left(\frac{x}{2}\right) dx = -\operatorname{arctanh}\left(\cos\left(\frac{x}{2}\right)\right) - \cot\left(\frac{x}{2}\right) \csc\left(\frac{x}{2}\right)$$

output `-arctanh(cos(1/2*x))-cot(1/2*x)*csc(1/2*x)`

3.344.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.71

$$\int \csc^3\left(\frac{x}{2}\right) dx = -\frac{1}{4} \csc^2\left(\frac{x}{4}\right) - \log\left(\cos\left(\frac{x}{4}\right)\right) + \log\left(\sin\left(\frac{x}{4}\right)\right) + \frac{1}{4} \sec^2\left(\frac{x}{4}\right)$$

input `Integrate[Csc[x/2]^3,x]`

output `-1/4*Csc[x/4]^2 - Log[Cos[x/4]] + Log[Sin[x/4]] + Sec[x/4]^2/4`

3.344.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \csc^3\left(\frac{x}{2}\right) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(\frac{x}{2}\right)^3 dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{1}{2} \int \csc\left(\frac{x}{2}\right) dx - \cot\left(\frac{x}{2}\right) \csc\left(\frac{x}{2}\right) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \csc\left(\frac{x}{2}\right) dx - \cot\left(\frac{x}{2}\right) \csc\left(\frac{x}{2}\right) \\
 & \quad \downarrow \text{4257} \\
 & -\operatorname{arctanh}\left(\cos\left(\frac{x}{2}\right)\right) - \cot\left(\frac{x}{2}\right) \csc\left(\frac{x}{2}\right)
 \end{aligned}$$

input `Int[Csc[x/2]^3,x]`

output `-ArcTanh[Cos[x/2]] - Cot[x/2]*Csc[x/2]`

3.344.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n_, x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.344.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

method	result	size
parallelrisc	$\frac{(\tan^2(\frac{x}{4}))}{4} + \ln(\tan(\frac{x}{4})) - \frac{(\cot^2(\frac{x}{4}))}{4}$	23
derivativdivides	$-\cot(\frac{x}{2}) \csc(\frac{x}{2}) + \ln(\csc(\frac{x}{2}) - \cot(\frac{x}{2}))$	24
default	$-\cot(\frac{x}{2}) \csc(\frac{x}{2}) + \ln(\csc(\frac{x}{2}) - \cot(\frac{x}{2}))$	24
norman	$-\frac{1}{4} + \frac{(\tan^4(\frac{x}{4}))}{4 \tan(\frac{x}{4})^2} + \ln(\tan(\frac{x}{4}))$	24
risc	$\frac{2e^{\frac{3ix}{2}} + 2e^{\frac{ix}{2}}}{(e^{ix} - 1)^2} + \ln(e^{\frac{ix}{2}} - 1) - \ln(e^{\frac{ix}{2}} + 1)$	42

input `int(csc(1/2*x)^3,x,method=_RETURNVERBOSE)`

output `1/4*tan(1/4*x)^2+ln(tan(1/4*x))-1/4*cot(1/4*x)^2`

3.344.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. $2(18) = 36$.

Time = 0.25 (sec) , antiderivative size = 56, normalized size of antiderivative = 2.33

$$\int \csc^3\left(\frac{x}{2}\right) dx = \frac{\left(\cos\left(\frac{1}{2}x\right)^2 - 1\right) \log\left(\frac{1}{2}\cos\left(\frac{1}{2}x\right) + \frac{1}{2}\right) - \left(\cos\left(\frac{1}{2}x\right)^2 - 1\right) \log\left(-\frac{1}{2}\cos\left(\frac{1}{2}x\right) + \frac{1}{2}\right) - 2\cos\left(\frac{1}{2}x\right)}{2\left(\cos\left(\frac{1}{2}x\right)^2 - 1\right)}$$

input `integrate(csc(1/2*x)^3,x, algorithm="fricas")`

output `-1/2*((cos(1/2*x)^2 - 1)*log(1/2*cos(1/2*x) + 1/2) - (cos(1/2*x)^2 - 1)*log(-1/2*cos(1/2*x) + 1/2) - 2*cos(1/2*x))/(cos(1/2*x)^2 - 1)`

3.344.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 36 vs. $2(17) = 34$.

Time = 0.06 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.50

$$\int \csc^3\left(\frac{x}{2}\right) dx = \frac{\log\left(\cos\left(\frac{x}{2}\right) - 1\right)}{2} - \frac{\log\left(\cos\left(\frac{x}{2}\right) + 1\right)}{2} + \frac{2 \cos\left(\frac{x}{2}\right)}{2 \cos^2\left(\frac{x}{2}\right) - 2}$$

input `integrate(csc(1/2*x)**3,x)`

output `log(cos(x/2) - 1)/2 - log(cos(x/2) + 1)/2 + 2*cos(x/2)/(2*cos(x/2)**2 - 2)`

3.344.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.42

$$\int \csc^3\left(\frac{x}{2}\right) dx = \frac{\cos\left(\frac{1}{2}x\right)}{\cos\left(\frac{1}{2}x\right)^2 - 1} - \frac{1}{2} \log\left(\cos\left(\frac{1}{2}x\right) + 1\right) + \frac{1}{2} \log\left(\cos\left(\frac{1}{2}x\right) - 1\right)$$

input `integrate(csc(1/2*x)^3,x, algorithm="maxima")`

output `cos(1/2*x)/(cos(1/2*x)^2 - 1) - 1/2*log(cos(1/2*x) + 1) + 1/2*log(cos(1/2*x) - 1)`

3.344.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 70 vs. $2(18) = 36$.

Time = 0.28 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.92

$$\int \csc^3\left(\frac{x}{2}\right) dx = -\frac{\left(\frac{2(\cos(\frac{1}{2}x)-1)}{\cos(\frac{1}{2}x)+1} - 1\right)(\cos(\frac{1}{2}x) + 1)}{4(\cos(\frac{1}{2}x) - 1)} - \frac{\cos(\frac{1}{2}x) - 1}{4(\cos(\frac{1}{2}x) + 1)} + \frac{1}{2} \log\left(-\frac{\cos(\frac{1}{2}x) - 1}{\cos(\frac{1}{2}x) + 1}\right)$$

input `integrate(csc(1/2*x)^3,x, algorithm="giac")`

output `-1/4*(2*(cos(1/2*x) - 1)/(cos(1/2*x) + 1) - 1)*(cos(1/2*x) + 1)/(cos(1/2*x) - 1) - 1/4*(cos(1/2*x) - 1)/(cos(1/2*x) + 1) + 1/2*log(-(cos(1/2*x) - 1)/(cos(1/2*x) + 1))`

3.344.9 Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.75

$$\int \csc^3\left(\frac{x}{2}\right) dx = \ln\left(\tan\left(\frac{x}{4}\right)\right) - \frac{\cos\left(\frac{x}{2}\right)}{\sin\left(\frac{x}{2}\right)^2}$$

input `int(1/sin(x/2)^3,x)`

output `log(tan(x/4)) - cos(x/2)/sin(x/2)^2`

3.345 $\int \frac{\sqrt{-1+9x^2}}{x^2} dx$

3.345.1 Optimal result	1735
3.345.2 Mathematica [A] (verified)	1735
3.345.3 Rubi [A] (verified)	1736
3.345.4 Maple [A] (verified)	1737
3.345.5 Fricas [A] (verification not implemented)	1737
3.345.6 Sympy [A] (verification not implemented)	1738
3.345.7 Maxima [A] (verification not implemented)	1738
3.345.8 Giac [A] (verification not implemented)	1738
3.345.9 Mupad [B] (verification not implemented)	1739

3.345.1 Optimal result

Integrand size = 15, antiderivative size = 34

$$\int \frac{\sqrt{-1+9x^2}}{x^2} dx = -\frac{\sqrt{-1+9x^2}}{x} + 3\operatorname{arctanh}\left(\frac{3x}{\sqrt{-1+9x^2}}\right)$$

output `3*arctanh(3*x/(9*x^2-1)^(1/2))- (9*x^2-1)^(1/2)/x`

3.345.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.03

$$\int \frac{\sqrt{-1+9x^2}}{x^2} dx = -\frac{\sqrt{-1+9x^2}}{x} - 3\log\left(-3x + \sqrt{-1+9x^2}\right)$$

input `Integrate[Sqrt[-1 + 9*x^2]/x^2,x]`

output `-(Sqrt[-1 + 9*x^2]/x) - 3*Log[-3*x + Sqrt[-1 + 9*x^2]]`

3.345.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {247, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{9x^2 - 1}}{x^2} dx \\
 & \quad \downarrow \text{247} \\
 & 9 \int \frac{1}{\sqrt{9x^2 - 1}} dx - \frac{\sqrt{9x^2 - 1}}{x} \\
 & \quad \downarrow \text{224} \\
 & 9 \int \frac{1}{1 - \frac{9x^2}{9x^2 - 1}} d \frac{x}{\sqrt{9x^2 - 1}} - \frac{\sqrt{9x^2 - 1}}{x} \\
 & \quad \downarrow \text{219} \\
 & 3 \operatorname{arctanh} \left(\frac{3x}{\sqrt{9x^2 - 1}} \right) - \frac{\sqrt{9x^2 - 1}}{x}
 \end{aligned}$$

input `Int[Sqrt[-1 + 9*x^2]/x^2,x]`

output `-(Sqrt[-1 + 9*x^2]/x) + 3*ArcTanh[(3*x)/Sqrt[-1 + 9*x^2]]`

3.345.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

```
rule 247 Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^p/(c*(m + 1))), x] - Simp[2*b*(p/(c^2*(m + 1))) Int[(c*x)^(m + 2)*(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && GtQ[p, 0] && LtQ[m, -1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

3.345.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.94

method	result	size
trager	$-\frac{\sqrt{9x^2-1}}{x} - 3 \ln(\sqrt{9x^2-1} - 3x)$	32
risch	$-\frac{\sqrt{9x^2-1}}{x} + \ln(\sqrt{9}x + \sqrt{9x^2-1})\sqrt{9}$	36
default	$\frac{(9x^2-1)^{\frac{3}{2}}}{x} - 9x\sqrt{9x^2-1} + \ln(\sqrt{9}x + \sqrt{9x^2-1})\sqrt{9}$	47
meijerg	$-\frac{3i\sqrt{\text{signum}(9x^2-1)}\left(-\frac{4i\sqrt{\pi}\sqrt{-9x^2+1}}{3x} - 4i\sqrt{\pi}\arcsin(3x)\right)}{4\sqrt{\pi}\sqrt{-\text{signum}(9x^2-1)}}$	58
pseudoelliptic	$\frac{3 \ln\left(\frac{\sqrt{9x^2-1}+3x}{x}\right)x - 3 \ln\left(\frac{\sqrt{9x^2-1}-3x}{x}\right)x - 2\sqrt{9x^2-1}}{2x}$	60

```
input int((9*x^2-1)^(1/2)/x^2,x,method=_RETURNVERBOSE)
```

```
output -(9*x^2-1)^(1/2)/x-3*ln((9*x^2-1)^(1/2)-3*x)
```

3.345.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.03

$$\int \frac{\sqrt{-1+9x^2}}{x^2} dx = -\frac{3x \log(-3x + \sqrt{9x^2-1}) + 3x + \sqrt{9x^2-1}}{x}$$

```
input integrate((9*x^2-1)^(1/2)/x^2,x, algorithm="fricas")
```

```
output -(3*x*log(-3*x + sqrt(9*x^2 - 1)) + 3*x + sqrt(9*x^2 - 1))/x
```

3.345.6 Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.79

$$\int \frac{\sqrt{-1+9x^2}}{x^2} dx = 3 \log(3x + \sqrt{9x^2 - 1}) - \frac{\sqrt{9x^2 - 1}}{x}$$

input `integrate((9*x**2-1)**(1/2)/x**2,x)`output `3*log(3*x + sqrt(9*x**2 - 1)) - sqrt(9*x**2 - 1)/x`**3.345.7 Maxima [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.97

$$\int \frac{\sqrt{-1+9x^2}}{x^2} dx = -\frac{\sqrt{9x^2-1}}{x} + 3 \log(18x + 6\sqrt{9x^2-1})$$

input `integrate((9*x^2-1)^(1/2)/x^2,x, algorithm="maxima")`output `-sqrt(9*x^2 - 1)/x + 3*log(18*x + 6*sqrt(9*x^2 - 1))`**3.345.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.29

$$\int \frac{\sqrt{-1+9x^2}}{x^2} dx = -\frac{6}{(3x - \sqrt{9x^2 - 1})^2 + 1} - \frac{3}{2} \log\left(\left(3x - \sqrt{9x^2 - 1}\right)^2\right)$$

input `integrate((9*x^2-1)^(1/2)/x^2,x, algorithm="giac")`output `-6/((3*x - sqrt(9*x^2 - 1))^2 + 1) - 3/2*log((3*x - sqrt(9*x^2 - 1))^2)`

3.345.9 Mupad [B] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.94

$$\int \frac{\sqrt{-1+9x^2}}{x^2} dx = -\frac{\left(\frac{3x \operatorname{asin}(3x)}{\sqrt{1-9x^2}} + 1\right) \sqrt{9x^2-1}}{x}$$

input `int((9*x^2 - 1)^(1/2)/x^2,x)`output `-(((3*x*asin(3*x))/(1 - 9*x^2)^(1/2) + 1)*(9*x^2 - 1)^(1/2))/x`

3.346 $\int \frac{\sqrt{4-3x^2}}{x} dx$

3.346.1 Optimal result	1740
3.346.2 Mathematica [A] (verified)	1740
3.346.3 Rubi [A] (verified)	1741
3.346.4 Maple [A] (verified)	1742
3.346.5 Fricas [A] (verification not implemented)	1743
3.346.6 Sympy [C] (verification not implemented)	1743
3.346.7 Maxima [A] (verification not implemented)	1743
3.346.8 Giac [A] (verification not implemented)	1744
3.346.9 Mupad [B] (verification not implemented)	1744

3.346.1 Optimal result

Integrand size = 15, antiderivative size = 30

$$\int \frac{\sqrt{4-3x^2}}{x} dx = \sqrt{4-3x^2} - 2\operatorname{arctanh}\left(\frac{1}{2}\sqrt{4-3x^2}\right)$$

output `-2*arctanh(1/2*(-3*x^2+4)^(1/2))+(-3*x^2+4)^(1/2)`

3.346.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{4-3x^2}}{x} dx = \sqrt{4-3x^2} - 2\operatorname{arctanh}\left(\frac{1}{2}\sqrt{4-3x^2}\right)$$

input `Integrate[Sqrt[4 - 3*x^2]/x,x]`

output `Sqrt[4 - 3*x^2] - 2*ArcTanh[Sqrt[4 - 3*x^2]/2]`

3.346.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.20, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {243, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{4-3x^2}}{x} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \int \frac{\sqrt{4-3x^2}}{x^2} dx^2 \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2} \left(4 \int \frac{1}{x^2 \sqrt{4-3x^2}} dx^2 + 2\sqrt{4-3x^2} \right) \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{2} \left(2\sqrt{4-3x^2} - \frac{8}{3} \int \frac{1}{\frac{4}{3} - \frac{x^4}{3}} d\sqrt{4-3x^2} \right) \\
 & \quad \downarrow \text{219} \\
 & \frac{1}{2} \left(2\sqrt{4-3x^2} - 4\operatorname{arctanh} \left(\frac{1}{2} \sqrt{4-3x^2} \right) \right)
 \end{aligned}$$

input `Int[Sqrt[4 - 3*x^2]/x,x]`

output `(2*Sqrt[4 - 3*x^2] - 4*ArcTanh[Sqrt[4 - 3*x^2]/2])/2`

3.346.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
 ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
 Q[a, 0] || LtQ[b, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[In
 t[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I
 ntegerQ[(m - 1)/2]`

3.346.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.83

method	result	size
default	$\sqrt{-3x^2 + 4} - 2 \operatorname{arctanh}\left(\frac{2}{\sqrt{-3x^2 + 4}}\right)$	25
trager	$\sqrt{-3x^2 + 4} - 2 \ln\left(\frac{\sqrt{-3x^2 + 4} + 2}{x}\right)$	29
pseudoelliptic	$\sqrt{-3x^2 + 4} + \ln(\sqrt{-3x^2 + 4} - 2) - \ln(\sqrt{-3x^2 + 4} + 2)$	37
meijerg	$-\frac{-2(2-4\ln(2)+2\ln(x)+\ln(3)+i\pi)\sqrt{\pi}+4\sqrt{\pi}-4\sqrt{\pi}\sqrt{1-\frac{3x^2}{4}}+4\sqrt{\pi}\ln\left(\frac{1}{2}+\frac{\sqrt{1-\frac{3x^2}{4}}}{2}\right)}{2\sqrt{\pi}}$	66

input `int((-3*x^2+4)^(1/2)/x,x,method=_RETURNVERBOSE)`

output `(-3*x^2+4)^(1/2)-2*arctanh(2/(-3*x^2+4)^(1/2))`

3.346.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{4-3x^2}}{x} dx = \sqrt{-3x^2+4} + 2 \log\left(\frac{\sqrt{-3x^2+4}-2}{x}\right)$$

input `integrate((-3*x^2+4)^(1/2)/x,x, algorithm="fricas")`

output `sqrt(-3*x^2 + 4) + 2*log((sqrt(-3*x^2 + 4) - 2)/x)`

3.346.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.84 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.43

$$\int \frac{\sqrt{4-3x^2}}{x} dx = \begin{cases} i\sqrt{3x^2-4} - 2\log(x) + \log(x^2) + 2i \operatorname{asin}\left(\frac{2\sqrt{3}}{3x}\right) & \text{for } |x^2| > \frac{4}{3} \\ \sqrt{4-3x^2} + \log(x^2) - 2\log\left(\sqrt{1-\frac{3x^2}{4}} + 1\right) & \text{otherwise} \end{cases}$$

input `integrate((-3*x**2+4)**(1/2)/x,x)`

output `Piecewise((I*sqrt(3*x**2 - 4) - 2*log(x) + log(x**2) + 2*I*asin(2*sqrt(3)/(3*x)), Abs(x**2) > 4/3), (sqrt(4 - 3*x**2) + log(x**2) - 2*log(sqrt(1 - 3*x**2/4) + 1), True))`

3.346.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.17

$$\int \frac{\sqrt{4-3x^2}}{x} dx = \sqrt{-3x^2+4} - 2 \log\left(\frac{4\sqrt{-3x^2+4}}{|x|} + \frac{8}{|x|}\right)$$

input `integrate((-3*x^2+4)^(1/2)/x,x, algorithm="maxima")`

output `sqrt(-3*x^2 + 4) - 2*log(4*sqrt(-3*x^2 + 4)/abs(x) + 8/abs(x))`

3.346.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.27

$$\int \frac{\sqrt{4-3x^2}}{x} dx = \sqrt{-3x^2+4} - \log\left(\sqrt{-3x^2+4}+2\right) + \log\left(-\sqrt{-3x^2+4}+2\right)$$

input `integrate((-3*x^2+4)^(1/2)/x,x, algorithm="giac")`output `sqrt(-3*x^2 + 4) - log(sqrt(-3*x^2 + 4) + 2) + log(-sqrt(-3*x^2 + 4) + 2)`**3.346.9 Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.23

$$\int \frac{\sqrt{4-3x^2}}{x} dx = 2 \ln\left(\sqrt{\frac{4}{3x^2}-1} - \frac{2\sqrt{3}\sqrt{\frac{1}{x^2}}}{3}\right) + \sqrt{3}\sqrt{\frac{4}{3}-x^2}$$

input `int((4 - 3*x^2)^(1/2)/x,x)`output `2*log((4/(3*x^2) - 1)^(1/2) - (2*3^(1/2)*(1/x^2)^(1/2))/3) + 3^(1/2)*(4/3 - x^2)^(1/2)`

3.347 $\int e^{3x} x^2 dx$

3.347.1 Optimal result	1745
3.347.2 Mathematica [A] (verified)	1745
3.347.3 Rubi [A] (verified)	1746
3.347.4 Maple [A] (verified)	1747
3.347.5 Fricas [A] (verification not implemented)	1747
3.347.6 Sympy [A] (verification not implemented)	1748
3.347.7 Maxima [A] (verification not implemented)	1748
3.347.8 Giac [A] (verification not implemented)	1748
3.347.9 Mupad [B] (verification not implemented)	1749

3.347.1 Optimal result

Integrand size = 9, antiderivative size = 32

$$\int e^{3x} x^2 dx = \frac{2e^{3x}}{27} - \frac{2}{9}e^{3x}x + \frac{1}{3}e^{3x}x^2$$

output `2/27*exp(3*x)-2/9*exp(3*x)*x+1/3*exp(3*x)*x^2`

3.347.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.59

$$\int e^{3x} x^2 dx = \frac{1}{27}e^{3x}(2 - 6x + 9x^2)$$

input `Integrate[E^(3*x)*x^2,x]`

output `(E^(3*x)*(2 - 6*x + 9*x^2))/27`

3.347.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.16, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2607, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^{3x} x^2 dx \\ & \quad \downarrow \text{2607} \\ & \frac{1}{3} e^{3x} x^2 - \frac{2}{3} \int e^{3x} x dx \\ & \quad \downarrow \text{2607} \\ & \frac{1}{3} e^{3x} x^2 - \frac{2}{3} \left(\frac{1}{3} e^{3x} x - \frac{\int e^{3x} dx}{3} \right) \\ & \quad \downarrow \text{2624} \\ & \frac{1}{3} e^{3x} x^2 - \frac{2}{3} \left(\frac{1}{3} e^{3x} x - \frac{e^{3x}}{9} \right) \end{aligned}$$

input `Int[E^(3*x)*x^2,x]`

output `(E^(3*x)*x^2)/3 - (2*(-1/9*E^(3*x) + (E^(3*x)*x)/3))/3`

3.347.3.1 Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2624 `Int[((F_)^(v_))^(n_.), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`

3.347.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.50

method	result	size
risch	$\left(\frac{1}{3}x^2 - \frac{2}{9}x + \frac{2}{27}\right)e^{3x}$	16
gospers	$\frac{(9x^2-6x+2)e^{3x}}{27}$	17
meijerg	$-\frac{2}{27} + \frac{(27x^2-18x+6)e^{3x}}{81}$	19
derivativedivides	$\frac{2e^{3x}}{27} - \frac{2e^{3x}x}{9} + \frac{e^{3x}x^2}{3}$	24
default	$\frac{2e^{3x}}{27} - \frac{2e^{3x}x}{9} + \frac{e^{3x}x^2}{3}$	24
norman	$\frac{2e^{3x}}{27} - \frac{2e^{3x}x}{9} + \frac{e^{3x}x^2}{3}$	24
parallelrisch	$\frac{2e^{3x}}{27} - \frac{2e^{3x}x}{9} + \frac{e^{3x}x^2}{3}$	24
parts	$\frac{2e^{3x}}{27} - \frac{2e^{3x}x}{9} + \frac{e^{3x}x^2}{3}$	24

input `int(exp(3*x)*x^2,x,method=_RETURNVERBOSE)`output `(1/3*x^2-2/9*x+2/27)*exp(3*x)`**3.347.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.50

$$\int e^{3x}x^2 dx = \frac{1}{27}(9x^2 - 6x + 2)e^{(3x)}$$

input `integrate(exp(3*x)*x^2,x, algorithm="fricas")`output `1/27*(9*x^2 - 6*x + 2)*e^(3*x)`

3.347.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.47

$$\int e^{3x} x^2 dx = \frac{(9x^2 - 6x + 2) e^{3x}}{27}$$

input `integrate(exp(3*x)*x**2,x)`output `(9*x**2 - 6*x + 2)*exp(3*x)/27`**3.347.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.50

$$\int e^{3x} x^2 dx = \frac{1}{27} (9x^2 - 6x + 2) e^{(3x)}$$

input `integrate(exp(3*x)*x^2,x, algorithm="maxima")`output `1/27*(9*x^2 - 6*x + 2)*e^(3*x)`**3.347.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.50

$$\int e^{3x} x^2 dx = \frac{1}{27} (9x^2 - 6x + 2) e^{(3x)}$$

input `integrate(exp(3*x)*x^2,x, algorithm="giac")`output `1/27*(9*x^2 - 6*x + 2)*e^(3*x)`

3.347.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.50

$$\int e^{3x} x^2 dx = \frac{e^{3x} (9x^2 - 6x + 2)}{27}$$

input `int(x^2*exp(3*x),x)`

output `(exp(3*x)*(9*x^2 - 6*x + 2))/27`

3.348 $\int \frac{\cos(x) \sin(x)}{\sqrt{1+\sin(x)}} dx$

3.348.1 Optimal result	1750
3.348.2 Mathematica [A] (verified)	1750
3.348.3 Rubi [A] (verified)	1751
3.348.4 Maple [A] (verified)	1752
3.348.5 Fricas [A] (verification not implemented)	1752
3.348.6 Sympy [A] (verification not implemented)	1753
3.348.7 Maxima [A] (verification not implemented)	1753
3.348.8 Giac [B] (verification not implemented)	1753
3.348.9 Mupad [B] (verification not implemented)	1754

3.348.1 Optimal result

Integrand size = 13, antiderivative size = 23

$$\int \frac{\cos(x) \sin(x)}{\sqrt{1 + \sin(x)}} dx = -2\sqrt{1 + \sin(x)} + \frac{2}{3}(1 + \sin(x))^{3/2}$$

output `2/3*(1+sin(x))^(3/2)-2*(1+sin(x))^(1/2)`

3.348.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.35

$$\int \frac{\cos(x) \sin(x)}{\sqrt{1 + \sin(x)}} dx = \frac{2(\cos(\frac{x}{2}) + \sin(\frac{x}{2}))^2 (-2 + \sin(x))}{3\sqrt{1 + \sin(x)}}$$

input `Integrate[(Cos[x]*Sin[x])/Sqrt[1 + Sin[x]],x]`

output `(2*(Cos[x/2] + Sin[x/2])^2*(-2 + Sin[x]))/(3*Sqrt[1 + Sin[x]])`

3.348.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3042, 3312, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin(x) \cos(x)}{\sqrt{\sin(x) + 1}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(x) \cos(x)}{\sqrt{\sin(x) + 1}} dx \\
 & \quad \downarrow \text{3312} \\
 & \int \frac{\sin(x)}{\sqrt{\sin(x) + 1}} d \sin(x) \\
 & \quad \downarrow \text{53} \\
 & \int \left(\sqrt{\sin(x) + 1} - \frac{1}{\sqrt{\sin(x) + 1}} \right) d \sin(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{2}{3}(\sin(x) + 1)^{3/2} - 2\sqrt{\sin(x) + 1}
 \end{aligned}$$

input `Int[(Cos[x]*Sin[x])/Sqrt[1 + Sin[x]],x]`

output `-2*Sqrt[1 + Sin[x]] + (2*(1 + Sin[x])^(3/2))/3`

3.348.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.348. $\int \frac{\cos(x) \sin(x)}{\sqrt{1+\sin(x)}} dx$


```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3312 Int[cos[(e_.) + (f_.)*(x_.)]*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.)*((
c_.) + (d_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[1/(b*f) Subst[Int[(a + x)^m*(c + (d/b)*x)^n, x], x, b*Sin[e + f*x]], x] /; FreeQ[{a,
b, c, d, e, f, m, n}, x]
```

3.348.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

method	result	size
derivativedivides	$\frac{2(\sin(x)+1)^{\frac{3}{2}}}{3} - 2\sqrt{\sin(x)+1}$	18
default	$\frac{2(\sin(x)+1)^{\frac{3}{2}}}{3} - 2\sqrt{\sin(x)+1}$	18

```
input int(cos(x)*sin(x)/(sin(x)+1)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 2/3*(sin(x)+1)^(3/2)-2*(sin(x)+1)^(1/2)
```

3.348.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.52

$$\int \frac{\cos(x) \sin(x)}{\sqrt{1 + \sin(x)}} dx = \frac{2}{3} \sqrt{\sin(x) + 1} (\sin(x) - 2)$$

```
input integrate(cos(x)*sin(x)/(1+sin(x))^(1/2),x, algorithm="fracas")
```

```
output 2/3*sqrt(sin(x) + 1)*(sin(x) - 2)
```

3.348.6 Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.13

$$\int \frac{\cos(x) \sin(x)}{\sqrt{1 + \sin(x)}} dx = \frac{2\sqrt{\sin(x) + 1} \sin(x)}{3} - \frac{4\sqrt{\sin(x) + 1}}{3}$$

input `integrate(cos(x)*sin(x)/(1+sin(x))**(1/2),x)`

output `2*sqrt(sin(x) + 1)*sin(x)/3 - 4*sqrt(sin(x) + 1)/3`

3.348.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{\cos(x) \sin(x)}{\sqrt{1 + \sin(x)}} dx = \frac{2}{3} (\sin(x) + 1)^{\frac{3}{2}} - 2\sqrt{\sin(x) + 1}$$

input `integrate(cos(x)*sin(x)/(1+sin(x))^(1/2),x, algorithm="maxima")`

output `2/3*(sin(x) + 1)^(3/2) - 2*sqrt(sin(x) + 1)`

3.348.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. $2(17) = 34$.

Time = 0.28 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.83

$$\int \frac{\cos(x) \sin(x)}{\sqrt{1 + \sin(x)}} dx = \frac{2 \left(2\sqrt{2} \cos\left(-\frac{1}{4}\pi + \frac{1}{2}x\right)^3 - 3\sqrt{2} \cos\left(-\frac{1}{4}\pi + \frac{1}{2}x\right) \right)}{3 \operatorname{sgn}\left(\cos\left(-\frac{1}{4}\pi + \frac{1}{2}x\right)\right)}$$

input `integrate(cos(x)*sin(x)/(1+sin(x))^(1/2),x, algorithm="giac")`

output `2/3*(2*sqrt(2)*cos(-1/4*pi + 1/2*x)^3 - 3*sqrt(2)*cos(-1/4*pi + 1/2*x))/sgn(cos(-1/4*pi + 1/2*x))`

3.348.9 Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.52

$$\int \frac{\cos(x) \sin(x)}{\sqrt{1 + \sin(x)}} dx = \frac{2 \sqrt{\sin(x) + 1} (\sin(x) - 2)}{3}$$

input `int((cos(x)*sin(x))/(sin(x) + 1)^(1/2),x)`

output `(2*(sin(x) + 1)^(1/2)*(sin(x) - 2))/3`

3.349 $\int x \arcsin(x^2) dx$

3.349.1 Optimal result	1755
3.349.2 Mathematica [A] (verified)	1755
3.349.3 Rubi [A] (verified)	1756
3.349.4 Maple [A] (verified)	1757
3.349.5 Fricas [A] (verification not implemented)	1757
3.349.6 Sympy [A] (verification not implemented)	1757
3.349.7 Maxima [A] (verification not implemented)	1758
3.349.8 Giac [A] (verification not implemented)	1758
3.349.9 Mupad [B] (verification not implemented)	1758

3.349.1 Optimal result

Integrand size = 6, antiderivative size = 27

$$\int x \arcsin(x^2) dx = \frac{\sqrt{1-x^4}}{2} + \frac{1}{2}x^2 \arcsin(x^2)$$

output `1/2*x^2*arcsin(x^2)+1/2*(-x^4+1)^(1/2)`

3.349.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

$$\int x \arcsin(x^2) dx = \frac{1}{2}(\sqrt{1-x^4} + x^2 \arcsin(x^2))$$

input `Integrate[x*ArcSin[x^2],x]`

output `(Sqrt[1 - x^4] + x^2*ArcSin[x^2])/2`

3.349.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {7266, 5130, 241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \arcsin(x^2) dx \\ & \quad \downarrow \text{7266} \\ & \frac{1}{2} \int \arcsin(x^2) dx^2 \\ & \quad \downarrow \text{5130} \\ & \frac{1}{2} \left(x^2 \arcsin(x^2) - \int \frac{x^2}{\sqrt{1-x^4}} dx^2 \right) \\ & \quad \downarrow \text{241} \\ & \frac{1}{2} \left(x^2 \arcsin(x^2) + \sqrt{1-x^4} \right) \end{aligned}$$

input `Int[x*ArcSin[x^2],x]`

output `(Sqrt[1 - x^4] + x^2*ArcSin[x^2])/2`

3.349.3.1 Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

rule 5130 `Int[((a_) + ArcSin[(c_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[x*(a + b*ArcSin[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcSin[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

rule 7266 `Int[(u_)*(x_)^(m_), x_Symbol] := Simp[1/(m + 1) Subst[Int[SubstFor[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^(m + 1), u, x]`

3.349.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

method	result	size
derivativeldivides	$\frac{x^2 \arcsin(x^2)}{2} + \frac{\sqrt{-x^4+1}}{2}$	22
default	$\frac{x^2 \arcsin(x^2)}{2} + \frac{\sqrt{-x^4+1}}{2}$	22
parts	$\frac{x^2 \arcsin(x^2)}{2} + \frac{\sqrt{-x^4+1}}{2}$	22

input `int(x*arcsin(x^2),x,method=_RETURNVERBOSE)`output `1/2*x^2*arcsin(x^2)+1/2*(-x^4+1)^(1/2)`**3.349.5 Fricas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int x \arcsin(x^2) dx = \frac{1}{2} x^2 \arcsin(x^2) + \frac{1}{2} \sqrt{-x^4 + 1}$$

input `integrate(x*arcsin(x^2),x, algorithm="fricas")`output `1/2*x^2*arcsin(x^2) + 1/2*sqrt(-x^4 + 1)`**3.349.6 Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int x \arcsin(x^2) dx = \frac{x^2 \operatorname{asin}(x^2)}{2} + \frac{\sqrt{1-x^4}}{2}$$

input `integrate(x*asin(x**2),x)`output `x**2*asin(x**2)/2 + sqrt(1 - x**4)/2`

3.349.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int x \arcsin(x^2) dx = \frac{1}{2} x^2 \arcsin(x^2) + \frac{1}{2} \sqrt{-x^4 + 1}$$

input `integrate(x*arcsin(x^2),x, algorithm="maxima")`output `1/2*x^2*arcsin(x^2) + 1/2*sqrt(-x^4 + 1)`**3.349.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int x \arcsin(x^2) dx = \frac{1}{2} x^2 \arcsin(x^2) + \frac{1}{2} \sqrt{-x^4 + 1}$$

input `integrate(x*arcsin(x^2),x, algorithm="giac")`output `1/2*x^2*arcsin(x^2) + 1/2*sqrt(-x^4 + 1)`**3.349.9 Mupad [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int x \arcsin(x^2) dx = \frac{x^2 \operatorname{asin}(x^2)}{2} + \frac{\sqrt{1-x^4}}{2}$$

input `int(x*asin(x^2),x)`output `(x^2*asin(x^2))/2 + (1 - x^4)^(1/2)/2`

3.350 $\int x^3 \arcsin(x^2) dx$

3.350.1 Optimal result	1759
3.350.2 Mathematica [A] (verified)	1759
3.350.3 Rubi [A] (verified)	1760
3.350.4 Maple [A] (verified)	1761
3.350.5 Fricas [A] (verification not implemented)	1762
3.350.6 Sympy [A] (verification not implemented)	1762
3.350.7 Maxima [A] (verification not implemented)	1762
3.350.8 Giac [A] (verification not implemented)	1763
3.350.9 Mupad [B] (verification not implemented)	1763

3.350.1 Optimal result

Integrand size = 8, antiderivative size = 38

$$\int x^3 \arcsin(x^2) dx = \frac{1}{8}x^2\sqrt{1-x^4} - \frac{\arcsin(x^2)}{8} + \frac{1}{4}x^4 \arcsin(x^2)$$

output `-1/8*arcsin(x^2)+1/4*x^4*arcsin(x^2)+1/8*x^2*(-x^4+1)^(1/2)`

3.350.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int x^3 \arcsin(x^2) dx = \frac{1}{8} \left(x^2 \sqrt{1-x^4} + (-1+2x^4) \arcsin(x^2) \right)$$

input `Integrate[x^3*ArcSin[x^2],x]`

output `(x^2*sqrt[1 - x^4] + (-1 + 2*x^4)*ArcSin[x^2])/8`

3.350.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.13, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {5341, 27, 807, 262, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \arcsin(x^2) dx \\
 & \quad \downarrow \text{5341} \\
 & \frac{1}{4}x^4 \arcsin(x^2) - \frac{1}{4} \int \frac{2x^5}{\sqrt{1-x^4}} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{4}x^4 \arcsin(x^2) - \frac{1}{2} \int \frac{x^5}{\sqrt{1-x^4}} dx \\
 & \quad \downarrow \text{807} \\
 & \frac{1}{4}x^4 \arcsin(x^2) - \frac{1}{4} \int \frac{x^4}{\sqrt{1-x^4}} dx^2 \\
 & \quad \downarrow \text{262} \\
 & \frac{1}{4} \left(\frac{1}{2}x^2 \sqrt{1-x^4} - \frac{1}{2} \int \frac{1}{\sqrt{1-x^4}} dx^2 \right) + \frac{1}{4}x^4 \arcsin(x^2) \\
 & \quad \downarrow \text{223} \\
 & \frac{1}{4}x^4 \arcsin(x^2) + \frac{1}{4} \left(\frac{1}{2}x^2 \sqrt{1-x^4} - \frac{\arcsin(x^2)}{2} \right)
 \end{aligned}$$

input `Int[x^3*ArcSin[x^2],x]`

output `((x^2*Sqrt[1 - x^4])/2 - ArcSin[x^2]/2)/4 + (x^4*ArcSin[x^2])/4`

3.350.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 262 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m-1)*((a+b*x^2)^(p+1)/(b*(m+2*p+1))), x] - Simp[a*c^2*((m-1)/(b*(m+2*p+1)) Int[(c*x)^(m-2)*(a+b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2-1] && NeQ[m+2*p+1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 807 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m+1, n]}, Simp[1/k Subst[Int[x^((m+1)/k-1)*(a+b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`
- rule 5341 `Int[((a_) + ArcSin[u_]*(b_))*((c_) + (d_)*(x_)^(m_)), x_Symbol] := Simp[(c+d*x)^(m+1)*((a+b*ArcSin[u])/(d*(m+1))), x] - Simp[b/(d*(m+1)) Int[SimplifyIntegrand[(c+d*x)^(m+1)*(D[u, x]/Sqrt[1-u^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c+d*x)^(m+1), u, x] && !FunctionOfExponentialQ[u, x]`

3.350.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$-\frac{\arcsin(x^2)}{8} + \frac{x^4 \arcsin(x^2)}{4} + \frac{x^2 \sqrt{-x^4+1}}{8}$	31
default	$-\frac{\arcsin(x^2)}{8} + \frac{x^4 \arcsin(x^2)}{4} + \frac{x^2 \sqrt{-x^4+1}}{8}$	31
parts	$-\frac{\arcsin(x^2)}{8} + \frac{x^4 \arcsin(x^2)}{4} + \frac{x^2 \sqrt{-x^4+1}}{8}$	31

input `int(x^3*arcsin(x^2),x,method=_RETURNVERBOSE)`

output `-1/8*arcsin(x^2)+1/4*x^4*arcsin(x^2)+1/8*x^2*(-x^4+1)^(1/2)`

3.350.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.74

$$\int x^3 \arcsin(x^2) dx = \frac{1}{8} \sqrt{-x^4 + 1} x^2 + \frac{1}{8} (2x^4 - 1) \arcsin(x^2)$$

input `integrate(x^3*arcsin(x^2),x, algorithm="fricas")`

output `1/8*sqrt(-x^4 + 1)*x^2 + 1/8*(2*x^4 - 1)*arcsin(x^2)`

3.350.6 Sympy [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.76

$$\int x^3 \arcsin(x^2) dx = \frac{x^4 \operatorname{asin}(x^2)}{4} + \frac{x^2 \sqrt{1-x^4}}{8} - \frac{\operatorname{asin}(x^2)}{8}$$

input `integrate(x**3*asin(x**2),x)`

output `x**4*asin(x**2)/4 + x**2*sqrt(1 - x**4)/8 - asin(x**2)/8`

3.350.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.39

$$\int x^3 \arcsin(x^2) dx = \frac{1}{4} x^4 \arcsin(x^2) - \frac{\sqrt{-x^4 + 1}}{8 x^2 \left(\frac{x^4 - 1}{x^4} - 1\right)} + \frac{1}{8} \arctan\left(\frac{\sqrt{-x^4 + 1}}{x^2}\right)$$

input `integrate(x^3*arcsin(x^2),x, algorithm="maxima")`

output `1/4*x^4*arcsin(x^2) - 1/8*sqrt(-x^4 + 1)/(x^2*((x^4 - 1)/x^4 - 1)) + 1/8*arctan(sqrt(-x^4 + 1)/x^2)`

3.350.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int x^3 \arcsin(x^2) dx = \frac{1}{8} \sqrt{-x^4 + 1} x^2 + \frac{1}{4} (x^4 - 1) \arcsin(x^2) + \frac{1}{8} \arcsin(x^2)$$

input `integrate(x^3*arcsin(x^2),x, algorithm="giac")`output `1/8*sqrt(-x^4 + 1)*x^2 + 1/4*(x^4 - 1)*arcsin(x^2) + 1/8*arcsin(x^2)`**3.350.9 Mupad [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.74

$$\int x^3 \arcsin(x^2) dx = \frac{x^2 \sqrt{1 - x^4}}{8} + \frac{\arcsin(x^2) (2x^4 - 1)}{8}$$

input `int(x^3*asin(x^2),x)`output `(x^2*(1 - x^4)^(1/2))/8 + (asin(x^2)*(2*x^4 - 1))/8`

3.351 $\int e^x \operatorname{sech}(e^x) dx$

3.351.1 Optimal result	1764
3.351.2 Mathematica [A] (verified)	1764
3.351.3 Rubi [A] (verified)	1765
3.351.4 Maple [A] (verified)	1766
3.351.5 Fricas [B] (verification not implemented)	1766
3.351.6 Sympy [A] (verification not implemented)	1767
3.351.7 Maxima [A] (verification not implemented)	1767
3.351.8 Giac [A] (verification not implemented)	1767
3.351.9 Mupad [B] (verification not implemented)	1768

3.351.1 Optimal result

Integrand size = 8, antiderivative size = 5

$$\int e^x \operatorname{sech}(e^x) dx = \arctan(\sinh(e^x))$$

output `arctan(sinh(exp(x)))`

3.351.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int e^x \operatorname{sech}(e^x) dx = \arctan(\sinh(e^x))$$

input `Integrate[E^x*Sech[E^x],x]`

output `ArcTan[Sinh[E^x]]`

3.351.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {2720, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^x \operatorname{sech}(e^x) dx \\ & \quad \downarrow \text{2720} \\ & \int \operatorname{sech}(e^x) de^x \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(\frac{\pi}{2} + ie^x\right) de^x \\ & \quad \downarrow \text{4257} \\ & \arctan(\sinh(e^x)) \end{aligned}$$

input `Int[E^x*Sech[E^x],x]`

output `ArcTan[Sinh[E^x]]`

3.351.3.1 Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.351.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

method	result	size
derivativdivides	$\arctan(\sinh(e^x))$	5
default	$\arctan(\sinh(e^x))$	5
risch	$i \ln(e^{e^x} + i) - i \ln(e^{e^x} - i)$	22
parallelrisc	$-i(\ln(\tanh(\frac{e^x}{2}) - i) - \ln(\tanh(\frac{e^x}{2}) + i))$	25

input `int(exp(x)*sech(exp(x)),x,method=_RETURNVERBOSE)`

output `arctan(sinh(exp(x)))`

3.351.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 16 vs. $2(4) = 8$.

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 3.20

$$\int e^x \operatorname{sech}(e^x) dx = 2 \arctan(\cosh(\cosh(x) + \sinh(x)) + \sinh(\cosh(x) + \sinh(x)))$$

input `integrate(exp(x)*sech(exp(x)),x, algorithm="fricas")`

output `2*arctan(cosh(cosh(x) + sinh(x)) + sinh(cosh(x) + sinh(x)))`

3.351.6 Sympy [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.60

$$\int e^x \operatorname{sech}(e^x) dx = 2 \operatorname{atan} \left(\tanh \left(\frac{e^x}{2} \right) \right)$$

input `integrate(exp(x)*sech(exp(x)),x)`output `2*atan(tanh(exp(x)/2))`**3.351.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.80

$$\int e^x \operatorname{sech}(e^x) dx = \operatorname{arctan}(\sinh(e^x))$$

input `integrate(exp(x)*sech(exp(x)),x, algorithm="maxima")`output `arctan(sinh(e^x))`**3.351.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

$$\int e^x \operatorname{sech}(e^x) dx = 2 \operatorname{arctan}(e^{(e^x)})$$

input `integrate(exp(x)*sech(exp(x)),x, algorithm="giac")`output `2*arctan(e^(e^x))`

3.351.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

$$\int e^x \operatorname{sech}(e^x) dx = 2 \operatorname{atan}(e^{e^x})$$

input `int(exp(x)/cosh(exp(x)),x)`

output `2*atan(exp(exp(x)))`

3.352 $\int x^2 \cos(3x) dx$

3.352.1 Optimal result	1769
3.352.2 Mathematica [A] (verified)	1769
3.352.3 Rubi [A] (verified)	1770
3.352.4 Maple [A] (verified)	1771
3.352.5 Fricas [A] (verification not implemented)	1772
3.352.6 Sympy [A] (verification not implemented)	1772
3.352.7 Maxima [A] (verification not implemented)	1772
3.352.8 Giac [A] (verification not implemented)	1773
3.352.9 Mupad [B] (verification not implemented)	1773

3.352.1 Optimal result

Integrand size = 8, antiderivative size = 29

$$\int x^2 \cos(3x) dx = \frac{2}{9}x \cos(3x) - \frac{2}{27} \sin(3x) + \frac{1}{3}x^2 \sin(3x)$$

output `2/9*x*cos(3*x)-2/27*sin(3*x)+1/3*x^2*sin(3*x)`

3.352.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.86

$$\int x^2 \cos(3x) dx = \frac{2}{9}x \cos(3x) + \frac{1}{27}(-2 + 9x^2) \sin(3x)$$

input `Integrate[x^2*Cos[3*x],x]`

output `(2*x*Cos[3*x])/9 + ((-2 + 9*x^2)*Sin[3*x])/27`

3.352.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.17, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {3042, 3777, 25, 3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \cos(3x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x^2 \sin\left(3x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3777} \\
 & \frac{2}{3} \int -x \sin(3x) dx + \frac{1}{3} x^2 \sin(3x) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{3} x^2 \sin(3x) - \frac{2}{3} \int x \sin(3x) dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} x^2 \sin(3x) - \frac{2}{3} \int x \sin(3x) dx \\
 & \quad \downarrow \text{3777} \\
 & \frac{1}{3} x^2 \sin(3x) - \frac{2}{3} \left(\frac{1}{3} \int \cos(3x) dx - \frac{1}{3} x \cos(3x) \right) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} x^2 \sin(3x) - \frac{2}{3} \left(\frac{1}{3} \int \sin\left(3x + \frac{\pi}{2}\right) dx - \frac{1}{3} x \cos(3x) \right) \\
 & \quad \downarrow \text{3117} \\
 & \frac{1}{3} x^2 \sin(3x) - \frac{2}{3} \left(\frac{1}{9} \sin(3x) - \frac{1}{3} x \cos(3x) \right)
 \end{aligned}$$

input `Int [x^2*Cos [3*x] , x]`

output `(-2*(-1/3*(x*Cos [3*x]) + Sin [3*x]/9))/3 + (x^2*Sin [3*x])/3`

3.352.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

3.352.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

method	result	size
risch	$\frac{2x \cos(3x)}{9} + \frac{(9x^2 - 2) \sin(3x)}{27}$	22
derivativedivides	$\frac{2x \cos(3x)}{9} - \frac{2 \sin(3x)}{27} + \frac{x^2 \sin(3x)}{3}$	24
default	$\frac{2x \cos(3x)}{9} - \frac{2 \sin(3x)}{27} + \frac{x^2 \sin(3x)}{3}$	24
parts	$\frac{2x \cos(3x)}{9} - \frac{2 \sin(3x)}{27} + \frac{x^2 \sin(3x)}{3}$	24
meijerg	$\frac{4\sqrt{\pi} \left(\frac{3x \cos(3x)}{2\sqrt{\pi}} - \frac{(-27x^2 + 3) \sin(3x)}{6\sqrt{\pi}} \right)}{27}$	33
norman	$\frac{\frac{2x}{9} - \frac{2x \tan^2\left(\frac{3x}{2}\right)}{9} + \frac{2x^2 \tan\left(\frac{3x}{2}\right)}{3} - \frac{4 \tan\left(\frac{3x}{2}\right)}{27}}{1 + \tan^2\left(\frac{3x}{2}\right)}$	40
parallelrisc	$\frac{18x^2 \tan\left(\frac{3x}{2}\right) - 6x \tan^2\left(\frac{3x}{2}\right) + 6x - 4 \tan\left(\frac{3x}{2}\right)}{27 \tan^2\left(\frac{3x}{2}\right) + 27}$	42

input `int(x^2*cos(3*x), x, method=_RETURNVERBOSE)`

output `2/9*x*cos(3*x)+1/27*(9*x^2-2)*sin(3*x)`

3.352.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int x^2 \cos(3x) dx = \frac{2}{9} x \cos(3x) + \frac{1}{27} (9x^2 - 2) \sin(3x)$$

input `integrate(x^2*cos(3*x),x, algorithm="fricas")`output `2/9*x*cos(3*x) + 1/27*(9*x^2 - 2)*sin(3*x)`**3.352.6 Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int x^2 \cos(3x) dx = \frac{x^2 \sin(3x)}{3} + \frac{2x \cos(3x)}{9} - \frac{2 \sin(3x)}{27}$$

input `integrate(x**2*cos(3*x),x)`output `x**2*sin(3*x)/3 + 2*x*cos(3*x)/9 - 2*sin(3*x)/27`**3.352.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int x^2 \cos(3x) dx = \frac{2}{9} x \cos(3x) + \frac{1}{27} (9x^2 - 2) \sin(3x)$$

input `integrate(x^2*cos(3*x),x, algorithm="maxima")`output `2/9*x*cos(3*x) + 1/27*(9*x^2 - 2)*sin(3*x)`

3.352.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int x^2 \cos(3x) dx = \frac{2}{9} x \cos(3x) + \frac{1}{27} (9x^2 - 2) \sin(3x)$$

input `integrate(x^2*cos(3*x),x, algorithm="giac")`output `2/9*x*cos(3*x) + 1/27*(9*x^2 - 2)*sin(3*x)`**3.352.9 Mupad [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

$$\int x^2 \cos(3x) dx = \frac{2x \cos(3x)}{9} - \frac{2 \sin(3x)}{27} + \frac{x^2 \sin(3x)}{3}$$

input `int(x^2*cos(3*x),x)`output `(2*x*cos(3*x))/9 - (2*sin(3*x))/27 + (x^2*sin(3*x))/3`

3.353 $\int \sqrt{5 - 4x - x^2} dx$

3.353.1 Optimal result	1774
3.353.2 Mathematica [A] (verified)	1774
3.353.3 Rubi [A] (verified)	1775
3.353.4 Maple [A] (verified)	1776
3.353.5 Fricas [A] (verification not implemented)	1776
3.353.6 Sympy [A] (verification not implemented)	1777
3.353.7 Maxima [A] (verification not implemented)	1777
3.353.8 Giac [A] (verification not implemented)	1777
3.353.9 Mupad [B] (verification not implemented)	1778

3.353.1 Optimal result

Integrand size = 14, antiderivative size = 36

$$\int \sqrt{5 - 4x - x^2} dx = \frac{1}{2}(2 + x)\sqrt{5 - 4x - x^2} - \frac{9}{2} \arcsin\left(\frac{1}{3}(-2 - x)\right)$$

output `9/2*arcsin(2/3+1/3*x)+1/2*(2+x)*(-x^2-4*x+5)^(1/2)`

3.353.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.25

$$\int \sqrt{5 - 4x - x^2} dx = \frac{1}{2}(2 + x)\sqrt{5 - 4x - x^2} - 9 \arctan\left(\frac{\sqrt{5 - 4x - x^2}}{5 + x}\right)$$

input `Integrate[Sqrt[5 - 4*x - x^2], x]`

output `((2 + x)*Sqrt[5 - 4*x - x^2])/2 - 9*ArcTan[Sqrt[5 - 4*x - x^2]/(5 + x)]`

3.353.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {1087, 1090, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{-x^2 - 4x + 5} dx$$

$$\downarrow \text{1087}$$

$$\frac{9}{2} \int \frac{1}{\sqrt{-x^2 - 4x + 5}} dx + \frac{1}{2} \sqrt{-x^2 - 4x + 5}(x + 2)$$

$$\downarrow \text{1090}$$

$$\frac{1}{2}(x + 2)\sqrt{-x^2 - 4x + 5} - \frac{3}{4} \int \frac{1}{\sqrt{1 - \frac{1}{36}(-2x - 4)^2}} d(-2x - 4)$$

$$\downarrow \text{223}$$

$$\frac{1}{2}(x + 2)\sqrt{-x^2 - 4x + 5} - \frac{9}{2} \arcsin\left(\frac{1}{6}(-2x - 4)\right)$$

input `Int[Sqrt[5 - 4*x - x^2], x]`

output `((2 + x)*Sqrt[5 - 4*x - x^2])/2 - (9*ArcSin[(-4 - 2*x)/6])/2`

3.353.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 1087 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] - Simp[p*(b^2 - 4*a*c)/(2*c*(2*p + 1)) Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[3*p])`

rule 1090 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

3.353.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.81

method	result	size
default	$-\frac{(-2x-4)\sqrt{-x^2-4x+5}}{4} + \frac{9 \arcsin(\frac{2}{3} + \frac{x}{3})}{2}$	29
risch	$-\frac{(2+x)(x^2+4x-5)}{2\sqrt{-x^2-4x+5}} + \frac{9 \arcsin(\frac{2}{3} + \frac{x}{3})}{2}$	35
trager	$(1 + \frac{x}{2}) \sqrt{-x^2 - 4x + 5} + \frac{9 \operatorname{RootOf}(_Z^2 + 1) \ln(-\operatorname{RootOf}(_Z^2 + 1)x + \sqrt{-x^2 - 4x + 5} - 2 \operatorname{RootOf}(_Z^2 + 1))}{2}$	59

input `int((-x^2-4*x+5)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/4*(-2*x-4)*(-x^2-4*x+5)^(1/2)+9/2*arcsin(2/3+1/3*x)`

3.353.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.31

$$\int \sqrt{5 - 4x - x^2} dx = \frac{1}{2} \sqrt{-x^2 - 4x + 5}(x + 2) - \frac{9}{2} \arctan\left(\frac{\sqrt{-x^2 - 4x + 5}(x + 2)}{x^2 + 4x - 5}\right)$$

input `integrate((-x^2-4*x+5)^(1/2),x, algorithm="fricas")`

output `1/2*sqrt(-x^2 - 4*x + 5)*(x + 2) - 9/2*arctan(sqrt(-x^2 - 4*x + 5)*(x + 2)/(x^2 + 4*x - 5))`

3.353.6 Sympy [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.75

$$\int \sqrt{5 - 4x - x^2} dx = \left(\frac{x}{2} + 1\right) \sqrt{-x^2 - 4x + 5} + \frac{9 \operatorname{asin}\left(\frac{x}{3} + \frac{2}{3}\right)}{2}$$

input `integrate((-x**2-4*x+5)**(1/2),x)`output `(x/2 + 1)*sqrt(-x**2 - 4*x + 5) + 9*asin(x/3 + 2/3)/2`**3.353.7 Maxima [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00

$$\int \sqrt{5 - 4x - x^2} dx = \frac{1}{2} \sqrt{-x^2 - 4x + 5} x + \sqrt{-x^2 - 4x + 5} - \frac{9}{2} \arcsin\left(-\frac{1}{3}x - \frac{2}{3}\right)$$

input `integrate((-x^2-4*x+5)^(1/2),x, algorithm="maxima")`output `1/2*sqrt(-x^2 - 4*x + 5)*x + sqrt(-x^2 - 4*x + 5) - 9/2*arcsin(-1/3*x - 2/3)`**3.353.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.72

$$\int \sqrt{5 - 4x - x^2} dx = \frac{1}{2} \sqrt{-x^2 - 4x + 5}(x + 2) + \frac{9}{2} \arcsin\left(\frac{1}{3}x + \frac{2}{3}\right)$$

input `integrate((-x^2-4*x+5)^(1/2),x, algorithm="giac")`output `1/2*sqrt(-x^2 - 4*x + 5)*(x + 2) + 9/2*arcsin(1/3*x + 2/3)`

3.353.9 Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.75

$$\int \sqrt{5 - 4x - x^2} dx = \frac{9 \operatorname{asin}\left(\frac{x}{3} + \frac{2}{3}\right)}{2} + \left(\frac{x}{2} + 1\right) \sqrt{-x^2 - 4x + 5}$$

input `int((5 - x^2 - 4*x)^(1/2),x)`

output `(9*asin(x/3 + 2/3))/2 + (x/2 + 1)*(5 - x^2 - 4*x)^(1/2)`

3.354 $\int \frac{x^5}{\sqrt{2+x^2}} dx$

3.354.1 Optimal result	1779
3.354.2 Mathematica [A] (verified)	1779
3.354.3 Rubi [A] (verified)	1780
3.354.4 Maple [A] (verified)	1781
3.354.5 Fricas [A] (verification not implemented)	1781
3.354.6 Sympy [A] (verification not implemented)	1781
3.354.7 Maxima [A] (verification not implemented)	1782
3.354.8 Giac [A] (verification not implemented)	1782
3.354.9 Mupad [B] (verification not implemented)	1782

3.354.1 Optimal result

Integrand size = 15, antiderivative size = 28

$$\int \frac{x^5}{\sqrt{2+x^2}} dx = -\frac{x^2}{\sqrt{2}} + \frac{x^4}{4} + \log(\sqrt{2} + x^2)$$

output `1/4*x^4+ln(x^2+2^(1/2))-1/2*x^2*2^(1/2)`

3.354.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.11

$$\int \frac{x^5}{\sqrt{2+x^2}} dx = \frac{1}{4} \left(-6 - 2\sqrt{2}x^2 + x^4 + 4 \log(\sqrt{2} + x^2) \right)$$

input `Integrate[x^5/(Sqrt[2] + x^2),x]`

output `(-6 - 2*Sqrt[2]*x^2 + x^4 + 4*Log[Sqrt[2] + x^2])/4`

3.354.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.21, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^5}{x^2 + \sqrt{2}} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int \frac{x^4}{x^2 + \sqrt{2}} dx^2 \\ & \quad \downarrow \text{49} \\ & \frac{1}{2} \int \left(x^2 + \frac{2}{x^2 + \sqrt{2}} - \sqrt{2} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \left(\frac{x^4}{2} - \sqrt{2}x^2 + 2 \log(x^2 + \sqrt{2}) \right) \end{aligned}$$

input `Int[x^5/(Sqrt[2] + x^2),x]`

output `(-(Sqrt[2]*x^2) + x^4/2 + 2*Log[Sqrt[2] + x^2])/2`

3.354.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.354.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{x^4}{4} + \ln(x^2 + \sqrt{2}) - \frac{x^2\sqrt{2}}{2}$	23
parallelrisc	$\frac{x^4}{4} + \ln(x^2 + \sqrt{2}) - \frac{x^2\sqrt{2}}{2}$	23
risc	$\frac{x^4}{4} - \frac{x^2\sqrt{2}}{2} + \frac{1}{2} + \ln(x^2 + \sqrt{2})$	24
meijerg	$-\frac{x^2\sqrt{2}\left(-\frac{3x^2\sqrt{2}}{2}+6\right)}{12} + \ln\left(1 + \frac{x^2\sqrt{2}}{2}\right)$	31

input `int(x^5/(x^2+2^(1/2)),x,method=_RETURNVERBOSE)`output `1/4*x^4+ln(x^2+2^(1/2))-1/2*x^2*2^(1/2)`**3.354.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int \frac{x^5}{\sqrt{2} + x^2} dx = \frac{1}{4}x^4 - \frac{1}{2}\sqrt{2}x^2 + \log(x^2 + \sqrt{2})$$

input `integrate(x^5/(x^2+2^(1/2)),x, algorithm="fracas")`output `1/4*x^4 - 1/2*sqrt(2)*x^2 + log(x^2 + sqrt(2))`**3.354.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int \frac{x^5}{\sqrt{2} + x^2} dx = \frac{x^4}{4} - \frac{\sqrt{2}x^2}{2} + \log(x^2 + \sqrt{2})$$

input `integrate(x**5/(x**2+2**(1/2)),x)`output `x**4/4 - sqrt(2)*x**2/2 + log(x**2 + sqrt(2))`

3.354.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int \frac{x^5}{\sqrt{2} + x^2} dx = \frac{1}{4}x^4 - \frac{1}{2}\sqrt{2}x^2 + \log(x^2 + \sqrt{2})$$

input `integrate(x^5/(x^2+2^(1/2)),x, algorithm="maxima")`output `1/4*x^4 - 1/2*sqrt(2)*x^2 + log(x^2 + sqrt(2))`**3.354.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int \frac{x^5}{\sqrt{2} + x^2} dx = \frac{1}{4}x^4 - \frac{1}{2}\sqrt{2}x^2 + \log(x^2 + \sqrt{2})$$

input `integrate(x^5/(x^2+2^(1/2)),x, algorithm="giac")`output `1/4*x^4 - 1/2*sqrt(2)*x^2 + log(x^2 + sqrt(2))`**3.354.9 Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int \frac{x^5}{\sqrt{2} + x^2} dx = \ln(x^2 + \sqrt{2}) - \frac{\sqrt{2}x^2}{2} + \frac{x^4}{4}$$

input `int(x^5/(2^(1/2) + x^2),x)`output `log(2^(1/2) + x^2) - (2^(1/2)*x^2)/2 + x^4/4`

3.355 $\int \sec^5(x) dx$

3.355.1 Optimal result	1783
3.355.2 Mathematica [A] (verified)	1783
3.355.3 Rubi [A] (verified)	1784
3.355.4 Maple [A] (verified)	1785
3.355.5 Fricas [B] (verification not implemented)	1786
3.355.6 Sympy [A] (verification not implemented)	1786
3.355.7 Maxima [B] (verification not implemented)	1786
3.355.8 Giac [A] (verification not implemented)	1787
3.355.9 Mupad [B] (verification not implemented)	1787

3.355.1 Optimal result

Integrand size = 4, antiderivative size = 26

$$\int \sec^5(x) dx = \frac{3}{8} \operatorname{arctanh}(\sin(x)) + \frac{3}{8} \sec(x) \tan(x) + \frac{1}{4} \sec^3(x) \tan(x)$$

output `3/8*arctanh(sin(x))+3/8*sec(x)*tan(x)+1/4*sec(x)^3*tan(x)`

3.355.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \sec^5(x) dx = \frac{3}{8} \operatorname{arctanh}(\sin(x)) + \frac{3}{8} \sec(x) \tan(x) + \frac{1}{4} \sec^3(x) \tan(x)$$

input `Integrate[Sec[x]^5,x]`

output `(3*ArcTanh[Sin[x]])/8 + (3*Sec[x]*Tan[x])/8 + (Sec[x]^3*Tan[x])/4`

3.355.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.19, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {3042, 4255, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^5(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(x + \frac{\pi}{2}\right)^5 dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{3}{4} \int \sec^3(x) dx + \frac{1}{4} \tan(x) \sec^3(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{4} \int \csc\left(x + \frac{\pi}{2}\right)^3 dx + \frac{1}{4} \tan(x) \sec^3(x) \\
 & \quad \downarrow \text{4255} \\
 & \frac{3}{4} \left(\frac{\int \sec(x) dx}{2} + \frac{1}{2} \tan(x) \sec(x) \right) + \frac{1}{4} \tan(x) \sec^3(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{4} \left(\frac{1}{2} \int \csc\left(x + \frac{\pi}{2}\right) dx + \frac{1}{2} \tan(x) \sec(x) \right) + \frac{1}{4} \tan(x) \sec^3(x) \\
 & \quad \downarrow \text{4257} \\
 & \frac{3}{4} \left(\frac{1}{2} \operatorname{arctanh}(\sin(x)) + \frac{1}{2} \tan(x) \sec(x) \right) + \frac{1}{4} \tan(x) \sec^3(x)
 \end{aligned}$$

input `Int[Sec[x]^5, x]`

output `(Sec[x]^3*Tan[x])/4 + (3*(ArcTanh[Sin[x]]/2 + (Sec[x]*Tan[x])/2))/4`

3.355.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.355.4 Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.96

method	result	size
default	$-\left(-\frac{\sec^3(x)}{4} - \frac{3\sec(x)}{8}\right)\tan(x) + \frac{3\ln(\sec(x)+\tan(x))}{8}$	25
parallelrisch	$\ln\left(-\cot(x) + 1 + \csc(x)\right)^{\frac{3}{8}} + \ln\left(\frac{1}{(-\cot(x)+\csc(x)-1)^{\frac{3}{8}}}\right) + \frac{3\sec(x)\tan(x)}{8} + \frac{(\sec^3(x))\tan(x)}{4}$	38
norman	$\frac{\frac{3(\tan^3(\frac{x}{2}))}{4} + \frac{3(\tan^5(\frac{x}{2}))}{4} + \frac{5(\tan^7(\frac{x}{2}))}{4} + \frac{5\tan(\frac{x}{2})}{4}}{(\tan^2(\frac{x}{2})-1)^4} - \frac{3\ln(\tan(\frac{x}{2})-1)}{8} + \frac{3\ln(1+\tan(\frac{x}{2}))}{8}$	62
risch	$-\frac{i(3e^{7ix}+11e^{5ix}-11e^{3ix}-3e^{ix})}{4(e^{2ix}+1)^4} + \frac{3\ln(i+e^{ix})}{8} - \frac{3\ln(e^{ix}-i)}{8}$	65

input `int(sec(x)^5,x,method=_RETURNVERBOSE)`

output `-(-1/4*sec(x)^3-3/8*sec(x))*tan(x)+3/8*ln(sec(x)+tan(x))`

3.355.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. $2(20) = 40$.

Time = 0.25 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.65

$$\int \sec^5(x) dx = \frac{3 \cos(x)^4 \log(\sin(x) + 1) - 3 \cos(x)^4 \log(-\sin(x) + 1) + 2(3 \cos(x)^2 + 2) \sin(x)}{16 \cos(x)^4}$$

input `integrate(sec(x)^5,x, algorithm="fricas")`

output `1/16*(3*cos(x)^4*log(sin(x) + 1) - 3*cos(x)^4*log(-sin(x) + 1) + 2*(3*cos(x)^2 + 2)*sin(x))/cos(x)^4`

3.355.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.77

$$\int \sec^5(x) dx = -\frac{3 \sin^3(x) - 5 \sin(x)}{8 \sin^4(x) - 16 \sin^2(x) + 8} - \frac{3 \log(\sin(x) - 1)}{16} + \frac{3 \log(\sin(x) + 1)}{16}$$

input `integrate(sec(x)**5,x)`

output `-(3*sin(x)**3 - 5*sin(x))/(8*sin(x)**4 - 16*sin(x)**2 + 8) - 3*log(sin(x) - 1)/16 + 3*log(sin(x) + 1)/16`

3.355.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. $2(20) = 40$.

Time = 0.21 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.62

$$\int \sec^5(x) dx = -\frac{3 \sin(x)^3 - 5 \sin(x)}{8 (\sin(x)^4 - 2 \sin(x)^2 + 1)} + \frac{3}{16} \log(\sin(x) + 1) - \frac{3}{16} \log(\sin(x) - 1)$$

input `integrate(sec(x)^5,x, algorithm="maxima")`

output
$$-1/8*(3*\sin(x)^3 - 5*\sin(x))/(\sin(x)^4 - 2*\sin(x)^2 + 1) + 3/16*\log(\sin(x) + 1) - 3/16*\log(\sin(x) - 1)$$

3.355.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.46

$$\int \sec^5(x) dx = -\frac{3 \sin(x)^3 - 5 \sin(x)}{8 (\sin(x)^2 - 1)^2} + \frac{3}{16} \log(\sin(x) + 1) - \frac{3}{16} \log(-\sin(x) + 1)$$

input `integrate(sec(x)^5,x, algorithm="giac")`

output
$$-1/8*(3*\sin(x)^3 - 5*\sin(x))/(\sin(x)^2 - 1)^2 + 3/16*\log(\sin(x) + 1) - 3/16*\log(-\sin(x) + 1)$$

3.355.9 Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.12

$$\int \sec^5(x) dx = \frac{3 \ln\left(\frac{\sin(x)+1}{\cos(x)}\right)}{8} + \sin(x) \left(\frac{3}{8 \cos(x)^2} + \frac{1}{4 \cos(x)^4} \right)$$

input `int(1/cos(x)^5,x)`

output
$$(3*\log((\sin(x) + 1)/\cos(x)))/8 + \sin(x)*(3/(8*\cos(x)^2) + 1/(4*\cos(x)^4))$$

3.356 $\int \sin^6(2x) dx$

3.356.1 Optimal result	1788
3.356.2 Mathematica [A] (verified)	1788
3.356.3 Rubi [A] (verified)	1789
3.356.4 Maple [A] (verified)	1790
3.356.5 Fricas [A] (verification not implemented)	1791
3.356.6 Sympy [A] (verification not implemented)	1791
3.356.7 Maxima [A] (verification not implemented)	1791
3.356.8 Giac [A] (verification not implemented)	1792
3.356.9 Mupad [B] (verification not implemented)	1792

3.356.1 Optimal result

Integrand size = 6, antiderivative size = 46

$$\int \sin^6(2x) dx = \frac{5x}{16} - \frac{5}{32} \cos(2x) \sin(2x) - \frac{5}{48} \cos(2x) \sin^3(2x) - \frac{1}{12} \cos(2x) \sin^5(2x)$$

output `5/16*x-5/32*cos(2*x)*sin(2*x)-5/48*cos(2*x)*sin(2*x)^3-1/12*cos(2*x)*sin(2*x)^5`

3.356.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.65

$$\int \sin^6(2x) dx = \frac{5x}{16} - \frac{15}{128} \sin(4x) + \frac{3}{128} \sin(8x) - \frac{1}{384} \sin(12x)$$

input `Integrate[Sin[2*x]^6,x]`

output `(5*x)/16 - (15*Sin[4*x])/128 + (3*Sin[8*x])/128 - Sin[12*x]/384`

3.356.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.22, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.167$, Rules used = {3042, 3115, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^6(2x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(2x)^6 dx \\
 & \quad \downarrow \text{3115} \\
 & \frac{5}{6} \int \sin^4(2x) dx - \frac{1}{12} \sin^5(2x) \cos(2x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{6} \int \sin(2x)^4 dx - \frac{1}{12} \sin^5(2x) \cos(2x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{5}{6} \left(\frac{3}{4} \int \sin^2(2x) dx - \frac{1}{8} \sin^3(2x) \cos(2x) \right) - \frac{1}{12} \sin^5(2x) \cos(2x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{6} \left(\frac{3}{4} \int \sin(2x)^2 dx - \frac{1}{8} \sin^3(2x) \cos(2x) \right) - \frac{1}{12} \sin^5(2x) \cos(2x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{5}{6} \left(\frac{3}{4} \left(\frac{\int 1 dx}{2} - \frac{1}{4} \sin(2x) \cos(2x) \right) - \frac{1}{8} \sin^3(2x) \cos(2x) \right) - \frac{1}{12} \sin^5(2x) \cos(2x) \\
 & \quad \downarrow \text{24} \\
 & \frac{5}{6} \left(\frac{3}{4} \left(\frac{x}{2} - \frac{1}{4} \sin(2x) \cos(2x) \right) - \frac{1}{8} \sin^3(2x) \cos(2x) \right) - \frac{1}{12} \sin^5(2x) \cos(2x)
 \end{aligned}$$

input `Int [Sin [2*x]^6, x]`

```
output -1/12*(Cos[2*x]*Sin[2*x]^5) + (5*(-1/8*(Cos[2*x]*Sin[2*x]^3) + (3*(x/2 - (Cos[2*x]*Sin[2*x])/4))/4))/6
```

3.356.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3115 Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sine[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sine[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

3.356.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.50

method	result
risch	$\frac{5x}{16} - \frac{\sin(12x)}{384} + \frac{3\sin(8x)}{128} - \frac{15\sin(4x)}{128}$
parallelrisch	$\frac{5x}{16} - \frac{\sin(12x)}{384} + \frac{3\sin(8x)}{128} - \frac{15\sin(4x)}{128}$
derivativedivides	$-\frac{\left(\sin^5(2x) + \frac{5\sin^3(2x)}{4} + \frac{15\sin(2x)}{8}\right)\cos(2x)}{12} + \frac{5x}{16}$
default	$-\frac{\left(\sin^5(2x) + \frac{5\sin^3(2x)}{4} + \frac{15\sin(2x)}{8}\right)\cos(2x)}{12} + \frac{5x}{16}$
norman	$\frac{5x}{16} - \frac{85(\tan^3(x))}{48} - \frac{33(\tan^5(x))}{8} + \frac{33(\tan^7(x))}{8} + \frac{85(\tan^9(x))}{48} + \frac{5(\tan^{11}(x))}{16} + \frac{15x(\tan^2(x))}{8} + \frac{75x(\tan^4(x))}{16} + \frac{25x(\tan^6(x))}{4} + \frac{7}{(1+\tan^2(x))^6}$

```
input int(sin(2*x)^6,x,method=_RETURNVERBOSE)
```

```
output 5/16*x-1/384*sin(12*x)+3/128*sin(8*x)-15/128*sin(4*x)
```

3.356.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.72

$$\int \sin^6(2x) dx = -\frac{1}{96} (8 \cos(2x)^5 - 26 \cos(2x)^3 + 33 \cos(2x)) \sin(2x) + \frac{5}{16} x$$

input `integrate(sin(2*x)^6,x, algorithm="fricas")`output `-1/96*(8*cos(2*x)^5 - 26*cos(2*x)^3 + 33*cos(2*x))*sin(2*x) + 5/16*x`**3.356.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \sin^6(2x) dx = \frac{5x}{16} - \frac{\sin^5(2x) \cos(2x)}{12} - \frac{5 \sin^3(2x) \cos(2x)}{48} - \frac{5 \sin(2x) \cos(2x)}{32}$$

input `integrate(sin(2*x)**6,x)`output `5*x/16 - sin(2*x)**5*cos(2*x)/12 - 5*sin(2*x)**3*cos(2*x)/48 - 5*sin(2*x)*cos(2*x)/32`**3.356.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.52

$$\int \sin^6(2x) dx = \frac{1}{96} \sin(4x)^3 + \frac{5}{16} x + \frac{3}{128} \sin(8x) - \frac{1}{8} \sin(4x)$$

input `integrate(sin(2*x)^6,x, algorithm="maxima")`output `1/96*sin(4*x)^3 + 5/16*x + 3/128*sin(8*x) - 1/8*sin(4*x)`

3.356.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.48

$$\int \sin^6(2x) dx = \frac{5}{16} x - \frac{1}{384} \sin(12x) + \frac{3}{128} \sin(8x) - \frac{15}{128} \sin(4x)$$

input `integrate(sin(2*x)^6,x, algorithm="giac")`output `5/16*x - 1/384*sin(12*x) + 3/128*sin(8*x) - 15/128*sin(4*x)`**3.356.9 Mupad [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.48

$$\int \sin^6(2x) dx = \frac{5x}{16} - \frac{15 \sin(4x)}{128} + \frac{3 \sin(8x)}{128} - \frac{\sin(12x)}{384}$$

input `int(sin(2*x)^6,x)`output `(5*x)/16 - (15*sin(4*x))/128 + (3*sin(8*x))/128 - sin(12*x)/384`

3.357 $\int \cos(x) \log(\sin(x)) \sin^2(x) dx$

3.357.1 Optimal result	1793
3.357.2 Mathematica [A] (verified)	1793
3.357.3 Rubi [A] (verified)	1794
3.357.4 Maple [A] (verified)	1795
3.357.5 Fricas [A] (verification not implemented)	1796
3.357.6 Sympy [A] (verification not implemented)	1796
3.357.7 Maxima [A] (verification not implemented)	1796
3.357.8 Giac [A] (verification not implemented)	1797
3.357.9 Mupad [B] (verification not implemented)	1797

3.357.1 Optimal result

Integrand size = 10, antiderivative size = 20

$$\int \cos(x) \log(\sin(x)) \sin^2(x) dx = -\frac{1}{9} \sin^3(x) + \frac{1}{3} \log(\sin(x)) \sin^3(x)$$

output `-1/9*sin(x)^3+1/3*ln(sin(x))*sin(x)^3`

3.357.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int \cos(x) \log(\sin(x)) \sin^2(x) dx = \frac{1}{9}(-1 + 3 \log(\sin(x))) \sin^3(x)$$

input `Integrate[Cos[x]*Log[Sin[x]]*Sin[x]^2,x]`

output `((-1 + 3*Log[Sin[x]])*Sin[x]^3)/9`

3.357.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3034, 27, 3042, 3044, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^2(x) \cos(x) \log(\sin(x)) dx \\
 & \quad \downarrow \text{3034} \\
 & \frac{1}{3} \sin^3(x) \log(\sin(x)) - \int \frac{1}{3} \cos(x) \sin^2(x) dx \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{3} \sin^3(x) \log(\sin(x)) - \frac{1}{3} \int \cos(x) \sin^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \sin^3(x) \log(\sin(x)) - \frac{1}{3} \int \cos(x) \sin(x)^2 dx \\
 & \quad \downarrow \text{3044} \\
 & \frac{1}{3} \sin^3(x) \log(\sin(x)) - \frac{1}{3} \int \sin^2(x) d \sin(x) \\
 & \quad \downarrow \text{15} \\
 & \frac{1}{3} \sin^3(x) \log(\sin(x)) - \frac{\sin^3(x)}{9}
 \end{aligned}$$

input `Int[Cos[x]*Log[Sin[x]]*Sin[x]^2,x]`

output `-1/9*Sin[x]^3 + (Log[Sin[x]]*Sin[x]^3)/3`

3.357.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3034 `Int[Log[u_]*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[Log[u] w, x] - Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w, x] /; InverseFunctionFreeQ[u, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3044 `Int[cos[(e_.) + (f_.)*(x_)^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Simp[1/(a*f) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])`

3.357.4 Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

method	result	size
derivativedivides	$-\frac{(\sin^3(x))}{9} + \frac{\ln(\sin(x))(\sin^3(x))}{3}$	17
default	$-\frac{(\sin^3(x))}{9} + \frac{\ln(\sin(x))(\sin^3(x))}{3}$	17
parallelrisc	$\frac{(3 \ln(\sin(x)) - 1)(-\sin(3x) + 3 \sin(x))}{36}$	21
risc	Expression too large to display	577

input `int(cos(x)*ln(sin(x))*sin(x)^2,x,method=_RETURNVERBOSE)`

output `-1/9*sin(x)^3+1/3*ln(sin(x))*sin(x)^3`

3.357.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.20

$$\int \cos(x) \log(\sin(x)) \sin^2(x) dx$$

$$= -\frac{1}{3} (\cos(x)^2 - 1) \log(\sin(x)) \sin(x) + \frac{1}{9} (\cos(x)^2 - 1) \sin(x)$$

input `integrate(cos(x)*log(sin(x))*sin(x)^2,x, algorithm="fricas")`output `-1/3*(cos(x)^2 - 1)*log(sin(x))*sin(x) + 1/9*(cos(x)^2 - 1)*sin(x)`**3.357.6 Sympy [A] (verification not implemented)**

Time = 0.64 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \cos(x) \log(\sin(x)) \sin^2(x) dx = \frac{\log(\sin(x)) \sin^3(x)}{3} - \frac{\sin^3(x)}{9}$$

input `integrate(cos(x)*ln(sin(x))*sin(x)**2,x)`output `log(sin(x))*sin(x)**3/3 - sin(x)**3/9`**3.357.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int \cos(x) \log(\sin(x)) \sin^2(x) dx = \frac{1}{3} \log(\sin(x)) \sin(x)^3 - \frac{1}{9} \sin(x)^3$$

input `integrate(cos(x)*log(sin(x))*sin(x)^2,x, algorithm="maxima")`output `1/3*log(sin(x))*sin(x)^3 - 1/9*sin(x)^3`

3.357.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int \cos(x) \log(\sin(x)) \sin^2(x) dx = \frac{1}{3} \log(\sin(x)) \sin^3(x) - \frac{1}{9} \sin^3(x)$$

input `integrate(cos(x)*log(sin(x))*sin(x)^2,x, algorithm="giac")`

output `1/3*log(sin(x))*sin(x)^3 - 1/9*sin(x)^3`

3.357.9 Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.55

$$\int \cos(x) \log(\sin(x)) \sin^2(x) dx = \frac{\sin^3(x) (\ln(\sin(x)) - \frac{1}{3})}{3}$$

input `int(log(sin(x))*cos(x)*sin(x)^2,x)`

output `(sin(x)^3*(log(sin(x)) - 1/3))/3`

3.358 $\int \frac{e^{-x}}{1+2e^x} dx$

3.358.1 Optimal result	1798
3.358.2 Mathematica [A] (verified)	1798
3.358.3 Rubi [A] (verified)	1799
3.358.4 Maple [A] (verified)	1800
3.358.5 Fricas [A] (verification not implemented)	1800
3.358.6 Sympy [A] (verification not implemented)	1801
3.358.7 Maxima [A] (verification not implemented)	1801
3.358.8 Giac [A] (verification not implemented)	1801
3.358.9 Mupad [B] (verification not implemented)	1802

3.358.1 Optimal result

Integrand size = 15, antiderivative size = 21

$$\int \frac{e^{-x}}{1+2e^x} dx = -e^{-x} - 2x + 2 \log(1+2e^x)$$

output `-1/exp(x)-2*x+2*ln(1+2*exp(x))`

3.358.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.14

$$\int \frac{e^{-x}}{1+2e^x} dx = -e^{-x} - 2 \log(e^x) + 2 \log(1+2e^x)$$

input `Integrate[1/(E^x*(1 + 2*E^x)),x]`

output `-E^(-x) - 2*Log[E^x] + 2*Log[1 + 2*E^x]`

3.358.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.14, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2678, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{e^{-x}}{2e^x + 1} dx \\ & \quad \downarrow \text{2678} \\ & \int \frac{e^{-2x}}{2e^x + 1} de^x \\ & \quad \downarrow \text{54} \\ & \int \left(e^{-2x} - 2e^{-x} + \frac{4}{2e^x + 1} \right) de^x \\ & \quad \downarrow \text{2009} \\ & -e^{-x} - 2 \log(e^x) + 2 \log(2e^x + 1) \end{aligned}$$

input `Int[1/(E^x*(1 + 2*E^x)),x]`

output `-E^(-x) - 2*Log[E^x] + 2*Log[1 + 2*E^x]`

3.358.3.1 Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`


```
rule 2678 Int[((a_) + (b_.)*(F_)^((e_.)*((c_.) + (d_.)*(x_)))^(p_.)*(G_)^((h_.)*((f_.) + (g_.)*(x_))), x_Symbol] := With[{m = FullSimplify[g*h*(Log[G]/(d*e*Log[F]))}], Simp[Denominator[m]*(G^(f*h - c*g*(h/d))/(d*e*Log[F])) Subst[Int[x^(Numerator[m] - 1)*(a + b*x^Denominator[m])^p, x], x, F^(e*((c + d*x)/Denominator[m]))], x] /; LeQ[m, -1] || GeQ[m, 1] /; FreeQ[{F, G, a, b, c, d, e, f, g, h, p}, x]
```

3.358.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

method	result	size
risch	$-e^{-x} - 2x + 2 \ln\left(\frac{1}{2} + e^x\right)$	18
derivativedivides	$2 \ln(1 + 2e^x) - e^{-x} - 2 \ln(e^x)$	22
default	$2 \ln(1 + 2e^x) - e^{-x} - 2 \ln(e^x)$	22
parallelrisch	$(-1 + 2 \ln\left(\frac{1}{2} + e^x\right) e^x - 2e^x x) e^{-x}$	22
norman	$(-1 - 2e^x x) e^{-x} + 2 \ln(1 + 2e^x)$	23

```
input int(1/exp(x)/(1+2*exp(x)),x,method=_RETURNVERBOSE)
```

```
output -exp(-x)-2*x+2*ln(1/2+exp(x))
```

3.358.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.14

$$\int \frac{e^{-x}}{1+2e^x} dx = -(2xe^x - 2e^x \log(2e^x + 1) + 1)e^{(-x)}$$

```
input integrate(1/exp(x)/(1+2*exp(x)),x, algorithm="fricas")
```

```
output -(2*x*e^x - 2*e^x*log(2*e^x + 1) + 1)*e^(-x)
```

3.358.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

$$\int \frac{e^{-x}}{1+2e^x} dx = 2 \log(2 + e^{-x}) - e^{-x}$$

input `integrate(1/exp(x)/(1+2*exp(x)),x)`output `2*log(2 + exp(-x)) - exp(-x)`**3.358.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.76

$$\int \frac{e^{-x}}{1+2e^x} dx = -e^{(-x)} + 2 \log(e^{(-x)} + 2)$$

input `integrate(1/exp(x)/(1+2*exp(x)),x, algorithm="maxima")`output `-e^(-x) + 2*log(e^(-x) + 2)`**3.358.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{e^{-x}}{1+2e^x} dx = -2x - e^{(-x)} + 2 \log(2e^x + 1)$$

input `integrate(1/exp(x)/(1+2*exp(x)),x, algorithm="giac")`output `-2*x - e^(-x) + 2*log(2*e^x + 1)`

3.358.9 Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{e^{-x}}{1+2e^x} dx = 2 \ln(2e^x + 1) - 2x - e^{-x}$$

input `int(exp(-x)/(2*exp(x) + 1),x)`

output `2*log(2*exp(x) + 1) - 2*x - exp(-x)`

3.359 $\int \sqrt{2 + 3 \cos(x)} \tan(x) dx$

3.359.1 Optimal result	1803
3.359.2 Mathematica [A] (verified)	1803
3.359.3 Rubi [A] (verified)	1804
3.359.4 Maple [B] (verified)	1806
3.359.5 Fricas [A] (verification not implemented)	1806
3.359.6 Sympy [F]	1807
3.359.7 Maxima [A] (verification not implemented)	1807
3.359.8 Giac [A] (verification not implemented)	1807
3.359.9 Mupad [F(-1)]	1808

3.359.1 Optimal result

Integrand size = 13, antiderivative size = 37

$$\int \sqrt{2 + 3 \cos(x)} \tan(x) dx = 2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2 + 3 \cos(x)}}{\sqrt{2}}\right) - 2\sqrt{2 + 3 \cos(x)}$$

output `2*arctanh(1/2*(2+3*cos(x))^(1/2)*2^(1/2))*2^(1/2)-2*(2+3*cos(x))^(1/2)`

3.359.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.89

$$\int \sqrt{2 + 3 \cos(x)} \tan(x) dx = 2\sqrt{2} \operatorname{arctanh}\left(\sqrt{1 + \frac{3 \cos(x)}{2}}\right) - 2\sqrt{2 + 3 \cos(x)}$$

input `Integrate[Sqrt[2 + 3*Cos[x]]*Tan[x], x]`

output `2*Sqrt[2]*ArcTanh[Sqrt[1 + (3*Cos[x])/2]] - 2*Sqrt[2 + 3*Cos[x]]`

3.359.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {3042, 25, 3200, 60, 73, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{3 \cos(x) + 2} \tan(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\sqrt{2 - 3 \sin\left(x - \frac{\pi}{2}\right)}}{\tan\left(x - \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{\sqrt{2 - 3 \sin\left(x - \frac{\pi}{2}\right)}}{\tan\left(x - \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{3200} \\
 & -\int \frac{1}{3} \sqrt{3 \cos(x) + 2} \sec(x) d(3 \cos(x)) \\
 & \quad \downarrow \text{60} \\
 & -2 \int \frac{\sec(x)}{3\sqrt{3 \cos(x) + 2}} d(3 \cos(x)) - 2\sqrt{3 \cos(x) + 2} \\
 & \quad \downarrow \text{73} \\
 & -4 \int \frac{1}{9 \cos^2(x) - 2} d\sqrt{3 \cos(x) + 2} - 2\sqrt{3 \cos(x) + 2} \\
 & \quad \downarrow \text{220} \\
 & 2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{3 \cos(x) + 2}}{\sqrt{2}}\right) - 2\sqrt{3 \cos(x) + 2}
 \end{aligned}$$

input `Int[Sqrt[2 + 3*Cos[x]]*Tan[x],x]`

output `2*Sqrt[2]*ArcTanh[Sqrt[2 + 3*Cos[x]]/Sqrt[2]] - 2*Sqrt[2 + 3*Cos[x]]`

3.359.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3200 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*tan[(e_.) + (f_.)*(x_)]^(p_.), x_Symbol] := Simp[1/f Subst[Int[(x^p*(a + x)^m)/(b^2 - x^2)^((p + 1)/2), x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && NeQ[a^2 - b^2, 0] && IntegerQ[(p + 1)/2]`

3.359.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 76 vs. $2(30) = 60$.

Time = 0.68 (sec) , antiderivative size = 77, normalized size of antiderivative = 2.08

method	result	size
default	$\sqrt{2} \operatorname{arctanh}\left(\frac{6 \cos\left(\frac{x}{2}\right) - \sqrt{2}}{2\sqrt{-6\left(\sin^2\left(\frac{x}{2}\right)\right) + 5}}\right) - \sqrt{2} \operatorname{arctanh}\left(\frac{6 \cos\left(\frac{x}{2}\right) + \sqrt{2}}{2\sqrt{-6\left(\sin^2\left(\frac{x}{2}\right)\right) + 5}}\right) - 2\sqrt{-6\left(\sin^2\left(\frac{x}{2}\right)\right) + 5}$	77

input `int((2+3*cos(x))^(1/2)*tan(x),x,method=_RETURNVERBOSE)`

output $2^{(1/2)} \operatorname{arctanh}\left(\frac{1}{2} / (-6 \sin(1/2 * x)^2 + 5)^{(1/2)} * (6 \cos(1/2 * x) - 2^{(1/2)})\right) - 2^{(1/2)} \operatorname{arctanh}\left(\frac{1}{2} / (-6 \sin(1/2 * x)^2 + 5)^{(1/2)} * (6 \cos(1/2 * x) + 2^{(1/2)})\right) - 2 * (-6 \sin(1/2 * x)^2 + 5)^{(1/2)}$

3.359.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.57

$$\int \sqrt{2 + 3 \cos(x)} \tan(x) dx$$

$$= \frac{1}{2} \sqrt{2} \log \left(-\frac{9 \cos(x)^2 + 4(3\sqrt{2} \cos(x) + 4\sqrt{2})\sqrt{3 \cos(x) + 2} + 48 \cos(x) + 32}{\cos(x)^2} \right) - 2\sqrt{3 \cos(x) + 2}$$

input `integrate((2+3*cos(x))^(1/2)*tan(x),x, algorithm="fracas")`

output $1/2 * \sqrt{2} * \log(-9 * \cos(x)^2 + 4 * (3 * \sqrt{2} * \cos(x) + 4 * \sqrt{2}) * \sqrt{3 * \cos(x) + 2} + 48 * \cos(x) + 32) / \cos(x)^2 - 2 * \sqrt{3 * \cos(x) + 2}$

3.359.6 Sympy [F]

$$\int \sqrt{2 + 3 \cos(x)} \tan(x) dx = \int \sqrt{3 \cos(x) + 2} \tan(x) dx$$

input `integrate((2+3*cos(x))**(1/2)*tan(x),x)`

output `Integral(sqrt(3*cos(x) + 2)*tan(x), x)`

3.359.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.27

$$\int \sqrt{2 + 3 \cos(x)} \tan(x) dx = -\sqrt{2} \log \left(-\frac{\sqrt{2} - \sqrt{3 \cos(x) + 2}}{\sqrt{2} + \sqrt{3 \cos(x) + 2}} \right) - 2 \sqrt{3 \cos(x) + 2}$$

input `integrate((2+3*cos(x))^(1/2)*tan(x),x, algorithm="maxima")`

output `-sqrt(2)*log(-(sqrt(2) - sqrt(3*cos(x) + 2))/(sqrt(2) + sqrt(3*cos(x) + 2))) - 2*sqrt(3*cos(x) + 2)`

3.359.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.35

$$\int \sqrt{2 + 3 \cos(x)} \tan(x) dx = -\sqrt{2} \log \left(\frac{|-2\sqrt{2} + 2\sqrt{3 \cos(x) + 2}|}{2(\sqrt{2} + \sqrt{3 \cos(x) + 2})} \right) - 2 \sqrt{3 \cos(x) + 2}$$

input `integrate((2+3*cos(x))^(1/2)*tan(x),x, algorithm="giac")`

output `-sqrt(2)*log(1/2*abs(-2*sqrt(2) + 2*sqrt(3*cos(x) + 2))/(sqrt(2) + sqrt(3*cos(x) + 2))) - 2*sqrt(3*cos(x) + 2)`

3.359.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{2 + 3 \cos(x)} \tan(x) dx = \int \tan(x) \sqrt{3 \cos(x) + 2} dx$$

input `int(tan(x)*(3*cos(x) + 2)^(1/2),x)`output `int(tan(x)*(3*cos(x) + 2)^(1/2), x)`

3.360 $\int \frac{x}{\sqrt{-4x+x^2}} dx$

3.360.1 Optimal result	1809
3.360.2 Mathematica [A] (verified)	1809
3.360.3 Rubi [A] (verified)	1810
3.360.4 Maple [A] (verified)	1811
3.360.5 Fricas [A] (verification not implemented)	1811
3.360.6 Sympy [A] (verification not implemented)	1812
3.360.7 Maxima [A] (verification not implemented)	1812
3.360.8 Giac [A] (verification not implemented)	1812
3.360.9 Mupad [B] (verification not implemented)	1813

3.360.1 Optimal result

Integrand size = 13, antiderivative size = 28

$$\int \frac{x}{\sqrt{-4x+x^2}} dx = \sqrt{-4x+x^2} + 4\operatorname{arctanh}\left(\frac{x}{\sqrt{-4x+x^2}}\right)$$

output `4*arctanh(x/(x^2-4*x)^(1/2))+(x^2-4*x)^(1/2)`

3.360.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.64

$$\int \frac{x}{\sqrt{-4x+x^2}} dx = \frac{(-4+x)x - 4\sqrt{-4+x}\sqrt{x}\log(\sqrt{-4+x} - \sqrt{x})}{\sqrt{(-4+x)x}}$$

input `Integrate[x/Sqrt[-4*x + x^2],x]`

output `((-4 + x)*x - 4*Sqrt[-4 + x]*Sqrt[x]*Log[Sqrt[-4 + x] - Sqrt[x]])/Sqrt[(-4 + x)*x]`

3.360.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {1160, 1091, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x}{\sqrt{x^2 - 4x}} dx \\ & \quad \downarrow \text{1160} \\ & 2 \int \frac{1}{\sqrt{x^2 - 4x}} dx + \sqrt{x^2 - 4x} \\ & \quad \downarrow \text{1091} \\ & 4 \int \frac{1}{1 - \frac{x^2}{x^2 - 4x}} d \frac{x}{\sqrt{x^2 - 4x}} + \sqrt{x^2 - 4x} \\ & \quad \downarrow \text{219} \\ & 4 \operatorname{arctanh} \left(\frac{x}{\sqrt{x^2 - 4x}} \right) + \sqrt{x^2 - 4x} \end{aligned}$$

input `Int[x/Sqrt[-4*x + x^2],x]`

output `Sqrt[-4*x + x^2] + 4*ArcTanh[x/Sqrt[-4*x + x^2]]`

3.360.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1091 `Int[1/Sqrt[(b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Simp[2 Subst[Int[1/(1 - c*x^2), x], x, x/Sqrt[b*x + c*x^2]], x] /; FreeQ[{b, c}, x]`

```
rule 1160 Int[((d._) + (e._)*(x_))*((a._) + (b._)*(x_) + (c._)*(x_)^2)^(p_), x_Symbol
] := Simp[e*((a + b*x + c*x^2)^(p + 1)/(2*c*(p + 1))), x] + Simp[(2*c*d - b
*e)/(2*c) Int[(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x]
&& NeQ[p, -1]
```

3.360.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

method	result	size
default	$\sqrt{x^2 - 4x} + 2 \ln(-2 + x + \sqrt{x^2 - 4x})$	26
trager	$\sqrt{x^2 - 4x} - 2 \ln(2 - x + \sqrt{x^2 - 4x})$	28
risch	$\frac{x(x-4)}{\sqrt{x(x-4)}} + 2 \ln(-2 + x + \sqrt{x^2 - 4x})$	29
pseudoelliptic	$\sqrt{x(x-4)} + 2 \ln\left(\frac{\sqrt{x(x-4)+x}}{x}\right) - 2 \ln\left(\frac{\sqrt{x(x-4)-x}}{x}\right)$	43
meijerg	$\frac{4i\sqrt{-\text{signum}(x-4)}\left(\frac{i\sqrt{\pi}\sqrt{x}\sqrt{-\frac{x}{4}+1}}{2} - i\sqrt{\pi}\arcsin\left(\frac{\sqrt{x}}{2}\right)\right)}{\sqrt{\pi}\sqrt{\text{signum}(x-4)}}$	50

```
input int(x/(x^2-4*x)^(1/2),x,method=_RETURNVERBOSE)
```

```
output (x^2-4*x)^(1/2)+2*ln(-2+x+(x^2-4*x)^(1/2))
```

3.360.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{x}{\sqrt{-4x+x^2}} dx = \sqrt{x^2 - 4x} - 2 \log(-x + \sqrt{x^2 - 4x} + 2)$$

```
input integrate(x/(x^2-4*x)^(1/2),x, algorithm="fricas")
```

```
output sqrt(x^2 - 4*x) - 2*log(-x + sqrt(x^2 - 4*x) + 2)
```

3.360.6 Sympy [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{x}{\sqrt{-4x+x^2}} dx = \sqrt{x^2-4x} + 2 \log(2x + 2\sqrt{x^2-4x} - 4)$$

input `integrate(x/(x**2-4*x)**(1/2),x)`output `sqrt(x**2 - 4*x) + 2*log(2*x + 2*sqrt(x**2 - 4*x) - 4)`**3.360.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{x}{\sqrt{-4x+x^2}} dx = \sqrt{x^2-4x} + 2 \log(2x + 2\sqrt{x^2-4x} - 4)$$

input `integrate(x/(x^2-4*x)^(1/2),x, algorithm="maxima")`output `sqrt(x^2 - 4*x) + 2*log(2*x + 2*sqrt(x^2 - 4*x) - 4)`**3.360.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x}{\sqrt{-4x+x^2}} dx = \sqrt{x^2-4x} - 2 \log\left(\left|-x + \sqrt{x^2-4x} + 2\right|\right)$$

input `integrate(x/(x^2-4*x)^(1/2),x, algorithm="giac")`output `sqrt(x^2 - 4*x) - 2*log(abs(-x + sqrt(x^2 - 4*x) + 2))`

3.360.9 Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

$$\int \frac{x}{\sqrt{-4x+x^2}} dx = 2 \ln \left(x + \sqrt{x(x-4)} - 2 \right) + \sqrt{x^2 - 4x}$$

input `int(x/(x^2 - 4*x)^(1/2),x)`

output `2*log(x + (x*(x - 4))^(1/2) - 2) + (x^2 - 4*x)^(1/2)`

3.361 $\int \cos^5(x) dx$

3.361.1 Optimal result	1814
3.361.2 Mathematica [A] (verified)	1814
3.361.3 Rubi [A] (verified)	1815
3.361.4 Maple [A] (verified)	1816
3.361.5 Fricas [A] (verification not implemented)	1816
3.361.6 Sympy [A] (verification not implemented)	1816
3.361.7 Maxima [A] (verification not implemented)	1817
3.361.8 Giac [A] (verification not implemented)	1817
3.361.9 Mupad [B] (verification not implemented)	1817

3.361.1 Optimal result

Integrand size = 4, antiderivative size = 19

$$\int \cos^5(x) dx = \sin(x) - \frac{2 \sin^3(x)}{3} + \frac{\sin^5(x)}{5}$$

output `sin(x)-2/3*sin(x)^3+1/5*sin(x)^5`

3.361.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \cos^5(x) dx = \sin(x) - \frac{2 \sin^3(x)}{3} + \frac{\sin^5(x)}{5}$$

input `Integrate[Cos[x]^5,x]`

output `Sin[x] - (2*Sin[x]^3)/3 + Sin[x]^5/5`

3.361.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^5(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin\left(x + \frac{\pi}{2}\right)^5 dx \\ & \quad \downarrow \text{3113} \\ & - \int (\sin^4(x) - 2\sin^2(x) + 1) d(-\sin(x)) \\ & \quad \downarrow \text{2009} \\ & \frac{\sin^5(x)}{5} - \frac{2\sin^3(x)}{3} + \sin(x) \end{aligned}$$

input `Int[Cos[x]^5,x]`

output `Sin[x] - (2*Sin[x]^3)/3 + Sin[x]^5/5`

3.361.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

3.361.4 Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

method	result	size
default	$\frac{\left(\frac{8}{3} + \cos^4(x) + \frac{4(\cos^2(x))}{3}\right) \sin(x)}{5}$	17
risch	$\frac{5 \sin(x)}{8} + \frac{\sin(5x)}{80} + \frac{5 \sin(3x)}{48}$	18
parallelrisc	$\frac{5 \sin(x)}{8} + \frac{\sin(5x)}{80} + \frac{5 \sin(3x)}{48}$	18

input `int(cos(x)^5,x,method=_RETURNVERBOSE)`output `1/5*(8/3+cos(x)^4+4/3*cos(x)^2)*sin(x)`**3.361.5 Fricas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

$$\int \cos^5(x) dx = \frac{1}{15} (3 \cos(x)^4 + 4 \cos(x)^2 + 8) \sin(x)$$

input `integrate(cos(x)^5,x, algorithm="fricas")`output `1/15*(3*cos(x)^4 + 4*cos(x)^2 + 8)*sin(x)`**3.361.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \cos^5(x) dx = \frac{\sin^5(x)}{5} - \frac{2 \sin^3(x)}{3} + \sin(x)$$

input `integrate(cos(x)**5,x)`output `sin(x)**5/5 - 2*sin(x)**3/3 + sin(x)`

3.361.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \cos^5(x) dx = \frac{1}{5} \sin(x)^5 - \frac{2}{3} \sin(x)^3 + \sin(x)$$

input `integrate(cos(x)^5,x, algorithm="maxima")`output `1/5*sin(x)^5 - 2/3*sin(x)^3 + sin(x)`**3.361.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \cos^5(x) dx = \frac{1}{5} \sin(x)^5 - \frac{2}{3} \sin(x)^3 + \sin(x)$$

input `integrate(cos(x)^5,x, algorithm="giac")`output `1/5*sin(x)^5 - 2/3*sin(x)^3 + sin(x)`**3.361.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \cos^5(x) dx = \frac{\sin(x) \cos(x)^4}{5} + \frac{4 \sin(x) \cos(x)^2}{15} + \frac{8 \sin(x)}{15}$$

input `int(cos(x)^5,x)`output `(8*sin(x))/15 + (4*cos(x)^2*sin(x))/15 + (cos(x)^4*sin(x))/5`

3.362 $\int e^{-x}x^4 dx$

3.362.1 Optimal result	1818
3.362.2 Mathematica [A] (verified)	1818
3.362.3 Rubi [A] (verified)	1819
3.362.4 Maple [A] (verified)	1820
3.362.5 Fricas [A] (verification not implemented)	1820
3.362.6 Sympy [A] (verification not implemented)	1821
3.362.7 Maxima [A] (verification not implemented)	1821
3.362.8 Giac [A] (verification not implemented)	1821
3.362.9 Mupad [B] (verification not implemented)	1822

3.362.1 Optimal result

Integrand size = 9, antiderivative size = 46

$$\int e^{-x}x^4 dx = -24e^{-x} - 24e^{-x}x - 12e^{-x}x^2 - 4e^{-x}x^3 - e^{-x}x^4$$

output `-24/exp(x)-24*x/exp(x)-12*x^2/exp(x)-4*x^3/exp(x)-x^4/exp(x)`

3.362.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.57

$$\int e^{-x}x^4 dx = e^{-x}(-24 - 24x - 12x^2 - 4x^3 - x^4)$$

input `Integrate[x^4/E^x,x]`

output `(-24 - 24*x - 12*x^2 - 4*x^3 - x^4)/E^x`

3.362.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.20, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {2607, 2607, 2607, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{-x} x^4 dx \\
 & \quad \downarrow \text{2607} \\
 & 4 \int e^{-x} x^3 dx - e^{-x} x^4 \\
 & \quad \downarrow \text{2607} \\
 & 4 \left(3 \int e^{-x} x^2 dx - e^{-x} x^3 \right) - e^{-x} x^4 \\
 & \quad \downarrow \text{2607} \\
 & 4 \left(3 \left(2 \int e^{-x} x dx - e^{-x} x^2 \right) - e^{-x} x^3 \right) - e^{-x} x^4 \\
 & \quad \downarrow \text{2607} \\
 & 4 \left(3 \left(2 \left(\int e^{-x} dx - e^{-x} x \right) - e^{-x} x^2 \right) - e^{-x} x^3 \right) - e^{-x} x^4 \\
 & \quad \downarrow \text{2624} \\
 & 4 \left(3 \left(2 \left(-e^{-x} x - e^{-x} \right) - e^{-x} x^2 \right) - e^{-x} x^3 \right) - e^{-x} x^4
 \end{aligned}$$

input `Int[x^4/E^x,x]`

output `-(x^4/E^x) + 4*(-(x^3/E^x) + 3*(-(x^2/E^x) + 2*(-E^(-x) - x/E^x)))`

3.362.3.1 Defintions of rubi rules used

```
rule 2607 Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m
_.), x_Symbol] := Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])),
x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^
n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*
m] && !TrueQ[$UseGamma]
```

```
rule 2624 Int[((F_)^(v_))^(n_), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;
FreeQ[{F, n}, x] && LinearQ[v, x]
```

3.362.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.54

method	result	size
gospers	$-(x^4 + 4x^3 + 12x^2 + 24x + 24)e^{-x}$	25
norman	$(-x^4 - 4x^3 - 12x^2 - 24x - 24)e^{-x}$	26
risch	$(-x^4 - 4x^3 - 12x^2 - 24x - 24)e^{-x}$	26
paralelrisch	$(-x^4 - 4x^3 - 12x^2 - 24x - 24)e^{-x}$	26
meijerg	$24 - \frac{(5x^4 + 20x^3 + 60x^2 + 120x + 120)e^{-x}}{5}$	29
default	$-24e^{-x} - 24xe^{-x} - 12x^2e^{-x} - 4x^3e^{-x} - x^4e^{-x}$	42

```
input int(x^4/exp(x),x,method=_RETURNVERBOSE)
```

```
output -(x^4+4*x^3+12*x^2+24*x+24)/exp(x)
```

3.362.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.52

$$\int e^{-x}x^4 dx = -(x^4 + 4x^3 + 12x^2 + 24x + 24)e^{(-x)}$$

```
input integrate(x^4/exp(x),x, algorithm="fracas")
```

output $-(x^4 + 4x^3 + 12x^2 + 24x + 24)e^{-x}$

3.362.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.48

$$\int e^{-x}x^4 dx = (-x^4 - 4x^3 - 12x^2 - 24x - 24)e^{-x}$$

input `integrate(x**4/exp(x),x)`

output $(-x^{**4} - 4*x^{**3} - 12*x^{**2} - 24*x - 24)*exp(-x)$

3.362.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.52

$$\int e^{-x}x^4 dx = -(x^4 + 4x^3 + 12x^2 + 24x + 24)e^{(-x)}$$

input `integrate(x^4/exp(x),x, algorithm="maxima")`

output $-(x^4 + 4x^3 + 12x^2 + 24x + 24)e^{-x}$

3.362.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.52

$$\int e^{-x}x^4 dx = -(x^4 + 4x^3 + 12x^2 + 24x + 24)e^{(-x)}$$

input `integrate(x^4/exp(x),x, algorithm="giac")`

output $-(x^4 + 4x^3 + 12x^2 + 24x + 24)e^{-x}$

3.362.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.52

$$\int e^{-x} x^4 dx = -e^{-x} (x^4 + 4x^3 + 12x^2 + 24x + 24)$$

input `int(x^4*exp(-x),x)`

output `-exp(-x)*(24*x + 12*x^2 + 4*x^3 + x^4 + 24)`

3.363 $\int \frac{x^4}{\sqrt{-2+x^{10}}} dx$

3.363.1 Optimal result	1823
3.363.2 Mathematica [A] (verified)	1823
3.363.3 Rubi [A] (verified)	1824
3.363.4 Maple [A] (verified)	1825
3.363.5 Fricas [A] (verification not implemented)	1825
3.363.6 Sympy [C] (verification not implemented)	1826
3.363.7 Maxima [B] (verification not implemented)	1826
3.363.8 Giac [B] (verification not implemented)	1826
3.363.9 Mupad [F(-1)]	1827

3.363.1 Optimal result

Integrand size = 13, antiderivative size = 18

$$\int \frac{x^4}{\sqrt{-2+x^{10}}} dx = \frac{1}{5} \operatorname{arctanh}\left(\frac{x^5}{\sqrt{-2+x^{10}}}\right)$$

output `1/5*arctanh(x^5/(x^10-2)^(1/2))`

3.363.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{x^4}{\sqrt{-2+x^{10}}} dx = \frac{1}{5} \log\left(x^5 + \sqrt{-2+x^{10}}\right)$$

input `Integrate[x^4/Sqrt[-2 + x^10],x]`

output `Log[x^5 + Sqrt[-2 + x^10]]/5`

3.363.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {807, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^4}{\sqrt{x^{10}-2}} dx \\ & \quad \downarrow 807 \\ & \frac{1}{5} \int \frac{1}{\sqrt{x^{10}-2}} dx^5 \\ & \quad \downarrow 224 \\ & \frac{1}{5} \int \frac{1}{1-x^{10}} d \frac{x^5}{\sqrt{x^{10}-2}} \\ & \quad \downarrow 219 \\ & \frac{1}{5} \operatorname{arctanh} \left(\frac{x^5}{\sqrt{x^{10}-2}} \right) \end{aligned}$$

input `Int[x^4/Sqrt[-2 + x^10],x]`

output `ArcTanh[x^5/Sqrt[-2 + x^10]]/5`

3.363.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

```
rule 807 Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m
+ 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x,
x^k], x] /; k != 1 /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

3.363.4 Maple [A] (verified)

Time = 0.74 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

method	result	size
trager	$\frac{\ln(x^5 + \sqrt{x^{10} - 2})}{5}$	15
pseudoelliptic	$\frac{\ln(x^5 + \sqrt{x^{10} - 2})}{5}$	15
meijerg	$\frac{\sqrt{-\operatorname{signum}\left(-1 + \frac{x^{10}}{2}\right)} \arcsin\left(\frac{x^5 \sqrt{2}}{2}\right)}{5 \sqrt{\operatorname{signum}\left(-1 + \frac{x^{10}}{2}\right)}}$	34

```
input int(x^4/(x^10-2)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/5*ln(x^5+(x^10-2)^(1/2))
```

3.363.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \frac{x^4}{\sqrt{-2 + x^{10}}} dx = -\frac{1}{5} \log\left(-x^5 + \sqrt{x^{10} - 2}\right)$$

```
input integrate(x^4/(x^10-2)^(1/2),x, algorithm="fracas")
```

```
output -1/5*log(-x^5 + sqrt(x^10 - 2))
```

3.363.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.53 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.78

$$\int \frac{x^4}{\sqrt{-2+x^{10}}} dx = \begin{cases} \frac{\operatorname{acosh}\left(\frac{\sqrt{2}x^5}{2}\right)}{5} & \text{for } |x^{10}| > 2 \\ -\frac{i \operatorname{asin}\left(\frac{\sqrt{2}x^5}{2}\right)}{5} & \text{otherwise} \end{cases}$$

input `integrate(x**4/(x**10-2)**(1/2),x)`

output `Piecewise((acosh(sqrt(2)*x**5/2)/5, Abs(x**10) > 2), (-I*asin(sqrt(2)*x**5/2)/5, True))`

3.363.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 33 vs. 2(14) = 28.

Time = 0.21 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.83

$$\int \frac{x^4}{\sqrt{-2+x^{10}}} dx = \frac{1}{10} \log\left(\frac{\sqrt{x^{10}-2}}{x^5} + 1\right) - \frac{1}{10} \log\left(\frac{\sqrt{x^{10}-2}}{x^5} - 1\right)$$

input `integrate(x^4/(x^10-2)^(1/2),x, algorithm="maxima")`

output `1/10*log(sqrt(x^10 - 2)/x^5 + 1) - 1/10*log(sqrt(x^10 - 2)/x^5 - 1)`

3.363.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 30 vs. 2(14) = 28.

Time = 0.30 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.67

$$\int \frac{x^4}{\sqrt{-2+x^{10}}} dx = \frac{1}{10} \sqrt{x^{10}-2}x^5 + \frac{1}{5} \log\left(\left|-x^5 + \sqrt{x^{10}-2}\right|\right)$$

input `integrate(x^4/(x^10-2)^(1/2),x, algorithm="giac")`

output `1/10*sqrt(x^10 - 2)*x^5 + 1/5*log(abs(-x^5 + sqrt(x^10 - 2)))`

3.363.9 Mupad [F(-1)]

Timed out.

$$\int \frac{x^4}{\sqrt{-2+x^{10}}} dx = \int \frac{x^4}{\sqrt{x^{10}-2}} dx$$

input `int(x^4/(x^10 - 2)^(1/2),x)`output `int(x^4/(x^10 - 2)^(1/2), x)`

3.364 $\int e^x \cos(4 + 3x) dx$

3.364.1 Optimal result	1828
3.364.2 Mathematica [A] (verified)	1828
3.364.3 Rubi [A] (verified)	1829
3.364.4 Maple [A] (verified)	1829
3.364.5 Fricas [A] (verification not implemented)	1830
3.364.6 Sympy [A] (verification not implemented)	1830
3.364.7 Maxima [A] (verification not implemented)	1830
3.364.8 Giac [A] (verification not implemented)	1831
3.364.9 Mupad [B] (verification not implemented)	1831

3.364.1 Optimal result

Integrand size = 10, antiderivative size = 27

$$\int e^x \cos(4 + 3x) dx = \frac{1}{10} e^x \cos(4 + 3x) + \frac{3}{10} e^x \sin(4 + 3x)$$

output `1/10*exp(x)*cos(4+3*x)+3/10*exp(x)*sin(4+3*x)`

3.364.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int e^x \cos(4 + 3x) dx = \frac{1}{10} e^x (\cos(4 + 3x) + 3 \sin(4 + 3x))$$

input `Integrate[E^x*Cos[4 + 3*x],x]`

output `(E^x*(Cos[4 + 3*x] + 3*Sin[4 + 3*x]))/10`

3.364.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {4933}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x \cos(3x + 4) dx$$

$$\downarrow \text{4933}$$

$$\frac{3}{10}e^x \sin(3x + 4) + \frac{1}{10}e^x \cos(3x + 4)$$

input `Int[E^x*Cos[4 + 3*x],x]`

output `(E^x*Cos[4 + 3*x])/10 + (3*E^x*Sin[4 + 3*x])/10`

3.364.3.1 Defintions of rubi rules used

rule 4933 `Int[Cos[(d_.) + (e_.)*(x_.)]*(F_)^((c_.)*((a_.) + (b_.)*(x_))), x_Symbol] :>
Simp[b*c*Log[F]*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x
] + Simp[e*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; F
reeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]`

3.364.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result	size
parallelrisch	$\frac{e^x (\cos(3x+4) + 3 \sin(3x+4))}{10}$	20
default	$\frac{e^x \cos(3x+4)}{10} + \frac{3 e^x \sin(3x+4)}{10}$	22
risch	$\left(\frac{1}{20} - \frac{3i}{20}\right) e^x e^{3ix} e^{4i} + \left(\frac{1}{20} + \frac{3i}{20}\right) e^x e^{-3ix} e^{-4i}$	30
norman	$\frac{3 e^x \tan\left(\frac{3x}{2}+2\right) - \frac{e^x \left(\tan^2\left(\frac{3x}{2}+2\right)\right)}{10} + \frac{e^x}{10}}{1 + \tan^2\left(\frac{3x}{2}+2\right)}$	41

input `int(exp(x)*cos(3*x+4),x,method=_RETURNVERBOSE)`

output `1/10*exp(x)*(cos(3*x+4)+3*sin(3*x+4))`

3.364.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int e^x \cos(4 + 3x) dx = \frac{1}{10} \cos(3x + 4) e^x + \frac{3}{10} e^x \sin(3x + 4)$$

input `integrate(exp(x)*cos(4+3*x),x, algorithm="fricas")`

output `1/10*cos(3*x + 4)*e^x + 3/10*e^x*sin(3*x + 4)`

3.364.6 Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

$$\int e^x \cos(4 + 3x) dx = \frac{3e^x \sin(3x + 4)}{10} + \frac{e^x \cos(3x + 4)}{10}$$

input `integrate(exp(x)*cos(4+3*x),x)`

output `3*exp(x)*sin(3*x + 4)/10 + exp(x)*cos(3*x + 4)/10`

3.364.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^x \cos(4 + 3x) dx = \frac{1}{10} (\cos(3x + 4) + 3 \sin(3x + 4)) e^x$$

input `integrate(exp(x)*cos(4+3*x),x, algorithm="maxima")`

output `1/10*(cos(3*x + 4) + 3*sin(3*x + 4))*e^x`

3.364.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^x \cos(4 + 3x) dx = \frac{1}{10} (\cos(3x + 4) + 3 \sin(3x + 4))e^x$$

input `integrate(exp(x)*cos(4+3*x),x, algorithm="giac")`

output `1/10*(cos(3*x + 4) + 3*sin(3*x + 4))*e^x`

3.364.9 Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^x \cos(4 + 3x) dx = \frac{e^x (\cos(3x + 4) + 3 \sin(3x + 4))}{10}$$

input `int(exp(x)*cos(3*x + 4),x)`

output `(exp(x)*(cos(3*x + 4) + 3*sin(3*x + 4)))/10`

3.365 $\int e^x \log(1 + e^x) dx$

3.365.1 Optimal result	1832
3.365.2 Mathematica [A] (verified)	1832
3.365.3 Rubi [A] (verified)	1833
3.365.4 Maple [A] (verified)	1834
3.365.5 Fracas [A] (verification not implemented)	1834
3.365.6 Sympy [F(-1)]	1835
3.365.7 Maxima [A] (verification not implemented)	1835
3.365.8 Giac [A] (verification not implemented)	1835
3.365.9 Mupad [B] (verification not implemented)	1836

3.365.1 Optimal result

Integrand size = 10, antiderivative size = 18

$$\int e^x \log(1 + e^x) dx = -e^x + (1 + e^x) \log(1 + e^x)$$

output `-exp(x)+(1+exp(x))*ln(1+exp(x))`

3.365.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int e^x \log(1 + e^x) dx = -e^x + (1 + e^x) \log(1 + e^x)$$

input `Integrate[E^x*Log[1 + E^x],x]`

output `-E^x + (1 + E^x)*Log[1 + E^x]`

3.365.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3034, 2678, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^x \log(e^x + 1) dx \\
 & \quad \downarrow \text{3034} \\
 & e^x \log(e^x + 1) - \int \frac{e^{2x}}{1 + e^x} dx \\
 & \quad \downarrow \text{2678} \\
 & e^x \log(e^x + 1) - \int \frac{e^x}{1 + e^x} de^x \\
 & \quad \downarrow \text{49} \\
 & e^x \log(e^x + 1) - \int \left(1 + \frac{1}{-1 - e^x}\right) de^x \\
 & \quad \downarrow \text{2009} \\
 & -e^x + e^x \log(e^x + 1) + \log(e^x + 1)
 \end{aligned}$$

input `Int[E^x*Log[1 + E^x],x]`

output `-E^x + Log[1 + E^x] + E^x*Log[1 + E^x]`

3.365.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 2678 Int[((a_) + (b_)*(F_)^((e_)*((c_) + (d_)*(x_)))^(p_))* (G_)^((h_)*((f_
.) + (g_)*(x_))), x_Symbol] := With[{m = FullSimplify[g*h*(Log[G]/(d*e*Log
[F]))}], Simp[Denominator[m]*(G^(f*h - c*g*(h/d))/(d*e*Log[F])) Subst[Int
[x^(Numerator[m] - 1)*(a + b*x^Denominator[m])^p, x], x, F^(e*((c + d*x)/De
nominator[m]))], x] /; LeQ[m, -1] || GeQ[m, 1]] /; FreeQ[{F, G, a, b, c, d,
e, f, g, h, p}, x]
```

```
rule 3034 Int[Log[u_]*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[Log[u] w, x
] - Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w,
x]] /; InverseFunctionFreeQ[u, x]
```

3.365.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

method	result	size
derivativedivides	$(1 + e^x) \ln(1 + e^x) - 1 - e^x$	17
default	$(1 + e^x) \ln(1 + e^x) - 1 - e^x$	17
norman	$e^x \ln(1 + e^x) - e^x + \ln(1 + e^x)$	19
risch	$e^x \ln(1 + e^x) - e^x + \ln(1 + e^x)$	19
parallelrisch	$e^x \ln(1 + e^x) - e^x + \ln(1 + e^x) + 1$	20

```
input int(exp(x)*ln(1+exp(x)),x,method=_RETURNVERBOSE)
```

```
output (1+exp(x))*ln(1+exp(x))-1-exp(x)
```

3.365.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int e^x \log(1 + e^x) dx = (e^x + 1) \log(e^x + 1) - e^x$$

```
input integrate(exp(x)*log(1+exp(x)),x, algorithm="fracas")
```

```
output (e^x + 1)*log(e^x + 1) - e^x
```

3.365.6 Sympy [F(-1)]

Timed out.

$$\int e^x \log(1 + e^x) dx = \text{Timed out}$$

input `integrate(exp(x)*ln(1+exp(x)),x)`output `Timed out`**3.365.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int e^x \log(1 + e^x) dx = (e^x + 1) \log(e^x + 1) - e^x - 1$$

input `integrate(exp(x)*log(1+exp(x)),x, algorithm="maxima")`output `(e^x + 1)*log(e^x + 1) - e^x - 1`**3.365.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int e^x \log(1 + e^x) dx = (e^x + 1) \log(e^x + 1) - e^x - 1$$

input `integrate(exp(x)*log(1+exp(x)),x, algorithm="giac")`output `(e^x + 1)*log(e^x + 1) - e^x - 1`

3.365.9 Mupad [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int e^x \log(1 + e^x) dx = \ln(e^x + 1) - e^x + e^x \ln(e^x + 1)$$

input `int(exp(x)*log(exp(x) + 1),x)`

output `log(exp(x) + 1) - exp(x) + exp(x)*log(exp(x) + 1)`

3.366 $\int x^2 \arctan(x) dx$

3.366.1 Optimal result	1837
3.366.2 Mathematica [A] (verified)	1837
3.366.3 Rubi [A] (verified)	1838
3.366.4 Maple [A] (verified)	1839
3.366.5 Fricas [A] (verification not implemented)	1839
3.366.6 Sympy [A] (verification not implemented)	1840
3.366.7 Maxima [A] (verification not implemented)	1840
3.366.8 Giac [A] (verification not implemented)	1840
3.366.9 Mupad [B] (verification not implemented)	1841

3.366.1 Optimal result

Integrand size = 6, antiderivative size = 27

$$\int x^2 \arctan(x) dx = -\frac{x^2}{6} + \frac{1}{3}x^3 \arctan(x) + \frac{1}{6} \log(1 + x^2)$$

output `-1/6*x^2+1/3*x^3*arctan(x)+1/6*ln(x^2+1)`

3.366.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int x^2 \arctan(x) dx = \frac{1}{6}(-x^2 + 2x^3 \arctan(x) + \log(1 + x^2))$$

input `Integrate[x^2*ArcTan[x],x]`

output `(-x^2 + 2*x^3*ArcTan[x] + Log[1 + x^2])/6`

3.366.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5361, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \arctan(x) dx \\
 & \quad \downarrow \text{5361} \\
 & \frac{1}{3}x^3 \arctan(x) - \frac{1}{3} \int \frac{x^3}{x^2+1} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{3}x^3 \arctan(x) - \frac{1}{6} \int \frac{x^2}{x^2+1} dx \\
 & \quad \downarrow \text{49} \\
 & \frac{1}{3}x^3 \arctan(x) - \frac{1}{6} \int \left(1 + \frac{1}{-x^2-1}\right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{3}x^3 \arctan(x) + \frac{1}{6}(\log(x^2+1) - x^2)
 \end{aligned}$$

input `Int[x^2*ArcTan[x],x]`

output `(x^3*ArcTan[x])/3 + (-x^2 + Log[1 + x^2])/6`

3.366.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5361 `Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m +
1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x],
x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] &
& IntegerQ[m])) && NeQ[m, -1]`

3.366.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

method	result	size
default	$-\frac{x^2}{6} + \frac{x^3 \arctan(x)}{3} + \frac{\ln(x^2+1)}{6}$	22
parts	$-\frac{x^2}{6} + \frac{x^3 \arctan(x)}{3} + \frac{\ln(x^2+1)}{6}$	22
parallelrisch	$\frac{x^3 \arctan(x)}{3} - \frac{x^2}{6} + \frac{\ln(x^2+1)}{6} + \frac{1}{6}$	23
meijerg	$-\frac{x^2}{6} + \frac{x^4 \arctan(\sqrt{x^2})}{3\sqrt{x^2}} + \frac{\ln(x^2+1)}{6}$	31
risch	$-\frac{ix^3 \ln(ix+1)}{6} + \frac{ix^3 \ln(-ix+1)}{6} - \frac{x^2}{6} + \frac{\ln(x^2+1)}{6}$	41

input `int(x^2*arctan(x),x,method=_RETURNVERBOSE)`

output `-1/6*x^2+1/3*x^3*arctan(x)+1/6*ln(x^2+1)`

3.366.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int x^2 \arctan(x) dx = \frac{1}{3} x^3 \arctan(x) - \frac{1}{6} x^2 + \frac{1}{6} \log(x^2 + 1)$$

input `integrate(x^2*arctan(x),x, algorithm="fricas")`

output `1/3*x^3*arctan(x) - 1/6*x^2 + 1/6*log(x^2 + 1)`

3.366.6 Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

$$\int x^2 \arctan(x) dx = \frac{x^3 \operatorname{atan}(x)}{3} - \frac{x^2}{6} + \frac{\log(x^2 + 1)}{6}$$

input `integrate(x**2*atan(x),x)`output `x**3*atan(x)/3 - x**2/6 + log(x**2 + 1)/6`**3.366.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int x^2 \arctan(x) dx = \frac{1}{3} x^3 \arctan(x) - \frac{1}{6} x^2 + \frac{1}{6} \log(x^2 + 1)$$

input `integrate(x^2*arctan(x),x, algorithm="maxima")`output `1/3*x^3*arctan(x) - 1/6*x^2 + 1/6*log(x^2 + 1)`**3.366.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int x^2 \arctan(x) dx = \frac{1}{3} x^3 \arctan(x) - \frac{1}{6} x^2 + \frac{1}{6} \log(x^2 + 1)$$

input `integrate(x^2*arctan(x),x, algorithm="giac")`output `1/3*x^3*arctan(x) - 1/6*x^2 + 1/6*log(x^2 + 1)`

3.366.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int x^2 \arctan(x) dx = \frac{\ln(x^2 + 1)}{6} + \frac{x^3 \operatorname{atan}(x)}{3} - \frac{x^2}{6}$$

input `int(x^2*atan(x),x)`

output `log(x^2 + 1)/6 + (x^3*atan(x))/3 - x^2/6`

3.367 $\int \sqrt{-1 + e^{2x}} dx$

3.367.1 Optimal result	1842
3.367.2 Mathematica [A] (verified)	1842
3.367.3 Rubi [A] (verified)	1843
3.367.4 Maple [A] (verified)	1844
3.367.5 Fricas [A] (verification not implemented)	1845
3.367.6 Sympy [A] (verification not implemented)	1845
3.367.7 Maxima [A] (verification not implemented)	1845
3.367.8 Giac [A] (verification not implemented)	1846
3.367.9 Mupad [B] (verification not implemented)	1846

3.367.1 Optimal result

Integrand size = 11, antiderivative size = 26

$$\int \sqrt{-1 + e^{2x}} dx = \sqrt{-1 + e^{2x}} - \arctan \left(\sqrt{-1 + e^{2x}} \right)$$

output `-arctan((-1+exp(2*x))^(1/2))+(-1+exp(2*x))^(1/2)`

3.367.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \sqrt{-1 + e^{2x}} dx = \sqrt{-1 + e^{2x}} - \arctan \left(\sqrt{-1 + e^{2x}} \right)$$

input `Integrate[Sqrt[-1 + E^(2*x)],x]`

output `Sqrt[-1 + E^(2*x)] - ArcTan[Sqrt[-1 + E^(2*x)]]`

3.367.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.23, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {2720, 60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{e^{2x} - 1} dx \\
 & \quad \downarrow \text{2720} \\
 & \frac{1}{2} \int e^{-2x} \sqrt{-1 + e^{2x}} de^{2x} \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2} \left(2\sqrt{e^{2x} - 1} - \int \frac{e^{-2x}}{\sqrt{-1 + e^{2x}}} de^{2x} \right) \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{2} \left(2\sqrt{e^{2x} - 1} - 2 \int \frac{1}{1 + e^{4x}} d\sqrt{-1 + e^{2x}} \right) \\
 & \quad \downarrow \text{216} \\
 & \frac{1}{2} \left(2\sqrt{e^{2x} - 1} - 2 \arctan \left(\sqrt{e^{2x} - 1} \right) \right)
 \end{aligned}$$

input `Int[Sqrt[-1 + E^(2*x)],x]`

output `(2*Sqrt[-1 + E^(2*x)] - 2*ArcTan[Sqrt[-1 + E^(2*x)]])/2`

3.367.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] -> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
  {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
  d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
  Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
  inearQ[a, b, c, d, m, n, x]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
  rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
  , 0] || GtQ[b, 0])
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
  ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
  [{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
  *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.367.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$-\arctan(\sqrt{e^{2x}-1}) + \sqrt{e^{2x}-1}$	21
default	$-\arctan(\sqrt{e^{2x}-1}) + \sqrt{e^{2x}-1}$	21
risch	$-\arctan(\sqrt{e^{2x}-1}) + \sqrt{e^{2x}-1}$	21

```
input int((exp(2*x)-1)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -arctan((exp(2*x)-1)^(1/2))+exp(2*x)-1)^(1/2)
```

3.367.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \sqrt{-1 + e^{2x}} dx = \sqrt{e^{(2x)} - 1} - \arctan\left(\sqrt{e^{(2x)} - 1}\right)$$

input `integrate((-1+exp(2*x))^(1/2),x, algorithm="fricas")`output `sqrt(e^(2*x) - 1) - arctan(sqrt(e^(2*x) - 1))`**3.367.6 Sympy [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \sqrt{-1 + e^{2x}} dx = \sqrt{e^{2x} - 1} - \operatorname{atan}\left(\sqrt{e^{2x} - 1}\right)$$

input `integrate((-1+exp(2*x))**(1/2),x)`output `sqrt(exp(2*x) - 1) - atan(sqrt(exp(2*x) - 1))`**3.367.7 Maxima [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \sqrt{-1 + e^{2x}} dx = \sqrt{e^{(2x)} - 1} - \arctan\left(\sqrt{e^{(2x)} - 1}\right)$$

input `integrate((-1+exp(2*x))^(1/2),x, algorithm="maxima")`output `sqrt(e^(2*x) - 1) - arctan(sqrt(e^(2*x) - 1))`

3.367.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \sqrt{-1 + e^{2x}} dx = \sqrt{e^{(2x)} - 1} - \arctan\left(\sqrt{e^{(2x)} - 1}\right)$$

input `integrate((-1+exp(2*x))^(1/2),x, algorithm="giac")`output `sqrt(e^(2*x) - 1) - arctan(sqrt(e^(2*x) - 1))`**3.367.9 Mupad [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.19

$$\int \sqrt{-1 + e^{2x}} dx = \sqrt{e^{2x} - 1} \left(\frac{e^{-x} \operatorname{asin}(e^{-x})}{\sqrt{1 - e^{-2x}}} + 1 \right)$$

input `int((exp(2*x) - 1)^(1/2),x)`output `(exp(2*x) - 1)^(1/2)*((exp(-x)*asin(exp(-x)))/(1 - exp(-2*x))^(1/2) + 1)`

3.368 $\int e^{\sin(x)} \sin(2x) dx$

3.368.1 Optimal result	1847
3.368.2 Mathematica [A] (verified)	1847
3.368.3 Rubi [A] (verified)	1848
3.368.4 Maple [A] (verified)	1849
3.368.5 Fricas [A] (verification not implemented)	1849
3.368.6 Sympy [A] (verification not implemented)	1850
3.368.7 Maxima [A] (verification not implemented)	1850
3.368.8 Giac [A] (verification not implemented)	1850
3.368.9 Mupad [B] (verification not implemented)	1851

3.368.1 Optimal result

Integrand size = 9, antiderivative size = 15

$$\int e^{\sin(x)} \sin(2x) dx = -2e^{\sin(x)} + 2e^{\sin(x)} \sin(x)$$

output `-2*exp(sin(x))+2*exp(sin(x))*sin(x)`

3.368.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int e^{\sin(x)} \sin(2x) dx = e^{\sin(x)}(-2 + 2 \sin(x))$$

input `Integrate[E^Sin[x]*Sin[2*x],x]`

output `E^Sin[x]*(-2 + 2*Sin[x])`

3.368.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {4878, 27, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{\sin(x)} \sin(2x) dx \\
 & \quad \downarrow 4878 \\
 & \int 2e^{\sin(x)} \sin(x) d \sin(x) \\
 & \quad \downarrow 27 \\
 & 2 \int e^{\sin(x)} \sin(x) d \sin(x) \\
 & \quad \downarrow 2607 \\
 & 2 \left(e^{\sin(x)} \sin(x) - \int e^{\sin(x)} d \sin(x) \right) \\
 & \quad \downarrow 2624 \\
 & 2 \left(e^{\sin(x)} \sin(x) - e^{\sin(x)} \right)
 \end{aligned}$$

input `Int[E^Sin[x]*Sin[2*x],x]`

output `2*(-E^Sin[x] + E^Sin[x]*Sin[x])`

3.368.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

```
rule 2607 Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m
_.), x_Symbol] := Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])),
x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^
n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*
m] && !TrueQ[$UseGamma]
```

```
rule 2624 Int[((F_)^(v_))^(n_), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;
FreeQ[{F, n}, x] && LinearQ[v, x]
```

```
rule 4878 Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFa
ctors[Sin[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Sin[v]/d,
u/Cos[v], x], x], x, Sin[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeF
actors[Sin[v], x], u/Cos[v], x]
```

3.368.4 Maple [A] (verified)

Time = 1.70 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
derivativedivides	$-2e^{\sin(x)} + 2e^{\sin(x)} \sin(x)$	14
default	$-2e^{\sin(x)} + 2e^{\sin(x)} \sin(x)$	14
risch	$-2e^{\sin(x)} + 2e^{\sin(x)} \sin(x)$	14

```
input int(exp(sin(x))*sin(2*x),x,method=_RETURNVERBOSE)
```

```
output -2*exp(sin(x))+2*exp(sin(x))*sin(x)
```

3.368.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.60

$$\int e^{\sin(x)} \sin(2x) dx = 2(\sin(x) - 1)e^{\sin(x)}$$

```
input integrate(exp(sin(x))*sin(2*x),x, algorithm="fricas")
```

output `2*(sin(x) - 1)*e^sin(x)`

3.368.6 Sympy [A] (verification not implemented)

Time = 0.78 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int e^{\sin(x)} \sin(2x) dx = 2e^{\sin(x)} \sin(x) - 2e^{\sin(x)}$$

input `integrate(exp(sin(x))*sin(2*x),x)`

output `2*exp(sin(x))*sin(x) - 2*exp(sin(x))`

3.368.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.60

$$\int e^{\sin(x)} \sin(2x) dx = 2(\sin(x) - 1)e^{\sin(x)}$$

input `integrate(exp(sin(x))*sin(2*x),x, algorithm="maxima")`

output `2*(sin(x) - 1)*e^sin(x)`

3.368.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.60

$$\int e^{\sin(x)} \sin(2x) dx = 2(\sin(x) - 1)e^{\sin(x)}$$

input `integrate(exp(sin(x))*sin(2*x),x, algorithm="giac")`

output `2*(sin(x) - 1)*e^sin(x)`

3.368.9 Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.60

$$\int e^{\sin(x)} \sin(2x) dx = 2 e^{\sin(x)} (\sin(x) - 1)$$

input `int(sin(2*x)*exp(sin(x)),x)`

output `2*exp(sin(x))*(sin(x) - 1)`

3.369 $\int x^2 \sqrt{5 - x^2} dx$

3.369.1 Optimal result	1852
3.369.2 Mathematica [A] (verified)	1852
3.369.3 Rubi [A] (verified)	1853
3.369.4 Maple [A] (verified)	1854
3.369.5 Fricas [A] (verification not implemented)	1854
3.369.6 Sympy [C] (verification not implemented)	1855
3.369.7 Maxima [A] (verification not implemented)	1855
3.369.8 Giac [A] (verification not implemented)	1855
3.369.9 Mupad [B] (verification not implemented)	1856

3.369.1 Optimal result

Integrand size = 15, antiderivative size = 47

$$\int x^2 \sqrt{5 - x^2} dx = -\frac{5}{8}x\sqrt{5 - x^2} + \frac{1}{4}x^3\sqrt{5 - x^2} + \frac{25}{8} \arcsin\left(\frac{x}{\sqrt{5}}\right)$$

output `25/8*arcsin(1/5*x*5^(1/2))-5/8*x*(-x^2+5)^(1/2)+1/4*x^3*(-x^2+5)^(1/2)`

3.369.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.06

$$\int x^2 \sqrt{5 - x^2} dx = \frac{1}{8}x\sqrt{5 - x^2}(-5 + 2x^2) + \frac{25}{4} \arctan\left(\frac{-\sqrt{5} + x}{\sqrt{5 - x^2}}\right)$$

input `Integrate[x^2*Sqrt[5 - x^2],x]`

output `(x*Sqrt[5 - x^2]*(-5 + 2*x^2))/8 + (25*ArcTan[(-Sqrt[5] + x)/Sqrt[5 - x^2]])/4`

3.369.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.11, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {248, 262, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \sqrt{5-x^2} dx$$

$$\downarrow 248$$

$$\frac{5}{4} \int \frac{x^2}{\sqrt{5-x^2}} dx + \frac{1}{4} \sqrt{5-x^2} x^3$$

$$\downarrow 262$$

$$\frac{5}{4} \left(\frac{5}{2} \int \frac{1}{\sqrt{5-x^2}} dx - \frac{1}{2} x \sqrt{5-x^2} \right) + \frac{1}{4} \sqrt{5-x^2} x^3$$

$$\downarrow 223$$

$$\frac{5}{4} \left(\frac{5}{2} \arcsin \left(\frac{x}{\sqrt{5}} \right) - \frac{1}{2} x \sqrt{5-x^2} \right) + \frac{1}{4} \sqrt{5-x^2} x^3$$

input `Int[x^2*Sqrt[5 - x^2],x]`

output `(x^3*Sqrt[5 - x^2])/4 + (5*(-1/2*(x*Sqrt[5 - x^2]) + (5*ArcSin[x/Sqrt[5]])) /2)/4`

3.369.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 248 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(c*x)^(m + 1)*((a + b*x^2)^p/(c*(m + 2*p + 1))), x] + Simp[2*a*(p/(m + 2*p + 1)) Int[(c*x)^m*(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c, m}, x] && GtQ[p, 0] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

```
rule 262 Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)
^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/
(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b
, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c
, 2, m, p, x]
```

3.369.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.74

method	result	size
default	$-\frac{x(-x^2+5)^{\frac{3}{2}}}{4} + \frac{5x\sqrt{-x^2+5}}{8} + \frac{25 \arcsin\left(\frac{x\sqrt{5}}{5}\right)}{8}$	35
risch	$-\frac{x(2x^2-5)(x^2-5)}{8\sqrt{-x^2+5}} + \frac{25 \arcsin\left(\frac{x\sqrt{5}}{5}\right)}{8}$	35
pseudoelliptic	$-\frac{25 \arctan\left(\frac{\sqrt{-x^2+5}}{x}\right)}{8} + \frac{(2x^3-5x)\sqrt{-x^2+5}}{8}$	38
meijerg	$-\frac{25i \left(-\frac{i\sqrt{\pi} x\sqrt{5} \left(-\frac{6x^2}{5} + 3 \right) \sqrt{-\frac{x^2}{5} + 1}}{30} + \frac{i\sqrt{\pi} \arcsin\left(\frac{x\sqrt{5}}{5}\right)}{2} \right)}{4\sqrt{\pi}}$	47
trager	$\frac{x(2x^2-5)\sqrt{-x^2+5}}{8} + \frac{25 \operatorname{RootOf}(_Z^2+1) \ln(\operatorname{RootOf}(_Z^2+1)\sqrt{-x^2+5+x})}{8}$	48

```
input int(x^2*(-x^2+5)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/4*x*(-x^2+5)^(3/2)+5/8*x*(-x^2+5)^(1/2)+25/8*arcsin(1/5*x*5^(1/2))
```

3.369.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.79

$$\int x^2 \sqrt{5-x^2} dx = \frac{1}{8} (2x^3 - 5x) \sqrt{-x^2+5} - \frac{25}{8} \arctan\left(\frac{\sqrt{-x^2+5}}{x}\right)$$

```
input integrate(x^2*(-x^2+5)^(1/2),x, algorithm="fracas")
```

```
output 1/8*(2*x^3 - 5*x)*sqrt(-x^2 + 5) - 25/8*arctan(sqrt(-x^2 + 5)/x)
```

3.369.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.71 (sec) , antiderivative size = 121, normalized size of antiderivative = 2.57

$$\int x^2 \sqrt{5-x^2} dx = \begin{cases} \frac{ix^5}{4\sqrt{x^2-5}} - \frac{15ix^3}{8\sqrt{x^2-5}} + \frac{25ix}{8\sqrt{x^2-5}} - \frac{25i \operatorname{acosh}\left(\frac{\sqrt{5}x}{5}\right)}{8} & \text{for } |x^2| > 5 \\ -\frac{x^5}{4\sqrt{5-x^2}} + \frac{15x^3}{8\sqrt{5-x^2}} - \frac{25x}{8\sqrt{5-x^2}} + \frac{25 \operatorname{asin}\left(\frac{\sqrt{5}x}{5}\right)}{8} & \text{otherwise} \end{cases}$$

input `integrate(x**2*(-x**2+5)**(1/2),x)`

output `Piecewise((I*x**5/(4*sqrt(x**2 - 5)) - 15*I*x**3/(8*sqrt(x**2 - 5)) + 25*I*x/(8*sqrt(x**2 - 5)) - 25*I*acosh(sqrt(5)*x/5)/8, Abs(x**2) > 5), (-x**5/(4*sqrt(5 - x**2)) + 15*x**3/(8*sqrt(5 - x**2)) - 25*x/(8*sqrt(5 - x**2)) + 25*asin(sqrt(5)*x/5)/8, True))`

3.369.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.72

$$\int x^2 \sqrt{5-x^2} dx = -\frac{1}{4} (-x^2 + 5)^{\frac{3}{2}} x + \frac{5}{8} \sqrt{-x^2 + 5} x + \frac{25}{8} \arcsin\left(\frac{1}{5} \sqrt{5} x\right)$$

input `integrate(x^2*(-x^2+5)^(1/2),x, algorithm="maxima")`

output `-1/4*(-x^2 + 5)^(3/2)*x + 5/8*sqrt(-x^2 + 5)*x + 25/8*arcsin(1/5*sqrt(5)*x)`

3.369.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.62

$$\int x^2 \sqrt{5-x^2} dx = \frac{1}{8} (2x^2 - 5) \sqrt{-x^2 + 5} x + \frac{25}{8} \arcsin\left(\frac{1}{5} \sqrt{5} x\right)$$

input `integrate(x^2*(-x^2+5)^(1/2),x, algorithm="giac")`

output `1/8*(2*x^2 - 5)*sqrt(-x^2 + 5)*x + 25/8*arcsin(1/5*sqrt(5)*x)`

3.369.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.64

$$\int x^2 \sqrt{5-x^2} dx = \frac{25 \operatorname{asin}\left(\frac{\sqrt{5}x}{5}\right)}{8} - \sqrt{5-x^2} \left(\frac{5x}{8} - \frac{x^3}{4}\right)$$

input `int(x^2*(5 - x^2)^(1/2),x)`output `(25*asin((5^(1/2)*x)/5))/8 - (5 - x^2)^(1/2)*((5*x)/8 - x^3/4)`

3.370 $\int x^2(1+x^3)^4 dx$

3.370.1 Optimal result	1857
3.370.2 Mathematica [B] (verified)	1857
3.370.3 Rubi [A] (verified)	1858
3.370.4 Maple [A] (verified)	1858
3.370.5 Fricas [B] (verification not implemented)	1859
3.370.6 Sympy [B] (verification not implemented)	1859
3.370.7 Maxima [A] (verification not implemented)	1859
3.370.8 Giac [A] (verification not implemented)	1860
3.370.9 Mupad [B] (verification not implemented)	1860

3.370.1 Optimal result

Integrand size = 11, antiderivative size = 11

$$\int x^2(1+x^3)^4 dx = \frac{1}{15}(1+x^3)^5$$

output `1/15*(x^3+1)^5`

3.370.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 36 vs. $2(11) = 22$.

Time = 0.00 (sec) , antiderivative size = 36, normalized size of antiderivative = 3.27

$$\int x^2(1+x^3)^4 dx = \frac{x^3}{3} + \frac{2x^6}{3} + \frac{2x^9}{3} + \frac{x^{12}}{3} + \frac{x^{15}}{15}$$

input `Integrate[x^2*(1 + x^3)^4,x]`

output `x^3/3 + (2*x^6)/3 + (2*x^9)/3 + x^12/3 + x^15/15`

3.370.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {793}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2(x^3 + 1)^4 dx$$

$$\downarrow 793$$

$$\frac{1}{15}(x^3 + 1)^5$$

input `Int[x^2*(1 + x^3)^4,x]`

output `(1 + x^3)^5/15`

3.370.3.1 Defintions of rubi rules used

rule 793 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(a + b*x^n)^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]`

3.370.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{(x^3+1)^5}{15}$	10
gospers	$\frac{x^3(x^{12}+5x^9+10x^6+10x^3+5)}{15}$	26
norman	$\frac{1}{3}x^3 + \frac{2}{3}x^6 + \frac{2}{3}x^9 + \frac{1}{3}x^{12} + \frac{1}{15}x^{15}$	27
paralelrisch	$\frac{1}{3}x^3 + \frac{2}{3}x^6 + \frac{2}{3}x^9 + \frac{1}{3}x^{12} + \frac{1}{15}x^{15}$	27
risch	$\frac{1}{15}x^{15} + \frac{1}{3}x^{12} + \frac{2}{3}x^9 + \frac{2}{3}x^6 + \frac{1}{3}x^3 + \frac{1}{15}$	28

input `int(x^2*(x^3+1)^4,x,method=_RETURNVERBOSE)`

output `1/15*(x^3+1)^5`

3.370.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. 2(9) = 18.

Time = 0.24 (sec) , antiderivative size = 26, normalized size of antiderivative = 2.36

$$\int x^2(1+x^3)^4 dx = \frac{1}{15}x^{15} + \frac{1}{3}x^{12} + \frac{2}{3}x^9 + \frac{2}{3}x^6 + \frac{1}{3}x^3$$

input `integrate(x^2*(x^3+1)^4,x, algorithm="fricas")`

output `1/15*x^15 + 1/3*x^12 + 2/3*x^9 + 2/3*x^6 + 1/3*x^3`

3.370.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(7) = 14.

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 2.45

$$\int x^2(1+x^3)^4 dx = \frac{x^{15}}{15} + \frac{x^{12}}{3} + \frac{2x^9}{3} + \frac{2x^6}{3} + \frac{x^3}{3}$$

input `integrate(x**2*(x**3+1)**4,x)`

output `x**15/15 + x**12/3 + 2*x**9/3 + 2*x**6/3 + x**3/3`

3.370.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int x^2(1+x^3)^4 dx = \frac{1}{15}(x^3+1)^5$$

input `integrate(x^2*(x^3+1)^4,x, algorithm="maxima")`

output `1/15*(x^3 + 1)^5`

3.370.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int x^2(1+x^3)^4 dx = \frac{1}{15}(x^3+1)^5$$

input `integrate(x^2*(x^3+1)^4,x, algorithm="giac")`output `1/15*(x^3 + 1)^5`**3.370.9 Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 2.36

$$\int x^2(1+x^3)^4 dx = \frac{x^{15}}{15} + \frac{x^{12}}{3} + \frac{2x^9}{3} + \frac{2x^6}{3} + \frac{x^3}{3}$$

input `int(x^2*(x^3 + 1)^4,x)`output `x^3/3 + (2*x^6)/3 + (2*x^9)/3 + x^12/3 + x^15/15`

3.371 $\int \cos^3(x) \sin^3(x) dx$

3.371.1 Optimal result1861
3.371.2 Mathematica [A] (verified)1861
3.371.3 Rubi [A] (verified)1862
3.371.4 Maple [A] (verified)1863
3.371.5 Fricas [A] (verification not implemented)1863
3.371.6 Sympy [A] (verification not implemented)1864
3.371.7 Maxima [A] (verification not implemented)1864
3.371.8 Giac [A] (verification not implemented)1864
3.371.9 Mupad [B] (verification not implemented)1865

3.371.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \cos^3(x) \sin^3(x) dx = \frac{\sin^4(x)}{4} - \frac{\sin^6(x)}{6}$$

output `1/4*sin(x)^4-1/6*sin(x)^6`

3.371.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cos^3(x) \sin^3(x) dx = -\frac{3}{64} \cos(2x) + \frac{1}{192} \cos(6x)$$

input `Integrate[Cos[x]^3*Sin[x]^3,x]`

output `(-3*Cos[2*x])/64 + Cos[6*x]/192`

3.371.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 3044, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^3(x) \cos^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^3 \cos(x)^3 dx \\
 & \quad \downarrow \text{3044} \\
 & \int \sin^3(x) (1 - \sin^2(x)) d \sin(x) \\
 & \quad \downarrow \text{244} \\
 & \int (\sin^3(x) - \sin^5(x)) d \sin(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sin^4(x)}{4} - \frac{\sin^6(x)}{6}
 \end{aligned}$$

input `Int[Cos[x]^3*Sin[x]^3,x]`

output `Sin[x]^4/4 - Sin[x]^6/6`

3.371.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] :> Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3044 Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_
Symbol] := Simp[1/(a*f) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a
*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(I
ntegerQ[(m - 1)/2] && LtQ[0, m, n])
```

3.371.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$\frac{\sin^4(x)}{4} - \frac{\sin^6(x)}{6}$	14
default	$\frac{\sin^4(x)}{4} - \frac{\sin^6(x)}{6}$	14
risch	$\frac{\cos(6x)}{192} - \frac{3 \cos(2x)}{64}$	14
parallelrisch	$\frac{7}{40} + \frac{\cos(6x)}{192} - \frac{3 \cos(2x)}{64}$	15
norman	$\frac{4(\tan^4(\frac{x}{2})) + 4(\tan^8(\frac{x}{2})) - \frac{8(\tan^6(\frac{x}{2}))}{3}}{(1 + \tan^2(\frac{x}{2}))^6}$	37

```
input int(sin(x)^3*cos(x)^3,x,method=_RETURNVERBOSE)
```

```
output 1/4*sin(x)^4-1/6*sin(x)^6
```

3.371.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos^3(x) \sin^3(x) dx = \frac{1}{6} \cos(x)^6 - \frac{1}{4} \cos(x)^4$$

```
input integrate(cos(x)^3*sin(x)^3,x, algorithm="fracas")
```

```
output 1/6*cos(x)^6 - 1/4*cos(x)^4
```


3.371.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int \cos^3(x) \sin^3(x) dx = -\frac{\sin^6(x)}{6} + \frac{\sin^4(x)}{4}$$

input `integrate(cos(x)**3*sin(x)**3,x)`output `-sin(x)**6/6 + sin(x)**4/4`**3.371.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos^3(x) \sin^3(x) dx = -\frac{1}{6} \sin(x)^6 + \frac{1}{4} \sin(x)^4$$

input `integrate(cos(x)^3*sin(x)^3,x, algorithm="maxima")`output `-1/6*sin(x)^6 + 1/4*sin(x)^4`**3.371.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos^3(x) \sin^3(x) dx = \frac{1}{6} \cos(x)^6 - \frac{1}{4} \cos(x)^4$$

input `integrate(cos(x)^3*sin(x)^3,x, algorithm="giac")`output `1/6*cos(x)^6 - 1/4*cos(x)^4`

3.371.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \cos^3(x) \sin^3(x) dx = -\frac{\sin(x)^4 (2 \sin(x)^2 - 3)}{12}$$

input `int(cos(x)^3*sin(x)^3,x)`

output `-(sin(x)^4*(2*sin(x)^2 - 3))/12`

3.372 $\int \sec^4(x) \tan^2(x) dx$

3.372.1 Optimal result	1866
3.372.2 Mathematica [A] (verified)	1866
3.372.3 Rubi [A] (verified)	1867
3.372.4 Maple [A] (verified)	1868
3.372.5 Fricas [A] (verification not implemented)	1868
3.372.6 Sympy [B] (verification not implemented)	1869
3.372.7 Maxima [A] (verification not implemented)	1869
3.372.8 Giac [A] (verification not implemented)	1869
3.372.9 Mupad [B] (verification not implemented)	1870

3.372.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \sec^4(x) \tan^2(x) dx = \frac{\tan^3(x)}{3} + \frac{\tan^5(x)}{5}$$

output `1/3*tan(x)^3+1/5*tan(x)^5`

3.372.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \sec^4(x) \tan^2(x) dx = -\frac{2 \tan(x)}{15} - \frac{1}{15} \sec^2(x) \tan(x) + \frac{1}{5} \sec^4(x) \tan(x)$$

input `Integrate[Sec[x]^4*Tan[x]^2,x]`

output `(-2*Tan[x])/15 - (Sec[x]^2*Tan[x])/15 + (Sec[x]^4*Tan[x])/5`

3.372.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 3087, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^2(x) \sec^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^2 \sec(x)^4 dx \\
 & \quad \downarrow \text{3087} \\
 & \int \tan^2(x) (\tan^2(x) + 1) d \tan(x) \\
 & \quad \downarrow \text{244} \\
 & \int (\tan^4(x) + \tan^2(x)) d \tan(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\tan^5(x)}{5} + \frac{\tan^3(x)}{3}
 \end{aligned}$$

input `Int[Sec[x]^4*Tan[x]^2,x]`

output `Tan[x]^3/3 + Tan[x]^5/5`

3.372.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] :> Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3087 `Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[1/f Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])`

3.372.4 Maple [A] (verified)

Time = 1.82 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$\frac{\tan^3(x)}{3} + \frac{\tan^5(x)}{5}$	14
default	$\frac{\tan^3(x)}{3} + \frac{\tan^5(x)}{5}$	14
risch	$-\frac{4i(15e^{6ix} - 5e^{4ix} + 5e^{2ix} + 1)}{15(e^{2ix} + 1)^5}$	36

input `int(sec(x)^4*tan(x)^2,x,method=_RETURNVERBOSE)`

output `1/3*tan(x)^3+1/5*tan(x)^5`

3.372.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \sec^4(x) \tan^2(x) dx = -\frac{(2 \cos(x)^4 + \cos(x)^2 - 3) \sin(x)}{15 \cos(x)^5}$$

input `integrate(sec(x)^4*tan(x)^2,x, algorithm="fracas")`

output `-1/15*(2*cos(x)^4 + cos(x)^2 - 3)*sin(x)/cos(x)^5`

3.372.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(12) = 24$.

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.71

$$\int \sec^4(x) \tan^2(x) dx = -\frac{2 \sin(x)}{15 \cos(x)} - \frac{\sin(x)}{15 \cos^3(x)} + \frac{\sin(x)}{5 \cos^5(x)}$$

input `integrate(sec(x)**4*tan(x)**2,x)`

output `-2*sin(x)/(15*cos(x)) - sin(x)/(15*cos(x)**3) + sin(x)/(5*cos(x)**5)`

3.372.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sec^4(x) \tan^2(x) dx = \frac{1}{5} \tan(x)^5 + \frac{1}{3} \tan(x)^3$$

input `integrate(sec(x)^4*tan(x)^2,x, algorithm="maxima")`

output `1/5*tan(x)^5 + 1/3*tan(x)^3`

3.372.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sec^4(x) \tan^2(x) dx = \frac{1}{5} \tan(x)^5 + \frac{1}{3} \tan(x)^3$$

input `integrate(sec(x)^4*tan(x)^2,x, algorithm="giac")`

output `1/5*tan(x)^5 + 1/3*tan(x)^3`

3.372.9 Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sec^4(x) \tan^2(x) dx = \frac{\tan(x)^5}{5} + \frac{\tan(x)^3}{3}$$

input `int(tan(x)^2/cos(x)^4,x)`

output `tan(x)^3/3 + tan(x)^5/5`

3.373 $\int x\sqrt{1+2x} dx$

3.373.1 Optimal result1871
3.373.2 Mathematica [A] (verified)1871
3.373.3 Rubi [A] (verified)1872
3.373.4 Maple [A] (verified)1873
3.373.5 Fricas [A] (verification not implemented)1873
3.373.6 Sympy [A] (verification not implemented)1874
3.373.7 Maxima [A] (verification not implemented)1874
3.373.8 Giac [A] (verification not implemented)1874
3.373.9 Mupad [B] (verification not implemented)1875

3.373.1 Optimal result

Integrand size = 11, antiderivative size = 27

$$\int x\sqrt{1+2x} dx = -\frac{1}{6}(1+2x)^{3/2} + \frac{1}{10}(1+2x)^{5/2}$$

output `-1/6*(1+2*x)^(3/2)+1/10*(1+2*x)^(5/2)`

3.373.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.67

$$\int x\sqrt{1+2x} dx = \frac{1}{15}(1+2x)^{3/2}(-1+3x)$$

input `Integrate[x*Sqrt[1 + 2*x],x]`

output `((1 + 2*x)^(3/2)*(-1 + 3*x))/15`

3.373.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x\sqrt{2x+1} dx$$

$$\downarrow \text{53}$$

$$\int \left(\frac{1}{2}(2x+1)^{3/2} - \frac{1}{2}\sqrt{2x+1} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{1}{10}(2x+1)^{5/2} - \frac{1}{6}(2x+1)^{3/2}$$

input `Int[x*Sqrt[1 + 2*x],x]`

output `-1/6*(1 + 2*x)^(3/2) + (1 + 2*x)^(5/2)/10`

3.373.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.373.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.56

method	result	size
gospers	$\frac{(1+2x)^{\frac{3}{2}}(-1+3x)}{15}$	15
risch	$\frac{(6x^2+x-1)\sqrt{1+2x}}{15}$	18
pseudoelliptic	$\frac{(6x^2+x-1)\sqrt{1+2x}}{15}$	18
trager	$\left(\frac{2}{5}x^2 + \frac{1}{15}x - \frac{1}{15}\right)\sqrt{1+2x}$	19
derivativedivides	$-\frac{(1+2x)^{\frac{3}{2}}}{6} + \frac{(1+2x)^{\frac{5}{2}}}{10}$	20
default	$-\frac{(1+2x)^{\frac{3}{2}}}{6} + \frac{(1+2x)^{\frac{5}{2}}}{10}$	20
meijerg	$-\frac{8\sqrt{\pi}}{15} + \frac{4\sqrt{\pi}(1+2x)^{\frac{3}{2}}(-6x+2)}{8\sqrt{\pi}15}$	29

input `int(x*(1+2*x)^(1/2),x,method=_RETURNVERBOSE)`output `1/15*(1+2*x)^(3/2)*(-1+3*x)`**3.373.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.63

$$\int x\sqrt{1+2x} dx = \frac{1}{15} (6x^2 + x - 1)\sqrt{2x+1}$$

input `integrate(x*(1+2*x)^(1/2),x, algorithm="fricas")`output `1/15*(6*x^2 + x - 1)*sqrt(2*x + 1)`

3.373.6 Sympy [A] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.33

$$\int x\sqrt{1+2x} dx = \frac{2x^2\sqrt{2x+1}}{5} + \frac{x\sqrt{2x+1}}{15} - \frac{\sqrt{2x+1}}{15}$$

input `integrate(x*(1+2*x)**(1/2),x)`output `2*x**2*sqrt(2*x + 1)/5 + x*sqrt(2*x + 1)/15 - sqrt(2*x + 1)/15`**3.373.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int x\sqrt{1+2x} dx = \frac{1}{10} (2x+1)^{\frac{5}{2}} - \frac{1}{6} (2x+1)^{\frac{3}{2}}$$

input `integrate(x*(1+2*x)^(1/2),x, algorithm="maxima")`output `1/10*(2*x + 1)^(5/2) - 1/6*(2*x + 1)^(3/2)`**3.373.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int x\sqrt{1+2x} dx = \frac{1}{10} (2x+1)^{\frac{5}{2}} - \frac{1}{6} (2x+1)^{\frac{3}{2}}$$

input `integrate(x*(1+2*x)^(1/2),x, algorithm="giac")`output `1/10*(2*x + 1)^(5/2) - 1/6*(2*x + 1)^(3/2)`

3.373.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.52

$$\int x\sqrt{1+2x} dx = \frac{(2x+1)^{3/2}(6x-2)}{30}$$

input `int(x*(2*x + 1)^(1/2),x)`

output `((2*x + 1)^(3/2)*(6*x - 2))/30`

3.374 $\int \sin^4(x) dx$

3.374.1 Optimal result	1876
3.374.2 Mathematica [A] (verified)	1876
3.374.3 Rubi [A] (verified)	1877
3.374.4 Maple [A] (verified)	1878
3.374.5 Fricas [A] (verification not implemented)	1879
3.374.6 Sympy [A] (verification not implemented)	1879
3.374.7 Maxima [A] (verification not implemented)	1879
3.374.8 Giac [A] (verification not implemented)	1880
3.374.9 Mupad [B] (verification not implemented)	1880

3.374.1 Optimal result

Integrand size = 4, antiderivative size = 24

$$\int \sin^4(x) dx = \frac{3x}{8} - \frac{3}{8} \cos(x) \sin(x) - \frac{1}{4} \cos(x) \sin^3(x)$$

output `3/8*x-3/8*cos(x)*sin(x)-1/4*cos(x)*sin(x)^3`

3.374.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \sin^4(x) dx = \frac{3x}{8} - \frac{1}{4} \sin(2x) + \frac{1}{32} \sin(4x)$$

input `Integrate[Sin[x]^4,x]`

output `(3*x)/8 - Sin[2*x]/4 + Sin[4*x]/32`

3.374.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^4 dx \\
 & \quad \downarrow \text{3115} \\
 & \frac{3}{4} \int \sin^2(x) dx - \frac{1}{4} \sin^3(x) \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{4} \int \sin(x)^2 dx - \frac{1}{4} \sin^3(x) \cos(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{3}{4} \left(\int \frac{1 dx}{2} - \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin^3(x) \cos(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{3}{4} \left(\frac{x}{2} - \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin^3(x) \cos(x)
 \end{aligned}$$

input `Int[Sin[x]^4,x]`

output `-1/4*(Cos[x]*Sin[x]^3) + (3*(x/2 - (Cos[x]*Sin[x])/2))/4`

3.374.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

3.374.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.71

method	result	size
risch	$\frac{3x}{8} + \frac{\sin(4x)}{32} - \frac{\sin(2x)}{4}$	17
parallelrisch	$\frac{3x}{8} + \frac{\sin(4x)}{32} - \frac{\sin(2x)}{4}$	17
default	$-\frac{(\sin^3(x) + \frac{3\sin(x)}{2})\cos(x)}{4} + \frac{3x}{8}$	18
norman	$\frac{3x}{8} - \frac{11(\tan^3(\frac{x}{2}))}{4} + \frac{11(\tan^5(\frac{x}{2}))}{4} + \frac{3(\tan^7(\frac{x}{2}))}{4} + \frac{3x(\tan^2(\frac{x}{2}))}{2} + \frac{9x(\tan^4(\frac{x}{2}))}{4} + \frac{3x(\tan^6(\frac{x}{2}))}{2} + \frac{3x(\tan^8(\frac{x}{2}))}{8} - \frac{3\tan(\frac{x}{2})}{4}$ $(1+\tan^2(\frac{x}{2}))^4$	82

input `int(sin(x)^4,x,method=_RETURNVERBOSE)`

output `3/8*x+1/32*sin(4*x)-1/4*sin(2*x)`

3.374.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \sin^4(x) dx = \frac{1}{8} (2 \cos(x)^3 - 5 \cos(x)) \sin(x) + \frac{3}{8} x$$

input `integrate(sin(x)^4,x, algorithm="fricas")`output `1/8*(2*cos(x)^3 - 5*cos(x))*sin(x) + 3/8*x`**3.374.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \sin^4(x) dx = \frac{3x}{8} - \frac{\sin^3(x) \cos(x)}{4} - \frac{3 \sin(x) \cos(x)}{8}$$

input `integrate(sin(x)**4,x)`output `3*x/8 - sin(x)**3*cos(x)/4 - 3*sin(x)*cos(x)/8`**3.374.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \sin^4(x) dx = \frac{3}{8} x + \frac{1}{32} \sin(4x) - \frac{1}{4} \sin(2x)$$

input `integrate(sin(x)^4,x, algorithm="maxima")`output `3/8*x + 1/32*sin(4*x) - 1/4*sin(2*x)`

3.374.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \sin^4(x) dx = \frac{3}{8}x + \frac{1}{32}\sin(4x) - \frac{1}{4}\sin(2x)$$

input `integrate(sin(x)^4,x, algorithm="giac")`

output `3/8*x + 1/32*sin(4*x) - 1/4*sin(2*x)`

3.374.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \sin^4(x) dx = \frac{3x}{8} - \frac{\sin(2x)}{4} + \frac{\sin(4x)}{32}$$

input `int(sin(x)^4,x)`

output `(3*x)/8 - sin(2*x)/4 + sin(4*x)/32`

3.375 $\int \tan^3(x) dx$

3.375.1 Optimal result	1881
3.375.2 Mathematica [A] (verified)	1881
3.375.3 Rubi [A] (verified)	1882
3.375.4 Maple [A] (verified)	1883
3.375.5 Fricas [A] (verification not implemented)	1883
3.375.6 Sympy [A] (verification not implemented)	1884
3.375.7 Maxima [A] (verification not implemented)	1884
3.375.8 Giac [A] (verification not implemented)	1884
3.375.9 Mupad [B] (verification not implemented)	1885

3.375.1 Optimal result

Integrand size = 4, antiderivative size = 12

$$\int \tan^3(x) dx = \log(\cos(x)) + \frac{\tan^2(x)}{2}$$

output `ln(cos(x))+1/2*tan(x)^2`

3.375.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \tan^3(x) dx = \log(\cos(x)) + \frac{\tan^2(x)}{2}$$

input `Integrate[Tan[x]^3,x]`

output `Log[Cos[x]] + Tan[x]^2/2`

3.375.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 3954, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \tan^3(x) dx \\
 \downarrow \text{3042} \\
 \int \tan(x)^3 dx \\
 \downarrow \text{3954} \\
 \frac{\tan^2(x)}{2} - \int \tan(x) dx \\
 \downarrow \text{3042} \\
 \frac{\tan^2(x)}{2} - \int \tan(x) dx \\
 \downarrow \text{3956} \\
 \frac{\tan^2(x)}{2} + \log(\cos(x))
 \end{array}$$

input `Int[Tan[x]^3,x]`

output `Log[Cos[x]] + Tan[x]^2/2`

3.375.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

3.375.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.42

method	result	size
derivativdivides	$\frac{\tan^2(x)}{2} - \frac{\ln(1+\tan^2(x))}{2}$	17
default	$\frac{\tan^2(x)}{2} - \frac{\ln(1+\tan^2(x))}{2}$	17
norman	$\frac{\tan^2(x)}{2} - \frac{\ln(1+\tan^2(x))}{2}$	17
parallelrisc	$\frac{\tan^2(x)}{2} - \frac{\ln(1+\tan^2(x))}{2}$	17
risc	$-ix + \frac{2e^{2ix}}{(e^{2ix}+1)^2} + \ln(e^{2ix} + 1)$	30

input `int(tan(x)^3,x,method=_RETURNVERBOSE)`

output `1/2*tan(x)^2-1/2*ln(1+tan(x)^2)`

3.375.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.50

$$\int \tan^3(x) dx = \frac{1}{2} \tan(x)^2 + \frac{1}{2} \log\left(\frac{1}{\tan(x)^2 + 1}\right)$$

input `integrate(tan(x)^3,x, algorithm="fricas")`

output `1/2*tan(x)^2 + 1/2*log(1/(tan(x)^2 + 1))`

3.375.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \tan^3(x) dx = \log(\cos(x)) + \frac{1}{2\cos^2(x)}$$

input `integrate(tan(x)**3,x)`output `log(cos(x)) + 1/(2*cos(x)**2)`**3.375.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.67

$$\int \tan^3(x) dx = -\frac{1}{2(\sin(x)^2 - 1)} + \frac{1}{2} \log(\sin(x)^2 - 1)$$

input `integrate(tan(x)^3,x, algorithm="maxima")`output `-1/2/(sin(x)^2 - 1) + 1/2*log(sin(x)^2 - 1)`**3.375.8 Giac [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.33

$$\int \tan^3(x) dx = \frac{1}{2} \tan(x)^2 - \frac{1}{2} \log(\tan(x)^2 + 1)$$

input `integrate(tan(x)^3,x, algorithm="giac")`output `1/2*tan(x)^2 - 1/2*log(tan(x)^2 + 1)`

3.375.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.33

$$\int \tan^3(x) dx = \ln(\cos(x)) - \frac{\cos(x)^2 - 1}{2\cos(x)^2}$$

input `int(tan(x)^3,x)`

output `log(cos(x)) - (cos(x)^2 - 1)/(2*cos(x)^2)`

3.376 $\int x^5 \sqrt{1+x^2} dx$

3.376.1 Optimal result	1886
3.376.2 Mathematica [A] (verified)	1886
3.376.3 Rubi [A] (verified)	1887
3.376.4 Maple [A] (verified)	1888
3.376.5 Fricas [A] (verification not implemented)	1888
3.376.6 Sympy [A] (verification not implemented)	1889
3.376.7 Maxima [A] (verification not implemented)	1889
3.376.8 Giac [A] (verification not implemented)	1889
3.376.9 Mupad [B] (verification not implemented)	1890

3.376.1 Optimal result

Integrand size = 13, antiderivative size = 40

$$\int x^5 \sqrt{1+x^2} dx = \frac{1}{3}(1+x^2)^{3/2} - \frac{2}{5}(1+x^2)^{5/2} + \frac{1}{7}(1+x^2)^{7/2}$$

output `1/3*(x^2+1)^(3/2)-2/5*(x^2+1)^(5/2)+1/7*(x^2+1)^(7/2)`

3.376.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.75

$$\int x^5 \sqrt{1+x^2} dx = \frac{1}{105} \sqrt{1+x^2} (8 - 4x^2 + 3x^4 + 15x^6)$$

input `Integrate[x^5*Sqrt[1 + x^2],x]`

output `(Sqrt[1 + x^2]*(8 - 4*x^2 + 3*x^4 + 15*x^6))/105`

3.376.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.10, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^5 \sqrt{x^2 + 1} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int x^4 \sqrt{x^2 + 1} dx^2 \\ & \quad \downarrow \text{53} \\ & \frac{1}{2} \int \left((x^2 + 1)^{5/2} - 2(x^2 + 1)^{3/2} + \sqrt{x^2 + 1} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \left(\frac{2}{7} (x^2 + 1)^{7/2} - \frac{4}{5} (x^2 + 1)^{5/2} + \frac{2}{3} (x^2 + 1)^{3/2} \right) \end{aligned}$$

input `Int[x^5*Sqrt[1 + x^2],x]`

output `((2*(1 + x^2)^(3/2))/3 - (4*(1 + x^2)^(5/2))/5 + (2*(1 + x^2)^(7/2))/7)/2`

3.376.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.376.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.55

method	result	size
gospers	$\frac{(x^2+1)^{\frac{3}{2}}(15x^4-12x^2+8)}{105}$	22
pseudoelliptic	$\frac{(x^2+1)^{\frac{3}{2}}(15x^4-12x^2+8)}{105}$	22
trager	$\left(\frac{1}{7}x^6 + \frac{1}{35}x^4 - \frac{4}{105}x^2 + \frac{8}{105}\right)\sqrt{x^2+1}$	26
risch	$\frac{(15x^6+3x^4-4x^2+8)\sqrt{x^2+1}}{105}$	27
default	$\frac{x^4(x^2+1)^{\frac{3}{2}}}{7} - \frac{4x^2(x^2+1)^{\frac{3}{2}}}{35} + \frac{8(x^2+1)^{\frac{3}{2}}}{105}$	35
meijerg	$-\frac{32\sqrt{\pi}}{105} - \frac{4\sqrt{\pi}(x^2+1)^{\frac{3}{2}}(15x^4-12x^2+8)}{105 \cdot 4\sqrt{\pi}}$	36

input `int(x^5*(x^2+1)^(1/2),x,method=_RETURNVERBOSE)`output `1/105*(x^2+1)^(3/2)*(15*x^4-12*x^2+8)`**3.376.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.65

$$\int x^5 \sqrt{1+x^2} dx = \frac{1}{105} (15x^6 + 3x^4 - 4x^2 + 8) \sqrt{x^2+1}$$

input `integrate(x^5*(x^2+1)^(1/2),x, algorithm="fracas")`output `1/105*(15*x^6 + 3*x^4 - 4*x^2 + 8)*sqrt(x^2 + 1)`

3.376.6 Sympy [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.32

$$\int x^5 \sqrt{1+x^2} dx = \frac{x^6 \sqrt{x^2+1}}{7} + \frac{x^4 \sqrt{x^2+1}}{35} - \frac{4x^2 \sqrt{x^2+1}}{105} + \frac{8\sqrt{x^2+1}}{105}$$

input `integrate(x**5*(x**2+1)**(1/2),x)`output `x**6*sqrt(x**2 + 1)/7 + x**4*sqrt(x**2 + 1)/35 - 4*x**2*sqrt(x**2 + 1)/105 + 8*sqrt(x**2 + 1)/105`**3.376.7 Maxima [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.85

$$\int x^5 \sqrt{1+x^2} dx = \frac{1}{7} (x^2+1)^{\frac{3}{2}} x^4 - \frac{4}{35} (x^2+1)^{\frac{3}{2}} x^2 + \frac{8}{105} (x^2+1)^{\frac{3}{2}}$$

input `integrate(x^5*(x^2+1)^(1/2),x, algorithm="maxima")`output `1/7*(x^2 + 1)^(3/2)*x^4 - 4/35*(x^2 + 1)^(3/2)*x^2 + 8/105*(x^2 + 1)^(3/2)`**3.376.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.70

$$\int x^5 \sqrt{1+x^2} dx = \frac{1}{7} (x^2+1)^{\frac{7}{2}} - \frac{2}{5} (x^2+1)^{\frac{5}{2}} + \frac{1}{3} (x^2+1)^{\frac{3}{2}}$$

input `integrate(x^5*(x^2+1)^(1/2),x, algorithm="giac")`output `1/7*(x^2 + 1)^(7/2) - 2/5*(x^2 + 1)^(5/2) + 1/3*(x^2 + 1)^(3/2)`

3.376.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.62

$$\int x^5 \sqrt{1+x^2} dx = \sqrt{x^2+1} \left(\frac{x^6}{7} + \frac{x^4}{35} - \frac{4x^2}{105} + \frac{8}{105} \right)$$

input `int(x^5*(x^2 + 1)^(1/2),x)`

output `(x^2 + 1)^(1/2)*(x^4/35 - (4*x^2)/105 + x^6/7 + 8/105)`

APPENDIX

4.1 Listing of Grading functions	1891
--	------

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.1.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
```

```

(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A"," "}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A"," "}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal.$
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
      ,
      finalresult={"F","Contains unresolved integral."}
    ]
  ];

  finalresult
]

```

```

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType,expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]],2]],
            Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3,ExpnType[expn[[1]]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
            If[Head[expn]===RootSum,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

4.1.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
      return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
            print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
      return "F","Result contains unresolved integral";
fi;

```



```

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
  if is_contains_complex(result) then
    if is_contains_complex(optimal) then
      if debug then
        print("both result and optimal complex");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        return "A"," ";
      else
        return "B",cat("Both result and optimal contain complex but leaf count of
                        convert(leaf_count_result,string)," vs. $2 (" ,
                        convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_
        end if
      else #result contains complex but optimal is not
        if debug then
          print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
      fi;
    else # result do not contain complex
      # this assumes optimal do not as well. No check is needed here.
      if debug then
        print("result do not contain complex, this assumes optimal do not as well");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        if debug then
          print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
      else
        if debug then
          print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string),"$ vs. $2(",
                        convert(leaf_count_optimal,string),"=" ,convert(2*leaf_cou
        fi;
      fi;
    fi;
  fi;

```

```

else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  end if
end if

```

```

elif type(expn, ``~`) then
  if type(op(2,expn), 'integer') then
    ExpnType(op(1,expn))
  elif type(op(2,expn), 'rational') then
    if type(op(1,expn), 'rational') then
      1
    else
      max(2, ExpnType(op(1,expn)))
    end if
  else
    max(3, ExpnType(op(1,expn)), ExpnType(op(2,expn)))
  end if
elif type(expn, ``+`) or type(expn, ``*`) then
  max(ExpnType(op(1,expn)), max(ExpnType(rest(expn))))
elif ElementaryFunctionQ(op(0,expn)) then
  max(3, ExpnType(op(1,expn)))
elif SpecialFunctionQ(op(0,expn)) then
  max(4, apply(max, map(ExpnType, [op(expn)])))
elif HypergeometricFunctionQ(op(0,expn)) then
  max(5, apply(max, map(ExpnType, [op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6, apply(max, map(ExpnType, [op(expn)])))
elif op(0,expn)='int' then
  max(8, apply(max, map(ExpnType, [op(expn)]))) else
  9
end if
end proc:

```

```

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

```

```

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,

```

```

        GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
        EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
    member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
    member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
    if nops(u)=2 then
        op(2,u)
    else
        apply(op(0,u),op(2..nops(u),u))
    end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma] [LeafCount] (u);
end proc:

```

4.1.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):

```

```
if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
        return True
    else:
        return False
else:
    return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
```

```

return 1
elif isinstance(expn,list):
    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
        return 1
    else:
        return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
elif isinstance(expn,Pow): #type(expn,``^`)
    if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
        return expnType(expn.args[0]) #ExpnType(op(1,expn))
    elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or type(expn,``*`)
    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

#main function

```

def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

    #print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```

4.1.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
```



```

        return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

```

```

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric', 'hypergeometric_M', 'hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=", expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception, AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list:    #isinstance(expn,list):

```

```

    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
elif expn.operator() == operator.pow: #isinstance(expn,Pow)
    if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
        return expnType(expn.operands()[0]) #expnType(expn.args[0])
    elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
        if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or inst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

```

```

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)",type(result))
    print("type(optimal)",type(optimal))

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal." + str(leaf_
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```