

Computer Algebra Independent Integration Tests

Summer 2023 edition with Rubi V 4.17.3

11-MIT

Nasser M. Abbasi

December 9, 2023

Compiled on December 9, 2023 at 10:45am

Contents

1	Introduction	2
2	detailed summary tables of results	20
3	Listing of integrals	118
4	Appendix	1606

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	3
1.2	Results	4
1.3	Time and leaf size Performance	7
1.4	Performance based on number of rules Rubi used	9
1.5	Performance based on number of steps Rubi used	10
1.6	Solved integrals histogram based on leaf size of result	11
1.7	Solved integrals histogram based on CPU time used	12
1.8	Leaf size vs. CPU time used	13
1.9	list of integrals with no known antiderivative	14
1.10	List of integrals solved by CAS but has no known antiderivative	14
1.11	list of integrals solved by CAS but failed verification	14
1.12	Timing	15
1.13	Verification	15
1.14	Important notes about some of the results	15
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [321]. This is test number [211].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 13.3.1 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Mathematica	99.38 (319)	0.62 (2)
Fricas	96.26 (309)	3.74 (12)
Maple	95.33 (306)	4.67 (15)
Rubi	94.70 (304)	5.30 (17)
Maxima	92.52 (297)	7.48 (24)
Giac	91.90 (295)	8.10 (26)
Mupad	90.03 (289)	9.97 (32)
Sympy	82.55 (265)	17.45 (56)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Mathematica	88.162	9.034	0.935	1.869
Rubi	85.358	5.919	2.181	6.542
Maple	83.801	4.673	5.607	5.919
Fricas	76.324	18.069	0.623	4.984
Maxima	74.143	13.396	3.738	8.723
Giac	71.963	16.822	1.869	9.346
Sympy	61.371	16.511	3.427	18.692
Mupad	0.000	88.785	0.000	11.215

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates

an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Mathematica	2	100.00	0.00	0.00
Fricas	12	41.67	25.00	33.33
Maple	15	86.67	13.33	0.00
Rubi	17	100.00	0.00	0.00
Maxima	24	91.67	4.17	4.17
Giac	26	96.15	3.85	0.00
Mupad	32	0.00	100.00	0.00
Sympy	56	87.50	12.50	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Mathematica	0.18
Rubi	0.22
Maxima	0.25
Fricas	0.27
Giac	0.41
Maple	0.47
Sympy	1.58
Mupad	9.08

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	20.48	1.23	13.00	0.86
Fricas	25.39	1.61	17.00	1.00
Rubi	28.70	1.81	18.00	1.00
Maple	88.41	4.21	14.00	0.92
Mathematica	97.55	4.50	18.00	1.00
Maxima	106.78	5.75	15.00	0.97
Giac	112.05	6.80	16.00	1.00
Sympy	132.82	6.21	15.00	0.94

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following gives the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{194, 225, 226, 236}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {7, 115, 219, 315}

Mathematica {}

Maple {81, 276}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

1.14.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.14.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

1.14.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

1.14.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer, 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer, Leaf size of result.
4. integer, Leaf size of the optimal antiderivative.
5. number, CPU time used to solve this integral. 0 if failed.
6. string, The integral in Latex format
7. string, The input used in CAS own syntax.
8. string, The result (antiderivative) produced by CAS in Latex format
9. string, The optimal antiderivative in Latex format.
10. integer, 0 or 1. Indicates if problem has known antiderivative or not
11. String, The result (antiderivative) in CAS own syntax.
12. String, The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String, Small string description of why the grade was given.
14. integer, 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer, Number of steps used.
16. integer, Number of rules used.
17. integer, Integrand leaf size.
18. real number, Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String, The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2023
Design v0.6

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	21
2.2	Detailed conclusion table per each integral for all CAS systems	27
2.3	Detailed conclusion table specific for Rubi results	108

2.1 List of integrals sorted by grade for each CAS

2.1.1	Rubi	21
2.1.2	Mma	22
2.1.3	Maple	22
2.1.4	Fricas	23
2.1.5	Maxima	23
2.1.6	Giac	24
2.1.7	Mupad	25
2.1.8	Sympy	25

2.1.1 Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 42, 43, 44, 45, 47, 48, 49, 50, 51, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 78, 79, 80, 81, 82, 83, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 141, 142, 143, 144, 145, 146, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 181, 182, 183, 184, 185, 186, 188, 189, 190, 191, 192, 193, 195, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 208, 209, 210, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 227, 228, 229, 230, 231, 232, 233, 235, 237, 238, 239, 240, 241, 247, 248, 249, 251, 252, 253, 255, 256, 259, 260, 261, 262, 263, 264, 267, 268, 269, 270, 271, 273, 274, 278, 279, 280, 281, 282, 283, 284, 285, 286, 289, 290, 291, 292, 295, 296, 297, 299, 300, 301, 302, 303, 304, 306, 308, 309, 310, 311, 312, 314, 315, 316, 318, 321 }

B grade { 41, 46, 52, 84, 85, 118, 119, 187, 196, 207, 242, 244, 254, 266, 272, 276, 287, 305, 319 }

C grade { 20, 234, 246, 257, 258, 288, 294 }

F normal fail { 77, 140, 147, 180, 211, 243, 245, 250, 265, 275, 277, 293, 298, 307, 313, 317, 320 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.2 Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 56, 58, 59, 60, 61, 62, 63, 65, 66, 68, 69, 70, 72, 73, 74, 75, 76, 77, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 90, 91, 92, 93, 95, 96, 98, 99, 100, 101, 103, 104, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 169, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 184, 185, 186, 187, 188, 189, 190, 191, 192, 195, 198, 199, 200, 201, 202, 203, 204, 205, 206, 208, 210, 211, 212, 214, 215, 216, 217, 218, 219, 221, 222, 223, 224, 228, 229, 230, 231, 232, 233, 234, 235, 237, 239, 240, 241, 242, 243, 244, 245, 246, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 259, 260, 261, 262, 263, 264, 265, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 291, 292, 293, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 311, 312, 313, 314, 316, 317, 318, 319, 320, 321 }

B grade { 20, 43, 55, 57, 64, 67, 71, 78, 89, 94, 97, 105, 129, 168, 170, 183, 193, 196, 197, 207, 213, 220, 227, 238, 247, 258, 290, 310, 315 }

C grade { 102, 147, 209 }

F normal fail { 266, 294 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.1.3 Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 47, 48, 49, 50, 51, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70, 72, 73, 74, 75, 76, 77, 78, 79, 80, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 133, 134, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 174, 176, 177, 178, 179, 181, 182, 183, 184, 185, 186, 187, 189, 190, 191, 192, 193, 195, 197, 198, 199, 200, 201, 202, 203, 204, 206, 208, 209, 210, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 228, 229, 230, 231, 232, 233, 234, 235, 237, 238, 239, 240, 241, 242, 243, 245, 246, 247, 248, 249, 251, 252, 253, 254, 255, 256, 260, 262, 264, 267, 269, 271, 272, 273, 274, 275, 277, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 292, 295, 296, 297, 298, 299, 300, 301, 303, 305, 306, 308, 310, 311, 314, 316, 321 }

B grade { 19, 46, 52, 64, 71, 105, 135, 180, 196, 211, 227, 304, 309, 312, 319 }

C grade { 20, 81, 116, 131, 132, 205, 207, 244, 257, 258, 261, 263, 268, 276, 278, 293, 313, 317 }

F normal fail { 173, 175, 250, 259, 265, 266, 270, 294, 302, 307, 315, 318, 320 }

F(-1) timeout fail { 188, 291 }

F(-2) exception fail { }

2.1.4 Fricas

A grade { 1, 2, 3, 4, 5, 6, 8, 9, 12, 13, 14, 15, 16, 17, 18, 19, 21, 22, 23, 24, 25, 26, 28, 29, 31, 32, 33, 34, 35, 37, 38, 39, 40, 41, 42, 44, 45, 47, 48, 49, 50, 51, 54, 56, 57, 58, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70, 72, 73, 74, 75, 76, 77, 79, 81, 82, 83, 84, 85, 86, 87, 88, 90, 91, 92, 93, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 122, 123, 124, 125, 126, 127, 128, 130, 131, 132, 133, 134, 136, 138, 139, 141, 142, 143, 144, 145, 146, 147, 148, 150, 151, 152, 153, 154, 155, 157, 158, 160, 161, 162, 163, 164, 165, 166, 167, 171, 172, 173, 174, 175, 176, 177, 179, 181, 183, 184, 185, 187, 188, 189, 191, 192, 195, 198, 201, 202, 203, 204, 206, 208, 209, 210, 212, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 229, 230, 231, 232, 233, 235, 237, 240, 241, 242, 244, 245, 246, 248, 249, 250, 251, 252, 254, 256, 258, 260, 261, 262, 263, 264, 265, 267, 269, 271, 272, 273, 274, 276, 277, 278, 279, 281, 282, 283, 284, 285, 286, 287, 288, 289, 293, 295, 296, 298, 299, 300, 301, 303, 305, 306, 307, 308, 309, 311, 314, 315, 316, 317, 319, 320, 321 }

B grade { 7, 27, 30, 36, 43, 52, 53, 55, 64, 71, 78, 80, 89, 94, 121, 129, 135, 137, 140, 149, 156, 159, 168, 169, 170, 178, 180, 182, 186, 190, 193, 196, 197, 200, 207, 211, 213, 224, 228, 234, 238, 239, 243, 247, 253, 255, 257, 275, 280, 290, 292, 294, 297, 304, 310, 312, 313, 318 }

C grade { 10, 20 }

F normal fail { 11, 205, 266, 268, 270 }

F(-1) timeout fail { 46, 199, 291 }

F(-2) exception fail { 105, 227, 259, 302 }

2.1.5 Maxima

A grade { 1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 21, 22, 24, 25, 28, 31, 33, 34, 35, 37, 38, 39, 40, 41, 44, 45, 47, 48, 49, 50, 51, 53, 54, 55, 56, 57, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70, 72, 73, 75, 76, 77, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 91, 92, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 106, 107, 108, 109, 110, 111, 112, 114, 115, 117, 118, 120, 121, 122, 123, 124, 125, 126, 127, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 146, 148, 149, 151, 152, 153, 154, 155, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 168, 169, 171, 172, 174, 175, 176, 177, 178, 179, 181, 182, 183, 185, 186, 189, 190, 192, 195, 198, 199, 200, 201, 202, 203, 204, 205, 206, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 228,

229, 230, 231, 232, 233, 234, 237, 241, 242, 243, 245, 247, 248, 249, 250, 251, 252, 253, 260, 262, 263, 264, 265, 267, 269, 271, 272, 273, 274, 275, 276, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 291, 292, 295, 296, 298, 300, 304, 306, 308, 310, 311, 312, 314, 316, 317, 318, 319, 321 }
 }

B grade { 7, 19, 26, 27, 29, 30, 32, 36, 43, 64, 71, 74, 78, 89, 90, 93, 94, 105, 113, 128, 145, 156, 180, 184, 187, 193, 196, 197, 207, 208, 227, 238, 239, 244, 254, 255, 258, 261, 290, 293, 297, 301, 303 }

C grade { 23, 119, 147, 150, 167, 173, 246, 256, 277, 289, 302, 309 }

F normal fail { 20, 42, 52, 58, 59, 116, 170, 188, 191, 235, 240, 257, 259, 266, 268, 270, 294, 299, 307, 313, 315, 320 }

F(-1) timeout fail { 305 }

F(-2) exception fail { 46 }

2.1.6 Giac

A grade { 1, 2, 3, 5, 6, 7, 8, 9, 10, 12, 13, 14, 15, 17, 18, 21, 22, 23, 24, 25, 26, 27, 28, 31, 33, 34, 35, 37, 39, 40, 41, 42, 44, 45, 47, 48, 49, 50, 51, 52, 53, 54, 56, 57, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70, 72, 73, 75, 76, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 90, 91, 92, 93, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 106, 107, 108, 109, 110, 111, 112, 114, 115, 118, 120, 122, 123, 124, 125, 126, 127, 130, 131, 132, 133, 134, 135, 136, 138, 139, 140, 141, 142, 143, 144, 146, 147, 148, 149, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 169, 171, 173, 174, 176, 177, 178, 179, 182, 183, 184, 185, 186, 188, 189, 190, 191, 192, 195, 198, 199, 200, 202, 203, 204, 206, 210, 212, 214, 215, 216, 218, 219, 220, 221, 222, 223, 224, 229, 230, 231, 233, 234, 235, 237, 241, 242, 243, 245, 248, 249, 251, 253, 254, 256, 260, 261, 262, 263, 264, 267, 269, 272, 273, 274, 276, 277, 279, 280, 281, 282, 283, 284, 286, 288, 289, 291, 292, 295, 296, 299, 300, 301, 303, 305, 306, 307, 308, 310, 311, 312, 314, 316, 317, 318, 319, 321 }

B grade { 4, 16, 19, 20, 29, 30, 36, 43, 46, 55, 58, 64, 71, 77, 78, 89, 94, 105, 113, 117, 121, 129, 137, 145, 168, 170, 180, 181, 187, 193, 196, 197, 207, 208, 209, 213, 217, 227, 228, 232, 238, 239, 244, 247, 255, 257, 258, 265, 285, 290, 293, 298, 304, 313 }

C grade { 119, 150, 246, 287, 302, 309 }

F normal fail { 11, 32, 38, 59, 74, 116, 128, 172, 175, 201, 205, 211, 240, 252, 259, 266, 268, 270, 271, 275, 278, 294, 297, 315, 320 }

F(-1) timeout fail { 250 }

F(-2) exception fail { }

2.1.7 Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 43, 44, 45, 46, 47, 48, 49, 50, 51, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 192, 193, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 224, 227, 228, 229, 230, 231, 232, 234, 235, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 251, 252, 253, 254, 255, 256, 257, 260, 261, 263, 264, 267, 269, 272, 273, 274, 276, 277, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 292, 293, 295, 296, 298, 299, 300, 301, 302, 303, 304, 305, 306, 308, 310, 311, 312, 313, 314, 316, 317, 319, 321 }

C grade { }

F normal fail { }

F(-1) timedout fail { 11, 42, 52, 63, 81, 93, 115, 116, 172, 191, 205, 223, 233, 250, 258, 259, 262, 265, 266, 268, 270, 271, 275, 278, 291, 294, 297, 307, 309, 315, 318, 320 }

F(-2) exception fail { }

2.1.8 Sympy

A grade { 4, 5, 6, 7, 9, 13, 15, 17, 19, 22, 23, 25, 26, 27, 28, 33, 34, 35, 40, 41, 44, 45, 47, 48, 50, 54, 55, 56, 57, 60, 61, 62, 63, 65, 66, 67, 68, 69, 72, 73, 75, 76, 77, 79, 80, 82, 83, 84, 85, 86, 88, 90, 91, 92, 95, 96, 97, 99, 100, 101, 102, 104, 106, 107, 108, 109, 110, 111, 114, 116, 117, 120, 123, 124, 125, 126, 127, 129, 130, 131, 133, 134, 135, 136, 138, 139, 140, 141, 143, 144, 145, 146, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 160, 162, 164, 165, 166, 167, 168, 171, 174, 176, 177, 178, 179, 180, 181, 182, 183, 185, 186, 187, 190, 192, 195, 198, 199, 200, 201, 202, 203, 205, 206, 208, 209, 210, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 224, 229, 230, 231, 232, 237, 240, 242, 245, 247, 248, 249, 251, 253, 254, 255, 256, 261, 262, 263, 264, 265, 269, 273, 274, 279, 281, 282, 283, 285, 286, 287, 289, 293, 295, 296, 298, 300, 301, 305, 306, 308, 309, 310, 311, 312, 314, 316, 321 }

B grade { 1, 2, 14, 16, 18, 29, 30, 31, 36, 38, 39, 43, 51, 53, 64, 70, 71, 78, 87, 89, 93, 94, 103, 105, 118, 119, 121, 137, 142, 159, 189, 193, 197, 204, 211, 227, 228, 234, 238, 239, 244, 260, 268, 272, 280, 284, 288, 292, 294, 302, 303, 304, 315 }

C grade { 3, 8, 24, 98, 112, 161, 223, 241, 252, 267, 278 }

F normal fail { 10, 11, 12, 20, 21, 32, 37, 42, 49, 52, 58, 59, 74, 81, 113, 115, 122, 128, 147, 163, 169, 170, 172, 173, 175, 184, 191, 233, 235, 243, 246, 250, 257, 258, 259, 266, 270, 271, 275, 276, 277, 290, 297, 299, 307, 313, 317, 318, 320 }

F(-1) timeout fail { 46, 132, 188, 196, 207, 291, 319 }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	20	19	17	114	13	14
N.S.	1	1.00	1.00	0.80	0.76	0.68	4.56	0.52	0.56
time (sec)	N/A	0.198	0.008	1.017	0.192	0.278	1.041	0.270	16.607

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	19	31	24	23	13	65	13	14
N.S.	1	1.12	1.82	1.41	1.35	0.76	3.82	0.76	0.82
time (sec)	N/A	0.232	0.021	0.212	0.185	0.272	0.591	0.266	0.056

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	32	21	18	22	22	144	22	19
N.S.	1	1.52	1.00	0.86	1.05	1.05	6.86	1.05	0.90
time (sec)	N/A	0.140	0.023	0.276	0.192	0.249	0.856	0.260	16.727

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	26	27	22	21	23	20	60	31
N.S.	1	0.96	1.00	0.81	0.78	0.85	0.74	2.22	1.15
time (sec)	N/A	0.161	0.012	0.069	0.188	0.259	0.044	0.269	16.963

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	26	18	20	21	37	21	24
N.S.	1	1.00	0.96	0.67	0.74	0.78	1.37	0.78	0.89
time (sec)	N/A	0.145	0.012	0.276	0.197	0.254	0.194	0.262	16.198

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	15	15	10	11	11	8	11	11
N.S.	1	1.25	1.25	0.83	0.92	0.92	0.67	0.92	0.92
time (sec)	N/A	0.139	0.018	0.023	0.184	0.272	0.033	0.267	0.062

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	A	A	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	31	33	50	69	46	46	30
N.S.	1	1.00	1.03	1.10	1.67	2.30	1.53	1.53	1.00
time (sec)	N/A	0.189	0.028	0.290	0.200	0.270	0.076	0.269	15.254

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	14	11	10	10	24	10	10
N.S.	1	1.00	1.00	0.79	0.71	0.71	1.71	0.71	0.71
time (sec)	N/A	0.138	0.005	0.383	0.304	0.249	0.445	0.260	15.308

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	17	13	12	15	11	10	13	11
N.S.	1	1.31	1.00	0.92	1.15	0.85	0.77	1.00	0.85
time (sec)	N/A	0.130	0.004	0.141	0.196	0.251	0.040	0.266	0.080

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	71	110	40	49	80	73	0	80	65
N.S.	1	1.55	0.56	0.69	1.13	1.03	0.00	1.13	0.92
time (sec)	N/A	0.302	0.049	0.125	0.283	0.270	0.000	0.270	0.154

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	89	89	70	56	0	0	0	0
N.S.	1	1.00	1.00	0.79	0.63	0.00	0.00	0.00	0.00
time (sec)	N/A	0.232	0.013	0.224	0.298	0.000	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	63	51	36	35	35	0	36	35
N.S.	1	1.19	0.96	0.68	0.66	0.66	0.00	0.68	0.66
time (sec)	N/A	0.174	0.027	0.185	0.185	0.240	0.000	0.286	0.063

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	3	3	2	3	3
N.S.	1	1.00	1.00	1.33	1.00	1.00	0.67	1.00	1.00
time (sec)	N/A	0.182	0.014	0.020	0.216	0.260	0.050	0.269	16.258

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	50	29	36	28	31	66	28	25
N.S.	1	1.72	1.00	1.24	0.97	1.07	2.28	0.97	0.86
time (sec)	N/A	0.156	0.060	0.182	0.194	0.296	84.094	0.277	16.617

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	16	16	15	19	12	15	16
N.S.	1	1.00	0.84	0.84	0.79	1.00	0.63	0.79	0.84
time (sec)	N/A	0.127	0.008	0.207	0.263	0.253	0.045	0.278	0.037

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	37	26	25	25	26	29	13
N.S.	1	1.00	1.76	1.24	1.19	1.19	1.24	1.38	0.62
time (sec)	N/A	0.135	0.006	0.023	0.194	0.253	0.078	0.273	15.978

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	11	12	11	11	10	11	8
N.S.	1	1.00	1.00	1.09	1.00	1.00	0.91	1.00	0.73
time (sec)	N/A	0.140	0.001	0.026	0.197	0.270	0.064	0.284	15.865

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	14	15	11	15	16	14
N.S.	1	1.00	1.00	1.75	1.88	1.38	1.88	2.00	1.75
time (sec)	N/A	0.177	0.135	0.062	0.206	0.286	0.084	0.279	16.238

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	10	25	65	10	8	79	10
N.S.	1	1.00	0.91	2.27	5.91	0.91	0.73	7.18	0.91
time (sec)	N/A	0.318	0.161	0.206	0.296	0.265	0.072	0.286	0.104

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	B	C	F	C	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	111	93	57	0	165	0	81	233
N.S.	1	3.96	3.32	2.04	0.00	5.89	0.00	2.89	8.32
time (sec)	N/A	0.424	1.251	0.213	0.000	0.288	0.000	0.294	16.058

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	23	20	18	37	0	37	21
N.S.	1	1.00	0.88	0.77	0.69	1.42	0.00	1.42	0.81
time (sec)	N/A	0.168	0.098	0.028	0.280	0.259	0.000	0.286	16.079

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	14	12	11	11	10	11	11
N.S.	1	1.00	0.64	0.55	0.50	0.50	0.45	0.50	0.50
time (sec)	N/A	0.170	0.026	0.024	0.188	0.257	0.033	0.279	15.787

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	6	7	12	6	3	6	6
N.S.	1	1.00	1.00	1.17	2.00	1.00	0.50	1.00	1.00
time (sec)	N/A	0.133	0.003	0.014	0.245	0.252	0.036	0.276	15.230

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	53	53	61	38	52	124	42	41
N.S.	1	1.00	1.00	1.15	0.72	0.98	2.34	0.79	0.77
time (sec)	N/A	0.166	0.118	0.217	0.284	0.251	1.886	0.287	0.122

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	13	12	12	10	12	13
N.S.	1	1.00	1.00	0.81	0.75	0.75	0.62	0.75	0.81
time (sec)	N/A	0.158	0.001	0.015	0.199	0.254	0.023	0.271	0.029

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	6	7	21	6	5	6	6
N.S.	1	1.00	1.00	1.17	3.50	1.00	0.83	1.00	1.00
time (sec)	N/A	0.133	0.001	0.009	0.195	0.261	0.033	0.276	14.715

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	4	4	4	5	16	18	20	4	14
N.S.	1	1.00	1.00	1.25	4.00	4.50	5.00	1.00	3.50
time (sec)	N/A	0.131	0.133	0.223	0.274	0.253	0.461	0.267	14.897

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	18	16	15	15	12	16	15
N.S.	1	1.00	1.06	0.94	0.88	0.88	0.71	0.94	0.88
time (sec)	N/A	0.141	0.005	0.182	0.203	0.255	0.028	0.265	0.035

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	31	12	301	29	162	39	39
N.S.	1	1.00	1.82	0.71	17.71	1.71	9.53	2.29	2.29
time (sec)	N/A	0.259	0.081	0.135	0.281	0.259	8.203	0.276	14.988

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	4	4	4	5	10	20	15	10	4
N.S.	1	1.00	1.00	1.25	2.50	5.00	3.75	2.50	1.00
time (sec)	N/A	0.162	0.004	0.226	0.191	0.235	0.228	0.254	0.019

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	27	14	13	20	29	13	13
N.S.	1	1.00	1.59	0.82	0.76	1.18	1.71	0.76	0.76
time (sec)	N/A	0.186	0.045	1.000	0.189	0.262	0.021	0.272	14.958

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	188	19	0	0	15
N.S.	1	1.00	1.00	0.92	14.46	1.46	0.00	0.00	1.15
time (sec)	N/A	0.292	0.029	0.598	0.367	0.265	0.000	0.000	15.457

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	44	30	23	24	25	36	22	22
N.S.	1	1.29	0.88	0.68	0.71	0.74	1.06	0.65	0.65
time (sec)	N/A	0.247	0.003	0.256	0.183	0.280	0.018	0.265	0.054

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	16	16	15	19	12	15	16
N.S.	1	1.00	0.84	0.84	0.79	1.00	0.63	0.79	0.84
time (sec)	N/A	0.137	0.006	0.019	0.280	0.252	0.041	0.265	0.033

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	11	10	13	15	13	13
N.S.	1	1.00	1.00	0.65	0.59	0.76	0.88	0.76	0.76
time (sec)	N/A	0.143	0.007	0.039	0.186	0.265	0.112	0.262	14.864

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	3	7	15	17	15	17	11
N.S.	1	1.00	1.00	2.33	5.00	5.67	5.00	5.67	3.67
time (sec)	N/A	0.140	0.001	0.053	0.212	0.257	0.058	0.266	0.055

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	11	11	8	7	21	0	20	16
N.S.	1	0.46	0.46	0.33	0.29	0.88	0.00	0.83	0.67
time (sec)	N/A	0.180	0.089	0.320	0.305	0.260	0.000	0.272	15.662

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	14	13	12	12	22	0	12
N.S.	1	1.00	1.00	0.93	0.86	0.86	1.57	0.00	0.86
time (sec)	N/A	0.127	0.004	0.175	0.200	0.269	0.325	0.000	15.016

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	31	24	23	30	66	23	27
N.S.	1	1.00	1.00	0.77	0.74	0.97	2.13	0.74	0.87
time (sec)	N/A	0.204	0.024	0.698	0.194	0.263	0.831	0.271	15.136

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	5	5	6	5	5	5	5	5
N.S.	1	1.00	1.00	1.20	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.129	0.003	0.019	0.229	0.259	0.085	0.269	15.168

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	19	8	12	11	11	10	11	11
N.S.	1	2.38	1.00	1.50	1.38	1.38	1.25	1.38	1.38
time (sec)	N/A	0.140	0.017	0.014	0.210	0.255	0.029	0.260	0.052

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	48	43	0	33	0	16	0
N.S.	1	1.00	1.17	1.05	0.00	0.80	0.00	0.39	0.00
time (sec)	N/A	0.205	0.277	0.784	0.000	0.278	0.000	0.272	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	25	7	15	15	12	16	6
N.S.	1	1.00	4.17	1.17	2.50	2.50	2.00	2.67	1.00
time (sec)	N/A	0.128	0.005	0.203	0.204	0.242	0.037	0.264	0.064

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	3	3	2	3	3
N.S.	1	1.00	1.00	1.33	1.00	1.00	0.67	1.00	1.00
time (sec)	N/A	0.181	0.003	0.000	0.232	0.250	0.051	0.260	0.003

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	37	47	27	36	35	26	25	26
N.S.	1	1.28	1.62	0.93	1.24	1.21	0.90	0.86	0.90
time (sec)	N/A	0.154	0.076	0.342	0.286	0.254	0.203	0.271	14.398

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	B	F(-2)	F(-1)	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	801	12	602	0	0	0	601	601
N.S.	1	66.75	1.00	50.17	0.00	0.00	0.00	50.08	50.08
time (sec)	N/A	0.972	0.008	1.155	0.000	0.000	0.000	0.286	21.211

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	10	11	7	7	7	7	7
N.S.	1	1.00	1.00	1.10	0.70	0.70	0.70	0.70	0.70
time (sec)	N/A	0.214	0.022	0.046	0.220	0.249	0.656	0.263	0.067

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	28	24	22	32	46	29	23	21
N.S.	1	1.00	0.86	0.79	1.14	1.64	1.04	0.82	0.75
time (sec)	N/A	0.204	0.016	0.024	0.206	0.254	0.040	0.273	0.064

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	57	64	39	43	32	0	29	43
N.S.	1	1.50	1.68	1.03	1.13	0.84	0.00	0.76	1.13
time (sec)	N/A	0.156	0.107	0.118	0.308	0.242	0.000	0.268	0.057

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	18	18	15	15	14	15	15	14
N.S.	1	0.90	0.90	0.75	0.75	0.70	0.75	0.75	0.70
time (sec)	N/A	0.147	0.015	0.151	0.213	0.240	0.049	0.271	0.071

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	15	14	13	14	105	13	9
N.S.	1	1.00	0.71	0.67	0.62	0.67	5.00	0.62	0.43
time (sec)	N/A	0.143	0.006	0.054	0.186	0.244	1.661	0.267	15.093

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	B	F	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	51	21	41	0	37	0	21	0
N.S.	1	2.55	1.05	2.05	0.00	1.85	0.00	1.05	0.00
time (sec)	N/A	0.226	0.121	0.155	0.000	0.260	0.000	0.279	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	29	22	20	19	186	138	16	53
N.S.	1	1.07	0.81	0.74	0.70	6.89	5.11	0.59	1.96
time (sec)	N/A	0.159	0.031	0.157	0.200	0.246	53.393	0.276	15.164

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	37	18	17	31	15	17	17
N.S.	1	1.00	1.61	0.78	0.74	1.35	0.65	0.74	0.74
time (sec)	N/A	0.136	0.064	0.260	0.277	0.248	0.070	0.277	0.032

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	17	23	10	11	21	10	29	11
N.S.	1	1.70	2.30	1.00	1.10	2.10	1.00	2.90	1.10
time (sec)	N/A	0.145	0.089	0.216	0.300	0.244	0.251	0.272	14.487

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	9	7	8	7	7	8
N.S.	1	1.00	1.00	1.12	0.88	1.00	0.88	0.88	1.00
time (sec)	N/A	0.146	0.001	0.010	0.191	0.252	0.035	0.271	0.021

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	23	11	15	18	8	10	10
N.S.	1	1.00	2.30	1.10	1.50	1.80	0.80	1.00	1.00
time (sec)	N/A	0.157	0.007	0.039	0.184	0.249	0.171	0.271	0.026

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	58	70	53	0	64	0	117	64
N.S.	1	1.09	1.32	1.00	0.00	1.21	0.00	2.21	1.21
time (sec)	N/A	0.272	0.208	0.022	0.000	0.240	0.000	0.308	16.545

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	46	18	0	18	0	0	179
N.S.	1	1.00	1.00	0.39	0.00	0.39	0.00	0.00	3.89
time (sec)	N/A	0.321	0.015	0.552	0.000	0.258	0.000	0.000	0.248

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	37	35	29	32	44	27	28	32
N.S.	1	1.00	0.95	0.78	0.86	1.19	0.73	0.76	0.86
time (sec)	N/A	0.205	0.014	0.023	0.190	0.236	0.037	0.267	13.811

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	5	5	6	5	5	5	5	5
N.S.	1	1.00	1.00	1.20	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.156	0.024	0.166	0.243	0.250	3.730	0.270	14.435

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	19	17	14	17	13	10	16	13
N.S.	1	1.12	1.00	0.82	1.00	0.76	0.59	0.94	0.76
time (sec)	N/A	0.148	0.005	0.159	0.201	0.236	0.041	0.263	0.075

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	16	15	15	12	15	0
N.S.	1	1.00	1.00	0.94	0.88	0.88	0.71	0.88	0.00
time (sec)	N/A	0.174	0.012	0.155	0.276	0.254	0.072	0.274	0.000

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	567	501	501	501	561	501	12
N.S.	1	1.00	24.65	21.78	21.78	21.78	24.39	21.78	0.52
time (sec)	N/A	0.169	0.010	0.188	0.198	0.238	0.079	0.269	13.917

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	11	12	12	11	11
N.S.	1	1.00	1.00	0.92	0.85	0.92	0.92	0.85	0.85
time (sec)	N/A	0.185	0.014	0.078	0.200	0.269	0.265	0.268	14.361

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	18	16	13	12	12	15	12	12
N.S.	1	1.12	1.00	0.81	0.75	0.75	0.94	0.75	0.75
time (sec)	N/A	0.142	0.036	0.156	0.270	0.244	0.408	0.272	14.481

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	54	22	24	28	26	36	21
N.S.	1	1.00	2.16	0.88	0.96	1.12	1.04	1.44	0.84
time (sec)	N/A	0.137	0.033	0.192	0.274	0.242	0.071	0.271	0.155

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	10	9	9	10	9	9
N.S.	1	1.00	1.00	0.83	0.75	0.75	0.83	0.75	0.75
time (sec)	N/A	0.149	0.030	0.041	0.282	0.232	0.221	0.272	0.135

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	9	8	8	7	8	8
N.S.	1	1.00	1.00	0.75	0.67	0.67	0.58	0.67	0.67
time (sec)	N/A	0.134	0.006	0.170	0.337	0.237	0.042	0.269	15.463

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	13	13	9	8	15	22	8	13
N.S.	1	1.62	1.62	1.12	1.00	1.88	2.75	1.00	1.62
time (sec)	N/A	0.169	0.031	0.140	0.196	0.252	0.327	0.299	15.118

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	39	27	26	80	22	49	30
N.S.	1	1.00	3.55	2.45	2.36	7.27	2.00	4.45	2.73
time (sec)	N/A	0.194	0.007	0.204	0.226	0.231	0.222	0.283	0.096

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	31	20	17	19	19	41	19	19
N.S.	1	1.15	0.74	0.63	0.70	0.70	1.52	0.70	0.70
time (sec)	N/A	0.153	0.024	0.231	0.183	0.245	0.093	0.277	0.051

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	10	16	11	9	10	11	10
N.S.	1	1.00	1.00	1.60	1.10	0.90	1.00	1.10	1.00
time (sec)	N/A	0.176	0.033	0.025	0.214	0.237	0.563	0.268	18.822

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	188	19	0	0	15
N.S.	1	1.00	1.00	0.92	14.46	1.46	0.00	0.00	1.15
time (sec)	N/A	0.283	0.022	0.434	0.363	0.258	0.000	0.000	0.002

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	14	15	14	5	7	14	16
N.S.	1	1.00	1.00	1.07	1.00	0.36	0.50	1.00	1.14
time (sec)	N/A	0.139	0.004	0.016	0.188	0.246	0.027	0.270	0.088

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	3	3	2	2	2	2	2
N.S.	1	1.00	1.00	1.00	0.67	0.67	0.67	0.67	0.67
time (sec)	N/A	0.126	0.000	0.013	0.198	0.260	0.025	0.278	0.009

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	A	A	A	A	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	7	0	7	8	7	7	5	20	7
N.S.	1	0.00	1.00	1.14	1.00	1.00	0.71	2.86	1.00
time (sec)	N/A	0.000	0.887	0.151	0.254	0.256	0.329	0.277	16.800

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	23	8	19	19	15	19	19
N.S.	1	1.00	2.09	0.73	1.73	1.73	1.36	1.73	1.73
time (sec)	N/A	0.144	0.000	0.009	0.183	0.242	0.015	0.268	0.033

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	43	23	22	36	27	23	25
N.S.	1	1.00	1.23	0.66	0.63	1.03	0.77	0.66	0.71
time (sec)	N/A	0.145	0.121	0.321	0.263	0.247	0.129	0.289	0.036

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	29	20	19	45	15	19	45
N.S.	1	1.00	1.07	0.74	0.70	1.67	0.56	0.70	1.67
time (sec)	N/A	0.150	0.040	0.193	0.196	0.260	0.048	0.264	16.237

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	15	15	15	36	13	5	0	6	0
N.S.	1	1.00	1.00	2.40	0.87	0.33	0.00	0.40	0.00
time (sec)	N/A	0.262	0.020	0.372	0.193	0.270	0.000	0.282	0.000

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	9	9	9	7	7	7	5	7	6
N.S.	1	1.00	1.00	0.78	0.78	0.78	0.56	0.78	0.67
time (sec)	N/A	0.141	0.000	0.010	0.191	0.224	0.019	0.272	0.020

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	9	8	8	5	8	6
N.S.	1	1.00	1.00	1.12	1.00	1.00	0.62	1.00	0.75
time (sec)	N/A	0.129	0.000	0.013	0.186	0.258	0.038	0.268	0.020

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	19	6	10	9	9	8	10	7
N.S.	1	3.17	1.00	1.67	1.50	1.50	1.33	1.67	1.17
time (sec)	N/A	0.153	0.016	0.019	0.203	0.246	0.036	0.282	15.700

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	29	14	11	10	19	14	10	18
N.S.	1	2.07	1.00	0.79	0.71	1.36	1.00	0.71	1.29
time (sec)	N/A	0.210	0.007	0.074	0.200	0.238	0.023	0.271	17.683

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	10	9	8	8	10	8	8
N.S.	1	1.00	1.00	0.90	0.80	0.80	1.00	0.80	0.80
time (sec)	N/A	0.187	0.033	0.073	0.202	0.281	0.106	0.274	17.252

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	8	7	6	6	7	6	6
N.S.	1	1.00	1.33	1.17	1.00	1.00	1.17	1.00	1.00
time (sec)	N/A	0.147	0.002	0.016	0.268	0.255	0.023	0.269	0.038

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	57	31	26	21	21	48	21	28
N.S.	1	1.90	1.03	0.87	0.70	0.70	1.60	0.70	0.93
time (sec)	N/A	0.239	0.014	0.016	0.208	0.238	0.139	0.282	0.028

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	19	12	17	21	19	19	18
N.S.	1	1.00	2.38	1.50	2.12	2.62	2.38	2.38	2.25
time (sec)	N/A	0.166	0.006	0.096	0.197	0.268	0.048	0.275	16.474

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	6	7	21	6	5	6	6
N.S.	1	1.00	1.00	1.17	3.50	1.00	0.83	1.00	1.00
time (sec)	N/A	0.134	0.001	0.010	0.199	0.250	0.032	0.266	0.002

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	16	18	15	14	14	12	14	14
N.S.	1	0.89	1.00	0.83	0.78	0.78	0.67	0.78	0.78
time (sec)	N/A	0.144	0.004	0.151	0.188	0.235	0.028	0.278	16.802

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	5	4	4	3	4	4
N.S.	1	1.00	1.00	0.62	0.50	0.50	0.38	0.50	0.50
time (sec)	N/A	0.131	0.005	0.299	0.286	0.259	0.039	0.275	16.309

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	B	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	26	20	89	28	105	26	0
N.S.	1	1.00	1.73	1.33	5.93	1.87	7.00	1.73	0.00
time (sec)	N/A	0.204	0.015	0.244	0.193	0.283	0.568	0.283	0.000

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	23	7	17	17	15	18	6
N.S.	1	1.00	2.88	0.88	2.12	2.12	1.88	2.25	0.75
time (sec)	N/A	0.128	0.006	0.184	0.188	0.240	0.040	0.269	16.744

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	43	23	22	36	27	23	25
N.S.	1	1.00	1.23	0.66	0.63	1.03	0.77	0.66	0.71
time (sec)	N/A	0.138	0.001	0.184	0.274	0.269	0.127	0.273	0.002

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	14	14	14	14	13
N.S.	1	1.00	1.00	0.82	0.82	0.82	0.82	0.82	0.76
time (sec)	N/A	0.183	0.028	3.794	0.198	0.259	0.047	0.275	16.449

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	25	11	15	17	8	10	10
N.S.	1	1.00	2.27	1.00	1.36	1.55	0.73	0.91	0.91
time (sec)	N/A	0.158	0.022	0.054	0.200	0.246	0.175	0.270	0.025

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	22	18	14	24	39	18	18
N.S.	1	1.00	1.00	0.82	0.64	1.09	1.77	0.82	0.82
time (sec)	N/A	0.141	0.007	0.234	0.275	0.251	0.502	0.272	17.029

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	10	11	10	10	8	10	8
N.S.	1	1.00	1.00	1.10	1.00	1.00	0.80	1.00	0.80
time (sec)	N/A	0.128	0.000	0.010	0.189	0.257	0.029	0.271	0.017

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	37	29	26	21	21	34	21	21
N.S.	1	1.06	0.83	0.74	0.60	0.60	0.97	0.60	0.60
time (sec)	N/A	0.274	0.028	0.025	0.186	0.253	0.150	0.277	16.283

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	10	10	9	9	7	9	11
N.S.	1	1.00	1.00	1.00	0.90	0.90	0.70	0.90	1.10
time (sec)	N/A	0.156	0.061	0.031	0.226	0.277	0.040	0.277	0.071

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	37	11	19	14	14	31	14	23
N.S.	1	1.09	0.32	0.56	0.41	0.41	0.91	0.41	0.68
time (sec)	N/A	0.224	0.004	0.010	0.194	0.252	0.072	0.268	17.718

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	12	11	13	26	11	13
N.S.	1	1.00	1.00	0.80	0.73	0.87	1.73	0.73	0.87
time (sec)	N/A	0.173	0.008	0.102	0.201	0.244	0.127	0.278	0.054

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	6	7	6	6	5	6	6
N.S.	1	1.00	1.00	1.17	1.00	1.00	0.83	1.00	1.00
time (sec)	N/A	0.133	0.003	0.036	0.212	0.249	0.019	0.273	18.421

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	F(-2)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	12138	10076	10076	0	12024	10076	15
N.S.	1	1.00	527.74	438.09	438.09	0.00	522.78	438.09	0.65
time (sec)	N/A	0.869	0.132	7.625	2.294	0.000	2.415	1.638	34.755

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	14	22	12	22	14
N.S.	1	1.00	1.00	0.94	0.88	1.38	0.75	1.38	0.88
time (sec)	N/A	0.154	0.004	0.008	0.185	0.265	0.049	0.277	19.877

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	16	12	11	11	10	12	11
N.S.	1	1.00	1.07	0.80	0.73	0.73	0.67	0.80	0.73
time (sec)	N/A	0.139	0.004	0.168	0.187	0.249	0.024	0.272	0.029

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	20	16	16	15	13	15	15	14
N.S.	1	0.95	0.76	0.76	0.71	0.62	0.71	0.71	0.67
time (sec)	N/A	0.160	0.006	0.038	0.270	0.237	0.079	0.270	0.029

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	14	13	13	12	15	8
N.S.	1	1.00	1.00	0.74	0.68	0.68	0.63	0.79	0.42
time (sec)	N/A	0.148	0.004	0.179	0.185	0.247	0.045	0.267	0.175

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	28	16	18	17	17	19	27	15
N.S.	1	1.47	0.84	0.95	0.89	0.89	1.00	1.42	0.79
time (sec)	N/A	0.398	0.046	0.198	0.339	0.245	0.789	0.267	21.299

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	26	25	24	23	23	22	23	22
N.S.	1	1.04	1.00	0.96	0.92	0.92	0.88	0.92	0.88
time (sec)	N/A	0.218	0.007	0.032	0.273	0.249	0.066	0.265	0.033

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	28	22	17	13	22	68	18	26
N.S.	1	1.27	1.00	0.77	0.59	1.00	3.09	0.82	1.18
time (sec)	N/A	0.147	0.006	0.245	0.274	0.255	0.704	0.263	0.105

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	16	74	28	0	103	15
N.S.	1	1.00	1.00	0.84	3.89	1.47	0.00	5.42	0.79
time (sec)	N/A	0.218	0.012	0.131	0.280	0.267	0.000	0.280	20.914

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	29	24	19	19	18	15	22	18
N.S.	1	1.16	0.96	0.76	0.76	0.72	0.60	0.88	0.72
time (sec)	N/A	0.176	0.007	0.023	0.195	0.254	0.050	0.267	19.619

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	6	11	17	16	11	0	12	0
N.S.	1	0.55	1.00	1.55	1.45	1.00	0.00	1.09	0.00
time (sec)	N/A	0.195	0.071	0.193	0.208	0.233	0.000	0.262	0.000

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	A	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	41	37	57	0	26	42	0	0
N.S.	1	1.11	1.00	1.54	0.00	0.70	1.14	0.00	0.00
time (sec)	N/A	0.171	0.010	0.034	0.000	0.260	0.118	0.000	0.000

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	30	20	16	19	23	32	40	32
N.S.	1	1.50	1.00	0.80	0.95	1.15	1.60	2.00	1.60
time (sec)	N/A	0.178	0.024	0.046	0.283	0.247	0.113	0.273	21.007

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	29	8	7	6	5	29	6	6
N.S.	1	5.80	1.60	1.40	1.20	1.00	5.80	1.20	1.20
time (sec)	N/A	0.165	0.004	0.240	0.190	0.252	0.022	0.263	20.292

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	A	C	A	B	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	37	16	13	19	12	31	22	14
N.S.	1	2.31	1.00	0.81	1.19	0.75	1.94	1.38	0.88
time (sec)	N/A	0.149	0.024	0.167	0.179	0.253	0.529	0.262	0.044

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	7	6	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.88	0.75	0.75
time (sec)	N/A	0.168	0.003	0.029	0.173	0.262	0.100	0.270	20.116

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	3	7	6	17	15	25	11
N.S.	1	1.00	1.00	2.33	2.00	5.67	5.00	8.33	3.67
time (sec)	N/A	0.145	0.000	0.022	0.177	0.250	0.060	0.264	0.004

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	4	4	4	4	3	3	0	3	3
N.S.	1	1.00	1.00	1.00	0.75	0.75	0.00	0.75	0.75
time (sec)	N/A	0.154	0.009	0.438	0.189	0.239	0.000	0.265	17.480

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	28	28	23	17	22	22	22	17
N.S.	1	1.00	1.00	0.82	0.61	0.79	0.79	0.79	0.61
time (sec)	N/A	0.162	0.001	0.013	0.185	0.262	0.046	0.260	0.079

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	34	22	19	18	18	20	18	18
N.S.	1	1.55	1.00	0.86	0.82	0.82	0.91	0.82	0.82
time (sec)	N/A	0.178	0.019	0.114	0.279	0.246	0.077	0.274	0.144

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	9	8	8	5	8	8
N.S.	1	1.00	1.00	1.12	1.00	1.00	0.62	1.00	1.00
time (sec)	N/A	0.133	0.001	0.033	0.183	0.247	0.031	0.262	0.103

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	28	28	21	20	27	22	21	22
N.S.	1	1.27	1.27	0.95	0.91	1.23	1.00	0.95	1.00
time (sec)	N/A	0.153	0.012	0.227	0.196	0.239	0.039	0.274	16.606

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	16	15	17	16	15	19	16	18
N.S.	1	1.07	1.00	1.13	1.07	1.00	1.27	1.07	1.20
time (sec)	N/A	0.176	0.013	0.038	0.201	0.245	0.078	0.266	15.224

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	188	19	0	0	15
N.S.	1	1.00	1.00	0.92	14.46	1.46	0.00	0.00	1.15
time (sec)	N/A	0.282	0.021	0.436	0.356	0.260	0.000	0.000	0.002

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	16	5	4	14	3	25	4
N.S.	1	1.00	2.67	0.83	0.67	2.33	0.50	4.17	0.67
time (sec)	N/A	0.118	0.020	0.204	0.264	0.256	0.057	0.266	0.034

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	7	7	7	8	7	11	7	7	7
N.S.	1	1.00	1.00	1.14	1.00	1.57	1.00	1.00	1.00
time (sec)	N/A	0.174	0.001	0.023	0.207	0.266	0.028	0.263	17.156

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	34	31	47	25	25	27	25	25
N.S.	1	0.97	0.89	1.34	0.71	0.71	0.77	0.71	0.71
time (sec)	N/A	0.230	0.018	0.222	0.199	0.256	3.329	0.268	15.271

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	71	37	32	40	39	47	0	48	44
N.S.	1	0.52	0.45	0.56	0.55	0.66	0.00	0.68	0.62
time (sec)	N/A	0.224	0.071	0.192	0.283	0.266	0.000	0.269	18.243

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	15	13	10	9	9	8	10	9
N.S.	1	1.25	1.08	0.83	0.75	0.75	0.67	0.83	0.75
time (sec)	N/A	0.166	0.028	0.025	0.197	0.254	0.033	0.277	0.061

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	16	13	12	12	19	12	12
N.S.	1	1.00	1.14	0.93	0.86	0.86	1.36	0.86	0.86
time (sec)	N/A	0.192	0.007	0.027	0.272	0.241	0.027	0.267	0.041

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	5	5	20	7	11	7	7	7
N.S.	1	1.00	1.00	4.00	1.40	2.20	1.40	1.40	1.40
time (sec)	N/A	0.176	0.005	0.224	0.184	0.257	0.031	0.267	15.419

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	11	11	10	11	11
N.S.	1	1.00	1.00	0.92	0.85	0.85	0.77	0.85	0.85
time (sec)	N/A	0.125	0.004	0.471	0.184	0.235	0.031	0.275	0.039

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	3	18	7	11	3
N.S.	1	1.00	1.00	1.33	1.00	6.00	2.33	3.67	1.00
time (sec)	N/A	0.140	0.002	0.034	0.187	0.234	0.051	0.279	0.027

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	11	11	11	8	11	10
N.S.	1	1.00	1.00	0.73	0.73	0.73	0.53	0.73	0.67
time (sec)	N/A	0.134	0.000	0.011	0.179	0.236	0.014	0.270	0.022

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	12	10	9	9	8	9	7
N.S.	1	1.00	0.86	0.71	0.64	0.64	0.57	0.64	0.50
time (sec)	N/A	0.131	0.000	0.012	0.179	0.232	0.032	0.266	0.025

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	A	A	B	A	A	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	0	11	9	8	20	14	12	8
N.S.	1	0.00	1.00	0.82	0.73	1.82	1.27	1.09	0.73
time (sec)	N/A	0.000	0.043	0.195	0.278	0.235	0.104	0.276	14.764

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	9	9	9	8	7	7	7	7	7
N.S.	1	1.00	1.00	0.89	0.78	0.78	0.78	0.78	0.78
time (sec)	N/A	0.173	0.006	0.030	0.191	0.249	0.068	0.263	15.846

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	14	11	16	20	36	16	10
N.S.	1	1.00	1.00	0.79	1.14	1.43	2.57	1.14	0.71
time (sec)	N/A	0.191	0.004	0.043	0.272	0.258	0.188	0.273	13.510

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	45	50	39	34	33	25	37	49	25
N.S.	1	1.11	0.87	0.76	0.73	0.56	0.82	1.09	0.56
time (sec)	N/A	0.193	0.052	0.053	0.276	4.288	0.166	0.279	14.143

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	9	8	8	5	8	6
N.S.	1	1.00	1.00	1.12	1.00	1.00	0.62	1.00	0.75
time (sec)	N/A	0.135	0.000	0.008	0.203	0.231	0.032	0.276	0.002

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	6	6	21	5	5	21	5
N.S.	1	1.00	1.00	1.00	3.50	0.83	0.83	3.50	0.83
time (sec)	N/A	0.142	0.007	0.086	0.187	0.241	0.086	0.291	0.054

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	16	14	13	13	12	13	13
N.S.	1	1.00	0.62	0.54	0.50	0.50	0.46	0.50	0.50
time (sec)	N/A	0.177	0.006	0.035	0.184	0.243	0.032	0.262	16.547

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	C	A	C	A	F	A	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	0	27	10	473	12	0	12	9
N.S.	1	0.00	2.70	1.00	47.30	1.20	0.00	1.20	0.90
time (sec)	N/A	0.000	0.627	0.212	0.279	0.247	0.000	0.266	0.084

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	56	62	50	35	34	32	51	34	34
N.S.	1	1.11	0.89	0.62	0.61	0.57	0.91	0.61	0.61
time (sec)	N/A	0.315	0.046	0.026	0.188	0.232	4.522	0.284	0.033

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	9	9	9	8	7	21	8	7	7
N.S.	1	1.00	1.00	0.89	0.78	2.33	0.89	0.78	0.78
time (sec)	N/A	0.197	2.049	0.220	0.182	0.238	0.274	0.279	0.148

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	A	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	27	10	19	29	14
N.S.	1	1.00	1.00	0.93	1.80	0.67	1.27	1.93	0.93
time (sec)	N/A	0.219	0.045	0.180	0.239	0.230	1.291	0.273	16.153

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	31	20	17	22	19	37	19	20
N.S.	1	1.15	0.74	0.63	0.81	0.70	1.37	0.70	0.74
time (sec)	N/A	0.151	0.006	0.164	0.277	0.238	0.125	0.271	0.038

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	20	19	18	18	26	18	20
N.S.	1	1.00	1.00	0.95	0.90	0.90	1.30	0.90	1.00
time (sec)	N/A	0.154	0.009	0.030	0.276	0.229	0.041	0.286	0.113

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	39	25	20	19	19	34	38	30
N.S.	1	0.93	0.60	0.48	0.45	0.45	0.81	0.90	0.71
time (sec)	N/A	0.212	0.017	0.018	0.199	0.260	0.413	0.266	15.592

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	7	7	12	8	7	9	5	8	26
N.S.	1	1.00	1.71	1.14	1.00	1.29	0.71	1.14	3.71
time (sec)	N/A	0.135	0.010	0.122	0.187	0.251	0.020	0.272	15.059

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	27	20	19	19	19	20	25
N.S.	1	1.00	1.00	0.74	0.70	0.70	0.70	0.74	0.93
time (sec)	N/A	0.166	0.008	0.024	0.271	0.238	0.057	0.273	0.061

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	30	20	16	30	39	36	26	42
N.S.	1	1.50	1.00	0.80	1.50	1.95	1.80	1.30	2.10
time (sec)	N/A	0.174	0.028	0.034	0.277	0.241	0.155	0.277	0.150

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	10	9	9	10	9	9
N.S.	1	1.00	1.00	0.77	0.69	0.69	0.77	0.69	0.69
time (sec)	N/A	0.131	0.015	0.194	0.198	0.224	0.059	0.278	0.047

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	11	13	9	8	13	9
N.S.	1	1.00	1.00	0.85	1.00	0.69	0.62	1.00	0.69
time (sec)	N/A	0.145	0.001	0.013	0.192	0.247	0.037	0.277	13.952

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	3	8	7	5	5
N.S.	1	1.00	1.00	1.33	1.00	2.67	2.33	1.67	1.67
time (sec)	N/A	0.147	0.002	0.089	0.187	0.235	0.168	0.271	0.026

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	14	12	11	11	10	11	11
N.S.	1	1.00	0.64	0.55	0.50	0.50	0.45	0.50	0.50
time (sec)	N/A	0.172	0.002	0.020	0.204	0.233	0.029	0.269	0.002

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	10	9	7	14	17	8	8
N.S.	1	1.00	1.00	0.90	0.70	1.40	1.70	0.80	0.80
time (sec)	N/A	0.136	0.021	0.190	0.278	0.239	0.482	0.266	0.098

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	17	13	12	15	11	10	15	11
N.S.	1	1.31	1.00	0.92	1.15	0.85	0.77	1.15	0.85
time (sec)	N/A	0.133	0.005	0.163	0.185	0.250	0.035	0.264	13.303

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	21	18	14	22	0	22	17
N.S.	1	1.00	1.00	0.86	0.67	1.05	0.00	1.05	0.81
time (sec)	N/A	0.155	0.005	0.014	0.181	0.244	0.000	0.275	0.072

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	34	34	23	22	21	32	21	22
N.S.	1	1.00	1.00	0.68	0.65	0.62	0.94	0.62	0.65
time (sec)	N/A	0.172	0.061	0.021	0.268	0.239	0.239	0.263	0.034

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	22	17	16	16	20	16	16
N.S.	1	1.00	1.00	0.77	0.73	0.73	0.91	0.73	0.73
time (sec)	N/A	0.218	0.029	0.030	0.194	0.239	0.096	0.277	13.609

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	16	15	21	12	15	17
N.S.	1	1.00	1.00	0.84	0.79	1.11	0.63	0.79	0.89
time (sec)	N/A	0.138	0.012	0.023	0.277	0.235	0.038	0.284	13.506

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	10	10	11	9	8	9	9
N.S.	1	1.00	1.00	1.00	1.10	0.90	0.80	0.90	0.90
time (sec)	N/A	0.154	0.034	0.041	0.233	0.236	0.035	0.275	0.064

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	40	7	6	16	5	25	6
N.S.	1	1.00	5.00	0.88	0.75	2.00	0.62	3.12	0.75
time (sec)	N/A	0.150	0.047	0.278	0.285	0.246	0.455	0.274	0.013

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	19	12	23	0	10	12
N.S.	1	1.00	1.00	1.58	1.00	1.92	0.00	0.83	1.00
time (sec)	N/A	0.150	0.017	0.035	0.288	0.238	0.000	0.286	0.049

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	40	17	0	24	0	22	16
N.S.	1	1.00	3.33	1.42	0.00	2.00	0.00	1.83	1.33
time (sec)	N/A	0.154	0.015	0.194	0.000	0.242	0.000	0.282	13.450

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	15	15	14	15	14	14
N.S.	1	1.00	1.00	0.83	0.83	0.78	0.83	0.78	0.78
time (sec)	N/A	0.143	0.016	0.150	0.186	0.231	0.055	0.275	0.062

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	52	52	38	35	26	43	0	0	0
N.S.	1	1.00	0.73	0.67	0.50	0.83	0.00	0.00	0.00
time (sec)	N/A	0.218	0.068	0.166	0.279	0.257	0.000	0.000	0.000

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	C	A	F	A	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	39	39	37	0	137	33	0	33	45
N.S.	1	1.00	0.95	0.00	3.51	0.85	0.00	0.85	1.15
time (sec)	N/A	0.378	0.697	0.000	0.335	0.239	0.000	0.279	14.609

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	22	20	19	21	26	19	19
N.S.	1	1.00	0.81	0.74	0.70	0.78	0.96	0.70	0.70
time (sec)	N/A	0.163	0.051	0.068	0.199	0.250	0.099	0.285	0.036

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	A	A	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	7	7	7	0	7	13	0	0	7
N.S.	1	1.00	1.00	0.00	1.00	1.86	0.00	0.00	1.00
time (sec)	N/A	0.244	0.037	0.000	0.229	0.247	0.000	0.000	17.652

Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	6	6	5	5	5	5	5
N.S.	1	1.00	1.00	1.00	0.83	0.83	0.83	0.83	0.83
time (sec)	N/A	0.157	0.016	0.026	0.200	0.229	0.029	0.268	0.032

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	7	6	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.88	0.75	0.75
time (sec)	N/A	0.151	0.029	0.042	0.192	0.239	0.035	0.266	13.962

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	12	8	7	10	16	10	6	6
N.S.	1	1.50	1.00	0.88	1.25	2.00	1.25	0.75	0.75
time (sec)	N/A	0.172	0.002	0.059	0.197	0.256	0.249	0.273	13.518

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	15	14	14	17	14	14
N.S.	1	1.00	1.00	0.79	0.74	0.74	0.89	0.74	0.74
time (sec)	N/A	0.137	0.009	0.189	0.280	0.235	0.056	0.275	0.040

Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	B	B	B	A	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	0	5	16	52	34	5	45	5
N.S.	1	0.00	1.00	3.20	10.40	6.80	1.00	9.00	1.00
time (sec)	N/A	0.000	0.018	0.871	0.218	0.250	0.145	0.274	0.070

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	7	7	7	7	6	6	7	77	6
N.S.	1	1.00	1.00	1.00	0.86	0.86	1.00	11.00	0.86
time (sec)	N/A	0.189	0.108	0.259	0.198	0.244	0.062	0.275	14.079

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	4	4	4	5	4	44	5	4	4
N.S.	1	1.00	1.00	1.25	1.00	11.00	1.25	1.00	1.00
time (sec)	N/A	0.225	9.252	8.366	0.189	0.257	0.901	0.281	14.974

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	23	11	15	18	8	10	10
N.S.	1	1.00	2.30	1.10	1.50	1.80	0.80	1.00	1.00
time (sec)	N/A	0.153	0.007	0.035	0.192	0.247	0.178	0.270	0.002

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	6	7	42	6	0	6	6
N.S.	1	1.00	1.00	1.17	7.00	1.00	0.00	1.00	1.00
time (sec)	N/A	0.181	0.004	0.174	0.194	0.240	0.000	0.277	15.081

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	6	6	5	5	5	5	5
N.S.	1	1.00	1.00	1.00	0.83	0.83	0.83	0.83	0.83
time (sec)	N/A	0.141	0.008	0.050	0.226	0.241	0.072	0.269	15.995

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	8	7	12	16	3	12	6
N.S.	1	1.00	1.33	1.17	2.00	2.67	0.50	2.00	1.00
time (sec)	N/A	0.152	0.009	0.021	0.199	0.238	0.057	0.263	16.601

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	7	16	12	8	36	9	8	21	26
N.S.	1	2.29	1.71	1.14	5.14	1.29	1.14	3.00	3.71
time (sec)	N/A	0.267	0.012	0.356	0.192	0.246	1.449	0.273	16.107

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F(-1)	F	A	F(-1)	A	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	10	0	0	6	0	6	6
N.S.	1	1.00	1.00	0.00	0.00	0.60	0.00	0.60	0.60
time (sec)	N/A	0.154	0.034	180.000	0.000	0.239	0.000	0.263	0.250

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	3	3	219	3	3
N.S.	1	1.00	1.00	1.33	1.00	1.00	73.00	1.00	1.00
time (sec)	N/A	0.173	0.013	0.240	0.274	0.251	9.125	0.266	16.484

Problem 190	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	11	10	9	22	8	9	9
N.S.	1	1.00	1.00	0.91	0.82	2.00	0.73	0.82	0.82
time (sec)	N/A	0.122	0.004	0.230	0.201	0.242	0.359	0.271	16.452

Problem 191	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	20	7	0	18	0	15	0
N.S.	1	1.00	1.00	0.35	0.00	0.90	0.00	0.75	0.00
time (sec)	N/A	0.142	0.113	0.251	0.000	0.235	0.000	0.286	0.000

Problem 192	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	7	7	12	8	11	11	7	12	12
N.S.	1	1.00	1.71	1.14	1.57	1.57	1.00	1.71	1.71
time (sec)	N/A	0.226	0.078	0.232	0.216	0.240	0.053	0.256	16.321

Problem 193	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	9	4	17	19	14	16	3
N.S.	1	1.00	3.00	1.33	5.67	6.33	4.67	5.33	1.00
time (sec)	N/A	0.149	0.003	0.122	0.192	0.252	0.051	0.265	0.047

Problem 194	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	3	3	5	3	5	5	5	5	5
N.S.	1	1.00	1.67	1.00	1.67	1.67	1.67	1.67	1.67
time (sec)	N/A	0.345	0.039	0.030	0.447	0.250	1.385	0.267	15.884

Problem 195	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	5	5	6	5	5	5	5	5
N.S.	1	1.00	1.00	1.20	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.189	0.061	0.288	0.220	0.244	0.050	0.269	16.576

Problem 196	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	B	B	B	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	15	15	12	11	27	0	11	5
N.S.	1	2.50	2.50	2.00	1.83	4.50	0.00	1.83	0.83
time (sec)	N/A	0.156	0.018	0.028	0.183	0.245	0.000	0.276	16.607

Problem 197	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	2	2	19	3	13	13	12	15	2
N.S.	1	1.00	9.50	1.50	6.50	6.50	6.00	7.50	1.00
time (sec)	N/A	0.120	0.004	0.186	0.195	0.231	0.048	0.275	16.288

Problem 198	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	13	12	13	24	14	10	14	13
N.S.	1	1.08	1.00	1.08	2.00	1.17	0.83	1.17	1.08
time (sec)	N/A	0.145	0.008	0.046	0.193	0.247	0.092	0.267	16.268

Problem 199	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	F(-1)	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	39	30	29	0	60	29	29
N.S.	1	1.00	1.00	0.77	0.74	0.00	1.54	0.74	0.74
time (sec)	N/A	0.188	0.067	1.460	0.209	0.000	0.246	0.268	17.420

Problem 200	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	4	4	4	5	4	41	5	4	4
N.S.	1	1.00	1.00	1.25	1.00	10.25	1.25	1.00	1.00
time (sec)	N/A	0.220	0.011	10.368	0.195	0.245	0.853	0.278	0.336

Problem 201	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	24	18	29	17	22	0	17
N.S.	1	1.00	1.00	0.75	1.21	0.71	0.92	0.00	0.71
time (sec)	N/A	0.177	0.010	0.042	0.231	0.250	0.394	0.000	15.657

Problem 202	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	22	17	16	16	20	16	16
N.S.	1	1.00	1.00	0.77	0.73	0.73	0.91	0.73	0.73
time (sec)	N/A	0.213	0.007	0.024	0.200	0.246	0.101	0.277	0.003

Problem 203	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	13	12	12	14	12	12
N.S.	1	1.00	1.00	0.81	0.75	0.75	0.88	0.75	0.75
time (sec)	N/A	0.132	0.005	0.188	0.285	0.240	0.066	0.278	15.914

Problem 204	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	30	23	22	25	114	22	22
N.S.	1	1.00	1.00	0.77	0.73	0.83	3.80	0.73	0.73
time (sec)	N/A	0.206	0.005	0.850	0.186	0.270	0.886	0.274	16.175

Problem 205	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	34	34	185	30	0	29	0	0
N.S.	1	1.48	1.48	8.04	1.30	0.00	1.26	0.00	0.00
time (sec)	N/A	0.151	0.009	0.055	0.284	0.000	0.379	0.000	0.000

Problem 206	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	3	9	4	4	3	4	4
N.S.	1	1.00	1.00	3.00	1.33	1.33	1.00	1.33	1.33
time (sec)	N/A	0.116	0.000	0.012	0.188	0.237	0.016	0.264	0.013

Problem 207	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	B	C	B	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	139	131	16	2527	123	0	134	15
N.S.	1	12.64	11.91	1.45	229.73	11.18	0.00	12.18	1.36
time (sec)	N/A	2.406	0.423	0.640	0.367	0.339	0.000	0.323	16.884

Problem 208	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	11	9	34	7	8	33	8
N.S.	1	1.00	1.38	1.12	4.25	0.88	1.00	4.12	1.00
time (sec)	N/A	0.154	0.015	0.231	0.193	0.242	0.144	0.274	15.331

Problem 209	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	29	27	33	21	22	44	46
N.S.	1	1.00	1.21	1.12	1.38	0.88	0.92	1.83	1.92
time (sec)	N/A	0.204	0.102	0.190	0.278	0.262	0.407	0.292	0.384

Problem 210	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	9	8	8	10	8	8
N.S.	1	1.00	1.00	0.75	0.67	0.67	0.83	0.67	0.67
time (sec)	N/A	0.141	0.016	0.166	0.272	0.240	0.065	0.279	0.258

Problem 211	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	B	A	B	B	F	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	0	5	12	5	11	7	0	5
N.S.	1	0.00	1.00	2.40	1.00	2.20	1.40	0.00	1.00
time (sec)	N/A	0.000	0.019	0.032	0.243	0.236	0.068	0.000	15.455

Problem 212	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	19	11	12	13	11	8	13	11
N.S.	1	1.73	1.00	1.09	1.18	1.00	0.73	1.18	1.00
time (sec)	N/A	0.160	0.007	0.170	0.176	0.243	0.039	0.267	15.515

Problem 213	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	2	2	16	3	2	14	2	25	2
N.S.	1	1.00	8.00	1.50	1.00	7.00	1.00	12.50	1.00
time (sec)	N/A	0.123	0.003	0.175	0.275	0.235	0.068	0.268	0.032

Problem 214	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	9	18	9	10	18	9	10	10	9
N.S.	1	2.00	1.00	1.11	2.00	1.00	1.11	1.11	1.00
time (sec)	N/A	0.200	0.014	0.052	0.184	0.252	0.048	0.273	15.230

Problem 215	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	13	10	10	9	9	7	9	9
N.S.	1	1.30	1.00	1.00	0.90	0.90	0.70	0.90	0.90
time (sec)	N/A	0.145	0.003	0.033	0.190	0.257	0.035	0.294	0.049

Problem 216	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	20	17	16	16	17	16	13
N.S.	1	1.00	1.00	0.85	0.80	0.80	0.85	0.80	0.65
time (sec)	N/A	0.178	0.011	0.039	0.190	0.249	0.118	0.277	14.933

Problem 217	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	22	22	20	15	107	20
N.S.	1	1.00	1.00	0.88	0.88	0.80	0.60	4.28	0.80
time (sec)	N/A	0.158	0.008	0.084	0.188	0.246	0.057	0.280	0.171

Problem 218	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	5	6	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.62	0.75	0.75
time (sec)	N/A	0.143	0.008	0.252	0.271	0.240	0.047	0.293	0.085

Problem 219	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	33	20	13	5	12	17	19	12
N.S.	1	1.65	1.00	0.65	0.25	0.60	0.85	0.95	0.60
time (sec)	N/A	0.261	0.008	0.098	0.333	0.244	0.512	0.289	15.403

Problem 220	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	16	25	20	8	19	10	19	16
N.S.	1	1.33	2.08	1.67	0.67	1.58	0.83	1.58	1.33
time (sec)	N/A	0.288	0.017	0.345	0.185	0.246	1.043	0.276	15.260

Problem 221	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	17	16	17	16	16	15	16	16
N.S.	1	1.06	1.00	1.06	1.00	1.00	0.94	1.00	1.00
time (sec)	N/A	0.157	0.004	0.051	0.276	0.245	0.050	0.269	0.092

Problem 222	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	10	13	13	12	13	9
N.S.	1	1.00	1.00	0.67	0.87	0.87	0.80	0.87	0.60
time (sec)	N/A	0.138	0.004	0.259	0.187	0.239	0.044	0.275	14.935

Problem 223	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	28	23	23	22	19	37	40	0
N.S.	1	1.00	0.82	0.82	0.79	0.68	1.32	1.43	0.00
time (sec)	N/A	0.167	0.011	0.013	0.271	0.256	0.776	0.263	0.000

Problem 224	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	12	13	21	13	47	17	13	24
N.S.	1	0.92	1.00	1.62	1.00	3.62	1.31	1.00	1.85
time (sec)	N/A	0.238	3.436	12.919	0.186	0.251	0.844	0.283	15.867

Problem 225	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	9	9	11	9	9	9	8	11	11
N.S.	1	1.00	1.22	1.00	1.00	1.00	0.89	1.22	1.22
time (sec)	N/A	0.170	1.895	0.040	0.400	0.249	0.495	0.267	15.397

Problem 226	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	14	14	16	12	1038	14	14	14	14
N.S.	1	1.00	1.14	0.86	74.14	1.00	1.00	1.00	1.00
time (sec)	N/A	0.190	9.778	0.105	0.736	0.247	0.800	0.278	15.380

Problem 227	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	F(-2)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	11128	10106	10106	0	11171	10106	15
N.S.	1	1.00	483.83	439.39	439.39	0.00	485.70	439.39	0.65
time (sec)	N/A	0.982	0.137	9.328	2.400	0.000	2.596	1.104	31.766

Problem 228	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	21	19	10	9	19	29	19	15
N.S.	1	1.91	1.73	0.91	0.82	1.73	2.64	1.73	1.36
time (sec)	N/A	0.225	0.020	3.911	0.187	0.257	0.252	0.261	15.045

Problem 229	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	5	5	6	5	5	3	5	5
N.S.	1	1.00	1.00	1.20	1.00	1.00	0.60	1.00	1.00
time (sec)	N/A	0.255	0.018	0.034	0.224	0.251	0.066	0.268	15.455

Problem 230	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	37	18	17	31	15	17	17
N.S.	1	1.00	1.61	0.78	0.74	1.35	0.65	0.74	0.74
time (sec)	N/A	0.133	0.006	0.236	0.277	0.237	0.083	0.293	0.029

Problem 231	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	28	28	22	21	21	20	35	21
N.S.	1	1.00	1.00	0.79	0.75	0.75	0.71	1.25	0.75
time (sec)	N/A	0.199	0.042	0.036	0.212	0.248	0.523	0.313	16.329

Problem 232	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	5	5	6	5	5	5	72	5
N.S.	1	1.00	1.00	1.20	1.00	1.00	1.00	14.40	1.00
time (sec)	N/A	0.171	0.066	0.188	0.201	0.250	0.073	0.276	0.093

Problem 233	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	57	57	57	12	9	5	0	5	0
N.S.	1	1.00	1.00	0.21	0.16	0.09	0.00	0.09	0.00
time (sec)	N/A	0.197	0.009	0.065	0.415	0.237	0.000	0.277	0.000

Problem 234	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	29	22	11	10	29	42	10	10
N.S.	1	2.07	1.57	0.79	0.71	2.07	3.00	0.71	0.71
time (sec)	N/A	0.186	0.006	0.075	0.193	0.243	0.266	0.277	0.091

Problem 235	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	45	55	34	30	0	33	0	33	51
N.S.	1	1.22	0.76	0.67	0.00	0.73	0.00	0.73	1.13
time (sec)	N/A	0.192	0.313	0.040	0.000	0.235	0.000	0.284	14.998

Problem 236	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	5	5	7	5	7	7	7	7	7
N.S.	1	1.00	1.40	1.00	1.40	1.40	1.40	1.40	1.40
time (sec)	N/A	0.164	1.613	0.065	0.410	0.249	0.591	0.273	15.895

Problem 237	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	18	20	17	16	16	15	16	16
N.S.	1	0.90	1.00	0.85	0.80	0.80	0.75	0.80	0.80
time (sec)	N/A	0.224	0.002	0.132	0.194	0.245	0.114	0.275	0.119

Problem 238	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	23	7	17	17	15	18	6
N.S.	1	1.00	2.88	0.88	2.12	2.12	1.88	2.25	0.75
time (sec)	N/A	0.130	0.004	0.204	0.187	0.254	0.046	0.279	0.004

Problem 239	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	10	20	14	10	2
N.S.	1	1.00	1.00	1.50	5.00	10.00	7.00	5.00	1.00
time (sec)	N/A	0.158	0.002	0.233	0.204	0.246	0.308	0.287	0.047

Problem 240	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	9	9	9	8	0	7	7	0	7
N.S.	1	1.00	1.00	0.89	0.00	0.78	0.78	0.00	0.78
time (sec)	N/A	0.165	0.014	0.042	0.000	0.262	0.070	0.000	0.068

Problem 241	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	37	26	24	23	21	306	30	10
N.S.	1	1.06	0.74	0.69	0.66	0.60	8.74	0.86	0.29
time (sec)	N/A	0.159	0.025	0.205	0.206	0.247	0.562	0.275	16.078

Problem 242	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	194	20	21	20	20	19	21	20
N.S.	1	8.08	0.83	0.88	0.83	0.83	0.79	0.88	0.83
time (sec)	N/A	0.637	0.007	0.026	0.264	0.239	0.051	0.275	16.006

Problem 243	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	A	A	B	F	A	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	0	5	6	5	65	0	5	5
N.S.	1	0.00	1.00	1.20	1.00	13.00	0.00	1.00	1.00
time (sec)	N/A	0.000	0.126	12.754	0.384	0.245	0.000	0.275	16.485

Problem 244	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	C	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	33	12	40	79	14	80	55	14
N.S.	1	2.75	1.00	3.33	6.58	1.17	6.67	4.58	1.17
time (sec)	N/A	0.227	0.031	6.327	0.207	0.250	0.138	0.270	16.361

Problem 245	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	A	A	A	A	A	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	0	18	12	16	11	10	11	11
N.S.	1	0.00	1.80	1.20	1.60	1.10	1.00	1.10	1.10
time (sec)	N/A	0.000	0.011	0.222	0.190	0.240	0.189	0.263	16.435

Problem 246	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	C	A	F	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	42	3	4	14	3	0	40	3
N.S.	1	14.00	1.00	1.33	4.67	1.00	0.00	13.33	1.00
time (sec)	N/A	0.243	0.002	0.605	0.284	0.225	0.000	0.287	15.736

Problem 247	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	40	7	6	16	5	25	6
N.S.	1	1.00	5.00	0.88	0.75	2.00	0.62	3.12	0.75
time (sec)	N/A	0.135	0.013	0.240	0.282	0.231	0.196	0.264	14.980

Problem 248	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	16	15	14	13	31	63	46	26
N.S.	1	0.47	0.44	0.41	0.38	0.91	1.85	1.35	0.76
time (sec)	N/A	0.171	0.012	0.106	0.270	0.248	0.327	0.270	14.835

Problem 249	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	16	18	19	14	20	18
N.S.	1	1.00	1.00	0.89	1.00	1.06	0.78	1.11	1.00
time (sec)	N/A	0.151	0.007	0.047	0.199	0.232	0.059	0.257	0.052

Problem 250	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	F	A	A	F	F(-1)	F(-1)
verified	N/A	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	56	0	56	0	51	69	0	0	0
N.S.	1	0.00	1.00	0.00	0.91	1.23	0.00	0.00	0.00
time (sec)	N/A	0.000	0.037	0.000	0.401	0.244	0.000	0.000	0.000

Problem 251	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	20	17	16	16	15	16	17
N.S.	1	1.00	1.00	0.85	0.80	0.80	0.75	0.80	0.85
time (sec)	N/A	0.191	0.003	0.010	0.193	0.222	0.015	0.273	0.035

Problem 252	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	13	12	32	11	65	0	11
N.S.	1	1.00	0.62	0.57	1.52	0.52	3.10	0.00	0.52
time (sec)	N/A	0.193	10.029	0.256	0.275	0.235	1.445	0.000	14.706

Problem 253	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	25	33	24	25	39	29	18	38
N.S.	1	1.47	1.94	1.41	1.47	2.29	1.71	1.06	2.24
time (sec)	N/A	0.203	0.054	0.392	0.192	0.242	0.026	0.270	15.193

Problem 254	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	31	8	7	61	10	5	6	6
N.S.	1	3.88	1.00	0.88	7.62	1.25	0.62	0.75	0.75
time (sec)	N/A	0.378	0.041	0.072	0.193	0.241	0.162	0.267	14.735

Problem 255	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	23	14	39	42	19	37	14
N.S.	1	1.00	1.35	0.82	2.29	2.47	1.12	2.18	0.82
time (sec)	N/A	0.192	0.015	9.814	0.197	0.239	0.170	0.273	14.884

Problem 256	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	30	22	23	31	22	34	51	22
N.S.	1	0.91	0.67	0.70	0.94	0.67	1.03	1.55	0.67
time (sec)	N/A	0.198	0.023	0.094	0.306	0.243	0.190	0.259	15.857

Problem 257	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	C	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	217	6	44	0	15	0	35	35
N.S.	1	36.17	1.00	7.33	0.00	2.50	0.00	5.83	5.83
time (sec)	N/A	0.921	0.084	0.489	0.000	0.257	0.000	0.276	16.005

Problem 258	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	B	C	B	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	9	81	21	28	45	16	0	5161	0
N.S.	1	9.00	2.33	3.11	5.00	1.78	0.00	573.44	0.00
time (sec)	N/A	0.224	0.879	1.581	0.277	0.255	0.000	1.841	0.000

Problem 259	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	67	78	56	0	0	0	0	0	0
N.S.	1	1.16	0.84	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.320	0.056	0.000	0.000	0.000	0.000	0.000	0.000

Problem 260	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	84	84	84	63	66	62	394	62	78
N.S.	1	1.00	1.00	0.75	0.79	0.74	4.69	0.74	0.93
time (sec)	N/A	0.377	0.103	1.304	0.236	0.264	4.266	0.261	16.535

Problem 261	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	20	21	58	16	19	23	32
N.S.	1	1.00	1.00	1.05	2.90	0.80	0.95	1.15	1.60
time (sec)	N/A	0.200	0.110	0.376	0.289	0.271	0.063	0.273	16.873

Problem 262	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	63	89	53	62	83	119	50	0
N.S.	1	1.00	1.41	0.84	0.98	1.32	1.89	0.79	0.00
time (sec)	N/A	0.185	0.184	0.237	0.273	0.252	1.065	0.267	0.000

Problem 263	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	21	26	14	17	20	17	14
N.S.	1	1.00	1.00	1.24	0.67	0.81	0.95	0.81	0.67
time (sec)	N/A	0.164	0.013	0.188	0.215	0.274	0.346	0.261	16.239

Problem 264	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	41	40	103	39	36	39	39
N.S.	1	1.00	0.54	0.53	1.36	0.51	0.47	0.51	0.51
time (sec)	N/A	0.411	0.624	0.079	0.198	0.244	0.049	0.259	16.514

Problem 265	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	F	A	A	A	B	F(-1)
verified	N/A	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	38	0	38	0	34	34	34	163	0
N.S.	1	0.00	1.00	0.00	0.89	0.89	0.89	4.29	0.00
time (sec)	N/A	0.000	0.027	0.000	0.296	0.246	0.099	0.283	0.000

Problem 266	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	52	115	0	0	0	0	0	0	0
N.S.	1	2.21	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.259	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 267	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	64	64	33	29	46	29	138	61	46
N.S.	1	1.00	0.52	0.45	0.72	0.45	2.16	0.95	0.72
time (sec)	N/A	0.177	0.027	0.212	0.204	0.236	1.700	0.270	0.034

Problem 268	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	77	70	47	0	0	56	0	0
N.S.	1	1.79	1.63	1.09	0.00	0.00	1.30	0.00	0.00
time (sec)	N/A	0.202	0.037	0.249	0.000	0.000	0.730	0.000	0.000

Problem 269	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	46	58	31	30	48	34	30	31
N.S.	1	1.12	1.41	0.76	0.73	1.17	0.83	0.73	0.76
time (sec)	N/A	0.162	0.137	0.194	0.306	0.232	0.106	0.268	15.214

Problem 270	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	99	99	142	0	0	0	0	0	0
N.S.	1	1.00	1.43	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.221	0.183	0.000	0.000	0.000	0.000	0.000	0.000

Problem 271	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	36	31	29	36	0	0	0
N.S.	1	1.00	1.00	0.86	0.81	1.00	0.00	0.00	0.00
time (sec)	N/A	0.253	0.182	0.043	0.234	0.244	0.000	0.000	0.000

Problem 272	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	104	47	42	41	45	85	41	55
N.S.	1	2.36	1.07	0.95	0.93	1.02	1.93	0.93	1.25
time (sec)	N/A	0.762	0.048	0.162	0.203	0.242	1.033	0.260	16.529

Problem 273	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	56	64	33	30	58	29	68	42	30
N.S.	1	1.14	0.59	0.54	1.04	0.52	1.21	0.75	0.54
time (sec)	N/A	0.279	0.039	0.239	0.193	0.231	0.847	0.273	0.224

Problem 274	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	36	25	23	28	37	26	23	28
N.S.	1	1.16	0.81	0.74	0.90	1.19	0.84	0.74	0.90
time (sec)	N/A	0.145	0.015	0.201	0.291	0.233	0.068	0.264	0.064

Problem 275	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	A	A	B	F	F	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	108	0	132	96	115	262	0	0	0
N.S.	1	0.00	1.22	0.89	1.06	2.43	0.00	0.00	0.00
time (sec)	N/A	0.000	0.050	0.644	0.337	0.272	0.000	0.000	0.000

Problem 276	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	C	A	A	F	A	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	32	83	29	17	8	8	0	32	8
N.S.	1	2.59	0.91	0.53	0.25	0.25	0.00	1.00	0.25
time (sec)	N/A	0.362	0.023	2.327	0.220	0.254	0.000	0.870	15.603

Problem 277	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	A	C	A	F	A	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	0	21	22	27	13	0	13	19
N.S.	1	0.00	1.00	1.05	1.29	0.62	0.00	0.62	0.90
time (sec)	N/A	0.000	0.057	0.114	0.241	0.239	0.000	0.256	0.035

Problem 278	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	106	114	13	117	114	31	0	0
N.S.	1	1.04	1.12	0.13	1.15	1.12	0.30	0.00	0.00
time (sec)	N/A	0.182	0.405	0.247	0.287	0.242	1.774	0.000	0.000

Problem 279	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	3	9	4	4	3	4	4
N.S.	1	1.00	1.00	3.00	1.33	1.33	1.00	1.33	1.33
time (sec)	N/A	0.118	0.000	0.017	0.200	0.227	0.013	0.248	0.004

Problem 280	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	3	8	7	5	5
N.S.	1	1.00	1.00	1.33	1.00	2.67	2.33	1.67	1.67
time (sec)	N/A	0.153	0.000	0.073	0.198	0.237	0.175	0.259	0.003

Problem 281	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	7	7	7	7	6	6	7	6	6
N.S.	1	1.00	1.00	1.00	0.86	0.86	1.00	0.86	0.86
time (sec)	N/A	0.240	0.022	0.054	0.198	0.243	0.068	0.269	0.182

Problem 282	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	30	23	22	22	20	22	22
N.S.	1	1.00	1.00	0.77	0.73	0.73	0.67	0.73	0.73
time (sec)	N/A	0.216	0.001	0.075	0.193	0.226	0.017	0.265	0.022

Problem 283	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	36	27	26	26	27	23	26
N.S.	1	1.00	1.00	0.75	0.72	0.72	0.75	0.64	0.72
time (sec)	N/A	0.218	0.003	0.015	0.205	0.221	0.017	0.263	0.024

Problem 284	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	20	20	17	16	16	15	16	8
N.S.	1	2.00	2.00	1.70	1.60	1.60	1.50	1.60	0.80
time (sec)	N/A	0.156	0.007	0.203	0.205	0.245	0.072	0.269	15.369

Problem 285	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	15	14	24	17	28	14
N.S.	1	1.00	1.00	1.25	1.17	2.00	1.42	2.33	1.17
time (sec)	N/A	0.168	0.261	1.191	0.289	0.252	0.030	0.292	16.002

Problem 286	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	8	8	8	8	7	7	7	7	7
N.S.	1	1.00	1.00	1.00	0.88	0.88	0.88	0.88	0.88
time (sec)	N/A	0.149	0.006	0.050	0.206	0.244	0.057	0.269	18.451

Problem 287	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	A	A	A	A	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	1	3	1	2	1	1	0	26	1
N.S.	1	3.00	1.00	2.00	1.00	1.00	0.00	26.00	1.00
time (sec)	N/A	0.158	0.000	0.250	0.225	0.231	0.020	0.267	0.005

Problem 288	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	1	50	1	2	1	1	58	1	1
N.S.	1	50.00	1.00	2.00	1.00	1.00	58.00	1.00	1.00
time (sec)	N/A	0.214	0.001	0.334	0.211	0.231	0.029	0.273	18.137

Problem 289	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	9	9	9	9	81	8	8	8	9
N.S.	1	1.00	1.00	1.00	9.00	0.89	0.89	0.89	1.00
time (sec)	N/A	0.158	0.033	0.383	0.232	0.241	0.166	0.267	18.143

Problem 290	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	49	49	100	61	79	114	0	73	238
N.S.	1	1.00	2.04	1.24	1.61	2.33	0.00	1.49	4.86
time (sec)	N/A	0.176	0.080	0.346	0.282	0.252	0.000	0.292	17.803

Problem 291	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F(-1)	A	F(-1)	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	13	15	13	0	11	0	0	15	0
N.S.	1	1.15	1.00	0.00	0.85	0.00	0.00	1.15	0.00
time (sec)	N/A	1.729	3.386	180.000	0.188	0.000	0.000	0.273	0.000

Problem 292	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	9	17	17	14	13	131	34	13	13
N.S.	1	1.89	1.89	1.56	1.44	14.56	3.78	1.44	1.44
time (sec)	N/A	0.149	0.103	2.044	0.197	1.626	0.126	0.287	17.314

Problem 293	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	C	B	A	A	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	0	21	30	91	16	17	119	22
N.S.	1	0.00	1.11	1.58	4.79	0.84	0.89	6.26	1.16
time (sec)	N/A	0.000	0.070	0.331	0.208	0.257	0.077	0.298	17.617

Problem 294	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	F	F	F	B	B	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	9	74	0	0	0	30	56	0	0
N.S.	1	8.22	0.00	0.00	0.00	3.33	6.22	0.00	0.00
time (sec)	N/A	0.371	0.000	0.000	0.000	0.252	18.125	0.000	0.000

Problem 295	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	32	24	24	24	26	24	24
N.S.	1	1.00	1.00	0.75	0.75	0.75	0.81	0.75	0.75
time (sec)	N/A	0.148	0.000	0.013	0.197	0.248	0.017	0.271	0.026

Problem 296	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	14	15	14	14	14	16	10
N.S.	1	1.00	1.00	1.07	1.00	1.00	1.00	1.14	0.71
time (sec)	N/A	0.144	0.007	0.209	0.200	0.239	0.041	0.269	0.047

Problem 297	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	47	34	28	80	65	0	0	0
N.S.	1	1.21	0.87	0.72	2.05	1.67	0.00	0.00	0.00
time (sec)	N/A	0.280	0.017	0.047	0.203	0.250	0.000	0.000	0.000

Problem 298	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	A	A	A	A	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	0	24	33	28	32	27	72	28
N.S.	1	0.00	1.00	1.38	1.17	1.33	1.12	3.00	1.17
time (sec)	N/A	0.000	0.047	0.121	0.247	0.237	0.051	0.276	18.187

Problem 299	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	23	20	19	0	34	0	22	17
N.S.	1	0.88	0.77	0.73	0.00	1.31	0.00	0.85	0.65
time (sec)	N/A	0.189	0.013	0.265	0.000	0.238	0.000	0.279	0.063

Problem 300	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	49	42	33	32	35	83	32	38
N.S.	1	1.11	0.95	0.75	0.73	0.80	1.89	0.73	0.86
time (sec)	N/A	0.237	0.019	0.369	0.206	0.243	0.178	0.269	16.927

Problem 301	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	17	11	12	54	17	15	16	11
N.S.	1	1.55	1.00	1.09	4.91	1.55	1.36	1.45	1.00
time (sec)	N/A	0.269	0.016	0.175	0.281	0.255	92.250	0.271	16.288

Problem 302	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	C	F(-2)	B	C	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	13	13	13	0	41	0	83	41	9
N.S.	1	1.00	1.00	0.00	3.15	0.00	6.38	3.15	0.69
time (sec)	N/A	0.170	0.019	0.000	0.286	0.000	0.243	0.272	16.432

Problem 303	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	13	94	22	83	12	35
N.S.	1	1.00	1.00	1.08	7.83	1.83	6.92	1.00	2.92
time (sec)	N/A	0.201	0.020	0.451	0.289	0.248	1.468	0.272	16.939

Problem 304	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	86	19	85	82	85	85
N.S.	1	1.00	1.00	4.53	1.00	4.47	4.32	4.47	4.47
time (sec)	N/A	0.233	0.051	0.264	0.195	0.233	0.466	0.289	17.036

Problem 305	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	A	F(-1)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	86	190	86	67	0	71	100	66	66
N.S.	1	2.21	1.00	0.78	0.00	0.83	1.16	0.77	0.77
time (sec)	N/A	1.243	0.028	0.306	0.000	0.362	69.501	0.272	0.197

Problem 306	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	26	23	22	22	19	22	22
N.S.	1	1.00	1.00	0.88	0.85	0.85	0.73	0.85	0.85
time (sec)	N/A	0.172	0.027	0.215	0.276	0.242	0.030	0.276	0.051

Problem 307	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	F	F	A	F	A	F(-1)
verified	N/A	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	23	0	25	0	0	12	0	12	0
N.S.	1	0.00	1.09	0.00	0.00	0.52	0.00	0.52	0.00
time (sec)	N/A	0.000	0.063	0.000	0.000	0.247	0.000	0.273	0.000

Problem 308	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	11	10	10	8	10	10
N.S.	1	1.00	1.00	0.92	0.83	0.83	0.67	0.83	0.83
time (sec)	N/A	0.132	0.056	0.219	0.199	0.232	0.232	0.264	0.104

Problem 309	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	C	A	A	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	22	34	46	15	27	42	0
N.S.	1	1.00	0.96	1.48	2.00	0.65	1.17	1.83	0.00
time (sec)	N/A	0.175	0.041	0.178	0.301	0.246	1.344	0.295	0.000

Problem 310	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	9	9	24	8	12	17	7	8	7
N.S.	1	1.00	2.67	0.89	1.33	1.89	0.78	0.89	0.78
time (sec)	N/A	0.174	0.027	0.167	0.206	0.258	0.099	0.267	0.097

Problem 311	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	22	23	34	24	20	22	18
N.S.	1	1.00	1.00	1.05	1.55	1.09	0.91	1.00	0.82
time (sec)	N/A	0.241	0.012	0.061	0.194	0.237	0.051	0.276	0.044

Problem 312	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	22	39	34	40	17	31	38
N.S.	1	1.00	1.00	1.77	1.55	1.82	0.77	1.41	1.73
time (sec)	N/A	0.150	0.042	0.112	0.185	0.244	0.764	0.283	0.102

Problem 313	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	C	F	B	F	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	0	21	52	0	84	0	136	15
N.S.	1	0.00	1.00	2.48	0.00	4.00	0.00	6.48	0.71
time (sec)	N/A	0.000	0.089	2.167	0.000	0.261	0.000	0.277	17.185

Problem 314	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	18	17	17	15	21	17
N.S.	1	1.00	1.00	1.06	1.00	1.00	0.88	1.24	1.00
time (sec)	N/A	0.290	0.065	0.315	0.248	0.244	0.061	0.272	17.009

Problem 315	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F	F	A	B	F	F(-1)
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	45	37	107	0	0	37	4362	0	0
N.S.	1	0.82	2.38	0.00	0.00	0.82	96.93	0.00	0.00
time (sec)	N/A	0.282	0.535	0.000	0.000	0.302	3.878	0.000	0.000

Problem 316	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	54	54	54	45	44	44	49	44	44
N.S.	1	1.00	1.00	0.83	0.81	0.81	0.91	0.81	0.81
time (sec)	N/A	0.221	0.005	0.198	0.214	0.230	0.023	0.272	0.111

Problem 317	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	C	A	A	F	A	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	0	18	34	23	23	0	23	18
N.S.	1	0.00	1.00	1.89	1.28	1.28	0.00	1.28	1.00
time (sec)	N/A	0.000	0.014	0.336	0.292	0.234	0.000	0.277	16.642

Problem 318	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	A	B	F	A	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	61	65	77	0	56	108	0	57	0
N.S.	1	1.07	1.26	0.00	0.92	1.77	0.00	0.93	0.00
time (sec)	N/A	0.269	0.284	0.000	0.277	0.249	0.000	0.290	0.000

Problem 319	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	B	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	64	136	64	228	52	55	0	61	66
N.S.	1	2.12	1.00	3.56	0.81	0.86	0.00	0.95	1.03
time (sec)	N/A	2.076	0.478	0.737	0.206	0.284	0.000	37.520	19.383

Problem 320	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	F	F	A	F	F	F(-1)
verified	N/A	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	105	0	105	0	0	148	0	0	0
N.S.	1	0.00	1.00	0.00	0.00	1.41	0.00	0.00	0.00
time (sec)	N/A	0.000	0.549	0.000	0.000	0.648	0.000	0.000	0.000

Problem 321	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	29	25	18	17	17	15	19	23
N.S.	1	1.16	1.00	0.72	0.68	0.68	0.60	0.76	0.92
time (sec)	N/A	0.182	0.006	0.055	0.202	0.236	0.080	0.335	0.105

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. The column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [305] had the largest ratio of [3.42857000000000012]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	3	3	1.00	11	0.273
2	A	7	6	1.12	9	0.667
3	A	2	2	1.52	13	0.154
4	A	4	4	0.96	8	0.500
5	A	2	2	1.00	5	0.400
6	A	5	4	1.25	9	0.444
7	A	6	5	1.00	9	0.556
8	A	4	3	1.00	13	0.231
9	A	5	4	1.31	11	0.364
10	A	12	11	1.55	6	1.833
11	A	2	2	1.00	12	0.167
12	A	6	5	1.19	21	0.238
13	A	2	2	1.00	8	0.250
14	A	4	3	1.72	17	0.176
15	A	2	2	1.00	7	0.286
16	A	2	2	1.00	12	0.167
17	A	1	1	1.00	7	0.143
18	A	2	2	1.00	13	0.154
19	A	3	2	1.00	17	0.118
20	C	5	4	3.96	7	0.571
21	A	4	3	1.00	14	0.214
22	A	2	2	1.00	9	0.222

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
23	A	1	1	1.00	11	0.091
24	A	6	5	1.00	17	0.294
25	A	2	2	1.00	19	0.105
26	A	1	1	1.00	9	0.111
27	A	4	3	1.00	14	0.214
28	A	2	2	1.00	11	0.182
29	A	3	3	1.00	27	0.111
30	A	5	4	1.00	4	1.000
31	A	5	4	1.00	9	0.444
32	A	6	6	1.00	11	0.545
33	A	7	7	1.29	4	1.750
34	A	3	3	1.00	12	0.250
35	A	1	1	1.00	3	0.333
36	A	2	2	1.00	2	1.000
37	A	4	3	0.46	15	0.200
38	A	1	1	1.00	13	0.077
39	A	3	3	1.00	11	0.273
40	A	1	1	1.00	8	0.125
41	B	5	4	2.38	7	0.571
42	A	5	4	1.00	15	0.267
43	A	4	3	1.00	12	0.250
44	A	2	2	1.00	8	0.250
45	A	4	3	1.28	13	0.231
46	B	3	3	66.75	9	0.333
47	A	4	3	1.00	9	0.333
48	A	4	4	1.00	34	0.118
49	A	4	3	1.50	15	0.200
50	A	5	4	0.90	9	0.444
51	A	1	1	1.00	8	0.125
52	B	5	4	2.55	21	0.190
53	A	4	3	1.07	15	0.200
54	A	2	2	1.00	11	0.182
55	A	3	2	1.70	14	0.143
56	A	1	1	1.00	4	0.250

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
57	A	2	2	1.00	6	0.333
58	A	11	10	1.09	23	0.435
59	A	6	5	1.00	21	0.238
60	A	4	4	1.00	34	0.118
61	A	1	1	1.00	12	0.083
62	A	7	6	1.12	9	0.667
63	A	2	2	1.00	15	0.133
64	A	2	2	1.00	9	0.222
65	A	4	3	1.00	7	0.429
66	A	5	4	1.12	15	0.267
67	A	3	2	1.00	11	0.182
68	A	4	3	1.00	9	0.333
69	A	3	2	1.00	9	0.222
70	A	3	3	1.62	11	0.273
71	A	4	4	1.00	6	0.667
72	A	4	3	1.15	13	0.231
73	A	2	2	1.00	10	0.200
74	A	6	6	1.00	11	0.545
75	A	1	1	1.00	11	0.091
76	A	1	1	1.00	3	0.333
77	F	0	0	N/A	0.000	N/A
78	A	2	2	1.00	13	0.154
79	A	2	2	1.00	11	0.182
80	A	1	1	1.00	15	0.067
81	A	6	6	1.00	13	0.462
82	A	2	2	1.00	27	0.074
83	A	1	1	1.00	2	0.500
84	B	5	4	3.17	11	0.364
85	B	5	5	2.07	9	0.556
86	A	5	4	1.00	15	0.267
87	A	3	3	1.00	4	0.750
88	A	6	5	1.90	7	0.714
89	A	6	5	1.00	5	1.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
90	A	1	1	1.00	9	0.111
91	A	4	3	0.89	11	0.273
92	A	3	2	1.00	10	0.200
93	A	8	7	1.00	6	1.167
94	A	3	2	1.00	11	0.182
95	A	2	2	1.00	11	0.182
96	A	6	5	1.00	9	0.556
97	A	2	2	1.00	8	0.250
98	A	4	3	1.00	13	0.231
99	A	1	1	1.00	4	0.250
100	A	9	8	1.06	6	1.333
101	A	1	1	1.00	16	0.062
102	A	5	4	1.09	13	0.308
103	A	2	2	1.00	9	0.222
104	A	1	1	1.00	6	0.167
105	A	2	2	1.00	9	0.222
106	A	2	2	1.00	2	1.000
107	A	2	2	1.00	9	0.222
108	A	3	3	0.95	4	0.750
109	A	2	2	1.00	10	0.200
110	A	2	2	1.47	18	0.111
111	A	3	3	1.04	4	0.750
112	A	5	4	1.27	13	0.308
113	A	5	5	1.00	8	0.625
114	A	3	3	1.16	19	0.158
115	A	4	3	0.55	12	0.250
116	A	4	3	1.11	15	0.200
117	A	2	2	1.50	6	0.333
118	B	1	1	5.80	11	0.091
119	B	2	2	2.31	11	0.182
120	A	4	3	1.00	12	0.250
121	A	2	2	1.00	2	1.000
122	A	3	2	1.00	7	0.286
123	A	2	2	1.00	6	0.333

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
124	A	7	6	1.55	12	0.500
125	A	1	1	1.00	3	0.333
126	A	2	2	1.27	12	0.167
127	A	4	3	1.07	9	0.333
128	A	6	6	1.00	11	0.545
129	A	1	1	1.00	9	0.111
130	A	5	4	1.00	14	0.286
131	A	6	5	0.97	10	0.500
132	A	6	5	0.52	11	0.455
133	A	4	3	1.25	15	0.200
134	A	5	5	1.00	4	1.250
135	A	5	4	1.00	7	0.571
136	A	1	1	1.00	14	0.071
137	A	3	3	1.00	2	1.500
138	A	1	1	1.00	7	0.143
139	A	1	1	1.00	6	0.167
140	F	0	0	N/A	0.000	N/A
141	A	3	2	1.00	11	0.182
142	A	5	5	1.00	8	0.625
143	A	6	5	1.11	10	0.500
144	A	1	1	1.00	2	0.500
145	A	1	1	1.00	11	0.091
146	A	2	2	1.00	11	0.182
147	F	0	0	N/A	0.000	N/A
148	A	4	3	1.11	25	0.120
149	A	5	4	1.00	9	0.444
150	A	1	1	1.00	21	0.048
151	A	4	3	1.15	13	0.231
152	A	4	3	1.00	12	0.250
153	A	5	4	0.93	11	0.364
154	A	1	1	1.00	6	0.167
155	A	2	2	1.00	15	0.133
156	A	2	2	1.50	6	0.333

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
157	A	1	1	1.00	13	0.077
158	A	1	1	1.00	6	0.167
159	A	2	2	1.00	2	1.000
160	A	2	2	1.00	9	0.222
161	A	4	3	1.00	13	0.231
162	A	5	4	1.31	11	0.364
163	A	2	2	1.00	2	1.000
164	A	2	2	1.00	12	0.167
165	A	6	5	1.00	6	0.833
166	A	3	3	1.00	7	0.429
167	A	1	1	1.00	12	0.083
168	A	4	3	1.00	11	0.273
169	A	5	4	1.00	8	0.500
170	A	2	2	1.00	8	0.250
171	A	4	3	1.00	9	0.333
172	A	2	2	1.00	18	0.111
173	A	6	5	1.00	25	0.200
174	A	1	1	1.00	10	0.100
175	A	4	3	1.00	13	0.231
176	A	3	2	1.00	11	0.182
177	A	2	2	1.00	4	0.500
178	A	4	3	1.50	7	0.429
179	A	3	2	1.00	11	0.182
180	F	0	0	N/A	0.000	N/A
181	A	1	1	1.00	15	0.067
182	A	5	4	1.00	10	0.400
183	A	2	2	1.00	6	0.333
184	A	4	3	1.00	13	0.231
185	A	1	1	1.00	10	0.100
186	A	4	4	1.00	4	1.000
187	B	6	5	2.29	25	0.200
188	A	5	4	1.00	13	0.308
189	A	4	3	1.00	13	0.231

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
190	A	1	1	1.00	9	0.111
191	A	3	2	1.00	15	0.133
192	A	2	2	1.00	14	0.143
193	A	4	3	1.00	5	0.600
194	N/A	4	0	1.00	3	0.000
195	A	1	1	1.00	13	0.077
196	B	1	1	2.50	17	0.059
197	A	1	1	1.00	9	0.111
198	A	2	2	1.08	4	0.500
199	A	1	1	1.00	21	0.048
200	A	5	4	1.00	10	0.400
201	A	3	2	1.00	13	0.154
202	A	6	5	1.00	6	0.833
203	A	4	3	1.00	11	0.273
204	A	3	3	1.00	11	0.273
205	A	1	1	1.48	9	0.111
206	A	1	1	1.00	1	1.000
207	B	3	3	12.64	21	0.143
208	A	1	1	1.00	20	0.050
209	A	5	5	1.00	11	0.455
210	A	2	2	1.00	9	0.222
211	F	0	0	N/A	0.000	N/A
212	A	5	4	1.73	18	0.222
213	A	1	1	1.00	9	0.111
214	A	2	2	2.00	12	0.167
215	A	5	4	1.30	7	0.571
216	A	3	2	1.00	9	0.222
217	A	2	2	1.00	12	0.167
218	A	4	3	1.00	11	0.273
219	A	5	4	1.65	15	0.267
220	A	4	3	1.33	21	0.143
221	A	3	3	1.06	6	0.500
222	A	1	1	1.00	18	0.056

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
223	A	2	2	1.00	6	0.333
224	A	8	7	0.92	8	0.875
225	N/A	2	0	1.00	9	0.000
226	N/A	2	0	1.00	14	0.000
227	A	2	2	1.00	9	0.222
228	A	7	6	1.91	15	0.400
229	A	4	4	1.00	12	0.333
230	A	2	2	1.00	11	0.182
231	A	4	3	1.00	11	0.273
232	A	1	1	1.00	11	0.091
233	A	2	2	1.00	9	0.222
234	C	4	4	2.07	12	0.333
235	A	3	3	1.22	19	0.158
236	N/A	1	0	1.00	5	0.000
237	A	6	5	0.90	8	0.625
238	A	3	2	1.00	11	0.182
239	A	4	3	1.00	4	0.750
240	A	1	1	1.00	17	0.059
241	A	4	3	1.06	13	0.231
242	B	6	5	8.08	23	0.217
243	F	0	0	N/A	0.000	N/A
244	B	1	1	2.75	19	0.053
245	F	0	0	N/A	0.000	N/A
246	C	1	1	14.00	37	0.027
247	A	3	2	1.00	11	0.182
248	A	4	3	0.47	8	0.375
249	A	4	4	1.00	8	0.500
250	F	0	0	N/A	0.000	N/A
251	A	2	2	1.00	14	0.143
252	A	2	2	1.00	22	0.091
253	A	5	4	1.47	9	0.444
254	B	11	10	3.88	11	0.909
255	A	6	5	1.00	9	0.556

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
256	A	4	3	0.91	9	0.333
257	C	3	3	36.17	16	0.188
258	C	2	2	9.00	19	0.105
259	A	6	5	1.16	12	0.417
260	A	3	3	1.00	13	0.231
261	A	2	2	1.00	21	0.095
262	A	1	1	1.00	34	0.029
263	A	1	1	1.00	7	0.143
264	A	3	3	1.00	20	0.150
265	F	0	0	N/A	0.000	N/A
266	B	4	3	2.21	22	0.136
267	A	2	2	1.00	13	0.154
268	A	2	2	1.79	6	0.333
269	A	2	2	1.12	19	0.105
270	A	2	2	1.00	12	0.167
271	A	2	2	1.00	16	0.125
272	B	4	4	2.36	11	0.364
273	A	5	4	1.14	29	0.138
274	A	3	3	1.16	7	0.429
275	F	0	0	N/A	0.000	N/A
276	B	6	5	2.59	39	0.128
277	F	0	0	N/A	0.000	N/A
278	A	3	3	1.04	15	0.200
279	A	1	1	1.00	1	1.000
280	A	2	2	1.00	2	1.000
281	A	5	4	1.00	19	0.211
282	A	2	2	1.00	29	0.069
283	A	3	3	1.00	17	0.176
284	A	1	1	2.00	13	0.077
285	A	1	1	1.00	25	0.040
286	A	1	1	1.00	13	0.077
287	B	1	1	3.00	43	0.023
288	C	1	1	50.00	19	0.053

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
289	A	1	1	1.00	11	0.091
290	A	1	1	1.00	31	0.032
291	A	5	4	1.15	15	0.267
292	A	1	1	1.89	17	0.059
293	F	0	0	N/A	0.000	N/A
294	C	6	5	8.22	16	0.312
295	A	1	1	1.00	20	0.050
296	A	2	2	1.00	17	0.118
297	A	8	7	1.21	4	1.750
298	F	0	0	N/A	0.000	N/A
299	A	2	2	0.88	13	0.154
300	A	5	5	1.11	6	0.833
301	A	7	6	1.55	17	0.353
302	A	1	1	1.00	25	0.040
303	A	5	5	1.00	8	0.625
304	A	1	1	1.00	40	0.025
305	B	25	24	2.21	7	3.429
306	A	6	5	1.00	18	0.278
307	F	0	0	N/A	0.000	N/A
308	A	1	1	1.00	16	0.062
309	A	2	2	1.00	13	0.154
310	A	4	3	1.00	8	0.375
311	A	6	6	1.00	4	1.500
312	A	5	4	1.00	9	0.444
313	F	0	0	N/A	0.000	N/A
314	A	2	2	1.00	25	0.080
315	A	7	6	0.82	17	0.353
316	A	1	1	1.00	17	0.059
317	F	0	0	N/A	0.000	N/A
318	A	8	7	1.07	14	0.500
319	B	19	18	2.12	47	0.383
320	F	0	0	N/A	0.000	N/A
321	A	4	3	1.16	16	0.188

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \sin(x) \sin(2x) \sin(3x) dx$	127
3.2	$\int \cos(x) \sin^3(2x) dx$	131
3.3	$\int \sqrt[3]{-1+x(1+x)^2} dx$	136
3.4	$\int x \log\left(1 + \frac{1}{x}\right) dx$	141
3.5	$\int \sin^2(\log(x)) dx$	146
3.6	$\int \frac{1}{1+3e^x} dx$	150
3.7	$\int \csc^3(x) \sec^5(x) dx$	155
3.8	$\int \frac{1}{x\sqrt{-1+x^4}} dx$	160
3.9	$\int \frac{1}{x(1+x^5)} dx$	165
3.10	$\int \sqrt{\tan(x)} dx$	170
3.11	$\int \frac{\log(1+x)}{1+x^2} dx$	177
3.12	$\int \frac{\sqrt{x}}{-\sqrt[3]{x+\sqrt{x}}} dx$	182
3.13	$\int x^x(1+\log(x)) dx$	187
3.14	$\int x^{13/2} \sqrt{1+x^{5/2}} dx$	191
3.15	$\int \frac{1}{(1+x^2)^2} dx$	196
3.16	$\int \frac{1}{36-13x^2+x^4} dx$	200
3.17	$\int \frac{\log(\log(x))}{x} dx$	204
3.18	$\int \frac{1+\cot(x)}{1-\cot(x)} dx$	208
3.19	$\int \frac{\cos(x)+x \sin(x)}{x(x+\cos(x))} dx$	212
3.20	$\int \frac{1}{\sec(x)+\sin(x)} dx$	217
3.21	$\int \frac{1}{\sqrt{1+e^x+e^{2x}}} dx$	223
3.22	$\int e^{x^2} x^3 dx$	228
3.23	$\int \left(-\frac{1}{\log^2(x)} + \frac{1}{\log(x)}\right) dx$	233
3.24	$\int \sqrt{2-x} \sqrt{-1+x} dx$	237
3.25	$\int \frac{-1+x^6}{-1-x+x^3+x^4} dx$	242
3.26	$\int (2 \log(x) + \log^2(x)) dx$	246
3.27	$\int \frac{2x}{\sqrt{1-x^4}} dx$	250

3.28	$\int \frac{1+x^2}{1+x} dx$	254
3.29	$\int \frac{-2-2\sin(x)+\sin^2(x)+\sin^3(x)}{1+2\sin(x)+\sin^2(x)} dx$	258
3.30	$\int \operatorname{csch}^2(x) dx$	264
3.31	$\int \sec^4(x) \tan^2(x) dx$	269
3.32	$\int \sqrt{\csc(x) - \sin(x)} dx$	274
3.33	$\int \cos^6(x) dx$	279
3.34	$\int \frac{1}{1+2x^2+x^4} dx$	284
3.35	$\int \cos(\log(x)) dx$	289
3.36	$\int \sec(x) dx$	293
3.37	$\int \frac{1}{9\cos^2(x)+4\sin^2(x)} dx$	297
3.38	$\int \frac{1}{x^2(1+x^4)^{3/4}} dx$	302
3.39	$\int \cos(x) \cos(3x) \cos(5x) dx$	306
3.40	$\int \left(\frac{1}{\log(x)} + \log(\log(x)) \right) dx$	310
3.41	$\int \frac{1}{2+e^x} dx$	314
3.42	$\int \sqrt{\frac{x}{1-x^3}} dx$	319
3.43	$\int \frac{4x}{1-x^4} dx$	324
3.44	$\int x^x (1 + \log(x)) dx$	328
3.45	$\int \sqrt{6x - x^2} dx$	332
3.46	$\int \sin^{99}(x) \sin(101x) dx$	337
3.47	$\int e^{e^{x^2}} x dx$	345
3.48	$\int \frac{-1+3x-3x^2+x^3}{1+4x+6x^2+4x^3+x^4} dx$	350
3.49	$\int \sqrt{\frac{1-x}{1+x}} dx$	355
3.50	$\int \frac{1}{-1+\sqrt{x}} dx$	360
3.51	$\int \sqrt[4]{x} \log(x) dx$	365
3.52	$\int \frac{1}{(1+\sqrt{x})\sqrt{x-x^2}} dx$	369
3.53	$\int \frac{1}{(1+\sqrt[4]{x})^{10} \sqrt{x}} dx$	374
3.54	$\int \sqrt{1-x^2} dx$	379
3.55	$\int \frac{1}{\sqrt{1-4x-x^2}} dx$	384
3.56	$\int \log\left(\frac{1}{x}\right) dx$	388
3.57	$\int \frac{1}{1+\sin(x)} dx$	392
3.58	$\int \frac{\sqrt{x}}{\sqrt{2012-x+\sqrt{x}}} dx$	396
3.59	$\int \frac{-1+x}{(1+x)\sqrt{x+x^2+x^3}} dx$	402
3.60	$\int \frac{1+4x+6x^2+4x^3+x^4}{-1+3x-3x^2+x^3} dx$	407
3.61	$\int \left(\cos(x) \log(x) + \frac{\sin(x)}{x} \right) dx$	412
3.62	$\int \frac{1}{-x+x^3} dx$	416
3.63	$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx$	421
3.64	$\int (1-x)^{99} x dx$	425

3.65	$\int \csc(x) \sin(4x) dx$	433
3.66	$\int \frac{1}{(1+\sqrt[3]{x})\sqrt{x}} dx$	437
3.67	$\int \frac{1}{\sqrt{-1+2x^2}} dx$	442
3.68	$\int \frac{1}{\sqrt{-1+e^x}} dx$	446
3.69	$\int \frac{x}{4+x^4} dx$	451
3.70	$\int \frac{2}{(\cos(x)-\sin(x))^2} dx$	455
3.71	$\int x \coth(x) \operatorname{csch}(x) dx$	459
3.72	$\int x^5 \sqrt{1+x^3} dx$	464
3.73	$\int \frac{-1+x^7}{\log(x)} dx$	469
3.74	$\int \sqrt{\csc(x) - \sin(x)} dx$	473
3.75	$\int (-2 \log(2x) + \log(x^2)) dx$	478
3.76	$\int e^x dx$	482
3.77	$\int \frac{\cos(x) \log(x) - \frac{\sin(x)}{x}}{\log^2(x)} dx$	486
3.78	$\int (-1 + 3x - 3x^2 + x^3) dx$	490
3.79	$\int \sqrt{12 - 3x^2} dx$	495
3.80	$\int ((-3+x)^7 + x - \sin(3-x)) dx$	500
3.81	$\int \sin(x) \sqrt{1 + \tan^2(x)} dx$	504
3.82	$\int \frac{-1+x^2-x^3+x^5}{-1+x-x^3+x^4} dx$	509
3.83	$\int \log(x) dx$	513
3.84	$\int \frac{1}{1-e^{-x}} dx$	517
3.85	$\int \cos^2(x) \sin^2(x) dx$	522
3.86	$\int \frac{\pi \sin(\pi\sqrt{x})}{\sqrt{x}} dx$	527
3.87	$\int \tan^2(x) dx$	532
3.88	$\int e^{\sqrt[4]{x}} dx$	536
3.89	$\int \cos(x) \cot(x) dx$	541
3.90	$\int (2 \log(x) + \log^2(x)) dx$	546
3.91	$\int \frac{x^3}{1+x^2} dx$	550
3.92	$\int \frac{1}{2-2x+x^2} dx$	554
3.93	$\int \log(\sin(x)) \sin(x) dx$	558
3.94	$\int \frac{x}{1-x^4} dx$	564
3.95	$\int \sqrt{12 - 3x^2} dx$	569
3.96	$\int \sec^5(x) \tan^3(x) dx$	574
3.97	$\int \frac{1}{1-\sin(x)} dx$	579
3.98	$\int \frac{1}{x\sqrt{-2+x^2}} dx$	583
3.99	$\int \log(x^2) dx$	588
3.100	$\int \sin(\sqrt[3]{x}) dx$	592
3.101	$\int e^{1+x-x^2} (1-2x) dx$	597
3.102	$\int e^{\sqrt{x}} \sqrt{x} dx$	601
3.103	$\int \cos(3x) \sin(2x) dx$	606

3.104	$\int (1 + 2 \sin(x)) dx$	610
3.105	$\int (1 - x)^{2014} x dx$	614
3.106	$\int \operatorname{arcsinh}(x) dx$	620
3.107	$\int \frac{x^2}{-1+x} dx$	624
3.108	$\int x \arctan(x) dx$	628
3.109	$\int \frac{1}{-2014-15x+x^2} dx$	633
3.110	$\int e^x (-2(1+x) \arctan(x) + \log(1+x^2)) dx$	637
3.111	$\int \arcsin(x)^2 dx$	641
3.112	$\int \frac{\sqrt{-1+x^2}}{x} dx$	645
3.113	$\int x \sec^2(4x) dx$	650
3.114	$\int \frac{2}{6-11x+6x^2-x^3} dx$	655
3.115	$\int \frac{1}{1-\log(1-x)} dx$	659
3.116	$\int \sqrt{x + \sqrt{1+x^2}} dx$	664
3.117	$\int \frac{1}{2+\cos(x)} dx$	669
3.118	$\int (\cos^4(x) - \sin^4(x)) dx$	673
3.119	$\int \frac{x}{\sqrt{2+4x}} dx$	677
3.120	$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx$	682
3.121	$\int \sec(x) dx$	687
3.122	$\int e^{\sin(x)} \cos(x) dx$	691
3.123	$\int x \log^2(x) dx$	695
3.124	$\int \frac{1}{5+4\sqrt{x+x}} dx$	699
3.125	$\int 2015^x dx$	704
3.126	$\int \frac{x}{(-3+x)(5+x)^2} dx$	708
3.127	$\int \frac{\log(1+\log(x))}{x} dx$	712
3.128	$\int \sqrt{\csc(x) - \sin(x)} dx$	716
3.129	$\int \frac{1}{\sqrt{25+x^2}} dx$	721
3.130	$\int \frac{-1+\log^2(x)}{x \log^2(x)} dx$	725
3.131	$\int e^{3x} \arctan(e^x) dx$	730
3.132	$\int \frac{1}{\cos^4(x)+\sin^4(x)} dx$	735
3.133	$\int \frac{1+e^x}{1-e^x} dx$	740
3.134	$\int \tan^4(x) dx$	745
3.135	$\int \sin(x) \tan^2(x) dx$	750
3.136	$\int \frac{1+x}{3+2x+x^2} dx$	755
3.137	$\int \tanh(x) dx$	759
3.138	$\int (-x + x^3) dx$	763
3.139	$\int \log(\sqrt{x}) dx$	767
3.140	$\int \left(-e^{e^{-x}+e^x-x} + e^{e^{-x}+e^x+x}\right) dx$	771
3.141	$\int \frac{\log(\log(x))}{x \log(x)} dx$	775

3.142	$\int \frac{1}{1+\tan^2(x)} dx$	779
3.143	$\int \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) dx$	784
3.144	$\int \log(x) dx$	789
3.145	$\int e^x(\cos(x) - \sin(x)) dx$	793
3.146	$\int e^{-x^2} x^3 dx$	797
3.147	$\int \left((1 + 2e^{x^2} x) \cos(x) - (e^{x^2} + x) \sin(x) \right) dx$	802
3.148	$\int \left(1 + \frac{1}{\sqrt{x}} + \frac{1}{\sqrt[3]{x}} \right) (1 + \sqrt[3]{x} + \sqrt{x}) dx$	807
3.149	$\int \cos(x) \cos(\sin(x)) \sin(\sin(x)) dx$	811
3.150	$\int \left(\frac{-\cos(x)+\sin(x)}{x} + \frac{\cos(x)+\sin(x)}{x^2} \right) dx$	816
3.151	$\int x^3 \sqrt{1+x^2} dx$	820
3.152	$\int \frac{x}{1+x^2+x^4} dx$	825
3.153	$\int e^{e^{2016x}+6048x} dx$	830
3.154	$\int (1 - \cot(x)) dx$	835
3.155	$\int \frac{1}{1-x+x^2-x^3} dx$	839
3.156	$\int \frac{1}{2+\cosh(x)} dx$	843
3.157	$\int \frac{x^2}{\sqrt{2+x^3}} dx$	847
3.158	$\int \frac{\log(x)}{x^2} dx$	852
3.159	$\int \operatorname{sech}(x) dx$	856
3.160	$\int e^{x^2} x^3 dx$	860
3.161	$\int \frac{1}{x\sqrt{-1+x^2}} dx$	865
3.162	$\int \frac{1}{x(1+x^2)} dx$	870
3.163	$\int \operatorname{arccosh}(x) dx$	875
3.164	$\int e^{-3-5x-2x^2} dx$	879
3.165	$\int \sin(\sqrt{x}) dx$	883
3.166	$\int \frac{1}{\left(\frac{1}{x}+x\right)^2} dx$	888
3.167	$\int \frac{e^{-x}(2+x)}{x^3} dx$	893
3.168	$\int \frac{1}{\sqrt{(1-x)x}} dx$	898
3.169	$\int e^{-x} \tanh(x) dx$	902
3.170	$\int \sqrt{1+\sin(x)} dx$	907
3.171	$\int \frac{1}{1+\sqrt{x}} dx$	911
3.172	$\int e^{-x^2} \sin^2\left(\frac{\pi}{4}+x\right) dx$	915
3.173	$\int 3e^{-2x^3-x^6} x^2(1+x^3)^2 dx$	919
3.174	$\int e^{2x} \cos(3x) dx$	924
3.175	$\int \cos^{\cos(x)}(x)(1+\log(\cos(x))) \sin(x) dx$	928
3.176	$\int \frac{e^x}{2+e^x} dx$	932
3.177	$\int \sin(2018x) dx$	936

3.178	$\int \frac{1}{\cot(x)+\tan(x)} dx$	940
3.179	$\int \frac{x^5}{2+x^{12}} dx$	945
3.180	$\int (\cos(x) \cosh(x) + \sin(x) \sinh(x)) dx$	949
3.181	$\int \frac{e^x + \cos(x)}{e^x + \sin(x)} dx$	953
3.182	$\int \cos(x) \sin(\cos(\sin(x))) \sin(\sin(x)) dx$	957
3.183	$\int \frac{1}{1+\sin(x)} dx$	962
3.184	$\int \frac{\cos(x)}{1-\cos(2x)} dx$	966
3.185	$\int e^x \left(\frac{1}{x} + \log(x) \right) dx$	971
3.186	$\int \tanh^2(x) dx$	975
3.187	$\int \frac{-\sin^2(x) + \sin(2x)}{-\cos^2(x) + \cos(2x)} dx$	980
3.188	$\int \frac{1}{x^{9/25} + x^{41/25}} dx$	985
3.189	$\int \frac{\cos(x)}{2 - \cos^2(x)} dx$	990
3.190	$\int \frac{1}{(1+x^2)^{3/2}} dx$	995
3.191	$\int \frac{1}{\sqrt{x^{3/2} - x^2}} dx$	999
3.192	$\int \frac{-1+x}{x+x^2 \log(x)} dx$	1003
3.193	$\int \csc(x) \sec(x) dx$	1007
3.194	$\int \tan(\cos(x)) dx$	1011
3.195	$\int \frac{1+x}{x(x+\log(x))} dx$	1016
3.196	$\int (e^{-e^x+x} + e^{e^x+x}) dx$	1020
3.197	$\int \frac{1}{1-x^2} dx$	1024
3.198	$\int 2^{\log(x)} dx$	1028
3.199	$\int (\cos(3x) + \sin(2x)(\cos(3x) - \sin(2019x))) dx$	1032
3.200	$\int \cos(x) \cos(\sin(x)) \cos(\sin(\sin(x))) dx$	1036
3.201	$\int \frac{e^{-\frac{2019}{4x^2}}}{x^2} dx$	1041
3.202	$\int \sin(\sqrt{x}) dx$	1045
3.203	$\int \frac{\sqrt{x}}{1+x} dx$	1050
3.204	$\int \cos(x) \cos(2x) \cos(3x) dx$	1055
3.205	$\int e^{-x^{2n}} dx$	1059
3.206	$\int e dx$	1063
3.207	$\int \frac{\sin(19x) + \sin(20x)}{\cos(19x) + \cos(20x)} dx$	1067
3.208	$\int e^x (\cos^2(x) + \cos(x) \sin(x) - \sin^2(x)) dx$	1073
3.209	$\int \csc\left(\frac{\pi}{4} + x\right) \sin(x) dx$	1077
3.210	$\int \frac{1}{\sqrt[3]{x+x}} dx$	1082
3.211	$\int x^{1+x^2} (1 + 2 \log(x)) dx$	1086
3.212	$\int \frac{-1+2x^3}{x(1+x^3)} dx$	1090
3.213	$\int \frac{1}{\sqrt{1+x^2}} dx$	1095
3.214	$\int \frac{\log(2x)}{x \log(x)} dx$	1099
3.215	$\int \frac{1}{1+e^x} dx$	1103

3.216	$\int \frac{\log(x) \log(\log(x))}{x} dx$	1108
3.217	$\int \log\left(\frac{1+x}{1-x}\right) dx$	1112
3.218	$\int \frac{1}{(-1+x)^2+x^2} dx$	1116
3.219	$\int \sqrt{x} \sqrt{x^{3/2}} dx$	1120
3.220	$\int \cos^4(x)(\cos(x) - \sin(x)) \sin^4(x)(\cos(x) + \sin(x)) dx$	1125
3.221	$\int \log(1+x^2) dx$	1130
3.222	$\int \frac{1+2x}{1+2x+2x^2} dx$	1135
3.223	$\int \frac{\arcsin(x)}{x^3} dx$	1139
3.224	$\int \cos(\cos(x)) \sin(2x) dx$	1143
3.225	$\int -\sin(x - \sin(x)) dx$	1148
3.226	$\int \frac{1}{1+\tan^2 \sqrt{505}(x)} dx$	1152
3.227	$\int (1-x)^{2020} x dx$	1157
3.228	$\int \frac{\sec^4(x) \tan(x)}{4+\sec^4(x)} dx$	1163
3.229	$\int x^{2x}(2+2\log(x)) dx$	1168
3.230	$\int \sqrt{1-x^2} dx$	1173
3.231	$\int e^{-x^4} x^5 dx$	1178
3.232	$\int \frac{1+\cos(x)}{x+\sin(x)} dx$	1183
3.233	$\int \frac{\cot^{-1}(x)+\arctan(x)}{x} dx$	1187
3.234	$\int \frac{\sinh(x)}{\cosh(x)-\sinh(x)} dx$	1191
3.235	$\int \frac{x}{\sqrt{-1+x+\sqrt{1+x}}} dx$	1196
3.236	$\int \cos(x + \cos(x)) dx$	1200
3.237	$\int x^3 \sin(x^2) dx$	1204
3.238	$\int \frac{x}{1-x^4} dx$	1209
3.239	$\int \operatorname{sech}^2(x) dx$	1214
3.240	$\int (e^{e^x} - e^{e^x-x}) dx$	1218
3.241	$\int \sqrt{1-\sqrt{x}} dx$	1222
3.242	$\int \frac{x^3}{1+x+\frac{x^2}{2}+\frac{x^3}{6}} dx$	1227
3.243	$\int (-\sin(x - \sin(x)) + \sin(x + \sin(x))) dx$	1233
3.244	$\int (\sec^5(x) \tan^2(x) + \sec^3(x) \tan^4(x)) dx$	1237
3.245	$\int (1 + \log(x)) \log(\log(x)) dx$	1242
3.246	$\int \left(\frac{1}{1+\cos(x)} + \frac{1}{1+\cot(x)} + \frac{1}{1+\csc(x)} + \frac{1}{1+\sec(x)} + \frac{1}{1+\sin(x)} + \frac{1}{1+\tan(x)} \right) dx$	1246
3.247	$\int \frac{1}{\sqrt{x-x^2}} dx$	1251
3.248	$\int \frac{1}{1+\cos^2(x)} dx$	1255
3.249	$\int \frac{\log(1+x)}{x^2} dx$	1260
3.250	$\int \sqrt{1 - \arccos(\sin(x))^2} dx$	1265
3.251	$\int (-2+x)(-1+x)x(1+x)(2+x) dx$	1269
3.252	$\int \frac{1+2x^2+3x^3}{\sqrt[3]{1+x^3}} dx$	1273
3.253	$\int \csc^4(x) \sec^4(x) dx$	1277

3.254	$\int \frac{x+\sin(x)}{1+\cos(x)} dx$	1282
3.255	$\int \cosh^2(x) \sinh^3(x) dx$	1288
3.256	$\int 3^{2^x} 4^x dx$	1293
3.257	$\int \frac{\cos(x)-\sin(x)}{2+\sin(2x)} dx$	1298
3.258	$\int \frac{\sec^2(1+\log(x))-\tan(1+\log(x))}{x^2} dx$	1303
3.259	$\int \sqrt{\frac{\log(\frac{1}{x})}{x}} dx$	1308
3.260	$\int (1-\cos(x))^5 \cos(5x) dx$	1313
3.261	$\int \frac{3\cos(x)+4\sin(x)}{4\cos(x)+3\sin(x)} dx$	1319
3.262	$\int \left(-\sqrt{3}-\sqrt{4-x^2}+\sqrt{4-(1+x)^2}\right) dx$	1324
3.263	$\int x^2 \sin(\log(x)) dx$	1329
3.264	$\int e^{-x}(36x^5-12x^6+x^7) dx$	1333
3.265	$\int \arccos(x) \arcsin(x) dx$	1338
3.266	$\int (1+6x-7x^2+4x^3-x^4)^n dx$	1342
3.267	$\int \frac{x^4}{\sqrt{1-x}} dx$	1347
3.268	$\int \sin(x^{-n}) dx$	1352
3.269	$\int \frac{1}{2}(-x+\sqrt{4-3x^2}) dx$	1356
3.270	$\int (1-3x^2+x^4)^n dx$	1360
3.271	$\int \frac{(1+e^{-x})x}{-1+e^x} dx$	1365
3.272	$\int e^{-x}x^4 \sin(x) dx$	1369
3.273	$\int \frac{-3(-3x+x^3)+(-3x+x^3)^3}{\sqrt{-4+x^2}} dx$	1374
3.274	$\int \frac{1}{(1+x^2)^3} dx$	1379
3.275	$\int \log(\sqrt{3}+\tan(x)) dx$	1383
3.276	$\int \sqrt{(\cos(20x)+3\cos(21x)+\cos(22x))^2+(\sin(20x)+3\sin(21x)+\sin(22x))^2} dx$	1388
3.277	$\int \frac{e^{-2x} \sin(3x)}{x} dx$	1393
3.278	$\int (1-x)^{2/3} \sqrt[3]{x} dx$	1397
3.279	$\int e dx$	1402
3.280	$\int \operatorname{sech}(x) dx$	1406
3.281	$\int \frac{e^x}{(1+e^x)\log(1+e^x)} dx$	1410
3.282	$\int (1-x+x^2-x^3+x^4)(1+x+x^2+x^3+x^4) dx$	1415
3.283	$\int \frac{1}{120}(-4+x)(-3+x)(-2+x)(-1+x)x dx$	1419
3.284	$\int (x+x\cos(x)+\sin(x)+\cos(x)\sin(x)) dx$	1423
3.285	$\int (\cos^2(x)+\cot^2(x)+\csc^2(x)+\sec^2(x)+\sin^2(x)+\tan^2(x)) dx$	1427
3.286	$\int e^{\log^2(x)}(1+2\log(x)) dx$	1431
3.287	$\int \left((1-x)^3+(x-x^2)^3-3(1-x)(x-x^2)(-1+x^2)+(-1+x^2)^3\right) dx$	1435
3.288	$\int (\cos^6(x)+3\cos^2(x)\sin^2(x)+\sin^6(x)) dx$	1439
3.289	$\int e^x x^e(1+e+x) dx$	1443
3.290	$\int \left(\sqrt{2}\sqrt{\frac{x}{1+x}+\frac{x^2}{2-x^2}}\right) dx$	1447

3.291	$\int \frac{1+2x^{2022}}{x+x^{2023}} dx$	1452
3.292	$\int (3 \cos(23x) \sin(20x) + 20 \sin(43x)) dx$	1457
3.293	$\int \frac{\sin(x)}{2e^x + \cos(x) + \sin(x)} dx$	1462
3.294	$\int \log^{-\log(e^\pi)}(x) \log\left(\frac{x}{\pi}\right) dx$	1466
3.295	$\int (x + 8x^2 + 10x^3 + 5x^4 + x^5) dx$	1471
3.296	$\int \frac{(1+x)(4+x)}{(2+x)(3+x)} dx$	1475
3.297	$\int x \cot(x) dx$	1479
3.298	$\int \frac{e^{\frac{1}{x}+x}(-1-x^2+x^4+x^6)}{x^4} dx$	1484
3.299	$\int \frac{1}{\sqrt{(-1+x)(1+x)^3}} dx$	1488
3.300	$\int x \sin^4(x) dx$	1492
3.301	$\int \cos(3x) \csc^2(x) \sec^3(x) \sin(2x) dx$	1497
3.302	$\int \left(\frac{1}{\sqrt{2}\sqrt{\log(x)}} + \sqrt{2}\sqrt{\log(x)} \right) dx$	1502
3.303	$\int \log(\cos(x)) \sec^2(x) dx$	1507
3.304	$\int \frac{\frac{5}{(-5+x)^6} + \frac{3}{(-3+x)^4} + \frac{1}{(-1+x)^2}}{\left(1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}\right)^2} dx$	1512
3.305	$\int \csc(x) \sin(23x) dx$	1517
3.306	$\int \frac{(1-x)^2 x^4}{1+x^2} dx$	1527
3.307	$\int x^{-\log(x)} dx$	1532
3.308	$\int \frac{1-2x}{x^{2/3}(1+x)^2} dx$	1536
3.309	$\int \cos^2\left(\frac{\pi x^2}{\sqrt{2}}\right) dx$	1541
3.310	$\int \frac{1}{1+\cos(x)+\sin(x)} dx$	1546
3.311	$\int \tan^5(x) dx$	1550
3.312	$\int \sqrt{1 + \frac{1}{x}} dx$	1555
3.313	$\int e^{\cos(x)} \cos(2x + \sin(x)) dx$	1560
3.314	$\int \frac{-1+2x+3 \log(x)}{x^2+2x^4+x \log(x)} dx$	1564
3.315	$\int (-\sqrt{x} + \sqrt{1+x})^\pi dx$	1568
3.316	$\int \left(-2 + \left(-2 + \left(-2 + (-2 + x^2)^2 \right)^2 \right)^2 \right) dx$	1574
3.317	$\int \sin(4 \arctan(x)) dx$	1579
3.318	$\int \frac{\sqrt[3]{\tan(x)}}{(\cos(x)+\sin(x))^2} dx$	1583
3.319	$\int \csc^2(x) \csc^2(6x) \csc^2(10x) \csc^2(15x) \sin^2(2x) \sin^2(3x) \sin^2(5x) \sin^2(30x) dx$	1589
3.320	$\int \sqrt{1+x^2} + \sqrt{1+x^2+x^4} dx$	1598
3.321	$\int \frac{x^9}{575-48x^{10}+x^{20}} dx$	1602

3.1 $\int \sin(x) \sin(2x) \sin(3x) dx$

3.1.1	Optimal result	127
3.1.2	Mathematica [A] (verified)	127
3.1.3	Rubi [A] (verified)	128
3.1.4	Maple [A] (verified)	129
3.1.5	Fricas [A] (verification not implemented)	129
3.1.6	Sympy [B] (verification not implemented)	129
3.1.7	Maxima [A] (verification not implemented)	130
3.1.8	Giac [A] (verification not implemented)	130
3.1.9	Mupad [B] (verification not implemented)	130

3.1.1 Optimal result

Integrand size = 11, antiderivative size = 25

$$\int \sin(x) \sin(2x) \sin(3x) dx = -\frac{1}{8} \cos(2x) - \frac{1}{16} \cos(4x) + \frac{1}{24} \cos(6x)$$

output `-1/8*cos(2*x)-1/16*cos(4*x)+1/24*cos(6*x)`

3.1.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \sin(x) \sin(2x) \sin(3x) dx = -\frac{1}{8} \cos(2x) - \frac{1}{16} \cos(4x) + \frac{1}{24} \cos(6x)$$

input `Integrate[Sin[x]*Sin[2*x]*Sin[3*x],x]`

output `-1/8*Cos[2*x] - Cos[4*x]/16 + Cos[6*x]/24`

3.1.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 4855, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin(x) \sin(2x) \sin(3x) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin(x) \sin(2x) \sin(3x) dx \\ & \quad \downarrow \text{4855} \\ & \int \left(\frac{1}{4} \sin(2x) + \frac{1}{4} \sin(4x) - \frac{1}{4} \sin(6x) \right) dx \\ & \quad \downarrow \text{2009} \\ & -\frac{1}{8} \cos(2x) - \frac{1}{16} \cos(4x) + \frac{1}{24} \cos(6x) \end{aligned}$$

input `Int[Sin[x]*Sin[2*x]*Sin[3*x],x]`

output `-1/8*Cos[2*x] - Cos[4*x]/16 + Cos[6*x]/24`

3.1.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4855 `Int[(F_)[(a_.) + (b_.)*(x_)]^(p_.)*(G_)[(c_.) + (d_.)*(x_)]^(q_.)*(H_)[(e_.) + (f_.)*(x_)]^(r_.), x_Symbol] := Int[ExpandTrigReduce[ActivateTrig[F[a + b*x]^p*G[c + d*x]^q*H[e + f*x]^r], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && (EqQ[F, sin] || EqQ[F, cos]) && (EqQ[G, sin] || EqQ[G, cos]) && (EqQ[H, sin] || EqQ[H, cos]) && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]`

3.1.4 Maple [A] (verified)

Time = 1.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

method	result	size
default	$-\frac{\cos(2x)}{8} - \frac{\cos(4x)}{16} + \frac{\cos(6x)}{24}$	20
risch	$-\frac{\cos(2x)}{8} - \frac{\cos(4x)}{16} + \frac{\cos(6x)}{24}$	20
parallelrisch	$\frac{1}{16} + \frac{\cos(6x)}{24} - \frac{\cos(4x)}{16} - \frac{\cos(2x)}{8}$	21

input `int(sin(x)*sin(2*x)*sin(3*x),x,method=_RETURNVERBOSE)`

output `-1/8*cos(2*x)-1/16*cos(4*x)+1/24*cos(6*x)`

3.1.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.68

$$\int \sin(x) \sin(2x) \sin(3x) dx = \frac{4}{3} \cos(x)^6 - \frac{5}{2} \cos(x)^4 + \cos(x)^2$$

input `integrate(sin(x)*sin(2*x)*sin(3*x),x, algorithm="fricas")`

output `4/3*cos(x)^6 - 5/2*cos(x)^4 + cos(x)^2`

3.1.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 114 vs. 2(19) = 38.

Time = 1.04 (sec) , antiderivative size = 114, normalized size of antiderivative = 4.56

$$\begin{aligned} \int \sin(x) \sin(2x) \sin(3x) dx = & \frac{x \sin(x) \sin(2x) \sin(3x)}{4} + \frac{x \sin(x) \cos(2x) \cos(3x)}{4} \\ & + \frac{x \sin(2x) \cos(x) \cos(3x)}{4} \\ & - \frac{x \sin(3x) \cos(x) \cos(2x)}{4} - \frac{3 \sin(x) \sin(2x) \cos(3x)}{8} \\ & + \frac{\sin(x) \sin(3x) \cos(2x)}{6} + \frac{\sin(2x) \sin(3x) \cos(x)}{24} \end{aligned}$$

input `integrate(sin(x)*sin(2*x)*sin(3*x),x)`

output `x*sin(x)*sin(2*x)*sin(3*x)/4 + x*sin(x)*cos(2*x)*cos(3*x)/4 + x*sin(2*x)*cos(x)*cos(3*x)/4 - x*sin(3*x)*cos(x)*cos(2*x)/4 - 3*sin(x)*sin(2*x)*cos(3*x)/8 + sin(x)*sin(3*x)*cos(2*x)/6 + sin(2*x)*sin(3*x)*cos(x)/24`

3.1.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \sin(x) \sin(2x) \sin(3x) dx = \frac{1}{24} \cos(6x) - \frac{1}{16} \cos(4x) - \frac{1}{8} \cos(2x)$$

input `integrate(sin(x)*sin(2*x)*sin(3*x),x, algorithm="maxima")`

output `1/24*cos(6*x) - 1/16*cos(4*x) - 1/8*cos(2*x)`

3.1.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.52

$$\int \sin(x) \sin(2x) \sin(3x) dx = -\frac{4}{3} \sin(x)^6 + \frac{3}{2} \sin(x)^4$$

input `integrate(sin(x)*sin(2*x)*sin(3*x),x, algorithm="giac")`

output `-4/3*sin(x)^6 + 3/2*sin(x)^4`

3.1.9 Mupad [B] (verification not implemented)

Time = 16.61 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.56

$$\int \sin(x) \sin(2x) \sin(3x) dx = -\frac{\sin(x)^4 (8 \sin(x)^2 - 9)}{6}$$

input `int(sin(2*x)*sin(3*x)*sin(x),x)`

output `-(sin(x)^4*(8*sin(x)^2 - 9))/6`

3.2 $\int \cos(x) \sin^3(2x) dx$

3.2.1	Optimal result	131
3.2.2	Mathematica [A] (verified)	131
3.2.3	Rubi [A] (verified)	132
3.2.4	Maple [A] (verified)	133
3.2.5	Fricas [A] (verification not implemented)	134
3.2.6	Sympy [B] (verification not implemented)	134
3.2.7	Maxima [A] (verification not implemented)	134
3.2.8	Giac [A] (verification not implemented)	135
3.2.9	Mupad [B] (verification not implemented)	135

3.2.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \cos(x) \sin^3(2x) dx = -\frac{8}{5} \cos^5(x) + \frac{8 \cos^7(x)}{7}$$

output `-8/5*cos(x)^5+8/7*cos(x)^7`

3.2.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.82

$$\int \cos(x) \sin^3(2x) dx = -\frac{3 \cos(x)}{8} - \frac{1}{8} \cos(3x) + \frac{1}{40} \cos(5x) + \frac{1}{56} \cos(7x)$$

input `Integrate[Cos[x]*Sin[2*x]^3,x]`

output `(-3*Cos[x])/8 - Cos[3*x]/8 + Cos[5*x]/40 + Cos[7*x]/56`

3.2.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4775, 3042, 3045, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^3(2x) \cos(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(2x)^3 \cos(x) dx \\
 & \quad \downarrow \text{4775} \\
 & 8 \int \cos^4(x) \sin^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & 8 \int \cos(x)^4 \sin(x)^3 dx \\
 & \quad \downarrow \text{3045} \\
 & -8 \int \cos^4(x) (1 - \cos^2(x)) d \cos(x) \\
 & \quad \downarrow \text{244} \\
 & -8 \int (\cos^4(x) - \cos^6(x)) d \cos(x) \\
 & \quad \downarrow \text{2009} \\
 & -8 \left(\frac{\cos^5(x)}{5} - \frac{\cos^7(x)}{7} \right)
 \end{aligned}$$

input `Int[Cos[x]*Sin[2*x]^3,x]`

output `-8*(Cos[x]^5/5 - Cos[x]^7/7)`

3.2.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3045 `Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[-(a*f)^(-1) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])`

rule 4775 `Int[(cos[(a_.) + (b_.)*(x_)]*(e_.))^(m_.)*sin[(c_.) + (d_.)*(x_)]^(p_.), x_Symbol] := Simp[2^p/e^p Int[(e*Cos[a + b*x])^(m + p)*Sin[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && EqQ[b*c - a*d, 0] && EqQ[d/b, 2] && IntegerQ[p]`

3.2.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.41

method	result	size
default	$-\frac{3 \cos(x)}{8} - \frac{\cos(3x)}{8} + \frac{\cos(5x)}{40} + \frac{\cos(7x)}{56}$	24
risch	$-\frac{3 \cos(x)}{8} - \frac{\cos(3x)}{8} + \frac{\cos(5x)}{40} + \frac{\cos(7x)}{56}$	24
parallelrisch	$\frac{8}{21} - \frac{3 \cos(x)}{8} - \frac{\cos(3x)}{8} + \frac{\cos(5x)}{40} + \frac{\cos(7x)}{56}$	25

input `int(sin(2*x)^3*cos(x),x,method=_RETURNVERBOSE)`

output `-3/8*cos(x)-1/8*cos(3*x)+1/40*cos(5*x)+1/56*cos(7*x)`

3.2.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(x) \sin^3(2x) dx = \frac{8}{7} \cos(x)^7 - \frac{8}{5} \cos(x)^5$$

input `integrate(sin(2*x)^3*cos(x),x, algorithm="fricas")`

output `8/7*cos(x)^7 - 8/5*cos(x)^5`

3.2.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. 2(15) = 30.

Time = 0.59 (sec) , antiderivative size = 65, normalized size of antiderivative = 3.82

$$\int \cos(x) \sin^3(2x) dx = -\frac{9 \sin(x) \sin^3(2x)}{35} - \frac{8 \sin(x) \sin(2x) \cos^2(2x)}{35} \\ - \frac{22 \sin^2(2x) \cos(x) \cos(2x)}{35} - \frac{16 \cos(x) \cos^3(2x)}{35}$$

input `integrate(sin(2*x)**3*cos(x),x)`

output `-9*sin(x)*sin(2*x)**3/35 - 8*sin(x)*sin(2*x)*cos(2*x)**2/35 - 22*sin(2*x)*
*2*cos(x)*cos(2*x)/35 - 16*cos(x)*cos(2*x)**3/35`

3.2.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.35

$$\int \cos(x) \sin^3(2x) dx = \frac{1}{56} \cos(7x) + \frac{1}{40} \cos(5x) - \frac{1}{8} \cos(3x) - \frac{3}{8} \cos(x)$$

input `integrate(sin(2*x)^3*cos(x),x, algorithm="maxima")`

output `1/56*cos(7*x) + 1/40*cos(5*x) - 1/8*cos(3*x) - 3/8*cos(x)`

3.2.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(x) \sin^3(2x) dx = \frac{8}{7} \cos(x)^7 - \frac{8}{5} \cos(x)^5$$

input `integrate(sin(2*x)^3*cos(x),x, algorithm="giac")`

output `8/7*cos(x)^7 - 8/5*cos(x)^5`

3.2.9 Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \cos(x) \sin^3(2x) dx = \frac{8 \cos(x)^5 (5 \cos(x)^2 - 7)}{35}$$

input `int(sin(2*x)^3*cos(x),x)`

output `(8*cos(x)^5*(5*cos(x)^2 - 7))/35`

3.3 $\int \sqrt[3]{-1+x}(1+x)^2 dx$

3.3.1	Optimal result	136
3.3.2	Mathematica [A] (verified)	136
3.3.3	Rubi [A] (verified)	137
3.3.4	Maple [A] (verified)	138
3.3.5	Fricas [A] (verification not implemented)	138
3.3.6	Sympy [C] (verification not implemented)	139
3.3.7	Maxima [A] (verification not implemented)	139
3.3.8	Giac [A] (verification not implemented)	140
3.3.9	Mupad [B] (verification not implemented)	140

3.3.1 Optimal result

Integrand size = 13, antiderivative size = 21

$$\int \sqrt[3]{-1+x}(1+x)^2 dx = \frac{3}{70}(-1+x)^{4/3} (37 + 26x + 7x^2)$$

output `3/70*(-1+x)^(4/3)*(7*x^2+26*x+37)`

3.3.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \sqrt[3]{-1+x}(1+x)^2 dx = \frac{3}{70}(-1+x)^{4/3} (37 + 26x + 7x^2)$$

input `Integrate[(-1 + x)^(1/3)*(1 + x)^2,x]`

output `(3*(-1 + x)^(4/3)*(37 + 26*x + 7*x^2))/70`

3.3.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.52, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt[3]{x-1}(x+1)^2 dx$$

$$\downarrow \text{53}$$

$$\int \left((x-1)^{7/3} + 4(x-1)^{4/3} + 4\sqrt[3]{x-1} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{3}{10}(x-1)^{10/3} + \frac{12}{7}(x-1)^{7/3} + 3(x-1)^{4/3}$$

input `Int[(-1 + x)^(1/3)*(1 + x)^2,x]`

output `3*(-1 + x)^(4/3) + (12*(-1 + x)^(7/3))/7 + (3*(-1 + x)^(10/3))/10`

3.3.3.1 Defintions of rubi rules used

rule 53 `Int[(a_.) + (b_.)*(x_)^(m_.)*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0])
|| LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.3.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

method	result
gosper	$\frac{3(-1+x)^{\frac{4}{3}}(7x^2+26x+37)}{70}$
trager	$\left(\frac{3}{10}x^3 + \frac{57}{70}x^2 + \frac{33}{70}x - \frac{111}{70}\right)(-1+x)^{\frac{1}{3}}$
derivativedivides	$\frac{3(-1+x)^{\frac{10}{3}}}{10} + \frac{12(-1+x)^{\frac{7}{3}}}{7} + 3(-1+x)^{\frac{4}{3}}$
default	$\frac{3(-1+x)^{\frac{10}{3}}}{10} + \frac{12(-1+x)^{\frac{7}{3}}}{7} + 3(-1+x)^{\frac{4}{3}}$
risch	$\frac{3(-1+x)^{\frac{1}{3}}(7x^3+19x^2+11x-37)}{70}$
meijerg	$\frac{\text{signum}(-1+x)^{\frac{1}{3}} \text{hypergeom}\left(\left[-\frac{1}{3}, 1\right], [2], x\right)}{\left(-\text{signum}(-1+x)\right)^{\frac{1}{3}}} + \frac{\text{signum}(-1+x)^{\frac{1}{3}} x^2 \text{hypergeom}\left(\left[-\frac{1}{3}, 2\right], [3], x\right)}{\left(-\text{signum}(-1+x)\right)^{\frac{1}{3}}} + \frac{\text{signum}(-1+x)^{\frac{1}{3}} x^3}{3(-\text{signum}(-1+x))^{\frac{1}{3}}}$

input `int((1+x)^2*(-1+x)^(1/3),x,method=_RETURNVERBOSE)`

output `3/70*(-1+x)^(4/3)*(7*x^2+26*x+37)`

3.3.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \sqrt[3]{-1+x}(1+x)^2 dx = \frac{3}{70} (7x^3 + 19x^2 + 11x - 37)(x-1)^{\frac{1}{3}}$$

input `integrate((1+x)^2*(-1+x)^(1/3),x, algorithm="fracas")`

output `3/70*(7*x^3 + 19*x^2 + 11*x - 37)*(x - 1)^(1/3)`

3.3.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.86 (sec) , antiderivative size = 144, normalized size of antiderivative = 6.86

$$\int \sqrt[3]{-1+x}(1+x)^2 dx = \begin{cases} \frac{3\sqrt[3]{x-1}(x+1)^3}{10} - \frac{3\sqrt[3]{x-1}(x+1)^2}{35} - \frac{9\sqrt[3]{x-1}(x+1)}{35} - \frac{54\sqrt[3]{x-1}}{35} & \text{for } |x+1| > 2 \\ -\frac{3\sqrt[3]{1-x}(x+1)^3 e^{-\frac{2i\pi}{3}}}{10} + \frac{3\sqrt[3]{1-x}(x+1)^2 e^{-\frac{2i\pi}{3}}}{35} + \frac{9\sqrt[3]{1-x}(x+1) e^{-\frac{2i\pi}{3}}}{35} + \frac{54\sqrt[3]{1-x} e^{-\frac{2i\pi}{3}}}{35} & \text{otherwise} \end{cases}$$

input `integrate((1+x)**2*(-1+x)**(1/3),x)`

output `Piecewise((3*(x - 1)**(1/3)*(x + 1)**3/10 - 3*(x - 1)**(1/3)*(x + 1)**2/35 - 9*(x - 1)**(1/3)*(x + 1)/35 - 54*(x - 1)**(1/3)/35, Abs(x + 1) > 2), (-3*(1 - x)**(1/3)*(x + 1)**3*exp(-2*I*pi/3)/10 + 3*(1 - x)**(1/3)*(x + 1)**2*exp(-2*I*pi/3)/35 + 9*(1 - x)**(1/3)*(x + 1)*exp(-2*I*pi/3)/35 + 54*(1 - x)**(1/3)*exp(-2*I*pi/3)/35, True))`

3.3.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \sqrt[3]{-1+x}(1+x)^2 dx = \frac{3}{10} (x-1)^{\frac{10}{3}} + \frac{12}{7} (x-1)^{\frac{7}{3}} + 3(x-1)^{\frac{4}{3}}$$

input `integrate((1+x)^2*(-1+x)^(1/3),x, algorithm="maxima")`

output `3/10*(x - 1)^(10/3) + 12/7*(x - 1)^(7/3) + 3*(x - 1)^(4/3)`

3.3.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \sqrt[3]{-1+x}(1+x)^2 dx = \frac{3}{10}(x-1)^{\frac{10}{3}} + \frac{12}{7}(x-1)^{\frac{7}{3}} + 3(x-1)^{\frac{4}{3}}$$

input `integrate((1+x)^2*(-1+x)^(1/3),x, algorithm="giac")`

output `3/10*(x - 1)^(10/3) + 12/7*(x - 1)^(7/3) + 3*(x - 1)^(4/3)`

3.3.9 Mupad [B] (verification not implemented)

Time = 16.73 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \sqrt[3]{-1+x}(1+x)^2 dx = \frac{3(x-1)^{4/3}(40x+7(x-1)^2+30)}{70}$$

input `int((x - 1)^(1/3)*(x + 1)^2,x)`

output `(3*(x - 1)^(4/3)*(40*x + 7*(x - 1)^2 + 30))/70`

3.4 $\int x \log \left(1 + \frac{1}{x}\right) dx$

3.4.1	Optimal result	141
3.4.2	Mathematica [A] (verified)	141
3.4.3	Rubi [A] (verified)	142
3.4.4	Maple [A] (verified)	143
3.4.5	Fricas [A] (verification not implemented)	143
3.4.6	Sympy [A] (verification not implemented)	144
3.4.7	Maxima [A] (verification not implemented)	144
3.4.8	Giac [B] (verification not implemented)	144
3.4.9	Mupad [B] (verification not implemented)	145

3.4.1 Optimal result

Integrand size = 8, antiderivative size = 27

$$\int x \log \left(1 + \frac{1}{x}\right) dx = \frac{x}{2} + \frac{1}{2}x^2 \log \left(1 + \frac{1}{x}\right) - \frac{1}{2} \log(1 + x)$$

output `1/2*x+1/2*x^2*ln(1+1/x)-1/2*ln(1+x)`

3.4.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int x \log \left(1 + \frac{1}{x}\right) dx = \frac{x}{2} + \frac{1}{2}x^2 \log \left(1 + \frac{1}{x}\right) - \frac{1}{2} \log(1 + x)$$

input `Integrate[x*Log[1 + x^(-1)],x]`

output `x/2 + (x^2*Log[1 + x^(-1)])/2 - Log[1 + x]/2`

3.4.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {2905, 772, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \log\left(\frac{1}{x} + 1\right) dx \\
 & \quad \downarrow \text{2905} \\
 & \frac{1}{2} \int \frac{1}{1 + \frac{1}{x}} dx + \frac{1}{2} x^2 \log\left(\frac{1}{x} + 1\right) \\
 & \quad \downarrow \text{772} \\
 & \frac{1}{2} \int \frac{x}{x+1} dx + \frac{1}{2} x^2 \log\left(\frac{1}{x} + 1\right) \\
 & \quad \downarrow \text{49} \\
 & \frac{1}{2} \int \left(1 + \frac{1}{-x-1}\right) dx + \frac{1}{2} x^2 \log\left(\frac{1}{x} + 1\right) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} x^2 \log\left(\frac{1}{x} + 1\right) + \frac{1}{2} (x - \log(x+1))
 \end{aligned}$$

input `Int[x*Log[1 + x^(-1)],x]`

output `(x^2*Log[1 + x^(-1)])/2 + (x - Log[1 + x])/2`

3.4.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.)), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 772 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Int[x^(n*p)*(b + a/x^n)^p, x] /; FreeQ[{a, b}, x] && ILtQ[n, 0] && IntegerQ[p]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2905 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_))^(p_.)]*(b_.))*((f_.)*(x_)^(m_.), x_Symbol] := Simp[(f*x)^(m + 1)*((a + b*Log[c*(d + e*x^n)^p])/(f*(m + 1))), x] - Simp[b*e*n*(p/(f*(m + 1))) Int[x^(n - 1)*((f*x)^(m + 1)/(d + e*x^n)), x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && NeQ[m, -1]`

3.4.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

method	result	size
risch	$\frac{x}{2} + \frac{x^2 \ln(1+\frac{1}{x})}{2} - \frac{\ln(1+x)}{2}$	22
parts	$\frac{x}{2} + \frac{x^2 \ln(1+\frac{1}{x})}{2} - \frac{\ln(1+x)}{2}$	22
derivativedivides	$\frac{x}{2} + \frac{\ln(\frac{1}{x})}{2} - \frac{\ln(1+\frac{1}{x})(1+\frac{1}{x})(\frac{1}{x}-1)x^2}{2}$	32
default	$\frac{x}{2} + \frac{\ln(\frac{1}{x})}{2} - \frac{\ln(1+\frac{1}{x})(1+\frac{1}{x})(\frac{1}{x}-1)x^2}{2}$	32
parallelrisch	$\frac{\ln(\frac{1+x}{x})x^2}{2} - \frac{1}{2} - \frac{\ln(x)}{2} + \frac{x}{2} - \frac{\ln(\frac{1+x}{x})}{2}$	33

input `int(x*ln(1+1/x),x,method=_RETURNVERBOSE)`

output `1/2*x+1/2*x^2*ln(1+1/x)-1/2*ln(1+x)`

3.4.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int x \log\left(1 + \frac{1}{x}\right) dx = \frac{1}{2} x^2 \log\left(\frac{x+1}{x}\right) + \frac{1}{2} x - \frac{1}{2} \log(x+1)$$

input `integrate(x*log(1+1/x),x, algorithm="fricas")`

output `1/2*x^2*log((x + 1)/x) + 1/2*x - 1/2*log(x + 1)`

3.4.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

$$\int x \log \left(1 + \frac{1}{x} \right) dx = \frac{x^2 \log \left(1 + \frac{1}{x} \right)}{2} + \frac{x}{2} - \frac{\log(x+1)}{2}$$

input `integrate(x*ln(1+1/x),x)`

output `x**2*log(1 + 1/x)/2 + x/2 - log(x + 1)/2`

3.4.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int x \log \left(1 + \frac{1}{x} \right) dx = \frac{1}{2} x^2 \log \left(\frac{1}{x} + 1 \right) + \frac{1}{2} x - \frac{1}{2} \log(x+1)$$

input `integrate(x*log(1+1/x),x, algorithm="maxima")`

output `1/2*x^2*log(1/x + 1) + 1/2*x - 1/2*log(x + 1)`

3.4.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. $2(21) = 42$.

Time = 0.27 (sec) , antiderivative size = 60, normalized size of antiderivative = 2.22

$$\int x \log \left(1 + \frac{1}{x} \right) dx = \frac{1}{2 \left(\frac{x+1}{x} - 1 \right)} + \frac{\log \left(\frac{x+1}{x} \right)}{2 \left(\frac{x+1}{x} - 1 \right)^2} - \frac{1}{2} \log \left(\frac{|x+1|}{|x|} \right) + \frac{1}{2} \log \left(\left| \frac{x+1}{x} - 1 \right| \right)$$

input `integrate(x*log(1+1/x),x, algorithm="giac")`

output `1/2/((x + 1)/x - 1) + 1/2*log((x + 1)/x)/((x + 1)/x - 1)^2 - 1/2*log(abs(x + 1)/abs(x)) + 1/2*log(abs((x + 1)/x - 1))`

3.4.9 Mupad [B] (verification not implemented)

Time = 16.96 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.15

$$\int x \log \left(1 + \frac{1}{x} \right) dx = \frac{x}{2} - \frac{\ln(x(x+1))}{4} - \frac{\ln\left(\frac{1}{x} + 1\right)}{4} + \frac{x^2 \ln\left(\frac{1}{x} + 1\right)}{2}$$

input `int(x*log(1/x + 1),x)`

output `x/2 - log(x*(x + 1))/4 - log(1/x + 1)/4 + (x^2*log(1/x + 1))/2`

3.5 $\int \sin^2(\log(x)) dx$

3.5.1	Optimal result	146
3.5.2	Mathematica [A] (verified)	146
3.5.3	Rubi [A] (verified)	147
3.5.4	Maple [A] (verified)	148
3.5.5	Fricas [A] (verification not implemented)	148
3.5.6	Sympy [A] (verification not implemented)	148
3.5.7	Maxima [A] (verification not implemented)	149
3.5.8	Giac [A] (verification not implemented)	149
3.5.9	Mupad [B] (verification not implemented)	149

3.5.1 Optimal result

Integrand size = 5, antiderivative size = 27

$$\int \sin^2(\log(x)) dx = \frac{2x}{5} - \frac{2}{5}x \cos(\log(x)) \sin(\log(x)) + \frac{1}{5}x \sin^2(\log(x))$$

output `2/5*x-2/5*x*cos(ln(x))*sin(ln(x))+1/5*x*sin(ln(x))^2`

3.5.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \sin^2(\log(x)) dx = \frac{x}{2} - \frac{1}{10}x \cos(2 \log(x)) - \frac{1}{5}x \sin(2 \log(x))$$

input `Integrate[Sin[Log[x]]^2,x]`

output `x/2 - (x*Cos[2*Log[x]])/10 - (x*Sin[2*Log[x]])/5`

3.5.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {4980, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sin^2(\log(x)) dx$$

$$\downarrow 4980$$

$$\frac{2 \int 1 dx}{5} + \frac{1}{5} x \sin^2(\log(x)) - \frac{2}{5} x \sin(\log(x)) \cos(\log(x))$$

$$\downarrow 24$$

$$\frac{2x}{5} + \frac{1}{5} x \sin^2(\log(x)) - \frac{2}{5} x \sin(\log(x)) \cos(\log(x))$$

input `Int[Sin[Log[x]]^2,x]`

output `(2*x)/5 - (2*x*Cos[Log[x]]*Sin[Log[x]])/5 + (x*Sin[Log[x]]^2)/5`

3.5.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 4980 `Int[Sin[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)]^(p_), x_Symbol] := Simp[x*(Sin[d*(a + b*Log[c*x^n])]^p/(b^2*d^2*n^2*p^2 + 1)), x] + (-Simp[b*d*n*p*x*Cos[d*(a + b*Log[c*x^n])]*(Sin[d*(a + b*Log[c*x^n])]^(p - 1)/(b^2*d^2*n^2*p^2 + 1)), x] + Simp[b^2*d^2*n^2*p*((p - 1)/(b^2*d^2*n^2*p^2 + 1)) Int[Sin[d*(a + b*Log[c*x^n])]^(p - 2), x], x]) /; FreeQ[{a, b, c, d, n}, x] && IGtQ[p, 1] && NeQ[b^2*d^2*n^2*p^2 + 1, 0]`

3.5.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.67

method	result	size
parallelrisch	$-\frac{x(-5+\cos(2\ln(x))+2\sin(2\ln(x)))}{10}$	18
default	$\frac{(\sin(\ln(x))-2\cos(\ln(x)))x\sin(\ln(x))}{5} + \frac{2x}{5}$	20
risch	$\frac{x}{2} + \left(-\frac{1}{20} + \frac{i}{10}\right) x x^{2i} + \left(-\frac{1}{20} - \frac{i}{10}\right) x x^{-2i}$	27
norman	$\frac{\frac{2x}{5} - \frac{4x \tan\left(\frac{\ln(x)}{2}\right)}{5} + \frac{8x \tan\left(\frac{\ln(x)}{2}\right)^2}{5} + \frac{4x \tan\left(\frac{\ln(x)}{2}\right)^3}{5} + \frac{2x \tan\left(\frac{\ln(x)}{2}\right)^4}{5}}{\left(1 + \tan\left(\frac{\ln(x)}{2}\right)^2\right)^2}$	55

input `int(sin(ln(x))^2,x,method=_RETURNVERBOSE)`

output `-1/10*x*(-5+cos(2*ln(x))+2*sin(2*ln(x)))`

3.5.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int \sin^2(\log(x)) dx = -\frac{1}{5} x \cos(\log(x))^2 - \frac{2}{5} x \cos(\log(x)) \sin(\log(x)) + \frac{3}{5} x$$

input `integrate(sin(log(x))^2,x, algorithm="fracas")`

output `-1/5*x*cos(log(x))^2 - 2/5*x*cos(log(x))*sin(log(x)) + 3/5*x`

3.5.6 Sympy [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.37

$$\int \sin^2(\log(x)) dx = \frac{3x \sin^2(\log(x))}{5} - \frac{2x \sin(\log(x)) \cos(\log(x))}{5} + \frac{2x \cos^2(\log(x))}{5}$$

input `integrate(sin(ln(x))**2,x)`

output `3*x*sin(log(x))**2/5 - 2*x*sin(log(x))*cos(log(x))/5 + 2*x*cos(log(x))**2/5`

3.5.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

$$\int \sin^2(\log(x)) dx = -\frac{1}{10} x \cos(2 \log(x)) - \frac{1}{5} x \sin(2 \log(x)) + \frac{1}{2} x$$

input `integrate(sin(log(x))^2,x, algorithm="maxima")`output `-1/10*x*cos(2*log(x)) - 1/5*x*sin(2*log(x)) + 1/2*x`**3.5.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int \sin^2(\log(x)) dx = -\frac{1}{5} x \cos(\log(x))^2 - \frac{2}{5} x \cos(\log(x)) \sin(\log(x)) + \frac{3}{5} x$$

input `integrate(sin(log(x))^2,x, algorithm="giac")`output `-1/5*x*cos(log(x))^2 - 2/5*x*cos(log(x))*sin(log(x)) + 3/5*x`**3.5.9 Mupad [B] (verification not implemented)**

Time = 16.20 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

$$\int \sin^2(\log(x)) dx = \frac{x}{2} + \frac{x(2 \sin(\ln(x))^2 - 1)}{10} - \frac{x \sin(2 \ln(x))}{5}$$

input `int(sin(log(x))^2,x)`output `x/2 + (x*(2*sin(log(x))^2 - 1))/10 - (x*sin(2*log(x)))/5`

3.6 $\int \frac{1}{1+3e^x} dx$

3.6.1	Optimal result	150
3.6.2	Mathematica [A] (verified)	150
3.6.3	Rubi [A] (verified)	151
3.6.4	Maple [A] (verified)	152
3.6.5	Fricas [A] (verification not implemented)	152
3.6.6	Sympy [A] (verification not implemented)	153
3.6.7	Maxima [A] (verification not implemented)	153
3.6.8	Giac [A] (verification not implemented)	153
3.6.9	Mupad [B] (verification not implemented)	154

3.6.1 Optimal result

Integrand size = 9, antiderivative size = 12

$$\int \frac{1}{1+3e^x} dx = x - \log(1+3e^x)$$

output `x-ln(1+3*exp(x))`

3.6.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.25

$$\int \frac{1}{1+3e^x} dx = \log(e^x) - \log(1+3e^x)$$

input `Integrate[(1 + 3*E^x)^(-1),x]`

output `Log[E^x] - Log[1 + 3*E^x]`

3.6.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.25, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {2720, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{1}{3e^x + 1} dx \\
 \downarrow 2720 \\
 \int \frac{e^{-x}}{3e^x + 1} de^x \\
 \downarrow 47 \\
 \int e^{-x} de^x - 3 \int \frac{1}{1 + 3e^x} de^x \\
 \downarrow 14 \\
 \log(e^x) - 3 \int \frac{1}{1 + 3e^x} de^x \\
 \downarrow 16 \\
 \log(e^x) - \log(3e^x + 1)
 \end{array}$$

input `Int[(1 + 3*E^x)^(-1),x]`

output `Log[E^x] - Log[1 + 3*E^x]`

3.6.3.1 Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`


```
rule 47 Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c
- a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x
], x] /; FreeQ[{a, b, c, d}, x]
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.6.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

method	result	size
risch	$x - \ln\left(\frac{1}{3} + e^x\right)$	10
parallelrisch	$x - \ln\left(\frac{1}{3} + e^x\right)$	10
norman	$x - \ln(1 + 3e^x)$	12
derivativedivides	$-\ln(1 + 3e^x) + \ln(e^x)$	14
default	$-\ln(1 + 3e^x) + \ln(e^x)$	14

```
input int(1/(1+3*exp(x)),x,method=_RETURNVERBOSE)
```

```
output x-ln(1/3+exp(x))
```

3.6.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

$$\int \frac{1}{1+3e^x} dx = x - \log(3e^x + 1)$$

```
input integrate(1/(1+3*exp(x)),x, algorithm="fricas")
```

```
output x - log(3*e^x + 1)
```

3.6.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{1+3e^x} dx = x - \log\left(e^x + \frac{1}{3}\right)$$

input `integrate(1/(1+3*exp(x)),x)`

output `x - log(exp(x) + 1/3)`

3.6.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

$$\int \frac{1}{1+3e^x} dx = x - \log(3e^x + 1)$$

input `integrate(1/(1+3*exp(x)),x, algorithm="maxima")`

output `x - log(3*e^x + 1)`

3.6.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

$$\int \frac{1}{1+3e^x} dx = x - \log(3e^x + 1)$$

input `integrate(1/(1+3*exp(x)),x, algorithm="giac")`

output `x - log(3*e^x + 1)`

3.6.9 Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

$$\int \frac{1}{1+3e^x} dx = x - \ln(3e^x + 1)$$

input `int(1/(3*exp(x) + 1),x)`

output `x - log(3*exp(x) + 1)`

3.7 $\int \csc^3(x) \sec^5(x) dx$

3.7.1	Optimal result	155
3.7.2	Mathematica [A] (verified)	155
3.7.3	Rubi [A] (warning: unable to verify)	156
3.7.4	Maple [A] (verified)	157
3.7.5	Fricas [B] (verification not implemented)	158
3.7.6	Sympy [A] (verification not implemented)	158
3.7.7	Maxima [B] (verification not implemented)	158
3.7.8	Giac [A] (verification not implemented)	159
3.7.9	Mupad [B] (verification not implemented)	159

3.7.1 Optimal result

Integrand size = 9, antiderivative size = 30

$$\int \csc^3(x) \sec^5(x) dx = -\frac{1}{2} \cot^2(x) + 3 \log(\tan(x)) + \frac{3 \tan^2(x)}{2} + \frac{\tan^4(x)}{4}$$

output `-1/2*cot(x)^2+3*ln(tan(x))+3/2*tan(x)^2+1/4*tan(x)^4`

3.7.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.03

$$\int \csc^3(x) \sec^5(x) dx = -\frac{1}{2} \csc^2(x) - 3 \log(\cos(x)) + 3 \log(\sin(x)) + \sec^2(x) + \frac{\sec^4(x)}{4}$$

input `Integrate[Csc[x]^3*Sec[x]^5,x]`

output `-1/2*Csc[x]^2 - 3*Log[Cos[x]] + 3*Log[Sin[x]] + Sec[x]^2 + Sec[x]^4/4`

3.7.3 Rubi [A] (warning: unable to verify)

Time = 0.19 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 3100, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \csc^3(x) \sec^5(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc(x)^3 \sec(x)^5 dx \\
 & \quad \downarrow \text{3100} \\
 & \int (\tan^2(x) + 1)^3 \cot^3(x) d \tan(x) \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \int \cot^2(x) (\tan^2(x) + 1)^3 d \tan^2(x) \\
 & \quad \downarrow \text{49} \\
 & \frac{1}{2} \int (\cot^2(x) + 3 \cot(x) + \tan^2(x) + 3) d \tan^2(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} \left(\frac{\tan^4(x)}{2} + 3 \tan^2(x) - \cot(x) + 3 \log(\tan^2(x)) \right)
 \end{aligned}$$

input `Int[Csc[x]^3*Sec[x]^5,x]`

output `(-Cot[x] + 3*Log[Tan[x]^2] + 3*Tan[x]^2 + Tan[x]^4/2)/2`

3.7.3.1 Defintions of rubi rules used

- rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int [ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

- rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int t[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

- rule 3100 `Int[csc[(e_.) + (f_.)*(x_)]^(m_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[1/f Subst[Int[(1 + x^2)^((m + n)/2 - 1)/x^m, x], x, Tan[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n)/2]`

3.7.4 Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.10

method	result
default	$\frac{1}{4 \sin(x)^2 \cos(x)^4} + \frac{3}{4 \sin(x)^2 \cos(x)^2} - \frac{3}{2 \sin(x)^2} + 3 \ln(\tan(x))$
parallelrisch	$-3 \ln(1 + \csc(x) - \cot(x)) - 3 \ln(\csc(x) - \cot(x) - 1) + 3 \ln(-\cot(x) + \csc(x)) + \frac{(8 \operatorname{se} \dots}{\dots}$
norman	$\frac{-\frac{1}{8} - 10 \tan(\frac{x}{2})^6 + \frac{57 \tan(\frac{x}{2})^4}{8} + \frac{57 \tan(\frac{x}{2})^8}{8} - \frac{\tan(\frac{x}{2})^{12}}{8}}{\tan(\frac{x}{2})^2 (\tan(\frac{x}{2})^2 - 1)^4} + 3 \ln(\tan(\frac{x}{2})) - 3 \ln(\tan(\frac{x}{2}) - 1) - 3 \ln(1 + \tan(\frac{x}{2}))$
risch	$\frac{6 e^{10ix} + 12 e^{8ix} - 4 e^{6ix} + 12 e^{4ix} + 6 e^{2ix}}{(e^{2ix} + 1)^4 (e^{2ix} - 1)^2} + 3 \ln(e^{2ix} - 1) - 3 \ln(e^{2ix} + 1)$

input `int(1/sin(x)^3/cos(x)^5,x,method=_RETURNVERBOSE)`

output `1/4/sin(x)^2/cos(x)^4+3/4/sin(x)^2/cos(x)^2-3/2/sin(x)^2+3*ln(tan(x))`

3.7.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 69 vs. $2(24) = 48$.

Time = 0.27 (sec) , antiderivative size = 69, normalized size of antiderivative = 2.30

$$\int \csc^3(x) \sec^5(x) dx = \frac{6 \cos(x)^4 - 3 \cos(x)^2 - 6 (\cos(x)^6 - \cos(x)^4) \log(\cos(x)^2) + 6 (\cos(x)^6 - \cos(x)^4) \log(-\frac{1}{4} \cos(x)^2 + 1)}{4 (\cos(x)^6 - \cos(x)^4)}$$

input `integrate(1/sin(x)^3/cos(x)^5,x, algorithm="fricas")`

output `1/4*(6*cos(x)^4 - 3*cos(x)^2 - 6*(cos(x)^6 - cos(x)^4)*log(cos(x)^2) + 6*(cos(x)^6 - cos(x)^4)*log(-1/4*cos(x)^2 + 1/4) - 1)/(cos(x)^6 - cos(x)^4)`

3.7.6 Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.53

$$\int \csc^3(x) \sec^5(x) dx = \frac{3 \log(\cos^2(x) - 1)}{2} - 3 \log(\cos(x)) - \frac{-6 \cos^4(x) + 3 \cos^2(x) + 1}{4 \cos^6(x) - 4 \cos^4(x)}$$

input `integrate(1/sin(x)**3/cos(x)**5,x)`

output `3*log(cos(x)**2 - 1)/2 - 3*log(cos(x)) - (-6*cos(x)**4 + 3*cos(x)**2 + 1)/(4*cos(x)**6 - 4*cos(x)**4)`

3.7.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 50 vs. $2(24) = 48$.

Time = 0.20 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.67

$$\int \csc^3(x) \sec^5(x) dx = -\frac{6 \sin(x)^4 - 9 \sin(x)^2 + 2}{4 (\sin(x)^6 - 2 \sin(x)^4 + \sin(x)^2)} - \frac{3}{2} \log(\sin(x)^2 - 1) + \frac{3}{2} \log(\sin(x)^2)$$

input `integrate(1/sin(x)^3/cos(x)^5,x, algorithm="maxima")`

output `-1/4*(6*sin(x)^4 - 9*sin(x)^2 + 2)/(sin(x)^6 - 2*sin(x)^4 + sin(x)^2) - 3/2*log(sin(x)^2 - 1) + 3/2*log(sin(x)^2)`

3.7.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.53

$$\int \csc^3(x) \sec^5(x) dx = \frac{6 \cos(x)^4 - 3 \cos(x)^2 - 1}{4 (\cos(x)^2 - 1) \cos(x)^4} + \frac{3}{2} \log(-\cos(x)^2 + 1) - 3 \log(|\cos(x)|)$$

input `integrate(1/sin(x)^3/cos(x)^5,x, algorithm="giac")`

output `1/4*(6*cos(x)^4 - 3*cos(x)^2 - 1)/((cos(x)^2 - 1)*cos(x)^4) + 3/2*log(-cos(x)^2 + 1) - 3*log(abs(cos(x)))`

3.7.9 Mupad [B] (verification not implemented)

Time = 15.25 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \csc^3(x) \sec^5(x) dx = 3 \ln(\tan(x)) + \frac{\frac{3}{4 \cos(x)^2} + \frac{1}{4 \cos(x)^4}}{\sin(x)^2} - \frac{3}{2 \sin(x)^2}$$

input `int(1/(cos(x)^5*sin(x)^3),x)`

output `3*log(tan(x)) + (3/(4*cos(x)^2) + 1/(4*cos(x)^4))/sin(x)^2 - 3/(2*sin(x)^2)`

3.8 $\int \frac{1}{x\sqrt{-1+x^4}} dx$

3.8.1	Optimal result	160
3.8.2	Mathematica [A] (verified)	160
3.8.3	Rubi [A] (verified)	161
3.8.4	Maple [A] (verified)	162
3.8.5	Fricas [A] (verification not implemented)	162
3.8.6	Sympy [C] (verification not implemented)	163
3.8.7	Maxima [A] (verification not implemented)	163
3.8.8	Giac [A] (verification not implemented)	163
3.8.9	Mupad [B] (verification not implemented)	164

3.8.1 Optimal result

Integrand size = 13, antiderivative size = 14

$$\int \frac{1}{x\sqrt{-1+x^4}} dx = \frac{1}{2} \arctan(\sqrt{-1+x^4})$$

output `1/2*arctan((x^4-1)^(1/2))`

3.8.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{-1+x^4}} dx = \frac{1}{2} \arctan(\sqrt{-1+x^4})$$

input `Integrate[1/(x*Sqrt[-1 + x^4]),x]`

output `ArcTan[Sqrt[-1 + x^4]]/2`

3.8.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {798, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x\sqrt{x^4-1}} dx \\ & \quad \downarrow \text{798} \\ & \frac{1}{4} \int \frac{1}{x^4\sqrt{x^4-1}} dx^4 \\ & \quad \downarrow \text{73} \\ & \frac{1}{2} \int \frac{1}{x^8+1} d\sqrt{x^4-1} \\ & \quad \downarrow \text{216} \\ & \frac{1}{2} \arctan(\sqrt{x^4-1}) \end{aligned}$$

input `Int[1/(x*sqrt[-1 + x^4]),x]`

output `ArcTan[Sqrt[-1 + x^4]]/2`

3.8.3.1 Defintions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

```
rule 798 Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst
[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a,
b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

3.8.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

method	result	size
default	$-\frac{\arctan\left(\frac{1}{\sqrt{x^4-1}}\right)}{2}$	11
elliptic	$-\frac{\arctan\left(\frac{1}{\sqrt{x^4-1}}\right)}{2}$	11
pseudoelliptic	$-\frac{\arctan\left(\frac{1}{\sqrt{x^4-1}}\right)}{2}$	11
trager	$\frac{\text{RootOf}(-Z^2+1) \ln\left(\frac{\text{RootOf}(-Z^2+1)+\sqrt{x^4-1}}{x^2}\right)}{2}$	28
meijerg	$\frac{\sqrt{-\text{signum}(x^4-1)}\left((-2\ln(2)+4\ln(x)+i\pi)\sqrt{\pi}-2\sqrt{\pi}\ln\left(\frac{1}{2}+\frac{\sqrt{-x^4+1}}{2}\right)\right)}{4\sqrt{\pi}\sqrt{\text{signum}(x^4-1)}}$	61

```
input int(1/x/(x^4-1)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/2*arctan(1/(x^4-1)^(1/2))
```

3.8.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{1}{x\sqrt{-1+x^4}} dx = \frac{1}{2} \arctan\left(\sqrt{x^4-1}\right)$$

```
input integrate(1/x/(x^4-1)^(1/2),x, algorithm="fricas")
```

```
output 1/2*arctan(sqrt(x^4 - 1))
```

3.8.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.44 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.71

$$\int \frac{1}{x\sqrt{-1+x^4}} dx = \begin{cases} \frac{i \operatorname{acosh}\left(\frac{1}{x^2}\right)}{2} & \text{for } \frac{1}{|x^4|} > 1 \\ -\frac{\operatorname{asin}\left(\frac{1}{x^2}\right)}{2} & \text{otherwise} \end{cases}$$

input `integrate(1/x/(x**4-1)**(1/2),x)`

output `Piecewise((I*acosh(x**(-2))/2, 1/Abs(x**4) > 1), (-asin(x**(-2))/2, True))`

3.8.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{1}{x\sqrt{-1+x^4}} dx = \frac{1}{2} \arctan\left(\sqrt{x^4-1}\right)$$

input `integrate(1/x/(x^4-1)^(1/2),x, algorithm="maxima")`

output `1/2*arctan(sqrt(x^4 - 1))`

3.8.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{1}{x\sqrt{-1+x^4}} dx = \frac{1}{2} \arctan\left(\sqrt{x^4-1}\right)$$

input `integrate(1/x/(x^4-1)^(1/2),x, algorithm="giac")`

output `1/2*arctan(sqrt(x^4 - 1))`

3.8.9 Mupad [B] (verification not implemented)

Time = 15.31 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{1}{x\sqrt{-1+x^4}} dx = \frac{\operatorname{atan}(\sqrt{x^4-1})}{2}$$

input `int(1/(x*(x^4 - 1)^(1/2)),x)`

output `atan((x^4 - 1)^(1/2))/2`

3.9 $\int \frac{1}{x(1+x^5)} dx$

3.9.1	Optimal result	165
3.9.2	Mathematica [A] (verified)	165
3.9.3	Rubi [A] (verified)	166
3.9.4	Maple [A] (verified)	167
3.9.5	Fricas [A] (verification not implemented)	167
3.9.6	Sympy [A] (verification not implemented)	168
3.9.7	Maxima [A] (verification not implemented)	168
3.9.8	Giac [A] (verification not implemented)	168
3.9.9	Mupad [B] (verification not implemented)	169

3.9.1 Optimal result

Integrand size = 11, antiderivative size = 13

$$\int \frac{1}{x(1+x^5)} dx = \log(x) - \frac{1}{5} \log(1+x^5)$$

output `ln(x)-1/5*ln(x^5+1)`

3.9.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+x^5)} dx = \log(x) - \frac{1}{5} \log(1+x^5)$$

input `Integrate[1/(x*(1 + x^5)),x]`

output `Log[x] - Log[1 + x^5]/5`

3.9.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {798, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x(x^5+1)} dx \\
 & \quad \downarrow 798 \\
 & \frac{1}{5} \int \frac{1}{x^5(x^5+1)} dx^5 \\
 & \quad \downarrow 47 \\
 & \frac{1}{5} \left(\int \frac{1}{x^5} dx^5 - \int \frac{1}{x^5+1} dx^5 \right) \\
 & \quad \downarrow 14 \\
 & \frac{1}{5} \left(\log(x^5) - \int \frac{1}{x^5+1} dx^5 \right) \\
 & \quad \downarrow 16 \\
 & \frac{1}{5} (\log(x^5) - \log(x^5+1))
 \end{aligned}$$

input `Int[1/(x*(1 + x^5)),x]`

output `(Log[x^5] - Log[1 + x^5])/5`

3.9.3.1 Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.)), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^(p_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

3.9.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
meijerg	$\ln(x) - \frac{\ln(x^5+1)}{5}$	12
risch	$\ln(x) - \frac{\ln(x^5+1)}{5}$	12
default	$-\frac{\ln(1+x)}{5} - \frac{\ln(x^4-x^3+x^2-x+1)}{5} + \ln(x)$	29
norman	$-\frac{\ln(1+x)}{5} - \frac{\ln(x^4-x^3+x^2-x+1)}{5} + \ln(x)$	29
parallelrisch	$-\frac{\ln(1+x)}{5} - \frac{\ln(x^4-x^3+x^2-x+1)}{5} + \ln(x)$	29

input `int(1/x/(x^5+1),x,method=_RETURNVERBOSE)`

output `ln(x)-1/5*ln(x^5+1)`

3.9.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1}{x(1+x^5)} dx = -\frac{1}{5} \log(x^5 + 1) + \log(x)$$

input `integrate(1/x/(x^5+1),x, algorithm="fricas")`

output `-1/5*log(x^5 + 1) + log(x)`

3.9.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \frac{1}{x(1+x^5)} dx = \log(x) - \frac{\log(x^5+1)}{5}$$

input `integrate(1/x/(x**5+1),x)`output `log(x) - log(x**5 + 1)/5`**3.9.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{1}{x(1+x^5)} dx = -\frac{1}{5} \log(x^5+1) + \frac{1}{5} \log(x^5)$$

input `integrate(1/x/(x^5+1),x, algorithm="maxima")`output `-1/5*log(x^5 + 1) + 1/5*log(x^5)`**3.9.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+x^5)} dx = -\frac{1}{5} \log(|x^5+1|) + \log(|x|)$$

input `integrate(1/x/(x^5+1),x, algorithm="giac")`output `-1/5*log(abs(x^5 + 1)) + log(abs(x))`

3.9.9 Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1}{x(1+x^5)} dx = \ln(x) - \frac{\ln(x^5+1)}{5}$$

input `int(1/(x*(x^5 + 1)),x)`

output `log(x) - log(x^5 + 1)/5`

3.10 $\int \sqrt{\tan(x)} dx$

3.10.1	Optimal result	170
3.10.2	Mathematica [A] (verified)	170
3.10.3	Rubi [A] (verified)	171
3.10.4	Maple [A] (verified)	174
3.10.5	Fricas [C] (verification not implemented)	174
3.10.6	Sympy [F]	175
3.10.7	Maxima [A] (verification not implemented)	175
3.10.8	Giac [A] (verification not implemented)	176
3.10.9	Mupad [B] (verification not implemented)	176

3.10.1 Optimal result

Integrand size = 6, antiderivative size = 71

$$\int \sqrt{\tan(x)} dx = -\frac{\arctan\left(1 - \sqrt{2}\sqrt{\tan(x)}\right)}{\sqrt{2}} + \frac{\arctan\left(1 + \sqrt{2}\sqrt{\tan(x)}\right)}{\sqrt{2}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{\tan(x)}}{1+\tan(x)}\right)}{\sqrt{2}}$$

```
output 1/2*arctan(-1+2^(1/2)*tan(x)^(1/2))*2^(1/2)+1/2*arctan(1+2^(1/2)*tan(x)^(1/2))*2^(1/2)-1/2*arctanh(2^(1/2)*tan(x)^(1/2)/(1+tan(x)))*2^(1/2)
```

3.10.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.56

$$\int \sqrt{\tan(x)} dx = \frac{\left(\arctan\left(\sqrt[4]{-\tan^2(x)}\right) - \operatorname{arctanh}\left(\sqrt[4]{-\tan^2(x)}\right)\right) \sqrt[4]{-\tan(x)}}{\sqrt[4]{\tan(x)}}$$

```
input Integrate[Sqrt[Tan[x]],x]
```

```
output ((ArcTan[(-Tan[x]^2)^(1/4)] - ArcTanh[(-Tan[x]^2)^(1/4)])*(-Tan[x])^(1/4))/Tan[x]^(1/4)
```

3.10.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.55, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 1.833$, Rules used = {3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\tan(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\tan(x)} dx \\
 & \quad \downarrow \text{3957} \\
 & \int \frac{\sqrt{\tan(x)}}{\tan^2(x) + 1} d \tan(x) \\
 & \quad \downarrow \text{266} \\
 & 2 \int \frac{\tan(x)}{\tan^2(x) + 1} d \sqrt{\tan(x)} \\
 & \quad \downarrow \text{826} \\
 & 2 \left(\frac{1}{2} \int \frac{\tan(x) + 1}{\tan^2(x) + 1} d \sqrt{\tan(x)} - \frac{1}{2} \int \frac{1 - \tan(x)}{\tan^2(x) + 1} d \sqrt{\tan(x)} \right) \\
 & \quad \downarrow \text{1476} \\
 & 2 \left(\frac{1}{2} \left(\frac{1}{2} \int \frac{1}{\tan(x) - \sqrt{2}\sqrt{\tan(x)} + 1} d \sqrt{\tan(x)} + \frac{1}{2} \int \frac{1}{\tan(x) + \sqrt{2}\sqrt{\tan(x)} + 1} d \sqrt{\tan(x)} \right) - \frac{1}{2} \int \frac{1 - \tan(x)}{\tan^2(x) + 1} d \sqrt{\tan(x)} \right) \\
 & \quad \downarrow \text{1082} \\
 & 2 \left(\frac{1}{2} \left(\frac{\int \frac{1}{-\tan(x)-1} d(1 - \sqrt{2}\sqrt{\tan(x)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(x)-1} d(\sqrt{2}\sqrt{\tan(x)} + 1)}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1 - \tan(x)}{\tan^2(x) + 1} d \sqrt{\tan(x)} \right) \\
 & \quad \downarrow \text{217} \\
 & 2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(x)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(x)})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1 - \tan(x)}{\tan^2(x) + 1} d \sqrt{\tan(x)} \right) \\
 & \quad \downarrow \text{1479}
 \end{aligned}$$

$$\begin{aligned}
& 2 \left(\frac{1}{2} \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(x)}}{\tan(x)-\sqrt{2}\sqrt{\tan(x)+1}} d\sqrt{\tan(x)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(x)+1})}{\tan(x)+\sqrt{2}\sqrt{\tan(x)+1}} d\sqrt{\tan(x)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(x)+1})}{\sqrt{2}} - \arctan(1 - \sqrt{2}\sqrt{\tan(x)}) \right) \right) \\
& \quad \downarrow \text{25} \\
& 2 \left(\frac{1}{2} \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(x)}}{\tan(x)-\sqrt{2}\sqrt{\tan(x)+1}} d\sqrt{\tan(x)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(x)+1})}{\tan(x)+\sqrt{2}\sqrt{\tan(x)+1}} d\sqrt{\tan(x)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(x)+1})}{\sqrt{2}} - \arctan(1 - \sqrt{2}\sqrt{\tan(x)}) \right) \right) \\
& \quad \downarrow \text{27} \\
& 2 \left(\frac{1}{2} \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(x)}}{\tan(x)-\sqrt{2}\sqrt{\tan(x)+1}} d\sqrt{\tan(x)}}{2\sqrt{2}} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(x)+1}}{\tan(x)+\sqrt{2}\sqrt{\tan(x)+1}} d\sqrt{\tan(x)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(x)+1})}{\sqrt{2}} - \arctan(1 - \sqrt{2}\sqrt{\tan(x)}) \right) \right) \\
& \quad \downarrow \text{1103} \\
& 2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(x)+1})}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(x)})}{\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\log(\tan(x) - \sqrt{2}\sqrt{\tan(x)+1})}{2\sqrt{2}} - \frac{\log(\tan(x) + \sqrt{2}\sqrt{\tan(x)+1})}{2\sqrt{2}} \right) \right)
\end{aligned}$$

input `Int[Sqrt[Tan[x]], x]`

output `2*((-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[x]]]/Sqrt[2])/2 + (Log[1 - Sqrt[2]*Sqrt[Tan[x]] + Tan[x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Tan[x]] + Tan[x]]/(2*Sqrt[2]))/2)`

3.10.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_.)*(x_)^m)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x])/b], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3957 Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int
[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] &&
!IntegerQ[n]
```

3.10.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.69

method	result	size
lookup	$\frac{\sqrt{\tan(x)} \cos(x) \sqrt{2} \arccos(\cos(x) - \sin(x))}{2\sqrt{\cos(x) \sin(x)}} - \frac{\sqrt{2} \ln(\cos(x) + \sqrt{2} \sqrt{\tan(x)} \cos(x) + \sin(x))}{2}$	49
default	$\frac{\sqrt{\tan(x)} \cos(x) \sqrt{2} \arccos(\cos(x) - \sin(x))}{2\sqrt{\cos(x) \sin(x)}} - \frac{\sqrt{2} \ln(\cos(x) + \sqrt{2} \sqrt{\tan(x)} \cos(x) + \sin(x))}{2}$	49
derivativedivides	$\frac{\sqrt{2} \left(\ln \left(\frac{\tan(x) - \sqrt{2} \sqrt{\tan(x) + 1}}{\tan(x) + \sqrt{2} \sqrt{\tan(x) + 1}} \right) + 2 \arctan(1 + \sqrt{2} \sqrt{\tan(x)}) + 2 \arctan(-1 + \sqrt{2} \sqrt{\tan(x)}) \right)}{4}$	62

```
input int(tan(x)^(1/2), x, method=_RETURNVERBOSE)
```

```
output 1/2*tan(x)^(1/2)/(cos(x)*sin(x))^(1/2)*cos(x)*2^(1/2)*arccos(cos(x)-sin(x))
)-1/2*2^(1/2)*ln(cos(x)+2^(1/2)*tan(x)^(1/2)*cos(x)+sin(x))
```

3.10.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.03

$$\begin{aligned} \int \sqrt{\tan(x)} dx &= \left(\frac{1}{4}i - \frac{1}{4}\right) \sqrt{2} \log\left((i+1)\sqrt{2} + 2\sqrt{\tan(x)}\right) \\ &\quad - \left(\frac{1}{4}i + \frac{1}{4}\right) \sqrt{2} \log\left(-(i-1)\sqrt{2} + 2\sqrt{\tan(x)}\right) \\ &\quad + \left(\frac{1}{4}i + \frac{1}{4}\right) \sqrt{2} \log\left((i-1)\sqrt{2} + 2\sqrt{\tan(x)}\right) \\ &\quad - \left(\frac{1}{4}i - \frac{1}{4}\right) \sqrt{2} \log\left(-(i+1)\sqrt{2} + 2\sqrt{\tan(x)}\right) \end{aligned}$$

input `integrate(tan(x)^(1/2),x, algorithm="fricas")`

output `(1/4*I - 1/4)*sqrt(2)*log((I + 1)*sqrt(2) + 2*sqrt(tan(x))) - (1/4*I + 1/4)*sqrt(2)*log(-(I - 1)*sqrt(2) + 2*sqrt(tan(x))) + (1/4*I + 1/4)*sqrt(2)*log((I - 1)*sqrt(2) + 2*sqrt(tan(x))) - (1/4*I - 1/4)*sqrt(2)*log(-(I + 1)*sqrt(2) + 2*sqrt(tan(x)))`

3.10.6 Sympy [F]

$$\int \sqrt{\tan(x)} dx = \int \sqrt{\tan(x)} dx$$

input `integrate(tan(x)**(1/2),x)`

output `Integral(sqrt(tan(x)), x)`

3.10.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.13

$$\begin{aligned} \int \sqrt{\tan(x)} dx = & \frac{1}{2} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} (\sqrt{2} + 2 \sqrt{\tan(x)}) \right) \\ & + \frac{1}{2} \sqrt{2} \arctan \left(-\frac{1}{2} \sqrt{2} (\sqrt{2} - 2 \sqrt{\tan(x)}) \right) \\ & - \frac{1}{4} \sqrt{2} \log \left(\sqrt{2} \sqrt{\tan(x)} + \tan(x) + 1 \right) \\ & + \frac{1}{4} \sqrt{2} \log \left(-\sqrt{2} \sqrt{\tan(x)} + \tan(x) + 1 \right) \end{aligned}$$

input `integrate(tan(x)^(1/2),x, algorithm="maxima")`

output `1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(x)))) + 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(x)))) - 1/4*sqrt(2)*log(sqrt(2)*sqrt(tan(x)) + tan(x) + 1) + 1/4*sqrt(2)*log(-sqrt(2)*sqrt(tan(x)) + tan(x) + 1)`

3.10.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.13

$$\int \sqrt{\tan(x)} dx = \frac{1}{2} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} (\sqrt{2} + 2 \sqrt{\tan(x)}) \right) + \frac{1}{2} \sqrt{2} \arctan \left(-\frac{1}{2} \sqrt{2} (\sqrt{2} - 2 \sqrt{\tan(x)}) \right) - \frac{1}{4} \sqrt{2} \log \left(\sqrt{2} \sqrt{\tan(x)} + \tan(x) + 1 \right) + \frac{1}{4} \sqrt{2} \log \left(-\sqrt{2} \sqrt{\tan(x)} + \tan(x) + 1 \right)$$

input `integrate(tan(x)^(1/2),x, algorithm="giac")`output `1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(x)))) + 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(x)))) - 1/4*sqrt(2)*log(sqrt(2)*sqrt(tan(x)) + tan(x) + 1) + 1/4*sqrt(2)*log(-sqrt(2)*sqrt(tan(x)) + tan(x) + 1)`**3.10.9 Mupad [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.92

$$\int \sqrt{\tan(x)} dx = \frac{\sqrt{2} \left(\ln \left(\sqrt{2} \sqrt{\tan(x)} - \tan(x) - 1 \right) - \ln \left(\tan(x) + \sqrt{2} \sqrt{\tan(x)} + 1 \right) \right)}{4} + \frac{\sqrt{2} \left(\operatorname{atan} \left(\sqrt{2} \sqrt{\tan(x)} - 1 \right) + \operatorname{atan} \left(\sqrt{2} \sqrt{\tan(x)} + 1 \right) \right)}{2}$$

input `int(tan(x)^(1/2),x)`output `(2^(1/2)*(log(2^(1/2)*tan(x)^(1/2) - tan(x) - 1) - log(tan(x) + 2^(1/2)*tan(x)^(1/2) + 1)))/4 + (2^(1/2)*(atan(2^(1/2)*tan(x)^(1/2) - 1) + atan(2^(1/2)*tan(x)^(1/2) + 1)))/2`

3.11 $\int \frac{\log(1+x)}{1+x^2} dx$

3.11.1	Optimal result	177
3.11.2	Mathematica [A] (verified)	177
3.11.3	Rubi [A] (verified)	178
3.11.4	Maple [A] (verified)	179
3.11.5	Fricas [F]	179
3.11.6	Sympy [F]	180
3.11.7	Maxima [A] (verification not implemented)	180
3.11.8	Giac [F]	180
3.11.9	Mupad [F(-1)]	181

3.11.1 Optimal result

Integrand size = 12, antiderivative size = 89

$$\int \frac{\log(1+x)}{1+x^2} dx = -\frac{1}{2}i \log\left(\left(\frac{1}{2} - \frac{i}{2}\right)(i-x)\right) \log(1+x) + \frac{1}{2}i \log\left(\left(-\frac{1}{2} - \frac{i}{2}\right)(i+x)\right) \log(1+x) - \frac{1}{2}i \operatorname{PolyLog}\left(2, \left(\frac{1}{2} - \frac{i}{2}\right)(1+x)\right) + \frac{1}{2}i \operatorname{PolyLog}\left(2, \left(\frac{1}{2} + \frac{i}{2}\right)(1+x)\right)$$

output `-1/2*I*ln((1/2-1/2*I)*(I-x))*ln(1+x)+1/2*I*ln((-1/2-1/2*I)*(I+x))*ln(1+x)-1/2*I*polylog(2,(1/2-1/2*I)*(1+x))+1/2*I*polylog(2,(1/2+1/2*I)*(1+x))`

3.11.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00

$$\int \frac{\log(1+x)}{1+x^2} dx = -\frac{1}{2}i \log\left(\left(\frac{1}{2} - \frac{i}{2}\right)(i-x)\right) \log(1+x) + \frac{1}{2}i \log\left(\left(-\frac{1}{2} - \frac{i}{2}\right)(i+x)\right) \log(1+x) - \frac{1}{2}i \operatorname{PolyLog}\left(2, \left(\frac{1}{2} - \frac{i}{2}\right)(1+x)\right) + \frac{1}{2}i \operatorname{PolyLog}\left(2, \left(\frac{1}{2} + \frac{i}{2}\right)(1+x)\right)$$

input `Integrate[Log[1 + x]/(1 + x^2),x]`

output `(-1/2*I)*Log[(1/2 - I/2)*(I - x)]*Log[1 + x] + (I/2)*Log[(-1/2 - I/2)*(I + x)]*Log[1 + x] - (I/2)*PolyLog[2, (1/2 - I/2)*(1 + x)] + (I/2)*PolyLog[2, (1/2 + I/2)*(1 + x)]`

3.11.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2856, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\log(x+1)}{x^2+1} dx$$

$$\downarrow \text{2856}$$

$$\int \left(\frac{i \log(x+1)}{2(-x+i)} + \frac{i \log(x+1)}{2(x+i)} \right) dx$$

$$\downarrow \text{2009}$$

$$-\frac{1}{2}i \text{PolyLog} \left(2, \left(\frac{1}{2} - \frac{i}{2} \right) (x+1) \right) + \frac{1}{2}i \text{PolyLog} \left(2, \left(\frac{1}{2} + \frac{i}{2} \right) (x+1) \right) - \frac{1}{2}i \log \left(\left(\frac{1}{2} - \frac{i}{2} \right) (-x+i) \right) \log(x+1) + \frac{1}{2}i \log \left(\left(-\frac{1}{2} - \frac{i}{2} \right) (x+i) \right) \log(x+1)$$

input `Int[Log[1 + x]/(1 + x^2),x]`

output `(-1/2*I)*Log[(1/2 - I/2)*(I - x)]*Log[1 + x] + (I/2)*Log[(-1/2 - I/2)*(I + x)]*Log[1 + x] - (I/2)*PolyLog[2, (1/2 - I/2)*(1 + x)] + (I/2)*PolyLog[2, (1/2 + I/2)*(1 + x)]`

3.11.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2856 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]*(b_.))^(p_.)*((f_) + (g_.)*(x_)^(r_.))^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log[c*(d + e*x)^n])^p, (f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, r}, x] && IntegerQ[p, 0] && IntegerQ[q] && (GtQ[q, 0] || (IntegerQ[r] && NeQ[r, 1]))`

3.11.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.79

method	result
derivativedivides	$-\frac{i \ln(1+x) \ln\left(\frac{1}{2}-\frac{x}{2}+\frac{i(1+x)}{2}\right)}{2} + \frac{i \ln(1+x) \ln\left(\frac{1}{2}-\frac{x}{2}-\frac{i(1+x)}{2}\right)}{2} - \frac{i \operatorname{dilog}\left(\frac{1}{2}-\frac{x}{2}+\frac{i(1+x)}{2}\right)}{2} + \frac{i \operatorname{dilog}\left(\frac{1}{2}-\frac{x}{2}-\frac{i(1+x)}{2}\right)}{2}$
default	$-\frac{i \ln(1+x) \ln\left(\frac{1}{2}-\frac{x}{2}+\frac{i(1+x)}{2}\right)}{2} + \frac{i \ln(1+x) \ln\left(\frac{1}{2}-\frac{x}{2}-\frac{i(1+x)}{2}\right)}{2} - \frac{i \operatorname{dilog}\left(\frac{1}{2}-\frac{x}{2}+\frac{i(1+x)}{2}\right)}{2} + \frac{i \operatorname{dilog}\left(\frac{1}{2}-\frac{x}{2}-\frac{i(1+x)}{2}\right)}{2}$
risch	$-\frac{i \ln(1+x) \ln\left(\frac{1}{2}-\frac{x}{2}+\frac{i(1+x)}{2}\right)}{2} + \frac{i \ln(1+x) \ln\left(\frac{1}{2}-\frac{x}{2}-\frac{i(1+x)}{2}\right)}{2} - \frac{i \operatorname{dilog}\left(\frac{1}{2}-\frac{x}{2}+\frac{i(1+x)}{2}\right)}{2} + \frac{i \operatorname{dilog}\left(\frac{1}{2}-\frac{x}{2}-\frac{i(1+x)}{2}\right)}{2}$
parts	$-\frac{i \ln(1+x) \ln\left(\frac{1}{2}-\frac{x}{2}+\frac{i(1+x)}{2}\right)}{2} + \frac{i \ln(1+x) \ln\left(\frac{1}{2}-\frac{x}{2}-\frac{i(1+x)}{2}\right)}{2} - \frac{i \operatorname{dilog}\left(\frac{1}{2}-\frac{x}{2}+\frac{i(1+x)}{2}\right)}{2} + \frac{i \operatorname{dilog}\left(\frac{1}{2}-\frac{x}{2}-\frac{i(1+x)}{2}\right)}{2}$

input `int(ln(1+x)/(x^2+1),x,method=_RETURNVERBOSE)`

output `-1/2*I*ln(1+x)*ln(1/2-1/2*x+1/2*I*(1+x))+1/2*I*ln(1+x)*ln(1/2-1/2*x-1/2*I*(1+x))-1/2*I*dilog(1/2-1/2*x+1/2*I*(1+x))+1/2*I*dilog(1/2-1/2*x-1/2*I*(1+x))`

3.11.5 Fracas [F]

$$\int \frac{\log(1+x)}{1+x^2} dx = \int \frac{\log(x+1)}{x^2+1} dx$$

input `integrate(log(1+x)/(x^2+1),x, algorithm="fracas")`

output `integral(log(x + 1)/(x^2 + 1), x)`

3.11.6 Sympy [F]

$$\int \frac{\log(1+x)}{1+x^2} dx = \int \frac{\log(x+1)}{x^2+1} dx$$

input `integrate(ln(1+x)/(x**2+1),x)`

output `Integral(log(x + 1)/(x**2 + 1), x)`

3.11.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.63

$$\begin{aligned} \int \frac{\log(1+x)}{1+x^2} dx &= \frac{1}{2} \arctan\left(\frac{1}{2}x + \frac{1}{2}, \frac{1}{2}x + \frac{1}{2}\right) \log(x^2+1) \\ &\quad - \frac{1}{2} \arctan(x) \log\left(\frac{1}{2}x^2 + x + \frac{1}{2}\right) + \arctan(x) \log(x+1) \\ &\quad + \frac{1}{2}i \operatorname{Li}_2\left(\left(\frac{1}{2}i - \frac{1}{2}\right)x + \frac{1}{2}i + \frac{1}{2}\right) - \frac{1}{2}i \operatorname{Li}_2\left(-\left(\frac{1}{2}i + \frac{1}{2}\right)x - \frac{1}{2}i + \frac{1}{2}\right) \end{aligned}$$

input `integrate(log(1+x)/(x^2+1),x, algorithm="maxima")`

output `1/2*arctan2(1/2*x + 1/2, 1/2*x + 1/2)*log(x^2 + 1) - 1/2*arctan(x)*log(1/2*x^2 + x + 1/2) + arctan(x)*log(x + 1) + 1/2*I*dilog((1/2*I - 1/2)*x + 1/2*I + 1/2) - 1/2*I*dilog(-(1/2*I + 1/2)*x - 1/2*I + 1/2)`

3.11.8 Giac [F]

$$\int \frac{\log(1+x)}{1+x^2} dx = \int \frac{\log(x+1)}{x^2+1} dx$$

input `integrate(log(1+x)/(x^2+1),x, algorithm="giac")`

output `integrate(log(x + 1)/(x^2 + 1), x)`

3.11.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\log(1+x)}{1+x^2} dx = \int \frac{\ln(x+1)}{x^2+1} dx$$

input `int(log(x + 1)/(x^2 + 1),x)`output `int(log(x + 1)/(x^2 + 1), x)`

3.12 $\int \frac{\sqrt{x}}{-\sqrt[3]{x} + \sqrt{x}} dx$

3.12.1	Optimal result	182
3.12.2	Mathematica [A] (verified)	182
3.12.3	Rubi [A] (verified)	183
3.12.4	Maple [A] (verified)	184
3.12.5	Fricas [A] (verification not implemented)	185
3.12.6	Sympy [F]	185
3.12.7	Maxima [A] (verification not implemented)	185
3.12.8	Giac [A] (verification not implemented)	186
3.12.9	Mupad [B] (verification not implemented)	186

3.12.1 Optimal result

Integrand size = 21, antiderivative size = 53

$$\int \frac{\sqrt{x}}{-\sqrt[3]{x} + \sqrt{x}} dx = 6\sqrt[6]{x} + 3\sqrt[3]{x} + 2\sqrt{x} + \frac{3x^{2/3}}{2} + \frac{6x^{5/6}}{5} + x + 6 \log(1 - \sqrt[6]{x})$$

output `6*x^(1/6)+3*x^(1/3)+2*x^(1/2)+3/2*x^(2/3)+6/5*x^(5/6)+x+6*ln(1-x^(1/6))`

3.12.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.96

$$\int \frac{\sqrt{x}}{-\sqrt[3]{x} + \sqrt{x}} dx = 6\sqrt[6]{x} + 3\sqrt[3]{x} + 2\sqrt{x} + \frac{3x^{2/3}}{2} + \frac{6x^{5/6}}{5} + x + 6 \log(-1 + \sqrt[6]{x})$$

input `Integrate[Sqrt[x]/(-x^(1/3) + Sqrt[x]),x]`

output `6*x^(1/6) + 3*x^(1/3) + 2*Sqrt[x] + (3*x^(2/3))/2 + (6*x^(5/6))/5 + x + 6*Log[-1 + x^(1/6)]`

3.12.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.19, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {10, 25, 798, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{x}}{\sqrt{x} - \sqrt[3]{x}} dx \\
 & \quad \downarrow 10 \\
 & \int -\frac{\sqrt[6]{x}}{1 - \sqrt[6]{x}} dx \\
 & \quad \downarrow 25 \\
 & -\int \frac{\sqrt[6]{x}}{1 - \sqrt[6]{x}} dx \\
 & \quad \downarrow 798 \\
 & -6 \int \frac{x}{1 - \sqrt[6]{x}} d\sqrt[6]{x} \\
 & \quad \downarrow 49 \\
 & -6 \int \left(-x^{5/6} - x^{2/3} - \sqrt{x} - \sqrt[3]{x} - \sqrt[6]{x} + \frac{1}{1 - \sqrt[6]{x}} - 1 \right) d\sqrt[6]{x} \\
 & \quad \downarrow 2009 \\
 & -6 \left(-\frac{x^{5/6}}{5} - \frac{x^{2/3}}{4} - \frac{x}{6} - \frac{\sqrt{x}}{3} - \frac{\sqrt[3]{x}}{2} - \sqrt[6]{x} - \log(1 - \sqrt[6]{x}) \right)
 \end{aligned}$$

input `Int[Sqrt[x]/(-x^(1/3) + Sqrt[x]),x]`

output `-6*(-x^(1/6) - x^(1/3)/2 - Sqrt[x]/3 - x^(2/3)/4 - x^(5/6)/5 - x/6 - Log[1 - x^(1/6)])`

3.12.3.1 Defintions of rubi rules used

- rule 10 `Int[(u_.)*((e_.)*(x_)^(m_.))*((a_.)*(x_)^(r_.) + (b_.)*(x_)^(s_.))^(p_.), x_Symbol] := Simp[1/e^(p*r) Int[u*(e*x)^(m + p*r)*(a + b*x^(s - r))^p, x], x] /; FreeQ[{a, b, e, m, r, s}, x] && IntegerQ[p] && (IntegerQ[p*r] || GtQ[e, 0]) && PosQ[s - r]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 49 `Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.)), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`
- rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^(p_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.12.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.68

method	result	size
derivativedivides	$x + \frac{6x^{\frac{5}{6}}}{5} + \frac{3x^{\frac{2}{3}}}{2} + 2\sqrt{x} + 3x^{\frac{1}{3}} + 6x^{\frac{1}{6}} + 6 \ln \left(x^{\frac{1}{6}} - 1 \right)$	36
default	$x + \frac{6x^{\frac{5}{6}}}{5} + \frac{3x^{\frac{2}{3}}}{2} + 2\sqrt{x} + 3x^{\frac{1}{3}} + 6x^{\frac{1}{6}} + 6 \ln \left(x^{\frac{1}{6}} - 1 \right)$	36
meijerg	$\frac{x^{\frac{1}{6}} \left(70x^{\frac{5}{6}} + 84x^{\frac{2}{3}} + 105\sqrt{x} + 140x^{\frac{1}{3}} + 210x^{\frac{1}{6}} + 420 \right)}{70} + 6 \ln \left(1 - x^{\frac{1}{6}} \right)$	44

input `int(x^(1/2)/(x^(1/2)-x^(1/3)),x,method=_RETURNVERBOSE)`

output `x+6/5*x^(5/6)+3/2*x^(2/3)+2*x^(1/2)+3*x^(1/3)+6*x^(1/6)+6*ln(x^(1/6)-1)`

3.12.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.66

$$\int \frac{\sqrt{x}}{-\sqrt[3]{x} + \sqrt{x}} dx = x + \frac{6}{5} x^{\frac{5}{6}} + \frac{3}{2} x^{\frac{2}{3}} + 2\sqrt{x} + 3x^{\frac{1}{3}} + 6x^{\frac{1}{6}} + 6 \log(x^{\frac{1}{6}} - 1)$$

input `integrate(x^(1/2)/(x^(1/2)-x^(1/3)),x, algorithm="fracas")`output `x + 6/5*x^(5/6) + 3/2*x^(2/3) + 2*sqrt(x) + 3*x^(1/3) + 6*x^(1/6) + 6*log(x^(1/6) - 1)`**3.12.6 Sympy [F]**

$$\int \frac{\sqrt{x}}{-\sqrt[3]{x} + \sqrt{x}} dx = \int \frac{\sqrt{x}}{-\sqrt[3]{x} + \sqrt{x}} dx$$

input `integrate(x**(1/2)/(x**(1/2)-x**(1/3)),x)`output `Integral(sqrt(x)/(-x**(1/3) + sqrt(x)), x)`**3.12.7 Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.66

$$\int \frac{\sqrt{x}}{-\sqrt[3]{x} + \sqrt{x}} dx = x + \frac{6}{5} x^{\frac{5}{6}} + \frac{3}{2} x^{\frac{2}{3}} + 2\sqrt{x} + 3x^{\frac{1}{3}} + 6x^{\frac{1}{6}} + 6 \log(x^{\frac{1}{6}} - 1)$$

input `integrate(x^(1/2)/(x^(1/2)-x^(1/3)),x, algorithm="maxima")`output `x + 6/5*x^(5/6) + 3/2*x^(2/3) + 2*sqrt(x) + 3*x^(1/3) + 6*x^(1/6) + 6*log(x^(1/6) - 1)`

3.12.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.68

$$\int \frac{\sqrt{x}}{-\sqrt[3]{x} + \sqrt{x}} dx = x + \frac{6}{5} x^{\frac{5}{6}} + \frac{3}{2} x^{\frac{2}{3}} + 2\sqrt{x} + 3x^{\frac{1}{3}} + 6x^{\frac{1}{6}} + 6 \log\left(\left|x^{\frac{1}{6}} - 1\right|\right)$$

input `integrate(x^(1/2)/(x^(1/2)-x^(1/3)),x, algorithm="giac")`output `x + 6/5*x^(5/6) + 3/2*x^(2/3) + 2*sqrt(x) + 3*x^(1/3) + 6*x^(1/6) + 6*log(abs(x^(1/6) - 1))`**3.12.9 Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.66

$$\int \frac{\sqrt{x}}{-\sqrt[3]{x} + \sqrt{x}} dx = x + 6 \ln(x^{1/6} - 1) + 2\sqrt{x} + 3x^{1/3} + \frac{3x^{2/3}}{2} + 6x^{1/6} + \frac{6x^{5/6}}{5}$$

input `int(x^(1/2)/(x^(1/2) - x^(1/3)),x)`output `x + 6*log(x^(1/6) - 1) + 2*x^(1/2) + 3*x^(1/3) + (3*x^(2/3))/2 + 6*x^(1/6) + (6*x^(5/6))/5`

3.13 $\int x^x(1 + \log(x)) dx$

3.13.1	Optimal result	187
3.13.2	Mathematica [A] (verified)	187
3.13.3	Rubi [A] (verified)	188
3.13.4	Maple [A] (verified)	188
3.13.5	Fricas [A] (verification not implemented)	189
3.13.6	Sympy [A] (verification not implemented)	189
3.13.7	Maxima [A] (verification not implemented)	190
3.13.8	Giac [A] (verification not implemented)	190
3.13.9	Mupad [B] (verification not implemented)	190

3.13.1 Optimal result

Integrand size = 8, antiderivative size = 3

$$\int x^x(1 + \log(x)) dx = x^x$$

output

x^x

3.13.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int x^x(1 + \log(x)) dx = x^x$$

input

`Integrate[x^x*(1 + Log[x]),x]`

output

x^x

3.13.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^x(\log(x) + 1) dx$$

$$\downarrow 7293$$

$$\int (x^x + x^x \log(x)) dx$$

$$\downarrow 2009$$

$$x^x$$

input `Int[x^x*(1 + Log[x]),x]`

output `x^x`

3.13.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] :> With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]`

3.13.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
derivativedivides	x^x	4
default	x^x	4
risch	x^x	4
parallelrisch	x^x	4
norman	$e^{x \ln(x)}$	6

input `int(x^x*(ln(x)+1),x,method=_RETURNVERBOSE)`

output `x^x`

3.13.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int x^x(1 + \log(x)) dx = x^x$$

input `integrate(x^x*(log(x)+1),x, algorithm="fricas")`

output `x^x`

3.13.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int x^x(1 + \log(x)) dx = x^x$$

input `integrate(x**x*(ln(x)+1),x)`

output `x**x`

3.13.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int x^x(1 + \log(x)) dx = x^x$$

input `integrate(x^x*(log(x)+1),x, algorithm="maxima")`output `x^x`**3.13.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int x^x(1 + \log(x)) dx = x^x$$

input `integrate(x^x*(log(x)+1),x, algorithm="giac")`output `x^x`**3.13.9 Mupad [B] (verification not implemented)**

Time = 16.26 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int x^x(1 + \log(x)) dx = x^x$$

input `int(x^x*(log(x) + 1),x)`output `x^x`

3.14 $\int x^{13/2} \sqrt{1 + x^{5/2}} dx$

3.14.1	Optimal result	191
3.14.2	Mathematica [A] (verified)	191
3.14.3	Rubi [A] (verified)	192
3.14.4	Maple [A] (verified)	193
3.14.5	Fricas [A] (verification not implemented)	193
3.14.6	Sympy [B] (verification not implemented)	193
3.14.7	Maxima [A] (verification not implemented)	194
3.14.8	Giac [A] (verification not implemented)	194
3.14.9	Mupad [B] (verification not implemented)	195

3.14.1 Optimal result

Integrand size = 17, antiderivative size = 29

$$\int x^{13/2} \sqrt{1 + x^{5/2}} dx = \frac{4}{525} (1 + x^{5/2})^{3/2} (8 - 12x^{5/2} + 15x^5)$$

output `4/525*(1+x^(5/2))^(3/2)*(8-12*x^(5/2)+15*x^5)`

3.14.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int x^{13/2} \sqrt{1 + x^{5/2}} dx = \frac{4}{525} (1 + x^{5/2})^{3/2} (8 - 12x^{5/2} + 15x^5)$$

input `Integrate[x^(13/2)*Sqrt[1 + x^(5/2)],x]`

output `(4*(1 + x^(5/2))^(3/2)*(8 - 12*x^(5/2) + 15*x^5))/525`

3.14.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.72, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {798, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^{13/2} \sqrt{x^{5/2} + 1} dx \\ & \quad \downarrow \text{798} \\ & \frac{2}{5} \int x^5 \sqrt{x^{5/2} + 1} dx^{5/2} \\ & \quad \downarrow \text{53} \\ & \frac{2}{5} \int \left((x^{5/2} + 1)^{5/2} - 2(x^{5/2} + 1)^{3/2} + \sqrt{x^{5/2} + 1} \right) dx^{5/2} \\ & \quad \downarrow \text{2009} \\ & \frac{2}{5} \left(\frac{2}{7} (x^{5/2} + 1)^{7/2} - \frac{4}{5} (x^{5/2} + 1)^{5/2} + \frac{2}{3} (x^{5/2} + 1)^{3/2} \right) \end{aligned}$$

input `Int[x^(13/2)*Sqrt[1 + x^(5/2)],x]`

output `(2*((2*(1 + x^(5/2))^(3/2))/3 - (4*(1 + x^(5/2))^(5/2))/5 + (2*(1 + x^(5/2))^(7/2))/7))/5`

3.14.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.14.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.24

method	result	size
meijerg	$-\frac{32\sqrt{\pi}}{105} - \frac{4\sqrt{\pi} \left(1+x^{\frac{5}{2}}\right)^{\frac{3}{2}} \left(8-12x^{\frac{5}{2}}+15x^5\right)}{5\sqrt{\pi} \cdot 105}$	36

input `int(x^(13/2)*(1+x^(5/2))^(1/2),x,method=_RETURNVERBOSE)`

output
$$-1/5\sqrt{\pi} \left(\frac{32}{105}\sqrt{\pi} - \frac{4}{105}\sqrt{\pi} \left(1+x^{5/2}\right)^{3/2} \left(8-12x^{5/2}+15x^5\right) \right)$$

3.14.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int x^{13/2} \sqrt{1+x^{5/2}} dx = \frac{4}{525} \left(3x^5 + (15x^7 - 4x^2)\sqrt{x} + 8 \right) \sqrt{x^{5/2} + 1}$$

input `integrate(x^(13/2)*(1+x^(5/2))^(1/2),x, algorithm="fracas")`

output
$$4/525 \cdot (3x^5 + (15x^7 - 4x^2)\sqrt{x} + 8)\sqrt{x^{5/2} + 1}$$

3.14.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 66 vs. $2(26) = 52$.

Time = 84.09 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.28

$$\int x^{13/2} \sqrt{1+x^{5/2}} dx = \frac{4x^{15/2} \sqrt{x^{5/2} + 1}}{35} - \frac{16x^{5/2} \sqrt{x^{5/2} + 1}}{525} + \frac{4x^5 \sqrt{x^{5/2} + 1}}{175} + \frac{32\sqrt{x^{5/2} + 1}}{525}$$

input `integrate(x**(13/2)*(1+x**(5/2))**(1/2),x)`

output `4*x**(15/2)*sqrt(x**(5/2) + 1)/35 - 16*x**(5/2)*sqrt(x**(5/2) + 1)/525 + 4*x**5*sqrt(x**(5/2) + 1)/175 + 32*sqrt(x**(5/2) + 1)/525`

3.14.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.97

$$\int x^{13/2} \sqrt{1 + x^{5/2}} dx = \frac{4}{35} \left(x^{5/2} + 1\right)^{7/2} - \frac{8}{25} \left(x^{5/2} + 1\right)^{5/2} + \frac{4}{15} \left(x^{5/2} + 1\right)^{3/2}$$

input `integrate(x^(13/2)*(1+x^(5/2))^(1/2),x, algorithm="maxima")`

output `4/35*(x^(5/2) + 1)^(7/2) - 8/25*(x^(5/2) + 1)^(5/2) + 4/15*(x^(5/2) + 1)^(3/2)`

3.14.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.97

$$\int x^{13/2} \sqrt{1 + x^{5/2}} dx = \frac{4}{35} \left(x^{5/2} + 1\right)^{7/2} - \frac{8}{25} \left(x^{5/2} + 1\right)^{5/2} + \frac{4}{15} \left(x^{5/2} + 1\right)^{3/2}$$

input `integrate(x^(13/2)*(1+x^(5/2))^(1/2),x, algorithm="giac")`

output `4/35*(x^(5/2) + 1)^(7/2) - 8/25*(x^(5/2) + 1)^(5/2) + 4/15*(x^(5/2) + 1)^(3/2)`

3.14.9 Mupad [B] (verification not implemented)

Time = 16.62 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.86

$$\int x^{13/2} \sqrt{1+x^{5/2}} dx = -\frac{4(x^{5/2}+1)^{3/2} (42x^{5/2} - 15(x^{5/2}+1)^2 + 7)}{525}$$

input `int(x^(13/2)*(x^(5/2) + 1)^(1/2),x)`

output `-(4*(x^(5/2) + 1)^(3/2)*(42*x^(5/2) - 15*(x^(5/2) + 1)^2 + 7))/525`

3.15 $\int \frac{1}{(1+x^2)^2} dx$

3.15.1	Optimal result	196
3.15.2	Mathematica [A] (verified)	196
3.15.3	Rubi [A] (verified)	197
3.15.4	Maple [A] (verified)	198
3.15.5	Fricas [A] (verification not implemented)	198
3.15.6	Sympy [A] (verification not implemented)	198
3.15.7	Maxima [A] (verification not implemented)	199
3.15.8	Giac [A] (verification not implemented)	199
3.15.9	Mupad [B] (verification not implemented)	199

3.15.1 Optimal result

Integrand size = 7, antiderivative size = 19

$$\int \frac{1}{(1+x^2)^2} dx = \frac{x}{2(1+x^2)} + \frac{\arctan(x)}{2}$$

output `x/(2*x^2+2)+1/2*arctan(x)`

3.15.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{1}{(1+x^2)^2} dx = \frac{1}{2} \left(\frac{x}{1+x^2} + \arctan(x) \right)$$

input `Integrate[(1 + x^2)^(-2),x]`

output `(x/(1 + x^2) + ArcTan[x])/2`

3.15.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {215, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x^2 + 1)^2} dx$$

↓ 215

$$\frac{1}{2} \int \frac{1}{x^2 + 1} dx + \frac{x}{2(x^2 + 1)}$$

↓ 216

$$\frac{\arctan(x)}{2} + \frac{x}{2(x^2 + 1)}$$

input `Int[(1 + x^2)^(-2), x]`

output `x/(2*(1 + x^2)) + ArcTan[x]/2`

3.15.3.1 Defintions of rubi rules used

rule 215 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

3.15.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

method	result	size
default	$\frac{x}{2x^2+2} + \frac{\arctan(x)}{2}$	16
risch	$\frac{x}{2x^2+2} + \frac{\arctan(x)}{2}$	16
meijerg	$\frac{x}{2x^2+2} + \frac{\arctan(x)}{2}$	17
parallelrisch	$-\frac{i \ln(x-i)x^2 - i \ln(i+x)x^2 + i \ln(x-i) - i \ln(i+x) - 2x}{4(x^2+1)}$	52

input `int(1/(x^2+1)^2,x,method=_RETURNVERBOSE)`output `1/2/(x^2+1)*x+1/2*arctan(x)`**3.15.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x^2)^2} dx = \frac{(x^2+1)\arctan(x) + x}{2(x^2+1)}$$

input `integrate(1/(x^2+1)^2,x, algorithm="fricas")`output `1/2*((x^2 + 1)*arctan(x) + x)/(x^2 + 1)`**3.15.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{1}{(1+x^2)^2} dx = \frac{x}{2x^2+2} + \frac{\operatorname{atan}(x)}{2}$$

input `integrate(1/(x**2+1)**2,x)`output `x/(2*x**2 + 2) + atan(x)/2`

3.15. $\int \frac{1}{(1+x^2)^2} dx$

3.15.7 Maxima [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{(1+x^2)^2} dx = \frac{x}{2(x^2+1)} + \frac{1}{2} \arctan(x)$$

input `integrate(1/(x^2+1)^2,x, algorithm="maxima")`output `1/2*x/(x^2 + 1) + 1/2*arctan(x)`**3.15.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{(1+x^2)^2} dx = \frac{x}{2(x^2+1)} + \frac{1}{2} \arctan(x)$$

input `integrate(1/(x^2+1)^2,x, algorithm="giac")`output `1/2*x/(x^2 + 1) + 1/2*arctan(x)`**3.15.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{1}{(1+x^2)^2} dx = \frac{\operatorname{atan}(x)}{2} + \frac{x}{2(x^2+1)}$$

input `int(1/(x^2 + 1)^2,x)`output `atan(x)/2 + x/(2*(x^2 + 1))`

3.16 $\int \frac{1}{36-13x^2+x^4} dx$

3.16.1	Optimal result	200
3.16.2	Mathematica [A] (verified)	200
3.16.3	Rubi [A] (verified)	201
3.16.4	Maple [A] (verified)	202
3.16.5	Fricas [A] (verification not implemented)	202
3.16.6	Sympy [B] (verification not implemented)	202
3.16.7	Maxima [A] (verification not implemented)	203
3.16.8	Giac [B] (verification not implemented)	203
3.16.9	Mupad [B] (verification not implemented)	203

3.16.1 Optimal result

Integrand size = 12, antiderivative size = 21

$$\int \frac{1}{36 - 13x^2 + x^4} dx = -\frac{1}{15} \operatorname{arctanh}\left(\frac{x}{3}\right) + \frac{1}{10} \operatorname{arctanh}\left(\frac{x}{2}\right)$$

output `-1/15*arctanh(1/3*x)+1/10*arctanh(1/2*x)`

3.16.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.76

$$\int \frac{1}{36 - 13x^2 + x^4} dx = -\frac{1}{20} \log(2 - x) + \frac{1}{30} \log(3 - x) + \frac{1}{20} \log(2 + x) - \frac{1}{30} \log(3 + x)$$

input `Integrate[(36 - 13*x^2 + x^4)^(-1), x]`

output `-1/20*Log[2 - x] + Log[3 - x]/30 + Log[2 + x]/20 - Log[3 + x]/30`

3.16.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {1406, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^4 - 13x^2 + 36} dx$$

$$\downarrow \text{1406}$$

$$\frac{1}{5} \int \frac{1}{x^2 - 9} dx - \frac{1}{5} \int \frac{1}{x^2 - 4} dx$$

$$\downarrow \text{220}$$

$$\frac{1}{10} \operatorname{arctanh}\left(\frac{x}{2}\right) - \frac{1}{15} \operatorname{arctanh}\left(\frac{x}{3}\right)$$

input `Int[(36 - 13*x^2 + x^4)^(-1),x]`

output `-1/15*ArcTanh[x/3] + ArcTanh[x/2]/10`

3.16.3.1 Defintions of rubi rules used

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 1406 `Int[((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c/q Int[1/(b/2 - q/2 + c*x^2), x], x] - Simp[c/q Int[1/(b/2 + q/2 + c*x^2), x], x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[b^2 - 4*a*c]`

3.16.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.24

method	result	size
default	$-\frac{\ln(3+x)}{30} + \frac{\ln(2+x)}{20} - \frac{\ln(-2+x)}{20} + \frac{\ln(-3+x)}{30}$	26
norman	$-\frac{\ln(3+x)}{30} + \frac{\ln(2+x)}{20} - \frac{\ln(-2+x)}{20} + \frac{\ln(-3+x)}{30}$	26
risch	$-\frac{\ln(3+x)}{30} + \frac{\ln(2+x)}{20} - \frac{\ln(-2+x)}{20} + \frac{\ln(-3+x)}{30}$	26
parallelrisc	$-\frac{\ln(3+x)}{30} + \frac{\ln(2+x)}{20} - \frac{\ln(-2+x)}{20} + \frac{\ln(-3+x)}{30}$	26

input `int(1/(x^4-13*x^2+36),x,method=_RETURNVERBOSE)`output `-1/30*ln(3+x)+1/20*ln(2+x)-1/20*ln(-2+x)+1/30*ln(-3+x)`**3.16.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.19

$$\int \frac{1}{36 - 13x^2 + x^4} dx = -\frac{1}{30} \log(x + 3) + \frac{1}{20} \log(x + 2) - \frac{1}{20} \log(x - 2) + \frac{1}{30} \log(x - 3)$$

input `integrate(1/(x^4-13*x^2+36),x, algorithm="fracas")`output `-1/30*log(x + 3) + 1/20*log(x + 2) - 1/20*log(x - 2) + 1/30*log(x - 3)`**3.16.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. 2(12) = 24.

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.24

$$\int \frac{1}{36 - 13x^2 + x^4} dx = \frac{\log(x - 3)}{30} - \frac{\log(x - 2)}{20} + \frac{\log(x + 2)}{20} - \frac{\log(x + 3)}{30}$$

input `integrate(1/(x**4-13*x**2+36),x)`output `log(x - 3)/30 - log(x - 2)/20 + log(x + 2)/20 - log(x + 3)/30`

3.16.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.19

$$\int \frac{1}{36 - 13x^2 + x^4} dx = -\frac{1}{30} \log(x + 3) + \frac{1}{20} \log(x + 2) - \frac{1}{20} \log(x - 2) + \frac{1}{30} \log(x - 3)$$

input `integrate(1/(x^4-13*x^2+36),x, algorithm="maxima")`

output `-1/30*log(x + 3) + 1/20*log(x + 2) - 1/20*log(x - 2) + 1/30*log(x - 3)`

3.16.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. 2(13) = 26.

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.38

$$\int \frac{1}{36 - 13x^2 + x^4} dx = -\frac{1}{30} \log(|x + 3|) + \frac{1}{20} \log(|x + 2|) - \frac{1}{20} \log(|x - 2|) + \frac{1}{30} \log(|x - 3|)$$

input `integrate(1/(x^4-13*x^2+36),x, algorithm="giac")`

output `-1/30*log(abs(x + 3)) + 1/20*log(abs(x + 2)) - 1/20*log(abs(x - 2)) + 1/30*log(abs(x - 3))`

3.16.9 Mupad [B] (verification not implemented)

Time = 15.98 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \frac{1}{36 - 13x^2 + x^4} dx = \frac{\operatorname{atanh}\left(\frac{x}{2}\right)}{10} - \frac{\operatorname{atanh}\left(\frac{x}{3}\right)}{15}$$

input `int(1/(x^4 - 13*x^2 + 36),x)`

output `atanh(x/2)/10 - atanh(x/3)/15`

3.17 $\int \frac{\log(\log(x))}{x} dx$

3.17.1	Optimal result	204
3.17.2	Mathematica [A] (verified)	204
3.17.3	Rubi [A] (verified)	205
3.17.4	Maple [A] (verified)	205
3.17.5	Fricas [A] (verification not implemented)	206
3.17.6	Sympy [A] (verification not implemented)	206
3.17.7	Maxima [A] (verification not implemented)	206
3.17.8	Giac [A] (verification not implemented)	207
3.17.9	Mupad [B] (verification not implemented)	207

3.17.1 Optimal result

Integrand size = 7, antiderivative size = 11

$$\int \frac{\log(\log(x))}{x} dx = -\log(x) + \log(x) \log(\log(x))$$

output `-ln(x)+ln(x)*ln(ln(x))`

3.17.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{\log(\log(x))}{x} dx = -\log(x) + \log(x) \log(\log(x))$$

input `Integrate[Log[Log[x]]/x,x]`

output `-Log[x] + Log[x]*Log[Log[x]]`

3.17.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3001}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\log(\log(x))}{x} dx$$

↓ 3001

$$\log(x) \log(\log(x)) - \log(x)$$

input `Int [Log [Log [x]] / x, x]`

output `-Log [x] + Log [x] * Log [Log [x]]`

3.17.3.1 Defintions of rubi rules used

rule 3001 `Int[((a_.) + Log[Log[(d_.)*(x_)^(n_.)]^(p_.)*(c_.)]*(b_.))/(x_), x_Symbol]
:> Simp[Log[d*x^n]*((a + b*Log[c*Log[d*x^n]^p])/n), x] - Simp[b*p*Log[x], x
] /; FreeQ[{a, b, c, d, n, p}, x]`

3.17.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
derivativedivides	$-\ln(x) + \ln(x) \ln(\ln(x))$	12
default	$-\ln(x) + \ln(x) \ln(\ln(x))$	12
norman	$-\ln(x) + \ln(x) \ln(\ln(x))$	12
risch	$-\ln(x) + \ln(x) \ln(\ln(x))$	12

input `int(ln(ln(x))/x,x,method=_RETURNVERBOSE)`

output `-ln(x)+ln(x)*ln(ln(x))`

3.17.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{\log(\log(x))}{x} dx = \log(x) \log(\log(x)) - \log(x)$$

input `integrate(log(log(x))/x,x, algorithm="fricas")`output `log(x)*log(log(x)) - log(x)`**3.17.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int \frac{\log(\log(x))}{x} dx = \log(x) \log(\log(x)) - \log(x)$$

input `integrate(ln(ln(x))/x,x)`output `log(x)*log(log(x)) - log(x)`**3.17.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{\log(\log(x))}{x} dx = \log(x) \log(\log(x)) - \log(x)$$

input `integrate(log(log(x))/x,x, algorithm="maxima")`output `log(x)*log(log(x)) - log(x)`

3.17.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{\log(\log(x))}{x} dx = \log(x) \log(\log(x)) - \log(x)$$

input `integrate(log(log(x))/x,x, algorithm="giac")`

output `log(x)*log(log(x)) - log(x)`

3.17.9 Mupad [B] (verification not implemented)

Time = 15.86 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \frac{\log(\log(x))}{x} dx = \ln(x) (\ln(\ln(x)) - 1)$$

input `int(log(log(x))/x,x)`

output `log(x)*(log(log(x)) - 1)`

3.18 $\int \frac{1+\cot(x)}{1-\cot(x)} dx$

3.18.1	Optimal result	208
3.18.2	Mathematica [A] (verified)	208
3.18.3	Rubi [A] (verified)	209
3.18.4	Maple [A] (verified)	210
3.18.5	Fricas [A] (verification not implemented)	210
3.18.6	Sympy [B] (verification not implemented)	210
3.18.7	Maxima [A] (verification not implemented)	211
3.18.8	Giac [A] (verification not implemented)	211
3.18.9	Mupad [B] (verification not implemented)	211

3.18.1 Optimal result

Integrand size = 13, antiderivative size = 8

$$\int \frac{1 + \cot(x)}{1 - \cot(x)} dx = \log(\cos(x) - \sin(x))$$

output `ln(cos(x)-sin(x))`

3.18.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{1 + \cot(x)}{1 - \cot(x)} dx = \log(\cos(x) - \sin(x))$$

input `Integrate[(1 + Cot[x])/(1 - Cot[x]), x]`

output `Log[Cos[x] - Sin[x]]`

3.18.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot(x) + 1}{1 - \cot(x)} dx$$

↓ 3042

$$\int \frac{1 - \tan\left(x + \frac{\pi}{2}\right)}{\tan\left(x + \frac{\pi}{2}\right) + 1} dx$$

↓ 4013

$$\log(\cos(x) - \sin(x))$$

input `Int[(1 + Cot[x])/(1 - Cot[x]),x]`

output `Log[Cos[x] - Sin[x]]`

3.18.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

3.18.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.75

method	result	size
parallelrisch	$\ln\left(\frac{1}{\sqrt{\sec(x)^2}}\right) + \ln(\tan(x) - 1)$	14
risch	$-ix + \ln(e^{2ix} - i)$	15
derivativedivides	$\ln(-1 + \cot(x)) - \frac{\ln(1 + \cot(x)^2)}{2}$	16
default	$\ln(-1 + \cot(x)) - \frac{\ln(1 + \cot(x)^2)}{2}$	16
norman	$-\frac{\ln(1 + \tan(x)^2)}{2} + \ln(\tan(x) - 1)$	16

input `int((1+cot(x))/(1-cot(x)),x,method=_RETURNVERBOSE)`

output `ln(1/(sec(x)^2)^(1/2))+ln(tan(x)-1)`

3.18.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.38

$$\int \frac{1 + \cot(x)}{1 - \cot(x)} dx = \frac{1}{2} \log(-\sin(2x) + 1)$$

input `integrate((1+cot(x))/(1-cot(x)),x, algorithm="fricas")`

output `1/2*log(-sin(2*x) + 1)`

3.18.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(7) = 14$.

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.88

$$\int \frac{1 + \cot(x)}{1 - \cot(x)} dx = \log(\tan(x) - 1) - \frac{\log(\tan^2(x) + 1)}{2}$$

input `integrate((1+cot(x))/(1-cot(x)),x)`

output `log(tan(x) - 1) - log(tan(x)**2 + 1)/2`

3.18.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.88

$$\int \frac{1 + \cot(x)}{1 - \cot(x)} dx = -\frac{1}{2} \log(\tan(x)^2 + 1) + \log(\tan(x) - 1)$$

input `integrate((1+cot(x))/(1-cot(x)),x, algorithm="maxima")`

output `-1/2*log(tan(x)^2 + 1) + log(tan(x) - 1)`

3.18.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 2.00

$$\int \frac{1 + \cot(x)}{1 - \cot(x)} dx = -\frac{1}{2} \log(\tan(x)^2 + 1) + \log(|\tan(x) - 1|)$$

input `integrate((1+cot(x))/(1-cot(x)),x, algorithm="giac")`

output `-1/2*log(tan(x)^2 + 1) + log(abs(tan(x) - 1))`

3.18.9 Mupad [B] (verification not implemented)

Time = 16.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.75

$$\int \frac{1 + \cot(x)}{1 - \cot(x)} dx = -x \operatorname{li} + \ln(e^{x2i} - i)$$

input `int(-(cot(x) + 1)/(cot(x) - 1),x)`

output `log(exp(x*2i) - 1i) - x*1i`

$$3.19 \quad \int \frac{\cos(x) + x \sin(x)}{x(x + \cos(x))} dx$$

3.19.1	Optimal result	212
3.19.2	Mathematica [A] (verified)	212
3.19.3	Rubi [A] (verified)	213
3.19.4	Maple [B] (verified)	214
3.19.5	Fricas [A] (verification not implemented)	214
3.19.6	Sympy [A] (verification not implemented)	214
3.19.7	Maxima [B] (verification not implemented)	215
3.19.8	Giac [B] (verification not implemented)	215
3.19.9	Mupad [B] (verification not implemented)	216

3.19.1 Optimal result

Integrand size = 17, antiderivative size = 11

$$\int \frac{\cos(x) + x \sin(x)}{x(x + \cos(x))} dx = -\log\left(1 + \frac{\cos(x)}{x}\right)$$

output `-ln(1+cos(x)/x)`

3.19.2 Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int \frac{\cos(x) + x \sin(x)}{x(x + \cos(x))} dx = \log(x) - \log(x + \cos(x))$$

input `Integrate[(Cos[x] + x*Sin[x])/(x*(x + Cos[x])),x]`

output `Log[x] - Log[x + Cos[x]]`

3.19.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {7263, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x \sin(x) + \cos(x)}{x(x + \cos(x))} dx$$

$$\downarrow 7263$$

$$- \int \frac{1}{\frac{\cos(x)}{x} + 1} d \frac{\cos(x)}{x}$$

$$\downarrow 16$$

$$- \log \left(\frac{\cos(x)}{x} + 1 \right)$$

input `Int[(Cos[x] + x*Sin[x])/(x*(x + Cos[x])),x]`

output `-Log[1 + Cos[x]/x]`

3.19.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 7263 `Int[(u_)*(v_)^(r_.)*((a_.)*(v_)^(p_.) + (b_.)*(w_)^(q_.))^(m_.), x_Symbol] := With[{c = Simplify[u/(p*w*D[v, x] - q*v*D[w, x])]}, Simp[(-c)*q Subst[Int[(a + b*x^q)^m, x], x, v^(m*p + r + 1)*w], x] /; FreeQ[c, x] /; FreeQ[{a, b, m, p, q, r}, x] && EqQ[p + q*(m*p + r + 1), 0] && IntegerQ[q] && IntegerQ[m]`

3.19.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 24 vs. $2(11) = 22$.

Time = 0.21 (sec) , antiderivative size = 25, normalized size of antiderivative = 2.27

method	result	size
parallelsch	$-\ln\left(\frac{x+\cos(x)}{1+\cos(x)}\right) + \ln(x) + \ln\left(\frac{1}{1+\cos(x)}\right)$	25
risch	$ix + \ln(x) - \ln(2x e^{ix} + e^{2ix} + 1)$	26
norman	$-\ln\left(x \tan\left(\frac{x}{2}\right)^2 - \tan\left(\frac{x}{2}\right)^2 + x + 1\right) + \ln(x) + \ln\left(1 + \tan\left(\frac{x}{2}\right)^2\right)$	35

input `int((cos(x)+x*sin(x))/x/(x+cos(x)),x,method=_RETURNVERBOSE)`

output `-ln((x+cos(x))/(1+cos(x)))+ln(x)+ln(1/(1+cos(x)))`

3.19.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int \frac{\cos(x) + x \sin(x)}{x(x + \cos(x))} dx = -\log(x + \cos(x)) + \log(x)$$

input `integrate((cos(x)+x*sin(x))/x/(x+cos(x)),x, algorithm="fricas")`

output `-log(x + cos(x)) + log(x)`

3.19.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \frac{\cos(x) + x \sin(x)}{x(x + \cos(x))} dx = \log(x) - \log(x + \cos(x))$$

input `integrate((cos(x)+x*sin(x))/x/(x+cos(x)),x)`

output `log(x) - log(x + cos(x))`

3.19. $\int \frac{\cos(x)+x \sin(x)}{x(x+\cos(x))} dx$

3.19.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. $2(11) = 22$.

Time = 0.30 (sec) , antiderivative size = 65, normalized size of antiderivative = 5.91

$$\int \frac{\cos(x) + x \sin(x)}{x(x + \cos(x))} dx = -\frac{1}{2} \log(4x^2 \cos(x)^2 + 4x^2 \sin(x)^2 + 4x \sin(2x) \sin(x) + 2(2x \cos(x) + 1) \cos(2x) + \cos(2x)^2 + 4x \cos(x) + \sin(2x)^2 + 1) + \log(x)$$

input `integrate((cos(x)+x*sin(x))/x/(x+cos(x)),x, algorithm="maxima")`

output `-1/2*log(4*x^2*cos(x)^2 + 4*x^2*sin(x)^2 + 4*x*sin(2*x)*sin(x) + 2*(2*x*cos(x) + 1)*cos(2*x) + cos(2*x)^2 + 4*x*cos(x) + sin(2*x)^2 + 1) + log(x)`

3.19.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 79 vs. $2(11) = 22$.

Time = 0.29 (sec) , antiderivative size = 79, normalized size of antiderivative = 7.18

$$\int \frac{\cos(x) + x \sin(x)}{x(x + \cos(x))} dx = -\frac{1}{2} \log\left(\frac{4\left(x^2 \tan\left(\frac{1}{2}x\right)^4 - 2x \tan\left(\frac{1}{2}x\right)^4 + 2x^2 \tan\left(\frac{1}{2}x\right)^2 + \tan\left(\frac{1}{2}x\right)^4 + x^2 - 2 \tan\left(\frac{1}{2}x\right)^2 + 2x + 1\right)}{\tan\left(\frac{1}{2}x\right)^4 + 2 \tan\left(\frac{1}{2}x\right)^2 + 1}\right) + \log(|x|)$$

input `integrate((cos(x)+x*sin(x))/x/(x+cos(x)),x, algorithm="giac")`

output `-1/2*log(4*(x^2*tan(1/2*x)^4 - 2*x*tan(1/2*x)^4 + 2*x^2*tan(1/2*x)^2 + tan(1/2*x)^4 + x^2 - 2*tan(1/2*x)^2 + 2*x + 1)/(tan(1/2*x)^4 + 2*tan(1/2*x)^2 + 1)) + log(abs(x))`

3.19.9 Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int \frac{\cos(x) + x \sin(x)}{x(x + \cos(x))} dx = \ln(x) - \ln(x + \cos(x))$$

input `int((cos(x) + x*sin(x))/(x*(x + cos(x))),x)`

output `log(x) - log(x + cos(x))`

3.20 $\int \frac{1}{\sec(x)+\sin(x)} dx$

3.20.1	Optimal result	217
3.20.2	Mathematica [B] (verified)	217
3.20.3	Rubi [C] (verified)	218
3.20.4	Maple [C] (verified)	219
3.20.5	Fricas [C] (verification not implemented)	220
3.20.6	Sympy [F]	220
3.20.7	Maxima [F]	221
3.20.8	Giac [B] (verification not implemented)	221
3.20.9	Mupad [B] (verification not implemented)	222

3.20.1 Optimal result

Integrand size = 7, antiderivative size = 28

$$\int \frac{1}{\sec(x) + \sin(x)} dx = \arctan(\cos(x) + \sin(x)) - \frac{\operatorname{arctanh}\left(\frac{\cos(x)-\sin(x)}{\sqrt{3}}\right)}{\sqrt{3}}$$

output `arctan(cos(x)+sin(x))-1/3*3^(1/2)*arctanh(1/3*(cos(x)-sin(x))*3^(1/2))`

3.20.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 93 vs. 2(28) = 56.

Time = 1.25 (sec) , antiderivative size = 93, normalized size of antiderivative = 3.32

$$\begin{aligned} & \int \frac{1}{\sec(x) + \sin(x)} dx \\ &= \arctan\left(1 - \left(-1 + \sqrt{3}\right) \tan\left(\frac{x}{2}\right)\right) + \arctan\left(1 + \left(1 + \sqrt{3}\right) \tan\left(\frac{x}{2}\right)\right) \\ & \quad + \frac{-\log\left(\sec^2\left(\frac{x}{2}\right) (\sqrt{3} + \cos(x) - \sin(x))\right) + \log\left(-\sec^2\left(\frac{x}{2}\right) (\sqrt{3} - \cos(x) + \sin(x))\right)}{2\sqrt{3}} \end{aligned}$$

input `Integrate[(Sec[x] + Sin[x])^(-1),x]`

output `ArcTan[1 - (-1 + Sqrt[3])*Tan[x/2]] + ArcTan[1 + (1 + Sqrt[3])*Tan[x/2]] + (-Log[Sec[x/2]^2*(Sqrt[3] + Cos[x] - Sin[x])] + Log[-(Sec[x/2]^2*(Sqrt[3] - Cos[x] + Sin[x]))])/(2*Sqrt[3])`

3.20.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.42 (sec) , antiderivative size = 111, normalized size of antiderivative = 3.96, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 4902, 2492, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sin(x) + \sec(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sin(x) + \sec(x)} dx \\
 & \quad \downarrow \text{4902} \\
 & 2 \int \frac{1 - \tan^2\left(\frac{x}{2}\right)}{\tan^4\left(\frac{x}{2}\right) - 2 \tan^3\left(\frac{x}{2}\right) + 2 \tan^2\left(\frac{x}{2}\right) + 2 \tan\left(\frac{x}{2}\right) + 1} d \tan\left(\frac{x}{2}\right) \\
 & \quad \downarrow \text{2492} \\
 & 2 \int \left(\frac{3i - \sqrt{3}}{6(-i \tan^2\left(\frac{x}{2}\right) + (i + \sqrt{3}) \tan\left(\frac{x}{2}\right) + i)} + \frac{3i + \sqrt{3}}{6(-i \tan^2\left(\frac{x}{2}\right) + (i - \sqrt{3}) \tan\left(\frac{x}{2}\right) + i)} \right) d \tan\left(\frac{x}{2}\right) \\
 & \quad \downarrow \text{2009} \\
 & 2 \left(\frac{1}{6} (3 + i\sqrt{3}) \arctan\left(\frac{-2i \tan\left(\frac{x}{2}\right) - \sqrt{3} + i}{\sqrt{2}(1 + i\sqrt{3})}\right) - \frac{1}{6} (3 - i\sqrt{3}) \arctan\left(\frac{-2i \tan\left(\frac{x}{2}\right) + \sqrt{3} + i}{\sqrt{2}(1 - i\sqrt{3})}\right) \right)
 \end{aligned}$$

input `Int[(Sec[x] + Sin[x])^(-1),x]`

output `2*(((3 + I*Sqrt[3])*ArcTan[(I - Sqrt[3] - (2*I)*Tan[x/2])/Sqrt[2*(1 + I*Sqrt[3])]])/6 - ((3 - I*Sqrt[3])*ArcTan[(I + Sqrt[3] - (2*I)*Tan[x/2])/Sqrt[2*(1 - I*Sqrt[3])]])/6)`

3.20.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2492 `Int[(Px_.)*((a_) + (b_.)*(x_) + (c_.)*(x_)^2 + (d_.)*(x_)^3 + (e_.)*(x_)^4
^(p_), x_Symbol] := Simp[e^p Int[ExpandIntegrand[Px*(b/d + ((d + Sqrt[e*(
(b^2 - 4*a*c)/a) + 8*a*d*(e/b)])/(2*e))*x + x^2)^p*(b/d + ((d - Sqrt[e*(b^
2 - 4*a*c)/a) + 8*a*d*(e/b)])/(2*e))*x + x^2]^p, x], x] /; FreeQ[{a, b,
c, d, e}, x] && PolyQ[Px, x] && ILtQ[p, 0] && EqQ[a*d^2 - b^2*e, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4902 `Int[u_, x_Symbol] := With[{w = Block[{$ShowSteps = False, $StepCounter = Nu
ll}, Int[SubstFor[1/(1 + FreeFactors[Tan[FunctionOfTrig[u, x]/2], x]^2*x^2)
, Tan[FunctionOfTrig[u, x]/2]/FreeFactors[Tan[FunctionOfTrig[u, x]/2], x],
u, x], x]}], Module[{v = FunctionOfTrig[u, x], d}, Simp[d = FreeFactors[Tan
[v/2], x]; 2*(d/Coefficient[v, x, 1]) Subst[Int[SubstFor[1/(1 + d^2*x^2),
Tan[v/2]/d, u, x], x], x, Tan[v/2]/d, x]] /; CalculusFreeQ[w, x]] /; Inve
rseFunctionFreeQ[u, x] && !FalseQ[FunctionOfTrig[u, x]]`

3.20.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.21 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.04

method	result
default	$\sum_{_R=\text{RootOf}(_Z^4-2_Z^3+2_Z^2+2_Z+1)} \frac{(-_R^2+1) \ln(\tan(\frac{x}{2})-_R)}{2_R^3-3_R^2+2_R+1}$
risch	$-\frac{i \ln\left(e^{ix} + \frac{1}{2} - \frac{i}{2} + \frac{i\sqrt{3} - \sqrt{3}}{2}\right)}{2} + \frac{\ln\left(e^{ix} + \frac{1}{2} - \frac{i}{2} + \frac{i\sqrt{3} - \sqrt{3}}{2}\right)\sqrt{3}}{6} - \frac{i \ln\left(e^{ix} + \frac{1}{2} - \frac{i}{2} - \frac{i\sqrt{3} + \sqrt{3}}{2}\right)}{2} - \frac{\ln\left(e^{ix} + \frac{1}{2} - \frac{i}{2} - \frac{i\sqrt{3} + \sqrt{3}}{2}\right)\sqrt{3}}{6} +$

input `int(1/(sin(x)+sec(x)),x,method=_RETURNVERBOSE)`

output `sum((-_R^2+1)/(2*_R^3-3*_R^2+2*_R+1)*ln(tan(1/2*x)-_R),_R=RootOf(_Z^4-2*_Z
^3+2*_Z^2+2*_Z+1))`

3.20. $\int \frac{1}{\sec(x)+\sin(x)} dx$

3.20.5 Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 165, normalized size of antiderivative = 5.89

$$\int \frac{1}{\sec(x) + \sin(x)} dx$$

$$= -\frac{1}{12} \sqrt{3} \sqrt{2i\sqrt{3} - 2} \log\left(-\frac{1}{2} \left((\sqrt{3} - i) \sin(x) + 2i \cos(x) \right) \sqrt{2i\sqrt{3} - 2} + 2\right)$$

$$+ \frac{1}{12} \sqrt{3} \sqrt{2i\sqrt{3} - 2} \log\left(-\frac{1}{2} \left((\sqrt{3} - i) \sin(x) + 2i \cos(x) \right) \sqrt{2i\sqrt{3} - 2} - 2\right)$$

$$+ \frac{1}{12} \sqrt{3} \sqrt{-2i\sqrt{3} - 2} \log\left(\frac{1}{2} \left((\sqrt{3} + i) \sin(x) - 2i \cos(x) \right) \sqrt{-2i\sqrt{3} - 2} + 2\right)$$

$$- \frac{1}{12} \sqrt{3} \sqrt{-2i\sqrt{3} - 2} \log\left(\frac{1}{2} \left((\sqrt{3} + i) \sin(x) - 2i \cos(x) \right) \sqrt{-2i\sqrt{3} - 2} - 2\right)$$

input `integrate(1/(sin(x)+sec(x)),x, algorithm="fricas")`

output `-1/12*sqrt(3)*sqrt(2*I*sqrt(3) - 2)*log(-1/2*((sqrt(3) - I)*sin(x) + 2*I*cos(x))*sqrt(2*I*sqrt(3) - 2) + 2) + 1/12*sqrt(3)*sqrt(2*I*sqrt(3) - 2)*log(-1/2*((sqrt(3) - I)*sin(x) + 2*I*cos(x))*sqrt(2*I*sqrt(3) - 2) - 2) + 1/12*sqrt(3)*sqrt(-2*I*sqrt(3) - 2)*log(1/2*((sqrt(3) + I)*sin(x) - 2*I*cos(x))*sqrt(-2*I*sqrt(3) - 2) + 2) - 1/12*sqrt(3)*sqrt(-2*I*sqrt(3) - 2)*log(1/2*((sqrt(3) + I)*sin(x) - 2*I*cos(x))*sqrt(-2*I*sqrt(3) - 2) - 2)`

3.20.6 Sympy [F]

$$\int \frac{1}{\sec(x) + \sin(x)} dx = \int \frac{1}{\sin(x) + \sec(x)} dx$$

input `integrate(1/(sin(x)+sec(x)),x)`

output `Integral(1/(sin(x) + sec(x)), x)`

3.20.7 Maxima [F]

$$\int \frac{1}{\sec(x) + \sin(x)} dx = \int \frac{1}{\sec(x) + \sin(x)} dx$$

input `integrate(1/(sin(x)+sec(x)),x, algorithm="maxima")`

output `integrate(1/(sec(x) + sin(x)), x)`

3.20.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 81 vs. $2(25) = 50$.

Time = 0.29 (sec) , antiderivative size = 81, normalized size of antiderivative = 2.89

$$\begin{aligned} \int \frac{1}{\sec(x) + \sin(x)} dx &= \frac{1}{2} \pi + \frac{1}{6} \sqrt{3} \log \left(\left(\sqrt{3} + \tan \left(\frac{1}{2} x \right) - 1 \right)^2 + \tan \left(\frac{1}{2} x \right)^2 \right) \\ &\quad - \frac{1}{6} \sqrt{3} \log \left(\left(\sqrt{3} - \tan \left(\frac{1}{2} x \right) + 1 \right)^2 + \tan \left(\frac{1}{2} x \right)^2 \right) \\ &\quad + \arctan \left(\left(\sqrt{3} + 1 \right) \tan \left(\frac{1}{2} x \right) + 1 \right) \\ &\quad + \arctan \left(- \left(\sqrt{3} - 1 \right) \tan \left(\frac{1}{2} x \right) + 1 \right) \end{aligned}$$

input `integrate(1/(sin(x)+sec(x)),x, algorithm="giac")`

output `1/2*pi + 1/6*sqrt(3)*log((sqrt(3) + tan(1/2*x) - 1)^2 + tan(1/2*x)^2) - 1/6*sqrt(3)*log((sqrt(3) - tan(1/2*x) + 1)^2 + tan(1/2*x)^2) + arctan((sqrt(3) + 1)*tan(1/2*x) + 1) + arctan(-(sqrt(3) - 1)*tan(1/2*x) + 1)`

3.20.9 Mupad [B] (verification not implemented)

Time = 16.06 (sec) , antiderivative size = 233, normalized size of antiderivative = 8.32

$$\int \frac{1}{\sec(x) + \sin(x)} dx = -\operatorname{atan}\left(\frac{96 \tan\left(\frac{x}{2}\right)}{64 \tan\left(\frac{x}{2}\right) + 64 - \sqrt{3} \tan\left(\frac{x}{2}\right) 64i}\right) + \frac{\sqrt{3} 32i}{64 \tan\left(\frac{x}{2}\right) + 64 - \sqrt{3} \tan\left(\frac{x}{2}\right) 64i} + \frac{32}{64 \tan\left(\frac{x}{2}\right) + 64 - \sqrt{3} \tan\left(\frac{x}{2}\right) 64i} + \frac{\sqrt{3} \tan\left(\frac{x}{2}\right) 32i}{64 \tan\left(\frac{x}{2}\right) + 64 - \sqrt{3} \tan\left(\frac{x}{2}\right) 64i} \left(-1 + \frac{\sqrt{3} 1i}{3}\right) - \operatorname{atan}\left(-\frac{96 \tan\left(\frac{x}{2}\right)}{64 \tan\left(\frac{x}{2}\right) + 64 + \sqrt{3} \tan\left(\frac{x}{2}\right) 64i}\right) + \frac{\sqrt{3} 32i}{64 \tan\left(\frac{x}{2}\right) + 64 + \sqrt{3} \tan\left(\frac{x}{2}\right) 64i} - \frac{32}{64 \tan\left(\frac{x}{2}\right) + 64 + \sqrt{3} \tan\left(\frac{x}{2}\right) 64i} + \frac{\sqrt{3} \tan\left(\frac{x}{2}\right) 32i}{64 \tan\left(\frac{x}{2}\right) + 64 + \sqrt{3} \tan\left(\frac{x}{2}\right) 64i} \left(1 + \frac{\sqrt{3} 1i}{3}\right)$$

input `int(1/(sin(x) + 1/cos(x)),x)`

output `- atan((96*tan(x/2))/(64*tan(x/2) - 3^(1/2)*tan(x/2)*64i + 64) + (3^(1/2)*32i)/(64*tan(x/2) - 3^(1/2)*tan(x/2)*64i + 64) + 32/(64*tan(x/2) - 3^(1/2)*tan(x/2)*64i + 64) + (3^(1/2)*tan(x/2)*32i)/(64*tan(x/2) - 3^(1/2)*tan(x/2)*64i + 64))*((3^(1/2)*1i)/3 - 1) - atan((3^(1/2)*32i)/(64*tan(x/2) + 3^(1/2)*tan(x/2)*64i + 64) - (96*tan(x/2))/(64*tan(x/2) + 3^(1/2)*tan(x/2)*64i + 64) - 32/(64*tan(x/2) + 3^(1/2)*tan(x/2)*64i + 64) + (3^(1/2)*tan(x/2)*32i)/(64*tan(x/2) + 3^(1/2)*tan(x/2)*64i + 64))*((3^(1/2)*1i)/3 + 1)`

3.21 $\int \frac{1}{\sqrt{1+e^x+e^{2x}}} dx$

3.21.1	Optimal result	223
3.21.2	Mathematica [A] (verified)	223
3.21.3	Rubi [A] (verified)	224
3.21.4	Maple [A] (verified)	225
3.21.5	Fricas [A] (verification not implemented)	225
3.21.6	Sympy [F]	226
3.21.7	Maxima [A] (verification not implemented)	226
3.21.8	Giac [A] (verification not implemented)	226
3.21.9	Mupad [B] (verification not implemented)	227

3.21.1 Optimal result

Integrand size = 14, antiderivative size = 26

$$\int \frac{1}{\sqrt{1+e^x+e^{2x}}} dx = -\operatorname{arctanh}\left(\frac{2+e^x}{2\sqrt{1+e^x+e^{2x}}}\right)$$

output `-arctanh(1/2*(exp(x)+2)/(1+exp(x)+exp(2*x))^(1/2))`

3.21.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.88

$$\int \frac{1}{\sqrt{1+e^x+e^{2x}}} dx = 2\operatorname{arctanh}\left(e^x - \sqrt{1+e^x+e^{2x}}\right)$$

input `Integrate[1/Sqrt[1 + E^x + E^(2*x)],x]`

output `2*ArcTanh[E^x - Sqrt[1 + E^x + E^(2*x)]]`

3.21.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {2720, 1154, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt{e^x + e^{2x} + 1}} dx \\ & \quad \downarrow \text{2720} \\ & \int \frac{e^{-x}}{\sqrt{e^x + e^{2x} + 1}} de^x \\ & \quad \downarrow \text{1154} \\ & -2 \int \frac{1}{4 - e^{2x}} d \frac{2 + e^x}{\sqrt{1 + e^x + e^{2x}}} \\ & \quad \downarrow \text{219} \\ & -\operatorname{arctanh} \left(\frac{e^x + 2}{2\sqrt{e^x + e^{2x} + 1}} \right) \end{aligned}$$

input `Int[1/Sqrt[1 + E^x + E^(2*x)], x]`

output `-ArcTanh[(2 + E^x)/(2*Sqrt[1 + E^x + E^(2*x)])]`

3.21.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1154 `Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] :> Simp[-2 Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x]`

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.21.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

method	result	size
default	$-\operatorname{arctanh}\left(\frac{e^x+2}{2\sqrt{1+e^x+e^{2x}}}\right)$	20

```
input int(1/(1+exp(x)+exp(2*x))^(1/2),x,method=_RETURNVERBOSE)
```

```
output -arctanh(1/2*(exp(x)+2)/(1+exp(x)+exp(x)^2)^(1/2))
```

3.21.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.42

$$\int \frac{1}{\sqrt{1+e^x+e^{2x}}} dx = -\log\left(\sqrt{e^{(2x)}+e^x+1}-e^x+1\right) + \log\left(\sqrt{e^{(2x)}+e^x+1}-e^x-1\right)$$

```
input integrate(1/(1+exp(x)+exp(2*x))^(1/2),x, algorithm="fricas")
```

```
output -log(sqrt(e^(2*x) + e^x + 1) - e^x + 1) + log(sqrt(e^(2*x) + e^x + 1) - e^
x - 1)
```

3.21.6 Sympy [F]

$$\int \frac{1}{\sqrt{1+e^x+e^{2x}}} dx = \int \frac{1}{\sqrt{e^{2x}+e^x+1}} dx$$

input `integrate(1/(1+exp(x)+exp(2*x))**(1/2),x)`

output `Integral(1/sqrt(exp(2*x) + exp(x) + 1), x)`

3.21.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.69

$$\int \frac{1}{\sqrt{1+e^x+e^{2x}}} dx = -\operatorname{arsinh}\left(\frac{2}{3}\sqrt{3}e^{-x} + \frac{1}{3}\sqrt{3}\right)$$

input `integrate(1/(1+exp(x)+exp(2*x))^(1/2),x, algorithm="maxima")`

output `-arcsinh(2/3*sqrt(3)*e^(-x) + 1/3*sqrt(3))`

3.21.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.42

$$\int \frac{1}{\sqrt{1+e^x+e^{2x}}} dx = -\log\left(\sqrt{e^{(2x)}+e^x+1}-e^x+1\right) + \log\left(-\sqrt{e^{(2x)}+e^x+1}+e^x+1\right)$$

input `integrate(1/(1+exp(x)+exp(2*x))^(1/2),x, algorithm="giac")`

output `-log(sqrt(e^(2*x) + e^x + 1) - e^x + 1) + log(-sqrt(e^(2*x) + e^x + 1) + e^x + 1)`

3.21.9 Mupad [B] (verification not implemented)

Time = 16.08 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.81

$$\int \frac{1}{\sqrt{1+e^x+e^{2x}}} dx = x - \ln \left(\frac{e^x}{2} + \sqrt{e^{2x} + e^x + 1} + 1 \right)$$

input `int(1/(exp(2*x) + exp(x) + 1)^(1/2),x)`

output `x - log(exp(x)/2 + (exp(2*x) + exp(x) + 1)^(1/2) + 1)`

3.22 $\int e^{x^2} x^3 dx$

3.22.1	Optimal result	228
3.22.2	Mathematica [A] (verified)	228
3.22.3	Rubi [A] (verified)	229
3.22.4	Maple [A] (verified)	230
3.22.5	Fricas [A] (verification not implemented)	230
3.22.6	Sympy [A] (verification not implemented)	231
3.22.7	Maxima [A] (verification not implemented)	231
3.22.8	Giac [A] (verification not implemented)	231
3.22.9	Mupad [B] (verification not implemented)	232

3.22.1 Optimal result

Integrand size = 9, antiderivative size = 22

$$\int e^{x^2} x^3 dx = -\frac{e^{x^2}}{2} + \frac{1}{2}e^{x^2} x^2$$

output `-1/2*exp(x^2)+1/2*x^2*exp(x^2)`

3.22.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.64

$$\int e^{x^2} x^3 dx = \frac{1}{2}e^{x^2}(-1 + x^2)$$

input `Integrate[E^x^2*x^3,x]`

output `(E^x^2*(-1 + x^2))/2`

3.22.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {2641, 2638}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{x^2} x^3 dx$$

$$\downarrow \text{2641}$$

$$\frac{1}{2} e^{x^2} x^2 - \int e^{x^2} x dx$$

$$\downarrow \text{2638}$$

$$\frac{1}{2} e^{x^2} x^2 - \frac{e^{x^2}}{2}$$

input `Int[E^x^2*x^3,x]`

output `-1/2*E^x^2 + (E^x^2*x^2)/2`

3.22.3.1 Defintions of rubi rules used

rule 2638 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[(e + f*x)^n*(F^(a + b*(c + d*x)^n)/(b*f*n*(c + d*x)^n *Log[F])), x] /; FreeQ[{F, a, b, c, d, e, f, n}, x] && EqQ[m, n - 1] && EqQ[d*e - c*f, 0]`

rule 2641 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^(m - n + 1)*(F^(a + b*(c + d*x)^n)/(b*d*n*Log[F])), x] - Simp[(m - n + 1)/(b*n*Log[F]) Int[(c + d*x)^(m - n)*F^(a + b*(c + d*x)^n), x], x] /; FreeQ[{F, a, b, c, d}, x] && IntegerQ[2*((m + 1)/n)] && LtQ[0, (m + 1)/n, 5] && IntegerQ[n] && (LtQ[0, n, m + 1] || LtQ[m, n, 0])`

3.22.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.55

method	result	size
gosper	$\frac{(x^2-1)e^{x^2}}{2}$	12
risch	$\left(\frac{x^2}{2} - \frac{1}{2}\right) e^{x^2}$	13
meijerg	$\frac{1}{2} - \frac{(-2x^2+2)e^{x^2}}{4}$	16
derivativedivides	$-\frac{e^{x^2}}{2} + \frac{x^2 e^{x^2}}{2}$	17
default	$-\frac{e^{x^2}}{2} + \frac{x^2 e^{x^2}}{2}$	17
norman	$-\frac{e^{x^2}}{2} + \frac{x^2 e^{x^2}}{2}$	17
parallelrisch	$-\frac{e^{x^2}}{2} + \frac{x^2 e^{x^2}}{2}$	17
parts	$\frac{\sqrt{\pi} \operatorname{erfi}(x)x^3}{2} - \frac{3\sqrt{\pi} \left(\frac{x^3 \operatorname{erfi}(x)}{3} - \frac{2 \left(-\frac{e^{x^2}}{2} + \frac{x^2 e^{x^2}}{2} \right)}{3\sqrt{\pi}} \right)}{2}$	46

input `int(x^3*exp(x^2),x,method=_RETURNVERBOSE)`output `1/2*(x^2-1)*exp(x^2)`**3.22.5 Fricas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.50

$$\int e^{x^2} x^3 dx = \frac{1}{2} (x^2 - 1) e^{(x^2)}$$

input `integrate(x^3*exp(x^2),x, algorithm="fricas")`output `1/2*(x^2 - 1)*e^(x^2)`

3.22.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.45

$$\int e^{x^2} x^3 dx = \frac{(x^2 - 1) e^{x^2}}{2}$$

input `integrate(x**3*exp(x**2),x)`output `(x**2 - 1)*exp(x**2)/2`**3.22.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.50

$$\int e^{x^2} x^3 dx = \frac{1}{2} (x^2 - 1) e^{(x^2)}$$

input `integrate(x^3*exp(x^2),x, algorithm="maxima")`output `1/2*(x^2 - 1)*e^(x^2)`**3.22.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.50

$$\int e^{x^2} x^3 dx = \frac{1}{2} (x^2 - 1) e^{(x^2)}$$

input `integrate(x^3*exp(x^2),x, algorithm="giac")`output `1/2*(x^2 - 1)*e^(x^2)`

3.22.9 Mupad [B] (verification not implemented)

Time = 15.79 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.50

$$\int e^{x^2} x^3 dx = \frac{e^{x^2} (x^2 - 1)}{2}$$

input `int(x^3*exp(x^2),x)`

output `(exp(x^2)*(x^2 - 1))/2`

$$3.23 \quad \int \left(-\frac{1}{\log^2(x)} + \frac{1}{\log(x)} \right) dx$$

3.23.1	Optimal result	233
3.23.2	Mathematica [A] (verified)	233
3.23.3	Rubi [A] (verified)	234
3.23.4	Maple [A] (verified)	234
3.23.5	Fricas [A] (verification not implemented)	235
3.23.6	Sympy [A] (verification not implemented)	235
3.23.7	Maxima [C] (verification not implemented)	235
3.23.8	Giac [A] (verification not implemented)	236
3.23.9	Mupad [B] (verification not implemented)	236

3.23.1 Optimal result

Integrand size = 11, antiderivative size = 6

$$\int \left(-\frac{1}{\log^2(x)} + \frac{1}{\log(x)} \right) dx = \frac{x}{\log(x)}$$

output `x/ln(x)`

3.23.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \left(-\frac{1}{\log^2(x)} + \frac{1}{\log(x)} \right) dx = \frac{x}{\log(x)}$$

input `Integrate[-Log[x]^(-2) + Log[x]^(-1), x]`

output `x/Log[x]`

3.23. $\int \left(-\frac{1}{\log^2(x)} + \frac{1}{\log(x)} \right) dx$

3.23.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\frac{1}{\log(x)} - \frac{1}{\log^2(x)} \right) dx$$

\downarrow 2009
 $\frac{x}{\log(x)}$

input `Int[-Log[x]^(-2) + Log[x]^(-1),x]`

output `x/Log[x]`

3.23.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.23.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

method	result	size
default	$\frac{x}{\ln(x)}$	7
norman	$\frac{x}{\ln(x)}$	7
risch	$\frac{x}{\ln(x)}$	7
parallelrisc	$\frac{x}{\ln(x)}$	7
parts	$\frac{x}{\ln(x)}$	7

input `int(1/ln(x)-1/ln(x)^2,x,method=_RETURNVERBOSE)`

output `x/ln(x)`

3.23. $\int \left(-\frac{1}{\log^2(x)} + \frac{1}{\log(x)} \right) dx$

3.23.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \left(-\frac{1}{\log^2(x)} + \frac{1}{\log(x)} \right) dx = \frac{x}{\log(x)}$$

input `integrate(1/log(x)-1/log(x)^2,x, algorithm="fracas")`

output `x/log(x)`

3.23.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.50

$$\int \left(-\frac{1}{\log^2(x)} + \frac{1}{\log(x)} \right) dx = \frac{x}{\log(x)}$$

input `integrate(1/ln(x)-1/ln(x)**2,x)`

output `x/log(x)`

3.23.7 Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 0.25 (sec) , antiderivative size = 12, normalized size of antiderivative = 2.00

$$\int \left(-\frac{1}{\log^2(x)} + \frac{1}{\log(x)} \right) dx = \text{Ei}(\log(x)) - \Gamma(-1, -\log(x))$$

input `integrate(1/log(x)-1/log(x)^2,x, algorithm="maxima")`

output `Ei(log(x)) - gamma(-1, -log(x))`

3.23.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \left(-\frac{1}{\log^2(x)} + \frac{1}{\log(x)} \right) dx = \frac{x}{\log(x)}$$

input `integrate(1/log(x)-1/log(x)^2,x, algorithm="giac")`

output `x/log(x)`

3.23.9 Mupad [B] (verification not implemented)

Time = 15.23 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \left(-\frac{1}{\log^2(x)} + \frac{1}{\log(x)} \right) dx = \frac{x}{\ln(x)}$$

input `int(1/log(x) - 1/log(x)^2,x)`

output `x/log(x)`

3.24 $\int \sqrt{2-x}\sqrt{-1+x} dx$

3.24.1	Optimal result	237
3.24.2	Mathematica [A] (verified)	237
3.24.3	Rubi [A] (verified)	238
3.24.4	Maple [A] (verified)	239
3.24.5	Fricas [A] (verification not implemented)	240
3.24.6	Sympy [C] (verification not implemented)	240
3.24.7	Maxima [A] (verification not implemented)	241
3.24.8	Giac [A] (verification not implemented)	241
3.24.9	Mupad [B] (verification not implemented)	241

3.24.1 Optimal result

Integrand size = 17, antiderivative size = 53

$$\int \sqrt{2-x}\sqrt{-1+x} dx = \frac{1}{4}\sqrt{2-x}\sqrt{-1+x} - \frac{1}{2}(2-x)^{3/2}\sqrt{-1+x} + \frac{1}{4}\arcsin(\sqrt{-1+x})$$

output `1/4*(-1+x)^(1/2)*(2-x)^(1/2)-1/2*(2-x)^(3/2)*(-1+x)^(1/2)+1/4*arcsin((-1+x)^(1/2))`

3.24.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00

$$\int \sqrt{2-x}\sqrt{-1+x} dx = \frac{1}{4}\sqrt{-2+3x-x^2} \left(-3+2x - \frac{\operatorname{arctanh}\left(\frac{1}{\sqrt{\frac{-2+x}{-1+x}}}\right)}{\sqrt{-2+x}\sqrt{-1+x}} \right)$$

input `Integrate[Sqrt[2 - x]*Sqrt[-1 + x],x]`

output `(Sqrt[-2 + 3*x - x^2]*(-3 + 2*x - ArcTanh[1/Sqrt[(-2 + x)/(-1 + x)]])/(Sqrt[-2 + x]*Sqrt[-1 + x]))/4`

3.24.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {60, 60, 62, 1090, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{2-x}\sqrt{x-1} dx \\
 & \quad \downarrow 60 \\
 & \frac{1}{4} \int \frac{\sqrt{2-x}}{\sqrt{x-1}} dx - \frac{1}{2}(2-x)^{3/2}\sqrt{x-1} \\
 & \quad \downarrow 60 \\
 & \frac{1}{4} \left(\frac{1}{2} \int \frac{1}{\sqrt{2-x}\sqrt{x-1}} dx + \sqrt{2-x}\sqrt{x-1} \right) - \frac{1}{2}(2-x)^{3/2}\sqrt{x-1} \\
 & \quad \downarrow 62 \\
 & \frac{1}{4} \left(\frac{1}{2} \int \frac{1}{\sqrt{-x^2+3x-2}} dx + \sqrt{2-x}\sqrt{x-1} \right) - \frac{1}{2}(2-x)^{3/2}\sqrt{x-1} \\
 & \quad \downarrow 1090 \\
 & \frac{1}{4} \left(\sqrt{2-x}\sqrt{x-1} - \frac{1}{2} \int \frac{1}{\sqrt{1-(3-2x)^2}} d(3-2x) \right) - \frac{1}{2}(2-x)^{3/2}\sqrt{x-1} \\
 & \quad \downarrow 223 \\
 & \frac{1}{4} \left(\sqrt{2-x}\sqrt{x-1} - \frac{1}{2} \arcsin(3-2x) \right) - \frac{1}{2}(2-x)^{3/2}\sqrt{x-1}
 \end{aligned}$$

input `Int[Sqrt[2 - x]*Sqrt[-1 + x],x]`

output `-1/2*((2 - x)^(3/2)*Sqrt[-1 + x]) + (Sqrt[2 - x]*Sqrt[-1 + x] - ArcSin[3 - 2*x])/4`

3.24.3.1 Defintions of rubi rules used

- rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 62 `Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b + d, 0] && GtQ[a + c, 0]`
- rule 223 `Int[1/Sqrt[(a_.) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 1090 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

3.24.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.15

method	result	size
default	$-\frac{(2-x)^{\frac{3}{2}}\sqrt{-1+x}}{2} + \frac{\sqrt{-1+x}\sqrt{2-x}}{4} + \frac{\sqrt{(2-x)(-1+x)} \arcsin(-3+2x)}{8\sqrt{2-x}\sqrt{-1+x}}$	61
risch	$-\frac{(-3+2x)\sqrt{-1+x}(-2+x)\sqrt{(2-x)(-1+x)}}{4\sqrt{-(-1+x)(-2+x)}\sqrt{2-x}} + \frac{\sqrt{(2-x)(-1+x)} \arcsin(-3+2x)}{8\sqrt{2-x}\sqrt{-1+x}}$	76

input `int((-1+x)^(1/2)*(2-x)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2*(2-x)^(3/2)*(-1+x)^(1/2)+1/4*(-1+x)^(1/2)*(2-x)^(1/2)+1/8*((2-x)*(-1+x))^(1/2)/(2-x)^(1/2)/(-1+x)^(1/2)*arcsin(-3+2*x)`

3.24.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.98

$$\int \sqrt{2-x}\sqrt{-1+x} dx$$

$$= \frac{1}{4}(2x-3)\sqrt{x-1}\sqrt{-x+2} - \frac{1}{8} \arctan\left(\frac{(2x-3)\sqrt{x-1}\sqrt{-x+2}}{2(x^2-3x+2)}\right)$$

input `integrate((-1+x)^(1/2)*(2-x)^(1/2),x, algorithm="fracas")`

output `1/4*(2*x - 3)*sqrt(x - 1)*sqrt(-x + 2) - 1/8*arctan(1/2*(2*x - 3)*sqrt(x - 1)*sqrt(-x + 2)/(x^2 - 3*x + 2))`

3.24.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.89 (sec) , antiderivative size = 124, normalized size of antiderivative = 2.34

$$\int \sqrt{2-x}\sqrt{-1+x} dx = \begin{cases} -\frac{i \operatorname{acosh}(\sqrt{x-1})}{4} + \frac{i(x-1)^{\frac{5}{2}}}{2\sqrt{x-2}} - \frac{3i(x-1)^{\frac{3}{2}}}{4\sqrt{x-2}} + \frac{i\sqrt{x-1}}{4\sqrt{x-2}} & \text{for } |x-1| > 1 \\ \frac{\operatorname{asin}(\sqrt{x-1})}{4} - \frac{(x-1)^{\frac{5}{2}}}{2\sqrt{2-x}} + \frac{3(x-1)^{\frac{3}{2}}}{4\sqrt{2-x}} - \frac{\sqrt{x-1}}{4\sqrt{2-x}} & \text{otherwise} \end{cases}$$

input `integrate((-1+x)**(1/2)*(2-x)**(1/2),x)`

output `Piecewise((-I*acosh(sqrt(x - 1))/4 + I*(x - 1)**(5/2)/(2*sqrt(x - 2)) - 3*I*(x - 1)**(3/2)/(4*sqrt(x - 2)) + I*sqrt(x - 1)/(4*sqrt(x - 2)), Abs(x - 1) > 1), (asin(sqrt(x - 1))/4 - (x - 1)**(5/2)/(2*sqrt(2 - x)) + 3*(x - 1)**(3/2)/(4*sqrt(2 - x)) - sqrt(x - 1)/(4*sqrt(2 - x)), True))`

3.24.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.72

$$\int \sqrt{2-x}\sqrt{-1+x} dx = \frac{1}{2} \sqrt{-x^2+3x-2}x - \frac{3}{4} \sqrt{-x^2+3x-2} + \frac{1}{8} \arcsin(2x-3)$$

input `integrate((-1+x)^(1/2)*(2-x)^(1/2),x, algorithm="maxima")`output `1/2*sqrt(-x^2 + 3*x - 2)*x - 3/4*sqrt(-x^2 + 3*x - 2) + 1/8*arcsin(2*x - 3)`**3.24.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.79

$$\int \sqrt{2-x}\sqrt{-1+x} dx = \frac{1}{4} (2x+1)\sqrt{x-1}\sqrt{-x+2} - \sqrt{x-1}\sqrt{-x+2} + \frac{1}{4} \arcsin(\sqrt{x-1})$$

input `integrate((-1+x)^(1/2)*(2-x)^(1/2),x, algorithm="giac")`output `1/4*(2*x + 1)*sqrt(x - 1)*sqrt(-x + 2) - sqrt(x - 1)*sqrt(-x + 2) + 1/4*arcsin(sqrt(x - 1))`**3.24.9 Mupad [B] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.77

$$\int \sqrt{2-x}\sqrt{-1+x} dx = \left(\frac{x}{2} - \frac{3}{4}\right) \sqrt{x-1}\sqrt{2-x} - \frac{\ln\left(x - \frac{3}{2} - \sqrt{x-1}\sqrt{2-x}\right)}{8}$$

input `int((x - 1)^(1/2)*(2 - x)^(1/2),x)`output `(x/2 - 3/4)*(x - 1)^(1/2)*(2 - x)^(1/2) - (log(x - (x - 1)^(1/2)*(2 - x)^(1/2)*i - 3/2)*i)/8`

$$3.25 \quad \int \frac{-1+x^6}{-1-x+x^3+x^4} dx$$

3.25.1	Optimal result	242
3.25.2	Mathematica [A] (verified)	242
3.25.3	Rubi [A] (verified)	243
3.25.4	Maple [A] (verified)	244
3.25.5	Fricas [A] (verification not implemented)	244
3.25.6	Sympy [A] (verification not implemented)	244
3.25.7	Maxima [A] (verification not implemented)	245
3.25.8	Giac [A] (verification not implemented)	245
3.25.9	Mupad [B] (verification not implemented)	245

3.25.1 Optimal result

Integrand size = 19, antiderivative size = 16

$$\int \frac{-1+x^6}{-1-x+x^3+x^4} dx = x - \frac{x^2}{2} + \frac{x^3}{3}$$

output `x-1/2*x^2+1/3*x^3`

3.25.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{-1+x^6}{-1-x+x^3+x^4} dx = x - \frac{x^2}{2} + \frac{x^3}{3}$$

input `Integrate[(-1 + x^6)/(-1 - x + x^3 + x^4), x]`

output `x - x^2/2 + x^3/3`

3.25.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2019, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^6 - 1}{x^4 + x^3 - x - 1} dx$$

↓ 2019

$$\int (x^2 - x + 1) dx$$

↓ 2009

$$\frac{x^3}{3} - \frac{x^2}{2} + x$$

input `Int[(-1 + x^6)/(-1 - x + x^3 + x^4), x]`

output `x - x^2/2 + x^3/3`

3.25.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2019 `Int[(u_.)*(Px_)^(p_.)*(Qx_)^(q_.), x_Symbol] := Int[u*PolynomialQuotient[Px, Qx, x]^p*Qx^(p + q), x] /; FreeQ[q, x] && PolyQ[Px, x] && PolyQ[Qx, x] && EqQ[PolynomialRemainder[Px, Qx, x], 0] && IntegerQ[p] && LtQ[p*q, 0]`

3.25.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

method	result	size
default	$x - \frac{1}{2}x^2 + \frac{1}{3}x^3$	13
norman	$x - \frac{1}{2}x^2 + \frac{1}{3}x^3$	13
risch	$x - \frac{1}{2}x^2 + \frac{1}{3}x^3$	13
parallelrisc	$x - \frac{1}{2}x^2 + \frac{1}{3}x^3$	13
parts	$x - \frac{1}{2}x^2 + \frac{1}{3}x^3$	13
gosper	$\frac{x(2x^2-3x+6)}{6}$	14

input `int((x^6-1)/(x^4+x^3-x-1),x,method=_RETURNVERBOSE)`

output `x-1/2*x^2+1/3*x^3`

3.25.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{-1+x^6}{-1-x+x^3+x^4} dx = \frac{1}{3}x^3 - \frac{1}{2}x^2 + x$$

input `integrate((x^6-1)/(x^4+x^3-x-1),x, algorithm="fricas")`

output `1/3*x^3 - 1/2*x^2 + x`

3.25.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

$$\int \frac{-1+x^6}{-1-x+x^3+x^4} dx = \frac{x^3}{3} - \frac{x^2}{2} + x$$

input `integrate((x**6-1)/(x**4+x**3-x-1),x)`

output `x**3/3 - x**2/2 + x`

3.25. $\int \frac{-1+x^6}{-1-x+x^3+x^4} dx$

3.25.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{-1 + x^6}{-1 - x + x^3 + x^4} dx = \frac{1}{3} x^3 - \frac{1}{2} x^2 + x$$

input `integrate((x^6-1)/(x^4+x^3-x-1),x, algorithm="maxima")`output `1/3*x^3 - 1/2*x^2 + x`**3.25.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{-1 + x^6}{-1 - x + x^3 + x^4} dx = \frac{1}{3} x^3 - \frac{1}{2} x^2 + x$$

input `integrate((x^6-1)/(x^4+x^3-x-1),x, algorithm="giac")`output `1/3*x^3 - 1/2*x^2 + x`**3.25.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

$$\int \frac{-1 + x^6}{-1 - x + x^3 + x^4} dx = \frac{x(2x^2 - 3x + 6)}{6}$$

input `int(-(x^6 - 1)/(x - x^3 - x^4 + 1),x)`output `(x*(2*x^2 - 3*x + 6))/6`

3.26 $\int (2 \log(x) + \log^2(x)) dx$

3.26.1	Optimal result	246
3.26.2	Mathematica [A] (verified)	246
3.26.3	Rubi [A] (verified)	247
3.26.4	Maple [A] (verified)	247
3.26.5	Fricas [A] (verification not implemented)	248
3.26.6	Sympy [A] (verification not implemented)	248
3.26.7	Maxima [B] (verification not implemented)	248
3.26.8	Giac [A] (verification not implemented)	249
3.26.9	Mupad [B] (verification not implemented)	249

3.26.1 Optimal result

Integrand size = 9, antiderivative size = 6

$$\int (2 \log(x) + \log^2(x)) dx = x \log^2(x)$$

output `x*ln(x)^2`

3.26.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int (2 \log(x) + \log^2(x)) dx = x \log^2(x)$$

input `Integrate[2*Log[x] + Log[x]^2,x]`

output `x*Log[x]^2`

3.26.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\log^2(x) + 2 \log(x)) dx$$

↓ 2009

$$x \log^2(x)$$

input `Int[2*Log[x] + Log[x]^2,x]`

output `x*Log[x]^2`

3.26.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.26.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

method	result	size
default	$x \ln(x)^2$	7
norman	$x \ln(x)^2$	7
risch	$x \ln(x)^2$	7
parallelrisch	$x \ln(x)^2$	7
parts	$x \ln(x)^2$	7

input `int(2*ln(x)+ln(x)^2,x,method=_RETURNVERBOSE)`

output `x*ln(x)^2`

3.26.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int (2\log(x) + \log^2(x)) dx = x \log(x)^2$$

input `integrate(2*log(x)+log(x)^2,x, algorithm="fracas")`

output `x*log(x)^2`

3.26.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int (2\log(x) + \log^2(x)) dx = x \log(x)^2$$

input `integrate(2*ln(x)+ln(x)**2,x)`

output `x*log(x)**2`

3.26.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. $2(6) = 12$.

Time = 0.20 (sec) , antiderivative size = 21, normalized size of antiderivative = 3.50

$$\int (2\log(x) + \log^2(x)) dx = (\log(x)^2 - 2\log(x) + 2)x + 2x\log(x) - 2x$$

input `integrate(2*log(x)+log(x)^2,x, algorithm="maxima")`

output `(log(x)^2 - 2*log(x) + 2)*x + 2*x*log(x) - 2*x`

3.26.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int (2\log(x) + \log^2(x)) dx = x \log(x)^2$$

input `integrate(2*log(x)+log(x)^2,x, algorithm="giac")`

output `x*log(x)^2`

3.26.9 Mupad [B] (verification not implemented)

Time = 14.72 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int (2\log(x) + \log^2(x)) dx = x \ln(x)^2$$

input `int(2*log(x) + log(x)^2,x)`

output `x*log(x)^2`

3.27 $\int \frac{2x}{\sqrt{1-x^4}} dx$

3.27.1	Optimal result	250
3.27.2	Mathematica [A] (verified)	250
3.27.3	Rubi [A] (verified)	251
3.27.4	Maple [A] (verified)	252
3.27.5	Fricas [B] (verification not implemented)	252
3.27.6	Sympy [A] (verification not implemented)	252
3.27.7	Maxima [B] (verification not implemented)	253
3.27.8	Giac [A] (verification not implemented)	253
3.27.9	Mupad [B] (verification not implemented)	253

3.27.1 Optimal result

Integrand size = 14, antiderivative size = 4

$$\int \frac{2x}{\sqrt{1-x^4}} dx = \arcsin(x^2)$$

output `arcsin(x^2)`

3.27.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{2x}{\sqrt{1-x^4}} dx = \arcsin(x^2)$$

input `Integrate[(2*x)/Sqrt[1 - x^4],x]`

output `ArcSin[x^2]`

3.27.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {27, 807, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{2x}{\sqrt{1-x^4}} dx \\ & \quad \downarrow \text{27} \\ & 2 \int \frac{x}{\sqrt{1-x^4}} dx \\ & \quad \downarrow \text{807} \\ & \int \frac{1}{\sqrt{1-x^4}} dx^2 \\ & \quad \downarrow \text{223} \\ & \arcsin(x^2) \end{aligned}$$

input `Int[(2*x)/Sqrt[1 - x^4],x]`

output `ArcSin[x^2]`

3.27.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 807 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

3.27.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

method	result	size
default	$\arcsin(x^2)$	5
meijerg	$\arcsin(x^2)$	5
elliptic	$\arcsin(x^2)$	5
pseudoelliptic	$\arcsin(x^2)$	5
trager	$\text{RootOf}(-Z^2 + 1) \ln(\text{RootOf}(-Z^2 + 1) \sqrt{-x^4 + 1} + x^2)$	29

input `int(2*x/(-x^4+1)^(1/2),x,method=_RETURNVERBOSE)`

output `arcsin(x^2)`

3.27.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18 vs. $2(4) = 8$.

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 4.50

$$\int \frac{2x}{\sqrt{1-x^4}} dx = -2 \arctan\left(\frac{\sqrt{-x^4+1}-1}{x^2}\right)$$

input `integrate(2*x/(-x^4+1)^(1/2),x, algorithm="fracas")`

output `-2*arctan((sqrt(-x^4 + 1) - 1)/x^2)`

3.27.6 Sympy [A] (verification not implemented)

Time = 0.46 (sec) , antiderivative size = 20, normalized size of antiderivative = 5.00

$$\int \frac{2x}{\sqrt{1-x^4}} dx = 2 \left(\begin{cases} -\frac{i \operatorname{acosh}(x^2)}{2} & \text{for } |x^4| > 1 \\ \frac{\operatorname{asin}(x^2)}{2} & \text{otherwise} \end{cases} \right)$$

input `integrate(2*x/(-x**4+1)**(1/2),x)`

output `2*Piecewise((-I*acosh(x**2)/2, Abs(x**4) > 1), (asin(x**2)/2, True))`

3.27.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 16 vs. $2(4) = 8$.

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 4.00

$$\int \frac{2x}{\sqrt{1-x^4}} dx = -\arctan\left(\frac{\sqrt{-x^4+1}}{x^2}\right)$$

input `integrate(2*x/(-x^4+1)^(1/2),x, algorithm="maxima")`

output `-arctan(sqrt(-x^4 + 1)/x^2)`

3.27.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{2x}{\sqrt{1-x^4}} dx = \arcsin(x^2)$$

input `integrate(2*x/(-x^4+1)^(1/2),x, algorithm="giac")`

output `arcsin(x^2)`

3.27.9 Mupad [B] (verification not implemented)

Time = 14.90 (sec) , antiderivative size = 14, normalized size of antiderivative = 3.50

$$\int \frac{2x}{\sqrt{1-x^4}} dx = \operatorname{atan}\left(\frac{x^2}{\sqrt{1-x^4}}\right)$$

input `int((2*x)/(1 - x^4)^(1/2),x)`

output `atan(x^2/(1 - x^4)^(1/2))`

3.28 $\int \frac{1+x^2}{1+x} dx$

3.28.1	Optimal result	254
3.28.2	Mathematica [A] (verified)	254
3.28.3	Rubi [A] (verified)	255
3.28.4	Maple [A] (verified)	256
3.28.5	Fricas [A] (verification not implemented)	256
3.28.6	Sympy [A] (verification not implemented)	256
3.28.7	Maxima [A] (verification not implemented)	257
3.28.8	Giac [A] (verification not implemented)	257
3.28.9	Mupad [B] (verification not implemented)	257

3.28.1 Optimal result

Integrand size = 11, antiderivative size = 17

$$\int \frac{1+x^2}{1+x} dx = -x + \frac{x^2}{2} + 2\log(1+x)$$

output `-x+1/2*x^2+2*ln(1+x)`

3.28.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

$$\int \frac{1+x^2}{1+x} dx = \frac{1}{2}(-3 - 2x + x^2 + 4\log(1+x))$$

input `Integrate[(1 + x^2)/(1 + x),x]`

output `(-3 - 2*x + x^2 + 4*Log[1 + x])/2`

3.28.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {476, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2 + 1}{x + 1} dx$$

↓ 476

$$\int \left(x + \frac{2}{x + 1} - 1 \right) dx$$

↓ 2009

$$\frac{x^2}{2} - x + 2 \log(x + 1)$$

input `Int[(1 + x^2)/(1 + x),x]`

output `-x + x^2/2 + 2*Log[1 + x]`

3.28.3.1 Defintions of rubi rules used

rule 476 `Int[((c_) + (d_.)*(x_)^(n_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.28.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

method	result	size
default	$-x + \frac{x^2}{2} + 2 \ln(1+x)$	16
norman	$-x + \frac{x^2}{2} + 2 \ln(1+x)$	16
meijerg	$-\frac{x(-3x+6)}{6} + 2 \ln(1+x)$	16
risch	$-x + \frac{x^2}{2} + 2 \ln(1+x)$	16
parallelrisc	$-x + \frac{x^2}{2} + 2 \ln(1+x)$	16

input `int((x^2+1)/(1+x),x,method=_RETURNVERBOSE)`output `-x+1/2*x^2+2*ln(1+x)`**3.28.5 Fricas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{1+x^2}{1+x} dx = \frac{1}{2}x^2 - x + 2 \log(x+1)$$

input `integrate((x^2+1)/(1+x),x, algorithm="fricas")`output `1/2*x^2 - x + 2*log(x + 1)`**3.28.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int \frac{1+x^2}{1+x} dx = \frac{x^2}{2} - x + 2 \log(x+1)$$

input `integrate((x**2+1)/(1+x),x)`output `x**2/2 - x + 2*log(x + 1)`

3.28.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{1+x^2}{1+x} dx = \frac{1}{2}x^2 - x + 2 \log(x+1)$$

input `integrate((x^2+1)/(1+x),x, algorithm="maxima")`output `1/2*x^2 - x + 2*log(x + 1)`**3.28.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

$$\int \frac{1+x^2}{1+x} dx = \frac{1}{2}x^2 - x + 2 \log(|x+1|)$$

input `integrate((x^2+1)/(1+x),x, algorithm="giac")`output `1/2*x^2 - x + 2*log(abs(x + 1))`**3.28.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{1+x^2}{1+x} dx = 2 \ln(x+1) - x + \frac{x^2}{2}$$

input `int((x^2 + 1)/(x + 1),x)`output `2*log(x + 1) - x + x^2/2`

$$3.29 \quad \int \frac{-2-2\sin(x)+\sin^2(x)+\sin^3(x)}{1+2\sin(x)+\sin^2(x)} dx$$

3.29.1	Optimal result	258
3.29.2	Mathematica [A] (verified)	258
3.29.3	Rubi [A] (verified)	259
3.29.4	Maple [A] (verified)	260
3.29.5	Fricas [A] (verification not implemented)	260
3.29.6	Sympy [B] (verification not implemented)	261
3.29.7	Maxima [B] (verification not implemented)	262
3.29.8	Giac [B] (verification not implemented)	262
3.29.9	Mupad [B] (verification not implemented)	263

3.29.1 Optimal result

Integrand size = 27, antiderivative size = 17

$$\int \frac{-2-2\sin(x)+\sin^2(x)+\sin^3(x)}{1+2\sin(x)+\sin^2(x)} dx = -x - \cos(x) + \frac{\cos(x)}{1+\sin(x)}$$

output `-x-cos(x)+cos(x)/(1+sin(x))`

3.29.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.82

$$\int \frac{-2-2\sin(x)+\sin^2(x)+\sin^3(x)}{1+2\sin(x)+\sin^2(x)} dx = -x - \cos(x) - \frac{2\sin\left(\frac{x}{2}\right)}{\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)}$$

input `Integrate[(-2 - 2*Sin[x] + Sin[x]^2 + Sin[x]^3)/(1 + 2*Sin[x] + Sin[x]^2), x]`

output `-x - Cos[x] - (2*Sin[x/2])/(Cos[x/2] + Sin[x/2])`

3.29.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4901, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sin^3(x) + \sin^2(x) - 2\sin(x) - 2}{\sin^2(x) + 2\sin(x) + 1} dx$$

$$\downarrow 3042$$

$$\int \frac{\sin(x)^3 + \sin(x)^2 - 2\sin(x) - 2}{\sin(x)^2 + 2\sin(x) + 1} dx$$

$$\downarrow 4901$$

$$\int \left(\sin(x) + \frac{1}{-\sin(x) - 1} - 1 \right) dx$$

$$\downarrow 2009$$

$$-x - \cos(x) + \frac{\cos(x)}{\sin(x) + 1}$$

input `Int[(-2 - 2*Sin[x] + Sin[x]^2 + Sin[x]^3)/(1 + 2*Sin[x] + Sin[x]^2),x]`

output `-x - Cos[x] + Cos[x]/(1 + Sin[x])`

3.29.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4901 `Int[u_, x_Symbol] := With[{v = ExpandTrig[u, x]}, Int[v, x] /; SumQ[v]] /; !InertTrigFreeQ[u]`

3.29. $\int \frac{-2-2\sin(x)+\sin^2(x)+\sin^3(x)}{1+2\sin(x)+\sin^2(x)} dx$

3.29.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

method	result
parallelrisc	$-x + \tan(x) (-1 + \sin(x))$
default	$\frac{2}{1+\tan(\frac{x}{2})} - \frac{2}{1+\tan(\frac{x}{2})^2} - 2 \arctan(\tan(\frac{x}{2}))$
risc	$-x - \frac{e^{ix}}{2} - \frac{e^{-ix}}{2} + \frac{2}{i+e^{ix}}$
norman	$\frac{-2 \tan(\frac{x}{2}) - 2 \tan(\frac{x}{2})^2 - 2 \tan(\frac{x}{2})^3 - 2 \tan(\frac{x}{2})^4 + 2 \tan(\frac{x}{2})^8 + 2 \tan(\frac{x}{2})^7 + 2 \tan(\frac{x}{2})^6 + 2 \tan(\frac{x}{2})^5 - x - 3x \tan(\frac{x}{2}) - 6x \tan(\frac{x}{2})^2 - 1}{(1+\tan(\frac{x}{2})^2)^3 (1+\tan(\frac{x}{2}))^3}$

input `int((sin(x)^3+sin(x)^2-2*sin(x)-2)/(1+sin(x)^2+2*sin(x)),x,method=_RETURNV
ERBOSE)`

output `-x+tan(x)*(-1+sin(x))`

3.29.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.71

$$\int \frac{-2 - 2 \sin(x) + \sin^2(x) + \sin^3(x)}{1 + 2 \sin(x) + \sin^2(x)} dx$$

$$= -\frac{x \cos(x) + \cos(x)^2 + (x + \cos(x) + 1) \sin(x) + x - 1}{\cos(x) + \sin(x) + 1}$$

input `integrate((sin(x)^3+sin(x)^2-2*sin(x)-2)/(1+sin(x)^2+2*sin(x)),x, algorithm
m="fricas")`

output `-(x*cos(x) + cos(x)^2 + (x + cos(x) + 1)*sin(x) + x - 1)/(cos(x) + sin(x)
+ 1)`

3.29.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 162 vs. $2(12) = 24$.

Time = 8.20 (sec) , antiderivative size = 162, normalized size of antiderivative = 9.53

$$\int \frac{-2 - 2 \sin(x) + \sin^2(x) + \sin^3(x)}{1 + 2 \sin(x) + \sin^2(x)} dx = -\frac{x \tan^3\left(\frac{x}{2}\right)}{\tan^3\left(\frac{x}{2}\right) + \tan^2\left(\frac{x}{2}\right) + \tan\left(\frac{x}{2}\right) + 1} - \frac{x \tan^2\left(\frac{x}{2}\right)}{\tan^3\left(\frac{x}{2}\right) + \tan^2\left(\frac{x}{2}\right) + \tan\left(\frac{x}{2}\right) + 1} - \frac{x \tan\left(\frac{x}{2}\right)}{\tan^3\left(\frac{x}{2}\right) + \tan^2\left(\frac{x}{2}\right) + \tan\left(\frac{x}{2}\right) + 1} - \frac{x}{\tan^3\left(\frac{x}{2}\right) + \tan^2\left(\frac{x}{2}\right) + \tan\left(\frac{x}{2}\right) + 1} + \frac{2 \tan^2\left(\frac{x}{2}\right)}{\tan^3\left(\frac{x}{2}\right) + \tan^2\left(\frac{x}{2}\right) + \tan\left(\frac{x}{2}\right) + 1} - \frac{2 \tan\left(\frac{x}{2}\right)}{\tan^3\left(\frac{x}{2}\right) + \tan^2\left(\frac{x}{2}\right) + \tan\left(\frac{x}{2}\right) + 1}$$

input `integrate((sin(x)**3+sin(x)**2-2*sin(x)-2)/(1+sin(x)**2+2*sin(x)),x)`

output `-x*tan(x/2)**3/(tan(x/2)**3 + tan(x/2)**2 + tan(x/2) + 1) - x*tan(x/2)**2/(tan(x/2)**3 + tan(x/2)**2 + tan(x/2) + 1) - x*tan(x/2)/(tan(x/2)**3 + tan(x/2)**2 + tan(x/2) + 1) - x/(tan(x/2)**3 + tan(x/2)**2 + tan(x/2) + 1) + 2*tan(x/2)**2/(tan(x/2)**3 + tan(x/2)**2 + tan(x/2) + 1) - 2*tan(x/2)/(tan(x/2)**3 + tan(x/2)**2 + tan(x/2) + 1)`

3.29.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 301 vs. $2(17) = 34$.

Time = 0.28 (sec) , antiderivative size = 301, normalized size of antiderivative = 17.71

$$\int \frac{-2 - 2 \sin(x) + \sin^2(x) + \sin^3(x)}{1 + 2 \sin(x) + \sin^2(x)} dx$$

$$= -\frac{4 \left(\frac{12 \sin(x)}{\cos(x)+1} + \frac{11 \sin(x)^2}{(\cos(x)+1)^2} + \frac{9 \sin(x)^3}{(\cos(x)+1)^3} + \frac{3 \sin(x)^4}{(\cos(x)+1)^4} + 5 \right)}{3 \left(\frac{3 \sin(x)}{\cos(x)+1} + \frac{4 \sin(x)^2}{(\cos(x)+1)^2} + \frac{4 \sin(x)^3}{(\cos(x)+1)^3} + \frac{3 \sin(x)^4}{(\cos(x)+1)^4} + \frac{\sin(x)^5}{(\cos(x)+1)^5} + 1 \right)}$$

$$+ \frac{2 \left(\frac{9 \sin(x)}{\cos(x)+1} + \frac{3 \sin(x)^2}{(\cos(x)+1)^2} + 4 \right)}{3 \left(\frac{3 \sin(x)}{\cos(x)+1} + \frac{3 \sin(x)^2}{(\cos(x)+1)^2} + \frac{\sin(x)^3}{(\cos(x)+1)^3} + 1 \right)} + \frac{4 \left(\frac{3 \sin(x)}{\cos(x)+1} + \frac{3 \sin(x)^2}{(\cos(x)+1)^2} + 2 \right)}{3 \left(\frac{3 \sin(x)}{\cos(x)+1} + \frac{3 \sin(x)^2}{(\cos(x)+1)^2} + \frac{\sin(x)^3}{(\cos(x)+1)^3} + 1 \right)}$$

$$+ \frac{4 \left(\frac{3 \sin(x)}{\cos(x)+1} + 1 \right)}{3 \left(\frac{3 \sin(x)}{\cos(x)+1} + \frac{3 \sin(x)^2}{(\cos(x)+1)^2} + \frac{\sin(x)^3}{(\cos(x)+1)^3} + 1 \right)} - 2 \arctan \left(\frac{\sin(x)}{\cos(x)+1} \right)$$

input `integrate((sin(x)^3+sin(x)^2-2*sin(x)-2)/(1+sin(x)^2+2*sin(x)),x, algorithm="maxima")`

output `-4/3*(12*sin(x)/(cos(x) + 1) + 11*sin(x)^2/(cos(x) + 1)^2 + 9*sin(x)^3/(cos(x) + 1)^3 + 3*sin(x)^4/(cos(x) + 1)^4 + 5)/(3*sin(x)/(cos(x) + 1) + 4*sin(x)^2/(cos(x) + 1)^2 + 4*sin(x)^3/(cos(x) + 1)^3 + 3*sin(x)^4/(cos(x) + 1)^4 + sin(x)^5/(cos(x) + 1)^5 + 1) + 2/3*(9*sin(x)/(cos(x) + 1) + 3*sin(x)^2/(cos(x) + 1)^2 + 4)/(3*sin(x)/(cos(x) + 1) + 3*sin(x)^2/(cos(x) + 1)^2 + sin(x)^3/(cos(x) + 1)^3 + 1) + 4/3*(3*sin(x)/(cos(x) + 1) + 3*sin(x)^2/(cos(x) + 1)^2 + 2)/(3*sin(x)/(cos(x) + 1) + 3*sin(x)^2/(cos(x) + 1)^2 + sin(x)^3/(cos(x) + 1)^3 + 1) + 4/3*(3*sin(x)/(cos(x) + 1) + 1)/(3*sin(x)/(cos(x) + 1) + 3*sin(x)^2/(cos(x) + 1)^2 + sin(x)^3/(cos(x) + 1)^3 + 1) - 2*arctan(sin(x)/(cos(x) + 1))`

3.29.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. $2(17) = 34$.

Time = 0.28 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.29

$$\int \frac{-2 - 2 \sin(x) + \sin^2(x) + \sin^3(x)}{1 + 2 \sin(x) + \sin^2(x)} dx = -x + \frac{2 \left(\tan \left(\frac{1}{2} x \right)^2 - \tan \left(\frac{1}{2} x \right) \right)}{\tan \left(\frac{1}{2} x \right)^3 + \tan \left(\frac{1}{2} x \right)^2 + \tan \left(\frac{1}{2} x \right) + 1}$$

input `integrate((sin(x)^3+sin(x)^2-2*sin(x)-2)/(1+sin(x)^2+2*sin(x)),x, algorithm="giac")`

output `-x + 2*(tan(1/2*x)^2 - tan(1/2*x))/(tan(1/2*x)^3 + tan(1/2*x)^2 + tan(1/2*x) + 1)`

3.29.9 Mupad [B] (verification not implemented)

Time = 14.99 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.29

$$\int \frac{-2 - 2 \sin(x) + \sin^2(x) + \sin^3(x)}{1 + 2 \sin(x) + \sin^2(x)} dx = -x - \frac{2 \tan\left(\frac{x}{2}\right) - 2 \tan\left(\frac{x}{2}\right)^2}{\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) \left(\tan\left(\frac{x}{2}\right) + 1\right)}$$

input `int(-(2*sin(x) - sin(x)^2 - sin(x)^3 + 2)/(2*sin(x) + sin(x)^2 + 1),x)`

output `- x - (2*tan(x/2) - 2*tan(x/2)^2)/((tan(x/2)^2 + 1)*(tan(x/2) + 1))`

3.30 $\int \operatorname{csch}^2(x) dx$

3.30.1	Optimal result	264
3.30.2	Mathematica [A] (verified)	264
3.30.3	Rubi [A] (verified)	265
3.30.4	Maple [A] (verified)	266
3.30.5	Fricas [B] (verification not implemented)	266
3.30.6	Sympy [B] (verification not implemented)	267
3.30.7	Maxima [B] (verification not implemented)	267
3.30.8	Giac [B] (verification not implemented)	267
3.30.9	Mupad [B] (verification not implemented)	268

3.30.1 Optimal result

Integrand size = 4, antiderivative size = 4

$$\int \operatorname{csch}^2(x) dx = -\operatorname{coth}(x)$$

output `-coth(x)`

3.30.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \operatorname{csch}^2(x) dx = -\operatorname{coth}(x)$$

input `Integrate[Csch[x]^2,x]`

output `-Coth[x]`

3.30.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 25, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \operatorname{csch}^2(x) dx \\
 \downarrow 3042 \\
 \int -\operatorname{csc}(ix)^2 dx \\
 \downarrow 25 \\
 -\int \operatorname{csc}(ix)^2 dx \\
 \downarrow 4254 \\
 -i \int 1d(-i \operatorname{coth}(x)) \\
 \downarrow 24 \\
 -\operatorname{coth}(x)
 \end{array}$$

input `Int [Csch[x]^2,x]`

output `-Coth[x]`

3.30.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

```
rule 4254 Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

3.30.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

method	result	size
default	$-\coth(x)$	5
parallelsch	$-\coth(x)$	5
risch	$-\frac{2}{e^{2x}-1}$	11

```
input int(1/sinh(x)^2,x,method=_RETURNVERBOSE)
```

```
output -coth(x)
```

3.30.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. $2(4) = 8$.

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 5.00

$$\int \operatorname{csch}^2(x) dx = -\frac{2}{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1}$$

```
input integrate(1/sinh(x)^2,x, algorithm="fricas")
```

```
output -2/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)
```

3.30.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(3) = 6$.

Time = 0.23 (sec) , antiderivative size = 15, normalized size of antiderivative = 3.75

$$\int \operatorname{csch}^2(x) dx = -\frac{\tanh\left(\frac{x}{2}\right)}{2} - \frac{1}{2 \tanh\left(\frac{x}{2}\right)}$$

input `integrate(1/sinh(x)**2,x)`

output `-tanh(x/2)/2 - 1/(2*tanh(x/2))`

3.30.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10 vs. $2(4) = 8$.

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 2.50

$$\int \operatorname{csch}^2(x) dx = \frac{2}{e^{(-2x)} - 1}$$

input `integrate(1/sinh(x)^2,x, algorithm="maxima")`

output `2/(e^(-2*x) - 1)`

3.30.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10 vs. $2(4) = 8$.

Time = 0.25 (sec) , antiderivative size = 10, normalized size of antiderivative = 2.50

$$\int \operatorname{csch}^2(x) dx = -\frac{2}{e^{(2x)} - 1}$$

input `integrate(1/sinh(x)^2,x, algorithm="giac")`

output `-2/(e^(2*x) - 1)`

3.30.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \operatorname{csch}^2(x) dx = -\operatorname{coth}(x)$$

input `int(1/sinh(x)^2,x)`

output `-coth(x)`

3.31 $\int \sec^4(x) \tan^2(x) dx$

3.31.1	Optimal result	269
3.31.2	Mathematica [A] (verified)	269
3.31.3	Rubi [A] (verified)	270
3.31.4	Maple [A] (verified)	271
3.31.5	Fricas [A] (verification not implemented)	271
3.31.6	Sympy [B] (verification not implemented)	272
3.31.7	Maxima [A] (verification not implemented)	272
3.31.8	Giac [A] (verification not implemented)	272
3.31.9	Mupad [B] (verification not implemented)	273

3.31.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \sec^4(x) \tan^2(x) dx = \frac{\tan^3(x)}{3} + \frac{\tan^5(x)}{5}$$

output `1/5*tan(x)^5+1/3*tan(x)^3`

3.31.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \sec^4(x) \tan^2(x) dx = -\frac{2 \tan(x)}{15} - \frac{1}{15} \sec^2(x) \tan(x) + \frac{1}{5} \sec^4(x) \tan(x)$$

input `Integrate[Sec[x]^4*Tan[x]^2,x]`

output `(-2*Tan[x])/15 - (Sec[x]^2*Tan[x])/15 + (Sec[x]^4*Tan[x])/5`

3.31.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 3087, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^2(x) \sec^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^2 \sec(x)^4 dx \\
 & \quad \downarrow \text{3087} \\
 & \int \tan^2(x) (\tan^2(x) + 1) d \tan(x) \\
 & \quad \downarrow \text{244} \\
 & \int (\tan^4(x) + \tan^2(x)) d \tan(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\tan^5(x)}{5} + \frac{\tan^3(x)}{3}
 \end{aligned}$$

input `Int[Sec[x]^4*Tan[x]^2,x]`

output `Tan[x]^3/3 + Tan[x]^5/5`

3.31.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] :> Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3087 `Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[1/f Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])`

3.31.4 Maple [A] (verified)

Time = 1.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$\frac{\tan(x)^5}{5} + \frac{\tan(x)^3}{3}$	14
default	$\frac{\tan(x)^5}{5} + \frac{\tan(x)^3}{3}$	14
risch	$-\frac{4i(15e^{6ix} - 5e^{4ix} + 5e^{2ix} + 1)}{15(e^{2ix} + 1)^5}$	36

input `int(sec(x)^4*tan(x)^2,x,method=_RETURNVERBOSE)`

output `1/5*tan(x)^5+1/3*tan(x)^3`

3.31.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \sec^4(x) \tan^2(x) dx = -\frac{(2 \cos(x)^4 + \cos(x)^2 - 3) \sin(x)}{15 \cos(x)^5}$$

input `integrate(sec(x)^4*tan(x)^2,x, algorithm="fricas")`

output `-1/15*(2*cos(x)^4 + cos(x)^2 - 3)*sin(x)/cos(x)^5`

3.31.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(12) = 24$.

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.71

$$\int \sec^4(x) \tan^2(x) dx = -\frac{2 \sin(x)}{15 \cos(x)} - \frac{\sin(x)}{15 \cos^3(x)} + \frac{\sin(x)}{5 \cos^5(x)}$$

input `integrate(sec(x)**4*tan(x)**2,x)`

output `-2*sin(x)/(15*cos(x)) - sin(x)/(15*cos(x)**3) + sin(x)/(5*cos(x)**5)`

3.31.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sec^4(x) \tan^2(x) dx = \frac{1}{5} \tan(x)^5 + \frac{1}{3} \tan(x)^3$$

input `integrate(sec(x)^4*tan(x)^2,x, algorithm="maxima")`

output `1/5*tan(x)^5 + 1/3*tan(x)^3`

3.31.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sec^4(x) \tan^2(x) dx = \frac{1}{5} \tan(x)^5 + \frac{1}{3} \tan(x)^3$$

input `integrate(sec(x)^4*tan(x)^2,x, algorithm="giac")`

output `1/5*tan(x)^5 + 1/3*tan(x)^3`

3.31.9 Mupad [B] (verification not implemented)

Time = 14.96 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sec^4(x) \tan^2(x) dx = \frac{\tan(x)^5}{5} + \frac{\tan(x)^3}{3}$$

input `int(tan(x)^2/cos(x)^4,x)`

output `tan(x)^3/3 + tan(x)^5/5`

3.32 $\int \sqrt{\csc(x) - \sin(x)} dx$

3.32.1	Optimal result	274
3.32.2	Mathematica [A] (verified)	274
3.32.3	Rubi [A] (verified)	275
3.32.4	Maple [A] (verified)	276
3.32.5	Fricas [A] (verification not implemented)	277
3.32.6	Sympy [F]	277
3.32.7	Maxima [B] (verification not implemented)	277
3.32.8	Giac [F]	278
3.32.9	Mupad [B] (verification not implemented)	278

3.32.1 Optimal result

Integrand size = 11, antiderivative size = 13

$$\int \sqrt{\csc(x) - \sin(x)} dx = 2\sqrt{\cos(x) \cot(x)} \tan(x)$$

output `2*(cos(x)*cot(x))^(1/2)*tan(x)`

3.32.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \sqrt{\csc(x) - \sin(x)} dx = 2\sqrt{\cos(x) \cot(x)} \tan(x)$$

input `Integrate[Sqrt[Csc[x] - Sin[x]],x]`

output `2*Sqrt[Cos[x]*Cot[x]]*Tan[x]`

3.32.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 4897, 3042, 4900, 3042, 3069}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\csc(x) - \sin(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\csc(x) - \sin(x)} dx \\
 & \quad \downarrow \text{4897} \\
 & \int \sqrt{\cos(x) \cot(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\cos(x) \cot(x)} dx \\
 & \quad \downarrow \text{4900} \\
 & \frac{\sqrt{\cos(x) \cot(x)} \int \sqrt{\cos(x)} \sqrt{\cot(x)} dx}{\sqrt{\cos(x)} \sqrt{\cot(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\cos(x) \cot(x)} \int \sqrt{\sin(x + \frac{\pi}{2})} \sqrt{-\tan(x + \frac{\pi}{2})} dx}{\sqrt{\cos(x)} \sqrt{\cot(x)}} \\
 & \quad \downarrow \text{3069} \\
 & 2 \tan(x) \sqrt{\cos(x) \cot(x)}
 \end{aligned}$$

input `Int[Sqrt[Csc[x] - Sin[x]],x]`

output `2*Sqrt[Cos[x]*Cot[x]]*Tan[x]`

3.32.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3069 `Int[((a_.)*sin[(e_.) + (f_.)*(x_)]^(m_))*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp[(-b)*(a*Sin[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*m)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 1, 0]`

rule 4897 `Int[u_, x_Symbol] := Int[TrigSimplify[u], x] /; TrigSimplifyQ[u]`

rule 4900 `Int[(u_.)*((v_)^(m_.)*(w_)^(n_.))^p, x_Symbol] := With[{uu = ActivateTrig[u], vv = ActivateTrig[v], ww = ActivateTrig[w]}, Simp[(vv^m*ww^n)^FracPart[p]/(vv^(m*FracPart[p])*ww^(n*FracPart[p])) Int[uu*vv^(m*p)*ww^(n*p), x], x] /; FreeQ[{m, n, p}, x] && !IntegerQ[p] && (!InertTrigFreeQ[v] || !InertTrigFreeQ[w])`

3.32.4 Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
default	$2\sqrt{\cos(x)\cot(x)}\tan(x)$	12
risch	$-\frac{i\sqrt{2}\sqrt{\frac{i(e^{2ix}+1)^2e^{-ix}}{e^{2ix}-1}}(e^{2ix}-1)}{e^{2ix}+1}$	51

input `int((csc(x)-sin(x))^(1/2),x,method=_RETURNVERBOSE)`

output `2*(cos(x)*cot(x))^(1/2)*tan(x)`

3.32.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.46

$$\int \sqrt{\csc(x) - \sin(x)} dx = \frac{2 \sqrt{\frac{\cos(x)^2}{\sin(x)}} \sin(x)}{\cos(x)}$$

input `integrate((csc(x)-sin(x))^(1/2),x, algorithm="fracas")`

output `2*sqrt(cos(x)^2/sin(x))*sin(x)/cos(x)`

3.32.6 Sympy [F]

$$\int \sqrt{\csc(x) - \sin(x)} dx = \int \sqrt{-\sin(x) + \csc(x)} dx$$

input `integrate((csc(x)-sin(x))**(1/2),x)`

output `Integral(sqrt(-sin(x) + csc(x)), x)`

3.32.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 188 vs. $2(11) = 22$.

Time = 0.37 (sec) , antiderivative size = 188, normalized size of antiderivative = 14.46

$$\int \sqrt{\csc(x) - \sin(x)} dx = \frac{((\cos(\frac{3}{2}x) - \cos(\frac{1}{2}x) + \sin(\frac{3}{2}x) + \sin(\frac{1}{2}x)) \cos(\frac{1}{2} \arctan(\sin(x), \cos(x) - 1)) - (\cos(\frac{3}{2}x) - \cos(\frac{1}{2}x)))}{\cos(x)}$$

input `integrate((csc(x)-sin(x))^(1/2),x, algorithm="maxima")`

output `((cos(3/2*x) - cos(1/2*x) + sin(3/2*x) + sin(1/2*x))*cos(1/2*arctan2(sin(x), cos(x) - 1)) - (cos(3/2*x) - cos(1/2*x) - sin(3/2*x) - sin(1/2*x))*sin(1/2*arctan2(sin(x), cos(x) - 1)))*cos(1/2*arctan2(sin(x), cos(x) + 1)) - ((cos(3/2*x) - cos(1/2*x) - sin(3/2*x) - sin(1/2*x))*cos(1/2*arctan2(sin(x), cos(x) - 1)) + (cos(3/2*x) - cos(1/2*x) + sin(3/2*x) + sin(1/2*x))*sin(1/2*arctan2(sin(x), cos(x) - 1)))*sin(1/2*arctan2(sin(x), cos(x) + 1)))/((cos(x)^2 + sin(x)^2 + 2*cos(x) + 1)^(1/4)*(cos(x)^2 + sin(x)^2 - 2*cos(x) + 1)^(1/4))`

3.32.8 Giac [F]

$$\int \sqrt{\csc(x) - \sin(x)} dx = \int \sqrt{\csc(x) - \sin(x)} dx$$

input `integrate((csc(x)-sin(x))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(csc(x) - sin(x)), x)`

3.32.9 Mupad [B] (verification not implemented)

Time = 15.46 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \sqrt{\csc(x) - \sin(x)} dx = \frac{2 |\cos(x)|}{\cos(x) \sqrt{\frac{1}{\sin(x)}}}$$

input `int((1/sin(x) - sin(x))^(1/2),x)`

output `(2*abs(cos(x)))/(cos(x)*(1/sin(x))^(1/2))`

3.33 $\int \cos^6(x) dx$

3.33.1	Optimal result	279
3.33.2	Mathematica [A] (verified)	279
3.33.3	Rubi [A] (verified)	280
3.33.4	Maple [A] (verified)	281
3.33.5	Fricas [A] (verification not implemented)	282
3.33.6	Sympy [A] (verification not implemented)	282
3.33.7	Maxima [A] (verification not implemented)	282
3.33.8	Giac [A] (verification not implemented)	283
3.33.9	Mupad [B] (verification not implemented)	283

3.33.1 Optimal result

Integrand size = 4, antiderivative size = 34

$$\int \cos^6(x) dx = \frac{5x}{16} + \frac{5}{16} \cos(x) \sin(x) + \frac{5}{24} \cos^3(x) \sin(x) + \frac{1}{6} \cos^5(x) \sin(x)$$

output `5/16*x+5/16*cos(x)*sin(x)+5/24*cos(x)^3*sin(x)+1/6*cos(x)^5*sin(x)`

3.33.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.88

$$\int \cos^6(x) dx = \frac{5x}{16} + \frac{15}{64} \sin(2x) + \frac{3}{64} \sin(4x) + \frac{1}{192} \sin(6x)$$

input `Integrate[Cos[x]^6,x]`

output `(5*x)/16 + (15*Sin[2*x])/64 + (3*Sin[4*x])/64 + Sin[6*x]/192`

3.33.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.29, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.750$, Rules used = {3042, 3115, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^6(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(x + \frac{\pi}{2}\right)^6 dx \\
 & \quad \downarrow \text{3115} \\
 & \frac{5}{6} \int \cos^4(x) dx + \frac{1}{6} \sin(x) \cos^5(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{6} \int \sin\left(x + \frac{\pi}{2}\right)^4 dx + \frac{1}{6} \sin(x) \cos^5(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{5}{6} \left(\frac{3}{4} \int \cos^2(x) dx + \frac{1}{4} \sin(x) \cos^3(x) \right) + \frac{1}{6} \sin(x) \cos^5(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{6} \left(\frac{3}{4} \int \sin\left(x + \frac{\pi}{2}\right)^2 dx + \frac{1}{4} \sin(x) \cos^3(x) \right) + \frac{1}{6} \sin(x) \cos^5(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{5}{6} \left(\frac{3}{4} \left(\frac{\int 1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \right) + \frac{1}{4} \sin(x) \cos^3(x) \right) + \frac{1}{6} \sin(x) \cos^5(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{1}{6} \sin(x) \cos^5(x) + \frac{5}{6} \left(\frac{1}{4} \sin(x) \cos^3(x) + \frac{3}{4} \left(\frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \right) \right)
 \end{aligned}$$

input `Int[Cos[x]^6,x]`

output $(\cos[x]^5 \sin[x])/6 + (5*((\cos[x]^3 \sin[x])/4 + (3*(x/2 + (\cos[x] \sin[x])/2))/4))/6$

3.33.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sine[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sine[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

3.33.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.68

method	result
risch	$\frac{5x}{16} + \frac{\sin(6x)}{192} + \frac{3 \sin(4x)}{64} + \frac{15 \sin(2x)}{64}$
parallelrisc	$\frac{5x}{16} + \frac{\sin(6x)}{192} + \frac{3 \sin(4x)}{64} + \frac{15 \sin(2x)}{64}$
default	$\frac{\left(\cos(x)^5 + \frac{5 \cos(x)^3}{4} + \frac{15 \cos(x)}{8}\right) \sin(x)}{6} + \frac{5x}{16}$
norman	$\frac{5x}{16} - \frac{5 \tan\left(\frac{x}{2}\right)^3}{24} + \frac{15 \tan\left(\frac{x}{2}\right)^5}{4} - \frac{15 \tan\left(\frac{x}{2}\right)^7}{4} + \frac{5 \tan\left(\frac{x}{2}\right)^9}{24} - \frac{11 \tan\left(\frac{x}{2}\right)^{11}}{8} + \frac{15x \tan\left(\frac{x}{2}\right)^2}{8} + \frac{75x \tan\left(\frac{x}{2}\right)^4}{16} + \frac{25x \tan\left(\frac{x}{2}\right)^6}{4} + \frac{75x \tan\left(\frac{x}{2}\right)^8}{16} + \frac{1}{\left(1 + \tan\left(\frac{x}{2}\right)^2\right)^6}$

input `int(cos(x)^6,x,method=_RETURNVERBOSE)`

output $5/16*x+1/192*\sin(6*x)+3/64*\sin(4*x)+15/64*\sin(2*x)$

3.33.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.74

$$\int \cos^6(x) dx = \frac{1}{48} (8 \cos(x)^5 + 10 \cos(x)^3 + 15 \cos(x)) \sin(x) + \frac{5}{16} x$$

input `integrate(cos(x)^6,x, algorithm="fricas")`

output `1/48*(8*cos(x)^5 + 10*cos(x)^3 + 15*cos(x))*sin(x) + 5/16*x`

3.33.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.06

$$\int \cos^6(x) dx = \frac{5x}{16} + \frac{\sin(x) \cos^5(x)}{6} + \frac{5 \sin(x) \cos^3(x)}{24} + \frac{5 \sin(x) \cos(x)}{16}$$

input `integrate(cos(x)**6,x)`

output `5*x/16 + sin(x)*cos(x)**5/6 + 5*sin(x)*cos(x)**3/24 + 5*sin(x)*cos(x)/16`

3.33.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.71

$$\int \cos^6(x) dx = -\frac{1}{48} \sin(2x)^3 + \frac{5}{16} x + \frac{3}{64} \sin(4x) + \frac{1}{4} \sin(2x)$$

input `integrate(cos(x)^6,x, algorithm="maxima")`

output `-1/48*sin(2*x)^3 + 5/16*x + 3/64*sin(4*x) + 1/4*sin(2*x)`

3.33.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.65

$$\int \cos^6(x) dx = \frac{5}{16} x + \frac{1}{192} \sin(6x) + \frac{3}{64} \sin(4x) + \frac{15}{64} \sin(2x)$$

input `integrate(cos(x)^6,x, algorithm="giac")`

output `5/16*x + 1/192*sin(6*x) + 3/64*sin(4*x) + 15/64*sin(2*x)`

3.33.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.65

$$\int \cos^6(x) dx = \frac{5x}{16} + \frac{15 \sin(2x)}{64} + \frac{3 \sin(4x)}{64} + \frac{\sin(6x)}{192}$$

input `int(cos(x)^6,x)`

output `(5*x)/16 + (15*sin(2*x))/64 + (3*sin(4*x))/64 + sin(6*x)/192`

3.34 $\int \frac{1}{1+2x^2+x^4} dx$

3.34.1	Optimal result	284
3.34.2	Mathematica [A] (verified)	284
3.34.3	Rubi [A] (verified)	285
3.34.4	Maple [A] (verified)	286
3.34.5	Fricas [A] (verification not implemented)	286
3.34.6	Sympy [A] (verification not implemented)	287
3.34.7	Maxima [A] (verification not implemented)	287
3.34.8	Giac [A] (verification not implemented)	287
3.34.9	Mupad [B] (verification not implemented)	288

3.34.1 Optimal result

Integrand size = 12, antiderivative size = 19

$$\int \frac{1}{1+2x^2+x^4} dx = \frac{x}{2(1+x^2)} + \frac{\arctan(x)}{2}$$

output `x/(2*x^2+2)+1/2*arctan(x)`

3.34.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{1}{1+2x^2+x^4} dx = \frac{1}{2} \left(\frac{x}{1+x^2} + \arctan(x) \right)$$

input `Integrate[(1 + 2*x^2 + x^4)^(-1),x]`

output `(x/(1 + x^2) + ArcTan[x])/2`

3.34.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {1379, 215, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x^4 + 2x^2 + 1} dx \\ & \quad \downarrow \text{1379} \\ & \int \frac{1}{(x^2 + 1)^2} dx \\ & \quad \downarrow \text{215} \\ & \frac{1}{2} \int \frac{1}{x^2 + 1} dx + \frac{x}{2(x^2 + 1)} \\ & \quad \downarrow \text{216} \\ & \frac{\arctan(x)}{2} + \frac{x}{2(x^2 + 1)} \end{aligned}$$

input `Int[(1 + 2*x^2 + x^4)^(-1),x]`

output `x/(2*(1 + x^2)) + ArcTan[x]/2`

3.34.3.1 Defintions of rubi rules used

rule 215 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

```
rule 1379 Int[((a_) + (c_.)*(x_)^(n2_.) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[1/
c^p Int[(b/2 + c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n
2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p] && NeQ[p, 1]
```

3.34.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

method	result	size
default	$\frac{x}{2x^2+2} + \frac{\arctan(x)}{2}$	16
risch	$\frac{x}{2x^2+2} + \frac{\arctan(x)}{2}$	16
parallelrisch	$-\frac{i \ln(x-i)x^2 - i \ln(i+x)x^2 + i \ln(x-i) - i \ln(i+x) - 2x}{4(x^2+1)}$	52

```
input int(1/(x^4+2*x^2+1),x,method=_RETURNVERBOSE)
```

```
output 1/2/(x^2+1)*x+1/2*arctan(x)
```

3.34.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+2x^2+x^4} dx = \frac{(x^2+1)\arctan(x)+x}{2(x^2+1)}$$

```
input integrate(1/(x^4+2*x^2+1),x, algorithm="fracas")
```

```
output 1/2*((x^2 + 1)*arctan(x) + x)/(x^2 + 1)
```

3.34.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{1}{1+2x^2+x^4} dx = \frac{x}{2x^2+2} + \frac{\operatorname{atan}(x)}{2}$$

input `integrate(1/(x**4+2*x**2+1),x)`output `x/(2*x**2 + 2) + atan(x)/2`**3.34.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{1+2x^2+x^4} dx = \frac{x}{2(x^2+1)} + \frac{1}{2} \arctan(x)$$

input `integrate(1/(x^4+2*x^2+1),x, algorithm="maxima")`output `1/2*x/(x^2 + 1) + 1/2*arctan(x)`**3.34.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{1+2x^2+x^4} dx = \frac{x}{2(x^2+1)} + \frac{1}{2} \arctan(x)$$

input `integrate(1/(x^4+2*x^2+1),x, algorithm="giac")`output `1/2*x/(x^2 + 1) + 1/2*arctan(x)`

3.34.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{1}{1+2x^2+x^4} dx = \frac{\operatorname{atan}(x)}{2} + \frac{x}{2(x^2+1)}$$

input `int(1/(2*x^2 + x^4 + 1),x)`

output `atan(x)/2 + x/(2*(x^2 + 1))`

3.35 $\int \cos(\log(x)) dx$

3.35.1	Optimal result	289
3.35.2	Mathematica [A] (verified)	289
3.35.3	Rubi [A] (verified)	290
3.35.4	Maple [A] (verified)	290
3.35.5	Fricas [A] (verification not implemented)	291
3.35.6	Sympy [A] (verification not implemented)	291
3.35.7	Maxima [A] (verification not implemented)	291
3.35.8	Giac [A] (verification not implemented)	292
3.35.9	Mupad [B] (verification not implemented)	292

3.35.1 Optimal result

Integrand size = 3, antiderivative size = 17

$$\int \cos(\log(x)) dx = \frac{1}{2}x \cos(\log(x)) + \frac{1}{2}x \sin(\log(x))$$

output `1/2*x*cos(ln(x))+1/2*x*sin(ln(x))`

3.35.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cos(\log(x)) dx = \frac{1}{2}x \cos(\log(x)) + \frac{1}{2}x \sin(\log(x))$$

input `Integrate[Cos[Log[x]],x]`

output `(x*Cos[Log[x]])/2 + (x*Sin[Log[x]])/2`

3.35.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {4979}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(\log(x)) dx$$

↓ 4979

$$\frac{1}{2}x \sin(\log(x)) + \frac{1}{2}x \cos(\log(x))$$

input `Int[Cos[Log[x]],x]`

output `(x*Cos[Log[x]])/2 + (x*Sin[Log[x]])/2`

3.35.3.1 Defintions of rubi rules used

rule 4979 `Int[Cos[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)], x_Symbol] := Simp[x*(Cos[d*(a + b*Log[c*x^n])]/(b^2*d^2*n^2 + 1)), x] + Simp[b*d*n*x*(Sin[d*(a + b*Log[c*x^n])]/(b^2*d^2*n^2 + 1)), x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b^2*d^2*n^2 + 1, 0]`

3.35.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

method	result	size
parallelrisch	$\frac{x(\cos(\ln(x)) + \sin(\ln(x)))}{2}$	11
lookup	$\frac{x \cos(\ln(x))}{2} + \frac{x \sin(\ln(x))}{2}$	14
default	$\frac{x \cos(\ln(x))}{2} + \frac{x \sin(\ln(x))}{2}$	14
risch	$(\frac{1}{4} - \frac{i}{4}) x x^i + (\frac{1}{4} + \frac{i}{4}) x x^{-i}$	22

input `int(cos(ln(x)),x,method=_RETURNVERBOSE)`

output `1/2*x*(cos(ln(x))+sin(ln(x)))`

3.35.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(\log(x)) dx = \frac{1}{2} x \cos(\log(x)) + \frac{1}{2} x \sin(\log(x))$$

input `integrate(cos(log(x)),x, algorithm="fricas")`

output `1/2*x*cos(log(x)) + 1/2*x*sin(log(x))`

3.35.6 Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \cos(\log(x)) dx = \frac{x \sin(\log(x))}{2} + \frac{x \cos(\log(x))}{2}$$

input `integrate(cos(ln(x)),x)`

output `x*sin(log(x))/2 + x*cos(log(x))/2`

3.35.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.59

$$\int \cos(\log(x)) dx = \frac{1}{2} x(\cos(\log(x)) + \sin(\log(x)))$$

input `integrate(cos(log(x)),x, algorithm="maxima")`

output `1/2*x*(cos(log(x)) + sin(log(x)))`

3.35.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(\log(x)) dx = \frac{1}{2} x \cos(\log(x)) + \frac{1}{2} x \sin(\log(x))$$

input `integrate(cos(log(x)),x, algorithm="giac")`

output `1/2*x*cos(log(x)) + 1/2*x*sin(log(x))`

3.35.9 Mupad [B] (verification not implemented)

Time = 14.86 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \cos(\log(x)) dx = \frac{\sqrt{2} x \sin\left(\frac{\pi}{4} + \ln(x)\right)}{2}$$

input `int(cos(log(x)),x)`

output `(2^(1/2)*x*sin(pi/4 + log(x)))/2`

3.36 $\int \sec(x) dx$

3.36.1	Optimal result	293
3.36.2	Mathematica [A] (verified)	293
3.36.3	Rubi [A] (verified)	294
3.36.4	Maple [A] (verified)	295
3.36.5	Fricas [B] (verification not implemented)	295
3.36.6	Sympy [B] (verification not implemented)	295
3.36.7	Maxima [B] (verification not implemented)	296
3.36.8	Giac [B] (verification not implemented)	296
3.36.9	Mupad [B] (verification not implemented)	296

3.36.1 Optimal result

Integrand size = 2, antiderivative size = 3

$$\int \sec(x) dx = \operatorname{arctanh}(\sin(x))$$

output `arctanh(sin(x))`

3.36.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \sec(x) dx = \operatorname{arctanh}(\sin(x))$$

input `Integrate[Sec[x],x]`

output `ArcTanh[Sin[x]]`

3.36.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sec(x) dx \\ \downarrow 3042 \\ \int \csc\left(x + \frac{\pi}{2}\right) dx \\ \downarrow 4257 \\ \operatorname{arctanh}(\sin(x)) \end{array}$$

input `Int [Sec [x] , x]`

output `ArcTanh [Sin [x]]`

3.36.3.1 Defintions of rubi rules used

rule 3042 `Int [u_ , x_Symbol] := Int [DeactivateTrig [u , x] , x] /; FunctionOfTrigOfLinear Q [u , x]`

rule 4257 `Int [csc [(c_) + (d_)*(x_)] , x_Symbol] := Simp [-ArcTanh [Cos [c + d*x]]/d , x] /; FreeQ [{c , d} , x]`

3.36.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 7, normalized size of antiderivative = 2.33

method	result	size
default	$\ln(\sec(x) + \tan(x))$	7
norman	$-\ln(\tan(\frac{x}{2}) - 1) + \ln(1 + \tan(\frac{x}{2}))$	18
parallelrisc	$-\ln(\tan(\frac{x}{2}) - 1) + \ln(1 + \tan(\frac{x}{2}))$	18
risc	$\ln(i + e^{ix}) - \ln(e^{ix} - i)$	22

input `int(1/cos(x),x,method=_RETURNVERBOSE)`

output `ln(sec(x)+tan(x))`

3.36.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(3) = 6$.

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 5.67

$$\int \sec(x) dx = \frac{1}{2} \log(\sin(x) + 1) - \frac{1}{2} \log(-\sin(x) + 1)$$

input `integrate(1/cos(x),x, algorithm="fricas")`

output `1/2*log(sin(x) + 1) - 1/2*log(-sin(x) + 1)`

3.36.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(3) = 6$.

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 5.00

$$\int \sec(x) dx = -\frac{\log(\sin(x) - 1)}{2} + \frac{\log(\sin(x) + 1)}{2}$$

input `integrate(1/cos(x),x)`

output `-log(sin(x) - 1)/2 + log(sin(x) + 1)/2`

3.36.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(3) = 6$.

Time = 0.21 (sec) , antiderivative size = 15, normalized size of antiderivative = 5.00

$$\int \sec(x) dx = \frac{1}{2} \log(\sin(x) + 1) - \frac{1}{2} \log(\sin(x) - 1)$$

input `integrate(1/cos(x),x, algorithm="maxima")`

output `1/2*log(sin(x) + 1) - 1/2*log(sin(x) - 1)`

3.36.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(3) = 6$.

Time = 0.27 (sec) , antiderivative size = 17, normalized size of antiderivative = 5.67

$$\int \sec(x) dx = \frac{1}{2} \log(\sin(x) + 1) - \frac{1}{2} \log(-\sin(x) + 1)$$

input `integrate(1/cos(x),x, algorithm="giac")`

output `1/2*log(sin(x) + 1) - 1/2*log(-sin(x) + 1)`

3.36.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 11, normalized size of antiderivative = 3.67

$$\int \sec(x) dx = \ln\left(\frac{1}{\cos(x)}\right) + \ln(\sin(x) + 1)$$

input `int(1/cos(x),x)`

output `log(1/cos(x)) + log(sin(x) + 1)`

$$3.37 \quad \int \frac{1}{9 \cos^2(x) + 4 \sin^2(x)} dx$$

3.37.1	Optimal result	297
3.37.2	Mathematica [A] (verified)	297
3.37.3	Rubi [A] (verified)	298
3.37.4	Maple [A] (verified)	299
3.37.5	Fricas [A] (verification not implemented)	299
3.37.6	Sympy [F]	300
3.37.7	Maxima [A] (verification not implemented)	300
3.37.8	Giac [A] (verification not implemented)	300
3.37.9	Mupad [B] (verification not implemented)	301

3.37.1 Optimal result

Integrand size = 15, antiderivative size = 24

$$\int \frac{1}{9 \cos^2(x) + 4 \sin^2(x)} dx = \frac{x}{6} - \frac{1}{6} \arctan\left(\frac{\cos(x) \sin(x)}{2 + \cos^2(x)}\right)$$

output `1/6*x-1/6*arctan(cos(x)*sin(x)/(2+cos(x)^2))`

3.37.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46

$$\int \frac{1}{9 \cos^2(x) + 4 \sin^2(x)} dx = \frac{1}{6} \arctan\left(\frac{2 \tan(x)}{3}\right)$$

input `Integrate[(9*Cos[x]^2 + 4*Sin[x]^2)^(-1),x]`

output `ArcTan[(2*Tan[x])/3]/6`

3.37.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.46, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 4889, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{4 \sin^2(x) + 9 \cos^2(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{4 \sin(x)^2 + 9 \cos(x)^2} dx \\ & \quad \downarrow \text{4889} \\ & \int \frac{1}{4 \tan^2(x) + 9} d \tan(x) \\ & \quad \downarrow \text{216} \\ & \frac{1}{6} \arctan\left(\frac{2 \tan(x)}{3}\right) \end{aligned}$$

input `Int[(9*Cos[x]^2 + 4*Sin[x]^2)^(-1),x]`

output `ArcTan[(2*Tan[x])/3]/6`

3.37.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4889 Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, With[{d = FreeFactors
[Tan[v], x]}, Simp[d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x
^2), Tan[v]/d, u, x], x], x, Tan[v]/d], x]] /; !FalseQ[v] && FunctionOfQ[N
onfreeFactors[Tan[v], x], u, x]] /; InverseFunctionFreeQ[u, x] && !MatchQ[
u, (v_.)*((c_.)*tan[w_]^(n_.)*tan[z_]^(n_.))^(p_.) /; FreeQ[{c, p}, x] && I
ntegerQ[n] && LinearQ[w, x] && EqQ[z, 2*w]]
```

3.37.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.33

method	result	size
default	$\frac{\arctan\left(\frac{2 \tan(x)}{3}\right)}{6}$	8
risch	$-\frac{i \ln\left(e^{2ix} + \frac{1}{5}\right)}{12} + \frac{i \ln(e^{2ix} + 5)}{12}$	24
parallelrisch	$-\frac{i \left(\ln\left(\frac{-2i \sin(x) - 3 \cos(x)}{1 + \cos(x)}\right) - \ln\left(\frac{2i \sin(x) - 3 \cos(x)}{1 + \cos(x)}\right) \right)}{12}$	43

input `int(1/(9*cos(x)^2+4*sin(x)^2),x,method=_RETURNVERBOSE)`

output `1/6*arctan(2/3*tan(x))`

3.37.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

$$\int \frac{1}{9 \cos^2(x) + 4 \sin^2(x)} dx = -\frac{1}{12} \arctan\left(\frac{13 \cos(x)^2 - 4}{12 \cos(x) \sin(x)}\right)$$

input `integrate(1/(9*cos(x)^2+4*sin(x)^2),x, algorithm="fricas")`

output `-1/12*arctan(1/12*(13*cos(x)^2 - 4)/(cos(x)*sin(x)))`

3.37.6 Sympy [F]

$$\int \frac{1}{9 \cos^2(x) + 4 \sin^2(x)} dx = \int \frac{1}{4 \sin^2(x) + 9 \cos^2(x)} dx$$

input `integrate(1/(9*cos(x)**2+4*sin(x)**2),x)`

output `Integral(1/(4*sin(x)**2 + 9*cos(x)**2), x)`

3.37.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.29

$$\int \frac{1}{9 \cos^2(x) + 4 \sin^2(x)} dx = \frac{1}{6} \arctan\left(\frac{2}{3} \tan(x)\right)$$

input `integrate(1/(9*cos(x)^2+4*sin(x)^2),x, algorithm="maxima")`

output `1/6*arctan(2/3*tan(x))`

3.37.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{1}{9 \cos^2(x) + 4 \sin^2(x)} dx = \frac{1}{6} x - \frac{1}{6} \arctan\left(\frac{\sin(2x)}{\cos(2x) + 5}\right)$$

input `integrate(1/(9*cos(x)^2+4*sin(x)^2),x, algorithm="giac")`

output `1/6*x - 1/6*arctan(sin(2*x)/(cos(2*x) + 5))`

3.37.9 Mupad [B] (verification not implemented)

Time = 15.66 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \frac{1}{9 \cos^2(x) + 4 \sin^2(x)} dx = \frac{x}{6} - \frac{\operatorname{atan}(\tan(x))}{6} + \frac{\operatorname{atan}\left(\frac{2 \tan(x)}{3}\right)}{6}$$

input `int(1/(9*cos(x)^2 + 4*sin(x)^2),x)`

output `x/6 - atan(tan(x))/6 + atan((2*tan(x))/3)/6`

3.38 $\int \frac{1}{x^2(1+x^4)^{3/4}} dx$

3.38.1	Optimal result	302
3.38.2	Mathematica [A] (verified)	302
3.38.3	Rubi [A] (verified)	303
3.38.4	Maple [A] (verified)	303
3.38.5	Fricas [A] (verification not implemented)	304
3.38.6	Sympy [B] (verification not implemented)	304
3.38.7	Maxima [A] (verification not implemented)	305
3.38.8	Giac [F]	305
3.38.9	Mupad [B] (verification not implemented)	305

3.38.1 Optimal result

Integrand size = 13, antiderivative size = 14

$$\int \frac{1}{x^2(1+x^4)^{3/4}} dx = -\frac{\sqrt[4]{1+x^4}}{x}$$

output

```
-(x^4+1)^(1/4)/x
```

3.38.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2(1+x^4)^{3/4}} dx = -\frac{\sqrt[4]{1+x^4}}{x}$$

input

```
Integrate[1/(x^2*(1 + x^4)^(3/4)),x]
```

output

```
-((1 + x^4)^(1/4)/x)
```

3.38.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {796}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 (x^4 + 1)^{3/4}} dx$$

↓ 796

$$-\frac{\sqrt[4]{x^4 + 1}}{x}$$

input `Int[1/(x^2*(1 + x^4)^(3/4)),x]`

output `-((1 + x^4)^(1/4)/x)`

3.38.3.1 Defintions of rubi rules used

rule 796 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, n, p}, x] && EqQ[(m + 1)/n + p + 1, 0] && NeQ[m, -1]`

3.38.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
gospers	$-\frac{(x^4+1)^{\frac{1}{4}}}{x}$	13
trager	$-\frac{(x^4+1)^{\frac{1}{4}}}{x}$	13
meijerg	$-\frac{(x^4+1)^{\frac{1}{4}}}{x}$	13
risch	$-\frac{(x^4+1)^{\frac{1}{4}}}{x}$	13
pseudoelliptic	$-\frac{(x^4+1)^{\frac{1}{4}}}{x}$	13

input `int(1/x^2/(x^4+1)^(3/4),x,method=_RETURNVERBOSE)`

output $-(x^4+1)^{1/4}/x$

3.38.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{x^2(1+x^4)^{3/4}} dx = -\frac{(x^4+1)^{1/4}}{x}$$

input `integrate(1/x^2/(x^4+1)^(3/4),x, algorithm="fricas")`

output $-(x^4 + 1)^{1/4}/x$

3.38.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 22 vs. $2(10) = 20$.

Time = 0.32 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.57

$$\int \frac{1}{x^2(1+x^4)^{3/4}} dx = \frac{\sqrt[4]{1+\frac{1}{x^4}}\Gamma(-\frac{1}{4})}{4\Gamma(\frac{3}{4})}$$

input `integrate(1/x**2/(x**4+1)**(3/4),x)`

output $(1 + x^{**(-4)})^{**}(1/4)*\text{gamma}(-1/4)/(4*\text{gamma}(3/4))$

3.38.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{x^2 (1+x^4)^{3/4}} dx = -\frac{(x^4+1)^{1/4}}{x}$$

input `integrate(1/x^2/(x^4+1)^(3/4),x, algorithm="maxima")`output `-(x^4 + 1)^(1/4)/x`**3.38.8 Giac [F]**

$$\int \frac{1}{x^2 (1+x^4)^{3/4}} dx = \int \frac{1}{(x^4+1)^{3/4} x^2} dx$$

input `integrate(1/x^2/(x^4+1)^(3/4),x, algorithm="giac")`output `integrate(1/((x^4 + 1)^(3/4)*x^2), x)`**3.38.9 Mupad [B] (verification not implemented)**

Time = 15.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{x^2 (1+x^4)^{3/4}} dx = -\frac{(x^4+1)^{1/4}}{x}$$

input `int(1/(x^2*(x^4 + 1)^(3/4)),x)`output `-(x^4 + 1)^(1/4)/x`

3.39 $\int \cos(x) \cos(3x) \cos(5x) dx$

3.39.1	Optimal result	306
3.39.2	Mathematica [A] (verified)	306
3.39.3	Rubi [A] (verified)	307
3.39.4	Maple [A] (verified)	308
3.39.5	Fricas [A] (verification not implemented)	308
3.39.6	Sympy [B] (verification not implemented)	308
3.39.7	Maxima [A] (verification not implemented)	309
3.39.8	Giac [A] (verification not implemented)	309
3.39.9	Mupad [B] (verification not implemented)	309

3.39.1 Optimal result

Integrand size = 11, antiderivative size = 31

$$\int \cos(x) \cos(3x) \cos(5x) dx = \frac{\sin(x)}{4} + \frac{1}{12} \sin(3x) + \frac{1}{28} \sin(7x) + \frac{1}{36} \sin(9x)$$

output `1/4*sin(x)+1/12*sin(3*x)+1/28*sin(7*x)+1/36*sin(9*x)`

3.39.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \cos(x) \cos(3x) \cos(5x) dx = \frac{\sin(x)}{4} + \frac{1}{12} \sin(3x) + \frac{1}{28} \sin(7x) + \frac{1}{36} \sin(9x)$$

input `Integrate[Cos[x]*Cos[3*x]*Cos[5*x],x]`

output `Sin[x]/4 + Sin[3*x]/12 + Sin[7*x]/28 + Sin[9*x]/36`

3.39.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 4855, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(x) \cos(3x) \cos(5x) dx$$

$$\downarrow 3042$$

$$\int \cos(x) \cos(3x) \cos(5x) dx$$

$$\downarrow 4855$$

$$\int \left(\frac{\cos(x)}{4} + \frac{1}{4} \cos(3x) + \frac{1}{4} \cos(7x) + \frac{1}{4} \cos(9x) \right) dx$$

$$\downarrow 2009$$

$$\frac{\sin(x)}{4} + \frac{1}{12} \sin(3x) + \frac{1}{28} \sin(7x) + \frac{1}{36} \sin(9x)$$

input `Int[Cos[x]*Cos[3*x]*Cos[5*x],x]`

output `Sin[x]/4 + Sin[3*x]/12 + Sin[7*x]/28 + Sin[9*x]/36`

3.39.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4855 `Int[(F_)[(a_.) + (b_.)*(x_)]^(p_.)*(G_)[(c_.) + (d_.)*(x_)]^(q_.)*(H_)[(e_.) + (f_.)*(x_)]^(r_.), x_Symbol] := Int[ExpandTrigReduce[ActivateTrig[F[a + b*x]^p*G[c + d*x]^q*H[e + f*x]^r], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && (EqQ[F, sin] || EqQ[F, cos]) && (EqQ[G, sin] || EqQ[G, cos]) && (EqQ[H, sin] || EqQ[H, cos]) && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]`

3.39.4 Maple [A] (verified)

Time = 0.70 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.77

method	result	size
default	$\frac{\sin(x)}{4} + \frac{\sin(3x)}{12} + \frac{\sin(7x)}{28} + \frac{\sin(9x)}{36}$	24
risch	$\frac{\sin(x)}{4} + \frac{\sin(3x)}{12} + \frac{\sin(7x)}{28} + \frac{\sin(9x)}{36}$	24
parallelrisch	$\frac{\sin(x)}{4} + \frac{\sin(3x)}{12} + \frac{\sin(7x)}{28} + \frac{\sin(9x)}{36}$	24

input `int(cos(x)*cos(3*x)*cos(5*x),x,method=_RETURNVERBOSE)`

output `1/4*sin(x)+1/12*sin(3*x)+1/28*sin(7*x)+1/36*sin(9*x)`

3.39.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.97

$$\int \cos(x) \cos(3x) \cos(5x) dx$$

$$= \frac{1}{63} (448 \cos(x)^8 - 640 \cos(x)^6 + 240 \cos(x)^4 + 5 \cos(x)^2 + 10) \sin(x)$$

input `integrate(cos(x)*cos(3*x)*cos(5*x),x, algorithm="fricas")`

output `1/63*(448*cos(x)^8 - 640*cos(x)^6 + 240*cos(x)^4 + 5*cos(x)^2 + 10)*sin(x)`

3.39.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 66 vs. $2(24) = 48$.

Time = 0.83 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.13

$$\int \cos(x) \cos(3x) \cos(5x) dx = -\frac{10 \sin(x) \sin(3x) \sin(5x)}{63} - \frac{11 \sin(x) \cos(3x) \cos(5x)}{63}$$

$$- \frac{17 \sin(3x) \cos(x) \cos(5x)}{63} + \frac{25 \sin(5x) \cos(x) \cos(3x)}{63}$$

input `integrate(cos(x)*cos(3*x)*cos(5*x),x)`

output `-10*sin(x)*sin(3*x)*sin(5*x)/63 - 11*sin(x)*cos(3*x)*cos(5*x)/63 - 17*sin(3*x)*cos(x)*cos(5*x)/63 + 25*sin(5*x)*cos(x)*cos(3*x)/63`

3.39.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int \cos(x) \cos(3x) \cos(5x) dx = \frac{1}{36} \sin(9x) + \frac{1}{28} \sin(7x) + \frac{1}{12} \sin(3x) + \frac{1}{4} \sin(x)$$

input `integrate(cos(x)*cos(3*x)*cos(5*x),x, algorithm="maxima")`

output `1/36*sin(9*x) + 1/28*sin(7*x) + 1/12*sin(3*x) + 1/4*sin(x)`

3.39.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int \cos(x) \cos(3x) \cos(5x) dx = \frac{1}{36} \sin(9x) + \frac{1}{28} \sin(7x) + \frac{1}{12} \sin(3x) + \frac{1}{4} \sin(x)$$

input `integrate(cos(x)*cos(3*x)*cos(5*x),x, algorithm="giac")`

output `1/36*sin(9*x) + 1/28*sin(7*x) + 1/12*sin(3*x) + 1/4*sin(x)`

3.39.9 Mupad [B] (verification not implemented)

Time = 15.14 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

$$\int \cos(x) \cos(3x) \cos(5x) dx = \frac{64 \sin(x)^9}{9} - \frac{128 \sin(x)^7}{7} + 16 \sin(x)^5 - \frac{17 \sin(x)^3}{3} + \sin(x)$$

input `int(cos(3*x)*cos(5*x)*cos(x),x)`

output `sin(x) - (17*sin(x)^3)/3 + 16*sin(x)^5 - (128*sin(x)^7)/7 + (64*sin(x)^9)/9`

$$\mathbf{3.40} \quad \int \left(\frac{1}{\log(x)} + \log(\log(x)) \right) dx$$

3.40.1	Optimal result	310
3.40.2	Mathematica [A] (verified)	310
3.40.3	Rubi [A] (verified)	311
3.40.4	Maple [A] (verified)	311
3.40.5	Fricas [A] (verification not implemented)	312
3.40.6	Sympy [A] (verification not implemented)	312
3.40.7	Maxima [A] (verification not implemented)	312
3.40.8	Giac [A] (verification not implemented)	313
3.40.9	Mupad [B] (verification not implemented)	313

3.40.1 Optimal result

Integrand size = 8, antiderivative size = 5

$$\int \left(\frac{1}{\log(x)} + \log(\log(x)) \right) dx = x \log(\log(x))$$

output `x*ln(ln(x))`

3.40.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \left(\frac{1}{\log(x)} + \log(\log(x)) \right) dx = x \log(\log(x))$$

input `Integrate[Log[x]^(-1) + Log[Log[x]], x]`

output `x*Log[Log[x]]`

$$3.40. \quad \int \left(\frac{1}{\log(x)} + \log(\log(x)) \right) dx$$

3.40.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\log(\log(x)) + \frac{1}{\log(x)} \right) dx$$

↓ 2009

$$x \log(\log(x))$$

input `Int [Log[x]^(-1) + Log[Log[x]], x]`

output `x*Log[Log[x]]`

3.40.3.1 Defintions of rubi rules used

rule 2009 `Int [u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.40.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
default	$x \ln(\ln(x))$	6
norman	$x \ln(\ln(x))$	6
risch	$x \ln(\ln(x))$	6
parallelrisch	$x \ln(\ln(x))$	6
parts	$x \ln(\ln(x))$	6

input `int(1/ln(x)+ln(ln(x)),x,method=_RETURNVERBOSE)`

output `x*ln(ln(x))`

3.40. $\int \left(\frac{1}{\log(x)} + \log(\log(x)) \right) dx$

3.40.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \left(\frac{1}{\log(x)} + \log(\log(x)) \right) dx = x \log(\log(x))$$

input `integrate(1/log(x)+log(log(x)),x, algorithm="fricas")`output `x*log(log(x))`**3.40.6 Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \left(\frac{1}{\log(x)} + \log(\log(x)) \right) dx = x \log(\log(x))$$

input `integrate(1/ln(x)+ln(ln(x)),x)`output `x*log(log(x))`**3.40.7 Maxima [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \left(\frac{1}{\log(x)} + \log(\log(x)) \right) dx = x \log(\log(x))$$

input `integrate(1/log(x)+log(log(x)),x, algorithm="maxima")`output `x*log(log(x))`

3.40.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \left(\frac{1}{\log(x)} + \log(\log(x)) \right) dx = x \log(\log(x))$$

input `integrate(1/log(x)+log(log(x)),x, algorithm="giac")`

output `x*log(log(x))`

3.40.9 Mupad [B] (verification not implemented)

Time = 15.17 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \left(\frac{1}{\log(x)} + \log(\log(x)) \right) dx = x \ln(\ln(x))$$

input `int(log(log(x)) + 1/log(x),x)`

output `x*log(log(x))`

3.41 $\int \frac{1}{2+e^x} dx$

3.41.1	Optimal result	314
3.41.2	Mathematica [A] (verified)	314
3.41.3	Rubi [B] (verified)	315
3.41.4	Maple [A] (verified)	316
3.41.5	Fricas [A] (verification not implemented)	316
3.41.6	Sympy [A] (verification not implemented)	317
3.41.7	Maxima [A] (verification not implemented)	317
3.41.8	Giac [A] (verification not implemented)	317
3.41.9	Mupad [B] (verification not implemented)	318

3.41.1 Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \frac{1}{2 + e^x} dx = -\operatorname{arctanh}(1 + e^x)$$

output `-arctanh(exp(x)+1)`

3.41.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{1}{2 + e^x} dx = -\operatorname{arctanh}(1 + e^x)$$

input `Integrate[(2 + E^x)^(-1),x]`

output `-ArcTanh[1 + E^x]`

3.41.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 19 vs. $2(8) = 16$.

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.38, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {2720, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{e^x + 2} dx \\ & \quad \downarrow \text{2720} \\ & \int \frac{e^{-x}}{e^x + 2} de^x \\ & \quad \downarrow \text{47} \\ & \frac{\int e^{-x} de^x}{2} - \frac{1}{2} \int \frac{1}{2 + e^x} de^x \\ & \quad \downarrow \text{14} \\ & \frac{\log(e^x)}{2} - \frac{1}{2} \int \frac{1}{2 + e^x} de^x \\ & \quad \downarrow \text{16} \\ & \frac{\log(e^x)}{2} - \frac{1}{2} \log(e^x + 2) \end{aligned}$$

input `Int[(2 + E^x)^(-1), x]`

output `Log[E^x]/2 - Log[2 + E^x]/2`

3.41.3.1 Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

```
rule 47 Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c
- a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x
], x] /; FreeQ[{a, b, c, d}, x]
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.41.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.50

method	result	size
norman	$\frac{x}{2} - \frac{\ln(e^x+2)}{2}$	12
risch	$\frac{x}{2} - \frac{\ln(e^x+2)}{2}$	12
parallelrisch	$\frac{x}{2} - \frac{\ln(e^x+2)}{2}$	12
derivativedivides	$\frac{\ln(e^x)}{2} - \frac{\ln(e^x+2)}{2}$	14
default	$\frac{\ln(e^x)}{2} - \frac{\ln(e^x+2)}{2}$	14

```
input int(1/(exp(x)+2),x,method=_RETURNVERBOSE)
```

```
output 1/2*x-1/2*ln(exp(x)+2)
```

3.41.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.38

$$\int \frac{1}{2+e^x} dx = \frac{1}{2}x - \frac{1}{2} \log(e^x + 2)$$

```
input integrate(1/(exp(x)+2),x, algorithm="fracas")
```

```
output 1/2*x - 1/2*log(e^x + 2)
```

3.41.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.25

$$\int \frac{1}{2+e^x} dx = \frac{x}{2} - \frac{\log(e^x + 2)}{2}$$

input `integrate(1/(exp(x)+2),x)`output `x/2 - log(exp(x) + 2)/2`**3.41.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.38

$$\int \frac{1}{2+e^x} dx = \frac{1}{2}x - \frac{1}{2}\log(e^x + 2)$$

input `integrate(1/(exp(x)+2),x, algorithm="maxima")`output `1/2*x - 1/2*log(e^x + 2)`**3.41.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.38

$$\int \frac{1}{2+e^x} dx = \frac{1}{2}x - \frac{1}{2}\log(e^x + 2)$$

input `integrate(1/(exp(x)+2),x, algorithm="giac")`output `1/2*x - 1/2*log(e^x + 2)`

3.41.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.38

$$\int \frac{1}{2 + e^x} dx = \frac{x}{2} - \frac{\ln(e^x + 2)}{2}$$

input `int(1/(exp(x) + 2),x)`

output `x/2 - log(exp(x) + 2)/2`

3.42 $\int \sqrt{\frac{x}{1-x^3}} dx$

3.42.1	Optimal result	319
3.42.2	Mathematica [A] (verified)	319
3.42.3	Rubi [A] (verified)	320
3.42.4	Maple [A] (verified)	321
3.42.5	Fricas [A] (verification not implemented)	322
3.42.6	Sympy [F]	322
3.42.7	Maxima [F]	322
3.42.8	Giac [A] (verification not implemented)	323
3.42.9	Mupad [F(-1)]	323

3.42.1 Optimal result

Integrand size = 15, antiderivative size = 41

$$\int \sqrt{\frac{x}{1-x^3}} dx = \frac{2\sqrt{\frac{x}{1-x^3}}\sqrt{1-x^3} \arcsin(x^{3/2})}{3\sqrt{x}}$$

output `2/3*(x/(-x^3+1))^(1/2)*(-x^3+1)^(1/2)*arcsin(x^(3/2))/x^(1/2)`

3.42.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.17

$$\int \sqrt{\frac{x}{1-x^3}} dx = \frac{2\sqrt{-\frac{x}{-1+x^3}}\sqrt{-1+x^3} \log(x^{3/2} + \sqrt{-1+x^3})}{3\sqrt{x}}$$

input `Integrate[Sqrt[x/(1 - x^3)],x]`

output `(2*Sqrt[-(x/(-1 + x^3))]*Sqrt[-1 + x^3]*Log[x^(3/2) + Sqrt[-1 + x^3]])/(3*Sqrt[x])`

3.42.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {7270, 851, 807, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \sqrt{\frac{x}{1-x^3}} dx \\
 \downarrow 7270 \\
 \frac{\sqrt{\frac{x}{1-x^3}} \sqrt{1-x^3} \int \frac{\sqrt{x}}{\sqrt{1-x^3}} dx}{\sqrt{x}} \\
 \downarrow 851 \\
 \frac{2\sqrt{\frac{x}{1-x^3}} \sqrt{1-x^3} \int \frac{x}{\sqrt{1-x^3}} d\sqrt{x}}{\sqrt{x}} \\
 \downarrow 807 \\
 \frac{2\sqrt{\frac{x}{1-x^3}} \sqrt{1-x^3} \int \frac{1}{\sqrt{1-x}} dx^{3/2}}{3\sqrt{x}} \\
 \downarrow 223 \\
 \frac{2\sqrt{\frac{x}{1-x^3}} \sqrt{1-x^3} \arcsin(x^{3/2})}{3\sqrt{x}}
 \end{array}$$

input `Int[Sqrt[x/(1 - x^3)],x]`

output `(2*Sqrt[x/(1 - x^3)]*Sqrt[1 - x^3]*ArcSin[x^(3/2)])/(3*Sqrt[x])`

3.42.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 851 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 7270 `Int[(u_.)*((a_.)*(v_)^(m_.)*(w_)^(n_.))^(p_), x_Symbol] := Simp[a^IntPart[p]*(a*v^m*w^n)^FracPart[p]/(v^(m*FracPart[p])*w^(n*FracPart[p])) Int[u*v^(m*p)*w^(n*p), x], x] /; FreeQ[{a, m, n, p}, x] && !IntegerQ[p] && !FreeQ[v, x] && !FreeQ[w, x]`

3.42.4 Maple [A] (verified)

Time = 0.78 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.05

method	result
default	$\frac{2\sqrt{-\frac{x}{x^3-1}}(x^3-1)\operatorname{arctanh}\left(\frac{\sqrt{x^4-x}}{x^2}\right)}{3\sqrt{(x^3-1)x}}$
trager	$\frac{\operatorname{RootOf}(_Z^2+1)\ln\left(-2\sqrt{-\frac{x}{x^3-1}}x^4-2\operatorname{RootOf}(_Z^2+1)x^3+2\sqrt{-\frac{x}{x^3-1}}x+\operatorname{RootOf}(_Z^2+1)\right)}{3}$
elliptic	$\frac{2\sqrt{-\frac{x}{x^3-1}}\sqrt{(x^3-1)x}\left(\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{\left(-\frac{3}{2}+\frac{i\sqrt{3}}{2}\right)x}{\left(-\frac{1}{2}+\frac{i\sqrt{3}}{2}\right)(-1+x)}}(-1+x)^2\sqrt{\frac{x+\frac{1}{2}+\frac{i\sqrt{3}}{2}}{\left(-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)(-1+x)}}\sqrt{\frac{x+\frac{1}{2}-\frac{i\sqrt{3}}{2}}{\left(-\frac{1}{2}+\frac{i\sqrt{3}}{2}\right)(-1+x)}}\left(\operatorname{EllipticF}\left(\sqrt{\frac{\left(-\frac{3}{2}+\frac{i\sqrt{3}}{2}\right)x}{\left(-\frac{1}{2}+\frac{i\sqrt{3}}{2}\right)(-1+x)}}}{x\left(-\frac{3}{2}+\frac{i\sqrt{3}}{2}\right)}\sqrt{x(-1+x)}\left(x+\frac{1}{2}+\frac{i\sqrt{3}}{2}\right)\right)}{x\left(-\frac{3}{2}+\frac{i\sqrt{3}}{2}\right)\sqrt{x(-1+x)}\left(x+\frac{1}{2}+\frac{i\sqrt{3}}{2}\right)}$

input `int((x/(-x^3+1))^(1/2),x,method=_RETURNVERBOSE)`

output `2/3*(-x/(x^3-1))^(1/2)*(x^3-1)/((x^3-1)*x)^(1/2)*arctanh((x^4-x)^(1/2)/x^2)`

3.42.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.80

$$\int \sqrt{\frac{x}{1-x^3}} dx = \frac{1}{3} \arctan \left(\frac{2(x^4-x)\sqrt{-\frac{x}{x^3-1}}}{2x^3-1} \right)$$

input `integrate((x/(-x^3+1))^(1/2),x, algorithm="fricas")`

output `1/3*arctan(2*(x^4 - x)*sqrt(-x/(x^3 - 1))/(2*x^3 - 1))`

3.42.6 Sympy [F]

$$\int \sqrt{\frac{x}{1-x^3}} dx = \int \sqrt{\frac{x}{1-x^3}} dx$$

input `integrate((x/(-x**3+1))**(1/2),x)`

output `Integral(sqrt(x/(1 - x**3)), x)`

3.42.7 Maxima [F]

$$\int \sqrt{\frac{x}{1-x^3}} dx = \int \sqrt{-\frac{x}{x^3-1}} dx$$

input `integrate((x/(-x^3+1))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(-x/(x^3 - 1)), x)`

3.42.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.39

$$\int \sqrt{\frac{x}{1-x^3}} dx = \frac{2}{3} \arctan \left(\sqrt{\frac{1}{x^3} - 1} \right) \operatorname{sgn}(x^3 - 1)$$

input `integrate((x/(-x^3+1))^(1/2),x, algorithm="giac")`

output `2/3*arctan(sqrt(1/x^3 - 1))*sgn(x^3 - 1)`

3.42.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\frac{x}{1-x^3}} dx = \int \sqrt{-\frac{x}{x^3-1}} dx$$

input `int((-x/(x^3 - 1))^(1/2),x)`

output `int((-x/(x^3 - 1))^(1/2), x)`

3.43 $\int \frac{4x}{1-x^4} dx$

3.43.1	Optimal result	324
3.43.2	Mathematica [B] (verified)	324
3.43.3	Rubi [A] (verified)	325
3.43.4	Maple [A] (verified)	326
3.43.5	Fricas [B] (verification not implemented)	326
3.43.6	Sympy [B] (verification not implemented)	326
3.43.7	Maxima [B] (verification not implemented)	327
3.43.8	Giac [B] (verification not implemented)	327
3.43.9	Mupad [B] (verification not implemented)	327

3.43.1 Optimal result

Integrand size = 12, antiderivative size = 6

$$\int \frac{4x}{1-x^4} dx = 2\operatorname{arctanh}(x^2)$$

output `2*arctanh(x^2)`

3.43.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 25 vs. $2(6) = 12$.

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 4.17

$$\int \frac{4x}{1-x^4} dx = -4 \left(\frac{1}{4} \log(1-x^2) - \frac{1}{4} \log(1+x^2) \right)$$

input `Integrate[(4*x)/(1 - x^4),x]`

output `-4*(Log[1 - x^2]/4 - Log[1 + x^2]/4)`

3.43.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {27, 807, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{4x}{1-x^4} dx \\ & \quad \downarrow \text{27} \\ & 4 \int \frac{x}{1-x^4} dx \\ & \quad \downarrow \text{807} \\ & 2 \int \frac{1}{1-x^4} dx^2 \\ & \quad \downarrow \text{219} \\ & 2\operatorname{arctanh}(x^2) \end{aligned}$$

input `Int[(4*x)/(1 - x^4),x]`

output `2*ArcTanh[x^2]`

3.43.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

3.43.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

method	result	size
meijerg	$2 \operatorname{arctanh}(x^2)$	7
risch	$\ln(x^2 + 1) - \ln(x^2 - 1)$	16
default	$-\ln(-1 + x) - \ln(1 + x) + \ln(x^2 + 1)$	20
norman	$-\ln(-1 + x) - \ln(1 + x) + \ln(x^2 + 1)$	20
parallelrisch	$-\ln(-1 + x) - \ln(1 + x) + \ln(x^2 + 1)$	20

input `int(4*x/(-x^4+1),x,method=_RETURNVERBOSE)`

output `2*arctanh(x^2)`

3.43.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(6) = 12$.

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 2.50

$$\int \frac{4x}{1-x^4} dx = \log(x^2 + 1) - \log(x^2 - 1)$$

input `integrate(4*x/(-x^4+1),x, algorithm="fricas")`

output `log(x^2 + 1) - log(x^2 - 1)`

3.43.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12 vs. $2(5) = 10$.

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 2.00

$$\int \frac{4x}{1-x^4} dx = -\log(x^2 - 1) + \log(x^2 + 1)$$

input `integrate(4*x/(-x**4+1),x)`

output `-log(x**2 - 1) + log(x**2 + 1)`

3.43.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(6) = 12$.

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 2.50

$$\int \frac{4x}{1-x^4} dx = \log(x^2 + 1) - \log(x^2 - 1)$$

input `integrate(4*x/(-x^4+1),x, algorithm="maxima")`

output `log(x^2 + 1) - log(x^2 - 1)`

3.43.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 16 vs. $2(6) = 12$.

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 2.67

$$\int \frac{4x}{1-x^4} dx = \log(x^2 + 1) - \log(|x^2 - 1|)$$

input `integrate(4*x/(-x^4+1),x, algorithm="giac")`

output `log(x^2 + 1) - log(abs(x^2 - 1))`

3.43.9 Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{4x}{1-x^4} dx = 2 \operatorname{atanh}(x^2)$$

input `int(-(4*x)/(x^4 - 1),x)`

output `2*atanh(x^2)`

3.44 $\int x^x(1 + \log(x)) dx$

3.44.1	Optimal result	328
3.44.2	Mathematica [A] (verified)	328
3.44.3	Rubi [A] (verified)	329
3.44.4	Maple [A] (verified)	329
3.44.5	Fricas [A] (verification not implemented)	330
3.44.6	Sympy [A] (verification not implemented)	330
3.44.7	Maxima [A] (verification not implemented)	331
3.44.8	Giac [A] (verification not implemented)	331
3.44.9	Mupad [B] (verification not implemented)	331

3.44.1 Optimal result

Integrand size = 8, antiderivative size = 3

$$\int x^x(1 + \log(x)) dx = x^x$$

output

x^x

3.44.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int x^x(1 + \log(x)) dx = x^x$$

input `Integrate[x^x*(1 + Log[x]),x]`

output

x^x

3.44.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^x(\log(x) + 1) dx$$

$$\downarrow 7293$$

$$\int (x^x + x^x \log(x)) dx$$

$$\downarrow 2009$$

$$x^x$$

input `Int[x^x*(1 + Log[x]),x]`

output `x^x`

3.44.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] :> With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

3.44.4 Maple [A] (verified)

Time = 0.00 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
derivativedivides	x^x	4
default	x^x	4
risch	x^x	4
parallelrisch	x^x	4
norman	$e^{x \ln(x)}$	6

input `int(x^x*(ln(x)+1),x,method=_RETURNVERBOSE)`

output `x^x`

3.44.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int x^x(1 + \log(x)) dx = x^x$$

input `integrate(x^x*(log(x)+1),x, algorithm="fricas")`

output `x^x`

3.44.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int x^x(1 + \log(x)) dx = x^x$$

input `integrate(x**x*(ln(x)+1),x)`

output `x**x`

3.44.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int x^x(1 + \log(x)) dx = x^x$$

input `integrate(x^x*(log(x)+1),x, algorithm="maxima")`output `x^x`**3.44.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int x^x(1 + \log(x)) dx = x^x$$

input `integrate(x^x*(log(x)+1),x, algorithm="giac")`output `x^x`**3.44.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int x^x(1 + \log(x)) dx = x^x$$

input `int(x^x*(log(x) + 1),x)`output `x^x`

3.45 $\int \sqrt{6x - x^2} dx$

3.45.1	Optimal result	332
3.45.2	Mathematica [A] (verified)	332
3.45.3	Rubi [A] (verified)	333
3.45.4	Maple [A] (verified)	334
3.45.5	Fricas [A] (verification not implemented)	334
3.45.6	Sympy [A] (verification not implemented)	335
3.45.7	Maxima [A] (verification not implemented)	335
3.45.8	Giac [A] (verification not implemented)	335
3.45.9	Mupad [B] (verification not implemented)	336

3.45.1 Optimal result

Integrand size = 13, antiderivative size = 29

$$\int \sqrt{6x - x^2} dx = \frac{1}{2} \left((-3 + x) \sqrt{-((-6 + x)x)} - 9 \arcsin \left(1 - \frac{x}{3} \right) \right)$$

output `1/2*(-3+x)*(-x*(-6+x))^(1/2)+9/2*arcsin(-1+1/3*x)`

3.45.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.62

$$\int \sqrt{6x - x^2} dx = \frac{1}{2} \sqrt{-((-6 + x)x)} \left(-3 + x + \frac{18 \log(\sqrt{-6 + x} - \sqrt{x})}{\sqrt{-6 + x}\sqrt{x}} \right)$$

input `Integrate[Sqrt[6*x - x^2],x]`

output `(Sqrt[-((-6 + x)*x)]*(-3 + x + (18*Log[Sqrt[-6 + x] - Sqrt[x]])/(Sqrt[-6 + x]*Sqrt[x])))/2`

3.45.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.28, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {1087, 1090, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{6x - x^2} dx$$

$$\downarrow 1087$$

$$\frac{9}{2} \int \frac{1}{\sqrt{6x - x^2}} dx - \frac{1}{2}(3 - x)\sqrt{6x - x^2}$$

$$\downarrow 1090$$

$$-\frac{3}{4} \int \frac{1}{\sqrt{1 - \frac{1}{36}(6 - 2x)^2}} d(6 - 2x) - \frac{1}{2}\sqrt{6x - x^2}(3 - x)$$

$$\downarrow 223$$

$$-\frac{9}{2} \arcsin\left(\frac{1}{6}(6 - 2x)\right) - \frac{1}{2}\sqrt{6x - x^2}(3 - x)$$

input `Int[Sqrt[6*x - x^2], x]`

output `-1/2*((3 - x)*Sqrt[6*x - x^2]) - (9*ArcSin[(6 - 2*x)/6])/2`

3.45.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 1087 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] - Simp[p*(b^2 - 4*a*c)/(2*c*(2*p + 1)) Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[3*p])`

rule 1090 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

3.45.4 Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

method	result	size
risch	$-\frac{(-3+x)x(-6+x)}{2\sqrt{-x(-6+x)}} + \frac{9 \arcsin(\frac{x}{3}-1)}{2}$	2
default	$-\frac{(-2x+6)\sqrt{-x^2+6x}}{4} + \frac{9 \arcsin(\frac{x}{3}-1)}{2}$	2
pseudoelliptic	$-9 \arctan\left(\frac{\sqrt{-x(-6+x)}}{x}\right) + \frac{(-3+x)\sqrt{-x(-6+x)}}{2}$	3
meijerg	$-\frac{18i\left(-\frac{i\sqrt{\pi}\sqrt{x}\sqrt{6(-x+3)}\sqrt{-\frac{x}{6}+1}}{36} + \frac{i\sqrt{\pi}\arcsin\left(\frac{\sqrt{6}\sqrt{x}}{6}\right)}{2}\right)}{\sqrt{\pi}}$	4
trager	$\left(-\frac{3}{2} + \frac{x}{2}\right)\sqrt{-x^2+6x} + \frac{9\text{RootOf}\left(_Z^2+1\right)\ln\left(-\text{RootOf}\left(_Z^2+1\right)x+\sqrt{-x^2+6x}+3\text{RootOf}\left(_Z^2+1\right)\right)}{2}$	5

input `int((-x^2+6*x)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2*(-3+x)*x*(-6+x)/(-x*(-6+x))^(1/2)+9/2*arcsin(1/3*x-1)`

3.45.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.21

$$\int \sqrt{6x - x^2} dx = \frac{1}{2} \sqrt{-x^2 + 6x}(x - 3) - 9 \arctan\left(\frac{\sqrt{-x^2 + 6x}}{x}\right)$$

input `integrate((-x^2+6*x)^(1/2),x, algorithm="fricas")`

output `1/2*sqrt(-x^2 + 6*x)*(x - 3) - 9*arctan(sqrt(-x^2 + 6*x)/x)`

3.45.6 Sympy [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.90

$$\int \sqrt{6x - x^2} dx = \left(\frac{x}{2} - \frac{3}{2}\right) \sqrt{-x^2 + 6x} + \frac{9 \operatorname{asin}\left(\frac{x}{3} - 1\right)}{2}$$

input `integrate((-x**2+6*x)**(1/2),x)`output `(x/2 - 3/2)*sqrt(-x**2 + 6*x) + 9*asin(x/3 - 1)/2`**3.45.7 Maxima [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.24

$$\int \sqrt{6x - x^2} dx = \frac{1}{2} \sqrt{-x^2 + 6x} x - \frac{3}{2} \sqrt{-x^2 + 6x} - \frac{9}{2} \arcsin\left(-\frac{1}{3}x + 1\right)$$

input `integrate((-x^2+6*x)^(1/2),x, algorithm="maxima")`output `1/2*sqrt(-x^2 + 6*x)*x - 3/2*sqrt(-x^2 + 6*x) - 9/2*arcsin(-1/3*x + 1)`**3.45.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.86

$$\int \sqrt{6x - x^2} dx = \frac{1}{2} \sqrt{-x^2 + 6x}(x - 3) + \frac{9}{2} \arcsin\left(\frac{1}{3}x - 1\right)$$

input `integrate((-x^2+6*x)^(1/2),x, algorithm="giac")`output `1/2*sqrt(-x^2 + 6*x)*(x - 3) + 9/2*arcsin(1/3*x - 1)`

3.45.9 Mupad [B] (verification not implemented)

Time = 14.40 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.90

$$\int \sqrt{6x - x^2} dx = \frac{9 \operatorname{asin}\left(\frac{x}{3} - 1\right)}{2} + \left(\frac{x}{2} - \frac{3}{2}\right) \sqrt{6x - x^2}$$

input `int((6*x - x^2)^(1/2),x)`

output `(9*asin(x/3 - 1))/2 + (x/2 - 3/2)*(6*x - x^2)^(1/2)`

3.46 $\int \sin^{99}(x) \sin(101x) dx$

3.46.1	Optimal result	337
3.46.2	Mathematica [A] (verified)	337
3.46.3	Rubi [B] (verified)	338
3.46.4	Maple [B] (verified)	341
3.46.5	Fricas [F(-1)]	341
3.46.6	Sympy [F(-1)]	342
3.46.7	Maxima [F(-2)]	342
3.46.8	Giac [B] (verification not implemented)	342
3.46.9	Mupad [B] (verification not implemented)	343

3.46.1 Optimal result

Integrand size = 9, antiderivative size = 12

$$\int \sin^{99}(x) \sin(101x) dx = \frac{1}{100} \sin^{100}(x) \sin(100x)$$

output `1/100*sin(x)^100*sin(100*x)`

3.46.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \sin^{99}(x) \sin(101x) dx = \frac{1}{100} \sin^{100}(x) \sin(100x)$$

input `Integrate[Sin[x]^99*Sin[101*x],x]`

output `(Sin[x]^100*Sin[100*x])/100`

3.46.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 801 vs. $2(12) = 24$.

Time = 0.97 (sec) , antiderivative size = 801, normalized size of antiderivative = 66.75, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 4854, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sin^{99}(x) \sin(101x) dx$$

$$\downarrow 3042$$

$$\int \sin(x)^{99} \sin(101x) dx$$

$$\downarrow 4854$$

$$\int \left(-\frac{\cos(2x)}{633825300114114700748351602688} + \frac{99 \cos(4x)}{633825300114114700748351602688} - \frac{4851 \cos(6x)}{633825300114114700748351602688} \right) dx$$

$$\downarrow 2009$$

$\frac{\sin(2x)}{1267650600228229401496703205376}$	+	$\frac{99 \sin(4x)}{2535301200456458802993406410752}$
$\frac{1617 \sin(6x)}{1267650600228229401496703205376}$	+	$\frac{156849 \sin(8x)}{5070602400912917605986812821504}$
$\frac{470547 \sin(10x)}{792281625142643375935439503360}$	+	$\frac{2980131 \sin(12x)}{316912650057057350374175801344}$
$\frac{20009451 \sin(14x)}{158456325028528675187087900672}$	+	$\frac{1860878943 \sin(16x)}{1267650600228229401496703205376}$
$\frac{4755579521 \sin(18x)}{316912650057057350374175801344}$	+	$\frac{432757736411 \sin(20x)}{3169126500570573503741758013440}$
$\frac{354074511609 \sin(22x)}{316912650057057350374175801344}$	+	$\frac{10504210511067 \sin(24x)}{1267650600228229401496703205376}$
$\frac{8888178124749 \sin(26x)}{158456325028528675187087900672}$	+	$\frac{110467356693309 \sin(28x)}{316912650057057350374175801344}$
$\frac{1583365445937429 \sin(30x)}{792281625142643375935439503360}$	+	$\frac{26917212580936293 \sin(32x)}{2535301200456458802993406410752}$
$\frac{33250674364686009 \sin(34x)}{633825300114114700748351602688}$	+	$\frac{306645108029882083 \sin(36x)}{1267650600228229401496703205376}$
$\frac{661707864696061337 \sin(38x)}{633825300114114700748351602688}$	+	$\frac{53598337040380968297 \sin(40x)}{12676506002282294014967032053760}$
$\frac{2552301763827665157 \sin(42x)}{158456325028528675187087900672}$	+	$\frac{18330167212944140673 \sin(44x)}{316912650057057350374175801344}$
$\frac{31081587882818325489 \sin(46x)}{158456325028528675187087900672}$	+	$\frac{797760755659003687551 \sin(48x)}{1267650600228229401496703205376}$
$\frac{15157454357521070063469 \sin(50x)}{7922816251426433759354395033600}$	+	$\frac{3497874082504862322339 \sin(52x)}{633825300114114700748351602688}$
$\frac{4793383001951107626909 \sin(54x)}{316912650057057350374175801344}$	+	$\frac{49988137020347265252051 \sin(56x)}{1267650600228229401496703205376}$
$\frac{15513559764935358181671 \sin(58x)}{158456325028528675187087900672}$	+	$\frac{367154247770136810299547 \sin(60x)}{5720500053966970302409071}$
$\frac{82905797883579279745059 \sin(62x)}{158456325028528675187087900672}$	+	$\frac{5720500053966970302409071 \sin(64x)}{5070602400912917605986812821504}$
$\frac{2946924270225408943665279 \sin(66x)}{1267650600228229401496703205376}$	+	$\frac{11614348594417788189739629 \sin(68x)}{79088183285797319577750807}$
$\frac{54753357659398144323058251 \sin(70x)}{6338253001141147007483516026880}$	+	$\frac{79088183285797319577750807 \sin(72x)}{5070602400912917605986812821504}$
$\frac{2137518467183711339939211 \sin(74x)}{79228162514264337593543950336}$	+	$\frac{7087561233293358653482647 \sin(76x)}{158456325028528675187087900672}$
$\frac{5633702518771644057896463 \sin(78x)}{79228162514264337593543950336}$	+	$\frac{343655853645070287531684243 \sin(80x)}{3169126500570573503741758013440}$
$\frac{25145550266712460063293969 \sin(82x)}{158456325028528675187087900672}$	+	$\frac{70647022177906435415921151 \sin(84x)}{246891178985368578736823811}$
$\frac{47645666119983409931667753 \sin(86x)}{158456325028528675187087900672}$	+	$\frac{246891178985368578736823811 \sin(88x)}{633825300114114700748351602688}$
$\frac{192026472544175561239751853 \sin(90x)}{396140812571321687967719751680}$	+	$\frac{91838747738518746679881321 \sin(92x)}{158456325028528675187087900672}$
$\frac{52758429551915024688442461 \sin(94x)}{79228162514264337593543950336}$	+	$\frac{932065588750498769495816811 \sin(96x)}{1267650600228229401496703205376}$
$\frac{247282707219520081702971807 \sin(98x)}{316912650057057350374175801344}$	+	$\frac{12611418068195524166851562157 \sin(100x)}{15845632502852867518708790067200}$
$\frac{247282707219520081702971807 \sin(102x)}{316912650057057350374175801344}$	+	$\frac{932065588750498769495816811 \sin(104x)}{1267650600228229401496703205376}$
$\frac{52758429551915024688442461 \sin(106x)}{79228162514264337593543950336}$	+	$\frac{91838747738518746679881321 \sin(108x)}{158456325028528675187087900672}$
$\frac{192026472544175561239751853 \sin(110x)}{396140812571321687967719751680}$	+	$\frac{246891178985368578736823811 \sin(112x)}{633825300114114700748351602688}$
$\frac{47645666119983409931667753 \sin(114x)}{158456325028528675187087900672}$	+	$\frac{70647022177906435415921151 \sin(116x)}{316912650057057350374175801344}$
$\frac{25145550266712460063293969 \sin(118x)}{158456325028528675187087900672}$	+	$\frac{246891178985368578736823811 \sin(120x)}{316912650057057350374175801344}$

input `Int [Sin[x]^99*Sin[101*x],x]`

output `-1/1267650600228229401496703205376*Sin[2*x] + (99*Sin[4*x])/2535301200456458802993406410752 - (1617*Sin[6*x])/1267650600228229401496703205376 + (156849*Sin[8*x])/5070602400912917605986812821504 - (470547*Sin[10*x])/792281625142643375935439503360 + (2980131*Sin[12*x])/316912650057057350374175801344 - (20009451*Sin[14*x])/158456325028528675187087900672 + (1860878943*Sin[16*x])/1267650600228229401496703205376 - (4755579521*Sin[18*x])/316912650057057350374175801344 + (432757736411*Sin[20*x])/3169126500570573503741758013440 - (354074511609*Sin[22*x])/316912650057057350374175801344 + (10504210511067*Sin[24*x])/1267650600228229401496703205376 - (8888178124749*Sin[26*x])/158456325028528675187087900672 + (110467356693309*Sin[28*x])/316912650057057350374175801344 - (1583365445937429*Sin[30*x])/792281625142643375935439503360 + (26917212580936293*Sin[32*x])/2535301200456458802993406410752 - (33250674364686009*Sin[34*x])/633825300114114700748351602688 + (306645108029882083*Sin[36*x])/1267650600228229401496703205376 - (661707864696061337*Sin[38*x])/633825300114114700748351602688 + (53598337040380968297*Sin[40*x])/12676506002282294014967032053760 - (2552301763827665157*Sin[42*x])/158456325028528675187087900672 + (18330167212944140673*Sin[44*x])/316912650057057350374175801344 - (31081587882818325489*Sin[46*x])/158456325028528675187087900672 + (797760755659003687551*Sin[48*x])/1267650600228229401496703205376 - (15157454357521070063469*Sin[50*x])/792281625142643375935439...`

3.46.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4854 `Int[(F_)[(a_.) + (b_.)*(x_)]^(p_.)*(G_)[(c_.) + (d_.)*(x_)]^(q_.), x_Symbol] := Int[ExpandTrigReduce[ActivateTrig[F[a + b*x]^p*G[c + d*x]^q], x], x] /; FreeQ[{a, b, c, d}, x] && (EqQ[F, sin] || EqQ[F, cos]) && (EqQ[G, sin] || EqQ[G, cos]) && IGtQ[p, 0] && IGtQ[q, 0]`

3.46.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 601 vs. $2(10) = 20$.

Time = 1.16 (sec) , antiderivative size = 602, normalized size of antiderivative = 50.17

Expression too large to display

input `int(sin(101*x)*sin(x)^99,x)`

output

```
-354074511609/316912650057057350374175801344*sin(178*x)+343655853645070287
531684243/3169126500570573503741758013440*sin(80*x)+1/12676506002282294014
9670320537600*sin(200*x)-1583365445937429/792281625142643375935439503360*s
in(170*x)-15157454357521070063469/7922816251426433759354395033600*sin(150*
x)-8888178124749/158456325028528675187087900672*sin(174*x)+2980131/3169126
50057057350374175801344*sin(12*x)-20009451/158456325028528675187087900672*
sin(14*x)+1860878943/1267650600228229401496703205376*sin(16*x)-4755579521/
316912650057057350374175801344*sin(18*x)-1583365445937429/7922816251426433
75935439503360*sin(30*x)-354074511609/316912650057057350374175801344*sin(2
2*x)+156849/5070602400912917605986812821504*sin(8*x)-470547/79228162514264
3375935439503360*sin(10*x)-1617/1267650600228229401496703205376*sin(6*x)+4
32757736411/3169126500570573503741758013440*sin(20*x)+99/25353012004564588
02993406410752*sin(4*x)+12611418068195524166851562157/15845632502852867518
708790067200*sin(100*x)-1/1267650600228229401496703205376*sin(2*x)-2472827
07219520081702971807/316912650057057350374175801344*sin(98*x)-192026472544
175561239751853/396140812571321687967719751680*sin(90*x)-54753357659398144
323058251/6338253001141147007483516026880*sin(130*x)+349787408250486232233
9/633825300114114700748351602688*sin(148*x)+26917212580936293/253530120045
6458802993406410752*sin(168*x)-2552301763827665157/15845632502852867518708
7900672*sin(158*x)+99/2535301200456458802993406410752*sin(196*x)-294692...
```

3.46.5 Fricas [F(-1)]

Timed out.

$$\int \sin^{99}(x) \sin(101x) dx = \text{Timed out}$$

input `integrate(sin(101*x)*sin(x)^99,x, algorithm="fricas")`

output `Timed out`

3.46.6 Sympy [F(-1)]

Timed out.

$$\int \sin^{99}(x) \sin(101x) dx = \text{Timed out}$$

input `integrate(sin(101*x)*sin(x)**99,x)`

output `Timed out`

3.46.7 Maxima [F(-2)]

Exception generated.

$$\int \sin^{99}(x) \sin(101x) dx = \text{Exception raised: RuntimeError}$$

input `integrate(sin(101*x)*sin(x)^99,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: Memory limit reached. Please jump to an outer pointer, quit program and enlarge thememory limits before executing the program again.`

3.46.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 601 vs. $2(10) = 20$.

Time = 0.29 (sec) , antiderivative size = 601, normalized size of antiderivative = 50.08

$$\int \sin^{99}(x) \sin(101x) dx = \text{Too large to display}$$

input `integrate(sin(101*x)*sin(x)^99,x, algorithm="giac")`

output

```

1/126765060022822940149670320537600*sin(200*x) - 1/12676506002282294014967
03205376*sin(198*x) + 99/2535301200456458802993406410752*sin(196*x) - 1617
/1267650600228229401496703205376*sin(194*x) + 156849/507060240091291760598
6812821504*sin(192*x) - 470547/792281625142643375935439503360*sin(190*x) +
2980131/316912650057057350374175801344*sin(188*x) - 20009451/158456325028
528675187087900672*sin(186*x) + 1860878943/1267650600228229401496703205376
*sin(184*x) - 4755579521/316912650057057350374175801344*sin(182*x) + 43275
7736411/3169126500570573503741758013440*sin(180*x) - 354074511609/31691265
0057057350374175801344*sin(178*x) + 10504210511067/12676506002282294014967
03205376*sin(176*x) - 8888178124749/158456325028528675187087900672*sin(174
*x) + 110467356693309/316912650057057350374175801344*sin(172*x) - 15833654
45937429/792281625142643375935439503360*sin(170*x) + 26917212580936293/253
5301200456458802993406410752*sin(168*x) - 33250674364686009/63382530011411
4700748351602688*sin(166*x) + 306645108029882083/1267650600228229401496703
205376*sin(164*x) - 661707864696061337/633825300114114700748351602688*sin(
162*x) + 53598337040380968297/12676506002282294014967032053760*sin(160*x)
- 2552301763827665157/158456325028528675187087900672*sin(158*x) + 18330167
212944140673/316912650057057350374175801344*sin(156*x) - 31081587882818325
489/158456325028528675187087900672*sin(154*x) + 797760755659003687551/1267
650600228229401496703205376*sin(152*x) - 15157454357521070063469/792281...

```

3.46.9 Mupad [B] (verification not implemented)

Time = 21.21 (sec) , antiderivative size = 601, normalized size of antiderivative = 50.08

$$\int \sin^{99}(x) \sin(101x) dx = \text{Too large to display}$$

input `int(sin(101*x)*sin(x)^99,x)`

output

```
(99*sin(4*x))/2535301200456458802993406410752 - sin(2*x)/12676506002282294
01496703205376 - (1617*sin(6*x))/1267650600228229401496703205376 + (156849
*sin(8*x))/5070602400912917605986812821504 - (470547*sin(10*x))/7922816251
42643375935439503360 + (2980131*sin(12*x))/316912650057057350374175801344
- (20009451*sin(14*x))/158456325028528675187087900672 + (1860878943*sin(16
*x))/1267650600228229401496703205376 - (4755579521*sin(18*x))/316912650057
057350374175801344 + (432757736411*sin(20*x))/3169126500570573503741758013
440 - (354074511609*sin(22*x))/316912650057057350374175801344 + (105042105
11067*sin(24*x))/1267650600228229401496703205376 - (8888178124749*sin(26*x
))/158456325028528675187087900672 + (110467356693309*sin(28*x))/3169126500
57057350374175801344 - (1583365445937429*sin(30*x))/7922816251426433759354
39503360 + (26917212580936293*sin(32*x))/2535301200456458802993406410752 -
(33250674364686009*sin(34*x))/633825300114114700748351602688 + (306645108
029882083*sin(36*x))/1267650600228229401496703205376 - (661707864696061337
*sin(38*x))/633825300114114700748351602688 + (53598337040380968297*sin(40*
x))/12676506002282294014967032053760 - (2552301763827665157*sin(42*x))/158
456325028528675187087900672 + (18330167212944140673*sin(44*x))/31691265005
7057350374175801344 - (31081587882818325489*sin(46*x))/1584563250285286751
87087900672 + (797760755659003687551*sin(48*x))/12676506002282294014967032
05376 - (15157454357521070063469*sin(50*x))/792281625142643375935439503...
```

3.47 $\int e^{e^{x^2}} x dx$

3.47.1	Optimal result	345
3.47.2	Mathematica [A] (verified)	345
3.47.3	Rubi [A] (verified)	346
3.47.4	Maple [A] (verified)	347
3.47.5	Fricas [A] (verification not implemented)	347
3.47.6	Sympy [A] (verification not implemented)	348
3.47.7	Maxima [A] (verification not implemented)	348
3.47.8	Giac [A] (verification not implemented)	348
3.47.9	Mupad [B] (verification not implemented)	349

3.47.1 Optimal result

Integrand size = 9, antiderivative size = 10

$$\int e^{e^{x^2}} x dx = \frac{\text{ExpIntegralEi}(e^{x^2})}{2}$$

output `1/2*Ei(exp(x^2))`

3.47.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int e^{e^{x^2}} x dx = \frac{\text{ExpIntegralEi}(e^{x^2})}{2}$$

input `Integrate[E^E^x^2*x,x]`

output `ExpIntegralEi[E^x^2]/2`

3.47.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {7266, 2720, 2609}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int e^{e^{x^2}} x dx \\ \downarrow 7266 \\ \frac{1}{2} \int e^{e^{x^2}} dx^2 \\ \downarrow 2720 \\ \frac{1}{2} \int \frac{e^{e^{x^2}}}{x^2} de^{x^2} \\ \downarrow 2609 \\ \frac{\text{ExpIntegralEi}(e^{x^2})}{2} \end{array}$$

input `Int [E^E^x^2*x, x]`

output `ExpIntegralEi [E^x^2]/2`

3.47.3.1 Defintions of rubi rules used

rule 2609 `Int[(F_)^((g_)*(e_) + (f_)*(x_)))/((c_) + (d_)*(x_)), x_Symbol] := Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*(a_) + (b_)*x)] *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

```
rule 7266 Int[(u_)*(x_)^(m_.), x_Symbol] := Simp[1/(m + 1) Subst[Int[SubstFor[x^(m
+ 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && Function
OfQ[x^(m + 1), u, x]
```

3.47.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

method	result	size
derivativedivides	$-\frac{\text{Ei}_1(-e^{x^2})}{2}$	11
default	$-\frac{\text{Ei}_1(-e^{x^2})}{2}$	11
risch	$-\frac{\text{Ei}_1(-e^{x^2})}{2}$	11

```
input int(exp(exp(x^2))*x,x,method=_RETURNVERBOSE)
```

```
output -1/2*Ei(1,-exp(x^2))
```

3.47.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int e^{e^{x^2}} x dx = \frac{1}{2} \text{Ei}(e^{(x^2)})$$

```
input integrate(exp(exp(x^2))*x,x, algorithm="fracas")
```

```
output 1/2*Ei(e^(x^2))
```

3.47.6 Sympy [A] (verification not implemented)

Time = 0.66 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int e^{e^{x^2}} x dx = \frac{\text{Ei}(e^{x^2})}{2}$$

input `integrate(exp(exp(x**2))*x,x)`output `Ei(exp(x**2))/2`**3.47.7 Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int e^{e^{x^2}} x dx = \frac{1}{2} \text{Ei}(e^{(x^2)})$$

input `integrate(exp(exp(x^2))*x,x, algorithm="maxima")`output `1/2*Ei(e^(x^2))`**3.47.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int e^{e^{x^2}} x dx = \frac{1}{2} \text{Ei}(e^{(x^2)})$$

input `integrate(exp(exp(x^2))*x,x, algorithm="giac")`output `1/2*Ei(e^(x^2))`

3.47.9 Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int e^{e^{x^2}} x dx = \frac{\text{ei}(e^{x^2})}{2}$$

input `int(x*exp(exp(x^2)),x)`

output `ei(exp(x^2))/2`

$$3.48 \quad \int \frac{-1+3x-3x^2+x^3}{1+4x+6x^2+4x^3+x^4} dx$$

3.48.1	Optimal result	350
3.48.2	Mathematica [A] (verified)	350
3.48.3	Rubi [A] (verified)	351
3.48.4	Maple [A] (verified)	352
3.48.5	Fricas [A] (verification not implemented)	353
3.48.6	Sympy [A] (verification not implemented)	353
3.48.7	Maxima [A] (verification not implemented)	353
3.48.8	Giac [A] (verification not implemented)	354
3.48.9	Mupad [B] (verification not implemented)	354

3.48.1 Optimal result

Integrand size = 34, antiderivative size = 28

$$\int \frac{-1+3x-3x^2+x^3}{1+4x+6x^2+4x^3+x^4} dx = \frac{8}{3(1+x)^3} - \frac{6}{(1+x)^2} + \frac{6}{1+x} + \log(1+x)$$

output `8/3/(1+x)^3-6/(1+x)^2+6/(1+x)+ln(1+x)`

3.48.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int \frac{-1+3x-3x^2+x^3}{1+4x+6x^2+4x^3+x^4} dx = \frac{2(4+9x+9x^2)}{3(1+x)^3} + \log(1+x)$$

input `Integrate[(-1 + 3*x - 3*x^2 + x^3)/(1 + 4*x + 6*x^2 + 4*x^3 + x^4),x]`

output `(2*(4 + 9*x + 9*x^2))/(3*(1 + x)^3) + Log[1 + x]`

3.48.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2006, 2007, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3 - 3x^2 + 3x - 1}{x^4 + 4x^3 + 6x^2 + 4x + 1} dx \\
 & \quad \downarrow \text{2006} \\
 & \int \frac{(x-1)^3}{x^4 + 4x^3 + 6x^2 + 4x + 1} dx \\
 & \quad \downarrow \text{2007} \\
 & \int \frac{(x-1)^3}{(x+1)^4} dx \\
 & \quad \downarrow \text{49} \\
 & \int \left(\frac{1}{x+1} - \frac{6}{(x+1)^2} + \frac{12}{(x+1)^3} - \frac{8}{(x+1)^4} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{6}{x+1} - \frac{6}{(x+1)^2} + \frac{8}{3(x+1)^3} + \log(x+1)
 \end{aligned}$$

input `Int[(-1 + 3*x - 3*x^2 + x^3)/(1 + 4*x + 6*x^2 + 4*x^3 + x^4), x]`

output `8/(3*(1 + x)^3) - 6/(1 + x)^2 + 6/(1 + x) + Log[1 + x]`

3.48.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`


```
rule 2006 Int[(u_.)*(Px_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px, x]],
b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Int[u*(a + b*x)^Expon[Px,
x], x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; PolyQ[Px, x] && GtQ[Expon[P
x, x], 1] && NeQ[Coeff[Px, x, 0], 0] && !MatchQ[Px, (a_.)*(v_)^Expon[Px, x
] /; FreeQ[a, x] && LinearQ[v, x]]
```

```
rule 2007 Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px,
x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Int[u*(a + b*x)^(Ex
pon[Px, x]*p), x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; IntegerQ[p] && Pol
yQ[Px, x] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

3.48.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

method	result	size
norman	$\frac{6x+6x^2+\frac{8}{3}}{(1+x)^3} + \ln(1+x)$	22
default	$\frac{8}{3(1+x)^3} - \frac{6}{(1+x)^2} + \frac{6}{1+x} + \ln(1+x)$	27
risch	$\frac{6x+6x^2+\frac{8}{3}}{x^3+3x^2+3x+1} + \ln(1+x)$	32
parallelrisc	$\frac{3 \ln(1+x)x^3+8+9 \ln(1+x)x^2+9 \ln(1+x)x+18x^2+3 \ln(1+x)+18x}{3x^3+9x^2+9x+3}$	59

```
input int((x^3-3*x^2+3*x-1)/(x^4+4*x^3+6*x^2+4*x+1),x,method=_RETURNVERBOSE)
```

```
output (6*x+6*x^2+8/3)/(1+x)^3+ln(1+x)
```

3.48.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.64

$$\int \frac{-1 + 3x - 3x^2 + x^3}{1 + 4x + 6x^2 + 4x^3 + x^4} dx = \frac{18x^2 + 3(x^3 + 3x^2 + 3x + 1)\log(x + 1) + 18x + 8}{3(x^3 + 3x^2 + 3x + 1)}$$

input `integrate((x^3-3*x^2+3*x-1)/(x^4+4*x^3+6*x^2+4*x+1),x, algorithm="fricas")`output `1/3*(18*x^2 + 3*(x^3 + 3*x^2 + 3*x + 1)*log(x + 1) + 18*x + 8)/(x^3 + 3*x^2 + 3*x + 1)`**3.48.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{-1 + 3x - 3x^2 + x^3}{1 + 4x + 6x^2 + 4x^3 + x^4} dx = \frac{18x^2 + 18x + 8}{3x^3 + 9x^2 + 9x + 3} + \log(x + 1)$$

input `integrate((x**3-3*x**2+3*x-1)/(x**4+4*x**3+6*x**2+4*x+1),x)`output `(18*x**2 + 18*x + 8)/(3*x**3 + 9*x**2 + 9*x + 3) + log(x + 1)`**3.48.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.14

$$\int \frac{-1 + 3x - 3x^2 + x^3}{1 + 4x + 6x^2 + 4x^3 + x^4} dx = \frac{2(9x^2 + 9x + 4)}{3(x^3 + 3x^2 + 3x + 1)} + \log(x + 1)$$

input `integrate((x^3-3*x^2+3*x-1)/(x^4+4*x^3+6*x^2+4*x+1),x, algorithm="maxima")`output `2/3*(9*x^2 + 9*x + 4)/(x^3 + 3*x^2 + 3*x + 1) + log(x + 1)`

3.48.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

$$\int \frac{-1 + 3x - 3x^2 + x^3}{1 + 4x + 6x^2 + 4x^3 + x^4} dx = \frac{2(9x^2 + 9x + 4)}{3(x+1)^3} + \log(|x+1|)$$

input `integrate((x^3-3*x^2+3*x-1)/(x^4+4*x^3+6*x^2+4*x+1),x, algorithm="giac")`output `2/3*(9*x^2 + 9*x + 4)/(x + 1)^3 + log(abs(x + 1))`**3.48.9 Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.75

$$\int \frac{-1 + 3x - 3x^2 + x^3}{1 + 4x + 6x^2 + 4x^3 + x^4} dx = \ln(x+1) + \frac{6x^2 + 6x + \frac{8}{3}}{(x+1)^3}$$

input `int((3*x - 3*x^2 + x^3 - 1)/(4*x + 6*x^2 + 4*x^3 + x^4 + 1),x)`output `log(x + 1) + (6*x + 6*x^2 + 8/3)/(x + 1)^3`

$$3.49 \quad \int \sqrt{\frac{1-x}{1+x}} dx$$

3.49.1	Optimal result	355
3.49.2	Mathematica [A] (verified)	355
3.49.3	Rubi [A] (verified)	356
3.49.4	Maple [A] (verified)	357
3.49.5	Fricas [A] (verification not implemented)	357
3.49.6	Sympy [F]	358
3.49.7	Maxima [A] (verification not implemented)	358
3.49.8	Giac [A] (verification not implemented)	358
3.49.9	Mupad [B] (verification not implemented)	359

3.49.1 Optimal result

Integrand size = 15, antiderivative size = 38

$$\int \sqrt{\frac{1-x}{1+x}} dx = \sqrt{\frac{1-x}{1+x}}(1+x) - 2 \arctan \left(\sqrt{\frac{1-x}{1+x}} \right)$$

output `((1-x)/(1+x))^(1/2)*(1+x)-2*arctan(((1-x)/(1+x))^(1/2))`

3.49.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.68

$$\int \sqrt{\frac{1-x}{1+x}} dx = \frac{\sqrt{\frac{1-x}{1+x}} \sqrt{1+x} \left(\sqrt{1-x^2} - 2 \arctan \left(\frac{\sqrt{1-x^2}}{-1+x} \right) \right)}{\sqrt{1-x}}$$

input `Integrate[Sqrt[(1 - x)/(1 + x)],x]`

output `(Sqrt[(1 - x)/(1 + x)]*Sqrt[1 + x]*(Sqrt[1 - x^2] - 2*ArcTan[Sqrt[1 - x^2]/(-1 + x)]))/Sqrt[1 - x]`

3.49.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.50, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2051, 252, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\frac{1-x}{x+1}} dx \\
 & \quad \downarrow \text{2051} \\
 & -4 \int \frac{1-x}{(x+1) \left(\frac{1-x}{x+1} + 1\right)^2} d\sqrt{\frac{1-x}{x+1}} \\
 & \quad \downarrow \text{252} \\
 & -4 \left(\frac{1}{2} \int \frac{1}{\frac{1-x}{x+1} + 1} d\sqrt{\frac{1-x}{x+1}} - \frac{\sqrt{\frac{1-x}{x+1}}}{2 \left(\frac{1-x}{x+1} + 1\right)} \right) \\
 & \quad \downarrow \text{216} \\
 & -4 \left(\frac{1}{2} \arctan \left(\sqrt{\frac{1-x}{x+1}} \right) - \frac{\sqrt{\frac{1-x}{x+1}}}{2 \left(\frac{1-x}{x+1} + 1\right)} \right)
 \end{aligned}$$

input `Int[Sqrt[(1 - x)/(1 + x)],x]`

output `-4*(-1/2*Sqrt[(1 - x)/(1 + x)]/(1 + (1 - x)/(1 + x)) + ArcTan[Sqrt[(1 - x)/(1 + x)]]/2)`

3.49.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

```
rule 252 Int[((c_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*((m - 1)/(2*b*(p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

```
rule 2051 Int[(((e_.)*((a_.) + (b_.)*(x_)^(n_.)))/((c_) + (d_.)*(x_)^(n_.)))^(p_), x_Symbol] :> With[{q = Denominator[p]}, Simp[q*e*((b*c - a*d)/n) Subst[Int[x^(q*(p + 1) - 1)*((-a)*e + c*x^q)^(1/n - 1)/(b*e - d*x^q)^(1/n + 1), x], x, (e*((a + b*x^n)/(c + d*x^n)))^(1/q)], x] /; FreeQ[{a, b, c, d, e}, x] && FractionQ[p] && IntegerQ[1/n]
```

3.49.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.03

method	result
default	$\frac{\sqrt{-\frac{-1+x}{1+x}}(1+x)(\sqrt{-x^2+1}+\arcsin(x))}{\sqrt{-(-1+x)(1+x)}}$
risch	$(1+x)\sqrt{-\frac{-1+x}{1+x}} - \frac{\arcsin(x)\sqrt{-\frac{-1+x}{1+x}}\sqrt{-(-1+x)(1+x)}}{-1+x}$
trager	$(1+x)\sqrt{-\frac{-1+x}{1+x}} + \text{RootOf}(_Z^2 + 1) \ln\left(\text{RootOf}(_Z^2 + 1)\sqrt{-\frac{-1+x}{1+x}}x + \text{RootOf}(_Z^2 + 1)\right)$

```
input int(((1-x)/(1+x))^(1/2),x,method=_RETURNVERBOSE)
```

```
output ((-(-1+x)/(1+x))^(1/2)*(1+x)/(-(-1+x)*(1+x))^(1/2)*((-x^2+1)^(1/2)+arcsin(x))
```

3.49.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int \sqrt{\frac{1-x}{1+x}} dx = (x+1)\sqrt{-\frac{x-1}{x+1}} - 2 \arctan\left(\sqrt{-\frac{x-1}{x+1}}\right)$$

```
input integrate(((1-x)/(1+x))^(1/2),x, algorithm="fricas")
```

3.49. $\int \sqrt{\frac{1-x}{1+x}} dx$

output `(x + 1)*sqrt(-(x - 1)/(x + 1)) - 2*arctan(sqrt(-(x - 1)/(x + 1)))`

3.49.6 Sympy [F]

$$\int \sqrt{\frac{1-x}{1+x}} dx = \int \sqrt{\frac{1-x}{x+1}} dx$$

input `integrate(((1-x)/(1+x))**(1/2),x)`

output `Integral(sqrt((1 - x)/(x + 1)), x)`

3.49.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.13

$$\int \sqrt{\frac{1-x}{1+x}} dx = -\frac{2\sqrt{-\frac{x-1}{x+1}}}{\frac{x-1}{x+1} - 1} - 2 \arctan\left(\sqrt{-\frac{x-1}{x+1}}\right)$$

input `integrate(((1-x)/(1+x))^(1/2),x, algorithm="maxima")`

output `-2*sqrt(-(x - 1)/(x + 1))/((x - 1)/(x + 1) - 1) - 2*arctan(sqrt(-(x - 1)/(x + 1)))`

3.49.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.76

$$\int \sqrt{\frac{1-x}{1+x}} dx = \frac{1}{2} \pi \operatorname{sgn}(x+1) + \arcsin(x) \operatorname{sgn}(x+1) + \sqrt{-x^2+1} \operatorname{sgn}(x+1)$$

input `integrate(((1-x)/(1+x))^(1/2),x, algorithm="giac")`

output `1/2*pi*sgn(x + 1) + arcsin(x)*sgn(x + 1) + sqrt(-x^2 + 1)*sgn(x + 1)`

3.49. $\int \sqrt{\frac{1-x}{1+x}} dx$

3.49.9 Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.13

$$\int \sqrt{\frac{1-x}{1+x}} dx = -2 \operatorname{atan}\left(\sqrt{-\frac{x-1}{x+1}}\right) - \frac{2\sqrt{-\frac{x-1}{x+1}}}{\frac{x-1}{x+1} - 1}$$

input `int((-x - 1)/(x + 1))^(1/2),x)`output `- 2*atan((-x - 1)/(x + 1))^(1/2) - (2*(-x - 1)/(x + 1))^(1/2)/((x - 1)/(x + 1) - 1)`

3.50 $\int \frac{1}{-1+\sqrt{x}} dx$

3.50.1	Optimal result	360
3.50.2	Mathematica [A] (verified)	360
3.50.3	Rubi [A] (verified)	361
3.50.4	Maple [A] (verified)	362
3.50.5	Fricas [A] (verification not implemented)	362
3.50.6	Sympy [A] (verification not implemented)	363
3.50.7	Maxima [A] (verification not implemented)	363
3.50.8	Giac [A] (verification not implemented)	363
3.50.9	Mupad [B] (verification not implemented)	364

3.50.1 Optimal result

Integrand size = 9, antiderivative size = 20

$$\int \frac{1}{-1+\sqrt{x}} dx = 2\sqrt{x} + 2 \log(1 - \sqrt{x})$$

output `2*x^(1/2)+2*ln(1-x^(1/2))`

3.50.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{1}{-1+\sqrt{x}} dx = 2\sqrt{x} + 2 \log(-1 + \sqrt{x})$$

input `Integrate[(-1 + Sqrt[x])^(-1), x]`

output `2*Sqrt[x] + 2*Log[-1 + Sqrt[x]]`

3.50.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {774, 25, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{x}-1} dx \\
 & \quad \downarrow 774 \\
 & 2 \int -\frac{\sqrt{x}}{1-\sqrt{x}} d\sqrt{x} \\
 & \quad \downarrow 25 \\
 & -2 \int \frac{\sqrt{x}}{1-\sqrt{x}} d\sqrt{x} \\
 & \quad \downarrow 49 \\
 & -2 \int \left(\frac{1}{1-\sqrt{x}} - 1 \right) d\sqrt{x} \\
 & \quad \downarrow 2009 \\
 & 2(\sqrt{x} + \log(1-\sqrt{x}))
 \end{aligned}$$

input `Int[(-1 + Sqrt[x])^(-1),x]`

output `2*(Sqrt[x] + Log[1 - Sqrt[x]])`

3.50.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

```
rule 774 Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[n]},
Simp[k Subst[Int[x^(k - 1)*(a + b*x^(k*n))^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, p}, x] && FractionQ[n]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

3.50.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

method	result	size
derivativedivides	$2\sqrt{x} + 2 \ln(-1 + \sqrt{x})$	15
meijerg	$2\sqrt{x} + 2 \ln(1 - \sqrt{x})$	17
trager	$2\sqrt{x} + \ln(2\sqrt{x} - 1 - x)$	18
default	$\ln(-1 + x) + 2\sqrt{x} + \ln(-1 + \sqrt{x}) - \ln(1 + \sqrt{x})$	25

```
input int(1/(-1+x^(1/2)),x,method=_RETURNVERBOSE)
```

```
output 2*x^(1/2)+2*ln(-1+x^(1/2))
```

3.50.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \frac{1}{-1 + \sqrt{x}} dx = 2\sqrt{x} + 2 \log(\sqrt{x} - 1)$$

```
input integrate(1/(-1+x^(1/2)),x, algorithm="fracas")
```

```
output 2*sqrt(x) + 2*log(sqrt(x) - 1)
```

3.50.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int \frac{1}{-1 + \sqrt{x}} dx = 2\sqrt{x} + 2 \log(\sqrt{x} - 1)$$

input `integrate(1/(-1+x**(1/2)),x)`output `2*sqrt(x) + 2*log(sqrt(x) - 1)`**3.50.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int \frac{1}{-1 + \sqrt{x}} dx = 2\sqrt{x} + 2 \log(\sqrt{x} - 1) - 2$$

input `integrate(1/(-1+x^(1/2)),x, algorithm="maxima")`output `2*sqrt(x) + 2*log(sqrt(x) - 1) - 2`**3.50.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int \frac{1}{-1 + \sqrt{x}} dx = 2\sqrt{x} + 2 \log(|\sqrt{x} - 1|)$$

input `integrate(1/(-1+x^(1/2)),x, algorithm="giac")`output `2*sqrt(x) + 2*log(abs(sqrt(x) - 1))`

3.50.9 Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \frac{1}{-1 + \sqrt{x}} dx = 2 \ln(\sqrt{x} - 1) + 2\sqrt{x}$$

input `int(1/(x^(1/2) - 1),x)`

output `2*log(x^(1/2) - 1) + 2*x^(1/2)`

3.51 $\int \sqrt[4]{x} \log(x) dx$

3.51.1	Optimal result	365
3.51.2	Mathematica [A] (verified)	365
3.51.3	Rubi [A] (verified)	366
3.51.4	Maple [A] (verified)	366
3.51.5	Fricas [A] (verification not implemented)	367
3.51.6	Sympy [B] (verification not implemented)	367
3.51.7	Maxima [A] (verification not implemented)	368
3.51.8	Giac [A] (verification not implemented)	368
3.51.9	Mupad [B] (verification not implemented)	368

3.51.1 Optimal result

Integrand size = 8, antiderivative size = 21

$$\int \sqrt[4]{x} \log(x) dx = -\frac{16x^{5/4}}{25} + \frac{4}{5}x^{5/4} \log(x)$$

output `-16/25*x^(5/4)+4/5*x^(5/4)*ln(x)`

3.51.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \sqrt[4]{x} \log(x) dx = \frac{4}{25}x^{5/4}(-4 + 5 \log(x))$$

input `Integrate[x^(1/4)*Log[x],x]`

output `(4*x^(5/4)*(-4 + 5*Log[x]))/25`

3.51.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt[4]{x} \log(x) dx$$

↓ 2741

$$\frac{4}{5}x^{5/4} \log(x) - \frac{16x^{5/4}}{25}$$

input `Int[x^(1/4)*Log[x],x]`

output `(-16*x^(5/4))/25 + (4*x^(5/4)*Log[x])/5`

3.51.3.1 Defintions of rubi rules used

rule 2741 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_))^(m_.), x_Symbol] :>
Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(m + 1))/(d*(m + 1)^2), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]`

3.51.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

method	result	size
derivativedivides	$-\frac{16x^{\frac{5}{4}}}{25} + \frac{4x^{\frac{5}{4}} \ln(x)}{5}$	14
default	$-\frac{16x^{\frac{5}{4}}}{25} + \frac{4x^{\frac{5}{4}} \ln(x)}{5}$	14
risch	$-\frac{16x^{\frac{5}{4}}}{25} + \frac{4x^{\frac{5}{4}} \ln(x)}{5}$	14

input `int(x^(1/4)*ln(x),x,method=_RETURNVERBOSE)`

output `-16/25*x^(5/4)+4/5*x^(5/4)*ln(x)`

3.51.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

$$\int \sqrt[4]{x} \log(x) dx = \frac{4}{25} (5x \log(x) - 4x)x^{\frac{1}{4}}$$

input `integrate(x^(1/4)*log(x),x, algorithm="fricas")`

output `4/25*(5*x*log(x) - 4*x)*x^(1/4)`

3.51.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. $2(19) = 38$.

Time = 1.66 (sec) , antiderivative size = 105, normalized size of antiderivative = 5.00

$$\int \sqrt[4]{x} \log(x) dx = \begin{cases} -\frac{4x^{\frac{5}{4}} \log(\frac{1}{x})}{5} + \frac{4x^{\frac{5}{4}} \log(x)}{5} - \frac{32x^{\frac{5}{4}}}{25} & \text{for } \frac{1}{|x|} < 1 \wedge |x| < 1 \\ \frac{4x^{\frac{5}{4}} \log(x)}{5} - \frac{16x^{\frac{5}{4}}}{25} & \text{for } |x| < 1 \\ -\frac{4x^{\frac{5}{4}} \log(\frac{1}{x})}{5} - \frac{16x^{\frac{5}{4}}}{25} & \text{for } \frac{1}{|x|} < 1 \\ -G_{3,3}^{2,1} \left(\begin{matrix} 1 & \frac{9}{4}, \frac{9}{4} \\ \frac{5}{4}, \frac{5}{4} \end{matrix} \middle| x \right) + G_{3,3}^{0,3} \left(\begin{matrix} \frac{9}{4}, \frac{9}{4}, 1 \\ \frac{5}{4}, \frac{5}{4}, 0 \end{matrix} \middle| x \right) & \text{otherwise} \end{cases}$$

input `integrate(x**(1/4)*ln(x),x)`

output `Piecewise((-4*x**(5/4)*log(1/x)/5 + 4*x**(5/4)*log(x)/5 - 32*x**(5/4)/25, (Abs(x) < 1) & (1/Abs(x) < 1)), (4*x**(5/4)*log(x)/5 - 16*x**(5/4)/25, Abs(x) < 1), (-4*x**(5/4)*log(1/x)/5 - 16*x**(5/4)/25, 1/Abs(x) < 1), (-meijerg(((1,), (9/4, 9/4)), ((5/4, 5/4), (0,)), x) + meijerg(((9/4, 9/4, 1), ()), ((, (5/4, 5/4, 0)), x), True))`

3.51.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \sqrt[4]{x} \log(x) dx = \frac{4}{5} x^{\frac{5}{4}} \log(x) - \frac{16}{25} x^{\frac{5}{4}}$$

input `integrate(x^(1/4)*log(x),x, algorithm="maxima")`output `4/5*x^(5/4)*log(x) - 16/25*x^(5/4)`**3.51.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \sqrt[4]{x} \log(x) dx = \frac{4}{5} x^{\frac{5}{4}} \log(x) - \frac{16}{25} x^{\frac{5}{4}}$$

input `integrate(x^(1/4)*log(x),x, algorithm="giac")`output `4/5*x^(5/4)*log(x) - 16/25*x^(5/4)`**3.51.9 Mupad [B] (verification not implemented)**

Time = 15.09 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.43

$$\int \sqrt[4]{x} \log(x) dx = \frac{4 x^{5/4} (\ln(x) - \frac{4}{5})}{5}$$

input `int(x^(1/4)*log(x),x)`output `(4*x^(5/4)*(log(x) - 4/5))/5`

$$3.52 \quad \int \frac{1}{(1+\sqrt{x})\sqrt{x-x^2}} dx$$

3.52.1	Optimal result	369
3.52.2	Mathematica [A] (verified)	369
3.52.3	Rubi [B] (verified)	370
3.52.4	Maple [B] (verified)	371
3.52.5	Fricas [B] (verification not implemented)	372
3.52.6	Sympy [F]	372
3.52.7	Maxima [F]	372
3.52.8	Giac [A] (verification not implemented)	373
3.52.9	Mupad [F(-1)]	373

3.52.1 Optimal result

Integrand size = 21, antiderivative size = 20

$$\int \frac{1}{(1+\sqrt{x})\sqrt{x-x^2}} dx = -\frac{2(1-\sqrt{x})}{\sqrt{1-x}}$$

output `-2*(1-x^(1/2))/(1-x)^(1/2)`

3.52.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.05

$$\int \frac{1}{(1+\sqrt{x})\sqrt{x-x^2}} dx = \frac{2(-\sqrt{x}+x)}{\sqrt{-((-1+x)x)}}$$

input `Integrate[1/((1+Sqrt[x])*Sqrt[x-x^2]),x]`

output `(2*(-Sqrt[x]+x))/Sqrt[-((-1+x)*x)]`

3.52.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 51 vs. $2(20) = 40$.

Time = 0.23 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.55, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2467, 1388, 946, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(\sqrt{x} + 1)\sqrt{x - x^2}} dx \\
 & \quad \downarrow \text{2467} \\
 & \frac{\sqrt{1-x}\sqrt{x} \int \frac{1}{(\sqrt{x+1})\sqrt{1-x}\sqrt{x}} dx}{\sqrt{x-x^2}} \\
 & \quad \downarrow \text{1388} \\
 & \frac{\sqrt{1-x}\sqrt{x} \int \frac{1}{\sqrt{1-\sqrt{x}}(\sqrt{x+1})^{3/2}\sqrt{x}} dx}{\sqrt{x-x^2}} \\
 & \quad \downarrow \text{946} \\
 & \frac{2\sqrt{1-x}\sqrt{x} \int \frac{1}{\sqrt{1-\sqrt{x}}(\sqrt{x+1})^{3/2}} d\sqrt{x}}{\sqrt{x-x^2}} \\
 & \quad \downarrow \text{48} \\
 & -\frac{2\sqrt{1-\sqrt{x}}\sqrt{1-x}\sqrt{x}}{\sqrt{\sqrt{x}+1}\sqrt{x-x^2}}
 \end{aligned}$$

input `Int[1/((1 + Sqrt[x])*Sqrt[x - x^2]),x]`

output `(-2*Sqrt[1 - Sqrt[x]]*Sqrt[1 - x]*Sqrt[x])/(Sqrt[1 + Sqrt[x]]*Sqrt[x - x^2])`

3.52.3.1 Defintions of rubi rules used

- rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`
- rule 946 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Simp[1/n Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n + 1, 0]`
- rule 1388 `Int[(u_.)*((a_) + (c_.)*(x_)^(n2_.))^(p_.)*((d_) + (e_.)*(x_)^(n_))^(q_.), x_Symbol] := Int[u*(d + e*x^n)^(p + q)*(a/d + (c/e)*x^n)^p, x] /; FreeQ[{a, c, d, e, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[c*d^2 + a*e^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[d, 0]))`
- rule 2467 `Int[(Fx_.)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Simp[Px^FracPart[p]/(x^(r*FracPart[p])*ExpandToSum[Px/x^r, x]^FracPart[p]) Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x], x] /; IGtQ[r, 0]] /; FreeQ[p, x] && PolyQ[Px, x] && !IntegerQ[p] && !MonomialQ[Px, x] && !PolyQ[Fx, x]`

3.52.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 40 vs. $2(16) = 32$.

Time = 0.16 (sec) , antiderivative size = 41, normalized size of antiderivative = 2.05

method	result	size
default	$\frac{2\sqrt{-(-1+x)x}}{\sqrt{x}(-1+x)} - \frac{2\sqrt{-(-1+x)^2-x+1}}{-1+x}$	41

input `int(1/(1+x^(1/2))/(-x^2+x)^(1/2),x,method=_RETURNVERBOSE)`

output `2*(-(-1+x)*x)^(1/2)/x^(1/2)/(-1+x)-2/(-1+x)*(-(-1+x)^2-x+1)^(1/2)`

3.52.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. $2(14) = 28$.

Time = 0.26 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.85

$$\int \frac{1}{(1 + \sqrt{x}) \sqrt{x - x^2}} dx = -\frac{2(\sqrt{-x^2 + xx} - \sqrt{-x^2 + x}\sqrt{x})}{x^2 - x}$$

input `integrate(1/(1+x^(1/2))/(-x^2+x)^(1/2),x, algorithm="fracas")`

output `-2*(sqrt(-x^2 + x)*x - sqrt(-x^2 + x)*sqrt(x))/(x^2 - x)`

3.52.6 Sympy [F]

$$\int \frac{1}{(1 + \sqrt{x}) \sqrt{x - x^2}} dx = \int \frac{1}{\sqrt{-x}(x - 1)(\sqrt{x} + 1)} dx$$

input `integrate(1/(1+x**(1/2))/(-x**2+x)**(1/2),x)`

output `Integral(1/(sqrt(-x*(x - 1))*(sqrt(x) + 1)), x)`

3.52.7 Maxima [F]

$$\int \frac{1}{(1 + \sqrt{x}) \sqrt{x - x^2}} dx = \int \frac{1}{\sqrt{-x^2 + x}(\sqrt{x} + 1)} dx$$

input `integrate(1/(1+x^(1/2))/(-x^2+x)^(1/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(-x^2 + x)*(sqrt(x) + 1)), x)`

3.52.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.05

$$\int \frac{1}{(1 + \sqrt{x}) \sqrt{x - x^2}} dx = \frac{4}{\frac{\sqrt{-x+1}-1}{\sqrt{x}} - 1} + 4$$

input `integrate(1/(1+x^(1/2))/(-x^2+x)^(1/2),x, algorithm="giac")`output `4/((sqrt(-x + 1) - 1)/sqrt(x) - 1) + 4`**3.52.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(1 + \sqrt{x}) \sqrt{x - x^2}} dx = \int \frac{1}{\sqrt{x - x^2} (\sqrt{x} + 1)} dx$$

input `int(1/((x - x^2)^(1/2)*(x^(1/2) + 1)),x)`output `int(1/((x - x^2)^(1/2)*(x^(1/2) + 1)), x)`

3.53
$$\int \frac{1}{(1 + \sqrt[4]{x})^{10} \sqrt{x}} dx$$

3.53.1 Optimal result 374
 3.53.2 Mathematica [A] (verified) 374
 3.53.3 Rubi [A] (verified) 375
 3.53.4 Maple [A] (verified) 376
 3.53.5 Fricas [B] (verification not implemented) 376
 3.53.6 Sympy [B] (verification not implemented) 377
 3.53.7 Maxima [A] (verification not implemented) 377
 3.53.8 Giac [A] (verification not implemented) 378
 3.53.9 Mupad [B] (verification not implemented) 378

3.53.1 Optimal result

Integrand size = 15, antiderivative size = 27

$$\int \frac{1}{(1 + \sqrt[4]{x})^{10} \sqrt{x}} dx = \frac{4}{9(1 + \sqrt[4]{x})^9} - \frac{1}{2(1 + \sqrt[4]{x})^8}$$

output `4/9/(x^(1/4)+1)^9-1/2/(x^(1/4)+1)^8`

3.53.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int \frac{1}{(1 + \sqrt[4]{x})^{10} \sqrt{x}} dx = \frac{-1 - 9\sqrt[4]{x}}{18(1 + \sqrt[4]{x})^9}$$

input `Integrate[1/((1 + x^(1/4))^10*Sqrt[x]),x]`

output `(-1 - 9*x^(1/4))/(18*(1 + x^(1/4))^9)`

3.53.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {798, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(\sqrt[4]{x} + 1)^{10} \sqrt{x}} dx$$

↓ 798

$$4 \int \frac{\sqrt[4]{x}}{(\sqrt[4]{x} + 1)^{10}} d\sqrt[4]{x}$$

↓ 53

$$4 \int \left(\frac{1}{(\sqrt[4]{x} + 1)^9} - \frac{1}{(\sqrt[4]{x} + 1)^{10}} \right) d\sqrt[4]{x}$$

↓ 2009

$$4 \left(\frac{1}{9(\sqrt[4]{x} + 1)^9} - \frac{1}{8(\sqrt[4]{x} + 1)^8} \right)$$

input `Int[1/((1 + x^(1/4))^10*Sqrt[x]),x]`

output `4*(1/(9*(1 + x^(1/4))^9) - 1/(8*(1 + x^(1/4))^8))`

3.53.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

3.53. $\int \frac{1}{(1 + \sqrt[4]{x})^{10} \sqrt{x}} dx$

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.53.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result	size
derivativedivides	$\frac{4}{9(x^{\frac{1}{4}}+1)^9} - \frac{1}{2(x^{\frac{1}{4}}+1)^8}$	20
default	$\frac{4}{9(x^{\frac{1}{4}}+1)^9} - \frac{1}{2(x^{\frac{1}{4}}+1)^8}$	20
meijerg	$\frac{\sqrt{x} (x^{\frac{7}{4}} + 9x^{\frac{3}{2}} + 36x^{\frac{5}{4}} + 84x + 126x^{\frac{3}{4}} + 126\sqrt{x} + 84x^{\frac{1}{4}} + 36)}{18(x^{\frac{1}{4}}+1)^9}$	46

input `int(1/x^(1/2)/(x^(1/4)+1)^10,x,method=_RETURNVERBOSE)`

output `4/9/(x^(1/4)+1)^9-1/2/(x^(1/4)+1)^8`

3.53.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 186 vs. $2(19) = 38$.

Time = 0.25 (sec) , antiderivative size = 186, normalized size of antiderivative = 6.89

$$\int \frac{1}{(1 + \sqrt[4]{x})^{10} \sqrt{x}} dx = \frac{9x^7 + 4209x^6 + 71109x^5 + 227277x^4 + 184587x^3 + 36099x^2 - 16(5x^6 + 648x^5 + 6813x^4 + 15288x^3 + 8847x^2 + 1152x + 15)x^{3/4} + 4(99x^6 + 5544x^5 + 38027x^4 + 60552x^3 + 24777x^2 + 2064x + 9)\sqrt{x} - 32(45x^6 + 1311x^5 + 6066x^4 + 6894x^3 + 1969x^2 + 99x)x^{1/4} + 999x - 1}{(x^9 - 9x^8 + 36x^7 - 84x^6 + 126x^5 - 126x^4 + 84x^3 - 36x^2 + 9x - 1)}$$

input `integrate(1/x^(1/2)/(x^(1/4)+1)^10,x, algorithm="fracas")`

output `-1/18*(9*x^7 + 4209*x^6 + 71109*x^5 + 227277*x^4 + 184587*x^3 + 36099*x^2 - 16*(5*x^6 + 648*x^5 + 6813*x^4 + 15288*x^3 + 8847*x^2 + 1152*x + 15)*x^(3/4) + 4*(99*x^6 + 5544*x^5 + 38027*x^4 + 60552*x^3 + 24777*x^2 + 2064*x + 9)*sqrt(x) - 32*(45*x^6 + 1311*x^5 + 6066*x^4 + 6894*x^3 + 1969*x^2 + 99*x)*x^(1/4) + 999*x - 1)/(x^9 - 9*x^8 + 36*x^7 - 84*x^6 + 126*x^5 - 126*x^4 + 84*x^3 - 36*x^2 + 9*x - 1)`

3.53. $\int \frac{1}{(1 + \sqrt[4]{x})^{10} \sqrt{x}} dx$

3.53.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 138 vs. $2(22) = 44$.

Time = 53.39 (sec) , antiderivative size = 138, normalized size of antiderivative = 5.11

$$\int \frac{1}{(1 + \sqrt[4]{x})^{10} \sqrt{x}} dx =$$

$$-\frac{\sqrt[4]{x}}{162x^{\frac{9}{4}} + 1512x^{\frac{7}{4}} + 2268x^{\frac{5}{4}} + 648x^{\frac{3}{4}} + 18\sqrt[4]{x} + 18x^{\frac{5}{2}} + 2268x^{\frac{3}{2}} + 162\sqrt{x} + 648x^2 + 1512x}$$

$$-\frac{9\sqrt{x}}{162x^{\frac{9}{4}} + 1512x^{\frac{7}{4}} + 2268x^{\frac{5}{4}} + 648x^{\frac{3}{4}} + 18\sqrt[4]{x} + 18x^{\frac{5}{2}} + 2268x^{\frac{3}{2}} + 162\sqrt{x} + 648x^2 + 1512x}$$

input `integrate(1/x**(1/2)/(x**(1/4)+1)**10,x)`

output `-x**(1/4)/(162*x**(9/4) + 1512*x**(7/4) + 2268*x**(5/4) + 648*x**(3/4) + 18*x**(1/4) + 18*x**(5/2) + 2268*x**(3/2) + 162*sqrt(x) + 648*x**2 + 1512*x) - 9*sqrt(x)/(162*x**(9/4) + 1512*x**(7/4) + 2268*x**(5/4) + 648*x**(3/4) + 18*x**(1/4) + 18*x**(5/2) + 2268*x**(3/2) + 162*sqrt(x) + 648*x**2 + 1512*x)`

3.53.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int \frac{1}{(1 + \sqrt[4]{x})^{10} \sqrt{x}} dx = -\frac{1}{2(x^{\frac{1}{4}} + 1)^8} + \frac{4}{9(x^{\frac{1}{4}} + 1)^9}$$

input `integrate(1/x^(1/2)/(x^(1/4)+1)^10,x, algorithm="maxima")`

output `-1/2/(x^(1/4) + 1)^8 + 4/9/(x^(1/4) + 1)^9`

3.53.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.59

$$\int \frac{1}{(1 + \sqrt[4]{x})^{10} \sqrt{x}} dx = -\frac{9x^{\frac{1}{4}} + 1}{18(x^{\frac{1}{4}} + 1)^9}$$

input `integrate(1/x^(1/2)/(x^(1/4)+1)^10,x, algorithm="giac")`output `-1/18*(9*x^(1/4) + 1)/(x^(1/4) + 1)^9`**3.53.9 Mupad [B] (verification not implemented)**

Time = 15.16 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.96

$$\int \frac{1}{(1 + \sqrt[4]{x})^{10} \sqrt{x}} dx =$$

$$-\frac{9x^{1/4} + 1}{2268x + \sqrt{x}(1512x + 648) + x^{3/4}(648x + 1512) + x^{1/4}(18x^2 + 2268x + 162) + 162x^2 + 18}$$

input `int(1/(x^(1/2)*(x^(1/4) + 1)^10),x)`output `-(9*x^(1/4) + 1)/(2268*x + x^(1/2)*(1512*x + 648) + x^(3/4)*(648*x + 1512) + x^(1/4)*(2268*x + 18*x^2 + 162) + 162*x^2 + 18)`

3.54 $\int \sqrt{1-x^2} dx$

3.54.1	Optimal result	379
3.54.2	Mathematica [A] (verified)	379
3.54.3	Rubi [A] (verified)	380
3.54.4	Maple [A] (verified)	381
3.54.5	Fricas [A] (verification not implemented)	381
3.54.6	Sympy [A] (verification not implemented)	382
3.54.7	Maxima [A] (verification not implemented)	382
3.54.8	Giac [A] (verification not implemented)	382
3.54.9	Mupad [B] (verification not implemented)	383

3.54.1 Optimal result

Integrand size = 11, antiderivative size = 23

$$\int \sqrt{1-x^2} dx = \frac{1}{2}x\sqrt{1-x^2} + \frac{\arcsin(x)}{2}$$

output `1/2*x*(-x^2+1)^(1/2)+1/2*arcsin(x)`

3.54.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.61

$$\int \sqrt{1-x^2} dx = \frac{1}{2}x\sqrt{1-x^2} - \arctan\left(\frac{\sqrt{1-x^2}}{1+x}\right)$$

input `Integrate[Sqrt[1 - x^2],x]`

output `(x*Sqrt[1 - x^2])/2 - ArcTan[Sqrt[1 - x^2]/(1 + x)]`

3.54.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {211, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{1-x^2} dx$$

$$\downarrow \text{211}$$

$$\frac{1}{2} \int \frac{1}{\sqrt{1-x^2}} dx + \frac{1}{2} \sqrt{1-x^2} x$$

$$\downarrow \text{223}$$

$$\frac{\arcsin(x)}{2} + \frac{1}{2} \sqrt{1-x^2} x$$

input `Int[Sqrt[1 - x^2], x]`

output `(x*Sqrt[1 - x^2])/2 + ArcSin[x]/2`

3.54.3.1 Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

3.54.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

method	result	size
default	$\frac{x\sqrt{-x^2+1}}{2} + \frac{\arcsin(x)}{2}$	18
risch	$-\frac{x(x^2-1)}{2\sqrt{-x^2+1}} + \frac{\arcsin(x)}{2}$	23
pseudoelliptic	$\frac{x\sqrt{-x^2+1}}{2} - \frac{\arctan\left(\frac{\sqrt{-x^2+1}}{x}\right)}{2}$	30
meijerg	$\frac{i(-2i\sqrt{\pi}x\sqrt{-x^2+1}-2i\sqrt{\pi}\arcsin(x))}{4\sqrt{\pi}}$	32
trager	$\frac{x\sqrt{-x^2+1}}{2} + \frac{\text{RootOf}(-Z^2+1)\ln(\text{RootOf}(-Z^2+1)\sqrt{-x^2+1}+x)}{2}$	41

input `int((-x^2+1)^(1/2),x,method=_RETURNVERBOSE)`output `1/2*x*(-x^2+1)^(1/2)+1/2*arcsin(x)`**3.54.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.35

$$\int \sqrt{1-x^2} dx = \frac{1}{2} \sqrt{-x^2+1} x - \arctan\left(\frac{\sqrt{-x^2+1}-1}{x}\right)$$

input `integrate((-x^2+1)^(1/2),x, algorithm="fracas")`output `1/2*sqrt(-x^2 + 1)*x - arctan((sqrt(-x^2 + 1) - 1)/x)`

3.54.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int \sqrt{1-x^2} dx = \frac{x\sqrt{1-x^2}}{2} + \frac{\operatorname{asin}(x)}{2}$$

input `integrate((-x**2+1)**(1/2),x)`output `x*sqrt(1 - x**2)/2 + asin(x)/2`**3.54.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \sqrt{1-x^2} dx = \frac{1}{2} \sqrt{-x^2+1}x + \frac{1}{2} \arcsin(x)$$

input `integrate((-x^2+1)^(1/2),x, algorithm="maxima")`output `1/2*sqrt(-x^2 + 1)*x + 1/2*arcsin(x)`**3.54.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \sqrt{1-x^2} dx = \frac{1}{2} \sqrt{-x^2+1}x + \frac{1}{2} \arcsin(x)$$

input `integrate((-x^2+1)^(1/2),x, algorithm="giac")`output `1/2*sqrt(-x^2 + 1)*x + 1/2*arcsin(x)`

3.54.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \sqrt{1-x^2} dx = \frac{\arcsin(x)}{2} + \frac{x\sqrt{1-x^2}}{2}$$

input `int((1 - x^2)^(1/2),x)`

output `asin(x)/2 + (x*(1 - x^2)^(1/2))/2`

3.55 $\int \frac{1}{\sqrt{1-4x-x^2}} dx$

3.55.1	Optimal result	384
3.55.2	Mathematica [B] (verified)	384
3.55.3	Rubi [A] (verified)	385
3.55.4	Maple [A] (verified)	386
3.55.5	Fricas [B] (verification not implemented)	386
3.55.6	Sympy [A] (verification not implemented)	386
3.55.7	Maxima [A] (verification not implemented)	387
3.55.8	Giac [B] (verification not implemented)	387
3.55.9	Mupad [B] (verification not implemented)	387

3.55.1 Optimal result

Integrand size = 14, antiderivative size = 10

$$\int \frac{1}{\sqrt{1-4x-x^2}} dx = \arcsin\left(\frac{2+x}{\sqrt{5}}\right)$$

output `arcsin(1/5*(2+x)*5^(1/2))`

3.55.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 23 vs. 2(10) = 20.

Time = 0.09 (sec) , antiderivative size = 23, normalized size of antiderivative = 2.30

$$\int \frac{1}{\sqrt{1-4x-x^2}} dx = 2 \arctan\left(\frac{x}{-1 + \sqrt{1-4x-x^2}}\right)$$

input `Integrate[1/Sqrt[1 - 4*x - x^2],x]`

output `2*ArcTan[x/(-1 + Sqrt[1 - 4*x - x^2])]`

3.55.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.70, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {1090, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{-x^2 - 4x + 1}} dx$$

↓ 1090

$$\frac{\int \frac{1}{\sqrt{1 - \frac{1}{20}(-2x-4)^2}} d(-2x-4)}{2\sqrt{5}}$$

↓ 223

$$-\arcsin\left(\frac{-2x-4}{2\sqrt{5}}\right)$$

input `Int[1/Sqrt[1 - 4*x - x^2],x]`

output `-ArcSin[(-4 - 2*x)/(2*Sqrt[5])]`

3.55.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 1090 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

3.55.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

method	result	size
default	$\arcsin\left(\frac{(2+x)\sqrt{5}}{5}\right)$	10
trager	$\text{RootOf}(-Z^2 + 1) \ln(-\text{RootOf}(-Z^2 + 1)x + \sqrt{-x^2 - 4x + 1}) - 2\text{RootOf}(-Z^2 + 1)$	39

input `int(1/(-x^2-4*x+1)^(1/2),x,method=_RETURNVERBOSE)`

output `arcsin(1/5*(2+x)*5^(1/2))`

3.55.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. 2(9) = 18.

Time = 0.24 (sec) , antiderivative size = 21, normalized size of antiderivative = 2.10

$$\int \frac{1}{\sqrt{1-4x-x^2}} dx = -2 \arctan\left(\frac{\sqrt{-x^2-4x+1}-1}{x}\right)$$

input `integrate(1/(-x^2-4*x+1)^(1/2),x, algorithm="fricas")`

output `-2*arctan((sqrt(-x^2 - 4*x + 1) - 1)/x)`

3.55.6 Sympy [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1-4x-x^2}} dx = \text{asin}\left(\frac{\sqrt{5}(x+2)}{5}\right)$$

input `integrate(1/(-x**2-4*x+1)**(1/2),x)`

output `asin(sqrt(5)*(x + 2)/5)`

3.55.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int \frac{1}{\sqrt{1-4x-x^2}} dx = -\arcsin\left(-\frac{1}{5}\sqrt{5}(x+2)\right)$$

input `integrate(1/(-x^2-4*x+1)^(1/2),x, algorithm="maxima")`

output `-arcsin(-1/5*sqrt(5)*(x + 2))`

3.55.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. 2(9) = 18.

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 2.90

$$\int \frac{1}{\sqrt{1-4x-x^2}} dx = \frac{1}{2}\sqrt{-x^2-4x+1}(x+2) + \frac{5}{2}\arcsin\left(\frac{1}{5}\sqrt{5}(x+2)\right)$$

input `integrate(1/(-x^2-4*x+1)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(-x^2 - 4*x + 1)*(x + 2) + 5/2*arcsin(1/5*sqrt(5)*(x + 2))`

3.55.9 Mupad [B] (verification not implemented)

Time = 14.49 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int \frac{1}{\sqrt{1-4x-x^2}} dx = \operatorname{asin}\left(\frac{\sqrt{20}(2x+4)}{20}\right)$$

input `int(1/(1 - x^2 - 4*x)^(1/2),x)`

output `asin((20^(1/2)*(2*x + 4))/20)`

3.56 $\int \log\left(\frac{1}{x}\right) dx$

3.56.1	Optimal result	388
3.56.2	Mathematica [A] (verified)	388
3.56.3	Rubi [A] (verified)	389
3.56.4	Maple [A] (verified)	389
3.56.5	Fricas [A] (verification not implemented)	390
3.56.6	Sympy [A] (verification not implemented)	390
3.56.7	Maxima [A] (verification not implemented)	390
3.56.8	Giac [A] (verification not implemented)	391
3.56.9	Mupad [B] (verification not implemented)	391

3.56.1 Optimal result

Integrand size = 4, antiderivative size = 8

$$\int \log\left(\frac{1}{x}\right) dx = x + x \log\left(\frac{1}{x}\right)$$

output `x+x*ln(1/x)`

3.56.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \log\left(\frac{1}{x}\right) dx = x + x \log\left(\frac{1}{x}\right)$$

input `Integrate[Log[x^(-1)],x]`

output `x + x*Log[x^(-1)]`

3.56.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2732}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \log\left(\frac{1}{x}\right) dx$$

↓ 2732

$$x + x \log\left(\frac{1}{x}\right)$$

input `Int [Log[x^(-1)], x]`

output `x + x*Log[x^(-1)]`

3.56.3.1 Defintions of rubi rules used

rule 2732 `Int [Log[(c_.)*(x_)^(n_.)], x_Symbol] :> Simp[x*Log[c*x^n], x] - Simp[n*x, x] /; FreeQ[{c, n}, x]`

3.56.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
derivativedivides	$x + x \ln\left(\frac{1}{x}\right)$	9
default	$x + x \ln\left(\frac{1}{x}\right)$	9
norman	$x + x \ln\left(\frac{1}{x}\right)$	9
risch	$x + x \ln\left(\frac{1}{x}\right)$	9
parallelrisch	$x + x \ln\left(\frac{1}{x}\right)$	9
parts	$x + x \ln\left(\frac{1}{x}\right)$	9

input `int(ln(1/x), x, method=_RETURNVERBOSE)`

output `x+x*ln(1/x)`

3.56.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \log\left(\frac{1}{x}\right) dx = x \log\left(\frac{1}{x}\right) + x$$

input `integrate(log(1/x),x, algorithm="fricas")`

output `x*log(1/x) + x`

3.56.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \log\left(\frac{1}{x}\right) dx = x \log\left(\frac{1}{x}\right) + x$$

input `integrate(ln(1/x),x)`

output `x*log(1/x) + x`

3.56.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \log\left(\frac{1}{x}\right) dx = -x \log(x) + x$$

input `integrate(log(1/x),x, algorithm="maxima")`

output `-x*log(x) + x`

3.56.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \log\left(\frac{1}{x}\right) dx = -x \log(x) + x$$

input `integrate(log(1/x),x, algorithm="giac")`

output `-x*log(x) + x`

3.56.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \log\left(\frac{1}{x}\right) dx = x \left(\ln\left(\frac{1}{x}\right) + 1 \right)$$

input `int(log(1/x),x)`

output `x*(log(1/x) + 1)`

3.57 $\int \frac{1}{1+\sin(x)} dx$

3.57.1	Optimal result	392
3.57.2	Mathematica [B] (verified)	392
3.57.3	Rubi [A] (verified)	393
3.57.4	Maple [A] (verified)	394
3.57.5	Fricas [A] (verification not implemented)	394
3.57.6	Sympy [A] (verification not implemented)	394
3.57.7	Maxima [A] (verification not implemented)	395
3.57.8	Giac [A] (verification not implemented)	395
3.57.9	Mupad [B] (verification not implemented)	395

3.57.1 Optimal result

Integrand size = 6, antiderivative size = 10

$$\int \frac{1}{1 + \sin(x)} dx = -\frac{\cos(x)}{1 + \sin(x)}$$

output `-cos(x)/(1+sin(x))`

3.57.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 23 vs. 2(10) = 20.

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 2.30

$$\int \frac{1}{1 + \sin(x)} dx = \frac{2 \sin\left(\frac{x}{2}\right)}{\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)}$$

input `Integrate[(1 + Sin[x])^(-1),x]`

output `(2*Sin[x/2])/(Cos[x/2] + Sin[x/2])`

3.57.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sin(x) + 1} dx$$

↓ 3042

$$\int \frac{1}{\sin(x) + 1} dx$$

↓ 3127

$$-\frac{\cos(x)}{\sin(x) + 1}$$

input `Int[(1 + Sin[x])^(-1),x]`

output `-(Cos[x]/(1 + Sin[x]))`

3.57.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

3.57.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

method	result	size
default	$-\frac{2}{1+\tan(\frac{x}{2})}$	11
norman	$-\frac{2}{1+\tan(\frac{x}{2})}$	11
parallelrisch	$-\frac{2}{1+\tan(\frac{x}{2})}$	11
risch	$-\frac{2}{i+e^{ix}}$	13

input `int(1/(1+sin(x)),x,method=_RETURNVERBOSE)`

output `-2/(1+tan(1/2*x))`

3.57.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.80

$$\int \frac{1}{1 + \sin(x)} dx = -\frac{\cos(x) - \sin(x) + 1}{\cos(x) + \sin(x) + 1}$$

input `integrate(1/(1+sin(x)),x, algorithm="fricas")`

output `-(cos(x) - sin(x) + 1)/(cos(x) + sin(x) + 1)`

3.57.6 Sympy [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{1}{1 + \sin(x)} dx = -\frac{2}{\tan(\frac{x}{2}) + 1}$$

input `integrate(1/(1+sin(x)),x)`

output `-2/(tan(x/2) + 1)`

3.57.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.50

$$\int \frac{1}{1 + \sin(x)} dx = -\frac{2}{\frac{\sin(x)}{\cos(x)+1} + 1}$$

input `integrate(1/(1+sin(x)),x, algorithm="maxima")`output `-2/(sin(x)/(cos(x) + 1) + 1)`**3.57.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \sin(x)} dx = -\frac{2}{\tan\left(\frac{1}{2}x\right) + 1}$$

input `integrate(1/(1+sin(x)),x, algorithm="giac")`output `-2/(tan(1/2*x) + 1)`**3.57.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \sin(x)} dx = -\frac{2}{\tan\left(\frac{x}{2}\right) + 1}$$

input `int(1/(sin(x) + 1),x)`output `-2/(tan(x/2) + 1)`

3.58 $\int \frac{\sqrt{x}}{\sqrt{2012-x+\sqrt{x}}} dx$

3.58.1	Optimal result	396
3.58.2	Mathematica [A] (verified)	396
3.58.3	Rubi [A] (verified)	397
3.58.4	Maple [A] (verified)	399
3.58.5	Fricas [A] (verification not implemented)	400
3.58.6	Sympy [F]	400
3.58.7	Maxima [F]	400
3.58.8	Giac [B] (verification not implemented)	401
3.58.9	Mupad [B] (verification not implemented)	401

3.58.1 Optimal result

Integrand size = 23, antiderivative size = 53

$$\int \frac{\sqrt{x}}{\sqrt{2012-x+\sqrt{x}}} dx = -\frac{1}{2}\sqrt{2012-x}\sqrt{x} + \frac{x}{2} + 503\operatorname{arctanh}\left(\frac{\sqrt{2012-x}\sqrt{x}}{1006}\right) + 503\log(1006-x)$$

output `-1/2*(2012-x)^(1/2)*x^(1/2)+1/2*x+503*arctanh(1/1006*(2012-x)^(1/2)*x^(1/2))+503*ln(1006-x)`

3.58.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.32

$$\int \frac{\sqrt{x}}{\sqrt{2012-x+\sqrt{x}}} dx = \frac{1}{2}\left(x - \sqrt{-((-2012+x)x)} + 2012\log\left(2\sqrt{503} - \sqrt{x}\right) + 2012\log\left(\sqrt{2012-x} + \sqrt{x}\right) - 2012\log\left(-1006 + \sqrt{503}\sqrt{x}\right)\right)$$

input `Integrate[Sqrt[x]/(Sqrt[2012 - x] + Sqrt[x]),x]`

output `(x - Sqrt[-((-2012 + x)*x)] + 2012*Log[2*Sqrt[503] - Sqrt[x]] + 2012*Log[Sqrt[2012 - x] + Sqrt[x]] - 2012*Log[-1006 + Sqrt[503]*Sqrt[x]])/2`

3.58.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.09, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.435$, Rules used = {2035, 2532, 27, 243, 49, 380, 27, 291, 219, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{x}}{\sqrt{2012-x} + \sqrt{x}} dx \\
 & \quad \downarrow \text{2035} \\
 & 2 \int \frac{x}{\sqrt{2012-x} + \sqrt{x}} d\sqrt{x} \\
 & \quad \downarrow \text{2532} \\
 & 2 \left(\int \frac{\sqrt{2012-xx}}{2(1006-x)} d\sqrt{x} - \int \frac{x^{3/2}}{2(1006-x)} d\sqrt{x} \right) \\
 & \quad \downarrow \text{27} \\
 & 2 \left(\frac{1}{2} \int \frac{\sqrt{2012-xx}}{1006-x} d\sqrt{x} - \frac{1}{2} \int \frac{x^{3/2}}{1006-x} d\sqrt{x} \right) \\
 & \quad \downarrow \text{243} \\
 & 2 \left(\frac{1}{2} \int \frac{\sqrt{2012-xx}}{1006-x} d\sqrt{x} - \frac{1}{4} \int \frac{x}{1006-x} dx \right) \\
 & \quad \downarrow \text{49} \\
 & 2 \left(\frac{1}{2} \int \frac{\sqrt{2012-xx}}{1006-x} d\sqrt{x} - \frac{1}{4} \int \left(-1 - \frac{1006}{x-1006} \right) dx \right) \\
 & \quad \downarrow \text{380} \\
 & 2 \left(\frac{1}{2} \left(\frac{1}{2} \int \frac{2024072}{(1006-x)\sqrt{2012-x}} d\sqrt{x} - \frac{1}{2} \sqrt{2012-x}\sqrt{x} \right) - \frac{1}{4} \int \left(-1 - \frac{1006}{x-1006} \right) dx \right) \\
 & \quad \downarrow \text{27} \\
 & 2 \left(\frac{1}{2} \left(1012036 \int \frac{1}{(1006-x)\sqrt{2012-x}} d\sqrt{x} - \frac{1}{2} \sqrt{2012-x}\sqrt{x} \right) - \frac{1}{4} \int \left(-1 - \frac{1006}{x-1006} \right) dx \right) \\
 & \quad \downarrow \text{291} \\
 & 2 \left(\frac{1}{2} \left(1012036 \int \frac{1}{1006-1006x} d\frac{\sqrt{x}}{\sqrt{2012-x}} - \frac{1}{2} \sqrt{2012-x}\sqrt{x} \right) - \frac{1}{4} \int \left(-1 - \frac{1006}{x-1006} \right) dx \right)
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow \text{219} \\
 2\left(\frac{1}{2}\left(1006\operatorname{arctanh}\left(\frac{\sqrt{x}}{\sqrt{2012-x}}\right) - \frac{1}{2}\sqrt{2012-x}\sqrt{x}\right) - \frac{1}{4}\int\left(-1 - \frac{1006}{x-1006}\right)dx\right) \\
 \downarrow \text{2009} \\
 2\left(\frac{1}{2}\left(1006\operatorname{arctanh}\left(\frac{\sqrt{x}}{\sqrt{2012-x}}\right) - \frac{1}{2}\sqrt{2012-x}\sqrt{x}\right) + \frac{1}{4}(x + 1006\log(1006 - x))\right)
 \end{array}$$

input `Int[Sqrt[x]/(Sqrt[2012 - x] + Sqrt[x]),x]`

output `2*((-1/2*(Sqrt[2012 - x]*Sqrt[x]) + 1006*ArcTanh[Sqrt[x]/Sqrt[2012 - x]])/2 + (x + 1006*Log[1006 - x])/4)`

3.58.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

```
rule 380 Int[((e._)*(x._))^(m._)*((a._) + (b._)*(x._)^2)^(p._)*((c._) + (d._)*(x._)^2)^(q._), x_Symbol] := Simp[e*(e*x)^(m - 1)*(a + b*x^2)^(p + 1)*((c + d*x^2)^q/(b*(m + 2*(p + q) + 1))), x] - Simp[e^2/(b*(m + 2*(p + q) + 1)) Int[(e*x)^(m - 2)*(a + b*x^2)^p*(c + d*x^2)^(q - 1)*Simp[a*c*(m - 1) + (a*d*(m - 1) - 2*q*(b*c - a*d))*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 0] && GtQ[m, 1] && IntBinomialQ[a, b, c, d, e, m, 2, p, q, x]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2035 Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x]] /; FractionQ[m] && AlgebraicFunctionQ[Fx, x]
```

```
rule 2532 Int[(x_)^(m._)/((d._)*(x_)^(n._) + (c._)*Sqrt[(a._) + (b._)*(x_)^(p._)]), x_Symbol] := Simp[-d Int[x^(m + n)/(a*c^2 + (b*c^2 - d^2)*x^(2*n)), x], x] + Simp[c Int[(x^m*Sqrt[a + b*x^(2*n)])/(a*c^2 + (b*c^2 - d^2)*x^(2*n)), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[p, 2*n] && NeQ[b*c^2 - d^2, 0]
```

3.58.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{x}{2} + 503 \ln(x - 1006) - \frac{\sqrt{x} \sqrt{2012-x} \left(\sqrt{-x(-2012+x)} - 1006 \operatorname{arctanh}\left(\frac{1006}{\sqrt{-x(-2012+x)}}\right) \right)}{2\sqrt{-x(-2012+x)}}$	53

```
input int(x^(1/2)/((2012-x)^(1/2)+x^(1/2)),x,method=_RETURNVERBOSE)
```

```
output 1/2*x+503*ln(x-1006)-1/2*x^(1/2)*(2012-x)^(1/2)*((-x*(-2012+x))^(1/2)-1006*arctanh(1006/(-x*(-2012+x))^(1/2)))/(-x*(-2012+x))^(1/2)
```

3.58. $\int \frac{\sqrt{x}}{\sqrt{2012-x}+\sqrt{x}} dx$

3.58.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.21

$$\int \frac{\sqrt{x}}{\sqrt{2012-x} + \sqrt{x}} dx = \frac{1}{2}x - \frac{1}{2}\sqrt{x}\sqrt{-x+2012} + 503 \log(x-1006) \\ + 503 \log\left(\frac{x + \sqrt{x}\sqrt{-x+2012}}{x}\right) \\ - 503 \log\left(-\frac{x - \sqrt{x}\sqrt{-x+2012}}{x}\right)$$

input `integrate(x^(1/2)/((2012-x)^(1/2)+x^(1/2)),x, algorithm="fricas")`output `1/2*x - 1/2*sqrt(x)*sqrt(-x + 2012) + 503*log(x - 1006) + 503*log((x + sqrt(x)*sqrt(-x + 2012))/x) - 503*log(-(x - sqrt(x)*sqrt(-x + 2012))/x)`**3.58.6 Sympy [F]**

$$\int \frac{\sqrt{x}}{\sqrt{2012-x} + \sqrt{x}} dx = \int \frac{\sqrt{x}}{\sqrt{x} + \sqrt{2012-x}} dx$$

input `integrate(x**(1/2)/((2012-x)**(1/2)+x**(1/2)),x)`output `Integral(sqrt(x)/(sqrt(x) + sqrt(2012 - x)), x)`**3.58.7 Maxima [F]**

$$\int \frac{\sqrt{x}}{\sqrt{2012-x} + \sqrt{x}} dx = \int \frac{\sqrt{x}}{\sqrt{x} + \sqrt{-x+2012}} dx$$

input `integrate(x^(1/2)/((2012-x)^(1/2)+x^(1/2)),x, algorithm="maxima")`output `integrate(sqrt(x)/(sqrt(x) + sqrt(-x + 2012)), x)`

3.58.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 117 vs. 2(39) = 78.

Time = 0.31 (sec) , antiderivative size = 117, normalized size of antiderivative = 2.21

$$\begin{aligned} & \int \frac{\sqrt{x}}{\sqrt{2012-x} + \sqrt{x}} dx \\ &= \frac{1}{2}x - \frac{1}{2}\sqrt{x}\sqrt{-x+2012} + 503 \log(|x-1006|) \\ &+ 503 \log\left(\left| -\frac{2\sqrt{503} - \sqrt{-x+2012}}{\sqrt{x}} + \frac{\sqrt{x}}{2\sqrt{503} - \sqrt{-x+2012}} + 2 \right|\right) \\ &- 503 \log\left(\left| -\frac{2\sqrt{503} - \sqrt{-x+2012}}{\sqrt{x}} + \frac{\sqrt{x}}{2\sqrt{503} - \sqrt{-x+2012}} - 2 \right|\right) \end{aligned}$$

input `integrate(x^(1/2)/((2012-x)^(1/2)+x^(1/2)),x, algorithm="giac")`

output `1/2*x - 1/2*sqrt(x)*sqrt(-x + 2012) + 503*log(abs(x - 1006)) + 503*log(abs(-2*sqrt(503) - sqrt(-x + 2012))/sqrt(x) + sqrt(x)/(2*sqrt(503) - sqrt(-x + 2012)) + 2)) - 503*log(abs(-2*sqrt(503) - sqrt(-x + 2012))/sqrt(x) + sqrt(x)/(2*sqrt(503) - sqrt(-x + 2012)) - 2))`

3.58.9 Mupad [B] (verification not implemented)

Time = 16.55 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.21

$$\begin{aligned} \int \frac{\sqrt{x}}{\sqrt{2012-x} + \sqrt{x}} dx &= \frac{x}{2} + 1006 \operatorname{atanh}\left(\frac{2\sqrt{503}\sqrt{x} - \sqrt{x}\sqrt{2012-x}}{x + 2\sqrt{503}\sqrt{2012-x} - 2012}\right) \\ &+ 503 \ln(x - 1006) - \frac{\sqrt{x}\sqrt{2012-x}}{2} \end{aligned}$$

input `int(x^(1/2)/((2012 - x)^(1/2) + x^(1/2)),x)`

output `x/2 + 1006*atanh((2*503^(1/2)*x^(1/2) - x^(1/2)*(2012 - x)^(1/2))/(x + 2*503^(1/2)*(2012 - x)^(1/2) - 2012)) + 503*log(x - 1006) - (x^(1/2)*(2012 - x)^(1/2))/2`

3.59 $\int \frac{-1+x}{(1+x)\sqrt{x+x^2+x^3}} dx$

3.59.1	Optimal result	402
3.59.2	Mathematica [A] (verified)	402
3.59.3	Rubi [A] (verified)	403
3.59.4	Maple [A] (verified)	404
3.59.5	Fricas [A] (verification not implemented)	405
3.59.6	Sympy [F]	405
3.59.7	Maxima [F]	405
3.59.8	Giac [F]	406
3.59.9	Mupad [B] (verification not implemented)	406

3.59.1 Optimal result

Integrand size = 21, antiderivative size = 46

$$\int \frac{-1+x}{(1+x)\sqrt{x+x^2+x^3}} dx = -\frac{2\sqrt{x}\sqrt{1+x+x^2} \arctan\left(\frac{\sqrt{x}}{\sqrt{1+x+x^2}}\right)}{\sqrt{x+x^2+x^3}}$$

output `-2*x^(1/2)*(x^2+x+1)^(1/2)*arctan(x^(1/2)/(x^2+x+1)^(1/2))/(x^3+x^2+x)^(1/2)`

3.59.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{-1+x}{(1+x)\sqrt{x+x^2+x^3}} dx = -\frac{2\sqrt{x}\sqrt{1+x+x^2} \arctan\left(\frac{\sqrt{x}}{\sqrt{1+x+x^2}}\right)}{\sqrt{x}(1+x+x^2)}$$

input `Integrate[(-1 + x)/((1 + x)*Sqrt[x + x^2 + x^3]),x]`

output `(-2*Sqrt[x]*Sqrt[1 + x + x^2]*ArcTan[Sqrt[x]/Sqrt[1 + x + x^2]])/Sqrt[x*(1 + x + x^2)]`

3.59.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {2467, 25, 2035, 2212, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x-1}{(x+1)\sqrt{x^3+x^2+x}} dx \\
 & \quad \downarrow \text{2467} \\
 & \frac{\sqrt{x}\sqrt{x^2+x+1} \int -\frac{1-x}{\sqrt{x(x+1)}\sqrt{x^2+x+1}} dx}{\sqrt{x^3+x^2+x}} \\
 & \quad \downarrow \text{25} \\
 & -\frac{\sqrt{x}\sqrt{x^2+x+1} \int \frac{1-x}{\sqrt{x(x+1)}\sqrt{x^2+x+1}} dx}{\sqrt{x^3+x^2+x}} \\
 & \quad \downarrow \text{2035} \\
 & -\frac{2\sqrt{x}\sqrt{x^2+x+1} \int \frac{1-x}{(x+1)\sqrt{x^2+x+1}} d\sqrt{x}}{\sqrt{x^3+x^2+x}} \\
 & \quad \downarrow \text{2212} \\
 & -\frac{2\sqrt{x}\sqrt{x^2+x+1} \int \frac{1}{x+1} d\frac{\sqrt{x}}{\sqrt{x^2+x+1}}}{\sqrt{x^3+x^2+x}} \\
 & \quad \downarrow \text{216} \\
 & -\frac{2\sqrt{x}\sqrt{x^2+x+1} \arctan\left(\frac{\sqrt{x}}{\sqrt{x^2+x+1}}\right)}{\sqrt{x^3+x^2+x}}
 \end{aligned}$$

input `Int[(-1 + x)/((1 + x)*Sqrt[x + x^2 + x^3]),x]`

output `(-2*Sqrt[x]*Sqrt[1 + x + x^2]*ArcTan[Sqrt[x]/Sqrt[1 + x + x^2]])/Sqrt[x + x^2 + x^3]`

3.59.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 2035 `Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[Fx, x]`
- rule 2212 `Int[((A_) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4]), x_Symbol] := Simp[A Subst[Int[1/(d - (b*d - 2*a*e)*x^2), x], x, x/Sqrt[a + b*x^2 + c*x^4]], x] /; FreeQ[{a, b, c, d, e, A, B}, x] && EqQ[c*d^2 - a*e^2, 0] && EqQ[B*d + A*e, 0]`
- rule 2467 `Int[(Fx_.)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Simp[Px^FracPart[p]/(x^(r*FracPart[p])*ExpandToSum[Px/x^r, x]^FracPart[p]) Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x], x] /; IGtQ[r, 0] /; FreeQ[p, x] && PolyQ[Px, x] && !IntegerQ[p] && !MonomialQ[Px, x] && !PolyQ[Fx, x]`

3.59.4 Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.39

method	result
default	$2 \arctan\left(\frac{\sqrt{(x^2+x+1)x}}{x}\right)$
pseudoelliptic	$2 \arctan\left(\frac{\sqrt{(x^2+x+1)x}}{x}\right)$
trager	$-\text{RootOf}(_Z^2 + 1) \ln\left(\frac{\text{RootOf}(_Z^2 + 1)x^2 + \text{RootOf}(_Z^2 + 1) + 2\sqrt{x^3+x^2+x}}{(1+x)^2}\right)$
elliptic	$\frac{2\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)\sqrt{\frac{x+\frac{1}{2}+\frac{i\sqrt{3}}{2}}{\frac{1}{2}+\frac{i\sqrt{3}}{2}}}\sqrt{3}\sqrt{i\left(x+\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)}\sqrt{3}\sqrt{\frac{x}{-\frac{1}{2}-\frac{i\sqrt{3}}{2}}}\text{EllipticF}\left(\sqrt{\frac{x+\frac{1}{2}+\frac{i\sqrt{3}}{2}}{\frac{1}{2}+\frac{i\sqrt{3}}{2}}}, \frac{\sqrt{3}\sqrt{i\left(-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)}\sqrt{3}}{3}\right)}{3\sqrt{x^3+x^2+x}} - 4\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)$

3.59. $\int \frac{-1+x}{(1+x)\sqrt{x+x^2+x^3}} dx$

input `int((-1+x)/(1+x)/(x^3+x^2+x)^(1/2),x,method=_RETURNVERBOSE)`

output `2*arctan(((x^2+x+1)*x)^(1/2)/x)`

3.59.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.39

$$\int \frac{-1+x}{(1+x)\sqrt{x+x^2+x^3}} dx = \arctan\left(\frac{x^2+1}{2\sqrt{x^3+x^2+x}}\right)$$

input `integrate((-1+x)/(1+x)/(x^3+x^2+x)^(1/2),x, algorithm="fricas")`

output `arctan(1/2*(x^2 + 1)/sqrt(x^3 + x^2 + x))`

3.59.6 Sympy [F]

$$\int \frac{-1+x}{(1+x)\sqrt{x+x^2+x^3}} dx = \int \frac{x-1}{\sqrt{x(x^2+x+1)}(x+1)} dx$$

input `integrate((-1+x)/(1+x)/(x**3+x**2+x)**(1/2),x)`

output `Integral((x - 1)/(sqrt(x*(x**2 + x + 1))*(x + 1)), x)`

3.59.7 Maxima [F]

$$\int \frac{-1+x}{(1+x)\sqrt{x+x^2+x^3}} dx = \int \frac{x-1}{\sqrt{x^3+x^2+x}(x+1)} dx$$

input `integrate((-1+x)/(1+x)/(x^3+x^2+x)^(1/2),x, algorithm="maxima")`

output `integrate((x - 1)/(sqrt(x^3 + x^2 + x)*(x + 1)), x)`

3.59.8 Giac [F]

$$\int \frac{-1+x}{(1+x)\sqrt{x+x^2+x^3}} dx = \int \frac{x-1}{\sqrt{x^3+x^2+x}(x+1)} dx$$

input `integrate((-1+x)/(1+x)/(x^3+x^2+x)^(1/2),x, algorithm="giac")`

output `integrate((x - 1)/(sqrt(x^3 + x^2 + x)*(x + 1)), x)`

3.59.9 Mupad [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 179, normalized size of antiderivative = 3.89

$$\int \frac{-1+x}{(1+x)\sqrt{x+x^2+x^3}} dx$$

$$= \frac{\sqrt{\frac{x}{-\frac{1}{2}+\frac{\sqrt{3}1i}{2}}} \sqrt{-\frac{x+\frac{1}{2}-\frac{\sqrt{3}1i}{2}}{-\frac{1}{2}+\frac{\sqrt{3}1i}{2}}} \sqrt{\frac{x+\frac{1}{2}+\frac{\sqrt{3}1i}{2}}{\frac{1}{2}+\frac{\sqrt{3}1i}{2}}} (\sqrt{3}+1i) \left(F\left(\operatorname{asin}\left(\sqrt{\frac{x}{-\frac{1}{2}+\frac{\sqrt{3}1i}{2}}}\right) \middle| -\frac{-\frac{1}{2}+\frac{\sqrt{3}1i}{2}}{\frac{1}{2}+\frac{\sqrt{3}1i}{2}}\right) - 2\Pi\left(\frac{1}{2}-\frac{\sqrt{3}1i}{2}; \operatorname{asin}\left(\sqrt{\frac{x}{-\frac{1}{2}+\frac{\sqrt{3}1i}{2}}}\right)\right) \right)}{\sqrt{x^3+x^2-\left(-\frac{1}{2}+\frac{\sqrt{3}1i}{2}\right)\left(\frac{1}{2}+\frac{\sqrt{3}1i}{2}\right)} x}$$

input `int((x - 1)/((x + 1)*(x + x^2 + x^3)^(1/2)),x)`

output `((x/((3^(1/2)*1i)/2 - 1/2))^(1/2)*(-(x - (3^(1/2)*1i)/2 + 1/2)/((3^(1/2)*1i)/2 - 1/2))^(1/2)*((x + (3^(1/2)*1i)/2 + 1/2)/((3^(1/2)*1i)/2 + 1/2))^(1/2)*(3^(1/2) + 1i)*(ellipticF(asin((x/((3^(1/2)*1i)/2 - 1/2))^(1/2)), -((3^(1/2)*1i)/2 - 1/2)/((3^(1/2)*1i)/2 + 1/2)) - 2*ellipticPi(1/2 - (3^(1/2)*1i)/2, asin((x/((3^(1/2)*1i)/2 - 1/2))^(1/2)), -((3^(1/2)*1i)/2 - 1/2)/((3^(1/2)*1i)/2 + 1/2))*1i)/(x^2 + x^3 - x*((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2))^(1/2)`

$$3.60 \quad \int \frac{1+4x+6x^2+4x^3+x^4}{-1+3x-3x^2+x^3} dx$$

3.60.1	Optimal result	407
3.60.2	Mathematica [A] (verified)	407
3.60.3	Rubi [A] (verified)	408
3.60.4	Maple [A] (verified)	409
3.60.5	Fricas [A] (verification not implemented)	410
3.60.6	Sympy [A] (verification not implemented)	410
3.60.7	Maxima [A] (verification not implemented)	410
3.60.8	Giac [A] (verification not implemented)	411
3.60.9	Mupad [B] (verification not implemented)	411

3.60.1 Optimal result

Integrand size = 34, antiderivative size = 37

$$\int \frac{1+4x+6x^2+4x^3+x^4}{-1+3x-3x^2+x^3} dx = -\frac{8}{(1-x)^2} + \frac{32}{1-x} + 7x + \frac{x^2}{2} + 24 \log(1-x)$$

output `-8/(1-x)^2+32/(1-x)+7*x+1/2*x^2+24*ln(1-x)`

3.60.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.95

$$\int \frac{1+4x+6x^2+4x^3+x^4}{-1+3x-3x^2+x^3} dx = -\frac{8}{(-1+x)^2} - \frac{32}{-1+x} + 8(-1+x) + \frac{1}{2}(-1+x)^2 + 24 \log(-1+x)$$

input `Integrate[(1 + 4*x + 6*x^2 + 4*x^3 + x^4)/(-1 + 3*x - 3*x^2 + x^3),x]`

output `-8/(-1 + x)^2 - 32/(-1 + x) + 8*(-1 + x) + (-1 + x)^2/2 + 24*Log[-1 + x]`

3.60.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2006, 2007, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4 + 4x^3 + 6x^2 + 4x + 1}{x^3 - 3x^2 + 3x - 1} dx \\
 & \quad \downarrow \text{2006} \\
 & \int \frac{(x+1)^4}{x^3 - 3x^2 + 3x - 1} dx \\
 & \quad \downarrow \text{2007} \\
 & \int \frac{(x+1)^4}{(x-1)^3} dx \\
 & \quad \downarrow \text{49} \\
 & \int \left(x + \frac{24}{x-1} + \frac{32}{(x-1)^2} + \frac{16}{(x-1)^3} + 7 \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{x^2}{2} + 7x + \frac{32}{1-x} - \frac{8}{(1-x)^2} + 24 \log(1-x)
 \end{aligned}$$

input `Int[(1 + 4*x + 6*x^2 + 4*x^3 + x^4)/(-1 + 3*x - 3*x^2 + x^3),x]`

output `-8/(1 - x)^2 + 32/(1 - x) + 7*x + x^2/2 + 24*Log[1 - x]`

3.60.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

```
rule 2006 Int[(u_.)*(Px_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px, x]],
b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Int[u*(a + b*x)^Expon[Px,
x], x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; PolyQ[Px, x] && GtQ[Expon[P
x, x], 1] && NeQ[Coeff[Px, x, 0], 0] && !MatchQ[Px, (a_.)*(v_)^Expon[Px, x
] /; FreeQ[a, x] && LinearQ[v, x]]
```

```
rule 2007 Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px,
x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Int[u*(a + b*x)^(Ex
pon[Px, x]*p), x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; IntegerQ[p] && Pol
yQ[Px, x] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

3.60.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.78

method	result	size
norman	$\frac{-52x+6x^3+\frac{1}{2}x^4+\frac{75}{2}}{(-1+x)^2} + 24 \ln(-1+x)$	29
default	$\frac{x^2}{2} + 7x - \frac{32}{-1+x} + 24 \ln(-1+x) - \frac{8}{(-1+x)^2}$	30
risch	$\frac{x^2}{2} + 7x + \frac{-32x+24}{x^2-2x+1} + 24 \ln(-1+x)$	32
parallelrisc	$\frac{x^4+48 \ln(-1+x)x^2+12x^3+75-96 \ln(-1+x)x+48 \ln(-1+x)-104x}{2x^2-4x+2}$	48

```
input int((x^4+4*x^3+6*x^2+4*x+1)/(x^3-3*x^2+3*x-1),x,method=_RETURNVERBOSE)
```

```
output (-52*x+6*x^3+1/2*x^4+75/2)/(-1+x)^2+24*ln(-1+x)
```

3.60.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.19

$$\int \frac{1 + 4x + 6x^2 + 4x^3 + x^4}{-1 + 3x - 3x^2 + x^3} dx$$

$$= \frac{x^4 + 12x^3 - 27x^2 + 48(x^2 - 2x + 1)\log(x - 1) - 50x + 48}{2(x^2 - 2x + 1)}$$

input `integrate((x^4+4*x^3+6*x^2+4*x+1)/(x^3-3*x^2+3*x-1),x, algorithm="fricas")`output `1/2*(x^4 + 12*x^3 - 27*x^2 + 48*(x^2 - 2*x + 1)*log(x - 1) - 50*x + 48)/(x^2 - 2*x + 1)`**3.60.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.73

$$\int \frac{1 + 4x + 6x^2 + 4x^3 + x^4}{-1 + 3x - 3x^2 + x^3} dx = \frac{x^2}{2} + 7x + \frac{24 - 32x}{x^2 - 2x + 1} + 24 \log(x - 1)$$

input `integrate((x**4+4*x**3+6*x**2+4*x+1)/(x**3-3*x**2+3*x-1),x)`output `x**2/2 + 7*x + (24 - 32*x)/(x**2 - 2*x + 1) + 24*log(x - 1)`**3.60.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.86

$$\int \frac{1 + 4x + 6x^2 + 4x^3 + x^4}{-1 + 3x - 3x^2 + x^3} dx = \frac{1}{2}x^2 + 7x - \frac{8(4x - 3)}{x^2 - 2x + 1} + 24 \log(x - 1)$$

input `integrate((x^4+4*x^3+6*x^2+4*x+1)/(x^3-3*x^2+3*x-1),x, algorithm="maxima")`output `1/2*x^2 + 7*x - 8*(4*x - 3)/(x^2 - 2*x + 1) + 24*log(x - 1)`

3.60.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.76

$$\int \frac{1 + 4x + 6x^2 + 4x^3 + x^4}{-1 + 3x - 3x^2 + x^3} dx = \frac{1}{2}x^2 + 7x - \frac{8(4x - 3)}{(x - 1)^2} + 24 \log(|x - 1|)$$

input `integrate((x^4+4*x^3+6*x^2+4*x+1)/(x^3-3*x^2+3*x-1),x, algorithm="giac")`output `1/2*x^2 + 7*x - 8*(4*x - 3)/(x - 1)^2 + 24*log(abs(x - 1))`**3.60.9 Mupad [B] (verification not implemented)**

Time = 13.81 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.86

$$\int \frac{1 + 4x + 6x^2 + 4x^3 + x^4}{-1 + 3x - 3x^2 + x^3} dx = 7x + 24 \ln(x - 1) - \frac{32x - 24}{x^2 - 2x + 1} + \frac{x^2}{2}$$

input `int((4*x + 6*x^2 + 4*x^3 + x^4 + 1)/(3*x - 3*x^2 + x^3 - 1),x)`output `7*x + 24*log(x - 1) - (32*x - 24)/(x^2 - 2*x + 1) + x^2/2`

$$\mathbf{3.61} \quad \int \left(\cos(x) \log(x) + \frac{\sin(x)}{x} \right) dx$$

3.61.1	Optimal result	412
3.61.2	Mathematica [A] (verified)	412
3.61.3	Rubi [A] (verified)	413
3.61.4	Maple [A] (verified)	413
3.61.5	Fricas [A] (verification not implemented)	414
3.61.6	Sympy [A] (verification not implemented)	414
3.61.7	Maxima [A] (verification not implemented)	414
3.61.8	Giac [A] (verification not implemented)	415
3.61.9	Mupad [B] (verification not implemented)	415

3.61.1 Optimal result

Integrand size = 12, antiderivative size = 5

$$\int \left(\cos(x) \log(x) + \frac{\sin(x)}{x} \right) dx = \log(x) \sin(x)$$

output `ln(x)*sin(x)`

3.61.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \left(\cos(x) \log(x) + \frac{\sin(x)}{x} \right) dx = \log(x) \sin(x)$$

input `Integrate[Cos[x]*Log[x] + Sin[x]/x,x]`

output `Log[x]*Sin[x]`

3.61. $\int \left(\cos(x) \log(x) + \frac{\sin(x)}{x} \right) dx$

3.61.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\frac{\sin(x)}{x} + \log(x) \cos(x) \right) dx$$

↓ 2009

$$\log(x) \sin(x)$$

input `Int[Cos[x]*Log[x] + Sin[x]/x,x]`

output `Log[x]*Sin[x]`

3.61.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.61.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
risch	$\ln(x) \sin(x)$	6
parallelrisch	$\ln(x) \sin(x)$	6
norman	$\frac{2 \ln(x) \tan(\frac{x}{2})}{1 + \tan(\frac{x}{2})^2}$	19

input `int(cos(x)*ln(x)+sin(x)/x,x,method=_RETURNVERBOSE)`

output `ln(x)*sin(x)`

3.61. $\int \left(\cos(x) \log(x) + \frac{\sin(x)}{x} \right) dx$

3.61.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \left(\cos(x) \log(x) + \frac{\sin(x)}{x} \right) dx = \log(x) \sin(x)$$

input `integrate(cos(x)*log(x)+sin(x)/x,x, algorithm="fricas")`output `log(x)*sin(x)`**3.61.6 Sympy [A] (verification not implemented)**

Time = 3.73 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \left(\cos(x) \log(x) + \frac{\sin(x)}{x} \right) dx = \log(x) \sin(x)$$

input `integrate(cos(x)*ln(x)+sin(x)/x,x)`output `log(x)*sin(x)`**3.61.7 Maxima [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \left(\cos(x) \log(x) + \frac{\sin(x)}{x} \right) dx = \log(x) \sin(x)$$

input `integrate(cos(x)*log(x)+sin(x)/x,x, algorithm="maxima")`output `log(x)*sin(x)`

3.61.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \left(\cos(x) \log(x) + \frac{\sin(x)}{x} \right) dx = \log(x) \sin(x)$$

input `integrate(cos(x)*log(x)+sin(x)/x,x, algorithm="giac")`

output `log(x)*sin(x)`

3.61.9 Mupad [B] (verification not implemented)

Time = 14.43 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \left(\cos(x) \log(x) + \frac{\sin(x)}{x} \right) dx = \ln(x) \sin(x)$$

input `int(cos(x)*log(x) + sin(x)/x,x)`

output `log(x)*sin(x)`

3.62 $\int \frac{1}{-x+x^3} dx$

3.62.1	Optimal result	416
3.62.2	Mathematica [A] (verified)	416
3.62.3	Rubi [A] (verified)	417
3.62.4	Maple [A] (verified)	418
3.62.5	Fricas [A] (verification not implemented)	419
3.62.6	Sympy [A] (verification not implemented)	419
3.62.7	Maxima [A] (verification not implemented)	419
3.62.8	Giac [A] (verification not implemented)	420
3.62.9	Mupad [B] (verification not implemented)	420

3.62.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \frac{1}{-x+x^3} dx = -\log(x) + \frac{1}{2} \log(1-x^2)$$

output `-ln(x)+1/2*ln(-x^2+1)`

3.62.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{1}{-x+x^3} dx = -\log(x) + \frac{1}{2} \log(1-x^2)$$

input `Integrate[(-x + x^3)^(-1),x]`

output `-Log[x] + Log[1 - x^2]/2`

3.62.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {2026, 243, 25, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^3 - x} dx \\
 & \quad \downarrow \text{2026} \\
 & \int \frac{1}{x(x^2 - 1)} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \int -\frac{1}{x^2(1 - x^2)} dx^2 \\
 & \quad \downarrow \text{25} \\
 & -\frac{1}{2} \int \frac{1}{x^2(1 - x^2)} dx^2 \\
 & \quad \downarrow \text{47} \\
 & \frac{1}{2} \left(-\int \frac{1}{x^2} dx^2 - \int \frac{1}{1 - x^2} dx^2 \right) \\
 & \quad \downarrow \text{14} \\
 & \frac{1}{2} \left(-\int \frac{1}{1 - x^2} dx^2 - \log(x^2) \right) \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{2} (\log(1 - x^2) - \log(x^2))
 \end{aligned}$$

input `Int[(-x + x^3)^(-1), x]`

output `(-Log[x^2] + Log[1 - x^2])/2`

3.62.3.1 Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2026 `Int[(Fx_.)*(Px_)^(p_.), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && IntegerQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])`

3.62.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
risch	$-\ln(x) + \frac{\ln(x^2-1)}{2}$	14
default	$\frac{\ln(1+x)}{2} - \ln(x) + \frac{\ln(-1+x)}{2}$	18
norman	$\frac{\ln(1+x)}{2} - \ln(x) + \frac{\ln(-1+x)}{2}$	18
parallelrisch	$\frac{\ln(1+x)}{2} - \ln(x) + \frac{\ln(-1+x)}{2}$	18
meijerg	$-\ln(x) - \frac{i\pi}{2} + \frac{\ln(-x^2+1)}{2}$	20

input `int(1/(x^3-x), x, method=_RETURNVERBOSE)`

output `-ln(x)+1/2*ln(x^2-1)`

3.62.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{1}{-x+x^3} dx = \frac{1}{2} \log(x^2-1) - \log(x)$$

input `integrate(1/(x^3-x),x, algorithm="fricas")`

output `1/2*log(x^2 - 1) - log(x)`

3.62.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.59

$$\int \frac{1}{-x+x^3} dx = -\log(x) + \frac{\log(x^2-1)}{2}$$

input `integrate(1/(x**3-x),x)`

output `-log(x) + log(x**2 - 1)/2`

3.62.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{1}{-x+x^3} dx = \frac{1}{2} \log(x+1) + \frac{1}{2} \log(x-1) - \log(x)$$

input `integrate(1/(x^3-x),x, algorithm="maxima")`

output `1/2*log(x + 1) + 1/2*log(x - 1) - log(x)`

3.62.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

$$\int \frac{1}{-x + x^3} dx = -\frac{1}{2} \log(x^2) + \frac{1}{2} \log(|x^2 - 1|)$$

input `integrate(1/(x^3-x),x, algorithm="giac")`output `-1/2*log(x^2) + 1/2*log(abs(x^2 - 1))`**3.62.9 Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{1}{-x + x^3} dx = \frac{\ln(x^2 - 1)}{2} - \ln(x)$$

input `int(-1/(x - x^3),x)`output `log(x^2 - 1)/2 - log(x)`

3.63 $\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx$

3.63.1	Optimal result	421
3.63.2	Mathematica [A] (verified)	421
3.63.3	Rubi [A] (verified)	422
3.63.4	Maple [A] (verified)	423
3.63.5	Fricas [A] (verification not implemented)	423
3.63.6	Sympy [A] (verification not implemented)	423
3.63.7	Maxima [A] (verification not implemented)	424
3.63.8	Giac [A] (verification not implemented)	424
3.63.9	Mupad [F(-1)]	424

3.63.1 Optimal result

Integrand size = 15, antiderivative size = 17

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx = x - \sqrt{1-x^2} \arcsin(x)$$

output `x-(-x^2+1)^(1/2)*arcsin(x)`

3.63.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx = x - \sqrt{1-x^2} \arcsin(x)$$

input `Integrate[(x*ArcSin[x])/Sqrt[1 - x^2],x]`

output `x - Sqrt[1 - x^2]*ArcSin[x]`

3.63.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {5182, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx$$

↓ 5182

$$\int 1 dx - \sqrt{1-x^2} \arcsin(x)$$

↓ 24

$$x - \sqrt{1-x^2} \arcsin(x)$$

input `Int[(x*ArcSin[x])/Sqrt[1 - x^2],x]`

output `x - Sqrt[1 - x^2]*ArcSin[x]`

3.63.3.1 Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 5182 `Int[((a_.) + ArcSin[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(d + e*x^2)^(p + 1)*((a + b*ArcSin[c*x])^n/(2*e*(p + 1))), x] + Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/(1 - c^2*x^2)^p] Int[(1 - c^2*x^2)^(p + 1/2)*(a + b*ArcSin[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

3.63.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

method	result	size
default	$x - \sqrt{-x^2 + 1} \arcsin(x)$	16

input `int(x*arcsin(x)/(-x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `x-(-x^2+1)^(1/2)*arcsin(x)`

3.63.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx = -\sqrt{-x^2 + 1} \arcsin(x) + x$$

input `integrate(x*arcsin(x)/(-x^2+1)^(1/2),x, algorithm="fricas")`

output `-sqrt(-x^2 + 1)*arcsin(x) + x`

3.63.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx = x - \sqrt{1-x^2} \arcsin(x)$$

input `integrate(x*asin(x)/(-x**2+1)**(1/2),x)`

output `x - sqrt(1 - x**2)*asin(x)`

3.63.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx = -\sqrt{-x^2+1} \arcsin(x) + x$$

input `integrate(x*arcsin(x)/(-x^2+1)^(1/2),x, algorithm="maxima")`output `-sqrt(-x^2 + 1)*arcsin(x) + x`**3.63.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx = -\sqrt{-x^2+1} \arcsin(x) + x$$

input `integrate(x*arcsin(x)/(-x^2+1)^(1/2),x, algorithm="giac")`output `-sqrt(-x^2 + 1)*arcsin(x) + x`**3.63.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx = \int \frac{x \operatorname{asin}(x)}{\sqrt{1-x^2}} dx$$

input `int((x*asin(x))/(1 - x^2)^(1/2),x)`output `int((x*asin(x))/(1 - x^2)^(1/2), x)`

3.64 $\int (1 - x)^{99} x dx$

3.64.1	Optimal result	425
3.64.2	Mathematica [B] (verified)	425
3.64.3	Rubi [A] (verified)	426
3.64.4	Maple [B] (verified)	427
3.64.5	Fricas [B] (verification not implemented)	428
3.64.6	Sympy [B] (verification not implemented)	429
3.64.7	Maxima [B] (verification not implemented)	430
3.64.8	Giac [B] (verification not implemented)	431
3.64.9	Mupad [B] (verification not implemented)	432

3.64.1 Optimal result

Integrand size = 9, antiderivative size = 23

$$\int (1 - x)^{99} x dx = -\frac{1}{100}(1 - x)^{100} + \frac{1}{101}(1 - x)^{101}$$

output `-1/100*(1-x)^100+1/101*(1-x)^101`

3.64.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 567 vs. $2(23) = 46$.

Time = 0.01 (sec) , antiderivative size = 567, normalized size of antiderivative = 24.65

$$\int (1 - x)^{99} x dx = \text{Too large to display}$$

input `Integrate[(1 - x)^99*x,x]`

output $x^2/2 - 33x^3 + (4851x^4)/4 - (156849x^5)/5 + 627396x^6 - 10217592x^7 + 140066157x^8 - 1654114616x^9 + (85600431378x^{10})/5 - 157366449604x^{11} + 1298273209233x^{12} - 9696194317908x^{13} + 66026466069564x^{14} - (2062057324941768x^{15})/5 + (4750096337812287x^{16})/2 - 12666923567499432x^{17} + 62806829355518017x^{18} - 290505891817783026x^{19} + (12572449429225165403x^{20})/10 - 5104603527655330314x^{21} + 19490304378320352108x^{22} - 70132813684308016488x^{23} + 238292173768273828749x^{24} - (19146258135816088501224x^{25})/25 + 2331916055003241548226x^{26} - 6736646381120475583764x^{27} + 18488763007525700846649x^{28} - 48264408157576669898532x^{29} + (599857644244167183024612x^{30})/5 - 284248449886557530554488x^{31} + (2570079734390957672096829x^{32})/4 - 1386787891870780679371896x^{33} + (5720500053966970302409071x^{34})/2 - (28206275157871771317939099x^{35})/5 + (4258594484619855669571973x^{36})/4 - 19237666204653402059452899x^{37} + 33300287699283081927474024x^{38} - 55246631151825154632274992x^{39} + (439428796464188236515924114x^{40})/5 - 134109601422466453670901168x^{41} + 196374773511468735732390996x^{42} - 276016272695076305811040776x^{43} + 372502480574415750374856978x^{44} - (2414047083412492769871166152x^{45})/5 + 601126348833940887359223192x^{46} - 719077854633508484642475024x^{47} + 826548729646668720118931889x^{48} - 913043842041304917057126672x^{49} + (24233705307512968006891237086x^{50})/25 - 989130828878080326811887228x^{51} + 970109082168886474373197089x^{52} - 9...$

3.64.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (1-x)^{99} x dx$$

$$\downarrow 49$$

$$\int ((1-x)^{99} - (1-x)^{100}) dx$$

$$\downarrow 2009$$

$$\frac{1}{101}(1-x)^{101} - \frac{1}{100}(1-x)^{100}$$

input `Int[(1 - x)^99*x, x]`

output $-1/100*(1 - x)^{100} + (1 - x)^{101}/101$

3.64.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.64.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 500 vs. $2(19) = 38$.

Time = 0.19 (sec) , antiderivative size = 501, normalized size of antiderivative = 21.78

method	result	size
gospers	Expression too large to display	501
default	Expression too large to display	502
risch	Expression too large to display	502
parallelrisch	Expression too large to display	502

input `int(x*(1-x)^99,x,method=_RETURNVERBOSE)`

```
output -1/10100*x^2*(100*x^99-9999*x^98+494900*x^97-16165050*x^96+391960800*x^95-
7524830775*x^94+119129952480*x^93-1599564027600*x^92+18592781869200*x^91-1
90037092945700*x^90+1729128713835600*x^89-14145670154903560*x^88+104900475
306026400*x^87-710003828928813300*x^86+4411583725232560800*x^85-2528966019
2321540400*x^84+134332724433331476360*x^83-663667626664673365350*x^82+3059
800945425883628200*x^81-13203492783118238531700*x^80+534659954674417560296
00*x^79-203648157735809402877030*x^78+731164847106703494794400*x^77-247919
4963740288382850800*x^76+7952742286283782215118800*x^75-241721508964678117
32795300*x^74+69714962380376909325764496*x^73-191035745261543332611892200*
x^72+497964017002692209469746400*x^71-1236085967492793928008474900*x^70+29
24823134349146195851039200*x^69-6603091490864731434780756240*x^68+14234925
496610562332226630300*x^67-29326230200904915179092563225*x^66+577770505450
66400054331617100*x^65-108925997889075399236629520550*x^64+196625391061305
336057915852480*x^63-340025750167248881400829989825*x^62+56358486911597475
4134709022400*x^61-895722354016828362340647962400*x^60+1365609490550246519
634102631200*x^59-1997897787191193993562250150280*x^58+2805764446902887448
586210864800*x^57-3783394480727510220367051779600*x^56+4899707046811384846
791142017600*x^55-6095468885616544243924694533800*x^54+7285651347867363554
793785087040*x^53-8367877730115588952806888703200*x^52+9236242400221923655
456660172400*x^51-9798101729905753391169290598900*x^50+9990221371668611...
```

3.64.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 501 vs. $2(15) = 30$.

Time = 0.24 (sec) , antiderivative size = 501, normalized size of antiderivative = 21.78

$$\int (1-x)^{99} x dx = \text{Too large to display}$$

```
input integrate(x*(1-x)^99,x, algorithm="fricas")
```

```
output -1/101*x^101 + 99/100*x^100 - 49*x^99 + 3201/2*x^98 - 38808*x^97 + 2980131
/4*x^96 - 58975224/5*x^95 + 158372676*x^94 - 1840869492*x^93 + 18815553757
*x^92 - 171200862756*x^91 + 7002807007378/5*x^90 - 10386185673864*x^89 + 7
0297408804833*x^88 - 436790467844808*x^87 + 2503926751715004*x^86 - 665013
48729372018/5*x^85 + 131419332012806607/2*x^84 - 302950588656028082*x^83 +
1307276513180023617*x^82 - 5293662917568490696*x^81 + 2016318393423855474
03/10*x^80 - 72392559119475593544*x^79 + 245464847895078057708*x^78 - 7874
00226364730912388*x^77 + 2393282266977011062653*x^76 - 1725617880702398745
68724/25*x^75 + 18914430223915181446722*x^74 - 49303368020068535591064*x^7
3 + 122384749256712270099849*x^72 - 289586448945460019391192*x^71 + 326885
7173695411601376612/5*x^70 - 1409398564020847755666003*x^69 + 116143485944
17788189739629/4*x^68 - 5720500053966970302409071*x^67 + 21569504532490178
066659311/2*x^66 - 97339302505596701018770224/5*x^65 + 1346636634325738144
16170293/4*x^64 - 55800482090690569716307824*x^63 + 8868538158582459033075
7224*x^62 - 135208860450519457389515112*x^61 + 989058310490690095822896114
/5*x^60 - 277798460089394796889723848*x^59 + 374593512943317843600698196*x
^58 - 485119509585285628395162576*x^57 + 603511770853123192467791538*x^56
- 3606758093003645324155339152/5*x^55 + 828502745555998906218503832*x^54 -
914479445566527094599669324*x^53 + 970109082168886474373197089*x^52 - 989
130828878080326811887228*x^51 + 24233705307512968006891237086/25*x^50 - ...
```

3.64.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 561 vs. $2(12) = 24$.

Time = 0.08 (sec) , antiderivative size = 561, normalized size of antiderivative = 24.39

$$\int (1-x)^{99} x dx = \text{Too large to display}$$

```
input integrate(x*(1-x)**99,x)
```

output

```
-x**101/101 + 99*x**100/100 - 49*x**99 + 3201*x**98/2 - 38808*x**97 + 2980
131*x**96/4 - 58975224*x**95/5 + 158372676*x**94 - 1840869492*x**93 + 1881
5553757*x**92 - 171200862756*x**91 + 7002807007378*x**90/5 - 1038618567386
4*x**89 + 70297408804833*x**88 - 436790467844808*x**87 + 2503926751715004*
x**86 - 66501348729372018*x**85/5 + 131419332012806607*x**84/2 - 302950588
656028082*x**83 + 1307276513180023617*x**82 - 5293662917568490696*x**81 +
201631839342385547403*x**80/10 - 72392559119475593544*x**79 + 245464847895
078057708*x**78 - 787400226364730912388*x**77 + 2393282266977011062653*x**
76 - 172561788070239874568724*x**75/25 + 18914430223915181446722*x**74 - 4
9303368020068535591064*x**73 + 122384749256712270099849*x**72 - 2895864489
45460019391192*x**71 + 3268857173695411601376612*x**70/5 - 140939856402084
7755666003*x**69 + 11614348594417788189739629*x**68/4 - 572050005396697030
2409071*x**67 + 21569504532490178066659311*x**66/2 - 973393025055967010187
70224*x**65/5 + 134663663432573814416170293*x**64/4 - 55800482090690569716
307824*x**63 + 88685381585824590330757224*x**62 - 135208860450519457389515
112*x**61 + 989058310490690095822896114*x**60/5 - 277798460089394796889723
848*x**59 + 374593512943317843600698196*x**58 - 48511950958528562839516257
6*x**57 + 603511770853123192467791538*x**56 - 3606758093003645324155339152
*x**55/5 + 828502745555998906218503832*x**54 - 914479445566527094599669324
*x**53 + 970109082168886474373197089*x**52 - 98913082887808032681188722...
```

3.64.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 501 vs. $2(15) = 30$.

Time = 0.20 (sec) , antiderivative size = 501, normalized size of antiderivative = 21.78

$$\int (1-x)^{99} x dx = \text{Too large to display}$$

input `integrate(x*(1-x)^99,x, algorithm="maxima")`

```
output -1/101*x^101 + 99/100*x^100 - 49*x^99 + 3201/2*x^98 - 38808*x^97 + 2980131
/4*x^96 - 58975224/5*x^95 + 158372676*x^94 - 1840869492*x^93 + 18815553757
*x^92 - 171200862756*x^91 + 7002807007378/5*x^90 - 10386185673864*x^89 + 7
0297408804833*x^88 - 436790467844808*x^87 + 2503926751715004*x^86 - 665013
48729372018/5*x^85 + 131419332012806607/2*x^84 - 302950588656028082*x^83 +
1307276513180023617*x^82 - 5293662917568490696*x^81 + 2016318393423855474
03/10*x^80 - 72392559119475593544*x^79 + 245464847895078057708*x^78 - 7874
00226364730912388*x^77 + 2393282266977011062653*x^76 - 1725617880702398745
68724/25*x^75 + 18914430223915181446722*x^74 - 49303368020068535591064*x^7
3 + 122384749256712270099849*x^72 - 289586448945460019391192*x^71 + 326885
7173695411601376612/5*x^70 - 1409398564020847755666003*x^69 + 116143485944
17788189739629/4*x^68 - 5720500053966970302409071*x^67 + 21569504532490178
066659311/2*x^66 - 97339302505596701018770224/5*x^65 + 1346636634325738144
16170293/4*x^64 - 55800482090690569716307824*x^63 + 8868538158582459033075
7224*x^62 - 135208860450519457389515112*x^61 + 989058310490690095822896114
/5*x^60 - 277798460089394796889723848*x^59 + 374593512943317843600698196*x
^58 - 485119509585285628395162576*x^57 + 603511770853123192467791538*x^56
- 3606758093003645324155339152/5*x^55 + 828502745555998906218503832*x^54 -
914479445566527094599669324*x^53 + 970109082168886474373197089*x^52 - 989
130828878080326811887228*x^51 + 24233705307512968006891237086/25*x^50 - ...
```

3.64.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 501 vs. $2(15) = 30$.

Time = 0.27 (sec) , antiderivative size = 501, normalized size of antiderivative = 21.78

$$\int (1-x)^{99} x dx = \text{Too large to display}$$

```
input integrate(x*(1-x)^99,x, algorithm="giac")
```


output

```
-1/101*x^101 + 99/100*x^100 - 49*x^99 + 3201/2*x^98 - 38808*x^97 + 2980131
/4*x^96 - 58975224/5*x^95 + 158372676*x^94 - 1840869492*x^93 + 18815553757
*x^92 - 171200862756*x^91 + 7002807007378/5*x^90 - 10386185673864*x^89 + 7
0297408804833*x^88 - 436790467844808*x^87 + 2503926751715004*x^86 - 665013
48729372018/5*x^85 + 131419332012806607/2*x^84 - 302950588656028082*x^83 +
1307276513180023617*x^82 - 5293662917568490696*x^81 + 2016318393423855474
03/10*x^80 - 72392559119475593544*x^79 + 245464847895078057708*x^78 - 7874
00226364730912388*x^77 + 2393282266977011062653*x^76 - 1725617880702398745
68724/25*x^75 + 18914430223915181446722*x^74 - 49303368020068535591064*x^7
3 + 122384749256712270099849*x^72 - 289586448945460019391192*x^71 + 326885
7173695411601376612/5*x^70 - 1409398564020847755666003*x^69 + 116143485944
17788189739629/4*x^68 - 5720500053966970302409071*x^67 + 21569504532490178
066659311/2*x^66 - 97339302505596701018770224/5*x^65 + 1346636634325738144
16170293/4*x^64 - 55800482090690569716307824*x^63 + 8868538158582459033075
7224*x^62 - 135208860450519457389515112*x^61 + 989058310490690095822896114
/5*x^60 - 277798460089394796889723848*x^59 + 374593512943317843600698196*x
^58 - 485119509585285628395162576*x^57 + 603511770853123192467791538*x^56
- 3606758093003645324155339152/5*x^55 + 828502745555998906218503832*x^54 -
914479445566527094599669324*x^53 + 970109082168886474373197089*x^52 - 989
130828878080326811887228*x^51 + 24233705307512968006891237086/25*x^50 ...
```

3.64.9 Mupad [B] (verification not implemented)

Time = 13.92 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.52

$$\int (1-x)^{99} x dx = -\frac{(100x+1)(x-1)^{100}}{10100}$$

input `int(-x*(x - 1)^99,x)`

output `-((100*x + 1)*(x - 1)^100)/10100`

3.65 $\int \csc(x) \sin(4x) dx$

3.65.1	Optimal result	433
3.65.2	Mathematica [A] (verified)	433
3.65.3	Rubi [A] (verified)	434
3.65.4	Maple [A] (verified)	435
3.65.5	Fricas [A] (verification not implemented)	435
3.65.6	Sympy [A] (verification not implemented)	435
3.65.7	Maxima [A] (verification not implemented)	436
3.65.8	Giac [A] (verification not implemented)	436
3.65.9	Mupad [B] (verification not implemented)	436

3.65.1 Optimal result

Integrand size = 7, antiderivative size = 13

$$\int \csc(x) \sin(4x) dx = 4 \sin(x) - \frac{8 \sin^3(x)}{3}$$

output `4*sin(x)-8/3*sin(x)^3`

3.65.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \csc(x) \sin(4x) dx = 4 \sin(x) - \frac{8 \sin^3(x)}{3}$$

input `Integrate[Csc[x]*Sin[4*x],x]`

output `4*Sin[x] - (8*Sin[x]^3)/3`

3.65.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 4878, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \sin(4x) \csc(x) dx \\
 \downarrow 3042 \\
 \int \frac{\sin(4x)}{\sin(x)} dx \\
 \downarrow 4878 \\
 \int (4 - 8 \sin^2(x)) d \sin(x) \\
 \downarrow 2009 \\
 4 \sin(x) - \frac{8 \sin^3(x)}{3}
 \end{array}$$

input `Int[Csc[x]*Sin[4*x],x]`

output `4*Sin[x] - (8*Sin[x]^3)/3`

3.65.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4878 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Sin[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Sin[v]/d, u/Cos[v], x], x], x, Sin[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Sin[v], x], u/Cos[v], x]]`

3.65.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
derivativedivides	$4 \sin(x) - \frac{8 \sin(x)^3}{3}$	12
default	$4 \sin(x) - \frac{8 \sin(x)^3}{3}$	12
risch	$2 \sin(x) + \frac{2 \sin(3x)}{3}$	12

input `int(sin(4*x)/sin(x),x,method=_RETURNVERBOSE)`output `4*sin(x)-8/3*sin(x)^3`**3.65.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

$$\int \csc(x) \sin(4x) dx = \frac{4}{3} (2 \cos(x)^2 + 1) \sin(x)$$

input `integrate(sin(4*x)/sin(x),x, algorithm="fricas")`output `4/3*(2*cos(x)^2 + 1)*sin(x)`**3.65.6 Sympy [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

$$\int \csc(x) \sin(4x) dx = -\frac{8 \sin^3(x)}{3} + 4 \sin(x)$$

input `integrate(sin(4*x)/sin(x),x)`output `-8*sin(x)**3/3 + 4*sin(x)`

3.65.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \csc(x) \sin(4x) dx = -\frac{8}{3} \sin(x)^3 + 4 \sin(x)$$

input `integrate(sin(4*x)/sin(x),x, algorithm="maxima")`output `-8/3*sin(x)^3 + 4*sin(x)`**3.65.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \csc(x) \sin(4x) dx = \frac{2}{3} \sin(3x) + 2 \sin(x)$$

input `integrate(sin(4*x)/sin(x),x, algorithm="giac")`output `2/3*sin(3*x) + 2*sin(x)`**3.65.9 Mupad [B] (verification not implemented)**

Time = 14.36 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \csc(x) \sin(4x) dx = 4 \sin(x) - \frac{8 \sin(x)^3}{3}$$

input `int(sin(4*x)/sin(x),x)`output `4*sin(x) - (8*sin(x)^3)/3`

3.66 $\int \frac{1}{(1 + \sqrt[3]{x})\sqrt{x}} dx$

3.66.1 Optimal result 437
 3.66.2 Mathematica [A] (verified) 437
 3.66.3 Rubi [A] (verified) 438
 3.66.4 Maple [A] (verified) 439
 3.66.5 Fricas [A] (verification not implemented) 440
 3.66.6 Sympy [A] (verification not implemented) 440
 3.66.7 Maxima [A] (verification not implemented) 440
 3.66.8 Giac [A] (verification not implemented) 441
 3.66.9 Mupad [B] (verification not implemented) 441

3.66.1 Optimal result

Integrand size = 15, antiderivative size = 16

$$\int \frac{1}{(1 + \sqrt[3]{x})\sqrt{x}} dx = 6\sqrt[6]{x} - 6 \arctan(\sqrt[6]{x})$$

output `6*x^(1/6)-6*arctan(x^(1/6))`

3.66.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1 + \sqrt[3]{x})\sqrt{x}} dx = 6\sqrt[6]{x} - 6 \arctan(\sqrt[6]{x})$$

input `Integrate[1/((1 + x^(1/3))*Sqrt[x]),x]`

output `6*x^(1/6) - 6*ArcTan[x^(1/6)]`

3.66.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {864, 60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(\sqrt[3]{x} + 1)\sqrt{x}} dx \\
 & \quad \downarrow 864 \\
 & 3 \int \frac{\sqrt[6]{x}}{\sqrt[3]{x} + 1} d\sqrt[3]{x} \\
 & \quad \downarrow 60 \\
 & 3 \left(2\sqrt[6]{x} - \int \frac{1}{(\sqrt[3]{x} + 1)\sqrt[6]{x}} d\sqrt[3]{x} \right) \\
 & \quad \downarrow 73 \\
 & 3 \left(2\sqrt[6]{x} - 2 \int \frac{1}{x^{2/3} + 1} d\sqrt[6]{x} \right) \\
 & \quad \downarrow 216 \\
 & 3(2\sqrt[6]{x} - 2 \arctan(\sqrt[6]{x}))
 \end{aligned}$$

input `Int[1/((1 + x^(1/3))*Sqrt[x]),x]`

output `3*(2*x^(1/6) - 2*ArcTan[x^(1/6)])`

3.66.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

3.66. $\int \frac{1}{(1 + \sqrt[3]{x})\sqrt{x}} dx$

- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 864 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a + b*x^(k*n))^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, m, p}, x] && FractionQ[n]`

3.66.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$6x^{\frac{1}{6}} - 6 \arctan\left(x^{\frac{1}{6}}\right)$	13
default	$6x^{\frac{1}{6}} - 6 \arctan\left(x^{\frac{1}{6}}\right)$	13
meijerg	$6x^{\frac{1}{6}} - 6 \arctan\left(x^{\frac{1}{6}}\right)$	13

input `int(1/x^(1/2)/(1+x^(1/3)),x,method=_RETURNVERBOSE)`

output `6*x^(1/6)-6*arctan(x^(1/6))`

3.66.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{(1 + \sqrt[3]{x})\sqrt{x}} dx = 6x^{\frac{1}{6}} - 6 \arctan\left(x^{\frac{1}{6}}\right)$$

input `integrate(1/x^(1/2)/(1+x^(1/3)),x, algorithm="fricas")`output `6*x^(1/6) - 6*arctan(x^(1/6))`**3.66.6 Sympy [A] (verification not implemented)**

Time = 0.41 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{1}{(1 + \sqrt[3]{x})\sqrt{x}} dx = 6\sqrt[6]{x} + 6 \operatorname{atan}\left(\frac{1}{\sqrt[6]{x}}\right)$$

input `integrate(1/x**(1/2)/(1+x**(1/3)),x)`output `6*x**(1/6) + 6*atan(x**(-1/6))`**3.66.7 Maxima [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{(1 + \sqrt[3]{x})\sqrt{x}} dx = 6x^{\frac{1}{6}} - 6 \arctan\left(x^{\frac{1}{6}}\right)$$

input `integrate(1/x^(1/2)/(1+x^(1/3)),x, algorithm="maxima")`output `6*x^(1/6) - 6*arctan(x^(1/6))`

3.66.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{(1 + \sqrt[3]{x})\sqrt{x}} dx = 6x^{\frac{1}{6}} - 6 \arctan\left(x^{\frac{1}{6}}\right)$$

input `integrate(1/x^(1/2)/(1+x^(1/3)),x, algorithm="giac")`output `6*x^(1/6) - 6*arctan(x^(1/6))`**3.66.9 Mupad [B] (verification not implemented)**

Time = 14.48 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{(1 + \sqrt[3]{x})\sqrt{x}} dx = 6x^{1/6} - 6 \operatorname{atan}(x^{1/6})$$

input `int(1/(x^(1/2)*(x^(1/3) + 1)),x)`output `6*x^(1/6) - 6*atan(x^(1/6))`

3.67 $\int \frac{1}{\sqrt{-1+2x^2}} dx$

3.67.1 Optimal result	442
3.67.2 Mathematica [B] (verified)	442
3.67.3 Rubi [A] (verified)	443
3.67.4 Maple [A] (verified)	444
3.67.5 Fricas [A] (verification not implemented)	444
3.67.6 Sympy [A] (verification not implemented)	444
3.67.7 Maxima [A] (verification not implemented)	445
3.67.8 Giac [A] (verification not implemented)	445
3.67.9 Mupad [B] (verification not implemented)	445

3.67.1 Optimal result

Integrand size = 11, antiderivative size = 25

$$\int \frac{1}{\sqrt{-1+2x^2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}x}{\sqrt{-1+2x^2}}\right)}{\sqrt{2}}$$

output `1/2*arctanh(2^(1/2)*x/(2*x^2-1)^(1/2))*2^(1/2)`

3.67.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 54 vs. 2(25) = 50.

Time = 0.03 (sec) , antiderivative size = 54, normalized size of antiderivative = 2.16

$$\int \frac{1}{\sqrt{-1+2x^2}} dx = \frac{-\log\left(\sqrt{2} - \frac{2x}{\sqrt{-1+2x^2}}\right) + \log\left(\sqrt{2} + \frac{2x}{\sqrt{-1+2x^2}}\right)}{2\sqrt{2}}$$

input `Integrate[1/Sqrt[-1 + 2*x^2], x]`

output `(-Log[Sqrt[2] - (2*x)/Sqrt[-1 + 2*x^2]] + Log[Sqrt[2] + (2*x)/Sqrt[-1 + 2*x^2]])/(2*Sqrt[2])`

3.67.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{2x^2 - 1}} dx$$

↓ 224

$$\int \frac{1}{1 - \frac{2x^2}{2x^2 - 1}} d \frac{x}{\sqrt{2x^2 - 1}}$$

↓ 219

$$\frac{\operatorname{arctanh}\left(\frac{\sqrt{2}x}{\sqrt{2x^2 - 1}}\right)}{\sqrt{2}}$$

input `Int[1/Sqrt[-1 + 2*x^2], x]`

output `ArcTanh[(Sqrt[2]*x)/Sqrt[-1 + 2*x^2]]/Sqrt[2]`

3.67.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

3.67.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

method	result	size
default	$\frac{\ln(x\sqrt{2}+\sqrt{2x^2-1})\sqrt{2}}{2}$	22
pseudoelliptic	$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2x^2-1}\sqrt{2}}{2x}\right)}{2}$	24
trager	$\frac{\operatorname{RootOf}(_Z^2-2) \ln(\operatorname{RootOf}(_Z^2-2)\sqrt{2x^2-1}+2x)}{2}$	30
meijerg	$\frac{\sqrt{-\operatorname{signum}(2x^2-1)}\sqrt{2} \arcsin(x\sqrt{2})}{2\sqrt{\operatorname{signum}(2x^2-1)}}$	34

input `int(1/(2*x^2-1)^(1/2),x,method=_RETURNVERBOSE)`output `1/2*ln(x*2^(1/2)+(2*x^2-1)^(1/2))*2^(1/2)`**3.67.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.12

$$\int \frac{1}{\sqrt{-1+2x^2}} dx = \frac{1}{4} \sqrt{2} \log(2\sqrt{2}\sqrt{2x^2-1}x + 4x^2 - 1)$$

input `integrate(1/(2*x^2-1)^(1/2),x, algorithm="fricas")`output `1/4*sqrt(2)*log(2*sqrt(2)*sqrt(2*x^2 - 1)*x + 4*x^2 - 1)`**3.67.6 Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

$$\int \frac{1}{\sqrt{-1+2x^2}} dx = \frac{\sqrt{2} \log(2x + \sqrt{2}\sqrt{2x^2-1})}{2}$$

input `integrate(1/(2*x**2-1)**(1/2),x)`output `sqrt(2)*log(2*x + sqrt(2)*sqrt(2*x**2 - 1))/2`

3.67.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{1}{\sqrt{-1+2x^2}} dx = \frac{1}{2} \sqrt{2} \log \left(2 \sqrt{2} \sqrt{2x^2 - 1} + 4x \right)$$

input `integrate(1/(2*x^2-1)^(1/2),x, algorithm="maxima")`output `1/2*sqrt(2)*log(2*sqrt(2)*sqrt(2*x^2 - 1) + 4*x)`**3.67.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.44

$$\int \frac{1}{\sqrt{-1+2x^2}} dx = \frac{1}{2} \sqrt{2x^2 - 1}x + \frac{1}{4} \sqrt{2} \log \left(\left| -\sqrt{2}x + \sqrt{2x^2 - 1} \right| \right)$$

input `integrate(1/(2*x^2-1)^(1/2),x, algorithm="giac")`output `1/2*sqrt(2*x^2 - 1)*x + 1/4*sqrt(2)*log(abs(-sqrt(2)*x + sqrt(2*x^2 - 1)))`**3.67.9 Mupad [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int \frac{1}{\sqrt{-1+2x^2}} dx = \frac{\sqrt{2} \ln \left(\sqrt{2}x + \sqrt{2x^2 - 1} \right)}{2}$$

input `int(1/(2*x^2 - 1)^(1/2),x)`output `(2^(1/2)*log(2^(1/2)*x + (2*x^2 - 1)^(1/2)))/2`

3.68 $\int \frac{1}{\sqrt{-1+e^x}} dx$

3.68.1	Optimal result	446
3.68.2	Mathematica [A] (verified)	446
3.68.3	Rubi [A] (verified)	447
3.68.4	Maple [A] (verified)	448
3.68.5	Fricas [A] (verification not implemented)	448
3.68.6	Sympy [A] (verification not implemented)	449
3.68.7	Maxima [A] (verification not implemented)	449
3.68.8	Giac [A] (verification not implemented)	449
3.68.9	Mupad [B] (verification not implemented)	450

3.68.1 Optimal result

Integrand size = 9, antiderivative size = 12

$$\int \frac{1}{\sqrt{-1+e^x}} dx = 2 \arctan(\sqrt{-1+e^x})$$

output `2*arctan((-1+exp(x))^(1/2))`

3.68.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{-1+e^x}} dx = 2 \arctan(\sqrt{-1+e^x})$$

input `Integrate[1/Sqrt[-1 + E^x],x]`

output `2*ArcTan[Sqrt[-1 + E^x]]`

3.68.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2720, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt{e^x - 1}} dx \\ & \quad \downarrow \text{2720} \\ & \int \frac{e^{-x}}{\sqrt{e^x - 1}} de^x \\ & \quad \downarrow \text{73} \\ & 2 \int \frac{1}{1 + e^{2x}} d\sqrt{-1 + e^x} \\ & \quad \downarrow \text{216} \\ & 2 \arctan(\sqrt{e^x - 1}) \end{aligned}$$

input `Int[1/Sqrt[-1 + E^x],x]`

output `2*ArcTan[Sqrt[-1 + E^x]]`

3.68.3.1 Defintions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`


```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.68.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$2 \arctan(\sqrt{e^x - 1})$	10
default	$2 \arctan(\sqrt{e^x - 1})$	10

```
input int(1/(exp(x)-1)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 2*arctan((exp(x)-1)^(1/2))
```

3.68.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{-1 + e^x}} dx = 2 \arctan(\sqrt{e^x - 1})$$

```
input integrate(1/(exp(x)-1)^(1/2),x, algorithm="fracas")
```

```
output 2*arctan(sqrt(e^x - 1))
```

3.68.6 Sympy [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1}{\sqrt{-1+e^x}} dx = 2 \operatorname{atan}(\sqrt{e^x-1})$$

input `integrate(1/(exp(x)-1)**(1/2),x)`output `2*atan(sqrt(exp(x) - 1))`**3.68.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{-1+e^x}} dx = 2 \operatorname{arctan}(\sqrt{e^x-1})$$

input `integrate(1/(exp(x)-1)^(1/2),x, algorithm="maxima")`output `2*arctan(sqrt(e^x - 1))`**3.68.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{-1+e^x}} dx = 2 \operatorname{arctan}(\sqrt{e^x-1})$$

input `integrate(1/(exp(x)-1)^(1/2),x, algorithm="giac")`output `2*arctan(sqrt(e^x - 1))`

3.68.9 Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{-1+e^x}} dx = 2 \operatorname{atan}(\sqrt{e^x - 1})$$

input `int(1/(exp(x) - 1)^(1/2),x)`

output `2*atan((exp(x) - 1)^(1/2))`

3.69 $\int \frac{x}{4+x^4} dx$

3.69.1	Optimal result	451
3.69.2	Mathematica [A] (verified)	451
3.69.3	Rubi [A] (verified)	452
3.69.4	Maple [A] (verified)	453
3.69.5	Fricas [A] (verification not implemented)	453
3.69.6	Sympy [A] (verification not implemented)	453
3.69.7	Maxima [A] (verification not implemented)	454
3.69.8	Giac [A] (verification not implemented)	454
3.69.9	Mupad [B] (verification not implemented)	454

3.69.1 Optimal result

Integrand size = 9, antiderivative size = 12

$$\int \frac{x}{4+x^4} dx = \frac{1}{4} \arctan\left(\frac{x^2}{2}\right)$$

output `1/4*arctan(1/2*x^2)`

3.69.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{x}{4+x^4} dx = \frac{1}{4} \arctan\left(\frac{x^2}{2}\right)$$

input `Integrate[x/(4 + x^4), x]`

output `ArcTan[x^2/2]/4`

3.69.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {807, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{x^4 + 4} dx$$

↓ 807

$$\frac{1}{2} \int \frac{1}{x^4 + 4} dx^2$$

↓ 216

$$\frac{1}{4} \arctan\left(\frac{x^2}{2}\right)$$

input `Int[x/(4 + x^4), x]`

output `ArcTan[x^2/2]/4`

3.69.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

3.69.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

method	result	size
default	$\frac{\arctan\left(\frac{x^2}{2}\right)}{4}$	9
meijerg	$\frac{\arctan\left(\frac{x^2}{2}\right)}{4}$	9
risch	$\frac{\arctan\left(\frac{x^2}{2}\right)}{4}$	9
parallelrisch	$-\frac{i \ln(x-1-i)}{8} + \frac{i \ln(x-1+i)}{8} + \frac{i \ln(x+1-i)}{8} - \frac{i \ln(x+1+i)}{8}$	38

input `int(x/(x^4+4),x,method=_RETURNVERBOSE)`output `1/4*arctan(1/2*x^2)`**3.69.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x}{4+x^4} dx = \frac{1}{4} \arctan\left(\frac{1}{2}x^2\right)$$

input `integrate(x/(x^4+4),x, algorithm="fricas")`output `1/4*arctan(1/2*x^2)`**3.69.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.58

$$\int \frac{x}{4+x^4} dx = \frac{\operatorname{atan}\left(\frac{x^2}{2}\right)}{4}$$

input `integrate(x/(x**4+4),x)`output `atan(x**2/2)/4`

3.69.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x}{4+x^4} dx = \frac{1}{4} \arctan\left(\frac{1}{2}x^2\right)$$

input `integrate(x/(x^4+4),x, algorithm="maxima")`output `1/4*arctan(1/2*x^2)`**3.69.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x}{4+x^4} dx = \frac{1}{4} \arctan\left(\frac{1}{2}x^2\right)$$

input `integrate(x/(x^4+4),x, algorithm="giac")`output `1/4*arctan(1/2*x^2)`**3.69.9 Mupad [B] (verification not implemented)**

Time = 15.46 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x}{4+x^4} dx = \frac{\operatorname{atan}\left(\frac{x^2}{2}\right)}{4}$$

input `int(x/(x^4 + 4),x)`output `atan(x^2/2)/4`

$$3.70 \quad \int \frac{2}{(\cos(x) - \sin(x))^2} dx$$

3.70.1	Optimal result	455
3.70.2	Mathematica [A] (verified)	455
3.70.3	Rubi [A] (verified)	456
3.70.4	Maple [A] (verified)	457
3.70.5	Fricas [A] (verification not implemented)	457
3.70.6	Sympy [B] (verification not implemented)	457
3.70.7	Maxima [A] (verification not implemented)	458
3.70.8	Giac [A] (verification not implemented)	458
3.70.9	Mupad [B] (verification not implemented)	458

3.70.1 Optimal result

Integrand size = 11, antiderivative size = 8

$$\int \frac{2}{(\cos(x) - \sin(x))^2} dx = \frac{2}{-1 + \cot(x)}$$

output `2/(-1+cot(x))`

3.70.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.62

$$\int \frac{2}{(\cos(x) - \sin(x))^2} dx = \frac{2 \sin(x)}{\cos(x) - \sin(x)}$$

input `Integrate[2/(Cos[x] - Sin[x])^2,x]`

output `(2*Sin[x])/(Cos[x] - Sin[x])`

3.70.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.62, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {27, 3042, 3554}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{2}{(\cos(x) - \sin(x))^2} dx \\ & \quad \downarrow 27 \\ & 2 \int \frac{1}{(\cos(x) - \sin(x))^2} dx \\ & \quad \downarrow 3042 \\ & 2 \int \frac{1}{(\cos(x) - \sin(x))^2} dx \\ & \quad \downarrow 3554 \\ & \frac{2 \sin(x)}{\cos(x) - \sin(x)} \end{aligned}$$

input `Int[2/(Cos[x] - Sin[x])^2,x]`

output `(2*Sin[x])/(Cos[x] - Sin[x])`

3.70.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3554 `Int[(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-2), x_Symbol] := Simp[Sin[c + d*x]/(a*d*(a*Cos[c + d*x] + b*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]`

3.70.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
default	$-\frac{2}{\tan(x)-1}$	9
risch	$\frac{2}{e^{2ix}-i}$	13
parallelrisch	$\frac{2\sin(x)}{\cos(x)-\sin(x)}$	14
norman	$-\frac{4\tan(\frac{x}{2})}{\tan^2(\frac{x}{2})+2\tan(\frac{x}{2})-1}$	23

input `int(2/(cos(x)-sin(x))^2,x,method=_RETURNVERBOSE)`

output `-2/(tan(x)-1)`

3.70.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.88

$$\int \frac{2}{(\cos(x) - \sin(x))^2} dx = \frac{\cos(x) + \sin(x)}{\cos(x) - \sin(x)}$$

input `integrate(2/(cos(x)-sin(x))^2,x, algorithm="fracas")`

output `(cos(x) + sin(x))/(cos(x) - sin(x))`

3.70.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 22 vs. $2(5) = 10$.

Time = 0.33 (sec) , antiderivative size = 22, normalized size of antiderivative = 2.75

$$\int \frac{2}{(\cos(x) - \sin(x))^2} dx = -\frac{4\tan\left(\frac{x}{2}\right)}{\tan^2\left(\frac{x}{2}\right) + 2\tan\left(\frac{x}{2}\right) - 1}$$

input `integrate(2/(cos(x)-sin(x))**2,x)`

output `-4*tan(x/2)/(tan(x/2)**2 + 2*tan(x/2) - 1)`

3.70. $\int \frac{2}{(\cos(x)-\sin(x))^2} dx$

3.70.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{2}{(\cos(x) - \sin(x))^2} dx = -\frac{2}{\tan(x) - 1}$$

input `integrate(2/(cos(x)-sin(x))^2,x, algorithm="maxima")`output `-2/(tan(x) - 1)`**3.70.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{2}{(\cos(x) - \sin(x))^2} dx = -\frac{2}{\tan(x) - 1}$$

input `integrate(2/(cos(x)-sin(x))^2,x, algorithm="giac")`output `-2/(tan(x) - 1)`**3.70.9 Mupad [B] (verification not implemented)**

Time = 15.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.62

$$\int \frac{2}{(\cos(x) - \sin(x))^2} dx = \frac{2 \sin(x)}{\cos(x) - \sin(x)}$$

input `int(2/(cos(x) - sin(x))^2,x)`output `(2*sin(x))/(cos(x) - sin(x))`

3.71 $\int x \coth(x) \operatorname{csch}(x) dx$

3.71.1	Optimal result	459
3.71.2	Mathematica [B] (verified)	459
3.71.3	Rubi [A] (verified)	460
3.71.4	Maple [B] (verified)	461
3.71.5	Fricas [B] (verification not implemented)	461
3.71.6	Sympy [B] (verification not implemented)	462
3.71.7	Maxima [B] (verification not implemented)	462
3.71.8	Giac [B] (verification not implemented)	463
3.71.9	Mupad [B] (verification not implemented)	463

3.71.1 Optimal result

Integrand size = 6, antiderivative size = 11

$$\int x \coth(x) \operatorname{csch}(x) dx = -\operatorname{arctanh}(\cosh(x)) - x \operatorname{csch}(x)$$

output `-arctanh(cosh(x))-x*csch(x)`

3.71.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 39 vs. $2(11) = 22$.

Time = 0.01 (sec) , antiderivative size = 39, normalized size of antiderivative = 3.55

$$\int x \coth(x) \operatorname{csch}(x) dx = -\frac{1}{2}x \coth\left(\frac{x}{2}\right) - \log\left(\cosh\left(\frac{x}{2}\right)\right) + \log\left(\sinh\left(\frac{x}{2}\right)\right) + \frac{1}{2}x \tanh\left(\frac{x}{2}\right)$$

input `Integrate[x*Coth[x]*Csch[x],x]`

output `-1/2*(x*Coth[x/2]) - Log[Cosh[x/2]] + Log[Sinh[x/2]] + (x*Tanh[x/2])/2`

3.71.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5942, 3042, 26, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \coth(x) \operatorname{csch}(x) dx \\
 & \quad \downarrow \text{5942} \\
 & \int \operatorname{csch}(x) dx - x \operatorname{csch}(x) \\
 & \quad \downarrow \text{3042} \\
 & -x \operatorname{csch}(x) + \int i \csc(ix) dx \\
 & \quad \downarrow \text{26} \\
 & -x \operatorname{csch}(x) + i \int \csc(ix) dx \\
 & \quad \downarrow \text{4257} \\
 & -\operatorname{arctanh}(\cosh(x)) - x \operatorname{csch}(x)
 \end{aligned}$$

input `Int[x*Coth[x]*Csch[x],x]`

output `-ArcTanh[Cosh[x]] - x*Csch[x]`

3.71.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 5942 `Int[Coth[(a_.) + (b_.)*(x_)^(n_.)]^(q_.)*Csch[(a_.) + (b_.)*(x_)^(n_.)]^(p_.)*(x_)^(m_.), x_Symbol] := Simp[(-x^(m - n + 1))*(Csch[a + b*x^n]^p/(b*n*p)), x] + Simp[(m - n + 1)/(b*n*p) Int[x^(m - n)*Csch[a + b*x^n]^p, x], x] /; FreeQ[{a, b, p}, x] && RationalQ[m] && IntegerQ[n] && GeQ[m - n, 0] && EqQ[q, 1]`

3.71.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. $2(11) = 22$.

Time = 0.20 (sec) , antiderivative size = 27, normalized size of antiderivative = 2.45

method	result	size
risch	$-\frac{2xe^x}{e^{2x}-1} + \ln(e^x - 1) - \ln(e^x + 1)$	27

input `int(x*cosh(x)/sinh(x)^2,x,method=_RETURNVERBOSE)`

output `-2*x*exp(x)/(exp(2*x)-1)+ln(exp(x)-1)-ln(exp(x)+1)`

3.71.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 80 vs. $2(11) = 22$.

Time = 0.23 (sec) , antiderivative size = 80, normalized size of antiderivative = 7.27

$$\int x \coth(x) \operatorname{csch}(x) dx = \frac{2x \cosh(x) + (\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1) \log(\cosh(x) + \sinh(x) + 1) - (\cosh(x) + \sinh(x)) \operatorname{csch}(x)}{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2}$$

input `integrate(x*cosh(x)/sinh(x)^2,x, algorithm="fricas")`

output $-(2*x*\cosh(x) + (\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 - 1)*\log(\cosh(x) + \sinh(x) + 1) - (\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 - 1)*\log(\cosh(x) + \sinh(x) - 1) + 2*x*\sinh(x))/(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 - 1)$

3.71.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 22 vs. $2(10) = 20$.

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 2.00

$$\int x \coth(x) \operatorname{csch}(x) dx = \frac{x \tanh\left(\frac{x}{2}\right)}{2} - \frac{x}{2 \tanh\left(\frac{x}{2}\right)} + \log\left(\tanh\left(\frac{x}{2}\right)\right)$$

input `integrate(x*cosh(x)/sinh(x)**2,x)`

output `x*tanh(x/2)/2 - x/(2*tanh(x/2)) + log(tanh(x/2))`

3.71.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. $2(11) = 22$.

Time = 0.23 (sec) , antiderivative size = 26, normalized size of antiderivative = 2.36

$$\int x \coth(x) \operatorname{csch}(x) dx = -\frac{2xe^x}{e^{2x} - 1} - \log(e^x + 1) + \log(e^x - 1)$$

input `integrate(x*cosh(x)/sinh(x)^2,x, algorithm="maxima")`

output `-2*x*e^x/(e^(2*x) - 1) - log(e^x + 1) + log(e^x - 1)`

3.71.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 49 vs. $2(11) = 22$.

Time = 0.28 (sec) , antiderivative size = 49, normalized size of antiderivative = 4.45

$$\int x \coth(x) \operatorname{csch}(x) dx$$

$$= -\frac{2xe^x + e^{(2x)} \log(e^x + 1) - e^{(2x)} \log(e^x - 1) - \log(e^x + 1) + \log(e^x - 1)}{e^{(2x)} - 1}$$

input `integrate(x*cosh(x)/sinh(x)^2,x, algorithm="giac")`

output `-(2*x*e^x + e^(2*x)*log(e^x + 1) - e^(2*x)*log(e^x - 1) - log(e^x + 1) + log(e^x - 1))/(e^(2*x) - 1)`

3.71.9 Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 30, normalized size of antiderivative = 2.73

$$\int x \coth(x) \operatorname{csch}(x) dx = \ln(2 - 2e^x) - \ln(-2e^x - 2) - \frac{2xe^x}{e^{2x} - 1}$$

input `int((x*cosh(x))/sinh(x)^2,x)`

output `log(2 - 2*exp(x)) - log(- 2*exp(x) - 2) - (2*x*exp(x))/(exp(2*x) - 1)`

3.72 $\int x^5 \sqrt{1 + x^3} dx$

3.72.1	Optimal result	464
3.72.2	Mathematica [A] (verified)	464
3.72.3	Rubi [A] (verified)	465
3.72.4	Maple [A] (verified)	466
3.72.5	Fricas [A] (verification not implemented)	466
3.72.6	Sympy [A] (verification not implemented)	467
3.72.7	Maxima [A] (verification not implemented)	467
3.72.8	Giac [A] (verification not implemented)	467
3.72.9	Mupad [B] (verification not implemented)	468

3.72.1 Optimal result

Integrand size = 13, antiderivative size = 27

$$\int x^5 \sqrt{1 + x^3} dx = -\frac{2}{9}(1 + x^3)^{3/2} + \frac{2}{15}(1 + x^3)^{5/2}$$

output `-2/9*(x^3+1)^(3/2)+2/15*(x^3+1)^(5/2)`

3.72.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

$$\int x^5 \sqrt{1 + x^3} dx = \frac{2}{45}(1 + x^3)^{3/2}(-2 + 3x^3)$$

input `Integrate[x^5*Sqrt[1 + x^3],x]`

output `(2*(1 + x^3)^(3/2)*(-2 + 3*x^3))/45`

3.72.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.15, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {798, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^5 \sqrt{x^3 + 1} dx \\ & \quad \downarrow \text{798} \\ & \frac{1}{3} \int x^3 \sqrt{x^3 + 1} dx^3 \\ & \quad \downarrow \text{53} \\ & \frac{1}{3} \int \left((x^3 + 1)^{3/2} - \sqrt{x^3 + 1} \right) dx^3 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{3} \left(\frac{2}{5} (x^3 + 1)^{5/2} - \frac{2}{3} (x^3 + 1)^{3/2} \right) \end{aligned}$$

input `Int[x^5*Sqrt[1 + x^3],x]`

output `((-2*(1 + x^3)^(3/2))/3 + (2*(1 + x^3)^(5/2))/5)/3`

3.72.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.72.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.63

method	result	size
pseudoelliptic	$\frac{2(x^3+1)^{\frac{3}{2}}(3x^3-2)}{45}$	17
risch	$\frac{2(3x^6+x^3-2)\sqrt{x^3+1}}{45}$	20
trager	$\left(\frac{2}{15}x^6 + \frac{2}{45}x^3 - \frac{4}{45}\right)\sqrt{x^3+1}$	21
gosper	$\frac{2(1+x)(x^2-x+1)(3x^3-2)\sqrt{x^3+1}}{45}$	28
meijerg	$-\frac{8\sqrt{\pi}}{15} + \frac{4\sqrt{\pi}(x^3+1)^{\frac{3}{2}}(-3x^3+2)}{6\sqrt{\pi} \cdot 15}$	31
default	$\frac{2x^6\sqrt{x^3+1}}{15} + \frac{2x^3\sqrt{x^3+1}}{45} - \frac{4\sqrt{x^3+1}}{45}$	35
elliptic	$\frac{2x^6\sqrt{x^3+1}}{15} + \frac{2x^3\sqrt{x^3+1}}{45} - \frac{4\sqrt{x^3+1}}{45}$	35

input `int(x^5*(x^3+1)^(1/2),x,method=_RETURNVERBOSE)`output `2/45*(x^3+1)^(3/2)*(3*x^3-2)`**3.72.5 Fricas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int x^5 \sqrt{1+x^3} dx = \frac{2}{45} (3x^6 + x^3 - 2) \sqrt{x^3 + 1}$$

input `integrate(x^5*(x^3+1)^(1/2),x, algorithm="fricas")`output `2/45*(3*x^6 + x^3 - 2)*sqrt(x^3 + 1)`

3.72.6 Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.52

$$\int x^5 \sqrt{1+x^3} dx = \frac{2x^6 \sqrt{x^3+1}}{15} + \frac{2x^3 \sqrt{x^3+1}}{45} - \frac{4\sqrt{x^3+1}}{45}$$

input `integrate(x**5*(x**3+1)**(1/2),x)`output `2*x**6*sqrt(x**3 + 1)/15 + 2*x**3*sqrt(x**3 + 1)/45 - 4*sqrt(x**3 + 1)/45`**3.72.7 Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int x^5 \sqrt{1+x^3} dx = \frac{2}{15} (x^3+1)^{\frac{5}{2}} - \frac{2}{9} (x^3+1)^{\frac{3}{2}}$$

input `integrate(x^5*(x^3+1)^(1/2),x, algorithm="maxima")`output `2/15*(x^3 + 1)^(5/2) - 2/9*(x^3 + 1)^(3/2)`**3.72.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int x^5 \sqrt{1+x^3} dx = \frac{2}{15} (x^3+1)^{\frac{5}{2}} - \frac{2}{9} (x^3+1)^{\frac{3}{2}}$$

input `integrate(x^5*(x^3+1)^(1/2),x, algorithm="giac")`output `2/15*(x^3 + 1)^(5/2) - 2/9*(x^3 + 1)^(3/2)`

3.72.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int x^5 \sqrt{1+x^3} dx = \frac{2(x^3+1)^{5/2}}{15} - \frac{2(x^3+1)^{3/2}}{9}$$

input `int(x^5*(x^3 + 1)^(1/2),x)`

output `(2*(x^3 + 1)^(5/2))/15 - (2*(x^3 + 1)^(3/2))/9`

3.73 $\int \frac{-1+x^7}{\log(x)} dx$

3.73.1	Optimal result	469
3.73.2	Mathematica [A] (verified)	469
3.73.3	Rubi [A] (verified)	470
3.73.4	Maple [A] (verified)	471
3.73.5	Fricas [A] (verification not implemented)	471
3.73.6	Sympy [A] (verification not implemented)	471
3.73.7	Maxima [A] (verification not implemented)	472
3.73.8	Giac [A] (verification not implemented)	472
3.73.9	Mupad [B] (verification not implemented)	472

3.73.1 Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{-1+x^7}{\log(x)} dx = \text{ExpIntegralEi}(8 \log(x)) - \text{LogIntegral}(x)$$

output `Ei(8*ln(x))-Li(x)`

3.73.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{-1+x^7}{\log(x)} dx = \text{ExpIntegralEi}(8 \log(x)) - \text{LogIntegral}(x)$$

input `Integrate[(-1 + x^7)/Log[x],x]`

output `ExpIntegralEi[8*Log[x]] - LogIntegral[x]`

3.73.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2767, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^7 - 1}{\log(x)} dx$$

↓ 2767

$$\int \left(\frac{x^7}{\log(x)} - \frac{1}{\log(x)} \right) dx$$

↓ 2009

$$\text{ExpIntegralEi}(8 \log(x)) - \text{LogIntegral}(x)$$

input `Int[(-1 + x^7)/Log[x],x]`

output `ExpIntegralEi[8*Log[x]] - LogIntegral[x]`

3.73.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2767 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((d_) + (e_.)*(x_)^(r_.))^(q_.), x_Symbol] := With[{u = ExpandIntegrand[(a + b*Log[c*x^n])^p, (d + e*x^r)^q, x]}, Int[u, x] /; SumQ[u] /; FreeQ[{a, b, c, d, e, n, p, q, r}, x] && IntegerQ[q] && (GtQ[q, 0] || (IGtQ[p, 0] && IntegerQ[r]))]`

3.73.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.60

method	result	size
default	$-Ei_1(-8 \ln(x)) + Ei_1(-\ln(x))$	16
risch	$-Ei_1(-8 \ln(x)) + Ei_1(-\ln(x))$	16

input `int((x^7-1)/ln(x),x,method=_RETURNVERBOSE)`output `-Ei(1,-8*ln(x))+Ei(1,-ln(x))`**3.73.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{-1 + x^7}{\log(x)} dx = \log_integral(x^8) - \log_integral(x)$$

input `integrate((x^7-1)/log(x),x, algorithm="fracas")`output `log_integral(x^8) - log_integral(x)`**3.73.6 Sympy [A] (verification not implemented)**

Time = 0.56 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{-1 + x^7}{\log(x)} dx = -Ei(\log(x)) + Ei(8 \log(x))$$

input `integrate((x**7-1)/ln(x),x)`output `-Ei(log(x)) + Ei(8*log(x))`

3.73.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int \frac{-1 + x^7}{\log(x)} dx = \text{Ei}(8 \log(x)) - \text{Ei}(\log(x))$$

input `integrate((x^7-1)/log(x),x, algorithm="maxima")`output `Ei(8*log(x)) - Ei(log(x))`**3.73.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int \frac{-1 + x^7}{\log(x)} dx = \text{Ei}(8 \log(x)) - \text{Ei}(\log(x))$$

input `integrate((x^7-1)/log(x),x, algorithm="giac")`output `Ei(8*log(x)) - Ei(log(x))`**3.73.9 Mupad [B] (verification not implemented)**

Time = 18.82 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{-1 + x^7}{\log(x)} dx = \text{ei}(8 \ln(x)) - \text{logint}(x)$$

input `int((x^7 - 1)/log(x),x)`output `ei(8*log(x)) - logint(x)`

3.74 $\int \sqrt{\csc(x) - \sin(x)} dx$

3.74.1	Optimal result	473
3.74.2	Mathematica [A] (verified)	473
3.74.3	Rubi [A] (verified)	474
3.74.4	Maple [A] (verified)	475
3.74.5	Fricas [A] (verification not implemented)	476
3.74.6	Sympy [F]	476
3.74.7	Maxima [B] (verification not implemented)	476
3.74.8	Giac [F]	477
3.74.9	Mupad [B] (verification not implemented)	477

3.74.1 Optimal result

Integrand size = 11, antiderivative size = 13

$$\int \sqrt{\csc(x) - \sin(x)} dx = 2\sqrt{\cos(x) \cot(x)} \tan(x)$$

output `2*(cos(x)*cot(x))^(1/2)*tan(x)`

3.74.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \sqrt{\csc(x) - \sin(x)} dx = 2\sqrt{\cos(x) \cot(x)} \tan(x)$$

input `Integrate[Sqrt[Csc[x] - Sin[x]],x]`

output `2*Sqrt[Cos[x]*Cot[x]]*Tan[x]`

3.74.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 4897, 3042, 4900, 3042, 3069}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\csc(x) - \sin(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\csc(x) - \sin(x)} dx \\
 & \quad \downarrow \text{4897} \\
 & \int \sqrt{\cos(x) \cot(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\cos(x) \cot(x)} dx \\
 & \quad \downarrow \text{4900} \\
 & \frac{\sqrt{\cos(x) \cot(x)} \int \sqrt{\cos(x)} \sqrt{\cot(x)} dx}{\sqrt{\cos(x)} \sqrt{\cot(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\cos(x) \cot(x)} \int \sqrt{\sin(x + \frac{\pi}{2})} \sqrt{-\tan(x + \frac{\pi}{2})} dx}{\sqrt{\cos(x)} \sqrt{\cot(x)}} \\
 & \quad \downarrow \text{3069} \\
 & 2 \tan(x) \sqrt{\cos(x) \cot(x)}
 \end{aligned}$$

input `Int[Sqrt[Csc[x] - Sin[x]],x]`

output `2*Sqrt[Cos[x]*Cot[x]]*Tan[x]`

3.74.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3069 `Int[((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*(a*Sin[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*m)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 1, 0]`

rule 4897 `Int[u_, x_Symbol] := Int[TrigSimplify[u], x] /; TrigSimplifyQ[u]`

rule 4900 `Int[(u_.)*((v_)^(m_.)*(w_)^(n_.))^p, x_Symbol] := With[{uu = ActivateTrig[u], vv = ActivateTrig[v], ww = ActivateTrig[w]}, Simp[(vv^m*ww^n)^FracPart[p]/(vv^(m*FracPart[p])*ww^(n*FracPart[p])) Int[uu*vv^(m*p)*ww^(n*p), x], x] /; FreeQ[{m, n, p}, x] && !IntegerQ[p] && (!InertTrigFreeQ[v] || !InertTrigFreeQ[w])`

3.74.4 Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
default	$2\sqrt{\cos(x)\cot(x)}\tan(x)$	12
risch	$-\frac{i\sqrt{2}\sqrt{\frac{i(e^{2ix}+1)^2e^{-ix}}{e^{2ix}-1}}(e^{2ix}-1)}{e^{2ix}+1}$	51

input `int((csc(x)-sin(x))^(1/2),x,method=_RETURNVERBOSE)`

output `2*(cos(x)*cot(x))^(1/2)*tan(x)`

3.74.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.46

$$\int \sqrt{\csc(x) - \sin(x)} dx = \frac{2 \sqrt{\frac{\cos(x)^2}{\sin(x)}} \sin(x)}{\cos(x)}$$

input `integrate((csc(x)-sin(x))^(1/2),x, algorithm="fracas")`

output `2*sqrt(cos(x)^2/sin(x))*sin(x)/cos(x)`

3.74.6 Sympy [F]

$$\int \sqrt{\csc(x) - \sin(x)} dx = \int \sqrt{-\sin(x) + \csc(x)} dx$$

input `integrate((csc(x)-sin(x))**(1/2),x)`

output `Integral(sqrt(-sin(x) + csc(x)), x)`

3.74.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 188 vs. 2(11) = 22.

Time = 0.36 (sec) , antiderivative size = 188, normalized size of antiderivative = 14.46

$$\int \sqrt{\csc(x) - \sin(x)} dx = \frac{((\cos(\frac{3}{2}x) - \cos(\frac{1}{2}x) + \sin(\frac{3}{2}x) + \sin(\frac{1}{2}x)) \cos(\frac{1}{2} \arctan(\sin(x), \cos(x) - 1)) - (\cos(\frac{3}{2}x) - \cos(\frac{1}{2}x)))}{\cos(x)}$$

input `integrate((csc(x)-sin(x))^(1/2),x, algorithm="maxima")`

```
output (((cos(3/2*x) - cos(1/2*x) + sin(3/2*x) + sin(1/2*x))*cos(1/2*arctan2(sin(x), cos(x) - 1)) - (cos(3/2*x) - cos(1/2*x) - sin(3/2*x) - sin(1/2*x))*sin(1/2*arctan2(sin(x), cos(x) - 1)))*cos(1/2*arctan2(sin(x), cos(x) + 1)) - ((cos(3/2*x) - cos(1/2*x) - sin(3/2*x) - sin(1/2*x))*cos(1/2*arctan2(sin(x), cos(x) - 1)) + (cos(3/2*x) - cos(1/2*x) + sin(3/2*x) + sin(1/2*x))*sin(1/2*arctan2(sin(x), cos(x) - 1)))*sin(1/2*arctan2(sin(x), cos(x) + 1)))/((cos(x)^2 + sin(x)^2 + 2*cos(x) + 1)^(1/4)*(cos(x)^2 + sin(x)^2 - 2*cos(x) + 1)^(1/4))
```

3.74.8 Giac [F]

$$\int \sqrt{\csc(x) - \sin(x)} dx = \int \sqrt{\csc(x) - \sin(x)} dx$$

```
input integrate((csc(x)-sin(x))^(1/2),x, algorithm="giac")
```

```
output integrate(sqrt(csc(x) - sin(x)), x)
```

3.74.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \sqrt{\csc(x) - \sin(x)} dx = \frac{2 |\cos(x)|}{\cos(x) \sqrt{\frac{1}{\sin(x)}}}$$

```
input int((1/sin(x) - sin(x))^(1/2),x)
```

```
output (2*abs(cos(x)))/(cos(x)*(1/sin(x))^(1/2))
```

3.75 $\int (-2 \log(2x) + \log(x^2)) dx$

3.75.1	Optimal result	478
3.75.2	Mathematica [A] (verified)	478
3.75.3	Rubi [A] (verified)	479
3.75.4	Maple [A] (verified)	479
3.75.5	Fricas [A] (verification not implemented)	480
3.75.6	Sympy [A] (verification not implemented)	480
3.75.7	Maxima [A] (verification not implemented)	480
3.75.8	Giac [A] (verification not implemented)	481
3.75.9	Mupad [B] (verification not implemented)	481

3.75.1 Optimal result

Integrand size = 11, antiderivative size = 14

$$\int (-2 \log(2x) + \log(x^2)) dx = -2x \log(2x) + x \log(x^2)$$

output `-2*x*ln(2*x)+x*ln(x^2)`

3.75.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int (-2 \log(2x) + \log(x^2)) dx = -2x \log(2x) + x \log(x^2)$$

input `Integrate[-2*Log[2*x] + Log[x^2],x]`

output `-2*x*Log[2*x] + x*Log[x^2]`

3.75.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\log(x^2) - 2\log(2x)) dx$$

$$\downarrow \text{2009}$$

$$x \log(x^2) - 2x \log(2x)$$

input `Int[-2*Log[2*x] + Log[x^2],x]`

output `-2*x*Log[2*x] + x*Log[x^2]`

3.75.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.75.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

method	result	size
default	$-2x \ln(2x) + x \ln(x^2)$	15
norman	$-2x \ln(2x) + x \ln(x^2)$	15
risch	$-2x \ln(2x) + x \ln(x^2)$	15
parallelrisch	$-2x \ln(2x) + x \ln(x^2)$	15
parts	$-2x \ln(2x) + x \ln(x^2)$	15

input `int(ln(x^2)-2*ln(2*x),x,method=_RETURNVERBOSE)`

output `-2*x*ln(2*x)+x*ln(x^2)`

3.75. $\int (-2\log(2x) + \log(x^2)) dx$

3.75.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.36

$$\int (-2 \log(2x) + \log(x^2)) dx = -2x \log(2)$$

input `integrate(log(x^2)-2*log(2*x),x, algorithm="fricas")`output `-2*x*log(2)`**3.75.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.50

$$\int (-2 \log(2x) + \log(x^2)) dx = -2x \log(2)$$

input `integrate(ln(x**2)-2*ln(2*x),x)`output `-2*x*log(2)`**3.75.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int (-2 \log(2x) + \log(x^2)) dx = x \log(x^2) - 2x \log(2x)$$

input `integrate(log(x^2)-2*log(2*x),x, algorithm="maxima")`output `x*log(x^2) - 2*x*log(2*x)`

3.75.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int (-2 \log(2x) + \log(x^2)) dx = x \log(x^2) - 2x \log(2x)$$

input `integrate(log(x^2)-2*log(2*x),x, algorithm="giac")`output `x*log(x^2) - 2*x*log(2*x)`**3.75.9 Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int (-2 \log(2x) + \log(x^2)) dx = -x (2 \ln(2x) - \ln(x^2))$$

input `int(log(x^2) - 2*log(2*x),x)`output `-x*(2*log(2*x) - log(x^2))`

3.76 $\int e^x dx$

3.76.1	Optimal result	482
3.76.2	Mathematica [A] (verified)	482
3.76.3	Rubi [A] (verified)	483
3.76.4	Maple [A] (verified)	483
3.76.5	Fricas [A] (verification not implemented)	484
3.76.6	Sympy [A] (verification not implemented)	484
3.76.7	Maxima [A] (verification not implemented)	484
3.76.8	Giac [A] (verification not implemented)	485
3.76.9	Mupad [B] (verification not implemented)	485

3.76.1 Optimal result

Integrand size = 3, antiderivative size = 3

$$\int e^x dx = e^x$$

output `exp(x)`

3.76.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int e^x dx = e^x$$

input `Integrate[E^x,x]`

output `E^x`

3.76.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x dx$$

$$\downarrow 2624$$

$$e^x$$

input `Int [E^x,x]`

output `E^x`

3.76.3.1 Defintions of rubi rules used

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;`
`FreeQ[{F, n}, x] && LinearQ[v, x]`

3.76.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

method	result	size
gospers	e^x	3
lookup	e^x	3
derivativedivides	e^x	3
default	e^x	3
norman	e^x	3
risch	e^x	3
parallelrisch	e^x	3
meijerg	$e^x - 1$	5

input `int(exp(x),x,method=_RETURNVERBOSE)`

output `exp(x)`

3.76.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int e^x dx = e^x$$

input `integrate(exp(x),x, algorithm="fricas")`

output `e^x`

3.76.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int e^x dx = e^x$$

input `integrate(exp(x),x)`

output `exp(x)`

3.76.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int e^x dx = e^x$$

input `integrate(exp(x),x, algorithm="maxima")`

output `e^x`

3.76.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int e^x dx = e^x$$

input `integrate(exp(x),x, algorithm="giac")`

output `e^x`

3.76.9 Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int e^x dx = e^x$$

input `int(exp(x),x)`

output `exp(x)`

3.77
$$\int \frac{\cos(x) \log(x) - \frac{\sin(x)}{x}}{\log^2(x)} dx$$

3.77.1	Optimal result	486
3.77.2	Mathematica [A] (verified)	486
3.77.3	Rubi [F]	487
3.77.4	Maple [A] (verified)	487
3.77.5	Fricas [A] (verification not implemented)	488
3.77.6	Sympy [A] (verification not implemented)	488
3.77.7	Maxima [A] (verification not implemented)	488
3.77.8	Giac [B] (verification not implemented)	489
3.77.9	Mupad [B] (verification not implemented)	489

3.77.1 Optimal result

Integrand size = 18, antiderivative size = 7

$$\int \frac{\cos(x) \log(x) - \frac{\sin(x)}{x}}{\log^2(x)} dx = \frac{\sin(x)}{\log(x)}$$

output `sin(x)/ln(x)`

3.77.2 Mathematica [A] (verified)

Time = 0.89 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x) \log(x) - \frac{\sin(x)}{x}}{\log^2(x)} dx = \frac{\sin(x)}{\log(x)}$$

input `Integrate[(Cos[x]*Log[x] - Sin[x]/x)/Log[x]^2,x]`

output `Sin[x]/Log[x]`

3.77.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\log(x) \cos(x) - \frac{\sin(x)}{x}}{\log^2(x)} dx$$

↓ 7293

$$\int \left(\frac{\cos(x)}{\log(x)} - \frac{\sin(x)}{x \log^2(x)} \right) dx$$

↓ 2009

$$\int \frac{\cos(x)}{\log(x)} dx - \int \frac{\sin(x)}{x \log^2(x)} dx$$

input `Int[(Cos[x]*Log[x] - Sin[x]/x)/Log[x]^2,x]`

output `$Aborted`

3.77.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

3.77.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

method	result	size
risch	$\frac{\sin(x)}{\ln(x)}$	8
parallelrisc	$\frac{\sin(x)}{\ln(x)}$	8
norman	$\frac{2 \tan(\frac{x}{2})}{(1 + \tan(\frac{x}{2})^2) \ln(x)}$	21

3.77. $\int \frac{\cos(x) \log(x) - \frac{\sin(x)}{x}}{\log^2(x)} dx$

input `int((cos(x)*ln(x)-sin(x)/x)/ln(x)^2,x,method=_RETURNVERBOSE)`

output `sin(x)/ln(x)`

3.77.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x) \log(x) - \frac{\sin(x)}{x}}{\log^2(x)} dx = \frac{\sin(x)}{\log(x)}$$

input `integrate((cos(x)*log(x)-sin(x)/x)/log(x)^2,x, algorithm="fricas")`

output `sin(x)/log(x)`

3.77.6 Sympy [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71

$$\int \frac{\cos(x) \log(x) - \frac{\sin(x)}{x}}{\log^2(x)} dx = \frac{\sin(x)}{\log(x)}$$

input `integrate((cos(x)*ln(x)-sin(x)/x)/ln(x)**2,x)`

output `sin(x)/log(x)`

3.77.7 Maxima [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x) \log(x) - \frac{\sin(x)}{x}}{\log^2(x)} dx = \frac{\sin(x)}{\log(x)}$$

input `integrate((cos(x)*log(x)-sin(x)/x)/log(x)^2,x, algorithm="maxima")`

output `sin(x)/log(x)`

3.77. $\int \frac{\cos(x) \log(x) - \frac{\sin(x)}{x}}{\log^2(x)} dx$

3.77.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. $2(7) = 14$.

Time = 0.28 (sec) , antiderivative size = 20, normalized size of antiderivative = 2.86

$$\int \frac{\cos(x) \log(x) - \frac{\sin(x)}{x}}{\log^2(x)} dx = \frac{2 \tan\left(\frac{1}{2}x\right)}{\log(x) \tan\left(\frac{1}{2}x\right)^2 + \log(x)}$$

input `integrate((cos(x)*log(x)-sin(x)/x)/log(x)^2,x, algorithm="giac")`

output `2*tan(1/2*x)/(log(x)*tan(1/2*x)^2 + log(x))`

3.77.9 Mupad [B] (verification not implemented)

Time = 16.80 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x) \log(x) - \frac{\sin(x)}{x}}{\log^2(x)} dx = \frac{\sin(x)}{\ln(x)}$$

input `int((cos(x)*log(x) - sin(x)/x)/log(x)^2,x)`

output `sin(x)/log(x)`

3.78 $\int (-1 + 3x - 3x^2 + x^3) dx$

3.78.1	Optimal result	490
3.78.2	Mathematica [B] (verified)	490
3.78.3	Rubi [A] (verified)	491
3.78.4	Maple [A] (verified)	492
3.78.5	Fricas [B] (verification not implemented)	492
3.78.6	Sympy [B] (verification not implemented)	493
3.78.7	Maxima [B] (verification not implemented)	493
3.78.8	Giac [B] (verification not implemented)	493
3.78.9	Mupad [B] (verification not implemented)	494

3.78.1 Optimal result

Integrand size = 13, antiderivative size = 11

$$\int (-1 + 3x - 3x^2 + x^3) dx = \frac{1}{4}(1 - x)^4$$

output `1/4*(1-x)^4`

3.78.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 23 vs. 2(11) = 22.

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 2.09

$$\int (-1 + 3x - 3x^2 + x^3) dx = -x + \frac{3x^2}{2} - x^3 + \frac{x^4}{4}$$

input `Integrate[-1 + 3*x - 3*x^2 + x^3,x]`

output `-x + (3*x^2)/2 - x^3 + x^4/4`

3.78.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {2006, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x^3 - 3x^2 + 3x - 1) dx$$

$$\downarrow \text{2006}$$

$$\int (x - 1)^3 dx$$

$$\downarrow \text{17}$$

$$\frac{1}{4}(1 - x)^4$$

input `Int[-1 + 3*x - 3*x^2 + x^3,x]`

output `(1 - x)^4/4`

3.78.3.1 Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] :> Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 2006 `Int[(u_.)*(Px_), x_Symbol] :> With[{a = Rt[Coeff[Px, x, 0], Expon[Px, x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Int[u*(a + b*x)^Expon[Px, x], x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; PolyQ[Px, x] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0] && !MatchQ[Px, (a_.)*(v_)^Expon[Px, x]] /; FreeQ[a, x] && LinearQ[v, x]]`

3.78.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

method	result	size
default	$\frac{(-1+x)^4}{4}$	8
gospers	$\frac{x(x^3-4x^2+6x-4)}{4}$	17
norman	$\frac{1}{4}x^4 - x^3 + \frac{3}{2}x^2 - x$	20
risch	$\frac{1}{4}x^4 - x^3 + \frac{3}{2}x^2 - x$	20
paralrelrisch	$\frac{1}{4}x^4 - x^3 + \frac{3}{2}x^2 - x$	20
parts	$\frac{1}{4}x^4 - x^3 + \frac{3}{2}x^2 - x$	20

input `int(x^3-3*x^2+3*x-1,x,method=_RETURNVERBOSE)`

output `1/4*(-1+x)^4`

3.78.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. 2(7) = 14.

Time = 0.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.73

$$\int (-1 + 3x - 3x^2 + x^3) dx = \frac{1}{4}x^4 - x^3 + \frac{3}{2}x^2 - x$$

input `integrate(x^3-3*x^2+3*x-1,x, algorithm="fracas")`

output `1/4*x^4 - x^3 + 3/2*x^2 - x`

3.78.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(5) = 10$.

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.36

$$\int (-1 + 3x - 3x^2 + x^3) dx = \frac{x^4}{4} - x^3 + \frac{3x^2}{2} - x$$

input `integrate(x**3-3*x**2+3*x-1,x)`

output `x**4/4 - x**3 + 3*x**2/2 - x`

3.78.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(7) = 14$.

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.73

$$\int (-1 + 3x - 3x^2 + x^3) dx = \frac{1}{4}x^4 - x^3 + \frac{3}{2}x^2 - x$$

input `integrate(x^3-3*x^2+3*x-1,x, algorithm="maxima")`

output `1/4*x^4 - x^3 + 3/2*x^2 - x`

3.78.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(7) = 14$.

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.73

$$\int (-1 + 3x - 3x^2 + x^3) dx = \frac{1}{4}x^4 - x^3 + \frac{3}{2}x^2 - x$$

input `integrate(x^3-3*x^2+3*x-1,x, algorithm="giac")`

output `1/4*x^4 - x^3 + 3/2*x^2 - x`

3.78. $\int (-1 + 3x - 3x^2 + x^3) dx$

3.78.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.73

$$\int (-1 + 3x - 3x^2 + x^3) dx = \frac{x^4}{4} - x^3 + \frac{3x^2}{2} - x$$

input `int(3*x - 3*x^2 + x^3 - 1,x)`

output `(3*x^2)/2 - x - x^3 + x^4/4`

3.79 $\int \sqrt{12 - 3x^2} dx$

3.79.1	Optimal result	495
3.79.2	Mathematica [A] (verified)	495
3.79.3	Rubi [A] (verified)	496
3.79.4	Maple [A] (verified)	497
3.79.5	Fricas [A] (verification not implemented)	497
3.79.6	Sympy [A] (verification not implemented)	498
3.79.7	Maxima [A] (verification not implemented)	498
3.79.8	Giac [A] (verification not implemented)	498
3.79.9	Mupad [B] (verification not implemented)	499

3.79.1 Optimal result

Integrand size = 11, antiderivative size = 35

$$\int \sqrt{12 - 3x^2} dx = \frac{1}{2}\sqrt{3}x\sqrt{4 - x^2} + 2\sqrt{3} \arcsin\left(\frac{x}{2}\right)$$

output `1/2*3^(1/2)*x*(-x^2+4)^(1/2)+2*3^(1/2)*arcsin(1/2*x)`

3.79.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.23

$$\int \sqrt{12 - 3x^2} dx = \frac{1}{2}\sqrt{3}\left(x\sqrt{4 - x^2} - 8 \arctan\left(\frac{\sqrt{4 - x^2}}{2 + x}\right)\right)$$

input `Integrate[Sqrt[12 - 3*x^2],x]`

output `(Sqrt[3]*(x*Sqrt[4 - x^2] - 8*ArcTan[Sqrt[4 - x^2]/(2 + x)]))/2`

3.79.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {211, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{12 - 3x^2} dx$$

$$\downarrow \text{211}$$

$$6 \int \frac{1}{\sqrt{12 - 3x^2}} dx + \frac{1}{2} \sqrt{3} \sqrt{4 - x^2} x$$

$$\downarrow \text{223}$$

$$2\sqrt{3} \arcsin\left(\frac{x}{2}\right) + \frac{1}{2} \sqrt{3} \sqrt{4 - x^2} x$$

input `Int[Sqrt[12 - 3*x^2], x]`

output `(Sqrt[3]*x*Sqrt[4 - x^2])/2 + 2*Sqrt[3]*ArcSin[x/2]`

3.79.3.1 Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

3.79.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.66

method	result	size
default	$\frac{x\sqrt{-3x^2+12}}{2} + 2\sqrt{3} \arcsin\left(\frac{x}{2}\right)$	23
risch	$-\frac{3x(x^2-4)}{2\sqrt{-3x^2+12}} + 2\sqrt{3} \arcsin\left(\frac{x}{2}\right)$	28
meijerg	$\frac{i\sqrt{3} \left(-i\sqrt{\pi} x \sqrt{-\frac{x^2}{4}+1} - 2i\sqrt{\pi} \arcsin\left(\frac{x}{2}\right) \right)}{\sqrt{\pi}}$	37
pseudoelliptic	$\frac{x\sqrt{-3x^2+12}}{2} - 2\sqrt{3} \arctan\left(\frac{\sqrt{-3x^2+12}\sqrt{3}}{3x}\right)$	37
trager	$\frac{x\sqrt{-3x^2+12}}{2} + 2 \operatorname{RootOf}(_Z^2 + 3) \ln(\operatorname{RootOf}(_Z^2 + 3) \sqrt{-3x^2 + 12} + 3x)$	43

input `int((-3*x^2+12)^(1/2),x,method=_RETURNVERBOSE)`output `1/2*x*(-3*x^2+12)^(1/2)+2*3^(1/2)*arcsin(1/2*x)`**3.79.5 Fricas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.03

$$\int \sqrt{12 - 3x^2} dx = \frac{1}{2} \sqrt{-3x^2 + 12}x - 2\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt{-3x^2 + 12}}{3x}\right)$$

input `integrate((-3*x^2+12)^(1/2),x, algorithm="fricas")`output `1/2*sqrt(-3*x^2 + 12)*x - 2*sqrt(3)*arctan(1/3*sqrt(3)*sqrt(-3*x^2 + 12)/x)`

3.79.6 Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.77

$$\int \sqrt{12 - 3x^2} dx = \frac{\sqrt{3}x\sqrt{4 - x^2}}{2} + 2\sqrt{3} \operatorname{asin}\left(\frac{x}{2}\right)$$

input `integrate((-3*x**2+12)**(1/2),x)`output `sqrt(3)*x*sqrt(4 - x**2)/2 + 2*sqrt(3)*asin(x/2)`**3.79.7 Maxima [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.63

$$\int \sqrt{12 - 3x^2} dx = \frac{1}{2} \sqrt{-3x^2 + 12}x + 2\sqrt{3} \arcsin\left(\frac{1}{2}x\right)$$

input `integrate((-3*x^2+12)^(1/2),x, algorithm="maxima")`output `1/2*sqrt(-3*x^2 + 12)*x + 2*sqrt(3)*arcsin(1/2*x)`**3.79.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.66

$$\int \sqrt{12 - 3x^2} dx = \frac{1}{2} \sqrt{3} \left(\sqrt{-x^2 + 4}x + 4 \arcsin\left(\frac{1}{2}x\right) \right)$$

input `integrate((-3*x^2+12)^(1/2),x, algorithm="giac")`output `1/2*sqrt(3)*(sqrt(-x^2 + 4)*x + 4*arcsin(1/2*x))`

3.79.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int \sqrt{12 - 3x^2} dx = 2\sqrt{3} \operatorname{asin}\left(\frac{x}{2}\right) + \frac{\sqrt{3}x\sqrt{4 - x^2}}{2}$$

input `int((12 - 3*x^2)^(1/2),x)`

output `2*3^(1/2)*asin(x/2) + (3^(1/2)*x*(4 - x^2)^(1/2))/2`

3.80 $\int ((-3 + x)^7 + x - \sin(3 - x)) dx$

3.80.1	Optimal result	500
3.80.2	Mathematica [A] (verified)	500
3.80.3	Rubi [A] (verified)	501
3.80.4	Maple [A] (verified)	501
3.80.5	Fricas [B] (verification not implemented)	502
3.80.6	Sympy [A] (verification not implemented)	502
3.80.7	Maxima [A] (verification not implemented)	503
3.80.8	Giac [A] (verification not implemented)	503
3.80.9	Mupad [B] (verification not implemented)	503

3.80.1 Optimal result

Integrand size = 15, antiderivative size = 27

$$\int ((-3 + x)^7 + x - \sin(3 - x)) dx = \frac{1}{8}(3 - x)^8 + \frac{x^2}{2} - \cos(3 - x)$$

output `1/8*(-x+3)^8+1/2*x^2-cos(-3+x)`

3.80.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07

$$\int ((-3 + x)^7 + x - \sin(3 - x)) dx = \frac{1}{8}(-3 + x)^8 + \frac{x^2}{2} - \cos(3) \cos(x) - \sin(3) \sin(x)$$

input `Integrate[(-3 + x)^7 + x - Sin[3 - x], x]`

output `(-3 + x)^8/8 + x^2/2 - Cos[3]*Cos[x] - Sin[3]*Sin[x]`

3.80.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int ((x - 3)^7 + x - \sin(3 - x)) dx$$

↓ 2009

$$\frac{x^2}{2} + \frac{1}{8}(3 - x)^8 - \cos(3 - x)$$

input `Int[(-3 + x)^7 + x - Sin[3 - x],x]`

output `(3 - x)^8/8 + x^2/2 - Cos[3 - x]`

3.80.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.80.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result
default	$\frac{x^2}{2} + \frac{(-3+x)^8}{8} - \cos(-3+x)$
derivativedivides	$-9 + 3x + \frac{(-3+x)^2}{2} + \frac{(-3+x)^8}{8} - \cos(-3+x)$
parts	$-2187x + 2552x^2 - 1701x^3 + \frac{2835x^4}{4} - 189x^5 + \frac{63x^6}{2} - 3x^7 + \frac{x^8}{8} - \cos(-3+x)$
risch	$2552x^2 + \frac{x^8}{8} - 3x^7 + \frac{63x^6}{2} - 189x^5 + \frac{2835x^4}{4} - 1701x^3 - 2187x + \frac{6561}{8} - \cos(-3+x)$
parallelrisch	$\frac{x^8}{8} - 3x^7 + \frac{63x^6}{2} - 189x^5 + \frac{2835x^4}{4} - 1701x^3 + 2552x^2 - 2187x - 1 - \cos(-3+x)$
norman	$\frac{-2187x+2552x^2-1701x^3+\frac{2835x^4}{4}-189x^5+\frac{63x^6}{2}-3x^7+\frac{x^8}{8}-2187x \tan(-\frac{3}{2}+\frac{x}{2})^2+2552x^2 \tan(-\frac{3}{2}+\frac{x}{2})^2-1701x^3 \tan(-\frac{3}{2}+\frac{x}{2})^2-1}{1+\tan(-\frac{3}{2}+\frac{x}{2})}$

3.80. $\int ((-3 + x)^7 + x - \sin(3 - x)) dx$

input `int(x+(-3+x)^7+sin(-3+x),x,method=_RETURNVERBOSE)`

output `1/2*x^2+1/8*(-3+x)^8-cos(-3+x)`

3.80.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 45 vs. $2(19) = 38$.

Time = 0.26 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.67

$$\int ((-3+x)^7 + x - \sin(3-x)) dx = \frac{1}{8}x^8 - 3x^7 + \frac{63}{2}x^6 - 189x^5 + \frac{2835}{4}x^4 - 1701x^3 + 2552x^2 - 2187x - \cos(x-3)$$

input `integrate(x+(-3+x)^7+sin(-3+x),x, algorithm="fricas")`

output `1/8*x^8 - 3*x^7 + 63/2*x^6 - 189*x^5 + 2835/4*x^4 - 1701*x^3 + 2552*x^2 - 2187*x - cos(x - 3)`

3.80.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.56

$$\int ((-3+x)^7 + x - \sin(3-x)) dx = \frac{x^2}{2} + \frac{(x-3)^8}{8} - \cos(x-3)$$

input `integrate(x+(-3+x)**7+sin(-3+x),x)`

output `x**2/2 + (x - 3)**8/8 - cos(x - 3)`

3.80.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int ((-3+x)^7 + x - \sin(3-x)) dx = \frac{1}{8}(x-3)^8 + \frac{1}{2}x^2 - \cos(x-3)$$

input `integrate(x+(-3+x)^7+sin(-3+x),x, algorithm="maxima")`output `1/8*(x - 3)^8 + 1/2*x^2 - cos(x - 3)`**3.80.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int ((-3+x)^7 + x - \sin(3-x)) dx = \frac{1}{8}(x-3)^8 + \frac{1}{2}x^2 - \cos(x-3)$$

input `integrate(x+(-3+x)^7+sin(-3+x),x, algorithm="giac")`output `1/8*(x - 3)^8 + 1/2*x^2 - cos(x - 3)`**3.80.9 Mupad [B] (verification not implemented)**

Time = 16.24 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.67

$$\int ((-3+x)^7 + x - \sin(3-x)) dx = 2552x^2 - \cos(x-3) - 2187x - 1701x^3 + \frac{2835x^4}{4} - 189x^5 + \frac{63x^6}{2} - 3x^7 + \frac{x^8}{8}$$

input `int(x + sin(x - 3) + (x - 3)^7,x)`output `2552*x^2 - cos(x - 3) - 2187*x - 1701*x^3 + (2835*x^4)/4 - 189*x^5 + (63*x^6)/2 - 3*x^7 + x^8/8`

3.81 $\int \sin(x) \sqrt{1 + \tan^2(x)} dx$

3.81.1	Optimal result	504
3.81.2	Mathematica [A] (verified)	504
3.81.3	Rubi [A] (verified)	505
3.81.4	Maple [C] (warning: unable to verify)	506
3.81.5	Fricas [A] (verification not implemented)	507
3.81.6	Sympy [F]	507
3.81.7	Maxima [A] (verification not implemented)	507
3.81.8	Giac [A] (verification not implemented)	508
3.81.9	Mupad [F(-1)]	508

3.81.1 Optimal result

Integrand size = 13, antiderivative size = 15

$$\int \sin(x) \sqrt{1 + \tan^2(x)} dx = -\cos(x) \log(\cos(x)) \sqrt{\sec^2(x)}$$

output `-cos(x)*ln(cos(x))*(sec(x)^2)^(1/2)`

3.81.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \sin(x) \sqrt{1 + \tan^2(x)} dx = -\cos(x) \log(\cos(x)) \sqrt{\sec^2(x)}$$

input `Integrate[Sin[x]*Sqrt[1 + Tan[x]^2],x]`

output `-(Cos[x]*Log[Cos[x]]*Sqrt[Sec[x]^2])`

3.81.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {3042, 4140, 3042, 4613, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(x) \sqrt{\tan^2(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x) \sqrt{\tan(x)^2 + 1} dx \\
 & \quad \downarrow \text{4140} \\
 & \int \sin(x) \sqrt{\sec^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x) \sqrt{\sec(x)^2} dx \\
 & \quad \downarrow \text{4613} \\
 & \cos(x) \sqrt{\sec^2(x)} \int \tan(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \cos(x) \sqrt{\sec^2(x)} \int \tan(x) dx \\
 & \quad \downarrow \text{3956} \\
 & -\cos(x) \sqrt{\sec^2(x)} \log(\cos(x))
 \end{aligned}$$

input `Int[Sin[x]*Sqrt[1 + Tan[x]^2], x]`

output `-(Cos[x]*Log[Cos[x]]*Sqrt[Sec[x]^2])`

3.81.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4140 `Int[(u_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^p, x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]`

rule 4613 `Int[(u_.)*((b_.)*sec[(e_.) + (f_.)*(x_)]^n)^p, x_Symbol] := With[{ff = FreeFactors[Sec[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Sec[e + f*x]^n)^FracPart[p]/(Sec[e + f*x]/ff)^(n*FracPart[p])) Int[ActivateTrig[u]*(Sec[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

3.81.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.37 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.40

method	result	size
default	$-\text{csgn}(\sec(x)) \left(\ln(1 + \csc(x) - \cot(x)) + \ln(\csc(x) - \cot(x) - 1) - \ln\left(\frac{2}{1 + \cos(x)}\right) \right)$	36
risch	$2i \sqrt{\frac{e^{2ix}}{(e^{2ix} + 1)^2}} x \cos(x) - 2 \sqrt{\frac{e^{2ix}}{(e^{2ix} + 1)^2}} \ln(e^{2ix} + 1) \cos(x)$	54

input `int(sin(x)*(1+tan(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

output `-csgn(sec(x))*(ln(1+csc(x)-cot(x))+ln(csc(x)-cot(x)-1)-ln(2/(1+cos(x))))`

3.81.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.33

$$\int \sin(x) \sqrt{1 + \tan^2(x)} dx = \log(-\cos(x))$$

input `integrate(sin(x)*(1+tan(x)^2)^(1/2),x, algorithm="fracas")`output `log(-cos(x))`**3.81.6 Sympy [F]**

$$\int \sin(x) \sqrt{1 + \tan^2(x)} dx = \int \sqrt{\tan^2(x) + 1} \sin(x) dx$$

input `integrate(sin(x)*(1+tan(x)**2)**(1/2),x)`output `Integral(sqrt(tan(x)**2 + 1)*sin(x), x)`**3.81.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \sin(x) \sqrt{1 + \tan^2(x)} dx = -\sqrt{\frac{1}{\cos(x)^2}} \cos(x) \log(\cos(x))$$

input `integrate(sin(x)*(1+tan(x)^2)^(1/2),x, algorithm="maxima")`output `-sqrt(cos(x)^(-2))*cos(x)*log(cos(x))`

3.81.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.40

$$\int \sin(x) \sqrt{1 + \tan^2(x)} dx = -\log(|\cos(x)|)$$

input `integrate(sin(x)*(1+tan(x)^2)^(1/2),x, algorithm="giac")`

output `-log(abs(cos(x)))`

3.81.9 Mupad [F(-1)]

Timed out.

$$\int \sin(x) \sqrt{1 + \tan^2(x)} dx = \int \sin(x) \sqrt{\tan(x)^2 + 1} dx$$

input `int(sin(x)*(tan(x)^2 + 1)^(1/2),x)`

output `int(sin(x)*(tan(x)^2 + 1)^(1/2), x)`

3.82 $\int \frac{-1+x^2-x^3+x^5}{-1+x-x^3+x^4} dx$

3.82.1	Optimal result	509
3.82.2	Mathematica [A] (verified)	509
3.82.3	Rubi [A] (verified)	510
3.82.4	Maple [A] (verified)	511
3.82.5	Fricas [A] (verification not implemented)	511
3.82.6	Sympy [A] (verification not implemented)	511
3.82.7	Maxima [A] (verification not implemented)	512
3.82.8	Giac [A] (verification not implemented)	512
3.82.9	Mupad [B] (verification not implemented)	512

3.82.1 Optimal result

Integrand size = 27, antiderivative size = 9

$$\int \frac{-1+x^2-x^3+x^5}{-1+x-x^3+x^4} dx = \frac{1}{2}(1+x)^2$$

output `1/2*(1+x)^2`

3.82.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{-1+x^2-x^3+x^5}{-1+x-x^3+x^4} dx = x + \frac{x^2}{2}$$

input `Integrate[(-1 + x^2 - x^3 + x^5)/(-1 + x - x^3 + x^4),x]`

output `x + x^2/2`

3.82.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.074$, Rules used = {2019, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^5 - x^3 + x^2 - 1}{x^4 - x^3 + x - 1} dx$$

↓ 2019

$$\int (x + 1) dx$$

↓ 17

$$\frac{1}{2}(x + 1)^2$$

input `Int[(-1 + x^2 - x^3 + x^5)/(-1 + x - x^3 + x^4), x]`

output `(1 + x)^2/2`

3.82.3.1 Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 2019 `Int[(u_.)*(Px_)^(p_.)*(Qx_)^(q_.), x_Symbol] := Int[u*PolynomialQuotient[Px, Qx, x]^p*Qx^(p + q), x] /; FreeQ[q, x] && PolyQ[Px, x] && PolyQ[Qx, x] && EqQ[PolynomialRemainder[Px, Qx, x], 0] && IntegerQ[p] && LtQ[p*q, 0]`

3.82.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

method	result	size
gospers	$\frac{x(2+x)}{2}$	7
default	$\frac{1}{2}x^2 + x$	8
norman	$\frac{1}{2}x^2 + x$	8
risch	$\frac{1}{2}x^2 + x$	8
parallelrisch	$\frac{1}{2}x^2 + x$	8
parts	$\frac{1}{2}x^2 + x$	8

input `int((x^5-x^3+x^2-1)/(x^4-x^3+x-1),x,method=_RETURNVERBOSE)`output `1/2*x*(2+x)`**3.82.5 Fricas [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{-1 + x^2 - x^3 + x^5}{-1 + x - x^3 + x^4} dx = \frac{1}{2}x^2 + x$$

input `integrate((x^5-x^3+x^2-1)/(x^4-x^3+x-1),x, algorithm="fricas")`output `1/2*x^2 + x`**3.82.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.56

$$\int \frac{-1 + x^2 - x^3 + x^5}{-1 + x - x^3 + x^4} dx = \frac{x^2}{2} + x$$

input `integrate((x**5-x**3+x**2-1)/(x**4-x**3+x-1),x)`output `x**2/2 + x`

3.82. $\int \frac{-1+x^2-x^3+x^5}{-1+x-x^3+x^4} dx$

3.82.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{-1 + x^2 - x^3 + x^5}{-1 + x - x^3 + x^4} dx = \frac{1}{2}x^2 + x$$

input `integrate((x^5-x^3+x^2-1)/(x^4-x^3+x-1),x, algorithm="maxima")`output `1/2*x^2 + x`**3.82.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{-1 + x^2 - x^3 + x^5}{-1 + x - x^3 + x^4} dx = \frac{1}{2}x^2 + x$$

input `integrate((x^5-x^3+x^2-1)/(x^4-x^3+x-1),x, algorithm="giac")`output `1/2*x^2 + x`**3.82.9 Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.67

$$\int \frac{-1 + x^2 - x^3 + x^5}{-1 + x - x^3 + x^4} dx = \frac{x(x+2)}{2}$$

input `int((x^2 - x^3 + x^5 - 1)/(x - x^3 + x^4 - 1),x)`output `(x*(x + 2))/2`

3.83 $\int \log(x) dx$

3.83.1	Optimal result	513
3.83.2	Mathematica [A] (verified)	513
3.83.3	Rubi [A] (verified)	514
3.83.4	Maple [A] (verified)	514
3.83.5	Fricas [A] (verification not implemented)	515
3.83.6	Sympy [A] (verification not implemented)	515
3.83.7	Maxima [A] (verification not implemented)	515
3.83.8	Giac [A] (verification not implemented)	516
3.83.9	Mupad [B] (verification not implemented)	516

3.83.1 Optimal result

Integrand size = 2, antiderivative size = 8

$$\int \log(x) dx = -x + x \log(x)$$

output `x*ln(x)-x`

3.83.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \log(x) dx = -x + x \log(x)$$

input `Integrate[Log[x],x]`

output `-x + x*Log[x]`

3.83.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {2732}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \log(x) dx$$

$$\downarrow \text{2732}$$

$$x \log(x) - x$$

input `Int [Log[x] , x]`

output `-x + x*Log[x]`

3.83.3.1 Defintions of rubi rules used

rule 2732 `Int [Log[(c_.)*(x_)^(n_.)], x_Symbol] :> Simp[x*Log[c*x^n], x] - Simp[n*x, x] /; FreeQ[{c, n}, x]`

3.83.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
lookup	$x \ln(x) - x$	9
default	$x \ln(x) - x$	9
norman	$x \ln(x) - x$	9
risch	$x \ln(x) - x$	9
parallelrisch	$x \ln(x) - x$	9
parts	$x \ln(x) - x$	9

input `int(ln(x), x, method=_RETURNVERBOSE)`

output `x*ln(x)-x`

3.83.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \log(x) dx = x \log(x) - x$$

input `integrate(log(x),x, algorithm="fricas")`

output `x*log(x) - x`

3.83.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \log(x) dx = x \log(x) - x$$

input `integrate(ln(x),x)`

output `x*log(x) - x`

3.83.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \log(x) dx = x \log(x) - x$$

input `integrate(log(x),x, algorithm="maxima")`

output `x*log(x) - x`

3.83.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \log(x) dx = x \log(x) - x$$

input `integrate(log(x),x, algorithm="giac")`

output `x*log(x) - x`

3.83.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \log(x) dx = x (\ln(x) - 1)$$

input `int(log(x),x)`

output `x*(log(x) - 1)`

3.84 $\int \frac{1}{1-e^{-x}} dx$

3.84.1	Optimal result	517
3.84.2	Mathematica [A] (verified)	517
3.84.3	Rubi [B] (verified)	518
3.84.4	Maple [A] (verified)	519
3.84.5	Fricas [A] (verification not implemented)	519
3.84.6	Sympy [A] (verification not implemented)	520
3.84.7	Maxima [A] (verification not implemented)	520
3.84.8	Giac [A] (verification not implemented)	520
3.84.9	Mupad [B] (verification not implemented)	521

3.84.1 Optimal result

Integrand size = 11, antiderivative size = 6

$$\int \frac{1}{1-e^{-x}} dx = \log(-1 + e^x)$$

output `ln(-1+exp(x))`

3.84.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{1}{1-e^{-x}} dx = \log(-1 + e^x)$$

input `Integrate[(1 - E^(-x))^(-1),x]`

output `Log[-1 + E^x]`

3.84.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 19 vs. $2(6) = 12$.

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 3.17, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {2720, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{1 - e^{-x}} dx \\
 & \quad \downarrow 2720 \\
 & - \int \frac{e^x}{1 - e^{-x}} de^{-x} \\
 & \quad \downarrow 47 \\
 & - \int e^x de^{-x} - \int \frac{1}{1 - e^{-x}} de^{-x} \\
 & \quad \downarrow 14 \\
 & - \int \frac{1}{1 - e^{-x}} de^{-x} - \log(e^{-x}) \\
 & \quad \downarrow 16 \\
 & \log(1 - e^{-x}) - \log(e^{-x})
 \end{aligned}$$

input `Int[(1 - E^(-x))^(-1),x]`

output `-Log[E^(-x)] + Log[1 - E^(-x)]`

3.84.3.1 Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

```
rule 47 Int[1/((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.)), x_Symbol] := Simp[b/(b*c
- a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x
], x] /; FreeQ[{a, b, c, d}, x]
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.84.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.67

method	result	size
norman	$x + \ln(-1 + e^{-x})$	10
risch	$x + \ln(-1 + e^{-x})$	10
parallelrisch	$x + \ln(-1 + e^{-x})$	10
derivativedivides	$-\ln(e^{-x}) + \ln(-1 + e^{-x})$	16
default	$-\ln(e^{-x}) + \ln(-1 + e^{-x})$	16

```
input int(1/(1-exp(-x)),x,method=_RETURNVERBOSE)
```

```
output x+ln(-1+exp(-x))
```

3.84.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.50

$$\int \frac{1}{1 - e^{-x}} dx = x + \log(e^{(-x)} - 1)$$

```
input integrate(1/(1-exp(-x)),x, algorithm="fricas")
```

```
output x + log(e^(-x) - 1)
```


3.84.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.33

$$\int \frac{1}{1 - e^{-x}} dx = x + \log(-1 + e^{-x})$$

input `integrate(1/(1-exp(-x)),x)`output `x + log(-1 + exp(-x))`**3.84.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.50

$$\int \frac{1}{1 - e^{-x}} dx = x + \log(e^{(-x)} - 1)$$

input `integrate(1/(1-exp(-x)),x, algorithm="maxima")`output `x + log(e^(-x) - 1)`**3.84.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.67

$$\int \frac{1}{1 - e^{-x}} dx = x + \log(|e^{(-x)} - 1|)$$

input `integrate(1/(1-exp(-x)),x, algorithm="giac")`output `x + log(abs(e^(-x) - 1))`

3.84.9 Mupad [B] (verification not implemented)

Time = 15.70 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

$$\int \frac{1}{1 - e^{-x}} dx = \ln(1 - e^x)$$

input `int(-1/(exp(-x) - 1),x)`

output `log(1 - exp(x))`

3.85 $\int \cos^2(x) \sin^2(x) dx$

3.85.1	Optimal result	522
3.85.2	Mathematica [A] (verified)	522
3.85.3	Rubi [B] (verified)	523
3.85.4	Maple [A] (verified)	524
3.85.5	Fricas [A] (verification not implemented)	525
3.85.6	Sympy [A] (verification not implemented)	525
3.85.7	Maxima [A] (verification not implemented)	525
3.85.8	Giac [A] (verification not implemented)	526
3.85.9	Mupad [B] (verification not implemented)	526

3.85.1 Optimal result

Integrand size = 9, antiderivative size = 14

$$\int \cos^2(x) \sin^2(x) dx = \frac{1}{32}(4x - \sin(4x))$$

output `1/8*x-1/32*sin(4*x)`

3.85.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \cos^2(x) \sin^2(x) dx = \frac{x}{8} - \frac{1}{32} \sin(4x)$$

input `Integrate[Cos[x]^2*Sin[x]^2,x]`

output `x/8 - Sin[4*x]/32`

3.85.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 29 vs. $2(14) = 28$.

Time = 0.21 (sec) , antiderivative size = 29, normalized size of antiderivative = 2.07, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 3048, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^2(x) \cos^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x)^2 \cos(x)^2 dx \\
 & \quad \downarrow \text{3048} \\
 & \frac{1}{4} \int \cos^2(x) dx - \frac{1}{4} \sin(x) \cos^3(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4} \int \sin\left(x + \frac{\pi}{2}\right)^2 dx - \frac{1}{4} \sin(x) \cos^3(x) \\
 & \quad \downarrow \text{3115} \\
 & \frac{1}{4} \left(\int \frac{1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin(x) \cos^3(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{1}{4} \left(\frac{x}{2} + \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin(x) \cos^3(x)
 \end{aligned}$$

input `Int[Cos[x]^2*Sin[x]^2,x]`

output `-1/4*(Cos[x]^3*Sin[x]) + (x/2 + (Cos[x]*Sin[x])/2)/4`

3.85.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3048 `Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^n]*((a_.)*sin[(e_.) + (f_.)*(x_)])^m, x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Simp[a^2*((m - 1)/(m + n)) Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

rule 3115 `Int[(b_.)*sin[(c_.) + (d_.)*(x_)]^n, x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

3.85.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

method	result	size
risch	$\frac{x}{8} - \frac{\sin(4x)}{32}$	11
parallelrisch	$\frac{x}{8} - \frac{\sin(4x)}{32}$	11
default	$-\frac{\cos(x)^3 \sin(x)}{4} + \frac{\cos(x) \sin(x)}{8} + \frac{x}{8}$	19
norman	$\frac{\frac{x}{8} + \frac{7 \tan(\frac{x}{2})^3}{4} - \frac{7 \tan(\frac{x}{2})^5}{4} + \frac{\tan(\frac{x}{2})^7}{4} + \frac{x \tan(\frac{x}{2})^2}{2} + \frac{3x \tan(\frac{x}{2})^4}{4} + \frac{x \tan(\frac{x}{2})^6}{2} + \frac{x \tan(\frac{x}{2})^8}{8} - \frac{\tan(\frac{x}{2})}{4}}{(1 + \tan(\frac{x}{2})^2)^4}$	82

input `int(sin(x)^2*cos(x)^2,x,method=_RETURNVERBOSE)`

output `1/8*x-1/32*sin(4*x)`

3.85.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.36

$$\int \cos^2(x) \sin^2(x) dx = -\frac{1}{8} (2 \cos(x)^3 - \cos(x)) \sin(x) + \frac{1}{8} x$$

input `integrate(sin(x)^2*cos(x)^2,x, algorithm="fracas")`output `-1/8*(2*cos(x)^3 - cos(x))*sin(x) + 1/8*x`**3.85.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \cos^2(x) \sin^2(x) dx = \frac{x}{8} - \frac{\sin(2x) \cos(2x)}{16}$$

input `integrate(sin(x)**2*cos(x)**2,x)`output `x/8 - sin(2*x)*cos(2*x)/16`**3.85.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) \sin^2(x) dx = \frac{1}{8} x - \frac{1}{32} \sin(4x)$$

input `integrate(sin(x)^2*cos(x)^2,x, algorithm="maxima")`output `1/8*x - 1/32*sin(4*x)`

3.85.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \cos^2(x) \sin^2(x) dx = \frac{1}{8} x - \frac{1}{32} \sin(4x)$$

input `integrate(sin(x)^2*cos(x)^2,x, algorithm="giac")`

output `1/8*x - 1/32*sin(4*x)`

3.85.9 Mupad [B] (verification not implemented)

Time = 17.68 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.29

$$\int \cos^2(x) \sin^2(x) dx = \frac{\cos(x) \sin(x)^3}{4} - \frac{\cos(x) \sin(x)}{8} + \frac{x}{8}$$

input `int(cos(x)^2*sin(x)^2,x)`

output `x/8 - (cos(x)*sin(x))/8 + (cos(x)*sin(x)^3)/4`

$$3.86 \quad \int \frac{\pi \sin(\pi\sqrt{x})}{\sqrt{x}} dx$$

3.86.1	Optimal result	527
3.86.2	Mathematica [A] (verified)	527
3.86.3	Rubi [A] (verified)	528
3.86.4	Maple [A] (verified)	529
3.86.5	Fricas [A] (verification not implemented)	529
3.86.6	Sympy [A] (verification not implemented)	530
3.86.7	Maxima [A] (verification not implemented)	530
3.86.8	Giac [A] (verification not implemented)	530
3.86.9	Mupad [B] (verification not implemented)	531

3.86.1 Optimal result

Integrand size = 15, antiderivative size = 10

$$\int \frac{\pi \sin(\pi\sqrt{x})}{\sqrt{x}} dx = -2 \cos(\pi\sqrt{x})$$

output `-2*cos(Pi*x^(1/2))`

3.86.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\pi \sin(\pi\sqrt{x})}{\sqrt{x}} dx = -2 \cos(\pi\sqrt{x})$$

input `Integrate[(Pi*Sin[Pi*Sqrt[x]])/Sqrt[x],x]`

output `-2*Cos[Pi*Sqrt[x]]`

3.86.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {27, 3860, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\pi \sin(\pi\sqrt{x})}{\sqrt{x}} dx \\
 & \quad \downarrow \text{27} \\
 & \pi \int \frac{\sin(\pi\sqrt{x})}{\sqrt{x}} dx \\
 & \quad \downarrow \text{3860} \\
 & 2\pi \int \sin(\pi\sqrt{x}) d\sqrt{x} \\
 & \quad \downarrow \text{3042} \\
 & 2\pi \int \sin(\pi\sqrt{x}) d\sqrt{x} \\
 & \quad \downarrow \text{3118} \\
 & -2 \cos(\pi\sqrt{x})
 \end{aligned}$$

input `Int[(Pi*Sin[Pi*Sqrt[x]])/Sqrt[x],x]`

output `-2*Cos[Pi*Sqrt[x]]`

3.86.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3118 Int[sin[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ
[{c, d}, x]
```

```
rule 3860 Int[(x_)^(m_.)*((a_.) + (b_.)*Sin[(c_.) + (d_.)*(x_)^(n_.)])^(p_.), x_Symbol
] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Sin[c + d*x])^
p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[
(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[
(m + 1)/n], 0]))
```

3.86.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

method	result	size
derivativedivides	$-2 \cos(\pi\sqrt{x})$	9
default	$-2 \cos(\pi\sqrt{x})$	9
meijerg	$2\sqrt{\pi} \left(\frac{1}{\sqrt{\pi}} - \frac{\cos(\pi\sqrt{x})}{\sqrt{\pi}} \right)$	21

```
input int(Pi*sin(Pi*x^(1/2))/x^(1/2),x,method=_RETURNVERBOSE)
```

```
output -2*cos(Pi*x^(1/2))
```

3.86.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\pi \sin(\pi\sqrt{x})}{\sqrt{x}} dx = -2 \cos(\pi\sqrt{x})$$

```
input integrate(pi*sin(pi*x^(1/2))/x^(1/2),x, algorithm="fracas")
```

```
output -2*cos(pi*sqrt(x))
```

3.86.6 Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\pi \sin(\pi\sqrt{x})}{\sqrt{x}} dx = -2 \cos(\pi\sqrt{x})$$

input `integrate(pi*sin(pi*x**(1/2))/x**(1/2),x)`output `-2*cos(pi*sqrt(x))`**3.86.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\pi \sin(\pi\sqrt{x})}{\sqrt{x}} dx = -2 \cos(\pi\sqrt{x})$$

input `integrate(pi*sin(pi*x^(1/2))/x^(1/2),x, algorithm="maxima")`output `-2*cos(pi*sqrt(x))`**3.86.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\pi \sin(\pi\sqrt{x})}{\sqrt{x}} dx = -2 \cos(\pi\sqrt{x})$$

input `integrate(pi*sin(pi*x^(1/2))/x^(1/2),x, algorithm="giac")`output `-2*cos(pi*sqrt(x))`

3.86.9 Mupad [B] (verification not implemented)

Time = 17.25 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\pi \sin(\pi\sqrt{x})}{\sqrt{x}} dx = -2 \cos(\Pi\sqrt{x})$$

input `int((Pi*sin(Pi*x^(1/2)))/x^(1/2),x)`

output `-2*cos(Pi*x^(1/2))`

3.87 $\int \tan^2(x) dx$

3.87.1	Optimal result	532
3.87.2	Mathematica [A] (verified)	532
3.87.3	Rubi [A] (verified)	533
3.87.4	Maple [A] (verified)	534
3.87.5	Fricas [A] (verification not implemented)	534
3.87.6	Sympy [B] (verification not implemented)	534
3.87.7	Maxima [A] (verification not implemented)	535
3.87.8	Giac [A] (verification not implemented)	535
3.87.9	Mupad [B] (verification not implemented)	535

3.87.1 Optimal result

Integrand size = 4, antiderivative size = 6

$$\int \tan^2(x) dx = -x + \tan(x)$$

output `-x+tan(x)`

3.87.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.33

$$\int \tan^2(x) dx = -\arctan(\tan(x)) + \tan(x)$$

input `Integrate[Tan[x]^2,x]`

output `-ArcTan[Tan[x]] + Tan[x]`

3.87.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \tan^2(x) dx \\
 \downarrow 3042 \\
 \int \tan(x)^2 dx \\
 \downarrow 3954 \\
 \tan(x) - \int 1 dx \\
 \downarrow 24 \\
 \tan(x) - x
 \end{array}$$

input `Int[Tan[x]^2,x]`

output `-x + Tan[x]`

3.87.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

3.87.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

method	result	size
norman	$-x + \tan(x)$	7
parallelrisch	$-x + \tan(x)$	7
derivativedivides	$\tan(x) - \arctan(\tan(x))$	9
default	$\tan(x) - \arctan(\tan(x))$	9
risch	$-x + \frac{2i}{e^{2ix}+1}$	17

input `int(tan(x)^2,x,method=_RETURNVERBOSE)`

output `-x+tan(x)`

3.87.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \tan^2(x) dx = -x + \tan(x)$$

input `integrate(tan(x)^2,x, algorithm="fricas")`

output `-x + tan(x)`

3.87.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7 vs. $2(3) = 6$.

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

$$\int \tan^2(x) dx = -x + \frac{\sin(x)}{\cos(x)}$$

input `integrate(tan(x)**2,x)`

output `-x + sin(x)/cos(x)`

3.87.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \tan^2(x) dx = -x + \tan(x)$$

input `integrate(tan(x)^2,x, algorithm="maxima")`

output `-x + tan(x)`

3.87.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \tan^2(x) dx = -x + \tan(x)$$

input `integrate(tan(x)^2,x, algorithm="giac")`

output `-x + tan(x)`

3.87.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \tan^2(x) dx = \tan(x) - x$$

input `int(tan(x)^2,x)`

output `tan(x) - x`

3.88 $\int e^{\sqrt[4]{x}} dx$

3.88.1	Optimal result	536
3.88.2	Mathematica [A] (verified)	536
3.88.3	Rubi [A] (verified)	537
3.88.4	Maple [A] (verified)	538
3.88.5	Fricas [A] (verification not implemented)	539
3.88.6	Sympy [A] (verification not implemented)	539
3.88.7	Maxima [A] (verification not implemented)	539
3.88.8	Giac [A] (verification not implemented)	540
3.88.9	Mupad [B] (verification not implemented)	540

3.88.1 Optimal result

Integrand size = 7, antiderivative size = 30

$$\int e^{\sqrt[4]{x}} dx = 4e^{\sqrt[4]{x}}(-6 + 6\sqrt[4]{x} - 3\sqrt{x} + x^{3/4})$$

output `4*exp(x^(1/4))*(-6+6*x^(1/4)-3*x^(1/2)+x^(3/4))`

3.88.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.03

$$\int e^{\sqrt[4]{x}} dx = e^{\sqrt[4]{x}}(-24 + 24\sqrt[4]{x} - 12\sqrt{x} + 4x^{3/4})$$

input `Integrate[E^x^(1/4),x]`

output `E^x^(1/4)*(-24 + 24*x^(1/4) - 12*Sqrt[x] + 4*x^(3/4))`

3.88.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.90, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {2636, 2607, 2607, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{\sqrt[4]{x}} dx \\
 & \quad \downarrow \text{2636} \\
 & 4 \int e^{\sqrt[4]{x}} x^{3/4} d\sqrt[4]{x} \\
 & \quad \downarrow \text{2607} \\
 & 4 \left(e^{\sqrt[4]{x}} x^{3/4} - 3 \int e^{\sqrt[4]{x}} \sqrt{x} d\sqrt[4]{x} \right) \\
 & \quad \downarrow \text{2607} \\
 & 4 \left(e^{\sqrt[4]{x}} x^{3/4} - 3 \left(e^{\sqrt[4]{x}} \sqrt{x} - 2 \int e^{\sqrt[4]{x}} \sqrt[4]{x} d\sqrt[4]{x} \right) \right) \\
 & \quad \downarrow \text{2607} \\
 & 4 \left(e^{\sqrt[4]{x}} x^{3/4} - 3 \left(e^{\sqrt[4]{x}} \sqrt{x} - 2 \left(e^{\sqrt[4]{x}} \sqrt[4]{x} - \int e^{\sqrt[4]{x}} d\sqrt[4]{x} \right) \right) \right) \\
 & \quad \downarrow \text{2624} \\
 & 4 \left(e^{\sqrt[4]{x}} x^{3/4} - 3 \left(e^{\sqrt[4]{x}} \sqrt{x} - 2 \left(e^{\sqrt[4]{x}} \sqrt[4]{x} - e^{\sqrt[4]{x}} \right) \right) \right)
 \end{aligned}$$

input `Int[E^x^(1/4),x]`

output `4*(-3*(-2*(-E^x^(1/4) + E^x^(1/4)*x^(1/4)) + E^x^(1/4)*Sqrt[x]) + E^x^(1/4)*x^(3/4))`

3.88.3.1 Defintions of rubi rules used

```
rule 2607 Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.))*((c_.) + (d_.)*(x_))^(m
_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])),
x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^
n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*
m] && !TrueQ[$UseGamma]
```

```
rule 2624 Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;
FreeQ[{F, n}, x] && LinearQ[v, x]
```

```
rule 2636 Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_)), x_Symbol] :> With[{k =
Denominator[n]}, Simp[k/d Subst[Int[x^(k - 1)*F^(a + b*x^(k*n)), x], x, (
c + d*x)^(1/k)], x] /; FreeQ[{F, a, b, c, d}, x] && IntegerQ[2/n] && !Int
egerQ[n]
```

3.88.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.87

method	result	size
meijerg	$24 - \left(-4x^{\frac{3}{4}} + 12\sqrt{x} - 24x^{\frac{1}{4}} + 24\right) e^{x^{\frac{1}{4}}}$	26
derivativedivides	$4e^{x^{\frac{1}{4}}} x^{\frac{3}{4}} - 12\sqrt{x} e^{x^{\frac{1}{4}}} + 24x^{\frac{1}{4}} e^{x^{\frac{1}{4}}} - 24e^{x^{\frac{1}{4}}}$	35
default	$4e^{x^{\frac{1}{4}}} x^{\frac{3}{4}} - 12\sqrt{x} e^{x^{\frac{1}{4}}} + 24x^{\frac{1}{4}} e^{x^{\frac{1}{4}}} - 24e^{x^{\frac{1}{4}}}$	35

```
input int(exp(x^(1/4)),x,method=_RETURNVERBOSE)
```

```
output 24-(-4*x^(3/4)+12*x^(1/2)-24*x^(1/4)+24)*exp(x^(1/4))
```

3.88.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.70

$$\int e^{\sqrt[4]{x}} dx = 4 \left(x^{\frac{3}{4}} - 3\sqrt{x} + 6x^{\frac{1}{4}} - 6 \right) e^{(x^{\frac{1}{4}})}$$

input `integrate(exp(x^(1/4)),x, algorithm="fracas")`output `4*(x^(3/4) - 3*sqrt(x) + 6*x^(1/4) - 6)*e^(x^(1/4))`**3.88.6 Sympy [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.60

$$\int e^{\sqrt[4]{x}} dx = 4x^{\frac{3}{4}}e^{\sqrt[4]{x}} + 24\sqrt[4]{x}e^{\sqrt[4]{x}} - 12\sqrt{x}e^{\sqrt[4]{x}} - 24e^{\sqrt[4]{x}}$$

input `integrate(exp(x**(1/4)),x)`output `4*x**(3/4)*exp(x**(1/4)) + 24*x**(1/4)*exp(x**(1/4)) - 12*sqrt(x)*exp(x**(1/4)) - 24*exp(x**(1/4))`**3.88.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.70

$$\int e^{\sqrt[4]{x}} dx = 4 \left(x^{\frac{3}{4}} - 3\sqrt{x} + 6x^{\frac{1}{4}} - 6 \right) e^{(x^{\frac{1}{4}})}$$

input `integrate(exp(x^(1/4)),x, algorithm="maxima")`output `4*(x^(3/4) - 3*sqrt(x) + 6*x^(1/4) - 6)*e^(x^(1/4))`

3.88.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.70

$$\int e^{\sqrt[4]{x}} dx = 4 \left(x^{\frac{3}{4}} - 3\sqrt{x} + 6x^{\frac{1}{4}} - 6 \right) e^{(x^{\frac{1}{4}})}$$

input `integrate(exp(x^(1/4)),x, algorithm="giac")`output `4*(x^(3/4) - 3*sqrt(x) + 6*x^(1/4) - 6)*e^(x^(1/4))`**3.88.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int e^{\sqrt[4]{x}} dx = -4x e^{x^{1/4}} \left(\frac{6}{x} + \frac{3}{\sqrt{x}} - \frac{1}{x^{1/4}} - \frac{6}{x^{3/4}} \right)$$

input `int(exp(x^(1/4)),x)`output `-4*x*exp(x^(1/4))*(6/x + 3/x^(1/2) - 1/x^(1/4) - 6/x^(3/4))`

3.89 $\int \cos(x) \cot(x) dx$

3.89.1	Optimal result	541
3.89.2	Mathematica [B] (verified)	541
3.89.3	Rubi [A] (verified)	542
3.89.4	Maple [A] (verified)	543
3.89.5	Fricas [B] (verification not implemented)	544
3.89.6	Sympy [B] (verification not implemented)	544
3.89.7	Maxima [B] (verification not implemented)	544
3.89.8	Giac [B] (verification not implemented)	545
3.89.9	Mupad [B] (verification not implemented)	545

3.89.1 Optimal result

Integrand size = 5, antiderivative size = 8

$$\int \cos(x) \cot(x) dx = -\operatorname{arctanh}(\cos(x)) + \cos(x)$$

output `-arctanh(cos(x))+cos(x)`

3.89.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 19 vs. $2(8) = 16$.

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.38

$$\int \cos(x) \cot(x) dx = \cos(x) - \log\left(\cos\left(\frac{x}{2}\right)\right) + \log\left(\sin\left(\frac{x}{2}\right)\right)$$

input `Integrate[Cos[x]*Cot[x],x]`

output `Cos[x] - Log[Cos[x/2]] + Log[Sin[x/2]]`

3.89.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 25, 3072, 262, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(x) \cot(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\sin\left(x + \frac{\pi}{2}\right) \tan\left(x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{25} \\
 & -\int \sin\left(x + \frac{\pi}{2}\right) \tan\left(x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3072} \\
 & -\int \frac{\cos^2(x)}{1 - \cos^2(x)} d\cos(x) \\
 & \quad \downarrow \text{262} \\
 & \cos(x) - \int \frac{1}{1 - \cos^2(x)} d\cos(x) \\
 & \quad \downarrow \text{219} \\
 & \cos(x) - \operatorname{arctanh}(\cos(x))
 \end{aligned}$$

input `Int[Cos[x]*Cot[x],x]`

output `-ArcTanh[Cos[x]] + Cos[x]`

3.89.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 262 `Int[((c_)*(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3072 `Int[((a_)*sin[(e_) + (f_)*(x_)])^(m_)*tan[(e_) + (f_)*(x_)]^(n_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Simp[ff/f Subst[Int[(ff*x)^(m + n)/(a^2 - ff^2*x^2)^((n + 1)/2), x], x, a*(Sin[e + f*x]/ff)], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2]`

3.89.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.50

method	result	size
default	$\cos(x) + \ln(-\cot(x) + \csc(x))$	12
risch	$\frac{e^{ix}}{2} + \frac{e^{-ix}}{2} + \ln(-1 + e^{ix}) - \ln(e^{ix} + 1)$	34

input `int(cos(x)*cot(x),x,method=_RETURNVERBOSE)`

output `cos(x)+ln(-cot(x)+csc(x))`

3.89.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. $2(8) = 16$.

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 2.62

$$\int \cos(x) \cot(x) dx = \cos(x) - \frac{1}{2} \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) + \frac{1}{2} \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right)$$

input `integrate(cos(x)*cot(x),x, algorithm="fracas")`

output `cos(x) - 1/2*log(1/2*cos(x) + 1/2) + 1/2*log(-1/2*cos(x) + 1/2)`

3.89.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(7) = 14$.

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.38

$$\int \cos(x) \cot(x) dx = \frac{\log(\cos(x) - 1)}{2} - \frac{\log(\cos(x) + 1)}{2} + \cos(x)$$

input `integrate(cos(x)*cot(x),x)`

output `log(cos(x) - 1)/2 - log(cos(x) + 1)/2 + cos(x)`

3.89.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(8) = 16$.

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.12

$$\int \cos(x) \cot(x) dx = \cos(x) - \frac{1}{2} \log(\cos(x) + 1) + \frac{1}{2} \log(\cos(x) - 1)$$

input `integrate(cos(x)*cot(x),x, algorithm="maxima")`

output `cos(x) - 1/2*log(cos(x) + 1) + 1/2*log(cos(x) - 1)`

3.89.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(8) = 16$.

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.38

$$\int \cos(x) \cot(x) dx = \cos(x) - \frac{1}{2} \log(\cos(x) + 1) + \frac{1}{2} \log(-\cos(x) + 1)$$

input `integrate(cos(x)*cot(x),x, algorithm="giac")`

output `cos(x) - 1/2*log(cos(x) + 1) + 1/2*log(-cos(x) + 1)`

3.89.9 Mupad [B] (verification not implemented)

Time = 16.47 (sec) , antiderivative size = 18, normalized size of antiderivative = 2.25

$$\int \cos(x) \cot(x) dx = \ln\left(\tan\left(\frac{x}{2}\right)\right) + \frac{2}{\tan\left(\frac{x}{2}\right)^2 + 1}$$

input `int(cos(x)*cot(x),x)`

output `log(tan(x/2)) + 2/(tan(x/2)^2 + 1)`

3.90 $\int (2 \log(x) + \log^2(x)) dx$

3.90.1	Optimal result	546
3.90.2	Mathematica [A] (verified)	546
3.90.3	Rubi [A] (verified)	547
3.90.4	Maple [A] (verified)	547
3.90.5	Fricas [A] (verification not implemented)	548
3.90.6	Sympy [A] (verification not implemented)	548
3.90.7	Maxima [B] (verification not implemented)	548
3.90.8	Giac [A] (verification not implemented)	549
3.90.9	Mupad [B] (verification not implemented)	549

3.90.1 Optimal result

Integrand size = 9, antiderivative size = 6

$$\int (2 \log(x) + \log^2(x)) dx = x \log^2(x)$$

output `x*ln(x)^2`

3.90.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int (2 \log(x) + \log^2(x)) dx = x \log^2(x)$$

input `Integrate[2*Log[x] + Log[x]^2,x]`

output `x*Log[x]^2`

3.90.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\log^2(x) + 2 \log(x)) dx$$

$$\downarrow \text{2009}$$

$$x \log^2(x)$$

input `Int[2*Log[x] + Log[x]^2,x]`

output `x*Log[x]^2`

3.90.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.90.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

method	result	size
default	$x \ln(x)^2$	7
norman	$x \ln(x)^2$	7
risch	$x \ln(x)^2$	7
parallelrisch	$x \ln(x)^2$	7
parts	$x \ln(x)^2$	7

input `int(2*ln(x)+ln(x)^2,x,method=_RETURNVERBOSE)`

output `x*ln(x)^2`

3.90.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int (2\log(x) + \log^2(x)) dx = x \log(x)^2$$

input `integrate(2*log(x)+log(x)^2,x, algorithm="fracas")`

output `x*log(x)^2`

3.90.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int (2\log(x) + \log^2(x)) dx = x \log(x)^2$$

input `integrate(2*ln(x)+ln(x)**2,x)`

output `x*log(x)**2`

3.90.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. $2(6) = 12$.

Time = 0.20 (sec) , antiderivative size = 21, normalized size of antiderivative = 3.50

$$\int (2\log(x) + \log^2(x)) dx = (\log(x)^2 - 2\log(x) + 2)x + 2x\log(x) - 2x$$

input `integrate(2*log(x)+log(x)^2,x, algorithm="maxima")`

output `(log(x)^2 - 2*log(x) + 2)*x + 2*x*log(x) - 2*x`

3.90.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int (2\log(x) + \log^2(x)) dx = x \log(x)^2$$

input `integrate(2*log(x)+log(x)^2,x, algorithm="giac")`

output `x*log(x)^2`

3.90.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int (2\log(x) + \log^2(x)) dx = x \ln(x)^2$$

input `int(2*log(x) + log(x)^2,x)`

output `x*log(x)^2`

3.91 $\int \frac{x^3}{1+x^2} dx$

3.91.1	Optimal result	550
3.91.2	Mathematica [A] (verified)	550
3.91.3	Rubi [A] (verified)	551
3.91.4	Maple [A] (verified)	552
3.91.5	Fricas [A] (verification not implemented)	552
3.91.6	Sympy [A] (verification not implemented)	552
3.91.7	Maxima [A] (verification not implemented)	553
3.91.8	Giac [A] (verification not implemented)	553
3.91.9	Mupad [B] (verification not implemented)	553

3.91.1 Optimal result

Integrand size = 11, antiderivative size = 18

$$\int \frac{x^3}{1+x^2} dx = \frac{x^2}{2} - \frac{1}{2} \log(1+x^2)$$

output `1/2*x^2-1/2*ln(x^2+1)`

3.91.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{x^3}{1+x^2} dx = \frac{x^2}{2} - \frac{1}{2} \log(1+x^2)$$

input `Integrate[x^3/(1 + x^2),x]`

output `x^2/2 - Log[1 + x^2]/2`

3.91.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^3}{x^2+1} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int \frac{x^2}{x^2+1} dx^2 \\ & \quad \downarrow \text{49} \\ & \frac{1}{2} \int \left(1 + \frac{1}{-x^2-1} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} (x^2 - \log(x^2+1)) \end{aligned}$$

input `Int[x^3/(1 + x^2), x]`

output `(x^2 - Log[1 + x^2])/2`

3.91.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.91.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{x^2}{2} - \frac{\ln(x^2+1)}{2}$	15
norman	$\frac{x^2}{2} - \frac{\ln(x^2+1)}{2}$	15
meijerg	$\frac{x^2}{2} - \frac{\ln(x^2+1)}{2}$	15
risch	$\frac{x^2}{2} - \frac{\ln(x^2+1)}{2}$	15
parallelrisch	$\frac{x^2}{2} - \frac{\ln(x^2+1)}{2}$	15

input `int(x^3/(x^2+1),x,method=_RETURNVERBOSE)`output `1/2*x^2-1/2*ln(x^2+1)`**3.91.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{x^3}{1+x^2} dx = \frac{1}{2} x^2 - \frac{1}{2} \log(x^2 + 1)$$

input `integrate(x^3/(x^2+1),x, algorithm="fricas")`output `1/2*x^2 - 1/2*log(x^2 + 1)`**3.91.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \frac{x^3}{1+x^2} dx = \frac{x^2}{2} - \frac{\log(x^2 + 1)}{2}$$

input `integrate(x**3/(x**2+1),x)`output `x**2/2 - log(x**2 + 1)/2`

3.91.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{x^3}{1+x^2} dx = \frac{1}{2}x^2 - \frac{1}{2}\log(x^2+1)$$

input `integrate(x^3/(x^2+1),x, algorithm="maxima")`output `1/2*x^2 - 1/2*log(x^2 + 1)`**3.91.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{x^3}{1+x^2} dx = \frac{1}{2}x^2 - \frac{1}{2}\log(x^2+1)$$

input `integrate(x^3/(x^2+1),x, algorithm="giac")`output `1/2*x^2 - 1/2*log(x^2 + 1)`**3.91.9 Mupad [B] (verification not implemented)**

Time = 16.80 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{x^3}{1+x^2} dx = \frac{x^2}{2} - \frac{\ln(x^2+1)}{2}$$

input `int(x^3/(x^2 + 1),x)`output `x^2/2 - log(x^2 + 1)/2`

3.92 $\int \frac{1}{2-2x+x^2} dx$

3.92.1	Optimal result	554
3.92.2	Mathematica [A] (verified)	554
3.92.3	Rubi [A] (verified)	555
3.92.4	Maple [A] (verified)	556
3.92.5	Fricas [A] (verification not implemented)	556
3.92.6	Sympy [A] (verification not implemented)	556
3.92.7	Maxima [A] (verification not implemented)	557
3.92.8	Giac [A] (verification not implemented)	557
3.92.9	Mupad [B] (verification not implemented)	557

3.92.1 Optimal result

Integrand size = 10, antiderivative size = 8

$$\int \frac{1}{2 - 2x + x^2} dx = -\arctan(1 - x)$$

output `arctan(-1+x)`

3.92.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{1}{2 - 2x + x^2} dx = -\arctan(1 - x)$$

input `Integrate[(2 - 2*x + x^2)^(-1),x]`

output `-ArcTan[1 - x]`

3.92.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 - 2x + 2} dx$$

↓ 1082

$$\int \frac{1}{-(1-x)^2 - 1} d(1-x)$$

↓ 217

$$-\arctan(1-x)$$

input `Int[(2 - 2*x + x^2)^(-1),x]`

output `-ArcTan[1 - x]`

3.92.3.1 Defintions of rubi rules used

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

3.92.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

method	result	size
default	$\arctan(-1 + x)$	5
risch	$\arctan(-1 + x)$	5
parallelrisc	$\frac{i \ln(x-1+i)}{2} - \frac{i \ln(x-1-i)}{2}$	20

input `int(1/(x^2-2*x+2),x,method=_RETURNVERBOSE)`

output `arctan(-1+x)`

3.92.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.50

$$\int \frac{1}{2 - 2x + x^2} dx = \arctan(x - 1)$$

input `integrate(1/(x^2-2*x+2),x, algorithm="fricas")`

output `arctan(x - 1)`

3.92.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.38

$$\int \frac{1}{2 - 2x + x^2} dx = \operatorname{atan}(x - 1)$$

input `integrate(1/(x**2-2*x+2),x)`

output `atan(x - 1)`

3.92.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.50

$$\int \frac{1}{2 - 2x + x^2} dx = \arctan(x - 1)$$

input `integrate(1/(x^2-2*x+2),x, algorithm="maxima")`output `arctan(x - 1)`**3.92.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.50

$$\int \frac{1}{2 - 2x + x^2} dx = \arctan(x - 1)$$

input `integrate(1/(x^2-2*x+2),x, algorithm="giac")`output `arctan(x - 1)`**3.92.9 Mupad [B] (verification not implemented)**

Time = 16.31 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.50

$$\int \frac{1}{2 - 2x + x^2} dx = \operatorname{atan}(x - 1)$$

input `int(1/(x^2 - 2*x + 2),x)`output `atan(x - 1)`

3.93 $\int \log(\sin(x)) \sin(x) dx$

3.93.1	Optimal result	558
3.93.2	Mathematica [A] (verified)	558
3.93.3	Rubi [A] (verified)	559
3.93.4	Maple [A] (verified)	561
3.93.5	Fricas [A] (verification not implemented)	561
3.93.6	Sympy [B] (verification not implemented)	562
3.93.7	Maxima [B] (verification not implemented)	562
3.93.8	Giac [A] (verification not implemented)	563
3.93.9	Mupad [F(-1)]	563

3.93.1 Optimal result

Integrand size = 6, antiderivative size = 15

$$\int \log(\sin(x)) \sin(x) dx = -\operatorname{arctanh}(\cos(x)) + \cos(x) - \cos(x) \log(\sin(x))$$

output `-arctanh(cos(x))+cos(x)-cos(x)*ln(sin(x))`

3.93.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.73

$$\int \log(\sin(x)) \sin(x) dx = \cos(x) - \log\left(\cos\left(\frac{x}{2}\right)\right) + \log\left(\sin\left(\frac{x}{2}\right)\right) - \cos(x) \log(\sin(x))$$

input `Integrate[Log[Sin[x]]*Sin[x],x]`

output `Cos[x] - Log[Cos[x/2]] + Log[Sin[x/2]] - Cos[x]*Log[Sin[x]]`

3.93.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.167$, Rules used = {3034, 25, 3042, 25, 3072, 262, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(x) \log(\sin(x)) dx \\
 & \quad \downarrow \text{3034} \\
 & - \int -\cos(x) \cot(x) dx - \cos(x) \log(\sin(x)) \\
 & \quad \downarrow \text{25} \\
 & \int \cos(x) \cot(x) dx - \cos(x) \log(\sin(x)) \\
 & \quad \downarrow \text{3042} \\
 & \int -\sin\left(x + \frac{\pi}{2}\right) \tan\left(x + \frac{\pi}{2}\right) dx - \cos(x) \log(\sin(x)) \\
 & \quad \downarrow \text{25} \\
 & - \int \sin\left(x + \frac{\pi}{2}\right) \tan\left(x + \frac{\pi}{2}\right) dx - \cos(x) \log(\sin(x)) \\
 & \quad \downarrow \text{3072} \\
 & - \int \frac{\cos^2(x)}{1 - \cos^2(x)} d \cos(x) - \cos(x) \log(\sin(x)) \\
 & \quad \downarrow \text{262} \\
 & - \int \frac{1}{1 - \cos^2(x)} d \cos(x) + \cos(x) + \cos(x)(-\log(\sin(x))) \\
 & \quad \downarrow \text{219} \\
 & -\operatorname{arctanh}(\cos(x)) + \cos(x) - \cos(x) \log(\sin(x))
 \end{aligned}$$

input `Int [Log [Sin [x]] *Sin [x] ,x]`

output `-ArcTanh [Cos [x]] + Cos [x] - Cos [x]*Log [Sin [x]]`

3.93.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 262 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*(m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 3034 `Int[Log[u_]*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[Log[u] w, x] - Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w, x]] /; InverseFunctionFreeQ[u, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3072 `Int[((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*tan[(e_.) + (f_.)*(x_)^(n_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Simp[ff/f Subst[Int[(ff*x)^(m + n)/(a^2 - ff^2*x^2)^((n + 1)/2), x], x, a*(Sin[e + f*x]/ff)], x]] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2]`

3.93.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.33

method	result
parallelrisc	$-\cos(x) \ln(\sin(x)) + \cos(x) + \ln(-\cot(x) + \csc(x)) + 1$
norman	$\frac{2 \tan(\frac{x}{2})^2 \ln\left(\frac{2 \tan(\frac{x}{2})}{1 + \tan(\frac{x}{2})^2}\right) + 2}{1 + \tan(\frac{x}{2})^2} + \ln\left(1 + \tan\left(\frac{x}{2}\right)^2\right)$
default	$-\frac{e^{ix} \ln(i(1-e^{2ix})e^{-ix})}{2} + \frac{e^{ix}}{2} + \ln(-1 + e^{ix}) - \ln(e^{ix} + 1) - \frac{e^{-ix} \ln(i(1-e^{2ix})e^{-ix})}{2} + \frac{e^{-ix}}{2} + \frac{\ln(2)(e^{ix})}{2}$
risch	$\ln(e^{ix}) \cos(x) + \frac{e^{-ix} \ln(2)}{2} - \frac{e^{-ix} \ln(e^{2ix}-1)}{2} - \frac{e^{ix} \ln(e^{2ix}-1)}{2} + \frac{e^{ix} \ln(2)}{2} + \frac{e^{ix}}{2} - \frac{ie^{-ix} \operatorname{csgn}(\sin(x))^2 \operatorname{csgn}(i)}{4}$

input `int(sin(x)*ln(sin(x)),x,method=_RETURNVERBOSE)`output `-cos(x)*ln(sin(x))+cos(x)+ln(-cot(x)+csc(x))+1`**3.93.5 Fricas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.87

$$\int \log(\sin(x)) \sin(x) dx = -\cos(x) \log(\sin(x)) + \cos(x) - \frac{1}{2} \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) + \frac{1}{2} \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right)$$

input `integrate(sin(x)*log(sin(x)),x, algorithm="fricas")`output `-cos(x)*log(sin(x)) + cos(x) - 1/2*log(1/2*cos(x) + 1/2) + 1/2*log(-1/2*cos(x) + 1/2)`

3.93.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. $2(15) = 30$.

Time = 0.57 (sec) , antiderivative size = 105, normalized size of antiderivative = 7.00

$$\int \log(\sin(x)) \sin(x) dx = \frac{2 \log\left(\frac{\tan\left(\frac{x}{2}\right)}{\tan^2\left(\frac{x}{2}\right)+1}\right) \tan^2\left(\frac{x}{2}\right)}{\tan^2\left(\frac{x}{2}\right)+1} + \frac{\log\left(\tan^2\left(\frac{x}{2}\right)+1\right) \tan^2\left(\frac{x}{2}\right)}{\tan^2\left(\frac{x}{2}\right)+1}$$

$$+ \frac{\log\left(\tan^2\left(\frac{x}{2}\right)+1\right)}{\tan^2\left(\frac{x}{2}\right)+1} + \frac{2 \log(2) \tan^2\left(\frac{x}{2}\right)}{\tan^2\left(\frac{x}{2}\right)+1} + \frac{2}{\tan^2\left(\frac{x}{2}\right)+1}$$

input `integrate(sin(x)*ln(sin(x)),x)`

output `2*log(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)**2/(tan(x/2)**2 + 1) + log(tan(x/2)**2 + 1)*tan(x/2)**2/(tan(x/2)**2 + 1) + log(tan(x/2)**2 + 1)/(tan(x/2)**2 + 1) + 2*log(2)*tan(x/2)**2/(tan(x/2)**2 + 1) + 2/(tan(x/2)**2 + 1)`

3.93.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 89 vs. $2(15) = 30$.

Time = 0.19 (sec) , antiderivative size = 89, normalized size of antiderivative = 5.93

$$\int \log(\sin(x)) \sin(x) dx = -\frac{2 \log\left(\frac{2 \sin(x)}{\left(\frac{\sin(x)^2}{(\cos(x)+1)^2+1}\right)(\cos(x)+1)}\right)}{\frac{\sin(x)^2}{(\cos(x)+1)^2}+1} + \frac{2}{\frac{\sin(x)^2}{(\cos(x)+1)^2}+1}$$

$$- \log\left(\frac{\sin(x)^2}{(\cos(x)+1)^2}+1\right) + \log\left(\frac{\sin(x)^2}{(\cos(x)+1)^2}\right)$$

input `integrate(sin(x)*log(sin(x)),x, algorithm="maxima")`

output `-2*log(2*sin(x)/((sin(x)^2/(cos(x) + 1)^2 + 1)*(cos(x) + 1)))/(sin(x)^2/(cos(x) + 1)^2 + 1) + 2/(sin(x)^2/(cos(x) + 1)^2 + 1) - log(sin(x)^2/(cos(x) + 1)^2 + 1) + log(sin(x)^2/(cos(x) + 1)^2)`

3.93.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.73

$$\int \log(\sin(x)) \sin(x) dx = -\cos(x) \log(\sin(x)) + \cos(x) - \frac{1}{2} \log(\cos(x) + 1) + \frac{1}{2} \log(-\cos(x) + 1)$$

input `integrate(sin(x)*log(sin(x)),x, algorithm="giac")`

output `-cos(x)*log(sin(x)) + cos(x) - 1/2*log(cos(x) + 1) + 1/2*log(-cos(x) + 1)`

3.93.9 Mupad [F(-1)]

Timed out.

$$\int \log(\sin(x)) \sin(x) dx = \int \ln(\sin(x)) \sin(x) dx$$

input `int(log(sin(x))*sin(x),x)`

output `int(log(sin(x))*sin(x), x)`

3.94 $\int \frac{x}{1-x^4} dx$

3.94.1	Optimal result	564
3.94.2	Mathematica [B] (verified)	564
3.94.3	Rubi [A] (verified)	565
3.94.4	Maple [A] (verified)	566
3.94.5	Fricas [B] (verification not implemented)	566
3.94.6	Sympy [B] (verification not implemented)	566
3.94.7	Maxima [B] (verification not implemented)	567
3.94.8	Giac [B] (verification not implemented)	567
3.94.9	Mupad [B] (verification not implemented)	568

3.94.1 Optimal result

Integrand size = 11, antiderivative size = 8

$$\int \frac{x}{1-x^4} dx = \frac{\operatorname{arctanh}(x^2)}{2}$$

output `1/2*arctanh(x^2)`

3.94.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 23 vs. $2(8) = 16$.

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 2.88

$$\int \frac{x}{1-x^4} dx = -\frac{1}{4} \log(1-x^2) + \frac{1}{4} \log(1+x^2)$$

input `Integrate[x/(1 - x^4),x]`

output `-1/4*Log[1 - x^2] + Log[1 + x^2]/4`

3.94.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {807, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{1-x^4} dx$$

↓ 807

$$\frac{1}{2} \int \frac{1}{1-x^4} dx^2$$

↓ 219

$$\frac{\operatorname{arctanh}(x^2)}{2}$$

input `Int[x/(1 - x^4), x]`

output `ArcTanh[x^2]/2`

3.94.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

3.94.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
meijerg	$\frac{\operatorname{arctanh}(x^2)}{2}$	7
risch	$\frac{\ln(x^2+1)}{4} - \frac{\ln(x^2-1)}{4}$	18
default	$-\frac{\ln(-1+x)}{4} - \frac{\ln(1+x)}{4} + \frac{\ln(x^2+1)}{4}$	22
norman	$-\frac{\ln(-1+x)}{4} - \frac{\ln(1+x)}{4} + \frac{\ln(x^2+1)}{4}$	22
parallelrisch	$-\frac{\ln(-1+x)}{4} - \frac{\ln(1+x)}{4} + \frac{\ln(x^2+1)}{4}$	22

input `int(x/(-x^4+1),x,method=_RETURNVERBOSE)`

output `1/2*arctanh(x^2)`

3.94.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. 2(6) = 12.

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.12

$$\int \frac{x}{1-x^4} dx = \frac{1}{4} \log(x^2+1) - \frac{1}{4} \log(x^2-1)$$

input `integrate(x/(-x^4+1),x, algorithm="fricas")`

output `1/4*log(x^2 + 1) - 1/4*log(x^2 - 1)`

3.94.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. 2(5) = 10.

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.88

$$\int \frac{x}{1-x^4} dx = -\frac{\log(x^2-1)}{4} + \frac{\log(x^2+1)}{4}$$

input `integrate(x/(-x**4+1),x)`

output `-log(x**2 - 1)/4 + log(x**2 + 1)/4`

3.94.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(6) = 12$.

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.12

$$\int \frac{x}{1-x^4} dx = \frac{1}{4} \log(x^2 + 1) - \frac{1}{4} \log(x^2 - 1)$$

input `integrate(x/(-x^4+1),x, algorithm="maxima")`

output `1/4*log(x^2 + 1) - 1/4*log(x^2 - 1)`

3.94.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18 vs. $2(6) = 12$.

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 2.25

$$\int \frac{x}{1-x^4} dx = \frac{1}{4} \log(x^2 + 1) - \frac{1}{4} \log(|x^2 - 1|)$$

input `integrate(x/(-x^4+1),x, algorithm="giac")`

output `1/4*log(x^2 + 1) - 1/4*log(abs(x^2 - 1))`

3.94.9 Mupad [B] (verification not implemented)

Time = 16.74 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{x}{1-x^4} dx = \frac{\operatorname{atanh}(x^2)}{2}$$

input `int(-x/(x^4 - 1),x)`

output `atanh(x^2)/2`

3.95 $\int \sqrt{12 - 3x^2} dx$

3.95.1	Optimal result	569
3.95.2	Mathematica [A] (verified)	569
3.95.3	Rubi [A] (verified)	570
3.95.4	Maple [A] (verified)	571
3.95.5	Fricas [A] (verification not implemented)	571
3.95.6	Sympy [A] (verification not implemented)	572
3.95.7	Maxima [A] (verification not implemented)	572
3.95.8	Giac [A] (verification not implemented)	572
3.95.9	Mupad [B] (verification not implemented)	573

3.95.1 Optimal result

Integrand size = 11, antiderivative size = 35

$$\int \sqrt{12 - 3x^2} dx = \frac{1}{2}\sqrt{3}x\sqrt{4 - x^2} + 2\sqrt{3} \arcsin\left(\frac{x}{2}\right)$$

output `1/2*3^(1/2)*x*(-x^2+4)^(1/2)+2*3^(1/2)*arcsin(1/2*x)`

3.95.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.23

$$\int \sqrt{12 - 3x^2} dx = \frac{1}{2}\sqrt{3}\left(x\sqrt{4 - x^2} - 8 \arctan\left(\frac{\sqrt{4 - x^2}}{2 + x}\right)\right)$$

input `Integrate[Sqrt[12 - 3*x^2],x]`

output `(Sqrt[3]*(x*Sqrt[4 - x^2] - 8*ArcTan[Sqrt[4 - x^2]/(2 + x)]))/2`

3.95.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {211, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{12 - 3x^2} dx$$

$$\downarrow \text{211}$$

$$6 \int \frac{1}{\sqrt{12 - 3x^2}} dx + \frac{1}{2} \sqrt{3} \sqrt{4 - x^2} x$$

$$\downarrow \text{223}$$

$$2\sqrt{3} \arcsin\left(\frac{x}{2}\right) + \frac{1}{2} \sqrt{3} \sqrt{4 - x^2} x$$

input `Int[Sqrt[12 - 3*x^2], x]`

output `(Sqrt[3]*x*Sqrt[4 - x^2])/2 + 2*Sqrt[3]*ArcSin[x/2]`

3.95.3.1 Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

3.95.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.66

method	result	size
default	$\frac{x\sqrt{-3x^2+12}}{2} + 2\sqrt{3} \arcsin\left(\frac{x}{2}\right)$	23
risch	$-\frac{3x(x^2-4)}{2\sqrt{-3x^2+12}} + 2\sqrt{3} \arcsin\left(\frac{x}{2}\right)$	28
meijerg	$\frac{i\sqrt{3} \left(-i\sqrt{\pi} x \sqrt{-\frac{x^2}{4}+1} - 2i\sqrt{\pi} \arcsin\left(\frac{x}{2}\right) \right)}{\sqrt{\pi}}$	37
pseudoelliptic	$\frac{x\sqrt{-3x^2+12}}{2} - 2\sqrt{3} \arctan\left(\frac{\sqrt{-3x^2+12}\sqrt{3}}{3x}\right)$	37
trager	$\frac{x\sqrt{-3x^2+12}}{2} + 2 \operatorname{RootOf}(_Z^2 + 3) \ln(\operatorname{RootOf}(_Z^2 + 3) \sqrt{-3x^2 + 12} + 3x)$	43

input `int((-3*x^2+12)^(1/2),x,method=_RETURNVERBOSE)`output `1/2*x*(-3*x^2+12)^(1/2)+2*3^(1/2)*arcsin(1/2*x)`**3.95.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.03

$$\int \sqrt{12 - 3x^2} dx = \frac{1}{2} \sqrt{-3x^2 + 12}x - 2\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt{-3x^2 + 12}}{3x}\right)$$

input `integrate((-3*x^2+12)^(1/2),x, algorithm="fracas")`output `1/2*sqrt(-3*x^2 + 12)*x - 2*sqrt(3)*arctan(1/3*sqrt(3)*sqrt(-3*x^2 + 12)/x)`

3.95.6 Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.77

$$\int \sqrt{12 - 3x^2} dx = \frac{\sqrt{3}x\sqrt{4 - x^2}}{2} + 2\sqrt{3} \operatorname{asin}\left(\frac{x}{2}\right)$$

input `integrate((-3*x**2+12)**(1/2),x)`output `sqrt(3)*x*sqrt(4 - x**2)/2 + 2*sqrt(3)*asin(x/2)`**3.95.7 Maxima [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.63

$$\int \sqrt{12 - 3x^2} dx = \frac{1}{2} \sqrt{-3x^2 + 12}x + 2\sqrt{3} \arcsin\left(\frac{1}{2}x\right)$$

input `integrate((-3*x^2+12)^(1/2),x, algorithm="maxima")`output `1/2*sqrt(-3*x^2 + 12)*x + 2*sqrt(3)*arcsin(1/2*x)`**3.95.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.66

$$\int \sqrt{12 - 3x^2} dx = \frac{1}{2} \sqrt{3} \left(\sqrt{-x^2 + 4}x + 4 \arcsin\left(\frac{1}{2}x\right) \right)$$

input `integrate((-3*x^2+12)^(1/2),x, algorithm="giac")`output `1/2*sqrt(3)*(sqrt(-x^2 + 4)*x + 4*arcsin(1/2*x))`

3.95.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int \sqrt{12 - 3x^2} dx = 2\sqrt{3} \operatorname{asin}\left(\frac{x}{2}\right) + \frac{\sqrt{3}x\sqrt{4 - x^2}}{2}$$

input `int((12 - 3*x^2)^(1/2),x)`

output `2*3^(1/2)*asin(x/2) + (3^(1/2)*x*(4 - x^2)^(1/2))/2`

3.96 $\int \sec^5(x) \tan^3(x) dx$

3.96.1	Optimal result	574
3.96.2	Mathematica [A] (verified)	574
3.96.3	Rubi [A] (verified)	575
3.96.4	Maple [A] (verified)	576
3.96.5	Fricas [A] (verification not implemented)	577
3.96.6	Sympy [A] (verification not implemented)	577
3.96.7	Maxima [A] (verification not implemented)	577
3.96.8	Giac [A] (verification not implemented)	578
3.96.9	Mupad [B] (verification not implemented)	578

3.96.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \sec^5(x) \tan^3(x) dx = -\frac{1}{5} \sec^5(x) + \frac{\sec^7(x)}{7}$$

output `-1/5*sec(x)^5+1/7*sec(x)^7`

3.96.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sec^5(x) \tan^3(x) dx = -\frac{1}{5} \sec^5(x) + \frac{\sec^7(x)}{7}$$

input `Integrate[Sec[x]^5*Tan[x]^3,x]`

output `-1/5*Sec[x]^5 + Sec[x]^7/7`

3.96.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 3086, 25, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^3(x) \sec^5(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^3 \sec(x)^5 dx \\
 & \quad \downarrow \text{3086} \\
 & \int -\sec^4(x) (1 - \sec^2(x)) d \sec(x) \\
 & \quad \downarrow \text{25} \\
 & - \int \sec^4(x) (1 - \sec^2(x)) d \sec(x) \\
 & \quad \downarrow \text{244} \\
 & - \int (\sec^4(x) - \sec^6(x)) d \sec(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sec^7(x)}{7} - \frac{\sec^5(x)}{5}
 \end{aligned}$$

input `Int [Sec [x] ^5*Tan [x] ^3,x]`

output `-1/5*Sec [x] ^5 + Sec [x] ^7/7`

3.96.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

3.96.4 Maple [A] (verified)

Time = 3.79 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$-\frac{\sec(x)^5}{5} + \frac{\sec(x)^7}{7}$	14
default	$-\frac{\sec(x)^5}{5} + \frac{\sec(x)^7}{7}$	14
risch	$-\frac{32(7e^{9ix} - 6e^{7ix} + 7e^{5ix})}{35(e^{2ix} + 1)^7}$	34

input `int(sec(x)^5*tan(x)^3,x,method=_RETURNVERBOSE)`

output `-1/5*sec(x)^5+1/7*sec(x)^7`

3.96.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^5(x) \tan^3(x) dx = -\frac{7 \cos(x)^2 - 5}{35 \cos(x)^7}$$

input `integrate(sec(x)^5*tan(x)^3,x, algorithm="fracas")`output `-1/35*(7*cos(x)^2 - 5)/cos(x)^7`**3.96.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^5(x) \tan^3(x) dx = \frac{5 - 7 \cos^2(x)}{35 \cos^7(x)}$$

input `integrate(sec(x)**5*tan(x)**3,x)`output `(5 - 7*cos(x)**2)/(35*cos(x)**7)`**3.96.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^5(x) \tan^3(x) dx = -\frac{7 \cos(x)^2 - 5}{35 \cos(x)^7}$$

input `integrate(sec(x)^5*tan(x)^3,x, algorithm="maxima")`output `-1/35*(7*cos(x)^2 - 5)/cos(x)^7`

3.96.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sec^5(x) \tan^3(x) dx = -\frac{7 \cos(x)^2 - 5}{35 \cos(x)^7}$$

input `integrate(sec(x)^5*tan(x)^3,x, algorithm="giac")`

output `-1/35*(7*cos(x)^2 - 5)/cos(x)^7`

3.96.9 Mupad [B] (verification not implemented)

Time = 16.45 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sec^5(x) \tan^3(x) dx = \frac{1}{7 \cos(x)^7} - \frac{1}{5 \cos(x)^5}$$

input `int(tan(x)^3/cos(x)^5,x)`

output `1/(7*cos(x)^7) - 1/(5*cos(x)^5)`

3.97 $\int \frac{1}{1-\sin(x)} dx$

3.97.1	Optimal result	579
3.97.2	Mathematica [B] (verified)	579
3.97.3	Rubi [A] (verified)	580
3.97.4	Maple [A] (verified)	581
3.97.5	Fricas [A] (verification not implemented)	581
3.97.6	Sympy [A] (verification not implemented)	581
3.97.7	Maxima [A] (verification not implemented)	582
3.97.8	Giac [A] (verification not implemented)	582
3.97.9	Mupad [B] (verification not implemented)	582

3.97.1 Optimal result

Integrand size = 8, antiderivative size = 11

$$\int \frac{1}{1-\sin(x)} dx = \frac{\cos(x)}{1-\sin(x)}$$

output `cos(x)/(1-sin(x))`

3.97.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 25 vs. 2(11) = 22.

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 2.27

$$\int \frac{1}{1-\sin(x)} dx = \frac{2 \sin\left(\frac{x}{2}\right)}{\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)}$$

input `Integrate[(1 - Sin[x])^(-1),x]`

output `(2*Sin[x/2])/(Cos[x/2] - Sin[x/2])`

3.97.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{1 - \sin(x)} dx$$

↓ 3042

$$\int \frac{1}{1 - \sin(x)} dx$$

↓ 3127

$$\frac{\cos(x)}{1 - \sin(x)}$$

input `Int[(1 - Sin[x])^(-1),x]`

output `Cos[x]/(1 - Sin[x])`

3.97.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*SIN[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

3.97.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

method	result	size
default	$-\frac{2}{\tan(\frac{x}{2})-1}$	11
norman	$-\frac{2}{\tan(\frac{x}{2})-1}$	11
parallelrisch	$-\frac{2}{\tan(\frac{x}{2})-1}$	11
risch	$\frac{2}{e^{ix}-i}$	13

input `int(1/(1-sin(x)),x,method=_RETURNVERBOSE)`output `-2/(tan(1/2*x)-1)`**3.97.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.55

$$\int \frac{1}{1-\sin(x)} dx = \frac{\cos(x) + \sin(x) + 1}{\cos(x) - \sin(x) + 1}$$

input `integrate(1/(1-sin(x)),x, algorithm="fricas")`output `(cos(x) + sin(x) + 1)/(cos(x) - sin(x) + 1)`**3.97.6 Sympy [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \frac{1}{1-\sin(x)} dx = -\frac{2}{\tan(\frac{x}{2})-1}$$

input `integrate(1/(1-sin(x)),x)`output `-2/(tan(x/2) - 1)`

3.97.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.36

$$\int \frac{1}{1 - \sin(x)} dx = -\frac{2}{\frac{\sin(x)}{\cos(x)+1} - 1}$$

input `integrate(1/(1-sin(x)),x, algorithm="maxima")`output `-2/(sin(x)/(cos(x) + 1) - 1)`**3.97.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int \frac{1}{1 - \sin(x)} dx = -\frac{2}{\tan\left(\frac{1}{2}x\right) - 1}$$

input `integrate(1/(1-sin(x)),x, algorithm="giac")`output `-2/(tan(1/2*x) - 1)`**3.97.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int \frac{1}{1 - \sin(x)} dx = -\frac{2}{\tan\left(\frac{x}{2}\right) - 1}$$

input `int(-1/(sin(x) - 1),x)`output `-2/(tan(x/2) - 1)`

3.98 $\int \frac{1}{x\sqrt{-2+x^2}} dx$

3.98.1 Optimal result	583
3.98.2 Mathematica [A] (verified)	583
3.98.3 Rubi [A] (verified)	584
3.98.4 Maple [A] (verified)	585
3.98.5 Fricas [A] (verification not implemented)	585
3.98.6 Sympy [C] (verification not implemented)	586
3.98.7 Maxima [A] (verification not implemented)	586
3.98.8 Giac [A] (verification not implemented)	586
3.98.9 Mupad [B] (verification not implemented)	587

3.98.1 Optimal result

Integrand size = 13, antiderivative size = 22

$$\int \frac{1}{x\sqrt{-2+x^2}} dx = \frac{\arctan\left(\frac{\sqrt{-2+x^2}}{\sqrt{2}}\right)}{\sqrt{2}}$$

output `1/2*arctan(1/2*(x^2-2)^(1/2)*2^(1/2))*2^(1/2)`

3.98.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{-2+x^2}} dx = \frac{\arctan\left(\frac{\sqrt{-2+x^2}}{\sqrt{2}}\right)}{\sqrt{2}}$$

input `Integrate[1/(x*Sqrt[-2 + x^2]),x]`

output `ArcTan[Sqrt[-2 + x^2]/Sqrt[2]]/Sqrt[2]`

3.98.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {243, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x\sqrt{x^2-2}} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int \frac{1}{x^2\sqrt{x^2-2}} dx^2 \\ & \quad \downarrow \text{73} \\ & \int \frac{1}{x^4+2} d\sqrt{x^2-2} \\ & \quad \downarrow \text{216} \\ & \frac{\arctan\left(\frac{\sqrt{x^2-2}}{\sqrt{2}}\right)}{\sqrt{2}} \end{aligned}$$

input `Int[1/(x*Sqrt[-2 + x^2]),x]`

output `ArcTan[Sqrt[-2 + x^2]/Sqrt[2]]/Sqrt[2]`

3.98.3.1 Defintions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

3.98. $\int \frac{1}{x\sqrt{-2+x^2}} dx$

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

3.98.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}}{\sqrt{x^2-2}}\right)}{2}$	18
pseudoelliptic	$\frac{\arctan\left(\frac{\sqrt{x^2-2}\sqrt{2}}{2}\right)\sqrt{2}}{2}$	19
trager	$-\frac{\text{RootOf}\left(-Z^2+2\right) \ln\left(\frac{-\text{RootOf}\left(-Z^2+2\right)+\sqrt{x^2-2}}{x}\right)}{2}$	30
meijerg	$\frac{\sqrt{2} \sqrt{-\text{signum}\left(-1+\frac{x^2}{2}\right)} \left((-3 \ln(2)+2 \ln(x)+i\pi)\sqrt{\pi}-2\sqrt{\pi} \ln\left(\frac{1}{2}+\frac{\sqrt{1-\frac{x^2}{2}}}{2}\right) \right)}{4\sqrt{\pi} \sqrt{\text{signum}\left(-1+\frac{x^2}{2}\right)}}$	68

input `int(1/x/(x^2-2)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2*2^(1/2)*arctan(1/(x^2-2)^(1/2)*2^(1/2))`

3.98.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{1}{x\sqrt{-2+x^2}} dx = \sqrt{2} \arctan\left(-\frac{1}{2}\sqrt{2}x + \frac{1}{2}\sqrt{2}\sqrt{x^2-2}\right)$$

input `integrate(1/x/(x^2-2)^(1/2),x, algorithm="fricas")`

output `sqrt(2)*arctan(-1/2*sqrt(2)*x + 1/2*sqrt(2)*sqrt(x^2 - 2))`

3.98.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.50 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.77

$$\int \frac{1}{x\sqrt{-2+x^2}} dx = \begin{cases} \frac{\sqrt{2}i \operatorname{acosh}\left(\frac{\sqrt{2}}{x}\right)}{2} & \text{for } \frac{1}{|x^2|} > \frac{1}{2} \\ -\frac{\sqrt{2} \operatorname{asin}\left(\frac{\sqrt{2}}{x}\right)}{2} & \text{otherwise} \end{cases}$$

input `integrate(1/x/(x**2-2)**(1/2),x)`

output `Piecewise((sqrt(2)*I*acosh(sqrt(2)/x)/2, 1/Abs(x**2) > 1/2), (-sqrt(2)*asin(sqrt(2)/x)/2, True))`

3.98.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.64

$$\int \frac{1}{x\sqrt{-2+x^2}} dx = -\frac{1}{2} \sqrt{2} \arcsin\left(\frac{\sqrt{2}}{|x|}\right)$$

input `integrate(1/x/(x^2-2)^(1/2),x, algorithm="maxima")`

output `-1/2*sqrt(2)*arcsin(sqrt(2)/abs(x))`

3.98.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{1}{x\sqrt{-2+x^2}} dx = \frac{1}{2} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2} \sqrt{x^2-2}\right)$$

input `integrate(1/x/(x^2-2)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(2)*arctan(1/2*sqrt(2)*sqrt(x^2 - 2))`

3.98.9 Mupad [B] (verification not implemented)

Time = 17.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{1}{x\sqrt{-2+x^2}} dx = \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}\sqrt{x^2-2}}{2}\right)}{2}$$

input `int(1/(x*(x^2 - 2)^(1/2)),x)`

output `(2^(1/2)*atan((2^(1/2)*(x^2 - 2)^(1/2))/2))/2`

3.99 $\int \log(x^2) dx$

3.99.1	Optimal result	588
3.99.2	Mathematica [A] (verified)	588
3.99.3	Rubi [A] (verified)	589
3.99.4	Maple [A] (verified)	589
3.99.5	Fricas [A] (verification not implemented)	590
3.99.6	Sympy [A] (verification not implemented)	590
3.99.7	Maxima [A] (verification not implemented)	590
3.99.8	Giac [A] (verification not implemented)	591
3.99.9	Mupad [B] (verification not implemented)	591

3.99.1 Optimal result

Integrand size = 4, antiderivative size = 10

$$\int \log(x^2) dx = -2x + x \log(x^2)$$

output `-2*x+x*ln(x^2)`

3.99.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \log(x^2) dx = -2x + x \log(x^2)$$

input `Integrate[Log[x^2],x]`

output `-2*x + x*Log[x^2]`

3.99.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2732}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \log(x^2) dx$$

$$\downarrow \text{2732}$$

$$x \log(x^2) - 2x$$

input `Int [Log [x^2] , x]`

output `-2*x + x*Log [x^2]`

3.99.3.1 Defintions of rubi rules used

rule 2732 `Int [Log [(c_.)*(x_)^(n_.)], x_Symbol] :> Simp [x*Log [c*x^n], x] - Simp [n*x, x] /; FreeQ [{c, n}, x]`

3.99.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

method	result	size
default	$-2x + x \ln(x^2)$	11
norman	$-2x + x \ln(x^2)$	11
risch	$-2x + x \ln(x^2)$	11
parallelrisch	$-2x + x \ln(x^2)$	11
parts	$-2x + x \ln(x^2)$	11

input `int (ln(x^2) , x, method=_RETURNVERBOSE)`

output `-2*x+x*ln(x^2)`

3.99.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \log(x^2) dx = x \log(x^2) - 2x$$

input `integrate(log(x^2),x, algorithm="fricas")`

output `x*log(x^2) - 2*x`

3.99.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \log(x^2) dx = x \log(x^2) - 2x$$

input `integrate(ln(x**2),x)`

output `x*log(x**2) - 2*x`

3.99.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \log(x^2) dx = x \log(x^2) - 2x$$

input `integrate(log(x^2),x, algorithm="maxima")`

output `x*log(x^2) - 2*x`

3.99.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \log(x^2) dx = x \log(x^2) - 2x$$

input `integrate(log(x^2),x, algorithm="giac")`

output `x*log(x^2) - 2*x`

3.99.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \log(x^2) dx = x (\ln(x^2) - 2)$$

input `int(log(x^2),x)`

output `x*(log(x^2) - 2)`

3.100 $\int \sin(\sqrt[3]{x}) dx$

3.100.1 Optimal result	592
3.100.2 Mathematica [A] (verified)	592
3.100.3 Rubi [A] (verified)	593
3.100.4 Maple [A] (verified)	594
3.100.5 Fricas [A] (verification not implemented)	595
3.100.6 Sympy [A] (verification not implemented)	595
3.100.7 Maxima [A] (verification not implemented)	595
3.100.8 Giac [A] (verification not implemented)	596
3.100.9 Mupad [B] (verification not implemented)	596

3.100.1 Optimal result

Integrand size = 6, antiderivative size = 35

$$\int \sin(\sqrt[3]{x}) dx = 6 \cos(\sqrt[3]{x}) - 3x^{2/3} \cos(\sqrt[3]{x}) + 6\sqrt[3]{x} \sin(\sqrt[3]{x})$$

output `6*cos(x^(1/3))-3*x^(2/3)*cos(x^(1/3))+6*x^(1/3)*sin(x^(1/3))`

3.100.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

$$\int \sin(\sqrt[3]{x}) dx = -3(-2 + x^{2/3}) \cos(\sqrt[3]{x}) + 6\sqrt[3]{x} \sin(\sqrt[3]{x})$$

input `Integrate[Sin[x^(1/3)],x]`

output `-3*(-2 + x^(2/3))*Cos[x^(1/3)] + 6*x^(1/3)*Sin[x^(1/3)]`

3.100.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.333$, Rules used = {3842, 3042, 3777, 3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(\sqrt[3]{x}) \, dx \\
 & \quad \downarrow \text{3842} \\
 & 3 \int x^{2/3} \sin(\sqrt[3]{x}) \, d\sqrt[3]{x} \\
 & \quad \downarrow \text{3042} \\
 & 3 \int x^{2/3} \sin(\sqrt[3]{x}) \, d\sqrt[3]{x} \\
 & \quad \downarrow \text{3777} \\
 & 3 \left(2 \int \sqrt[3]{x} \cos(\sqrt[3]{x}) \, d\sqrt[3]{x} - x^{2/3} \cos(\sqrt[3]{x}) \right) \\
 & \quad \downarrow \text{3042} \\
 & 3 \left(2 \int \sqrt[3]{x} \sin\left(\sqrt[3]{x} + \frac{\pi}{2}\right) \, d\sqrt[3]{x} - x^{2/3} \cos(\sqrt[3]{x}) \right) \\
 & \quad \downarrow \text{3777} \\
 & 3 \left(2 \left(\int -\sin(\sqrt[3]{x}) \, d\sqrt[3]{x} + \sqrt[3]{x} \sin(\sqrt[3]{x}) \right) - x^{2/3} \cos(\sqrt[3]{x}) \right) \\
 & \quad \downarrow \text{25} \\
 & 3 \left(2 \left(\sqrt[3]{x} \sin(\sqrt[3]{x}) - \int \sin(\sqrt[3]{x}) \, d\sqrt[3]{x} \right) - x^{2/3} \cos(\sqrt[3]{x}) \right) \\
 & \quad \downarrow \text{3042} \\
 & 3 \left(2 \left(\sqrt[3]{x} \sin(\sqrt[3]{x}) - \int \sin(\sqrt[3]{x}) \, d\sqrt[3]{x} \right) - x^{2/3} \cos(\sqrt[3]{x}) \right) \\
 & \quad \downarrow \text{3118} \\
 & 3 \left(2 \left(\sqrt[3]{x} \sin(\sqrt[3]{x}) + \cos(\sqrt[3]{x}) \right) - x^{2/3} \cos(\sqrt[3]{x}) \right)
 \end{aligned}$$

input `Int[Sin[x^(1/3)],x]`

output `3*(-(x^(2/3)*Cos[x^(1/3)]) + 2*(Cos[x^(1/3)] + x^(1/3)*Sin[x^(1/3)]))`

3.100.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3842 `Int[((a_.) + (b_.)*Sin[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)])^(p_.), x_Symbol] := Simp[1/(n*f) Subst[Int[x^(1/n - 1)*(a + b*Sin[c + d*x])^p, x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]`

3.100.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

method	result	size
derivativedivides	$6 \cos\left(x^{\frac{1}{3}}\right) - 3x^{\frac{2}{3}} \cos\left(x^{\frac{1}{3}}\right) + 6x^{\frac{1}{3}} \sin\left(x^{\frac{1}{3}}\right)$	26
default	$6 \cos\left(x^{\frac{1}{3}}\right) - 3x^{\frac{2}{3}} \cos\left(x^{\frac{1}{3}}\right) + 6x^{\frac{1}{3}} \sin\left(x^{\frac{1}{3}}\right)$	26
meijerg	$12\sqrt{\pi} \left(-\frac{1}{2\sqrt{\pi}} + \frac{\left(-\frac{x^{\frac{2}{3}}}{2} + 1\right) \cos\left(x^{\frac{1}{3}}\right)}{2\sqrt{\pi}} + \frac{x^{\frac{1}{3}} \sin\left(x^{\frac{1}{3}}\right)}{2\sqrt{\pi}} \right)$	40

input `int(sin(x^(1/3)),x,method=_RETURNVERBOSE)`

output `6*cos(x^(1/3))-3*x^(2/3)*cos(x^(1/3))+6*x^(1/3)*sin(x^(1/3))`

3.100.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.60

$$\int \sin(\sqrt[3]{x}) dx = -3 \left(x^{\frac{2}{3}} - 2 \right) \cos \left(x^{\frac{1}{3}} \right) + 6 x^{\frac{1}{3}} \sin \left(x^{\frac{1}{3}} \right)$$

input `integrate(sin(x^(1/3)),x, algorithm="fricas")`

output `-3*(x^(2/3) - 2)*cos(x^(1/3)) + 6*x^(1/3)*sin(x^(1/3))`

3.100.6 Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int \sin(\sqrt[3]{x}) dx = -3x^{\frac{2}{3}} \cos(\sqrt[3]{x}) + 6\sqrt[3]{x} \sin(\sqrt[3]{x}) + 6 \cos(\sqrt[3]{x})$$

input `integrate(sin(x**(1/3)),x)`

output `-3*x**(2/3)*cos(x**(1/3)) + 6*x**(1/3)*sin(x**(1/3)) + 6*cos(x**(1/3))`

3.100.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.60

$$\int \sin(\sqrt[3]{x}) dx = -3 \left(x^{\frac{2}{3}} - 2 \right) \cos \left(x^{\frac{1}{3}} \right) + 6 x^{\frac{1}{3}} \sin \left(x^{\frac{1}{3}} \right)$$

input `integrate(sin(x^(1/3)),x, algorithm="maxima")`

output `-3*(x^(2/3) - 2)*cos(x^(1/3)) + 6*x^(1/3)*sin(x^(1/3))`

3.100.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.60

$$\int \sin(\sqrt[3]{x}) dx = -3 \left(x^{\frac{2}{3}} - 2 \right) \cos \left(x^{\frac{1}{3}} \right) + 6 x^{\frac{1}{3}} \sin \left(x^{\frac{1}{3}} \right)$$

input `integrate(sin(x^(1/3)),x, algorithm="giac")`output `-3*(x^(2/3) - 2)*cos(x^(1/3)) + 6*x^(1/3)*sin(x^(1/3))`**3.100.9 Mupad [B] (verification not implemented)**

Time = 16.28 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.60

$$\int \sin(\sqrt[3]{x}) dx = 6 x^{1/3} \sin(x^{1/3}) - 3 \cos(x^{1/3}) (x^{2/3} - 2)$$

input `int(sin(x^(1/3)),x)`output `6*x^(1/3)*sin(x^(1/3)) - 3*cos(x^(1/3))*(x^(2/3) - 2)`

3.101 $\int e^{1+x-x^2}(1-2x) dx$

3.101.1 Optimal result	597
3.101.2 Mathematica [A] (verified)	597
3.101.3 Rubi [A] (verified)	598
3.101.4 Maple [A] (verified)	598
3.101.5 Fricas [A] (verification not implemented)	599
3.101.6 Sympy [A] (verification not implemented)	599
3.101.7 Maxima [A] (verification not implemented)	599
3.101.8 Giac [A] (verification not implemented)	600
3.101.9 Mupad [B] (verification not implemented)	600

3.101.1 Optimal result

Integrand size = 16, antiderivative size = 10

$$\int e^{1+x-x^2}(1-2x) dx = e^{1+x-x^2}$$

output `exp(-x^2+x+1)`

3.101.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int e^{1+x-x^2}(1-2x) dx = e^{1+x-x^2}$$

input `Integrate[E^(1 + x - x^2)*(1 - 2*x), x]`

output `E^(1 + x - x^2)`

3.101.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.062$, Rules used = {2666}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{-x^2+x+1}(1-2x) dx$$

$$\downarrow \text{2666}$$

$$e^{-x^2+x+1}$$

input `Int[E^(1 + x - x^2)*(1 - 2*x),x]`

output `E^(1 + x - x^2)`

3.101.3.1 Defintions of rubi rules used

rule 2666 `Int[(F_)^((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)*((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[e*(F^(a + b*x + c*x^2)/(2*c*Log[F])), x] /; FreeQ[{F, a, b, c, d, e}, x] && EqQ[b*e - 2*c*d, 0]`

3.101.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

method	result	size
gospers	e^{-x^2+x+1}	10
derivativedivides	e^{-x^2+x+1}	10
default	e^{-x^2+x+1}	10
norman	e^{-x^2+x+1}	10
risch	e^{-x^2+x+1}	10
parallelrisch	e^{-x^2+x+1}	10
parts	$-\sqrt{\pi} e^{\frac{5}{4}} \operatorname{erf}\left(x - \frac{1}{2}\right) x + \frac{\sqrt{\pi} e^{\frac{5}{4}} \operatorname{erf}\left(x - \frac{1}{2}\right)}{2} + \frac{e^{\frac{5}{4}} \left(2 \operatorname{erf}\left(x - \frac{1}{2}\right) x \sqrt{\pi} - \operatorname{erf}\left(x - \frac{1}{2}\right) \sqrt{\pi} + 2 e^{-(x - \frac{1}{2})^2}\right)}{2}$	59

input `int(exp(-x^2+x+1)*(1-2*x),x,method=_RETURNVERBOSE)`

output `exp(-x^2+x+1)`

3.101.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int e^{1+x-x^2}(1-2x) dx = e^{(-x^2+x+1)}$$

input `integrate(exp(-x^2+x+1)*(1-2*x),x, algorithm="fricas")`

output `e^(-x^2 + x + 1)`

3.101.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int e^{1+x-x^2}(1-2x) dx = e^{-x^2+x+1}$$

input `integrate(exp(-x**2+x+1)*(1-2*x),x)`

output `exp(-x**2 + x + 1)`

3.101.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int e^{1+x-x^2}(1-2x) dx = e^{(-x^2+x+1)}$$

input `integrate(exp(-x^2+x+1)*(1-2*x),x, algorithm="maxima")`

output `e^(-x^2 + x + 1)`

3.101.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int e^{1+x-x^2}(1-2x) dx = e^{(-x^2+x+1)}$$

input `integrate(exp(-x^2+x+1)*(1-2*x),x, algorithm="giac")`output `e^(-x^2 + x + 1)`**3.101.9 Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int e^{1+x-x^2}(1-2x) dx = e e^{-x^2} e^x$$

input `int(-exp(x - x^2 + 1)*(2*x - 1),x)`output `exp(1)*exp(-x^2)*exp(x)`

3.102 $\int e^{\sqrt{x}} \sqrt{x} dx$

3.102.1 Optimal result	601
3.102.2 Mathematica [C] (verified)	601
3.102.3 Rubi [A] (verified)	602
3.102.4 Maple [A] (verified)	603
3.102.5 Fricas [A] (verification not implemented)	603
3.102.6 Sympy [A] (verification not implemented)	604
3.102.7 Maxima [A] (verification not implemented)	604
3.102.8 Giac [A] (verification not implemented)	604
3.102.9 Mupad [B] (verification not implemented)	605

3.102.1 Optimal result

Integrand size = 13, antiderivative size = 34

$$\int e^{\sqrt{x}} \sqrt{x} dx = 4e^{\sqrt{x}} - 4e^{\sqrt{x}} \sqrt{x} + 2e^{\sqrt{x}} x$$

output `4*exp(x^(1/2))-4*x^(1/2)*exp(x^(1/2))+2*exp(x^(1/2))*x`

3.102.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3 in optimal.

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.32

$$\int e^{\sqrt{x}} \sqrt{x} dx = 2\Gamma(3, -\sqrt{x})$$

input `Integrate[E^Sqrt[x]*Sqrt[x],x]`

output `2*Gamma[3, -Sqrt[x]]`

3.102.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {2645, 2607, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{\sqrt{x}} \sqrt{x} dx \\
 & \quad \downarrow \text{2645} \\
 & 2 \int e^{\sqrt{x}} x d\sqrt{x} \\
 & \quad \downarrow \text{2607} \\
 & 2 \left(e^{\sqrt{x}} x - 2 \int e^{\sqrt{x}} \sqrt{x} d\sqrt{x} \right) \\
 & \quad \downarrow \text{2607} \\
 & 2 \left(e^{\sqrt{x}} x - 2 \left(e^{\sqrt{x}} \sqrt{x} - \int e^{\sqrt{x}} d\sqrt{x} \right) \right) \\
 & \quad \downarrow \text{2624} \\
 & 2 \left(e^{\sqrt{x}} x - 2 \left(e^{\sqrt{x}} \sqrt{x} - e^{\sqrt{x}} \right) \right)
 \end{aligned}$$

input `Int[E^Sqrt[x]*Sqrt[x],x]`

output `2*(-2*(-E^Sqrt[x] + E^Sqrt[x]*Sqrt[x]) + E^Sqrt[x]*x)`

3.102.3.1 Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

```
rule 2624 Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;
FreeQ[{F, n}, x] && LinearQ[v, x]
```

```
rule 2645 Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((c_.) + (d_.)*(x_))^(m_
.), x_Symbol] :> With[{k = Denominator[n]}, Simp[k/d Subst[Int[x^(k*(m +
1) - 1)*F^(a + b*x^(k*n)), x], x, (c + d*x)^(1/k)], x]] /; FreeQ[{F, a, b,
c, d, m, n}, x] && IntegerQ[2*((m + 1)/n)] && LtQ[0, (m + 1)/n, 5] && !Int
egerQ[n]
```

3.102.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.56

method	result	size
meijerg	$-4 + \frac{2(3x-6\sqrt{x}+6)e^{\sqrt{x}}}{3}$	19
derivativedivides	$4e^{\sqrt{x}} - 4\sqrt{x}e^{\sqrt{x}} + 2e^{\sqrt{x}}x$	24
default	$4e^{\sqrt{x}} - 4\sqrt{x}e^{\sqrt{x}} + 2e^{\sqrt{x}}x$	24

```
input int(x^(1/2)*exp(x^(1/2)),x,method=_RETURNVERBOSE)
```

```
output -4+2/3*(3*x-6*x^(1/2)+6)*exp(x^(1/2))
```

3.102.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.41

$$\int e^{\sqrt{x}} \sqrt{x} dx = 2(x - 2\sqrt{x} + 2)e^{\sqrt{x}}$$

```
input integrate(x^(1/2)*exp(x^(1/2)),x, algorithm="fricas")
```

```
output 2*(x - 2*sqrt(x) + 2)*e^sqrt(x)
```

3.102.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.91

$$\int e^{\sqrt{x}} \sqrt{x} dx = -4\sqrt{x}e^{\sqrt{x}} + 2xe^{\sqrt{x}} + 4e^{\sqrt{x}}$$

input `integrate(x**(1/2)*exp(x**(1/2)),x)`output `-4*sqrt(x)*exp(sqrt(x)) + 2*x*exp(sqrt(x)) + 4*exp(sqrt(x))`**3.102.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.41

$$\int e^{\sqrt{x}} \sqrt{x} dx = 2(x - 2\sqrt{x} + 2)e^{\sqrt{x}}$$

input `integrate(x^(1/2)*exp(x^(1/2)),x, algorithm="maxima")`output `2*(x - 2*sqrt(x) + 2)*e^sqrt(x)`**3.102.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.41

$$\int e^{\sqrt{x}} \sqrt{x} dx = 2(x - 2\sqrt{x} + 2)e^{\sqrt{x}}$$

input `integrate(x^(1/2)*exp(x^(1/2)),x, algorithm="giac")`output `2*(x - 2*sqrt(x) + 2)*e^sqrt(x)`

3.102.9 Mupad [B] (verification not implemented)

Time = 17.72 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.68

$$\int e^{\sqrt{x}} \sqrt{x} dx = 4e^{\sqrt{x}} + 2xe^{\sqrt{x}} - 4\sqrt{x}e^{\sqrt{x}}$$

input `int(x^(1/2)*exp(x^(1/2)),x)`

output `4*exp(x^(1/2)) + 2*x*exp(x^(1/2)) - 4*x^(1/2)*exp(x^(1/2))`

3.103 $\int \cos(3x) \sin(2x) dx$

3.103.1 Optimal result	606
3.103.2 Mathematica [A] (verified)	606
3.103.3 Rubi [A] (verified)	607
3.103.4 Maple [A] (verified)	608
3.103.5 Fricas [A] (verification not implemented)	608
3.103.6 Sympy [B] (verification not implemented)	608
3.103.7 Maxima [A] (verification not implemented)	609
3.103.8 Giac [A] (verification not implemented)	609
3.103.9 Mupad [B] (verification not implemented)	609

3.103.1 Optimal result

Integrand size = 9, antiderivative size = 15

$$\int \cos(3x) \sin(2x) dx = \frac{\cos(x)}{2} - \frac{1}{10} \cos(5x)$$

output `1/2*cos(x)-1/10*cos(5*x)`

3.103.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \cos(3x) \sin(2x) dx = \frac{\cos(x)}{2} - \frac{1}{10} \cos(5x)$$

input `Integrate[Cos[3*x]*Sin[2*x],x]`

output `Cos[x]/2 - Cos[5*x]/10`

3.103.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4772}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sin(2x) \cos(3x) dx \\ \downarrow \text{3042} \\ \int \sin(2x) \cos(3x) dx \\ \downarrow \text{4772} \\ \frac{\cos(x)}{2} - \frac{1}{10} \cos(5x) \end{array}$$

input `Int[Cos[3*x]*Sin[2*x],x]`

output `Cos[x]/2 - Cos[5*x]/10`

3.103.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4772 `Int[cos[(c_.) + (d_.)*(x_)]*sin[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[-Cos[a - c + (b - d)*x]/(2*(b - d)), x] - Simp[Cos[a + c + (b + d)*x]/(2*(b + d)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

3.103.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

method	result	size
default	$\frac{\cos(x)}{2} - \frac{\cos(5x)}{10}$	12
risch	$\frac{\cos(x)}{2} - \frac{\cos(5x)}{10}$	12
parallelrisch	$\frac{\cos(x)}{2} - \frac{\cos(5x)}{10} - \frac{2}{5}$	13
norman	$\frac{-\frac{4 \tan(x)^2}{5} - \frac{4 \tan(\frac{3x}{2})^2}{5} + \frac{12 \tan(x) \tan(\frac{3x}{2})}{5}}{(1+\tan(x)^2)(1+\tan(\frac{3x}{2})^2)}$	43

input `int(sin(2*x)*cos(3*x),x,method=_RETURNVERBOSE)`output `1/2*cos(x)-1/10*cos(5*x)`**3.103.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \cos(3x) \sin(2x) dx = -\frac{8}{5} \cos(x)^5 + 2 \cos(x)^3$$

input `integrate(sin(2*x)*cos(3*x),x, algorithm="fracas")`output `-8/5*cos(x)^5 + 2*cos(x)^3`**3.103.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. 2(10) = 20.

Time = 0.13 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.73

$$\int \cos(3x) \sin(2x) dx = \frac{3 \sin(2x) \sin(3x)}{5} + \frac{2 \cos(2x) \cos(3x)}{5}$$

input `integrate(sin(2*x)*cos(3*x),x)`output `3*sin(2*x)*sin(3*x)/5 + 2*cos(2*x)*cos(3*x)/5`

3.103.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \cos(3x) \sin(2x) dx = -\frac{1}{10} \cos(5x) + \frac{1}{2} \cos(x)$$

input `integrate(sin(2*x)*cos(3*x),x, algorithm="maxima")`output `-1/10*cos(5*x) + 1/2*cos(x)`**3.103.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \cos(3x) \sin(2x) dx = -\frac{1}{10} \cos(5x) + \frac{1}{2} \cos(x)$$

input `integrate(sin(2*x)*cos(3*x),x, algorithm="giac")`output `-1/10*cos(5*x) + 1/2*cos(x)`**3.103.9 Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \cos(3x) \sin(2x) dx = 2 \cos(x)^3 - \frac{8 \cos(x)^5}{5}$$

input `int(cos(3*x)*sin(2*x),x)`output `2*cos(x)^3 - (8*cos(x)^5)/5`

3.104 $\int (1 + 2 \sin(x)) dx$

3.104.1 Optimal result	610
3.104.2 Mathematica [A] (verified)	610
3.104.3 Rubi [A] (verified)	611
3.104.4 Maple [A] (verified)	611
3.104.5 Fricas [A] (verification not implemented)	612
3.104.6 Sympy [A] (verification not implemented)	612
3.104.7 Maxima [A] (verification not implemented)	612
3.104.8 Giac [A] (verification not implemented)	613
3.104.9 Mupad [B] (verification not implemented)	613

3.104.1 Optimal result

Integrand size = 6, antiderivative size = 6

$$\int (1 + 2 \sin(x)) dx = x - 2 \cos(x)$$

output `x-2*cos(x)`

3.104.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int (1 + 2 \sin(x)) dx = x - 2 \cos(x)$$

input `Integrate[1 + 2*Sin[x],x]`

output `x - 2*Cos[x]`

3.104.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (2 \sin(x) + 1) dx$$

$$\downarrow \text{2009}$$

$$x - 2 \cos(x)$$

input `Int[1 + 2*Sin[x],x]`

output `x - 2*Cos[x]`

3.104.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.104.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

method	result	size
default	$x - 2 \cos(x)$	7
risch	$x - 2 \cos(x)$	7
parts	$x - 2 \cos(x)$	7
parallelrisch	$-2 - 2 \cos(x) + x$	8
norman	$\frac{x + x \tan(\frac{x}{2})^2 - 4}{1 + \tan(\frac{x}{2})^2}$	23

input `int(1+2*sin(x),x,method=_RETURNVERBOSE)`

output `x-2*cos(x)`

3.104.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int (1 + 2 \sin(x)) dx = x - 2 \cos(x)$$

input `integrate(1+2*sin(x),x, algorithm="fricas")`

output `x - 2*cos(x)`

3.104.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int (1 + 2 \sin(x)) dx = x - 2 \cos(x)$$

input `integrate(1+2*sin(x),x)`

output `x - 2*cos(x)`

3.104.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int (1 + 2 \sin(x)) dx = x - 2 \cos(x)$$

input `integrate(1+2*sin(x),x, algorithm="maxima")`

output `x - 2*cos(x)`

3.104.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int (1 + 2 \sin(x)) dx = x - 2 \cos(x)$$

input `integrate(1+2*sin(x),x, algorithm="giac")`

output `x - 2*cos(x)`

3.104.9 Mupad [B] (verification not implemented)

Time = 18.42 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int (1 + 2 \sin(x)) dx = x - 2 \cos(x)$$

input `int(2*sin(x) + 1,x)`

output `x - 2*cos(x)`

3.105 $\int (1 - x)^{2014} x dx$

3.105.1 Optimal result	614
3.105.2 Mathematica [B] (verified)	614
3.105.3 Rubi [A] (verified)	615
3.105.4 Maple [B] (verified)	616
3.105.5 Fricas [F(-2)]	616
3.105.6 Sympy [B] (verification not implemented)	616
3.105.7 Maxima [B] (verification not implemented)	617
3.105.8 Giac [B] (verification not implemented)	618
3.105.9 Mupad [B] (verification not implemented)	619

3.105.1 Optimal result

Integrand size = 9, antiderivative size = 23

$$\int (1 - x)^{2014} x dx = -\frac{(1 - x)^{2015}}{2015} + \frac{(1 - x)^{2016}}{2016}$$

output `-1/2015*(1-x)^2015+1/2016*(1-x)^2016`

3.105.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 12138 vs. 2(23) = 46.

Time = 0.13 (sec) , antiderivative size = 12138, normalized size of antiderivative = 527.74

$$\int (1 - x)^{2014} x dx = \text{Result too large to show}$$

input `Integrate[(1 - x)^2014*x,x]`

output `Result too large to show`

3.105.3 Rubi [A] (verified)

Time = 0.87 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (1-x)^{2014} x dx$$

$$\downarrow 49$$

$$\int ((1-x)^{2014} - (1-x)^{2015}) dx$$

$$\downarrow 2009$$

$$\frac{(1-x)^{2016}}{2016} - \frac{(1-x)^{2015}}{2015}$$

input `Int[(1 - x)^2014*x,x]`

output `-1/2015*(1 - x)^2015 + (1 - x)^2016/2016`

3.105.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.105.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 10075 vs. $2(19) = 38$.

Time = 7.62 (sec) , antiderivative size = 10076, normalized size of antiderivative = 438.09

method	result	size
gospers	Expression too large to display	10076
default	Expression too large to display	10077
risch	Expression too large to display	10077
parallelrisch	Expression too large to display	10077

input `int(x*(1-x)^2014,x,method=_RETURNVERBOSE)`

output `result too large to display`

3.105.5 Fracas [F(-2)]

Exception generated.

$$\int (1-x)^{2014} x dx = \text{Exception raised: RuntimeError}$$

input `integrate(x*(1-x)^2014,x, algorithm="fricas")`

output `Exception raised: RuntimeError >> System error: Heap exhausted (no more space for allocation).2293760 bytes available, 2557536 requested.PROCEED WITH CAUTION.`

3.105.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12024 vs. $2(12) = 24$.

Time = 2.41 (sec) , antiderivative size = 12024, normalized size of antiderivative = 522.78

$$\int (1-x)^{2014} x dx = \text{Too large to display}$$

input `integrate(x*(1-x)**2014,x)`

output `x**2016/2016 - 2014*x**2015/2015 + 2013*x**2014/2 - 2026084*x**2013/3 + 1358826667*x**2012/4 - 136629987582*x**2011 + 1373131035323509*x**2010/30 - 91953985549170536*x**2009/7 + 26377651026988133103*x**2008/8 - 6617491558542444915874*x**2007/9 + 294992994835264731661117*x**2006/2 - 134422910799097740606580164*x**2005/5 + 53876689232818214844524454823*x**2004/12 - 691762702790623489451562620638*x**2003 + 2571973087266166342850029070691063*x**2002/26 - 39588591248274824267756569403940400*x**2001/3 + 131961937837090040663501674882660163383*x**2000/80 - 193964593913505209402992927651733959950*x**1999 + 387541161559807075301489477559812273935075*x**1998/18 - 2262922530915969121458419278661196219696900*x**1997 + 903358447595910292527771972191922410542659825*x**1996/4 - 21454757739792727450218123378990236305630977010*x**1995 + 3889161470317168646281025630946632817062982660525*x**1994/2 - 168502105628944447818705199670044187130493533353400*x**1993 + 111885369941148961252482398045233357984870743212666075*x**1992/8 - 1113818576786312843600553552092286915421060813044865230*x**1991 + 170499877545918079073432876733170209565984828166917578371*x**1990/2 - 734769036554419859459683303160816263037237362099897570964140*x**1989/117 + 1783541483599579507954055928245847409394570340210724349640535*x**1988/4 - 30550827847475767604775649853836796394905428348789090875481890*x**1987 + 12134785744398256314996786936077927810352871698504752093214631735*x**1986/6 - 129502922523593280...`

3.105.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10076 vs. $2(15) = 30$.

Time = 2.29 (sec) , antiderivative size = 10076, normalized size of antiderivative = 438.09

$$\int (1-x)^{2014} x dx = \text{Too large to display}$$

input `integrate(x*(1-x)^2014,x, algorithm="maxima")`

```
output 1/2016*x^2016 - 2014/2015*x^2015 + 2013/2*x^2014 - 2026084/3*x^2013 + 1358
826667/4*x^2012 - 136629987582*x^2011 + 1373131035323509/30*x^2010 - 91953
985549170536/7*x^2009 + 26377651026988133103/8*x^2008 - 661749155854244491
5874/9*x^2007 + 294992994835264731661117/2*x^2006 - 1344229107990977406065
80164/5*x^2005 + 53876689232818214844524454823/12*x^2004 - 691762702790623
489451562620638*x^2003 + 2571973087266166342850029070691063/26*x^2002 - 39
588591248274824267756569403940400/3*x^2001 + 13196193783709004066350167488
2660163383/80*x^2000 - 193964593913505209402992927651733959950*x^1999 + 38
7541161559807075301489477559812273935075/18*x^1998 - 226292253091596912145
8419278661196219696900*x^1997 + 903358447595910292527771972191922410542659
825/4*x^1996 - 21454757739792727450218123378990236305630977010*x^1995 + 38
89161470317168646281025630946632817062982660525/2*x^1994 - 168502105628944
447818705199670044187130493533353400*x^1993 + 1118853699411489612524823980
45233357984870743212666075/8*x^1992 - 111381857678631284360055355209228691
5421060813044865230*x^1991 + 170499877545918079073432876733170209565984828
166917578371/2*x^1990 - 73476903655441985945968330316081626303723736209989
7570964140/117*x^1989 + 17835414835995795079540559282458474093945703402107
24349640535/4*x^1988 - 305508278474757676047756498538367963949054283487890
90875481890*x^1987 + 12134785744398256314996786936077927810352871698504752
093214631735/6*x^1986 - 12950292252359328049213699468709059079824445040...
```

3.105.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10076 vs. $2(15) = 30$.

Time = 1.64 (sec) , antiderivative size = 10076, normalized size of antiderivative = 438.09

$$\int (1-x)^{2014} x dx = \text{Too large to display}$$

```
input integrate(x*(1-x)^2014,x, algorithm="giac")
```

output

```

1/2016*x^2016 - 2014/2015*x^2015 + 2013/2*x^2014 - 2026084/3*x^2013 + 1358
826667/4*x^2012 - 136629987582*x^2011 + 1373131035323509/30*x^2010 - 91953
985549170536/7*x^2009 + 26377651026988133103/8*x^2008 - 661749155854244491
5874/9*x^2007 + 294992994835264731661117/2*x^2006 - 1344229107990977406065
80164/5*x^2005 + 53876689232818214844524454823/12*x^2004 - 691762702790623
489451562620638*x^2003 + 2571973087266166342850029070691063/26*x^2002 - 39
588591248274824267756569403940400/3*x^2001 + 13196193783709004066350167488
2660163383/80*x^2000 - 193964593913505209402992927651733959950*x^1999 + 38
7541161559807075301489477559812273935075/18*x^1998 - 226292253091596912145
8419278661196219696900*x^1997 + 903358447595910292527771972191922410542659
825/4*x^1996 - 21454757739792727450218123378990236305630977010*x^1995 + 38
89161470317168646281025630946632817062982660525/2*x^1994 - 168502105628944
447818705199670044187130493533353400*x^1993 + 1118853699411489612524823980
45233357984870743212666075/8*x^1992 - 111381857678631284360055355209228691
5421060813044865230*x^1991 + 170499877545918079073432876733170209565984828
166917578371/2*x^1990 - 73476903655441985945968330316081626303723736209989
7570964140/117*x^1989 + 17835414835995795079540559282458474093945703402107
24349640535/4*x^1988 - 305508278474757676047756498538367963949054283487890
90875481890*x^1987 + 12134785744398256314996786936077927810352871698504752
093214631735/6*x^1986 - 12950292252359328049213699468709059079824445040...

```

3.105.9 Mupad [B] (verification not implemented)

Time = 34.75 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int (1-x)^{2014} x dx = \frac{(x-1)^{2015}}{2015} + \frac{(x-1)^{2016}}{2016}$$

input `int(x*(x - 1)^2014,x)`

output `(x - 1)^2015/2015 + (x - 1)^2016/2016`

3.106 $\int \operatorname{arcsinh}(x) dx$

3.106.1 Optimal result	620
3.106.2 Mathematica [A] (verified)	620
3.106.3 Rubi [A] (verified)	621
3.106.4 Maple [A] (verified)	622
3.106.5 Fricas [A] (verification not implemented)	622
3.106.6 Sympy [A] (verification not implemented)	622
3.106.7 Maxima [A] (verification not implemented)	623
3.106.8 Giac [A] (verification not implemented)	623
3.106.9 Mupad [B] (verification not implemented)	623

3.106.1 Optimal result

Integrand size = 2, antiderivative size = 16

$$\int \operatorname{arcsinh}(x) dx = -\sqrt{1+x^2} + x \operatorname{arcsinh}(x)$$

output `x*arcsinh(x)-(x^2+1)^(1/2)`

3.106.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \operatorname{arcsinh}(x) dx = -\sqrt{1+x^2} + x \operatorname{arcsinh}(x)$$

input `Integrate[ArcSinh[x],x]`

output `-Sqrt[1 + x^2] + x*ArcSinh[x]`

3.106.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {6187, 241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arcsinh}(x) dx$$

$$\downarrow \text{6187}$$

$$x \operatorname{arcsinh}(x) - \int \frac{x}{\sqrt{x^2 + 1}} dx$$

$$\downarrow \text{241}$$

$$x \operatorname{arcsinh}(x) - \sqrt{x^2 + 1}$$

input `Int[ArcSinh[x], x]`

output `-Sqrt[1 + x^2] + x*ArcSinh[x]`

3.106.3.1 Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

rule 6187 `Int[((a_) + ArcSinh[(c_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[x*(a + b*ArcSinh[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcSinh[c*x])^(n - 1)/Sqrt[1 + c^2*x^2]), x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

3.106.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result	size
lookup	$x \operatorname{arcsinh}(x) - \sqrt{x^2 + 1}$	15
default	$x \operatorname{arcsinh}(x) - \sqrt{x^2 + 1}$	15
parts	$x \operatorname{arcsinh}(x) - \sqrt{x^2 + 1}$	15

input `int(arcsinh(x),x,method=_RETURNVERBOSE)`output `x*arcsinh(x)-(x^2+1)^(1/2)`**3.106.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int \operatorname{arcsinh}(x) dx = x \log(x + \sqrt{x^2 + 1}) - \sqrt{x^2 + 1}$$

input `integrate(arcsinh(x),x, algorithm="fricas")`output `x*log(x + sqrt(x^2 + 1)) - sqrt(x^2 + 1)`**3.106.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \operatorname{arcsinh}(x) dx = x \operatorname{asinh}(x) - \sqrt{x^2 + 1}$$

input `integrate(asinh(x),x)`output `x*asinh(x) - sqrt(x**2 + 1)`

3.106.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \operatorname{arcsinh}(x) dx = x \operatorname{arsinh}(x) - \sqrt{x^2 + 1}$$

input `integrate(arcsinh(x),x, algorithm="maxima")`output `x*arcsinh(x) - sqrt(x^2 + 1)`**3.106.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int \operatorname{arcsinh}(x) dx = x \log(x + \sqrt{x^2 + 1}) - \sqrt{x^2 + 1}$$

input `integrate(arcsinh(x),x, algorithm="giac")`output `x*log(x + sqrt(x^2 + 1)) - sqrt(x^2 + 1)`**3.106.9 Mupad [B] (verification not implemented)**

Time = 19.88 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \operatorname{arcsinh}(x) dx = x \operatorname{asinh}(x) - \sqrt{x^2 + 1}$$

input `int(asinh(x),x)`output `x*asinh(x) - (x^2 + 1)^(1/2)`

3.107 $\int \frac{x^2}{-1+x} dx$

3.107.1 Optimal result	624
3.107.2 Mathematica [A] (verified)	624
3.107.3 Rubi [A] (verified)	625
3.107.4 Maple [A] (verified)	626
3.107.5 Fricas [A] (verification not implemented)	626
3.107.6 Sympy [A] (verification not implemented)	626
3.107.7 Maxima [A] (verification not implemented)	627
3.107.8 Giac [A] (verification not implemented)	627
3.107.9 Mupad [B] (verification not implemented)	627

3.107.1 Optimal result

Integrand size = 9, antiderivative size = 15

$$\int \frac{x^2}{-1+x} dx = x + \frac{x^2}{2} + \log(1-x)$$

output `x+1/2*x^2+ln(1-x)`

3.107.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

$$\int \frac{x^2}{-1+x} dx = -\frac{3}{2} + x + \frac{x^2}{2} + \log(-1+x)$$

input `Integrate[x^2/(-1 + x),x]`

output `-3/2 + x + x^2/2 + Log[-1 + x]`

3.107.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{x-1} dx$$

$$\downarrow 49$$

$$\int \left(x + \frac{1}{x-1} + 1 \right) dx$$

$$\downarrow 2009$$

$$\frac{x^2}{2} + x + \log(1-x)$$

input `Int[x^2/(-1 + x), x]`

output `x + x^2/2 + Log[1 - x]`

3.107.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.107.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

method	result	size
default	$x + \frac{x^2}{2} + \ln(-1 + x)$	12
norman	$x + \frac{x^2}{2} + \ln(-1 + x)$	12
risch	$x + \frac{x^2}{2} + \ln(-1 + x)$	12
parallelrisch	$x + \frac{x^2}{2} + \ln(-1 + x)$	12
meijerg	$\frac{x(3x+6)}{6} + \ln(1 - x)$	16

input `int(x^2/(-1+x),x,method=_RETURNVERBOSE)`output `x+1/2*x^2+ln(-1+x)`**3.107.5 Fricas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \frac{x^2}{-1+x} dx = \frac{1}{2}x^2 + x + \log(x-1)$$

input `integrate(x^2/(-1+x),x, algorithm="fricas")`output `1/2*x^2 + x + log(x - 1)`**3.107.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int \frac{x^2}{-1+x} dx = \frac{x^2}{2} + x + \log(x-1)$$

input `integrate(x**2/(-1+x),x)`output `x**2/2 + x + log(x - 1)`

3.107.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \frac{x^2}{-1+x} dx = \frac{1}{2}x^2 + x + \log(x-1)$$

input `integrate(x^2/(-1+x),x, algorithm="maxima")`output `1/2*x^2 + x + log(x - 1)`**3.107.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \frac{x^2}{-1+x} dx = \frac{1}{2}x^2 + x + \log(|x-1|)$$

input `integrate(x^2/(-1+x),x, algorithm="giac")`output `1/2*x^2 + x + log(abs(x - 1))`**3.107.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \frac{x^2}{-1+x} dx = x + \ln(x-1) + \frac{x^2}{2}$$

input `int(x^2/(x - 1),x)`output `x + log(x - 1) + x^2/2`

3.108 $\int x \arctan(x) dx$

3.108.1 Optimal result	628
3.108.2 Mathematica [A] (verified)	628
3.108.3 Rubi [A] (verified)	629
3.108.4 Maple [A] (verified)	630
3.108.5 Fricas [A] (verification not implemented)	630
3.108.6 Sympy [A] (verification not implemented)	631
3.108.7 Maxima [A] (verification not implemented)	631
3.108.8 Giac [A] (verification not implemented)	631
3.108.9 Mupad [B] (verification not implemented)	632

3.108.1 Optimal result

Integrand size = 4, antiderivative size = 21

$$\int x \arctan(x) dx = -\frac{x}{2} + \frac{\arctan(x)}{2} + \frac{1}{2}x^2 \arctan(x)$$

output `-1/2*x+1/2*arctan(x)+1/2*x^2*arctan(x)`

3.108.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.76

$$\int x \arctan(x) dx = \frac{1}{2}(-x + (1 + x^2) \arctan(x))$$

input `Integrate[x*ArcTan[x],x]`

output `(-x + (1 + x^2)*ArcTan[x])/2`

3.108.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.95, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5361, 262, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \arctan(x) dx$$

$$\downarrow \text{5361}$$

$$\frac{1}{2}x^2 \arctan(x) - \frac{1}{2} \int \frac{x^2}{x^2 + 1} dx$$

$$\downarrow \text{262}$$

$$\frac{1}{2} \left(\int \frac{1}{x^2 + 1} dx - x \right) + \frac{1}{2}x^2 \arctan(x)$$

$$\downarrow \text{216}$$

$$\frac{1}{2}x^2 \arctan(x) + \frac{1}{2}(\arctan(x) - x)$$

input `Int[x*ArcTan[x],x]`

output `(x^2*ArcTan[x])/2 + (-x + ArcTan[x])/2`

3.108.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 262 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m-1)*((a + b*x^2)^(p+1)/(b*(m+2*p+1))), x] - Simp[a*c^2*((m-1)/(b*(m+2*p+1)) Int[(c*x)^(m-2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2-1] && NeQ[m+2*p+1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

```
rule 5361 Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :>
  Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m +
  1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x],
  x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] &
  & IntegerQ[m])) && NeQ[m, -1]
```

3.108.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.76

method	result	size
default	$-\frac{x}{2} + \frac{\arctan(x)}{2} + \frac{x^2 \arctan(x)}{2}$	16
meijerg	$-\frac{x}{2} + \frac{(3x^2+3) \arctan(x)}{6}$	16
parallelrisch	$-\frac{x}{2} + \frac{\arctan(x)}{2} + \frac{x^2 \arctan(x)}{2}$	16
parts	$-\frac{x}{2} + \frac{\arctan(x)}{2} + \frac{x^2 \arctan(x)}{2}$	16
risch	$-\frac{ix^2 \ln(ix+1)}{4} + \frac{ix^2 \ln(-ix+1)}{4} - \frac{x}{2} + \frac{\arctan(x)}{2}$	35

```
input int(x*arctan(x), x, method=_RETURNVERBOSE)
```

```
output -1/2*x+1/2*arctan(x)+1/2*x^2*arctan(x)
```

3.108.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int x \arctan(x) dx = \frac{1}{2} (x^2 + 1) \arctan(x) - \frac{1}{2} x$$

```
input integrate(x*arctan(x), x, algorithm="fricas")
```

```
output 1/2*(x^2 + 1)*arctan(x) - 1/2*x
```

3.108.6 Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int x \arctan(x) dx = \frac{x^2 \operatorname{atan}(x)}{2} - \frac{x}{2} + \frac{\operatorname{atan}(x)}{2}$$

input `integrate(x*atan(x),x)`output `x**2*atan(x)/2 - x/2 + atan(x)/2`**3.108.7 Maxima [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int x \arctan(x) dx = \frac{1}{2} x^2 \arctan(x) - \frac{1}{2} x + \frac{1}{2} \arctan(x)$$

input `integrate(x*arctan(x),x, algorithm="maxima")`output `1/2*x^2*arctan(x) - 1/2*x + 1/2*arctan(x)`**3.108.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int x \arctan(x) dx = \frac{1}{2} x^2 \arctan(x) - \frac{1}{2} x + \frac{1}{2} \arctan(x)$$

input `integrate(x*arctan(x),x, algorithm="giac")`output `1/2*x^2*arctan(x) - 1/2*x + 1/2*arctan(x)`

3.108.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

$$\int x \arctan(x) dx = \operatorname{atan}(x) \left(\frac{x^2}{2} + \frac{1}{2} \right) - \frac{x}{2}$$

input `int(x*atan(x),x)`

output `atan(x)*(x^2/2 + 1/2) - x/2`

3.109 $\int \frac{1}{-2014-15x+x^2} dx$

3.109.1 Optimal result	633
3.109.2 Mathematica [A] (verified)	633
3.109.3 Rubi [A] (verified)	634
3.109.4 Maple [A] (verified)	635
3.109.5 Fricas [A] (verification not implemented)	635
3.109.6 Sympy [A] (verification not implemented)	635
3.109.7 Maxima [A] (verification not implemented)	636
3.109.8 Giac [A] (verification not implemented)	636
3.109.9 Mupad [B] (verification not implemented)	636

3.109.1 Optimal result

Integrand size = 10, antiderivative size = 19

$$\int \frac{1}{-2014 - 15x + x^2} dx = \frac{1}{91} \log(53 - x) - \frac{1}{91} \log(38 + x)$$

output `1/91*ln(53-x)-1/91*ln(38+x)`

3.109.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{1}{-2014 - 15x + x^2} dx = \frac{1}{91} \log(53 - x) - \frac{1}{91} \log(38 + x)$$

input `Integrate[(-2014 - 15*x + x^2)^(-1), x]`

output `Log[53 - x]/91 - Log[38 + x]/91`

3.109.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 - 15x - 2014} dx$$

↓ 1081

$$\int \left(-\frac{1}{91(x+38)} - \frac{1}{91(53-x)} \right) dx$$

↓ 2009

$$\frac{1}{91} \log(53-x) - \frac{1}{91} \log(x+38)$$

input `Int[(-2014 - 15*x + x^2)^(-1), x]`

output `Log[53 - x]/91 - Log[38 + x]/91`

3.109.3.1 Defintions of rubi rules used

rule 1081 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.109.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

method	result	size
default	$\frac{\ln(x-53)}{91} - \frac{\ln(38+x)}{91}$	14
norman	$\frac{\ln(x-53)}{91} - \frac{\ln(38+x)}{91}$	14
risch	$\frac{\ln(x-53)}{91} - \frac{\ln(38+x)}{91}$	14
parallelrisch	$\frac{\ln(x-53)}{91} - \frac{\ln(38+x)}{91}$	14

input `int(1/(x^2-15*x-2014),x,method=_RETURNVERBOSE)`output `1/91*ln(x-53)-1/91*ln(38+x)`**3.109.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int \frac{1}{-2014 - 15x + x^2} dx = -\frac{1}{91} \log(x + 38) + \frac{1}{91} \log(x - 53)$$

input `integrate(1/(x^2-15*x-2014),x, algorithm="fricas")`output `-1/91*log(x + 38) + 1/91*log(x - 53)`**3.109.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{1}{-2014 - 15x + x^2} dx = \frac{\log(x - 53)}{91} - \frac{\log(x + 38)}{91}$$

input `integrate(1/(x**2-15*x-2014),x)`output `log(x - 53)/91 - log(x + 38)/91`

3.109.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int \frac{1}{-2014 - 15x + x^2} dx = -\frac{1}{91} \log(x + 38) + \frac{1}{91} \log(x - 53)$$

input `integrate(1/(x^2-15*x-2014),x, algorithm="maxima")`output `-1/91*log(x + 38) + 1/91*log(x - 53)`**3.109.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{-2014 - 15x + x^2} dx = -\frac{1}{91} \log(|x + 38|) + \frac{1}{91} \log(|x - 53|)$$

input `integrate(1/(x^2-15*x-2014),x, algorithm="giac")`output `-1/91*log(abs(x + 38)) + 1/91*log(abs(x - 53))`**3.109.9 Mupad [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.42

$$\int \frac{1}{-2014 - 15x + x^2} dx = -\frac{2 \operatorname{atanh}\left(\frac{2x}{91} - \frac{15}{91}\right)}{91}$$

input `int(-1/(15*x - x^2 + 2014),x)`output `-(2*atanh((2*x)/91 - 15/91))/91`

3.110 $\int e^x(-2(1+x)\arctan(x) + \log(1+x^2)) dx$

3.110.1 Optimal result	637
3.110.2 Mathematica [A] (verified)	637
3.110.3 Rubi [A] (verified)	638
3.110.4 Maple [A] (verified)	638
3.110.5 Fricas [A] (verification not implemented)	639
3.110.6 Sympy [A] (verification not implemented)	639
3.110.7 Maxima [A] (verification not implemented)	639
3.110.8 Giac [A] (verification not implemented)	640
3.110.9 Mupad [B] (verification not implemented)	640

3.110.1 Optimal result

Integrand size = 18, antiderivative size = 19

$$\int e^x(-2(1+x)\arctan(x) + \log(1+x^2)) dx = -2e^x x \arctan(x) + e^x \log(1+x^2)$$

output `-2*exp(x)*x*arctan(x)+exp(x)*ln(x^2+1)`

3.110.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int e^x(-2(1+x)\arctan(x) + \log(1+x^2)) dx = e^x(-2x \arctan(x) + \log(1+x^2))$$

input `Integrate[E^x*(-2*(1+x)*ArcTan[x] + Log[1+x^2]),x]`

output `E^x*(-2*x*ArcTan[x] + Log[1+x^2])`

3.110.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.47, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x (\log(x^2 + 1) - 2(x + 1) \arctan(x)) dx$$

$$\downarrow \text{7293}$$

$$\int (e^x \log(x^2 + 1) - 2e^x(x + 1) \arctan(x)) dx$$

$$\downarrow \text{2009}$$

$$2e^x \arctan(x) - 2e^x(x + 1) \arctan(x) + e^x \log(x^2 + 1)$$

input `Int[E^x*(-2*(1 + x)*ArcTan[x] + Log[1 + x^2]),x]`

output `2*E^x*ArcTan[x] - 2*E^x*(1 + x)*ArcTan[x] + E^x*Log[1 + x^2]`

3.110.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

3.110.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

method	result
parallelrisch	$-2e^x x \arctan(x) + e^x \ln(x^2 + 1)$
risch	$\frac{i\pi \operatorname{csgn}(i(i+x)) \operatorname{csgn}(i(x-i)(i+x))^2 e^x}{2} + e^x \ln(x - i) - i(i + x) e^x \ln(i + x) - \frac{i\pi \operatorname{csgn}(i(x-i)) \operatorname{csgn}(i(i+x))}{2}$

3.110. $\int e^x (-2(1 + x) \arctan(x) + \log(1 + x^2)) dx$

input `int(exp(x)*(ln(x^2+1)-2*(1+x)*arctan(x)),x,method=_RETURNVERBOSE)`

output `-2*exp(x)*x*arctan(x)+exp(x)*ln(x^2+1)`

3.110.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int e^x(-2(1+x)\arctan(x) + \log(1+x^2)) dx = -2x\arctan(x)e^x + e^x\log(x^2+1)$$

input `integrate(exp(x)*(log(x^2+1)-2*(1+x)*arctan(x)),x, algorithm="fricas")`

output `-2*x*arctan(x)*e^x + e^x*log(x^2 + 1)`

3.110.6 Sympy [A] (verification not implemented)

Time = 0.79 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int e^x(-2(1+x)\arctan(x) + \log(1+x^2)) dx = -2xe^x\operatorname{atan}(x) + e^x\log(x^2+1)$$

input `integrate(exp(x)*(ln(x**2+1)-2*(1+x)*atan(x)),x)`

output `-2*x*exp(x)*atan(x) + exp(x)*log(x**2 + 1)`

3.110.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int e^x(-2(1+x)\arctan(x) + \log(1+x^2)) dx = -2x\arctan(x)e^x + e^x\log(x^2+1)$$

input `integrate(exp(x)*(log(x^2+1)-2*(1+x)*arctan(x)),x, algorithm="maxima")`

output `-2*x*arctan(x)*e^x + e^x*log(x^2 + 1)`

3.110. $\int e^x(-2(1+x)\arctan(x) + \log(1+x^2)) dx$

3.110.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.42

$$\int e^x (-2(1+x) \arctan(x) + \log(1+x^2)) dx$$

$$= -\pi x e^x \operatorname{sgn}(x) + 2x \arctan\left(\frac{1}{x}\right) e^x + e^x \log(x^2 + 1)$$

input `integrate(exp(x)*(log(x^2+1)-2*(1+x)*arctan(x)),x, algorithm="giac")`output `-pi*x*e^x*sgn(x) + 2*x*arctan(1/x)*e^x + e^x*log(x^2 + 1)`**3.110.9 Mupad [B] (verification not implemented)**

Time = 21.30 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int e^x (-2(1+x) \arctan(x) + \log(1+x^2)) dx = e^x (\ln(x^2 + 1) - 2x \operatorname{atan}(x))$$

input `int(exp(x)*(log(x^2 + 1) - 2*atan(x)*(x + 1)),x)`output `exp(x)*(log(x^2 + 1) - 2*x*atan(x))`

3.111 $\int \arcsin(x)^2 dx$

3.111.1 Optimal result	641
3.111.2 Mathematica [A] (verified)	641
3.111.3 Rubi [A] (verified)	642
3.111.4 Maple [A] (verified)	643
3.111.5 Fricas [A] (verification not implemented)	643
3.111.6 Sympy [A] (verification not implemented)	643
3.111.7 Maxima [A] (verification not implemented)	644
3.111.8 Giac [A] (verification not implemented)	644
3.111.9 Mupad [B] (verification not implemented)	644

3.111.1 Optimal result

Integrand size = 4, antiderivative size = 25

$$\int \arcsin(x)^2 dx = -2x + 2\sqrt{1-x^2} \arcsin(x) + x \arcsin(x)^2$$

output `-2*x+2*(-x^2+1)^(1/2)*arcsin(x)+x*arcsin(x)^2`

3.111.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \arcsin(x)^2 dx = -2x + 2\sqrt{1-x^2} \arcsin(x) + x \arcsin(x)^2$$

input `Integrate[ArcSin[x]^2,x]`

output `-2*x + 2*Sqrt[1 - x^2]*ArcSin[x] + x*ArcSin[x]^2`

3.111.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5130, 5182, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \arcsin(x)^2 dx \\
 & \quad \downarrow \text{5130} \\
 & x \arcsin(x)^2 - 2 \int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx \\
 & \quad \downarrow \text{5182} \\
 & x \arcsin(x)^2 - 2 \left(\int 1 dx - \sqrt{1-x^2} \arcsin(x) \right) \\
 & \quad \downarrow \text{24} \\
 & x \arcsin(x)^2 - 2 \left(x - \sqrt{1-x^2} \arcsin(x) \right)
 \end{aligned}$$

input `Int[ArcSin[x]^2,x]`

output `x*ArcSin[x]^2 - 2*(x - Sqrt[1 - x^2]*ArcSin[x])`

3.111.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 5130 `Int[((a_.) + ArcSin[(c_.)*(x_.)]*(b_.))^n, x_Symbol] :> Simp[x*(a + b*ArcSin[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcSin[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

```
rule 5182 Int[((a_.) + ArcSin[(c_.)*(x_.)]*(b_.))^(n_.)*(x_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(d + e*x^2)^(p + 1)*((a + b*ArcSin[c*x])^n/(2*e*(p + 1))), x] + Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/(1 - c^2*x^2)^p] Int[(1 - c^2*x^2)^(p + 1/2)*(a + b*ArcSin[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]
```

3.111.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

method	result	size
default	$-2x + 2\sqrt{-x^2 + 1} \arcsin(x) + x \arcsin(x)^2$	24

input `int(arcsin(x)^2,x,method=_RETURNVERBOSE)`

output $-2*x+2*(-x^2+1)^(1/2)*arcsin(x)+x*arcsin(x)^2$

3.111.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \arcsin(x)^2 dx = x \arcsin(x)^2 + 2\sqrt{-x^2 + 1} \arcsin(x) - 2x$$

input `integrate(arcsin(x)^2,x, algorithm="fricas")`

output $x*arcsin(x)^2 + 2*sqrt(-x^2 + 1)*arcsin(x) - 2*x$

3.111.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \arcsin(x)^2 dx = x \arcsin^2(x) - 2x + 2\sqrt{1 - x^2} \arcsin(x)$$

input `integrate(asin(x)**2,x)`

output `x*asin(x)**2 - 2*x + 2*sqrt(1 - x**2)*asin(x)`

3.111.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \arcsin(x)^2 dx = x \arcsin(x)^2 + 2\sqrt{-x^2 + 1} \arcsin(x) - 2x$$

input `integrate(arcsin(x)^2,x, algorithm="maxima")`

output `x*arcsin(x)^2 + 2*sqrt(-x^2 + 1)*arcsin(x) - 2*x`

3.111.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \arcsin(x)^2 dx = x \arcsin(x)^2 + 2\sqrt{-x^2 + 1} \arcsin(x) - 2x$$

input `integrate(arcsin(x)^2,x, algorithm="giac")`

output `x*arcsin(x)^2 + 2*sqrt(-x^2 + 1)*arcsin(x) - 2*x`

3.111.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \arcsin(x)^2 dx = 2 \arcsin(x) \sqrt{1 - x^2} + x (\arcsin(x)^2 - 2)$$

input `int(asin(x)^2,x)`

output `2*asin(x)*(1 - x^2)^(1/2) + x*(asin(x)^2 - 2)`

3.112 $\int \frac{\sqrt{-1+x^2}}{x} dx$

3.112.1 Optimal result	645
3.112.2 Mathematica [A] (verified)	645
3.112.3 Rubi [A] (verified)	646
3.112.4 Maple [A] (verified)	647
3.112.5 Fricas [A] (verification not implemented)	648
3.112.6 Sympy [C] (verification not implemented)	648
3.112.7 Maxima [A] (verification not implemented)	648
3.112.8 Giac [A] (verification not implemented)	649
3.112.9 Mupad [B] (verification not implemented)	649

3.112.1 Optimal result

Integrand size = 13, antiderivative size = 22

$$\int \frac{\sqrt{-1+x^2}}{x} dx = \sqrt{-1+x^2} - \arctan\left(\sqrt{-1+x^2}\right)$$

output $(x^2-1)^{(1/2)}-\arctan((x^2-1)^{(1/2)})$

3.112.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{-1+x^2}}{x} dx = \sqrt{-1+x^2} - \arctan\left(\sqrt{-1+x^2}\right)$$

input `Integrate[Sqrt[-1 + x^2]/x,x]`

output `Sqrt[-1 + x^2] - ArcTan[Sqrt[-1 + x^2]]`

3.112.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.27, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {243, 60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{x^2-1}}{x} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \int \frac{\sqrt{x^2-1}}{x^2} dx^2 \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2} \left(2\sqrt{x^2-1} - \int \frac{1}{x^2\sqrt{x^2-1}} dx^2 \right) \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{2} \left(2\sqrt{x^2-1} - 2 \int \frac{1}{x^4+1} d\sqrt{x^2-1} \right) \\
 & \quad \downarrow \text{216} \\
 & \frac{1}{2} \left(2\sqrt{x^2-1} - 2 \arctan(\sqrt{x^2-1}) \right)
 \end{aligned}$$

input `Int[Sqrt[-1 + x^2]/x,x]`

output `(2*Sqrt[-1 + x^2] - 2*ArcTan[Sqrt[-1 + x^2]])/2`

3.112.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
 rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
 , 0] || GtQ[b, 0])`
- rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[In
 t[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I
 ntegerQ[(m - 1)/2]`

3.112.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

method	result	size
default	$\sqrt{x^2 - 1} + \arctan\left(\frac{1}{\sqrt{x^2 - 1}}\right)$	17
pseudoelliptic	$\sqrt{x^2 - 1} - \arctan(\sqrt{x^2 - 1})$	19
trager	$\sqrt{x^2 - 1} + \text{RootOf}(_Z^2 + 1) \ln\left(\frac{-\text{RootOf}(_Z^2 + 1) + \sqrt{x^2 - 1}}{x}\right)$	37
meijerg	$-\frac{\sqrt{\text{signum}(x^2 - 1)} \left(-2(2 - 2\ln(2) + 2\ln(x) + i\pi)\sqrt{\pi} + 4\sqrt{\pi} - 4\sqrt{\pi}\sqrt{-x^2 + 1} + 4\sqrt{\pi} \ln\left(\frac{1}{2} + \frac{\sqrt{-x^2 + 1}}{2}\right)\right)}{4\sqrt{\pi}\sqrt{-\text{signum}(x^2 - 1)}}$	82

input `int((x^2-1)^(1/2)/x,x,method=_RETURNVERBOSE)`

output `(x^2-1)^(1/2)+arctan(1/(x^2-1)^(1/2))`

3.112.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{-1+x^2}}{x} dx = \sqrt{x^2-1} - 2 \arctan(-x + \sqrt{x^2-1})$$

input `integrate((x^2-1)^(1/2)/x,x, algorithm="fricas")`

output `sqrt(x^2 - 1) - 2*arctan(-x + sqrt(x^2 - 1))`

3.112.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.70 (sec) , antiderivative size = 68, normalized size of antiderivative = 3.09

$$\int \frac{\sqrt{-1+x^2}}{x} dx = \begin{cases} -\frac{ix}{\sqrt{-1+\frac{1}{x^2}}} - i \operatorname{acosh}\left(\frac{1}{x}\right) + \frac{i}{x\sqrt{-1+\frac{1}{x^2}}} & \text{for } \frac{1}{|x^2|} > 1 \\ \frac{x}{\sqrt{1-\frac{1}{x^2}}} + \operatorname{asin}\left(\frac{1}{x}\right) - \frac{1}{x\sqrt{1-\frac{1}{x^2}}} & \text{otherwise} \end{cases}$$

input `integrate((x**2-1)**(1/2)/x,x)`

output `Piecewise((-I*x/sqrt(-1 + x**(-2)) - I*acosh(1/x) + I/(x*sqrt(-1 + x**(-2))), 1/Abs(x**2) > 1), (x/sqrt(1 - 1/x**2) + asin(1/x) - 1/(x*sqrt(1 - 1/x**2))), True))`

3.112.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.59

$$\int \frac{\sqrt{-1+x^2}}{x} dx = \sqrt{x^2-1} + \arcsin\left(\frac{1}{|x|}\right)$$

input `integrate((x^2-1)^(1/2)/x,x, algorithm="maxima")`

output `sqrt(x^2 - 1) + arcsin(1/abs(x))`

3.112.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{\sqrt{-1+x^2}}{x} dx = \sqrt{x^2-1} - \arctan(\sqrt{x^2-1})$$

input `integrate((x^2-1)^(1/2)/x,x, algorithm="giac")`output `sqrt(x^2 - 1) - arctan(sqrt(x^2 - 1))`**3.112.9 Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.18

$$\int \frac{\sqrt{-1+x^2}}{x} dx = \sqrt{x^2-1} - \ln\left(\frac{\sqrt{x^2-1} + 1i}{x}\right) 1i$$

input `int((x^2 - 1)^(1/2)/x,x)`output `(x^2 - 1)^(1/2) - log(((x^2 - 1)^(1/2) + 1i)/x)*1i`

3.113 $\int x \sec^2(4x) dx$

3.113.1 Optimal result	650
3.113.2 Mathematica [A] (verified)	650
3.113.3 Rubi [A] (verified)	651
3.113.4 Maple [A] (verified)	652
3.113.5 Fricas [A] (verification not implemented)	653
3.113.6 Sympy [F]	653
3.113.7 Maxima [B] (verification not implemented)	653
3.113.8 Giac [B] (verification not implemented)	654
3.113.9 Mupad [B] (verification not implemented)	654

3.113.1 Optimal result

Integrand size = 8, antiderivative size = 19

$$\int x \sec^2(4x) dx = \frac{1}{16} \log(\cos(4x)) + \frac{1}{4} x \tan(4x)$$

output `1/16*ln(cos(4*x))+1/4*x*tan(4*x)`

3.113.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int x \sec^2(4x) dx = \frac{1}{16} \log(\cos(4x)) + \frac{1}{4} x \tan(4x)$$

input `Integrate[x*Sec[4*x]^2,x]`

output `Log[Cos[4*x]]/16 + (x*Tan[4*x])/4`

3.113.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3042, 4672, 25, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \sec^2(4x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x \csc\left(4x + \frac{\pi}{2}\right)^2 dx \\
 & \quad \downarrow \text{4672} \\
 & \frac{1}{4} \int -\tan(4x) dx + \frac{1}{4} x \tan(4x) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{4} x \tan(4x) - \frac{1}{4} \int \tan(4x) dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4} x \tan(4x) - \frac{1}{4} \int \tan(4x) dx \\
 & \quad \downarrow \text{3956} \\
 & \frac{1}{4} x \tan(4x) + \frac{1}{16} \log(\cos(4x))
 \end{aligned}$$

input `Int[x*Sec[4*x]^2,x]`

output `Log[Cos[4*x]]/16 + (x*Tan[4*x])/4`

3.113.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]

rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

rule 3956 Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]

rule 4672 Int[csc[(e_.) + (f_.)*(x_.)]^2*((c_.) + (d_.)*(x_.))^(m_.), x_Symbol] := Simp
[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)
*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

3.113.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$\frac{\ln(\cos(4x))}{16} + \frac{x \tan(4x)}{4}$	16
default	$\frac{\ln(\cos(4x))}{16} + \frac{x \tan(4x)}{4}$	16
risch	$-\frac{ix}{2} + \frac{ix}{2e^{8ix}+2} + \frac{\ln(e^{8ix}+1)}{16}$	29
norman	$-\frac{x \tan(2x)}{2(\tan(2x)^2-1)} + \frac{\ln(\tan(2x)-1)}{16} + \frac{\ln(\tan(2x)+1)}{16} - \frac{\ln(1+\tan(2x)^2)}{16}$	48
parallelrisch	$\frac{\ln(\tan(2x)-1) \cos(4x) + \ln(\tan(2x)+1) \cos(4x) - \ln(\sec(2x)^2) \cos(4x) + 4x \sin(4x)}{16 \cos(4x)}$	54

```
input int(x*sec(4*x)^2,x,method=_RETURNVERBOSE)
```

```
output 1/16*ln(cos(4*x))+1/4*x*tan(4*x)
```

3.113.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.47

$$\int x \sec^2(4x) dx = \frac{\cos(4x) \log(-\cos(4x)) + 4x \sin(4x)}{16 \cos(4x)}$$

input `integrate(x*sec(4*x)^2,x, algorithm="fricas")`

output `1/16*(cos(4*x)*log(-cos(4*x)) + 4*x*sin(4*x))/cos(4*x)`

3.113.6 Sympy [F]

$$\int x \sec^2(4x) dx = \int x \sec^2(4x) dx$$

input `integrate(x*sec(4*x)**2,x)`

output `Integral(x*sec(4*x)**2, x)`

3.113.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 74 vs. 2(15) = 30.

Time = 0.28 (sec) , antiderivative size = 74, normalized size of antiderivative = 3.89

$$\int x \sec^2(4x) dx = \frac{(\cos(8x)^2 + \sin(8x)^2 + 2 \cos(8x) + 1) \log(\cos(8x)^2 + \sin(8x)^2 + 2 \cos(8x) + 1) + 16x \sin(8x)}{32 (\cos(8x)^2 + \sin(8x)^2 + 2 \cos(8x) + 1)}$$

input `integrate(x*sec(4*x)^2,x, algorithm="maxima")`

output `1/32*((cos(8*x)^2 + sin(8*x)^2 + 2*cos(8*x) + 1)*log(cos(8*x)^2 + sin(8*x)^2 + 2*cos(8*x) + 1) + 16*x*sin(8*x))/(cos(8*x)^2 + sin(8*x)^2 + 2*cos(8*x) + 1)`

3.113.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 103 vs. $2(15) = 30$.

Time = 0.28 (sec) , antiderivative size = 103, normalized size of antiderivative = 5.42

$$\int x \sec^2(4x) dx$$

$$= \frac{\log\left(\frac{4(\tan(2x)^4 - 2\tan(2x)^2 + 1)}{\tan(2x)^4 + 2\tan(2x)^2 + 1}\right) \tan(2x)^2 - 16x \tan(2x) - \log\left(\frac{4(\tan(2x)^4 - 2\tan(2x)^2 + 1)}{\tan(2x)^4 + 2\tan(2x)^2 + 1}\right)}{32(\tan(2x)^2 - 1)}$$

input `integrate(x*sec(4*x)^2,x, algorithm="giac")`

output `1/32*(log(4*(tan(2*x)^4 - 2*tan(2*x)^2 + 1)/(tan(2*x)^4 + 2*tan(2*x)^2 + 1))*tan(2*x)^2 - 16*x*tan(2*x) - log(4*(tan(2*x)^4 - 2*tan(2*x)^2 + 1)/(tan(2*x)^4 + 2*tan(2*x)^2 + 1)))/(tan(2*x)^2 - 1)`

3.113.9 Mupad [B] (verification not implemented)

Time = 20.91 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int x \sec^2(4x) dx = \frac{\ln(\cos(4x))}{16} + \frac{x \tan(4x)}{4}$$

input `int(x/cos(4*x)^2,x)`

output `log(cos(4*x))/16 + (x*tan(4*x))/4`

3.114 $\int \frac{2}{6-11x+6x^2-x^3} dx$

3.114.1 Optimal result	655
3.114.2 Mathematica [A] (verified)	655
3.114.3 Rubi [A] (verified)	656
3.114.4 Maple [A] (verified)	657
3.114.5 Fricas [A] (verification not implemented)	657
3.114.6 Sympy [A] (verification not implemented)	657
3.114.7 Maxima [A] (verification not implemented)	658
3.114.8 Giac [A] (verification not implemented)	658
3.114.9 Mupad [B] (verification not implemented)	658

3.114.1 Optimal result

Integrand size = 19, antiderivative size = 25

$$\int \frac{2}{6-11x+6x^2-x^3} dx = -\log(1-x) + 2\log(2-x) - \log(3-x)$$

output `-ln(1-x)+2*ln(2-x)-ln(-x+3)`

3.114.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{2}{6-11x+6x^2-x^3} dx = -2\left(-\log(2-x) + \frac{1}{2}\log(3-4x+x^2)\right)$$

input `Integrate[2/(6 - 11*x + 6*x^2 - x^3),x]`

output `-2*(-Log[2 - x] + Log[3 - 4*x + x^2]/2)`

3.114.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.16, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {27, 2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{2}{-x^3 + 6x^2 - 11x + 6} dx \\ & \quad \downarrow \text{27} \\ & 2 \int \frac{1}{-x^3 + 6x^2 - 11x + 6} dx \\ & \quad \downarrow \text{2462} \\ & 2 \int \left(\frac{1}{x-2} - \frac{1}{2(x-1)} - \frac{1}{2(x-3)} \right) dx \\ & \quad \downarrow \text{2009} \\ & 2 \left(-\frac{1}{2} \log(1-x) + \log(2-x) - \frac{1}{2} \log(3-x) \right) \end{aligned}$$

input `Int[2/(6 - 11*x + 6*x^2 - x^3),x]`

output `2*(-1/2*Log[1 - x] + Log[2 - x] - Log[3 - x]/2)`

3.114.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2462 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegrand[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

3.114.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

method	result	size
risch	$2 \ln(-2 + x) - \ln(x^2 - 4x + 3)$	19
default	$-\ln(-1 + x) + 2 \ln(-2 + x) - \ln(-3 + x)$	20
norman	$-\ln(-1 + x) + 2 \ln(-2 + x) - \ln(-3 + x)$	20
parallelsch	$-\ln(-1 + x) + 2 \ln(-2 + x) - \ln(-3 + x)$	20

input `int(2/(-x^3+6*x^2-11*x+6),x,method=_RETURNVERBOSE)`output `2*ln(-2+x)-ln(x^2-4*x+3)`**3.114.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.72

$$\int \frac{2}{6 - 11x + 6x^2 - x^3} dx = -\log(x^2 - 4x + 3) + 2 \log(x - 2)$$

input `integrate(2/(-x^3+6*x^2-11*x+6),x, algorithm="fracas")`output `-log(x^2 - 4*x + 3) + 2*log(x - 2)`**3.114.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.60

$$\int \frac{2}{6 - 11x + 6x^2 - x^3} dx = 2 \log(x - 2) - \log(x^2 - 4x + 3)$$

input `integrate(2/(-x**3+6*x**2-11*x+6),x)`output `2*log(x - 2) - log(x**2 - 4*x + 3)`

3.114.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{2}{6 - 11x + 6x^2 - x^3} dx = -\log(x - 1) + 2 \log(x - 2) - \log(x - 3)$$

input `integrate(2/(-x^3+6*x^2-11*x+6),x, algorithm="maxima")`output `-log(x - 1) + 2*log(x - 2) - log(x - 3)`**3.114.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{2}{6 - 11x + 6x^2 - x^3} dx = -\log(|x - 1|) + 2 \log(|x - 2|) - \log(|x - 3|)$$

input `integrate(2/(-x^3+6*x^2-11*x+6),x, algorithm="giac")`output `-log(abs(x - 1)) + 2*log(abs(x - 2)) - log(abs(x - 3))`**3.114.9 Mupad [B] (verification not implemented)**

Time = 19.62 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.72

$$\int \frac{2}{6 - 11x + 6x^2 - x^3} dx = 2 \ln(x - 2) - \ln(x^2 - 4x + 3)$$

input `int(-2/(11*x - 6*x^2 + x^3 - 6),x)`output `2*log(x - 2) - log(x^2 - 4*x + 3)`

3.115 $\int \frac{1}{1-\log(1-x)} dx$

3.115.1 Optimal result	659
3.115.2 Mathematica [A] (verified)	659
3.115.3 Rubi [A] (warning: unable to verify)	660
3.115.4 Maple [A] (verified)	661
3.115.5 Fricas [A] (verification not implemented)	661
3.115.6 Sympy [F]	662
3.115.7 Maxima [A] (verification not implemented)	662
3.115.8 Giac [A] (verification not implemented)	662
3.115.9 Mupad [F(-1)]	663

3.115.1 Optimal result

Integrand size = 12, antiderivative size = 11

$$\int \frac{1}{1-\log(1-x)} dx = e \operatorname{ExpIntegralEi}(-1 + \log(1-x))$$

output `exp(1)*Ei(-1+ln(1-x))`

3.115.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{1}{1-\log(1-x)} dx = e \operatorname{ExpIntegralEi}(-1 + \log(1-x))$$

input `Integrate[(1 - Log[1 - x])^(-1), x]`

output `E*ExpIntegralEi[-1 + Log[1 - x]]`

3.115.3 Rubi [A] (warning: unable to verify)

Time = 0.20 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.55, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2836, 2736, 2609}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{1 - \log(1 - x)} dx \\ & \quad \downarrow \text{2836} \\ & - \int \frac{1}{1 - \log(1 - x)} d(1 - x) \\ & \quad \downarrow \text{2736} \\ & - \int \frac{1 - x}{x} d \log(1 - x) \\ & \quad \downarrow \text{2609} \\ & e \operatorname{ExpIntegralEi}(-x) \end{aligned}$$

input `Int[(1 - Log[1 - x])^(-1),x]`

output `E*ExpIntegralEi[-x]`

3.115.3.1 Defintions of rubi rules used

rule 2609 `Int[(F_)^((g_.)*((e_.) + (f_.)*(x_)))/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[(F^(g*(e - c*(f/d)))/d)*ExpIntegralEi[f*g*(c + d*x)*(Log[F]/d)], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`

rule 2736 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_), x_Symbol] := Simp[1/(n*c^(1/n)) Subst[Int[E^(x/n)*(a + b*x)^p, x], x, Log[c*x^n]], x] /; FreeQ[{a, b, c, p}, x] && IntegerQ[1/n]`

```
rule 2836 Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]*(b_.))^(p_.), x_Symbol] :
> Simp[1/e Subst[Int[(a + b*Log[c*x^n])^p, x], x, d + e*x], x] /; FreeQ[{
a, b, c, d, e, n, p}, x]
```

3.115.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.55

method	result	size
derivativedivides	$-e \operatorname{Ei}_1(1 - \ln(1 - x))$	17
default	$-e \operatorname{Ei}_1(1 - \ln(1 - x))$	17
risch	$-e \operatorname{Ei}_1(1 - \ln(1 - x))$	17

```
input int(1/(1-ln(1-x)),x,method=_RETURNVERBOSE)
```

```
output -exp(1)*Ei(1,1-ln(1-x))
```

3.115.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 - \log(1 - x)} dx = e \log_integral(-(x - 1)e^{(-1)})$$

```
input integrate(1/(1-log(1-x)),x, algorithm="fricas")
```

```
output e*log_integral(-(x - 1)*e^(-1))
```

3.115.6 Sympy [F]

$$\int \frac{1}{1 - \log(1 - x)} dx = - \int \frac{1}{\log(1 - x) - 1} dx$$

input `integrate(1/(1-ln(1-x)),x)`

output `-Integral(1/(log(1 - x) - 1), x)`

3.115.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.45

$$\int \frac{1}{1 - \log(1 - x)} dx = -eE_1(-\log(-x + 1) + 1)$$

input `integrate(1/(1-log(1-x)),x, algorithm="maxima")`

output `-e*exp_integral_e(1, -log(-x + 1) + 1)`

3.115.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

$$\int \frac{1}{1 - \log(1 - x)} dx = \text{Ei}(\log(-x + 1) - 1) e$$

input `integrate(1/(1-log(1-x)),x, algorithm="giac")`

output `Ei(log(-x + 1) - 1)*e`

3.115.9 Mupad [F(-1)]

Timed out.

$$\int \frac{1}{1 - \log(1 - x)} dx = - \int \frac{1}{\ln(1 - x) - 1} dx$$

input `int(-1/(log(1 - x) - 1),x)`output `-int(1/(log(1 - x) - 1), x)`

3.116 $\int \sqrt{x + \sqrt{1 + x^2}} dx$

3.116.1 Optimal result	664
3.116.2 Mathematica [A] (verified)	664
3.116.3 Rubi [A] (verified)	665
3.116.4 Maple [C] (verified)	666
3.116.5 Fricas [A] (verification not implemented)	666
3.116.6 Sympy [A] (verification not implemented)	667
3.116.7 Maxima [F]	667
3.116.8 Giac [F]	667
3.116.9 Mupad [F(-1)]	668

3.116.1 Optimal result

Integrand size = 15, antiderivative size = 37

$$\int \sqrt{x + \sqrt{1 + x^2}} dx = -\frac{1}{\sqrt{x + \sqrt{1 + x^2}}} + \frac{1}{3} \left(x + \sqrt{1 + x^2}\right)^{3/2}$$

output `-1/(x+(x^2+1)^(1/2))^(1/2)+1/3*(x+(x^2+1)^(1/2))^(3/2)`

3.116.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

$$\int \sqrt{x + \sqrt{1 + x^2}} dx = -\frac{1}{\sqrt{x + \sqrt{1 + x^2}}} + \frac{1}{3} \left(x + \sqrt{1 + x^2}\right)^{3/2}$$

input `Integrate[Sqrt[x + Sqrt[1 + x^2]],x]`

output `-(1/Sqrt[x + Sqrt[1 + x^2]]) + (x + Sqrt[1 + x^2])^(3/2)/3`

3.116.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.11, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2542, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\sqrt{x^2+1}+x} dx$$

$$\downarrow 2542$$

$$\frac{1}{2} \int \frac{(x + \sqrt{x^2+1})^2 + 1}{(x + \sqrt{x^2+1})^{3/2}} d(x + \sqrt{x^2+1})$$

$$\downarrow 244$$

$$\frac{1}{2} \int \left(\sqrt{x + \sqrt{x^2+1}} + \frac{1}{(x + \sqrt{x^2+1})^{3/2}} \right) d(x + \sqrt{x^2+1})$$

$$\downarrow 2009$$

$$\frac{1}{2} \left(\frac{2}{3} (\sqrt{x^2+1} + x)^{3/2} - \frac{2}{\sqrt{\sqrt{x^2+1} + x}} \right)$$

input `Int[Sqrt[x + Sqrt[1 + x^2]],x]`

output `(-2/Sqrt[x + Sqrt[1 + x^2]] + (2*(x + Sqrt[1 + x^2])^(3/2))/3)/2`

3.116.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 2542 Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(
n_))^(p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f
^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; Fr
eeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]
```

3.116.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 3 vs. order 2.

Time = 0.03 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.54

method	result	size
meijerg	$\frac{16\sqrt{\pi}\sqrt{2}x^{\frac{3}{2}}\left(-\frac{1}{x^2}+1\right)\cosh\left(\frac{\operatorname{arcsinh}\left(\frac{1}{x}\right)}{2}\right)}{3} + \frac{16\sqrt{\pi}\sqrt{2}\sqrt{x}\sqrt{\frac{1}{x^2}+1}\sinh\left(\frac{\operatorname{arcsinh}\left(\frac{1}{x}\right)}{2}\right)}{8\sqrt{\pi}}$	57

```
input int((x+(x^2+1)^(1/2))^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/8/Pi^(1/2)*(16/3*Pi^(1/2)*2^(1/2)*x^(3/2)*(-1/x^2+1)*cosh(1/2*arcsinh(1/
x))+16/3*Pi^(1/2)*2^(1/2)*x^(1/2)*(1/x^2+1)^(1/2)*sinh(1/2*arcsinh(1/x)))
```

3.116.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.70

$$\int \sqrt{x + \sqrt{1 + x^2}} dx = \frac{2}{3} \left(2x - \sqrt{x^2 + 1} \right) \sqrt{x + \sqrt{x^2 + 1}}$$

```
input integrate((x+(x^2+1)^(1/2))^(1/2),x, algorithm="fracas")
```

```
output 2/3*(2*x - sqrt(x^2 + 1))*sqrt(x + sqrt(x^2 + 1))
```

3.116.6 Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.14

$$\int \sqrt{x + \sqrt{1 + x^2}} dx = \frac{4x\sqrt{x + \sqrt{x^2 + 1}}}{3} - \frac{2\sqrt{x + \sqrt{x^2 + 1}}\sqrt{x^2 + 1}}{3}$$

input `integrate((x+(x**2+1)**(1/2))**(1/2),x)`output `4*x*sqrt(x + sqrt(x**2 + 1))/3 - 2*sqrt(x + sqrt(x**2 + 1))*sqrt(x**2 + 1)/3`**3.116.7 Maxima [F]**

$$\int \sqrt{x + \sqrt{1 + x^2}} dx = \int \sqrt{x + \sqrt{x^2 + 1}} dx$$

input `integrate((x+(x^2+1)^(1/2))^(1/2),x, algorithm="maxima")`output `integrate(sqrt(x + sqrt(x^2 + 1)), x)`**3.116.8 Giac [F]**

$$\int \sqrt{x + \sqrt{1 + x^2}} dx = \int \sqrt{x + \sqrt{x^2 + 1}} dx$$

input `integrate((x+(x^2+1)^(1/2))^(1/2),x, algorithm="giac")`output `integrate(sqrt(x + sqrt(x^2 + 1)), x)`

3.116.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{x + \sqrt{1 + x^2}} dx = \int \sqrt{x + \sqrt{x^2 + 1}} dx$$

input `int((x + (x^2 + 1)^(1/2))^(1/2), x)`output `int((x + (x^2 + 1)^(1/2))^(1/2), x)`

3.117 $\int \frac{1}{2+\cos(x)} dx$

3.117.1 Optimal result	669
3.117.2 Mathematica [A] (verified)	669
3.117.3 Rubi [A] (verified)	670
3.117.4 Maple [A] (verified)	671
3.117.5 Fricas [A] (verification not implemented)	671
3.117.6 Sympy [A] (verification not implemented)	671
3.117.7 Maxima [A] (verification not implemented)	672
3.117.8 Giac [B] (verification not implemented)	672
3.117.9 Mupad [B] (verification not implemented)	672

3.117.1 Optimal result

Integrand size = 6, antiderivative size = 20

$$\int \frac{1}{2 + \cos(x)} dx = \frac{2 \arctan\left(\frac{\tan\left(\frac{x}{2}\right)}{\sqrt{3}}\right)}{\sqrt{3}}$$

output `2/3*arctan(1/3*tan(1/2*x)*3^(1/2))*3^(1/2)`

3.117.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{2 + \cos(x)} dx = \frac{2 \arctan\left(\frac{\tan\left(\frac{x}{2}\right)}{\sqrt{3}}\right)}{\sqrt{3}}$$

input `Integrate[(2 + Cos[x])^(-1),x]`

output `(2*ArcTan[Tan[x/2]/Sqrt[3]])/Sqrt[3]`

3.117.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.50, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\cos(x) + 2} dx$$

↓ 3042

$$\int \frac{1}{\sin\left(x + \frac{\pi}{2}\right) + 2} dx$$

↓ 3136

$$\frac{x}{\sqrt{3}} - \frac{2 \arctan\left(\frac{\sin(x)}{\cos(x) + \sqrt{3} + 2}\right)}{\sqrt{3}}$$

input `Int[(2 + Cos[x])^(-1),x]`

output `x/Sqrt[3] - (2*ArcTan[Sin[x]/(2 + Sqrt[3] + Cos[x]))/Sqrt[3]`

3.117.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

3.117.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

method	result	size
default	$\frac{2 \arctan\left(\frac{\tan\left(\frac{x}{2}\right)\sqrt{3}}{3}\right)\sqrt{3}}{3}$	16
risch	$\frac{i\sqrt{3} \ln(e^{ix}+2+\sqrt{3})}{3} - \frac{i\sqrt{3} \ln(e^{ix}+2-\sqrt{3})}{3}$	38

input `int(1/(2+cos(x)),x,method=_RETURNVERBOSE)`output `2/3*arctan(1/3*tan(1/2*x)*3^(1/2))*3^(1/2)`**3.117.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.15

$$\int \frac{1}{2 + \cos(x)} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{2\sqrt{3} \cos(x) + \sqrt{3}}{3 \sin(x)}\right)$$

input `integrate(1/(2+cos(x)),x, algorithm="fricas")`output `-1/3*sqrt(3)*arctan(1/3*(2*sqrt(3)*cos(x) + sqrt(3))/sin(x))`**3.117.6 Sympy [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.60

$$\int \frac{1}{2 + \cos(x)} dx = \frac{2\sqrt{3}\left(\operatorname{atan}\left(\frac{\sqrt{3}\tan\left(\frac{x}{2}\right)}{3}\right) + \pi\left\lfloor\frac{\frac{x}{2}-\frac{\pi}{2}}{\pi}\right\rfloor\right)}{3}$$

input `integrate(1/(2+cos(x)),x)`output `2*sqrt(3)*(atan(sqrt(3)*tan(x/2)/3) + pi*floor((x/2 - pi/2)/pi))/3`

3.117.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{1}{2 + \cos(x)} dx = \frac{2}{3} \sqrt{3} \arctan \left(\frac{\sqrt{3} \sin(x)}{3(\cos(x) + 1)} \right)$$

input `integrate(1/(2+cos(x)),x, algorithm="maxima")`

output `2/3*sqrt(3)*arctan(1/3*sqrt(3)*sin(x)/(cos(x) + 1))`

3.117.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 40 vs. 2(15) = 30.

Time = 0.27 (sec) , antiderivative size = 40, normalized size of antiderivative = 2.00

$$\int \frac{1}{2 + \cos(x)} dx = \frac{1}{3} \sqrt{3} \left(x + 2 \arctan \left(-\frac{\sqrt{3} \sin(x) - \sin(x)}{\sqrt{3} \cos(x) + \sqrt{3} - \cos(x) + 1} \right) \right)$$

input `integrate(1/(2+cos(x)),x, algorithm="giac")`

output `1/3*sqrt(3)*(x + 2*arctan(-(sqrt(3)*sin(x) - sin(x))/(sqrt(3)*cos(x) + sqrt(3) - cos(x) + 1)))`

3.117.9 Mupad [B] (verification not implemented)

Time = 21.01 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.60

$$\int \frac{1}{2 + \cos(x)} dx = \frac{2\sqrt{3} \left(\frac{x}{2} - \operatorname{atan}(\tan(\frac{x}{2})) \right)}{3} + \frac{2\sqrt{3} \operatorname{atan}\left(\frac{\sqrt{3} \tan(\frac{x}{2})}{3}\right)}{3}$$

input `int(1/(cos(x) + 2),x)`

output `(2*3^(1/2)*(x/2 - atan(tan(x/2)))/3 + (2*3^(1/2)*atan((3^(1/2)*tan(x/2))/3))/3`

3.118 $\int (\cos^4(x) - \sin^4(x)) dx$

3.118.1 Optimal result	673
3.118.2 Mathematica [A] (verified)	673
3.118.3 Rubi [B] (verified)	674
3.118.4 Maple [A] (verified)	674
3.118.5 Fricas [A] (verification not implemented)	675
3.118.6 Sympy [B] (verification not implemented)	675
3.118.7 Maxima [A] (verification not implemented)	675
3.118.8 Giac [A] (verification not implemented)	676
3.118.9 Mupad [B] (verification not implemented)	676

3.118.1 Optimal result

Integrand size = 11, antiderivative size = 5

$$\int (\cos^4(x) - \sin^4(x)) dx = \cos(x) \sin(x)$$

output `cos(x)*sin(x)`

3.118.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.60

$$\int (\cos^4(x) - \sin^4(x)) dx = \frac{1}{2} \sin(2x)$$

input `Integrate[Cos[x]^4 - Sin[x]^4,x]`

output `Sin[2*x]/2`

3.118.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 29 vs. $2(5) = 10$.

Time = 0.17 (sec) , antiderivative size = 29, normalized size of antiderivative = 5.80, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\cos^4(x) - \sin^4(x)) dx$$

↓ 2009

$$\frac{1}{4} \sin(x) \cos^3(x) + \frac{1}{4} \sin^3(x) \cos(x) + \frac{3}{4} \sin(x) \cos(x)$$

input `Int[Cos[x]^4 - Sin[x]^4,x]`

output `(3*Cos[x]*Sin[x])/4 + (Cos[x]^3*Sin[x])/4 + (Cos[x]*Sin[x]^3)/4`

3.118.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.118.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.40

method	result	size
risch	$\frac{\sin(2x)}{2}$	7
parallelrisch	$\frac{\sin(2x)}{2}$	7
default	$\frac{(\cos(x)^3 + \frac{3\cos(x)}{2}) \sin(x)}{4} + \frac{(\sin(x)^3 + \frac{3\sin(x)}{2}) \cos(x)}{4}$	28
parts	$\frac{(\cos(x)^3 + \frac{3\cos(x)}{2}) \sin(x)}{4} + \frac{(\sin(x)^3 + \frac{3\sin(x)}{2}) \cos(x)}{4}$	28
norman	$\frac{2 \tan(\frac{x}{2})^3 - 2 \tan(\frac{x}{2})^5 - 2 \tan(\frac{x}{2})^7 + 2 \tan(\frac{x}{2})}{(1 + \tan(\frac{x}{2})^2)^4}$	43

input `int(cos(x)^4-sin(x)^4,x,method=_RETURNVERBOSE)`

output `1/2*sin(2*x)`

3.118.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int (\cos^4(x) - \sin^4(x)) dx = \cos(x) \sin(x)$$

input `integrate(cos(x)^4-sin(x)^4,x, algorithm="fricas")`

output `cos(x)*sin(x)`

3.118.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(5) = 10$.

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 5.80

$$\int (\cos^4(x) - \sin^4(x)) dx = \frac{\sin^3(x) \cos(x)}{4} + \frac{\sin(x) \cos^3(x)}{4} + \frac{3 \sin(x) \cos(x)}{4}$$

input `integrate(cos(x)**4-sin(x)**4,x)`

output `sin(x)**3*cos(x)/4 + sin(x)*cos(x)**3/4 + 3*sin(x)*cos(x)/4`

3.118.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

$$\int (\cos^4(x) - \sin^4(x)) dx = \frac{1}{2} \sin(2x)$$

input `integrate(cos(x)^4-sin(x)^4,x, algorithm="maxima")`

output `1/2*sin(2*x)`

3.118.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

$$\int (\cos^4(x) - \sin^4(x)) dx = \frac{1}{2} \sin(2x)$$

input `integrate(cos(x)^4-sin(x)^4,x, algorithm="giac")`

output `1/2*sin(2*x)`

3.118.9 Mupad [B] (verification not implemented)

Time = 20.29 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

$$\int (\cos^4(x) - \sin^4(x)) dx = \frac{\sin(2x)}{2}$$

input `int(cos(x)^4 - sin(x)^4,x)`

output `sin(2*x)/2`

3.119 $\int \frac{x}{\sqrt{2+4x}} dx$

3.119.1 Optimal result	677
3.119.2 Mathematica [A] (verified)	677
3.119.3 Rubi [B] (verified)	678
3.119.4 Maple [A] (verified)	679
3.119.5 Fricas [A] (verification not implemented)	679
3.119.6 Sympy [B] (verification not implemented)	680
3.119.7 Maxima [C] (verification not implemented)	680
3.119.8 Giac [C] (verification not implemented)	680
3.119.9 Mupad [B] (verification not implemented)	681

3.119.1 Optimal result

Integrand size = 11, antiderivative size = 16

$$\int \frac{x}{\sqrt{2+4x}} dx = \frac{1}{6}(-1+x)\sqrt{2+4x}$$

output `1/6*(-1+x)*(4*x+2)^(1/2)`

3.119.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{x}{\sqrt{2+4x}} dx = \frac{1}{3}(-1+x)\sqrt{\frac{1}{2}+x}$$

input `Integrate[x/Sqrt[2 + 4*x],x]`

output `((-1 + x)*Sqrt[1/2 + x])/3`

3.119.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 37 vs. $2(16) = 32$.

Time = 0.15 (sec) , antiderivative size = 37, normalized size of antiderivative = 2.31, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt{4x+2}} dx$$

↓ 53

$$\int \left(\frac{1}{4}\sqrt{4x+2} - \frac{1}{2\sqrt{4x+2}} \right) dx$$

↓ 2009

$$\frac{(2x+1)^{3/2}}{6\sqrt{2}} - \frac{\sqrt{2x+1}}{2\sqrt{2}}$$

input `Int[x/Sqrt[2 + 4*x],x]`

output `-1/2*Sqrt[1 + 2*x]/Sqrt[2] + (1 + 2*x)^(3/2)/(6*Sqrt[2])`

3.119.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.119.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

method	result	size
pseudoelliptic	$\frac{(-1+x)\sqrt{4x+2}}{6}$	13
trager	$\frac{(\frac{x}{3}-\frac{1}{3})\sqrt{4x+2}}{2}$	15
gosper	$\frac{(1+2x)(-1+x)}{3\sqrt{4x+2}}$	18
risch	$\frac{(1+2x)(-1+x)}{3\sqrt{4x+2}}$	18
derivativdivides	$\frac{(4x+2)^{\frac{3}{2}}}{24} - \frac{\sqrt{4x+2}}{4}$	20
default	$\frac{(4x+2)^{\frac{3}{2}}}{24} - \frac{\sqrt{4x+2}}{4}$	20
meijerg	$\frac{\sqrt{2} \left(\frac{4\sqrt{\pi}}{3} - \frac{\sqrt{\pi}(-8x+8)\sqrt{1+2x}}{6} \right)}{8\sqrt{\pi}}$	32

input `int(x/(4*x+2)^(1/2),x,method=_RETURNVERBOSE)`output `1/6*(-1+x)*(4*x+2)^(1/2)`**3.119.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{x}{\sqrt{2+4x}} dx = \frac{1}{6} \sqrt{4x+2}(x-1)$$

input `integrate(x/(4*x+2)^(1/2),x, algorithm="fracas")`output `1/6*sqrt(4*x + 2)*(x - 1)`

3.119.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. $2(14) = 28$.

Time = 0.53 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.94

$$\int \frac{x}{\sqrt{2+4x}} dx = \frac{\sqrt{2x}\sqrt{2x+1}}{6} - \frac{\sqrt{2}\sqrt{2x+1}}{6}$$

input `integrate(x/(4*x+2)**(1/2),x)`

output `sqrt(2)*x*sqrt(2*x + 1)/6 - sqrt(2)*sqrt(2*x + 1)/6`

3.119.7 Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 3 vs. order 2.

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.19

$$\int \frac{x}{\sqrt{2+4x}} dx = \frac{1}{24} (4x+2)^{\frac{3}{2}} - \frac{1}{4} \sqrt{4x+2}$$

input `integrate(x/(4*x+2)^(1/2),x, algorithm="maxima")`

output `1/24*(4*x + 2)^(3/2) - 1/4*sqrt(4*x + 2)`

3.119.8 Giac [C] (verification not implemented)

Result contains higher order function than in optimal. Order 3 vs. order 2.

Time = 0.26 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int \frac{x}{\sqrt{2+4x}} dx = \frac{1}{12} \sqrt{2} \left((2x+1)^{\frac{3}{2}} - 3\sqrt{2x+1} \right)$$

input `integrate(x/(4*x+2)^(1/2),x, algorithm="giac")`

output `1/12*sqrt(2)*((2*x + 1)^(3/2) - 3*sqrt(2*x + 1))`

3.119.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{x}{\sqrt{2+4x}} dx = \frac{\sqrt{4x+2}(4x-4)}{24}$$

input `int(x/(4*x + 2)^(1/2),x)`

output `((4*x + 2)^(1/2)*(4*x - 4))/24`

3.120 $\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx$

3.120.1 Optimal result	682
3.120.2 Mathematica [A] (verified)	682
3.120.3 Rubi [A] (verified)	683
3.120.4 Maple [A] (verified)	684
3.120.5 Fricas [A] (verification not implemented)	684
3.120.6 Sympy [A] (verification not implemented)	685
3.120.7 Maxima [A] (verification not implemented)	685
3.120.8 Giac [A] (verification not implemented)	685
3.120.9 Mupad [B] (verification not implemented)	686

3.120.1 Optimal result

Integrand size = 12, antiderivative size = 8

$$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = 2 \sin(\sqrt{x})$$

output `2*sin(x^(1/2))`

3.120.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = 2 \sin(\sqrt{x})$$

input `Integrate[Cos[Sqrt[x]]/Sqrt[x],x]`

output `2*Sin[Sqrt[x]]`

3.120.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3861, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx \\ & \quad \downarrow \text{3861} \\ & 2 \int \cos(\sqrt{x}) d\sqrt{x} \\ & \quad \downarrow \text{3042} \\ & 2 \int \sin\left(\sqrt{x} + \frac{\pi}{2}\right) d\sqrt{x} \\ & \quad \downarrow \text{3117} \\ & 2 \sin(\sqrt{x}) \end{aligned}$$

input `Int[Cos[Sqrt[x]]/Sqrt[x],x]`

output `2*Sin[Sqrt[x]]`

3.120.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

```
rule 3861 Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol
] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^
p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[
(m + 1)/n] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[
(m + 1)/n], 0]))
```

3.120.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$2 \sin(\sqrt{x})$	7
default	$2 \sin(\sqrt{x})$	7
meijerg	$2 \sin(\sqrt{x})$	7

```
input int(cos(x^(1/2))/x^(1/2),x,method=_RETURNVERBOSE)
```

```
output 2*sin(x^(1/2))
```

3.120.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = 2 \sin(\sqrt{x})$$

```
input integrate(cos(x^(1/2))/x^(1/2),x, algorithm="fricas")
```

```
output 2*sin(sqrt(x))
```

3.120.6 Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = 2 \sin(\sqrt{x})$$

input `integrate(cos(x**(1/2))/x**(1/2),x)`output `2*sin(sqrt(x))`**3.120.7 Maxima [A] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = 2 \sin(\sqrt{x})$$

input `integrate(cos(x^(1/2))/x^(1/2),x, algorithm="maxima")`output `2*sin(sqrt(x))`**3.120.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = 2 \sin(\sqrt{x})$$

input `integrate(cos(x^(1/2))/x^(1/2),x, algorithm="giac")`output `2*sin(sqrt(x))`

3.120.9 Mupad [B] (verification not implemented)

Time = 20.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = 2 \sin(\sqrt{x})$$

input `int(cos(x^(1/2))/x^(1/2),x)`

output `2*sin(x^(1/2))`

3.121 $\int \sec(x) dx$

3.121.1 Optimal result	687
3.121.2 Mathematica [A] (verified)	687
3.121.3 Rubi [A] (verified)	688
3.121.4 Maple [A] (verified)	689
3.121.5 Fricas [B] (verification not implemented)	689
3.121.6 Sympy [B] (verification not implemented)	689
3.121.7 Maxima [A] (verification not implemented)	690
3.121.8 Giac [B] (verification not implemented)	690
3.121.9 Mupad [B] (verification not implemented)	690

3.121.1 Optimal result

Integrand size = 2, antiderivative size = 3

$$\int \sec(x) dx = \operatorname{arctanh}(\sin(x))$$

output `arctanh(sin(x))`

3.121.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \sec(x) dx = \operatorname{arctanh}(\sin(x))$$

input `Integrate[Sec[x],x]`

output `ArcTanh[Sin[x]]`

3.121.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sec(x) dx \\ \downarrow 3042 \\ \int \csc\left(x + \frac{\pi}{2}\right) dx \\ \downarrow 4257 \\ \operatorname{arctanh}(\sin(x)) \end{array}$$

input `Int[Sec[x], x]`

output `ArcTanh[Sin[x]]`

3.121.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.121.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 2.33

method	result	size
lookup	$\ln(\sec(x) + \tan(x))$	7
default	$\ln(\sec(x) + \tan(x))$	7
norman	$-\ln(\tan(\frac{x}{2}) - 1) + \ln(1 + \tan(\frac{x}{2}))$	18
parallelrisc	$-\ln(\tan(\frac{x}{2}) - 1) + \ln(1 + \tan(\frac{x}{2}))$	18
risc	$\ln(i + e^{ix}) - \ln(e^{ix} - i)$	22

input `int(sec(x),x,method=_RETURNVERBOSE)`

output `ln(sec(x)+tan(x))`

3.121.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(3) = 6$.

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 5.67

$$\int \sec(x) dx = \frac{1}{2} \log(\sin(x) + 1) - \frac{1}{2} \log(-\sin(x) + 1)$$

input `integrate(sec(x),x, algorithm="fricas")`

output `1/2*log(sin(x) + 1) - 1/2*log(-sin(x) + 1)`

3.121.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(3) = 6$.

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 5.00

$$\int \sec(x) dx = -\frac{\log(\sin(x) - 1)}{2} + \frac{\log(\sin(x) + 1)}{2}$$

input `integrate(sec(x),x)`

output `-log(sin(x) - 1)/2 + log(sin(x) + 1)/2`

3.121.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 6, normalized size of antiderivative = 2.00

$$\int \sec(x) dx = \log(\sec(x) + \tan(x))$$

input `integrate(sec(x),x, algorithm="maxima")`

output `log(sec(x) + tan(x))`

3.121.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. 2(3) = 6.

Time = 0.26 (sec) , antiderivative size = 25, normalized size of antiderivative = 8.33

$$\int \sec(x) dx = \frac{1}{4} \log\left(\left|\frac{1}{\sin(x)} + \sin(x) + 2\right|\right) - \frac{1}{4} \log\left(\left|\frac{1}{\sin(x)} + \sin(x) - 2\right|\right)$$

input `integrate(sec(x),x, algorithm="giac")`

output `1/4*log(abs(1/sin(x) + sin(x) + 2)) - 1/4*log(abs(1/sin(x) + sin(x) - 2))`

3.121.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 3.67

$$\int \sec(x) dx = \ln\left(\frac{1}{\cos(x)}\right) + \ln(\sin(x) + 1)$$

input `int(1/cos(x),x)`

output `log(1/cos(x)) + log(sin(x) + 1)`

3.122 $\int e^{\sin(x)} \cos(x) dx$

3.122.1 Optimal result	691
3.122.2 Mathematica [A] (verified)	691
3.122.3 Rubi [A] (verified)	692
3.122.4 Maple [A] (verified)	693
3.122.5 Fricas [A] (verification not implemented)	693
3.122.6 Sympy [F]	693
3.122.7 Maxima [A] (verification not implemented)	694
3.122.8 Giac [A] (verification not implemented)	694
3.122.9 Mupad [B] (verification not implemented)	694

3.122.1 Optimal result

Integrand size = 7, antiderivative size = 4

$$\int e^{\sin(x)} \cos(x) dx = e^{\sin(x)}$$

output `exp(sin(x))`

3.122.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int e^{\sin(x)} \cos(x) dx = e^{\sin(x)}$$

input `Integrate[E^Sin[x]*Cos[x],x]`

output `E^Sin[x]`

3.122.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {4834, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{\sin(x)} \cos(x) dx$$

$$\downarrow 4834$$

$$\int e^{\sin(x)} d \sin(x)$$

$$\downarrow 2624$$

$$e^{\sin(x)}$$

input `Int [E^Sin[x]*Cos[x],x]`

output `E^Sin[x]`

3.122.3.1 Defintions of rubi rules used

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;`
`FreeQ[{F, n}, x] && LinearQ[v, x]`

rule 4834 `Int[(u_)*(F_)[(c_)*((a_) + (b_)*(x_))], x_Symbol] :> With[{d = FreeFacto`
`rs[Sin[c*(a + b*x)], x]}, Simp[d/(b*c) Subst[Int[SubstFor[1, Sin[c*(a + b`
`*x)]/d, u, x], x], x, Sin[c*(a + b*x)]/d, x] /;` `FunctionOfQ[Sin[c*(a + b*x`
`)]/d, u, x, True]] /;` `FreeQ[{a, b, c}, x] && (EqQ[F, Cos] || EqQ[F, cos])`

3.122.4 Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

method	result	size
derivativedivides	$e^{\sin(x)}$	4
default	$e^{\sin(x)}$	4
risch	$e^{\sin(x)}$	4

input `int(exp(sin(x))/tan(x)/csc(x),x,method=_RETURNVERBOSE)`

output `exp(sin(x))`

3.122.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int e^{\sin(x)} \cos(x) dx = e^{\sin(x)}$$

input `integrate(exp(sin(x))/tan(x)/csc(x),x, algorithm="fricas")`

output `e^sin(x)`

3.122.6 Sympy [F]

$$\int e^{\sin(x)} \cos(x) dx = \int \frac{e^{\sin(x)}}{\tan(x) \csc(x)} dx$$

input `integrate(exp(sin(x))/tan(x)/csc(x),x)`

output `Integral(exp(sin(x))/(tan(x)*csc(x)), x)`

3.122.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int e^{\sin(x)} \cos(x) dx = e^{\sin(x)}$$

input `integrate(exp(sin(x))/tan(x)/csc(x),x, algorithm="maxima")`output `e^sin(x)`**3.122.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int e^{\sin(x)} \cos(x) dx = e^{\sin(x)}$$

input `integrate(exp(sin(x))/tan(x)/csc(x),x, algorithm="giac")`output `e^sin(x)`**3.122.9 Mupad [B] (verification not implemented)**

Time = 17.48 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int e^{\sin(x)} \cos(x) dx = e^{\sin(x)}$$

input `int((exp(sin(x))*sin(x))/tan(x),x)`output `exp(sin(x))`

3.123 $\int x \log^2(x) dx$

3.123.1 Optimal result	695
3.123.2 Mathematica [A] (verified)	695
3.123.3 Rubi [A] (verified)	696
3.123.4 Maple [A] (verified)	697
3.123.5 Fricas [A] (verification not implemented)	697
3.123.6 Sympy [A] (verification not implemented)	697
3.123.7 Maxima [A] (verification not implemented)	698
3.123.8 Giac [A] (verification not implemented)	698
3.123.9 Mupad [B] (verification not implemented)	698

3.123.1 Optimal result

Integrand size = 6, antiderivative size = 28

$$\int x \log^2(x) dx = \frac{x^2}{4} - \frac{1}{2}x^2 \log(x) + \frac{1}{2}x^2 \log^2(x)$$

output `1/4*x^2-1/2*x^2*ln(x)+1/2*x^2*ln(x)^2`

3.123.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int x \log^2(x) dx = \frac{x^2}{4} - \frac{1}{2}x^2 \log(x) + \frac{1}{2}x^2 \log^2(x)$$

input `Integrate[x*Log[x]^2,x]`

output `x^2/4 - (x^2*Log[x])/2 + (x^2*Log[x]^2)/2`

3.123.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2742, 2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \log^2(x) dx$$

$$\downarrow \text{2742}$$

$$\frac{1}{2}x^2 \log^2(x) - \int x \log(x) dx$$

$$\downarrow \text{2741}$$

$$\frac{x^2}{4} + \frac{1}{2}x^2 \log^2(x) - \frac{1}{2}x^2 \log(x)$$

input `Int[x*Log[x]^2,x]`

output `x^2/4 - (x^2*Log[x])/2 + (x^2*Log[x]^2)/2`

3.123.3.1 Defintions of rubi rules used

rule 2741 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_.), x_Symbol] :=
Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(
m + 1)/(d*(m + 1)^2)), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]`

rule 2742 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((d_.)*(x_)^(m_.), x_Symbol
1] := Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])^p/(d*(m + 1))), x] - Simp[b*n*
(p/(m + 1)) Int[(d*x)^m*(a + b*Log[c*x^n])^(p - 1), x], x] /; FreeQ[{a, b
, c, d, m, n}, x] && NeQ[m, -1] && GtQ[p, 0]`

3.123.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{x^2}{4} - \frac{x^2 \ln(x)}{2} + \frac{x^2 \ln(x)^2}{2}$	23
norman	$\frac{x^2}{4} - \frac{x^2 \ln(x)}{2} + \frac{x^2 \ln(x)^2}{2}$	23
risch	$\frac{x^2}{4} - \frac{x^2 \ln(x)}{2} + \frac{x^2 \ln(x)^2}{2}$	23
parallelrisch	$\frac{x^2}{4} - \frac{x^2 \ln(x)}{2} + \frac{x^2 \ln(x)^2}{2}$	23
parts	$\frac{x^2}{4} - \frac{x^2 \ln(x)}{2} + \frac{x^2 \ln(x)^2}{2}$	23

input `int(x*ln(x)^2,x,method=_RETURNVERBOSE)`output `1/4*x^2-1/2*x^2*ln(x)+1/2*x^2*ln(x)^2`**3.123.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int x \log^2(x) dx = \frac{1}{2} x^2 \log(x)^2 - \frac{1}{2} x^2 \log(x) + \frac{1}{4} x^2$$

input `integrate(x*log(x)^2,x, algorithm="fricas")`output `1/2*x^2*log(x)^2 - 1/2*x^2*log(x) + 1/4*x^2`**3.123.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int x \log^2(x) dx = \frac{x^2 \log(x)^2}{2} - \frac{x^2 \log(x)}{2} + \frac{x^2}{4}$$

input `integrate(x*ln(x)**2,x)`output `x**2*log(x)**2/2 - x**2*log(x)/2 + x**2/4`

3.123.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.61

$$\int x \log^2(x) dx = \frac{1}{4} (2 \log(x)^2 - 2 \log(x) + 1) x^2$$

input `integrate(x*log(x)^2,x, algorithm="maxima")`output `1/4*(2*log(x)^2 - 2*log(x) + 1)*x^2`**3.123.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int x \log^2(x) dx = \frac{1}{2} x^2 \log(x)^2 - \frac{1}{2} x^2 \log(x) + \frac{1}{4} x^2$$

input `integrate(x*log(x)^2,x, algorithm="giac")`output `1/2*x^2*log(x)^2 - 1/2*x^2*log(x) + 1/4*x^2`**3.123.9 Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.61

$$\int x \log^2(x) dx = \frac{x^2 (2 \ln(x)^2 - 2 \ln(x) + 1)}{4}$$

input `int(x*log(x)^2,x)`output `(x^2*(2*log(x)^2 - 2*log(x) + 1))/4`

3.124 $\int \frac{1}{5+4\sqrt{x}+x} dx$

3.124.1 Optimal result	699
3.124.2 Mathematica [A] (verified)	699
3.124.3 Rubi [A] (verified)	700
3.124.4 Maple [A] (verified)	701
3.124.5 Fricas [A] (verification not implemented)	702
3.124.6 Sympy [A] (verification not implemented)	702
3.124.7 Maxima [A] (verification not implemented)	702
3.124.8 Giac [A] (verification not implemented)	703
3.124.9 Mupad [B] (verification not implemented)	703

3.124.1 Optimal result

Integrand size = 12, antiderivative size = 22

$$\int \frac{1}{5 + 4\sqrt{x} + x} dx = -4 \arctan(2 + \sqrt{x}) + \log(5 + 4\sqrt{x} + x)$$

output `-4*arctan(2+x^(1/2))+ln(5+4*x^(1/2)+x)`

3.124.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{5 + 4\sqrt{x} + x} dx = -4 \arctan(2 + \sqrt{x}) + \log(5 + 4\sqrt{x} + x)$$

input `Integrate[(5 + 4*Sqrt[x] + x)^(-1), x]`

output `-4*ArcTan[2 + Sqrt[x]] + Log[5 + 4*Sqrt[x] + x]`

3.124.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.55, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {1680, 1142, 27, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x + 4\sqrt{x} + 5} dx \\
 & \quad \downarrow \text{1680} \\
 & 2 \int \frac{\sqrt{x}}{x + 4\sqrt{x} + 5} d\sqrt{x} \\
 & \quad \downarrow \text{1142} \\
 & 2 \left(\frac{1}{2} \int \frac{2(\sqrt{x} + 2)}{x + 4\sqrt{x} + 5} d\sqrt{x} - 2 \int \frac{1}{x + 4\sqrt{x} + 5} d\sqrt{x} \right) \\
 & \quad \downarrow \text{27} \\
 & 2 \left(\int \frac{\sqrt{x} + 2}{x + 4\sqrt{x} + 5} d\sqrt{x} - 2 \int \frac{1}{x + 4\sqrt{x} + 5} d\sqrt{x} \right) \\
 & \quad \downarrow \text{1083} \\
 & 2 \left(4 \int \frac{1}{-x - 4} d(2\sqrt{x} + 4) + \int \frac{\sqrt{x} + 2}{x + 4\sqrt{x} + 5} d\sqrt{x} \right) \\
 & \quad \downarrow \text{217} \\
 & 2 \left(\int \frac{\sqrt{x} + 2}{x + 4\sqrt{x} + 5} d\sqrt{x} - 2 \arctan \left(\frac{1}{2} (2\sqrt{x} + 4) \right) \right) \\
 & \quad \downarrow \text{1103} \\
 & 2 \left(\frac{1}{2} \log(x + 4\sqrt{x} + 5) - 2 \arctan \left(\frac{1}{2} (2\sqrt{x} + 4) \right) \right)
 \end{aligned}$$

input `Int[(5 + 4*Sqrt[x] + x)^(-1),x]`

output `2*(-2*ArcTan[(4 + 2*Sqrt[x])/2] + Log[5 + 4*Sqrt[x] + x]/2)`

3.124.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

- rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

- rule 1142 `Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

- rule 1680 `Int[((a_) + (c_.)*(x_)^(n2_.) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[n]}, Simp[k Subst[Int[x^(k - 1)*(a + b*x^(k*n) + c*x^(2*k*n))^p, x], x, x^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && EqQ[n2, 2*n] && FractionQ[n]`

3.124.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.86

method	result
derivativedivides	$-4 \arctan(2 + \sqrt{x}) + \ln(5 + 4\sqrt{x} + x)$
default	$2 \arctan\left(\frac{x}{4} - \frac{3}{4}\right) + \frac{\ln(x^2 - 6x + 25)}{2} - \frac{\ln(5 + x - 4\sqrt{x})}{2} - 2 \arctan(-2 + \sqrt{x}) + \frac{\ln(5 + 4\sqrt{x} + x)}{2} - 2$
trager	$\text{RootOf}(_Z^2 - 2_Z + 5) \ln(5 + 4\sqrt{x} + x) - \ln\left(79 \text{RootOf}(_Z^2 - 2_Z + 5)^2 x - 9\right)$

3.124. $\int \frac{1}{5+4\sqrt{x}+x} dx$

input `int(1/(5+4*x^(1/2)+x),x,method=_RETURNVERBOSE)`

output `-4*arctan(2+x^(1/2))+ln(5+4*x^(1/2)+x)`

3.124.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{1}{5 + 4\sqrt{x} + x} dx = -4 \arctan(\sqrt{x} + 2) + \log(x + 4\sqrt{x} + 5)$$

input `integrate(1/(5+4*x^(1/2)+x),x, algorithm="fricas")`

output `-4*arctan(sqrt(x) + 2) + log(x + 4*sqrt(x) + 5)`

3.124.6 Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{5 + 4\sqrt{x} + x} dx = \log(4\sqrt{x} + x + 5) - 4 \operatorname{atan}(\sqrt{x} + 2)$$

input `integrate(1/(5+4*x**(1/2)+x),x)`

output `log(4*sqrt(x) + x + 5) - 4*atan(sqrt(x) + 2)`

3.124.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{1}{5 + 4\sqrt{x} + x} dx = -4 \arctan(\sqrt{x} + 2) + \log(x + 4\sqrt{x} + 5)$$

input `integrate(1/(5+4*x^(1/2)+x),x, algorithm="maxima")`

output `-4*arctan(sqrt(x) + 2) + log(x + 4*sqrt(x) + 5)`

3.124.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{1}{5 + 4\sqrt{x} + x} dx = -4 \arctan(\sqrt{x} + 2) + \log(x + 4\sqrt{x} + 5)$$

input `integrate(1/(5+4*x^(1/2)+x),x, algorithm="giac")`output `-4*arctan(sqrt(x) + 2) + log(x + 4*sqrt(x) + 5)`**3.124.9 Mupad [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{1}{5 + 4\sqrt{x} + x} dx = \ln(x + 4\sqrt{x} + 5) - 4 \operatorname{atan}(\sqrt{x} + 2)$$

input `int(1/(x + 4*x^(1/2) + 5),x)`output `log(x + 4*x^(1/2) + 5) - 4*atan(x^(1/2) + 2)`

3.125 $\int 2015^x dx$

3.125.1 Optimal result	704
3.125.2 Mathematica [A] (verified)	704
3.125.3 Rubi [A] (verified)	705
3.125.4 Maple [A] (verified)	705
3.125.5 Fricas [A] (verification not implemented)	706
3.125.6 Sympy [A] (verification not implemented)	706
3.125.7 Maxima [A] (verification not implemented)	706
3.125.8 Giac [A] (verification not implemented)	707
3.125.9 Mupad [B] (verification not implemented)	707

3.125.1 Optimal result

Integrand size = 3, antiderivative size = 8

$$\int 2015^x dx = \frac{2015^x}{\log(2015)}$$

output `2015^x/ln(2015)`

3.125.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int 2015^x dx = \frac{2015^x}{\log(2015)}$$

input `Integrate[2015^x,x]`

output `2015^x/Log[2015]`

3.125.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int 2015^x dx$$

$$\downarrow 2624$$

$$\frac{2015^x}{\log(2015)}$$

input `Int[2015^x,x]`

output `2015^x/Log[2015]`

3.125.3.1 Defintions of rubi rules used

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;`
`FreeQ[{F, n}, x] && LinearQ[v, x]`

3.125.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
gospers	$\frac{2015^x}{\ln(2015)}$	9
derivativedivides	$\frac{2015^x}{\ln(2015)}$	9
default	$\frac{2015^x}{\ln(2015)}$	9
parallelrisch	$\frac{2015^x}{\ln(2015)}$	9
norman	$\frac{e^{x \ln(2015)}}{\ln(2015)}$	11
meijerg	$-\frac{1-e^{x \ln(2015)}}{\ln(2015)}$	16
risch	$\frac{31^x 13^x 5^x}{\ln(5)+\ln(13)+\ln(31)}$	20

input `int(2015^x,x,method=_RETURNVERBOSE)`

output `2015^x/ln(2015)`

3.125.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int 2015^x dx = \frac{2015^x}{\log(2015)}$$

input `integrate(2015^x,x, algorithm="fricas")`

output `2015^x/log(2015)`

3.125.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int 2015^x dx = \frac{2015^x}{\log(2015)}$$

input `integrate(2015**x,x)`

output `2015**x/log(2015)`

3.125.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int 2015^x dx = \frac{2015^x}{\log(2015)}$$

input `integrate(2015^x,x, algorithm="maxima")`

output `2015^x/log(2015)`

3.125.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int 2015^x dx = \frac{2015^x}{\log(2015)}$$

input `integrate(2015^x,x, algorithm="giac")`

output `2015^x/log(2015)`

3.125.9 Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int 2015^x dx = \frac{2015^x}{\ln(2015)}$$

input `int(2015^x,x)`

output `2015^x/log(2015)`

3.126 $\int \frac{x}{(-3+x)(5+x)^2} dx$

3.126.1 Optimal result	708
3.126.2 Mathematica [A] (verified)	708
3.126.3 Rubi [A] (verified)	709
3.126.4 Maple [A] (verified)	710
3.126.5 Fricas [A] (verification not implemented)	710
3.126.6 Sympy [A] (verification not implemented)	710
3.126.7 Maxima [A] (verification not implemented)	711
3.126.8 Giac [A] (verification not implemented)	711
3.126.9 Mupad [B] (verification not implemented)	711

3.126.1 Optimal result

Integrand size = 12, antiderivative size = 22

$$\int \frac{x}{(-3+x)(5+x)^2} dx = -\frac{5}{8(5+x)} - \frac{3}{32} \operatorname{arctanh}\left(\frac{1+x}{4}\right)$$

output `-5/(8*x+40)-3/32*arctanh(1/4+1/4*x)`

3.126.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.27

$$\int \frac{x}{(-3+x)(5+x)^2} dx = -\frac{5}{8(5+x)} + \frac{3}{64} \log(3-x) - \frac{3}{64} \log(5+x)$$

input `Integrate[x/((-3 + x)*(5 + x)^2),x]`

output `-5/(8*(5 + x)) + (3*Log[3 - x])/64 - (3*Log[5 + x])/64`

3.126.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.27, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(x-3)(x+5)^2} dx$$

↓ 86

$$\int \left(-\frac{3}{64(x+5)} + \frac{5}{8(x+5)^2} + \frac{3}{64(x-3)} \right) dx$$

↓ 2009

$$-\frac{5}{8(x+5)} + \frac{3}{64} \log(3-x) - \frac{3}{64} \log(x+5)$$

input `Int[x/((-3 + x)*(5 + x)^2),x]`

output `-5/(8*(5 + x)) + (3*Log[3 - x])/64 - (3*Log[5 + x])/64`

3.126.3.1 Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.126.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.95

method	result	size
default	$-\frac{5}{8(5+x)} - \frac{3 \ln(5+x)}{64} + \frac{3 \ln(-3+x)}{64}$	21
norman	$-\frac{5}{8(5+x)} - \frac{3 \ln(5+x)}{64} + \frac{3 \ln(-3+x)}{64}$	21
risch	$-\frac{5}{8(5+x)} - \frac{3 \ln(5+x)}{64} + \frac{3 \ln(-3+x)}{64}$	21
parallelrisc	$-\frac{3 \ln(5+x)x - 3 \ln(-3+x)x + 40 + 15 \ln(5+x) - 15 \ln(-3+x)}{64(5+x)}$	36

input `int(x/(-3+x)/(5+x)^2,x,method=_RETURNVERBOSE)`output `-5/8/(5+x)-3/64*ln(5+x)+3/64*ln(-3+x)`**3.126.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.23

$$\int \frac{x}{(-3+x)(5+x)^2} dx = -\frac{3(x+5)\log(x+5) - 3(x+5)\log(x-3) + 40}{64(x+5)}$$

input `integrate(x/(-3+x)/(5+x)^2,x, algorithm="fricas")`output `-1/64*(3*(x + 5)*log(x + 5) - 3*(x + 5)*log(x - 3) + 40)/(x + 5)`**3.126.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{x}{(-3+x)(5+x)^2} dx = \frac{3 \log(x-3)}{64} - \frac{3 \log(x+5)}{64} - \frac{5}{8x+40}$$

input `integrate(x/(-3+x)/(5+x)**2,x)`output `3*log(x - 3)/64 - 3*log(x + 5)/64 - 5/(8*x + 40)`

3.126.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{x}{(-3+x)(5+x)^2} dx = -\frac{5}{8(x+5)} - \frac{3}{64} \log(x+5) + \frac{3}{64} \log(x-3)$$

input `integrate(x/(-3+x)/(5+x)^2,x, algorithm="maxima")`output `-5/8/(x + 5) - 3/64*log(x + 5) + 3/64*log(x - 3)`**3.126.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.95

$$\int \frac{x}{(-3+x)(5+x)^2} dx = -\frac{5}{8(x+5)} + \frac{3}{64} \log\left(\left|-\frac{8}{x+5} + 1\right|\right)$$

input `integrate(x/(-3+x)/(5+x)^2,x, algorithm="giac")`output `-5/8/(x + 5) + 3/64*log(abs(-8/(x + 5) + 1))`**3.126.9 Mupad [B] (verification not implemented)**

Time = 16.61 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{x}{(-3+x)(5+x)^2} dx = -\frac{3 \ln\left(\frac{x+5}{x-3}\right)}{64} - \frac{5}{8(x+5)}$$

input `int(x/((x - 3)*(x + 5)^2),x)`output `-(3*log((x + 5)/(x - 3)))/64 - 5/(8*(x + 5))`

3.127 $\int \frac{\log(1+\log(x))}{x} dx$

3.127.1 Optimal result	712
3.127.2 Mathematica [A] (verified)	712
3.127.3 Rubi [A] (verified)	713
3.127.4 Maple [A] (verified)	714
3.127.5 Fricas [A] (verification not implemented)	714
3.127.6 Sympy [A] (verification not implemented)	714
3.127.7 Maxima [A] (verification not implemented)	715
3.127.8 Giac [A] (verification not implemented)	715
3.127.9 Mupad [B] (verification not implemented)	715

3.127.1 Optimal result

Integrand size = 9, antiderivative size = 15

$$\int \frac{\log(1 + \log(x))}{x} dx = -\log(x) + (1 + \log(x)) \log(1 + \log(x))$$

output `-ln(x)+(ln(x)+1)*ln(ln(x)+1)`

3.127.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\log(1 + \log(x))}{x} dx = -\log(x) + (1 + \log(x)) \log(1 + \log(x))$$

input `Integrate[Log[1 + Log[x]]/x,x]`

output `-Log[x] + (1 + Log[x])*Log[1 + Log[x]]`

3.127.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3039, 2836, 2732}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\log(\log(x) + 1)}{x} dx \\ & \quad \downarrow \text{3039} \\ & \int \log(\log(x) + 1) d\log(x) \\ & \quad \downarrow \text{2836} \\ & \int \log(\log(x) + 1) d(\log(x) + 1) \\ & \quad \downarrow \text{2732} \\ & -\log(x) + (\log(x) + 1) \log(\log(x) + 1) - 1 \end{aligned}$$

input `Int[Log[1 + Log[x]]/x,x]`

output `-1 - Log[x] + (1 + Log[x])*Log[1 + Log[x]]`

3.127.3.1 Defintions of rubi rules used

rule 2732 `Int[Log[(c_.)*(x_)^(n_.)], x_Symbol] := Simp[x*Log[c*x^n], x] - Simp[n*x, x] /; FreeQ[{c, n}, x]`

rule 2836 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]*(b_.))^(p_.), x_Symbol] := Simp[1/e Subst[Int[(a + b*Log[c*x^n])^p, x], x, d + e*x], x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

rule 3039 `Int[u_, x_Symbol] := With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst[[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst] /; NonsumQ[u]`

3.127. $\int \frac{\log(1+\log(x))}{x} dx$

3.127.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

method	result	size
derivativedivides	$(\ln(x) + 1) \ln(\ln(x) + 1) - \ln(x) - 1$	17
default	$(\ln(x) + 1) \ln(\ln(x) + 1) - \ln(x) - 1$	17
norman	$-\ln(x) + \ln(\ln(x) + 1) \ln(x) + \ln(\ln(x) + 1)$	19
risch	$-\ln(x) + \ln(\ln(x) + 1) \ln(x) + \ln(\ln(x) + 1)$	19

input `int(ln(ln(x)+1)/x,x,method=_RETURNVERBOSE)`output `(ln(x)+1)*ln(ln(x)+1)-ln(x)-1`**3.127.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\log(1 + \log(x))}{x} dx = (\log(x) + 1) \log(\log(x) + 1) - \log(x)$$

input `integrate(log(log(x)+1)/x,x, algorithm="fracas")`output `(log(x) + 1)*log(log(x) + 1) - log(x)`**3.127.6 Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.27

$$\int \frac{\log(1 + \log(x))}{x} dx = \log(x) \log(\log(x) + 1) - \log(x) + \log(\log(x) + 1)$$

input `integrate(ln(ln(x)+1)/x,x)`output `log(x)*log(log(x) + 1) - log(x) + log(log(x) + 1)`

3.127.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

$$\int \frac{\log(1 + \log(x))}{x} dx = (\log(x) + 1) \log(\log(x) + 1) - \log(x) - 1$$

input `integrate(log(log(x)+1)/x,x, algorithm="maxima")`output `(log(x) + 1)*log(log(x) + 1) - log(x) - 1`**3.127.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

$$\int \frac{\log(1 + \log(x))}{x} dx = (\log(x) + 1) \log(\log(x) + 1) - \log(x) - 1$$

input `integrate(log(log(x)+1)/x,x, algorithm="giac")`output `(log(x) + 1)*log(log(x) + 1) - log(x) - 1`**3.127.9 Mupad [B] (verification not implemented)**

Time = 15.22 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.20

$$\int \frac{\log(1 + \log(x))}{x} dx = \ln(\ln(x) + 1) - \ln(x) + \ln(\ln(x) + 1) \ln(x)$$

input `int(log(log(x) + 1)/x,x)`output `log(log(x) + 1) - log(x) + log(log(x) + 1)*log(x)`

3.128 $\int \sqrt{\csc(x) - \sin(x)} dx$

3.128.1 Optimal result	716
3.128.2 Mathematica [A] (verified)	716
3.128.3 Rubi [A] (verified)	717
3.128.4 Maple [A] (verified)	718
3.128.5 Fricas [A] (verification not implemented)	719
3.128.6 Sympy [F]	719
3.128.7 Maxima [B] (verification not implemented)	719
3.128.8 Giac [F]	720
3.128.9 Mupad [B] (verification not implemented)	720

3.128.1 Optimal result

Integrand size = 11, antiderivative size = 13

$$\int \sqrt{\csc(x) - \sin(x)} dx = 2\sqrt{\cos(x) \cot(x)} \tan(x)$$

output `2*(cos(x)*cot(x))^(1/2)*tan(x)`

3.128.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \sqrt{\csc(x) - \sin(x)} dx = 2\sqrt{\cos(x) \cot(x)} \tan(x)$$

input `Integrate[Sqrt[Csc[x] - Sin[x]],x]`

output `2*Sqrt[Cos[x]*Cot[x]]*Tan[x]`

3.128.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 4897, 3042, 4900, 3042, 3069}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\csc(x) - \sin(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\csc(x) - \sin(x)} dx \\
 & \quad \downarrow \text{4897} \\
 & \int \sqrt{\cos(x) \cot(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\cos(x) \cot(x)} dx \\
 & \quad \downarrow \text{4900} \\
 & \frac{\sqrt{\cos(x) \cot(x)} \int \sqrt{\cos(x)} \sqrt{\cot(x)} dx}{\sqrt{\cos(x)} \sqrt{\cot(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\cos(x) \cot(x)} \int \sqrt{\sin(x + \frac{\pi}{2})} \sqrt{-\tan(x + \frac{\pi}{2})} dx}{\sqrt{\cos(x)} \sqrt{\cot(x)}} \\
 & \quad \downarrow \text{3069} \\
 & 2 \tan(x) \sqrt{\cos(x) \cot(x)}
 \end{aligned}$$

input `Int[Sqrt[Csc[x] - Sin[x]],x]`

output `2*Sqrt[Cos[x]*Cot[x]]*Tan[x]`

3.128.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3069 `Int[((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*(a*Sin[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*m)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 1, 0]`

rule 4897 `Int[u_, x_Symbol] := Int[TrigSimplify[u], x] /; TrigSimplifyQ[u]`

rule 4900 `Int[(u_.)*((v_)^(m_.)*(w_)^(n_.))^p_, x_Symbol] := With[{uu = ActivateTrig[u], vv = ActivateTrig[v], ww = ActivateTrig[w]}, Simp[(vv^m*ww^n)^FracPart[p]/(vv^(m*FracPart[p])*ww^(n*FracPart[p])) Int[uu*vv^(m*p)*ww^(n*p), x], x] /; FreeQ[{m, n, p}, x] && !IntegerQ[p] && (!InertTrigFreeQ[v] || !InertTrigFreeQ[w])`

3.128.4 Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
default	$2\sqrt{\cos(x)\cot(x)}\tan(x)$	12
risch	$-\frac{i\sqrt{2}\sqrt{\frac{i(e^{2ix}+1)^2e^{-ix}}{e^{2ix}-1}}(e^{2ix}-1)}{e^{2ix}+1}$	51

input `int((csc(x)-sin(x))^(1/2),x,method=_RETURNVERBOSE)`

output `2*(cos(x)*cot(x))^(1/2)*tan(x)`

3.128.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.46

$$\int \sqrt{\csc(x) - \sin(x)} dx = \frac{2 \sqrt{\frac{\cos(x)^2}{\sin(x)}} \sin(x)}{\cos(x)}$$

input `integrate((csc(x)-sin(x))^(1/2),x, algorithm="fricas")`

output `2*sqrt(cos(x)^2/sin(x))*sin(x)/cos(x)`

3.128.6 Sympy [F]

$$\int \sqrt{\csc(x) - \sin(x)} dx = \int \sqrt{-\sin(x) + \csc(x)} dx$$

input `integrate((csc(x)-sin(x))**(1/2),x)`

output `Integral(sqrt(-sin(x) + csc(x)), x)`

3.128.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 188 vs. $2(11) = 22$.

Time = 0.36 (sec) , antiderivative size = 188, normalized size of antiderivative = 14.46

$$\int \sqrt{\csc(x) - \sin(x)} dx = \frac{((\cos(\frac{3}{2}x) - \cos(\frac{1}{2}x) + \sin(\frac{3}{2}x) + \sin(\frac{1}{2}x)) \cos(\frac{1}{2} \arctan(\sin(x), \cos(x) - 1)) - (\cos(\frac{3}{2}x) - \cos(\frac{1}{2}x)))}{\cos(x)}$$

input `integrate((csc(x)-sin(x))^(1/2),x, algorithm="maxima")`

output $((\cos(3/2*x) - \cos(1/2*x) + \sin(3/2*x) + \sin(1/2*x))*\cos(1/2*\arctan2(\sin(x), \cos(x) - 1)) - (\cos(3/2*x) - \cos(1/2*x) - \sin(3/2*x) - \sin(1/2*x))*\sin(1/2*\arctan2(\sin(x), \cos(x) - 1)))*\cos(1/2*\arctan2(\sin(x), \cos(x) + 1)) - ((\cos(3/2*x) - \cos(1/2*x) - \sin(3/2*x) - \sin(1/2*x))*\cos(1/2*\arctan2(\sin(x), \cos(x) - 1)) + (\cos(3/2*x) - \cos(1/2*x) + \sin(3/2*x) + \sin(1/2*x))*\sin(1/2*\arctan2(\sin(x), \cos(x) - 1)))*\sin(1/2*\arctan2(\sin(x), \cos(x) + 1)))/((\cos(x)^2 + \sin(x)^2 + 2*\cos(x) + 1)^{(1/4)}*(\cos(x)^2 + \sin(x)^2 - 2*\cos(x) + 1)^{(1/4)})$

3.128.8 Giac [F]

$$\int \sqrt{\csc(x) - \sin(x)} dx = \int \sqrt{\csc(x) - \sin(x)} dx$$

input `integrate((csc(x)-sin(x))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(csc(x) - sin(x)), x)`

3.128.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \sqrt{\csc(x) - \sin(x)} dx = \frac{2 |\cos(x)|}{\cos(x) \sqrt{\frac{1}{\sin(x)}}}$$

input `int((1/sin(x) - sin(x))^(1/2),x)`

output `(2*abs(cos(x)))/(cos(x)*(1/sin(x))^(1/2))`

3.129 $\int \frac{1}{\sqrt{25+x^2}} dx$

3.129.1 Optimal result	721
3.129.2 Mathematica [B] (verified)	721
3.129.3 Rubi [A] (verified)	722
3.129.4 Maple [A] (verified)	722
3.129.5 Fricas [B] (verification not implemented)	723
3.129.6 Sympy [A] (verification not implemented)	723
3.129.7 Maxima [A] (verification not implemented)	723
3.129.8 Giac [B] (verification not implemented)	724
3.129.9 Mupad [B] (verification not implemented)	724

3.129.1 Optimal result

Integrand size = 9, antiderivative size = 6

$$\int \frac{1}{\sqrt{25+x^2}} dx = \operatorname{arcsinh}\left(\frac{x}{5}\right)$$

output `arcsinh(1/5*x)`

3.129.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 16 vs. $2(6) = 12$.

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 2.67

$$\int \frac{1}{\sqrt{25+x^2}} dx = -\log\left(-x + \sqrt{25+x^2}\right)$$

input `Integrate[1/Sqrt[25 + x^2],x]`

output `-Log[-x + Sqrt[25 + x^2]]`

3.129.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x^2 + 25}} dx$$

↓ 222

$$\operatorname{arcsinh}\left(\frac{x}{5}\right)$$

input `Int [1/Sqrt [25 + x^2], x]`

output `ArcSinh[x/5]`

3.129.3.1 Defintions of rubi rules used

rule 222 `Int [1/Sqrt [(a_) + (b_.)*(x_)^2], x_Symbol] := Simp [ArcSinh [Rt [b, 2]*(x/Sqrt [a])]/Rt [b, 2], x] /; FreeQ [{a, b}, x] && GtQ [a, 0] && PosQ [b]`

3.129.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

method	result	size
default	$\operatorname{arcsinh}\left(\frac{x}{5}\right)$	5
meijerg	$\operatorname{arcsinh}\left(\frac{x}{5}\right)$	5
pseudoelliptic	$\operatorname{arctanh}\left(\frac{\sqrt{x^2+25}}{x}\right)$	13
trager	$-\ln\left(x - \sqrt{x^2 + 25}\right)$	15

input `int (1/(x^2+25)^(1/2), x, method=_RETURNVERBOSE)`

output `arcsinh(1/5*x)`

3.129.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 14 vs. $2(4) = 8$.

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 2.33

$$\int \frac{1}{\sqrt{25 + x^2}} dx = -\log(-x + \sqrt{x^2 + 25})$$

input `integrate(1/(x^2+25)^(1/2),x, algorithm="fricas")`

output `-log(-x + sqrt(x^2 + 25))`

3.129.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.50

$$\int \frac{1}{\sqrt{25 + x^2}} dx = \operatorname{asinh}\left(\frac{x}{5}\right)$$

input `integrate(1/(x**2+25)**(1/2),x)`

output `asinh(x/5)`

3.129.7 Maxima [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt{25 + x^2}} dx = \operatorname{arsinh}\left(\frac{1}{5}x\right)$$

input `integrate(1/(x^2+25)^(1/2),x, algorithm="maxima")`

output `arcsinh(1/5*x)`

3.129.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. $2(4) = 8$.

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 4.17

$$\int \frac{1}{\sqrt{25+x^2}} dx = \frac{1}{2} \sqrt{x^2+25}x - \frac{25}{2} \log(-x + \sqrt{x^2+25})$$

input `integrate(1/(x^2+25)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(x^2 + 25)*x - 25/2*log(-x + sqrt(x^2 + 25))`

3.129.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt{25+x^2}} dx = \operatorname{asinh}\left(\frac{x}{5}\right)$$

input `int(1/(x^2 + 25)^(1/2),x)`

output `asinh(x/5)`

3.130 $\int \frac{-1+\log^2(x)}{x \log^2(x)} dx$

3.130.1 Optimal result	725
3.130.2 Mathematica [A] (verified)	725
3.130.3 Rubi [A] (verified)	726
3.130.4 Maple [A] (verified)	727
3.130.5 Fricas [A] (verification not implemented)	727
3.130.6 Sympy [A] (verification not implemented)	728
3.130.7 Maxima [A] (verification not implemented)	728
3.130.8 Giac [A] (verification not implemented)	728
3.130.9 Mupad [B] (verification not implemented)	729

3.130.1 Optimal result

Integrand size = 14, antiderivative size = 7

$$\int \frac{-1 + \log^2(x)}{x \log^2(x)} dx = \frac{1}{\log(x)} + \log(x)$$

output 1/ln(x)+ln(x)

3.130.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{-1 + \log^2(x)}{x \log^2(x)} dx = \frac{1}{\log(x)} + \log(x)$$

input Integrate[(-1 + Log[x]^2)/(x*Log[x]^2), x]

output Log[x]^(-1) + Log[x]

3.130.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3039, 25, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\log^2(x) - 1}{x \log^2(x)} dx \\
 & \quad \downarrow \text{3039} \\
 & \int -\frac{1 - \log^2(x)}{\log^2(x)} d\log(x) \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{1 - \log^2(x)}{\log^2(x)} d\log(x) \\
 & \quad \downarrow \text{244} \\
 & -\int \left(\frac{1}{\log^2(x)} - 1 \right) d\log(x) \\
 & \quad \downarrow \text{2009} \\
 & \log(x) + \frac{1}{\log(x)}
 \end{aligned}$$

input `Int[(-1 + Log[x]^2)/(x*Log[x]^2), x]`

output `Log[x]^(-1) + Log[x]`

3.130.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3039 `Int[u_, x_Symbol] := With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst
[[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst]] /;
NonsumQ[u]`

3.130.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

method	result	size
derivativedivides	$\frac{1}{\ln(x)} + \ln(x)$	8
default	$\frac{1}{\ln(x)} + \ln(x)$	8
risch	$\frac{1}{\ln(x)} + \ln(x)$	8
parts	$\frac{1}{\ln(x)} + \ln(x)$	8
norman	$\frac{1+\ln(x)^2}{\ln(x)}$	12
parallelrisch	$\frac{1+\ln(x)^2}{\ln(x)}$	12

input `int((ln(x)^2-1)/x/ln(x)^2,x,method=_RETURNVERBOSE)`

output `1/ln(x)+ln(x)`

3.130.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.57

$$\int \frac{-1 + \log^2(x)}{x \log^2(x)} dx = \frac{\log(x)^2 + 1}{\log(x)}$$

input `integrate((log(x)^2-1)/x/log(x)^2,x, algorithm="fracas")`

output `(log(x)^2 + 1)/log(x)`

3.130.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{-1 + \log^2(x)}{x \log^2(x)} dx = \log(x) + \frac{1}{\log(x)}$$

input `integrate((ln(x)**2-1)/x/ln(x)**2,x)`output `log(x) + 1/log(x)`**3.130.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{-1 + \log^2(x)}{x \log^2(x)} dx = \frac{1}{\log(x)} + \log(x)$$

input `integrate((log(x)^2-1)/x/log(x)^2,x, algorithm="maxima")`output `1/log(x) + log(x)`**3.130.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{-1 + \log^2(x)}{x \log^2(x)} dx = \frac{1}{\log(x)} + \log(x)$$

input `integrate((log(x)^2-1)/x/log(x)^2,x, algorithm="giac")`output `1/log(x) + log(x)`

3.130.9 Mupad [B] (verification not implemented)

Time = 17.16 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{-1 + \log^2(x)}{x \log^2(x)} dx = \ln(x) + \frac{1}{\ln(x)}$$

input `int((log(x)^2 - 1)/(x*log(x)^2),x)`

output `log(x) + 1/log(x)`

3.131 $\int e^{3x} \arctan(e^x) dx$

3.131.1 Optimal result	730
3.131.2 Mathematica [A] (verified)	730
3.131.3 Rubi [A] (verified)	731
3.131.4 Maple [C] (verified)	732
3.131.5 Fricas [A] (verification not implemented)	733
3.131.6 Sympy [A] (verification not implemented)	733
3.131.7 Maxima [A] (verification not implemented)	733
3.131.8 Giac [A] (verification not implemented)	734
3.131.9 Mupad [B] (verification not implemented)	734

3.131.1 Optimal result

Integrand size = 10, antiderivative size = 35

$$\int e^{3x} \arctan(e^x) dx = -\frac{e^{2x}}{6} + \frac{1}{3}e^{3x} \arctan(e^x) + \frac{1}{6} \log(1 + e^{2x})$$

output `-1/6*exp(2*x)+1/3*exp(3*x)*arctan(exp(x))+1/6*ln(exp(2*x)+1)`

3.131.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int e^{3x} \arctan(e^x) dx = \frac{1}{6}(-e^{2x} + 2e^{3x} \arctan(e^x) + \log(1 + e^{2x}))$$

input `Integrate[E^(3*x)*ArcTan[E^x],x]`

output `(-E^(2*x) + 2*E^(3*x)*ArcTan[E^x] + Log[1 + E^(2*x)])/6`

3.131.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5730, 27, 2678, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{3x} \arctan(e^x) dx \\
 & \quad \downarrow \text{5730} \\
 & \frac{1}{3} e^{3x} \arctan(e^x) - \int \frac{e^{4x}}{3(1+e^{2x})} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{3} e^{3x} \arctan(e^x) - \frac{1}{3} \int \frac{e^{4x}}{1+e^{2x}} dx \\
 & \quad \downarrow \text{2678} \\
 & \frac{1}{3} e^{3x} \arctan(e^x) - \frac{1}{6} \int \frac{e^{2x}}{1+e^{2x}} de^{2x} \\
 & \quad \downarrow \text{49} \\
 & \frac{1}{3} e^{3x} \arctan(e^x) - \frac{1}{6} \int \left(1 + \frac{1}{-1-e^{2x}}\right) de^{2x} \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{3} e^{3x} \arctan(e^x) + \frac{1}{6} (\log(e^{2x} + 1) - e^{2x})
 \end{aligned}$$

input `Int[E^(3*x)*ArcTan[E^x],x]`

output `(E^(3*x)*ArcTan[E^x])/3 + (-E^(2*x) + Log[1 + E^(2*x)])/6`

3.131.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2678 `Int[((a_) + (b_.)*(F_)^((e_.)*((c_.) + (d_.)*(x_))))^(p_.)*(G_)^((h_.)*((f_.) + (g_.)*(x_))), x_Symbol] := With[{m = FullSimplify[g*h*(Log[G]/(d*e*Log[F]))}], Simp[Denominator[m]*(G^(f*h - c*g*(h/d))/(d*e*Log[F])) Subst[Int[x^(Numerator[m] - 1)*(a + b*x^Denominator[m])^p, x], x, F^(e*((c + d*x)/Denominator[m]))], x] /; LeQ[m, -1] || GeQ[m, 1] /; FreeQ[{F, G, a, b, c, d, e, f, g, h, p}, x]`
- rule 5730 `Int[((a_.) + ArcTan[u_]*(b_.))*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[(a + b*ArcTan[u]) w, x] - Simp[b Int[SimplifyIntegrand[w*(D[u, x]/(1 + u^2)), x], x], x] /; InverseFunctionFreeQ[w, x] /; FreeQ[{a, b}, x] && InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_.) + (d_.)*x)^(m_.)] /; FreeQ[{c, d, m}, x] && FalseQ[FunctionOfLinear[v*(a + b*ArcTan[u]), x]]`

3.131.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.22 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.34

method	result	size
risch	$-\frac{ie^{3x} \ln(1+ie^x)}{6} + \frac{ie^{3x} \ln(1-ie^x)}{6} - \frac{e^{2x}}{6} + \frac{\ln(e^{2x}+1)}{6}$	47

input `int(exp(3*x)*arctan(exp(x)),x,method=_RETURNVERBOSE)`

output $-1/6*I*\exp(3*x)*\ln(1+I*\exp(x))+1/6*I*\exp(3*x)*\ln(1-I*\exp(x))-1/6*\exp(2*x)+1/6*\ln(\exp(2*x)+1)$

3.131.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int e^{3x} \arctan(e^x) dx = \frac{1}{3} \arctan(e^x) e^{(3x)} - \frac{1}{6} e^{(2x)} + \frac{1}{6} \log(e^{(2x)} + 1)$$

input `integrate(exp(3*x)*arctan(exp(x)),x, algorithm="fricas")`

output $1/3*\arctan(e^x)*e^{(3*x)} - 1/6*e^{(2*x)} + 1/6*\log(e^{(2*x)} + 1)$

3.131.6 Sympy [A] (verification not implemented)

Time = 3.33 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.77

$$\int e^{3x} \arctan(e^x) dx = \frac{e^{3x} \operatorname{atan}(e^x)}{3} - \frac{e^{2x}}{6} + \frac{\log(e^{2x} + 1)}{6}$$

input `integrate(exp(3*x)*atan(exp(x)),x)`

output $\exp(3*x)*\operatorname{atan}(\exp(x))/3 - \exp(2*x)/6 + \log(\exp(2*x) + 1)/6$

3.131.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int e^{3x} \arctan(e^x) dx = \frac{1}{3} \arctan(e^x) e^{(3x)} - \frac{1}{6} e^{(2x)} + \frac{1}{6} \log(e^{(2x)} + 1)$$

input `integrate(exp(3*x)*arctan(exp(x)),x, algorithm="maxima")`

output $1/3*\arctan(e^x)*e^{(3*x)} - 1/6*e^{(2*x)} + 1/6*\log(e^{(2*x)} + 1)$

3.131.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int e^{3x} \arctan(e^x) dx = \frac{1}{3} \arctan(e^x) e^{(3x)} - \frac{1}{6} e^{(2x)} + \frac{1}{6} \log(e^{(2x)} + 1)$$

input `integrate(exp(3*x)*arctan(exp(x)),x, algorithm="giac")`output `1/3*arctan(e^x)*e^(3*x) - 1/6*e^(2*x) + 1/6*log(e^(2*x) + 1)`**3.131.9 Mupad [B] (verification not implemented)**

Time = 15.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int e^{3x} \arctan(e^x) dx = \frac{\ln(e^{2x} + 1)}{6} - \frac{e^{2x}}{6} + \frac{\operatorname{atan}(e^x) e^{3x}}{3}$$

input `int(atan(exp(x))*exp(3*x),x)`output `log(exp(2*x) + 1)/6 - exp(2*x)/6 + (atan(exp(x))*exp(3*x))/3`

3.132 $\int \frac{1}{\cos^4(x)+\sin^4(x)} dx$

3.132.1 Optimal result	735
3.132.2 Mathematica [A] (verified)	735
3.132.3 Rubi [A] (verified)	736
3.132.4 Maple [C] (verified)	737
3.132.5 Fricas [A] (verification not implemented)	738
3.132.6 Sympy [F(-1)]	738
3.132.7 Maxima [A] (verification not implemented)	738
3.132.8 Giac [A] (verification not implemented)	739
3.132.9 Mupad [B] (verification not implemented)	739

3.132.1 Optimal result

Integrand size = 11, antiderivative size = 71

$$\int \frac{1}{\cos^4(x) + \sin^4(x)} dx = \sqrt{2}x + \frac{\arctan\left(\frac{1-2\cos^2(x)}{1+\sqrt{2}-2\cos(x)\sin(x)}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-2\cos^2(x)}{1+\sqrt{2}+2\cos(x)\sin(x)}\right)}{\sqrt{2}}$$

```
output x*2^(1/2)+1/2*arctan((1-2*cos(x)^2)/(1+2^(1/2)-2*cos(x)*sin(x)))*2^(1/2)-1/2*arctan((1-2*cos(x)^2)/(1+2^(1/2)+2*cos(x)*sin(x)))*2^(1/2)
```

3.132.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.45

$$\int \frac{1}{\cos^4(x) + \sin^4(x)} dx = \frac{-\arctan(1 - \sqrt{2}\tan(x)) + \arctan(1 + \sqrt{2}\tan(x))}{\sqrt{2}}$$

```
input Integrate[(Cos[x]^4 + Sin[x]^4)^(-1),x]
```

```
output (-ArcTan[1 - Sqrt[2]*Tan[x]] + ArcTan[1 + Sqrt[2]*Tan[x]])/Sqrt[2]
```


3.132.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.52, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 4889, 1476, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sin^4(x) + \cos^4(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sin(x)^4 + \cos(x)^4} dx \\
 & \quad \downarrow \text{4889} \\
 & \int \frac{\tan^2(x) + 1}{\tan^4(x) + 1} d \tan(x) \\
 & \quad \downarrow \text{1476} \\
 & \frac{1}{2} \int \frac{1}{\tan^2(x) - \sqrt{2} \tan(x) + 1} d \tan(x) + \frac{1}{2} \int \frac{1}{\tan^2(x) + \sqrt{2} \tan(x) + 1} d \tan(x) \\
 & \quad \downarrow \text{1082} \\
 & \frac{\int \frac{1}{-(1-\sqrt{2} \tan(x))^2 - 1} d(1 - \sqrt{2} \tan(x))}{\sqrt{2}} - \frac{\int \frac{1}{-(\sqrt{2} \tan(x)+1)^2 - 1} d(\sqrt{2} \tan(x) + 1)}{\sqrt{2}} \\
 & \quad \downarrow \text{217} \\
 & \frac{\arctan(\sqrt{2} \tan(x) + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2} \tan(x))}{\sqrt{2}}
 \end{aligned}$$

input `Int[(Cos[x]^4 + Sin[x]^4)^(-1), x]`

output `-(ArcTan[1 - Sqrt[2]*Tan[x]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Tan[x]]/Sqrt[2]`

3.132.3.1 Defintions of rubi rules used

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]
```

```
rule 1476 Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4889 Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, With[{d = FreeFactors[Tan[v], x]}, Simp[d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x^2), Tan[v]/d, u, x], x], x, Tan[v]/d, x]] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Tan[v], x], u, x]] /; InverseFunctionFreeQ[u, x] && !MatchQ[u, (v_.)*((c_.)*tan[w_]^(n_.)*tan[z_]^(n_.))^ (p_.) /; FreeQ[{c, p}, x] && IntegerQ[n] && LinearQ[w, x] && EqQ[z, 2*w]]
```

3.132.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.56

method	result
risch	$\frac{i\sqrt{2} \ln(e^{4ix} + 2\sqrt{2} + 3)}{4} - \frac{i\sqrt{2} \ln(e^{4ix} - 2\sqrt{2} + 3)}{4}$
default	$\frac{\sqrt{2} \left(\ln\left(\frac{\tan(x)^2 + \tan(x)\sqrt{2} + 1}{\tan(x)^2 - \tan(x)\sqrt{2} + 1}\right) + 2 \arctan(\tan(x)\sqrt{2} + 1) + 2 \arctan(\tan(x)\sqrt{2} - 1) \right)}{8} + \frac{\sqrt{2} \left(\ln\left(\frac{\tan(x)^2 - \tan(x)\sqrt{2} + 1}{\tan(x)^2 + \tan(x)\sqrt{2} + 1}\right) + 2 \arctan(\tan(x)\sqrt{2} + 1) + 2 \arctan(\tan(x)\sqrt{2} - 1) \right)}{8}$

3.132. $\int \frac{1}{\cos^4(x) + \sin^4(x)} dx$

input `int(1/(sin(x)^4+cos(x)^4),x,method=_RETURNVERBOSE)`

output `1/4*I*2^(1/2)*ln(exp(4*I*x)+2*2^(1/2)+3)-1/4*I*2^(1/2)*ln(exp(4*I*x)-2*2^(1/2)+3)`

3.132.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.66

$$\int \frac{1}{\cos^4(x) + \sin^4(x)} dx = -\frac{1}{4} \sqrt{2} \arctan \left(\frac{6 \sqrt{2} \cos(x)^4 - 6 \sqrt{2} \cos(x)^2 + \sqrt{2}}{4 (2 \cos(x)^3 - \cos(x)) \sin(x)} \right)$$

input `integrate(1/(sin(x)^4+cos(x)^4),x, algorithm="fricas")`

output `-1/4*sqrt(2)*arctan(1/4*(6*sqrt(2)*cos(x)^4 - 6*sqrt(2)*cos(x)^2 + sqrt(2))/(2*cos(x)^3 - cos(x))*sin(x))`

3.132.6 Sympy [F(-1)]

Timed out.

$$\int \frac{1}{\cos^4(x) + \sin^4(x)} dx = \text{Timed out}$$

input `integrate(1/(sin(x)**4+cos(x)**4),x)`

output `Timed out`

3.132.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.55

$$\int \frac{1}{\cos^4(x) + \sin^4(x)} dx = \frac{1}{2} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} (\sqrt{2} + 2 \tan(x)) \right) + \frac{1}{2} \sqrt{2} \arctan \left(-\frac{1}{2} \sqrt{2} (\sqrt{2} - 2 \tan(x)) \right)$$

input `integrate(1/(sin(x)^4+cos(x)^4),x, algorithm="maxima")`

output `1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*tan(x))) + 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*tan(x)))`

3.132.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.68

$$\int \frac{1}{\cos^4(x) + \sin^4(x)} dx = \frac{1}{2} \sqrt{2} \left(2x + \arctan \left(-\frac{\sqrt{2} \sin(4x) - \sin(4x)}{\sqrt{2} \cos(4x) + \sqrt{2} - \cos(4x) + 1} \right) \right)$$

input `integrate(1/(sin(x)^4+cos(x)^4),x, algorithm="giac")`

output `1/2*sqrt(2)*(2*x + arctan(-(sqrt(2)*sin(4*x) - sin(4*x))/(sqrt(2)*cos(4*x) + sqrt(2) - cos(4*x) + 1)))`

3.132.9 Mupad [B] (verification not implemented)

Time = 18.24 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.62

$$\int \frac{1}{\cos^4(x) + \sin^4(x)} dx = \sqrt{2} (x - \operatorname{atan}(\tan(x))) + \frac{\sqrt{2} \left(\operatorname{atan}\left(\frac{\sqrt{2}\tan(x)}{2}\right) + \operatorname{atan}\left(\frac{\sqrt{2}\tan(x)^3}{2} + \frac{\sqrt{2}\tan(x)}{2}\right) \right)}{2}$$

input `int(1/(cos(x)^4 + sin(x)^4),x)`

output `2^(1/2)*(x - atan(tan(x))) + (2^(1/2)*(atan((2^(1/2)*tan(x))/2) + atan((2^(1/2)*tan(x)^3)/2 + (2^(1/2)*tan(x))/2)))/2`

3.133 $\int \frac{1+e^x}{1-e^x} dx$

3.133.1 Optimal result	740
3.133.2 Mathematica [A] (verified)	740
3.133.3 Rubi [A] (verified)	741
3.133.4 Maple [A] (verified)	742
3.133.5 Fricas [A] (verification not implemented)	742
3.133.6 Sympy [A] (verification not implemented)	743
3.133.7 Maxima [A] (verification not implemented)	743
3.133.8 Giac [A] (verification not implemented)	743
3.133.9 Mupad [B] (verification not implemented)	744

3.133.1 Optimal result

Integrand size = 15, antiderivative size = 12

$$\int \frac{1+e^x}{1-e^x} dx = x - 2 \log(1 - e^x)$$

output `x-2*ln(1-exp(x))`

3.133.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

$$\int \frac{1+e^x}{1-e^x} dx = \log(e^x) - 2 \log(-1 + e^x)$$

input `Integrate[(1 + E^x)/(1 - E^x), x]`

output `Log[E^x] - 2*Log[-1 + E^x]`

3.133.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.25, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2720, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{e^x + 1}{1 - e^x} dx \\
 & \quad \downarrow \text{2720} \\
 & \int \frac{e^{-x}(e^x + 1)}{1 - e^x} de^x \\
 & \quad \downarrow \text{86} \\
 & \int \left(e^{-x} - \frac{2}{e^x - 1} \right) de^x \\
 & \quad \downarrow \text{2009} \\
 & \log(e^x) - 2 \log(1 - e^x)
 \end{aligned}$$

input `Int[(1 + E^x)/(1 - E^x),x]`

output `Log[E^x] - 2*Log[1 - E^x]`

3.133.3.1 Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.133.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

method	result	size
norman	$x - 2 \ln(e^x - 1)$	10
risch	$x - 2 \ln(e^x - 1)$	10
parallelrisch	$x - 2 \ln(e^x - 1)$	10
derivativedivides	$-2 \ln(e^x - 1) + \ln(e^x)$	12
default	$-2 \ln(e^x - 1) + \ln(e^x)$	12

```
input int((exp(x)+1)/(1-exp(x)),x,method=_RETURNVERBOSE)
```

```
output x-2*ln(exp(x)-1)
```

3.133.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{1+e^x}{1-e^x} dx = x - 2 \log(e^x - 1)$$

```
input integrate((exp(x)+1)/(1-exp(x)),x, algorithm="fricas")
```

```
output x - 2*log(e^x - 1)
```

3.133.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1 + e^x}{1 - e^x} dx = x - 2 \log(e^x - 1)$$

input `integrate((exp(x)+1)/(1-exp(x)),x)`output `x - 2*log(exp(x) - 1)`**3.133.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{1 + e^x}{1 - e^x} dx = x - 2 \log(e^x - 1)$$

input `integrate((exp(x)+1)/(1-exp(x)),x, algorithm="maxima")`output `x - 2*log(e^x - 1)`**3.133.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1 + e^x}{1 - e^x} dx = x - 2 \log(|e^x - 1|)$$

input `integrate((exp(x)+1)/(1-exp(x)),x, algorithm="giac")`output `x - 2*log(abs(e^x - 1))`

3.133.9 Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{1+e^x}{1-e^x} dx = x - 2 \ln(e^x - 1)$$

input `int(-(exp(x) + 1)/(exp(x) - 1),x)`

output `x - 2*log(exp(x) - 1)`

3.134 $\int \tan^4(x) dx$

3.134.1 Optimal result	745
3.134.2 Mathematica [A] (verified)	745
3.134.3 Rubi [A] (verified)	746
3.134.4 Maple [A] (verified)	747
3.134.5 Fricas [A] (verification not implemented)	748
3.134.6 Sympy [A] (verification not implemented)	748
3.134.7 Maxima [A] (verification not implemented)	748
3.134.8 Giac [A] (verification not implemented)	749
3.134.9 Mupad [B] (verification not implemented)	749

3.134.1 Optimal result

Integrand size = 4, antiderivative size = 14

$$\int \tan^4(x) dx = x - \tan(x) + \frac{\tan^3(x)}{3}$$

output `x-tan(x)+1/3*tan(x)^3`

3.134.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \tan^4(x) dx = \arctan(\tan(x)) - \tan(x) + \frac{\tan^3(x)}{3}$$

input `Integrate[Tan[x]^4,x]`

output `ArcTan[Tan[x]] - Tan[x] + Tan[x]^3/3`

3.134.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {3042, 3954, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \tan^4(x) dx \\
 \downarrow 3042 \\
 \int \tan(x)^4 dx \\
 \downarrow 3954 \\
 \frac{\tan^3(x)}{3} - \int \tan^2(x) dx \\
 \downarrow 3042 \\
 \frac{\tan^3(x)}{3} - \int \tan(x)^2 dx \\
 \downarrow 3954 \\
 \int 1 dx + \frac{\tan^3(x)}{3} - \tan(x) \\
 \downarrow 24 \\
 x + \frac{\tan^3(x)}{3} - \tan(x)
 \end{array}$$

input `Int [Tan [x] ^4, x]`

output `x - Tan [x] + Tan [x] ^3/3`

3.134.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

3.134.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
norman	$x - \tan(x) + \frac{\tan(x)^3}{3}$	13
parallelrisc	$x - \tan(x) + \frac{\tan(x)^3}{3}$	13
derivativedivides	$\frac{\tan(x)^3}{3} - \tan(x) + \arctan(\tan(x))$	15
default	$\frac{\tan(x)^3}{3} - \tan(x) + \arctan(\tan(x))$	15
risc	$x - \frac{4i(3e^{4ix} + 3e^{2ix} + 2)}{3(e^{2ix} + 1)^3}$	31

input `int(tan(x)^4,x,method=_RETURNVERBOSE)`

output `x-tan(x)+1/3*tan(x)^3`

3.134.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(x) dx = \frac{1}{3} \tan(x)^3 + x - \tan(x)$$

input `integrate(tan(x)^4,x, algorithm="fricas")`output `1/3*tan(x)^3 + x - tan(x)`**3.134.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.36

$$\int \tan^4(x) dx = x + \frac{\sin^3(x)}{3 \cos^3(x)} - \frac{\sin(x)}{\cos(x)}$$

input `integrate(tan(x)**4,x)`output `x + sin(x)**3/(3*cos(x)**3) - sin(x)/cos(x)`**3.134.7 Maxima [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(x) dx = \frac{1}{3} \tan(x)^3 + x - \tan(x)$$

input `integrate(tan(x)^4,x, algorithm="maxima")`output `1/3*tan(x)^3 + x - tan(x)`

3.134.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(x) dx = \frac{1}{3} \tan(x)^3 + x - \tan(x)$$

input `integrate(tan(x)^4,x, algorithm="giac")`

output `1/3*tan(x)^3 + x - tan(x)`

3.134.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \tan^4(x) dx = \frac{\tan(x)^3}{3} - \tan(x) + x$$

input `int(tan(x)^4,x)`

output `x - tan(x) + tan(x)^3/3`

3.135 $\int \sin(x) \tan^2(x) dx$

3.135.1 Optimal result	750
3.135.2 Mathematica [A] (verified)	750
3.135.3 Rubi [A] (verified)	751
3.135.4 Maple [B] (verified)	752
3.135.5 Fricas [B] (verification not implemented)	752
3.135.6 Sympy [A] (verification not implemented)	753
3.135.7 Maxima [A] (verification not implemented)	753
3.135.8 Giac [A] (verification not implemented)	753
3.135.9 Mupad [B] (verification not implemented)	754

3.135.1 Optimal result

Integrand size = 7, antiderivative size = 5

$$\int \sin(x) \tan^2(x) dx = \cos(x) + \sec(x)$$

output `cos(x)+sec(x)`

3.135.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \sin(x) \tan^2(x) dx = \cos(x) + \sec(x)$$

input `Integrate[Sin[x]*Tan[x]^2,x]`

output `Cos[x] + Sec[x]`

3.135.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 3070, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(x) \tan^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(x) \tan(x)^2 dx \\
 & \quad \downarrow \text{3070} \\
 & - \int (1 - \cos^2(x)) \sec^2(x) d \cos(x) \\
 & \quad \downarrow \text{244} \\
 & - \int (\sec^2(x) - 1) d \cos(x) \\
 & \quad \downarrow \text{2009} \\
 & \cos(x) + \sec(x)
 \end{aligned}$$

input `Int[Sin[x]*Tan[x]^2,x]`

output `Cos[x] + Sec[x]`

3.135.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`


```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3070 Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol]
:= Simp[-f^(-1) Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f
*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]
```

3.135.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(5) = 10$.

Time = 0.22 (sec) , antiderivative size = 20, normalized size of antiderivative = 4.00

method	result	size
default	$\frac{\sin(x)^4}{\cos(x)} + (2 + \sin(x)^2) \cos(x)$	20
risch	$\frac{e^{3ix} + 7 \cos(x) + 5i \sin(x)}{2e^{2ix} + 2}$	27

```
input int(sin(x)*tan(x)^2,x,method=_RETURNVERBOSE)
```

```
output sin(x)^4/cos(x)+(2+sin(x)^2)*cos(x)
```

3.135.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. $2(5) = 10$.

Time = 0.26 (sec) , antiderivative size = 11, normalized size of antiderivative = 2.20

$$\int \sin(x) \tan^2(x) dx = \frac{\cos(x)^2 + 1}{\cos(x)}$$

```
input integrate(sin(x)*tan(x)^2,x, algorithm="fricas")
```

```
output (cos(x)^2 + 1)/cos(x)
```

3.135.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.40

$$\int \sin(x) \tan^2(x) dx = \cos(x) + \frac{1}{\cos(x)}$$

input `integrate(sin(x)*tan(x)**2,x)`output `cos(x) + 1/cos(x)`**3.135.7 Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.40

$$\int \sin(x) \tan^2(x) dx = \frac{1}{\cos(x)} + \cos(x)$$

input `integrate(sin(x)*tan(x)^2,x, algorithm="maxima")`output `1/cos(x) + cos(x)`**3.135.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.40

$$\int \sin(x) \tan^2(x) dx = \frac{1}{\cos(x)} + \cos(x)$$

input `integrate(sin(x)*tan(x)^2,x, algorithm="giac")`output `1/cos(x) + cos(x)`

3.135.9 Mupad [B] (verification not implemented)

Time = 15.42 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.40

$$\int \sin(x) \tan^2(x) dx = \cos(x) + \frac{1}{\cos(x)}$$

input `int(sin(x)*tan(x)^2,x)`

output `cos(x) + 1/cos(x)`

3.136 $\int \frac{1+x}{3+2x+x^2} dx$

3.136.1 Optimal result	755
3.136.2 Mathematica [A] (verified)	755
3.136.3 Rubi [A] (verified)	756
3.136.4 Maple [A] (verified)	756
3.136.5 Fricas [A] (verification not implemented)	757
3.136.6 Sympy [A] (verification not implemented)	757
3.136.7 Maxima [A] (verification not implemented)	757
3.136.8 Giac [A] (verification not implemented)	758
3.136.9 Mupad [B] (verification not implemented)	758

3.136.1 Optimal result

Integrand size = 14, antiderivative size = 13

$$\int \frac{1+x}{3+2x+x^2} dx = \frac{1}{2} \log(3+2x+x^2)$$

output `1/2*ln(x^2+2*x+3)`

3.136.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1+x}{3+2x+x^2} dx = \frac{1}{2} \log(3+2x+x^2)$$

input `Integrate[(1 + x)/(3 + 2*x + x^2), x]`

output `Log[3 + 2*x + x^2]/2`

3.136.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$, Rules used = {1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x+1}{x^2+2x+3} dx$$

↓ 1103

$$\frac{1}{2} \log(x^2+2x+3)$$

input `Int[(1 + x)/(3 + 2*x + x^2),x]`

output `Log[3 + 2*x + x^2]/2`

3.136.3.1 Defintions of rubi rules used

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] :> Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

3.136.4 Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
default	$\frac{\ln(x^2+2x+3)}{2}$	12
norman	$\frac{\ln(x^2+2x+3)}{2}$	12
risch	$\frac{\ln(x^2+2x+3)}{2}$	12
parallelrisc	$\frac{\ln(x^2+2x+3)}{2}$	12

input `int((1+x)/(x^2+2*x+3),x,method=_RETURNVERBOSE)`

output `1/2*ln(x^2+2*x+3)`

3.136.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1+x}{3+2x+x^2} dx = \frac{1}{2} \log(x^2 + 2x + 3)$$

input `integrate((1+x)/(x^2+2*x+3),x, algorithm="fricas")`

output `1/2*log(x^2 + 2*x + 3)`

3.136.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \frac{1+x}{3+2x+x^2} dx = \frac{\log(x^2 + 2x + 3)}{2}$$

input `integrate((1+x)/(x**2+2*x+3),x)`

output `log(x**2 + 2*x + 3)/2`

3.136.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1+x}{3+2x+x^2} dx = \frac{1}{2} \log(x^2 + 2x + 3)$$

input `integrate((1+x)/(x^2+2*x+3),x, algorithm="maxima")`

output `1/2*log(x^2 + 2*x + 3)`

3.136.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1+x}{3+2x+x^2} dx = \frac{1}{2} \log(x^2 + 2x + 3)$$

input `integrate((1+x)/(x^2+2*x+3),x, algorithm="giac")`

output `1/2*log(x^2 + 2*x + 3)`

3.136.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1+x}{3+2x+x^2} dx = \frac{\ln(x^2 + 2x + 3)}{2}$$

input `int((x + 1)/(2*x + x^2 + 3),x)`

output `log(2*x + x^2 + 3)/2`

3.137 $\int \tanh(x) dx$

3.137.1 Optimal result	759
3.137.2 Mathematica [A] (verified)	759
3.137.3 Rubi [A] (verified)	760
3.137.4 Maple [A] (verified)	761
3.137.5 Fricas [B] (verification not implemented)	761
3.137.6 Sympy [B] (verification not implemented)	761
3.137.7 Maxima [A] (verification not implemented)	762
3.137.8 Giac [B] (verification not implemented)	762
3.137.9 Mupad [B] (verification not implemented)	762

3.137.1 Optimal result

Integrand size = 2, antiderivative size = 3

$$\int \tanh(x) dx = \log(\cosh(x))$$

output `ln(cosh(x))`

3.137.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \tanh(x) dx = \log(\cosh(x))$$

input `Integrate[Tanh[x],x]`

output `Log[Cosh[x]]`

3.137.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \tanh(x) dx \\ \downarrow 3042 \\ \int -i \tan(ix) dx \\ \downarrow 26 \\ -i \int \tan(ix) dx \\ \downarrow 3956 \\ \log(\cosh(x)) \end{array}$$

input `Int [Tanh[x], x]`

output `Log[Cosh[x]]`

3.137.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

3.137.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
lookup	$\ln(\cosh(x))$	4
derivativedivides	$\ln(\cosh(x))$	4
default	$\ln(\cosh(x))$	4
risch	$-x + \ln(e^{2x} + 1)$	12
parallelrisc	$-\ln(1 - \tanh(x)) - x$	14

input `int(tanh(x),x,method=_RETURNVERBOSE)`

output `ln(cosh(x))`

3.137.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18 vs. $2(3) = 6$.

Time = 0.23 (sec) , antiderivative size = 18, normalized size of antiderivative = 6.00

$$\int \tanh(x) dx = -x + \log\left(\frac{2 \cosh(x)}{\cosh(x) - \sinh(x)}\right)$$

input `integrate(tanh(x),x, algorithm="fricas")`

output `-x + log(2*cosh(x)/(cosh(x) - sinh(x)))`

3.137.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7 vs. $2(3) = 6$.

Time = 0.05 (sec) , antiderivative size = 7, normalized size of antiderivative = 2.33

$$\int \tanh(x) dx = x - \log(\tanh(x) + 1)$$

input `integrate(tanh(x),x)`

output `x - log(tanh(x) + 1)`

3.137.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \tanh(x) dx = \log(\cosh(x))$$

input `integrate(tanh(x),x, algorithm="maxima")`

output `log(cosh(x))`

3.137.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. 2(3) = 6.

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 3.67

$$\int \tanh(x) dx = -x + \log(e^{2x} + 1)$$

input `integrate(tanh(x),x, algorithm="giac")`

output `-x + log(e^(2*x) + 1)`

3.137.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \tanh(x) dx = \ln(\cosh(x))$$

input `int(tanh(x),x)`

output `log(cosh(x))`

3.138 $\int (-x + x^3) dx$

3.138.1 Optimal result	763
3.138.2 Mathematica [A] (verified)	763
3.138.3 Rubi [A] (verified)	764
3.138.4 Maple [A] (verified)	764
3.138.5 Fricas [A] (verification not implemented)	765
3.138.6 Sympy [A] (verification not implemented)	765
3.138.7 Maxima [A] (verification not implemented)	765
3.138.8 Giac [A] (verification not implemented)	766
3.138.9 Mupad [B] (verification not implemented)	766

3.138.1 Optimal result

Integrand size = 7, antiderivative size = 15

$$\int (-x + x^3) dx = -\frac{x^2}{2} + \frac{x^4}{4}$$

output `-1/2*x^2+1/4*x^4`

3.138.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int (-x + x^3) dx = -\frac{x^2}{2} + \frac{x^4}{4}$$

input `Integrate[-x + x^3,x]`

output `-1/2*x^2 + x^4/4`

3.138.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x^3 - x) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^4}{4} - \frac{x^2}{2}$$

input `Int[-x + x^3,x]`

output `-1/2*x^2 + x^4/4`

3.138.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.138.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

method	result	size
gosper	$\frac{x^2(x^2-2)}{4}$	11
default	$-\frac{1}{2}x^2 + \frac{1}{4}x^4$	12
norman	$-\frac{1}{2}x^2 + \frac{1}{4}x^4$	12
risch	$-\frac{1}{2}x^2 + \frac{1}{4}x^4$	12
parallelrisc	$-\frac{1}{2}x^2 + \frac{1}{4}x^4$	12
parts	$-\frac{1}{2}x^2 + \frac{1}{4}x^4$	12

input `int(x^3-x,x,method=_RETURNVERBOSE)`

output `1/4*x^2*(x^2-2)`

3.138.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int (-x + x^3) dx = \frac{1}{4}x^4 - \frac{1}{2}x^2$$

input `integrate(x^3-x,x, algorithm="fricas")`

output `1/4*x^4 - 1/2*x^2`

3.138.6 Sympy [A] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.53

$$\int (-x + x^3) dx = \frac{x^4}{4} - \frac{x^2}{2}$$

input `integrate(x**3-x,x)`

output `x**4/4 - x**2/2`

3.138.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int (-x + x^3) dx = \frac{1}{4}x^4 - \frac{1}{2}x^2$$

input `integrate(x^3-x,x, algorithm="maxima")`

output `1/4*x^4 - 1/2*x^2`

3.138.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int (-x + x^3) dx = \frac{1}{4}x^4 - \frac{1}{2}x^2$$

input `integrate(x^3-x,x, algorithm="giac")`

output `1/4*x^4 - 1/2*x^2`

3.138.9 Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int (-x + x^3) dx = \frac{x^2(x^2 - 2)}{4}$$

input `int(x^3 - x,x)`

output `(x^2*(x^2 - 2))/4`

3.139 $\int \log(\sqrt{x}) dx$

3.139.1 Optimal result	767
3.139.2 Mathematica [A] (verified)	767
3.139.3 Rubi [A] (verified)	768
3.139.4 Maple [A] (verified)	768
3.139.5 Fricas [A] (verification not implemented)	769
3.139.6 Sympy [A] (verification not implemented)	769
3.139.7 Maxima [A] (verification not implemented)	769
3.139.8 Giac [A] (verification not implemented)	770
3.139.9 Mupad [B] (verification not implemented)	770

3.139.1 Optimal result

Integrand size = 6, antiderivative size = 14

$$\int \log(\sqrt{x}) dx = -\frac{x}{2} + x \log(\sqrt{x})$$

output `-1/2*x+1/2*x*ln(x)`

3.139.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \log(\sqrt{x}) dx = \frac{1}{2}(-x + x \log(x))$$

input `Integrate[Log[Sqrt[x]],x]`

output `(-x + x*Log[x])/2`

3.139.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2732}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \log(\sqrt{x}) dx$$

$$\downarrow \text{2732}$$

$$x \log(\sqrt{x}) - \frac{x}{2}$$

input `Int[Log[Sqrt[x]], x]`

output `-1/2*x + x*Log[Sqrt[x]]`

3.139.3.1 Defintions of rubi rules used

rule 2732 `Int[Log[(c_.)*(x_)^(n_.)], x_Symbol] :> Simp[x*Log[c*x^n], x] - Simp[n*x, x] /; FreeQ[{c, n}, x]`

3.139.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

method	result	size
lookup	$-\frac{x}{2} + \frac{x \ln(x)}{2}$	10
default	$-\frac{x}{2} + \frac{x \ln(x)}{2}$	10
norman	$-\frac{x}{2} + \frac{x \ln(x)}{2}$	10
risch	$-\frac{x}{2} + \frac{x \ln(x)}{2}$	10
parallelrisch	$-\frac{x}{2} + \frac{x \ln(x)}{2}$	10
parts	$-\frac{x}{2} + \frac{x \ln(x)}{2}$	10

input `int(1/2*ln(x), x, method=_RETURNVERBOSE)`

output `-1/2*x+1/2*x*ln(x)`

3.139.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.64

$$\int \log(\sqrt{x}) dx = \frac{1}{2} x \log(x) - \frac{1}{2} x$$

input `integrate(1/2*log(x),x, algorithm="fricas")`

output `1/2*x*log(x) - 1/2*x`

3.139.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.57

$$\int \log(\sqrt{x}) dx = \frac{x \log(x)}{2} - \frac{x}{2}$$

input `integrate(1/2*ln(x),x)`

output `x*log(x)/2 - x/2`

3.139.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.64

$$\int \log(\sqrt{x}) dx = \frac{1}{2} x \log(x) - \frac{1}{2} x$$

input `integrate(1/2*log(x),x, algorithm="maxima")`

output `1/2*x*log(x) - 1/2*x`

3.139.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.64

$$\int \log(\sqrt{x}) dx = \frac{1}{2} x \log(x) - \frac{1}{2} x$$

input `integrate(1/2*log(x),x, algorithm="giac")`

output `1/2*x*log(x) - 1/2*x`

3.139.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.50

$$\int \log(\sqrt{x}) dx = \frac{x(\ln(x) - 1)}{2}$$

input `int(log(x)/2,x)`

output `(x*(log(x) - 1))/2`

3.140 $\int \left(-e^{e^{-x}+e^x-x} + e^{e^{-x}+e^x+x} \right) dx$

3.140.1 Optimal result 771
 3.140.2 Mathematica [A] (verified) 771
 3.140.3 Rubi [F] 772
 3.140.4 Maple [A] (verified) 772
 3.140.5 Fricas [B] (verification not implemented) 773
 3.140.6 Sympy [A] (verification not implemented) 773
 3.140.7 Maxima [A] (verification not implemented) 773
 3.140.8 Giac [A] (verification not implemented) 774
 3.140.9 Mupad [B] (verification not implemented) 774

3.140.1 Optimal result

Integrand size = 29, antiderivative size = 11

$$\int \left(-e^{e^{-x}+e^x-x} + e^{e^{-x}+e^x+x} \right) dx = e^{e^{-x}+e^x}$$

output `exp(exp(x)+exp(-x))`

3.140.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \left(-e^{e^{-x}+e^x-x} + e^{e^{-x}+e^x+x} \right) dx = e^{e^{-x}+e^x}$$

input `Integrate[-E^(E^(-x) + E^x - x) + E^(E^(-x) + E^x + x), x]`

output `E^(E^(-x) + E^x)`

3.140. $\int \left(-e^{e^{-x}+e^x-x} + e^{e^{-x}+e^x+x} \right) dx$

3.140.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(e^{x+e^{-x}+e^x} - e^{-x+e^{-x}+e^x} \right) dx$$

↓ 2009

$$\text{Subst} \left(\int e^{x+\frac{1}{x}} dx, x, e^x \right) - \text{Subst} \left(\int \frac{e^{x+\frac{1}{x}}}{x^2} dx, x, e^x \right)$$

input `Int[-E^(E^(-x) + E^x - x) + E^(E^(-x) + E^x + x),x]`

output `$Aborted`

3.140.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.140.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

method	result	size
risch	$e^{e^x+e^{-x}}$	9
norman	$e^{e^x+e^{-x}-x}e^x$	15

input `int(exp(exp(x)+exp(-x)+x)-exp(exp(x)+exp(-x)-x),x,method=_RETURNVERBOSE)`

output `exp(exp(x)+exp(-x))`

3.140. $\int \left(-e^{-x+e^x-x} + e^{-x+e^x+x} \right) dx$

3.140.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. 2(8) = 16.

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.82

$$\int \left(-e^{e^{-x}+e^x-x} + e^{e^{-x}+e^x+x} \right) dx = e^{((xe^x+e^{2x})+1)e^{(-x)-x}}$$

input `integrate(exp(exp(x)+exp(-x)+x)-exp(exp(x)+exp(-x)-x),x, algorithm="fracas")`

output `e^((x*e^x + e^(2*x) + 1)*e^(-x) - x)`

3.140.6 Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.27

$$\int \left(-e^{e^{-x}+e^x-x} + e^{e^{-x}+e^x+x} \right) dx = e^x e^{-x+e^x+e^{-x}}$$

input `integrate(exp(exp(x)+exp(-x)+x)-exp(exp(x)+exp(-x)-x),x)`

output `exp(x)*exp(-x + exp(x) + exp(-x))`

3.140.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \left(-e^{e^{-x}+e^x-x} + e^{e^{-x}+e^x+x} \right) dx = e^{(e^{(-x)}+e^x)}$$

input `integrate(exp(exp(x)+exp(-x)+x)-exp(exp(x)+exp(-x)-x),x, algorithm="maxima")`

output `e^(e^(-x) + e^x)`

3.140. $\int \left(-e^{e^{-x}+e^x-x} + e^{e^{-x}+e^x+x} \right) dx$

3.140.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

$$\int \left(-e^{e^{-x}+e^x-x} + e^{e^{-x}+e^x+x} \right) dx = e^{(e^{2x}+1)e^{-x}}$$

input `integrate(exp(exp(x)+exp(-x)+x)-exp(exp(x)+exp(-x)-x),x, algorithm="giac")`output `e^((e^(2*x) + 1)*e^(-x))`**3.140.9 Mupad [B] (verification not implemented)**

Time = 14.76 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \left(-e^{e^{-x}+e^x-x} + e^{e^{-x}+e^x+x} \right) dx = e^{e^{-x}+e^x}$$

input `int(exp(x + exp(-x) + exp(x)) - exp(exp(-x) - x + exp(x)),x)`output `exp(exp(-x) + exp(x))`

3.141 $\int \frac{\log(\log(x))}{x \log(x)} dx$

3.141.1 Optimal result	775
3.141.2 Mathematica [A] (verified)	775
3.141.3 Rubi [A] (verified)	776
3.141.4 Maple [A] (verified)	777
3.141.5 Fricas [A] (verification not implemented)	777
3.141.6 Sympy [A] (verification not implemented)	777
3.141.7 Maxima [A] (verification not implemented)	778
3.141.8 Giac [A] (verification not implemented)	778
3.141.9 Mupad [B] (verification not implemented)	778

3.141.1 Optimal result

Integrand size = 11, antiderivative size = 9

$$\int \frac{\log(\log(x))}{x \log(x)} dx = \frac{1}{2} \log^2(\log(x))$$

output `1/2*ln(ln(x))^2`

3.141.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{\log(\log(x))}{x \log(x)} dx = \frac{1}{2} \log^2(\log(x))$$

input `Integrate[Log[Log[x]]/(x*Log[x]),x]`

output `Log[Log[x]]^2/2`

3.141.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3039, 2738}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{\log(\log(x))}{x \log(x)} dx \\ \downarrow \text{3039} \\ \int \frac{\log(\log(x))}{\log(x)} d\log(x) \\ \downarrow \text{2738} \\ \frac{1}{2} \log^2(\log(x)) \end{array}$$

input `Int[Log[Log[x]]/(x*Log[x]),x]`

output `Log[Log[x]]^2/2`

3.141.3.1 Defintions of rubi rules used

rule 2738 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))/(x_), x_Symbol] :> Simp[(a + b*Log[c*x^n])^2/(2*b*n), x] /; FreeQ[{a, b, c, n}, x]`

rule 3039 `Int[u_, x_Symbol] :> With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst[[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst] /; NonsumQ[u]`

3.141.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

method	result	size
derivativedivides	$\frac{\ln(\ln(x))^2}{2}$	8
default	$\frac{\ln(\ln(x))^2}{2}$	8
norman	$\frac{\ln(\ln(x))^2}{2}$	8
risch	$\frac{\ln(\ln(x))^2}{2}$	8

input `int(ln(ln(x))/x/ln(x),x,method=_RETURNVERBOSE)`output `1/2*ln(ln(x))^2`**3.141.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{\log(\log(x))}{x \log(x)} dx = \frac{1}{2} \log(\log(x))^2$$

input `integrate(log(log(x))/x/log(x),x, algorithm="fracas")`output `1/2*log(log(x))^2`**3.141.6 Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{\log(\log(x))}{x \log(x)} dx = \frac{\log(\log(x))^2}{2}$$

input `integrate(ln(ln(x))/x/ln(x),x)`output `log(log(x))**2/2`

3.141.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{\log(\log(x))}{x \log(x)} dx = \frac{1}{2} \log(\log(x))^2$$

input `integrate(log(log(x))/x/log(x),x, algorithm="maxima")`output `1/2*log(log(x))^2`**3.141.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{\log(\log(x))}{x \log(x)} dx = \frac{1}{2} \log(\log(x))^2$$

input `integrate(log(log(x))/x/log(x),x, algorithm="giac")`output `1/2*log(log(x))^2`**3.141.9 Mupad [B] (verification not implemented)**

Time = 15.85 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{\log(\log(x))}{x \log(x)} dx = \frac{\ln(\ln(x))^2}{2}$$

input `int(log(log(x))/(x*log(x)),x)`output `log(log(x))^2/2`

3.142 $\int \frac{1}{1+\tan^2(x)} dx$

3.142.1 Optimal result	779
3.142.2 Mathematica [A] (verified)	779
3.142.3 Rubi [A] (verified)	780
3.142.4 Maple [A] (verified)	781
3.142.5 Fricas [A] (verification not implemented)	782
3.142.6 Sympy [B] (verification not implemented)	782
3.142.7 Maxima [A] (verification not implemented)	782
3.142.8 Giac [A] (verification not implemented)	783
3.142.9 Mupad [B] (verification not implemented)	783

3.142.1 Optimal result

Integrand size = 8, antiderivative size = 14

$$\int \frac{1}{1 + \tan^2(x)} dx = \frac{x}{2} + \frac{1}{2} \cos(x) \sin(x)$$

output `1/2*x+1/2*cos(x)*sin(x)`

3.142.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \tan^2(x)} dx = \frac{x}{2} + \frac{1}{4} \sin(2x)$$

input `Integrate[(1 + Tan[x]^2)^(-1),x]`

output `x/2 + Sin[2*x]/4`

3.142.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3042, 4140, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\tan^2(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\tan(x)^2 + 1} dx \\
 & \quad \downarrow \text{4140} \\
 & \int \cos^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(x + \frac{\pi}{2}\right)^2 dx \\
 & \quad \downarrow \text{3115} \\
 & \frac{\int 1 dx}{2} + \frac{1}{2} \sin(x) \cos(x) \\
 & \quad \downarrow \text{24} \\
 & \frac{x}{2} + \frac{1}{2} \sin(x) \cos(x)
 \end{aligned}$$

input `Int[(1 + Tan[x]^2)^(-1), x]`

output `x/2 + (Cos[x]*Sin[x])/2`

3.142.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4140 `Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)^2]^(p_)), x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]`

3.142.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

method	result	size
risch	$\frac{x}{2} + \frac{\sin(2x)}{4}$	11
derivativedivides	$\frac{\tan(x)}{2+2\tan(x)^2} + \frac{\arctan(\tan(x))}{2}$	19
default	$\frac{\tan(x)}{2+2\tan(x)^2} + \frac{\arctan(\tan(x))}{2}$	19
parallelrisch	$\frac{x \tan(x)^2 + x + \tan(x)}{2+2\tan(x)^2}$	21
norman	$\frac{\frac{x}{2} + \frac{x \tan(x)^2}{2} + \frac{\tan(x)}{2}}{1+\tan(x)^2}$	25

input `int(1/(1+tan(x)^2),x,method=_RETURNVERBOSE)`

output `1/2*x+1/4*sin(2*x)`

3.142.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.43

$$\int \frac{1}{1 + \tan^2(x)} dx = \frac{x \tan(x)^2 + x + \tan(x)}{2(\tan(x)^2 + 1)}$$

input `integrate(1/(1+tan(x)^2),x, algorithm="fricas")`

output `1/2*(x*tan(x)^2 + x + tan(x))/(tan(x)^2 + 1)`

3.142.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 36 vs. $2(10) = 20$.

Time = 0.19 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.57

$$\int \frac{1}{1 + \tan^2(x)} dx = \frac{x \tan^2(x)}{2 \tan^2(x) + 2} + \frac{x}{2 \tan^2(x) + 2} + \frac{\tan(x)}{2 \tan^2(x) + 2}$$

input `integrate(1/(1+tan(x)**2),x)`

output `x*tan(x)**2/(2*tan(x)**2 + 2) + x/(2*tan(x)**2 + 2) + tan(x)/(2*tan(x)**2 + 2)`

3.142.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{1}{1 + \tan^2(x)} dx = \frac{1}{2} x + \frac{\tan(x)}{2(\tan(x)^2 + 1)}$$

input `integrate(1/(1+tan(x)^2),x, algorithm="maxima")`

output `1/2*x + 1/2*tan(x)/(tan(x)^2 + 1)`

3.142.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{1}{1 + \tan^2(x)} dx = \frac{1}{2}x + \frac{\tan(x)}{2(\tan(x)^2 + 1)}$$

input `integrate(1/(1+tan(x)^2),x, algorithm="giac")`

output `1/2*x + 1/2*tan(x)/(tan(x)^2 + 1)`

3.142.9 Mupad [B] (verification not implemented)

Time = 13.51 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{1}{1 + \tan^2(x)} dx = \frac{x}{2} + \frac{\sin(2x)}{4}$$

input `int(1/(tan(x)^2 + 1),x)`

output `x/2 + sin(2*x)/4`

3.143 $\int \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) dx$

3.143.1 Optimal result 784
 3.143.2 Mathematica [A] (verified) 784
 3.143.3 Rubi [A] (verified) 785
 3.143.4 Maple [A] (verified) 786
 3.143.5 Fricas [A] (verification not implemented) 787
 3.143.6 Sympy [A] (verification not implemented) 787
 3.143.7 Maxima [A] (verification not implemented) 787
 3.143.8 Giac [A] (verification not implemented) 788
 3.143.9 Mupad [B] (verification not implemented) 788

3.143.1 Optimal result

Integrand size = 10, antiderivative size = 45

$$\int \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) dx = 9\sqrt{9 - x^{2/3}} - \frac{1}{3}(9 - x^{2/3})^{3/2} + x \arcsin\left(\frac{\sqrt[3]{x}}{3}\right)$$

output `9*(9-x^(2/3))^(1/2)-1/3*(9-x^(2/3))^(3/2)+x*arcsin(1/3*x^(1/3))`

3.143.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.87

$$\int \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) dx = -\frac{1}{3}(-18 - x^{2/3}) \sqrt{9 - x^{2/3}} + x \arcsin\left(\frac{\sqrt[3]{x}}{3}\right)$$

input `Integrate[ArcSin[x^(1/3)/3],x]`

output `-1/3*((-18 - x^(2/3))*Sqrt[9 - x^(2/3)]) + x*ArcSin[x^(1/3)/3]`

3.143.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.11, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5339, 27, 798, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) dx \\
 & \quad \downarrow \text{5339} \\
 & x \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) - \int \frac{\sqrt[3]{x}}{3\sqrt{9-x^{2/3}}} dx \\
 & \quad \downarrow \text{27} \\
 & x \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) - \frac{1}{3} \int \frac{\sqrt[3]{x}}{\sqrt{9-x^{2/3}}} dx \\
 & \quad \downarrow \text{798} \\
 & x \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) - \frac{1}{2} \int \frac{x^{2/3}}{\sqrt{9-x^{2/3}}} dx^{2/3} \\
 & \quad \downarrow \text{53} \\
 & x \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) - \frac{1}{2} \int \left(\frac{9}{\sqrt{9-x^{2/3}}} - \sqrt{9-x^{2/3}} \right) dx^{2/3} \\
 & \quad \downarrow \text{2009} \\
 & x \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) + \frac{1}{2} \left(18\sqrt{9-x^{2/3}} - \frac{2}{3}(9-x^{2/3})^{3/2} \right)
 \end{aligned}$$

input `Int[ArcSin[x^(1/3)/3],x]`

output `(18*Sqrt[9 - x^(2/3)] - (2*(9 - x^(2/3))^(3/2))/3)/2 + x*ArcSin[x^(1/3)/3]`

3.143.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 53 `Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.)), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^(p_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5339 `Int[ArcSin[u_], x_Symbol] := Simp[x*ArcSin[u], x] - Int[SimplifyIntegrand[x*(D[u, x]/Sqrt[1 - u^2]), x], x] /; InverseFunctionFreeQ[u, x] && !FunctionOfExponentialQ[u, x]`

3.143.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.76

method	result	size
derivativedivides	$x \arcsin\left(\frac{x^{\frac{1}{3}}}{3}\right) + x^{\frac{2}{3}} \sqrt{-\frac{x^{\frac{2}{3}}}{9} + 1} + 18 \sqrt{-\frac{x^{\frac{2}{3}}}{9} + 1}$	34
default	$x \arcsin\left(\frac{x^{\frac{1}{3}}}{3}\right) + x^{\frac{2}{3}} \sqrt{-\frac{x^{\frac{2}{3}}}{9} + 1} + 18 \sqrt{-\frac{x^{\frac{2}{3}}}{9} + 1}$	34
parts	$x \arcsin\left(\frac{x^{\frac{1}{3}}}{3}\right) + \frac{x^{\frac{2}{3}} \sqrt{9 - x^{\frac{2}{3}}}}{3} + 6 \sqrt{9 - x^{\frac{2}{3}}}$	35

input `int(arcsin(1/3*x^(1/3)),x,method=_RETURNVERBOSE)`

output `x*arcsin(1/3*x^(1/3))+x^(2/3)*(-1/9*x^(2/3)+1)^(1/2)+18*(-1/9*x^(2/3)+1)^(1/2)`

3.143. $\int \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) dx$

3.143.5 Fricas [A] (verification not implemented)

Time = 4.29 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.56

$$\int \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) dx = x \arcsin\left(\frac{1}{3}x^{\frac{1}{3}}\right) + \frac{1}{3}\left(x^{\frac{2}{3}} + 18\right)\sqrt{-x^{\frac{2}{3}} + 9}$$

input `integrate(arcsin(1/3*x^(1/3)),x, algorithm="fricas")`output `x*arcsin(1/3*x^(1/3)) + 1/3*(x^(2/3) + 18)*sqrt(-x^(2/3) + 9)`**3.143.6 Sympy [A] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.82

$$\int \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) dx = \frac{x^{\frac{2}{3}}\sqrt{9 - x^{\frac{2}{3}}}}{3} + x \operatorname{asin}\left(\frac{\sqrt[3]{x}}{3}\right) + 6\sqrt{9 - x^{\frac{2}{3}}}$$

input `integrate(asin(1/3*x**(1/3)),x)`output `x**(2/3)*sqrt(9 - x**(2/3))/3 + x*asin(x**(1/3)/3) + 6*sqrt(9 - x**(2/3))`**3.143.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.73

$$\int \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) dx = x \arcsin\left(\frac{1}{3}x^{\frac{1}{3}}\right) + x^{\frac{2}{3}}\sqrt{-\frac{1}{9}x^{\frac{2}{3}} + 1} + 18\sqrt{-\frac{1}{9}x^{\frac{2}{3}} + 1}$$

input `integrate(arcsin(1/3*x^(1/3)),x, algorithm="maxima")`output `x*arcsin(1/3*x^(1/3)) + x^(2/3)*sqrt(-1/9*x^(2/3) + 1) + 18*sqrt(-1/9*x^(2/3) + 1)`

3.143.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.09

$$\int \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) dx = x^{\frac{1}{3}}\left(x^{\frac{2}{3}} - 9\right) \arcsin\left(\frac{1}{3}x^{\frac{1}{3}}\right) - 9\left(-\frac{1}{9}x^{\frac{2}{3}} + 1\right)^{\frac{3}{2}} + 9x^{\frac{1}{3}} \arcsin\left(\frac{1}{3}x^{\frac{1}{3}}\right) + 27\sqrt{-\frac{1}{9}x^{\frac{2}{3}} + 1}$$

input `integrate(arcsin(1/3*x^(1/3)),x, algorithm="giac")`output `x^(1/3)*(x^(2/3) - 9)*arcsin(1/3*x^(1/3)) - 9*(-1/9*x^(2/3) + 1)^(3/2) + 9*x^(1/3)*arcsin(1/3*x^(1/3)) + 27*sqrt(-1/9*x^(2/3) + 1)`**3.143.9 Mupad [B] (verification not implemented)**

Time = 14.14 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.56

$$\int \arcsin\left(\frac{\sqrt[3]{x}}{3}\right) dx = x \operatorname{asin}\left(\frac{x^{1/3}}{3}\right) + \frac{\sqrt{9 - x^{2/3}}(x^{2/3} + 18)}{3}$$

input `int(asin(x^(1/3)/3),x)`output `x*asin(x^(1/3)/3) + ((9 - x^(2/3))^(1/2)*(x^(2/3) + 18))/3`

3.144 $\int \log(x) dx$

3.144.1 Optimal result	789
3.144.2 Mathematica [A] (verified)	789
3.144.3 Rubi [A] (verified)	790
3.144.4 Maple [A] (verified)	790
3.144.5 Fricas [A] (verification not implemented)	791
3.144.6 Sympy [A] (verification not implemented)	791
3.144.7 Maxima [A] (verification not implemented)	791
3.144.8 Giac [A] (verification not implemented)	792
3.144.9 Mupad [B] (verification not implemented)	792

3.144.1 Optimal result

Integrand size = 2, antiderivative size = 8

$$\int \log(x) dx = -x + x \log(x)$$

output `x*ln(x)-x`

3.144.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \log(x) dx = -x + x \log(x)$$

input `Integrate[Log[x],x]`

output `-x + x*Log[x]`

3.144.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {2732}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \log(x) dx$$

$$\downarrow \text{2732}$$

$$x \log(x) - x$$

input `Int [Log[x] , x]`

output `-x + x*Log[x]`

3.144.3.1 Defintions of rubi rules used

rule 2732 `Int [Log[(c_.)*(x_)^(n_.)], x_Symbol] :> Simp[x*Log[c*x^n], x] - Simp[n*x, x] /; FreeQ[{c, n}, x]`

3.144.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
lookup	$x \ln(x) - x$	9
default	$x \ln(x) - x$	9
norman	$x \ln(x) - x$	9
risch	$x \ln(x) - x$	9
parallelrisch	$x \ln(x) - x$	9
parts	$x \ln(x) - x$	9

input `int(ln(x), x, method=_RETURNVERBOSE)`

output `x*ln(x)-x`

3.144.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \log(x) dx = x \log(x) - x$$

input `integrate(log(x),x, algorithm="fricas")`

output `x*log(x) - x`

3.144.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \log(x) dx = x \log(x) - x$$

input `integrate(ln(x),x)`

output `x*log(x) - x`

3.144.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \log(x) dx = x \log(x) - x$$

input `integrate(log(x),x, algorithm="maxima")`

output `x*log(x) - x`

3.144.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \log(x) dx = x \log(x) - x$$

input `integrate(log(x),x, algorithm="giac")`

output `x*log(x) - x`

3.144.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \log(x) dx = x (\ln(x) - 1)$$

input `int(log(x),x)`

output `x*(log(x) - 1)`

3.145 $\int e^x(\cos(x) - \sin(x)) dx$

3.145.1 Optimal result	793
3.145.2 Mathematica [A] (verified)	793
3.145.3 Rubi [A] (verified)	794
3.145.4 Maple [A] (verified)	794
3.145.5 Fricas [A] (verification not implemented)	795
3.145.6 Sympy [A] (verification not implemented)	795
3.145.7 Maxima [B] (verification not implemented)	795
3.145.8 Giac [B] (verification not implemented)	796
3.145.9 Mupad [B] (verification not implemented)	796

3.145.1 Optimal result

Integrand size = 11, antiderivative size = 6

$$\int e^x(\cos(x) - \sin(x)) dx = e^x \cos(x)$$

output `exp(x)*cos(x)`

3.145.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int e^x(\cos(x) - \sin(x)) dx = e^x \cos(x)$$

input `Integrate[E^x*(Cos[x] - Sin[x]),x]`

output `E^x*Cos[x]`

3.145.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {2726}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x(\cos(x) - \sin(x)) dx$$

↓ 2726

$$e^x \cos(x)$$

input `Int[E^x*(Cos[x] - Sin[x]),x]`

output `E^x*cos[x]`

3.145.3.1 Defintions of rubi rules used

rule 2726 `Int[(y_.)*(F_)^(u_)*((v_) + (w_)), x_Symbol] := With[{z = v*(y/(Log[F]*D[u, x]))}, Simp[F^u*z, x] /; EqQ[D[z, x], w*y]] /; FreeQ[F, x]`

3.145.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

method	result	size
default	$e^x \cos(x)$	6
parallelrisc	$e^x \cos(x)$	6
parts	$e^x \cos(x)$	6
risc	$\frac{e^{(1+i)x}}{2} + \frac{e^{(1-i)x}}{2}$	18
norman	$\frac{-e^x \tan(\frac{x}{2})^2 + e^x}{1 + \tan(\frac{x}{2})^2}$	25

input `int(exp(x)*(cos(x)-sin(x)),x,method=_RETURNVERBOSE)`

output `exp(x)*cos(x)`

3.145.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int e^x(\cos(x) - \sin(x)) dx = \cos(x) e^x$$

input `integrate(exp(x)*(cos(x)-sin(x)),x, algorithm="fricas")`

output `cos(x)*e^x`

3.145.6 Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int e^x(\cos(x) - \sin(x)) dx = e^x \cos(x)$$

input `integrate(exp(x)*(cos(x)-sin(x)),x)`

output `exp(x)*cos(x)`

3.145.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. $2(5) = 10$.

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 3.50

$$\int e^x(\cos(x) - \sin(x)) dx = \frac{1}{2}(\cos(x) + \sin(x))e^x + \frac{1}{2}(\cos(x) - \sin(x))e^x$$

input `integrate(exp(x)*(cos(x)-sin(x)),x, algorithm="maxima")`

output `1/2*(cos(x) + sin(x))*e^x + 1/2*(cos(x) - sin(x))*e^x`

3.145.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. 2(5) = 10.

Time = 0.29 (sec) , antiderivative size = 21, normalized size of antiderivative = 3.50

$$\int e^x(\cos(x) - \sin(x)) dx = \frac{1}{2}(\cos(x) + \sin(x))e^x + \frac{1}{2}(\cos(x) - \sin(x))e^x$$

input `integrate(exp(x)*(cos(x)-sin(x)),x, algorithm="giac")`

output `1/2*(cos(x) + sin(x))*e^x + 1/2*(cos(x) - sin(x))*e^x`

3.145.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int e^x(\cos(x) - \sin(x)) dx = e^x \cos(x)$$

input `int(exp(x)*(cos(x) - sin(x)),x)`

output `exp(x)*cos(x)`

3.146 $\int e^{-x^2} x^3 dx$

3.146.1 Optimal result	797
3.146.2 Mathematica [A] (verified)	797
3.146.3 Rubi [A] (verified)	798
3.146.4 Maple [A] (verified)	799
3.146.5 Fricas [A] (verification not implemented)	799
3.146.6 Sympy [A] (verification not implemented)	800
3.146.7 Maxima [A] (verification not implemented)	800
3.146.8 Giac [A] (verification not implemented)	800
3.146.9 Mupad [B] (verification not implemented)	801

3.146.1 Optimal result

Integrand size = 11, antiderivative size = 26

$$\int e^{-x^2} x^3 dx = -\frac{e^{-x^2}}{2} - \frac{1}{2}e^{-x^2} x^2$$

output `-1/2/exp(x^2)-1/2*x^2/exp(x^2)`

3.146.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.62

$$\int e^{-x^2} x^3 dx = -\frac{1}{2}e^{-x^2} (1 + x^2)$$

input `Integrate[x^3/E^x^2,x]`

output `-1/2*(1 + x^2)/E^x^2`

3.146.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2641, 2638}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{-x^2} x^3 dx$$

$$\downarrow \text{2641}$$

$$\int e^{-x^2} x dx - \frac{1}{2} e^{-x^2} x^2$$

$$\downarrow \text{2638}$$

$$-\frac{1}{2} e^{-x^2} x^2 - \frac{e^{-x^2}}{2}$$

input `Int[x^3/E^x^2,x]`

output `-1/2*1/E^x^2 - x^2/(2*E^x^2)`

3.146.3.1 Defintions of rubi rules used

rule 2638 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[(e + f*x)^n*(F^(a + b*(c + d*x)^n)/(b*f*n*(c + d*x)^n *Log[F])), x] /; FreeQ[{F, a, b, c, d, e, f, n}, x] && EqQ[m, n - 1] && EqQ[d*e - c*f, 0]`

rule 2641 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^(m - n + 1)*(F^(a + b*(c + d*x)^n)/(b*d*n*Log[F])), x] - Simp[(m - n + 1)/(b*n*Log[F]) Int[(c + d*x)^(m - n)*F^(a + b*(c + d*x)^n), x], x] /; FreeQ[{F, a, b, c, d}, x] && IntegerQ[2*((m + 1)/n)] && LtQ[0, (m + 1)/n, 5] && IntegerQ[n] && (LtQ[0, n, m + 1] || LtQ[m, n, 0])`

3.146.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.54

method	result	size
gospers	$-\frac{(x^2+1)e^{-x^2}}{2}$	14
risch	$\left(-\frac{x^2}{2} - \frac{1}{2}\right) e^{-x^2}$	15
meijerg	$\frac{1}{2} - \frac{(2x^2+2)e^{-x^2}}{4}$	18
default	$-\frac{x^2 e^{-x^2}}{2} - \frac{e^{-x^2}}{2}$	21
norman	$-\frac{x^2 e^{-x^2}}{2} - \frac{e^{-x^2}}{2}$	21
parallelrisch	$-\frac{x^2 e^{-x^2}}{2} - \frac{e^{-x^2}}{2}$	21
parts	$\frac{\sqrt{\pi} \operatorname{erf}(x)x^3}{2} - \frac{3\sqrt{\pi} \left(\frac{\operatorname{erf}(x)x^3}{3} - \frac{2 \left(-\frac{x^2 e^{-x^2}}{2} - \frac{e^{-x^2}}{2} \right)}{3\sqrt{\pi}} \right)}{2}$	50

input `int(x^3*exp(-x^2),x,method=_RETURNVERBOSE)`output `-1/2*(x^2+1)*exp(-x^2)`**3.146.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.50

$$\int e^{-x^2} x^3 dx = -\frac{1}{2} (x^2 + 1) e^{(-x^2)}$$

input `integrate(x^3*exp(-x^2),x, algorithm="fricas")`output `-1/2*(x^2 + 1)*e^(-x^2)`

3.146.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.46

$$\int e^{-x^2} x^3 dx = \frac{(-x^2 - 1) e^{-x^2}}{2}$$

input `integrate(x**3*exp(-x**2),x)`output `(-x**2 - 1)*exp(-x**2)/2`**3.146.7 Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.50

$$\int e^{-x^2} x^3 dx = -\frac{1}{2} (x^2 + 1) e^{(-x^2)}$$

input `integrate(x^3*exp(-x^2),x, algorithm="maxima")`output `-1/2*(x^2 + 1)*e^(-x^2)`**3.146.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.50

$$\int e^{-x^2} x^3 dx = -\frac{1}{2} (x^2 + 1) e^{(-x^2)}$$

input `integrate(x^3*exp(-x^2),x, algorithm="giac")`output `-1/2*(x^2 + 1)*e^(-x^2)`

3.146.9 Mupad [B] (verification not implemented)

Time = 16.55 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.50

$$\int e^{-x^2} x^3 dx = -\frac{e^{-x^2} (x^2 + 1)}{2}$$

input `int(x^3*exp(-x^2),x)`

output `-(exp(-x^2)*(x^2 + 1))/2`

$$3.147 \quad \int \left((1 + 2e^{x^2}x) \cos(x) - (e^{x^2} + x) \sin(x) \right) dx$$

3.147.1 Optimal result	802
3.147.2 Mathematica [C] (verified)	802
3.147.3 Rubi [F]	803
3.147.4 Maple [A] (verified)	803
3.147.5 Fricas [A] (verification not implemented)	804
3.147.6 Sympy [F]	804
3.147.7 Maxima [C] (verification not implemented)	804
3.147.8 Giac [A] (verification not implemented)	805
3.147.9 Mupad [B] (verification not implemented)	806

3.147.1 Optimal result

Integrand size = 25, antiderivative size = 10

$$\int \left((1 + 2e^{x^2}x) \cos(x) - (e^{x^2} + x) \sin(x) \right) dx = (e^{x^2} + x) \cos(x)$$

output `(exp(x^2)+x)*cos(x)`

3.147.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.63 (sec) , antiderivative size = 27, normalized size of antiderivative = 2.70

$$\int \left((1 + 2e^{x^2}x) \cos(x) - (e^{x^2} + x) \sin(x) \right) dx = \frac{1}{2}e^{-ix}(1 + e^{2ix})(e^{x^2} + x)$$

input `Integrate[(1 + 2*E^x^2*x)*Cos[x] - (E^x^2 + x)*Sin[x],x]`

output `((1 + E^((2*I)*x))*(E^x^2 + x))/(2*E^(I*x))`

$$3.147. \quad \int \left((1 + 2e^{x^2}x) \cos(x) - (e^{x^2} + x) \sin(x) \right) dx$$

3.147.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left((2e^{x^2}x + 1) \cos(x) - (e^{x^2} + x) \sin(x) \right) dx$$

↓ 2009

$$2 \int e^{x^2} x \cos(x) dx - \frac{1}{4} i \sqrt[4]{e} \sqrt{\pi} \operatorname{erfi} \left(\frac{1}{2} (2x - i) \right) + \frac{1}{4} i \sqrt[4]{e} \sqrt{\pi} \operatorname{erfi} \left(\frac{1}{2} (2x + i) \right) + x \cos(x)$$

input `Int[(1 + 2*E^x^2*x)*Cos[x] - (E^x^2 + x)*Sin[x],x]`

output `$Aborted`

3.147.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.147.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

method	result	size
parallelrisch	$(e^{x^2} + x) \cos(x)$	10
risch	$\frac{e^{x(x-i)}}{2} + \frac{e^{x(i+x)}}{2} + x \cos(x)$	24
norman	$\frac{x - x \tan(\frac{x}{2})^2 - e^{x^2} \tan(\frac{x}{2})^2 + e^{x^2}}{1 + \tan(\frac{x}{2})^2}$	39

input `int((2*exp(x^2)*x+1)*cos(x)-(exp(x^2)+x)*sin(x),x,method=_RETURNVERBOSE)`

output `(exp(x^2)+x)*cos(x)`

3.147. $\int \left((1 + 2e^{x^2}x) \cos(x) - (e^{x^2} + x) \sin(x) \right) dx$

3.147.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \left((1 + 2e^{x^2} x) \cos(x) - (e^{x^2} + x) \sin(x) \right) dx = x \cos(x) + \cos(x) e^{(x^2)}$$

input `integrate((2*exp(x^2)*x+1)*cos(x)-(exp(x^2)+x)*sin(x),x, algorithm="fricas")`

output `x*cos(x) + cos(x)*e^(x^2)`

3.147.6 Sympy [F]

$$\begin{aligned} & \int \left((1 + 2e^{x^2} x) \cos(x) - (e^{x^2} + x) \sin(x) \right) dx \\ &= \int \left(-(x + e^{x^2}) \sin(x) + (2xe^{x^2} + 1) \cos(x) \right) dx \end{aligned}$$

input `integrate((2*exp(x**2)*x+1)*cos(x)-(exp(x**2)+x)*sin(x),x)`

output `Integral(-(x + exp(x**2))*sin(x) + (2*x*exp(x**2) + 1)*cos(x), x)`

3.147.7 Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.28 (sec) , antiderivative size = 473, normalized size of antiderivative = 47.30

$$\begin{aligned} & \int \left((1 + 2e^{x^2} x) \cos(x) - (e^{x^2} + x) \sin(x) \right) dx \\ &= -\frac{1}{4} \sqrt{\pi} \left(\operatorname{erf} \left(ix + \frac{1}{2} \right) - \operatorname{erf} \left(ix - \frac{1}{2} \right) \right) e^{\frac{1}{4}} + x \cos(x) \\ & \quad + \frac{2(16x^4 + 8x^2 + 1)^{\frac{1}{4}} \left(e^{(x^2+ix-\frac{1}{4})} + e^{(x^2-ix-\frac{1}{4})} + e^{(\bar{x}^2+i\bar{x}-\frac{1}{4})} + e^{(\bar{x}^2-i\bar{x}-\frac{1}{4})} \right) e^{\frac{1}{4}} - \left(2 \left(i \sqrt{\pi} \left(\operatorname{erf} \left(\sqrt{-x^2} \right) \right) \right)} \right)}{4} \end{aligned}$$

3.147. $\int \left((1 + 2e^{x^2} x) \cos(x) - (e^{x^2} + x) \sin(x) \right) dx$

input `integrate((2*exp(x^2)*x+1)*cos(x)-(exp(x^2)+x)*sin(x),x, algorithm="maxima")`

output `-1/4*sqrt(pi)*(erf(I*x + 1/2) - erf(I*x - 1/2))*e^(1/4) + x*cos(x) + 1/8*(2*(16*x^4 + 8*x^2 + 1)^(1/4)*(e^(x^2 + I*x - 1/4) + e^(x^2 - I*x - 1/4) + e^(conjugate(x)^2 + I*conjugate(x) - 1/4) + e^(conjugate(x)^2 - I*conjugate(x) - 1/4))*e^(1/4) - ((2*(I*sqrt(pi)*(conjugate(erf(sqrt(-x^2 + I*x + 1/4)))) - 1) - I*sqrt(pi)*(conjugate(erf(sqrt(-x^2 - I*x + 1/4)))) - 1) - I*sqrt(pi)*(erf(sqrt(-x^2 + I*x + 1/4)) - 1) + I*sqrt(pi)*(erf(sqrt(-x^2 - I*x + 1/4)) - 1))*x - sqrt(pi)*(conjugate(erf(sqrt(-x^2 + I*x + 1/4)))) - 1) - sqrt(pi)*(conjugate(erf(sqrt(-x^2 - I*x + 1/4)))) - 1) - sqrt(pi)*(erf(sqrt(-x^2 + I*x + 1/4)) - 1) - sqrt(pi)*(erf(sqrt(-x^2 - I*x + 1/4)) - 1))*cos(1/2*arctan2(4*x, -4*x^2 + 1)) - (2*(sqrt(pi)*(conjugate(erf(sqrt(-x^2 + I*x + 1/4)))) - 1) + sqrt(pi)*(conjugate(erf(sqrt(-x^2 - I*x + 1/4)))) - 1) + sqrt(pi)*(erf(sqrt(-x^2 + I*x + 1/4)) - 1) + sqrt(pi)*(erf(sqrt(-x^2 - I*x + 1/4)) - 1))*x + I*sqrt(pi)*(conjugate(erf(sqrt(-x^2 + I*x + 1/4)))) - 1) - I*sqrt(pi)*(conjugate(erf(sqrt(-x^2 - I*x + 1/4)))) - 1) - I*sqrt(pi)*(erf(sqrt(-x^2 + I*x + 1/4)) - 1) + I*sqrt(pi)*(erf(sqrt(-x^2 - I*x + 1/4)) - 1))*sin(1/2*arctan2(4*x, -4*x^2 + 1))*e^(1/4))/(16*x^4 + 8*x^2 + 1)^(1/4)`

3.147.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \left((1 + 2e^{x^2}x) \cos(x) - (e^{x^2} + x) \sin(x) \right) dx = x \cos(x) + \cos(x) e^{(x^2)}$$

input `integrate((2*exp(x^2)*x+1)*cos(x)-(exp(x^2)+x)*sin(x),x, algorithm="giac")`

output `x*cos(x) + cos(x)*e^(x^2)`

3.147. $\int \left((1 + 2e^{x^2}x) \cos(x) - (e^{x^2} + x) \sin(x) \right) dx$

3.147.9 Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \left((1 + 2e^{x^2}x) \cos(x) - (e^{x^2} + x) \sin(x) \right) dx = \cos(x) (x + e^{x^2})$$

input `int(cos(x)*(2*x*exp(x^2) + 1) - sin(x)*(x + exp(x^2)),x)`

output `cos(x)*(x + exp(x^2))`

3.148 $\int \left(1 + \frac{1}{\sqrt{x}} + \frac{1}{\sqrt[3]{x}} \right) (1 + \sqrt[3]{x} + \sqrt{x}) dx$

3.148.1 Optimal result	807
3.148.2 Mathematica [A] (verified)	807
3.148.3 Rubi [A] (verified)	808
3.148.4 Maple [A] (verified)	809
3.148.5 Fricas [A] (verification not implemented)	809
3.148.6 Sympy [A] (verification not implemented)	809
3.148.7 Maxima [A] (verification not implemented)	810
3.148.8 Giac [A] (verification not implemented)	810
3.148.9 Mupad [B] (verification not implemented)	810

3.148.1 Optimal result

Integrand size = 25, antiderivative size = 56

$$\int \left(1 + \frac{1}{\sqrt{x}} + \frac{1}{\sqrt[3]{x}} \right) (1 + \sqrt[3]{x} + \sqrt{x}) dx = 2\sqrt{x} + \frac{3x^{2/3}}{2} + \frac{6x^{5/6}}{5} + 3x + \frac{6x^{7/6}}{7} + \frac{3x^{4/3}}{4} + \frac{2x^{3/2}}{3}$$

output `2*x^(1/2)+3/2*x^(2/3)+6/5*x^(5/6)+3*x+6/7*x^(7/6)+3/4*x^(4/3)+2/3*x^(3/2)`

3.148.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.89

$$\int \left(1 + \frac{1}{\sqrt{x}} + \frac{1}{\sqrt[3]{x}} \right) (1 + \sqrt[3]{x} + \sqrt{x}) dx = \frac{1}{420} (840\sqrt{x} + 630x^{2/3} + 504x^{5/6} + 1260x + 360x^{7/6} + 315x^{4/3} + 280x^{3/2})$$

input `Integrate[(1 + 1/Sqrt[x] + x^(-1/3))*(1 + x^(1/3) + Sqrt[x]),x]`

output `(840*Sqrt[x] + 630*x^(2/3) + 504*x^(5/6) + 1260*x + 360*x^(7/6) + 315*x^(4/3) + 280*x^(3/2))/420`

3.148. $\int \left(1 + \frac{1}{\sqrt{x}} + \frac{1}{\sqrt[3]{x}} \right) (1 + \sqrt[3]{x} + \sqrt{x}) dx$

3.148.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.11, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {7267, 7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\frac{1}{\sqrt[3]{x}} + \frac{1}{\sqrt{x}} + 1 \right) (\sqrt{x} + \sqrt[3]{x} + 1) dx$$

$$\downarrow \text{7267}$$

$$6 \int (\sqrt{x} + \sqrt[6]{x} + 1) (\sqrt{x} + \sqrt[3]{x} + 1) \sqrt[3]{x} d\sqrt[6]{x}$$

$$\downarrow \text{7293}$$

$$6 \int \left(x^{4/3} + x^{7/6} + x + 3x^{5/6} + x^{2/3} + \sqrt{x} + \sqrt[3]{x} \right) d\sqrt[6]{x}$$

$$\downarrow \text{2009}$$

$$6 \left(\frac{x^{3/2}}{9} + \frac{x^{4/3}}{8} + \frac{x^{7/6}}{7} + \frac{x^{5/6}}{5} + \frac{x^{2/3}}{4} + \frac{x}{2} + \frac{\sqrt{x}}{3} \right)$$

input `Int[(1 + 1/Sqrt[x] + x^(-1/3))*(1 + x^(1/3) + Sqrt[x]),x]`

output `6*(Sqrt[x]/3 + x^(2/3)/4 + x^(5/6)/5 + x/2 + x^(7/6)/7 + x^(4/3)/8 + x^(3/2)/9)`

3.148.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7267 `Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Simp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2]])], x] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

3.148. $\int \left(1 + \frac{1}{\sqrt{x}} + \frac{1}{\sqrt[3]{x}} \right) (1 + \sqrt[3]{x} + \sqrt{x}) dx$

3.148.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.62

$$2\sqrt{x} + \frac{3x^{\frac{2}{3}}}{2} + \frac{6x^{\frac{5}{6}}}{5} + 3x + \frac{6x^{\frac{7}{6}}}{7} + \frac{3x^{\frac{4}{3}}}{4} + \frac{2x^{\frac{3}{2}}}{3}$$

input `int((1+x^(1/2)+x^(1/3))*(1+1/x^(1/2)+1/x^(1/3)),x)`output `2*x^(1/2)+3/2*x^(2/3)+6/5*x^(5/6)+3*x+6/7*x^(7/6)+3/4*x^(4/3)+2/3*x^(3/2)`**3.148.5 Fricas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.57

$$\int \left(1 + \frac{1}{\sqrt{x}} + \frac{1}{\sqrt[3]{x}}\right) (1 + \sqrt[3]{x} + \sqrt{x}) dx = \frac{2}{3}(x+3)\sqrt{x} + \frac{3}{4}x^{\frac{4}{3}} + \frac{6}{7}x^{\frac{7}{6}} + 3x + \frac{6}{5}x^{\frac{5}{6}} + \frac{3}{2}x^{\frac{2}{3}}$$

input `integrate((1+x^(1/2)+x^(1/3))*(1+1/x^(1/2)+1/x^(1/3)),x, algorithm="fricas")`output `2/3*(x + 3)*sqrt(x) + 3/4*x^(4/3) + 6/7*x^(7/6) + 3*x + 6/5*x^(5/6) + 3/2*x^(2/3)`**3.148.6 Sympy [A] (verification not implemented)**

Time = 4.52 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.91

$$\int \left(1 + \frac{1}{\sqrt{x}} + \frac{1}{\sqrt[3]{x}}\right) (1 + \sqrt[3]{x} + \sqrt{x}) dx = \frac{6x^{\frac{7}{6}}}{7} + \frac{6x^{\frac{5}{6}}}{5} + \frac{3x^{\frac{4}{3}}}{4} + \frac{3x^{\frac{2}{3}}}{2} + \frac{2x^{\frac{3}{2}}}{3} + 2\sqrt{x} + 3x$$

input `integrate((1+x**(1/2)+x**(1/3))*(1+1/x**(1/2)+1/x**(1/3)),x)`output `6*x**(7/6)/7 + 6*x**(5/6)/5 + 3*x**(4/3)/4 + 3*x**(2/3)/2 + 2*x**(3/2)/3 + 2*sqrt(x) + 3*x`

3.148. $\int \left(1 + \frac{1}{\sqrt{x}} + \frac{1}{\sqrt[3]{x}}\right) (1 + \sqrt[3]{x} + \sqrt{x}) dx$

3.148.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.61

$$\int \left(1 + \frac{1}{\sqrt{x}} + \frac{1}{\sqrt[3]{x}}\right) (1 + \sqrt[3]{x} + \sqrt{x}) dx = \frac{2}{3} x^{\frac{3}{2}} + \frac{3}{4} x^{\frac{4}{3}} + \frac{6}{7} x^{\frac{7}{6}} + 3x + \frac{6}{5} x^{\frac{5}{6}} + \frac{3}{2} x^{\frac{2}{3}} + 2\sqrt{x}$$

input `integrate((1+x^(1/2)+x^(1/3))*(1+1/x^(1/2)+1/x^(1/3)),x, algorithm="maxima")`

output `2/3*x^(3/2) + 3/4*x^(4/3) + 6/7*x^(7/6) + 3*x + 6/5*x^(5/6) + 3/2*x^(2/3) + 2*sqrt(x)`

3.148.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.61

$$\int \left(1 + \frac{1}{\sqrt{x}} + \frac{1}{\sqrt[3]{x}}\right) (1 + \sqrt[3]{x} + \sqrt{x}) dx = \frac{2}{3} x^{\frac{3}{2}} + \frac{3}{4} x^{\frac{4}{3}} + \frac{6}{7} x^{\frac{7}{6}} + 3x + \frac{6}{5} x^{\frac{5}{6}} + \frac{3}{2} x^{\frac{2}{3}} + 2\sqrt{x}$$

input `integrate((1+x^(1/2)+x^(1/3))*(1+1/x^(1/2)+1/x^(1/3)),x, algorithm="giac")`

output `2/3*x^(3/2) + 3/4*x^(4/3) + 6/7*x^(7/6) + 3*x + 6/5*x^(5/6) + 3/2*x^(2/3) + 2*sqrt(x)`

3.148.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.61

$$\int \left(1 + \frac{1}{\sqrt{x}} + \frac{1}{\sqrt[3]{x}}\right) (1 + \sqrt[3]{x} + \sqrt{x}) dx = 3x + 2\sqrt{x} + \frac{3x^{2/3}}{2} + \frac{2x^{3/2}}{3} + \frac{3x^{4/3}}{4} + \frac{6x^{5/6}}{5} + \frac{6x^{7/6}}{7}$$

input `int((1/x^(1/2) + 1/x^(1/3) + 1)*(x^(1/2) + x^(1/3) + 1),x)`

output `3*x + 2*x^(1/2) + (3*x^(2/3))/2 + (2*x^(3/2))/3 + (3*x^(4/3))/4 + (6*x^(5/6))/5 + (6*x^(7/6))/7`

3.148. $\int \left(1 + \frac{1}{\sqrt{x}} + \frac{1}{\sqrt[3]{x}}\right) (1 + \sqrt[3]{x} + \sqrt{x}) dx$

3.149 $\int \cos(x) \cos(\sin(x)) \sin(\sin(x)) dx$

3.149.1 Optimal result	811
3.149.2 Mathematica [A] (verified)	811
3.149.3 Rubi [A] (verified)	812
3.149.4 Maple [A] (verified)	813
3.149.5 Fricas [B] (verification not implemented)	814
3.149.6 Sympy [A] (verification not implemented)	814
3.149.7 Maxima [A] (verification not implemented)	814
3.149.8 Giac [A] (verification not implemented)	815
3.149.9 Mupad [B] (verification not implemented)	815

3.149.1 Optimal result

Integrand size = 9, antiderivative size = 9

$$\int \cos(x) \cos(\sin(x)) \sin(\sin(x)) dx = \frac{1}{2} \sin^2(\sin(x))$$

output `1/2*sin(sin(x))^2`

3.149.2 Mathematica [A] (verified)

Time = 2.05 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \cos(x) \cos(\sin(x)) \sin(\sin(x)) dx = -\frac{1}{2} \cos^2(\sin(x))$$

input `Integrate[Cos[x]*Cos[Sin[x]]*Sin[Sin[x]],x]`

output `-1/2*Cos[Sin[x]]^2`

3.149.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {4834, 3042, 3044, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(\sin(x)) \cos(x) \cos(\sin(x)) dx \\
 & \quad \downarrow \text{4834} \\
 & \int \sin(\sin(x)) \cos(\sin(x)) d \sin(x) \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(\sin(x)) \cos(\sin(x)) d \sin(x) \\
 & \quad \downarrow \text{3044} \\
 & \int \sin(\sin(x)) d \sin(\sin(x)) \\
 & \quad \downarrow \text{15} \\
 & \frac{1}{2} \sin^2(\sin(x))
 \end{aligned}$$

input `Int[Cos[x]*Cos[Sin[x]]*Sin[Sin[x]],x]`

output `Sin[Sin[x]]^2/2`

3.149.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3044 Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)]^(m_.), x_
Symbol] :> Simp[1/(a*f) Subst[Int[x^m*(1 - x^2/a^2)^(n - 1)/2], x], x, a
*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(I
ntegerQ[(m - 1)/2] && LtQ[0, m, n])
```

```
rule 4834 Int[(u_)*(F_)[(c_.)*((a_.) + (b_.)*(x_))], x_Symbol] :> With[{d = FreeFacto
rs[Sin[c*(a + b*x)], x]}, Simp[d/(b*c) Subst[Int[SubstFor[1, Sin[c*(a + b
*x)]/d, u, x], x], x, Sin[c*(a + b*x)]/d], x] /; FunctionOfQ[Sin[c*(a + b*x
)]/d, u, x, True]] /; FreeQ[{a, b, c}, x] && (EqQ[F, Cos] || EqQ[F, cos])
```

3.149.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

method	result	size
derivativedivides	$\frac{\sin(\sin(x))^2}{2}$	8
default	$\frac{\sin(\sin(x))^2}{2}$	8
risch	$-\frac{\cos(2 \sin(x))}{4}$	8
parallelrisch	$-\frac{3}{4} - \frac{\cos(2 \sin(x))}{4}$	10
norman	$\frac{2 \tan\left(\frac{\tan\left(\frac{x}{2}\right)}{1+\tan\left(\frac{x}{2}\right)^2}\right)^2 + 2 \tan\left(\frac{x}{2}\right)^2 \tan\left(\frac{\tan\left(\frac{x}{2}\right)}{1+\tan\left(\frac{x}{2}\right)^2}\right)^2}{\left(1+\tan\left(\frac{\tan\left(\frac{x}{2}\right)}{1+\tan\left(\frac{x}{2}\right)^2}\right)^2\right)^2 (1+\tan\left(\frac{x}{2}\right)^2)}$	81

```
input int(sin(sin(x))*cos(sin(x))*cos(x), x, method=_RETURNVERBOSE)
```

```
output 1/2*sin(sin(x))^2
```

3.149.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. $2(7) = 14$.

Time = 0.24 (sec) , antiderivative size = 21, normalized size of antiderivative = 2.33

$$\int \cos(x) \cos(\sin(x)) \sin(\sin(x)) dx = -\frac{1}{2} \cos\left(\frac{2 \tan\left(\frac{1}{2}x\right)}{\tan\left(\frac{1}{2}x\right)^2 + 1}\right)^2$$

input `integrate(sin(sin(x))*cos(sin(x))*cos(x),x, algorithm="fricas")`

output `-1/2*cos(2*tan(1/2*x)/(tan(1/2*x)^2 + 1))^2`

3.149.6 Sympy [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int \cos(x) \cos(\sin(x)) \sin(\sin(x)) dx = -\frac{\cos^2(\sin(x))}{2}$$

input `integrate(sin(sin(x))*cos(sin(x))*cos(x),x)`

output `-cos(sin(x))**2/2`

3.149.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \cos(x) \cos(\sin(x)) \sin(\sin(x)) dx = -\frac{1}{2} \cos(\sin(x))^2$$

input `integrate(sin(sin(x))*cos(sin(x))*cos(x),x, algorithm="maxima")`

output `-1/2*cos(sin(x))^2`

3.149.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \cos(x) \cos(\sin(x)) \sin(\sin(x)) dx = -\frac{1}{2} \cos(\sin(x))^2$$

input `integrate(sin(sin(x))*cos(sin(x))*cos(x),x, algorithm="giac")`

output `-1/2*cos(sin(x))^2`

3.149.9 Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \cos(x) \cos(\sin(x)) \sin(\sin(x)) dx = \frac{\sin(\sin(x))^2}{2}$$

input `int(cos(sin(x))*sin(sin(x))*cos(x),x)`

output `sin(sin(x))^2/2`

3.150 $\int \left(\frac{-\cos(x)+\sin(x)}{x} + \frac{\cos(x)+\sin(x)}{x^2} \right) dx$

3.150.1 Optimal result 816
 3.150.2 Mathematica [A] (verified) 816
 3.150.3 Rubi [A] (verified) 817
 3.150.4 Maple [A] (verified) 817
 3.150.5 Fricas [A] (verification not implemented) 818
 3.150.6 Sympy [A] (verification not implemented) 818
 3.150.7 Maxima [C] (verification not implemented) 818
 3.150.8 Giac [C] (verification not implemented) 819
 3.150.9 Mupad [B] (verification not implemented) 819

3.150.1 Optimal result

Integrand size = 21, antiderivative size = 15

$$\int \left(\frac{-\cos(x) + \sin(x)}{x} + \frac{\cos(x) + \sin(x)}{x^2} \right) dx = -\frac{\cos(x)}{x} - \frac{\sin(x)}{x}$$

output -cos(x)/x-sin(x)/x

3.150.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \left(\frac{-\cos(x) + \sin(x)}{x} + \frac{\cos(x) + \sin(x)}{x^2} \right) dx = -\frac{\cos(x)}{x} - \frac{\sin(x)}{x}$$

input Integrate[(-Cos[x] + Sin[x])/x + (Cos[x] + Sin[x])/x^2,x]

output -(Cos[x]/x) - Sin[x]/x

3.150.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\frac{\sin(x) + \cos(x)}{x^2} + \frac{\sin(x) - \cos(x)}{x} \right) dx$$

↓ 2009

$$-\frac{\sin(x)}{x} - \frac{\cos(x)}{x}$$

input `Int[(-Cos[x] + Sin[x])/x + (Cos[x] + Sin[x])/x^2,x]`

output `-(Cos[x]/x) - Sin[x]/x`

3.150.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.150.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result
parallelrisch	$-\frac{\cos(x) - \sin(x)}{x}$
default	$-\frac{\cos(x)}{x} - \frac{\sin(x)}{x}$
parts	$-\frac{\cos(x)}{x} - \frac{\sin(x)}{x}$
risch	$\frac{i(2i \cos(x) + 2i \sin(x))}{2x}$
norman	$\frac{-1 + \tan(\frac{x}{2})^2 - 2 \tan(\frac{x}{2})}{(1 + \tan(\frac{x}{2})^2)x}$
meijerg	$\frac{\sqrt{\pi} \left(-\frac{4 \cos(x)}{x\sqrt{\pi}} - \frac{4 \operatorname{Si}(x)}{\sqrt{\pi}} \right)}{4} + \frac{\sqrt{\pi} \left(\frac{4\gamma - 4 + 4 \ln(x)}{\sqrt{\pi}} + \frac{4}{\sqrt{\pi}} - \frac{4\gamma}{\sqrt{\pi}} - \frac{4 \ln(2)}{\sqrt{\pi}} - \frac{4 \ln(\frac{x}{2})}{\sqrt{\pi}} - \frac{4 \sin(x)}{\sqrt{\pi} x} + \frac{4 \operatorname{Ci}(x)}{\sqrt{\pi}} \right)}{4} + \operatorname{Si}(x) - \frac{\sqrt{\pi} \left(\frac{2\gamma + 2 \ln(x)}{\sqrt{\pi}} \right)}{4}$

3.150. $\int \left(\frac{-\cos(x) + \sin(x)}{x} + \frac{\cos(x) + \sin(x)}{x^2} \right) dx$

input `int((cos(x)+sin(x))/x^2+(sin(x)-cos(x))/x,x,method=_RETURNVERBOSE)`

output `1/x*(-cos(x)-sin(x))`

3.150.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int \left(\frac{-\cos(x) + \sin(x)}{x} + \frac{\cos(x) + \sin(x)}{x^2} \right) dx = -\frac{\cos(x) + \sin(x)}{x}$$

input `integrate((cos(x)+sin(x))/x^2+(sin(x)-cos(x))/x,x, algorithm="fricas")`

output `-(cos(x) + sin(x))/x`

3.150.6 Sympy [A] (verification not implemented)

Time = 1.29 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.27

$$\int \left(\frac{-\cos(x) + \sin(x)}{x} + \frac{\cos(x) + \sin(x)}{x^2} \right) dx = -\log(x) + \frac{\log(x^2)}{2} - \frac{\sin(x)}{x} - \frac{\cos(x)}{x}$$

input `integrate((cos(x)+sin(x))/x**2+(sin(x)-cos(x))/x,x)`

output `-log(x) + log(x**2)/2 - sin(x)/x - cos(x)/x`

3.150.7 Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 0.24 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.80

$$\int \left(\frac{-\cos(x) + \sin(x)}{x} + \frac{\cos(x) + \sin(x)}{x^2} \right) dx = -\left(\frac{1}{2}i + \frac{1}{2}\right) \text{Ei}(ix) + \left(\frac{1}{2}i - \frac{1}{2}\right) \text{Ei}(-ix) - \left(\frac{1}{2}i - \frac{1}{2}\right) \Gamma(-1, ix) + \left(\frac{1}{2}i + \frac{1}{2}\right) \Gamma(-1, -ix)$$

3.150. $\int \left(\frac{-\cos(x)+\sin(x)}{x} + \frac{\cos(x)+\sin(x)}{x^2} \right) dx$

input `integrate((cos(x)+sin(x))/x^2+(sin(x)-cos(x))/x,x, algorithm="maxima")`

output `-(1/2*I + 1/2)*Ei(I*x) + (1/2*I - 1/2)*Ei(-I*x) - (1/2*I - 1/2)*gamma(-1, I*x) + (1/2*I + 1/2)*gamma(-1, -I*x)`

3.150.8 Giac [C] (verification not implemented)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.93

$$\int \left(\frac{-\cos(x) + \sin(x)}{x} + \frac{\cos(x) + \sin(x)}{x^2} \right) dx$$

$$= \frac{x \operatorname{Ci}(x) - x \operatorname{Si}(x) - \cos(x) - \sin(x)}{x} - \operatorname{Ci}(x) + \operatorname{Si}(x)$$

input `integrate((cos(x)+sin(x))/x^2+(sin(x)-cos(x))/x,x, algorithm="giac")`

output `(x*cos_integral(x) - x*sin_integral(x) - cos(x) - sin(x))/x - cos_integral(x) + sin_integral(x)`

3.150.9 Mupad [B] (verification not implemented)

Time = 16.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \left(\frac{-\cos(x) + \sin(x)}{x} + \frac{\cos(x) + \sin(x)}{x^2} \right) dx = -\frac{\sqrt{2} \sin\left(x + \frac{\pi}{4}\right)}{x}$$

input `int((cos(x) + sin(x))/x^2 - (cos(x) - sin(x))/x,x)`

output `-(2^(1/2)*sin(x + pi/4))/x`

3.151 $\int x^3 \sqrt{1+x^2} dx$

3.151.1 Optimal result	820
3.151.2 Mathematica [A] (verified)	820
3.151.3 Rubi [A] (verified)	821
3.151.4 Maple [A] (verified)	822
3.151.5 Fricas [A] (verification not implemented)	822
3.151.6 Sympy [A] (verification not implemented)	823
3.151.7 Maxima [A] (verification not implemented)	823
3.151.8 Giac [A] (verification not implemented)	823
3.151.9 Mupad [B] (verification not implemented)	824

3.151.1 Optimal result

Integrand size = 13, antiderivative size = 27

$$\int x^3 \sqrt{1+x^2} dx = -\frac{1}{3}(1+x^2)^{3/2} + \frac{1}{5}(1+x^2)^{5/2}$$

output `-1/3*(x^2+1)^(3/2)+1/5*(x^2+1)^(5/2)`

3.151.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

$$\int x^3 \sqrt{1+x^2} dx = \frac{1}{15}(1+x^2)^{3/2}(-2+3x^2)$$

input `Integrate[x^3*Sqrt[1+x^2],x]`

output `((1+x^2)^(3/2)*(-2+3*x^2))/15`

3.151.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.15, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^3 \sqrt{x^2 + 1} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int x^2 \sqrt{x^2 + 1} dx^2 \\ & \quad \downarrow \text{53} \\ & \frac{1}{2} \int \left((x^2 + 1)^{3/2} - \sqrt{x^2 + 1} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \left(\frac{2}{5} (x^2 + 1)^{5/2} - \frac{2}{3} (x^2 + 1)^{3/2} \right) \end{aligned}$$

input `Int[x^3*Sqrt[1 + x^2],x]`

output `((-2*(1 + x^2)^(3/2))/3 + (2*(1 + x^2)^(5/2))/5)/2`

3.151.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.151.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.63

method	result	size
gospers	$\frac{(x^2+1)^{\frac{3}{2}}(3x^2-2)}{15}$	17
pseudoelliptic	$\frac{(x^2+1)^{\frac{3}{2}}(3x^2-2)}{15}$	17
risch	$\frac{(3x^4+x^2-2)\sqrt{x^2+1}}{15}$	20
trager	$\left(\frac{1}{5}x^4 + \frac{1}{15}x^2 - \frac{2}{15}\right)\sqrt{x^2+1}$	21
default	$\frac{x^2(x^2+1)^{\frac{3}{2}}}{5} - \frac{2(x^2+1)^{\frac{3}{2}}}{15}$	23
meijerg	$-\frac{8\sqrt{\pi}}{15} + \frac{4\sqrt{\pi}(x^2+1)^{\frac{3}{2}}(-3x^2+2)}{15 \cdot 4\sqrt{\pi}}$	31

input `int(x^3*(x^2+1)^(1/2),x,method=_RETURNVERBOSE)`output `1/15*(x^2+1)^(3/2)*(3*x^2-2)`**3.151.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int x^3\sqrt{1+x^2} dx = \frac{1}{15}(3x^4+x^2-2)\sqrt{x^2+1}$$

input `integrate(x^3*(x^2+1)^(1/2),x, algorithm="fricas")`output `1/15*(3*x^4 + x^2 - 2)*sqrt(x^2 + 1)`

3.151.6 Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.37

$$\int x^3 \sqrt{1+x^2} dx = \frac{x^4 \sqrt{x^2+1}}{5} + \frac{x^2 \sqrt{x^2+1}}{15} - \frac{2\sqrt{x^2+1}}{15}$$

input `integrate(x**3*(x**2+1)**(1/2),x)`output `x**4*sqrt(x**2 + 1)/5 + x**2*sqrt(x**2 + 1)/15 - 2*sqrt(x**2 + 1)/15`**3.151.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int x^3 \sqrt{1+x^2} dx = \frac{1}{5} (x^2+1)^{\frac{3}{2}} x^2 - \frac{2}{15} (x^2+1)^{\frac{3}{2}}$$

input `integrate(x^3*(x^2+1)^(1/2),x, algorithm="maxima")`output `1/5*(x^2 + 1)^(3/2)*x^2 - 2/15*(x^2 + 1)^(3/2)`**3.151.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int x^3 \sqrt{1+x^2} dx = \frac{1}{5} (x^2+1)^{\frac{5}{2}} - \frac{1}{3} (x^2+1)^{\frac{3}{2}}$$

input `integrate(x^3*(x^2+1)^(1/2),x, algorithm="giac")`output `1/5*(x^2 + 1)^(5/2) - 1/3*(x^2 + 1)^(3/2)`

3.151.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

$$\int x^3 \sqrt{1+x^2} dx = \sqrt{x^2+1} \left(\frac{x^4}{5} + \frac{x^2}{15} - \frac{2}{15} \right)$$

input `int(x^3*(x^2 + 1)^(1/2),x)`

output `(x^2 + 1)^(1/2)*(x^2/15 + x^4/5 - 2/15)`

3.152 $\int \frac{x}{1+x^2+x^4} dx$

3.152.1 Optimal result	825
3.152.2 Mathematica [A] (verified)	825
3.152.3 Rubi [A] (verified)	826
3.152.4 Maple [A] (verified)	827
3.152.5 Fricas [A] (verification not implemented)	827
3.152.6 Sympy [A] (verification not implemented)	828
3.152.7 Maxima [A] (verification not implemented)	828
3.152.8 Giac [A] (verification not implemented)	828
3.152.9 Mupad [B] (verification not implemented)	829

3.152.1 Optimal result

Integrand size = 12, antiderivative size = 20

$$\int \frac{x}{1+x^2+x^4} dx = \frac{\arctan\left(\frac{1+2x^2}{\sqrt{3}}\right)}{\sqrt{3}}$$

output `1/3*arctan(1/3*(2*x^2+1)*3^(1/2))*3^(1/2)`

3.152.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{x}{1+x^2+x^4} dx = \frac{\arctan\left(\frac{1+2x^2}{\sqrt{3}}\right)}{\sqrt{3}}$$

input `Integrate[x/(1 + x^2 + x^4),x]`

output `ArcTan[(1 + 2*x^2)/Sqrt[3]]/Sqrt[3]`

3.152.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {1432, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x}{x^4 + x^2 + 1} dx \\ & \quad \downarrow \text{1432} \\ & \frac{1}{2} \int \frac{1}{x^4 + x^2 + 1} dx^2 \\ & \quad \downarrow \text{1083} \\ & - \int \frac{1}{-x^4 - 3} d(2x^2 + 1) \\ & \quad \downarrow \text{217} \\ & \frac{\arctan\left(\frac{2x^2+1}{\sqrt{3}}\right)}{\sqrt{3}} \end{aligned}$$

input `Int[x/(1 + x^2 + x^4),x]`

output `ArcTan[(1 + 2*x^2)/Sqrt[3]]/Sqrt[3]`

3.152.3.1 Defintions of rubi rules used

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])`

rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] :> Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 1432 `Int[(x_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := Simp[1/2
Subst[Int[(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, p}, x]`

3.152.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

method	result	size
default	$\frac{\arctan\left(\frac{(2x^2+1)\sqrt{3}}{3}\right)\sqrt{3}}{3}$	19
risch	$\frac{\arctan\left(\frac{(2x^2+1)\sqrt{3}}{3}\right)\sqrt{3}}{3}$	19

input `int(x/(x^4+x^2+1),x,method=_RETURNVERBOSE)`

output `1/3*arctan(1/3*(2*x^2+1)*3^(1/2))*3^(1/2)`

3.152.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{x}{1+x^2+x^4} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x^2+1)\right)$$

input `integrate(x/(x^4+x^2+1),x, algorithm="fricas")`

output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x^2 + 1))`

3.152.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.30

$$\int \frac{x}{1+x^2+x^4} dx = \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x^2}{3} + \frac{\sqrt{3}}{3}\right)}{3}$$

input `integrate(x/(x**4+x**2+1),x)`output `sqrt(3)*atan(2*sqrt(3)*x**2/3 + sqrt(3)/3)/3`**3.152.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{x}{1+x^2+x^4} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x^2+1)\right)$$

input `integrate(x/(x^4+x^2+1),x, algorithm="maxima")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x^2 + 1))`**3.152.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{x}{1+x^2+x^4} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x^2+1)\right)$$

input `integrate(x/(x^4+x^2+1),x, algorithm="giac")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x^2 + 1))`

3.152.9 Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{x}{1+x^2+x^4} dx = \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x^2}{3} + \frac{\sqrt{3}}{3}\right)}{3}$$

input `int(x/(x^2 + x^4 + 1),x)`

output `(3^(1/2)*atan(3^(1/2)/3 + (2*3^(1/2)*x^2)/3))/3`

3.153 $\int e^{e^{2016x}+6048x} dx$

3.153.1 Optimal result	830
3.153.2 Mathematica [A] (verified)	830
3.153.3 Rubi [A] (verified)	831
3.153.4 Maple [A] (verified)	832
3.153.5 Fricas [A] (verification not implemented)	833
3.153.6 Sympy [A] (verification not implemented)	833
3.153.7 Maxima [A] (verification not implemented)	833
3.153.8 Giac [A] (verification not implemented)	834
3.153.9 Mupad [B] (verification not implemented)	834

3.153.1 Optimal result

Integrand size = 11, antiderivative size = 42

$$\int e^{e^{2016x}+6048x} dx = \frac{e^{e^{2016x}}}{1008} - \frac{e^{e^{2016x}+2016x}}{1008} + \frac{e^{e^{2016x}+4032x}}{2016}$$

output `1/1008*exp(exp(1)^(2016*x))-1/1008*exp(exp(1)^(2016*x)+2016*x)+1/2016*exp(exp(1)^(2016*x)+4032*x)`

3.153.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.60

$$\int e^{e^{2016x}+6048x} dx = \frac{e^{e^{2016x}}(2 - 2e^{2016x} + e^{4032x})}{2016}$$

input `Integrate[E^(E^(2016*x) + 6048*x),x]`

output `(E^E^(2016*x)*(2 - 2*E^(2016*x) + E^(4032*x)))/2016`

3.153.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.93, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {2720, 2607, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int e^{6048x+e^{2016x}} dx \\
 \downarrow \text{2720} \\
 \frac{\int e^{4032x+e^{2016x}} de^{2016x}}{2016} \\
 \downarrow \text{2607} \\
 \frac{e^{4032x+e^{2016x}} - 2 \int e^{2016x+e^{2016x}} de^{2016x}}{2016} \\
 \downarrow \text{2607} \\
 \frac{e^{4032x+e^{2016x}} - 2 \left(e^{2016x+e^{2016x}} - \int e^{e^{2016x}} de^{2016x} \right)}{2016} \\
 \downarrow \text{2624} \\
 \frac{e^{4032x+e^{2016x}} - 2 \left(e^{2016x+e^{2016x}} - e^{e^{2016x}} \right)}{2016}
 \end{array}$$

input `Int [E^(E^(2016*x) + 6048*x), x]`

output `(E^(E^(2016*x) + 4032*x) - 2*(-E^E^(2016*x) + E^(E^(2016*x) + 2016*x)))/2016`

3.153.3.1 Defintions of rubi rules used

```
rule 2607 Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_)))^(n_.))*((c_.) + (d_.)*(x_))^(m
_.), x_Symbol] := Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])),
x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^
n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*
m] && !TrueQ[$UseGamma]
```

```
rule 2624 Int[((F_)^(v_))^(n_), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;
FreeQ[{F, n}, x] && LinearQ[v, x]
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.153.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.48

method	result	size
risch	$\frac{(e^{4032x} - 2e^{2016x} + 2)e^{2016x}}{2016}$	20

```
input int(exp(exp(2016*x)+6048*x),x,method=_RETURNVERBOSE)
```

```
output 1/2016*(exp(4032*x)-2*exp(2016*x)+2)*exp(exp(2016*x))
```

3.153.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.45

$$\int e^{e^{2016x}+6048x} dx = \frac{1}{2016} (e^{(4032x)} - 2e^{(2016x)} + 2)e^{(e^{2016x})}$$

input `integrate(exp(exp(2016*x)+6048*x),x, algorithm="fricas")`output `1/2016*(e^(4032*x) - 2*e^(2016*x) + 2)*e^(e^(2016*x))`**3.153.6 Sympy [A] (verification not implemented)**

Time = 0.41 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.81

$$\int e^{e^{2016x}+6048x} dx = \frac{e^{4032x}e^{e^{2016x}}}{2016} - \frac{e^{2016x}e^{e^{2016x}}}{1008} + \frac{e^{e^{2016x}}}{1008}$$

input `integrate(exp(exp(2016*x)+6048*x),x)`output `exp(4032*x)*exp(exp(2016*x))/2016 - exp(2016*x)*exp(exp(2016*x))/1008 + exp(exp(2016*x))/1008`**3.153.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.45

$$\int e^{e^{2016x}+6048x} dx = \frac{1}{2016} (e^{(4032x)} - 2e^{(2016x)} + 2)e^{(e^{2016x})}$$

input `integrate(exp(exp(2016*x)+6048*x),x, algorithm="maxima")`output `1/2016*(e^(4032*x) - 2*e^(2016*x) + 2)*e^(e^(2016*x))`

3.153.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.90

$$\int e^{e^{2016x}+6048x} dx = \frac{1}{2016} \left(e^{(10080x+e^{(2016x)})} - 2e^{(8064x+e^{(2016x)})} + 2e^{(6048x+e^{(2016x)})} \right) e^{(-6048x)}$$

input `integrate(exp(exp(2016*x)+6048*x),x, algorithm="giac")`output `1/2016*(e^(10080*x + e^(2016*x)) - 2*e^(8064*x + e^(2016*x)) + 2*e^(6048*x + e^(2016*x)))*e^(-6048*x)`**3.153.9 Mupad [B] (verification not implemented)**

Time = 15.59 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.71

$$\int e^{e^{2016x}+6048x} dx = \frac{e^{e^{2016x}}}{1008} - \frac{e^{2016x} e^{e^{2016x}}}{1008} + \frac{e^{4032x} e^{e^{2016x}}}{2016}$$

input `int(exp(6048*x + exp(2016*x)),x)`output `exp(exp(2016*x))/1008 - (exp(2016*x)*exp(exp(2016*x)))/1008 + (exp(4032*x)*exp(exp(2016*x)))/2016`

3.154 $\int (1 - \cot(x)) dx$

3.154.1 Optimal result	835
3.154.2 Mathematica [A] (verified)	835
3.154.3 Rubi [A] (verified)	836
3.154.4 Maple [A] (verified)	836
3.154.5 Fricas [A] (verification not implemented)	837
3.154.6 Sympy [A] (verification not implemented)	837
3.154.7 Maxima [A] (verification not implemented)	837
3.154.8 Giac [A] (verification not implemented)	838
3.154.9 Mupad [B] (verification not implemented)	838

3.154.1 Optimal result

Integrand size = 6, antiderivative size = 7

$$\int (1 - \cot(x)) dx = x - \log(\sin(x))$$

output `x-ln(sin(x))`

3.154.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.71

$$\int (1 - \cot(x)) dx = x - \log(\cos(x)) - \log(\tan(x))$$

input `Integrate[1 - Cot[x],x]`

output `x - Log[Cos[x]] - Log[Tan[x]]`

3.154.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (1 - \cot(x)) dx$$

$$\downarrow \text{2009}$$

$$x - \log(\sin(x))$$

input `Int[1 - Cot[x], x]`

output `x - Log[Sin[x]]`

3.154.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.154.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

method	result	size
default	$x - \ln(\sin(x))$	8
parts	$x - \ln(\sin(x))$	8
risch	$x + ix - \ln(e^{2ix} - 1)$	17
norman	$x - \ln\left(\tan\left(\frac{x}{2}\right)\right) + \ln\left(1 + \tan\left(\frac{x}{2}\right)^2\right)$	19
parallelrisc	$-\ln\left(-\frac{\cot(x)}{2} + \frac{\csc(x)}{2}\right) + \ln\left(\frac{1}{1+\cos(x)}\right) + x$	22

input `int(1-cos(x)/sin(x), x, method=_RETURNVERBOSE)`

output `x-ln(sin(x))`

3.154.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.29

$$\int (1 - \cot(x)) dx = x - \log\left(\frac{1}{2} \sin(x)\right)$$

input `integrate(1-cos(x)/sin(x),x, algorithm="fricas")`output `x - log(1/2*sin(x))`**3.154.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71

$$\int (1 - \cot(x)) dx = x - \log(\sin(x))$$

input `integrate(1-cos(x)/sin(x),x)`output `x - log(sin(x))`**3.154.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int (1 - \cot(x)) dx = x - \log(\sin(x))$$

input `integrate(1-cos(x)/sin(x),x, algorithm="maxima")`output `x - log(sin(x))`

3.154.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

$$\int (1 - \cot(x)) dx = x - \log(|\sin(x)|)$$

input `integrate(1-cos(x)/sin(x),x, algorithm="giac")`

output `x - log(abs(sin(x)))`

3.154.9 Mupad [B] (verification not implemented)

Time = 15.06 (sec) , antiderivative size = 26, normalized size of antiderivative = 3.71

$$\int (1 - \cot(x)) dx = -\ln(\tan(x)) + \ln(\tan(x) - i) \left(\frac{1}{2} - \frac{1}{2}i\right) + \ln(\tan(x) + i) \left(\frac{1}{2} + \frac{1}{2}i\right)$$

input `int(1 - cos(x)/sin(x),x)`

output `log(tan(x) - 1i)*(1/2 - 1i/2) - log(tan(x)) + log(tan(x) + 1i)*(1/2 + 1i/2)`
`)`

3.155 $\int \frac{1}{1-x+x^2-x^3} dx$

3.155.1 Optimal result	839
3.155.2 Mathematica [A] (verified)	839
3.155.3 Rubi [A] (verified)	840
3.155.4 Maple [A] (verified)	841
3.155.5 Fricas [A] (verification not implemented)	841
3.155.6 Sympy [A] (verification not implemented)	841
3.155.7 Maxima [A] (verification not implemented)	842
3.155.8 Giac [A] (verification not implemented)	842
3.155.9 Mupad [B] (verification not implemented)	842

3.155.1 Optimal result

Integrand size = 15, antiderivative size = 27

$$\int \frac{1}{1-x+x^2-x^3} dx = \frac{\arctan(x)}{2} - \frac{1}{2} \log(1-x) + \frac{1}{4} \log(1+x^2)$$

output `1/2*arctan(x)-1/2*ln(1-x)+1/4*ln(x^2+1)`

3.155.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{1}{1-x+x^2-x^3} dx = \frac{\arctan(x)}{2} - \frac{1}{2} \log(1-x) + \frac{1}{4} \log(1+x^2)$$

input `Integrate[(1 - x + x^2 - x^3)^(-1), x]`

output `ArcTan[x]/2 - Log[1 - x]/2 + Log[1 + x^2]/4`

3.155.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{-x^3 + x^2 - x + 1} dx$$

↓ 2462

$$\int \left(\frac{x+1}{2(x^2+1)} - \frac{1}{2(x-1)} \right) dx$$

↓ 2009

$$\frac{\arctan(x)}{2} + \frac{1}{4} \log(x^2+1) - \frac{1}{2} \log(1-x)$$

input `Int[(1 - x + x^2 - x^3)^(-1),x]`

output `ArcTan[x]/2 - Log[1 - x]/2 + Log[1 + x^2]/4`

3.155.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2462 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegrand[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

3.155.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result	size
default	$-\frac{\ln(-1+x)}{2} + \frac{\ln(x^2+1)}{4} + \frac{\arctan(x)}{2}$	20
risch	$-\frac{\ln(-1+x)}{2} + \frac{\ln(x^2+1)}{4} + \frac{\arctan(x)}{2}$	20
parallelrisc	$-\frac{\ln(-1+x)}{2} + \frac{\ln(x-i)}{4} - \frac{i \ln(x-i)}{4} + \frac{\ln(i+x)}{4} + \frac{i \ln(i+x)}{4}$	38

input `int(1/(-x^3+x^2-x+1),x,method=_RETURNVERBOSE)`output `-1/2*ln(-1+x)+1/4*ln(x^2+1)+1/2*arctan(x)`**3.155.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int \frac{1}{1-x+x^2-x^3} dx = \frac{1}{2} \arctan(x) + \frac{1}{4} \log(x^2+1) - \frac{1}{2} \log(x-1)$$

input `integrate(1/(-x^3+x^2-x+1),x, algorithm="fricas")`output `1/2*arctan(x) + 1/4*log(x^2 + 1) - 1/2*log(x - 1)`**3.155.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int \frac{1}{1-x+x^2-x^3} dx = -\frac{\log(x-1)}{2} + \frac{\log(x^2+1)}{4} + \frac{\operatorname{atan}(x)}{2}$$

input `integrate(1/(-x**3+x**2-x+1),x)`output `-log(x - 1)/2 + log(x**2 + 1)/4 + atan(x)/2`

3.155.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int \frac{1}{1-x+x^2-x^3} dx = \frac{1}{2} \arctan(x) + \frac{1}{4} \log(x^2+1) - \frac{1}{2} \log(x-1)$$

input `integrate(1/(-x^3+x^2-x+1),x, algorithm="maxima")`output `1/2*arctan(x) + 1/4*log(x^2 + 1) - 1/2*log(x - 1)`**3.155.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

$$\int \frac{1}{1-x+x^2-x^3} dx = \frac{1}{2} \arctan(x) + \frac{1}{4} \log(x^2+1) - \frac{1}{2} \log(|x-1|)$$

input `integrate(1/(-x^3+x^2-x+1),x, algorithm="giac")`output `1/2*arctan(x) + 1/4*log(x^2 + 1) - 1/2*log(abs(x - 1))`**3.155.9 Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.93

$$\int \frac{1}{1-x+x^2-x^3} dx = -\frac{\ln(x-1)}{2} + \ln(x-i) \left(\frac{1}{4} - \frac{1}{4}i\right) + \ln(x+1i) \left(\frac{1}{4} + \frac{1}{4}i\right)$$

input `int(-1/(x - x^2 + x^3 - 1),x)`output `log(x - 1i)*(1/4 - 1i/4) - log(x - 1)/2 + log(x + 1i)*(1/4 + 1i/4)`

3.156 $\int \frac{1}{2+\cosh(x)} dx$

3.156.1 Optimal result	843
3.156.2 Mathematica [A] (verified)	843
3.156.3 Rubi [A] (verified)	844
3.156.4 Maple [A] (verified)	845
3.156.5 Fricas [B] (verification not implemented)	845
3.156.6 Sympy [A] (verification not implemented)	845
3.156.7 Maxima [B] (verification not implemented)	846
3.156.8 Giac [A] (verification not implemented)	846
3.156.9 Mupad [B] (verification not implemented)	846

3.156.1 Optimal result

Integrand size = 6, antiderivative size = 20

$$\int \frac{1}{2 + \cosh(x)} dx = \frac{2 \coth^{-1}(\sqrt{3} \coth(\frac{x}{2}))}{\sqrt{3}}$$

output `2/3*arccoth(3^(1/2)*coth(1/2*x))*3^(1/2)`

3.156.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{2 + \cosh(x)} dx = \frac{2 \operatorname{arctanh}\left(\frac{\tanh(\frac{x}{2})}{\sqrt{3}}\right)}{\sqrt{3}}$$

input `Integrate[(2 + Cosh[x])^(-1), x]`

output `(2*ArcTanh[Tanh[x/2]/Sqrt[3]])/Sqrt[3]`

3.156.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.50, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\cosh(x) + 2} dx$$

↓ 3042

$$\int \frac{1}{2 + \sin\left(\frac{\pi}{2} + ix\right)} dx$$

↓ 3136

$$\frac{x}{\sqrt{3}} - \frac{2\operatorname{arctanh}\left(\frac{\sinh(x)}{\cosh(x) + \sqrt{3} + 2}\right)}{\sqrt{3}}$$

input `Int[(2 + Cosh[x])^(-1),x]`

output `x/Sqrt[3] - (2*ArcTanh[Sinh[x]/(2 + Sqrt[3] + Cosh[x])])/Sqrt[3]`

3.156.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

3.156.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

method	result	size
default	$\frac{2\sqrt{3} \operatorname{arctanh}\left(\frac{\tanh\left(\frac{x}{2}\right)\sqrt{3}}{3}\right)}{3}$	16
risch	$\frac{\sqrt{3} \ln(e^x + 2 - \sqrt{3})}{3} - \frac{\sqrt{3} \ln(e^x + 2 + \sqrt{3})}{3}$	30

input `int(1/(2+cosh(x)),x,method=_RETURNVERBOSE)`

output `2/3*3^(1/2)*arctanh(1/3*tanh(1/2*x)*3^(1/2))`

3.156.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. $2(14) = 28$.

Time = 0.24 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.95

$$\int \frac{1}{2 + \cosh(x)} dx = \frac{1}{3} \sqrt{3} \log \left(-\frac{2(\sqrt{3} - 2) \cosh(x) - (2\sqrt{3} - 3) \sinh(x) + \sqrt{3} - 2}{\cosh(x) + 2} \right)$$

input `integrate(1/(2+cosh(x)),x, algorithm="fricas")`

output `1/3*sqrt(3)*log(-(2*(sqrt(3) - 2)*cosh(x) - (2*sqrt(3) - 3)*sinh(x) + sqrt(3) - 2)/(cosh(x) + 2))`

3.156.6 Sympy [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.80

$$\int \frac{1}{2 + \cosh(x)} dx = -\frac{\sqrt{3} \log\left(\tanh\left(\frac{x}{2}\right) - \sqrt{3}\right)}{3} + \frac{\sqrt{3} \log\left(\tanh\left(\frac{x}{2}\right) + \sqrt{3}\right)}{3}$$

input `integrate(1/(2+cosh(x)),x)`

output `-sqrt(3)*log(tanh(x/2) - sqrt(3))/3 + sqrt(3)*log(tanh(x/2) + sqrt(3))/3`

3.156.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 30 vs. $2(14) = 28$.

Time = 0.28 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.50

$$\int \frac{1}{2 + \cosh(x)} dx = -\frac{1}{3} \sqrt{3} \log \left(-\frac{\sqrt{3} - e^{(-x)} - 2}{\sqrt{3} + e^{(-x)} + 2} \right)$$

input `integrate(1/(2+cosh(x)),x, algorithm="maxima")`

output `-1/3*sqrt(3)*log(-(sqrt(3) - e^(-x) - 2)/(sqrt(3) + e^(-x) + 2))`

3.156.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.30

$$\int \frac{1}{2 + \cosh(x)} dx = \frac{1}{3} \sqrt{3} \log \left(-\frac{\sqrt{3} - e^x - 2}{\sqrt{3} + e^x + 2} \right)$$

input `integrate(1/(2+cosh(x)),x, algorithm="giac")`

output `1/3*sqrt(3)*log(-(sqrt(3) - e^x - 2)/(sqrt(3) + e^x + 2))`

3.156.9 Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.10

$$\int \frac{1}{2 + \cosh(x)} dx = \frac{\sqrt{3} \left(\ln \left(-2e^x - \frac{\sqrt{3}(4e^x+2)}{3} \right) - \ln \left(\frac{\sqrt{3}(4e^x+2)}{3} - 2e^x \right) \right)}{3}$$

input `int(1/(cosh(x) + 2),x)`

output `(3^(1/2)*(log(- 2*exp(x) - (3^(1/2)*(4*exp(x) + 2))/3) - log((3^(1/2)*(4*exp(x) + 2))/3 - 2*exp(x))))/3`

3.157 $\int \frac{x^2}{\sqrt{2+x^3}} dx$

3.157.1 Optimal result	847
3.157.2 Mathematica [A] (verified)	847
3.157.3 Rubi [A] (verified)	848
3.157.4 Maple [A] (verified)	849
3.157.5 Fricas [A] (verification not implemented)	849
3.157.6 Sympy [A] (verification not implemented)	850
3.157.7 Maxima [A] (verification not implemented)	850
3.157.8 Giac [A] (verification not implemented)	850
3.157.9 Mupad [B] (verification not implemented)	851

3.157.1 Optimal result

Integrand size = 13, antiderivative size = 13

$$\int \frac{x^2}{\sqrt{2+x^3}} dx = \frac{2\sqrt{2+x^3}}{3}$$

output `2/3*(x^3+2)^(1/2)`

3.157.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{\sqrt{2+x^3}} dx = \frac{2\sqrt{2+x^3}}{3}$$

input `Integrate[x^2/Sqrt[2 + x^3],x]`

output `(2*Sqrt[2 + x^3])/3`

3.157.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {793}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{\sqrt{x^3 + 2}} dx$$

↓ 793

$$\frac{2\sqrt{x^3 + 2}}{3}$$

input `Int[x^2/Sqrt[2 + x^3],x]`

output `(2*Sqrt[2 + x^3])/3`

3.157.3.1 Defintions of rubi rules used

rule 793 `Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(a + b*x^n)^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]`

3.157.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

method	result	size
gospers	$\frac{2\sqrt{x^3+2}}{3}$	10
derivativedivides	$\frac{2\sqrt{x^3+2}}{3}$	10
default	$\frac{2\sqrt{x^3+2}}{3}$	10
trager	$\frac{2\sqrt{x^3+2}}{3}$	10
risch	$\frac{2\sqrt{x^3+2}}{3}$	10
elliptic	$\frac{2\sqrt{x^3+2}}{3}$	10
pseudoelliptic	$\frac{2\sqrt{x^3+2}}{3}$	10
meijerg	$\frac{\sqrt{2} \left(-2\sqrt{\pi} + 2\sqrt{\pi} \sqrt{1 + \frac{x^3}{2}} \right)}{3\sqrt{\pi}}$	29

input `int(x^2/(x^3+2)^(1/2),x,method=_RETURNVERBOSE)`output `2/3*(x^3+2)^(1/2)`**3.157.5 Fracas [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \frac{x^2}{\sqrt{2+x^3}} dx = \frac{2}{3} \sqrt{x^3+2}$$

input `integrate(x^2/(x^3+2)^(1/2),x, algorithm="fracas")`output `2/3*sqrt(x^3 + 2)`

3.157.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \frac{x^2}{\sqrt{2+x^3}} dx = \frac{2\sqrt{x^3+2}}{3}$$

input `integrate(x**2/(x**3+2)**(1/2),x)`output `2*sqrt(x**3 + 2)/3`**3.157.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \frac{x^2}{\sqrt{2+x^3}} dx = \frac{2}{3} \sqrt{x^3+2}$$

input `integrate(x^2/(x^3+2)^(1/2),x, algorithm="maxima")`output `2/3*sqrt(x^3 + 2)`**3.157.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \frac{x^2}{\sqrt{2+x^3}} dx = \frac{2}{3} \sqrt{x^3+2}$$

input `integrate(x^2/(x^3+2)^(1/2),x, algorithm="giac")`output `2/3*sqrt(x^3 + 2)`

3.157.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \frac{x^2}{\sqrt{2+x^3}} dx = \frac{2\sqrt{x^3+2}}{3}$$

input `int(x^2/(x^3 + 2)^(1/2),x)`

output `(2*(x^3 + 2)^(1/2))/3`

3.158 $\int \frac{\log(x)}{x^2} dx$

3.158.1 Optimal result	852
3.158.2 Mathematica [A] (verified)	852
3.158.3 Rubi [A] (verified)	853
3.158.4 Maple [A] (verified)	853
3.158.5 Fricas [A] (verification not implemented)	854
3.158.6 Sympy [A] (verification not implemented)	854
3.158.7 Maxima [A] (verification not implemented)	854
3.158.8 Giac [A] (verification not implemented)	855
3.158.9 Mupad [B] (verification not implemented)	855

3.158.1 Optimal result

Integrand size = 6, antiderivative size = 13

$$\int \frac{\log(x)}{x^2} dx = -\frac{1}{x} - \frac{\log(x)}{x}$$

output `-1/x-ln(x)/x`

3.158.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\log(x)}{x^2} dx = -\frac{1}{x} - \frac{\log(x)}{x}$$

input `Integrate[Log[x]/x^2,x]`

output `-x^(-1) - Log[x]/x`

3.158.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\log(x)}{x^2} dx$$

↓ 2741

$$-\frac{1}{x} - \frac{\log(x)}{x}$$

input `Int [Log[x]/x^2,x]`

output `-x^(-1) - Log[x]/x`

3.158.3.1 Defintions of rubi rules used

rule 2741 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_)^(m_.), x_Symbol] :=
Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(
m + 1)/(d*(m + 1)^2)), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]`

3.158.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

method	result	size
norman	$\frac{-\ln(x)-1}{x}$	11
parallelrisc	$\frac{-\ln(x)-1}{x}$	11
default	$-\frac{1}{x} - \frac{\ln(x)}{x}$	14
risc	$-\frac{1}{x} - \frac{\ln(x)}{x}$	14
parts	$-\frac{1}{x} - \frac{\ln(x)}{x}$	14

input `int(ln(x)/x^2,x,method=_RETURNVERBOSE)`

output $(-\ln(x)-1)/x$

3.158.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \frac{\log(x)}{x^2} dx = -\frac{\log(x) + 1}{x}$$

input `integrate(log(x)/x^2,x, algorithm="fricas")`

output $-(\log(x) + 1)/x$

3.158.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.62

$$\int \frac{\log(x)}{x^2} dx = -\frac{\log(x)}{x} - \frac{1}{x}$$

input `integrate(ln(x)/x**2,x)`

output $-\log(x)/x - 1/x$

3.158.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\log(x)}{x^2} dx = -\frac{\log(x)}{x} - \frac{1}{x}$$

input `integrate(log(x)/x^2,x, algorithm="maxima")`

output $-\log(x)/x - 1/x$

3.158.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\log(x)}{x^2} dx = -\frac{\log(x)}{x} - \frac{1}{x}$$

input `integrate(log(x)/x^2,x, algorithm="giac")`

output `-log(x)/x - 1/x`

3.158.9 Mupad [B] (verification not implemented)

Time = 13.95 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \frac{\log(x)}{x^2} dx = -\frac{\ln(x) + 1}{x}$$

input `int(log(x)/x^2,x)`

output `-(log(x) + 1)/x`

3.159 $\int \operatorname{sech}(x) dx$

3.159.1 Optimal result	856
3.159.2 Mathematica [A] (verified)	856
3.159.3 Rubi [A] (verified)	857
3.159.4 Maple [A] (verified)	858
3.159.5 Fricas [B] (verification not implemented)	858
3.159.6 Sympy [B] (verification not implemented)	858
3.159.7 Maxima [A] (verification not implemented)	859
3.159.8 Giac [A] (verification not implemented)	859
3.159.9 Mupad [B] (verification not implemented)	859

3.159.1 Optimal result

Integrand size = 2, antiderivative size = 3

$$\int \operatorname{sech}(x) dx = \arctan(\sinh(x))$$

output `arctan(sinh(x))`

3.159.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \operatorname{sech}(x) dx = \arctan(\sinh(x))$$

input `Integrate[Sech[x],x]`

output `ArcTan[Sinh[x]]`

3.159.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \operatorname{sech}(x) dx \\ \downarrow 3042 \\ \int \csc\left(\frac{\pi}{2} + ix\right) dx \\ \downarrow 4257 \\ \arctan(\sinh(x)) \end{array}$$

input `Int [Sech [x] , x]`

output `ArcTan [Sinh [x]]`

3.159.3.1 Defintions of rubi rules used

rule 3042 `Int [u_ , x_Symbol] :> Int [DeactivateTrig [u , x] , x] /; FunctionOfTrigOfLinear Q [u , x]`

rule 4257 `Int [csc [(c_.) + (d_.)*(x_)] , x_Symbol] :> Simp [-ArcTanh [Cos [c + d*x]]/d , x] /; FreeQ [{c , d} , x]`

3.159.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
lookup	$\arctan(\sinh(x))$	4
default	$\arctan(\sinh(x))$	4
risch	$i \ln(e^x + i) - i \ln(e^x - i)$	20
parallelrisch	$-i(\ln(\tanh(\frac{x}{2}) - i) - \ln(\tanh(\frac{x}{2}) + i))$	23

input `int(sech(x), x, method=_RETURNVERBOSE)`

output `arctan(sinh(x))`

3.159.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8 vs. $2(3) = 6$.

Time = 0.23 (sec) , antiderivative size = 8, normalized size of antiderivative = 2.67

$$\int \operatorname{sech}(x) dx = 2 \arctan(\cosh(x) + \sinh(x))$$

input `integrate(sech(x), x, algorithm="fricas")`

output `2*arctan(cosh(x) + sinh(x))`

3.159.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7 vs. $2(3) = 6$.

Time = 0.17 (sec) , antiderivative size = 7, normalized size of antiderivative = 2.33

$$\int \operatorname{sech}(x) dx = 2 \operatorname{atan}\left(\tanh\left(\frac{x}{2}\right)\right)$$

input `integrate(sech(x), x)`

output `2*atan(tanh(x/2))`

3.159.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \operatorname{sech}(x) dx = \arctan(\sinh(x))$$

input `integrate(sech(x),x, algorithm="maxima")`output `arctan(sinh(x))`**3.159.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.67

$$\int \operatorname{sech}(x) dx = 2 \arctan(e^x)$$

input `integrate(sech(x),x, algorithm="giac")`output `2*arctan(e^x)`**3.159.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.67

$$\int \operatorname{sech}(x) dx = 2 \operatorname{atan}(e^x)$$

input `int(1/cosh(x),x)`output `2*atan(exp(x))`

3.160 $\int e^{x^2} x^3 dx$

3.160.1 Optimal result	860
3.160.2 Mathematica [A] (verified)	860
3.160.3 Rubi [A] (verified)	861
3.160.4 Maple [A] (verified)	862
3.160.5 Fricas [A] (verification not implemented)	862
3.160.6 Sympy [A] (verification not implemented)	863
3.160.7 Maxima [A] (verification not implemented)	863
3.160.8 Giac [A] (verification not implemented)	863
3.160.9 Mupad [B] (verification not implemented)	864

3.160.1 Optimal result

Integrand size = 9, antiderivative size = 22

$$\int e^{x^2} x^3 dx = -\frac{e^{x^2}}{2} + \frac{1}{2}e^{x^2} x^2$$

output `-1/2*exp(x^2)+1/2*x^2*exp(x^2)`

3.160.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.64

$$\int e^{x^2} x^3 dx = \frac{1}{2}e^{x^2}(-1 + x^2)$$

input `Integrate[E^x^2*x^3,x]`

output `(E^x^2*(-1 + x^2))/2`

3.160.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {2641, 2638}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^{x^2} x^3 dx \\ & \quad \downarrow \text{2641} \\ & \frac{1}{2} e^{x^2} x^2 - \int e^{x^2} x dx \\ & \quad \downarrow \text{2638} \\ & \frac{1}{2} e^{x^2} x^2 - \frac{e^{x^2}}{2} \end{aligned}$$

input `Int[E^x^2*x^3,x]`

output `-1/2*E^x^2 + (E^x^2*x^2)/2`

3.160.3.1 Defintions of rubi rules used

rule 2638 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[(e + f*x)^n*(F^(a + b*(c + d*x)^n)/(b*f*n*(c + d*x)^n *Log[F])), x] /; FreeQ[{F, a, b, c, d, e, f, n}, x] && EqQ[m, n - 1] && EqQ[d*e - c*f, 0]`

rule 2641 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^(m - n + 1)*(F^(a + b*(c + d*x)^n)/(b*d*n*Log[F])), x] - Simp[(m - n + 1)/(b*n*Log[F]) Int[(c + d*x)^(m - n)*F^(a + b*(c + d*x)^n), x], x] /; FreeQ[{F, a, b, c, d}, x] && IntegerQ[2*((m + 1)/n)] && LtQ[0, (m + 1)/n, 5] && IntegerQ[n] && (LtQ[0, n, m + 1] || LtQ[m, n, 0])`

3.160.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.55

method	result	size
gospers	$\frac{(x^2-1)e^{x^2}}{2}$	12
risch	$\left(\frac{x^2}{2} - \frac{1}{2}\right) e^{x^2}$	13
meijerg	$\frac{1}{2} - \frac{(-2x^2+2)e^{x^2}}{4}$	16
derivativedivides	$-\frac{e^{x^2}}{2} + \frac{x^2 e^{x^2}}{2}$	17
default	$-\frac{e^{x^2}}{2} + \frac{x^2 e^{x^2}}{2}$	17
norman	$-\frac{e^{x^2}}{2} + \frac{x^2 e^{x^2}}{2}$	17
parallelrisch	$-\frac{e^{x^2}}{2} + \frac{x^2 e^{x^2}}{2}$	17
parts	$\frac{\sqrt{\pi} \operatorname{erfi}(x) x^3}{2} - \frac{3\sqrt{\pi} \left(\frac{x^3 \operatorname{erfi}(x)}{3} - \frac{2 \left(-\frac{e^{x^2}}{2} + \frac{x^2 e^{x^2}}{2} \right)}{3\sqrt{\pi}} \right)}{2}$	46

input `int(x^3*exp(x^2),x,method=_RETURNVERBOSE)`output `1/2*(x^2-1)*exp(x^2)`**3.160.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.50

$$\int e^{x^2} x^3 dx = \frac{1}{2} (x^2 - 1) e^{(x^2)}$$

input `integrate(x^3*exp(x^2),x, algorithm="fricas")`output `1/2*(x^2 - 1)*e^(x^2)`

3.160.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.45

$$\int e^{x^2} x^3 dx = \frac{(x^2 - 1) e^{x^2}}{2}$$

input `integrate(x**3*exp(x**2),x)`output `(x**2 - 1)*exp(x**2)/2`**3.160.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.50

$$\int e^{x^2} x^3 dx = \frac{1}{2} (x^2 - 1) e^{(x^2)}$$

input `integrate(x^3*exp(x^2),x, algorithm="maxima")`output `1/2*(x^2 - 1)*e^(x^2)`**3.160.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.50

$$\int e^{x^2} x^3 dx = \frac{1}{2} (x^2 - 1) e^{(x^2)}$$

input `integrate(x^3*exp(x^2),x, algorithm="giac")`output `1/2*(x^2 - 1)*e^(x^2)`

3.160.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.50

$$\int e^{x^2} x^3 dx = \frac{e^{x^2} (x^2 - 1)}{2}$$

input `int(x^3*exp(x^2),x)`

output `(exp(x^2)*(x^2 - 1))/2`

3.161 $\int \frac{1}{x\sqrt{-1+x^2}} dx$

3.161.1 Optimal result	865
3.161.2 Mathematica [A] (verified)	865
3.161.3 Rubi [A] (verified)	866
3.161.4 Maple [A] (verified)	867
3.161.5 Fricas [A] (verification not implemented)	867
3.161.6 Sympy [C] (verification not implemented)	868
3.161.7 Maxima [A] (verification not implemented)	868
3.161.8 Giac [A] (verification not implemented)	868
3.161.9 Mupad [B] (verification not implemented)	869

3.161.1 Optimal result

Integrand size = 13, antiderivative size = 10

$$\int \frac{1}{x\sqrt{-1+x^2}} dx = \arctan\left(\sqrt{-1+x^2}\right)$$

output `arctan((x^2-1)^(1/2))`

3.161.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{-1+x^2}} dx = \arctan\left(\sqrt{-1+x^2}\right)$$

input `Integrate[1/(x*Sqrt[-1 + x^2]),x]`

output `ArcTan[Sqrt[-1 + x^2]]`

3.161.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {243, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x\sqrt{x^2-1}} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int \frac{1}{x^2\sqrt{x^2-1}} dx^2 \\ & \quad \downarrow \text{73} \\ & \int \frac{1}{x^4+1} d\sqrt{x^2-1} \\ & \quad \downarrow \text{216} \\ & \arctan\left(\sqrt{x^2-1}\right) \end{aligned}$$

input `Int[1/(x*sqrt[-1 + x^2]),x]`

output `ArcTan[Sqrt[-1 + x^2]]`

3.161.3.1 Defintions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

3.161.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

method	result	size
pseudoelliptic	$\arctan(\sqrt{x^2 - 1})$	9
default	$-\arctan\left(\frac{1}{\sqrt{x^2 - 1}}\right)$	11
trager	$\text{RootOf}(_Z^2 + 1) \ln\left(\frac{\text{RootOf}(_Z^2 + 1) + \sqrt{x^2 - 1}}{x}\right)$	27
meijerg	$\frac{\sqrt{-\text{signum}(x^2 - 1)} \left((2 \ln(x) - 2 \ln(2) + i\pi) \sqrt{\pi} - 2\sqrt{\pi} \ln\left(\frac{1}{2} + \frac{\sqrt{-x^2 + 1}}{2}\right) \right)}{2\sqrt{\pi} \sqrt{\text{signum}(x^2 - 1)}}$	61

input `int(1/x/(x^2-1)^(1/2),x,method=_RETURNVERBOSE)`

output `arctan((x^2-1)^(1/2))`

3.161.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.40

$$\int \frac{1}{x\sqrt{-1+x^2}} dx = 2 \arctan(-x + \sqrt{x^2 - 1})$$

input `integrate(1/x/(x^2-1)^(1/2),x, algorithm="fracas")`

output `2*arctan(-x + sqrt(x^2 - 1))`

3.161.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.48 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.70

$$\int \frac{1}{x\sqrt{-1+x^2}} dx = \begin{cases} i \operatorname{acosh}\left(\frac{1}{x}\right) & \text{for } \frac{1}{|x^2|} > 1 \\ -\operatorname{asin}\left(\frac{1}{x}\right) & \text{otherwise} \end{cases}$$

input `integrate(1/x/(x**2-1)**(1/2),x)`

output `Piecewise((I*acosh(1/x), 1/Abs(x**2) > 1), (-asin(1/x), True))`

3.161.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \frac{1}{x\sqrt{-1+x^2}} dx = -\operatorname{arcsin}\left(\frac{1}{|x|}\right)$$

input `integrate(1/x/(x^2-1)^(1/2),x, algorithm="maxima")`

output `-arcsin(1/abs(x))`

3.161.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{1}{x\sqrt{-1+x^2}} dx = \operatorname{arctan}\left(\sqrt{x^2-1}\right)$$

input `integrate(1/x/(x^2-1)^(1/2),x, algorithm="giac")`

output `arctan(sqrt(x^2 - 1))`

3.161.9 Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{1}{x\sqrt{-1+x^2}} dx = \operatorname{atan}\left(\sqrt{x^2-1}\right)$$

input `int(1/(x*(x^2 - 1)^(1/2)),x)`

output `atan((x^2 - 1)^(1/2))`

3.162 $\int \frac{1}{x(1+x^2)} dx$

3.162.1 Optimal result	870
3.162.2 Mathematica [A] (verified)	870
3.162.3 Rubi [A] (verified)	871
3.162.4 Maple [A] (verified)	872
3.162.5 Fricas [A] (verification not implemented)	872
3.162.6 Sympy [A] (verification not implemented)	873
3.162.7 Maxima [A] (verification not implemented)	873
3.162.8 Giac [A] (verification not implemented)	873
3.162.9 Mupad [B] (verification not implemented)	874

3.162.1 Optimal result

Integrand size = 11, antiderivative size = 13

$$\int \frac{1}{x(1+x^2)} dx = \log(x) - \frac{1}{2} \log(1+x^2)$$

output `ln(x)-1/2*ln(x^2+1)`

3.162.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1+x^2)} dx = \log(x) - \frac{1}{2} \log(1+x^2)$$

input `Integrate[1/(x*(1 + x^2)),x]`

output `Log[x] - Log[1 + x^2]/2`

3.162.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {243, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x(x^2+1)} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \int \frac{1}{x^2(x^2+1)} dx^2 \\
 & \quad \downarrow \text{47} \\
 & \frac{1}{2} \left(\int \frac{1}{x^2} dx^2 - \int \frac{1}{x^2+1} dx^2 \right) \\
 & \quad \downarrow \text{14} \\
 & \frac{1}{2} \left(\log(x^2) - \int \frac{1}{x^2+1} dx^2 \right) \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{2} (\log(x^2) - \log(x^2+1))
 \end{aligned}$$

input `Int[1/(x*(1 + x^2)),x]`

output `(Log[x^2] - Log[1 + x^2])/2`

3.162.3.1 Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`


```
rule 47 Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c
- a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x
], x] /; FreeQ[{a, b, c, d}, x]
```

```
rule 243 Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int
t[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I
ntegerQ[(m - 1)/2]
```

3.162.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
default	$\ln(x) - \frac{\ln(x^2+1)}{2}$	12
norman	$\ln(x) - \frac{\ln(x^2+1)}{2}$	12
meijerg	$\ln(x) - \frac{\ln(x^2+1)}{2}$	12
risch	$\ln(x) - \frac{\ln(x^2+1)}{2}$	12
parallelrisch	$\ln(x) - \frac{\ln(x^2+1)}{2}$	12

```
input int(1/x/(x^2+1),x,method=_RETURNVERBOSE)
```

```
output ln(x)-1/2*ln(x^2+1)
```

3.162.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1}{x(1+x^2)} dx = -\frac{1}{2} \log(x^2 + 1) + \log(x)$$

```
input integrate(1/x/(x^2+1),x, algorithm="fricas")
```

```
output -1/2*log(x^2 + 1) + log(x)
```

3.162.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \frac{1}{x(1+x^2)} dx = \log(x) - \frac{\log(x^2+1)}{2}$$

input `integrate(1/x/(x**2+1),x)`output `log(x) - log(x**2 + 1)/2`**3.162.7 Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{1}{x(1+x^2)} dx = -\frac{1}{2} \log(x^2+1) + \frac{1}{2} \log(x^2)$$

input `integrate(1/x/(x^2+1),x, algorithm="maxima")`output `-1/2*log(x^2 + 1) + 1/2*log(x^2)`**3.162.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{1}{x(1+x^2)} dx = -\frac{1}{2} \log(x^2+1) + \frac{1}{2} \log(x^2)$$

input `integrate(1/x/(x^2+1),x, algorithm="giac")`output `-1/2*log(x^2 + 1) + 1/2*log(x^2)`

3.162.9 Mupad [B] (verification not implemented)

Time = 13.30 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1}{x(1+x^2)} dx = \ln(x) - \frac{\ln(x^2+1)}{2}$$

input `int(1/(x*(x^2 + 1)),x)`

output `log(x) - log(x^2 + 1)/2`

3.163 $\int \operatorname{arccosh}(x) dx$

3.163.1 Optimal result	875
3.163.2 Mathematica [A] (verified)	875
3.163.3 Rubi [A] (verified)	876
3.163.4 Maple [A] (verified)	877
3.163.5 Fricas [A] (verification not implemented)	877
3.163.6 Sympy [F]	877
3.163.7 Maxima [A] (verification not implemented)	878
3.163.8 Giac [A] (verification not implemented)	878
3.163.9 Mupad [B] (verification not implemented)	878

3.163.1 Optimal result

Integrand size = 2, antiderivative size = 21

$$\int \operatorname{arccosh}(x) dx = -\sqrt{-1+x}\sqrt{1+x} + x\operatorname{arccosh}(x)$$

output `-(-1+x)^(1/2)*(1+x)^(1/2)+x*arccosh(x)`

3.163.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \operatorname{arccosh}(x) dx = -\sqrt{-1+x}\sqrt{1+x} + x\operatorname{arccosh}(x)$$

input `Integrate[ArcCosh[x],x]`

output `-(Sqrt[-1 + x]*Sqrt[1 + x]) + x*ArcCosh[x]`

3.163.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {6294, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arccosh}(x) dx$$

$$\downarrow 6294$$

$$x \operatorname{arccosh}(x) - \int \frac{x}{\sqrt{x-1}\sqrt{x+1}} dx$$

$$\downarrow 83$$

$$x \operatorname{arccosh}(x) - \sqrt{x-1}\sqrt{x+1}$$

input `Int[ArcCosh[x], x]`

output `-(Sqrt[-1 + x]*Sqrt[1 + x]) + x*ArcCosh[x]`

3.163.3.1 Defintions of rubi rules used

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] :> Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

rule 6294 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.), x_Symbol] :> Simp[x*(a + b*ArcCosh[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

3.163.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

method	result	size
lookup	$-\sqrt{-1+x}\sqrt{1+x}+x \operatorname{arccosh}(x)$	18
default	$-\sqrt{-1+x}\sqrt{1+x}+x \operatorname{arccosh}(x)$	18
parts	$-\sqrt{-1+x}\sqrt{1+x}+x \operatorname{arccosh}(x)$	18

input `int(arccosh(x),x,method=_RETURNVERBOSE)`output `-(-1+x)^(1/2)*(1+x)^(1/2)+x*arccosh(x)`**3.163.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \operatorname{arccosh}(x) dx = x \log(x + \sqrt{x^2 - 1}) - \sqrt{x^2 - 1}$$

input `integrate(arccosh(x),x, algorithm="fricas")`output `x*log(x + sqrt(x^2 - 1)) - sqrt(x^2 - 1)`**3.163.6 Sympy [F]**

$$\int \operatorname{arccosh}(x) dx = \int \operatorname{acosh}(x) dx$$

input `integrate(acosh(x),x)`output `Integral(acosh(x), x)`

3.163.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

$$\int \operatorname{arccosh}(x) dx = x \operatorname{arccosh}(x) - \sqrt{x^2 - 1}$$

input `integrate(arccosh(x),x, algorithm="maxima")`output `x*arccosh(x) - sqrt(x^2 - 1)`**3.163.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \operatorname{arccosh}(x) dx = x \log(x + \sqrt{x^2 - 1}) - \sqrt{x^2 - 1}$$

input `integrate(arccosh(x),x, algorithm="giac")`output `x*log(x + sqrt(x^2 - 1)) - sqrt(x^2 - 1)`**3.163.9 Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \operatorname{arccosh}(x) dx = x \operatorname{acosh}(x) - \sqrt{x - 1} \sqrt{x + 1}$$

input `int(acosh(x),x)`output `x*acosh(x) - (x - 1)^(1/2)*(x + 1)^(1/2)`

3.164 $\int e^{-3-5x-2x^2} dx$

3.164.1 Optimal result	879
3.164.2 Mathematica [A] (verified)	879
3.164.3 Rubi [A] (verified)	880
3.164.4 Maple [A] (verified)	881
3.164.5 Fricas [A] (verification not implemented)	881
3.164.6 Sympy [A] (verification not implemented)	881
3.164.7 Maxima [A] (verification not implemented)	882
3.164.8 Giac [A] (verification not implemented)	882
3.164.9 Mupad [B] (verification not implemented)	882

3.164.1 Optimal result

Integrand size = 12, antiderivative size = 34

$$\int e^{-3-5x-2x^2} dx = \frac{\sqrt[8]{e}\sqrt{\pi}\operatorname{erf}\left(\frac{5+4x}{2\sqrt{2}}\right)}{2\sqrt{2}}$$

output `1/4*exp(1/8)*Pi^(1/2)*erf(1/4*(5+4*x)*2^(1/2))*2^(1/2)`

3.164.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00

$$\int e^{-3-5x-2x^2} dx = \frac{\sqrt[8]{e}\sqrt{\pi}\operatorname{erf}\left(\frac{5+4x}{2\sqrt{2}}\right)}{2\sqrt{2}}$$

input `Integrate[E^(-3 - 5*x - 2*x^2),x]`

output `(E^(1/8)*Sqrt[Pi]*Erf[(5 + 4*x)/(2*Sqrt[2])])/(2*Sqrt[2])`

3.164.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2664, 2634}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{-2x^2-5x-3} dx$$

$$\downarrow \text{2664}$$

$$\sqrt[8]{e} \int e^{-\frac{1}{8}(4x+5)^2} dx$$

$$\downarrow \text{2634}$$

$$\frac{\sqrt[8]{e} \sqrt{\pi} \operatorname{erf}\left(\frac{4x+5}{2\sqrt{2}}\right)}{2\sqrt{2}}$$

input `Int[E^(-3 - 5*x - 2*x^2), x]`

output `(E^(1/8)*Sqrt[Pi]*Erf[(5 + 4*x)/(2*Sqrt[2])])/(2*Sqrt[2])`

3.164.3.1 Defintions of rubi rules used

rule 2634 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)) ^2), x_Symbol] :> Simp[F^a*Sqrt[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]])/(2*d*Rt[(-b)*Log[F], 2])], x] /; FreeQ[{F, a, b, c, d}, x] && NegQ[b]`

rule 2664 `Int[(F_)^((a_.) + (b_.)*(x_) + (c_.)*(x_) ^2), x_Symbol] :> Simp[F^(a - b^2/(4*c)) Int[F^((b + 2*c*x)^2/(4*c)), x], x] /; FreeQ[{F, a, b, c}, x]`

3.164.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.68

method	result	size
default	$\frac{\sqrt{\pi} e^{\frac{1}{8}} \sqrt{2} \operatorname{erf}\left(x\sqrt{2} + \frac{5\sqrt{2}}{4}\right)}{4}$	23
risch	$\frac{\sqrt{\pi} e^{\frac{1}{8}} \sqrt{2} \operatorname{erf}\left(x\sqrt{2} + \frac{5\sqrt{2}}{4}\right)}{4}$	23

input `int(exp(-2*x^2-5*x-3),x,method=_RETURNVERBOSE)`output `1/4*Pi^(1/2)*exp(1/8)*2^(1/2)*erf(x*2^(1/2)+5/4*2^(1/2))`**3.164.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.62

$$\int e^{-3-5x-2x^2} dx = \frac{1}{4} \sqrt{2} \sqrt{\pi} \operatorname{erf}\left(\frac{1}{4} \sqrt{2}(4x+5)\right) e^{\frac{1}{8}}$$

input `integrate(exp(-2*x^2-5*x-3),x, algorithm="fricas")`output `1/4*sqrt(2)*sqrt(pi)*erf(1/4*sqrt(2)*(4*x + 5))*e^(1/8)`**3.164.6 Sympy [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.94

$$\int e^{-3-5x-2x^2} dx = \frac{\sqrt{2} \sqrt{\pi} e^{\frac{1}{8}} \operatorname{erf}\left(\sqrt{2}x + \frac{5\sqrt{2}}{4}\right)}{4}$$

input `integrate(exp(-2*x**2-5*x-3),x)`output `sqrt(2)*sqrt(pi)*exp(1/8)*erf(sqrt(2)*x + 5*sqrt(2)/4)/4`

3.164.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.65

$$\int e^{-3-5x-2x^2} dx = \frac{1}{4} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\sqrt{2}x + \frac{5}{4} \sqrt{2} \right) e^{\frac{1}{8}}$$

input `integrate(exp(-2*x^2-5*x-3),x, algorithm="maxima")`output `1/4*sqrt(2)*sqrt(pi)*erf(sqrt(2)*x + 5/4*sqrt(2))*e^(1/8)`**3.164.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.62

$$\int e^{-3-5x-2x^2} dx = \frac{1}{4} \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\frac{1}{4} \sqrt{2}(4x + 5) \right) e^{\frac{1}{8}}$$

input `integrate(exp(-2*x^2-5*x-3),x, algorithm="giac")`output `1/4*sqrt(2)*sqrt(pi)*erf(1/4*sqrt(2)*(4*x + 5))*e^(1/8)`**3.164.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.65

$$\int e^{-3-5x-2x^2} dx = \frac{\sqrt{2} \sqrt{\pi} e^{1/8} \operatorname{erf} \left(\sqrt{2}x + \frac{5\sqrt{2}}{4} \right)}{4}$$

input `int(exp(- 5*x - 2*x^2 - 3),x)`output `(2^(1/2)*pi^(1/2)*exp(1/8)*erf(2^(1/2)*x + (5*2^(1/2))/4))/4`

3.165 $\int \sin(\sqrt{x}) dx$

3.165.1 Optimal result	883
3.165.2 Mathematica [A] (verified)	883
3.165.3 Rubi [A] (verified)	884
3.165.4 Maple [A] (verified)	885
3.165.5 Fricas [A] (verification not implemented)	886
3.165.6 Sympy [A] (verification not implemented)	886
3.165.7 Maxima [A] (verification not implemented)	886
3.165.8 Giac [A] (verification not implemented)	887
3.165.9 Mupad [B] (verification not implemented)	887

3.165.1 Optimal result

Integrand size = 6, antiderivative size = 22

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

output `-2*x^(1/2)*cos(x^(1/2))+2*sin(x^(1/2))`

3.165.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

input `Integrate[Sin[Sqrt[x]],x]`

output `-2*Sqrt[x]*Cos[Sqrt[x]] + 2*Sin[Sqrt[x]]`

3.165.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {3842, 3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(\sqrt{x}) \, dx \\
 & \quad \downarrow \text{3842} \\
 & 2 \int \sqrt{x} \sin(\sqrt{x}) \, d\sqrt{x} \\
 & \quad \downarrow \text{3042} \\
 & 2 \int \sqrt{x} \sin(\sqrt{x}) \, d\sqrt{x} \\
 & \quad \downarrow \text{3777} \\
 & 2 \left(\int \cos(\sqrt{x}) \, d\sqrt{x} - \sqrt{x} \cos(\sqrt{x}) \right) \\
 & \quad \downarrow \text{3042} \\
 & 2 \left(\int \sin\left(\sqrt{x} + \frac{\pi}{2}\right) \, d\sqrt{x} - \sqrt{x} \cos(\sqrt{x}) \right) \\
 & \quad \downarrow \text{3117} \\
 & 2(\sin(\sqrt{x}) - \sqrt{x} \cos(\sqrt{x}))
 \end{aligned}$$

input `Int[Sin[Sqrt[x]], x]`

output `2*(-(Sqrt[x]*Cos[Sqrt[x]]) + Sin[Sqrt[x]])`

3.165.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3842 `Int[((a_.) + (b_.)*Sin[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)])^(p_.), x_Symbol] := Simp[1/(n*f) Subst[Int[x^(1/n - 1)*(a + b*SIN[c + d*x])^p, x], (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]`

3.165.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

method	result	size
derivativedivides	$-2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$	17
default	$-2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$	17
meijerg	$4\sqrt{\pi} \left(-\frac{\sqrt{x} \cos(\sqrt{x})}{2\sqrt{\pi}} + \frac{\sin(\sqrt{x})}{2\sqrt{\pi}} \right)$	28

input `int(sin(x^(1/2)),x,method=_RETURNVERBOSE)`

output `-2*x^(1/2)*cos(x^(1/2))+2*sin(x^(1/2))`

3.165.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

input `integrate(sin(x^(1/2)),x, algorithm="fricas")`

output `-2*sqrt(x)*cos(sqrt(x)) + 2*sin(sqrt(x))`

3.165.6 Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

input `integrate(sin(x**(1/2)),x)`

output `-2*sqrt(x)*cos(sqrt(x)) + 2*sin(sqrt(x))`

3.165.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

input `integrate(sin(x^(1/2)),x, algorithm="maxima")`

output `-2*sqrt(x)*cos(sqrt(x)) + 2*sin(sqrt(x))`

3.165.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

input `integrate(sin(x^(1/2)),x, algorithm="giac")`

output `-2*sqrt(x)*cos(sqrt(x)) + 2*sin(sqrt(x))`

3.165.9 Mupad [B] (verification not implemented)

Time = 13.61 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \sin(\sqrt{x}) dx = 2 \sin(\sqrt{x}) - 2\sqrt{x} \cos(\sqrt{x})$$

input `int(sin(x^(1/2)),x)`

output `2*sin(x^(1/2)) - 2*x^(1/2)*cos(x^(1/2))`

3.166 $\int \frac{1}{\left(\frac{1}{x}+x\right)^2} dx$

3.166.1 Optimal result	888
3.166.2 Mathematica [A] (verified)	888
3.166.3 Rubi [A] (verified)	889
3.166.4 Maple [A] (verified)	890
3.166.5 Fricas [A] (verification not implemented)	890
3.166.6 Sympy [A] (verification not implemented)	891
3.166.7 Maxima [A] (verification not implemented)	891
3.166.8 Giac [A] (verification not implemented)	891
3.166.9 Mupad [B] (verification not implemented)	892

3.166.1 Optimal result

Integrand size = 7, antiderivative size = 19

$$\int \frac{1}{\left(\frac{1}{x}+x\right)^2} dx = -\frac{x}{2(1+x^2)} + \frac{\arctan(x)}{2}$$

output -x/(2*x^2+2)+1/2*arctan(x)

3.166.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{1}{\left(\frac{1}{x}+x\right)^2} dx = -\frac{x}{2(1+x^2)} + \frac{\arctan(x)}{2}$$

input Integrate[(x^(-1) + x)^(-2), x]

output -1/2*x/(1 + x^2) + ArcTan[x]/2

3.166.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {2027, 252, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\left(x + \frac{1}{x}\right)^2} dx \\ & \quad \downarrow \text{2027} \\ & \int \frac{x^2}{(x^2 + 1)^2} dx \\ & \quad \downarrow \text{252} \\ & \frac{1}{2} \int \frac{1}{x^2 + 1} dx - \frac{x}{2(x^2 + 1)} \\ & \quad \downarrow \text{216} \\ & \frac{\arctan(x)}{2} - \frac{x}{2(x^2 + 1)} \end{aligned}$$

input `Int[(x^(-1) + x)^(-2),x]`

output `-1/2*x/(1 + x^2) + ArcTan[x]/2`

3.166.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 252 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] - Simp[c^2*((m - 1)/(2*b*(p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m + 2*p + 3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2027 `Int[(Fx_.)*((a_.)*(x_)^(r_.) + (b_.)*(x_)^(s_.))^p_.], x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r))^p*Fx, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] && PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])`

3.166.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

method	result	size
default	$-\frac{x}{2(x^2+1)} + \frac{\arctan(x)}{2}$	16
risch	$-\frac{x}{2(x^2+1)} + \frac{\arctan(x)}{2}$	16
parallelrisc	$-\frac{i \ln(x-i)x^2 - i \ln(i+x)x^2 + i \ln(x-i) - i \ln(i+x) + 2x}{4(x^2+1)}$	52

input `int(1/(x+1/x)^2,x,method=_RETURNVERBOSE)`

output `-1/2/(x^2+1)*x+1/2*arctan(x)`

3.166.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{1}{\left(\frac{1}{x} + x\right)^2} dx = \frac{(x^2 + 1) \arctan(x) - x}{2(x^2 + 1)}$$

input `integrate(1/(x+1/x)^2,x, algorithm="fricas")`

output `1/2*((x^2 + 1)*arctan(x) - x)/(x^2 + 1)`

3.166.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{1}{\left(\frac{1}{x} + x\right)^2} dx = -\frac{x}{2x^2 + 2} + \frac{\operatorname{atan}(x)}{2}$$

input `integrate(1/(x+1/x)**2,x)`output `-x/(2*x**2 + 2) + atan(x)/2`**3.166.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{\left(\frac{1}{x} + x\right)^2} dx = -\frac{x}{2(x^2 + 1)} + \frac{1}{2} \arctan(x)$$

input `integrate(1/(x+1/x)^2,x, algorithm="maxima")`output `-1/2*x/(x^2 + 1) + 1/2*arctan(x)`**3.166.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{\left(\frac{1}{x} + x\right)^2} dx = -\frac{x}{2(x^2 + 1)} + \frac{1}{2} \arctan(x)$$

input `integrate(1/(x+1/x)^2,x, algorithm="giac")`output `-1/2*x/(x^2 + 1) + 1/2*arctan(x)`

3.166.9 Mupad [B] (verification not implemented)

Time = 13.51 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{1}{\left(\frac{1}{x} + x\right)^2} dx = \frac{\operatorname{atan}(x)}{2} - \frac{x}{2(x^2 + 1)}$$

input `int(1/(x + 1/x)^2,x)`

output `atan(x)/2 - x/(2*(x^2 + 1))`

3.167 $\int \frac{e^{-x}(2+x)}{x^3} dx$

3.167.1 Optimal result	893
3.167.2 Mathematica [A] (verified)	893
3.167.3 Rubi [A] (verified)	894
3.167.4 Maple [A] (verified)	895
3.167.5 Fricas [A] (verification not implemented)	895
3.167.6 Sympy [A] (verification not implemented)	896
3.167.7 Maxima [C] (verification not implemented)	896
3.167.8 Giac [A] (verification not implemented)	896
3.167.9 Mupad [B] (verification not implemented)	897

3.167.1 Optimal result

Integrand size = 12, antiderivative size = 10

$$\int \frac{e^{-x}(2+x)}{x^3} dx = -\frac{e^{-x}}{x^2}$$

output `-1/exp(x)/x^2`

3.167.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{e^{-x}(2+x)}{x^3} dx = -\frac{e^{-x}}{x^2}$$

input `Integrate[(2 + x)/(E^x*x^3),x]`

output `-(1/(E^x*x^2))`

3.167.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2627}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{-x}(x+2)}{x^3} dx$$

↓ 2627

$$-\frac{e^{-x}}{x^2}$$

input `Int[(2 + x)/(E^x*x^3),x]`

output `-(1/(E^x*x^2))`

3.167.3.1 Defintions of rubi rules used

rule 2627 `Int[(F_)^(v_)*((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_)), x_Symbol] :>
Simp[g*(d + e*x)^(m + 1)*(F^v/(D[v, x]*e*Log[F])), x] /; FreeQ[{F, d, e, f
, g, m}, x] && LinearQ[v, x] && EqQ[e*g*(m + 1) - D[v, x]*(e*f - d*g)*Log[F
, 0]`

3.167.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

method	result	size
gosper	$-\frac{e^{-x}}{x^2}$	10
derivativdivides	$-\frac{e^{-x}}{x^2}$	10
default	$-\frac{e^{-x}}{x^2}$	10
norman	$-\frac{e^{-x}}{x^2}$	10
risch	$-\frac{e^{-x}}{x^2}$	10
parallelrisch	$-\frac{e^{-x}}{x^2}$	10
meijerg	$-\frac{1}{x^2} + \frac{1}{x} - \frac{1}{2} + \frac{9x^2-12x+6}{6x^2} - \frac{(3-3x)e^{-x}}{3x^2} + \frac{-2x+2}{2x} - \frac{e^{-x}}{x}$	59

input `int((2+x)*exp(-x)/x^3,x,method=_RETURNVERBOSE)`output `-1/x^2*exp(-x)`**3.167.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{e^{-x}(2+x)}{x^3} dx = -\frac{e^{(-x)}}{x^2}$$

input `integrate((2+x)*exp(-x)/x^3,x, algorithm="fricas")`output `-e^(-x)/x^2`

3.167.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{e^{-x}(2+x)}{x^3} dx = -\frac{e^{-x}}{x^2}$$

input `integrate((2+x)*exp(-x)/x**3,x)`

output `-exp(-x)/x**2`

3.167.7 Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 0.23 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int \frac{e^{-x}(2+x)}{x^3} dx = -\Gamma(-1, x) - 2\Gamma(-2, x)$$

input `integrate((2+x)*exp(-x)/x^3,x, algorithm="maxima")`

output `-gamma(-1, x) - 2*gamma(-2, x)`

3.167.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{e^{-x}(2+x)}{x^3} dx = -\frac{e^{(-x)}}{x^2}$$

input `integrate((2+x)*exp(-x)/x^3,x, algorithm="giac")`

output `-e^(-x)/x^2`

3.167.9 Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{e^{-x}(2+x)}{x^3} dx = -\frac{e^{-x}}{x^2}$$

input `int((exp(-x)*(x + 2))/x^3,x)`

output `-exp(-x)/x^2`

3.168 $\int \frac{1}{\sqrt{(1-x)x}} dx$

3.168.1 Optimal result	898
3.168.2 Mathematica [B] (verified)	898
3.168.3 Rubi [A] (verified)	899
3.168.4 Maple [A] (verified)	900
3.168.5 Fricas [B] (verification not implemented)	900
3.168.6 Sympy [A] (verification not implemented)	900
3.168.7 Maxima [A] (verification not implemented)	901
3.168.8 Giac [B] (verification not implemented)	901
3.168.9 Mupad [B] (verification not implemented)	901

3.168.1 Optimal result

Integrand size = 11, antiderivative size = 8

$$\int \frac{1}{\sqrt{(1-x)x}} dx = -\arcsin(1-2x)$$

output `arcsin(-1+2*x)`

3.168.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 40 vs. 2(8) = 16.

Time = 0.05 (sec) , antiderivative size = 40, normalized size of antiderivative = 5.00

$$\int \frac{1}{\sqrt{(1-x)x}} dx = -\frac{2\sqrt{-1+x}\sqrt{x} \log(\sqrt{-1+x} - \sqrt{x})}{\sqrt{-((-1+x)x)}}$$

input `Integrate[1/Sqrt[(1-x)*x],x]`

output `(-2*Sqrt[-1+x]*Sqrt[x]*Log[Sqrt[-1+x]-Sqrt[x]])/Sqrt[-((-1+x)*x)]`

3.168.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {2048, 1090, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt{(1-x)x}} dx \\ & \quad \downarrow \text{2048} \\ & \int \frac{1}{\sqrt{x-x^2}} dx \\ & \quad \downarrow \text{1090} \\ & - \int \frac{1}{\sqrt{1-(1-2x)^2}} d(1-2x) \\ & \quad \downarrow \text{223} \\ & - \arcsin(1-2x) \end{aligned}$$

input `Int[1/Sqrt[(1 - x)*x],x]`

output `-ArcSin[1 - 2*x]`

3.168.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 1090 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

rule 2048 `Int[(u_.)*((e_.)*((a_.) + (b_.)*(x_)^(n_.))*((c_) + (d_.)*(x_)^(n_.)))^(p_) , x_Symbol] := Int[u*(a*c*e + (b*c + a*d)*e*x^n + b*d*e*x^(2*n))^p, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

3.168.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
default	$\arcsin(-1 + 2x)$	7
meijerg	$2 \arcsin(\sqrt{x})$	7
pseudoelliptic	$-2 \arctan\left(\frac{\sqrt{-(-1+x)x}}{x}\right)$	16
trager	$\text{RootOf}(-Z^2 + 1) \ln(-2 \text{RootOf}(-Z^2 + 1)x + 2\sqrt{-x^2 + x} + \text{RootOf}(-Z^2 + 1))$	36

input `int(1/(x*(1-x))^(1/2),x,method=_RETURNVERBOSE)`output `arcsin(-1+2*x)`**3.168.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 16 vs. 2(6) = 12.

Time = 0.25 (sec) , antiderivative size = 16, normalized size of antiderivative = 2.00

$$\int \frac{1}{\sqrt{(1-x)x}} dx = -2 \arctan\left(\frac{\sqrt{-x^2 + x}}{x}\right)$$

input `integrate(1/(x*(1-x))^(1/2),x, algorithm="fricas")`output `-2*arctan(sqrt(-x^2 + x)/x)`**3.168.6 Sympy [A] (verification not implemented)**

Time = 0.45 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \frac{1}{\sqrt{(1-x)x}} dx = \text{asin}(2x - 1)$$

input `integrate(1/(x*(1-x))**(1/2),x)`output `asin(2*x - 1)`

3.168.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{(1-x)x}} dx = \arcsin(2x - 1)$$

input `integrate(1/(x*(1-x))^(1/2),x, algorithm="maxima")`

output `arcsin(2*x - 1)`

3.168.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. 2(6) = 12.

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 3.12

$$\int \frac{1}{\sqrt{(1-x)x}} dx = \frac{1}{4} \sqrt{-x^2 + x}(2x - 1) + \frac{1}{8} \arcsin(2x - 1)$$

input `integrate(1/(x*(1-x))^(1/2),x, algorithm="giac")`

output `1/4*sqrt(-x^2 + x)*(2*x - 1) + 1/8*arcsin(2*x - 1)`

3.168.9 Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{(1-x)x}} dx = \operatorname{asin}(2x - 1)$$

input `int(1/(-x*(x - 1))^(1/2),x)`

output `asin(2*x - 1)`

3.169 $\int e^{-x} \tanh(x) dx$

3.169.1 Optimal result	902
3.169.2 Mathematica [A] (verified)	902
3.169.3 Rubi [A] (verified)	903
3.169.4 Maple [A] (verified)	904
3.169.5 Fricas [B] (verification not implemented)	904
3.169.6 Sympy [F]	905
3.169.7 Maxima [A] (verification not implemented)	905
3.169.8 Giac [A] (verification not implemented)	905
3.169.9 Mupad [B] (verification not implemented)	906

3.169.1 Optimal result

Integrand size = 8, antiderivative size = 12

$$\int e^{-x} \tanh(x) dx = e^{-x} + 2 \arctan(e^x)$$

output `exp(-x)+2*arctan(exp(x))`

3.169.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int e^{-x} \tanh(x) dx = e^{-x} + 2 \arctan(e^x)$$

input `Integrate[Tanh[x]/E^x,x]`

output `E^(-x) + 2*ArcTan[E^x]`

3.169.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {2720, 25, 359, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int e^{-x} \tanh(x) dx \\
 \downarrow \text{2720} \\
 \int -\frac{e^{-2x}(1-e^{2x})}{e^{2x}+1} de^x \\
 \downarrow \text{25} \\
 -\int \frac{e^{-2x}(1-e^{2x})}{1+e^{2x}} de^x \\
 \downarrow \text{359} \\
 2 \int \frac{1}{1+e^{2x}} de^x + e^{-x} \\
 \downarrow \text{216} \\
 2 \arctan(e^x) + e^{-x}
 \end{array}$$

input `Int[Tanh[x]/E^x,x]`

output `E^(-x) + 2*ArcTan[E^x]`

3.169.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`


```
rule 359 Int[((e._)*(x._))^(m._)*((a._) + (b._)*(x._)^2)^(p._)*((c._) + (d._)*(x._)^2), x
_Symbol] := Simp[c*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*e*(m + 1))), x] +
Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(a*e^2*(m + 1)) Int[(e*x)^(m + 2)*
(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0]
&& LtQ[m, -1] && !LtQ[p, -1]
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a._)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c._)*((a._) + (b._)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.169.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.58

method	result	size
default	$\frac{2}{\tanh(\frac{x}{2})+1} + 2 \arctan\left(\tanh\left(\frac{x}{2}\right)\right)$	19
risch	$e^{-x} + i \ln(e^x + i) - i \ln(e^x - i)$	24

```
input int(tanh(x)/exp(x),x,method=_RETURNVERBOSE)
```

```
output 2/(tanh(1/2*x)+1)+2*arctan(tanh(1/2*x))
```

3.169.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 23 vs. $2(10) = 20$.

Time = 0.24 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.92

$$\int e^{-x} \tanh(x) dx = \frac{2(\cosh(x) + \sinh(x)) \arctan(\cosh(x) + \sinh(x)) + 1}{\cosh(x) + \sinh(x)}$$

```
input integrate(tanh(x)/exp(x),x, algorithm="fricas")
```

```
output (2*(cosh(x) + sinh(x))*arctan(cosh(x) + sinh(x)) + 1)/(cosh(x) + sinh(x))
```

3.169.6 Sympy [F]

$$\int e^{-x} \tanh(x) dx = \int e^{-x} \tanh(x) dx$$

input `integrate(tanh(x)/exp(x),x)`

output `Integral(exp(-x)*tanh(x), x)`

3.169.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int e^{-x} \tanh(x) dx = -2 \arctan(e^{(-x)}) + e^{(-x)}$$

input `integrate(tanh(x)/exp(x),x, algorithm="maxima")`

output `-2*arctan(e^(-x)) + e^(-x)`

3.169.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int e^{-x} \tanh(x) dx = 2 \arctan(e^x) + e^{(-x)}$$

input `integrate(tanh(x)/exp(x),x, algorithm="giac")`

output `2*arctan(e^x) + e^(-x)`

3.169.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int e^{-x} \tanh(x) dx = e^{-x} - 2 \operatorname{atan}(e^{-x})$$

input `int(exp(-x)*tanh(x),x)`

output `exp(-x) - 2*atan(exp(-x))`

3.170 $\int \sqrt{1 + \sin(x)} dx$

3.170.1 Optimal result	907
3.170.2 Mathematica [B] (verified)	907
3.170.3 Rubi [A] (verified)	908
3.170.4 Maple [A] (verified)	909
3.170.5 Fricas [B] (verification not implemented)	909
3.170.6 Sympy [F]	909
3.170.7 Maxima [F]	910
3.170.8 Giac [B] (verification not implemented)	910
3.170.9 Mupad [B] (verification not implemented)	910

3.170.1 Optimal result

Integrand size = 8, antiderivative size = 12

$$\int \sqrt{1 + \sin(x)} dx = -\frac{2 \cos(x)}{\sqrt{1 + \sin(x)}}$$

output `-2*cos(x)/(1+sin(x))^(1/2)`

3.170.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 40 vs. 2(12) = 24.

Time = 0.02 (sec) , antiderivative size = 40, normalized size of antiderivative = 3.33

$$\int \sqrt{1 + \sin(x)} dx = \frac{2(-\cos(\frac{x}{2}) + \sin(\frac{x}{2})) \sqrt{1 + \sin(x)}}{\cos(\frac{x}{2}) + \sin(\frac{x}{2})}$$

input `Integrate[Sqrt[1 + Sin[x]],x]`

output `(2*(-Cos[x/2] + Sin[x/2])*Sqrt[1 + Sin[x]])/(Cos[x/2] + Sin[x/2])`

3.170.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\sin(x) + 1} dx$$

↓ 3042

$$\int \sqrt{\sin(x) + 1} dx$$

↓ 3125

$$-\frac{2 \cos(x)}{\sqrt{\sin(x) + 1}}$$

input `Int[Sqrt[1 + Sin[x]],x]`

output `(-2*Cos[x])/Sqrt[1 + Sin[x]]`

3.170.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

3.170.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.42

method	result	size
default	$\frac{2(-1+\sin(x))\sqrt{1+\sin(x)}}{\cos(x)}$	17
risch	$-\frac{i\sqrt{2}\sqrt{2+2\sin(x)}(e^{ix}-i)(i+e^{ix})}{e^{2ix}-1+2ie^{ix}}$	48

input `int((1+sin(x))^(1/2),x,method=_RETURNVERBOSE)`

output `2*(-1+sin(x))*(1+sin(x))^(1/2)/cos(x)`

3.170.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 24 vs. $2(10) = 20$.

Time = 0.24 (sec) , antiderivative size = 24, normalized size of antiderivative = 2.00

$$\int \sqrt{1 + \sin(x)} dx = -\frac{2(\cos(x) - \sin(x) + 1)\sqrt{\sin(x) + 1}}{\cos(x) + \sin(x) + 1}$$

input `integrate((1+sin(x))^(1/2),x, algorithm="fracas")`

output `-2*(cos(x) - sin(x) + 1)*sqrt(sin(x) + 1)/(cos(x) + sin(x) + 1)`

3.170.6 Sympy [F]

$$\int \sqrt{1 + \sin(x)} dx = \int \sqrt{\sin(x) + 1} dx$$

input `integrate((1+sin(x))**(1/2),x)`

output `Integral(sqrt(sin(x) + 1), x)`

3.170.7 Maxima [F]

$$\int \sqrt{1 + \sin(x)} dx = \int \sqrt{\sin(x) + 1} dx$$

input `integrate((1+sin(x))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(sin(x) + 1), x)`

3.170.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 22 vs. 2(10) = 20.

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.83

$$\int \sqrt{1 + \sin(x)} dx = 2\sqrt{2}\operatorname{sgn}\left(\cos\left(-\frac{1}{4}\pi + \frac{1}{2}x\right)\right)\sin\left(-\frac{1}{4}\pi + \frac{1}{2}x\right)$$

input `integrate((1+sin(x))^(1/2),x, algorithm="giac")`

output `2*sqrt(2)*sgn(cos(-1/4*pi + 1/2*x))*sin(-1/4*pi + 1/2*x)`

3.170.9 Mupad [B] (verification not implemented)

Time = 13.45 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.33

$$\int \sqrt{1 + \sin(x)} dx = \frac{2(\sin(x) - 1)\sqrt{\sin(x) + 1}}{\cos(x)}$$

input `int((sin(x) + 1)^(1/2),x)`

output `(2*(sin(x) - 1)*(sin(x) + 1)^(1/2))/cos(x)`

3.171 $\int \frac{1}{1+\sqrt{x}} dx$

3.171.1 Optimal result	911
3.171.2 Mathematica [A] (verified)	911
3.171.3 Rubi [A] (verified)	912
3.171.4 Maple [A] (verified)	913
3.171.5 Fricas [A] (verification not implemented)	913
3.171.6 Sympy [A] (verification not implemented)	913
3.171.7 Maxima [A] (verification not implemented)	914
3.171.8 Giac [A] (verification not implemented)	914
3.171.9 Mupad [B] (verification not implemented)	914

3.171.1 Optimal result

Integrand size = 9, antiderivative size = 18

$$\int \frac{1}{1+\sqrt{x}} dx = 2\sqrt{x} - 2\log(1+\sqrt{x})$$

output `2*x^(1/2)-2*ln(1+x^(1/2))`

3.171.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+\sqrt{x}} dx = 2\sqrt{x} - 2\log(1+\sqrt{x})$$

input `Integrate[(1 + Sqrt[x])^(-1),x]`

output `2*Sqrt[x] - 2*Log[1 + Sqrt[x]]`

3.171.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {774, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt{x}+1} dx \\ & \quad \downarrow \text{774} \\ & 2 \int \frac{\sqrt{x}}{\sqrt{x}+1} d\sqrt{x} \\ & \quad \downarrow \text{49} \\ & 2 \int \left(1 + \frac{1}{-\sqrt{x}-1}\right) d\sqrt{x} \\ & \quad \downarrow \text{2009} \\ & 2(\sqrt{x} - \log(\sqrt{x}+1)) \end{aligned}$$

input `Int[(1 + Sqrt[x])^(-1),x]`

output `2*(Sqrt[x] - Log[1 + Sqrt[x]])`

3.171.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 774 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[n]}, Simp[k Subst[Int[x^(k - 1)*(a + b*x^(k*n))^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, p}, x] && FractionQ[n]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.171.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$2\sqrt{x} - 2\ln(1 + \sqrt{x})$	15
meijerg	$2\sqrt{x} - 2\ln(1 + \sqrt{x})$	15
trager	$2\sqrt{x} - \ln(2\sqrt{x} + 1 + x)$	18
default	$2\sqrt{x} + \ln(-1 + \sqrt{x}) - \ln(1 + \sqrt{x}) - \ln(-1 + x)$	27

input `int(1/(1+x^(1/2)),x,method=_RETURNVERBOSE)`output `2*x^(1/2)-2*ln(1+x^(1/2))`**3.171.5 Fricas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{1}{1 + \sqrt{x}} dx = 2\sqrt{x} - 2 \log(\sqrt{x} + 1)$$

input `integrate(1/(1+x^(1/2)),x, algorithm="fricas")`output `2*sqrt(x) - 2*log(sqrt(x) + 1)`**3.171.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{1}{1 + \sqrt{x}} dx = 2\sqrt{x} - 2 \log(\sqrt{x} + 1)$$

input `integrate(1/(1+x**(1/2)),x)`output `2*sqrt(x) - 2*log(sqrt(x) + 1)`

3.171.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{1}{1+\sqrt{x}} dx = 2\sqrt{x} - 2\log(\sqrt{x}+1) + 2$$

input `integrate(1/(1+x^(1/2)),x, algorithm="maxima")`output `2*sqrt(x) - 2*log(sqrt(x) + 1) + 2`**3.171.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{1}{1+\sqrt{x}} dx = 2\sqrt{x} - 2\log(\sqrt{x}+1)$$

input `integrate(1/(1+x^(1/2)),x, algorithm="giac")`output `2*sqrt(x) - 2*log(sqrt(x) + 1)`**3.171.9 Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{1}{1+\sqrt{x}} dx = 2\sqrt{x} - 2\ln(\sqrt{x}+1)$$

input `int(1/(x^(1/2) + 1),x)`output `2*x^(1/2) - 2*log(x^(1/2) + 1)`

3.172 $\int e^{-x^2} \sin^2\left(\frac{\pi}{4} + x\right) dx$

3.172.1 Optimal result	915
3.172.2 Mathematica [A] (verified)	915
3.172.3 Rubi [A] (verified)	916
3.172.4 Maple [A] (verified)	917
3.172.5 Fricas [A] (verification not implemented)	917
3.172.6 Sympy [F]	917
3.172.7 Maxima [A] (verification not implemented)	918
3.172.8 Giac [F]	918
3.172.9 Mupad [F(-1)]	918

3.172.1 Optimal result

Integrand size = 18, antiderivative size = 52

$$\int e^{-x^2} \sin^2\left(\frac{\pi}{4} + x\right) dx = \frac{1}{4}\sqrt{\pi}\operatorname{erf}(x) - \frac{\sqrt{\pi}\operatorname{erfi}(1 - ix)}{8e} - \frac{\sqrt{\pi}\operatorname{erfi}(1 + ix)}{8e}$$

output `1/4*Pi^(1/2)*erf(x)+1/8*Pi^(1/2)*erfi(-1+I*x)*exp(-1)-1/8*Pi^(1/2)*erfi(1+I*x)*exp(-1)`

3.172.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.73

$$\int e^{-x^2} \sin^2\left(\frac{\pi}{4} + x\right) dx = \frac{\sqrt{\pi}(2e\operatorname{erf}(x) - \operatorname{erfi}(1 - ix) - \operatorname{erfi}(1 + ix))}{8e}$$

input `Integrate[Sin[Pi/4 + x]^2/E^x^2,x]`

output `(Sqrt[Pi]*(2*E*Erf[x] - Erfi[1 - I*x] - Erfi[1 + I*x]))/(8*E)`

3.172.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {4975, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{-x^2} \sin^2\left(x + \frac{\pi}{4}\right) dx$$

↓ 4975

$$\int \left(\frac{e^{-x^2}}{2} + \frac{1}{4}ie^{-x^2-2ix} - \frac{1}{4}ie^{-x^2+2ix} \right) dx$$

↓ 2009

$$\frac{1}{4}\sqrt{\pi}\text{erf}(x) - \frac{\sqrt{\pi}\text{erfi}(1-ix)}{8e} - \frac{\sqrt{\pi}\text{erfi}(1+ix)}{8e}$$

input `Int[Sin[Pi/4 + x]^2/E^x^2,x]`

output `(Sqrt[Pi]*Erf[x])/4 - (Sqrt[Pi]*Erfi[1 - I*x])/(8*E) - (Sqrt[Pi]*Erfi[1 + I*x])/(8*E)`

3.172.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 4975 `Int[(F_)^(u_)*Sin[v_]^(n_), x_Symbol] := Int[ExpandTrigToExp[F^u, Sin[v]^n, x], x] /; FreeQ[F, x] && (LinearQ[u, x] || PolyQ[u, x, 2]) && (LinearQ[v, x] || PolyQ[v, x, 2]) && IGtQ[n, 0]`

3.172.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.67

method	result	size
risch	$\frac{i\sqrt{\pi}e^{-1}\operatorname{erf}(i+x)}{8} - \frac{i\sqrt{\pi}e^{-1}\operatorname{erf}(x-i)}{8} + \frac{\sqrt{\pi}\operatorname{erf}(x)}{4}$	35

input `int(sin(x+1/4*Pi)^2/exp(x^2),x,method=_RETURNVERBOSE)`output `1/8*I*Pi^(1/2)*exp(-1)*erf(I+x)-1/8*I*Pi^(1/2)*exp(-1)*erf(x-I)+1/4*Pi^(1/2)*erf(x)`**3.172.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.83

$$\int e^{-x^2} \sin^2\left(\frac{\pi}{4} + x\right) dx$$

$$= \frac{1}{8} \sqrt{\pi} \left((\operatorname{erf}(-x+i) e^{\frac{1}{2}i\pi-1} + 2 \operatorname{erf}(x)) e^{\frac{1}{2}i\pi+1} - \operatorname{erf}(x+i) \right) e^{-\frac{1}{2}i\pi-1}$$

input `integrate(sin(x+1/4*pi)^2/exp(x^2),x, algorithm="fracas")`output `1/8*sqrt(pi)*((erf(-x + I))*e^(1/2*I*pi - 1) + 2*erf(x))*e^(1/2*I*pi + 1) - erf(x + I))*e^(-1/2*I*pi - 1)`**3.172.6 Sympy [F]**

$$\int e^{-x^2} \sin^2\left(\frac{\pi}{4} + x\right) dx = \int e^{-x^2} \sin^2\left(x + \frac{\pi}{4}\right) dx$$

input `integrate(sin(x+1/4*pi)**2/exp(x**2),x)`output `Integral(exp(-x**2)*sin(x + pi/4)**2, x)`

3.172.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.50

$$\int e^{-x^2} \sin^2\left(\frac{\pi}{4} + x\right) dx = \frac{1}{8} \sqrt{\pi} (2 \operatorname{erf}(x) e + i \operatorname{erf}(x + i) - i \operatorname{erf}(x - i)) e^{(-1)}$$

input `integrate(sin(x+1/4*pi)^2/exp(x^2),x, algorithm="maxima")`output `1/8*sqrt(pi)*(2*erf(x)*e + I*erf(x + I) - I*erf(x - I))*e^(-1)`**3.172.8 Giac [F]**

$$\int e^{-x^2} \sin^2\left(\frac{\pi}{4} + x\right) dx = \int e^{(-x^2)} \sin\left(\frac{1}{4} \pi + x\right)^2 dx$$

input `integrate(sin(x+1/4*pi)^2/exp(x^2),x, algorithm="giac")`output `integrate(e^(-x^2)*sin(1/4*pi + x)^2, x)`**3.172.9 Mupad [F(-1)]**

Timed out.

$$\int e^{-x^2} \sin^2\left(\frac{\pi}{4} + x\right) dx = \int e^{-x^2} \sin\left(\frac{\pi}{4} + x\right)^2 dx$$

input `int(exp(-x^2)*sin(Pi/4 + x)^2,x)`output `int(exp(-x^2)*sin(Pi/4 + x)^2, x)`

3.173 $\int 3e^{-2x^3-x^6} x^2(1+x^3)^2 dx$

3.173.1 Optimal result	919
3.173.2 Mathematica [A] (verified)	919
3.173.3 Rubi [A] (verified)	920
3.173.4 Maple [F]	921
3.173.5 Fricas [A] (verification not implemented)	922
3.173.6 Sympy [F]	922
3.173.7 Maxima [C] (verification not implemented)	922
3.173.8 Giac [A] (verification not implemented)	923
3.173.9 Mupad [B] (verification not implemented)	923

3.173.1 Optimal result

Integrand size = 25, antiderivative size = 39

$$\int 3e^{-2x^3-x^6} x^2(1+x^3)^2 dx = -\frac{1}{2}e^{-2x^3-x^6}(1+x^3) + \frac{1}{4}e\sqrt{\pi}\operatorname{erf}(1+x^3)$$

output `-1/2*exp(-x^6-2*x^3)*(x^3+1)+1/4*exp(1)*Pi^(1/2)*erf(x^3+1)`

3.173.2 Mathematica [A] (verified)

Time = 0.70 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.95

$$\int 3e^{-2x^3-x^6} x^2(1+x^3)^2 dx = \frac{1}{4}\left(-2e^{-x^3(2+x^3)}(1+x^3) + e\sqrt{\pi}\operatorname{erf}(1+x^3)\right)$$

input `Integrate[3*E^(-2*x^3 - x^6)*x^2*(1 + x^3)^2,x]`

output `((-2*(1 + x^3))/E^(x^3*(2 + x^3)) + E*Sqrt[Pi]*Erf[1 + x^3])/4`

3.173.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {27, 7266, 2667, 2664, 2634}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int 3e^{-x^6-2x^3} x^2 (x^3 + 1)^2 dx \\
 & \quad \downarrow \text{27} \\
 & 3 \int e^{-x^6-2x^3} x^2 (x^3 + 1)^2 dx \\
 & \quad \downarrow \text{7266} \\
 & \int e^{-x^6-2x^3} (x^3 + 1)^2 dx^3 \\
 & \quad \downarrow \text{2667} \\
 & \frac{1}{2} \int e^{-x^6-2x^3} dx^3 - \frac{1}{2} e^{-x^6-2x^3} (x^3 + 1) \\
 & \quad \downarrow \text{2664} \\
 & \frac{1}{2} e \int e^{-(x^3+1)^2} dx^3 - \frac{1}{2} e^{-x^6-2x^3} (x^3 + 1) \\
 & \quad \downarrow \text{2634} \\
 & \frac{1}{4} e \sqrt{\pi} \operatorname{erf}(x^3 + 1) - \frac{1}{2} e^{-x^6-2x^3} (x^3 + 1)
 \end{aligned}$$

input `Int[3*E^(-2*x^3 - x^6)*x^2*(1 + x^3)^2,x]`

output `-1/2*(E^(-2*x^3 - x^6)*(1 + x^3)) + (E*sqrt[Pi]*Erf[1 + x^3])/4`

3.173.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 2634 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))2), x_Symbol] := Simp[Fa*Sqrt[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && NegQ[b]`
- rule 2664 `Int[(F_)^((a_.) + (b_.)*(x_) + (c_.)*(x_)2), x_Symbol] := Simp[F(a - b2/(4*c)) Int[F((b + 2*c*x)2/(4*c)), x], x] /; FreeQ[{F, a, b, c}, x]`
- rule 2667 `Int[(F_)^((a_.) + (b_.)*(x_) + (c_.)*(x_)2)*((d_.) + (e_.)*(x_))(m_), x_Symbol] := Simp[e*(d + e*x)(m - 1)*F(a + b*x + c*x2)/(2*c*Log[F]), x] - Simp[(m - 1)*(e2/(2*c*Log[F])) Int[(d + e*x)(m - 2)*F(a + b*x + c*x2), x], x] /; FreeQ[{F, a, b, c, d, e}, x] && EqQ[b*e - 2*c*d, 0] && GtQ[m, 1]`
- rule 7266 `Int[(u_)*(x_)(m_.), x_Symbol] := Simp[1/(m + 1) Subst[Int[SubstFor[x(m + 1), u, x], x, x(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x(m + 1), u, x]`

3.173.4 Maple [F]

$$\int 3x^2(x^3 + 1)^2 e^{-x^6 - 2x^3} dx$$

input `int(3*x^2*(x^3+1)^2*exp(-x^6-2*x^3),x)`

output `int(3*x^2*(x^3+1)^2*exp(-x^6-2*x^3),x)`

3.173.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.85

$$\int 3e^{-2x^3-x^6} x^2(1+x^3)^2 dx = \frac{1}{4} \sqrt{\pi} \operatorname{erf}(x^3+1) e - \frac{1}{2} (x^3+1) e^{-(x^6-2x^3)}$$

input `integrate(3*x^2*(x^3+1)^2*exp(-x^6-2*x^3),x, algorithm="fricas")`

output `1/4*sqrt(pi)*erf(x^3 + 1)*e - 1/2*(x^3 + 1)*e^(-x^6 - 2*x^3)`

3.173.6 Sympy [F]

$$\int 3e^{-2x^3-x^6} x^2(1+x^3)^2 dx = 3 \left(\int x^2 e^{-2x^3} e^{-x^6} dx + \int 2x^5 e^{-2x^3} e^{-x^6} dx + \int x^8 e^{-2x^3} e^{-x^6} dx \right)$$

input `integrate(3*x**2*(x**3+1)**2*exp(-x**6-2*x**3),x)`

output `3*(Integral(x**2*exp(-2*x**3)*exp(-x**6), x) + Integral(2*x**5*exp(-2*x**3)*exp(-x**6), x) + Integral(x**8*exp(-2*x**3)*exp(-x**6), x))`

3.173.7 Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 137, normalized size of antiderivative = 3.51

$$\int 3e^{-2x^3-x^6} x^2(1+x^3)^2 dx = \frac{1}{2} \sqrt{\pi} \operatorname{erf}(x^3+1) e$$

$$+ \frac{1}{2} i \left(\frac{i(x^3+1)^3 \Gamma\left(\frac{3}{2}, (x^3+1)^2\right)}{((x^3+1)^2)^{\frac{3}{2}}} - \frac{i \sqrt{\pi} (x^3+1) \left(\operatorname{erf}\left(\sqrt{(x^3+1)^2}\right) - 1 \right)}{\sqrt{(x^3+1)^2}} - 2i e^{-(x^3+1)^2} \right) e$$

$$+ i \left(\frac{i \sqrt{\pi} (x^3+1) \left(\operatorname{erf}\left(\sqrt{(x^3+1)^2}\right) - 1 \right)}{\sqrt{(x^3+1)^2}} + i e^{-(x^3+1)^2} \right) e$$

input `integrate(3*x^2*(x^3+1)^2*exp(-x^6-2*x^3),x, algorithm="maxima")`

output $\frac{1}{2}\sqrt{\pi}\operatorname{erf}(x^3 + 1)e + \frac{1}{2}I*(I*(x^3 + 1)^3\operatorname{gamma}(3/2, (x^3 + 1)^2) / ((x^3 + 1)^2)^{(3/2)} - I*\sqrt{\pi}*(x^3 + 1)*(\operatorname{erf}(\sqrt{(x^3 + 1)^2}) - 1)/\sqrt{(x^3 + 1)^2} - 2*I*e^{-(x^3 + 1)^2})*e + I*(I*\sqrt{\pi}*(x^3 + 1)*(\operatorname{erf}(\sqrt{(x^3 + 1)^2}) - 1)/\sqrt{(x^3 + 1)^2} + I*e^{-(x^3 + 1)^2})*e$

3.173.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.85

$$\int 3e^{-2x^3-x^6} x^2(1+x^3)^2 dx = \frac{1}{4}\sqrt{\pi}\operatorname{erf}(x^3 + 1)e - \frac{1}{2}(x^3 + 1)e^{(-x^6-2x^3)}$$

input `integrate(3*x^2*(x^3+1)^2*exp(-x^6-2*x^3),x, algorithm="giac")`

output $\frac{1}{4}\sqrt{\pi}\operatorname{erf}(x^3 + 1)e - \frac{1}{2}*(x^3 + 1)*e^{(-x^6 - 2*x^3)}$

3.173.9 Mupad [B] (verification not implemented)

Time = 14.61 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.15

$$\int 3e^{-2x^3-x^6} x^2(1+x^3)^2 dx = \frac{\sqrt{\pi}e\operatorname{erf}(x^3 + 1)}{4} - \frac{x^3 e^{-x^6-2x^3}}{2} - \frac{e^{-x^6-2x^3}}{2}$$

input `int(3*x^2*exp(- 2*x^3 - x^6)*(x^3 + 1)^2,x)`

output $(\pi^{(1/2)}*\exp(1)*\operatorname{erf}(x^3 + 1))/4 - (x^3*\exp(- 2*x^3 - x^6))/2 - \exp(- 2*x^3 - x^6)/2$

3.174 $\int e^{2x} \cos(3x) dx$

3.174.1 Optimal result	924
3.174.2 Mathematica [A] (verified)	924
3.174.3 Rubi [A] (verified)	925
3.174.4 Maple [A] (verified)	925
3.174.5 Fricas [A] (verification not implemented)	926
3.174.6 Sympy [A] (verification not implemented)	926
3.174.7 Maxima [A] (verification not implemented)	926
3.174.8 Giac [A] (verification not implemented)	927
3.174.9 Mupad [B] (verification not implemented)	927

3.174.1 Optimal result

Integrand size = 10, antiderivative size = 27

$$\int e^{2x} \cos(3x) dx = \frac{2}{13} e^{2x} \cos(3x) + \frac{3}{13} e^{2x} \sin(3x)$$

output `2/13*exp(2*x)*cos(3*x)+3/13*exp(2*x)*sin(3*x)`

3.174.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int e^{2x} \cos(3x) dx = \frac{1}{13} e^{2x} (2 \cos(3x) + 3 \sin(3x))$$

input `Integrate[E^(2*x)*Cos[3*x],x]`

output `(E^(2*x)*(2*Cos[3*x] + 3*Sin[3*x]))/13`

3.174.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {4933}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{2x} \cos(3x) dx$$

$$\downarrow 4933$$

$$\frac{3}{13} e^{2x} \sin(3x) + \frac{2}{13} e^{2x} \cos(3x)$$

input `Int[E^(2*x)*Cos[3*x],x]`

output `(2*E^(2*x)*Cos[3*x])/13 + (3*E^(2*x)*Sin[3*x])/13`

3.174.3.1 Defintions of rubi rules used

rule 4933 `Int[Cos[(d_.) + (e_.)*(x_.)]*(F_)^((c_.)*((a_.) + (b_.)*(x_.))), x_Symbol] :=
Simp[b*c*Log[F]*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x
] + Simp[e*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; F
reeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]`

3.174.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result	size
parallelrisc	$\frac{e^{2x}(2 \cos(3x) + 3 \sin(3x))}{13}$	20
default	$\frac{2 e^{2x} \cos(3x)}{13} + \frac{3 e^{2x} \sin(3x)}{13}$	22
risch	$\frac{e^{(2+3i)x}}{13} - \frac{3ie^{(2+3i)x}}{26} + \frac{e^{(2-3i)x}}{13} + \frac{3ie^{(2-3i)x}}{26}$	36
norman	$\frac{6 e^{2x} \tan\left(\frac{3x}{2}\right) - 2 e^{2x} \tan\left(\frac{3x}{2}\right)^2}{13} + \frac{2 e^{2x}}{13}$ $1 + \tan\left(\frac{3x}{2}\right)^2$	41

input `int(exp(2*x)*cos(3*x),x,method=_RETURNVERBOSE)`

output `1/13*exp(2*x)*(2*cos(3*x)+3*sin(3*x))`

3.174.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int e^{2x} \cos(3x) dx = \frac{2}{13} \cos(3x) e^{(2x)} + \frac{3}{13} e^{(2x)} \sin(3x)$$

input `integrate(exp(2*x)*cos(3*x),x, algorithm="fricas")`

output `2/13*cos(3*x)*e^(2*x) + 3/13*e^(2*x)*sin(3*x)`

3.174.6 Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int e^{2x} \cos(3x) dx = \frac{3e^{2x} \sin(3x)}{13} + \frac{2e^{2x} \cos(3x)}{13}$$

input `integrate(exp(2*x)*cos(3*x),x)`

output `3*exp(2*x)*sin(3*x)/13 + 2*exp(2*x)*cos(3*x)/13`

3.174.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^{2x} \cos(3x) dx = \frac{1}{13} (2 \cos(3x) + 3 \sin(3x)) e^{(2x)}$$

input `integrate(exp(2*x)*cos(3*x),x, algorithm="maxima")`

output `1/13*(2*cos(3*x) + 3*sin(3*x))*e^(2*x)`

3.174.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^{2x} \cos(3x) dx = \frac{1}{13} (2 \cos(3x) + 3 \sin(3x))e^{(2x)}$$

input `integrate(exp(2*x)*cos(3*x),x, algorithm="giac")`

output `1/13*(2*cos(3*x) + 3*sin(3*x))*e^(2*x)`

3.174.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int e^{2x} \cos(3x) dx = \frac{e^{2x} (2 \cos(3x) + 3 \sin(3x))}{13}$$

input `int(cos(3*x)*exp(2*x),x)`

output `(exp(2*x)*(2*cos(3*x) + 3*sin(3*x)))/13`

3.175 $\int \cos^{\cos(x)}(x)(1 + \log(\cos(x))) \sin(x) dx$

3.175.1 Optimal result	928
3.175.2 Mathematica [A] (verified)	928
3.175.3 Rubi [A] (verified)	929
3.175.4 Maple [F]	930
3.175.5 Fricas [A] (verification not implemented)	930
3.175.6 Sympy [F]	930
3.175.7 Maxima [A] (verification not implemented)	931
3.175.8 Giac [F]	931
3.175.9 Mupad [B] (verification not implemented)	931

3.175.1 Optimal result

Integrand size = 13, antiderivative size = 7

$$\int \cos^{\cos(x)}(x)(1 + \log(\cos(x))) \sin(x) dx = -\cos^{\cos(x)}(x)$$

output

`-cos(x)^cos(x)`

3.175.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \cos^{\cos(x)}(x)(1 + \log(\cos(x))) \sin(x) dx = -\cos^{\cos(x)}(x)$$

input

`Integrate[Cos[x]^Cos[x]*(1 + Log[Cos[x]])*Sin[x],x]`

output

`-Cos[x]^Cos[x]`

3.175.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {4835, 7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin(x) \cos^{\cos(x)}(x) (\log(\cos(x)) + 1) dx \\ & \quad \downarrow \text{4835} \\ & - \int \cos^{\cos(x)}(x) (\log(\cos(x)) + 1) d \cos(x) \\ & \quad \downarrow \text{7293} \\ & - \int \left(\log(\cos(x)) \cos^{\cos(x)}(x) + \cos^{\cos(x)}(x) \right) d \cos(x) \\ & \quad \downarrow \text{2009} \\ & - \cos^{\cos(x)}(x) \end{aligned}$$

input `Int[Cos[x]^Cos[x]*(1 + Log[Cos[x]])*Sin[x],x]`

output `-Cos[x]^Cos[x]`

3.175.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 4835 `Int[(u_)*(F_)[(c_.)*((a_.) + (b_.)*(x_))], x_Symbol] := With[{d = FreeFactors[Cos[c*(a + b*x)], x]}, Simp[-d/(b*c) Subst[Int[SubstFor[1, Cos[c*(a + b*x)]]/d, u, x], x], x, Cos[c*(a + b*x)]/d, x] /; FunctionOfQ[Cos[c*(a + b*x)]/d, u, x, True]] /; FreeQ[{a, b, c}, x] && (EqQ[F, Sin] || EqQ[F, sin])`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

3.175.4 Maple [F]

$$\int \cos(x)^{1+\cos(x)} \tan(x) (1 + \ln(\cos(x))) dx$$

input `int(cos(x)^(1+cos(x))*tan(x)*(1+ln(cos(x))),x)`

output `int(cos(x)^(1+cos(x))*tan(x)*(1+ln(cos(x))),x)`

3.175.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.86

$$\int \cos^{\cos(x)}(x)(1 + \log(\cos(x))) \sin(x) dx = -\frac{\cos(x)^{\cos(x)+1}}{\cos(x)}$$

input `integrate(cos(x)^(1+cos(x))*tan(x)*(1+log(cos(x))),x, algorithm="fricas")`

output `-cos(x)^(cos(x) + 1)/cos(x)`

3.175.6 Sympy [F]

$$\int \cos^{\cos(x)}(x)(1 + \log(\cos(x))) \sin(x) dx = \int (\log(\cos(x)) + 1) \cos^{\cos(x)+1}(x) \tan(x) dx$$

input `integrate(cos(x)**(1+cos(x))*tan(x)*(1+ln(cos(x))),x)`

output `Integral((log(cos(x)) + 1)*cos(x)**(cos(x) + 1)*tan(x), x)`

3.175.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \cos^{\cos(x)}(x)(1 + \log(\cos(x))) \sin(x) dx = -\cos(x)^{\cos(x)}$$

input `integrate(cos(x)^(1+cos(x))*tan(x)*(1+log(cos(x))),x, algorithm="maxima")`output `-cos(x)^cos(x)`**3.175.8 Giac [F]**

$$\int \cos^{\cos(x)}(x)(1 + \log(\cos(x))) \sin(x) dx = \int \cos(x)^{\cos(x)+1} (\log(\cos(x)) + 1) \tan(x) dx$$

input `integrate(cos(x)^(1+cos(x))*tan(x)*(1+log(cos(x))),x, algorithm="giac")`output `integrate(cos(x)^(cos(x) + 1)*(log(cos(x)) + 1)*tan(x), x)`**3.175.9 Mupad [B] (verification not implemented)**

Time = 17.65 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \cos^{\cos(x)}(x)(1 + \log(\cos(x))) \sin(x) dx = -\cos(x)^{\cos(x)}$$

input `int(cos(x)^(cos(x) + 1)*tan(x)*(log(cos(x)) + 1),x)`output `-cos(x)^cos(x)`

3.176 $\int \frac{e^x}{2+e^x} dx$

3.176.1 Optimal result	932
3.176.2 Mathematica [A] (verified)	932
3.176.3 Rubi [A] (verified)	933
3.176.4 Maple [A] (verified)	934
3.176.5 Fricas [A] (verification not implemented)	934
3.176.6 Sympy [A] (verification not implemented)	934
3.176.7 Maxima [A] (verification not implemented)	935
3.176.8 Giac [A] (verification not implemented)	935
3.176.9 Mupad [B] (verification not implemented)	935

3.176.1 Optimal result

Integrand size = 11, antiderivative size = 6

$$\int \frac{e^x}{2+e^x} dx = \log(2+e^x)$$

output `ln(exp(x)+2)`

3.176.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{2+e^x} dx = \log(2+e^x)$$

input `Integrate[E^x/(2 + E^x), x]`

output `Log[2 + E^x]`

3.176.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2676, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^x}{e^x + 2} dx$$

↓ 2676

$$\int \frac{1}{e^x + 2} de^x$$

↓ 16

$$\log(e^x + 2)$$

input `Int[E^x/(2 + E^x), x]`

output `Log[2 + E^x]`

3.176.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 2676 `Int[((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)*((a_.) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.))^p_., x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[(a + b*x)^p, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n, p}, x]`

3.176.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

method	result	size
derivativdivides	$\ln(e^x + 2)$	6
default	$\ln(e^x + 2)$	6
norman	$\ln(e^x + 2)$	6
risch	$\ln(e^x + 2)$	6
parallelrisc	$\ln(e^x + 2)$	6

input `int(exp(x)/(exp(x)+2),x,method=_RETURNVERBOSE)`

output `ln(exp(x)+2)`

3.176.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{2 + e^x} dx = \log(e^x + 2)$$

input `integrate(exp(x)/(exp(x)+2),x, algorithm="fricas")`

output `log(e^x + 2)`

3.176.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{2 + e^x} dx = \log(e^x + 2)$$

input `integrate(exp(x)/(exp(x)+2),x)`

output `log(exp(x) + 2)`

3.176.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{2 + e^x} dx = \log(e^x + 2)$$

input `integrate(exp(x)/(exp(x)+2),x, algorithm="maxima")`output `log(e^x + 2)`**3.176.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{2 + e^x} dx = \log(e^x + 2)$$

input `integrate(exp(x)/(exp(x)+2),x, algorithm="giac")`output `log(e^x + 2)`**3.176.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int \frac{e^x}{2 + e^x} dx = \ln(e^x + 2)$$

input `int(exp(x)/(exp(x) + 2),x)`output `log(exp(x) + 2)`

3.177 $\int \sin(2018x) dx$

3.177.1 Optimal result	936
3.177.2 Mathematica [A] (verified)	936
3.177.3 Rubi [A] (verified)	937
3.177.4 Maple [A] (verified)	938
3.177.5 Fricas [A] (verification not implemented)	938
3.177.6 Sympy [A] (verification not implemented)	938
3.177.7 Maxima [A] (verification not implemented)	939
3.177.8 Giac [A] (verification not implemented)	939
3.177.9 Mupad [B] (verification not implemented)	939

3.177.1 Optimal result

Integrand size = 4, antiderivative size = 8

$$\int \sin(2018x) dx = -\frac{\cos(2018x)}{2018}$$

output `-1/2018*cos(2018*x)`

3.177.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \sin(2018x) dx = -\frac{\cos(2018x)}{2018}$$

input `Integrate[Sin[2018*x],x]`

output `-1/2018*Cos[2018*x]`

3.177.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sin(2018x) dx \\ \downarrow \text{3042} \\ \int \sin(2018x) dx \\ \downarrow \text{3118} \\ -\frac{\cos(2018x)}{2018} \end{array}$$

input `Int[Sin[2018*x],x]`

output `-1/2018*Cos[2018*x]`

3.177.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

3.177.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$-\frac{\cos(2018x)}{2018}$	7
default	$-\frac{\cos(2018x)}{2018}$	7
risch	$-\frac{\cos(2018x)}{2018}$	7
parallelrisch	$-\frac{\cos(2018x)}{2018} - \frac{1}{2018}$	9
norman	$-\frac{1}{1009(1+\tan(1009x)^2)}$	13
meijerg	$\frac{\sqrt{\pi} \left(\frac{1}{\sqrt{\pi}} - \frac{\cos(2018x)}{\sqrt{\pi}} \right)}{2018}$	19

input `int(sin(2018*x),x,method=_RETURNVERBOSE)`output `-1/2018*cos(2018*x)`**3.177.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sin(2018x) dx = -\frac{1}{2018} \cos(2018x)$$

input `integrate(sin(2018*x),x, algorithm="fricas")`output `-1/2018*cos(2018*x)`**3.177.6 Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \sin(2018x) dx = -\frac{\cos(2018x)}{2018}$$

input `integrate(sin(2018*x),x)`

output `-cos(2018*x)/2018`

3.177.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sin(2018x) dx = -\frac{1}{2018} \cos(2018x)$$

input `integrate(sin(2018*x),x, algorithm="maxima")`

output `-1/2018*cos(2018*x)`

3.177.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sin(2018x) dx = -\frac{1}{2018} \cos(2018x)$$

input `integrate(sin(2018*x),x, algorithm="giac")`

output `-1/2018*cos(2018*x)`

3.177.9 Mupad [B] (verification not implemented)

Time = 13.96 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sin(2018x) dx = -\frac{\cos(2018x)}{2018}$$

input `int(sin(2018*x),x)`

output `-cos(2018*x)/2018`

3.178 $\int \frac{1}{\cot(x)+\tan(x)} dx$

3.178.1 Optimal result	940
3.178.2 Mathematica [A] (verified)	940
3.178.3 Rubi [A] (verified)	941
3.178.4 Maple [A] (verified)	942
3.178.5 Fricas [B] (verification not implemented)	942
3.178.6 Sympy [A] (verification not implemented)	943
3.178.7 Maxima [A] (verification not implemented)	943
3.178.8 Giac [A] (verification not implemented)	943
3.178.9 Mupad [B] (verification not implemented)	944

3.178.1 Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \frac{1}{\cot(x) + \tan(x)} dx = \frac{\sin^2(x)}{2}$$

output `1/2*sin(x)^2`

3.178.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{1}{\cot(x) + \tan(x)} dx = -\frac{1}{2} \cos^2(x)$$

input `Integrate[(Cot[x] + Tan[x])^(-1),x]`

output `-1/2*Cos[x]^2`

3.178.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.50, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 4853, 241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\tan(x) + \cot(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{\tan(x) + \cot(x)} dx \\ & \quad \downarrow \text{4853} \\ & \int \frac{\tan(x)}{(\tan^2(x) + 1)^2} d \tan(x) \\ & \quad \downarrow \text{241} \\ & -\frac{1}{2(\tan^2(x) + 1)} \end{aligned}$$

input `Int[(Cot[x] + Tan[x])^(-1),x]`

output `-1/2*1/(1 + Tan[x]^2)`

3.178.3.1 Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4853 Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Tan[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x^2), Tan[v]/d, u, x], x], x, Tan[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Tan[v], x], u, x, True] && TryPureTanSubst[ActivateTrig[u], x]]
```

3.178.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
default	$-\frac{\cos(x)^2}{2}$	7
risch	$-\frac{\cos(2x)}{4}$	7
parallelrisch	$-\frac{\cos(2x)}{4} - \frac{1}{4}$	9
norman	$-\frac{1}{2(1+\tan(x)^2)}$	11

input `int(1/(tan(x)+cot(x)),x,method=_RETURNVERBOSE)`

output `-1/2*cos(x)^2`

3.178.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 16 vs. 2(6) = 12.

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 2.00

$$\int \frac{1}{\cot(x) + \tan(x)} dx = \frac{\tan(x)^2 - 1}{4(\tan(x)^2 + 1)}$$

input `integrate(1/(tan(x)+cot(x)),x, algorithm="fracas")`

output `1/4*(tan(x)^2 - 1)/(tan(x)^2 + 1)`

3.178.6 Sympy [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.25

$$\int \frac{1}{\cot(x) + \tan(x)} dx = -\frac{1}{2 \tan^2(x) + 2}$$

input `integrate(1/(tan(x)+cot(x)),x)`output `-1/(2*tan(x)**2 + 2)`**3.178.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.25

$$\int \frac{1}{\cot(x) + \tan(x)} dx = -\frac{1}{2 (\tan(x)^2 + 1)}$$

input `integrate(1/(tan(x)+cot(x)),x, algorithm="maxima")`output `-1/2/(tan(x)^2 + 1)`**3.178.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{\cot(x) + \tan(x)} dx = -\frac{1}{2} \cos(x)^2$$

input `integrate(1/(tan(x)+cot(x)),x, algorithm="giac")`output `-1/2*cos(x)^2`

3.178.9 Mupad [B] (verification not implemented)

Time = 13.52 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{\cot(x) + \tan(x)} dx = \frac{\sin(x)^2}{2}$$

input `int(1/(cot(x) + tan(x)),x)`

output `sin(x)^2/2`

3.179 $\int \frac{x^5}{2+x^{12}} dx$

3.179.1 Optimal result	945
3.179.2 Mathematica [A] (verified)	945
3.179.3 Rubi [A] (verified)	946
3.179.4 Maple [A] (verified)	947
3.179.5 Fricas [A] (verification not implemented)	947
3.179.6 Sympy [A] (verification not implemented)	947
3.179.7 Maxima [A] (verification not implemented)	948
3.179.8 Giac [A] (verification not implemented)	948
3.179.9 Mupad [B] (verification not implemented)	948

3.179.1 Optimal result

Integrand size = 11, antiderivative size = 19

$$\int \frac{x^5}{2+x^{12}} dx = \frac{\arctan\left(\frac{x^6}{\sqrt{2}}\right)}{6\sqrt{2}}$$

output `1/12*arctan(1/2*2^(1/2)*x^6)*2^(1/2)`

3.179.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{x^5}{2+x^{12}} dx = \frac{\arctan\left(\frac{x^6}{\sqrt{2}}\right)}{6\sqrt{2}}$$

input `Integrate[x^5/(2 + x^12),x]`

output `ArcTan[x^6/Sqrt[2]]/(6*Sqrt[2])`

3.179.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {807, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^5}{x^{12} + 2} dx$$

↓ 807

$$\frac{1}{6} \int \frac{1}{x^{12} + 2} dx^6$$

↓ 216

$$\frac{\arctan\left(\frac{x^6}{\sqrt{2}}\right)}{6\sqrt{2}}$$

input `Int[x^5/(2 + x^12), x]`

output `ArcTan[x^6/Sqrt[2]]/(6*Sqrt[2])`

3.179.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

3.179.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{\arctan\left(\frac{x^6\sqrt{2}}{2}\right)\sqrt{2}}{12}$	15
meijerg	$\frac{\arctan\left(\frac{x^6\sqrt{2}}{2}\right)\sqrt{2}}{12}$	15
risch	$\frac{\arctan\left(\frac{x^6\sqrt{2}}{2}\right)\sqrt{2}}{12}$	15

input `int(x^5/(x^12+2),x,method=_RETURNVERBOSE)`output `1/12*arctan(1/2*x^6*2^(1/2))*2^(1/2)`**3.179.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int \frac{x^5}{2+x^{12}} dx = \frac{1}{12} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2} x^6\right)$$

input `integrate(x^5/(x^12+2),x, algorithm="fricas")`output `1/12*sqrt(2)*arctan(1/2*sqrt(2)*x^6)`**3.179.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{x^5}{2+x^{12}} dx = \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}x^6}{2}\right)}{12}$$

input `integrate(x**5/(x**12+2),x)`output `sqrt(2)*atan(sqrt(2)*x**6/2)/12`

3.179.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int \frac{x^5}{2+x^{12}} dx = \frac{1}{12} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} x^6 \right)$$

input `integrate(x^5/(x^12+2),x, algorithm="maxima")`output `1/12*sqrt(2)*arctan(1/2*sqrt(2)*x^6)`**3.179.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int \frac{x^5}{2+x^{12}} dx = \frac{1}{12} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} x^6 \right)$$

input `integrate(x^5/(x^12+2),x, algorithm="giac")`output `1/12*sqrt(2)*arctan(1/2*sqrt(2)*x^6)`**3.179.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int \frac{x^5}{2+x^{12}} dx = \frac{\sqrt{2} \operatorname{atan} \left(\frac{\sqrt{2} x^6}{2} \right)}{12}$$

input `int(x^5/(x^12 + 2),x)`output `(2^(1/2)*atan((2^(1/2)*x^6)/2))/12`

3.180 $\int (\cos(x) \cosh(x) + \sin(x) \sinh(x)) dx$

3.180.1 Optimal result	949
3.180.2 Mathematica [A] (verified)	949
3.180.3 Rubi [F]	950
3.180.4 Maple [B] (verified)	950
3.180.5 Fricas [B] (verification not implemented)	951
3.180.6 Sympy [A] (verification not implemented)	951
3.180.7 Maxima [B] (verification not implemented)	951
3.180.8 Giac [B] (verification not implemented)	952
3.180.9 Mupad [B] (verification not implemented)	952

3.180.1 Optimal result

Integrand size = 11, antiderivative size = 5

$$\int (\cos(x) \cosh(x) + \sin(x) \sinh(x)) dx = \cosh(x) \sin(x)$$

output `cosh(x)*sin(x)`

3.180.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int (\cos(x) \cosh(x) + \sin(x) \sinh(x)) dx = \cosh(x) \sin(x)$$

input `Integrate[Cos[x]*Cosh[x] + Sin[x]*Sinh[x],x]`

output `Cosh[x]*Sin[x]`

3.180.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\sin(x) \sinh(x) + \cos(x) \cosh(x)) dx$$

$$\downarrow \text{2009}$$

$$\int \sin(x) \sinh(x) dx + \int \cos(x) \cosh(x) dx$$

input `Int[Cos[x]*Cosh[x] + Sin[x]*Sinh[x],x]`

output `$Aborted`

3.180.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.180.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(5) = 10$.

Time = 0.87 (sec) , antiderivative size = 16, normalized size of antiderivative = 3.20

method	result
default	$\frac{e^x \sin(x)}{2} + \frac{e^{-x} \sin(x)}{2}$
parts	$\frac{e^x \sin(x)}{2} + \frac{e^{-x} \sin(x)}{2}$
risch	$\frac{ie^{(1-i)x}}{4} + \frac{ie^{(-1-i)x}}{4} - \frac{ie^{(1+i)x}}{4} - \frac{ie^{(-1+i)x}}{4}$
meijerg	$\pi^{\frac{3}{2}} \left(\frac{e^x \cos(x)}{4\pi^{\frac{3}{2}}} + \frac{e^x \sin(x)}{4\pi^{\frac{3}{2}}} - \frac{e^{-x} \cos(x)}{4\pi^{\frac{3}{2}}} + \frac{e^{-x} \sin(x)}{4\pi^{\frac{3}{2}}} \right) + \pi^{\frac{3}{2}} \left(-\frac{e^x \cos(x)}{4\pi^{\frac{3}{2}}} + \frac{e^x \sin(x)}{4\pi^{\frac{3}{2}}} + \frac{e^{-x} \cos(x)}{4\pi^{\frac{3}{2}}} + \frac{e^{-x} \sin(x)}{4\pi^{\frac{3}{2}}} \right)$

input `int(cos(x)*cosh(x)+sin(x)*sinh(x),x,method=_RETURNVERBOSE)`

output `1/2*exp(x)*sin(x)+1/2*exp(-x)*sin(x)`

3.180.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. $2(5) = 10$.

Time = 0.25 (sec) , antiderivative size = 34, normalized size of antiderivative = 6.80

$$\int (\cos(x) \cosh(x) + \sin(x) \sinh(x)) dx$$

$$= \frac{2 \cosh(x) \sin(x) \sinh(x) + \sin(x) \sinh(x)^2 + (\cosh(x)^2 + 1) \sin(x)}{2 (\cosh(x) + \sinh(x))}$$

input `integrate(cos(x)*cosh(x)+sin(x)*sinh(x),x, algorithm="fricas")`

output `1/2*(2*cosh(x)*sin(x)*sinh(x) + sin(x)*sinh(x)^2 + (cosh(x)^2 + 1)*sin(x))
/(cosh(x) + sinh(x))`

3.180.6 Sympy [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int (\cos(x) \cosh(x) + \sin(x) \sinh(x)) dx = \sin(x) \cosh(x)$$

input `integrate(cos(x)*cosh(x)+sin(x)*sinh(x),x)`

output `sin(x)*cosh(x)`

3.180.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 52 vs. $2(5) = 10$.

Time = 0.22 (sec) , antiderivative size = 52, normalized size of antiderivative = 10.40

$$\int (\cos(x) \cosh(x) + \sin(x) \sinh(x)) dx = \frac{1}{4} ((e^{2x} - 1) \cos(x) + (e^{2x} + 1) \sin(x)) e^{-x}$$

$$- \frac{1}{4} ((e^{2x} - 1) \cos(x) - (e^{2x} + 1) \sin(x)) e^{-x}$$

input `integrate(cos(x)*cosh(x)+sin(x)*sinh(x),x, algorithm="maxima")`

output `1/4*((e^(2*x) - 1)*cos(x) + (e^(2*x) + 1)*sin(x))*e^(-x) - 1/4*((e^(2*x) - 1)*cos(x) - (e^(2*x) + 1)*sin(x))*e^(-x)`

3.180.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 45 vs. $2(5) = 10$.

Time = 0.27 (sec) , antiderivative size = 45, normalized size of antiderivative = 9.00

$$\int (\cos(x) \cosh(x) + \sin(x) \sinh(x)) dx = \frac{1}{4} (\cos(x) + \sin(x))e^{(-x)} - \frac{1}{4} (\cos(x) - \sin(x))e^{(-x)} + \frac{1}{4} (\cos(x) + \sin(x))e^x - \frac{1}{4} (\cos(x) - \sin(x))e^x$$

input `integrate(cos(x)*cosh(x)+sin(x)*sinh(x),x, algorithm="giac")`

output `1/4*(cos(x) + sin(x))*e^(-x) - 1/4*(cos(x) - sin(x))*e^(-x) + 1/4*(cos(x) + sin(x))*e^x - 1/4*(cos(x) - sin(x))*e^x`

3.180.9 Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int (\cos(x) \cosh(x) + \sin(x) \sinh(x)) dx = \cosh(x) \sin(x)$$

input `int(cos(x)*cosh(x) + sin(x)*sinh(x),x)`

output `cosh(x)*sin(x)`

$$\mathbf{3.181} \quad \int \frac{e^x + \cos(x)}{e^x + \sin(x)} dx$$

3.181.1 Optimal result	953
3.181.2 Mathematica [A] (verified)	953
3.181.3 Rubi [A] (verified)	954
3.181.4 Maple [A] (verified)	954
3.181.5 Fricas [A] (verification not implemented)	955
3.181.6 Sympy [A] (verification not implemented)	955
3.181.7 Maxima [A] (verification not implemented)	955
3.181.8 Giac [B] (verification not implemented)	956
3.181.9 Mupad [B] (verification not implemented)	956

3.181.1 Optimal result

Integrand size = 15, antiderivative size = 7

$$\int \frac{e^x + \cos(x)}{e^x + \sin(x)} dx = \log(e^x + \sin(x))$$

output `ln(exp(x)+sin(x))`

3.181.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{e^x + \cos(x)}{e^x + \sin(x)} dx = \log(e^x + \sin(x))$$

input `Integrate[(E^x + Cos[x])/(E^x + Sin[x]),x]`

output `Log[E^x + Sin[x]]`

3.181.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$, Rules used = {7235}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^x + \cos(x)}{e^x + \sin(x)} dx$$

↓ 7235

$$\log(e^x + \sin(x))$$

input `Int[(E^x + Cos[x])/(E^x + Sin[x]), x]`

output `Log[E^x + Sin[x]]`

3.181.3.1 Defintions of rubi rules used

rule 7235 `Int[(u_)/(y_), x_Symbol] := With[{q = DerivativeDivides[y, u, x]}, Simp[q*Log[RemoveContent[y, x]], x] /; !FalseQ[q]]`

3.181.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

method	result	size
derivativedivides	$\ln(e^x + \sin(x))$	7
default	$\ln(e^x + \sin(x))$	7
risch	$-ix + \ln(e^{2ix} + 2ie^{(1+i)x} - 1)$	23
parallelrisch	$-\ln\left(\frac{1}{1+\cos(x)}\right) + \ln\left(\frac{e^x + \sin(x)}{1+\cos(x)}\right)$	24
norman	$-\ln\left(1 + \tan\left(\frac{x}{2}\right)^2\right) + \ln\left(e^x \tan\left(\frac{x}{2}\right)^2 + e^x + 2 \tan\left(\frac{x}{2}\right)\right)$	32

input `int((exp(x)+cos(x))/(exp(x)+sin(x)), x, method=_RETURNVERBOSE)`

output `ln(exp(x)+sin(x))`

3.181.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int \frac{e^x + \cos(x)}{e^x + \sin(x)} dx = \log(e^x + \sin(x))$$

input `integrate((exp(x)+cos(x))/(exp(x)+sin(x)),x, algorithm="fricas")`

output `log(e^x + sin(x))`

3.181.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{e^x + \cos(x)}{e^x + \sin(x)} dx = \log(e^x + \sin(x))$$

input `integrate((exp(x)+cos(x))/(exp(x)+sin(x)),x)`

output `log(exp(x) + sin(x))`

3.181.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int \frac{e^x + \cos(x)}{e^x + \sin(x)} dx = \log(e^x + \sin(x))$$

input `integrate((exp(x)+cos(x))/(exp(x)+sin(x)),x, algorithm="maxima")`

output `log(e^x + sin(x))`

3.181.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 77 vs. $2(6) = 12$.

Time = 0.28 (sec) , antiderivative size = 77, normalized size of antiderivative = 11.00

$$\int \frac{e^x + \cos(x)}{e^x + \sin(x)} dx$$

$$= \frac{1}{2} \log \left(\frac{4 \left(e^{(2x)} \tan \left(\frac{1}{2} x \right)^4 + 4 e^x \tan \left(\frac{1}{2} x \right)^3 + 2 e^{(2x)} \tan \left(\frac{1}{2} x \right)^2 + 4 e^x \tan \left(\frac{1}{2} x \right) + 4 \tan \left(\frac{1}{2} x \right)^2 + e^{(2x)} \right)}{\tan \left(\frac{1}{2} x \right)^4 + 2 \tan \left(\frac{1}{2} x \right)^2 + 1} \right)$$

input `integrate((exp(x)+cos(x))/(exp(x)+sin(x)),x, algorithm="giac")`

output `1/2*log(4*(e^(2*x))*tan(1/2*x)^4 + 4*e^x*tan(1/2*x)^3 + 2*e^(2*x)*tan(1/2*x)^2 + 4*e^x*tan(1/2*x) + 4*tan(1/2*x)^2 + e^(2*x))/(tan(1/2*x)^4 + 2*tan(1/2*x)^2 + 1))`

3.181.9 Mupad [B] (verification not implemented)

Time = 14.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int \frac{e^x + \cos(x)}{e^x + \sin(x)} dx = \ln(e^x + \sin(x))$$

input `int((cos(x) + exp(x))/(exp(x) + sin(x)),x)`

output `log(exp(x) + sin(x))`

3.182 $\int \cos(x) \sin(\cos(\sin(x))) \sin(\sin(x)) dx$

3.182.1 Optimal result	957
3.182.2 Mathematica [A] (verified)	957
3.182.3 Rubi [A] (verified)	958
3.182.4 Maple [A] (verified)	959
3.182.5 Fricas [B] (verification not implemented)	959
3.182.6 Sympy [A] (verification not implemented)	960
3.182.7 Maxima [A] (verification not implemented)	960
3.182.8 Giac [A] (verification not implemented)	960
3.182.9 Mupad [B] (verification not implemented)	961

3.182.1 Optimal result

Integrand size = 10, antiderivative size = 4

$$\int \cos(x) \sin(\cos(\sin(x))) \sin(\sin(x)) dx = \cos(\cos(\sin(x)))$$

output `cos(cos(sin(x)))`

3.182.2 Mathematica [A] (verified)

Time = 9.25 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \cos(x) \sin(\cos(\sin(x))) \sin(\sin(x)) dx = \cos(\cos(\sin(x)))$$

input `Integrate[Cos[x]*Sin[Cos[Sin[x]]]*Sin[Sin[x]],x]`

output `Cos[Cos[Sin[x]]]`

3.182.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {4834, 4835, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(\sin(x)) \cos(x) \sin(\cos(\sin(x))) dx \\
 & \quad \downarrow \text{4834} \\
 & \int \sin(\sin(x)) \sin(\cos(\sin(x))) d \sin(x) \\
 & \quad \downarrow \text{4835} \\
 & - \int \sin(\cos(\sin(x))) d \cos(\sin(x)) \\
 & \quad \downarrow \text{3042} \\
 & - \int \sin(\cos(\sin(x))) d \cos(\sin(x)) \\
 & \quad \downarrow \text{3118} \\
 & \cos(\cos(\sin(x)))
 \end{aligned}$$

input `Int[Cos[x]*Sin[Cos[Sin[x]]]*Sin[Sin[x]],x]`

output `Cos[Cos[Sin[x]]]`

3.182.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

```
rule 4834 Int[(u_)*(F_)[(c_.)*((a_.) + (b_.)*(x_))], x_Symbol] := With[{d = FreeFactors[Sin[c*(a + b*x)], x]}, Simp[d/(b*c) Subst[Int[SubstFor[1, Sin[c*(a + b*x)]]/d, u, x], x], x, Sin[c*(a + b*x)]/d, x] /; FunctionOfQ[Sin[c*(a + b*x)]/d, u, x, True]] /; FreeQ[{a, b, c}, x] && (EqQ[F, Cos] || EqQ[F, cos])
```

```
rule 4835 Int[(u_)*(F_)[(c_.)*((a_.) + (b_.)*(x_))], x_Symbol] := With[{d = FreeFactors[Cos[c*(a + b*x)], x]}, Simp[-d/(b*c) Subst[Int[SubstFor[1, Cos[c*(a + b*x)]]/d, u, x], x], x, Cos[c*(a + b*x)]/d, x] /; FunctionOfQ[Cos[c*(a + b*x)]/d, u, x, True]] /; FreeQ[{a, b, c}, x] && (EqQ[F, Sin] || EqQ[F, sin])
```

3.182.4 Maple [A] (verified)

Time = 8.37 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

method	result	size
derivativedivides	$\cos(\cos(\sin(x)))$	5
default	$\cos(\cos(\sin(x)))$	5
risch	$\cos(\cos(\sin(x)))$	5
parallelrisch	$-1 + \cos(\cos(\sin(x)))$	7

```
input int(sin(cos(sin(x)))*sin(sin(x))*cos(x),x,method=_RETURNVERBOSE)
```

```
output cos(cos(sin(x)))
```

3.182.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 44 vs. $2(4) = 8$.

Time = 0.26 (sec) , antiderivative size = 44, normalized size of antiderivative = 11.00

$$\int \cos(x) \sin(\cos(\sin(x))) \sin(\sin(x)) dx = \cos \left(\frac{\tan \left(\frac{\tan(\frac{1}{2}x)}{\tan(\frac{1}{2}x)^2 + 1} \right)^2 - 1}{\tan \left(\frac{\tan(\frac{1}{2}x)}{\tan(\frac{1}{2}x)^2 + 1} \right)^2 + 1} \right)$$

```
input integrate(sin(cos(sin(x)))*sin(sin(x))*cos(x),x, algorithm="fricas")
```


output `cos((tan(tan(1/2*x)/(tan(1/2*x)^2 + 1))^2 - 1)/(tan(tan(1/2*x)/(tan(1/2*x)^2 + 1))^2 + 1))`

3.182.6 Sympy [A] (verification not implemented)

Time = 0.90 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

$$\int \cos(x) \sin(\cos(\sin(x))) \sin(\sin(x)) dx = \cos(\cos(\sin(x)))$$

input `integrate(sin(cos(sin(x)))*sin(sin(x))*cos(x),x)`

output `cos(cos(sin(x)))`

3.182.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \cos(x) \sin(\cos(\sin(x))) \sin(\sin(x)) dx = \cos(\cos(\sin(x)))$$

input `integrate(sin(cos(sin(x)))*sin(sin(x))*cos(x),x, algorithm="maxima")`

output `cos(cos(sin(x)))`

3.182.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \cos(x) \sin(\cos(\sin(x))) \sin(\sin(x)) dx = \cos(\cos(\sin(x)))$$

input `integrate(sin(cos(sin(x)))*sin(sin(x))*cos(x),x, algorithm="giac")`

output `cos(cos(sin(x)))`

3.182.9 Mupad [B] (verification not implemented)

Time = 14.97 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \cos(x) \sin(\cos(\sin(x))) \sin(\sin(x)) dx = \cos(\cos(\sin(x)))$$

input `int(sin(sin(x))*sin(cos(sin(x)))*cos(x),x)`

output `cos(cos(sin(x)))`

3.183 $\int \frac{1}{1+\sin(x)} dx$

3.183.1 Optimal result	962
3.183.2 Mathematica [B] (verified)	962
3.183.3 Rubi [A] (verified)	963
3.183.4 Maple [A] (verified)	964
3.183.5 Fricas [A] (verification not implemented)	964
3.183.6 Sympy [A] (verification not implemented)	964
3.183.7 Maxima [A] (verification not implemented)	965
3.183.8 Giac [A] (verification not implemented)	965
3.183.9 Mupad [B] (verification not implemented)	965

3.183.1 Optimal result

Integrand size = 6, antiderivative size = 10

$$\int \frac{1}{1+\sin(x)} dx = -\frac{\cos(x)}{1+\sin(x)}$$

output `-cos(x)/(1+sin(x))`

3.183.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 23 vs. $2(10) = 20$.

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 2.30

$$\int \frac{1}{1+\sin(x)} dx = \frac{2 \sin\left(\frac{x}{2}\right)}{\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)}$$

input `Integrate[(1 + Sin[x])^(-1),x]`

output `(2*Sin[x/2])/(Cos[x/2] + Sin[x/2])`

3.183.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sin(x) + 1} dx$$

↓ 3042

$$\int \frac{1}{\sin(x) + 1} dx$$

↓ 3127

$$-\frac{\cos(x)}{\sin(x) + 1}$$

input `Int[(1 + Sin[x])^(-1),x]`

output `-(Cos[x]/(1 + Sin[x]))`

3.183.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

3.183.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

method	result	size
default	$-\frac{2}{1+\tan(\frac{x}{2})}$	11
norman	$-\frac{2}{1+\tan(\frac{x}{2})}$	11
parallelrisch	$-\frac{2}{1+\tan(\frac{x}{2})}$	11
risch	$-\frac{2}{i+e^{ix}}$	13

input `int(1/(1+sin(x)),x,method=_RETURNVERBOSE)`output `-2/(1+tan(1/2*x))`**3.183.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.80

$$\int \frac{1}{1 + \sin(x)} dx = -\frac{\cos(x) - \sin(x) + 1}{\cos(x) + \sin(x) + 1}$$

input `integrate(1/(1+sin(x)),x, algorithm="fricas")`output `-(cos(x) - sin(x) + 1)/(cos(x) + sin(x) + 1)`**3.183.6 Sympy [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{1}{1 + \sin(x)} dx = -\frac{2}{\tan\left(\frac{x}{2}\right) + 1}$$

input `integrate(1/(1+sin(x)),x)`output `-2/(tan(x/2) + 1)`

3.183.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.50

$$\int \frac{1}{1 + \sin(x)} dx = -\frac{2}{\frac{\sin(x)}{\cos(x)+1} + 1}$$

input `integrate(1/(1+sin(x)),x, algorithm="maxima")`output `-2/(sin(x)/(cos(x) + 1) + 1)`**3.183.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \sin(x)} dx = -\frac{2}{\tan\left(\frac{1}{2}x\right) + 1}$$

input `integrate(1/(1+sin(x)),x, algorithm="giac")`output `-2/(tan(1/2*x) + 1)`**3.183.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \sin(x)} dx = -\frac{2}{\tan\left(\frac{x}{2}\right) + 1}$$

input `int(1/(sin(x) + 1),x)`output `-2/(tan(x/2) + 1)`

3.184 $\int \frac{\cos(x)}{1-\cos(2x)} dx$

3.184.1 Optimal result	966
3.184.2 Mathematica [A] (verified)	966
3.184.3 Rubi [A] (verified)	967
3.184.4 Maple [A] (verified)	968
3.184.5 Fricas [A] (verification not implemented)	968
3.184.6 Sympy [F]	969
3.184.7 Maxima [B] (verification not implemented)	969
3.184.8 Giac [A] (verification not implemented)	969
3.184.9 Mupad [B] (verification not implemented)	970

3.184.1 Optimal result

Integrand size = 13, antiderivative size = 6

$$\int \frac{\cos(x)}{1-\cos(2x)} dx = -\frac{\csc(x)}{2}$$

output `-1/2*csc(x)`

3.184.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{1-\cos(2x)} dx = -\frac{\csc(x)}{2}$$

input `Integrate[Cos[x]/(1 - Cos[2*x]),x]`

output `-1/2*Csc[x]`

3.184.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3042, 4856, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(x)}{1 - \cos(2x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\cos(x)}{1 - \cos(2x)} dx \\ & \quad \downarrow \text{4856} \\ & \int \frac{\csc^2(x)}{2} d \sin(x) \\ & \quad \downarrow \text{15} \\ & -\frac{\csc(x)}{2} \end{aligned}$$

input `Int[Cos[x]/(1 - Cos[2*x]),x]`

output `-1/2*Csc[x]`

3.184.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`


```
rule 4856 Int[(u_)*(F_)[(c_.)*((a_.) + (b_.)*(x_))], x_Symbol] :> With[{d = FreeFactors[Sin[c*(a + b*x)], x]}, Simp[d/(b*c) Subst[Int[SubstFor[1, Sin[c*(a + b*x)]]/d, u, x], x], x, Sin[c*(a + b*x)]/d, x] /; FunctionOfQ[Sin[c*(a + b*x)]/d, u, x] /; FreeQ[{a, b, c}, x] && (EqQ[F, Cos] || EqQ[F, cos])
```

3.184.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

method	result	size
default	$-\frac{1}{2\sin(x)}$	7
risch	$-\frac{ie^{ix}}{e^{2ix}-1}$	18

```
input int(cos(x)/(1-cos(2*x)),x,method=_RETURNVERBOSE)
```

```
output -1/2/sin(x)
```

3.184.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{1 - \cos(2x)} dx = -\frac{1}{2\sin(x)}$$

```
input integrate(cos(x)/(1-cos(2*x)),x, algorithm="fricas")
```

```
output -1/2/sin(x)
```

3.184.6 Sympy [F]

$$\int \frac{\cos(x)}{1 - \cos(2x)} dx = - \int \frac{\cos(x)}{\cos(2x) - 1} dx$$

input `integrate(cos(x)/(1-cos(2*x)),x)`

output `-Integral(cos(x)/(cos(2*x) - 1), x)`

3.184.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. 2(4) = 8.

Time = 0.19 (sec) , antiderivative size = 42, normalized size of antiderivative = 7.00

$$\int \frac{\cos(x)}{1 - \cos(2x)} dx = - \frac{\cos(x) \sin(2x) - \cos(2x) \sin(x) + \sin(x)}{\cos(2x)^2 + \sin(2x)^2 - 2 \cos(2x) + 1}$$

input `integrate(cos(x)/(1-cos(2*x)),x, algorithm="maxima")`

output `-(cos(x)*sin(2*x) - cos(2*x)*sin(x) + sin(x))/(cos(2*x)^2 + sin(2*x)^2 - 2*cos(2*x) + 1)`

3.184.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{1 - \cos(2x)} dx = - \frac{1}{2 \sin(x)}$$

input `integrate(cos(x)/(1-cos(2*x)),x, algorithm="giac")`

output `-1/2/sin(x)`

3.184.9 Mupad [B] (verification not implemented)

Time = 15.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{1 - \cos(2x)} dx = -\frac{1}{2 \sin(x)}$$

input `int(-cos(x)/(cos(2*x) - 1),x)`

output `-1/(2*sin(x))`

3.185 $\int e^x \left(\frac{1}{x} + \log(x) \right) dx$

3.185.1 Optimal result	971
3.185.2 Mathematica [A] (verified)	971
3.185.3 Rubi [A] (verified)	972
3.185.4 Maple [A] (verified)	972
3.185.5 Fracas [A] (verification not implemented)	973
3.185.6 Sympy [A] (verification not implemented)	973
3.185.7 Maxima [A] (verification not implemented)	973
3.185.8 Giac [A] (verification not implemented)	974
3.185.9 Mupad [B] (verification not implemented)	974

3.185.1 Optimal result

Integrand size = 10, antiderivative size = 6

$$\int e^x \left(\frac{1}{x} + \log(x) \right) dx = e^x \log(x)$$

output `exp(x)*ln(x)`

3.185.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int e^x \left(\frac{1}{x} + \log(x) \right) dx = e^x \log(x)$$

input `Integrate[E^x*(x^(-1) + Log[x]),x]`

output `E^x*Log[x]`

3.185.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2726}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x \left(\frac{1}{x} + \log(x) \right) dx$$

↓ 2726

$$e^x \log(x)$$

input `Int[E^x*(x^(-1) + Log[x]),x]`

output `E^x*Log[x]`

3.185.3.1 Defintions of rubi rules used

rule 2726 `Int[(y_.)*(F_)^(u_)*((v_) + (w_)), x_Symbol] := With[{z = v*(y/(Log[F]*D[u, x]))}, Simp[F^u*z, x] /; EqQ[D[z, x], w*y]] /; FreeQ[F, x]`

3.185.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

method	result	size
norman	$e^x \ln(x)$	6
risch	$e^x \ln(x)$	6
parallelrisk	$e^x \ln(x)$	6

input `int(exp(x)*(1/x+ln(x)),x,method=_RETURNVERBOSE)`

output `exp(x)*ln(x)`

3.185.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int e^x \left(\frac{1}{x} + \log(x) \right) dx = e^x \log(x)$$

input `integrate(exp(x)*(1/x+log(x)),x, algorithm="fricas")`output `e^x*log(x)`**3.185.6 Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int e^x \left(\frac{1}{x} + \log(x) \right) dx = e^x \log(x)$$

input `integrate(exp(x)*(1/x+ln(x)),x)`output `exp(x)*log(x)`**3.185.7 Maxima [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int e^x \left(\frac{1}{x} + \log(x) \right) dx = e^x \log(x)$$

input `integrate(exp(x)*(1/x+log(x)),x, algorithm="maxima")`output `e^x*log(x)`

3.185.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int e^x \left(\frac{1}{x} + \log(x) \right) dx = e^x \log(x)$$

input `integrate(exp(x)*(1/x+log(x)),x, algorithm="giac")`output `e^x*log(x)`**3.185.9 Mupad [B] (verification not implemented)**

Time = 15.99 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int e^x \left(\frac{1}{x} + \log(x) \right) dx = e^x \ln(x)$$

input `int(exp(x)*(log(x) + 1/x),x)`output `exp(x)*log(x)`

3.186 $\int \tanh^2(x) dx$

3.186.1 Optimal result	975
3.186.2 Mathematica [A] (verified)	975
3.186.3 Rubi [A] (verified)	976
3.186.4 Maple [A] (verified)	977
3.186.5 Fricas [B] (verification not implemented)	977
3.186.6 Sympy [A] (verification not implemented)	978
3.186.7 Maxima [A] (verification not implemented)	978
3.186.8 Giac [A] (verification not implemented)	978
3.186.9 Mupad [B] (verification not implemented)	979

3.186.1 Optimal result

Integrand size = 4, antiderivative size = 6

$$\int \tanh^2(x) dx = x - \tanh(x)$$

output `x-tanh(x)`

3.186.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.33

$$\int \tanh^2(x) dx = \operatorname{arctanh}(\tanh(x)) - \tanh(x)$$

input `Integrate[Tanh[x]^2,x]`

output `ArcTanh[Tanh[x]] - Tanh[x]`

3.186.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 25, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \tanh^2(x) dx \\
 \downarrow \text{3042} \\
 \int -\tan(ix)^2 dx \\
 \downarrow \text{25} \\
 -\int \tan(ix)^2 dx \\
 \downarrow \text{3954} \\
 \int 1 dx - \tanh(x) \\
 \downarrow \text{24} \\
 x - \tanh(x)
 \end{array}$$

input `Int [Tanh [x]^2, x]`

output `x - Tanh [x]`

3.186.3.1 Defintions of rubi rules used

rule 24 `Int [a_, x_Symbol] :> Simp [a*x, x] /; FreeQ [a, x]`

rule 25 `Int [-(Fx_), x_Symbol] :> Simp [Identity [-1] Int [Fx, x], x]`

rule 3042 `Int [u_, x_Symbol] :> Int [DeactivateTrig [u, x], x] /; FunctionOfTrigOfLinear Q [u, x]`

```
rule 3954 Int[((b_.)*tan[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] :> Simp[b*((b*Tan[c + d
*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x]
, x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

3.186.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

method	result	size
parallelrisch	$x - \tanh(x)$	7
risch	$x + \frac{2}{e^{2x}+1}$	13
derivativedivides	$-\tanh(x) - \frac{\ln(\tanh(x)-1)}{2} + \frac{\ln(\tanh(x)+1)}{2}$	20
default	$-\tanh(x) - \frac{\ln(\tanh(x)-1)}{2} + \frac{\ln(\tanh(x)+1)}{2}$	20

```
input int(tanh(x)^2,x,method=_RETURNVERBOSE)
```

```
output x-tanh(x)
```

3.186.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 16 vs. $2(6) = 12$.

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 2.67

$$\int \tanh^2(x) dx = \frac{(x+1) \cosh(x) - \sinh(x)}{\cosh(x)}$$

```
input integrate(tanh(x)^2,x, algorithm="fricas")
```

```
output ((x + 1)*cosh(x) - sinh(x))/cosh(x)
```

3.186.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.50

$$\int \tanh^2(x) dx = x - \tanh(x)$$

input `integrate(tanh(x)**2,x)`output `x - tanh(x)`**3.186.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 2.00

$$\int \tanh^2(x) dx = x - \frac{2}{e^{(-2x)} + 1}$$

input `integrate(tanh(x)^2,x, algorithm="maxima")`output `x - 2/(e^(-2*x) + 1)`**3.186.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 12, normalized size of antiderivative = 2.00

$$\int \tanh^2(x) dx = x + \frac{2}{e^{(2x)} + 1}$$

input `integrate(tanh(x)^2,x, algorithm="giac")`output `x + 2/(e^(2*x) + 1)`

3.186.9 Mupad [B] (verification not implemented)

Time = 16.60 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \tanh^2(x) dx = x - \tanh(x)$$

input `int(tanh(x)^2,x)`

output `x - tanh(x)`

$$3.187 \quad \int \frac{-\sin^2(x) + \sin(2x)}{-\cos^2(x) + \cos(2x)} dx$$

3.187.1 Optimal result	980
3.187.2 Mathematica [A] (verified)	980
3.187.3 Rubi [B] (verified)	981
3.187.4 Maple [A] (verified)	982
3.187.5 Fricas [A] (verification not implemented)	983
3.187.6 Sympy [A] (verification not implemented)	983
3.187.7 Maxima [B] (verification not implemented)	983
3.187.8 Giac [B] (verification not implemented)	984
3.187.9 Mupad [B] (verification not implemented)	984

3.187.1 Optimal result

Integrand size = 25, antiderivative size = 7

$$\int \frac{-\sin^2(x) + \sin(2x)}{-\cos^2(x) + \cos(2x)} dx = x - 2 \log(\sin(x))$$

output `x-2*ln(sin(x))`

3.187.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.71

$$\int \frac{-\sin^2(x) + \sin(2x)}{-\cos^2(x) + \cos(2x)} dx = x - 2 \log(\cos(x)) - 2 \log(\tan(x))$$

input `Integrate[(-Sin[x]^2 + Sin[2*x])/(-Cos[x]^2 + Cos[2*x]),x]`

output `x - 2*Log[Cos[x]] - 2*Log[Tan[x]]`

3.187.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 16 vs. $2(7) = 14$.

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 2.29, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 4889, 25, 523, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin(2x) - \sin^2(x)}{\cos(2x) - \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(2x) - \sin(x)^2}{\cos(2x) - \cos(x)^2} dx \\
 & \quad \downarrow \text{4889} \\
 & \int -\frac{(2 - \tan(x)) \cot(x)}{\tan^2(x) + 1} d \tan(x) \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{\cot(x)(2 - \tan(x))}{\tan^2(x) + 1} d \tan(x) \\
 & \quad \downarrow \text{523} \\
 & - \int \left(2 \cot(x) + \frac{-2 \tan(x) - 1}{\tan^2(x) + 1} \right) d \tan(x) \\
 & \quad \downarrow \text{2009} \\
 & \arctan(\tan(x)) + \log(\tan^2(x) + 1) - 2 \log(\tan(x))
 \end{aligned}$$

input `Int[(-Sin[x]^2 + Sin[2*x])/(-Cos[x]^2 + Cos[2*x]),x]`

output `ArcTan[Tan[x]] - 2*Log[Tan[x]] + Log[1 + Tan[x]^2]`

3.187.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 523 `Int[((x_)^(m_.)*((c_) + (d_.)*(x_)))/((a_) + (b_.)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[x^m*((c + d*x)/(a + b*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && IntegerQ[m]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4889 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, With[{d = FreeFactors[Tan[v], x]}, Simp[d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x^2), Tan[v]/d, u, x], x], x, Tan[v]/d], x]] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Tan[v], x], u, x]] /; InverseFunctionFreeQ[u, x] && !MatchQ[u, (v_.)*((c_.)*tan[w_]^(n_.)*tan[z_]^(n_.))^ (p_.) /; FreeQ[{c, p}, x] && IntegerQ[n] && LinearQ[w, x] && EqQ[z, 2*w]]`

3.187.4 Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

method	result	size
default	$x - 2 \ln(\sin(x))$	8
risch	$2ix - 2 \ln(e^{2ix} - 1) + x$	17

input `int((sin(2*x)-sin(x)^2)/(cos(2*x)-cos(x)^2),x,method=_RETURNVERBOSE)`

output `x-2*ln(sin(x))`

3.187.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.29

$$\int \frac{-\sin^2(x) + \sin(2x)}{-\cos^2(x) + \cos(2x)} dx = x - 2 \log \left(\frac{1}{2} \sin(x) \right)$$

input `integrate((sin(2*x)-sin(x)^2)/(cos(2*x)-cos(x)^2),x, algorithm="fricas")`

output `x - 2*log(1/2*sin(x))`

3.187.6 Sympy [A] (verification not implemented)

Time = 1.45 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

$$\int \frac{-\sin^2(x) + \sin(2x)}{-\cos^2(x) + \cos(2x)} dx = x - \log(\cos^2(x) - 1)$$

input `integrate((sin(2*x)-sin(x)**2)/(cos(2*x)-cos(x)**2),x)`

output `x - log(cos(x)**2 - 1)`

3.187.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 36 vs. 2(7) = 14.

Time = 0.19 (sec) , antiderivative size = 36, normalized size of antiderivative = 5.14

$$\int \frac{-\sin^2(x) + \sin(2x)}{-\cos^2(x) + \cos(2x)} dx = x - \log(\cos(x)^2 + \sin(x)^2 + 2 \cos(x) + 1) - \log(\cos(x)^2 + \sin(x)^2 - 2 \cos(x) + 1)$$

input `integrate((sin(2*x)-sin(x)^2)/(cos(2*x)-cos(x)^2),x, algorithm="maxima")`

output `x - log(cos(x)^2 + sin(x)^2 + 2*cos(x) + 1) - log(cos(x)^2 + sin(x)^2 - 2*cos(x) + 1)`

3.187.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. 2(7) = 14.

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 3.00

$$\int \frac{-\sin^2(x) + \sin(2x)}{-\cos^2(x) + \cos(2x)} dx = x + 2 \log \left(\tan \left(\frac{1}{2} x \right)^2 + 1 \right) - 2 \log \left(\left| \tan \left(\frac{1}{2} x \right) \right| \right)$$

input `integrate((sin(2*x)-sin(x)^2)/(cos(2*x)-cos(x)^2),x, algorithm="giac")`

output `x + 2*log(tan(1/2*x)^2 + 1) - 2*log(abs(tan(1/2*x)))`

3.187.9 Mupad [B] (verification not implemented)

Time = 16.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 3.71

$$\int \frac{-\sin^2(x) + \sin(2x)}{-\cos^2(x) + \cos(2x)} dx = -2 \ln(\tan(x)) + \ln(\tan(x) - i) \left(1 - \frac{1}{2}i \right) + \ln(\tan(x) + i) \left(1 + \frac{1}{2}i \right)$$

input `int((sin(2*x) - sin(x)^2)/(cos(2*x) - cos(x)^2),x)`

output `log(tan(x) - 1i)*(1 - 1i/2) - 2*log(tan(x)) + log(tan(x) + 1i)*(1 + 1i/2)`

3.188 $\int \frac{1}{x^{9/25} + x^{41/25}} dx$

3.188.1 Optimal result	985
3.188.2 Mathematica [A] (verified)	985
3.188.3 Rubi [A] (verified)	986
3.188.4 Maple [F(-1)]	987
3.188.5 Fricas [A] (verification not implemented)	987
3.188.6 Sympy [F(-1)]	988
3.188.7 Maxima [F]	988
3.188.8 Giac [A] (verification not implemented)	988
3.188.9 Mupad [B] (verification not implemented)	989

3.188.1 Optimal result

Integrand size = 13, antiderivative size = 10

$$\int \frac{1}{x^{9/25} + x^{41/25}} dx = \frac{25}{16} \arctan(x^{16/25})$$

output 25/16*arctan(x^(16/25))

3.188.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^{9/25} + x^{41/25}} dx = \frac{25}{16} \arctan(x^{16/25})$$

input Integrate[(x^(9/25) + x^(41/25))^(-1), x]

output (25*ArcTan[x^(16/25)])/16

3.188.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {2027, 864, 807, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^{41/25} + x^{9/25}} dx \\
 & \quad \downarrow \text{2027} \\
 & \int \frac{1}{x^{9/25} (x^{32/25} + 1)} dx \\
 & \quad \downarrow \text{864} \\
 & 25 \int \frac{x^{3/5}}{x^{32/25} + 1} d \sqrt[25]{x} \\
 & \quad \downarrow \text{807} \\
 & \frac{25}{16} \int \frac{1}{x^{2/25} + 1} dx^{16/25} \\
 & \quad \downarrow \text{216} \\
 & \frac{25}{16} \arctan \left(x^{16/25} \right)
 \end{aligned}$$

input `Int[(x^(9/25) + x^(41/25))^-1], x]`

output `(25*ArcTan[x^(16/25)])/16`

3.188.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 864 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a + b*x^(k*n))^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, m, p}, x] && FractionQ[n]`

rule 2027 `Int[(Fx_)*((a_)*(x_)^(r_) + (b_)*(x_)^(s_))^(p_), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r))^p*Fx, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] && PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])`

3.188.4 Maple **[F(-1)]**

Timed out.

hanged

input `int(1/(x^(41/25)+x^(9/25)),x)`

output `int(1/(x^(41/25)+x^(9/25)),x)`

3.188.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.60

$$\int \frac{1}{x^{9/25} + x^{41/25}} dx = \frac{25}{16} \arctan\left(x^{16/25}\right)$$

input `integrate(1/(x^(41/25)+x^(9/25)),x, algorithm="fricas")`

output `25/16*arctan(x^(16/25))`

3.188.6 Sympy [F(-1)]

Timed out.

$$\int \frac{1}{x^{9/25} + x^{41/25}} dx = \text{Timed out}$$

input `integrate(1/(x**(41/25)+x**(9/25)),x)`output `Timed out`**3.188.7 Maxima [F]**

$$\int \frac{1}{x^{9/25} + x^{41/25}} dx = \int \frac{1}{x^{41/25} + x^{9/25}} dx$$

input `integrate(1/(x^(41/25)+x^(9/25)),x, algorithm="maxima")`output `25/16*x^(16/25) - integrate(x^(23/25)/(x^(32/25) + 1), x)`**3.188.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.60

$$\int \frac{1}{x^{9/25} + x^{41/25}} dx = \frac{25}{16} \arctan \left(x^{16/25} \right)$$

input `integrate(1/(x^(41/25)+x^(9/25)),x, algorithm="giac")`output `25/16*arctan(x^(16/25))`

3.188.9 Mupad [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.60

$$\int \frac{1}{x^{9/25} + x^{41/25}} dx = \frac{25 \operatorname{atan}(x^{16/25})}{16}$$

input `int(1/(x^(9/25) + x^(41/25)),x)`

output `(25*atan(x^(16/25)))/16`

$$3.189 \quad \int \frac{\cos(x)}{2 - \cos^2(x)} dx$$

3.189.1 Optimal result	990
3.189.2 Mathematica [A] (verified)	990
3.189.3 Rubi [A] (verified)	991
3.189.4 Maple [A] (verified)	992
3.189.5 Fricas [A] (verification not implemented)	992
3.189.6 Sympy [B] (verification not implemented)	993
3.189.7 Maxima [A] (verification not implemented)	993
3.189.8 Giac [A] (verification not implemented)	994
3.189.9 Mupad [B] (verification not implemented)	994

3.189.1 Optimal result

Integrand size = 13, antiderivative size = 3

$$\int \frac{\cos(x)}{2 - \cos^2(x)} dx = \arctan(\sin(x))$$

output `arctan(sin(x))`

3.189.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{2 - \cos^2(x)} dx = \arctan(\sin(x))$$

input `Integrate[Cos[x]/(2 - Cos[x]^2), x]`

output `ArcTan[Sin[x]]`

3.189.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3042, 3665, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(x)}{2 - \cos^2(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sin(x + \frac{\pi}{2})}{2 - \sin(x + \frac{\pi}{2})^2} dx \\ & \quad \downarrow \text{3665} \\ & \int \frac{1}{\sin^2(x) + 1} d\sin(x) \\ & \quad \downarrow \text{216} \\ & \arctan(\sin(x)) \end{aligned}$$

input `Int[Cos[x]/(2 - Cos[x]^2),x]`

output `ArcTan[Sin[x]]`

3.189.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`


```
rule 3665 Int[sin[(e_.) + (f_.)*(x_.)]^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_.)]^2)^(p_.), x_Symbol] :> With[{ff = FreeFactors[Cos[e + f*x], x]}, Simp[-ff/f Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b - b*ff^2*x^2)^p, x], x, Cos[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

3.189.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
default	$\arctan(\sin(x))$	4
parallelrisc	$\frac{i \left(\ln\left(-\frac{2i(\sin(x)+i)}{1+\cos(x)}\right) - \ln\left(\frac{2+2i\sin(x)}{1+\cos(x)}\right) \right)}{2}$	37
risc	$\frac{i \ln(e^{2ix} - 2e^{ix} - 1)}{2} - \frac{i \ln(e^{2ix} + 2e^{ix} - 1)}{2}$	38

```
input int(cos(x)/(2-cos(x)^2),x,method=_RETURNVERBOSE)
```

```
output arctan(sin(x))
```

3.189.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{2 - \cos^2(x)} dx = \arctan(\sin(x))$$

```
input integrate(cos(x)/(2-cos(x)^2),x, algorithm="fracas")
```

```
output arctan(sin(x))
```

3.189.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 219 vs. $2(3) = 6$.

Time = 9.12 (sec) , antiderivative size = 219, normalized size of antiderivative = 73.00

$$\int \frac{\cos(x)}{2 - \cos^2(x)} dx = \frac{1607521\sqrt{3 - 2\sqrt{2}} \left(\operatorname{atan} \left(\frac{\tan(\frac{x}{2})}{\sqrt{3 - 2\sqrt{2}}} \right) + \pi \left\lfloor \frac{\frac{x}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{470832\sqrt{2} + 665857} + \frac{1136689\sqrt{2}\sqrt{3 - 2\sqrt{2}} \left(\operatorname{atan} \left(\frac{\tan(\frac{x}{2})}{\sqrt{3 - 2\sqrt{2}}} \right) + \pi \left\lfloor \frac{\frac{x}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{470832\sqrt{2} + 665857} - \frac{195025\sqrt{2}\sqrt{2\sqrt{2} + 3} \left(\operatorname{atan} \left(\frac{\tan(\frac{x}{2})}{\sqrt{2\sqrt{2} + 3}} \right) + \pi \left\lfloor \frac{\frac{x}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{470832\sqrt{2} + 665857} - \frac{275807\sqrt{2\sqrt{2} + 3} \left(\operatorname{atan} \left(\frac{\tan(\frac{x}{2})}{\sqrt{2\sqrt{2} + 3}} \right) + \pi \left\lfloor \frac{\frac{x}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{470832\sqrt{2} + 665857}$$

input `integrate(cos(x)/(2-cos(x)**2),x)`

output `1607521*sqrt(3 - 2*sqrt(2))*(atan(tan(x/2)/sqrt(3 - 2*sqrt(2)))) + pi*floor((x/2 - pi/2)/pi)/(470832*sqrt(2) + 665857) + 1136689*sqrt(2)*sqrt(3 - 2*sqrt(2))*(atan(tan(x/2)/sqrt(3 - 2*sqrt(2)))) + pi*floor((x/2 - pi/2)/pi)/(470832*sqrt(2) + 665857) - 195025*sqrt(2)*sqrt(2*sqrt(2) + 3)*(atan(tan(x/2)/sqrt(2*sqrt(2) + 3)) + pi*floor((x/2 - pi/2)/pi))/(470832*sqrt(2) + 665857) - 275807*sqrt(2*sqrt(2) + 3)*(atan(tan(x/2)/sqrt(2*sqrt(2) + 3)) + pi*floor((x/2 - pi/2)/pi))/(470832*sqrt(2) + 665857)`

3.189.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{2 - \cos^2(x)} dx = \arctan(\sin(x))$$

input `integrate(cos(x)/(2-cos(x)^2),x, algorithm="maxima")`

output `arctan(sin(x))`

3.189.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{2 - \cos^2(x)} dx = \arctan(\sin(x))$$

input `integrate(cos(x)/(2-cos(x)^2),x, algorithm="giac")`

output `arctan(sin(x))`

3.189.9 Mupad [B] (verification not implemented)

Time = 16.48 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{2 - \cos^2(x)} dx = \operatorname{atan}(\sin(x))$$

input `int(-cos(x)/(cos(x)^2 - 2),x)`

output `atan(sin(x))`

3.190 $\int \frac{1}{(1+x^2)^{3/2}} dx$

3.190.1 Optimal result 995
 3.190.2 Mathematica [A] (verified) 995
 3.190.3 Rubi [A] (verified) 996
 3.190.4 Maple [A] (verified) 996
 3.190.5 Fricas [B] (verification not implemented) 997
 3.190.6 Sympy [A] (verification not implemented) 997
 3.190.7 Maxima [A] (verification not implemented) 997
 3.190.8 Giac [A] (verification not implemented) 998
 3.190.9 Mupad [B] (verification not implemented) 998

3.190.1 Optimal result

Integrand size = 9, antiderivative size = 11

$$\int \frac{1}{(1+x^2)^{3/2}} dx = \frac{x}{\sqrt{1+x^2}}$$

output `x/(x^2+1)^(1/2)`

3.190.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x^2)^{3/2}} dx = \frac{x}{\sqrt{1+x^2}}$$

input `Integrate[(1 + x^2)^(-3/2),x]`

output `x/Sqrt[1 + x^2]`

3.190.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {208}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x^2 + 1)^{3/2}} dx$$

↓ 208

$$\frac{x}{\sqrt{x^2 + 1}}$$

input `Int[(1 + x^2)^(-3/2),x]`

output `x/Sqrt[1 + x^2]`

3.190.3.1 Defintions of rubi rules used

rule 208 `Int[((a_) + (b_.)*(x_)^2)^(-3/2), x_Symbol] :> Simp[x/(a*Sqrt[a + b*x^2]), x] /; FreeQ[{a, b}, x]`

3.190.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

method	result	size
gospers	$\frac{x}{\sqrt{x^2+1}}$	10
default	$\frac{x}{\sqrt{x^2+1}}$	10
trager	$\frac{x}{\sqrt{x^2+1}}$	10
meijerg	$\frac{x}{\sqrt{x^2+1}}$	10
risch	$\frac{x}{\sqrt{x^2+1}}$	10
pseudoelliptic	$\frac{x}{\sqrt{x^2+1}}$	10

input `int(1/(x^2+1)^(3/2),x,method=_RETURNVERBOSE)`

output $x/(x^2+1)^{(1/2)}$

3.190.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 22 vs. 2(9) = 18.

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 2.00

$$\int \frac{1}{(1+x^2)^{3/2}} dx = \frac{x^2 + \sqrt{x^2 + 1}x + 1}{x^2 + 1}$$

input `integrate(1/(x^2+1)^(3/2),x, algorithm="fricas")`

output $(x^2 + \sqrt{x^2 + 1}x + 1)/(x^2 + 1)$

3.190.6 Sympy [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \frac{1}{(1+x^2)^{3/2}} dx = \frac{x}{\sqrt{x^2 + 1}}$$

input `integrate(1/(x**2+1)**(3/2),x)`

output $x/\sqrt{x^2 + 1}$

3.190.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \frac{1}{(1+x^2)^{3/2}} dx = \frac{x}{\sqrt{x^2 + 1}}$$

input `integrate(1/(x^2+1)^(3/2),x, algorithm="maxima")`

output $x/\sqrt{x^2 + 1}$

3.190.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \frac{1}{(1+x^2)^{3/2}} dx = \frac{x}{\sqrt{x^2+1}}$$

input `integrate(1/(x^2+1)^(3/2),x, algorithm="giac")`output `x/sqrt(x^2 + 1)`**3.190.9 Mupad [B] (verification not implemented)**

Time = 16.45 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \frac{1}{(1+x^2)^{3/2}} dx = \frac{x}{\sqrt{x^2+1}}$$

input `int(1/(x^2 + 1)^(3/2),x)`output `x/(x^2 + 1)^(1/2)`

3.191 $\int \frac{1}{\sqrt{x^{3/2}-x^2}} dx$

3.191.1 Optimal result 999
 3.191.2 Mathematica [A] (verified) 999
 3.191.3 Rubi [A] (verified) 1000
 3.191.4 Maple [A] (verified) 1001
 3.191.5 Fricas [A] (verification not implemented) 1001
 3.191.6 Sympy [F] 1001
 3.191.7 Maxima [F] 1002
 3.191.8 Giac [A] (verification not implemented) 1002
 3.191.9 Mupad [F(-1)] 1002

3.191.1 Optimal result

Integrand size = 15, antiderivative size = 20

$$\int \frac{1}{\sqrt{x^{3/2}-x^2}} dx = 4 \arctan \left(\frac{x}{\sqrt{x^{3/2}-x^2}} \right)$$

output `4*arctan(x/(x^(3/2)-x^2)^(1/2))`

3.191.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{x^{3/2}-x^2}} dx = 4 \arctan \left(\frac{x}{\sqrt{x^{3/2}-x^2}} \right)$$

input `Integrate[1/Sqrt[x^(3/2) - x^2],x]`

output `4*ArcTan[x/Sqrt[x^(3/2) - x^2]]`

3.191.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {1914, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x^{3/2} - x^2}} dx$$

↓ 1914

$$4 \int \frac{1}{\frac{x^2}{x^{3/2} - x^2} + 1} d \frac{x}{\sqrt{x^{3/2} - x^2}}$$

↓ 216

$$4 \arctan \left(\frac{x}{\sqrt{x^{3/2} - x^2}} \right)$$

input `Int[1/Sqrt[x^(3/2) - x^2],x]`

output `4*ArcTan[x/Sqrt[x^(3/2) - x^2]]`

3.191.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 1914 `Int[1/Sqrt[(a_.)*(x_)^2 + (b_.)*(x_)^(n_.)], x_Symbol] := Simp[2/(2 - n) Subst[Int[1/(1 - a*x^2), x], x, x/Sqrt[a*x^2 + b*x^n]], x] /; FreeQ[{a, b, n}, x] && NeQ[n, 2]`

3.191.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.35

method	result	size
meijerg	$4 \arcsin \left(x^{\frac{1}{4}} \right)$	7
derivativedivides	$\frac{2\sqrt{x} \sqrt{-\sqrt{x}(-1+\sqrt{x})} \arcsin(2\sqrt{x}-1)}{\sqrt{x^{\frac{3}{2}}-x^2}}$	37
default	$\frac{2x(-1+\sqrt{x}) \ln\left(\sqrt{x}-\frac{1}{2}+\sqrt{x-\sqrt{x}}\right)}{\sqrt{x^{\frac{3}{2}}-x^2} \sqrt{\sqrt{x}(-1+\sqrt{x})}}$	46

input `int(1/(x^(3/2)-x^2)^(1/2),x,method=_RETURNVERBOSE)`output `4*arcsin(x^(1/4))`**3.191.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{1}{\sqrt{x^{3/2}-x^2}} dx = -4 \arctan \left(\frac{\sqrt{-x^2+x^{3/2}}}{x} \right)$$

input `integrate(1/(x^(3/2)-x^2)^(1/2),x, algorithm="fracas")`output `-4*arctan(sqrt(-x^2 + x^(3/2))/x)`**3.191.6 Sympy [F]**

$$\int \frac{1}{\sqrt{x^{3/2}-x^2}} dx = \int \frac{1}{\sqrt{x^{\frac{3}{2}}-x^2}} dx$$

input `integrate(1/(x**(3/2)-x**2)**(1/2),x)`output `Integral(1/sqrt(x**(3/2) - x**2), x)`

3.191.7 Maxima [F]

$$\int \frac{1}{\sqrt{x^{3/2} - x^2}} dx = \int \frac{1}{\sqrt{-x^2 + x^{\frac{3}{2}}}} dx$$

input `integrate(1/(x^(3/2)-x^2)^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(-x^2 + x^(3/2)), x)`

3.191.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{x^{3/2} - x^2}} dx = (\pi + 2 \arcsin(2\sqrt{x} - 1)) \operatorname{sgn}(x)$$

input `integrate(1/(x^(3/2)-x^2)^(1/2),x, algorithm="giac")`

output `(pi + 2*arcsin(2*sqrt(x) - 1))*sgn(x)`

3.191.9 Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{x^{3/2} - x^2}} dx = \int \frac{1}{\sqrt{x^{3/2} - x^2}} dx$$

input `int(1/(x^(3/2) - x^2)^(1/2),x)`

output `int(1/(x^(3/2) - x^2)^(1/2), x)`

3.192 $\int \frac{-1+x}{x+x^2 \log(x)} dx$

3.192.1 Optimal result	1003
3.192.2 Mathematica [A] (verified)	1003
3.192.3 Rubi [A] (verified)	1004
3.192.4 Maple [A] (verified)	1005
3.192.5 Fricas [A] (verification not implemented)	1005
3.192.6 Sympy [A] (verification not implemented)	1005
3.192.7 Maxima [A] (verification not implemented)	1006
3.192.8 Giac [A] (verification not implemented)	1006
3.192.9 Mupad [B] (verification not implemented)	1006

3.192.1 Optimal result

Integrand size = 14, antiderivative size = 7

$$\int \frac{-1+x}{x+x^2 \log(x)} dx = \log\left(\frac{1}{x} + \log(x)\right)$$

output `ln(1/x+ln(x))`

3.192.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.71

$$\int \frac{-1+x}{x+x^2 \log(x)} dx = -\log(x) + \log(1+x \log(x))$$

input `Integrate[(-1 + x)/(x + x^2*Log[x]),x]`

output `-Log[x] + Log[1 + x*Log[x]]`

3.192.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3041, 7235}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x-1}{x^2 \log(x) + x} dx$$

↓ 3041

$$\int \frac{x-1}{x^2 \left(\frac{1}{x} + \log(x)\right)} dx$$

↓ 7235

$$\log\left(\frac{1}{x} + \log(x)\right)$$

input `Int[(-1 + x)/(x + x^2*Log[x]), x]`

output `Log[x^(-1) + Log[x]]`

3.192.3.1 Defintions of rubi rules used

rule 3041 `Int[(u_.)*((a_.)*(x_)^(m_.) + Log[(c_.)*(x_)^(n_.)]^(q_.)*(b_.)*(x_)^(r_.))^(p_.), x_Symbol] :=> Int[u*x^(p*r)*(a*x^(m - r) + b*Log[c*x^n]^q)^p, x] /;`
`FreeQ[{a, b, c, m, n, p, q, r}, x] && IntegerQ[p]`

rule 7235 `Int[(u_)/(y_), x_Symbol] :=> With[{q = DerivativeDivides[y, u, x]}, Simp[q*Log[RemoveContent[y, x]], x] /;`
`!FalseQ[q]`

3.192.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

method	result	size
risch	$\ln\left(\frac{1}{x} + \ln(x)\right)$	8
default	$-\ln(x) + \ln(x \ln(x) + 1)$	13
norman	$-\ln(x) + \ln(x \ln(x) + 1)$	13
parallelrisch	$-\ln(x) + \ln(x \ln(x) + 1)$	13

input `int((-1+x)/(x+x^2*ln(x)),x,method=_RETURNVERBOSE)`output `ln(1/x+ln(x))`**3.192.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.57

$$\int \frac{-1+x}{x+x^2 \log(x)} dx = \log\left(\frac{x \log(x) + 1}{x}\right)$$

input `integrate((-1+x)/(x+x^2*log(x)),x, algorithm="fricas")`output `log((x*log(x) + 1)/x)`**3.192.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{-1+x}{x+x^2 \log(x)} dx = \log\left(\log(x) + \frac{1}{x}\right)$$

input `integrate((-1+x)/(x+x**2*ln(x)),x)`output `log(log(x) + 1/x)`

3.192.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.57

$$\int \frac{-1+x}{x+x^2 \log(x)} dx = \log\left(\frac{x \log(x)+1}{x}\right)$$

input `integrate((-1+x)/(x+x^2*log(x)),x, algorithm="maxima")`output `log((x*log(x) + 1)/x)`**3.192.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.71

$$\int \frac{-1+x}{x+x^2 \log(x)} dx = \log(x \log(x)+1) - \log(x)$$

input `integrate((-1+x)/(x+x^2*log(x)),x, algorithm="giac")`output `log(x*log(x) + 1) - log(x)`**3.192.9 Mupad [B] (verification not implemented)**

Time = 16.32 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.71

$$\int \frac{-1+x}{x+x^2 \log(x)} dx = \ln(x \ln(x)+1) - \ln(x)$$

input `int((x - 1)/(x + x^2*log(x)),x)`output `log(x*log(x) + 1) - log(x)`

3.193 $\int \csc(x) \sec(x) dx$

3.193.1 Optimal result	1007
3.193.2 Mathematica [B] (verified)	1007
3.193.3 Rubi [A] (verified)	1008
3.193.4 Maple [A] (verified)	1009
3.193.5 Fricas [B] (verification not implemented)	1009
3.193.6 Sympy [B] (verification not implemented)	1009
3.193.7 Maxima [B] (verification not implemented)	1010
3.193.8 Giac [B] (verification not implemented)	1010
3.193.9 Mupad [B] (verification not implemented)	1010

3.193.1 Optimal result

Integrand size = 5, antiderivative size = 3

$$\int \csc(x) \sec(x) dx = \log(\tan(x))$$

output `ln(tan(x))`

3.193.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 9 vs. $2(3) = 6$.

Time = 0.00 (sec) , antiderivative size = 9, normalized size of antiderivative = 3.00

$$\int \csc(x) \sec(x) dx = -\log(\cos(x)) + \log(\sin(x))$$

input `Integrate[Csc[x]*Sec[x],x]`

output `-Log[Cos[x]] + Log[Sin[x]]`

3.193.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 3100, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \csc(x) \sec(x) dx \\ \downarrow \text{3042} \\ \int \csc(x) \sec(x) dx \\ \downarrow \text{3100} \\ \int \cot(x) d \tan(x) \\ \downarrow \text{14} \\ \log(\tan(x)) \end{array}$$

input `Int[Csc[x]*Sec[x],x]`

output `Log[Tan[x]]`

3.193.3.1 Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3100 `Int[csc[(e_.) + (f_.)*(x_)]^(m_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[1/f Subst[Int[(1 + x^2)^((m + n)/2 - 1)/x^m, x], x, Tan[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n)/2]`

3.193.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
default	$\ln(\tan(x))$	4
risch	$\ln(e^{2ix} - 1) - \ln(e^{2ix} + 1)$	20
norman	$-\ln(\tan(\frac{x}{2}) - 1) - \ln(1 + \tan(\frac{x}{2})) + \ln(\tan(\frac{x}{2}))$	25
parallelrisc	$-\ln(\tan(\frac{x}{2}) - 1) - \ln(1 + \tan(\frac{x}{2})) + \ln(\tan(\frac{x}{2}))$	25

input `int(csc(x)*sec(x),x,method=_RETURNVERBOSE)`

output `ln(tan(x))`

3.193.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(3) = 6$.

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 6.33

$$\int \csc(x) \sec(x) dx = -\frac{1}{2} \log(\cos(x)^2) + \frac{1}{2} \log\left(-\frac{1}{4} \cos(x)^2 + \frac{1}{4}\right)$$

input `integrate(csc(x)*sec(x),x, algorithm="fricas")`

output `-1/2*log(cos(x)^2) + 1/2*log(-1/4*cos(x)^2 + 1/4)`

3.193.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 14 vs. $2(3) = 6$.

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 4.67

$$\int \csc(x) \sec(x) dx = -\frac{\log(\sin^2(x) - 1)}{2} + \log(\sin(x))$$

input `integrate(csc(x)*sec(x),x)`

output `-log(sin(x)**2 - 1)/2 + log(sin(x))`

3.193.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(3) = 6$.

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 5.67

$$\int \csc(x) \sec(x) dx = -\frac{1}{2} \log(\sin(x)^2 - 1) + \frac{1}{2} \log(\sin(x)^2)$$

input `integrate(csc(x)*sec(x),x, algorithm="maxima")`

output `-1/2*log(sin(x)^2 - 1) + 1/2*log(sin(x)^2)`

3.193.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 16 vs. $2(3) = 6$.

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 5.33

$$\int \csc(x) \sec(x) dx = -\frac{1}{2} \log(-\sin(x)^2 + 1) + \log(|\sin(x)|)$$

input `integrate(csc(x)*sec(x),x, algorithm="giac")`

output `-1/2*log(-sin(x)^2 + 1) + log(abs(sin(x)))`

3.193.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \csc(x) \sec(x) dx = \ln(\tan(x))$$

input `int(1/(cos(x)*sin(x)),x)`

output `log(tan(x))`

3.194 $\int \tan(\cos(x)) dx$

3.194.1 Optimal result1011
3.194.2 Mathematica [N/A]1011
3.194.3 Rubi [N/A]	1012
3.194.4 Maple [N/A] (verified)	1013
3.194.5 Fricas [N/A]	1013
3.194.6 Sympy [N/A]	1014
3.194.7 Maxima [N/A]	1014
3.194.8 Giac [N/A]	1014
3.194.9 Mupad [N/A]	1015

3.194.1 Optimal result

Integrand size = 3, antiderivative size = 3

$$\int \tan(\cos(x)) dx = \text{Int}(\tan(\cos(x)), x)$$

output `CannotIntegrate(tan(cos(x)),x)`

3.194.2 Mathematica [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.67

$$\int \tan(\cos(x)) dx = \int \tan(\cos(x)) dx$$

input `Integrate[Tan[Cos[x]],x]`

output `Integrate[Tan[Cos[x]], x]`

3.194.3 Rubi [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {4902, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan(\cos(x)) dx \\
 & \quad \downarrow \text{4902} \\
 & 2 \int \frac{\tan\left(\frac{1-\tan^2\left(\frac{x}{2}\right)}{\tan^2\left(\frac{x}{2}\right)+1}\right)}{\tan^2\left(\frac{x}{2}\right)+1} d \tan\left(\frac{x}{2}\right) \\
 & \quad \downarrow \text{7276} \\
 & 2 \int \left(\frac{i \tan\left(\frac{1-\tan^2\left(\frac{x}{2}\right)}{\tan^2\left(\frac{x}{2}\right)+1}\right)}{2(i-\tan\left(\frac{x}{2}\right))} + \frac{i \tan\left(\frac{1-\tan^2\left(\frac{x}{2}\right)}{\tan^2\left(\frac{x}{2}\right)+1}\right)}{2(\tan\left(\frac{x}{2}\right)+i)} \right) d \tan\left(\frac{x}{2}\right) \\
 & \quad \downarrow \text{2009} \\
 & 2 \left(\frac{1}{2} i \int \frac{\tan\left(\frac{1-\tan^2\left(\frac{x}{2}\right)}{\tan^2\left(\frac{x}{2}\right)+1}\right)}{i-\tan\left(\frac{x}{2}\right)} d \tan\left(\frac{x}{2}\right) + \frac{1}{2} i \int \frac{\tan\left(\frac{1-\tan^2\left(\frac{x}{2}\right)}{\tan^2\left(\frac{x}{2}\right)+1}\right)}{\tan\left(\frac{x}{2}\right)+i} d \tan\left(\frac{x}{2}\right) \right)
 \end{aligned}$$

input `Int[Tan[Cos[x]], x]`

output `$Aborted`

3.194.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 4902 Int[u_, x_Symbol] := With[{w = Block[{$ShowSteps = False, $StepCounter = Null}, Int[SubstFor[1/(1 + FreeFactors[Tan[FunctionOfTrig[u, x]/2], x]^2*x^2), Tan[FunctionOfTrig[u, x]/2]/FreeFactors[Tan[FunctionOfTrig[u, x]/2], x], u, x], x]}], Module[{v = FunctionOfTrig[u, x], d}, Simp[d = FreeFactors[Tan[v/2], x]; 2*(d/Coefficient[v, x, 1]) Subst[Int[SubstFor[1/(1 + d^2*x^2), Tan[v/2]/d, u, x], x], x, Tan[v/2]/d], x]] /; CalculusFreeQ[w, x]] /; InverseFunctionFreeQ[u, x] && !FalseQ[FunctionOfTrig[u, x]]
```

```
rule 7276 Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]
```

3.194.4 Maple [N/A] (verified)

Not integrable

Time = 0.03 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \tan(\cos(x)) dx$$

input `int(tan(cos(x)),x)`

output `int(tan(cos(x)),x)`

3.194.5 Fracas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.67

$$\int \tan(\cos(x)) dx = \int \tan(\cos(x)) dx$$

input `integrate(tan(cos(x)),x, algorithm="fricas")`

output `integral(tan(cos(x)), x)`

3.194.6 Sympy [N/A]

Not integrable

Time = 1.38 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.67

$$\int \tan(\cos(x)) dx = \int \tan(\cos(x)) dx$$

input `integrate(tan(cos(x)),x)`output `Integral(tan(cos(x)), x)`**3.194.7 Maxima [N/A]**

Not integrable

Time = 0.45 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.67

$$\int \tan(\cos(x)) dx = \int \tan(\cos(x)) dx$$

input `integrate(tan(cos(x)),x, algorithm="maxima")`output `integrate(tan(cos(x)), x)`**3.194.8 Giac [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.67

$$\int \tan(\cos(x)) dx = \int \tan(\cos(x)) dx$$

input `integrate(tan(cos(x)),x, algorithm="giac")`output `integrate(tan(cos(x)), x)`

3.194.9 Mupad [N/A]

Not integrable

Time = 15.88 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.67

$$\int \tan(\cos(x)) dx = \int \tan(\cos(x)) dx$$

input `int(tan(cos(x)), x)`

output `int(tan(cos(x)), x)`

$$\mathbf{3.195} \quad \int \frac{1+x}{x(x+\log(x))} dx$$

3.195.1 Optimal result	1016
3.195.2 Mathematica [A] (verified)	1016
3.195.3 Rubi [A] (verified)	1017
3.195.4 Maple [A] (verified)	1017
3.195.5 Fricas [A] (verification not implemented)	1018
3.195.6 Sympy [A] (verification not implemented)	1018
3.195.7 Maxima [A] (verification not implemented)	1018
3.195.8 Giac [A] (verification not implemented)	1019
3.195.9 Mupad [B] (verification not implemented)	1019

3.195.1 Optimal result

Integrand size = 13, antiderivative size = 5

$$\int \frac{1+x}{x(x+\log(x))} dx = \log(x + \log(x))$$

output `ln(x+ln(x))`

3.195.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1+x}{x(x+\log(x))} dx = \log(x + \log(x))$$

input `Integrate[(1 + x)/(x*(x + Log[x])), x]`

output `Log[x + Log[x]]`

3.195.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {7235}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x+1}{x(x+\log(x))} dx$$

↓ 7235

$$\log(x+\log(x))$$

input `Int[(1 + x)/(x*(x + Log[x])),x]`

output `Log[x + Log[x]]`

3.195.3.1 Defintions of rubi rules used

rule 7235 `Int[(u_)/(y_), x_Symbol] :=> With[{q = DerivativeDivides[y, u, x]}, Simp[q*Log[RemoveContent[y, x]], x] /; !FalseQ[q]]`

3.195.4 Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
default	$\ln(x + \ln(x))$	6
norman	$\ln(x + \ln(x))$	6
risch	$\ln(x + \ln(x))$	6
parallelrisk	$\ln(x + \ln(x))$	6

input `int((1+x)/x/(x+ln(x)),x,method=_RETURNVERBOSE)`

output `ln(x+ln(x))`

3.195.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1+x}{x(x+\log(x))} dx = \log(x + \log(x))$$

input `integrate((1+x)/x/(x+log(x)),x, algorithm="fricas")`output `log(x + log(x))`**3.195.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1+x}{x(x+\log(x))} dx = \log(x + \log(x))$$

input `integrate((1+x)/x/(x+ln(x)),x)`output `log(x + log(x))`**3.195.7 Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1+x}{x(x+\log(x))} dx = \log(x + \log(x))$$

input `integrate((1+x)/x/(x+log(x)),x, algorithm="maxima")`output `log(x + log(x))`

3.195.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1+x}{x(x+\log(x))} dx = \log(x + \log(x))$$

input `integrate((1+x)/x/(x+log(x)),x, algorithm="giac")`

output `log(x + log(x))`

3.195.9 Mupad [B] (verification not implemented)

Time = 16.58 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1+x}{x(x+\log(x))} dx = \ln(x + \ln(x))$$

input `int((x + 1)/(x*(x + log(x))),x)`

output `log(x + log(x))`

3.196 $\int (e^{-e^x+x} + e^{e^x+x}) dx$

3.196.1 Optimal result	1020
3.196.2 Mathematica [B] (verified)	1020
3.196.3 Rubi [B] (verified)	1021
3.196.4 Maple [B] (verified)	1021
3.196.5 Fracas [B] (verification not implemented)	1022
3.196.6 Sympy [F(-1)]	1022
3.196.7 Maxima [B] (verification not implemented)	1022
3.196.8 Giac [B] (verification not implemented)	1023
3.196.9 Mupad [B] (verification not implemented)	1023

3.196.1 Optimal result

Integrand size = 17, antiderivative size = 6

$$\int (e^{-e^x+x} + e^{e^x+x}) dx = 2 \sinh(e^x)$$

output `2*sinh(exp(x))`

3.196.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 15 vs. 2(6) = 12.

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 2.50

$$\int (e^{-e^x+x} + e^{e^x+x}) dx = -e^{-e^x} + e^{e^x}$$

input `Integrate[E^(-E^x + x) + E^(E^x + x), x]`

output `-E^(-E^x) + E^E^x`

3.196.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 15 vs. $2(6) = 12$.

Time = 0.16 (sec) , antiderivative size = 15, normalized size of antiderivative = 2.50, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e^{x-e^x} + e^{x+e^x}) dx$$

$$\downarrow \text{2009}$$

$$e^{e^x} - e^{-e^x}$$

input `Int[E^(-E^x + x) + E^(E^x + x),x]`

output `-E^(-E^x) + E^E^x`

3.196.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.196.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. $2(5) = 10$.

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 2.00

method	result	size
default	$-e^{-e^x} + e^{e^x}$	12
parts	$-e^{-e^x} + e^{e^x}$	12
risch	$(e^{e^x+x} - e^{x-e^x}) e^{-x}$	21

input `int(exp(exp(x)+x)+exp(x-exp(x)),x,method=_RETURNVERBOSE)`

output `-1/exp(exp(x))+exp(exp(x))`

3.196.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. $2(5) = 10$.

Time = 0.25 (sec) , antiderivative size = 27, normalized size of antiderivative = 4.50

$$\int (e^{-e^x+x} + e^{e^x+x}) dx = -(e^{(2x)} - e^{(2x+2e^x)})e^{(-2x-e^x)}$$

input `integrate(exp(exp(x)+x)+exp(x-exp(x)),x, algorithm="fricas")`

output `-(e^(2*x) - e^(2*x + 2*e^x))*e^(-2*x - e^x)`

3.196.6 Sympy [F(-1)]

Timed out.

$$\int (e^{-e^x+x} + e^{e^x+x}) dx = \text{Timed out}$$

input `integrate(exp(exp(x)+x)+exp(x-exp(x)),x)`

output `Timed out`

3.196.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. $2(5) = 10$.

Time = 0.18 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.83

$$\int (e^{-e^x+x} + e^{e^x+x}) dx = -e^{(-e^x)} + e^{(e^x)}$$

input `integrate(exp(exp(x)+x)+exp(x-exp(x)),x, algorithm="maxima")`

output `-e^(-e^x) + e^(e^x)`

3.196.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. $2(5) = 10$.

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.83

$$\int (e^{-e^x+x} + e^{e^x+x}) dx = -e^{(-e^x)} + e^{(e^x)}$$

input `integrate(exp(exp(x)+x)+exp(x-exp(x)),x, algorithm="giac")`

output `-e^(-e^x) + e^(e^x)`

3.196.9 Mupad [B] (verification not implemented)

Time = 16.61 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.83

$$\int (e^{-e^x+x} + e^{e^x+x}) dx = 2 \sinh(e^x)$$

input `int(exp(x + exp(x)) + exp(x - exp(x)),x)`

output `2*sinh(exp(x))`

3.197 $\int \frac{1}{1-x^2} dx$

3.197.1 Optimal result	1024
3.197.2 Mathematica [B] (verified)	1024
3.197.3 Rubi [A] (verified)	1025
3.197.4 Maple [A] (verified)	1025
3.197.5 Fricas [B] (verification not implemented)	1026
3.197.6 Sympy [B] (verification not implemented)	1026
3.197.7 Maxima [B] (verification not implemented)	1026
3.197.8 Giac [B] (verification not implemented)	1027
3.197.9 Mupad [B] (verification not implemented)	1027

3.197.1 Optimal result

Integrand size = 9, antiderivative size = 2

$$\int \frac{1}{1-x^2} dx = \operatorname{arctanh}(x)$$

output `arctanh(x)`

3.197.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 19 vs. $2(2) = 4$.

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 9.50

$$\int \frac{1}{1-x^2} dx = -\frac{1}{2} \log(1-x) + \frac{1}{2} \log(1+x)$$

input `Integrate[(1 - x^2)^(-1), x]`

output `-1/2*Log[1 - x] + Log[1 + x]/2`

3.197.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{1-x^2} dx$$

↓ 219

$$\operatorname{arctanh}(x)$$

input `Int[(1 - x^2)^(-1), x]`

output `ArcTanh[x]`

3.197.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

3.197.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\operatorname{arctanh}(x)$	3
meijerg	$\operatorname{arctanh}(x)$	3
norman	$-\frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$	14
risch	$-\frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$	14
parallelrisc	$-\frac{\ln(-1+x)}{2} + \frac{\ln(1+x)}{2}$	14

input `int(1/(-x^2+1), x, method=_RETURNVERBOSE)`

output `arctanh(x)`

3.197.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 13 vs. $2(2) = 4$.

Time = 0.23 (sec) , antiderivative size = 13, normalized size of antiderivative = 6.50

$$\int \frac{1}{1-x^2} dx = \frac{1}{2} \log(x+1) - \frac{1}{2} \log(x-1)$$

input `integrate(1/(-x^2+1),x, algorithm="fricas")`

output `1/2*log(x + 1) - 1/2*log(x - 1)`

3.197.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12 vs. $2(2) = 4$.

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 6.00

$$\int \frac{1}{1-x^2} dx = -\frac{\log(x-1)}{2} + \frac{\log(x+1)}{2}$$

input `integrate(1/(-x**2+1),x)`

output `-log(x - 1)/2 + log(x + 1)/2`

3.197.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 13 vs. $2(2) = 4$.

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 6.50

$$\int \frac{1}{1-x^2} dx = \frac{1}{2} \log(x+1) - \frac{1}{2} \log(x-1)$$

input `integrate(1/(-x^2+1),x, algorithm="maxima")`

output `1/2*log(x + 1) - 1/2*log(x - 1)`

3.197.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(2) = 4$.

Time = 0.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 7.50

$$\int \frac{1}{1-x^2} dx = \frac{1}{2} \log(|x+1|) - \frac{1}{2} \log(|x-1|)$$

input `integrate(1/(-x^2+1),x, algorithm="giac")`

output `1/2*log(abs(x + 1)) - 1/2*log(abs(x - 1))`

3.197.9 Mupad [B] (verification not implemented)

Time = 16.29 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1-x^2} dx = \operatorname{atanh}(x)$$

input `int(-1/(x^2 - 1),x)`

output `atanh(x)`

3.198 $\int 2^{\log(x)} dx$

3.198.1 Optimal result	1028
3.198.2 Mathematica [A] (verified)	1028
3.198.3 Rubi [A] (verified)	1029
3.198.4 Maple [A] (verified)	1030
3.198.5 Fricas [A] (verification not implemented)	1030
3.198.6 Sympy [A] (verification not implemented)	1030
3.198.7 Maxima [A] (verification not implemented)	1031
3.198.8 Giac [A] (verification not implemented)	1031
3.198.9 Mupad [B] (verification not implemented)	1031

3.198.1 Optimal result

Integrand size = 4, antiderivative size = 12

$$\int 2^{\log(x)} dx = \frac{2^{\log(x)} x}{1 + \log(2)}$$

output `2ln(x)*x/(1+ln(2))`

3.198.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int 2^{\log(x)} dx = \frac{2^{\log(x)} x}{1 + \log(2)}$$

input `Integrate[2Log[x],x]`

output `(2Log[x]*x)/(1 + Log[2])`

3.198.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {2704, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int 2^{\log(x)} dx \\ \downarrow 2704 \\ \int x^{\log(2)} dx \\ \downarrow 15 \\ \frac{x^{1+\log(2)}}{1+\log(2)} \end{array}$$

input `Int[2^Log[x], x]`

output `x^(1 + Log[2])/(1 + Log[2])`

3.198.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2704 `Int[(u_.)*(F_)^((a_.)*(Log[z_]*(b_.) + (v_.))), x_Symbol] := Int[u*F^(a*v)*z^(a*b*Log[F]), x] /; FreeQ[{F, a, b}, x]`

3.198.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

method	result	size
gospers	$\frac{2^{\ln(x)} x}{1+\ln(2)}$	13
risch	$\frac{x x^{\ln(2)}}{1+\ln(2)}$	13
paralelrisch	$\frac{2^{\ln(x)} x}{1+\ln(2)}$	13
norman	$\frac{x e^{\ln(x) \ln(2)}}{1+\ln(2)}$	15

input `int(2^ln(x),x,method=_RETURNVERBOSE)`output `2^ln(x)*x/(1+ln(2))`**3.198.5 Fricas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int 2^{\log(x)} dx = \frac{x e^{(\log(2) \log(x))}}{\log(2) + 1}$$

input `integrate(2^log(x),x, algorithm="fricas")`output `x*e^(log(2)*log(x))/(log(2) + 1)`**3.198.6 Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int 2^{\log(x)} dx = \frac{2^{\log(x)} x}{\log(2) + 1}$$

input `integrate(2**ln(x),x)`output `2**log(x)*x/(log(2) + 1)`

3.198.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 24, normalized size of antiderivative = 2.00

$$\int 2^{\log(x)} dx = \frac{2^{\left(\frac{1}{\log(2)}+1\right)\log(x)}}{\left(\frac{1}{\log(2)}+1\right)\log(2)}$$

input `integrate(2^log(x),x, algorithm="maxima")`output `2^((1/log(2) + 1)*log(x))/((1/log(2) + 1)*log(2))`**3.198.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int 2^{\log(x)} dx = \frac{x e^{(\log(2)\log(x))}}{\log(2) + 1}$$

input `integrate(2^log(x),x, algorithm="giac")`output `x*e^(log(2)*log(x))/(log(2) + 1)`**3.198.9 Mupad [B] (verification not implemented)**

Time = 16.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

$$\int 2^{\log(x)} dx = \frac{x^{\ln(2)+1}}{\ln(2) + 1}$$

input `int(2^log(x),x)`output `x^(log(2) + 1)/(log(2) + 1)`

3.199 $\int (\cos(3x) + \sin(2x)(\cos(3x) - \sin(2019x))) dx$

3.199.1 Optimal result	1032
3.199.2 Mathematica [A] (verified)	1032
3.199.3 Rubi [A] (verified)	1033
3.199.4 Maple [A] (verified)	1033
3.199.5 Fricas [F(-1)]	1034
3.199.6 Sympy [A] (verification not implemented)	1034
3.199.7 Maxima [A] (verification not implemented)	1034
3.199.8 Giac [A] (verification not implemented)	1035
3.199.9 Mupad [B] (verification not implemented)	1035

3.199.1 Optimal result

Integrand size = 21, antiderivative size = 39

$$\begin{aligned} & \int (\cos(3x) + \sin(2x)(\cos(3x) - \sin(2019x))) dx \\ &= \frac{\cos(x)}{2} - \frac{1}{10} \cos(5x) + \frac{1}{3} \sin(3x) - \frac{\sin(2017x)}{4034} + \frac{\sin(2021x)}{4042} \end{aligned}$$

output `1/2*cos(x)-1/10*cos(5*x)+1/3*sin(3*x)-1/4034*sin(2017*x)+1/4042*sin(2021*x)`

3.199.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

$$\begin{aligned} & \int (\cos(3x) + \sin(2x)(\cos(3x) - \sin(2019x))) dx \\ &= \frac{\cos(x)}{2} - \frac{1}{10} \cos(5x) + \frac{1}{3} \sin(3x) - \frac{\sin(2017x)}{4034} + \frac{\sin(2021x)}{4042} \end{aligned}$$

input `Integrate[Cos[3*x] + Sin[2*x]*(Cos[3*x] - Sin[2019*x]),x]`

output `Cos[x]/2 - Cos[5*x]/10 + Sin[3*x]/3 - Sin[2017*x]/4034 + Sin[2021*x]/4042`

3.199.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\cos(3x) + \sin(2x)(\cos(3x) - \sin(2019x))) dx$$

↓ 2009

$$\frac{1}{3} \sin(3x) - \frac{\sin(2017x)}{4034} + \frac{\sin(2021x)}{4042} + \frac{\cos(x)}{2} - \frac{1}{10} \cos(5x)$$

input `Int[Cos[3*x] + Sin[2*x]*(Cos[3*x] - Sin[2019*x]),x]`

output `Cos[x]/2 - Cos[5*x]/10 + Sin[3*x]/3 - Sin[2017*x]/4034 + Sin[2021*x]/4042`

3.199.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.199.4 Maple [A] (verified)

Time = 1.46 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.77

method	result	size
default	$\frac{\cos(x)}{2} - \frac{\cos(5x)}{10} + \frac{\sin(3x)}{3} - \frac{\sin(2017x)}{4034} + \frac{\sin(2021x)}{4042}$	30
risch	$\frac{\cos(x)}{2} - \frac{\cos(5x)}{10} + \frac{\sin(3x)}{3} - \frac{\sin(2017x)}{4034} + \frac{\sin(2021x)}{4042}$	30
parts	$\frac{\cos(x)}{2} - \frac{\cos(5x)}{10} + \frac{\sin(3x)}{3} - \frac{\sin(2017x)}{4034} + \frac{\sin(2021x)}{4042}$	30
parallelrisch	$-\frac{2}{5} - \frac{\sin(2017x)}{4034} + \frac{\sin(2021x)}{4042} - \frac{\cos(5x)}{10} + \frac{\sin(3x)}{3} + \frac{\cos(x)}{2}$	31

input `int(cos(3*x)+sin(2*x)*(-sin(2019*x)+cos(3*x)),x,method=_RETURNVERBOSE)`

output `1/2*cos(x)-1/10*cos(5*x)+1/3*sin(3*x)-1/4034*sin(2017*x)+1/4042*sin(2021*x)`
`)`

3.199.5 Fricas [F(-1)]

Timed out.

$$\int (\cos(3x) + \sin(2x)(\cos(3x) - \sin(2019x))) dx = \text{Timed out}$$

input `integrate(cos(3*x)+sin(2*x)*(-sin(2019*x)+cos(3*x)),x, algorithm="fricas")`

output `Timed out`

3.199.6 Sympy [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.54

$$\begin{aligned} & \int (\cos(3x) + \sin(2x)(\cos(3x) - \sin(2019x))) dx \\ &= \frac{3 \sin(2x) \sin(3x)}{5} + \frac{2019 \sin(2x) \cos(2019x)}{4076357} \\ &+ \frac{\sin(3x)}{3} - \frac{2 \sin(2019x) \cos(2x)}{4076357} + \frac{2 \cos(2x) \cos(3x)}{5} \end{aligned}$$

input `integrate(cos(3*x)+sin(2*x)*(-sin(2019*x)+cos(3*x)),x)`

output `3*sin(2*x)*sin(3*x)/5 + 2019*sin(2*x)*cos(2019*x)/4076357 + sin(3*x)/3 - 2*sin(2019*x)*cos(2*x)/4076357 + 2*cos(2*x)*cos(3*x)/5`

3.199.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.74

$$\begin{aligned} & \int (\cos(3x) + \sin(2x)(\cos(3x) - \sin(2019x))) dx \\ &= -\frac{1}{10} \cos(5x) + \frac{1}{2} \cos(x) + \frac{1}{4042} \sin(2021x) - \frac{1}{4034} \sin(2017x) + \frac{1}{3} \sin(3x) \end{aligned}$$

input `integrate(cos(3*x)+sin(2*x)*(-sin(2019*x)+cos(3*x)),x, algorithm="maxima")`

output `-1/10*cos(5*x) + 1/2*cos(x) + 1/4042*sin(2021*x) - 1/4034*sin(2017*x) + 1/3*sin(3*x)`

3.199.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.74

$$\int (\cos(3x) + \sin(2x)(\cos(3x) - \sin(2019x))) dx$$

$$= -\frac{1}{10} \cos(5x) + \frac{1}{2} \cos(x) + \frac{1}{4042} \sin(2021x) - \frac{1}{4034} \sin(2017x) + \frac{1}{3} \sin(3x)$$

input `integrate(cos(3*x)+sin(2*x)*(-sin(2019*x)+cos(3*x)),x, algorithm="giac")`output `-1/10*cos(5*x) + 1/2*cos(x) + 1/4042*sin(2021*x) - 1/4034*sin(2017*x) + 1/3*sin(3*x)`**3.199.9 Mupad [B] (verification not implemented)**

Time = 17.42 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.74

$$\int (\cos(3x) + \sin(2x)(\cos(3x) - \sin(2019x))) dx$$

$$= \frac{\sin(3x)}{3} - \frac{\cos(5x)}{10} - \frac{\sin(2017x)}{4034} + \frac{\sin(2021x)}{4042} + \frac{\cos(x)}{2}$$

input `int(cos(3*x) + sin(2*x)*(cos(3*x) - sin(2019*x)),x)`output `sin(3*x)/3 - cos(5*x)/10 - sin(2017*x)/4034 + sin(2021*x)/4042 + cos(x)/2`

3.200 $\int \cos(x) \cos(\sin(x)) \cos(\sin(\sin(x))) dx$

3.200.1 Optimal result	1036
3.200.2 Mathematica [A] (verified)	1036
3.200.3 Rubi [A] (verified)	1037
3.200.4 Maple [A] (verified)	1038
3.200.5 Fricas [B] (verification not implemented)	1038
3.200.6 Sympy [A] (verification not implemented)	1039
3.200.7 Maxima [A] (verification not implemented)	1039
3.200.8 Giac [A] (verification not implemented)	1039
3.200.9 Mupad [B] (verification not implemented)	1040

3.200.1 Optimal result

Integrand size = 10, antiderivative size = 4

$$\int \cos(x) \cos(\sin(x)) \cos(\sin(\sin(x))) dx = \sin(\sin(\sin(x)))$$

output `sin(sin(sin(x)))`

3.200.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \cos(x) \cos(\sin(x)) \cos(\sin(\sin(x))) dx = \sin(\sin(\sin(x)))$$

input `Integrate[Cos[x]*Cos[Sin[x]]*Cos[Sin[Sin[x]]],x]`

output `Sin[Sin[Sin[x]]]`

3.200.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {4834, 4834, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(x) \cos(\sin(x)) \cos(\sin(\sin(x))) dx \\
 & \quad \downarrow \text{4834} \\
 & \int \cos(\sin(x)) \cos(\sin(\sin(x))) d \sin(x) \\
 & \quad \downarrow \text{4834} \\
 & \int \cos(\sin(\sin(x))) d \sin(\sin(x)) \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(\sin(\sin(x)) + \frac{\pi}{2}\right) d \sin(\sin(x)) \\
 & \quad \downarrow \text{3117} \\
 & \sin(\sin(\sin(x)))
 \end{aligned}$$

input `Int[Cos[x]*Cos[Sin[x]]*Cos[Sin[Sin[x]]],x]`

output `Sin[Sin[Sin[x]]]`

3.200.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

```
rule 4834 Int[(u_)*(F_)[(c_.)*((a_.) + (b_.)*(x_))], x_Symbol] := With[{d = FreeFactors[Sin[c*(a + b*x)], x]}, Simp[d/(b*c) Subst[Int[SubstFor[1, Sin[c*(a + b*x)]]/d, u, x], x], x, Sin[c*(a + b*x)]/d, x] /; FunctionOfQ[Sin[c*(a + b*x)]/d, u, x, True]] /; FreeQ[{a, b, c}, x] && (EqQ[F, Cos] || EqQ[F, cos])
```

3.200.4 Maple [A] (verified)

Time = 10.37 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

method	result	size
derivativedivides	$\sin(\sin(\sin(x)))$	5
default	$\sin(\sin(\sin(x)))$	5
risch	$\sin(\sin(\sin(x)))$	5
parallelrisc	$\sin(\sin(\sin(x)))$	5

```
input int(cos(x)*cos(sin(x))*cos(sin(sin(x))),x,method=_RETURNVERBOSE)
```

```
output sin(sin(sin(x)))
```

3.200.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 41 vs. $2(4) = 8$.

Time = 0.25 (sec) , antiderivative size = 41, normalized size of antiderivative = 10.25

$$\int \cos(x) \cos(\sin(x)) \cos(\sin(\sin(x))) dx = \sin \left(\frac{2 \tan \left(\frac{\tan(\frac{1}{2}x)}{\tan(\frac{1}{2}x)^2 + 1} \right)}{\tan \left(\frac{\tan(\frac{1}{2}x)}{\tan(\frac{1}{2}x)^2 + 1} \right)^2 + 1} \right)$$

```
input integrate(cos(x)*cos(sin(x))*cos(sin(sin(x))),x, algorithm="fricas")
```

```
output sin(2*tan(tan(1/2*x)/(tan(1/2*x)^2 + 1))/(tan(tan(1/2*x)/(tan(1/2*x)^2 + 1))^2 + 1))
```

3.200.6 Sympy [A] (verification not implemented)

Time = 0.85 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

$$\int \cos(x) \cos(\sin(x)) \cos(\sin(\sin(x))) dx = \sin(\sin(\sin(x)))$$

input `integrate(cos(x)*cos(sin(x))*cos(sin(sin(x))),x)`

output `sin(sin(sin(x)))`

3.200.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \cos(x) \cos(\sin(x)) \cos(\sin(\sin(x))) dx = \sin(\sin(\sin(x)))$$

input `integrate(cos(x)*cos(sin(x))*cos(sin(sin(x))),x, algorithm="maxima")`

output `sin(sin(sin(x)))`

3.200.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \cos(x) \cos(\sin(x)) \cos(\sin(\sin(x))) dx = \sin(\sin(\sin(x)))$$

input `integrate(cos(x)*cos(sin(x))*cos(sin(sin(x))),x, algorithm="giac")`

output `sin(sin(sin(x)))`

3.200.9 Mupad [B] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \cos(x) \cos(\sin(x)) \cos(\sin(\sin(x))) dx = \sin(\sin(\sin(x)))$$

input `int(cos(sin(x))*cos(sin(sin(x)))*cos(x),x)`

output `sin(sin(sin(x)))`

$$\mathbf{3.201} \quad \int \frac{e^{-\frac{2019}{4x^2}}}{x^2} dx$$

3.201.1 Optimal result1041
3.201.2 Mathematica [A] (verified)1041
3.201.3 Rubi [A] (verified)	1042
3.201.4 Maple [A] (verified)	1043
3.201.5 Fricas [A] (verification not implemented)	1043
3.201.6 Sympy [A] (verification not implemented)	1043
3.201.7 Maxima [A] (verification not implemented)	1044
3.201.8 Giac [F]	1044
3.201.9 Mupad [B] (verification not implemented)	1044

3.201.1 Optimal result

Integrand size = 13, antiderivative size = 24

$$\int \frac{e^{-\frac{2019}{4x^2}}}{x^2} dx = -\sqrt{\frac{\pi}{2019}} \operatorname{erf}\left(\frac{\sqrt{2019}}{2x}\right)$$

output `-1/2019*2019^(1/2)*Pi^(1/2)*erf(1/2*2019^(1/2)/x)`

3.201.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{e^{-\frac{2019}{4x^2}}}{x^2} dx = -\sqrt{\frac{\pi}{2019}} \operatorname{erf}\left(\frac{\sqrt{2019}}{2x}\right)$$

input `Integrate[1/(E^(2019/(4*x^2))*x^2),x]`

output `-(Sqrt[Pi/2019]*Erf[Sqrt[2019]/(2*x)])`

3.201.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {2640, 2634}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{-\frac{2019}{4x^2}}}{x^2} dx$$

↓ 2640

$$-\int e^{-\frac{2019}{4x^2}} d\frac{1}{x}$$

↓ 2634

$$-\sqrt{\frac{\pi}{2019}} \operatorname{erf}\left(\frac{\sqrt{2019}}{2x}\right)$$

input `Int[1/(E^(2019/(4*x^2)))*x^2),x]`

output `-(Sqrt[Pi/2019]*Erf[Sqrt[2019]/(2*x)])`

3.201.3.1 Defintions of rubi rules used

rule 2634 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(2)), x_Symbol] := Simp[F^a*Sqrt[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && NegQ[b]`

rule 2640 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[1/(d*(m + 1)) Subst[Int[F^(a + b*x^2), x], x, (c + d*x)^(m + 1)], x] /; FreeQ[{F, a, b, c, d, m, n}, x] && EqQ[n, 2*(m + 1)]`

3.201.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.75

method	result	size
derivativedivides	$-\frac{\sqrt{2019}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{2019}}{2x}\right)}{2019}$	18
default	$-\frac{\sqrt{2019}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{2019}}{2x}\right)}{2019}$	18
meijerg	$-\frac{\sqrt{2019}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{2019}}{2x}\right)}{2019}$	18
risch	$-\frac{\sqrt{2019}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{2019}}{2x}\right)}{2019}$	18

input `int(exp(-2019/4/x^2)/x^2,x,method=_RETURNVERBOSE)`output `-1/2019*2019^(1/2)*Pi^(1/2)*erf(1/2*2019^(1/2)/x)`**3.201.5 Fricas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.71

$$\int \frac{e^{-\frac{2019}{4x^2}}}{x^2} dx = -\frac{1}{2019} \sqrt{2019}\sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{2019}}{2x}\right)$$

input `integrate(exp(-2019/4/x^2)/x^2,x, algorithm="fricas")`output `-1/2019*sqrt(2019)*sqrt(pi)*erf(1/2*sqrt(2019)/x)`**3.201.6 Sympy [A] (verification not implemented)**

Time = 0.39 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{e^{-\frac{2019}{4x^2}}}{x^2} dx = -\frac{\sqrt{2019}\sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{2019}}{2x}\right)}{2019}$$

input `integrate(exp(-2019/4/x**2)/x**2,x)`output `-sqrt(2019)*sqrt(pi)*erf(sqrt(2019)/(2*x))/2019`

3.201. $\int \frac{e^{-\frac{2019}{4x^2}}}{x^2} dx$

3.201.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21

$$\int \frac{e^{-\frac{2019}{4x^2}}}{x^2} dx = -\frac{\sqrt{2019}\sqrt{\pi}\sqrt{x^2}\left(\operatorname{erf}\left(\frac{1}{2}\sqrt{2019}\sqrt{\frac{1}{x^2}}\right) - 1\right)}{2019x}$$

input `integrate(exp(-2019/4/x^2)/x^2,x, algorithm="maxima")`output `-1/2019*sqrt(2019)*sqrt(pi)*sqrt(x^2)*(erf(1/2*sqrt(2019)*sqrt(x^(-2)))) - 1)/x`**3.201.8 Giac [F]**

$$\int \frac{e^{-\frac{2019}{4x^2}}}{x^2} dx = \int \frac{e^{\left(-\frac{2019}{4x^2}\right)}}{x^2} dx$$

input `integrate(exp(-2019/4/x^2)/x^2,x, algorithm="giac")`output `integrate(e^(-2019/4/x^2)/x^2, x)`**3.201.9 Mupad [B] (verification not implemented)**

Time = 15.66 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.71

$$\int \frac{e^{-\frac{2019}{4x^2}}}{x^2} dx = -\frac{\sqrt{2019}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{2019}}{2x}\right)}{2019}$$

input `int(exp(-2019/(4*x^2))/x^2,x)`output `-(2019^(1/2)*pi^(1/2)*erf(2019^(1/2)/(2*x)))/2019`

3.202 $\int \sin(\sqrt{x}) dx$

3.202.1 Optimal result	1045
3.202.2 Mathematica [A] (verified)	1045
3.202.3 Rubi [A] (verified)	1046
3.202.4 Maple [A] (verified)	1047
3.202.5 Fricas [A] (verification not implemented)	1048
3.202.6 Sympy [A] (verification not implemented)	1048
3.202.7 Maxima [A] (verification not implemented)	1048
3.202.8 Giac [A] (verification not implemented)	1049
3.202.9 Mupad [B] (verification not implemented)	1049

3.202.1 Optimal result

Integrand size = 6, antiderivative size = 22

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

output `-2*x^(1/2)*cos(x^(1/2))+2*sin(x^(1/2))`

3.202.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

input `Integrate[Sin[Sqrt[x]],x]`

output `-2*Sqrt[x]*Cos[Sqrt[x]] + 2*Sin[Sqrt[x]]`

3.202.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {3842, 3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(\sqrt{x}) \, dx \\
 & \quad \downarrow \text{3842} \\
 & 2 \int \sqrt{x} \sin(\sqrt{x}) \, d\sqrt{x} \\
 & \quad \downarrow \text{3042} \\
 & 2 \int \sqrt{x} \sin(\sqrt{x}) \, d\sqrt{x} \\
 & \quad \downarrow \text{3777} \\
 & 2 \left(\int \cos(\sqrt{x}) \, d\sqrt{x} - \sqrt{x} \cos(\sqrt{x}) \right) \\
 & \quad \downarrow \text{3042} \\
 & 2 \left(\int \sin\left(\sqrt{x} + \frac{\pi}{2}\right) \, d\sqrt{x} - \sqrt{x} \cos(\sqrt{x}) \right) \\
 & \quad \downarrow \text{3117} \\
 & 2(\sin(\sqrt{x}) - \sqrt{x} \cos(\sqrt{x}))
 \end{aligned}$$

input `Int[Sin[Sqrt[x]], x]`

output `2*(-(Sqrt[x]*Cos[Sqrt[x]]) + Sin[Sqrt[x]])`

3.202.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3842 `Int[((a_.) + (b_.)*Sin[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)])^(p_.), x_Symbol] := Simp[1/(n*f) Subst[Int[x^(1/n - 1)*(a + b*SIN[c + d*x])^p, x], (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]`

3.202.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

method	result	size
derivativedivides	$-2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$	17
default	$-2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$	17
meijerg	$4\sqrt{\pi} \left(-\frac{\sqrt{x} \cos(\sqrt{x})}{2\sqrt{\pi}} + \frac{\sin(\sqrt{x})}{2\sqrt{\pi}} \right)$	28

input `int(sin(x^(1/2)),x,method=_RETURNVERBOSE)`

output $-2*x^{(1/2)}*\cos(x^{(1/2)})+2*\sin(x^{(1/2)})$

3.202.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

input `integrate(sin(x^(1/2)),x, algorithm="fricas")`

output `-2*sqrt(x)*cos(sqrt(x)) + 2*sin(sqrt(x))`

3.202.6 Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

input `integrate(sin(x**(1/2)),x)`

output `-2*sqrt(x)*cos(sqrt(x)) + 2*sin(sqrt(x))`

3.202.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

input `integrate(sin(x^(1/2)),x, algorithm="maxima")`

output `-2*sqrt(x)*cos(sqrt(x)) + 2*sin(sqrt(x))`

3.202.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \sin(\sqrt{x}) dx = -2\sqrt{x} \cos(\sqrt{x}) + 2 \sin(\sqrt{x})$$

input `integrate(sin(x^(1/2)),x, algorithm="giac")`

output `-2*sqrt(x)*cos(sqrt(x)) + 2*sin(sqrt(x))`

3.202.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \sin(\sqrt{x}) dx = 2 \sin(\sqrt{x}) - 2\sqrt{x} \cos(\sqrt{x})$$

input `int(sin(x^(1/2)),x)`

output `2*sin(x^(1/2)) - 2*x^(1/2)*cos(x^(1/2))`

3.203 $\int \frac{\sqrt{x}}{1+x} dx$

3.203.1 Optimal result	1050
3.203.2 Mathematica [A] (verified)	1050
3.203.3 Rubi [A] (verified)	1051
3.203.4 Maple [A] (verified)	1052
3.203.5 Fricas [A] (verification not implemented)	1052
3.203.6 Sympy [A] (verification not implemented)	1053
3.203.7 Maxima [A] (verification not implemented)	1053
3.203.8 Giac [A] (verification not implemented)	1053
3.203.9 Mupad [B] (verification not implemented)	1054

3.203.1 Optimal result

Integrand size = 11, antiderivative size = 16

$$\int \frac{\sqrt{x}}{1+x} dx = 2\sqrt{x} - 2 \arctan(\sqrt{x})$$

output `2*x^(1/2)-2*arctan(x^(1/2))`

3.203.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{x}}{1+x} dx = 2\sqrt{x} - 2 \arctan(\sqrt{x})$$

input `Integrate[Sqrt[x]/(1 + x),x]`

output `2*Sqrt[x] - 2*ArcTan[Sqrt[x]]`

3.203.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{x}}{x+1} dx \\ & \quad \downarrow \text{60} \\ & 2\sqrt{x} - \int \frac{1}{\sqrt{x}(x+1)} dx \\ & \quad \downarrow \text{73} \\ & 2\sqrt{x} - 2 \int \frac{1}{x+1} d\sqrt{x} \\ & \quad \downarrow \text{216} \\ & 2\sqrt{x} - 2 \arctan(\sqrt{x}) \end{aligned}$$

input `Int[Sqrt[x]/(1 + x),x]`

output `2*Sqrt[x] - 2*ArcTan[Sqrt[x]]`

3.203.3.1 Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

3.203.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$2\sqrt{x} - 2 \arctan(\sqrt{x})$	13
default	$2\sqrt{x} - 2 \arctan(\sqrt{x})$	13
meijerg	$2\sqrt{x} - 2 \arctan(\sqrt{x})$	13
risch	$2\sqrt{x} - 2 \arctan(\sqrt{x})$	13
trager	$2\sqrt{x} + \text{RootOf}(-Z^2 + 1) \ln\left(-\frac{2\text{RootOf}(-Z^2 + 1)\sqrt{x-x+1}}{1+x}\right)$	38

```
input int(x^(1/2)/(1+x),x,method=_RETURNVERBOSE)
```

```
output 2*x^(1/2)-2*arctan(x^(1/2))
```

3.203.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{x}}{1+x} dx = 2\sqrt{x} - 2 \arctan(\sqrt{x})$$

```
input integrate(x^(1/2)/(1+x),x, algorithm="fricas")
```

```
output 2*sqrt(x) - 2*arctan(sqrt(x))
```

3.203.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{\sqrt{x}}{1+x} dx = 2\sqrt{x} - 2 \operatorname{atan}(\sqrt{x})$$

input `integrate(x**(1/2)/(1+x),x)`

output `2*sqrt(x) - 2*atan(sqrt(x))`

3.203.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{x}}{1+x} dx = 2\sqrt{x} - 2 \operatorname{arctan}(\sqrt{x})$$

input `integrate(x^(1/2)/(1+x),x, algorithm="maxima")`

output `2*sqrt(x) - 2*arctan(sqrt(x))`

3.203.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{x}}{1+x} dx = 2\sqrt{x} - 2 \operatorname{arctan}(\sqrt{x})$$

input `integrate(x^(1/2)/(1+x),x, algorithm="giac")`

output `2*sqrt(x) - 2*arctan(sqrt(x))`

3.203.9 Mupad [B] (verification not implemented)

Time = 15.91 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{x}}{1+x} dx = 2\sqrt{x} - 2 \operatorname{atan}(\sqrt{x})$$

input `int(x^(1/2)/(x + 1),x)`

output `2*x^(1/2) - 2*atan(x^(1/2))`

3.204 $\int \cos(x) \cos(2x) \cos(3x) dx$

3.204.1 Optimal result	1055
3.204.2 Mathematica [A] (verified)	1055
3.204.3 Rubi [A] (verified)	1056
3.204.4 Maple [A] (verified)	1057
3.204.5 Fricas [A] (verification not implemented)	1057
3.204.6 Sympy [B] (verification not implemented)	1057
3.204.7 Maxima [A] (verification not implemented)	1058
3.204.8 Giac [A] (verification not implemented)	1058
3.204.9 Mupad [B] (verification not implemented)	1058

3.204.1 Optimal result

Integrand size = 11, antiderivative size = 30

$$\int \cos(x) \cos(2x) \cos(3x) dx = \frac{x}{4} + \frac{1}{8} \sin(2x) + \frac{1}{16} \sin(4x) + \frac{1}{24} \sin(6x)$$

output `1/4*x+1/8*sin(2*x)+1/16*sin(4*x)+1/24*sin(6*x)`

3.204.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \cos(x) \cos(2x) \cos(3x) dx = \frac{x}{4} + \frac{1}{8} \sin(2x) + \frac{1}{16} \sin(4x) + \frac{1}{24} \sin(6x)$$

input `Integrate[Cos[x]*Cos[2*x]*Cos[3*x],x]`

output `x/4 + Sin[2*x]/8 + Sin[4*x]/16 + Sin[6*x]/24`

3.204.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 4855, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(x) \cos(2x) \cos(3x) dx$$

$$\downarrow \text{3042}$$

$$\int \cos(x) \cos(2x) \cos(3x) dx$$

$$\downarrow \text{4855}$$

$$\int \left(\frac{1}{4} \cos(2x) + \frac{1}{4} \cos(4x) + \frac{1}{4} \cos(6x) + \frac{1}{4} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{x}{4} + \frac{1}{8} \sin(2x) + \frac{1}{16} \sin(4x) + \frac{1}{24} \sin(6x)$$

input `Int[Cos[x]*Cos[2*x]*Cos[3*x],x]`

output `x/4 + Sin[2*x]/8 + Sin[4*x]/16 + Sin[6*x]/24`

3.204.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4855 `Int[(F_)[(a_.) + (b_.)*(x_)]^(p_.)*(G_)[(c_.) + (d_.)*(x_)]^(q_.)*(H_)[(e_.) + (f_.)*(x_)]^(r_.), x_Symbol] := Int[ExpandTrigReduce[ActivateTrig[F[a + b*x]^p*G[c + d*x]^q*H[e + f*x]^r], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && (EqQ[F, sin] || EqQ[F, cos]) && (EqQ[G, sin] || EqQ[G, cos]) && (EqQ[H, sin] || EqQ[H, cos]) && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]`

3.204.4 Maple [A] (verified)

Time = 0.85 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.77

method	result	size
default	$\frac{x}{4} + \frac{\sin(2x)}{8} + \frac{\sin(4x)}{16} + \frac{\sin(6x)}{24}$	23
risch	$\frac{x}{4} + \frac{\sin(2x)}{8} + \frac{\sin(4x)}{16} + \frac{\sin(6x)}{24}$	23
parallelrisch	$\frac{x}{4} + \frac{\sin(2x)}{8} + \frac{\sin(4x)}{16} + \frac{\sin(6x)}{24}$	23

input `int(cos(x)*cos(2*x)*cos(3*x),x,method=_RETURNVERBOSE)`output `1/4*x+1/8*sin(2*x)+1/16*sin(4*x)+1/24*sin(6*x)`**3.204.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.83

$$\int \cos(x) \cos(2x) \cos(3x) dx = \frac{1}{12} (16 \cos(x)^5 - 10 \cos(x)^3 + 3 \cos(x)) \sin(x) + \frac{1}{4} x$$

input `integrate(cos(x)*cos(2*x)*cos(3*x),x, algorithm="fricas")`output `1/12*(16*cos(x)^5 - 10*cos(x)^3 + 3*cos(x))*sin(x) + 1/4*x`**3.204.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 114 vs. 2(22) = 44.

Time = 0.89 (sec) , antiderivative size = 114, normalized size of antiderivative = 3.80

$$\begin{aligned} \int \cos(x) \cos(2x) \cos(3x) dx = & -\frac{x \sin(x) \sin(2x) \cos(3x)}{4} + \frac{x \sin(x) \sin(3x) \cos(2x)}{4} \\ & + \frac{x \sin(2x) \sin(3x) \cos(x)}{4} + \frac{x \cos(x) \cos(2x) \cos(3x)}{4} \\ & + \frac{\sin(x) \sin(2x) \sin(3x)}{6} + \frac{\sin(x) \cos(2x) \cos(3x)}{8} \\ & + \frac{5 \sin(3x) \cos(x) \cos(2x)}{24} \end{aligned}$$

input `integrate(cos(x)*cos(2*x)*cos(3*x),x)`

output `-x*sin(x)*sin(2*x)*cos(3*x)/4 + x*sin(x)*sin(3*x)*cos(2*x)/4 + x*sin(2*x)*sin(3*x)*cos(x)/4 + x*cos(x)*cos(2*x)*cos(3*x)/4 + sin(x)*sin(2*x)*sin(3*x)/6 + sin(x)*cos(2*x)*cos(3*x)/8 + 5*sin(3*x)*cos(x)*cos(2*x)/24`

3.204.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \cos(x) \cos(2x) \cos(3x) dx = \frac{1}{4}x + \frac{1}{24} \sin(6x) + \frac{1}{16} \sin(4x) + \frac{1}{8} \sin(2x)$$

input `integrate(cos(x)*cos(2*x)*cos(3*x),x, algorithm="maxima")`

output `1/4*x + 1/24*sin(6*x) + 1/16*sin(4*x) + 1/8*sin(2*x)`

3.204.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \cos(x) \cos(2x) \cos(3x) dx = \frac{1}{4}x + \frac{1}{24} \sin(6x) + \frac{1}{16} \sin(4x) + \frac{1}{8} \sin(2x)$$

input `integrate(cos(x)*cos(2*x)*cos(3*x),x, algorithm="giac")`

output `1/4*x + 1/24*sin(6*x) + 1/16*sin(4*x) + 1/8*sin(2*x)`

3.204.9 Mupad [B] (verification not implemented)

Time = 16.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \cos(x) \cos(2x) \cos(3x) dx = \frac{x}{4} + \frac{\sin(2x)}{8} + \frac{\sin(4x)}{16} + \frac{\sin(6x)}{24}$$

input `int(cos(2*x)*cos(3*x)*cos(x),x)`

output `x/4 + sin(2*x)/8 + sin(4*x)/16 + sin(6*x)/24`

3.205 $\int e^{-x^{2n}} dx$

3.205.1 Optimal result	1059
3.205.2 Mathematica [A] (verified)	1059
3.205.3 Rubi [A] (verified)	1060
3.205.4 Maple [C] (verified)	1060
3.205.5 Fricas [F]	1061
3.205.6 Sympy [A] (verification not implemented)	1061
3.205.7 Maxima [A] (verification not implemented)	1061
3.205.8 Giac [F]	1062
3.205.9 Mupad [F(-1)]	1062

3.205.1 Optimal result

Integrand size = 9, antiderivative size = 23

$$\int e^{-x^{2n}} dx = -\frac{x \operatorname{ExpIntegralE}\left(1 - \frac{1}{2n}, x^{2n}\right)}{2n}$$

output `-1/2*x*Ei(1-1/2/n,x^(2*n))/n`

3.205.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.48

$$\int e^{-x^{2n}} dx = -\frac{x(x^{2n})^{-\frac{1}{2}/n} \Gamma\left(\frac{1}{2n}, x^{2n}\right)}{2n}$$

input `Integrate[E^(-x^(2*n)),x]`

output `-1/2*(x*Gamma[1/(2*n), x^(2*n)])/(n*(x^(2*n))^(1/(2*n)))`

3.205.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.48, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {2637}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{-x^{2n}} dx$$

↓ 2637

$$-\frac{x(x^{2n})^{-\frac{1}{2}/n} \Gamma(\frac{1}{2n}, x^{2n})}{2n}$$

input `Int [E^(-x^(2*n)), x]`

output `-1/2*(x*Gamma[1/(2*n), x^(2*n)])/(n*(x^(2*n))^(1/(2*n)))`

3.205.3.1 Defintions of rubi rules used

rule 2637 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(-F^a)*(c + d*x)*(Gamma[1/n, (-b)*(c + d*x)^n*Log[F]]/(d*n*((-b)*(c + d*x)^n*Log[F]))^(1/n)), x] /; FreeQ[{F, a, b, c, d, n}, x] && !IntegerQ[2/n]`

3.205.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.06 (sec) , antiderivative size = 185, normalized size of antiderivative = 8.04

method	result
meijerg	$\frac{4n^2 x^{-2n+1} (2n x^{2n} + 2n + 1) (x^{2n})^{-\frac{2n+1}{4n}} e^{-\frac{x^{2n}}{2}} \text{WhittakerM}\left(\frac{1}{2n} - \frac{2n+1}{4n}, \frac{2n+1}{4n} + \frac{1}{2}, x^{2n}\right) + 2n x^{-2n+1} (2n+1) (x^{2n})^{-\frac{2n+1}{4n}} e^{-\frac{x^{2n}}{2}} \text{WhittakerM}\left(\frac{1}{2n} - \frac{2n+1}{4n}, \frac{2n+1}{4n} + \frac{1}{2}, x^{2n}\right)}{(2n+1)(4n+1) 2n 4n+1}$

input `int(exp(-x^(2*n)), x, method=_RETURNVERBOSE)`

```
output 1/2/n*(4*n^2*x^(-2*n+1)*(2*n*x^(2*n)+2*n+1)/(2*n+1)/(4*n+1)*(x^(2*n))^(-1/
4*(2*n+1)/n)*exp(-1/2*x^(2*n))*WhittakerM(1/2/n-1/4*(2*n+1)/n,1/4*(2*n+1)/
n+1/2,x^(2*n))+2*n*x^(-2*n+1)*(2*n+1)/(4*n+1)*(x^(2*n))^(-1/4*(2*n+1)/n)*e
xp(-1/2*x^(2*n))*WhittakerM(1/2/n-1/4*(2*n+1)/n+1,1/4*(2*n+1)/n+1/2,x^(2*n
)))
```

3.205.5 Fricas [F]

$$\int e^{-x^{2n}} dx = \int e^{(-x^{2n})} dx$$

```
input integrate(exp(-x^(2*n)),x, algorithm="fricas")
```

```
output integral(e^(-x^(2*n)), x)
```

3.205.6 Sympy [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.26

$$\int e^{-x^{2n}} dx = \frac{\Gamma\left(\frac{1}{2n}\right) \gamma\left(\frac{1}{2n}, x^{2n}\right)}{4n^2 \Gamma\left(1 + \frac{1}{2n}\right)}$$

```
input integrate(exp(-x**(2*n)),x)
```

```
output gamma(1/(2*n))*lowergamma(1/(2*n), x**(2*n))/(4*n**2*gamma(1 + 1/(2*n)))
```

3.205.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.30

$$\int e^{-x^{2n}} dx = -\frac{x \Gamma\left(\frac{1}{2n}, x^{2n}\right)}{2n(x^{2n})^{\frac{1}{2n}}}$$

```
input integrate(exp(-x^(2*n)),x, algorithm="maxima")
```

```
output -1/2*x*gamma(1/2/n, x^(2*n))/(n*(x^(2*n))^(1/2/n))
```

3.205.8 Giac [F]

$$\int e^{-x^{2n}} dx = \int e^{(-x^{2n})} dx$$

input `integrate(exp(-x^(2*n)),x, algorithm="giac")`

output `integrate(e^(-x^(2*n)), x)`

3.205.9 Mupad [F(-1)]

Timed out.

$$\int e^{-x^{2n}} dx = \int e^{-x^{2n}} dx$$

input `int(exp(-x^(2*n)),x)`

output `int(exp(-x^(2*n)), x)`

3.206 $\int e dx$

3.206.1 Optimal result	1063
3.206.2 Mathematica [A] (verified)	1063
3.206.3 Rubi [A] (verified)	1064
3.206.4 Maple [A] (verified)	1064
3.206.5 Fricas [A] (verification not implemented)	1065
3.206.6 Sympy [A] (verification not implemented)	1065
3.206.7 Maxima [A] (verification not implemented)	1065
3.206.8 Giac [A] (verification not implemented)	1066
3.206.9 Mupad [B] (verification not implemented)	1066

3.206.1 Optimal result

Integrand size = 1, antiderivative size = 3

$$\int e dx = ex$$

output `x*exp(1)`

3.206.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int e dx = ex$$

input `Integrate[E,x]`

output `E*x`

3.206.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e dx$$

$$\downarrow 24$$

$$ex$$

input `Int [E, x]`

output `E*x`

3.206.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

3.206.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 3.00

method	result	size
parallelrisc	$x^{\frac{1}{\ln(x)}} x$	9

input `int(x^(1/ln(x)), x, method=_RETURNVERBOSE)`

output `x^(1/ln(x))*x`

3.206.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

$$\int e dx = xe$$

input `integrate(x^(1/log(x)),x, algorithm="fricas")`

output `x*e`

3.206.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int e dx = ex$$

input `integrate(x**(1/ln(x)),x)`

output `E*x`

3.206.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

$$\int e dx = xe$$

input `integrate(x^(1/log(x)),x, algorithm="maxima")`

output `x*e`

3.206.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

$$\int e dx = x e$$

input `integrate(x^(1/log(x)),x, algorithm="giac")`

output `x*e`

3.206.9 Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

$$\int e dx = x e$$

input `int(x^(1/log(x)),x)`

output `x*exp(1)`

3.207 $\int \frac{\sin(19x)+\sin(20x)}{\cos(19x)+\cos(20x)} dx$

3.207.1 Optimal result	1067
3.207.2 Mathematica [B] (verified)	1067
3.207.3 Rubi [B] (verified)	1068
3.207.4 Maple [C] (verified)	1069
3.207.5 Fricas [B] (verification not implemented)	1070
3.207.6 Sympy [F(-1)]	1070
3.207.7 Maxima [B] (verification not implemented)	1071
3.207.8 Giac [B] (verification not implemented)	1072
3.207.9 Mupad [B] (verification not implemented)	1072

3.207.1 Optimal result

Integrand size = 21, antiderivative size = 11

$$\int \frac{\sin(19x) + \sin(20x)}{\cos(19x) + \cos(20x)} dx = -\frac{2}{39} \log\left(\cos\left(\frac{39x}{2}\right)\right)$$

output `-2/39*ln(cos(39/2*x))`

3.207.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 131 vs. 2(11) = 22.

Time = 0.42 (sec) , antiderivative size = 131, normalized size of antiderivative = 11.91

$$\begin{aligned} \int \frac{\sin(19x) + \sin(20x)}{\cos(19x) + \cos(20x)} dx = & -\frac{2}{39} \log\left(\cos\left(\frac{x}{2}\right)\right) - \frac{2}{39} \log(1 - 2 \cos(x) + 2 \cos(2x)) \\ & - 2 \cos(3x) + 2 \cos(4x) - 2 \cos(5x) + 2 \cos(6x) - 2 \cos(7x) \\ & + 2 \cos(8x) - 2 \cos(9x) + 2 \cos(10x) - 2 \cos(11x) + 2 \cos(12x) \\ & - 2 \cos(13x) + 2 \cos(14x) - 2 \cos(15x) + 2 \cos(16x) \\ & - 2 \cos(17x) + 2 \cos(18x) - 2 \cos(19x) \end{aligned}$$

input `Integrate[(Sin[19*x] + Sin[20*x])/(Cos[19*x] + Cos[20*x]),x]`

output `(-2*Log[Cos[x/2]])/39 - (2*Log[1 - 2*Cos[x] + 2*Cos[2*x] - 2*Cos[3*x] + 2*Cos[4*x] - 2*Cos[5*x] + 2*Cos[6*x] - 2*Cos[7*x] + 2*Cos[8*x] - 2*Cos[9*x] + 2*Cos[10*x] - 2*Cos[11*x] + 2*Cos[12*x] - 2*Cos[13*x] + 2*Cos[14*x] - 2*Cos[15*x] + 2*Cos[16*x] - 2*Cos[17*x] + 2*Cos[18*x] - 2*Cos[19*x]])/39`

3.207.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 139 vs. $2(11) = 22$.

Time = 2.41 (sec) , antiderivative size = 139, normalized size of antiderivative = 12.64, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 4901, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sin(19x) + \sin(20x)}{\cos(19x) + \cos(20x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sin(19x) + \sin(20x)}{\cos(19x) + \cos(20x)} dx \\ & \quad \downarrow \text{4901} \\ & \int \left(\frac{\sin(19x)}{\cos(19x) + \cos(20x)} + \frac{\sin(20x)}{\cos(19x) + \cos(20x)} \right) dx \\ & \quad \downarrow \text{2009} \\ & -\frac{2}{39} \log(-64 \cos^6(x) + 32 \cos^5(x) + 80 \cos^4(x) - 32 \cos^3(x) - 24 \cos^2(x) + 6 \cos(x) + 1) - \\ & \quad \frac{2}{39} \log(1 - 2 \cos(x)) - \frac{1}{39} \log(\cos(x) + 1) - \\ & \quad \frac{2}{39} \log(4096 \cos^{12}(x) + 2048 \cos^{11}(x) - 12288 \cos^{10}(x) - 6144 \cos^9(x) + 13568 \cos^8(x) + 6784 \cos^7(x) - 6592 \cos^6(x) \end{aligned}$$

input `Int[(Sin[19*x] + Sin[20*x])/(Cos[19*x] + Cos[20*x]),x]`

output `(-2*Log[1 - 2*Cos[x]])/39 - Log[1 + Cos[x]]/39 - (2*Log[1 + 6*Cos[x] - 24*Cos[x]^2 - 32*Cos[x]^3 + 80*Cos[x]^4 + 32*Cos[x]^5 - 64*Cos[x]^6])/39 - (2*Log[1 - 24*Cos[x] - 48*Cos[x]^2 + 632*Cos[x]^3 + 1264*Cos[x]^4 - 3296*Cos[x]^5 - 6592*Cos[x]^6 + 6784*Cos[x]^7 + 13568*Cos[x]^8 - 6144*Cos[x]^9 - 12288*Cos[x]^10 + 2048*Cos[x]^11 + 4096*Cos[x]^12])/39`

3.207.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4901 `Int[u_, x_Symbol] := With[{v = ExpandTrig[u, x]}, Int[v, x] /; SumQ[v]] /; !InertTrigFreeQ[u]`

3.207.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.64 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.45

method	result	size
risch	$ix - \frac{2 \ln(e^{39ix} + 1)}{39}$	16
parallelrisc	$\ln\left(\left(\sec(10x)^2\right)^{\frac{1}{39}}\right) + \ln\left(\left(\sec\left(\frac{19x}{2}\right)^2\right)^{\frac{1}{39}}\right) + \ln\left(\frac{1}{\left(\tan\left(\frac{19x}{2}\right)\tan(10x)-1\right)^{\frac{2}{39}}}\right)$	34

input `int((sin(20*x)+sin(19*x))/(cos(20*x)+cos(19*x)),x,method=_RETURNVERBOSE)`

output `I*x-2/39*ln(exp(39*I*x)+1)`

3.207.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 123 vs. $2(7) = 14$.

Time = 0.34 (sec) , antiderivative size = 123, normalized size of antiderivative = 11.18

$$\int \frac{\sin(19x) + \sin(20x)}{\cos(19x) + \cos(20x)} dx = -\frac{1}{39} \log \left(137438953472 \cos(x)^{39} - 1340029796352 \cos(x)^{37} \right. \\ \left. + 6030134083584 \cos(x)^{35} - 16610786017280 \cos(x)^{33} \right. \\ \left. + 31323196489728 \cos(x)^{31} - 42839077552128 \cos(x)^{29} \right. \\ \left. + 43920872439808 \cos(x)^{27} - 34411219255296 \cos(x)^{25} \right. \\ \left. + 20813237452800 \cos(x)^{23} - 9751387176960 \cos(x)^{21} \right. \\ \left. + 3530674667520 \cos(x)^{19} - 980106117120 \cos(x)^{17} \right. \\ \left. + 205701283840 \cos(x)^{15} - 31950643200 \cos(x)^{13} \right. \\ \left. + 3560214528 \cos(x)^{11} - 271960832 \cos(x)^9 \right. \\ \left. + 13302432 \cos(x)^7 - 373464 \cos(x)^5 + 4940 \cos(x)^3 \right. \\ \left. - \frac{39}{2} \cos(x) + \frac{1}{2} \right)$$

```
input integrate((sin(20*x)+sin(19*x))/(cos(20*x)+cos(19*x)),x, algorithm="fricas")
```

```
output -1/39*log(137438953472*cos(x)^39 - 1340029796352*cos(x)^37 + 6030134083584
*cos(x)^35 - 16610786017280*cos(x)^33 + 31323196489728*cos(x)^31 - 4283907
7552128*cos(x)^29 + 43920872439808*cos(x)^27 - 34411219255296*cos(x)^25 +
20813237452800*cos(x)^23 - 9751387176960*cos(x)^21 + 3530674667520*cos(x)^
19 - 980106117120*cos(x)^17 + 205701283840*cos(x)^15 - 31950643200*cos(x)^
13 + 3560214528*cos(x)^11 - 271960832*cos(x)^9 + 13302432*cos(x)^7 - 37346
4*cos(x)^5 + 4940*cos(x)^3 - 39/2*cos(x) + 1/2)
```

3.207.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sin(19x) + \sin(20x)}{\cos(19x) + \cos(20x)} dx = \text{Timed out}$$

```
input integrate((sin(20*x)+sin(19*x))/(cos(20*x)+cos(19*x)),x)
```

```
output Timed out
```

3.207.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2527 vs. $2(7) = 14$.

Time = 0.37 (sec) , antiderivative size = 2527, normalized size of antiderivative = 229.73

$$\int \frac{\sin(19x) + \sin(20x)}{\cos(19x) + \cos(20x)} dx = \text{Too large to display}$$

```
input integrate((sin(20*x)+sin(19*x))/(cos(20*x)+cos(19*x)),x, algorithm="maxima")
```

```
output -1/39*log(2*(cos(23*x) - cos(21*x) - cos(20*x) + cos(18*x) + cos(17*x) - c
os(15*x) - cos(14*x) + cos(12*x) - cos(10*x) - cos(9*x) + cos(7*x) + cos(6
*x) - cos(4*x) - cos(3*x) + cos(x) + 1)*cos(24*x) + cos(24*x)^2 - 2*(cos(2
1*x) + cos(20*x) - cos(18*x) - cos(17*x) + cos(15*x) + cos(14*x) - cos(12*
x) + cos(10*x) + cos(9*x) - cos(7*x) - cos(6*x) + cos(4*x) + cos(3*x) - co
s(x) - 1)*cos(23*x) + cos(23*x)^2 + 2*(cos(20*x) - cos(18*x) - cos(17*x) +
cos(15*x) + cos(14*x) - cos(12*x) + cos(10*x) + cos(9*x) - cos(7*x) - cos
(6*x) + cos(4*x) + cos(3*x) - cos(x) - 1)*cos(21*x) + cos(21*x)^2 - 2*(cos
(18*x) + cos(17*x) - cos(15*x) - cos(14*x) + cos(12*x) - cos(10*x) - cos(9
*x) + cos(7*x) + cos(6*x) - cos(4*x) - cos(3*x) + cos(x) + 1)*cos(20*x) +
cos(20*x)^2 + 2*(cos(17*x) - cos(15*x) - cos(14*x) + cos(12*x) - cos(10*x)
- cos(9*x) + cos(7*x) + cos(6*x) - cos(4*x) - cos(3*x) + cos(x) + 1)*cos(
18*x) + cos(18*x)^2 - 2*(cos(15*x) + cos(14*x) - cos(12*x) + cos(10*x) + c
os(9*x) - cos(7*x) - cos(6*x) + cos(4*x) + cos(3*x) - cos(x) - 1)*cos(17*x
) + cos(17*x)^2 + 2*(cos(14*x) - cos(12*x) + cos(10*x) + cos(9*x) - cos(7*
x) - cos(6*x) + cos(4*x) + cos(3*x) - cos(x) - 1)*cos(15*x) + cos(15*x)^2
- 2*(cos(12*x) - cos(10*x) - cos(9*x) + cos(7*x) + cos(6*x) - cos(4*x) - c
os(3*x) + cos(x) + 1)*cos(14*x) + cos(14*x)^2 - 2*(cos(10*x) + cos(9*x) -
cos(7*x) - cos(6*x) + cos(4*x) + cos(3*x) - cos(x) - 1)*cos(12*x) + cos(12
*x)^2 + 2*(cos(9*x) - cos(7*x) - cos(6*x) + cos(4*x) + cos(3*x) - cos(x)...
```


3.207.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 134 vs. $2(7) = 14$.

Time = 0.32 (sec) , antiderivative size = 134, normalized size of antiderivative = 12.18

$$\int \frac{\sin(19x) + \sin(20x)}{\cos(19x) + \cos(20x)} dx = -\frac{1}{39} \log(\cos(x) + 1) - \frac{2}{39} \log(|4096 \cos(x)^{12} + 2048 \cos(x)^{11} - 12288 \cos(x)^{10} - 6144 \cos(x)^9 + 13568 \cos(x)^8 + 6784 \cos(x)^7 - 6592 \cos(x)^6 - 3296 \cos(x)^5 + 1264 \cos(x)^4 + 632 \cos(x)^3 - 48 \cos(x)^2 - 24 \cos(x) + 1|) - \frac{2}{39} \log(|64 \cos(x)^6 - 32 \cos(x)^5 - 80 \cos(x)^4 + 32 \cos(x)^3 + 24 \cos(x)^2 - 6 \cos(x) - 1|) - \frac{2}{39} \log(|2 \cos(x) - 1|)$$

input `integrate((sin(20*x)+sin(19*x))/(cos(20*x)+cos(19*x)),x, algorithm="giac")`

output `-1/39*log(cos(x) + 1) - 2/39*log(abs(4096*cos(x)^12 + 2048*cos(x)^11 - 12288*cos(x)^10 - 6144*cos(x)^9 + 13568*cos(x)^8 + 6784*cos(x)^7 - 6592*cos(x)^6 - 3296*cos(x)^5 + 1264*cos(x)^4 + 632*cos(x)^3 - 48*cos(x)^2 - 24*cos(x) + 1)) - 2/39*log(abs(64*cos(x)^6 - 32*cos(x)^5 - 80*cos(x)^4 + 32*cos(x)^3 + 24*cos(x)^2 - 6*cos(x) - 1)) - 2/39*log(abs(2*cos(x) - 1))`

3.207.9 Mupad [B] (verification not implemented)

Time = 16.88 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.36

$$\int \frac{\sin(19x) + \sin(20x)}{\cos(19x) + \cos(20x)} dx = x \operatorname{li} - \frac{2 \ln(e^{x 39i} + 1)}{39}$$

input `int((sin(19*x) + sin(20*x))/(cos(19*x) + cos(20*x)),x)`

output `x*i - (2*log(exp(x*39i) + 1))/39`

3.208 $\int e^x (\cos^2(x) + \cos(x) \sin(x) - \sin^2(x)) dx$

3.208.1 Optimal result	1073
3.208.2 Mathematica [A] (verified)	1073
3.208.3 Rubi [A] (verified)	1074
3.208.4 Maple [A] (verified)	1074
3.208.5 Fricas [A] (verification not implemented)	1075
3.208.6 Sympy [A] (verification not implemented)	1075
3.208.7 Maxima [B] (verification not implemented)	1075
3.208.8 Giac [B] (verification not implemented)	1076
3.208.9 Mupad [B] (verification not implemented)	1076

3.208.1 Optimal result

Integrand size = 20, antiderivative size = 8

$$\int e^x (\cos^2(x) + \cos(x) \sin(x) - \sin^2(x)) dx = e^x \cos(x) \sin(x)$$

output `exp(x)*cos(x)*sin(x)`

3.208.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.38

$$\int e^x (\cos^2(x) + \cos(x) \sin(x) - \sin^2(x)) dx = \frac{1}{2} e^x \sin(2x)$$

input `Integrate[E^x*(Cos[x]^2 + Cos[x]*Sin[x] - Sin[x]^2),x]`

output `(E^x*Sin[2*x])/2`

3.208.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {2726}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x (-\sin^2(x) + \cos^2(x) + \sin(x) \cos(x)) dx$$

\downarrow 2726
 $e^x \sin(x) \cos(x)$

input `Int[E^x*(Cos[x]^2 + Cos[x]*Sin[x] - Sin[x]^2),x]`

output `E^x*Cos[x]*Sin[x]`

3.208.3.1 Defintions of rubi rules used

rule 2726 `Int[(y_.)*(F_)^(u_)*((v_) + (w_)), x_Symbol] := With[{z = v*(y/(Log[F]*D[u, x]))}, Simp[F^u*z, x] /; EqQ[D[z, x], w*y]] /; FreeQ[F, x]`

3.208.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
parallelrisch	$\frac{e^x \sin(2x)}{2}$	9
risch	$-\frac{ie^{(1+2i)x}}{4} + \frac{ie^{(1-2i)x}}{4}$	20
norman	$\frac{2e^x \tan(\frac{x}{2}) - 2e^x \tan(\frac{x}{2})^3}{(1 + \tan(\frac{x}{2})^2)^2}$	31
default	$\frac{(\cos(x) + 2 \sin(x))e^x \cos(x)}{5} + \frac{e^x (\sin(2x) - 2 \cos(2x))}{10} - \frac{(-2 \cos(x) + \sin(x))e^x \sin(x)}{5}$	43
parts	$\frac{(\cos(x) + 2 \sin(x))e^x \cos(x)}{5} + \frac{e^x (\sin(2x) - 2 \cos(2x))}{10} - \frac{(-2 \cos(x) + \sin(x))e^x \sin(x)}{5}$	43

input `int(exp(x)*(cos(x)^2+cos(x)*sin(x)-sin(x)^2),x,method=_RETURNVERBOSE)`

3.208. $\int e^x (\cos^2(x) + \cos(x) \sin(x) - \sin^2(x)) dx$

output `1/2*exp(x)*sin(2*x)`

3.208.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int e^x (\cos^2(x) + \cos(x) \sin(x) - \sin^2(x)) dx = \cos(x) e^x \sin(x)$$

input `integrate(exp(x)*(cos(x)^2+cos(x)*sin(x)-sin(x)^2),x, algorithm="fricas")`

output `cos(x)*e^x*sin(x)`

3.208.6 Sympy [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int e^x (\cos^2(x) + \cos(x) \sin(x) - \sin^2(x)) dx = e^x \sin(x) \cos(x)$$

input `integrate(exp(x)*(cos(x)**2+cos(x)*sin(x)-sin(x)**2),x)`

output `exp(x)*sin(x)*cos(x)`

3.208.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. $2(7) = 14$.

Time = 0.19 (sec) , antiderivative size = 34, normalized size of antiderivative = 4.25

$$\int e^x (\cos^2(x) + \cos(x) \sin(x) - \sin^2(x)) dx = -\frac{1}{10} (2 \cos(2x) - \sin(2x))e^x + \frac{1}{5} \cos(2x) e^x + \frac{2}{5} e^x \sin(2x)$$

input `integrate(exp(x)*(cos(x)^2+cos(x)*sin(x)-sin(x)^2),x, algorithm="maxima")`

output `-1/10*(2*cos(2*x) - sin(2*x))*e^x + 1/5*cos(2*x)*e^x + 2/5*e^x*sin(2*x)`

3.208.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 33 vs. $2(7) = 14$.

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 4.12

$$\int e^x (\cos^2(x) + \cos(x) \sin(x) - \sin^2(x)) dx = -\frac{1}{10} (2 \cos(2x) - \sin(2x))e^x + \frac{1}{5} (\cos(2x) + 2 \sin(2x))e^x$$

input `integrate(exp(x)*(cos(x)^2+cos(x)*sin(x)-sin(x)^2),x, algorithm="giac")`

output `-1/10*(2*cos(2*x) - sin(2*x))*e^x + 1/5*(cos(2*x) + 2*sin(2*x))*e^x`

3.208.9 Mupad [B] (verification not implemented)

Time = 15.33 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int e^x (\cos^2(x) + \cos(x) \sin(x) - \sin^2(x)) dx = \frac{\sin(2x) e^x}{2}$$

input `int(exp(x)*(cos(x)*sin(x) + cos(x)^2 - sin(x)^2),x)`

output `(sin(2*x)*exp(x))/2`

3.209 $\int \csc\left(\frac{\pi}{4} + x\right) \sin(x) dx$

3.209.1 Optimal result	1077
3.209.2 Mathematica [C] (verified)	1077
3.209.3 Rubi [A] (verified)	1078
3.209.4 Maple [A] (verified)	1079
3.209.5 Fricas [A] (verification not implemented)	1080
3.209.6 Sympy [A] (verification not implemented)	1080
3.209.7 Maxima [A] (verification not implemented)	1080
3.209.8 Giac [B] (verification not implemented)	1081
3.209.9 Mupad [B] (verification not implemented)	1081

3.209.1 Optimal result

Integrand size = 11, antiderivative size = 24

$$\int \csc\left(\frac{\pi}{4} + x\right) \sin(x) dx = \frac{x}{\sqrt{2}} - \frac{\log\left(\sin\left(\frac{\pi}{4} + x\right)\right)}{\sqrt{2}}$$

output `1/2*x*2^(1/2)-1/2*ln(sin(x+1/4*Pi))*2^(1/2)`

3.209.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21

$$\int \csc\left(\frac{\pi}{4} + x\right) \sin(x) dx = \left(-\frac{1}{4} - \frac{i}{4}\right) (-1)^{3/4} (2x - 2\operatorname{arctanh}(\cot(x)) - \log(\cos(2x)))$$

input `Integrate[Csc[Pi/4 + x]*Sin[x],x]`

output `(-1/4 - I/4)*(-1)^(3/4)*(2*x - 2*ArcTanh[Cot[x]] - Log[Cos[2*x]])`

3.209.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {5093, 24, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(x) \csc\left(x + \frac{\pi}{4}\right) dx \\
 & \quad \downarrow \text{5093} \\
 & \frac{\int 1 dx}{\sqrt{2}} - \frac{\int \cot\left(x + \frac{\pi}{4}\right) dx}{\sqrt{2}} \\
 & \quad \downarrow \text{24} \\
 & \frac{x}{\sqrt{2}} - \frac{\int \cot\left(x + \frac{\pi}{4}\right) dx}{\sqrt{2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{x}{\sqrt{2}} - \frac{\int -\tan\left(x + \frac{3\pi}{4}\right) dx}{\sqrt{2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \tan\left(x + \frac{3\pi}{4}\right) dx}{\sqrt{2}} + \frac{x}{\sqrt{2}} \\
 & \quad \downarrow \text{3956} \\
 & \frac{x}{\sqrt{2}} - \frac{\log\left(\sin\left(x + \frac{\pi}{4}\right)\right)}{\sqrt{2}}
 \end{aligned}$$

input `Int[Csc[Pi/4 + x]*Sin[x],x]`

output `x/Sqrt[2] - Log[Sin[Pi/4 + x]]/Sqrt[2]`

3.209.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 5093 `Int[Csc[w_]^(n_.)*Sin[v_], x_Symbol] := Simp[Sin[v - w] Int[Cot[w]*Csc[w]^(n - 1), x], x] + Simp[Cos[v - w] Int[Csc[w]^(n - 1), x], x] /; GtQ[n, 0] && FreeQ[v - w, x] && NeQ[w, v]`

3.209.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.12

method	result	size
default	$\sqrt{2} \left(-\frac{\ln(1+\tan(x))}{2} + \frac{\ln(1+\tan(x)^2)}{4} + \frac{\arctan(\tan(x))}{2} \right)$	27
risch	$\frac{x\sqrt{2}}{2} + \frac{i\sqrt{2}x}{2} - \frac{\sqrt{2} \ln(e^{2ix}+i)}{2}$	29

input `int(sin(x)/sin(x+1/4*Pi),x,method=_RETURNVERBOSE)`

output `2^(1/2)*(-1/2*ln(1+tan(x))+1/4*ln(1+tan(x)^2)+1/2*arctan(tan(x)))`

3.209.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

$$\int \csc\left(\frac{\pi}{4} + x\right) \sin(x) dx = \frac{1}{2} \sqrt{2}x - \frac{1}{2} \sqrt{2} \log\left(\frac{1}{2} \sin\left(\frac{1}{4} \pi + x\right)\right)$$

input `integrate(sin(x)/sin(x+1/4*pi),x, algorithm="fracas")`output `1/2*sqrt(2)*x - 1/2*sqrt(2)*log(1/2*sin(1/4*pi + x))`**3.209.6 Sympy [A] (verification not implemented)**

Time = 0.41 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \csc\left(\frac{\pi}{4} + x\right) \sin(x) dx = \frac{\sqrt{2}x}{2} - \frac{\sqrt{2} \log(\sin(x) + \cos(x))}{2}$$

input `integrate(sin(x)/sin(x+1/4*pi),x)`output `sqrt(2)*x/2 - sqrt(2)*log(sin(x) + cos(x))/2`**3.209.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.38

$$\int \csc\left(\frac{\pi}{4} + x\right) \sin(x) dx = \frac{1}{2} \sqrt{2}x - \frac{1}{4} \sqrt{2} \log(\cos(2x)^2 + \sin(2x)^2 + 2 \sin(2x) + 1)$$

input `integrate(sin(x)/sin(x+1/4*pi),x, algorithm="maxima")`output `1/2*sqrt(2)*x - 1/4*sqrt(2)*log(cos(2*x)^2 + sin(2*x)^2 + 2*sin(2*x) + 1)`

3.209.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 44 vs. $2(19) = 38$.

Time = 0.29 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.83

$$\int \csc\left(\frac{\pi}{4} + x\right) \sin(x) dx = \frac{1}{8}\sqrt{2}(\pi + 4x) + \frac{1}{2}\sqrt{2}\log\left(\tan\left(\frac{1}{8}\pi + \frac{1}{2}x\right)^2 + 1\right) - \frac{1}{2}\sqrt{2}\log\left(\left|\tan\left(\frac{1}{8}\pi + \frac{1}{2}x\right)\right|\right)$$

input `integrate(sin(x)/sin(x+1/4*pi),x, algorithm="giac")`

output `1/8*sqrt(2)*(pi + 4*x) + 1/2*sqrt(2)*log(tan(1/8*pi + 1/2*x)^2 + 1) - 1/2*sqrt(2)*log(abs(tan(1/8*pi + 1/2*x)))`

3.209.9 Mupad [B] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.92

$$\int \csc\left(\frac{\pi}{4} + x\right) \sin(x) dx = x e^{\frac{\pi 1i}{4}} - \frac{e^{-\frac{\pi 1i}{2}} \ln\left(e^{\frac{\pi 1i}{2}} e^{x 2i} - 1\right) \left(e^{\frac{\pi 1i}{4}} 2i - e^{\frac{\pi 3i}{4}} 2i\right)}{4}$$

input `int(sin(x)/sin(Pi/4 + x),x)`

output `x*exp((Pi*1i)/4) - (exp(-(Pi*1i)/2)*log(exp((Pi*1i)/2)*exp(x*2i) - 1)*(exp((Pi*1i)/4)*2i - exp((Pi*3i)/4)*2i))/4`

$$3.210 \quad \int \frac{1}{\sqrt[3]{x+x}} dx$$

3.210.1 Optimal result	1082
3.210.2 Mathematica [A] (verified)	1082
3.210.3 Rubi [A] (verified)	1083
3.210.4 Maple [A] (verified)	1084
3.210.5 Fricas [A] (verification not implemented)	1084
3.210.6 Sympy [A] (verification not implemented)	1084
3.210.7 Maxima [A] (verification not implemented)	1085
3.210.8 Giac [A] (verification not implemented)	1085
3.210.9 Mupad [B] (verification not implemented)	1085

3.210.1 Optimal result

Integrand size = 9, antiderivative size = 12

$$\int \frac{1}{\sqrt[3]{x+x}} dx = \frac{3}{2} \log(1+x^{2/3})$$

output `3/2*ln(1+x^(2/3))`

3.210.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt[3]{x+x}} dx = \frac{3}{2} \log(1+x^{2/3})$$

input `Integrate[(x^(1/3) + x)^(-1),x]`

output `(3*Log[1 + x^(2/3)])/2`

3.210.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {2027, 792}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x + \sqrt[3]{x}} dx$$

↓ 2027

$$\int \frac{1}{(x^{2/3} + 1)\sqrt[3]{x}} dx$$

↓ 792

$$\frac{3}{2} \log(x^{2/3} + 1)$$

input `Int[(x^(1/3) + x)^(-1),x]`

output `(3*Log[1 + x^(2/3)])/2`

3.210.3.1 Defintions of rubi rules used

rule 792 `Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]`

rule 2027 `Int[(Fx_)*((a_)*(x_)^(r_) + (b_)*(x_)^(s_))^(p_), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r))^p*Fx, x] /; FreeQ[{a, b, r, s}, x] && IntegerQ[p] && PosQ[s - r] && !(EqQ[p, 1] && EqQ[u, 1])`

3.210.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

method	result	size
derivativdivides	$\frac{3 \ln(1+x^{\frac{2}{3}})}{2}$	9
meijerg	$\frac{3 \ln(1+x^{\frac{2}{3}})}{2}$	9
trager	$\frac{\ln(3x^{\frac{2}{3}}+3x^{\frac{4}{3}}+x^2+1)}{2}$	19
default	$\ln\left(1+x^{\frac{2}{3}}\right) - \frac{\ln\left(x^{\frac{4}{3}}-x^{\frac{2}{3}}+1\right)}{2} + \frac{\ln(x^2+1)}{2}$	29

input `int(1/(x+x^(1/3)),x,method=_RETURNVERBOSE)`output `3/2*ln(1+x^(2/3))`**3.210.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt[3]{x}+x} dx = \frac{3}{2} \log\left(x^{\frac{2}{3}}+1\right)$$

input `integrate(1/(x+x^(1/3)),x, algorithm="fricas")`output `3/2*log(x^(2/3) + 1)`**3.210.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1}{\sqrt[3]{x}+x} dx = \frac{3 \log\left(x^{\frac{2}{3}}+1\right)}{2}$$

input `integrate(1/(x+x**(1/3)),x)`output `3*log(x**(2/3) + 1)/2`

3.210.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt[3]{x} + x} dx = \frac{3}{2} \log(x^{2/3} + 1)$$

input `integrate(1/(x+x^(1/3)),x, algorithm="maxima")`output `3/2*log(x^(2/3) + 1)`**3.210.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt[3]{x} + x} dx = \frac{3}{2} \log(x^{2/3} + 1)$$

input `integrate(1/(x+x^(1/3)),x, algorithm="giac")`output `3/2*log(x^(2/3) + 1)`**3.210.9 Mupad [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt[3]{x} + x} dx = \frac{3 \ln(x^{2/3} + 1)}{2}$$

input `int(1/(x + x^(1/3)),x)`output `(3*log(x^(2/3) + 1))/2`

3.211 $\int x^{1+x^2}(1 + 2 \log(x)) dx$

3.211.1 Optimal result	1086
3.211.2 Mathematica [A] (verified)	1086
3.211.3 Rubi [F]	1087
3.211.4 Maple [B] (verified)	1087
3.211.5 Fricas [B] (verification not implemented)	1088
3.211.6 Sympy [B] (verification not implemented)	1088
3.211.7 Maxima [A] (verification not implemented)	1089
3.211.8 Giac [F]	1089
3.211.9 Mupad [B] (verification not implemented)	1089

3.211.1 Optimal result

Integrand size = 14, antiderivative size = 5

$$\int x^{1+x^2}(1 + 2 \log(x)) dx = x^{x^2}$$

output `x^(x^2)`

3.211.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int x^{1+x^2}(1 + 2 \log(x)) dx = x^{x^2}$$

input `Integrate[x^(1 + x^2)*(1 + 2*Log[x]),x]`

output `x^x^2`

3.211.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{x^2+1}(2\log(x) + 1) dx$$

$$\downarrow \text{7293}$$

$$\int (x^{x^2+1} + 2x^{x^2+1}\log(x)) dx$$

$$\downarrow \text{2009}$$

$$\int x^{x^2+1} dx - 2 \int \frac{x^{x^2+1} dx}{x} + 2\log(x) \int x^{x^2+1} dx$$

input `Int[x^(1 + x^2)*(1 + 2*Log[x]),x]`

output `$Aborted`

3.211.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

3.211.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. $2(5) = 10$.

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 2.40

method	result	size
risch	$\frac{x^{x^2+1}}{x}$	12
parallelrisch	$\frac{x^{x^2+1}}{x}$	12

input `int(x^(x^2+1)*(1+2*ln(x)),x,method=_RETURNVERBOSE)`

output `x^(x^2+1)/x`

3.211.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. $2(5) = 10$.

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 2.20

$$\int x^{1+x^2} (1 + 2 \log(x)) dx = \frac{x^{x^2+1}}{x}$$

input `integrate(x^(x^2+1)*(1+2*log(x)),x, algorithm="fricas")`

output `x^(x^2 + 1)/x`

3.211.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7 vs. $2(3) = 6$.

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.40

$$\int x^{1+x^2} (1 + 2 \log(x)) dx = \frac{x^{x^2+1}}{x}$$

input `integrate(x**(x**2+1)*(1+2*ln(x)),x)`

output `x**(x**2 + 1)/x`

3.211.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int x^{1+x^2}(1+2\log(x)) dx = x^{(x^2)}$$

input `integrate(x^(x^2+1)*(1+2*log(x)),x, algorithm="maxima")`output `x^(x^2)`**3.211.8 Giac [F]**

$$\int x^{1+x^2}(1+2\log(x)) dx = \int x^{x^2+1}(2\log(x)+1) dx$$

input `integrate(x^(x^2+1)*(1+2*log(x)),x, algorithm="giac")`output `integrate(x^(x^2 + 1)*(2*log(x) + 1), x)`**3.211.9 Mupad [B] (verification not implemented)**

Time = 15.46 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int x^{1+x^2}(1+2\log(x)) dx = x^{x^2}$$

input `int(x^(x^2 + 1)*(2*log(x) + 1),x)`output `x^(x^2)`

$$3.212 \quad \int \frac{-1+2x^3}{x(1+x^3)} dx$$

3.212.1 Optimal result	1090
3.212.2 Mathematica [A] (verified)	1090
3.212.3 Rubi [A] (verified)	1091
3.212.4 Maple [A] (verified)	1092
3.212.5 Fricas [A] (verification not implemented)	1093
3.212.6 Sympy [A] (verification not implemented)	1093
3.212.7 Maxima [A] (verification not implemented)	1093
3.212.8 Giac [A] (verification not implemented)	1094
3.212.9 Mupad [B] (verification not implemented)	1094

3.212.1 Optimal result

Integrand size = 18, antiderivative size = 11

$$\int \frac{-1+2x^3}{x(1+x^3)} dx = -\log(x) + \log(1+x^3)$$

output `-ln(x)+ln(x^3+1)`

3.212.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{-1+2x^3}{x(1+x^3)} dx = -\log(x) + \log(1+x^3)$$

input `Integrate[(-1 + 2*x^3)/(x*(1 + x^3)),x]`

output `-Log[x] + Log[1 + x^3]`

3.212.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.73, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {948, 25, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{2x^3 - 1}{x(x^3 + 1)} dx \\ & \quad \downarrow \text{948} \\ & \frac{1}{3} \int -\frac{1 - 2x^3}{x^3(x^3 + 1)} dx^3 \\ & \quad \downarrow \text{25} \\ & -\frac{1}{3} \int \frac{1 - 2x^3}{x^3(x^3 + 1)} dx^3 \\ & \quad \downarrow \text{86} \\ & -\frac{1}{3} \int \left(\frac{1}{x^3} - \frac{3}{x^3 + 1} \right) dx^3 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{3} (3 \log(x^3 + 1) - \log(x^3)) \end{aligned}$$

input `Int[(-1 + 2*x^3)/(x*(1 + x^3)),x]`

output `(-Log[x^3] + 3*Log[1 + x^3])/3`

3.212.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 948 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^(p_.)*((c_) + (d_.)*(x_)^(n_.))^(q_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.212.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
meijerg	$-\ln(x) + \ln(x^3 + 1)$	12
risch	$-\ln(x) + \ln(x^3 + 1)$	12
default	$\ln(1 + x) + \ln(x^2 - x + 1) - \ln(x)$	19
norman	$\ln(1 + x) + \ln(x^2 - x + 1) - \ln(x)$	19
parallelrisc	$\ln(1 + x) + \ln(x^2 - x + 1) - \ln(x)$	19

input `int((2*x^3-1)/x/(x^3+1),x,method=_RETURNVERBOSE)`

output `-ln(x)+ln(x^3+1)`

3.212.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{-1 + 2x^3}{x(1 + x^3)} dx = \log(x^3 + 1) - \log(x)$$

input `integrate((2*x^3-1)/x/(x^3+1),x, algorithm="fricas")`output `log(x^3 + 1) - log(x)`**3.212.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.73

$$\int \frac{-1 + 2x^3}{x(1 + x^3)} dx = -\log(x) + \log(x^3 + 1)$$

input `integrate((2*x**3-1)/x/(x**3+1),x)`output `-log(x) + log(x**3 + 1)`**3.212.7 Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.18

$$\int \frac{-1 + 2x^3}{x(1 + x^3)} dx = \log(x^3 + 1) - \frac{1}{3} \log(x^3)$$

input `integrate((2*x^3-1)/x/(x^3+1),x, algorithm="maxima")`output `log(x^3 + 1) - 1/3*log(x^3)`

3.212.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.18

$$\int \frac{-1 + 2x^3}{x(1 + x^3)} dx = \log(|x^3 + 1|) - \log(|x|)$$

input `integrate((2*x^3-1)/x/(x^3+1),x, algorithm="giac")`output `log(abs(x^3 + 1)) - log(abs(x))`**3.212.9 Mupad [B] (verification not implemented)**

Time = 15.51 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{-1 + 2x^3}{x(1 + x^3)} dx = \ln(x^3 + 1) - \ln(x)$$

input `int((2*x^3 - 1)/(x*(x^3 + 1)),x)`output `log(x^3 + 1) - log(x)`

3.213 $\int \frac{1}{\sqrt{1+x^2}} dx$

3.213.1 Optimal result	1095
3.213.2 Mathematica [B] (verified)	1095
3.213.3 Rubi [A] (verified)	1096
3.213.4 Maple [A] (verified)	1096
3.213.5 Fricas [B] (verification not implemented)	1097
3.213.6 Sympy [A] (verification not implemented)	1097
3.213.7 Maxima [A] (verification not implemented)	1097
3.213.8 Giac [B] (verification not implemented)	1098
3.213.9 Mupad [B] (verification not implemented)	1098

3.213.1 Optimal result

Integrand size = 9, antiderivative size = 2

$$\int \frac{1}{\sqrt{1+x^2}} dx = \operatorname{arcsinh}(x)$$

output `arcsinh(x)`

3.213.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 16 vs. $2(2) = 4$.

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 8.00

$$\int \frac{1}{\sqrt{1+x^2}} dx = -\log(-x + \sqrt{1+x^2})$$

input `Integrate[1/Sqrt[1 + x^2],x]`

output `-Log[-x + Sqrt[1 + x^2]]`

3.213.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x^2 + 1}} dx$$

↓ 222

$$\operatorname{arcsinh}(x)$$

input `Int[1/Sqrt[1 + x^2], x]`

output `ArcSinh[x]`

3.213.3.1 Defintions of rubi rules used

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

3.213.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\operatorname{arcsinh}(x)$	3
meijerg	$\operatorname{arcsinh}(x)$	3
trager	$\ln(x + \sqrt{x^2 + 1})$	11
pseudoelliptic	$\operatorname{arctanh}\left(\frac{\sqrt{x^2+1}}{x}\right)$	13

input `int(1/(x^2+1)^(1/2), x, method=_RETURNVERBOSE)`

output `arcsinh(x)`

3.213.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 14 vs. $2(2) = 4$.

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 7.00

$$\int \frac{1}{\sqrt{1+x^2}} dx = -\log\left(-x + \sqrt{x^2+1}\right)$$

input `integrate(1/(x^2+1)^(1/2),x, algorithm="fricas")`

output `-log(-x + sqrt(x^2 + 1))`

3.213.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1+x^2}} dx = \operatorname{asinh}(x)$$

input `integrate(1/(x**2+1)**(1/2),x)`

output `asinh(x)`

3.213.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1+x^2}} dx = \operatorname{arsinh}(x)$$

input `integrate(1/(x^2+1)^(1/2),x, algorithm="maxima")`

output `arcsinh(x)`

3.213.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. $2(2) = 4$.

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 12.50

$$\int \frac{1}{\sqrt{1+x^2}} dx = \frac{1}{2} \sqrt{x^2+1}x - \frac{1}{2} \log(-x + \sqrt{x^2+1})$$

input `integrate(1/(x^2+1)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(x^2 + 1)*x - 1/2*log(-x + sqrt(x^2 + 1))`

3.213.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1+x^2}} dx = \operatorname{asinh}(x)$$

input `int(1/(x^2 + 1)^(1/2),x)`

output `asinh(x)`

3.214 $\int \frac{\log(2x)}{x \log(x)} dx$

3.214.1 Optimal result	1099
3.214.2 Mathematica [A] (verified)	1099
3.214.3 Rubi [A] (verified)	1100
3.214.4 Maple [A] (verified)1101
3.214.5 Fricas [A] (verification not implemented)1101
3.214.6 Sympy [A] (verification not implemented)1101
3.214.7 Maxima [A] (verification not implemented)	1102
3.214.8 Giac [A] (verification not implemented)	1102
3.214.9 Mupad [B] (verification not implemented)	1102

3.214.1 Optimal result

Integrand size = 12, antiderivative size = 9

$$\int \frac{\log(2x)}{x \log(x)} dx = \log(x) + \log(2) \log(\log(x))$$

output `ln(x)+ln(2)*ln(ln(x))`

3.214.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{\log(2x)}{x \log(x)} dx = \log(x) + \log(2) \log(\log(x))$$

input `Integrate[Log[2*x]/(x*Log[x]),x]`

output `Log[x] + Log[2]*Log[Log[x]]`

3.214.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 2.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2813, 3001}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\log(2x)}{x \log(x)} dx$$

↓ 2813

$$\log(2x) \log(\log(x)) - \int \frac{\log(\log(x))}{x} dx$$

↓ 3001

$$-\log(\log(x)) \log(x) + \log(x) + \log(2x) \log(\log(x))$$

input `Int [Log [2*x] / (x*Log [x]), x]`

output `Log [x] - Log [x]*Log [Log [x]] + Log [2*x]*Log [Log [x]]`

3.214.3.1 Defintions of rubi rules used

rule 2813 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((d_.) + Log[(f_.)*(x_)^(r_.)]*(e_.))*((g_.)*(x_)^(m_.), x_Symbol] := With[{u = IntHide[(g*x)^m*(a + b*Log[c*x^n])^p, x]}, Simp[(d + e*Log[f*x^r]) u, x] - Simp[e*r Int[SimplifyIntegrand[u/x, x], x], x]] /; FreeQ[{a, b, c, d, e, f, g, m, n, p, r}, x] && !(EqQ[p, 1] && EqQ[a, 0] && NeQ[d, 0])`

rule 3001 `Int[((a_.) + Log[Log[(d_.)*(x_)^(n_.)]^(p_.)*(c_.)]*(b_.))/(x_), x_Symbol] := Simp[Log[d*x^n]*((a + b*Log[c*Log[d*x^n]^p])/n), x] - Simp[b*p*Log[x], x] /; FreeQ[{a, b, c, d, n, p}, x]`

3.214.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.11

method	result	size
default	$\ln(x) + \ln(2) \ln(\ln(x))$	10
risch	$\ln(x) + \ln(2) \ln(\ln(x))$	10

input `int(ln(2*x)/x/ln(x),x,method=_RETURNVERBOSE)`

output `ln(x)+ln(2)*ln(ln(x))`

3.214.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{\log(2x)}{x \log(x)} dx = \log(2) \log(\log(x)) + \log(x)$$

input `integrate(log(2*x)/x/log(x),x, algorithm="fracas")`

output `log(2)*log(log(x)) + log(x)`

3.214.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.11

$$\int \frac{\log(2x)}{x \log(x)} dx = \log(x) + \log(2) \log(\log(x))$$

input `integrate(ln(2*x)/x/ln(x),x)`

output `log(x) + log(2)*log(log(x))`

3.214.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 18, normalized size of antiderivative = 2.00

$$\int \frac{\log(2x)}{x \log(x)} dx = \log(2x) \log(\log(x)) - \log(x) \log(\log(x)) + \log(x)$$

input `integrate(log(2*x)/x/log(x),x, algorithm="maxima")`output `log(2*x)*log(log(x)) - log(x)*log(log(x)) + log(x)`**3.214.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.11

$$\int \frac{\log(2x)}{x \log(x)} dx = \log(2) \log(|\log(x)|) + \log(x)$$

input `integrate(log(2*x)/x/log(x),x, algorithm="giac")`output `log(2)*log(abs(log(x))) + log(x)`**3.214.9 Mupad [B] (verification not implemented)**

Time = 15.23 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{\log(2x)}{x \log(x)} dx = \ln(x) + \ln(\ln(x)) \ln(2)$$

input `int(log(2*x)/(x*log(x)),x)`output `log(x) + log(log(x))*log(2)`

3.215 $\int \frac{1}{1+e^x} dx$

3.215.1 Optimal result	1103
3.215.2 Mathematica [A] (verified)	1103
3.215.3 Rubi [A] (verified)	1104
3.215.4 Maple [A] (verified)	1105
3.215.5 Fricas [A] (verification not implemented)	1105
3.215.6 Sympy [A] (verification not implemented)	1106
3.215.7 Maxima [A] (verification not implemented)	1106
3.215.8 Giac [A] (verification not implemented)	1106
3.215.9 Mupad [B] (verification not implemented)	1107

3.215.1 Optimal result

Integrand size = 7, antiderivative size = 10

$$\int \frac{1}{1+e^x} dx = x - \log(1+e^x)$$

output `x-ln(exp(x)+1)`

3.215.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+e^x} dx = -2\operatorname{arctanh}(1+2e^x)$$

input `Integrate[(1 + E^x)^(-1),x]`

output `-2*ArcTanh[1 + 2*E^x]`

3.215.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.30, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {2720, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{1}{e^x + 1} dx \\
 \downarrow 2720 \\
 \int \frac{e^{-x}}{e^x + 1} de^x \\
 \downarrow 47 \\
 \int e^{-x} de^x - \int \frac{1}{1 + e^x} de^x \\
 \downarrow 14 \\
 \log(e^x) - \int \frac{1}{1 + e^x} de^x \\
 \downarrow 16 \\
 \log(e^x) - \log(e^x + 1)
 \end{array}$$

input `Int[(1 + E^x)^(-1), x]`

output `Log[E^x] - Log[1 + E^x]`

3.215.3.1 Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

```
rule 47 Int[1/(((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))), x_Symbol] := Simp[b/(b*c
- a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x
], x] /; FreeQ[{a, b, c, d}, x]
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.215.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

method	result	size
norman	$x - \ln(e^x + 1)$	10
risch	$x - \ln(e^x + 1)$	10
parallelrisch	$x - \ln(e^x + 1)$	10
derivativedivides	$-\ln(e^x + 1) + \ln(e^x)$	12
default	$-\ln(e^x + 1) + \ln(e^x)$	12

```
input int(1/(exp(x)+1),x,method=_RETURNVERBOSE)
```

```
output x-ln(exp(x)+1)
```

3.215.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{1}{1+e^x} dx = x - \log(e^x + 1)$$

```
input integrate(1/(exp(x)+1),x, algorithm="fricas")
```

```
output x - log(e^x + 1)
```

3.215.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \frac{1}{1+e^x} dx = x - \log(e^x + 1)$$

input `integrate(1/(exp(x)+1),x)`

output `x - log(exp(x) + 1)`

3.215.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{1}{1+e^x} dx = x - \log(e^x + 1)$$

input `integrate(1/(exp(x)+1),x, algorithm="maxima")`

output `x - log(e^x + 1)`

3.215.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{1}{1+e^x} dx = x - \log(e^x + 1)$$

input `integrate(1/(exp(x)+1),x, algorithm="giac")`

output `x - log(e^x + 1)`

3.215.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{1}{1+e^x} dx = x - \ln(e^x + 1)$$

input `int(1/(exp(x) + 1),x)`

output `x - log(exp(x) + 1)`

3.216 $\int \frac{\log(x) \log(\log(x))}{x} dx$

3.216.1 Optimal result	1108
3.216.2 Mathematica [A] (verified)	1108
3.216.3 Rubi [A] (verified)	1109
3.216.4 Maple [A] (verified)	1110
3.216.5 Fricas [A] (verification not implemented)	1110
3.216.6 Sympy [A] (verification not implemented)	1110
3.216.7 Maxima [A] (verification not implemented)	1111
3.216.8 Giac [A] (verification not implemented)	1111
3.216.9 Mupad [B] (verification not implemented)	1111

3.216.1 Optimal result

Integrand size = 9, antiderivative size = 20

$$\int \frac{\log(x) \log(\log(x))}{x} dx = -\frac{1}{4} \log^2(x) + \frac{1}{2} \log^2(x) \log(\log(x))$$

output `-1/4*ln(x)^2+1/2*ln(x)^2*ln(ln(x))`

3.216.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{\log(x) \log(\log(x))}{x} dx = -\frac{1}{4} \log^2(x) + \frac{1}{2} \log^2(x) \log(\log(x))$$

input `Integrate[(Log[x]*Log[Log[x]])/x,x]`

output `-1/4*Log[x]^2 + (Log[x]^2*Log[Log[x]])/2`

3.216.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3039, 2741}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\log(x) \log(\log(x))}{x} dx$$

↓ 3039

$$\int \log(x) \log(\log(x)) d \log(x)$$

↓ 2741

$$\frac{1}{2} \log^2(x) \log(\log(x)) - \frac{\log^2(x)}{4}$$

input `Int[(Log[x]*Log[Log[x]])/x,x]`

output `-1/4*Log[x]^2 + (Log[x]^2*Log[Log[x]])/2`

3.216.3.1 Defintions of rubi rules used

rule 2741 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((d_.)*(x_))^(m_.), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Simp[b*n*((d*x)^(m + 1)/(d*(m + 1)^2)), x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[m, -1]`

rule 3039 `Int[u_, x_Symbol] :> With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst[[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst] /; NonsumQ[u]`

3.216.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

method	result	size
derivativedivides	$-\frac{\ln(x)^2}{4} + \frac{\ln(x)^2 \ln(\ln(x))}{2}$	17
default	$-\frac{\ln(x)^2}{4} + \frac{\ln(x)^2 \ln(\ln(x))}{2}$	17
norman	$-\frac{\ln(x)^2}{4} + \frac{\ln(x)^2 \ln(\ln(x))}{2}$	17
risch	$-\frac{\ln(x)^2}{4} + \frac{\ln(x)^2 \ln(\ln(x))}{2}$	17

input `int(ln(x)*ln(ln(x))/x,x,method=_RETURNVERBOSE)`output `-1/4*ln(x)^2+1/2*ln(x)^2*ln(ln(x))`**3.216.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int \frac{\log(x) \log(\log(x))}{x} dx = \frac{1}{2} \log(x)^2 \log(\log(x)) - \frac{1}{4} \log(x)^2$$

input `integrate(log(x)*log(log(x))/x,x, algorithm="fracas")`output `1/2*log(x)^2*log(log(x)) - 1/4*log(x)^2`**3.216.6 Sympy [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{\log(x) \log(\log(x))}{x} dx = \frac{\log(x)^2 \log(\log(x))}{2} - \frac{\log(x)^2}{4}$$

input `integrate(ln(x)*ln(ln(x))/x,x)`output `log(x)**2*log(log(x))/2 - log(x)**2/4`

3.216.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int \frac{\log(x) \log(\log(x))}{x} dx = \frac{1}{2} \log(x)^2 \log(\log(x)) - \frac{1}{4} \log(x)^2$$

input `integrate(log(x)*log(log(x))/x,x, algorithm="maxima")`output `1/2*log(x)^2*log(log(x)) - 1/4*log(x)^2`**3.216.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int \frac{\log(x) \log(\log(x))}{x} dx = \frac{1}{2} \log(x)^2 \log(\log(x)) - \frac{1}{4} \log(x)^2$$

input `integrate(log(x)*log(log(x))/x,x, algorithm="giac")`output `1/2*log(x)^2*log(log(x)) - 1/4*log(x)^2`**3.216.9 Mupad [B] (verification not implemented)**

Time = 14.93 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.65

$$\int \frac{\log(x) \log(\log(x))}{x} dx = \frac{\ln(x)^2 (2 \ln(\ln(x)) - 1)}{4}$$

input `int((log(log(x))*log(x))/x,x)`output `(log(x)^2*(2*log(log(x)) - 1))/4`

3.217 $\int \log\left(\frac{1+x}{1-x}\right) dx$

3.217.1 Optimal result	1112
3.217.2 Mathematica [A] (verified)	1112
3.217.3 Rubi [A] (verified)	1113
3.217.4 Maple [A] (verified)	1114
3.217.5 Fracas [A] (verification not implemented)	1114
3.217.6 Sympy [A] (verification not implemented)	1114
3.217.7 Maxima [A] (verification not implemented)	1115
3.217.8 Giac [B] (verification not implemented)	1115
3.217.9 Mupad [B] (verification not implemented)	1115

3.217.1 Optimal result

Integrand size = 12, antiderivative size = 25

$$\int \log\left(\frac{1+x}{1-x}\right) dx = 2\log(1-x) + (1+x)\log\left(\frac{1+x}{1-x}\right)$$

output `2*ln(1-x)+(1+x)*ln((1+x)/(1-x))`

3.217.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \log\left(\frac{1+x}{1-x}\right) dx = 2\log(1-x) + (1+x)\log\left(\frac{1+x}{1-x}\right)$$

input `Integrate[Log[(1 + x)/(1 - x)], x]`

output `2*Log[1 - x] + (1 + x)*Log[(1 + x)/(1 - x)]`

3.217.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2935, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \log\left(\frac{x+1}{1-x}\right) dx$$

↓ 2935

$$(x+1) \log\left(\frac{x+1}{1-x}\right) - 2 \int \frac{1}{1-x} dx$$

↓ 16

$$2 \log(1-x) + (x+1) \log\left(\frac{x+1}{1-x}\right)$$

input `Int[Log[(1 + x)/(1 - x)],x]`

output `2*Log[1 - x] + (1 + x)*Log[(1 + x)/(1 - x)]`

3.217.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 2935 `Int[((A_.) + Log[(e_.)*((a_.) + (b_.)*(x_))/((c_.) + (d_.)*(x_))]^(n_.)]*(B_.)^(p_.), x_Symbol] :> Simp[(a + b*x)*((A + B*Log[e*((a + b*x)/(c + d*x))]^n)]^p/b, x] - Simp[B*n*p*((b*c - a*d)/b) Int[(A + B*Log[e*((a + b*x)/(c + d*x))]^n)]^(p - 1)/(c + d*x), x], x] /; FreeQ[{a, b, c, d, e, A, B, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0]`

3.217.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

method	result	size
risch	$x \ln\left(\frac{1+x}{1-x}\right) + \ln(x^2 - 1)$	22
parts	$x \ln\left(\frac{1+x}{1-x}\right) + \ln((-1+x)(1+x))$	24
meijerg	$\frac{(2+2x)\ln(1+x)}{2} + \frac{(-2x+2)\ln(1-x)}{2}$	26
parallelrisc	$\ln\left(-\frac{1+x}{-1+x}\right) x + 2 \ln(-1+x) + \ln\left(-\frac{1+x}{-1+x}\right)$	32
derivativedivides	$-2 \ln\left(-\frac{2}{-1+x}\right) - \ln\left(-1 - \frac{2}{-1+x}\right) \left(-1 - \frac{2}{-1+x}\right) (-1+x)$	36
default	$-2 \ln\left(-\frac{2}{-1+x}\right) - \ln\left(-1 - \frac{2}{-1+x}\right) \left(-1 - \frac{2}{-1+x}\right) (-1+x)$	36

input `int(ln((1+x)/(1-x)),x,method=_RETURNVERBOSE)`output `x*ln((1+x)/(1-x))+ln(x^2-1)`**3.217.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \log\left(\frac{1+x}{1-x}\right) dx = x \log\left(-\frac{x+1}{x-1}\right) + \log(x^2 - 1)$$

input `integrate(log((1+x)/(1-x)),x, algorithm="fricas")`output `x*log(-(x + 1)/(x - 1)) + log(x^2 - 1)`**3.217.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.60

$$\int \log\left(\frac{1+x}{1-x}\right) dx = x \log\left(\frac{x+1}{1-x}\right) + \log(x^2 - 1)$$

input `integrate(ln((1+x)/(1-x)),x)`output `x*log((x + 1)/(1 - x)) + log(x**2 - 1)`

3.217.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \log\left(\frac{1+x}{1-x}\right) dx = x \log\left(-\frac{x+1}{x-1}\right) + \log(x+1) + \log(x-1)$$

input `integrate(log((1+x)/(1-x)),x, algorithm="maxima")`output `x*log(-(x + 1)/(x - 1)) + log(x + 1) + log(x - 1)`**3.217.8 Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 107 vs. 2(24) = 48.

Time = 0.28 (sec) , antiderivative size = 107, normalized size of antiderivative = 4.28

$$\int \log\left(\frac{1+x}{1-x}\right) dx = \frac{2 \log\left(-\frac{\frac{\frac{x+1}{x-1}+1}{\frac{x+1}{x-1}-1}+1}{\frac{\frac{x+1}{x-1}+1}{\frac{x+1}{x-1}-1}-1}\right)}{\frac{x+1}{x-1}-1} + 2 \log\left(\frac{|-x-1|}{|x-1|}\right) - 2 \log\left(\left|-\frac{x+1}{x-1}+1\right|\right)$$

input `integrate(log((1+x)/(1-x)),x, algorithm="giac")`output `2*log(-(((x + 1)/(x - 1) + 1)/((x + 1)/(x - 1) - 1) + 1)/(((x + 1)/(x - 1) + 1)/((x + 1)/(x - 1) - 1) - 1))/((x + 1)/(x - 1) - 1) + 2*log(abs(-x - 1)/abs(x - 1)) - 2*log(abs(-(x + 1)/(x - 1) + 1))`**3.217.9 Mupad [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \log\left(\frac{1+x}{1-x}\right) dx = \ln(x^2 - 1) + x \ln\left(-\frac{x+1}{x-1}\right)$$

input `int(log(-(x + 1)/(x - 1)),x)`output `log(x^2 - 1) + x*log(-(x + 1)/(x - 1))`

$$3.218 \quad \int \frac{1}{(-1+x)^2+x^2} dx$$

3.218.1 Optimal result	1116
3.218.2 Mathematica [A] (verified)	1116
3.218.3 Rubi [A] (verified)	1117
3.218.4 Maple [A] (verified)	1118
3.218.5 Fricas [A] (verification not implemented)	1118
3.218.6 Sympy [A] (verification not implemented)	1118
3.218.7 Maxima [A] (verification not implemented)	1119
3.218.8 Giac [A] (verification not implemented)	1119
3.218.9 Mupad [B] (verification not implemented)	1119

3.218.1 Optimal result

Integrand size = 11, antiderivative size = 8

$$\int \frac{1}{(-1+x)^2+x^2} dx = -\arctan(1-2x)$$

output `arctan(-1+2*x)`

3.218.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{1}{(-1+x)^2+x^2} dx = \arctan(1+2(-1+x))$$

input `Integrate[((-1 + x)^2 + x^2)^(-1), x]`

output `ArcTan[1 + 2*(-1 + x)]`

3.218.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {2080, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x^2 + (x-1)^2} dx \\ & \quad \downarrow \text{2080} \\ & \int \frac{1}{2x^2 - 2x + 1} dx \\ & \quad \downarrow \text{1082} \\ & \int \frac{1}{-(1-2x)^2 - 1} d(1-2x) \\ & \quad \downarrow \text{217} \\ & -\arctan(1-2x) \end{aligned}$$

input `Int[((-1 + x)^2 + x^2)^(-1),x]`

output `-ArcTan[1 - 2*x]`

3.218.3.1 Defintions of rubi rules used

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 2080 `Int[(u_)^(p_), x_Symbol] := Int[ExpandToSum[u, x]^p, x] /; FreeQ[p, x] && Q
uadraticQ[u, x] && !QuadraticMatchQ[u, x]`

3.218.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
default	$\arctan(-1 + 2x)$	7
risch	$\arctan(-1 + 2x)$	7
parallelrisch	$-\frac{i \ln(x - \frac{1}{2} - \frac{i}{2})}{2} + \frac{i \ln(x - \frac{1}{2} + \frac{i}{2})}{2}$	20

input `int(1/(x^2+(-1+x)^2),x,method=_RETURNVERBOSE)`

output `arctan(-1+2*x)`

3.218.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{(-1+x)^2+x^2} dx = \arctan(2x-1)$$

input `integrate(1/(x^2+(-1+x)^2),x, algorithm="fricas")`

output `arctan(2*x - 1)`

3.218.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \frac{1}{(-1+x)^2+x^2} dx = \operatorname{atan}(2x-1)$$

input `integrate(1/(x**2+(-1+x)**2),x)`

output `atan(2*x - 1)`

3.218.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{(-1+x)^2+x^2} dx = \arctan(2x-1)$$

input `integrate(1/(x^2+(-1+x)^2),x, algorithm="maxima")`output `arctan(2*x - 1)`**3.218.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{(-1+x)^2+x^2} dx = \arctan(2x-1)$$

input `integrate(1/(x^2+(-1+x)^2),x, algorithm="giac")`output `arctan(2*x - 1)`**3.218.9 Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{(-1+x)^2+x^2} dx = \operatorname{atan}(2x-1)$$

input `int(1/((x-1)^2+x^2),x)`output `atan(2*x - 1)`

3.219 $\int \sqrt{x\sqrt{x^{3/2}}} dx$

3.219.1 Optimal result	1120
3.219.2 Mathematica [A] (verified)	1120
3.219.3 Rubi [A] (warning: unable to verify)	1121
3.219.4 Maple [A] (verified)	1122
3.219.5 Fricas [A] (verification not implemented)	1123
3.219.6 Sympy [A] (verification not implemented)	1123
3.219.7 Maxima [A] (verification not implemented)	1123
3.219.8 Giac [A] (verification not implemented)	1124
3.219.9 Mupad [B] (verification not implemented)	1124

3.219.1 Optimal result

Integrand size = 15, antiderivative size = 20

$$\int \sqrt{x\sqrt{x^{3/2}}} dx = \frac{8}{15}x\sqrt{x\sqrt{x^{3/2}}}$$

output `8/15*x*(x*(x^(3/2))^(1/2))^(1/2)`

3.219.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \sqrt{x\sqrt{x^{3/2}}} dx = \frac{8}{15}x\sqrt{x\sqrt{x^{3/2}}}$$

input `Integrate[Sqrt[x*Sqrt[x^(3/2)]],x]`

output `(8*x*Sqrt[x*Sqrt[x^(3/2)]])/15`

3.219.3 Rubi [A] (warning: unable to verify)

Time = 0.26 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.65, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {7267, 7270, 21, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \sqrt{x\sqrt{x^{3/2}}} dx \\
 \downarrow \text{7267} \\
 2 \int \sqrt{x} \sqrt{x\sqrt{x^{3/2}}} d\sqrt{x} \\
 \downarrow \text{7270} \\
 \frac{2\sqrt{x\sqrt{x^{3/2}}} \int x^4 \sqrt{x^{3/2}} d\sqrt{x}}{\sqrt{x} \sqrt[4]{x^{3/2}}} \\
 \downarrow \text{21} \\
 \frac{2\sqrt{x\sqrt{x^{3/2}}} \int \sqrt[8]{x} dx^{3/2}}{3\sqrt{x} \sqrt[4]{x^{3/2}}} \\
 \downarrow \text{15} \\
 \frac{8\sqrt[8]{x} \sqrt{x\sqrt{x^{3/2}}}}{15 \sqrt[4]{x^{3/2}}}
 \end{array}$$

input `Int[Sqrt[x*Sqrt[x^(3/2)]],x]`

output `(8*x^(1/8)*Sqrt[x*Sqrt[x^(3/2)]])/(15*(x^(3/2))^(1/4))`

3.219.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 21 `Int[(x_)^(m_.)*((a_.)*(x_)^(n_.))^p_, x_Symbol] := Simp[1/(n*a^(Simplify[(m + 1)/n] - 1)) Subst[Int[(a*x)^(Simplify[(m + 1)/n] + p - 1), x], x, x^n], x] /; FreeQ[{a, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`
- rule 7267 `Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Simp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2]])], x] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]`
- rule 7270 `Int[(u_.)*((a_.)*(v_)^(m_.)*(w_)^(n_.))^p_, x_Symbol] := Simp[a^IntPart[p]*((a*v^m*w^n)^FracPart[p]/(v^(m*FracPart[p])*w^(n*FracPart[p]))) Int[u*v^(m*p)*w^(n*p), x], x] /; FreeQ[{a, m, n, p}, x] && !IntegerQ[p] && !FreeQ[v, x] && !FreeQ[w, x]`

3.219.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.65

method	result	size
gospers	$\frac{8x\sqrt{x}\sqrt{x^{\frac{3}{2}}}}{15}$	13
derivativedivides	$\frac{8x\sqrt{x}\sqrt{x^{\frac{3}{2}}}}{15}$	13
default	$\frac{8x\sqrt{x}\sqrt{x^{\frac{3}{2}}}}{15}$	13

input `int((x*(x^(3/2))^(1/2))^(1/2),x,method=_RETURNVERBOSE)`

output `8/15*x*(x*(x^(3/2))^(1/2))^(1/2)`

3.219.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.60

$$\int \sqrt{x\sqrt{x^{3/2}}} dx = \frac{8}{15} \sqrt{\sqrt{x^{\frac{3}{2}}}x}$$

input `integrate((x*(x^(3/2))^(1/2))^(1/2),x, algorithm="fricas")`output `8/15*sqrt(sqrt(x^(3/2))*x)*x`**3.219.6 Sympy [A] (verification not implemented)**

Time = 0.51 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \sqrt{x\sqrt{x^{3/2}}} dx = \frac{8x\sqrt{x\sqrt{x^{\frac{3}{2}}}}}{15}$$

input `integrate((x*(x**(3/2))**(1/2))**(1/2),x)`output `8*x*sqrt(x*sqrt(x**(3/2)))/15`**3.219.7 Maxima [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.25

$$\int \sqrt{x\sqrt{x^{3/2}}} dx = \frac{8}{15} x^{\frac{15}{8}}$$

input `integrate((x*(x^(3/2))^(1/2))^(1/2),x, algorithm="maxima")`output `8/15*x^(15/8)`

3.219.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \sqrt{x\sqrt{x^{3/2}}} dx = \frac{8}{15} x^{15/8} \operatorname{sgn}\left(x^{7/4} + 4x^{3/4}\right) \operatorname{sgn}(x + 4)$$

input `integrate((x*(x^(3/2))^(1/2))^(1/2),x, algorithm="giac")`output `8/15*x^(15/8)*sgn(x^(7/4) + 4*x^(3/4))*sgn(x + 4)`**3.219.9 Mupad [B] (verification not implemented)**

Time = 15.40 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.60

$$\int \sqrt{x\sqrt{x^{3/2}}} dx = \frac{8x\sqrt{x\sqrt{x^{3/2}}}}{15}$$

input `int((x*(x^(3/2))^(1/2))^(1/2),x)`output `(8*x*(x*(x^(3/2))^(1/2))^(1/2))/15`

3.220 $\int \cos^4(x)(\cos(x) - \sin(x)) \sin^4(x)(\cos(x) + \sin(x)) dx$

3.220.1 Optimal result	1125
3.220.2 Mathematica [B] (verified)	1125
3.220.3 Rubi [A] (verified)	1126
3.220.4 Maple [A] (verified)	1127
3.220.5 Fracas [A] (verification not implemented)	1127
3.220.6 Sympy [A] (verification not implemented)	1128
3.220.7 Maxima [A] (verification not implemented)	1128
3.220.8 Giac [A] (verification not implemented)	1128
3.220.9 Mupad [B] (verification not implemented)	1129

3.220.1 Optimal result

Integrand size = 21, antiderivative size = 12

$$\int \cos^4(x)(\cos(x) - \sin(x)) \sin^4(x)(\cos(x) + \sin(x)) dx = \frac{1}{5} \cos^5(x) \sin^5(x)$$

output `1/5*cos(x)^5*sin(x)^5`

3.220.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 25 vs. 2(12) = 24.

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 2.08

$$\int \cos^4(x)(\cos(x) - \sin(x)) \sin^4(x)(\cos(x) + \sin(x)) dx = \frac{1}{256} \sin(2x) - \frac{1}{512} \sin(6x) + \frac{\sin(10x)}{2560}$$

input `Integrate[Cos[x]^4*(Cos[x] - Sin[x])*Sin[x]^4*(Cos[x] + Sin[x]),x]`

output `Sin[2*x]/256 - Sin[6*x]/512 + Sin[10*x]/2560`

3.220.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.33, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 4889, 356}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin^4(x) \cos^4(x) (\cos(x) - \sin(x)) (\sin(x) + \cos(x)) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin(x)^4 \cos(x)^4 (\cos(x) - \sin(x)) (\sin(x) + \cos(x)) dx \\ & \quad \downarrow \text{4889} \\ & \int \frac{\tan^4(x) (1 - \tan^2(x))}{(\tan^2(x) + 1)^6} d \tan(x) \\ & \quad \downarrow \text{356} \\ & \frac{\tan^5(x)}{5 (\tan^2(x) + 1)^5} \end{aligned}$$

input `Int[Cos[x]^4*(Cos[x] - Sin[x])*Sin[x]^4*(Cos[x] + Sin[x]),x]`

output `Tan[x]^5/(5*(1 + Tan[x]^2)^5)`

3.220.3.1 Defintions of rubi rules used

rule 356 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2), x_Symbol] :> Simp[c*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*e*(m + 1))), x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d, 0] && EqQ[a*d*(m + 1) - b*c*(m + 2*p + 3), 0] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4889 Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, With[{d = FreeFactors
[Tan[v], x]}, Simp[d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x
^2), Tan[v]/d, u, x], x], x, Tan[v]/d, x]] /; !FalseQ[v] && FunctionOfQ[N
onfreeFactors[Tan[v], x], u, x]] /; InverseFunctionFreeQ[u, x] && !MatchQ[
u, (v_.)*((c_.)*tan[w_]^(n_.)*tan[z_]^(n_.))^(p_.) /; FreeQ[{c, p}, x] && I
ntegerQ[n] && LinearQ[w, x] && EqQ[z, 2*w]]
```

3.220.4 Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.67

method	result
risch	$\frac{\sin(10x)}{2560} - \frac{\sin(6x)}{512} + \frac{\sin(2x)}{256}$
parallelrisch	$\frac{\sin(10x)}{2560} - \frac{\sin(6x)}{512} + \frac{\sin(2x)}{256}$
default	$-\frac{\sin(x)^3 \cos(x)^7}{10} - \frac{3 \sin(x) \cos(x)^7}{80} + \frac{\left(\cos(x)^5 + \frac{5 \cos(x)^3}{4} + \frac{15 \cos(x)}{8}\right) \sin(x)}{160} + \frac{\cos(x)^5 \sin(x)^5}{10} + \frac{\sin(x)^3 \cos(x)^5}{16} + \dots$
parts	$-\frac{\sin(x)^3 \cos(x)^7}{10} - \frac{3 \sin(x) \cos(x)^7}{80} + \frac{\left(\cos(x)^5 + \frac{5 \cos(x)^3}{4} + \frac{15 \cos(x)}{8}\right) \sin(x)}{160} + \frac{\cos(x)^5 \sin(x)^5}{10} + \frac{\sin(x)^3 \cos(x)^5}{16} + \dots$

```
input int(sin(x)^4*cos(x)^4*(cos(x)+sin(x))*(cos(x)-sin(x)),x,method=_RETURNVERB
OSE)
```

```
output 1/2560*sin(10*x)-1/512*sin(6*x)+1/256*sin(2*x)
```

3.220.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.58

$$\int \cos^4(x)(\cos(x) - \sin(x)) \sin^4(x)(\cos(x) + \sin(x)) dx$$

$$= \frac{1}{5} (\cos(x)^9 - 2 \cos(x)^7 + \cos(x)^5) \sin(x)$$

```
input integrate(sin(x)^4*cos(x)^4*(cos(x)+sin(x))*(cos(x)-sin(x)),x, algorithm="
fracas")
```

```
output 1/5*(cos(x)^9 - 2*cos(x)^7 + cos(x)^5)*sin(x)
```

3.220. $\int \cos^4(x)(\cos(x) - \sin(x)) \sin^4(x)(\cos(x) + \sin(x)) dx$

3.220.6 Sympy [A] (verification not implemented)

Time = 1.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \cos^4(x)(\cos(x) - \sin(x)) \sin^4(x)(\cos(x) + \sin(x)) dx = \frac{\sin^5(x) \cos^5(x)}{5}$$

input `integrate(sin(x)**4*cos(x)**4*(cos(x)+sin(x))*(cos(x)-sin(x)),x)`output `sin(x)**5*cos(x)**5/5`**3.220.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \cos^4(x)(\cos(x) - \sin(x)) \sin^4(x)(\cos(x) + \sin(x)) dx = \frac{1}{160} \sin(2x)^5$$

input `integrate(sin(x)^4*cos(x)^4*(cos(x)+sin(x))*(cos(x)-sin(x)),x, algorithm="maxima")`output `1/160*sin(2*x)^5`**3.220.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.58

$$\begin{aligned} & \int \cos^4(x)(\cos(x) - \sin(x)) \sin^4(x)(\cos(x) + \sin(x)) dx \\ &= \frac{1}{2560} \sin(10x) - \frac{1}{512} \sin(6x) + \frac{1}{256} \sin(2x) \end{aligned}$$

input `integrate(sin(x)^4*cos(x)^4*(cos(x)+sin(x))*(cos(x)-sin(x)),x, algorithm="giac")`output `1/2560*sin(10*x) - 1/512*sin(6*x) + 1/256*sin(2*x)`

3.220.9 Mupad [B] (verification not implemented)

Time = 15.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.33

$$\int \cos^4(x)(\cos(x) - \sin(x)) \sin^4(x)(\cos(x) + \sin(x)) dx = \frac{\cos(x)^5 \sin(x) (\cos(x)^2 - 1)^2}{5}$$

input `int(cos(x)^4*sin(x)^4*(cos(x) + sin(x))*(cos(x) - sin(x)),x)`

output `(cos(x)^5*sin(x)*(cos(x)^2 - 1)^2)/5`

3.221 $\int \log(1 + x^2) dx$

3.221.1 Optimal result	1130
3.221.2 Mathematica [A] (verified)	1130
3.221.3 Rubi [A] (verified)	1131
3.221.4 Maple [A] (verified)	1132
3.221.5 Fricas [A] (verification not implemented)	1132
3.221.6 Sympy [A] (verification not implemented)	1133
3.221.7 Maxima [A] (verification not implemented)	1133
3.221.8 Giac [A] (verification not implemented)	1133
3.221.9 Mupad [B] (verification not implemented)	1134

3.221.1 Optimal result

Integrand size = 6, antiderivative size = 16

$$\int \log(1 + x^2) dx = -2x + 2 \arctan(x) + x \log(1 + x^2)$$

output `-2*x+2*arctan(x)+x*ln(x^2+1)`

3.221.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \log(1 + x^2) dx = -2x + 2 \arctan(x) + x \log(1 + x^2)$$

input `Integrate[Log[1 + x^2],x]`

output `-2*x + 2*ArcTan[x] + x*Log[1 + x^2]`

3.221.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {2898, 262, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \log(x^2 + 1) dx \\ & \quad \downarrow \text{2898} \\ & x \log(x^2 + 1) - 2 \int \frac{x^2}{x^2 + 1} dx \\ & \quad \downarrow \text{262} \\ & x \log(x^2 + 1) - 2 \left(x - \int \frac{1}{x^2 + 1} dx \right) \\ & \quad \downarrow \text{216} \\ & x \log(x^2 + 1) - 2(x - \arctan(x)) \end{aligned}$$

input `Int[Log[1 + x^2], x]`

output `-2*(x - ArcTan[x]) + x*Log[1 + x^2]`

3.221.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 262 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2898 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_))^(p_.)], x_Symbol] := Simp[x*Log[c*(d + e*x^n)^p], x] - Simp[e*n*p Int[x^n/(d + e*x^n), x], x] /; FreeQ[{c, d, e, n, p}, x]`

3.221.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

method	result	size
default	$-2x + 2 \arctan(x) + x \ln(x^2 + 1)$	17
risch	$-2x + 2 \arctan(x) + x \ln(x^2 + 1)$	17
parts	$-2x + 2 \arctan(x) + x \ln(x^2 + 1)$	17
meijerg	$-2x + \frac{2x \arctan(\sqrt{x^2})}{\sqrt{x^2}} + x \ln(x^2 + 1)$	27
parallelrisc	$-2i \ln(x - i) + i \ln(x^2 + 1) + x \ln(x^2 + 1) - 2x$	30

input `int(ln(x^2+1),x,method=_RETURNVERBOSE)`

output `-2*x+2*arctan(x)+x*ln(x^2+1)`

3.221.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \log(1 + x^2) dx = x \log(x^2 + 1) - 2x + 2 \arctan(x)$$

input `integrate(log(x^2+1),x, algorithm="fricas")`

output `x*log(x^2 + 1) - 2*x + 2*arctan(x)`

3.221.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \log(1+x^2) dx = x \log(x^2+1) - 2x + 2 \operatorname{atan}(x)$$

input `integrate(ln(x**2+1),x)`output `x*log(x**2 + 1) - 2*x + 2*atan(x)`**3.221.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \log(1+x^2) dx = x \log(x^2+1) - 2x + 2 \operatorname{arctan}(x)$$

input `integrate(log(x^2+1),x, algorithm="maxima")`output `x*log(x^2 + 1) - 2*x + 2*arctan(x)`**3.221.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \log(1+x^2) dx = x \log(x^2+1) - 2x + 2 \operatorname{arctan}(x)$$

input `integrate(log(x^2+1),x, algorithm="giac")`output `x*log(x^2 + 1) - 2*x + 2*arctan(x)`

3.221.9 Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \log(1 + x^2) dx = 2 \operatorname{atan}(x) - 2x + x \ln(x^2 + 1)$$

input `int(log(x^2 + 1),x)`

output `2*atan(x) - 2*x + x*log(x^2 + 1)`

3.222 $\int \frac{1+2x}{1+2x+2x^2} dx$

3.222.1 Optimal result	1135
3.222.2 Mathematica [A] (verified)	1135
3.222.3 Rubi [A] (verified)	1136
3.222.4 Maple [A] (verified)	1136
3.222.5 Fricas [A] (verification not implemented)	1137
3.222.6 Sympy [A] (verification not implemented)	1137
3.222.7 Maxima [A] (verification not implemented)	1137
3.222.8 Giac [A] (verification not implemented)	1138
3.222.9 Mupad [B] (verification not implemented)	1138

3.222.1 Optimal result

Integrand size = 18, antiderivative size = 15

$$\int \frac{1+2x}{1+2x+2x^2} dx = \frac{1}{2} \log(1+2x+2x^2)$$

output `1/2*ln(2*x^2+2*x+1)`

3.222.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{1+2x}{1+2x+2x^2} dx = \frac{1}{2} \log(1+2x+2x^2)$$

input `Integrate[(1 + 2*x)/(1 + 2*x + 2*x^2), x]`

output `Log[1 + 2*x + 2*x^2]/2`

3.222.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x + 1}{2x^2 + 2x + 1} dx$$

↓ 1103

$$\frac{1}{2} \log(2x^2 + 2x + 1)$$

input `Int[(1 + 2*x)/(1 + 2*x + 2*x^2),x]`

output `Log[1 + 2*x + 2*x^2]/2`

3.222.3.1 Defintions of rubi rules used

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] :> Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

3.222.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

method	result	size
parallelrisch	$\frac{\ln(x^2+x+\frac{1}{2})}{2}$	10
default	$\frac{\ln(2x^2+2x+1)}{2}$	14
norman	$\frac{\ln(2x^2+2x+1)}{2}$	14
risch	$\frac{\ln(2x^2+2x+1)}{2}$	14

input `int((1+2*x)/(2*x^2+2*x+1),x,method=_RETURNVERBOSE)`

output $1/2*\ln(x^2+x+1/2)$

3.222.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{1+2x}{1+2x+2x^2} dx = \frac{1}{2} \log(2x^2+2x+1)$$

input `integrate((1+2*x)/(2*x^2+2*x+1),x, algorithm="fricas")`

output $1/2*\log(2*x^2 + 2*x + 1)$

3.222.6 Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \frac{1+2x}{1+2x+2x^2} dx = \frac{\log(2x^2+2x+1)}{2}$$

input `integrate((1+2*x)/(2*x**2+2*x+1),x)`

output $\log(2*x**2 + 2*x + 1)/2$

3.222.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{1+2x}{1+2x+2x^2} dx = \frac{1}{2} \log(2x^2+2x+1)$$

input `integrate((1+2*x)/(2*x^2+2*x+1),x, algorithm="maxima")`

output $1/2*\log(2*x^2 + 2*x + 1)$

3.222.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{1+2x}{1+2x+2x^2} dx = \frac{1}{2} \log(2x^2+2x+1)$$

input `integrate((1+2*x)/(2*x^2+2*x+1),x, algorithm="giac")`

output `1/2*log(2*x^2 + 2*x + 1)`

3.222.9 Mupad [B] (verification not implemented)

Time = 14.94 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.60

$$\int \frac{1+2x}{1+2x+2x^2} dx = \frac{\ln(x^2+x+\frac{1}{2})}{2}$$

input `int((2*x + 1)/(2*x + 2*x^2 + 1),x)`

output `log(x + x^2 + 1/2)/2`

3.223 $\int \frac{\arcsin(x)}{x^3} dx$

3.223.1 Optimal result	1139
3.223.2 Mathematica [A] (verified)	1139
3.223.3 Rubi [A] (verified)	1140
3.223.4 Maple [A] (verified)	1141
3.223.5 Fricas [A] (verification not implemented)	1141
3.223.6 Sympy [C] (verification not implemented)	1141
3.223.7 Maxima [A] (verification not implemented)	1142
3.223.8 Giac [A] (verification not implemented)	1142
3.223.9 Mupad [F(-1)]	1142

3.223.1 Optimal result

Integrand size = 6, antiderivative size = 28

$$\int \frac{\arcsin(x)}{x^3} dx = -\frac{\sqrt{1-x^2}}{2x} - \frac{\arcsin(x)}{2x^2}$$

output `-1/2*(-x^2+1)^(1/2)/x-1/2*arcsin(x)/x^2`

3.223.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

$$\int \frac{\arcsin(x)}{x^3} dx = -\frac{x\sqrt{1-x^2} + \arcsin(x)}{2x^2}$$

input `Integrate[ArcSin[x]/x^3,x]`

output `-1/2*(x*Sqrt[1 - x^2] + ArcSin[x])/x^2`

3.223.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5138, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arcsin(x)}{x^3} dx$$

↓ 5138

$$\frac{1}{2} \int \frac{1}{x^2 \sqrt{1-x^2}} dx - \frac{\arcsin(x)}{2x^2}$$

↓ 242

$$-\frac{\arcsin(x)}{2x^2} - \frac{\sqrt{1-x^2}}{2x}$$

input `Int[ArcSin[x]/x^3,x]`

output `-1/2*Sqrt[1 - x^2]/x - ArcSin[x]/(2*x^2)`

3.223.3.1 Defintions of rubi rules used

rule 242 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

rule 5138 `Int[((a_.) + ArcSin[(c_.)*(x_)])*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

3.223.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{\sqrt{-x^2+1}}{2x} - \frac{\arcsin(x)}{2x^2}$	23
parts	$-\frac{\sqrt{-x^2+1}}{2x} - \frac{\arcsin(x)}{2x^2}$	23

input `int(arcsin(x)/x^3,x,method=_RETURNVERBOSE)`

output `-1/2*(-x^2+1)^(1/2)/x-1/2*arcsin(x)/x^2`

3.223.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.68

$$\int \frac{\arcsin(x)}{x^3} dx = -\frac{\sqrt{-x^2+1}x + \arcsin(x)}{2x^2}$$

input `integrate(arcsin(x)/x^3,x, algorithm="fracas")`

output `-1/2*(sqrt(-x^2 + 1)*x + arcsin(x))/x^2`

3.223.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.78 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.32

$$\int \frac{\arcsin(x)}{x^3} dx = \frac{\begin{cases} -\frac{i\sqrt{x^2-1}}{x} & \text{for } |x^2| > 1 \\ -\frac{\sqrt{1-x^2}}{x} & \text{otherwise} \end{cases}}{2} - \frac{\operatorname{asin}(x)}{2x^2}$$

input `integrate(asin(x)/x**3,x)`

output `Piecewise((-I*sqrt(x**2 - 1)/x, Abs(x**2) > 1), (-sqrt(1 - x**2)/x, True)) /2 - asin(x)/(2*x**2)`

3.223.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

$$\int \frac{\arcsin(x)}{x^3} dx = -\frac{\sqrt{-x^2+1}}{2x} - \frac{\arcsin(x)}{2x^2}$$

input `integrate(arcsin(x)/x^3,x, algorithm="maxima")`output `-1/2*sqrt(-x^2 + 1)/x - 1/2*arcsin(x)/x^2`**3.223.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.43

$$\int \frac{\arcsin(x)}{x^3} dx = \frac{x}{4(\sqrt{-x^2+1}-1)} - \frac{\sqrt{-x^2+1}-1}{4x} - \frac{\arcsin(x)}{2x^2}$$

input `integrate(arcsin(x)/x^3,x, algorithm="giac")`output `1/4*x/(sqrt(-x^2 + 1) - 1) - 1/4*(sqrt(-x^2 + 1) - 1)/x - 1/2*arcsin(x)/x^2`**3.223.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\arcsin(x)}{x^3} dx = \int \frac{\operatorname{asin}(x)}{x^3} dx$$

input `int(asin(x)/x^3,x)`output `int(asin(x)/x^3, x)`

3.224 $\int \cos(\cos(x)) \sin(2x) dx$

3.224.1 Optimal result	1143
3.224.2 Mathematica [A] (verified)	1143
3.224.3 Rubi [A] (verified)	1144
3.224.4 Maple [A] (verified)	1145
3.224.5 Fricas [B] (verification not implemented)	1146
3.224.6 Sympy [A] (verification not implemented)	1146
3.224.7 Maxima [A] (verification not implemented)	1146
3.224.8 Giac [A] (verification not implemented)	1147
3.224.9 Mupad [B] (verification not implemented)	1147

3.224.1 Optimal result

Integrand size = 8, antiderivative size = 13

$$\int \cos(\cos(x)) \sin(2x) dx = -2 \cos(\cos(x)) - 2 \cos(x) \sin(\cos(x))$$

output `-2*cos(cos(x))-2*cos(x)*sin(cos(x))`

3.224.2 Mathematica [A] (verified)

Time = 3.44 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \cos(\cos(x)) \sin(2x) dx = -2 \cos(\cos(x)) - 2 \cos(x) \sin(\cos(x))$$

input `Integrate[Cos[Cos[x]]*Sin[2*x],x]`

output `-2*Cos[Cos[x]] - 2*Cos[x]*Sin[Cos[x]]`

3.224.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {4879, 27, 3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(2x) \cos(\cos(x)) dx \\
 & \quad \downarrow 4879 \\
 & - \int 2 \cos(x) \cos(\cos(x)) d \cos(x) \\
 & \quad \downarrow 27 \\
 & -2 \int \cos(x) \cos(\cos(x)) d \cos(x) \\
 & \quad \downarrow 3042 \\
 & -2 \int \cos(x) \sin\left(\cos(x) + \frac{\pi}{2}\right) d \cos(x) \\
 & \quad \downarrow 3777 \\
 & -2\left(\int -\sin(\cos(x)) d \cos(x) + \cos(x) \sin(\cos(x))\right) \\
 & \quad \downarrow 25 \\
 & -2(\cos(x) \sin(\cos(x)) - \int \sin(\cos(x)) d \cos(x)) \\
 & \quad \downarrow 3042 \\
 & -2(\cos(x) \sin(\cos(x)) - \int \sin(\cos(x)) d \cos(x)) \\
 & \quad \downarrow 3118 \\
 & -2(\cos(\cos(x)) + \cos(x) \sin(\cos(x)))
 \end{aligned}$$

input `Int[Cos[Cos[x]]*Sin[2*x],x]`

output `-2*(Cos[Cos[x]] + Cos[x]*Sin[Cos[x]])`

3.224.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 4879 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Cos[v], x]}, -d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Cos[v]/d, u/Sin[v], x], x], x, Cos[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Cos[v], x], u/Sin[v], x]`

3.224.4 Maple [A] (verified)

Time = 12.92 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.62

method	result	size
risch	$\sin(x - \cos(x)) - \sin(x + \cos(x)) - 2 \cos(\cos(x))$	21

input `int(sin(2*x)*cos(cos(x)),x,method=_RETURNVERBOSE)`

output `sin(x-cos(x))-sin(x+cos(x))-2*cos(cos(x))`

3.224.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 47 vs. $2(13) = 26$.

Time = 0.25 (sec) , antiderivative size = 47, normalized size of antiderivative = 3.62

$$\int \cos(\cos(x)) \sin(2x) dx = 2 \cos(x) \sin\left(\frac{\tan\left(\frac{1}{2}x\right)^2 - 1}{\tan\left(\frac{1}{2}x\right)^2 + 1}\right) - 2 \cos\left(\frac{\tan\left(\frac{1}{2}x\right)^2 - 1}{\tan\left(\frac{1}{2}x\right)^2 + 1}\right)$$

input `integrate(sin(2*x)*cos(cos(x)),x, algorithm="fricas")`

output `2*cos(x)*sin((tan(1/2*x)^2 - 1)/(tan(1/2*x)^2 + 1)) - 2*cos((tan(1/2*x)^2 - 1)/(tan(1/2*x)^2 + 1))`

3.224.6 Sympy [A] (verification not implemented)

Time = 0.84 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31

$$\int \cos(\cos(x)) \sin(2x) dx = -2 \sin(\cos(x)) \cos(x) - 2 \cos(\cos(x))$$

input `integrate(sin(2*x)*cos(cos(x)),x)`

output `-2*sin(cos(x))*cos(x) - 2*cos(cos(x))`

3.224.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \cos(\cos(x)) \sin(2x) dx = -2 \cos(x) \sin(\cos(x)) - 2 \cos(\cos(x))$$

input `integrate(sin(2*x)*cos(cos(x)),x, algorithm="maxima")`

output `-2*cos(x)*sin(cos(x)) - 2*cos(cos(x))`

3.224.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \cos(\cos(x)) \sin(2x) dx = -2 \cos(x) \sin(\cos(x)) - 2 \cos(\cos(x))$$

input `integrate(sin(2*x)*cos(cos(x)),x, algorithm="giac")`output `-2*cos(x)*sin(cos(x)) - 2*cos(cos(x))`**3.224.9 Mupad [B] (verification not implemented)**

Time = 15.87 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.85

$$\int \cos(\cos(x)) \sin(2x) dx = -4 \cos\left(\frac{\cos(x)}{2}\right)^2 - 4 \sin\left(\frac{\cos(x)}{2}\right) \cos(x) \cos\left(\frac{\cos(x)}{2}\right)$$

input `int(cos(cos(x))*sin(2*x),x)`output `- 4*cos(cos(x)/2)^2 - 4*cos(cos(x)/2)*sin(cos(x)/2)*cos(x)`

3.225 $\int -\sin(x - \sin(x)) dx$

3.225.1 Optimal result	1148
3.225.2 Mathematica [N/A]	1148
3.225.3 Rubi [N/A]	1149
3.225.4 Maple [N/A] (verified)	1150
3.225.5 Fricas [N/A]	1150
3.225.6 Sympy [N/A]	1150
3.225.7 Maxima [N/A]	1151
3.225.8 Giac [N/A]	1151
3.225.9 Mupad [N/A]	1151

3.225.1 Optimal result

Integrand size = 9, antiderivative size = 9

$$\int -\sin(x - \sin(x)) dx = -\text{Int}(\sin(x - \sin(x)), x)$$

output `-CannotIntegrate(sin(x-sin(x)),x)`

3.225.2 Mathematica [N/A]

Not integrable

Time = 1.89 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.22

$$\int -\sin(x - \sin(x)) dx = \int -\sin(x - \sin(x)) dx$$

input `Integrate[-Sin[x - Sin[x]],x]`

output `-Integrate[Sin[x - Sin[x]], x]`

3.225.3 Rubi [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {25, 7299}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int -\sin(x - \sin(x)) dx \\ \downarrow 25 \\ - \int \sin(x - \sin(x)) dx \\ \downarrow 7299 \\ - \int \sin(x - \sin(x)) dx \end{array}$$

input `Int[-Sin[x - Sin[x]],x]`

output `$Aborted`

3.225.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 7299 `Int[u_, x_] := CannotIntegrate[u, x]`

3.225.4 Maple [N/A] (verified)

Not integrable

Time = 0.04 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int -\sin(x - \sin(x)) dx$$

input `int(-sin(x-sin(x)),x)`output `int(-sin(x-sin(x)),x)`**3.225.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int -\sin(x - \sin(x)) dx = \int -\sin(x - \sin(x)) dx$$

input `integrate(-sin(x-sin(x)),x, algorithm="fricas")`output `integral(sin(-x + sin(x)), x)`**3.225.6 Sympy [N/A]**

Not integrable

Time = 0.49 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int -\sin(x - \sin(x)) dx = -\int \sin(x - \sin(x)) dx$$

input `integrate(-sin(x-sin(x)),x)`output `-Integral(sin(x - sin(x)), x)`

3.225.7 Maxima [N/A]

Not integrable

Time = 0.40 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int -\sin(x - \sin(x)) dx = \int -\sin(x - \sin(x)) dx$$

input `integrate(-sin(x-sin(x)),x, algorithm="maxima")`output `integrate(sin(-x + sin(x)), x)`**3.225.8 Giac [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.22

$$\int -\sin(x - \sin(x)) dx = \int -\sin(x - \sin(x)) dx$$

input `integrate(-sin(x-sin(x)),x, algorithm="giac")`output `integrate(-sin(x - sin(x)), x)`**3.225.9 Mupad [N/A]**

Not integrable

Time = 15.40 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.22

$$\int -\sin(x - \sin(x)) dx = \int -\sin(x - \sin(x)) dx$$

input `int(-sin(x - sin(x)),x)`output `int(-sin(x - sin(x)), x)`

3.226 $\int \frac{1}{1+\tan^{2\sqrt{505}}(x)} dx$

3.226.1 Optimal result	1152
3.226.2 Mathematica [N/A]	1152
3.226.3 Rubi [N/A]	1153
3.226.4 Maple [N/A] (verified)	1154
3.226.5 Fricas [N/A]	1154
3.226.6 Sympy [N/A]	1154
3.226.7 Maxima [N/A]	1155
3.226.8 Giac [N/A]	1155
3.226.9 Mupad [N/A]	1156

3.226.1 Optimal result

Integrand size = 14, antiderivative size = 14

$$\int \frac{1}{1 + \tan^{2\sqrt{505}}(x)} dx = \text{Int}\left(\frac{1}{1 + \tan^{2\sqrt{505}}(x)}, x\right)$$

output `Unintegrable(1/(tan(x)^(2*505^(1/2))+1), x)`

3.226.2 Mathematica [N/A]

Not integrable

Time = 9.78 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{1}{1 + \tan^{2\sqrt{505}}(x)} dx = \int \frac{1}{1 + \tan^{2\sqrt{505}}(x)} dx$$

input `Integrate[(1 + Tan[x]^(2*Sqrt[505]))^(-1), x]`

output `Integrate[(1 + Tan[x]^(2*Sqrt[505]))^(-1), x]`

3.226.3 Rubi [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3042, 4145}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\tan^{2\sqrt{505}}(x) + 1} dx$$

↓ 3042

$$\int \frac{1}{\tan(x)^{2\sqrt{505}} + 1} dx$$

↓ 4145

$$\int \frac{1}{\tan^{2\sqrt{505}}(x) + 1} dx$$

input `Int[(1 + Tan[x]^(2*Sqrt[505]))^(-1), x]`

output `$Aborted`

3.226.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4145 `Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_.), x_Symbol] := Unintegrable[(a + b*(c*Tan[e + f*x])^n)^p, x] /; FreeQ[{a, b, c, e, f, n, p}, x]`

3.226.4 Maple [N/A] (verified)

Not integrable

Time = 0.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{\tan(x)^{2\sqrt{505}} + 1} dx$$

input `int(1/(tan(x)^(2*505^(1/2))+1),x)`output `int(1/(tan(x)^(2*505^(1/2))+1),x)`**3.226.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \tan^{2\sqrt{505}}(x)} dx = \int \frac{1}{\tan(x)^{2\sqrt{505}} + 1} dx$$

input `integrate(1/(tan(x)^(2*505^(1/2))+1),x, algorithm="fricas")`output `integral(1/(tan(x)^(2*sqrt(505)) + 1), x)`**3.226.6 Sympy [N/A]**

Not integrable

Time = 0.80 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \tan^{2\sqrt{505}}(x)} dx = \int \frac{1}{\tan^{2\sqrt{505}}(x) + 1} dx$$

input `integrate(1/(tan(x)**(2*505**(1/2))+1),x)`output `Integral(1/(tan(x)**(2*sqrt(505)) + 1), x)`

3.226.7 Maxima [N/A]

Not integrable

Time = 0.74 (sec) , antiderivative size = 1038, normalized size of antiderivative = 74.14

$$\int \frac{1}{1 + \tan^{2\sqrt{505}}(x)} dx = \int \frac{1}{\tan(x)^{2\sqrt{505}} + 1} dx$$

```
input integrate(1/(tan(x)^(2*505^(1/2))+1),x, algorithm="maxima")
```

```
output -(-1)^(sqrt(101)*sqrt(5))*integrate(((1)^(sqrt(101)*sqrt(5))*cos(2*sqrt(101)*sqrt(5)*arctan2(sin(x), cos(x) + 1) - 2*sqrt(101)*sqrt(5)*arctan2(sin(x), -cos(x) + 1))^2*e^(2*sqrt(101)*sqrt(5)*log(cos(x)^2 + sin(x)^2 + 2*cos(x) + 1) + 2*sqrt(101)*sqrt(5)*log(cos(x)^2 + sin(x)^2 - 2*cos(x) + 1)) + (-1)^(sqrt(101)*sqrt(5))*e^(2*sqrt(101)*sqrt(5)*log(cos(x)^2 + sin(x)^2 + 2*cos(x) + 1) + 2*sqrt(101)*sqrt(5)*log(cos(x)^2 + sin(x)^2 - 2*cos(x) + 1)))*sin(2*sqrt(101)*sqrt(5)*arctan2(sin(x), cos(x) + 1) - 2*sqrt(101)*sqrt(5)*arctan2(sin(x), -cos(x) + 1))^2 + cos(2*sqrt(101)*sqrt(5)*arctan2(sin(2*x), cos(2*x) + 1))*cos(2*sqrt(101)*sqrt(5)*arctan2(sin(x), cos(x) + 1) - 2*sqrt(101)*sqrt(5)*arctan2(sin(x), -cos(x) + 1))*e^(sqrt(101)*sqrt(5)*log(cos(2*x)^2 + sin(2*x)^2 + 2*cos(2*x) + 1) + sqrt(101)*sqrt(5)*log(cos(x)^2 + sin(x)^2 + 2*cos(x) + 1) + sqrt(101)*sqrt(5)*log(cos(x)^2 + sin(x)^2 - 2*cos(x) + 1)) + e^(sqrt(101)*sqrt(5)*log(cos(2*x)^2 + sin(2*x)^2 + 2*cos(2*x) + 1) + sqrt(101)*sqrt(5)*log(cos(x)^2 + sin(x)^2 + 2*cos(x) + 1) + sqrt(101)*sqrt(5)*log(cos(x)^2 + sin(x)^2 - 2*cos(x) + 1))*sin(2*sqrt(101)*sqrt(5)*arctan2(sin(2*x), cos(2*x) + 1))*sin(2*sqrt(101)*sqrt(5)*arctan2(sin(x), cos(x) + 1) - 2*sqrt(101)*sqrt(5)*arctan2(sin(x), -cos(x) + 1)))/(2*(-1)^(sqrt(101)*sqrt(5))*cos(2*sqrt(101)*sqrt(5)*arctan2(sin(2*x), cos(2*x) + 1))*cos(2*sqrt(101)*sqrt(5)*arctan2(sin(x), cos(x) + 1) - 2*sqrt(101)*sqrt(5)*arctan2(sin(x), -cos(x) + 1))*e^(sqrt(101)*sqrt(5)*log(cos(2*x)...
```

3.226.8 Giac [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \tan^{2\sqrt{505}}(x)} dx = \int \frac{1}{\tan(x)^{2\sqrt{505}} + 1} dx$$

input `integrate(1/(tan(x)^(2*505^(1/2))+1),x, algorithm="giac")`

output `integrate(1/(tan(x)^(2*sqrt(505)) + 1), x)`

3.226.9 Mupad [N/A]

Not integrable

Time = 15.38 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \tan^{2\sqrt{505}}(x)} dx = \int \frac{1}{\tan(x)^{2\sqrt{505}} + 1} dx$$

input `int(1/(tan(x)^(2*505^(1/2)) + 1),x)`

output `int(1/(tan(x)^(2*505^(1/2)) + 1), x)`

3.227 $\int (1 - x)^{2020} x dx$

3.227.1 Optimal result	1157
3.227.2 Mathematica [B] (verified)	1157
3.227.3 Rubi [A] (verified)	1158
3.227.4 Maple [B] (verified)	1159
3.227.5 Fricas [F(-2)]	1159
3.227.6 Sympy [B] (verification not implemented)	1159
3.227.7 Maxima [B] (verification not implemented)	1160
3.227.8 Giac [B] (verification not implemented)	1161
3.227.9 Mupad [B] (verification not implemented)	1162

3.227.1 Optimal result

Integrand size = 9, antiderivative size = 23

$$\int (1 - x)^{2020} x dx = -\frac{(1 - x)^{2021}}{2021} + \frac{(1 - x)^{2022}}{2022}$$

output `-1/2021*(1-x)^2021+1/2022*(1-x)^2022`

3.227.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 11128 vs. 2(23) = 46.

Time = 0.14 (sec) , antiderivative size = 11128, normalized size of antiderivative = 483.83

$$\int (1 - x)^{2020} x dx = \text{Result too large to show}$$

input `Integrate[(1 - x)^2020*x,x]`

output `Result too large to show`

3.227.3 Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (1-x)^{2020} x dx$$

$$\downarrow 49$$

$$\int ((1-x)^{2020} - (1-x)^{2021}) dx$$

$$\downarrow 2009$$

$$\frac{(1-x)^{2022}}{2022} - \frac{(1-x)^{2021}}{2021}$$

input `Int[(1 - x)^2020*x,x]`

output `-1/2021*(1 - x)^2021 + (1 - x)^2022/2022`

3.227.3.1 Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.227.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 10105 vs. $2(19) = 38$.

Time = 9.33 (sec) , antiderivative size = 10106, normalized size of antiderivative = 439.39

method	result	size
gospers	Expression too large to display	10106
default	Expression too large to display	10107
risch	Expression too large to display	10107
parallelrisch	Expression too large to display	10107

input `int(x*(1-x)^2020,x,method=_RETURNVERBOSE)`

output `result too large to display`

3.227.5 Fracas [F(-2)]

Exception generated.

$$\int (1-x)^{2020} x dx = \text{Exception raised: RuntimeError}$$

input `integrate(x*(1-x)^2020,x, algorithm="fricas")`

output `Exception raised: RuntimeError >> System error: Heap exhausted (no more space for allocation).1998848 bytes available, 2179456 requested.PROCEED WITH CAUTION.`

3.227.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11171 vs. $2(12) = 24$.

Time = 2.60 (sec) , antiderivative size = 11171, normalized size of antiderivative = 485.70

$$\int (1-x)^{2020} x dx = \text{Too large to display}$$

input `integrate(x*(1-x)**2020,x)`

output `x**2022/2022 - 2020*x**2021/2021 + 2019*x**2020/2 - 2038180*x**2019/3 + 685507705*x**2018/2 - 138266870112*x**2017 + 278745941561035*x**2016/6 - 13373165093095968*x**2015 + 3366693482174180730*x**2014 - 2259050769046992147760*x**2013/3 + 151506967484463186448138*x**2012 - 27698221479285170723685360*x**2011 + 13918352848288375491988868066*x**2010/3 - 716973434466946772290161333280*x**2009 + 102834449953653272749024785712470*x**2008 - 41277737963817952963643626311862816*x**2007/3 + 1725065037047525925910348951124302995*x**2006 - 203456149346983815836345155695893355528*x**2005 + 67954336961055157115435769332338918346705*x**2004/3 - 2387938649240603820483180755599043532230840*x**2003 + 239032599150157038818882365583996835022772313*x**2002 - 68329158777556135034683015049498345643691036000*x**2001/3 + 2070580051371710322595432011477108583912261365977*x**2000 - 179960368994961708008163784454456051967579363996000*x**1999 + 44945090897709842545903018770867237413834629868000750*x**1998/3 - 1196737653552288657999009221047362441958082554435826640*x**1997 + 91872606035477133069946098385058726668676396410333163850*x**1996 - 20365089221038364860461105557164855112824957167732656091280*x**1995/3 + 483428305875789720518480870193807402556007179708207427412550*x**1994 - 33223185208528449708566031169292044668543059434670067575119200*x**1993 + 6618056825708284424494058802194502943955470785618233551823806130*x**1992/3 - 14168330984944992315661556907483236515975115240006657516...`

3.227.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10106 vs. $2(15) = 30$.

Time = 2.40 (sec) , antiderivative size = 10106, normalized size of antiderivative = 439.39

$$\int (1-x)^{2020} x dx = \text{Too large to display}$$

input `integrate(x*(1-x)^2020,x, algorithm="maxima")`

```
output 1/2022*x^2022 - 2020/2021*x^2021 + 2019/2*x^2020 - 2038180/3*x^2019 + 6855
07705/2*x^2018 - 138266870112*x^2017 + 278745941561035/6*x^2016 - 13373165
093095968*x^2015 + 3366693482174180730*x^2014 - 2259050769046992147760/3*x
^2013 + 151506967484463186448138*x^2012 - 27698221479285170723685360*x^201
1 + 13918352848288375491988868066/3*x^2010 - 71697343446694677229016133328
0*x^2009 + 102834449953653272749024785712470*x^2008 - 41277737963817952963
643626311862816/3*x^2007 + 1725065037047525925910348951124302995*x^2006 -
203456149346983815836345155695893355528*x^2005 + 6795433696105515711543576
9332338918346705/3*x^2004 - 2387938649240603820483180755599043532230840*x^
2003 + 239032599150157038818882365583996835022772313*x^2002 - 683291587775
56135034683015049498345643691036000/3*x^2001 + 207058005137171032259543201
1477108583912261365977*x^2000 - 179960368994961708008163784454456051967579
363996000*x^1999 + 44945090897709842545903018770867237413834629868000750/3
*x^1998 - 1196737653552288657999009221047362441958082554435826640*x^1997 +
91872606035477133069946098385058726668676396410333163850*x^1996 - 2036508
9221038364860461105557164855112824957167732656091280/3*x^1995 + 4834283058
75789720518480870193807402556007179708207427412550*x^1994 - 33223185208528
449708566031169292044668543059434670067575119200*x^1993 + 6618056825708284
424494058802194502943955470785618233551823806130/3*x^1992 - 14168330984944
9923156615569074832365159751152400066575169519360800*x^1991 + 176218572...
```

3.227.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10106 vs. $2(15) = 30$.

Time = 1.10 (sec) , antiderivative size = 10106, normalized size of antiderivative = 439.39

$$\int (1-x)^{2020} x dx = \text{Too large to display}$$

```
input integrate(x*(1-x)^2020,x, algorithm="giac")
```

output

```

1/2022*x^2022 - 2020/2021*x^2021 + 2019/2*x^2020 - 2038180/3*x^2019 + 6855
07705/2*x^2018 - 138266870112*x^2017 + 278745941561035/6*x^2016 - 13373165
093095968*x^2015 + 3366693482174180730*x^2014 - 2259050769046992147760/3*x
^2013 + 151506967484463186448138*x^2012 - 27698221479285170723685360*x^201
1 + 13918352848288375491988868066/3*x^2010 - 71697343446694677229016133328
0*x^2009 + 102834449953653272749024785712470*x^2008 - 41277737963817952963
643626311862816/3*x^2007 + 1725065037047525925910348951124302995*x^2006 -
203456149346983815836345155695893355528*x^2005 + 6795433696105515711543576
9332338918346705/3*x^2004 - 2387938649240603820483180755599043532230840*x^
2003 + 239032599150157038818882365583996835022772313*x^2002 - 683291587775
56135034683015049498345643691036000/3*x^2001 + 207058005137171032259543201
1477108583912261365977*x^2000 - 179960368994961708008163784454456051967579
363996000*x^1999 + 44945090897709842545903018770867237413834629868000750/3
*x^1998 - 1196737653552288657999009221047362441958082554435826640*x^1997 +
91872606035477133069946098385058726668676396410333163850*x^1996 - 2036508
9221038364860461105557164855112824957167732656091280/3*x^1995 + 4834283058
75789720518480870193807402556007179708207427412550*x^1994 - 33223185208528
449708566031169292044668543059434670067575119200*x^1993 + 6618056825708284
424494058802194502943955470785618233551823806130/3*x^1992 - 14168330984944
9923156615569074832365159751152400066575169519360800*x^1991 + 176218572...

```

3.227.9 Mupad [B] (verification not implemented)

Time = 31.77 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int (1-x)^{2020} x dx = \frac{(x-1)^{2021}}{2021} + \frac{(x-1)^{2022}}{2022}$$

input `int(x*(x - 1)^2020,x)`

output `(x - 1)^2021/2021 + (x - 1)^2022/2022`

$$3.228 \quad \int \frac{\sec^4(x) \tan(x)}{4 + \sec^4(x)} dx$$

3.228.1 Optimal result	1163
3.228.2 Mathematica [A] (verified)	1163
3.228.3 Rubi [A] (verified)	1164
3.228.4 Maple [A] (verified)	1165
3.228.5 Fricas [B] (verification not implemented)	1166
3.228.6 Sympy [B] (verification not implemented)	1166
3.228.7 Maxima [A] (verification not implemented)	1167
3.228.8 Giac [B] (verification not implemented)	1167
3.228.9 Mupad [B] (verification not implemented)	1167

3.228.1 Optimal result

Integrand size = 15, antiderivative size = 11

$$\int \frac{\sec^4(x) \tan(x)}{4 + \sec^4(x)} dx = \frac{1}{4} \log(4 + \sec^4(x))$$

output `1/4*ln(sec(x)^4+4)`

3.228.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.73

$$\int \frac{\sec^4(x) \tan(x)}{4 + \sec^4(x)} dx = -\log(\cos(x)) + \frac{1}{4} \log(1 + 4 \cos^4(x))$$

input `Integrate[(Sec[x]^4*Tan[x])/(4 + Sec[x]^4),x]`

output `-Log[Cos[x]] + Log[1 + 4*Cos[x]^4]/4`

3.228.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.91, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 4839, 798, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(x) \sec^4(x)}{\sec^4(x) + 4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(x) \sec(x)^4}{\sec(x)^4 + 4} dx \\
 & \quad \downarrow \text{4839} \\
 & - \int \frac{\sec(x)}{4 \cos^4(x) + 1} d \cos(x) \\
 & \quad \downarrow \text{798} \\
 & - \frac{1}{4} \int \frac{\sec(x)}{4 \cos^4(x) + 1} d \cos^4(x) \\
 & \quad \downarrow \text{47} \\
 & \frac{1}{4} \left(4 \int \frac{1}{4 \cos^4(x) + 1} d \cos^4(x) - \int \sec(x) d \cos^4(x) \right) \\
 & \quad \downarrow \text{14} \\
 & \frac{1}{4} \left(4 \int \frac{1}{4 \cos^4(x) + 1} d \cos^4(x) - \log(\cos^4(x)) \right) \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{4} (\log(4 \cos^4(x) + 1) - \log(\cos^4(x)))
 \end{aligned}$$

input `Int[(Sec[x]^4*Tan[x])/(4 + Sec[x]^4), x]`

output `(-Log[Cos[x]^4] + Log[1 + 4*Cos[x]^4])/4`

3.228.3.1 Defintions of rubi rules used

- rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`
- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`
- rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4839 `Int[(u_)*(F_)[(c_.)*((a_.) + (b_.)*(x_))], x_Symbol] := With[{d = FreeFactors[Cos[c*(a + b*x)], x]}, Simp[-(b*c)^(-1) Subst[Int[SubstFor[1/x, Cos[c*(a + b*x)]]/d, u, x], x], x, Cos[c*(a + b*x)]]/d, u, x, True]] /; FreeQ[{a, b, c}, x] && (EqQ[F, Tan] || EqQ[F, tan])`

3.228.4 Maple [A] (verified)

Time = 3.91 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

method	result	size
derivativedivides	$\frac{\ln(\sec(x)^4+4)}{4}$	10
default	$\frac{\ln(\sec(x)^4+4)}{4}$	10
risch	$-\ln(e^{2ix} + 1) + \frac{\ln(e^{8ix}+4e^{6ix}+10e^{4ix}+4e^{2ix}+1)}{4}$	43

input `int(sec(x)^4*tan(x)/(sec(x)^4+4),x,method=_RETURNVERBOSE)`

output `1/4*ln(sec(x)^4+4)`

3.228.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(9) = 18$.

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.73

$$\int \frac{\sec^4(x) \tan(x)}{4 + \sec^4(x)} dx = \frac{1}{4} \log(4 \cos(x)^4 + 1) - \log(-\cos(x))$$

input `integrate(sec(x)^4*tan(x)/(sec(x)^4+4),x, algorithm="fricas")`

output `1/4*log(4*cos(x)^4 + 1) - log(-cos(x))`

3.228.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(8) = 16$.

Time = 0.25 (sec) , antiderivative size = 29, normalized size of antiderivative = 2.64

$$\int \frac{\sec^4(x) \tan(x)}{4 + \sec^4(x)} dx = \frac{\log(\sec^2(x) - 2 \sec(x) + 2)}{4} + \frac{\log(\sec^2(x) + 2 \sec(x) + 2)}{4}$$

input `integrate(sec(x)**4*tan(x)/(sec(x)**4+4),x)`

output `log(sec(x)**2 - 2*sec(x) + 2)/4 + log(sec(x)**2 + 2*sec(x) + 2)/4`

3.228.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.82

$$\int \frac{\sec^4(x) \tan(x)}{4 + \sec^4(x)} dx = \frac{1}{4} \log(\sec(x)^4 + 4)$$

input `integrate(sec(x)^4*tan(x)/(sec(x)^4+4),x, algorithm="maxima")`

output `1/4*log(sec(x)^4 + 4)`

3.228.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. 2(9) = 18.

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.73

$$\int \frac{\sec^4(x) \tan(x)}{4 + \sec^4(x)} dx = \frac{1}{4} \log(4 \cos(x)^4 + 1) - \frac{1}{4} \log(\cos(x)^4)$$

input `integrate(sec(x)^4*tan(x)/(sec(x)^4+4),x, algorithm="giac")`

output `1/4*log(4*cos(x)^4 + 1) - 1/4*log(cos(x)^4)`

3.228.9 Mupad [B] (verification not implemented)

Time = 15.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.36

$$\int \frac{\sec^4(x) \tan(x)}{4 + \sec^4(x)} dx = \frac{\ln(\tan(x)^4 + 2 \tan(x)^2 + 5)}{4}$$

input `int(tan(x)/(cos(x)^4*(1/cos(x)^4 + 4)),x)`

output `log(2*tan(x)^2 + tan(x)^4 + 5)/4`

3.229 $\int x^{2x}(2 + 2 \log(x)) dx$

3.229.1 Optimal result	1168
3.229.2 Mathematica [A] (verified)	1168
3.229.3 Rubi [A] (verified)	1169
3.229.4 Maple [A] (verified)	1170
3.229.5 Fricas [A] (verification not implemented)	1170
3.229.6 Sympy [A] (verification not implemented)	1171
3.229.7 Maxima [A] (verification not implemented)	1171
3.229.8 Giac [A] (verification not implemented)	1171
3.229.9 Mupad [B] (verification not implemented)	1172

3.229.1 Optimal result

Integrand size = 12, antiderivative size = 5

$$\int x^{2x}(2 + 2 \log(x)) dx = x^{2x}$$

output `x^(2*x)`

3.229.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int x^{2x}(2 + 2 \log(x)) dx = x^{2x}$$

input `Integrate[x^(2*x)*(2 + 2*Log[x]),x]`

output `x^(2*x)`

3.229.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {7292, 27, 7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^{2x}(2\log(x) + 2) dx \\ & \quad \downarrow \text{7292} \\ & \int 2x^{2x}(\log(x) + 1) dx \\ & \quad \downarrow \text{27} \\ & 2 \int x^{2x}(\log(x) + 1) dx \\ & \quad \downarrow \text{7293} \\ & 2 \int (\log(x)x^{2x} + x^{2x}) dx \\ & \quad \downarrow \text{2009} \\ & x^{2x} \end{aligned}$$

input `Int[x^(2*x)*(2 + 2*Log[x]),x]`

output `x^(2*x)`

3.229.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7292 `Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v != u]`

```
rule 7293 Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

3.229.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
derivativedivides	x^{2x}	6
risch	x^{2x}	6
parallelrisch	x^{2x}	6
norman	$e^{2x \ln(x)}$	7

```
input int(x^(2*x)*(2*ln(x)+2),x,method=_RETURNVERBOSE)
```

```
output x^(2*x)
```

3.229.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int x^{2x}(2 + 2 \log(x)) dx = x^{2x}$$

```
input integrate(x^(2*x)*(2*log(x)+2),x, algorithm="fricas")
```

```
output x^(2*x)
```

3.229.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.60

$$\int x^{2x}(2 + 2\log(x)) dx = x^{2x}$$

input `integrate(x**(2*x)*(2*ln(x)+2),x)`output `x**(2*x)`**3.229.7 Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int x^{2x}(2 + 2\log(x)) dx = x^{2x}$$

input `integrate(x^(2*x)*(2*log(x)+2),x, algorithm="maxima")`output `x^(2*x)`**3.229.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int x^{2x}(2 + 2\log(x)) dx = x^{2x}$$

input `integrate(x^(2*x)*(2*log(x)+2),x, algorithm="giac")`output `x^(2*x)`

3.229.9 Mupad [B] (verification not implemented)

Time = 15.46 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int x^{2x}(2 + 2\log(x)) dx = x^{2x}$$

input `int(x^(2*x)*(2*log(x) + 2),x)`

output `x^(2*x)`

3.230 $\int \sqrt{1-x^2} dx$

3.230.1 Optimal result	1173
3.230.2 Mathematica [A] (verified)	1173
3.230.3 Rubi [A] (verified)	1174
3.230.4 Maple [A] (verified)	1175
3.230.5 Fricas [A] (verification not implemented)	1175
3.230.6 Sympy [A] (verification not implemented)	1176
3.230.7 Maxima [A] (verification not implemented)	1176
3.230.8 Giac [A] (verification not implemented)	1176
3.230.9 Mupad [B] (verification not implemented)	1177

3.230.1 Optimal result

Integrand size = 11, antiderivative size = 23

$$\int \sqrt{1-x^2} dx = \frac{1}{2}x\sqrt{1-x^2} + \frac{\arcsin(x)}{2}$$

output `1/2*x*(-x^2+1)^(1/2)+1/2*arcsin(x)`

3.230.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.61

$$\int \sqrt{1-x^2} dx = \frac{1}{2}x\sqrt{1-x^2} - \arctan\left(\frac{\sqrt{1-x^2}}{1+x}\right)$$

input `Integrate[Sqrt[1 - x^2],x]`

output `(x*Sqrt[1 - x^2])/2 - ArcTan[Sqrt[1 - x^2]/(1 + x)]`

3.230.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {211, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{1-x^2} dx$$

$$\downarrow \text{211}$$

$$\frac{1}{2} \int \frac{1}{\sqrt{1-x^2}} dx + \frac{1}{2} \sqrt{1-x^2} x$$

$$\downarrow \text{223}$$

$$\frac{\arcsin(x)}{2} + \frac{1}{2} \sqrt{1-x^2} x$$

input `Int[Sqrt[1 - x^2], x]`

output `(x*Sqrt[1 - x^2])/2 + ArcSin[x]/2`

3.230.3.1 Defintions of rubi rules used

rule 211 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[x*((a + b*x^2)^p/(2*p + 1)), x] + Simp[2*a*(p/(2*p + 1)) Int[(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b}, x] && GtQ[p, 0] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

3.230.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

method	result	size
default	$\frac{x\sqrt{-x^2+1}}{2} + \frac{\arcsin(x)}{2}$	18
risch	$-\frac{x(x^2-1)}{2\sqrt{-x^2+1}} + \frac{\arcsin(x)}{2}$	23
pseudoelliptic	$\frac{x\sqrt{-x^2+1}}{2} - \frac{\arctan\left(\frac{\sqrt{-x^2+1}}{x}\right)}{2}$	30
meijerg	$\frac{i(-2i\sqrt{\pi}x\sqrt{-x^2+1}-2i\sqrt{\pi}\arcsin(x))}{4\sqrt{\pi}}$	32
trager	$\frac{x\sqrt{-x^2+1}}{2} + \frac{\text{RootOf}(-Z^2+1)\ln(\text{RootOf}(-Z^2+1)\sqrt{-x^2+1}+x)}{2}$	41

input `int((-x^2+1)^(1/2),x,method=_RETURNVERBOSE)`output `1/2*x*(-x^2+1)^(1/2)+1/2*arcsin(x)`**3.230.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.35

$$\int \sqrt{1-x^2} dx = \frac{1}{2} \sqrt{-x^2+1}x - \arctan\left(\frac{\sqrt{-x^2+1}-1}{x}\right)$$

input `integrate((-x^2+1)^(1/2),x, algorithm="fricas")`output `1/2*sqrt(-x^2 + 1)*x - arctan((sqrt(-x^2 + 1) - 1)/x)`

3.230.6 Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int \sqrt{1-x^2} dx = \frac{x\sqrt{1-x^2}}{2} + \frac{\arcsin(x)}{2}$$

input `integrate((-x**2+1)**(1/2),x)`output `x*sqrt(1 - x**2)/2 + asin(x)/2`**3.230.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \sqrt{1-x^2} dx = \frac{1}{2} \sqrt{-x^2+1}x + \frac{1}{2} \arcsin(x)$$

input `integrate((-x^2+1)^(1/2),x, algorithm="maxima")`output `1/2*sqrt(-x^2 + 1)*x + 1/2*arcsin(x)`**3.230.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \sqrt{1-x^2} dx = \frac{1}{2} \sqrt{-x^2+1}x + \frac{1}{2} \arcsin(x)$$

input `integrate((-x^2+1)^(1/2),x, algorithm="giac")`output `1/2*sqrt(-x^2 + 1)*x + 1/2*arcsin(x)`

3.230.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \sqrt{1-x^2} dx = \frac{\arcsin(x)}{2} + \frac{x\sqrt{1-x^2}}{2}$$

input `int((1 - x^2)^(1/2),x)`

output `asin(x)/2 + (x*(1 - x^2)^(1/2))/2`

3.231 $\int e^{-x^4} x^5 dx$

3.231.1 Optimal result	1178
3.231.2 Mathematica [A] (verified)	1178
3.231.3 Rubi [A] (verified)	1179
3.231.4 Maple [A] (verified)	1180
3.231.5 Fricas [A] (verification not implemented)	1180
3.231.6 Sympy [A] (verification not implemented)	1181
3.231.7 Maxima [A] (verification not implemented)	1181
3.231.8 Giac [A] (verification not implemented)	1181
3.231.9 Mupad [B] (verification not implemented)	1182

3.231.1 Optimal result

Integrand size = 11, antiderivative size = 28

$$\int e^{-x^4} x^5 dx = -\frac{1}{4}e^{-x^4} x^2 + \frac{1}{8}\sqrt{\pi}\operatorname{erf}(x^2)$$

output `-1/4*x^2/exp(x^4)+1/8*Pi^(1/2)*erf(x^2)`

3.231.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int e^{-x^4} x^5 dx = -\frac{1}{4}e^{-x^4} x^2 + \frac{1}{8}\sqrt{\pi}\operatorname{erf}(x^2)$$

input `Integrate[x^5/E^x^4,x]`

output `-1/4*x^2/E^x^4 + (Sqrt[Pi]*Erf[x^2])/8`

3.231.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {2641, 2640, 2634}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^{-x^4} x^5 dx \\ & \quad \downarrow \text{2641} \\ & \frac{1}{2} \int e^{-x^4} x dx - \frac{1}{4} e^{-x^4} x^2 \\ & \quad \downarrow \text{2640} \\ & \frac{1}{4} \int e^{-x^4} dx^2 - \frac{1}{4} e^{-x^4} x^2 \\ & \quad \downarrow \text{2634} \\ & \frac{1}{8} \sqrt{\pi} \operatorname{erf}(x^2) - \frac{1}{4} e^{-x^4} x^2 \end{aligned}$$

input `Int[x^5/E^x^4,x]`

output `-1/4*x^2/E^x^4 + (Sqrt[Pi]*Erf[x^2])/8`

3.231.3.1 Defintions of rubi rules used

rule 2634 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(2)), x_Symbol] :> Simp[F^a*Sqrt[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && NegQ[b]`

rule 2640 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[1/(d*(m + 1)) Subst[Int[F^(a + b*x^2), x], x, (c + d*x)^(m + 1)], x] /; FreeQ[{F, a, b, c, d, m, n}, x] && EqQ[n, 2*(m + 1)]`

```
rule 2641 Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))^(n_))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m - n + 1)*(F^(a + b*(c + d*x)^n)/(b*d*n*Log[F])), x] - Simp[(m - n + 1)/(b*n*Log[F]) Int[(c + d*x)^(m - n)*F^(a + b*(c + d*x)^n), x], x] /; FreeQ[{F, a, b, c, d}, x] && IntegerQ[2*(m + 1)/n] && LtQ[0, (m + 1)/n, 5] && IntegerQ[n] && (LtQ[0, n, m + 1] || LtQ[m, n, 0])
```

3.231.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.79

method	result	size
meijerg	$-\frac{x^2 e^{-x^4}}{4} + \frac{\sqrt{\pi} \operatorname{erf}(x^2)}{8}$	22

```
input int(x^5*exp(-x^4),x,method=_RETURNVERBOSE)
```

```
output -1/4*x^2*exp(-x^4)+1/8*Pi^(1/2)*erf(x^2)
```

3.231.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.75

$$\int e^{-x^4} x^5 dx = -\frac{1}{4} x^2 e^{-x^4} + \frac{1}{8} \sqrt{\pi} \operatorname{erf}(x^2)$$

```
input integrate(x^5*exp(-x^4),x, algorithm="fricas")
```

```
output -1/4*x^2*e^(-x^4) + 1/8*sqrt(pi)*erf(x^2)
```

3.231.6 Sympy [A] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.71

$$\int e^{-x^4} x^5 dx = -\frac{x^2 e^{-x^4}}{4} + \frac{\sqrt{\pi} \operatorname{erf}(x^2)}{8}$$

input `integrate(x**5*exp(-x**4),x)`output `-x**2*exp(-x**4)/4 + sqrt(pi)*erf(x**2)/8`**3.231.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.75

$$\int e^{-x^4} x^5 dx = -\frac{1}{4} x^2 e^{-x^4} + \frac{1}{8} \sqrt{\pi} \operatorname{erf}(x^2)$$

input `integrate(x^5*exp(-x^4),x, algorithm="maxima")`output `-1/4*x^2*e^(-x^4) + 1/8*sqrt(pi)*erf(x^2)`**3.231.8 Giac [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.25

$$\int e^{-x^4} x^5 dx = -\frac{\sqrt{\pi} \left(\frac{2\sqrt{x^4} e^{-x^4}}{\sqrt{\pi}} - \operatorname{erf}(\sqrt{x^4}) \right) |x|}{8x}$$

input `integrate(x^5*exp(-x^4),x, algorithm="giac")`output `-1/8*sqrt(pi)*(2*sqrt(x^4)*e^(-x^4)/sqrt(pi) - erf(sqrt(x^4)))*abs(x)/x`

3.231.9 Mupad [B] (verification not implemented)

Time = 16.33 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.75

$$\int e^{-x^4} x^5 dx = \frac{\sqrt{\pi} \operatorname{erf}(x^2)}{8} - \frac{x^2 e^{-x^4}}{4}$$

input `int(x^5*exp(-x^4),x)`

output `(pi^(1/2)*erf(x^2))/8 - (x^2*exp(-x^4))/4`

$$\mathbf{3.232} \quad \int \frac{1+\cos(x)}{x+\sin(x)} dx$$

3.232.1 Optimal result	1183
3.232.2 Mathematica [A] (verified)	1183
3.232.3 Rubi [A] (verified)	1184
3.232.4 Maple [A] (verified)	1184
3.232.5 Fricas [A] (verification not implemented)	1185
3.232.6 Sympy [A] (verification not implemented)	1185
3.232.7 Maxima [A] (verification not implemented)	1185
3.232.8 Giac [B] (verification not implemented)	1186
3.232.9 Mupad [B] (verification not implemented)	1186

3.232.1 Optimal result

Integrand size = 11, antiderivative size = 5

$$\int \frac{1 + \cos(x)}{x + \sin(x)} dx = \log(x + \sin(x))$$

output `ln(x+sin(x))`

3.232.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1 + \cos(x)}{x + \sin(x)} dx = \log(x + \sin(x))$$

input `Integrate[(1 + Cos[x])/(x + Sin[x]), x]`

output `Log[x + Sin[x]]`

3.232.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {7235}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(x) + 1}{x + \sin(x)} dx$$

↓ 7235

$$\log(x + \sin(x))$$

input `Int[(1 + Cos[x])/(x + Sin[x]),x]`

output `Log[x + Sin[x]]`

3.232.3.1 Defintions of rubi rules used

rule 7235 `Int[(u_)/(y_), x_Symbol] := With[{q = DerivativeDivides[y, u, x]}, Simp[q*Log[RemoveContent[y, x]], x] /; !FalseQ[q]]`

3.232.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
derivativedivides	$\ln(x + \sin(x))$	6
default	$\ln(x + \sin(x))$	6
risch	$-ix + \ln(e^{2ix} + 2ix e^{ix} - 1)$	23
parallelrisc	$-\ln\left(\frac{1}{1+\cos(x)}\right) + \ln\left(\frac{x+\sin(x)}{1+\cos(x)}\right)$	23
norman	$-\ln\left(1 + \tan\left(\frac{x}{2}\right)^2\right) + \ln\left(x \tan\left(\frac{x}{2}\right)^2 + x + 2 \tan\left(\frac{x}{2}\right)\right)$	30

input `int((1+cos(x))/(x+sin(x)),x,method=_RETURNVERBOSE)`

output `ln(x+sin(x))`

3.232.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1 + \cos(x)}{x + \sin(x)} dx = \log(x + \sin(x))$$

input `integrate((1+cos(x))/(x+sin(x)),x, algorithm="fricas")`

output `log(x + sin(x))`

3.232.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1 + \cos(x)}{x + \sin(x)} dx = \log(x + \sin(x))$$

input `integrate((1+cos(x))/(x+sin(x)),x)`

output `log(x + sin(x))`

3.232.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1 + \cos(x)}{x + \sin(x)} dx = \log(x + \sin(x))$$

input `integrate((1+cos(x))/(x+sin(x)),x, algorithm="maxima")`

output `log(x + sin(x))`

3.232.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 72 vs. $2(5) = 10$.

Time = 0.28 (sec) , antiderivative size = 72, normalized size of antiderivative = 14.40

$$\int \frac{1 + \cos(x)}{x + \sin(x)} dx$$

$$= \frac{1}{2} \log \left(\frac{4 \left(x^2 \tan \left(\frac{1}{2} x \right)^4 + 2 x^2 \tan \left(\frac{1}{2} x \right)^2 + 4 x \tan \left(\frac{1}{2} x \right)^3 + x^2 + 4 x \tan \left(\frac{1}{2} x \right) + 4 \tan \left(\frac{1}{2} x \right)^2 \right)}{\tan \left(\frac{1}{2} x \right)^4 + 2 \tan \left(\frac{1}{2} x \right)^2 + 1} \right)$$

input `integrate((1+cos(x))/(x+sin(x)),x, algorithm="giac")`

output `1/2*log(4*(x^2*tan(1/2*x)^4 + 2*x^2*tan(1/2*x)^2 + 4*x*tan(1/2*x)^3 + x^2 + 4*x*tan(1/2*x) + 4*tan(1/2*x)^2)/(tan(1/2*x)^4 + 2*tan(1/2*x)^2 + 1))`

3.232.9 Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1 + \cos(x)}{x + \sin(x)} dx = \ln(x + \sin(x))$$

input `int((cos(x) + 1)/(x + sin(x)),x)`

output `log(x + sin(x))`

3.233 $\int \frac{\cot^{-1}(x)+\arctan(x)}{x} dx$

3.233.1 Optimal result	1187
3.233.2 Mathematica [A] (verified)	1187
3.233.3 Rubi [A] (verified)	1188
3.233.4 Maple [A] (verified)	1189
3.233.5 Fricas [A] (verification not implemented)	1189
3.233.6 Sympy [F]	1189
3.233.7 Maxima [A] (verification not implemented)	1190
3.233.8 Giac [A] (verification not implemented)	1190
3.233.9 Mupad [F(-1)]	1190

3.233.1 Optimal result

Integrand size = 9, antiderivative size = 57

$$\int \frac{\cot^{-1}(x) + \arctan(x)}{x} dx = -\frac{1}{2}i \operatorname{PolyLog}\left(2, -\frac{i}{x}\right) + \frac{1}{2}i \operatorname{PolyLog}\left(2, \frac{i}{x}\right) + \frac{1}{2}i \operatorname{PolyLog}(2, -ix) - \frac{1}{2}i \operatorname{PolyLog}(2, ix)$$

output `-1/2*I*polylog(2,-I/x)+1/2*I*polylog(2,I/x)+1/2*I*polylog(2,-I*x)-1/2*I*polylog(2,I*x)`

3.233.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00

$$\int \frac{\cot^{-1}(x) + \arctan(x)}{x} dx = -\frac{1}{2}i \operatorname{PolyLog}\left(2, -\frac{i}{x}\right) + \frac{1}{2}i \operatorname{PolyLog}\left(2, \frac{i}{x}\right) + \frac{1}{2}i \operatorname{PolyLog}(2, -ix) - \frac{1}{2}i \operatorname{PolyLog}(2, ix)$$

input `Integrate[(ArcCot[x] + ArcTan[x])/x,x]`

output `(-1/2*I)*PolyLog[2, (-I)/x] + (I/2)*PolyLog[2, I/x] + (I/2)*PolyLog[2, (-I)*x] - (I/2)*PolyLog[2, I*x]`

3.233.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {2010, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(x) + \cot^{-1}(x)}{x} dx$$

↓ 2010

$$\int \left(\frac{\arctan(x)}{x} + \frac{\cot^{-1}(x)}{x} \right) dx$$

↓ 2009

$$-\frac{1}{2}i \operatorname{PolyLog}\left(2, -\frac{i}{x}\right) + \frac{1}{2}i \operatorname{PolyLog}\left(2, \frac{i}{x}\right) + \frac{1}{2}i \operatorname{PolyLog}(2, -ix) - \frac{1}{2}i \operatorname{PolyLog}(2, ix)$$

input `Int[(ArcCot[x] + ArcTan[x])/x,x]`

output `(-1/2*I)*PolyLog[2, (-I)/x] + (I/2)*PolyLog[2, I/x] + (I/2)*PolyLog[2, (-I)*x] - (I/2)*PolyLog[2, I*x]`

3.233.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2010 `Int[(u_)*((c_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]`

3.233.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.21

method	result	size
default	$\ln(x) \operatorname{arccot}(x) + \ln(x) \operatorname{arctan}(x)$	12
parts	$\ln(x) \operatorname{arccot}(x) + \ln(x) \operatorname{arctan}(x)$	12

input `int((arctan(x)+arccot(x))/x,x,method=_RETURNVERBOSE)`output `ln(x)*arccot(x)+ln(x)*arctan(x)`**3.233.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.09

$$\int \frac{\cot^{-1}(x) + \operatorname{arctan}(x)}{x} dx = -\frac{1}{2} \pi \log(x)$$

input `integrate((arctan(x)+arccot(x))/x,x, algorithm="fricas")`output `-1/2*pi*log(x)`**3.233.6 Sympy [F]**

$$\int \frac{\cot^{-1}(x) + \operatorname{arctan}(x)}{x} dx = \int \frac{\operatorname{acot}(x) + \operatorname{atan}(x)}{x} dx$$

input `integrate((atan(x)+acot(x))/x,x)`output `Integral((acot(x) + atan(x))/x, x)`

3.233.7 Maxima [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.16

$$\int \frac{\cot^{-1}(x) + \arctan(x)}{x} dx = (\arctan(x) + \arctan(1, x)) \log(x)$$

input `integrate((arctan(x)+arccot(x))/x,x, algorithm="maxima")`output `(arctan(x) + arctan2(1, x))*log(x)`**3.233.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.09

$$\int \frac{\cot^{-1}(x) + \arctan(x)}{x} dx = -\frac{1}{2} \pi \log(x)$$

input `integrate((arctan(x)+arccot(x))/x,x, algorithm="giac")`output `-1/2*pi*log(x)`**3.233.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\cot^{-1}(x) + \arctan(x)}{x} dx = \int \frac{\operatorname{atan}(x) + \operatorname{acot}(x)}{x} dx$$

input `int((atan(x) + acot(x))/x,x)`output `int((atan(x) + acot(x))/x, x)`

3.234 $\int \frac{\sinh(x)}{\cosh(x) - \sinh(x)} dx$

3.234.1 Optimal result	1191
3.234.2 Mathematica [A] (verified)	1191
3.234.3 Rubi [C] (verified)	1192
3.234.4 Maple [A] (verified)	1193
3.234.5 Fricas [B] (verification not implemented)	1194
3.234.6 Sympy [B] (verification not implemented)	1194
3.234.7 Maxima [A] (verification not implemented)	1194
3.234.8 Giac [A] (verification not implemented)	1195
3.234.9 Mupad [B] (verification not implemented)	1195

3.234.1 Optimal result

Integrand size = 12, antiderivative size = 14

$$\int \frac{\sinh(x)}{\cosh(x) - \sinh(x)} dx = \frac{1}{2}(-x + e^x \sinh(x))$$

output `-1/2*x+1/2*exp(x)*sinh(x)`

3.234.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.57

$$\int \frac{\sinh(x)}{\cosh(x) - \sinh(x)} dx = -\frac{x}{2} + \frac{\cosh^2(x)}{2} + \frac{1}{4} \sinh(2x)$$

input `Integrate[Sinh[x]/(Cosh[x] - Sinh[x]),x]`

output `-1/2*x + Cosh[x]^2/2 + Sinh[2*x]/4`

3.234.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 29, normalized size of antiderivative = 2.07, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 26, 3560, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sinh(x)}{\cosh(x) - \sinh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \sin(ix)}{i \sin(ix) + \cos(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\sin(ix)}{\cos(ix) + i \sin(ix)} dx \\
 & \quad \downarrow \text{3560} \\
 & -i \left(\frac{i \sinh(x)}{2(\cosh(x) - \sinh(x))} - \frac{i \int 1 dx}{2} \right) \\
 & \quad \downarrow \text{24} \\
 & -i \left(\frac{i \sinh(x)}{2(\cosh(x) - \sinh(x))} - \frac{ix}{2} \right)
 \end{aligned}$$

input `Int[Sinh[x]/(Cosh[x] - Sinh[x]),x]`

output `(-I)*((-1/2*I)*x + ((I/2)*Sinh[x])/(Cosh[x] - Sinh[x]))`

3.234.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3560 `Int[sin[(c_.) + (d_.)*(x_)]^(m_.)*(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a*cos[c + d*x] + b*sin[c + d*x])^n/(2*b*d*n*sin[c + d*x]^n)), x] + Simp[1/(2*b) Int[(a*cos[c + d*x] + b*sin[c + d*x])^(n + 1)/sin[c + d*x]^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[m + n, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

3.234.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

method	result	size
risch	$-\frac{x}{2} + \frac{e^{2x}}{4}$	11
parallelrisch	$\frac{-\tanh(x)x+x-1}{2\tanh(x)-2}$	18
default	$-\frac{\ln(\tanh(\frac{x}{2})+1)}{2} + \frac{1}{(\tanh(\frac{x}{2})-1)^2} + \frac{1}{\tanh(\frac{x}{2})-1} + \frac{\ln(\tanh(\frac{x}{2})-1)}{2}$	36

input `int(sinh(x)/(cosh(x)-sinh(x)),x,method=_RETURNVERBOSE)`

output `-1/2*x+1/4*exp(2*x)`

3.234.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(10) = 20$.

Time = 0.24 (sec) , antiderivative size = 29, normalized size of antiderivative = 2.07

$$\int \frac{\sinh(x)}{\cosh(x) - \sinh(x)} dx = -\frac{(2x - 1)\cosh(x) - (2x + 1)\sinh(x)}{4(\cosh(x) - \sinh(x))}$$

input `integrate(sinh(x)/(cosh(x)-sinh(x)),x, algorithm="fricas")`

output `-1/4*((2*x - 1)*cosh(x) - (2*x + 1)*sinh(x))/(cosh(x) - sinh(x))`

3.234.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. $2(10) = 20$.

Time = 0.27 (sec) , antiderivative size = 42, normalized size of antiderivative = 3.00

$$\int \frac{\sinh(x)}{\cosh(x) - \sinh(x)} dx = \frac{x \sinh(x)}{-2 \sinh(x) + 2 \cosh(x)} - \frac{x \cosh(x)}{-2 \sinh(x) + 2 \cosh(x)} + \frac{\cosh(x)}{-2 \sinh(x) + 2 \cosh(x)}$$

input `integrate(sinh(x)/(cosh(x)-sinh(x)),x)`

output `x*sinh(x)/(-2*sinh(x) + 2*cosh(x)) - x*cosh(x)/(-2*sinh(x) + 2*cosh(x)) + cosh(x)/(-2*sinh(x) + 2*cosh(x))`

3.234.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{\sinh(x)}{\cosh(x) - \sinh(x)} dx = -\frac{1}{2}x + \frac{1}{4}e^{(2x)}$$

input `integrate(sinh(x)/(cosh(x)-sinh(x)),x, algorithm="maxima")`

output `-1/2*x + 1/4*e^(2*x)`

3.234.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{\sinh(x)}{\cosh(x) - \sinh(x)} dx = -\frac{1}{2}x + \frac{1}{4}e^{(2x)}$$

input `integrate(sinh(x)/(cosh(x)-sinh(x)),x, algorithm="giac")`output `-1/2*x + 1/4*e^(2*x)`**3.234.9 Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{\sinh(x)}{\cosh(x) - \sinh(x)} dx = \frac{e^{2x}}{4} - \frac{x}{2}$$

input `int(sinh(x)/(cosh(x) - sinh(x)),x)`output `exp(2*x)/4 - x/2`

3.235 $\int \frac{x}{\sqrt{-1+x}+\sqrt{1+x}} dx$

3.235.1 Optimal result	1196
3.235.2 Mathematica [A] (verified)	1196
3.235.3 Rubi [A] (verified)	1197
3.235.4 Maple [A] (verified)	1198
3.235.5 Fricas [A] (verification not implemented)	1198
3.235.6 Sympy [F]	1198
3.235.7 Maxima [F]	1199
3.235.8 Giac [A] (verification not implemented)	1199
3.235.9 Mupad [B] (verification not implemented)	1199

3.235.1 Optimal result

Integrand size = 19, antiderivative size = 45

$$\int \frac{x}{\sqrt{-1+x}+\sqrt{1+x}} dx = -\frac{1}{3}(-1+x)^{3/2} - \frac{1}{5}(-1+x)^{5/2} - \frac{1}{3}(1+x)^{3/2} + \frac{1}{5}(1+x)^{5/2}$$

output `-1/3*(-1+x)^(3/2)-1/5*(-1+x)^(5/2)-1/3*(1+x)^(3/2)+1/5*(1+x)^(5/2)`

3.235.2 Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.76

$$\int \frac{x}{\sqrt{-1+x}+\sqrt{1+x}} dx = \frac{1}{15}((1+x)^{3/2}(-2+3x) + \sqrt{-1+x}(2+x-3x^2))$$

input `Integrate[x/(Sqrt[-1 + x] + Sqrt[1 + x]),x]`

output `((1 + x)^(3/2)*(-2 + 3*x) + Sqrt[-1 + x]*(2 + x - 3*x^2))/15`

3.235.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.22, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2529, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x}{\sqrt{x-1} + \sqrt{x+1}} dx \\ & \quad \downarrow \text{2529} \\ & \frac{1}{2} \int x\sqrt{x+1} dx - \frac{1}{2} \int \sqrt{x-1} x dx \\ & \quad \downarrow \text{53} \\ & \frac{1}{2} \int \left((x+1)^{3/2} - \sqrt{x+1} \right) dx - \frac{1}{2} \int \left((x-1)^{3/2} + \sqrt{x-1} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \left(-\frac{2}{5}(x-1)^{5/2} - \frac{2}{3}(x-1)^{3/2} \right) + \frac{1}{2} \left(\frac{2}{5}(x+1)^{5/2} - \frac{2}{3}(x+1)^{3/2} \right) \end{aligned}$$

input `Int[x/(Sqrt[-1 + x] + Sqrt[1 + x]), x]`

output `((-2*(-1 + x)^(3/2))/3 - (2*(-1 + x)^(5/2))/5)/2 + ((-2*(1 + x)^(3/2))/3 + (2*(1 + x)^(5/2))/5)/2`

3.235.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 2529 Int[(u_)/((e_)*Sqrt[(a_.) + (b_.)*(x_)] + (f_.)*Sqrt[(c_.) + (d_.)*(x_)]),
  x_Symbol] :> Simp[-d/(e*(b*c - a*d)) Int[u*Sqrt[a + b*x], x], x] + Simp[
  b/(f*(b*c - a*d)) Int[u*Sqrt[c + d*x], x], x] /; FreeQ[{a, b, c, d, e, f}
  , x] && NeQ[b*c - a*d, 0] && EqQ[b*e^2 - d*f^2, 0]
```

3.235.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.67

method	result	size
default	$-\frac{(-1+x)^{\frac{3}{2}}}{3} - \frac{(-1+x)^{\frac{5}{2}}}{5} - \frac{(1+x)^{\frac{3}{2}}}{3} + \frac{(1+x)^{\frac{5}{2}}}{5}$	30

```
input int(x/((-1+x)^(1/2)+(1+x)^(1/2)),x,method=_RETURNVERBOSE)
```

```
output -1/3*(-1+x)^(3/2)-1/5*(-1+x)^(5/2)-1/3*(1+x)^(3/2)+1/5*(1+x)^(5/2)
```

3.235.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.73

$$\int \frac{x}{\sqrt{-1+x} + \sqrt{1+x}} dx = \frac{1}{15} (3x^2 + x - 2)\sqrt{x+1} - \frac{1}{15} (3x^2 - x - 2)\sqrt{x-1}$$

```
input integrate(x/((-1+x)^(1/2)+(1+x)^(1/2)),x, algorithm="fracas")
```

```
output 1/15*(3*x^2 + x - 2)*sqrt(x + 1) - 1/15*(3*x^2 - x - 2)*sqrt(x - 1)
```

3.235.6 Sympy [F]

$$\int \frac{x}{\sqrt{-1+x} + \sqrt{1+x}} dx = \int \frac{x}{\sqrt{x-1} + \sqrt{x+1}} dx$$

```
input integrate(x/((-1+x)**(1/2)+(1+x)**(1/2)),x)
```

```
output Integral(x/(sqrt(x - 1) + sqrt(x + 1)), x)
```

3.235.7 Maxima [F]

$$\int \frac{x}{\sqrt{-1+x} + \sqrt{1+x}} dx = \int \frac{x}{\sqrt{x+1} + \sqrt{x-1}} dx$$

input `integrate(x/((-1+x)^(1/2)+(1+x)^(1/2)),x, algorithm="maxima")`

output `integrate(x/(sqrt(x + 1) + sqrt(x - 1)), x)`

3.235.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.73

$$\int \frac{x}{\sqrt{-1+x} + \sqrt{1+x}} dx = \frac{1}{5} (x+1)^{\frac{5}{2}} - \frac{1}{3} (x+1)^{\frac{3}{2}} - \frac{1}{15} ((3x-4)(x+1) + 2)\sqrt{x-1}$$

input `integrate(x/((-1+x)^(1/2)+(1+x)^(1/2)),x, algorithm="giac")`

output `1/5*(x + 1)^(5/2) - 1/3*(x + 1)^(3/2) - 1/15*((3*x - 4)*(x + 1) + 2)*sqrt(x - 1)`

3.235.9 Mupad [B] (verification not implemented)

Time = 15.00 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.13

$$\int \frac{x}{\sqrt{-1+x} + \sqrt{1+x}} dx = \frac{x\sqrt{x-1}}{15} + \frac{x\sqrt{x+1}}{15} + \frac{2\sqrt{x-1}}{15} - \frac{2\sqrt{x+1}}{15} - \frac{x^2\sqrt{x-1}}{5} + \frac{x^2\sqrt{x+1}}{5}$$

input `int(x/((x - 1)^(1/2) + (x + 1)^(1/2)),x)`

output `(x*(x - 1)^(1/2))/15 + (x*(x + 1)^(1/2))/15 + (2*(x - 1)^(1/2))/15 - (2*(x + 1)^(1/2))/15 - (x^2*(x - 1)^(1/2))/5 + (x^2*(x + 1)^(1/2))/5`

3.236 $\int \cos(x + \cos(x)) dx$

3.236.1 Optimal result	1200
3.236.2 Mathematica [N/A]	1200
3.236.3 Rubi [N/A]	1201
3.236.4 Maple [N/A] (verified)	1201
3.236.5 Fricas [N/A]	1202
3.236.6 Sympy [N/A]	1202
3.236.7 Maxima [N/A]	1202
3.236.8 Giac [N/A]	1203
3.236.9 Mupad [N/A]	1203

3.236.1 Optimal result

Integrand size = 5, antiderivative size = 5

$$\int \cos(x + \cos(x)) dx = \text{Int}(\cos(x + \cos(x)), x)$$

output `CannotIntegrate(cos(x+cos(x)),x)`

3.236.2 Mathematica [N/A]

Not integrable

Time = 1.61 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.40

$$\int \cos(x + \cos(x)) dx = \int \cos(x + \cos(x)) dx$$

input `Integrate[Cos[x + Cos[x]],x]`

output `Integrate[Cos[x + Cos[x]], x]`

3.236.3 Rubi [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {7299}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(x + \cos(x)) dx$$

↓ 7299

$$\int \cos(x + \cos(x)) dx$$

input `Int[Cos[x + Cos[x]], x]`

output `$Aborted`

3.236.3.1 Defintions of rubi rules used

rule 7299 `Int[u_, x_] :> CannotIntegrate[u, x]`

3.236.4 Maple [N/A] (verified)

Not integrable

Time = 0.06 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \cos(x + \cos(x)) dx$$

input `int(cos(x+cos(x)), x)`

output `int(cos(x+cos(x)), x)`

3.236.5 Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.40

$$\int \cos(x + \cos(x)) dx = \int \cos(x + \cos(x)) dx$$

input `integrate(cos(x+cos(x)),x, algorithm="fricas")`output `integral(cos(x + cos(x)), x)`**3.236.6 Sympy [N/A]**

Not integrable

Time = 0.59 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.40

$$\int \cos(x + \cos(x)) dx = \int \cos(x + \cos(x)) dx$$

input `integrate(cos(x+cos(x)),x)`output `Integral(cos(x + cos(x)), x)`**3.236.7 Maxima [N/A]**

Not integrable

Time = 0.41 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.40

$$\int \cos(x + \cos(x)) dx = \int \cos(x + \cos(x)) dx$$

input `integrate(cos(x+cos(x)),x, algorithm="maxima")`output `integrate(cos(x + cos(x)), x)`

3.236.8 Giac [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.40

$$\int \cos(x + \cos(x)) dx = \int \cos(x + \cos(x)) dx$$

input `integrate(cos(x+cos(x)),x, algorithm="giac")`output `integrate(cos(x + cos(x)), x)`**3.236.9 Mupad [N/A]**

Not integrable

Time = 15.89 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.40

$$\int \cos(x + \cos(x)) dx = \int \cos(x + \cos(x)) dx$$

input `int(cos(x + cos(x)),x)`output `int(cos(x + cos(x)), x)`

3.237 $\int x^3 \sin(x^2) dx$

3.237.1 Optimal result	1204
3.237.2 Mathematica [A] (verified)	1204
3.237.3 Rubi [A] (verified)	1205
3.237.4 Maple [A] (verified)	1206
3.237.5 Fricas [A] (verification not implemented)	1207
3.237.6 Sympy [A] (verification not implemented)	1207
3.237.7 Maxima [A] (verification not implemented)	1207
3.237.8 Giac [A] (verification not implemented)	1208
3.237.9 Mupad [B] (verification not implemented)	1208

3.237.1 Optimal result

Integrand size = 8, antiderivative size = 20

$$\int x^3 \sin(x^2) dx = -\frac{1}{2}x^2 \cos(x^2) + \frac{\sin(x^2)}{2}$$

output `-1/2*x^2*cos(x^2)+1/2*sin(x^2)`

3.237.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int x^3 \sin(x^2) dx = -\frac{1}{2}x^2 \cos(x^2) + \frac{\sin(x^2)}{2}$$

input `Integrate[x^3*Sin[x^2],x]`

output `-1/2*(x^2*Cos[x^2]) + Sin[x^2]/2`

3.237.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3860, 3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \sin(x^2) dx \\
 & \quad \downarrow \text{3860} \\
 & \frac{1}{2} \int x^2 \sin(x^2) dx^2 \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int x^2 \sin(x^2) dx^2 \\
 & \quad \downarrow \text{3777} \\
 & \frac{1}{2} \left(\int \cos(x^2) dx^2 - x^2 \cos(x^2) \right) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \left(\int \sin\left(x^2 + \frac{\pi}{2}\right) dx^2 - x^2 \cos(x^2) \right) \\
 & \quad \downarrow \text{3117} \\
 & \frac{1}{2} (\sin(x^2) - x^2 \cos(x^2))
 \end{aligned}$$

input `Int[x^3*Sin[x^2],x]`

output `(-x^2*Cos[x^2]) + Sin[x^2])/2`

3.237.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3860 `Int[(x_)^(m_.)*((a_.) + (b_.)*Sin[(c_.) + (d_.)*(x_)^(n_)])^(p_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Sin[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))`

3.237.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

method	result	size
derivativedivides	$-\frac{x^2 \cos(x^2)}{2} + \frac{\sin(x^2)}{2}$	17
default	$-\frac{x^2 \cos(x^2)}{2} + \frac{\sin(x^2)}{2}$	17
risch	$-\frac{x^2 \cos(x^2)}{2} + \frac{\sin(x^2)}{2}$	17
parallelrisch	$-\frac{x^2 \cos(x^2)}{2} + \frac{\sin(x^2)}{2}$	17
meijerg	$\sqrt{\pi} \left(-\frac{x^2 \cos(x^2)}{2\sqrt{\pi}} + \frac{\sin(x^2)}{2\sqrt{\pi}} \right)$	27
norman	$\frac{-\frac{x^2}{2} + \frac{x^2 \tan\left(\frac{x^2}{2}\right)^2}{2} + \tan\left(\frac{x^2}{2}\right)}{1 + \tan\left(\frac{x^2}{2}\right)^2}$	39
parts	$\frac{\sqrt{2}\sqrt{\pi} \operatorname{FresnelS}\left(\frac{\sqrt{2}x}{\sqrt{\pi}}\right)x^3}{2} - \frac{3\pi^2 \left(\frac{2 \operatorname{FresnelS}\left(\frac{\sqrt{2}x}{\sqrt{\pi}}\right)\sqrt{2}x^3}{3\pi^{\frac{3}{2}}} + \frac{2x^2 \cos(x^2)}{3\pi^2} - \frac{2 \sin(x^2)}{3\pi^2} \right)}{4}$	69

input `int(x^3*sin(x^2),x,method=_RETURNVERBOSE)`

output `-1/2*x^2*cos(x^2)+1/2*sin(x^2)`

3.237.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int x^3 \sin(x^2) dx = -\frac{1}{2} x^2 \cos(x^2) + \frac{1}{2} \sin(x^2)$$

input `integrate(x^3*sin(x^2),x, algorithm="fricas")`

output `-1/2*x^2*cos(x^2) + 1/2*sin(x^2)`

3.237.6 Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int x^3 \sin(x^2) dx = -\frac{x^2 \cos(x^2)}{2} + \frac{\sin(x^2)}{2}$$

input `integrate(x**3*sin(x**2),x)`

output `-x**2*cos(x**2)/2 + sin(x**2)/2`

3.237.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int x^3 \sin(x^2) dx = -\frac{1}{2} x^2 \cos(x^2) + \frac{1}{2} \sin(x^2)$$

input `integrate(x^3*sin(x^2),x, algorithm="maxima")`

output `-1/2*x^2*cos(x^2) + 1/2*sin(x^2)`

3.237.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int x^3 \sin(x^2) dx = -\frac{1}{2} x^2 \cos(x^2) + \frac{1}{2} \sin(x^2)$$

input `integrate(x^3*sin(x^2),x, algorithm="giac")`

output `-1/2*x^2*cos(x^2) + 1/2*sin(x^2)`

3.237.9 Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int x^3 \sin(x^2) dx = \frac{\sin(x^2)}{2} - \frac{x^2 \cos(x^2)}{2}$$

input `int(x^3*sin(x^2),x)`

output `sin(x^2)/2 - (x^2*cos(x^2))/2`

3.238 $\int \frac{x}{1-x^4} dx$

3.238.1 Optimal result	1209
3.238.2 Mathematica [B] (verified)	1209
3.238.3 Rubi [A] (verified)	1210
3.238.4 Maple [A] (verified)	1211
3.238.5 Fricas [B] (verification not implemented)	1211
3.238.6 Sympy [B] (verification not implemented)	1211
3.238.7 Maxima [B] (verification not implemented)	1212
3.238.8 Giac [B] (verification not implemented)	1212
3.238.9 Mupad [B] (verification not implemented)	1213

3.238.1 Optimal result

Integrand size = 11, antiderivative size = 8

$$\int \frac{x}{1-x^4} dx = \frac{\operatorname{arctanh}(x^2)}{2}$$

output `1/2*arctanh(x^2)`

3.238.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 23 vs. $2(8) = 16$.

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 2.88

$$\int \frac{x}{1-x^4} dx = -\frac{1}{4} \log(1-x^2) + \frac{1}{4} \log(1+x^2)$$

input `Integrate[x/(1 - x^4), x]`

output `-1/4*Log[1 - x^2] + Log[1 + x^2]/4`

3.238.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {807, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{1-x^4} dx$$

↓ 807

$$\frac{1}{2} \int \frac{1}{1-x^4} dx^2$$

↓ 219

$$\frac{\operatorname{arctanh}(x^2)}{2}$$

input `Int[x/(1 - x^4), x]`

output `ArcTanh[x^2]/2`

3.238.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

3.238.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
meijerg	$\frac{\operatorname{arctanh}(x^2)}{2}$	7
risch	$\frac{\ln(x^2+1)}{4} - \frac{\ln(x^2-1)}{4}$	18
default	$-\frac{\ln(-1+x)}{4} - \frac{\ln(1+x)}{4} + \frac{\ln(x^2+1)}{4}$	22
norman	$-\frac{\ln(-1+x)}{4} - \frac{\ln(1+x)}{4} + \frac{\ln(x^2+1)}{4}$	22
parallelrisch	$-\frac{\ln(-1+x)}{4} - \frac{\ln(1+x)}{4} + \frac{\ln(x^2+1)}{4}$	22

input `int(x/(-x^4+1),x,method=_RETURNVERBOSE)`

output `1/2*arctanh(x^2)`

3.238.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. 2(6) = 12.

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.12

$$\int \frac{x}{1-x^4} dx = \frac{1}{4} \log(x^2+1) - \frac{1}{4} \log(x^2-1)$$

input `integrate(x/(-x^4+1),x, algorithm="fricas")`

output `1/4*log(x^2 + 1) - 1/4*log(x^2 - 1)`

3.238.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. 2(5) = 10.

Time = 0.05 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.88

$$\int \frac{x}{1-x^4} dx = -\frac{\log(x^2-1)}{4} + \frac{\log(x^2+1)}{4}$$

input `integrate(x/(-x**4+1),x)`

output `-log(x**2 - 1)/4 + log(x**2 + 1)/4`

3.238.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(6) = 12$.

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.12

$$\int \frac{x}{1-x^4} dx = \frac{1}{4} \log(x^2 + 1) - \frac{1}{4} \log(x^2 - 1)$$

input `integrate(x/(-x^4+1),x, algorithm="maxima")`

output `1/4*log(x^2 + 1) - 1/4*log(x^2 - 1)`

3.238.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18 vs. $2(6) = 12$.

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 2.25

$$\int \frac{x}{1-x^4} dx = \frac{1}{4} \log(x^2 + 1) - \frac{1}{4} \log(|x^2 - 1|)$$

input `integrate(x/(-x^4+1),x, algorithm="giac")`

output `1/4*log(x^2 + 1) - 1/4*log(abs(x^2 - 1))`

3.238.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{x}{1-x^4} dx = \frac{\operatorname{atanh}(x^2)}{2}$$

input `int(-x/(x^4 - 1),x)`

output `atanh(x^2)/2`

3.239 $\int \operatorname{sech}^2(x) dx$

3.239.1 Optimal result	1214
3.239.2 Mathematica [A] (verified)	1214
3.239.3 Rubi [A] (verified)	1215
3.239.4 Maple [A] (verified)	1216
3.239.5 Fricas [B] (verification not implemented)	1216
3.239.6 Sympy [B] (verification not implemented)	1216
3.239.7 Maxima [B] (verification not implemented)	1217
3.239.8 Giac [B] (verification not implemented)	1217
3.239.9 Mupad [B] (verification not implemented)	1217

3.239.1 Optimal result

Integrand size = 4, antiderivative size = 2

$$\int \operatorname{sech}^2(x) dx = \tanh(x)$$

output `tanh(x)`

3.239.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \operatorname{sech}^2(x) dx = \tanh(x)$$

input `Integrate[Sech[x]^2,x]`

output `Tanh[x]`

3.239.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \operatorname{sech}^2(x) dx \\ \downarrow 3042 \\ \int \csc\left(\frac{\pi}{2} + ix\right)^2 dx \\ \downarrow 4254 \\ i \int 1d(-i \tanh(x)) \\ \downarrow 24 \\ \tanh(x) \end{array}$$

input `Int[Sech[x]^2,x]`

output `Tanh[x]`

3.239.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

3.239.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\tanh(x)$	3
parallelrisch	$\tanh(x)$	3
risch	$-\frac{2}{e^{2x}+1}$	11

input `int(1/cosh(x)^2,x,method=_RETURNVERBOSE)`

output `tanh(x)`

3.239.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. $2(2) = 4$.

Time = 0.25 (sec) , antiderivative size = 20, normalized size of antiderivative = 10.00

$$\int \operatorname{sech}^2(x) dx = -\frac{2}{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1}$$

input `integrate(1/cosh(x)^2,x, algorithm="fricas")`

output `-2/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)`

3.239.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 14 vs. $2(2) = 4$.

Time = 0.31 (sec) , antiderivative size = 14, normalized size of antiderivative = 7.00

$$\int \operatorname{sech}^2(x) dx = \frac{2 \tanh\left(\frac{x}{2}\right)}{\tanh^2\left(\frac{x}{2}\right) + 1}$$

input `integrate(1/cosh(x)**2,x)`

output `2*tanh(x/2)/(tanh(x/2)**2 + 1)`

3.239.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10 vs. $2(2) = 4$.

Time = 0.20 (sec) , antiderivative size = 10, normalized size of antiderivative = 5.00

$$\int \operatorname{sech}^2(x) dx = \frac{2}{e^{(-2x)} + 1}$$

input `integrate(1/cosh(x)^2,x, algorithm="maxima")`

output `2/(e^(-2*x) + 1)`

3.239.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10 vs. $2(2) = 4$.

Time = 0.29 (sec) , antiderivative size = 10, normalized size of antiderivative = 5.00

$$\int \operatorname{sech}^2(x) dx = -\frac{2}{e^{(2x)} + 1}$$

input `integrate(1/cosh(x)^2,x, algorithm="giac")`

output `-2/(e^(2*x) + 1)`

3.239.9 Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \operatorname{sech}^2(x) dx = \tanh(x)$$

input `int(1/cosh(x)^2,x)`

output `tanh(x)`

3.240 $\int (e^{e^x} - e^{e^x-x}) dx$

3.240.1 Optimal result	1218
3.240.2 Mathematica [A] (verified)	1218
3.240.3 Rubi [A] (verified)	1219
3.240.4 Maple [A] (verified)	1219
3.240.5 Fricas [A] (verification not implemented)	1220
3.240.6 Sympy [A] (verification not implemented)	1220
3.240.7 Maxima [F]	1220
3.240.8 Giac [F]	1221
3.240.9 Mupad [B] (verification not implemented)	1221

3.240.1 Optimal result

Integrand size = 17, antiderivative size = 9

$$\int (e^{e^x} - e^{e^x-x}) dx = e^{e^x-x}$$

output `exp(exp(x)-x)`

3.240.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int (e^{e^x} - e^{e^x-x}) dx = e^{e^x-x}$$

input `Integrate[E^E^x - E^(E^x - x), x]`

output `E^(E^x - x)`

3.240.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e^{e^x} - e^{e^x-x}) dx$$

↓ 2009

$$e^{e^x-x}$$

input `Int[E^E^x - E^(E^x - x), x]`

output `E^(E^x - x)`

3.240.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.240.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

method	result	size
risch	e^{e^x-x}	8
default	$e^{e^x}e^{-x}$	9
norman	$e^{e^x}e^{-x}$	9
parts	$e^{e^x}e^{-x}$	9

input `int(exp(exp(x))-exp(exp(x)-x), x, method=_RETURNVERBOSE)`

output `exp(exp(x)-x)`

3.240.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int (e^{e^x} - e^{e^x-x}) dx = e^{(-x+e^x)}$$

input `integrate(exp(exp(x))-exp(exp(x)-x),x, algorithm="fricas")`output `e^(-x + e^x)`**3.240.6 Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int (e^{e^x} - e^{e^x-x}) dx = e^{-x}e^{e^x}$$

input `integrate(exp(exp(x))-exp(exp(x)-x),x)`output `exp(-x)*exp(exp(x))`**3.240.7 Maxima [F]**

$$\int (e^{e^x} - e^{e^x-x}) dx = \int -e^{(-x+e^x)} + e^{(e^x)} dx$$

input `integrate(exp(exp(x))-exp(exp(x)-x),x, algorithm="maxima")`output `Ei(e^x) - integrate(e^(-x + e^x), x)`

3.240.8 Giac [F]

$$\int (e^{e^x} - e^{e^x-x}) dx = \int -e^{(-x+e^x)} + e^{(e^x)} dx$$

input `integrate(exp(exp(x))-exp(exp(x)-x),x, algorithm="giac")`

output `integrate(-e^(-x + e^x) + e^(e^x), x)`

3.240.9 Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int (e^{e^x} - e^{e^x-x}) dx = e^{e^x-x}$$

input `int(exp(exp(x)) - exp(exp(x) - x),x)`

output `exp(exp(x) - x)`

3.241 $\int \sqrt{1 - \sqrt{x}} dx$

3.241.1 Optimal result	1222
3.241.2 Mathematica [A] (verified)	1222
3.241.3 Rubi [A] (verified)	1223
3.241.4 Maple [A] (verified)	1224
3.241.5 Fricas [A] (verification not implemented)	1224
3.241.6 Sympy [C] (verification not implemented)	1225
3.241.7 Maxima [A] (verification not implemented)	1225
3.241.8 Giac [A] (verification not implemented)	1226
3.241.9 Mupad [B] (verification not implemented)	1226

3.241.1 Optimal result

Integrand size = 13, antiderivative size = 35

$$\int \sqrt{1 - \sqrt{x}} dx = -\frac{4}{3}(1 - \sqrt{x})^{3/2} + \frac{4}{5}(1 - \sqrt{x})^{5/2}$$

output `-4/3*(1-x^(1/2))^(3/2)+4/5*(1-x^(1/2))^(5/2)`

3.241.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

$$\int \sqrt{1 - \sqrt{x}} dx = -\frac{4}{15}(1 - \sqrt{x})^{3/2} (2 + 3\sqrt{x})$$

input `Integrate[Sqrt[1 - Sqrt[x]], x]`

output `(-4*(1 - Sqrt[x])^(3/2)*(2 + 3*Sqrt[x]))/15`

3.241.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {774, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{1 - \sqrt{x}} dx \\ & \quad \downarrow 774 \\ & 2 \int \sqrt{1 - \sqrt{x}} \sqrt{x} d\sqrt{x} \\ & \quad \downarrow 53 \\ & 2 \int \left(\sqrt{1 - \sqrt{x}} - (1 - \sqrt{x})^{3/2} \right) d\sqrt{x} \\ & \quad \downarrow 2009 \\ & 2 \left(\frac{2}{5} (1 - \sqrt{x})^{5/2} - \frac{2}{3} (1 - \sqrt{x})^{3/2} \right) \end{aligned}$$

input `Int[Sqrt[1 - Sqrt[x]],x]`

output `2*((-2*(1 - Sqrt[x])^(3/2))/3 + (2*(1 - Sqrt[x])^(5/2))/5)`

3.241.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 774 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[n]}, Simp[k Subst[Int[x^(k - 1)*(a + b*x^(k*n))^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, p}, x] && FractionQ[n]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.241.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.69

method	result	size
derivativedivides	$-\frac{4(1-\sqrt{x})^{\frac{3}{2}}}{3} + \frac{4(1-\sqrt{x})^{\frac{5}{2}}}{5}$	24
default	$-\frac{4(1-\sqrt{x})^{\frac{3}{2}}}{3} + \frac{4(1-\sqrt{x})^{\frac{5}{2}}}{5}$	24
meijerg	$-\frac{-\frac{8\sqrt{\pi}}{15} + \frac{4\sqrt{\pi}(1-\sqrt{x})^{\frac{3}{2}}(3\sqrt{x}+2)}{15}}{\sqrt{\pi}}$	33

input `int((1-x^(1/2))^(1/2),x,method=_RETURNVERBOSE)`

output `-4/3*(1-x^(1/2))^(3/2)+4/5*(1-x^(1/2))^(5/2)`

3.241.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.60

$$\int \sqrt{1 - \sqrt{x}} dx = \frac{4}{15} (3x - \sqrt{x} - 2) \sqrt{-\sqrt{x} + 1}$$

input `integrate((1-x^(1/2))^(1/2),x, algorithm="fracas")`

output `4/15*(3*x - sqrt(x) - 2)*sqrt(-sqrt(x) + 1)`

3.241.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.56 (sec) , antiderivative size = 306, normalized size of antiderivative = 8.74

$$\int \sqrt{1 - \sqrt{x}} dx$$

$$= \begin{cases} -\frac{12ix^{\frac{7}{2}}\sqrt{\sqrt{x}-1}}{-15x^{\frac{5}{2}}+15x^2} + \frac{4ix^{\frac{5}{2}}\sqrt{\sqrt{x}-1}}{-15x^{\frac{5}{2}}+15x^2} - \frac{8x^{\frac{5}{2}}}{-15x^{\frac{5}{2}}+15x^2} + \frac{16ix^3\sqrt{\sqrt{x}-1}}{-15x^{\frac{5}{2}}+15x^2} - \frac{8ix^2\sqrt{\sqrt{x}-1}}{-15x^{\frac{5}{2}}+15x^2} + \frac{8x^2}{-15x^{\frac{5}{2}}+15x^2} & \text{for } |\sqrt{x}| > 1 \\ -\frac{12x^{\frac{7}{2}}\sqrt{1-\sqrt{x}}}{-15x^{\frac{5}{2}}+15x^2} + \frac{4x^{\frac{5}{2}}\sqrt{1-\sqrt{x}}}{-15x^{\frac{5}{2}}+15x^2} - \frac{8x^{\frac{5}{2}}}{-15x^{\frac{5}{2}}+15x^2} + \frac{16x^3\sqrt{1-\sqrt{x}}}{-15x^{\frac{5}{2}}+15x^2} - \frac{8x^2\sqrt{1-\sqrt{x}}}{-15x^{\frac{5}{2}}+15x^2} + \frac{8x^2}{-15x^{\frac{5}{2}}+15x^2} & \text{otherwise} \end{cases}$$

input `integrate((1-x**(1/2))**(1/2),x)`

output `Piecewise((-12*I*x**(7/2)*sqrt(sqrt(x) - 1)/(-15*x**(5/2) + 15*x**2) + 4*I*x**(5/2)*sqrt(sqrt(x) - 1)/(-15*x**(5/2) + 15*x**2) - 8*x**(5/2)/(-15*x**(5/2) + 15*x**2) + 16*I*x**3*sqrt(sqrt(x) - 1)/(-15*x**(5/2) + 15*x**2) - 8*I*x**2*sqrt(sqrt(x) - 1)/(-15*x**(5/2) + 15*x**2) + 8*x**2/(-15*x**(5/2) + 15*x**2), Abs(sqrt(x)) > 1, (-12*x**(7/2)*sqrt(1 - sqrt(x))/(-15*x**(5/2) + 15*x**2) + 4*x**(5/2)*sqrt(1 - sqrt(x))/(-15*x**(5/2) + 15*x**2) - 8*x**(5/2)/(-15*x**(5/2) + 15*x**2) + 16*x**3*sqrt(1 - sqrt(x))/(-15*x**(5/2) + 15*x**2) - 8*x**2*sqrt(1 - sqrt(x))/(-15*x**(5/2) + 15*x**2) + 8*x**2/(-15*x**(5/2) + 15*x**2), True))`

3.241.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.66

$$\int \sqrt{1 - \sqrt{x}} dx = \frac{4}{5} (-\sqrt{x} + 1)^{\frac{5}{2}} - \frac{4}{3} (-\sqrt{x} + 1)^{\frac{3}{2}}$$

input `integrate((1-x^(1/2))^(1/2),x, algorithm="maxima")`

output `4/5*(-sqrt(x) + 1)^(5/2) - 4/3*(-sqrt(x) + 1)^(3/2)`

3.241.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

$$\int \sqrt{1 - \sqrt{x}} dx = \frac{4}{5} (\sqrt{x} - 1)^2 \sqrt{-\sqrt{x} + 1} - \frac{4}{3} (-\sqrt{x} + 1)^{\frac{3}{2}}$$

input `integrate((1-x^(1/2))^(1/2),x, algorithm="giac")`output `4/5*(sqrt(x) - 1)^2*sqrt(-sqrt(x) + 1) - 4/3*(-sqrt(x) + 1)^(3/2)`**3.241.9 Mupad [B] (verification not implemented)**

Time = 16.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.29

$$\int \sqrt{1 - \sqrt{x}} dx = x {}_2F_1\left(-\frac{1}{2}, 2; 3; \sqrt{x}\right)$$

input `int((1 - x^(1/2))^(1/2),x)`output `x*hypergeom([-1/2, 2], 3, x^(1/2))`

$$3.242 \quad \int \frac{x^3}{1+x+\frac{x^2}{2}+\frac{x^3}{6}} dx$$

3.242.1 Optimal result	1227
3.242.2 Mathematica [A] (verified)	1227
3.242.3 Rubi [B] (verified)	1228
3.242.4 Maple [A] (verified)	1230
3.242.5 Fricas [A] (verification not implemented)	1230
3.242.6 Sympy [A] (verification not implemented)	1231
3.242.7 Maxima [A] (verification not implemented)	1231
3.242.8 Giac [A] (verification not implemented)	1231
3.242.9 Mupad [B] (verification not implemented)	1232

3.242.1 Optimal result

Integrand size = 23, antiderivative size = 24

$$\int \frac{x^3}{1+x+\frac{x^2}{2}+\frac{x^3}{6}} dx = 6x - 6 \log \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} \right)$$

output `6*x-6*ln(1+x+1/2*x^2+1/6*x^3)`

3.242.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{x^3}{1+x+\frac{x^2}{2}+\frac{x^3}{6}} dx = 6(x - \log(6 + 6x + 3x^2 + x^3))$$

input `Integrate[x^3/(1 + x + x^2/2 + x^3/6), x]`

output `6*(x - Log[6 + 6*x + 3*x^2 + x^3])`

3.242.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 194 vs. $2(24) = 48$.

Time = 0.64 (sec) , antiderivative size = 194, normalized size of antiderivative = 8.08, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {2490, 2485, 27, 1200, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{\frac{x^3}{6} + \frac{x^2}{2} + x + 1} dx$$

$$\downarrow \text{2490}$$

$$\int \frac{x^3}{\frac{1}{6}(x+1)^3 + \frac{x+1}{2} + \frac{1}{3}} d(x+1)$$

$$\downarrow \text{2485}$$

$$\frac{1}{36} \int \frac{216x^3}{\left(x + \frac{1 - (-1+\sqrt{2})^{2/3}}{\sqrt[3]{-1+\sqrt{2}}} + 1\right) \left(\frac{(x+1)^2 - \frac{(1 - (-1+\sqrt{2})^{2/3})(x+1)}{\sqrt[3]{-1+\sqrt{2}}} + (-1+\sqrt{2})^{2/3} + \frac{1}{(-1+\sqrt{2})^{2/3}} + 1\right)} d(x+1)$$

$$\downarrow \text{27}$$

$$-6 \int -\frac{x^3}{\left(x - \sqrt[3]{-1+\sqrt{2}} + \frac{1}{\sqrt[3]{-1+\sqrt{2}}} + 1\right) \left(\frac{(x+1)^2 - \frac{(1 - (-1+\sqrt{2})^{2/3})(x+1)}{\sqrt[3]{-1+\sqrt{2}}} + (-1+\sqrt{2})^{2/3} + \frac{1}{(-1+\sqrt{2})^{2/3}} + 1\right)} d(x+1)$$

$$\downarrow \text{1200}$$

$$-6 \int \left(\frac{2 \left((3 - 2\sqrt{2} + (-1 + \sqrt{2})^{2/3} - (-1 + \sqrt{2})^{4/3}) (x+1) - (1 - \sqrt{2}) \left(1 - \sqrt[3]{-1 + \sqrt{2}} - \frac{1}{\sqrt[3]{-1 + \sqrt{2}}} \right) \right)}{\left((1 - (-1 + \sqrt{2})^{2/3} + (-1 + \sqrt{2})^{4/3}) \left((-1 + \sqrt{2})^{2/3} (x+1)^2 - \left(1 - \sqrt{2} + \sqrt[3]{-1 + \sqrt{2}} \right) (x+1) + (-1 + \sqrt{2})^{2/3} \right) \right)} \right) d(x+1)$$

$$\downarrow \text{2009}$$

$$-6 \left(-x + \log \left(\sqrt[3]{\sqrt{2}-1}(x+1) - (\sqrt{2}-1)^{2/3} + 1 \right) + \frac{2(1-\sqrt{2}) \left(1 - \sqrt[3]{\sqrt{2}-1} - (\sqrt{2}-1)^{2/3} \right) \log \left((\sqrt{2}-1 - \sqrt{2} + \sqrt[3]{\sqrt{2}-1}) \right)}{\left(1 - \sqrt{2} + \sqrt[3]{\sqrt{2}-1} \right)} \right)$$

input `Int[x^3/(1 + x + x^2/2 + x^3/6), x]`

output `-6*(-1 - x + Log[1 - (-1 + Sqrt[2])^(2/3) + (-1 + Sqrt[2])^(1/3)*(1 + x)] + (2*(1 - Sqrt[2])*(1 - (-1 + Sqrt[2])^(1/3) - (-1 + Sqrt[2])^(2/3))*Log[1 + (-1 + Sqrt[2])^(2/3) + (-1 + Sqrt[2])^(4/3) - (1 - Sqrt[2] + (-1 + Sqrt[2])^(1/3))*(1 + x) + (-1 + Sqrt[2])^(2/3)*(1 + x)^2])/((1 - Sqrt[2] + (-1 + Sqrt[2])^(1/3))*(1 - (-1 + Sqrt[2])^(2/3) + (-1 + Sqrt[2])^(4/3))))`

3.242.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 1200 `Int[(((d_) + (e_)*(x_)^(m_))*((f_) + (g_)*(x_)^(n_)))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)^n/(a + b*x + c*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && IntegerQ[n]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2485 `Int[((e_) + (f_)*(x_)^(m_))*((a_) + (b_)*(x_) + (d_)*(x_)^3)^(p_), x_Symbol] := With[{r = Rt[-9*a*d^2 + Sqrt[3]*d*Sqrt[4*b^3*d + 27*a^2*d^2], 3]}, Simp[1/d^(2*p) Int[(e + f*x)^m*Simp[18^(1/3)*b*(d/(3*r)) - r/18^(1/3) + d*x, x]^p*Simp[b*(d/3) + 12^(1/3)*b^2*(d^2/(3*r^2)) + r^2/(3*12^(1/3)) - d*(2^(1/3)*b*(d/(3^(1/3)*r)) - r/18^(1/3))*x + d^2*x^2, x]^p, x]] /; FreeQ[{a, b, d, e, f, m}, x] && NeQ[4*b^3 + 27*a^2*d, 0] && ILtQ[p, 0]`

```
rule 2490 Int[(P3_)^(p_.)*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol] := With[{a = Coeff[P3
, x, 0], b = Coeff[P3, x, 1], c = Coeff[P3, x, 2], d = Coeff[P3, x, 3]}, Su
bst[Int[((3*d*e - c*f)/(3*d) + f*x)^m*Simp[(2*c^3 - 9*b*c*d + 27*a*d^2)/(27
*d^2) - (c^2 - 3*b*d)*(x/(3*d)) + d*x^3, x]^p, x], x, x + c/(3*d)] /; NeQ[c
, 0]] /; FreeQ[{e, f, m, p}, x] && PolyQ[P3, x, 3]
```

3.242.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

method	result	size
default	$6x - 6 \ln(x^3 + 3x^2 + 6x + 6)$	21
norman	$6x - 6 \ln(x^3 + 3x^2 + 6x + 6)$	21
risch	$6x - 6 \ln(x^3 + 3x^2 + 6x + 6)$	21
parallelrisk	$6x - 6 \ln(x^3 + 3x^2 + 6x + 6)$	21

```
input int(x^3/(1+x+1/2*x^2+1/6*x^3),x,method=_RETURNVERBOSE)
```

```
output 6*x-6*ln(x^3+3*x^2+6*x+6)
```

3.242.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{x^3}{1+x+\frac{x^2}{2}+\frac{x^3}{6}} dx = 6x - 6 \log(x^3 + 3x^2 + 6x + 6)$$

```
input integrate(x^3/(1+x+1/2*x^2+1/6*x^3),x, algorithm="fricas")
```

```
output 6*x - 6*log(x^3 + 3*x^2 + 6*x + 6)
```

3.242.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \frac{x^3}{1+x+\frac{x^2}{2}+\frac{x^3}{6}} dx = 6x - 6 \log(x^3 + 3x^2 + 6x + 6)$$

input `integrate(x**3/(1+x+1/2*x**2+1/6*x**3),x)`output `6*x - 6*log(x**3 + 3*x**2 + 6*x + 6)`**3.242.7 Maxima [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{x^3}{1+x+\frac{x^2}{2}+\frac{x^3}{6}} dx = 6x - 6 \log(x^3 + 3x^2 + 6x + 6)$$

input `integrate(x^3/(1+x+1/2*x^2+1/6*x^3),x, algorithm="maxima")`output `6*x - 6*log(x^3 + 3*x^2 + 6*x + 6)`**3.242.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

$$\int \frac{x^3}{1+x+\frac{x^2}{2}+\frac{x^3}{6}} dx = 6x - 6 \log(|x^3 + 3x^2 + 6x + 6|)$$

input `integrate(x^3/(1+x+1/2*x^2+1/6*x^3),x, algorithm="giac")`output `6*x - 6*log(abs(x^3 + 3*x^2 + 6*x + 6))`

3.242.9 Mupad [B] (verification not implemented)

Time = 16.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{x^3}{1 + x + \frac{x^2}{2} + \frac{x^3}{6}} dx = 6x - 6 \ln(x^3 + 3x^2 + 6x + 6)$$

input `int(x^3/(x + x^2/2 + x^3/6 + 1),x)`

output `6*x - 6*log(6*x + 3*x^2 + x^3 + 6)`

3.243 $\int (-\sin(x - \sin(x)) + \sin(x + \sin(x))) dx$

3.243.1 Optimal result	1233
3.243.2 Mathematica [A] (verified)	1233
3.243.3 Rubi [F]	1234
3.243.4 Maple [A] (verified)	1234
3.243.5 Fricas [B] (verification not implemented)	1235
3.243.6 Sympy [F]	1235
3.243.7 Maxima [A] (verification not implemented)	1235
3.243.8 Giac [A] (verification not implemented)	1236
3.243.9 Mupad [B] (verification not implemented)	1236

3.243.1 Optimal result

Integrand size = 15, antiderivative size = 5

$$\int (-\sin(x - \sin(x)) + \sin(x + \sin(x))) dx = -2 \cos(\sin(x))$$

output `-2*cos(sin(x))`

3.243.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int (-\sin(x - \sin(x)) + \sin(x + \sin(x))) dx = -2 \cos(\sin(x))$$

input `Integrate[-Sin[x - Sin[x]] + Sin[x + Sin[x]], x]`

output `-2*Cos[Sin[x]]`

3.243.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\sin(x + \sin(x)) - \sin(x - \sin(x))) dx$$

$$\downarrow \text{2009}$$

$$\int \sin(x + \sin(x)) dx - \int \sin(x - \sin(x)) dx$$

input `Int[-Sin[x - Sin[x]] + Sin[x + Sin[x]], x]`

output `$Aborted`

3.243.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.243.4 Maple [A] (verified)

Time = 12.75 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
derivativedivides	$-2 \cos(\sin(x))$	6
default	$-2 \cos(\sin(x))$	6
risch	$-2 \cos(\sin(x))$	6

input `int(sin(x+sin(x))-sin(x-sin(x)),x,method=_RETURNVERBOSE)`

output `-2*cos(sin(x))`

3.243.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. $2(5) = 10$.

Time = 0.24 (sec) , antiderivative size = 65, normalized size of antiderivative = 13.00

$$\begin{aligned} & \int (-\sin(x - \sin(x)) + \sin(x + \sin(x))) dx \\ &= -2 \cos(x) \cos\left(\frac{x \tan\left(\frac{1}{2}x\right)^2 + x + 2 \tan\left(\frac{1}{2}x\right)}{\tan\left(\frac{1}{2}x\right)^2 + 1}\right) \\ & \quad - 2 \sin(x) \sin\left(\frac{x \tan\left(\frac{1}{2}x\right)^2 + x + 2 \tan\left(\frac{1}{2}x\right)}{\tan\left(\frac{1}{2}x\right)^2 + 1}\right) \end{aligned}$$

input `integrate(sin(x+sin(x))-sin(x-sin(x)),x, algorithm="fracas")`

output `-2*cos(x)*cos((x*tan(1/2*x)^2 + x + 2*tan(1/2*x))/(tan(1/2*x)^2 + 1)) - 2*
sin(x)*sin((x*tan(1/2*x)^2 + x + 2*tan(1/2*x))/(tan(1/2*x)^2 + 1))`

3.243.6 Sympy [F]

$$\int (-\sin(x - \sin(x)) + \sin(x + \sin(x))) dx = \int (-\sin(x - \sin(x)) + \sin(x + \sin(x))) dx$$

input `integrate(sin(x+sin(x))-sin(x-sin(x)),x)`

output `Integral(-sin(x - sin(x)) + sin(x + sin(x)), x)`

3.243.7 Maxima [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int (-\sin(x - \sin(x)) + \sin(x + \sin(x))) dx = -2 \cos(\sin(x))$$

input `integrate(sin(x+sin(x))-sin(x-sin(x)),x, algorithm="maxima")`

output `-2*cos(sin(x))`

3.243.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int (-\sin(x - \sin(x)) + \sin(x + \sin(x))) dx = -2 \cos(\sin(x))$$

input `integrate(sin(x+sin(x))-sin(x-sin(x)),x, algorithm="giac")`

output `-2*cos(sin(x))`

3.243.9 Mupad [B] (verification not implemented)

Time = 16.49 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int (-\sin(x - \sin(x)) + \sin(x + \sin(x))) dx = -2 \cos(\sin(x))$$

input `int(sin(x + sin(x)) - sin(x - sin(x)),x)`

output `-2*cos(sin(x))`

3.244 $\int (\sec^5(x) \tan^2(x) + \sec^3(x) \tan^4(x)) dx$

3.244.1 Optimal result	1237
3.244.2 Mathematica [A] (verified)	1237
3.244.3 Rubi [B] (verified)	1238
3.244.4 Maple [C] (verified)	1238
3.244.5 Fricas [A] (verification not implemented)	1239
3.244.6 Sympy [B] (verification not implemented)	1239
3.244.7 Maxima [B] (verification not implemented)	1240
3.244.8 Giac [B] (verification not implemented)	1240
3.244.9 Mupad [B] (verification not implemented)	1241

3.244.1 Optimal result

Integrand size = 19, antiderivative size = 12

$$\int (\sec^5(x) \tan^2(x) + \sec^3(x) \tan^4(x)) dx = \frac{1}{3} \sec^3(x) \tan^3(x)$$

output `1/3*sec(x)^3*tan(x)^3`

3.244.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int (\sec^5(x) \tan^2(x) + \sec^3(x) \tan^4(x)) dx = \frac{1}{3} \sec^3(x) \tan^3(x)$$

input `Integrate[Sec[x]^5*Tan[x]^2 + Sec[x]^3*Tan[x]^4,x]`

output `(Sec[x]^3*Tan[x]^3)/3`

3.244.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 33 vs. $2(12) = 24$.

Time = 0.23 (sec) , antiderivative size = 33, normalized size of antiderivative = 2.75, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\tan^2(x) \sec^5(x) + \tan^4(x) \sec^3(x)) dx$$

↓ 2009

$$\frac{1}{6} \tan(x) \sec^5(x) + \frac{1}{6} \tan^3(x) \sec^3(x) - \frac{1}{6} \tan(x) \sec^3(x)$$

input `Int[Sec[x]^5*Tan[x]^2 + Sec[x]^3*Tan[x]^4,x]`

output `-1/6*(Sec[x]^3*Tan[x]) + (Sec[x]^5*Tan[x])/6 + (Sec[x]^3*Tan[x]^3)/6`

3.244.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.244.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 6.33 (sec) , antiderivative size = 40, normalized size of antiderivative = 3.33

method	result	size
risch	$\frac{8i(e^{9ix} - 3e^{7ix} + 3e^{5ix} - e^{3ix})}{3(e^{2ix} + 1)^6}$	40
default	$\frac{\sin(x)^3}{6 \cos(x)^6} + \frac{\sin(x)^3}{8 \cos(x)^4} + \frac{\sin(x)^3}{16 \cos(x)^2} + \frac{\sin(x)^5}{6 \cos(x)^6} + \frac{\sin(x)^5}{24 \cos(x)^4} - \frac{\sin(x)^5}{48 \cos(x)^2} - \frac{\sin(x)^3}{48}$	68
parts	$\frac{\sin(x)^3}{6 \cos(x)^6} + \frac{\sin(x)^3}{8 \cos(x)^4} + \frac{\sin(x)^3}{16 \cos(x)^2} + \frac{\sin(x)^5}{6 \cos(x)^6} + \frac{\sin(x)^5}{24 \cos(x)^4} - \frac{\sin(x)^5}{48 \cos(x)^2} - \frac{\sin(x)^3}{48}$	68

input `int(tan(x)^4*sec(x)^3+tan(x)^2*sec(x)^5,x,method=_RETURNVERBOSE)`

output `8/3*I/(exp(2*I*x)+1)^6*(exp(9*I*x)-3*exp(7*I*x)+3*exp(5*I*x)-exp(3*I*x))`

3.244.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int (\sec^5(x) \tan^2(x) + \sec^3(x) \tan^4(x)) dx = -\frac{(\cos(x)^2 - 1) \sin(x)}{3 \cos(x)^6}$$

input `integrate(tan(x)^4*sec(x)^3+tan(x)^2*sec(x)^5,x, algorithm="fricas")`

output `-1/3*(cos(x)^2 - 1)*sin(x)/cos(x)^6`

3.244.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 80 vs. $2(10) = 20$.

Time = 0.14 (sec) , antiderivative size = 80, normalized size of antiderivative = 6.67

$$\int (\sec^5(x) \tan^2(x) + \sec^3(x) \tan^4(x)) dx = \frac{-3 \sin^5(x) - 8 \sin^3(x) + 3 \sin(x)}{48 \sin^6(x) - 144 \sin^4(x) + 144 \sin^2(x) - 48} + \frac{3 \sin^5(x) - 8 \sin^3(x) - 3 \sin(x)}{48 \sin^6(x) - 144 \sin^4(x) + 144 \sin^2(x) - 48}$$

input `integrate(tan(x)**4*sec(x)**3+tan(x)**2*sec(x)**5,x)`

output `(-3*sin(x)**5 - 8*sin(x)**3 + 3*sin(x))/(48*sin(x)**6 - 144*sin(x)**4 + 144*sin(x)**2 - 48) + (3*sin(x)**5 - 8*sin(x)**3 - 3*sin(x))/(48*sin(x)**6 - 144*sin(x)**4 + 144*sin(x)**2 - 48)`

3.244.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 79 vs. $2(10) = 20$.

Time = 0.21 (sec) , antiderivative size = 79, normalized size of antiderivative = 6.58

$$\int (\sec^5(x) \tan^2(x) + \sec^3(x) \tan^4(x)) dx = -\frac{3 \sin(x)^5 + 8 \sin(x)^3 - 3 \sin(x)}{48 (\sin(x)^6 - 3 \sin(x)^4 + 3 \sin(x)^2 - 1)} + \frac{3 \sin(x)^5 - 8 \sin(x)^3 - 3 \sin(x)}{48 (\sin(x)^6 - 3 \sin(x)^4 + 3 \sin(x)^2 - 1)}$$

input `integrate(tan(x)^4*sec(x)^3+tan(x)^2*sec(x)^5,x, algorithm="maxima")`

output `-1/48*(3*sin(x)^5 + 8*sin(x)^3 - 3*sin(x))/(sin(x)^6 - 3*sin(x)^4 + 3*sin(x)^2 - 1) + 1/48*(3*sin(x)^5 - 8*sin(x)^3 - 3*sin(x))/(sin(x)^6 - 3*sin(x)^4 + 3*sin(x)^2 - 1)`

3.244.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 55 vs. $2(10) = 20$.

Time = 0.27 (sec) , antiderivative size = 55, normalized size of antiderivative = 4.58

$$\int (\sec^5(x) \tan^2(x) + \sec^3(x) \tan^4(x)) dx = -\frac{3 \sin(x)^5 + 8 \sin(x)^3 - 3 \sin(x)}{48 (\sin(x)^2 - 1)^3} + \frac{3 \sin(x)^5 - 8 \sin(x)^3 - 3 \sin(x)}{48 (\sin(x)^2 - 1)^3}$$

input `integrate(tan(x)^4*sec(x)^3+tan(x)^2*sec(x)^5,x, algorithm="giac")`

output `-1/48*(3*sin(x)^5 + 8*sin(x)^3 - 3*sin(x))/(sin(x)^2 - 1)^3 + 1/48*(3*sin(x)^5 - 8*sin(x)^3 - 3*sin(x))/(sin(x)^2 - 1)^3`

3.244.9 Mupad [B] (verification not implemented)

Time = 16.36 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int (\sec^5(x) \tan^2(x) + \sec^3(x) \tan^4(x)) dx = -\frac{\sin(x)^3}{3(\sin(x)^2 - 1)^3}$$

input `int(tan(x)^4/cos(x)^3 + tan(x)^2/cos(x)^5,x)`

output `-sin(x)^3/(3*(sin(x)^2 - 1)^3)`

3.245 $\int (1 + \log(x)) \log(\log(x)) dx$

3.245.1 Optimal result	1242
3.245.2 Mathematica [A] (verified)	1242
3.245.3 Rubi [F]	1243
3.245.4 Maple [A] (verified)	1243
3.245.5 Fricas [A] (verification not implemented)	1244
3.245.6 Sympy [A] (verification not implemented)	1244
3.245.7 Maxima [A] (verification not implemented)	1244
3.245.8 Giac [A] (verification not implemented)	1245
3.245.9 Mupad [B] (verification not implemented)	1245

3.245.1 Optimal result

Integrand size = 8, antiderivative size = 10

$$\int (1 + \log(x)) \log(\log(x)) dx = x(-1 + \log(x) \log(\log(x)))$$

output `x*(-1+ln(x)*ln(ln(x)))`

3.245.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.80

$$\int (1 + \log(x)) \log(\log(x)) dx = -x + x \log(\log(x)) + x(-1 + \log(x)) \log(\log(x))$$

input `Integrate[(1 + Log[x])*Log[Log[x]], x]`

output `-x + x*Log[Log[x]] + x*(-1 + Log[x])*Log[Log[x]]`

3.245.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\log(x) + 1) \log(\log(x)) dx$$

$$\downarrow \text{7293}$$

$$\int (\log(x) \log(\log(x)) + \log(\log(x))) dx$$

$$\downarrow \text{2009}$$

$$\int \log(x) \log(\log(x)) dx - \text{LogIntegral}(x) + x \log(\log(x))$$

input `Int[(1 + Log[x])*Log[Log[x]],x]`

output `$Aborted`

3.245.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

3.245.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

method	result	size
norman	$x \ln(x) \ln(\ln(x)) - x$	12
risch	$x \ln(x) \ln(\ln(x)) - x$	12
parallelrisc	$x \ln(x) \ln(\ln(x)) - x$	12
default	$(-1 + \ln(x)) x \ln(\ln(x)) - x + x \ln(\ln(x))$	19
parts	$(-1 + \ln(x)) x \ln(\ln(x)) - x + x \ln(\ln(x))$	19

input `int((ln(x)+1)*ln(ln(x)),x,method=_RETURNVERBOSE)`

output `x*ln(x)*ln(ln(x))-x`

3.245.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int (1 + \log(x)) \log(\log(x)) dx = x \log(x) \log(\log(x)) - x$$

input `integrate((log(x)+1)*log(log(x)),x, algorithm="fricas")`

output `x*log(x)*log(log(x)) - x`

3.245.6 Sympy [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int (1 + \log(x)) \log(\log(x)) dx = x \log(x) \log(\log(x)) - x$$

input `integrate((ln(x)+1)*ln(ln(x)),x)`

output `x*log(x)*log(log(x)) - x`

3.245.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.60

$$\int (1 + \log(x)) \log(\log(x)) dx = (x(\log(x) - 1) + x) \log(\log(x)) - x$$

input `integrate((log(x)+1)*log(log(x)),x, algorithm="maxima")`

output `(x*(log(x) - 1) + x)*log(log(x)) - x`

3.245.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int (1 + \log(x)) \log(\log(x)) dx = x \log(x) \log(\log(x)) - x$$

input `integrate((log(x)+1)*log(log(x)),x, algorithm="giac")`

output `x*log(x)*log(log(x)) - x`

3.245.9 Mupad [B] (verification not implemented)

Time = 16.43 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int (1 + \log(x)) \log(\log(x)) dx = x \ln(\ln(x)) \ln(x) - x$$

input `int(log(log(x))*(log(x) + 1),x)`

output `x*log(log(x))*log(x) - x`

3.246 $\int \left(\frac{1}{1+\cos(x)} + \frac{1}{1+\cot(x)} + \frac{1}{1+\csc(x)} + \frac{1}{1+\sec(x)} + \frac{1}{1+\sin(x)} + \frac{1}{1+\tan(x)} \right) dx = 3x$

3.246.1 Optimal result	1246
3.246.2 Mathematica [A] (verified)	1246
3.246.3 Rubi [C] (verified)	1247
3.246.4 Maple [A] (verified)	1247
3.246.5 Fricas [A] (verification not implemented)	1248
3.246.6 Sympy [F]	1248
3.246.7 Maxima [C] (verification not implemented)	1249
3.246.8 Giac [C] (verification not implemented)	1250
3.246.9 Mupad [B] (verification not implemented)	1250

3.246.1 Optimal result

Integrand size = 37, antiderivative size = 3

$$\int \left(\frac{1}{1+\cos(x)} + \frac{1}{1+\cot(x)} + \frac{1}{1+\csc(x)} + \frac{1}{1+\sec(x)} + \frac{1}{1+\sin(x)} + \frac{1}{1+\tan(x)} \right) dx = 3x$$

output 3*x

3.246.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \left(\frac{1}{1+\cos(x)} + \frac{1}{1+\cot(x)} + \frac{1}{1+\csc(x)} + \frac{1}{1+\sec(x)} + \frac{1}{1+\sin(x)} + \frac{1}{1+\tan(x)} \right) dx = 3x$$

input Integrate[(1 + Cos[x])^(-1) + (1 + Cot[x])^(-1) + (1 + Csc[x])^(-1) + (1 + Sec[x])^(-1) + (1 + Sin[x])^(-1) + (1 + Tan[x])^(-1), x]

output 3*x

3.246. $\int \left(\frac{1}{1+\cos(x)} + \frac{1}{1+\cot(x)} + \frac{1}{1+\csc(x)} + \frac{1}{1+\sec(x)} + \frac{1}{1+\sin(x)} + \frac{1}{1+\tan(x)} \right) dx$

3.246.3 Rubi [C] (verified)

Result contains higher order function than in optimal. Order 3 vs. order 1 in optimal.

Time = 0.24 (sec) , antiderivative size = 42, normalized size of antiderivative = 14.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.027$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\frac{1}{\sin(x)+1} + \frac{1}{\cos(x)+1} + \frac{1}{\tan(x)+1} + \frac{1}{\cot(x)+1} + \frac{1}{\csc(x)+1} + \frac{1}{\sec(x)+1} \right) dx$$

↓ 2009

$$3x + \frac{\sin(x)}{\cos(x)+1} - \frac{\cos(x)}{\sin(x)+1} + \frac{\cot(x)}{\csc(x)+1} - \frac{\tan(x)}{\sec(x)+1}$$

input `Int[(1 + Cos[x])^(-1) + (1 + Cot[x])^(-1) + (1 + Csc[x])^(-1) + (1 + Sec[x])^(-1) + (1 + Sin[x])^(-1) + (1 + Tan[x])^(-1), x]`

output `3*x + Cot[x]/(1 + Csc[x]) + Sin[x]/(1 + Cos[x]) - Cos[x]/(1 + Sin[x]) - Tan[x]/(1 + Sec[x])`

3.246.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.246.4 Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result
risch	$3x$
norman	$\frac{3x+3x \tan(\frac{x}{2})}{1+\tan(\frac{x}{2})}$
default	$-\frac{\ln(1+\cot(x))}{2} + \frac{\ln(1+\cot(x)^2)}{4} - \frac{\pi}{4} + \frac{\operatorname{arccot}(\cot(x))}{2} + 4 \arctan\left(\tan\left(\frac{x}{2}\right)\right) + \frac{\ln(1+\tan(x))}{2} - \frac{\ln(1+\tan(x)^2)}{4} +$

3.246. $\int \left(\frac{1}{1+\cos(x)} + \frac{1}{1+\cot(x)} + \frac{1}{1+\csc(x)} + \frac{1}{1+\sec(x)} + \frac{1}{1+\sin(x)} + \frac{1}{1+\tan(x)} \right) dx$

input `int(1/(1+sin(x))+1/(1+cos(x))+1/(1+tan(x))+1/(1+cot(x))+1/(1+sec(x))+1/(1+csc(x)),x,method=_RETURNVERBOSE)`

output `3*x`

3.246.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \left(\frac{1}{1 + \cos(x)} + \frac{1}{1 + \cot(x)} + \frac{1}{1 + \csc(x)} + \frac{1}{1 + \sec(x)} + \frac{1}{1 + \sin(x)} + \frac{1}{1 + \tan(x)} \right) dx = 3x$$

input `integrate(1/(1+sin(x))+1/(1+cos(x))+1/(1+tan(x))+1/(1+cot(x))+1/(1+sec(x))+1/(1+csc(x)),x, algorithm="fricas")`

output `3*x`

3.246.6 Sympy [F]

$$\int \left(\frac{1}{1 + \cos(x)} + \frac{1}{1 + \cot(x)} + \frac{1}{1 + \csc(x)} + \frac{1}{1 + \sec(x)} + \frac{1}{1 + \sin(x)} + \frac{1}{1 + \tan(x)} \right) dx$$

= Too large to display

input `integrate(1/(1+sin(x))+1/(1+cos(x))+1/(1+tan(x))+1/(1+cot(x))+1/(1+sec(x))+1/(1+csc(x)),x)`

3.246. $\int \left(\frac{1}{1 + \cos(x)} + \frac{1}{1 + \cot(x)} + \frac{1}{1 + \csc(x)} + \frac{1}{1 + \sec(x)} + \frac{1}{1 + \sin(x)} + \frac{1}{1 + \tan(x)} \right) dx$

output `Integral((sin(x)*cos(x)*tan(x)*cot(x)*csc(x) + sin(x)*cos(x)*tan(x)*cot(x)*sec(x) + 2*sin(x)*cos(x)*tan(x)*cot(x) + sin(x)*cos(x)*tan(x)*csc(x)*sec(x) + 2*sin(x)*cos(x)*tan(x)*csc(x) + 2*sin(x)*cos(x)*tan(x)*sec(x) + 3*sin(x)*cos(x)*tan(x) + sin(x)*cos(x)*cot(x)*csc(x)*sec(x) + 2*sin(x)*cos(x)*cot(x)*csc(x) + 2*sin(x)*cos(x)*cot(x)*sec(x) + 3*sin(x)*cos(x)*cot(x) + 2*sin(x)*cos(x)*csc(x)*sec(x) + 3*sin(x)*cos(x)*csc(x) + 3*sin(x)*cos(x)*sec(x) + 4*sin(x)*cos(x) + sin(x)*tan(x)*cot(x)*csc(x)*sec(x) + 2*sin(x)*tan(x)*cot(x)*csc(x) + 2*sin(x)*tan(x)*cot(x)*sec(x) + 3*sin(x)*tan(x)*cot(x) + 2*sin(x)*tan(x)*csc(x)*sec(x) + 3*sin(x)*tan(x)*csc(x) + 3*sin(x)*tan(x)*sec(x) + 4*sin(x)*tan(x) + 2*sin(x)*cot(x)*csc(x)*sec(x) + 3*sin(x)*cot(x)*csc(x) + 3*sin(x)*cot(x)*sec(x) + 4*sin(x)*cot(x) + 3*sin(x)*csc(x)*sec(x) + 4*sin(x)*csc(x) + 4*sin(x)*sec(x) + 5*sin(x) + cos(x)*tan(x)*cot(x)*csc(x)*sec(x) + 2*cos(x)*tan(x)*cot(x)*csc(x) + 2*cos(x)*tan(x)*cot(x)*sec(x) + 3*cos(x)*tan(x)*cot(x) + 2*cos(x)*tan(x)*csc(x)*sec(x) + 3*cos(x)*tan(x)*csc(x) + 3*cos(x)*tan(x)*sec(x) + 4*cos(x)*tan(x) + 2*cos(x)*cot(x)*csc(x)*sec(x) + 3*cos(x)*cot(x)*csc(x) + 3*cos(x)*cot(x)*sec(x) + 4*cos(x)*cot(x) + 3*cos(x)*csc(x)*sec(x) + 4*cos(x)*csc(x) + 4*cos(x)*sec(x) + 5*cos(x) + 2*tan(x)*cot(x)*csc(x)*sec(x) + 3*tan(x)*cot(x)*csc(x) + 3*tan(x)*cot(x)*sec(x) + 4*tan(x)*cot(x) + 3*tan(x)*csc(x)*sec(x) + 4*tan(x)*csc(x) + 4*tan(x)*sec(x) + 5*tan(x) + 3*cot(x)*csc(x)*sec(x) + 4*cot(x)*csc(x) ...`

3.246.7 Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 3 vs. order 1.

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 4.67

$$\int \left(\frac{1}{1 + \cos(x)} + \frac{1}{1 + \cot(x)} + \frac{1}{1 + \csc(x)} + \frac{1}{1 + \sec(x)} + \frac{1}{1 + \sin(x)} + \frac{1}{1 + \tan(x)} \right) dx$$

$$= x + 4 \arctan \left(\frac{\sin(x)}{\cos(x) + 1} \right)$$

input `integrate(1/(1+sin(x))+1/(1+cos(x))+1/(1+tan(x))+1/(1+cot(x))+1/(1+sec(x))+1/(1+csc(x)),x, algorithm="maxima")`

output `x + 4*arctan(sin(x)/(cos(x) + 1))`

3.246. $\int \left(\frac{1}{1 + \cos(x)} + \frac{1}{1 + \cot(x)} + \frac{1}{1 + \csc(x)} + \frac{1}{1 + \sec(x)} + \frac{1}{1 + \sin(x)} + \frac{1}{1 + \tan(x)} \right) dx$

3.246.8 Giac [C] (verification not implemented)

Result contains higher order function than in optimal. Order 3 vs. order 1.

Time = 0.29 (sec) , antiderivative size = 40, normalized size of antiderivative = 13.33

$$\int \left(\frac{1}{1 + \cos(x)} + \frac{1}{1 + \cot(x)} + \frac{1}{1 + \csc(x)} + \frac{1}{1 + \sec(x)} + \frac{1}{1 + \sin(x)} + \frac{1}{1 + \tan(x)} \right) dx$$

$$= 3x - \frac{2 \tan\left(\frac{1}{2}x\right)}{(x^2 + 1)\left(\frac{x^2 - 1}{x^2 + 1} - 1\right)} - \tan\left(\frac{1}{2}x\right)$$

input `integrate(1/(1+sin(x))+1/(1+cos(x))+1/(1+tan(x))+1/(1+cot(x))+1/(1+sec(x))+1/(1+csc(x)),x, algorithm="giac")`

output `3*x - 2*tan(1/2*x)/((x^2 + 1)*((x^2 - 1)/(x^2 + 1) - 1)) - tan(1/2*x)`

3.246.9 Mupad [B] (verification not implemented)

Time = 15.74 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \left(\frac{1}{1 + \cos(x)} + \frac{1}{1 + \cot(x)} + \frac{1}{1 + \csc(x)} + \frac{1}{1 + \sec(x)} + \frac{1}{1 + \sin(x)} + \frac{1}{1 + \tan(x)} \right) dx = 3x$$

input `int(1/(1/sin(x) + 1) + 1/(cos(x) + 1) + 1/(cot(x) + 1) + 1/(sin(x) + 1) + 1/(tan(x) + 1) + 1/(1/cos(x) + 1),x)`

output `3*x`

3.246. $\int \left(\frac{1}{1 + \cos(x)} + \frac{1}{1 + \cot(x)} + \frac{1}{1 + \csc(x)} + \frac{1}{1 + \sec(x)} + \frac{1}{1 + \sin(x)} + \frac{1}{1 + \tan(x)} \right) dx$

3.247 $\int \frac{1}{\sqrt{x-x^2}} dx$

3.247.1 Optimal result	1251
3.247.2 Mathematica [B] (verified)	1251
3.247.3 Rubi [A] (verified)	1252
3.247.4 Maple [A] (verified)	1253
3.247.5 Fricas [B] (verification not implemented)	1253
3.247.6 Sympy [A] (verification not implemented)	1253
3.247.7 Maxima [A] (verification not implemented)	1254
3.247.8 Giac [B] (verification not implemented)	1254
3.247.9 Mupad [B] (verification not implemented)	1254

3.247.1 Optimal result

Integrand size = 11, antiderivative size = 8

$$\int \frac{1}{\sqrt{x-x^2}} dx = -\arcsin(1-2x)$$

output `arcsin(-1+2*x)`

3.247.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 40 vs. $2(8) = 16$.

Time = 0.01 (sec) , antiderivative size = 40, normalized size of antiderivative = 5.00

$$\int \frac{1}{\sqrt{x-x^2}} dx = -\frac{2\sqrt{-1+x}\sqrt{x}\log(\sqrt{-1+x}-\sqrt{x})}{\sqrt{-((-1+x)x)}}$$

input `Integrate[1/Sqrt[x - x^2],x]`

output `(-2*Sqrt[-1 + x]*Sqrt[x]*Log[Sqrt[-1 + x] - Sqrt[x]])/Sqrt[-((-1 + x)*x)]`

3.247.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {1090, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{1}{\sqrt{x-x^2}} dx \\ \downarrow 1090 \\ - \int \frac{1}{\sqrt{1-(1-2x)^2}} d(1-2x) \\ \downarrow 223 \\ - \arcsin(1-2x) \end{array}$$

input `Int[1/Sqrt[x - x^2],x]`

output `-ArcSin[1 - 2*x]`

3.247.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 1090 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> Simp[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p) Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

3.247.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
default	$\arcsin(-1 + 2x)$	7
meijerg	$2 \arcsin(\sqrt{x})$	7
pseudoelliptic	$-2 \arctan\left(\frac{\sqrt{-(-1+x)x}}{x}\right)$	16
trager	$\text{RootOf}(-Z^2 + 1) \ln(-2 \text{RootOf}(-Z^2 + 1)x + 2\sqrt{-x^2 + x} + \text{RootOf}(-Z^2 + 1))$	36

input `int(1/(-x^2+x)^(1/2),x,method=_RETURNVERBOSE)`output `arcsin(-1+2*x)`**3.247.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 16 vs. 2(6) = 12.

Time = 0.23 (sec) , antiderivative size = 16, normalized size of antiderivative = 2.00

$$\int \frac{1}{\sqrt{x-x^2}} dx = -2 \arctan\left(\frac{\sqrt{-x^2+x}}{x}\right)$$

input `integrate(1/(-x^2+x)^(1/2),x, algorithm="fracas")`output `-2*arctan(sqrt(-x^2 + x)/x)`**3.247.6 Sympy [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \frac{1}{\sqrt{x-x^2}} dx = \text{asin}(2x - 1)$$

input `integrate(1/(-x**2+x)**(1/2),x)`output `asin(2*x - 1)`

3.247.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{x-x^2}} dx = \arcsin(2x-1)$$

input `integrate(1/(-x^2+x)^(1/2),x, algorithm="maxima")`

output `arcsin(2*x - 1)`

3.247.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. 2(6) = 12.

Time = 0.26 (sec) , antiderivative size = 25, normalized size of antiderivative = 3.12

$$\int \frac{1}{\sqrt{x-x^2}} dx = \frac{1}{4} \sqrt{-x^2+x}(2x-1) + \frac{1}{8} \arcsin(2x-1)$$

input `integrate(1/(-x^2+x)^(1/2),x, algorithm="giac")`

output `1/4*sqrt(-x^2 + x)*(2*x - 1) + 1/8*arcsin(2*x - 1)`

3.247.9 Mupad [B] (verification not implemented)

Time = 14.98 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{x-x^2}} dx = \operatorname{asin}(2x-1)$$

input `int(1/(x - x^2)^(1/2),x)`

output `asin(2*x - 1)`

3.248 $\int \frac{1}{1+\cos^2(x)} dx$

3.248.1 Optimal result	1255
3.248.2 Mathematica [A] (verified)	1255
3.248.3 Rubi [A] (verified)	1256
3.248.4 Maple [A] (verified)	1257
3.248.5 Fricas [A] (verification not implemented)	1257
3.248.6 Sympy [A] (verification not implemented)	1258
3.248.7 Maxima [A] (verification not implemented)	1258
3.248.8 Giac [A] (verification not implemented)	1258
3.248.9 Mupad [B] (verification not implemented)	1259

3.248.1 Optimal result

Integrand size = 8, antiderivative size = 34

$$\int \frac{1}{1+\cos^2(x)} dx = \frac{x}{\sqrt{2}} - \frac{\arctan\left(\frac{\cos(x)\sin(x)}{1+\sqrt{2}+\cos^2(x)}\right)}{\sqrt{2}}$$

output `1/2*x*2^(1/2)-1/2*arctan(cos(x)*sin(x)/(1+2^(1/2)+cos(x)^2))*2^(1/2)`

3.248.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.44

$$\int \frac{1}{1+\cos^2(x)} dx = \frac{\arctan\left(\frac{\tan(x)}{\sqrt{2}}\right)}{\sqrt{2}}$$

input `Integrate[(1 + Cos[x]^2)^(-1), x]`

output `ArcTan[Tan[x]/Sqrt[2]]/Sqrt[2]`

3.248.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.47, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3042, 3660, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\cos^2(x) + 1} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{\sin(x + \frac{\pi}{2})^2 + 1} dx \\ & \quad \downarrow \text{3660} \\ & - \int \frac{1}{2 \cot^2(x) + 1} d \cot(x) \\ & \quad \downarrow \text{216} \\ & - \frac{\arctan(\sqrt{2} \cot(x))}{\sqrt{2}} \end{aligned}$$

input `Int[(1 + Cos[x]^2)^(-1),x]`

output `-(ArcTan[Sqrt[2]*Cot[x]]/Sqrt[2])`

3.248.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3660 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(-1), x_Symbol] := With[{ff =
FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[1/(a + (a + b)*ff^2*x^
2), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, e, f}, x]
```

3.248.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.41

method	result	size
default	$\frac{\sqrt{2} \arctan\left(\frac{\tan(x)\sqrt{2}}{2}\right)}{2}$	14
risch	$\frac{i\sqrt{2} \ln(e^{2ix} + 2\sqrt{2} + 3)}{4} - \frac{i\sqrt{2} \ln(e^{2ix} - 2\sqrt{2} + 3)}{4}$	40

```
input int(1/(1+cos(x)^2),x,method=_RETURNVERBOSE)
```

```
output 1/2*2^(1/2)*arctan(1/2*tan(x)*2^(1/2))
```

3.248.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.91

$$\int \frac{1}{1 + \cos^2(x)} dx = -\frac{1}{4} \sqrt{2} \arctan\left(\frac{3\sqrt{2} \cos(x)^2 - \sqrt{2}}{4 \cos(x) \sin(x)}\right)$$

```
input integrate(1/(1+cos(x)^2),x, algorithm="fracas")
```

```
output -1/4*sqrt(2)*arctan(1/4*(3*sqrt(2)*cos(x)^2 - sqrt(2))/(cos(x)*sin(x)))
```

3.248.6 Sympy [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.85

$$\int \frac{1}{1 + \cos^2(x)} dx = \frac{\sqrt{2} \left(\operatorname{atan} \left(\sqrt{2} \tan \left(\frac{x}{2} \right) - 1 \right) + \pi \left\lfloor \frac{\frac{x}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{2} + \frac{\sqrt{2} \left(\operatorname{atan} \left(\sqrt{2} \tan \left(\frac{x}{2} \right) + 1 \right) + \pi \left\lfloor \frac{\frac{x}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{2}$$

input `integrate(1/(1+cos(x)**2),x)`output `sqrt(2)*(atan(sqrt(2)*tan(x/2) - 1) + pi*floor((x/2 - pi/2)/pi))/2 + sqrt(2)*(atan(sqrt(2)*tan(x/2) + 1) + pi*floor((x/2 - pi/2)/pi))/2`**3.248.7 Maxima [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.38

$$\int \frac{1}{1 + \cos^2(x)} dx = \frac{1}{2} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} \tan(x) \right)$$

input `integrate(1/(1+cos(x)^2),x, algorithm="maxima")`output `1/2*sqrt(2)*arctan(1/2*sqrt(2)*tan(x))`**3.248.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.35

$$\int \frac{1}{1 + \cos^2(x)} dx = \frac{1}{2} \sqrt{2} \left(x + \arctan \left(-\frac{\sqrt{2} \sin(2x) - \sin(2x)}{\sqrt{2} \cos(2x) + \sqrt{2} - \cos(2x) + 1} \right) \right)$$

input `integrate(1/(1+cos(x)^2),x, algorithm="giac")`output `1/2*sqrt(2)*(x + arctan(-(sqrt(2)*sin(2*x) - sin(2*x))/(sqrt(2)*cos(2*x) + sqrt(2) - cos(2*x) + 1)))`

3.248.9 Mupad [B] (verification not implemented)

Time = 14.83 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.76

$$\int \frac{1}{1 + \cos^2(x)} dx = \frac{\sqrt{2}(x - \operatorname{atan}(\tan(x)))}{2} + \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2} \tan(x)}{2}\right)}{2}$$

input `int(1/(cos(x)^2 + 1),x)`

output `(2^(1/2)*(x - atan(tan(x))))/2 + (2^(1/2)*atan((2^(1/2)*tan(x))/2))/2`

3.249 $\int \frac{\log(1+x)}{x^2} dx$

3.249.1 Optimal result	1260
3.249.2 Mathematica [A] (verified)	1260
3.249.3 Rubi [A] (verified)	1261
3.249.4 Maple [A] (verified)	1262
3.249.5 Fricas [A] (verification not implemented)	1263
3.249.6 Sympy [A] (verification not implemented)	1263
3.249.7 Maxima [A] (verification not implemented)	1263
3.249.8 Giac [A] (verification not implemented)	1264
3.249.9 Mupad [B] (verification not implemented)	1264

3.249.1 Optimal result

Integrand size = 8, antiderivative size = 18

$$\int \frac{\log(1+x)}{x^2} dx = \log(x) - \log(1+x) - \frac{\log(1+x)}{x}$$

output `ln(x)-ln(1+x)-ln(1+x)/x`

3.249.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{\log(1+x)}{x^2} dx = \log(x) - \log(1+x) - \frac{\log(1+x)}{x}$$

input `Integrate[Log[1 + x]/x^2,x]`

output `Log[x] - Log[1 + x] - Log[1 + x]/x`

3.249.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {2842, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\log(x+1)}{x^2} dx \\
 & \quad \downarrow \text{2842} \\
 & \int \frac{1}{x(x+1)} dx - \frac{\log(x+1)}{x} \\
 & \quad \downarrow \text{47} \\
 & \int \frac{1}{x} dx - \int \frac{1}{x+1} dx - \frac{\log(x+1)}{x} \\
 & \quad \downarrow \text{14} \\
 & - \int \frac{1}{x+1} dx + \log(x) - \frac{\log(x+1)}{x} \\
 & \quad \downarrow \text{16} \\
 & \log(x) - \frac{\log(x+1)}{x} - \log(x+1)
 \end{aligned}$$

input `Int[Log[1 + x]/x^2,x]`

output `Log[x] - Log[1 + x] - Log[1 + x]/x`

3.249.3.1 Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.)), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

rule 2842 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_.))^(n_.)]*(b_.))*((f_.) + (g_.)*(x_.))^(q_.), x_Symbol] := Simp[(f + g*x)^(q + 1)*((a + b*Log[c*(d + e*x)^n])/(g*(q + 1))), x] - Simp[b*e*(n/(g*(q + 1))) Int[(f + g*x)^(q + 1)/(d + e*x), x], x] /; FreeQ[{a, b, c, d, e, f, g, n, q}, x] && NeQ[e*f - d*g, 0] && NeQ[q, -1]`

3.249.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

method	result	size
derivativedivides	$\ln(x) - \frac{\ln(1+x)(1+x)}{x}$	16
default	$\ln(x) - \frac{\ln(1+x)(1+x)}{x}$	16
meijerg	$\ln(x) - \frac{(2+2x)\ln(1+x)}{2x}$	18
risch	$\ln(x) - \ln(1+x) - \frac{\ln(1+x)}{x}$	19
parts	$\ln(x) - \ln(1+x) - \frac{\ln(1+x)}{x}$	19
norman	$\frac{-\ln(1+x)x - \ln(1+x)}{x} + \ln(x)$	22
parallelrisch	$\frac{x \ln(x) - \ln(1+x)x - \ln(1+x)}{x}$	23

input `int(ln(1+x)/x^2,x,method=_RETURNVERBOSE)`

output `ln(x)-ln(1+x)*(1+x)/x`

3.249.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.06

$$\int \frac{\log(1+x)}{x^2} dx = -\frac{(x+1)\log(x+1) - x\log(x)}{x}$$

input `integrate(log(1+x)/x^2,x, algorithm="fricas")`output `-((x + 1)*log(x + 1) - x*log(x))/x`**3.249.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{\log(1+x)}{x^2} dx = \log(x) - \log(x+1) - \frac{\log(x+1)}{x}$$

input `integrate(ln(1+x)/x**2,x)`output `log(x) - log(x + 1) - log(x + 1)/x`**3.249.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{\log(1+x)}{x^2} dx = -\frac{\log(x+1)}{x} - \log(x+1) + \log(x)$$

input `integrate(log(1+x)/x^2,x, algorithm="maxima")`output `-log(x + 1)/x - log(x + 1) + log(x)`

3.249.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{\log(1+x)}{x^2} dx = -\frac{\log(x+1)}{x} - \log(|x+1|) + \log(|x|)$$

input `integrate(log(1+x)/x^2,x, algorithm="giac")`output `-log(x + 1)/x - log(abs(x + 1)) + log(abs(x))`**3.249.9 Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{\log(1+x)}{x^2} dx = -\ln\left(\frac{1}{x} + 1\right) - \frac{\ln(x+1)}{x}$$

input `int(log(x + 1)/x^2,x)`output `- log(1/x + 1) - log(x + 1)/x`

3.250 $\int \sqrt{1 - \arccos(\sin(x))^2} dx$

3.250.1 Optimal result	1265
3.250.2 Mathematica [A] (verified)	1265
3.250.3 Rubi [F]	1266
3.250.4 Maple [F]	1266
3.250.5 Fricas [A] (verification not implemented)	1267
3.250.6 Sympy [F]	1267
3.250.7 Maxima [A] (verification not implemented)	1267
3.250.8 Giac [F(-1)]	1268
3.250.9 Mupad [F(-1)]	1268

3.250.1 Optimal result

Integrand size = 13, antiderivative size = 56

$$\int \sqrt{1 - \arccos(\sin(x))^2} dx = -\frac{1}{2} \left(\arccos(\sin(x)) \sqrt{1 - \arccos(\sin(x))^2} - 2 \arctan \left(\frac{\sqrt{1 - \arccos(\sin(x))^2}}{1 + \arccos(\sin(x))} \right) \right) \sqrt{\cos^2(x)} \sec(x)$$

output `-1/2*((1/2*Pi-arcsin(sin(x)))*(1-(1/2*Pi-arcsin(sin(x)))^2)^(1/2)-2*arctan((1-(1/2*Pi-arcsin(sin(x)))^2)^(1/2)/(1+1/2*Pi-arcsin(sin(x)))))*(cos(x)^2)^(1/2)*sec(x)`

3.250.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00

$$\int \sqrt{1 - \arccos(\sin(x))^2} dx = -\frac{1}{2} \left(\arccos(\sin(x)) \sqrt{1 - \arccos(\sin(x))^2} - 2 \arctan \left(\frac{\sqrt{1 - \arccos(\sin(x))^2}}{1 + \arccos(\sin(x))} \right) \right) \sqrt{\cos^2(x)} \sec(x)$$

input `Integrate[Sqrt[1 - ArcCos[Sin[x]]^2],x]`

output `-1/2*((ArcCos[Sin[x]]*Sqrt[1 - ArcCos[Sin[x]]^2] - 2*ArcTan[Sqrt[1 - ArcCos[Sin[x]]^2]/(1 + ArcCos[Sin[x]])])*Sqrt[Cos[x]^2]*Sec[x])`

3.250.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{1 - \arccos(\sin(x))^2} dx$$

↓ 7299

$$\int \sqrt{1 - \arccos(\sin(x))^2} dx$$

input `Int[Sqrt[1 - ArcCos[Sin[x]]^2],x]`

output `$Aborted`

3.250.3.1 Defintions of rubi rules used

rule 7299 `Int[u_, x_] :> CannotIntegrate[u, x]`

3.250.4 Maple [F]

$$\int \sqrt{1 - \left(-\frac{\pi}{2} + \arcsin(\sin(x))\right)^2} dx$$

input `int((1-(-1/2*Pi+arcsin(sin(x)))^2)^(1/2),x)`

output `int((1-(-1/2*Pi+arcsin(sin(x)))^2)^(1/2),x)`

3.250.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.23

$$\int \sqrt{1 - \arccos(\sin(x))^2} dx = \frac{1}{8} (\pi + 2x) \sqrt{-\pi^2 - 4\pi x - 4x^2 + 4} - \frac{1}{2} \arctan \left(\frac{(\pi + 2x) \sqrt{-\pi^2 - 4\pi x - 4x^2 + 4}}{\pi^2 + 4\pi x + 4x^2 - 4} \right)$$

input `integrate((1-(-1/2*pi+arcsin(sin(x)))^2)^(1/2),x, algorithm="fricas")`output `1/8*(pi + 2*x)*sqrt(-pi^2 - 4*pi*x - 4*x^2 + 4) - 1/2*arctan((pi + 2*x)*sqrt(-pi^2 - 4*pi*x - 4*x^2 + 4)/(pi^2 + 4*pi*x + 4*x^2 - 4))`**3.250.6 Sympy [F]**

$$\int \sqrt{1 - \arccos(\sin(x))^2} dx = \frac{\int \sqrt{-4 \operatorname{asin}^2(\sin(x)) + 4\pi \operatorname{asin}(\sin(x)) - \pi^2 + 4} dx}{2}$$

input `integrate((1-(-1/2*pi+asin(sin(x)))**2)**(1/2),x)`output `Integral(sqrt(-4*asin(sin(x))**2 + 4*pi*asin(sin(x)) - pi**2 + 4), x)/2`**3.250.7 Maxima [A] (verification not implemented)**

Time = 0.40 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.91

$$\int \sqrt{1 - \arccos(\sin(x))^2} dx = -\frac{1}{8} (\pi - 2x) \sqrt{-\pi^2 + 4\pi x - 4x^2 + 4} + \frac{1}{2} \arctan \left(-\frac{1}{2} \pi + x, \sqrt{-\frac{1}{4} \pi^2 + \pi x - x^2 + 1} \right)$$

input `integrate((1-(-1/2*pi+arcsin(sin(x)))^2)^(1/2),x, algorithm="maxima")`output `-1/8*(pi - 2*x)*sqrt(-pi^2 + 4*pi*x - 4*x^2 + 4) + 1/2*arctan2(-1/2*pi + x, sqrt(-1/4*pi^2 + pi*x - x^2 + 1))`

3.250.8 Giac [F(-1)]

Timed out.

$$\int \sqrt{1 - \arccos(\sin(x))^2} dx = \text{Timed out}$$

input `integrate((1-(-1/2*pi+arcsin(sin(x)))^2)^(1/2),x, algorithm="giac")`

output `Timed out`

3.250.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{1 - \arccos(\sin(x))^2} dx = \int \sqrt{1 - \left(\frac{\Pi}{2} - \text{asin}(\sin(x))\right)^2} dx$$

input `int((1 - (Pi/2 - asin(sin(x)))^2)^(1/2),x)`

output `int((1 - (Pi/2 - asin(sin(x)))^2)^(1/2), x)`

3.251 $\int(-2+x)(-1+x)x(1+x)(2+x) dx$

3.251.1 Optimal result	1269
3.251.2 Mathematica [A] (verified)	1269
3.251.3 Rubi [A] (verified)	1270
3.251.4 Maple [A] (verified)	1271
3.251.5 Fricas [A] (verification not implemented)	1271
3.251.6 Sympy [A] (verification not implemented)	1271
3.251.7 Maxima [A] (verification not implemented)	1272
3.251.8 Giac [A] (verification not implemented)	1272
3.251.9 Mupad [B] (verification not implemented)	1272

3.251.1 Optimal result

Integrand size = 14, antiderivative size = 20

$$\int(-2+x)(-1+x)x(1+x)(2+x) dx = 2x^2 - \frac{5x^4}{4} + \frac{x^6}{6}$$

output `2*x^2-5/4*x^4+1/6*x^6`

3.251.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int(-2+x)(-1+x)x(1+x)(2+x) dx = 2x^2 - \frac{5x^4}{4} + \frac{x^6}{6}$$

input `Integrate[(-2 + x)*(-1 + x)*x*(1 + x)*(2 + x),x]`

output `2*x^2 - (5*x^4)/4 + x^6/6`

3.251.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2109, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x-2)(x-1)x(x+1)(x+2) dx$$

$$\downarrow \text{2109}$$

$$\int (x^5 - 5x^3 + 4x) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^6}{6} - \frac{5x^4}{4} + 2x^2$$

input `Int[(-2 + x)*(-1 + x)*x*(1 + x)*(2 + x), x]`

output `2*x^2 - (5*x^4)/4 + x^6/6`

3.251.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2109 `Int[(Px_)*((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.)*((g_.) + (h_.)*(x_))^(q_.), x_Symbol] := Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n*(e + f*x)^p*(g + h*x)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n, p, q}, x] && PolyQ[Px, x] && IntegersQ[m, n]`

3.251.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

method	result	size
gospers	$2x^2 - \frac{5}{4}x^4 + \frac{1}{6}x^6$	17
default	$2x^2 - \frac{5}{4}x^4 + \frac{1}{6}x^6$	17
norman	$2x^2 - \frac{5}{4}x^4 + \frac{1}{6}x^6$	17
risch	$2x^2 - \frac{5}{4}x^4 + \frac{1}{6}x^6$	17
parallelrisch	$2x^2 - \frac{5}{4}x^4 + \frac{1}{6}x^6$	17

input `int((-2+x)*(-1+x)*x*(1+x)*(2+x),x,method=_RETURNVERBOSE)`output `2*x^2-5/4*x^4+1/6*x^6`**3.251.5 Fracas [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int (-2+x)(-1+x)x(1+x)(2+x) dx = \frac{1}{6}x^6 - \frac{5}{4}x^4 + 2x^2$$

input `integrate((-2+x)*(-1+x)*x*(1+x)*(2+x),x, algorithm="fracas")`output `1/6*x^6 - 5/4*x^4 + 2*x^2`**3.251.6 Sympy [A] (verification not implemented)**

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int (-2+x)(-1+x)x(1+x)(2+x) dx = \frac{x^6}{6} - \frac{5x^4}{4} + 2x^2$$

input `integrate((-2+x)*(-1+x)*x*(1+x)*(2+x),x)`output `x**6/6 - 5*x**4/4 + 2*x**2`

3.251.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int (-2+x)(-1+x)x(1+x)(2+x) dx = \frac{1}{6}x^6 - \frac{5}{4}x^4 + 2x^2$$

input `integrate((-2+x)*(-1+x)*x*(1+x)*(2+x),x, algorithm="maxima")`output `1/6*x^6 - 5/4*x^4 + 2*x^2`**3.251.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int (-2+x)(-1+x)x(1+x)(2+x) dx = \frac{1}{6}x^6 - \frac{5}{4}x^4 + 2x^2$$

input `integrate((-2+x)*(-1+x)*x*(1+x)*(2+x),x, algorithm="giac")`output `1/6*x^6 - 5/4*x^4 + 2*x^2`**3.251.9 Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int (-2+x)(-1+x)x(1+x)(2+x) dx = \frac{x^2(2x^4 - 15x^2 + 24)}{12}$$

input `int(x*(x - 1)*(x + 1)*(x - 2)*(x + 2),x)`output `(x^2*(2*x^4 - 15*x^2 + 24))/12`

$$3.252 \quad \int \frac{1+2x^2+3x^3}{\sqrt[3]{1+x^3}} dx$$

3.252.1 Optimal result	1273
3.252.2 Mathematica [A] (verified)	1273
3.252.3 Rubi [A] (verified)	1274
3.252.4 Maple [A] (verified)	1275
3.252.5 Fricas [A] (verification not implemented)	1275
3.252.6 Sympy [C] (verification not implemented)	1275
3.252.7 Maxima [A] (verification not implemented)	1276
3.252.8 Giac [F]	1276
3.252.9 Mupad [B] (verification not implemented)	1276

3.252.1 Optimal result

Integrand size = 22, antiderivative size = 21

$$\int \frac{1+2x^2+3x^3}{\sqrt[3]{1+x^3}} dx = (1+x^3)^{2/3} + x(1+x^3)^{2/3}$$

output $(x^3+1)^{(2/3)}+x*(x^3+1)^{(2/3)}$

3.252.2 Mathematica [A] (verified)

Time = 10.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \frac{1+2x^2+3x^3}{\sqrt[3]{1+x^3}} dx = (1+x)(1+x^3)^{2/3}$$

input `Integrate[(1 + 2*x^2 + 3*x^3)/(1 + x^3)^(1/3),x]`

output $(1+x)*(1+x^3)^{(2/3)}$

3.252.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {2432, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{3x^3 + 2x^2 + 1}{\sqrt[3]{x^3 + 1}} dx$$

↓ 2432

$$\int \left(\frac{3x^3}{\sqrt[3]{x^3 + 1}} + \frac{1}{\sqrt[3]{x^3 + 1}} + \frac{2x^2}{\sqrt[3]{x^3 + 1}} \right) dx$$

↓ 2009

$$(x^3 + 1)^{2/3} x + (x^3 + 1)^{2/3}$$

input `Int[(1 + 2*x^2 + 3*x^3)/(1 + x^3)^(1/3),x]`

output `(1 + x^3)^(2/3) + x*(1 + x^3)^(2/3)`

3.252.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2432 `Int[(Pq_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Int[ExpandIntegrand[Pq*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, n, p}, x] && (PolyQ[Pq, x] || PolyQ[Pq, x^n])`

3.252.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.57

method	result	size
trager	$(1+x)(x^3+1)^{\frac{2}{3}}$	12
risch	$(1+x)(x^3+1)^{\frac{2}{3}}$	12
gosper	$\frac{(1+x)^2(x^2-x+1)}{(x^3+1)^{\frac{1}{3}}}$	22
meijerg	$x \operatorname{hypergeom}\left(\left[\frac{1}{3}, \frac{1}{3}\right], \left[\frac{4}{3}\right], -x^3\right) + \frac{3x^4 \operatorname{hypergeom}\left(\left[\frac{1}{3}, \frac{4}{3}\right], \left[\frac{7}{3}\right], -x^3\right)}{4} + \frac{2x^3 \operatorname{hypergeom}\left(\left[\frac{1}{3}, 1\right], [2], -x^3\right)}{3}$	47

input `int((3*x^3+2*x^2+1)/(x^3+1)^(1/3),x,method=_RETURNVERBOSE)`output `(1+x)*(x^3+1)^(2/3)`**3.252.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.52

$$\int \frac{1+2x^2+3x^3}{\sqrt[3]{1+x^3}} dx = (x^3+1)^{\frac{2}{3}}(x+1)$$

input `integrate((3*x^3+2*x^2+1)/(x^3+1)^(1/3),x, algorithm="fricas")`output `(x^3 + 1)^(2/3)*(x + 1)`**3.252.6 Sympy [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 1.44 (sec) , antiderivative size = 65, normalized size of antiderivative = 3.10

$$\int \frac{1+2x^2+3x^3}{\sqrt[3]{1+x^3}} dx = \frac{x^4 \Gamma\left(\frac{4}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{4}{3} \middle| \frac{7}{3}; x^3 e^{i\pi}\right)}{\Gamma\left(\frac{7}{3}\right)} + \frac{x \Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{1}{3} \middle| \frac{4}{3}; x^3 e^{i\pi}\right)}{3 \Gamma\left(\frac{4}{3}\right)} + (x^3+1)^{\frac{2}{3}}$$

3.252. $\int \frac{1+2x^2+3x^3}{\sqrt[3]{1+x^3}} dx$

input `integrate((3*x**3+2*x**2+1)/(x**3+1)**(1/3),x)`

output `x**4*gamma(4/3)*hyper((1/3, 4/3), (7/3,), x**3*exp_polar(I*pi))/gamma(7/3) + x*gamma(1/3)*hyper((1/3, 1/3), (4/3,), x**3*exp_polar(I*pi))/(3*gamma(4/3)) + (x**3 + 1)**(2/3)`

3.252.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.52

$$\int \frac{1 + 2x^2 + 3x^3}{\sqrt[3]{1 + x^3}} dx = (x^3 + 1)^{\frac{2}{3}} + \frac{(x^3 + 1)^{\frac{2}{3}}}{x^2 \left(\frac{x^3 + 1}{x^3} - 1 \right)}$$

input `integrate((3*x^3+2*x^2+1)/(x^3+1)^(1/3),x, algorithm="maxima")`

output `(x^3 + 1)^(2/3) + (x^3 + 1)^(2/3)/(x^2*((x^3 + 1)/x^3 - 1))`

3.252.8 Giac [F]

$$\int \frac{1 + 2x^2 + 3x^3}{\sqrt[3]{1 + x^3}} dx = \int \frac{3x^3 + 2x^2 + 1}{(x^3 + 1)^{\frac{1}{3}}} dx$$

input `integrate((3*x^3+2*x^2+1)/(x^3+1)^(1/3),x, algorithm="giac")`

output `integrate((3*x^3 + 2*x^2 + 1)/(x^3 + 1)^(1/3), x)`

3.252.9 Mupad [B] (verification not implemented)

Time = 14.71 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.52

$$\int \frac{1 + 2x^2 + 3x^3}{\sqrt[3]{1 + x^3}} dx = (x^3 + 1)^{2/3} (x + 1)$$

input `int((2*x^2 + 3*x^3 + 1)/(x^3 + 1)^(1/3),x)`

output `(x^3 + 1)^(2/3)*(x + 1)`

3.252. $\int \frac{1+2x^2+3x^3}{\sqrt[3]{1+x^3}} dx$

3.253 $\int \csc^4(x) \sec^4(x) dx$

3.253.1 Optimal result	1277
3.253.2 Mathematica [A] (verified)	1277
3.253.3 Rubi [A] (verified)	1278
3.253.4 Maple [A] (verified)	1279
3.253.5 Fricas [B] (verification not implemented)	1279
3.253.6 Sympy [A] (verification not implemented)	1280
3.253.7 Maxima [A] (verification not implemented)	1280
3.253.8 Giac [A] (verification not implemented)	1280
3.253.9 Mupad [B] (verification not implemented)	1281

3.253.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \csc^4(x) \sec^4(x) dx = -8 \cot(2x) - \frac{8}{3} \cot^3(2x)$$

output `-8*cot(2*x)-8/3*cot(2*x)^3`

3.253.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.94

$$\int \csc^4(x) \sec^4(x) dx = -\frac{8 \cot(x)}{3} - \frac{1}{3} \cot(x) \csc^2(x) + \frac{8 \tan(x)}{3} + \frac{1}{3} \sec^2(x) \tan(x)$$

input `Integrate[Csc[x]^4*Sec[x]^4,x]`

output `(-8*Cot[x])/3 - (Cot[x]*Csc[x]^2)/3 + (8*Tan[x])/3 + (Sec[x]^2*Tan[x])/3`

3.253.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.47, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 3100, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \csc^4(x) \sec^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc(x)^4 \sec(x)^4 dx \\
 & \quad \downarrow \text{3100} \\
 & \int (\tan^2(x) + 1)^3 \cot^4(x) d \tan(x) \\
 & \quad \downarrow \text{244} \\
 & \int (\tan^2(x) + \cot^4(x) + 3 \cot^2(x) + 3) d \tan(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\tan^3(x)}{3} + 3 \tan(x) - \frac{1}{3} \cot^3(x) - 3 \cot(x)
 \end{aligned}$$

input `Int[Csc[x]^4*Sec[x]^4,x]`

output `-3*Cot[x] - Cot[x]^3/3 + 3*Tan[x] + Tan[x]^3/3`

3.253.3.1 Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3100 `Int[csc[(e_.) + (f_.)*(x_.)]^(m_.)*sec[(e_.) + (f_.)*(x_.)]^(n_.), x_Symbol] := Simp[1/f Subst[Int[(1 + x^2)^((m + n)/2 - 1)/x^m, x], x, Tan[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n)/2]`

3.253.4 Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.41

method	result	size
parallelrisch	$-\frac{(-\cos(6x)+3\cos(2x))\sec(x)^3\csc(x)^3}{6}$	24
risch	$\frac{32i(3e^{4ix}-1)}{3(e^{2ix}-1)^3(e^{2ix}+1)^3}$	31
default	$\frac{1}{3\sin(x)^3\cos(x)^3} - \frac{2}{3\sin(x)^3\cos(x)} + \frac{8}{3\cos(x)\sin(x)} - \frac{16\cot(x)}{3}$	36
norman	$\frac{\frac{1}{24} + \frac{5\tan(\frac{x}{2})^2}{4} - \frac{91\tan(\frac{x}{2})^4}{8} + \frac{35\tan(\frac{x}{2})^6}{2} - \frac{91\tan(\frac{x}{2})^8}{8} + \frac{5\tan(\frac{x}{2})^{10}}{4} + \frac{\tan(\frac{x}{2})^{12}}{24}}{\tan(\frac{x}{2})^3(\tan(\frac{x}{2})^2-1)^3}$	68

input `int(1/sin(x)^4/cos(x)^4,x,method=_RETURNVERBOSE)`

output `-1/6*(-cos(6*x)+3*cos(2*x))*sec(x)^3*csc(x)^3`

3.253.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. $2(15) = 30$.

Time = 0.24 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.29

$$\int \csc^4(x) \sec^4(x) dx = -\frac{16 \cos(x)^6 - 24 \cos(x)^4 + 6 \cos(x)^2 + 1}{3(\cos(x)^5 - \cos(x)^3) \sin(x)}$$

input `integrate(1/sin(x)^4/cos(x)^4,x, algorithm="fracas")`

output `-1/3*(16*cos(x)^6 - 24*cos(x)^4 + 6*cos(x)^2 + 1)/((cos(x)^5 - cos(x)^3)*sin(x))`

3.253.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.71

$$\int \csc^4(x) \sec^4(x) dx = -\frac{16 \cos(2x)}{3 \sin(2x)} - \frac{8 \cos(2x)}{3 \sin^3(2x)}$$

input `integrate(1/sin(x)**4/cos(x)**4,x)`output `-16*cos(2*x)/(3*sin(2*x)) - 8*cos(2*x)/(3*sin(2*x)**3)`**3.253.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.47

$$\int \csc^4(x) \sec^4(x) dx = \frac{1}{3} \tan(x)^3 - \frac{9 \tan(x)^2 + 1}{3 \tan(x)^3} + 3 \tan(x)$$

input `integrate(1/sin(x)^4/cos(x)^4,x, algorithm="maxima")`output `1/3*tan(x)^3 - 1/3*(9*tan(x)^2 + 1)/tan(x)^3 + 3*tan(x)`**3.253.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

$$\int \csc^4(x) \sec^4(x) dx = -\frac{8(3 \tan(2x)^2 + 1)}{3 \tan(2x)^3}$$

input `integrate(1/sin(x)^4/cos(x)^4,x, algorithm="giac")`output `-8/3*(3*tan(2*x)^2 + 1)/tan(2*x)^3`

3.253.9 Mupad [B] (verification not implemented)

Time = 15.19 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.24

$$\int \csc^4(x) \sec^4(x) dx = -\frac{24 \cos(2x) - 16 \cos(2x)^3}{3 \sin(2x) - 3 \cos(2x)^2 \sin(2x)}$$

input `int(1/(cos(x)^4*sin(x)^4),x)`

output `-(24*cos(2*x) - 16*cos(2*x)^3)/(3*sin(2*x) - 3*cos(2*x)^2*sin(2*x))`

3.254 $\int \frac{x+\sin(x)}{1+\cos(x)} dx$

3.254.1 Optimal result	1282
3.254.2 Mathematica [A] (verified)	1282
3.254.3 Rubi [B] (verified)	1283
3.254.4 Maple [A] (verified)	1285
3.254.5 Fricas [A] (verification not implemented)	1285
3.254.6 Sympy [A] (verification not implemented)	1286
3.254.7 Maxima [B] (verification not implemented)	1286
3.254.8 Giac [A] (verification not implemented)	1286
3.254.9 Mupad [B] (verification not implemented)	1287

3.254.1 Optimal result

Integrand size = 11, antiderivative size = 8

$$\int \frac{x + \sin(x)}{1 + \cos(x)} dx = x \tan\left(\frac{x}{2}\right)$$

output `x*tan(1/2*x)`

3.254.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{x + \sin(x)}{1 + \cos(x)} dx = x \tan\left(\frac{x}{2}\right)$$

input `Integrate[(x + Sin[x])/(1 + Cos[x]), x]`

output `x*Tan[x/2]`

3.254.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 31 vs. $2(8) = 16$.

Time = 0.38 (sec) , antiderivative size = 31, normalized size of antiderivative = 3.88, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.909$, Rules used = {4877, 3042, 3146, 16, 3799, 3042, 4672, 25, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x + \sin(x)}{\cos(x) + 1} dx \\
 & \quad \downarrow \text{4877} \\
 & \int \frac{x}{\cos(x) + 1} dx + \int \frac{\sin(x)}{\cos(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{x}{\sin(x + \frac{\pi}{2}) + 1} dx + \int \frac{\cos(x - \frac{\pi}{2})}{1 - \sin(x - \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3146} \\
 & \int \frac{x}{\sin(x + \frac{\pi}{2}) + 1} dx - \int \frac{1}{\cos(x) + 1} d \cos(x) \\
 & \quad \downarrow \text{16} \\
 & \int \frac{x}{\sin(x + \frac{\pi}{2}) + 1} dx - \log(\cos(x) + 1) \\
 & \quad \downarrow \text{3799} \\
 & \frac{1}{2} \int x \sec^2\left(\frac{x}{2}\right) dx - \log(\cos(x) + 1) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int x \csc\left(\frac{x}{2} + \frac{\pi}{2}\right)^2 dx - \log(\cos(x) + 1) \\
 & \quad \downarrow \text{4672} \\
 & \frac{1}{2} \left(2 \int -\tan\left(\frac{x}{2}\right) dx + 2x \tan\left(\frac{x}{2}\right) \right) - \log(\cos(x) + 1) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{2} \left(2x \tan\left(\frac{x}{2}\right) - 2 \int \tan\left(\frac{x}{2}\right) dx \right) - \log(\cos(x) + 1)
 \end{aligned}$$

$$\begin{array}{c} \downarrow \text{3042} \\ \frac{1}{2} \left(2x \tan \left(\frac{x}{2} \right) - 2 \int \tan \left(\frac{x}{2} \right) dx \right) - \log(\cos(x) + 1) \\ \downarrow \text{3956} \\ \frac{1}{2} \left(2x \tan \left(\frac{x}{2} \right) + 4 \log \left(\cos \left(\frac{x}{2} \right) \right) \right) - \log(\cos(x) + 1) \end{array}$$

input `Int[(x + Sin[x])/(1 + Cos[x]),x]`

output `-Log[1 + Cos[x]] + (4*Log[Cos[x/2]] + 2*x*Tan[x/2])/2`

3.254.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3146 `Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1)/2, x], x, b*Sin[e + f*x], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])]`

rule 3799 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(2*a)^n Int[(c + d*x)^m*Sin[(1/2)*(e + Pi*(a/(2*b))) + f*(x/2)]^(2*n), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 - b^2, 0] && IntegerQ[n] && (GtQ[n, 0] || IGtQ[m, 0])`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4672 `Int[csc[(e_.) + (f_.)*(x_.)]^2*((c_.) + (d_.)*(x_.))^(m_.), x_Symbol] := Simp [(-c + d*x)^m*(Cot[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1) *Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 4877 `Int[(u_)*((v_) + (d_.)*(F_)[(c_.)*((a_.) + (b_.)*(x_.))]^(n_.)), x_Symbol] :> With[{e = FreeFactors[Cos[c*(a + b*x)], x]}, Int[ActivateTrig[u*v], x] + Simp[d Int[ActivateTrig[u]*Sin[c*(a + b*x)]^n, x], x] /; FunctionOfQ[Cos[c*(a + b*x)]/e, u, x] /; FreeQ[{a, b, c, d}, x] && !FreeQ[v, x] && IntegerQ[(n - 1)/2] && NonsumQ[u] && (EqQ[F, Sin] || EqQ[F, sin])`

3.254.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
lookup	$x \tan\left(\frac{x}{2}\right)$	7
default	$x \tan\left(\frac{x}{2}\right)$	7
norman	$x \tan\left(\frac{x}{2}\right)$	7
paralletrisch	$x \tan\left(\frac{x}{2}\right)$	7
risch	$-ix + \frac{2ix}{e^{ix}+1}$	19

input `int((x+sin(x))/(1+cos(x)),x,method=_RETURNVERBOSE)`

output `x*tan(1/2*x)`

3.254.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.25

$$\int \frac{x + \sin(x)}{1 + \cos(x)} dx = \frac{x \sin(x)}{\cos(x) + 1}$$

input `integrate((x+sin(x))/(1+cos(x)),x, algorithm="fricas")`

output `x*sin(x)/(cos(x) + 1)`

3.254. $\int \frac{x + \sin(x)}{1 + \cos(x)} dx$

3.254.6 Sympy [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \frac{x + \sin(x)}{1 + \cos(x)} dx = x \tan\left(\frac{x}{2}\right)$$

input `integrate((x+sin(x))/(1+cos(x)),x)`

output `x*tan(x/2)`

3.254.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs. $2(6) = 12$.

Time = 0.19 (sec) , antiderivative size = 61, normalized size of antiderivative = 7.62

$$\begin{aligned} & \int \frac{x + \sin(x)}{1 + \cos(x)} dx \\ &= \frac{(\cos(x)^2 + \sin(x)^2 + 2 \cos(x) + 1) \log(\cos(x)^2 + \sin(x)^2 + 2 \cos(x) + 1) + 2x \sin(x)}{\cos(x)^2 + \sin(x)^2 + 2 \cos(x) + 1} \\ & \quad - \log(\cos(x) + 1) \end{aligned}$$

input `integrate((x+sin(x))/(1+cos(x)),x, algorithm="maxima")`

output `((cos(x)^2 + sin(x)^2 + 2*cos(x) + 1)*log(cos(x)^2 + sin(x)^2 + 2*cos(x) + 1) + 2*x*sin(x))/(cos(x)^2 + sin(x)^2 + 2*cos(x) + 1) - log(cos(x) + 1)`

3.254.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{x + \sin(x)}{1 + \cos(x)} dx = x \tan\left(\frac{1}{2}x\right)$$

input `integrate((x+sin(x))/(1+cos(x)),x, algorithm="giac")`

output `x*tan(1/2*x)`

3.254.9 Mupad [B] (verification not implemented)

Time = 14.73 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{x + \sin(x)}{1 + \cos(x)} dx = x \tan\left(\frac{x}{2}\right)$$

input `int((x + sin(x))/(cos(x) + 1),x)`

output `x*tan(x/2)`

3.255 $\int \cosh^2(x) \sinh^3(x) dx$

3.255.1 Optimal result	1288
3.255.2 Mathematica [A] (verified)	1288
3.255.3 Rubi [A] (verified)	1289
3.255.4 Maple [A] (verified)	1290
3.255.5 Fricas [B] (verification not implemented)	1291
3.255.6 Sympy [A] (verification not implemented)	1291
3.255.7 Maxima [B] (verification not implemented)	1291
3.255.8 Giac [B] (verification not implemented)	1292
3.255.9 Mupad [B] (verification not implemented)	1292

3.255.1 Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \cosh^2(x) \sinh^3(x) dx = -\frac{1}{3} \cosh^3(x) + \frac{\cosh^5(x)}{5}$$

output `-1/3*cosh(x)^3+1/5*cosh(x)^5`

3.255.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.35

$$\int \cosh^2(x) \sinh^3(x) dx = -\frac{\cosh(x)}{8} - \frac{1}{48} \cosh(3x) + \frac{1}{80} \cosh(5x)$$

input `Integrate[Cosh[x]^2*Sinh[x]^3,x]`

output `-1/8*Cosh[x] - Cosh[3*x]/48 + Cosh[5*x]/80`

3.255.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 26, 3045, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh^3(x) \cosh^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int i \sin(ix)^3 \cos(ix)^2 dx \\
 & \quad \downarrow \text{26} \\
 & i \int \cos(ix)^2 \sin(ix)^3 dx \\
 & \quad \downarrow \text{3045} \\
 & - \int \cosh^2(x) (1 - \cosh^2(x)) d \cosh(x) \\
 & \quad \downarrow \text{244} \\
 & - \int (\cosh^2(x) - \cosh^4(x)) d \cosh(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\cosh^5(x)}{5} - \frac{\cosh^3(x)}{3}
 \end{aligned}$$

input `Int[Cosh[x]^2*Sinh[x]^3,x]`

output `-1/3*Cosh[x]^3 + Cosh[x]^5/5`

3.255.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 244 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3045 `Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Simp[-(a*f)^(-1) Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])`

3.255.4 Maple [A] (verified)

Time = 9.81 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$-\frac{\cosh(x)^3}{3} + \frac{\cosh(x)^5}{5}$	14
default	$-\frac{\cosh(x)^3}{3} + \frac{\cosh(x)^5}{5}$	14
risch	$\frac{e^{5x}}{160} - \frac{e^{3x}}{96} - \frac{e^x}{16} - \frac{e^{-x}}{16} - \frac{e^{-3x}}{96} + \frac{e^{-5x}}{160}$	36

input `int(sinh(x)^3*cosh(x)^2,x,method=_RETURNVERBOSE)`

output `-1/3*cosh(x)^3+1/5*cosh(x)^5`

3.255.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. $2(13) = 26$.

Time = 0.24 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.47

$$\int \cosh^2(x) \sinh^3(x) dx = \frac{1}{80} \cosh(x)^5 + \frac{1}{16} \cosh(x) \sinh(x)^4 - \frac{1}{48} \cosh(x)^3 + \frac{1}{16} (2 \cosh(x)^3 - \cosh(x)) \sinh(x)^2 - \frac{1}{8} \cosh(x)$$

input `integrate(sinh(x)^3*cosh(x)^2,x, algorithm="fricas")`

output `1/80*cosh(x)^5 + 1/16*cosh(x)*sinh(x)^4 - 1/48*cosh(x)^3 + 1/16*(2*cosh(x)^3 - cosh(x))*sinh(x)^2 - 1/8*cosh(x)`

3.255.6 Sympy [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \cosh^2(x) \sinh^3(x) dx = \frac{\sinh^2(x) \cosh^3(x)}{3} - \frac{2 \cosh^5(x)}{15}$$

input `integrate(sinh(x)**3*cosh(x)**2,x)`

output `sinh(x)**2*cosh(x)**3/3 - 2*cosh(x)**5/15`

3.255.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. $2(13) = 26$.

Time = 0.20 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.29

$$\int \cosh^2(x) \sinh^3(x) dx = -\frac{1}{480} (5 e^{(-2x)} + 30 e^{(-4x)} - 3) e^{(5x)} - \frac{1}{16} e^{(-x)} - \frac{1}{96} e^{(-3x)} + \frac{1}{160} e^{(-5x)}$$

input `integrate(sinh(x)^3*cosh(x)^2,x, algorithm="maxima")`

output `-1/480*(5*e^(-2*x) + 30*e^(-4*x) - 3)*e^(5*x) - 1/16*e^(-x) - 1/96*e^(-3*x) + 1/160*e^(-5*x)`

3.255.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. $2(13) = 26$.

Time = 0.27 (sec) , antiderivative size = 37, normalized size of antiderivative = 2.18

$$\int \cosh^2(x) \sinh^3(x) dx = -\frac{1}{480} (30 e^{(4x)} + 5 e^{(2x)} - 3) e^{(-5x)} + \frac{1}{160} e^{(5x)} - \frac{1}{96} e^{(3x)} - \frac{1}{16} e^x$$

input `integrate(sinh(x)^3*cosh(x)^2,x, algorithm="giac")`

output `-1/480*(30*e^(4*x) + 5*e^(2*x) - 3)*e^(-5*x) + 1/160*e^(5*x) - 1/96*e^(3*x) - 1/16*e^x`

3.255.9 Mupad [B] (verification not implemented)

Time = 14.88 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \cosh^2(x) \sinh^3(x) dx = \frac{\cosh(x)^3 (3 \cosh(x)^2 - 5)}{15}$$

input `int(cosh(x)^2*sinh(x)^3,x)`

output `(cosh(x)^3*(3*cosh(x)^2 - 5))/15`

3.256 $\int 3^{2^x} 4^x dx$

3.256.1 Optimal result	1293
3.256.2 Mathematica [A] (verified)	1293
3.256.3 Rubi [A] (verified)	1294
3.256.4 Maple [A] (verified)	1295
3.256.5 Fricas [A] (verification not implemented)	1295
3.256.6 Sympy [A] (verification not implemented)	1296
3.256.7 Maxima [C] (verification not implemented)	1296
3.256.8 Giac [A] (verification not implemented)	1296
3.256.9 Mupad [B] (verification not implemented)	1297

3.256.1 Optimal result

Integrand size = 9, antiderivative size = 33

$$\int 3^{2^x} 4^x dx = -\frac{3^{2^x}}{\log(2) \log^2(3)} + \frac{2^x 3^{2^x}}{\log(2) \log(3)}$$

output `-3^(2^x)/ln(2)/ln(3)^2+2^x*3^(2^x)/ln(2)/ln(3)`

3.256.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.67

$$\int 3^{2^x} 4^x dx = \frac{3^{2^x}(-1 + 2^x \log(3))}{\log(2) \log^2(3)}$$

input `Integrate[3^2^x*4^x,x]`

output `(3^2^x*(-1 + 2^x*Log[3]))/(Log[2]*Log[3]^2)`

3.256.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.91, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2720, 2607, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int 3^{2^x} 4^x dx \\ & \quad \downarrow \text{2720} \\ & \frac{\int 2^x 3^{2^x} d2^x}{\log(2)} \\ & \quad \downarrow \text{2607} \\ & \frac{\frac{2^x 3^{2^x}}{\log(3)} - \frac{\int 3^{2^x} d2^x}{\log(3)}}{\log(2)} \\ & \quad \downarrow \text{2624} \\ & \frac{\frac{2^x 3^{2^x}}{\log(3)} - \frac{3^{2^x}}{\log^2(3)}}{\log(2)} \end{aligned}$$

input `Int[3^2^x*4^x,x]`

output `(-(3^2^x/Log[3]^2) + (2^x*3^2^x)/Log[3])/Log[2]`

3.256.3.1 Defintions of rubi rules used

rule 2607 `Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Simp[d*(m/(f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]`

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /; FreeQ[{F, n}, x] && LinearQ[v, x]`

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

3.256.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.70

method	result	size
risch	$\frac{(2^x \ln(3) - 1)3^{2x}}{\ln(2) \ln(3)^2}$	23
norman	$\frac{e^{x \ln(2)} e^{e^x \ln(2) \ln(3)}}{\ln(2) \ln(3)} - \frac{e^{e^x \ln(2) \ln(3)}}{\ln(2) \ln(3)^2}$	44

```
input int(4^x*3^(2^x),x,method=_RETURNVERBOSE)
```

```
output (2^x*ln(3)-1)/ln(2)/ln(3)^2*3^(2^x)
```

3.256.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.67

$$\int 3^{2^x} 4^x dx = \frac{(2^x \log(3) - 1)3^{(2^x)}}{\log(3)^2 \log(2)}$$

```
input integrate(4^x*3^(2^x),x, algorithm="fracas")
```

```
output (2^x*log(3) - 1)*3^(2^x)/(log(3)^2*log(2))
```


3.256.6 Sympy [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.03

$$\int 3^{2^x} 4^x dx = \frac{\left(e^{\frac{x \log(4)}{2}} \log(3) - 1\right) e^{e^{\frac{x \log(4)}{2}} \log(3)}}{\log(2) \log(3)^2}$$

input `integrate(4**x*3**(2**x),x)`

output `(exp(x*log(4)/2)*log(3) - 1)*exp(exp(x*log(4)/2)*log(3))/(log(2)*log(3)**2)`

3.256.7 Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 0.31 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.94

$$\int 3^{2^x} 4^x dx = -\frac{4^x \Gamma\left(2, -4^{\frac{1}{2}x} \log(3)\right)}{4^x \log(3)^2 \log(2)}$$

input `integrate(4^x*3^(2^x),x, algorithm="maxima")`

output `-4^x*gamma(2, -4^(1/2*x)*log(3))/(4^x*log(3)^2*log(2))`

3.256.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.55

$$\int 3^{2^x} 4^x dx = \frac{2^x e^{(2^x \log(3) + 2^x \log(2))} \log(3) - e^{(2^x \log(3) + 2^x \log(2))}}{2^{2^x} \log(3)^2 \log(2)}$$

input `integrate(4^x*3^(2^x),x, algorithm="giac")`

output `(2^x*e^(2^x*log(3) + 2*x*log(2))*log(3) - e^(2^x*log(3) + 2*x*log(2)))/(2^(2*x)*log(3)^2*log(2))`

3.256.9 Mupad [B] (verification not implemented)

Time = 15.86 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.67

$$\int 3^{2^x} 4^x dx = \frac{3^{2^x} (2^x \ln(3) - 1)}{\ln(2) \ln(3)^2}$$

input `int(3^(2^x)*4^x,x)`

output `(3^(2^x)*(2^x*log(3) - 1))/(log(2)*log(3)^2)`

$$3.257 \quad \int \frac{\cos(x) - \sin(x)}{2 + \sin(2x)} dx$$

3.257.1 Optimal result	1298
3.257.2 Mathematica [A] (verified)	1298
3.257.3 Rubi [C] (verified)	1299
3.257.4 Maple [C] (verified)	1300
3.257.5 Fricas [B] (verification not implemented)	1301
3.257.6 Sympy [F]	1301
3.257.7 Maxima [F]	1301
3.257.8 Giac [B] (verification not implemented)	1302
3.257.9 Mupad [B] (verification not implemented)	1302

3.257.1 Optimal result

Integrand size = 16, antiderivative size = 6

$$\int \frac{\cos(x) - \sin(x)}{2 + \sin(2x)} dx = \arctan(\cos(x) + \sin(x))$$

output `arctan(cos(x)+sin(x))`

3.257.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x) - \sin(x)}{2 + \sin(2x)} dx = \arctan(\cos(x) + \sin(x))$$

input `Integrate[(Cos[x] - Sin[x])/(2 + Sin[2*x]),x]`

output `ArcTan[Cos[x] + Sin[x]]`

3.257.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.92 (sec) , antiderivative size = 217, normalized size of antiderivative = 36.17, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3042, 4901, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(x) - \sin(x)}{\sin(2x) + 2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\cos(x) - \sin(x)}{\sin(2x) + 2} dx \\ & \quad \downarrow \text{4901} \\ & \int \left(\frac{\cos(x)}{\sin(2x) + 2} - \frac{\sin(x)}{\sin(2x) + 2} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{1}{6} (3 + i\sqrt{3}) \arctan \left(\frac{-2i \tan(\frac{x}{2}) - \sqrt{3} + i}{\sqrt{2}(1 + i\sqrt{3})} \right) + \frac{1}{6} (3 - i\sqrt{3}) \arctan \left(\frac{-2i \tan(\frac{x}{2}) - \sqrt{3} + i}{\sqrt{2}(1 + i\sqrt{3})} \right) - \\ & \frac{1}{6} (3 + i\sqrt{3}) \arctan \left(\frac{-2i \tan(\frac{x}{2}) + \sqrt{3} + i}{\sqrt{2}(1 - i\sqrt{3})} \right) - \frac{1}{6} (3 - i\sqrt{3}) \arctan \left(\frac{-2i \tan(\frac{x}{2}) + \sqrt{3} + i}{\sqrt{2}(1 - i\sqrt{3})} \right) \end{aligned}$$

input `Int[(Cos[x] - Sin[x])/(2 + Sin[2*x]),x]`

output `((3 - I*Sqrt[3])*ArcTan[(I - Sqrt[3] - (2*I)*Tan[x/2])/Sqrt[2*(1 + I*Sqrt[3])]])/6 + ((3 + I*Sqrt[3])*ArcTan[(I - Sqrt[3] - (2*I)*Tan[x/2])/Sqrt[2*(1 + I*Sqrt[3])]])/6 - ((3 - I*Sqrt[3])*ArcTan[(I + Sqrt[3] - (2*I)*Tan[x/2])/Sqrt[2*(1 - I*Sqrt[3])]])/6 - ((3 + I*Sqrt[3])*ArcTan[(I + Sqrt[3] - (2*I)*Tan[x/2])/Sqrt[2*(1 - I*Sqrt[3])]])/6`

3.257.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4901 `Int[u_, x_Symbol] := With[{v = ExpandTrig[u, x]}, Int[v, x] /; SumQ[v]] /; !InertTrigFreeQ[u]`

3.257.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.49 (sec) , antiderivative size = 44, normalized size of antiderivative = 7.33

method	result
risch	$-\frac{i \ln(e^{2ix} + (1-i)e^{ix} + i)}{2} + \frac{i \ln(e^{2ix} + (-1+i)e^{ix} + i)}{2}$
default	$\left(\frac{\sum_{R=\text{RootOf}(_Z^4 - 2_Z^3 + 2_Z^2 + 2_Z + 1)} \frac{(-R^2 + 1) \ln(\tan(\frac{x}{2}) - R)}{2R^3 - 3R^2 + 2R + 1}}{2} \right) - \left(\frac{\sum_{R=\text{RootOf}(_Z^4 - 2_Z^3 + 2_Z^2 + 2_Z + 1)} \frac{(-R^2 + 1) \ln(\tan(\frac{x}{2}) - R)}{2R^3 - 3R^2 + 2R + 1}}{2} \right)$
parts	$\left(\frac{\sum_{R=\text{RootOf}(_Z^4 - 2_Z^3 + 2_Z^2 + 2_Z + 1)} \frac{(-R^2 + 1) \ln(\tan(\frac{x}{2}) - R)}{2R^3 - 3R^2 + 2R + 1}}{2} \right) - \left(\frac{\sum_{R=\text{RootOf}(_Z^4 - 2_Z^3 + 2_Z^2 + 2_Z + 1)} \frac{(-R^2 + 1) \ln(\tan(\frac{x}{2}) - R)}{2R^3 - 3R^2 + 2R + 1}}{2} \right)$

input `int((cos(x)-sin(x))/(2+sin(2*x)),x,method=_RETURNVERBOSE)`

output `-1/2*I*ln(exp(2*I*x)+(1-I)*exp(I*x)+I)+1/2*I*ln(exp(2*I*x)+(-1+I)*exp(I*x)+I)`

3.257.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(6) = 12$.

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 2.50

$$\int \frac{\cos(x) - \sin(x)}{2 + \sin(2x)} dx = \frac{1}{2} \arctan \left(\frac{\cos(x) \sin(x)}{\cos(x) + \sin(x)} \right)$$

input `integrate((cos(x)-sin(x))/(2+sin(2*x)),x, algorithm="fricas")`

output `1/2*arctan(cos(x)*sin(x)/(cos(x) + sin(x)))`

3.257.6 Sympy [F]

$$\int \frac{\cos(x) - \sin(x)}{2 + \sin(2x)} dx = \int \frac{-\sin(x) + \cos(x)}{\sin(2x) + 2} dx$$

input `integrate((cos(x)-sin(x))/(2+sin(2*x)),x)`

output `Integral((-sin(x) + cos(x))/(sin(2*x) + 2), x)`

3.257.7 Maxima [F]

$$\int \frac{\cos(x) - \sin(x)}{2 + \sin(2x)} dx = \int \frac{\cos(x) - \sin(x)}{\sin(2x) + 2} dx$$

input `integrate((cos(x)-sin(x))/(2+sin(2*x)),x, algorithm="maxima")`

output `integrate((cos(x) - sin(x))/(sin(2*x) + 2), x)`

3.257.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 35 vs. $2(6) = 12$.

Time = 0.28 (sec) , antiderivative size = 35, normalized size of antiderivative = 5.83

$$\int \frac{\cos(x) - \sin(x)}{2 + \sin(2x)} dx = \arctan\left(\frac{1}{2} \tan\left(\frac{1}{2}x\right)^3 - \frac{3}{2} \tan\left(\frac{1}{2}x\right)^2 + \frac{3}{2} \tan\left(\frac{1}{2}x\right) + \frac{1}{2}\right) - \arctan\left(\tan\left(\frac{1}{2}x\right) - 2\right)$$

input `integrate((cos(x)-sin(x))/(2+sin(2*x)),x, algorithm="giac")`

output `arctan(1/2*tan(1/2*x)^3 - 3/2*tan(1/2*x)^2 + 3/2*tan(1/2*x) + 1/2) - arctan(tan(1/2*x) - 2)`

3.257.9 Mupad [B] (verification not implemented)

Time = 16.00 (sec) , antiderivative size = 35, normalized size of antiderivative = 5.83

$$\int \frac{\cos(x) - \sin(x)}{2 + \sin(2x)} dx = \operatorname{atan}\left(\frac{\tan\left(\frac{x}{2}\right)^3}{2} - \frac{3 \tan\left(\frac{x}{2}\right)^2}{2} + \frac{3 \tan\left(\frac{x}{2}\right)}{2} + \frac{1}{2}\right) - \operatorname{atan}\left(\tan\left(\frac{x}{2}\right) - 2\right)$$

input `int((cos(x) - sin(x))/(sin(2*x) + 2),x)`

output `atan((3*tan(x/2))/2 - (3*tan(x/2)^2)/2 + tan(x/2)^3/2 + 1/2) - atan(tan(x/2) - 2)`

$$3.258 \quad \int \frac{\sec^2(1+\log(x)) - \tan(1+\log(x))}{x^2} dx$$

3.258.1 Optimal result	1303
3.258.2 Mathematica [B] (verified)	1303
3.258.3 Rubi [C] (verified)	1304
3.258.4 Maple [C] (verified)	1305
3.258.5 Fracas [A] (verification not implemented)	1305
3.258.6 Sympy [F]	1305
3.258.7 Maxima [B] (verification not implemented)	1306
3.258.8 Giac [B] (verification not implemented)	1306
3.258.9 Mupad [F(-1)]	1307

3.258.1 Optimal result

Integrand size = 19, antiderivative size = 9

$$\int \frac{\sec^2(1 + \log(x)) - \tan(1 + \log(x))}{x^2} dx = \frac{\tan(1 + \log(x))}{x}$$

output `tan(ln(x)+1)/x`

3.258.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 21 vs. $2(9) = 18$.

Time = 0.88 (sec) , antiderivative size = 21, normalized size of antiderivative = 2.33

$$\int \frac{\sec^2(1 + \log(x)) - \tan(1 + \log(x))}{x^2} dx = \frac{\sec(1) \sec(1 + \log(x)) \sin(\log(x))}{x} + \frac{\tan(1)}{x}$$

input `Integrate[(Sec[1 + Log[x]]^2 - Tan[1 + Log[x]])/x^2,x]`

output `(Sec[1]*Sec[1 + Log[x]]*Sin[Log[x]])/x + Tan[1]/x`

3.258.3 Rubi [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.22 (sec) , antiderivative size = 81, normalized size of antiderivative = 9.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2010, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^2(\log(x) + 1) - \tan(\log(x) + 1)}{x^2} dx$$

↓ 2010

$$\int \left(\frac{\sec^2(\log(x) + 1)}{x^2} - \frac{\tan(\log(x) + 1)}{x^2} \right) dx$$

↓ 2009

$$\frac{2i \operatorname{Hypergeometric2F1}\left(\frac{i}{2}, 1, 1 + \frac{i}{2}, -e^{2i} x^{2i}\right)}{x} - \left(\frac{4}{5} + \frac{8i}{5}\right) e^{2i} x^{-1+2i} \operatorname{Hypergeometric2F1}\left(1 + \frac{i}{2}, 2, 2 + \frac{i}{2}, -e^{2i} x^{2i}\right) - \frac{i}{x}$$

input `Int[(Sec[1 + Log[x]]^2 - Tan[1 + Log[x]])/x^2,x]`

output `(-I)/x + ((2*I)*Hypergeometric2F1[I/2, 1, 1 + I/2, -(E^(2*I))*x^(2*I)])/x - ((4/5 + (8*I)/5)*E^(2*I)*Hypergeometric2F1[1 + I/2, 2, 2 + I/2, -(E^(2*I))*x^(2*I)])/x^(1 - 2*I)`

3.258.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2010 `Int[(u_)*((c_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]`

3.258.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.58 (sec) , antiderivative size = 28, normalized size of antiderivative = 3.11

method	result	size
risch	$-\frac{i}{x} + \frac{2i}{x(x^{2i}e^{2i}+1)}$	28

input `int((sec(ln(x)+1)^2-tan(ln(x)+1))/x^2,x,method=_RETURNVERBOSE)`

output `-I/x+2*I/x/((x^I)^2*exp(2*I)+1)`

3.258.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.78

$$\int \frac{\sec^2(1 + \log(x)) - \tan(1 + \log(x))}{x^2} dx = \frac{\sin(\log(x) + 1)}{x \cos(\log(x) + 1)}$$

input `integrate((sec(log(x)+1)^2-tan(log(x)+1))/x^2,x, algorithm="fricas")`

output `sin(log(x) + 1)/(x*cos(log(x) + 1))`

3.258.6 Sympy [F]

$$\int \frac{\sec^2(1 + \log(x)) - \tan(1 + \log(x))}{x^2} dx = \int \frac{-\tan(\log(x) + 1) + \sec^2(\log(x) + 1)}{x^2} dx$$

input `integrate((sec(ln(x)+1)**2-tan(ln(x)+1))/x**2,x)`

output `Integral((-tan(log(x) + 1) + sec(log(x) + 1)**2)/x**2, x)`

3.258.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 45 vs. $2(9) = 18$.

Time = 0.28 (sec) , antiderivative size = 45, normalized size of antiderivative = 5.00

$$\int \frac{\sec^2(1 + \log(x)) - \tan(1 + \log(x))}{x^2} dx$$

$$= \frac{2 \sin(2 \log(x) + 2)}{x \cos(2 \log(x) + 2)^2 + x \sin(2 \log(x) + 2)^2 + 2x \cos(2 \log(x) + 2) + x}$$

input `integrate((sec(log(x)+1)^2-tan(log(x)+1))/x^2,x, algorithm="maxima")`

output `2*sin(2*log(x) + 2)/(x*cos(2*log(x) + 2)^2 + x*sin(2*log(x) + 2)^2 + 2*x*cos(2*log(x) + 2) + x)`

3.258.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5161 vs. $2(9) = 18$.

Time = 1.84 (sec) , antiderivative size = 5161, normalized size of antiderivative = 573.44

$$\int \frac{\sec^2(1 + \log(x)) - \tan(1 + \log(x))}{x^2} dx = \text{Too large to display}$$

input `integrate((sec(log(x)+1)^2-tan(log(x)+1))/x^2,x, algorithm="giac")`

output $1/4*(13*\tan(1)^4*\tan(\log(x))^9/((\tan(\log(x))^2 - 1)*x) + 42*\tan(1)^4*\tan(\log(x))^10/((\tan(\log(x))^2 - 1)^2*x) + 84*\tan(1)^4*\tan(\log(x))^11/((\tan(\log(x))^2 - 1)^3*x) - 88*\tan(1)^4*\tan(\log(x))^12/((\tan(\log(x))^2 - 1)^4*x) - 17*\tan(1)^4*\tan(\log(x))^7/x - 4*\tan(1)^3*\tan(\log(x))^8/x - 62*\tan(1)^4*\tan(\log(x))^8/((\tan(\log(x))^2 - 1)*x) + 22*\tan(1)^3*\tan(\log(x))^9/((\tan(\log(x))^2 - 1)*x) - 102*\tan(1)^4*\tan(\log(x))^9/((\tan(\log(x))^2 - 1)^2*x) - 6*\tan(1)^3*\tan(\log(x))^10/((\tan(\log(x))^2 - 1)^2*x) + 520*\tan(1)^4*\tan(\log(x))^10/((\tan(\log(x))^2 - 1)^3*x) - 168*\tan(1)^3*\tan(\log(x))^11/((\tan(\log(x))^2 - 1)^3*x) + 248*\tan(1)^4*\tan(\log(x))^11/((\tan(\log(x))^2 - 1)^4*x) - 88*\tan(1)^3*\tan(\log(x))^12/((\tan(\log(x))^2 - 1)^4*x) + 20*\tan(1)^4*\tan(\log(x))^6/x - 22*\tan(1)^3*\tan(\log(x))^7/x - 17*\tan(1)^4*\tan(\log(x))^7/((\tan(\log(x))^2 - 1)*x) - 70*\tan(1)^3*\tan(\log(x))^8/((\tan(\log(x))^2 - 1)*x) - 740*\tan(1)^4*\tan(\log(x))^8/((\tan(\log(x))^2 - 1)^2*x) + 17*\tan(1)^2*\tan(\log(x))^9/((\tan(\log(x))^2 - 1)*x) + 488*\tan(1)^3*\tan(\log(x))^9/((\tan(\log(x))^2 - 1)^2*x) - 548*\tan(1)^4*\tan(\log(x))^9/((\tan(\log(x))^2 - 1)^3*x) - 56*\tan(1)^2*\tan(\log(x))^10/((\tan(\log(x))^2 - 1)^2*x) + 40*\tan(1)^3*\tan(\log(x))^10/((\tan(\log(x))^2 - 1)^3*x) + 560*\tan(1)^4*\tan(\log(x))^10/((\tan(\log(x))^2 - 1)^4*x) + 36*\tan(1)^2*\tan(\log(x))^11/((\tan(\log(x))^2 - 1)^3*x) - 256*\tan(1)^3*\tan(\log(x))^11/((\tan(\log(x))^2 - 1)^4*x) + 32*\tan(1)^2*\tan(\log(x))^12/((\tan(\log(x))^2 - 1)^4*x) - 22*\tan(1)^4*\tan(\log(x))^5/x + 36*\tan(1)^3*\tan(1...$

3.258.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^2(1 + \log(x)) - \tan(1 + \log(x))}{x^2} dx = \int -\frac{\tan(\ln(x) + 1) - \frac{1}{\cos(\ln(x)+1)^2}}{x^2} dx$$

input `int(-(tan(log(x) + 1) - 1/cos(log(x) + 1)^2)/x^2,x)`

output `int(-(tan(log(x) + 1) - 1/cos(log(x) + 1)^2)/x^2, x)`

3.259 $\int \sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}} dx$

3.259.1 Optimal result 1308
 3.259.2 Mathematica [A] (verified) 1308
 3.259.3 Rubi [A] (verified) 1309
 3.259.4 Maple [F] 1310
 3.259.5 Fricas [F(-2)] 1311
 3.259.6 Sympy [F] 1311
 3.259.7 Maxima [F] 1311
 3.259.8 Giac [F] 1312
 3.259.9 Mupad [F(-1)] 1312

3.259.1 Optimal result

Integrand size = 12, antiderivative size = 67

$$\int \sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}} dx = 2x\sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}} - \frac{\sqrt{2\pi}\operatorname{erf}\left(\frac{\sqrt{\log\left(\frac{1}{x}\right)}}{\sqrt{2}}\right)\sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}}}{\sqrt{\frac{1}{x}}\sqrt{\log\left(\frac{1}{x}\right)}}$$

output `2*x*(ln(1/x)/x)^(1/2)-2^(1/2)*Pi^(1/2)*erf(1/2*ln(1/x)^(1/2)*2^(1/2))*(ln(1/x)/x)^(1/2)/(1/x)^(1/2)/ln(1/x)^(1/2)`

3.259.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.84

$$\int \sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}} dx = \left(2x - \frac{\sqrt{2\pi}\operatorname{erf}\left(\frac{\sqrt{\log\left(\frac{1}{x}\right)}}{\sqrt{2}}\right)}{\sqrt{\frac{1}{x}}\sqrt{\log\left(\frac{1}{x}\right)}}\right)\sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}}$$

input `Integrate[Sqrt[Log[x^(-1)]/x],x]`

output `(2*x - (Sqrt[2*Pi]*Erf[Sqrt[Log[x^(-1)]/Sqrt[2]]])/(Sqrt[x^(-1)]*Sqrt[Log[x^(-1)]]))*Sqrt[Log[x^(-1)]/x]`

3.259. $\int \sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}} dx$

3.259.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.16, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {7270, 2742, 2747, 2611, 2634}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}} dx \\
 & \quad \downarrow \text{7270} \\
 & \frac{\sqrt{x} \sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}} \int \frac{\sqrt{\log\left(\frac{1}{x}\right)}}{\sqrt{x}} dx}{\sqrt{\log\left(\frac{1}{x}\right)}} \\
 & \quad \downarrow \text{2742} \\
 & \frac{\sqrt{x} \sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}} \left(\int \frac{1}{\sqrt{x} \sqrt{\log\left(\frac{1}{x}\right)}} dx + 2\sqrt{x} \sqrt{\log\left(\frac{1}{x}\right)} \right)}{\sqrt{\log\left(\frac{1}{x}\right)}} \\
 & \quad \downarrow \text{2747} \\
 & \frac{\sqrt{x} \sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}} \left(2\sqrt{x} \sqrt{\log\left(\frac{1}{x}\right)} - \sqrt{\frac{1}{x}} \sqrt{x} \int \frac{1}{\sqrt{\frac{1}{x}} \sqrt{\log\left(\frac{1}{x}\right)}} d\log\left(\frac{1}{x}\right) \right)}{\sqrt{\log\left(\frac{1}{x}\right)}} \\
 & \quad \downarrow \text{2611} \\
 & \frac{\sqrt{x} \sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}} \left(2\sqrt{x} \sqrt{\log\left(\frac{1}{x}\right)} - 2\sqrt{\frac{1}{x}} \sqrt{x} \int \frac{1}{\sqrt{\frac{1}{x}}} d\sqrt{\log\left(\frac{1}{x}\right)} \right)}{\sqrt{\log\left(\frac{1}{x}\right)}} \\
 & \quad \downarrow \text{2634} \\
 & \frac{\sqrt{x} \sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}} \left(2\sqrt{x} \sqrt{\log\left(\frac{1}{x}\right)} - \sqrt{2\pi} \sqrt{\frac{1}{x}} \sqrt{x} \operatorname{erf}\left(\frac{\sqrt{\log\left(\frac{1}{x}\right)}}{\sqrt{2}}\right) \right)}{\sqrt{\log\left(\frac{1}{x}\right)}}
 \end{aligned}$$

input `Int[Sqrt[Log[x^(-1)]/x], x]`

3.259. $\int \sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}} dx$

output $(\sqrt{x} * (-\sqrt{2\pi} * \sqrt{x^{-1}} * \sqrt{x} * \operatorname{Erf}[\sqrt{\log[x^{-1}]}] / \sqrt{2}] + 2 * \sqrt{x} * \sqrt{\log[x^{-1}]}) * \sqrt{\log[x^{-1}] / x} / \sqrt{\log[x^{-1}]}$

3.259.3.1 Defintions of rubi rules used

rule 2611 $\operatorname{Int}[(F_)^{(g_)*((e_)+(f_)*(x_))}/\sqrt{(c_)+(d_)*(x_)}, x_Symbol] :> \operatorname{Simp}[2/d \operatorname{Subst}[\operatorname{Int}[F^{(g*(e-c*(f/d))+f*g*(x^2/d))}, x], x, \sqrt{c+dx}], x] /;$ $\operatorname{FreeQ}\{F, c, d, e, f, g\}, x\} \&\& \operatorname{!TrueQ}\{\$UseGamma\}$

rule 2634 $\operatorname{Int}[(F_)^{(a_)+(b_)*((c_)+(d_)*(x_))^2}, x_Symbol] :> \operatorname{Simp}[F^a * \sqrt{\pi} * (\operatorname{Erf}[(c+dx)*\operatorname{Rt}[-b]*\log[F], 2]) / (2*d*\operatorname{Rt}[-b]*\log[F], 2)), x] /;$ $\operatorname{FreeQ}\{F, a, b, c, d\}, x\} \&\& \operatorname{NegQ}[b]$

rule 2742 $\operatorname{Int}[(a_)+\log[(c_)*(x_)^{(n_)}]*(b_)]^{(p_)*((d_)*(x_))^m}, x_Symbol] :> \operatorname{Simp}[(d*x)^{(m+1)}*((a+b*\log[c*x^n])^p/(d*(m+1))), x] - \operatorname{Simp}[b*n*(p/(m+1)) \operatorname{Int}[(d*x)^m*(a+b*\log[c*x^n])^{(p-1)}, x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, m, n\}, x\} \&\& \operatorname{NeQ}[m, -1] \&\& \operatorname{GtQ}[p, 0]$

rule 2747 $\operatorname{Int}[(a_)+\log[(c_)*(x_)^{(n_)}]*(b_)]^{(p_)*((d_)*(x_))^m}, x_Symbol] :> \operatorname{Simp}[(d*x)^{(m+1)}/(d*n*(c*x^n)^{((m+1)/n)}) \operatorname{Subst}[\operatorname{Int}[E^{((m+1)/n)*x}*(a+b*x)^p, x], x, \log[c*x^n]], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, m, n, p\}, x\}$

rule 7270 $\operatorname{Int}[(u_)*((a_)*(v_)^{(m_)}*(w_)^{(n_)})^p], x_Symbol] :> \operatorname{Simp}[a^{\operatorname{IntPart}[p]}*((a*v^m*w^n)^{\operatorname{FracPart}[p]}/(v^{(m*\operatorname{FracPart}[p])}*w^{(n*\operatorname{FracPart}[p])})) \operatorname{Int}[u*v^{(m*p)}*w^{(n*p)}, x], x] /;$ $\operatorname{FreeQ}\{a, m, n, p\}, x\} \&\& \operatorname{!IntegerQ}[p] \&\& \operatorname{!FreeQ}[v, x] \&\& \operatorname{!FreeQ}[w, x]$

3.259.4 Maple [F]

$$\int \sqrt{\frac{\ln(\frac{1}{x})}{x}} dx$$

input $\operatorname{int}((\ln(1/x)/x)^{(1/2)}, x)$

output $\operatorname{int}((\ln(1/x)/x)^{(1/2)}, x)$

3.259. $\int \sqrt{\frac{\log(\frac{1}{x})}{x}} dx$

3.259.5 Fracas [F(-2)]

Exception generated.

$$\int \sqrt{\frac{\log(\frac{1}{x})}{x}} dx = \text{Exception raised: TypeError}$$

input `integrate((log(1/x)/x)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

3.259.6 Sympy [F]

$$\int \sqrt{\frac{\log(\frac{1}{x})}{x}} dx = \int \sqrt{\frac{\log(\frac{1}{x})}{x}} dx$$

input `integrate((ln(1/x)/x)**(1/2),x)`

output `Integral(sqrt(log(1/x)/x), x)`

3.259.7 Maxima [F]

$$\int \sqrt{\frac{\log(\frac{1}{x})}{x}} dx = \int \sqrt{\frac{\log(\frac{1}{x})}{x}} dx$$

input `integrate((log(1/x)/x)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(-log(x)/x), x)`

3.259.8 Giac [F]

$$\int \sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}} dx = \int \sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}} dx$$

input `integrate((log(1/x)/x)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(log(1/x)/x), x)`

3.259.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\frac{\log\left(\frac{1}{x}\right)}{x}} dx = \int \sqrt{\frac{\ln\left(\frac{1}{x}\right)}{x}} dx$$

input `int((log(1/x)/x)^(1/2),x)`

output `int((log(1/x)/x)^(1/2), x)`

3.260 $\int (1 - \cos(x))^5 \cos(5x) dx$

3.260.1 Optimal result	1313
3.260.2 Mathematica [A] (verified)	1313
3.260.3 Rubi [A] (verified)	1314
3.260.4 Maple [A] (verified)	1315
3.260.5 Fricas [A] (verification not implemented)	1315
3.260.6 Sympy [B] (verification not implemented)	1316
3.260.7 Maxima [A] (verification not implemented)	1317
3.260.8 Giac [A] (verification not implemented)	1317
3.260.9 Mupad [B] (verification not implemented)	1318

3.260.1 Optimal result

Integrand size = 13, antiderivative size = 84

$$\int (1 - \cos(x))^5 \cos(5x) dx = -\frac{x}{32} + \frac{5 \sin(x)}{16} - \frac{45}{64} \sin(2x) + \frac{5}{4} \sin(3x) - \frac{105}{64} \sin(4x) + \frac{63}{40} \sin(5x) - \frac{35}{32} \sin(6x) + \frac{15}{28} \sin(7x) - \frac{45}{256} \sin(8x) + \frac{5}{144} \sin(9x) - \frac{1}{320} \sin(10x)$$

output `-1/32*x+5/16*sin(x)-45/64*sin(2*x)+5/4*sin(3*x)-105/64*sin(4*x)+63/40*sin(5*x)-35/32*sin(6*x)+15/28*sin(7*x)-45/256*sin(8*x)+5/144*sin(9*x)-1/320*sin(10*x)`

3.260.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.00

$$\int (1 - \cos(x))^5 \cos(5x) dx = -\frac{x}{32} + \frac{5 \sin(x)}{16} - \frac{45}{64} \sin(2x) + \frac{5}{4} \sin(3x) - \frac{105}{64} \sin(4x) + \frac{63}{40} \sin(5x) - \frac{35}{32} \sin(6x) + \frac{15}{28} \sin(7x) - \frac{45}{256} \sin(8x) + \frac{5}{144} \sin(9x) - \frac{1}{320} \sin(10x)$$

input `Integrate[(1 - Cos[x])^5*Cos[5*x], x]`

output
$$-1/32*x + (5*\text{Sin}[x])/16 - (45*\text{Sin}[2*x])/64 + (5*\text{Sin}[3*x])/4 - (105*\text{Sin}[4*x])/64 + (63*\text{Sin}[5*x])/40 - (35*\text{Sin}[6*x])/32 + (15*\text{Sin}[7*x])/28 - (45*\text{Sin}[8*x])/256 + (5*\text{Sin}[9*x])/144 - \text{Sin}[10*x]/320$$

3.260.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3042, 4901, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (1 - \cos(x))^5 \cos(5x) dx \\ & \quad \downarrow 3042 \\ & \int (1 - \cos(x))^5 \cos(5x) dx \\ & \quad \downarrow 4901 \\ & \int (-\cos(5x) \cos^5(x) + 5 \cos(5x) \cos^4(x) - 10 \cos(5x) \cos^3(x) + 10 \cos(5x) \cos^2(x) - 5 \cos(5x) \cos(x) + \cos(5x)) dx \\ & \quad \downarrow 2009 \\ & -\frac{x}{32} + \frac{5 \sin(x)}{16} - \frac{45}{64} \sin(2x) + \frac{5}{4} \sin(3x) - \frac{105}{64} \sin(4x) + \frac{63}{40} \sin(5x) - \frac{35}{32} \sin(6x) + \frac{15}{28} \sin(7x) - \\ & \quad \frac{45}{256} \sin(8x) + \frac{5}{144} \sin(9x) - \frac{1}{320} \sin(10x) \end{aligned}$$

input $\text{Int}[(1 - \text{Cos}[x])^5 * \text{Cos}[5*x], x]$

output
$$-1/32*x + (5*\text{Sin}[x])/16 - (45*\text{Sin}[2*x])/64 + (5*\text{Sin}[3*x])/4 - (105*\text{Sin}[4*x])/64 + (63*\text{Sin}[5*x])/40 - (35*\text{Sin}[6*x])/32 + (15*\text{Sin}[7*x])/28 - (45*\text{Sin}[8*x])/256 + (5*\text{Sin}[9*x])/144 - \text{Sin}[10*x]/320$$

3.260.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4901 `Int[u_, x_Symbol] := With[{v = ExpandTrig[u, x]}, Int[v, x] /; SumQ[v]] /; !InertTrigFreeQ[u]`

3.260.4 Maple [A] (verified)

Time = 1.30 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.75

method	result
default	$-\frac{x}{32} + \frac{5 \sin(x)}{16} - \frac{45 \sin(2x)}{64} + \frac{5 \sin(3x)}{4} - \frac{105 \sin(4x)}{64} + \frac{63 \sin(5x)}{40} - \frac{35 \sin(6x)}{32} + \frac{15 \sin(7x)}{28} - \frac{45 \sin(8x)}{256} + \frac{5 \sin(9x)}{144} - \frac{1}{320} \sin(10x)$
risch	$-\frac{x}{32} + \frac{5 \sin(x)}{16} - \frac{45 \sin(2x)}{64} + \frac{5 \sin(3x)}{4} - \frac{105 \sin(4x)}{64} + \frac{63 \sin(5x)}{40} - \frac{35 \sin(6x)}{32} + \frac{15 \sin(7x)}{28} - \frac{45 \sin(8x)}{256} + \frac{5 \sin(9x)}{144} - \frac{1}{320} \sin(10x)$
parallelrisch	$-\frac{x}{32} + \frac{5 \sin(x)}{16} - \frac{45 \sin(2x)}{64} + \frac{5 \sin(3x)}{4} - \frac{105 \sin(4x)}{64} + \frac{63 \sin(5x)}{40} - \frac{35 \sin(6x)}{32} + \frac{15 \sin(7x)}{28} - \frac{45 \sin(8x)}{256} + \frac{5 \sin(9x)}{144} - \frac{1}{320} \sin(10x)$
parts	$-\frac{x}{32} + \frac{5 \sin(x)}{16} - \frac{45 \sin(2x)}{64} + \frac{5 \sin(3x)}{4} - \frac{105 \sin(4x)}{64} + \frac{63 \sin(5x)}{40} - \frac{35 \sin(6x)}{32} + \frac{15 \sin(7x)}{28} - \frac{45 \sin(8x)}{256} + \frac{5 \sin(9x)}{144} - \frac{1}{320} \sin(10x)$

input `int((1-cos(x))^5*cos(5*x),x,method=_RETURNVERBOSE)`

output `-1/32*x+5/16*sin(x)-45/64*sin(2*x)+5/4*sin(3*x)-105/64*sin(4*x)+63/40*sin(5*x)-35/32*sin(6*x)+15/28*sin(7*x)-45/256*sin(8*x)+5/144*sin(9*x)-1/320*sin(10*x)`

3.260.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.74

$$\int (1 - \cos(x))^5 \cos(5x) dx =$$

$$-\frac{1}{10080} (16128 \cos(x)^9 - 89600 \cos(x)^8 + 194544 \cos(x)^7 - 188800 \cos(x)^6 + 33768 \cos(x)^5 + 93984 \cos(x)^4 - 14400 \cos(x)^3 + 1440 \cos(x)^2 - 144 \cos(x) + 144) - \frac{1}{32} x$$

3.260. $\int (1 - \cos(x))^5 \cos(5x) dx$

input `integrate((1-cos(x))^5*cos(5*x),x, algorithm="fricas")`

output `-1/10080*(16128*cos(x)^9 - 89600*cos(x)^8 + 194544*cos(x)^7 - 188800*cos(x)^6 + 33768*cos(x)^5 + 93984*cos(x)^4 - 83790*cos(x)^3 + 24512*cos(x)^2 + 315*cos(x) - 1376)*sin(x) - 1/32*x`

3.260.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 394 vs. $2(83) = 166$.

Time = 4.27 (sec) , antiderivative size = 394, normalized size of antiderivative = 4.69

$$\begin{aligned} \int (1 - \cos(x))^5 \cos(5x) dx = & -\frac{x \sin^5(x) \sin(5x)}{32} - \frac{5x \sin^4(x) \cos(x) \cos(5x)}{32} \\ & + \frac{5x \sin^3(x) \sin(5x) \cos^2(x)}{16} + \frac{5x \sin^2(x) \cos^3(x) \cos(5x)}{16} \\ & - \frac{5x \sin(x) \sin(5x) \cos^4(x)}{32} - \frac{x \cos^5(x) \cos(5x)}{32} \\ & - \frac{\sin^5(x) \cos(5x)}{64} + \frac{3 \sin^4(x) \sin(5x) \cos(x)}{64} \\ & + \frac{8 \sin^4(x) \sin(5x)}{63} + \frac{40 \sin^3(x) \cos(x) \cos(5x)}{63} \\ & - \frac{5 \sin^3(x) \cos(5x)}{32} + \frac{\sin^2(x) \sin(5x) \cos^3(x)}{6} \\ & - \frac{4 \sin^2(x) \sin(5x) \cos^2(x)}{21} + \frac{25 \sin^2(x) \sin(5x) \cos(x)}{32} \\ & - \frac{4 \sin^2(x) \sin(5x)}{21} + \frac{55 \sin(x) \cos^4(x) \cos(5x)}{192} \\ & - \frac{100 \sin(x) \cos^3(x) \cos(5x)}{63} + \frac{55 \sin(x) \cos^2(x) \cos(5x)}{32} \\ & - \frac{20 \sin(x) \cos(x) \cos(5x)}{21} + \frac{5 \sin(x) \cos(5x)}{24} \\ & - \frac{241 \sin(5x) \cos^5(x)}{960} + \frac{83 \sin(5x) \cos^4(x)}{63} \\ & - \frac{75 \sin(5x) \cos^3(x)}{32} + \frac{46 \sin(5x) \cos^2(x)}{21} \\ & - \frac{25 \sin(5x) \cos(x)}{24} + \frac{\sin(5x)}{5} \end{aligned}$$

input `integrate((1-cos(x))**5*cos(5*x),x)`

output

```
-x*sin(x)**5*sin(5*x)/32 - 5*x*sin(x)**4*cos(x)*cos(5*x)/32 + 5*x*sin(x)**3*sin(5*x)*cos(x)**2/16 + 5*x*sin(x)**2*cos(x)**3*cos(5*x)/16 - 5*x*sin(x)**sin(5*x)*cos(x)**4/32 - x*cos(x)**5*cos(5*x)/32 - sin(x)**5*cos(5*x)/64 + 3*sin(x)**4*sin(5*x)*cos(x)/64 + 8*sin(x)**4*sin(5*x)/63 + 40*sin(x)**3*cos(x)*cos(5*x)/63 - 5*sin(x)**3*cos(5*x)/32 + sin(x)**2*sin(5*x)*cos(x)**3/6 - 4*sin(x)**2*sin(5*x)*cos(x)**2/3 + 25*sin(x)**2*sin(5*x)*cos(x)/32 - 4*sin(x)**2*sin(5*x)/21 + 55*sin(x)*cos(x)**4*cos(5*x)/192 - 100*sin(x)*cos(x)**3*cos(5*x)/63 + 55*sin(x)*cos(x)**2*cos(5*x)/32 - 20*sin(x)*cos(x)*cos(5*x)/21 + 5*sin(x)*cos(5*x)/24 - 241*sin(5*x)*cos(x)**5/960 + 83*sin(5*x)*cos(x)**4/63 - 75*sin(5*x)*cos(x)**3/32 + 46*sin(5*x)*cos(x)**2/21 - 25*sin(5*x)*cos(x)/24 + sin(5*x)/5
```

3.260.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.79

$$\int (1 - \cos(x))^5 \cos(5x) dx = \frac{80}{9} \sin(x)^9 - \frac{380}{7} \sin(x)^7 - \frac{1}{20} \sin(2x)^5 + \frac{501}{5} \sin(x)^5 + \frac{71}{16} \sin(2x)^3 - \frac{212}{3} \sin(x)^3 - \frac{1}{32} x - \frac{45}{256} \sin(8x) - \frac{105}{64} \sin(4x) - 4 \sin(2x) + 16 \sin(x)$$

input `integrate((1-cos(x))^5*cos(5*x),x, algorithm="maxima")`

output `80/9*sin(x)^9 - 380/7*sin(x)^7 - 1/20*sin(2*x)^5 + 501/5*sin(x)^5 + 71/16*sin(2*x)^3 - 212/3*sin(x)^3 - 1/32*x - 45/256*sin(8*x) - 105/64*sin(4*x) - 4*sin(2*x) + 16*sin(x)`

3.260.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.74

$$\int (1 - \cos(x))^5 \cos(5x) dx = -\frac{1}{32} x - \frac{1}{320} \sin(10x) + \frac{5}{144} \sin(9x) - \frac{45}{256} \sin(8x) + \frac{15}{28} \sin(7x) - \frac{35}{32} \sin(6x) + \frac{63}{40} \sin(5x) - \frac{105}{64} \sin(4x) + \frac{5}{4} \sin(3x) - \frac{45}{64} \sin(2x) + \frac{5}{16} \sin(x)$$

input `integrate((1-cos(x))^5*cos(5*x),x, algorithm="giac")`

output `-1/32*x - 1/320*sin(10*x) + 5/144*sin(9*x) - 45/256*sin(8*x) + 15/28*sin(7*x) - 35/32*sin(6*x) + 63/40*sin(5*x) - 105/64*sin(4*x) + 5/4*sin(3*x) - 45/64*sin(2*x) + 5/16*sin(x)`

3.260.9 Mupad [B] (verification not implemented)

Time = 16.54 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.93

$$\begin{aligned} \int (1 - \cos(x))^5 \cos(5x) dx = & -\frac{8 \sin(x) \cos(x)^9}{5} + \frac{80 \sin(x) \cos(x)^8}{9} - \frac{193 \sin(x) \cos(x)^7}{10} \\ & + \frac{1180 \sin(x) \cos(x)^6}{63} - \frac{67 \sin(x) \cos(x)^5}{20} \\ & - \frac{979 \sin(x) \cos(x)^4}{105} + \frac{133 \sin(x) \cos(x)^3}{16} \\ & - \frac{766 \sin(x) \cos(x)^2}{315} - \frac{\sin(x) \cos(x)}{32} - \frac{x}{32} + \frac{43 \sin(x)}{315} \end{aligned}$$

input `int(-cos(5*x)*(cos(x) - 1)^5,x)`

output `(43*sin(x))/315 - x/32 - (cos(x)*sin(x))/32 - (766*cos(x)^2*sin(x))/315 + (133*cos(x)^3*sin(x))/16 - (979*cos(x)^4*sin(x))/105 - (67*cos(x)^5*sin(x))/20 + (1180*cos(x)^6*sin(x))/63 - (193*cos(x)^7*sin(x))/10 + (80*cos(x)^8*sin(x))/9 - (8*cos(x)^9*sin(x))/5`

3.261 $\int \frac{3 \cos(x)+4 \sin(x)}{4 \cos(x)+3 \sin(x)} dx$

3.261.1 Optimal result 1319
 3.261.2 Mathematica [A] (verified) 1319
 3.261.3 Rubi [A] (verified) 1320
 3.261.4 Maple [C] (verified) 1321
 3.261.5 Fricas [A] (verification not implemented) 1321
 3.261.6 Sympy [A] (verification not implemented) 1322
 3.261.7 Maxima [B] (verification not implemented) 1322
 3.261.8 Giac [A] (verification not implemented) 1322
 3.261.9 Mupad [B] (verification not implemented) 1323

3.261.1 Optimal result

Integrand size = 21, antiderivative size = 20

$$\int \frac{3 \cos(x) + 4 \sin(x)}{4 \cos(x) + 3 \sin(x)} dx = \frac{24x}{25} - \frac{7}{25} \log(4 \cos(x) + 3 \sin(x))$$

output `24/25*x-7/25*ln(3*sin(x)+4*cos(x))`

3.261.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{3 \cos(x) + 4 \sin(x)}{4 \cos(x) + 3 \sin(x)} dx = \frac{24x}{25} - \frac{7}{25} \log(4 \cos(x) + 3 \sin(x))$$

input `Integrate[(3*Cos[x] + 4*Sin[x])/(4*Cos[x] + 3*Sin[x]),x]`

output `(24*x)/25 - (7*Log[4*Cos[x] + 3*Sin[x]])/25`

3.261.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {3042, 3612}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{4 \sin(x) + 3 \cos(x)}{3 \sin(x) + 4 \cos(x)} dx$$

↓ 3042

$$\int \frac{4 \sin(x) + 3 \cos(x)}{3 \sin(x) + 4 \cos(x)} dx$$

↓ 3612

$$\frac{24x}{25} - \frac{7}{25} \log(3 \sin(x) + 4 \cos(x))$$

input `Int[(3*Cos[x] + 4*Sin[x])/(4*Cos[x] + 3*Sin[x]),x]`

output `(24*x)/25 - (7*Log[4*Cos[x] + 3*Sin[x]])/25`

3.261.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3612 `Int[((A_.) + cos[(d_.) + (e_.)*(x_)])*(B_.) + (C_.)*sin[(d_.) + (e_.)*(x_)]) / ((a_.) + cos[(d_.) + (e_.)*(x_)])*(b_.) + (c_.)*sin[(d_.) + (e_.)*(x_)]) , x_Symbol] :> Simp[(b*B + c*C)*(x/(b^2 + c^2)), x] + Simp[(c*B - b*C)*(Log[a + b*Cos[d + e*x] + c*Sin[d + e*x]]/(e*(b^2 + c^2))), x] /; FreeQ[{a, b, c, d, e, A, B, C}, x] && NeQ[b^2 + c^2, 0] && EqQ[A*(b^2 + c^2) - a*(b*B + c*C), 0]`

3.261.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.38 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.05

method	result	size
risch	$\frac{24x}{25} + \frac{7ix}{25} - \frac{7 \ln(e^{2ix} + \frac{7}{25} + \frac{24i}{25})}{25}$	21
default	$\frac{7 \ln(1 + \tan(x)^2)}{50} + \frac{24 \arctan(\tan(x))}{25} - \frac{7 \ln(3 \tan(x) + 4)}{25}$	25
parallelrisch	$\ln\left(\frac{2}{(-2048(\cot(x) - \csc(x) + 2)^7)^{\frac{1}{25}}}\right) + \ln\left(\frac{1}{(-2 \cot(x) + 2 \csc(x) + 1)^{\frac{7}{25}}}\right) + \ln\left(\left(\frac{1}{1 + \cos(x)}\right)^{\frac{7}{25}}\right) + \frac{24x}{25}$	44
norman	$\frac{\frac{24x}{25} + \frac{24x \tan(\frac{x}{2})^2}{25}}{1 + \tan(\frac{x}{2})^2} - \frac{7 \ln(\tan(\frac{x}{2}) - 2)}{25} - \frac{7 \ln(2 \tan(\frac{x}{2}) + 1)}{25} + \frac{7 \ln(1 + \tan(\frac{x}{2})^2)}{25}$	57

input `int((4*sin(x)+3*cos(x))/(3*sin(x)+4*cos(x)),x,method=_RETURNVERBOSE)`

output `24/25*x+7/25*I*x-7/25*ln(exp(2*I*x)+7/25+24/25*I)`

3.261.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int \frac{3 \cos(x) + 4 \sin(x)}{4 \cos(x) + 3 \sin(x)} dx = \frac{24}{25} x - \frac{7}{25} \log\left(-2 \cos(x) - \frac{3}{2} \sin(x)\right)$$

input `integrate((4*sin(x)+3*cos(x))/(3*sin(x)+4*cos(x)),x, algorithm="fricas")`

output `24/25*x - 7/25*log(-2*cos(x) - 3/2*sin(x))`

3.261.6 Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{3 \cos(x) + 4 \sin(x)}{4 \cos(x) + 3 \sin(x)} dx = \frac{24x}{25} - \frac{7 \log(3 \sin(x) + 4 \cos(x))}{25}$$

input `integrate((4*sin(x)+3*cos(x))/(3*sin(x)+4*cos(x)),x)`

output `24*x/25 - 7*log(3*sin(x) + 4*cos(x))/25`

3.261.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 58 vs. $2(16) = 32$.

Time = 0.29 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.90

$$\begin{aligned} \int \frac{3 \cos(x) + 4 \sin(x)}{4 \cos(x) + 3 \sin(x)} dx = & \frac{48}{25} \arctan\left(\frac{\sin(x)}{\cos(x) + 1}\right) - \frac{7}{25} \log\left(\frac{2 \sin(x)}{\cos(x) + 1} + 1\right) \\ & - \frac{7}{25} \log\left(\frac{\sin(x)}{\cos(x) + 1} - 2\right) + \frac{7}{25} \log\left(\frac{\sin(x)^2}{(\cos(x) + 1)^2} + 1\right) \end{aligned}$$

input `integrate((4*sin(x)+3*cos(x))/(3*sin(x)+4*cos(x)),x, algorithm="maxima")`

output `48/25*arctan(sin(x)/(cos(x) + 1)) - 7/25*log(2*sin(x)/(cos(x) + 1) + 1) - 7/25*log(sin(x)/(cos(x) + 1) - 2) + 7/25*log(sin(x)^2/(cos(x) + 1)^2 + 1)`

3.261.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.15

$$\int \frac{3 \cos(x) + 4 \sin(x)}{4 \cos(x) + 3 \sin(x)} dx = \frac{24}{25} x + \frac{7}{50} \log(\tan(x)^2 + 1) - \frac{7}{25} \log(|3 \tan(x) + 4|)$$

input `integrate((4*sin(x)+3*cos(x))/(3*sin(x)+4*cos(x)),x, algorithm="giac")`

output `24/25*x + 7/50*log(tan(x)^2 + 1) - 7/25*log(abs(3*tan(x) + 4))`

3.261.9 Mupad [B] (verification not implemented)

Time = 16.87 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.60

$$\int \frac{3 \cos(x) + 4 \sin(x)}{4 \cos(x) + 3 \sin(x)} dx = \frac{24x}{25} - \frac{7 \ln \left(\tan\left(\frac{x}{2}\right)^2 - \frac{3 \tan\left(\frac{x}{2}\right)}{2} - 1 \right)}{25} + \frac{7 \ln \left(\tan\left(\frac{x}{2}\right)^2 + 1 \right)}{25}$$

input `int((3*cos(x) + 4*sin(x))/(4*cos(x) + 3*sin(x)),x)`

output `(24*x)/25 - (7*log(tan(x/2)^2 - (3*tan(x/2))/2 - 1))/25 + (7*log(tan(x/2)^2 + 1))/25`

$$3.262 \quad \int \left(-\sqrt{3} - \sqrt{4 - x^2} + \sqrt{4 - (1 + x)^2} \right) dx$$

3.262.1 Optimal result	1324
3.262.2 Mathematica [A] (verified)	1324
3.262.3 Rubi [A] (verified)	1325
3.262.4 Maple [A] (verified)	1325
3.262.5 Fricas [A] (verification not implemented)	1326
3.262.6 Sympy [A] (verification not implemented)	1326
3.262.7 Maxima [A] (verification not implemented)	1327
3.262.8 Giac [A] (verification not implemented)	1327
3.262.9 Mupad [F(-1)]	1328

3.262.1 Optimal result

Integrand size = 34, antiderivative size = 63

$$\int \left(-\sqrt{3} - \sqrt{4 - x^2} + \sqrt{4 - (1 + x)^2} \right) dx = -\sqrt{3}x - \frac{1}{2}x\sqrt{4 - x^2} + \frac{1}{2}(1 + x)\sqrt{4 - (1 + x)^2} \\ - 2 \arcsin\left(\frac{x}{2}\right) + 2 \arcsin\left(\frac{1 + x}{2}\right)$$

output `-x*3^(1/2)-1/2*x*(-x^2+4)^(1/2)+1/2*(1+x)*(4-(1+x)^2)^(1/2)-2*arcsin(1/2*x)+2*arcsin(1/2+1/2*x)`

3.262.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.41

$$\int \left(-\sqrt{3} - \sqrt{4 - x^2} + \sqrt{4 - (1 + x)^2} \right) dx = -\sqrt{3}x - \frac{1}{2}x\sqrt{4 - x^2} + \frac{1}{2}(1 + x)\sqrt{3 - 2x - x^2} \\ + 4 \arctan\left(\frac{\sqrt{4 - x^2}}{2 + x}\right) \\ - 4 \arctan\left(\frac{\sqrt{3 - 2x - x^2}}{3 + x}\right)$$

input `Integrate[-Sqrt[3] - Sqrt[4 - x^2] + Sqrt[4 - (1 + x)^2], x]`

3.262. $\int \left(-\sqrt{3} - \sqrt{4 - x^2} + \sqrt{4 - (1 + x)^2} \right) dx$

output $-(\text{Sqrt}[3]*x) - (x*\text{Sqrt}[4 - x^2])/2 + ((1 + x)*\text{Sqrt}[3 - 2*x - x^2])/2 + 4*\text{ArcTan}[\text{Sqrt}[4 - x^2]/(2 + x)] - 4*\text{ArcTan}[\text{Sqrt}[3 - 2*x - x^2]/(3 + x)]$

3.262.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.029$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(-\sqrt{4-x^2} + \sqrt{4-(x+1)^2} - \sqrt{3} \right) dx$$

↓ 2009

$$-2 \arcsin\left(\frac{x}{2}\right) + 2 \arcsin\left(\frac{x+1}{2}\right) - \frac{1}{2}\sqrt{4-x^2} - \sqrt{3}x + \frac{1}{2}(x+1)\sqrt{4-(x+1)^2}$$

input $\text{Int}[-\text{Sqrt}[3] - \text{Sqrt}[4 - x^2] + \text{Sqrt}[4 - (1 + x)^2], x]$

output $-(\text{Sqrt}[3]*x) - (x*\text{Sqrt}[4 - x^2])/2 + ((1 + x)*\text{Sqrt}[4 - (1 + x)^2])/2 - 2*\text{ArcSin}[x/2] + 2*\text{ArcSin}[(1 + x)/2]$

3.262.3.1 Defintions of rubi rules used

rule 2009 $\text{Int}[u_, x_Symbol] \text{ :> } \text{Simp}[\text{IntSum}[u, x], x] \text{ /; } \text{SumQ}[u]$

3.262.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.84

method	result	size
default	$-\frac{(-2-2x)\sqrt{-x^2-2x+3}}{4} + 2 \arcsin\left(\frac{x}{2} + \frac{1}{2}\right) - x\sqrt{3} - \frac{x\sqrt{-x^2+4}}{2} - 2 \arcsin\left(\frac{x}{2}\right)$	53
parts	$-\frac{(-2-2x)\sqrt{-x^2-2x+3}}{4} + 2 \arcsin\left(\frac{x}{2} + \frac{1}{2}\right) - x\sqrt{3} - \frac{x\sqrt{-x^2+4}}{2} - 2 \arcsin\left(\frac{x}{2}\right)$	53

input $\text{int}((4-(1+x)^2)^{(1/2)}-3^{(1/2)}-(-x^2+4)^{(1/2)}, x, \text{method}=_RETURNVERBOSE)$

3.262. $\int \left(-\sqrt{3} - \sqrt{4-x^2} + \sqrt{4-(1+x)^2} \right) dx$

output `-1/4*(-2-2*x)*(-x^2-2*x+3)^(1/2)+2*arcsin(1/2*x+1/2)-x*3^(1/2)-1/2*x*(-x^2+4)^(1/2)-2*arcsin(1/2*x)`

3.262.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.32

$$\int \left(-\sqrt{3} - \sqrt{4-x^2} + \sqrt{4-(1+x)^2} \right) dx = \frac{1}{2} \sqrt{-x^2-2x+3}(x+1) - \sqrt{3}x - \frac{1}{2} \sqrt{-x^2+4}x - 2 \arctan \left(\frac{\sqrt{-x^2-2x+3}(x+1)}{x^2+2x-3} \right) + 4 \arctan \left(\frac{\sqrt{-x^2+4}-2}{x} \right)$$

input `integrate((4-(1+x)^2)^(1/2)-3^(1/2)-(-x^2+4)^(1/2),x, algorithm="fricas")`

output `1/2*sqrt(-x^2 - 2*x + 3)*(x + 1) - sqrt(3)*x - 1/2*sqrt(-x^2 + 4)*x - 2*arctan(sqrt(-x^2 - 2*x + 3)*(x + 1)/(x^2 + 2*x - 3)) + 4*arctan((sqrt(-x^2 + 4) - 2)/x)`

3.262.6 Sympy [A] (verification not implemented)

Time = 1.06 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.89

$$\int \left(-\sqrt{3} - \sqrt{4-x^2} + \sqrt{4-(1+x)^2} \right) dx = -\frac{x\sqrt{4-x^2}}{2} - \sqrt{3}x + \begin{cases} \frac{i(x+1)^3}{2\sqrt{(x+1)^2-4}} - \frac{2i(x+1)}{\sqrt{(x+1)^2-4}} - 2i \operatorname{acosh} \left(\frac{x}{2} + \frac{1}{2} \right) & \text{for } |(x+1)^2| > 4 \\ 2 \operatorname{asin} \left(\frac{x}{2} + \frac{1}{2} \right) - \frac{(x+1)^3}{2\sqrt{4-(x+1)^2}} + \frac{2(x+1)}{\sqrt{4-(x+1)^2}} & \text{otherwise} \end{cases} - 2 \operatorname{asin} \left(\frac{x}{2} \right)$$

input `integrate((4-(1+x)**2)**(1/2)-3**(1/2)-(-x**2+4)**(1/2),x)`

output `-x*sqrt(4 - x**2)/2 - sqrt(3)*x + Piecewise((I*(x + 1)**3/(2*sqrt((x + 1)*
*2 - 4)) - 2*I*(x + 1)/sqrt((x + 1)**2 - 4) - 2*I*acosh(x/2 + 1/2), Abs((x
+ 1)**2) > 4), (2*asin(x/2 + 1/2) - (x + 1)**3/(2*sqrt(4 - (x + 1)**2)) +
2*(x + 1)/sqrt(4 - (x + 1)**2), True)) - 2*asin(x/2)`

3.262.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.98

$$\int \left(-\sqrt{3} - \sqrt{4 - x^2} + \sqrt{4 - (1 + x)^2} \right) dx = -\sqrt{3}x + \frac{1}{2} \sqrt{-x^2 - 2x + 3} \\ - \frac{1}{2} \sqrt{-x^2 + 4x} + \frac{1}{2} \sqrt{-x^2 - 2x + 3} \\ - 2 \arcsin\left(\frac{1}{2}x\right) - 2 \arcsin\left(-\frac{1}{2}x - \frac{1}{2}\right)$$

input `integrate((4-(1+x)^2)^(1/2)-3^(1/2)-(-x^2+4)^(1/2),x, algorithm="maxima")`

output `-sqrt(3)*x + 1/2*sqrt(-x^2 - 2*x + 3)*x - 1/2*sqrt(-x^2 + 4)*x + 1/2*sqrt(
-x^2 - 2*x + 3) - 2*arcsin(1/2*x) - 2*arcsin(-1/2*x - 1/2)`

3.262.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.79

$$\int \left(-\sqrt{3} - \sqrt{4 - x^2} + \sqrt{4 - (1 + x)^2} \right) dx = \frac{1}{2} \sqrt{-x^2 - 2x + 3}(x + 1) - \sqrt{3}x - \frac{1}{2} \sqrt{-x^2 + 4}x \\ - 2 \arcsin\left(\frac{1}{2}x\right) + 2 \arcsin\left(\frac{1}{2}x + \frac{1}{2}\right)$$

input `integrate((4-(1+x)^2)^(1/2)-3^(1/2)-(-x^2+4)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(-x^2 - 2*x + 3)*(x + 1) - sqrt(3)*x - 1/2*sqrt(-x^2 + 4)*x - 2*ar
csin(1/2*x) + 2*arcsin(1/2*x + 1/2)`

3.262.9 Mupad [F(-1)]

Timed out.

$$\int \left(-\sqrt{3} - \sqrt{4-x^2} + \sqrt{4-(1+x)^2} \right) dx = \int \sqrt{4-(x+1)^2} - \sqrt{3} - \sqrt{4-x^2} dx$$

input `int((4 - (x + 1)^2)^(1/2) - 3^(1/2) - (4 - x^2)^(1/2), x)`output `int((4 - (x + 1)^2)^(1/2) - 3^(1/2) - (4 - x^2)^(1/2), x)`

3.263 $\int x^2 \sin(\log(x)) dx$

3.263.1 Optimal result	1329
3.263.2 Mathematica [A] (verified)	1329
3.263.3 Rubi [A] (verified)	1330
3.263.4 Maple [C] (verified)	1330
3.263.5 Fricas [A] (verification not implemented)	1331
3.263.6 Sympy [A] (verification not implemented)	1331
3.263.7 Maxima [A] (verification not implemented)	1331
3.263.8 Giac [A] (verification not implemented)	1332
3.263.9 Mupad [B] (verification not implemented)	1332

3.263.1 Optimal result

Integrand size = 7, antiderivative size = 21

$$\int x^2 \sin(\log(x)) dx = -\frac{1}{10}x^3 \cos(\log(x)) + \frac{3}{10}x^3 \sin(\log(x))$$

output `-1/10*x^3*cos(ln(x))+3/10*x^3*sin(ln(x))`

3.263.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int x^2 \sin(\log(x)) dx = -\frac{1}{10}x^3 \cos(\log(x)) + \frac{3}{10}x^3 \sin(\log(x))$$

input `Integrate[x^2*Sin[Log[x]],x]`

output `-1/10*(x^3*Cos[Log[x]]) + (3*x^3*Sin[Log[x]])/10`

3.263.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {4988}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \sin(\log(x)) dx$$

↓ 4988

$$\frac{3}{10}x^3 \sin(\log(x)) - \frac{1}{10}x^3 \cos(\log(x))$$

input `Int[x^2*Sin[Log[x]],x]`

output `-1/10*(x^3*Cos[Log[x]]) + (3*x^3*Sin[Log[x]])/10`

3.263.3.1 Defintions of rubi rules used

rule 4988 `Int[((e_.)*(x_)^(m_.)*Sin[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)], x_Symbol] :> Simp[(m + 1)*(e*x)^(m + 1)*(Sin[d*(a + b*Log[c*x^n])]/(b^2*d^2*e*n^2 + e*(m + 1)^2)), x] - Simp[b*d*n*(e*x)^(m + 1)*(Cos[d*(a + b*Log[c*x^n])]/(b^2*d^2*e*n^2 + e*(m + 1)^2)), x] /; FreeQ[{a, b, c, d, e, m, n}, x] & NeQ[b^2*d^2*n^2 + (m + 1)^2, 0]`

3.263.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.24

method	result	size
risch	$\left(-\frac{1}{20} - \frac{3i}{20}\right) x^3 x^i + \left(-\frac{1}{20} + \frac{3i}{20}\right) x^3 x^{-i}$	26
norman	$\frac{-\frac{x^3}{10} + \frac{3x^3 \tan\left(\frac{\ln(x)}{2}\right)}{5} + \frac{x^3 \tan\left(\frac{\ln(x)}{2}\right)^2}{10}}{1 + \tan\left(\frac{\ln(x)}{2}\right)^2}$	41

input `int(x^2*sin(ln(x)),x,method=_RETURNVERBOSE)`

output `(-1/20-3/20*I)*x^3*x^I+(-1/20+3/20*I)*x^3/(x^I)`

3.263.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int x^2 \sin(\log(x)) dx = -\frac{1}{10} x^3 \cos(\log(x)) + \frac{3}{10} x^3 \sin(\log(x))$$

input `integrate(x^2*sin(log(x)),x, algorithm="fricas")`

output `-1/10*x^3*cos(log(x)) + 3/10*x^3*sin(log(x))`

3.263.6 Sympy [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.95

$$\int x^2 \sin(\log(x)) dx = \frac{3x^3 \sin(\log(x))}{10} - \frac{x^3 \cos(\log(x))}{10}$$

input `integrate(x**2*sin(ln(x)),x)`

output `3*x**3*sin(log(x))/10 - x**3*cos(log(x))/10`

3.263.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

$$\int x^2 \sin(\log(x)) dx = -\frac{1}{10} x^3 (\cos(\log(x)) - 3 \sin(\log(x)))$$

input `integrate(x^2*sin(log(x)),x, algorithm="maxima")`

output `-1/10*x^3*(cos(log(x)) - 3*sin(log(x)))`

3.263.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int x^2 \sin(\log(x)) dx = -\frac{1}{10} x^3 \cos(\log(x)) + \frac{3}{10} x^3 \sin(\log(x))$$

input `integrate(x^2*sin(log(x)),x, algorithm="giac")`output `-1/10*x^3*cos(log(x)) + 3/10*x^3*sin(log(x))`**3.263.9 Mupad [B] (verification not implemented)**

Time = 16.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

$$\int x^2 \sin(\log(x)) dx = -\frac{\sqrt{10} x^3 \cos(\operatorname{atan}(3) + \ln(x))}{10}$$

input `int(x^2*sin(log(x)),x)`output `-(10^(1/2)*x^3*cos(atan(3) + log(x)))/10`

3.264 $\int e^{-x}(36x^5 - 12x^6 + x^7) dx$

3.264.1 Optimal result	1333
3.264.2 Mathematica [A] (verified)	1333
3.264.3 Rubi [A] (verified)	1334
3.264.4 Maple [A] (verified)	1335
3.264.5 Fricas [A] (verification not implemented)	1335
3.264.6 Sympy [A] (verification not implemented)	1336
3.264.7 Maxima [A] (verification not implemented)	1336
3.264.8 Giac [A] (verification not implemented)	1336
3.264.9 Mupad [B] (verification not implemented)	1337

3.264.1 Optimal result

Integrand size = 20, antiderivative size = 76

$$\int e^{-x}(36x^5 - 12x^6 + x^7) dx = -720e^{-x} - 720e^{-x}x - 360e^{-x}x^2 - 120e^{-x}x^3 - 30e^{-x}x^4 - 6e^{-x}x^5 + 5e^{-x}x^6 - e^{-x}x^7$$

output $-720/\exp(x)-720*x/\exp(x)-360*x^2/\exp(x)-120*x^3/\exp(x)-30*x^4/\exp(x)-6*x^5/\exp(x)+5*x^6/\exp(x)-x^7/\exp(x)$

3.264.2 Mathematica [A] (verified)

Time = 0.62 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.54

$$\int e^{-x}(36x^5 - 12x^6 + x^7) dx = e^{-x}(-720 - 720x - 360x^2 - 120x^3 - 30x^4 - 6x^5 + 5x^6 - x^7)$$

input `Integrate[(36*x^5 - 12*x^6 + x^7)/E^x,x]`

output $(-720 - 720*x - 360*x^2 - 120*x^3 - 30*x^4 - 6*x^5 + 5*x^6 - x^7)/E^x$

3.264.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {2028, 2626, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^{-x}(x^7 - 12x^6 + 36x^5) dx \\ & \quad \downarrow \text{2028} \\ & \int e^{-x}x^5(x^2 - 12x + 36) dx \\ & \quad \downarrow \text{2626} \\ & \int (e^{-x}x^7 - 12e^{-x}x^6 + 36e^{-x}x^5) dx \\ & \quad \downarrow \text{2009} \\ & -e^{-x}x^7 + 5e^{-x}x^6 - 6e^{-x}x^5 - 30e^{-x}x^4 - 120e^{-x}x^3 - 360e^{-x}x^2 - 720e^{-x}x - 720e^{-x} \end{aligned}$$

input `Int[(36*x^5 - 12*x^6 + x^7)/E^x,x]`

output `-720/E^x - (720*x)/E^x - (360*x^2)/E^x - (120*x^3)/E^x - (30*x^4)/E^x - (6*x^5)/E^x + (5*x^6)/E^x - x^7/E^x`

3.264.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2028 `Int[(F_x_.)*((a_.)*(x_)^(r_.) + (b_.)*(x_)^(s_.) + (c_.)*(x_)^(t_.))^p_.), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r) + c*x^(t - r))^p*F_x, x] /; FreeQ[{a, b, c, r, s, t}, x] && IntegerQ[p] && PosQ[s - r] && PosQ[t - r] && !(EqQ[p, 1] && EqQ[u, 1])`

rule 2626 `Int[(F_)^(v_)*(P_x_), x_Symbol] := Int[ExpandIntegrand[F^v, P_x, x], x] /; FreeQ[F, x] && PolynomialQ[P_x, x] && LinearQ[v, x] && !TrueQ[$UseGamma]`

3.264.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.53

method	result
gosper	$-e^{-x}(x^7 - 5x^6 + 6x^5 + 30x^4 + 120x^3 + 360x^2 + 720x + 720)$
risch	$(-x^7 + 5x^6 - 6x^5 - 30x^4 - 120x^3 - 360x^2 - 720x - 720)e^{-x}$
derivativedivides	$-6e^{-x}x^5 - 30x^4e^{-x} - 120x^3e^{-x} - 360x^2e^{-x} - 720xe^{-x} - 720e^{-x} + 5e^{-x}x^6 - e^{-x}x^7$
default	$-6e^{-x}x^5 - 30x^4e^{-x} - 120x^3e^{-x} - 360x^2e^{-x} - 720xe^{-x} - 720e^{-x} + 5e^{-x}x^6 - e^{-x}x^7$
norman	$-6e^{-x}x^5 - 30x^4e^{-x} - 120x^3e^{-x} - 360x^2e^{-x} - 720xe^{-x} - 720e^{-x} + 5e^{-x}x^6 - e^{-x}x^7$
parallelrisch	$-6e^{-x}x^5 - 30x^4e^{-x} - 120x^3e^{-x} - 360x^2e^{-x} - 720xe^{-x} - 720e^{-x} + 5e^{-x}x^6 - e^{-x}x^7$
parts	$-6e^{-x}x^5 - 30x^4e^{-x} - 120x^3e^{-x} - 360x^2e^{-x} - 720xe^{-x} - 720e^{-x} + 5e^{-x}x^6 - e^{-x}x^7$
meijerg	$720 - \frac{(8x^7+56x^6+336x^5+1680x^4+6720x^3+20160x^2+40320x+40320)e^{-x}}{8} + \frac{12(7x^6+42x^5+210x^4+840x^3+2520x^2+2016x+5040)e^{-x}}{7}$

input `int(exp(-x)*(x^7-12*x^6+36*x^5),x,method=_RETURNVERBOSE)`output `-exp(-x)*(x^7-5*x^6+6*x^5+30*x^4+120*x^3+360*x^2+720*x+720)`**3.264.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.51

$$\int e^{-x}(36x^5 - 12x^6 + x^7) dx$$

$$= -(x^7 - 5x^6 + 6x^5 + 30x^4 + 120x^3 + 360x^2 + 720x + 720)e^{(-x)}$$

input `integrate(exp(-x)*(x^7-12*x^6+36*x^5),x, algorithm="fricas")`output `-(x^7 - 5*x^6 + 6*x^5 + 30*x^4 + 120*x^3 + 360*x^2 + 720*x + 720)*e^(-x)`

3.264.6 Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.47

$$\int e^{-x}(36x^5 - 12x^6 + x^7) dx = (-x^7 + 5x^6 - 6x^5 - 30x^4 - 120x^3 - 360x^2 - 720x - 720)e^{-x}$$

input `integrate(exp(-x)*(x**7-12*x**6+36*x**5),x)`output `(-x**7 + 5*x**6 - 6*x**5 - 30*x**4 - 120*x**3 - 360*x**2 - 720*x - 720)*exp(-x)`**3.264.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.36

$$\begin{aligned} & \int e^{-x}(36x^5 - 12x^6 + x^7) dx \\ &= -(x^7 + 7x^6 + 42x^5 + 210x^4 + 840x^3 + 2520x^2 + 5040x + 5040)e^{(-x)} \\ & \quad + 12(x^6 + 6x^5 + 30x^4 + 120x^3 + 360x^2 + 720x + 720)e^{(-x)} \\ & \quad - 36(x^5 + 5x^4 + 20x^3 + 60x^2 + 120x + 120)e^{(-x)} \end{aligned}$$

input `integrate(exp(-x)*(x^7-12*x^6+36*x^5),x, algorithm="maxima")`output `-(x^7 + 7*x^6 + 42*x^5 + 210*x^4 + 840*x^3 + 2520*x^2 + 5040*x + 5040)*e^(-x) + 12*(x^6 + 6*x^5 + 30*x^4 + 120*x^3 + 360*x^2 + 720*x + 720)*e^(-x) - 36*(x^5 + 5*x^4 + 20*x^3 + 60*x^2 + 120*x + 120)*e^(-x)`**3.264.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.51

$$\begin{aligned} & \int e^{-x}(36x^5 - 12x^6 + x^7) dx \\ &= -(x^7 - 5x^6 + 6x^5 + 30x^4 + 120x^3 + 360x^2 + 720x + 720)e^{(-x)} \end{aligned}$$

input `integrate(exp(-x)*(x^7-12*x^6+36*x^5),x, algorithm="giac")`output `-(x^7 - 5*x^6 + 6*x^5 + 30*x^4 + 120*x^3 + 360*x^2 + 720*x + 720)*e^(-x)`

3.264.9 Mupad [B] (verification not implemented)

Time = 16.51 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.51

$$\int e^{-x}(36x^5 - 12x^6 + x^7) dx = -e^{-x}(x^7 - 5x^6 + 6x^5 + 30x^4 + 120x^3 + 360x^2 + 720x + 720)$$

input `int(exp(-x)*(36*x^5 - 12*x^6 + x^7),x)`

output `-exp(-x)*(720*x + 360*x^2 + 120*x^3 + 30*x^4 + 6*x^5 - 5*x^6 + x^7 + 720)`

3.265 $\int \arccos(x) \arcsin(x) dx$

3.265.1 Optimal result	1338
3.265.2 Mathematica [A] (verified)	1338
3.265.3 Rubi [F]	1339
3.265.4 Maple [F]	1339
3.265.5 Fricas [A] (verification not implemented)	1340
3.265.6 Sympy [A] (verification not implemented)	1340
3.265.7 Maxima [A] (verification not implemented)	1340
3.265.8 Giac [B] (verification not implemented)	1341
3.265.9 Mupad [F(-1)]	1341

3.265.1 Optimal result

Integrand size = 5, antiderivative size = 38

$$\int \arccos(x) \arcsin(x) dx = 2x - \sqrt{1 - x^2} \arcsin(x) + \arccos(x) (\sqrt{1 - x^2} + x \arcsin(x))$$

output `2*x-(-x^2+1)^(1/2)*arcsin(x)+arccos(x)*((-x^2+1)^(1/2)+x*arcsin(x))`

3.265.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \arccos(x) \arcsin(x) dx = 2x - \sqrt{1 - x^2} \arcsin(x) + \arccos(x) (\sqrt{1 - x^2} + x \arcsin(x))$$

input `Integrate[ArcCos[x]*ArcSin[x],x]`

output `2*x - Sqrt[1 - x^2]*ArcSin[x] + ArcCos[x]*(Sqrt[1 - x^2] + x*ArcSin[x])`

3.265.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arccos(x) \arcsin(x) dx$$

↓ 5300

$$\int \arccos(x) \arcsin(x) dx$$

input `Int[ArcCos[x]*ArcSin[x],x]`

output `$Aborted`

3.265.3.1 Defintions of rubi rules used

rule 5300 `Int[((a_.) + ArcSin[(c_.)*(x_.)]*(b_.))^n_.*(u_.), x_Symbol] := Unintegrate[u*(a + b*ArcSin[c*x])^n, x] /; FreeQ[{a, b, c, n}, x]`

3.265.4 Maple [F]

$$\int \arcsin(x) \arccos(x) dx$$

input `int(arcsin(x)*arccos(x),x)`

output `int(arcsin(x)*arccos(x),x)`

3.265.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.89

$$\int \arccos(x) \arcsin(x) dx = \frac{1}{2} \pi x \arccos(x) - x \arccos(x)^2 - \frac{1}{2} (\pi - 4 \arccos(x)) \sqrt{-x^2 + 1} + 2x$$

input `integrate(arcsin(x)*arccos(x),x, algorithm="fricas")`output `1/2*pi*x*arccos(x) - x*arccos(x)^2 - 1/2*(pi - 4*arccos(x))*sqrt(-x^2 + 1) + 2*x`**3.265.6 Sympy [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.89

$$\int \arccos(x) \arcsin(x) dx = x \arccos(x) \arcsin(x) + 2x + \sqrt{1-x^2} \arccos(x) - \sqrt{1-x^2} \arcsin(x)$$

input `integrate(asin(x)*acos(x),x)`output `x*acos(x)*asin(x) + 2*x + sqrt(1 - x**2)*acos(x) - sqrt(1 - x**2)*asin(x)`**3.265.7 Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.89

$$\int \arccos(x) \arcsin(x) dx = \left(x \arcsin(x) + \sqrt{-x^2 + 1} \right) \arccos(x) - \sqrt{-x^2 + 1} \arcsin(x) + 2x$$

input `integrate(arcsin(x)*arccos(x),x, algorithm="maxima")`output `(x*arcsin(x) + sqrt(-x^2 + 1))*arccos(x) - sqrt(-x^2 + 1)*arcsin(x) + 2*x`

3.265.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 163 vs. $2(34) = 68$.

Time = 0.28 (sec) , antiderivative size = 163, normalized size of antiderivative = 4.29

$$\int \arccos(x) \arcsin(x) dx = -\pi(-1)^{\lfloor -\frac{\arccos(x)}{\pi} + 1 \rfloor} x \arccos(x) \left[-\frac{\arccos(x)}{\pi} + 1 \right] \\ + \frac{1}{2} \pi(-1)^{\lfloor -\frac{\arccos(x)}{\pi} + 1 \rfloor} x \arccos(x) \\ - (-1)^{\lfloor -\frac{\arccos(x)}{\pi} + 1 \rfloor} x \arccos(x)^2 \\ + \pi \sqrt{-x^2 + 1} (-1)^{\lfloor -\frac{\arccos(x)}{\pi} + 1 \rfloor} \left[-\frac{\arccos(x)}{\pi} + 1 \right] \\ - \frac{1}{2} \pi \sqrt{-x^2 + 1} (-1)^{\lfloor -\frac{\arccos(x)}{\pi} + 1 \rfloor} \\ + 2 \sqrt{-x^2 + 1} (-1)^{\lfloor -\frac{\arccos(x)}{\pi} + 1 \rfloor} \arccos(x) + 2 (-1)^{\lfloor -\frac{\arccos(x)}{\pi} + 1 \rfloor} x$$

input `integrate(arcsin(x)*arccos(x),x, algorithm="giac")`

output `-pi*(-1)^floor(-arccos(x)/pi + 1)*x*arccos(x)*floor(-arccos(x)/pi + 1) + 1/2*pi*(-1)^floor(-arccos(x)/pi + 1)*x*arccos(x) - (-1)^floor(-arccos(x)/pi + 1)*x*arccos(x)^2 + pi*sqrt(-x^2 + 1)*(-1)^floor(-arccos(x)/pi + 1)*floor(-arccos(x)/pi + 1) - 1/2*pi*sqrt(-x^2 + 1)*(-1)^floor(-arccos(x)/pi + 1) + 2*sqrt(-x^2 + 1)*(-1)^floor(-arccos(x)/pi + 1)*arccos(x) + 2*(-1)^floor(-arccos(x)/pi + 1)*x`

3.265.9 Mupad [F(-1)]

Timed out.

$$\int \arccos(x) \arcsin(x) dx = \int \operatorname{acos}(x) \operatorname{asin}(x) dx$$

input `int(acos(x)*asin(x),x)`

output `int(acos(x)*asin(x), x)`

3.266 $\int (1 + 6x - 7x^2 + 4x^3 - x^4)^n dx$

3.266.1 Optimal result	1342
3.266.2 Mathematica [F]	1342
3.266.3 Rubi [B] (verified)	1343
3.266.4 Maple [F]	1344
3.266.5 Fricas [F]	1344
3.266.6 Sympy [F]	1345
3.266.7 Maxima [F]	1345
3.266.8 Giac [F]	1345
3.266.9 Mupad [F(-1)]	1346

3.266.1 Optimal result

Integrand size = 22, antiderivative size = 52

$$\int (1 + 6x - 7x^2 + 4x^3 - x^4)^n dx = 3^n(-1 + x) \operatorname{AppellF1}\left(\frac{1}{2}, -n, -n, \frac{3}{2}, \frac{2(-1 + x)^2}{1 + \sqrt{13}}, \frac{2(-1 + x)^2}{-1 + \sqrt{13}}\right)$$

output `3^n*(-1+x)*AppellF1(1/2,-n,-n,3/2,2*(-1+x)^2/(-1+13^(1/2)),-2*(-1+x)^2/(1+13^(1/2)))`

3.266.2 Mathematica [F]

$$\int (1 + 6x - 7x^2 + 4x^3 - x^4)^n dx = \int (1 + 6x - 7x^2 + 4x^3 - x^4)^n dx$$

input `Integrate[(1 + 6*x - 7*x^2 + 4*x^3 - x^4)^n,x]`

output `Integrate[(1 + 6*x - 7*x^2 + 4*x^3 - x^4)^n, x]`

3.266.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 115 vs. $2(52) = 104$.

Time = 0.26 (sec) , antiderivative size = 115, normalized size of antiderivative = 2.21, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$, Rules used = {2458, 1418, 333}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (-x^4 + 4x^3 - 7x^2 + 6x + 1)^n dx$$

$$\downarrow \text{2458}$$

$$\int (-(x-1)^4 - (x-1)^2 + 3)^n d(x-1)$$

$$\downarrow \text{1418}$$

$$\left(\frac{2(x-1)^2}{1-\sqrt{13}} + 1\right)^{-n} \left(\frac{2(x-1)^2}{1+\sqrt{13}} + 1\right)^{-n} (-(x-1)^4 - (x-1)^2 + 3)^n \int \left(\frac{2(x-1)^2}{1-\sqrt{13}} + 1\right)^n \left(\frac{2(x-1)^2}{1+\sqrt{13}} + 1\right)^n d(x-1)$$

$$\downarrow \text{333}$$

$$1) \left(\frac{2(x-1)^2}{1-\sqrt{13}} + 1\right)^{-n} \left(\frac{2(x-1)^2}{1+\sqrt{13}} + 1\right)^{-n} (-(x-1)^4 - (x-1)^2 + 3)^n \text{AppellF1}\left(\frac{1}{2}, -n, -n, \frac{3}{2}, -\frac{2(x-1)^2}{1-\sqrt{13}}, -\frac{2(x-1)^2}{1+\sqrt{13}}\right)$$

input `Int[(1 + 6*x - 7*x^2 + 4*x^3 - x^4)^n, x]`

output `((3 - (-1 + x)^2 - (-1 + x)^4)^n*(-1 + x)*AppellF1[1/2, -n, -n, 3/2, (-2*(-1 + x)^2)/(1 - Sqrt[13]), (-2*(-1 + x)^2)/(1 + Sqrt[13])])/((1 + (2*(-1 + x)^2)/(1 - Sqrt[13]))^n*(1 + (2*(-1 + x)^2)/(1 + Sqrt[13]))^n)`

3.266.3.1 Defintions of rubi rules used

rule 333 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 1418 `Int[((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[a^IntPart[p]*((a + b*x^2 + c*x^4)^FracPart[p]/((1 + 2*c*(x^2/(b + q)))^FracPart[p]*(1 + 2*c*(x^2/(b - q)))^FracPart[p])) Int[(1 + 2*c*(x^2/(b + q)))^p*(1 + 2*c*(x^2/(b - q)))^p, x], x] /; FreeQ[{a, b, c, p}, x] && NeQ[b^2 - 4*a*c, 0]`

rule 2458 `Int[(Pn_)^(p_.), x_Symbol] := With[{S = Coeff[Pn, x, Expon[Pn, x] - 1]/(Expon[Pn, x]*Coeff[Pn, x, Expon[Pn, x]])}, Subst[Int[ExpandToSum[Pn /. x -> x - S, x]^p, x], x, x + S] /; BinomialQ[Pn /. x -> x - S, x] || (IntegerQ[Expon[Pn, x]/2] && TrinomialQ[Pn /. x -> x - S, x]) /; FreeQ[p, x] && PolyQ[Pn, x] && GtQ[Expon[Pn, x], 2] && NeQ[Coeff[Pn, x, Expon[Pn, x] - 1], 0]`

3.266.4 Maple [F]

$$\int (-x^4 + 4x^3 - 7x^2 + 6x + 1)^n dx$$

input `int((-x^4+4*x^3-7*x^2+6*x+1)^n,x)`

output `int((-x^4+4*x^3-7*x^2+6*x+1)^n,x)`

3.266.5 Fracas [F]

$$\int (1 + 6x - 7x^2 + 4x^3 - x^4)^n dx = \int (-x^4 + 4x^3 - 7x^2 + 6x + 1)^n dx$$

input `integrate((-x^4+4*x^3-7*x^2+6*x+1)^n,x, algorithm="fracas")`

output `integral((-x^4 + 4*x^3 - 7*x^2 + 6*x + 1)^n, x)`

3.266.6 Sympy [F]

$$\int (1 + 6x - 7x^2 + 4x^3 - x^4)^n dx = \int (-x^4 + 4x^3 - 7x^2 + 6x + 1)^n dx$$

input `integrate((-x**4+4*x**3-7*x**2+6*x+1)**n,x)`

output `Integral((-x**4 + 4*x**3 - 7*x**2 + 6*x + 1)**n, x)`

3.266.7 Maxima [F]

$$\int (1 + 6x - 7x^2 + 4x^3 - x^4)^n dx = \int (-x^4 + 4x^3 - 7x^2 + 6x + 1)^n dx$$

input `integrate((-x^4+4*x^3-7*x^2+6*x+1)^n,x, algorithm="maxima")`

output `integrate((-x^4 + 4*x^3 - 7*x^2 + 6*x + 1)^n, x)`

3.266.8 Giac [F]

$$\int (1 + 6x - 7x^2 + 4x^3 - x^4)^n dx = \int (-x^4 + 4x^3 - 7x^2 + 6x + 1)^n dx$$

input `integrate((-x^4+4*x^3-7*x^2+6*x+1)^n,x, algorithm="giac")`

output `integrate((-x^4 + 4*x^3 - 7*x^2 + 6*x + 1)^n, x)`

3.266.9 Mupad [F(-1)]

Timed out.

$$\int (1 + 6x - 7x^2 + 4x^3 - x^4)^n dx = \int (-x^4 + 4x^3 - 7x^2 + 6x + 1)^n dx$$

input `int((6*x - 7*x^2 + 4*x^3 - x^4 + 1)^n,x)`output `int((6*x - 7*x^2 + 4*x^3 - x^4 + 1)^n, x)`

3.267 $\int \frac{x^4}{\sqrt{1-x}} dx$

3.267.1 Optimal result	1347
3.267.2 Mathematica [A] (verified)	1347
3.267.3 Rubi [A] (verified)	1348
3.267.4 Maple [A] (verified)	1349
3.267.5 Fricas [A] (verification not implemented)	1349
3.267.6 Sympy [C] (verification not implemented)	1350
3.267.7 Maxima [A] (verification not implemented)	1350
3.267.8 Giac [A] (verification not implemented)	1350
3.267.9 Mupad [B] (verification not implemented)	1351

3.267.1 Optimal result

Integrand size = 13, antiderivative size = 64

$$\int \frac{x^4}{\sqrt{1-x}} dx = -2\sqrt{1-x} + \frac{8}{3}(1-x)^{3/2} - \frac{12}{5}(1-x)^{5/2} + \frac{8}{7}(1-x)^{7/2} - \frac{2}{9}(1-x)^{9/2}$$

output `-2*(1-x)^(1/2)+8/3*(1-x)^(3/2)-12/5*(1-x)^(5/2)+8/7*(1-x)^(7/2)-2/9*(1-x)^(9/2)`

3.267.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.52

$$\int \frac{x^4}{\sqrt{1-x}} dx = -\frac{2}{315}\sqrt{1-x}(128 + 64x + 48x^2 + 40x^3 + 35x^4)$$

input `Integrate[x^4/Sqrt[1 - x],x]`

output `(-2*Sqrt[1 - x]*(128 + 64*x + 48*x^2 + 40*x^3 + 35*x^4))/315`

3.267.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4}{\sqrt{1-x}} dx$$

↓ 53

$$\int \left((1-x)^{7/2} - 4(1-x)^{5/2} + 6(1-x)^{3/2} - 4\sqrt{1-x} + \frac{1}{\sqrt{1-x}} \right) dx$$

↓ 2009

$$-\frac{2}{9}(1-x)^{9/2} + \frac{8}{7}(1-x)^{7/2} - \frac{12}{5}(1-x)^{5/2} + \frac{8}{3}(1-x)^{3/2} - 2\sqrt{1-x}$$

input `Int[x^4/Sqrt[1 - x],x]`

output `-2*Sqrt[1 - x] + (8*(1 - x)^(3/2))/3 - (12*(1 - x)^(5/2))/5 + (8*(1 - x)^(7/2))/7 - (2*(1 - x)^(9/2))/9`

3.267.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.267.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.45

method	result	size
trager	$\left(-\frac{2}{9}x^4 - \frac{16}{63}x^3 - \frac{32}{105}x^2 - \frac{128}{315}x - \frac{256}{315}\right) \sqrt{1-x}$	29
gosper	$-\frac{2\sqrt{1-x}(35x^4+40x^3+48x^2+64x+128)}{315}$	30
pseudoelliptic	$-\frac{2\sqrt{1-x}(35x^4+40x^3+48x^2+64x+128)}{315}$	30
risch	$\frac{2(-1+x)(35x^4+40x^3+48x^2+64x+128)}{315\sqrt{1-x}}$	33
meijerg	$-\frac{\frac{256\sqrt{\pi}}{315} + \frac{\sqrt{\pi}(70x^4+80x^3+96x^2+128x+256)\sqrt{1-x}}{315}}{\sqrt{\pi}}$	44
derivativedivides	$-2\sqrt{1-x} + \frac{8(1-x)^{\frac{3}{2}}}{3} - \frac{12(1-x)^{\frac{5}{2}}}{5} + \frac{8(1-x)^{\frac{7}{2}}}{7} - \frac{2(1-x)^{\frac{9}{2}}}{9}$	47
default	$-2\sqrt{1-x} + \frac{8(1-x)^{\frac{3}{2}}}{3} - \frac{12(1-x)^{\frac{5}{2}}}{5} + \frac{8(1-x)^{\frac{7}{2}}}{7} - \frac{2(1-x)^{\frac{9}{2}}}{9}$	47

input `int(x^4/(1-x)^(1/2),x,method=_RETURNVERBOSE)`output $(-2/9*x^4-16/63*x^3-32/105*x^2-128/315*x-256/315)*(1-x)^(1/2)$ **3.267.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.45

$$\int \frac{x^4}{\sqrt{1-x}} dx = -\frac{2}{315} (35x^4 + 40x^3 + 48x^2 + 64x + 128) \sqrt{-x+1}$$

input `integrate(x^4/(1-x)^(1/2),x, algorithm="fricas")`output $-2/315*(35*x^4 + 40*x^3 + 48*x^2 + 64*x + 128)*sqrt(-x + 1)$

3.267.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.70 (sec) , antiderivative size = 138, normalized size of antiderivative = 2.16

$$\int \frac{x^4}{\sqrt{1-x}} dx = \begin{cases} -\frac{2ix^4\sqrt{x-1}}{9} - \frac{16ix^3\sqrt{x-1}}{63} - \frac{32ix^2\sqrt{x-1}}{105} - \frac{128ix\sqrt{x-1}}{315} - \frac{256i\sqrt{x-1}}{315} & \text{for } |x| > 1 \\ -\frac{2x^4\sqrt{1-x}}{9} - \frac{16x^3\sqrt{1-x}}{63} - \frac{32x^2\sqrt{1-x}}{105} - \frac{128x\sqrt{1-x}}{315} - \frac{256\sqrt{1-x}}{315} & \text{otherwise} \end{cases}$$

input `integrate(x**4/(1-x)**(1/2),x)`

output `Piecewise((-2*I*x**4*sqrt(x - 1)/9 - 16*I*x**3*sqrt(x - 1)/63 - 32*I*x**2*sqrt(x - 1)/105 - 128*I*x*sqrt(x - 1)/315 - 256*I*sqrt(x - 1)/315, Abs(x) > 1), (-2*x**4*sqrt(1 - x)/9 - 16*x**3*sqrt(1 - x)/63 - 32*x**2*sqrt(1 - x)/105 - 128*x*sqrt(1 - x)/315 - 256*sqrt(1 - x)/315, True))`

3.267.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.72

$$\int \frac{x^4}{\sqrt{1-x}} dx = -\frac{2}{9}(-x+1)^{\frac{9}{2}} + \frac{8}{7}(-x+1)^{\frac{7}{2}} - \frac{12}{5}(-x+1)^{\frac{5}{2}} + \frac{8}{3}(-x+1)^{\frac{3}{2}} - 2\sqrt{-x+1}$$

input `integrate(x^4/(1-x)^(1/2),x, algorithm="maxima")`

output `-2/9*(-x + 1)^(9/2) + 8/7*(-x + 1)^(7/2) - 12/5*(-x + 1)^(5/2) + 8/3*(-x + 1)^(3/2) - 2*sqrt(-x + 1)`

3.267.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.95

$$\int \frac{x^4}{\sqrt{1-x}} dx = -\frac{2}{9}(x-1)^4\sqrt{-x+1} - \frac{8}{7}(x-1)^3\sqrt{-x+1} - \frac{12}{5}(x-1)^2\sqrt{-x+1} + \frac{8}{3}(-x+1)^{\frac{3}{2}} - 2\sqrt{-x+1}$$

input `integrate(x^4/(1-x)^(1/2),x, algorithm="giac")`

output `-2/9*(x - 1)^4*sqrt(-x + 1) - 8/7*(x - 1)^3*sqrt(-x + 1) - 12/5*(x - 1)^2*sqrt(-x + 1) + 8/3*(-x + 1)^(3/2) - 2*sqrt(-x + 1)`

3.267.9 Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.72

$$\int \frac{x^4}{\sqrt{1-x}} dx = \frac{8(1-x)^{3/2}}{3} - 2\sqrt{1-x} - \frac{12(1-x)^{5/2}}{5} + \frac{8(1-x)^{7/2}}{7} - \frac{2(1-x)^{9/2}}{9}$$

input `int(x^4/(1 - x)^(1/2),x)`

output `(8*(1 - x)^(3/2))/3 - 2*(1 - x)^(1/2) - (12*(1 - x)^(5/2))/5 + (8*(1 - x)^(7/2))/7 - (2*(1 - x)^(9/2))/9`

3.268 $\int \sin(x^{-n}) dx$

3.268.1 Optimal result	1352
3.268.2 Mathematica [A] (verified)	1352
3.268.3 Rubi [A] (verified)	1353
3.268.4 Maple [C] (verified)	1354
3.268.5 Fricas [F]	1354
3.268.6 Sympy [B] (verification not implemented)	1354
3.268.7 Maxima [F]	1355
3.268.8 Giac [F]	1355
3.268.9 Mupad [F(-1)]	1355

3.268.1 Optimal result

Integrand size = 6, antiderivative size = 43

$$\int \sin(x^{-n}) dx = \frac{ix(-\text{ExpIntegralE}(1 + \frac{1}{n}, -ix^{-n}) + \text{ExpIntegralE}(1 + \frac{1}{n}, ix^{-n}))}{2n}$$

output `1/2*I*x*(-Ei(1+1/n,-I/(x^n))+Ei(1+1/n,I/(x^n)))/n`

3.268.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.63

$$\int \sin(x^{-n}) dx = -\frac{ix\left((-ix^{-n})^{\frac{1}{n}} \Gamma(-\frac{1}{n}, -ix^{-n}) - (ix^{-n})^{\frac{1}{n}} \Gamma(-\frac{1}{n}, ix^{-n})\right)}{2n}$$

input `Integrate[Sin[x^(-n)],x]`

output `((-1/2*I)*x*(((I/x^n)^n*(-1)*Gamma[-n^(-1), (-I)/x^n] - (I/x^n)^n*(-1)*Gamma[-n^(-1), I/x^n]))/n`

3.268.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.79, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3846, 2637}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sin(x^{-n}) dx$$

$$\downarrow \text{3846}$$

$$\frac{1}{2}i \int e^{-ix^{-n}} dx - \frac{1}{2}i \int e^{ix^{-n}} dx$$

$$\downarrow \text{2637}$$

$$\frac{ix(ix^{-n})^{\frac{1}{n}} \Gamma(-\frac{1}{n}, ix^{-n})}{2n} - \frac{ix(-ix^{-n})^{\frac{1}{n}} \Gamma(-\frac{1}{n}, -ix^{-n})}{2n}$$

input `Int[Sin[x^(-n)], x]`

output `((-1/2*I)*x*((-I)/x^n)^n^(-1)*Gamma[-n^(-1), (-I)/x^n])/n + ((I/2)*x*(I/x^n)^n^(-1)*Gamma[-n^(-1), I/x^n])/n`

3.268.3.1 Defintions of rubi rules used

rule 2637 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_))), x_Symbol] := Simp[(-F^a)*(c + d*x)*(Gamma[1/n, (-b)*(c + d*x)^n*Log[F]]/(d*n*((-b)*(c + d*x)^n*Log[F])^(1/n))), x] /; FreeQ[{F, a, b, c, d, n}, x] && !IntegerQ[2/n]`

rule 3846 `Int[Sin[(c_.) + (d_.)*((e_.) + (f_.)*(x_)^(n_))], x_Symbol] := Simp[I/2 Int[E^((-c)*I - d*I*(e + f*x)^n), x], x] - Simp[I/2 Int[E^(c*I + d*I*(e + f*x)^n), x], x] /; FreeQ[{c, d, e, f, n}, x]`

3.268.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.25 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.09

method	result	size
meijerg	$-\frac{x^{-n+1} \operatorname{hypergeom}\left(\left[\frac{1}{2}-\frac{1}{2n}\right], \left[\frac{3}{2}, \frac{3}{2}-\frac{1}{2n}\right], -\frac{x^{-4n}x^{2n}}{4}\right)}{n-1}$	47

input `int(sin(1/(x^n)),x,method=_RETURNVERBOSE)`

output `-1/(n-1)*x^(-n+1)*hypergeom([1/2-1/2/n],[3/2,3/2-1/2/n],-1/4*(x^(-2*n))^2*x^(2*n))`

3.268.5 Fracas [F]

$$\int \sin(x^{-n}) dx = \int \sin\left(\frac{1}{x^n}\right) dx$$

input `integrate(sin(1/(x^n)),x, algorithm="fricas")`

output `integral(sin(1/(x^n)), x)`

3.268.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. $2(27) = 54$.

Time = 0.73 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.30

$$\int \sin(x^{-n}) dx = -\frac{x^{1-n} \Gamma\left(\frac{1}{2} - \frac{1}{2n}\right) {}_1F_2\left(\frac{1}{2} - \frac{1}{2n} \mid -\frac{x^{-2n}}{4}\right)}{2n \Gamma\left(\frac{3}{2} - \frac{1}{2n}\right)}$$

input `integrate(sin(1/(x**n)),x)`

output `-x**(1-n)*gamma(1/2-1/(2*n))*hyper((1/2-1/(2*n)),(3/2,3/2-1/(2*n)),-1/(4*x**(2*n)))/(2*n*gamma(3/2-1/(2*n)))`

3.268.7 Maxima [F]

$$\int \sin(x^{-n}) dx = \int \sin\left(\frac{1}{x^n}\right) dx$$

input `integrate(sin(1/(x^n)),x, algorithm="maxima")`

output `integrate(sin(1/(x^n)), x)`

3.268.8 Giac [F]

$$\int \sin(x^{-n}) dx = \int \sin\left(\frac{1}{x^n}\right) dx$$

input `integrate(sin(1/(x^n)),x, algorithm="giac")`

output `integrate(sin(1/(x^n)), x)`

3.268.9 Mupad [F(-1)]

Timed out.

$$\int \sin(x^{-n}) dx = \int \sin\left(\frac{1}{x^n}\right) dx$$

input `int(sin(1/x^n),x)`

output `int(sin(1/x^n), x)`

3.269 $\int \frac{1}{2}(-x + \sqrt{4 - 3x^2}) dx$

3.269.1 Optimal result	1356
3.269.2 Mathematica [A] (verified)	1356
3.269.3 Rubi [A] (verified)	1357
3.269.4 Maple [A] (verified)	1358
3.269.5 Fricas [A] (verification not implemented)	1358
3.269.6 Sympy [A] (verification not implemented)	1358
3.269.7 Maxima [A] (verification not implemented)	1359
3.269.8 Giac [A] (verification not implemented)	1359
3.269.9 Mupad [B] (verification not implemented)	1359

3.269.1 Optimal result

Integrand size = 19, antiderivative size = 41

$$\int \frac{1}{2}(-x + \sqrt{4 - 3x^2}) dx = -\frac{x^2}{4} + \frac{1}{4}x\sqrt{4 - 3x^2} + \frac{\arcsin\left(\frac{\sqrt{3}x}{2}\right)}{\sqrt{3}}$$

output `-1/4*x^2+1/4*x*(-3*x^2+4)^(1/2)+1/3*arcsin(1/2*x*3^(1/2))*3^(1/2)`

3.269.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.41

$$\int \frac{1}{2}(-x + \sqrt{4 - 3x^2}) dx = \frac{1}{2} \left(-\frac{x^2}{2} + \frac{1}{2}x\sqrt{4 - 3x^2} + \frac{4 \arctan\left(\frac{\sqrt{3}x}{-2 + \sqrt{4 - 3x^2}}\right)}{\sqrt{3}} \right)$$

input `Integrate[(-x + Sqrt[4 - 3*x^2])/2,x]`

output `(-1/2*x^2 + (x*Sqrt[4 - 3*x^2])/2 + (4*ArcTan[(Sqrt[3]*x)/(-2 + Sqrt[4 - 3*x^2])])/Sqrt[3])/2`

3.269.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.12, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {27, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{2} (\sqrt{4-3x^2} - x) dx$$

$$\downarrow 27$$

$$\frac{1}{2} \int (\sqrt{4-3x^2} - x) dx$$

$$\downarrow 2009$$

$$\frac{1}{2} \left(\frac{2 \arcsin\left(\frac{\sqrt{3}x}{2}\right)}{\sqrt{3}} - \frac{x^2}{2} + \frac{1}{2} \sqrt{4-3x^2} \right)$$

input `Int[(-x + Sqrt[4 - 3*x^2])/2,x]`

output `(-1/2*x^2 + (x*Sqrt[4 - 3*x^2])/2 + (2*ArcSin[(Sqrt[3]*x)/2])/Sqrt[3])/2`

3.269.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.269.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

method	result	size
default	$-\frac{x^2}{4} + \frac{x\sqrt{-3x^2+4}}{4} + \frac{\arcsin\left(\frac{x\sqrt{3}}{2}\right)\sqrt{3}}{3}$	31
parts	$-\frac{x^2}{4} + \frac{x\sqrt{-3x^2+4}}{4} + \frac{\arcsin\left(\frac{x\sqrt{3}}{2}\right)\sqrt{3}}{3}$	31
trager	$-\frac{x^2}{4} + \frac{x\sqrt{-3x^2+4}}{4} + \frac{\text{RootOf}(-Z^2+3)\ln(\text{RootOf}(-Z^2+3)\sqrt{-3x^2+4+3x})}{3}$	48

input `int(-1/2*x+1/2*(-3*x^2+4)^(1/2),x,method=_RETURNVERBOSE)`output `-1/4*x^2+1/4*x*(-3*x^2+4)^(1/2)+1/3*arcsin(1/2*x*3^(1/2))*3^(1/2)`**3.269.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.17

$$\int \frac{1}{2}(-x + \sqrt{4 - 3x^2}) dx = -\frac{1}{4}x^2 + \frac{1}{4}\sqrt{-3x^2 + 4}x - \frac{2}{3}\sqrt{3}\arctan\left(\frac{\sqrt{3}\sqrt{-3x^2 + 4} - 2\sqrt{3}}{3x}\right)$$

input `integrate(-1/2*x+1/2*(-3*x^2+4)^(1/2),x, algorithm="fricas")`output `-1/4*x^2 + 1/4*sqrt(-3*x^2 + 4)*x - 2/3*sqrt(3)*arctan(1/3*(sqrt(3)*sqrt(-3*x^2 + 4) - 2*sqrt(3))/x)`**3.269.6 Sympy [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \frac{1}{2}(-x + \sqrt{4 - 3x^2}) dx = -\frac{x^2}{4} + \frac{x\sqrt{4 - 3x^2}}{4} + \frac{\sqrt{3}\operatorname{asin}\left(\frac{\sqrt{3}x}{2}\right)}{3}$$

input `integrate(-1/2*x+1/2*(-3*x**2+4)**(1/2),x)`output `-x**2/4 + x*sqrt(4 - 3*x**2)/4 + sqrt(3)*asin(sqrt(3)*x/2)/3`

3.269.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.73

$$\int \frac{1}{2}(-x + \sqrt{4 - 3x^2}) dx = -\frac{1}{4}x^2 + \frac{1}{4}\sqrt{-3x^2 + 4x} + \frac{1}{3}\sqrt{3}\arcsin\left(\frac{1}{2}\sqrt{3}x\right)$$

input `integrate(-1/2*x+1/2*(-3*x^2+4)^(1/2),x, algorithm="maxima")`output `-1/4*x^2 + 1/4*sqrt(-3*x^2 + 4)*x + 1/3*sqrt(3)*arcsin(1/2*sqrt(3)*x)`**3.269.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.73

$$\int \frac{1}{2}(-x + \sqrt{4 - 3x^2}) dx = -\frac{1}{4}x^2 + \frac{1}{4}\sqrt{-3x^2 + 4x} + \frac{1}{3}\sqrt{3}\arcsin\left(\frac{1}{2}\sqrt{3}x\right)$$

input `integrate(-1/2*x+1/2*(-3*x^2+4)^(1/2),x, algorithm="giac")`output `-1/4*x^2 + 1/4*sqrt(-3*x^2 + 4)*x + 1/3*sqrt(3)*arcsin(1/2*sqrt(3)*x)`**3.269.9 Mupad [B] (verification not implemented)**

Time = 15.21 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

$$\int \frac{1}{2}(-x + \sqrt{4 - 3x^2}) dx = \frac{\sqrt{3}\left(\operatorname{asin}\left(\frac{\sqrt{3}x}{2}\right) + \frac{3x\sqrt{\frac{4}{3}-x^2}}{4}\right)}{3} - \frac{x^2}{4}$$

input `int((4 - 3*x^2)^(1/2)/2 - x/2,x)`output `(3^(1/2)*(asin((3^(1/2)*x)/2) + (3*x*(4/3 - x^2)^(1/2))/4)/3 - x^2/4`

3.270 $\int (1 - 3x^2 + x^4)^n dx$

3.270.1 Optimal result	1360
3.270.2 Mathematica [A] (verified)	1360
3.270.3 Rubi [A] (verified)	1361
3.270.4 Maple [F]	1362
3.270.5 Fricas [F]	1362
3.270.6 Sympy [F]	1363
3.270.7 Maxima [F]	1363
3.270.8 Giac [F]	1363
3.270.9 Mupad [F(-1)]	1364

3.270.1 Optimal result

Integrand size = 12, antiderivative size = 99

$$\int (1 - 3x^2 + x^4)^n dx = x \left(1 - \frac{2x^2}{3 - \sqrt{5}}\right)^{-n} \left(1 - \frac{2x^2}{3 + \sqrt{5}}\right)^{-n} (1 - 3x^2 + x^4)^n \operatorname{AppellF1}\left(\frac{1}{2}, -n, -n, \frac{3}{2}, \frac{2x^2}{3 + \sqrt{5}}, \frac{2x^2}{3 - \sqrt{5}}\right)$$

```
output x*(x^4-3*x^2+1)^n*AppellF1(1/2,-n,-n,3/2,2*x^2/(3-5^(1/2)),2*x^2/(3+5^(1/2)))
/((1-2*x^2/(3-5^(1/2)))^n)/((1-2*x^2/(3+5^(1/2)))^n)
```

3.270.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.43

$$\int (1 - 3x^2 + x^4)^n dx = (3 + \sqrt{5})^n x \left(-\left(3 + \sqrt{5} - 2x^2\right)^2\right)^{-n} (-3 - \sqrt{5} + 2x^2)^n (-3 + \sqrt{5} + 2x^2)^n \left(\frac{(-3 + \sqrt{5} + 2x^2)^2}{-3 + \sqrt{5}}\right)^{-n} (1 - 3x^2 + x^4)^n \operatorname{AppellF1}\left(\frac{1}{2}, -n, -n, \frac{3}{2}, -\frac{2x^2}{-3 + \sqrt{5}}, \frac{2x^2}{3 + \sqrt{5}}\right)$$

```
input Integrate[(1 - 3*x^2 + x^4)^n,x]
```

output $((3 + \sqrt{5})^n x (-3 - \sqrt{5} + 2x^2)^n (-3 + \sqrt{5} + 2x^2)^n (1 - 3x^2 + x^4)^n \text{AppellF1}[1/2, -n, -n, 3/2, (-2x^2)/(-3 + \sqrt{5}), (2x^2)/(3 + \sqrt{5})]) / ((-3 + \sqrt{5} - 2x^2)^2)^n ((-3 + \sqrt{5} + 2x^2)^2 / (-3 + \sqrt{5}))^n$

3.270.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {1418, 333}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x^4 - 3x^2 + 1)^n dx$$

$$\downarrow 1418$$

$$\left(1 - \frac{2x^2}{3 - \sqrt{5}}\right)^{-n} \left(1 - \frac{2x^2}{3 + \sqrt{5}}\right)^{-n} (x^4 - 3x^2 + 1)^n \int \left(1 - \frac{2x^2}{3 - \sqrt{5}}\right)^n \left(1 - \frac{2x^2}{3 + \sqrt{5}}\right)^n dx$$

$$\downarrow 333$$

$$x \left(1 - \frac{2x^2}{3 - \sqrt{5}}\right)^{-n} \left(1 - \frac{2x^2}{3 + \sqrt{5}}\right)^{-n} (x^4 - 3x^2 + 1)^n \text{AppellF1}\left(\frac{1}{2}, -n, -n, \frac{3}{2}, \frac{2x^2}{3 - \sqrt{5}}, \frac{2x^2}{3 + \sqrt{5}}\right)$$

input $\text{Int}[(1 - 3x^2 + x^4)^n, x]$

output $(x(1 - 3x^2 + x^4)^n \text{AppellF1}[1/2, -n, -n, 3/2, (2x^2)/(3 - \sqrt{5}), (2x^2)/(3 + \sqrt{5})]) / ((1 - (2x^2)/(3 - \sqrt{5}))^n (1 - (2x^2)/(3 + \sqrt{5}))^n)$

3.270.3.1 Defintions of rubi rules used

rule 333 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp`
`p[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /;`
`FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a,`
`0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 1418 `Int[((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := With[{q = Rt[b^`
`2 - 4*a*c, 2]}, Simp[a^IntPart[p]*((a + b*x^2 + c*x^4)^FracPart[p]/((1 + 2*`
`c*(x^2/(b + q)))^FracPart[p]*(1 + 2*c*(x^2/(b - q)))^FracPart[p])) Int[(1`
`+ 2*c*(x^2/(b + q)))^p*(1 + 2*c*(x^2/(b - q)))^p, x], x] /;`
`FreeQ[{a, b, c, p}, x] && NeQ[b^2 - 4*a*c, 0]`

3.270.4 Maple [F]

$$\int (x^4 - 3x^2 + 1)^n dx$$

input `int((x^4-3*x^2+1)^n,x)`

output `int((x^4-3*x^2+1)^n,x)`

3.270.5 Fracas [F]

$$\int (1 - 3x^2 + x^4)^n dx = \int (x^4 - 3x^2 + 1)^n dx$$

input `integrate((x^4-3*x^2+1)^n,x, algorithm="fricas")`

output `integral((x^4 - 3*x^2 + 1)^n, x)`

3.270.6 Sympy [F]

$$\int (1 - 3x^2 + x^4)^n dx = \int (x^4 - 3x^2 + 1)^n dx$$

input `integrate((x**4-3*x**2+1)**n,x)`

output `Integral((x**4 - 3*x**2 + 1)**n, x)`

3.270.7 Maxima [F]

$$\int (1 - 3x^2 + x^4)^n dx = \int (x^4 - 3x^2 + 1)^n dx$$

input `integrate((x^4-3*x^2+1)^n,x, algorithm="maxima")`

output `integrate((x^4 - 3*x^2 + 1)^n, x)`

3.270.8 Giac [F]

$$\int (1 - 3x^2 + x^4)^n dx = \int (x^4 - 3x^2 + 1)^n dx$$

input `integrate((x^4-3*x^2+1)^n,x, algorithm="giac")`

output `integrate((x^4 - 3*x^2 + 1)^n, x)`

3.270.9 Mupad [F(-1)]

Timed out.

$$\int (1 - 3x^2 + x^4)^n dx = \int (x^4 - 3x^2 + 1)^n dx$$

input `int((x^4 - 3*x^2 + 1)^n,x)`output `int((x^4 - 3*x^2 + 1)^n, x)`

3.271 $\int \frac{(1+e^{-x})x}{-1+e^x} dx$

3.271.1 Optimal result	1365
3.271.2 Mathematica [A] (verified)	1365
3.271.3 Rubi [A] (verified)	1366
3.271.4 Maple [A] (verified)	1367
3.271.5 Fricas [A] (verification not implemented)	1367
3.271.6 Sympy [F]	1367
3.271.7 Maxima [A] (verification not implemented)	1368
3.271.8 Giac [F]	1368
3.271.9 Mupad [F(-1)]	1368

3.271.1 Optimal result

Integrand size = 16, antiderivative size = 36

$$\int \frac{(1 + e^{-x})x}{-1 + e^x} dx = e^{-x} + e^{-x}x - x^2 + 2x \log(1 - e^x) + 2 \text{PolyLog}(2, e^x)$$

output `exp(-x)+x/exp(x)-x^2+2*x*ln(1-exp(x))+2*polylog(2,exp(x))`

3.271.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00

$$\int \frac{(1 + e^{-x})x}{-1 + e^x} dx = e^{-x}(1 + x - e^x x^2) + 2x \log(1 - e^x) + 2 \text{PolyLog}(2, e^x)$$

input `Integrate[((1 + E^(-x))*x)/(-1 + E^x),x]`

output `(1 + x - E^x*x^2)/E^x + 2*x*Log[1 - E^x] + 2*PolyLog[2, E^x]`

3.271.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2684, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(e^{-x} + 1)x}{e^x - 1} dx$$

↓ 2684

$$\int \left(\frac{2x}{e^x - 1} - e^{-x}x \right) dx$$

↓ 2009

$$2 \text{PolyLog}(2, e^x) - x^2 + e^{-x}x + e^{-x} + 2x \log(1 - e^x)$$

input `Int[((1 + E^(-x))*x)/(-1 + E^x),x]`

output `E^(-x) + x/E^x - x^2 + 2*x*Log[1 - E^x] + 2*PolyLog[2, E^x]`

3.271.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2684 `Int[((a_.) + (b_.)*(F_)^(u_))^(p_.)*((c_.) + (d_.)*(F_)^(v_))^(q_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := With[{w = ExpandIntegrand[(e + f*x)^m, (a + b*F^u)^p*(c + d*F^v)^q, x]}, Int[w, x] /; SumQ[w]] /; FreeQ[{F, a, b, c, d, e, f, m}, x] && IntegersQ[p, q] && LinearQ[{u, v}, x] && RationalQ[Simplify[u/v]]`

3.271.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

method	result	size
risch	$(1+x)e^{-x} - x^2 + 2x \ln(1 - e^x) + 2 \operatorname{polylog}(2, e^x)$	31
default	$xe^{-x} + 2x \ln(1 - e^x) - x^2 + e^{-x} + 2 \operatorname{polylog}(2, e^x)$	33

input `int(x*(exp(-x)+1)/(exp(x)-1),x,method=_RETURNVERBOSE)`output `(1+x)/exp(x)-x^2+2*x*ln(1-exp(x))+2*polylog(2,exp(x))`**3.271.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00

$$\int \frac{(1+e^{-x})x}{-1+e^x} dx = -(x^2 e^x - 2x e^x \log(-e^x + 1) - 2 \operatorname{Li}_2(e^x) e^x - x - 1) e^{(-x)}$$

input `integrate(x*(exp(-x)+1)/(exp(x)-1),x, algorithm="fricas")`output `-(x^2*e^x - 2*x*e^x*log(-e^x + 1) - 2*dilog(e^x)*e^x - x - 1)*e^(-x)`**3.271.6 Sympy [F]**

$$\int \frac{(1+e^{-x})x}{-1+e^x} dx = \int \frac{x(e^x + 1)e^{-x}}{e^x - 1} dx$$

input `integrate(x*(exp(-x)+1)/(exp(x)-1),x)`output `Integral(x*(exp(x) + 1)*exp(-x)/(exp(x) - 1), x)`

3.271.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.81

$$\int \frac{(1 + e^{-x})x}{-1 + e^x} dx = -x^2 + (x + 1)e^{(-x)} + 2x \log(-e^x + 1) + 2\text{Li}_2(e^x)$$

input `integrate(x*(exp(-x)+1)/(exp(x)-1),x, algorithm="maxima")`output `-x^2 + (x + 1)*e^(-x) + 2*x*log(-e^x + 1) + 2*dilog(e^x)`**3.271.8 Giac [F]**

$$\int \frac{(1 + e^{-x})x}{-1 + e^x} dx = \int \frac{x(e^{-x} + 1)}{e^x - 1} dx$$

input `integrate(x*(exp(-x)+1)/(exp(x)-1),x, algorithm="giac")`output `integrate(x*(e^(-x) + 1)/(e^x - 1), x)`**3.271.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(1 + e^{-x})x}{-1 + e^x} dx = \int \frac{x(e^{-x} + 1)}{e^x - 1} dx$$

input `int((x*(exp(-x) + 1))/(exp(x) - 1),x)`output `int((x*(exp(-x) + 1))/(exp(x) - 1), x)`

3.272 $\int e^{-x} x^4 \sin(x) dx$

3.272.1 Optimal result	1369
3.272.2 Mathematica [A] (verified)	1369
3.272.3 Rubi [B] (verified)	1370
3.272.4 Maple [A] (verified)	1371
3.272.5 Fricas [A] (verification not implemented)	1372
3.272.6 Sympy [B] (verification not implemented)	1372
3.272.7 Maxima [A] (verification not implemented)	1372
3.272.8 Giac [A] (verification not implemented)	1373
3.272.9 Mupad [B] (verification not implemented)	1373

3.272.1 Optimal result

Integrand size = 11, antiderivative size = 44

$$\int e^{-x} x^4 \sin(x) dx = \frac{1}{2} e^{-x} (- ((-6 + x^2(6 + x(4 + x))) \cos(x)) + (6 + x(12 + 6x - x^3)) \sin(x))$$

output `1/2*(-(-6+x^2*(6+(4+x)*x))*cos(x)+(6+x*(-x^3+6*x+12))*sin(x))/exp(x)`

3.272.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.07

$$\int e^{-x} x^4 \sin(x) dx = \frac{1}{2} e^{-x} (- ((-6 + 6x^2 + 4x^3 + x^4) \cos(x)) + (6 + 12x + 6x^2 - x^4) \sin(x))$$

input `Integrate[(x^4*Sin[x])/E^x,x]`

output `(-((-6 + 6*x^2 + 4*x^3 + x^4)*Cos[x]) + (6 + 12*x + 6*x^2 - x^4)*Sin[x])/ (2*E^x)`

3.272.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 104 vs. $2(44) = 88$.

Time = 0.76 (sec) , antiderivative size = 104, normalized size of antiderivative = 2.36, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {4968, 27, 2010, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{-x} x^4 \sin(x) dx \\
 & \quad \downarrow \text{4968} \\
 & -4 \int -\frac{1}{2} x^3 (e^{-x} \cos(x) + e^{-x} \sin(x)) dx - \frac{1}{2} e^{-x} x^4 \sin(x) - \frac{1}{2} e^{-x} x^4 \cos(x) \\
 & \quad \downarrow \text{27} \\
 & 2 \int x^3 (e^{-x} \cos(x) + e^{-x} \sin(x)) dx - \frac{1}{2} e^{-x} x^4 \sin(x) - \frac{1}{2} e^{-x} x^4 \cos(x) \\
 & \quad \downarrow \text{2010} \\
 & 2 \int (e^{-x} \cos(x) x^3 + e^{-x} \sin(x) x^3) dx - \frac{1}{2} e^{-x} x^4 \sin(x) - \frac{1}{2} e^{-x} x^4 \cos(x) \\
 & \quad \downarrow \text{2009} \\
 & -\frac{1}{2} e^{-x} x^4 \sin(x) - \frac{1}{2} e^{-x} x^4 \cos(x) + \\
 & 2 \left(-e^{-x} x^3 \cos(x) + \frac{3}{2} e^{-x} x^2 \sin(x) - \frac{3}{2} e^{-x} x^2 \cos(x) + 3e^{-x} x \sin(x) + \frac{3}{2} e^{-x} \sin(x) + \frac{3}{2} e^{-x} \cos(x) \right)
 \end{aligned}$$

input `Int[(x^4*Sin[x])/E^x,x]`

output `-1/2*(x^4*Cos[x])/E^x - (x^4*Sin[x])/(2*E^x) + 2*((3*Cos[x])/(2*E^x) - (3*x^2*Cos[x])/(2*E^x) - (x^3*Cos[x])/E^x + (3*Sin[x])/(2*E^x) + (3*x*Sin[x])/E^x + (3*x^2*Sin[x])/(2*E^x))`

3.272.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2010 `Int[(u_)*((c_)*(x_)^(m_)), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]`

rule 4968 `Int[(F_)^((c_)*((a_) + (b_)*(x_)))*((f_)*(x_)^(m_)*Sin[(d_) + (e_)*(x_)^(n_)], x_Symbol] := Module[{u = IntHide[F^(c*(a + b*x))*Sin[d + e*x]^n, x]}, Simp[(f*x)^m u, x] - Simp[f^m Int[(f*x)^(m - 1)*u, x], x] /; FreeQ[{F, a, b, c, d, e, f}, x] && IGtQ[n, 0] && GtQ[m, 0]`

3.272.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.95

method	result
parallelrisch	$-\frac{e^{-x}((x^4+4x^3+6x^2-6)\cos(x)+\sin(x)(x^4-6x^2-12x-6))}{2}$
default	$\left(-\frac{1}{2}x^4 - 2x^3 - 3x^2 + 3\right)e^{-x}\cos(x) + \left(-\frac{1}{2}x^4 + 3x^2 + 6x + 3\right)e^{-x}\sin(x)$
risch	$\left(-\frac{1}{4} + \frac{i}{4}\right)(x^4 + 2ix^3 + 2x^3 + 6ix^2 + 6ix - 6x - 6)e^{(-1+i)x} + \left(-\frac{1}{4} - \frac{i}{4}\right)(x^4 - 2ix^3 + 2x^3 - 6ix^2 - 6ix + 6x + 6)e^{(-1-i)x}$
norman	$\frac{-3x^2e^{-x} - 2x^3e^{-x} - \frac{x^4e^{-x}}{2} + 6e^{-x}\tan\left(\frac{x}{2}\right) - 3e^{-x}\tan\left(\frac{x}{2}\right)^2 + 12xe^{-x}\tan\left(\frac{x}{2}\right) + 6x^2e^{-x}\tan\left(\frac{x}{2}\right) + 3x^2e^{-x}\tan\left(\frac{x}{2}\right)^2 + 2x^3e^{-x}\tan\left(\frac{x}{2}\right)}{1 + \tan\left(\frac{x}{2}\right)^2}$

input `int(x^4*exp(-x)*sin(x),x,method=_RETURNVERBOSE)`

output `-1/2*exp(-x)*((x^4+4*x^3+6*x^2-6)*cos(x)+sin(x)*(x^4-6*x^2-12*x-6))`

3.272.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.02

$$\int e^{-x} x^4 \sin(x) dx = -\frac{1}{2} (x^4 + 4x^3 + 6x^2 - 6) \cos(x) e^{(-x)} - \frac{1}{2} (x^4 - 6x^2 - 12x - 6) e^{(-x)} \sin(x)$$

input `integrate(x^4*exp(-x)*sin(x),x, algorithm="fricas")`

output `-1/2*(x^4 + 4*x^3 + 6*x^2 - 6)*cos(x)*e^(-x) - 1/2*(x^4 - 6*x^2 - 12*x - 6)*e^(-x)*sin(x)`

3.272.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 85 vs. 2(36) = 72.

Time = 1.03 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.93

$$\int e^{-x} x^4 \sin(x) dx = -\frac{x^4 e^{-x} \sin(x)}{2} - \frac{x^4 e^{-x} \cos(x)}{2} - 2x^3 e^{-x} \cos(x) + 3x^2 e^{-x} \sin(x) - 3x^2 e^{-x} \cos(x) + 6x e^{-x} \sin(x) + 3e^{-x} \sin(x) + 3e^{-x} \cos(x)$$

input `integrate(x**4*exp(-x)*sin(x),x)`

output `-x**4*exp(-x)*sin(x)/2 - x**4*exp(-x)*cos(x)/2 - 2*x**3*exp(-x)*cos(x) + 3*x**2*exp(-x)*sin(x) - 3*x**2*exp(-x)*cos(x) + 6*x*exp(-x)*sin(x) + 3*exp(-x)*sin(x) + 3*exp(-x)*cos(x)`

3.272.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.93

$$\int e^{-x} x^4 \sin(x) dx = -\frac{1}{2} ((x^4 + 4x^3 + 6x^2 - 6) \cos(x) + (x^4 - 6x^2 - 12x - 6) \sin(x)) e^{(-x)}$$

input `integrate(x^4*exp(-x)*sin(x),x, algorithm="maxima")`

output `-1/2*((x^4 + 4*x^3 + 6*x^2 - 6)*cos(x) + (x^4 - 6*x^2 - 12*x - 6)*sin(x))*e^(-x)`

3.272.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.93

$$\int e^{-x} x^4 \sin(x) dx = -\frac{1}{2} \left((x^4 + 4x^3 + 6x^2 - 6) \cos(x) + (x^4 - 6x^2 - 12x - 6) \sin(x) \right) e^{-x}$$

input `integrate(x^4*exp(-x)*sin(x),x, algorithm="giac")`output `-1/2*((x^4 + 4*x^3 + 6*x^2 - 6)*cos(x) + (x^4 - 6*x^2 - 12*x - 6)*sin(x))*
e^(-x)`**3.272.9 Mupad [B] (verification not implemented)**

Time = 16.53 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.25

$$\int e^{-x} x^4 \sin(x) dx = \frac{e^{-x} (6 \cos(x) + 6 \sin(x) - 6x^2 \cos(x) - 4x^3 \cos(x) - x^4 \cos(x) + 6x^2 \sin(x) - x^4 \sin(x) + 12x \sin(x))}{2}$$

input `int(x^4*exp(-x)*sin(x),x)`output `(exp(-x)*(6*cos(x) + 6*sin(x) - 6*x^2*cos(x) - 4*x^3*cos(x) - x^4*cos(x) +
6*x^2*sin(x) - x^4*sin(x) + 12*x*sin(x)))/2`

$$3.273 \quad \int \frac{-3(-3x+x^3)+(-3x+x^3)^3}{\sqrt{-4+x^2}} dx$$

3.273.1 Optimal result	1374
3.273.2 Mathematica [A] (verified)	1374
3.273.3 Rubi [A] (verified)	1375
3.273.4 Maple [A] (verified)	1376
3.273.5 Fricas [A] (verification not implemented)	1377
3.273.6 Sympy [A] (verification not implemented)	1377
3.273.7 Maxima [A] (verification not implemented)	1377
3.273.8 Giac [A] (verification not implemented)	1378
3.273.9 Mupad [B] (verification not implemented)	1378

3.273.1 Optimal result

Integrand size = 29, antiderivative size = 56

$$\int \frac{-3(-3x+x^3)+(-3x+x^3)^3}{\sqrt{-4+x^2}} dx = \sqrt{-4+x^2} + \frac{10}{3}(-4+x^2)^{3/2} + 3(-4+x^2)^{5/2} + (-4+x^2)^{7/2} + \frac{1}{9}(-4+x^2)^{9/2}$$

output $(x^2-4)^{(1/2)}+10/3*(x^2-4)^{(3/2)}+3*(x^2-4)^{(5/2)}+(x^2-4)^{(7/2)}+1/9*(x^2-4)^{(9/2)}$

3.273.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.59

$$\int \frac{-3(-3x+x^3)+(-3x+x^3)^3}{\sqrt{-4+x^2}} dx = \frac{1}{9}\sqrt{-4+x^2}(1-10x^2+15x^4-7x^6+x^8)$$

input `Integrate[(-3*(-3*x + x^3) + (-3*x + x^3)^3)/Sqrt[-4 + x^2], x]`

output $(\text{Sqrt}[-4 + x^2]*(1 - 10*x^2 + 15*x^4 - 7*x^6 + x^8))/9$

$$3.273. \quad \int \frac{-3(-3x+x^3)+(-3x+x^3)^3}{\sqrt{-4+x^2}} dx$$

3.273.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.14, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.138$, Rules used = {2342, 2331, 2389, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(x^3 - 3x)^3 - 3(x^3 - 3x)}{\sqrt{x^2 - 4}} dx \\
 & \quad \downarrow \text{2342} \\
 & \int \frac{x(x^8 - 9x^6 + 27x^4 - 30x^2 + 9)}{\sqrt{x^2 - 4}} dx \\
 & \quad \downarrow \text{2331} \\
 & \frac{1}{2} \int \frac{x^8 - 9x^6 + 27x^4 - 30x^2 + 9}{\sqrt{x^2 - 4}} dx^2 \\
 & \quad \downarrow \text{2389} \\
 & \frac{1}{2} \int \left((x^2 - 4)^{7/2} + 7(x^2 - 4)^{5/2} + 15(x^2 - 4)^{3/2} + 10\sqrt{x^2 - 4} + \frac{1}{\sqrt{x^2 - 4}} \right) dx^2 \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} \left(\frac{2}{9}(x^2 - 4)^{9/2} + 2(x^2 - 4)^{7/2} + 6(x^2 - 4)^{5/2} + \frac{20}{3}(x^2 - 4)^{3/2} + 2\sqrt{x^2 - 4} \right)
 \end{aligned}$$

input `Int[(-3*(-3*x + x^3) + (-3*x + x^3)^3)/Sqrt[-4 + x^2],x]`

output `(2*Sqrt[-4 + x^2] + (20*(-4 + x^2)^(3/2))/3 + 6*(-4 + x^2)^(5/2) + 2*(-4 + x^2)^(7/2) + (2*(-4 + x^2)^(9/2))/9)/2`

3.273.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2331 Int[(Pq_)*(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 S
ubst[Int[x^((m - 1)/2)*SubstFor[x^2, Pq, x]*(a + b*x)^p, x], x, x^2], x] /;
FreeQ[{a, b, p}, x] && PolyQ[Pq, x^2] && IntegerQ[(m - 1)/2]
```

```
rule 2342 Int[(Pq_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[x*PolynomialQuotient
[Pq, x, x]*(a + b*x^2)^p, x] /; FreeQ[{a, b, p}, x] && PolyQ[Pq, x] && EqQ[
Coeff[Pq, x, 0], 0] && !MatchQ[Pq, x^(m_)*(u_)] /; IntegerQ[m]
```

```
rule 2389 Int[(Pq_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Int[ExpandIntegrand
[Pq*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, n}, x] && PolyQ[Pq, x] && (IGtQ[p
, 0] || EqQ[n, 1])
```

3.273.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.54

method	result
risch	$\frac{(x^8-7x^6+15x^4-10x^2+1)\sqrt{x^2-4}}{9}$
pseudoelliptic	$\frac{(x^8-7x^6+15x^4-10x^2+1)\sqrt{x^2-4}}{9}$
trager	$\left(\frac{1}{9}x^8 - \frac{7}{9}x^6 + \frac{5}{3}x^4 - \frac{10}{9}x^2 + \frac{1}{9}\right)\sqrt{x^2-4}$
gosper	$\frac{(x^8-7x^6+15x^4-10x^2+1)(-2+x)(2+x)}{9\sqrt{x^2-4}}$
default	$\frac{x^8\sqrt{x^2-4}}{9} - \frac{7x^6\sqrt{x^2-4}}{9} + \frac{5x^4\sqrt{x^2-4}}{3} - \frac{10x^2\sqrt{x^2-4}}{9} + \frac{\sqrt{x^2-4}}{9}$
meijerg	$-\frac{256\sqrt{-\operatorname{signum}\left(-1+\frac{x^2}{4}\right)}\left(-\frac{256\sqrt{\pi}}{315} + \frac{\sqrt{\pi}\left(\frac{35}{128}x^8 + \frac{5}{4}x^6 + 6x^4 + 32x^2 + 256\right)\sqrt{-\frac{x^2}{4}+1}}{315}\right)}{\sqrt{\pi}\sqrt{\operatorname{signum}\left(-1+\frac{x^2}{4}\right)}} - \frac{576\sqrt{-\operatorname{signum}\left(-1+\frac{x^2}{4}\right)}\left(\frac{32\sqrt{\pi}}{35}\right)}{\sqrt{\pi}\sqrt{\operatorname{signum}\left(-1+\frac{x^2}{4}\right)}}$

```
input int(((x^3-3*x)^3-3*x^3+9*x)/(x^2-4)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/9*(x^8-7*x^6+15*x^4-10*x^2+1)*(x^2-4)^(1/2)
```

3.273. $\int \frac{-3(-3x+x^3)+(-3x+x^3)^3}{\sqrt{-4+x^2}} dx$

3.273.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.52

$$\int \frac{-3(-3x + x^3) + (-3x + x^3)^3}{\sqrt{-4 + x^2}} dx = \frac{1}{9} (x^8 - 7x^6 + 15x^4 - 10x^2 + 1)\sqrt{x^2 - 4}$$

input `integrate(((x^3-3*x)^3-3*x^3+9*x)/(x^2-4)^(1/2),x, algorithm="fracas")`output `1/9*(x^8 - 7*x^6 + 15*x^4 - 10*x^2 + 1)*sqrt(x^2 - 4)`**3.273.6 Sympy [A] (verification not implemented)**

Time = 0.85 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.21

$$\int \frac{-3(-3x + x^3) + (-3x + x^3)^3}{\sqrt{-4 + x^2}} dx = \frac{x^8\sqrt{x^2 - 4}}{9} - \frac{7x^6\sqrt{x^2 - 4}}{9} + \frac{5x^4\sqrt{x^2 - 4}}{3} - \frac{10x^2\sqrt{x^2 - 4}}{9} + \frac{\sqrt{x^2 - 4}}{9}$$

input `integrate(((x**3-3*x)**3-3*x**3+9*x)/(x**2-4)**(1/2),x)`output `x**8*sqrt(x**2 - 4)/9 - 7*x**6*sqrt(x**2 - 4)/9 + 5*x**4*sqrt(x**2 - 4)/3 - 10*x**2*sqrt(x**2 - 4)/9 + sqrt(x**2 - 4)/9`**3.273.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.04

$$\int \frac{-3(-3x + x^3) + (-3x + x^3)^3}{\sqrt{-4 + x^2}} dx = \frac{1}{9} \sqrt{x^2 - 4} x^8 - \frac{7}{9} \sqrt{x^2 - 4} x^6 + \frac{5}{3} \sqrt{x^2 - 4} x^4 - \frac{10}{9} \sqrt{x^2 - 4} x^2 + \frac{1}{9} \sqrt{x^2 - 4}$$

input `integrate(((x^3-3*x)^3-3*x^3+9*x)/(x^2-4)^(1/2),x, algorithm="maxima")`output `1/9*sqrt(x^2 - 4)*x^8 - 7/9*sqrt(x^2 - 4)*x^6 + 5/3*sqrt(x^2 - 4)*x^4 - 10/9*sqrt(x^2 - 4)*x^2 + 1/9*sqrt(x^2 - 4)`

3.273. $\int \frac{-3(-3x+x^3)+(-3x+x^3)^3}{\sqrt{-4+x^2}} dx$

3.273.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.75

$$\int \frac{-3(-3x + x^3) + (-3x + x^3)^3}{\sqrt{-4 + x^2}} dx = \frac{1}{9} (x^2 - 4)^{\frac{9}{2}} + (x^2 - 4)^{\frac{7}{2}} + 3 (x^2 - 4)^{\frac{5}{2}} + \frac{10}{3} (x^2 - 4)^{\frac{3}{2}} + \sqrt{x^2 - 4}$$

input `integrate(((x^3-3*x)^3-3*x^3+9*x)/(x^2-4)^(1/2),x, algorithm="giac")`output `1/9*(x^2 - 4)^(9/2) + (x^2 - 4)^(7/2) + 3*(x^2 - 4)^(5/2) + 10/3*(x^2 - 4)^(3/2) + sqrt(x^2 - 4)`**3.273.9 Mupad [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.54

$$\int \frac{-3(-3x + x^3) + (-3x + x^3)^3}{\sqrt{-4 + x^2}} dx = \sqrt{x^2 - 4} \left(\frac{x^8}{9} - \frac{7x^6}{9} + \frac{5x^4}{3} - \frac{10x^2}{9} + \frac{1}{9} \right)$$

input `int(-((3*x - x^3)^3 - 9*x + 3*x^3)/(x^2 - 4)^(1/2),x)`output `(x^2 - 4)^(1/2)*((5*x^4)/3 - (10*x^2)/9 - (7*x^6)/9 + x^8/9 + 1/9)`

$$3.274 \quad \int \frac{1}{(1+x^2)^3} dx$$

3.274.1 Optimal result	1379
3.274.2 Mathematica [A] (verified)	1379
3.274.3 Rubi [A] (verified)	1380
3.274.4 Maple [A] (verified)	1381
3.274.5 Fricas [A] (verification not implemented)	1381
3.274.6 Sympy [A] (verification not implemented)	1381
3.274.7 Maxima [A] (verification not implemented)	1382
3.274.8 Giac [A] (verification not implemented)	1382
3.274.9 Mupad [B] (verification not implemented)	1382

3.274.1 Optimal result

Integrand size = 7, antiderivative size = 31

$$\int \frac{1}{(1+x^2)^3} dx = \frac{x}{4(1+x^2)^2} + \frac{3x}{8(1+x^2)} + \frac{3 \arctan(x)}{8}$$

output `1/4*x/(x^2+1)^2+3*x/(8*x^2+8)+3/8*arctan(x)`

3.274.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.81

$$\int \frac{1}{(1+x^2)^3} dx = \frac{1}{8} \left(\frac{x(5+3x^2)}{(1+x^2)^2} + 3 \arctan(x) \right)$$

input `Integrate[(1 + x^2)^(-3),x]`

output `((x*(5 + 3*x^2))/(1 + x^2)^2 + 3*ArcTan[x])/8`

3.274.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.16, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {215, 215, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(x^2 + 1)^3} dx \\ & \quad \downarrow \text{215} \\ & \frac{3}{4} \int \frac{1}{(x^2 + 1)^2} dx + \frac{x}{4(x^2 + 1)^2} \\ & \quad \downarrow \text{215} \\ & \frac{3}{4} \left(\frac{1}{2} \int \frac{1}{x^2 + 1} dx + \frac{x}{2(x^2 + 1)} \right) + \frac{x}{4(x^2 + 1)^2} \\ & \quad \downarrow \text{216} \\ & \frac{3}{4} \left(\frac{\arctan(x)}{2} + \frac{x}{2(x^2 + 1)} \right) + \frac{x}{4(x^2 + 1)^2} \end{aligned}$$

input `Int[(1 + x^2)^(-3), x]`

output `x/(4*(1 + x^2)^2) + (3*(x/(2*(1 + x^2)) + ArcTan[x]/2))/4`

3.274.3.1 Defintions of rubi rules used

rule 215 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

3.274.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

method	result	size
meijerg	$\frac{x(3x^2+5)}{8(x^2+1)^2} + \frac{3 \arctan(x)}{8}$	23
risch	$\frac{\frac{3}{8}x^3 + \frac{5}{8}x}{(x^2+1)^2} + \frac{3 \arctan(x)}{8}$	23
default	$\frac{x}{4(x^2+1)^2} + \frac{3x}{8(x^2+1)} + \frac{3 \arctan(x)}{8}$	26
parallelrisc	$-\frac{3i \ln(x-i)x^4 - 3i \ln(i+x)x^4 + 6i \ln(x-i)x^2 - 6i \ln(i+x)x^2 - 6x^3 + 3i \ln(x-i) - 3i \ln(i+x) - 10x}{16(x^2+1)^2}$	79

input `int(1/(x^2+1)^3,x,method=_RETURNVERBOSE)`output `1/8*x*(3*x^2+5)/(x^2+1)^2+3/8*arctan(x)`**3.274.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.19

$$\int \frac{1}{(1+x^2)^3} dx = \frac{3x^3 + 3(x^4 + 2x^2 + 1) \arctan(x) + 5x}{8(x^4 + 2x^2 + 1)}$$

input `integrate(1/(x^2+1)^3,x, algorithm="fricas")`output `1/8*(3*x^3 + 3*(x^4 + 2*x^2 + 1)*arctan(x) + 5*x)/(x^4 + 2*x^2 + 1)`**3.274.6 Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{1}{(1+x^2)^3} dx = \frac{3x^3 + 5x}{8x^4 + 16x^2 + 8} + \frac{3 \operatorname{atan}(x)}{8}$$

input `integrate(1/(x**2+1)**3,x)`output `(3*x**3 + 5*x)/(8*x**4 + 16*x**2 + 8) + 3*atan(x)/8`

3.274.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

$$\int \frac{1}{(1+x^2)^3} dx = \frac{3x^3 + 5x}{8(x^4 + 2x^2 + 1)} + \frac{3}{8} \arctan(x)$$

input `integrate(1/(x^2+1)^3,x, algorithm="maxima")`output `1/8*(3*x^3 + 5*x)/(x^4 + 2*x^2 + 1) + 3/8*arctan(x)`**3.274.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int \frac{1}{(1+x^2)^3} dx = \frac{3x^3 + 5x}{8(x^2 + 1)^2} + \frac{3}{8} \arctan(x)$$

input `integrate(1/(x^2+1)^3,x, algorithm="giac")`output `1/8*(3*x^3 + 5*x)/(x^2 + 1)^2 + 3/8*arctan(x)`**3.274.9 Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

$$\int \frac{1}{(1+x^2)^3} dx = \frac{3 \operatorname{atan}(x)}{8} + x \left(\frac{1}{4(x^2 + 1)^2} + \frac{3}{8x^2 + 8} \right)$$

input `int(1/(x^2 + 1)^3,x)`output `(3*atan(x))/8 + x*(1/(4*(x^2 + 1)^2) + 3/(8*x^2 + 8))`

3.275 $\int \log(\sqrt{3} + \tan(x)) dx$

3.275.1 Optimal result	1383
3.275.2 Mathematica [A] (verified)	1383
3.275.3 Rubi [F]	1384
3.275.4 Maple [A] (verified)	1385
3.275.5 Fricas [B] (verification not implemented)	1385
3.275.6 Sympy [F]	1386
3.275.7 Maxima [A] (verification not implemented)	1386
3.275.8 Giac [F]	1387
3.275.9 Mupad [F(-1)]	1387

3.275.1 Optimal result

Integrand size = 9, antiderivative size = 108

$$\int \log(\sqrt{3} + \tan(x)) dx = -\frac{1}{2}i \left(\left(\log\left(\frac{i - \tan(x)}{i + \sqrt{3}}\right) - \log\left(\frac{i + \tan(x)}{i - \sqrt{3}}\right) \right) \log(\sqrt{3} + \tan(x)) \right. \\ \left. - \text{PolyLog}\left(2, \frac{\sqrt{3} + \tan(x)}{-i + \sqrt{3}}\right) + \text{PolyLog}\left(2, \frac{\sqrt{3} + \tan(x)}{i + \sqrt{3}}\right) \right)$$

output `-1/2*I*((ln((I-tan(x))/(3^(1/2)+I))-ln((I+tan(x))/(I-3^(1/2))))*ln(3^(1/2)+tan(x))-polylog(2,(3^(1/2)+tan(x))/(-I+3^(1/2)))+polylog(2,(3^(1/2)+tan(x))/(3^(1/2)+I)))`

3.275.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.22

$$\int \log(\sqrt{3} + \tan(x)) dx = -\frac{1}{2}i \log\left(\frac{i - \tan(x)}{i + \sqrt{3}}\right) \log(\sqrt{3} + \tan(x)) \\ + \frac{1}{2}i \log\left(\frac{i + \tan(x)}{i - \sqrt{3}}\right) \log(\sqrt{3} + \tan(x)) \\ + \frac{1}{2}i \text{PolyLog}\left(2, -\frac{\sqrt{3} + \tan(x)}{i - \sqrt{3}}\right) \\ - \frac{1}{2}i \text{PolyLog}\left(2, \frac{\sqrt{3} + \tan(x)}{i + \sqrt{3}}\right)$$

input `Integrate[Log[Sqrt[3] + Tan[x]], x]`

output `(-1/2*I)*Log[(I - Tan[x])/(I + Sqrt[3])]*Log[Sqrt[3] + Tan[x]] + (I/2)*Log[(I + Tan[x])/(I - Sqrt[3])]*Log[Sqrt[3] + Tan[x]] + (I/2)*PolyLog[2, -(Sqrt[3] + Tan[x])/(I - Sqrt[3])] - (I/2)*PolyLog[2, (Sqrt[3] + Tan[x])/(I + Sqrt[3])]`

3.275.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \log(\tan(x) + \sqrt{3}) dx$$

$$\downarrow \text{3028}$$

$$x \log(\tan(x) + \sqrt{3}) - \int \frac{x \sec^2(x)}{\tan(x) + \sqrt{3}} dx$$

$$\downarrow \text{7299}$$

$$x \log(\tan(x) + \sqrt{3}) - \int \frac{x \sec^2(x)}{\tan(x) + \sqrt{3}} dx$$

input `Int[Log[Sqrt[3] + Tan[x]], x]`

output `$Aborted`

3.275.3.1 Defintions of rubi rules used

rule 3028 `Int[Log[u_], x_Symbol] := Simp[x*Log[u], x] - Int[SimplifyIntegrand[x*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[u, x]`

rule 7299 `Int[u_, x_] := CannotIntegrate[u, x]`

3.275.4 Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.89

method	result
derivativedivides	$-\frac{i \ln(\sqrt{3} + \tan(x)) \ln\left(\frac{i - \tan(x)}{\sqrt{3} + i}\right)}{2} + \frac{i \ln(\sqrt{3} + \tan(x)) \ln\left(\frac{i + \tan(x)}{i - \sqrt{3}}\right)}{2} - \frac{i \operatorname{dilog}\left(\frac{i - \tan(x)}{\sqrt{3} + i}\right)}{2} + \frac{i \operatorname{dilog}\left(\frac{i + \tan(x)}{i - \sqrt{3}}\right)}{2}$
default	$-\frac{i \ln(\sqrt{3} + \tan(x)) \ln\left(\frac{i - \tan(x)}{\sqrt{3} + i}\right)}{2} + \frac{i \ln(\sqrt{3} + \tan(x)) \ln\left(\frac{i + \tan(x)}{i - \sqrt{3}}\right)}{2} - \frac{i \operatorname{dilog}\left(\frac{i - \tan(x)}{\sqrt{3} + i}\right)}{2} + \frac{i \operatorname{dilog}\left(\frac{i + \tan(x)}{i - \sqrt{3}}\right)}{2}$
risch	$-i \ln(e^{ix}) \ln(1 - ie^{ix}) + i \ln(e^{ix}) \ln(e^{2ix} + 1) - \frac{i\pi \operatorname{csgn}\left(\frac{i}{e^{2ix} + 1}\right) \operatorname{csgn}\left(i\left(\sqrt{3} e^{2ix} - ie^{2ix} + \sqrt{3} + i\right)\right)}{2}$

input `int(ln(3^(1/2)+tan(x)),x,method=_RETURNVERBOSE)`output `-1/2*I*ln(3^(1/2)+tan(x))*ln((I-tan(x))/(3^(1/2)+I))+1/2*I*ln(3^(1/2)+tan(x))*ln((I+tan(x))/(I-3^(1/2)))-1/2*I*dilog((I-tan(x))/(3^(1/2)+I))+1/2*I*dilog((I+tan(x))/(I-3^(1/2)))`**3.275.5 Fracas [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 262 vs. 2(75) = 150.

Time = 0.27 (sec) , antiderivative size = 262, normalized size of antiderivative = 2.43

$$\begin{aligned} & \int \log(\sqrt{3} + \tan(x)) \, dx \\ &= x \log(\sqrt{3} + \tan(x)) - \frac{1}{2} x \log\left(\frac{(i\sqrt{3} + 1)\tan(x)^2 + 2(\sqrt{3} + i)\tan(x) - i\sqrt{3} + 3}{2(\tan(x)^2 + 1)}\right) \\ & \quad - \frac{1}{2} x \log\left(\frac{(-i\sqrt{3} + 1)\tan(x)^2 + 2(\sqrt{3} - i)\tan(x) + i\sqrt{3} + 3}{2(\tan(x)^2 + 1)}\right) \\ & \quad + \frac{1}{2} x \log\left(-\frac{2(i\tan(x) - 1)}{\tan(x)^2 + 1}\right) + \frac{1}{2} x \log\left(-\frac{2(-i\tan(x) - 1)}{\tan(x)^2 + 1}\right) \\ & \quad + \frac{1}{4} i \operatorname{Li}_2\left(-\frac{(i\sqrt{3} + 1)\tan(x)^2 + 2(\sqrt{3} + i)\tan(x) - i\sqrt{3} + 3}{2(\tan(x)^2 + 1)} + 1\right) \\ & \quad - \frac{1}{4} i \operatorname{Li}_2\left(-\frac{(-i\sqrt{3} + 1)\tan(x)^2 + 2(\sqrt{3} - i)\tan(x) + i\sqrt{3} + 3}{2(\tan(x)^2 + 1)} + 1\right) \\ & \quad + \frac{1}{4} i \operatorname{Li}_2\left(\frac{2(i\tan(x) - 1)}{\tan(x)^2 + 1} + 1\right) - \frac{1}{4} i \operatorname{Li}_2\left(\frac{2(-i\tan(x) - 1)}{\tan(x)^2 + 1} + 1\right) \end{aligned}$$

input `integrate(log(3^(1/2)+tan(x)),x, algorithm="fricas")`

output `x*log(sqrt(3) + tan(x)) - 1/2*x*log(1/2*((I*sqrt(3) + 1)*tan(x)^2 + 2*(sqrt(3) + I)*tan(x) - I*sqrt(3) + 3)/(tan(x)^2 + 1)) - 1/2*x*log(1/2*((-I*sqrt(3) + 1)*tan(x)^2 + 2*(sqrt(3) - I)*tan(x) + I*sqrt(3) + 3)/(tan(x)^2 + 1)) + 1/2*x*log(-2*(I*tan(x) - 1)/(tan(x)^2 + 1)) + 1/2*x*log(-2*(-I*tan(x) - 1)/(tan(x)^2 + 1)) + 1/4*I*dilog(-1/2*((I*sqrt(3) + 1)*tan(x)^2 + 2*(sqrt(3) + I)*tan(x) - I*sqrt(3) + 3)/(tan(x)^2 + 1) + 1) - 1/4*I*dilog(-1/2*((-I*sqrt(3) + 1)*tan(x)^2 + 2*(sqrt(3) - I)*tan(x) + I*sqrt(3) + 3)/(tan(x)^2 + 1) + 1) + 1/4*I*dilog(2*(I*tan(x) - 1)/(tan(x)^2 + 1) + 1) - 1/4*I*dilog(2*(-I*tan(x) - 1)/(tan(x)^2 + 1) + 1)`

3.275.6 Sympy [F]

$$\int \log(\sqrt{3} + \tan(x)) dx = \int \log(\tan(x) + \sqrt{3}) dx$$

input `integrate(ln(3**(1/2)+tan(x)),x)`

output `Integral(log(tan(x) + sqrt(3)), x)`

3.275.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.06

$$\begin{aligned} \int \log(\sqrt{3} + \tan(x)) dx = & \frac{1}{2} \arctan\left(\frac{1}{4}\sqrt{3} + \frac{1}{4}\tan(x), \frac{1}{4}\sqrt{3}\tan(x) + \frac{3}{4}\right) \log(\tan(x)^2 + 1) \\ & - \frac{1}{2}x \log\left(\frac{1}{4}\tan(x)^2 + \frac{1}{2}\sqrt{3}\tan(x) + \frac{3}{4}\right) + x \log(\sqrt{3} + \tan(x)) \\ & + \frac{1}{2}i \operatorname{Li}_2\left(-\frac{(\sqrt{3} + i)\tan(x) - i\sqrt{3} + 1}{2i\sqrt{3} + 2}\right) \\ & - \frac{1}{2}i \operatorname{Li}_2\left(\frac{(\sqrt{3} - i)\tan(x) + i\sqrt{3} + 1}{2i\sqrt{3} - 2}\right) \end{aligned}$$

input `integrate(log(3^(1/2)+tan(x)),x, algorithm="maxima")`

output $\frac{1}{2} \arctan 2\left(\frac{1}{4}\sqrt{3} + \frac{1}{4}\tan(x), \frac{1}{4}\sqrt{3}\tan(x) + \frac{3}{4}\right) \log(\tan(x)^2 + 1) - \frac{1}{2}x \log\left(\frac{1}{4}\tan(x)^2 + \frac{1}{2}\sqrt{3}\tan(x) + \frac{3}{4}\right) + x \log(\sqrt{3} + \tan(x)) + \frac{1}{2}I \operatorname{dilog}\left(-\frac{(\sqrt{3} + I)\tan(x) - I\sqrt{3} + 1}{2I\sqrt{3} + 2}\right) - \frac{1}{2}I \operatorname{dilog}\left(\frac{(\sqrt{3} - I)\tan(x) + I\sqrt{3} + 1}{2I\sqrt{3} - 2}\right)$

3.275.8 Giac [F]

$$\int \log(\sqrt{3} + \tan(x)) dx = \int \log(\sqrt{3} + \tan(x)) dx$$

input `integrate(log(3^(1/2)+tan(x)),x, algorithm="giac")`

output `integrate(log(sqrt(3) + tan(x)), x)`

3.275.9 Mupad [F(-1)]

Timed out.

$$\int \log(\sqrt{3} + \tan(x)) dx = \int \ln(\tan(x) + \sqrt{3}) dx$$

input `int(log(tan(x) + 3^(1/2)),x)`

output `int(log(tan(x) + 3^(1/2)), x)`

3.276 $\int \sqrt{(\cos(20x) + 3 \cos(21x) + \cos(22x))^2 + (\sin(20x) + 3 \sin(21x) + \sin(22x))^2} dx$

3.276.1 Optimal result	1388
3.276.2 Mathematica [A] (verified)	1388
3.276.3 Rubi [B] (verified)	1389
3.276.4 Maple [C] (warning: unable to verify)	1390
3.276.5 Fricas [A] (verification not implemented)	1391
3.276.6 Sympy [F]	1391
3.276.7 Maxima [A] (verification not implemented)	1392
3.276.8 Giac [A] (verification not implemented)	1392
3.276.9 Mupad [B] (verification not implemented)	1392

3.276.1 Optimal result

Integrand size = 39, antiderivative size = 32

$$\int \sqrt{(\cos(20x) + 3 \cos(21x) + \cos(22x))^2 + (\sin(20x) + 3 \sin(21x) + \sin(22x))^2} dx$$

$$= \frac{2(3 + 2 \cos(x)) (3 \arctan(\tan(\frac{x}{2})) + \sin(x))}{\sqrt{(3 + 2 \cos(x))^2}}$$

```
output 2*(3+2*cos(x))*(3*arctan(tan(1/2*x))+sin(x))/((3+2*cos(x))^2)^(1/2)
```

3.276.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91

$$\int \sqrt{(\cos(20x) + 3 \cos(21x) + \cos(22x))^2 + (\sin(20x) + 3 \sin(21x) + \sin(22x))^2} dx$$

$$= \frac{\sqrt{(3 + 2 \cos(x))^2}(3x + 2 \sin(x))}{3 + 2 \cos(x)}$$

```
input Integrate[Sqrt[(Cos[20*x] + 3*Cos[21*x] + Cos[22*x])^2 + (Sin[20*x] + 3*Sin[21*x] + Sin[22*x])^2],x]
```

```
output (Sqrt[(3 + 2*Cos[x])^2]*(3*x + 2*Sin[x]))/(3 + 2*Cos[x])
```

3.276.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 83 vs. $2(32) = 64$.

Time = 0.36 (sec) , antiderivative size = 83, normalized size of antiderivative = 2.59, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.128$, Rules used = {3042, 4902, 2058, 298, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{(\sin(20x) + 3\sin(21x) + \sin(22x))^2 + (\cos(20x) + 3\cos(21x) + \cos(22x))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{(\sin(20x) + 3\sin(21x) + \sin(22x))^2 + (\cos(20x) + 3\cos(21x) + \cos(22x))^2} dx \\
 & \quad \downarrow \text{4902} \\
 & 2 \int \frac{\sqrt{\frac{(\tan^2(\frac{x}{2})+5)^2}{(\tan^2(\frac{x}{2})+1)^2}}}{\tan^2(\frac{x}{2})+1} d \tan\left(\frac{x}{2}\right) \\
 & \quad \downarrow \text{2058} \\
 & \frac{2(\tan^2(\frac{x}{2})+1) \sqrt{\frac{(\tan^2(\frac{x}{2})+5)^2}{(\tan^2(\frac{x}{2})+1)^2}} \int \frac{\tan^2(\frac{x}{2})+5}{(\tan^2(\frac{x}{2})+1)^2} d \tan\left(\frac{x}{2}\right)}{\tan^2(\frac{x}{2})+5} \\
 & \quad \downarrow \text{298} \\
 & \frac{2(\tan^2(\frac{x}{2})+1) \sqrt{\frac{(\tan^2(\frac{x}{2})+5)^2}{(\tan^2(\frac{x}{2})+1)^2}} \left(3 \int \frac{1}{\tan^2(\frac{x}{2})+1} d \tan\left(\frac{x}{2}\right) + \frac{2 \tan(\frac{x}{2})}{\tan^2(\frac{x}{2})+1}\right)}{\tan^2(\frac{x}{2})+5} \\
 & \quad \downarrow \text{216} \\
 & \frac{2(\tan^2(\frac{x}{2})+1) \sqrt{\frac{(\tan^2(\frac{x}{2})+5)^2}{(\tan^2(\frac{x}{2})+1)^2}} \left(3 \arctan(\tan(\frac{x}{2})) + \frac{2 \tan(\frac{x}{2})}{\tan^2(\frac{x}{2})+1}\right)}{\tan^2(\frac{x}{2})+5}
 \end{aligned}$$

input `Int[Sqrt[(Cos[20*x] + 3*Cos[21*x] + Cos[22*x])^2 + (Sin[20*x] + 3*Sin[21*x] + Sin[22*x])^2], x]`

output `(2*(1 + Tan[x/2]^2)*Sqrt[(5 + Tan[x/2]^2)^2/(1 + Tan[x/2]^2)^2]*(3*ArcTan[Tan[x/2]] + (2*Tan[x/2])/(1 + Tan[x/2]^2)))/(5 + Tan[x/2]^2)`

3.276. $\int \sqrt{(\cos(20x) + 3\cos(21x) + \cos(22x))^2 + (\sin(20x) + 3\sin(21x) + \sin(22x))^2} dx$

3.276.3.1 Defintions of rubi rules used

- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

- rule 298 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[(-(b*c - a*d))*x*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] - Simp[(a*d - b*c*(2*p + 3))/(2*a*b*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/2 + p, 0])`

- rule 2058 `Int[(u_.)*((e_.)*((a_.) + (b_.)*(x_)^(n_.))^(q_.)*((c_) + (d_.)*(x_)^(n_.))^(r_.))^(p_), x_Symbol] := Simp[Simp[(e*(a + b*x^n)^q*(c + d*x^n)^r]^p/((a + b*x^n)^(p*q)*(c + d*x^n)^(p*r))] Int[u*(a + b*x^n)^(p*q)*(c + d*x^n)^(p*r), x], x] /; FreeQ[{a, b, c, d, e, n, p, q, r}, x]`

- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

- rule 4902 `Int[u_, x_Symbol] := With[{w = Block[{$ShowSteps = False, $StepCounter = Null}, Int[SubstFor[1/(1 + FreeFactors[Tan[FunctionOfTrig[u, x]/2], x]^2*x^2), Tan[FunctionOfTrig[u, x]/2]/FreeFactors[Tan[FunctionOfTrig[u, x]/2], x], u, x], x]}], Module[{v = FunctionOfTrig[u, x], d}, Simp[d = FreeFactors[Tan[v/2], x]; 2*(d/Coefficient[v, x, 1]) Subst[Int[SubstFor[1/(1 + d^2*x^2), Tan[v/2]/d, u, x], x], x, Tan[v/2]/d, x]] /; CalculusFreeQ[w, x]] /; InverseFunctionFreeQ[u, x] && !FalseQ[FunctionOfTrig[u, x]]`

3.276.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 2.33 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.53

method	result	size
default	$\text{csgn}(3 + 2 \cos(x))(3x + 2 \sin(x))$	17
risch	$\frac{3\sqrt{(e^{2ix}+3e^{ix}+1)^2e^{-2ix}e^{ix}x}}{e^{2ix}+3e^{ix}+1} - \frac{i\sqrt{(e^{2ix}+3e^{ix}+1)^2e^{-2ix}e^{2ix}}}{e^{2ix}+3e^{ix}+1} + \frac{i\sqrt{(e^{2ix}+3e^{ix}+1)^2e^{-2ix}}}{e^{2ix}+3e^{ix}+1}$	141

3.276. $\int \sqrt{(\cos(20x) + 3 \cos(21x) + \cos(22x))^2 + (\sin(20x) + 3 \sin(21x) + \sin(22x))^2} dx$

input `int(((sin(20*x)+3*sin(21*x)+sin(22*x))^2+(cos(20*x)+3*cos(21*x)+cos(22*x))^2)^(1/2),x,method=_RETURNVERBOSE)`

output `csgn(3+2*cos(x))*(3*x+2*sin(x))`

3.276.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.25

$$\int \sqrt{(\cos(20x) + 3 \cos(21x) + \cos(22x))^2 + (\sin(20x) + 3 \sin(21x) + \sin(22x))^2} dx$$

$$= 3x + 2 \sin(x)$$

input `integrate(((sin(20*x)+3*sin(21*x)+sin(22*x))^2+(cos(20*x)+3*cos(21*x)+cos(22*x))^2)^(1/2),x,algorithm="fricas")`

output `3*x + 2*sin(x)`

3.276.6 Sympy [F]

$$\int \sqrt{(\cos(20x) + 3 \cos(21x) + \cos(22x))^2 + (\sin(20x) + 3 \sin(21x) + \sin(22x))^2} dx$$

$$= \int \sqrt{(\sin(20x) + 3 \sin(21x) + \sin(22x))^2 + (\cos(20x) + 3 \cos(21x) + \cos(22x))^2} dx$$

input `integrate(((sin(20*x)+3*sin(21*x)+sin(22*x))**2+(cos(20*x)+3*cos(21*x)+cos(22*x))**2)**(1/2),x)`

output `Integral(sqrt((sin(20*x) + 3*sin(21*x) + sin(22*x))**2 + (cos(20*x) + 3*cos(21*x) + cos(22*x))**2), x)`

3.276.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.25

$$\int \sqrt{(\cos(20x) + 3 \cos(21x) + \cos(22x))^2 + (\sin(20x) + 3 \sin(21x) + \sin(22x))^2} dx$$

$$= 3x + 2 \sin(x)$$

input `integrate(((sin(20*x)+3*sin(21*x)+sin(22*x))^2+(cos(20*x)+3*cos(21*x)+cos(22*x))^2)^(1/2),x, algorithm="maxima")`

output `3*x + 2*sin(x)`

3.276.8 Giac [A] (verification not implemented)

Time = 0.87 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \sqrt{(\cos(20x) + 3 \cos(21x) + \cos(22x))^2 + (\sin(20x) + 3 \sin(21x) + \sin(22x))^2} dx$$

$$= -6\pi \left[\frac{x}{2\pi} + \frac{1}{2} \right] + 3x + \frac{4 \tan\left(\frac{1}{2}x\right)}{\tan\left(\frac{1}{2}x\right)^2 + 1}$$

input `integrate(((sin(20*x)+3*sin(21*x)+sin(22*x))^2+(cos(20*x)+3*cos(21*x)+cos(22*x))^2)^(1/2),x, algorithm="giac")`

output `-6*pi*floor(1/2*x/pi + 1/2) + 3*x + 4*tan(1/2*x)/(tan(1/2*x)^2 + 1)`

3.276.9 Mupad [B] (verification not implemented)

Time = 15.60 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.25

$$\int \sqrt{(\cos(20x) + 3 \cos(21x) + \cos(22x))^2 + (\sin(20x) + 3 \sin(21x) + \sin(22x))^2} dx$$

$$= 3x + 2 \sin(x)$$

input `int(((cos(20*x) + 3*cos(21*x) + cos(22*x))^2 + (sin(20*x) + 3*sin(21*x) + sin(22*x))^2)^(1/2),x)`

output `3*x + 2*sin(x)`

$$3.277 \quad \int \frac{e^{-2x} \sin(3x)}{x} dx$$

3.277.1 Optimal result	1393
3.277.2 Mathematica [A] (verified)	1393
3.277.3 Rubi [F]	1394
3.277.4 Maple [A] (verified)	1394
3.277.5 Fricas [A] (verification not implemented)	1395
3.277.6 Sympy [F]	1395
3.277.7 Maxima [C] (verification not implemented)	1395
3.277.8 Giac [A] (verification not implemented)	1396
3.277.9 Mupad [B] (verification not implemented)	1396

3.277.1 Optimal result

Integrand size = 13, antiderivative size = 21

$$\int \frac{e^{-2x} \sin(3x)}{x} dx = \frac{1}{2}i(\text{ExpIntegralEi}((-2 - 3i)x) - \text{ExpIntegralEi}((-2 + 3i)x))$$

output `1/2*I*(Ei((-2-3*I)*x)-Ei((-2+3*I)*x))`

3.277.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{e^{-2x} \sin(3x)}{x} dx = -\frac{1}{2}i(-\text{ExpIntegralEi}((-2 - 3i)x) + \text{ExpIntegralEi}((-2 + 3i)x))$$

input `Integrate[Sin[3*x]/(E^(2*x)*x),x]`

output `(-1/2*I)*(-ExpIntegralEi[(-2 - 3*I)*x] + ExpIntegralEi[(-2 + 3*I)*x])`

3.277.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{-2x} \sin(3x)}{x} dx$$

↓ 7299

$$\int \frac{e^{-2x} \sin(3x)}{x} dx$$

input `Int [Sin [3*x] / (E^(2*x)*x) , x]`

output `$Aborted`

3.277.3.1 Defintions of rubi rules used

rule 7299 `Int [u_ , x_] :=> CannotIntegrate [u , x]`

3.277.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

method	result	size
risch	$\frac{i \operatorname{Ei}_1((2-3i)x)}{2} - \frac{i \operatorname{Ei}_1((2+3i)x)}{2}$	22

input `int (exp (-2*x) *sin (3*x) /x ,x ,method =_RETURNVERBOSE)`

output `1/2*I*Ei (1 , (2-3*I) *x) -1/2*I*Ei (1 , (2+3*I) *x)`

3.277.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \frac{e^{-2x} \sin(3x)}{x} dx = -\frac{1}{2}i \operatorname{Ei}((3i - 2)x) + \frac{1}{2}i \operatorname{Ei}(-(3i + 2)x)$$

input `integrate(exp(-2*x)*sin(3*x)/x,x, algorithm="fricas")`

output `-1/2*I*Ei((3*I - 2)*x) + 1/2*I*Ei(-(3*I + 2)*x)`

3.277.6 Sympy [F]

$$\int \frac{e^{-2x} \sin(3x)}{x} dx = \int \frac{e^{-2x} \sin(3x)}{x} dx$$

input `integrate(exp(-2*x)*sin(3*x)/x,x)`

output `Integral(exp(-2*x)*sin(3*x)/x, x)`

3.277.7 Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.24 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.29

$$\int \frac{e^{-2x} \sin(3x)}{x} dx = -\frac{1}{4}i \operatorname{Ei}((3i - 2)x) + \frac{1}{4}i \operatorname{Ei}(-(3i + 2)x) \\ + \frac{1}{4}i \overline{\operatorname{Ei}((3i - 2)x)} - \frac{1}{4}i \overline{\operatorname{Ei}(-(3i + 2)x)}$$

input `integrate(exp(-2*x)*sin(3*x)/x,x, algorithm="maxima")`

output `-1/4*I*Ei((3*I - 2)*x) + 1/4*I*Ei(-(3*I + 2)*x) + 1/4*I*conjugate(Ei((3*I - 2)*x)) - 1/4*I*conjugate(Ei(-(3*I + 2)*x))`

3.277.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \frac{e^{-2x} \sin(3x)}{x} dx = -\frac{1}{2}i \operatorname{Ei}((3i - 2)x) + \frac{1}{2}i \operatorname{Ei}(-(3i + 2)x)$$

input `integrate(exp(-2*x)*sin(3*x)/x,x, algorithm="giac")`output `-1/2*I*Ei((3*I - 2)*x) + 1/2*I*Ei(-(3*I + 2)*x)`**3.277.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{e^{-2x} \sin(3x)}{x} dx = \frac{\operatorname{ei}(x(-2 - 3i)) \operatorname{li}}{2} - \frac{\operatorname{ei}(x(-2 + 3i)) \operatorname{li}}{2}$$

input `int((sin(3*x)*exp(-2*x))/x,x)`output `(ei(x*(- 2 - 3i))*1i)/2 - (ei(x*(- 2 + 3i))*1i)/2`

3.278 $\int (1 - x)^{2/3} \sqrt[3]{x} dx$

3.278.1 Optimal result	1397
3.278.2 Mathematica [A] (verified)	1397
3.278.3 Rubi [A] (verified)	1398
3.278.4 Maple [C] (verified)	1399
3.278.5 Fracas [A] (verification not implemented)	1400
3.278.6 Sympy [C] (verification not implemented)	1400
3.278.7 Maxima [A] (verification not implemented)	1401
3.278.8 Giac [F]	1401
3.278.9 Mupad [F(-1)]	1401

3.278.1 Optimal result

Integrand size = 15, antiderivative size = 102

$$\int (1 - x)^{2/3} \sqrt[3]{x} dx = \frac{1}{6}(1 - x)^{2/3} \sqrt[3]{x} - \frac{1}{2}(1 - x)^{5/3} \sqrt[3]{x} + \frac{\arctan\left(\frac{1}{\sqrt{3}} - \frac{2\sqrt[3]{1-x}}{\sqrt{3}\sqrt[3]{x}}\right)}{3\sqrt{3}} + \frac{1}{6} \log\left(1 + \frac{\sqrt[3]{1-x}}{\sqrt[3]{x}}\right) + \frac{\log(x)}{18}$$

```
output 1/6*x^(1/3)*(1-x)^(2/3)-1/2*(1-x)^(5/3)*x^(1/3)-1/9*arctan(-1/3*3^(1/2)+2/3*(1-x)^(1/3)*3^(1/2)/x^(1/3))*3^(1/2)+1/6*ln(1+(1-x)^(1/3)/x^(1/3))+1/18*ln(x)
```

3.278.2 Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.12

$$\int (1 - x)^{2/3} \sqrt[3]{x} dx = \frac{1}{18} \left(3(1-x)^{2/3} \sqrt[3]{x}(-2+3x) + 2\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{x}}{2\sqrt[3]{1-x} - \sqrt[3]{x}}\right) + 2 \log(\sqrt[3]{1-x} + \sqrt[3]{x}) - \log\left(\frac{\sqrt{3}\sqrt[3]{x}}{2\sqrt[3]{1-x} - \sqrt[3]{x}}\right) \right)$$

```
input Integrate[(1 - x)^(2/3)*x^(1/3), x]
```

output $(3*(1-x)^{2/3}*x^{1/3}*(-2+3*x) + 2*\text{Sqrt}[3]*\text{ArcTan}[\text{Sqrt}[3]*x^{1/3}]/(2*(1-x)^{1/3} - x^{1/3})) + 2*\text{Log}[(1-x)^{1/3} + x^{1/3}] - \text{Log}[(1-x)^{2/3} + x^{2/3} - (-((-1+x)*x))^{1/3}]/18$

3.278.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.04, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {60, 60, 72}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (1-x)^{2/3} \sqrt[3]{x} dx$$

$$\downarrow 60$$

$$\frac{1}{6} \int \frac{(1-x)^{2/3}}{x^{2/3}} dx - \frac{1}{2} (1-x)^{5/3} \sqrt[3]{x}$$

$$\downarrow 60$$

$$\frac{1}{6} \left(\frac{2}{3} \int \frac{1}{\sqrt[3]{1-xx^{2/3}}} dx + (1-x)^{2/3} \sqrt[3]{x} \right) - \frac{1}{2} (1-x)^{5/3} \sqrt[3]{x}$$

$$\downarrow 72$$

$$\frac{1}{6} \left(\frac{2}{3} \left(\sqrt{3} \arctan \left(\frac{1}{\sqrt{3}} - \frac{2\sqrt[3]{1-x}}{\sqrt{3}\sqrt[3]{x}} \right) + \frac{3}{2} \log \left(\frac{\sqrt[3]{1-x}}{\sqrt[3]{x}} + 1 \right) + \frac{\log(x)}{2} \right) + (1-x)^{2/3} \sqrt[3]{x} \right) - \frac{1}{2} (1-x)^{5/3} \sqrt[3]{x}$$

input $\text{Int}[(1-x)^{2/3}*x^{1/3},x]$

output $-1/2*((1-x)^{5/3}*x^{1/3}) + ((1-x)^{2/3}*x^{1/3} + (2*(\text{Sqrt}[3]*\text{ArcTan}[1/\text{Sqrt}[3] - (2*(1-x)^{1/3})/(\text{Sqrt}[3]*x^{1/3})]) + (3*\text{Log}[1 + (1-x)^{1/3}]/x^{1/3}))/2 + \text{Log}[x]/2))/3/6$

3.278.3.1 Defintions of rubi rules used

```
rule 60 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(
b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]
```

```
rule 72 Int[1/(((a_.) + (b_.)*(x_))^(1/3)*((c_.) + (d_.)*(x_))^(2/3)), x_Symbol] :=
With[{q = Rt[-d/b, 3]}, Simp[Sqrt[3]*(q/d)*ArcTan[1/Sqrt[3] - 2*q*((a + b*
x)^(1/3)/(Sqrt[3]*(c + d*x)^(1/3))], x] + (Simp[3*(q/(2*d))*Log[q*((a + b*
x)^(1/3)/(c + d*x)^(1/3)) + 1], x] + Simp[(q/(2*d))*Log[c + d*x], x])] /; F
reeQ[{a, b, c, d}, x] && NegQ[d/b]
```

3.278.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3.

Time = 0.25 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.13

method	result	size
meijerg	$\frac{3x^{\frac{4}{3}} \operatorname{hypergeom}\left(\left[-\frac{2}{3}, \frac{4}{3}\right], \left[\frac{7}{3}\right], x\right)}{4}$	13
risch	$-\frac{(-2+3x)(-1+x)x^{\frac{1}{3}}(x^2(1-x))^{\frac{1}{3}}}{6(-x^2(-1+x))^{\frac{1}{3}}(1-x)^{\frac{1}{3}}} + \frac{\operatorname{hypergeom}\left(\left[\frac{1}{3}, \frac{1}{3}\right], \left[\frac{4}{3}\right], x\right)(x^2(1-x))^{\frac{1}{3}}}{3x^{\frac{1}{3}}(1-x)^{\frac{1}{3}}}$	73

```
input int(x^(1/3)*(1-x)^(2/3), x, method=_RETURNVERBOSE)
```

```
output 3/4*x^(4/3)*hypergeom([-2/3, 4/3], [7/3], x)
```


3.278.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.12

$$\int (1-x)^{2/3} \sqrt[3]{x} dx = \frac{1}{6} (3x-2)x^{1/3}(-x+1)^{2/3} - \frac{1}{9} \sqrt{3} \arctan\left(\frac{\sqrt{3}(x-1) + 2\sqrt{3}x^{1/3}(-x+1)^{2/3}}{3(x-1)}\right) - \frac{1}{18} \log\left(\frac{x - x^{2/3}(-x+1)^{1/3} + x^{1/3}(-x+1)^{2/3} - 1}{x-1}\right) + \frac{1}{9} \log\left(-\frac{x - x^{1/3}(-x+1)^{2/3} - 1}{x-1}\right)$$

input `integrate(x^(1/3)*(1-x)^(2/3),x, algorithm="fricas")`output `1/6*(3*x - 2)*x^(1/3)*(-x + 1)^(2/3) - 1/9*sqrt(3)*arctan(1/3*(sqrt(3)*(x - 1) + 2*sqrt(3)*x^(1/3)*(-x + 1)^(2/3))/(x - 1)) - 1/18*log((x - x^(2/3)*(-x + 1)^(1/3) + x^(1/3)*(-x + 1)^(2/3) - 1)/(x - 1)) + 1/9*log(-(x - x^(1/3)*(-x + 1)^(2/3) - 1)/(x - 1))`**3.278.6 Sympy [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 1.77 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.30

$$\int (1-x)^{2/3} \sqrt[3]{x} dx = \frac{x^{4/3} \Gamma\left(\frac{4}{3}\right) {}_2F_1\left(\begin{matrix} -\frac{2}{3}, \frac{4}{3} \\ \frac{7}{3} \end{matrix} \middle| xe^{2i\pi}\right)}{\Gamma\left(\frac{7}{3}\right)}$$

input `integrate(x**(1/3)*(1-x)**(2/3),x)`output `x**(4/3)*gamma(4/3)*hyper((-2/3, 4/3), (7/3,), x*exp_polar(2*I*pi))/gamma(7/3)`

3.278.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.15

$$\int (1-x)^{2/3} \sqrt[3]{x} dx =$$

$$-\frac{1}{9} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} \left(\frac{2(-x+1)^{1/3}}{x^{1/3}} - 1 \right) \right) + \frac{\frac{(-x+1)^{2/3}}{x^{2/3}} - \frac{2(-x+1)^{5/3}}{x^{5/3}}}{6 \left(\frac{(x-1)^2}{x^2} - \frac{2(x-1)}{x} + 1 \right)}$$

$$+ \frac{1}{9} \log \left(\frac{(-x+1)^{1/3}}{x^{1/3}} + 1 \right) - \frac{1}{18} \log \left(-\frac{(-x+1)^{1/3}}{x^{1/3}} + \frac{(-x+1)^{2/3}}{x^{2/3}} + 1 \right)$$

input `integrate(x^(1/3)*(1-x)^(2/3),x, algorithm="maxima")`output `-1/9*sqrt(3)*arctan(1/3*sqrt(3)*(2*(-x + 1)^(1/3)/x^(1/3) - 1)) + 1/6*((-x + 1)^(2/3)/x^(2/3) - 2*(-x + 1)^(5/3)/x^(5/3))/((x - 1)^2/x^2 - 2*(x - 1)/x + 1) + 1/9*log((-x + 1)^(1/3)/x^(1/3) + 1) - 1/18*log(-(-x + 1)^(1/3)/x^(1/3) + (-x + 1)^(2/3)/x^(2/3) + 1)`**3.278.8 Giac [F]**

$$\int (1-x)^{2/3} \sqrt[3]{x} dx = \int x^{1/3} (-x+1)^{2/3} dx$$

input `integrate(x^(1/3)*(1-x)^(2/3),x, algorithm="giac")`output `integrate(x^(1/3)*(-x + 1)^(2/3), x)`**3.278.9 Mupad [F(-1)]**

Timed out.

$$\int (1-x)^{2/3} \sqrt[3]{x} dx = \int x^{1/3} (1-x)^{2/3} dx$$

input `int(x^(1/3)*(1 - x)^(2/3),x)`output `int(x^(1/3)*(1 - x)^(2/3), x)`

3.279 $\int e dx$

3.279.1 Optimal result	1402
3.279.2 Mathematica [A] (verified)	1402
3.279.3 Rubi [A] (verified)	1403
3.279.4 Maple [A] (verified)	1403
3.279.5 Fricas [A] (verification not implemented)	1404
3.279.6 Sympy [A] (verification not implemented)	1404
3.279.7 Maxima [A] (verification not implemented)	1404
3.279.8 Giac [A] (verification not implemented)	1405
3.279.9 Mupad [B] (verification not implemented)	1405

3.279.1 Optimal result

Integrand size = 1, antiderivative size = 3

$$\int e dx = ex$$

output `x*exp(1)`

3.279.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int e dx = ex$$

input `Integrate[E,x]`

output `E*x`

3.279.3 Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e dx$$

$$\downarrow 24$$

$$ex$$

input `Int [E, x]`

output `E*x`

3.279.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

3.279.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 3.00

method	result	size
parallelrisc	$x^{\frac{1}{\ln(x)}} x$	9

input `int(x^(1/ln(x)), x, method=_RETURNVERBOSE)`

output `x^(1/ln(x))*x`

3.279.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

$$\int e dx = xe$$

input `integrate(x^(1/log(x)),x, algorithm="fricas")`

output `x*e`

3.279.6 Sympy [A] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int e dx = ex$$

input `integrate(x**(1/ln(x)),x)`

output `E*x`

3.279.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

$$\int e dx = xe$$

input `integrate(x^(1/log(x)),x, algorithm="maxima")`

output `x*e`

3.279.8 Giac [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

$$\int e dx = xe$$

input `integrate(x^(1/log(x)),x, algorithm="giac")`

output `x*e`

3.279.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

$$\int e dx = xe$$

input `int(x^(1/log(x)),x)`

output `x*exp(1)`

3.280 $\int \operatorname{sech}(x) dx$

3.280.1 Optimal result	1406
3.280.2 Mathematica [A] (verified)	1406
3.280.3 Rubi [A] (verified)	1407
3.280.4 Maple [A] (verified)	1408
3.280.5 Fricas [B] (verification not implemented)	1408
3.280.6 Sympy [B] (verification not implemented)	1408
3.280.7 Maxima [A] (verification not implemented)	1409
3.280.8 Giac [A] (verification not implemented)	1409
3.280.9 Mupad [B] (verification not implemented)	1409

3.280.1 Optimal result

Integrand size = 2, antiderivative size = 3

$$\int \operatorname{sech}(x) dx = \arctan(\sinh(x))$$

output `arctan(sinh(x))`

3.280.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \operatorname{sech}(x) dx = \arctan(\sinh(x))$$

input `Integrate[Sech[x],x]`

output `ArcTan[Sinh[x]]`

3.280.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \operatorname{sech}(x) dx \\ \downarrow 3042 \\ \int \csc\left(\frac{\pi}{2} + ix\right) dx \\ \downarrow 4257 \\ \arctan(\sinh(x)) \end{array}$$

input `Int [Sech [x] , x]`

output `ArcTan [Sinh [x]]`

3.280.3.1 Defintions of rubi rules used

rule 3042 `Int [u_ , x_Symbol] := Int [DeactivateTrig [u , x] , x] /; FunctionOfTrigOfLinear Q [u , x]`

rule 4257 `Int [csc [(c_.) + (d_.)*(x_)] , x_Symbol] := Simp [-ArcTanh [Cos [c + d*x]] / d , x] /; FreeQ [{c , d} , x]`

3.280.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
lookup	$\arctan(\sinh(x))$	4
default	$\arctan(\sinh(x))$	4
risch	$i \ln(e^x + i) - i \ln(e^x - i)$	20
parallelrisch	$-i(\ln(\tanh(\frac{x}{2}) - i) - \ln(\tanh(\frac{x}{2}) + i))$	23

input `int(sech(x), x, method=_RETURNVERBOSE)`

output `arctan(sinh(x))`

3.280.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8 vs. $2(3) = 6$.

Time = 0.24 (sec) , antiderivative size = 8, normalized size of antiderivative = 2.67

$$\int \operatorname{sech}(x) dx = 2 \arctan(\cosh(x) + \sinh(x))$$

input `integrate(sech(x), x, algorithm="fricas")`

output `2*arctan(cosh(x) + sinh(x))`

3.280.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7 vs. $2(3) = 6$.

Time = 0.17 (sec) , antiderivative size = 7, normalized size of antiderivative = 2.33

$$\int \operatorname{sech}(x) dx = 2 \operatorname{atan}\left(\tanh\left(\frac{x}{2}\right)\right)$$

input `integrate(sech(x), x)`

output `2*atan(tanh(x/2))`

3.280.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \operatorname{sech}(x) dx = \arctan(\sinh(x))$$

input `integrate(sech(x),x, algorithm="maxima")`

output `arctan(sinh(x))`

3.280.8 Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.67

$$\int \operatorname{sech}(x) dx = 2 \arctan(e^x)$$

input `integrate(sech(x),x, algorithm="giac")`

output `2*arctan(e^x)`

3.280.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.67

$$\int \operatorname{sech}(x) dx = 2 \operatorname{atan}(e^x)$$

input `int(1/cosh(x),x)`

output `2*atan(exp(x))`

$$3.281 \quad \int \frac{e^x}{(1+e^x)\log(1+e^x)} dx$$

3.281.1 Optimal result	1410
3.281.2 Mathematica [A] (verified)	1410
3.281.3 Rubi [A] (verified)	1411
3.281.4 Maple [A] (verified)	1412
3.281.5 Fricas [A] (verification not implemented)	1412
3.281.6 Sympy [A] (verification not implemented)	1413
3.281.7 Maxima [A] (verification not implemented)	1413
3.281.8 Giac [A] (verification not implemented)	1413
3.281.9 Mupad [B] (verification not implemented)	1414

3.281.1 Optimal result

Integrand size = 19, antiderivative size = 7

$$\int \frac{e^x}{(1+e^x)\log(1+e^x)} dx = \log(\log(1+e^x))$$

output `ln(ln(exp(x)+1))`

3.281.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{(1+e^x)\log(1+e^x)} dx = \log(\log(1+e^x))$$

input `Integrate[E^x/((1 + E^x)*Log[1 + E^x]), x]`

output `Log[Log[1 + E^x]]`

3.281.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {2720, 2837, 2739, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{e^x}{(e^x + 1) \log(e^x + 1)} dx \\
 \downarrow \text{2720} \\
 \int \frac{1}{(e^x + 1) \log(e^x + 1)} de^x \\
 \downarrow \text{2837} \\
 \int \frac{e^{-x}}{\log(e^x + 1)} d(e^x + 1) \\
 \downarrow \text{2739} \\
 \int e^{-x} d \log(e^x + 1) \\
 \downarrow \text{14} \\
 \log(\log(e^x + 1))
 \end{array}$$

input `Int[E^x/((1 + E^x)*Log[1 + E^x]),x]`

output `Log[Log[1 + E^x]]`

3.281.3.1 Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_)^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 2739 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/(x_), x_Symbol] := Simp[1/(b*n) Subst[Int[x^p, x], x, a + b*Log[c*x^n]], x] /; FreeQ[{a, b, c, n, p}, x]`

rule 2837 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.)]*(b_.))^(p_.)*((f_) + (g_.)*(x_)^(q_.), x_Symbol] := Simp[1/e Subst[Int[(f*(x/d))^q*(a + b*Log[c*x^n])^p, x], x, d + e*x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p, q}, x] && EqQ[e*f - d*g, 0]`

3.281.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

method	result	size
derivativedivides	$\ln(\ln(e^x + 1))$	7
default	$\ln(\ln(e^x + 1))$	7
norman	$\ln(\ln(e^x + 1))$	7
risch	$\ln(\ln(e^x + 1))$	7
parallelrisc	$\ln(\ln(e^x + 1))$	7

input `int(exp(x)/(exp(x)+1)/ln(exp(x)+1),x,method=_RETURNVERBOSE)`

output `ln(ln(exp(x)+1))`

3.281.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int \frac{e^x}{(1+e^x)\log(1+e^x)} dx = \log(\log(e^x + 1))$$

input `integrate(exp(x)/(exp(x)+1)/log(exp(x)+1),x, algorithm="fricas")`

output `log(log(e^x + 1))`

3.281.6 Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{e^x}{(1+e^x)\log(1+e^x)} dx = \log(\log(e^x + 1))$$

input `integrate(exp(x)/(exp(x)+1)/ln(exp(x)+1),x)`output `log(log(exp(x) + 1))`**3.281.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int \frac{e^x}{(1+e^x)\log(1+e^x)} dx = \log(\log(e^x + 1))$$

input `integrate(exp(x)/(exp(x)+1)/log(exp(x)+1),x, algorithm="maxima")`output `log(log(e^x + 1))`**3.281.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int \frac{e^x}{(1+e^x)\log(1+e^x)} dx = \log(\log(e^x + 1))$$

input `integrate(exp(x)/(exp(x)+1)/log(exp(x)+1),x, algorithm="giac")`output `log(log(e^x + 1))`

3.281.9 Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int \frac{e^x}{(1+e^x)\log(1+e^x)} dx = \ln(\ln(e^x + 1))$$

input `int(exp(x)/(log(exp(x) + 1)*(exp(x) + 1)),x)`

output `log(log(exp(x) + 1))`

3.282 $\int (1 - x + x^2 - x^3 + x^4) (1 + x + x^2 + x^3 + x^4) dx$

3.282.1 Optimal result	1415
3.282.2 Mathematica [A] (verified)	1415
3.282.3 Rubi [A] (verified)	1416
3.282.4 Maple [A] (verified)	1417
3.282.5 Fricas [A] (verification not implemented)	1417
3.282.6 Sympy [A] (verification not implemented)	1417
3.282.7 Maxima [A] (verification not implemented)	1418
3.282.8 Giac [A] (verification not implemented)	1418
3.282.9 Mupad [B] (verification not implemented)	1418

3.282.1 Optimal result

Integrand size = 29, antiderivative size = 30

$$\int (1 - x + x^2 - x^3 + x^4) (1 + x + x^2 + x^3 + x^4) dx = x + \frac{x^3}{3} + \frac{x^5}{5} + \frac{x^7}{7} + \frac{x^9}{9}$$

output `x+1/3*x^3+1/5*x^5+1/7*x^7+1/9*x^9`

3.282.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int (1 - x + x^2 - x^3 + x^4) (1 + x + x^2 + x^3 + x^4) dx = x + \frac{x^3}{3} + \frac{x^5}{5} + \frac{x^7}{7} + \frac{x^9}{9}$$

input `Integrate[(1 - x + x^2 - x^3 + x^4)*(1 + x + x^2 + x^3 + x^4),x]`

output `x + x^3/3 + x^5/5 + x^7/7 + x^9/9`

3.282.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.069$, Rules used = {7239, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x^4 - x^3 + x^2 - x + 1)(x^4 + x^3 + x^2 + x + 1) dx$$

$$\downarrow \text{7239}$$

$$\int (x^8 + x^6 + x^4 + x^2 + 1) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^9}{9} + \frac{x^7}{7} + \frac{x^5}{5} + \frac{x^3}{3} + x$$

input `Int[(1 - x + x^2 - x^3 + x^4)*(1 + x + x^2 + x^3 + x^4),x]`

output `x + x^3/3 + x^5/5 + x^7/7 + x^9/9`

3.282.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7239 `Int[u_, x_Symbol] := With[{v = SimplifyIntegrand[u, x]}, Int[v, x] /; SimplifierIntegrandQ[v, u, x]]`

3.282.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.77

method	result	size
gosper	$x + \frac{1}{3}x^3 + \frac{1}{5}x^5 + \frac{1}{7}x^7 + \frac{1}{9}x^9$	23
default	$x + \frac{1}{3}x^3 + \frac{1}{5}x^5 + \frac{1}{7}x^7 + \frac{1}{9}x^9$	23
norman	$x + \frac{1}{3}x^3 + \frac{1}{5}x^5 + \frac{1}{7}x^7 + \frac{1}{9}x^9$	23
risch	$x + \frac{1}{3}x^3 + \frac{1}{5}x^5 + \frac{1}{7}x^7 + \frac{1}{9}x^9$	23
parallelrisc	$x + \frac{1}{3}x^3 + \frac{1}{5}x^5 + \frac{1}{7}x^7 + \frac{1}{9}x^9$	23

input `int((x^4+x^3+x^2+x+1)*(x^4-x^3+x^2-x+1),x,method=_RETURNVERBOSE)`output `x+1/3*x^3+1/5*x^5+1/7*x^7+1/9*x^9`**3.282.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int (1 - x + x^2 - x^3 + x^4) (1 + x + x^2 + x^3 + x^4) dx = \frac{1}{9}x^9 + \frac{1}{7}x^7 + \frac{1}{5}x^5 + \frac{1}{3}x^3 + x$$

input `integrate((x^4+x^3+x^2+x+1)*(x^4-x^3+x^2-x+1),x, algorithm="fracas")`output `1/9*x^9 + 1/7*x^7 + 1/5*x^5 + 1/3*x^3 + x`**3.282.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.67

$$\int (1 - x + x^2 - x^3 + x^4) (1 + x + x^2 + x^3 + x^4) dx = \frac{x^9}{9} + \frac{x^7}{7} + \frac{x^5}{5} + \frac{x^3}{3} + x$$

input `integrate((x**4+x**3+x**2+x+1)*(x**4-x**3+x**2-x+1),x)`output `x**9/9 + x**7/7 + x**5/5 + x**3/3 + x`

3.282.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int (1 - x + x^2 - x^3 + x^4) (1 + x + x^2 + x^3 + x^4) dx = \frac{1}{9} x^9 + \frac{1}{7} x^7 + \frac{1}{5} x^5 + \frac{1}{3} x^3 + x$$

input `integrate((x^4+x^3+x^2+x+1)*(x^4-x^3+x^2-x+1),x, algorithm="maxima")`output `1/9*x^9 + 1/7*x^7 + 1/5*x^5 + 1/3*x^3 + x`**3.282.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int (1 - x + x^2 - x^3 + x^4) (1 + x + x^2 + x^3 + x^4) dx = \frac{1}{9} x^9 + \frac{1}{7} x^7 + \frac{1}{5} x^5 + \frac{1}{3} x^3 + x$$

input `integrate((x^4+x^3+x^2+x+1)*(x^4-x^3+x^2-x+1),x, algorithm="giac")`output `1/9*x^9 + 1/7*x^7 + 1/5*x^5 + 1/3*x^3 + x`**3.282.9 Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int (1 - x + x^2 - x^3 + x^4) (1 + x + x^2 + x^3 + x^4) dx = \frac{x^9}{9} + \frac{x^7}{7} + \frac{x^5}{5} + \frac{x^3}{3} + x$$

input `int((x^2 - x - x^3 + x^4 + 1)*(x + x^2 + x^3 + x^4 + 1),x)`output `x + x^3/3 + x^5/5 + x^7/7 + x^9/9`

3.283 $\int \frac{1}{120}(-4+x)(-3+x)(-2+x)(-1+x)x dx$

3.283.1 Optimal result	1419
3.283.2 Mathematica [A] (verified)	1419
3.283.3 Rubi [A] (verified)	1420
3.283.4 Maple [A] (verified)	1421
3.283.5 Fricas [A] (verification not implemented)	1421
3.283.6 Sympy [A] (verification not implemented)	1421
3.283.7 Maxima [A] (verification not implemented)	1422
3.283.8 Giac [A] (verification not implemented)	1422
3.283.9 Mupad [B] (verification not implemented)	1422

3.283.1 Optimal result

Integrand size = 17, antiderivative size = 36

$$\int \frac{1}{120}(-4+x)(-3+x)(-2+x)(-1+x)x dx = \frac{x^2}{10} - \frac{5x^3}{36} + \frac{7x^4}{96} - \frac{x^5}{60} + \frac{x^6}{720}$$

output `1/10*x^2-5/36*x^3+7/96*x^4-1/60*x^5+1/720*x^6`

3.283.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00

$$\int \frac{1}{120}(-4+x)(-3+x)(-2+x)(-1+x)x dx = \frac{1}{120} \left(12x^2 - \frac{50x^3}{3} + \frac{35x^4}{4} - 2x^5 + \frac{x^6}{6} \right)$$

input `Integrate[((-4 + x)*(-3 + x)*(-2 + x)*(-1 + x)*x)/120,x]`

output `(12*x^2 - (50*x^3)/3 + (35*x^4)/4 - 2*x^5 + x^6/6)/120`

3.283.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {27, 2109, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{120}(x-4)(x-3)(x-2)(x-1)x dx$$

$$\downarrow 27$$

$$\frac{1}{120} \int (1-x)(2-x)(3-x)(4-x)x dx$$

$$\downarrow 2109$$

$$\frac{1}{120} \int (x^5 - 10x^4 + 35x^3 - 50x^2 + 24x) dx$$

$$\downarrow 2009$$

$$\frac{1}{120} \left(\frac{x^6}{6} - 2x^5 + \frac{35x^4}{4} - \frac{50x^3}{3} + 12x^2 \right)$$

input `Int[((-4 + x)*(-3 + x)*(-2 + x)*(-1 + x)*x)/120,x]`

output `(12*x^2 - (50*x^3)/3 + (35*x^4)/4 - 2*x^5 + x^6/6)/120`

3.283.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2109 `Int[(P_x)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_)*((g_) + (h_)*(x_))^(q_), x_Symbol] := Int[ExpandIntegrand[P_x*(a + b*x)^m*(c + d*x)^n*(e + f*x)^p*(g + h*x)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n, p, q}, x] && PolyQ[P_x, x] && IntegersQ[m, n]`

3.283. $\int \frac{1}{120}(-4+x)(-3+x)(-2+x)(-1+x)x dx$

3.283.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.75

method	result	size
gosper	$\frac{1}{10}x^2 - \frac{5}{36}x^3 + \frac{7}{96}x^4 - \frac{1}{60}x^5 + \frac{1}{720}x^6$	27
default	$\frac{1}{10}x^2 - \frac{5}{36}x^3 + \frac{7}{96}x^4 - \frac{1}{60}x^5 + \frac{1}{720}x^6$	27
norman	$\frac{1}{10}x^2 - \frac{5}{36}x^3 + \frac{7}{96}x^4 - \frac{1}{60}x^5 + \frac{1}{720}x^6$	27
risch	$\frac{1}{10}x^2 - \frac{5}{36}x^3 + \frac{7}{96}x^4 - \frac{1}{60}x^5 + \frac{1}{720}x^6$	27
parallelrisch	$\frac{1}{10}x^2 - \frac{5}{36}x^3 + \frac{7}{96}x^4 - \frac{1}{60}x^5 + \frac{1}{720}x^6$	27

input `int(1/120*(x-4)*(-3+x)*(-2+x)*(-1+x)*x,x,method=_RETURNVERBOSE)`output `1/10*x^2-5/36*x^3+7/96*x^4-1/60*x^5+1/720*x^6`**3.283.5 Fracas [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.72

$$\int \frac{1}{120}(-4+x)(-3+x)(-2+x)(-1+x)x dx = \frac{1}{720}x^6 - \frac{1}{60}x^5 + \frac{7}{96}x^4 - \frac{5}{36}x^3 + \frac{1}{10}x^2$$

input `integrate(1/120*(x-4)*(-3+x)*(-2+x)*(-1+x)*x,x, algorithm="fricas")`output `1/720*x^6 - 1/60*x^5 + 7/96*x^4 - 5/36*x^3 + 1/10*x^2`**3.283.6 Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.75

$$\int \frac{1}{120}(-4+x)(-3+x)(-2+x)(-1+x)x dx = \frac{x^6}{720} - \frac{x^5}{60} + \frac{7x^4}{96} - \frac{5x^3}{36} + \frac{x^2}{10}$$

input `integrate(1/120*(x-4)*(-3+x)*(-2+x)*(-1+x)*x,x)`output `x**6/720 - x**5/60 + 7*x**4/96 - 5*x**3/36 + x**2/10`

3.283. $\int \frac{1}{120}(-4+x)(-3+x)(-2+x)(-1+x)x dx$

3.283.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.72

$$\int \frac{1}{120}(-4+x)(-3+x)(-2+x)(-1+x)x dx = \frac{1}{720}x^6 - \frac{1}{60}x^5 + \frac{7}{96}x^4 - \frac{5}{36}x^3 + \frac{1}{10}x^2$$

input `integrate(1/120*(x-4)*(-3+x)*(-2+x)*(-1+x)*x,x, algorithm="maxima")`output `1/720*x^6 - 1/60*x^5 + 7/96*x^4 - 5/36*x^3 + 1/10*x^2`**3.283.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.64

$$\int \frac{1}{120}(-4+x)(-3+x)(-2+x)(-1+x)x dx = \frac{1}{720}(x^2 - 4x)^3 + \frac{1}{160}(x^2 - 4x)^2$$

input `integrate(1/120*(x-4)*(-3+x)*(-2+x)*(-1+x)*x,x, algorithm="giac")`output `1/720*(x^2 - 4*x)^3 + 1/160*(x^2 - 4*x)^2`**3.283.9 Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.72

$$\int \frac{1}{120}(-4+x)(-3+x)(-2+x)(-1+x)x dx = \frac{x^6}{720} - \frac{x^5}{60} + \frac{7x^4}{96} - \frac{5x^3}{36} + \frac{x^2}{10}$$

input `int((x*(x - 1)*(x - 2)*(x - 3)*(x - 4))/120,x)`output `x^2/10 - (5*x^3)/36 + (7*x^4)/96 - x^5/60 + x^6/720`

3.284 $\int (x + x \cos(x) + \sin(x) + \cos(x) \sin(x)) dx$

3.284.1 Optimal result	1423
3.284.2 Mathematica [A] (verified)	1423
3.284.3 Rubi [A] (verified)	1424
3.284.4 Maple [A] (verified)	1424
3.284.5 Fricas [A] (verification not implemented)	1425
3.284.6 Sympy [B] (verification not implemented)	1425
3.284.7 Maxima [A] (verification not implemented)	1425
3.284.8 Giac [A] (verification not implemented)	1426
3.284.9 Mupad [B] (verification not implemented)	1426

3.284.1 Optimal result

Integrand size = 13, antiderivative size = 10

$$\int (x + x \cos(x) + \sin(x) + \cos(x) \sin(x)) dx = \frac{1}{2}(x + \sin(x))^2$$

output `1/2*(x+sin(x))^2`

3.284.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 2.00

$$\int (x + x \cos(x) + \sin(x) + \cos(x) \sin(x)) dx = \frac{x^2}{2} - \frac{\cos^2(x)}{2} + x \sin(x)$$

input `Integrate[x + x*Cos[x] + Sin[x] + Cos[x]*Sin[x],x]`

output `x^2/2 - Cos[x]^2/2 + x*Sin[x]`

3.284.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 2.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x + \sin(x) + x \cos(x) + \sin(x) \cos(x)) dx$$

$$\downarrow \text{2009}$$

$$\frac{x^2}{2} + \frac{\sin^2(x)}{2} + x \sin(x)$$

input `Int[x + x*Cos[x] + Sin[x] + Cos[x]*Sin[x],x]`

output `x^2/2 + x*Sin[x] + Sin[x]^2/2`

3.284.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.284.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.70

method	result	size
default	$\frac{x^2}{2} + x \sin(x) + \frac{\sin(x)^2}{2}$	17
risch	$\frac{x^2}{2} + x \sin(x) - \frac{\cos(2x)}{4}$	17
parts	$\frac{x^2}{2} - \frac{\cos(x)^2}{2} + x \sin(x)$	17
norman	$\frac{x^2 \tan(\frac{x}{2})^2 + 2 \tan(\frac{x}{2})^2 + \frac{x^2}{2} + 2x \tan(\frac{x}{2}) + 2x \tan(\frac{x}{2})^3 + \frac{x^2 \tan(\frac{x}{2})^4}{2}}{(1 + \tan(\frac{x}{2})^2)^2}$	63

input `int(x+sin(x)+x*cos(x)+cos(x)*sin(x),x,method=_RETURNVERBOSE)`

output `1/2*x^2+x*sin(x)+1/2*sin(x)^2`

3.284.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.60

$$\int (x + x \cos(x) + \sin(x) + \cos(x) \sin(x)) dx = \frac{1}{2} x^2 - \frac{1}{2} \cos(x)^2 + x \sin(x)$$

input `integrate(x+sin(x)+x*cos(x)+cos(x)*sin(x),x, algorithm="fricas")`

output `1/2*x^2 - 1/2*cos(x)^2 + x*sin(x)`

3.284.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. 2(7) = 14.

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.50

$$\int (x + x \cos(x) + \sin(x) + \cos(x) \sin(x)) dx = \frac{x^2}{2} + x \sin(x) + \frac{\sin^2(x)}{2}$$

input `integrate(x+sin(x)+x*cos(x)+cos(x)*sin(x),x)`

output `x**2/2 + x*sin(x) + sin(x)**2/2`

3.284.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.60

$$\int (x + x \cos(x) + \sin(x) + \cos(x) \sin(x)) dx = \frac{1}{2} x^2 - \frac{1}{2} \cos(x)^2 + x \sin(x)$$

input `integrate(x+sin(x)+x*cos(x)+cos(x)*sin(x),x, algorithm="maxima")`

output `1/2*x^2 - 1/2*cos(x)^2 + x*sin(x)`

3.284.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.60

$$\int (x + x \cos(x) + \sin(x) + \cos(x) \sin(x)) dx = \frac{1}{2} x^2 - \frac{1}{2} \cos(x)^2 + x \sin(x)$$

input `integrate(x+sin(x)+x*cos(x)+cos(x)*sin(x),x, algorithm="giac")`output `1/2*x^2 - 1/2*cos(x)^2 + x*sin(x)`**3.284.9 Mupad [B] (verification not implemented)**

Time = 15.37 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int (x + x \cos(x) + \sin(x) + \cos(x) \sin(x)) dx = \frac{(x + \sin(x))^2}{2}$$

input `int(x + sin(x) + cos(x)*sin(x) + x*cos(x),x)`output `(x + sin(x))^2/2`

3.285 $\int (\cos^2(x) + \cot^2(x) + \csc^2(x) + \sec^2(x) + \sin^2(x) + \tan^2(x) + \dots)$

3.285.1 Optimal result	1427
3.285.2 Mathematica [A] (verified)	1427
3.285.3 Rubi [A] (verified)	1428
3.285.4 Maple [A] (verified)	1428
3.285.5 Fricas [A] (verification not implemented)	1429
3.285.6 Sympy [A] (verification not implemented)	1429
3.285.7 Maxima [A] (verification not implemented)	1429
3.285.8 Giac [B] (verification not implemented)	1430
3.285.9 Mupad [B] (verification not implemented)	1430

3.285.1 Optimal result

Integrand size = 25, antiderivative size = 12

$$\int (\cos^2(x) + \cot^2(x) + \csc^2(x) + \sec^2(x) + \sin^2(x) + \tan^2(x)) dx = -x - 2 \cot(x) + 2 \tan(x)$$

output `-x-2*cot(x)+2*tan(x)`

3.285.2 Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int (\cos^2(x) + \cot^2(x) + \csc^2(x) + \sec^2(x) + \sin^2(x) + \tan^2(x)) dx = -x - 2 \cot(x) + 2 \tan(x)$$

input `Integrate[Cos[x]^2 + Cot[x]^2 + Csc[x]^2 + Sec[x]^2 + Sin[x]^2 + Tan[x]^2, x]`

output `-x - 2*Cot[x] + 2*Tan[x]`

3.285.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\sin^2(x) + \cos^2(x) + \tan^2(x) + \cot^2(x) + \csc^2(x) + \sec^2(x)) dx$$

↓ 2009

$$-x + 2 \tan(x) - 2 \cot(x)$$

input `Int[Cos[x]^2 + Cot[x]^2 + Csc[x]^2 + Sec[x]^2 + Sin[x]^2 + Tan[x]^2,x]`

output `-x - 2*Cot[x] + 2*Tan[x]`

3.285.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.285.4 Maple [A] (verified)

Time = 1.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.25

method	result	size
parallelrisc	$-x - 4 \cot(x) + 2 \csc(x) \sec(x)$	15
default	$x - 2 \cot(x) + \frac{\pi}{2} - \operatorname{arccot}(\cot(x)) + 2 \tan(x) - \arctan(\tan(x))$	24
parts	$x - 2 \cot(x) + \frac{\pi}{2} - \operatorname{arccot}(\cot(x)) + 2 \tan(x) - \arctan(\tan(x))$	24
risc	$\frac{4i}{e^{2ix}+1} - x - \frac{4i}{e^{2ix}-1}$	29

input `int(sin(x)^2+cos(x)^2+tan(x)^2+cot(x)^2+sec(x)^2+csc(x)^2,x,method=_RETURN
VERBOSE)`

output `-x-4*cot(x)+2*csc(x)*sec(x)`

3.285. $\int (\cos^2(x) + \cot^2(x) + \csc^2(x) + \sec^2(x) + \sin^2(x) + \tan^2(x)) dx$

3.285.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 2.00

$$\int (\cos^2(x) + \cot^2(x) + \csc^2(x) + \sec^2(x) + \sin^2(x) + \tan^2(x)) dx$$

$$= -\frac{x \cos(x) \sin(x) + 4 \cos(x)^2 - 2}{\cos(x) \sin(x)}$$

```
input integrate(sin(x)^2+cos(x)^2+tan(x)^2+cot(x)^2+sec(x)^2+csc(x)^2,x, algorit
hm="fricas")
```

```
output -(x*cos(x)*sin(x) + 4*cos(x)^2 - 2)/(cos(x)*sin(x))
```

3.285.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.42

$$\int (\cos^2(x) + \cot^2(x) + \csc^2(x) + \sec^2(x) + \sin^2(x) + \tan^2(x)) dx = -x + \frac{2 \sin(x)}{\cos(x)} - \frac{2 \cos(x)}{\sin(x)}$$

```
input integrate(sin(x)**2+cos(x)**2+tan(x)**2+cot(x)**2+sec(x)**2+csc(x)**2,x)
```

```
output -x + 2*sin(x)/cos(x) - 2*cos(x)/sin(x)
```

3.285.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int (\cos^2(x) + \cot^2(x) + \csc^2(x) + \sec^2(x) + \sin^2(x) + \tan^2(x)) dx = -x - \frac{2}{\tan(x)} + 2 \tan(x)$$

```
input integrate(sin(x)^2+cos(x)^2+tan(x)^2+cot(x)^2+sec(x)^2+csc(x)^2,x, algorit
hm="maxima")
```

```
output -x - 2/tan(x) + 2*tan(x)
```

3.285.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 28 vs. $2(12) = 24$.

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 2.33

$$\int (\cos^2(x) + \cot^2(x) + \csc^2(x) + \sec^2(x) + \sin^2(x) + \tan^2(x)) dx$$

$$= -x - \frac{1}{2 \tan\left(\frac{1}{2}x\right)} - \frac{1}{\tan(x)} + \frac{1}{2} \tan\left(\frac{1}{2}x\right) + 2 \tan(x)$$

input `integrate(sin(x)^2+cos(x)^2+tan(x)^2+cot(x)^2+sec(x)^2+csc(x)^2,x, algorith
hm="giac")`

output `-x - 1/2/tan(1/2*x) - 1/tan(x) + 1/2*tan(1/2*x) + 2*tan(x)`

3.285.9 Mupad [B] (verification not implemented)

Time = 16.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int (\cos^2(x) + \cot^2(x) + \csc^2(x) + \sec^2(x) + \sin^2(x) + \tan^2(x)) dx = 2 \tan(x) - x - \frac{2}{\tan(x)}$$

input `int(1/cos(x)^2 + cos(x)^2 + cot(x)^2 + 1/sin(x)^2 + sin(x)^2 + tan(x)^2,x)`

output `2*tan(x) - x - 2/tan(x)`

3.286 $\int e^{\log^2(x)}(1 + 2 \log(x)) dx$

3.286.1 Optimal result	1431
3.286.2 Mathematica [A] (verified)	1431
3.286.3 Rubi [A] (verified)	1432
3.286.4 Maple [A] (verified)	1432
3.286.5 Fricas [A] (verification not implemented)	1433
3.286.6 Sympy [A] (verification not implemented)	1433
3.286.7 Maxima [A] (verification not implemented)	1433
3.286.8 Giac [A] (verification not implemented)	1434
3.286.9 Mupad [B] (verification not implemented)	1434

3.286.1 Optimal result

Integrand size = 13, antiderivative size = 8

$$\int e^{\log^2(x)}(1 + 2 \log(x)) dx = e^{\log^2(x)}x$$

output `x*exp(ln(x)^2)`

3.286.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int e^{\log^2(x)}(1 + 2 \log(x)) dx = e^{\log^2(x)}x$$

input `Integrate[E^Log[x]^2*(1 + 2*Log[x]),x]`

output `E^Log[x]^2*x`

3.286.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {2726}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{\log^2(x)}(2 \log(x) + 1) dx$$

$$\downarrow \text{2726}$$

$$x e^{\log^2(x)}$$

input `Int[E^Log[x]^2*(1 + 2*Log[x]),x]`

output `E^Log[x]^2*x`

3.286.3.1 Defintions of rubi rules used

rule 2726 `Int[(y_.)*(F_)^(u_)*((v_) + (w_)), x_Symbol] := With[{z = v*(y/(Log[F]*D[u, x]))}, Simp[F^u*z, x] /; EqQ[D[z, x], w*y]] /; FreeQ[F, x]`

3.286.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

method	result	size
norman	$x e^{\ln(x)^2}$	8
risch	$x e^{\ln(x)^2}$	8
parallelrisch	$x e^{\ln(x)^2}$	8

input `int((1+2*ln(x))*exp(ln(x)^2),x,method=_RETURNVERBOSE)`

output `x*exp(ln(x)^2)`

3.286.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int e^{\log^2(x)}(1 + 2\log(x)) dx = xe^{(\log(x)^2)}$$

input `integrate((1+2*log(x))*exp(log(x)^2),x, algorithm="fracas")`output `x*e^(log(x)^2)`**3.286.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int e^{\log^2(x)}(1 + 2\log(x)) dx = xe^{\log(x)^2}$$

input `integrate((1+2*ln(x))*exp(ln(x)**2),x)`output `x*exp(log(x)**2)`**3.286.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int e^{\log^2(x)}(1 + 2\log(x)) dx = xe^{(\log(x)^2)}$$

input `integrate((1+2*log(x))*exp(log(x)^2),x, algorithm="maxima")`output `x*e^(log(x)^2)`

3.286.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int e^{\log^2(x)}(1 + 2 \log(x)) dx = x e^{(\log(x)^2)}$$

input `integrate((1+2*log(x))*exp(log(x)^2),x, algorithm="giac")`

output `x*e^(log(x)^2)`

3.286.9 Mupad [B] (verification not implemented)

Time = 18.45 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int e^{\log^2(x)}(1 + 2 \log(x)) dx = x e^{\ln(x)^2}$$

input `int(exp(log(x)^2)*(2*log(x) + 1),x)`

output `x*exp(log(x)^2)`

$$3.287 \quad \int \left((1-x)^3 + (x-x^2)^3 - 3(1-x)(x-x^2)(-1+x^2) + (-1+x^2)^3 \right) dx = 0$$

3.287.1 Optimal result	1435
3.287.2 Mathematica [A] (verified)	1435
3.287.3 Rubi [B] (verified)	1436
3.287.4 Maple [A] (verified)	1436
3.287.5 Fricas [A] (verification not implemented)	1437
3.287.6 Sympy [A] (verification not implemented)	1437
3.287.7 Maxima [A] (verification not implemented)	1437
3.287.8 Giac [C] (verification not implemented)	1438
3.287.9 Mupad [B] (verification not implemented)	1438

3.287.1 Optimal result

Integrand size = 43, antiderivative size = 1

$$\int \left((1-x)^3 + (x-x^2)^3 - 3(1-x)(x-x^2)(-1+x^2) + (-1+x^2)^3 \right) dx = 0$$

output 0

3.287.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 1, normalized size of antiderivative = 1.00

$$\int \left((1-x)^3 + (x-x^2)^3 - 3(1-x)(x-x^2)(-1+x^2) + (-1+x^2)^3 \right) dx = 0$$

input `Integrate[(1 - x)^3 + (x - x^2)^3 - 3*(1 - x)*(x - x^2)*(-1 + x^2) + (-1 + x^2)^3, x]`

output 0

$$3.287. \quad \int \left((1-x)^3 + (x-x^2)^3 - 3(1-x)(x-x^2)(-1+x^2) + (-1+x^2)^3 \right) dx$$

3.287.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 3 vs. $2(1) = 2$.

Time = 0.16 (sec) , antiderivative size = 3, normalized size of antiderivative = 3.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.023$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(-3(x-x^2)(x^2-1)(1-x) + (x-x^2)^3 + (x^2-1)^3 + (1-x)^3 \right) dx$$

$$\downarrow \text{2009}$$

$$-\frac{1}{4}$$

input `Int[(1 - x)^3 + (x - x^2)^3 - 3*(1 - x)*(x - x^2)*(-1 + x^2) + (-1 + x^2)^3, x]`

output `-1/4`

3.287.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.287.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 2, normalized size of antiderivative = 2.00

method	result	size
default	0	2
norman	0	2
meijerg	0	2
risch	$-\frac{1}{4}$	2
parallelrisch	0	2

input `int((1-x)^3+(-x^2+x)^3+(x^2-1)^3-3*(1-x)*(-x^2+x)*(x^2-1), x, method=_RETURN VERBOSE)`

3.287. $\int \left((1-x)^3 + (x-x^2)^3 - 3(1-x)(x-x^2)(-1+x^2) + (-1+x^2)^3 \right) dx$

output 0

3.287.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 1, normalized size of antiderivative = 1.00

$$\int \left((1-x)^3 + (x-x^2)^3 - 3(1-x)(x-x^2)(-1+x^2) + (-1+x^2)^3 \right) dx = 0$$

```
input integrate((1-x)^3+(-x^2+x)^3+(x^2-1)^3-3*(1-x)*(-x^2+x)*(x^2-1),x, algorith
hm="fricas")
```

output 0

3.287.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 0, normalized size of antiderivative = 0.00

$$\int \left((1-x)^3 + (x-x^2)^3 - 3(1-x)(x-x^2)(-1+x^2) + (-1+x^2)^3 \right) dx = 0$$

```
input integrate((1-x)**3+(-x**2+x)**3+(x**2-1)**3-3*(1-x)*(-x**2+x)*(x**2-1),x)
```

output 0

3.287.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 1, normalized size of antiderivative = 1.00

$$\int \left((1-x)^3 + (x-x^2)^3 - 3(1-x)(x-x^2)(-1+x^2) + (-1+x^2)^3 \right) dx = 0$$

```
input integrate((1-x)^3+(-x^2+x)^3+(x^2-1)^3-3*(1-x)*(-x^2+x)*(x^2-1),x, algorith
hm="maxima")
```

output 0

3.287. $\int \left((1-x)^3 + (x-x^2)^3 - 3(1-x)(x-x^2)(-1+x^2) + (-1+x^2)^3 \right) dx$

3.287.8 Giac [C] (verification not implemented)

Result contains higher order function than in optimal. Order 3 vs. order 1.

Time = 0.27 (sec) , antiderivative size = 26, normalized size of antiderivative = 26.00

$$\int \left((1-x)^3 + (x-x^2)^3 - 3(1-x)(x-x^2)(-1+x^2) + (-1+x^2)^3 \right) dx$$

$$= -\frac{1}{4}(x-1)^4 + \frac{1}{4}x^4 - x^3 + \frac{3}{2}x^2 - x$$

input `integrate((1-x)^3+(-x^2+x)^3+(x^2-1)^3-3*(1-x)*(-x^2+x)*(x^2-1),x, algorithm="giac")`

output `-1/4*(x - 1)^4 + 1/4*x^4 - x^3 + 3/2*x^2 - x`

3.287.9 Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 1, normalized size of antiderivative = 1.00

$$\int \left((1-x)^3 + (x-x^2)^3 - 3(1-x)(x-x^2)(-1+x^2) + (-1+x^2)^3 \right) dx = 0$$

input `int((x - x^2)^3 - (x - 1)^3 + (x^2 - 1)^3 + 3*(x - x^2)*(x^2 - 1)*(x - 1), x)`

output `0`

3.288 $\int (\cos^6(x) + 3 \cos^2(x) \sin^2(x) + \sin^6(x)) dx$

3.288.1 Optimal result	1439
3.288.2 Mathematica [A] (verified)	1439
3.288.3 Rubi [C] (verified)	1440
3.288.4 Maple [A] (verified)	1440
3.288.5 Fricas [A] (verification not implemented)	1441
3.288.6 Sympy [B] (verification not implemented)	1441
3.288.7 Maxima [A] (verification not implemented)	1442
3.288.8 Giac [A] (verification not implemented)	1442
3.288.9 Mupad [B] (verification not implemented)	1442

3.288.1 Optimal result

Integrand size = 19, antiderivative size = 1

$$\int (\cos^6(x) + 3 \cos^2(x) \sin^2(x) + \sin^6(x)) dx = x$$

output

x

3.288.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 1, normalized size of antiderivative = 1.00

$$\int (\cos^6(x) + 3 \cos^2(x) \sin^2(x) + \sin^6(x)) dx = x$$

input

Integrate[Cos[x]^6 + 3*Cos[x]^2*Sin[x]^2 + Sin[x]^6,x]

output

x

3.288.3 Rubi [C] (verified)

Result contains higher order function than in optimal. Order 3 vs. order 1 in optimal.

Time = 0.21 (sec) , antiderivative size = 50, normalized size of antiderivative = 50.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\sin^6(x) + \cos^6(x) + 3 \sin^2(x) \cos^2(x)) dx$$

↓ 2009

$$x + \frac{1}{6} \sin(x) \cos^5(x) - \frac{13}{24} \sin(x) \cos^3(x) - \frac{1}{6} \sin^5(x) \cos(x) - \frac{5}{24} \sin^3(x) \cos(x) + \frac{3}{8} \sin(x) \cos(x)$$

input `Int[Cos[x]^6 + 3*Cos[x]^2*Sin[x]^2 + Sin[x]^6,x]`

output `x + (3*Cos[x]*Sin[x])/8 - (13*Cos[x]^3*Sin[x])/24 + (Cos[x]^5*Sin[x])/6 - (5*Cos[x]*Sin[x]^3)/24 - (Cos[x]*Sin[x]^5)/6`

3.288.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.288.4 Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 2, normalized size of antiderivative = 2.00

method	result	si
risch	x	2
default	$\frac{(\cos(x)^5 + \frac{5 \cos(x)^3}{4} + \frac{15 \cos(x)}{8}) \sin(x)}{6} + x - \frac{(\sin(x)^5 + \frac{5 \sin(x)^3}{4} + \frac{15 \sin(x)}{8}) \cos(x)}{6} - \frac{3 \cos(x)^3 \sin(x)}{4} + \frac{3 \cos(x) \sin(x)}{8}$	5
parts	$\frac{(\cos(x)^5 + \frac{5 \cos(x)^3}{4} + \frac{15 \cos(x)}{8}) \sin(x)}{6} + x - \frac{(\sin(x)^5 + \frac{5 \sin(x)^3}{4} + \frac{15 \sin(x)}{8}) \cos(x)}{6} - \frac{3 \cos(x)^3 \sin(x)}{4} + \frac{3 \cos(x) \sin(x)}{8}$	5
norman	$\frac{x + x \tan(\frac{x}{2})^{12} + 6x \tan(\frac{x}{2})^2 + 15x \tan(\frac{x}{2})^4 + 20x \tan(\frac{x}{2})^6 + 15x \tan(\frac{x}{2})^8 + 6x \tan(\frac{x}{2})^{10}}{(1 + \tan(\frac{x}{2})^2)^6}$	6

3.288. $\int (\cos^6(x) + 3 \cos^2(x) \sin^2(x) + \sin^6(x)) dx$

input `int(sin(x)^6+cos(x)^6+3*sin(x)^2*cos(x)^2,x,method=_RETURNVERBOSE)`

output `x`

3.288.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 1, normalized size of antiderivative = 1.00

$$\int (\cos^6(x) + 3 \cos^2(x) \sin^2(x) + \sin^6(x)) dx = x$$

input `integrate(sin(x)^6+cos(x)^6+3*sin(x)^2*cos(x)^2,x, algorithm="fricas")`

output `x`

3.288.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 58 vs. 2(0) = 0.

Time = 0.03 (sec) , antiderivative size = 58, normalized size of antiderivative = 58.00

$$\int (\cos^6(x) + 3 \cos^2(x) \sin^2(x) + \sin^6(x)) dx = x - \frac{\sin^5(x) \cos(x)}{6} - \frac{5 \sin^3(x) \cos(x)}{24} + \frac{\sin(x) \cos^5(x)}{6} + \frac{5 \sin(x) \cos^3(x)}{24} - \frac{3 \sin(2x) \cos(2x)}{16}$$

input `integrate(sin(x)**6+cos(x)**6+3*sin(x)**2*cos(x)**2,x)`

output `x - sin(x)**5*cos(x)/6 - 5*sin(x)**3*cos(x)/24 + sin(x)*cos(x)**5/6 + 5*sin(x)*cos(x)**3/24 - 3*sin(2*x)*cos(2*x)/16`

3.288.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 1, normalized size of antiderivative = 1.00

$$\int (\cos^6(x) + 3 \cos^2(x) \sin^2(x) + \sin^6(x)) dx = x$$

input `integrate(sin(x)^6+cos(x)^6+3*sin(x)^2*cos(x)^2,x, algorithm="maxima")`output `x`**3.288.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 1, normalized size of antiderivative = 1.00

$$\int (\cos^6(x) + 3 \cos^2(x) \sin^2(x) + \sin^6(x)) dx = x$$

input `integrate(sin(x)^6+cos(x)^6+3*sin(x)^2*cos(x)^2,x, algorithm="giac")`output `x`**3.288.9 Mupad [B] (verification not implemented)**

Time = 18.14 (sec) , antiderivative size = 1, normalized size of antiderivative = 1.00

$$\int (\cos^6(x) + 3 \cos^2(x) \sin^2(x) + \sin^6(x)) dx = x$$

input `int(cos(x)^6 + sin(x)^6 + 3*cos(x)^2*sin(x)^2,x)`output `x`

3.289 $\int e^x x^e (1 + e + x) dx$

3.289.1 Optimal result	1443
3.289.2 Mathematica [A] (verified)	1443
3.289.3 Rubi [A] (verified)	1444
3.289.4 Maple [A] (verified)	1444
3.289.5 Fricas [A] (verification not implemented)	1445
3.289.6 Sympy [A] (verification not implemented)	1445
3.289.7 Maxima [C] (verification not implemented)	1445
3.289.8 Giac [A] (verification not implemented)	1446
3.289.9 Mupad [B] (verification not implemented)	1446

3.289.1 Optimal result

Integrand size = 11, antiderivative size = 9

$$\int e^x x^e (1 + e + x) dx = e^x x^{1+e}$$

output `exp(x)*x^(1+exp(1))`

3.289.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int e^x x^e (1 + e + x) dx = e^x x^{1+e}$$

input `Integrate[E^x*x^E*(1 + E + x),x]`

output `E^x*x^(1 + E)`

3.289.3 Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {2627}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x x^e (x + e + 1) dx$$

↓ 2627

$$e^x x^{1+e}$$

input `Int [E^x*x^E*(1 + E + x), x]`

output `E^x*x^(1 + E)`

3.289.3.1 Defintions of rubi rules used

rule 2627 `Int[(F_)^(v_)*((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_)), x_Symbol] :> Simp[g*(d + e*x)^(m + 1)*(F^v/(D[v, x]*e*Log[F])), x] /; FreeQ[{F, d, e, f, g, m}, x] && LinearQ[v, x] && EqQ[e*g*(m + 1) - D[v, x]*(e*f - d*g)*Log[F], 0]`

3.289.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

method	result
risch	$x x^e e^x$
parallelrisch	$x x^e e^x$
gospers	$e^x x^{1+e}$
norman	$x e^x e^{e \ln(x)}$
meijerg	$(-1)^{-e} (x^e (-1)^e (1 + e) e \Gamma(e) (-x)^{-e} - x^e (-1)^e (e - x + 1) e^x - x^e (-1)^e (1 + e) e (-x)^{-e} \Gamma(e))$

input `int((x+exp(1)+1)*x^exp(1)*exp(x), x, method=_RETURNVERBOSE)`

output `x*x^exp(1)*exp(x)`

3.289.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int e^x x^e (1 + e + x) dx = x x^e e^x$$

input `integrate((x+exp(1)+1)*x^exp(1)*exp(x),x, algorithm="fricas")`

output `x*x^e*e^x`

3.289.6 Sympy [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int e^x x^e (1 + e + x) dx = x x^e e^x$$

input `integrate((x+exp(1)+1)*x**exp(1)*exp(x),x)`

output `x*x**E*exp(x)`

3.289.7 Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 0.23 (sec) , antiderivative size = 81, normalized size of antiderivative = 9.00

$$\int e^x x^e (1 + e + x) dx = -(-x)^{-e-1} x^{e+1} e \Gamma(e+1, -x) \\ - (-x)^{-e-2} x^{e+2} \Gamma(e+2, -x) - (-x)^{-e-1} x^{e+1} \Gamma(e+1, -x)$$

input `integrate((x+exp(1)+1)*x^exp(1)*exp(x),x, algorithm="maxima")`

output `-(-x)^(-e - 1)*x^(e + 1)*e*gamma(e + 1, -x) - (-x)^(-e - 2)*x^(e + 2)*gamma(e + 2, -x) - (-x)^(-e - 1)*x^(e + 1)*gamma(e + 1, -x)`

3.289.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int e^x x^e (1 + e + x) dx = x x^e e^x$$

input `integrate((x+exp(1)+1)*x^exp(1)*exp(x),x, algorithm="giac")`

output `x*x^e*e^x`

3.289.9 Mupad [B] (verification not implemented)

Time = 18.14 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int e^x x^e (1 + e + x) dx = x^{e+1} e^x$$

input `int(x^exp(1)*exp(x)*(x + exp(1) + 1),x)`

output `x^(exp(1) + 1)*exp(x)`

3.290 $\int \left(\sqrt{2} \sqrt{\frac{x}{1+x}} + \frac{x^2}{2-x^2} \right) dx$

3.290.1 Optimal result 1447
 3.290.2 Mathematica [B] (verified) 1447
 3.290.3 Rubi [A] (verified) 1448
 3.290.4 Maple [A] (verified) 1448
 3.290.5 Fricas [B] (verification not implemented) 1449
 3.290.6 Sympy [F] 1449
 3.290.7 Maxima [B] (verification not implemented) 1450
 3.290.8 Giac [B] (verification not implemented) 1450
 3.290.9 Mupad [B] (verification not implemented) 1451

3.290.1 Optimal result

Integrand size = 31, antiderivative size = 49

$$\int \left(\sqrt{2} \sqrt{\frac{x}{1+x}} + \frac{x^2}{2-x^2} \right) dx = -x + \sqrt{2} \sqrt{x} \sqrt{1+x} - \sqrt{2} \operatorname{arcsinh}(\sqrt{x}) + \sqrt{2} \operatorname{arctanh}\left(\frac{x}{\sqrt{2}}\right)$$

output `-x+2^(1/2)*x^(1/2)*(1+x)^(1/2)-2^(1/2)*arcsinh(x^(1/2))+2^(1/2)*arctanh(1/2*x*2^(1/2))`

3.290.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 100 vs. 2(49) = 98.

Time = 0.08 (sec) , antiderivative size = 100, normalized size of antiderivative = 2.04

$$\int \left(\sqrt{2} \sqrt{\frac{x}{1+x}} + \frac{x^2}{2-x^2} \right) dx = -x + \sqrt{2} \sqrt{\frac{x}{1+x}} (1+x) - \frac{\log(\sqrt{2}-x)}{\sqrt{2}} + \frac{\log(\sqrt{2}+x)}{\sqrt{2}} + \frac{\sqrt{2} \sqrt{\frac{x}{1+x}} \sqrt{1+x} \log(-\sqrt{x} + \sqrt{1+x})}{\sqrt{x}}$$

input `Integrate[Sqrt[2]*Sqrt[x/(1+x)] + x^2/(2-x^2),x]`

output `-x + Sqrt[2]*Sqrt[x/(1+x)]*(1+x) - Log[Sqrt[2]-x]/Sqrt[2] + Log[Sqrt[2]+x]/Sqrt[2] + (Sqrt[2]*Sqrt[x/(1+x)]*Sqrt[1+x]*Log[-Sqrt[x]+Sqrt[1+x]])/Sqrt[x]`

3.290. $\int \left(\sqrt{2} \sqrt{\frac{x}{1+x}} + \frac{x^2}{2-x^2} \right) dx$

3.290.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.032$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\frac{x^2}{2-x^2} + \sqrt{2} \sqrt{\frac{x}{x+1}} \right) dx$$

↓ 2009

$$-\sqrt{2} \operatorname{arcsinh}(\sqrt{x}) + \sqrt{2} \operatorname{arctanh}\left(\frac{x}{\sqrt{2}}\right) - x + \sqrt{2} \sqrt{x+1} \sqrt{x}$$

input `Int[Sqrt[2]*Sqrt[x/(1+x)] + x^2/(2-x^2),x]`

output `-x + Sqrt[2]*Sqrt[x]*Sqrt[1+x] - Sqrt[2]*ArcSinh[Sqrt[x]] + Sqrt[2]*ArcTanh[x/Sqrt[2]]`

3.290.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.290.4 Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.24

method	result
default	$-\frac{\sqrt{2} \sqrt{\frac{x}{1+x}} (1+x) \left(-2\sqrt{x^2+x} + \ln\left(\frac{1}{2} + x + \sqrt{x^2+x}\right) \right)}{2\sqrt{x(1+x)}} - x + \sqrt{2} \operatorname{arctanh}\left(\frac{x\sqrt{2}}{2}\right)$
trager	$1 - x + \sqrt{2} (1+x) \sqrt{\frac{x}{1+x}} - \frac{\sqrt{2} \ln\left(-\frac{2\sqrt{2} \sqrt{\frac{x}{1+x}} x^2 + 2\sqrt{2} x^2 + 2\sqrt{\frac{x}{1+x}} x^2 - 2\sqrt{2} \sqrt{\frac{x}{1+x}} - x\sqrt{2} - 2x\sqrt{\frac{x}{1+x}} + 2x^2 - \sqrt{2} - 4\sqrt{\frac{x}{1+x}} - 3x}{x\sqrt{2} - \sqrt{2} - x + 2}\right)}{2}$

input `int(x^2/(-x^2+2)+2^(1/2)*(x/(1+x))^(1/2),x,method=_RETURNVERBOSE)`

3.290. $\int \left(\sqrt{2} \sqrt{\frac{x}{1+x}} + \frac{x^2}{2-x^2} \right) dx$

output
$$\frac{-1/2 \cdot 2^{1/2} \cdot (x/(1+x))^{1/2} \cdot (1+x) \cdot (-2 \cdot (x^2+x)^{1/2} + \ln(1/2+x+(x^2+x)^{1/2}))}{(x \cdot (1+x))^{1/2} - x + 2^{1/2} \cdot \operatorname{arctanh}(1/2 \cdot x \cdot 2^{1/2})}$$

3.290.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 114 vs. $2(36) = 72$.

Time = 0.25 (sec) , antiderivative size = 114, normalized size of antiderivative = 2.33

$$\int \left(\sqrt{2} \sqrt{\frac{x}{1+x}} + \frac{x^2}{2-x^2} \right) dx = \sqrt{2}(x+1) \sqrt{\frac{x}{x+1}} + \frac{1}{2} \sqrt{2} \log \left(-\frac{6x^3 + 19x^2 + 2\sqrt{2}(2x^3 + 7x^2 + 7x + 2) - 2(3x^3 + 11x^2 + 2\sqrt{2}(x^3 + 4x^2 + 5x + 2) + 14x + 6)}{x^2 - 2} \right) - x$$

input `integrate(x^2/(-x^2+2)+2^(1/2)*(x/(1+x))^(1/2),x, algorithm="fricas")`

output
$$\sqrt{2} \cdot (x + 1) \cdot \sqrt{x/(x + 1)} + 1/2 \cdot \sqrt{2} \cdot \log(-6 \cdot x^3 + 19 \cdot x^2 + 2 \cdot \sqrt{2} \cdot (2 \cdot x^3 + 7 \cdot x^2 + 7 \cdot x + 2) - 2 \cdot (3 \cdot x^3 + 11 \cdot x^2 + 2 \cdot \sqrt{2} \cdot (x^3 + 4 \cdot x^2 + 5 \cdot x + 2) + 14 \cdot x + 6) \cdot \sqrt{x/(x + 1)} + 20 \cdot x + 6)/(x^2 - 2)) - x$$

3.290.6 Sympy [F]

$$\int \left(\sqrt{2} \sqrt{\frac{x}{1+x}} + \frac{x^2}{2-x^2} \right) dx = \int \frac{\sqrt{2} x^2 \sqrt{\frac{x}{x+1}} - x^2 - 2\sqrt{2} \sqrt{\frac{x}{x+1}}}{x^2 - 2} dx$$

input `integrate(x**2/(-x**2+2)+2**(1/2)*(x/(1+x))**(1/2),x)`

output
$$\operatorname{Integral}((\sqrt{2} \cdot x^{**2} \cdot \sqrt{x/(x + 1)} - x^{**2} - 2 \cdot \sqrt{2} \cdot \sqrt{x/(x + 1)})/(x^{**2} - 2), x)$$

3.290.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 79 vs. $2(36) = 72$.

Time = 0.28 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.61

$$\begin{aligned} & \int \left(\sqrt{2} \sqrt{\frac{x}{1+x}} + \frac{x^2}{2-x^2} \right) dx \\ &= -\frac{1}{2} \sqrt{2} \left(\frac{2 \sqrt{\frac{x}{x+1}}}{\frac{x}{x+1} - 1} + \log \left(\sqrt{\frac{x}{x+1}} + 1 \right) - \log \left(\sqrt{\frac{x}{x+1}} - 1 \right) \right) \\ & \quad - \frac{1}{2} \sqrt{2} \log \left(\frac{x - \sqrt{2}}{x + \sqrt{2}} \right) - x \end{aligned}$$

input `integrate(x^2/(-x^2+2)+2^(1/2)*(x/(1+x))^(1/2),x, algorithm="maxima")`

output `-1/2*sqrt(2)*(2*sqrt(x/(x + 1))/(x/(x + 1) - 1) + log(sqrt(x/(x + 1)) + 1) - log(sqrt(x/(x + 1)) - 1)) - 1/2*sqrt(2)*log((x - sqrt(2))/(x + sqrt(2))) - x`

3.290.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. $2(36) = 72$.

Time = 0.29 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.49

$$\begin{aligned} & \int \left(\sqrt{2} \sqrt{\frac{x}{1+x}} + \frac{x^2}{2-x^2} \right) dx \\ &= \frac{1}{2} \sqrt{2} \left(\log \left(\left| -2x + 2\sqrt{x^2 + x} - 1 \right| \right) \operatorname{sgn}(x + 1) + 2\sqrt{x^2 + x} \operatorname{sgn}(x + 1) \right) \\ & \quad - \frac{1}{2} \sqrt{2} \log \left(\frac{|2x - 2\sqrt{2}|}{|2x + 2\sqrt{2}|} \right) - x \end{aligned}$$

input `integrate(x^2/(-x^2+2)+2^(1/2)*(x/(1+x))^(1/2),x, algorithm="giac")`

output `1/2*sqrt(2)*(log(abs(-2*x + 2*sqrt(x^2 + x) - 1))*sgn(x + 1) + 2*sqrt(x^2 + x)*sgn(x + 1)) - 1/2*sqrt(2)*log(abs(2*x - 2*sqrt(2))/abs(2*x + 2*sqrt(2)))) - x`

3.290.9 Mupad [B] (verification not implemented)

Time = 17.80 (sec) , antiderivative size = 238, normalized size of antiderivative = 4.86

$$\int \left(\sqrt{2} \sqrt{\frac{x}{1+x}} + \frac{x^2}{2-x^2} \right) dx$$

$$= -\sqrt{2} \operatorname{atanh} \left(\frac{96 \sqrt{\frac{2x}{x+1}}}{64 \sqrt{2} \sqrt{\frac{2x}{x+1}} - 56 \sqrt{2} \left(\frac{2x}{x+1}\right)^{3/2} + 96 \sqrt{2} - \frac{160 \sqrt{2} x}{x+1}} \right) - \frac{80 \left(\frac{2x}{x+1}\right)^{3/2}}{64 \sqrt{2} \sqrt{\frac{2x}{x+1}} - 56 \sqrt{2} \left(\frac{2x}{x+1}\right)^{3/2} + 96 \sqrt{2} - \frac{160 \sqrt{2} x}{x+1}} + \frac{128}{64 \sqrt{2} \sqrt{\frac{2x}{x+1}} - 56 \sqrt{2} \left(\frac{2x}{x+1}\right)^{3/2} + 96 \sqrt{2} - \frac{160 \sqrt{2} x}{x+1}} - \frac{224 x}{(x+1) \left(64 \sqrt{2} \sqrt{\frac{2x}{x+1}} - 56 \sqrt{2} \left(\frac{2x}{x+1}\right)^{3/2} + 96 \sqrt{2} - \frac{160 \sqrt{2} x}{x+1}\right)} - \frac{\sqrt{\frac{2x}{x+1}} - 1}{\frac{x}{x+1} - 1}$$

input `int(2^(1/2)*(x/(x + 1))^(1/2) - x^2/(x^2 - 2),x)`

output

```
- 2^(1/2)*atanh((96*((2*x)/(x + 1))^(1/2))/(64*2^(1/2)*((2*x)/(x + 1))^(1/2) - 56*2^(1/2)*((2*x)/(x + 1))^(3/2) + 96*2^(1/2) - (160*2^(1/2)*x)/(x + 1)) - (80*((2*x)/(x + 1))^(3/2))/(64*2^(1/2)*((2*x)/(x + 1))^(1/2) - 56*2^(1/2)*((2*x)/(x + 1))^(3/2) + 96*2^(1/2) - (160*2^(1/2)*x)/(x + 1)) + 128/(64*2^(1/2)*((2*x)/(x + 1))^(1/2) - 56*2^(1/2)*((2*x)/(x + 1))^(3/2) + 96*2^(1/2) - (160*2^(1/2)*x)/(x + 1)) - (224*x)/((x + 1)*(64*2^(1/2)*((2*x)/(x + 1))^(1/2) - 56*2^(1/2)*((2*x)/(x + 1))^(3/2) + 96*2^(1/2) - (160*2^(1/2)*x)/(x + 1))) - (((2*x)/(x + 1))^(1/2) - 1)/(x/(x + 1) - 1)
```

$$\mathbf{3.291} \quad \int \frac{1+2x^{2022}}{x+x^{2023}} dx$$

3.291.1 Optimal result	1452
3.291.2 Mathematica [A] (verified)	1452
3.291.3 Rubi [A] (verified)	1453
3.291.4 Maple [F(-1)]	1454
3.291.5 Fricas [F(-1)]	1454
3.291.6 Sympy [F(-1)]	1455
3.291.7 Maxima [A] (verification not implemented)	1455
3.291.8 Giac [A] (verification not implemented)	1455
3.291.9 Mupad [F(-1)]	1456

3.291.1 Optimal result

Integrand size = 15, antiderivative size = 13

$$\int \frac{1+2x^{2022}}{x+x^{2023}} dx = \log(x) + \frac{\log(1+x^{2022})}{2022}$$

output `ln(x)+1/2022*ln(x^2022+1)`

3.291.2 Mathematica [A] (verified)

Time = 3.39 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1+2x^{2022}}{x+x^{2023}} dx = \log(x) + \frac{\log(1+x^{2022})}{2022}$$

input `Integrate[(1 + 2*x^2022)/(x + x^2023),x]`

output `Log[x] + Log[1 + x^2022]/2022`

3.291.3 Rubi [A] (verified)

Time = 1.73 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {2026, 948, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{2x^{2022} + 1}{x^{2023} + x} dx \\
 \downarrow \text{2026} \\
 \int \frac{2x^{2022} + 1}{x(x^{2022} + 1)} dx \\
 \downarrow \text{948} \\
 \int \frac{2x^{2022} + 1}{x^{2022}(x^{2022} + 1)} dx \\
 \frac{2022}{2022} \\
 \downarrow \text{86} \\
 \int \left(\frac{1}{x^{2022}} + \frac{1}{x^{2022} + 1} \right) dx \\
 \frac{2022}{2022} \\
 \downarrow \text{2009} \\
 \frac{\log(x^{2022}) + \log(x^{2022} + 1)}{2022}
 \end{array}$$

input `Int[(1 + 2*x^2022)/(x + x^2023),x]`

output `(Log[x^2022] + Log[1 + x^2022])/2022`

3.291.3.1 Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] :> Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

```
rule 948 Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_
), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^
p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ
[b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2026 Int[(Fx_)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Int[x^(p
*r)*ExpandToSum[Px/x^r, x]^p*Fx, x] /; IGtQ[r, 0]] /; PolyQ[Px, x] && Integ
erQ[p] && !MonomialQ[Px, x] && (ILtQ[p, 0] || !PolyQ[u, x])
```

3.291.4 Maple [F(-1)]

Timed out.

hanged

```
input int((2*x^2022+1)/(x^2023+x),x)
```

```
output int((2*x^2022+1)/(x^2023+x),x)
```

3.291.5 Fracas [F(-1)]

Timed out.

$$\int \frac{1 + 2x^{2022}}{x + x^{2023}} dx = \text{Timed out}$$

```
input integrate((2*x^2022+1)/(x^2023+x),x, algorithm="fricas")
```

```
output Timed out
```

3.291.6 Sympy [F(-1)]

Timed out.

$$\int \frac{1 + 2x^{2022}}{x + x^{2023}} dx = \text{Timed out}$$

input `integrate((2*x**2022+1)/(x**2023+x),x)`output `Timed out`**3.291.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1 + 2x^{2022}}{x + x^{2023}} dx = \frac{1}{2022} \log(x^{2022} + 1) + \log(x)$$

input `integrate((2*x^2022+1)/(x^2023+x),x, algorithm="maxima")`output `1/2022*log(x^2022 + 1) + log(x)`**3.291.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{1 + 2x^{2022}}{x + x^{2023}} dx = \frac{1}{2022} \log(x^{2022} + 1) + \frac{1}{2022} \log(x^{2022})$$

input `integrate((2*x^2022+1)/(x^2023+x),x, algorithm="giac")`output `1/2022*log(x^2022 + 1) + 1/2022*log(x^2022)`

3.291.9 Mupad [F(-1)]

Timed out.

$$\int \frac{1 + 2x^{2022}}{x + x^{2023}} dx = \int \frac{2x^{2022} + 1}{x^{2023} + x} dx$$

input `int((2*x^2022 + 1)/(x + x^2023),x)`output `int((2*x^2022 + 1)/(x + x^2023), x)`

3.292 $\int (3 \cos(23x) \sin(20x) + 20 \sin(43x)) dx$

3.292.1 Optimal result	1457
3.292.2 Mathematica [A] (verified)	1457
3.292.3 Rubi [A] (verified)	1458
3.292.4 Maple [A] (verified)	1458
3.292.5 Fricas [B] (verification not implemented)	1459
3.292.6 Sympy [B] (verification not implemented)	1460
3.292.7 Maxima [A] (verification not implemented)	1460
3.292.8 Giac [A] (verification not implemented)	1460
3.292.9 Mupad [B] (verification not implemented)	1461

3.292.1 Optimal result

Integrand size = 17, antiderivative size = 9

$$\int (3 \cos(23x) \sin(20x) + 20 \sin(43x)) dx = \sin(20x) \sin(23x)$$

output `sin(20*x)*sin(23*x)`

3.292.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.89

$$\int (3 \cos(23x) \sin(20x) + 20 \sin(43x)) dx = \frac{1}{2} \cos(3x) - \frac{1}{2} \cos(43x)$$

input `Integrate[3*Cos[23*x]*Sin[20*x] + 20*Sin[43*x],x]`

output `Cos[3*x]/2 - Cos[43*x]/2`

3.292.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.89, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (20 \sin(43x) + 3 \sin(20x) \cos(23x)) dx$$

↓ 2009

$$\frac{1}{2} \cos(3x) - \frac{1}{2} \cos(43x)$$

input `Int[3*Cos[23*x]*Sin[20*x] + 20*Sin[43*x],x]`

output `Cos[3*x]/2 - Cos[43*x]/2`

3.292.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.292.4 Maple [A] (verified)

Time = 2.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.56

method	result
default	$\frac{\cos(3x)}{2} - \frac{\cos(43x)}{2}$
risch	$\frac{\cos(3x)}{2} - \frac{\cos(43x)}{2}$
parts	$\frac{\cos(3x)}{2} - \frac{\cos(43x)}{2}$
parallelrisc	$-\frac{40}{43} + \frac{40(-\tan(10x)^2 - \tan(\frac{43x}{2})^2 - 2)\tan(\frac{23x}{2})^2}{43} + \frac{92 \tan(\frac{23x}{2}) \tan(10x) (1 + \tan(\frac{43x}{2})^2)}{43} - \frac{40 \tan(\frac{43x}{2})^2 \tan(10x)^2}{43} - \frac{80 \tan(10x)^2}{43}$ $(1 + \tan(10x)^2) (1 + \tan(\frac{23x}{2})^2) (1 + \tan(\frac{43x}{2})^2)$

input `int(3*sin(20*x)*cos(23*x)+20*sin(43*x),x,method=_RETURNVERBOSE)`

output `1/2*cos(3*x)-1/2*cos(43*x)`

3.292.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 131 vs. $2(9) = 18$.

Time = 1.63 (sec) , antiderivative size = 131, normalized size of antiderivative = 14.56

$$\int (3 \cos(23x) \sin(20x) + 20 \sin(43x)) dx = -2199023255552 \cos(x)^{43} \\ + 23639499997184 \cos(x)^{41} \\ - 118197499985920 \cos(x)^{39} \\ + 364934781206528 \cos(x)^{37} \\ - 778995398344704 \cos(x)^{35} \\ + 1219742794776576 \cos(x)^{33} \\ - 1450504945139712 \cos(x)^{31} \\ + 1338263491051520 \cos(x)^{29} \\ - 970241031012352 \cos(x)^{27} \\ + 556461767786496 \cos(x)^{25} \\ - 252937167175680 \cos(x)^{23} \\ + 90899294453760 \cos(x)^{21} \\ - 25657058918400 \cos(x)^{19} \\ + 5624816762880 \cos(x)^{17} \\ - 942087536640 \cos(x)^{15} \\ + 117760942080 \cos(x)^{13} \\ - 10631196160 \cos(x)^{11} + 661443200 \cos(x)^9 \\ - 26457728 \cos(x)^7 + 609224 \cos(x)^5 \\ - 6620 \cos(x)^3 + 20 \cos(x)$$

input `integrate(3*sin(20*x)*cos(23*x)+20*sin(43*x),x, algorithm="fracas")`

output `-2199023255552*cos(x)^43 + 23639499997184*cos(x)^41 - 118197499985920*cos(x)^39 + 364934781206528*cos(x)^37 - 778995398344704*cos(x)^35 + 1219742794776576*cos(x)^33 - 1450504945139712*cos(x)^31 + 1338263491051520*cos(x)^29 - 970241031012352*cos(x)^27 + 556461767786496*cos(x)^25 - 252937167175680*cos(x)^23 + 90899294453760*cos(x)^21 - 25657058918400*cos(x)^19 + 5624816762880*cos(x)^17 - 942087536640*cos(x)^15 + 117760942080*cos(x)^13 - 10631196160*cos(x)^11 + 661443200*cos(x)^9 - 26457728*cos(x)^7 + 609224*cos(x)^5 - 6620*cos(x)^3 + 20*cos(x)`

3.292.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. $2(8) = 16$.

Time = 0.13 (sec) , antiderivative size = 34, normalized size of antiderivative = 3.78

$$\int (3 \cos(23x) \sin(20x) + 20 \sin(43x)) dx = \frac{23 \sin(20x) \sin(23x)}{43} + \frac{20 \cos(20x) \cos(23x)}{43} - \frac{20 \cos(43x)}{43}$$

input `integrate(3*sin(20*x)*cos(23*x)+20*sin(43*x),x)`

output `23*sin(20*x)*sin(23*x)/43 + 20*cos(20*x)*cos(23*x)/43 - 20*cos(43*x)/43`

3.292.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.44

$$\int (3 \cos(23x) \sin(20x) + 20 \sin(43x)) dx = -\frac{1}{2} \cos(43x) + \frac{1}{2} \cos(3x)$$

input `integrate(3*sin(20*x)*cos(23*x)+20*sin(43*x),x, algorithm="maxima")`

output `-1/2*cos(43*x) + 1/2*cos(3*x)`

3.292.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.44

$$\int (3 \cos(23x) \sin(20x) + 20 \sin(43x)) dx = -\frac{1}{2} \cos(43x) + \frac{1}{2} \cos(3x)$$

input `integrate(3*sin(20*x)*cos(23*x)+20*sin(43*x),x, algorithm="giac")`

output `-1/2*cos(43*x) + 1/2*cos(3*x)`

3.292.9 Mupad [B] (verification not implemented)

Time = 17.31 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.44

$$\int (3 \cos(23x) \sin(20x) + 20 \sin(43x)) dx = \frac{\cos(3x)}{2} - \frac{\cos(43x)}{2}$$

input `int(20*sin(43*x) + 3*cos(23*x)*sin(20*x),x)`

output `cos(3*x)/2 - cos(43*x)/2`

3.293 $\int \frac{\sin(x)}{2e^x + \cos(x) + \sin(x)} dx$

3.293.1 Optimal result 1462
 3.293.2 Mathematica [A] (verified) 1462
 3.293.3 Rubi [F] 1463
 3.293.4 Maple [C] (verified) 1463
 3.293.5 Fricas [A] (verification not implemented) 1464
 3.293.6 Sympy [A] (verification not implemented) 1464
 3.293.7 Maxima [B] (verification not implemented) 1464
 3.293.8 Giac [B] (verification not implemented) 1465
 3.293.9 Mupad [B] (verification not implemented) 1465

3.293.1 Optimal result

Integrand size = 15, antiderivative size = 19

$$\int \frac{\sin(x)}{2e^x + \cos(x) + \sin(x)} dx = \frac{1}{2}(x - \log(2e^x + \cos(x) + \sin(x)))$$

output `1/2*x-1/2*ln(2*exp(x)+cos(x)+sin(x))`

3.293.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\sin(x)}{2e^x + \cos(x) + \sin(x)} dx = \frac{x}{2} - \frac{1}{2} \log(2e^x + \cos(x) + \sin(x))$$

input `Integrate[Sin[x]/(2*E^x + Cos[x] + Sin[x]),x]`

output `x/2 - Log[2*E^x + Cos[x] + Sin[x]]/2`

3.293.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sin(x)}{2e^x + \sin(x) + \cos(x)} dx$$

↓ 7299

$$\int \frac{\sin(x)}{2e^x + \sin(x) + \cos(x)} dx$$

input `Int[Sin[x]/(2*E^x + Cos[x] + Sin[x]),x]`

output `$Aborted`

3.293.3.1 Defintions of rubi rules used

rule 7299 `Int[u_, x_] :=> CannotIntegrate[u, x]`

3.293.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.58

method	result	size
risch	$\frac{x}{2} + \frac{ix}{2} - \frac{\ln(e^{2ix} + (2+2i)e^{(1+i)x} + i)}{2}$	30
parallelrisc	$\ln\left(\frac{\sqrt{2}}{\sqrt{2e^x + \cos(x) + \sin(x)}}\right) + \ln\left(\sqrt{\frac{1}{1 + \cos(x)}}\right) + \frac{x}{2}$	37
norman	$\frac{\frac{x}{2} + \frac{x \tan(\frac{x}{2})^2}{2}}{1 + \tan(\frac{x}{2})^2} + \frac{\ln(1 + \tan(\frac{x}{2})^2)}{2} - \frac{\ln(2e^x \tan(\frac{x}{2})^2 - \tan(\frac{x}{2})^2 + 2e^x + 2 \tan(\frac{x}{2}) + 1)}{2}$	70

input `int(sin(x)/(2*exp(x)+cos(x)+sin(x)),x,method=_RETURNVERBOSE)`

output `1/2*x+1/2*I*x-1/2*ln(exp(2*I*x)+(2+2*I)*exp((1+I)*x)+I)`

3.293.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{\sin(x)}{2e^x + \cos(x) + \sin(x)} dx = \frac{1}{2}x - \frac{1}{2} \log(\cos(x) + 2e^x + \sin(x))$$

input `integrate(sin(x)/(2*exp(x)+cos(x)+sin(x)),x, algorithm="fricas")`output `1/2*x - 1/2*log(cos(x) + 2*e^x + sin(x))`**3.293.6 Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{\sin(x)}{2e^x + \cos(x) + \sin(x)} dx = \frac{x}{2} - \frac{\log(2e^x + \sin(x) + \cos(x))}{2}$$

input `integrate(sin(x)/(2*exp(x)+cos(x)+sin(x)),x)`output `x/2 - log(2*exp(x) + sin(x) + cos(x))/2`**3.293.7 Maxima [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 91 vs. $2(16) = 32$.

Time = 0.21 (sec) , antiderivative size = 91, normalized size of antiderivative = 4.79

$$\begin{aligned} \int \frac{\sin(x)}{2e^x + \cos(x) + \sin(x)} dx = & \frac{1}{2}x - \frac{1}{4} \log(8 \cos(x)^2 e^{(2x)} + 8 e^{(2x)} \sin(x)^2 \\ & + 4(\cos(x) e^x - e^x \sin(x)) \cos(2x) + \cos(2x)^2 \\ & + 4 \cos(x) e^x + 2(2 \cos(x) e^x + 2 e^x \sin(x) + 1) \sin(2x) \\ & + \sin(2x)^2 + 4 e^x \sin(x) + 1) \end{aligned}$$

input `integrate(sin(x)/(2*exp(x)+cos(x)+sin(x)),x, algorithm="maxima")`output `1/2*x - 1/4*log(8*cos(x)^2*e^(2*x) + 8*e^(2*x)*sin(x)^2 + 4*(cos(x)*e^x - e^x*sin(x))*cos(2*x) + cos(2*x)^2 + 4*cos(x)*e^x + 2*(2*cos(x)*e^x + 2*e^x*sin(x) + 1)*sin(2*x) + sin(2*x)^2 + 4*e^x*sin(x) + 1)`

3.293.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs. $2(16) = 32$.

Time = 0.30 (sec) , antiderivative size = 119, normalized size of antiderivative = 6.26

$$\int \frac{\sin(x)}{2e^x + \cos(x) + \sin(x)} dx = \frac{1}{2}x - \frac{1}{4} \log \left(\frac{2 \left(4e^{(2x)} \tan\left(\frac{1}{2}x\right)^4 - 4e^x \tan\left(\frac{1}{2}x\right)^4 + 8e^x \tan\left(\frac{1}{2}x\right)^3 + \tan\left(\frac{1}{2}x\right)^4 + 8e^{(2x)} \tan\left(\frac{1}{2}x\right)^2 - 4 \tan\left(\frac{1}{2}x\right) + 1 \right)}{\tan\left(\frac{1}{2}x\right)^4 + 2 \tan\left(\frac{1}{2}x\right)^2 + 1} \right)$$

input `integrate(sin(x)/(2*exp(x)+cos(x)+sin(x)),x, algorithm="giac")`

output `1/2*x - 1/4*log(2*(4*e^(2*x))*tan(1/2*x)^4 - 4*e^x*tan(1/2*x)^4 + 8*e^x*tan(1/2*x)^3 + tan(1/2*x)^4 + 8*e^(2*x)*tan(1/2*x)^2 - 4*tan(1/2*x)^3 + 8*e^x*tan(1/2*x) + 2*tan(1/2*x)^2 + 4*e^(2*x) + 4*e^x + 4*tan(1/2*x) + 1)/(tan(1/2*x)^4 + 2*tan(1/2*x)^2 + 1))`

3.293.9 Mupad [B] (verification not implemented)

Time = 17.62 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.16

$$\int \frac{\sin(x)}{2e^x + \cos(x) + \sin(x)} dx = \frac{x}{2} - \frac{\ln(2e^x + \sqrt{2} \cos(x - \frac{\pi}{4}))}{2}$$

input `int(sin(x)/(cos(x) + 2*exp(x) + sin(x)),x)`

output `x/2 - log(2*exp(x) + 2^(1/2)*cos(x - pi/4))/2`

3.294 $\int \log^{-\log(e\pi)}(x) \log\left(\frac{x}{\pi}\right) dx$

3.294.1 Optimal result	1466
3.294.2 Mathematica [F]	1466
3.294.3 Rubi [C] (verified)	1467
3.294.4 Maple [F]	1468
3.294.5 Fracas [B] (verification not implemented)	1469
3.294.6 Sympy [B] (verification not implemented)	1469
3.294.7 Maxima [F]	1469
3.294.8 Giac [F]	1470
3.294.9 Mupad [F(-1)]	1470

3.294.1 Optimal result

Integrand size = 16, antiderivative size = 9

$$\int \log^{-\log(e\pi)}(x) \log\left(\frac{x}{\pi}\right) dx = x \log^{-\log(\pi)}(x)$$

output `x/(ln(x)^ln(Pi))`

3.294.2 Mathematica [F]

$$\int \log^{-\log(e\pi)}(x) \log\left(\frac{x}{\pi}\right) dx = \int \log^{-\log(e\pi)}(x) \log\left(\frac{x}{\pi}\right) dx$$

input `Integrate[Log[x/Pi]/Log[x]^Log[E*Pi], x]`

output `Integrate[Log[x/Pi]/Log[x]^Log[E*Pi], x]`

3.294.3 Rubi [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3 in optimal.

Time = 0.37 (sec) , antiderivative size = 74, normalized size of antiderivative = 8.22, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {2808, 25, 2033, 3039, 7111}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \log^{-\log(e\pi)}(x) \log\left(\frac{x}{\pi}\right) dx \\
 & \quad \downarrow \text{2808} \\
 & (-\log(x))^{1+\log(\pi)} \log^{-\log(e\pi)}(x) \log\left(\frac{x}{\pi}\right) \Gamma(-\log(\pi), -\log(x)) - \\
 & \quad \int -\frac{\Gamma(-\log(\pi), -\log(x))(-\log(x))^{\log(\pi)} \log^{-\log(\pi)}(x)}{x} dx \\
 & \quad \downarrow \text{25} \\
 & \int \frac{\Gamma(-\log(\pi), -\log(x))(-\log(x))^{\log(\pi)} \log^{-\log(\pi)}(x)}{x} dx + \\
 & (-\log(x))^{1+\log(\pi)} \log\left(\frac{x}{\pi}\right) \log^{-\log(e\pi)}(x) \Gamma(-\log(\pi), -\log(x)) \\
 & \quad \downarrow \text{2033} \\
 & (-\log(x))^{\log(\pi)} \log^{-\log(\pi)}(x) \int \frac{\Gamma(-\log(\pi), -\log(x))}{x} dx + \\
 & (-\log(x))^{1+\log(\pi)} \log\left(\frac{x}{\pi}\right) \log^{-\log(e\pi)}(x) \Gamma(-\log(\pi), -\log(x)) \\
 & \quad \downarrow \text{3039} \\
 & (-\log(x))^{\log(\pi)} \log^{-\log(\pi)}(x) \int \Gamma(-\log(\pi), -\log(x)) d\log(x) + \\
 & (-\log(x))^{1+\log(\pi)} \log\left(\frac{x}{\pi}\right) \log^{-\log(e\pi)}(x) \Gamma(-\log(\pi), -\log(x)) \\
 & \quad \downarrow \text{7111} \\
 & (-\log(x))^{1+\log(\pi)} \log\left(\frac{x}{\pi}\right) \log^{-\log(e\pi)}(x) \Gamma(-\log(\pi), -\log(x)) + \\
 & (-\log(x))^{\log(\pi)} \log^{-\log(\pi)}(x) (\Gamma(1 - \log(\pi), -\log(x)) + \log(x) \Gamma(-\log(\pi), -\log(x)))
 \end{aligned}$$

input `Int [Log [x/Pi] /Log [x] ^Log [E*Pi] , x]`

```
output ((-Log[x])^Log[Pi]*(Gamma[1 - Log[Pi], -Log[x]] + Gamma[-Log[Pi], -Log[x]]
*Log[x]))/Log[x]^Log[Pi] + (Gamma[-Log[Pi], -Log[x]]*(-Log[x])^(1 + Log[Pi]
])*Log[x/Pi])/Log[x]^Log[E*Pi]
```

3.294.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 2033 Int[(Fx_)*((a_)*(v_)^(m_))*((b_)*(v_)^(n_), x_Symbol] := Simp[a^(m + n)
*((b*v)^n/(a*v)^n) Int[v^(m + n)*Fx, x], x] /; FreeQ[{a, b, m, n}, x] &&
!IntegerQ[m] && !IntegerQ[n] && IntegerQ[m + n]
```

```
rule 2808 Int[((a_) + Log[(c_)*(x_)^(n_)])*(b_)^(p_)*((d_) + Log[(f_)*(x_)^(r_
.)*e_]), x_Symbol] := With[{u = IntHide[(a + b*Log[c*x^n])^p, x]}, Simp[
(d + e*Log[f*x^r]) u, x] - Simp[e*r Int[SimplifyIntegrand[u/x, x], x],
x]] /; FreeQ[{a, b, c, d, e, f, n, p, r}, x]
```

```
rule 3039 Int[u_, x_Symbol] := With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst
[[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst] /;
NonsumQ[u]
```

```
rule 7111 Int[Gamma[n_, (a_) + (b_)*(x_)], x_Symbol] := Simp[(a + b*x)*(Gamma[n, a
+ b*x]/b), x] - Simp[Gamma[n + 1, a + b*x]/b, x] /; FreeQ[{a, b, n}, x]
```

3.294.4 Maple [F]

$$\int \ln\left(\frac{x}{\pi}\right) \ln(x)^{-\ln(\pi e)} dx$$

```
input int(ln(x/Pi)/(ln(x)^ln(Pi*exp(1))),x)
```

```
output int(ln(x/Pi)/(ln(x)^ln(Pi*exp(1))),x)
```

3.294.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 30 vs. $2(9) = 18$.

Time = 0.25 (sec) , antiderivative size = 30, normalized size of antiderivative = 3.33

$$\int \log^{-\log(e\pi)}(x) \log\left(\frac{x}{\pi}\right) dx = \frac{x \log(\pi) + x \log\left(\frac{x}{\pi}\right)}{(\log(\pi) + \log\left(\frac{x}{\pi}\right))^{\log(\pi)+1}}$$

input `integrate(log(x/pi)/(log(x)^log(pi*exp(1))),x, algorithm="fricas")`

output `(x*log(pi) + x*log(x/pi))/(log(pi) + log(x/pi))^(log(pi) + 1)`

3.294.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. $2(7) = 14$.

Time = 18.12 (sec) , antiderivative size = 56, normalized size of antiderivative = 6.22

$$\int \log^{-\log(e\pi)}(x) \log\left(\frac{x}{\pi}\right) dx = -\frac{(-\log(x))^{1+\log(\pi)} \log(\pi) \Gamma(-\log(\pi), -\log(x))}{\log(x)^{1+\log(\pi)}} + \frac{(-\log(x))^{\log(\pi)} \Gamma(1 - \log(\pi), -\log(x))}{\log(x)^{\log(\pi)}}$$

input `integrate(ln(x/pi)/(ln(x)**ln(pi*exp(1))),x)`

output `-(-log(x))**(1 + log(pi))*log(pi)*log(x)**(-log(pi) - 1)*uppergamma(-log(pi), -log(x)) + (-log(x))**log(pi)*uppergamma(1 - log(pi), -log(x))/log(x)*log(pi)`

3.294.7 Maxima [F]

$$\int \log^{-\log(e\pi)}(x) \log\left(\frac{x}{\pi}\right) dx = \int \frac{\log\left(\frac{x}{\pi}\right)}{\log(x)^{\log(\pi e)}} dx$$

input `integrate(log(x/pi)/(log(x)^log(pi*exp(1))),x, algorithm="maxima")`

output `integrate(log(x)^(-log(pi*e))*log(x/pi), x)`

3.294.8 Giac [F]

$$\int \log^{-\log(e\pi)}(x) \log\left(\frac{x}{\pi}\right) dx = \int \frac{\log\left(\frac{x}{\pi}\right)}{\log(x)^{\log(\pi e)}} dx$$

input `integrate(log(x/pi)/(log(x)^log(pi*exp(1))),x, algorithm="giac")`

output `integrate(log(x/pi)/log(x)^log(pi*e), x)`

3.294.9 Mupad [F(-1)]

Timed out.

$$\int \log^{-\log(e\pi)}(x) \log\left(\frac{x}{\pi}\right) dx = \int \frac{\ln\left(\frac{x}{\pi}\right)}{\ln(x)^{\ln(\pi e)}} dx$$

input `int(log(x/Pi)/log(x)^log(Pi*exp(1)),x)`

output `int(log(x/Pi)/log(x)^log(Pi*exp(1)), x)`

3.295 $\int (x + 8x^2 + 10x^3 + 5x^4 + x^5) dx$

3.295.1 Optimal result1471
3.295.2 Mathematica [A] (verified)1471
3.295.3 Rubi [A] (verified)1472
3.295.4 Maple [A] (verified)1472
3.295.5 Fricas [A] (verification not implemented)1473
3.295.6 Sympy [A] (verification not implemented)1473
3.295.7 Maxima [A] (verification not implemented)1473
3.295.8 Giac [A] (verification not implemented)1474
3.295.9 Mupad [B] (verification not implemented)1474

3.295.1 Optimal result

Integrand size = 20, antiderivative size = 32

$$\int (x + 8x^2 + 10x^3 + 5x^4 + x^5) dx = \frac{x^2}{2} + \frac{8x^3}{3} + \frac{5x^4}{2} + x^5 + \frac{x^6}{6}$$

output `1/2*x^2+8/3*x^3+5/2*x^4+x^5+1/6*x^6`

3.295.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int (x + 8x^2 + 10x^3 + 5x^4 + x^5) dx = \frac{x^2}{2} + \frac{8x^3}{3} + \frac{5x^4}{2} + x^5 + \frac{x^6}{6}$$

input `Integrate[x + 8*x^2 + 10*x^3 + 5*x^4 + x^5,x]`

output `x^2/2 + (8*x^3)/3 + (5*x^4)/2 + x^5 + x^6/6`

3.295.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x^5 + 5x^4 + 10x^3 + 8x^2 + x) dx$$

↓ 2009

$$\frac{x^6}{6} + x^5 + \frac{5x^4}{2} + \frac{8x^3}{3} + \frac{x^2}{2}$$

input `Int[x + 8*x^2 + 10*x^3 + 5*x^4 + x^5,x]`

output `x^2/2 + (8*x^3)/3 + (5*x^4)/2 + x^5 + x^6/6`

3.295.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.295.4 Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75

method	result	size
gospers	$\frac{x^2(x^4+6x^3+15x^2+16x+3)}{6}$	24
default	$\frac{1}{2}x^2 + \frac{8}{3}x^3 + \frac{5}{2}x^4 + x^5 + \frac{1}{6}x^6$	25
norman	$\frac{1}{2}x^2 + \frac{8}{3}x^3 + \frac{5}{2}x^4 + x^5 + \frac{1}{6}x^6$	25
risch	$\frac{1}{2}x^2 + \frac{8}{3}x^3 + \frac{5}{2}x^4 + x^5 + \frac{1}{6}x^6$	25
parallelrisch	$\frac{1}{2}x^2 + \frac{8}{3}x^3 + \frac{5}{2}x^4 + x^5 + \frac{1}{6}x^6$	25
parts	$\frac{1}{2}x^2 + \frac{8}{3}x^3 + \frac{5}{2}x^4 + x^5 + \frac{1}{6}x^6$	25

input `int(x^5+5*x^4+10*x^3+8*x^2+x,x,method=_RETURNVERBOSE)`

output `1/6*x^2*(x^4+6*x^3+15*x^2+16*x+3)`

3.295.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75

$$\int (x + 8x^2 + 10x^3 + 5x^4 + x^5) dx = \frac{1}{6}x^6 + x^5 + \frac{5}{2}x^4 + \frac{8}{3}x^3 + \frac{1}{2}x^2$$

input `integrate(x^5+5*x^4+10*x^3+8*x^2+x,x,algorithm="fricas")`

output `1/6*x^6 + x^5 + 5/2*x^4 + 8/3*x^3 + 1/2*x^2`

3.295.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

$$\int (x + 8x^2 + 10x^3 + 5x^4 + x^5) dx = \frac{x^6}{6} + x^5 + \frac{5x^4}{2} + \frac{8x^3}{3} + \frac{x^2}{2}$$

input `integrate(x**5+5*x**4+10*x**3+8*x**2+x,x)`

output `x**6/6 + x**5 + 5*x**4/2 + 8*x**3/3 + x**2/2`

3.295.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75

$$\int (x + 8x^2 + 10x^3 + 5x^4 + x^5) dx = \frac{1}{6}x^6 + x^5 + \frac{5}{2}x^4 + \frac{8}{3}x^3 + \frac{1}{2}x^2$$

input `integrate(x^5+5*x^4+10*x^3+8*x^2+x,x,algorithm="maxima")`

output `1/6*x^6 + x^5 + 5/2*x^4 + 8/3*x^3 + 1/2*x^2`

3.295.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75

$$\int (x + 8x^2 + 10x^3 + 5x^4 + x^5) dx = \frac{1}{6}x^6 + x^5 + \frac{5}{2}x^4 + \frac{8}{3}x^3 + \frac{1}{2}x^2$$

input `integrate(x^5+5*x^4+10*x^3+8*x^2+x,x, algorithm="giac")`output `1/6*x^6 + x^5 + 5/2*x^4 + 8/3*x^3 + 1/2*x^2`**3.295.9 Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75

$$\int (x + 8x^2 + 10x^3 + 5x^4 + x^5) dx = \frac{x^6}{6} + x^5 + \frac{5x^4}{2} + \frac{8x^3}{3} + \frac{x^2}{2}$$

input `int(x + 8*x^2 + 10*x^3 + 5*x^4 + x^5,x)`output `x^2/2 + (8*x^3)/3 + (5*x^4)/2 + x^5 + x^6/6`

$$3.296 \quad \int \frac{(1+x)(4+x)}{(2+x)(3+x)} dx$$

3.296.1 Optimal result	1475
3.296.2 Mathematica [A] (verified)	1475
3.296.3 Rubi [A] (verified)	1476
3.296.4 Maple [A] (verified)	1477
3.296.5 Fricas [A] (verification not implemented)	1477
3.296.6 Sympy [A] (verification not implemented)	1477
3.296.7 Maxima [A] (verification not implemented)	1478
3.296.8 Giac [A] (verification not implemented)	1478
3.296.9 Mupad [B] (verification not implemented)	1478

3.296.1 Optimal result

Integrand size = 17, antiderivative size = 14

$$\int \frac{(1+x)(4+x)}{(2+x)(3+x)} dx = x - 2 \log(2+x) + 2 \log(3+x)$$

output `x-2*ln(2+x)+2*ln(3+x)`

3.296.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{(1+x)(4+x)}{(2+x)(3+x)} dx = x - 2 \log(2+x) + 2 \log(3+x)$$

input `Integrate[((1 + x)*(4 + x))/((2 + x)*(3 + x)),x]`

output `x - 2*Log[2 + x] + 2*Log[3 + x]`

3.296.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {159, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(x+1)(x+4)}{(x+2)(x+3)} dx$$

↓ 159

$$\int \left(\frac{2}{x+3} - \frac{2}{x+2} + 1 \right) dx$$

↓ 2009

$$x - 2 \log(x+2) + 2 \log(x+3)$$

input `Int[((1 + x)*(4 + x))/((2 + x)*(3 + x)),x]`

output `x - 2*Log[2 + x] + 2*Log[3 + x]`

3.296.3.1 Defintions of rubi rules used

rule 159 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))*(g_.) + (h_.)*(x_)), x_] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)*(g + h*x), x], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x] && (IGtQ[m, 0] || IntegersQ[m, n])`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.296.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

method	result	size
default	$x - 2 \ln(2 + x) + 2 \ln(3 + x)$	15
norman	$x - 2 \ln(2 + x) + 2 \ln(3 + x)$	15
risch	$x - 2 \ln(2 + x) + 2 \ln(3 + x)$	15
parallelrisc	$x - 2 \ln(2 + x) + 2 \ln(3 + x)$	15

input `int((1+x)/(2+x)/(3+x)*(4+x),x,method=_RETURNVERBOSE)`output `x-2*ln(2+x)+2*ln(3+x)`**3.296.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{(1+x)(4+x)}{(2+x)(3+x)} dx = x + 2 \log(x+3) - 2 \log(x+2)$$

input `integrate((1+x)/(2+x)/(3+x)*(4+x),x, algorithm="fricas")`output `x + 2*log(x + 3) - 2*log(x + 2)`**3.296.6 Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{(1+x)(4+x)}{(2+x)(3+x)} dx = x - 2 \log(x+2) + 2 \log(x+3)$$

input `integrate((1+x)/(2+x)/(3+x)*(4+x),x)`output `x - 2*log(x + 2) + 2*log(x + 3)`

3.296.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{(1+x)(4+x)}{(2+x)(3+x)} dx = x + 2 \log(x+3) - 2 \log(x+2)$$

input `integrate((1+x)/(2+x)/(3+x)*(4+x),x, algorithm="maxima")`output `x + 2*log(x + 3) - 2*log(x + 2)`**3.296.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{(1+x)(4+x)}{(2+x)(3+x)} dx = x + 2 \log(|x+3|) - 2 \log(|x+2|)$$

input `integrate((1+x)/(2+x)/(3+x)*(4+x),x, algorithm="giac")`output `x + 2*log(abs(x + 3)) - 2*log(abs(x + 2))`**3.296.9 Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{(1+x)(4+x)}{(2+x)(3+x)} dx = x + 4 \operatorname{atanh}(2x+5)$$

input `int(((x + 1)*(x + 4))/((x + 2)*(x + 3)),x)`output `x + 4*atanh(2*x + 5)`

3.297 $\int x \cot(x) dx$

3.297.1 Optimal result	1479
3.297.2 Mathematica [A] (verified)	1479
3.297.3 Rubi [A] (verified)	1480
3.297.4 Maple [A] (verified)	1482
3.297.5 Fricas [B] (verification not implemented)	1482
3.297.6 Sympy [F]	1482
3.297.7 Maxima [B] (verification not implemented)	1483
3.297.8 Giac [F]	1483
3.297.9 Mupad [F(-1)]	1483

3.297.1 Optimal result

Integrand size = 4, antiderivative size = 39

$$\int x \cot(x) dx = -\frac{ix^2}{2} + x \log(1 - e^{2ix}) - \frac{1}{2}i \operatorname{PolyLog}(2, e^{2ix})$$

output `-1/2*I*x^2+x*ln(1-exp(2*I*x))-1/2*I*polylog(2,exp(2*I*x))`

3.297.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.87

$$\int x \cot(x) dx = x \log(1 - e^{2ix}) - \frac{1}{2}i(x^2 + \operatorname{PolyLog}(2, e^{2ix}))$$

input `Integrate[x*Cot[x],x]`

output `x*Log[1 - E^((2*I)*x)] - (I/2)*(x^2 + PolyLog[2, E^((2*I)*x)])`

3.297.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.21, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.750$, Rules used = {3042, 25, 4200, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \cot(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -x \tan\left(x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{25} \\
 & - \int x \tan\left(x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{4200} \\
 & 2i \int -\frac{e^{2ix} x}{1 - e^{2ix}} dx - \frac{ix^2}{2} \\
 & \quad \downarrow \text{25} \\
 & -2i \int \frac{e^{2ix} x}{1 - e^{2ix}} dx - \frac{ix^2}{2} \\
 & \quad \downarrow \text{2620} \\
 & -2i \left(\frac{1}{2} ix \log(1 - e^{2ix}) - \frac{1}{2} i \int \log(1 - e^{2ix}) dx \right) - \frac{ix^2}{2} \\
 & \quad \downarrow \text{2715} \\
 & -2i \left(\frac{1}{2} ix \log(1 - e^{2ix}) - \frac{1}{4} \int e^{-2ix} \log(1 - e^{2ix}) de^{2ix} \right) - \frac{ix^2}{2} \\
 & \quad \downarrow \text{2838} \\
 & -2i \left(\frac{1}{4} \text{PolyLog}(2, e^{2ix}) + \frac{1}{2} ix \log(1 - e^{2ix}) \right) - \frac{ix^2}{2}
 \end{aligned}$$

input `Int [x*Cot [x] , x]`

output $(-1/2*I)*x^2 - (2*I)*((I/2)*x*\text{Log}[1 - E^{((2*I)*x)}] + \text{PolyLog}[2, E^{((2*I)*x)}])/4$

3.297.3.1 Defintions of rubi rules used

rule 25 $\text{Int}[-(F x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F x, x], x]$

rule 2620 $\text{Int}[\frac{((F_)^{((g_)*(e_) + (f_)*(x_)))^{(n_)*((c_) + (d_)*(x_))^{(m_))}}}{((a_) + (b_)*((F_)^{((g_)*(e_) + (f_)*(x_)))^{(n_))}}), x_Symbol] \rightarrow \text{Simp}[\frac{((c + d*x)^m/(b*f*g*n*\text{Log}[F]))*\text{Log}[1 + b*((F^{(g*(e + f*x)))^n/a})]}{((c + d*x)^m/(b*f*g*n*\text{Log}[F]))} \quad \text{Int}[(c + d*x)^{(m-1)}*\text{Log}[1 + b*((F^{(g*(e + f*x)))^n/a})], x], x] /; \text{FreeQ}\{F, a, b, c, d, e, f, g, n\}, x\} \&\& \text{IGtQ}[m, 0]$

rule 2715 $\text{Int}[\text{Log}[(a_) + (b_)*((F_)^{((e_)*(c_) + (d_)*(x_))})^{(n_)}], x_Symbol] \rightarrow \text{Simp}[1/(d*e*n*\text{Log}[F]) \quad \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^{(e*(c + d*x)})^n)], x] /; \text{FreeQ}\{F, a, b, c, d, e, n\}, x\} \&\& \text{GtQ}[a, 0]$

rule 2838 $\text{Int}[\text{Log}[(c_)*(d_) + (e_)*(x_)^{(n_)}]/(x_), x_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n]/n, x] /; \text{FreeQ}\{c, d, e, n\}, x\} \&\& \text{EqQ}[c*d, 1]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4200 $\text{Int}[\frac{((c_) + (d_)*(x_))^{(m_)*\tan[(e_) + \text{Pi}*(k_) + (f_)*(x_)]}{(c + d*x)^{(m+1)}/(d*(m+1))}, x_Symbol] \rightarrow \text{Simp}[I*((c + d*x)^{(m+1)}/(d*(m+1))), x] - \text{Simp}[2*I \quad \text{Int}[(c + d*x)^m * E^{(2*I*k*Pi)} * (E^{(2*I*(e + f*x)})/(1 + E^{(2*I*k*Pi)} * E^{(2*I*(e + f*x)}))], x], x] /; \text{FreeQ}\{c, d, e, f\}, x\} \&\& \text{IntegerQ}[4*k] \&\& \text{IGtQ}[m, 0]$

3.297.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.72

method	result	size
parts	$x \ln(1 - e^{2ix}) - \frac{i(x^2 + \text{polylog}(2, e^{2ix}))}{2}$	28
risch	$-\frac{ix^2}{2} + x \ln(e^{ix} + 1) - i \text{polylog}(2, -e^{ix}) + x \ln(1 - e^{ix}) - i \text{polylog}(2, e^{ix})$	52

input `int(x*cot(x),x,method=_RETURNVERBOSE)`output `x*ln(1-exp(2*I*x))-1/2*I*(x^2+polylog(2,exp(2*I*x)))`**3.297.5 Fracas [B] (verification not implemented)**Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 65 vs. $2(24) = 48$.

Time = 0.25 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.67

$$\int x \cot(x) dx = \frac{1}{2} x \log(-\cos(2x) + i \sin(2x) + 1) + \frac{1}{2} x \log(-\cos(2x) - i \sin(2x) + 1) \\ - \frac{1}{4} i \text{Li}_2(\cos(2x) + i \sin(2x)) + \frac{1}{4} i \text{Li}_2(\cos(2x) - i \sin(2x))$$

input `integrate(x*cot(x),x, algorithm="fracas")`output `1/2*x*log(-cos(2*x) + I*sin(2*x) + 1) + 1/2*x*log(-cos(2*x) - I*sin(2*x) + 1) - 1/4*I*dilog(cos(2*x) + I*sin(2*x)) + 1/4*I*dilog(cos(2*x) - I*sin(2*x))`**3.297.6 Sympy [F]**

$$\int x \cot(x) dx = \int x \cot(x) dx$$

input `integrate(x*cot(x),x)`output `Integral(x*cot(x), x)`

3.297.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 80 vs. $2(24) = 48$.

Time = 0.20 (sec) , antiderivative size = 80, normalized size of antiderivative = 2.05

$$\int x \cot(x) dx = -\frac{1}{2}i x^2 + i x \arctan(\sin(x), \cos(x) + 1) - i x \arctan(\sin(x), -\cos(x) + 1) \\ + \frac{1}{2} x \log(\cos(x)^2 + \sin(x)^2 + 2 \cos(x) + 1) \\ + \frac{1}{2} x \log(\cos(x)^2 + \sin(x)^2 - 2 \cos(x) + 1) - i \operatorname{Li}_2(-e^{ix}) - i \operatorname{Li}_2(e^{ix})$$

input `integrate(x*cot(x),x, algorithm="maxima")`

output `-1/2*I*x^2 + I*x*arctan2(sin(x), cos(x) + 1) - I*x*arctan2(sin(x), -cos(x) + 1) + 1/2*x*log(cos(x)^2 + sin(x)^2 + 2*cos(x) + 1) + 1/2*x*log(cos(x)^2 + sin(x)^2 - 2*cos(x) + 1) - I*dilog(-e^(I*x)) - I*dilog(e^(I*x))`

3.297.8 Giac [F]

$$\int x \cot(x) dx = \int x \cot(x) dx$$

input `integrate(x*cot(x),x, algorithm="giac")`

output `integrate(x*cot(x), x)`

3.297.9 Mupad [F(-1)]

Timed out.

$$\int x \cot(x) dx = \int x \cot(x) dx$$

input `int(x*cot(x),x)`

output `int(x*cot(x), x)`

3.298
$$\int \frac{e^{\frac{1}{x}+x}(-1-x^2+x^4+x^6)}{x^4} dx$$

3.298.1 Optimal result	1484
3.298.2 Mathematica [A] (verified)	1484
3.298.3 Rubi [F]	1485
3.298.4 Maple [A] (verified)	1486
3.298.5 Fricas [A] (verification not implemented)	1486
3.298.6 Sympy [A] (verification not implemented)	1486
3.298.7 Maxima [A] (verification not implemented)	1487
3.298.8 Giac [B] (verification not implemented)	1487
3.298.9 Mupad [B] (verification not implemented)	1487

3.298.1 Optimal result

Integrand size = 24, antiderivative size = 24

$$\int \frac{e^{\frac{1}{x}+x}(-1-x^2+x^4+x^6)}{x^4} dx = e^{\frac{1}{x}+x} \left(4 + \frac{1}{x^2} - \frac{2}{x} - 2x + x^2 \right)$$

output `exp(x+1/x)*(4+1/x^2-2/x-2*x+x^2)`

3.298.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{e^{\frac{1}{x}+x}(-1-x^2+x^4+x^6)}{x^4} dx = e^{\frac{1}{x}+x} \left(4 + \frac{1}{x^2} - \frac{2}{x} - 2x + x^2 \right)$$

input `Integrate[(E^(x^(-1) + x))*(-1 - x^2 + x^4 + x^6))/x^4,x]`

output `E^(x^(-1) + x)*(4 + x^(-2) - 2/x - 2*x + x^2)`

3.298.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{x+\frac{1}{x}}(x^6 + x^4 - x^2 - 1)}{x^4} dx$$

↓ 7292

$$\int \frac{e^{x+\frac{1}{x}}(x^2 - 1)(x^2 + 1)^2}{x^4} dx$$

↓ 7293

$$\int \left(-\frac{e^{x+\frac{1}{x}}}{x^4} + e^{x+\frac{1}{x}}x^2 - \frac{e^{x+\frac{1}{x}}}{x^2} + e^{x+\frac{1}{x}} \right) dx$$

↓ 2009

$$-\int \frac{e^{x+\frac{1}{x}}}{x^4} dx - \int \frac{e^{x+\frac{1}{x}}}{x^2} dx + \int e^{x+\frac{1}{x}}x^2 dx + \int e^{x+\frac{1}{x}} dx$$

input `Int[(E^(x^(-1) + x)*(-1 - x^2 + x^4 + x^6))/x^4,x]`

output `$Aborted`

3.298.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7292 `Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v != u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

3.298.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.38

method	result	size
gospers	$\frac{e^{\frac{x^2+1}{x}}(x^4-2x^3+4x^2-2x+1)}{x^2}$	33
risch	$\frac{e^{\frac{x^2+1}{x}}(x^4-2x^3+4x^2-2x+1)}{x^2}$	33
norman	$\frac{x e^{x+\frac{1}{x}}+x^5 e^{x+\frac{1}{x}}+4x^3 e^{x+\frac{1}{x}}-2x^4 e^{x+\frac{1}{x}}-2 e^{x+\frac{1}{x}} x^2}{x^3}$	57
parallelrisch	$\frac{e^{\frac{x^2+1}{x}} x^4 - 2 e^{\frac{x^2+1}{x}} x^3 + 4 e^{\frac{x^2+1}{x}} x^2 - 2 e^{\frac{x^2+1}{x}} x + e^{\frac{x^2+1}{x}}}{x^2}$	73

input `int(exp(x+1/x)*(x^6+x^4-x^2-1)/x^4,x,method=_RETURNVERBOSE)`output `exp((x^2+1)/x)*(x^4-2*x^3+4*x^2-2*x+1)/x^2`**3.298.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.33

$$\int \frac{e^{\frac{1}{x}+x}(-1-x^2+x^4+x^6)}{x^4} dx = \frac{(x^4-2x^3+4x^2-2x+1)e^{\left(\frac{x^2+1}{x}\right)}}{x^2}$$

input `integrate(exp(x+1/x)*(x^6+x^4-x^2-1)/x^4,x, algorithm="fricas")`output `(x^4 - 2*x^3 + 4*x^2 - 2*x + 1)*e^((x^2 + 1)/x)/x^2`**3.298.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.12

$$\int \frac{e^{\frac{1}{x}+x}(-1-x^2+x^4+x^6)}{x^4} dx = \frac{(x^4-2x^3+4x^2-2x+1)e^{x+\frac{1}{x}}}{x^2}$$

input `integrate(exp(x+1/x)*(x**6+x**4-x**2-1)/x**4,x)`output `(x**4 - 2*x**3 + 4*x**2 - 2*x + 1)*exp(x + 1/x)/x**2`

3.298. $\int \frac{e^{\frac{1}{x}+x}(-1-x^2+x^4+x^6)}{x^4} dx$

3.298.7 Maxima [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17

$$\int \frac{e^{\frac{1}{x}+x}(-1-x^2+x^4+x^6)}{x^4} dx = \frac{(x^4-2x^3+4x^2-2x+1)e^{(x+\frac{1}{x})}}{x^2}$$

input `integrate(exp(x+1/x)*(x^6+x^4-x^2-1)/x^4,x, algorithm="maxima")`

output `(x^4 - 2*x^3 + 4*x^2 - 2*x + 1)*e^(x + 1/x)/x^2`

3.298.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 72 vs. 2(23) = 46.

Time = 0.28 (sec) , antiderivative size = 72, normalized size of antiderivative = 3.00

$$\begin{aligned} \int \frac{e^{\frac{1}{x}+x}(-1-x^2+x^4+x^6)}{x^4} dx \\ = \frac{x^4 e^{\left(\frac{x^2+1}{x}\right)} - 2x^3 e^{\left(\frac{x^2+1}{x}\right)} + 4x^2 e^{\left(\frac{x^2+1}{x}\right)} - 2x e^{\left(\frac{x^2+1}{x}\right)} + e^{\left(\frac{x^2+1}{x}\right)}}{x^2} \end{aligned}$$

input `integrate(exp(x+1/x)*(x^6+x^4-x^2-1)/x^4,x, algorithm="giac")`

output `(x^4*e^((x^2 + 1)/x) - 2*x^3*e^((x^2 + 1)/x) + 4*x^2*e^((x^2 + 1)/x) - 2*x*e^((x^2 + 1)/x) + e^((x^2 + 1)/x))/x^2`

3.298.9 Mupad [B] (verification not implemented)

Time = 18.19 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17

$$\int \frac{e^{\frac{1}{x}+x}(-1-x^2+x^4+x^6)}{x^4} dx = \frac{e^{x+\frac{1}{x}}(x^4-2x^3+4x^2-2x+1)}{x^2}$$

input `int(-(exp(x + 1/x)*(x^2 - x^4 - x^6 + 1))/x^4,x)`

output `(exp(x + 1/x)*(4*x^2 - 2*x - 2*x^3 + x^4 + 1))/x^2`

3.298. $\int \frac{e^{\frac{1}{x}+x}(-1-x^2+x^4+x^6)}{x^4} dx$

3.299 $\int \frac{1}{\sqrt{(-1+x)(1+x)^3}} dx$

3.299.1 Optimal result 1488
 3.299.2 Mathematica [A] (verified) 1488
 3.299.3 Rubi [A] (verified) 1489
 3.299.4 Maple [A] (verified) 1490
 3.299.5 Fricas [A] (verification not implemented) 1490
 3.299.6 Sympy [F] 1490
 3.299.7 Maxima [F] 1491
 3.299.8 Giac [A] (verification not implemented) 1491
 3.299.9 Mupad [B] (verification not implemented) 1491

3.299.1 Optimal result

Integrand size = 13, antiderivative size = 26

$$\int \frac{1}{\sqrt{(-1+x)(1+x)^3}} dx = -\frac{(1-x)(1+x)}{\sqrt{-((1-x)(1+x)^3)}}$$

output `-(1-x)*(1+x)/(-(1-x)*(1+x)^3)^(1/2)`

3.299.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \frac{1}{\sqrt{(-1+x)(1+x)^3}} dx = \frac{(-1+x)(1+x)}{\sqrt{(-1+x)(1+x)^3}}$$

input `Integrate[1/Sqrt[(-1 + x)*(1 + x)^3], x]`

output `((-1 + x)*(1 + x))/Sqrt[(-1 + x)*(1 + x)^3]`

3.299.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.88, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {7270, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{(x-1)(x+1)^3}} dx$$

↓ 7270

$$\frac{\sqrt{x-1}(x+1)^{3/2} \int \frac{1}{\sqrt{x-1}(x+1)^{3/2}} dx}{\sqrt{-((1-x)(x+1)^3)}}$$

↓ 48

$$\frac{(x-1)(x+1)}{\sqrt{-((1-x)(x+1)^3)}}$$

input `Int[1/Sqrt[(-1 + x)*(1 + x)^3],x]`

output `((-1 + x)*(1 + x))/Sqrt[-((1 - x)*(1 + x)^3)]`

3.299.3.1 Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 7270 `Int[(u_.)*((a_.)*(v_)^(m_.)*(w_)^(n_.))^(p_), x_Symbol] := Simp[a^IntPart[p] * ((a*v^m*w^n)^FracPart[p]/(v^(m*FracPart[p])*w^(n*FracPart[p])))] Int[u*v^(m*p)*w^(n*p), x], x] /; FreeQ[{a, m, n, p}, x] && !IntegerQ[p] && !FreeQ[v, x] && !FreeQ[w, x]`

3.299.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.73

method	result	size
gospers	$\frac{(-1+x)(1+x)}{\sqrt{(1+x)^3(-1+x)}}$	19
risch	$\frac{(-1+x)(1+x)}{\sqrt{(1+x)^3(-1+x)}}$	19
trager	$\frac{\sqrt{x^4+2x^3-2x-1}}{(1+x)^2}$	22
default	$\frac{\sqrt{(-1+x)(1+x)}\sqrt{x^2-1}}{\sqrt{(1+x)^3(-1+x)}}$	29

input `int(1/((1+x)^3*(-1+x))^(1/2),x,method=_RETURNVERBOSE)`output `(-1+x)*(1+x)/((1+x)^3*(-1+x))^(1/2)`**3.299.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.31

$$\int \frac{1}{\sqrt{(-1+x)(1+x)^3}} dx = \frac{x^2 + 2x + \sqrt{x^4 + 2x^3 - 2x - 1} + 1}{x^2 + 2x + 1}$$

input `integrate(1/((1+x)^3*(-1+x))^(1/2),x, algorithm="fracas")`output `(x^2 + 2*x + sqrt(x^4 + 2*x^3 - 2*x - 1) + 1)/(x^2 + 2*x + 1)`**3.299.6 Sympy [F]**

$$\int \frac{1}{\sqrt{(-1+x)(1+x)^3}} dx = \int \frac{1}{\sqrt{(x-1)(x+1)^3}} dx$$

input `integrate(1/((1+x)**3*(-1+x))^(1/2),x)`output `Integral(1/sqrt((x - 1)*(x + 1)**3), x)`

3.299.7 Maxima [F]

$$\int \frac{1}{\sqrt{(-1+x)(1+x)^3}} dx = \int \frac{1}{\sqrt{(x+1)^3(x-1)}} dx$$

input `integrate(1/((1+x)^3*(-1+x))^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt((x + 1)^3*(x - 1)), x)`

3.299.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{1}{\sqrt{(-1+x)(1+x)^3}} dx = \frac{2}{(x - \sqrt{x^2 - 1} + 1)\operatorname{sgn}(x + 1)}$$

input `integrate(1/((1+x)^3*(-1+x))^(1/2),x, algorithm="giac")`

output `2/((x - sqrt(x^2 - 1) + 1)*sgn(x + 1))`

3.299.9 Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.65

$$\int \frac{1}{\sqrt{(-1+x)(1+x)^3}} dx = \frac{x^2 - 1}{\sqrt{(x - 1)(x + 1)^3}}$$

input `int(1/((x - 1)*(x + 1)^3)^(1/2),x)`

output `(x^2 - 1)/((x - 1)*(x + 1)^3)^(1/2)`

3.300 $\int x \sin^4(x) dx$

3.300.1 Optimal result	1492
3.300.2 Mathematica [A] (verified)	1492
3.300.3 Rubi [A] (verified)	1493
3.300.4 Maple [A] (verified)	1494
3.300.5 Fricas [A] (verification not implemented)	1495
3.300.6 Sympy [A] (verification not implemented)	1495
3.300.7 Maxima [A] (verification not implemented)	1495
3.300.8 Giac [A] (verification not implemented)	1496
3.300.9 Mupad [B] (verification not implemented)	1496

3.300.1 Optimal result

Integrand size = 6, antiderivative size = 44

$$\int x \sin^4(x) dx = \frac{3x^2}{16} - \frac{3}{8}x \cos(x) \sin(x) + \frac{3 \sin^2(x)}{16} - \frac{1}{4}x \cos(x) \sin^3(x) + \frac{\sin^4(x)}{16}$$

output `3/16*x^2-3/8*x*cos(x)*sin(x)+3/16*sin(x)^2-1/4*x*cos(x)*sin(x)^3+1/16*sin(x)^4`

3.300.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.95

$$\int x \sin^4(x) dx = \frac{3x^2}{16} - \frac{1}{8} \cos(2x) + \frac{1}{128} \cos(4x) - \frac{1}{4}x \sin(2x) + \frac{1}{32}x \sin(4x)$$

input `Integrate[x*Sin[x]^4,x]`

output `(3*x^2)/16 - Cos[2*x]/8 + Cos[4*x]/128 - (x*Sin[2*x])/4 + (x*Sin[4*x])/32`

3.300.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.11, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {3042, 3791, 3042, 3791, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \sin^4(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int x \sin(x)^4 dx \\
 & \quad \downarrow \text{3791} \\
 & \frac{3}{4} \int x \sin^2(x) dx + \frac{\sin^4(x)}{16} - \frac{1}{4} x \sin^3(x) \cos(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{4} \int x \sin(x)^2 dx + \frac{\sin^4(x)}{16} - \frac{1}{4} x \sin^3(x) \cos(x) \\
 & \quad \downarrow \text{3791} \\
 & \frac{3}{4} \left(\frac{\int x dx}{2} + \frac{\sin^2(x)}{4} - \frac{1}{2} x \sin(x) \cos(x) \right) + \frac{\sin^4(x)}{16} - \frac{1}{4} x \sin^3(x) \cos(x) \\
 & \quad \downarrow \text{15} \\
 & \frac{3}{4} \left(\frac{x^2}{4} + \frac{\sin^2(x)}{4} - \frac{1}{2} x \sin(x) \cos(x) \right) + \frac{\sin^4(x)}{16} - \frac{1}{4} x \sin^3(x) \cos(x)
 \end{aligned}$$

input `Int[x*Sin[x]^4,x]`

output `-1/4*(x*Cos[x]*Sin[x]^3) + Sin[x]^4/16 + (3*(x^2/4 - (x*Cos[x]*Sin[x])/2 + Sin[x]^2/4))/4`

3.300.3.1 Defintions of rubi rules used

```
rule 15 Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[
{a, m}, x] && NeQ[m, -1]

rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

rule 3791 Int[((c_.) + (d_.)*(x_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :=
Simp[d*((b*Sin[e + f*x])^n/(f^n)), x] + (-Simp[b*(c + d*x)*Cos[e + f*x]
]*((b*Sin[e + f*x])^(n - 1)/(f^n)), x] + Simp[b^2*((n - 1)/n) Int[(c + d*
x)*(b*Sin[e + f*x])^(n - 2), x], x] /; FreeQ[{b, c, d, e, f}, x] && GtQ[n,
1]
```

3.300.4 Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.75

method	result
risch	$\frac{3x^2}{16} + \frac{\cos(4x)}{128} + \frac{x \sin(4x)}{32} - \frac{\cos(2x)}{8} - \frac{x \sin(2x)}{4}$
default	$x \left(-\frac{(\sin(x)^3 + \frac{3 \sin(x)}{2}) \cos(x)}{4} + \frac{3x}{8} \right) - \frac{3x^2}{16} + \frac{(2 \cos(x)^2 - 5)^2}{64}$
norman	$\frac{\frac{3 \tan(\frac{x}{2})^2}{4} + \frac{3 \tan(\frac{x}{2})^6}{4} + \frac{5 \tan(\frac{x}{2})^4}{2} + \frac{3x^2}{16} - \frac{3x \tan(\frac{x}{2})}{4} - \frac{11x \tan(\frac{x}{2})^3}{4} + \frac{11x \tan(\frac{x}{2})^5}{4} + \frac{3x \tan(\frac{x}{2})^7}{4} + \frac{3x^2 \tan(\frac{x}{2})^2}{4} + \frac{9x^2 \tan(\frac{x}{2})^4}{8} + \frac{3x^2 \tan(\frac{x}{2})^6}{4}}{(1 + \tan(\frac{x}{2})^2)^4}$

```
input int(x*sin(x)^4,x,method=_RETURNVERBOSE)

output 3/16*x^2+1/128*cos(4*x)+1/32*x*sin(4*x)-1/8*cos(2*x)-1/4*x*sin(2*x)
```

3.300.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.80

$$\int x \sin^4(x) dx = \frac{1}{16} \cos(x)^4 + \frac{3}{16} x^2 - \frac{5}{16} \cos(x)^2 + \frac{1}{8} (2x \cos(x)^3 - 5x \cos(x)) \sin(x)$$

input `integrate(x*sin(x)^4,x, algorithm="fricas")`output `1/16*cos(x)^4 + 3/16*x^2 - 5/16*cos(x)^2 + 1/8*(2*x*cos(x)^3 - 5*x*cos(x))
*sin(x)`**3.300.6 Sympy [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.89

$$\int x \sin^4(x) dx = \frac{3x^2 \sin^4(x)}{16} + \frac{3x^2 \sin^2(x) \cos^2(x)}{8} + \frac{3x^2 \cos^4(x)}{16} - \frac{5x \sin^3(x) \cos(x)}{8} - \frac{3x \sin(x) \cos^3(x)}{8} + \frac{5 \sin^4(x)}{32} - \frac{3 \cos^4(x)}{32}$$

input `integrate(x*sin(x)**4,x)`output `3*x**2*sin(x)**4/16 + 3*x**2*sin(x)**2*cos(x)**2/8 + 3*x**2*cos(x)**4/16 -
5*x*sin(x)**3*cos(x)/8 - 3*x*sin(x)*cos(x)**3/8 + 5*sin(x)**4/32 - 3*cos(x)
4/32`3.300.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.73

$$\int x \sin^4(x) dx = \frac{3}{16} x^2 + \frac{1}{32} x \sin(4x) - \frac{1}{4} x \sin(2x) + \frac{1}{128} \cos(4x) - \frac{1}{8} \cos(2x)$$

input `integrate(x*sin(x)^4,x, algorithm="maxima")`output `3/16*x^2 + 1/32*x*sin(4*x) - 1/4*x*sin(2*x) + 1/128*cos(4*x) - 1/8*cos(2*x)
)`

3.300.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.73

$$\int x \sin^4(x) dx = \frac{3}{16} x^2 + \frac{1}{32} x \sin(4x) - \frac{1}{4} x \sin(2x) + \frac{1}{128} \cos(4x) - \frac{1}{8} \cos(2x)$$

input `integrate(x*sin(x)^4,x, algorithm="giac")`output `3/16*x^2 + 1/32*x*sin(4*x) - 1/4*x*sin(2*x) + 1/128*cos(4*x) - 1/8*cos(2*x)`
`)`**3.300.9 Mupad [B] (verification not implemented)**

Time = 16.93 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.86

$$\int x \sin^4(x) dx = \frac{\cos(2x)^2}{64} - \frac{x \sin(2x)}{4} - \frac{\cos(2x)}{8} + \frac{3x^2}{16} + \frac{x \cos(2x) \sin(2x)}{16}$$

input `int(x*sin(x)^4,x)`output `cos(2*x)^2/64 - (x*sin(2*x))/4 - cos(2*x)/8 + (3*x^2)/16 + (x*cos(2*x)*sin(2*x))/16`

3.301 $\int \cos(3x) \csc^2(x) \sec^3(x) \sin(2x) dx$

3.301.1 Optimal result	1497
3.301.2 Mathematica [A] (verified)	1497
3.301.3 Rubi [A] (verified)	1498
3.301.4 Maple [A] (verified)	1499
3.301.5 Fricas [A] (verification not implemented)	1500
3.301.6 Sympy [A] (verification not implemented)	1500
3.301.7 Maxima [B] (verification not implemented)	1500
3.301.8 Giac [A] (verification not implemented)	1501
3.301.9 Mupad [B] (verification not implemented)	1501

3.301.1 Optimal result

Integrand size = 17, antiderivative size = 11

$$\int \cos(3x) \csc^2(x) \sec^3(x) \sin(2x) dx = 6 \log(\cos(x)) + 2 \log(\sin(x))$$

output `6*ln(cos(x))+2*ln(sin(x))`

3.301.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \cos(3x) \csc^2(x) \sec^3(x) \sin(2x) dx = 8 \log(\cos(x)) + 2 \log(\tan(x))$$

input `Integrate[Cos[3*x]*Csc[x]^2*Sec[x]^3*Sin[2*x],x]`

output `8*Log[Cos[x]] + 2*Log[Tan[x]]`

3.301.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.55, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3042, 4865, 27, 354, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(2x) \cos(3x) \csc^2(x) \sec^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin(2x) \cos(3x) \csc(x)^2 \sec(x)^3 dx \\
 & \quad \downarrow \text{4865} \\
 & \int \frac{2(1 - 4 \sin^2(x)) \csc(x)}{1 - \sin^2(x)} d \sin(x) \\
 & \quad \downarrow \text{27} \\
 & 2 \int \frac{\csc(x) (1 - 4 \sin^2(x))}{1 - \sin^2(x)} d \sin(x) \\
 & \quad \downarrow \text{354} \\
 & \int \frac{(1 - 4 \sin^2(x)) \csc(x)}{1 - \sin^2(x)} d \sin^2(x) \\
 & \quad \downarrow \text{86} \\
 & \int \left(\frac{3}{\sin^2(x) - 1} + \csc(x) \right) d \sin^2(x) \\
 & \quad \downarrow \text{2009} \\
 & \log(\sin^2(x)) + 3 \log(1 - \sin^2(x))
 \end{aligned}$$

input `Int[Cos[3*x]*Csc[x]^2*Sec[x]^3*Sin[2*x],x]`

output `Log[Sin[x]^2] + 3*Log[1 - Sin[x]^2]`

3.301.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 354 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4865 `Int[(u_)*(F_)[(c_.)*((a_.) + (b_.)*(x_))]^(n_), x_Symbol] := With[{d = FreeFactors[Sin[c*(a + b*x)], x]}, Simp[d/(b*c) Subst[Int[SubstFor[(1 - d^2*x^2)^((-n - 1)/2), Sin[c*(a + b*x)]/d, u, x], x], x, Sin[c*(a + b*x)]/d, x] /; FunctionOfQ[Sin[c*(a + b*x)]/d, u, x] /; FreeQ[{a, b, c}, x] && IntegerQ[(n - 1)/2] && NonsumQ[u] && (EqQ[F, Sec] || EqQ[F, sec])`

3.301.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

$$-6 \ln(\tan(x)) + 8 \ln(\sin(x))$$

input `int(sin(2*x)*cos(3*x)/sin(x)^2/cos(x)^3,x)`

output `-6*ln(tan(x))+8*ln(sin(x))`

3.301.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.55

$$\int \cos(3x) \csc^2(x) \sec^3(x) \sin(2x) dx = 3 \log(\cos(x)^2) + \log\left(-\frac{1}{4} \cos(x)^2 + \frac{1}{4}\right)$$

input `integrate(sin(2*x)*cos(3*x)/sin(x)^2/cos(x)^3,x, algorithm="fricas")`

output `3*log(cos(x)^2) + log(-1/4*cos(x)^2 + 1/4)`

3.301.6 Sympy [A] (verification not implemented)

Time = 92.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.36

$$\int \cos(3x) \csc^2(x) \sec^3(x) \sin(2x) dx = 3 \log(\sin^2(x) - 1) + 2 \log(\sin(x))$$

input `integrate(sin(2*x)*cos(3*x)/sin(x)**2/cos(x)**3,x)`

output `3*log(sin(x)**2 - 1) + 2*log(sin(x))`

3.301.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 54 vs. 2(11) = 22.

Time = 0.28 (sec) , antiderivative size = 54, normalized size of antiderivative = 4.91

$$\begin{aligned} \int \cos(3x) \csc^2(x) \sec^3(x) \sin(2x) dx = & 3 \log(\cos(2x)^2 + \sin(2x)^2 + 2 \cos(2x) + 1) \\ & + \log(\cos(x)^2 + \sin(x)^2 + 2 \cos(x) + 1) \\ & + \log(\cos(x)^2 + \sin(x)^2 - 2 \cos(x) + 1) \end{aligned}$$

input `integrate(sin(2*x)*cos(3*x)/sin(x)^2/cos(x)^3,x, algorithm="maxima")`

output `3*log(cos(2*x)^2 + sin(2*x)^2 + 2*cos(2*x) + 1) + log(cos(x)^2 + sin(x)^2 + 2*cos(x) + 1) + log(cos(x)^2 + sin(x)^2 - 2*cos(x) + 1)`

3.301.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.45

$$\int \cos(3x) \csc^2(x) \sec^3(x) \sin(2x) dx = \log(-\cos(x)^2 + 1) + 6 \log(|\cos(x)|)$$

input `integrate(sin(2*x)*cos(3*x)/sin(x)^2/cos(x)^3,x, algorithm="giac")`

output `log(-cos(x)^2 + 1) + 6*log(abs(cos(x)))`

3.301.9 Mupad [B] (verification not implemented)

Time = 16.29 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \cos(3x) \csc^2(x) \sec^3(x) \sin(2x) dx = 6 \ln(\cos(x)) + \ln(\sin(x)^2)$$

input `int((cos(3*x)*sin(2*x))/(cos(x)^3*sin(x)^2),x)`

output `6*log(cos(x)) + log(sin(x)^2)`

$$\mathbf{3.302} \quad \int \left(\frac{1}{\sqrt{2}\sqrt{\log(x)}} + \sqrt{2}\sqrt{\log(x)} \right) dx$$

3.302.1 Optimal result	1502
3.302.2 Mathematica [A] (verified)	1502
3.302.3 Rubi [A] (verified)	1503
3.302.4 Maple [F]	1503
3.302.5 Fricas [F(-2)]	1504
3.302.6 Sympy [B] (verification not implemented)	1504
3.302.7 Maxima [C] (verification not implemented)	1505
3.302.8 Giac [C] (verification not implemented)	1505
3.302.9 Mupad [B] (verification not implemented)	1506

3.302.1 Optimal result

Integrand size = 25, antiderivative size = 13

$$\int \left(\frac{1}{\sqrt{2}\sqrt{\log(x)}} + \sqrt{2}\sqrt{\log(x)} \right) dx = \sqrt{2}x\sqrt{\log(x)}$$

output `2^(1/2)*x*ln(x)^(1/2)`

3.302.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \left(\frac{1}{\sqrt{2}\sqrt{\log(x)}} + \sqrt{2}\sqrt{\log(x)} \right) dx = \sqrt{2}x\sqrt{\log(x)}$$

input `Integrate[1/(Sqrt[2]*Sqrt[Log[x]]) + Sqrt[2]*Sqrt[Log[x]], x]`

output `Sqrt[2]*x*Sqrt[Log[x]]`

3.302.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\sqrt{2}\sqrt{\log(x)} + \frac{1}{\sqrt{2}\sqrt{\log(x)}} \right) dx$$

↓ 2009

$$\sqrt{2x}\sqrt{\log(x)}$$

input `Int[1/(Sqrt[2]*Sqrt[Log[x]]) + Sqrt[2]*Sqrt[Log[x]],x]`

output `Sqrt[2]*x*Sqrt[Log[x]]`

3.302.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.302.4 Maple [F]

$$\int \left(\sqrt{2}\sqrt{\ln(x)} + \frac{\sqrt{2}}{2\sqrt{\ln(x)}} \right) dx$$

input `int(2^(1/2)*ln(x)^(1/2)+1/2*2^(1/2)/ln(x)^(1/2),x)`

output `int(2^(1/2)*ln(x)^(1/2)+1/2*2^(1/2)/ln(x)^(1/2),x)`

3.302.5 Fricas [F(-2)]

Exception generated.

$$\int \left(\frac{1}{\sqrt{2}\sqrt{\log(x)}} + \sqrt{2}\sqrt{\log(x)} \right) dx = \text{Exception raised: TypeError}$$

input `integrate(2^(1/2)*log(x)^(1/2)+1/2*2^(1/2)/log(x)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

3.302.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 83 vs. $2(12) = 24$.

Time = 0.24 (sec) , antiderivative size = 83, normalized size of antiderivative = 6.38

$$\int \left(\frac{1}{\sqrt{2}\sqrt{\log(x)}} + \sqrt{2}\sqrt{\log(x)} \right) dx = \frac{\sqrt{2}\sqrt{\pi}\sqrt{-\log(x)} \operatorname{erfc}\left(\sqrt{-\log(x)}\right)}{2\sqrt{\log(x)}} + \frac{\sqrt{2}\left(x\sqrt{-\log(x)} + \frac{\sqrt{\pi}\operatorname{erfc}\left(\frac{\sqrt{-\log(x)}}{2}\right)}{2}\right)\sqrt{\log(x)}}{\sqrt{-\log(x)}}$$

input `integrate(2**(1/2)*ln(x)**(1/2)+1/2*2**(1/2)/ln(x)**(1/2),x)`

output `sqrt(2)*sqrt(pi)*sqrt(-log(x))*erfc(sqrt(-log(x)))/(2*sqrt(log(x))) + sqrt(2)*(x*sqrt(-log(x)) + sqrt(pi)*erfc(sqrt(-log(x)))/2)*sqrt(log(x))/sqrt(-log(x))`

3.302.7 Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 0.29 (sec) , antiderivative size = 41, normalized size of antiderivative = 3.15

$$\int \left(\frac{1}{\sqrt{2}\sqrt{\log(x)}} + \sqrt{2}\sqrt{\log(x)} \right) dx = -\frac{1}{2}i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(i\sqrt{\log(x)}\right) - \frac{1}{2}\sqrt{2}\left(-i\sqrt{\pi} \operatorname{erf}\left(i\sqrt{\log(x)}\right) - 2x\sqrt{\log(x)}\right)$$

input `integrate(2^(1/2)*log(x)^(1/2)+1/2*2^(1/2)/log(x)^(1/2),x, algorithm="maxima")`

output `-1/2*I*sqrt(2)*sqrt(pi)*erf(I*sqrt(log(x))) - 1/2*sqrt(2)*(-I*sqrt(pi)*erf(I*sqrt(log(x))) - 2*x*sqrt(log(x)))`

3.302.8 Giac [C] (verification not implemented)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 0.27 (sec) , antiderivative size = 41, normalized size of antiderivative = 3.15

$$\int \left(\frac{1}{\sqrt{2}\sqrt{\log(x)}} + \sqrt{2}\sqrt{\log(x)} \right) dx = \frac{1}{2}i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-i\sqrt{\log(x)}\right) - \frac{1}{2}\sqrt{2}\left(i\sqrt{\pi} \operatorname{erf}\left(-i\sqrt{\log(x)}\right) - 2x\sqrt{\log(x)}\right)$$

input `integrate(2^(1/2)*log(x)^(1/2)+1/2*2^(1/2)/log(x)^(1/2),x, algorithm="giac")`

output `1/2*I*sqrt(2)*sqrt(pi)*erf(-I*sqrt(log(x))) - 1/2*sqrt(2)*(I*sqrt(pi)*erf(-I*sqrt(log(x))) - 2*x*sqrt(log(x)))`

3.302.9 Mupad [B] (verification not implemented)

Time = 16.43 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \left(\frac{1}{\sqrt{2}\sqrt{\log(x)}} + \sqrt{2}\sqrt{\log(x)} \right) dx = \sqrt{2}x \sqrt{\ln(x)}$$

input `int(2^(1/2)/(2*log(x)^(1/2)) + 2^(1/2)*log(x)^(1/2),x)`output `2^(1/2)*x*log(x)^(1/2)`

3.303 $\int \log(\cos(x)) \sec^2(x) dx$

3.303.1 Optimal result	1507
3.303.2 Mathematica [A] (verified)	1507
3.303.3 Rubi [A] (verified)	1508
3.303.4 Maple [A] (verified)	1509
3.303.5 Fricas [A] (verification not implemented)	1510
3.303.6 Sympy [B] (verification not implemented)	1510
3.303.7 Maxima [B] (verification not implemented)	1510
3.303.8 Giac [A] (verification not implemented)	1511
3.303.9 Mupad [B] (verification not implemented)	1511

3.303.1 Optimal result

Integrand size = 8, antiderivative size = 12

$$\int \log(\cos(x)) \sec^2(x) dx = -x + \tan(x) + \log(\cos(x)) \tan(x)$$

output `-x+tan(x)+ln(cos(x))*tan(x)`

3.303.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \log(\cos(x)) \sec^2(x) dx = -x + \tan(x) + \log(\cos(x)) \tan(x)$$

input `Integrate[Log[Cos[x]]*Sec[x]^2,x]`

output `-x + Tan[x] + Log[Cos[x]]*Tan[x]`

3.303.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3034, 25, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(x) \log(\cos(x)) dx \\
 & \quad \downarrow \text{3034} \\
 & \tan(x) \log(\cos(x)) - \int -\tan^2(x) dx \\
 & \quad \downarrow \text{25} \\
 & \int \tan^2(x) dx + \tan(x) \log(\cos(x)) \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^2 dx + \tan(x) \log(\cos(x)) \\
 & \quad \downarrow \text{3954} \\
 & - \int 1 dx + \tan(x) + \tan(x) \log(\cos(x)) \\
 & \quad \downarrow \text{24} \\
 & -x + \tan(x) + \tan(x) \log(\cos(x))
 \end{aligned}$$

input `Int [Log [Cos [x]] *Sec [x]^2, x]`

output `-x + Tan[x] + Log[Cos[x]]*Tan[x]`

3.303.3.1 Defintions of rubi rules used

rule 244 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 255 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3034 `Int[Log[u_]*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[Log[u] w, x] - Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w, x]] /; InverseFunctionFreeQ[u, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

3.303.4 Maple [A] (verified)

Time = 0.45 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

method	result
parallelrisch	$-x + \tan(x) + \ln(\cos(x)) \tan(x)$
norman	$\frac{x - x \tan\left(\frac{x}{2}\right)^2 - 2 \tan\left(\frac{x}{2}\right) \ln\left(\frac{1 - \tan\left(\frac{x}{2}\right)^2}{1 + \tan\left(\frac{x}{2}\right)^2}\right) - 2 \tan\left(\frac{x}{2}\right)}{\tan\left(\frac{x}{2}\right)^2 - 1}$
default	$-4i \left(\frac{e^{2ix} \ln\left(\frac{e^{2ix} + 1}{e^{-ix}}\right) - \frac{1}{2}}{e^{2ix} + 1} - \frac{\ln(e^{2ix} + 1)}{4} + \frac{\ln(2)}{2e^{2ix} + 2} \right)$
risch	$-\frac{2i \ln(e^{ix})}{e^{2ix} + 1} + \frac{-i \ln(e^{2ix} + 1) e^{2ix} + \pi \operatorname{csgn}(ie^{-ix}) \operatorname{csgn}(i(e^{2ix} + 1)) \operatorname{csgn}(i \cos(x)) - \pi \operatorname{csgn}(ie^{-ix}) \operatorname{csgn}(i \cos(x))^2 - \pi \operatorname{csgn}(i)}{e^{2ix} + 1}$

input `int(ln(cos(x))/cos(x)^2,x,method=_RETURNVERBOSE)`

output `-x+tan(x)+ln(cos(x))*tan(x)`

3.303.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.83

$$\int \log(\cos(x)) \sec^2(x) dx = -\frac{x \cos(x) - \log(\cos(x)) \sin(x) - \sin(x)}{\cos(x)}$$

input `integrate(log(cos(x))/cos(x)^2,x, algorithm="fricas")`

output `-(x*cos(x) - log(cos(x))*sin(x) - sin(x))/cos(x)`

3.303.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 83 vs. 2(12) = 24.

Time = 1.47 (sec) , antiderivative size = 83, normalized size of antiderivative = 6.92

$$\int \log(\cos(x)) \sec^2(x) dx = -\frac{x \tan^2\left(\frac{x}{2}\right)}{\tan^2\left(\frac{x}{2}\right) - 1} + \frac{x}{\tan^2\left(\frac{x}{2}\right) - 1} - \frac{2 \log\left(-\frac{\tan^2\left(\frac{x}{2}\right)}{\tan^2\left(\frac{x}{2}\right)+1} + \frac{1}{\tan^2\left(\frac{x}{2}\right)+1}\right) \tan\left(\frac{x}{2}\right)}{\tan^2\left(\frac{x}{2}\right) - 1} - \frac{2 \tan\left(\frac{x}{2}\right)}{\tan^2\left(\frac{x}{2}\right) - 1}$$

input `integrate(ln(cos(x))/cos(x)**2,x)`

output `-x*tan(x/2)**2/(tan(x/2)**2 - 1) + x/(tan(x/2)**2 - 1) - 2*log(-tan(x/2)**2/(tan(x/2)**2 + 1) + 1/(tan(x/2)**2 + 1))*tan(x/2)/(tan(x/2)**2 - 1) - 2*tan(x/2)/(tan(x/2)**2 - 1)`

3.303.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 94 vs. 2(12) = 24.

Time = 0.29 (sec) , antiderivative size = 94, normalized size of antiderivative = 7.83

$$\int \log(\cos(x)) \sec^2(x) dx = -\frac{2 \log\left(-\frac{\frac{\sin(x)^2}{(\cos(x)+1)^2}-1}{\frac{\sin(x)^2}{(\cos(x)+1)^2}+1}\right) \sin(x)}{\left(\frac{\sin(x)^2}{(\cos(x)+1)^2} - 1\right) (\cos(x) + 1)} - \frac{2 \sin(x)}{\left(\frac{\sin(x)^2}{(\cos(x)+1)^2} - 1\right) (\cos(x) + 1)} - 2 \arctan\left(\frac{\sin(x)}{\cos(x) + 1}\right)$$

input `integrate(log(cos(x))/cos(x)^2,x, algorithm="maxima")`

output `-2*log(-(sin(x)^2/(cos(x) + 1)^2 - 1)/(sin(x)^2/(cos(x) + 1)^2 + 1))*sin(x)
)/((sin(x)^2/(cos(x) + 1)^2 - 1)*(cos(x) + 1)) - 2*sin(x)/((sin(x)^2/(cos(x)
+ 1)^2 - 1)*(cos(x) + 1)) - 2*arctan(sin(x)/(cos(x) + 1))`

3.303.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \log(\cos(x)) \sec^2(x) dx = \log(\cos(x)) \tan(x) - x + \tan(x)$$

input `integrate(log(cos(x))/cos(x)^2,x, algorithm="giac")`

output `log(cos(x))*tan(x) - x + tan(x)`

3.303.9 Mupad [B] (verification not implemented)

Time = 16.94 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.92

$$\int \log(\cos(x)) \sec^2(x) dx = \tan(x) - 2x + \ln(\cos(x)) \tan(x) + \ln(\cos(x)) \operatorname{li} \\ - \ln(\cos(2x) + 1 + \sin(2x) \operatorname{li}) \operatorname{li}$$

input `int(log(cos(x))/cos(x)^2,x)`

output `log(cos(x))*1i - 2*x - log(cos(2*x) + sin(2*x)*1i + 1)*1i + tan(x) + log(c
os(x))*tan(x)`

3.304
$$\int \frac{\frac{5}{(-5+x)^6} + \frac{3}{(-3+x)^4} + \frac{1}{(-1+x)^2}}{\left(1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}\right)^2} dx$$

3.304.1 Optimal result	1512
3.304.2 Mathematica [A] (verified)	1512
3.304.3 Rubi [A] (verified)	1513
3.304.4 Maple [B] (verified)	1514
3.304.5 Fricas [B] (verification not implemented)	1514
3.304.6 Sympy [B] (verification not implemented)	1515
3.304.7 Maxima [A] (verification not implemented)	1515
3.304.8 Giac [B] (verification not implemented)	1516
3.304.9 Mupad [B] (verification not implemented)	1516

3.304.1 Optimal result

Integrand size = 40, antiderivative size = 19

$$\int \frac{\frac{5}{(-5+x)^6} + \frac{3}{(-3+x)^4} + \frac{1}{(-1+x)^2}}{\left(1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}\right)^2} dx = \frac{1}{1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}}$$

output `1/(1+1/(-1+x)+1/(-3+x)^3+1/(-5+x)^5)`

3.304.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\frac{5}{(-5+x)^6} + \frac{3}{(-3+x)^4} + \frac{1}{(-1+x)^2}}{\left(1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}\right)^2} dx = \frac{1}{1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}}$$

input `Integrate[(5/(-5 + x)^6 + 3/(-3 + x)^4 + (-1 + x)^(-2))/(1 + (-5 + x)^(-5) + (-3 + x)^(-3) + (-1 + x)^(-1))^2,x]`

output `(1 + (-5 + x)^(-5) + (-3 + x)^(-3) + (-1 + x)^(-1))^(-1)`

3.304.
$$\int \frac{\frac{5}{(-5+x)^6} + \frac{3}{(-3+x)^4} + \frac{1}{(-1+x)^2}}{\left(1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}\right)^2} dx$$

3.304.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.025$, Rules used = {7237}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\frac{3}{(x-3)^4} + \frac{5}{(x-5)^6} + \frac{1}{(x-1)^2}}{\left(\frac{1}{(x-3)^3} + \frac{1}{(x-5)^5} + \frac{1}{x-1} + 1\right)^2} dx$$

↓ 7237

$$\frac{1}{\frac{1}{(x-3)^3} + \frac{1}{(x-5)^5} + \frac{1}{x-1} + 1}$$

input `Int[(5/(-5 + x)^6 + 3/(-3 + x)^4 + (-1 + x)^(-2))/(1 + (-5 + x)^(-5) + (-3 + x)^(-3) + (-1 + x)^(-1))^2,x]`

output `(1 + (-5 + x)^(-5) + (-3 + x)^(-3) + (-1 + x)^(-1))^(-1)`

3.304.3.1 Defintions of rubi rules used

rule 7237 `Int[(u_)*(y_)^(m_.), x_Symbol] :> With[{q = DerivativeDivides[y, u, x]}, Si mp[q*(y^(m + 1)/(m + 1)), x] /; !FalseQ[q]] /; FreeQ[m, x] && NeQ[m, -1]`

3.304. $\int \frac{\frac{5}{(-5+x)^6} + \frac{3}{(-3+x)^4} + \frac{1}{(-1+x)^2}}{\left(1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}\right)^2} dx$

3.304.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 85 vs. $2(19) = 38$.

Time = 0.26 (sec) , antiderivative size = 86, normalized size of antiderivative = 4.53

method	result
gospers	$-\frac{x^8-34x^7+503x^6-4228x^5+22076x^4-73260x^3+150661x^2-175054x+87527}{x^9-34x^8+502x^7-4201x^6+21774x^5-71474x^4+144740x^3-164339x^2+78071x+3152}$
default	$\frac{-x^8+34x^7-503x^6+4228x^5-22076x^4+73260x^3-150661x^2+175054x-87527}{x^9-34x^8+502x^7-4201x^6+21774x^5-71474x^4+144740x^3-164339x^2+78071x+3152}$
risch	$\frac{-x^8+34x^7-503x^6+4228x^5-22076x^4+73260x^3-150661x^2+175054x-87527}{x^9-34x^8+502x^7-4201x^6+21774x^5-71474x^4+144740x^3-164339x^2+78071x+3152}$
parallelrisch	$\frac{-x^8+34x^7-503x^6+4228x^5-22076x^4+73260x^3-150661x^2+175054x-87527}{x^9-34x^8+502x^7-4201x^6+21774x^5-71474x^4+144740x^3-164339x^2+78071x+3152}$
norman	$\frac{-x^{17}+69x^{16}-2229x^{15}+44761x^{14}-625602x^{13}+6455858x^{12}-50913715x^{11}+313280159x^{10}-1521750430x^9+5864557678x^8-1521750430x^7+5864557678x^6-1521750430x^5+5864557678x^4-1521750430x^3+5864557678x^2-1521750430x+5864557678}{(-1+x)(-3+x)^3(-5+x)^5(x^9-34x^8+502x^7-4201x^6+21774x^5-71474x^4+144740x^3-164339x^2+78071x+3152)}$

input `int((1/(-1+x)^2+3/(-3+x)^4+5/(-5+x)^6)/(1+1/(-1+x)+1/(-3+x)^3+1/(-5+x)^5)^2,x,method=_RETURNVERBOSE)`

output `-(x^8-34*x^7+503*x^6-4228*x^5+22076*x^4-73260*x^3+150661*x^2-175054*x+87527)/(x^9-34*x^8+502*x^7-4201*x^6+21774*x^5-71474*x^4+144740*x^3-164339*x^2+78071*x+3152)`

3.304.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 85 vs. $2(19) = 38$.

Time = 0.23 (sec) , antiderivative size = 85, normalized size of antiderivative = 4.47

$$\int \frac{\frac{5}{(-5+x)^6} + \frac{3}{(-3+x)^4} + \frac{1}{(-1+x)^2}}{\left(1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}\right)^2} dx =$$

$$-\frac{x^8 - 34x^7 + 503x^6 - 4228x^5 + 22076x^4 - 73260x^3 + 150661x^2 - 175054x + 87527}{x^9 - 34x^8 + 502x^7 - 4201x^6 + 21774x^5 - 71474x^4 + 144740x^3 - 164339x^2 + 78071x + 3152}$$

input `integrate((1/(-1+x)^2+3/(-3+x)^4+5/(-5+x)^6)/(1+1/(-1+x)+1/(-3+x)^3+1/(-5+x)^5)^2,x, algorithm="fracas")`

output `-(x^8 - 34*x^7 + 503*x^6 - 4228*x^5 + 22076*x^4 - 73260*x^3 + 150661*x^2 - 175054*x + 87527)/(x^9 - 34*x^8 + 502*x^7 - 4201*x^6 + 21774*x^5 - 71474*x^4 + 144740*x^3 - 164339*x^2 + 78071*x + 3152)`

3.304.
$$\int \frac{\frac{5}{(-5+x)^6} + \frac{3}{(-3+x)^4} + \frac{1}{(-1+x)^2}}{\left(1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}\right)^2} dx$$

3.304.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 82 vs. $2(20) = 40$.

Time = 0.47 (sec) , antiderivative size = 82, normalized size of antiderivative = 4.32

$$\int \frac{\frac{5}{(-5+x)^6} + \frac{3}{(-3+x)^4} + \frac{1}{(-1+x)^2}}{\left(1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}\right)^2} dx$$

$$= \frac{-x^8 + 34x^7 - 503x^6 + 4228x^5 - 22076x^4 + 73260x^3 - 150661x^2 + 175054x - 87527}{x^9 - 34x^8 + 502x^7 - 4201x^6 + 21774x^5 - 71474x^4 + 144740x^3 - 164339x^2 + 78071x + 3152}$$

input `integrate((1/(-1+x)**2+3/(-3+x)**4+5/(-5+x)**6)/(1+1/(-1+x)+1/(-3+x)**3+1/(-5+x)**5)**2,x)`

output `(-x**8 + 34*x**7 - 503*x**6 + 4228*x**5 - 22076*x**4 + 73260*x**3 - 150661*x**2 + 175054*x - 87527)/(x**9 - 34*x**8 + 502*x**7 - 4201*x**6 + 21774*x**5 - 71474*x**4 + 144740*x**3 - 164339*x**2 + 78071*x + 3152)`

3.304.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\frac{5}{(-5+x)^6} + \frac{3}{(-3+x)^4} + \frac{1}{(-1+x)^2}}{\left(1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}\right)^2} dx = \frac{1}{\frac{1}{x-1} + \frac{1}{(x-3)^3} + \frac{1}{(x-5)^5} + 1}$$

input `integrate((1/(-1+x)^2+3/(-3+x)^4+5/(-5+x)^6)/(1+1/(-1+x)+1/(-3+x)^3+1/(-5+x)^5)^2,x, algorithm="maxima")`

output `1/(1/(x - 1) + 1/(x - 3)^3 + 1/(x - 5)^5 + 1)`

3.304. $\int \frac{\frac{5}{(-5+x)^6} + \frac{3}{(-3+x)^4} + \frac{1}{(-1+x)^2}}{\left(1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}\right)^2} dx$

3.304.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 85 vs. $2(19) = 38$.

Time = 0.29 (sec) , antiderivative size = 85, normalized size of antiderivative = 4.47

$$\int \frac{\frac{5}{(-5+x)^6} + \frac{3}{(-3+x)^4} + \frac{1}{(-1+x)^2}}{\left(1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}\right)^2} dx = \frac{x^8 - 34x^7 + 503x^6 - 4228x^5 + 22076x^4 - 73260x^3 + 150661x^2 - 175054x + 87527}{x^9 - 34x^8 + 502x^7 - 4201x^6 + 21774x^5 - 71474x^4 + 144740x^3 - 164339x^2 + 78071x + 3152}$$

input `integrate((1/(-1+x)^2+3/(-3+x)^4+5/(-5+x)^6)/(1+1/(-1+x)+1/(-3+x)^3+1/(-5+x)^5)^2,x, algorithm="giac")`

output `-(x^8 - 34*x^7 + 503*x^6 - 4228*x^5 + 22076*x^4 - 73260*x^3 + 150661*x^2 - 175054*x + 87527)/(x^9 - 34*x^8 + 502*x^7 - 4201*x^6 + 21774*x^5 - 71474*x^4 + 144740*x^3 - 164339*x^2 + 78071*x + 3152)`

3.304.9 Mupad [B] (verification not implemented)

Time = 17.04 (sec) , antiderivative size = 85, normalized size of antiderivative = 4.47

$$\int \frac{\frac{5}{(-5+x)^6} + \frac{3}{(-3+x)^4} + \frac{1}{(-1+x)^2}}{\left(1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}\right)^2} dx = \frac{x^8 - 34x^7 + 503x^6 - 4228x^5 + 22076x^4 - 73260x^3 + 150661x^2 - 175054x + 87527}{x^9 - 34x^8 + 502x^7 - 4201x^6 + 21774x^5 - 71474x^4 + 144740x^3 - 164339x^2 + 78071x + 3152}$$

input `int((1/(x - 1)^2 + 3/(x - 3)^4 + 5/(x - 5)^6)/(1/(x - 1) + 1/(x - 3)^3 + 1/(x - 5)^5 + 1)^2,x)`

output `-(150661*x^2 - 175054*x - 73260*x^3 + 22076*x^4 - 4228*x^5 + 503*x^6 - 34*x^7 + x^8 + 87527)/(78071*x - 164339*x^2 + 144740*x^3 - 71474*x^4 + 21774*x^5 - 4201*x^6 + 502*x^7 - 34*x^8 + x^9 + 3152)`

3.304. $\int \frac{\frac{5}{(-5+x)^6} + \frac{3}{(-3+x)^4} + \frac{1}{(-1+x)^2}}{\left(1 + \frac{1}{(-5+x)^5} + \frac{1}{(-3+x)^3} + \frac{1}{-1+x}\right)^2} dx$

3.305 $\int \csc(x) \sin(23x) dx$

3.305.1 Optimal result	1517
3.305.2 Mathematica [A] (verified)	1517
3.305.3 Rubi [B] (verified)	1518
3.305.4 Maple [A] (verified)	1523
3.305.5 Fricas [A] (verification not implemented)	1524
3.305.6 Sympy [A] (verification not implemented)	1524
3.305.7 Maxima [F(-1)]	1525
3.305.8 Giac [A] (verification not implemented)	1525
3.305.9 Mupad [B] (verification not implemented)	1525

3.305.1 Optimal result

Integrand size = 7, antiderivative size = 86

$$\int \csc(x) \sin(23x) dx = x + \sin(2x) + \frac{1}{2} \sin(4x) + \frac{1}{3} \sin(6x) + \frac{1}{4} \sin(8x) + \frac{1}{5} \sin(10x) + \frac{1}{6} \sin(12x) \\ + \frac{1}{7} \sin(14x) + \frac{1}{8} \sin(16x) + \frac{1}{9} \sin(18x) + \frac{1}{10} \sin(20x) + \frac{1}{11} \sin(22x)$$

output `x+sin(2*x)+1/2*sin(4*x)+1/3*sin(6*x)+1/4*sin(8*x)+1/5*sin(10*x)+1/6*sin(12*x)+1/7*sin(14*x)+1/8*sin(16*x)+1/9*sin(18*x)+1/10*sin(20*x)+1/11*sin(22*x)`

3.305.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.00

$$\int \csc(x) \sin(23x) dx = x + \sin(2x) + \frac{1}{2} \sin(4x) + \frac{1}{3} \sin(6x) + \frac{1}{4} \sin(8x) + \frac{1}{5} \sin(10x) + \frac{1}{6} \sin(12x) \\ + \frac{1}{7} \sin(14x) + \frac{1}{8} \sin(16x) + \frac{1}{9} \sin(18x) + \frac{1}{10} \sin(20x) + \frac{1}{11} \sin(22x)$$

input `Integrate[Csc[x]*Sin[23*x],x]`

output `x + Sin[2*x] + Sin[4*x]/2 + Sin[6*x]/3 + Sin[8*x]/4 + Sin[10*x]/5 + Sin[12*x]/6 + Sin[14*x]/7 + Sin[16*x]/8 + Sin[18*x]/9 + Sin[20*x]/10 + Sin[22*x]/11`

3.305.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 190 vs. $2(86) = 172$.

Time = 1.24 (sec) , antiderivative size = 190, normalized size of antiderivative = 2.21, number of steps used = 25, number of rules used = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 3.429$, Rules used = {3042, 4889, 2345, 27, 2345, 27, 2345, 27, 2345, 27, 2345, 27, 2345, 27, 2345, 27, 2345, 27, 2345, 27, 2345, 27, 1471, 27, 298, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(23x) \csc(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(23x)}{\sin(x)} dx \\
 & \quad \downarrow \text{4889} \\
 & \int \frac{-\tan^{22}(x) + 253 \tan^{20}(x) - 8855 \tan^{18}(x) + 100947 \tan^{16}(x) - 490314 \tan^{14}(x) + 1144066 \tan^{12}(x) - 1352070 \tan^{10}(x) + 1144066 \tan^8(x) - 490314 \tan^6(x) + 100947 \tan^4(x) - 253 \tan^2(x) + 1}{(\tan^2(x) + 1)^{12}} dx \\
 & \quad \downarrow \text{2345} \\
 & \frac{1}{22} \int \frac{2(11 \tan^{20}(x) - 2794 \tan^{18}(x) + 100199 \tan^{16}(x) - 1210616 \tan^{14}(x) + 6604070 \tan^{12}(x) - 19188796 \tan^{10}(x) + 1144066 \tan^8(x) - 490314 \tan^6(x) + 100947 \tan^4(x) - 253 \tan^2(x) + 1)}{(\tan^2(x) + 1)^{11}} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{11} \int \frac{11 \tan^{20}(x) - 2794 \tan^{18}(x) + 100199 \tan^{16}(x) - 1210616 \tan^{14}(x) + 6604070 \tan^{12}(x) - 19188796 \tan^{10}(x) + 1144066 \tan^8(x) - 490314 \tan^6(x) + 100947 \tan^4(x) - 253 \tan^2(x) + 1}{(\tan^2(x) + 1)^{11}} dx \\
 & \quad \downarrow \text{2345} \\
 & \frac{1}{11} \left(\frac{1}{20} \int \frac{4(-55 \tan^{18}(x) + 14025 \tan^{16}(x) - 515020 \tan^{14}(x) + 6568100 \tan^{12}(x) - 39588450 \tan^{10}(x) + 13553200 \tan^8(x) - 2980000 \tan^6(x) + 298000 \tan^4(x) - 20 \tan^2(x) + 1)}{(\tan^2(x) + 1)^{11}} dx \right. \\
 & \quad \left. \frac{2097152 \tan(x)}{11 (\tan^2(x) + 1)^{11}} \right) \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$\frac{1}{11} \left(\frac{1}{5} \int \frac{-55 \tan^{18}(x) + 14025 \tan^{16}(x) - 515020 \tan^{14}(x) + 6568100 \tan^{12}(x) - 39588450 \tan^{10}(x) + 135532430 \tan^8(x) - 29852100 \tan^6(x) + 2345 \tan^4(x) - 110 \tan^2(x) + 1}{(\tan^2(x) + 1)^{10}} dx \right)$$

$$\frac{2097152 \tan(x)}{11 (\tan^2(x) + 1)^{11}}$$

↓ 2345

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{899022848 \tan(x)}{9 (\tan^2(x) + 1)^9} - \frac{1}{18} \int \frac{2(495 \tan^{16}(x) - 126720 \tan^{14}(x) + 4761900 \tan^{12}(x) - 63874800 \tan^{10}(x) + 42072000 \tan^8(x) - 110 \tan^6(x) + 1)}{(\tan^2(x) + 1)^8} dx \right) \right)$$

$$\frac{2097152 \tan(x)}{11 (\tan^2(x) + 1)^{11}}$$

↓ 27

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{899022848 \tan(x)}{9 (\tan^2(x) + 1)^9} - \frac{1}{9} \int \frac{495 \tan^{16}(x) - 126720 \tan^{14}(x) + 4761900 \tan^{12}(x) - 63874800 \tan^{10}(x) + 42072000 \tan^8(x) - 110 \tan^6(x) + 1}{(\tan^2(x) + 1)^8} dx \right) \right)$$

$$\frac{2097152 \tan(x)}{11 (\tan^2(x) + 1)^{11}}$$

↓ 2345

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(\frac{1}{16} \int \frac{240(-33 \tan^{14}(x) + 8481 \tan^{12}(x) - 325941 \tan^{10}(x) + 4584261 \tan^8(x) - 32595651 \tan^6(x) + 1419240 \tan^4(x) - 110 \tan^2(x) + 1)}{(\tan^2(x) + 1)^8} dx \right) \right) \right)$$

$$\frac{2097152 \tan(x)}{11 (\tan^2(x) + 1)^{11}}$$

↓ 27

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(15 \int \frac{-33 \tan^{14}(x) + 8481 \tan^{12}(x) - 325941 \tan^{10}(x) + 4584261 \tan^8(x) - 32595651 \tan^6(x) + 1419240 \tan^4(x) - 110 \tan^2(x) + 1}{(\tan^2(x) + 1)^8} dx \right) \right) \right)$$

$$\frac{2097152 \tan(x)}{11 (\tan^2(x) + 1)^{11}}$$

↓ 2345

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(15 \left(\frac{322500608 \tan(x)}{7 (\tan^2(x) + 1)^7} - \frac{1}{14} \int \frac{2(231 \tan^{12}(x) - 59598 \tan^{10}(x) + 2341185 \tan^8(x) - 34431012 \tan^6(x) + 110 \tan^4(x) - 1)}{(\tan^2(x) + 1)^8} dx \right) \right) \right) \right)$$

$$\frac{2097152 \tan(x)}{11 (\tan^2(x) + 1)^{11}}$$

↓ 27

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(15 \left(\frac{322500608 \tan(x)}{7 (\tan^2(x) + 1)^7} - \frac{1}{7} \int \frac{231 \tan^{12}(x) - 59598 \tan^{10}(x) + 2341185 \tan^8(x) - 34431012 \tan^6(x) + 2097152 \tan(x)}{(\tan^2(x) + 1)^7} dx \right. \right. \right. \right. \\ \left. \left. \left. \left. \frac{2097152 \tan(x)}{11 (\tan^2(x) + 1)^{11}} \right. \right. \right. \right. \\ \left. \left. \left. \left. \downarrow \quad 2345 \right. \right. \right. \right.$$

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(15 \left(\frac{1}{7} \left(\frac{1}{12} \int \frac{44(-63 \tan^{10}(x) + 16317 \tan^8(x) - 654822 \tan^6(x) + 10045098 \tan^4(x) - 81663435 \tan^2(x) + 2097152 \tan(x)}{(\tan^2(x) + 1)^6} dx \right. \right. \right. \right. \\ \left. \left. \left. \left. \frac{2097152 \tan(x)}{11 (\tan^2(x) + 1)^{11}} \right. \right. \right. \right. \\ \left. \left. \left. \left. \downarrow \quad 27 \right. \right. \right. \right.$$

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(15 \left(\frac{1}{7} \left(\frac{11}{3} \int \frac{-63 \tan^{10}(x) + 16317 \tan^8(x) - 654822 \tan^6(x) + 10045098 \tan^4(x) - 81663435 \tan^2(x) + 2097152 \tan(x)}{(\tan^2(x) + 1)^6} dx \right. \right. \right. \right. \\ \left. \left. \left. \left. \frac{2097152 \tan(x)}{11 (\tan^2(x) + 1)^{11}} \right. \right. \right. \right. \\ \left. \left. \left. \left. \downarrow \quad 2345 \right. \right. \right. \right.$$

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(15 \left(\frac{1}{7} \left(\frac{11}{3} \left(\frac{10100480 \tan(x)}{(\tan^2(x) + 1)^5} - \frac{1}{10} \int \frac{90(7 \tan^8(x) - 1820 \tan^6(x) + 74578 \tan^4(x) - 1190700 \tan^2(x) + 2097152 \tan(x)}{(\tan^2(x) + 1)^5} dx \right. \right. \right. \right. \\ \left. \left. \left. \left. \frac{2097152 \tan(x)}{11 (\tan^2(x) + 1)^{11}} \right. \right. \right. \right. \\ \left. \left. \left. \left. \downarrow \quad 27 \right. \right. \right. \right.$$

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(15 \left(\frac{1}{7} \left(\frac{11}{3} \left(\frac{10100480 \tan(x)}{(\tan^2(x) + 1)^5} - 9 \int \frac{7 \tan^8(x) - 1820 \tan^6(x) + 74578 \tan^4(x) - 1190700 \tan^2(x) + 2097152 \tan(x)}{(\tan^2(x) + 1)^5} dx \right. \right. \right. \right. \\ \left. \left. \left. \left. \frac{2097152 \tan(x)}{11 (\tan^2(x) + 1)^{11}} \right. \right. \right. \right. \\ \left. \left. \left. \left. \downarrow \quad 2345 \right. \right. \right. \right.$$

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(15 \left(\frac{1}{7} \left(\frac{11}{3} \left(\frac{10100480 \tan(x)}{(\tan^2(x) + 1)^5} - 9 \left(\frac{178880 \tan(x)}{(\tan^2(x) + 1)^4} - \frac{1}{8} \int \frac{56(-\tan^6(x) + 261 \tan^4(x) - 10915 \tan^2(x) + 2097152 \tan(x)}{(\tan^2(x) + 1)^4} dx \right. \right. \right. \right. \\ \left. \left. \left. \left. \frac{2097152 \tan(x)}{11 (\tan^2(x) + 1)^{11}} \right. \right. \right. \right. \\ \left. \left. \left. \left. \downarrow \quad 27 \right. \right. \right. \right.$$

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(15 \left(\frac{1}{7} \left(\frac{11}{3} \left(\frac{10100480 \tan(x)}{(\tan^2(x) + 1)^5} - 9 \left(\frac{178880 \tan(x)}{(\tan^2(x) + 1)^4} - 7 \int \frac{-\tan^6(x) + 261 \tan^4(x) - 10915 \tan^2(x) + 2097152 \tan(x)}{(\tan^2(x) + 1)^4} \right) \right) \right) \right) \right) \right)$$

↓ 2345

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(15 \left(\frac{1}{7} \left(\frac{11}{3} \left(\frac{10100480 \tan(x)}{(\tan^2(x) + 1)^5} - 9 \left(\frac{178880 \tan(x)}{(\tan^2(x) + 1)^4} - 7 \left(\frac{6656 \tan(x)}{3(\tan^2(x) + 1)^3} - \frac{1}{6} \int \frac{2(3 \tan^4(x) - 786 \tan^2(x) + 2097152 \tan(x))}{(\tan^2(x) + 1)^2} \right) \right) \right) \right) \right) \right)$$

↓ 27

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(15 \left(\frac{1}{7} \left(\frac{11}{3} \left(\frac{10100480 \tan(x)}{(\tan^2(x) + 1)^5} - 9 \left(\frac{178880 \tan(x)}{(\tan^2(x) + 1)^4} - 7 \left(\frac{6656 \tan(x)}{3(\tan^2(x) + 1)^3} - \frac{1}{3} \int \frac{3 \tan^4(x) - 786 \tan^2(x) + 2097152 \tan(x)}{(\tan^2(x) + 1)} \right) \right) \right) \right) \right) \right)$$

↓ 1471

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(15 \left(\frac{1}{7} \left(\frac{11}{3} \left(\frac{10100480 \tan(x)}{(\tan^2(x) + 1)^5} - 9 \left(\frac{178880 \tan(x)}{(\tan^2(x) + 1)^4} - 7 \left(\frac{1}{3} \left(\frac{1}{4} \int \frac{12(3 - \tan^2(x))}{(\tan^2(x) + 1)^2} d \tan(x) - \frac{260 \tan(x)}{(\tan^2(x) + 1)} \right) \right) \right) \right) \right) \right)$$

↓ 27

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(15 \left(\frac{1}{7} \left(\frac{11}{3} \left(\frac{10100480 \tan(x)}{(\tan^2(x) + 1)^5} - 9 \left(\frac{178880 \tan(x)}{(\tan^2(x) + 1)^4} - 7 \left(\frac{1}{3} \left(3 \int \frac{3 - \tan^2(x)}{(\tan^2(x) + 1)^2} d \tan(x) - \frac{260 \tan(x)}{(\tan^2(x) + 1)} \right) \right) \right) \right) \right) \right)$$

↓ 298

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(15 \left(\frac{1}{7} \left(\frac{11}{3} \left(\frac{10100480 \tan(x)}{(\tan^2(x) + 1)^5} - 9 \left(\frac{178880 \tan(x)}{(\tan^2(x) + 1)^4} - 7 \left(\frac{1}{3} \left(3 \left(\int \frac{1}{\tan^2(x) + 1} d \tan(x) + \frac{2 \tan(x)}{\tan^2(x) + 1} \right) \right) \right) \right) \right) \right) \right)$$

↓ 216

$$\frac{1}{11} \left(\frac{1}{5} \left(\frac{1}{9} \left(15 \left(\frac{1}{7} \left(\frac{11}{3} \left(\frac{10100480 \tan(x)}{(\tan^2(x) + 1)^5} - 9 \left(\frac{178880 \tan(x)}{(\tan^2(x) + 1)^4} - 7 \left(\frac{1}{3} \left(3 \left(\arctan(\tan(x)) + \frac{2 \tan(x)}{\tan^2(x) + 1} \right) - \frac{2097152 \tan(x)}{11 (\tan^2(x) + 1)^{11}} \right) \right) \right) \right) \right) \right) \right)$$

input `Int[Csc[x]*Sin[23*x],x]`

output `(2097152*Tan[x])/((11*(1 + Tan[x]^2)^11) + ((-49545216*Tan[x])/(5*(1 + Tan[x]^2)^10) + ((899022848*Tan[x])/(9*(1 + Tan[x]^2)^9) + ((-1009455104*Tan[x])/(1 + Tan[x]^2)^8 + 15*((322500608*Tan[x])/(7*(1 + Tan[x]^2)^7) + ((-415607296*Tan[x])/(3*(1 + Tan[x]^2)^6) + (11*((10100480*Tan[x])/(1 + Tan[x]^2)^5 - 9*((178880*Tan[x])/(1 + Tan[x]^2)^4 - 7*((6656*Tan[x])/(3*(1 + Tan[x]^2)^3) + ((-260*Tan[x])/(1 + Tan[x]^2)^2 + 3*(ArcTan[Tan[x]] + (2*Tan[x])/(1 + Tan[x]^2))))/3))))/3)/7)/9)/5)/11`

3.305.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 298 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[(-(b*c - a*d))*x*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] - Simp[(a*d - b*c*(2*p + 3))/(2*a*b*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/2 + p, 0])`

rule 1471 `Int[((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], R = Coeff[PolynomialRemainder[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], x, 0]}, Simp[(-R)*x*((d + e*x^2)^(q + 1)/(2*d*(q + 1))), x] + Simp[1/(2*d*(q + 1)) Int[(d + e*x^2)^(q + 1)*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]`

```
rule 2345 Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(a*g - b*f*x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[1/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1)*ExpandToSum[2*a*(p + 1)*Q + f*(2*p + 3), x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && LtQ[p, -1]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4889 Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, With[{d = FreeFactors[Tan[v], x]}, Simp[d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x^2), Tan[v]/d, u, x], x], x, Tan[v]/d, x]] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Tan[v], x], u, x]] /; InverseFunctionFreeQ[u, x] && !MatchQ[u, (v_.)*((c_.)*tan[w_]^(n_.)*tan[z_]^(n_.))^(p_.) /; FreeQ[{c, p}, x] && IntegerQ[n] && LinearQ[w, x] && EqQ[z, 2*w]]
```

3.305.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.78

method	result
risch	$x + \sin(2x) + \frac{\sin(4x)}{2} + \frac{\sin(6x)}{3} + \frac{\sin(8x)}{4} + \frac{\sin(10x)}{5} + \frac{\sin(12x)}{6} + \frac{\sin(14x)}{7} + \frac{\sin(16x)}{8} + \frac{\sin(18x)}{9} + \frac{\sin(20x)}{10}$

```
input int(sin(23*x)/sin(x),x,method=_RETURNVERBOSE)
```

```
output x+sin(2*x)+1/2*sin(4*x)+1/3*sin(6*x)+1/4*sin(8*x)+1/5*sin(10*x)+1/6*sin(12*x)+1/7*sin(14*x)+1/8*sin(16*x)+1/9*sin(18*x)+1/10*sin(20*x)+1/11*sin(22*x)
```

3.305.5 Fracas [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.83

$$\int \csc(x) \sin(23x) dx$$

$$= \frac{2}{3465} (330301440 \cos(x)^{21} - 1560674304 \cos(x)^{19} + 3146579968 \cos(x)^{17} - 3533092864 \cos(x)^{15} + 241763200 \cos(x)^9 - 44272800 \cos(x)^7 + 3843840 \cos(x)^5 - 150150 \cos(x)^3 + 3465 \cos(x)) \sin(x) + x$$

input `integrate(sin(23*x)/sin(x),x, algorithm="fricas")`output `2/3465*(330301440*cos(x)^21 - 1560674304*cos(x)^19 + 3146579968*cos(x)^17 - 3533092864*cos(x)^15 + 241763200*cos(x)^9 - 44272800*cos(x)^7 + 3843840*cos(x)^5 - 150150*cos(x)^3 + 3465*cos(x))*sin(x) + x`**3.305.6 Sympy [A] (verification not implemented)**

Time = 69.50 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.16

$$\int \csc(x) \sin(23x) dx = x - \frac{1024 \sin^{11}(2x)}{11} + \frac{2560 \sin^9(2x)}{9} - \frac{2304 \sin^7(2x)}{7} + \frac{896 \sin^5(2x)}{5} - \frac{140 \sin^3(2x)}{3} + 6 \sin(2x) + \frac{8 \sin^5(4x)}{5} - \frac{8 \sin^3(4x)}{3} + \frac{3 \sin(4x)}{2} + \frac{\sin(8x)}{4} + \frac{\sin(16x)}{8}$$

input `integrate(sin(23*x)/sin(x),x)`output `x - 1024*sin(2*x)**11/11 + 2560*sin(2*x)**9/9 - 2304*sin(2*x)**7/7 + 896*sin(2*x)**5/5 - 140*sin(2*x)**3/3 + 6*sin(2*x) + 8*sin(4*x)**5/5 - 8*sin(4*x)**3/3 + 3*sin(4*x)/2 + sin(8*x)/4 + sin(16*x)/8`

3.305.7 Maxima [F(-1)]

Timed out.

$$\int \csc(x) \sin(23x) dx = \text{Timed out}$$

input `integrate(sin(23*x)/sin(x),x, algorithm="maxima")`output `Timed out`**3.305.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.77

$$\begin{aligned} \int \csc(x) \sin(23x) dx &= x + \frac{1}{11} \sin(22x) + \frac{1}{10} \sin(20x) + \frac{1}{9} \sin(18x) \\ &+ \frac{1}{8} \sin(16x) + \frac{1}{7} \sin(14x) + \frac{1}{6} \sin(12x) + \frac{1}{5} \sin(10x) \\ &+ \frac{1}{4} \sin(8x) + \frac{1}{3} \sin(6x) + \frac{1}{2} \sin(4x) + \sin(2x) \end{aligned}$$

input `integrate(sin(23*x)/sin(x),x, algorithm="giac")`output `x + 1/11*sin(22*x) + 1/10*sin(20*x) + 1/9*sin(18*x) + 1/8*sin(16*x) + 1/7*
sin(14*x) + 1/6*sin(12*x) + 1/5*sin(10*x) + 1/4*sin(8*x) + 1/3*sin(6*x) +
1/2*sin(4*x) + sin(2*x)`**3.305.9 Mupad [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.77

$$\begin{aligned} \int \csc(x) \sin(23x) dx &= x + \sin(2x) + \frac{\sin(4x)}{2} + \frac{\sin(6x)}{3} + \frac{\sin(8x)}{4} + \frac{\sin(10x)}{5} + \frac{\sin(12x)}{6} \\ &+ \frac{\sin(14x)}{7} + \frac{\sin(16x)}{8} + \frac{\sin(18x)}{9} + \frac{\sin(20x)}{10} + \frac{\sin(22x)}{11} \end{aligned}$$

input `int(sin(23*x)/sin(x),x)`

output $x + \sin(2*x) + \sin(4*x)/2 + \sin(6*x)/3 + \sin(8*x)/4 + \sin(10*x)/5 + \sin(12*x)/6 + \sin(14*x)/7 + \sin(16*x)/8 + \sin(18*x)/9 + \sin(20*x)/10 + \sin(22*x)/11$

$$\mathbf{3.306} \quad \int \frac{(1-x)^2 x^4}{1+x^2} dx$$

3.306.1 Optimal result	1527
3.306.2 Mathematica [A] (verified)	1527
3.306.3 Rubi [A] (verified)	1528
3.306.4 Maple [A] (verified)	1529
3.306.5 Fricas [A] (verification not implemented)	1530
3.306.6 Sympy [A] (verification not implemented)	1530
3.306.7 Maxima [A] (verification not implemented)	1530
3.306.8 Giac [A] (verification not implemented)	1531
3.306.9 Mupad [B] (verification not implemented)	1531

3.306.1 Optimal result

Integrand size = 18, antiderivative size = 26

$$\int \frac{(1-x)^2 x^4}{1+x^2} dx = x^2 - \frac{x^4}{2} + \frac{x^5}{5} - \log(1+x^2)$$

output `x^2-1/2*x^4+1/5*x^5-ln(x^2+1)`

3.306.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{(1-x)^2 x^4}{1+x^2} dx = x^2 - \frac{x^4}{2} + \frac{x^5}{5} - \log(1+x^2)$$

input `Integrate[((1 - x)^2*x^4)/(1 + x^2), x]`

output `x^2 - x^4/2 + x^5/5 - Log[1 + x^2]`

3.306.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {525, 27, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(1-x)^2 x^4}{x^2+1} dx \\
 & \quad \downarrow \text{525} \\
 & \int -\frac{2x^5}{x^2+1} dx + \frac{x^5}{5} \\
 & \quad \downarrow \text{27} \\
 & \frac{x^5}{5} - 2 \int \frac{x^5}{x^2+1} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{x^5}{5} - \int \frac{x^4}{x^2+1} dx^2 \\
 & \quad \downarrow \text{49} \\
 & \frac{x^5}{5} - \int \left(x^2 + \frac{1}{x^2+1} - 1 \right) dx^2 \\
 & \quad \downarrow \text{2009} \\
 & \frac{x^5}{5} - \frac{x^4}{2} + x^2 - \log(x^2+1)
 \end{aligned}$$

input `Int[((1 - x)^2*x^4)/(1 + x^2),x]`

output `x^2 - x^4/2 + x^5/5 - Log[1 + x^2]`

3.306.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 525 `Int[((x_)^(m_.)*((c_) + (d_.)*(x_)^n_))/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[d^n*(x^(m + n - 1)/(b*(m + n - 1))), x] + Simp[1/b Int[x^m*(ExpandToSum[b*(c + d*x)^n - b*d^n*x^n - a*d^n*x^(n - 2), x]/(a + b*x^2)), x] /; FreeQ[{a, b, c, d}, x] && IGtQ[n, 1] && IGtQ[m, -2] && NeQ[m + n - 1, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.306.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.88

method	result	size
default	$x^2 - \frac{x^4}{2} + \frac{x^5}{5} - \ln(x^2 + 1)$	23
norman	$x^2 - \frac{x^4}{2} + \frac{x^5}{5} - \ln(x^2 + 1)$	23
risch	$x^2 - \frac{x^4}{2} + \frac{x^5}{5} - \ln(x^2 + 1)$	23
paralelrisch	$x^2 - \frac{x^4}{2} + \frac{x^5}{5} - \ln(x^2 + 1)$	23
meijerg	$-\frac{x(-5x^2+15)}{15} + \frac{x^2(-3x^2+6)}{6} - \ln(x^2 + 1) + \frac{x(21x^4-35x^2+105)}{105}$	47

input `int(x^4*(1-x)^2/(x^2+1),x,method=_RETURNVERBOSE)`

output $x^2 - 1/2x^4 + 1/5x^5 - \ln(x^2 + 1)$

3.306.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{(1-x)^2 x^4}{1+x^2} dx = \frac{1}{5}x^5 - \frac{1}{2}x^4 + x^2 - \log(x^2 + 1)$$

input `integrate(x^4*(1-x)^2/(x^2+1),x, algorithm="fricas")`

output $1/5x^5 - 1/2x^4 + x^2 - \log(x^2 + 1)$

3.306.6 Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.73

$$\int \frac{(1-x)^2 x^4}{1+x^2} dx = \frac{x^5}{5} - \frac{x^4}{2} + x^2 - \log(x^2 + 1)$$

input `integrate(x**4*(1-x)**2/(x**2+1),x)`

output $x**5/5 - x**4/2 + x**2 - \log(x**2 + 1)$

3.306.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{(1-x)^2 x^4}{1+x^2} dx = \frac{1}{5}x^5 - \frac{1}{2}x^4 + x^2 - \log(x^2 + 1)$$

input `integrate(x^4*(1-x)^2/(x^2+1),x, algorithm="maxima")`

output $1/5x^5 - 1/2x^4 + x^2 - \log(x^2 + 1)$

3.306.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{(1-x)^2 x^4}{1+x^2} dx = \frac{1}{5} x^5 - \frac{1}{2} x^4 + x^2 - \log(x^2 + 1)$$

input `integrate(x^4*(1-x)^2/(x^2+1),x, algorithm="giac")`output `1/5*x^5 - 1/2*x^4 + x^2 - log(x^2 + 1)`**3.306.9 Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{(1-x)^2 x^4}{1+x^2} dx = x^2 - \ln(x^2 + 1) - \frac{x^4}{2} + \frac{x^5}{5}$$

input `int((x^4*(x - 1)^2)/(x^2 + 1),x)`output `x^2 - log(x^2 + 1) - x^4/2 + x^5/5`

3.307 $\int x^{-\log(x)} dx$

3.307.1 Optimal result	1532
3.307.2 Mathematica [A] (verified)	1532
3.307.3 Rubi [F]	1533
3.307.4 Maple [F]	1533
3.307.5 Fracas [A] (verification not implemented)	1533
3.307.6 Sympy [F]	1534
3.307.7 Maxima [F]	1534
3.307.8 Giac [A] (verification not implemented)	1534
3.307.9 Mupad [F(-1)]	1535

3.307.1 Optimal result

Integrand size = 6, antiderivative size = 23

$$\int x^{-\log(x)} dx = -\frac{1}{2} \sqrt[4]{e} \sqrt{\pi} \operatorname{erf}\left(\frac{1}{2} - \log(x)\right)$$

output `1/2*exp(1/4)*Pi^(1/2)*erf(-1/2+ln(x))`

3.307.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int x^{-\log(x)} dx = \frac{1}{2} \sqrt[4]{e} \sqrt{\pi} \operatorname{erf}\left(\frac{1}{2}(-1 + 2 \log(x))\right)$$

input `Integrate[x^(-Log[x]),x]`

output `(E^(1/4)*Sqrt[Pi]*Erf[(-1 + 2*Log[x])/2])/2`

3.307.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{-\log(x)} dx$$

$$\downarrow 7299$$

$$\int x^{-\log(x)} dx$$

input `Int[x^(-Log[x]),x]`

output `$Aborted`

3.307.3.1 Defintions of rubi rules used

rule 7299 `Int[u_, x_] :> CannotIntegrate[u, x]`

3.307.4 Maple [F]

$$\int x^{-\ln(x)} dx$$

input `int(x^(-ln(x)),x)`

output `int(x^(-ln(x)),x)`

3.307.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.52

$$\int x^{-\log(x)} dx = \frac{1}{2} \sqrt{\pi} \operatorname{erf} \left(\log(x) - \frac{1}{2} \right) e^{\frac{1}{4}}$$

input `integrate(x^(-log(x)),x, algorithm="fricas")`

output `1/2*sqrt(pi)*erf(log(x) - 1/2)*e^(1/4)`

3.307.6 Sympy [F]

$$\int x^{-\log(x)} dx = \int x^{-\log(x)} dx$$

input `integrate(x**(-ln(x)),x)`

output `Integral(x**(-log(x)), x)`

3.307.7 Maxima [F]

$$\int x^{-\log(x)} dx = \int \frac{1}{x^{\log(x)}} dx$$

input `integrate(x^(-log(x)),x, algorithm="maxima")`

output `integrate(1/(x^log(x)), x)`

3.307.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.52

$$\int x^{-\log(x)} dx = \frac{1}{2} \sqrt{\pi} \operatorname{erf} \left(\log(x) - \frac{1}{2} \right) e^{\frac{1}{4}}$$

input `integrate(x^(-log(x)),x, algorithm="giac")`

output `1/2*sqrt(pi)*erf(log(x) - 1/2)*e^(1/4)`

3.307.9 Mupad [F(-1)]

Timed out.

$$\int x^{-\log(x)} dx = \int e^{-\ln(x)^2} dx$$

input `int(1/x^log(x), x)`output `int(exp(-log(x)^2), x)`

$$\mathbf{3.308} \quad \int \frac{1-2x}{x^{2/3}(1+x)^2} dx$$

3.308.1 Optimal result	1536
3.308.2 Mathematica [A] (verified)	1536
3.308.3 Rubi [A] (verified)	1537
3.308.4 Maple [A] (verified)	1538
3.308.5 Fricas [A] (verification not implemented)	1538
3.308.6 Sympy [A] (verification not implemented)	1539
3.308.7 Maxima [A] (verification not implemented)	1539
3.308.8 Giac [A] (verification not implemented)	1539
3.308.9 Mupad [B] (verification not implemented)	1540

3.308.1 Optimal result

Integrand size = 16, antiderivative size = 12

$$\int \frac{1-2x}{x^{2/3}(1+x)^2} dx = \frac{3\sqrt[3]{x}}{1+x}$$

output `3*x^(1/3)/(1+x)`

3.308.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1-2x}{x^{2/3}(1+x)^2} dx = \frac{3\sqrt[3]{x}}{1+x}$$

input `Integrate[(1 - 2*x)/(x^(2/3)*(1 + x)^2), x]`

output `(3*x^(1/3))/(1 + x)`

3.308.3 Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.062$, Rules used = {83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1-2x}{x^{2/3}(x+1)^2} dx$$

↓ 83

$$\frac{3\sqrt[3]{x}}{x+1}$$

input `Int[(1 - 2*x)/(x^(2/3)*(1 + x)^2),x]`

output `(3*x^(1/3))/(1 + x)`

3.308.3.1 Defintions of rubi rules used

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

3.308.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result
gospers	$\frac{3x^{\frac{1}{3}}}{1+x}$
trager	$\frac{3x^{\frac{1}{3}}}{1+x}$
risch	$\frac{3x^{\frac{1}{3}}}{1+x}$
derivativedivides	$-\frac{-x^{\frac{1}{3}}-1}{x^{\frac{2}{3}}-x^{\frac{1}{3}}+1} - \frac{1}{1+x^{\frac{1}{3}}}$
default	$-\frac{-x^{\frac{1}{3}}-1}{x^{\frac{2}{3}}-x^{\frac{1}{3}}+1} - \frac{1}{1+x^{\frac{1}{3}}}$
meijerg	$\frac{3x^{\frac{1}{3}}}{3+3x} + \frac{2\ln(1+x^{\frac{1}{3}})}{3} - \frac{\ln(x^{\frac{2}{3}}-x^{\frac{1}{3}}+1)}{3} + \frac{2\sqrt{3}\arctan\left(\frac{\sqrt{3}x^{\frac{1}{3}}}{2-x^{\frac{1}{3}}}\right)}{3} + \frac{2x^{\frac{1}{3}}}{1+x} - \frac{2x^{\frac{1}{3}}\left(\frac{\ln(1+x^{\frac{1}{3}})}{x^{\frac{1}{3}}} - \frac{\ln(x^{\frac{2}{3}}-x^{\frac{1}{3}}+1)}{2x^{\frac{1}{3}}}\right)}{3}$

input `int((1-2*x)/(1+x)^2/x^(2/3),x,method=_RETURNVERBOSE)`output `3*x^(1/3)/(1+x)`**3.308.5 Fracas [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1-2x}{x^{2/3}(1+x)^2} dx = \frac{3x^{\frac{1}{3}}}{x+1}$$

input `integrate((1-2*x)/(1+x)^2/x^(2/3),x, algorithm="fricas")`output `3*x^(1/3)/(x + 1)`

3.308.6 Sympy [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1-2x}{x^{2/3}(1+x)^2} dx = \frac{3\sqrt[3]{x}}{x+1}$$

input `integrate((1-2*x)/(1+x)**2/x**(2/3),x)`output `3*x**(1/3)/(x + 1)`**3.308.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1-2x}{x^{2/3}(1+x)^2} dx = \frac{3x^{1/3}}{x+1}$$

input `integrate((1-2*x)/(1+x)^2/x^(2/3),x, algorithm="maxima")`output `3*x^(1/3)/(x + 1)`**3.308.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1-2x}{x^{2/3}(1+x)^2} dx = \frac{3x^{1/3}}{x+1}$$

input `integrate((1-2*x)/(1+x)^2/x^(2/3),x, algorithm="giac")`output `3*x^(1/3)/(x + 1)`

3.308.9 Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1 - 2x}{x^{2/3}(1+x)^2} dx = \frac{3x^{1/3}}{x+1}$$

input `int(-(2*x - 1)/(x^(2/3)*(x + 1)^2),x)`

output `(3*x^(1/3))/(x + 1)`

3.309 $\int \cos^2\left(\frac{\pi x^2}{\sqrt{2}}\right) dx$

3.309.1 Optimal result	1541
3.309.2 Mathematica [A] (verified)	1541
3.309.3 Rubi [A] (verified)	1542
3.309.4 Maple [B] (verified)	1543
3.309.5 Fricas [A] (verification not implemented)	1543
3.309.6 Sympy [A] (verification not implemented)	1543
3.309.7 Maxima [C] (verification not implemented)	1544
3.309.8 Giac [C] (verification not implemented)	1544
3.309.9 Mupad [F(-1)]	1545

3.309.1 Optimal result

Integrand size = 13, antiderivative size = 23

$$\int \cos^2\left(\frac{\pi x^2}{\sqrt{2}}\right) dx = \frac{x}{2} + \frac{\text{FresnelC}\left(2^{3/4}x\right)}{2 \cdot 2^{3/4}}$$

output `1/2*x+1/4*FresnelC(2^(3/4)*x)*2^(1/4)`

3.309.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int \cos^2\left(\frac{\pi x^2}{\sqrt{2}}\right) dx = \frac{1}{4}\left(2x + \sqrt[4]{2} \text{FresnelC}\left(2^{3/4}x\right)\right)$$

input `Integrate[Cos[(Pi*x^2)/Sqrt[2]]^2,x]`

output `(2*x + 2^(1/4)*FresnelC[2^(3/4)*x])/4`

3.309.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3839, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^2\left(\frac{\pi x^2}{\sqrt{2}}\right) dx$$

↓ 3839

$$\int \left(\frac{1}{2} \cos(\sqrt{2}\pi x^2) + \frac{1}{2}\right) dx$$

↓ 2009

$$\frac{\text{FresnelC}(2^{3/4}x)}{2 \cdot 2^{3/4}} + \frac{x}{2}$$

input `Int[Cos[(Pi*x^2)/Sqrt[2]]^2,x]`

output `x/2 + FresnelC[2^(3/4)*x]/(2*2^(3/4))`

3.309.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3839 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)]*(b_.))^(p_), x_Symbol] :> Int[ExpandTrigReduce[(a + b*Cos[c + d*(e + f*x)^n])^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 1] && IGtQ[n, 1]`

3.309.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 33 vs. $2(15) = 30$.

Time = 0.18 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.48

method	result	size
default	$\frac{x}{2} + \frac{\sqrt{2}\sqrt{\pi} \operatorname{FresnelC}\left(\frac{2\sqrt{\pi}x}{\sqrt{\pi\sqrt{2}}}\right)}{4\sqrt{\pi\sqrt{2}}}$	34
risch	$\frac{x}{2} + \frac{\sqrt{\pi} \operatorname{erf}\left(\sqrt{i\pi\sqrt{2}}x\right)}{8\sqrt{i\pi\sqrt{2}}} + \frac{\sqrt{\pi} \operatorname{erf}\left(\sqrt{-i\pi\sqrt{2}}x\right)}{8\sqrt{-i\pi\sqrt{2}}}$	57

input `int(cos(1/2*Pi*x^2*2^(1/2))^2,x,method=_RETURNVERBOSE)`

output `1/2*x+1/4*2^(1/2)*Pi^(1/2)/(Pi*2^(1/2))^(1/2)*FresnelC(2*Pi^(1/2)/(Pi*2^(1/2))^(1/2)*x)`

3.309.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.65

$$\int \cos^2\left(\frac{\pi x^2}{\sqrt{2}}\right) dx = \frac{1}{4} \cdot 2^{\frac{1}{4}} C\left(2^{\frac{3}{4}}x\right) + \frac{1}{2}x$$

input `integrate(cos(1/2*pi*x^2*2^(1/2))^2,x, algorithm="fracas")`

output `1/4*2^(1/4)*fresnel_cos(2^(3/4)*x) + 1/2*x`

3.309.6 Sympy [A] (verification not implemented)

Time = 1.34 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \cos^2\left(\frac{\pi x^2}{\sqrt{2}}\right) dx = \frac{x}{2} + \frac{\sqrt[4]{2}C\left(2^{\frac{3}{4}}x\right)\Gamma\left(\frac{1}{4}\right)}{16\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(cos(1/2*pi*x**2*2**(1/2))**2,x)`

output `x/2 + 2**(1/4)*fresnelc(2**(3/4)*x)*gamma(1/4)/(16*gamma(5/4))`

3.309. $\int \cos^2\left(\frac{\pi x^2}{\sqrt{2}}\right) dx$

3.309.7 Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 46, normalized size of antiderivative = 2.00

$$\int \cos^2\left(\frac{\pi x^2}{\sqrt{2}}\right) dx = -\frac{2^{\frac{1}{4}}\pi^2\left((i-1)\operatorname{erf}\left(\sqrt{i}\sqrt{2}\pi x\right) - (i+1)\operatorname{erf}\left(\sqrt{-i}\sqrt{2}\pi x\right)\right) - 8\pi^2x}{16\pi^2}$$

input `integrate(cos(1/2*pi*x^2*2^(1/2))^2,x, algorithm="maxima")`

output `-1/16*(2^(1/4)*pi^2*((I - 1)*erf(sqrt(I*sqrt(2)*pi)*x) - (I + 1)*erf(sqrt(-I*sqrt(2)*pi)*x)) - 8*pi^2*x)/pi^2`

3.309.8 Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.83

$$\int \cos^2\left(\frac{\pi x^2}{\sqrt{2}}\right) dx = -\left(\frac{1}{16}i + \frac{1}{16}\right) \cdot 2^{\frac{1}{4}} \operatorname{erf}\left(\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2}\sqrt{\sqrt{2}\pi x}\right) + \left(\frac{1}{16}i - \frac{1}{16}\right) \cdot 2^{\frac{1}{4}} \operatorname{erf}\left(-\left(\frac{1}{2}i + \frac{1}{2}\right) \sqrt{2}\sqrt{\sqrt{2}\pi x}\right) + \frac{1}{2}x$$

input `integrate(cos(1/2*pi*x^2*2^(1/2))^2,x, algorithm="giac")`

output `-(1/16*I + 1/16)*2^(1/4)*erf((1/2*I - 1/2)*sqrt(2)*sqrt(sqrt(2)*pi)*x) + (1/16*I - 1/16)*2^(1/4)*erf(-(1/2*I + 1/2)*sqrt(2)*sqrt(sqrt(2)*pi)*x) + 1/2*x`

3.309.9 Mupad [F(-1)]

Timed out.

$$\int \cos^2\left(\frac{\pi x^2}{\sqrt{2}}\right) dx = \int \cos\left(\frac{\sqrt{2}\pi x^2}{2}\right)^2 dx$$

input `int(cos((2^(1/2)*Pi*x^2)/2)^2,x)`output `int(cos((2^(1/2)*Pi*x^2)/2)^2, x)`

3.310 $\int \frac{1}{1+\cos(x)+\sin(x)} dx$

3.310.1 Optimal result	1546
3.310.2 Mathematica [B] (verified)	1546
3.310.3 Rubi [A] (verified)	1547
3.310.4 Maple [A] (verified)	1548
3.310.5 Fricas [B] (verification not implemented)	1548
3.310.6 Sympy [A] (verification not implemented)	1548
3.310.7 Maxima [A] (verification not implemented)	1549
3.310.8 Giac [A] (verification not implemented)	1549
3.310.9 Mupad [B] (verification not implemented)	1549

3.310.1 Optimal result

Integrand size = 8, antiderivative size = 9

$$\int \frac{1}{1 + \cos(x) + \sin(x)} dx = \log \left(1 + \tan \left(\frac{x}{2} \right) \right)$$

output `ln(1+tan(1/2*x))`

3.310.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 24 vs. 2(9) = 18.

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 2.67

$$\int \frac{1}{1 + \cos(x) + \sin(x)} dx = -\log \left(\cos \left(\frac{x}{2} \right) \right) + \log \left(\cos \left(\frac{x}{2} \right) + \sin \left(\frac{x}{2} \right) \right)$$

input `Integrate[(1 + Cos[x] + Sin[x])^(-1),x]`

output `-Log[Cos[x/2]] + Log[Cos[x/2] + Sin[x/2]]`

3.310.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3042, 3603, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sin(x) + \cos(x) + 1} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{\sin(x) + \cos(x) + 1} dx \\ & \quad \downarrow \text{3603} \\ & 2 \int \frac{1}{2 \tan\left(\frac{x}{2}\right) + 2} d \tan\left(\frac{x}{2}\right) \\ & \quad \downarrow \text{16} \\ & \log\left(\tan\left(\frac{x}{2}\right) + 1\right) \end{aligned}$$

input `Int[(1 + Cos[x] + Sin[x])^(-1),x]`

output `Log[1 + Tan[x/2]]`

3.310.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_.)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3603 `Int[(cos[(d_.) + (e_.)*(x_.)]*(b_.) + (a_.) + (c_.)*sin[(d_.) + (e_.)*(x_.)])^(-1), x_Symbol] := Module[{f = FreeFactors[Tan[(d + e*x)/2], x]}, Simp[2*(f/e) Subst[Int[1/(a + b + 2*c*f*x + (a - b)*f^2*x^2), x], x, Tan[(d + e*x)/2]/f], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[a^2 - b^2 - c^2, 0]`

3.310.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

method	result	size
default	$\ln\left(1 + \tan\left(\frac{x}{2}\right)\right)$	8
parallelrisch	$\ln\left(1 + \tan\left(\frac{x}{2}\right)\right)$	8
norman	$\ln\left(2 + 2 \tan\left(\frac{x}{2}\right)\right)$	10
risch	$\ln(i + e^{ix}) - \ln(e^{ix} + 1)$	21

input `int(1/(1+cos(x)+sin(x)),x,method=_RETURNVERBOSE)`

output `ln(1+tan(1/2*x))`

3.310.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(7) = 14$.

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.89

$$\int \frac{1}{1 + \cos(x) + \sin(x)} dx = -\frac{1}{2} \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) + \frac{1}{2} \log(\sin(x) + 1)$$

input `integrate(1/(1+cos(x)+sin(x)),x, algorithm="fracas")`

output `-1/2*log(1/2*cos(x) + 1/2) + 1/2*log(sin(x) + 1)`

3.310.6 Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{1}{1 + \cos(x) + \sin(x)} dx = \log\left(\tan\left(\frac{x}{2}\right) + 1\right)$$

input `integrate(1/(1+cos(x)+sin(x)),x)`

output `log(tan(x/2) + 1)`

3.310.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.33

$$\int \frac{1}{1 + \cos(x) + \sin(x)} dx = \log \left(\frac{\sin(x)}{\cos(x) + 1} + 1 \right)$$

input `integrate(1/(1+cos(x)+sin(x)),x, algorithm="maxima")`output `log(sin(x)/(cos(x) + 1) + 1)`**3.310.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int \frac{1}{1 + \cos(x) + \sin(x)} dx = \log \left(\left| \tan \left(\frac{1}{2} x \right) + 1 \right| \right)$$

input `integrate(1/(1+cos(x)+sin(x)),x, algorithm="giac")`output `log(abs(tan(1/2*x) + 1))`**3.310.9 Mupad [B] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.78

$$\int \frac{1}{1 + \cos(x) + \sin(x)} dx = \ln \left(\tan \left(\frac{x}{2} \right) + 1 \right)$$

input `int(1/(cos(x) + sin(x) + 1),x)`output `log(tan(x/2) + 1)`

3.311 $\int \tan^5(x) dx$

3.311.1 Optimal result	1550
3.311.2 Mathematica [A] (verified)	1550
3.311.3 Rubi [A] (verified)	1551
3.311.4 Maple [A] (verified)	1552
3.311.5 Fricas [A] (verification not implemented)	1553
3.311.6 Sympy [A] (verification not implemented)	1553
3.311.7 Maxima [A] (verification not implemented)	1553
3.311.8 Giac [A] (verification not implemented)	1554
3.311.9 Mupad [B] (verification not implemented)	1554

3.311.1 Optimal result

Integrand size = 4, antiderivative size = 22

$$\int \tan^5(x) dx = -\log(\cos(x)) - \frac{\tan^2(x)}{2} + \frac{\tan^4(x)}{4}$$

output `-ln(cos(x))-1/2*tan(x)^2+1/4*tan(x)^4`

3.311.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \tan^5(x) dx = -\log(\cos(x)) - \frac{\tan^2(x)}{2} + \frac{\tan^4(x)}{4}$$

input `Integrate[Tan[x]^5,x]`

output `-Log[Cos[x]] - Tan[x]^2/2 + Tan[x]^4/4`

3.311.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {3042, 3954, 3042, 3954, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^5(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x)^5 dx \\
 & \quad \downarrow \text{3954} \\
 & \frac{\tan^4(x)}{4} - \int \tan^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{\tan^4(x)}{4} - \int \tan(x)^3 dx \\
 & \quad \downarrow \text{3954} \\
 & \int \tan(x) dx + \frac{\tan^4(x)}{4} - \frac{\tan^2(x)}{2} \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(x) dx + \frac{\tan^4(x)}{4} - \frac{\tan^2(x)}{2} \\
 & \quad \downarrow \text{3956} \\
 & \frac{\tan^4(x)}{4} - \frac{\tan^2(x)}{2} - \log(\cos(x))
 \end{aligned}$$

input `Int [Tan [x] ^5, x]`

output `-Log [Cos [x]] - Tan [x] ^2/2 + Tan [x] ^4/4`

3.311.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

3.311.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.05

method	result	size
derivativedivides	$\frac{\tan(x)^4}{4} - \frac{\tan(x)^2}{2} + \frac{\ln(1+\tan(x)^2)}{2}$	23
default	$\frac{\tan(x)^4}{4} - \frac{\tan(x)^2}{2} + \frac{\ln(1+\tan(x)^2)}{2}$	23
norman	$\frac{\tan(x)^4}{4} - \frac{\tan(x)^2}{2} + \frac{\ln(1+\tan(x)^2)}{2}$	23
parallelrisch	$\frac{\tan(x)^4}{4} - \frac{\tan(x)^2}{2} + \frac{\ln(1+\tan(x)^2)}{2}$	23
risch	$i x - \frac{4(e^{6ix} + e^{4ix} + e^{2ix})}{(e^{2ix} + 1)^4} - \ln(e^{2ix} + 1)$	43

input `int(tan(x)^5,x,method=_RETURNVERBOSE)`

output `1/4*tan(x)^4-1/2*tan(x)^2+1/2*ln(1+tan(x)^2)`

3.311.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \tan^5(x) dx = \frac{1}{4} \tan(x)^4 - \frac{1}{2} \tan(x)^2 - \frac{1}{2} \log\left(\frac{1}{\tan(x)^2 + 1}\right)$$

input `integrate(tan(x)^5,x, algorithm="fricas")`output `1/4*tan(x)^4 - 1/2*tan(x)^2 - 1/2*log(1/(tan(x)^2 + 1))`**3.311.6 Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \tan^5(x) dx = -\frac{4 \cos^2(x) - 1}{4 \cos^4(x)} - \log(\cos(x))$$

input `integrate(tan(x)**5,x)`output `-(4*cos(x)**2 - 1)/(4*cos(x)**4) - log(cos(x))`**3.311.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.55

$$\int \tan^5(x) dx = \frac{4 \sin(x)^2 - 3}{4 (\sin(x)^4 - 2 \sin(x)^2 + 1)} - \frac{1}{2} \log(\sin(x)^2 - 1)$$

input `integrate(tan(x)^5,x, algorithm="maxima")`output `1/4*(4*sin(x)^2 - 3)/(sin(x)^4 - 2*sin(x)^2 + 1) - 1/2*log(sin(x)^2 - 1)`

3.311.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \tan^5(x) dx = \frac{1}{4} \tan(x)^4 - \frac{1}{2} \tan(x)^2 + \frac{1}{2} \log(\tan(x)^2 + 1)$$

input `integrate(tan(x)^5,x, algorithm="giac")`

output `1/4*tan(x)^4 - 1/2*tan(x)^2 + 1/2*log(tan(x)^2 + 1)`

3.311.9 Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \tan^5(x) dx = \frac{\tan(x)^4}{4} - \frac{\tan(x)^2}{2} - \ln(\cos(x))$$

input `int(tan(x)^5,x)`

output `tan(x)^4/4 - tan(x)^2/2 - log(cos(x))`

$$\mathbf{3.312} \quad \int \sqrt{1 + \frac{1}{x}} dx$$

3.312.1 Optimal result	1555
3.312.2 Mathematica [A] (verified)	1555
3.312.3 Rubi [A] (verified)	1556
3.312.4 Maple [B] (verified)	1557
3.312.5 Fricas [B] (verification not implemented)	1558
3.312.6 Sympy [A] (verification not implemented)	1558
3.312.7 Maxima [A] (verification not implemented)	1558
3.312.8 Giac [A] (verification not implemented)	1559
3.312.9 Mupad [B] (verification not implemented)	1559

3.312.1 Optimal result

Integrand size = 9, antiderivative size = 22

$$\int \sqrt{1 + \frac{1}{x}} dx = \sqrt{1 + \frac{1}{x}}x + \operatorname{arctanh}\left(\sqrt{1 + \frac{1}{x}}\right)$$

output `(1+1/x)^(1/2)*x+arctanh((1+1/x)^(1/2))`

3.312.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \sqrt{1 + \frac{1}{x}} dx = \sqrt{1 + \frac{1}{x}}x + \operatorname{arctanh}\left(\sqrt{1 + \frac{1}{x}}\right)$$

input `Integrate[Sqrt[1 + x^(-1)],x]`

output `Sqrt[1 + x^(-1)]*x + ArcTanh[Sqrt[1 + x^(-1)]]`

3.312.3 Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {773, 51, 73, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\frac{1}{x} + 1} dx \\
 & \quad \downarrow \text{773} \\
 & - \int \sqrt{1 + \frac{1}{x}} x^2 d\frac{1}{x} \\
 & \quad \downarrow \text{51} \\
 & \sqrt{\frac{1}{x} + 1} x - \frac{1}{2} \int \frac{x}{\sqrt{1 + \frac{1}{x}}} d\frac{1}{x} \\
 & \quad \downarrow \text{73} \\
 & \sqrt{\frac{1}{x} + 1} x - \int \frac{1}{\frac{1}{x^2} - 1} d\sqrt{1 + \frac{1}{x}} \\
 & \quad \downarrow \text{220} \\
 & \operatorname{arctanh}\left(\sqrt{\frac{1}{x} + 1}\right) + \sqrt{\frac{1}{x} + 1} x
 \end{aligned}$$

input `Int[Sqrt[1 + x^(-1)], x]`

output `Sqrt[1 + x^(-1)]*x + ArcTanh[Sqrt[1 + x^(-1)]]`

3.312.3.1 Defintions of rubi rules used

```
rule 51 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x
] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 220 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-
1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] &&
(LtQ[a, 0] || GtQ[b, 0])
```

```
rule 773 Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^
2, x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0] && !IntegerQ[p]
```

3.312.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 38 vs. $2(18) = 36$.

Time = 0.11 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.77

method	result	size
trager	$\sqrt{-\frac{-1-x}{x}} x - \frac{\ln\left(2\sqrt{-\frac{-1-x}{x}} x - 2x - 1\right)}{2}$	39
default	$\frac{\sqrt{\frac{1+x}{x}} x \left(2\sqrt{x^2+x} + \ln\left(\frac{1}{2} + x + \sqrt{x^2+x}\right)\right)}{2\sqrt{x(1+x)}}$	41
risch	$x\sqrt{\frac{1+x}{x}} + \frac{\ln\left(\frac{1}{2} + x + \sqrt{x^2+x}\right)\sqrt{\frac{1+x}{x}}\sqrt{x(1+x)}}{2+2x}$	47

```
input int((1+1/x)^(1/2), x, method=_RETURNVERBOSE)
```

```
output (-(-1-x)/x)^(1/2)*x-1/2*ln(2*(-(-1-x)/x)^(1/2)*x-2*x-1)
```

3.312.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 40 vs. $2(18) = 36$.

Time = 0.24 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.82

$$\int \sqrt{1 + \frac{1}{x}} dx = x\sqrt{\frac{x+1}{x}} + \frac{1}{2} \log\left(\sqrt{\frac{x+1}{x}} + 1\right) - \frac{1}{2} \log\left(\sqrt{\frac{x+1}{x}} - 1\right)$$

input `integrate((1+1/x)^(1/2),x, algorithm="fricas")`

output `x*sqrt((x + 1)/x) + 1/2*log(sqrt((x + 1)/x) + 1) - 1/2*log(sqrt((x + 1)/x) - 1)`

3.312.6 Sympy [A] (verification not implemented)

Time = 0.76 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

$$\int \sqrt{1 + \frac{1}{x}} dx = \sqrt{x}\sqrt{x+1} + \operatorname{asinh}(\sqrt{x})$$

input `integrate((1+1/x)**(1/2),x)`

output `sqrt(x)*sqrt(x + 1) + asinh(sqrt(x))`

3.312.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.55

$$\int \sqrt{1 + \frac{1}{x}} dx = x\sqrt{\frac{1}{x} + 1} + \frac{1}{2} \log\left(\sqrt{\frac{1}{x} + 1} + 1\right) - \frac{1}{2} \log\left(\sqrt{\frac{1}{x} + 1} - 1\right)$$

input `integrate((1+1/x)^(1/2),x, algorithm="maxima")`

output `x*sqrt(1/x + 1) + 1/2*log(sqrt(1/x + 1) + 1) - 1/2*log(sqrt(1/x + 1) - 1)`

3.312.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.41

$$\int \sqrt{1 + \frac{1}{x}} dx = -\frac{1}{2} \log \left(\left| -2x + 2\sqrt{x^2 + x} - 1 \right| \right) \operatorname{sgn}(x) + \sqrt{x^2 + x} \operatorname{sgn}(x)$$

input `integrate((1+1/x)^(1/2),x, algorithm="giac")`output `-1/2*log(abs(-2*x + 2*sqrt(x^2 + x) - 1))*sgn(x) + sqrt(x^2 + x)*sgn(x)`**3.312.9 Mupad [B] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.73

$$\int \sqrt{1 + \frac{1}{x}} dx = x \sqrt{\frac{1}{x} + 1} + \frac{x \ln \left(x + \sqrt{x^2 + x} + \frac{1}{2} \right) \sqrt{\frac{1}{x} + 1}}{2\sqrt{x^2 + x}}$$

input `int((1/x + 1)^(1/2),x)`output `x*(1/x + 1)^(1/2) + (x*log(x + (x + x^2)^(1/2) + 1/2)*(1/x + 1)^(1/2))/(2*(x + x^2)^(1/2))`

3.313 $\int e^{\cos(x)} \cos(2x + \sin(x)) dx$

3.313.1 Optimal result	1560
3.313.2 Mathematica [A] (verified)	1560
3.313.3 Rubi [F]	1561
3.313.4 Maple [C] (verified)	1561
3.313.5 Fricas [B] (verification not implemented)	1562
3.313.6 Sympy [F]	1562
3.313.7 Maxima [F]	1563
3.313.8 Giac [B] (verification not implemented)	1563
3.313.9 Mupad [B] (verification not implemented)	1563

3.313.1 Optimal result

Integrand size = 12, antiderivative size = 21

$$\int e^{\cos(x)} \cos(2x + \sin(x)) dx = 2e^{\cos(x)} \cos\left(\frac{x}{2} + \sin(x)\right) \sin\left(\frac{x}{2}\right)$$

output `2*exp(cos(x))*cos(1/2*x+sin(x))*sin(1/2*x)`

3.313.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int e^{\cos(x)} \cos(2x + \sin(x)) dx = 2e^{\cos(x)} \cos\left(\frac{x}{2} + \sin(x)\right) \sin\left(\frac{x}{2}\right)$$

input `Integrate[E^Cos[x]*Cos[2*x + Sin[x]],x]`

output `2*E^Cos[x]*Cos[x/2 + Sin[x]]*Sin[x/2]`

3.313.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{\cos(x)} \cos(2x + \sin(x)) dx$$

↓ 7299

$$\int e^{\cos(x)} \cos(2x + \sin(x)) dx$$

input `Int[E^Cos[x]*Cos[2*x + Sin[x]],x]`

output `$Aborted`

3.313.3.1 Defintions of rubi rules used

rule 7299 `Int[u_, x_] :> CannotIntegrate[u, x]`

3.313.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.17 (sec) , antiderivative size = 52, normalized size of antiderivative = 2.48

method	result	size
risch	$-\frac{ie^{ix}e^{ix}}{2} + \frac{ie^{ix}}{2} + \frac{ie^{-ix}e^{-ix}}{2} - \frac{ie^{-ix}}{2}$	52

input `int(exp(cos(x))*cos(2*x+sin(x)),x,method=_RETURNVERBOSE)`

output `-1/2*I*exp(I*x)*exp(exp(I*x))+1/2*I*exp(exp(I*x))+1/2*I*exp(1/exp(I*x))*exp(-I*x)-1/2*I*exp(1/exp(I*x))`

3.313.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 84 vs. $2(16) = 32$.

Time = 0.26 (sec) , antiderivative size = 84, normalized size of antiderivative = 4.00

$$\begin{aligned} & \int e^{\cos(x)} \cos(2x + \sin(x)) dx \\ &= (2 \cos(x) - 1) \cos\left(\frac{2\left(x \tan\left(\frac{1}{2}x\right)^2 + x + \tan\left(\frac{1}{2}x\right)\right)}{\tan\left(\frac{1}{2}x\right)^2 + 1}\right) e^{\cos(x)} \sin(x) \\ &\quad - (2 \cos(x)^2 - \cos(x) - 1) e^{\cos(x)} \sin\left(\frac{2\left(x \tan\left(\frac{1}{2}x\right)^2 + x + \tan\left(\frac{1}{2}x\right)\right)}{\tan\left(\frac{1}{2}x\right)^2 + 1}\right) \end{aligned}$$

input `integrate(exp(cos(x))*cos(2*x+sin(x)),x, algorithm="fricas")`

output `(2*cos(x) - 1)*cos(2*(x*tan(1/2*x)^2 + x + tan(1/2*x))/(tan(1/2*x)^2 + 1)) *e^cos(x)*sin(x) - (2*cos(x)^2 - cos(x) - 1)*e^cos(x)*sin(2*(x*tan(1/2*x)^2 + x + tan(1/2*x))/(tan(1/2*x)^2 + 1))`

3.313.6 Sympy [F]

$$\int e^{\cos(x)} \cos(2x + \sin(x)) dx = \int e^{\cos(x)} \cos(2x + \sin(x)) dx$$

input `integrate(exp(cos(x))*cos(2*x+sin(x)),x)`

output `Integral(exp(cos(x))*cos(2*x + sin(x)), x)`

3.313.7 Maxima [F]

$$\int e^{\cos(x)} \cos(2x + \sin(x)) dx = \int \cos(2x + \sin(x)) e^{\cos(x)} dx$$

input `integrate(exp(cos(x))*cos(2*x+sin(x)),x, algorithm="maxima")`

output `1/4*e^cos(x)*sin(2*x + sin(x)) + 1/2*e^cos(x)*sin(x + sin(x)) - 1/2*e^cos(x)*sin(sin(x)) - 1/4*integrate(cos(3*x + sin(x))*e^cos(x), x)`

3.313.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 136 vs. $2(16) = 32$.

Time = 0.28 (sec) , antiderivative size = 136, normalized size of antiderivative = 6.48

$$\int e^{\cos(x)} \cos(2x + \sin(x)) dx$$

$$= \frac{\cos(x)^3 e^{\cos(x)} \sin(2x + \sin(x)) - \cos(2x + \sin(x)) \cos(x)^2 e^{\cos(x)} \sin(x) + \cos(x) e^{\cos(x)} \sin(2x + \sin(x))}{\cos(x)^4 + 2\cos(x)^2 \sin(x)^2 + \sin(x)^4}$$

input `integrate(exp(cos(x))*cos(2*x+sin(x)),x, algorithm="giac")`

output `(cos(x)^3*e^cos(x)*sin(2*x + sin(x)) - cos(2*x + sin(x))*cos(x)^2*e^cos(x)*sin(x) + cos(x)*e^cos(x)*sin(2*x + sin(x))*sin(x)^2 - cos(2*x + sin(x))*e^cos(x)*sin(x)^3 - cos(x)^2*e^cos(x)*sin(2*x + sin(x)) + 2*cos(2*x + sin(x))*cos(x)*e^cos(x)*sin(x) + e^cos(x)*sin(2*x + sin(x))*sin(x)^2)/(cos(x)^4 + 2*cos(x)^2*sin(x)^2 + sin(x)^4)`

3.313.9 Mupad [B] (verification not implemented)

Time = 17.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int e^{\cos(x)} \cos(2x + \sin(x)) dx = e^{\cos(x)} (\sin(x + \sin(x)) - \sin(\sin(x)))$$

input `int(exp(cos(x))*cos(2*x + sin(x)),x)`

output `exp(cos(x))*(sin(x + sin(x)) - sin(sin(x)))`

$$\mathbf{3.314} \quad \int \frac{-1+2x+3\log(x)}{x^2+2x^4+x\log(x)} dx$$

3.314.1 Optimal result	1564
3.314.2 Mathematica [A] (verified)	1564
3.314.3 Rubi [A] (verified)	1565
3.314.4 Maple [A] (verified)	1566
3.314.5 Fricas [A] (verification not implemented)	1566
3.314.6 Sympy [A] (verification not implemented)	1566
3.314.7 Maxima [A] (verification not implemented)	1567
3.314.8 Giac [A] (verification not implemented)	1567
3.314.9 Mupad [B] (verification not implemented)	1567

3.314.1 Optimal result

Integrand size = 25, antiderivative size = 17

$$\int \frac{-1+2x+3\log(x)}{x^2+2x^4+x\log(x)} dx = 3\log(x) - \log(x+2x^3+\log(x))$$

output `3*ln(x)-ln(x+2*x^3+ln(x))`

3.314.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{-1+2x+3\log(x)}{x^2+2x^4+x\log(x)} dx = 3\log(x) - \log(x+2x^3+\log(x))$$

input `Integrate[(-1 + 2*x + 3*Log[x])/(x^2 + 2*x^4 + x*Log[x]),x]`

output `3*Log[x] - Log[x + 2*x^3 + Log[x]]`

3.314.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2x + 3 \log(x) - 1}{2x^4 + x^2 + x \log(x)} dx$$

↓ 7293

$$\int \left(\frac{-6x^3 - x - 1}{x(2x^3 + x + \log(x))} + \frac{3}{x} \right) dx$$

↓ 2009

$$3 \log(x) - \log(2x^3 + x + \log(x))$$

input `Int[(-1 + 2*x + 3*Log[x])/(x^2 + 2*x^4 + x*Log[x]),x]`

output `3*Log[x] - Log[x + 2*x^3 + Log[x]]`

3.314.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] :> With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

3.314.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

method	result	size
default	$3 \ln(x) - \ln(x + 2x^3 + \ln(x))$	18
norman	$3 \ln(x) - \ln(x + 2x^3 + \ln(x))$	18
risch	$3 \ln(x) - \ln(x + 2x^3 + \ln(x))$	18
parallelrisch	$-\ln\left(x^3 + \frac{x}{2} + \frac{\ln(x)}{2}\right) + 3 \ln(x)$	20

input `int((3*ln(x)+2*x-1)/(x*ln(x)+x^2+2*x^4),x,method=_RETURNVERBOSE)`output `3*ln(x)-ln(x+2*x^3+ln(x))`**3.314.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{-1 + 2x + 3 \log(x)}{x^2 + 2x^4 + x \log(x)} dx = -\log(2x^3 + x + \log(x)) + 3 \log(x)$$

input `integrate((3*log(x)+2*x-1)/(x*log(x)+x^2+2*x^4),x, algorithm="fricas")`output `-log(2*x^3 + x + log(x)) + 3*log(x)`**3.314.6 Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{-1 + 2x + 3 \log(x)}{x^2 + 2x^4 + x \log(x)} dx = 3 \log(x) - \log(2x^3 + x + \log(x))$$

input `integrate((3*ln(x)+2*x-1)/(x*ln(x)+x**2+2*x**4),x)`output `3*log(x) - log(2*x**3 + x + log(x))`

3.314.7 Maxima [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{-1 + 2x + 3 \log(x)}{x^2 + 2x^4 + x \log(x)} dx = -\log(2x^3 + x + \log(x)) + 3 \log(x)$$

input `integrate((3*log(x)+2*x-1)/(x*log(x)+x^2+2*x^4),x, algorithm="maxima")`output `-log(2*x^3 + x + log(x)) + 3*log(x)`**3.314.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.24

$$\int \frac{-1 + 2x + 3 \log(x)}{x^2 + 2x^4 + x \log(x)} dx = -\log(-2x^3 - x - \log(x)) + 3 \log(x)$$

input `integrate((3*log(x)+2*x-1)/(x*log(x)+x^2+2*x^4),x, algorithm="giac")`output `-log(-2*x^3 - x - log(x)) + 3*log(x)`**3.314.9 Mupad [B] (verification not implemented)**

Time = 17.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{-1 + 2x + 3 \log(x)}{x^2 + 2x^4 + x \log(x)} dx = 3 \ln(x) - \ln(x + \ln(x) + 2x^3)$$

input `int((2*x + 3*log(x) - 1)/(x*log(x) + x^2 + 2*x^4),x)`output `3*log(x) - log(x + log(x) + 2*x^3)`

3.315 $\int (-\sqrt{x} + \sqrt{1+x})^\pi dx$

3.315.1 Optimal result	1568
3.315.2 Mathematica [B] (verified)	1568
3.315.3 Rubi [A] (warning: unable to verify)	1569
3.315.4 Maple [F]	1570
3.315.5 Fricas [A] (verification not implemented)	1571
3.315.6 Sympy [B] (verification not implemented)	1571
3.315.7 Maxima [F]	1572
3.315.8 Giac [F]	1573
3.315.9 Mupad [F(-1)]	1573

3.315.1 Optimal result

Integrand size = 17, antiderivative size = 45

$$\int (-\sqrt{x} + \sqrt{1+x})^\pi dx = -\frac{2(-\sqrt{x} + \sqrt{1+x})^\pi (1 + 2x + \pi\sqrt{x}\sqrt{1+x})}{-4 + \pi^2}$$

output `-2*(-x^(1/2)+(1+x)^(1/2))^Pi*(1+2*x+Pi*x^(1/2)*(1+x)^(1/2))/(Pi^2-4)`

3.315.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 107 vs. 2(45) = 90.

Time = 0.54 (sec) , antiderivative size = 107, normalized size of antiderivative = 2.38

$$\int (-\sqrt{x} + \sqrt{1+x})^\pi dx = \frac{2\sqrt{1+x}(-\sqrt{x} + \sqrt{1+x})^{-1+\pi} (1 - 2(-2 + \pi)x - 2(-2 + \pi)x^2 + (-2 + \pi)\sqrt{x}\sqrt{1+x} + 2(-2 + \pi)x^{3/2})}{(-2 + \pi)(2 + \pi) (-1 - x + \sqrt{x}\sqrt{1+x})}$$

input `Integrate[(-Sqrt[x] + Sqrt[1 + x])^Pi,x]`

output `(2*Sqrt[1 + x]*(-Sqrt[x] + Sqrt[1 + x])^(-1 + Pi)*(1 - 2*(-2 + Pi)*x - 2*(-2 + Pi)*x^2 + (-2 + Pi)*Sqrt[x]*Sqrt[1 + x] + 2*(-2 + Pi)*x^(3/2)*Sqrt[1 + x]))/((-2 + Pi)*(2 + Pi)*(-1 - x + Sqrt[x]*Sqrt[1 + x]))`

3.315.3 Rubi [A] (warning: unable to verify)

Time = 0.28 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.82, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {7296, 2544, 25, 335, 802, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (\sqrt{x+1} - \sqrt{x})^\pi dx \\
 & \quad \downarrow \text{7296} \\
 & 2 \int \sqrt{x} (\sqrt{x+1} - \sqrt{x})^\pi d\sqrt{x} \\
 & \quad \downarrow \text{2544} \\
 & \frac{1}{2} \int -((1-x)x^{\frac{1}{2}(-3+\pi)}(x+1)) d(\sqrt{x+1} - \sqrt{x}) \\
 & \quad \downarrow \text{25} \\
 & -\frac{1}{2} \int (1-x)x^{\frac{1}{2}(-3+\pi)}(x+1) d(\sqrt{x+1} - \sqrt{x}) \\
 & \quad \downarrow \text{335} \\
 & -\frac{1}{2} \int x^{\frac{1}{2}(-3+\pi)}(1-x^2) d(\sqrt{x+1} - \sqrt{x}) \\
 & \quad \downarrow \text{802} \\
 & -\frac{1}{2} \int (x^{\frac{1}{2}(-3+\pi)} - x^{\frac{1+\pi}{2}}) d(\sqrt{x+1} - \sqrt{x}) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} \left(\frac{x^{\frac{1}{2}(\pi-2)}}{2-\pi} + \frac{x^{\frac{2+\pi}{2}}}{2+\pi} \right)
 \end{aligned}$$

input `Int[(-Sqrt[x] + Sqrt[1 + x])^Pi, x]`

output `(x^((-2 + Pi)/2)/(2 - Pi) + x^((2 + Pi)/2)/(2 + Pi))/2`

3.315.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 335 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] := Int[(e*x)^m*(a*c + b*d*x^4)^p, x] /; FreeQ[{a, b, c, d, e, m, p}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[c, 0]))`

rule 802 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2544 `Int[((g_.) + (h_.)*(x_))^(m_.)*((e_.)*(x_) + (f_.)*Sqrt[(a_.) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[1/(2^(m + 1)*e^(m + 1)) Subst[Int[x^(n - m - 2)*(a*f^2 + x^2)*((-a)*f^2*h + 2*e*g*x + h*x^2)^m, x], x, e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[m]`

rule 7296 `Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Simp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x] /; !FalseQ[lst]]`

3.315.4 Maple [F]

$$\int (\sqrt{1+x} - \sqrt{x})^\pi dx$$

input `int(((1+x)^(1/2)-x^(1/2))^Pi,x)`

output `int(((1+x)^(1/2)-x^(1/2))^Pi,x)`

3.315.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.82

$$\int \left(-\sqrt{x} + \sqrt{1+x}\right)^\pi dx = -\frac{2(\pi\sqrt{x+1}\sqrt{x} + 2x + 1)(\sqrt{x+1} - \sqrt{x})^\pi}{\pi^2 - 4}$$

input `integrate(((1+x)^(1/2)-x^(1/2))^pi,x, algorithm="fricas")`

output `-2*(pi*sqrt(x + 1)*sqrt(x) + 2*x + 1)*(sqrt(x + 1) - sqrt(x))^pi/(pi^2 - 4)`

3.315.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4362 vs. 2(39) = 78.

Time = 3.88 (sec) , antiderivative size = 4362, normalized size of antiderivative = 96.93

$$\int \left(-\sqrt{x} + \sqrt{1+x}\right)^\pi dx = \text{Too large to display}$$

input `integrate(((1+x)**(1/2)-x**(1/2))**pi,x)`

output `Piecewise((4*x**(13/2)*sqrt(1 + 1/x)*sinh(asinh(sqrt(x)) + pi*asinh(sqrt(x))) * gamma(1 + pi/2) / (-4*x**5*gamma(1 + pi/2) + pi**2*x**5*gamma(1 + pi/2) - 4*x**4*gamma(1 + pi/2) + pi**2*x**4*gamma(1 + pi/2)) + 2*pi*x**(13/2)*sqrt(1 + 1/x)*sinh(asinh(sqrt(x)) + pi*asinh(sqrt(x))) * gamma(1 + pi/2) / (-4*x**5*gamma(1 + pi/2) + pi**2*x**5*gamma(1 + pi/2) - 4*x**4*gamma(1 + pi/2) + pi**2*x**4*gamma(1 + pi/2)) - 2*pi*x**(13/2)*cosh(asinh(sqrt(x)) + pi*asinh(sqrt(x))) * gamma(1 + pi/2) / (-4*x**5*gamma(1 + pi/2) + pi**2*x**5*gamma(1 + pi/2) - 4*x**4*gamma(1 + pi/2) + pi**2*x**4*gamma(1 + pi/2)) - 4*x**(13/2)*cosh(asinh(sqrt(x)) + pi*asinh(sqrt(x))) * gamma(1 + pi/2) / (-4*x**5*gamma(1 + pi/2) + pi**2*x**5*gamma(1 + pi/2) - 4*x**4*gamma(1 + pi/2) + pi**2*x**4*gamma(1 + pi/2)) + 6*x**(11/2)*sqrt(1 + 1/x)*sinh(asinh(sqrt(x)) + pi*asinh(sqrt(x))) * gamma(1 + pi/2) / (-4*x**5*gamma(1 + pi/2) + pi**2*x**5*gamma(1 + pi/2) - 4*x**4*gamma(1 + pi/2) + pi**2*x**4*gamma(1 + pi/2)) + 2*pi*x**(11/2)*sqrt(1 + 1/x)*sinh(asinh(sqrt(x)) + pi*asinh(sqrt(x))) * gamma(1 + pi/2) / (-4*x**5*gamma(1 + pi/2) + pi**2*x**5*gamma(1 + pi/2) - 4*x**4*gamma(1 + pi/2) + pi**2*x**4*gamma(1 + pi/2)) - 4*pi*x**(11/2)*cosh(asinh(sqrt(x)) + pi*asinh(sqrt(x))) * gamma(1 + pi/2) / (-4*x**5*gamma(1 + pi/2) + pi**2*x**5*gamma(1 + pi/2) - 4*x**4*gamma(1 + pi/2) + pi**2*x**4*gamma(1 + pi/2)) - 6*x**(11/2)*cosh(asinh(sqrt(x)) + pi*asinh(sqrt(x))) * gamma(1 + pi/2) / (-4*x**5*gamma(1 + pi/2) + pi**2*x**5*gamma(1 + pi/2) - 4*x**4*gamma(...`

3.315.7 Maxima [F]

$$\int (-\sqrt{x} + \sqrt{1+x})^\pi dx = \int (\sqrt{x+1} - \sqrt{x})^\pi dx$$

input `integrate(((1+x)^(1/2)-x^(1/2))^pi,x, algorithm="maxima")`

output `integrate((sqrt(x + 1) - sqrt(x))^pi, x)`

3.315.8 Giac [F]

$$\int \left(-\sqrt{x} + \sqrt{1+x}\right)^\pi dx = \int \left(\sqrt{x+1} - \sqrt{x}\right)^\pi dx$$

input `integrate(((1+x)^(1/2)-x^(1/2))^pi,x, algorithm="giac")`

output `integrate((sqrt(x + 1) - sqrt(x))^pi, x)`

3.315.9 Mupad [F(-1)]

Timed out.

$$\int \left(-\sqrt{x} + \sqrt{1+x}\right)^\pi dx = \int \left(\sqrt{x+1} - \sqrt{x}\right)^\pi dx$$

input `int(((x + 1)^(1/2) - x^(1/2))^Pi,x)`

output `int(((x + 1)^(1/2) - x^(1/2))^Pi, x)`

3.316 $\int \left(-2 + \left(-2 + \left(-2 + (-2 + x^2)^2 \right)^2 \right)^2 \right) dx$

3.316.1 Optimal result	1574
3.316.2 Mathematica [A] (verified)	1574
3.316.3 Rubi [A] (verified)	1575
3.316.4 Maple [A] (verified)	1575
3.316.5 Fricas [A] (verification not implemented)	1576
3.316.6 Sympy [A] (verification not implemented)	1576
3.316.7 Maxima [A] (verification not implemented)	1577
3.316.8 Giac [A] (verification not implemented)	1577
3.316.9 Mupad [B] (verification not implemented)	1578

3.316.1 Optimal result

Integrand size = 17, antiderivative size = 54

$$\int \left(-2 + \left(-2 + \left(-2 + (-2 + x^2)^2 \right)^2 \right)^2 \right) dx$$

$$= 2x - \frac{64x^3}{3} + \frac{336x^5}{5} - 96x^7 + \frac{220x^9}{3} - 32x^{11} + 8x^{13} - \frac{16x^{15}}{15} + \frac{x^{17}}{17}$$

output `2*x-64/3*x^3+336/5*x^5-96*x^7+220/3*x^9-32*x^11+8*x^13-16/15*x^15+1/17*x^17`
7

3.316.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00

$$\int \left(-2 + \left(-2 + \left(-2 + (-2 + x^2)^2 \right)^2 \right)^2 \right) dx$$

$$= 2x - \frac{64x^3}{3} + \frac{336x^5}{5} - 96x^7 + \frac{220x^9}{3} - 32x^{11} + 8x^{13} - \frac{16x^{15}}{15} + \frac{x^{17}}{17}$$

input `Integrate[-2 + (-2 + (-2 + (-2 + x^2)^2)^2)^2,x]`

3.316. $\int \left(-2 + \left(-2 + \left(-2 + (-2 + x^2)^2 \right)^2 \right)^2 \right) dx$

output $2*x - (64*x^3)/3 + (336*x^5)/5 - 96*x^7 + (220*x^9)/3 - 32*x^{11} + 8*x^{13} - (16*x^{15})/15 + x^{17}/17$

3.316.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\left(\left((x^2 - 2)^2 - 2 \right)^2 - 2 \right)^2 - 2 \right) dx$$

↓ 2009

$$\frac{x^{17}}{17} - \frac{16x^{15}}{15} + 8x^{13} - 32x^{11} + \frac{220x^9}{3} - 96x^7 + \frac{336x^5}{5} - \frac{64x^3}{3} + 2x$$

input `Int[-2 + (-2 + (-2 + (-2 + x^2)^2)^2)^2, x]`

output $2*x - (64*x^3)/3 + (336*x^5)/5 - 96*x^7 + (220*x^9)/3 - 32*x^{11} + 8*x^{13} - (16*x^{15})/15 + x^{17}/17$

3.316.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

3.316.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.83

3.316. $\int \left(-2 + \left(-2 + \left(-2 + (-2 + x^2)^2 \right)^2 \right)^2 \right) dx$

method	result	size
default	$2x - \frac{64}{3}x^3 + \frac{336}{5}x^5 - 96x^7 + \frac{220}{3}x^9 - 32x^{11} + 8x^{13} - \frac{16}{15}x^{15} + \frac{1}{17}x^{17}$	45
norman	$2x - \frac{64}{3}x^3 + \frac{336}{5}x^5 - 96x^7 + \frac{220}{3}x^9 - 32x^{11} + 8x^{13} - \frac{16}{15}x^{15} + \frac{1}{17}x^{17}$	45
risch	$2x - \frac{64}{3}x^3 + \frac{336}{5}x^5 - 96x^7 + \frac{220}{3}x^9 - 32x^{11} + 8x^{13} - \frac{16}{15}x^{15} + \frac{1}{17}x^{17}$	45
parallelrisch	$2x - \frac{64}{3}x^3 + \frac{336}{5}x^5 - 96x^7 + \frac{220}{3}x^9 - 32x^{11} + 8x^{13} - \frac{16}{15}x^{15} + \frac{1}{17}x^{17}$	45
parts	$2x - \frac{64}{3}x^3 + \frac{336}{5}x^5 - 96x^7 + \frac{220}{3}x^9 - 32x^{11} + 8x^{13} - \frac{16}{15}x^{15} + \frac{1}{17}x^{17}$	45
gosper	$\frac{x(15x^{16} - 272x^{14} + 2040x^{12} - 8160x^{10} + 18700x^8 - 24480x^6 + 17136x^4 - 5440x^2 + 510)}{255}$	46

input `int(((x^2-2)^2-2)^2-2,x,method=_RETURNVERBOSE)`

output $2*x-64/3*x^3+336/5*x^5-96*x^7+220/3*x^9-32*x^{11}+8*x^{13}-16/15*x^{15}+1/17*x^{17}$
7

3.316.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.81

$$\int \left(-2 + \left(-2 + \left(-2 + (-2 + x^2)^2 \right)^2 \right)^2 \right) dx$$

$$= \frac{1}{17}x^{17} - \frac{16}{15}x^{15} + 8x^{13} - 32x^{11} + \frac{220}{3}x^9 - 96x^7 + \frac{336}{5}x^5 - \frac{64}{3}x^3 + 2x$$

input `integrate(((x^2-2)^2-2)^2-2,x, algorithm="fricas")`

output $1/17*x^{17} - 16/15*x^{15} + 8*x^{13} - 32*x^{11} + 220/3*x^9 - 96*x^7 + 336/5*x^5 - 64/3*x^3 + 2*x$

3.316.6 Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.91

$$\int \left(-2 + \left(-2 + \left(-2 + (-2 + x^2)^2 \right)^2 \right)^2 \right) dx$$

$$= \frac{x^{17}}{17} - \frac{16x^{15}}{15} + 8x^{13} - 32x^{11} + \frac{220x^9}{3} - 96x^7 + \frac{336x^5}{5} - \frac{64x^3}{3} + 2x$$

3.316. $\int \left(-2 + \left(-2 + \left(-2 + (-2 + x^2)^2 \right)^2 \right)^2 \right) dx$

input `integrate((((x**2-2)**2-2)**2-2)**2-2,x)`

output `x**17/17 - 16*x**15/15 + 8*x**13 - 32*x**11 + 220*x**9/3 - 96*x**7 + 336*x**5/5 - 64*x**3/3 + 2*x`

3.316.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.81

$$\int \left(-2 + \left(-2 + \left(-2 + (-2 + x^2)^2 \right)^2 \right)^2 \right) dx$$

$$= \frac{1}{17} x^{17} - \frac{16}{15} x^{15} + 8 x^{13} - 32 x^{11} + \frac{220}{3} x^9 - 96 x^7 + \frac{336}{5} x^5 - \frac{64}{3} x^3 + 2 x$$

input `integrate((((x^2-2)^2-2)^2-2,x, algorithm="maxima")`

output `1/17*x^17 - 16/15*x^15 + 8*x^13 - 32*x^11 + 220/3*x^9 - 96*x^7 + 336/5*x^5 - 64/3*x^3 + 2*x`

3.316.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.81

$$\int \left(-2 + \left(-2 + \left(-2 + (-2 + x^2)^2 \right)^2 \right)^2 \right) dx$$

$$= \frac{1}{17} x^{17} - \frac{16}{15} x^{15} + 8 x^{13} - 32 x^{11} + \frac{220}{3} x^9 - 96 x^7 + \frac{336}{5} x^5 - \frac{64}{3} x^3 + 2 x$$

input `integrate((((x^2-2)^2-2)^2-2,x, algorithm="giac")`

output `1/17*x^17 - 16/15*x^15 + 8*x^13 - 32*x^11 + 220/3*x^9 - 96*x^7 + 336/5*x^5 - 64/3*x^3 + 2*x`

3.316. $\int \left(-2 + \left(-2 + \left(-2 + (-2 + x^2)^2 \right)^2 \right)^2 \right) dx$

3.316.9 Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.81

$$\int \left(-2 + \left(-2 + \left(-2 + (-2 + x^2)^2 \right)^2 \right)^2 \right) dx$$

$$= \frac{x^{17}}{17} - \frac{16x^{15}}{15} + 8x^{13} - 32x^{11} + \frac{220x^9}{3} - 96x^7 + \frac{336x^5}{5} - \frac{64x^3}{3} + 2x$$

input `int((((x^2 - 2)^2 - 2)^2 - 2),x)`

output `2*x - (64*x^3)/3 + (336*x^5)/5 - 96*x^7 + (220*x^9)/3 - 32*x^11 + 8*x^13 - (16*x^15)/15 + x^17/17`

3.316. $\int \left(-2 + \left(-2 + \left(-2 + (-2 + x^2)^2 \right)^2 \right)^2 \right) dx$

3.317 $\int \sin(4 \arctan(x)) dx$

3.317.1 Optimal result	1579
3.317.2 Mathematica [A] (verified)	1579
3.317.3 Rubi [F]	1580
3.317.4 Maple [C] (verified)	1580
3.317.5 Fricas [A] (verification not implemented)	1581
3.317.6 Sympy [F]	1581
3.317.7 Maxima [A] (verification not implemented)	1581
3.317.8 Giac [A] (verification not implemented)	1582
3.317.9 Mupad [B] (verification not implemented)	1582

3.317.1 Optimal result

Integrand size = 5, antiderivative size = 18

$$\int \sin(4 \arctan(x)) dx = -\frac{4}{1+x^2} - 2 \log(1+x^2)$$

output `-4/(x^2+1)-2*ln(x^2+1)`

3.317.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \sin(4 \arctan(x)) dx = -\frac{4}{1+x^2} - 2 \log(1+x^2)$$

input `Integrate[Sin[4*ArcTan[x]],x]`

output `-4/(1+x^2)-2*Log[1+x^2]`

3.317.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sin(4 \arctan(x)) dx$$

↓ 7299

$$\int \sin(4 \arctan(x)) dx$$

input `Int[Sin[4*ArcTan[x]],x]`

output `$Aborted`

3.317.3.1 Defintions of rubi rules used

rule 7299 `Int[u_, x_] :> CannotIntegrate[u, x]`

3.317.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.34 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.89

method	result	size
default	$\frac{2i}{x-i} - 2 \ln(x-i) - \frac{2i}{i+x} - 2 \ln(i+x)$	34

input `int(sin(4*arctan(x)),x,method=_RETURNVERBOSE)`

output `2*I/(x-I)-2*ln(x-I)-2*I/(I+x)-2*ln(I+x)`

3.317.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.28

$$\int \sin(4 \arctan(x)) dx = -\frac{2((x^2 + 1) \log(x^2 + 1) + 2)}{x^2 + 1}$$

input `integrate(sin(4*arctan(x)),x, algorithm="fricas")`output `-2*((x^2 + 1)*log(x^2 + 1) + 2)/(x^2 + 1)`**3.317.6 Sympy [F]**

$$\int \sin(4 \arctan(x)) dx = \int \sin(4 \operatorname{atan}(x)) dx$$

input `integrate(sin(4*atan(x)),x)`output `Integral(sin(4*atan(x)), x)`**3.317.7 Maxima [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.28

$$\int \sin(4 \arctan(x)) dx = -\frac{2((x^2 + 1) \log(x^2 + 1) + 2)}{x^2 + 1}$$

input `integrate(sin(4*arctan(x)),x, algorithm="maxima")`output `-2*((x^2 + 1)*log(x^2 + 1) + 2)/(x^2 + 1)`

3.317.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.28

$$\int \sin(4 \arctan(x)) dx = \frac{2(x^2 - 1)}{x^2 + 1} - 2 \log(x^2 + 1)$$

input `integrate(sin(4*arctan(x)),x, algorithm="giac")`output `2*(x^2 - 1)/(x^2 + 1) - 2*log(x^2 + 1)`**3.317.9 Mupad [B] (verification not implemented)**

Time = 16.64 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \sin(4 \arctan(x)) dx = -2 \ln(x^2 + 1) - \frac{4}{x^2 + 1}$$

input `int(sin(4*atan(x)),x)`output `- 2*log(x^2 + 1) - 4/(x^2 + 1)`

3.318
$$\int \frac{\sqrt[3]{\tan(x)}}{(\cos(x)+\sin(x))^2} dx$$

3.318.1 Optimal result 1583
 3.318.2 Mathematica [A] (verified) 1583
 3.318.3 Rubi [A] (verified) 1584
 3.318.4 Maple [F] 1586
 3.318.5 Fricas [B] (verification not implemented) 1586
 3.318.6 Sympy [F] 1587
 3.318.7 Maxima [A] (verification not implemented) 1587
 3.318.8 Giac [A] (verification not implemented) 1588
 3.318.9 Mupad [F(-1)] 1588

3.318.1 Optimal result

Integrand size = 14, antiderivative size = 61

$$\int \frac{\sqrt[3]{\tan(x)}}{(\cos(x) + \sin(x))^2} dx = -\frac{\arctan\left(\frac{1-2\sqrt[3]{\tan(x)}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{2} \log\left(1 + \sqrt[3]{\tan(x)}\right) - \frac{1}{6} \log(1 + \tan(x)) - \frac{\sqrt[3]{\tan(x)}}{1 + \tan(x)}$$

output `-1/3*arctan(1/3*(1-2*tan(x)^(1/3))*3^(1/2))*3^(1/2)+1/2*ln(1+tan(x)^(1/3))-1/6*ln(1+tan(x))-tan(x)^(1/3)/(1+tan(x))`

3.318.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.26

$$\int \frac{\sqrt[3]{\tan(x)}}{(\cos(x) + \sin(x))^2} dx = \frac{\arctan\left(\frac{-1+2\sqrt[3]{\tan(x)}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{3} \log\left(1 + \sqrt[3]{\tan(x)}\right) - \frac{1}{6} \log\left(1 - \sqrt[3]{\tan(x)} + \tan^{\frac{2}{3}}(x)\right) + \left(-1 + \frac{\sin(x)}{\cos(x) + \sin(x)}\right) \sqrt[3]{\tan(x)}$$

input `Integrate[Tan[x]^(1/3)/(Cos[x] + Sin[x])^2,x]`

output `ArcTan[(-1 + 2*Tan[x]^(1/3))/Sqrt[3]]/Sqrt[3] + Log[1 + Tan[x]^(1/3)]/3 - Log[1 - Tan[x]^(1/3) + Tan[x]^(2/3)]/6 + (-1 + Sin[x]/(Cos[x] + Sin[x]))*Tan[x]^(1/3)`

3.318.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4889, 51, 70, 16, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt[3]{\tan(x)}}{(\sin(x) + \cos(x))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt[3]{\tan(x)}}{(\sin(x) + \cos(x))^2} dx \\
 & \quad \downarrow \text{4889} \\
 & \int \frac{\sqrt[3]{\tan(x)}}{(\tan(x) + 1)^2} d \tan(x) \\
 & \quad \downarrow \text{51} \\
 & \frac{1}{3} \int \frac{1}{\tan^{\frac{2}{3}}(x)(\tan(x) + 1)} d \tan(x) - \frac{\sqrt[3]{\tan(x)}}{\tan(x) + 1} \\
 & \quad \downarrow \text{70} \\
 & \frac{1}{3} \left(\frac{3}{2} \int \frac{1}{\tan^{\frac{2}{3}}(x) - \sqrt[3]{\tan(x)} + 1} d \sqrt[3]{\tan(x)} + \frac{3}{2} \int \frac{1}{\sqrt[3]{\tan(x)} + 1} d \sqrt[3]{\tan(x)} - \frac{1}{2} \log(\tan(x) + 1) \right) - \\
 & \quad \frac{\sqrt[3]{\tan(x)}}{\tan(x) + 1} \\
 & \quad \downarrow \text{16}
 \end{aligned}$$

3.318. $\int \frac{\sqrt[3]{\tan(x)}}{(\cos(x) + \sin(x))^2} dx$

$$\frac{1}{3} \left(\frac{3}{2} \int \frac{1}{\tan^{\frac{2}{3}}(x) - \sqrt[3]{\tan(x)} + 1} d\sqrt[3]{\tan(x)} + \frac{3}{2} \log(\sqrt[3]{\tan(x)} + 1) - \frac{1}{2} \log(\tan(x) + 1) \right) - \frac{\sqrt[3]{\tan(x)}}{\tan(x) + 1}$$

↓ 1083

$$\frac{1}{3} \left(-3 \int \frac{1}{-\tan^{\frac{2}{3}}(x) - 3} d(2\sqrt[3]{\tan(x)} - 1) + \frac{3}{2} \log(\sqrt[3]{\tan(x)} + 1) - \frac{1}{2} \log(\tan(x) + 1) \right) - \frac{\sqrt[3]{\tan(x)}}{\tan(x) + 1}$$

↓ 217

$$\frac{1}{3} \left(\sqrt{3} \arctan\left(\frac{2\sqrt[3]{\tan(x)} - 1}{\sqrt{3}}\right) + \frac{3}{2} \log(\sqrt[3]{\tan(x)} + 1) - \frac{1}{2} \log(\tan(x) + 1) \right) - \frac{\sqrt[3]{\tan(x)}}{\tan(x) + 1}$$

input `Int[Tan[x]^(1/3)/(Cos[x] + Sin[x])^2,x]`

output `(Sqrt[3]*ArcTan[(-1 + 2*Tan[x]^(1/3))/Sqrt[3]] + (3*Log[1 + Tan[x]^(1/3)])/2 - Log[1 + Tan[x]]/2)/3 - Tan[x]^(1/3)/(1 + Tan[x])`

3.318.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 51 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

rule 70 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(2/3)), x_Symbol] := With[{q = Rt[-(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q^2), x] + (Simp[3/(2*b*q) Subst[Int[1/(q^2 - q*x + x^2), x], x, (c + d*x)^(1/3)], x] + Simp[3/(2*b*q^2) Subst[Int[1/(q + x), x], x, (c + d*x)^(1/3)], x])] /; FreeQ[{a, b, c, d}, x] && NegQ[(b*c - a*d)/b]`

3.318. $\int \frac{\sqrt[3]{\tan(x)}}{(\cos(x)+\sin(x))^2} dx$

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & (LtQ[a, 0] || LtQ[b, 0])`

rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4889 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, With[{d = FreeFactors[Tan[v], x]}, Simp[d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x^2), Tan[v]/d, u, x], x], x, Tan[v]/d, x]] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Tan[v], x], u, x]] /; InverseFunctionFreeQ[u, x] && !MatchQ[u, (v_.)*((c_.)*tan[w_]^(n_.)*tan[z_]^(n_.))^ (p_.) /; FreeQ[{c, p}, x] && IntegerQ[n] && LinearQ[w, x] && EqQ[z, 2*w]]`

3.318.4 Maple [F]

$$\int \frac{\tan(x)^{\frac{1}{3}}}{(\cos(x) + \sin(x))^2} dx$$

input `int(tan(x)^(1/3)/(cos(x)+sin(x))^2,x)`

output `int(tan(x)^(1/3)/(cos(x)+sin(x))^2,x)`

3.318.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 108 vs. $2(48) = 96$.

Time = 0.25 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.77

$$\int \frac{\sqrt[3]{\tan(x)}}{(\cos(x) + \sin(x))^2} dx$$

$$= \frac{2(\sqrt{3}\cos(x) + \sqrt{3}\sin(x)) \arctan\left(\frac{2}{3}\sqrt{3}\left(\frac{\sin(x)}{\cos(x)}\right)^{\frac{1}{3}} - \frac{1}{3}\sqrt{3}\right) - (\cos(x) + \sin(x)) \log\left(\left(\frac{\sin(x)}{\cos(x)}\right)^{\frac{2}{3}} - \left(\frac{\sin(x)}{\cos(x)}\right)\right)}{6(\cos(x) + \sin(x))}$$

3.318. $\int \frac{\sqrt[3]{\tan(x)}}{(\cos(x) + \sin(x))^2} dx$

input `integrate(tan(x)^(1/3)/(cos(x)+sin(x))^2,x, algorithm="fricas")`

output `1/6*(2*(sqrt(3)*cos(x) + sqrt(3)*sin(x))*arctan(2/3*sqrt(3)*(sin(x)/cos(x))^(1/3) - 1/3*sqrt(3)) - (cos(x) + sin(x))*log((sin(x)/cos(x))^(2/3) - (sin(x)/cos(x))^(1/3) + 1) + 2*(cos(x) + sin(x))*log((sin(x)/cos(x))^(1/3) + 1) - 6*(sin(x)/cos(x))^(1/3)*cos(x))/(cos(x) + sin(x))`

3.318.6 Sympy [F]

$$\int \frac{\sqrt[3]{\tan(x)}}{(\cos(x) + \sin(x))^2} dx = \int \frac{\sqrt[3]{\tan(x)}}{(\sin(x) + \cos(x))^2} dx$$

input `integrate(tan(x)**(1/3)/(cos(x)+sin(x))**2,x)`

output `Integral(tan(x)**(1/3)/(sin(x) + cos(x))**2, x)`

3.318.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt[3]{\tan(x)}}{(\cos(x) + \sin(x))^2} dx = \frac{1}{3} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} (2 \tan(x)^{\frac{1}{3}} - 1) \right) - \frac{\tan(x)^{\frac{1}{3}}}{\tan(x) + 1} - \frac{1}{6} \log \left(\tan(x)^{\frac{2}{3}} - \tan(x)^{\frac{1}{3}} + 1 \right) + \frac{1}{3} \log \left(\tan(x)^{\frac{1}{3}} + 1 \right)$$

input `integrate(tan(x)^(1/3)/(cos(x)+sin(x))^2,x, algorithm="maxima")`

output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*tan(x)^(1/3) - 1)) - tan(x)^(1/3)/(tan(x) + 1) - 1/6*log(tan(x)^(2/3) - tan(x)^(1/3) + 1) + 1/3*log(tan(x)^(1/3) + 1)`

3.318.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt[3]{\tan(x)}}{(\cos(x) + \sin(x))^2} dx = \frac{1}{3} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} (2 \tan(x)^{\frac{1}{3}} - 1) \right) - \frac{\tan(x)^{\frac{1}{3}}}{\tan(x) + 1} - \frac{1}{6} \log \left(\tan(x)^{\frac{2}{3}} - \tan(x)^{\frac{1}{3}} + 1 \right) + \frac{1}{3} \log \left(\left| \tan(x)^{\frac{1}{3}} + 1 \right| \right)$$

input `integrate(tan(x)^(1/3)/(cos(x)+sin(x))^2,x, algorithm="giac")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*tan(x)^(1/3) - 1)) - tan(x)^(1/3)/(tan(x) + 1) - 1/6*log(tan(x)^(2/3) - tan(x)^(1/3) + 1) + 1/3*log(abs(tan(x)^(1/3) + 1))`**3.318.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt[3]{\tan(x)}}{(\cos(x) + \sin(x))^2} dx = \int \frac{\tan(x)^{1/3}}{(\cos(x) + \sin(x))^2} dx$$

input `int(tan(x)^(1/3)/(cos(x) + sin(x))^2,x)`output `int(tan(x)^(1/3)/(cos(x) + sin(x))^2, x)`

3.319 $\int \csc^2(x) \csc^2(6x) \csc^2(10x) \csc^2(15x) \sin^2(2x) \sin^2(3x)$

3.319.1 Optimal result	1589
3.319.2 Mathematica [A] (verified)	1589
3.319.3 Rubi [B] (verified)	1590
3.319.4 Maple [B] (verified)	1594
3.319.5 Fricas [A] (verification not implemented)	1595
3.319.6 Sympy [F(-1)]	1595
3.319.7 Maxima [A] (verification not implemented)	1596
3.319.8 Giac [A] (verification not implemented)	1596
3.319.9 Mupad [B] (verification not implemented)	1597

3.319.1 Optimal result

Integrand size = 47, antiderivative size = 64

$$\int \csc^2(x) \csc^2(6x) \csc^2(10x) \csc^2(15x) \sin^2(2x) \sin^2(3x) \sin^2(5x) \sin^2(30x) dx$$

$$= 7x + 4 \sin(2x) - \frac{1}{2} \sin(4x) - \frac{4}{3} \sin(6x) - \sin(8x)$$

$$- \frac{2}{5} \sin(10x) + \frac{1}{6} \sin(12x) + \frac{2}{7} \sin(14x) + \frac{1}{8} \sin(16x)$$

output `7*x+4*sin(2*x)-1/2*sin(4*x)-4/3*sin(6*x)-sin(8*x)-2/5*sin(10*x)+1/6*sin(12*x)+2/7*sin(14*x)+1/8*sin(16*x)`

3.319.2 Mathematica [A] (verified)

Time = 0.48 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00

$$\int \csc^2(x) \csc^2(6x) \csc^2(10x) \csc^2(15x) \sin^2(2x) \sin^2(3x) \sin^2(5x) \sin^2(30x) dx$$

$$= 7x + 4 \sin(2x) - \frac{1}{2} \sin(4x) - \frac{4}{3} \sin(6x) - \sin(8x)$$

$$- \frac{2}{5} \sin(10x) + \frac{1}{6} \sin(12x) + \frac{2}{7} \sin(14x) + \frac{1}{8} \sin(16x)$$

input `Integrate[Csc[x]^2*Csc[6*x]^2*Csc[10*x]^2*Csc[15*x]^2*Sin[2*x]^2*Sin[3*x]^2*Sin[5*x]^2*Sin[30*x]^2,x]`

output $7*x + 4*\text{Sin}[2*x] - \text{Sin}[4*x]/2 - (4*\text{Sin}[6*x])/3 - \text{Sin}[8*x] - (2*\text{Sin}[10*x])/5 + \text{Sin}[12*x]/6 + (2*\text{Sin}[14*x])/7 + \text{Sin}[16*x]/8$

3.319.3 Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 136 vs. $2(64) = 128$.

Time = 2.08 (sec) , antiderivative size = 136, normalized size of antiderivative = 2.12, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.383$, Rules used = {3042, 4889, 2345, 27, 2345, 27, 2345, 27, 2345, 27, 2345, 27, 2345, 27, 1471, 27, 298, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^2(2x) \sin^2(3x) \sin^2(5x) \sin^2(30x) \csc^2(x) \csc^2(6x) \csc^2(10x) \csc^2(15x) dx \\
 & \quad \downarrow 3042 \\
 & \int \sin(2x)^2 \sin(3x)^2 \sin(5x)^2 \sin(30x)^2 \csc(x)^2 \csc(6x)^2 \csc(10x)^2 \csc(15x)^2 dx \\
 & \quad \downarrow 4889 \\
 & \int \frac{(\tan^8(x) - 28 \tan^6(x) + 134 \tan^4(x) - 92 \tan^2(x) + 1)^2}{(\tan^2(x) + 1)^9} d \tan(x) \\
 & \quad \downarrow 2345 \\
 & \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8} - \\
 & \frac{1}{16} \int \frac{16(-\tan^{14}(x) + 57 \tan^{12}(x) - 1109 \tan^{10}(x) + 8797 \tan^8(x) - 31907 \tan^6(x) + 56619 \tan^4(x) - 65351 \tan^2(x) + 4)}{(\tan^2(x) + 1)^8} dx \\
 & \quad \downarrow 27 \\
 & \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8} - \\
 & \int \frac{-\tan^{14}(x) + 57 \tan^{12}(x) - 1109 \tan^{10}(x) + 8797 \tan^8(x) - 31907 \tan^6(x) + 56619 \tan^4(x) - 65351 \tan^2(x) + 4}{(\tan^2(x) + 1)^8} dx \\
 & \quad \downarrow 2345
 \end{aligned}$$

3.319. $\int \csc^2(x) \csc^2(6x) \csc^2(10x) \csc^2(15x) \sin^2(2x) \sin^2(3x) \sin^2(5x) \sin^2(30x) dx$

$$\frac{1}{14} \int \frac{2(7 \tan^{12}(x) - 406 \tan^{10}(x) + 8169 \tan^8(x) - 69748 \tan^6(x) + 293097 \tan^4(x) - 689430 \tan^2(x) + 55303)}{(\tan^2(x) + 1)^7} d \tan(x) + \frac{83968 \tan(x)}{7(\tan^2(x) + 1)^7} + \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8}$$

↓ 27

$$\frac{1}{7} \int \frac{7 \tan^{12}(x) - 406 \tan^{10}(x) + 8169 \tan^8(x) - 69748 \tan^6(x) + 293097 \tan^4(x) - 689430 \tan^2(x) + 55303}{(\tan^2(x) + 1)^7} d \tan(x) + \frac{83968 \tan(x)}{7(\tan^2(x) + 1)^7} + \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8}$$

↓ 2345

$$\frac{1}{7} \left(\frac{279040 \tan(x)}{3(\tan^2(x) + 1)^6} - \frac{1}{12} \int \frac{4(-21 \tan^{10}(x) + 1239 \tan^8(x) - 25746 \tan^6(x) + 234990 \tan^4(x) - 1114281 \tan^2(x) + 55303)}{(\tan^2(x) + 1)^6} d \tan(x) + \frac{83968 \tan(x)}{7(\tan^2(x) + 1)^7} + \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8} \right)$$

↓ 27

$$\frac{1}{7} \left(\frac{279040 \tan(x)}{3(\tan^2(x) + 1)^6} - \frac{1}{3} \int \frac{-21 \tan^{10}(x) + 1239 \tan^8(x) - 25746 \tan^6(x) + 234990 \tan^4(x) - 1114281 \tan^2(x) + 55303}{(\tan^2(x) + 1)^6} d \tan(x) + \frac{83968 \tan(x)}{7(\tan^2(x) + 1)^7} + \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8} \right)$$

↓ 2345

$$\frac{1}{7} \left(\frac{1}{3} \left(\frac{1}{10} \int \frac{6(35 \tan^8(x) - 2100 \tan^6(x) + 45010 \tan^4(x) - 436660 \tan^2(x) + 59683)}{(\tan^2(x) + 1)^5} d \tan(x) - \frac{744704 \tan(x)}{5(\tan^2(x) + 1)^5} \right) + \frac{83968 \tan(x)}{7(\tan^2(x) + 1)^7} + \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8} \right)$$

↓ 27

$$\frac{1}{7} \left(\frac{1}{3} \left(\frac{3}{5} \int \frac{35 \tan^8(x) - 2100 \tan^6(x) + 45010 \tan^4(x) - 436660 \tan^2(x) + 59683}{(\tan^2(x) + 1)^5} d \tan(x) - \frac{744704 \tan(x)}{5(\tan^2(x) + 1)^5} \right) + \frac{83968 \tan(x)}{7(\tan^2(x) + 1)^7} + \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8} \right)$$

↓ 2345

3.319. $\int \csc^2(x) \csc^2(6x) \csc^2(10x) \csc^2(15x) \sin^2(2x) \sin^2(3x) \sin^2(5x) \sin^2(30x) dx$

$$\frac{1}{7} \left(\frac{1}{3} \left(\frac{3}{5} \left(\frac{67936 \tan(x)}{(\tan^2(x) + 1)^4} - \frac{1}{8} \int \frac{56(-5 \tan^6(x) + 305 \tan^4(x) - 6735 \tan^2(x) + 1179)}{(\tan^2(x) + 1)^4} d \tan(x) \right) - \frac{744704 \tan(x)}{5(\tan^2(x) + 1)^5} \right) + \frac{83968 \tan(x)}{7(\tan^2(x) + 1)^7} + \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8} \right) \downarrow 27$$

$$\frac{1}{7} \left(\frac{1}{3} \left(\frac{3}{5} \left(\frac{67936 \tan(x)}{(\tan^2(x) + 1)^4} - 7 \int \frac{-5 \tan^6(x) + 305 \tan^4(x) - 6735 \tan^2(x) + 1179}{(\tan^2(x) + 1)^4} d \tan(x) \right) - \frac{744704 \tan(x)}{5(\tan^2(x) + 1)^5} \right) + \frac{83968 \tan(x)}{7(\tan^2(x) + 1)^7} + \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8} \right) \downarrow 2345$$

$$\frac{1}{7} \left(\frac{1}{3} \left(\frac{3}{5} \left(\frac{67936 \tan(x)}{(\tan^2(x) + 1)^4} - 7 \left(\frac{4112 \tan(x)}{3(\tan^2(x) + 1)^3} - \frac{1}{6} \int \frac{10(3 \tan^4(x) - 186 \tan^2(x) + 115)}{(\tan^2(x) + 1)^3} d \tan(x) \right) \right) - \frac{744704 \tan(x)}{5(\tan^2(x) + 1)^5} \right) + \frac{83968 \tan(x)}{7(\tan^2(x) + 1)^7} + \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8} \right) \downarrow 27$$

$$\frac{1}{7} \left(\frac{1}{3} \left(\frac{3}{5} \left(\frac{67936 \tan(x)}{(\tan^2(x) + 1)^4} - 7 \left(\frac{4112 \tan(x)}{3(\tan^2(x) + 1)^3} - \frac{5}{3} \int \frac{3 \tan^4(x) - 186 \tan^2(x) + 115}{(\tan^2(x) + 1)^3} d \tan(x) \right) \right) - \frac{744704 \tan(x)}{5(\tan^2(x) + 1)^5} \right) + \frac{83968 \tan(x)}{7(\tan^2(x) + 1)^7} + \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8} \right) \downarrow 1471$$

$$\frac{1}{7} \left(\frac{1}{3} \left(\frac{3}{5} \left(\frac{67936 \tan(x)}{(\tan^2(x) + 1)^4} - 7 \left(\frac{4112 \tan(x)}{3(\tan^2(x) + 1)^3} - \frac{5}{3} \left(\frac{76 \tan(x)}{(\tan^2(x) + 1)^2} - \frac{1}{4} \int -\frac{12(\tan^2(x) + 13)}{(\tan^2(x) + 1)^2} d \tan(x) \right) \right) \right) - \frac{744704 \tan(x)}{5(\tan^2(x) + 1)^5} \right) + \frac{83968 \tan(x)}{7(\tan^2(x) + 1)^7} + \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8} \right) \downarrow 27$$

$$\frac{1}{7} \left(\frac{1}{3} \left(\frac{3}{5} \left(\frac{67936 \tan(x)}{(\tan^2(x) + 1)^4} - 7 \left(\frac{4112 \tan(x)}{3(\tan^2(x) + 1)^3} - \frac{5}{3} \left(3 \int \frac{\tan^2(x) + 13}{(\tan^2(x) + 1)^2} d \tan(x) + \frac{76 \tan(x)}{(\tan^2(x) + 1)^2} \right) \right) \right) - \frac{744704 \tan(x)}{5(\tan^2(x) + 1)^5} \right) + \frac{83968 \tan(x)}{7(\tan^2(x) + 1)^7} + \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8} \right) \downarrow 298$$

3.319. $\int \csc^2(x) \csc^2(6x) \csc^2(10x) \csc^2(15x) \sin^2(2x) \sin^2(3x) \sin^2(5x) \sin^2(30x) dx$

$$\frac{1}{7} \left(\frac{1}{3} \left(\frac{3}{5} \left(\frac{67936 \tan(x)}{(\tan^2(x) + 1)^4} - 7 \left(\frac{4112 \tan(x)}{3(\tan^2(x) + 1)^3} - \frac{5}{3} \left(7 \int \frac{1}{\tan^2(x) + 1} d \tan(x) + \frac{6 \tan(x)}{\tan^2(x) + 1} \right) + \frac{76 \tan(x)}{(\tan^2(x) + 1)^2} \right) \right) \right) \right. \\ \left. + \frac{83968 \tan(x)}{7(\tan^2(x) + 1)^7} + \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8} \right)$$

↓ 216

$$\frac{1}{7} \left(\frac{1}{3} \left(\frac{3}{5} \left(\frac{67936 \tan(x)}{(\tan^2(x) + 1)^4} - 7 \left(\frac{4112 \tan(x)}{3(\tan^2(x) + 1)^3} - \frac{5}{3} \left(7 \arctan(\tan(x)) + \frac{6 \tan(x)}{\tan^2(x) + 1} \right) + \frac{76 \tan(x)}{(\tan^2(x) + 1)^2} \right) \right) \right) \right. \\ \left. + \frac{83968 \tan(x)}{7(\tan^2(x) + 1)^7} + \frac{4096 \tan(x)}{(\tan^2(x) + 1)^8} \right)$$

input `Int[Csc[x]^2*Csc[6*x]^2*Csc[10*x]^2*Csc[15*x]^2*Sin[2*x]^2*Sin[3*x]^2*Sin[5*x]^2*Sin[30*x]^2,x]`

output `(4096*Tan[x])/(1 + Tan[x]^2)^8 - (83968*Tan[x])/(7*(1 + Tan[x]^2)^7) + ((279040*Tan[x])/(3*(1 + Tan[x]^2)^6) + ((-744704*Tan[x])/(5*(1 + Tan[x]^2)^5) + (3*((67936*Tan[x])/(1 + Tan[x]^2)^4 - 7*((4112*Tan[x])/(3*(1 + Tan[x]^2)^3) - (5*((76*Tan[x])/(1 + Tan[x]^2)^2 + 3*(7*ArcTan[Tan[x]] + (6*Tan[x])/(1 + Tan[x]^2))))/3)))/5)/3)/7`

3.319.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 298 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[(-(b*c - a*d))*x*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] - Simp[(a*d - b*c*(2*p + 3))/(2*a*b*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/2 + p, 0])`

```
rule 1471 Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_),
x_Symbol] := With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, d + e*x^2
, x], R = Coeff[PolynomialRemainder[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], x
, 0]}, Simp[(-R)*x*((d + e*x^2)^(q + 1)/(2*d*(q + 1))), x] + Simp[1/(2*d*(q
+ 1)) Int[(d + e*x^2)^(q + 1)*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3),
x], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^
2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]
```

```
rule 2345 Int[(Pq_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuot
ient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x,
0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(a*g - b
*f*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Simp[1/(2*a*(p + 1)) In
t[(a + b*x^2)^(p + 1)*ExpandToSum[2*a*(p + 1)*Q + f*(2*p + 3), x], x], x]
/; FreeQ[{a, b}, x] && PolyQ[Pq, x] && LtQ[p, -1]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4889 Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, With[{d = FreeFactors
[Tan[v], x]}, Simp[d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x
^2), Tan[v]/d, u, x], x], x, Tan[v]/d, x]] /; !FalseQ[v] && FunctionOfQ[N
onfreeFactors[Tan[v], x], u, x]] /; InverseFunctionFreeQ[u, x] && !MatchQ[
u, (v_)*((c_)*tan[w_]^(n_)*tan[z_]^(n_))^(p_) /; FreeQ[{c, p}, x] && I
ntegerQ[n] && LinearQ[w, x] && EqQ[z, 2*w]]
```

3.319.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 227 vs. $2(52) = 104$.

Time = 0.74 (sec) , antiderivative size = 228, normalized size of antiderivative = 3.56

$$4096 \left(\cos(x)^{15} + \frac{15 \cos(x)^{13}}{14} + \frac{65 \cos(x)^{11}}{56} + \frac{143 \cos(x)^9}{112} + \frac{1287 \cos(x)^7}{896} + \frac{429 \cos(x)^5}{256} + \frac{2145 \cos(x)^3}{1024} \right)$$

```
input int(sin(2*x)^2*sin(3*x)^2*sin(5*x)^2*sin(30*x)^2/sin(x)^2/sin(6*x)^2/sin(1
0*x)^2/sin(15*x)^2,x)
```

output `4096*(cos(x)^15+15/14*cos(x)^13+65/56*cos(x)^11+143/112*cos(x)^9+1287/896*cos(x)^7+429/256*cos(x)^5+2145/1024*cos(x)^3+6435/2048*cos(x))*sin(x)+7*x-16384*(cos(x)^13+13/12*cos(x)^11+143/120*cos(x)^9+429/320*cos(x)^7+1001/640*cos(x)^5+1001/512*cos(x)^3+3003/1024*cos(x))*sin(x)+78848/3*(cos(x)^11+11/10*cos(x)^9+99/80*cos(x)^7+231/160*cos(x)^5+231/128*cos(x)^3+693/256*cos(x))*sin(x)-108544/5*(cos(x)^9+9/8*cos(x)^7+21/16*cos(x)^5+105/64*cos(x)^3+315/128*cos(x))*sin(x)+9920*(cos(x)^7+7/6*cos(x)^5+35/24*cos(x)^3+35/16*cos(x))*sin(x)-7616/3*(cos(x)^5+5/4*cos(x)^3+15/8*cos(x))*sin(x)+368*(cos(x)^3+3/2*cos(x))*sin(x)-32*cos(x)*sin(x)`

3.319.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.86

$$\int \csc^2(x) \csc^2(6x) \csc^2(10x) \csc^2(15x) \sin^2(2x) \sin^2(3x) \sin^2(5x) \sin^2(30x) dx$$

$$= \frac{2}{105} (215040 \cos(x)^{15} - 629760 \cos(x)^{13} + 697600 \cos(x)^{11} - 372352 \cos(x)^9 + 101904 \cos(x)^7 - 14392 \cos(x)^5 + 1330 \cos(x)^3 + 315 \cos(x)) \sin(x) + 7x$$

input `integrate(sin(2*x)^2*sin(3*x)^2*sin(5*x)^2*sin(30*x)^2/sin(x)^2/sin(6*x)^2/sin(10*x)^2/sin(15*x)^2,x, algorithm="fricas")`

output `2/105*(215040*cos(x)^15 - 629760*cos(x)^13 + 697600*cos(x)^11 - 372352*cos(x)^9 + 101904*cos(x)^7 - 14392*cos(x)^5 + 1330*cos(x)^3 + 315*cos(x))*sin(x) + 7*x`

3.319.6 Sympy [F(-1)]

Timed out.

$$\int \csc^2(x) \csc^2(6x) \csc^2(10x) \csc^2(15x) \sin^2(2x) \sin^2(3x) \sin^2(5x) \sin^2(30x) dx = \text{Timed out}$$

input `integrate(sin(2*x)**2*sin(3*x)**2*sin(5*x)**2*sin(30*x)**2/sin(x)**2/sin(6*x)**2/sin(10*x)**2/sin(15*x)**2,x)`

output `Timed out`

3.319. $\int \csc^2(x) \csc^2(6x) \csc^2(10x) \csc^2(15x) \sin^2(2x) \sin^2(3x) \sin^2(5x) \sin^2(30x) dx$

3.319.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.81

$$\int \csc^2(x) \csc^2(6x) \csc^2(10x) \csc^2(15x) \sin^2(2x) \sin^2(3x) \sin^2(5x) \sin^2(30x) dx$$

$$= 7x + \frac{1}{8} \sin(16x) + \frac{2}{7} \sin(14x) + \frac{1}{6} \sin(12x) - \frac{2}{5} \sin(10x)$$

$$- \sin(8x) - \frac{4}{3} \sin(6x) - \frac{1}{2} \sin(4x) + 4 \sin(2x)$$

```
input integrate(sin(2*x)^2*sin(3*x)^2*sin(5*x)^2*sin(30*x)^2/sin(x)^2/sin(6*x)^2
/sin(10*x)^2/sin(15*x)^2,x, algorithm="maxima")
```

```
output 7*x + 1/8*sin(16*x) + 2/7*sin(14*x) + 1/6*sin(12*x) - 2/5*sin(10*x) - sin(
8*x) - 4/3*sin(6*x) - 1/2*sin(4*x) + 4*sin(2*x)
```

3.319.8 Giac [A] (verification not implemented)

Time = 37.52 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.95

$$\int \csc^2(x) \csc^2(6x) \csc^2(10x) \csc^2(15x) \sin^2(2x) \sin^2(3x) \sin^2(5x) \sin^2(30x) dx = 7x$$

$$+ \frac{2(315 \tan(x)^{15} + 3535 \tan(x)^{13} + 203 \tan(x)^{11} + 60919 \tan(x)^9 - 71031 \tan(x)^7 + 74613 \tan(x)^5 - 5775 \tan(x)^3 - 315 \tan(x))}{105(\tan(x)^2 + 1)^8}$$

```
input integrate(sin(2*x)^2*sin(3*x)^2*sin(5*x)^2*sin(30*x)^2/sin(x)^2/sin(6*x)^2
/sin(10*x)^2/sin(15*x)^2,x, algorithm="giac")
```

```
output 7*x + 2/105*(315*tan(x)^15 + 3535*tan(x)^13 + 203*tan(x)^11 + 60919*tan(x)
^9 - 71031*tan(x)^7 + 74613*tan(x)^5 - 5775*tan(x)^3 - 315*tan(x))/(tan(x)
^2 + 1)^8
```

3.319.9 Mupad [B] (verification not implemented)

Time = 19.38 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.03

$$\int \csc^2(x) \csc^2(6x) \csc^2(10x) \csc^2(15x) \sin^2(2x) \sin^2(3x) \sin^2(5x) \sin^2(30x) dx$$

$$= 4096 \sin(x) \cos(x)^{15} - \frac{83968 \sin(x) \cos(x)^{13}}{7} + \frac{279040 \sin(x) \cos(x)^{11}}{21}$$

$$- \frac{744704 \sin(x) \cos(x)^9}{105} + \frac{67936 \sin(x) \cos(x)^7}{35}$$

$$- \frac{4112 \sin(x) \cos(x)^5}{15} + \frac{76 \sin(x) \cos(x)^3}{3} + 6 \sin(x) \cos(x) + 7x$$

input `int((sin(2*x)^2*sin(3*x)^2*sin(5*x)^2*sin(30*x)^2)/(sin(6*x)^2*sin(10*x)^2*sin(15*x)^2*sin(x)^2),x)`

output `7*x + 6*cos(x)*sin(x) + (76*cos(x)^3*sin(x))/3 - (4112*cos(x)^5*sin(x))/15 + (67936*cos(x)^7*sin(x))/35 - (744704*cos(x)^9*sin(x))/105 + (279040*cos(x)^11*sin(x))/21 - (83968*cos(x)^13*sin(x))/7 + 4096*cos(x)^15*sin(x)`

3.320 $\int \sqrt{1+x^2} + \sqrt{1+x^2+x^4} dx$

3.320.1 Optimal result	1598
3.320.2 Mathematica [A] (verified)	1598
3.320.3 Rubi [F]	1599
3.320.4 Maple [F]	1599
3.320.5 Fricas [A] (verification not implemented)	1600
3.320.6 Sympy [F]	1600
3.320.7 Maxima [F]	1600
3.320.8 Giac [F]	1601
3.320.9 Mupad [F(-1)]	1601

3.320.1 Optimal result

Integrand size = 21, antiderivative size = 105

$$\int \sqrt{1+x^2} + \sqrt{1+x^2+x^4} dx = \frac{1}{8} \left(\frac{2x(5+x^2+5\sqrt{1+x^2+x^4})}{(1-x^2+\sqrt{1+x^2+x^4})\sqrt{1+x^2} + \sqrt{1+x^2+x^4}} + 3\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{2}x}{\sqrt{1+x^2} + \sqrt{1+x^2+x^4}}\right) \right)$$

```
output 1/4*x*(5+x^2+5*(x^4+x^2+1)^(1/2))/(1-x^2+(x^4+x^2+1)^(1/2))/(1+x^2+(x^4+x^2+1)^(1/2))^(1/2)+3/8*2^(1/2)*arctanh(2^(1/2)*x/(1+x^2+(x^4+x^2+1)^(1/2))^(1/2))
```

3.320.2 Mathematica [A] (verified)

Time = 0.55 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.00

$$\int \sqrt{1+x^2} + \sqrt{1+x^2+x^4} dx = \frac{1}{8} \left(\frac{2x(5+x^2+5\sqrt{1+x^2+x^4})}{(1-x^2+\sqrt{1+x^2+x^4})\sqrt{1+x^2} + \sqrt{1+x^2+x^4}} + 3\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{2}x}{\sqrt{1+x^2} + \sqrt{1+x^2+x^4}}\right) \right)$$

```
input Integrate[Sqrt[1 + x^2 + Sqrt[1 + x^2 + x^4]], x]
```

output $((2*x*(5 + x^2 + 5*\text{Sqrt}[1 + x^2 + x^4]))/((1 - x^2 + \text{Sqrt}[1 + x^2 + x^4])* \text{Sqrt}[1 + x^2 + \text{Sqrt}[1 + x^2 + x^4]]) + 3*\text{Sqrt}[2]*\text{ArcTanh}[(\text{Sqrt}[2]*x)/\text{Sqrt}[1 + x^2 + \text{Sqrt}[1 + x^2 + x^4]]])/8$

3.320.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x^2 + \sqrt{x^4 + x^2 + 1} + 1} dx$$

↓ 7299

$$\int \sqrt{x^2 + \sqrt{x^4 + x^2 + 1} + 1} dx$$

input `Int[Sqrt[1 + x^2 + Sqrt[1 + x^2 + x^4]],x]`

output `$Aborted`

3.320.3.1 Defintions of rubi rules used

rule 7299 `Int[u_, x_] :> CannotIntegrate[u, x]`

3.320.4 Maple [F]

$$\int \sqrt{1 + x^2 + \sqrt{x^4 + x^2 + 1}} dx$$

input `int((1+x^2+(x^4+x^2+1)^(1/2))^(1/2),x)`

output `int((1+x^2+(x^4+x^2+1)^(1/2))^(1/2),x)`

3.320.5 Fracas [A] (verification not implemented)

Time = 0.65 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.41

$$\int \sqrt{1+x^2} + \sqrt{1+x^2+x^4} dx$$

$$= \frac{3\sqrt{2}x \log\left(-\frac{64x^5+16x^3+16\sqrt{x^4+x^2+1}(4x^3-x)+4\left(\sqrt{2}\sqrt{x^4+x^2+1}(8x^2-5)+\sqrt{2}(8x^4-x^2+5)\right)\sqrt{x^2+\sqrt{x^4+x^2+1}+1+25x}}{x}\right)+8(3\sqrt{x^2+\sqrt{x^4+x^2+1}+1})}{32x} + 8(3\sqrt{x^2+\sqrt{x^4+x^2+1}+1})$$

input `integrate((1+x^2+(x^4+x^2+1)^(1/2))^(1/2),x, algorithm="fricas")`output `1/32*(3*sqrt(2)*x*log(-(64*x^5 + 16*x^3 + 16*sqrt(x^4 + x^2 + 1)*(4*x^3 - x) + 4*(sqrt(2)*sqrt(x^4 + x^2 + 1)*(8*x^2 - 5) + sqrt(2)*(8*x^4 - x^2 + 5))*sqrt(x^2 + sqrt(x^4 + x^2 + 1) + 1) + 25*x)/x) + 8*(3*x^2 - sqrt(x^4 + x^2 + 1) + 1)*sqrt(x^2 + sqrt(x^4 + x^2 + 1) + 1))/x`**3.320.6 Sympy [F]**

$$\int \sqrt{1+x^2} + \sqrt{1+x^2+x^4} dx = \int \sqrt{x^2 + \sqrt{x^4 + x^2 + 1} + 1} dx$$

input `integrate((1+x**2+(x**4+x**2+1)**(1/2))**(1/2),x)`output `Integral(sqrt(x**2 + sqrt(x**4 + x**2 + 1) + 1), x)`**3.320.7 Maxima [F]**

$$\int \sqrt{1+x^2} + \sqrt{1+x^2+x^4} dx = \int \sqrt{x^2 + \sqrt{x^4 + x^2 + 1} + 1} dx$$

input `integrate((1+x^2+(x^4+x^2+1)^(1/2))^(1/2),x, algorithm="maxima")`output `integrate(sqrt(x^2 + sqrt(x^4 + x^2 + 1) + 1), x)`

3.320.8 Giac [F]

$$\int \sqrt{1+x^2} + \sqrt{1+x^2+x^4} dx = \int \sqrt{x^2 + \sqrt{x^4+x^2+1} + 1} dx$$

input `integrate((1+x^2+(x^4+x^2+1)^(1/2))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(x^2 + sqrt(x^4 + x^2 + 1) + 1), x)`

3.320.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{1+x^2} + \sqrt{1+x^2+x^4} dx = \int \sqrt{\sqrt{x^4+x^2+1} + x^2 + 1} dx$$

input `int(((x^2 + x^4 + 1)^(1/2) + x^2 + 1)^(1/2),x)`

output `int(((x^2 + x^4 + 1)^(1/2) + x^2 + 1)^(1/2), x)`

3.321 $\int \frac{x^9}{575-48x^{10}+x^{20}} dx$

3.321.1 Optimal result	1602
3.321.2 Mathematica [A] (verified)	1602
3.321.3 Rubi [A] (verified)	1603
3.321.4 Maple [A] (verified)	1604
3.321.5 Fricas [A] (verification not implemented)	1604
3.321.6 Sympy [A] (verification not implemented)	1604
3.321.7 Maxima [A] (verification not implemented)	1605
3.321.8 Giac [A] (verification not implemented)	1605
3.321.9 Mupad [B] (verification not implemented)	1605

3.321.1 Optimal result

Integrand size = 16, antiderivative size = 25

$$\int \frac{x^9}{575 - 48x^{10} + x^{20}} dx = -\frac{1}{20} \log(23 - x^{10}) + \frac{1}{20} \log(25 - x^{10})$$

output `-1/20*ln(-x^10+23)+1/20*ln(-x^10+25)`

3.321.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{x^9}{575 - 48x^{10} + x^{20}} dx = -\frac{1}{20} \log(23 - x^{10}) + \frac{1}{20} \log(25 - x^{10})$$

input `Integrate[x^9/(575 - 48*x^10 + x^20),x]`

output `-1/20*Log[23 - x^10] + Log[25 - x^10]/20`

3.321.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.16, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {1690, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^9}{x^{20} - 48x^{10} + 575} dx$$

$$\downarrow 1690$$

$$\frac{1}{10} \int \frac{1}{x^{20} - 48x^{10} + 575} dx^{10}$$

$$\downarrow 1081$$

$$\frac{1}{10} \int \left(\frac{1}{2(23 - x^{10})} - \frac{1}{2(25 - x^{10})} \right) dx^{10}$$

$$\downarrow 2009$$

$$\frac{1}{10} \left(\frac{1}{2} \log(25 - x^{10}) - \frac{1}{2} \log(23 - x^{10}) \right)$$

input `Int[x^9/(575 - 48*x^10 + x^20), x]`

output `(-1/2*Log[23 - x^10] + Log[25 - x^10]/2)/10`

3.321.3.1 Defintions of rubi rules used

rule 1081 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 1690 `Int[(x_)^(m_.)*((a_) + (c_.)*(x_)^(n2_.) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] := Simp[1/n Subst[Int[(a + b*x + c*x^2)^p, x], x, x^n], x] /; FreeQ[{a, b, c, m, n, p}, x] && EqQ[n2, 2*n] && EqQ[Simplify[m - n + 1], 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.321.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.72

method	result	size
default	$\frac{\ln(x^{10}-25)}{20} - \frac{\ln(x^{10}-23)}{20}$	18
risch	$\frac{\ln(x^{10}-25)}{20} - \frac{\ln(x^{10}-23)}{20}$	18
norman	$\frac{\ln(x^5-5)}{20} + \frac{\ln(x^5+5)}{20} - \frac{\ln(x^{10}-23)}{20}$	26
parallelrisc	$\frac{\ln(x^5-5)}{20} + \frac{\ln(x^5+5)}{20} - \frac{\ln(x^{10}-23)}{20}$	26

input `int(x^9/(x^20-48*x^10+575),x,method=_RETURNVERBOSE)`output `1/20*ln(x^10-25)-1/20*ln(x^10-23)`**3.321.5 Fricas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.68

$$\int \frac{x^9}{575 - 48x^{10} + x^{20}} dx = -\frac{1}{20} \log(x^{10} - 23) + \frac{1}{20} \log(x^{10} - 25)$$

input `integrate(x^9/(x^20-48*x^10+575),x, algorithm="fricas")`output `-1/20*log(x^10 - 23) + 1/20*log(x^10 - 25)`**3.321.6 Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.60

$$\int \frac{x^9}{575 - 48x^{10} + x^{20}} dx = \frac{\log(x^{10} - 25)}{20} - \frac{\log(x^{10} - 23)}{20}$$

input `integrate(x**9/(x**20-48*x**10+575),x)`output `log(x**10 - 25)/20 - log(x**10 - 23)/20`

3.321. $\int \frac{x^9}{575-48x^{10}+x^{20}} dx$

3.321.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.68

$$\int \frac{x^9}{575 - 48x^{10} + x^{20}} dx = -\frac{1}{20} \log(x^{10} - 23) + \frac{1}{20} \log(x^{10} - 25)$$

input `integrate(x^9/(x^20-48*x^10+575),x, algorithm="maxima")`output `-1/20*log(x^10 - 23) + 1/20*log(x^10 - 25)`**3.321.8 Giac [A] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{x^9}{575 - 48x^{10} + x^{20}} dx = -\frac{1}{20} \log(|x^{10} - 23|) + \frac{1}{20} \log(|x^{10} - 25|)$$

input `integrate(x^9/(x^20-48*x^10+575),x, algorithm="giac")`output `-1/20*log(abs(x^10 - 23)) + 1/20*log(abs(x^10 - 25))`**3.321.9 Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{x^9}{575 - 48x^{10} + x^{20}} dx = \frac{\operatorname{atan}\left(\frac{x^{10} 1006772302081i - 21354610286400i}{2807924963544 x^{10} - 66383426822975}\right)}{10} 1i$$

input `int(x^9/(x^20 - 48*x^10 + 575),x)`output `(atan((x^10*1006772302081i - 21354610286400i)/(2807924963544*x^10 - 66383426822975))*1i)/10`

APPENDIX

4.1 Listing of Grading functions	1606
--------------------------------------------	------

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.1.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
```

```

(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A"," "}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A"," "}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal.$
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
      ,
      finalresult={"F","Contains unresolved integral."}
    ]
  ];

  finalresult
]

```



```

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType,expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]],2]],
            Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3,ExpnType[expn[[1]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
            If[Head[expn]===RootSum,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, CsCh,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsCh
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

4.1.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

```

```

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
  if is_contains_complex(result) then
    if is_contains_complex(optimal) then
      if debug then
        print("both result and optimal complex");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        return "A"," ";
      else
        return "B",cat("Both result and optimal contain complex but leaf count of
                        convert(leaf_count_result,string)," vs. $2 (" ,
                        convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_
        end if
      else #result contains complex but optimal is not
        if debug then
          print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
      fi;
    else # result do not contain complex
      # this assumes optimal do not as well. No check is needed here.
      if debug then
        print("result do not contain complex, this assumes optimal do not as well");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        if debug then
          print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
      else
        if debug then
          print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string)," $ vs. $2(" ,
                        convert(leaf_count_optimal,string)," )=" ,convert(2*leaf_cou
        fi;
      fi;
    fi;
  fi;

```

```

else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  end if
end if

```

```

elif type(expn, ``~`) then
  if type(op(2,expn), 'integer') then
    ExpnType(op(1,expn))
  elif type(op(2,expn), 'rational') then
    if type(op(1,expn), 'rational') then
      1
    else
      max(2, ExpnType(op(1,expn)))
    end if
  else
    max(3, ExpnType(op(1,expn)), ExpnType(op(2,expn)))
  end if
elif type(expn, ``+`) or type(expn, ``*`) then
  max(ExpnType(op(1,expn)), max(ExpnType(rest(expn))))
elif ElementaryFunctionQ(op(0,expn)) then
  max(3, ExpnType(op(1,expn)))
elif SpecialFunctionQ(op(0,expn)) then
  max(4, apply(max, map(ExpnType, [op(expn)])))
elif HypergeometricFunctionQ(op(0,expn)) then
  max(5, apply(max, map(ExpnType, [op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6, apply(max, map(ExpnType, [op(expn)])))
elif op(0,expn)='int' then
  max(8, apply(max, map(ExpnType, [op(expn)]))) else
  9
end if
end proc:

```

```

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

```

```

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,

```

```

        GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
        EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
    member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
    member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
    if nops(u)=2 then
        op(2,u)
    else
        apply(op(0,u),op(2..nops(u),u))
    end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma] [LeafCount] (u);
end proc:

```

4.1.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):

```

```
if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
        return True
    else:
        return False
else:
    return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
```



```

return 1
elif isinstance(expn,list):
    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
        return 1
    else:
        return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
elif isinstance(expn,Pow): #type(expn,``^`)
    if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
        return expnType(expn.args[0]) #ExpnType(op(1,expn))
    elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or type(expn,``*`)
    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

#main function

```

def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

    #print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```

4.1.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
```

```

        return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func ," is special_function")
        else:
            print ("func ", func ," is NOT special_function")

    return m

```

```

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric', 'hypergeometric_M', 'hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=", expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception, AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list:  #isinstance(expn,list):

```

```

    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
elif expn.operator() == operator.pow: #isinstance(expn,Pow)
    if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
        return expnType(expn.operands()[0]) #expnType(expn.args[0])
    elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
        if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or inst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

```

```

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)",type(result))
    print("type(optimal)",type(optimal))

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal." + str(leaf
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```