

# Computer Algebra Independent Integration Tests

Summer 2023 edition with Rubi V 4.17.3

1-Algebraic-functions/1.2-Trinomial-products/1.2.2-Quartic/44-  
1.2.2.7-P-x-d+e-x<sup>2</sup>-<sup>q</sup>-a+b-x<sup>2</sup>+c-x<sup>4</sup>-<sup>p</sup>

Nasser M. Abbasi

December 9, 2023

Compiled on December 9, 2023 at 4:56am

# Contents

|          |   |            |
|----------|---|------------|
| <b>1</b> | <b>Introduction</b>                       | <b>2</b>   |
| <b>2</b> | <b>detailed summary tables of results</b> | <b>20</b>  |
| <b>3</b> | <b>Listing of integrals</b>               | <b>37</b>  |
| <b>4</b> | <b>Appendix</b>                           | <b>349</b> |

# CHAPTER 1

## INTRODUCTION

|      |   |    |
|------|---|----|
| 1.1  | Listing of CAS systems tested . . . . .                                   | 3  |
| 1.2  | Results . . . . .   | 4  |
| 1.3  | Time and leaf size Performance . . . . .                                  | 7  |
| 1.4  | Performance based on number of rules Rubi used . . . . .                  | 9  |
| 1.5  | Performance based on number of steps Rubi used . . . . .                  | 10 |
| 1.6  | Solved integrals histogram based on leaf size of result . . . . .         | 11 |
| 1.7  | Solved integrals histogram based on CPU time used . . . . .               | 12 |
| 1.8  | Leaf size vs. CPU time used . . . . .                                     | 13 |
| 1.9  | list of integrals with no known antiderivative . . . . .                  | 14 |
| 1.10 | List of integrals solved by CAS but has no known antiderivative . . . . . | 14 |
| 1.11 | list of integrals solved by CAS but failed verification . . . . .         | 14 |
| 1.12 | Timing . . . . .  | 15 |
| 1.13 | Verification . . . . .  | 15 |
| 1.14 | Important notes about some of the results . . . . .                       | 15 |
| 1.15 | Design of the test system . . . . .                                       | 19 |

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [ 42 ]. This is test number [ 44 ].

## 1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 13.3.1 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

## 1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

| System      | % solved      | % Failed      |
|-------------|---------------|---------------|
| Rubi        | 100.00 ( 42 ) | 0.00 ( 0 )    |
| Mathematica | 95.24 ( 40 )  | 4.76 ( 2 )    |
| Maple       | 95.24 ( 40 )  | 4.76 ( 2 )    |
| Fricas      | 61.90 ( 26 )  | 38.10 ( 16 )  |
| Giac        | 14.29 ( 6 )   | 85.71 ( 36 )  |
| Sympy       | 14.29 ( 6 )   | 85.71 ( 36 )  |
| Mupad       | 2.38 ( 1 )    | 97.62 ( 41 )  |
| Maxima      | 0.00 ( 0 )    | 100.00 ( 42 ) |

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

| grade | description   |
|-------|---|
| A     | Integral was solved and antiderivative is optimal in quality and leaf size.   |
| B     | Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.  |
| C     | Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> <li>1. antiderivative contains a hypergeometric function and the optimal antiderivative does not.</li> <li>2. antiderivative contains a special function and the optimal antiderivative does not.</li> <li>3. antiderivative contains the imaginary unit and the optimal antiderivative does not.</li> </ol> |
| F     | Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.  |


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


| System      | % A grade | % B grade | % C grade | % F grade |
|-------------|-----------|-----------|-----------|-----------|
| Rubi        | 100.000   | 0.000     | 0.000     | 0.000     |
| Maple       | 47.619    | 9.524     | 38.095    | 4.762     |
| Fricas      | 45.238    | 14.286    | 2.381     | 38.095    |
| Mathematica | 16.667    | 0.000     | 78.571    | 4.762     |
| Giac        | 11.905    | 2.381     | 0.000     | 85.714    |
| Mupad       | 0.000     | 2.381     | 0.000     | 97.619    |
| Maxima      | 0.000     | 0.000     | 0.000     | 100.000   |
| Sympy       | 0.000     | 0.000     | 14.286    | 85.714    |

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima

and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

| System      | Number failed | Percentage normal failure | Percentage time-out failure | Percentage exception failure |
|-------------|---------------|---------------------------|-----------------------------|------------------------------|
| Rubi        | 0             | 0.00                      | 0.00                        | 0.00                         |
| Mathematica | 2             | 100.00                    | 0.00                        | 0.00                         |
| Maple       | 2             | 100.00                    | 0.00                        | 0.00                         |
| Fricas      | 16            | 31.25                     | 68.75                       | 0.00                         |
| Giac        | 36            | 97.22                     | 0.00                        | 2.78                         |
| Sympy       | 36            | 80.56                     | 19.44                       | 0.00                         |
| Mupad       | 41            | 0.00                      | 100.00                      | 0.00                         |
| Maxima      | 42            | 83.33                     | 0.00                        | 16.67                        |

Table 1.4: Failure statistics for each CAS

## 1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.



| System      | Mean time (sec) |
|-------------|-----------------|
| Giac        | 0.33            |
| Fricas      | 0.34            |
| Rubi        | 0.96            |
| Maple       | 2.58            |
| Sympy       | 2.61            |
| Mupad       | 8.63            |
| Mathematica | 8.73            |
| Maxima      | -nan(ind)       |

Table 1.5: Time performance for each CAS

| System      | Mean size | Normalized mean | Median size | Normalized median |
|-------------|-----------|-----------------|-------------|-------------------|
| Sympy       | 186.00    | 0.53            | 167.00      | 0.51              |
| Giac        | 330.17    | 1.20            | 310.00      | 1.04              |
| Mupad       | 397.00    | 3.75            | 397.00      | 3.75              |
| Mathematica | 423.88    | 0.83            | 259.50      | 0.68              |
| Rubi        | 551.14    | 1.03            | 388.50      | 1.00              |
| Fricas      | 603.31    | 1.63            | 458.50      | 1.33              |
| Maple       | 863.42    | 1.26            | 360.50      | 0.88              |
| Maxima      | -nan(ind) | -nan(ind)       | nan         | nan               |

Table 1.6: Leaf size performance for each CAS

## 1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the  $y$  axis is the percentage solved which Rubi itself needed the number of rules given the  $x$  axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

# 1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

## 1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

## 1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

## 1.8 Leaf size vs. CPU time used

The following gives the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time.




Figure 1.5: Leaf size vs. CPU time. Full range

## 1.9 list of integrals with no known antiderivative

{}

## 1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

## 1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {34}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

**Giac** Verification phase not currently implemented.

**Mupad** Verification phase not currently implemented.

## 1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

## 1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

## 1.14 Important notes about some of the results

### 1.14.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.



The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

### 1.14.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

### 1.14.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at [https://ask.sagemath.org/question/57123/could-we-have-a-leaf\\_count-function-in-base-sagemath/](https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/)

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

### 1.14.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives  $\sin(x)^2/2$

# 1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



**High level overview of the CAS independent integration test build system**

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer, 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer, Leaf size of result.
4. integer, Leaf size of the optimal antiderivative.
5. number, CPU time used to solve this integral. 0 if failed.
6. string, The integral in Latex format
7. string, The input used in CAS own syntax.
8. string, The result (antiderivative) produced by CAS in Latex format
9. string, The optimal antiderivative in Latex format.
10. integer, 0 or 1. Indicates if problem has known antiderivative or not
11. String, The result (antiderivative) in CAS own syntax.
12. String, The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String, Small string description of why the grade was given.
14. integer, 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

*The following fields are present only in Rubi Table file*

15. integer, Number of steps used.
16. integer, Number of rules used.
17. integer, Integrand leaf size.
18. real number, Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String, The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi  
June 27, 2023  
Design v0.6

# CHAPTER 2

## DETAILED SUMMARY TABLES OF RESULTS

|     |   |    |
|-----|---|----|
| 2.1 | List of integrals sorted by grade for each CAS . . . . .                  | 21 |
| 2.2 | Detailed conclusion table per each integral for all CAS systems . . . . . | 24 |
| 2.3 | Detailed conclusion table specific for Rubi results . . . . .             | 35 |

## 2.1 List of integrals sorted by grade for each CAS

|       |                  |    |
|-------|------------------|----|
| 2.1.1 | Rubi . . . . .   | 21 |
| 2.1.2 | Mma . . . . .    | 21 |
| 2.1.3 | Maple . . . . .  | 22 |
| 2.1.4 | Fricas . . . . . | 22 |
| 2.1.5 | Maxima . . . . . | 22 |
| 2.1.6 | Giac . . . . .   | 23 |
| 2.1.7 | Mupad . . . . .  | 23 |
| 2.1.8 | Sympy . . . . .  | 23 |

### 2.1.1 Rubi

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42 }

**B grade** { }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

### 2.1.2 Mma

**A grade** { 35, 37, 38, 39, 40, 41, 42 }

**B grade** { }

**C grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 34, 36 }

**F normal fail** { 15, 33 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

### 2.1.3 Maple

**A grade** { 17, 18, 19, 20, 21, 24, 25, 26, 27, 30, 31, 34, 35, 36, 37, 38, 39, 40, 41, 42 }

**B grade** { 22, 23, 28, 29 }

**C grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 32 }

**F normal fail** { 15, 33 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

### 2.1.4 Fricas

**A grade** { 1, 2, 3, 4, 8, 9, 10, 11, 17, 18, 19, 20, 27, 37, 38, 39, 40, 41, 42 }

**B grade** { 24, 25, 26, 34, 35, 36 }

**C grade** { 16 }

**F normal fail** { 5, 15, 22, 32, 33 }

**F(-1) timeout fail** { 6, 7, 12, 13, 14, 21, 23, 28, 29, 30, 31 }

**F(-2) exception fail** { }

### 2.1.5 Maxima

**A grade** { }

**B grade** { }

**C grade** { }

**F normal fail** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 31, 32, 33, 34, 35, 36 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { 30, 37, 38, 39, 40, 41, 42 }

### 2.1.6 Giac

A grade { 34, 37, 38, 40, 41 }

B grade { 42 }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 35, 36 }

F(-1) timedout fail { }

F(-2) exception fail { 39 }

### 2.1.7 Mupad

A grade { }

B grade { 34 }

C grade { }

F normal fail { }

F(-1) timedout fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 35, 36, 37, 38, 39, 40, 41, 42 }

F(-2) exception fail { }

### 2.1.8 Sympy

A grade { }

B grade { }

C grade { 1, 2, 3, 4, 10, 11 }

F normal fail { 5, 6, 7, 8, 9, 12, 13, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 30, 31, 32, 34, 35, 36, 37, 38, 39, 40 }

F(-1) timedout fail { 14, 15, 28, 29, 33, 41, 42 }

F(-2) exception fail { }



## 2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column N.S. means **normalized size** and is defined as  $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$ . To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

| Problem 1  | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas | Sympy | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------|-------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | A      | C     | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD    | TBD   | TBD      | TBD          |
| size       | 453     | 878   | 217    | 327   | 0        | 283    | 364   | 0        | 0            |
| N.S.       | 1       | 1.94  | 0.48   | 0.72  | 0.00     | 0.62   | 0.80  | 0.00     | 0.00         |
| time (sec) | N/A     | 0.902 | 10.207 | 5.690 | 0.000    | 0.088  | 3.491 | 0.000    | 0.000        |

| Problem 2  | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas | Sympy | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------|-------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | A      | C     | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD    | TBD   | TBD      | TBD          |
| size       | 367     | 706   | 159    | 258   | 0        | 206    | 262   | 0        | 0            |
| N.S.       | 1       | 1.92  | 0.43   | 0.70  | 0.00     | 0.56   | 0.71  | 0.00     | 0.00         |
| time (sec) | N/A     | 0.713 | 10.120 | 3.023 | 0.000    | 0.092  | 2.590 | 0.000    | 0.000        |

| Problem 3  | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas | Sympy | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------|-------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | A      | C     | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD    | TBD   | TBD      | TBD          |
| size       | 277     | 432   | 121    | 203   | 0        | 127    | 167   | 0        | 0            |
| N.S.       | 1       | 1.56  | 0.44   | 0.73  | 0.00     | 0.46   | 0.60  | 0.00     | 0.00         |
| time (sec) | N/A     | 0.493 | 10.074 | 1.205 | 0.000    | 0.090  | 1.767 | 0.000    | 0.000        |

| Problem 4  | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas | Sympy | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------|-------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | A      | C     | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD    | TBD   | TBD      | TBD          |
| size       | 226     | 228   | 77     | 169   | 0        | 90     | 78    | 0        | 0            |
| N.S.       | 1       | 1.01  | 0.34   | 0.75  | 0.00     | 0.40   | 0.35  | 0.00     | 0.00         |
| time (sec) | N/A     | 0.294 | 10.029 | 0.530 | 0.000    | 0.091  | 0.856 | 0.000    | 0.000        |

| Problem 5  | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas   | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|----------|----------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | <b>F</b> | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD      | TBD      | TBD      | TBD          |
| size       | 369     | 382   | 138    | 192   | 0        | 0        | 0        | 0        | 0            |
| N.S.       | 1       | 1.04  | 0.37   | 0.52  | 0.00     | 0.00     | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 0.594 | 10.274 | 0.815 | 0.000    | 0.000    | 0.000    | 0.000    | 0.000        |

| Problem 6  | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas       | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------------|----------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | <b>F(-1)</b> | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD          | TBD      | TBD      | TBD          |
| size       | 641     | 600   | 297    | 679   | 0        | 0            | 0        | 0        | 0            |
| N.S.       | 1       | 0.94  | 0.46   | 1.06  | 0.00     | 0.00         | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 1.293 | 10.785 | 1.418 | 0.000    | 0.000        | 0.000    | 0.000    | 0.000        |

| Problem 7  | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas       | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------------|----------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | <b>F(-1)</b> | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD          | TBD      | TBD      | TBD          |
| size       | 875     | 788   | 453    | 1591  | 0        | 0            | 0        | 0        | 0            |
| N.S.       | 1       | 0.90  | 0.52   | 1.82  | 0.00     | 0.00         | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 2.191 | 11.327 | 3.007 | 0.000    | 0.000        | 0.000    | 0.000    | 0.000        |

| Problem 8  | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------|----------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | A      | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD    | TBD      | TBD      | TBD          |
| size       | 912     | 912   | 222    | 426   | 0        | 448    | 0        | 0        | 0            |
| N.S.       | 1       | 1.00  | 0.24   | 0.47  | 0.00     | 0.49   | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 1.016 | 10.188 | 8.289 | 0.000    | 0.117  | 0.000    | 0.000    | 0.000        |

| Problem 9  | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------|----------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | A      | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD    | TBD      | TBD      | TBD          |
| size       | 694     | 694   | 166    | 323   | 0        | 332    | 0        | 0        | 0            |
| N.S.       | 1       | 1.00  | 0.24   | 0.47  | 0.00     | 0.48   | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 0.785 | 10.138 | 1.549 | 0.000    | 0.101  | 0.000    | 0.000    | 0.000        |

| Problem 10 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas | Sympy | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------|-------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | A      | C     | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD    | TBD   | TBD      | TBD          |
| size       | 395     | 395   | 126    | 253   | 0        | 188    | 167   | 0        | 0            |
| N.S.       | 1       | 1.00  | 0.32   | 0.64  | 0.00     | 0.48   | 0.42  | 0.00     | 0.00         |
| time (sec) | N/A     | 0.523 | 10.083 | 0.690 | 0.000    | 0.102  | 4.471 | 0.000    | 0.000        |

| Problem 11 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas | Sympy | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------|-------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | A      | C     | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD    | TBD   | TBD      | TBD          |
| size       | 262     | 262   | 99     | 212   | 0        | 115    | 78    | 0        | 0            |
| N.S.       | 1       | 1.00  | 0.38   | 0.81  | 0.00     | 0.44   | 0.30  | 0.00     | 0.00         |
| time (sec) | N/A     | 0.325 | 10.041 | 0.559 | 0.000    | 0.085  | 2.465 | 0.000    | 0.000        |

| Problem 12 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas       | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------------|----------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | <b>F(-1)</b> | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD          | TBD      | TBD      | TBD          |
| size       | 732     | 732   | 432    | 564   | 0        | 0            | 0        | 0        | 0            |
| N.S.       | 1       | 1.00  | 0.59   | 0.77  | 0.00     | 0.00         | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 1.031 | 10.660 | 0.865 | 0.000    | 0.000        | 0.000    | 0.000    | 0.000        |

| Problem 13 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas       | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------------|----------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | <b>F(-1)</b> | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD          | TBD      | TBD      | TBD          |
| size       | 1494    | 1494  | 427    | 1384  | 0        | 0            | 0        | 0        | 0            |
| N.S.       | 1       | 1.00  | 0.29   | 0.93  | 0.00     | 0.00         | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 2.224 | 11.135 | 1.503 | 0.000    | 0.000        | 0.000    | 0.000    | 0.000        |

| Problem 14 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas       | Sympy        | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------------|--------------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | <b>F(-1)</b> | <b>F(-1)</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD          | TBD          | TBD      | TBD          |
| size       | 2452    | 2452  | 630    | 2326  | 0        | 0            | 0            | 0        | 0            |
| N.S.       | 1       | 1.00  | 0.26   | 0.95  | 0.00     | 0.00         | 0.00         | 0.00     | 0.00         |
| time (sec) | N/A     | 3.990 | 11.740 | 3.429 | 0.000    | 0.000        | 0.000        | 0.000    | 0.000        |

| Problem 15 | Optimal | Rubi  | MMA      | Maple    | Maxima   | Fricas   | Sympy        | Giac     | Mupad        |
|------------|---------|-------|----------|----------|----------|----------|--------------|----------|--------------|
| grade      | N/A     | A     | <b>F</b> | <b>F</b> | <b>F</b> | <b>F</b> | <b>F(-1)</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | N/A      | N/A      | TBD      | TBD      | TBD          | TBD      | TBD          |
| size       | 169     | 169   | 0        | 0        | 0        | 0        | 0            | 0        | 0            |
| N.S.       | 1       | 1.00  | 0.00     | 0.00     | 0.00     | 0.00     | 0.00         | 0.00     | 0.00         |
| time (sec) | N/A     | 0.382 | 0.000    | 0.000    | 0.000    | 0.000    | 0.000        | 0.000    | 0.000        |

| Problem 16 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------|----------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | C      | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD    | TBD      | TBD      | TBD          |
| size       | 48      | 48    | 94     | 80    | 0        | 59     | 0        | 0        | 0            |
| N.S.       | 1       | 1.00  | 1.96   | 1.67  | 0.00     | 1.23   | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 0.174 | 10.169 | 1.564 | 0.000    | 0.136  | 0.000    | 0.000    | 0.000        |

| Problem 17 | Optimal | Rubi  | MMA    | Maple  | Maxima   | Fricas | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|--------|----------|--------|----------|----------|--------------|
| grade      | N/A     | A     | C      | A      | <b>F</b> | A      | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes    | TBD      | TBD    | TBD      | TBD      | TBD          |
| size       | 755     | 690   | 901    | 673    | 0        | 1041   | 0        | 0        | 0            |
| N.S.       | 1       | 0.91  | 1.19   | 0.89   | 0.00     | 1.38   | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 1.447 | 14.513 | 10.893 | 0.000    | 0.148  | 0.000    | 0.000    | 0.000        |

| Problem 18 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------|----------|----------|--------------|
| grade      | N/A     | A     | C      | A     | <b>F</b> | A      | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD    | TBD      | TBD      | TBD          |
| size       | 528     | 498   | 674    | 508   | 0        | 681    | 0        | 0        | 0            |
| N.S.       | 1       | 0.94  | 1.28   | 0.96  | 0.00     | 1.29   | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 0.887 | 12.807 | 4.278 | 0.000    | 0.114  | 0.000    | 0.000    | 0.000        |

| Problem 19 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------|----------|----------|--------------|
| grade      | N/A     | A     | C      | A     | <b>F</b> | A      | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD    | TBD      | TBD      | TBD          |
| size       | 368     | 358   | 521    | 409   | 0        | 435    | 0        | 0        | 0            |
| N.S.       | 1       | 0.97  | 1.42   | 1.11  | 0.00     | 1.18   | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 0.507 | 11.347 | 3.066 | 0.000    | 0.123  | 0.000    | 0.000    | 0.000        |

| Problem 20 | Optimal | Rubi  | MMA   | Maple | Maxima   | Fricas | Sympy    | Giac     | Mupad        |
|------------|---------|-------|-------|-------|----------|--------|----------|----------|--------------|
| grade      | N/A     | A     | C     | A     | <b>F</b> | A      | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes   | Yes   | TBD      | TBD    | TBD      | TBD      | TBD          |
| size       | 283     | 285   | 302   | 362   | 0        | 300    | 0        | 0        | 0            |
| N.S.       | 1       | 1.01  | 1.07  | 1.28  | 0.00     | 1.06   | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 0.333 | 0.117 | 0.931 | 0.000    | 0.113  | 0.000    | 0.000    | 0.000        |

| Problem 21 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas       | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------------|----------|----------|--------------|
| grade      | N/A     | A     | C      | A     | <b>F</b> | <b>F(-1)</b> | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD          | TBD      | TBD      | TBD          |
| size       | 436     | 449   | 298    | 359   | 0        | 0            | 0        | 0        | 0            |
| N.S.       | 1       | 1.03  | 0.68   | 0.82  | 0.00     | 0.00         | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 0.664 | 10.428 | 1.734 | 0.000    | 0.000        | 0.000    | 0.000    | 0.000        |

| Problem 22 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas   | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|----------|----------|----------|--------------|
| grade      | N/A     | A     | C      | B     | <b>F</b> | <b>F</b> | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD      | TBD      | TBD      | TBD          |
| size       | 782     | 727   | 1853   | 1495  | 0        | 0        | 0        | 0        | 0            |
| N.S.       | 1       | 0.93  | 2.37   | 1.91  | 0.00     | 0.00     | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 1.590 | 12.375 | 3.167 | 0.000    | 0.000    | 0.000    | 0.000    | 0.000        |

| Problem 23 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas       | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------------|----------|----------|--------------|
| grade      | N/A     | A     | C      | B     | <b>F</b> | <b>F(-1)</b> | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD          | TBD      | TBD      | TBD          |
| size       | 1125    | 985   | 781    | 4476  | 0        | 0            | 0        | 0        | 0            |
| N.S.       | 1       | 0.88  | 0.69   | 3.98  | 0.00     | 0.00         | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 3.514 | 13.909 | 4.712 | 0.000    | 0.000        | 0.000    | 0.000    | 0.000        |

| Problem 24 | Optimal | Rubi  | MMA    | Maple  | Maxima   | Fricas | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|--------|----------|--------|----------|----------|--------------|
| grade      | N/A     | A     | C      | A      | <b>F</b> | B      | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes    | TBD      | TBD    | TBD      | TBD      | TBD          |
| size       | 859     | 765   | 1058   | 1141   | 0        | 2623   | 0        | 0        | 0            |
| N.S.       | 1       | 0.89  | 1.23   | 1.33   | 0.00     | 3.05   | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 1.716 | 15.269 | 12.568 | 0.000    | 0.134  | 0.000    | 0.000    | 0.000        |

| Problem 25 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------|----------|----------|--------------|
| grade      | N/A     | A     | C      | A     | <b>F</b> | B      | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD    | TBD      | TBD      | TBD          |
| size       | 628     | 576   | 766    | 799   | 0        | 1669   | 0        | 0        | 0            |
| N.S.       | 1       | 0.92  | 1.22   | 1.27  | 0.00     | 2.66   | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 0.918 | 12.987 | 2.938 | 0.000    | 0.108  | 0.000    | 0.000    | 0.000        |

| Problem 26 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------|----------|----------|--------------|
| grade      | N/A     | A     | C      | A     | <b>F</b> | B      | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD    | TBD      | TBD      | TBD          |
| size       | 481     | 446   | 597    | 593   | 0        | 1031   | 0        | 0        | 0            |
| N.S.       | 1       | 0.93  | 1.24   | 1.23  | 0.00     | 2.14   | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 0.660 | 11.657 | 1.671 | 0.000    | 0.100  | 0.000    | 0.000    | 0.000        |

| Problem 27 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------|----------|----------|--------------|
| grade      | N/A     | A     | C      | A     | <b>F</b> | A      | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD    | TBD      | TBD      | TBD          |
| size       | 398     | 382   | 497    | 509   | 0        | 656    | 0        | 0        | 0            |
| N.S.       | 1       | 0.96  | 1.25   | 1.28  | 0.00     | 1.65   | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 0.462 | 10.911 | 1.035 | 0.000    | 0.093  | 0.000    | 0.000    | 0.000        |

| Problem 28 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas       | Sympy        | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------------|--------------|----------|--------------|
| grade      | N/A     | A     | C      | B     | <b>F</b> | <b>F(-1)</b> | <b>F(-1)</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD          | TBD          | TBD      | TBD          |
| size       | 867     | 1045  | 1736   | 3241  | 0        | 0            | 0            | 0        | 0            |
| N.S.       | 1       | 1.21  | 2.00   | 3.74  | 0.00     | 0.00         | 0.00         | 0.00     | 0.00         |
| time (sec) | N/A     | 1.574 | 13.305 | 1.759 | 0.000    | 0.000        | 0.000        | 0.000    | 0.000        |

| Problem 29 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas       | Sympy        | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------------|--------------|----------|--------------|
| grade      | N/A     | A     | C      | B     | <b>F</b> | <b>F(-1)</b> | <b>F(-1)</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD          | TBD          | TBD      | TBD          |
| size       | 1301    | 2112  | 1116   | 8276  | 0        | 0            | 0            | 0        | 0            |
| N.S.       | 1       | 1.62  | 0.86   | 6.36  | 0.00     | 0.00         | 0.00         | 0.00     | 0.00         |
| time (sec) | N/A     | 3.431 | 15.222 | 3.607 | 0.000    | 0.000        | 0.000        | 0.000    | 0.000        |

| Problem 30 | Optimal | Rubi  | MMA    | Maple | Maxima       | Fricas       | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|--------------|--------------|----------|----------|--------------|
| grade      | N/A     | A     | C      | A     | <b>F(-2)</b> | <b>F(-1)</b> | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD          | TBD          | TBD      | TBD      | TBD          |
| size       | 273     | 273   | 310    | 365   | 0            | 0            | 0        | 0        | 0            |
| N.S.       | 1       | 1.00  | 1.14   | 1.34  | 0.00         | 0.00         | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 0.373 | 10.417 | 1.798 | 0.000        | 0.000        | 0.000    | 0.000    | 0.000        |

| Problem 31 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas       | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|--------------|----------|----------|--------------|
| grade      | N/A     | A     | C      | A     | <b>F</b> | <b>F(-1)</b> | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD          | TBD      | TBD      | TBD          |
| size       | 271     | 271   | 312    | 369   | 0        | 0            | 0        | 0        | 0            |
| N.S.       | 1       | 1.00  | 1.15   | 1.36  | 0.00     | 0.00         | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 0.423 | 10.408 | 4.428 | 0.000    | 0.000        | 0.000    | 0.000    | 0.000        |



| Problem 32 | Optimal | Rubi  | MMA    | Maple | Maxima   | Fricas   | Sympy    | Giac     | Mupad        |
|------------|---------|-------|--------|-------|----------|----------|----------|----------|--------------|
| grade      | N/A     | A     | C      | C     | <b>F</b> | <b>F</b> | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes    | Yes   | TBD      | TBD      | TBD      | TBD      | TBD          |
| size       | 106     | 106   | 74     | 93    | 0        | 0        | 0        | 0        | 0            |
| N.S.       | 1       | 1.00  | 0.70   | 0.88  | 0.00     | 0.00     | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 0.305 | 10.251 | 1.544 | 0.000    | 0.000    | 0.000    | 0.000    | 0.000        |

| Problem 33 | Optimal | Rubi  | MMA      | Maple    | Maxima   | Fricas   | Sympy        | Giac     | Mupad        |
|------------|---------|-------|----------|----------|----------|----------|--------------|----------|--------------|
| grade      | N/A     | A     | <b>F</b> | <b>F</b> | <b>F</b> | <b>F</b> | <b>F(-1)</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | N/A      | N/A      | TBD      | TBD      | TBD          | TBD      | TBD          |
| size       | 218     | 218   | 0        | 0        | 0        | 0        | 0            | 0        | 0            |
| N.S.       | 1       | 1.00  | 0.00     | 0.00     | 0.00     | 0.00     | 0.00         | 0.00     | 0.00         |
| time (sec) | N/A     | 0.482 | 0.000    | 0.000    | 0.000    | 0.000    | 0.000        | 0.000    | 0.000        |

| Problem 34 | Optimal | Rubi      | MMA   | Maple | Maxima   | Fricas | Sympy    | Giac  | Mupad |
|------------|---------|-----------|-------|-------|----------|--------|----------|-------|-------|
| grade      | N/A     | A         | C     | A     | <b>F</b> | B      | <b>F</b> | A     | B     |
| verified   | N/A     | <b>No</b> | Yes   | Yes   | TBD      | TBD    | TBD      | TBD   | TBD   |
| size       | 106     | 114       | 81    | 81    | 0        | 286    | 0        | 80    | 397   |
| N.S.       | 1       | 1.08      | 0.76  | 0.76  | 0.00     | 2.70   | 0.00     | 0.75  | 3.75  |
| time (sec) | N/A     | 0.378     | 0.024 | 1.168 | 0.000    | 0.331  | 0.000    | 0.309 | 8.632 |

| Problem 35 | Optimal | Rubi  | MMA   | Maple | Maxima   | Fricas | Sympy    | Giac     | Mupad        |
|------------|---------|-------|-------|-------|----------|--------|----------|----------|--------------|
| grade      | N/A     | A     | A     | A     | <b>F</b> | B      | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes   | Yes   | TBD      | TBD    | TBD      | TBD      | TBD          |
| size       | 145     | 191   | 139   | 126   | 0        | 603    | 0        | 0        | 0            |
| N.S.       | 1       | 1.32  | 0.96  | 0.87  | 0.00     | 4.16   | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 0.366 | 0.908 | 1.668 | 0.000    | 1.782  | 0.000    | 0.000    | 0.000        |

| Problem 36 | Optimal | Rubi  | MMA   | Maple | Maxima   | Fricas | Sympy    | Giac     | Mupad        |
|------------|---------|-------|-------|-------|----------|--------|----------|----------|--------------|
| grade      | N/A     | A     | C     | A     | <b>F</b> | B      | <b>F</b> | <b>F</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes   | Yes   | TBD      | TBD    | TBD      | TBD      | TBD          |
| size       | 239     | 239   | 151   | 351   | 0        | 669    | 0        | 0        | 0            |
| N.S.       | 1       | 1.00  | 0.63  | 1.47  | 0.00     | 2.80   | 0.00     | 0.00     | 0.00         |
| time (sec) | N/A     | 0.394 | 0.643 | 2.488 | 0.000    | 1.677  | 0.000    | 0.000    | 0.000        |

| Problem 37 | Optimal | Rubi  | MMA   | Maple | Maxima       | Fricas | Sympy    | Giac  | Mupad        |
|------------|---------|-------|-------|-------|--------------|--------|----------|-------|--------------|
| grade      | N/A     | A     | A     | A     | <b>F(-2)</b> | A      | <b>F</b> | A     | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes   | Yes   | TBD          | TBD    | TBD      | TBD   | TBD          |
| size       | 309     | 215   | 212   | 245   | 0            | 469    | 0        | 358   | 0            |
| N.S.       | 1       | 0.70  | 0.69  | 0.79  | 0.00         | 1.52   | 0.00     | 1.16  | 0.00         |
| time (sec) | N/A     | 0.438 | 0.689 | 0.726 | 0.000        | 0.325  | 0.000    | 0.310 | 0.000        |

| Problem 38 | Optimal | Rubi  | MMA   | Maple | Maxima       | Fricas | Sympy    | Giac  | Mupad        |
|------------|---------|-------|-------|-------|--------------|--------|----------|-------|--------------|
| grade      | N/A     | A     | A     | A     | <b>F(-2)</b> | A      | <b>F</b> | A     | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes   | Yes   | TBD          | TBD    | TBD      | TBD   | TBD          |
| size       | 283     | 189   | 214   | 191   | 0            | 361    | 0        | 262   | 0            |
| N.S.       | 1       | 0.67  | 0.76  | 0.67  | 0.00         | 1.28   | 0.00     | 0.93  | 0.00         |
| time (sec) | N/A     | 0.376 | 1.190 | 0.581 | 0.000        | 0.301  | 0.000    | 0.333 | 0.000        |

| Problem 39 | Optimal | Rubi  | MMA   | Maple | Maxima       | Fricas | Sympy    | Giac         | Mupad        |
|------------|---------|-------|-------|-------|--------------|--------|----------|--------------|--------------|
| grade      | N/A     | A     | A     | A     | <b>F(-2)</b> | A      | <b>F</b> | <b>F(-2)</b> | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes   | Yes   | TBD          | TBD    | TBD      | TBD          | TBD          |
| size       | 286     | 210   | 174   | 251   | 0            | 743    | 0        | 0            | 0            |
| N.S.       | 1       | 0.73  | 0.61  | 0.88  | 0.00         | 2.60   | 0.00     | 0.00         | 0.00         |
| time (sec) | N/A     | 0.525 | 0.537 | 0.574 | 0.000        | 0.733  | 0.000    | 0.000        | 0.000        |

| Problem 40 | Optimal | Rubi  | MMA   | Maple | Maxima       | Fricas | Sympy    | Giac  | Mupad        |
|------------|---------|-------|-------|-------|--------------|--------|----------|-------|--------------|
| grade      | N/A     | A     | A     | A     | <b>F(-2)</b> | A      | <b>F</b> | A     | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes   | Yes   | TBD          | TBD    | TBD      | TBD   | TBD          |
| size       | 294     | 212   | 171   | 229   | 0            | 731    | 0        | 238   | 0            |
| N.S.       | 1       | 0.72  | 0.58  | 0.78  | 0.00         | 2.49   | 0.00     | 0.81  | 0.00         |
| time (sec) | N/A     | 0.543 | 0.607 | 0.590 | 0.000        | 0.659  | 0.000    | 0.327 | 0.000        |

| Problem 41 | Optimal | Rubi  | MMA   | Maple | Maxima       | Fricas | Sympy        | Giac  | Mupad        |
|------------|---------|-------|-------|-------|--------------|--------|--------------|-------|--------------|
| grade      | N/A     | A     | A     | A     | <b>F(-2)</b> | A      | <b>F(-1)</b> | A     | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes   | Yes   | TBD          | TBD    | TBD          | TBD   | TBD          |
| size       | 288     | 200   | 173   | 214   | 0            | 749    | 0            | 360   | 0            |
| N.S.       | 1       | 0.69  | 0.60  | 0.74  | 0.00         | 2.60   | 0.00         | 1.25  | 0.00         |
| time (sec) | N/A     | 0.518 | 0.641 | 0.607 | 0.000        | 0.585  | 0.000        | 0.343 | 0.000        |

| Problem 42 | Optimal | Rubi  | MMA   | Maple | Maxima       | Fricas | Sympy        | Giac  | Mupad        |
|------------|---------|-------|-------|-------|--------------|--------|--------------|-------|--------------|
| grade      | N/A     | A     | A     | A     | <b>F(-2)</b> | A      | <b>F(-1)</b> | B     | <b>F(-1)</b> |
| verified   | N/A     | Yes   | Yes   | Yes   | TBD          | TBD    | TBD          | TBD   | TBD          |
| size       | 294     | 220   | 186   | 194   | 0            | 791    | 0            | 683   | 0            |
| N.S.       | 1       | 0.75  | 0.63  | 0.66  | 0.00         | 2.69   | 0.00         | 2.32  | 0.00         |
| time (sec) | N/A     | 0.541 | 0.877 | 0.614 | 0.000        | 0.595  | 0.000        | 0.351 | 0.000        |

## 2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. The column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio  $\frac{\text{number of rules}}{\text{integrand size}}$  is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [7] had the largest ratio of [.392857000000000012]

Table 2.1: Rubi specific breakdown of results for each integral

| #  | grade | number of steps used | number of unique rules | normalized antiderivative leaf size | integrand leaf size | $\frac{\text{number of rules}}{\text{integrand leaf size}}$ |
|----|-------|----------------------|------------------------|-------------------------------------|---------------------|---|
| 1  | A     | 2                    | 2                      | 1.94                                | 28                  | 0.071   |
| 2  | A     | 2                    | 2                      | 1.92                                | 28                  | 0.071   |
| 3  | A     | 2                    | 2                      | 1.56                                | 26                  | 0.077   |
| 4  | A     | 4                    | 4                      | 1.01                                | 19                  | 0.211   |
| 5  | A     | 4                    | 4                      | 1.04                                | 28                  | 0.143   |
| 6  | A     | 9                    | 9                      | 0.94                                | 28                  | 0.321   |
| 7  | A     | 11                   | 11                     | 0.90                                | 28                  | 0.393   |
| 8  | A     | 2                    | 2                      | 1.00                                | 28                  | 0.071   |
| 9  | A     | 2                    | 2                      | 1.00                                | 28                  | 0.071   |
| 10 | A     | 2                    | 2                      | 1.00                                | 26                  | 0.077   |
| 11 | A     | 6                    | 6                      | 1.00                                | 19                  | 0.316   |
| 12 | A     | 2                    | 2                      | 1.00                                | 28                  | 0.071   |
| 13 | A     | 2                    | 2                      | 1.00                                | 28                  | 0.071   |
| 14 | A     | 2                    | 2                      | 1.00                                | 28                  | 0.071   |
| 15 | A     | 2                    | 2                      | 1.00                                | 26                  | 0.077   |
| 16 | A     | 2                    | 2                      | 1.00                                | 27                  | 0.074   |
| 17 | A     | 7                    | 7                      | 0.91                                | 33                  | 0.212   |
| 18 | A     | 7                    | 7                      | 0.94                                | 33                  | 0.212   |
| 19 | A     | 5                    | 5                      | 0.97                                | 31                  | 0.161   |
| 20 | A     | 4                    | 4                      | 1.01                                | 24                  | 0.167   |
| 21 | A     | 4                    | 4                      | 1.03                                | 33                  | 0.121   |
| 22 | A     | 9                    | 9                      | 0.93                                | 33                  | 0.273   |

Continued on next page

Table 2.1 – continued from previous page

| #  | grade | number of steps used | number of unique rules | normalized antiderivative leaf size | integrand leaf size | $\frac{\text{number of rules}}{\text{integrand leaf size}}$ |
|----|-------|----------------------|------------------------|-------------------------------------|---------------------|---|
| 23 | A     | 11                   | 11                     | 0.88                                | 33                  | 0.333   |
| 24 | A     | 7                    | 7                      | 0.89                                | 33                  | 0.212   |
| 25 | A     | 6                    | 6                      | 0.92                                | 33                  | 0.182   |
| 26 | A     | 6                    | 6                      | 0.93                                | 31                  | 0.194   |
| 27 | A     | 6                    | 6                      | 0.96                                | 24                  | 0.250   |
| 28 | A     | 2                    | 2                      | 1.21                                | 33                  | 0.061   |
| 29 | A     | 2                    | 2                      | 1.62                                | 33                  | 0.061   |
| 30 | A     | 1                    | 1                      | 1.00                                | 41                  | 0.024   |
| 31 | A     | 1                    | 1                      | 1.00                                | 41                  | 0.024   |
| 32 | A     | 5                    | 5                      | 1.00                                | 31                  | 0.161   |
| 33 | A     | 2                    | 2                      | 1.00                                | 31                  | 0.065   |
| 34 | A     | 11                   | 10                     | 1.08                                | 28                  | 0.357   |
| 35 | A     | 4                    | 3                      | 1.32                                | 30                  | 0.100   |
| 36 | A     | 1                    | 1                      | 1.00                                | 30                  | 0.033   |
| 37 | A     | 9                    | 8                      | 0.70                                | 38                  | 0.211   |
| 38 | A     | 9                    | 8                      | 0.67                                | 37                  | 0.216   |
| 39 | A     | 12                   | 11                     | 0.73                                | 40                  | 0.275   |
| 40 | A     | 13                   | 12                     | 0.72                                | 40                  | 0.300   |
| 41 | A     | 11                   | 10                     | 0.69                                | 40                  | 0.250   |
| 42 | A     | 12                   | 11                     | 0.75                                | 40                  | 0.275   |

# CHAPTER 3

## LISTING OF INTEGRALS

|      |   |     |
|------|---|-----|
| 3.1  | $\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+cx^4}} dx$      | 39  |
| 3.2  | $\int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+cx^4}} dx$      | 48  |
| 3.3  | $\int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+cx^4}} dx$        | 55  |
| 3.4  | $\int \frac{A+Bx^2}{\sqrt{a+cx^4}} dx$                  | 61  |
| 3.5  | $\int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+cx^4}} dx$          | 67  |
| 3.6  | $\int \frac{A+Bx^2}{(d+ex^2)^2\sqrt{a+cx^4}} dx$        | 73  |
| 3.7  | $\int \frac{A+Bx^2}{(d+ex^2)^3\sqrt{a+cx^4}} dx$        | 82  |
| 3.8  | $\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+cx^4)^{3/2}} dx$     | 92  |
| 3.9  | $\int \frac{(A+Bx^2)(d+ex^2)^2}{(a+cx^4)^{3/2}} dx$     | 100 |
| 3.10 | $\int \frac{(A+Bx^2)(d+ex^2)}{(a+cx^4)^{3/2}} dx$       | 107 |
| 3.11 | $\int \frac{A+Bx^2}{(a+cx^4)^{3/2}} dx$                 | 113 |
| 3.12 | $\int \frac{A+Bx^2}{(d+ex^2)(a+cx^4)^{3/2}} dx$         | 119 |
| 3.13 | $\int \frac{A+Bx^2}{(d+ex^2)^2(a+cx^4)^{3/2}} dx$       | 126 |
| 3.14 | $\int \frac{A+Bx^2}{(d+ex^2)^3(a+cx^4)^{3/2}} dx$       | 133 |
| 3.15 | $\int \frac{(A+Bx^2)(d+ex^2)^q}{a+cx^4} dx$             | 141 |
| 3.16 | $\int \frac{2+x^2}{(1+x^2)\sqrt{2+3x^2+x^4}} dx$        | 145 |
| 3.17 | $\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+bx^2+cx^4}} dx$ | 150 |
| 3.18 | $\int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+bx^2+cx^4}} dx$ | 159 |
| 3.19 | $\int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+bx^2+cx^4}} dx$   | 167 |
| 3.20 | $\int \frac{A+Bx^2}{\sqrt{a+bx^2+cx^4}} dx$             | 175 |
| 3.21 | $\int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$     | 182 |
| 3.22 | $\int \frac{A+Bx^2}{(d+ex^2)^2\sqrt{a+bx^2+cx^4}} dx$   | 189 |

|      |  |     |
|------|--|-----|
| 3.23 | $\int \frac{A+Bx^2}{(d+ex^2)^3 \sqrt{a+bx^2+cx^4}} dx$               | 199 |
| 3.24 | $\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+bx^2+cx^4)^{3/2}} dx$             | 209 |
| 3.25 | $\int \frac{(A+Bx^2)(d+ex^2)^2}{(a+bx^2+cx^4)^{3/2}} dx$             | 218 |
| 3.26 | $\int \frac{(A+Bx^2)(d+ex^2)}{(a+bx^2+cx^4)^{3/2}} dx$               | 227 |
| 3.27 | $\int \frac{A+Bx^2}{(a+bx^2+cx^4)^{3/2}} dx$                         | 235 |
| 3.28 | $\int \frac{A+Bx^2}{(d+ex^2)(a+bx^2+cx^4)^{3/2}} dx$                 | 242 |
| 3.29 | $\int \frac{A+Bx^2}{(d+ex^2)^2 (a+bx^2+cx^4)^{3/2}} dx$              | 250 |
| 3.30 | $\int \frac{\sqrt{a+\sqrt{cx^2}}}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$    | 257 |
| 3.31 | $\int \frac{1+\sqrt{\frac{c}{a}}x^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$ | 263 |
| 3.32 | $\int \frac{946+315x^2}{(7+5x^2)\sqrt{2+3x^2+x^4}} dx$               | 269 |
| 3.33 | $\int \frac{(A+Bx^2)(d+ex^2)^9}{a+bx^2+cx^4} dx$                     | 275 |
| 3.34 | $\int \frac{x(1+2x^2)}{\sqrt{1+x^2(1+x^2+x^4)}} dx$                  | 279 |
| 3.35 | $\int \frac{\sqrt{a+bx^2+cx^4}}{ad-cdx^4} dx$                        | 287 |
| 3.36 | $\int \frac{\sqrt{a+bx^2-cx^4}}{ad+cdx^4} dx$                        | 293 |
| 3.37 | $\int x\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4} dx$                  | 299 |
| 3.38 | $\int \sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4} dx$                   | 307 |
| 3.39 | $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x} dx$         | 314 |
| 3.40 | $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^2} dx$       | 322 |
| 3.41 | $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx$       | 331 |
| 3.42 | $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^4} dx$       | 340 |

**3.1** 
$$\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+cx^4}} dx$$

|       |   |    |
|-------|---|----|
| 3.1.1 | Optimal result . . . . .                            | 39 |
| 3.1.2 | Mathematica [C] (verified) . . . . .                | 40 |
| 3.1.3 | Rubi [A] (verified) . . . . .                       | 41 |
| 3.1.4 | Maple [C] (verified) . . . . .                      | 43 |
| 3.1.5 | Fricas [A] (verification not implemented) . . . . . | 43 |
| 3.1.6 | Sympy [C] (verification not implemented) . . . . .  | 44 |
| 3.1.7 | Maxima [F] . . . . .                                | 46 |
| 3.1.8 | Giac [F] . . . . .                                  | 46 |
| 3.1.9 | Mupad [F(-1)] . . . . .                             | 47 |

**3.1.1 Optimal result**

Integrand size = 28, antiderivative size = 453

$$\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+cx^4}} dx$$

$$= \frac{e(21Bcd^2 + 21Acde - 5aBe^2) x\sqrt{a+cx^4}}{21c^2} + \frac{e^2(3Bd + Ae)x^3\sqrt{a+cx^4}}{5c}$$

$$+ \frac{Be^3x^5\sqrt{a+cx^4}}{7c} + \frac{(5Bcd^3 + 15Acd^2e - 9aBde^2 - 3aAe^3) x\sqrt{a+cx^4}}{5c^{3/2}(\sqrt{a} + \sqrt{cx^2})}$$

$$- \frac{\sqrt[4]{a}(5Bcd^3 + 15Acd^2e - 9aBde^2 - 3aAe^3) (\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{5c^{7/4}\sqrt{a+cx^4}}$$

$$+ \frac{(105Ac^2d^3 + 25a^2Be^3 - 105acde(Bd + Ae) - 63a^{3/2}\sqrt{ce^2(3Bd + Ae)} + 105\sqrt{ac^3/2}d^2(Bd + 3Ae)) (\sqrt{a+cx^4})}{210\sqrt[4]{ac^9/4}\sqrt{a+cx^4}}$$

---

3.1. 
$$\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+cx^4}} dx$$



output  $\frac{1}{21}e^{21Ac*d*e-5B*a*e^2+21B*c*d^2}xx(c*x^4+a)^{(1/2)}/c^2+1/5e^{2*(Ae+3B*d)}x^3(c*x^4+a)^{(1/2)}/c+1/7B*e^3x^5(c*x^4+a)^{(1/2)}/c+1/5*(-3A*a^3+15A*c*d^2e-9B*a*d*e^2+5B*c*d^3)xx(c*x^4+a)^{(1/2)}/c^{(3/2)}/(a^{(1/2)}+x^2*c^{(1/2)})-1/5a^{(1/4)}*(-3A*a^3+15A*c*d^2e-9B*a*d*e^2+5B*c*d^3)*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)})))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/c^{(7/4)}/(c*x^4+a)^{(1/2)}+1/210*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)})))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(105A*c^2*d^3+25a^2B*e^3-105a*c*d*e*(Ae+B*d)+105c^{(3/2)}*d^2*(3Ae+B*d)*a^{(1/2)}-63a^{(3/2)}*e^2*(Ae+3B*d)*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/a^{(1/4)}/c^{(9/4)}/(c*x^4+a)^{(1/2)}$

### 3.1.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.21 (sec) , antiderivative size = 217, normalized size of antiderivative = 0.48

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + cx^4}} dx$$

$$= \frac{ex(a + cx^4)(-25aBe^2 + 21Ace(5d + ex^2) + 3Bc(35d^2 + 21dex^2 + 5e^2x^4)) + 5(21Acd(cd^2 - ae^2) + aBe^2)}{\dots}$$

input `Integrate[((A + B*x^2)*(d + e*x^2)^3)/Sqrt[a + c*x^4],x]`

output  $(e*x*(a + c*x^4)*(-25*a*B*e^2 + 21*A*c*e*(5*d + e*x^2) + 3*B*c*(35*d^2 + 21*d*e*x^2 + 5*e^2*x^4)) + 5*(21*A*c*d*(c*d^2 - a*e^2) + a*B*e*(-21*c*d^2 + 5*a*e^2))*x*\text{Sqrt}[1 + (c*x^4)/a]*\text{Hypergeometric2F1}[1/4, 1/2, 5/4, -((c*x^4)/a)] + 7*c*(5*B*c*d^3 + 15*A*c*d^2*e - 9*a*B*d*e^2 - 3*a*A*e^3)*x^3*\text{Sqrt}[1 + (c*x^4)/a]*\text{Hypergeometric2F1}[1/2, 3/4, 7/4, -((c*x^4)/a)]/(105*c^2*\text{Sqrt}[a + c*x^4])$

---

3.1.  $\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+cx^4}} dx$

### 3.1.3 Rubi [A] (verified)

Time = 0.90 (sec) , antiderivative size = 878, normalized size of antiderivative = 1.94, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {2259, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + cx^4}} dx \\
 & \quad \downarrow \text{2259} \\
 & \int \left( \frac{d^2 x^2 (3Ae + Bd)}{\sqrt{a + cx^4}} + \frac{e^2 x^6 (Ae + 3Bd)}{\sqrt{a + cx^4}} + \frac{3dex^4 (Ae + Bd)}{\sqrt{a + cx^4}} + \frac{Ad^3}{\sqrt{a + cx^4}} + \frac{Be^3 x^8}{\sqrt{a + cx^4}} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{Be^3 \sqrt{cx^4 + ax^5}}{7c} + \frac{e^2 (3Bd + Ae) \sqrt{cx^4 + ax^3}}{5c} - \frac{5aBe^3 \sqrt{cx^4 + ax}}{21c^2} + \frac{de(Bd + Ae) \sqrt{cx^4 + ax}}{c} - \\
 & \quad \frac{3ae^2 (3Bd + Ae) \sqrt{cx^4 + ax}}{5c^{3/2} (\sqrt{cx^2 + \sqrt{a}})} + \frac{d^2 (Bd + 3Ae) \sqrt{cx^4 + ax}}{\sqrt{c} (\sqrt{cx^2 + \sqrt{a}})} + \\
 & \quad \frac{3a^{5/4} e^2 (3Bd + Ae) (\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{5c^{7/4} \sqrt{cx^4 + a}} - \\
 & \quad \frac{\sqrt[4]{a} d^2 (Bd + 3Ae) (\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{c^{3/4} \sqrt{cx^4 + a}} + \\
 & \quad \frac{Ad^3 (\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2\sqrt[4]{a} \sqrt[4]{c} \sqrt{cx^4 + a}} + \\
 & \quad \frac{5a^{7/4} Be^3 (\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{42c^{9/4} \sqrt{cx^4 + a}} - \\
 & \quad \frac{a^{3/4} de (Bd + Ae) (\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2c^{5/4} \sqrt{cx^4 + a}} \\
 & \quad \frac{3a^{5/4} e^2 (3Bd + Ae) (\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{10c^{7/4} \sqrt{cx^4 + a}} + \\
 & \quad \frac{\sqrt[4]{a} d^2 (Bd + 3Ae) (\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2c^{3/4} \sqrt{cx^4 + a}}
 \end{aligned}$$

---

3.1.  $\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+cx^4}} dx$

input `Int[((A + B*x^2)*(d + e*x^2)^3)/Sqrt[a + c*x^4],x]`

output `(-5*a*B*e^3*x*Sqrt[a + c*x^4])/(21*c^2) + (d*e*(B*d + A*e)*x*Sqrt[a + c*x^4])/c + (e^2*(3*B*d + A*e)*x^3*Sqrt[a + c*x^4])/(5*c) + (B*e^3*x^5*Sqrt[a + c*x^4])/(7*c) - (3*a*e^2*(3*B*d + A*e)*x*Sqrt[a + c*x^4])/(5*c^(3/2)*(Sqrt[a + Sqrt[c]*x^2])) + (d^2*(B*d + 3*A*e)*x*Sqrt[a + c*x^4])/(Sqrt[c]*(Sqrt[a + Sqrt[c]*x^2])) + (3*a^(5/4)*e^2*(3*B*d + A*e)*(Sqrt[a + Sqrt[c]*x^2])*Sqrt[(a + c*x^4)/(Sqrt[a + Sqrt[c]*x^2]^2)*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]])/(5*c^(7/4)*Sqrt[a + c*x^4]) - (a^(1/4)*d^2*(B*d + 3*A*e)*(Sqrt[a + Sqrt[c]*x^2])*Sqrt[(a + c*x^4)/(Sqrt[a + Sqrt[c]*x^2]^2)*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]])/(c^(3/4)*Sqrt[a + c*x^4]) + (A*d^3*(Sqrt[a + Sqrt[c]*x^2])*Sqrt[(a + c*x^4)/(Sqrt[a + Sqrt[c]*x^2]^2)*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]])/(2*a^(1/4)*c^(1/4)*Sqrt[a + c*x^4]) + (5*a^(7/4)*B*e^3*(Sqrt[a + Sqrt[c]*x^2])*Sqrt[(a + c*x^4)/(Sqrt[a + Sqrt[c]*x^2]^2)*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]])/(42*c^(9/4)*Sqrt[a + c*x^4]) - (a^(3/4)*d*e*(B*d + A*e)*(Sqrt[a + Sqrt[c]*x^2])*Sqrt[(a + c*x^4)/(Sqrt[a + Sqrt[c]*x^2]^2)*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]])/(2*c^(5/4)*Sqrt[a + c*x^4]) - (3*a^(5/4)*e^2*(3*B*d + A*e)*(Sqrt[a + Sqrt[c]*x^2])*Sqrt[(a + c*x^4)/(Sqrt[a + Sqrt[c]*x^2]^2)*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]])/(10*c^(7/4)*Sqrt[a + c*x^4]) + (a^(1/4)*d^2*(B*d + 3*A*e)*(Sqrt[a + Sqrt[c]*x^2])*Sqrt[(a + c*x^4)/(Sqrt[a + Sqrt[c]*x^2]^2)*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]])/(2*...`

### 3.1.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2259 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x] && IntegerQ[p + 1/2] && IntegerQ[q]`

$$3.1. \int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+cx^4}} dx$$

### 3.1.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 5.69 (sec) , antiderivative size = 327, normalized size of antiderivative = 0.72

| method   | result  |
|----------|---|
| elliptic | $\frac{B e^3 x^5 \sqrt{c x^4 + a}}{7c} + \frac{(e^3 A + 3B e^2 d) x^3 \sqrt{c x^4 + a}}{5c} + \frac{(3A d e^2 + 3B d^2 e - \frac{5B e^3 a}{7c}) x \sqrt{c x^4 + a}}{3c} + \frac{\left( A d^3 - \frac{(3A d e^2 + 3B d^2 e - \frac{5B e^3 a}{7c}) a}{3c} \right) \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}}$  |
| risch    | $\frac{e x (15 B e^2 x^4 c + 21 A c e^2 x^2 + 63 B c d e x^2 + 105 A c d e - 25 B a e^2 + 105 B c d^2) \sqrt{c x^4 + a}}{105 c^2} - \frac{25 a^2 B e^3 \sqrt{1 - \frac{i\sqrt{c} x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c} x^2}{\sqrt{a}}} F\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{c x^4 + a}}$   |
| default  | $\frac{A d^3 \sqrt{1 - \frac{i\sqrt{c} x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c} x^2}{\sqrt{a}}} F\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{c x^4 + a}} + B e^3 \left( \frac{x^5 \sqrt{c x^4 + a}}{7c} - \frac{5 a x \sqrt{c x^4 + a}}{21 c^2} + \frac{5 a^2 \sqrt{1 - \frac{i\sqrt{c} x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c} x^2}{\sqrt{a}}} F\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{21 c^2 \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{c x^4 + a}} \right)$ |

input `int((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(1/2),x,method=_RETURNVERBOSE)`

output `1/7*B*e^3*x^5*(c*x^4+a)^(1/2)/c+1/5*(A*e^3+3*B*d*e^2)/c*x^3*(c*x^4+a)^(1/2)+1/3*(3*A*d*e^2+3*B*d^2*e-5/7*B*e^3/c*a)/c*x*(c*x^4+a)^(1/2)+(A*d^3-1/3*(3*A*d*e^2+3*B*d^2*e-5/7*B*e^3/c*a)/c*a)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+I*(3*A*d^2*e+d^3*B-3/5*(A*e^3+3*B*d*e^2)/c*a)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)-EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I))`

### 3.1.5 Fracas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 283, normalized size of antiderivative = 0.62

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + cx^4}} dx$$

$$= \frac{21(5Bacd^3 + 15Aacd^2e - 9Ba^2de^2 - 3Aa^2e^3)\sqrt{cx}\left(-\frac{a}{c}\right)^{\frac{3}{4}} E\left(\arcsin\left(\frac{\left(-\frac{a}{c}\right)^{\frac{1}{4}}}{x}\right) \mid -1\right) - (105(3A + B)ac)}{\dots}$$

input `integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(1/2),x, algorithm="fracas")`

3.1.  $\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+cx^4}} dx$

```
output 1/105*(21*(5*B*a*c*d^3 + 15*A*a*c*d^2*e - 9*B*a^2*d*e^2 - 3*A*a^2*e^3)*sqrt(c)*x*(-a/c)^(3/4)*elliptic_e(arcsin((-a/c)^(1/4)/x), -1) - (105*(3*A + B)*a*c*d^2*e - (63*A + 25*B)*a^2*e^3 + 105*(B*a*c - A*c^2)*d^3 - 21*(9*B*a^2 - 5*A*a*c)*d*e^2)*sqrt(c)*x*(-a/c)^(3/4)*elliptic_f(arcsin((-a/c)^(1/4)/x), -1) + (15*B*a*c*e^3*x^6 + 105*B*a*c*d^3 + 315*A*a*c*d^2*e - 189*B*a^2*d*e^2 - 63*A*a^2*e^3 + 21*(3*B*a*c*d*e^2 + A*a*c*e^3)*x^4 + 5*(21*B*a*c*d^2*e + 21*A*a*c*d*e^2 - 5*B*a^2*e^3)*x^2)*sqrt(c*x^4 + a))/(a*c^2*x)
```

### 3.1.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

---

3.1.  $\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+cx^4}} dx$

Time = 3.49 (sec) , antiderivative size = 364, normalized size of antiderivative = 0.80

$$\begin{aligned}
 \int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + cx^4}} dx = & \frac{Ad^3 x \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{5}{4}\right)} \\
 & + \frac{3Ad^2 ex^3 \Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{7}{4}\right)} \\
 & + \frac{3Ade^2 x^5 \Gamma\left(\frac{5}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{5}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{9}{4}\right)} \\
 & + \frac{Ae^3 x^7 \Gamma\left(\frac{7}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{7}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{11}{4}\right)} \\
 & + \frac{Bd^3 x^3 \Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{7}{4}\right)} \\
 & + \frac{3Bd^2 ex^5 \Gamma\left(\frac{5}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{5}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{9}{4}\right)} \\
 & + \frac{3Bde^2 x^7 \Gamma\left(\frac{7}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{7}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{11}{4}\right)} \\
 & + \frac{Be^3 x^9 \Gamma\left(\frac{9}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{9}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{13}{4}\right)}
 \end{aligned}$$

input `integrate((B*x**2+A)*(e*x**2+d)**3/(c*x**4+a)**(1/2),x)`

---

3.1.  $\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+cx^4}} dx$

```
output A*d**3*x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4
*sqrt(a)*gamma(5/4)) + 3*A*d**2*e*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,)
, c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4)) + 3*A*d*e**2*x**5*gamma
(5/4)*hyper((1/2, 5/4), (9/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma
(9/4)) + A*e**3*x**7*gamma(7/4)*hyper((1/2, 7/4), (11/4,), c*x**4*exp_pola
r(I*pi)/a)/(4*sqrt(a)*gamma(11/4)) + B*d**3*x**3*gamma(3/4)*hyper((1/2, 3/
4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4)) + 3*B*d**2*e*
x**5*gamma(5/4)*hyper((1/2, 5/4), (9/4,), c*x**4*exp_polar(I*pi)/a)/(4*sq
r(a)*gamma(9/4)) + 3*B*d*e**2*x**7*gamma(7/4)*hyper((1/2, 7/4), (11/4,), c
*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(11/4)) + B*e**3*x**9*gamma(9/4)*
hyper((1/2, 9/4), (13/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(13/4
))
```

### 3.1.7 Maxima [F]

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^3}{\sqrt{cx^4 + a}} dx$$

```
input integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(1/2),x, algorithm="maxima")
```

```
output integrate((B*x^2 + A)*(e*x^2 + d)^3/sqrt(c*x^4 + a), x)
```

### 3.1.8 Giac [F]

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^3}{\sqrt{cx^4 + a}} dx$$

```
input integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(1/2),x, algorithm="giac")
```

```
output integrate((B*x^2 + A)*(e*x^2 + d)^3/sqrt(c*x^4 + a), x)
```

**3.1.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^3}{\sqrt{cx^4 + a}} dx$$

input `int(((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(1/2),x)`output `int(((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(1/2), x)`



### 3.2 $\int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+cx^4}} dx$

|       |   |    |
|-------|---|----|
| 3.2.1 | Optimal result . . . . .                            | 48 |
| 3.2.2 | Mathematica [C] (verified) . . . . .                | 49 |
| 3.2.3 | Rubi [A] (verified) . . . . .                       | 49 |
| 3.2.4 | Maple [C] (verified) . . . . .                      | 51 |
| 3.2.5 | Fricas [A] (verification not implemented) . . . . . | 52 |
| 3.2.6 | Sympy [C] (verification not implemented) . . . . .  | 52 |
| 3.2.7 | Maxima [F] . . . . .                                | 54 |
| 3.2.8 | Giac [F] . . . . .                                  | 54 |
| 3.2.9 | Mupad [F(-1)] . . . . .                             | 54 |

#### 3.2.1 Optimal result

Integrand size = 28, antiderivative size = 367

$$\int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+cx^4}} dx$$

$$= \frac{e(2Bd+ Ae)x\sqrt{a+cx^4}}{3c} + \frac{Be^2x^3\sqrt{a+cx^4}}{5c} + \frac{(5Bcd^2+ 10Acde- 3aBe^2)x\sqrt{a+cx^4}}{5c^{3/2}(\sqrt{a}+\sqrt{cx^2})}$$

$$- \frac{\sqrt[4]{a}(5Bcd^2+ 10Acde- 3aBe^2)(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)}{5c^{7/4}\sqrt{a+cx^4}}$$

$$+ \frac{(15Ac^{3/2}d^2- 9a^{3/2}Be^2- 5a\sqrt{ce}(2Bd+ Ae)+ 15\sqrt{acd}(Bd+ 2Ae))(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}\text{EllipticE}}{30\sqrt[4]{ac^{7/4}}\sqrt{a+cx^4}}$$

output

```
1/3*e*(A*e+2*B*d)*x*(c*x^4+a)^(1/2)/c+1/5*B*e^2*x^3*(c*x^4+a)^(1/2)/c+1/5*
(10*A*c*d*e-3*B*a*e^2+5*B*c*d^2)*x*(c*x^4+a)^(1/2)/c^(3/2)/(a^(1/2)+x^2*c^(
1/2))-1/5*a^(1/4)*(10*A*c*d*e-3*B*a*e^2+5*B*c*d^2)*(cos(2*arctan(c^(1/4)*
x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arct
an(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1
/2)+x^2*c^(1/2)))^(1/2)/c^(7/4)/(c*x^4+a)^(1/2)+1/30*(cos(2*arctan(c^(1/
4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*a
rctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(15*A*c^(3/2)*d^2-9*a^(3/2)*B*e^2+1
5*c*d*(2*A*e+B*d)*a^(1/2)-5*a*e*(A*e+2*B*d)*c^(1/2))*(a^(1/2)+x^2*c^(1/2)
)*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(1/4)/c^(7/4)/(c*x^4+a)^(1/2)
```

### 3.2.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.12 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.43

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + cx^4}} dx$$

$$= \frac{ex(10Bd + 5Ae + 3Bex^2)(a + cx^4) - 5(-3Acd^2 + 2aBde + aAe^2)x\sqrt{1 + \frac{cx^4}{a}} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -\left(\frac{cx^4}{a}\right)\right) + (5Bc^2d^2 + 10Acd^2e - 3a^2B^2e^2)x^3\sqrt{1 + \frac{cx^4}{a}} \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -\left(\frac{cx^4}{a}\right)\right]}{15c\sqrt{a + cx^4}}$$

input `Integrate[((A + B*x^2)*(d + e*x^2)^2)/Sqrt[a + c*x^4],x]`

output `(e*x*(10*B*d + 5*A*e + 3*B*e*x^2)*(a + c*x^4) - 5*(-3*A*c*d^2 + 2*a*B*d*e + a*A*e^2)*x*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + (5*B*c*d^2 + 10*A*c*d*e - 3*a*B*e^2)*x^3*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/2, 3/4, 7/4, -((c*x^4)/a)])/(15*c*Sqrt[a + c*x^4])`

### 3.2.3 Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 706, normalized size of antiderivative = 1.92, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {2259, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + cx^4}} dx$$

$$\downarrow 2259$$

$$\int \left( \frac{ex^4(Ae + 2Bd)}{\sqrt{a + cx^4}} + \frac{dx^2(2Ae + Bd)}{\sqrt{a + cx^4}} + \frac{Ad^2}{\sqrt{a + cx^4}} + \frac{Be^2x^6}{\sqrt{a + cx^4}} \right) dx$$

$$\downarrow 2009$$

---

3.2.  $\int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+cx^4}} dx$

$$\begin{aligned}
& \frac{a^{3/4}e(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} (Ae + 2Bd) \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{6c^{5/4}\sqrt{a+cx^4}} \\
& \frac{3a^{5/4}Be^2(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{10c^{7/4}\sqrt{a+cx^4}} + \\
& \frac{3a^{5/4}Be^2(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{5c^{7/4}\sqrt{a+cx^4}} + \\
& \frac{\sqrt[4]{ad}(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} (2Ae + Bd) \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2c^{3/4}\sqrt{a+cx^4}} - \\
& \frac{\sqrt[4]{ad}(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} (2Ae + Bd) E\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{c^{3/4}\sqrt{a+cx^4}} + \\
& \frac{Ad^2(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2\sqrt[4]{a}\sqrt[4]{c}\sqrt{a+cx^4}} + \frac{ex\sqrt{a+cx^4}(Ae + 2Bd)}{3c} + \\
& \frac{dx\sqrt{a+cx^4}(2Ae + Bd)}{\sqrt{c}(\sqrt{a} + \sqrt{cx^2})} - \frac{3aBe^2x\sqrt{a+cx^4}}{5c^{3/2}(\sqrt{a} + \sqrt{cx^2})} + \frac{Be^2x^3\sqrt{a+cx^4}}{5c}
\end{aligned}$$

input `Int[((A + B*x^2)*(d + e*x^2)^2)/Sqrt[a + c*x^4], x]`

output `(e*(2*B*d + A*e)*x*Sqrt[a + c*x^4])/(3*c) + (B*e^2*x^3*Sqrt[a + c*x^4])/(5*c) - (3*a*B*e^2*x*Sqrt[a + c*x^4])/(5*c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2)) + (d*(B*d + 2*A*e)*x*Sqrt[a + c*x^4])/(Sqrt[c]*(Sqrt[a] + Sqrt[c]*x^2)) + (3*a^(5/4)*B*e^2*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(5*c^(7/4)*Sqrt[a + c*x^4]) - (a^(1/4)*d*(B*d + 2*A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(c^(3/4)*Sqrt[a + c*x^4]) + (A*d^2*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*c^(1/4)*Sqrt[a + c*x^4]) - (3*a^(5/4)*B*e^2*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(10*c^(7/4)*Sqrt[a + c*x^4]) - (a^(3/4)*e*(2*B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(6*c^(5/4)*Sqrt[a + c*x^4]) + (a^(1/4)*d*(B*d + 2*A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*c^(3/4)*Sqrt[a + c*x^4])`

3.2.  $\int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+cx^4}} dx$

### 3.2.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2259 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x] && IntegerQ[p + 1/2] && IntegerQ[q]`

### 3.2.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 3.02 (sec) , antiderivative size = 258, normalized size of antiderivative = 0.70

| method   | result   |
|----------|--|
| elliptic | $\frac{B e^2 x^3 \sqrt{c x^4 + a}}{5c} + \frac{(A e^2 + 2B e d) x \sqrt{c x^4 + a}}{3c} + \frac{\left( d^2 A - \frac{a(A e^2 + 2B e d)}{3c} \right) \sqrt{1 - \frac{i\sqrt{c} x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c} x^2}{\sqrt{a}}} F\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{c x^4 + a}} + \frac{i(2A e d + B d^2)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{c x^4 + a}}$  |
| risch    | $\frac{e x (3B e x^2 + 5A e + 10B d) \sqrt{c x^4 + a}}{15c} - \frac{5a A e^2 \sqrt{1 - \frac{i\sqrt{c} x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c} x^2}{\sqrt{a}}} F\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{c x^4 + a}} - \frac{15A c d^2 \sqrt{1 - \frac{i\sqrt{c} x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c} x^2}{\sqrt{a}}} F\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{c x^4 + a}} + \frac{10a}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{c x^4 + a}}$                       |
| default  | $\frac{d^2 A \sqrt{1 - \frac{i\sqrt{c} x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c} x^2}{\sqrt{a}}} F\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{c x^4 + a}} + B e^2 \left( \frac{x^3 \sqrt{c x^4 + a}}{5c} - \frac{3i a^{\frac{3}{2}} \sqrt{1 - \frac{i\sqrt{c} x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c} x^2}{\sqrt{a}}} \left( F\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) - E\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) \right)}{5c^{\frac{3}{2}} \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{c x^4 + a}} \right)$ |

input `int((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(1/2),x,method=_RETURNVERBOSE)`

output `1/5*B*e^2*x^3*(c*x^4+a)^(1/2)/c+1/3*(A*e^2+2*B*d*e)/c*x*(c*x^4+a)^(1/2)+(d^2*A-1/3*a/c*(A*e^2+2*B*d*e))/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+I*(2*A*e*d+B*d^2-3/5*a/c*B*e^2)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)-EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I))`

3.2. 
$$\int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+cx^4}} dx$$

### 3.2.5 Fracas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 206, normalized size of antiderivative = 0.56

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + cx^4}} dx$$

$$= \frac{3(5Bacd^2 + 10Aacde - 3Ba^2e^2)\sqrt{cx}\left(-\frac{a}{c}\right)^{\frac{3}{4}} E\left(\arcsin\left(\frac{\left(-\frac{a}{c}\right)^{\frac{1}{4}}}{x}\right) \mid -1\right) - (10(3A + B)acde + 15(Bac -$$

```
input integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(1/2),x, algorithm="fricas")
```

```
output 1/15*(3*(5*B*a*c*d^2 + 10*A*a*c*d*e - 3*B*a^2*e^2)*sqrt(c)*x*(-a/c)^(3/4)*
elliptic_e(arcsin((-a/c)^(1/4)/x), -1) - (10*(3*A + B)*a*c*d*e + 15*(B*a*c
- A*c^2)*d^2 - (9*B*a^2 - 5*A*a*c)*e^2)*sqrt(c)*x*(-a/c)^(3/4)*elliptic_f
(arcsin((-a/c)^(1/4)/x), -1) + (3*B*a*c*e^2*x^4 + 15*B*a*c*d^2 + 30*A*a*c*
d*e - 9*B*a^2*e^2 + 5*(2*B*a*c*d*e + A*a*c*e^2)*x^2)*sqrt(c*x^4 + a))/(a*c
^2*x)
```

### 3.2.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

---

3.2.  $\int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+cx^4}} dx$

Time = 2.59 (sec) , antiderivative size = 262, normalized size of antiderivative = 0.71

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + cx^4}} dx = \frac{Ad^2x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{5}{4}\right)} + \frac{Ade^2x^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{2\sqrt{a}\Gamma\left(\frac{7}{4}\right)}$$

$$+ \frac{Ae^2x^5\Gamma\left(\frac{5}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{5}{4} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{9}{4}\right)}$$

$$+ \frac{Bd^2x^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{7}{4}\right)}$$

$$+ \frac{Bde^2x^5\Gamma\left(\frac{5}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{5}{4} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{2\sqrt{a}\Gamma\left(\frac{9}{4}\right)}$$

$$+ \frac{Be^2x^7\Gamma\left(\frac{7}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{7}{4} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{11}{4}\right)}$$

input `integrate((B*x**2+A)*(e*x**2+d)**2/(c*x**4+a)**(1/2),x)`

output `A*d**2*x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(5/4)) + A*d*e*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(2*sqrt(a)*gamma(7/4)) + A*e**2*x**5*gamma(5/4)*hyper((1/2, 5/4), (9/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(9/4)) + B*d**2*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4)) + B*d*e*x**5*gamma(5/4)*hyper((1/2, 5/4), (9/4,), c*x**4*exp_polar(I*pi)/a)/(2*sqrt(a)*gamma(9/4)) + B*e**2*x**7*gamma(7/4)*hyper((1/2, 7/4), (11/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(11/4))`

### 3.2.7 Maxima [F]

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^2}{\sqrt{cx^4 + a}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(1/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^2/sqrt(c*x^4 + a), x)`

### 3.2.8 Giac [F]

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^2}{\sqrt{cx^4 + a}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(1/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^2/sqrt(c*x^4 + a), x)`

### 3.2.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^2}{\sqrt{cx^4 + a}} dx$$

input `int(((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(1/2),x)`

output `int(((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(1/2), x)`

### 3.3 $\int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+cx^4}} dx$

|       |   |    |
|-------|---|----|
| 3.3.1 | Optimal result . . . . .                            | 55 |
| 3.3.2 | Mathematica [C] (verified) . . . . .                | 56 |
| 3.3.3 | Rubi [A] (verified) . . . . .                       | 56 |
| 3.3.4 | Maple [C] (verified) . . . . .                      | 58 |
| 3.3.5 | Fricas [A] (verification not implemented) . . . . . | 58 |
| 3.3.6 | Sympy [C] (verification not implemented) . . . . .  | 59 |
| 3.3.7 | Maxima [F] . . . . .                                | 59 |
| 3.3.8 | Giac [F] . . . . .                                  | 60 |
| 3.3.9 | Mupad [F(-1)] . . . . .                             | 60 |

#### 3.3.1 Optimal result

Integrand size = 26, antiderivative size = 277

$$\int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+cx^4}} dx = \frac{Bex\sqrt{a+cx^4}}{3c} + \frac{(Bd+ Ae)x\sqrt{a+cx^4}}{\sqrt{c}(\sqrt{a}+\sqrt{cx^2})}$$

$$- \frac{\sqrt[4]{a}(Bd+ Ae)(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)}{c^{3/4}\sqrt{a+cx^4}}$$

$$+ \frac{\sqrt[4]{a}\left(3\sqrt{c}(Bd+ Ae) + \frac{3Acd-aBe}{\sqrt{a}}\right)(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{6c^{5/4}\sqrt{a+cx^4}}$$

```
output 1/3*B*e*x*(c*x^4+a)^(1/2)/c+(A*e+B*d)*x*(c*x^4+a)^(1/2)/c^(1/2)/(a^(1/2)+x
^2*c^(1/2))-a^(1/4)*(A*e+B*d)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/c
os(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4)))
,1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1
/2)/c^(3/4)/(c*x^4+a)^(1/2)+1/6*a^(1/4)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^
2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x
/a^(1/4))),1/2*2^(1/2))*((3*A*c*d-B*a*e)/a^(1/2)+3*(A*e+B*d)*c^(1/2))*(a^(
1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(5/4)/(c*x^4
+a)^(1/2)
```



### 3.3.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.07 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.44

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + cx^4}} dx$$

$$= \frac{Bex(a + cx^4) + (3Acd - aBe)x\sqrt{1 + \frac{cx^4}{a}} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -\frac{cx^4}{a}\right) + c(Bd + Ae)x^3\sqrt{1 + \frac{cx^4}{a}}}{3c\sqrt{a + cx^4}}$$

input `Integrate[((A + B*x^2)*(d + e*x^2))/Sqrt[a + c*x^4],x]`

output `(B*e*x*(a + c*x^4) + (3*A*c*d - a*B*e)*x*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + c*(B*d + A*e)*x^3*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/2, 3/4, 7/4, -((c*x^4)/a)]/(3*c*Sqrt[a + c*x^4])`

### 3.3.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 432, normalized size of antiderivative = 1.56, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$ , Rules used = {2259, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + cx^4}} dx$$

$$\downarrow \text{2259}$$

$$\int \left( \frac{x^2(Ae + Bd)}{\sqrt{a + cx^4}} + \frac{Ad}{\sqrt{a + cx^4}} + \frac{Bex^4}{\sqrt{a + cx^4}} \right) dx$$

$$\downarrow \text{2009}$$

$$\begin{aligned}
& \frac{a^{3/4} B e(\sqrt{a} + \sqrt{c x^2}) \sqrt{\frac{a+c x^4}{(\sqrt{a}+\sqrt{c x^2})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{c x}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{6 c^{5/4} \sqrt{a+c x^4}} + \\
& \frac{\sqrt[4]{a}(\sqrt{a} + \sqrt{c x^2}) \sqrt{\frac{a+c x^4}{(\sqrt{a}+\sqrt{c x^2})^2}} (A e+B d) \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{c x}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2 c^{3/4} \sqrt{a+c x^4}} - \\
& \frac{\sqrt[4]{a}(\sqrt{a} + \sqrt{c x^2}) \sqrt{\frac{a+c x^4}{(\sqrt{a}+\sqrt{c x^2})^2}} (A e+B d) E\left(2 \arctan\left(\frac{\sqrt[4]{c x}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{c^{3/4} \sqrt{a+c x^4}} + \\
& \frac{A d(\sqrt{a} + \sqrt{c x^2}) \sqrt{\frac{a+c x^4}{(\sqrt{a}+\sqrt{c x^2})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{c x}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2 \sqrt[4]{a} \sqrt[4]{c} \sqrt{a+c x^4}} + \frac{x \sqrt{a+c x^4} (A e+B d)}{\sqrt{c}(\sqrt{a} + \sqrt{c x^2})} + \\
& \frac{B e x \sqrt{a+c x^4}}{3 c}
\end{aligned}$$

input `Int[((A + B*x^2)*(d + e*x^2))/Sqrt[a + c*x^4],x]`

output `(B*e*x*Sqrt[a + c*x^4])/(3*c) + ((B*d + A*e)*x*Sqrt[a + c*x^4])/(Sqrt[c]*(Sqrt[a] + Sqrt[c]*x^2)) - (a^(1/4)*(B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(c^(3/4)*Sqrt[a + c*x^4]) + (A*d*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*c^(1/4)*Sqrt[a + c*x^4]) - (a^(3/4)*B*e*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(6*c^(5/4)*Sqrt[a + c*x^4]) + (a^(1/4)*(B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*c^(3/4)*Sqrt[a + c*x^4])`

### 3.3.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2259 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x] && IntegerQ[p + 1/2] && IntegerQ[q]`

### 3.3.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.20 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.73

| method   | result   |
|----------|--|
| elliptic | $\frac{Bex\sqrt{cx^4+a}}{3c} + \frac{(Ad - \frac{aeB}{3c})\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}} F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) + \frac{i(Ae+Bd)\sqrt{a}\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}\sqrt{c}} \left(F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) - E\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)\right)$              |
| default  | $\frac{Ad\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}} F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) + eB\left(\frac{x\sqrt{cx^4+a}}{3c} - \frac{a\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}}{3c\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}} F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)\right) + \frac{i(Ae+Bd)\sqrt{a}}{\sqrt{c}}$  |
| risch    | $\frac{Bex\sqrt{cx^4+a}}{3c} + \frac{3Acd\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}} F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) - \frac{Bae\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}} F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) + \frac{i(3Ace+3Bcd)\sqrt{a}\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}}{3c\sqrt{c}}$ |

input `int((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(1/2),x,method=_RETURNVERBOSE)`

output `1/3*B*e*x*(c*x^4+a)^(1/2)/c+(A*d-1/3*a/c*e*B)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+I*(A*e+B*d)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)-EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I))`

### 3.3.5 Fracas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.46

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + cx^4}} dx$$

$$= \frac{3(Bad + Aae)\sqrt{cx}\left(-\frac{a}{c}\right)^{\frac{3}{4}} E\left(\arcsin\left(\frac{\left(-\frac{a}{c}\right)^{\frac{1}{4}}}{x}\right) \mid -1\right) - ((3A + B)ae + 3(Ba - Ac)d)\sqrt{cx}\left(-\frac{a}{c}\right)^{\frac{3}{4}} F\left(\arcsin\left(\frac{\left(-\frac{a}{c}\right)^{\frac{1}{4}}}{x}\right) \mid -1\right)}{3ac}$$

input `integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="fricas")`

```
output 1/3*(3*(B*a*d + A*a*e)*sqrt(c)*x*(-a/c)^(3/4)*elliptic_e(arcsin((-a/c)^(1/4)/x), -1) - ((3*A + B)*a*e + 3*(B*a - A*c)*d)*sqrt(c)*x*(-a/c)^(3/4)*elliptic_f(arcsin((-a/c)^(1/4)/x), -1) + (B*a*e*x^2 + 3*B*a*d + 3*A*a*e)*sqrt(c*x^4 + a)/(a*c*x)
```

### 3.3.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.77 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.60

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + cx^4}} dx = \frac{Adx\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{5}{4}\right)} + \frac{Aex^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{7}{4}\right)} + \frac{Bdx^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{7}{4}\right)} + \frac{Bex^5\Gamma\left(\frac{5}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{5}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{9}{4}\right)}$$

```
input integrate((B*x**2+A)*(e*x**2+d)/(c*x**4+a)**(1/2),x)
```

```
output A*d*x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(5/4)) + A*e*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4)) + B*d*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4)) + B*e*x**5*gamma(5/4)*hyper((1/2, 5/4), (9/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(9/4))
```

### 3.3.7 Maxima [F]

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)}{\sqrt{cx^4 + a}} dx$$

```
input integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="maxima")
```

```
output integrate((B*x^2 + A)*(e*x^2 + d)/sqrt(c*x^4 + a), x)
```

---

3.3.  $\int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+cx^4}} dx$

**3.3.8 Giac [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)}{\sqrt{cx^4 + a}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)*(e*x^2 + d)/sqrt(c*x^4 + a), x)`

**3.3.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)}{\sqrt{cx^4 + a}} dx$$

input `int(((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(1/2),x)`

output `int(((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(1/2), x)`

### 3.4 $\int \frac{A+Bx^2}{\sqrt{a+cx^4}} dx$

|       |   |    |
|-------|---|----|
| 3.4.1 | Optimal result . . . . .                            | 61 |
| 3.4.2 | Mathematica [C] (verified) . . . . .                | 62 |
| 3.4.3 | Rubi [A] (verified) . . . . .                       | 62 |
| 3.4.4 | Maple [C] (verified) . . . . .                      | 64 |
| 3.4.5 | Fricas [A] (verification not implemented) . . . . . | 64 |
| 3.4.6 | Sympy [C] (verification not implemented) . . . . .  | 65 |
| 3.4.7 | Maxima [F] . . . . .                                | 65 |
| 3.4.8 | Giac [F] . . . . .                                  | 66 |
| 3.4.9 | Mupad [F(-1)] . . . . .                             | 66 |

#### 3.4.1 Optimal result

Integrand size = 19, antiderivative size = 226

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4}} dx$$

$$= \frac{Bx\sqrt{a + cx^4}}{\sqrt{c}(\sqrt{a} + \sqrt{cx^2})} - \frac{\sqrt[4]{a}B(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{c^{3/4}\sqrt{a + cx^4}}$$

$$+ \frac{\sqrt[4]{a}\left(B + \frac{A\sqrt{c}}{\sqrt{a}}\right) (\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2c^{3/4}\sqrt{a + cx^4}}$$

```
output B**x*(c*x^4+a)^(1/2)/c^(1/2)/(a^(1/2)+x^2*c^(1/2))-a^(1/4)*B*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(3/4)/(c*x^4+a)^(1/2)+1/2*a^(1/4)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*(B+A*c^(1/2)/a^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(3/4)/(c*x^4+a)^(1/2)
```

### 3.4.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.03 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.34

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4}} dx$$

$$= \frac{\sqrt{1 + \frac{cx^4}{a}} \left( 3Ax \operatorname{Hypergeometric2F1} \left( \frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -\frac{cx^4}{a} \right) + Bx^3 \operatorname{Hypergeometric2F1} \left( \frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -\frac{cx^4}{a} \right) \right)}{3\sqrt{a + cx^4}}$$

input `Integrate[(A + B*x^2)/Sqrt[a + c*x^4],x]`

output `(Sqrt[1 + (c*x^4)/a]*(3*A*x*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + B*x^3*Hypergeometric2F1[1/2, 3/4, 7/4, -((c*x^4)/a)]))/(3*Sqrt[a + c*x^4])`

### 3.4.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.01, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {1512, 27, 761, 1510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4}} dx$$

$$\downarrow \text{1512}$$

$$\left( \frac{\sqrt{a}B}{\sqrt{c}} + A \right) \int \frac{1}{\sqrt{cx^4 + a}} dx - \frac{\sqrt{a}B \int \frac{\sqrt{a} - \sqrt{cx^2}}{\sqrt{a}\sqrt{cx^4 + a}} dx}{\sqrt{c}}$$

$$\downarrow \text{27}$$

$$\left( \frac{\sqrt{a}B}{\sqrt{c}} + A \right) \int \frac{1}{\sqrt{cx^4 + a}} dx - \frac{B \int \frac{\sqrt{a} - \sqrt{cx^2}}{\sqrt{cx^4 + a}} dx}{\sqrt{c}}$$

$$\downarrow \text{761}$$

$$\frac{(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \left(\frac{\sqrt{a}B}{\sqrt{c}} + A\right) \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2\sqrt[4]{a}\sqrt[4]{c}\sqrt{a+cx^4}} - \frac{B \int \frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{cx^4+a}} dx}{\sqrt{c}}$$

↓ 1510

$$\frac{(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \left(\frac{\sqrt{a}B}{\sqrt{c}} + A\right) \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2\sqrt[4]{a}\sqrt[4]{c}\sqrt{a+cx^4}} - \frac{B \left( \frac{\sqrt[4]{a}(\sqrt{a}+\sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{\sqrt[4]{c}\sqrt{a+cx^4}} - \frac{x\sqrt{a+cx^4}}{\sqrt{a+\sqrt{cx^2}}} \right)}{\sqrt{c}}$$

input `Int[(A + B*x^2)/Sqrt[a + c*x^4], x]`

output `-(B*(-(x*Sqrt[a + c*x^4])/(Sqrt[a] + Sqrt[c]*x^2)) + (a^(1/4)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(c^(1/4)*Sqrt[a + c*x^4]))/Sqrt[c]) + ((A + (Sqrt[a]*B)/Sqrt[c])*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*c^(1/4)*Sqrt[a + c*x^4])`

### 3.4.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 1510 `Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2])]/(q*Sqrt[a + c*x^4])*EllipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]`



```
rule 1512 Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] := With[{q =
  Rt[c/a, 2]}, Simp[(e + d*q)/q Int[1/Sqrt[a + c*x^4], x], x] - Simp[e/q
  Int[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, c
  , d, e}, x] && PosQ[c/a]
```

### 3.4.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.53 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.75

| method   | result   | size |
|----------|--|------|
| default  | $\frac{A\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}},i\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}} + \frac{iB\sqrt{a}\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}\left(F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}},i\right)-E\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}},i\right)\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}\sqrt{c}}$ | 169  |
| elliptic | $\frac{A\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}},i\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}} + \frac{iB\sqrt{a}\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}\left(F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}},i\right)-E\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}},i\right)\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}\sqrt{c}}$ | 169  |

```
input int((B*x^2+A)/(c*x^4+a)^(1/2),x,method=_RETURNVERBOSE)
```

```
output A/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c
^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)
+I*B*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+
I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF(x*(I/a^(1/
2)*c^(1/2))^(1/2),I)-EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I))
```

### 3.4.5 Fracas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.40

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4}} dx$$

$$= \frac{Ba\sqrt{cx}\left(-\frac{a}{c}\right)^{\frac{3}{4}} E\left(\arcsin\left(\frac{\left(-\frac{a}{c}\right)^{\frac{1}{4}}}{x}\right) \mid -1\right) - (Ba - Ac)\sqrt{cx}\left(-\frac{a}{c}\right)^{\frac{3}{4}} F\left(\arcsin\left(\frac{\left(-\frac{a}{c}\right)^{\frac{1}{4}}}{x}\right) \mid -1\right) + \sqrt{cx^4 + a}Ba}{acx}$$

```
input integrate((B*x^2+A)/(c*x^4+a)^(1/2),x, algorithm="fricas")
```

---

3.4.  $\int \frac{A+Bx^2}{\sqrt{a+cx^4}} dx$

output `(B*a*sqrt(c)*x*(-a/c)^(3/4)*elliptic_e(arcsin((-a/c)^(1/4)/x), -1) - (B*a - A*c)*sqrt(c)*x*(-a/c)^(3/4)*elliptic_f(arcsin((-a/c)^(1/4)/x), -1) + sqrt(c*x^4 + a)*B*a)/(a*c*x)`

### 3.4.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.86 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.35

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4}} dx = \frac{Ax\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{5}{4}\right)} + \frac{Bx^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{7}{4}\right)}$$

input `integrate((B*x**2+A)/(c*x**4+a)**(1/2),x)`

output `A*x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(5/4)) + B*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4))`

### 3.4.7 Maxima [F]

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + a}} dx$$

input `integrate((B*x^2+A)/(c*x^4+a)^(1/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/sqrt(c*x^4 + a), x)`

### 3.4.8 Giac [F]

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + a}} dx$$

input `integrate((B*x^2+A)/(c*x^4+a)^(1/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/sqrt(c*x^4 + a), x)`

### 3.4.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + a}} dx$$

input `int((A + B*x^2)/(a + c*x^4)^(1/2),x)`

output `int((A + B*x^2)/(a + c*x^4)^(1/2), x)`

### 3.5 $\int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+cx^4}} dx$

|       |                            |    |
|-------|----------------------------|----|
| 3.5.1 | Optimal result             | 67 |
| 3.5.2 | Mathematica [C] (verified) | 68 |
| 3.5.3 | Rubi [A] (verified)        | 68 |
| 3.5.4 | Maple [C] (verified)       | 70 |
| 3.5.5 | Fricas [F]                 | 71 |
| 3.5.6 | Sympy [F]                  | 71 |
| 3.5.7 | Maxima [F]                 | 71 |
| 3.5.8 | Giac [F]                   | 72 |
| 3.5.9 | Mupad [F(-1)]              | 72 |

#### 3.5.1 Optimal result

Integrand size = 28, antiderivative size = 369

$$\int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+cx^4}} dx = -\frac{(Bd-Ae)\arctan\left(\frac{\sqrt{cd^2+ae^2x}}{\sqrt{d}\sqrt{e}\sqrt{a+cx^4}}\right)}{2\sqrt{d}\sqrt{e}\sqrt{cd^2+ae^2}} - \frac{(\sqrt{a}B-A\sqrt{c})(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right),\frac{1}{2}\right)}{2\sqrt[4]{a}\sqrt[4]{c}(\sqrt{cd}-\sqrt{ae})\sqrt{a+cx^4}} + \frac{a^{3/4}\left(\frac{\sqrt{cd}}{\sqrt{a}}+e\right)^2(Bd-Ae)(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}\text{EllipticPi}\left(-\frac{(\sqrt{cd}-\sqrt{ae})^2}{4\sqrt{a}\sqrt{cde}},2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right),\frac{1}{2}\right)}{4\sqrt[4]{cde}(cd^2-ae^2)\sqrt{a+cx^4}}$$

output

```
-1/2*(-A*e+B*d)*arctan(x*(a*e^2+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+a)^(1/2))/d^(1/2)/e^(1/2)/(a*e^2+c*d^2)^(1/2)-1/2*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(B*a^(1/2)-A*c^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(1/4)/c^(1/4)/(-e*a^(1/2)+d*c^(1/2))/(c*x^4+a)^(1/2)+1/4*a^(3/4)*(-A*e+B*d)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticPi(sin(2*arctan(c^(1/4)*x/a^(1/4))),-1/4*(-e*a^(1/2)+d*c^(1/2))^2/d/e/a^(1/2)/c^(1/2),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*(e*d*c^(1/2)/a^(1/2))^2*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(1/4)/d/e/(-a*e^2+c*d^2)/(c*x^4+a)^(1/2)
```

### 3.5.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.27 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.37

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + cx^4}} dx = \frac{i\sqrt{1 + \frac{cx^4}{a}} \left( Bd \operatorname{EllipticF} \left( \operatorname{arcsinh} \left( \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} x \right), -1 \right) + (-Bd + Ae) \operatorname{EllipticPi} \left( -\frac{i\sqrt{ae}}{\sqrt{cd}}, \operatorname{arcsinh} \left( \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} x \right) \right) \right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} de \sqrt{a + cx^4}}$$

input `Integrate[(A + B*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]),x]`

output `((-I)*Sqrt[1 + (c*x^4)/a]*(B*d*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + (-B*d) + A*e)*EllipticPi[(-I)*Sqrt[a]*e)/(Sqrt[c]*d), I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1)]/(Sqrt[(I*Sqrt[c])/Sqrt[a]]*d*e*Sqrt[a + c*x^4])`

### 3.5.3 Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 382, normalized size of antiderivative = 1.04, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {2227, 27, 761, 2221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + Bx^2}{\sqrt{a + cx^4}(d + ex^2)} dx \\ & \quad \downarrow \text{2227} \\ & \frac{\sqrt{a}(Bd - Ae) \int \frac{\sqrt{cx^2 + \sqrt{a}}}{\sqrt{a}(ex^2 + d)\sqrt{cx^4 + a}} dx}{\sqrt{cd} - \sqrt{ae}} - \frac{(\sqrt{a}B - A\sqrt{c}) \int \frac{1}{\sqrt{cx^4 + a}} dx}{\sqrt{cd} - \sqrt{ae}} \\ & \quad \downarrow \text{27} \\ & \frac{(Bd - Ae) \int \frac{\sqrt{cx^2 + \sqrt{a}}}{(ex^2 + d)\sqrt{cx^4 + a}} dx}{\sqrt{cd} - \sqrt{ae}} - \frac{(\sqrt{a}B - A\sqrt{c}) \int \frac{1}{\sqrt{cx^4 + a}} dx}{\sqrt{cd} - \sqrt{ae}} \\ & \quad \downarrow \text{761} \end{aligned}$$

---

3.5.  $\int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+cx^4}} dx$

$$\frac{(Bd - Ae) \int \frac{\sqrt{cx^2 + \sqrt{a}}}{(ex^2 + d)\sqrt{cx^4 + a}} dx}{\sqrt{cd} - \sqrt{ae}} - \frac{(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} (\sqrt{a}B - A\sqrt{c}) \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2\sqrt[4]{a}\sqrt[4]{c}\sqrt{a+cx^4}(\sqrt{cd} - \sqrt{ae})}$$

↓ 2221

$$(Bd - Ae) \left( \frac{(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} (\sqrt{ae} + \sqrt{cd}) \operatorname{EllipticPi}\left(-\frac{\sqrt{a}\left(\frac{\sqrt{cd}}{\sqrt{a}} - e\right)^2}{4\sqrt{cde}}, 2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{4\sqrt[4]{a}\sqrt[4]{cde}\sqrt{a+cx^4}} - \frac{(\sqrt{cd} - \sqrt{ae}) \arctan\left(\frac{x\sqrt{ae^2 + cd}}{\sqrt{d}\sqrt{e}\sqrt{a+cx^2}}\right)}{2\sqrt{d}\sqrt{e}\sqrt{ae^2 + cd^2}} \right)$$


---


$$\frac{(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} (\sqrt{a}B - A\sqrt{c}) \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2\sqrt[4]{a}\sqrt[4]{c}\sqrt{a+cx^4}(\sqrt{cd} - \sqrt{ae})}$$

input `Int[(A + B*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]),x]`

output `-1/2*((Sqrt[a]*B - A*Sqrt[c])*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/((a^(1/4)*c^(1/4)*(Sqrt[c]*d - Sqrt[a]*e)*Sqrt[a + c*x^4]) + ((B*d - A*e)*(-1/2*((Sqrt[c]*d - Sqrt[a]*e)*ArcTan[(Sqrt[c*d^2 + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + c*x^4])])/(Sqrt[d]*Sqrt[e]*Sqrt[c*d^2 + a*e^2]) + ((Sqrt[c]*d + Sqrt[a]*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-1/4*(Sqrt[a]*((Sqrt[c]*d)/Sqrt[a] - e)^2]/(Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2))/(4*a^(1/4)*c^(1/4)*d*e*Sqrt[a + c*x^4]))/(Sqrt[c]*d - Sqrt[a]*e)`

### 3.5.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

```
rule 2221 Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4])
, x_Symbol] :> With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[c*(d/e)
+ a*(e/d), 2]*(x/Sqrt[a + c*x^4])]/(2*d*e*Rt[c*(d/e) + a*(e/d), 2]))], x]
+ Simp[(B*d + A*e)*(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]/(4*
d*e*q*Sqrt[a + c*x^4]))*EllipticPi[-(e - d*q^2)^2/(4*d*e*q^2), 2*ArcTan[q*x
], 1/2], x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && Po
sQ[c/a] && EqQ[c*A^2 - a*B^2, 0] && PosQ[B/A] && PosQ[c*(d/e) + a*(e/d)]
```

```
rule 2227 Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4])
, x_Symbol] :> With[{q = Rt[c/a, 2]}, Simp[(A*(c*d + a*e*q) - a*B*(e + d*q)
)/(c*d^2 - a*e^2) Int[1/Sqrt[a + c*x^4], x], x] + Simp[a*(B*d - A*e)*((e
+ d*q)/(c*d^2 - a*e^2)) Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]), x]
, x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]
&& NeQ[c*A^2 - a*B^2, 0]
```

### 3.5.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.82 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.52

| method   | result  |
|----------|---|
| default  | $\frac{B\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{e\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}} + \frac{(Ae-Bd)\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}\Pi\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, \frac{i\sqrt{a}e}{\sqrt{cd}}, \sqrt{\frac{-i\sqrt{c}}{\sqrt{a}}}\right)}{ed\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}}$   |
| elliptic | $\frac{B\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{e\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}} + \frac{\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}\Pi\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, \frac{i\sqrt{a}e}{\sqrt{cd}}, \sqrt{\frac{-i\sqrt{c}}{\sqrt{a}}}\right)A}{d\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}} - \frac{\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}\Pi\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, \frac{i\sqrt{a}e}{\sqrt{cd}}, \sqrt{\frac{-i\sqrt{c}}{\sqrt{a}}}\right)}{e\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}}$ |

```
input int((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(1/2), x, method=_RETURNVERBOSE)
```

```
output B/e/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)
*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),
I)+(A*e-B*d)/e/d/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)
*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticPi(x*(I/a^(1/2)*c
^(1/2))^(1/2), I*a^(1/2)/c^(1/2)*e/d, (-I/a^(1/2)*c^(1/2))^(1/2)/(I/a^(1/2)*
c^(1/2))^(1/2))
```

$$3.5. \int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+cx^4}} dx$$

### 3.5.5 Fricas [F]

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + a}(ex^2 + d)} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="fricas")`

output `integral(sqrt(c*x^4 + a)*(B*x^2 + A)/(c*e*x^6 + c*d*x^4 + a*e*x^2 + a*d), x)`

### 3.5.6 Sympy [F]

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + cx^4}} dx = \int \frac{A + Bx^2}{\sqrt{a + cx^4}(d + ex^2)} dx$$

input `integrate((B*x**2+A)/(e*x**2+d)/(c*x**4+a)**(1/2),x)`

output `Integral((A + B*x**2)/(sqrt(a + c*x**4)*(d + e*x**2)), x)`

### 3.5.7 Maxima [F]

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + a}(ex^2 + d)} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(e*x^2 + d)), x)`



### 3.5.8 Giac [F]

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + a}(ex^2 + d)} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(e*x^2 + d)), x)`

### 3.5.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + a}(ex^2 + d)} dx$$

input `int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)),x)`

output `int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)), x)`

## 3.6 $\int \frac{A+Bx^2}{(d+ex^2)^2 \sqrt{a+cx^4}} dx$

|       |                            |    |
|-------|----------------------------|----|
| 3.6.1 | Optimal result             | 73 |
| 3.6.2 | Mathematica [C] (verified) | 74 |
| 3.6.3 | Rubi [A] (verified)        | 75 |
| 3.6.4 | Maple [C] (verified)       | 79 |
| 3.6.5 | Fricas [F(-1)]             | 80 |
| 3.6.6 | Sympy [F]                  | 80 |
| 3.6.7 | Maxima [F]                 | 80 |
| 3.6.8 | Giac [F]                   | 81 |
| 3.6.9 | Mupad [F(-1)]              | 81 |

### 3.6.1 Optimal result

Integrand size = 28, antiderivative size = 641

$$\int \frac{A+Bx^2}{(d+ex^2)^2 \sqrt{a+cx^4}} dx = \frac{\sqrt{c}(Bd-Ae)x\sqrt{a+cx^4}}{2d(cd^2+ae^2)(\sqrt{a}+\sqrt{cx^2})} - \frac{e(Bd-Ae)x\sqrt{a+cx^4}}{2d(cd^2+ae^2)(d+ex^2)}$$

$$- \frac{(Bcd^3-3Acd^2e-aBde^2-aAe^3) \arctan\left(\frac{\sqrt{cd^2+ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{a+cx^4}}\right)}{4d^{3/2}\sqrt{e}(cd^2+ae^2)^{3/2}}$$

$$- \frac{\sqrt[4]{a}\sqrt[4]{c}(Bd-Ae)(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} E\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)}{2d(cd^2+ae^2)\sqrt{a+cx^4}}$$

$$+ \frac{A\sqrt[4]{c}(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2\sqrt[4]{ad}(\sqrt{cd}-\sqrt{ae})\sqrt{a+cx^4}}$$

$$+ \frac{(\sqrt{cd}+\sqrt{ae})(Bcd^3-3Acd^2e-aBde^2-aAe^3)(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \text{EllipticPi}\left(-\frac{(\sqrt{cd}-\sqrt{ae})^2}{4\sqrt{a}\sqrt{cde}}, 2\right)}{8\sqrt[4]{a}\sqrt[4]{cd^2e}(\sqrt{cd}-\sqrt{ae})(cd^2+ae^2)\sqrt{a+cx^4}}$$

output

$$\begin{aligned}
& -1/4*(-A*a*e^3-3*A*c*d^2*e-B*a*d*e^2+B*c*d^3)*\arctan(x*(a*e^2+c*d^2)^{(1/2)} \\
& /d^{(1/2)}/e^{(1/2)}/(c*x^4+a)^{(1/2)})/d^{(3/2)}/(a*e^2+c*d^2)^{(3/2)}/e^{(1/2)}-1/2* \\
& e*(-A*e+B*d)*x*(c*x^4+a)^{(1/2)}/d/(a*e^2+c*d^2)/(e*x^2+d)+1/2*(-A*e+B*d)*x* \\
& c^{(1/2)}*(c*x^4+a)^{(1/2)}/d/(a*e^2+c*d^2)/(a^{(1/2)}+x^2*c^{(1/2)})-1/2*a^{(1/4)}* \\
& c^{(1/4)}*(-A*e+B*d)*(cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/cos(2*\arctan \\
& (c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)} \\
& ))*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/d/(a*e^ \\
& 2+c*d^2)/(c*x^4+a)^{(1/2)}+1/2*A*c^{(1/4)}*(cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2 \\
& )^{(1/2)}/cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/ \\
& a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/ \\
& 2)}))^2)^{(1/2)}/a^{(1/4)}/d/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+a)^{(1/2)}+1/8*(-A*a*e^ \\
& 3-3*A*c*d^2*e-B*a*d*e^2+B*c*d^3)*(cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/ \\
& cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticPi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/ \\
& 4)})), -1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)},1/2*2^{(1/2)})*(e*a^{( \\
& 1/2)}+d*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2 \\
& )^{(1/2)}/a^{(1/4)}/c^{(1/4)}/d^2/e/(a*e^2+c*d^2)/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+a) \\
& ^{(1/2)}
\end{aligned}$$

### 3.6.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.79 (sec) , antiderivative size = 297, normalized size of antiderivative = 0.46

$$\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + cx^4}} dx$$

$$\begin{aligned}
& \frac{de(-Bd+ Ae)x(a+cx^4)}{(cd^2+ae^2)(d+ex^2)} - \frac{i\sqrt{1+\frac{cx^4}{a}}\left(i\sqrt{a}\sqrt{cde}(Bd-Ae)E\left(i\operatorname{arcsinh}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x\right)\middle| -1\right)+\sqrt{cd}(\sqrt{cd}-i\sqrt{ae})(Bd-Ae)\operatorname{EllipticF}\left(i\operatorname{arcsinh}\left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}x\right)\middle| -1\right)\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}(cd^2e+ae^3)} \\
& = \frac{\hspace{15em}}{2d^2\sqrt{a+cx^4}}
\end{aligned}$$

input `Integrate[(A + B*x^2)/((d + e*x^2)^2*Sqrt[a + c*x^4]),x]`

output

$$\begin{aligned}
& ((d*e*(-(B*d) + A*e)*x*(a + c*x^4))/((c*d^2 + a*e^2)*(d + e*x^2)) - (I*\text{Sqrt}[1 + (c*x^4)/a]*(I*\text{Sqrt}[a]*\text{Sqrt}[c]*d*e*(B*d - A*e)*\text{EllipticE}[I*\text{ArcSinh}[\text{Sqrt}[(I*\text{Sqrt}[c])/ \text{Sqrt}[a]]*x], -1] + \text{Sqrt}[c]*d*(\text{Sqrt}[c]*d - I*\text{Sqrt}[a]*e)*(B*d - A*e)*\text{EllipticF}[I*\text{ArcSinh}[\text{Sqrt}[(I*\text{Sqrt}[c])/ \text{Sqrt}[a]]*x], -1] + (-(B*c*d^3) + 3*A*c*d^2*e + a*B*d*e^2 + a*A*e^3)*\text{EllipticPi}[((-I)*\text{Sqrt}[a]*e)/(\text{Sqrt}[c]*d), I*\text{ArcSinh}[\text{Sqrt}[(I*\text{Sqrt}[c])/ \text{Sqrt}[a]]*x], -1)))/(\text{Sqrt}[(I*\text{Sqrt}[c])/ \text{Sqrt}[a]]*(c*d^2*e + a*e^3)))/(2*d^2*\text{Sqrt}[a + c*x^4])
\end{aligned}$$

$$3.6. \int \frac{A+Bx^2}{(d+ex^2)^2\sqrt{a+cx^4}} dx$$

### 3.6.3 Rubi [A] (verified)

Time = 1.29 (sec) , antiderivative size = 600, normalized size of antiderivative = 0.94, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.321$ , Rules used = {2211, 25, 2233, 27, 1510, 2227, 27, 761, 2221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + Bx^2}{\sqrt{a + cx^4} (d + ex^2)^2} dx \\
 & \quad \downarrow \text{2211} \\
 & - \frac{\int -\frac{ce(Bd - Ae)x^4 + 2cd(Bd - Ae)x^2 + 2Acd^2 + aAe^2 + aBde}{(ex^2 + d)\sqrt{cx^4 + a}} dx}{2d(ae^2 + cd^2)} - \frac{ex\sqrt{a + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 + cd^2)} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{ce(Bd - Ae)x^4 + 2cd(Bd - Ae)x^2 + 2Acd^2 + aAe^2 + aBde}{(ex^2 + d)\sqrt{cx^4 + a}} dx}{2d(ae^2 + cd^2)} - \frac{ex\sqrt{a + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 + cd^2)} \\
 & \quad \downarrow \text{2233} \\
 & \frac{\int \frac{ce(2Acd^2 + \sqrt{a}\sqrt{c}(Bd - Ae)d + \sqrt{c}(\sqrt{cd} + \sqrt{ae})(Bd - Ae)x^2 + ae(Bd + Ae))}{(ex^2 + d)\sqrt{cx^4 + a}} dx}{ce} - \frac{\sqrt{a}\sqrt{c}(Bd - Ae) \int \frac{\sqrt{a} - \sqrt{cx^2}}{\sqrt{a}\sqrt{cx^4 + a}} dx}{2d(ae^2 + cd^2)} - \frac{ex\sqrt{a + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 + cd^2)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{2Acd^2 + \sqrt{a}\sqrt{c}(Bd - Ae)d + \sqrt{c}(\sqrt{cd} + \sqrt{ae})(Bd - Ae)x^2 + ae(Bd + Ae)}{(ex^2 + d)\sqrt{cx^4 + a}} dx}{2d(ae^2 + cd^2)} - \frac{\sqrt{c}(Bd - Ae) \int \frac{\sqrt{a} - \sqrt{cx^2}}{\sqrt{cx^4 + a}} dx}{2d(d + ex^2)(ae^2 + cd^2)} \\
 & \quad \downarrow \text{1510} \\
 & \frac{\int \frac{2Acd^2 + \sqrt{a}\sqrt{c}(Bd - Ae)d + \sqrt{c}(\sqrt{cd} + \sqrt{ae})(Bd - Ae)x^2 + ae(Bd + Ae)}{(ex^2 + d)\sqrt{cx^4 + a}} dx}{2d(ae^2 + cd^2)} - \frac{\sqrt{c}(Bd - Ae) \left( \frac{\sqrt[4]{a}(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a + cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt{a} - \sqrt{cx^2}}{\sqrt{a}\sqrt{cx^4 + a}}\right)\right)}{\sqrt[4]{c}\sqrt{a + cx^4}} \right)}{2d(d + ex^2)(ae^2 + cd^2)} \\
 & \quad \frac{ex\sqrt{a + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 + cd^2)}
 \end{aligned}$$

---

3.6.  $\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + cx^4}} dx$

↓ 2227

$$\frac{\sqrt{a}(-aAe^3 - aBde^2 - 3Acd^2e + Bcd^3) \int \frac{\sqrt{cx^2 + \sqrt{a}}}{\sqrt{a}(ex^2 + d)\sqrt{cx^4 + a}} dx + \frac{2A\sqrt{c}(ae^2 + cd^2) \int \frac{1}{\sqrt{cx^4 + a}} dx}{\sqrt{cd - \sqrt{ae}}} - \sqrt{c}(Bd - Ae) \left( \frac{\sqrt[4]{a}(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}}}{\sqrt[4]{c\sqrt{a}}} \right)}{\sqrt{cd - \sqrt{ae}}}$$


---


$$\frac{ex\sqrt{a + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 + cd^2)}$$

↓ 27

$$\frac{(-aAe^3 - aBde^2 - 3Acd^2e + Bcd^3) \int \frac{\sqrt{cx^2 + \sqrt{a}}}{(ex^2 + d)\sqrt{cx^4 + a}} dx + \frac{2A\sqrt{c}(ae^2 + cd^2) \int \frac{1}{\sqrt{cx^4 + a}} dx}{\sqrt{cd - \sqrt{ae}}} - \sqrt{c}(Bd - Ae) \left( \frac{\sqrt[4]{a}(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}}}{\sqrt[4]{c\sqrt{a}}} \right)}{\sqrt{cd - \sqrt{ae}}}$$


---


$$\frac{ex\sqrt{a + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 + cd^2)}$$

↓ 761

$$\frac{(-aAe^3 - aBde^2 - 3Acd^2e + Bcd^3) \int \frac{\sqrt{cx^2 + \sqrt{a}}}{(ex^2 + d)\sqrt{cx^4 + a}} dx - \sqrt{c}(Bd - Ae) \left( \frac{\sqrt[4]{a}(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right) - \frac{x}{\sqrt{d}}}{\sqrt[4]{c\sqrt{a+cx^4}}} \right)}{\sqrt{cd - \sqrt{ae}}}$$


---


$$\frac{ex\sqrt{a + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 + cd^2)}$$

↓ 2221

$$\frac{(-aAe^3 - aBde^2 - 3Acd^2e + Bcd^3) \left( \frac{(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} (\sqrt{ae} + \sqrt{cd}) \text{EllipticPi}\left(-\frac{\sqrt{a}\left(\frac{\sqrt{cd}}{\sqrt{a}} - e\right)^2}{4\sqrt{cde}}, 2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{4\sqrt[4]{a}\sqrt[4]{Cde}\sqrt{a+cx^4}} - \frac{(\sqrt{cd} - \sqrt{ae}) \arctan\left(\frac{x}{\sqrt{d}}\right)}{2\sqrt{d}\sqrt{e}\sqrt{ae^2 + cd^2}} \right)}{\sqrt{cd - \sqrt{ae}}}$$


---


$$\frac{ex\sqrt{a + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 + cd^2)}$$

input `Int[(A + B*x^2)/((d + e*x^2)^2*sqrt[a + c*x^4]),x]`

3.6.  $\int \frac{A+Bx^2}{(d+ex^2)^2\sqrt{a+cx^4}} dx$

```
output -1/2*(e*(B*d - A*e)*x*Sqrt[a + c*x^4])/(d*(c*d^2 + a*e^2)*(d + e*x^2)) + (
-(Sqrt[c]*(B*d - A*e)*(-(x*Sqrt[a + c*x^4])/(Sqrt[a] + Sqrt[c]*x^2)) + (a
^(1/4)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]
*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(c^(1/4)*Sqrt[a + c*x^4]))
) + (A*c^(1/4)*(c*d^2 + a*e^2)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(S
qrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(a
^(1/4)*(Sqrt[c]*d - Sqrt[a]*e)*Sqrt[a + c*x^4]) + ((B*c*d^3 - 3*A*c*d^2*e
- a*B*d*e^2 - a*A*e^3)*(-1/2*((Sqrt[c]*d - Sqrt[a]*e)*ArcTan[(Sqrt[c*d^2 +
a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + c*x^4])])/(Sqrt[d]*Sqrt[e]*Sqrt[c*d^2
+ a*e^2]) + ((Sqrt[c]*d + Sqrt[a]*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*
x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-1/4*(Sqrt[a]*((Sqrt[c]*d)/Sqrt
[a] - e)^2)/(Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(4*a^(1/4)
*c^(1/4)*d*e*Sqrt[a + c*x^4]))/(Sqrt[c]*d - Sqrt[a]*e))/(2*d*(c*d^2 + a*e
^2))
```

### 3.6.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 761 Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(
1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^4]))*
EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

```
rule 1510 Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*
(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2])/(q*Sqrt[a + c*x^4]))*E
llipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e
}, x] && PosQ[c/a]
```

rule 2211 `Int[((P4x_)*((d_) + (e_)*(x_)^2)^(q_))/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol]
:> With[{A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]
}, Simp[(-(C*d^2 - B*d*e + A*e^2))*x*(d + e*x^2)^(q + 1)*(Sqrt[a + c*x^4]/(
2*d*(q + 1)*(c*d^2 + a*e^2))), x] + Simp[1/(2*d*(q + 1)*(c*d^2 + a*e^2))
Int[((d + e*x^2)^(q + 1)/Sqrt[a + c*x^4])*Simp[a*d*(C*d - B*e) + A*(a*e^2*(
2*q + 3) + 2*c*d^2*(q + 1)) + 2*d*(B*c*d - A*c*e + a*C*e)*(q + 1)*x^2 + c*(
C*d^2 - B*d*e + A*e^2)*(2*q + 5)*x^4, x], x], x]] /; FreeQ[{a, c, d, e}, x]
&& PolyQ[P4x, x^2] && LeQ[Expon[P4x, x], 4] && ILtQ[q, -1]`

rule 2221 `Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4])
, x_Symbol] :> With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[c*(d/e)
+ a*(e/d), 2]*(x/Sqrt[a + c*x^4])]/(2*d*e*Rt[c*(d/e) + a*(e/d), 2])), x]
+ Simp[(B*d + A*e)*(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]/(4*
d*e*q*Sqrt[a + c*x^4]))*EllipticPi[-(e - d*q^2)^2/(4*d*e*q^2), 2*ArcTan[q*x
], 1/2], x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && Po
sQ[c/a] && EqQ[c*A^2 - a*B^2, 0] && PosQ[B/A] && PosQ[c*(d/e) + a*(e/d)]`

rule 2227 `Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4])
, x_Symbol] :> With[{q = Rt[c/a, 2]}, Simp[(A*(c*d + a*e*q) - a*B*(e + d*q)
)/(c*d^2 - a*e^2) Int[1/Sqrt[a + c*x^4], x], x] + Simp[a*(B*d - A*e)*((e
+ d*q)/(c*d^2 - a*e^2)) Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]), x]
, x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]
&& NeQ[c*A^2 - a*B^2, 0]`

rule 2233 `Int[(P4x_)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4]), x_Symbol] :>
With[{q = Rt[c/a, 2], A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff
[P4x, x, 4]}, Simp[-C/(e*q) Int[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] + Sim
p[1/(c*e) Int[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a*e*q))*x^2)/((d + e*x
^2)*Sqrt[a + c*x^4]), x], x]] /; FreeQ[{a, c, d, e}, x] && PolyQ[P4x, x^2,
2] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]`

### 3.6.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.42 (sec) , antiderivative size = 679, normalized size of antiderivative = 1.06

| method   | result  |
|----------|---|
| default  | $\frac{B\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}\Pi\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}},\frac{i\sqrt{a}e}{\sqrt{c}d},\sqrt{\frac{-i\sqrt{c}}{\sqrt{a}}}\right)}{ed\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}} + (Ae-Bd)\left(\frac{e^2x\sqrt{cx^4+a}}{2d(ae^2+cd^2)(ex^2+d)} - \frac{c\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}},i\right)}{2(ae^2+cd^2)\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}}\right)$ |
| elliptic | Expression too large to display   |

```
input int((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(1/2),x,method=_RETURNVERBOSE)
```

```
output B/e/d/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticPi(x*(I/a^(1/2)*c^(1/2))^(1/2),I*a^(1/2)/c^(1/2)*e/d,(-I/a^(1/2)*c^(1/2))^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2))+ (A*e-B*d)/e*(1/2*e^2*x*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)/(e*x^2+d)-1/2*c/(a*e^2+c*d^2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)-1/2*I*c^(1/2)*e/d/(a*e^2+c*d^2)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+1/2*I*c^(1/2)*e/d/(a*e^2+c*d^2)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+1/2/d^2/(a*e^2+c*d^2)*e^2/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticPi(x*(I/a^(1/2)*c^(1/2))^(1/2),I*a^(1/2)/c^(1/2)*e/d,(-I/a^(1/2)*c^(1/2))^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2))*a+3/2/(a*e^2+c*d^2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticPi(x*(I/a^(1/2)*c^(1/2))^(1/2),I*a^(1/2)/c^(1/2)*e/d,(-I/a^(1/2)*c^(1/2))^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2))*c
```

3.6.  $\int \frac{A+Bx^2}{(d+ex^2)^2\sqrt{a+cx^4}} dx$



### 3.6.5 Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + cx^4}} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(1/2),x, algorithm="fricas")`

output `Timed out`

### 3.6.6 Sympy [F]

$$\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + cx^4}} dx = \int \frac{A + Bx^2}{\sqrt{a + cx^4} (d + ex^2)^2} dx$$

input `integrate((B*x**2+A)/(e*x**2+d)**2/(c*x**4+a)**(1/2),x)`

output `Integral((A + B*x**2)/(sqrt(a + c*x**4)*(d + e*x**2)**2), x)`

### 3.6.7 Maxima [F]

$$\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + a}(ex^2 + d)^2} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(1/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(e*x^2 + d)^2), x)`

**3.6.8 Giac [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + a}(ex^2 + d)^2} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(1/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(e*x^2 + d)^2), x)`

**3.6.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + a}(ex^2 + d)^2} dx$$

input `int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)^2),x)`

output `int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)^2), x)`

### 3.7 $\int \frac{A+Bx^2}{(d+ex^2)^3 \sqrt{a+cx^4}} dx$

|       |                            |    |
|-------|----------------------------|----|
| 3.7.1 | Optimal result             | 82 |
| 3.7.2 | Mathematica [C] (verified) | 83 |
| 3.7.3 | Rubi [A] (verified)        | 84 |
| 3.7.4 | Maple [C] (verified)       | 89 |
| 3.7.5 | Fricas [F(-1)]             | 90 |
| 3.7.6 | Sympy [F]                  | 90 |
| 3.7.7 | Maxima [F]                 | 90 |
| 3.7.8 | Giac [F]                   | 91 |
| 3.7.9 | Mupad [F(-1)]              | 91 |

#### 3.7.1 Optimal result

Integrand size = 28, antiderivative size = 875

$$\int \frac{A+Bx^2}{(d+ex^2)^3 \sqrt{a+cx^4}} dx = \frac{\sqrt{c}(5Bcd^3 - 9Acd^2e - aBde^2 - 3aAe^3)x\sqrt{a+cx^4}}{8d^2(cd^2+ae^2)^2(\sqrt{a}+\sqrt{cx^2})} - \frac{e(Bd-Ae)x\sqrt{a+cx^4}}{4d(cd^2+ae^2)(d+ex^2)^2} - \frac{e(5Bcd^3 - 9Acd^2e - aBde^2 - 3aAe^3)x\sqrt{a+cx^4}}{8d^2(cd^2+ae^2)^2(d+ex^2)} + \frac{(3Ae(5c^2d^4 + 2acd^2e^2 + a^2e^4) - B(3c^2d^5 - 10acd^3e^2 - a^2de^4)) \arctan\left(\frac{\sqrt{cd^2+ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{a+cx^4}}\right)}{16d^{5/2}\sqrt{e}(cd^2+ae^2)^{5/2}} - \frac{\sqrt[4]{a}\sqrt[4]{c}(5Bcd^3 - 9Acd^2e - aBde^2 - 3aAe^3)(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)}{8d^2(cd^2+ae^2)^2\sqrt{a+cx^4}} + \frac{\sqrt[4]{c}(4Acd^2 + \sqrt{a}\sqrt{cd}(Bd-Ae) + ae(Bd+3Ae))(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right)\right)}{8\sqrt[4]{ad^2}(\sqrt{cd}-\sqrt{ae})(cd^2+ae^2)\sqrt{a+cx^4}} - \frac{(\sqrt{cd}+\sqrt{ae})(3Ae(5c^2d^4 + 2acd^2e^2 + a^2e^4) - B(3c^2d^5 - 10acd^3e^2 - a^2de^4))(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}}{32\sqrt[4]{a}\sqrt[4]{cd^3e}(\sqrt{cd}-\sqrt{ae})(cd^2+ae^2)^2\sqrt{a+cx^4}}$$

```
output 1/16*(3*A*e*(a^2*e^4+2*a*c*d^2*e^2+5*c^2*d^4)-B*(-a^2*d*e^4-10*a*c*d^3*e^2
+3*c^2*d^5))*arctan(x*(a*e^2+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+a)^(1/2))
/d^(5/2)/(a*e^2+c*d^2)^(5/2)/e^(1/2)-1/4*e*(-A*e+B*d)*x*(c*x^4+a)^(1/2)/d/
(a*e^2+c*d^2)/(e*x^2+d)^2-1/8*e*(-3*A*a*e^3-9*A*c*d^2*e-B*a*d*e^2+5*B*c*d^
3)*x*(c*x^4+a)^(1/2)/d^2/(a*e^2+c*d^2)^2/(e*x^2+d)+1/8*(-3*A*a*e^3-9*A*c*d
^2*e-B*a*d*e^2+5*B*c*d^3)*x*c^(1/2)*(c*x^4+a)^(1/2)/d^2/(a*e^2+c*d^2)^2/(a
^(1/2)+x^2*c^(1/2))-1/8*a^(1/4)*c^(1/4)*(-3*A*a*e^3-9*A*c*d^2*e-B*a*d*e^2+
5*B*c*d^3)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)
*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1
/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/d^2/(a*e^2+c*d^
2)^2/(c*x^4+a)^(1/2)-1/32*(3*A*e*(a^2*e^4+2*a*c*d^2*e^2+5*c^2*d^4)-B*(-a^2
*d*e^4-10*a*c*d^3*e^2+3*c^2*d^5))*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/
2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticPi(sin(2*arctan(c^(1/4)*x/a^(1
/4))),-1/4*(-e*a^(1/2)+d*c^(1/2))^2/d/e/a^(1/2)/c^(1/2),1/2*2^(1/2))*(e*a^
(1/2)+d*c^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2
^(1/2)/a^(1/4)/c^(1/4)/d^3/e/(a*e^2+c*d^2)^2/(-e*a^(1/2)+d*c^(1/2))/(c*x^4
+a)^(1/2)+1/8*c^(1/4)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arc
tan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(
1/2))*(a^(1/2)+x^2*c^(1/2))*(4*A*c*d^2+a*e*(3*A*e+B*d)+d*(-A*e+B*d)*a^(1/2
))*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(1/4)/d^2/(a*e^2...
```

### 3.7.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 11.33 (sec) , antiderivative size = 453, normalized size of antiderivative = 0.52

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + cx^4}} dx$$

$$= \frac{de^2x(a+cx^4)(2d(Bd-Ae)(cd^2+ae^2)+(5Bcd^3-9Acd^2e-aBde^2-3aAe^3)(d+ex^2))}{(d+ex^2)^2} - \frac{i\sqrt{1+\frac{cx^4}{a}}(-i\sqrt{a}\sqrt{cde}(-5Bcd^3+9Acd^2e+aBde^2+3aAe^3))}{(d+ex^2)^2}$$

```
input Integrate[(A + B*x^2)/((d + e*x^2)^3*Sqrt[a + c*x^4]),x]
```

output  $(-((d*e^2*x*(a + c*x^4)*(2*d*(B*d - A*e)*(c*d^2 + a*e^2) + (5*B*c*d^3 - 9*A*c*d^2*e - a*B*d*e^2 - 3*a*A*e^3)*(d + e*x^2)))/(d + e*x^2)^2) - (I*\text{Sqrt}[1 + (c*x^4)/a]*((-I)*\text{Sqrt}[a]*\text{Sqrt}[c]*d*e*(-5*B*c*d^3 + 9*A*c*d^2*e + a*B*d*e^2 + 3*a*A*e^3)*\text{EllipticE}[I*\text{ArcSinh}[\text{Sqrt}[(I*\text{Sqrt}[c])/ \text{Sqrt}[a]]*x], -1] + \text{Sqrt}[c]*d*(\text{Sqrt}[c]*d - I*\text{Sqrt}[a]*e)*(A*e*(-7*c*d^2 + (2*I)*\text{Sqrt}[a]*\text{Sqrt}[c]*d*e - 3*a*e^2) + B*d*(3*c*d^2 - (2*I)*\text{Sqrt}[a]*\text{Sqrt}[c]*d*e - a*e^2))*\text{EllipticF}[I*\text{ArcSinh}[\text{Sqrt}[(I*\text{Sqrt}[c])/ \text{Sqrt}[a]]*x], -1] + (3*A*e*(5*c^2*d^4 + 2*a*c*d^2*e^2 + a^2*e^4) + B*(-3*c^2*d^5 + 10*a*c*d^3*e^2 + a^2*d*e^4))*\text{EllipticPi}[((-I)*\text{Sqrt}[a]*e)/(\text{Sqrt}[c]*d), I*\text{ArcSinh}[\text{Sqrt}[(I*\text{Sqrt}[c])/ \text{Sqrt}[a]]*x], -1]))/\text{Sqrt}[(I*\text{Sqrt}[c])/ \text{Sqrt}[a]]/(8*d^3*e*(c*d^2 + a*e^2)^2*\text{Sqrt}[a + c*x^4])$

### 3.7.3 Rubi [A] (verified)

Time = 2.19 (sec) , antiderivative size = 788, normalized size of antiderivative = 0.90, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.393$ , Rules used = {2211, 25, 2211, 25, 2233, 27, 1510, 2227, 27, 761, 2221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4} (d + ex^2)^3} dx$$

↓ 2211

$$-\frac{\int -\frac{ce(Bd - Ae)x^4 + 4cd(Bd - Ae)x^2 + 4Acd^2 + 3aAe^2 + aBde}{(ex^2 + d)^2 \sqrt{cx^4 + a}} dx}{4d(ae^2 + cd^2)} - \frac{ex\sqrt{a + cx^4}(Bd - Ae)}{4d(d + ex^2)^2(ae^2 + cd^2)}$$

↓ 25

$$\frac{\int -\frac{ce(Bd - Ae)x^4 + 4cd(Bd - Ae)x^2 + 4Acd^2 + 3aAe^2 + aBde}{(ex^2 + d)^2 \sqrt{cx^4 + a}} dx}{4d(ae^2 + cd^2)} - \frac{ex\sqrt{a + cx^4}(Bd - Ae)}{4d(d + ex^2)^2(ae^2 + cd^2)}$$

↓ 2211

$$-\frac{\int -\frac{ce(5Bcd^3 - 9Aced^2 - aBe^2d - 3aAe^3)x^4 + 4cd(2Bcd^3 - 4Aced^2 - aBe^2d - aAe^3)x^2 + aBde(7cd^2 + ae^2) + A(8e^2d^4 + 5ace^2d^2 + 3a^2e^4)}{(ex^2 + d)\sqrt{cx^4 + a}} dx}{2d(ae^2 + cd^2)} - \frac{ex\sqrt{a + cx^4}(-3a^2e^4 + 2a^2e^2d + 2a^2d^2)}{2a^2d(ae^2 + cd^2)}$$

↓ 25

$$\frac{ex\sqrt{a + cx^4}(Bd - Ae)}{4d(d + ex^2)^2(ae^2 + cd^2)}$$

---

3.7.  $\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + cx^4}} dx$

$$\int \frac{ce(5Bcd^3 - 9Aced^2 - aBe^2d - 3aAe^3)x^4 + 4cd(2Bcd^3 - 4Aced^2 - aBe^2d - aAe^3)x^2 + aBde(7cd^2 + ae^2) + A(8c^2d^4 + 5ace^2d^2 + 3a^2e^4)}{(ex^2 + d)\sqrt{cx^4 + a}} dx - \frac{ex\sqrt{a + cx^4}(-3aAe)}{2d(d + ex^2)}$$

$$\frac{4d(ae^2 + cd^2)ex\sqrt{a + cx^4}(Bd - Ae)}{4d(d + ex^2)^2(ae^2 + cd^2)}$$

↓ 2233

$$\int \frac{ce(8Ac^2d^4 + \sqrt{a}c^{3/2}(5Bd - 9Ae)d^3 + ace(7Bd + 5Ae)d^2 - a^{3/2}\sqrt{ce^2}(Bd + 3Ae)d - ((cd - \sqrt{a}\sqrt{ce})(5Bcd^3 - 9Aced^2 - aBe^2d - 3aAe^3) - 4cd(2Bcd^3 - 4Aced^2 - aBe^2d - aAe^3))}{(ex^2 + d)\sqrt{cx^4 + a}} dx - \frac{ex\sqrt{a + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 + cd^2)}$$

$$\frac{ex\sqrt{a + cx^4}(Bd - Ae)}{4d(d + ex^2)^2(ae^2 + cd^2)}$$

↓ 27

$$\int \frac{8Ac^2d^4 + \sqrt{a}c^{3/2}(5Bd - 9Ae)d^3 + ace(7Bd + 5Ae)d^2 - a^{3/2}\sqrt{ce^2}(Bd + 3Ae)d - ((cd - \sqrt{a}\sqrt{ce})(5Bcd^3 - 9Aced^2 - aBe^2d - 3aAe^3) - 4cd(2Bcd^3 - 4Aced^2 - aBe^2d - aAe^3))}{(ex^2 + d)\sqrt{cx^4 + a}} dx - \frac{ex\sqrt{a + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 + cd^2)}$$

$$\frac{ex\sqrt{a + cx^4}(Bd - Ae)}{4d(d + ex^2)^2(ae^2 + cd^2)}$$

↓ 1510

$$\int \frac{8Ac^2d^4 + \sqrt{a}c^{3/2}(5Bd - 9Ae)d^3 + ace(7Bd + 5Ae)d^2 - a^{3/2}\sqrt{ce^2}(Bd + 3Ae)d - ((cd - \sqrt{a}\sqrt{ce})(5Bcd^3 - 9Aced^2 - aBe^2d - 3aAe^3) - 4cd(2Bcd^3 - 4Aced^2 - aBe^2d - aAe^3))}{(ex^2 + d)\sqrt{cx^4 + a}} dx - \frac{ex\sqrt{a + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 + cd^2)}$$

$$\frac{ex\sqrt{a + cx^4}(Bd - Ae)}{4d(d + ex^2)^2(ae^2 + cd^2)}$$

↓ 2227

$$\frac{\sqrt{a}(3Ae(a^2e^4 + 2acd^2e^2 + 5c^2d^4) - B(-a^2de^4 - 10acd^3e^2 + 3c^2d^5))}{\sqrt{cd - \sqrt{ae}}} \int \frac{\sqrt{cx^2 + \sqrt{a}}}{\sqrt{a}(ex^2 + d)\sqrt{cx^4 + a}} dx + \frac{2\sqrt{c}(ae^2 + cd^2)(\sqrt{a}\sqrt{cd}(Bd - Ae) + ae(3Ae + Bd) + 4Acd^2)}{\sqrt{cd - \sqrt{ae}}} \int \frac{1}{\sqrt{cx^4 + a}} dx - \frac{ex\sqrt{a + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 + cd^2)}$$

$$\frac{ex\sqrt{a + cx^4}(Bd - Ae)}{4d(d + ex^2)^2(ae^2 + cd^2)}$$

3.7.  $\int \frac{A+Bx^2}{(d+ex^2)^3\sqrt{a+cx^4}} dx$

↓ 27

$$-\frac{(3Ae(a^2e^4+2acd^2e^2+5c^2d^4)-B(-a^2de^4-10acd^3e^2+3c^2d^5)) \int \frac{\sqrt{cx^2+\sqrt{a}}}{(ex^2+d)\sqrt{cx^4+a}} dx}{\sqrt{cd}-\sqrt{ae}} + \frac{2\sqrt{c}(ae^2+cd^2)(\sqrt{a}\sqrt{cd}(Bd-Ae)+ae(3Ae+Bd)+4Acd^2) \int \frac{1}{\sqrt{cx^4+a}} dx}{\sqrt{cd}-\sqrt{ae}}$$


---


$$2d(ae^2+cd^2)$$

$$\frac{ex\sqrt{a+cx^4}(Bd-Ae)}{4d(d+ex^2)^2(ae^2+cd^2)}$$

↓ 761

$$-\frac{(3Ae(a^2e^4+2acd^2e^2+5c^2d^4)-B(-a^2de^4-10acd^3e^2+3c^2d^5)) \int \frac{\sqrt{cx^2+\sqrt{a}}}{(ex^2+d)\sqrt{cx^4+a}} dx}{\sqrt{cd}-\sqrt{ae}} + \frac{\sqrt[4]{c}(\sqrt{a}+\sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} (ae^2+cd^2) \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right), \frac{1}{2}\right)}{\sqrt[4]{a}\sqrt{a+cx^4}(\sqrt{cd}-\sqrt{ae})}$$


---

$$\frac{ex\sqrt{a+cx^4}(Bd-Ae)}{4d(d+ex^2)^2(ae^2+cd^2)}$$

↓ 2221

$$-\frac{(3Ae(a^2e^4+2acd^2e^2+5c^2d^4)-B(-a^2de^4-10acd^3e^2+3c^2d^5)) \left( \frac{(\sqrt{a}+\sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} (\sqrt{ae}+\sqrt{cd}) \text{EllipticPi}\left(-\frac{\sqrt{a}\left(\frac{\sqrt{cd}}{\sqrt{a}}-e\right)^2}{4\sqrt{cde}}, 2 \arctan\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right), \frac{1}{2}\right)}{4\sqrt[4]{a}\sqrt[4]{cde}\sqrt{a+cx^4}} \right)}{\sqrt{cd}-\sqrt{ae}}$$


---

$$\frac{ex\sqrt{a+cx^4}(Bd-Ae)}{4d(d+ex^2)^2(ae^2+cd^2)}$$

input `Int[(A + B*x^2)/((d + e*x^2)^3*Sqrt[a + c*x^4]),x]`

```

output -1/4*(e*(B*d - A*e)*x*Sqrt[a + c*x^4])/(d*(c*d^2 + a*e^2)*(d + e*x^2)^2) +
(-1/2*(e*(5*B*c*d^3 - 9*A*c*d^2*e - a*B*d*e^2 - 3*a*A*e^3)*x*Sqrt[a + c*x
^4])/(d*(c*d^2 + a*e^2)*(d + e*x^2)) + (-Sqrt[c]*(5*B*c*d^3 - 9*A*c*d^2*e
- a*B*d*e^2 - 3*a*A*e^3)*(-(x*Sqrt[a + c*x^4])/(Sqrt[a + Sqrt[c]*x^2))
+ (a^(1/4)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2
)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(c^(1/4)*Sqrt[a + c*x^
4]))) + (c^(1/4)*(c*d^2 + a*e^2)*(4*A*c*d^2 + Sqrt[a]*Sqrt[c]*d*(B*d - A*e
) + a*e*(B*d + 3*A*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] +
Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(a^(1/4)*(
Sqrt[c]*d - Sqrt[a]*e)*Sqrt[a + c*x^4]) - ((3*A*e*(5*c^2*d^4 + 2*a*c*d^2*e
^2 + a^2*e^4) - B*(3*c^2*d^5 - 10*a*c*d^3*e^2 - a^2*d*e^4))*(-1/2*((Sqrt[c
]*d - Sqrt[a]*e)*ArcTan[(Sqrt[c*d^2 + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a +
c*x^4])])/(Sqrt[d]*Sqrt[e]*Sqrt[c*d^2 + a*e^2]) + ((Sqrt[c]*d + Sqrt[a]*e
)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*Ellip
ticPi[-1/4*(Sqrt[a]*((Sqrt[c]*d)/Sqrt[a] - e)^2)/(Sqrt[c]*d*e), 2*ArcTan[(
c^(1/4)*x)/a^(1/4)], 1/2])/(4*a^(1/4)*c^(1/4)*d*e*Sqrt[a + c*x^4]))/(Sqrt
[c]*d - Sqrt[a]*e))/(2*d*(c*d^2 + a*e^2))/(4*d*(c*d^2 + a*e^2))

```

### 3.7.3.1 Defintions of rubi rules used

```

rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]

```

```

rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]

```

```

rule 761 Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(
1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^4]))*
EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

```

```

rule 1510 Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*
(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2])/(q*Sqrt[a + c*x^4]))*E
llipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e
}, x] && PosQ[c/a]

```



rule 2211 `Int[((P4x_)*((d_) + (e_)*(x_)^2)^(q_))/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol]
:> With[{A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]
}, Simp[(-(C*d^2 - B*d*e + A*e^2))*x*(d + e*x^2)^(q + 1)*(Sqrt[a + c*x^4]/(
2*d*(q + 1)*(c*d^2 + a*e^2))), x] + Simp[1/(2*d*(q + 1)*(c*d^2 + a*e^2))
Int[((d + e*x^2)^(q + 1)/Sqrt[a + c*x^4])*Simp[a*d*(C*d - B*e) + A*(a*e^2*(
2*q + 3) + 2*c*d^2*(q + 1)) + 2*d*(B*c*d - A*c*e + a*C*e)*(q + 1)*x^2 + c*(
C*d^2 - B*d*e + A*e^2)*(2*q + 5)*x^4, x], x], x]] /; FreeQ[{a, c, d, e}, x]
&& PolyQ[P4x, x^2] && LeQ[Expon[P4x, x], 4] && ILtQ[q, -1]`

rule 2221 `Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4])
, x_Symbol] :> With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[c*(d/e)
+ a*(e/d), 2]*(x/Sqrt[a + c*x^4])]/(2*d*e*Rt[c*(d/e) + a*(e/d), 2])), x]
+ Simp[(B*d + A*e)*(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]/(4*
d*e*q*Sqrt[a + c*x^4]))*EllipticPi[-(e - d*q^2)^2/(4*d*e*q^2), 2*ArcTan[q*x
], 1/2], x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && Po
sQ[c/a] && EqQ[c*A^2 - a*B^2, 0] && PosQ[B/A] && PosQ[c*(d/e) + a*(e/d)]`

rule 2227 `Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4])
, x_Symbol] :> With[{q = Rt[c/a, 2]}, Simp[(A*(c*d + a*e*q) - a*B*(e + d*q)
)/(c*d^2 - a*e^2) Int[1/Sqrt[a + c*x^4], x], x] + Simp[a*(B*d - A*e)*((e
+ d*q)/(c*d^2 - a*e^2)) Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]), x]
, x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]
&& NeQ[c*A^2 - a*B^2, 0]`

rule 2233 `Int[(P4x_)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4]), x_Symbol] :>
With[{q = Rt[c/a, 2], A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff
[P4x, x, 4]}, Simp[-C/(e*q) Int[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] + Sim
p[1/(c*e) Int[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a*e*q))*x^2)/((d + e*x
^2)*Sqrt[a + c*x^4]), x], x]] /; FreeQ[{a, c, d, e}, x] && PolyQ[P4x, x^2,
2] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]`

### 3.7.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 3.01 (sec) , antiderivative size = 1591, normalized size of antiderivative = 1.82

| method   | result                          | size |
|----------|---------------------------------|------|
| default  | Expression too large to display | 1591 |
| elliptic | Expression too large to display | 1982 |

```
input int((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(1/2),x,method=_RETURNVERBOSE)
```

```
output B/e*(1/2*e^2*x*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)/(e*x^2+d)-1/2*c/(a*e^2+c*d^2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)-1/2*I*c^(1/2)*e/d/(a*e^2+c*d^2)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+1/2*I*c^(1/2)*e/d/(a*e^2+c*d^2)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+1/2/d^2/(a*e^2+c*d^2)*e^2/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticPi(x*(I/a^(1/2)*c^(1/2))^(1/2),I*a^(1/2)/c^(1/2)*e/d,(-I/a^(1/2)*c^(1/2))^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2))*a+3/2/(a*e^2+c*d^2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticPi(x*(I/a^(1/2)*c^(1/2))^(1/2),I*a^(1/2)/c^(1/2)*e/d,(-I/a^(1/2)*c^(1/2))^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2))*c)+(A*e-B*d)/e*(1/4*e^2*x*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)/(e*x^2+d)^2+3/8*e^2*(a*e^2+3*c*d^2)*x*(c*x^4+a)^(1/2)/d^2/(a*e^2+c*d^2)^2/(e*x^2+d)-1/8*c/d/(a*e^2+c*d^2)^2/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)*a*e^2-7/8*c^2*d/(a*e^2+c*d^2)^2/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2))...
```

### 3.7.5 Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + cx^4}} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(1/2),x, algorithm="fricas")`

output `Timed out`

### 3.7.6 Sympy [F]

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + cx^4}} dx = \int \frac{A + Bx^2}{\sqrt{a + cx^4} (d + ex^2)^3} dx$$

input `integrate((B*x**2+A)/(e*x**2+d)**3/(c*x**4+a)**(1/2),x)`

output `Integral((A + B*x**2)/(sqrt(a + c*x**4)*(d + e*x**2)**3), x)`

### 3.7.7 Maxima [F]

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + a} (ex^2 + d)^3} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(1/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(e*x^2 + d)^3), x)`

### 3.7.8 Giac [F]

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + a}(ex^2 + d)^3} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(1/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(e*x^2 + d)^3), x)`

### 3.7.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + a}(ex^2 + d)^3} dx$$

input `int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)^3),x)`

output `int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)^3), x)`

$$3.8 \quad \int \frac{(A+Bx^2)(d+ex^2)^3}{(a+cx^4)^{3/2}} dx$$

|       |   |    |
|-------|---|----|
| 3.8.1 | Optimal result . . . . .                            | 93 |
| 3.8.2 | Mathematica [C] (verified) . . . . .                | 94 |
| 3.8.3 | Rubi [A] (verified) . . . . .                       | 95 |
| 3.8.4 | Maple [C] (verified) . . . . .                      | 97 |
| 3.8.5 | Fricas [A] (verification not implemented) . . . . . | 97 |
| 3.8.6 | Sympy [F] . . . . .                                 | 98 |
| 3.8.7 | Maxima [F] . . . . .                                | 98 |
| 3.8.8 | Giac [F] . . . . .                                  | 99 |
| 3.8.9 | Mupad [F(-1)] . . . . .                             | 99 |

### 3.8.1 Optimal result

Integrand size = 28, antiderivative size = 912

$$\begin{aligned}
& \int \frac{(A+Bx^2)(d+ex^2)^3}{(a+cx^4)^{3/2}} dx = \frac{x(Actd^2 - 3ae^2) - aBe(3cd^2 - ae^2) + c(Bcd^3 + 3Acd^2e - 3aBde^2 - aAe^3)}{2ac^2\sqrt{a+cx^4}} \\
& + \frac{Be^3x\sqrt{a+cx^4}}{3c^2} + \frac{e^2(3Bd+ Ae)x\sqrt{a+cx^4}}{c^{3/2}(\sqrt{a} + \sqrt{cx^2})} \\
& - \frac{(Bcd^3 + 3Acd^2e - 3aBde^2 - aAe^3)x\sqrt{a+cx^4}}{2ac^{3/2}(\sqrt{a} + \sqrt{cx^2})} \\
& - \frac{\sqrt[4]{ae^2}(3Bd+ Ae)(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{c^{7/4}\sqrt{a+cx^4}} \\
& + \frac{(Bcd^3 + 3Acd^2e - 3aBde^2 - aAe^3)(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{2a^{3/4}c^{7/4}\sqrt{a+cx^4}} \\
& - \frac{a^{3/4}Be^3(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{6c^{9/4}\sqrt{a+cx^4}} \\
& + \frac{\sqrt[4]{ae^2}(3Bd+ Ae)(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2c^{7/4}\sqrt{a+cx^4}} \\
& + \frac{e(3Bcd^2 + 3Acde - aBe^2)(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2^4\sqrt{ac^9}\sqrt{a+cx^4}} \\
& + \frac{(Ac^2d^3 + a^2Be^3 - 3acde(Bd+ Ae) + a^{3/2}\sqrt{ce^2}(3Bd+ Ae) - \sqrt{ac^3/2}d^2(Bd+ 3Ae))(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}}}{4a^{5/4}c^{9/4}\sqrt{a+cx^4}}
\end{aligned}$$

output

```

1/2*x*(A*c*d*(-3*a*e^2+c*d^2)-a*B*e*(-a*e^2+3*c*d^2)+c*(-A*a*e^3+3*A*c*d^2
*e-3*B*a*d*e^2+B*c*d^3)*x^2)/a/c^2/(c*x^4+a)^(1/2)+1/3*B*e^3*x*(c*x^4+a)^(
1/2)/c^2+e^2*(A*e+3*B*d)*x*(c*x^4+a)^(1/2)/c^(3/2)/(a^(1/2)+x^2*c^(1/2))-1
/2*(-A*a*e^3+3*A*c*d^2*e-3*B*a*d*e^2+B*c*d^3)*x*(c*x^4+a)^(1/2)/a/c^(3/2)/
(a^(1/2)+x^2*c^(1/2))-a^(1/4)*e^2*(A*e+3*B*d)*(cos(2*arctan(c^(1/4)*x/a^(1
/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(
1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^
2*c^(1/2))^2)^(1/2)/c^(7/4)/(c*x^4+a)^(1/2)+1/2*(-A*a*e^3+3*A*c*d^2*e-3*B*
a*d*e^2+B*c*d^3)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c
^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))
*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2))^2)^(1/2)/a^(3/4)/c
^(7/4)/(c*x^4+a)^(1/2)-1/6*a^(3/4)*B*e^3*(cos(2*arctan(c^(1/4)*x/a^(1/4)))
^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*
x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(
1/2))^2)^(1/2)/c^(9/4)/(c*x^4+a)^(1/2)+1/2*a^(1/4)*e^2*(A*e+3*B*d)*(cos(2*
arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*Ellip
ticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((
c*x^4+a)/(a^(1/2)+x^2*c^(1/2))^2)^(1/2)/c^(7/4)/(c*x^4+a)^(1/2)+1/2*e*(3*
A*c*d*e-B*a*e^2+3*B*c*d^2)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(
2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4)))...

```

### 3.8.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.19 (sec) , antiderivative size = 222, normalized size of antiderivative = 0.24

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + cx^4)^{3/2}} dx = \frac{3Acx(cd^3 + ae^2(-3d + 2ex^2)) + aBex(5ae^2 + c(-9d^2 + 18dex^2 + 2e^2x^4)) +$$

input `Integrate[((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(3/2),x]`

output

```

(3*A*c*x*(c*d^3 + a*e^2*(-3*d + 2*e*x^2)) + a*B*e*x*(5*a*e^2 + c*(-9*d^2 +
18*d*e*x^2 + 2*e^2*x^4)) + (a*B*e*(9*c*d^2 - 5*a*e^2) + 3*A*c*d*(c*d^2 +
3*a*e^2))*x*sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)
/a)] + 2*c*(B*c*d^3 + 3*A*c*d^2*e - 9*a*B*d*e^2 - 3*a*A*e^3)*x^3*sqrt[1 +
(c*x^4)/a]*Hypergeometric2F1[3/4, 3/2, 7/4, -((c*x^4)/a)]/(6*a*c^2*sqrt[a
+ c*x^4])

```

---

3.8.  $\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+cx^4)^{3/2}} dx$

### 3.8.3 Rubi [A] (verified)

Time = 1.02 (sec) , antiderivative size = 912, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {2259, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + cx^4)^{3/2}} dx$$

↓ 2259

$$\int \left( \frac{e(-aBe^2 + 3Acde + 3Bcd^2)}{c^2\sqrt{a + cx^4}} + \frac{cx^2(-aAe^3 - 3aBde^2 + 3Acd^2e + Bcd^3) + Acd(cd^2 - 3ae^2) - aBe(3cd^2 - ae^2)}{c^2(a + cx^4)^{3/2}} \right) dx$$

↓ 2009

$$\frac{a^{3/4}B(\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4+a}{(\sqrt{cx^2+\sqrt{a}})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right) e^3}{6c^{9/4}\sqrt{cx^4 + a}} + \frac{Bx\sqrt{cx^4 + ae^3}}{3c^2} -$$

$$\frac{\sqrt[4]{a}(3Bd + Ae)(\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4+a}{(\sqrt{cx^2+\sqrt{a}})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right) e^2}{c^{7/4}\sqrt{cx^4 + a}} +$$

$$\frac{\sqrt[4]{a}(3Bd + Ae)(\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4+a}{(\sqrt{cx^2+\sqrt{a}})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right) e^2}{2c^{7/4}\sqrt{cx^4 + a}} +$$

$$\frac{(3Bd + Ae)x\sqrt{cx^4 + ae^2}}{c^{3/2}(\sqrt{cx^2 + \sqrt{a}})} +$$

$$\frac{(3Bcd^2 + 3Aced - aBe^2)(\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4+a}{(\sqrt{cx^2+\sqrt{a}})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right) e}{2\sqrt[4]{ac^9}\sqrt{cx^4 + a}} +$$

$$\frac{(Bcd^3 + 3Aced^2 - 3aBe^2d - aAe^3)(\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4+a}{(\sqrt{cx^2+\sqrt{a}})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{2a^{3/4}c^{7/4}\sqrt{cx^4 + a}} +$$

$$\frac{(Ac^2d^3 - \sqrt{ac^3/2}(Bd + 3Ae)d^2 - 3ace(Bd + Ae)d + a^2Be^3 + a^{3/2}\sqrt{ce^2}(3Bd + Ae))(\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4+a}{(\sqrt{cx^2+\sqrt{a}})^2}}}{4a^{5/4}c^{9/4}\sqrt{cx^4 + a}}$$

$$\frac{(Bcd^3 + 3Aced^2 - 3aBe^2d - aAe^3)x\sqrt{cx^4 + a}}{2ac^{3/2}(\sqrt{cx^2 + \sqrt{a}})} +$$

$$\frac{x(c(Bcd^3 + 3Aced^2 - 3aBe^2d - aAe^3)x^2 + Acd(cd^2 - 3ae^2) - aBe(3cd^2 - ae^2))}{2ac^2\sqrt{cx^4 + a}}$$

---

3.8.  $\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+cx^4)^{3/2}} dx$



input `Int[((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(3/2),x]`

output `(x*(A*c*d*(c*d^2 - 3*a*e^2) - a*B*e*(3*c*d^2 - a*e^2) + c*(B*c*d^3 + 3*A*c*d^2*e - 3*a*B*d*e^2 - a*A*e^3)*x^2))/(2*a*c^2*Sqrt[a + c*x^4]) + (B*e^3*x*Sqrt[a + c*x^4])/(3*c^2) + (e^2*(3*B*d + A*e)*x*Sqrt[a + c*x^4])/(c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2)) - ((B*c*d^3 + 3*A*c*d^2*e - 3*a*B*d*e^2 - a*A*e^3)*x*Sqrt[a + c*x^4])/(2*a*c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2)) - (a^(1/4)*e^2*(3*B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(c^(7/4)*Sqrt[a + c*x^4]) + ((B*c*d^3 + 3*A*c*d^2*e - 3*a*B*d*e^2 - a*A*e^3)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(3/4)*c^(7/4)*Sqrt[a + c*x^4]) - (a^(3/4)*B*e^3*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(6*c^(9/4)*Sqrt[a + c*x^4]) + (a^(1/4)*e^2*(3*B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*c^(7/4)*Sqrt[a + c*x^4]) + (e*(3*B*c*d^2 + 3*A*c*d*e - a*B*e^2)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*c^(9/4)*Sqrt[a + c*x^4]) + ((A*c^2*d^3 + a^2*B*e^3 - 3*a*c*d*e*(B*d + A*e) + a^(3/2)*Sqrt[c]*e^2*(3*B*d + A*e) - Sqrt[a]*c^(3/2)*d^2*(B*d + 3*A*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*c^(9/4)*Sqrt[a + c*x^4])`

### 3.8.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2259 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x] && IntegerQ[p + 1/2] && IntegerQ[q]`

---

3.8.  $\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+cx^4)^{3/2}} dx$

### 3.8.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 8.29 (sec) , antiderivative size = 426, normalized size of antiderivative = 0.47

| method   | result  |
|----------|---|
| elliptic | $-\frac{2c \left( \frac{(Aae^3 - 3Ac d^2 e + 3Bad e^2 - Bcd^3)x^3 + (3Aacd e^2 - A c^2 d^3 - a^2 B e^3 + 3Bac d^2 e)x}{4c^2 a} \right)}{\sqrt{(x^4 + \frac{a}{c})c}} + \frac{B e^3 x \sqrt{c x^4 + a}}{3c^2} + \frac{\left( \frac{e(3Acde - Ba e^2 + 3Bc d^2)}{c^2} \right)}{\sqrt{(x^4 + \frac{a}{c})c}}$  |
| default  | $A d^3 \left( \frac{x}{2a \sqrt{(x^4 + \frac{a}{c})c}} + \frac{\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}} F\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{2a \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{c x^4 + a}} \right) + B e^3 \left( \frac{xa}{2c^2 \sqrt{(x^4 + \frac{a}{c})c}} + \frac{x \sqrt{c x^4 + a}}{3c^2} - \frac{5a \sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}}{6c} \right)$  |
| risch    | $\frac{B e^3 x \sqrt{c x^4 + a}}{3c^2} + \frac{3A c^2 d^3 \left( \frac{x}{2a \sqrt{(x^4 + \frac{a}{c})c}} + \frac{\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}} F\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{2a \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{c x^4 + a}} \right) - a^2 B e^3 \left( \frac{x}{2a \sqrt{(x^4 + \frac{a}{c})c}} + \frac{\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}}{2a \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{c x^4 + a}} \right)}{\sqrt{(x^4 + \frac{a}{c})c}}$ |

input `int((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(3/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -2*c*(1/4/c^2*(A*a*e^3-3*A*c*d^2*e+3*B*a*d*e^2-B*c*d^3)/a*x^3+1/4/a/c^3*(3 \\ & *A*a*c*d*e^2-A*c^2*d^3-B*a^2*e^3+3*B*a*c*d^2*e)*x)/((x^4+a/c)*c)^(1/2)+1/3 \\ & *B*e^3*x*(c*x^4+a)^(1/2)/c^2+(e*(3*A*c*d*e-B*a*e^2+3*B*c*d^2)/c^2-1/2/c^2/ \\ & a*(3*A*a*c*d*e^2-A*c^2*d^3-B*a^2*e^3+3*B*a*c*d^2*e)-1/3*B*e^3/c^2*a)/(I/a^ \\ & (1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)* \\ & x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+I*(1/c \\ & *e^2*(A*e+3*B*d)+1/2/c*(A*a*e^3-3*A*c*d^2*e+3*B*a*d*e^2-B*c*d^3)/a)*a^(1/2 \\ & )/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c \\ & ^{(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF(x*(I/a^(1/2)*c^(1/2)) \\ & ^{(1/2),I)-EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I)) \end{aligned}$$

### 3.8.5 Fracas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 448, normalized size of antiderivative = 0.49

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + cx^4)^{3/2}} dx =$$

$$3((Bac^2d^3 + 3Aac^2d^2e - 9Ba^2cde^2 - 3Aa^2ce^3)x^5 + (Ba^2cd^3 + 3Aa^2cd^2e - 9Ba^3de^2 - 3Aa^3e^3)x)\sqrt{c}(-$$

3.8. 
$$\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+cx^4)^{3/2}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="fricas")`

output `-1/6*(3*((B*a*c^2*d^3 + 3*A*a*c^2*d^2*e - 9*B*a^2*c*d*e^2 - 3*A*a^2*c*e^3)*x^5 + (B*a^2*c*d^3 + 3*A*a^2*c*d^2*e - 9*B*a^3*d*e^2 - 3*A*a^3*e^3)*x)*sqrt(c)*(-a/c)^(3/4)*elliptic_e(arcsin((-a/c)^(1/4)/x), -1) - ((9*(A + B)*a*c^2*d^2*e - (9*A + 5*B)*a^2*c*e^3 + 3*(B*a*c^2 + A*c^3)*d^3 - 9*(3*B*a^2*c - A*a*c^2)*d*e^2)*x^5 + (9*(A + B)*a^2*c*d^2*e - (9*A + 5*B)*a^3*e^3 + 3*(B*a^2*c + A*a*c^2)*d^3 - 9*(3*B*a^3 - A*a^2*c)*d*e^2)*x)*sqrt(c)*(-a/c)^(3/4)*elliptic_f(arcsin((-a/c)^(1/4)/x), -1) - (2*B*a^2*c*e^3*x^6 - 3*B*a^2*c*d^3 - 9*A*a^2*c*d^2*e + 27*B*a^3*d*e^2 + 9*A*a^3*e^3 + 6*(3*B*a^2*c*d*e^2 + A*a^2*c*e^3)*x^4 + (3*A*a*c^2*d^3 - 9*B*a^2*c*d^2*e - 9*A*a^2*c*d*e^2 + 5*B*a^3*e^3)*x^2)*sqrt(c*x^4 + a))/(a^2*c^3*x^5 + a^3*c^2*x)`

### 3.8.6 Sympy [F]

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + cx^4)^{3/2}} dx = \int \frac{(A + Bx^2)(d + ex^2)^3}{(a + cx^4)^{\frac{3}{2}}} dx$$

input `integrate((B*x**2+A)*(e*x**2+d)**3/(c*x**4+a)**(3/2),x)`

output `Integral((A + B*x**2)*(d + e*x**2)**3/(a + c*x**4)**(3/2), x)`

### 3.8.7 Maxima [F]

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^3}{(cx^4 + a)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^3/(c*x^4 + a)^(3/2), x)`

**3.8.8 Giac [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^3}{(cx^4 + a)^{3/2}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^3/(c*x^4 + a)^(3/2), x)`

**3.8.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^3}{(cx^4 + a)^{3/2}} dx$$

input `int(((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(3/2),x)`

output `int(((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(3/2), x)`

$$3.9 \quad \int \frac{(A+Bx^2)(d+ex^2)^2}{(a+cx^4)^{3/2}} dx$$

|       |   |     |
|-------|---|-----|
| 3.9.1 | Optimal result                            | 100 |
| 3.9.2 | Mathematica [C] (verified)                | 101 |
| 3.9.3 | Rubi [A] (verified)                       | 102 |
| 3.9.4 | Maple [C] (verified)                      | 103 |
| 3.9.5 | Fricas [A] (verification not implemented) | 104 |
| 3.9.6 | Sympy [F]                                 | 105 |
| 3.9.7 | Maxima [F]                                | 105 |
| 3.9.8 | Giac [F]                                  | 105 |
| 3.9.9 | Mupad [F(-1)]                             | 106 |

### 3.9.1 Optimal result

Integrand size = 28, antiderivative size = 694

$$\begin{aligned} & \int \frac{(A+Bx^2)(d+ex^2)^2}{(a+cx^4)^{3/2}} dx = \frac{x(Acd^2 - 2aBde - aAe^2 + (Bcd^2 + 2Acde - aBe^2)x^2)}{2ac\sqrt{a+cx^4}} \\ & + \frac{Be^2x\sqrt{a+cx^4}}{c^{3/2}(\sqrt{a} + \sqrt{cx^2})} - \frac{(Bcd^2 + 2Acde - aBe^2)x\sqrt{a+cx^4}}{2ac^{3/2}(\sqrt{a} + \sqrt{cx^2})} \\ & - \frac{\sqrt[4]{a}Be^2(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{c^{7/4}\sqrt{a+cx^4}} \\ & + \frac{(Bcd^2 + 2Acde - aBe^2)(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{2a^{3/4}c^{7/4}\sqrt{a+cx^4}} \\ & + \frac{\sqrt[4]{a}Be^2(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2c^{7/4}\sqrt{a+cx^4}} \\ & + \frac{e(2Bd + Ae)(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2^4\sqrt{ac^5}\sqrt{a+cx^4}} \\ & - \frac{\left(Bcd^2 + 2Acde - aBe^2 - \frac{\sqrt{c}(Acd^2 - 2aBde - aAe^2)}{\sqrt{a}}\right)(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{4a^{3/4}c^{7/4}\sqrt{a+cx^4}} \end{aligned}$$

---


$$3.9. \quad \int \frac{(A+Bx^2)(d+ex^2)^2}{(a+cx^4)^{3/2}} dx$$

output  $\frac{1}{2}x(Acd^2 - 2aBde - aAe^2 + (2Acd - Bae^2 + Bcd^2)x^2)/a/c/(cx^4+a)^{(1/2)} + B^2e^2x^2/(cx^4+a)^{(1/2)}/c^{(3/2)}/(a^{(1/2)}+x^2c^{(1/2)}) - \frac{1}{2}(2Acd - Bae^2 + Bcd^2)x^2/(cx^4+a)^{(1/2)}/a/c^{(3/2)}/(a^{(1/2)}+x^2c^{(1/2)}) - a^{(1/4)}B^2e^2(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)})) * \text{EllipticE}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})), 1/2, 2^{(1/2)}) * (a^{(1/2)}+x^2c^{(1/2)}) * ((cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)}))^2)^{(1/2)}/c^{(7/4)}/(cx^4+a)^{(1/2)} + \frac{1}{2}(2Acd - Bae^2 + Bcd^2)(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)})) * \text{EllipticE}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})), 1/2, 2^{(1/2)}) * (a^{(1/2)}+x^2c^{(1/2)}) * ((cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)}))^2)^{(1/2)}/a^{(3/4)}/c^{(7/4)}/(cx^4+a)^{(1/2)} + \frac{1}{2}a^{(1/4)}B^2e^2(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)})) * \text{EllipticF}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})), 1/2, 2^{(1/2)}) * (a^{(1/2)}+x^2c^{(1/2)}) * ((cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)}))^2)^{(1/2)}/c^{(7/4)}/(cx^4+a)^{(1/2)} + \frac{1}{2}e^2(Ae^2 + 2Bd)(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)})) * \text{EllipticF}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})), 1/2, 2^{(1/2)}) * (a^{(1/2)}+x^2c^{(1/2)}) * ((cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)}))^2)^{(1/2)}/a^{(1/4)}/c^{(5/4)}/(cx^4+a)^{(1/2)} - \frac{1}{4}(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)})) * \text{EllipticF}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})), 1/2, 2^{(1/2)}) * (a^{(1/2)}+x^2c^{(1/2)}) * (Bcd^2 + 2Acd - Bae^2 - (-Aae^2 + Acd^2 - 2Bde)) * c^{(1/2)}/a^{(1/2)}) * ((cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)}))^2)^{(1/2)}/a^{(3...}$

### 3.9.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.14 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.24

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + cx^4)^{3/2}} dx = \frac{3A(cd^2 - ae^2)x + 6aBex(-d + ex^2) + 3(Acd^2 + 2aBde + aAe^2)x\sqrt{1 + \frac{cx^4}{a}}}{(a + cx^4)^{3/2}}$$

input `Integrate[((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(3/2), x]`

output  $(3A(c*d^2 - a*e^2)*x + 6*a*B*e*x*(-d + e*x^2) + 3*(A*c*d^2 + 2*a*B*d*e + a*A*e^2)*x*\text{Sqrt}[1 + (c*x^4)/a]*\text{Hypergeometric2F1}[1/4, 1/2, 5/4, -((c*x^4)/a)] + 2*(B*c*d^2 + 2*A*c*d*e - 3*a*B*e^2)*x^3*\text{Sqrt}[1 + (c*x^4)/a]*\text{Hypergeometric2F1}[3/4, 3/2, 7/4, -((c*x^4)/a)]/(6*a*c*\text{Sqrt}[a + c*x^4])$

### 3.9.3 Rubi [A] (verified)

Time = 0.79 (sec) , antiderivative size = 694, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {2259, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + cx^4)^{3/2}} dx$$

↓ 2259

$$\int \left( \frac{x^2(-aBe^2 + 2Acde + Bcd^2) - aAe^2 - 2aBde + Acd^2}{c(a + cx^4)^{3/2}} + \frac{e(Ae + 2Bd)}{c\sqrt{a + cx^4}} + \frac{Be^2x^2}{c\sqrt{a + cx^4}} \right) dx$$

↓ 2009

$$\frac{(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \text{EllipticF} \left( 2 \arctan \left( \frac{\sqrt[4]{cx^2}}{\sqrt[4]{a}} \right), \frac{1}{2} \right) \left( -\frac{\sqrt{c}(-aAe^2 - 2aBde + Acd^2)}{\sqrt{a}} - aBe^2 + 2Acde + Bcd^2 \right)}{4a^{3/4}c^{7/4}\sqrt{a + cx^4}} + \frac{(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} E \left( 2 \arctan \left( \frac{\sqrt[4]{cx^2}}{\sqrt[4]{a}} \right) \middle| \frac{1}{2} \right) (-aBe^2 + 2Acde + Bcd^2)}{2a^{3/4}c^{7/4}\sqrt{a + cx^4}} + \frac{e(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} (Ae + 2Bd) \text{EllipticF} \left( 2 \arctan \left( \frac{\sqrt[4]{cx^2}}{\sqrt[4]{a}} \right), \frac{1}{2} \right)}{2\sqrt[4]{ac^5}\sqrt{a + cx^4}} - \frac{x\sqrt{a + cx^4}(-aBe^2 + 2Acde + Bcd^2)}{2ac^{3/2}(\sqrt{a} + \sqrt{cx^2})} + \frac{x(x^2(-aBe^2 + 2Acde + Bcd^2) - aAe^2 - 2aBde + Acd^2)}{2ac\sqrt{a + cx^4}} + \frac{\sqrt[4]{a}Be^2(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \text{EllipticF} \left( 2 \arctan \left( \frac{\sqrt[4]{cx^2}}{\sqrt[4]{a}} \right), \frac{1}{2} \right)}{2c^{7/4}\sqrt{a + cx^4}} - \frac{\sqrt[4]{a}Be^2(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} E \left( 2 \arctan \left( \frac{\sqrt[4]{cx^2}}{\sqrt[4]{a}} \right) \middle| \frac{1}{2} \right)}{c^{7/4}\sqrt{a + cx^4}} + \frac{Be^2x\sqrt{a + cx^4}}{c^{3/2}(\sqrt{a} + \sqrt{cx^2})}$$

input `Int[((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(3/2), x]`

---

3.9.  $\int \frac{(A+Bx^2)(d+ex^2)^2}{(a+cx^4)^{3/2}} dx$

```
output (x*(A*c*d^2 - 2*a*B*d*e - a*A*e^2 + (B*c*d^2 + 2*A*c*d*e - a*B*e^2)*x^2))/
(2*a*c*Sqrt[a + c*x^4]) + (B*e^2*x*Sqrt[a + c*x^4])/(c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2)) -
((B*c*d^2 + 2*A*c*d*e - a*B*e^2)*x*Sqrt[a + c*x^4])/(2*a*c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2)) -
(a^(1/4)*B*e^2*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(c^(7/4)*Sqrt[a + c*x^4]) +
((B*c*d^2 + 2*A*c*d*e - a*B*e^2)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(3/4)*c^(7/4)*Sqrt[a + c*x^4]) +
(a^(1/4)*B*e^2*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*c^(7/4)*Sqrt[a + c*x^4]) +
(e*(2*B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*c^(5/4)*Sqrt[a + c*x^4]) -
((B*c*d^2 + 2*A*c*d*e - a*B*e^2 - (Sqrt[c]*(A*c*d^2 - 2*a*B*d*e - a*A*e^2))/Sqrt[a])*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(4*a^(3/4)*c^(7/4)*Sqrt[a + c*x^4])
```

### 3.9.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2259 Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol]
:= Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x] && IntegerQ[p + 1/2] && IntegerQ[q]
```

### 3.9.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.55 (sec) , antiderivative size = 323, normalized size of antiderivative = 0.47

| method   | result  |
|----------|---|
| elliptic | $-\frac{2c \left( -\frac{(2Acde - Ba e^2 + Bc d^2)x^3 + (aA e^2 - Ac d^2 + 2aBde)x}{4a c^2} \right)}{\sqrt{(x^4 + \frac{a}{c})c}} + \frac{\left( \frac{e(Ae + 2Bd) - aA e^2 - Ac d^2 + 2aBde}{2ac} \right) \sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}} F\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a}}$   |
| default  | $d^2 A \left( \frac{x}{2a \sqrt{(x^4 + \frac{a}{c})c}} + \frac{\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}} F\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\right)}{2a \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a}} \right) + B e^2 \left( -\frac{x^3}{2c \sqrt{(x^4 + \frac{a}{c})c}} + \frac{3i\sqrt{a} \sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}}{2c^{\frac{3}{2}} \sqrt{cx^4 + a}} \right)$ |

3.9.  $\int \frac{(A+Bx^2)(d+ex^2)^2}{(a+cx^4)^{3/2}} dx$



input `int((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(3/2),x,method=_RETURNVERBOSE)`

output `-2*c*(-1/4*(2*A*c*d*e-B*a*e^2+B*c*d^2)/a/c^2*x^3+1/4*(A*a*e^2-A*c*d^2+2*B*a*d*e)/c^2/a*x)/((x^4+a/c)*c)^(1/2)+(e*(A*e+2*B*d)/c-1/2*(A*a*e^2-A*c*d^2+2*B*a*d*e)/a/c)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+I*(1/c*B*e^2-1/2*(2*A*c*d*e-B*a*e^2+B*c*d^2)/a/c)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)-EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I))`

### 3.9.5 Fracas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 332, normalized size of antiderivative = 0.48

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + cx^4)^{3/2}} dx =$$

$$\left( (Bac^2d^2 + 2Aac^2de - 3Ba^2ce^2)x^5 + (Ba^2cd^2 + 2Aa^2cde - 3Ba^3e^2)x \right) \sqrt{c} \left( -\frac{a}{c} \right)^{\frac{3}{4}} E\left( \arcsin\left( \frac{(-a/c)^{\frac{1}{4}}}{x} \right) \mid - \right)$$

input `integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="fricas")`

output `-1/2*(((B*a*c^2*d^2 + 2*A*a*c^2*d*e - 3*B*a^2*c*e^2)*x^5 + (B*a^2*c*d^2 + 2*A*a^2*c*d*e - 3*B*a^3*e^2)*x)*sqrt(c)*(-a/c)^(3/4)*elliptic_e(arcsin((-a/c)^(1/4)/x), -1) - ((2*(A + B)*a*c^2*d*e + (B*a*c^2 + A*c^3)*d^2 - (3*B*a^2*c - A*a*c^2)*e^2)*x^5 + (2*(A + B)*a^2*c*d*e + (B*a^2*c + A*a*c^2)*d^2 - (3*B*a^3 - A*a^2*c)*e^2)*x)*sqrt(c)*(-a/c)^(3/4)*elliptic_f(arcsin((-a/c)^(1/4)/x), -1) - (2*B*a^2*c*e^2*x^4 - B*a^2*c*d^2 - 2*A*a^2*c*d*e + 3*B*a^3*e^2 + (A*a*c^2*d^2 - 2*B*a^2*c*d*e - A*a^2*c*e^2)*x^2)*sqrt(c*x^4 + a)/(a^2*c^3*x^5 + a^3*c^2*x)`

---

3.9.  $\int \frac{(A+Bx^2)(d+ex^2)^2}{(a+cx^4)^{3/2}} dx$

**3.9.6 Sympy [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + cx^4)^{3/2}} dx = \int \frac{(A + Bx^2)(d + ex^2)^2}{(a + cx^4)^{\frac{3}{2}}} dx$$

input `integrate((B*x**2+A)*(e*x**2+d)**2/(c*x**4+a)**(3/2),x)`

output `Integral((A + B*x**2)*(d + e*x**2)**2/(a + c*x**4)**(3/2), x)`

**3.9.7 Maxima [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^2}{(cx^4 + a)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^2/(c*x^4 + a)^(3/2), x)`

**3.9.8 Giac [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^2}{(cx^4 + a)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^2/(c*x^4 + a)^(3/2), x)`

**3.9.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^2}{(cx^4 + a)^{3/2}} dx$$

input `int(((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(3/2),x)`output `int(((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(3/2), x)`

**3.10** 
$$\int \frac{(A+Bx^2)(d+ex^2)}{(a+cx^4)^{3/2}} dx$$

3.10.1 Optimal result . . . . . 107  
 3.10.2 Mathematica [C] (verified) . . . . . 108  
 3.10.3 Rubi [A] (verified) . . . . . 108  
 3.10.4 Maple [C] (verified) . . . . . 110  
 3.10.5 Fricas [A] (verification not implemented) . . . . . 111  
 3.10.6 Sympy [C] (verification not implemented) . . . . . 111  
 3.10.7 Maxima [F] . . . . . 112  
 3.10.8 Giac [F] . . . . . 112  
 3.10.9 Mupad [F(-1)] . . . . . 112

**3.10.1 Optimal result**

Integrand size = 26, antiderivative size = 395

$$\int \frac{(A+Bx^2)(d+ex^2)}{(a+cx^4)^{3/2}} dx = \frac{x(Acd - aBe + c(Bd + Ae)x^2)}{2ac\sqrt{a+cx^4}} - \frac{(Bd + Ae)x\sqrt{a+cx^4}}{2a\sqrt{c}(\sqrt{a} + \sqrt{cx^2})}$$

$$+ \frac{(Bd + Ae)(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{2a^{3/4}c^{3/4}\sqrt{a+cx^4}}$$

$$+ \frac{Be(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2\sqrt[4]{ac^5}\sqrt{a+cx^4}}$$

$$+ \frac{(Acd - aBe - \sqrt{a}\sqrt{c}(Bd + Ae))(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{4a^{5/4}c^{5/4}\sqrt{a+cx^4}}$$

output  $\frac{1}{2}x(Ac*d - B*a*e + c*(A*e + B*d)*x^2)/a/c/(c*x^4+a)^{(1/2)} - \frac{1}{2}*(A*e + B*d)*x*(c*x^4+a)^{(1/2)}/a/c^{(1/2)}/(a^{(1/2)}+x^2*c^{(1/2)}) + \frac{1}{2}*(A*e + B*d)*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), 1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/a^{(3/4)}/c^{(3/4)}/(c*x^4+a)^{(1/2)} + \frac{1}{2}*B*e*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), 1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/a^{(1/4)}/c^{(5/4)}/(c*x^4+a)^{(1/2)} + \frac{1}{4}*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), 1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*(Ac*d - B*a*e - (A*e + B*d)*a^{(1/2)}*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/a^{(5/4)}/c^{(5/4)}/(c*x^4+a)^{(1/2)}$

### 3.10.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.08 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.32

$$\int \frac{(A + Bx^2)(d + ex^2)}{(a + cx^4)^{3/2}} dx = \frac{3(Acd - aBe)x + 3(Acd + aBe)x\sqrt{1 + \frac{cx^4}{a}} \text{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -\frac{cx^4}{a}\right)}{6ac\sqrt{a + cx^4}}$$

input `Integrate[((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(3/2), x]`

output  $(3*(A*c*d - a*B*e)*x + 3*(A*c*d + a*B*e)*x*\text{Sqrt}[1 + (c*x^4)/a]*\text{Hypergeometric2F1}[1/4, 1/2, 5/4, -((c*x^4)/a)] + 2*c*(B*d + A*e)*x^3*\text{Sqrt}[1 + (c*x^4)/a]*\text{Hypergeometric2F1}[3/4, 3/2, 7/4, -((c*x^4)/a)]/(6*a*c*\text{Sqrt}[a + c*x^4])$

### 3.10.3 Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 395, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$ , Rules used = {2259, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.10.  $\int \frac{(A+Bx^2)(d+ex^2)}{(a+cx^4)^{3/2}} dx$

$$\begin{aligned}
& \int \frac{(A + Bx^2)(d + ex^2)}{(a + cx^4)^{3/2}} dx \\
& \quad \downarrow \text{2259} \\
& \int \left( \frac{-aBe + cx^2(Ae + Bd) + Acd}{c(a + cx^4)^{3/2}} + \frac{Be}{c\sqrt{a + cx^4}} \right) dx \\
& \quad \downarrow \text{2009} \\
& \frac{(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \operatorname{EllipticF} \left( 2 \arctan \left( \frac{\sqrt[4]{cx}}{\sqrt{a}} \right), \frac{1}{2} \right) (-\sqrt{a}\sqrt{c}(Ae + Bd) - aBe + Acd)}{4a^{5/4}c^{5/4}\sqrt{a + cx^4}} + \\
& \frac{(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} (Ae + Bd) E \left( 2 \arctan \left( \frac{\sqrt[4]{cx}}{\sqrt{a}} \right) \middle| \frac{1}{2} \right)}{2a^{3/4}c^{3/4}\sqrt{a + cx^4}} - \frac{x\sqrt{a + cx^4}(Ae + Bd)}{2a\sqrt{c}(\sqrt{a} + \sqrt{cx^2})} + \\
& \frac{x(-aBe + cx^2(Ae + Bd) + Acd)}{2ac\sqrt{a + cx^4}} + \\
& \frac{Be(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \operatorname{EllipticF} \left( 2 \arctan \left( \frac{\sqrt[4]{cx}}{\sqrt{a}} \right), \frac{1}{2} \right)}{2\sqrt[4]{ac^5}\sqrt{a + cx^4}}
\end{aligned}$$

input `Int[((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(3/2),x]`

output `(x*(A*c*d - a*B*e + c*(B*d + A*e)*x^2))/(2*a*c*Sqrt[a + c*x^4]) - ((B*d + A*e)*x*Sqrt[a + c*x^4])/(2*a*Sqrt[c]*(Sqrt[a] + Sqrt[c]*x^2)) + ((B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(3/4)*c^(3/4)*Sqrt[a + c*x^4]) + (B*e*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*c^(5/4)*Sqrt[a + c*x^4]) + ((A*c*d - a*B*e - Sqrt[a]*Sqrt[c]*(B*d + A*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(4*a^(5/4)*c^(5/4)*Sqrt[a + c*x^4])`

## 3.10.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2259 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x] && IntegerQ[p + 1/2] && IntegerQ[q]`

## 3.10.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.69 (sec) , antiderivative size = 253, normalized size of antiderivative = 0.64

| method   | result   |
|----------|--|
| elliptic | $-\frac{2c\left(-\frac{(Ae+Bd)x^3}{4ca}-\frac{(Acd-Bae)x}{4ac^2}\right)}{\sqrt{(x^4+\frac{a}{c})c}} + \frac{\left(\frac{eB}{c} + \frac{Acd-Bae}{2ac}\right)\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}} - \frac{i(Ae+Bd)\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}}{2\sqrt{a}\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}}$  |
| default  | $Ad\left(\frac{x}{2a\sqrt{(x^4+\frac{a}{c})c}} + \frac{\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{2a\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}}\right) + eB\left(-\frac{x}{2c\sqrt{(x^4+\frac{a}{c})c}} + \frac{\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{2c\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}}\right)$ |

input `int((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(3/2), x, method=_RETURNVERBOSE)`

output `-2*c*(-1/4*(A*e+B*d)/c/a*x^3-1/4*(A*c*d-B*a*e)/a/c^2*x)/((x^4+a/c)*c)^(1/2)+(e*B/c+1/2*(A*c*d-B*a*e)/a/c)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2), I)-1/2*I*(A*e+B*d)/a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2), I)-EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2), I))`

$$3.10. \int \frac{(A+Bx^2)(d+ex^2)}{(a+cx^4)^{3/2}} dx$$

### 3.10.5 Fracas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 188, normalized size of antiderivative = 0.48

$$\int \frac{(A + Bx^2)(d + ex^2)}{(a + cx^4)^{3/2}} dx = \frac{((Bc^2d + Ac^2e)x^4 + Bacd + Aace)\sqrt{a}\left(-\frac{c}{a}\right)^{\frac{3}{4}} E(\arcsin\left(x\left(-\frac{c}{a}\right)^{\frac{1}{4}}\right) | -1) - ((A + B)c^2d + (B^2ac + A^2c^2)e)x^4 + (A + B)ac^2d + (B^2a^2 + A^2ac)e\sqrt{a}\left(-\frac{c}{a}\right)^{\frac{3}{4}} \text{elliptic}_f(\arcsin(x(-c/a)^{1/4}), -1) + \sqrt{cx^4 + a}((Bc^2d + Ac^2e)x^3 + (A^2c^2d - B^2ac^2e)x)}{(a^2c^3x^4 + a^2c^2)}$$

input `integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="fracas")`

output `1/2*((B*c^2*d + A*c^2*e)*x^4 + B*a*c*d + A*a*c*e)*sqrt(a)*(-c/a)^(3/4)*elliptic_e(arcsin(x*(-c/a)^(1/4)), -1) - (((A + B)*c^2*d + (B*a*c + A*c^2)*e)*x^4 + (A + B)*a*c*d + (B*a^2 + A*a*c)*e)*sqrt(a)*(-c/a)^(3/4)*elliptic_f(arcsin(x*(-c/a)^(1/4)), -1) + sqrt(c*x^4 + a)*((B*c^2*d + A*c^2*e)*x^3 + (A*c^2*d - B*a*c*e)*x)/(a*c^3*x^4 + a^2*c^2)`

### 3.10.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 4.47 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.42

$$\int \frac{(A + Bx^2)(d + ex^2)}{(a + cx^4)^{3/2}} dx = \frac{Adx\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}}\Gamma\left(\frac{5}{4}\right)} + \frac{Aex^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{3}{4}, \frac{3}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}}\Gamma\left(\frac{7}{4}\right)} + \frac{Bdx^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{3}{4}, \frac{3}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}}\Gamma\left(\frac{7}{4}\right)} + \frac{Bex^5\Gamma\left(\frac{5}{4}\right) {}_2F_1\left(\frac{5}{4}, \frac{3}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}}\Gamma\left(\frac{9}{4}\right)}$$

input `integrate((B*x**2+A)*(e*x**2+d)/(c*x**4+a)**(3/2),x)`

output `A*d*x*gamma(1/4)*hyper((1/4, 3/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*a**3/2)*gamma(5/4) + A*e*x**3*gamma(3/4)*hyper((3/4, 3/2), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*a**3/2)*gamma(7/4) + B*d*x**3*gamma(3/4)*hyper((3/4, 3/2), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*a**3/2)*gamma(7/4) + B*e*x**5*gamma(5/4)*hyper((5/4, 3/2), (9/4,), c*x**4*exp_polar(I*pi)/a)/(4*a**3/2)*gamma(9/4)`

---

3.10.  $\int \frac{(A+Bx^2)(d+ex^2)}{(a+cx^4)^{3/2}} dx$



**3.10.7 Maxima [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)}{(a + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)}{(cx^4 + a)^{3/2}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)*(e*x^2 + d)/(c*x^4 + a)^(3/2), x)`

**3.10.8 Giac [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)}{(a + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)}{(cx^4 + a)^{3/2}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)*(e*x^2 + d)/(c*x^4 + a)^(3/2), x)`

**3.10.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)}{(a + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)}{(cx^4 + a)^{3/2}} dx$$

input `int(((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(3/2),x)`

output `int(((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(3/2), x)`

### 3.11 $\int \frac{A+Bx^2}{(a+cx^4)^{3/2}} dx$

|        |   |     |
|--------|---|-----|
| 3.11.1 | Optimal result                            | 113 |
| 3.11.2 | Mathematica [C] (verified)                | 114 |
| 3.11.3 | Rubi [A] (verified)                       | 114 |
| 3.11.4 | Maple [C] (verified)                      | 116 |
| 3.11.5 | Fricas [A] (verification not implemented) | 117 |
| 3.11.6 | Sympy [C] (verification not implemented)  | 117 |
| 3.11.7 | Maxima [F]                                | 118 |
| 3.11.8 | Giac [F]                                  | 118 |
| 3.11.9 | Mupad [F(-1)]                             | 118 |

#### 3.11.1 Optimal result

Integrand size = 19, antiderivative size = 262

$$\int \frac{A+Bx^2}{(a+cx^4)^{3/2}} dx = \frac{x(A+Bx^2)}{2a\sqrt{a+cx^4}} - \frac{Bx\sqrt{a+cx^4}}{2a\sqrt{c}(\sqrt{a}+\sqrt{cx^2})} + \frac{B(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)}{2a^{3/4}c^{3/4}\sqrt{a+cx^4}} - \frac{(\sqrt{a}B-A\sqrt{c})(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right),\frac{1}{2}\right)}{4a^{5/4}c^{3/4}\sqrt{a+cx^4}}$$

output

```
1/2*x*(B*x^2+A)/a/(c*x^4+a)^(1/2)-1/2*B*x*(c*x^4+a)^(1/2)/a/c^(1/2)/(a^(1/2)+x^2*c^(1/2))+1/2*B*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(3/4)/c^(3/4)/(c*x^4+a)^(1/2)-1/4*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(B*a^(1/2)-A*c^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(5/4)/c^(3/4)/(c*x^4+a)^(1/2)
```

### 3.11.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.04 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.38

$$\int \frac{A + Bx^2}{(a + cx^4)^{3/2}} dx = \frac{3Ax + 3Ax\sqrt{1 + \frac{cx^4}{a}} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -\frac{cx^4}{a}\right) + 2Bx^3\sqrt{1 + \frac{cx^4}{a}} \operatorname{Hypergeometric2F1}\left(\frac{3}{4}, \frac{3}{2}, \frac{7}{4}, -\frac{cx^4}{a}\right)}{6a\sqrt{a + cx^4}}$$

input `Integrate[(A + B*x^2)/(a + c*x^4)^(3/2), x]`

output `(3*A*x + 3*A*x*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + 2*B*x^3*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[3/4, 3/2, 7/4, -((c*x^4)/a)])/(6*a*Sqrt[a + c*x^4])`

### 3.11.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$ , Rules used = {1493, 25, 1512, 27, 761, 1510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + Bx^2}{(a + cx^4)^{3/2}} dx \\ & \quad \downarrow \text{1493} \\ & \frac{x(A + Bx^2)}{2a\sqrt{a + cx^4}} - \frac{\int -\frac{A - Bx^2}{\sqrt{cx^4 + a}} dx}{2a} \\ & \quad \downarrow \text{25} \\ & \frac{\int \frac{A - Bx^2}{\sqrt{cx^4 + a}} dx}{2a} + \frac{x(A + Bx^2)}{2a\sqrt{a + cx^4}} \\ & \quad \downarrow \text{1512} \\ & \frac{\left(A - \frac{\sqrt{a}B}{\sqrt{c}}\right) \int \frac{1}{\sqrt{cx^4 + a}} dx + \frac{\sqrt{a}B \int \frac{\sqrt{a} - \sqrt{cx^2}}{\sqrt{a}\sqrt{cx^4 + a}} dx}{\sqrt{c}}}{2a} + \frac{x(A + Bx^2)}{2a\sqrt{a + cx^4}} \end{aligned}$$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{\left(A - \frac{\sqrt{a}B}{\sqrt{c}}\right) \int \frac{1}{\sqrt{cx^4+a}} dx + \frac{B \int \frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{cx^4+a}} dx}{\sqrt{c}}}{2a} + \frac{x(A+Bx^2)}{2a\sqrt{a+cx^4}} \\
 & \downarrow 761 \\
 & \frac{\frac{B \int \frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{cx^4+a}} dx}{\sqrt{c}} + \frac{(\sqrt{a}+\sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \left(A - \frac{\sqrt{a}B}{\sqrt{c}}\right) \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2\sqrt[4]{a}\sqrt[4]{c}\sqrt{a+cx^4}}}{2a} + \frac{x(A+Bx^2)}{2a\sqrt{a+cx^4}} \\
 & \downarrow 1510 \\
 & \frac{(\sqrt{a}+\sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \left(A - \frac{\sqrt{a}B}{\sqrt{c}}\right) \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2\sqrt[4]{a}\sqrt[4]{c}\sqrt{a+cx^4}} + \frac{B \left(\frac{\sqrt[4]{a}(\sqrt{a}+\sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{\sqrt[4]{c}\sqrt{a+cx^4}} - \frac{x\sqrt{a+cx^2}}{\sqrt{a+cx^4}}\right)}{\sqrt{c}}}{2a} \\
 & \frac{x(A+Bx^2)}{2a\sqrt{a+cx^4}}
 \end{aligned}$$

input `Int[(A + B*x^2)/(a + c*x^4)^(3/2),x]`

output `(x*(A + B*x^2))/(2*a*Sqrt[a + c*x^4]) + ((B*(-((x*Sqrt[a + c*x^4]))/(Sqrt[a] + Sqrt[c]*x^2)) + (a^(1/4)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]))/(c^(1/4)*Sqrt[a + c*x^4]))/Sqrt[c] + ((A - (Sqrt[a]*B)/Sqrt[c])*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]]/(2*a^(1/4)*c^(1/4)*Sqrt[a + c*x^4]))/(2*a)`

### 3.11.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 1493 `Int[((d_) + (e_.)*(x_)^2)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)*((a + c*x^4)^(p + 1)/(4*a*(p + 1))), x] + Simp[1/(4*a*(p + 1)) Int[Simp[d*(4*p + 5) + e*(4*p + 7)*x^2, x]*(a + c*x^4)^(p + 1), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && IntegerQ[2*p]`

rule 1510 `Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2])]/(q*Sqrt[a + c*x^4))*EllipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]`

rule 1512 `Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 2]}, Simp[(e + d*q)/q Int[1/Sqrt[a + c*x^4], x], x] - Simp[e/q Int[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]`

### 3.11.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.56 (sec) , antiderivative size = 212, normalized size of antiderivative = 0.81

| method   | result   |
|----------|--|
| elliptic | $-\frac{2c\left(-\frac{Bx^3}{4ac} - \frac{Ax}{4ac}\right)}{\sqrt{\left(x^4 + \frac{a}{c}\right)c}} + \frac{A\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}\operatorname{F}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{2a\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}} - \frac{iB\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}\left(\operatorname{F}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) - E\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)\right)}{2\sqrt{a}\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}\sqrt{c}}$                                   |
| default  | $A\left(\frac{x}{2a\sqrt{\left(x^4 + \frac{a}{c}\right)c}} + \frac{\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}\operatorname{F}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{2a\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}}\right) + B\left(\frac{x^3}{2a\sqrt{\left(x^4 + \frac{a}{c}\right)c}} - \frac{i\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}\left(\operatorname{F}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) - E\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)\right)}{2\sqrt{a}\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}\sqrt{c}}\right)$ |

input `int((B*x^2+A)/(c*x^4+a)^(3/2),x,method=_RETURNVERBOSE)`

output 
$$-2*c*(-1/4/a*B/c*x^3-1/4*A/a/c*x)/((x^4+a/c)*c)^(1/2)+1/2*A/a/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)-1/2*I/a^(1/2)*B/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)-EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I))$$

### 3.11.5 Fracas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.44

$$\int \frac{A + Bx^2}{(a + cx^4)^{3/2}} dx = \frac{(Bcx^4 + Ba)\sqrt{a}\left(-\frac{c}{a}\right)^{\frac{3}{4}} E\left(\arcsin\left(x\left(-\frac{c}{a}\right)^{\frac{1}{4}}\right) \mid -1\right) - ((A + B)cx^4 + (A + B)a)\sqrt{a}\left(-\frac{c}{a}\right)^{\frac{1}{4}}}{2(ac^2x^4 + a^2c)}$$

input `integrate((B*x^2+A)/(c*x^4+a)^(3/2),x, algorithm="fricas")`

output 
$$1/2*((B*c*x^4 + B*a)*sqrt(a)*(-c/a)^(3/4)*elliptic_e(arcsin(x*(-c/a)^(1/4)), -1) - ((A + B)*c*x^4 + (A + B)*a)*sqrt(a)*(-c/a)^(3/4)*elliptic_f(arcsin(x*(-c/a)^(1/4)), -1) + (B*c*x^3 + A*c*x)*sqrt(c*x^4 + a)/(a*c^2*x^4 + a^2*c)$$

### 3.11.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.47 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.30

$$\int \frac{A + Bx^2}{(a + cx^4)^{3/2}} dx = \frac{Ax\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{2} \mid \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}}\Gamma\left(\frac{5}{4}\right)} + \frac{Bx^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{3}{4}, \frac{3}{2} \mid \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}}\Gamma\left(\frac{7}{4}\right)}$$

input `integrate((B*x**2+A)/(c*x**4+a)**(3/2),x)`

output 
$$A*x*\gamma(1/4)*hyper((1/4, 3/2), (5/4, ), c*x**4*exp_polar(I*pi)/a)/(4*a**(3/2)*\gamma(5/4)) + B*x**3*\gamma(3/4)*hyper((3/4, 3/2), (7/4, ), c*x**4*exp_polar(I*pi)/a)/(4*a**(3/2)*\gamma(7/4))$$

---

3.11. 
$$\int \frac{A+Bx^2}{(a+cx^4)^{3/2}} dx$$

**3.11.7 Maxima [F]**

$$\int \frac{A + Bx^2}{(a + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + a)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(c*x^4+a)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(c*x^4 + a)^(3/2), x)`

**3.11.8 Giac [F]**

$$\int \frac{A + Bx^2}{(a + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + a)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(c*x^4+a)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(c*x^4 + a)^(3/2), x)`

**3.11.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{(a + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + a)^{3/2}} dx$$

input `int((A + B*x^2)/(a + c*x^4)^(3/2),x)`

output `int((A + B*x^2)/(a + c*x^4)^(3/2), x)`

### 3.12 $\int \frac{A+Bx^2}{(d+ex^2)(a+cx^4)^{3/2}} dx$

|        |                            |     |
|--------|----------------------------|-----|
| 3.12.1 | Optimal result             | 119 |
| 3.12.2 | Mathematica [C] (verified) | 120 |
| 3.12.3 | Rubi [A] (verified)        | 121 |
| 3.12.4 | Maple [C] (verified)       | 123 |
| 3.12.5 | Fricas [F(-1)]             | 124 |
| 3.12.6 | Sympy [F]                  | 124 |
| 3.12.7 | Maxima [F]                 | 125 |
| 3.12.8 | Giac [F]                   | 125 |
| 3.12.9 | Mupad [F(-1)]              | 125 |

#### 3.12.1 Optimal result

Integrand size = 28, antiderivative size = 732

$$\int \frac{A+Bx^2}{(d+ex^2)(a+cx^4)^{3/2}} dx = \frac{x(Acd+aBe+c(Bd-Ae)x^2)}{2a(cd^2+ae^2)\sqrt{a+cx^4}} - \frac{\sqrt{c}(Bd-Ae)x\sqrt{a+cx^4}}{2a(cd^2+ae^2)(\sqrt{a}+\sqrt{cx^2})} - \frac{e^{3/2}(Bd-Ae)\arctan\left(\frac{\sqrt{cd^2+ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{a+cx^4}}\right)}{2\sqrt{d}(cd^2+ae^2)^{3/2}} + \frac{\sqrt[4]{c}(Bd-Ae)(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)}{2a^{3/4}(cd^2+ae^2)\sqrt{a+cx^4}} - \frac{\sqrt[4]{ce}(Bd-Ae)(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right),\frac{1}{2}\right)}{2\sqrt[4]{a}(\sqrt{cd}-\sqrt{ae})(cd^2+ae^2)\sqrt{a+cx^4}} + \frac{(Acd+aBe-\sqrt{a}\sqrt{c}(Bd-Ae))(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right),\frac{1}{2}\right)}{4a^{5/4}\sqrt[4]{c}(cd^2+ae^2)\sqrt{a+cx^4}} + \frac{a^{3/4}e\left(\frac{\sqrt{cd}}{\sqrt{a}}+e\right)^2(Bd-Ae)(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}\text{EllipticPi}\left(-\frac{(\sqrt{cd}-\sqrt{ae})^2}{4\sqrt{a}\sqrt{cde}},2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right),\frac{1}{2}\right)}{4\sqrt[4]{cd}(c^2d^4-a^2e^4)\sqrt{a+cx^4}}$$



output

```

-1/2*e^(3/2)*(-A*e+B*d)*arctan(x*(a*e^2+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^
4+a)^(1/2))/(a*e^2+c*d^2)^(3/2)/d^(1/2)+1/2*x*(A*c*d+B*a*e+c*(-A*e+B*d)*x^
2)/a/(a*e^2+c*d^2)/(c*x^4+a)^(1/2)-1/2*(-A*e+B*d)*x*c^(1/2)*(c*x^4+a)^(1/2
)/a/(a*e^2+c*d^2)/(a^(1/2)+x^2*c^(1/2))+1/2*c^(1/4)*(-A*e+B*d)*(cos(2*arct
an(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE
(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x
^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(3/4)/(a*e^2+c*d^2)/(c*x^4+a)^(1/2)
-1/2*c^(1/4)*e*(-A*e+B*d)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2
*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2
*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/
a^(1/4)/(a*e^2+c*d^2)/(-e*a^(1/2)+d*c^(1/2))/(c*x^4+a)^(1/2)+1/4*a^(3/4)*e
*(-A*e+B*d)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4
)*x/a^(1/4)))*EllipticPi(sin(2*arctan(c^(1/4)*x/a^(1/4))),-1/4*(-e*a^(1/2)
+d*c^(1/2))^2/d/e/a^(1/2)/c^(1/2),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*(e+d*
c^(1/2)/a^(1/2))^2*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(1/4)/d/(-a
^2*e^4+c^2*d^4)/(c*x^4+a)^(1/2)+1/4*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(
1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(
1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*(A*c*d+B*a*e-(-A*e+B*d)*a^(1/2)*
c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(5/4)/c^(1/4)/(a*e^2+
c*d^2)/(c*x^4+a)^(1/2)

```

### 3.12.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.66 (sec) , antiderivative size = 432, normalized size of antiderivative = 0.59

$$\int \frac{A + Bx^2}{(d + ex^2)(a + cx^4)^{3/2}} dx = \frac{A\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}cd^2x + aB\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}dex + B\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}cd^2x^3 - A\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}cdex^3 - \sqrt{a}\sqrt{cd}(Bd -$$

input `Integrate[(A + B*x^2)/((d + e*x^2)*(a + c*x^4)^(3/2)),x]`

output  $(A\sqrt{(I\sqrt{c})/\sqrt{a}}*c*d^2*x + a*B*\sqrt{(I\sqrt{c})/\sqrt{a}}*d*e*x + B*\sqrt{(I\sqrt{c})/\sqrt{a}}*c*d^2*x^3 - A*\sqrt{(I\sqrt{c})/\sqrt{a}}*c*d*e*x^3 - \sqrt{a}*\sqrt{c}*d*(B*d - A*e)*\sqrt{1 + (c*x^4)/a}*\text{EllipticE}[I*\text{ArcSinh}[\sqrt{(I\sqrt{c})/\sqrt{a}}*x], -1] + (\sqrt{a}*B - I*A*\sqrt{c})*d*(\sqrt{c}*d - I*\sqrt{a}*e)*\sqrt{1 + (c*x^4)/a}*\text{EllipticF}[I*\text{ArcSinh}[\sqrt{(I\sqrt{c})/\sqrt{a}}*x], -1] + (2*I)*a*B*d*e*\sqrt{1 + (c*x^4)/a}*\text{EllipticPi}[((-I)*\sqrt{a}*e)/(\sqrt{c}*d), I*\text{ArcSinh}[\sqrt{(I\sqrt{c})/\sqrt{a}}*x], -1] - (2*I)*a*A*e^2*\sqrt{1 + (c*x^4)/a}*\text{EllipticPi}[((-I)*\sqrt{a}*e)/(\sqrt{c}*d), I*\text{ArcSinh}[\sqrt{(I\sqrt{c})/\sqrt{a}}*x], -1])/(2*a*\sqrt{(I\sqrt{c})/\sqrt{a}}*d*(c*d^2 + a*e^2)*\sqrt{a + c*x^4})$

### 3.12.3 Rubi [A] (verified)

Time = 1.03 (sec) , antiderivative size = 732, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {2259, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(a + cx^4)^{3/2} (d + ex^2)} dx$$

↓ 2259

$$\int \left( \frac{e(Ae - Bd)}{\sqrt{a + cx^4} (d + ex^2) (ae^2 + cd^2)} + \frac{aBe + cx^2(Bd - Ae) + Acd}{(a + cx^4)^{3/2} (ae^2 + cd^2)} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right) (-\sqrt{a}\sqrt{c}(Bd - Ae) + aBe + Acd)}{4a^{5/4}\sqrt[4]{c}\sqrt{a+cx^4}(ae^2+cd^2)} + \\
& \frac{\sqrt[4]{c}(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} (Bd - Ae) E\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{2}\right)}{2a^{3/4}\sqrt{a+cx^4}(ae^2+cd^2)} + \\
& \frac{a^{3/4}e(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \left(\frac{\sqrt{cd}}{\sqrt{a}} + e\right)^2 (Bd - Ae) \operatorname{EllipticPi}\left(-\frac{(\sqrt{cd}-\sqrt{ae})^2}{4\sqrt{a}\sqrt{cde}}, 2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{4\sqrt[4]{cd}\sqrt{a+cx^4}(c^2d^4 - a^2e^4)} - \\
& \frac{\sqrt[4]{ce}(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} (Bd - Ae) \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{2\sqrt[4]{a}\sqrt{a+cx^4}(\sqrt{cd} - \sqrt{ae})(ae^2+cd^2)} - \\
& \frac{e^{3/2}(Bd - Ae) \arctan\left(\frac{x\sqrt{ae^2+cd^2}}{\sqrt{d}\sqrt{e}\sqrt{a+cx^4}}\right)}{2\sqrt{d}(ae^2+cd^2)^{3/2}} - \frac{\sqrt{cx}\sqrt{a+cx^4}(Bd - Ae)}{2a(\sqrt{a} + \sqrt{cx^2})(ae^2+cd^2)} + \\
& \frac{x(aBe + cx^2(Bd - Ae) + Acd)}{2a\sqrt{a+cx^4}(ae^2+cd^2)}
\end{aligned}$$

input `Int[(A + B*x^2)/((d + e*x^2)*(a + c*x^4)^(3/2)),x]`

output `(x*(A*c*d + a*B*e + c*(B*d - A*e)*x^2))/(2*a*(c*d^2 + a*e^2)*Sqrt[a + c*x^4]) - (Sqrt[c]*(B*d - A*e)*x*Sqrt[a + c*x^4])/(2*a*(c*d^2 + a*e^2)*(Sqrt[a] + Sqrt[c]*x^2)) - (e^(3/2)*(B*d - A*e)*ArcTan[(Sqrt[c*d^2 + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + c*x^4])])/(2*Sqrt[d]*(c*d^2 + a*e^2)^(3/2)) + (c^(1/4)*(B*d - A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(3/4)*(c*d^2 + a*e^2)*Sqrt[a + c*x^4]) - (c^(1/4)*e*(B*d - A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*(Sqrt[c]*d - Sqrt[a]*e)*(c*d^2 + a*e^2)*Sqrt[a + c*x^4]) + ((A*c*d + a*B*e - Sqrt[a]*Sqrt[c]*(B*d - A*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(4*a^(5/4)*c^(1/4)*(c*d^2 + a*e^2)*Sqrt[a + c*x^4]) + (a^(3/4)*e*((Sqrt[c]*d)/Sqrt[a] + e)^2*(B*d - A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-1/4*(Sqrt[c]*d - Sqrt[a]*e)^2/(Sqrt[a]*Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(4*c^(1/4)*d*(c^2*d^4 - a^2*e^4)*Sqrt[a + c*x^4])`

## 3.12.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2259 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x] && IntegerQ[p + 1/2] && IntegerQ[q]`

## 3.12.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.86 (sec) , antiderivative size = 564, normalized size of antiderivative = 0.77

| method   | result   |
|----------|--|
| default  | $\frac{B \left( \frac{x}{2a\sqrt{(x^4 + \frac{a}{c})c}} + \frac{\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}}{2a\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}} F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) \right) (Ae - Bd)}{e} - \frac{2c \left( \frac{ex^3}{4a(ae^2 + cd^2)} - \frac{dx}{4a(ae^2 + cd^2)} \right)}{\sqrt{(x^4 + \frac{a}{c})c}} + \frac{cd\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}}{2a(ae^2 + cd^2)}$  |
| elliptic | $-\frac{2c \left( \frac{(Ae - Bd)x^3}{4a(ae^2 + cd^2)} - \frac{(Acd + Bae)x}{4a(ae^2 + cd^2)c} \right)}{\sqrt{(x^4 + \frac{a}{c})c}} + \frac{\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}}{2a(ae^2 + cd^2)} \frac{F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) Acd}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}} + \frac{\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}}{2(ae^2 + cd^2)} \frac{F\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) E}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}}$ |

input `int((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(3/2), x, method=_RETURNVERBOSE)`

output  $B/e*(1/2*x/a/((x^4+a/c)*c)^{(1/2)}+1/2/a/(I/a^{(1/2)*c^{(1/2)}})^{(1/2)}*(1-I/a^{(1/2)*c^{(1/2)*x^2}})^{(1/2)}*(1+I/a^{(1/2)*c^{(1/2)*x^2}})^{(1/2)}/(c*x^4+a)^{(1/2)}*EllipticF(x*(I/a^{(1/2)*c^{(1/2)}})^{(1/2)},I))+(A*e-B*d)/e*(-2*c*(1/4/a*e/(a*e^2+c*d^2)*x^3-1/4*d/a/(a*e^2+c*d^2)*x)/((x^4+a/c)*c)^{(1/2)}+1/2*c*d/a/(a*e^2+c*d^2)/(I/a^{(1/2)*c^{(1/2)}})^{(1/2)}*(1-I/a^{(1/2)*c^{(1/2)*x^2}})^{(1/2)}*(1+I/a^{(1/2)*c^{(1/2)*x^2}})^{(1/2)}/(c*x^4+a)^{(1/2)}*EllipticF(x*(I/a^{(1/2)*c^{(1/2)}})^{(1/2)},I)+1/2*I/a^{(1/2)*c^{(1/2)}}*e/(a*e^2+c*d^2)/(I/a^{(1/2)*c^{(1/2)}})^{(1/2)}*(1-I/a^{(1/2)*c^{(1/2)*x^2}})^{(1/2)}*(1+I/a^{(1/2)*c^{(1/2)*x^2}})^{(1/2)}/(c*x^4+a)^{(1/2)}*EllipticE(x*(I/a^{(1/2)*c^{(1/2)}})^{(1/2)},I)+1/(a*e^2+c*d^2)*e^2/d/(I/a^{(1/2)*c^{(1/2)}})^{(1/2)}*(1-I/a^{(1/2)*c^{(1/2)*x^2}})^{(1/2)}*(1+I/a^{(1/2)*c^{(1/2)*x^2}})^{(1/2)}/(c*x^4+a)^{(1/2)}*EllipticPi(x*(I/a^{(1/2)*c^{(1/2)}})^{(1/2)},I*a^{(1/2)}/c^{(1/2)}*e/d,(-I/a^{(1/2)*c^{(1/2)}})^{(1/2)}/(I/a^{(1/2)*c^{(1/2)}})^{(1/2))}$

### 3.12.5 Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)(a + cx^4)^{3/2}} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="fricas")`

output `Timed out`

### 3.12.6 Sympy [F]

$$\int \frac{A + Bx^2}{(d + ex^2)(a + cx^4)^{3/2}} dx = \int \frac{A + Bx^2}{(a + cx^4)^{\frac{3}{2}}(d + ex^2)} dx$$

input `integrate((B*x**2+A)/(e*x**2+d)/(c*x**4+a)**(3/2),x)`

output `Integral((A + B*x**2)/((a + c*x**4)**(3/2)*(d + e*x**2)), x)`

**3.12.7 Maxima [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)(a + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + a)^{\frac{3}{2}}(ex^2 + d)} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(e*x^2 + d)), x)`

**3.12.8 Giac [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)(a + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + a)^{\frac{3}{2}}(ex^2 + d)} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(e*x^2 + d)), x)`

**3.12.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)(a + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + a)^{3/2}(ex^2 + d)} dx$$

input `int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)),x)`

output `int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)), x)`

### 3.13 $\int \frac{A+Bx^2}{(d+ex^2)^2(a+cx^4)^{3/2}} dx$

|        |                            |     |
|--------|----------------------------|-----|
| 3.13.1 | Optimal result             | 126 |
| 3.13.2 | Mathematica [C] (verified) | 127 |
| 3.13.3 | Rubi [A] (verified)        | 127 |
| 3.13.4 | Maple [C] (verified)       | 129 |
| 3.13.5 | Fricas [F(-1)]             | 130 |
| 3.13.6 | Sympy [F]                  | 131 |
| 3.13.7 | Maxima [F]                 | 131 |
| 3.13.8 | Giac [F]                   | 131 |
| 3.13.9 | Mupad [F(-1)]              | 132 |

#### 3.13.1 Optimal result

Integrand size = 28, antiderivative size = 1494

$$\int \frac{A+Bx^2}{(d+ex^2)^2(a+cx^4)^{3/2}} dx = \text{Too large to display}$$

output

```
-1/4*e^(3/2)*(-A*e+B*d)*(a*e^2+3*c*d^2)*arctan(x*(a*e^2+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+a)^(1/2))/d^(3/2)/(a*e^2+c*d^2)^(5/2)-1/2*e^(3/2)*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*arctan(x*(a*e^2+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+a)^(1/2))/(a*e^2+c*d^2)^(5/2)/d^(1/2)+1/2*c*x*(A*c*d^2+2*a*B*d*e-a*A*e^2+(-2*A*c*d*e-B*a*e^2+B*c*d^2)*x^2)/a/(a*e^2+c*d^2)^2/(c*x^4+a)^(1/2)-1/2*e^3*(-A*e+B*d)*x*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)^2/(e*x^2+d)+1/2*e^2*(-A*e+B*d)*x*c^(1/2)*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)^2/(a^(1/2)+x^2*c^(1/2))-1/2*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*x*c^(1/2)*(c*x^4+a)^(1/2)/a/(a*e^2+c*d^2)^2/(a^(1/2)+x^2*c^(1/2))-1/2*a^(1/4)*c^(1/4)*e^2*(-A*e+B*d)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/d/(a*e^2+c*d^2)^2/(c*x^4+a)^(1/2)+1/2*c^(1/4)*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/a^(3/4)/(a*e^2+c*d^2)^2/(c*x^4+a)^(1/2)-1/2*c^(1/4)*e*(-A*e+B*d)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/a^(1/4)/d/(a*e^2+c*d^2)/(-e*a^(1/2)+d*c^(1/2))/(c*x^4+a)^(1/2)-1/2*c^(1/4)*e*(-2*A*c*d*e-B*a*e^2+B*c*...
```

### 3.13.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 11.14 (sec) , antiderivative size = 427, normalized size of antiderivative = 0.29

$$\int \frac{A + Bx^2}{(d + ex^2)^2 (a + cx^4)^{3/2}} dx = \frac{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} d (ae^3 (-Bd + Ae)x(a + cx^4) + cdx(d + ex^2) (-aAe^2 + Bcd^2x^2 + Acd^2))}{(d + ex^2)^2 (a + cx^4)^{3/2}}$$

input `Integrate[(A + B*x^2)/((d + e*x^2)^2*(a + c*x^4)^(3/2)),x]`

output `(Sqrt[(I*Sqrt[c])/Sqrt[a]]*d*(a*e^3*(-(B*d) + A*e)*x*(a + c*x^4) + c*d*x*(d + e*x^2)*(-(a*A*e^2) + B*c*d^2*x^2 + A*c*d*(d - 2*e*x^2) + a*B*e*(2*d - e*x^2))) - (d + e*x^2)*Sqrt[1 + (c*x^4)/a]*(-(Sqrt[a]*Sqrt[c]*d*(-(B*c*d^3) + 2*A*c*d^2*e + 2*a*B*d*e^2 - a*A*e^3)*EllipticE[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1]) + I*(Sqrt[c]*d*(Sqrt[c]*d - I*Sqrt[a]*e)*(A*c*d^2 + I*Sqrt[a]*Sqrt[c]*d*(B*d - A*e) + a*e*(2*B*d - A*e))*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + a*e*(-5*B*c*d^3 + 7*A*c*d^2*e + a*B*d*e^2 + a*A*e^3)*EllipticPi[(-I)*Sqrt[a]*e/(Sqrt[c]*d), I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1])))/(2*a*Sqrt[(I*Sqrt[c])/Sqrt[a]]*(c*d^3 + a*d*e^2)^2*(d + e*x^2)*Sqrt[a + c*x^4])`

### 3.13.3 Rubi [A] (verified)

Time = 2.22 (sec) , antiderivative size = 1494, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {2259, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(a + cx^4)^{3/2} (d + ex^2)^2} dx$$

↓ 2259

$$\int \left( \frac{e(Ae - Bd)}{\sqrt{a + cx^4} (d + ex^2)^2 (ae^2 + cd^2)} + \frac{e(aBe^2 + 2Acde - Bcd^2)}{\sqrt{a + cx^4} (d + ex^2) (ae^2 + cd^2)^2} + \frac{c(x^2(-aBe^2 - 2Acde + Bcd^2) - aAe^2)}{(a + cx^4)^{3/2} (ae^2 + cd^2)} \right) dx$$

↓ 2009

---

3.13.  $\int \frac{A+Bx^2}{(d+ex^2)^2(a+cx^4)^{3/2}} dx$



$$\begin{aligned}
& \frac{(Bd - Ae)x\sqrt{cx^4 + ae^3}}{2d(cd^2 + ae^2)^2(e^2x^2 + d)} - \\
& \frac{\sqrt[4]{a}\sqrt[4]{c}(Bd - Ae)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4+a}{(\sqrt{cx^2+\sqrt{a}})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)e^2}{2d(cd^2 + ae^2)^2\sqrt{cx^4 + a}} + \\
& \frac{\sqrt{c}(Bd - Ae)x\sqrt{cx^4 + ae^2}}{2d(cd^2 + ae^2)^2(\sqrt{cx^2 + \sqrt{a}})} - \frac{(Bd - Ae)(3cd^2 + ae^2)\arctan\left(\frac{\sqrt{cd^2+ae^2x}}{\sqrt{d}\sqrt{e}\sqrt{cx^4+a}}\right)e^{3/2}}{4d^{3/2}(cd^2 + ae^2)^{5/2}} - \\
& \frac{(Bcd^2 - 2Aced - aBe^2)\arctan\left(\frac{\sqrt{cd^2+ae^2x}}{\sqrt{d}\sqrt{e}\sqrt{cx^4+a}}\right)e^{3/2}}{2\sqrt{d}(cd^2 + ae^2)^{5/2}} - \\
& \frac{\sqrt[4]{c}(Bcd^2 - 2Aced - aBe^2)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4+a}{(\sqrt{cx^2+\sqrt{a}})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)e}{2^4\sqrt{a}(\sqrt{cd} - \sqrt{ae})(cd^2 + ae^2)^2\sqrt{cx^4 + a}} - \\
& \frac{\sqrt[4]{c}(Bd - Ae)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4+a}{(\sqrt{cx^2+\sqrt{a}})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)e}{2^4\sqrt{ad}(\sqrt{cd} - \sqrt{ae})(cd^2 + ae^2)\sqrt{cx^4 + a}} + \\
& \frac{(\sqrt{cd} + \sqrt{ae})(Bd - Ae)(3cd^2 + ae^2)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4+a}{(\sqrt{cx^2+\sqrt{a}})^2}}\text{EllipticPi}\left(-\frac{(\sqrt{cd}-\sqrt{ae})^2}{4\sqrt{a}\sqrt{cde}}, 2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{8^4\sqrt{a}\sqrt[4]{cd^2}(\sqrt{cd} - \sqrt{ae})(cd^2 + ae^2)^2\sqrt{cx^4 + a}} \\
& \frac{a^{3/4}\left(\frac{\sqrt{cd}}{\sqrt{a}} + e\right)^2(Bcd^2 - 2Aced - aBe^2)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4+a}{(\sqrt{cx^2+\sqrt{a}})^2}}\text{EllipticPi}\left(-\frac{(\sqrt{cd}-\sqrt{ae})^2}{4\sqrt{a}\sqrt{cde}}, 2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{4^4\sqrt{cd}(cd^2 - ae^2)(cd^2 + ae^2)^2\sqrt{cx^4 + a}} \\
& \frac{\sqrt[4]{c}(Bcd^2 - 2Aced - aBe^2)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4+a}{(\sqrt{cx^2+\sqrt{a}})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)}{2a^{3/4}(cd^2 + ae^2)^2\sqrt{cx^4 + a}} - \\
& \frac{\sqrt[4]{c}\left(Bcd^2 - 2Aced - aBe^2 - \frac{\sqrt{c}(Acd^2+2aBed-aAe^2)}{\sqrt{a}}\right)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4+a}{(\sqrt{cx^2+\sqrt{a}})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)}{4a^{3/4}(cd^2 + ae^2)^2\sqrt{cx^4 + a}} \\
& \frac{\sqrt{c}(Bcd^2 - 2Aced - aBe^2)x\sqrt{cx^4 + a}}{2a(cd^2 + ae^2)^2(\sqrt{cx^2 + \sqrt{a}})} + \frac{cx(Acd^2 + 2aBed - aAe^2 + (Bcd^2 - 2Aced - aBe^2)x^2)}{2a(cd^2 + ae^2)^2\sqrt{cx^4 + a}}
\end{aligned}$$

input `Int[(A + B*x^2)/((d + e*x^2)^2*(a + c*x^4)^(3/2)),x]`

```

output (c*x*(A*c*d^2 + 2*a*B*d*e - a*A*e^2 + (B*c*d^2 - 2*A*c*d*e - a*B*e^2)*x^2)
)/(2*a*(c*d^2 + a*e^2)^2*Sqrt[a + c*x^4]) + (Sqrt[c]*e^2*(B*d - A*e)*x*Sqr
t[a + c*x^4])/(2*d*(c*d^2 + a*e^2)^2*(Sqrt[a] + Sqrt[c]*x^2)) - (Sqrt[c]*(
B*c*d^2 - 2*A*c*d*e - a*B*e^2)*x*Sqrt[a + c*x^4])/(2*a*(c*d^2 + a*e^2)^2*(
Sqrt[a] + Sqrt[c]*x^2)) - (e^3*(B*d - A*e)*x*Sqrt[a + c*x^4])/(2*d*(c*d^2
+ a*e^2)^2*(d + e*x^2)) - (e^(3/2)*(B*d - A*e)*(3*c*d^2 + a*e^2)*ArcTan[(S
qrt[c*d^2 + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + c*x^4])])/(4*d^(3/2)*(c*d^
2 + a*e^2)^(5/2)) - (e^(3/2)*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*ArcTan[(Sqrt[
c*d^2 + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + c*x^4])])/(2*Sqrt[d]*(c*d^2 +
a*e^2)^(5/2)) - (a^(1/4)*c^(1/4)*e^2*(B*d - A*e)*(Sqrt[a] + Sqrt[c]*x^2)*S
qrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/
a^(1/4)], 1/2])/(2*d*(c*d^2 + a*e^2)^2*Sqrt[a + c*x^4]) + (c^(1/4)*(B*c*d^
2 - 2*A*c*d*e - a*B*e^2)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a]
+ Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(3/
4)*(c*d^2 + a*e^2)^2*Sqrt[a + c*x^4]) - (c^(1/4)*e*(B*d - A*e)*(Sqrt[a] +
Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTa
n[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*d*(Sqrt[c]*d - Sqrt[a]*e)*(c*d^2
+ a*e^2)*Sqrt[a + c*x^4]) - (c^(1/4)*e*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*(Sq
rt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF
[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*(Sqrt[c]*d - Sqrt[a]*e...

```

### 3.13.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2259 Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol]
:= Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p
+ 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x] && IntegerQ[p + 1/
2] && IntegerQ[q]
```

### 3.13.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.50 (sec) , antiderivative size = 1384, normalized size of antiderivative = 0.93

| method   | result                          | size |
|----------|---------------------------------|------|
| default  | Expression too large to display | 1384 |
| elliptic | Expression too large to display | 1664 |

$$3.13. \int \frac{A+Bx^2}{(d+ex^2)^2(a+cx^4)^{3/2}} dx$$

input `int((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(3/2),x,method=_RETURNVERBOSE)`

output `B/e*(-2*c*(1/4/a*e/(a*e^2+c*d^2)*x^3-1/4*d/a/(a*e^2+c*d^2)*x)/((x^4+a/c)*c)^(1/2)+1/2*c*d/a/(a*e^2+c*d^2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+1/2*I/a^(1/2)*c^(1/2)*e/(a*e^2+c*d^2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)-1/2*I/a^(1/2)*c^(1/2)*e/(a*e^2+c*d^2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+1/(a*e^2+c*d^2)*e^2/d/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticPi(x*(I/a^(1/2)*c^(1/2))^(1/2),I*a^(1/2)/c^(1/2)*e/d,(-I/a^(1/2)*c^(1/2))^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2))+A*e-B*d)/e*(-2*c*(1/2/a*d*c*e/(a*e^2+c*d^2)^2*x^3+1/4/a*(a*e^2-c*d^2)/(a*e^2+c*d^2)^2*x)/((x^4+a/c)*c)^(1/2)+1/2*e^4/(a*e^2+c*d^2)^2/d*x*(c*x^4+a)^(1/2)/(e*x^2+d)-1/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)*e^2*c/(a*e^2+c*d^2)^2+1/2/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)*c^2/a/(a*e^2+c*d^2)^2*d^2+I/a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)...`

### 3.13.5 Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)^2 (a + cx^4)^{3/2}} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="fricas")`

output `Timed out`

**3.13.6 Sympy [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)^2 (a + cx^4)^{3/2}} dx = \int \frac{A + Bx^2}{(a + cx^4)^{\frac{3}{2}} (d + ex^2)^2} dx$$

input `integrate((B*x**2+A)/(e*x**2+d)**2/(c*x**4+a)**(3/2),x)`

output `Integral((A + B*x**2)/((a + c*x**4)**(3/2)*(d + e*x**2)**2), x)`

**3.13.7 Maxima [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)^2 (a + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + a)^{\frac{3}{2}} (ex^2 + d)^2} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(e*x^2 + d)^2), x)`

**3.13.8 Giac [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)^2 (a + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + a)^{\frac{3}{2}} (ex^2 + d)^2} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(e*x^2 + d)^2), x)`

**3.13.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)^2 (a + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + a)^{3/2} (ex^2 + d)^2} dx$$

input `int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)^2),x)`output `int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)^2), x)`

$$3.14 \quad \int \frac{A+Bx^2}{(d+ex^2)^3(a+cx^4)^{3/2}} dx$$

|        |                            |     |
|--------|----------------------------|-----|
| 3.14.1 | Optimal result             | 133 |
| 3.14.2 | Mathematica [C] (verified) | 134 |
| 3.14.3 | Rubi [A] (verified)        | 134 |
| 3.14.4 | Maple [C] (verified)       | 138 |
| 3.14.5 | Fricas [F(-1)]             | 139 |
| 3.14.6 | Sympy [F(-1)]              | 139 |
| 3.14.7 | Maxima [F]                 | 139 |
| 3.14.8 | Giac [F]                   | 140 |
| 3.14.9 | Mupad [F(-1)]              | 140 |

### 3.14.1 Optimal result

Integrand size = 28, antiderivative size = 2452

$$\int \frac{A+Bx^2}{(d+ex^2)^3(a+cx^4)^{3/2}} dx = \text{Too large to display}$$

output

```
-1/4*e^(3/2)*(a*e^2+3*c*d^2)*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*arctan(x*(a*e^2+
c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+a)^(1/2))/d^(3/2)/(a*e^2+c*d^2)^(7/2)-
3/16*e^(3/2)*(-A*e+B*d)*(a^2*e^4+2*a*c*d^2*e^2+5*c^2*d^4)*arctan(x*(a*e^2+
c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+a)^(1/2))/d^(5/2)/(a*e^2+c*d^2)^(7/2)-
1/2*c*e^(3/2)*(A*a*e^3-3*A*c*d^2*e-3*B*a*d*e^2+B*c*d^3)*arctan(x*(a*e^2+c*
d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+a)^(1/2))/(a*e^2+c*d^2)^(7/2)/d^(1/2)+1/
2*c*x*(A*c*d*(-3*a*e^2+c*d^2)+a*B*e*(-a*e^2+3*c*d^2)+c*(A*a*e^3-3*A*c*d^2*
e-3*B*a*d*e^2+B*c*d^3)*x^2)/a/(a*e^2+c*d^2)^3/(c*x^4+a)^(1/2)-1/4*e^3*(-A*
e+B*d)*x*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)^2/(e*x^2+d)^2-3/8*e^3*(-A*e+B*d)*
(a*e^2+3*c*d^2)*x*(c*x^4+a)^(1/2)/d^2/(a*e^2+c*d^2)^3/(e*x^2+d)-1/2*e^3*(-
2*A*c*d*e-B*a*e^2+B*c*d^2)*x*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)^3/(e*x^2+d)-1
/2*c^(3/2)*(A*a*e^3-3*A*c*d^2*e-3*B*a*d*e^2+B*c*d^3)*x*(c*x^4+a)^(1/2)/a/(
a*e^2+c*d^2)^3/(a^(1/2)+x^2*c^(1/2))+3/8*e^2*(-A*e+B*d)*(a*e^2+3*c*d^2)*x*
c^(1/2)*(c*x^4+a)^(1/2)/d^2/(a*e^2+c*d^2)^3/(a^(1/2)+x^2*c^(1/2))+1/2*e^2*
(-2*A*c*d*e-B*a*e^2+B*c*d^2)*x*c^(1/2)*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)^3/(
a^(1/2)+x^2*c^(1/2))-3/8*a^(1/4)*c^(1/4)*e^2*(-A*e+B*d)*(a*e^2+3*c*d^2)*(c
os(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*
EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/
2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/d^2/(a*e^2+c*d^2)^3/(c*x^4+a
)^(1/2)-1/2*a^(1/4)*c^(1/4)*e^2*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*(cos(2*arc...
```

### 3.14.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 11.74 (sec) , antiderivative size = 630, normalized size of antiderivative = 0.26

$$\int \frac{A + Bx^2}{(d + ex^2)^3 (a + cx^4)^{3/2}} dx = \frac{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} dx \left( -2ade^3(Bd - Ae)(cd^2 + ae^2)(a + cx^4) + ae^3(-13Bcd^3 + 17Acd^2) \right)}{\dots}$$

input `Integrate[(A + B*x^2)/((d + e*x^2)^3*(a + c*x^4)^(3/2)),x]`

output `(Sqrt[(I*Sqrt[c])/Sqrt[a]]*d*x*(-2*a*d*e^3*(B*d - A*e)*(c*d^2 + a*e^2)*(a + c*x^4) + a*e^3*(-13*B*c*d^3 + 17*A*c*d^2*e + a*B*d*e^2 + 3*a*A*e^3)*(d + e*x^2)*(a + c*x^4) + 4*c*d^2*(d + e*x^2)^2*(B*(-a^2*e^3) + c^2*d^3*x^2 + 3*a*c*d*e*(d - e*x^2)) + A*c*(c*d^2*(d - 3*e*x^2) + a*e^2*(-3*d + e*x^2))) - (d + e*x^2)^2*Sqrt[1 + (c*x^4)/a]*(Sqrt[a]*Sqrt[c]*d*(3*A*e*(-4*c^2*d^4 + 7*a*c*d^2*e^2 + a^2*e^4) + B*(4*c^2*d^5 - 25*a*c*d^3*e^2 + a^2*d*e^4))*EllipticE[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + I*(Sqrt[c]*d*(Sqrt[c]*d - I*Sqrt[a]*e)*(4*A*c^2*d^4 + (4*I)*Sqrt[a]*c^(3/2)*d^3*(B*d - 2*A*e) + 19*a*c*d^2*e*(B*d - A*e) - (2*I)*a^(3/2)*Sqrt[c]*d*e^2*(3*B*d - A*e) - a^2*e^3*(B*d + 3*A*e))*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + a*e*(3*A*e*(21*c^2*d^4 + 2*a*c*d^2*e^2 + a^2*e^4) + B*(-35*c^2*d^5 + 26*a*c*d^3*e^2 + a^2*d*e^4))*EllipticPi[((-I)*Sqrt[a]*e)/(Sqrt[c]*d), I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1]))/(8*a*Sqrt[(I*Sqrt[c])/Sqrt[a]])*(c*d^3 + a*d*e^2)^3*(d + e*x^2)^2*Sqrt[a + c*x^4])`

### 3.14.3 Rubi [A] (verified)

Time = 3.99 (sec) , antiderivative size = 2452, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {2259, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(a + cx^4)^{3/2} (d + ex^2)^3} dx$$

↓ 2259

---

3.14.  $\int \frac{A+Bx^2}{(d+ex^2)^3(a+cx^4)^{3/2}} dx$

$$\int \left( \frac{e(Ae - Bd)}{\sqrt{a + cx^4} (d + ex^2)^3 (ae^2 + cd^2)} + \frac{e(aBe^2 + 2Acde - Bcd^2)}{\sqrt{a + cx^4} (d + ex^2)^2 (ae^2 + cd^2)^2} + \frac{ce(-aAe^3 + 3aBde^2 + 3Acd^2e - Bcd^3)}{\sqrt{a + cx^4} (d + ex^2) (ae^2 + cd^2)^3} \right) dx$$

↓ 2009



$$\begin{aligned}
& - \frac{3(Bd - Ae)(3cd^2 + ae^2)x\sqrt{cx^4 + ae^3}}{8d^2(cd^2 + ae^2)^3(ex^2 + d)} - \frac{(Bcd^2 - 2Aced - aBe^2)x\sqrt{cx^4 + ae^3}}{2d(cd^2 + ae^2)^3(ex^2 + d)} - \\
& \frac{(Bd - Ae)x\sqrt{cx^4 + ae^3}}{4d(cd^2 + ae^2)^2(ex^2 + d)^2} - \\
& \frac{3\sqrt[4]{a}\sqrt[4]{c}(Bd - Ae)(3cd^2 + ae^2)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)e^2}{8d^2(cd^2 + ae^2)^3\sqrt{cx^4 + a}} - \\
& \frac{\sqrt[4]{a}\sqrt[4]{c}(Bcd^2 - 2Aced - aBe^2)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)e^2}{2d(cd^2 + ae^2)^3\sqrt{cx^4 + a}} + \\
& \frac{3\sqrt{c}(Bd - Ae)(3cd^2 + ae^2)x\sqrt{cx^4 + ae^2}}{8d^2(cd^2 + ae^2)^3(\sqrt{cx^2 + \sqrt{a}})} + \frac{\sqrt{c}(Bcd^2 - 2Aced - aBe^2)x\sqrt{cx^4 + ae^2}}{2d(cd^2 + ae^2)^3(\sqrt{cx^2 + \sqrt{a}})} - \\
& \frac{(3cd^2 + ae^2)(Bcd^2 - 2Aced - aBe^2)\arctan\left(\frac{\sqrt{cd^2 + ae^2x}}{\sqrt{d}\sqrt{e}\sqrt{cx^4 + a}}\right)e^{3/2}}{4d^{3/2}(cd^2 + ae^2)^{7/2}} - \\
& \frac{c(Bcd^3 - 3Aced^2 - 3aBe^2d + aAe^3)\arctan\left(\frac{\sqrt{cd^2 + ae^2x}}{\sqrt{d}\sqrt{e}\sqrt{cx^4 + a}}\right)e^{3/2}}{2\sqrt{d}(cd^2 + ae^2)^{7/2}} - \\
& \frac{3(Bd - Ae)(5c^2d^4 + 2ace^2d^2 + a^2e^4)\arctan\left(\frac{\sqrt{cd^2 + ae^2x}}{\sqrt{d}\sqrt{e}\sqrt{cx^4 + a}}\right)e^{3/2}}{16d^{5/2}(cd^2 + ae^2)^{7/2}} - \\
& \frac{\sqrt[4]{c}(Bd - Ae)(4cd^2 - \sqrt{a}\sqrt{ced} + 3ae^2)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)e}{8\sqrt[4]{ad^2}(\sqrt{cd} - \sqrt{ae})(cd^2 + ae^2)^2\sqrt{cx^4 + a}} - \\
& \frac{\sqrt[4]{c}(Bcd^2 - 2Aced - aBe^2)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)e}{2\sqrt[4]{ad}(\sqrt{cd} - \sqrt{ae})(cd^2 + ae^2)^2\sqrt{cx^4 + a}} - \\
& \frac{c^{5/4}(Bcd^3 - 3Aced^2 - 3aBe^2d + aAe^3)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{2}\right)e}{2\sqrt[4]{a}(\sqrt{cd} - \sqrt{ae})(cd^2 + ae^2)^3\sqrt{cx^4 + a}} + \\
& \frac{(\sqrt{cd} + \sqrt{ae})(3cd^2 + ae^2)(Bcd^2 - 2Aced - aBe^2)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}\text{EllipticPi}\left(-\frac{(\sqrt{cd} - \sqrt{ae})^2}{4\sqrt{a}\sqrt{cde}}, 2\arctan\right)}{8\sqrt[4]{a}\sqrt[4]{cd^2}(\sqrt{cd} - \sqrt{ae})(cd^2 + ae^2)^3\sqrt{cx^4 + a}} - \\
& \frac{a^{3/4}c^{3/4}\left(\frac{\sqrt{cd}}{\sqrt{a}} + e\right)^2(Bcd^3 - 3Aced^2 - 3aBe^2d + aAe^3)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}\text{EllipticPi}\left(-\frac{(\sqrt{cd} - \sqrt{ae})^2}{4\sqrt{a}\sqrt{cde}}, 2\arctan\right)}{4d(cd^2 - ae^2)(cd^2 + ae^2)^3\sqrt{cx^4 + a}} - \\
& \frac{3(\sqrt{cd} + \sqrt{ae})(Bd - Ae)(5c^2d^4 + 2ace^2d^2 + a^2e^4)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}\text{EllipticPi}\left(-\frac{(\sqrt{cd} - \sqrt{ae})^2}{4\sqrt{a}\sqrt{cde}}, 2\arctan\right)}{32\sqrt[4]{a}\sqrt[4]{cd^3}(\sqrt{cd} - \sqrt{ae})(cd^2 + ae^2)^3\sqrt{cx^4 + a}} - \\
& \frac{c^{5/4}(Bcd^3 - 3Aced^2 - 3aBe^2d + aAe^3)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)}{2a^{3/4}(cd^2 + ae^2)^3\sqrt{cx^4 + a}} + \\
& \frac{c^{3/4}(Ac^2d^3 - \sqrt{ac}^{3/2}(Bd - 3Ae)d^2 + 3ace(Bd - Ae)d - a^2Be^3 + a^{3/2}\sqrt{ce^2}(3Bd - Ae))(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}}{4a^{5/4}(cd^2 + ae^2)^3\sqrt{cx^4 + a}}
\end{aligned}$$

3.14.  $\int \frac{A+Bx^2}{(d+ex^2)^3(a+cx^4)^{3/2}} \frac{c^{3/2}(Bcd^3 - 3Aced^2 - 3aBe^2d + aAe^3)x\sqrt{cx^4 + a}}{2a(cd^2 + ae^2)^3(\sqrt{cx^2 + \sqrt{a}})} dx +$   
 $cx(c(Bcd^3 - 3Aced^2 - 3aBe^2d + aAe^3)x^2 + Acd(cd^2 - 3ae^2) + aBe(3cd^2 - ae^2))$

input `Int[(A + B*x^2)/((d + e*x^2)^3*(a + c*x^4)^(3/2)),x]`

output `(c*x*(A*c*d*(c*d^2 - 3*a*e^2) + a*B*e*(3*c*d^2 - a*e^2) + c*(B*c*d^3 - 3*A*c*d^2*e - 3*a*B*d*e^2 + a*A*e^3)*x^2))/(2*a*(c*d^2 + a*e^2)^3*sqrt[a + c*x^4]) + (3*sqrt[c]*e^2*(B*d - A*e)*(3*c*d^2 + a*e^2)*x*sqrt[a + c*x^4])/(8*d^2*(c*d^2 + a*e^2)^3*(sqrt[a] + sqrt[c]*x^2)) + (sqrt[c]*e^2*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*x*sqrt[a + c*x^4])/(2*d*(c*d^2 + a*e^2)^3*(sqrt[a] + sqrt[c]*x^2)) - (c^(3/2)*(B*c*d^3 - 3*A*c*d^2*e - 3*a*B*d*e^2 + a*A*e^3)*x*sqrt[a + c*x^4])/(2*a*(c*d^2 + a*e^2)^3*(sqrt[a] + sqrt[c]*x^2)) - (e^3*(B*d - A*e)*x*sqrt[a + c*x^4])/(4*d*(c*d^2 + a*e^2)^2*(d + e*x^2)^2) - (3*e^3*(B*d - A*e)*(3*c*d^2 + a*e^2)*x*sqrt[a + c*x^4])/(8*d^2*(c*d^2 + a*e^2)^3*(d + e*x^2)) - (e^3*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*x*sqrt[a + c*x^4])/(2*d*(c*d^2 + a*e^2)^3*(d + e*x^2)) - (e^(3/2)*(3*c*d^2 + a*e^2)*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*ArcTan[(sqrt[c*d^2 + a*e^2]*x)/(sqrt[d]*sqrt[e]*sqrt[a + c*x^4])])/(4*d^(3/2)*(c*d^2 + a*e^2)^(7/2)) - (c*e^(3/2)*(B*c*d^3 - 3*A*c*d^2*e - 3*a*B*d*e^2 + a*A*e^3)*ArcTan[(sqrt[c*d^2 + a*e^2]*x)/(sqrt[d]*sqrt[e]*sqrt[a + c*x^4])])/(2*sqrt[d]*(c*d^2 + a*e^2)^(7/2)) - (3*e^(3/2)*(B*d - A*e)*(5*c^2*d^4 + 2*a*c*d^2*e^2 + a^2*e^4)*ArcTan[(sqrt[c*d^2 + a*e^2]*x)/(sqrt[d]*sqrt[e]*sqrt[a + c*x^4])])/(16*d^(5/2)*(c*d^2 + a*e^2)^(7/2)) - (3*a^(1/4)*c^(1/4)*e^2*(B*d - A*e)*(3*c*d^2 + a*e^2)*(sqrt[a] + sqrt[c]*x^2)*sqrt[(a + c*x^4)/(sqrt[a] + sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(8*d^2*(c*d^2 + a*e^2)^3*sqrt[a + c*x^4]) ...`

### 3.14.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2259 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x] && IntegerQ[p + 1/2] && IntegerQ[q]`

### 3.14.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 3.43 (sec) , antiderivative size = 2326, normalized size of antiderivative = 0.95

| method   | result                          | size |
|----------|---------------------------------|------|
| default  | Expression too large to display | 2326 |
| elliptic | Expression too large to display | 2596 |

```
input int((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(3/2),x,method=_RETURNVERBOSE)
```

```
output B/e*(-2*c*(1/2/a*d*c*e/(a*e^2+c*d^2)^2*x^3+1/4/a*(a*e^2-c*d^2)/(a*e^2+c*d^2)^2*x)/((x^4+a/c)*c)^(1/2)+1/2*e^4/(a*e^2+c*d^2)^2/d*x*(c*x^4+a)^(1/2)/(e*x^2+d)-1/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)*e^2*c/(a*e^2+c*d^2)^2+1/2/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)*c^2/a/(a*e^2+c*d^2)^2*d^2+I/a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*c^(3/2)*d*e/(a*e^2+c*d^2)^2*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)-I/a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*c^(3/2)*d*e/(a*e^2+c*d^2)^2*EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I)-1/2*I*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*c^(1/2)*e^3/d/(a*e^2+c*d^2)^2*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+1/2*I*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*c^(1/2)*e^3/d/(a*e^2+c*d^2)^2*EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+1/2*e^4/d^2/(a*e^2+c*d^2)^2/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticPi(x*(I/a^(1/2)*c^(1/2))^(1/2),I*a^(1/2)/c^(1/2)*e/d,(-I/a^(1/2)*c^(1/2))^(1/2)...
```

**3.14.5 Fracas [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)^3 (a + cx^4)^{3/2}} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="fracas")`

output `Timed out`

**3.14.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)^3 (a + cx^4)^{3/2}} dx = \text{Timed out}$$

input `integrate((B*x**2+A)/(e*x**2+d)**3/(c*x**4+a)**(3/2),x)`

output `Timed out`

**3.14.7 Maxima [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)^3 (a + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + a)^{3/2} (ex^2 + d)^3} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(e*x^2 + d)^3), x)`

**3.14.8 Giac [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)^3 (a + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + a)^{\frac{3}{2}} (ex^2 + d)^3} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(e*x^2 + d)^3), x)`

**3.14.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)^3 (a + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + a)^{3/2} (ex^2 + d)^3} dx$$

input `int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)^3),x)`

output `int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)^3), x)`

$$3.15 \quad \int \frac{(A+Bx^2)(d+ex^2)^q}{a+cx^4} dx$$

|        |                     |     |
|--------|---------------------|-----|
| 3.15.1 | Optimal result      | 141 |
| 3.15.2 | Mathematica [F]     | 141 |
| 3.15.3 | Rubi [A] (verified) | 142 |
| 3.15.4 | Maple [F]           | 143 |
| 3.15.5 | Fricas [F]          | 143 |
| 3.15.6 | Sympy [F(-1)]       | 143 |
| 3.15.7 | Maxima [F]          | 144 |
| 3.15.8 | Giac [F]            | 144 |
| 3.15.9 | Mupad [F(-1)]       | 144 |

### 3.15.1 Optimal result

Integrand size = 26, antiderivative size = 169

$$\begin{aligned} & \int \frac{(A+Bx^2)(d+ex^2)^q}{a+cx^4} dx \\ &= \frac{\left(A - \frac{\sqrt{-aB}}{\sqrt{c}}\right) x(d+ex^2)^q \left(1 + \frac{ex^2}{d}\right)^{-q} \operatorname{AppellF1}\left(\frac{1}{2}, 1, -q, \frac{3}{2}, -\frac{\sqrt{cx^2}}{\sqrt{-a}}, -\frac{ex^2}{d}\right)}{2a} \\ &+ \frac{\left(A + \frac{\sqrt{-aB}}{\sqrt{c}}\right) x(d+ex^2)^q \left(1 + \frac{ex^2}{d}\right)^{-q} \operatorname{AppellF1}\left(\frac{1}{2}, 1, -q, \frac{3}{2}, \frac{\sqrt{cx^2}}{\sqrt{-a}}, -\frac{ex^2}{d}\right)}{2a} \end{aligned}$$

output  $1/2*x*(e*x^2+d)^q*\operatorname{AppellF1}(1/2, -q, 1, 3/2, -e*x^2/d, -x^2*c^{(1/2)}/(-a)^{(1/2)})*(A-B*(-a)^{(1/2)}/c^{(1/2)})/a/((1+e*x^2/d)^q)+1/2*x*(e*x^2+d)^q*\operatorname{AppellF1}(1/2, 1, -q, 3/2, x^2*c^{(1/2)}/(-a)^{(1/2)}, -e*x^2/d)*(A+B*(-a)^{(1/2)}/c^{(1/2)})/a/((1+e*x^2/d)^q)$

### 3.15.2 Mathematica [F]

$$\int \frac{(A+Bx^2)(d+ex^2)^q}{a+cx^4} dx = \int \frac{(A+Bx^2)(d+ex^2)^q}{a+cx^4} dx$$

input `Integrate[((A + B*x^2)*(d + e*x^2)^q)/(a + c*x^4), x]`

output `Integrate[((A + B*x^2)*(d + e*x^2)^q)/(a + c*x^4), x]`

---


$$3.15. \quad \int \frac{(A+Bx^2)(d+ex^2)^q}{a+cx^4} dx$$

### 3.15.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$ , Rules used = {2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + cx^4} dx$$

↓ 2257

$$\int \left( \frac{(\sqrt{-a}B - A\sqrt{c})(d + ex^2)^q}{2\sqrt{-a}\sqrt{c}(\sqrt{-a} + \sqrt{cx^2})} - \frac{(\sqrt{-a}B + A\sqrt{c})(d + ex^2)^q}{2\sqrt{-a}\sqrt{c}(\sqrt{-a} - \sqrt{cx^2})} \right) dx$$

↓ 2009

$$\frac{x \left( A - \frac{\sqrt{-a}B}{\sqrt{c}} \right) (d + ex^2)^q \left( \frac{ex^2}{d} + 1 \right)^{-q} \text{AppellF1} \left( \frac{1}{2}, 1, -q, \frac{3}{2}, -\frac{\sqrt{cx^2}}{\sqrt{-a}}, -\frac{ex^2}{d} \right)}{2a} +$$

$$\frac{x \left( \frac{\sqrt{-a}B}{\sqrt{c}} + A \right) (d + ex^2)^q \left( \frac{ex^2}{d} + 1 \right)^{-q} \text{AppellF1} \left( \frac{1}{2}, 1, -q, \frac{3}{2}, \frac{\sqrt{cx^2}}{\sqrt{-a}}, -\frac{ex^2}{d} \right)}{2a}$$

input `Int[(A + B*x^2)*(d + e*x^2)^q]/(a + c*x^4),x]`

output `((A - (Sqrt[-a]*B)/Sqrt[c])*x*(d + e*x^2)^q*AppellF1[1/2, 1, -q, 3/2, -(Sqrt[c]*x^2)/Sqrt[-a]], -((e*x^2)/d))/(2*a*(1 + (e*x^2)/d)^q) + ((A + (Sqrt[-a]*B)/Sqrt[c])*x*(d + e*x^2)^q*AppellF1[1/2, 1, -q, 3/2, (Sqrt[c]*x^2)/Sqrt[-a], -((e*x^2)/d)]/(2*a*(1 + (e*x^2)/d)^q)`

#### 3.15.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2257 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`

---

3.15.  $\int \frac{(A+Bx^2)(d+ex^2)^q}{a+cx^4} dx$

**3.15.4 Maple [F]**

$$\int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + a} dx$$

input `int((B*x^2+A)*(e*x^2+d)^q/(c*x^4+a),x)`

output `int((B*x^2+A)*(e*x^2+d)^q/(c*x^4+a),x)`

**3.15.5 Fricas [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + cx^4} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + a} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+a),x, algorithm="fricas")`

output `integral((B*x^2 + A)*(e*x^2 + d)^q/(c*x^4 + a), x)`

**3.15.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + cx^4} dx = \text{Timed out}$$

input `integrate((B*x**2+A)*(e*x**2+d)**q/(c*x**4+a),x)`

output `Timed out`



**3.15.7 Maxima [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + cx^4} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + a} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+a),x, algorithm="maxima")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^q/(c*x^4 + a), x)`

**3.15.8 Giac [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + cx^4} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + a} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+a),x, algorithm="giac")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^q/(c*x^4 + a), x)`

**3.15.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + cx^4} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + a} dx$$

input `int(((A + B*x^2)*(d + e*x^2)^q)/(a + c*x^4),x)`

output `int(((A + B*x^2)*(d + e*x^2)^q)/(a + c*x^4), x)`

### 3.16 $\int \frac{2+x^2}{(1+x^2)\sqrt{2+3x^2+x^4}} dx$

|        |   |     |
|--------|---|-----|
| 3.16.1 | Optimal result                            | 145 |
| 3.16.2 | Mathematica [C] (verified)                | 145 |
| 3.16.3 | Rubi [A] (verified)                       | 146 |
| 3.16.4 | Maple [C] (verified)                      | 147 |
| 3.16.5 | Fricas [C] (verification not implemented) | 147 |
| 3.16.6 | Sympy [F]                                 | 148 |
| 3.16.7 | Maxima [F]                                | 148 |
| 3.16.8 | Giac [F]                                  | 148 |
| 3.16.9 | Mupad [F(-1)]                             | 149 |

#### 3.16.1 Optimal result

Integrand size = 27, antiderivative size = 48

$$\int \frac{2+x^2}{(1+x^2)\sqrt{2+3x^2+x^4}} dx = \frac{\sqrt{2}(2+x^2) E(\arctan(x) \mid \frac{1}{2})}{\sqrt{\frac{2+x^2}{1+x^2}} \sqrt{2+3x^2+x^4}}$$

output  $(x^2+2)*(1/(x^2+1))^{(1/2)}*(x^2+1)^{(1/2)}*EllipticE(x/(x^2+1)^{(1/2)},1/2)*2^{(1/2)}*(1/2))^{(1/2)}/((x^2+2)/(x^2+1))^{(1/2)}/(x^4+3*x^2+2)^{(1/2)}$

#### 3.16.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.17 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.96

$$\int \frac{2+x^2}{(1+x^2)\sqrt{2+3x^2+x^4}} dx = \frac{2x+x^3+i\sqrt{1+x^2}\sqrt{2+x^2}E\left(i\operatorname{arcsinh}\left(\frac{x}{\sqrt{2}}\right)\mid 2\right)-i\sqrt{1+x^2}\sqrt{2+x^2}\operatorname{EllipticF}\left(i\operatorname{arcsinh}\left(\frac{x}{\sqrt{2}}\right),2\right)}{\sqrt{2+3x^2+x^4}}$$

input `Integrate[(2 + x^2)/((1 + x^2)*Sqrt[2 + 3*x^2 + x^4]),x]`

output  $(2*x + x^3 + I*Sqrt[1 + x^2]*Sqrt[2 + x^2]*EllipticE[I*ArcSinh[x/Sqrt[2]]], 2) - I*Sqrt[1 + x^2]*Sqrt[2 + x^2]*EllipticF[I*ArcSinh[x/Sqrt[2]], 2])/Sqrt[2 + 3*x^2 + x^4]$

### 3.16.3 Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.074$ , Rules used = {1395, 313}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2 + 2}{(x^2 + 1)\sqrt{x^4 + 3x^2 + 2}} dx$$

↓ 1395

$$\frac{\sqrt{x^2 + 1}\sqrt{x^2 + 2} \int \frac{\sqrt{x^2 + 2}}{(x^2 + 1)^{3/2}} dx}{\sqrt{x^4 + 3x^2 + 2}}$$

↓ 313

$$\frac{\sqrt{2}(x^2 + 2) E(\arctan(x) | \frac{1}{2})}{\sqrt{\frac{x^2 + 2}{x^2 + 1}} \sqrt{x^4 + 3x^2 + 2}}$$

input `Int[(2 + x^2)/((1 + x^2)*Sqrt[2 + 3*x^2 + x^4]),x]`

output `(Sqrt[2]*(2 + x^2)*EllipticE[ArcTan[x], 1/2])/(Sqrt[(2 + x^2)/(1 + x^2)]*Sqrt[2 + 3*x^2 + x^4])`

#### 3.16.3.1 Defintions of rubi rules used

rule 313 `Int[Sqrt[(a_) + (b_)*(x_)^2]/((c_) + (d_)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[c*((a + b*x^2)/(a*(c + d*x^2))]))*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]`

rule 1395 `Int[(u_)*((a_) + (c_)*(x_)^(n2_)) + (b_)*(x_)^(n_)]^(p_)*((d_) + (e_)*(x_)^(n_)]^(q_), x_Symbol] := Simp[(a + b*x^n + c*x^(2*n))^FracPart[p]/((d + e*x^n)^FracPart[p]*(a/d + c*(x^n/e))^FracPart[p]) Int[u*(d + e*x^n)^(p + q)*(a/d + (c/e)*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, n, p, q}, x] && Eqq[n2, 2*n] && EqQ[c*d^2 - b*d*e + a*e^2, 0] && !IntegerQ[p] && !(EqQ[q, 1] && EqQ[n, 2])`

### 3.16.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.56 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.67

| method   | result   | size |
|----------|--|------|
| risch    | $\frac{x(x^2+2)}{\sqrt{x^4+3x^2+2}} - \frac{i\sqrt{2}\sqrt{2x^2+4}\sqrt{x^2+1}\left(F\left(\frac{i\sqrt{2}x}{2},\sqrt{2}\right)-E\left(\frac{i\sqrt{2}x}{2},\sqrt{2}\right)\right)}{2\sqrt{x^4+3x^2+2}}$     | 80   |
| default  | $\frac{(x^2+2)x}{\sqrt{(x^2+1)(x^2+2)}} - \frac{i\sqrt{2}\sqrt{2x^2+4}\sqrt{x^2+1}\left(F\left(\frac{i\sqrt{2}x}{2},\sqrt{2}\right)-E\left(\frac{i\sqrt{2}x}{2},\sqrt{2}\right)\right)}{2\sqrt{x^4+3x^2+2}}$ | 81   |
| elliptic | $\frac{(x^2+2)x}{\sqrt{(x^2+1)(x^2+2)}} - \frac{i\sqrt{2}\sqrt{2x^2+4}\sqrt{x^2+1}\left(F\left(\frac{i\sqrt{2}x}{2},\sqrt{2}\right)-E\left(\frac{i\sqrt{2}x}{2},\sqrt{2}\right)\right)}{2\sqrt{x^4+3x^2+2}}$ | 81   |

input `int((x^2+2)/(x^2+1)/(x^4+3*x^2+2)^(1/2),x,method=_RETURNVERBOSE)`

output `x*(x^2+2)/(x^4+3*x^2+2)^(1/2)-1/2*I*2^(1/2)*(2*x^2+4)^(1/2)*(x^2+1)^(1/2)/(x^4+3*x^2+2)^(1/2)*(EllipticF(1/2*I*2^(1/2)*x,2^(1/2))-EllipticE(1/2*I*2^(1/2)*x,2^(1/2)))`

### 3.16.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.23

$$\int \frac{2+x^2}{(1+x^2)\sqrt{2+3x^2+x^4}} dx$$

$$= \frac{(-ix^2-i)E(\arcsin(\frac{1}{2}i\sqrt{2}x)|2) + (ix^2+i)F(\arcsin(\frac{1}{2}i\sqrt{2}x)|2) + 2\sqrt{x^4+3x^2+2}x}{2(x^2+1)}$$

input `integrate((x^2+2)/(x^2+1)/(x^4+3*x^2+2)^(1/2),x, algorithm="fracas")`

output `1/2*((-I*x^2 - I)*elliptic_e(arcsin(1/2*I*sqrt(2)*x), 2) + (I*x^2 + I)*elliptic_f(arcsin(1/2*I*sqrt(2)*x), 2) + 2*sqrt(x^4 + 3*x^2 + 2)*x)/(x^2 + 1)`

**3.16.6 Sympy [F]**

$$\int \frac{2+x^2}{(1+x^2)\sqrt{2+3x^2+x^4}} dx = \int \frac{x^2+2}{\sqrt{(x^2+1)(x^2+2)(x^2+1)}} dx$$

input `integrate((x**2+2)/(x**2+1)/(x**4+3*x**2+2)**(1/2),x)`

output `Integral((x**2 + 2)/(sqrt((x**2 + 1)*(x**2 + 2))*(x**2 + 1)), x)`

**3.16.7 Maxima [F]**

$$\int \frac{2+x^2}{(1+x^2)\sqrt{2+3x^2+x^4}} dx = \int \frac{x^2+2}{\sqrt{x^4+3x^2+2}(x^2+1)} dx$$

input `integrate((x^2+2)/(x^2+1)/(x^4+3*x^2+2)^(1/2),x, algorithm="maxima")`

output `integrate((x^2 + 2)/(sqrt(x^4 + 3*x^2 + 2)*(x^2 + 1)), x)`

**3.16.8 Giac [F]**

$$\int \frac{2+x^2}{(1+x^2)\sqrt{2+3x^2+x^4}} dx = \int \frac{x^2+2}{\sqrt{x^4+3x^2+2}(x^2+1)} dx$$

input `integrate((x^2+2)/(x^2+1)/(x^4+3*x^2+2)^(1/2),x, algorithm="giac")`

output `integrate((x^2 + 2)/(sqrt(x^4 + 3*x^2 + 2)*(x^2 + 1)), x)`

**3.16.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{2+x^2}{(1+x^2)\sqrt{2+3x^2+x^4}} dx = \int \frac{x^2+2}{(x^2+1)\sqrt{x^4+3x^2+2}} dx$$

input `int((x^2 + 2)/((x^2 + 1)*(3*x^2 + x^4 + 2)^(1/2)),x)`output `int((x^2 + 2)/((x^2 + 1)*(3*x^2 + x^4 + 2)^(1/2)), x)`

**3.17** 
$$\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+bx^2+cx^4}} dx$$

|        |   |     |
|--------|---|-----|
| 3.17.1 | Optimal result . . . . .                            | 150 |
| 3.17.2 | Mathematica [C] (verified) . . . . .                | 151 |
| 3.17.3 | Rubi [A] (verified) . . . . .                       | 152 |
| 3.17.4 | Maple [A] (verified) . . . . .                      | 155 |
| 3.17.5 | Fricas [A] (verification not implemented) . . . . . | 156 |
| 3.17.6 | Sympy [F] . . . . .                                 | 157 |
| 3.17.7 | Maxima [F] . . . . .                                | 158 |
| 3.17.8 | Giac [F] . . . . .                                  | 158 |
| 3.17.9 | Mupad [F(-1)] . . . . .                             | 158 |

**3.17.1 Optimal result**

Integrand size = 33, antiderivative size = 755

$$\begin{aligned} & \int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+bx^2+cx^4}} dx \\ &= \frac{e(7Ace(15cd-4be)+B(105c^2d^2+24b^2e^2-ce(84bd+25ae)))x\sqrt{a+bx^2+cx^4}}{105c^3} \\ &+ \frac{e^2(21Bcd-6bBe+7Ace)x^3\sqrt{a+bx^2+cx^4}}{35c^2} + \frac{Be^3x^5\sqrt{a+bx^2+cx^4}}{7c} \\ &+ \frac{(7Ace(45c^2d^2+8b^2e^2-3ce(10bd+3ae))+B(105c^3d^3-48b^3e^3-21c^2de(10bd+9ae))+8bce^2(21bd+10bd^2+3ae^2))\sqrt{a+bx^2+cx^4}}{105c^{7/2}(\sqrt{a}+\sqrt{cx^2})} \\ &- \frac{\sqrt[4]{a}(7Ace(45c^2d^2+8b^2e^2-3ce(10bd+3ae))+B(105c^3d^3-48b^3e^3-21c^2de(10bd+9ae))+8bce^2(21bd+10bd^2+3ae^2))}{105c^{15/4}\sqrt{a+bx^2+cx^4}} \\ &+ \frac{\sqrt[4]{a}(7Ace(45c^2d^2+8b^2e^2-3ce(10bd+3ae))+B(105c^3d^3-48b^3e^3-21c^2de(10bd+9ae))+8bce^2(21bd+10bd^2+3ae^2))}{105c^{15/4}\sqrt{a+bx^2+cx^4}} \end{aligned}$$

---

3.17. 
$$\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+bx^2+cx^4}} dx$$

output

```

1/105*e*(7*A*c*e*(-4*b*e+15*c*d)+B*(105*c^2*d^2+24*b^2*e^2-c*e*(25*a*e+84*
b*d)))*x*(c*x^4+b*x^2+a)^(1/2)/c^3+1/35*e^2*(7*A*c*e-6*B*b*e+21*B*c*d)*x^3
*(c*x^4+b*x^2+a)^(1/2)/c^2+1/7*B*e^3*x^5*(c*x^4+b*x^2+a)^(1/2)/c+1/105*(7*
A*c*e*(45*c^2*d^2+8*b^2*e^2-3*c*e*(3*a*e+10*b*d))+B*(105*c^3*d^3-48*b^3*e^
3-21*c^2*d*e*(9*a*e+10*b*d)+8*b*c*e^2*(13*a*e+21*b*d)))*x*(c*x^4+b*x^2+a)^(
1/2)/c^(7/2)/(a^(1/2)+x^2*c^(1/2))-1/105*a^(1/4)*(7*A*c*e*(45*c^2*d^2+8*b
^2*e^2-3*c*e*(3*a*e+10*b*d))+B*(105*c^3*d^3-48*b^3*e^3-21*c^2*d*e*(9*a*e+1
0*b*d)+8*b*c*e^2*(13*a*e+21*b*d)))*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1
/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1
/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2
+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(15/4)/(c*x^4+b*x^2+a)^(1/2)+1/210*a^(
1/4)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(
1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2
)))^(1/2))*(a^(1/2)+x^2*c^(1/2))*(7*A*c*e*(45*c^2*d^2+8*b^2*e^2-3*c*e*(3*a*
e+10*b*d))+B*(105*c^3*d^3-48*b^3*e^3-21*c^2*d*e*(9*a*e+10*b*d)+8*b*c*e^2*(
13*a*e+21*b*d)))+(7*A*c*(4*a*b*e^3-15*a*c*d*e^2+15*c^2*d^3)-a*B*e*(105*c^2*
d^2+24*b^2*e^2-c*e*(25*a*e+84*b*d)))*c^(1/2)/a^(1/2))*((c*x^4+b*x^2+a)/(a^(
1/2)+x^2*c^(1/2)))^(1/2)/c^(15/4)/(c*x^4+b*x^2+a)^(1/2)

```

### 3.17.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 14.51 (sec) , antiderivative size = 901, normalized size of antiderivative = 1.19

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + bx^2 + cx^4}} dx$$

$$= \frac{4c\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}ex(a + bx^2 + cx^4)(7Ace(-4be + 3c(5d + ex^2)) + B(24b^2e^2 - ce(84bd + 25ae + 18bex^2)) + (A + Bx^2)(d + ex^2)^3)}{\sqrt{a + bx^2 + cx^4}}$$

input `Integrate[((A + B*x^2)*(d + e*x^2)^3)/Sqrt[a + b*x^2 + c*x^4], x]`

---

3.17.  $\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+bx^2+cx^4}} dx$



output  $(4*c*\text{Sqrt}[c/(b + \text{Sqrt}[b^2 - 4*a*c])]*e*x*(a + b*x^2 + c*x^4)*(7*A*c*e*(-4*b*e + 3*c*(5*d + e*x^2)) + B*(24*b^2*e^2 - c*e*(84*b*d + 25*a*e + 18*b*e*x^2) + 3*c^2*(35*d^2 + 21*d*e*x^2 + 5*e^2*x^4))) - I*(-b + \text{Sqrt}[b^2 - 4*a*c])*(7*A*c*e*(-45*c^2*d^2 - 8*b^2*e^2 + 3*c*e*(10*b*d + 3*a*e)) + B*(-105*c^3*d^3 + 48*b^3*e^3 + 21*c^2*d*e*(10*b*d + 9*a*e) - 8*b*c*e^2*(21*b*d + 13*a*e)))*\text{Sqrt}[(b + \text{Sqrt}[b^2 - 4*a*c] + 2*c*x^2)/(b + \text{Sqrt}[b^2 - 4*a*c])]*\text{Sqrt}[(2*b - 2*\text{Sqrt}[b^2 - 4*a*c] + 4*c*x^2)/(b - \text{Sqrt}[b^2 - 4*a*c])]*\text{EllipticE}[I*\text{ArcSinh}[\text{Sqrt}[2]*\text{Sqrt}[c/(b + \text{Sqrt}[b^2 - 4*a*c])]]*x], (b + \text{Sqrt}[b^2 - 4*a*c])/(b - \text{Sqrt}[b^2 - 4*a*c])] + I*(-48*b^4*B*e^3 + 8*b^3*e^2*(21*B*c*d + 7*A*c*e + 6*B*\text{Sqrt}[b^2 - 4*a*c]*e) - 2*b^2*c*e*(7*A*e*(15*c*d + 4*\text{Sqrt}[b^2 - 4*a*c]*e) + B*(105*c*d^2 + 84*\text{Sqrt}[b^2 - 4*a*c]*d*e - 76*a*e^2)) + b*c*(B*(105*c^2*d^3 - 104*a*\text{Sqrt}[b^2 - 4*a*c]*e^3 + 21*c*d*e*(10*\text{Sqrt}[b^2 - 4*a*c]*d - 17*a*e)) + 7*A*c*e*(45*c*d^2 + e*(30*\text{Sqrt}[b^2 - 4*a*c]*d - 17*a*e))) + c^2*(B*(a*e^2*(189*\text{Sqrt}[b^2 - 4*a*c]*d - 50*a*e) - 105*c*d^2*(\text{Sqrt}[b^2 - 4*a*c]*d - 2*a*e)) - 21*A*(10*c^2*d^3 - 3*a*\text{Sqrt}[b^2 - 4*a*c]*e^3 + 5*c*d*e*(3*\text{Sqrt}[b^2 - 4*a*c]*d - 2*a*e)))*\text{Sqrt}[(b + \text{Sqrt}[b^2 - 4*a*c] + 2*c*x^2)/(b + \text{Sqrt}[b^2 - 4*a*c])]*\text{Sqrt}[(2*b - 2*\text{Sqrt}[b^2 - 4*a*c] + 4*c*x^2)/(b - \text{Sqrt}[b^2 - 4*a*c])]*\text{EllipticF}[I*\text{ArcSinh}[\text{Sqrt}[2]*\text{Sqrt}[c/(b + \text{Sqrt}[b^2 - 4*a*c])]]*x], (b + \text{Sqrt}[b^2 - 4*a*c])/(b - \text{Sqrt}[b^2 - 4*a*c])]/(420*c^4*\text{Sqrt}[c/(b + \text{Sqrt}[b^2 - 4*a*c])]*\text{Sqrt}[a + b*x^2 + c*x^4])$

### 3.17.3 Rubi [A] (verified)

Time = 1.45 (sec) , antiderivative size = 690, normalized size of antiderivative = 0.91, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.212$ , Rules used = {2207, 2207, 2207, 1511, 27, 1416, 1509}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + bx^2 + cx^4}} dx$$

↓ 2207

$$\frac{\int \frac{e^2(21Bcd - 6bBe + 7Ace)x^6 + e(21Bcd^2 + 21Aced - 5aBe^2)x^4 + 7cd^2(Bd + 3Ae)x^2 + 7Acd^3}{\sqrt{cx^4 + bx^2 + a}} dx}{7c} + \frac{Be^3x^5\sqrt{a + bx^2 + cx^4}}{7c}$$

↓ 2207

---

3.17.  $\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+bx^2+cx^4}} dx$

$$\int \frac{e \left( 7Ac(15cd-4be) + B(105c^2d^2+24b^2e^2-ce(84bd+25ae)) \right) x^4 + (21Ace(5cd^2-ae^2) + B(35c^2d^3-63ace^2d+18abe^3)) x^2 + 35Ac^2d^3}{\sqrt{cx^4+bx^2+a}} dx + \frac{e^2 x^3 \sqrt{a+bx^2+cx^4}}{5c}$$

$$\frac{Be^3 x^5 \sqrt{a+bx^2+cx^4}}{7c}$$

↓ 2207

$$\int \frac{(7Ace(45c^2d^2+8b^2e^2-3ce(10bd+3ae)) + B(105c^3d^3-21c^2e(10bd+9ae)d-48b^3e^3+8bce^2(21bd+13ae))) x^2 + 7Ac(15c^2d^3-15ace^2d+4abe^3) - aBe(105c^2d^2+24b^2e^2)}{\sqrt{cx^4+bx^2+a}} dx$$

$$\frac{Be^3 x^5 \sqrt{a+bx^2+cx^4}}{7c}$$

↓ 1511

$$\sqrt{a} \left( \frac{\sqrt{c} (7Ac(4abe^3-15acde^2+15c^2d^3) - aBe(-ce(25ae+84bd)+24b^2e^2+105c^2d^2))}{\sqrt{a}} + 7Ace(-3ce(3ae+10bd)+8b^2e^2+45c^2d^2) + B(-21c^2de(9ae+10bd)+8bce^2(105c^2d^2+24b^2e^2)) \right) / \sqrt{c}$$

$$\frac{Be^3 x^5 \sqrt{a+bx^2+cx^4}}{7c}$$

↓ 27

$$\sqrt{a} \left( \frac{\sqrt{c} (7Ac(4abe^3-15acde^2+15c^2d^3) - aBe(-ce(25ae+84bd)+24b^2e^2+105c^2d^2))}{\sqrt{a}} + 7Ace(-3ce(3ae+10bd)+8b^2e^2+45c^2d^2) + B(-21c^2de(9ae+10bd)+8bce^2(105c^2d^2+24b^2e^2)) \right) / \sqrt{c}$$

$$\frac{Be^3 x^5 \sqrt{a+bx^2+cx^4}}{7c}$$

↓ 1416

$$\sqrt[4]{a} (\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \text{EllipticF} \left( 2 \arctan \left( \frac{\sqrt[4]{Cx}}{\sqrt{a}} \right), \frac{1}{4} \left( 2 - \frac{b}{\sqrt{a}\sqrt{c}} \right) \right) \left( \frac{\sqrt{c} (7Ac(4abe^3-15acde^2+15c^2d^3) - aBe(-ce(25ae+84bd)+24b^2e^2+105c^2d^2))}{\sqrt{a}} + 7Ace(-3ce(3ae+10bd)+8b^2e^2+45c^2d^2) + B(-21c^2de(9ae+10bd)+8bce^2(105c^2d^2+24b^2e^2)) \right) / (2c^{3/4} \sqrt{a+bx^2+cx^4})$$

$$\frac{Be^3 x^5 \sqrt{a+bx^2+cx^4}}{7c}$$

↓ 1509

3.17.  $\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+bx^2+cx^4}} dx$

$$\frac{4\sqrt{a}(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}\text{EllipticF}\left(2\arctan\left(\frac{4\sqrt{Cx}}{4\sqrt{a}}\right),\frac{1}{4}\left(2-\frac{b}{\sqrt{a}\sqrt{c}}\right)\right)\left(\frac{\sqrt{c}(7Ac(4abe^3-15acde^2+15c^2d^3)-aBe(-ce(25ae+84bd)+24b^2e^2+105c^2d^2)}{\sqrt{a}}\right)}{2c^{3/4}\sqrt{a+bx^2+cx^4}}$$

$$\frac{Be^3x^5\sqrt{a+bx^2+cx^4}}{7c}$$

input `Int[(A + B*x^2)*(d + e*x^2)^3/Sqrt[a + b*x^2 + c*x^4], x]`

output `(B*e^3*x^5*Sqrt[a + b*x^2 + c*x^4])/(7*c) + ((e^2*(21*B*c*d - 6*b*B*e + 7*A*c*e)*x^3*Sqrt[a + b*x^2 + c*x^4])/(5*c) + ((e*(7*A*c*e*(15*c*d - 4*b*e) + B*(105*c^2*d^2 + 24*b^2*e^2 - c*e*(84*b*d + 25*a*e)))*x*Sqrt[a + b*x^2 + c*x^4])/(3*c) + (-(((7*A*c*e*(45*c^2*d^2 + 8*b^2*e^2 - 3*c*e*(10*b*d + 3*a*e)) + B*(105*c^3*d^3 - 48*b^3*e^3 - 21*c^2*d*e*(10*b*d + 9*a*e) + 8*b*c*e^2*(21*b*d + 13*a*e)))*(-(x*Sqrt[a + b*x^2 + c*x^4])/(Sqrt[a] + Sqrt[c]*x^2)) + (a^(1/4)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(c^(1/4)*Sqrt[a + b*x^2 + c*x^4]))/Sqrt[c] + (a^(1/4)*(7*A*c*e*(45*c^2*d^2 + 8*b^2*e^2 - 3*c*e*(10*b*d + 3*a*e)) + B*(105*c^3*d^3 - 48*b^3*e^3 - 21*c^2*d*e*(10*b*d + 9*a*e) + 8*b*c*e^2*(21*b*d + 13*a*e)) + (Sqrt[c]*(7*A*c*(15*c^2*d^3 - 15*a*c*d*e^2 + 4*a*b*e^3) - a*B*e*(105*c^2*d^2 + 24*b^2*e^2 - c*e*(84*b*d + 25*a*e))))/Sqrt[a])*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*c^(3/4)*Sqrt[a + b*x^2 + c*x^4]))/(3*c))/(5*c))/(7*c)`

### 3.17.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 1416 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`

$$3.17. \int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+bx^2+cx^4}} dx$$

```
rule 1509 Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol]
:= With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

```
rule 1511 Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol]
:= With[{q = Rt[c/a, 2]}, Simp[(e + d*q)/q Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Simp[e/q Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

```
rule 2207 Int[(Px_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol]
:= With[{n = Expon[Px, x^2], e = Coeff[Px, x^2, Expon[Px, x^2]]}, Simp[e*x^(2*n - 3)*((a + b*x^2 + c*x^4)^(p + 1)/(c*(2*n + 4*p + 1))), x] + Simp[1/(c*(2*n + 4*p + 1)) Int[(a + b*x^2 + c*x^4)^p*ExpandToSum[c*(2*n + 4*p + 1)*Px - a*e*(2*n - 3)*x^(2*n - 4) - b*e*(2*n + 2*p - 1)*x^(2*n - 2) - c*e*(2*n + 4*p + 1)*x^(2*n), x], x], x] /; FreeQ[{a, b, c, p}, x] && PolyQ[Px, x^2] && Expon[Px, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && !LtQ[p, -1]
```

### 3.17.4 Maple [A] (verified)

Time = 10.89 (sec) , antiderivative size = 673, normalized size of antiderivative = 0.89

| method   | result   |
|----------|--|
| elliptic | $\frac{B e^3 x^5 \sqrt{c x^4 + b x^2 + a}}{7c} + \frac{(e^3 A + 3B e^2 d - \frac{6B e^3 b}{7c}) x^3 \sqrt{c x^4 + b x^2 + a}}{5c} + \frac{\left(3A d e^2 + 3B d^2 e - \frac{5B e^3 a}{7c} - \frac{4(e^3 A + 3B e^2 d - \frac{6B e^3 b}{7c}) b}{5c}\right) x \sqrt{c x^4 + b x^2 + a}}{3c}$ |
| risch    | Expression too large to display  |
| default  | Expression too large to display  |

```
input int((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURNVERBOSE)
```

$$3.17. \int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+bx^2+cx^4}} dx$$

output  $\frac{1}{7}B^3e^3x^5(c^2x^4+bx^2+a)^{1/2}/c+1/5(e^3A+3B^2e^2d-6/7B^3/c^2b)/c^2x^3(c^2x^4+bx^2+a)^{1/2}+1/3(3Ade^2+3Bd^2e-5/7B^3/c^2a-4/5(e^3A+3B^2e^2d-6/7B^3/c^2b)/c^2b)/c^2x(c^2x^4+bx^2+a)^{1/2}+1/4(A^3d^3-1/3(3Ade^2+3Bd^2e-5/7B^3/c^2a-4/5(e^3A+3B^2e^2d-6/7B^3/c^2b)/c^2b)/c^2a)^2^{1/2}/((-b+(-4ac+b^2)^{1/2})/a)^{1/2}(4-2(-b+(-4ac+b^2)^{1/2})/ax^2)^{1/2}(4+2(b+(-4ac+b^2)^{1/2})/ax^2)^{1/2}/(c^2x^4+bx^2+a)^{1/2}*\text{EllipticF}(1/2*x^2^{1/2}*((-b+(-4ac+b^2)^{1/2})/a)^{1/2},1/2*(-4+2*b*(b+(-4ac+b^2)^{1/2})/a/c)^{1/2})-1/2(3Ade^2+3Bd^2e-5/7B^3/c^2a-4/5(e^3A+3B^2e^2d-6/7B^3/c^2b)/c^2a-2/3(3Ade^2+3Bd^2e-5/7B^3/c^2a-4/5(e^3A+3B^2e^2d-6/7B^3/c^2b)/c^2b)/c^2b)*a^2^{1/2}/((-b+(-4ac+b^2)^{1/2})/a)^{1/2}(4-2(-b+(-4ac+b^2)^{1/2})/ax^2)^{1/2}(4+2(b+(-4ac+b^2)^{1/2})/ax^2)^{1/2}/(c^2x^4+bx^2+a)^{1/2}/(b+(-4ac+b^2)^{1/2})*(\text{EllipticF}(1/2*x^2^{1/2}*((-b+(-4ac+b^2)^{1/2})/a)^{1/2},1/2*(-4+2*b*(b+(-4ac+b^2)^{1/2})/a/c)^{1/2})-\text{EllipticE}(1/2*x^2^{1/2}*((-b+(-4ac+b^2)^{1/2})/a)^{1/2}),1/2*(-4+2*b*(b+(-4ac+b^2)^{1/2})/a/c)^{1/2}))$

### 3.17.5 Fracas [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 1041, normalized size of antiderivative = 1.38

$$\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+bx^2+cx^4}} dx = \text{Too large to display}$$

input `integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fracas")`

```

output 1/210*(sqrt(1/2)*((105*B*a*c^4*d^3 - 105*(2*B*a*b*c^3 - 3*A*a*c^4)*d^2*e +
  21*(8*B*a*b^2*c^2 - (9*B*a^2 + 10*A*a*b)*c^3)*d*e^2 - (48*B*a*b^3*c + 63*
  A*a^2*c^3 - 8*(13*B*a^2*b + 7*A*a*b^2)*c^2)*e^3)*x*sqrt((b^2 - 4*a*c)/c^2)
  - (105*B*a*b*c^3*d^3 - 105*(2*B*a*b^2*c^2 - 3*A*a*b*c^3)*d^2*e + 21*(8*B*
  a*b^3*c - (9*B*a^2*b + 10*A*a*b^2)*c^2)*d*e^2 - (48*B*a*b^4 + 63*A*a^2*b*c
  ^2 - 8*(13*B*a^2*b^2 + 7*A*a*b^3)*c)*e^3)*x)*sqrt(c)*sqrt((c*sqrt((b^2 - 4
  *a*c)/c^2) - b)/c)*elliptic_e(arcsin(sqrt(1/2)*sqrt((c*sqrt((b^2 - 4*a*c)/
  c^2) - b)/c)/x), 1/2*(b*c*sqrt((b^2 - 4*a*c)/c^2) + b^2 - 2*a*c)/(a*c)) -
  sqrt(1/2)*((105*(B*a*c^4 - A*c^5)*d^3 - 105*(2*B*a*b*c^3 - (3*A + B)*a*c^4
  )*d^2*e + 21*(8*B*a*b^2*c^2 + 5*A*a*c^4 - (9*B*a^2 + 2*(5*A + 2*B)*a*b)*c^
  3)*d*e^2 - (48*B*a*b^3*c + ((63*A + 25*B)*a^2 + 28*A*a*b)*c^3 - 8*(13*B*a^
  2*b + (7*A + 3*B)*a*b^2)*c^2)*e^3)*x*sqrt((b^2 - 4*a*c)/c^2) - (105*(B*a*b
  *c^3 + A*b*c^4)*d^3 - 105*(2*B*a*b^2*c^2 - (3*A - B)*a*b*c^3)*d^2*e + 21*(
  8*B*a*b^3*c - 5*A*a*b*c^3 - (9*B*a^2*b + 2*(5*A - 2*B)*a*b^2)*c^2)*d*e^2 -
  (48*B*a*b^4 + ((63*A - 25*B)*a^2*b - 28*A*a*b^2)*c^2 - 8*(13*B*a^2*b^2 +
  (7*A - 3*B)*a*b^3)*c)*e^3)*x)*sqrt(c)*sqrt((c*sqrt((b^2 - 4*a*c)/c^2) - b)
  /c)*elliptic_f(arcsin(sqrt(1/2)*sqrt((c*sqrt((b^2 - 4*a*c)/c^2) - b)/c)/x)
  , 1/2*(b*c*sqrt((b^2 - 4*a*c)/c^2) + b^2 - 2*a*c)/(a*c)) + 2*(15*B*a*c^4*e
  ^3*x^6 + 105*B*a*c^4*d^3 + 3*(21*B*a*c^4*d*e^2 - (6*B*a*b*c^3 - 7*A*a*c^4)
  *e^3)*x^4 - 105*(2*B*a*b*c^3 - 3*A*a*c^4)*d^2*e + 21*(8*B*a*b^2*c^2 - (...

```

### 3.17.6 Sympy [F]

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + bx^2 + cx^4}} dx$$

```
input integrate((B*x**2+A)*(e*x**2+d)**3/(c*x**4+b*x**2+a)**(1/2), x)
```

```
output Integral((A + B*x**2)*(d + e*x**2)**3/sqrt(a + b*x**2 + c*x**4), x)
```

**3.17.7 Maxima [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^3}{\sqrt{cx^4 + bx^2 + a}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^3/sqrt(c*x^4 + b*x^2 + a), x)`

**3.17.8 Giac [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^3}{\sqrt{cx^4 + bx^2 + a}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^3/sqrt(c*x^4 + b*x^2 + a), x)`

**3.17.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^3}{\sqrt{cx^4 + bx^2 + a}} dx$$

input `int(((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(1/2),x)`

output `int(((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(1/2), x)`

**3.18** 
$$\int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+bx^2+cx^4}} dx$$

|        |   |     |
|--------|---|-----|
| 3.18.1 | Optimal result . . . . .                            | 159 |
| 3.18.2 | Mathematica [C] (verified) . . . . .                | 160 |
| 3.18.3 | Rubi [A] (verified) . . . . .                       | 161 |
| 3.18.4 | Maple [A] (verified) . . . . .                      | 164 |
| 3.18.5 | Fricas [A] (verification not implemented) . . . . . | 165 |
| 3.18.6 | Sympy [F] . . . . .                                 | 165 |
| 3.18.7 | Maxima [F] . . . . .                                | 166 |
| 3.18.8 | Giac [F] . . . . .                                  | 166 |
| 3.18.9 | Mupad [F(-1)] . . . . .                             | 166 |

**3.18.1 Optimal result**

Integrand size = 33, antiderivative size = 528

$$\begin{aligned} & \int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+bx^2+cx^4}} dx \\ &= \frac{e(10Bcd-4bBe+5Ace)x\sqrt{a+bx^2+cx^4}}{15c^2} + \frac{Be^2x^3\sqrt{a+bx^2+cx^4}}{5c} \\ &+ \frac{(10Ace(3cd-be)+B(15c^2d^2+8b^2e^2-ce(20bd+9ae)))x\sqrt{a+bx^2+cx^4}}{15c^{5/2}(\sqrt{a}+\sqrt{cx^2})} \\ &- \frac{\sqrt[4]{a}(10Ace(3cd-be)+B(15c^2d^2+8b^2e^2-ce(20bd+9ae)))(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}E\left(2\arctan\left(\frac{\sqrt{a+bx^2+cx^4}}{\sqrt{a}+\sqrt{cx^2}}\right)\right)}{15c^{11/4}\sqrt{a+bx^2+cx^4}} \\ &+ \frac{\sqrt[4]{a}\left(10Ace(3cd-be)+B(15c^2d^2+8b^2e^2-ce(20bd+9ae))-\frac{\sqrt{c}(2aBe(5cd-2be)-5Ac(3cd^2-ae^2))}{\sqrt{a}}\right)(\sqrt{a}+\sqrt{cx^2})}{30c^{11/4}\sqrt{a+bx^2+cx^4}} \end{aligned}$$



output 
$$\begin{aligned} & 1/15*e*(5*A*c*e-4*B*b*e+10*B*c*d)*x*(c*x^4+b*x^2+a)^{(1/2)}/c^2+1/5*B*e^2*x^3*(c*x^4+b*x^2+a)^{(1/2)}/c+1/15*(10*A*c*e*(-b*e+3*c*d)+B*(15*c^2*d^2+8*b^2*e^2-c*e*(9*a*e+20*b*d)))*x*(c*x^4+b*x^2+a)^{(1/2)}/c^{(5/2)}/(a^{(1/2)}+x^2*c^{(1/2)})-1/15*a^{(1/4)}*(10*A*c*e*(-b*e+3*c*d)+B*(15*c^2*d^2+8*b^2*e^2-c*e*(9*a*e+20*b*d)))*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/c^{(11/4)}/(c*x^4+b*x^2+a)^{(1/2)}+1/30*a^{(1/4)}*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/c^{(11/4)}/(c*x^4+b*x^2+a)^{(1/2)} \\ & - (2*a*B*e*(-2*b*e+5*c*d)-5*A*c*(-a*e^2+3*c*d^2))*c^{(1/2)}/a^{(1/2)}*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/c^{(11/4)}/(c*x^4+b*x^2+a)^{(1/2)} \end{aligned}$$

### 3.18.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 12.81 (sec) , antiderivative size = 674, normalized size of antiderivative = 1.28

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + bx^2 + cx^4}} dx$$

$$= \frac{4c\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}ex(a + bx^2 + cx^4)(5Ace + B(10cd - 4be + 3cex^2)) + i(-b + \sqrt{b^2 - 4ac})(10Ace(3cd - be))}{\dots}$$

input `Integrate[((A + B*x^2)*(d + e*x^2)^2)/Sqrt[a + b*x^2 + c*x^4],x]`

```
output (4*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*e*x*(a + b*x^2 + c*x^4)*(5*A*c*e + B*(10*c*d - 4*b*e + 3*c*e*x^2)) + I*(-b + Sqrt[b^2 - 4*a*c])*(10*A*c*e*(3*c*d - b*e) + B*(15*c^2*d^2 + 8*b^2*e^2 - c*e*(20*b*d + 9*a*e)))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] - I*(-8*b^3*B*e^2 + 2*b^2*e*(10*B*c*d + 5*A*c*e + 4*B*Sqrt[b^2 - 4*a*c]*e) - b*c*(15*B*c*d^2 + B*e*(20*Sqrt[b^2 - 4*a*c]*d - 17*a*e) + 10*A*e*(3*c*d + Sqrt[b^2 - 4*a*c]*e)) + c*(B*(-9*a*Sqrt[b^2 - 4*a*c]*e^2 + 5*c*d*(3*Sqrt[b^2 - 4*a*c]*d - 4*a*e)) + 10*A*c*(3*c*d^2 + e*(3*Sqrt[b^2 - 4*a*c]*d - a*e))))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])])]/(60*c^3*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[a + b*x^2 + c*x^4])
```

### 3.18.3 Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 498, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.212$ , Rules used = {2207, 2207, 25, 1511, 27, 1416, 1509}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + bx^2 + cx^4}} dx$$

↓ 2207

$$\int \frac{e(10Bcd - 4bBe + 5Ace)x^4 + (5Bcd^2 + 10Aced - 3aBe^2)x^2 + 5Acd^2}{\sqrt{cx^4 + bx^2 + a}} dx + \frac{Be^2x^3\sqrt{a + bx^2 + cx^4}}{5c}$$

↓ 2207

$$\int \frac{-((10Ace(3cd - be) + B(15c^2d^2 + 8b^2e^2 - ce(20bd + 9ae)))x^2 + 2aBe(5cd - 2be) - 5Ac(3cd^2 - ae^2))}{\sqrt{cx^4 + bx^2 + a}} dx + \frac{ex\sqrt{a + bx^2 + cx^4}(5Ace - 4bBe + 10Bcd)}{3c}$$


---


$$\frac{Be^2x^3\sqrt{a + bx^2 + cx^4}}{5c}$$

↓ 25

---

3.18.  $\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + bx^2 + cx^4}} dx$

$$\frac{ex\sqrt{a+bx^2+cx^4}(5Ace-4bBe+10Bcd)}{3c} - \frac{\int \frac{-((10Ace(3cd-be)+B(15c^2d^2+8b^2e^2-ce(20bd+9ae)))x^2)+2aBe(5cd-2be)-5Ac(3cd^2-ae^2)}{\sqrt{cx^4+bx^2+a}} dx}{3c} +$$

$$\frac{Be^2x^3\sqrt{a+bx^2+cx^4}}{5c}$$

↓ 1511

$$\frac{ex\sqrt{a+bx^2+cx^4}(5Ace-4bBe+10Bcd)}{3c} - \frac{\sqrt{a}(B(-ce(9ae+20bd)+8b^2e^2+15c^2d^2)+10Ace(3cd-be)) \int \frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{a}\sqrt{cx^4+bx^2+a}} dx}{\sqrt{c}} - \frac{(-9a^{3/2}Bce^2+\sqrt{a}(10Ace(3cd-be)))}{3c}$$

$$\frac{Be^2x^3\sqrt{a+bx^2+cx^4}}{5c}$$

↓ 27

$$\frac{ex\sqrt{a+bx^2+cx^4}(5Ace-4bBe+10Bcd)}{3c} - \frac{(B(-ce(9ae+20bd)+8b^2e^2+15c^2d^2)+10Ace(3cd-be)) \int \frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{cx^4+bx^2+a}} dx}{\sqrt{c}} - \frac{(-9a^{3/2}Bce^2+\sqrt{a}(10Ace(3cd-be)))}{3c}$$

$$\frac{Be^2x^3\sqrt{a+bx^2+cx^4}}{5c}$$

↓ 1416

$$\frac{ex\sqrt{a+bx^2+cx^4}(5Ace-4bBe+10Bcd)}{3c} - \frac{(B(-ce(9ae+20bd)+8b^2e^2+15c^2d^2)+10Ace(3cd-be)) \int \frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{cx^4+bx^2+a}} dx}{\sqrt{c}} - \frac{(\sqrt{a}+\sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \text{EllipticE}}{3c}$$

$$\frac{Be^2x^3\sqrt{a+bx^2+cx^4}}{5c}$$

↓ 1509

$$\frac{ex\sqrt{a+bx^2+cx^4}(5Ace-4bBe+10Bcd)}{3c} - \frac{\left( \frac{\sqrt[4]{a}(\sqrt{a}+\sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{4}\left(2-\frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{\sqrt[4]{C}\sqrt{a+bx^2+cx^4}} - \frac{x\sqrt{a+bx^2+cx^4}}{\sqrt{a}+\sqrt{cx^2}} \right) (B(-ce(9ae+20bd)+8b^2e^2+15c^2d^2)+10Ace(3cd-be))}{\sqrt{c}}$$

$$\frac{Be^2x^3\sqrt{a+bx^2+cx^4}}{5c}$$

input `Int[((A + B*x^2)*(d + e*x^2)^2)/Sqrt[a + b*x^2 + c*x^4], x]`

3.18.  $\int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+bx^2+cx^4}} dx$

```
output (B*e^2*x^3*Sqrt[a + b*x^2 + c*x^4])/(5*c) + ((e*(10*B*c*d - 4*b*B*e + 5*A*
c*e))*x*Sqrt[a + b*x^2 + c*x^4])/(3*c) - (((10*A*c*e*(3*c*d - b*e) + B*(15*
c^2*d^2 + 8*b^2*e^2 - c*e*(20*b*d + 9*a*e)))*(-(x*Sqrt[a + b*x^2 + c*x^4]
)/(Sqrt[a] + Sqrt[c]*x^2)) + (a^(1/4)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*
x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(
1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(c^(1/4)*Sqrt[a + b*x^2 + c*x^4])))/S
qrt[c] - ((15*A*c^(5/2)*d^2 - 9*a^(3/2)*B*c*e^2 - a*Sqrt[c]*e*(10*B*c*d -
4*b*B*e + 5*A*c*e) + Sqrt[a]*(10*A*c*e*(3*c*d - b*e) + B*(15*c^2*d^2 - 20*
b*c*d*e + 8*b^2*e^2)))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(S
qrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(
Sqrt[a]*Sqrt[c]))/4])/(2*a^(1/4)*c^(3/4)*Sqrt[a + b*x^2 + c*x^4]))/(3*c))/
(5*c)
```

### 3.18.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 1416 Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c
/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/
(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))
], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

```
rule 1509 Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbo
l] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q
^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*
x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2
/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2
- 4*a*c, 0] && PosQ[c/a]
```

```
rule 1511 Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbo
l] := With[{q = Rt[c/a, 2]}, Simp[(e + d*q)/q Int[1/Sqrt[a + b*x^2 + c*x^
4], x], x] - Simp[e/q Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /;
NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && Pos
Q[c/a]
```

$$3.18. \int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+bx^2+cx^4}} dx$$

```
rule 2207 Int[(Px_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := With[{n =
  Expon[Px, x^2], e = Coeff[Px, x^2, Expon[Px, x^2]]}, Simp[e*x^(2*n - 3)*((
  a + b*x^2 + c*x^4)^(p + 1)/(c*(2*n + 4*p + 1))), x] + Simp[1/(c*(2*n + 4*p
  + 1)) Int[(a + b*x^2 + c*x^4)^p*ExpandToSum[c*(2*n + 4*p + 1)*Px - a*e*(2
  *n - 3)*x^(2*n - 4) - b*e*(2*n + 2*p - 1)*x^(2*n - 2) - c*e*(2*n + 4*p + 1)
  *x^(2*n), x], x] /; FreeQ[{a, b, c, p}, x] && PolyQ[Px, x^2] && Expon[
  Px, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && !LtQ[p, -1]
```

### 3.18.4 Maple [A] (verified)

Time = 4.28 (sec) , antiderivative size = 508, normalized size of antiderivative = 0.96

| method   | result   |
|----------|--|
| elliptic | $\frac{B e^2 x^3 \sqrt{c x^4 + b x^2 + a}}{5c} + \frac{(A e^2 + 2 B e d - \frac{4 b B e^2}{5c}) x \sqrt{c x^4 + b x^2 + a}}{3c} + \frac{\left( d^2 A - \frac{a(A e^2 + 2 B e d - \frac{4 b B e^2}{5c})}{3c} \right) \sqrt{2} \sqrt{4 - \frac{2(-b + \sqrt{-4ac + b^2})x^2}{a}}}{4\sqrt{-b + \sqrt{-4ac + b^2}}}$                                       |
| risch    | $\frac{15 A c^2 d^2 \sqrt{2} \sqrt{4 - \frac{2(-b + \sqrt{-4ac + b^2})x^2}{a}} \sqrt{4 + \frac{2(b + \sqrt{-4ac + b^2})x^2}{a}} F\left(\frac{x \sqrt{2} \sqrt{-b + \sqrt{-4ac + b^2}}}{2}\right)}{15c^2} - \frac{e x (3 B e x^2 c + 5 A c e - 4 B b e + 10 B c d) \sqrt{c x^4 + b x^2 + a}}{4 \sqrt{-b + \sqrt{-4ac + b^2}} \sqrt{c x^4 + b x^2 + a}}$ |
| default  | Expression too large to display  |

```
input int((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/5*B*e^2*x^3*(c*x^4+b*x^2+a)^(1/2)/c+1/3*(A*e^2+2*B*e*d-4/5*b/c*B*e^2)/c*
x*(c*x^4+b*x^2+a)^(1/2)+1/4*(d^2*A-1/3*a/c*(A*e^2+2*B*e*d-4/5*b/c*B*e^2))*
2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x
^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*E
llipticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-
4*a*c+b^2)^(1/2))/a/c)^(1/2))-1/2*(2*A*e*d+B*d^2-3/5*a/c*B*e^2-2/3*b/c*(A*
e^2+2*B*e*d-4/5*b/c*B*e^2))*a*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4
-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)
^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*(EllipticF(1/2*x*2^(1/
2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/
c)^(1/2))-EllipticE(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-
4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2)))
```

$$3.18. \int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+bx^2+cx^4}} dx$$

### 3.18.5 Fracas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 681, normalized size of antiderivative = 1.29

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + bx^2 + cx^4}} dx$$

$$\sqrt{\frac{1}{2}} \left( (15 Bac^3 d^2 - 10(2 Babc^2 - 3 Aac^3)de + (8 Bab^2c - (9 Ba^2 + 10 Aab)c^2)e^2)x \sqrt{\frac{b^2 - 4ac}{c^2}} - (15 Babc^2 d \right.$$

=

```
input integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fracas")
```

```
output 1/30*(sqrt(1/2)*((15*B*a*c^3*d^2 - 10*(2*B*a*b*c^2 - 3*A*a*c^3)*d*e + (8*B
*a*b^2*c - (9*B*a^2 + 10*A*a*b)*c^2)*e^2)*x*sqrt((b^2 - 4*a*c)/c^2) - (15*
B*a*b*c^2*d^2 - 10*(2*B*a*b^2*c - 3*A*a*b*c^2)*d*e + (8*B*a*b^3 - (9*B*a^2
*b + 10*A*a*b^2)*c)*e^2)*x)*sqrt(c)*sqrt((c*sqrt((b^2 - 4*a*c)/c^2) - b)/c
)*elliptic_e(arcsin(sqrt(1/2)*sqrt((c*sqrt((b^2 - 4*a*c)/c^2) - b)/c)/x),
1/2*(b*c*sqrt((b^2 - 4*a*c)/c^2) + b^2 - 2*a*c)/(a*c)) - sqrt(1/2)*((15*(B
*a*c^3 - A*c^4)*d^2 - 10*(2*B*a*b*c^2 - (3*A + B)*a*c^3)*d*e + (8*B*a*b^2*
c + 5*A*a*c^3 - (9*B*a^2 + 2*(5*A + 2*B)*a*b)*c^2)*e^2)*x*sqrt((b^2 - 4*a*
c)/c^2) - (15*(B*a*b*c^2 + A*b*c^3)*d^2 - 10*(2*B*a*b^2*c - (3*A - B)*a*b*
c^2)*d*e + (8*B*a*b^3 - 5*A*a*b*c^2 - (9*B*a^2*b + 2*(5*A - 2*B)*a*b^2)*c
)*e^2)*x)*sqrt(c)*sqrt((c*sqrt((b^2 - 4*a*c)/c^2) - b)/c)*elliptic_f(arcsin
(sqrt(1/2)*sqrt((c*sqrt((b^2 - 4*a*c)/c^2) - b)/c)/x), 1/2*(b*c*sqrt((b^2
- 4*a*c)/c^2) + b^2 - 2*a*c)/(a*c)) + 2*(3*B*a*c^3*e^2*x^4 + 15*B*a*c^3*d^
2 - 10*(2*B*a*b*c^2 - 3*A*a*c^3)*d*e + (8*B*a*b^2*c - (9*B*a^2 + 10*A*a*b)
*c^2)*e^2 + (10*B*a*c^3*d*e - (4*B*a*b*c^2 - 5*A*a*c^3)*e^2)*x^2)*sqrt(c*x
^4 + b*x^2 + a))/(a*c^4*x)
```

### 3.18.6 Sympy [F]

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + bx^2 + cx^4}} dx$$

```
input integrate((B*x**2+A)*(e*x**2+d)**2/(c*x**4+b*x**2+a)**(1/2),x)
```

```
output Integral((A + B*x**2)*(d + e*x**2)**2/sqrt(a + b*x**2 + c*x**4), x)
```

---

3.18.  $\int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+bx^2+cx^4}} dx$

**3.18.7 Maxima [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^2}{\sqrt{cx^4 + bx^2 + a}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^2/sqrt(c*x^4 + b*x^2 + a), x)`

**3.18.8 Giac [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^2}{\sqrt{cx^4 + bx^2 + a}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^2/sqrt(c*x^4 + b*x^2 + a), x)`

**3.18.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^2}{\sqrt{cx^4 + bx^2 + a}} dx$$

input `int(((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(1/2),x)`

output `int(((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(1/2), x)`

### 3.19 $\int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+bx^2+cx^4}} dx$

|        |   |     |
|--------|---|-----|
| 3.19.1 | Optimal result                            | 167 |
| 3.19.2 | Mathematica [C] (verified)                | 168 |
| 3.19.3 | Rubi [A] (verified)                       | 168 |
| 3.19.4 | Maple [A] (verified)                      | 171 |
| 3.19.5 | Fricas [A] (verification not implemented) | 172 |
| 3.19.6 | Sympy [F]                                 | 173 |
| 3.19.7 | Maxima [F]                                | 173 |
| 3.19.8 | Giac [F]                                  | 173 |
| 3.19.9 | Mupad [F(-1)]                             | 174 |

#### 3.19.1 Optimal result

Integrand size = 31, antiderivative size = 368

$$\int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+bx^2+cx^4}} dx = \frac{Bex\sqrt{a+bx^2+cx^4}}{3c} + \frac{(3Bcd-2bBe+3Ace)x\sqrt{a+bx^2+cx^4}}{3c^{3/2}(\sqrt{a}+\sqrt{cx^2})}$$

$$- \frac{\sqrt{a}(3Bcd-2bBe+3Ace)(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} E\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{4}\left(2-\frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{3c^{7/4}\sqrt{a+bx^2+cx^4}}$$

$$+ \frac{\sqrt{a}\left(3Bcd-2bBe+3Ace+\frac{\sqrt{c}(3Acd-aBe)}{\sqrt{a}}\right)(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right),\frac{1}{4}\right)}{6c^{7/4}\sqrt{a+bx^2+cx^4}}$$

output

```
1/3*B*e*x*(c*x^4+b*x^2+a)^(1/2)/c+1/3*(3*A*c*e-2*B*b*e+3*B*c*d)*x*(c*x^4+b*x^2+a)^(1/2)/c^(3/2)/(a^(1/2)+x^2*c^(1/2))-1/3*a^(1/4)*(3*A*c*e-2*B*b*e+3*B*c*d)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2)))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(7/4)/(c*x^4+b*x^2+a)^(1/2)+1/6*a^(1/4)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2)))^(1/2)*(a^(1/2)+x^2*c^(1/2))*(3*B*c*d-2*B*b*e+3*A*c*e+(3*A*c*d-B*a*e)*c^(1/2)/a^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(7/4)/(c*x^4+b*x^2+a)^(1/2)
```



### 3.19.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 11.35 (sec) , antiderivative size = 521, normalized size of antiderivative = 1.42

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + bx^2 + cx^4}} dx$$

$$= \frac{4Bc\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}ex(a + bx^2 + cx^4) - i(-b + \sqrt{b^2 - 4ac})(-3Bcd + 2bBe - 3Ace)\sqrt{\frac{b+\sqrt{b^2-4ac}+2cx^2}{b+\sqrt{b^2-4ac}}}\sqrt{\frac{2b-2\sqrt{b^2-4ac}}{b}}}{\dots}$$

input `Integrate[((A + B*x^2)*(d + e*x^2))/Sqrt[a + b*x^2 + c*x^4],x]`

output `(4*B*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*e*x*(a + b*x^2 + c*x^4) - I*(-b + Sqrt[b^2 - 4*a*c])*(-3*B*c*d + 2*b*B*e - 3*A*c*e)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] + I*(-2*b^2*B*e - c*(6*A*c*d + 3*B*Sqrt[b^2 - 4*a*c]*d - 2*a*B*e + 3*A*Sqrt[b^2 - 4*a*c]*e) + b*(3*B*c*d + 3*A*c*e + 2*B*Sqrt[b^2 - 4*a*c]*e))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])]/(12*c^2*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[a + b*x^2 + c*x^4])`

### 3.19.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 358, normalized size of antiderivative = 0.97, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.161$ , Rules used = {2207, 1511, 27, 1416, 1509}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + bx^2 + cx^4}} dx$$

↓ 2207

---

3.19.  $\int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+bx^2+cx^4}} dx$

$$\begin{aligned}
 & \int \frac{(3Bcd-2bBe+3Ace)x^2+3Acd-aBe}{\sqrt{cx^4+bx^2+a}} dx + \frac{Bex\sqrt{a+bx^2+cx^4}}{3c} \\
 & \quad \downarrow \text{1511} \\
 & \frac{\sqrt{a}\left(\frac{\sqrt{c}(3Acd-aBe)}{\sqrt{a}}+3Ace-2bBe+3Bcd\right) \int \frac{1}{\sqrt{cx^4+bx^2+a}} dx - \sqrt{a}(3Ace-2bBe+3Bcd) \int \frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{a}\sqrt{cx^4+bx^2+a}} dx}{\sqrt{c}} + \\
 & \quad \frac{3c}{3c} \frac{Bex\sqrt{a+bx^2+cx^4}}{3c} \\
 & \quad \downarrow \text{27} \\
 & \frac{\sqrt{a}\left(\frac{\sqrt{c}(3Acd-aBe)}{\sqrt{a}}+3Ace-2bBe+3Bcd\right) \int \frac{1}{\sqrt{cx^4+bx^2+a}} dx - (3Ace-2bBe+3Bcd) \int \frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{cx^4+bx^2+a}} dx}{\sqrt{c}} + \\
 & \quad \frac{3c}{3c} \frac{Bex\sqrt{a+bx^2+cx^4}}{3c} \\
 & \quad \downarrow \text{1416} \\
 & \frac{\sqrt[4]{a}(\sqrt{a}+\sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt{a}}\right), \frac{1}{4}\left(2-\frac{b}{\sqrt{a}\sqrt{c}}\right)\right) \left(\frac{\sqrt{c}(3Acd-aBe)}{\sqrt{a}}+3Ace-2bBe+3Bcd\right)}{2c^{3/4}\sqrt{a+bx^2+cx^4}} - \frac{(3Ace-2bBe+3Bcd) \int \frac{1}{\sqrt{cx^4+bx^2+a}} dx}{\sqrt{c}} \\
 & \quad \frac{3c}{3c} \frac{Bex\sqrt{a+bx^2+cx^4}}{3c} \\
 & \quad \downarrow \text{1509} \\
 & \frac{\sqrt[4]{a}(\sqrt{a}+\sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt{a}}\right), \frac{1}{4}\left(2-\frac{b}{\sqrt{a}\sqrt{c}}\right)\right) \left(\frac{\sqrt{c}(3Acd-aBe)}{\sqrt{a}}+3Ace-2bBe+3Bcd\right)}{2c^{3/4}\sqrt{a+bx^2+cx^4}} - \left(\frac{\sqrt[4]{a}(\sqrt{a}+\sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}}{\sqrt{c}}\right) \\
 & \quad \frac{3c}{3c} \frac{Bex\sqrt{a+bx^2+cx^4}}{3c}
 \end{aligned}$$

input `Int[((A + B*x^2)*(d + e*x^2))/Sqrt[a + b*x^2 + c*x^4], x]`

```
output (B*e*x*Sqrt[a + b*x^2 + c*x^4])/(3*c) + (-(((3*B*c*d - 2*b*B*e + 3*A*c*e)*
(-(x*Sqrt[a + b*x^2 + c*x^4])/(Sqrt[a] + Sqrt[c]*x^2)) + (a^(1/4)*(Sqrt[a
] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*Ellip
ticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4]))/(c^(1/4)
*Sqrt[a + b*x^2 + c*x^4]))/Sqrt[c]) + (a^(1/4)*(3*B*c*d - 2*b*B*e + 3*A*c
*e + (Sqrt[c]*(3*A*c*d - a*B*e))/Sqrt[a])*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a
+ b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x
/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*c^(3/4)*Sqrt[a + b*x^2 + c*x^4
]))/(3*c)
```

### 3.19.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 1416 Int[1/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c
/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/
(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))
], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

```
rule 1509 Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbo
l] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q
^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*
x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2
/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2
- 4*a*c, 0] && PosQ[c/a]
```

```
rule 1511 Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbo
l] := With[{q = Rt[c/a, 2]}, Simp[(e + d*q)/q Int[1/Sqrt[a + b*x^2 + c*x^
4], x], x] - Simp[e/q Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /;
NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && Pos
Q[c/a]
```

```
rule 2207 Int[(Px_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := With[{n =
  Expon[Px, x^2], e = Coeff[Px, x^2, Expon[Px, x^2]]}, Simp[e*x^(2*n - 3)*((
  a + b*x^2 + c*x^4)^(p + 1)/(c*(2*n + 4*p + 1))), x] + Simp[1/(c*(2*n + 4*p
  + 1)) Int[(a + b*x^2 + c*x^4)^p*ExpandToSum[c*(2*n + 4*p + 1)*Px - a*e*(2
  *n - 3)*x^(2*n - 4) - b*e*(2*n + 2*p - 1)*x^(2*n - 2) - c*e*(2*n + 4*p + 1)
  *x^(2*n), x], x], x]] /; FreeQ[{a, b, c, p}, x] && PolyQ[Px, x^2] && Expon[
  Px, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && !LtQ[p, -1]
```

### 3.19.4 Maple [A] (verified)

Time = 3.07 (sec) , antiderivative size = 409, normalized size of antiderivative = 1.11

| method   | result   |
|----------|--|
| elliptic | $\frac{Bex\sqrt{cx^4+bx^2+a}}{3c} + \frac{\left(Ad - \frac{aeB}{3c}\right)\sqrt{2}\sqrt{4 - \frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}}\sqrt{4 + \frac{2(b+\sqrt{-4ac+b^2})x^2}{a}}F\left(\frac{x\sqrt{2}\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}, \sqrt{-4 + \frac{2b(b+\sqrt{-4ac+b^2})}{ac}}}\right)}{4\sqrt{-b+\sqrt{-4ac+b^2}}\sqrt{cx^4+bx^2+a}}$ |
| risch    | $\frac{Bex\sqrt{cx^4+bx^2+a}}{3c} + \frac{3Acd\sqrt{2}\sqrt{4 - \frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}}\sqrt{4 + \frac{2(b+\sqrt{-4ac+b^2})x^2}{a}}F\left(\frac{x\sqrt{2}\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}, \sqrt{-4 + \frac{2b(b+\sqrt{-4ac+b^2})}{ac}}}\right)}{4\sqrt{-b+\sqrt{-4ac+b^2}}\sqrt{cx^4+bx^2+a}}$                             |
| default  | $\frac{Ad\sqrt{2}\sqrt{4 - \frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}}\sqrt{4 + \frac{2(b+\sqrt{-4ac+b^2})x^2}{a}}F\left(\frac{x\sqrt{2}\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}, \sqrt{-4 + \frac{2b(b+\sqrt{-4ac+b^2})}{ac}}}\right)}{4\sqrt{-b+\sqrt{-4ac+b^2}}\sqrt{cx^4+bx^2+a}} + eB\left(\frac{x\sqrt{cx^4+bx^2+a}}{3c}\right)$                  |

```
input int((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURNVERBOSE)
```

3.19.  $\int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+bx^2+cx^4}} dx$

output  $\frac{1}{3}B^*e^*x^*(c^*x^4+b^*x^2+a)^{(1/2)}/c+1/4*(A*d-1/3*a/c^*e^*B)^*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a^*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a^*x^2)^{(1/2)}/(c^*x^4+b^*x^2+a)^{(1/2)}*EllipticF(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-1/2*(A^*e+B^*d-2/3*b/c^*e^*B)^*a^*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a^*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a^*x^2)^{(1/2)}/(c^*x^4+b^*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*(EllipticF(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-EllipticE(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2))}$

### 3.19.5 Fracas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 435, normalized size of antiderivative = 1.18

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + bx^2 + cx^4}} dx$$

$$\sqrt{\frac{1}{2}} \left( (3Bac^2d - (2Babc - 3Aac^2)e)x\sqrt{\frac{b^2-4ac}{c^2}} - (3Babcd - (2Bab^2 - 3Aabc)e)x \right) \sqrt{c} \sqrt{\frac{c\sqrt{\frac{b^2-4ac}{c^2}} - b}{c}} E(a$$

=

input `integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")`

output  $\frac{1}{6}*(\text{sqrt}(1/2)*((3*B^*a^*c^2*d - (2*B^*a^*b^*c - 3*A^*a^*c^2)*e)^*x*\text{sqrt}((b^2 - 4*a*c)/c^2) - (3*B^*a^*b^*c*d - (2*B^*a^*b^2 - 3*A^*a^*b^*c)*e)^*x)*\text{sqrt}(c)*\text{sqrt}((c*\text{sqrt}((b^2 - 4*a*c)/c^2) - b)/c)*\text{elliptic}_e(\arcsin(\text{sqrt}(1/2)*\text{sqrt}((c*\text{sqrt}((b^2 - 4*a*c)/c^2) - b)/c)/x), 1/2*(b^*c*\text{sqrt}((b^2 - 4*a*c)/c^2) + b^2 - 2*a*c)/(a^*c)) - \text{sqrt}(1/2)*((3*(B^*a^*c^2 - A^*c^3)*d - (2*B^*a^*b^*c - (3*A + B)^*a^*c^2)*e)^*x*\text{sqrt}((b^2 - 4*a*c)/c^2) - (3*(B^*a^*b^*c + A^*b^*c^2)*d - (2*B^*a^*b^2 - (3*A - B)^*a^*b^*c)*e)^*x)*\text{sqrt}(c)*\text{sqrt}((c*\text{sqrt}((b^2 - 4*a*c)/c^2) - b)/c)*\text{elliptic}_f(\arcsin(\text{sqrt}(1/2)*\text{sqrt}((c*\text{sqrt}((b^2 - 4*a*c)/c^2) - b)/c)/x), 1/2*(b^*c*\text{sqrt}((b^2 - 4*a*c)/c^2) + b^2 - 2*a*c)/(a^*c)) + 2*(B^*a^*c^2*e^*x^2 + 3*B^*a^*c^2*d - (2*B^*a^*b^*c - 3*A^*a^*c^2)*e)*\text{sqrt}(c^*x^4 + b^*x^2 + a))/(a^*c^3*x)$

**3.19.6 Sympy [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + bx^2 + cx^4}} dx$$

input `integrate((B*x**2+A)*(e*x**2+d)/(c*x**4+b*x**2+a)**(1/2),x)`

output `Integral((A + B*x**2)*(d + e*x**2)/sqrt(a + b*x**2 + c*x**4), x)`

**3.19.7 Maxima [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)}{\sqrt{cx^4 + bx^2 + a}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)*(e*x^2 + d)/sqrt(c*x^4 + b*x^2 + a), x)`

**3.19.8 Giac [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)}{\sqrt{cx^4 + bx^2 + a}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)*(e*x^2 + d)/sqrt(c*x^4 + b*x^2 + a), x)`

**3.19.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)}{\sqrt{cx^4 + bx^2 + a}} dx$$

input `int(((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(1/2), x)`output `int(((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(1/2), x)`

### 3.20 $\int \frac{A+Bx^2}{\sqrt{a+bx^2+cx^4}} dx$

|        |   |     |
|--------|---|-----|
| 3.20.1 | Optimal result                            | 175 |
| 3.20.2 | Mathematica [C] (verified)                | 176 |
| 3.20.3 | Rubi [A] (verified)                       | 176 |
| 3.20.4 | Maple [A] (verified)                      | 178 |
| 3.20.5 | Fricas [A] (verification not implemented) | 179 |
| 3.20.6 | Sympy [F]                                 | 180 |
| 3.20.7 | Maxima [F]                                | 180 |
| 3.20.8 | Giac [F]                                  | 180 |
| 3.20.9 | Mupad [F(-1)]                             | 181 |

#### 3.20.1 Optimal result

Integrand size = 24, antiderivative size = 283

$$\int \frac{A+Bx^2}{\sqrt{a+bx^2+cx^4}} dx = \frac{Bx\sqrt{a+bx^2+cx^4}}{\sqrt{c}(\sqrt{a}+\sqrt{cx^2})} - \frac{\sqrt[4]{a}B(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{4}\left(2-\frac{b}{\sqrt{a\sqrt{c}}}\right)\right)}{c^{3/4}\sqrt{a+bx^2+cx^4}} + \frac{\sqrt[4]{a}\left(B+\frac{A\sqrt{c}}{\sqrt{a}}\right)(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right),\frac{1}{4}\left(2-\frac{b}{\sqrt{a\sqrt{c}}}\right)\right)}{2c^{3/4}\sqrt{a+bx^2+cx^4}}$$

output

```
B*x*(c*x^4+b*x^2+a)^(1/2)/c^(1/2)/(a^(1/2)+x^2*c^(1/2))-a^(1/4)*B*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(3/4)/(c*x^4+b*x^2+a)^(1/2)+1/2*a^(1/4)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*(B+A*c^(1/2)/a^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(3/4)/(c*x^4+b*x^2+a)^(1/2)
```



### 3.20.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 302, normalized size of antiderivative = 1.07

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2 + cx^4}} dx$$

$$= \frac{i \sqrt{\frac{b + \sqrt{b^2 - 4ac} + 2cx^2}{b + \sqrt{b^2 - 4ac}}} \sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}} \left( B(-b + \sqrt{b^2 - 4ac}) E \left( \operatorname{iarcsinh} \left( \sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x \right) \middle| \frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}} \right) + (b + \sqrt{b^2 - 4ac}) \operatorname{EllipticF} \left( \operatorname{iarcsinh} \left( \sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x \right) \middle| \frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}} \right) \right)}{2\sqrt{2}c \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} \sqrt{a + bx^2 + cx^4}}$$

input `Integrate[(A + B*x^2)/Sqrt[a + b*x^2 + c*x^4],x]`

output `((I/2)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c]])*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*(B*(-b + Sqrt[b^2 - 4*a*c])*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + (b*B - 2*A*c - B*Sqrt[b^2 - 4*a*c])*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])))]/(Sqrt[2]*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])])*Sqrt[a + b*x^2 + c*x^4])`

### 3.20.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.01, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {1511, 27, 1416, 1509}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2 + cx^4}} dx$$

$$\downarrow \text{1511}$$

$$\left( \frac{\sqrt{a}B}{\sqrt{c}} + A \right) \int \frac{1}{\sqrt{cx^4 + bx^2 + a}} dx - \frac{\sqrt{a}B \int \frac{\sqrt{a} - \sqrt{cx^2}}{\sqrt{a}\sqrt{cx^4 + bx^2 + a}} dx}{\sqrt{c}}$$

$$\downarrow \text{27}$$

$$\begin{aligned}
& \left( \frac{\sqrt{a}B}{\sqrt{c}} + A \right) \int \frac{1}{\sqrt{cx^4 + bx^2 + a}} dx - \frac{B \int \frac{\sqrt{a} - \sqrt{cx^2}}{\sqrt{cx^4 + bx^2 + a}} dx}{\sqrt{c}} \\
& \quad \downarrow \text{1416} \\
& \frac{(\sqrt{a} + \sqrt{cx^2}) \left( \frac{\sqrt{a}B}{\sqrt{c}} + A \right) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \operatorname{EllipticF} \left( 2 \arctan \left( \frac{\sqrt[4]{c}x}{\sqrt[4]{a}} \right), \frac{1}{4} \left( 2 - \frac{b}{\sqrt{a}\sqrt{c}} \right) \right)}{2\sqrt[4]{a}\sqrt[4]{c}\sqrt{a+bx^2+cx^4}} \\
& \quad \frac{B \int \frac{\sqrt{a} - \sqrt{cx^2}}{\sqrt{cx^4 + bx^2 + a}} dx}{\sqrt{c}} \\
& \quad \downarrow \text{1509} \\
& \frac{(\sqrt{a} + \sqrt{cx^2}) \left( \frac{\sqrt{a}B}{\sqrt{c}} + A \right) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \operatorname{EllipticF} \left( 2 \arctan \left( \frac{\sqrt[4]{c}x}{\sqrt[4]{a}} \right), \frac{1}{4} \left( 2 - \frac{b}{\sqrt{a}\sqrt{c}} \right) \right)}{2\sqrt[4]{a}\sqrt[4]{c}\sqrt{a+bx^2+cx^4}} \\
& \quad B \left( \frac{\sqrt[4]{a}(\sqrt{a}+\sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} E \left( 2 \arctan \left( \frac{\sqrt[4]{c}x}{\sqrt[4]{a}} \right) \middle| \frac{1}{4} \left( 2 - \frac{b}{\sqrt{a}\sqrt{c}} \right) \right)}{\sqrt[4]{c}\sqrt{a+bx^2+cx^4}} - \frac{x\sqrt{a+bx^2+cx^4}}{\sqrt{a+\sqrt{cx^2}}} \right) \\
& \quad \sqrt{c}
\end{aligned}$$

input `Int[(A + B*x^2)/Sqrt[a + b*x^2 + c*x^4],x]`

output `-(B*(-((x*Sqrt[a + b*x^2 + c*x^4])/(Sqrt[a] + Sqrt[c]*x^2)) + (a^(1/4)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4]))/(c^(1/4)*Sqrt[a + b*x^2 + c*x^4]))/Sqrt[c] + ((A + (Sqrt[a]*B)/Sqrt[c])*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*a^(1/4)*c^(1/4)*Sqrt[a + b*x^2 + c*x^4])`

3.20.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1416 `Int[1/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`

rule 1509 `Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`

rule 1511 `Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 2]}, Simp[(e + d*q)/q Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Simp[e/q Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`

3.20.4 Maple [A] (verified)

Time = 0.93 (sec) , antiderivative size = 362, normalized size of antiderivative = 1.28

| method   | result   |
|----------|--|
| default  | $\frac{A\sqrt{2}\sqrt{4-\frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}}\sqrt{4+\frac{2(b+\sqrt{-4ac+b^2})x^2}{a}}F\left(\frac{x\sqrt{2}\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}}{2}, \sqrt{-4+\frac{2b(b+\sqrt{-4ac+b^2})}{ac}}\right)}{4\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}\sqrt{cx^4+bx^2+a}} - \frac{Ba\sqrt{2}\sqrt{4-\frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}}}{4\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}\sqrt{cx^4+bx^2+a}}$ |
| elliptic | $\frac{A\sqrt{2}\sqrt{4-\frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}}\sqrt{4+\frac{2(b+\sqrt{-4ac+b^2})x^2}{a}}F\left(\frac{x\sqrt{2}\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}}{2}, \sqrt{-4+\frac{2b(b+\sqrt{-4ac+b^2})}{ac}}\right)}{4\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}\sqrt{cx^4+bx^2+a}} - \frac{Ba\sqrt{2}\sqrt{4-\frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}}}{4\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}\sqrt{cx^4+bx^2+a}}$ |

3.20.  $\int \frac{A+Bx^2}{\sqrt{a+bx^2+cx^4}} dx$

```
input int((B*x^2+A)/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/4*A*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2)))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))-1/2*B*a*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*(EllipticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))-EllipticE(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2)))
```

### 3.20.5 Fracas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.06

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2 + cx^4}} dx$$

$$2\sqrt{cx^4 + bx^2 + a}Bac + \sqrt{\frac{1}{2}} \left( Bacx\sqrt{\frac{b^2 - 4ac}{c^2}} - Babx \right) \sqrt{c} \sqrt{\frac{c\sqrt{\frac{b^2 - 4ac}{c^2}} - b}{c}} E\left(\arcsin\left(\frac{\sqrt{\frac{1}{2}} \sqrt{\frac{c\sqrt{\frac{b^2 - 4ac}{c^2}} - b}}{x}}}{x}\right) \mid \frac{bc\sqrt{b^2 - 4ac}}{c^2}\right)$$

$$=$$

```
input integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")
```

```
output 1/2*(2*sqrt(c*x^4 + b*x^2 + a)*B*a*c + sqrt(1/2)*(B*a*c*x*sqrt((b^2 - 4*a*c)/c^2) - B*a*b*x)*sqrt(c)*sqrt((c*sqrt((b^2 - 4*a*c)/c^2) - b)/c)*elliptic_e(arcsin(sqrt(1/2)*sqrt((c*sqrt((b^2 - 4*a*c)/c^2) - b)/c)/x), 1/2*(b*c*sqrt((b^2 - 4*a*c)/c^2) + b^2 - 2*a*c)/(a*c)) - sqrt(1/2)*((B*a*c - A*c^2)*x*sqrt((b^2 - 4*a*c)/c^2) - (B*a*b + A*b*c)*x)*sqrt(c)*sqrt((c*sqrt((b^2 - 4*a*c)/c^2) - b)/c)*elliptic_f(arcsin(sqrt(1/2)*sqrt((c*sqrt((b^2 - 4*a*c)/c^2) - b)/c)/x), 1/2*(b*c*sqrt((b^2 - 4*a*c)/c^2) + b^2 - 2*a*c)/(a*c))/a*c^2*x)
```

**3.20.6 Sympy [F]**

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{A + Bx^2}{\sqrt{a + bx^2 + cx^4}} dx$$

input `integrate((B*x**2+A)/(c*x**4+b*x**2+a)**(1/2),x)`

output `Integral((A + B*x**2)/sqrt(a + b*x**2 + c*x**4), x)`

**3.20.7 Maxima [F]**

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}} dx$$

input `integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/sqrt(c*x^4 + b*x^2 + a), x)`

**3.20.8 Giac [F]**

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}} dx$$

input `integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/sqrt(c*x^4 + b*x^2 + a), x)`

**3.20.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2 + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}} dx$$

input `int((A + B*x^2)/(a + b*x^2 + c*x^4)^(1/2), x)`output `int((A + B*x^2)/(a + b*x^2 + c*x^4)^(1/2), x)`

### 3.21 $\int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$

|        |                            |     |
|--------|----------------------------|-----|
| 3.21.1 | Optimal result             | 182 |
| 3.21.2 | Mathematica [C] (verified) | 183 |
| 3.21.3 | Rubi [A] (verified)        | 183 |
| 3.21.4 | Maple [A] (verified)       | 185 |
| 3.21.5 | Fricas [F(-1)]             | 186 |
| 3.21.6 | Sympy [F]                  | 187 |
| 3.21.7 | Maxima [F]                 | 187 |
| 3.21.8 | Giac [F]                   | 187 |
| 3.21.9 | Mupad [F(-1)]              | 188 |

#### 3.21.1 Optimal result

Integrand size = 33, antiderivative size = 436

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = -\frac{(Bd - Ae) \arctan\left(\frac{\sqrt{cd^2 - bde + ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{a + bx^2 + cx^4}}\right)}{2\sqrt{d}\sqrt{e}\sqrt{cd^2 - bde + ae^2}} - \frac{(\sqrt{a}B - A\sqrt{c})(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{2\sqrt[4]{a}\sqrt[4]{c}(\sqrt{cd} - \sqrt{ae})\sqrt{a + bx^2 + cx^4}} + \frac{a^{3/4}\left(\frac{\sqrt{cd}}{\sqrt{a}} + e\right)^2 (Bd - Ae)(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \operatorname{EllipticPi}\left(-\frac{(\sqrt{cd} - \sqrt{ae})^2}{4\sqrt{a}\sqrt{cde}}, 2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{4\sqrt[4]{cde}(cd^2 - ae^2)\sqrt{a + bx^2 + cx^4}}$$

```
output -1/2*(-A*e+B*d)*arctan(x*(a*e^2-b*d*e+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+b*x^2+a)^(1/2))/d^(1/2)/e^(1/2)/(a*e^2-b*d*e+c*d^2)^(1/2)-1/2*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(B*a^(1/2)-A*c^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(1/4)/c^(1/4)/(-e*a^(1/2)+d*c^(1/2))/(c*x^4+b*x^2+a)^(1/2)+1/4*a^(3/4)*(-A*e+B*d)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticPi(sin(2*arctan(c^(1/4)*x/a^(1/4))),-1/4*(-e*a^(1/2)+d*c^(1/2))^2/d/e/a^(1/2)/c^(1/2),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*(e+d*c^(1/2)/a^(1/2))^2*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(1/4)/d/e/(-a*e^2+c*d^2)/(c*x^4+b*x^2+a)^(1/2)
```

### 3.21.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.43 (sec) , antiderivative size = 298, normalized size of antiderivative = 0.68

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx =$$

$$\frac{i\sqrt{\frac{b+\sqrt{b^2-4ac}+2cx^2}{b+\sqrt{b^2-4ac}}}\sqrt{1+\frac{2cx^2}{b-\sqrt{b^2-4ac}}}\left(Bd\text{EllipticF}\left(\text{iarcsinh}\left(\sqrt{2}\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}x\right),\frac{b+\sqrt{b^2-4ac}}{b-\sqrt{b^2-4ac}}\right)+(-Bd+Ae)\text{EllipticPi}\left(\frac{b+\sqrt{b^2-4ac}}{b-\sqrt{b^2-4ac}}\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}\right),\frac{b+\sqrt{b^2-4ac}}{b-\sqrt{b^2-4ac}}\right)}{\sqrt{2}\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}de\sqrt{a+bx^2+cx^4}}$$

input `Integrate[(A + B*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]),x]`

output `((-I)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c]])*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*(B*d*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + (-B*d) + A*e)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])))/(Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])])*d*e*Sqrt[a + b*x^2 + c*x^4])`

### 3.21.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 449, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.121$ , Rules used = {2226, 27, 1416, 2220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx$$

$$\downarrow 2226$$

$$\frac{\sqrt{a}(Bd - Ae) \int \frac{\sqrt{cx^2 + \sqrt{a}}}{\sqrt{a}(ex^2 + d)\sqrt{cx^4 + bx^2 + a}} dx}{\sqrt{cd} - \sqrt{ae}} - \frac{(\sqrt{a}B - A\sqrt{c}) \int \frac{1}{\sqrt{cx^4 + bx^2 + a}} dx}{\sqrt{cd} - \sqrt{ae}}$$

$$\downarrow 27$$

---

3.21.  $\int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$



$$\begin{aligned}
 & \frac{(Bd - Ae) \int \frac{\sqrt{cx^2 + \sqrt{a}}}{(ex^2 + d)\sqrt{cx^4 + bx^2 + a}} dx}{\sqrt{cd} - \sqrt{ae}} - \frac{(\sqrt{a}B - A\sqrt{c}) \int \frac{1}{\sqrt{cx^4 + bx^2 + a}} dx}{\sqrt{cd} - \sqrt{ae}} \\
 & \quad \downarrow \text{1416} \\
 & \frac{(Bd - Ae) \int \frac{\sqrt{cx^2 + \sqrt{a}}}{(ex^2 + d)\sqrt{cx^4 + bx^2 + a}} dx}{\sqrt{cd} - \sqrt{ae}} - \\
 & \frac{(\sqrt{a} + \sqrt{cx^2}) (\sqrt{a}B - A\sqrt{c}) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt{a}}\right), \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{2\sqrt[4]{a}\sqrt[4]{c}\sqrt{a + bx^2 + cx^4} (\sqrt{cd} - \sqrt{ae})} \\
 & \quad \downarrow \text{2220} \\
 & (Bd - Ae) \left( \frac{(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} (\sqrt{ae} + \sqrt{cd}) \text{EllipticPi}\left(-\frac{\sqrt{a}\left(\frac{\sqrt{cd}}{\sqrt{a}} - e\right)^2}{4\sqrt{cde}}, 2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt{a}}\right), \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{4\sqrt[4]{a}\sqrt[4]{cde}\sqrt{a + bx^2 + cx^4}} - \frac{(\sqrt{cd} - \sqrt{ae}) \arctan\left(\frac{\sqrt{cd} - \sqrt{ae}}{2\sqrt{d}\sqrt{e}\sqrt{a + bx^2 + cx^4}}\right)}{2\sqrt{d}\sqrt{e}\sqrt{a + bx^2 + cx^4}} \right) \\
 & \frac{(\sqrt{a} + \sqrt{cx^2}) (\sqrt{a}B - A\sqrt{c}) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt{a}}\right), \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{2\sqrt[4]{a}\sqrt[4]{c}\sqrt{a + bx^2 + cx^4} (\sqrt{cd} - \sqrt{ae})}
 \end{aligned}$$

input `Int[(A + B*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]),x]`

output `-1/2*((Sqrt[a]*B - A*Sqrt[c])*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(a^(1/4)*c^(1/4)*(Sqrt[c]*d - Sqrt[a]*e)*Sqrt[a + b*x^2 + c*x^4]) + ((B*d - A*e)*(-1/2*((Sqrt[c]*d - Sqrt[a]*e)*ArcTan[(Sqrt[c*d^2 - b*d*e + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + b*x^2 + c*x^4])])/(Sqrt[d]*Sqrt[e]*Sqrt[c*d^2 - b*d*e + a*e^2]) + ((Sqrt[c]*d + Sqrt[a]*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-1/4*(Sqrt[a]*((Sqrt[c]*d)/Sqrt[a] - e)^2/(Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(4*a^(1/4)*c^(1/4)*d*e*Sqrt[a + b*x^2 + c*x^4]))/(Sqrt[c]*d - Sqrt[a]*e)`

## 3.21.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 1416 `Int[1/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`
- rule 2220 `Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[-b + c*(d/e) + a*(e/d), 2]*(x/Sqrt[a + b*x^2 + c*x^4])]/(2*d*e*Rt[-b + c*(d/e) + a*(e/d), 2]))], x] + Simp[(B*d + A*e)*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(4*d*e*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticPi[-(e - d*q^2)^2/(4*d*e*q^2), 2*ArcTan[q*x], 1/2 - b/(4*a*q^2)], x] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0] && PosQ[B/A] && PosQ[-b + c*(d/e) + a*(e/d)]`
- rule 2226 `Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2]}, Simp[(A*(c*d + a*e*q) - a*B*(e + d*q))/(c*d^2 - a*e^2) Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] + Simp[a*(B*d - A*e)*((e + d*q)/(c*d^2 - a*e^2)) Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]`

## 3.21.4 Maple [A] (verified)

Time = 1.73 (sec) , antiderivative size = 359, normalized size of antiderivative = 0.82

$$3.21. \int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$$

| method   | result  |
|----------|---|
| default  | $\frac{B\sqrt{2}\sqrt{4-\frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}}\sqrt{4+\frac{2(b+\sqrt{-4ac+b^2})x^2}{a}}F\left(\frac{x\sqrt{2}\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}},\sqrt{-4+\frac{2b(b+\sqrt{-4ac+b^2})}{ac}}}{2},\sqrt{-4+\frac{2b(b+\sqrt{-4ac+b^2})}{ac}}\right)}{4e\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}\sqrt{cx^4+bx^2+a}} + \frac{(Ae-Bd)\sqrt{2}\sqrt{1+\frac{bx^2}{2a}}}{\sqrt{2}\sqrt{1+\frac{bx^2}{2a}}}$                           |
| elliptic | $\frac{B\sqrt{2}\sqrt{4+\frac{2bx^2}{a}-\frac{2x^2\sqrt{-4ac+b^2}}{a}}\sqrt{4+\frac{2bx^2}{a}+\frac{2x^2\sqrt{-4ac+b^2}}{a}}F\left(\frac{x\sqrt{2}\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}},\sqrt{-4+\frac{2b(b+\sqrt{-4ac+b^2})}{ac}}}{2},\sqrt{-4+\frac{2b(b+\sqrt{-4ac+b^2})}{ac}}\right)}{4e\sqrt{-\frac{b}{a}+\frac{\sqrt{-4ac+b^2}}{a}}\sqrt{cx^4+bx^2+a}} + \frac{\sqrt{2}\sqrt{1+\frac{bx^2}{2a}}}{\sqrt{2}\sqrt{1+\frac{bx^2}{2a}}}$ |

```
input int((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/4*B/e*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2)+(A*e-B*d)/e/d*2^(1/2)/(-b/a+1/a*(-4*a*c+b^2)^(1/2))^(1/2)*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2)*(1+1/2*b*x^2/a+1/2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticPi(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),-2/(-b+(-4*a*c+b^2)^(1/2))*a*e/d,(-1/2*(b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2))
```

### 3.21.5 Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = \text{Timed out}$$

```
input integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")
```

```
output Timed out
```

---

3.21.  $\int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$

**3.21.6 Sympy [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = \int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx$$

input `integrate((B*x**2+A)/(e*x**2+d)/(c*x**4+b*x**2+a)**(1/2),x)`

output `Integral((A + B*x**2)/((d + e*x**2)*sqrt(a + b*x**2 + c*x**4)), x)`

**3.21.7 Maxima [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}(ex^2 + d)} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)), x)`

**3.21.8 Giac [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}(ex^2 + d)} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)), x)`

**3.21.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = \int \frac{Bx^2 + A}{(ex^2 + d)\sqrt{cx^4 + bx^2 + a}} dx$$

input `int((A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)), x)`output `int((A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)), x)`

### 3.22 $\int \frac{A+Bx^2}{(d+ex^2)^2 \sqrt{a+bx^2+cx^4}} dx$

|        |                            |     |
|--------|----------------------------|-----|
| 3.22.1 | Optimal result             | 189 |
| 3.22.2 | Mathematica [C] (verified) | 190 |
| 3.22.3 | Rubi [A] (verified)        | 191 |
| 3.22.4 | Maple [B] (verified)       | 195 |
| 3.22.5 | Fricas [F]                 | 196 |
| 3.22.6 | Sympy [F]                  | 197 |
| 3.22.7 | Maxima [F]                 | 197 |
| 3.22.8 | Giac [F]                   | 197 |
| 3.22.9 | Mupad [F(-1)]              | 198 |

#### 3.22.1 Optimal result

Integrand size = 33, antiderivative size = 782

$$\int \frac{A+Bx^2}{(d+ex^2)^2 \sqrt{a+bx^2+cx^4}} dx$$

$$= \frac{\sqrt{c}(Bd - Ae)x\sqrt{a+bx^2+cx^4}}{2d(cd^2 - bde + ae^2)(\sqrt{a} + \sqrt{cx^2})} - \frac{e(Bd - Ae)x\sqrt{a+bx^2+cx^4}}{2d(cd^2 - bde + ae^2)(d + ex^2)}$$

$$- \frac{(B(cd^3 - ade^2) - Ae(3cd^2 - e(2bd - ae))) \arctan\left(\frac{\sqrt{cd^2 - bde + ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{a+bx^2+cx^4}}\right)}{4d^{3/2}\sqrt{e}(cd^2 - bde + ae^2)^{3/2}}$$

$$- \frac{\sqrt[4]{a}\sqrt[4]{c}(Bd - Ae)(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right) \middle| \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{2d(cd^2 - bde + ae^2)\sqrt{a+bx^2+cx^4}}$$

$$+ \frac{A\sqrt[4]{c}(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right), \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{2\sqrt[4]{ad}(\sqrt{cd} - \sqrt{ae})\sqrt{a+bx^2+cx^4}}$$

$$+ \frac{(\sqrt{cd} + \sqrt{ae})(B(cd^3 - ade^2) - Ae(3cd^2 - e(2bd - ae))) (\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \text{EllipticPi}\left(-\frac{\sqrt{cd}}{4\sqrt{a}}\right)}{8\sqrt[4]{a}\sqrt[4]{cd^2}e(\sqrt{cd} - \sqrt{ae})(cd^2 - bde + ae^2)\sqrt{a+bx^2+cx^4}}$$

output

```

-1/4*(B*(-a*d*e^2+c*d^3)-A*e*(3*c*d^2-e*(-a*e+2*b*d)))*arctan(x*(a*e^2-b*d
*e+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+b*x^2+a)^(1/2))/d^(3/2)/(a*e^2-b*d*
e+c*d^2)^(3/2)/e^(1/2)-1/2*e*(-A*e+B*d)*x*(c*x^4+b*x^2+a)^(1/2)/d/(a*e^2-b
*d*e+c*d^2)/(e*x^2+d)+1/2*(-A*e+B*d)*x*c^(1/2)*(c*x^4+b*x^2+a)^(1/2)/d/(a
e^2-b*d*e+c*d^2)/(a^(1/2)+x^2*c^(1/2))-1/2*a^(1/4)*c^(1/4)*(-A*e+B*d)*(cos
(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*El
lipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*
(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/d/(a
*e^2-b*d*e+c*d^2)/(c*x^4+b*x^2+a)^(1/2)+1/2*A*c^(1/4)*(cos(2*arctan(c^(1/4)
)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*ar
ctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(
1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(1/4)/d/(-e*a^(1/2)
+d*c^(1/2))/(c*x^4+b*x^2+a)^(1/2)+1/8*(B*(-a*d*e^2+c*d^3)-A*e*(3*c*d^2-e
(-a*e+2*b*d)))*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(
1/4)*x/a^(1/4)))*EllipticPi(sin(2*arctan(c^(1/4)*x/a^(1/4))),-1/4*(-e*a^(1
/2)+d*c^(1/2))^2/d/e/a^(1/2)/c^(1/2),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(e*a
^(1/2)+d*c^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1
/2)))^(1/2)/a^(1/4)/c^(1/4)/d^2/e/(a*e^2-b*d*e+c*d^2)/(-e*a^(1/2)+d*c^(1
/2))/(c*x^4+b*x^2+a)^(1/2)

```

### 3.22.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 12.38 (sec) , antiderivative size = 1853, normalized size of antiderivative = 2.37

$$\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + bx^2 + cx^4}} dx = \text{Too large to display}$$

input `Integrate[(A + B*x^2)/((d + e*x^2)^2*Sqrt[a + b*x^2 + c*x^4]),x]`

output  $((-1/8*I)*((-4*I)*\text{Sqrt}[c/(b + \text{Sqrt}[b^2 - 4*a*c])])*d*e^2*(B*d - A*e)*x*(a + b*x^2 + c*x^4) + \text{Sqrt}[2]*B*(b - \text{Sqrt}[b^2 - 4*a*c])*d^2*e*\text{Sqrt}[(b + \text{Sqrt}[b^2 - 4*a*c] + 2*c*x^2)/(b + \text{Sqrt}[b^2 - 4*a*c])]*\text{Sqrt}[1 + (2*c*x^2)/(b - \text{Sqrt}[b^2 - 4*a*c])]*(d + e*x^2)*(EllipticE[I*ArcSinh[\text{Sqrt}[2]*\text{Sqrt}[c/(b + \text{Sqrt}[b^2 - 4*a*c])]]*x], (b + \text{Sqrt}[b^2 - 4*a*c])/(b - \text{Sqrt}[b^2 - 4*a*c])) - \text{EllipticF}[I*ArcSinh[\text{Sqrt}[2]*\text{Sqrt}[c/(b + \text{Sqrt}[b^2 - 4*a*c])]]*x], (b + \text{Sqrt}[b^2 - 4*a*c])/(b - \text{Sqrt}[b^2 - 4*a*c])) + \text{Sqrt}[2]*A*(-b + \text{Sqrt}[b^2 - 4*a*c])*d*e^2*\text{Sqrt}[(b + \text{Sqrt}[b^2 - 4*a*c] + 2*c*x^2)/(b + \text{Sqrt}[b^2 - 4*a*c])]*\text{Sqrt}[1 + (2*c*x^2)/(b - \text{Sqrt}[b^2 - 4*a*c])]*(d + e*x^2)*(EllipticE[I*ArcSinh[\text{Sqrt}[2]*\text{Sqrt}[c/(b + \text{Sqrt}[b^2 - 4*a*c])]]*x], (b + \text{Sqrt}[b^2 - 4*a*c])/(b - \text{Sqrt}[b^2 - 4*a*c])) - \text{EllipticF}[I*ArcSinh[\text{Sqrt}[2]*\text{Sqrt}[c/(b + \text{Sqrt}[b^2 - 4*a*c])]]*x], (b + \text{Sqrt}[b^2 - 4*a*c])/(b - \text{Sqrt}[b^2 - 4*a*c])) + 2*\text{Sqrt}[2]*B*c*d^3*\text{Sqrt}[(b + \text{Sqrt}[b^2 - 4*a*c] + 2*c*x^2)/(b + \text{Sqrt}[b^2 - 4*a*c])]*\text{Sqrt}[1 + (2*c*x^2)/(b - \text{Sqrt}[b^2 - 4*a*c])]*(d + e*x^2)*\text{EllipticF}[I*ArcSinh[\text{Sqrt}[2]*\text{Sqrt}[c/(b + \text{Sqrt}[b^2 - 4*a*c])]]*x], (b + \text{Sqrt}[b^2 - 4*a*c])/(b - \text{Sqrt}[b^2 - 4*a*c])) - 2*\text{Sqrt}[2]*A*c*d^2*e*\text{Sqrt}[(b + \text{Sqrt}[b^2 - 4*a*c] + 2*c*x^2)/(b + \text{Sqrt}[b^2 - 4*a*c])]*\text{Sqrt}[1 + (2*c*x^2)/(b - \text{Sqrt}[b^2 - 4*a*c])]*(d + e*x^2)*\text{EllipticF}[I*ArcSinh[\text{Sqrt}[2]*\text{Sqrt}[c/(b + \text{Sqrt}[b^2 - 4*a*c])]]*x], (b + \text{Sqrt}[b^2 - 4*a*c])/(b - \text{Sqrt}[b^2 - 4*a*c])) - 2*\text{Sqrt}[2]*B*c*d^3*\text{Sqrt}[(b + \text{Sqrt}[b^2 - 4*a*c] + 2*c*x^2)/(b + \text{Sqrt}[b^2 - 4*a*c])]*\text{Sqrt}[1 + (...$

### 3.22.3 Rubi [A] (verified)

Time = 1.59 (sec) , antiderivative size = 727, normalized size of antiderivative = 0.93, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {2210, 25, 2232, 27, 1509, 2226, 27, 1416, 2220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + bx^2 + cx^4}} dx$$

↓ 2210

$$-\frac{\int -\frac{ce(Bd-Ae)x^4 + 2cd(Bd-Ae)x^2 + aBde + A(2cd^2 - e(2bd-ae))}{(ex^2+d)\sqrt{cx^4+bx^2+a}} dx}{2d(ae^2 - bde + cd^2)} - \frac{ex\sqrt{a + bx^2 + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 - bde + cd^2)}$$

↓ 25

$$\frac{\int \frac{ce(Bd-Ae)x^4 + 2cd(Bd-Ae)x^2 + 2Acd^2 + aBde - Ae(2bd-ae)}{(ex^2+d)\sqrt{cx^4+bx^2+a}} dx}{2d(ae^2 - bde + cd^2)} - \frac{ex\sqrt{a + bx^2 + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 - bde + cd^2)}$$

---

3.22.  $\int \frac{A+Bx^2}{(d+ex^2)^2\sqrt{a+bx^2+cx^4}} dx$



$$\int \frac{ce(\sqrt{c}(\sqrt{cd}+\sqrt{ae})(Bd-Ae)x^2+\sqrt{a}\sqrt{cd}(Bd-Ae)+ae(Bd+ Ae)+2Ad(cd-be))}{(ex^2+d)\sqrt{cx^4+bx^2+a}} dx - \sqrt{a}\sqrt{c}(Bd-Ae) \int \frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{a}\sqrt{cx^4+bx^2+a}} dx$$

$$\frac{2d(ae^2 - bde + cd^2)}{ex\sqrt{a + bx^2 + cx^4}(Bd - Ae)} - \frac{2d(d + ex^2)(ae^2 - bde + cd^2)}{2d(d + ex^2)(ae^2 - bde + cd^2)}$$

27

$$\int \frac{\sqrt{c}(\sqrt{cd}+\sqrt{ae})(Bd-Ae)x^2+\sqrt{a}\sqrt{cd}(Bd-Ae)+ae(Bd+ Ae)+2Ad(cd-be)}{(ex^2+d)\sqrt{cx^4+bx^2+a}} dx - \sqrt{c}(Bd - Ae) \int \frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{cx^4+bx^2+a}} dx$$

$$\frac{2d(ae^2 - bde + cd^2)}{ex\sqrt{a + bx^2 + cx^4}(Bd - Ae)} - \frac{2d(d + ex^2)(ae^2 - bde + cd^2)}{2d(d + ex^2)(ae^2 - bde + cd^2)}$$

1509

$$\int \frac{\sqrt{c}(\sqrt{cd}+\sqrt{ae})(Bd-Ae)x^2+\sqrt{a}\sqrt{cd}(Bd-Ae)+ae(Bd+ Ae)+2Ad(cd-be)}{(ex^2+d)\sqrt{cx^4+bx^2+a}} dx - \sqrt{c}(Bd - Ae) \left( \frac{\sqrt[4]{a}(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt{a}+\sqrt{cx^2}}{\sqrt{a}}\right)\right)}{\sqrt[4]{c}\sqrt{a+bx^2+cx^4}} \right)$$

$$\frac{2d(ae^2 - bde + cd^2)}{ex\sqrt{a + bx^2 + cx^4}(Bd - Ae)} - \frac{2d(d + ex^2)(ae^2 - bde + cd^2)}{2d(d + ex^2)(ae^2 - bde + cd^2)}$$

2226

$$\frac{\sqrt{a}(B(cd^3-ade^2)-Ae(3cd^2-e(2bd-ae)))}{\sqrt{cd}-\sqrt{ae}} \int \frac{\sqrt{cx^2}+\sqrt{a}}{\sqrt{a}(ex^2+d)\sqrt{cx^4+bx^2+a}} dx + \frac{2A\sqrt{c}(ae^2-bde+cd^2)}{\sqrt{cd}-\sqrt{ae}} \int \frac{1}{\sqrt{cx^4+bx^2+a}} dx - \sqrt{c}(Bd - Ae) \left( \frac{\sqrt[4]{a}(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt{a}+\sqrt{cx^2}}{\sqrt{a}}\right)\right)}{\sqrt[4]{c}\sqrt{a+bx^2+cx^4}} \right)$$

$$\frac{2d(ae^2 - bde + cd^2)}{ex\sqrt{a + bx^2 + cx^4}(Bd - Ae)} - \frac{2d(d + ex^2)(ae^2 - bde + cd^2)}{2d(d + ex^2)(ae^2 - bde + cd^2)}$$

27

$$\frac{(B(cd^3-ade^2)-Ae(3cd^2-e(2bd-ae)))}{\sqrt{cd}-\sqrt{ae}} \int \frac{\sqrt{cx^2}+\sqrt{a}}{(ex^2+d)\sqrt{cx^4+bx^2+a}} dx + \frac{2A\sqrt{c}(ae^2-bde+cd^2)}{\sqrt{cd}-\sqrt{ae}} \int \frac{1}{\sqrt{cx^4+bx^2+a}} dx - \sqrt{c}(Bd - Ae) \left( \frac{\sqrt[4]{a}(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt{a}+\sqrt{cx^2}}{\sqrt{a}}\right)\right)}{\sqrt[4]{c}\sqrt{a+bx^2+cx^4}} \right)$$

$$\frac{2d(ae^2 - bde + cd^2)}{ex\sqrt{a + bx^2 + cx^4}(Bd - Ae)} - \frac{2d(d + ex^2)(ae^2 - bde + cd^2)}{2d(d + ex^2)(ae^2 - bde + cd^2)}$$

3.22.  $\int \frac{A+Bx^2}{(d+ex^2)^2\sqrt{a+bx^2+cx^4}} dx$

↓ 1416

$$\frac{(B(cd^3 - ade^2) - Ae(3cd^2 - e(2bd - ae))) \int \frac{\sqrt{cx^2 + \sqrt{a}}}{(ex^2 + d)\sqrt{cx^4 + bx^2 + a}} dx}{\sqrt{cd} - \sqrt{ae}} - \sqrt{c}(Bd - Ae) \left( \frac{\sqrt[4]{a}(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} E \left( 2 \arctan \left( \frac{\sqrt[4]{cx}}{\sqrt[4]{a}} \right) \right)}{\sqrt[4]{c}\sqrt{a + bx^2 + cx^4}} \right)$$

$2d(ae^2 - bde)$

$$\frac{ex\sqrt{a + bx^2 + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 - bde + cd^2)}$$

↓ 2220

$$\frac{(B(cd^3 - ade^2) - Ae(3cd^2 - e(2bd - ae))) \left( \frac{(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} (\sqrt{ae} + \sqrt{cd}) \operatorname{EllipticPi} \left( -\frac{\sqrt{a} \left( \frac{\sqrt{cd}}{\sqrt{a}} - e \right)^2}{4\sqrt{cde}}, 2 \arctan \left( \frac{\sqrt[4]{cx}}{\sqrt[4]{a}} \right), \frac{1}{4} \left( 2 - \frac{b}{\sqrt{a}\sqrt{c}} \right) \right)}{4\sqrt[4]{a}\sqrt[4]{cde}\sqrt{a + bx^2 + cx^4}} \right)}{\sqrt{cd} - \sqrt{ae}}$$

$$\frac{ex\sqrt{a + bx^2 + cx^4}(Bd - Ae)}{2d(d + ex^2)(ae^2 - bde + cd^2)}$$

input `Int[(A + B*x^2)/((d + e*x^2)^2*Sqrt[a + b*x^2 + c*x^4]),x]`

output

```
-1/2*(e*(B*d - A*e)*x*Sqrt[a + b*x^2 + c*x^4])/(d*(c*d^2 - b*d*e + a*e^2)*(d + e*x^2)) + (- (Sqrt[c]*(B*d - A*e)*(-(x*Sqrt[a + b*x^2 + c*x^4])/(Sqrt[a] + Sqrt[c]*x^2)) + (a^(1/4)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(c^(1/4)*Sqrt[a + b*x^2 + c*x^4]))) + (A*c^(1/4)*(c*d^2 - b*d*e + a*e^2)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(a^(1/4)*(Sqrt[c]*d - Sqrt[a]*e)*Sqrt[a + b*x^2 + c*x^4]) + ((B*(c*d^3 - a*d*e^2) - A*e*(3*c*d^2 - e*(2*b*d - a*e)))*(-1/2*((Sqrt[c]*d - Sqrt[a]*e)*ArcTan[(Sqrt[c*d^2 - b*d*e + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + b*x^2 + c*x^4])])/(Sqrt[d]*Sqrt[e]*Sqrt[c*d^2 - b*d*e + a*e^2]) + ((Sqrt[c]*d + Sqrt[a]*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-1/4*(Sqrt[a]*((Sqrt[c]*d)/Sqrt[a] - e)^2/(Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(4*a^(1/4)*c^(1/4)*d*e*Sqrt[a + b*x^2 + c*x^4]))/(Sqrt[c]*d - Sqrt[a]*e))/(2*d*(c*d^2 - b*d*e + a*e^2))
```

## 3.22.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 1416 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`
- rule 1509 `Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`
- rule 2210 `Int[((P4x_)*((d_) + (e_.)*(x_)^2)^(q_))/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]}, Simp[(-(C*d^2 - B*d*e + A*e^2))*x*(d + e*x^2)^(q + 1)*(Sqrt[a + b*x^2 + c*x^4]/(2*d*(q + 1)*(c*d^2 - b*d*e + a*e^2))), x] + Simp[1/(2*d*(q + 1)*(c*d^2 - b*d*e + a*e^2)) Int[((d + e*x^2)^(q + 1)/Sqrt[a + b*x^2 + c*x^4])*Simp[a*d*(C*d - B*e) + A*(a*e^2*(2*q + 3) + 2*d*(c*d - b*e)*(q + 1)) - 2*((B*d - A*e)*(b*e*(q + 2) - c*d*(q + 1)) - C*d*(b*d + a*e*(q + 1)))*x^2 + c*(C*d^2 - B*d*e + A*e^2)*(2*q + 5)*x^4, x], x]] /; FreeQ[{a, b, c, d, e}, x] && PolyQ[P4x, x^2] && LeQ[Expon[P4x, x], 4] && ILtQ[q, -1]`

```
rule 2220 Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 +
(c_)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(A
rcTan[Rt[-b + c*(d/e) + a*(e/d), 2]*(x/Sqrt[a + b*x^2 + c*x^4])]/(2*d*e*Rt[
-b + c*(d/e) + a*(e/d), 2]))], x] + Simp[(B*d + A*e)*(1 + q^2*x^2)*(Sqrt[(a
+ b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(4*d*e*q*Sqrt[a + b*x^2 + c*x^4]))*El
lipticPi[-(e - d*q^2)^2/(4*d*e*q^2), 2*ArcTan[q*x], 1/2 - b/(4*a*q^2)], x]]
/; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] &
& EqQ[c*A^2 - a*B^2, 0] && PosQ[B/A] && PosQ[-b + c*(d/e) + a*(e/d)]
```

```
rule 2226 Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 +
(c_)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2]}, Simp[(A*(c*d + a*e*q)
- a*B*(e + d*q))/(c*d^2 - a*e^2) Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] +
Simp[a*(B*d - A*e)*((e + d*q)/(c*d^2 - a*e^2)) Int[(1 + q*x^2)/((d + e*x^
2)*Sqrt[a + b*x^2 + c*x^4]), x], x]] /; FreeQ[{a, b, c, d, e, A, B}, x] &&
NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]
```

```
rule 2232 Int[(P4x_)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4])
, x_Symbol] := With[{q = Rt[c/a, 2], A = Coeff[P4x, x, 0], B = Coeff[P4x, x
, 2], C = Coeff[P4x, x, 4]}, Simp[-C/(e*q) Int[(1 - q*x^2)/Sqrt[a + b*x^2
+ c*x^4], x], x] + Simp[1/(c*e) Int[(A*c*e + a*C*d*q + (B*c*e - C*(c*d -
a*e*q))*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x]] /; FreeQ[{a, b
, c, d, e}, x] && PolyQ[P4x, x^2, 2] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]
&& !GtQ[b^2 - 4*a*c, 0]
```

### 3.22.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1494 vs.  $2(745) = 1490$ .

Time = 3.17 (sec) , antiderivative size = 1495, normalized size of antiderivative = 1.91

| method   | result                          | size |
|----------|---------------------------------|------|
| default  | Expression too large to display | 1495 |
| elliptic | Expression too large to display | 2301 |

```
input int((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURNVERBOSE)
```

$$3.22. \int \frac{A+Bx^2}{(d+ex^2)^2\sqrt{a+bx^2+cx^4}} dx$$

output  $B/e/d*2^{(1/2)}/(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(1+1/2*b*x^2/a+1/2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticPi(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},-2/(-b+(-4*a*c+b^2)^{(1/2)})*a*e/d,(-1/2*(b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}+(A*e-B*d)/e*(1/2*e^2*x*(c*x^4+b*x^2+a)^{(1/2)}/d/(a*e^2-b*d*e+c*d^2)/(e*x^2+d)-1/8*c/(a*e^2-b*d*e+c*d^2)*2^{(1/2)}/(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a-2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)}+1/4*c*e/d/(a*e^2-b*d*e+c*d^2)*a*2^{(1/2)}/(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a-2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-1/4*c*e/d/(a*e^2-b*d*e+c*d^2)*a*2^{(1/2)}/(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a-2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*EllipticE(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})+1/2/(a*e^2-b*d*e+c*d^2)/d^2*e^2*2^{(1/2)}/(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}$

### 3.22.5 Fracas [F]

$$\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + bx^2 + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}(ex^2 + d)^2} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fracas")`

output `integral(sqrt(c*x^4 + b*x^2 + a)*(B*x^2 + A)/(c*e^2*x^8 + (2*c*d*e + b*e^2)*x^6 + (c*d^2 + 2*b*d*e + a*e^2)*x^4 + a*d^2 + (b*d^2 + 2*a*d*e)*x^2), x)`

## 3.22.6 Sympy [F]

$$\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + bx^2 + cx^4}} dx = \int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + bx^2 + cx^4}} dx$$

input `integrate((B*x**2+A)/(e*x**2+d)**2/(c*x**4+b*x**2+a)**(1/2),x)`

output `Integral((A + B*x**2)/((d + e*x**2)**2*sqrt(a + b*x**2 + c*x**4)), x)`

## 3.22.7 Maxima [F]

$$\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + bx^2 + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}(ex^2 + d)^2} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)^2), x)`

## 3.22.8 Giac [F]

$$\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + bx^2 + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}(ex^2 + d)^2} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)^2), x)`

**3.22.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + bx^2 + cx^4}} dx = \int \frac{Bx^2 + A}{(ex^2 + d)^2 \sqrt{cx^4 + bx^2 + a}} dx$$

input `int((A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(1/2)), x)`output `int((A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(1/2)), x)`

### 3.23 $\int \frac{A+Bx^2}{(d+ex^2)^3 \sqrt{a+bx^2+cx^4}} dx$

|        |                                      |     |
|--------|--------------------------------------|-----|
| 3.23.1 | Optimal result . . . . .             | 199 |
| 3.23.2 | Mathematica [C] (verified) . . . . . | 200 |
| 3.23.3 | Rubi [A] (verified) . . . . .        | 201 |
| 3.23.4 | Maple [B] (verified) . . . . .       | 206 |
| 3.23.5 | Fricas [F(-1)] . . . . .             | 207 |
| 3.23.6 | Sympy [F] . . . . .                  | 207 |
| 3.23.7 | Maxima [F] . . . . .                 | 207 |
| 3.23.8 | Giac [F] . . . . .                   | 208 |
| 3.23.9 | Mupad [F(-1)] . . . . .              | 208 |

#### 3.23.1 Optimal result

Integrand size = 33, antiderivative size = 1125

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + bx^2 + cx^4}} dx$$

$$= -\frac{\sqrt{c}(3Ae(3cd^2 - e(2bd - ae)) - Bd(5cd^2 - e(2bd + ae))) x \sqrt{a + bx^2 + cx^4}}{8d^2 (cd^2 - bde + ae^2)^2 (\sqrt{a} + \sqrt{cx^2})}$$

$$- \frac{e(Bd - Ae)x \sqrt{a + bx^2 + cx^4}}{4d (cd^2 - bde + ae^2) (d + ex^2)^2}$$

$$+ \frac{e(3Ae(3cd^2 - e(2bd - ae)) - Bd(5cd^2 - e(2bd + ae))) x \sqrt{a + bx^2 + cx^4}}{8d^2 (cd^2 - bde + ae^2)^2 (d + ex^2)}$$

$$- \frac{(Bd(3c^2d^4 - 10acd^2e^2 + ae^3(4bd - ae)) - Ae(15c^2d^4 - 2cd^2e(10bd - 3ae)) + e^2(8b^2d^2 - 8abde + 3a^2e^2))}{16d^{5/2}\sqrt{e} (cd^2 - bde + ae^2)^{5/2}}$$

$$+ \frac{\sqrt[4]{a}\sqrt[4]{c}(3Ae(3cd^2 - e(2bd - ae)) - Bd(5cd^2 - e(2bd + ae))) (\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt{a+bx^2+cx^4}}{\sqrt{a}+\sqrt{cx^2}}\right)\right)}{8d^2 (cd^2 - bde + ae^2)^2 \sqrt{a + bx^2 + cx^4}}$$

$$+ \frac{\sqrt[4]{c}(\sqrt{a}\sqrt{cd}(Bd - Ae) + ae(Bd + 3Ae) + 4Ad(cd - be)) (\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt{a+bx^2+cx^4}}{\sqrt{a}+\sqrt{cx^2}}\right)\right)}{8\sqrt[4]{ad^2} (\sqrt{cd} - \sqrt{ae}) (cd^2 - bde + ae^2) \sqrt{a + bx^2 + cx^4}}$$

$$+ \frac{(\sqrt{cd} + \sqrt{ae}) (Bd(3c^2d^4 - 10acd^2e^2 + ae^3(4bd - ae)) - Ae(15c^2d^4 - 2cd^2e(10bd - 3ae)) + e^2(8b^2d^2 - 8abde + 3a^2e^2))}{32\sqrt[4]{a}\sqrt[4]{c}d^3e (\sqrt{cd} - \sqrt{ae}) (cd^2 - bde + ae^2)}$$



```
output -1/16*(B*d*(3*c^2*d^4-10*a*c*d^2*e^2+a*e^3*(-a*e+4*b*d))-A*e*(15*c^2*d^4-2
*c*d^2*e*(-3*a*e+10*b*d)+e^2*(3*a^2*e^2-8*a*b*d*e+8*b^2*d^2)))*arctan(x*(a
*e^2-b*d*e+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+b*x^2+a)^(1/2))/d^(5/2)/(a*
e^2-b*d*e+c*d^2)^(5/2)/e^(1/2)-1/4*e*(-A*e+B*d)*x*(c*x^4+b*x^2+a)^(1/2)/d/
(a*e^2-b*d*e+c*d^2)/(e*x^2+d)^2+1/8*e*(3*A*e*(3*c*d^2-e*(-a*e+2*b*d))-B*d*
(5*c*d^2-e*(a*e+2*b*d)))*x*(c*x^4+b*x^2+a)^(1/2)/d^2/(a*e^2-b*d*e+c*d^2)^2
/(e*x^2+d)-1/8*(3*A*e*(3*c*d^2-e*(-a*e+2*b*d))-B*d*(5*c*d^2-e*(a*e+2*b*d))
)*x*c^(1/2)*(c*x^4+b*x^2+a)^(1/2)/d^2/(a*e^2-b*d*e+c*d^2)^2/(a^(1/2)+x^2*c
^(1/2))+1/8*a^(1/4)*c^(1/4)*(3*A*e*(3*c*d^2-e*(-a*e+2*b*d))-B*d*(5*c*d^2-e
*(a*e+2*b*d)))*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(
1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/
2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(
1/2)))^(1/2)/d^2/(a*e^2-b*d*e+c*d^2)^2/(c*x^4+b*x^2+a)^(1/2)+1/32*(B*d*(
3*c^2*d^4-10*a*c*d^2*e^2+a*e^3*(-a*e+4*b*d))-A*e*(15*c^2*d^4-2*c*d^2*e*(-3
*a*e+10*b*d)+e^2*(3*a^2*e^2-8*a*b*d*e+8*b^2*d^2)))*(cos(2*arctan(c^(1/4)*x
/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticPi(sin(2*arct
an(c^(1/4)*x/a^(1/4))),-1/4*(-e*a^(1/2)+d*c^(1/2))^2/d/e/a^(1/2)/c^(1/2),1
/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(e*a^(1/2)+d*c^(1/2))*(a^(1/2)+x^2*c^(1/2
))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(1/4)/c^(1/4)/d^3/e/(a
*e^2-b*d*e+c*d^2)^2/(-e*a^(1/2)+d*c^(1/2))/(c*x^4+b*x^2+a)^(1/2)+1/8*c...
```

### 3.23.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 13.91 (sec) , antiderivative size = 781, normalized size of antiderivative = 0.69

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + bx^2 + cx^4}} dx$$

$$= \frac{-4de^2x(a+bx^2+cx^4)(2d(Bd-Ae)(cd^2+e(-bd+ae))+(-3Ae(3cd^2+e(-2bd+ae))+B(5cd^3-de(2bd+ae)))(d+ex^2))}{(d+ex^2)^2} - \frac{i\sqrt{2}\sqrt{\frac{b+\sqrt{b^2-4ac+2c}}{b+\sqrt{b^2-4ac}}}}{...}$$

```
input Integrate[(A + B*x^2)/((d + e*x^2)^3*Sqrt[a + b*x^2 + c*x^4]),x]
```

output  $((-4*d*e^2*x*(a + b*x^2 + c*x^4)*(2*d*(B*d - A*e)*(c*d^2 + e*(-(b*d) + a*e)) + (-3*A*e*(3*c*d^2 + e*(-2*b*d + a*e)) + B*(5*c*d^3 - d*e*(2*b*d + a*e)))*(d + e*x^2))/(d + e*x^2)^2 - (I*sqrt[2]*sqrt[(b + sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + sqrt[b^2 - 4*a*c]])*sqrt[1 + (2*c*x^2)/(b - sqrt[b^2 - 4*a*c]])]*((-b + sqrt[b^2 - 4*a*c])*d*e*(3*A*e*(3*c*d^2 + e*(-2*b*d + a*e)) + B*(-5*c*d^3 + d*e*(2*b*d + a*e)))*EllipticE[I*ArcSinh[sqrt[2]*sqrt[c/(b + sqrt[b^2 - 4*a*c])]]*x], (b + sqrt[b^2 - 4*a*c])/(b - sqrt[b^2 - 4*a*c])) + d*(B*d*(6*c^2*d^3 + c*d*e*(-5*b*d + 5*sqrt[b^2 - 4*a*c]*d - 6*a*e) - (-b + sqrt[b^2 - 4*a*c])*e^2*(2*b*d + a*e)) - A*e*(14*c^2*d^3 - 3*(-b + sqrt[b^2 - 4*a*c])*e^2*(2*b*d - a*e) + c*d*e*(-17*b*d + 9*sqrt[b^2 - 4*a*c]*d + 2*a*e))*EllipticF[I*ArcSinh[sqrt[2]*sqrt[c/(b + sqrt[b^2 - 4*a*c])]]*x], (b + sqrt[b^2 - 4*a*c])/(b - sqrt[b^2 - 4*a*c])) + 2*(B*(-3*c^2*d^5 + 10*a*c*d^3*e^2 + a*d*e^3*(-4*b*d + a*e)) + A*e*(15*c^2*d^4 + 2*c*d^2*e*(-10*b*d + 3*a*e) + e^2*(8*b^2*d^2 - 8*a*b*d*e + 3*a^2*e^2)))*EllipticPi[((b + sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[sqrt[2]*sqrt[c/(b + sqrt[b^2 - 4*a*c])]]*x], (b + sqrt[b^2 - 4*a*c])/(b - sqrt[b^2 - 4*a*c])))/sqrt[c/(b + sqrt[b^2 - 4*a*c])]/(32*d^3*e*(c*d^2 + e*(-(b*d) + a*e))^2*sqrt[a + b*x^2 + c*x^4])$

### 3.23.3 Rubi [A] (verified)

Time = 3.51 (sec) , antiderivative size = 985, normalized size of antiderivative = 0.88, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2210, 25, 2210, 25, 2232, 27, 1509, 2226, 27, 1416, 2220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + bx^2 + cx^4}} dx$$

↓ 2210

$$\frac{\int -\frac{ce(Bd - Ae)x^4 + 2(Bd - Ae)(2cd - be)x^2 + 4Acd^2 + aBde - Ae(4bd - 3ae)}{(ex^2 + d)^2 \sqrt{cx^4 + bx^2 + a}} dx}{4d(ae^2 - bde + cd^2)} - \frac{ex\sqrt{a + bx^2 + cx^4}(Bd - Ae)}{4d(d + ex^2)^2(ae^2 - bde + cd^2)}$$

↓ 25

$$\frac{\int -\frac{ce(Bd - Ae)x^4 + 2(Bd - Ae)(2cd - be)x^2 + 4Acd^2 + aBde - Ae(4bd - 3ae)}{(ex^2 + d)^2 \sqrt{cx^4 + bx^2 + a}} dx}{4d(ae^2 - bde + cd^2)} - \frac{ex\sqrt{a + bx^2 + cx^4}(Bd - Ae)}{4d(d + ex^2)^2(ae^2 - bde + cd^2)}$$

↓ 2210

---

3.23.  $\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + bx^2 + cx^4}} dx$

$$\frac{ex\sqrt{a+bx^2+cx^4}(3Ae(3cd^2-e(2bd-ae))-B(5cd^3-de(ae+2bd)))}{2d(d+ex^2)(ae^2-bde+cd^2)} - \int \frac{-ce(3Ae(3cd^2-e(2bd-ae))-B(5cd^3-de(2bd+ae)))x^4-2cd(Ae(8cd^2-e(5bd-ae)))}{4d(ae^2-bde+cd^2)}$$

$$\frac{ex\sqrt{a+bx^2+cx^4}(Bd-Ae)}{4d(d+ex^2)^2(ae^2-bde+cd^2)}$$

↓ 25

$$\int \frac{-ce(3Ae(3cd^2-e(2bd-ae))-B(5cd^3-de(2bd+ae)))x^4-2cd(Ae(8cd^2-e(5bd-2ae))-B(4cd^3-de(bd+2ae)))x^2+ade(Bd-Ae)(5cd-2be)+(4Acd^2+aBed-Ae)}{(ex^2+d)\sqrt{cx^4+bx^2+a}}}{2d(ae^2-bde+cd^2)}$$

$$\frac{ex\sqrt{a+bx^2+cx^4}(Bd-Ae)}{4d(d+ex^2)^2(ae^2-bde+cd^2)}$$

↓ 2232

$$\sqrt{a}\sqrt{c}(3Ae(3cd^2-e(2bd-ae))-B(5cd^3-de(ae+2bd))) \int \frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{a}\sqrt{cx^4+bx^2+a}} dx + \int \frac{ce((cd-\sqrt{a}\sqrt{ce})(3Ae(3cd^2-e(2bd-ae))-B(5cd^3-de(2bd+ae))))-2cd}{(ex^2+d)\sqrt{cx^4+bx^2+a}}$$

$$\frac{ex\sqrt{a+bx^2+cx^4}(Bd-Ae)}{4d(d+ex^2)^2(ae^2-bde+cd^2)}$$

↓ 27

$$\sqrt{c}(3Ae(3cd^2-e(2bd-ae))-B(5cd^3-de(ae+2bd))) \int \frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{a}\sqrt{cx^4+bx^2+a}} dx + \int \frac{((cd-\sqrt{a}\sqrt{ce})(3Ae(3cd^2-e(2bd-ae))-B(5cd^3-de(2bd+ae))))-2cd(Ae(8cd^2-e(5bd-ae)))}{(ex^2+d)\sqrt{cx^4+bx^2+a}}$$

$$\frac{ex\sqrt{a+bx^2+cx^4}(Bd-Ae)}{4d(d+ex^2)^2(ae^2-bde+cd^2)}$$

↓ 1509

$$\int \frac{((cd-\sqrt{a}\sqrt{ce})(3Ae(3cd^2-e(2bd-ae))-B(5cd^3-de(2bd+ae))))-2cd(Ae(8cd^2-e(5bd-2ae))-B(4cd^3-de(bd+2ae)))x^2+ade(Bd-Ae)(5cd-2be)+(4Acd^2+a)}{(ex^2+d)\sqrt{cx^4+bx^2+a}}$$

$$\frac{ex\sqrt{a+bx^2+cx^4}(Bd-Ae)}{4d(d+ex^2)^2(ae^2-bde+cd^2)}$$

↓ 2226

3.23.  $\int \frac{A+Bx^2}{(d+ex^2)^3\sqrt{a+bx^2+cx^4}} dx$

$$\frac{\sqrt{a} \left( B \left( ade^3(4bd-ae) - 10acd^3e^2 + 3c^2d^5 \right) - Ae \left( e^2(3a^2e^2 - 8abde + 8b^2d^2) - 2cd^2e(10bd - 3ae) + 15c^2d^4 \right) \right) \int \frac{\sqrt{cx^2 + \sqrt{a}}}{\sqrt{a}(ex^2 + d)\sqrt{cx^4 + bx^2 + a}} dx}{\sqrt{cd} - \sqrt{ae}} + \frac{2\sqrt{c}(ae^2 - bde + cd^2)(\sqrt{a}\sqrt{cd})}{\sqrt{cd} - \sqrt{ae}}$$

$$\frac{ex\sqrt{a + bx^2 + cx^4}(Bd - Ae)}{4d(d + ex^2)^2(ae^2 - bde + cd^2)}$$

↓ 27

$$\frac{\left( B \left( ade^3(4bd-ae) - 10acd^3e^2 + 3c^2d^5 \right) - Ae \left( e^2(3a^2e^2 - 8abde + 8b^2d^2) - 2cd^2e(10bd - 3ae) + 15c^2d^4 \right) \right) \int \frac{\sqrt{cx^2 + \sqrt{a}}}{(ex^2 + d)\sqrt{cx^4 + bx^2 + a}} dx}{\sqrt{cd} - \sqrt{ae}} + \frac{2\sqrt{c}(ae^2 - bde + cd^2)(\sqrt{a}\sqrt{cd})}{\sqrt{cd} - \sqrt{ae}}$$

$$\frac{ex\sqrt{a + bx^2 + cx^4}(Bd - Ae)}{4d(d + ex^2)^2(ae^2 - bde + cd^2)}$$

↓ 1416

$$\frac{\left( B \left( ade^3(4bd-ae) - 10acd^3e^2 + 3c^2d^5 \right) - Ae \left( e^2(3a^2e^2 - 8abde + 8b^2d^2) - 2cd^2e(10bd - 3ae) + 15c^2d^4 \right) \right) \int \frac{\sqrt{cx^2 + \sqrt{a}}}{(ex^2 + d)\sqrt{cx^4 + bx^2 + a}} dx}{\sqrt{cd} - \sqrt{ae}} + \frac{{}^4\sqrt{c}(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}}}{\sqrt{cd} - \sqrt{ae}}$$

$$\frac{ex\sqrt{a + bx^2 + cx^4}(Bd - Ae)}{4d(d + ex^2)^2(ae^2 - bde + cd^2)}$$

↓ 2220

$$\frac{e(3Ae(3cd^2 - e(2bd - ae)) - B(5cd^3 - de(2bd + ae)))\sqrt{cx^4 + bx^2 + ax}}{2d(cd^2 - bed + ae^2)(ex^2 + d)} + \frac{\sqrt{c}(3Ae(3cd^2 - e(2bd - ae)) - B(5cd^3 - de(2bd + ae))) \left( \frac{{}^4\sqrt{a}(\sqrt{cx^2 + \sqrt{a}})}{\sqrt{cx^4 + bx^2 + ax}} \sqrt{\frac{cx^4 + bx^2 + ax}{(\sqrt{a} + \sqrt{cx^2})^2}} \right)}{2d(cd^2 - bed + ae^2)(ex^2 + d)}$$

$$\frac{e(Bd - Ae)x\sqrt{cx^4 + bx^2 + a}}{4d(cd^2 - bed + ae^2)(ex^2 + d)^2}$$

3.23.  $\int \frac{A+Bx^2}{(d+ex^2)^3\sqrt{a+bx^2+cx^4}} dx$

input `Int[(A + B*x^2)/((d + e*x^2)^3*Sqrt[a + b*x^2 + c*x^4]),x]`

output `-1/4*(e*(B*d - A*e)*x*Sqrt[a + b*x^2 + c*x^4])/(d*(c*d^2 - b*d*e + a*e^2)*(d + e*x^2)^2) + ((e*(3*A*e*(3*c*d^2 - e*(2*b*d - a*e)) - B*(5*c*d^3 - d*e*(2*b*d + a*e)))*x*Sqrt[a + b*x^2 + c*x^4])/(2*d*(c*d^2 - b*d*e + a*e^2)*(d + e*x^2) + (Sqrt[c]*(3*A*e*(3*c*d^2 - e*(2*b*d - a*e)) - B*(5*c*d^3 - d*e*(2*b*d + a*e)))*(-(x*Sqrt[a + b*x^2 + c*x^4])/(Sqrt[a] + Sqrt[c]*x^2)) + (a^(1/4)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(c^(1/4)*Sqrt[a + b*x^2 + c*x^4])) + (c^(1/4)*(c*d^2 - b*d*e + a*e^2)*(Sqrt[a]*Sqrt[c]*d*(B*d - A*e) + a*e*(B*d + 3*A*e) + 4*A*d*(c*d - b*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(a^(1/4)*(Sqrt[c]*d - Sqrt[a]*e)*Sqrt[a + b*x^2 + c*x^4]) + ((B*(3*c^2*d^5 - 10*a*c*d^3*e^2 + a*d*e^3*(4*b*d - a*e)) - A*e*(15*c^2*d^4 - 2*c*d^2*e*(10*b*d - 3*a*e) + e^2*(8*b^2*d^2 - 8*a*b*d*e + 3*a^2*e^2)))*(-1/2*(Sqrt[c]*d - Sqrt[a]*e)*ArcTan[(Sqrt[c*d^2 - b*d*e + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + b*x^2 + c*x^4])])/(Sqrt[d]*Sqrt[e]*Sqrt[c*d^2 - b*d*e + a*e^2]) + ((Sqrt[c]*d + Sqrt[a]*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-1/4*(Sqrt[a]*((Sqrt[c]*d)/Sqrt[a] - e)^2)/(Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(4*a^(1/4)*c^(1/4)*d*e*Sqrt[a + b*x^2 + c*x^4]))/(Sqrt[c]...`

### 3.23.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 1416 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`

rule 1509 `Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`

rule 2210 `Int[((P4x_)*((d_) + (e_)*(x_)^2)^(q_))/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]}, Simp[(-(C*d^2 - B*d*e + A*e^2))*x*(d + e*x^2)^(q + 1)*(Sqrt[a + b*x^2 + c*x^4]/(2*d*(q + 1)*(c*d^2 - b*d*e + a*e^2))), x] + Simp[1/(2*d*(q + 1)*(c*d^2 - b*d*e + a*e^2)) Int[((d + e*x^2)^(q + 1)/Sqrt[a + b*x^2 + c*x^4])*Simp[a*d*(C*d - B*e) + A*(a*e^2*(2*q + 3) + 2*d*(c*d - b*e)*(q + 1)) - 2*((B*d - A*e)*(b*e*(q + 2) - c*d*(q + 1)) - C*d*(b*d + a*e*(q + 1)))*x^2 + c*(C*d^2 - B*d*e + A*e^2)*(2*q + 5)*x^4, x], x] /; FreeQ[{a, b, c, d, e}, x] && PolyQ[P4x, x^2] && LeQ[Expon[P4x, x], 4] && ILtQ[q, -1]`

rule 2220 `Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[-b + c*(d/e) + a*(e/d), 2]*(x/Sqrt[a + b*x^2 + c*x^4])]/(2*d*e*Rt[-b + c*(d/e) + a*(e/d), 2])), x] + Simp[(B*d + A*e)*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(4*d*e*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticPi[-(e - d*q^2)^2/(4*d*e*q^2), 2*ArcTan[q*x], 1/2 - b/(4*a*q^2)], x] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0] && PosQ[B/A] && PosQ[-b + c*(d/e) + a*(e/d)]`

rule 2226 `Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2]}, Simp[(A*(c*d + a*e*q) - a*B*(e + d*q))/(c*d^2 - a*e^2) Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] + Simp[a*(B*d - A*e)*((e + d*q)/(c*d^2 - a*e^2)) Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]`

```
rule 2232 Int[(P4x_)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4])
, x_Symbol] :> With[{q = Rt[c/a, 2], A = Coeff[P4x, x, 0], B = Coeff[P4x, x
, 2], C = Coeff[P4x, x, 4]}, Simp[-C/(e*q) Int[(1 - q*x^2)/Sqrt[a + b*x^2
+ c*x^4], x], x] + Simp[1/(c*e) Int[(A*c*e + a*C*d*q + (B*c*e - C*(c*d -
a*e*q))*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x]] /; FreeQ[{a, b
, c, d, e}, x] && PolyQ[P4x, x^2, 2] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]
&& !GtQ[b^2 - 4*a*c, 0]
```

### 3.23.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4475 vs. 2(1080) = 2160.

Time = 4.71 (sec) , antiderivative size = 4476, normalized size of antiderivative = 3.98

| method   | result                          | size |
|----------|---------------------------------|------|
| default  | Expression too large to display | 4476 |
| elliptic | Expression too large to display | 5671 |

```
input int((B*x^2+A)/(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURNVERBOSE)
```

```
output B/e*(1/2*e^2*x*(c*x^4+b*x^2+a)^(1/2)/d/(a*e^2-b*d*e+c*d^2)/(e*x^2+d)-1/8*c
/(a*e^2-b*d*e+c*d^2)*2^(1/2)/(-b/a+1/a*(-4*a*c+b^2)^(1/2))^(1/2)*(4+2*b*x^
2/a-2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2)*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^(1
/2))^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)
^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))+1/4*c*e/d/
(a*e^2-b*d*e+c*d^2)*a*2^(1/2)/(-b/a+1/a*(-4*a*c+b^2)^(1/2))^(1/2)*(4+2*b*x
^2/a-2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2)*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^(
1/2))^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*EllipticF(1/2*x*2
^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2)
)/a/c)^(1/2))-1/4*c*e/d/(a*e^2-b*d*e+c*d^2)*a*2^(1/2)/(-b/a+1/a*(-4*a*c+b^
2)^(1/2))^(1/2)*(4+2*b*x^2/a-2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2)*(4+2*b*x^2/
a+2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(
1/2))*EllipticE(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2
*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))+1/2/(a*e^2-b*d*e+c*d^2)/d^2*e^2*2^(1
/2)/(-b/a+1/a*(-4*a*c+b^2)^(1/2))^(1/2)*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c+b
^2)^(1/2))^(1/2)*(1+1/2*b*x^2/a+1/2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2)/(c*x^4
+b*x^2+a)^(1/2)*EllipticPi(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)
,-2/(-b+(-4*a*c+b^2)^(1/2))*a*e/d,(-1/2*(b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*2^
(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2))*a-1/(a*e^2-b*d*e+c*d^2)*e/d*2^(1/
2)/(-b/a+1/a*(-4*a*c+b^2)^(1/2))^(1/2)*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c...
```

$$3.23. \int \frac{A+Bx^2}{(d+ex^2)^3\sqrt{a+bx^2+cx^4}} dx$$

**3.23.5 Fracas [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + bx^2 + cx^4}} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fracas")`

output `Timed out`

**3.23.6 Sympy [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + bx^2 + cx^4}} dx = \int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + bx^2 + cx^4}} dx$$

input `integrate((B*x**2+A)/(e*x**2+d)**3/(c*x**4+b*x**2+a)**(1/2),x)`

output `Integral((A + B*x**2)/((d + e*x**2)**3*sqrt(a + b*x**2 + c*x**4)), x)`

**3.23.7 Maxima [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + bx^2 + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}(ex^2 + d)^3} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)^3), x)`



**3.23.8 Giac [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + bx^2 + cx^4}} dx = \int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}(ex^2 + d)^3} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)^3), x)`

**3.23.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + bx^2 + cx^4}} dx = \int \frac{Bx^2 + A}{(ex^2 + d)^3 \sqrt{cx^4 + bx^2 + a}} dx$$

input `int((A + B*x^2)/((d + e*x^2)^3*(a + b*x^2 + c*x^4)^(1/2)),x)`

output `int((A + B*x^2)/((d + e*x^2)^3*(a + b*x^2 + c*x^4)^(1/2)), x)`

**3.24** 
$$\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+bx^2+cx^4)^{3/2}} dx$$

3.24.1 Optimal result . . . . . 209  
 3.24.2 Mathematica [C] (verified) . . . . . 210  
 3.24.3 Rubi [A] (verified) . . . . . 211  
 3.24.4 Maple [A] (verified) . . . . . 215  
 3.24.5 Fricas [B] (verification not implemented) . . . . . 215  
 3.24.6 Sympy [F] . . . . . 216  
 3.24.7 Maxima [F] . . . . . 217  
 3.24.8 Giac [F] . . . . . 217  
 3.24.9 Mupad [F(-1)] . . . . . 217

**3.24.1 Optimal result**

Integrand size = 33, antiderivative size = 859

$$\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+bx^2+cx^4)^{3/2}} dx = \frac{x(Ac(b^2cd^3 - 2acd(cd^2 - 3ae^2) - abe(3cd^2 + ae^2)) + aB(ab^2e^3 + 2ace(3cd^2 + ae^2))}{(a+bx^2+cx^4)^{3/2}} + \frac{Be^3x\sqrt{a+bx^2+cx^4}}{3c^2} + \frac{(aB(6c^3d^3 - 8b^3e^3 - 9c^2de(bd + 6ae)) + bce^2(18bd + 29ae)) + 3Ac(2ab^2e^3 + 6ace(cd^2 - ae^2) - bcd(cd^2 + ae^2))}{3ac^{5/2}(b^2 - 4ac)(\sqrt{a} + \sqrt{cx^2})} - \frac{(aB(6c^3d^3 - 8b^3e^3 - 9c^2de(bd + 6ae)) + bce^2(18bd + 29ae)) + 3Ac(2ab^2e^3 + 6ace(cd^2 - ae^2) - bcd(cd^2 + ae^2))}{3a^{3/4}c^{11/4}(b^2 - 4ac)\sqrt{a+bx^2+cx^4}} - \frac{(3Ac^3d^3 - 5a^2Bce^3 - 3\sqrt{ac}^{5/2}d^2(Bd + 3Ae) + ae(3cd - 2be)(3Bcd - 4bBe + 3Ace) + 3a^{3/2}\sqrt{ce^2}(9Bcd - 4bBe + 3Ace))}{6a^{3/4}(b - 2\sqrt{a}\sqrt{c})c^{11/4}\sqrt{a}}$$

---

3.24. 
$$\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+bx^2+cx^4)^{3/2}} dx$$

output

```
x*(A*c*(b^2*c*d^3-2*a*c*d*(-3*a*e^2+c*d^2)-a*b*e*(a*e^2+3*c*d^2))+a*B*(a*b^2*e^3+2*a*c*e*(-a*e^2+3*c*d^2)-b*c*d*(3*a*e^2+c*d^2))-(a*B*(-b*e+2*c*d)*(c^2*d^2+b^2*e^2-c*e*(3*a*e+b*d))+A*c*(a*b^2*e^3+2*a*c*e*(-a*e^2+3*c*d^2)-b*c*d*(3*a*e^2+c*d^2)))*x^2)/a/c^2/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^(1/2)+1/3*B*e^3*x*(c*x^4+b*x^2+a)^(1/2)/c^2+1/3*(a*B*(6*c^3*d^3-8*b^3*e^3-9*c^2*d*e*(6*a*e+b*d)+b*c*e^2*(29*a*e+18*b*d))+3*A*c*(2*a*b^2*e^3+6*a*c*e*(-a*e^2+c*d^2)-b*c*d*(3*a*e^2+c*d^2)))*x*(c*x^4+b*x^2+a)^(1/2)/a/c^(5/2)/(-4*a*c+b^2)/(a^(1/2)+x^2*c^(1/2))-1/3*(a*B*(6*c^3*d^3-8*b^3*e^3-9*c^2*d*e*(6*a*e+b*d)+b*c*e^2*(29*a*e+18*b*d))+3*A*c*(2*a*b^2*e^3+6*a*c*e*(-a*e^2+c*d^2)-b*c*d*(3*a*e^2+c*d^2)))*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2))^2)^(1/2)/a^(3/4)/c^(11/4)/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^(1/2)-1/6*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(3*A*c^3*d^3-5*a^2*B*c*e^3+a*e*(-2*b*e+3*c*d))*(3*A*c*e-4*B*b*e+3*B*c*d)-3*c^(5/2)*d^2*(3*A*e+B*d)*a^(1/2)+3*a^(3/2)*e^2*(3*A*c*e-4*B*b*e+9*B*c*d)*c^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2))^2)^(1/2)/a^(3/4)/c^(11/4)/(b-2*a^(1/2)*c^(1/2))/(c*x^4+b*x^2+a)^(1/2)
```

### 3.24.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 15.27 (sec) , antiderivative size = 1058, normalized size of antiderivative = 1.23

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + bx^2 + cx^4)^{3/2}} dx = \frac{-4c \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x (3Ac(b^2(cd^3 - ae^3x^2) + b(-a^2e^3 + c^2d^3x^2 - 3acde(d - ex^2)))}{(a + bx^2 + cx^4)^{3/2}}$$

input `Integrate[((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(3/2),x]`

output

```
(-4*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x*(3*A*c*(b^2*(c*d^3 - a*e^3*x^2) +
b*(-a^2*e^3) + c^2*d^3*x^2 - 3*a*c*d*e*(d - e*x^2)) + 2*a*c*(a*e^2*(3*d +
e*x^2) - c*d^2*(d + 3*e*x^2))) + a*B*(4*b^3*e^3*x^2 + b^2*e^2*(4*a*e - 9*
c*d*x^2 + c*e*x^4) - b*c*(3*c*d^2*(d - 3*e*x^2) + a*e^2*(9*d + 13*e*x^2))
- 2*c*(5*a^2*e^3 + 3*c^2*d^3*x^2 + a*c*e*(-9*d^2 - 9*d*e*x^2 + 2*e^2*x^4))
)) + I*(-b + Sqrt[b^2 - 4*a*c])*(a*B*(-6*c^3*d^3 + 8*b^3*e^3 + 9*c^2*d*e*(
b*d + 6*a*e) - b*c*e^2*(18*b*d + 29*a*e)) + 3*A*c*(-2*a*b^2*e^3 + 6*a*c*e*
(-(c*d^2) + a*e^2) + b*c*d*(c*d^2 + 3*a*e^2))*Sqrt[(b + Sqrt[b^2 - 4*a*c]
+ 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c
*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqr
t[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] - I*
(a*B*(8*b^3*(-b + Sqrt[b^2 - 4*a*c])*e^3 - 6*c^3*d^2*(Sqrt[b^2 - 4*a*c]*d
- 6*a*e) + b*c*e^2*(18*b^2*d - 18*b*Sqrt[b^2 - 4*a*c]*d + 37*a*b*e - 29*a*
Sqrt[b^2 - 4*a*c]*e) + c^2*e*(-9*b^2*d^2 + 2*a*e*(27*Sqrt[b^2 - 4*a*c]*d -
10*a*e) + 9*b*d*(Sqrt[b^2 - 4*a*c]*d - 8*a*e))) + 3*A*c*(2*a*b^3*e^3 - b^
2*(c^2*d^3 + 3*a*c*d*e^2 + 2*a*Sqrt[b^2 - 4*a*c]*e^3) + b*c*(c*Sqrt[b^2 -
4*a*c]*d^3 + a*e^2*(3*Sqrt[b^2 - 4*a*c]*d - 8*a*e)) + 2*a*c*(2*c^2*d^3 + 3
*a*Sqrt[b^2 - 4*a*c]*e^3 - 3*c*d*e*(Sqrt[b^2 - 4*a*c]*d - 2*a*e))))*Sqrt[(
b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sq
rt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticF[I*ArcSinh...
```

### 3.24.3 Rubi [A] (verified)

Time = 1.72 (sec) , antiderivative size = 765, normalized size of antiderivative = 0.89, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.212$ , Rules used = {2206, 2207, 27, 1511, 27, 1416, 1509}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + bx^2 + cx^4)^{3/2}} dx$$

↓ 2206

$$x(-x^2(Ac(ab^2e^3 - bcd(3ae^2 + cd^2)) + 2ace(3cd^2 - ae^2)) + aB(2cd - be)(-ce(3ae + bd) + b^2e^2 + c^2d^2)) + Ac$$

$$\int \frac{aB(4a - \frac{b^2}{c})e^3x^4 - \frac{(aB(2c^3d^3 - 3c^2e(bd + 6ae)d - 2b^3e^3 + bce^2(6bd + 7ae)) + Ac(2ab^2e^3 + 6ac(cd^2 - ae^2)e - bcd(cd^2 + 3ae^2)))x^2}{c^2} + \frac{ac^2(b^2 - 4ac)\sqrt{a + \sqrt{cx^4 + bx^2 + a}}}{a(ab^2Be^3 - bc(Bcd^3 + 3Ace))}}{a(b^2 - 4ac)\sqrt{cx^4 + bx^2 + a}}$$

↓ 2207

---

3.24.  $\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+bx^2+cx^4)^{3/2}} dx$

$$\frac{x\left(-\left(x^2\left(Ac\left(ab^2e^3-bcd\left(3ae^2+cd^2\right)+2ace\left(3cd^2-ae^2\right)\right)+aB\left(2cd-be\right)\left(-ce\left(3ae+bd\right)+b^2e^2+c^2d^2\right)\right)\right)+Ac\left(2ab^2e^3-3bc\left(Bcd^3+3Aced^2+3aBe^2d+aAe^3\right)+2c\left(aBe\left(9cd^2-5ae^2\right)+3Acd\left(cd^2+3ae^2\right)\right)\right)-\left(aB\left(6c^3d^3-9c^2e\left(bd+6ae\right)d-8b^3e^3+bce^2\left(18bd+29ae\right)\right)+3Ac\left(2ab^2e^3-bcd\left(3ae^2+cd^2\right)+2ace\left(3cd^2-ae^2\right)\right)\right)}{c\sqrt{cx^4+bx^2+a}}}{\frac{ac^2\left(b^2-4ac\right)\sqrt{a}}{3c}}+a\left(b^2-4ac\right)$$

↓ 27

$$\frac{x\left(-\left(x^2\left(Ac\left(ab^2e^3-bcd\left(3ae^2+cd^2\right)+2ace\left(3cd^2-ae^2\right)\right)+aB\left(2cd-be\right)\left(-ce\left(3ae+bd\right)+b^2e^2+c^2d^2\right)\right)\right)+Ac\left(2ab^2e^3-3bc\left(Bcd^3+3Aced^2+3aBe^2d+aAe^3\right)+2c\left(aBe\left(9cd^2-5ae^2\right)+3Acd\left(cd^2+3ae^2\right)\right)\right)-\left(aB\left(6c^3d^3-9c^2e\left(bd+6ae\right)d-8b^3e^3+bce^2\left(18bd+29ae\right)\right)+3Ac\left(2ab^2e^3-bcd\left(3ae^2+cd^2\right)+2ace\left(3cd^2-ae^2\right)\right)\right)}{\sqrt{cx^4+bx^2+a}}}{\frac{ac^2\left(b^2-4ac\right)\sqrt{a}}{3c^2}}+a\left(b^2-4ac\right)$$

↓ 1511

$$\frac{x\left(-\left(x^2\left(Ac\left(ab^2e^3-bcd\left(3ae^2+cd^2\right)+2ace\left(3cd^2-ae^2\right)\right)+aB\left(2cd-be\right)\left(-ce\left(3ae+bd\right)+b^2e^2+c^2d^2\right)\right)\right)+Ac\left(2ab^2e^3-3bc\left(Bcd^3+3Aced^2+3aBe^2d+aAe^3\right)+2c\left(aBe\left(9cd^2-5ae^2\right)+3Acd\left(cd^2+3ae^2\right)\right)\right)-\left(aB\left(6c^3d^3-9c^2e\left(bd+6ae\right)d-8b^3e^3+bce^2\left(18bd+29ae\right)\right)+3Ac\left(2ab^2e^3-bcd\left(3ae^2+cd^2\right)+2ace\left(3cd^2-ae^2\right)\right)\right)}{\sqrt{c}}}{\frac{ac^2\left(b^2-4ac\right)\sqrt{a}}{3c^2}}+\frac{\sqrt{a}\left(2\sqrt{a}\sqrt{c}+b\right)\left(3a^{3/2}\sqrt{c}e^2\left(3Ace-4bBe+9Bcd\right)-5a^2Bce^3+ae\left(3cd-2be\right)\left(3Ace-4bBe+3Bcd\right)-3\sqrt{ac}^{5/2}d^2\left(3Ae+Bd\right)+3Ac^3d^3\right)}{3c^2}}+\frac{\int\frac{1}{\sqrt{cx^4+bx^2+a}}dx}{\sqrt{c}}+\frac{\sqrt{a}\left(3Ac\left(2ab^2e^3-bcd\left(3ae^2+cd^2\right)+2ace\left(3cd^2-ae^2\right)\right)+Ac\left(2ab^2e^3-3bc\left(Bcd^3+3Aced^2+3aBe^2d+aAe^3\right)+2c\left(aBe\left(9cd^2-5ae^2\right)+3Acd\left(cd^2+3ae^2\right)\right)\right)-\left(aB\left(6c^3d^3-9c^2e\left(bd+6ae\right)d-8b^3e^3+bce^2\left(18bd+29ae\right)\right)+3Ac\left(2ab^2e^3-bcd\left(3ae^2+cd^2\right)+2ace\left(3cd^2-ae^2\right)\right)\right)}{3c^2}}}{a\left(b^2-4ac\right)}$$

↓ 27

$$\frac{x\left(-\left(x^2\left(Ac\left(ab^2e^3-bcd\left(3ae^2+cd^2\right)+2ace\left(3cd^2-ae^2\right)\right)+aB\left(2cd-be\right)\left(-ce\left(3ae+bd\right)+b^2e^2+c^2d^2\right)\right)\right)+Ac\left(2ab^2e^3-3bc\left(Bcd^3+3Aced^2+3aBe^2d+aAe^3\right)+2c\left(aBe\left(9cd^2-5ae^2\right)+3Acd\left(cd^2+3ae^2\right)\right)\right)-\left(aB\left(6c^3d^3-9c^2e\left(bd+6ae\right)d-8b^3e^3+bce^2\left(18bd+29ae\right)\right)+3Ac\left(2ab^2e^3-bcd\left(3ae^2+cd^2\right)+2ace\left(3cd^2-ae^2\right)\right)\right)}{\sqrt{c}}}{\frac{ac^2\left(b^2-4ac\right)\sqrt{a}}{3c^2}}+\frac{\sqrt{a}\left(2\sqrt{a}\sqrt{c}+b\right)\left(3a^{3/2}\sqrt{c}e^2\left(3Ace-4bBe+9Bcd\right)-5a^2Bce^3+ae\left(3cd-2be\right)\left(3Ace-4bBe+3Bcd\right)-3\sqrt{ac}^{5/2}d^2\left(3Ae+Bd\right)+3Ac^3d^3\right)}{3c^2}}+\frac{\int\frac{1}{\sqrt{cx^4+bx^2+a}}dx}{\sqrt{c}}+\frac{\sqrt{a}\left(3Ac\left(2ab^2e^3-bcd\left(3ae^2+cd^2\right)+2ace\left(3cd^2-ae^2\right)\right)+Ac\left(2ab^2e^3-3bc\left(Bcd^3+3Aced^2+3aBe^2d+aAe^3\right)+2c\left(aBe\left(9cd^2-5ae^2\right)+3Acd\left(cd^2+3ae^2\right)\right)\right)-\left(aB\left(6c^3d^3-9c^2e\left(bd+6ae\right)d-8b^3e^3+bce^2\left(18bd+29ae\right)\right)+3Ac\left(2ab^2e^3-bcd\left(3ae^2+cd^2\right)+2ace\left(3cd^2-ae^2\right)\right)\right)}{3c^2}}}{a\left(b^2-4ac\right)}$$

↓ 1416

$$\frac{x\left(-\left(x^2\left(Ac\left(ab^2e^3-bcd\left(3ae^2+cd^2\right)+2ace\left(3cd^2-ae^2\right)\right)+aB\left(2cd-be\right)\left(-ce\left(3ae+bd\right)+b^2e^2+c^2d^2\right)\right)\right)+Ac\left(2ab^2e^3-3bc\left(Bcd^3+3Aced^2+3aBe^2d+aAe^3\right)+2c\left(aBe\left(9cd^2-5ae^2\right)+3Acd\left(cd^2+3ae^2\right)\right)\right)-\left(aB\left(6c^3d^3-9c^2e\left(bd+6ae\right)d-8b^3e^3+bce^2\left(18bd+29ae\right)\right)+3Ac\left(2ab^2e^3-bcd\left(3ae^2+cd^2\right)+2ace\left(3cd^2-ae^2\right)\right)\right)}{\sqrt{c}}}{\frac{ac^2\left(b^2-4ac\right)\sqrt{a}}{3c^2}}+\frac{\sqrt{a}\left(2\sqrt{a}\sqrt{c}+b\right)\left(3a^{3/2}\sqrt{c}e^2\left(3Ace-4bBe+9Bcd\right)-5a^2Bce^3+ae\left(3cd-2be\right)\left(3Ace-4bBe+3Bcd\right)-3\sqrt{ac}^{5/2}d^2\left(3Ae+Bd\right)+3Ac^3d^3\right)}{3c^2}}+\frac{\int\frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{cx^4+bx^2+a}}dx}{\sqrt{c}}+\frac{\sqrt[4]{a}\left(2\sqrt{a}\sqrt{c}+b\right)\left(\sqrt{a}+\sqrt{cx^2}\right)}{\sqrt{c}}+\frac{\sqrt{a}\left(3Ac\left(2ab^2e^3-bcd\left(3ae^2+cd^2\right)+6ace\left(cd^2-ae^2\right)\right)+aB\left(-9c^2de\left(6ae+bd\right)+bce^2\left(29ae+18bd\right)-8b^3e^3+6c^3d^3\right)\right)}{\sqrt{c}}}{a\left(b^2-4ac\right)}$$

↓ 1509

3.24.  $\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+bx^2+cx^4)^{3/2}} dx$

$$x \left( - \left( x^2 \left( Ac(ab^2e^3 - bcd(3ae^2 + cd^2)) + 2ace(3cd^2 - ae^2) \right) + aB(2cd - be) \left( -ce(3ae + bd) + b^2e^2 + c^2d^2 \right) \right) + Ac \right) / (ac^2(b^2 - 4ac)\sqrt{a +$$

$$\sqrt[4]{a} (2\sqrt{a}\sqrt{c+b}) (\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \text{EllipticF} \left( 2 \arctan \left( \frac{\sqrt[4]{Cx}}{\sqrt[4]{a}} \right), \frac{1}{4} \left( 2 - \frac{b}{\sqrt{a}\sqrt{c}} \right) \right) (3a^{3/2}\sqrt{ce^2}(3Ace-4bBe+9Bcd)-5a^2Bce^3+ae(3cd-2be)(3Ace-4bBc) - 2c^{3/4}\sqrt{a+bx^2+cx^4}}$$

input `Int[((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(3/2), x]`

output `(x*(A*c*(b^2*c*d^3 - 2*a*c*d*(c*d^2 - 3*a*e^2) - a*b*e*(3*c*d^2 + a*e^2)) + a*B*(a*b^2*e^3 + 2*a*c*e*(3*c*d^2 - a*e^2) - b*c*d*(c*d^2 + 3*a*e^2)) - (a*B*(2*c*d - b*e)*(c^2*d^2 + b^2*e^2 - c*e*(b*d + 3*a*e)) + A*c*(a*b^2*e^3 + 2*a*c*e*(3*c*d^2 - a*e^2) - b*c*d*(c*d^2 + 3*a*e^2)))*x^2)/(a*c^2*(b^2 - 4*a*c)*Sqrt[a + b*x^2 + c*x^4]) - (-1/3*(a*B*(b^2 - 4*a*c)*e^3*x*Sqrt[a + b*x^2 + c*x^4])/c^2 + (((a*B*(6*c^3*d^3 - 8*b^3*e^3 - 9*c^2*d*e*(b*d + 6*a*e)) + b*c*e^2*(18*b*d + 29*a*e)) + 3*A*c*(2*a*b^2*e^3 + 6*a*c*e*(c*d^2 - a*e^2) - b*c*d*(c*d^2 + 3*a*e^2)))*(-(x*Sqrt[a + b*x^2 + c*x^4])/(Sqrt[a] + Sqrt[c]*x^2)) + (a^(1/4)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(c^(1/4)*Sqrt[a + b*x^2 + c*x^4]))/Sqrt[c] + (a^(1/4)*(b + 2*Sqrt[a]*Sqrt[c])*(3*A*c^3*d^3 - 5*a^2*B*c*e^3 - 3*Sqrt[a]*c^(5/2)*d^2*(B*d + 3*A*e) + a*e*(3*c*d - 2*b*e)*(3*B*c*d - 4*b*B*e + 3*A*c*e) + 3*a^(3/2)*Sqrt[c]*e^2*(9*B*c*d - 4*b*B*e + 3*A*c*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*c^(3/4)*Sqrt[a + b*x^2 + c*x^4]))/(3*c^2))/(a*(b^2 - 4*a*c))`

### 3.24.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 1416 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`

$$3.24. \int \frac{(A+Bx^2)(d+ex^2)^3}{(a+bx^2+cx^4)^{3/2}} dx$$

rule 1509 `Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`

rule 1511 `Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 2]}, Simp[(e + d*q)/q Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Simp[e/q Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`

rule 2206 `Int[(Px_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := With[{d = Coeff[PolynomialRemainder[Px, a + b*x^2 + c*x^4, x], x, 0], e = Coeff[PolynomialRemainder[Px, a + b*x^2 + c*x^4, x], x, 2]}, Simp[x*(a + b*x^2 + c*x^4)^(p + 1)*((a*b*e - d*(b^2 - 2*a*c) - c*(b*d - 2*a*e)*x^2)/(2*a*(p + 1)*(b^2 - 4*a*c)), x] + Simp[1/(2*a*(p + 1)*(b^2 - 4*a*c)) Int[(a + b*x^2 + c*x^4)^(p + 1)*ExpandToSum[2*a*(p + 1)*(b^2 - 4*a*c)*PolynomialQuotient[Px, a + b*x^2 + c*x^4, x] + b^2*d*(2*p + 3) - 2*a*c*d*(4*p + 5) - a*b*e + c*(4*p + 7)*(b*d - 2*a*e)*x^2, x], x]] /; FreeQ[{a, b, c}, x] && PolyQ[Px, x^2] && Expon[Px, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1]`

rule 2207 `Int[(Px_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := With[{n = Expon[Px, x^2], e = Coeff[Px, x^2, Expon[Px, x^2]]}, Simp[e*x^(2*n - 3)*((a + b*x^2 + c*x^4)^(p + 1)/(c*(2*n + 4*p + 1))), x] + Simp[1/(c*(2*n + 4*p + 1)) Int[(a + b*x^2 + c*x^4)^p*ExpandToSum[c*(2*n + 4*p + 1)*Px - a*e*(2*n - 3)*x^(2*n - 4) - b*e*(2*n + 2*p - 1)*x^(2*n - 2) - c*e*(2*n + 4*p + 1)*x^(2*n), x], x]] /; FreeQ[{a, b, c, p}, x] && PolyQ[Px, x^2] && Expon[Px, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && !LtQ[p, -1]`

### 3.24.4 Maple [A] (verified)

Time = 12.57 (sec) , antiderivative size = 1141, normalized size of antiderivative = 1.33

| method   | result                          | size |
|----------|---------------------------------|------|
| elliptic | Expression too large to display | 1141 |
| default  | Expression too large to display | 2445 |
| risch    | Expression too large to display | 2482 |

```
input int((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(3/2),x,method=_RETURNVERBOSE)
```

```
output -2*c*(1/2/c^3*(2*A*a^2*c^2*e^3-A*a*b^2*c*e^3+3*A*a*b*c^2*d*e^2-6*A*a*c^3*d^2*e+A*b*c^3*d^3-3*B*a^2*b*c*e^3+6*B*a^2*c^2*d*e^2+B*a*b^3*e^3-3*B*a*b^2*c*d*e^2+3*B*a*b*c^2*d^2*e-2*B*a*c^3*d^3)/a/(4*a*c-b^2)*x^3-1/2/c^3*(A*a^2*b*c*e^3-6*A*a^2*c^2*d*e^2+3*A*a*b*c^2*d^2*e+2*A*a*c^3*d^3-A*b^2*c^2*d^3+2*B*a^3*c*e^3-B*a^2*b^2*e^3+3*B*a^2*b*c*d*e^2-6*B*a^2*c^2*d^2*e+B*a*b*c^2*d^3)/a/(4*a*c-b^2)*x/((x^4+1/c*b*x^2+a/c)*c)^(1/2)+1/3*B*e^3*x*(c*x^4+b*x^2+a)^(1/2)/c^2+1/4*(-e*(A*b*c*e^2-3*A*c^2*d*e+B*a*c*e^2-B*b^2*e^2+3*B*b*c*d*e-3*B*c^2*d^2)/c^3+1/c^3*(A*a*b*c*e^3-3*A*a*c^2*d*e^2+A*c^3*d^3+B*a^2*c*e^3-B*a*b^2*e^3+3*B*a*b*c*d*e^2-3*B*a*c^2*d^2*e)/a-1/c^2*(A*a^2*b*c*e^3-6*A*a^2*c^2*d*e^2+3*A*a*b*c^2*d^2*e+2*A*a*c^3*d^3-A*b^2*c^2*d^3+2*B*a^3*c*e^3-B*a^2*b^2*e^3+3*B*a^2*b*c*d*e^2-6*B*a^2*c^2*d^2*e+B*a*b*c^2*d^3)/a/(4*a*c-b^2)-1/3*B*e^3/c^2*a)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*x^2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))-1/2*(1/c^2*e^2*(A*c*e-B*b*e+3*B*c*d)+1/c^2*(2*A*a^2*c^2*e^3-A*a*b^2*c*e^3+3*A*a*b*c^2*d*e^2-6*A*a*c^3*d^2*e+A*b*c^3*d^3-3*B*a^2*b*c*e^3+6*B*a^2*c^2*d*e^2+B*a*b^3*e^3-3*B*a*b^2*c*d*e^2+3*B*a*b*c^2*d^2*e-2*B*a*c^3*d^3)/a/(4*a*c-b^2)-2/3*B/c^2*e^3*b)*a*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a...
```

### 3.24.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2623 vs. 2(841) = 1682.

Time = 0.13 (sec) , antiderivative size = 2623, normalized size of antiderivative = 3.05

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + bx^2 + cx^4)^{3/2}} dx = \text{Too large to display}$$

3.24.  $\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+bx^2+cx^4)^{3/2}} dx$



input `integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fricas")`

output `-1/6*(sqrt(1/2)*((3*(2*B*a*b - A*b^2)*c^4*d^3 - 9*(B*a*b^2*c^3 - 2*A*a*b*c^4)*d^2*e + 9*(2*B*a*b^3*c^2 - (6*B*a^2*b + A*a*b^2)*c^3)*d*e^2 - (8*B*a*b^4*c + 18*A*a^2*b*c^3 - (29*B*a^2*b^2 + 6*A*a*b^3)*c^2)*e^3)*x^5 + (3*(2*B*a*b^2 - A*b^3)*c^3*d^3 - 9*(B*a*b^3*c^2 - 2*A*a*b^2*c^3)*d^2*e + 9*(2*B*a*b^4*c - (6*B*a^2*b^2 + A*a*b^3)*c^2)*d*e^2 - (8*B*a*b^5 + 18*A*a^2*b^2*c^2 - (29*B*a^2*b^3 + 6*A*a*b^4)*c)*e^3)*x^3 + (3*(2*B*a^2*b - A*a*b^2)*c^3*d^3 - 9*(B*a^2*b^2*c^2 - 2*A*a^2*b*c^3)*d^2*e + 9*(2*B*a^2*b^3*c - (6*B*a^3*b + A*a^2*b^2)*c^2)*d*e^2 - (8*B*a^2*b^4 + 18*A*a^3*b*c^2 - (29*B*a^3*b^2 + 6*A*a^2*b^3)*c)*e^3)*x - ((3*(2*B*a - A*b)*c^5*d^3 - 9*(B*a*b*c^4 - 2*A*a*c^5)*d^2*e + 9*(2*B*a*b^2*c^3 - (6*B*a^2 + A*a*b)*c^4)*d*e^2 - (8*B*a*b^3*c^2 + 18*A*a^2*c^4 - (29*B*a^2*b + 6*A*a*b^2)*c^3)*e^3)*x^5 + (3*(2*B*a*b - A*b^2)*c^4*d^3 - 9*(B*a*b^2*c^3 - 2*A*a*b*c^4)*d^2*e + 9*(2*B*a*b^3*c^2 - (6*B*a^2*b + A*a*b^2)*c^3)*d*e^2 - (8*B*a*b^4*c + 18*A*a^2*b*c^3 - (29*B*a^2*b^2 + 6*A*a*b^3)*c^2)*e^3)*x^3 + (3*(2*B*a^2 - A*a*b)*c^4*d^3 - 9*(B*a^2*b*c^3 - 2*A*a^2*c^4)*d^2*e + 9*(2*B*a^2*b^2*c^2 - (6*B*a^3 + A*a^2*b)*c^3)*d*e^2 - (8*B*a^2*b^3*c + 18*A*a^3*c^3 - (29*B*a^3*b + 6*A*a^2*b^2)*c^2)*e^3)*x)*sqrt((b^2 - 4*a*c)/c^2))*sqrt(c)*sqrt((c*sqrt((b^2 - 4*a*c)/c^2) - b)/c)*elliptic_e(arcsin(sqrt(1/2)*sqrt((c*sqrt((b^2 - 4*a*c)/c^2) - b)/c)/x), 1/2*(b*c*sqrt((b^2 - 4*a*c)/c^2) + b^2 - 2*a*c)/(a*c)) + sqrt(1/2)*((3*(2*A*b*c^5 - (2*B*a*b - (A - B)*b^2)*c^4)*d^3 + 9*(B*a*b^2*c^3...`

### 3.24.6 Sympy [F]

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{(A + Bx^2)(d + ex^2)^3}{(a + bx^2 + cx^4)^{3/2}} dx$$

input `integrate((B*x**2+A)*(e*x**2+d)**3/(c*x**4+b*x**2+a)**(3/2),x)`

output `Integral((A + B*x**2)*(d + e*x**2)**3/(a + b*x**2 + c*x**4)**(3/2), x)`

**3.24.7 Maxima [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^3}{(cx^4 + bx^2 + a)^{3/2}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^3/(c*x^4 + b*x^2 + a)^(3/2), x)`

**3.24.8 Giac [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^3}{(cx^4 + bx^2 + a)^{3/2}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^3/(c*x^4 + b*x^2 + a)^(3/2), x)`

**3.24.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^3}{(cx^4 + bx^2 + a)^{3/2}} dx$$

input `int(((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(3/2), x)`

output `int(((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(3/2), x)`

$$3.25 \quad \int \frac{(A+Bx^2)(d+ex^2)^2}{(a+bx^2+cx^4)^{3/2}} dx$$

|        |   |     |
|--------|---|-----|
| 3.25.1 | Optimal result                            | 218 |
| 3.25.2 | Mathematica [C] (verified)                | 219 |
| 3.25.3 | Rubi [A] (verified)                       | 220 |
| 3.25.4 | Maple [A] (verified)                      | 223 |
| 3.25.5 | Fricas [B] (verification not implemented) | 224 |
| 3.25.6 | Sympy [F]                                 | 225 |
| 3.25.7 | Maxima [F]                                | 226 |
| 3.25.8 | Giac [F]                                  | 226 |
| 3.25.9 | Mupad [F(-1)]                             | 226 |

### 3.25.1 Optimal result

Integrand size = 33, antiderivative size = 628

$$\int \frac{(A+Bx^2)(d+ex^2)^2}{(a+bx^2+cx^4)^{3/2}} dx =$$

$$\frac{x(aB(bcd^2 - 4acde + abe^2) - Ac(b^2d^2 - 2abde - 2a(cd^2 - ae^2)) - (Ac(bcd^2 - 4acde + abe^2) - aB(2c^2d^2 - ac(b^2 - 4ac)\sqrt{a+bx^2+cx^4}))}{ac^3/2(b^2 - 4ac)(\sqrt{a} + \sqrt{cx^2})} + \frac{(Ac(bcd^2 - 4acde + abe^2) - 2aB(c^2d^2 + b^2e^2 - ce(bd + 3ae)))x\sqrt{a+bx^2+cx^4}}{ac^3/2(b^2 - 4ac)(\sqrt{a} + \sqrt{cx^2})} + \frac{(Ac(bcd^2 - 4acde + abe^2) - 2aB(c^2d^2 + b^2e^2 - ce(bd + 3ae)))(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{a+bx^2+cx^4}}{\sqrt{a}+\sqrt{cx^2}}\right)\right)}{a^{3/4}c^{7/4}(b^2 - 4ac)\sqrt{a+bx^2+cx^4}} + \frac{(Ac^2d^2 + 3a^{3/2}B\sqrt{ce^2} - \sqrt{ac^3/2}d(Bd + 2Ae) + ae(2Bcd - 2bBe + Ace))(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}\text{EllipE}}{2a^{3/4}(b - 2\sqrt{a}\sqrt{c})c^{7/4}\sqrt{a+bx^2+cx^4}}$$

---

3.25.  $\int \frac{(A+Bx^2)(d+ex^2)^2}{(a+bx^2+cx^4)^{3/2}} dx$

output

```

-x*(a*B*(a*b*e^2-4*a*c*d*e+b*c*d^2)-A*c*(b^2*d^2-2*a*b*d*e-2*a*(-a*e^2+c*d
^2))-(A*c*(a*b*e^2-4*a*c*d*e+b*c*d^2)-a*B*(2*c^2*d^2+b^2*e^2-2*c*e*(a*e+b*
d)))*x^2)/a/c/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^(1/2)-(A*c*(a*b*e^2-4*a*c*d*e+b
*c*d^2)-2*a*B*(c^2*d^2+b^2*e^2-c*e*(3*a*e+b*d)))*x*(c*x^4+b*x^2+a)^(1/2)/a
/c^(3/2)/(-4*a*c+b^2)/(a^(1/2)+x^2*c^(1/2))+(A*c*(a*b*e^2-4*a*c*d*e+b*c*d^
2)-2*a*B*(c^2*d^2+b^2*e^2-c*e*(3*a*e+b*d)))*(cos(2*arctan(c^(1/4)*x/a^(1/4
)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/
4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*
x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2))^2)^(1/2)/a^(3/4)/c^(7/4)/(-4*a*c+b^2)/(
c*x^4+b*x^2+a)^(1/2)-1/2*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*
arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*
(2-b/a^(1/2)/c^(1/2))^(1/2))*(A*c^2*d^2+a*e*(A*c*e-2*B*b*e+2*B*c*d)-c^(3/2
)*d*(2*A*e+B*d)*a^(1/2)+3*a^(3/2)*B*e^2*c^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c
*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2))^2)^(1/2)/a^(3/4)/c^(7/4)/(b-2*a^(1/2)*
c^(1/2))/(c*x^4+b*x^2+a)^(1/2)

```

### 3.25.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 12.99 (sec) , antiderivative size = 766, normalized size of antiderivative = 1.22

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + bx^2 + cx^4)^{3/2}} dx = \frac{-4c\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}x(-aB(abe^2 + 2c^2d^2x^2 + b^2e^2x^2 + bcd(d - 2ex^2) - 2ace(2d - ex^2)) + (A + Bx^2)(d + ex^2)^2)}{(a + bx^2 + cx^4)^{3/2}}$$

input `Integrate[((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(3/2), x]`

---

3.25.  $\int \frac{(A+Bx^2)(d+ex^2)^2}{(a+bx^2+cx^4)^{3/2}} dx$

```
output (-4*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])] * x * (- (a*B*(a*b*e^2 + 2*c^2*d^2*x^2 +
b^2*e^2*x^2 + b*c*d*(d - 2*e*x^2) - 2*a*c*e*(2*d + e*x^2))) + A*c*(b^2*d^2
+ 2*a^2*e^2 + b*c*d^2*x^2 + a*b*e*(-2*d + e*x^2) - 2*a*c*d*(d + 2*e*x^2))
) - I*(-b + Sqrt[b^2 - 4*a*c]) * (- (A*c*(b*c*d^2 - 4*a*c*d*e + a*b*e^2)) + 2
*a*B*(c^2*d^2 + b^2*e^2 - c*e*(b*d + 3*a*e))) * Sqrt[(b + Sqrt[b^2 - 4*a*c]
+ 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])] * Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*
x^2)/(b - Sqrt[b^2 - 4*a*c])] * EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt
[b^2 - 4*a*c])] * x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] + I*(
2*a*B*(b^2*(-b + Sqrt[b^2 - 4*a*c])*e^2 + c^2*d*(Sqrt[b^2 - 4*a*c]*d - 4*a
*e) + c*e*(b^2*d - b*Sqrt[b^2 - 4*a*c]*d + 4*a*b*e - 3*a*Sqrt[b^2 - 4*a*c]
*e)) + A*c*(b^2*(c*d^2 + a*e^2) - b*Sqrt[b^2 - 4*a*c]*(c*d^2 + a*e^2) - 4*
a*c*(c*d^2 - Sqrt[b^2 - 4*a*c]*d*e + a*e^2))) * Sqrt[(b + Sqrt[b^2 - 4*a*c]
+ 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])] * Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*
x^2)/(b - Sqrt[b^2 - 4*a*c])] * EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt
[b^2 - 4*a*c])] * x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])])]/(4*a
*c^2*(-b^2 + 4*a*c)*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])] * Sqrt[a + b*x^2 + c*x^4
])
```

### 3.25.3 Rubi [A] (verified)

Time = 0.92 (sec) , antiderivative size = 576, normalized size of antiderivative = 0.92, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2206, 27, 1511, 27, 1416, 1509}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + bx^2 + cx^4)^{3/2}} dx$$

↓ 2206

$$\int \frac{(Ac(bcd^2 - 4acde + abe^2) - 2aB(c^2d^2 + b^2e^2 - ce(bd + 3ae)))x^2 + a(ae(4Bcd - bBe + 2Ace) + cd(2Acd - b(Bd + 2Ae)))}{c\sqrt{cx^4 + bx^2 + a}} dx$$


---


$$x \left( \frac{aB(abe^2 - 4acde + bcd^2)}{c} - A(-2abde - 2a(cd^2 - ae^2) + b^2d^2) \right) - \frac{x^2(Ac(abe^2 - 4acde + bcd^2) - aB(-2ce(ae + b^2d^2) + cd(2Acd - b(Bd + 2Ae))))}{ac(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}}$$

↓ 27

---

3.25.  $\int \frac{(A+Bx^2)(d+ex^2)^2}{(a+bx^2+cx^4)^{3/2}} dx$

$$\frac{\int \frac{(Ac(bcd^2 - 4aced + abe^2) - 2aB(c^2d^2 + b^2e^2 - ce(bd + 3ae)))x^2 + a(ae(4Bcd - bBe + 2Ace) + cd(2Ac d - b(Bd + 2Ae)))}{\sqrt{cx^4 + bx^2 + a}} dx}{ac(b^2 - 4ac)} -$$

$$x \left( c \left( \frac{aB(abe^2 - 4acde + bcd^2)}{c} - A(-2abde - 2a(cd^2 - ae^2) + b^2d^2) \right) - x^2 (Ac(abe^2 - 4acde + bcd^2) - aB(-2ce(ae + bcd))) \right) \frac{1}{ac(b^2 - 4ac) \sqrt{a + bx^2 + cx^4}}$$

↓ 1511

$$\frac{\sqrt{a}(2\sqrt{a}\sqrt{c}+b)(3a^{3/2}B\sqrt{ce^2}+ae(Ace-2bBe+2Bcd)-\sqrt{ac^{3/2}d(2Ae+Bd)+Ac^2d^2}) \int \frac{1}{\sqrt{cx^4+bx^2+a}} dx}{\sqrt{c}} - \frac{\sqrt{a}(Ac(abe^2-4acde+bcd^2)-2aB(-2ce(ae+bcd)))}{ac(b^2-4ac)}$$

$$x \left( c \left( \frac{aB(abe^2 - 4acde + bcd^2)}{c} - A(-2abde - 2a(cd^2 - ae^2) + b^2d^2) \right) - x^2 (Ac(abe^2 - 4acde + bcd^2) - aB(-2ce(ae + bcd))) \right) \frac{1}{ac(b^2 - 4ac) \sqrt{a + bx^2 + cx^4}}$$

↓ 27

$$\frac{\sqrt{a}(2\sqrt{a}\sqrt{c}+b)(3a^{3/2}B\sqrt{ce^2}+ae(Ace-2bBe+2Bcd)-\sqrt{ac^{3/2}d(2Ae+Bd)+Ac^2d^2}) \int \frac{1}{\sqrt{cx^4+bx^2+a}} dx}{\sqrt{c}} - \frac{(Ac(abe^2-4acde+bcd^2)-2aB(-2ce(ae+bcd)))}{ac(b^2-4ac)}$$

$$x \left( c \left( \frac{aB(abe^2 - 4acde + bcd^2)}{c} - A(-2abde - 2a(cd^2 - ae^2) + b^2d^2) \right) - x^2 (Ac(abe^2 - 4acde + bcd^2) - aB(-2ce(ae + bcd))) \right) \frac{1}{ac(b^2 - 4ac) \sqrt{a + bx^2 + cx^4}}$$

↓ 1416

$$\frac{4\sqrt{a}(2\sqrt{a}\sqrt{c}+b)(\sqrt{a}+\sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{4\sqrt{Cx}}{4\sqrt{a}}\right), \frac{1}{4}\left(2-\frac{b}{\sqrt{a}\sqrt{c}}\right)\right) (3a^{3/2}B\sqrt{ce^2}+ae(Ace-2bBe+2Bcd)-\sqrt{ac^{3/2}d(2Ae+Bd)+Ac^2d^2})}{2c^{3/4}\sqrt{a+bx^2+cx^4}}$$

$$x \left( c \left( \frac{aB(abe^2 - 4acde + bcd^2)}{c} - A(-2abde - 2a(cd^2 - ae^2) + b^2d^2) \right) - x^2 (Ac(abe^2 - 4acde + bcd^2) - aB(-2ce(ae + bcd))) \right) \frac{1}{ac(b^2 - 4ac) \sqrt{a + bx^2 + cx^4}}$$

↓ 1509

$$\frac{4\sqrt{a}(2\sqrt{a}\sqrt{c}+b)(\sqrt{a}+\sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \text{EllipticF}\left(2 \arctan\left(\frac{4\sqrt{Cx}}{4\sqrt{a}}\right), \frac{1}{4}\left(2-\frac{b}{\sqrt{a}\sqrt{c}}\right)\right) (3a^{3/2}B\sqrt{ce^2}+ae(Ace-2bBe+2Bcd)-\sqrt{ac^{3/2}d(2Ae+Bd)+Ac^2d^2})}{2c^{3/4}\sqrt{a+bx^2+cx^4}}$$

$$x \left( c \left( \frac{aB(abe^2 - 4acde + bcd^2)}{c} - A(-2abde - 2a(cd^2 - ae^2) + b^2d^2) \right) - x^2 (Ac(abe^2 - 4acde + bcd^2) - aB(-2ce(ae + bcd))) \right) \frac{1}{ac(b^2 - 4ac) \sqrt{a + bx^2 + cx^4}}$$

---

3.25.  $\int \frac{(A+Bx^2)(d+ex^2)^2}{(a+bx^2+cx^4)^{3/2}} dx$

input `Int[((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(3/2),x]`

output `-((x*(c*((a*B*(b*c*d^2 - 4*a*c*d*e + a*b*e^2))/c - A*(b^2*d^2 - 2*a*b*d*e - 2*a*(c*d^2 - a*e^2))) - (A*c*(b*c*d^2 - 4*a*c*d*e + a*b*e^2) - a*B*(2*c^2*d^2 + b^2*e^2 - 2*c*e*(b*d + a*e)))*x^2))/(a*c*(b^2 - 4*a*c)*Sqrt[a + b*x^2 + c*x^4]) - (-(((A*c*(b*c*d^2 - 4*a*c*d*e + a*b*e^2) - 2*a*B*(c^2*d^2 + b^2*e^2 - c*e*(b*d + 3*a*e)))*(-(x*Sqrt[a + b*x^2 + c*x^4])/(Sqrt[a + Sqrt[c]*x^2)) + (a^(1/4)*(Sqrt[a + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a + Sqrt[c]*x^2)]^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(c^(1/4)*Sqrt[a + b*x^2 + c*x^4])))/Sqrt[c]) + (a^(1/4)*(b + 2*Sqrt[a]*Sqrt[c])*(A*c^2*d^2 + 3*a^(3/2)*B*Sqrt[c]*e^2 - Sqrt[a]*c^(3/2)*d*(B*d + 2*A*e) + a*e*(2*B*c*d - 2*b*B*e + A*c*e))*(Sqrt[a + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a + Sqrt[c]*x^2)]^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*c^(3/4)*Sqrt[a + b*x^2 + c*x^4]))/(a*c*(b^2 - 4*a*c))`

### 3.25.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1416 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`

rule 1509 `Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2])/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`

```
rule 1511 Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol]
  := With[{q = Rt[c/a, 2]}, Simp[(e + d*q)/q Int[1/Sqrt[a + b*x^2 + c*x^4], x], x]
  - Simp[e/q Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /;
  NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

```
rule 2206 Int[(Px_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := With[{d =
  Coeff[PolynomialRemainder[Px, a + b*x^2 + c*x^4, x], x, 0], e = Coeff[PolynomialRemainder[Px,
  a + b*x^2 + c*x^4, x], x, 2]}, Simp[x*(a + b*x^2 + c*x^4)^(p + 1)*((a*b*e - d*(b^2 - 2*a*c) -
  c*(b*d - 2*a*e)*x^2)/(2*a*(p + 1)*(b^2 - 4*a*c)), x] + Simp[1/(2*a*(p + 1)*(b^2 - 4*a*c))
  Int[(a + b*x^2 + c*x^4)^(p + 1)*ExpandToSum[2*a*(p + 1)*(b^2 - 4*a*c)*PolynomialQuotient[Px,
  a + b*x^2 + c*x^4, x] + b^2*d*(2*p + 3) - 2*a*c*d*(4*p + 5) - a*b*e + c*(4*p + 7)*(b*d -
  2*a*e)*x^2, x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Px, x^2] && Expon[Px, x^2] > 1 &&
  NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1]
```

### 3.25.4 Maple [A] (verified)

Time = 2.94 (sec) , antiderivative size = 799, normalized size of antiderivative = 1.27

| method   | result  |
|----------|---|
| elliptic | $2c \frac{\left( \frac{Aabc e^2 - 4Aa c^2 de + Ab c^2 d^2 + 2B a^2 c e^2 - Ba b^2 e^2 + 2Babcde - 2Ba c^2 d^2}{2c^2 a (4ac - b^2)} x^3 + \frac{(2A a^2 c e^2 - 2Aabcde - 2Aa c^2 d^2 + A b^2 c d^2 - B a^2 b e^2 + 4B a^2 c^2 d^2 - 2Aabcde - 2Aa c^2 d^2 + A b^2 c d^2 - B a^2 b e^2 + 4B a^2 c^2 d^2)}{2c^2 (4ac - b^2) a} \right)}{\sqrt{\left(x^4 + \frac{bx^2}{c} + \frac{a}{c}\right)c}}$ |
| default  | Expression too large to display   |

```
input int((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x,method=_RETURNVERBOSE)
```

$$3.25. \int \frac{(A+Bx^2)(d+ex^2)^2}{(a+bx^2+cx^4)^{3/2}} dx$$



output

```
-2*c*(1/2/c^2*(A*a*b*c*e^2-4*A*a*c^2*d*e+A*b*c^2*d^2+2*B*a^2*c*e^2-B*a*b^2
*e^2+2*B*a*b*c*d*e-2*B*a*c^2*d^2)/a/(4*a*c-b^2)*x^3+1/2*(2*A*a^2*c*e^2-2*A
*a*b*c*d*e-2*A*a*c^2*d^2+A*b^2*c*d^2-B*a^2*b*e^2+4*B*a^2*c*d*e-B*a*b*c*d^2
)/c^2/(4*a*c-b^2)/a*x)/((x^4+1/c*b*x^2+a/c)*c)^(1/2)+1/4*(e*(A*c*e-B*b*e+2
*B*c*d)/c^2-1/c^2*(A*a*c*e^2-A*c^2*d^2-B*a*b*e^2+2*B*a*c*d*e)/a+1/a*(2*A*a
^2*c*e^2-2*A*a*b*c*d*e-2*A*a*c^2*d^2+A*b^2*c*d^2-B*a^2*b*e^2+4*B*a^2*c*d*e
-B*a*b*c*d^2)/c/(4*a*c-b^2))*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-
2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(
1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2)
))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))-1/2*(1/c*B*e^2+
1/a*(A*a*b*c*e^2-4*A*a*c^2*d*e+A*b*c^2*d^2+2*B*a^2*c*e^2-B*a*b^2*e^2+2*B*a
*b*c*d*e-2*B*a*c^2*d^2)/c/(4*a*c-b^2))*a*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/
a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1
/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*(EllipticF(
1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2
)^(1/2))/a/c)^(1/2))-EllipticE(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(
1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2)))
```

### 3.25.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1669 vs.  $2(619) = 1238$ .

Time = 0.11 (sec) , antiderivative size = 1669, normalized size of antiderivative = 2.66

$$\int \frac{(A+Bx^2)(d+ex^2)^2}{(a+bx^2+cx^4)^{3/2}} dx = \text{Too large to display}$$

input

```
integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fricas
")
```

output

```

-1/2*(sqrt(1/2)*(((2*B*a*b - A*b^2)*c^3*d^2 - 2*(B*a*b^2*c^2 - 2*A*a*b*c^3)
)*d*e + (2*B*a*b^3*c - (6*B*a^2*b + A*a*b^2)*c^2)*e^2)*x^5 + ((2*B*a*b^2 -
A*b^3)*c^2*d^2 - 2*(B*a*b^3*c - 2*A*a*b^2*c^2)*d*e + (2*B*a*b^4 - (6*B*a^
2*b^2 + A*a*b^3)*c)*e^2)*x^3 + ((2*B*a^2*b - A*a*b^2)*c^2*d^2 - 2*(B*a^2*b
^2*c - 2*A*a^2*b*c^2)*d*e + (2*B*a^2*b^3 - (6*B*a^3*b + A*a^2*b^2)*c)*e^2)
*x - (((2*B*a - A*b)*c^4*d^2 - 2*(B*a*b*c^3 - 2*A*a*c^4)*d*e + (2*B*a*b^2*
c^2 - (6*B*a^2 + A*a*b)*c^3)*e^2)*x^5 + ((2*B*a*b - A*b^2)*c^3*d^2 - 2*(B*
a*b^2*c^2 - 2*A*a*b*c^3)*d*e + (2*B*a*b^3*c - (6*B*a^2*b + A*a*b^2)*c^2)*e
^2)*x^3 + ((2*B*a^2 - A*a*b)*c^3*d^2 - 2*(B*a^2*b*c^2 - 2*A*a^2*c^3)*d*e +
(2*B*a^2*b^2*c - (6*B*a^3 + A*a^2*b)*c^2)*e^2)*x)*sqrt((b^2 - 4*a*c)/c^2)
)*sqrt(c)*sqrt((c*sqrt((b^2 - 4*a*c)/c^2) - b)/c)*elliptic_e(arcsin(sqrt(1
/2)*sqrt((c*sqrt((b^2 - 4*a*c)/c^2) - b)/c)/x), 1/2*(b*c*sqrt((b^2 - 4*a*c
)/c^2) + b^2 - 2*a*c)/(a*c)) + sqrt(1/2)*(((2*A*b*c^4 - (2*B*a*b - (A - B)
)*b^2)*c^3)*d^2 + 2*(B*a*b^2*c^2 - (2*(A - B)*a*b + A*b^2)*c^3)*d*e - (2*B*
a*b^3*c - 2*A*a*b*c^3 - (6*B*a^2*b + (A - B)*a*b^2)*c^2)*e^2)*x^5 + ((2*A*
b^2*c^3 - (2*B*a*b^2 - (A - B)*b^3)*c^2)*d^2 + 2*(B*a*b^3*c - (2*(A - B)*a
*b^2 + A*b^3)*c^2)*d*e - (2*B*a*b^4 - 2*A*a*b^2*c^2 - (6*B*a^2*b^2 + (A -
B)*a*b^3)*c)*e^2)*x^3 + ((2*A*a*b*c^3 - (2*B*a^2*b - (A - B)*a*b^2)*c^2)*d
^2 + 2*(B*a^2*b^2*c - (2*(A - B)*a^2*b + A*a*b^2)*c^2)*d*e - (2*B*a^2*b^3
- 2*A*a^2*b*c^2 - (6*B*a^3*b + (A - B)*a^2*b^2)*c)*e^2)*x + (((2*A*c^5 ...

```

### 3.25.6 Sympy [F]

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{(A + Bx^2)(d + ex^2)^2}{(a + bx^2 + cx^4)^{3/2}} dx$$

input `integrate((B*x**2+A)*(e*x**2+d)**2/(c*x**4+b*x**2+a)**(3/2),x)`

output `Integral((A + B*x**2)*(d + e*x**2)**2/(a + b*x**2 + c*x**4)**(3/2), x)`

## 3.25.7 Maxima [F]

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^2}{(cx^4 + bx^2 + a)^{3/2}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^2/(c*x^4 + b*x^2 + a)^(3/2), x)`

## 3.25.8 Giac [F]

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^2}{(cx^4 + bx^2 + a)^{3/2}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^2/(c*x^4 + b*x^2 + a)^(3/2), x)`

## 3.25.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^2}{(cx^4 + bx^2 + a)^{3/2}} dx$$

input `int(((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(3/2), x)`

output `int(((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(3/2), x)`

**3.26** 
$$\int \frac{(A+Bx^2)(d+ex^2)}{(a+bx^2+cx^4)^{3/2}} dx$$

3.26.1 Optimal result . . . . . 227  
 3.26.2 Mathematica [C] (verified) . . . . . 228  
 3.26.3 Rubi [A] (verified) . . . . . 229  
 3.26.4 Maple [A] (verified) . . . . . 231  
 3.26.5 Fricas [B] (verification not implemented) . . . . . 232  
 3.26.6 Sympy [F] . . . . . 233  
 3.26.7 Maxima [F] . . . . . 234  
 3.26.8 Giac [F] . . . . . 234  
 3.26.9 Mupad [F(-1)] . . . . . 234

**3.26.1 Optimal result**

Integrand size = 31, antiderivative size = 481

$$\int \frac{(A+Bx^2)(d+ex^2)}{(a+bx^2+cx^4)^{3/2}} dx =$$

$$\frac{x(aB(bd-2ae) - A(b^2d - 2acd - abe) - (Ac(bd - 2ae) - aB(2cd - be))x^2)}{a(b^2 - 4ac)\sqrt{a+bx^2+cx^4}}$$

$$- \frac{(Ac(bd - 2ae) - aB(2cd - be))x\sqrt{a+bx^2+cx^4}}{a\sqrt{c}(b^2 - 4ac)(\sqrt{a} + \sqrt{cx^2})}$$

$$+ \frac{(Ac(bd - 2ae) - aB(2cd - be))(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{4}\left(2 - \frac{b}{\sqrt{a\sqrt{c}}}\right)\right)}{a^{3/4}c^{3/4}(b^2 - 4ac)\sqrt{a+bx^2+cx^4}}$$

$$+ \frac{(\sqrt{a}B - A\sqrt{c})(\sqrt{cd} - \sqrt{ae})(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{4}\left(2 - \frac{b}{\sqrt{a\sqrt{c}}}\right)\right)}{2a^{3/4}(b - 2\sqrt{a}\sqrt{c})c^{3/4}\sqrt{a+bx^2+cx^4}}$$

---

3.26. 
$$\int \frac{(A+Bx^2)(d+ex^2)}{(a+bx^2+cx^4)^{3/2}} dx$$

output

```

-x*(a*B*(-2*a*e+b*d)-A*(-a*b*e-2*a*c*d+b^2*d)-(A*c*(-2*a*e+b*d)-a*B*(-b*e+
2*c*d))*x^2)/a/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^(1/2)-(A*c*(-2*a*e+b*d)-a*B*(-
b*e+2*c*d))*x*(c*x^4+b*x^2+a)^(1/2)/a/(-4*a*c+b^2)/c^(1/2)/(a^(1/2)+x^2*c^
(1/2))+(A*c*(-2*a*e+b*d)-a*B*(-b*e+2*c*d))*(cos(2*arctan(c^(1/4)*x/a^(1/4)
))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)
)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x
^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2))^2)^(1/2)/a^(3/4)/c^(3/4)/(-4*a*c+b^2)/(c
*x^4+b*x^2+a)^(1/2)+1/2*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*a
rctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(
2-b/a^(1/2)/c^(1/2))^(1/2))*(B*a^(1/2)-A*c^(1/2))*(-e*a^(1/2)+d*c^(1/2))*
(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2))^2)^(1/2)/a^(3/
4)/c^(3/4)/(b-2*a^(1/2)*c^(1/2))/(c*x^4+b*x^2+a)^(1/2)

```

### 3.26.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 11.66 (sec) , antiderivative size = 597, normalized size of antiderivative = 1.24

$$\int \frac{(A + Bx^2)(d + ex^2)}{(a + bx^2 + cx^4)^{3/2}} dx = \frac{4c\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}x(aB(-2ae + 2cdx^2 + b(d - ex^2)) + A(-b^2d + b(ae - cdx^2) +$$

input `Integrate[((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(3/2),x]`

output

```

(4*c*sqrt[c/(b + sqrt[b^2 - 4*a*c])]*x*(a*B*(-2*a*e + 2*c*d*x^2 + b*(d - e
*x^2)) + A*(-(b^2*d) + b*(a*e - c*d*x^2) + 2*a*c*(d + e*x^2))) + I*(-b + S
qrt[b^2 - 4*a*c])*(A*c*(b*d - 2*a*e) + a*B*(-2*c*d + b*e))*sqrt[(b + sqrt[
b^2 - 4*a*c] + 2*c*x^2)/(b + sqrt[b^2 - 4*a*c])]*sqrt[(2*b - 2*sqrt[b^2 -
4*a*c] + 4*c*x^2)/(b - sqrt[b^2 - 4*a*c])]*EllipticE[I*ArcSinh[Sqrt[2]*Sqr
t[c/(b + sqrt[b^2 - 4*a*c])]*x], (b + sqrt[b^2 - 4*a*c])/(b - sqrt[b^2 - 4
*a*c])] - I*(A*c*(-(b^2*d) + 4*a*c*d + b*sqrt[b^2 - 4*a*c]*d - 2*a*sqrt[b^
2 - 4*a*c]*e) + a*B*(b*(-b + sqrt[b^2 - 4*a*c])*e + c*(-2*sqrt[b^2 - 4*a*c
]*d + 4*a*e))*sqrt[(b + sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + sqrt[b^2 - 4*a*
c])]*sqrt[(2*b - 2*sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - sqrt[b^2 - 4*a*c])]*E
llipticF[I*ArcSinh[Sqrt[2]*sqrt[c/(b + sqrt[b^2 - 4*a*c])]*x], (b + sqrt[b
^2 - 4*a*c])/(b - sqrt[b^2 - 4*a*c])]/(4*a*c*(-b^2 + 4*a*c)*sqrt[c/(b + S
qrt[b^2 - 4*a*c])]*sqrt[a + b*x^2 + c*x^4])

```

---

3.26.  $\int \frac{(A+Bx^2)(d+ex^2)}{(a+bx^2+cx^4)^{3/2}} dx$

### 3.26.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 446, normalized size of antiderivative = 0.93, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.194$ , Rules used = {2206, 25, 1511, 27, 1416, 1509}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(A+Bx^2)(d+ex^2)}{(a+bx^2+cx^4)^{3/2}} dx \\
 & \quad \downarrow \text{2206} \\
 & \int \frac{-\frac{a(bBd-2Acd+Abe-2aBe)-(Ac(bd-2ae)-aB(2cd-be))x^2}{\sqrt{cx^4+bx^2+a}}}{a(b^2-4ac)} dx \\
 & \frac{x(-A(-abe-2acd+b^2d)-(x^2(Ac(bd-2ae)-aB(2cd-be))))+aB(bd-2ae)}{a(b^2-4ac)\sqrt{a+bx^2+cx^4}} \\
 & \quad \downarrow \text{25} \\
 & \int \frac{\frac{a(bBd-2Acd+Abe-2aBe)-(Ac(bd-2ae)-aB(2cd-be))x^2}{\sqrt{cx^4+bx^2+a}}}{a(b^2-4ac)} dx \\
 & \frac{x(-A(-abe-2acd+b^2d)-(x^2(Ac(bd-2ae)-aB(2cd-be))))+aB(bd-2ae)}{a(b^2-4ac)\sqrt{a+bx^2+cx^4}} \\
 & \quad \downarrow \text{1511} \\
 & \frac{\frac{\sqrt{a}(2\sqrt{a}\sqrt{c}+b)(\sqrt{a}B-A\sqrt{c})(\sqrt{cd}-\sqrt{ae})}{\sqrt{c}} \int \frac{1}{\sqrt{cx^4+bx^2+a}} dx + \frac{\sqrt{a}(-2aAce+abBe-2aBcd+Abcd)}{\sqrt{c}} \int \frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{a}\sqrt{cx^4+bx^2+a}} dx}{a(b^2-4ac)}}{x(-A(-abe-2acd+b^2d)-(x^2(Ac(bd-2ae)-aB(2cd-be))))+aB(bd-2ae)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\frac{\sqrt{a}(2\sqrt{a}\sqrt{c}+b)(\sqrt{a}B-A\sqrt{c})(\sqrt{cd}-\sqrt{ae})}{\sqrt{c}} \int \frac{1}{\sqrt{cx^4+bx^2+a}} dx + \frac{(-2aAce+abBe-2aBcd+Abcd)}{\sqrt{c}} \int \frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{cx^4+bx^2+a}} dx}{a(b^2-4ac)}}{x(-A(-abe-2acd+b^2d)-(x^2(Ac(bd-2ae)-aB(2cd-be))))+aB(bd-2ae)} \\
 & \quad \downarrow \text{1416}
 \end{aligned}$$

---

3.26.  $\int \frac{(A+Bx^2)(d+ex^2)}{(a+bx^2+cx^4)^{3/2}} dx$

$$\frac{(-2aAce+abBe-2aBcd+Abcd) \int \frac{\sqrt{a}-\sqrt{cx^2}}{\sqrt{cx^4+bx^2+a}} dx + \frac{\sqrt[4]{a}(2\sqrt{a}\sqrt{c}+b)(\sqrt{a}+\sqrt{cx^2})(\sqrt{a}B-A\sqrt{c})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}(\sqrt{cd}-\sqrt{ae}) \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt{a}}{\sqrt{a+bx^2+cx^4}}\right)\right)}{2c^{3/4}\sqrt{a+bx^2+cx^4}}}{\frac{x(-A(-abe-2acd+b^2d) - (x^2(AC(bd-2ae) - aB(2cd-be))) + aB(bd-2ae))}{a(b^2-4ac)\sqrt{a+bx^2+cx^4}}}$$

↓ 1509

$$\frac{\frac{\sqrt[4]{a}(2\sqrt{a}\sqrt{c}+b)(\sqrt{a}+\sqrt{cx^2})(\sqrt{a}B-A\sqrt{c})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}(\sqrt{cd}-\sqrt{ae}) \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt{a}}\right), \frac{1}{4}\left(2-\frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{2c^{3/4}\sqrt{a+bx^2+cx^4}} + \frac{\left(\frac{\sqrt[4]{a}(\sqrt{a}+\sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}}{\sqrt{a+bx^2+cx^4}}\right)}{a(b^2-4ac)}}{x(-A(-abe-2acd+b^2d) - (x^2(AC(bd-2ae) - aB(2cd-be))) + aB(bd-2ae))}{a(b^2-4ac)\sqrt{a+bx^2+cx^4}}}$$

input `Int[(A + B*x^2)*(d + e*x^2)/(a + b*x^2 + c*x^4)^(3/2), x]`

output `-((x*(a*B*(b*d - 2*a*e) - A*(b^2*d - 2*a*c*d - a*b*e) - (A*c*(b*d - 2*a*e) - a*B*(2*c*d - b*e))*x^2))/(a*(b^2 - 4*a*c)*Sqrt[a + b*x^2 + c*x^4])) + ((A*b*c*d - 2*a*B*c*d + a*b*B*e - 2*a*A*c*e)*(-(x*Sqrt[a + b*x^2 + c*x^4])/(Sqrt[a] + Sqrt[c]*x^2)) + (a^(1/4)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(c^(1/4)*Sqrt[a + b*x^2 + c*x^4])))/Sqrt[c] + (a^(1/4)*(b + 2*Sqrt[a]*Sqrt[c])*(Sqrt[a]*B - A*Sqrt[c])*(Sqrt[c]*d - Sqrt[a]*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*c^(3/4)*Sqrt[a + b*x^2 + c*x^4]))/(a*(b^2 - 4*a*c))`

### 3.26.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

3.26.  $\int \frac{(A+Bx^2)(d+ex^2)}{(a+bx^2+cx^4)^{3/2}} dx$

rule 1416 `Int[1/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`

rule 1509 `Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`

rule 1511 `Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 2]}, Simp[(e + d*q)/q Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Simp[e/q Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`

rule 2206 `Int[(Px_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := With[{d = Coeff[PolynomialRemainder[Px, a + b*x^2 + c*x^4, x], x, 0], e = Coeff[PolynomialRemainder[Px, a + b*x^2 + c*x^4, x], x, 2]}, Simp[x*(a + b*x^2 + c*x^4)^(p + 1)*((a*b*e - d*(b^2 - 2*a*c) - c*(b*d - 2*a*e)*x^2)/(2*a*(p + 1)*(b^2 - 4*a*c))), x] + Simp[1/(2*a*(p + 1)*(b^2 - 4*a*c)) Int[(a + b*x^2 + c*x^4)^(p + 1)*ExpandToSum[2*a*(p + 1)*(b^2 - 4*a*c)*PolynomialQuotient[Px, a + b*x^2 + c*x^4, x] + b^2*d*(2*p + 3) - 2*a*c*d*(4*p + 5) - a*b*e + c*(4*p + 7)*(b*d - 2*a*e)*x^2, x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Px, x^2] && Expon[Px, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1]`

### 3.26.4 Maple [A] (verified)

Time = 1.67 (sec) , antiderivative size = 593, normalized size of antiderivative = 1.23

| method   | result   |
|----------|--|
| elliptic | $-\frac{2c \left( -\frac{(2Aace - Abcd - Babe + 2Bacd)x^3 - (Aabe + 2Acda - Ab^2d - 2eBa^2 + Babd)x}{2ca(4ac - b^2)} \right)}{\sqrt{\left(x^4 + \frac{bx^2}{c} + \frac{a}{c}\right)c}} + \frac{\left(\frac{eB}{c} + \frac{Acd - Bae}{ac} - \frac{Aabe + 2Acda - Ab^2d - 2eBa^2 + Babd}{a(4ac - b^2)}\right)}{\sqrt{\left(x^4 + \frac{bx^2}{c} + \frac{a}{c}\right)c}}$ |
| default  | Expression too large to display  |

3.26.  $\int \frac{(A+Bx^2)(d+ex^2)}{(a+bx^2+cx^4)^{3/2}} dx$



input `int((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -2*c*(-1/2/c*(2*A*a*c*e-A*b*c*d-B*a*b*e+2*B*a*c*d)/a/(4*a*c-b^2)*x^3-1/2/c \\ & *(A*a*b*e+2*A*a*c*d-A*b^2*d-2*B*a^2*e+B*a*b*d)/a/(4*a*c-b^2)*x)/((x^4+1/c* \\ & b*x^2+a/c)*c)^(1/2)+1/4*(e*B/c+(A*c*d-B*a*e)/a/c-(A*a*b*e+2*A*a*c*d-A*b^2* \\ & d-2*B*a^2*e+B*a*b*d)/a/(4*a*c-b^2))*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2) \\ & *(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/ \\ & a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2) \\ & ^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))+1/2*(2*A \\ & *a*c*e-A*b*c*d-B*a*b*e+2*B*a*c*d)/(4*a*c-b^2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2) \\ & )/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2) \\ & ^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*(Ellip \\ & ticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a* \\ & c+b^2)^(1/2))/a/c)^(1/2))-EllipticE(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2) \\ & )/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))) \end{aligned}$$

### 3.26.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1031 vs.  $2(470) = 940$ .

Time = 0.10 (sec) , antiderivative size = 1031, normalized size of antiderivative = 2.14

$$\int \frac{(A+Bx^2)(d+ex^2)}{(a+bx^2+cx^4)^{3/2}} dx = \text{Too large to display}$$

input `integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fricas")`

output

```

1/2*(sqrt(1/2)*(((2*B*a*b - A*b^2)*c^2*d - (B*a*b^2*c - 2*A*a*b*c^2)*e)*x^
4 + (2*B*a^2*b - A*a*b^2)*c*d + ((2*B*a*b^2 - A*b^3)*c*d - (B*a*b^3 - 2*A*
a*b^2*c)*e)*x^2 - (B*a^2*b^2 - 2*A*a^2*b*c)*e - (((2*B*a^2 - A*a*b)*c^2*d
- (B*a^2*b*c - 2*A*a^2*c^2)*e)*x^4 + (2*B*a^3 - A*a^2*b)*c*d + ((2*B*a^2*b
- A*a*b^2)*c*d - (B*a^2*b^2 - 2*A*a^2*b*c)*e)*x^2 - (B*a^3*b - 2*A*a^3*c)
*e)*sqrt((b^2 - 4*a*c)/a^2))*sqrt(a)*sqrt((a*sqrt((b^2 - 4*a*c)/a^2) - b)/
a)*elliptic_e(arcsin(sqrt(1/2)*x*sqrt((a*sqrt((b^2 - 4*a*c)/a^2) - b)/a)),
1/2*(a*b*sqrt((b^2 - 4*a*c)/a^2) + b^2 - 2*a*c)/(a*c)) - sqrt(1/2)*(((B*a
*b^2*c - (2*(A - B)*a*b + A*b^2)*c^2)*d + (2*A*a*b*c^2 - (2*B*a^2*b - (A -
B)*a*b^2)*c)*e)*x^4 + ((B*a*b^3 - (2*(A - B)*a*b^2 + A*b^3)*c)*d - (2*B*a
^2*b^2 - (A - B)*a*b^3 - 2*A*a*b^2*c)*e)*x^2 + (B*a^2*b^2 - (2*(A - B)*a^2
*b + A*a*b^2)*c)*d - (2*B*a^3*b - (A - B)*a^2*b^2 - 2*A*a^2*b*c)*e + (((B*
a^2*b*c - (2*(A + B)*a^2 - A*a*b)*c^2)*d - (2*A*a^2*c^2 + (2*B*a^3 - (A +
B)*a^2*b)*c)*e)*x^4 + ((B*a^2*b^2 - (2*(A + B)*a^2*b - A*a*b^2)*c)*d - (2*
B*a^3*b - (A + B)*a^2*b^2 + 2*A*a^2*b*c)*e)*x^2 + (B*a^3*b - (2*(A + B)*a^
3 - A*a^2*b)*c)*d - (2*B*a^4 - (A + B)*a^3*b + 2*A*a^3*c)*e)*sqrt((b^2 - 4
*a*c)/a^2))*sqrt(a)*sqrt((a*sqrt((b^2 - 4*a*c)/a^2) - b)/a)*elliptic_f(arc
sin(sqrt(1/2)*x*sqrt((a*sqrt((b^2 - 4*a*c)/a^2) - b)/a)), 1/2*(a*b*sqrt((b
^2 - 4*a*c)/a^2) + b^2 - 2*a*c)/(a*c)) - 2*sqrt(c*x^4 + b*x^2 + a)*(((2*B*
a^2 - A*a*b)*c^2*d - (B*a^2*b*c - 2*A*a^2*c^2)*e)*x^3 - ((2*B*a^3 - A*a...

```

### 3.26.6 Sympy [F]

$$\int \frac{(A + Bx^2)(d + ex^2)}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{(A + Bx^2)(d + ex^2)}{(a + bx^2 + cx^4)^{\frac{3}{2}}} dx$$

input `integrate((B*x**2+A)*(e*x**2+d)/(c*x**4+b*x**2+a)**(3/2), x)`

output `Integral((A + B*x**2)*(d + e*x**2)/(a + b*x**2 + c*x**4)**(3/2), x)`

**3.26.7 Maxima [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)}{(cx^4 + bx^2 + a)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)*(e*x^2 + d)/(c*x^4 + b*x^2 + a)^(3/2), x)`

**3.26.8 Giac [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)}{(cx^4 + bx^2 + a)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)*(e*x^2 + d)/(c*x^4 + b*x^2 + a)^(3/2), x)`

**3.26.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{(Bx^2 + A)(ex^2 + d)}{(cx^4 + bx^2 + a)^{3/2}} dx$$

input `int(((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(3/2),x)`

output `int(((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(3/2), x)`

### 3.27 $\int \frac{A+Bx^2}{(a+bx^2+cx^4)^{3/2}} dx$

|        |   |     |
|--------|---|-----|
| 3.27.1 | Optimal result                            | 235 |
| 3.27.2 | Mathematica [C] (verified)                | 236 |
| 3.27.3 | Rubi [A] (verified)                       | 236 |
| 3.27.4 | Maple [A] (verified)                      | 239 |
| 3.27.5 | Fricas [A] (verification not implemented) | 240 |
| 3.27.6 | Sympy [F]                                 | 240 |
| 3.27.7 | Maxima [F]                                | 241 |
| 3.27.8 | Giac [F]                                  | 241 |
| 3.27.9 | Mupad [F(-1)]                             | 241 |

#### 3.27.1 Optimal result

Integrand size = 24, antiderivative size = 398

$$\int \frac{A + Bx^2}{(a + bx^2 + cx^4)^{3/2}} dx = \frac{x(Ab^2 - abB - 2aAc + (Ab - 2aB)cx^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} - \frac{(Ab - 2aB)\sqrt{cx}\sqrt{a + bx^2 + cx^4}}{a(b^2 - 4ac)(\sqrt{a} + \sqrt{cx^2})} + \frac{(Ab - 2aB)\sqrt[4]{c}(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{a^{3/4}(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} + \frac{(\sqrt{a}B - A\sqrt{c})(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{cx}}{\sqrt[4]{a}}\right), \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{2a^{3/4}(b - 2\sqrt{a}\sqrt{c})\sqrt[4]{c}\sqrt{a + bx^2 + cx^4}}$$

```
output x*(A*b^2-a*b*B-2*a*A*c+(A*b-2*B*a)*c*x^2)/a/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^(1/2)-(A*b-2*B*a)*x*c^(1/2)*(c*x^4+b*x^2+a)^(1/2)/a/(-4*a*c+b^2)/(a^(1/2)+x^2*c^(1/2))+(A*b-2*B*a)*c^(1/4)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(3/4)/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^(1/2)+1/2*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(B*a^(1/2)-A*c^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(3/4)/c^(1/4)/(b-2*a^(1/2)*c^(1/2))/(c*x^4+b*x^2+a)^(1/2)
```

### 3.27.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.91 (sec) , antiderivative size = 497, normalized size of antiderivative = 1.25

$$\int \frac{A + Bx^2}{(a + bx^2 + cx^4)^{3/2}} dx =$$

$$4\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}x(aB(b+2cx^2) - A(b^2 - 2ac + bcx^2)) + i(Ab - 2aB) (-b + \sqrt{b^2 - 4ac}) \sqrt{\frac{b+\sqrt{b^2-4ac}+2cx^2}{b+\sqrt{b^2-4ac}}}$$

input `Integrate[(A + B*x^2)/(a + b*x^2 + c*x^4)^(3/2),x]`

output `-1/4*(4*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])] * x*(a*B*(b + 2*c*x^2) - A*(b^2 - 2*a*c + b*c*x^2)) + I*(A*b - 2*a*B)*(-b + Sqrt[b^2 - 4*a*c])*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])] * Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])] * EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] - I*(-2*a*B*Sqrt[b^2 - 4*a*c] + A*(-b^2 + 4*a*c + b*Sqrt[b^2 - 4*a*c])) * Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])] * Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])] * EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])])/(a*(b^2 - 4*a*c)*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])])*Sqrt[a + b*x^2 + c*x^4])`

### 3.27.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 382, normalized size of antiderivative = 0.96, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {1492, 25, 1511, 27, 1416, 1509}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(a + bx^2 + cx^4)^{3/2}} dx$$

↓ 1492

$$\begin{aligned}
& \frac{x(cx^2(Ab - 2aB) - 2aAc - abB + Ab^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} - \frac{\int -\frac{a(bB - 2Ac) - (Ab - 2aB)cx^2}{\sqrt{cx^4 + bx^2 + a}} dx}{a(b^2 - 4ac)} \\
& \quad \downarrow \text{25} \\
& \frac{\int \frac{a(bB - 2Ac) - (Ab - 2aB)cx^2}{\sqrt{cx^4 + bx^2 + a}} dx}{a(b^2 - 4ac)} + \frac{x(cx^2(Ab - 2aB) - 2aAc - abB + Ab^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} \\
& \quad \downarrow \text{1511} \\
& \frac{\sqrt{a}\sqrt{c}(Ab - 2aB) \int \frac{\sqrt{a} - \sqrt{cx^2}}{\sqrt{a}\sqrt{cx^4 + bx^2 + a}} dx - \sqrt{a}(\sqrt{c}(Ab - 2aB) - \sqrt{a}(bB - 2Ac)) \int \frac{1}{\sqrt{cx^4 + bx^2 + a}} dx}{a(b^2 - 4ac)} + \\
& \quad \frac{x(cx^2(Ab - 2aB) - 2aAc - abB + Ab^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} \\
& \quad \downarrow \text{27} \\
& \frac{\sqrt{c}(Ab - 2aB) \int \frac{\sqrt{a} - \sqrt{cx^2}}{\sqrt{cx^4 + bx^2 + a}} dx - \sqrt{a}(\sqrt{c}(Ab - 2aB) - \sqrt{a}(bB - 2Ac)) \int \frac{1}{\sqrt{cx^4 + bx^2 + a}} dx}{a(b^2 - 4ac)} + \\
& \quad \frac{x(cx^2(Ab - 2aB) - 2aAc - abB + Ab^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} \\
& \quad \downarrow \text{1416} \\
& \frac{\sqrt{c}(Ab - 2aB) \int \frac{\sqrt{a} - \sqrt{cx^2}}{\sqrt{cx^4 + bx^2 + a}} dx - \frac{\sqrt[4]{a}(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} (\sqrt{c}(Ab - 2aB) - \sqrt{a}(bB - 2Ac)) \text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt{a}}\right), \frac{1}{4}\left(2 - \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}}\right)\right)}{2\sqrt[4]{c}\sqrt{a + bx^2 + cx^4}}}{a(b^2 - 4ac)} \\
& \quad \frac{x(cx^2(Ab - 2aB) - 2aAc - abB + Ab^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} \\
& \quad \downarrow \text{1509} \\
& \frac{\sqrt{c}(Ab - 2aB) \left( \frac{\sqrt[4]{a}(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt{a}}\right) \middle| \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{\sqrt[4]{c}\sqrt{a + bx^2 + cx^4}} - \frac{x\sqrt{a + bx^2 + cx^4}}{\sqrt{a} + \sqrt{cx^2}} \right) - \frac{\sqrt[4]{a}(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}}}{\sqrt{a} + \sqrt{cx^2}}}{a(b^2 - 4ac)} \\
& \quad \frac{x(cx^2(Ab - 2aB) - 2aAc - abB + Ab^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}}
\end{aligned}$$

input `Int[(A + B*x^2)/(a + b*x^2 + c*x^4)^(3/2), x]`

```
output (x*(A*b^2 - a*b*B - 2*a*A*c + (A*b - 2*a*B)*c*x^2))/(a*(b^2 - 4*a*c)*Sqrt[
a + b*x^2 + c*x^4]) + ((A*b - 2*a*B)*Sqrt[c]*(-(x*Sqrt[a + b*x^2 + c*x^4]
)/(Sqrt[a] + Sqrt[c]*x^2)) + (a^(1/4)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*
x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(
1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(c^(1/4)*Sqrt[a + b*x^2 + c*x^4])) -
(a^(1/4)*((A*b - 2*a*B)*Sqrt[c] - Sqrt[a]*(b*B - 2*A*c))*(Sqrt[a] + Sqrt[c]
]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*Arc
Tan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*c^(1/4)*Sqrt[a
+ b*x^2 + c*x^4]))/(a*(b^2 - 4*a*c))
```

### 3.27.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 1416 Int[1/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c
/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/
(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))
], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

```
rule 1492 Int[((d_) + (e_)*(x_)^2)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symb
ol] := Simp[x*(a*b*e - d*(b^2 - 2*a*c) - c*(b*d - 2*a*e)*x^2)*((a + b*x^2 +
c*x^4)^(p + 1)/(2*a*(p + 1)*(b^2 - 4*a*c))), x] + Simp[1/(2*a*(p + 1)*(b^2
- 4*a*c)) Int[Simp[(2*p + 3)*d*b^2 - a*b*e - 2*a*c*d*(4*p + 5) + (4*p +
7)*(d*b - 2*a*e)*c*x^2, x]*(a + b*x^2 + c*x^4)^(p + 1), x], x] /; FreeQ[{a,
b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] &&
LtQ[p, -1] && IntegerQ[2*p]
```

```
rule 1509 Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symb
ol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q
^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*
x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2
/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2
- 4*a*c, 0] && PosQ[c/a]
```

```
rule 1511 Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol]
:= With[{q = Rt[c/a, 2]}, Simp[(e + d*q)/q Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Simp[e/q Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /;
NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

### 3.27.4 Maple [A] (verified)

Time = 1.04 (sec) , antiderivative size = 509, normalized size of antiderivative = 1.28

| method   | result   |
|----------|--|
| elliptic | $-\frac{2c \left( \frac{(Ab-2Ba)x^3}{2a(4ac-b^2)} - \frac{(2Aac-Ab^2+abB)x}{2a(4ac-b^2)} \right)}{\sqrt{\left(x^4 + \frac{bx^2}{c} + \frac{a}{c}\right)c}} + \frac{\left(\frac{A}{a} - \frac{2Aac-Ab^2+abB}{(4ac-b^2)a}\right) \sqrt{2} \sqrt{4 - \frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}} \sqrt{4 + \frac{2(b+\sqrt{-4ac+b^2})x^2}{a}} F\left(x\sqrt{2}\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}\right)}{4\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}} \sqrt{cx^4+bx^2+a}}$ |
| default  | $A \left( -\frac{2c \left( \frac{bx^3}{2a(4ac-b^2)} - \frac{(2ac-b^2)x}{2a(4ac-b^2)c} \right)}{\sqrt{\left(x^4 + \frac{bx^2}{c} + \frac{a}{c}\right)c}} + \frac{\left(\frac{1}{a} - \frac{2ac-b^2}{a(4ac-b^2)}\right) \sqrt{2} \sqrt{4 - \frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}} \sqrt{4 + \frac{2(b+\sqrt{-4ac+b^2})x^2}{a}} F\left(x\sqrt{2}\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}\right)}{4\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}} \sqrt{cx^4+bx^2+a}} \right)$  |

```
input int((B*x^2+A)/(c*x^4+b*x^2+a)^(3/2),x,method=_RETURNVERBOSE)
```

```
output -2*c*(1/2*(A*b-2*B*a)/a/(4*a*c-b^2)*x^3-1/2*(2*A*a*c-A*b^2+B*a*b)/a/c/(4*a*c-b^2)*x)/((x^4+1/c*b*x^2+a/c)*c)^(1/2)+1/4*(A/a-(2*A*a*c-A*b^2+B*a*b)/(4*a*c-b^2)/a)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*x^2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))-1/2*c*(A*b-2*B*a)/(4*a*c-b^2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*(EllipticF(1/2*x^2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))-EllipticE(1/2*x^2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2)))
```

3.27.  $\int \frac{A+Bx^2}{(a+bx^2+cx^4)^{3/2}} dx$



### 3.27.5 Fracas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 656, normalized size of antiderivative = 1.65

$$\int \frac{A + Bx^2}{(a + bx^2 + cx^4)^{3/2}} dx = \frac{\sqrt{\frac{1}{2}} \left( (2 Bab - Ab^2)c^2x^4 + (2 Bab^2 - Ab^3)cx^2 + (2 Ba^2b - Aab^2)c - ((2 Ba^2 - A$$

```
input integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fracas")
```

```
output 1/2*(sqrt(1/2)*((2*B*a*b - A*b^2)*c^2*x^4 + (2*B*a*b^2 - A*b^3)*c*x^2 + (2
*B*a^2*b - A*a*b^2)*c - ((2*B*a^2 - A*a*b)*c^2*x^4 + (2*B*a^2*b - A*a*b^2)
*c*x^2 + (2*B*a^3 - A*a^2*b)*c)*sqrt((b^2 - 4*a*c)/a^2))*sqrt(a)*sqrt((a*s
qrt((b^2 - 4*a*c)/a^2) - b)/a)*elliptic_e(arcsin(sqrt(1/2)*x*sqrt((a*sqrt(
(b^2 - 4*a*c)/a^2) - b)/a)), 1/2*(a*b*sqrt((b^2 - 4*a*c)/a^2) + b^2 - 2*a*
c)/(a*c)) - sqrt(1/2)*(B*a^2*b^2 + (B*a*b^2*c - (2*(A - B)*a*b + A*b^2)*c^
2)*x^4 + (B*a*b^3 - (2*(A - B)*a*b^2 + A*b^3)*c)*x^2 - (2*(A - B)*a^2*b +
A*a*b^2)*c + (B*a^3*b + (B*a^2*b*c - (2*(A + B)*a^2 - A*a*b)*c^2)*x^4 + (B
*a^2*b^2 - (2*(A + B)*a^2*b - A*a*b^2)*c)*x^2 - (2*(A + B)*a^3 - A*a^2*b)*
c)*sqrt((b^2 - 4*a*c)/a^2))*sqrt(a)*sqrt((a*sqrt((b^2 - 4*a*c)/a^2) - b)/a
)*elliptic_f(arcsin(sqrt(1/2)*x*sqrt((a*sqrt((b^2 - 4*a*c)/a^2) - b)/a)),
1/2*(a*b*sqrt((b^2 - 4*a*c)/a^2) + b^2 - 2*a*c)/(a*c)) - 2*((2*B*a^2 - A*a
*b)*c^2*x^3 + (2*A*a^2*c^2 + (B*a^2*b - A*a*b^2)*c)*x)*sqrt(c*x^4 + b*x^2
+ a))/(a^3*b^2*c - 4*a^4*c^2 + (a^2*b^2*c^2 - 4*a^3*c^3)*x^4 + (a^2*b^3*c
- 4*a^3*b*c^2)*x^2)
```

### 3.27.6 Sympy [F]

$$\int \frac{A + Bx^2}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{A + Bx^2}{(a + bx^2 + cx^4)^{\frac{3}{2}}} dx$$

```
input integrate((B*x**2+A)/(c*x**4+b*x**2+a)**(3/2),x)
```

```
output Integral((A + B*x**2)/(a + b*x**2 + c*x**4)**(3/2), x)
```

**3.27.7 Maxima [F]**

$$\int \frac{A + Bx^2}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + bx^2 + a)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/(c*x^4 + b*x^2 + a)^(3/2), x)`

**3.27.8 Giac [F]**

$$\int \frac{A + Bx^2}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + bx^2 + a)^{\frac{3}{2}}} dx$$

input `integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/(c*x^4 + b*x^2 + a)^(3/2), x)`

**3.27.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + bx^2 + a)^{3/2}} dx$$

input `int((A + B*x^2)/(a + b*x^2 + c*x^4)^(3/2),x)`

output `int((A + B*x^2)/(a + b*x^2 + c*x^4)^(3/2), x)`

$$3.28 \quad \int \frac{A+Bx^2}{(d+ex^2)(a+bx^2+cx^4)^{3/2}} dx$$

|        |                            |     |
|--------|----------------------------|-----|
| 3.28.1 | Optimal result             | 242 |
| 3.28.2 | Mathematica [C] (verified) | 243 |
| 3.28.3 | Rubi [A] (verified)        | 244 |
| 3.28.4 | Maple [B] (verified)       | 246 |
| 3.28.5 | Fricas [F(-1)]             | 247 |
| 3.28.6 | Sympy [F(-1)]              | 248 |
| 3.28.7 | Maxima [F]                 | 248 |
| 3.28.8 | Giac [F]                   | 248 |
| 3.28.9 | Mupad [F(-1)]              | 249 |

### 3.28.1 Optimal result

Integrand size = 33, antiderivative size = 867

$$\int \frac{A+Bx^2}{(d+ex^2)(a+bx^2+cx^4)^{3/2}} dx =$$

$$\frac{x(abc(Bd - Ae) - (b^2 - 2ac)(Acd - Abe + aBe) + c(aB(2cd - be) - A(bcd - b^2e + 2ace))x^2)}{a(b^2 - 4ac)(cd^2 - bde + ae^2)\sqrt{a+bx^2+cx^4}}$$

$$+ \frac{\sqrt{c}(aB(2cd - be) - A(bcd - b^2e + 2ace))x\sqrt{a+bx^2+cx^4}}{a(b^2 - 4ac)(cd^2 - bde + ae^2)(\sqrt{a} + \sqrt{cx^2})}$$

$$- \frac{e^{3/2}(Bd - Ae) \arctan\left(\frac{\sqrt{cd^2 - bde + ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{a+bx^2+cx^4}}\right)}{2\sqrt{d}(cd^2 - bde + ae^2)^{3/2}}$$

$$- \frac{\sqrt[4]{c}(aB(2cd - be) - A(bcd - b^2e + 2ace))(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}}E\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt{a}}\right) \middle| \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{a^{3/4}(b^2 - 4ac)(cd^2 - bde + ae^2)\sqrt{a+bx^2+cx^4}}$$

$$+ \frac{(\sqrt{a}B - A\sqrt{c})\sqrt[4]{c}(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}}\text{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt{a}}\right), \frac{1}{2} - \frac{b}{4\sqrt{a}\sqrt{c}}\right)}{2a^{3/4}(b - 2\sqrt{a}\sqrt{c})(\sqrt{cd} - \sqrt{ae})\sqrt{a+bx^2+cx^4}}$$

$$+ \frac{a^{3/4}e\left(\frac{\sqrt{cd}}{\sqrt{a}} + e\right)^2(Bd - Ae)(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}}\text{EllipticPi}\left(-\frac{(\sqrt{cd} - \sqrt{ae})^2}{4\sqrt{a}\sqrt{cde}}, 2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt{a}}\right), \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{4\sqrt[4]{cd}(cd^2 - ae^2)(cd^2 - bde + ae^2)\sqrt{a+bx^2+cx^4}}$$

---

3.28.  $\int \frac{A+Bx^2}{(d+ex^2)(a+bx^2+cx^4)^{3/2}} dx$

output

```
-1/2*e^(3/2)*(-A*e+B*d)*arctan(x*(a*e^2-b*d*e+c*d^2)^(1/2)/d^(1/2)/e^(1/2)
/(c*x^4+b*x^2+a)^(1/2))/(a*e^2-b*d*e+c*d^2)^(3/2)/d^(1/2)-x*(a*b*c*(-A*e+B
*d)-(-2*a*c+b^2)*(-A*b*e+A*c*d+B*a*e)+c*(a*B*(-b*e+2*c*d)-A*(2*a*c*e-b^2*e
+b*c*d))*x^2)/a/(-4*a*c+b^2)/(a*e^2-b*d*e+c*d^2)/(c*x^4+b*x^2+a)^(1/2)+(a*
B*(-b*e+2*c*d)-A*(2*a*c*e-b^2*e+b*c*d))*x*c^(1/2)*(c*x^4+b*x^2+a)^(1/2)/a/
(-4*a*c+b^2)/(a*e^2-b*d*e+c*d^2)/(a^(1/2)+x^2*c^(1/2))-c^(1/4)*(a*B*(-b*e+
2*c*d)-A*(2*a*c*e-b^2*e+b*c*d))*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)
/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4)
)),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)
/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(3/4)/(-4*a*c+b^2)/(a*e^2-b*d*e+c*d^2)/(
c*x^4+b*x^2+a)^(1/2)+1/4*a^(3/4)*e*(-A*e+B*d)*(cos(2*arctan(c^(1/4)*x/a^(1
/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticPi(sin(2*arctan(c^
(1/4)*x/a^(1/4))),-1/4*(-e*a^(1/2)+d*c^(1/2))^2/d/e/a^(1/2)/c^(1/2),1/2*(2
-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*(e+d*c^(1/2)/a^(1/2))^2*(
(c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(1/4)/d/(-a*e^2+c*d^2)/(a
*e^2-b*d*e+c*d^2)/(c*x^4+b*x^2+a)^(1/2)+1/2*c^(1/4)*(cos(2*arctan(c^(1/4)*
x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arct
an(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(B*a^(1/2)-A*c^(1/
2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/
a^(3/4)/(-e*a^(1/2)+d*c^(1/2))/(b-2*a^(1/2)*c^(1/2))/(c*x^4+b*x^2+a)^(1...
```

### 3.28.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 13.30 (sec) , antiderivative size = 1736, normalized size of antiderivative = 2.00

$$\int \frac{A + Bx^2}{(d + ex^2)(a + bx^2 + cx^4)^{3/2}} dx = \text{Too large to display}$$

input `Integrate[(A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(3/2)),x]`

output

```
(-4*A*b^2*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d^2*x + 4*a*b*B*c*Sqrt[c/(b +
Sqrt[b^2 - 4*a*c])]*d^2*x + 8*a*A*c^2*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d^2*
x + 4*A*b^3*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x - 4*a*b^2*B*Sqrt[c/(b +
Sqrt[b^2 - 4*a*c])]*d*e*x - 12*a*A*b*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e
*x + 8*a^2*B*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x - 4*A*b*c^2*Sqrt[c/(b
+ Sqrt[b^2 - 4*a*c])]*d^2*x^3 + 8*a*B*c^2*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])])
*d^2*x^3 + 4*A*b^2*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x^3 - 4*a*b*B*c*S
qrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x^3 - 8*a*A*c^2*Sqrt[c/(b + Sqrt[b^2 -
4*a*c])]*d*e*x^3 - I*(-b + Sqrt[b^2 - 4*a*c])*d*(a*B*(2*c*d - b*e) + A*(-(
b*c*d) + b^2*e - 2*a*c*e))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqr
t[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2
- 4*a*c])]*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x],
(b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] + I*d*(a*B*(b*(b - Sqrt[
b^2 - 4*a*c])*e + 2*c*(Sqrt[b^2 - 4*a*c]*d - 2*a*e)) + A*(-(b^3*e) + b*c*(
-(Sqrt[b^2 - 4*a*c]*d) + 4*a*e) + b^2*(c*d + Sqrt[b^2 - 4*a*c]*e) - 2*a*c*
(2*c*d + Sqrt[b^2 - 4*a*c]*e))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b
+ Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt
[b^2 - 4*a*c])]*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])
]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] - (2*I)*a*b^2*B*d*e
*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(...
```

### 3.28.3 Rubi [A] (verified)

Time = 1.57 (sec) , antiderivative size = 1045, normalized size of antiderivative = 1.21, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.061$ , Rules used = {2258, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(d + ex^2)(a + bx^2 + cx^4)^{3/2}} dx$$

↓ 2258

$$\int \left( \frac{e(Ae - Bd)}{(d + ex^2)\sqrt{a + bx^2 + cx^4}(ae^2 - bde + cd^2)} + \frac{aBe - Abe + cx^2(Bd - Ae) + Acd}{(a + bx^2 + cx^4)^{3/2}(ae^2 - bde + cd^2)} \right) dx$$

↓ 2009

---

3.28.  $\int \frac{A+Bx^2}{(d+ex^2)(a+bx^2+cx^4)^{3/2}} dx$

$$\begin{aligned}
& \frac{a^{3/4}e(Bd - Ae) (\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4+bx^2+a}{(\sqrt{cx^2+\sqrt{a}})^2}} \operatorname{EllipticPi} \left( -\frac{(\sqrt{cd}-\sqrt{ae})^2}{4\sqrt{a}\sqrt{cde}}, 2 \arctan \left( \frac{\sqrt[4]{Cx}}{\sqrt[4]{a}} \right), \frac{1}{4} \left( 2 - \frac{b}{\sqrt{a}\sqrt{c}} \right) \right) (\frac{\sqrt{cd}}{\sqrt{a}} + e)}{4\sqrt[4]{cd} (cd^2 - ae^2) (cd^2 - bed + ae^2) \sqrt{cx^4 + bx^2 + a}} \\
& - \frac{e^{3/2}(Bd - Ae) \arctan \left( \frac{\sqrt{cd^2-bed+ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{cx^4+bx^2+a}} \right)}{2\sqrt{d} (cd^2 - bed + ae^2)^{3/2}} \\
& \frac{\sqrt[4]{c}(aB(2cd - be) - A(-eb^2 + cdb + 2ace)) (\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4+bx^2+a}{(\sqrt{cx^2+\sqrt{a}})^2}} E \left( 2 \arctan \left( \frac{\sqrt[4]{Cx}}{\sqrt[4]{a}} \right) \middle| \frac{1}{4} \left( 2 - \frac{b}{\sqrt{a}\sqrt{c}} \right) \right)}{a^{3/4} (b^2 - 4ac) (cd^2 - bed + ae^2) \sqrt{cx^4 + bx^2 + a}} \\
& \frac{\sqrt[4]{c}(aBe - \sqrt{a}\sqrt{c}(Bd - Ae) + A(cd - be)) (\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4+bx^2+a}{(\sqrt{cx^2+\sqrt{a}})^2}} \operatorname{EllipticF} \left( 2 \arctan \left( \frac{\sqrt[4]{Cx}}{\sqrt[4]{a}} \right), \frac{1}{4} \left( 2 - \frac{b}{\sqrt{a}\sqrt{c}} \right) \right)}{2a^{3/4} (b - 2\sqrt{a}\sqrt{c}) (cd^2 - bed + ae^2) \sqrt{cx^4 + bx^2 + a}} \\
& + \frac{\sqrt[4]{ce}(Bd - Ae) (\sqrt{cx^2 + \sqrt{a}}) \sqrt{\frac{cx^4+bx^2+a}{(\sqrt{cx^2+\sqrt{a}})^2}} \operatorname{EllipticF} \left( 2 \arctan \left( \frac{\sqrt[4]{Cx}}{\sqrt[4]{a}} \right), \frac{1}{4} \left( 2 - \frac{b}{\sqrt{a}\sqrt{c}} \right) \right)}{2\sqrt[4]{a} (\sqrt{cd} - \sqrt{ae}) (cd^2 - bed + ae^2) \sqrt{cx^4 + bx^2 + a}} \\
& - \frac{\sqrt{c}(aB(2cd - be) - A(-eb^2 + cdb + 2ace)) x \sqrt{cx^4 + bx^2 + a}}{a (b^2 - 4ac) (cd^2 - bed + ae^2) (\sqrt{cx^2 + \sqrt{a}})} \\
& \frac{x(c(aB(2cd - be) - A(-eb^2 + cdb + 2ace)) x^2 + abc(Bd - Ae) - (b^2 - 2ac) (Acd - Abe + aBe))}{a (b^2 - 4ac) (cd^2 - bed + ae^2) \sqrt{cx^4 + bx^2 + a}}
\end{aligned}$$

input `Int[(A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(3/2)), x]`

output

$$\begin{aligned}
& -((x*(a*b*c*(B*d - A*e) - (b^2 - 2*a*c)*(A*c*d - A*b*e + a*B*e) + c*(a*B*(2*c*d - b*e) - A*(b*c*d - b^2*e + 2*a*c*e))*x^2))/(a*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)*\text{Sqrt}[a + b*x^2 + c*x^4]) + (\text{Sqrt}[c]*(a*B*(2*c*d - b*e) - A*(b*c*d - b^2*e + 2*a*c*e))*x*\text{Sqrt}[a + b*x^2 + c*x^4])/(a*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) - (e^{3/2}*(B*d - A*e)*\text{ArcTan}[(\text{Sqrt}[c*d^2 - b*d*e + a*e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + b*x^2 + c*x^4])])/(2*\text{Sqrt}[d]*(c*d^2 - b*d*e + a*e^2)^{3/2}) - (c^{1/4}*(a*B*(2*c*d - b*e) - A*(b*c*d - b^2*e + 2*a*c*e))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[c^{1/4}*x/a^{1/4}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(a^{3/4}*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)*\text{Sqrt}[a + b*x^2 + c*x^4]) - (c^{1/4}*e*(B*d - A*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{1/4}*x)/a^{1/4}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(2*a^{1/4}*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 - b*d*e + a*e^2)*\text{Sqrt}[a + b*x^2 + c*x^4]) - (c^{1/4}*(a*B*e - \text{Sqrt}[a]*\text{Sqrt}[c]*(B*d - A*e) + A*(c*d - b*e))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{1/4}*x)/a^{1/4}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(2*a^{3/4}*(b - 2*\text{Sqrt}[a]*\text{Sqrt}[c])*(c*d^2 - b*d*e + a*e^2)*\text{Sqrt}[a + b*x^2 + c*x^4]) + (a^{3/4}*e*((\text{Sqrt}[c]*d)/\text{Sqrt}[a] + e)^2*(B*d - A*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticPi}[-1/4*(\text{Sqrt}[c]...
\end{aligned}$$

### 3.28.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2258 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[1/Sqrt[a + b*x^2 + c*x^4], Px*(d + e*x^2)^q*(a + b*x^2 + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e}, x] && PolyQ[Px, x] && IntegerQ[p + 1/2] && IntegerQ[q]`

### 3.28.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3240 vs.  $2(832) = 1664$ .

Time = 1.76 (sec) , antiderivative size = 3241, normalized size of antiderivative = 3.74

| method   | result                          | size |
|----------|---------------------------------|------|
| default  | Expression too large to display | 3241 |
| elliptic | Expression too large to display | 4597 |

$$3.28. \int \frac{A+Bx^2}{(d+ex^2)(a+bx^2+cx^4)^{3/2}} dx$$

input `int((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x,method=_RETURNVERBOSE)`

output 
$$\frac{B}{e} \cdot \frac{-2c \cdot \left(\frac{1}{2} \sqrt{\frac{b}{4ac-b^2}} x^3 - \frac{1}{2} \sqrt{\frac{2ac-b^2}{a(4ac-b^2)}} \frac{1}{cx}\right)}{\left(x^4 + \frac{1}{c} \sqrt{\frac{b}{4ac-b^2}} x^2 + \frac{a}{c}\right)^{1/2}} + \frac{1}{4} \cdot \frac{\left(\frac{1}{a} - \sqrt{\frac{2ac-b^2}{a(4ac-b^2)}}\right) x^2}{\left(-b + (-4ac+b^2)^{1/2}\right)^{1/2}} \cdot \frac{\left(4 - 2 \sqrt{\frac{-b + (-4ac+b^2)^{1/2}}{ax^2}}\right)^{1/2}}{\left(4 + 2 \sqrt{\frac{b + (-4ac+b^2)^{1/2}}{ax^2}}\right)^{1/2}} \cdot \frac{1}{\left(c x^4 + b x^2 + a\right)^{1/2}} \cdot \text{EllipticF}\left(\frac{1}{2} \sqrt{x^2} \cdot \frac{\left(-b + (-4ac+b^2)^{1/2}\right)^{1/2}}{a}, \frac{1}{2} \sqrt{-4 + 2 \sqrt{\frac{b + (-4ac+b^2)^{1/2}}{a/c}}}\right) - \frac{1}{2} \sqrt{\frac{b}{4ac-b^2}} \cdot \frac{x^2}{\left(-b + (-4ac+b^2)^{1/2}\right)^{1/2}} \cdot \frac{\left(4 - 2 \sqrt{\frac{-b + (-4ac+b^2)^{1/2}}{ax^2}}\right)^{1/2}}{\left(4 + 2 \sqrt{\frac{b + (-4ac+b^2)^{1/2}}{ax^2}}\right)^{1/2}} \cdot \frac{1}{\left(c x^4 + b x^2 + a\right)^{1/2}} \cdot \frac{1}{\left(b + (-4ac+b^2)^{1/2}\right)} \cdot \left(\text{EllipticF}\left(\frac{1}{2} \sqrt{x^2} \cdot \frac{\left(-b + (-4ac+b^2)^{1/2}\right)^{1/2}}{a}, \frac{1}{2} \sqrt{-4 + 2 \sqrt{\frac{b + (-4ac+b^2)^{1/2}}{a/c}}}\right) - \text{EllipticE}\left(\frac{1}{2} \sqrt{x^2} \cdot \frac{\left(-b + (-4ac+b^2)^{1/2}\right)^{1/2}}{a}, \frac{1}{2} \sqrt{-4 + 2 \sqrt{\frac{b + (-4ac+b^2)^{1/2}}{a/c}}}\right)\right) + \frac{(Ae - Bd)}{e} \cdot \frac{-2c \cdot \left(\frac{1}{2} \sqrt{\frac{b}{4ac-b^2}} x^3 + \frac{1}{2} \sqrt{\frac{3abce - 2ac^2d - b^3e + b^2cd}{a(4ac-b^2)}} \frac{1}{(ae^2 - bd + ce + cd^2)} x^3\right)}{\left(x^4 + \frac{1}{c} \sqrt{\frac{b}{4ac-b^2}} x^2 + \frac{a}{c}\right)^{1/2}} - \frac{1}{4} \cdot \frac{x^2}{\left(-b/a + 1/a \sqrt{-4ac+b^2}\right)^{1/2}} \cdot \frac{\left(4 + 2 \sqrt{\frac{bx^2/a - 2x^2/a \sqrt{-4ac+b^2}}{ax^2}}\right)^{1/2}}{\left(4 + 2 \sqrt{\frac{bx^2/a + 2x^2/a \sqrt{-4ac+b^2}}{ax^2}}\right)^{1/2}} \cdot \frac{1}{\left(c x^4 + b x^2 + a\right)^{1/2}} \cdot \text{EllipticF}\left(\frac{1}{2} \sqrt{x^2} \cdot \frac{\left(-b + (-4ac+b^2)^{1/2}\right)^{1/2}}{a}, \frac{1}{2} \sqrt{-4 + 2 \sqrt{\frac{b + (-4ac+b^2)^{1/2}}{a/c}}}\right) / \frac{1}{\left(ae^2 - bd + ce + cd^2\right)} \cdot \frac{b + e + \frac{1}{4} x^2}{\left(-b/a + 1/a \sqrt{-4ac+b^2}\right)^{1/2}} \cdot \frac{\left(4 + 2 \sqrt{\frac{bx^2/a - 2x^2/a \sqrt{-4ac+b^2}}{ax^2}}\right)^{1/2}}{\left(4 + 2 \sqrt{\frac{bx^2/a + 2x^2/a \sqrt{-4ac+b^2}}{ax^2}}\right)^{1/2}} \cdot \frac{1}{\left(c x^4 + b x^2 + a\right)^{1/2}} \cdot \text{EllipticF}\left(\frac{1}{2} \sqrt{x^2} \cdot \frac{\left(-b + (-4ac+b^2)^{1/2}\right)^{1/2}}{a}, \frac{1}{2} \sqrt{-4 + 2 \sqrt{\frac{b + (-4ac+b^2)^{1/2}}{a/c}}}\right)$$

### 3.28.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)(a + bx^2 + cx^4)^{3/2}} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fricas")`

output `Timed out`



**3.28.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)(a + bx^2 + cx^4)^{3/2}} dx = \text{Timed out}$$

input `integrate((B*x**2+A)/(e*x**2+d)/(c*x**4+b*x**2+a)**(3/2),x)`output `Timed out`**3.28.7 Maxima [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + bx^2 + a)^{\frac{3}{2}}(ex^2 + d)} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")`output `integrate((B*x^2 + A)/((c*x^4 + b*x^2 + a)^(3/2)*(e*x^2 + d)), x)`**3.28.8 Giac [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + bx^2 + a)^{\frac{3}{2}}(ex^2 + d)} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")`output `integrate((B*x^2 + A)/((c*x^4 + b*x^2 + a)^(3/2)*(e*x^2 + d)), x)`

**3.28.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)(a + bx^2 + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(ex^2 + d)(cx^4 + bx^2 + a)^{3/2}} dx$$

input `int((A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(3/2)),x)`output `int((A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(3/2)), x)`

$$3.29 \quad \int \frac{A+Bx^2}{(d+ex^2)^2(a+bx^2+cx^4)^{3/2}} dx$$

|        |                                      |     |
|--------|--------------------------------------|-----|
| 3.29.1 | Optimal result . . . . .             | 250 |
| 3.29.2 | Mathematica [C] (verified) . . . . . | 251 |
| 3.29.3 | Rubi [A] (verified) . . . . .        | 252 |
| 3.29.4 | Maple [B] (verified) . . . . .       | 254 |
| 3.29.5 | Fricas [F(-1)] . . . . .             | 255 |
| 3.29.6 | Sympy [F(-1)] . . . . .              | 255 |
| 3.29.7 | Maxima [F] . . . . .                 | 255 |
| 3.29.8 | Giac [F] . . . . .                   | 256 |
| 3.29.9 | Mupad [F(-1)] . . . . .              | 256 |

### 3.29.1 Optimal result

Integrand size = 33, antiderivative size = 1301

$$\int \frac{A+Bx^2}{(d+ex^2)^2(a+bx^2+cx^4)^{3/2}} dx = \frac{x(abc(Ae(2cd-be) - B(cd^2 - ae^2)) + (b^2 - 2ac)(aBe(2cd-be) + Acd^2 - b^2cd)) + \sqrt{c}(aBd(-4c^2d^2 - 3b^2e^2 + 4ce(bd + 2ae)) + A(2b^3de^2 + 2bcd(cd^2 - 3ae^2) - 4ace(-2cd^2 + ae^2) + b^2(-4cd^2 + ae^2)) - 2a(-b^2 + 4ac)d(cd^2 + e(-bd + ae))^2(\sqrt{a} + \sqrt{cx^2}))}{4d^{3/2}(cd^2 - bde + ae^2)^{5/2}} - \frac{e^3(Bd - Ae)x\sqrt{a+bx^2+cx^4}}{2d(cd^2 - bde + ae^2)^2(d+ex^2)} + \frac{e^{3/2}(Ae(7cd^2 - e(4bd - ae)) - Bd(5cd^2 - e(2bd + ae))) \arctan\left(\frac{\sqrt{cd^2 - bde + ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{a+bx^2+cx^4}}\right)}{4d^{3/2}(cd^2 - bde + ae^2)^{5/2}} - \frac{\sqrt[4]{c}(aBd(4c^2d^2 + 3b^2e^2 - 4ce(bd + 2ae)) - A(2b^3de^2 + 2bcd(cd^2 - 3ae^2) - 4ace(-2cd^2 + ae^2) + b^2(-4cd^2 + ae^2)) + 2a^{3/4}(b^2 - 4ac)d(cd^2 + e(-bd + ae))^2\sqrt{a} + \sqrt[4]{c}(a\sqrt{ce}(Bd - 2Ae) + \sqrt{a}(Bd - Ae)(cd - be) + A\sqrt{cd}(-cd + be))(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \text{EllipticF}}{2a^{3/4}(b - 2\sqrt{a}\sqrt{c})d(-\sqrt{cd} + \sqrt{ae})(-cd^2 + e(bd - ae))\sqrt{a+bx^2+cx^4}} + \frac{e(\sqrt{cd} + \sqrt{ae})(Ae(7cd^2 - e(4bd - ae)) - Bd(5cd^2 - e(2bd + ae)))(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} \text{EllipticPi}}{8\sqrt[4]{a}\sqrt[4]{cd^2}(\sqrt{cd} - \sqrt{ae})(cd^2 - bde + ae^2)^2\sqrt{a+bx^2+cx^4}}$$

---

3.29.  $\int \frac{A+Bx^2}{(d+ex^2)^2(a+bx^2+cx^4)^{3/2}} dx$

output

```

1/4*e^(3/2)*(A*e*(7*c*d^2-e*(-a*e+4*b*d))-B*d*(5*c*d^2-e*(a*e+2*b*d)))*arc
tan(x*(a*e^2-b*d*e+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+b*x^2+a)^(1/2))/d^(
3/2)/(a*e^2-b*d*e+c*d^2)^(5/2)+x*(a*b*c*(A*e*(-b*e+2*c*d)-B*(-a*e^2+c*d^2)
)+(-2*a*c+b^2)*(a*B*e*(-b*e+2*c*d)+A*(c^2*d^2+b^2*e^2-c*e*(a*e+2*b*d)))-c*
(a*B*(2*c^2*d^2+b^2*e^2-2*c*e*(a*e+b*d))+A*(2*b^2*c*d*e-4*a*c^2*d*e-b^3*e^
2-b*c*(-3*a*e^2+c*d^2)))*x^2)/a/(-4*a*c+b^2)/(a*e^2-b*d*e+c*d^2)^(c*x^4+
b*x^2+a)^(1/2)-1/2*e^3*(-A*e+B*d)*x*(c*x^4+b*x^2+a)^(1/2)/d/(a*e^2-b*d*e+c
*d^2)^(e*x^2+d)+1/2*(a*B*d*(-4*c^2*d^2-3*b^2*e^2+4*c*e*(2*a*e+b*d))+A*(2
*b^3*d*e^2+2*b*c*d*(-3*a*e^2+c*d^2)-4*a*c*e*(a*e^2-2*c*d^2)+b^2*(a*e^3-4*c
*d^2*e)))*x*c^(1/2)*(c*x^4+b*x^2+a)^(1/2)/a/(4*a*c-b^2)/d/(c*d^2+e*(a*e-b*
d))^2/(a^(1/2)+x^2*c^(1/2))-1/2*c^(1/4)*(a*B*d*(4*c^2*d^2+3*b^2*e^2-4*c*e*
(2*a*e+b*d))-A*(2*b^3*d*e^2+2*b*c*d*(-3*a*e^2+c*d^2)-4*a*c*e*(a*e^2-2*c*d^
2)+b^2*(a*e^3-4*c*d^2*e)))*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(
2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/
2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(
1/2)+x^2*c^(1/2))^2)^(1/2)/a^(3/4)/(-4*a*c+b^2)/d/(c*d^2+e*(a*e-b*d))^2/(c
*x^4+b*x^2+a)^(1/2)-1/8*e*(A*e*(7*c*d^2-e*(-a*e+4*b*d))-B*d*(5*c*d^2-e*(a*
e+2*b*d)))*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)
*x/a^(1/4)))*EllipticPi(sin(2*arctan(c^(1/4)*x/a^(1/4))),-1/4*(-e*a^(1/2)+
d*c^(1/2))^2/d/e/a^(1/2)/c^(1/2),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(e*a^...

```

### 3.29.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 15.22 (sec) , antiderivative size = 1116, normalized size of antiderivative = 0.86

$$\int \frac{A + Bx^2}{(d + ex^2)^2 (a + bx^2 + cx^4)^{3/2}} dx = \frac{4 \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} d (-a(b^2 - 4ac) e^3 (Bd - Ae) x (a + bx^2 + cx^4) + 2dx (d +$$

input `Integrate[(A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(3/2)),x]`

```
output (4*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*(-(a*(b^2 - 4*a*c)*e^3*(B*d - A*e)*x*
(a + b*x^2 + c*x^4)) + 2*d*x*(d + e*x^2)*(a*B*(-(b^3*e^2) + b^2*c*e*(2*d -
e*x^2) + b*c*(3*a*e^2 - c*d*(d - 2*e*x^2)) - 2*c^2*(c*d^2*x^2 + a*e*(2*d
- e*x^2))) + A*(b^4*e^2 + b^3*c*e*(-2*d + e*x^2) + 2*a*c^2*(a*e^2 - c*d*(d
- 2*e*x^2)) + b^2*c*(-4*a*e^2 + c*d*(d - 2*e*x^2)) + b*c^2*(c*d^2*x^2 - 3
*a*e*(-2*d + e*x^2)))) - I*Sqrt[2]*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)
/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*(d +
e*x^2)*((-b + Sqrt[b^2 - 4*a*c])*d*(a*B*d*(-4*c^2*d^2 - 3*b^2*e^2 + 4*c*e
*(b*d + 2*a*e)) + A*(2*b^3*d*e^2 + 2*b*c*d*(c*d^2 - 3*a*e^2) - 4*a*c*e*(-2
*c*d^2 + a*e^2) + b^2*(-4*c*d^2*e + a*e^3)))*EllipticE[I*ArcSinh[Sqrt[2]*S
qrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 -
4*a*c])] - d*(a*B*d*(3*b^2*(b - Sqrt[b^2 - 4*a*c])*e^2 - 4*c^2*d*(Sqrt[b^
2 - 4*a*c]*d - 6*a*e) + 2*c*(-3*b + 2*Sqrt[b^2 - 4*a*c])*e*(b*d + 2*a*e))
+ A*(-2*b^4*d*e^2 + b^2*(-2*c^2*d^3 + a*Sqrt[b^2 - 4*a*c])*e^3 - 4*c*d*e*(S
qrt[b^2 - 4*a*c]*d - 3*a*e)) - 4*a*c*(-2*c^2*d^3 + a*Sqrt[b^2 - 4*a*c])*e^3
- 2*c*d*e*(Sqrt[b^2 - 4*a*c]*d - 2*a*e)) + b^3*e*(4*c*d^2 + e*(2*Sqrt[b^2
- 4*a*c]*d - a*e)) + 2*b*c*(c*d^2*(Sqrt[b^2 - 4*a*c]*d - 8*a*e) + a*e^2*(
-3*Sqrt[b^2 - 4*a*c]*d + 2*a*e)))*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b +
Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])]
- 2*a*(-b^2 + 4*a*c)*e*(A*e*(7*c*d^2 + e*(-4*b*d + a*e)) + B*(-5*c*d^3 ...
```

### 3.29.3 Rubi [A] (verified)

Time = 3.43 (sec) , antiderivative size = 2112, normalized size of antiderivative = 1.62, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.061$ , Rules used = {2258, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx^2}{(d + ex^2)^2 (a + bx^2 + cx^4)^{3/2}} dx$$

↓ 2258

$$\int \left( \frac{A(-ce(ae + 2bd) + b^2e^2 + c^2d^2) - cx^2(Ae(2cd - be) - B(cd^2 - ae^2)) + aBe(2cd - be)}{(a + bx^2 + cx^4)^{3/2} (ae^2 - bde + cd^2)^2} + \frac{e}{(d + ex^2)^2 \sqrt{a + b}}

↓ 2009$$

$$\begin{aligned}
& \frac{(Bd - Ae)x\sqrt{cx^4 + bx^2 + ae^3}}{2d(cd^2 - bed + ae^2)^2(ex^2 + d)} - \\
& \frac{\sqrt[4]{a}\sqrt[4]{c}(Bd - Ae)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + bx^2 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right)\middle|\frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)e^2}{2d(cd^2 - bed + ae^2)^2\sqrt{cx^4 + bx^2 + a}} + \\
& \frac{\sqrt{c}(Bd - Ae)x\sqrt{cx^4 + bx^2 + ae^2}}{2d(cd^2 - bed + ae^2)^2(\sqrt{cx^2 + \sqrt{a}})} - \\
& \frac{(Bd - Ae)(3cd^2 - e(2bd - ae))\arctan\left(\frac{\sqrt{cd^2 - bed + ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{cx^4 + bx^2 + a}}\right)e^{3/2}}{4d^{3/2}(cd^2 - bed + ae^2)^{5/2}} + \\
& \frac{(Ae(2cd - be) - B(cd^2 - ae^2))\arctan\left(\frac{\sqrt{cd^2 - bed + ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{cx^4 + bx^2 + a}}\right)e^{3/2}}{2\sqrt{d}(cd^2 - bed + ae^2)^{5/2}} + \\
& \frac{\sqrt[4]{c}(Ae(2cd - be) - B(cd^2 - ae^2))(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + bx^2 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)e}{2^4\sqrt{a}(\sqrt{cd} - \sqrt{ae})(cd^2 - bed + ae^2)^2\sqrt{cx^4 + bx^2 + a}} - \\
& \frac{\sqrt[4]{c}(Bd - Ae)(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + bx^2 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)e}{2^4\sqrt{ad}(\sqrt{cd} - \sqrt{ae})(cd^2 - bed + ae^2)\sqrt{cx^4 + bx^2 + a}} + \\
& \frac{(\sqrt{cd} + \sqrt{ae})(Bd - Ae)(3cd^2 - e(2bd - ae))(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + bx^2 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}\text{EllipticPi}\left(-\frac{(\sqrt{cd} - \sqrt{ae})^2}{4\sqrt{a}\sqrt{cde}}, 2\arctan\left(\frac{\sqrt[4]{c}}{\sqrt[4]{a}}\right)\right)}{8^4\sqrt{a}\sqrt[4]{cd^2}(\sqrt{cd} - \sqrt{ae})(cd^2 - bed + ae^2)^2\sqrt{cx^4 + bx^2 + a}} \\
& \frac{a^{3/4}\left(\frac{\sqrt{cd}}{\sqrt{a}} + e\right)^2(Ae(2cd - be) - B(cd^2 - ae^2))(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + bx^2 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}\text{EllipticPi}\left(-\frac{(\sqrt{cd} - \sqrt{ae})^2}{4\sqrt{a}\sqrt{cde}}, 2\arctan\left(\frac{\sqrt[4]{c}}{\sqrt[4]{a}}\right)\right)}{4^4\sqrt{cd}(cd^2 - ae^2)(cd^2 - bed + ae^2)^2\sqrt{cx^4 + bx^2 + a}} \\
& \frac{\sqrt[4]{c}(aB(2c^2d^2 + b^2e^2 - 2ce(bd + ae)) + A(-e^2b^3 + 2cdeb^2 - c(cd^2 - 3ae^2)b - 4ac^2de))(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + bx^2 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}}{a^{3/4}(b^2 - 4ac)(cd^2 - bed + ae^2)^2\sqrt{cx^4 + bx^2 + a}} \\
& \frac{\sqrt[4]{c}(a^{3/2}B\sqrt{ce^2} + a(2Bcd - bBe - Ace)e + A(cd - be)^2 - \sqrt{a}\sqrt{c}(Bcd^2 - Ae(2cd - be)))(\sqrt{cx^2 + \sqrt{a}})\sqrt{\frac{cx^4 + bx^2 + a}{(\sqrt{cx^2 + \sqrt{a}})^2}}}{2a^{3/4}(b - 2\sqrt{a}\sqrt{c})(cd^2 - bed + ae^2)^2\sqrt{cx^4 + bx^2 + a}} \\
& \frac{\sqrt{c}(aB(2c^2d^2 + b^2e^2 - 2ce(bd + ae)) + A(-e^2b^3 + 2cdeb^2 - c(cd^2 - 3ae^2)b - 4ac^2de))x\sqrt{cx^4 + bx^2 + a}}{a(b^2 - 4ac)(cd^2 - bed + ae^2)^2(\sqrt{cx^2 + \sqrt{a}})} + \\
& \frac{x(-c(aB(2c^2d^2 + b^2e^2 - 2ce(bd + ae)) + A(-e^2b^3 + 2cdeb^2 - c(cd^2 - 3ae^2)b - 4ac^2de))x^2 + abc(Ae(2cd - be))}{a(b^2 - 4ac)(cd^2 - bed + ae^2)^2\sqrt{cx^4 + a}}
\end{aligned}$$

input `Int[(A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(3/2)),x]`

output

```
(x*(a*b*c*(A*e*(2*c*d - b*e) - B*(c*d^2 - a*e^2)) + (b^2 - 2*a*c)*(a*B*e*(
2*c*d - b*e) + A*(c^2*d^2 + b^2*e^2 - c*e*(2*b*d + a*e))) - c*(a*B*(2*c^2*
d^2 + b^2*e^2 - 2*c*e*(b*d + a*e)) + A*(2*b^2*c*d*e - 4*a*c^2*d*e - b^3*e^
2 - b*c*(c*d^2 - 3*a*e^2)))*x^2)/(a*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)
^2*Sqrt[a + b*x^2 + c*x^4]) + (Sqrt[c]*e^2*(B*d - A*e)*x*Sqrt[a + b*x^2 +
c*x^4])/(2*d*(c*d^2 - b*d*e + a*e^2)^2*(Sqrt[a] + Sqrt[c]*x^2)) + (Sqrt[c]
*(a*B*(2*c^2*d^2 + b^2*e^2 - 2*c*e*(b*d + a*e)) + A*(2*b^2*c*d*e - 4*a*c^2
*d*e - b^3*e^2 - b*c*(c*d^2 - 3*a*e^2)))*x*Sqrt[a + b*x^2 + c*x^4])/(a*(b^
2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)^2*(Sqrt[a] + Sqrt[c]*x^2)) - (e^3*(B*d
- A*e)*x*Sqrt[a + b*x^2 + c*x^4])/(2*d*(c*d^2 - b*d*e + a*e^2)^2*(d + e*x^
2)) - (e^(3/2)*(B*d - A*e)*(3*c*d^2 - e*(2*b*d - a*e))*ArcTan[(Sqrt[c*d^2
- b*d*e + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + b*x^2 + c*x^4])])/(4*d^(3/2)
*(c*d^2 - b*d*e + a*e^2)^(5/2)) + (e^(3/2)*(A*e*(2*c*d - b*e) - B*(c*d^2 -
a*e^2))*ArcTan[(Sqrt[c*d^2 - b*d*e + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a +
b*x^2 + c*x^4])])/(2*Sqrt[d]*(c*d^2 - b*d*e + a*e^2)^(5/2)) - (a^(1/4)*c^(
1/4)*e^2*(B*d - A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqr
t[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqr
t[a]*Sqrt[c]))/4])/(2*d*(c*d^2 - b*d*e + a*e^2)^2*Sqrt[a + b*x^2 + c*x^4]
) - (c^(1/4)*(a*B*(2*c^2*d^2 + b^2*e^2 - 2*c*e*(b*d + a*e)) + A*(2*b^2*c*d
*e - 4*a*c^2*d*e - b^3*e^2 - b*c*(c*d^2 - 3*a*e^2)))*(Sqrt[a] + Sqrt[c]...
```

### 3.29.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2258 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[1/Sqrt[a + b*x^2 + c*x^4], Px*(d + e*x^2)^q*(a + b*x^2 + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e}, x] && PolyQ[Px, x] && IntegerQ[p + 1/2] && IntegerQ[q]`

### 3.29.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 8275 vs.  $2(1252) = 2504$ .

Time = 3.61 (sec) , antiderivative size = 8276, normalized size of antiderivative = 6.36

| method   | result                          | size |
|----------|---------------------------------|------|
| default  | Expression too large to display | 8276 |
| elliptic | Expression too large to display | 9725 |

$$3.29. \int \frac{A+Bx^2}{(d+ex^2)^2(a+bx^2+cx^4)^{3/2}} dx$$

input `int((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x,method=_RETURNVERBOSE)`

output `result too large to display`

### 3.29.5 Fricas [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)^2 (a + bx^2 + cx^4)^{3/2}} dx = \text{Timed out}$$

input `integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fricas")`

output `Timed out`

### 3.29.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)^2 (a + bx^2 + cx^4)^{3/2}} dx = \text{Timed out}$$

input `integrate((B*x**2+A)/(e*x**2+d)**2/(c*x**4+b*x**2+a)**(3/2),x)`

output `Timed out`

### 3.29.7 Maxima [F]

$$\int \frac{A + Bx^2}{(d + ex^2)^2 (a + bx^2 + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + bx^2 + a)^{\frac{3}{2}} (ex^2 + d)^2} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")`

output `integrate((B*x^2 + A)/((c*x^4 + b*x^2 + a)^(3/2)*(e*x^2 + d)^2), x)`

---

3.29.  $\int \frac{A+Bx^2}{(d+ex^2)^2(a+bx^2+cx^4)^{3/2}} dx$



**3.29.8 Giac [F]**

$$\int \frac{A + Bx^2}{(d + ex^2)^2 (a + bx^2 + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(cx^4 + bx^2 + a)^{\frac{3}{2}} (ex^2 + d)^2} dx$$

input `integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")`

output `integrate((B*x^2 + A)/((c*x^4 + b*x^2 + a)^(3/2)*(e*x^2 + d)^2), x)`

**3.29.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{A + Bx^2}{(d + ex^2)^2 (a + bx^2 + cx^4)^{3/2}} dx = \int \frac{Bx^2 + A}{(ex^2 + d)^2 (cx^4 + bx^2 + a)^{3/2}} dx$$

input `int((A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(3/2)),x)`

output `int((A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(3/2)), x)`

### 3.30 $\int \frac{\sqrt{a} + \sqrt{cx^2}}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$

|        |                            |     |
|--------|----------------------------|-----|
| 3.30.1 | Optimal result             | 257 |
| 3.30.2 | Mathematica [C] (verified) | 258 |
| 3.30.3 | Rubi [A] (verified)        | 258 |
| 3.30.4 | Maple [A] (verified)       | 260 |
| 3.30.5 | Fricas [F(-1)]             | 260 |
| 3.30.6 | Sympy [F]                  | 261 |
| 3.30.7 | Maxima [F(-2)]             | 261 |
| 3.30.8 | Giac [F]                   | 261 |
| 3.30.9 | Mupad [F(-1)]              | 262 |

#### 3.30.1 Optimal result

Integrand size = 41, antiderivative size = 273

$$\int \frac{\sqrt{a} + \sqrt{cx^2}}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx = -\frac{(\sqrt{cd} - \sqrt{ae}) \arctan\left(\frac{\sqrt{cd^2 - bde + ae^2x}}{\sqrt{d}\sqrt{e}\sqrt{a+bx^2+cx^4}}\right)}{2\sqrt{d}\sqrt{e}\sqrt{cd^2 - bde + ae^2}} + \frac{(\sqrt{cd} + \sqrt{ae})(\sqrt{a} + \sqrt{cx^2})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a} + \sqrt{cx^2})^2}} \operatorname{EllipticPi}\left(-\frac{(\sqrt{cd} - \sqrt{ae})^2}{4\sqrt{a}\sqrt{cde}}, 2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{4\sqrt[4]{a}\sqrt[4]{cde}\sqrt{a+bx^2+cx^4}}$$

```
output -1/2*arctan(x*(a*e^2-b*d*e+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+b*x^2+a)^(1/2))*(-e*a^(1/2)+d*c^(1/2))/d^(1/2)/e^(1/2)/(a*e^2-b*d*e+c*d^2)^(1/2)+1/4*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticPi(sin(2*arctan(c^(1/4)*x/a^(1/4))),-1/4*(-e*a^(1/2)+d*c^(1/2))^2/d/e/a^(1/2)/c^(1/2),1/2*(2-b/a^(1/2)/c^(1/2))^2)^(1/2))*(e*a^(1/2)+d*c^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2))^2)^(1/2)/a^(1/4)/c^(1/4)/d/e/(c*x^4+b*x^2+a)^(1/2)
```

### 3.30.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.42 (sec) , antiderivative size = 310, normalized size of antiderivative = 1.14

$$\int \frac{\sqrt{a} + \sqrt{cx^2}}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx =$$

$$\frac{i\sqrt{\frac{b+\sqrt{b^2-4ac}+2cx^2}{b+\sqrt{b^2-4ac}}}\sqrt{1+\frac{2cx^2}{b-\sqrt{b^2-4ac}}}\left(\sqrt{cd}\operatorname{EllipticF}\left(\operatorname{iarcsinh}\left(\sqrt{2}\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}x\right),\frac{b+\sqrt{b^2-4ac}}{b-\sqrt{b^2-4ac}}\right)+(-\sqrt{cd}\right)}{\sqrt{2}\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}de\sqrt{a+bx^2+cx^4}}$$

input `Integrate[(Sqrt[a] + Sqrt[c]*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]),x]`

output `((-I)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c]])*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*(Sqrt[c]*d*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x, (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])]) + (-Sqrt[c]*d) + Sqrt[a]*e)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x, (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])])]/(Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])])*d*e*Sqrt[a + b*x^2 + c*x^4])`

### 3.30.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.024$ , Rules used = {2220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a} + \sqrt{cx^2}}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx$$

↓ 2220

$$\frac{(\sqrt{a} + \sqrt{cx^2}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{cx^2})^2}} (\sqrt{ae} + \sqrt{cd}) \operatorname{EllipticPi}\left(-\frac{\sqrt{a}\left(\frac{\sqrt{cd}}{\sqrt{a}}-e\right)^2}{4\sqrt{cde}}, 2 \arctan\left(\frac{\sqrt[4]{Cx}}{\sqrt[4]{a}}\right), \frac{1}{4}\left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{4\sqrt[4]{a}\sqrt[4]{cde}\sqrt{a+bx^2+cx^4} (\sqrt{cd} - \sqrt{ae}) \arctan\left(\frac{x\sqrt{ae^2-bde+cd^2}}{\sqrt{d}\sqrt{e}\sqrt{a+bx^2+cx^4}}\right)} \\ \frac{2\sqrt{d}\sqrt{e}\sqrt{ae^2-bde+cd^2}}$$

input `Int[(Sqrt[a] + Sqrt[c]*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]),x]`

output `-1/2*((Sqrt[c]*d - Sqrt[a]*e)*ArcTan[(Sqrt[c*d^2 - b*d*e + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + b*x^2 + c*x^4])]/(Sqrt[d]*Sqrt[e]*Sqrt[c*d^2 - b*d*e + a*e^2]) + ((Sqrt[c]*d + Sqrt[a]*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-1/4*(Sqrt[a]*((Sqrt[c]*d)/Sqrt[a] - e)^2)/(Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4)]/(4*a^(1/4)*c^(1/4)*d*e*Sqrt[a + b*x^2 + c*x^4])`

### 3.30.3.1 Defintions of rubi rules used

rule 2220 `Int[((A_) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[-b + c*(d/e) + a*(e/d), 2]*(x/Sqrt[a + b*x^2 + c*x^4])]/(2*d*e*Rt[-b + c*(d/e) + a*(e/d), 2])), x] + Simp[(B*d + A*e)*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(4*d*e*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticPi[-(e - d*q^2)^2/(4*d*e*q^2), 2*ArcTan[q*x], 1/2 - b/(4*a*q^2)], x] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0] && PosQ[B/A] && PosQ[-b + c*(d/e) + a*(e/d)]`

### 3.30.4 Maple [A] (verified)

Time = 1.80 (sec) , antiderivative size = 365, normalized size of antiderivative = 1.34

| method   | result   |
|----------|--|
| default  | $\frac{\sqrt{c}\sqrt{2}\sqrt{4-\frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}}\sqrt{4+\frac{2(b+\sqrt{-4ac+b^2})x^2}{a}}F\left(\frac{x\sqrt{2}\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}},\sqrt{-4+\frac{2b(b+\sqrt{-4ac+b^2})}{ac}}\right)}{4e\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}\sqrt{cx^4+bx^2+a}} + \frac{(e\sqrt{a}-d\sqrt{c})\sqrt{2}\sqrt{c}}{4e\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}\sqrt{cx^4+bx^2+a}}$   |
| elliptic | $\frac{\sqrt{(cx^4+bx^2+a)c}\sqrt{(cx^4+bx^2+a)a}(\sqrt{a+x^2}\sqrt{c})}{d\sqrt{-\frac{b}{a}+\frac{\sqrt{-4ac+b^2}}{a}}\sqrt{acx^4+abx^2+a^2}} \left( \frac{a\sqrt{2}\sqrt{1+\frac{bx^2}{2a}-\frac{x^2\sqrt{-4ac+b^2}}{2a}}\sqrt{1+\frac{bx^2}{2a}+\frac{x^2\sqrt{-4ac+b^2}}{2a}}\Pi\left(\frac{x\sqrt{2}\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}}{2},-\frac{1}{2}\right)}{d\sqrt{-\frac{b}{a}+\frac{\sqrt{-4ac+b^2}}{a}}\sqrt{acx^4+abx^2+a^2}} \right)$ |

```
input int((a^(1/2)+x^2*c^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURN
VERBOSE)
```

```
output 1/4*c^(1/2)/e*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b
^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*
x^2+a)^(1/2)*EllipticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2
*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))+1/e*(e*a^(1/2)-d*c^(1/2))/d*2^
(1/2)/(-b/a+1/a*(-4*a*c+b^2)^(1/2))^(1/2)*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c
+b^2)^(1/2))^(1/2)*(1+1/2*b*x^2/a+1/2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2)/(c*x
^4+b*x^2+a)^(1/2)*EllipticPi(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/
2),-2/(-b+(-4*a*c+b^2)^(1/2))*a*e/d,(-1/2*(b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*
2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2))
```

### 3.30.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{a} + \sqrt{cx^2}}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = \text{Timed out}$$

```
input integrate((a^(1/2)+x^2*c^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorith
m="fricas")
```

```
output Timed out
```

3.30.  $\int \frac{\sqrt{a} + \sqrt{cx^2}}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx$

### 3.30.6 Sympy [F]

$$\int \frac{\sqrt{a} + \sqrt{cx^2}}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = \int \frac{\sqrt{a} + \sqrt{cx^2}}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx$$

input `integrate((a**(1/2)+x**2*c**(1/2))/(e*x**2+d)/(c*x**4+b*x**2+a)**(1/2),x)`

output `Integral((sqrt(a) + sqrt(c)*x**2)/((d + e*x**2)*sqrt(a + b*x**2 + c*x**4)), x)`

### 3.30.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a} + \sqrt{cx^2}}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a^(1/2)+x^2*c^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

### 3.30.8 Giac [F]

$$\int \frac{\sqrt{a} + \sqrt{cx^2}}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = \int \frac{\sqrt{cx^2} + \sqrt{a}}{\sqrt{cx^4 + bx^2 + a}(ex^2 + d)} dx$$

input `integrate((a^(1/2)+x^2*c^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")`

output `integrate((sqrt(c)*x^2 + sqrt(a))/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)), x)`

**3.30.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{a} + \sqrt{cx^2}}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = \int \frac{\sqrt{a} + \sqrt{cx^2}}{(ex^2 + d)\sqrt{cx^4 + bx^2 + a}} dx$$

input `int((a^(1/2) + c^(1/2)*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)),x)`output `int((a^(1/2) + c^(1/2)*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)), x)`

$$3.31 \quad \int \frac{1 + \sqrt{\frac{c}{a}}x^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$$

|        |                            |     |
|--------|----------------------------|-----|
| 3.31.1 | Optimal result             | 263 |
| 3.31.2 | Mathematica [C] (verified) | 264 |
| 3.31.3 | Rubi [A] (verified)        | 264 |
| 3.31.4 | Maple [A] (verified)       | 266 |
| 3.31.5 | Fricas [F(-1)]             | 266 |
| 3.31.6 | Sympy [F]                  | 267 |
| 3.31.7 | Maxima [F]                 | 267 |
| 3.31.8 | Giac [F]                   | 267 |
| 3.31.9 | Mupad [F(-1)]              | 268 |

### 3.31.1 Optimal result

Integrand size = 41, antiderivative size = 271

$$\int \frac{1 + \sqrt{\frac{c}{a}}x^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx = -\frac{(\sqrt{\frac{c}{a}}d - e) \arctan\left(\frac{\sqrt{cd^2 - bde + ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{a+bx^2+cx^4}}\right)}{2\sqrt{d}\sqrt{e}\sqrt{cd^2 - bde + ae^2}} + \frac{(\sqrt{\frac{c}{a}}d + e)(1 + \sqrt{\frac{c}{a}}x^2) \sqrt{\frac{a+bx^2+cx^4}{a(1+\sqrt{\frac{c}{a}}x^2)^2}} \text{EllipticPi}\left(-\frac{(\sqrt{\frac{c}{a}}d - e)^2}{4\sqrt{\frac{c}{a}}de}, 2 \arctan\left(\sqrt[4]{\frac{c}{a}}x\right), \frac{1}{4}\left(2 - \frac{b\sqrt{\frac{c}{a}}}{c}\right)\right)}{4\sqrt[4]{\frac{c}{a}}de\sqrt{a+bx^2+cx^4}}$$

output

```
-1/2*arctan(x*(a*e^2-b*d*e+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+b*x^2+a)^(1/2))*(-e+d*(c/a)^(1/2))/d^(1/2)/e^(1/2)/(a*e^2-b*d*e+c*d^2)^(1/2)+1/4*(cos(2*arctan((c/a)^(1/4)*x))^2)^(1/2)/cos(2*arctan((c/a)^(1/4)*x))*EllipticPi(sin(2*arctan((c/a)^(1/4)*x)), -1/4*(-e+d*(c/a)^(1/2))^2/d/e/(c/a)^(1/2), 1/2*(2-b*(c/a)^(1/2)/c)^(1/2))*(e+d*(c/a)^(1/2))*(1+x^2*(c/a)^(1/2))*((c*x^4+b*x^2+a)/a/(1+x^2*(c/a)^(1/2))^2)^(1/2)/(c/a)^(1/4)/d/e/(c*x^4+b*x^2+a)^(1/2)
```

---

3.31.  $\int \frac{1 + \sqrt{\frac{c}{a}}x^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$



### 3.31.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.41 (sec) , antiderivative size = 312, normalized size of antiderivative = 1.15

$$\int \frac{1 + \sqrt{\frac{c}{a}}x^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx =$$

$$\frac{i\sqrt{\frac{b+\sqrt{b^2-4ac}+2cx^2}{b+\sqrt{b^2-4ac}}}\sqrt{1+\frac{2cx^2}{b-\sqrt{b^2-4ac}}}\left(\sqrt{\frac{c}{a}}d\operatorname{EllipticF}\left(i\operatorname{arcsinh}\left(\sqrt{2}\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}x\right),\frac{b+\sqrt{b^2-4ac}}{b-\sqrt{b^2-4ac}}\right)+\left(-\sqrt{\frac{c}{a}}\right)\right)}{\sqrt{2}\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}de\sqrt{a+bx^2+cx^4}}$$

input `Integrate[(1 + Sqrt[c/a]*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]),x]`

output `((-I)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c]])*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*(Sqrt[c/a]*d*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x, (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] + (-Sqrt[c/a]*d) + e)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x, (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])])]/(Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*Sqrt[a + b*x^2 + c*x^4])`

### 3.31.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.024$ , Rules used = {2220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2 \sqrt{\frac{c}{a}} + 1}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx$$

↓ 2220

---

3.31.  $\int \frac{1 + \sqrt{\frac{c}{a}}x^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$

$$\frac{(x^2 \sqrt{\frac{c}{a}} + 1) \sqrt{\frac{a+bx^2+cx^4}{a(x^2 \sqrt{\frac{c}{a}} + 1)^2}} (d\sqrt{\frac{c}{a}} + e) \operatorname{EllipticPi} \left( -\frac{(\sqrt{\frac{c}{a}}d - e)^2}{4\sqrt{\frac{c}{a}}de}, 2 \arctan \left( \sqrt[4]{\frac{c}{a}}x \right), \frac{1}{4} \left( 2 - \frac{b\sqrt{\frac{c}{a}}}{c} \right) \right)}{4de \sqrt[4]{\frac{c}{a}} \sqrt{a+bx^2+cx^4} \frac{(d\sqrt{\frac{c}{a}} - e) \arctan \left( \frac{x\sqrt{ae^2 - bde + cd^2}}{\sqrt{d}\sqrt{e}\sqrt{a+bx^2+cx^4}} \right)}{2\sqrt{d}\sqrt{e}\sqrt{ae^2 - bde + cd^2}}}$$

input `Int[(1 + Sqrt[c/a]*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]),x]`

output `-1/2*((Sqrt[c/a]*d - e)*ArcTan[(Sqrt[c*d^2 - b*d*e + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + b*x^2 + c*x^4])])/(Sqrt[d]*Sqrt[e]*Sqrt[c*d^2 - b*d*e + a*e^2]) + ((Sqrt[c/a]*d + e)*(1 + Sqrt[c/a]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + Sqrt[c/a]*x^2)^2)]*EllipticPi[-1/4*(Sqrt[c/a]*d - e)^2/(Sqrt[c/a]*d*e), 2*ArcTan[(c/a)^(1/4)*x], (2 - (b*Sqrt[c/a])/c)/4])/(4*(c/a)^(1/4)*d*e*Sqrt[a + b*x^2 + c*x^4])`

### 3.31.3.1 Defintions of rubi rules used

rule 2220 `Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[-b + c*(d/e) + a*(e/d), 2]*(x/Sqrt[a + b*x^2 + c*x^4])]/(2*d*e*Rt[-b + c*(d/e) + a*(e/d), 2]))], x] + Simp[(B*d + A*e)*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2])]/(4*d*e*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticPi[-(e - d*q^2)^2/(4*d*e*q^2), 2*ArcTan[q*x], 1/2 - b/(4*a*q^2)], x] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0] && PosQ[B/A] && PosQ[-b + c*(d/e) + a*(e/d)]`

$$3.31. \int \frac{1 + \sqrt{\frac{c}{a}}x^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$$

### 3.31.4 Maple [A] (verified)

Time = 4.43 (sec) , antiderivative size = 369, normalized size of antiderivative = 1.36

| method   | result   |
|----------|--|
| default  | $\frac{\sqrt{\frac{c}{a}} \sqrt{2} \sqrt{4 - \frac{2(-b + \sqrt{-4ac + b^2})x^2}{a}} \sqrt{4 + \frac{2(b + \sqrt{-4ac + b^2})x^2}{a}} F\left(\frac{x\sqrt{2} \sqrt{\frac{-b + \sqrt{-4ac + b^2}}{a}}, \sqrt{-4 + \frac{2b(b + \sqrt{-4ac + b^2})}{ac}}\right)}{4e\sqrt{\frac{-b + \sqrt{-4ac + b^2}}{a}} \sqrt{cx^4 + bx^2 + a}} + \frac{(e - d\sqrt{\frac{c}{a}})\sqrt{2} \sqrt{4 - \frac{2(-b + \sqrt{-4ac + b^2})x^2}{a}}}{4e\sqrt{\frac{-b + \sqrt{-4ac + b^2}}{a}} \sqrt{cx^4 + bx^2 + a}}$ |
| elliptic | $\sqrt{\frac{(cx^4 + bx^2 + a)c}{a}} a \left(1 + x^2 \sqrt{\frac{c}{a}}\right) \left( \frac{c\sqrt{2} \sqrt{4 + \frac{2bx^2}{a} - \frac{2x^2\sqrt{-4ac + b^2}}{a}} \sqrt{4 + \frac{2bx^2}{a} + \frac{2x^2\sqrt{-4ac + b^2}}{a}} F\left(\frac{x\sqrt{2} \sqrt{\frac{-b + \sqrt{-4ac + b^2}}{a}}, \sqrt{-4 + \frac{2b(b + \sqrt{-4ac + b^2})}{ac}}\right)}{4ae\sqrt{-\frac{b}{a} + \frac{\sqrt{-4ac + b^2}}{a}} \sqrt{\frac{c^2x^4}{a} + \frac{bcx^2}{a} + c}} \right)$                            |

input `int((1+x^2*(c/a)^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2), x, method=_RETURNVE  
RBOSE)`

output `1/4*(c/a)^(1/2)/e*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a  
*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*x^2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)  
, 1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))+1/e*(e-d*(c/a)^(1/2))/d*2^(1/2)/(-b/a+1/a*(-4*a*c+b^2)^(1/2))^(1/2)*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2)*(1+1/2*b*x^2/a+1/2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticPi(1/2*x^2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2), -2/(-b+(-4*a*c+b^2)^(1/2))*a*e/d, (-1/2*(b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)`

### 3.31.5 Fricas [F(-1)]

Timed out.

$$\int \frac{1 + \sqrt{\frac{c}{a}}x^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = \text{Timed out}$$

input `integrate((1+x^2*(c/a)^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2), x, algorithm  
="fricas")`

output Timed out

3.31.  $\int \frac{1 + \sqrt{\frac{c}{a}}x^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx$

**3.31.6 Sympy [F]**

$$\int \frac{1 + \sqrt{\frac{c}{a}}x^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = \int \frac{x^2\sqrt{\frac{c}{a}} + 1}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx$$

input `integrate((1+x**2*(c/a)**(1/2))/(e*x**2+d)/(c*x**4+b*x**2+a)**(1/2),x)`

output `Integral((x**2*sqrt(c/a) + 1)/((d + e*x**2)*sqrt(a + b*x**2 + c*x**4)), x)`

**3.31.7 Maxima [F]**

$$\int \frac{1 + \sqrt{\frac{c}{a}}x^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = \int \frac{x^2\sqrt{\frac{c}{a}} + 1}{\sqrt{cx^4 + bx^2 + a}(ex^2 + d)} dx$$

input `integrate((1+x^2*(c/a)^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")`

output `integrate((x^2*sqrt(c/a) + 1)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)), x)`

**3.31.8 Giac [F]**

$$\int \frac{1 + \sqrt{\frac{c}{a}}x^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = \int \frac{x^2\sqrt{\frac{c}{a}} + 1}{\sqrt{cx^4 + bx^2 + a}(ex^2 + d)} dx$$

input `integrate((1+x^2*(c/a)^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")`

output `integrate((x^2*sqrt(c/a) + 1)/(sqrt(c*x^4 + b*x^2 + a)*(e*x^2 + d)), x)`

**3.31.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1 + \sqrt{\frac{c}{a}}x^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = \int \frac{x^2 \sqrt{\frac{c}{a}} + 1}{(ex^2 + d)\sqrt{cx^4 + bx^2 + a}} dx$$

input `int((x^2*(c/a)^(1/2) + 1)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)),x)`

output `int((x^2*(c/a)^(1/2) + 1)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)), x)`

**3.32**  $\int \frac{946+315x^2}{(7+5x^2)\sqrt{2+3x^2+x^4}} dx$

|        |                            |     |
|--------|----------------------------|-----|
| 3.32.1 | Optimal result             | 269 |
| 3.32.2 | Mathematica [C] (verified) | 269 |
| 3.32.3 | Rubi [A] (verified)        | 270 |
| 3.32.4 | Maple [C] (verified)       | 272 |
| 3.32.5 | Fricas [F]                 | 272 |
| 3.32.6 | Sympy [F]                  | 273 |
| 3.32.7 | Maxima [F]                 | 273 |
| 3.32.8 | Giac [F]                   | 273 |
| 3.32.9 | Mupad [F(-1)]              | 274 |

**3.32.1 Optimal result**

Integrand size = 31, antiderivative size = 106

$$\int \frac{946 + 315x^2}{(7 + 5x^2)\sqrt{2 + 3x^2 + x^4}} dx = \frac{631(1 + x^2)\sqrt{\frac{2+x^2}{1+x^2}} \text{EllipticF}(\arctan(x), \frac{1}{2})}{2\sqrt{2}\sqrt{2 + 3x^2 + x^4}} - \frac{2525(2 + x^2) \text{EllipticPi}(\frac{2}{7}, \arctan(x), \frac{1}{2})}{14\sqrt{2}\sqrt{\frac{2+x^2}{1+x^2}}\sqrt{2 + 3x^2 + x^4}}$$

```
output -2525/28*(x^2+2)*(1/(x^2+1))^(1/2)*(x^2+1)^(1/2)*EllipticPi(x/(x^2+1)^(1/2), 2/7, 1/2*2^(1/2))*2^(1/2)/((x^2+2)/(x^2+1))^(1/2)/(x^4+3*x^2+2)^(1/2)+631/4*(x^2+1)^(3/2)*(1/(x^2+1))^(1/2)*EllipticF(x/(x^2+1)^(1/2), 1/2*2^(1/2))*((x^2+2)/(x^2+1))^(1/2)*2^(1/2)/(x^4+3*x^2+2)^(1/2)
```

**3.32.2 Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 10.25 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.70

$$\int \frac{946 + 315x^2}{(7 + 5x^2)\sqrt{2 + 3x^2 + x^4}} dx = \frac{i\sqrt{1 + x^2}\sqrt{2 + x^2}\left(441 \text{EllipticF}\left(i\text{arcsinh}\left(\frac{x}{\sqrt{2}}\right), 2\right) + 505 \text{EllipticPi}\left(\frac{10}{7}, i\text{arcsinh}\left(\frac{x}{\sqrt{2}}\right), 2\right)\right)}{7\sqrt{2 + 3x^2 + x^4}}$$

input `Integrate[(946 + 315*x^2)/((7 + 5*x^2)*Sqrt[2 + 3*x^2 + x^4]),x]`

output `((-1/7*I)*Sqrt[1 + x^2]*Sqrt[2 + x^2]*(441*EllipticF[I*ArcSinh[x/Sqrt[2]], 2] + 505*EllipticPi[10/7, I*ArcSinh[x/Sqrt[2]], 2]))/Sqrt[2 + 3*x^2 + x^4]`

### 3.32.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.161$ , Rules used = {2218, 27, 1412, 1786, 414}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{315x^2 + 946}{(5x^2 + 7)\sqrt{x^4 + 3x^2 + 2}} dx \\
 & \quad \downarrow \text{2218} \\
 & \frac{631}{2} \int \frac{1}{\sqrt{x^4 + 3x^2 + 2}} dx - \frac{2525}{8} \int \frac{4(x^2 + 1)}{(5x^2 + 7)\sqrt{x^4 + 3x^2 + 2}} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{631}{2} \int \frac{1}{\sqrt{x^4 + 3x^2 + 2}} dx - \frac{2525}{2} \int \frac{x^2 + 1}{(5x^2 + 7)\sqrt{x^4 + 3x^2 + 2}} dx \\
 & \quad \downarrow \text{1412} \\
 & \frac{631(x^2 + 1)\sqrt{\frac{x^2+2}{x^2+1}} \text{EllipticF}\left(\arctan(x), \frac{1}{2}\right)}{2\sqrt{2}\sqrt{x^4 + 3x^2 + 2}} - \frac{2525}{2} \int \frac{x^2 + 1}{(5x^2 + 7)\sqrt{x^4 + 3x^2 + 2}} dx \\
 & \quad \downarrow \text{1786} \\
 & \frac{631(x^2 + 1)\sqrt{\frac{x^2+2}{x^2+1}} \text{EllipticF}\left(\arctan(x), \frac{1}{2}\right)}{2\sqrt{2}\sqrt{x^4 + 3x^2 + 2}} - \frac{2525\sqrt{x^2 + 1}\sqrt{x^2 + 2} \int \frac{\sqrt{x^2+1}}{\sqrt{x^2+2}(5x^2+7)} dx}{2\sqrt{x^4 + 3x^2 + 2}} \\
 & \quad \downarrow \text{414} \\
 & \frac{631(x^2 + 1)\sqrt{\frac{x^2+2}{x^2+1}} \text{EllipticF}\left(\arctan(x), \frac{1}{2}\right)}{2\sqrt{2}\sqrt{x^4 + 3x^2 + 2}} - \frac{2525(x^2 + 2) \text{EllipticPi}\left(\frac{2}{7}, \arctan(x), \frac{1}{2}\right)}{14\sqrt{2}\sqrt{\frac{x^2+2}{x^2+1}}\sqrt{x^4 + 3x^2 + 2}}
 \end{aligned}$$

---

3.32.  $\int \frac{946+315x^2}{(7+5x^2)\sqrt{2+3x^2+x^4}} dx$

input `Int[(946 + 315*x^2)/((7 + 5*x^2)*Sqrt[2 + 3*x^2 + x^4]),x]`

output `(631*(1 + x^2)*Sqrt[(2 + x^2)/(1 + x^2)]*EllipticF[ArcTan[x], 1/2])/(2*Sqrt[2]*Sqrt[2 + 3*x^2 + x^4]) - (2525*(2 + x^2)*EllipticPi[2/7, ArcTan[x], 1/2])/(14*Sqrt[2]*Sqrt[(2 + x^2)/(1 + x^2)]*Sqrt[2 + 3*x^2 + x^4])`

### 3.32.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 414 `Int[Sqrt[(c_) + (d_)*(x_)^2]/(((a_) + (b_)*(x_)^2)*Sqrt[(e_) + (f_)*(x_)^2]), x_Symbol] := Simp[c*(Sqrt[e + f*x^2]/(a*e*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[c*(e + f*x^2)/(e*(c + d*x^2))]))*EllipticPi[1 - b*(c/(a*d)), ArcTan[Rt[d/c, 2]*x], 1 - c*(f/(d*e))], x] /; FreeQ[{a, b, c, d, e, f}, x] && PosQ[d/c]`

rule 1412 `Int[1/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[(2*a + (b + q)*x^2)*(Sqrt[(2*a + (b - q)*x^2)/(2*a + (b + q)*x^2)])/(2*a*Rt[(b + q)/(2*a), 2]*Sqrt[a + b*x^2 + c*x^4])*EllipticF[ArcTan[Rt[(b + q)/(2*a), 2]*x], 2*(q/(b + q))], x] /; PosQ[(b + q)/a] && !(PosQ[(b - q)/a] && SimplifierSqrtQ[(b - q)/(2*a), (b + q)/(2*a)])] /; FreeQ[{a, b, c}, x] && GtQ[b^2 - 4*a*c, 0]`

rule 1786 `Int[((d_) + (e_)*(x_)^(n_))^(q_)*((f_) + (g_)*(x_)^(n_))^(r_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Simp[(a + b*x^n + c*x^(2*n))^FracPart[p]/((d + e*x^n)^FracPart[p]*(a/d + (c*x^n)/e)^FracPart[p]) Int[(d + e*x^n)^(p + q)*(f + g*x^n)^r*(a/d + (c/e)*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p, q, r}, x] && EqQ[n2, 2*n] && NeQ[b^2 - 4*a*c, 0] && EqQ[c*d^2 - b*d*e + a*e^2, 0] && !IntegerQ[p]`



```
rule 2218 Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 +
(c_)*(x_)^4]), x_Symbol] :> With[{q = Sqrt[b^2 - 4*a*c]}, Simp[(2*a*B - A
*(b + q))/(2*a*e - d*(b + q)) Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Sim
p[(B*d - A*e)/(2*a*e - d*(b + q)) Int[(2*a + (b + q)*x^2)/((d + e*x^2)*Sq
rt[a + b*x^2 + c*x^4]), x], x] /; RationalQ[q]] /; FreeQ[{a, b, c, d, e, A,
B}, x] && GtQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*A^
2 - b*A*B + a*B^2, 0]
```

### 3.32.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.54 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.88

| method   | result  | size |
|----------|---|------|
| default  | $-\frac{63i\sqrt{2}\sqrt{2x^2+4}\sqrt{x^2+1}F\left(\frac{i\sqrt{2}x}{2},\sqrt{2}\right)}{2\sqrt{x^4+3x^2+2}} - \frac{505i\sqrt{2}\sqrt{1+\frac{x^2}{2}}\sqrt{x^2+1}\Pi\left(\frac{i\sqrt{2}x}{2},\frac{10}{7},\sqrt{2}\right)}{7\sqrt{x^4+3x^2+2}}$ | 93   |
| elliptic | $-\frac{63i\sqrt{2}\sqrt{2x^2+4}\sqrt{x^2+1}F\left(\frac{i\sqrt{2}x}{2},\sqrt{2}\right)}{2\sqrt{x^4+3x^2+2}} - \frac{505i\sqrt{2}\sqrt{1+\frac{x^2}{2}}\sqrt{x^2+1}\Pi\left(\frac{i\sqrt{2}x}{2},\frac{10}{7},\sqrt{2}\right)}{7\sqrt{x^4+3x^2+2}}$ | 93   |

```
input int((315*x^2+946)/(5*x^2+7)/(x^4+3*x^2+2)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -63/2*I*2^(1/2)*(2*x^2+4)^(1/2)*(x^2+1)^(1/2)/(x^4+3*x^2+2)^(1/2)*Elliptic
F(1/2*I*2^(1/2)*x,2^(1/2))-505/7*I*2^(1/2)*(1+1/2*x^2)^(1/2)*(x^2+1)^(1/2)
/(x^4+3*x^2+2)^(1/2)*EllipticPi(1/2*I*2^(1/2)*x,10/7,2^(1/2))
```

### 3.32.5 Fricas [F]

$$\int \frac{946 + 315x^2}{(7 + 5x^2)\sqrt{2 + 3x^2 + x^4}} dx = \int \frac{315x^2 + 946}{\sqrt{x^4 + 3x^2 + 2}(5x^2 + 7)} dx$$

```
input integrate((315*x^2+946)/(5*x^2+7)/(x^4+3*x^2+2)^(1/2),x, algorithm="fricas
")
```

```
output integral(sqrt(x^4 + 3*x^2 + 2)*(315*x^2 + 946)/(5*x^6 + 22*x^4 + 31*x^2 +
14), x)
```

**3.32.6 Sympy [F]**

$$\int \frac{946 + 315x^2}{(7 + 5x^2)\sqrt{2 + 3x^2 + x^4}} dx = \int \frac{315x^2 + 946}{\sqrt{(x^2 + 1)(x^2 + 2)}(5x^2 + 7)} dx$$

input `integrate((315*x**2+946)/(5*x**2+7)/(x**4+3*x**2+2)**(1/2),x)`

output `Integral((315*x**2 + 946)/(sqrt((x**2 + 1)*(x**2 + 2))*(5*x**2 + 7)), x)`

**3.32.7 Maxima [F]**

$$\int \frac{946 + 315x^2}{(7 + 5x^2)\sqrt{2 + 3x^2 + x^4}} dx = \int \frac{315x^2 + 946}{\sqrt{x^4 + 3x^2 + 2}(5x^2 + 7)} dx$$

input `integrate((315*x^2+946)/(5*x^2+7)/(x^4+3*x^2+2)^(1/2),x, algorithm="maxima")`

output `integrate((315*x^2 + 946)/(sqrt(x^4 + 3*x^2 + 2)*(5*x^2 + 7)), x)`

**3.32.8 Giac [F]**

$$\int \frac{946 + 315x^2}{(7 + 5x^2)\sqrt{2 + 3x^2 + x^4}} dx = \int \frac{315x^2 + 946}{\sqrt{x^4 + 3x^2 + 2}(5x^2 + 7)} dx$$

input `integrate((315*x^2+946)/(5*x^2+7)/(x^4+3*x^2+2)^(1/2),x, algorithm="giac")`

output `integrate((315*x^2 + 946)/(sqrt(x^4 + 3*x^2 + 2)*(5*x^2 + 7)), x)`

**3.32.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{946 + 315x^2}{(7 + 5x^2)\sqrt{2 + 3x^2 + x^4}} dx = \int \frac{315x^2 + 946}{(5x^2 + 7)\sqrt{x^4 + 3x^2 + 2}} dx$$

input `int((315*x^2 + 946)/((5*x^2 + 7)*(3*x^2 + x^4 + 2)^(1/2)), x)`output `int((315*x^2 + 946)/((5*x^2 + 7)*(3*x^2 + x^4 + 2)^(1/2)), x)`

### 3.33 $\int \frac{(A+Bx^2)(d+ex^2)^q}{a+bx^2+cx^4} dx$

|        |                     |     |
|--------|---------------------|-----|
| 3.33.1 | Optimal result      | 275 |
| 3.33.2 | Mathematica [F]     | 276 |
| 3.33.3 | Rubi [A] (verified) | 276 |
| 3.33.4 | Maple [F]           | 277 |
| 3.33.5 | Fricas [F]          | 277 |
| 3.33.6 | Sympy [F(-1)]       | 277 |
| 3.33.7 | Maxima [F]          | 278 |
| 3.33.8 | Giac [F]            | 278 |
| 3.33.9 | Mupad [F(-1)]       | 278 |

#### 3.33.1 Optimal result

Integrand size = 31, antiderivative size = 218

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + bx^2 + cx^4} dx$$

$$= \frac{\left(B - \frac{bB-2Ac}{\sqrt{b^2-4ac}}\right) x(d + ex^2)^q \left(1 + \frac{ex^2}{d}\right)^{-q} \text{AppellF1}\left(\frac{1}{2}, 1, -q, \frac{3}{2}, -\frac{2cx^2}{b-\sqrt{b^2-4ac}}, -\frac{ex^2}{d}\right)}{b - \sqrt{b^2 - 4ac}}$$

$$+ \frac{\left(B + \frac{bB-2Ac}{\sqrt{b^2-4ac}}\right) x(d + ex^2)^q \left(1 + \frac{ex^2}{d}\right)^{-q} \text{AppellF1}\left(\frac{1}{2}, 1, -q, \frac{3}{2}, -\frac{2cx^2}{b+\sqrt{b^2-4ac}}, -\frac{ex^2}{d}\right)}{b + \sqrt{b^2 - 4ac}}$$

```
output x*(e*x^2+d)^q*AppellF1(1/2,1,-q,3/2,-2*c*x^2/(b-(-4*a*c+b^2)^(1/2)),-e*x^2/d)*(B+(2*A*c-B*b)/(-4*a*c+b^2)^(1/2))/((1+e*x^2/d)^q)/(b-(-4*a*c+b^2)^(1/2))+x*(e*x^2+d)^q*AppellF1(1/2,1,-q,3/2,-2*c*x^2/(b+(-4*a*c+b^2)^(1/2)),-e*x^2/d)*(B+(-2*A*c+B*b)/(-4*a*c+b^2)^(1/2))/((1+e*x^2/d)^q)/(b+(-4*a*c+b^2)^(1/2))
```

### 3.33.2 Mathematica [F]

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + bx^2 + cx^4} dx = \int \frac{(A + Bx^2)(d + ex^2)^q}{a + bx^2 + cx^4} dx$$

input `Integrate[((A + B*x^2)*(d + e*x^2)^q)/(a + b*x^2 + c*x^4), x]`

output `Integrate[((A + B*x^2)*(d + e*x^2)^q)/(a + b*x^2 + c*x^4), x]`

### 3.33.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.065$ , Rules used = {2256, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(A + Bx^2)(d + ex^2)^q}{a + bx^2 + cx^4} dx \\ & \quad \downarrow \text{2256} \\ & \int \left( \frac{(d + ex^2)^q \left( \frac{2Ac - bB}{\sqrt{b^2 - 4ac}} + B \right)}{-\sqrt{b^2 - 4ac} + b + 2cx^2} + \frac{(d + ex^2)^q \left( B - \frac{2Ac - bB}{\sqrt{b^2 - 4ac}} \right)}{\sqrt{b^2 - 4ac} + b + 2cx^2} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{x(d + ex^2)^q \left( \frac{ex^2}{d} + 1 \right)^{-q} \left( B - \frac{bB - 2Ac}{\sqrt{b^2 - 4ac}} \right) \text{AppellF1} \left( \frac{1}{2}, 1, -q, \frac{3}{2}, -\frac{2cx^2}{b - \sqrt{b^2 - 4ac}}, -\frac{ex^2}{d} \right)}{b - \sqrt{b^2 - 4ac}} + \\ & \frac{x(d + ex^2)^q \left( \frac{ex^2}{d} + 1 \right)^{-q} \left( \frac{bB - 2Ac}{\sqrt{b^2 - 4ac}} + B \right) \text{AppellF1} \left( \frac{1}{2}, 1, -q, \frac{3}{2}, -\frac{2cx^2}{b + \sqrt{b^2 - 4ac}}, -\frac{ex^2}{d} \right)}{\sqrt{b^2 - 4ac} + b} \end{aligned}$$

input `Int[((A + B*x^2)*(d + e*x^2)^q)/(a + b*x^2 + c*x^4), x]`

output `((B - (b*B - 2*A*c)/Sqrt[b^2 - 4*a*c])*x*(d + e*x^2)^q*AppellF1[1/2, 1, -q, 3/2, (-2*c*x^2)/(b - Sqrt[b^2 - 4*a*c]), -(e*x^2)/d])/((b - Sqrt[b^2 - 4*a*c])*(1 + (e*x^2)/d)^q) + ((B + (b*B - 2*A*c)/Sqrt[b^2 - 4*a*c])*x*(d + e*x^2)^q*AppellF1[1/2, 1, -q, 3/2, (-2*c*x^2)/(b + Sqrt[b^2 - 4*a*c]), -(e*x^2)/d])/((b + Sqrt[b^2 - 4*a*c])*(1 + (e*x^2)/d)^q)`

---

3.33.  $\int \frac{(A+Bx^2)(d+ex^2)^q}{a+bx^2+cx^4} dx$

## 3.33.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2256 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`

## 3.33.4 Maple [F]

$$\int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + bx^2 + a} dx$$

input `int((B*x^2+A)*(e*x^2+d)^q/(c*x^4+b*x^2+a),x)`

output `int((B*x^2+A)*(e*x^2+d)^q/(c*x^4+b*x^2+a),x)`

## 3.33.5 Fricas [F]

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + bx^2 + cx^4} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + bx^2 + a} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+b*x^2+a),x, algorithm="fricas")`

output `integral((B*x^2 + A)*(e*x^2 + d)^q/(c*x^4 + b*x^2 + a), x)`

## 3.33.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + bx^2 + cx^4} dx = \text{Timed out}$$

input `integrate((B*x**2+A)*(e*x**2+d)**q/(c*x**4+b*x**2+a),x)`

output `Timed out`

---

3.33.  $\int \frac{(A+Bx^2)(d+ex^2)^q}{a+bx^2+cx^4} dx$

**3.33.7 Maxima [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + bx^2 + cx^4} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + bx^2 + a} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+b*x^2+a),x, algorithm="maxima")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^q/(c*x^4 + b*x^2 + a), x)`

**3.33.8 Giac [F]**

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + bx^2 + cx^4} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + bx^2 + a} dx$$

input `integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+b*x^2+a),x, algorithm="giac")`

output `integrate((B*x^2 + A)*(e*x^2 + d)^q/(c*x^4 + b*x^2 + a), x)`

**3.33.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + bx^2 + cx^4} dx = \int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + bx^2 + a} dx$$

input `int(((A + B*x^2)*(d + e*x^2)^q)/(a + b*x^2 + c*x^4),x)`

output `int(((A + B*x^2)*(d + e*x^2)^q)/(a + b*x^2 + c*x^4), x)`

**3.34** 
$$\int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx$$

|        |   |     |
|--------|---|-----|
| 3.34.1 | Optimal result                            | 279 |
| 3.34.2 | Mathematica [C] (verified)                | 279 |
| 3.34.3 | Rubi [A] (warning: unable to verify)      | 280 |
| 3.34.4 | Maple [A] (verified)                      | 283 |
| 3.34.5 | Fricas [B] (verification not implemented) | 284 |
| 3.34.6 | Sympy [F]                                 | 285 |
| 3.34.7 | Maxima [F]                                | 285 |
| 3.34.8 | Giac [A] (verification not implemented)   | 285 |
| 3.34.9 | Mupad [B] (verification not implemented)  | 286 |

**3.34.1 Optimal result**

Integrand size = 28, antiderivative size = 106

$$\int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx = -\frac{1}{2} \arctan(\sqrt{3}-2\sqrt{1+x^2}) + \frac{1}{2} \arctan(\sqrt{3}+2\sqrt{1+x^2}) + \frac{1}{4}\sqrt{3} \log(2+x^2-\sqrt{3}\sqrt{1+x^2}) - \frac{1}{4}\sqrt{3} \log(2+x^2+\sqrt{3}\sqrt{1+x^2})$$

output `1/2*arctan(-3^(1/2)+2*(x^2+1)^(1/2))+1/2*arctan(3^(1/2)+2*(x^2+1)^(1/2))+1/4*ln(2+x^2-3^(1/2)*(x^2+1)^(1/2))*3^(1/2)-1/4*ln(2+x^2+3^(1/2)*(x^2+1)^(1/2))*3^(1/2)`

**3.34.2 Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.02 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.76

$$\int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx = \frac{1}{2}(1-i\sqrt{3}) \arctan\left(\frac{1}{2}(1-i\sqrt{3})\sqrt{1+x^2}\right) + \frac{1}{2}(1+i\sqrt{3}) \arctan\left(\frac{1}{2}(1+i\sqrt{3})\sqrt{1+x^2}\right)$$



input `Integrate[(x*(1 + 2*x^2))/(Sqrt[1 + x^2]*(1 + x^2 + x^4)),x]`

output `((1 - I*Sqrt[3])*ArcTan[((1 - I*Sqrt[3])*Sqrt[1 + x^2])/2])/2 + ((1 + I*Sqrt[3])*ArcTan[((1 + I*Sqrt[3])*Sqrt[1 + x^2])/2])/2`

### 3.34.3 Rubi [A] (warning: unable to verify)

Time = 0.38 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$ , Rules used = {2238, 1197, 25, 1483, 27, 1142, 25, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x(2x^2 + 1)}{\sqrt{x^2 + 1}(x^4 + x^2 + 1)} dx \\
 & \quad \downarrow \text{2238} \\
 & \frac{1}{2} \int \frac{2x^2 + 1}{\sqrt{x^2 + 1}(x^4 + x^2 + 1)} dx^2 \\
 & \quad \downarrow \text{1197} \\
 & \int -\frac{1 - 2x^4}{x^8 - x^4 + 1} d\sqrt{x^2 + 1} \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{1 - 2x^4}{x^8 - x^4 + 1} d\sqrt{x^2 + 1} \\
 & \quad \downarrow \text{1483} \\
 & -\frac{\int \frac{\sqrt{3}-3\sqrt{x^2+1}}{x^4-\sqrt{3}\sqrt{x^2+1}+1} d\sqrt{x^2+1}}{2\sqrt{3}} - \frac{\int \frac{\sqrt{3}(\sqrt{3}\sqrt{x^2+1}+1)}{x^4+\sqrt{3}\sqrt{x^2+1}+1} d\sqrt{x^2+1}}{2\sqrt{3}} \\
 & \quad \downarrow \text{27} \\
 & -\frac{\int \frac{\sqrt{3}-3\sqrt{x^2+1}}{x^4-\sqrt{3}\sqrt{x^2+1}+1} d\sqrt{x^2+1}}{2\sqrt{3}} - \frac{1}{2} \int \frac{\sqrt{3}\sqrt{x^2+1}+1}{x^4+\sqrt{3}\sqrt{x^2+1}+1} d\sqrt{x^2+1} \\
 & \quad \downarrow \text{1142}
 \end{aligned}$$

---

3.34.  $\int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx$

$$\begin{aligned}
& \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{x^4 + \sqrt{3}\sqrt{x^2+1} + 1} d\sqrt{x^2+1} - \frac{1}{2} \sqrt{3} \int \frac{2\sqrt{x^2+1} + \sqrt{3}}{x^4 + \sqrt{3}\sqrt{x^2+1} + 1} d\sqrt{x^2+1} \right) - \\
& \quad \frac{-\frac{1}{2} \sqrt{3} \int \frac{1}{x^4 - \sqrt{3}\sqrt{x^2+1} + 1} d\sqrt{x^2+1} - \frac{3}{2} \int \frac{-\sqrt{3} - 2\sqrt{x^2+1}}{x^4 - \sqrt{3}\sqrt{x^2+1} + 1} d\sqrt{x^2+1}}{2\sqrt{3}} \\
& \quad \downarrow \text{25} \\
& \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{x^4 + \sqrt{3}\sqrt{x^2+1} + 1} d\sqrt{x^2+1} - \frac{1}{2} \sqrt{3} \int \frac{2\sqrt{x^2+1} + \sqrt{3}}{x^4 + \sqrt{3}\sqrt{x^2+1} + 1} d\sqrt{x^2+1} \right) - \\
& \quad \frac{\frac{3}{2} \int \frac{\sqrt{3} - 2\sqrt{x^2+1}}{x^4 - \sqrt{3}\sqrt{x^2+1} + 1} d\sqrt{x^2+1} - \frac{1}{2} \sqrt{3} \int \frac{1}{x^4 - \sqrt{3}\sqrt{x^2+1} + 1} d\sqrt{x^2+1}}{2\sqrt{3}} \\
& \quad \downarrow \text{1083} \\
& \frac{1}{2} \left( - \int \frac{1}{-x^4 - 1} d(2\sqrt{x^2+1} + \sqrt{3}) - \frac{1}{2} \sqrt{3} \int \frac{2\sqrt{x^2+1} + \sqrt{3}}{x^4 + \sqrt{3}\sqrt{x^2+1} + 1} d\sqrt{x^2+1} \right) - \\
& \quad \frac{\sqrt{3} \int \frac{1}{-x^4 - 1} d(2\sqrt{x^2+1} - \sqrt{3}) + \frac{3}{2} \int \frac{\sqrt{3} - 2\sqrt{x^2+1}}{x^4 - \sqrt{3}\sqrt{x^2+1} + 1} d\sqrt{x^2+1}}{2\sqrt{3}} \\
& \quad \downarrow \text{217} \\
& \frac{1}{2} \left( \arctan(2\sqrt{x^2+1} + \sqrt{3}) - \frac{1}{2} \sqrt{3} \int \frac{2\sqrt{x^2+1} + \sqrt{3}}{x^4 + \sqrt{3}\sqrt{x^2+1} + 1} d\sqrt{x^2+1} \right) - \\
& \quad \frac{\frac{3}{2} \int \frac{\sqrt{3} - 2\sqrt{x^2+1}}{x^4 - \sqrt{3}\sqrt{x^2+1} + 1} d\sqrt{x^2+1} + \sqrt{3} \arctan(\sqrt{3} - 2\sqrt{x^2+1})}{2\sqrt{3}} \\
& \quad \downarrow \text{1103} \\
& \frac{1}{2} \left( \arctan(2\sqrt{x^2+1} + \sqrt{3}) - \frac{1}{2} \sqrt{3} \log(x^4 + \sqrt{3}\sqrt{x^2+1} + 1) \right) - \\
& \quad \frac{\sqrt{3} \arctan(\sqrt{3} - 2\sqrt{x^2+1}) - \frac{3}{2} \log(x^4 - \sqrt{3}\sqrt{x^2+1} + 1)}{2\sqrt{3}}
\end{aligned}$$

input `Int[(x*(1 + 2*x^2))/(Sqrt[1 + x^2]*(1 + x^2 + x^4)),x]`

output `-1/2*(Sqrt[3]*ArcTan[Sqrt[3] - 2*Sqrt[1 + x^2]] - (3*Log[1 + x^4 - Sqrt[3]*Sqrt[1 + x^2]])/2)/Sqrt[3] + (ArcTan[Sqrt[3] + 2*Sqrt[1 + x^2]] - (Sqrt[3]*Log[1 + x^4 + Sqrt[3]*Sqrt[1 + x^2]])/2)/2`

## 3.34.3.1 Defintions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, x], x]$
- rule 27  $\text{Int}[(a_)*(\text{Fx}_), x\_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[\text{Fx}, (b_)*(\text{Gx}_) /; \text{FreeQ}[b, x]]$
- rule 217  $\text{Int}[(a_) + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 1083  $\text{Int}[(a_) + (b_)*(x_) + (c_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[-2 \quad \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103  $\text{Int}[(d_) + (e_)*(x_)] / [(a_) + (b_)*(x_) + (c_)*(x_)^2], x\_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$
- rule 1142  $\text{Int}[(d_) + (e_)*(x_)] / [(a_) + (b_)*(x_) + (c_)*(x_)^2], x\_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \quad \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \quad \text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$
- rule 1197  $\text{Int}[(f_) + (g_)*(x_)] / (\text{Sqrt}[(d_) + (e_)*(x_)] * [(a_) + (b_)*(x_) + (c_)*(x_)^2]), x\_Symbol] \rightarrow \text{Simp}[2 \quad \text{Subst}[\text{Int}[(e*f - d*g + g*x^2)/(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4), x], x, \text{Sqrt}[d + e*x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g\}, x]$
- rule 1483  $\text{Int}[(d_) + (e_)*(x_)^2] / [(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a/c, 2]\}, \text{With}[\{r = \text{Rt}[2*q - b/c, 2]\}, \text{Simp}[1/(2*c*q*r) \quad \text{Int}[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + \text{Simp}[1/(2*c*q*r) \quad \text{Int}[(d*r + (d - e*q)*x)/(q + r*x + x^2), x], x]]] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \ \&\& \ \text{NegQ}[b^2 - 4*a*c]$

rule 2238 `Int[(Px_)*(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[(Px /. x -> Sqrt[x])*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && PolyQ[Px, x^2]`

### 3.34.4 Maple [A] (verified)

Time = 1.17 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.76

| method         | result  |
|----------------|---|
| pseudoelliptic | $\frac{\arctan\left(\frac{-\sqrt{3}+2\sqrt{x^2+1}}{2}\right)}{2} + \frac{\arctan\left(\frac{\sqrt{3}+2\sqrt{x^2+1}}{2}\right)}{2} + \frac{\ln\left(2+x^2-\sqrt{3}\sqrt{x^2+1}\right)\sqrt{3}}{4} - \frac{\ln\left(2+x^2+\sqrt{3}\sqrt{x^2+1}\right)\sqrt{3}}{4}$  |
| default        | $\frac{\sqrt{2}\sqrt{\frac{2(x-1)^2}{(-x-1)^2}+2}\left(\sqrt{3}\operatorname{arctanh}\left(\frac{\sqrt{\frac{2(x-1)^2}{(-x-1)^2}+2}\sqrt{3}}{2}\right)+\operatorname{arctan}\left(\frac{\sqrt{\frac{2(x-1)^2}{(-x-1)^2}+2}(x-1)}{\left(\frac{(x-1)^2}{(-x-1)^2}+1\right)(-x-1)}\right)\right)}{4\sqrt{\frac{(x-1)^2}{(-x-1)^2}+1}\left(\frac{x-1}{(-x-1)+1}\right)} - \frac{\sqrt{2}\sqrt{\frac{2(x+1)^2}{(1-x)^2}+2}\left(\sqrt{3}\operatorname{arctanh}\left(\frac{\sqrt{\frac{2(x+1)^2}{(1-x)^2}+2}\sqrt{3}}{2}\right)+\operatorname{arctan}\left(\frac{\sqrt{\frac{2(x+1)^2}{(1-x)^2}+2}(x+1)}{\left(\frac{(x+1)^2}{(1-x)^2}+1\right)(x-1)}\right)\right)}{4\sqrt{\frac{(x+1)^2}{(1-x)^2}+1}\left(\frac{x+1}{(-x-1)+1}\right)}$ |
| trager         | $-4\ln\left(\frac{-16\operatorname{RootOf}\left(16\_Z^4-4\_Z^2+1\right)^5x^2+16\operatorname{RootOf}\left(16\_Z^4-4\_Z^2+1\right)^3x^2+12\operatorname{RootOf}\left(16\_Z^4-4\_Z^2+1\right)^3}{\left(4\operatorname{RootOf}\left(16\_Z^4-4\_Z^2+1\right)^2x-x+1\right)\left(4\operatorname{RootOf}\left(16\_Z^4-4\_Z^2+1\right)^2x-x+1\right)}\right)$  |

input `int(x*(2*x^2+1)/(x^4+x^2+1)/(x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{2}\arctan(-3^{(1/2)}+2*(x^2+1)^{(1/2)})+\frac{1}{2}\arctan(3^{(1/2)}+2*(x^2+1)^{(1/2)})+\frac{1}{4}\ln(2+x^2-3^{(1/2)}*(x^2+1)^{(1/2)})*\frac{3^{(1/2)}}{2}-\frac{1}{4}\ln(2+x^2+3^{(1/2)}*(x^2+1)^{(1/2)})*\frac{3^{(1/2)}}{2}$$

**3.34.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 286 vs. 2(80) = 160.

Time = 0.33 (sec) , antiderivative size = 286, normalized size of antiderivative = 2.70

$$\int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx = \frac{1}{4} \sqrt{2} \sqrt{-\sqrt{-3}+1} \log \left( 4x^2 + \sqrt{2}(\sqrt{-3}x+x) \sqrt{-\sqrt{-3}+1} \right. \\ \left. - \sqrt{x^2+1} \left( \sqrt{2}(\sqrt{-3}+1) \sqrt{-\sqrt{-3}+1+4x} \right) + 4 \right) \\ - \frac{1}{4} \sqrt{2} \sqrt{-\sqrt{-3}+1} \log \left( 4x^2 \right. \\ \left. - \sqrt{2}(\sqrt{-3}x+x) \sqrt{-\sqrt{-3}+1} \right. \\ \left. + \sqrt{x^2+1} \left( \sqrt{2}(\sqrt{-3}+1) \sqrt{-\sqrt{-3}+1-4x} \right) + 4 \right) \\ + \frac{1}{4} \sqrt{2} \sqrt{\sqrt{-3}+1} \log \left( 4x^2 - 4\sqrt{x^2+1}x \right. \\ \left. + \left( \sqrt{2}\sqrt{x^2+1}(\sqrt{-3}-1) - \sqrt{2}(\sqrt{-3}x-x) \right) \sqrt{\sqrt{-3}+1} \right. \\ \left. + 4 \right) - \frac{1}{4} \sqrt{2} \sqrt{\sqrt{-3}+1} \log \left( 4x^2 - 4\sqrt{x^2+1}x \right. \\ \left. - \left( \sqrt{2}\sqrt{x^2+1}(\sqrt{-3}-1) - \sqrt{2}(\sqrt{-3}x-x) \right) \sqrt{\sqrt{-3}+1} \right. \\ \left. + 4 \right)$$

input `integrate(x*(2*x^2+1)/(x^4+x^2+1)/(x^2+1)^(1/2),x, algorithm="fracas")`

output `1/4*sqrt(2)*sqrt(-sqrt(-3)+1)*log(4*x^2+sqrt(2)*(sqrt(-3)*x+x)*sqrt(-sqrt(-3)+1)-sqrt(x^2+1)*(sqrt(2)*(sqrt(-3)+1)*sqrt(-sqrt(-3)+1)+4*x)+4)-1/4*sqrt(2)*sqrt(-sqrt(-3)+1)*log(4*x^2-sqrt(2)*(sqrt(-3)*x+x)*sqrt(-sqrt(-3)+1)+sqrt(x^2+1)*(sqrt(2)*(sqrt(-3)+1)*sqrt(-sqrt(-3)+1)-4*x)+4)+1/4*sqrt(2)*sqrt(sqrt(-3)+1)*log(4*x^2-4*sqrt(x^2+1)*x+(sqrt(2)*sqrt(x^2+1)*(sqrt(-3)-1)-sqrt(2)*(sqrt(-3)*x-x))*sqrt(sqrt(-3)+1)+4)-1/4*sqrt(2)*sqrt(sqrt(-3)+1)*log(4*x^2-4*sqrt(x^2+1)*x-(sqrt(2)*sqrt(x^2+1)*(sqrt(-3)-1)-sqrt(2)*(sqrt(-3)*x-x))*sqrt(sqrt(-3)+1)+4)`

**3.34.6 Sympy [F]**

$$\int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx = \int \frac{x(2x^2+1)}{\sqrt{x^2+1}(x^2-x+1)(x^2+x+1)} dx$$

input `integrate(x*(2*x**2+1)/(x**4+x**2+1)/(x**2+1)**(1/2),x)`

output `Integral(x*(2*x**2 + 1)/(sqrt(x**2 + 1)*(x**2 - x + 1)*(x**2 + x + 1)), x)`

**3.34.7 Maxima [F]**

$$\int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx = \int \frac{(2x^2+1)x}{(x^4+x^2+1)\sqrt{x^2+1}} dx$$

input `integrate(x*(2*x^2+1)/(x^4+x^2+1)/(x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate((2*x^2 + 1)*x/((x^4 + x^2 + 1)*sqrt(x^2 + 1)), x)`

**3.34.8 Giac [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.75

$$\begin{aligned} \int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx = & -\frac{1}{4} \sqrt{3} \log(x^2 + \sqrt{3}\sqrt{x^2+1} + 2) \\ & + \frac{1}{4} \sqrt{3} \log(x^2 - \sqrt{3}\sqrt{x^2+1} + 2) \\ & + \frac{1}{2} \arctan(\sqrt{3} + 2\sqrt{x^2+1}) + \frac{1}{2} \arctan(-\sqrt{3} + 2\sqrt{x^2+1}) \end{aligned}$$

input `integrate(x*(2*x^2+1)/(x^4+x^2+1)/(x^2+1)^(1/2),x, algorithm="giac")`

output `-1/4*sqrt(3)*log(x^2 + sqrt(3)*sqrt(x^2 + 1) + 2) + 1/4*sqrt(3)*log(x^2 - sqrt(3)*sqrt(x^2 + 1) + 2) + 1/2*arctan(sqrt(3) + 2*sqrt(x^2 + 1)) + 1/2*arctan(-sqrt(3) + 2*sqrt(x^2 + 1))`

---

3.34.  $\int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx$

**3.34.9 Mupad [B] (verification not implemented)**

Time = 8.63 (sec) , antiderivative size = 397, normalized size of antiderivative = 3.75

$$\begin{aligned}
& \int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx \\
&= \frac{\left(\ln\left(x - \frac{1}{2} - \frac{\sqrt{3}li}{2}\right) - \ln\left(\frac{x}{2} + \left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) \sqrt{x^2+1} + 1 + \frac{\sqrt{3}xli}{2}\right)\right) \left(2\left(\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)^3 + \frac{1}{2} + \frac{\sqrt{3}li}{2}\right)}{\sqrt{\left(\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)^2 + 1} \left(4\left(\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)^3 + 1 + \sqrt{3}li\right)} \\
&+ \frac{\left(\ln\left(x + \frac{1}{2} - \frac{\sqrt{3}li}{2}\right) - \ln\left(1 + \left(\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) \sqrt{x^2+1} - \frac{x}{2} + \frac{\sqrt{3}xli}{2}\right)\right) \left(2\left(-\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)^3 - \frac{1}{2} + \frac{\sqrt{3}li}{2}\right)}{\sqrt{\left(-\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)^2 + 1} \left(4\left(-\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)^3 - 1 + \sqrt{3}li\right)} \\
&+ \frac{\left(\ln\left(x - \frac{1}{2} + \frac{\sqrt{3}li}{2}\right) - \ln\left(\frac{x}{2} + \left(\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) \sqrt{x^2+1} + 1 - \frac{\sqrt{3}xli}{2}\right)\right) \left(2\left(-\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)^3 - \frac{1}{2} + \frac{\sqrt{3}li}{2}\right)}{\sqrt{\left(-\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)^2 + 1} \left(4\left(-\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)^3 - 1 + \sqrt{3}li\right)} \\
&+ \frac{\left(\ln\left(x + \frac{1}{2} + \frac{\sqrt{3}li}{2}\right) - \ln\left(1 + \left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) \sqrt{x^2+1} - \frac{x}{2} - \frac{\sqrt{3}xli}{2}\right)\right) \left(2\left(\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)^3 + \frac{1}{2} + \frac{\sqrt{3}li}{2}\right)}{\sqrt{\left(\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)^2 + 1} \left(4\left(\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)^3 + 1 + \sqrt{3}li\right)}
\end{aligned}$$

input `int((x*(2*x^2 + 1))/((x^2 + 1)^(1/2)*(x^2 + x^4 + 1)),x)`

```

output ((log(x - (3^(1/2)*1i)/2 - 1/2) - log(x/2 + (3^(1/2)/2 + 1i/2)*(x^2 + 1)^(
1/2) + (3^(1/2)*x*1i)/2 + 1))*((3^(1/2)*1i)/2 + 2*((3^(1/2)*1i)/2 + 1/2)^3
+ 1/2))/((((3^(1/2)*1i)/2 + 1/2)^2 + 1)^(1/2)*(3^(1/2)*1i + 4*((3^(1/2)*1
i)/2 + 1/2)^3 + 1)) + ((log(x - (3^(1/2)*1i)/2 + 1/2) - log((3^(1/2)/2 - 1
i/2)*(x^2 + 1)^(1/2) - x/2 + (3^(1/2)*x*1i)/2 + 1))*((3^(1/2)*1i)/2 + 2*((
3^(1/2)*1i)/2 - 1/2)^3 - 1/2))/((((3^(1/2)*1i)/2 - 1/2)^2 + 1)^(1/2)*(3^(1
/2)*1i + 4*((3^(1/2)*1i)/2 - 1/2)^3 - 1)) + ((log(x + (3^(1/2)*1i)/2 - 1/2
) - log(x/2 + (3^(1/2)/2 - 1i/2)*(x^2 + 1)^(1/2) - (3^(1/2)*x*1i)/2 + 1))*
((3^(1/2)*1i)/2 + 2*((3^(1/2)*1i)/2 - 1/2)^3 - 1/2))/((((3^(1/2)*1i)/2 - 1
/2)^2 + 1)^(1/2)*(3^(1/2)*1i + 4*((3^(1/2)*1i)/2 - 1/2)^3 - 1)) + ((log(x
+ (3^(1/2)*1i)/2 + 1/2) - log((3^(1/2)/2 + 1i/2)*(x^2 + 1)^(1/2) - x/2 - (
3^(1/2)*x*1i)/2 + 1))*((3^(1/2)*1i)/2 + 2*((3^(1/2)*1i)/2 + 1/2)^3 + 1/2))
/((((3^(1/2)*1i)/2 + 1/2)^2 + 1)^(1/2)*(3^(1/2)*1i + 4*((3^(1/2)*1i)/2 + 1
/2)^3 + 1))

```

### 3.35 $\int \frac{\sqrt{a+bx^2+cx^4}}{ad-cdx^4} dx$

|        |   |     |
|--------|---|-----|
| 3.35.1 | Optimal result                            | 287 |
| 3.35.2 | Mathematica [A] (verified)                | 287 |
| 3.35.3 | Rubi [A] (verified)                       | 288 |
| 3.35.4 | Maple [A] (verified)                      | 289 |
| 3.35.5 | Fricas [B] (verification not implemented) | 290 |
| 3.35.6 | Sympy [F]                                 | 291 |
| 3.35.7 | Maxima [F]                                | 291 |
| 3.35.8 | Giac [F]                                  | 291 |
| 3.35.9 | Mupad [F(-1)]                             | 292 |

#### 3.35.1 Optimal result

Integrand size = 30, antiderivative size = 145

$$\int \frac{\sqrt{a+bx^2+cx^4}}{ad-cdx^4} dx = -\frac{\sqrt{b-2\sqrt{a}\sqrt{c}} \operatorname{arctanh}\left(\frac{\sqrt{b-2\sqrt{a}\sqrt{c}x}}{\sqrt{a+bx^2+cx^4}}\right)}{4\sqrt{a}\sqrt{cd}} + \frac{\sqrt{b+2\sqrt{a}\sqrt{c}} \operatorname{arctanh}\left(\frac{\sqrt{b+2\sqrt{a}\sqrt{c}x}}{\sqrt{a+bx^2+cx^4}}\right)}{4\sqrt{a}\sqrt{cd}}$$

output  $-1/4*\operatorname{arctanh}(x*(b-2*a^{(1/2)}*c^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)})*(b-2*a^{(1/2)}*c^{(1/2)})^{(1/2)}/d/a^{(1/2)}/c^{(1/2)}+1/4*\operatorname{arctanh}(x*(b+2*a^{(1/2)}*c^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)})*(b+2*a^{(1/2)}*c^{(1/2)})^{(1/2)}/d/a^{(1/2)}/c^{(1/2)}$

#### 3.35.2 Mathematica [A] (verified)

Time = 0.91 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.96

$$\int \frac{\sqrt{a+bx^2+cx^4}}{ad-cdx^4} dx = \frac{-\sqrt{-b-2\sqrt{a}\sqrt{c}} \operatorname{arctan}\left(\frac{\sqrt{-b-2\sqrt{a}\sqrt{c}x}}{\sqrt{a+bx^2+cx^4}}\right) + \sqrt{-b+2\sqrt{a}\sqrt{c}} \operatorname{arctan}\left(\frac{\sqrt{-b+2\sqrt{a}\sqrt{c}x}}{\sqrt{a+bx^2+cx^4}}\right)}{4\sqrt{a}\sqrt{cd}}$$

input `Integrate[Sqrt[a + b*x^2 + c*x^4]/(a*d - c*d*x^4),x]`



output  $(-\text{Sqrt}[-b - 2\text{Sqrt}[a]\text{Sqrt}[c]]\text{ArcTan}[(\text{Sqrt}[-b - 2\text{Sqrt}[a]\text{Sqrt}[c]]x)/\text{Sqrt}[a + b x^2 + c x^4]]) + \text{Sqrt}[-b + 2\text{Sqrt}[a]\text{Sqrt}[c]]\text{ArcTan}[(\text{Sqrt}[-b + 2\text{Sqrt}[a]\text{Sqrt}[c]]x)/\text{Sqrt}[a + b x^2 + c x^4]])/(4\text{Sqrt}[a]\text{Sqrt}[c]d)$

### 3.35.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.32, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {2517, 1406, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + bx^2 + cx^4}}{ad - cd x^4} dx$$

$$\downarrow 2517$$

$$\int \frac{1}{\frac{(b^2 - 4ac)x^4}{(cx^4 + bx^2 + a)^2} - \frac{2bx^2}{cx^4 + bx^2 + a} + 1} d \frac{x}{\sqrt{cx^4 + bx^2 + a}}$$

$$\downarrow 1406$$

$$\frac{(b^2 - 4ac) \int \frac{1}{\frac{(b^2 - 4ac)x^2}{cx^4 + bx^2 + a} - b - 2\sqrt{a}\sqrt{c}} d \frac{x}{\sqrt{cx^4 + bx^2 + a}}}{4\sqrt{a}\sqrt{c}} - \frac{(b^2 - 4ac) \int \frac{1}{\frac{(b^2 - 4ac)x^2}{cx^4 + bx^2 + a} - b + 2\sqrt{a}\sqrt{c}} d \frac{x}{\sqrt{cx^4 + bx^2 + a}}}{4\sqrt{a}\sqrt{c}}$$

$$\downarrow 221$$

$$\frac{(b^2 - 4ac) \text{arctanh}\left(\frac{x\sqrt{2\sqrt{a}\sqrt{c}+b}}{\sqrt{a+bx^2+cx^4}}\right)}{4\sqrt{a}\sqrt{c}(b-2\sqrt{a}\sqrt{c})\sqrt{2\sqrt{a}\sqrt{c}+b}} - \frac{(b^2 - 4ac) \text{arctanh}\left(\frac{x\sqrt{b-2\sqrt{a}\sqrt{c}}}{\sqrt{a+bx^2+cx^4}}\right)}{4\sqrt{a}\sqrt{c}\sqrt{b-2\sqrt{a}\sqrt{c}}(2\sqrt{a}\sqrt{c}+b)}$$

$$d$$

input `Int[Sqrt[a + b*x^2 + c*x^4]/(a*d - c*d*x^4), x]`

output  $(-1/4*((b^2 - 4*a*c)*\text{ArcTanh}[(\text{Sqrt}[b - 2\text{Sqrt}[a]\text{Sqrt}[c]]x)/\text{Sqrt}[a + b x^2 + c x^4]])/(\text{Sqrt}[a]\text{Sqrt}[b - 2\text{Sqrt}[a]\text{Sqrt}[c]]*(b + 2\text{Sqrt}[a]\text{Sqrt}[c])*\text{Sqrt}[c]) + ((b^2 - 4*a*c)*\text{ArcTanh}[(\text{Sqrt}[b + 2\text{Sqrt}[a]\text{Sqrt}[c]]x)/\text{Sqrt}[a + b x^2 + c x^4]])/(4\text{Sqrt}[a]*(b - 2\text{Sqrt}[a]\text{Sqrt}[c])*\text{Sqrt}[b + 2\text{Sqrt}[a]\text{Sqrt}[c]]*\text{Sqrt}[c]))/d$

## 3.35.3.1 Defintions of rubi rules used

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 1406 `Int[((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c/q Int[1/(b/2 - q/2 + c*x^2), x], x] - Simp[c/q Int[1/(b/2 + q/2 + c*x^2), x], x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[b^2 - 4*a*c]`

rule 2517 `Int[Sqrt[v_] / ((d_) + (e_.)*(x_)^4), x_Symbol] := With[{a = Coeff[v, x, 0], b = Coeff[v, x, 2], c = Coeff[v, x, 4]}, Simp[a/d Subst[Int[1/(1 - 2*b*x^2 + (b^2 - 4*a*c)*x^4), x], x, x/Sqrt[v]], x] /; EqQ[c*d + a*e, 0] && PosQ[a*c]] /; FreeQ[{d, e}, x] && PolyQ[v, x^2, 2]`

## 3.35.4 Maple [A] (verified)

Time = 1.67 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.87

| method         | result   | size |
|----------------|--|------|
| pseudoelliptic | $\frac{(2\sqrt{ac}+b) \arctan\left(\frac{\sqrt{cx^4+bx^2+a}}{x\sqrt{-2\sqrt{ac}-b}}\right) + \sqrt{2\sqrt{ac}-b} \arctan\left(\frac{\sqrt{cx^4+bx^2+a}}{x\sqrt{2\sqrt{ac}-b}}\right) \sqrt{-2\sqrt{ac}-b}}{4\sqrt{-2\sqrt{ac}-b}\sqrt{ac}d}$   | 126  |
| default        | $\frac{\left(-\frac{(2\sqrt{ac}+b) \arctan\left(\frac{\sqrt{cx^4+bx^2+a}\sqrt{2}}{x\sqrt{-4\sqrt{ac}-2b}}\right)}{2\sqrt{ac}\sqrt{-4\sqrt{ac}-2b}} - \frac{(2\sqrt{ac}-b) \arctan\left(\frac{\sqrt{cx^4+bx^2+a}\sqrt{2}}{x\sqrt{4\sqrt{ac}-2b}}\right)}{2\sqrt{ac}\sqrt{4\sqrt{ac}-2b}}\right)\sqrt{2}}{2d}$ | 140  |
| elliptic       | $2\frac{\left(-\frac{(2\sqrt{ac}+b) \arctan\left(\frac{\sqrt{cx^4+bx^2+a}\sqrt{2}}{x\sqrt{-4\sqrt{ac}-2b}}\right)}{8\sqrt{ac}\sqrt{-4\sqrt{ac}-2b}} - \frac{(2\sqrt{ac}-b) \arctan\left(\frac{\sqrt{cx^4+bx^2+a}\sqrt{2}}{x\sqrt{4\sqrt{ac}-2b}}\right)}{8\sqrt{ac}\sqrt{4\sqrt{ac}-2b}}\right)\sqrt{2}}{d}$ | 140  |

input `int((c*x^4+b*x^2+a)^(1/2)/(-c*d*x^4+a*d), x, method=_RETURNVERBOSE)`

output 
$$-1/4/(-2*(a*c)^(1/2)-b)^(1/2)/(a*c)^(1/2)*((2*(a*c)^(1/2)+b)*\arctan((c*x^4+b*x^2+a)^(1/2)/x/(-2*(a*c)^(1/2)-b)^(1/2))+(2*(a*c)^(1/2)-b)^(1/2)*\arctan((c*x^4+b*x^2+a)^(1/2)/x/(2*(a*c)^(1/2)-b)^(1/2))*(-2*(a*c)^(1/2)-b)^(1/2))/d$$

### 3.35.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 603 vs.  $2(105) = 210$ .

Time = 1.78 (sec) , antiderivative size = 603, normalized size of antiderivative = 4.16

$$\int \frac{\sqrt{a+bx^2+cx^4}}{ad-cdx^4} dx$$

$$= \frac{1}{8} \sqrt{\frac{2acd^2\sqrt{\frac{1}{acd^4}}+b}{acd^2}} \log\left(\frac{\sqrt{cx^4+bx^2+a}\left(ad^2\sqrt{\frac{1}{acd^4}}+x^2\right)+\left(acd^3x^3\sqrt{\frac{1}{acd^4}}+adx\right)\sqrt{\frac{2acd^2\sqrt{\frac{1}{acd^4}}+b}{acd^2}}}{cx^4-a}\right)$$

$$- \frac{1}{8} \sqrt{\frac{2acd^2\sqrt{\frac{1}{acd^4}}+b}{acd^2}} \log\left(\frac{\sqrt{cx^4+bx^2+a}\left(ad^2\sqrt{\frac{1}{acd^4}}+x^2\right)-\left(acd^3x^3\sqrt{\frac{1}{acd^4}}+adx\right)\sqrt{\frac{2acd^2\sqrt{\frac{1}{acd^4}}+b}{acd^2}}}{cx^4-a}\right)$$

$$+ \frac{1}{8} \sqrt{-\frac{2acd^2\sqrt{\frac{1}{acd^4}}-b}{acd^2}} \log\left(-\frac{\sqrt{cx^4+bx^2+a}\left(ad^2\sqrt{\frac{1}{acd^4}}-x^2\right)+\left(acd^3x^3\sqrt{\frac{1}{acd^4}}-adx\right)\sqrt{-\frac{2acd^2\sqrt{\frac{1}{acd^4}}-b}{acd^2}}}{cx^4-a}\right)$$

$$- \frac{1}{8} \sqrt{-\frac{2acd^2\sqrt{\frac{1}{acd^4}}-b}{acd^2}} \log\left(-\frac{\sqrt{cx^4+bx^2+a}\left(ad^2\sqrt{\frac{1}{acd^4}}-x^2\right)-\left(acd^3x^3\sqrt{\frac{1}{acd^4}}-adx\right)\sqrt{-\frac{2acd^2\sqrt{\frac{1}{acd^4}}-b}{acd^2}}}{cx^4-a}\right)$$

```
input integrate((c*x^4+b*x^2+a)^(1/2)/(-c*d*x^4+a*d),x, algorithm="fracas")
```

```
output 1/8*sqrt((2*a*c*d^2*sqrt(1/(a*c*d^4)) + b)/(a*c*d^2))*log((sqrt(c*x^4 + b
x^2 + a)*(a*d^2*sqrt(1/(a*c*d^4)) + x^2) + (a*c*d^3*x^3*sqrt(1/(a*c*d^4))
+ a*d*x)*sqrt((2*a*c*d^2*sqrt(1/(a*c*d^4)) + b)/(a*c*d^2)))/(c*x^4 - a)) -
1/8*sqrt((2*a*c*d^2*sqrt(1/(a*c*d^4)) + b)/(a*c*d^2))*log((sqrt(c*x^4 + b
*x^2 + a)*(a*d^2*sqrt(1/(a*c*d^4)) + x^2) - (a*c*d^3*x^3*sqrt(1/(a*c*d^4))
+ a*d*x)*sqrt((2*a*c*d^2*sqrt(1/(a*c*d^4)) + b)/(a*c*d^2)))/(c*x^4 - a))
+ 1/8*sqrt(-(2*a*c*d^2*sqrt(1/(a*c*d^4)) - b)/(a*c*d^2))*log(-(sqrt(c*x^4
+ b*x^2 + a)*(a*d^2*sqrt(1/(a*c*d^4)) - x^2) + (a*c*d^3*x^3*sqrt(1/(a*c*d^
4)) - a*d*x)*sqrt(-(2*a*c*d^2*sqrt(1/(a*c*d^4)) - b)/(a*c*d^2)))/(c*x^4 -
a)) - 1/8*sqrt(-(2*a*c*d^2*sqrt(1/(a*c*d^4)) - b)/(a*c*d^2))*log(-(sqrt(c*
x^4 + b*x^2 + a)*(a*d^2*sqrt(1/(a*c*d^4)) - x^2) - (a*c*d^3*x^3*sqrt(1/(a*
c*d^4)) - a*d*x)*sqrt(-(2*a*c*d^2*sqrt(1/(a*c*d^4)) - b)/(a*c*d^2)))/(c*x^
4 - a))
```

**3.35.6 Sympy [F]**

$$\int \frac{\sqrt{a + bx^2 + cx^4}}{ad - cdx^4} dx = -\int \frac{\sqrt{a+bx^2+cx^4}}{-a+cx^4} dx$$

input `integrate((c*x**4+b*x**2+a)**(1/2)/(-c*d*x**4+a*d), x)`

output `-Integral(sqrt(a + b*x**2 + c*x**4)/(-a + c*x**4), x)/d`

**3.35.7 Maxima [F]**

$$\int \frac{\sqrt{a + bx^2 + cx^4}}{ad - cdx^4} dx = \int -\frac{\sqrt{cx^4 + bx^2 + a}}{cdx^4 - ad} dx$$

input `integrate((c*x^4+b*x^2+a)^(1/2)/(-c*d*x^4+a*d), x, algorithm="maxima")`

output `-integrate(sqrt(c*x^4 + b*x^2 + a)/(c*d*x^4 - a*d), x)`

**3.35.8 Giac [F]**

$$\int \frac{\sqrt{a + bx^2 + cx^4}}{ad - cdx^4} dx = \int -\frac{\sqrt{cx^4 + bx^2 + a}}{cdx^4 - ad} dx$$

input `integrate((c*x^4+b*x^2+a)^(1/2)/(-c*d*x^4+a*d), x, algorithm="giac")`

output `integrate(-sqrt(c*x^4 + b*x^2 + a)/(c*d*x^4 - a*d), x)`

**3.35.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{a + bx^2 + cx^4}}{ad - cd x^4} dx = \int \frac{\sqrt{cx^4 + bx^2 + a}}{ad - cd x^4} dx$$

input `int((a + b*x^2 + c*x^4)^(1/2)/(a*d - c*d*x^4),x)`output `int((a + b*x^2 + c*x^4)^(1/2)/(a*d - c*d*x^4), x)`

### 3.36 $\int \frac{\sqrt{a+bx^2-cx^4}}{ad+cdx^4} dx$

|        |   |     |
|--------|---|-----|
| 3.36.1 | Optimal result                            | 293 |
| 3.36.2 | Mathematica [C] (verified)                | 293 |
| 3.36.3 | Rubi [A] (verified)                       | 294 |
| 3.36.4 | Maple [A] (verified)                      | 295 |
| 3.36.5 | Fricas [B] (verification not implemented) | 296 |
| 3.36.6 | Sympy [F]                                 | 297 |
| 3.36.7 | Maxima [F]                                | 297 |
| 3.36.8 | Giac [F]                                  | 298 |
| 3.36.9 | Mupad [F(-1)]                             | 298 |

#### 3.36.1 Optimal result

Integrand size = 30, antiderivative size = 239

$$\int \frac{\sqrt{a+bx^2-cx^4}}{ad+cdx^4} dx = -\frac{\sqrt{b+\sqrt{b^2+4ac}} \arctan\left(\frac{\sqrt{b+\sqrt{b^2+4ac}}(b-\sqrt{b^2+4ac}-2cx^2)}{2\sqrt{2}\sqrt{a}\sqrt{c}\sqrt{a+bx^2-cx^4}}\right)}{2\sqrt{2}\sqrt{a}\sqrt{cd}} + \frac{\sqrt{-b+\sqrt{b^2+4ac}} \operatorname{arctanh}\left(\frac{\sqrt{-b+\sqrt{b^2+4ac}}(b+\sqrt{b^2+4ac}-2cx^2)}{2\sqrt{2}\sqrt{a}\sqrt{c}\sqrt{a+bx^2-cx^4}}\right)}{2\sqrt{2}\sqrt{a}\sqrt{cd}}$$

output  $\frac{1}{4} \operatorname{arctanh}\left(\frac{1}{4} x (b - 2c x^2 + (4ac + b^2)^{1/2})^{1/2}\right) (-b + (4ac + b^2)^{1/2})^{1/2} (1/2) * 2^{1/2} / a^{1/2} / c^{1/2} / (-cx^4 + bx^2 + a)^{1/2} * (-b + (4ac + b^2)^{1/2})^{1/2} (1/2) / d * 2^{1/2} / a^{1/2} / c^{1/2} - \frac{1}{4} \operatorname{arctan}\left(\frac{1}{4} x (b - 2c x^2 - (4ac + b^2)^{1/2})^{1/2}\right) (b + (4ac + b^2)^{1/2})^{1/2} (1/2) * 2^{1/2} / a^{1/2} / c^{1/2} / (-cx^4 + bx^2 + a)^{1/2} * (b + (4ac + b^2)^{1/2})^{1/2} (1/2) / d * 2^{1/2} / a^{1/2} / c^{1/2}$

#### 3.36.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.64 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.63

$$\int \frac{\sqrt{a+bx^2-cx^4}}{ad+cdx^4} dx = \frac{i\left(\sqrt{-b-2i\sqrt{a}\sqrt{c}} \arctan\left(\frac{\sqrt{-b-2i\sqrt{a}\sqrt{c}}}{\sqrt{a+bx^2-cx^4}}\right) - \sqrt{-b+2i\sqrt{a}\sqrt{c}} \arctan\left(\frac{\sqrt{-b+2i\sqrt{a}\sqrt{c}}}{\sqrt{a+bx^2-cx^4}}\right)\right)}{4\sqrt{a}\sqrt{cd}}$$

input `Integrate[Sqrt[a + b*x^2 - c*x^4]/(a*d + c*d*x^4),x]`

output `((I/4)*(Sqrt[-b - (2*I)*Sqrt[a]*Sqrt[c]]*ArcTan[(Sqrt[-b - (2*I)*Sqrt[a]*Sqrt[c]]*x)/Sqrt[a + b*x^2 - c*x^4]] - Sqrt[-b + (2*I)*Sqrt[a]*Sqrt[c]]*ArcTan[(Sqrt[-b + (2*I)*Sqrt[a]*Sqrt[c]]*x)/Sqrt[a + b*x^2 - c*x^4]])/(Sqrt[a]*Sqrt[c]*d)`

### 3.36.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.033$ , Rules used = {2518}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + bx^2 - cx^4}}{ad + cd x^4} dx$$

↓ 2518

$$\frac{\sqrt{\sqrt{4ac + b^2} - b} \operatorname{arctanh}\left(\frac{x\sqrt{\sqrt{4ac + b^2} - b}(\sqrt{4ac + b^2 + b - 2cx^2})}{2\sqrt{2}\sqrt{a}\sqrt{c}\sqrt{a + bx^2 - cx^4}}\right)}{2\sqrt{2}\sqrt{a}\sqrt{cd}} - \frac{\sqrt{\sqrt{4ac + b^2} + b} \operatorname{arctan}\left(\frac{x\sqrt{\sqrt{4ac + b^2} + b}(-\sqrt{4ac + b^2 + b - 2cx^2})}{2\sqrt{2}\sqrt{a}\sqrt{c}\sqrt{a + bx^2 - cx^4}}\right)}{2\sqrt{2}\sqrt{a}\sqrt{cd}}$$

input `Int[Sqrt[a + b*x^2 - c*x^4]/(a*d + c*d*x^4),x]`

output `-1/2*(Sqrt[b + Sqrt[b^2 + 4*a*c]]*ArcTan[(Sqrt[b + Sqrt[b^2 + 4*a*c]]*x*(b - Sqrt[b^2 + 4*a*c] - 2*c*x^2))/(2*Sqrt[2]*Sqrt[a]*Sqrt[c]*Sqrt[a + b*x^2 - c*x^4])]/(Sqrt[2]*Sqrt[a]*Sqrt[c]*d) + (Sqrt[-b + Sqrt[b^2 + 4*a*c]]*ArcTanh[(Sqrt[-b + Sqrt[b^2 + 4*a*c]]*x*(b + Sqrt[b^2 + 4*a*c] - 2*c*x^2))/(2*Sqrt[2]*Sqrt[a]*Sqrt[c]*Sqrt[a + b*x^2 - c*x^4])]/(2*Sqrt[2]*Sqrt[a]*Sqrt[c]*d)`

### 3.36.3.1 Defintions of rubi rules used

```
rule 2518 Int[Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4]/((d_) + (e_.)*(x_)^4), x_Symbol]
:= With[{q = Sqrt[b^2 - 4*a*c]}, Simp[(-a)*(Sqrt[b + q]/(2*Sqrt[2]*Rt[(-a)*c, 2]*d))
*ArcTan[Sqrt[b + q]*x*((b - q + 2*c*x^2)/(2*Sqrt[2]*Rt[(-a)*c, 2]*Sqrt[a + b*x^2 + c*x^4]))], x]
+ Simp[a*(Sqrt[-b + q]/(2*Sqrt[2]*Rt[(-a)*c, 2]*d))*ArcTanh[Sqrt[-b + q]*x*((b + q + 2*c*x^2)/(2*Sqrt[2]*Rt[(-a)*c, 2]*Sqrt[a + b*x^2 + c*x^4]))], x]]
/; FreeQ[{a, b, c, d, e}, x] && EqQ[c*d + a*e, 0] && NegQ[a*c]
```

### 3.36.4 Maple [A] (verified)

Time = 2.49 (sec) , antiderivative size = 351, normalized size of antiderivative = 1.47

| method         | result  |
|----------------|---|
| pseudoelliptic | $\frac{\sqrt{2\sqrt{4ac+b^2}+2b} (b-\sqrt{4ac+b^2}) \left( \ln \left( \frac{-cx^4 + \sqrt{-cx^4 + bx^2 + a} \sqrt{2\sqrt{4ac+b^2} + 2bx + \sqrt{4ac+b^2} x^2 + bx^2 + a}}{x^2} \right) - \ln \left( \frac{cx^4 + \sqrt{-cx^4 + bx^2 + a}}{x^2} \right) \right)}{\dots}$   |
| elliptic       | $\left( \frac{\sqrt{b + \sqrt{4ac + b^2}} b \ln \left( \frac{-cx^4 + bx^2 + a}{x^2} - \frac{\sqrt{-cx^4 + bx^2 + a} \sqrt{2} \sqrt{b + \sqrt{4ac + b^2}}}{x} + \sqrt{4ac + b^2} \right)}{16dac} \right) + \frac{b^2 \arctan \left( \frac{2\sqrt{-cx^4 + bx^2 + a} \sqrt{2} - 2\sqrt{b + \sqrt{4ac + b^2}}}{2\sqrt{-b + \sqrt{4ac + b^2}}} \right)}{8dac\sqrt{-b + \sqrt{4ac + b^2}}}$ |
| default        | $\left( \frac{\sqrt{b + \sqrt{4ac + b^2}} b \ln \left( \frac{-cx^4 + bx^2 + a}{x^2} - \frac{\sqrt{-cx^4 + bx^2 + a} \sqrt{2} \sqrt{b + \sqrt{4ac + b^2}}}{x} + \sqrt{4ac + b^2} \right)}{16ac} \right) + \frac{(b + \sqrt{4ac + b^2}) b \arctan \left( \frac{2\sqrt{-cx^4 + bx^2 + a} \sqrt{2}}{2\sqrt{-b + \sqrt{4ac + b^2}}} \right)}{8ac\sqrt{-b + \sqrt{4ac + b^2}}}$             |

```
input int((-c*x^4+b*x^2+a)^(1/2)/(c*d*x^4+a*d),x,method=_RETURNVERBOSE)
```

```
output -1/32/(2*(4*a*c+b^2)^(1/2)-2*b)^(1/2)*((2*(4*a*c+b^2)^(1/2)+2*b)^(1/2)*(b-
(4*a*c+b^2)^(1/2))*(ln((-c*x^4+(-c*x^4+b*x^2+a)^(1/2)*(2*(4*a*c+b^2)^(1/2)
+2*b)^(1/2)*x+(4*a*c+b^2)^(1/2)*x^2+b*x^2+a)/x^2)-ln((c*x^4+(-c*x^4+b*x^2+
a)^(1/2)*(2*(4*a*c+b^2)^(1/2)+2*b)^(1/2)*x-(4*a*c+b^2)^(1/2)*x^2-b*x^2-a)/
x^2))*(2*(4*a*c+b^2)^(1/2)-2*b)^(1/2)-16*a*c*(arctan(((2*(4*a*c+b^2)^(1/2)
+2*b)^(1/2)*x-2*(-c*x^4+b*x^2+a)^(1/2))/x/(2*(4*a*c+b^2)^(1/2)-2*b)^(1/2))
-arctan(((2*(4*a*c+b^2)^(1/2)+2*b)^(1/2)*x+2*(-c*x^4+b*x^2+a)^(1/2))/x/(2*
(4*a*c+b^2)^(1/2)-2*b)^(1/2))))/d/a/c
```



**3.36.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 669 vs.  $2(187) = 374$ .

Time = 1.68 (sec) , antiderivative size = 669, normalized size of antiderivative = 2.80

$$\int \frac{\sqrt{a+bx^2-cx^4}}{ad+cdx^4} dx =$$

$$-\frac{1}{8} \sqrt{\frac{2acd^2\sqrt{-\frac{1}{acd^4}}-b}{acd^2}} \log \left( -\frac{\sqrt{-cx^4+bx^2+a}ad^2\sqrt{-\frac{1}{acd^4}}+\sqrt{-cx^4+bx^2+ax^2}+(acd^3x^3\sqrt{-\frac{1}{acd^4}})}{cx^4+a} \right)$$

$$+\frac{1}{8} \sqrt{\frac{2acd^2\sqrt{-\frac{1}{acd^4}}-b}{acd^2}} \log \left( -\frac{\sqrt{-cx^4+bx^2+a}ad^2\sqrt{-\frac{1}{acd^4}}+\sqrt{-cx^4+bx^2+ax^2}-(acd^3x^3\sqrt{-\frac{1}{acd^4}})}{cx^4+a} \right)$$

$$-\frac{1}{8} \sqrt{-\frac{2acd^2\sqrt{-\frac{1}{acd^4}}+b}{acd^2}} \log \left( \frac{\sqrt{-cx^4+bx^2+a}ad^2\sqrt{-\frac{1}{acd^4}}-\sqrt{-cx^4+bx^2+ax^2}+(acd^3x^3\sqrt{-\frac{1}{acd^4}})}{cx^4+a} \right)$$

$$+\frac{1}{8} \sqrt{-\frac{2acd^2\sqrt{-\frac{1}{acd^4}}+b}{acd^2}} \log \left( \frac{\sqrt{-cx^4+bx^2+a}ad^2\sqrt{-\frac{1}{acd^4}}-\sqrt{-cx^4+bx^2+ax^2}-(acd^3x^3\sqrt{-\frac{1}{acd^4}})}{cx^4+a} \right)$$

input `integrate((-c*x^4+b*x^2+a)^(1/2)/(c*d*x^4+a*d),x, algorithm="fracas")`

output 
$$\begin{aligned} & -1/8*\sqrt{((2*a*c*d^2*\sqrt{-1/(a*c*d^4)} - b)/(a*c*d^2))*\log(-(\sqrt{-c*x^4 + b*x^2 + a})*a*d^2*\sqrt{-1/(a*c*d^4)} + \sqrt{-c*x^4 + b*x^2 + a})*x^2 + (a*c*d^3*x^3*\sqrt{-1/(a*c*d^4)} - a*d*x)*\sqrt{((2*a*c*d^2*\sqrt{-1/(a*c*d^4)} - b)/(a*c*d^2)))/(c*x^4 + a)} + 1/8*\sqrt{((2*a*c*d^2*\sqrt{-1/(a*c*d^4)} - b)/(a*c*d^2))*\log(-(\sqrt{-c*x^4 + b*x^2 + a})*a*d^2*\sqrt{-1/(a*c*d^4)} + \sqrt{-c*x^4 + b*x^2 + a})*x^2 - (a*c*d^3*x^3*\sqrt{-1/(a*c*d^4)} - a*d*x)*\sqrt{((2*a*c*d^2*\sqrt{-1/(a*c*d^4)} - b)/(a*c*d^2)))/(c*x^4 + a)} - 1/8*\sqrt{-(2*a*c*d^2*\sqrt{-1/(a*c*d^4)} + b)/(a*c*d^2))*\log((\sqrt{-c*x^4 + b*x^2 + a})*a*d^2*\sqrt{-1/(a*c*d^4)} - \sqrt{-c*x^4 + b*x^2 + a})*x^2 + (a*c*d^3*x^3*\sqrt{-1/(a*c*d^4)} + a*d*x)*\sqrt{-(2*a*c*d^2*\sqrt{-1/(a*c*d^4)} + b)/(a*c*d^2)))/(c*x^4 + a)} + 1/8*\sqrt{-(2*a*c*d^2*\sqrt{-1/(a*c*d^4)} + b)/(a*c*d^2))*\log((\sqrt{-c*x^4 + b*x^2 + a})*a*d^2*\sqrt{-1/(a*c*d^4)} - \sqrt{-c*x^4 + b*x^2 + a})*x^2 - (a*c*d^3*x^3*\sqrt{-1/(a*c*d^4)} + a*d*x)*\sqrt{-(2*a*c*d^2*\sqrt{-1/(a*c*d^4)} + b)/(a*c*d^2)))/(c*x^4 + a)} \end{aligned}$$

### 3.36.6 Sympy [F]

$$\int \frac{\sqrt{a + bx^2 - cx^4}}{ad + cdx^4} dx = \int \frac{\sqrt{a+bx^2-cx^4}}{a+cx^4} \frac{dx}{d}$$

input `integrate((-c*x**4+b*x**2+a)**(1/2)/(c*d*x**4+a*d), x)`

output `Integral(sqrt(a + b*x**2 - c*x**4)/(a + c*x**4), x)/d`

### 3.36.7 Maxima [F]

$$\int \frac{\sqrt{a + bx^2 - cx^4}}{ad + cdx^4} dx = \int \frac{\sqrt{-cx^4 + bx^2 + a}}{cdx^4 + ad} dx$$

input `integrate((-c*x^4+b*x^2+a)^(1/2)/(c*d*x^4+a*d), x, algorithm="maxima")`

output `integrate(sqrt(-c*x^4 + b*x^2 + a)/(c*d*x^4 + a*d), x)`

**3.36.8 Giac [F]**

$$\int \frac{\sqrt{a + bx^2 - cx^4}}{ad + cdx^4} dx = \int \frac{\sqrt{-cx^4 + bx^2 + a}}{cdx^4 + ad} dx$$

input `integrate((-c*x^4+b*x^2+a)^(1/2)/(c*d*x^4+a*d),x, algorithm="giac")`

output `integrate(sqrt(-c*x^4 + b*x^2 + a)/(c*d*x^4 + a*d), x)`

**3.36.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{a + bx^2 - cx^4}}{ad + cdx^4} dx = \int \frac{\sqrt{-cx^4 + bx^2 + a}}{cdx^4 + ad} dx$$

input `int((a + b*x^2 - c*x^4)^(1/2)/(a*d + c*d*x^4),x)`

output `int((a + b*x^2 - c*x^4)^(1/2)/(a*d + c*d*x^4), x)`

### 3.37 $\int x\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4} dx$

|        |   |     |
|--------|---|-----|
| 3.37.1 | Optimal result                            | 299 |
| 3.37.2 | Mathematica [A] (verified)                | 300 |
| 3.37.3 | Rubi [A] (verified)                       | 300 |
| 3.37.4 | Maple [A] (verified)                      | 303 |
| 3.37.5 | Fricas [A] (verification not implemented) | 304 |
| 3.37.6 | Sympy [F]                                 | 304 |
| 3.37.7 | Maxima [F(-2)]                            | 305 |
| 3.37.8 | Giac [A] (verification not implemented)   | 305 |
| 3.37.9 | Mupad [F(-1)]                             | 306 |

#### 3.37.1 Optimal result

Integrand size = 38, antiderivative size = 309

$$\int x\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4} dx$$

$$= \frac{e(12bcd - 16ad^2 - 7be^2)(e + 2dx)\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{128d^4(a+bx^2)}$$

$$+ \frac{bx^2(c+ex+dx^2)^{3/2}\sqrt{a^2+2abx^2+b^2x^4}}{5d(a+bx^2)}$$

$$- \frac{(32bcd - 80ad^2 - 35be^2 + 42bdex)(c+ex+dx^2)^{3/2}\sqrt{a^2+2abx^2+b^2x^4}}{240d^3(a+bx^2)}$$

$$+ \frac{e(4cd - e^2)(12bcd - 16ad^2 - 7be^2)\sqrt{a^2+2abx^2+b^2x^4}\operatorname{arctanh}\left(\frac{e+2dx}{2\sqrt{d}\sqrt{c+ex+dx^2}}\right)}{256d^{9/2}(a+bx^2)}$$

output

```
1/5*b*x^2*(d*x^2+e*x+c)^(3/2)*((b*x^2+a)^2)^(1/2)/d/(b*x^2+a)-1/240*(42*b*
d*e*x-80*a*d^2+32*b*c*d-35*b*e^2)*(d*x^2+e*x+c)^(3/2)*((b*x^2+a)^2)^(1/2)/
d^3/(b*x^2+a)+1/256*e*(4*c*d-e^2)*(-16*a*d^2+12*b*c*d-7*b*e^2)*arctanh(1/2
*(2*d*x+e)/d^(1/2)/(d*x^2+e*x+c)^(1/2))*((b*x^2+a)^2)^(1/2)/d^(9/2)/(b*x^2
+a)+1/128*e*(-16*a*d^2+12*b*c*d-7*b*e^2)*(2*d*x+e)*(d*x^2+e*x+c)^(1/2)*((b
*x^2+a)^2)^(1/2)/d^4/(b*x^2+a)
```

### 3.37.2 Mathematica [A] (verified)

Time = 0.69 (sec) , antiderivative size = 212, normalized size of antiderivative = 0.69

$$\int x\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4} dx$$

$$= \frac{\sqrt{(a+bx^2)^2} \left( 2\sqrt{d}\sqrt{c+x(e+dx)}(80ad^2(8cd-3e^2+2dex+8d^2x^2) + b(-256c^2d^2-105e^4+70de^3x - \dots \right)}{3840d^{9/2}(a+bx^2)}$$

input `Integrate[x*Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4],x]`

output `(Sqrt[(a + b*x^2)^2]*(2*Sqrt[d]*Sqrt[c + x*(e + d*x)]*(80*a*d^2*(8*c*d - 3*e^2 + 2*d*e*x + 8*d^2*x^2) + b*(-256*c^2*d^2 - 105*e^4 + 70*d*e^3*x - 56*d^2*e^2*x^2 + 48*d^3*e*x^3 + 384*d^4*x^4 + 4*c*d*(115*e^2 - 58*d*e*x + 32*d^2*x^2))) - 15*e*(-4*c*d + e^2)*(-12*b*c*d + 16*a*d^2 + 7*b*e^2)*Log[e + 2*d*x - 2*Sqrt[d]*Sqrt[c + x*(e + d*x)]])/(3840*d^(9/2)*(a + b*x^2))`

### 3.37.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 215, normalized size of antiderivative = 0.70, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {1384, 27, 2184, 27, 1225, 1087, 1092, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x\sqrt{a^2+2abx^2+b^2x^4}\sqrt{c+dx^2+ex} dx$$

$$\downarrow \text{1384}$$

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \int bx(bx^2+a)\sqrt{dx^2+ex+cdx}}{b(a+bx^2)}$$

$$\downarrow \text{27}$$

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \int x(bx^2+a)\sqrt{dx^2+ex+cdx}}{a+bx^2}$$

$$\downarrow \text{2184}$$

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \left( \frac{\int -\frac{1}{2}x(4bc-10ad+7bex)\sqrt{dx^2+ex+cdx}}{5d} + \frac{bx^2(c+dx^2+ex)^{3/2}}{5d} \right)}{a+bx^2}$$

$$\begin{array}{c}
 \downarrow 27 \\
 \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{bx^2(c+dx^2+ex)^{3/2}}{5d} - \frac{\int x(2(2bc-5ad)+7be)x\sqrt{dx^2+ex+cdx}}{10d} \right)}{a + bx^2} \\
 \downarrow 1225 \\
 \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{bx^2(c+dx^2+ex)^{3/2}}{5d} - \frac{(c+dx^2+ex)^{3/2} (16d(2bc-5ad)+42bdex-35be^2)}{24d^2} - \frac{5e(-16ad^2+12bcd-7be^2) \int \sqrt{dx^2+ex+cdx}}{16d^2} \right)}{a + bx^2} \\
 \downarrow 1087 \\
 \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{bx^2(c+dx^2+ex)^{3/2}}{5d} - \frac{(c+dx^2+ex)^{3/2} (16d(2bc-5ad)+42bdex-35be^2)}{24d^2} - \frac{5e(-16ad^2+12bcd-7be^2)}{10d} \left( \frac{(4cd-e^2) \int \frac{1}{\sqrt{dx^2+ex+cdx}}}{8d} \right) \right)}{a + bx^2} \\
 \downarrow 1092 \\
 \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{bx^2(c+dx^2+ex)^{3/2}}{5d} - \frac{(c+dx^2+ex)^{3/2} (16d(2bc-5ad)+42bdex-35be^2)}{24d^2} - \frac{5e(-16ad^2+12bcd-7be^2)}{10d} \left( \frac{(4cd-e^2) \int \frac{1}{4d - \frac{(c+2cd+dx^2+ex)}{4d}}}{8d} \right) \right)}{a + bx^2} \\
 \downarrow 219 \\
 \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{bx^2(c+dx^2+ex)^{3/2}}{5d} - \frac{(c+dx^2+ex)^{3/2} (16d(2bc-5ad)+42bdex-35be^2)}{24d^2} - \frac{5e(-16ad^2+12bcd-7be^2)}{10d} \left( \frac{(4cd-e^2) \operatorname{arctanh}\left(\frac{1}{8d^3/2}\right)}{16d^2} \right) \right)}{a + bx^2}
 \end{array}$$

input `Int[x*Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4], x]`

output  $(\sqrt{a^2 + 2abx + b^2x^2} * ((bx^2(c + ex + dx^2)^{3/2}) / (5d) - ((16d(2bc - 5ad) - 35b^2e + 42bde)(c + ex + dx^2)^{3/2}) / (24d^2) - (5e(12bcd - 16ad^2 - 7b^2e)(e + 2dx)\sqrt{c + ex + dx^2}) / (4d) + ((4cd - e^2)\text{ArcTanh}[(e + 2dx) / (2\sqrt{d}\sqrt{c + ex + dx^2})]) / (8d^{3/2}))) / (16d^2) / (10d) / (a + bx^2)$

### 3.37.3.1 Defintions of rubi rules used

rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$

rule 219  $\text{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 1087  $\text{Int}[(a_*) + (b_*)(x_) + (c_*)(x_)^2)^{p_}, x\_Symbol] \rightarrow \text{Simp}[(b + 2cx) * ((a + bx + cx^2)^p / (2c(2p + 1))), x] - \text{Simp}[p * ((b^2 - 4ac) / (2c(2p + 1))) \text{Int}[(a + bx + cx^2)^{p-1}, x], x] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ (\text{IntegerQ}[4p] \ || \ \text{IntegerQ}[3p])$

rule 1092  $\text{Int}[1/\sqrt{(a_*) + (b_*)(x_) + (c_*)(x_)^2}, x\_Symbol] \rightarrow \text{Simp}[2 \text{ Subst}[\text{Int}[1/(4c - x^2), x], x, (b + 2cx)/\sqrt{a + bx + cx^2}], x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1225  $\text{Int}[(d_*) + (e_*)(x_))*((f_*) + (g_*)(x_))*((a_*) + (b_*)(x_) + (c_*)(x_)^2)^{p_}, x\_Symbol] \rightarrow \text{Simp}[(-b*eg*(p + 2) - c*(ef + dg)*(2p + 3) - 2c*eg*(p + 1)*x) * ((a + bx + cx^2)^{p+1} / (2c^2*(p + 1)*(2p + 3))), x] + \text{Simp}[(b^2*eg*(p + 2) - 2a*c*eg + c*(2c*d*f - b*(ef + dg))*(2p + 3)) / (2c^2*(2p + 3)) \text{Int}[(a + bx + cx^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, p\}, x] \ \&\& \ !\text{LeQ}[p, -1]$

rule 1384  $\text{Int}[(u_*)((a_*) + (c_*)(x_)^{n2_}) + (b_*)(x_)^{n_})^{p_}, x\_Symbol] \rightarrow \text{Simp}[(a + bx^n + cx^{2n})^{\text{FracPart}[p]} / (c^{\text{IntPart}[p]} * (b/2 + cx^n)^{2*\text{FracPart}[p]}) \text{Int}[u * (b/2 + cx^n)^{2p}, x], x] /; \text{FreeQ}[\{a, b, c, n, p\}, x] \ \&\& \ \text{EqQ}[n2, 2*n] \ \&\& \ \text{EqQ}[b^2 - 4ac, 0] \ \&\& \ \text{IntegerQ}[p - 1/2] \ \&\& \ \text{NeQ}[u, x^{(n - 1)}] \ \&\& \ \text{NeQ}[u, x^{(2*n - 1)}] \ \&\& \ !(\text{EqQ}[p, 1/2] \ \&\& \ \text{EqQ}[u, x^{(-2*n - 1)}])$

```
rule 2184 Int[(Pq_)*((d_.) + (e_.)*(x_))^(m_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p
_), x_Symbol] := With[{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, S
imp[f*(d + e*x)^(m + q - 1)*((a + b*x + c*x^2)^(p + 1)/(c*e^(q - 1)*(m + q
+ 2*p + 1))), x] + Simp[1/(c*e^q*(m + q + 2*p + 1)) Int[(d + e*x)^m*(a +
b*x + c*x^2)^p*ExpandToSum[c*e^q*(m + q + 2*p + 1)*Pq - c*f*(m + q + 2*p +
1)*(d + e*x)^q - f*(d + e*x)^(q - 2)*(b*d*e*(p + 1) + a*e^2*(m + q - 1) - c
*d^2*(m + q + 2*p + 1) - e*(2*c*d - b*e)*(m + q + p)*x), x], x] /; GtQ[
q, 1] && NeQ[m + q + 2*p + 1, 0] /; FreeQ[{a, b, c, d, e, m, p}, x] && Pol
yQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !(IGt
Q[m, 0] && RationalQ[a, b, c, d, e] && (IntegerQ[p] || ILtQ[p + 1/2, 0]))
```

### 3.37.4 Maple [A] (verified)

Time = 0.73 (sec) , antiderivative size = 245, normalized size of antiderivative = 0.79

| method  | result  |
|---------|---|
| risch   | $\frac{(384b x^4 d^4 + 48e x^3 b d^3 + 640a d^4 x^2 + 128bc d^3 x^2 - 56b d^2 e^2 x^2 + 160a d^3 e x - 232bc d^2 e x + 70bd e^3 x + 640ac d^3 - 240e^2 d^2 a - 256b c^2 d^2 + 460c^2 d^2 a^2)}{1920d^4(bx^2+a)}$  |
| default | $\frac{\sqrt{(bx^2+a)^2} \left( 768d^{\frac{9}{2}}(dx^2+ex+c)^{\frac{3}{2}}bx^2 - 672d^{\frac{7}{2}}(dx^2+ex+c)^{\frac{3}{2}}bcx + 1280d^{\frac{9}{2}}(dx^2+ex+c)^{\frac{3}{2}}a - 512d^{\frac{7}{2}}(dx^2+ex+c)^{\frac{3}{2}}bc + 560d^{\frac{5}{2}}(dx^2+ex+c)^{\frac{3}{2}} \right)}{\sqrt{(bx^2+a)^2}}$ |

```
input int(x*(d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/1920*(384*b*d^4*x^4+48*b*d^3*e*x^3+640*a*d^4*x^2+128*b*c*d^3*x^2-56*b*d^
2*e^2*x^2+160*a*d^3*e*x-232*b*c*d^2*e*x+70*b*d*e^3*x+640*a*c*d^3-240*a*d^2
*e^2-256*b*c^2*d^2+460*b*c*d*e^2-105*b*e^4)*(d*x^2+e*x+c)^(1/2)/d^4*((b*x^
2+a)^2)^(1/2)/(b*x^2+a)-1/256*e*(64*a*c*d^3-16*a*d^2*e^2-48*b*c^2*d^2+40*b
*c*d*e^2-7*b*e^4)/d^(9/2)*ln((1/2*e+d*x)/d^(1/2)+(d*x^2+e*x+c)^(1/2))*((b
x^2+a)^2)^(1/2)/(b*x^2+a)
```

---

3.37.  $\int x\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}dx$



**3.37.5 Fracas [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 469, normalized size of antiderivative = 1.52

$$\int x\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4} dx$$

$$= \left[ \frac{15(7be^5 - 8(5bcd - 2ad^2)e^3 + 16(3bc^2d^2 - 4acd^3)e)\sqrt{d} \log\left(8d^2x^2 + 8dex + 4\sqrt{dx^2+ex+c}(2dx + e)\sqrt{d} + 4c*d + e^2\right) + 4*(384*b*d^5*x^4 + 48*b*d^4*e*x^3 - 256*b*c^2*d^3 + 640*a*c*d^4 - 105*b*d*e^4 + 20*(23*b*c*d^2 - 12*a*d^3)*e^2 + 8*(16*b*c*d^4 + 80*a*d^5 - 7*b*d^3*e^2)*x^2 + 2*(35*b*d^2*e^3 - 4*(29*b*c*d^3 - 20*a*d^4)*e)*x)\sqrt{d^2x^2 + dex + cd}}{15(7be^5 - 8(5bcd - 2ad^2)e^3 + 16(3bc^2d^2 - 4acd^3)e)\sqrt{-d} \arctan\left(\frac{\sqrt{dx^2+ex+c}(2dx+e)\sqrt{-d}}{2(d^2x^2+dex+cd)}\right) - 2(384bd^5x^4 + 48bd^4ex^3 - 256b^2c^2d^3 + 640acd^4 - 105bde^4 + 20(23b^2cd^2 - 12ad^3)e^2 + 8(16b^2cd^4 + 80ad^5 - 7bd^3e^2)x^2 + 2(35bd^2e^3 - 4(29b^2cd^3 - 20ad^4)e)x)\sqrt{d^2x^2 + dex + c})/d^5} \right]$$

```
input integrate(x*(d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x, algorithm="fracas")
```

```
output [1/7680*(15*(7*b*e^5 - 8*(5*b*c*d - 2*a*d^2)*e^3 + 16*(3*b*c^2*d^2 - 4*a*c*d^3)*e)*sqrt(d)*log(8*d^2*x^2 + 8*d*e*x + 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 4*(384*b*d^5*x^4 + 48*b*d^4*e*x^3 - 256*b*c^2*d^3 + 640*a*c*d^4 - 105*b*d*e^4 + 20*(23*b*c*d^2 - 12*a*d^3)*e^2 + 8*(16*b*c*d^4 + 80*a*d^5 - 7*b*d^3*e^2)*x^2 + 2*(35*b*d^2*e^3 - 4*(29*b*c*d^3 - 20*a*d^4)*e)*x)*sqrt(d*x^2 + e*x + c))/d^5, -1/3840*(15*(7*b*e^5 - 8*(5*b*c*d - 2*a*d^2)*e^3 + 16*(3*b*c^2*d^2 - 4*a*c*d^3)*e)*sqrt(-d)*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) - 2*(384*b*d^5*x^4 + 48*b*d^4*e*x^3 - 256*b*c^2*d^3 + 640*a*c*d^4 - 105*b*d*e^4 + 20*(23*b*c*d^2 - 12*a*d^3)*e^2 + 8*(16*b*c*d^4 + 80*a*d^5 - 7*b*d^3*e^2)*x^2 + 2*(35*b*d^2*e^3 - 4*(29*b*c*d^3 - 20*a*d^4)*e)*x)*sqrt(d*x^2 + e*x + c))/d^5]
```

**3.37.6 Sympy [F]**

$$\int x\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4} dx = \int x\sqrt{c+dx^2+ex}\sqrt{(a+bx^2)^2} dx$$

```
input integrate(x*(d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2),x)
```

```
output Integral(x*sqrt(c + d*x**2 + e*x)*sqrt((a + b*x**2)**2), x)
```

**3.37.7 Maxima [F(-2)]**

Exception generated.

$$\int x\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}dx = \text{Exception raised: ValueError}$$

```
input integrate(x*(d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**3.37.8 Giac [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 358, normalized size of antiderivative = 1.16

$$\int x\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}dx$$

$$= \frac{1}{1920} \sqrt{dx^2+ex+c} \left( 2 \left( 4 \left( 6 \left( 8bx\operatorname{sgn}(bx^2+a) + \frac{b\operatorname{sgn}(bx^2+a)}{d} \right) x + \frac{16bcd^3\operatorname{sgn}(bx^2+a) + 80ad^4\operatorname{sgn}(bx^2+a)}{d^4} \right) \right. \right.$$

$$\left. \left. - \frac{(48bc^2d^2\operatorname{esgn}(bx^2+a) - 64acd^3\operatorname{esgn}(bx^2+a) - 40bcde^3\operatorname{sgn}(bx^2+a) + 16ad^2e^3\operatorname{sgn}(bx^2+a) + 7be^5\operatorname{sgn}(bx^2+a))}{256d^{\frac{9}{2}}} \right) \right)$$

```
input integrate(x*(d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x, algorithm="giac")
```

```
output 1/1920*sqrt(d*x^2 + e*x + c)*(2*(4*(6*(8*b*x*sgn(b*x^2 + a) + b*e*sgn(b*x^
2 + a)/d)*x + (16*b*c*d^3*sgn(b*x^2 + a) + 80*a*d^4*sgn(b*x^2 + a) - 7*b*d
^2*e^2*sgn(b*x^2 + a))/d^4)*x - (116*b*c*d^2*e*sgn(b*x^2 + a) - 80*a*d^3*e
*sgn(b*x^2 + a) - 35*b*d*e^3*sgn(b*x^2 + a))/d^4)*x - (256*b*c^2*d^2*sgn(b
*x^2 + a) - 640*a*c*d^3*sgn(b*x^2 + a) - 460*b*c*d*e^2*sgn(b*x^2 + a) + 24
0*a*d^2*e^2*sgn(b*x^2 + a) + 105*b*e^4*sgn(b*x^2 + a))/d^4) - 1/256*(48*b*
c^2*d^2*e*sgn(b*x^2 + a) - 64*a*c*d^3*e*sgn(b*x^2 + a) - 40*b*c*d*e^3*sgn(
b*x^2 + a) + 16*a*d^2*e^3*sgn(b*x^2 + a) + 7*b*e^5*sgn(b*x^2 + a))*log(abs
(2*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))*sqrt(d) + e))/d^(9/2)
```

**3.37.9 Mupad [F(-1)]**

Timed out.

$$\int x\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}dx = \int x\sqrt{(bx^2+a)^2}\sqrt{dx^2+ex+c}dx$$

input `int(x*((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2),x)`output `int(x*((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2), x)`

### 3.38 $\int \sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4} dx$

|        |   |     |
|--------|---|-----|
| 3.38.1 | Optimal result . . . . .                            | 307 |
| 3.38.2 | Mathematica [A] (verified) . . . . .                | 308 |
| 3.38.3 | Rubi [A] (verified) . . . . .                       | 308 |
| 3.38.4 | Maple [A] (verified) . . . . .                      | 311 |
| 3.38.5 | Fricas [A] (verification not implemented) . . . . . | 311 |
| 3.38.6 | Sympy [F] . . . . .                                 | 312 |
| 3.38.7 | Maxima [F(-2)] . . . . .                            | 312 |
| 3.38.8 | Giac [A] (verification not implemented) . . . . .   | 313 |
| 3.38.9 | Mupad [F(-1)] . . . . .                             | 313 |

#### 3.38.1 Optimal result

Integrand size = 37, antiderivative size = 283

$$\int \sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4} dx$$

$$= -\frac{(4bcd - 16ad^2 - 5be^2)(e + 2dx)\sqrt{c + ex + dx^2}\sqrt{a^2 + 2abx^2 + b^2x^4}}{64d^3(a + bx^2)}$$

$$- \frac{5be(c + ex + dx^2)^{3/2}\sqrt{a^2 + 2abx^2 + b^2x^4}}{24d^2(a + bx^2)} + \frac{bx(c + ex + dx^2)^{3/2}\sqrt{a^2 + 2abx^2 + b^2x^4}}{4d(a + bx^2)}$$

$$- \frac{(4cd - e^2)(4bcd - 16ad^2 - 5be^2)\sqrt{a^2 + 2abx^2 + b^2x^4}\operatorname{arctanh}\left(\frac{e+2dx}{2\sqrt{d}\sqrt{c+ex+dx^2}}\right)}{128d^{7/2}(a + bx^2)}$$

output

```
-5/24*b*e*(d*x^2+e*x+c)^(3/2)*((b*x^2+a)^2)^(1/2)/d^2/(b*x^2+a)+1/4*b*x*(d*x^2+e*x+c)^(3/2)*((b*x^2+a)^2)^(1/2)/d/(b*x^2+a)-1/128*(4*c*d-e^2)*(-16*a*d^2+4*b*c*d-5*b*e^2)*arctanh(1/2*(2*d*x+e)/d^(1/2)/(d*x^2+e*x+c)^(1/2))*((b*x^2+a)^2)^(1/2)/d^(7/2)/(b*x^2+a)-1/64*(-16*a*d^2+4*b*c*d-5*b*e^2)*(2*d*x+e)*(d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/d^3/(b*x^2+a)
```

### 3.38.2 Mathematica [A] (verified)

Time = 1.19 (sec) , antiderivative size = 214, normalized size of antiderivative = 0.76

$$\int \sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4} dx$$

$$= \frac{\sqrt{(a+bx^2)^2}\left(\sqrt{d}\sqrt{c+x(e+dx)}(48ad^2(e+2dx)+b(15e^3-10de^2x+8d^2ex^2+48d^3x^3+4cd(-13e+6d^2x)))\right)}{192d^7}$$

input `Integrate[Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4], x]`

output `(Sqrt[(a + b*x^2)^2]*(Sqrt[d]*Sqrt[c + x*(e + d*x)]*(48*a*d^2*(e + 2*d*x) + b*(15*e^3 - 10*d*e^2*x + 8*d^2*e*x^2 + 48*d^3*x^3 + 4*c*d*(-13*e + 6*d*x))) + 3*(16*b*c^2*d^2 + 16*a*d^2*e^2 + 5*b*e^4)*ArcTanh[(Sqrt[d]*x)/(Sqrt[c] - Sqrt[c + x*(e + d*x)])] + 24*c*d*(8*a*d^2 + 3*b*e^2)*ArcTanh[(Sqrt[d]*x)/(-Sqrt[c] + Sqrt[c + x*(e + d*x)])])/(192*d^(7/2)*(a + b*x^2))`

### 3.38.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 189, normalized size of antiderivative = 0.67, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.216$ , Rules used = {1384, 27, 2192, 27, 1160, 1087, 1092, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a^2+2abx^2+b^2x^4}\sqrt{c+dx^2+ex} dx$$

$$\downarrow \text{1384}$$

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \int b(bx^2+a)\sqrt{dx^2+ex+cdx}}{b(a+bx^2)}$$

$$\downarrow \text{27}$$

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \int (bx^2+a)\sqrt{dx^2+ex+cdx}}{a+bx^2}$$

$$\downarrow \text{2192}$$

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \left( \frac{\int -\frac{1}{2}(2bc-8ad+5bex)\sqrt{dx^2+ex+cdx}}{4d} + \frac{bx(c+dx^2+ex)^{3/2}}{4d} \right)}{a+bx^2}$$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{bx(c+dx^2+ex)^{3/2}}{4d} - \frac{\int (2(bc-4ad)+5bex)\sqrt{dx^2+ex+cdx}}{8d} \right)}{a + bx^2} \\
 & \downarrow 1160 \\
 & \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{bx(c+dx^2+ex)^{3/2}}{4d} - \frac{(4d(bc-4ad)-5be^2) \int \sqrt{dx^2+ex+cdx}}{2d \cdot 8d} + \frac{5be(c+dx^2+ex)^{3/2}}{3d} \right)}{a + bx^2} \\
 & \downarrow 1087 \\
 & \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{bx(c+dx^2+ex)^{3/2}}{4d} - \frac{(4d(bc-4ad)-5be^2) \left( \frac{(4cd-e^2) \int \frac{1}{\sqrt{dx^2+ex+c}} dx}{8d} + \frac{(2dx+e)\sqrt{c+dx^2+ex}}{4d} \right)}{2d \cdot 8d} + \frac{5be(c+dx^2+ex)^{3/2}}{3d} \right)}{a + bx^2} \\
 & \downarrow 1092 \\
 & \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{bx(c+dx^2+ex)^{3/2}}{4d} - \frac{(4d(bc-4ad)-5be^2) \left( \frac{(4cd-e^2) \int \frac{1}{4d - \frac{(e+2dx)^2}{dx^2+ex+c}} d \frac{e+2dx}{\sqrt{dx^2+ex+c}}}{8d} + \frac{(2dx+e)\sqrt{c+dx^2+ex}}{4d} \right)}{2d \cdot 8d} + \frac{5be(c+dx^2+ex)^{3/2}}{3d} \right)}{a + bx^2} \\
 & \downarrow 219 \\
 & \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{bx(c+dx^2+ex)^{3/2}}{4d} - \frac{(4d(bc-4ad)-5be^2) \left( \frac{(4cd-e^2) \operatorname{arctanh}\left(\frac{2dx+e}{2\sqrt{d}\sqrt{c+dx^2+ex}}\right)}{8d^{3/2}} + \frac{(2dx+e)\sqrt{c+dx^2+ex}}{4d} \right)}{2d \cdot 8d} + \frac{5be(c+dx^2+ex)^{3/2}}{3d} \right)}{a + bx^2}
 \end{aligned}$$

input `Int[Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4], x]`

output  $(\sqrt{a^2 + 2abx + b^2x^2} * ((bx(c + ex + dx^2)^{3/2}) / (4d) - ((5b^2e(c + ex + dx^2)^{3/2}) / (3d) + ((4d(bc - 4ad) - 5b^2e^2) * ((e + 2dx) * \sqrt{c + ex + dx^2}) / (4d) + ((4cd - e^2) * \text{ArcTanh}[(e + 2dx) / (2\sqrt{d} * \sqrt{c + ex + dx^2})]) / (8d^{3/2}))) / (2d)) / (8d)) / (a + bx^2)$

### 3.38.3.1 Defintions of rubi rules used

rule 27  $\text{Int}[(a_*)(F_x), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_*)(G_x) /; \text{FreeQ}[b, x]]$

rule 219  $\text{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] * \text{Rt}[-b, 2])) * \text{ArcTanh}[\text{Rt}[-b, 2] * (x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 1087  $\text{Int}[(a_*) + (b_*)(x_) + (c_*)(x_)^2)^{p_}, x\_Symbol] \rightarrow \text{Simp}[(b + 2cx) * ((a + bx + cx^2)^p / (2c(2p + 1))), x] - \text{Simp}[p * (b^2 - 4ac) / (2c(2p + 1)) \text{ Int}[(a + bx + cx^2)^{p-1}, x], x] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ (\text{IntegerQ}[4p] \ || \ \text{IntegerQ}[3p])$

rule 1092  $\text{Int}[1/\sqrt{(a_*) + (b_*)(x_) + (c_*)(x_)^2}], x\_Symbol] \rightarrow \text{Simp}[2 \text{ Subst}[\text{Int}[1/(4c - x^2), x], x, (b + 2cx)/\sqrt{a + bx + cx^2}], x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1160  $\text{Int}[(d_*) + (e_*)(x_)) * ((a_*) + (b_*)(x_) + (c_*)(x_)^2)^{p_}, x\_Symbol] \rightarrow \text{Simp}[e * ((a + bx + cx^2)^{p+1} / (2c(p+1))), x] + \text{Simp}[(2cd - be) / (2c) \text{ Int}[(a + bx + cx^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, p\}, x] \ \&\& \ \text{NeQ}[p, -1]$

rule 1384  $\text{Int}[(u_*) * ((a_*) + (c_*)(x_)^{n2_}) + (b_*)(x_)^{n_})^{p_}, x\_Symbol] \rightarrow \text{Simp}[(a + bx^n + cx^{2n})^{\text{FracPart}[p]} / (c^{\text{IntPart}[p]} * (b/2 + cx^n)^{2 * \text{FracPart}[p]}) \text{ Int}[u * (b/2 + cx^n)^{2p}, x], x] /; \text{FreeQ}[\{a, b, c, n, p\}, x] \ \&\& \ \text{EqQ}[n2, 2n] \ \&\& \ \text{EqQ}[b^2 - 4ac, 0] \ \&\& \ \text{IntegerQ}[p - 1/2] \ \&\& \ \text{NeQ}[u, x^{(n-1)}] \ \&\& \ \text{NeQ}[u, x^{(2n-1)}] \ \&\& \ !(\text{EqQ}[p, 1/2] \ \&\& \ \text{EqQ}[u, x^{(-2n-1)}])$

```
rule 2192 Int[(Pq_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{q =
Expon[Pq, x], e = Coeff[Pq, x, Expon[Pq, x]]}, Simp[e*x^(q - 1)*((a + b*x +
c*x^2)^(p + 1)/(c*(q + 2*p + 1))), x] + Simp[1/(c*(q + 2*p + 1)) Int[(a
+ b*x + c*x^2)^p*ExpandToSum[c*(q + 2*p + 1)*Pq - a*e*(q - 1)*x^(q - 2) - b
*e*(q + p)*x^(q - 1) - c*e*(q + 2*p + 1)*x^q, x], x]] /; FreeQ[{a, b, c
, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && !LeQ[p, -1]
```

### 3.38.4 Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.67

| method  | result  |
|---------|---|
| risch   | $\frac{(48bx^3d^3+8ex^2bd^2+96ad^3x+24bcd^2x-10bde^2x+48ad^2e-52bcde+15be^3)\sqrt{dx^2+ex+c}\sqrt{(bx^2+a)^2}}{192d^3(bx^2+a)} + \frac{(64acd^3-16e^2d^2a-16b^2e^2d^2)}{192d^3(bx^2+a)}$  |
| default | $\frac{\sqrt{(bx^2+a)^2} \left( 96d^{\frac{7}{2}}(dx^2+ex+c)^{\frac{3}{2}}bx - 80d^{\frac{5}{2}}(dx^2+ex+c)^{\frac{3}{2}}be + 192d^{\frac{9}{2}}\sqrt{dx^2+ex+c}ax - 48d^{\frac{7}{2}}\sqrt{dx^2+ex+c}bcx + 60d^{\frac{5}{2}}\sqrt{dx^2+ex+c}b^2x \right)}{192d^3(bx^2+a)}$ |

```
input int((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/192*(48*b*d^3*x^3+8*b*d^2*e*x^2+96*a*d^3*x+24*b*c*d^2*x-10*b*d*e^2*x+48*
a*d^2*e-52*b*c*d*e+15*b*e^3)*(d*x^2+e*x+c)^(1/2)/d^3*((b*x^2+a)^2)^(1/2)/(
b*x^2+a)+1/128*(64*a*c*d^3-16*a*d^2*e^2-16*b*c^2*d^2+24*b*c*d*e^2-5*b*e^4)
/d^(7/2)*ln((1/2*e+d*x)/d^(1/2)+(d*x^2+e*x+c)^(1/2))*((b*x^2+a)^2)^(1/2)/(
b*x^2+a)
```

### 3.38.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 361, normalized size of antiderivative = 1.28

$$\int \sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4} dx$$

$$= \left[ \frac{3(16bc^2d^2-64acd^3+5be^4-8(3bcd-2ad^2)e^2)\sqrt{d}\log\left(8d^2x^2+8dex-4\sqrt{dx^2+ex+c}(2dx+e)\right)}{\dots} \right]$$

```
input integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x, algorithm="fracas")
```



```
output [1/768*(3*(16*b*c^2*d^2 - 64*a*c*d^3 + 5*b*e^4 - 8*(3*b*c*d - 2*a*d^2)*e^2
)*sqrt(d)*log(8*d^2*x^2 + 8*d*e*x - 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sq
rt(d) + 4*c*d + e^2) + 4*(48*b*d^4*x^3 + 8*b*d^3*e*x^2 + 15*b*d*e^3 - 4*(1
3*b*c*d^2 - 12*a*d^3)*e + 2*(12*b*c*d^3 + 48*a*d^4 - 5*b*d^2*e^2)*x)*sqrt(
d*x^2 + e*x + c))/d^4, 1/384*(3*(16*b*c^2*d^2 - 64*a*c*d^3 + 5*b*e^4 - 8*(
3*b*c*d - 2*a*d^2)*e^2)*sqrt(-d)*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x +
e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) + 2*(48*b*d^4*x^3 + 8*b*d^3*e*x^2 +
15*b*d*e^3 - 4*(13*b*c*d^2 - 12*a*d^3)*e + 2*(12*b*c*d^3 + 48*a*d^4 - 5*b*
d^2*e^2)*x)*sqrt(d*x^2 + e*x + c))/d^4]
```

### 3.38.6 Sympy [F]

$$\int \sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4} dx = \int \sqrt{c + dx^2 + ex} \sqrt{(a + bx^2)^2} dx$$

```
input integrate((d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2), x)
```

```
output Integral(sqrt(c + d*x**2 + e*x)*sqrt((a + b*x**2)**2), x)
```

### 3.38.7 Maxima [F(-2)]

Exception generated.

$$\int \sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4} dx = \text{Exception raised: ValueError}$$

```
input integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2), x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**3.38.8 Giac [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 262, normalized size of antiderivative = 0.93

$$\int \sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4} dx$$

$$= \frac{1}{192} \sqrt{dx^2+ex+c} \left( 2 \left( 4 \left( 6bx\operatorname{sgn}(bx^2+a) + \frac{b\operatorname{sgn}(bx^2+a)}{d} \right) x + \frac{12bcd^2\operatorname{sgn}(bx^2+a) + 48ad^3\operatorname{sgn}(bx^2+a)}{d^3} \right. \right.$$

$$\left. \left. + \frac{(16bc^2d^2\operatorname{sgn}(bx^2+a) - 64acd^3\operatorname{sgn}(bx^2+a) - 24bcde^2\operatorname{sgn}(bx^2+a) + 16ad^2e^2\operatorname{sgn}(bx^2+a) + 5be^4\operatorname{sgn}(bx^2+a))}{128d^{7/2}} \right) \right)$$

```
input integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x, algorithm="giac")
```

```
output 1/192*sqrt(d*x^2 + e*x + c)*(2*(4*(6*b*x*sgn(b*x^2 + a) + b*e*sgn(b*x^2 + a)/d)*x + (12*b*c*d^2*sgn(b*x^2 + a) + 48*a*d^3*sgn(b*x^2 + a) - 5*b*d*e^2*sgn(b*x^2 + a))/d^3)*x - (52*b*c*d*e*sgn(b*x^2 + a) - 48*a*d^2*e*sgn(b*x^2 + a) - 15*b*e^3*sgn(b*x^2 + a))/d^3) + 1/128*(16*b*c^2*d^2*sgn(b*x^2 + a) - 64*a*c*d^3*sgn(b*x^2 + a) - 24*b*c*d*e^2*sgn(b*x^2 + a) + 16*a*d^2*e^2*sgn(b*x^2 + a) + 5*b*e^4*sgn(b*x^2 + a))*log(abs(2*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))*sqrt(d) + e))/d^(7/2)
```

**3.38.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4} dx = \int \sqrt{(bx^2+a)^2}\sqrt{dx^2+ex+c} dx$$

```
input int(((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2),x)
```

```
output int(((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2), x)
```

**3.39**  $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x} dx$

3.39.1 Optimal result . . . . . 314  
 3.39.2 Mathematica [A] (verified) . . . . . 315  
 3.39.3 Rubi [A] (verified) . . . . . 315  
 3.39.4 Maple [A] (verified) . . . . . 319  
 3.39.5 Fricas [A] (verification not implemented) . . . . . 319  
 3.39.6 Sympy [F] . . . . . 320  
 3.39.7 Maxima [F(-2)] . . . . . 320  
 3.39.8 Giac [F(-2)] . . . . . 321  
 3.39.9 Mupad [F(-1)] . . . . . 321

**3.39.1 Optimal result**

Integrand size = 40, antiderivative size = 286

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x} dx$$

$$= \frac{(8ad^2 - be^2 - 2bdex)\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{8d^2(a+bx^2)}$$

$$+ \frac{b(c+ex+dx^2)^{3/2}\sqrt{a^2+2abx^2+b^2x^4}}{3d(a+bx^2)}$$

$$+ \frac{e(8ad^2 - b(4cd - e^2))\sqrt{a^2+2abx^2+b^2x^4}\operatorname{arctanh}\left(\frac{e+2dx}{2\sqrt{d}\sqrt{c+ex+dx^2}}\right)}{16d^{5/2}(a+bx^2)}$$

$$- \frac{a\sqrt{c}\sqrt{a^2+2abx^2+b^2x^4}\operatorname{arctanh}\left(\frac{2c+ex}{2\sqrt{c}\sqrt{c+ex+dx^2}}\right)}{a+bx^2}$$

output

```
1/3*b*(d*x^2+e*x+c)^(3/2)*((b*x^2+a)^2)^(1/2)/d/(b*x^2+a)+1/16*e*(8*a*d^2-
b*(4*c*d-e^2))*arctanh(1/2*(2*d*x+e)/d^(1/2)/(d*x^2+e*x+c)^(1/2))*((b*x^2+
a)^2)^(1/2)/d^(5/2)/(b*x^2+a)-a*arctanh(1/2*(e*x+2*c)/c^(1/2)/(d*x^2+e*x+c
)^(1/2))*c^(1/2)*((b*x^2+a)^2)^(1/2)/(b*x^2+a)+1/8*(-2*b*d*e*x+8*a*d^2-b*e
^2)*(d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/d^2/(b*x^2+a)
```

### 3.39.2 Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.61

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x} dx$$

$$= \frac{\sqrt{(a+bx^2)^2} \left( 2\sqrt{d}\sqrt{c+x(e+dx)}(24ad^2+b(8cd-3e^2+2dex+8d^2x^2)) + 96a\sqrt{cd}^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{dx}-\sqrt{c}}{\sqrt{c}}\right) \right)}{48d^{5/2}(a+bx^2)}$$

input `Integrate[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x,x]`

output `(Sqrt[(a + b*x^2)^2]*(2*Sqrt[d]*Sqrt[c + x*(e + d*x)]*(24*a*d^2 + b*(8*c*d - 3*e^2 + 2*d*e*x + 8*d^2*x^2)) + 96*a*Sqrt[c]*d^(5/2)*ArcTanh[(Sqrt[d]*x - Sqrt[c + x*(e + d*x)])/Sqrt[c]] - 3*e*(8*a*d^2 + b*(-4*c*d + e^2))*Log[e + 2*d*x - 2*Sqrt[d]*Sqrt[c + x*(e + d*x)]])/(48*d^(5/2)*(a + b*x^2))`

### 3.39.3 Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 210, normalized size of antiderivative = 0.73, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.275$ , Rules used = {1384, 27, 2184, 27, 1231, 27, 1269, 1092, 219, 1154, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a^2+2abx^2+b^2x^4}\sqrt{c+dx^2+ex}}{x} dx$$

$$\downarrow 1384$$

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{b(bx^2+a)\sqrt{dx^2+ex+c}}{x} dx}{b(a+bx^2)}$$

$$\downarrow 27$$

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{(bx^2+a)\sqrt{dx^2+ex+c}}{x} dx}{a+bx^2}$$

$$\downarrow 2184$$

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \left( \int \frac{3(2ad-bex)\sqrt{dx^2+ex+c}}{2x} dx + \frac{b(c+dx^2+ex)^{3/2}}{3d} \right)}{a+bx^2}$$

---

3.39.  $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x} dx$

$$\begin{array}{c}
 \downarrow 27 \\
 \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{\int \frac{(2ad - be)x \sqrt{dx^2 + ex + c}}{2d} dx + \frac{b(c + dx^2 + ex)^{3/2}}{3d}}{a + bx^2} \right)}{a + bx^2} \\
 \downarrow 1231 \\
 \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{\frac{\sqrt{c + dx^2 + ex} (8ad^2 - 2bdex - be^2)}{4d} - \frac{\int -\frac{16acd^2 + e(8ad^2 - b(4cd - e^2))x}{2x\sqrt{dx^2 + ex + c}} dx}{2d} + \frac{b(c + dx^2 + ex)^{3/2}}{3d}}{a + bx^2} \right)}{a + bx^2} \\
 \downarrow 27 \\
 \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{\frac{\int \frac{16acd^2 + e(8ad^2 - b(4cd - e^2))x}{x\sqrt{dx^2 + ex + c}} dx}{8d} + \frac{\sqrt{c + dx^2 + ex} (8ad^2 - 2bdex - be^2)}{4d} + \frac{b(c + dx^2 + ex)^{3/2}}{3d}}{a + bx^2} \right)}{a + bx^2} \\
 \downarrow 1269 \\
 \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{\frac{e(8ad^2 - b(4cd - e^2)) \int \frac{1}{\sqrt{dx^2 + ex + c}} dx + 16acd^2 \int \frac{1}{x\sqrt{dx^2 + ex + c}} dx + \frac{\sqrt{c + dx^2 + ex} (8ad^2 - 2bdex - be^2)}{4d}}{2d} + \frac{b(c + dx^2 + ex)^{3/2}}{3d}}{a + bx^2} \right)}{a + bx^2} \\
 \downarrow 1092 \\
 \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{\frac{2e(8ad^2 - b(4cd - e^2)) \int \frac{1}{4d - \frac{(e + 2dx)^2}{dx^2 + ex + c}} dx + \frac{e + 2dx}{\sqrt{dx^2 + ex + c}} + 16acd^2 \int \frac{1}{x\sqrt{dx^2 + ex + c}} dx}{8d} + \frac{\sqrt{c + dx^2 + ex} (8ad^2 - 2bdex - be^2)}{4d} + \frac{b(c + dx^2 + ex)^{3/2}}{3d}}{a + bx^2} \right)}{a + bx^2} \\
 \downarrow 219 \\
 \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{\frac{16acd^2 \int \frac{1}{x\sqrt{dx^2 + ex + c}} dx + \frac{e(8ad^2 - b(4cd - e^2)) \operatorname{arctanh}\left(\frac{2dx + e}{2\sqrt{d}\sqrt{c + dx^2 + ex}}\right)}{\sqrt{d}}}{8d} + \frac{\sqrt{c + dx^2 + ex} (8ad^2 - 2bdex - be^2)}{4d} + \frac{b(c + dx^2 + ex)^{3/2}}{3d}}{a + bx^2} \right)}{a + bx^2} \\
 \downarrow 1154
 \end{array}$$

---

3.39.  $\int \frac{\sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{x} dx$

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4}}{a + bx^2} \left( \frac{e(8ad^2 - b(4cd - e^2)) \operatorname{arctanh}\left(\frac{2dx + e}{2\sqrt{d}\sqrt{c + dx^2 + ex}}\right) - 32acd^2 \int \frac{1}{4c - \frac{(2c + ex)^2}{dx^2 + ex + c}} d \frac{2c + ex}{\sqrt{dx^2 + ex + c}}}{8d} + \frac{\sqrt{c + dx^2 + ex}(8ad^2 - 2bdex - be^2)}{4d} \right)$$


---

219

---


$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4}}{a + bx^2} \left( \frac{e(8ad^2 - b(4cd - e^2)) \operatorname{arctanh}\left(\frac{2dx + e}{2\sqrt{d}\sqrt{c + dx^2 + ex}}\right) - 16a\sqrt{cd}^2 \operatorname{arctanh}\left(\frac{2c + ex}{2\sqrt{c}\sqrt{c + dx^2 + ex}}\right)}{8d} + \frac{\sqrt{c + dx^2 + ex}(8ad^2 - 2bdex - be^2)}{4d} \right)$$

```
input Int[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x,x]
```

```
output (Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4]*((b*(c + e*x + d*x^2)^(3/2))/(3*d) + (((8*a*d^2 - b*e^2 - 2*b*d*e*x)*Sqrt[c + e*x + d*x^2])/(4*d) + ((e*(8*a*d^2 - b*(4*c*d - e^2))*ArcTanh[(e + 2*d*x)/(2*Sqrt[d]*Sqrt[c + e*x + d*x^2])])/Sqrt[d] - 16*a*Sqrt[c]*d^2*ArcTanh[(2*c + e*x)/(2*Sqrt[c]*Sqrt[c + e*x + d*x^2])])/(8*d))/(2*d)))/(a + b*x^2)
```

3.39.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

```
rule 1092 Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Simp[2 Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x]
```

rule 1154 `Int[1/(((d_.) + (e_.)*(x_.))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] := Simp[-2 Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 1231 `Int[((d_.) + (e_.)*(x_.))^(m_.)*((f_.) + (g_.)*(x_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x)^(m + 1)*(c*e*f*(m + 2*p + 2) - g*(c*d + 2*c*d*p - b*e*p) + g*c*e*(m + 2*p + 1)*x)*((a + b*x + c*x^2)^p/(c*e^2*(m + 2*p + 1)*(m + 2*p + 2))), x] - Simp[p/(c*e^2*(m + 2*p + 1)*(m + 2*p + 2)) Int[(d + e*x)^m*(a + b*x + c*x^2)^(p - 1)*Simp[c*e*f*(b*d - 2*a*e)*(m + 2*p + 2) + g*(a*e*(b*e - 2*c*d*m + b*e*m) + b*d*(b*e*p - c*d - 2*c*d*p)) + (c*e*f*(2*c*d - b*e)*(m + 2*p + 2) + g*(b^2*e^2*(p + m + 1) - 2*c^2*d^2*(1 + 2*p) - c*e*(b*d*(m - 2*p) + 2*a*e*(m + 2*p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && GtQ[p, 0] && (IntegerQ[p] || !RationalQ[m] || (GeQ[m, -1] && LtQ[m, 0])) && !ILtQ[m + 2*p, 0] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])`

rule 1269 `Int[((d_.) + (e_.)*(x_.))^(m_.)*((f_.) + (g_.)*(x_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[g/e Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Simp[(e*f - d*g)/e Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && !IGtQ[m, 0]`

rule 1384 `Int[(u_.)*((a_.) + (c_.)*(x_)^(n2_.) + (b_.)*(x_)^(n_.))^p, x_Symbol] := Simp[(a + b*x^n + c*x^(2*n))^FracPart[p]/(c^IntPart[p]*(b/2 + c*x^n)^(2*FracPart[p])) Int[u*(b/2 + c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p - 1/2] && NeQ[u, x^(n - 1)] && NeQ[u, x^(2*n - 1)] && !(EqQ[p, 1/2] && EqQ[u, x^(-2*n - 1)])`

rule 2184 `Int[(Pq_.)*((d_.) + (e_.)*(x_.))^(m_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := With[{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[f*(d + e*x)^(m + q - 1)*((a + b*x + c*x^2)^(p + 1)/(c*e^(q - 1)*(m + q + 2*p + 1))), x] + Simp[1/(c*e^q*(m + q + 2*p + 1)) Int[(d + e*x)^m*(a + b*x + c*x^2)^p*ExpandToSum[c*e^q*(m + q + 2*p + 1)*Pq - c*f*(m + q + 2*p + 1)*(d + e*x)^q - f*(d + e*x)^(q - 2)*(b*d*e*(p + 1) + a*e^2*(m + q - 1) - c*d^2*(m + q + 2*p + 1) - e*(2*c*d - b*e)*(m + q + p)*x), x], x] /; GtQ[q, 1] && NeQ[m + q + 2*p + 1, 0] /; FreeQ[{a, b, c, d, e, m, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !(IGtQ[m, 0] && RationalQ[a, b, c, d, e] && (IntegerQ[p] || ILtQ[p + 1/2, 0]))`

### 3.39.4 Maple [A] (verified)

Time = 0.57 (sec) , antiderivative size = 251, normalized size of antiderivative = 0.88

| method  | result  |
|---------|---|
| default | $-\frac{\sqrt{(bx^2+a)^2} \left( 48\sqrt{c}d^{\frac{7}{2}} \ln\left(\frac{2c+ex+2\sqrt{c}\sqrt{dx^2+ex+c}}{x}\right) a - 16d^{\frac{5}{2}}(dx^2+ex+c)^{\frac{3}{2}} b + 12d^{\frac{5}{2}}\sqrt{dx^2+ex+c} bex - 48d^{\frac{7}{2}}\sqrt{dx^2+ex+c} a + 6d^{\frac{3}{2}} \right)}{48(b$ |

input `int((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x,x,method=_RETURNVERBOSE)`

output 
$$-1/48*((b*x^2+a)^2)^(1/2)*(48*c^(1/2)*d^(7/2)*\ln((2*c+e*x+2*c^(1/2)*(d*x^2+e*x+c)^(1/2))/x)*a-16*d^(5/2)*(d*x^2+e*x+c)^(3/2)*b+12*d^(5/2)*(d*x^2+e*x+c)^(1/2)*b*e*x-48*d^(7/2)*(d*x^2+e*x+c)^(1/2)*a+6*d^(3/2)*(d*x^2+e*x+c)^(1/2)*b*e^2-24*d^3*\ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*a*e+12*\ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*b*c*d^2*e-3*\ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*b*d*e^3)/(b*x^2+a)/d^(7/2)$$

### 3.39.5 Fracas [A] (verification not implemented)

Time = 0.73 (sec) , antiderivative size = 743, normalized size of antiderivative = 2.60

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x} dx$$

$$= \left[ \frac{48 a \sqrt{c} d^3 \log\left(\frac{8 c e x+(4 c d+e^2) x^2-4 \sqrt{d x^2+e x+c}(e x+2 c) \sqrt{c+8 c^2}}{x^2}\right)}{48 a \sqrt{c} d^3} + 3 (b e^3 - 4 (b c d - 2 a d^2) e) \sqrt{d} \log\left(8 d^2 x^2 + 8 d\right) \right]$$

input `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x,x, algorithm="fricas")`



output `[1/96*(48*a*sqrt(c)*d^3*log((8*c*e*x + (4*c*d + e^2)*x^2 - 4*sqrt(d*x^2 + e*x + c))*(e*x + 2*c)*sqrt(c) + 8*c^2)/x^2) + 3*(b*e^3 - 4*(b*c*d - 2*a*d^2)*e)*sqrt(d)*log(8*d^2*x^2 + 8*d*e*x + 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 4*(8*b*d^3*x^2 + 2*b*d^2*e*x + 8*b*c*d^2 + 24*a*d^3 - 3*b*d*e^2)*sqrt(d*x^2 + e*x + c))/d^3, 1/48*(24*a*sqrt(c)*d^3*log((8*c*e*x + (4*c*d + e^2)*x^2 - 4*sqrt(d*x^2 + e*x + c))*(e*x + 2*c)*sqrt(c) + 8*c^2)/x^2) - 3*(b*e^3 - 4*(b*c*d - 2*a*d^2)*e)*sqrt(-d)*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) + 2*(8*b*d^3*x^2 + 2*b*d^2*e*x + 8*b*c*d^2 + 24*a*d^3 - 3*b*d*e^2)*sqrt(d*x^2 + e*x + c))/d^3, 1/96*(96*a*sqrt(-c)*d^3*arctan(1/2*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(-c)/(c*d*x^2 + c*e*x + c^2)) + 3*(b*e^3 - 4*(b*c*d - 2*a*d^2)*e)*sqrt(d)*log(8*d^2*x^2 + 8*d*e*x + 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 4*(8*b*d^3*x^2 + 2*b*d^2*e*x + 8*b*c*d^2 + 24*a*d^3 - 3*b*d*e^2)*sqrt(d*x^2 + e*x + c))/d^3, 1/48*(48*a*sqrt(-c)*d^3*arctan(1/2*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(-c)/(c*d*x^2 + c*e*x + c^2)) - 3*(b*e^3 - 4*(b*c*d - 2*a*d^2)*e)*sqrt(-d)*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) + 2*(8*b*d^3*x^2 + 2*b*d^2*e*x + 8*b*c*d^2 + 24*a*d^3 - 3*b*d*e^2)*sqrt(d*x^2 + e*x + c))/d^3]`

### 3.39.6 Sympy [F]

$$\int \frac{\sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{x} dx = \int \frac{\sqrt{c + dx^2 + ex} \sqrt{(a + bx^2)^2}}{x} dx$$

input `integrate((d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2)/x,x)`

output `Integral(sqrt(c + d*x**2 + e*x)*sqrt((a + b*x**2)**2)/x, x)`

### 3.39.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{x} dx = \text{Exception raised: ValueError}$$

input `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x,x, algorithm="maxima")`

---

3.39.  $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x} dx$

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation \*may\* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e

### 3.39.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x} dx = \text{Exception raised: TypeError}$$

input `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x,x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:Error: Bad Argument Type

### 3.39.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x} dx = \int \frac{\sqrt{(bx^2+a)^2}\sqrt{dx^2+ex+c}}{x} dx$$

input `int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x,x)`

output `int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x, x)`

**3.40**  $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^2} dx$

3.40.1 Optimal result . . . . . 322  
 3.40.2 Mathematica [A] (verified) . . . . . 323  
 3.40.3 Rubi [A] (verified) . . . . . 323  
 3.40.4 Maple [A] (verified) . . . . . 327  
 3.40.5 Fricas [A] (verification not implemented) . . . . . 328  
 3.40.6 Sympy [F] . . . . . 328  
 3.40.7 Maxima [F(-2)] . . . . . 329  
 3.40.8 Giac [A] (verification not implemented) . . . . . 329  
 3.40.9 Mupad [F(-1)] . . . . . 330

**3.40.1 Optimal result**

Integrand size = 40, antiderivative size = 294

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^2} dx$$

$$= \frac{((bc+4ad)e+2d(bc+2ad)x)\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{4cd(a+bx^2)}$$

$$- \frac{a(c+ex+dx^2)^{3/2}\sqrt{a^2+2abx^2+b^2x^4}}{cx(a+bx^2)}$$

$$+ \frac{(4bcd+8ad^2-be^2)\sqrt{a^2+2abx^2+b^2x^4}\operatorname{arctanh}\left(\frac{e+2dx}{2\sqrt{d}\sqrt{c+ex+dx^2}}\right)}{8d^{3/2}(a+bx^2)}$$

$$- \frac{ae\sqrt{a^2+2abx^2+b^2x^4}\operatorname{arctanh}\left(\frac{2c+ex}{2\sqrt{c}\sqrt{c+ex+dx^2}}\right)}{2\sqrt{c}(a+bx^2)}$$

```
output -a*(d*x^2+e*x+c)^(3/2)*((b*x^2+a)^2)^(1/2)/c/x/(b*x^2+a)+1/8*(8*a*d^2+4*b*
c*d-b*e^2)*arctanh(1/2*(2*d*x+e)/d^(1/2)/(d*x^2+e*x+c)^(1/2))*((b*x^2+a)^2
)^(1/2)/d^(3/2)/(b*x^2+a)-1/2*a*e*arctanh(1/2*(e*x+2*c)/c^(1/2)/(d*x^2+e*x
+c)^(1/2))*((b*x^2+a)^2)^(1/2)/(b*x^2+a)/c^(1/2)+1/4*((4*a*d+b*c)*e+2*d*(2
*a*d+b*c)*x)*(d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/c/d/(b*x^2+a)
```

### 3.40.2 Mathematica [A] (verified)

Time = 0.61 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.58

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^2} dx$$

$$= \frac{\sqrt{(a+bx^2)^2} \left( \sqrt{c}(4bcd+8ad^2-be^2) x \operatorname{arctanh}\left(\frac{e+2dx}{2\sqrt{d}\sqrt{c+x(e+dx)}}\right) + 2\sqrt{d} \left( \sqrt{c}\sqrt{c+x(e+dx)}(-4ad+bx(e+dx)) \right) \right)}{8\sqrt{cd}^{3/2}x(a+bx^2)}$$

input `Integrate[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^2,x]`

output `(Sqrt[(a + b*x^2)^2]*(Sqrt[c]*(4*b*c*d + 8*a*d^2 - b*e^2)*x*ArcTanh[(e + 2*d*x)/(2*Sqrt[d]*Sqrt[c + x*(e + d*x)])] + 2*Sqrt[d]*(Sqrt[c]*Sqrt[c + x*(e + d*x)]*(-4*a*d + b*x*(e + 2*d*x)) + 4*a*d*e*x*ArcTanh[(Sqrt[d]*x - Sqrt[c + x*(e + d*x)]/Sqrt[c])])/(8*Sqrt[c]*d^(3/2)*x*(a + b*x^2))`

### 3.40.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 212, normalized size of antiderivative = 0.72, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$ , Rules used = {1384, 27, 2181, 27, 1231, 25, 27, 1269, 1092, 219, 1154, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a^2+2abx^2+b^2x^4}\sqrt{c+dx^2+ex}}{x^2} dx$$

$$\downarrow \text{1384}$$

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{b(bx^2+a)\sqrt{dx^2+ex+c}}{x^2} dx}{b(a+bx^2)}$$

$$\downarrow \text{27}$$

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{(bx^2+a)\sqrt{dx^2+ex+c}}{x^2} dx}{a+bx^2}$$

$$\downarrow \text{2181}$$

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \left( -\frac{\int -\frac{(ae+2(bc+2ad)x)\sqrt{dx^2+ex+c}}{2x} dx}{c} - \frac{a(c+dx^2+ex)^{3/2}}{cx} \right)}{a+bx^2}$$

---

3.40.  $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^2} dx$

$$\begin{array}{c}
\downarrow 27 \\
\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{\int \frac{(ae+2(bc+2ad)x)\sqrt{dx^2+ex+c}}{2c} dx - \frac{a(c+dx^2+ex)^{3/2}}{cx}}{a + bx^2} \right)}{a + bx^2} \\
\downarrow 1231 \\
\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{\frac{\sqrt{c+dx^2+ex}(e(4ad+bc)+2dx(2ad+bc))}{2d} - \frac{\int -\frac{c(4ade+(8ad^2+4bcd-be^2)x)}{x\sqrt{dx^2+ex+c}} dx}{2c} - \frac{a(c+dx^2+ex)^{3/2}}{cx}}{a + bx^2} \right)}{a + bx^2} \\
\downarrow 25 \\
\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{\frac{\int \frac{c(4ade+(8ad^2+4bcd-be^2)x}{x\sqrt{dx^2+ex+c}} dx}{4d} + \frac{\sqrt{c+dx^2+ex}(e(4ad+bc)+2dx(2ad+bc))}{2c} - \frac{a(c+dx^2+ex)^{3/2}}{cx}}{a + bx^2} \right)}{a + bx^2} \\
\downarrow 27 \\
\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{\frac{c \int \frac{4ade+(8ad^2+4bcd-be^2)x}{x\sqrt{dx^2+ex+c}} dx}{4d} + \frac{\sqrt{c+dx^2+ex}(e(4ad+bc)+2dx(2ad+bc))}{2c} - \frac{a(c+dx^2+ex)^{3/2}}{cx}}{a + bx^2} \right)}{a + bx^2} \\
\downarrow 1269 \\
\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{\frac{c \left( (8ad^2+4bcd-be^2) \int \frac{1}{\sqrt{dx^2+ex+c}} dx + 4ade \int \frac{1}{x\sqrt{dx^2+ex+c}} dx \right) + \sqrt{c+dx^2+ex}(e(4ad+bc)+2dx(2ad+bc))}{2c} - \frac{a(c+dx^2+ex)^{3/2}}{cx}}{a + bx^2} \right)}{a + bx^2} \\
\downarrow 1092 \\
\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{\frac{c \left( 2(8ad^2+4bcd-be^2) \int \frac{1}{4d - \frac{(e+2dx)^2}{dx^2+ex+c}} dx + 4ade \int \frac{1}{x\sqrt{dx^2+ex+c}} dx \right) + \sqrt{c+dx^2+ex}(e(4ad+bc)+2dx(2ad+bc))}{2c} - \frac{a(c+dx^2+ex)^{3/2}}{cx}}{a + bx^2} \right)}{a + bx^2} \\
\downarrow 219
\end{array}$$

---

3.40.  $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^2} dx$

$$\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{c \left( 4ade \int \frac{1}{x\sqrt{dx^2+ex+c}} dx + \frac{(8ad^2+4bcd-be^2) \operatorname{arctanh}\left(\frac{2dx+e}{2\sqrt{d}\sqrt{c+dx^2+ex}}\right)}{\sqrt{d}} \right)}{4d} + \frac{\sqrt{c+dx^2+ex}(e(4ad+bc)+2dx(2ad+bc))}{2c} \right) \frac{1}{2c} + \frac{\sqrt{c+dx^2+ex}(e(4ad+bc)+2dx(2ad+bc))}{2d}$$


---

$a + bx^2$

↓ 1154

$$\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{c \left( \frac{(8ad^2+4bcd-be^2) \operatorname{arctanh}\left(\frac{2dx+e}{2\sqrt{d}\sqrt{c+dx^2+ex}}\right)}{\sqrt{d}} - 8ade \int \frac{1}{4c - \frac{(2c+ex)^2}{dx^2+ex+c}} d - \frac{2c+ex}{\sqrt{dx^2+ex+c}} \right)}{4d} + \frac{\sqrt{c+dx^2+ex}(e(4ad+bc)+2dx(2ad+bc))}{2c} \right) \frac{1}{2c} + \frac{\sqrt{c+dx^2+ex}(e(4ad+bc)+2dx(2ad+bc))}{2d}$$


---

$a + bx^2$

↓ 219

$$\sqrt{a^2 + 2abx^2 + b^2x^4} \left( \frac{c \left( \frac{(8ad^2+4bcd-be^2) \operatorname{arctanh}\left(\frac{2dx+e}{2\sqrt{d}\sqrt{c+dx^2+ex}}\right)}{\sqrt{d}} - \frac{4ade \operatorname{arctanh}\left(\frac{2c+ex}{2\sqrt{c}\sqrt{c+dx^2+ex}}\right)}{\sqrt{c}} \right)}{4d} + \frac{\sqrt{c+dx^2+ex}(e(4ad+bc)+2dx(2ad+bc))}{2c} \right) \frac{1}{2c} + \frac{\sqrt{c+dx^2+ex}(e(4ad+bc)+2dx(2ad+bc))}{2d}$$


---

$a + bx^2$

input `Int[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^2,x]`

output `(Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4]*(-(a*(c + e*x + d*x^2)^(3/2))/(c*x)) + ((b*c + 4*a*d)*e + 2*d*(b*c + 2*a*d)*x)*Sqrt[c + e*x + d*x^2])/(2*d) + (c*((4*b*c*d + 8*a*d^2 - b*e^2)*ArcTanh[(e + 2*d*x)/(2*Sqrt[d]*Sqrt[c + e*x + d*x^2])])/Sqrt[d] - (4*a*d*e*ArcTanh[(2*c + e*x)/(2*Sqrt[c]*Sqrt[c + e*x + d*x^2])])/Sqrt[c])/(4*d))/(2*c))/(a + b*x^2)`

## 3.40.3.1 Defintions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, x], x]$
- rule 27  $\text{Int}[(a_)*(\text{Fx}_), x\_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[\text{Fx}, (b_)*(\text{Gx}_) /; \text{FreeQ}[b, x]]$
- rule 219  $\text{Int}[(a_) + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))* \text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 1092  $\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_) + (c_)*(x_)^2], x\_Symbol] \rightarrow \text{Simp}[2 \quad \text{Subst}[\text{Int}[1/(4*c - x^2), x], x, (b + 2*c*x)/\text{Sqrt}[a + b*x + c*x^2]], x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1154  $\text{Int}[1/(((d_) + (e_)*(x_))*\text{Sqrt}[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x\_Symbol] \rightarrow \text{Simp}[-2 \quad \text{Subst}[\text{Int}[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/\text{Sqrt}[a + b*x + c*x^2]], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$
- rule 1231  $\text{Int}[(d_) + (e_)*(x_))^{(m_)*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^{(p_)}, x\_Symbol] \rightarrow \text{Simp}[(d + e*x)^{(m+1)}*(c*e*f*(m+2*p+2) - g*(c*d + 2*c*d*p - b*e*p) + g*c*e*(m+2*p+1)*x)*((a + b*x + c*x^2)^p / (c*e^2*(m+2*p+1)*(m+2*p+2))), x] - \text{Simp}[p/(c*e^2*(m+2*p+1)*(m+2*p+2)) \quad \text{Int}[(d + e*x)^m*(a + b*x + c*x^2)^{(p-1)}*\text{Simp}[c*e*f*(b*d - 2*a*e)*(m+2*p+2) + g*(a*e*(b*e - 2*c*d*m + b*e*m) + b*d*(b*e*p - c*d - 2*c*d*p)) + (c*e*f*(2*c*d - b*e)*(m+2*p+2) + g*(b^2*e^2*(p+m+1) - 2*c^2*d^2*(1+2*p) - c*e*(b*d*(m-2*p) + 2*a*e*(m+2*p+1)))]*x, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, m\}, x] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ (\text{IntegerQ}[p] \ || \ !\text{RationalQ}[m] \ || \ (\text{GeQ}[m, -1] \ \&\& \ \text{LtQ}[m, 0])) \ \&\& \ !\text{ILtQ}[m+2*p, 0] \ \&\& \ (\text{IntegerQ}[m] \ || \ \text{IntegerQ}[p] \ || \ \text{IntegersQ}[2*m, 2*p])$
- rule 1269  $\text{Int}[(d_) + (e_)*(x_))^{(m_)*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^{(p_)}, x\_Symbol] \rightarrow \text{Simp}[g/e \quad \text{Int}[(d + e*x)^{(m+1)}*(a + b*x + c*x^2)^p, x], x] + \text{Simp}[(e*f - d*g)/e \quad \text{Int}[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, m, p\}, x] \ \&\& \ !\text{IGtQ}[m, 0]$

```
rule 1384 Int[(u_.)*((a_.) + (c_.)*(x_)^(n2_.) + (b_.)*(x_)^(n_.))^(p_), x_Symbol] := S
imp[(a + b*x^n + c*x^(2*n))^FracPart[p]/(c^IntPart[p]*(b/2 + c*x^n)^(2*Frac
Part[p])) Int[u*(b/2 + c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x]
&& EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p - 1/2] && NeQ[u, x^(n
- 1)] && NeQ[u, x^(2*n - 1)] && !(EqQ[p, 1/2] && EqQ[u, x^(-2*n - 1)])
```

```
rule 2181 Int[(Pq_)*((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_
), x_Symbol] := With[{Qx = PolynomialQuotient[Pq, d + e*x, x], R = Polynomi
alRemainder[Pq, d + e*x, x]}, Simp[(e*R*(d + e*x)^(m + 1)*(a + b*x + c*x^2)
^(p + 1))/((m + 1)*(c*d^2 - b*d*e + a*e^2)), x] + Simp[1/((m + 1)*(c*d^2 -
b*d*e + a*e^2)) Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p*ExpandToSum[(m
+ 1)*(c*d^2 - b*d*e + a*e^2)*Qx + c*d*R*(m + 1) - b*e*R*(m + p + 2) - c*e*R
*(m + 2*p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Pq,
x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && LtQ[m, -1]
```

### 3.40.4 Maple [A] (verified)

Time = 0.59 (sec) , antiderivative size = 229, normalized size of antiderivative = 0.78

| method  | result   |
|---------|--|
| risch   | $-\frac{a\sqrt{dx^2+ex+c}\sqrt{bx^2+a}}{x(bx^2+a)} + \frac{\left(\sqrt{d}a\ln\left(\frac{\frac{5}{2}+dx}{\sqrt{d}}+\sqrt{dx^2+ex+c}\right) + \frac{bc\ln\left(\frac{\frac{5}{2}+dx}{\sqrt{d}}+\sqrt{dx^2+ex+c}\right)}{2\sqrt{d}} - \frac{ae\ln\left(\frac{2c+ex+2\sqrt{c}\sqrt{dx^2+ex+c}}{x}\right)}{2\sqrt{c}}\right)}{bx^2+a}$ |
| default | $\frac{\sqrt{(bx^2+a)^2}\left(8d^{\frac{7}{2}}\sqrt{dx^2+ex+c}ax^2-4d^{\frac{5}{2}}\sqrt{c}\ln\left(\frac{2c+ex+2\sqrt{c}\sqrt{dx^2+ex+c}}{x}\right)ae+4d^{\frac{5}{2}}\sqrt{dx^2+ex+c}bcx^2-8d^{\frac{5}{2}}(dx^2+ex+c)^{\frac{3}{2}}a+8d^{\frac{5}{2}}\right)}{bx^2+a}$  |

```
input int((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^2,x,method=_RETURNVERBOSE)
```

```
output -a*(d*x^2+e*x+c)^(1/2)/x*((b*x^2+a)^2)^(1/2)/(b*x^2+a)+(d^(1/2)*a*ln((1/2*
e+d*x)/d^(1/2)+(d*x^2+e*x+c)^(1/2))+1/2*b*c*ln((1/2*e+d*x)/d^(1/2)+(d*x^2+
e*x+c)^(1/2))/d^(1/2)-1/2*a*e/c^(1/2)*ln((2*c+e*x+2*c^(1/2)*(d*x^2+e*x+c)^(
1/2))/x)+1/2*b*x*(d*x^2+e*x+c)^(1/2)+1/4*b/d*e*(d*x^2+e*x+c)^(1/2)-1/8*b/
d^(3/2)*e^2*ln((1/2*e+d*x)/d^(1/2)+(d*x^2+e*x+c)^(1/2)))*((b*x^2+a)^2)^(1/
2)/(b*x^2+a)
```



### 3.40.5 Fracas [A] (verification not implemented)

Time = 0.66 (sec) , antiderivative size = 731, normalized size of antiderivative = 2.49

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^2} dx$$

$$= \left[ \frac{4a\sqrt{cd^2ex} \log\left(\frac{8cex+(4cd+e^2)x^2-4\sqrt{dx^2+ex+c}(ex+2c)\sqrt{c+8c^2}}{x^2}\right) - (4bc^2d+8acd^2-bce^2)\sqrt{dx} \log\left(\frac{8d^2x^2+8cd^2x}{16cd^2x}\right)}{16cd^2x} \right]$$

input `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^2,x, algorithm="fracas")`

output `[1/16*(4*a*sqrt(c)*d^2*e*x*log((8*c*e*x + (4*c*d + e^2)*x^2 - 4*sqrt(d*x^2 + e*x + c)*(e*x + 2*c))*sqrt(c) + 8*c^2)/x^2) - (4*b*c^2*d + 8*a*c*d^2 - b*c*e^2)*sqrt(d)*x*log(8*d^2*x^2 + 8*d*e*x - 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 4*(2*b*c*d^2*x^2 + b*c*d*e*x - 4*a*c*d^2)*sqrt(d*x^2 + e*x + c)/(c*d^2*x), 1/8*(2*a*sqrt(c)*d^2*e*x*log((8*c*e*x + (4*c*d + e^2)*x^2 - 4*sqrt(d*x^2 + e*x + c)*(e*x + 2*c))*sqrt(c) + 8*c^2)/x^2) - (4*b*c^2*d + 8*a*c*d^2 - b*c*e^2)*sqrt(-d)*x*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) + 2*(2*b*c*d^2*x^2 + b*c*d*e*x - 4*a*c*d^2)*sqrt(d*x^2 + e*x + c)/(c*d^2*x), 1/16*(8*a*sqrt(-c)*d^2*e*x*arctan(1/2*sqrt(d*x^2 + e*x + c)*(e*x + 2*c))*sqrt(-c)/(c*d*x^2 + c*e*x + c^2)) - (4*b*c^2*d + 8*a*c*d^2 - b*c*e^2)*sqrt(d)*x*log(8*d^2*x^2 + 8*d*e*x - 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 4*(2*b*c*d^2*x^2 + b*c*d*e*x - 4*a*c*d^2)*sqrt(d*x^2 + e*x + c)/(c*d^2*x), 1/8*(4*a*sqrt(-c)*d^2*e*x*arctan(1/2*sqrt(d*x^2 + e*x + c)*(e*x + 2*c))*sqrt(-c)/(c*d*x^2 + c*e*x + c^2)) - (4*b*c^2*d + 8*a*c*d^2 - b*c*e^2)*sqrt(-d)*x*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) + 2*(2*b*c*d^2*x^2 + b*c*d*e*x - 4*a*c*d^2)*sqrt(d*x^2 + e*x + c)/(c*d^2*x)]`

### 3.40.6 Sympy [F]

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^2} dx = \int \frac{\sqrt{c+dx^2+ex}\sqrt{(a+bx^2)^2}}{x^2} dx$$

input `integrate((d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2)/x**2,x)`

---

3.40.  $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^2} dx$

output `Integral(sqrt(c + d*x**2 + e*x)*sqrt((a + b*x**2)**2)/x**2, x)`

### 3.40.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{x^2} dx = \text{Exception raised: ValueError}$$

input `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

### 3.40.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 238, normalized size of antiderivative = 0.81

$$\begin{aligned} \int \frac{\sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{x^2} dx &= \frac{ae \arctan\left(-\frac{\sqrt{dx} - \sqrt{dx^2 + ex + c}}{\sqrt{-c}}\right) \operatorname{sgn}(bx^2 + a)}{\sqrt{-c}} \\ &+ \frac{1}{4} \sqrt{dx^2 + ex + c} \left( 2bx \operatorname{sgn}(bx^2 + a) + \frac{b \operatorname{sgn}(bx^2 + a)}{d} \right) \\ &- \frac{(4bcd \operatorname{sgn}(bx^2 + a) + 8ad^2 \operatorname{sgn}(bx^2 + a) - be^2 \operatorname{sgn}(bx^2 + a)) \log\left(\left| 2\left(\sqrt{dx} - \sqrt{dx^2 + ex + c}\right) \sqrt{d} + e \right|\right)}{8d^{\frac{3}{2}}} \\ &+ \frac{\left(\sqrt{dx} - \sqrt{dx^2 + ex + c}\right) a e \operatorname{sgn}(bx^2 + a) + 2ac \sqrt{d} \operatorname{sgn}(bx^2 + a)}{\left(\sqrt{dx} - \sqrt{dx^2 + ex + c}\right)^2 - c} \end{aligned}$$

input `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^2,x, algorithm="giac")`

output  $a*e*\arctan(-(\sqrt{d}*x - \sqrt{d*x^2 + e*x + c})/\sqrt{-c})*\text{sgn}(b*x^2 + a)/\sqrt{-c} + 1/4*\sqrt{d*x^2 + e*x + c}*(2*b*x*\text{sgn}(b*x^2 + a) + b*e*\text{sgn}(b*x^2 + a)/d) - 1/8*(4*b*c*d*\text{sgn}(b*x^2 + a) + 8*a*d^2*\text{sgn}(b*x^2 + a) - b*e^2*\text{sgn}(b*x^2 + a))*\log(\text{abs}(2*(\sqrt{d}*x - \sqrt{d*x^2 + e*x + c})*\sqrt{d} + e))/d^{3/2} + ((\sqrt{d}*x - \sqrt{d*x^2 + e*x + c})*a*e*\text{sgn}(b*x^2 + a) + 2*a*c*\sqrt{d}*\text{sgn}(b*x^2 + a))/((\sqrt{d}*x - \sqrt{d*x^2 + e*x + c})^2 - c)$

### 3.40.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^2} dx = \int \frac{\sqrt{(bx^2+a)^2}\sqrt{dx^2+ex+c}}{x^2} dx$$

input `int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^2, x)`

output `int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^2, x)`

**3.41**  $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx$

3.41.1 Optimal result . . . . . 331  
 3.41.2 Mathematica [A] (verified) . . . . . 332  
 3.41.3 Rubi [A] (verified) . . . . . 332  
 3.41.4 Maple [A] (verified) . . . . . 336  
 3.41.5 Fricas [A] (verification not implemented) . . . . . 336  
 3.41.6 Sympy [F(-1)] . . . . . 337  
 3.41.7 Maxima [F(-2)] . . . . . 337  
 3.41.8 Giac [A] (verification not implemented) . . . . . 338  
 3.41.9 Mupad [F(-1)] . . . . . 339

**3.41.1 Optimal result**

Integrand size = 40, antiderivative size = 288

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx$$

$$= \frac{(ae+2(2bc+ad)x)\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{4cx(a+bx^2)}$$

$$- \frac{a(c+ex+dx^2)^{3/2}\sqrt{a^2+2abx^2+b^2x^4}}{2cx^2(a+bx^2)}$$

$$+ \frac{be\sqrt{a^2+2abx^2+b^2x^4}\operatorname{arctanh}\left(\frac{e+2dx}{2\sqrt{d}\sqrt{c+ex+dx^2}}\right)}{2\sqrt{d}(a+bx^2)}$$

$$- \frac{(8bc^2+4acd-ae^2)\sqrt{a^2+2abx^2+b^2x^4}\operatorname{arctanh}\left(\frac{2c+ex}{2\sqrt{c}\sqrt{c+ex+dx^2}}\right)}{8c^{3/2}(a+bx^2)}$$

output

```
-1/2*a*(d*x^2+e*x+c)^(3/2)*((b*x^2+a)^2)^(1/2)/c/x^2/(b*x^2+a)-1/8*(4*a*c*d-a*e^2+8*b*c^2)*arctanh(1/2*(e*x+2*c)/c^(1/2)/(d*x^2+e*x+c)^(1/2))*((b*x^2+a)^2)^(1/2)/c^(3/2)/(b*x^2+a)+1/2*b*e*arctanh(1/2*(2*d*x+e)/d^(1/2)/(d*x^2+e*x+c)^(1/2))*((b*x^2+a)^2)^(1/2)/(b*x^2+a)/d^(1/2)+1/4*(a*e+2*(a*d+2*b*c)*x)*(d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/c/x/(b*x^2+a)
```

### 3.41.2 Mathematica [A] (verified)

Time = 0.64 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.60

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx = \frac{\sqrt{(a+bx^2)^2}\left(\sqrt{d}(8bc^2+4acd-ae^2)x^2\operatorname{arctanh}\left(\frac{-\sqrt{dx+\sqrt{c+x(e+dx)}}}{\sqrt{c}}\right)\right) + \sqrt{c}\left(\sqrt{d}(2ac+ae^2-4bcx^2)\sqrt{c}\right)}{4c^{3/2}\sqrt{dx^2(a+bx^2)}}$$

input `Integrate[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^3,x]`

output `-1/4*(Sqrt[(a + b*x^2)^2]*(Sqrt[d]*(8*b*c^2 + 4*a*c*d - a*e^2)*x^2*ArcTanh[(-(Sqrt[d]*x) + Sqrt[c + x*(e + d*x)])/Sqrt[c]] + Sqrt[c]*(Sqrt[d]*(2*a*c + a*e*x - 4*b*c*x^2)*Sqrt[c + x*(e + d*x)] + 2*b*c*e*x^2*Log[e + 2*d*x - 2*Sqrt[d]*Sqrt[c + x*(e + d*x)]])))/(c^(3/2)*Sqrt[d]*x^2*(a + b*x^2))`

### 3.41.3 Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 200, normalized size of antiderivative = 0.69, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {1384, 27, 2181, 27, 1230, 1269, 1092, 219, 1154, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{a^2+2abx^2+b^2x^4}\sqrt{c+dx^2+ex}}{x^3} dx \\ & \quad \downarrow \text{1384} \\ & \frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{b(bx^2+a)\sqrt{dx^2+ex+c}}{x^3} dx}{b(a+bx^2)} \\ & \quad \downarrow \text{27} \\ & \frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{(bx^2+a)\sqrt{dx^2+ex+c}}{x^3} dx}{a+bx^2} \\ & \quad \downarrow \text{2181} \end{aligned}$$

---

3.41.  $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx$

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( -\frac{\int \frac{(ae-2(2bc+ad)x)\sqrt{dx^2+ex+c}}{2x^2} dx}{2c} - \frac{a(c+dx^2+ex)^{3/2}}{2cx^2} \right)}{a + bx^2}$$

↓ 27

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( -\frac{\int \frac{(ae-2(2bc+ad)x)\sqrt{dx^2+ex+c}}{4c} dx}{4c} - \frac{a(c+dx^2+ex)^{3/2}}{2cx^2} \right)}{a + bx^2}$$

↓ 1230

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( -\frac{\frac{1}{2} \int \frac{8bc^2+4adc+4bexc-ae^2}{x\sqrt{dx^2+ex+c}} dx - \frac{\sqrt{c+dx^2+ex}(2x(ad+2bc)+ae)}{x}}{4c} - \frac{a(c+dx^2+ex)^{3/2}}{2cx^2} \right)}{a + bx^2}$$

↓ 1269

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( -\frac{\frac{1}{2} \left( -(4acd-ae^2+8bc^2) \int \frac{1}{x\sqrt{dx^2+ex+c}} dx - 4bce \int \frac{1}{\sqrt{dx^2+ex+c}} dx \right) - \frac{\sqrt{c+dx^2+ex}(2x(ad+2bc)+ae)}{x}}{4c} - \frac{a(c+dx^2+ex)^{3/2}}{2cx^2} \right)}{a + bx^2}$$

↓ 1092

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( -\frac{\frac{1}{2} \left( -\left( (4acd-ae^2+8bc^2) \int \frac{1}{x\sqrt{dx^2+ex+c}} dx \right) - 8bce \int \frac{1}{4d - \frac{(e+2dx)^2}{dx^2+ex+c}} d \frac{e+2dx}{\sqrt{dx^2+ex+c}} \right) - \frac{\sqrt{c+dx^2+ex}(2x(ad+2bc)+ae)}{x}}{4c} - \frac{a(c+dx^2+ex)^{3/2}}{2cx^2} \right)}{a + bx^2}$$

↓ 219

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( -\frac{\frac{1}{2} \left( -(4acd-ae^2+8bc^2) \int \frac{1}{x\sqrt{dx^2+ex+c}} dx - \frac{4bce \operatorname{arctanh} \left( \frac{2dx+e}{2\sqrt{d}\sqrt{c+dx^2+ex}} \right)}{\sqrt{d}} \right) - \frac{\sqrt{c+dx^2+ex}(2x(ad+2bc)+ae)}{x}}{4c} - \frac{a(c+dx^2+ex)^{3/2}}{2cx^2} \right)}{a + bx^2}$$

↓ 1154

---

3.41.  $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx$

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4}}{a + bx^2} \left( -\frac{\frac{1}{2} \left( 2(4acd - ae^2 + 8bc^2) \int \frac{1}{4c - \frac{(2c+ex)^2}{dx^2+ex+c}} d \frac{2c+ex}{\sqrt{dx^2+ex+c}} - \frac{4bce \operatorname{arctanh}\left(\frac{2dx+e}{2\sqrt{d}\sqrt{c+dx^2+ex}}\right)}{\sqrt{d}} \right)}{4c} - \frac{\sqrt{c+dx^2+ex}(2x(ad+2bc)+a)}{x} \right)$$

↓ 219

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4}}{a + bx^2} \left( -\frac{\frac{1}{2} \left( \frac{(4acd - ae^2 + 8bc^2) \operatorname{arctanh}\left(\frac{2c+ex}{2\sqrt{c}\sqrt{c+dx^2+ex}}\right) - \frac{4bce \operatorname{arctanh}\left(\frac{2dx+e}{2\sqrt{d}\sqrt{c+dx^2+ex}}\right)}{\sqrt{d}} \right)}{\sqrt{c}} - \frac{\sqrt{c+dx^2+ex}(2x(ad+2bc)+a)}{x} \right)$$

input `Int[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^3,x]`

output `(Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4]*(-1/2*(a*(c + e*x + d*x^2)^(3/2))/(c*x^2) - (-(((a*e + 2*(2*b*c + a*d)*x)*Sqrt[c + e*x + d*x^2])/x) + ((-4*b*c*e*ArcTanh[(e + 2*d*x)/(2*Sqrt[d]*Sqrt[c + e*x + d*x^2])])/Sqrt[d] + ((8*b*c^2 + 4*a*c*d - a*e^2)*ArcTanh[(2*c + e*x)/(2*Sqrt[c]*Sqrt[c + e*x + d*x^2])])/Sqrt[c])/2)/(4*c))/(a + b*x^2)`

### 3.41.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1092 `Int[1/Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] := Simp[2 Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x]`

rule 1154 `Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] := Simp[-2 Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 1230 `Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*(e*f*(m + 2*p + 2) - d*g*(2*p + 1) + e*g*(m + 1)*x)*((a + b*x + c*x^2)^p/(e^2*(m + 1)*(m + 2*p + 2))), x] + Simp[p/(e^2*(m + 1)*(m + 2*p + 2)) Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^(p - 1)*Simp[g*(b*d + 2*a*e + 2*a*e*m + 2*b*d*p) - f*b*e*(m + 2*p + 2) + (g*(2*c*d + b*e + b*e*m + 4*c*d*p) - 2*c*e*f*(m + 2*p + 2))*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && GtQ[p, 0] && (LtQ[m, -1] || EqQ[p, 1] || (IntegerQ[p] && !RationalQ[m])) && NeQ[m, -1] && !ILtQ[m + 2*p + 1, 0] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])`

rule 1269 `Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[g/e Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Simp[(e*f - d*g)/e Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && !IGtQ[m, 0]`

rule 1384 `Int[(u_.)*((a_) + (c_.)*(x_)^(n2_.) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n + c*x^(2*n))^FracPart[p]/(c^IntPart[p]*(b/2 + c*x^n)^(2*FracPart[p])) Int[u*(b/2 + c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p - 1/2] && NeQ[u, x^(n - 1)] && NeQ[u, x^(2*n - 1)] && !(EqQ[p, 1/2] && EqQ[u, x^(-2*n - 1)])`

rule 2181 `Int[(Pq_)*((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{Qx = PolynomialQuotient[Pq, d + e*x, x], R = PolynomialRemainder[Pq, d + e*x, x]}, Simp[(e*R*(d + e*x)^(m + 1)*(a + b*x + c*x^2)^(p + 1))/((m + 1)*(c*d^2 - b*d*e + a*e^2)), x] + Simp[1/((m + 1)*(c*d^2 - b*d*e + a*e^2)) Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p*ExpandToSum[(m + 1)*(c*d^2 - b*d*e + a*e^2)*Qx + c*d*R*(m + 1) - b*e*R*(m + p + 2) - c*e*R*(m + 2*p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && LtQ[m, -1]`



### 3.41.4 Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 214, normalized size of antiderivative = 0.74

| method  | result   |
|---------|--|
| risch   | $-\frac{a\sqrt{dx^2+ex+c}(ex+2c)\sqrt{(bx^2+a)^2}}{4x^2c(bx^2+a)} + \left( \frac{8ebc \ln\left(\frac{\frac{5}{2}+dx}{\sqrt{d}} + \sqrt{dx^2+ex+c}\right)}{\sqrt{d}} + 8bcd \left( \frac{\sqrt{dx^2+ex+c}}{d} - \frac{e \ln\left(\frac{\frac{5}{2}+dx}{\sqrt{d}} + \sqrt{dx^2+ex+c}\right)}{2d^{\frac{3}{2}}}\right) \right) - \frac{(4acd)}{8c(bx^2+a)}$ |
| default | $\frac{\sqrt{(bx^2+a)^2} \left( -4d^{\frac{5}{2}}c^{\frac{3}{2}} \ln\left(\frac{2c+ex+2\sqrt{c}\sqrt{dx^2+ex+c}}{x}\right) ax^2 - 8d^{\frac{3}{2}}c^{\frac{5}{2}} \ln\left(\frac{2c+ex+2\sqrt{c}\sqrt{dx^2+ex+c}}{x}\right) bx^2 - 2d^{\frac{5}{2}}\sqrt{dx^2+ex+c} aex^3 + 4d^{\frac{5}{2}}\sqrt{dx^2+ex+c} \right)}{8c(bx^2+a)}$                       |

input `int((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^3,x,method=_RETURNVERBOSE)`

output 
$$-1/4*a*(d*x^2+e*x+c)^(1/2)*(e*x+2*c)/x^2/c*((b*x^2+a)^2)^(1/2)/(b*x^2+a)+1/8/c*(8*e*b*c*\ln((1/2*e+d*x)/d^(1/2)+(d*x^2+e*x+c)^(1/2))/d^(1/2)+8*b*c*d*(1/d*(d*x^2+e*x+c)^(1/2)-1/2*e/d^(3/2)*\ln((1/2*e+d*x)/d^(1/2)+(d*x^2+e*x+c)^(1/2)))-(4*a*c*d-a*e^2+8*b*c^2)/c^(1/2)*\ln((2*c+e*x+2*c^(1/2)*(d*x^2+e*x+c)^(1/2))/x))*((b*x^2+a)^2)^(1/2)/(b*x^2+a)$$

### 3.41.5 Fracas [A] (verification not implemented)

Time = 0.59 (sec) , antiderivative size = 749, normalized size of antiderivative = 2.60

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx$$

$$= \frac{4bc^2\sqrt{dex^2} \log\left(8d^2x^2+8dex+4\sqrt{dx^2+ex+c}(2dx+e)\sqrt{d}+4cd+e^2\right) - (8bc^2d+4acd^2-ade^2)}{16c^2dx^2} + \frac{8bc^2\sqrt{-dex^2} \arctan\left(\frac{\sqrt{dx^2+ex+c}(2dx+e)\sqrt{-d}}{2(d^2x^2+dex+cd)}\right) + (8bc^2d+4acd^2-ade^2)\sqrt{cx^2} \log\left(\frac{8cex+(4cd+e^2)x^2+4\sqrt{dx^2+ex+c}}{x^2}\right)}{16c^2dx^2} - \frac{4bc^2\sqrt{-dex^2} \arctan\left(\frac{\sqrt{dx^2+ex+c}(2dx+e)\sqrt{-d}}{2(d^2x^2+dex+cd)}\right) - (8bc^2d+4acd^2-ade^2)\sqrt{-cx^2} \arctan\left(\frac{\sqrt{dx^2+ex+c}(ex+2c)}{2(cd^2+ce^2+c^2)}\right)}{8c^2dx^2}$$

input `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^3,x, algorithm="fracas")`

output `[1/16*(4*b*c^2*sqrt(d)*e*x^2*log(8*d^2*x^2 + 8*d*e*x + 4*sqrt(d*x^2 + e*x + c))*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) - (8*b*c^2*d + 4*a*c*d^2 - a*d*e^2)*sqrt(c)*x^2*log((8*c*e*x + (4*c*d + e^2)*x^2 + 4*sqrt(d*x^2 + e*x + c))*(e*x + 2*c)*sqrt(c) + 8*c^2)/x^2) + 4*(4*b*c^2*d*x^2 - a*c*d*e*x - 2*a*c^2*d)*sqrt(d*x^2 + e*x + c)/(c^2*d*x^2), -1/16*(8*b*c^2*sqrt(-d)*e*x^2*arctan(1/2*sqrt(d*x^2 + e*x + c))*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) + (8*b*c^2*d + 4*a*c*d^2 - a*d*e^2)*sqrt(c)*x^2*log((8*c*e*x + (4*c*d + e^2)*x^2 + 4*sqrt(d*x^2 + e*x + c))*(e*x + 2*c)*sqrt(c) + 8*c^2)/x^2) - 4*(4*b*c^2*d*x^2 - a*c*d*e*x - 2*a*c^2*d)*sqrt(d*x^2 + e*x + c)/(c^2*d*x^2), 1/8*(2*b*c^2*sqrt(d)*e*x^2*log(8*d^2*x^2 + 8*d*e*x + 4*sqrt(d*x^2 + e*x + c))*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + (8*b*c^2*d + 4*a*c*d^2 - a*d*e^2)*sqrt(-c)*x^2*arctan(1/2*sqrt(d*x^2 + e*x + c))*(e*x + 2*c)*sqrt(-c)/(c*d*x^2 + c*e*x + c^2)) + 2*(4*b*c^2*d*x^2 - a*c*d*e*x - 2*a*c^2*d)*sqrt(d*x^2 + e*x + c)/(c^2*d*x^2), -1/8*(4*b*c^2*sqrt(-d)*e*x^2*arctan(1/2*sqrt(d*x^2 + e*x + c))*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) - (8*b*c^2*d + 4*a*c*d^2 - a*d*e^2)*sqrt(-c)*x^2*arctan(1/2*sqrt(d*x^2 + e*x + c))*(e*x + 2*c)*sqrt(-c)/(c*d*x^2 + c*e*x + c^2)) - 2*(4*b*c^2*d*x^2 - a*c*d*e*x - 2*a*c^2*d)*sqrt(d*x^2 + e*x + c)/(c^2*d*x^2)]`

### 3.41.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx = \text{Timed out}$$

input `integrate((d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2)/x**3,x)`

output `Timed out`

### 3.41.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx = \text{Exception raised: ValueError}$$

input `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^3,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation \*may\* help (example of legal syntax is 'assume(e^2-4\*c\*d>0)', see 'assume?' for more deta

### 3.41.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 360, normalized size of antiderivative = 1.25

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx$$

$$= -\frac{be \log\left(\left|-2\left(\sqrt{dx}-\sqrt{dx^2+ex+c}\right)\sqrt{d}-e\right|\right) \operatorname{sgn}(bx^2+a)}{2\sqrt{d}} + \sqrt{dx^2+ex+c} b \operatorname{sgn}(bx^2+a)$$

$$+ \frac{(8bc^2 \operatorname{sgn}(bx^2+a) + 4ac d \operatorname{sgn}(bx^2+a) - ae^2 \operatorname{sgn}(bx^2+a)) \arctan\left(-\frac{\sqrt{dx}-\sqrt{dx^2+ex+c}}{\sqrt{-c}}\right)}{4\sqrt{-c}}$$

$$+ \frac{4\left(\sqrt{dx}-\sqrt{dx^2+ex+c}\right)^3 ac d \operatorname{sgn}(bx^2+a) + \left(\sqrt{dx}-\sqrt{dx^2+ex+c}\right)^3 ae^2 \operatorname{sgn}(bx^2+a) + 8\left(\sqrt{dx}-\sqrt{dx^2+ex+c}\right)^2 ac \sqrt{d} \operatorname{sgn}(bx^2+a) + 4\left(\sqrt{d}x - \sqrt{dx^2+ex+c}\right) ac^2 d \operatorname{sgn}(bx^2+a) + \left(\sqrt{d}x - \sqrt{dx^2+ex+c}\right) ac^2 e^2 \operatorname{sgn}(bx^2+a)}{\left(\left(\sqrt{d}x - \sqrt{dx^2+ex+c}\right)^2 - c\right)^2 c}$$

input `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^3,x, algorithm="giac")`

output `-1/2*b*e*log(abs(-2*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))*sqrt(d) - e))*sgn(b*x^2 + a)/sqrt(d) + sqrt(d*x^2 + e*x + c)*b*sgn(b*x^2 + a) + 1/4*(8*b*c^2*sgn(b*x^2 + a) + 4*a*c*d*sgn(b*x^2 + a) - a*e^2*sgn(b*x^2 + a))*arctan(-(sqrt(d)*x - sqrt(d*x^2 + e*x + c))/sqrt(-c))/(sqrt(-c)*c) + 1/4*(4*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))^3*a*c*d*sgn(b*x^2 + a) + (sqrt(d)*x - sqrt(d*x^2 + e*x + c))^3*a*e^2*sgn(b*x^2 + a) + 8*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))^2*a*c*sqrt(d)*e*sgn(b*x^2 + a) + 4*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))*a*c^2*d*sgn(b*x^2 + a) + (sqrt(d)*x - sqrt(d*x^2 + e*x + c))*a*c^2*e^2*sgn(b*x^2 + a))/(((sqrt(d)*x - sqrt(d*x^2 + e*x + c))^2 - c)^2*c)`

**3.41.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx = \int \frac{\sqrt{(bx^2+a)^2}\sqrt{dx^2+ex+c}}{x^3} dx$$

input `int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^3,x)`output `int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^3, x)`

**3.42**  $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^4} dx$

3.42.1 Optimal result . . . . . 340  
 3.42.2 Mathematica [A] (verified) . . . . . 341  
 3.42.3 Rubi [A] (verified) . . . . . 341  
 3.42.4 Maple [A] (verified) . . . . . 345  
 3.42.5 Fricas [A] (verification not implemented) . . . . . 345  
 3.42.6 Sympy [F(-1)] . . . . . 346  
 3.42.7 Maxima [F(-2)] . . . . . 346  
 3.42.8 Giac [B] (verification not implemented) . . . . . 347  
 3.42.9 Mupad [F(-1)] . . . . . 348

**3.42.1 Optimal result**

Integrand size = 40, antiderivative size = 294

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^4} dx$$

$$= \frac{(2ace - (8bc^2 - ae^2)x)\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{8c^2x^2(a+bx^2)}$$

$$- \frac{a(c+ex+dx^2)^{3/2}\sqrt{a^2+2abx^2+b^2x^4}}{3cx^3(a+bx^2)}$$

$$+ \frac{b\sqrt{d}\sqrt{a^2+2abx^2+b^2x^4}\operatorname{arctanh}\left(\frac{e+2dx}{2\sqrt{d}\sqrt{c+ex+dx^2}}\right)}{a+bx^2}$$

$$- \frac{e(8bc^2 - a(4cd - e^2))\sqrt{a^2+2abx^2+b^2x^4}\operatorname{arctanh}\left(\frac{2c+ex}{2\sqrt{c}\sqrt{c+ex+dx^2}}\right)}{16c^{5/2}(a+bx^2)}$$

output

```
-1/3*a*(d*x^2+e*x+c)^(3/2)*((b*x^2+a)^2)^(1/2)/c/x^3/(b*x^2+a)-1/16*e*(8*b
*c^2-a*(4*c*d-e^2))*arctanh(1/2*(e*x+2*c)/c^(1/2)/(d*x^2+e*x+c)^(1/2))*((b
*x^2+a)^2)^(1/2)/c^(5/2)/(b*x^2+a)+b*arctanh(1/2*(2*d*x+e)/d^(1/2)/(d*x^2+
e*x+c)^(1/2))*d^(1/2)*((b*x^2+a)^2)^(1/2)/(b*x^2+a)+1/8*(2*a*c*e-(-a*e^2+8
*b*c^2)*x)*(d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/c^2/x^2/(b*x^2+a)
```

### 3.42.2 Mathematica [A] (verified)

Time = 0.88 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.63

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^4} dx$$

$$= \frac{\sqrt{(a+bx^2)^2} \left( -3e(8bc^2 + a(-4cd + e^2)) x^3 \operatorname{arctanh}\left(\frac{-\sqrt{dx+\sqrt{c+x(e+dx)}}}{\sqrt{c}}\right) - \sqrt{c} \left( \sqrt{c+x(e+dx)}(24bc^2x^2 + 24c^{5/2}x^3(a+bx^2) \right) \right)}{24c^{5/2}x^3(a+bx^2)}$$

input `Integrate[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^4,x]`

output `(Sqrt[(a + b*x^2)^2]*(-3*e*(8*b*c^2 + a*(-4*c*d + e^2))*x^3*ArcTanh[(-(Sqrt[d]*x) + Sqrt[c + x*(e + d*x)])/Sqrt[c]] - Sqrt[c]*(Sqrt[c + x*(e + d*x)]*(24*b*c^2*x^2 + a*(8*c^2 - 3*e^2*x^2 + 2*c*x*(e + 4*d*x))) + 24*b*c^2*Sqrt[d]*x^3*Log[e + 2*d*x - 2*Sqrt[d]*Sqrt[c + x*(e + d*x)]]))/(24*c^(5/2)*x^3*(a + b*x^2))`

### 3.42.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 220, normalized size of antiderivative = 0.75, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.275$ , Rules used = {1384, 27, 2181, 27, 1229, 27, 1269, 1092, 219, 1154, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a^2+2abx^2+b^2x^4}\sqrt{c+dx^2+ex}}{x^4} dx$$

$$\downarrow \text{1384}$$

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{b(bx^2+a)\sqrt{dx^2+ex+c}}{x^4} dx}{b(a+bx^2)}$$

$$\downarrow \text{27}$$

$$\frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{(bx^2+a)\sqrt{dx^2+ex+c}}{x^4} dx}{a+bx^2}$$

$$\downarrow \text{2181}$$

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( -\frac{\int \frac{3(ae-2bcx)\sqrt{dx^2+ex+c}}{2x^3} dx}{3c} - \frac{a(c+dx^2+ex)^{3/2}}{3cx^3} \right)}{a + bx^2}$$

↓ 27

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( -\frac{\int \frac{(ae-2bcx)\sqrt{dx^2+ex+c}}{x^3} dx}{2c} - \frac{a(c+dx^2+ex)^{3/2}}{3cx^3} \right)}{a + bx^2}$$

↓ 1229

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( -\frac{\int \frac{16bdxc^2+e(8bc^2-a(4cd-e^2))}{2x\sqrt{dx^2+ex+c}} dx}{4c} - \frac{\sqrt{c+dx^2+ex}(2ace-x(8bc^2-ae^2))}{4cx^2} - \frac{a(c+dx^2+ex)^{3/2}}{3cx^3} \right)}{a + bx^2}$$

↓ 27

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( -\frac{\int \frac{16bdxc^2+e(8bc^2-a(4cd-e^2))}{x\sqrt{dx^2+ex+c}} dx}{8c} - \frac{\sqrt{c+dx^2+ex}(2ace-x(8bc^2-ae^2))}{4cx^2} - \frac{a(c+dx^2+ex)^{3/2}}{3cx^3} \right)}{a + bx^2}$$

↓ 1269

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( -\frac{e(8bc^2-a(4cd-e^2)) \int \frac{1}{x\sqrt{dx^2+ex+c}} dx + 16bc^2 d \int \frac{1}{\sqrt{dx^2+ex+c}} dx}{2c} - \frac{\sqrt{c+dx^2+ex}(2ace-x(8bc^2-ae^2))}{4cx^2} - \frac{a(c+dx^2+ex)^{3/2}}{3cx^3} \right)}{a + bx^2}$$

↓ 1092

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( -\frac{e(8bc^2-a(4cd-e^2)) \int \frac{1}{x\sqrt{dx^2+ex+c}} dx + 32bc^2 d \int \frac{1}{4d - \frac{(e+2dx)^2}{dx^2+ex+c}} d \frac{e+2dx}{\sqrt{dx^2+ex+c}}}{8c} - \frac{\sqrt{c+dx^2+ex}(2ace-x(8bc^2-ae^2))}{4cx^2} \right)}{a + bx^2}$$

↓ 219

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \left( -\frac{e(8bc^2-a(4cd-e^2)) \int \frac{1}{x\sqrt{dx^2+ex+c}} dx + 16bc^2 \sqrt{d} \operatorname{arctanh} \left( \frac{2dx+e}{2\sqrt{d}\sqrt{c+dx^2+ex}} \right)}{8c} - \frac{\sqrt{c+dx^2+ex}(2ace-x(8bc^2-ae^2))}{4cx^2} \right)}{a + bx^2}$$

---

3.42.  $\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^4} dx$

$$\begin{array}{c} \downarrow 1154 \\ \sqrt{a^2 + 2abx^2 + b^2x^4} \left( -\frac{16bc^2\sqrt{d}\operatorname{arctanh}\left(\frac{2dx+e}{2\sqrt{d}\sqrt{c+dx^2+ex}}\right) - 2e(8bc^2 - a(4cd - e^2))}{8c} \int \frac{1}{4c - \frac{(2c+ex)^2}{dx^2+ex+c}} d \frac{2c+ex}{\sqrt{dx^2+ex+c}} - \frac{\sqrt{c+dx^2+ex}(2ace - x(8bc^2 - a(4cd - e^2)))}{4cx^2} \right) \\ \hline a + bx^2 \\ \downarrow 219 \\ \sqrt{a^2 + 2abx^2 + b^2x^4} \left( -\frac{16bc^2\sqrt{d}\operatorname{arctanh}\left(\frac{2dx+e}{2\sqrt{d}\sqrt{c+dx^2+ex}}\right) - \frac{e(8bc^2 - a(4cd - e^2))\operatorname{arctanh}\left(\frac{2c+ex}{2\sqrt{c}\sqrt{c+dx^2+ex}}\right)}{\sqrt{c}}}{8c} - \frac{\sqrt{c+dx^2+ex}(2ace - x(8bc^2 - a(4cd - e^2)))}{4cx^2} \right) \\ \hline a + bx^2 \end{array}$$

input `Int[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^4,x]`

output `(Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4]*(-1/3*(a*(c + e*x + d*x^2)^(3/2))/(c*x^3) - (-1/4*((2*a*c*e - (8*b*c^2 - a*e^2)*x)*Sqrt[c + e*x + d*x^2])/(c*x^2) - (16*b*c^2*Sqrt[d]*ArcTanh[(e + 2*d*x)/(2*Sqrt[d]*Sqrt[c + e*x + d*x^2])]) - (e*(8*b*c^2 - a*(4*c*d - e^2))*ArcTanh[(2*c + e*x)/(2*Sqrt[c]*Sqrt[c + e*x + d*x^2])]))/Sqrt[c])/(8*c))/(2*c))/(a + b*x^2)`

### 3.42.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1092 `Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Simp[2 Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x]`



rule 1154 `Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] := Simp[-2 Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 1229 `Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(- (d + e*x)^(m + 1))*((a + b*x + c*x^2)^p/(e^2*(m + 1)*(m + 2)*(c*d^2 - b*d*e + a*e^2)))*((d*g - e*f*(m + 2))*(c*d^2 - b*d*e + a*e^2) - d*p*(2*c*d - b*e)*(e*f - d*g) - e*(g*(m + 1)*(c*d^2 - b*d*e + a*e^2) + p*(2*c*d - b*e)*(e*f - d*g))*x), x] - Simp[p/(e^2*(m + 1)*(m + 2)*(c*d^2 - b*d*e + a*e^2)) Int[(d + e*x)^(m + 2)*(a + b*x + c*x^2)^(p - 1)*Simp[2*a*c*e*(e*f - d*g)*(m + 2) + b^2*e*(d*g*(p + 1) - e*f*(m + p + 2)) + b*(a*e^2*g*(m + 1) - c*d*(d*g*(2*p + 1) - e*f*(m + 2*p + 2))) - c*(2*c*d*(d*g*(2*p + 1) - e*f*(m + 2*p + 2)) - e*(2*a*e*g*(m + 1) - b*(d*g*(m - 2*p) + e*f*(m + 2*p + 2)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && GtQ[p, 0] && LtQ[m, -2] && LtQ[m + 2*p, 0] && !ILtQ[m + 2*p + 3, 0]`

rule 1269 `Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[g/e Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Simp[(e*f - d*g)/e Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && !IGtQ[m, 0]`

rule 1384 `Int[(u_.)*((a_) + (c_.)*(x_)^(n2_.) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n + c*x^(2*n))^FracPart[p]/(c^IntPart[p]*(b/2 + c*x^n)^(2*FracPart[p])) Int[u*(b/2 + c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p - 1/2] && NeQ[u, x^(n - 1)] && NeQ[u, x^(2*n - 1)] && !(EqQ[p, 1/2] && EqQ[u, x^(-2*n - 1)])`

rule 2181 `Int[(Pq_)*((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{Qx = PolynomialQuotient[Pq, d + e*x, x], R = PolynomialRemainder[Pq, d + e*x, x]}, Simp[(e*R*(d + e*x)^(m + 1)*(a + b*x + c*x^2)^(p + 1))/((m + 1)*(c*d^2 - b*d*e + a*e^2)), x] + Simp[1/((m + 1)*(c*d^2 - b*d*e + a*e^2)) Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p*ExpandToSum[(m + 1)*(c*d^2 - b*d*e + a*e^2)*Qx + c*d*R*(m + 1) - b*e*R*(m + p + 2) - c*e*R*(m + 2*p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && LtQ[m, -1]`

### 3.42.4 Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.66

| method  | result   |
|---------|--|
| risch   | $-\frac{\sqrt{dx^2+ex+c}(8acd^2x^2-3ae^2x^2+24b^2c^2x^2+2acex+8a^2c^2)\sqrt{(bx^2+a)^2}}{24x^3c^2(bx^2+a)} - \frac{\left(-16bc^2\sqrt{d}\ln\left(\frac{\frac{e}{2}+dx}{\sqrt{d}}+\sqrt{dx^2+ex+c}\right)-\frac{e^{(4acd-ae^2)}}{16c^2}\right)}{16c^2(bx^2+a)}$                               |
| default | $\frac{\sqrt{(bx^2+a)^2}\left(12d^{\frac{5}{2}}c^{\frac{3}{2}}\ln\left(\frac{2c+ex+2\sqrt{c}\sqrt{dx^2+ex+c}}{x}\right)ae^x-24d^{\frac{3}{2}}c^{\frac{5}{2}}\ln\left(\frac{2c+ex+2\sqrt{c}\sqrt{dx^2+ex+c}}{x}\right)be^x+6d^{\frac{5}{2}}\sqrt{dx^2+ex+c}ae^2x^4+48\right)}{16c^2(bx^2+a)}$ |

input `int((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^4,x,method=_RETURNVERBOSE)`

output 
$$-1/24*(d*x^2+e*x+c)^(1/2)*(8*a*c*d*x^2-3*a*e^2*x^2+24*b*c^2*x^2+2*a*c*e*x+8*a*c^2)/x^3/c^2*((b*x^2+a)^2)^(1/2)/(b*x^2+a)-1/16/c^2*(-16*b*c^2*d^(1/2)*\ln((1/2*e+d*x)/d^(1/2)+(d*x^2+e*x+c)^(1/2))-e*(4*a*c*d-a*e^2-8*b*c^2)/c^(1/2)*\ln((2*c+e*x+2*c^(1/2)*(d*x^2+e*x+c)^(1/2))/x))*((b*x^2+a)^2)^(1/2)/(b*x^2+a)$$

### 3.42.5 Fracas [A] (verification not implemented)

Time = 0.59 (sec) , antiderivative size = 791, normalized size of antiderivative = 2.69

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^4} dx$$

$$= \frac{48bc^3\sqrt{dx^3}\log\left(8d^2x^2+8dex+4\sqrt{dx^2+ex+c}(2dx+e)\sqrt{d}+4cd+e^2\right)+3(ae^3+4(2bc^2-acd)e)}{96bc^3\sqrt{-dx^3}\arctan\left(\frac{\sqrt{dx^2+ex+c}(2dx+e)\sqrt{-d}}{2(d^2x^2+dex+cd)}\right)-3(ae^3+4(2bc^2-acd)e)\sqrt{cx^3}\log\left(\frac{8cex+(4cd+e^2)x^2-4\sqrt{dx^2+ex+c}}{x^2}\right)}{48c^3x^3}$$

$$48bc^3\sqrt{-dx^3}\arctan\left(\frac{\sqrt{dx^2+ex+c}(2dx+e)\sqrt{-d}}{2(d^2x^2+dex+cd)}\right)-3(ae^3+4(2bc^2-acd)e)\sqrt{-cx^3}\arctan\left(\frac{\sqrt{dx^2+ex+c}(ex+2c)}{2(cd^2x^2+ce^2x+c^2)}\right)$$

input `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^4,x,algorithm="fracas")`

output

```
[1/96*(48*b*c^3*sqrt(d)*x^3*log(8*d^2*x^2 + 8*d*e*x + 4*sqrt(d*x^2 + e*x +
c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 3*(a*e^3 + 4*(2*b*c^2 - a*c*d)*e)
*sqrt(c)*x^3*log((8*c*e*x + (4*c*d + e^2)*x^2 - 4*sqrt(d*x^2 + e*x + c)*(e
*x + 2*c)*sqrt(c) + 8*c^2)/x^2) - 4*(2*a*c^2*e*x + 8*a*c^3 + (24*b*c^3 + 8
*a*c^2*d - 3*a*c*e^2)*x^2)*sqrt(d*x^2 + e*x + c))/(c^3*x^3), -1/96*(96*b*c
^3*sqrt(-d)*x^3*arctan(1/2*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(-d)/(d^2
*x^2 + d*e*x + c*d)) - 3*(a*e^3 + 4*(2*b*c^2 - a*c*d)*e)*sqrt(c)*x^3*log((
8*c*e*x + (4*c*d + e^2)*x^2 - 4*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt(c)
+ 8*c^2)/x^2) + 4*(2*a*c^2*e*x + 8*a*c^3 + (24*b*c^3 + 8*a*c^2*d - 3*a*c*e
^2)*x^2)*sqrt(d*x^2 + e*x + c))/(c^3*x^3), 1/48*(24*b*c^3*sqrt(d)*x^3*log(
8*d^2*x^2 + 8*d*e*x + 4*sqrt(d*x^2 + e*x + c)*(2*d*x + e)*sqrt(d) + 4*c*d
+ e^2) + 3*(a*e^3 + 4*(2*b*c^2 - a*c*d)*e)*sqrt(-c)*x^3*arctan(1/2*sqrt(d*
x^2 + e*x + c)*(e*x + 2*c)*sqrt(-c)/(c*d*x^2 + c*e*x + c^2)) - 2*(2*a*c^2*
e*x + 8*a*c^3 + (24*b*c^3 + 8*a*c^2*d - 3*a*c*e^2)*x^2)*sqrt(d*x^2 + e*x +
c))/(c^3*x^3), -1/48*(48*b*c^3*sqrt(-d)*x^3*arctan(1/2*sqrt(d*x^2 + e*x +
c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*e*x + c*d)) - 3*(a*e^3 + 4*(2*b*c^2
- a*c*d)*e)*sqrt(-c)*x^3*arctan(1/2*sqrt(d*x^2 + e*x + c)*(e*x + 2*c)*sqrt
(-c)/(c*d*x^2 + c*e*x + c^2)) + 2*(2*a*c^2*e*x + 8*a*c^3 + (24*b*c^3 + 8*a
*c^2*d - 3*a*c*e^2)*x^2)*sqrt(d*x^2 + e*x + c))/(c^3*x^3)]
```

### 3.42.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^4} dx = \text{Timed out}$$

input `integrate((d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2)/x**4,x)`

output Timed out

### 3.42.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^4} dx = \text{Exception raised: ValueError}$$

input `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^4,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation \*may\* help (example of legal syntax is 'assume(e^2-4\*c\*d>0)', see `assume?` for more deta

### 3.42.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 683 vs.  $2(224) = 448$ .

Time = 0.35 (sec) , antiderivative size = 683, normalized size of antiderivative = 2.32

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^4} dx$$

$$= -b\sqrt{d} \log \left( \left| -2 \left( \sqrt{dx} - \sqrt{dx^2+ex+c} \right) \sqrt{d} - e \right| \right) \operatorname{sgn}(bx^2+a)$$

$$+ \frac{(8bc^2 \operatorname{esgn}(bx^2+a) - 4ac \operatorname{desgn}(bx^2+a) + ae^3 \operatorname{sgn}(bx^2+a)) \arctan \left( -\frac{\sqrt{dx} - \sqrt{dx^2+ex+c}}{\sqrt{-c}} \right)}{8\sqrt{-cc^2}}$$

$$+ \frac{24 \left( \sqrt{dx} - \sqrt{dx^2+ex+c} \right)^5 bc^2 \operatorname{esgn}(bx^2+a) + 12 \left( \sqrt{dx} - \sqrt{dx^2+ex+c} \right)^5 ac \operatorname{desgn}(bx^2+a) - 3 \left( \sqrt{dx} - \sqrt{dx^2+ex+c} \right)^5}{8\sqrt{-cc^2}}$$

input `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^4,x, algorithm="giac")`

output

```
-b*sqrt(d)*log(abs(-2*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))*sqrt(d) - e))*sgn(b*x^2 + a) + 1/8*(8*b*c^2*e*sgn(b*x^2 + a) - 4*a*c*d*e*sgn(b*x^2 + a) + a*e^3*sgn(b*x^2 + a))*arctan(-(sqrt(d)*x - sqrt(d*x^2 + e*x + c))/sqrt(-c))/(sqrt(-c)*c^2) + 1/24*(24*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))^5*b*c^2*e*sgn(b*x^2 + a) + 12*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))^5*a*c*d*e*sgn(b*x^2 + a) - 3*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))^5*a*e^3*sgn(b*x^2 + a) + 48*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))^4*b*c^3*sqrt(d)*sgn(b*x^2 + a) + 48*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))^4*a*c^2*d^(3/2)*sgn(b*x^2 + a) - 48*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))^3*b*c^3*e*sgn(b*x^2 + a) + 48*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))^3*a*c^2*d*e*sgn(b*x^2 + a) + 8*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))^3*a*c*e^3*sgn(b*x^2 + a) - 96*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))^2*b*c^4*sqrt(d)*sgn(b*x^2 + a) + 48*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))^2*a*c^2*sqrt(d)*e^2*sgn(b*x^2 + a) + 24*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))*b*c^4*e*sgn(b*x^2 + a) + 36*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))*a*c^3*d*e*sgn(b*x^2 + a) + 3*(sqrt(d)*x - sqrt(d*x^2 + e*x + c))*a*c^2*e^3*sgn(b*x^2 + a) + 48*b*c^5*sqrt(d)*sgn(b*x^2 + a) + 16*a*c^4*d^(3/2)*sgn(b*x^2 + a)/(((sqrt(d)*x - sqrt(d*x^2 + e*x + c))^2 - c)^3*c^2)
```

### 3.42.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{c+ex+dx^2}\sqrt{a^2+2abx^2+b^2x^4}}{x^4} dx = \int \frac{\sqrt{(bx^2+a)^2}\sqrt{dx^2+ex+c}}{x^4} dx$$

input `int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^4,x)`

output `int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^4, x)`

## APPENDIX

|  |     |
|--|-----|
| 4.1 Listing of Grading functions . . . . . | 349 |
|--|-----|

## 4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

### 4.1.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
```

```

(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A"," "}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A"," "}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal.$
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
      ,
      finalresult={"F","Contains unresolved integral."}
    ]
  ];

  finalresult
]

```

```

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType,expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]],2]],
            Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3,ExpnType[expn[[1]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
            If[Head[expn]===RootSum,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
            9]]]]]]]]]]

```



```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, CsCh,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsCh
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

### 4.1.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
      return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
            print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
      return "F","Result contains unresolved integral";
fi;

```

```

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
  if is_contains_complex(result) then
    if is_contains_complex(optimal) then
      if debug then
        print("both result and optimal complex");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        return "A"," ";
      else
        return "B",cat("Both result and optimal contain complex but leaf count of
                        convert(leaf_count_result,string)," vs. $2 (" ,
                        convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_
        end if
      else #result contains complex but optimal is not
        if debug then
          print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
      fi;
    else # result do not contain complex
      # this assumes optimal do not as well. No check is needed here.
      if debug then
        print("result do not contain complex, this assumes optimal do not as well");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        if debug then
          print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
      else
        if debug then
          print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string),"$ vs. $2(",
                        convert(leaf_count_optimal,string),"=" ,convert(2*leaf_cou
        fi;
      fi;
    fi;
  fi;

```

```

else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  end if
end if

```

```

elif type(expn, ``~`) then
  if type(op(2,expn), 'integer') then
    ExpnType(op(1,expn))
  elif type(op(2,expn), 'rational') then
    if type(op(1,expn), 'rational') then
      1
    else
      max(2, ExpnType(op(1,expn)))
    end if
  else
    max(3, ExpnType(op(1,expn)), ExpnType(op(2,expn)))
  end if
elif type(expn, ``+`) or type(expn, ``*`) then
  max(ExpnType(op(1,expn)), max(ExpnType(rest(expn))))
elif ElementaryFunctionQ(op(0,expn)) then
  max(3, ExpnType(op(1,expn)))
elif SpecialFunctionQ(op(0,expn)) then
  max(4, apply(max, map(ExpnType, [op(expn)])))
elif HypergeometricFunctionQ(op(0,expn)) then
  max(5, apply(max, map(ExpnType, [op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6, apply(max, map(ExpnType, [op(expn)])))
elif op(0,expn)='int' then
  max(8, apply(max, map(ExpnType, [op(expn)]))) else
  9
end if
end proc:

```

```

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

```

```

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,

```

```

        GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
        EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
    member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
    member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
    if nops(u)=2 then
        op(2,u)
    else
        apply(op(0,u),op(2..nops(u),u))
    end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma] [LeafCount] (u);
end proc:

```

### 4.1.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):

```

```

if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
        return True
    else:
        return False
else:
    return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):

```

```

return 1
elif isinstance(expn,list):
    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
        return 1
    else:
        return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
elif isinstance(expn,Pow): #type(expn,``^`)
    if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
        return expnType(expn.args[0]) #ExpnType(op(1,expn))
    elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or type(expn,``*`)
    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

*#main function*



```

def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

    #print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```

#### 4.1.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
```

```

        return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

```

```

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric', 'hypergeometric_M', 'hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=", expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception, AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list:  #isinstance(expn,list):

```

```

    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
elif expn.operator() == operator.pow: #isinstance(expn,Pow)
    if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
        return expnType(expn.operands()[0]) #expnType(expn.args[0])
    elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
        if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or inst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

```

```

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)",type(result))
    print("type(optimal)",type(optimal))

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal." + str(leaf
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```