

# Computer Algebra Independent Integration Tests

Summer 2023 edition with Rubi V 4.17.3

4-Trig-functions/4.1-Sine/78-4.1.4.2-a+b-sin<sup>m</sup>-c+d-sin<sup>n</sup>-A+B-  
sin+C-sin<sup>2</sup>-

Nasser M. Abbasi

December 9, 2023

Compiled on December 9, 2023 at 5:02am

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>detailed summary tables of results</b>	<b>20</b>
<b>3</b>	<b>Listing of integrals</b>	<b>35</b>
<b>4</b>	<b>Appendix</b>	<b>303</b>

# CHAPTER 1

## INTRODUCTION

1.1	Listing of CAS systems tested . . . . .	3
1.2	Results . . . . .	4
1.3	Time and leaf size Performance . . . . .	7
1.4	Performance based on number of rules Rubi used . . . . .	9
1.5	Performance based on number of steps Rubi used . . . . .	10
1.6	Solved integrals histogram based on leaf size of result . . . . .	11
1.7	Solved integrals histogram based on CPU time used . . . . .	12
1.8	Leaf size vs. CPU time used . . . . .	13
1.9	list of integrals with no known antiderivative . . . . .	14
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	14
1.11	list of integrals solved by CAS but failed verification . . . . .	14
1.12	Timing . . . . .	15
1.13	Verification . . . . .	15
1.14	Important notes about some of the results . . . . .	15
1.15	Design of the test system . . . . .	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [ 34 ]. This is test number [ 78 ].

## 1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 13.3.1 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

## 1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 ( 34 )	0.00 ( 0 )
Mathematica	52.94 ( 18 )	47.06 ( 16 )
Fricas	26.47 ( 9 )	73.53 ( 25 )
Mupad	26.47 ( 9 )	73.53 ( 25 )
Maxima	20.59 ( 7 )	79.41 ( 27 )
Maple	14.71 ( 5 )	85.29 ( 29 )
Giac	8.82 ( 3 )	91.18 ( 31 )
Sympy	2.94 ( 1 )	97.06 ( 33 )

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> <li>1. antiderivative contains a hypergeometric function and the optimal antiderivative does not.</li> <li>2. antiderivative contains a special function and the optimal antiderivative does not.</li> <li>3. antiderivative contains the imaginary unit and the optimal antiderivative does not.</li> </ol>
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	97.059	0.000	0.000	2.941
Mathematica	44.118	0.000	5.882	50.000
Fricas	14.706	5.882	2.941	76.471
Giac	8.824	0.000	0.000	91.176
Maple	2.941	8.824	0.000	88.235
Maxima	2.941	17.647	0.000	79.412
Mupad	0.000	23.529	0.000	76.471
Sympy	0.000	2.941	0.000	97.059

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima

and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	16	100.00	0.00	0.00
Fricas	25	100.00	0.00	0.00
Mupad	25	0.00	100.00	0.00
Maxima	27	96.30	3.70	0.00
Maple	29	100.00	0.00	0.00
Giac	31	77.42	9.68	12.90
Sympy	33	51.52	48.48	0.00

Table 1.4: Failure statistics for each CAS

## 1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.



System	Mean time (sec)
Sympy	0.14
Maxima	0.36
Fricas	0.42
Giac	0.42
Rubi	1.12
Maple	2.74
Mathematica	16.34
Mupad	25.79

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Giac	164.33	1.13	197.00	1.18
Sympy	189.00	2.33	189.00	2.33
Mathematica	252.56	1.07	169.50	0.93
Rubi	302.32	1.05	309.00	1.06
Maple	361.80	2.58	396.00	3.24
Fricas	407.33	1.63	310.00	1.62
Mupad	541.22	2.00	510.00	2.45
Maxima	714.43	2.51	648.00	2.45

Table 1.6: Leaf size performance for each CAS

## 1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the  $y$  axis is the percentage solved which Rubi itself needed the number of rules given the  $x$  axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

# 1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

## 1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

## 1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

## 1.8 Leaf size vs. CPU time used

The following gives the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time.




Figure 1.5: Leaf size vs. CPU time. Full range

## 1.9 list of integrals with no known antiderivative

{34}

## 1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

## 1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

**Giac** Verification phase not currently implemented.

**Mupad** Verification phase not currently implemented.

## 1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

## 1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

## 1.14 Important notes about some of the results

### 1.14.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.



The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

### 1.14.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

### 1.14.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at [https://ask.sagemath.org/question/57123/could-we-have-a-leaf\\_count-function-in-base-sagemath/](https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/)

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

### 1.14.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand,the_variable)
```

Which gives  $\sin(x)^2/2$

## 1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer, 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer, Leaf size of result.
4. integer, Leaf size of the optimal antiderivative.
5. number, CPU time used to solve this integral. 0 if failed.
6. string, The integral in Latex format
7. string, The input used in CAS own syntax.
8. string, The result (antiderivative) produced by CAS in Latex format
9. string, The optimal antiderivative in Latex format.
10. integer, 0 or 1. Indicates if problem has known antiderivative or not
11. String, The result (antiderivative) in CAS own syntax.
12. String, The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String, Small string description of why the grade was given.
14. integer, 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

*The following fields are present only in Rubi Table file*

15. integer, Number of steps used.
16. integer, Number of rules used.
17. integer, Integrand leaf size.
18. real number, Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String, The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi  
June 27, 2013  
Design v0.01

# CHAPTER 2

## DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS . . . . .	21
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	24
2.3	Detailed conclusion table specific for Rubi results . . . . .	33

## 2.1 List of integrals sorted by grade for each CAS

2.1.1	Rubi . . . . .	21
2.1.2	Mma . . . . .	21
2.1.3	Maple . . . . .	22
2.1.4	Fricas . . . . .	22
2.1.5	Maxima . . . . .	22
2.1.6	Giac . . . . .	23
2.1.7	Mupad . . . . .	23
2.1.8	Sympy . . . . .	23

### 2.1.1 Rubi

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33 }

**B grade** { }

**C grade** { }

**F normal fail** { }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

### 2.1.2 Mma

**A grade** { 2, 3, 4, 5, 6, 7, 16, 19, 20, 21, 22, 23, 24, 32, 33 }

**B grade** { }

**C grade** { 1, 18 }

**F normal fail** { 8, 9, 10, 11, 12, 13, 14, 15, 17, 25, 26, 27, 28, 29, 30, 31 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

### 2.1.3 Maple

A grade { 32 }

B grade { 7, 16, 33 }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13, 14, 15, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 }

F(-1) timedout fail { }

F(-2) exception fail { }

### 2.1.4 Fricas

A grade { 2, 3, 19, 20, 32 }

B grade { 1, 18 }

C grade { 33 }

F normal fail { 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 }

F(-1) timedout fail { }

F(-2) exception fail { }

### 2.1.5 Maxima

A grade { 32 }

B grade { 1, 2, 3, 18, 19, 20 }

C grade { }

F normal fail { 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 33 }

F(-1) timedout fail { 34 }

F(-2) exception fail { }

### 2.1.6 Giac

A grade { 7, 16, 32 }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 8, 9, 10, 11, 12, 13, 14, 15, 17, 18, 19, 20, 24, 25, 26, 27, 28, 29, 30, 31, 33 }

F(-1) timeout fail { 4, 21, 34 }

F(-2) exception fail { 5, 6, 22, 23 }

### 2.1.7 Mupad

A grade { }

B grade { 1, 2, 3, 18, 19, 20, 32, 33 }

C grade { }

F normal fail { }

F(-1) timeout fail { 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 }

F(-2) exception fail { }

### 2.1.8 Sympy

A grade { }

B grade { 32 }

C grade { }

F normal fail { 3, 4, 5, 7, 8, 12, 13, 14, 16, 20, 21, 22, 24, 28, 29, 30, 33 }

F(-1) timeout fail { 1, 2, 6, 9, 10, 11, 15, 17, 18, 19, 23, 25, 26, 27, 31, 34 }

F(-2) exception fail { }



## 2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as  $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$ . To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	<b>F</b>	B	B	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	384	312	899	0	892	777	0	0	1110
N.S.	1	0.81	2.34	0.00	2.32	2.02	0.00	0.00	2.89
time (sec)	N/A	1.471	13.448	0.000	0.394	0.321	0.000	0.000	22.874

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	B	A	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	285	255	264	0	648	461	0	0	714
N.S.	1	0.89	0.93	0.00	2.27	1.62	0.00	0.00	2.51
time (sec)	N/A	1.188	5.871	0.000	0.361	0.305	0.000	0.000	20.948

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	B	A	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	180	183	160	0	441	261	0	0	185
N.S.	1	1.02	0.89	0.00	2.45	1.45	0.00	0.00	1.03
time (sec)	N/A	0.949	1.992	0.000	0.370	0.294	0.000	0.000	17.566

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	123	128	103	0	0	0	0	0	0
N.S.	1	1.04	0.84	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.640	32.981	0.000	0.000	0.000	0.000	0.000	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	202	214	110	0	0	0	0	0	0
N.S.	1	1.06	0.54	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.109	52.516	0.000	0.000	0.000	0.000	0.000	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	207	223	131	0	0	0	0	0	0
N.S.	1	1.08	0.63	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.144	36.826	0.000	0.000	0.000	0.000	0.000	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	167	181	190	541	0	0	0	197	0
N.S.	1	1.08	1.14	3.24	0.00	0.00	0.00	1.18	0.00
time (sec)	N/A	1.123	2.958	3.852	0.000	0.000	0.000	0.451	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	257	306	0	0	0	0	0	0	0
N.S.	1	1.19	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.184	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	366	388	0	0	0	0	0	0	0
N.S.	1	1.06	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.101	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	392	423	0	0	0	0	0	0	0
N.S.	1	1.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.288	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	385	406	0	0	0	0	0	0	0
N.S.	1	1.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.162	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	375	396	0	0	0	0	0	0	0
N.S.	1	1.06	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.125	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	365	395	0	0	0	0	0	0	0
N.S.	1	1.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.158	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	413	431	0	0	0	0	0	0	0
N.S.	1	1.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.269	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	424	451	0	0	0	0	0	0	0
N.S.	1	1.06	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.288	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	174	188	196	747	0	0	0	220	0
N.S.	1	1.08	1.13	4.29	0.00	0.00	0.00	1.26	0.00
time (sec)	N/A	1.136	3.129	3.905	0.000	0.000	0.000	0.467	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	269	318	0	0	0	0	0	0	0
N.S.	1	1.18	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.205	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	<b>F</b>	B	B	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	435	336	1029	0	1324	937	0	0	1253
N.S.	1	0.77	2.37	0.00	3.04	2.15	0.00	0.00	2.88
time (sec)	N/A	1.486	13.210	0.000	0.423	0.377	0.000	0.000	23.241

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	B	A	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	322	279	306	0	950	563	0	0	790
N.S.	1	0.87	0.95	0.00	2.95	1.75	0.00	0.00	2.45
time (sec)	N/A	1.220	6.435	0.000	0.380	0.340	0.000	0.000	24.214

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	B	A	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	197	200	177	0	644	310	0	0	510
N.S.	1	1.02	0.90	0.00	3.27	1.57	0.00	0.00	2.59
time (sec)	N/A	0.996	2.066	0.000	0.375	0.293	0.000	0.000	19.684

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	170	175	159	0	0	0	0	0	0
N.S.	1	1.03	0.94	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.931	31.181	0.000	0.000	0.000	0.000	0.000	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	216	228	162	0	0	0	0	0	0
N.S.	1	1.06	0.75	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.168	34.092	0.000	0.000	0.000	0.000	0.000	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	230	246	228	0	0	0	0	0	0
N.S.	1	1.07	0.99	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.247	29.424	0.000	0.000	0.000	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	232	299	196	0	0	0	0	0	0
N.S.	1	1.29	0.84	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.278	1.161	0.000	0.000	0.000	0.000	0.000	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	383	410	0	0	0	0	0	0	0
N.S.	1	1.07	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.172	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	410	441	0	0	0	0	0	0	0
N.S.	1	1.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.331	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	406	425	0	0	0	0	0	0	0
N.S.	1	1.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.216	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	396	415	0	0	0	0	0	0	0
N.S.	1	1.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.167	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	389	415	0	0	0	0	0	0	0
N.S.	1	1.07	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.215	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	433	451	0	0	0	0	0	0	0
N.S.	1	1.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.344	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	451	478	0	0	0	0	0	0	0
N.S.	1	1.06	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.410	0.000	0.000	0.000	0.000	0.000	0.000	0.000



Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	81	118	92	80	102	71	189	76	93
N.S.	1	1.46	1.14	0.99	1.26	0.88	2.33	0.94	1.15
time (sec)	N/A	0.349	0.278	1.126	0.224	0.270	0.135	0.340	14.098

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	C	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	120	97	396	0	232	0	0	169
N.S.	1	1.03	0.83	3.38	0.00	1.98	0.00	0.00	1.44
time (sec)	N/A	0.655	1.567	2.800	0.000	0.108	0.000	0.000	15.719

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	<b>F(-1)</b>	N/A	<b>F(-1)</b>	<b>F(-1)</b>	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	45	45	47	45	0	54	0	0	47
N.S.	1	1.00	1.04	1.00	0.00	1.20	0.00	0.00	1.04
time (sec)	N/A	0.292	24.925	2.037	0.000	1.429	0.000	0.000	73.796

## 2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio  $\frac{\text{number of rules}}{\text{integrand size}}$  is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [10] had the largest ratio of [.292683000000000026]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	11	11	0.81	40	0.275
2	A	9	9	0.89	40	0.225
3	A	7	7	1.02	40	0.175
4	A	8	7	1.04	40	0.175
5	A	11	10	1.06	40	0.250
6	A	11	10	1.08	40	0.250
7	A	11	10	1.08	42	0.238
8	A	12	11	1.19	38	0.289
9	A	11	10	1.06	37	0.270
10	A	13	12	1.08	41	0.293
11	A	12	11	1.05	39	0.282
12	A	12	11	1.06	39	0.282
13	A	12	11	1.08	39	0.282
14	A	12	11	1.04	39	0.282
15	A	12	11	1.06	39	0.282
16	A	11	10	1.08	50	0.200
17	A	12	11	1.18	46	0.239
18	A	11	11	0.77	48	0.229
19	A	9	9	0.87	48	0.188
20	A	7	7	1.02	48	0.146
21	A	10	9	1.03	48	0.188
22	A	11	10	1.06	48	0.208

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
23	A	11	10	1.07	48	0.208
24	A	11	10	1.29	50	0.200
25	A	11	10	1.07	45	0.222
26	A	13	12	1.08	49	0.245
27	A	12	11	1.05	47	0.234
28	A	12	11	1.05	47	0.234
29	A	12	11	1.07	47	0.234
30	A	12	11	1.04	47	0.234
31	A	12	11	1.06	47	0.234
32	A	4	4	1.46	31	0.129
33	A	11	11	1.03	41	0.268
34	N/A	2	0	1.00	45	0.000

LISTING OF INTEGRALS

3.1  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx \dots\dots\dots 37$

3.2  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx \dots\dots\dots 47$

3.3  $\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx \dots\dots\dots 56$

3.4  $\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx \dots\dots\dots 63$

3.5  $\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx \dots\dots\dots 69$

3.6  $\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx \dots\dots\dots 76$

3.7  $\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)} (c - c \sin(e + fx))^{3/2}} dx \dots\dots\dots 83$

3.8  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx \dots\dots\dots 90$

3.9  $\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx \dots\dots\dots 98$

3.10  $\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx \dots\dots\dots 106$

3.11  $\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx \dots\dots\dots 115$

3.12  $\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx \dots\dots\dots 123$

3.13  $\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx \dots\dots\dots 131$

3.14  $\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx \dots\dots\dots 139$

3.15  $\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx \dots\dots\dots 147$

3.16  $\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)} (c - c \sin(e + fx))^{3/2}} dx \dots\dots\dots 155$

3.17  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx 163$

3.18  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx 171$

3.19  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx 181$

3.20  $\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx 190$

3.21  $\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx \dots\dots\dots 198$

3.22  $\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx \dots\dots\dots 205$

3.23  $\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx \dots\dots\dots 212$

3.24  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx 219$

3.25  $\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx 227$

3.26	$\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{-2-m} (A+B \sin(e+fx) + C \sin^2(e+fx)) dx$	235
3.27	$\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{3/2} (A+B \sin(e+fx) + C \sin^2(e+fx)) dx$	244
3.28	$\int (a+a \sin(e+fx))^m \sqrt{c+d \sin(e+fx)} (A+B \sin(e+fx) + C \sin^2(e+fx)) dx$	252
3.29	$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx$	260
3.30	$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx$	268
3.31	$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{5/2}} dx$	276
3.32	$\int (a+b \sin(c+dx)) (A+B \sin(c+dx) + C \sin^2(c+dx)) dx$	284
3.33	$\int \frac{(a+b \sin(e+fx))(A+B \sin(e+fx)+C \sin^2(e+fx))}{\sin^{3/2}(e+fx)} dx$	290
3.34	$\int (a+b \sin(e+fx))^m (c+d \sin(e+fx))^n (A+B \sin(e+fx) + C \sin^2(e+fx)) dx$	298

### 3.1 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^{5/2} (A + C \sin^2(e +$

3.1.1	Optimal result . . . . .	37
3.1.2	Mathematica [C] (verified) . . . . .	38
3.1.3	Rubi [A] (verified) . . . . .	39
3.1.4	Maple [F] . . . . .	42
3.1.5	Fricas [B] (verification not implemented) . . . . .	43
3.1.6	Sympy [F(-1)] . . . . .	44
3.1.7	Maxima [B] (verification not implemented) . . . . .	44
3.1.8	Giac [F] . . . . .	45
3.1.9	Mupad [B] (verification not implemented) . . . . .	46

#### 3.1.1 Optimal result

Integrand size = 40, antiderivative size = 384

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx = \frac{64c^3(C(39 - 16m + 4m^2) + A(63 + 32m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m}{f(5 + 2m)(7 + 2m)(9 + 2m)(3 + 8m + 4m^2) \sqrt{c - c \sin(e + fx)}} + \frac{16c^2(C(39 - 16m + 4m^2) + A(63 + 32m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)}}{f(7 + 2m)(9 + 2m)(15 + 16m + 4m^2)} + \frac{2c(C(39 - 16m + 4m^2) + A(63 + 32m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2}}{f(5 + 2m)(7 + 2m)(9 + 2m)} - \frac{4C(1 + 2m) \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2}}{f(7 + 2m)(9 + 2m)} + \frac{2C \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{7/2}}{cf(9 + 2m)}$$

output

```
2*c*(C*(4*m^2-16*m+39)+A*(4*m^2+32*m+63))*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)/f/(9+2*m)/(4*m^2+24*m+35)-4*C*(1+2*m)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)/f/(4*m^2+32*m+63)+2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(7/2)/c/f/(9+2*m)+64*c^3*(C*(4*m^2-16*m+39)+A*(4*m^2+32*m+63))*cos(f*x+e)*(a+a*sin(f*x+e))^m/f/(9+2*m)/(16*m^4+128*m^3+344*m^2+352*m+105)/(c-c*sin(f*x+e))^(1/2)+16*c^2*(C*(4*m^2-16*m+39)+A*(4*m^2+32*m+63))*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)/f/(9+2*m)/(8*m^3+60*m^2+142*m+105)
```

### 3.1.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 13.45 (sec) , antiderivative size = 899, normalized size of antiderivative = 2.34

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx = \frac{(a(1 + \sin(e + fx)))^m (c - c \sin(e + fx))^{5/2} \left( \frac{18900A + 12285C + 15648Am + 648Cm + 5280Am^2 + 1416Cm^2 + 896Am^3 + 224Cm^3 + 64Am^4 + 16Cm^4}{(1 + 2m)(3 + 2m)(5 + 2m)(7 + 2m)(9 + 2m)} + \frac{18900A + 12285C + 15648Am + 648Cm + 5280Am^2 + 1416Cm^2 + 896Am^3 + 224Cm^3 + 64Am^4 + 16Cm^4}{(1 + 2m)(3 + 2m)(5 + 2m)(7 + 2m)(9 + 2m)} \right)}{(1 + 2m)(3 + 2m)(5 + 2m)(7 + 2m)(9 + 2m)}$$

input `Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(5/2)*(A + C*Sin[e + f*x]^2),x]`

output `((a*(1 + Sin[e + f*x]))^m*(c - c*Sin[e + f*x])^(5/2)*(((18900*A + 12285*C + 15648*A*m + 648*C*m + 5280*A*m^2 + 1416*C*m^2 + 896*A*m^3 + 224*C*m^3 + 64*A*m^4 + 16*C*m^4)*((1/8 + I/8)*Cos[(e + f*x)/2] + (1/8 - I/8)*Sin[(e + f*x)/2])))/((1 + 2*m)*(3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((18900*A + 12285*C + 15648*A*m + 648*C*m + 5280*A*m^2 + 1416*C*m^2 + 896*A*m^3 + 224*C*m^3 + 64*A*m^4 + 16*C*m^4)*((1/8 - I/8)*Cos[(e + f*x)/2] + (1/8 + I/8)*Sin[(e + f*x)/2])))/((1 + 2*m)*(3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((1575*A + 1575*C + 1178*A*m + 414*C*m + 292*A*m^2 + 100*C*m^2 + 24*A*m^3 + 8*C*m^3)*((1/4 - I/4)*Cos[(3*(e + f*x))/2] - (1/4 + I/4)*Sin[(3*(e + f*x))/2])))/((3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((1575*A + 1575*C + 1178*A*m + 414*C*m + 292*A*m^2 + 100*C*m^2 + 24*A*m^3 + 8*C*m^3)*((1/4 + I/4)*Cos[(3*(e + f*x))/2] - (1/4 - I/4)*Sin[(3*(e + f*x))/2])))/((3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((63*A + 189*C + 32*A*m + 44*C*m + 4*A*m^2 + 4*C*m^2)*((-1/4 + I/4)*Cos[(5*(e + f*x))/2] - (1/4 + I/4)*Sin[(5*(e + f*x))/2])))/((5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((63*A + 189*C + 32*A*m + 44*C*m + 4*A*m^2 + 4*C*m^2)*((-1/4 - I/4)*Cos[(5*(e + f*x))/2] - (1/4 - I/4)*Sin[(5*(e + f*x))/2])))/((5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((15 + 2*m)*((-3/16 - (3*I)/16)*C*Cos[(7*(e + f*x))/2] + (3/16 - (3*I)/16)*C*Sin[(7*(e + f*x))/2])))/((7 + 2*m)*(9 + 2*m)) + ((15 + 2*m)*((-3/16 + (3*I)/16)*C*Cos[(7*(e + f*x))/2] + (3/16 + (3*I)/16)*C*Sin[(7*(e + f*x))/2])))/((7 + 2*m)*(9 + ...`

### 3.1.3 Rubi [A] (verified)

Time = 1.47 (sec) , antiderivative size = 312, normalized size of antiderivative = 0.81, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.275$ , Rules used = {3042, 3519, 27, 3042, 3452, 3042, 3219, 3042, 3219, 3042, 3217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2) dx$$

$$\downarrow \text{3519}$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2} (a \sin(e + fx) + a)^m}{cf(2m + 9)} -$$

$$\frac{2 \int -\frac{1}{2} (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{5/2} (ac(C(7 - 2m) + A(2m + 9)) + 2acC(2m + 1) \sin(e + fx)) dx}{ac(2m + 9)}}{ac(2m + 9)}$$

$$\downarrow \text{27}$$

$$\frac{\int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{5/2} (ac(C(7 - 2m) + A(2m + 9)) + 2acC(2m + 1) \sin(e + fx)) dx}{ac(2m + 9)} +$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2} (a \sin(e + fx) + a)^m}{cf(2m + 9)}$$

$$\downarrow \text{3042}$$

$$\frac{\int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{5/2} (ac(C(7 - 2m) + A(2m + 9)) + 2acC(2m + 1) \sin(e + fx)) dx}{ac(2m + 9)} +$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2} (a \sin(e + fx) + a)^m}{cf(2m + 9)}$$

$$\downarrow \text{3452}$$

$$\frac{\frac{ac(A(4m^2 + 32m + 63) + C(4m^2 - 16m + 39))}{2m + 7} \int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{5/2} dx}{ac(2m + 9)} - \frac{4acC(2m + 1) \cos(e + fx)(c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m}{f(2m + 7)}}{cf(2m + 9)}$$

$$\downarrow \text{3042}$$

---

3.1.  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx$



$$\frac{ac(A(4m^2+32m+63)+C(4m^2-16m+39)) \int (\sin(e+fx)a+a)^m (c-c\sin(e+fx))^{5/2} dx}{2m+7} - \frac{4acC(2m+1) \cos(e+fx)(c-c\sin(e+fx))^{5/2} (a\sin(e+fx))^{2m}}{f(2m+7)}$$

$$\frac{ac(2m+9)}{cf(2m+9)} \frac{2C \cos(e+fx)(c-c\sin(e+fx))^{7/2} (a\sin(e+fx)+a)^m}{cf(2m+9)}$$

↓ 3219

$$\frac{ac(A(4m^2+32m+63)+C(4m^2-16m+39)) \left( \frac{8c \int (\sin(e+fx)a+a)^m (c-c\sin(e+fx))^{3/2} dx}{2m+5} + \frac{2c \cos(e+fx)(c-c\sin(e+fx))^{3/2} (a\sin(e+fx)+a)^m}{f(2m+5)} \right)}{2m+7} - \frac{4acC(2m+1) \cos(e+fx)(c-c\sin(e+fx))^{3/2} (a\sin(e+fx))^{2m}}{f(2m+5)}$$

$$\frac{ac(2m+9)}{cf(2m+9)} \frac{2C \cos(e+fx)(c-c\sin(e+fx))^{7/2} (a\sin(e+fx)+a)^m}{cf(2m+9)}$$

↓ 3042

$$\frac{ac(A(4m^2+32m+63)+C(4m^2-16m+39)) \left( \frac{8c \int (\sin(e+fx)a+a)^m (c-c\sin(e+fx))^{3/2} dx}{2m+5} + \frac{2c \cos(e+fx)(c-c\sin(e+fx))^{3/2} (a\sin(e+fx)+a)^m}{f(2m+5)} \right)}{2m+7} - \frac{4acC(2m+1) \cos(e+fx)(c-c\sin(e+fx))^{3/2} (a\sin(e+fx))^{2m}}{f(2m+5)}$$

$$\frac{ac(2m+9)}{cf(2m+9)} \frac{2C \cos(e+fx)(c-c\sin(e+fx))^{7/2} (a\sin(e+fx)+a)^m}{cf(2m+9)}$$

↓ 3219

$$\frac{ac(A(4m^2+32m+63)+C(4m^2-16m+39)) \left( \frac{8c \left( \frac{4c \int (\sin(e+fx)a+a)^m \sqrt{c-c\sin(e+fx)} dx}{2m+3} + \frac{2c \cos(e+fx) \sqrt{c-c\sin(e+fx)} (a\sin(e+fx)+a)^m}{f(2m+3)} \right)}{2m+5} + \frac{2c \cos(e+fx)(c-c\sin(e+fx))^{3/2} (a\sin(e+fx)+a)^m}{f(2m+5)} \right)}{2m+7} - \frac{4acC(2m+1) \cos(e+fx)(c-c\sin(e+fx))^{3/2} (a\sin(e+fx))^{2m}}{f(2m+5)}$$

$$\frac{ac(2m+9)}{cf(2m+9)} \frac{2C \cos(e+fx)(c-c\sin(e+fx))^{7/2} (a\sin(e+fx)+a)^m}{cf(2m+9)}$$

↓ 3042

$$\frac{ac(A(4m^2+32m+63)+C(4m^2-16m+39)) \left( \frac{8c \left( \frac{4c \int (\sin(e+fx)a+a)^m \sqrt{c-c\sin(e+fx)} dx}{2m+3} + \frac{2c \cos(e+fx) \sqrt{c-c\sin(e+fx)} (a\sin(e+fx)+a)^m}{f(2m+3)} \right)}{2m+5} + \frac{2c \cos(e+fx)(c-c\sin(e+fx))^{3/2} (a\sin(e+fx)+a)^m}{f(2m+5)} \right)}{2m+7} - \frac{4acC(2m+1) \cos(e+fx)(c-c\sin(e+fx))^{3/2} (a\sin(e+fx))^{2m}}{f(2m+5)}$$

$$\frac{ac(2m+9)}{cf(2m+9)} \frac{2C \cos(e+fx)(c-c\sin(e+fx))^{7/2} (a\sin(e+fx)+a)^m}{cf(2m+9)}$$

↓ 3217

---

3.1.  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx$

$$\frac{ac(A(4m^2+32m+63)+C(4m^2-16m+39)) \left( \frac{8c \left( \frac{8c^2 \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)(2m+3)\sqrt{c-c \sin(e+fx)}} + \frac{2c \cos(e+fx)\sqrt{c-c \sin(e+fx)}(a \sin(e+fx)+a)^m}{f(2m+3)} \right)}{2m+5} + \frac{2c \cos(e+fx)(c-c \sin(e+fx))}{2m+7} \right)}{ac(2m+9)} \\ \frac{2C \cos(e+fx)(c-c \sin(e+fx))^{7/2}(a \sin(e+fx)+a)^m}{cf(2m+9)}$$

input `Int[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(5/2)*(A + C*Sin[e + f*x]^2),x]`

output `(2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(7/2))/(c*f*(9 + 2*m)) + ((-4*a*c*C*(1 + 2*m)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(5/2))/(f*(7 + 2*m)) + (a*c*(C*(39 - 16*m + 4*m^2) + A*(63 + 32*m + 4*m^2))*((2*c*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2))/(f*(5 + 2*m)) + (8*c*((8*c^2*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) + (2*c*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*Sqrt[c - c*Sin[e + f*x]])/(f*(3 + 2*m)))))/(5 + 2*m)))/(7 + 2*m))/(a*c*(9 + 2*m))`

### 3.1.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3217 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[n, -2^(-1)]`

```
rule 3219 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^n/(f*(m + n))), x] + Simp[a*((2*m - 1)/(m + n)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[m - 1/2, 0] && !LtQ[n, -1] && !(IGtQ[n - 1/2, 0] && LtQ[n, m]) && !(ILtQ[m + n, 0] && GtQ[2*m + n + 1, 0])
```

```
rule 3452 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 1, 0]
```

```
rule 3519 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1))/(d*f*(m + n + 2)), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) - b*c*C*(2*m + 1)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]
```

### 3.1.4 Maple [F]

$$\int (a + a \sin(fx + e))^m (c - c \sin(fx + e))^{5/2} (A + C(\sin^2(fx + e))) dx$$

```
input int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+C*sin(f*x+e)^2),x)
```

```
output int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+C*sin(f*x+e)^2),x)
```

### 3.1.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 777 vs.  $2(364) = 728$ .

Time = 0.32 (sec) , antiderivative size = 777, normalized size of antiderivative = 2.02

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx = \frac{2((16Cc^2m^4 + 128Cc^2m^3 + 344Cc^2m^2 + 352Cc^2m + 105Cc^2) \cos(fx + e)^5 + 128$$

```
input integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+C*sin(f*x+e)^2),x,
algorithm="fricas")
```

```
output 2*((16*C*c^2*m^4 + 128*C*c^2*m^3 + 344*C*c^2*m^2 + 352*C*c^2*m + 105*C*c^2
)*cos(f*x + e)^5 + 128*(A + C)*c^2*m^2 - (16*C*c^2*m^4 + 224*C*c^2*m^3 + 7
76*C*c^2*m^2 + 904*C*c^2*m + 285*C*c^2)*cos(f*x + e)^4 + 512*(2*A - C)*c^2
*m - (16*(A + 3*C)*c^2*m^4 + 32*(5*A + 16*C)*c^2*m^3 + 8*(65*A + 253*C)*c^
2*m^2 + 8*(75*A + 328*C)*c^2*m + 3*(63*A + 289*C)*c^2)*cos(f*x + e)^3 + 96
*(21*A + 13*C)*c^2 + (16*(A + C)*c^2*m^4 + 224*(A + C)*c^2*m^3 + 8*(133*A
+ 85*C)*c^2*m^2 + 1864*(A + C)*c^2*m + 3*(231*A + 263*C)*c^2)*cos(f*x + e)
^2 + 2*(16*(A + C)*c^2*m^4 + 192*(A + C)*c^2*m^3 + 856*(A + C)*c^2*m^2 + 1
6*(109*A + 85*C)*c^2*m + 3*(483*A + 419*C)*c^2)*cos(f*x + e) + (128*(A + C
)*c^2*m^2 + (16*C*c^2*m^4 + 128*C*c^2*m^3 + 344*C*c^2*m^2 + 352*C*c^2*m +
105*C*c^2)*cos(f*x + e)^4 + 512*(2*A - C)*c^2*m + 2*(16*C*c^2*m^4 + 176*C*
c^2*m^3 + 560*C*c^2*m^2 + 628*C*c^2*m + 195*C*c^2)*cos(f*x + e)^3 + 96*(21
*A + 13*C)*c^2 - (16*(A + C)*c^2*m^4 + 160*(A + C)*c^2*m^3 + 8*(65*A + 113
*C)*c^2*m^2 + 24*(25*A + 57*C)*c^2*m + 9*(21*A + 53*C)*c^2)*cos(f*x + e)^2
- 2*(16*(A + C)*c^2*m^4 + 192*(A + C)*c^2*m^3 + 792*(A + C)*c^2*m^2 + 16*
(77*A + 101*C)*c^2*m + 3*(147*A + 211*C)*c^2)*cos(f*x + e))*sin(f*x + e))*
sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(32*f*m^5 + 400*f*m^4 + 1
840*f*m^3 + 3800*f*m^2 + 3378*f*m + (32*f*m^5 + 400*f*m^4 + 1840*f*m^3 + 3
800*f*m^2 + 3378*f*m + 945*f)*cos(f*x + e) - (32*f*m^5 + 400*f*m^4 + 1840*
f*m^3 + 3800*f*m^2 + 3378*f*m + 945*f)*sin(f*x + e) + 945*f)
```

---

3.1.  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx$

**3.1.6 Sympy [F(-1)]**

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(c-c*sin(f*x+e))**(5/2)*(A+C*sin(f*x+e)**2), x)`

output `Timed out`

**3.1.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 892 vs.  $2(364) = 728$ .

Time = 0.39 (sec) , antiderivative size = 892, normalized size of antiderivative = 2.32

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+C*sin(f*x+e)^2), x, algorithm="maxima")`

output

```

-2*((4*m^2 + 24*m + 43)*a^m*c^(5/2) - (12*m^2 + 40*m - 15)*a^m*c^(5/2)*si
n(f*x + e)/(cos(f*x + e) + 1) + 2*(4*m^2 + 8*m + 35)*a^m*c^(5/2)*sin(f*x +
e)^2/(cos(f*x + e) + 1)^2 + 2*(4*m^2 + 8*m + 35)*a^m*c^(5/2)*sin(f*x + e)
^3/(cos(f*x + e) + 1)^3 - (12*m^2 + 40*m - 15)*a^m*c^(5/2)*sin(f*x + e)^4/
(cos(f*x + e) + 1)^4 + (4*m^2 + 24*m + 43)*a^m*c^(5/2)*sin(f*x + e)^5/(cos
(f*x + e) + 1)^5)*A*e^(2*m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1) - m*lo
g(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/((8*m^3 + 36*m^2 + 46*m + 15)*
(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1)^(5/2)) + 4*(2*(4*m^2 + 56*m + 21
9)*a^m*c^(5/2) - 4*(4*m^3 + 56*m^2 + 219*m)*a^m*c^(5/2)*sin(f*x + e)/(cos(
f*x + e) + 1) + (16*m^4 + 240*m^3 + 1136*m^2 + 1380*m + 1971)*a^m*c^(5/2)*
sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - (48*m^4 + 496*m^3 + 1568*m^2 + 3108*
m - 315)*a^m*c^(5/2)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 4*(8*m^4 + 68*m
^3 + 290*m^2 + 111*m + 567)*a^m*c^(5/2)*sin(f*x + e)^4/(cos(f*x + e) + 1)^
4 + 4*(8*m^4 + 68*m^3 + 290*m^2 + 111*m + 567)*a^m*c^(5/2)*sin(f*x + e)^5/
(cos(f*x + e) + 1)^5 - (48*m^4 + 496*m^3 + 1568*m^2 + 3108*m - 315)*a^m*c^
(5/2)*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 + (16*m^4 + 240*m^3 + 1136*m^2 +
1380*m + 1971)*a^m*c^(5/2)*sin(f*x + e)^7/(cos(f*x + e) + 1)^7 - 4*(4*m^3
+ 56*m^2 + 219*m)*a^m*c^(5/2)*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 + 2*(4*
m^2 + 56*m + 219)*a^m*c^(5/2)*sin(f*x + e)^9/(cos(f*x + e) + 1)^9)*C*e^(2*
m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(...

```

### 3.1.8 Giac [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e)^2 + A) (-c \sin(fx + e) + c)^{5/2} (a \sin(fx + e) + a)^m dx$$

input

```

integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+C*sin(f*x+e)^2),x,
algorithm="giac")

```

output

```

integrate((C*sin(f*x + e)^2 + A)*(-c*sin(f*x + e) + c)^(5/2)*(a*sin(f*x +
e) + a)^m, x)

```

### 3.1.9 Mupad [B] (verification not implemented)

Time = 22.87 (sec) , antiderivative size = 1110, normalized size of antiderivative = 2.89

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^(5/2),x)`

output `((c - c*sin(e + f*x))^(1/2)*((C*c^2*(a + a*sin(e + f*x))^m*(m^3*352i + m^2*344i + m^3*128i + m^4*16i + 105i))/(8*f*(m^3*3378i + m^2*3800i + m^3*1840i + m^4*400i + m^5*32i + 945i)) + (c^2*exp(e*5i + f*x*5i)*(a + a*sin(e + f*x))^m*(18900*A + 12285*C + 15648*A*m + 648*C*m + 5280*A*m^2 + 896*A*m^3 + 64*A*m^4 + 1416*C*m^2 + 224*C*m^3 + 16*C*m^4))/(4*f*(m^3*3378i + m^2*3800i + m^3*1840i + m^4*400i + m^5*32i + 945i)) + (c^2*exp(e*4i + f*x*4i)*(a + a*sin(e + f*x))^m*(A*18900i + C*12285i + A*m*15648i + C*m*648i + A*m^2*5280i + A*m^3*896i + A*m^4*64i + C*m^2*1416i + C*m^3*224i + C*m^4*16i))/(4*f*(m^3*3378i + m^2*3800i + m^3*1840i + m^4*400i + m^5*32i + 945i)) + (c^2*exp(e*3i + f*x*3i)*(2*m + 1)*(a + a*sin(e + f*x))^m*(1575*A + 1575*C + 1178*A*m + 414*C*m + 292*A*m^2 + 24*A*m^3 + 100*C*m^2 + 8*C*m^3))/(2*f*(m^3*3378i + m^2*3800i + m^3*1840i + m^4*400i + m^5*32i + 945i)) + (c^2*exp(e*6i + f*x*6i)*(2*m + 1)*(a + a*sin(e + f*x))^m*(A*1575i + C*1575i + A*m*1178i + C*m*414i + A*m^2*292i + A*m^3*24i + C*m^2*100i + C*m^3*8i))/(2*f*(m^3*3378i + m^2*3800i + m^3*1840i + m^4*400i + m^5*32i + 945i)) + (C*c^2*exp(e*9i + f*x*9i)*(a + a*sin(e + f*x))^m*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105))/(8*f*(m^3*3378i + m^2*3800i + m^3*1840i + m^4*400i + m^5*32i + 945i)) - (3*C*c^2*exp(e*1i + f*x*1i)*(a + a*sin(e + f*x))^m*(720*m + 632*m^2 + 192*m^3 + 16*m^4 + 225))/(8*f*(m^3*3378i + m^2*3800i + m^3*1840i + m^4*400i + m^5*32i + 945i)) - (3*C*c^2*exp(e*8i + f*x*8i)*(a + a*sin(e + f*x))^m*(m*720i + m^2*...`

### 3.2 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^{3/2} (A + C \sin^2(e +$

3.2.1	Optimal result . . . . .	47
3.2.2	Mathematica [A] (verified) . . . . .	48
3.2.3	Rubi [A] (verified) . . . . .	48
3.2.4	Maple [F] . . . . .	51
3.2.5	Fricas [A] (verification not implemented) . . . . .	52
3.2.6	Sympy [F(-1)] . . . . .	52
3.2.7	Maxima [B] (verification not implemented) . . . . .	53
3.2.8	Giac [F] . . . . .	54
3.2.9	Mupad [B] (verification not implemented) . . . . .	54

#### 3.2.1 Optimal result

Integrand size = 40, antiderivative size = 285

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \frac{8c^2(C(19 - 8m + 4m^2) + A(35 + 24m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m}{f(5 + 2m)(7 + 2m)(3 + 8m + 4m^2) \sqrt{c - c \sin(e + fx)}} + \frac{2c(C(19 - 8m + 4m^2) + A(35 + 24m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)}}{f(3 + 2m)(5 + 2m)(7 + 2m)} - \frac{4C(1 + 2m) \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2}}{f(5 + 2m)(7 + 2m)} + \frac{2C \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2}}{cf(7 + 2m)}$$

output

```
-4*C*(1+2*m)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)/f/(4*m^2
+24*m+35)+2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)/c/f/(7+
2*m)+8*c^2*(C*(4*m^2-8*m+19)+A*(4*m^2+24*m+35))*cos(f*x+e)*(a+a*sin(f*x+e)
)^m/f/(7+2*m)/(8*m^3+36*m^2+46*m+15)/(c-c*sin(f*x+e))^(1/2)+2*c*(C*(4*m^2-
8*m+19)+A*(4*m^2+24*m+35))*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(
1/2)/f/(7+2*m)/(4*m^2+16*m+15)
```



### 3.2.2 Mathematica [A] (verified)

Time = 5.87 (sec) , antiderivative size = 264, normalized size of antiderivative = 0.93

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \frac{c(\cos(\frac{1}{2}(e + fx)) + \sin(\frac{1}{2}(e + fx))) (a(1 + \sin(e + fx)))^m \sqrt{c - c \sin(e + fx)} (700A + C \sin^2(e + fx))}{(700A + C \sin^2(e + fx))}$$

input `Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2)*(A + C*Sin[e + f*x]^2),x]`

output `(c*(Cos[(e + f*x)/2] + Sin[(e + f*x)/2])*(a*(1 + Sin[e + f*x]))^m*Sqrt[c - c*Sin[e + f*x]]*(700*A + 494*C + 760*A*m + 284*C*m + 272*A*m^2 + 136*C*m^2 + 32*A*m^3 + 16*C*m^3 - 2*C*(39 + 110*m + 68*m^2 + 8*m^3)*Cos[2*(e + f*x)] - (1 + 2*m)*(4*A*(35 + 24*m + 4*m^2) + C*(253 + 80*m + 12*m^2))*Sin[e + f*x] + 15*C*Sin[3*(e + f*x)] + 46*C*m*Sin[3*(e + f*x)] + 36*C*m^2*Sin[3*(e + f*x)] + 8*C*m^3*Sin[3*(e + f*x)])/(2*f*(1 + 2*m)*(3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2]))`

### 3.2.3 Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 255, normalized size of antiderivative = 0.89, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.225$ , Rules used = {3042, 3519, 27, 3042, 3452, 3042, 3219, 3042, 3217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (c - c \sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (c - c \sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2) dx \\ & \quad \downarrow \text{3519} \\ & \frac{2C \cos(e + fx) (c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m}{cf(2m + 7)} - \\ & \frac{2 \int -\frac{1}{2} (\sin(e + fx) a + a)^m (c - c \sin(e + fx))^{3/2} (ac(C(5 - 2m) + A(2m + 7)) + 2acC(2m + 1) \sin(e + fx)) dx}{ac(2m + 7)}} \end{aligned}$$

---

3.2.  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx$

↓ 27

$$\frac{\int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{3/2} (ac(C(5 - 2m) + A(2m + 7)) + 2acC(2m + 1) \sin(e + fx)) dx}{ac(2m + 7)} + \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{3/2} (ac(C(5 - 2m) + A(2m + 7)) + 2acC(2m + 1) \sin(e + fx)) dx}{ac(2m + 7)} + \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

↓ 3452

$$\frac{\frac{ac(A(4m^2 + 24m + 35) + C(4m^2 - 8m + 19))}{2m + 5} \int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{3/2} dx - \frac{4acC(2m + 1) \cos(e + fx)(c - c \sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{f(2m + 5)}}{ac(2m + 7)} + \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

↓ 3042

$$\frac{\frac{ac(A(4m^2 + 24m + 35) + C(4m^2 - 8m + 19))}{2m + 5} \int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{3/2} dx - \frac{4acC(2m + 1) \cos(e + fx)(c - c \sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{f(2m + 5)}}{ac(2m + 7)} + \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

↓ 3219

$$\frac{\frac{ac(A(4m^2 + 24m + 35) + C(4m^2 - 8m + 19))}{2m + 5} \left( \frac{4c \int (\sin(e + fx)a + a)^m \sqrt{c - c \sin(e + fx)} dx}{2m + 3} + \frac{2c \cos(e + fx) \sqrt{c - c \sin(e + fx)} (a \sin(e + fx) + a)^m}{f(2m + 3)} \right) - \frac{4acC(2m + 1) \cos(e + fx)(c - c \sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{f(2m + 5)}}{ac(2m + 7)} + \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

↓ 3042

$$\frac{\frac{ac(A(4m^2 + 24m + 35) + C(4m^2 - 8m + 19))}{2m + 5} \left( \frac{4c \int (\sin(e + fx)a + a)^m \sqrt{c - c \sin(e + fx)} dx}{2m + 3} + \frac{2c \cos(e + fx) \sqrt{c - c \sin(e + fx)} (a \sin(e + fx) + a)^m}{f(2m + 3)} \right) - \frac{4acC(2m + 1) \cos(e + fx)(c - c \sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{f(2m + 5)}}{ac(2m + 7)} + \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

---

3.2.  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx$

↓ 3217

$$\frac{ac(A(4m^2+24m+35)+C(4m^2-8m+19))\left(\frac{8c^2 \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)(2m+3)\sqrt{c-c \sin(e+fx)}} + \frac{2c \cos(e+fx)\sqrt{c-c \sin(e+fx)}(a \sin(e+fx)+a)^m}{f(2m+3)}\right)}{2m+5} - \frac{4acC(2m+1) \cos(e+fx)}{cf(2m+7)}$$

input `Int[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2)*(A + C*Sin[e + f*x]^2),x]`

output `(2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(5/2))/(c*f*(7 + 2*m)) + ((-4*a*c*C*(1 + 2*m)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2))/(f*(5 + 2*m)) + (a*c*(C*(19 - 8*m + 4*m^2) + A*(35 + 24*m + 4*m^2))*((8*c^2*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m))*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) + (2*c*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*Sqrt[c - c*Sin[e + f*x]))/(f*(3 + 2*m)))/(5 + 2*m)/(a*c*(7 + 2*m))`

### 3.2.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3217 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[n, -2^(-1)]`

```
rule 3219 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^n/(f*(m + n))), x] + Simp[a*((2*m - 1)/(m + n)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[m - 1/2, 0] && !LtQ[n, -1] && !(IGtQ[n - 1/2, 0] && LtQ[n, m]) && !(ILtQ[m + n, 0] && GtQ[2*m + n + 1, 0])
```

```
rule 3452 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 1, 0]
```

```
rule 3519 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1))/(d*f*(m + n + 2)), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) - b*c*C*(2*m + 1)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]
```

### 3.2.4 Maple [F]

$$\int (a + a \sin(fx + e))^m (c - c \sin(fx + e))^{3/2} (A + C(\sin^2(fx + e))) dx$$

```
input int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x)
```

```
output int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x)
```

### 3.2.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 461, normalized size of antiderivative = 1.62

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \frac{2((8Ccm^3 + 36Ccm^2 + 46Ccm + 15Cc) \cos(fx + e)^4 - 16(A + C)cm^2 + (8Ccm^3 + 68Ccm^2 + 110C$$

```
input integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x,
algorithm="fricas")
```

```
output -2*((8*C*c*m^3 + 36*C*c*m^2 + 46*C*c*m + 15*C*c)*cos(f*x + e)^4 - 16*(A +
C)*c*m^2 + (8*C*c*m^3 + 68*C*c*m^2 + 110*C*c*m + 39*C*c)*cos(f*x + e)^3 -
32*(3*A - C)*c*m - (8*(A + C)*c*m^3 + 4*(13*A + 5*C)*c*m^2 + 94*(A + C)*c*
m + (35*A + 43*C)*c)*cos(f*x + e)^2 - 4*(35*A + 19*C)*c - (8*(A + C)*c*m^3
+ 68*(A + C)*c*m^2 + 2*(95*A + 63*C)*c*m + (175*A + 143*C)*c)*cos(f*x + e
) - (16*(A + C)*c*m^2 + (8*C*c*m^3 + 36*C*c*m^2 + 46*C*c*m + 15*C*c)*cos(f
*x + e)^3 + 32*(3*A - C)*c*m - 8*(4*C*c*m^2 + 8*C*c*m + 3*C*c)*cos(f*x + e
)^2 + 4*(35*A + 19*C)*c - (8*(A + C)*c*m^3 + 52*(A + C)*c*m^2 + 2*(47*A +
79*C)*c*m + (35*A + 67*C)*c)*cos(f*x + e))*sin(f*x + e)*sqrt(-c*sin(f*x +
e) + c)*(a*sin(f*x + e) + a)^m/(16*f*m^4 + 128*f*m^3 + 344*f*m^2 + 352*f*
m + (16*f*m^4 + 128*f*m^3 + 344*f*m^2 + 352*f*m + 105*f)*cos(f*x + e) - (1
6*f*m^4 + 128*f*m^3 + 344*f*m^2 + 352*f*m + 105*f)*sin(f*x + e) + 105*f)
```

### 3.2.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \text{Timed out}$$

```
input integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)**2),
x)
```

```
output Timed out
```

### 3.2.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 648 vs.  $2(271) = 542$ .

Time = 0.36 (sec) , antiderivative size = 648, normalized size of antiderivative = 2.27

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx =$$

$$2 \left( \frac{\left( a^m c^{\frac{3}{2}} (2m+5) - \frac{a^m c^{\frac{3}{2}} (2m-3) \sin(fx+e)}{\cos(fx+e)+1} - \frac{a^m c^{\frac{3}{2}} (2m-3) \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{a^m c^{\frac{3}{2}} (2m+5) \sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right) A e^{\left( 2m \log\left( \frac{\sin(fx+e)}{\cos(fx+e)+1} + 1 \right) - m \log\left( \frac{\sin(fx+e)}{\cos(fx+e)+1} \right) \right)}}{(4m^2 + 8m + 3) \left( \frac{\sin(fx+e)^2}{(\cos(fx+e)+1)^2} + 1 \right)^{\frac{3}{2}}} \right)$$

```
input integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x,
algorithm="maxima")
```

```
output -2*((a^m*c^(3/2)*(2*m + 5) - a^m*c^(3/2)*(2*m - 3)*sin(f*x + e)/(cos(f*x + e) + 1) - a^m*c^(3/2)*(2*m - 3)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + a^m*c^(3/2)*(2*m + 5)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3)*A*e^(2*m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/(4*m^2 + 8*m + 3)*(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1)^(3/2) + 4*(2*a^m*c^(3/2)*(2*m + 13) - 4*(2*m^2 + 13*m)*a^m*c^(3/2)*sin(f*x + e)/(cos(f*x + e) + 1) + (8*m^3 + 60*m^2 + 66*m + 91)*a^m*c^(3/2)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - (8*m^3 + 20*m^2 + 82*m - 35)*a^m*c^(3/2)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 - (8*m^3 + 20*m^2 + 82*m - 35)*a^m*c^(3/2)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + (8*m^3 + 60*m^2 + 66*m + 91)*a^m*c^(3/2)*sin(f*x + e)^5/(cos(f*x + e) + 1)^5 - 4*(2*m^2 + 13*m)*a^m*c^(3/2)*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 + 2*a^m*c^(3/2)*(2*m + 13)*sin(f*x + e)^7/(cos(f*x + e) + 1)^7)*C*e^(2*m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/(16*m^4 + 128*m^3 + 344*m^2 + 352*m + 2*(16*m^4 + 128*m^3 + 344*m^2 + 352*m + 105)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + (16*m^4 + 128*m^3 + 344*m^2 + 352*m + 105)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 105*(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1)^(3/2))/f
```

---

3.2.  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx$

### 3.2.8 Giac [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e) + A) (-c \sin(fx + e) + c)^{3/2} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x,algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + A)*(-c*sin(f*x + e) + c)^(3/2)*(a*sin(f*x + e) + a)^m, x)`

### 3.2.9 Mupad [B] (verification not implemented)

Time = 20.95 (sec) , antiderivative size = 714, normalized size of antiderivative = 2.51

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \frac{\sqrt{c - c \sin(e + fx)}}{4 f (16 m^4 + 128 m^3 + 344 m^2 + 352 m + 105)} \left( \frac{C c (a + a \sin(e + fx))^m (m^3 8i + m^2 36i + m 46i + 15i)}{4 f (16 m^4 + 128 m^3 + 344 m^2 + 352 m + 105)} + \frac{c e^{3i + f x 3i} (a + a \sin(e + fx))}{4 f (16 m^4 + 128 m^3 + 344 m^2 + 352 m + 105)} \right)$$

input `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^(3/2),x)`

output

$$\begin{aligned}
& ((c - c\sin(e + fx))^{1/2} * ((C*c*(a + a\sin(e + fx))^m * (m^4 6i + m^2 * 36i \\
& + m^3 * 8i + 15i)) / (4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) + (c*\exp \\
& (e*3i + f*x*3i)*(a + a\sin(e + fx))^m * (1260*A + 735*C + 1144*A*m - 18*C*m \\
& + 336*A*m^2 + 32*A*m^3 + 100*C*m^2 + 8*C*m^3)) / (4*f*(352*m + 344*m^2 + 12 \\
& 8*m^3 + 16*m^4 + 105)) - (c*\exp(e*4i + f*x*4i)*(a + a\sin(e + fx))^m * (A*1 \\
& 260i + C*735i + A*m*1144i - C*m*18i + A*m^2*336i + A*m^3*32i + C*m^2*100i \\
& + C*m^3*8i)) / (4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) - (C*c*\exp(e \\
& *7i + f*x*7i)*(a + a\sin(e + fx))^m * (46*m + 36*m^2 + 8*m^3 + 15)) / (4*f*(3 \\
& 52*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) - (C*c*\exp(e*1i + f*x*1i)*(a + a \\
& *sin(e + fx))^m * (174*m + 100*m^2 + 8*m^3 + 63)) / (4*f*(352*m + 344*m^2 + 1 \\
& 28*m^3 + 16*m^4 + 105)) + (C*c*\exp(e*6i + f*x*6i)*(a + a\sin(e + fx))^m * ( \\
& m*174i + m^2*100i + m^3*8i + 63i)) / (4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^ \\
& 4 + 105)) + (c*\exp(e*5i + f*x*5i)*(2*m + 1)*(a + a\sin(e + fx))^m * (140*A \\
& + 175*C + 96*A*m + 16*C*m + 16*A*m^2 + 4*C*m^2)) / (4*f*(352*m + 344*m^2 + 1 \\
& 28*m^3 + 16*m^4 + 105)) - (c*\exp(e*2i + f*x*2i)*(2*m + 1)*(a + a\sin(e + f \\
& *x))^m * (A*140i + C*175i + A*m*96i + C*m*16i + A*m^2*16i + C*m^2*4i)) / (4*f* \\
& (352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105))) / (\exp(e*4i + f*x*4i) - (\exp(e \\
& *3i + f*x*3i)*(m^3*352i + m^2*344i + m^3*128i + m^4*16i + 105i)) / (352*m + 34 \\
& 4*m^2 + 128*m^3 + 16*m^4 + 105))
\end{aligned}$$



### 3.3 $\int (a+a \sin(e+fx))^m \sqrt{c-c \sin(e+fx)}(A+C \sin^2(e+fx)) dx$

3.3.1	Optimal result . . . . .	56
3.3.2	Mathematica [A] (verified) . . . . .	57
3.3.3	Rubi [A] (verified) . . . . .	57
3.3.4	Maple [F] . . . . .	59
3.3.5	Fricas [A] (verification not implemented) . . . . .	60
3.3.6	Sympy [F] . . . . .	60
3.3.7	Maxima [B] (verification not implemented) . . . . .	61
3.3.8	Giac [F] . . . . .	61
3.3.9	Mupad [B] (verification not implemented) . . . . .	62

#### 3.3.1 Optimal result

Integrand size = 40, antiderivative size = 180

$$\int (a+a \sin(e+fx))^m \sqrt{c-c \sin(e+fx)}(A+C \sin^2(e+fx)) dx$$

$$= \frac{2c(C-6Cm+A(5+2m)) \cos(e+fx)(a+a \sin(e+fx))^m}{f(1+2m)(5+2m)\sqrt{c-c \sin(e+fx)}} + \frac{4cC(1+2m) \cos(e+fx)(a+a \sin(e+fx))^{1+m}}{af(3+2m)(5+2m)\sqrt{c-c \sin(e+fx)}} + \frac{2C \cos(e+fx)(a+a \sin(e+fx))^m(c-c \sin(e+fx))^{3/2}}{cf(5+2m)}$$

output

```
2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)/c/f/(5+2*m)+2*c*(
C-6*C*m+A*(5+2*m))*cos(f*x+e)*(a+a*sin(f*x+e))^m/f/(4*m^2+12*m+5)/(c-c*sin
(f*x+e))^(1/2)+4*c*C*(1+2*m)*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)/a/f/(4*m^2+
16*m+15)/(c-c*sin(f*x+e))^(1/2)
```

### 3.3.2 Mathematica [A] (verified)

Time = 1.99 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.89

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx = \frac{(\cos(\frac{1}{2}(e + fx)) + \sin(\frac{1}{2}(e + fx))) (a(1 + \sin(e + fx)))^m \sqrt{c - c \sin(e + fx)} (-30A - 19C - 32Am - 8Cm - 8Am^2 - 4Cm^2 + C(3 + 8m + 4m^2) \cos[2(e + fx)] + 8C(1 + 2m) \sin[e + fx])}{f(1 + 2m)(3 + 2m)(5 + 2m) (\cos(\frac{1}{2}(e + fx)) - \sin(\frac{1}{2}(e + fx)))}$$

input `Integrate[(a + a*Sin[e + f*x])^m*Sqrt[c - c*Sin[e + f*x]]*(A + C*Sin[e + f*x]^2), x]`

output `-(((Cos[(e + f*x)/2] + Sin[(e + f*x)/2])*(a*(1 + Sin[e + f*x]))^m*Sqrt[c - c*Sin[e + f*x]]*(-30*A - 19*C - 32*A*m - 8*C*m - 8*A*m^2 - 4*C*m^2 + C*(3 + 8*m + 4*m^2)*Cos[2*(e + f*x)] + 8*C*(1 + 2*m)*Sin[e + f*x]))/(f*(1 + 2*m)*(3 + 2*m)*(5 + 2*m)*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2]))`

### 3.3.3 Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.02, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.175$ , Rules used = {3042, 3519, 27, 3042, 3450, 3042, 3217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{c - c \sin(e + fx)} (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) dx$$

↓ 3042

$$\int \sqrt{c - c \sin(e + fx)} (a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2) dx$$

↓ 3519

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{cf(2m + 5)} - \frac{2 \int -\frac{1}{2}(\sin(e + fx)a + a)^m \sqrt{c - c \sin(e + fx)}(ac(C(3 - 2m) + A(2m + 5)) + 2acC(2m + 1) \sin(e + fx))dx}{ac(2m + 5)}$$

↓ 27

---

3.3.  $\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx$

$$\frac{\int (\sin(e + fx)a + a)^m \sqrt{c - c\sin(e + fx)} (ac(C(3 - 2m) + A(2m + 5)) + 2acC(2m + 1) \sin(e + fx)) dx}{ac(2m + 5)} + \frac{2C \cos(e + fx)(c - c\sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{cf(2m + 5)}$$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^m \sqrt{c - c\sin(e + fx)} (ac(C(3 - 2m) + A(2m + 5)) + 2acC(2m + 1) \sin(e + fx)) dx}{ac(2m + 5)} + \frac{2C \cos(e + fx)(c - c\sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{cf(2m + 5)}$$

↓ 3450

$$\frac{ac(A(2m + 5) - 6Cm + C) \int (\sin(e + fx)a + a)^m \sqrt{c - c\sin(e + fx)} dx + 2cC(2m + 1) \int (\sin(e + fx)a + a)^{m+1} dx}{ac(2m + 5)} + \frac{2C \cos(e + fx)(c - c\sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{cf(2m + 5)}$$

↓ 3042

$$\frac{ac(A(2m + 5) - 6Cm + C) \int (\sin(e + fx)a + a)^m \sqrt{c - c\sin(e + fx)} dx + 2cC(2m + 1) \int (\sin(e + fx)a + a)^{m+1} dx}{ac(2m + 5)} + \frac{2C \cos(e + fx)(c - c\sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{cf(2m + 5)}$$

↓ 3217

$$\frac{\frac{2ac^2(A(2m+5)-6Cm+C) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}} + \frac{4c^2C(2m+1) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}}{f(2m+3)\sqrt{c-c\sin(e+fx)}}}{ac(2m + 5)} + \frac{2C \cos(e + fx)(c - c\sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{cf(2m + 5)}$$

input `Int[(a + a*Sin[e + f*x])^m*Sqrt[c - c*Sin[e + f*x]]*(A + C*Sin[e + f*x]^2),x]`

output `(2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2))/(c*f*(5 + 2*m)) + ((2*a*c^2*(C - 6*C*m + A*(5 + 2*m))*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) + (4*c^2*C*(1 + 2*m)*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m))/(f*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]])/(a*c*(5 + 2*m))`

---

3.3.  $\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx$

## 3.3.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3217 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[n, -2^(-1)]`
- rule 3450 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[B/d Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] - Simp[(B*c - A*d)/d Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`
- rule 3519 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1))/(d*f*(m + n + 2)), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) - b*c*C*(2*m + 1)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]`

## 3.3.4 Maple [F]

$$\int (a + a \sin(fx + e))^m \sqrt{c - c \sin(fx + e)} (A + C(\sin^2(fx + e))) dx$$

input `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x)`

output `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x)`

---

3.3.  $\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx$

### 3.3.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.45

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx =$$

$$\frac{2((4Cm^2 + 8Cm + 3C) \cos(fx + e)^3 - 4(A + C)m^2 + (4Cm^2 - C) \cos(fx + e)^2 - 16Am - (4(A$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x,  
algorithm="fricas")`

output `-2*((4*C*m^2 + 8*C*m + 3*C)*cos(f*x + e)^3 - 4*(A + C)*m^2 + (4*C*m^2 - C)  
*cos(f*x + e)^2 - 16*A*m - (4*(A + C)*m^2 + 8*(2*A + C)*m + 15*A + 11*C)*c  
os(f*x + e) - (4*(A + C)*m^2 - (4*C*m^2 + 8*C*m + 3*C)*cos(f*x + e)^2 + 16  
*A*m - 4*(2*C*m + C)*cos(f*x + e) + 15*A + 7*C)*sin(f*x + e) - 15*A - 7*C)  
*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(8*f*m^3 + 36*f*m^2 + 46  
*f*m + (8*f*m^3 + 36*f*m^2 + 46*f*m + 15*f)*cos(f*x + e) - (8*f*m^3 + 36*f  
*m^2 + 46*f*m + 15*f)*sin(f*x + e) + 15*f)`

### 3.3.6 Sympy [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \int (a(\sin(e + fx) + 1))^m \sqrt{-c(\sin(e + fx) - 1)} (A + C \sin^2(e + fx)) dx$$

input `integrate((a+a*sin(f*x+e))**m*(c-c*sin(f*x+e))**(1/2)*(A+C*sin(f*x+e)**2),  
x)`

output `Integral((a*(sin(e + f*x) + 1))**m*sqrt(-c*(sin(e + f*x) - 1))*(A + C*sin(  
e + f*x)**2), x)`

### 3.3.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 441 vs.  $2(170) = 340$ .

Time = 0.37 (sec) , antiderivative size = 441, normalized size of antiderivative = 2.45

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \frac{2 \left( \frac{4 \left( \frac{4 a^m \sqrt{c m} \sin(fx+e)}{\cos(fx+e)+1} - \frac{(4 m^2+4 m+5) a^m \sqrt{c} \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{(4 m^2+4 m+5) a^m \sqrt{c} \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{4 a^m \sqrt{c m} \sin(fx+e)^4}{(\cos(fx+e)+1)^4} - 2 a^m \sqrt{c} - \frac{2 a^m \sqrt{c} \sin(fx+e)^5}{(\cos(fx+e)+1)^5} \right)}{\left( 8 m^3+36 m^2+46 m+ \frac{2(8 m^3+36 m^2+46 m+15) \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{(8 m^3+36 m^2+46 m+15) \sin(fx+e)^4}{(\cos(fx+e)+1)^4} + 15 \right)} \right)}{}$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x,  
algorithm="maxima")`

output `2*(4*(4*a^m*sqrt(c)*m*sin(f*x + e)/(cos(f*x + e) + 1) - (4*m^2 + 4*m + 5)*  
a^m*sqrt(c)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - (4*m^2 + 4*m + 5)*a^m*sq  
rt(c)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 4*a^m*sqrt(c)*m*sin(f*x + e)^4  
/(cos(f*x + e) + 1)^4 - 2*a^m*sqrt(c) - 2*a^m*sqrt(c)*sin(f*x + e)^5/(cos(  
f*x + e) + 1)^5)*C*e^(2*m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1) - m*log  
(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/((8*m^3 + 36*m^2 + 46*m + 2*(8*  
m^3 + 36*m^2 + 46*m + 15)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + (8*m^3 + 3  
6*m^2 + 46*m + 15)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 15)*sqrt(sin(f*x  
+ e)^2/(cos(f*x + e) + 1)^2 + 1)) - (a^m*sqrt(c) + a^m*sqrt(c)*sin(f*x + e  
)/(cos(f*x + e) + 1))*A*e^(2*m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1) -  
m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/((2*m + 1)*sqrt(sin(f*x +  
e)^2/(cos(f*x + e) + 1)^2 + 1)))/f`

### 3.3.8 Giac [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(fx + e) + A) \sqrt{-c \sin(fx + e) + c} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x,  
algorithm="giac")`

---

3.3.  $\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx$

output `integrate((C*sin(f*x + e)^2 + A)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m, x)`

### 3.3.9 Mupad [B] (verification not implemented)

Time = 17.57 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.03

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx =$$

$$\frac{(a(\sin(e + fx) + 1))^m \sqrt{-c(\sin(e + fx) - 1)} (60A \cos(e + fx) + 35C \cos(e + fx) - 3C \cos(3e + 3fx))}{(2f(\sin(e + fx) - 1)(46m + 36m^2 + 8m^3 + 15))}$$

input `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^(1/2),x)`

output `-((a*(sin(e + f*x) + 1))^m*(-c*(sin(e + f*x) - 1))^(1/2)*(60*A*cos(e + f*x) + 35*C*cos(e + f*x) - 3*C*cos(3*e + 3*f*x) - 8*C*sin(2*e + 2*f*x) - 4*C*m^2*cos(3*e + 3*f*x) + 64*A*m*cos(e + f*x) + 8*C*m*cos(e + f*x) + 16*A*m^2*cos(e + f*x) - 8*C*m*cos(3*e + 3*f*x) + 4*C*m^2*cos(e + f*x) - 16*C*m*sin(2*e + 2*f*x)))/(2*f*(sin(e + f*x) - 1)*(46*m + 36*m^2 + 8*m^3 + 15))`

$$3.4 \quad \int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{\sqrt{c-c \sin(e+fx)}} dx$$

3.4.1	Optimal result . . . . .	63
3.4.2	Mathematica [A] (verified) . . . . .	63
3.4.3	Rubi [A] (verified) . . . . .	64
3.4.4	Maple [F] . . . . .	66
3.4.5	Fricas [F] . . . . .	66
3.4.6	Sympy [F] . . . . .	67
3.4.7	Maxima [F] . . . . .	67
3.4.8	Giac [F(-1)] . . . . .	67
3.4.9	Mupad [F(-1)] . . . . .	68

### 3.4.1 Optimal result

Integrand size = 40, antiderivative size = 123

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= \frac{(A + C) \cos(e + fx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e + fx))\right) (a + a \sin(e + fx))^m}{f(1 + 2m)\sqrt{c - c \sin(e + fx)}} - \frac{2C \cos(e + fx)(a + a \sin(e + fx))^{1+m}}{af(3 + 2m)\sqrt{c - c \sin(e + fx)}}$$

```
output (A+C)*cos(f*x+e)*hypergeom([1, 1/2+m], [3/2+m], 1/2+1/2*sin(f*x+e))*(a+a*sin
(f*x+e))^m/f/(1+2*m)/(c-c*sin(f*x+e))^(1/2)-2*C*cos(f*x+e)*(a+a*sin(f*x+e)
)^(1+m)/a/f/(3+2*m)/(c-c*sin(f*x+e))^(1/2)
```

### 3.4.2 Mathematica [A] (verified)

Time = 32.98 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.84

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx =$$

$$\frac{\cos(e + fx)(a(1 + \sin(e + fx)))^m \left(-((A + C)(3 + 2m) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e + fx))\right)\right)}{f(1 + 2m)(3 + 2m)\sqrt{c - c \sin(e + fx)}}$$

---

3.4.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{\sqrt{c-c \sin(e+fx)}} dx$



input `Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/Sqrt[c - c*Sin[e + f*x]],x]`

output `-((Cos[e + f*x]*(a*(1 + Sin[e + f*x]))^m*(-((A + C)*(3 + 2*m)*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2]) + 2*C*(1 + 2*m)*(1 + Sin[e + f*x])))/(f*(1 + 2*m)*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]])`

### 3.4.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.04, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.175$ , Rules used = {3042, 3517, 3042, 3224, 3042, 3146, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a \sin(e + fx) + a)^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2)}{\sqrt{c - c \sin(e + fx)}} dx \\
 & \quad \downarrow \text{3517} \\
 & (A + C) \int \frac{(\sin(e + fx)a + a)^m}{\sqrt{c - c \sin(e + fx)}} dx - \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^{m+1}}{af(2m + 3)\sqrt{c - c \sin(e + fx)}} \\
 & \quad \downarrow \text{3042} \\
 & (A + C) \int \frac{(\sin(e + fx)a + a)^m}{\sqrt{c - c \sin(e + fx)}} dx - \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^{m+1}}{af(2m + 3)\sqrt{c - c \sin(e + fx)}} \\
 & \quad \downarrow \text{3224} \\
 & \frac{(A + C) \cos(e + fx) \int \sec(e + fx)(\sin(e + fx)a + a)^{m+\frac{1}{2}} dx}{\sqrt{a \sin(e + fx) + a} \sqrt{c - c \sin(e + fx)}} - \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^{m+1}}{af(2m + 3)\sqrt{c - c \sin(e + fx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A + C) \cos(e + fx) \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}}{\cos(e+fx)} dx}{\sqrt{a \sin(e + fx) + a} \sqrt{c - c \sin(e + fx)}} - \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^{m+1}}{af(2m + 3)\sqrt{c - c \sin(e + fx)}}
 \end{aligned}$$

---

3.4.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{\sqrt{c-c \sin(e+fx)}} dx$

$$\begin{aligned}
 & \downarrow \text{3146} \\
 & \frac{a(A+C)\cos(e+fx)\int\frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}d(a\sin(e+fx))}{a-a\sin(e+fx)}}{f\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} - \frac{2C\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}} \\
 & \downarrow \text{78} \\
 & \frac{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^m \operatorname{Hypergeometric2F1}\left(1, m+\frac{1}{2}, m+\frac{3}{2}, \frac{\sin(e+fx)a+a}{2a}\right)}{f(2m+1)\sqrt{c-c\sin(e+fx)}} - \frac{2C\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}}
 \end{aligned}$$

input `Int[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/Sqrt[c - c*Sin[e + f*x]],x]`

output `((A + C)*Cos[e + f*x]*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (a + a*Sin[e + f*x])/(2*a)]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) - (2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m))/(a*f*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]])`

### 3.4.3.1 Defintions of rubi rules used

rule 78 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3146 `Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1)/2], x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])`

---

3.4.  $\int \frac{(a+a\sin(e+fx))^m (A+C\sin^2(e+fx))}{\sqrt{c-c\sin(e+fx)}} dx$

rule 3224 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*Sin[e + f*x])^FracPart[m]*((c + d*Sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*Sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])`

rule 3517 `Int((((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)]^2))/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] :> Simp[-2*C*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(2*m + 3)*Sqrt[c + d*Sin[e + f*x]]), x] + Simp[(A + C) Int[(a + b*Sin[e + f*x])^m/Sqrt[c + d*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

### 3.4.4 Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + C(\sin^2(fx + e)))}{\sqrt{c - c \sin(fx + e)}} dx$$

input `int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x)`

output `int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x)`

### 3.4.5 Fricas [F]

$$\begin{aligned} & \int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx \\ &= \int \frac{(C \sin(fx + e)^2 + A)(a \sin(fx + e) + a)^m}{\sqrt{-c \sin(fx + e) + c}} dx \end{aligned}$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x, algorithm="fricas")`

output `integral((C*cos(f*x + e)^2 - A - C)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(c*sin(f*x + e) - c), x)`

---

3.4.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{\sqrt{c-c \sin(e+fx)}} dx$

### 3.4.6 Sympy [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= \int \frac{(a(\sin(e + fx) + 1))^m (A + C \sin^2(e + fx))}{\sqrt{-c(\sin(e + fx) - 1)}} dx$$

input `integrate((a+a*sin(f*x+e))**m*(A+C*sin(f*x+e)**2)/(c-c*sin(f*x+e))**(1/2), x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + C*sin(e + f*x)**2)/sqrt(-c*(sin(e + f*x) - 1)), x)`

### 3.4.7 Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{\sqrt{-c \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/sqrt(-c*sin(f*x + e) + c), x)`

### 3.4.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x,  
algorithm="giac")`

output `Timed out`

### 3.4.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin(e + fx)^2 + A) (a + a \sin(e + fx))^m}{\sqrt{c - c \sin(e + fx)}} dx$$

input `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c - c*sin(e + f*x))^(  
1/2),x)`

output `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c - c*sin(e + f*x))^(  
1/2), x)`

$$3.5 \quad \int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{3/2}} dx$$

3.5.1	Optimal result . . . . .	69
3.5.2	Mathematica [A] (verified) . . . . .	69
3.5.3	Rubi [A] (verified) . . . . .	70
3.5.4	Maple [F] . . . . .	73
3.5.5	Fricas [F] . . . . .	74
3.5.6	Sympy [F] . . . . .	74
3.5.7	Maxima [F] . . . . .	74
3.5.8	Giac [F(-2)] . . . . .	75
3.5.9	Mupad [F(-1)] . . . . .	75

### 3.5.1 Optimal result

Integrand size = 40, antiderivative size = 202

$$\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{3/2}} dx = \frac{(A+C) \cos(e+fx)(a+a \sin(e+fx))^{1+m}}{4af(c-c \sin(e+fx))^{3/2}} + \frac{(A+2Am+C(9+2m)) \cos(e+fx)(a+a \sin(e+fx))^m}{4cf(1+2m)\sqrt{c-c \sin(e+fx)}} + \frac{(A(1-2m)-C(7+2m)) \cos(e+fx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}+m, \frac{3}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right)}{4cf(1+2m)\sqrt{c-c \sin(e+fx)}} (a+a \sin(e+fx))^{1+m}$$

output

```
1/4*(A+C)*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)/a/f/(c-c*sin(f*x+e))^(3/2)+1/4
*(A+2*A*m+C*(9+2*m))*cos(f*x+e)*(a+a*sin(f*x+e))^m/c/f/(1+2*m)/(c-c*sin(f*
x+e))^(1/2)+1/4*(A*(1-2*m)-C*(7+2*m))*cos(f*x+e)*hypergeom([1, 1/2+m],[3/2
+m],1/2+1/2*sin(f*x+e))*(a+a*sin(f*x+e))^m/c/f/(1+2*m)/(c-c*sin(f*x+e))^(1
/2)
```

### 3.5.2 Mathematica [A] (verified)

Time = 52.52 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.54

$$\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{3/2}} dx = \frac{\cos(e+fx) \left(-4C + 4C \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}+m, \frac{3}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right) - (A+C) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}+m, \frac{3}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right)\right)}{2cf(1+2m)\sqrt{c-c \sin(e+fx)}} (a+a \sin(e+fx))^{1+m}$$

---

3.5.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{3/2}} dx$

input `Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c - c*Sin[e + f*x])^(3/2),x]`

output `-1/2*(Cos[e + f*x]*(-4*C + 4*C*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2] - (A + C)*Hypergeometric2F1[2, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2])*(a*(1 + Sin[e + f*x])^m)/(c*f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]])`

### 3.5.3 Rubi [A] (verified)

Time = 1.11 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.06, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {3042, 3515, 27, 3042, 3452, 3042, 3224, 3042, 3146, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a \sin(e + fx) + a)^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2)}{(c - c \sin(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{3515} \\
 & \frac{(A + C) \cos(e + fx) (a \sin(e + fx) + a)^{m+1}}{4af(c - c \sin(e + fx))^{3/2}} - \\
 & \frac{\int \frac{(\sin(e + fx)a + a)^m (a^2(A(3 - 2m) - C(2m + 5)) - a^2(2mA + A + C(2m + 9)) \sin(e + fx))}{2\sqrt{c - c \sin(e + fx)}} dx}{4a^2c} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{(\sin(e + fx)a + a)^m (a^2(A(3 - 2m) - C(2m + 5)) - a^2(2mA + A + C(2m + 9)) \sin(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx}{8a^2c} + \\
 & \frac{(A + C) \cos(e + fx) (a \sin(e + fx) + a)^{m+1}}{4af(c - c \sin(e + fx))^{3/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.5.  $\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx$

$$\begin{aligned}
& \frac{\int \frac{(\sin(e+fx)a+a)^m (a^2(A(3-2m)-C(2m+5))-a^2(2mA+A+C(2m+9)) \sin(e+fx))}{\sqrt{c-c \sin(e+fx)}} dx}{\frac{8a^2c}{(A+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} + \frac{4af(c-c \sin(e+fx))^{3/2}}{}} \\
& \quad \downarrow \text{3452} \\
& \frac{2a^2(A(1-2m)-C(2m+7)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c \sin(e+fx)}} dx + \frac{2a^2(2Am+A+C(2m+9)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c \sin(e+fx)}}}{\frac{8a^2c}{(A+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} + \frac{4af(c-c \sin(e+fx))^{3/2}}{}} + \\
& \quad \downarrow \text{3042} \\
& \frac{2a^2(A(1-2m)-C(2m+7)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c \sin(e+fx)}} dx + \frac{2a^2(2Am+A+C(2m+9)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c \sin(e+fx)}}}{\frac{8a^2c}{(A+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} + \frac{4af(c-c \sin(e+fx))^{3/2}}{}} + \\
& \quad \downarrow \text{3224} \\
& \frac{2a^2(A(1-2m)-C(2m+7)) \cos(e+fx) \int \frac{\sec(e+fx)(\sin(e+fx)a+a)^{m+\frac{1}{2}}}{\sqrt{a \sin(e+fx)+a\sqrt{c-c \sin(e+fx)}}} dx + \frac{2a^2(2Am+A+C(2m+9)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c \sin(e+fx)}}}{\frac{8a^2c}{(A+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} + \frac{4af(c-c \sin(e+fx))^{3/2}}{}} + \\
& \quad \downarrow \text{3042} \\
& \frac{2a^2(A(1-2m)-C(2m+7)) \cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}}{\cos(e+fx)\sqrt{a \sin(e+fx)+a\sqrt{c-c \sin(e+fx)}}} dx + \frac{2a^2(2Am+A+C(2m+9)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c \sin(e+fx)}}}{\frac{8a^2c}{(A+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} + \frac{4af(c-c \sin(e+fx))^{3/2}}{}} + \\
& \quad \downarrow \text{3146} \\
& \frac{2a^3(A(1-2m)-C(2m+7)) \cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{a-a \sin(e+fx)} d(a \sin(e+fx)) + \frac{2a^2(2Am+A+C(2m+9)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c \sin(e+fx)}}}{\frac{8a^2c}{(A+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} + \frac{4af(c-c \sin(e+fx))^{3/2}}{}} + \\
& \quad \downarrow \text{78}
\end{aligned}$$

---

3.5.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{3/2}} dx$



$$\frac{2a^2(A(1-2m)-C(2m+7))\cos(e+fx)(a\sin(e+fx)+a)^m \operatorname{Hypergeometric2F1}\left(1, m+\frac{1}{2}, m+\frac{3}{2}, \frac{\sin(e+fx)a+a}{2a}\right)}{f(2m+1)\sqrt{c-c\sin(e+fx)}} + \frac{2a^2(2Am+A+C(2m+9))\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{f(2m+1)\sqrt{c-c\sin(e+fx)}} \\ \frac{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{4af(c-c\sin(e+fx))^{3/2}}$$

input `Int[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c - c*Sin[e + f*x])^(3/2), x]`

output `((A + C)*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m))/(4*a*f*(c - c*Sin[e + f*x])^(3/2)) + (((2*a^2*(A + 2*A*m + C*(9 + 2*m))*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) + (2*a^2*(A*(1 - 2*m) - C*(7 + 2*m))*Cos[e + f*x]*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (a + a*Sin[e + f*x])/(2*a)]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]))/(8*a^2*c)`

### 3.5.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 78 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3146 `Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1/2), x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])`

```
rule 3224 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*Sin[e + f*x])^FracPart[m]*((c + d*Sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*Sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])
```

```
rule 3452 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 1, 0]
```

```
rule 3515 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(a*A + a*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(2*b*c*f*(2*m + 1))), x] - Simp[1/(2*b*c*d*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[A*(c^2*(m + 1) + d^2*(2*m + n + 2)) - C*(c^2*m - d^2*(n + 1)) + d*(A*c*(m + n + 2) - c*C*(3*m - n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (EqQ[m + n + 2, 0] && NeQ[2*m + 1, 0]))
```

### 3.5.4 Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + C \sin^2(fx + e))}{(c - c \sin(fx + e))^{\frac{3}{2}}} dx$$

```
input int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x)
```

```
output int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x)
```

---

3.5.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{\frac{3}{2}}} dx$

### 3.5.5 Fracas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{(-c \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x,  
algorithm="fricas")`

output `integral((C*cos(f*x + e)^2 - A - C)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x +  
e) + a)^m/(c^2*cos(f*x + e)^2 + 2*c^2*sin(f*x + e) - 2*c^2), x)`

### 3.5.6 Sympy [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(a(\sin(e + fx) + 1))^m (A + C \sin^2(e + fx))}{(-c(\sin(e + fx) - 1))^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))**m*(A+C*sin(f*x+e)**2)/(c-c*sin(f*x+e))**(3/2),  
x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + C*sin(e + f*x)**2)/(-c*(sin(e + f*  
x) - 1))**(3/2), x)`

### 3.5.7 Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{(-c \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x,  
algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/(-c*sin(f*x + e) +  
c)^(3/2), x)`

---

3.5.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{3/2}} dx$

### 3.5.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x,  
algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:Unable to divide, perhaps due to ro  
unding error%%{1,[0,1,1,1,0,0,0,0,0]%%}+%%{1,[0,0,1,1,1,0,0,0,0]%%} /  
%%{16,[0`

### 3.5.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin(e + fx)^2 + A) (a + a \sin(e + fx))^m}{(c - c \sin(e + fx))^{3/2}} dx$$

input `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c - c*sin(e + f*x))^(  
3/2),x)`

output `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c - c*sin(e + f*x))^(  
3/2), x)`

$$3.6 \quad \int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx$$

3.6.1	Optimal result . . . . .	76
3.6.2	Mathematica [A] (verified) . . . . .	76
3.6.3	Rubi [A] (verified) . . . . .	77
3.6.4	Maple [F] . . . . .	80
3.6.5	Fricas [F] . . . . .	80
3.6.6	Sympy [F(-1)] . . . . .	81
3.6.7	Maxima [F] . . . . .	81
3.6.8	Giac [F(-2)] . . . . .	81
3.6.9	Mupad [F(-1)] . . . . .	82

### 3.6.1 Optimal result

Integrand size = 40, antiderivative size = 207

$$\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx = \frac{(A+C) \cos(e+fx)(a+a \sin(e+fx))^{1+m}}{8af(c-c \sin(e+fx))^{5/2}} + \frac{(A(5-2m)-C(11+2m)) \cos(e+fx)(a+a \sin(e+fx))^m}{16cf(c-c \sin(e+fx))^{3/2}} + \frac{(A(3-8m+4m^2)+C(19+24m+4m^2)) \cos(e+fx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}+m, \frac{3}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right)}{32c^2f(1+2m)\sqrt{c-c \sin(e+fx)}}$$

```
output 1/8*(A+C)*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)/a/f/(c-c*sin(f*x+e))^(5/2)+1/16*(A*(5-2*m)-C*(11+2*m))*cos(f*x+e)*(a+a*sin(f*x+e))^m/c/f/(c-c*sin(f*x+e))^(3/2)+1/32*(A*(4*m^2-8*m+3)+C*(4*m^2+24*m+19))*cos(f*x+e)*hypergeom([1, 1/2+m], [3/2+m], 1/2+1/2*sin(f*x+e))*(a+a*sin(f*x+e))^m/c^2/f/(1+2*m)/(c-c*sin(f*x+e))^(1/2)
```

### 3.6.2 Mathematica [A] (verified)

Time = 36.83 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.63

$$\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx = \frac{\cos(e+fx) (4C \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}+m, \frac{3}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right) + (A(5-2m)-C(11+2m)) (a+a \sin(e+fx))^m}{32c^2f(1+2m)\sqrt{c-c \sin(e+fx)}}$$

---

3.6.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx$

input `Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c - c*Sin[e + f*x])^(5/2),x]`

output `(Cos[e + f*x]*(4*C*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2] - 4*C*Hypergeometric2F1[2, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2] + (A + C)*Hypergeometric2F1[3, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2])*(a*(1 + Sin[e + f*x]))^m)/(4*c^2*(f + 2*f*m)*Sqrt[c - c*Sin[e + f*x]])]`

### 3.6.3 Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {3042, 3515, 27, 3042, 3451, 3042, 3224, 3042, 3146, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a \sin(e + fx) + a)^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2)}{(c - c \sin(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{3515} \\
 & \frac{(A + C) \cos(e + fx)(a \sin(e + fx) + a)^{m+1}}{8af(c - c \sin(e + fx))^{5/2}} - \\
 & \frac{\int -\frac{(\sin(e+fx)a+a)^m((A(9-2m)-C(2m+7))a^2+(A(1-2m)-C(2m+15))\sin(e+fx)a^2)}{2(c-c\sin(e+fx))^{3/2}} dx}{8a^2c} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{(\sin(e+fx)a+a)^m((A(9-2m)-C(2m+7))a^2+(A(1-2m)-C(2m+15))\sin(e+fx)a^2)}{(c-c\sin(e+fx))^{3/2}} dx}{16a^2c} + \\
 & \frac{(A + C) \cos(e + fx)(a \sin(e + fx) + a)^{m+1}}{8af(c - c \sin(e + fx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{(\sin(e+fx)a+a)^m((A(9-2m)-C(2m+7))a^2+(A(1-2m)-C(2m+15))\sin(e+fx)a^2)}{(c-c\sin(e+fx))^{3/2}} dx}{16a^2c} + \\
 & \frac{(A + C) \cos(e + fx)(a \sin(e + fx) + a)^{m+1}}{8af(c - c \sin(e + fx))^{5/2}}
 \end{aligned}$$

---

3.6.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx$

$$\begin{aligned}
& \downarrow \text{3451} \\
& \frac{a^2(A(4m^2-8m+3)+C(4m^2+24m+19)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx}{2c} + \frac{a^2(A(5-2m)-C(2m+11)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} \\
& \frac{16a^2c}{(A+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} \\
& \frac{8af(c-c\sin(e+fx))^{5/2}}{+} \\
& \downarrow \text{3042} \\
& \frac{a^2(A(4m^2-8m+3)+C(4m^2+24m+19)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx}{2c} + \frac{a^2(A(5-2m)-C(2m+11)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} \\
& \frac{16a^2c}{(A+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} \\
& \frac{8af(c-c\sin(e+fx))^{5/2}}{+} \\
& \downarrow \text{3224} \\
& \frac{a^2(A(4m^2-8m+3)+C(4m^2+24m+19)) \cos(e+fx) \int \sec(e+fx)(\sin(e+fx)a+a)^{m+\frac{1}{2}} dx}{2c\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \frac{a^2(A(5-2m)-C(2m+11)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} \\
& \frac{16a^2c}{(A+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} \\
& \frac{8af(c-c\sin(e+fx))^{5/2}}{+} \\
& \downarrow \text{3042} \\
& \frac{a^2(A(4m^2-8m+3)+C(4m^2+24m+19)) \cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}}{\cos(e+fx)} dx}{2c\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \frac{a^2(A(5-2m)-C(2m+11)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} \\
& \frac{16a^2c}{(A+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} \\
& \frac{8af(c-c\sin(e+fx))^{5/2}}{+} \\
& \downarrow \text{3146} \\
& \frac{a^3(A(4m^2-8m+3)+C(4m^2+24m+19)) \cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{a-a\sin(e+fx)} d(a \sin(e+fx))}{2cf\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \frac{a^2(A(5-2m)-C(2m+11)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} \\
& \frac{16a^2c}{(A+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} \\
& \frac{8af(c-c\sin(e+fx))^{5/2}}{+} \\
& \downarrow \text{78} \\
& \frac{a^2(A(4m^2-8m+3)+C(4m^2+24m+19)) \cos(e+fx)(a \sin(e+fx)+a)^m \text{Hypergeometric2F1}\left(1, m+\frac{1}{2}, m+\frac{3}{2}, \frac{\sin(e+fx)a+a}{2a}\right)}{2cf(2m+1)\sqrt{c-c\sin(e+fx)}} + \frac{a^2(A(5-2m)-C(2m+11)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} \\
& \frac{16a^2c}{(A+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} \\
& \frac{8af(c-c\sin(e+fx))^{5/2}}{+}
\end{aligned}$$

---

3.6.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx$

input `Int[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c - c*Sin[e + f*x])^(5/2),x]`

output `((A + C)*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m))/(8*a*f*(c - c*Sin[e + f*x])^(5/2)) + ((a^2*(A*(5 - 2*m) - C*(11 + 2*m))*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(c - c*Sin[e + f*x])^(3/2)) + (a^2*(A*(3 - 8*m + 4*m^2) + C*(19 + 24*m + 4*m^2))*Cos[e + f*x]*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (a + a*Sin[e + f*x])/(2*a)]*(a + a*Sin[e + f*x])^m)/(2*c*f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]))/(16*a^2*c)`

### 3.6.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 78 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3146 `Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1)/2], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])`

rule 3224 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*Sin[e + f*x])^FracPart[m]*((c + d*Sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*Sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])`

---

3.6.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx$



```
rule 3451 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(
a*f*(2*m + 1))), x] + Simp[(a*B*(m - n) + A*b*(m + n + 1))/(a*b*(2*m + 1))
  Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[
{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0
] && (LtQ[m, -2^(-1)] || (ILtQ[m + n, 0] && !SumSimplerQ[n, 1])) && NeQ[2*
m + 1, 0]
```

```
rule 3515 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] :=
Simp[(a*A + a*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^
(n + 1)/(2*b*c*f*(2*m + 1))), x] - Simp[1/(2*b*c*d*(2*m + 1))  Int[(a + b*
Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[A*(c^2*(m + 1) + d^2*(2*m
+ n + 2)) - C*(c^2*m - d^2*(n + 1)) + d*(A*c*(m + n + 2) - c*C*(3*m - n))*
Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ
[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (EqQ[m + n + 2,
0] && NeQ[2*m + 1, 0]))
```

### 3.6.4 Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + C(\sin^2(fx + e)))}{(c - c \sin(fx + e))^{\frac{5}{2}}} dx$$

```
input int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x)
```

```
output int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x)
```

### 3.6.5 Fracas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin(fx + e)^2 + A)(a \sin(fx + e) + a)^m}{(-c \sin(fx + e) + c)^{\frac{5}{2}}} dx$$

```
input integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x,
algorithm="fricas")
```

---

3.6.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx$

output `integral((C*cos(f*x + e)^2 - A - C)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(3*c^3*cos(f*x + e)^2 - 4*c^3 - (c^3*cos(f*x + e)^2 - 4*c^3)*sin(f*x + e)), x)`

### 3.6.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(A+C*sin(f*x+e)**2)/(c-c*sin(f*x+e))**(5/2), x)`

output `Timed out`

### 3.6.7 Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{(-c \sin(fx + e) + c)^{5/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2), x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/(-c*sin(f*x + e) + c)^(5/2), x)`

### 3.6.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x,  
algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:Error index.cc index\_gcd Error: Bad  
Argument Value

### 3.6.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin(e + fx)^2 + A) (a + a \sin(e + fx))^m}{(c - c \sin(e + fx))^{5/2}} dx$$

input `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c - c*sin(e + f*x))^(  
5/2),x)`

output `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c - c*sin(e + f*x))^(  
5/2), x)`

$$3.7 \quad \int \frac{A+C \sin^2(e+fx)}{\sqrt{a+a \sin(e+fx)}(c-c \sin(e+fx))^{3/2}} dx$$

3.7.1	Optimal result . . . . .	83
3.7.2	Mathematica [A] (verified) . . . . .	83
3.7.3	Rubi [A] (verified) . . . . .	84
3.7.4	Maple [B] (verified) . . . . .	87
3.7.5	Fricas [F] . . . . .	88
3.7.6	Sympy [F] . . . . .	88
3.7.7	Maxima [F] . . . . .	88
3.7.8	Giac [A] (verification not implemented) . . . . .	89
3.7.9	Mupad [F(-1)] . . . . .	89

### 3.7.1 Optimal result

Integrand size = 42, antiderivative size = 167

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \frac{(A + C) \cos(e + fx) \sqrt{a + a \sin(e + fx)}}{4af(c - c \sin(e + fx))^{3/2}} - \frac{(A - 3C) \cos(e + fx) \log(1 - \sin(e + fx))}{4cf \sqrt{a + a \sin(e + fx)} \sqrt{c - c \sin(e + fx)}} + \frac{(A + C) \cos(e + fx) \log(1 + \sin(e + fx))}{4cf \sqrt{a + a \sin(e + fx)} \sqrt{c - c \sin(e + fx)}}$$

output `1/4*(A+C)*cos(f*x+e)*(a+a*sin(f*x+e))^(1/2)/a/f/(c-c*sin(f*x+e))^(3/2)-1/4*(A-3*C)*cos(f*x+e)*ln(1-sin(f*x+e))/c/f/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f*x+e))^(1/2)+1/4*(A+C)*cos(f*x+e)*ln(1+sin(f*x+e))/c/f/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f*x+e))^(1/2)`

### 3.7.2 Mathematica [A] (verified)

Time = 2.96 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.14

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \frac{(A + C - (A - 3C) \log(\cos(\frac{1}{2}(e + fx)) - \sin(\frac{1}{2}(e + fx)))}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}}$$

input `Integrate[(A + C*Sin[e + f*x]^2)/(Sqrt[a + a*Sin[e + f*x]]*(c - c*Sin[e + f*x])^(3/2)),x]`

output  $((A + C - (A - 3C) \cdot \text{Log}[\text{Cos}[(e + f \cdot x)/2] - \text{Sin}[(e + f \cdot x)/2]] \cdot (\text{Cos}[(e + f \cdot x)/2] - \text{Sin}[(e + f \cdot x)/2])^2 + (A + C) \cdot \text{Log}[\text{Cos}[(e + f \cdot x)/2] + \text{Sin}[(e + f \cdot x)/2]] \cdot (\text{Cos}[(e + f \cdot x)/2] - \text{Sin}[(e + f \cdot x)/2])^2) \cdot (\text{Cos}[(e + f \cdot x)/2] - \text{Sin}[(e + f \cdot x)/2]) \cdot (\text{Cos}[(e + f \cdot x)/2] + \text{Sin}[(e + f \cdot x)/2])) / (2 \cdot f \cdot \text{Sqrt}[a \cdot (1 + \text{Sin}[e + f \cdot x])] \cdot (c - c \cdot \text{Sin}[e + f \cdot x])^{(3/2)})$

### 3.7.3 Rubi [A] (verified)

Time = 1.12 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {3042, 3515, 27, 3042, 3448, 3042, 3216, 3042, 3146, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a \sin(e + fx) + a(c - c \sin(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{A + C \sin(e + fx)^2}{\sqrt{a \sin(e + fx) + a(c - c \sin(e + fx))^{3/2}} dx$$

↓ 3515

$$\frac{(A + C) \cos(e + fx) \sqrt{a \sin(e + fx) + a}}{4af(c - c \sin(e + fx))^{3/2}} - \frac{\int -\frac{2(a^2(A - C) - 2a^2C \sin(e + fx))}{\sqrt{\sin(e + fx)a + a\sqrt{c - c \sin(e + fx)}} dx}{4a^2c}$$

↓ 27

$$\frac{\int \frac{a^2(A - C) - 2a^2C \sin(e + fx)}{\sqrt{\sin(e + fx)a + a\sqrt{c - c \sin(e + fx)}} dx}{2a^2c} + \frac{(A + C) \cos(e + fx) \sqrt{a \sin(e + fx) + a}}{4af(c - c \sin(e + fx))^{3/2}}$$

↓ 3042

$$\frac{\int \frac{a^2(A - C) - 2a^2C \sin(e + fx)}{\sqrt{\sin(e + fx)a + a\sqrt{c - c \sin(e + fx)}} dx}{2a^2c} + \frac{(A + C) \cos(e + fx) \sqrt{a \sin(e + fx) + a}}{4af(c - c \sin(e + fx))^{3/2}}$$

↓ 3448

$$\frac{\frac{a^2(A + C) \int \frac{\sqrt{c - c \sin(e + fx)}}{\sqrt{\sin(e + fx)a + a}} dx}{2c} + \frac{1}{2}a(A - 3C) \int \frac{\sqrt{\sin(e + fx)a + a}}{\sqrt{c - c \sin(e + fx)}} dx}{2a^2c} + \frac{(A + C) \cos(e + fx) \sqrt{a \sin(e + fx) + a}}{4af(c - c \sin(e + fx))^{3/2}}$$

---

3.7.  $\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)(c - c \sin(e + fx))^{3/2}} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{a^2(A+C) \int \frac{\sqrt{c-c \sin(e+fx)}}{\sqrt{\sin(e+fx)a+a}} dx + \frac{1}{2}a(A-3C) \int \frac{\sqrt{\sin(e+fx)a+a}}{\sqrt{c-c \sin(e+fx)}} dx}{\frac{2a^2c}{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}} + \frac{4af(c-c \sin(e+fx))^{3/2}}{}} \\
& \downarrow 3216 \\
& \frac{a^3(A+C) \cos(e+fx) \int \frac{\cos(e+fx)}{\sin(e+fx)a+a} dx + \frac{a^2c(A-3C) \cos(e+fx) \int \frac{\cos(e+fx)}{c-c \sin(e+fx)} dx}{\frac{2\sqrt{a \sin(e+fx)+a} \sqrt{c-c \sin(e+fx)}}{2a^2c}} + \frac{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c \sin(e+fx))^{3/2}}}{\frac{2\sqrt{a \sin(e+fx)+a} \sqrt{c-c \sin(e+fx)}}{2a^2c}} + \frac{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c \sin(e+fx))^{3/2}} \\
& \downarrow 3042 \\
& \frac{a^3(A+C) \cos(e+fx) \int \frac{\cos(e+fx)}{\sin(e+fx)a+a} dx + \frac{a^2c(A-3C) \cos(e+fx) \int \frac{\cos(e+fx)}{c-c \sin(e+fx)} dx}{\frac{2\sqrt{a \sin(e+fx)+a} \sqrt{c-c \sin(e+fx)}}{2a^2c}} + \frac{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c \sin(e+fx))^{3/2}}}{\frac{2\sqrt{a \sin(e+fx)+a} \sqrt{c-c \sin(e+fx)}}{2a^2c}} + \frac{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c \sin(e+fx))^{3/2}} \\
& \downarrow 3146 \\
& \frac{a^2(A+C) \cos(e+fx) \int \frac{1}{\sin(e+fx)a+a} d(a \sin(e+fx)) - \frac{a^2(A-3C) \cos(e+fx) \int \frac{1}{c-c \sin(e+fx)} d(-c \sin(e+fx))}{\frac{2f \sqrt{a \sin(e+fx)+a} \sqrt{c-c \sin(e+fx)}}{2f \sqrt{a \sin(e+fx)+a} \sqrt{c-c \sin(e+fx)}}} + \frac{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c \sin(e+fx))^{3/2}}}{\frac{2f \sqrt{a \sin(e+fx)+a} \sqrt{c-c \sin(e+fx)}}{2f \sqrt{a \sin(e+fx)+a} \sqrt{c-c \sin(e+fx)}}} + \frac{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c \sin(e+fx))^{3/2}} \\
& \downarrow 16 \\
& \frac{a^2(A+C) \cos(e+fx) \log(a \sin(e+fx)+a) - \frac{a^2(A-3C) \cos(e+fx) \log(c-c \sin(e+fx))}{\frac{2f \sqrt{a \sin(e+fx)+a} \sqrt{c-c \sin(e+fx)}}{2f \sqrt{a \sin(e+fx)+a} \sqrt{c-c \sin(e+fx)}}} + \frac{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c \sin(e+fx))^{3/2}}}{\frac{2f \sqrt{a \sin(e+fx)+a} \sqrt{c-c \sin(e+fx)}}{2f \sqrt{a \sin(e+fx)+a} \sqrt{c-c \sin(e+fx)}}} + \frac{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c \sin(e+fx))^{3/2}}
\end{aligned}$$

input `Int[(A + C*Sin[e + f*x]^2)/(Sqrt[a + a*Sin[e + f*x]]*(c - c*Sin[e + f*x])^(3/2)),x]`

output `((A + C)*Cos[e + f*x]*Sqrt[a + a*Sin[e + f*x]]/(4*a*f*(c - c*Sin[e + f*x])^(3/2)) + ((a^2*(A + C)*Cos[e + f*x]*Log[a + a*Sin[e + f*x]])/(2*f*Sqrt[a + a*Sin[e + f*x]]*Sqrt[c - c*Sin[e + f*x]]) - (a^2*(A - 3*C)*Cos[e + f*x]*Log[c - c*Sin[e + f*x]])/(2*f*Sqrt[a + a*Sin[e + f*x]]*Sqrt[c - c*Sin[e + f*x]]))/(2*a^2*c)`

$$3.7. \int \frac{A+C \sin^2(e+fx)}{\sqrt{a+a \sin(e+fx)}(c-c \sin(e+fx))^{3/2}} dx$$

## 3.7.3.1 Defintions of rubi rules used

- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3146 `Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1)/2, x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])`
- rule 3216 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[a*c*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])) Int[Cos[e + f*x]/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`
- rule 3448 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A*b + a*B)/(2*a*b) Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[(B*c + A*d)/(2*c*d) Int[Sqrt[c + d*Sin[e + f*x]]/Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

```
rule 3515 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] :=
Simp[(a*A + a*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^
(n + 1)/(2*b*c*f*(2*m + 1))), x] - Simp[1/(2*b*c*d*(2*m + 1)) Int[(a + b*
Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[A*(c^2*(m + 1) + d^2*(2*m
+ n + 2)) - C*(c^2*m - d^2*(n + 1)) + d*(A*c*(m + n + 2) - c*C*(3*m - n))*
Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ
[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (EqQ[m + n + 2,
0] && NeQ[2*m + 1, 0]))
```

### 3.7.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 540 vs.  $2(149) = 298$ .

Time = 3.85 (sec) , antiderivative size = 541, normalized size of antiderivative = 3.24

method	result
default	$\frac{A \sin(fx+e) \cos(fx+e) \ln(-\cot(fx+e)+\csc(fx+e)+1)-A \ln(\csc(fx+e)-\cot(fx+e)-1) \sin(fx+e) \cos(fx+e)-A(\cos^2(fx+e)) \ln(\csc(fx+e)-\cot(fx+e)-1)}{A((\cos^2(fx+e)) \ln(\csc(fx+e)-\cot(fx+e)-1)-\cos(fx+e) \sin(fx+e) \ln(\csc(fx+e)-\cot(fx+e)-1)-(\cos^2(fx+e)) \ln(-\cot(fx+e)))}$
parts	

```
input int((A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2)/(a+a*sin(f*x+e))^(1/2),x,met
hod=_RETURNVERBOSE)
```

```
output 1/2/c/f*(A*sin(f*x+e)*cos(f*x+e)*ln(-cot(f*x+e)+csc(f*x+e)+1)-A*ln(csc(f*x
+e)-cot(f*x+e)-1)*sin(f*x+e)*cos(f*x+e)-A*cos(f*x+e)^2*ln(-cot(f*x+e)+csc(
f*x+e)+1)+A*cos(f*x+e)^2*ln(csc(f*x+e)-cot(f*x+e)-1)+C*ln(-cot(f*x+e)+csc(
f*x+e)+1)*sin(f*x+e)*cos(f*x+e)+3*C*ln(csc(f*x+e)-cot(f*x+e)-1)*sin(f*x+e)
*cos(f*x+e)-2*C*ln(2/(1+cos(f*x+e)))*sin(f*x+e)*cos(f*x+e)-C*cos(f*x+e)^2*
ln(-cot(f*x+e)+csc(f*x+e)+1)-3*C*cos(f*x+e)^2*ln(csc(f*x+e)-cot(f*x+e)-1)+
2*C*cos(f*x+e)^2*ln(2/(1+cos(f*x+e)))-A*sin(f*x+e)*cos(f*x+e)+A*cos(f*x+e)
^2-A*cos(f*x+e)*ln(-cot(f*x+e)+csc(f*x+e)+1)+A*cos(f*x+e)*ln(csc(f*x+e)-co
t(f*x+e)-1)-C*sin(f*x+e)*cos(f*x+e)+C*cos(f*x+e)^2-C*cos(f*x+e)*ln(-cot(f*
x+e)+csc(f*x+e)+1)-3*C*cos(f*x+e)*ln(csc(f*x+e)-cot(f*x+e)-1)+2*C*cos(f*x+
e)*ln(2/(1+cos(f*x+e)))-A*sin(f*x+e)-C*sin(f*x+e)-A-C)/(-cos(f*x+e)+sin(f*
x+e)-1)/(-c*(sin(f*x+e)-1))^(1/2)/(a*(1+sin(f*x+e)))^(1/2)
```

$$3.7. \int \frac{A+C \sin^2(e+fx)}{\sqrt{a+a \sin(e+fx)}(c-c \sin(e+fx))^{3/2}} dx$$



### 3.7.5 Fricas [F]

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{C \sin(fx + e)^2 + A}{\sqrt{a \sin(fx + e) + a}(-c \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2)/(a+a*sin(f*x+e))^(1/2),x, algorithm="fricas")`

output `integral((C*cos(f*x + e)^2 - A - C)*sqrt(a*sin(f*x + e) + a)*sqrt(-c*sin(f*x + e) + c)/(a*c^2*cos(f*x + e)^2*sin(f*x + e) - a*c^2*cos(f*x + e)^2), x)`

### 3.7.6 Sympy [F]

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{A + C \sin^2(e + fx)}{\sqrt{a(\sin(e + fx) + 1)}(-c(\sin(e + fx) - 1))^{3/2}} dx$$

input `integrate((A+C*sin(f*x+e)**2)/(c-c*sin(f*x+e))**(3/2)/(a+a*sin(f*x+e))**(1/2),x)`

output `Integral((A + C*sin(e + f*x)**2)/(sqrt(a*(sin(e + f*x) + 1))*(-c*(sin(e + f*x) - 1))**(3/2)), x)`

### 3.7.7 Maxima [F]

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{C \sin(fx + e)^2 + A}{\sqrt{a \sin(fx + e) + a}(-c \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2)/(a+a*sin(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)/(sqrt(a*sin(f*x + e) + a)*(-c*sin(f*x + e) + c)^(3/2)), x)`

### 3.7.8 Giac [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.18

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx =$$

$$\frac{2(A+C) \log(|\cos(-\frac{1}{4}\pi + \frac{1}{2}fx + \frac{1}{2}e)|)}{\sqrt{ac}^{\frac{3}{2}} \operatorname{sgn}(\cos(-\frac{1}{4}\pi + \frac{1}{2}fx + \frac{1}{2}e)) \operatorname{sgn}(\sin(-\frac{1}{4}\pi + \frac{1}{2}fx + \frac{1}{2}e))} - \frac{(A\sqrt{a} - 3C\sqrt{a}) \log(-\cos(-\frac{1}{4}\pi + \frac{1}{2}fx + \frac{1}{2}e)^2 + 1)}{ac^{\frac{3}{2}} \operatorname{sgn}(\cos(-\frac{1}{4}\pi + \frac{1}{2}fx + \frac{1}{2}e)) \operatorname{sgn}(\sin(-\frac{1}{4}\pi + \frac{1}{2}fx + \frac{1}{2}e))} - \frac{1}{(\cos(-\frac{1}{4}\pi + \frac{1}{2}fx + \frac{1}{2}e))^{3/2}} - \frac{1}{4f}$$

input `integrate((A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2)/(a+a*sin(f*x+e))^(1/2),x, algorithm="giac")`

output `-1/4*(2*(A + C)*log(abs(cos(-1/4*pi + 1/2*f*x + 1/2*e)))/(sqrt(a)*c^(3/2)*sgn(cos(-1/4*pi + 1/2*f*x + 1/2*e))*sgn(sin(-1/4*pi + 1/2*f*x + 1/2*e))) - (A*sqrt(a) - 3*C*sqrt(a))*log(-cos(-1/4*pi + 1/2*f*x + 1/2*e)^2 + 1)/(a*c^(3/2)*sgn(cos(-1/4*pi + 1/2*f*x + 1/2*e))*sgn(sin(-1/4*pi + 1/2*f*x + 1/2*e))) - (A*sqrt(a) + C*sqrt(a))/((cos(-1/4*pi + 1/2*f*x + 1/2*e)^2 - 1)*a*c^(3/2)*sgn(cos(-1/4*pi + 1/2*f*x + 1/2*e))*sgn(sin(-1/4*pi + 1/2*f*x + 1/2*e))))/f`

### 3.7.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{C \sin(e + fx)^2 + A}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx$$

input `int((A + C*sin(e + f*x)^2)/((a + a*sin(e + f*x))^(1/2)*(c - c*sin(e + f*x))^(3/2)),x)`

output `int((A + C*sin(e + f*x)^2)/((a + a*sin(e + f*x))^(1/2)*(c - c*sin(e + f*x))^(3/2)), x)`

### 3.8 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^n (A + C \sin^2(e + fx)) dx$

3.8.1	Optimal result	90
3.8.2	Mathematica [F]	91
3.8.3	Rubi [A] (verified)	91
3.8.4	Maple [F]	95
3.8.5	Fricas [F]	95
3.8.6	Sympy [F]	96
3.8.7	Maxima [F]	96
3.8.8	Giac [F]	96
3.8.9	Mupad [F(-1)]	97

#### 3.8.1 Optimal result

Integrand size = 38, antiderivative size = 257

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= \frac{2^{\frac{1}{2}+n} c (C(1 + 2m)(m - n) + (1 + m + n)(C(1 - m + n) + A(2 + m + n))) \cos(e + fx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2} + n, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2} + \frac{1}{2} \sin(fx + e)\right) (1 - \sin(fx + e))^{1/2-n} (a + a \sin(fx + e))^m (c - c \sin(fx + e))^{-1+n}}{f(1 + m + n)(2 + m + n)}$$

$$- \frac{C(1 + 2m) \cos(e + fx) (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n}{f(1 + m + n)(2 + m + n)}$$

$$+ \frac{C \cos(e + fx) (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{1+n}}{cf(2 + m + n)}$$

```
output 2^(1/2+n)*c*(C*(1+2*m)*(m-n)+(1+m+n)*(C*(1-m+n)+A*(2+m+n)))*cos(f*x+e)*hypergeom([1/2-n, 1/2+m], [3/2+m], 1/2+1/2*sin(f*x+e))*(1-sin(f*x+e))^(1/2-n)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(-1+n)/f/(1+2*m)/(1+m+n)/(2+m+n)-C*(1+2*m)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n/f/(1+m+n)/(2+m+n)+C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1+n)/c/f/(2+m+n)
```

### 3.8.2 Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

input `Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n*(A + C*Sin[e + f*x]^2),x]`

output `Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n*(A + C*Sin[e + f*x]^2), x]`

### 3.8.3 Rubi [A] (verified)

Time = 1.18 (sec) , antiderivative size = 306, normalized size of antiderivative = 1.19, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$ , Rules used = {3042, 3519, 25, 3042, 3452, 3042, 3224, 3042, 3168, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) (c - c \sin(e + fx))^n dx$$

$$\downarrow \text{3042}$$

$$\int (a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2) (c - c \sin(e + fx))^n dx$$

$$\downarrow \text{3519}$$

$$\frac{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{n+1}}{cf(m + n + 2)} - \frac{\int -(\sin(e + fx)a + a)^m (c - c \sin(e + fx))^n (ac(C(-m + n + 1) + A(m + n + 2)) + acC(2m + 1) \sin(e + fx)) dx}{ac(m + n + 2)}$$

$$\downarrow \text{25}$$

$$\frac{\int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^n (ac(C(-m + n + 1) + A(m + n + 2)) + acC(2m + 1) \sin(e + fx)) dx}{ac(m + n + 2)} + \frac{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{n+1}}{cf(m + n + 2)}$$

---

3.8.  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^m (c - c\sin(e + fx))^n (ac(C(-m + n + 1) + A(m + n + 2)) + acC(2m + 1)\sin(e + fx)) dx}{\frac{ac(m + n + 2)}{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c\sin(e + fx))^{n+1}} \frac{cf(m + n + 2)}}{cf(m + n + 2)}$$

↓ 3452

$$\frac{\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+C(2m+1)(m-n))}{m+n+1} \int (\sin(e+fx)a+a)^m (c-c\sin(e+fx))^n dx - \frac{acC(2m+1) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(m+n+1)}}{\frac{ac(m + n + 2)}{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c\sin(e + fx))^{n+1}} \frac{cf(m + n + 2)}}{cf(m + n + 2)}}$$

↓ 3042

$$\frac{\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+C(2m+1)(m-n))}{m+n+1} \int (\sin(e+fx)a+a)^m (c-c\sin(e+fx))^n dx - \frac{acC(2m+1) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(m+n+1)}}{\frac{ac(m + n + 2)}{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c\sin(e + fx))^{n+1}} \frac{cf(m + n + 2)}}{cf(m + n + 2)}}$$

↓ 3224

$$\frac{\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+C(2m+1)(m-n)) \cos^{-2m}(e+fx)(a \sin(e+fx)+a)^m (c-c\sin(e+fx))^m}{m+n+1} \int \cos^{2m}(e+fx)(c-c\sin(e+fx))^{n+1} dx}{\frac{ac(m + n + 2)}{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c\sin(e + fx))^{n+1}} \frac{cf(m + n + 2)}}{cf(m + n + 2)}}$$

↓ 3042

$$\frac{\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+C(2m+1)(m-n)) \cos^{-2m}(e+fx)(a \sin(e+fx)+a)^m (c-c\sin(e+fx))^m}{m+n+1} \int \cos(e+fx)^{2m}(c-c\sin(e+fx))^{n+1} dx}{\frac{ac(m + n + 2)}{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c\sin(e + fx))^{n+1}} \frac{cf(m + n + 2)}}{cf(m + n + 2)}}$$

↓ 3168

$$\frac{\frac{ac^3((m+n+1)(A(m+n+2)+C(-m+n+1))+C(2m+1)(m-n)) \cos(e+fx)(a \sin(e+fx)+a)^m (c-c\sin(e+fx))^{\frac{1}{2}(-2m-1)+m} (c \sin(e+fx)+c)^{\frac{1}{2}(-2m-1)+m}}{f(m+n+1)}}{\frac{ac(m + n + 2)}{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c\sin(e + fx))^{n+1}} \frac{cf(m + n + 2)}}{cf(m + n + 2)}}$$

↓ 80

---

3.8.  $\int (a + a \sin(e + fx))^m (c - c\sin(e + fx))^n (A + C \sin^2(e + fx)) dx$

$$\frac{ac^3 2^{n-\frac{1}{2}}((m+n+1)(A(m+n+2)+C(-m+n+1))+C(2m+1)(m-n)) \cos(e+fx)(1-\sin(e+fx))^{\frac{1}{2}-n} (a \sin(e+fx)+a)^m (c \sin(e+fx)+c)^{\frac{1}{2}(-2m-1)}}{f(m+n+1)}$$

$$\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c \sin(e+fx))^{n+1}}{cf(m+n+2)}$$

↓ 79

$$\frac{ac^2 2^{n+\frac{1}{2}}((m+n+1)(A(m+n+2)+C(-m+n+1))+C(2m+1)(m-n)) \cos(e+fx)(1-\sin(e+fx))^{\frac{1}{2}-n} (a \sin(e+fx)+a)^m (c \sin(e+fx)+c)^{\frac{1}{2}(-2m-1)}}{f(2m+1)(m+n+1)}$$

$$\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c \sin(e+fx))^{n+1}}{cf(m+n+2)}$$

input `Int[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n*(A + C*Sin[e + f*x]^2),x]`

output `(C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(1 + n))/(c*f*(2 + m + n)) + (-((a*c*C*(1 + 2*m)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n)/(f*(1 + m + n))) + (2^(1/2 + n)*a*c^2*(C*(1 + 2*m)*(m - n) + (1 + m + n)*(C*(1 - m + n) + A*(2 + m + n)))*Cos[e + f*x]*Hypergeometric2F1[(1 + 2*m)/2, (1 - 2*n)/2, (3 + 2*m)/2, (1 + Sin[e + f*x])/2]*(1 - Sin[e + f*x])^(1/2 - n)*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(-1/2 + (-1 - 2*m)/2 + m + n)*(c + c*Sin[e + f*x])^((-1 - 2*m)/2 + (1 + 2*m)/2)))/(f*(1 + 2*m)*(1 + m + n))/(a*c*(2 + m + n))`

### 3.8.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 79 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))`

- rule 80 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n, x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3168 `Int[(cos[(e_.) + (f_.)*(x_)]*(g_.))^(p_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Simp[a^2*((g*cos[e + f*x])^(p + 1)/(f*g*(a + b*sin[e + f*x])^((p + 1)/2)*(a - b*sin[e + f*x])^((p + 1)/2))) Subst[Int[(a + b*x)^(m + (p - 1)/2)*(a - b*x)^((p - 1)/2), x], x, Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, g, m, p}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[m]`
- rule 3224 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*sin[e + f*x])^FracPart[m]*((c + d*sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])`
- rule 3452 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*sin[e + f*x])^m*((c + d*sin[e + f*x])^n/(f*(m + n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(a + b*sin[e + f*x])^m*(c + d*sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 1, 0]`

```
rule 3519 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)^2), x_Symbol] :>
Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)
)/(d*f*(m + n + 2)), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*
x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)
) - b*c*C*(2*m + 1)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A
, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1
)] && NeQ[m + n + 2, 0]
```

### 3.8.4 Maple [F]

$$\int (a + a \sin(fx + e))^m (c - c \sin(fx + e))^n (A + C(\sin^2(fx + e))) dx$$

```
input int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x)
```

```
output int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x)
```

### 3.8.5 Fracas [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx \\ &= \int (C \sin(fx + e)^2 + A) (a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^n dx \end{aligned}$$

```
input integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x, algo
rithm="fracas")
```

```
output integral(-(C*cos(f*x + e)^2 - A - C)*(a*sin(f*x + e) + a)^m*(-c*sin(f*x +
e) + c)^n, x)
```



**3.8.6 Sympy [F]**

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= \int (a(\sin(e + fx) + 1))^m (-c(\sin(e + fx) - 1))^n (A + C \sin^2(e + fx)) dx$$

input `integrate((a+a*sin(f*x+e))**m*(c-c*sin(f*x+e))**n*(A+C*sin(f*x+e)**2),x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(-c*(sin(e + f*x) - 1))**n*(A + C*sin(e + f*x)**2), x)`

**3.8.7 Maxima [F]**

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^n dx$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m*(-c*sin(f*x + e) + c)^n, x)`

**3.8.8 Giac [F]**

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^n dx$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m*(-c*sin(f*x + e) + c)^n, x)`

**3.8.9 Mupad [F(-1)]**

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(e + fx)^2 + A) (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n dx$$

input `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^n,x)`

output `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^n,x)`

### 3.9 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^n (A + C \sin^2(e +$

3.9.1	Optimal result	98
3.9.2	Mathematica [F]	99
3.9.3	Rubi [A] (verified)	99
3.9.4	Maple [F]	103
3.9.5	Fricas [F]	103
3.9.6	Sympy [F(-1)]	104
3.9.7	Maxima [F]	104
3.9.8	Giac [F]	104
3.9.9	Mupad [F(-1)]	105

#### 3.9.1 Optimal result

Integrand size = 37, antiderivative size = 366

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= -\frac{C \cos(e + fx)(a + a \sin(e + fx))^m (c + d \sin(e + fx))^{1+n}}{df(2 + m + n)}$$

$$+ \frac{\sqrt{2}(c(C + 2Cm) + d(C(1 - m + n) + A(2 + m + n))) \operatorname{AppellF1}\left(\frac{1}{2} + m, \frac{1}{2}, -n, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e + fx))\right)}{df(1 + 2m)(2 + m + n)\sqrt{1 - \sin(e + fx)}}$$

$$+ \frac{\sqrt{2}C(dm - c(1 + m)) \operatorname{AppellF1}\left(\frac{3}{2} + m, \frac{1}{2}, -n, \frac{5}{2} + m, \frac{1}{2}(1 + \sin(e + fx))\right), -\frac{d(1 + \sin(e + fx))}{c - d}}{adf(3 + 2m)(2 + m + n)\sqrt{1 - \sin(e + fx)}}$$

output

```
-C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1+n)/d/f/(2+m+n)+(c*(2*C*m+C)+d*(C*(1-m+n)+A*(2+m+n))*AppellF1(1/2+m,-n,1/2,3/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*2^(1/2)/d/f/(1+2*m)/(2+m+n)/(((c+d*sin(f*x+e))/(c-d))^n)/(1-sin(f*x+e))^(1/2)+C*(d*m-c*(1+m))*AppellF1(3/2+m,-n,1/2,5/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)*(c+d*sin(f*x+e))^n*2^(1/2)/a/d/f/(3+2*m)/(2+m+n)/(((c+d*sin(f*x+e))/(c-d))^n)/(1-sin(f*x+e))^(1/2)
```

### 3.9.2 Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

input `Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + C*Sin[e + f*x]^2),x]`

output `Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + C*Sin[e + f*x]^2), x]`

### 3.9.3 Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 388, normalized size of antiderivative = 1.06, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.270$ , Rules used = {3042, 3525, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) (c + d \sin(e + fx))^n dx$$

$$\downarrow \text{3042}$$

$$\int (a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2) (c + d \sin(e + fx))^n dx$$

$$\downarrow \text{3525}$$

$$\frac{\int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^n (a(Ad(m + n + 2) + C(nd + d + cm)) + aC(dm - c(m + 1)) \sin(e + fx) + \frac{ad(m + n + 2)}{df(m + n + 2)} C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{n+1}) dx}{df(m + n + 2)}$$

$$\downarrow \text{3042}$$

$$\frac{\int (\sin(e+fx)a+a)^m (c+d\sin(e+fx))^n (a(Ad(m+n+2)+C(nd+d+cm)) + aC(dm-c(m+1)) \sin(e+fx) dx}{\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d\sin(e+fx))^{n+1}}{df(m+n+2)}} \quad \downarrow \quad 3466$$

$$\frac{a(Ad(m+n+2)+c(2Cm+C)+Cd(-m+n+1)) \int (\sin(e+fx)a+a)^m (c+d\sin(e+fx))^n dx + C(dm-c(m+1)) \sin(e+fx) dx}{\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d\sin(e+fx))^{n+1}}{df(m+n+2)}} \quad \downarrow \quad 3042$$

$$\frac{a(Ad(m+n+2)+c(2Cm+C)+Cd(-m+n+1)) \int (\sin(e+fx)a+a)^m (c+d\sin(e+fx))^n dx + C(dm-c(m+1)) \sin(e+fx) dx}{\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d\sin(e+fx))^{n+1}}{df(m+n+2)}} \quad \downarrow \quad 3267$$

$$\frac{a^3 \cos(e+fx)(Ad(m+n+2)+c(2Cm+C)+Cd(-m+n+1)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} (c+d\sin(e+fx))^n d\sin(e+fx)}{f\sqrt{a-a\sin(e+fx)}\sqrt{a\sin(e+fx)+a}} + \frac{a^2 C(dm-c(m+1)) \cos(e+fx) dx}{f\sqrt{a-a\sin(e+fx)}}}{\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d\sin(e+fx))^{n+1}}{df(m+n+2)}} \quad \downarrow \quad 157$$

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)(Ad(m+n+2)+c(2Cm+C)+Cd(-m+n+1)) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m-\frac{1}{2}} (c+d\sin(e+fx))^n d\sin(e+fx)}{\sqrt{2}f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}} + \frac{a^2 C(dm-c(m+1)) \cos(e+fx) dx}{f\sqrt{1-\sin(e+fx)}}}{\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d\sin(e+fx))^{n+1}}{df(m+n+2)}} \quad \downarrow \quad 27$$

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)(Ad(m+n+2)+c(2Cm+C)+Cd(-m+n+1)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} (c+d\sin(e+fx))^n d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}} + \frac{a^2 C(dm-c(m+1)) \cos(e+fx) dx}{f\sqrt{1-\sin(e+fx)}}}{\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d\sin(e+fx))^{n+1}}{df(m+n+2)}} \quad \downarrow \quad 156$$

---

3.9.  $\int (a+a\sin(e+fx))^m (c+d\sin(e+fx))^n (A+C\sin^2(e+fx)) dx$

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (Ad(m+n+2)+c(2Cm+C)+Cd(-m+n+1))(c+d \sin(e+fx))^n \left(\frac{c+d \sin(e+fx)}{c-d}\right)^{-n} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} \left(\frac{c}{c-d} + \frac{d \sin(e+fx)}{c}\right)}{\sqrt{1-\sin(e+fx)}}}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}}$$

$$\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{n+1}}{df(m+n+2)}$$

↓ 155

$$\frac{\sqrt{2}a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx)(a \sin(e+fx)+a)^m (Ad(m+n+2)+c(2Cm+C)+Cd(-m+n+1))(c+d \sin(e+fx))^n \left(\frac{c+d \sin(e+fx)}{c-d}\right)^{-n} \text{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -n, \frac{3}{2}+m, \frac{1+\sin(e+fx)}{2}, -\left(\frac{d(1+\sin(e+fx))}{c-d}\right)\right)}{f(2m+1)(a-a \sin(e+fx))}$$

$$\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{n+1}}{df(m+n+2)}$$

input `Int[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + C*Sin[e + f*x]^2),x]`

output `-((C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(1 + n))/(d*f*(2 + m + n))) + ((Sqrt[2]*a^2*(c*(C + 2*C*m) + C*d*(1 - m + n) + A*d*(2 + m + n))*AppellF1[1/2 + m, 1/2, -n, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n)/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*((c + d*Sin[e + f*x])/(c - d))^n) + (Sqrt[2]*a*C*(d*m - c*(1 + m))*AppellF1[3/2 + m, 1/2, -n, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*(c + d*Sin[e + f*x])^n)/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*((c + d*Sin[e + f*x])/(c - d))^n))/(a*d*(2 + m + n))`

### 3.9.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplifierQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplifierQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplifierQ[c + d*x, a + b*x] && !SimplifierQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3267 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]])*Sqrt[a - b*Sin[e + f*x]]) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]`

```
rule 3466 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x]
+ Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x]
, x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ
[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]
```

```
rule 3525 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] :=
Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)
)/(d*f*(m + n + 2)), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*
x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)
) + C*(a*d*m - b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d,
e, f, A, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2
- d^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]
```

### 3.9.4 Maple [F]

$$\int (a + a \sin(fx + e))^m (c + d \sin(fx + e))^n (A + C(\sin^2(fx + e))) dx$$

```
input int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x)
```

```
output int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x)
```

### 3.9.5 Fracas [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx \\ &= \int (C \sin(fx + e)^2 + A)(a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^n dx \end{aligned}$$

```
input integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x, algo
rithm="fracas")
```

```
output integral(-(C*cos(f*x + e)^2 - A - C)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e)
) + c)^n, x)
```

---


$$3.9. \quad \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$



### 3.9.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(c+d*sin(f*x+e))**n*(A+C*sin(f*x+e)**2),x)`

output `Timed out`

### 3.9.7 Maxima [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx \\ &= \int (C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^n dx \end{aligned}$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e) + c)^n, x)`

### 3.9.8 Giac [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx \\ &= \int (C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^n dx \end{aligned}$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e) + c)^n, x)`

**3.9.9 Mupad [F(-1)]**

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(e + fx)^2 + A) (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n dx$$

input `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^n,x)`

output `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^n,x)`

### 3.10 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{-2-m} (A + C \sin^2$

3.10.1	Optimal result	106
3.10.2	Mathematica [F]	107
3.10.3	Rubi [A] (verified)	107
3.10.4	Maple [F]	112
3.10.5	Fricas [F]	112
3.10.6	Sympy [F(-1)]	113
3.10.7	Maxima [F]	113
3.10.8	Giac [F]	113
3.10.9	Mupad [F(-1)]	114

#### 3.10.1 Optimal result

Integrand size = 41, antiderivative size = 392

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx$$

$$= \frac{(c^2C + Ad^2) \cos(e + fx)(a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-1-m}}{d(c^2 - d^2) f(1 + m)}$$

$$- \frac{2^{\frac{1}{2}+m} a(c(A + C)d(1 + m) + d^2(C - Am + Cm) - c^2(C + 2Cm)) \cos(e + fx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, (c - d)d(c + d)}{(c - d)d(c + d)}\right)}{(c - d)d(c + d)}$$

$$+ \frac{\sqrt{2}C \operatorname{AppellF1}\left(\frac{3}{2} + m, \frac{1}{2}, 1 + m, \frac{5}{2} + m, \frac{1}{2}(1 + \sin(e + fx)), -\frac{d(1 + \sin(e + fx))}{c - d}\right) \cos(e + fx)(a + a \sin(e + fx))^{-1-m}}{a(c - d)df(3 + 2m)\sqrt{1 - \sin(e + fx)}}$$

output

```
(A*d^2+C*c^2)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1-m)/d/(c^2-d^2)/f/(1+m)-2^(1/2+m)*a*(c*(A+C)*d*(1+m)+d^2*(-A*m+C*m+C)-c^2*(2*C*m+C))*cos(f*x+e)*hypergeom([1/2, 1/2-m], [3/2], 1/2*(c-d)*(1-sin(f*x+e))/(c+d*sin(f*x+e)))*(a+a*sin(f*x+e))^(1-m)*((c+d)*(1+sin(f*x+e))/(c+d*sin(f*x+e)))^(1/2-m)/(c-d)/d/(c+d)^2/f/(1+m)/((c+d*sin(f*x+e))^m)+C*AppellF1(3/2+m, 1+m, 1/2, 5/2+m, -d*(1+sin(f*x+e))/(c-d), 1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)*((c+d*sin(f*x+e))/(c-d))^m*2^(1/2)/a/(c-d)/d/f/(3+2*m)/((c+d*sin(f*x+e))^m)/(1-sin(f*x+e))^(1/2)
```

### 3.10.2 Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx$$

input `Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(-2 - m)*(A + C*Sin[e + f*x]^2), x]`

output `Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(-2 - m)*(A + C*Sin[e + f*x]^2), x]`

### 3.10.3 Rubi [A] (verified)

Time = 1.29 (sec) , antiderivative size = 423, normalized size of antiderivative = 1.08, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.293$ , Rules used = {3042, 3523, 25, 3042, 3466, 3042, 3267, 142, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) (c + d \sin(e + fx))^{-m-2} dx$$

$$\downarrow \text{3042}$$

$$\int (a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2) (c + d \sin(e + fx))^{-m-2} dx$$

$$\downarrow \text{3523}$$

$$\frac{(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}}{df(m+1)(c^2 - d^2)} -$$

$$\frac{\int -(\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1} (a(Ad(mc + c - dm) + cC(md + d - cm)) + aC(c^2 - d^2)(m + 1))}{ad(m+1)(c^2 - d^2)}$$

$$\downarrow \text{25}$$

$$\frac{\int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1} (a(Ad(mc + c - dm) + cC(md + d - cm)) + aC(c^2 - d^2)(m + 1))}{ad(m+1)(c^2 - d^2)} -$$

$$\frac{(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}}{df(m+1)(c^2 - d^2)}$$

---

3.10.  $\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1} (a(Ad(mc + c - dm) + cC(md + d - cm)) + aC(c^2 - d^2)(m + 1))}{ad(m + 1)(c^2 - d^2)} \\ \frac{(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}}{df(m + 1)(c^2 - d^2)}$$

↓ 3466

$$\frac{a(cd(m + 1)(A + C) + d^2(-Am + Cm + C) - (c^2(2Cm + C))) \int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1} dx}{ad(m + 1)(c^2 - d^2)} \\ \frac{(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}}{df(m + 1)(c^2 - d^2)}$$

↓ 3042

$$\frac{a(cd(m + 1)(A + C) + d^2(-Am + Cm + C) - (c^2(2Cm + C))) \int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1} dx}{ad(m + 1)(c^2 - d^2)} \\ \frac{(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}}{df(m + 1)(c^2 - d^2)}$$

↓ 3267

$$\frac{a^3 \cos(e + fx)(cd(m + 1)(A + C) + d^2(-Am + Cm + C) - (c^2(2Cm + C))) \int \frac{(\sin(e + fx)a + a)^{m-\frac{1}{2}} (c + d \sin(e + fx))^{-m-1} d \sin(e + fx)}{\sqrt{a - a \sin(e + fx)}} + \frac{a^2 C(m + 1)(c^2 - d^2)}{f \sqrt{a - a \sin(e + fx)} \sqrt{a \sin(e + fx) + a}}}{ad(m + 1)(c^2 - d^2)} \\ \frac{(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}}{df(m + 1)(c^2 - d^2)}$$

↓ 142

$$\frac{a^2 C(m + 1)(c^2 - d^2) \cos(e + fx) \int \frac{(\sin(e + fx)a + a)^{m+\frac{1}{2}} (c + d \sin(e + fx))^{-m-1} d \sin(e + fx)}{\sqrt{a - a \sin(e + fx)}} - \frac{a^2 2^{m+\frac{1}{2}} \cos(e + fx)(cd(m + 1)(A + C) + d^2(-Am + Cm + C))}{f \sqrt{a - a \sin(e + fx)} \sqrt{a \sin(e + fx) + a}}}{ad(m + 1)(c^2 - d^2)} \\ \frac{(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}}{df(m + 1)(c^2 - d^2)}$$

↓ 157

---

3.10.  $\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx$

$$\frac{a^2 C(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)\int\frac{\sqrt{2}(\sin(e+fx)a+a)^{m+\frac{1}{2}}(c+d\sin(e+fx))^{-m-1}d\sin(e+fx)}{\sqrt{2f(a-a\sin(e+fx))}\sqrt{a\sin(e+fx)+a}}}{a^2 2^{m+\frac{1}{2}}\cos(e+fx)(cd(m+1)(A+C)+ad(m+1))} - \frac{(Ad^2+c^2C)\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

↓ 27

$$\frac{a^2 C(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)\int\frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}(c+d\sin(e+fx))^{-m-1}d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}}}{a^2 2^{m+\frac{1}{2}}\cos(e+fx)(cd(m+1)(A+C)+ad(m+1))} - \frac{(Ad^2+c^2C)\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

↓ 156

$$\frac{a^2 C(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)(c+d\sin(e+fx))^{-m}\left(\frac{c+d\sin(e+fx)}{c-d}\right)^m\int\frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}\left(\frac{c}{c-d}+\frac{d\sin(e+fx)}{c-d}\right)^{-m-1}d\sin(e+fx)}{f(c-d)(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}}}{(Ad^2+c^2C)\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}} - \frac{(Ad^2+c^2C)\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

↓ 155

$$\frac{\sqrt{2}aC(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)(a\sin(e+fx)+a)^{m+1}(c+d\sin(e+fx))^{-m}\left(\frac{c+d\sin(e+fx)}{c-d}\right)^m\text{AppellF1}\left(m+\frac{3}{2},\frac{1}{2},m+1,m+\frac{5}{2},\frac{1}{2}(\sin(e+fx)+1)\right)}{f(2m+3)(c-d)(a-a\sin(e+fx))} - \frac{(Ad^2+c^2C)\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

$$\frac{(Ad^2+c^2C)\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

input `Int[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(-2 - m)*(A + C*Sin[e + f*x])^2, x]`

```
output ((c^2*C + A*d^2)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^
(-1 - m))/(d*(c^2 - d^2)*f*(1 + m) + (-((2^(1/2 + m)*a^2*(c*(A + C)*d*(1
+ m) + d^2*(C - A*m + C*m) - c^2*(C + 2*C*m))*Cos[e + f*x]*Hypergeometric2
F1[1/2, 1/2 - m, 3/2, ((c - d)*(1 - Sin[e + f*x]))/(2*(c + d*Sin[e + f*x])
])* (a + a*Sin[e + f*x])^(-1 + m)*(((c + d)*(1 + Sin[e + f*x]))/(c + d*Sin[
e + f*x]))^(1/2 - m))/((c + d)*f*(c + d*Sin[e + f*x])^m) + (Sqrt[2]*a*C*(
c^2 - d^2)*(1 + m)*AppellF1[3/2 + m, 1/2, 1 + m, 5/2 + m, (1 + Sin[e + f*x
])/2, -((d*(1 + Sin[e + f*x]))/(c - d))*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x
]]*(a + a*Sin[e + f*x])^(1 + m)*((c + d*Sin[e + f*x])/(c - d))^m)/((c - d)
*f*(3 + 2*m)*(a - a*Sin[e + f*x])*(c + d*Sin[e + f*x])^m)/(a*d*(c^2 - d^2
)*(1 + m))
```

### 3.10.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 142 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^(p_), x_] := Simp[((a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/((b*e
- a*f)*(m + 1)))*Hypergeometric2F1[m + 1, -n, m + 2, (-d*e - c*f))*((a +
b*x)/((b*c - a*d)*(e + f*x)))]/((b*e - a*f)*((c + d*x)/((b*c - a*d)*(e +
f*x))))^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[m + n + p + 2,
0] && !IntegerQ[n]
```

```
rule 155 Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simp
lify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplerQ[e + f*x, a + b*x])
```

rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p])*((b*(e + f*x)/(b*e - a*f))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`

rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n])*((b*(c + d*x)/(b*c - a*d))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]])*Sqrt[a - b*Sin[e + f*x]]) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]`

rule 3466 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]`



```
rule 3523 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
  (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)^2], x_Symbol] :=
  Simp[(-(c^2*C + A*d^2))*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e +
  f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*d*(n + 1)*(c^2 -
  d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(a
  *d*m + b*c*(n + 1)) + c*C*(a*c*m + b*d*(n + 1)) - b*(A*d^2*(m + n + 2) + C*
  (c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d,
  e, f, A, C, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 -
  d^2, 0] && !LtQ[m, -2^(-1)] && (LtQ[n, -1] || EqQ[m + n + 2, 0])
```

### 3.10.4 Maple [F]

$$\int (a + a \sin(fx + e))^m (c + d \sin(fx + e))^{-2-m} (A + C(\sin^2(fx + e))) dx$$

```
input int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+C*sin(f*x+e)^2),x)
```

```
output int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+C*sin(f*x+e)^2),x)
```

### 3.10.5 Fricas [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + A)(a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^{-m-2} dx$$

```
input integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+C*sin(f*x+e)^2),x,
  algorithm="fricas")
```

```
output integral(-(C*cos(f*x + e)^2 - A - C)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e)
  ) + c)^(-m - 2), x)
```

**3.10.6 Sympy [F(-1)]**

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx = \text{Timed out}$$

```
input integrate((a+a*sin(f*x+e))**m*(c+d*sin(f*x+e))**(-2-m)*(A+C*sin(f*x+e)**2),x)
```

```
output Timed out
```

**3.10.7 Maxima [F]**

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx \\ &= \int (C \sin^2(fx + e) + A) (a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^{-m-2} dx \end{aligned}$$

```
input integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))**(-2-m)*(A+C*sin(f*x+e)^2),x,
algorithm="maxima")
```

```
output integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e) +
c)**(-m - 2), x)
```

**3.10.8 Giac [F]**

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx \\ &= \int (C \sin^2(fx + e) + A) (a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^{-m-2} dx \end{aligned}$$

```
input integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))**(-2-m)*(A+C*sin(f*x+e)^2),x,
algorithm="giac")
```

```
output integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e) +
c)**(-m - 2), x)
```

**3.10.9 Mupad [F(-1)]**

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx$$

$$= \int \frac{(C \sin(e + fx)^2 + A) (a + a \sin(e + fx))^m}{(c + d \sin(e + fx))^{m+2}} dx$$

input `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c + d*sin(e + f*x))^(m + 2),x)`

output `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c + d*sin(e + f*x))^(m + 2), x)`

### 3.11 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{3/2} (A + C \sin^2(e$

3.11.1	Optimal result	115
3.11.2	Mathematica [F]	116
3.11.3	Rubi [A] (verified)	116
3.11.4	Maple [F]	120
3.11.5	Fricas [F]	121
3.11.6	Sympy [F(-1)]	121
3.11.7	Maxima [F]	121
3.11.8	Giac [F]	122
3.11.9	Mupad [F(-1)]	122

#### 3.11.1 Optimal result

Integrand size = 39, antiderivative size = 385

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx =$$

$$\frac{2C \cos(e + fx)(a + a \sin(e + fx))^m (c + d \sin(e + fx))^{5/2}}{df(7 + 2m)}$$

$$+ \frac{\sqrt{2}(c - d)(2c(C + 2Cm) + d(C(5 - 2m) + A(7 + 2m))) \operatorname{AppellF1}\left(\frac{1}{2} + m, \frac{1}{2}, -\frac{3}{2}, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e + fx))\right)}{df(1 + 2m)(7 + 2m)\sqrt{1 - \sin(e + fx)}\sqrt{\frac{c + d \sin(e + fx)}{c - d}}}$$

$$+ \frac{2\sqrt{2}C(c - d)(dm - c(1 + m)) \operatorname{AppellF1}\left(\frac{3}{2} + m, \frac{1}{2}, -\frac{3}{2}, \frac{5}{2} + m, \frac{1}{2}(1 + \sin(e + fx))\right) \cos(e + fx)}{adf(3 + 2m)(7 + 2m)\sqrt{1 - \sin(e + fx)}\sqrt{\frac{c + d \sin(e + fx)}{c - d}}}$$

output

```
-2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(5/2)/d/f/(7+2*m)+(c-d)
)*(2*c*(2*C*m+C)+d*(C*(5-2*m)+A*(7+2*m)))*AppellF1(1/2+m,-3/2,1/2,3/2+m,-d
*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^m*2^
(1/2)*(c+d*sin(f*x+e))^(1/2)/d/f/(1+2*m)/(7+2*m)/(1-sin(f*x+e))^(1/2)/((c+
d*sin(f*x+e))/(c-d))^(1/2)+2*C*(c-d)*(d*m-c*(1+m))*AppellF1(3/2+m,-3/2,1/2
,5/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x
+e))^(1+m)*2^(1/2)*(c+d*sin(f*x+e))^(1/2)/a/d/f/(3+2*m)/(7+2*m)/(1-sin(f*x
+e))^(1/2)/((c+d*sin(f*x+e))/(c-d))^(1/2)
```

### 3.11.2 Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx$$

input `Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2)*(A + C*Sin[e + f*x]^2),x]`

output `Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2)*(A + C*Sin[e + f*x]^2), x]`

### 3.11.3 Rubi [A] (verified)

Time = 1.16 (sec) , antiderivative size = 406, normalized size of antiderivative = 1.05, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.282$ , Rules used = {3042, 3525, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) (c + d \sin(e + fx))^{3/2} dx$$

↓ 3042

$$\int (a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2) (c + d \sin(e + fx))^{3/2} dx$$

↓ 3525

$$\frac{2 \int \frac{1}{2} (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{3/2} (a(Ad(2m + 7) + C(5d + 2cm)) + 2aC(dm - c(m + 1)) \sin(e + fx)) dx}{ad(2m + 7)} + \frac{2C \cos(e + fx) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}$$

↓ 27

$$\frac{\int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{3/2} (a(Ad(2m + 7) + C(5d + 2cm)) + 2aC(dm - c(m + 1)) \sin(e + fx)) dx}{ad(2m + 7)} + \frac{2C \cos(e + fx) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}$$

---

3.11.  $\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{3/2} (a(Ad(2m + 7) + C(5d + 2cm)) + 2aC(dm - c(m + 1)) \sin(e + fx))}{ad(2m + 7)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}$$

↓ 3466

$$\frac{a(Ad(2m + 7) + 2c(2Cm + C) + Cd(5 - 2m)) \int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{3/2} dx + 2C(dm - c(m + 1)) \sin(e + fx)}{ad(2m + 7)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}$$

↓ 3042

$$\frac{a(Ad(2m + 7) + 2c(2Cm + C) + Cd(5 - 2m)) \int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{3/2} dx + 2C(dm - c(m + 1)) \sin(e + fx)}{ad(2m + 7)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}$$

↓ 3267

$$\frac{a^3 \cos(e + fx)(Ad(2m + 7) + 2c(2Cm + C) + Cd(5 - 2m)) \int \frac{(\sin(e + fx)a + a)^{m - \frac{1}{2}} (c + d \sin(e + fx))^{3/2} d \sin(e + fx)}{\sqrt{a - a \sin(e + fx)} \sqrt{a \sin(e + fx) + a}} + \frac{2a^2 C(dm - c(m + 1)) \cos(e + fx) \int \frac{dx}{\sqrt{a - a \sin(e + fx)}}}{f \sqrt{a - a \sin(e + fx)}}}{ad(2m + 7)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}$$

↓ 157

$$\frac{a^3 \sqrt{1 - \sin(e + fx)} \cos(e + fx)(Ad(2m + 7) + 2c(2Cm + C) + Cd(5 - 2m)) \int \frac{\sqrt{2}(\sin(e + fx)a + a)^{m - \frac{1}{2}} (c + d \sin(e + fx))^{3/2} d \sin(e + fx)}{\sqrt{2}f(a - a \sin(e + fx)) \sqrt{a \sin(e + fx) + a}} + \frac{\sqrt{2}a^2 C(dm - c(m + 1)) \cos(e + fx) \int \frac{dx}{\sqrt{1 - \sin(e + fx)}}}{f \sqrt{1 - \sin(e + fx)}}}{ad(2m + 7)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}$$

↓ 27

---

3.11.  $\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx$

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (Ad(2m+7)+2c(2Cm+C)+Cd(5-2m)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} (c+d \sin(e+fx))^{3/2}}{\sqrt{1-\sin(e+fx)}} d \sin(e+fx)}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} + \frac{2a^2 C(dm-c(m+1))}{ad(2m+7)}$$

$$\frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{5/2}}{df(2m+7)}$$

↓ 156

$$\frac{a^3(c-d) \sqrt{1-\sin(e+fx)} \cos(e+fx) (Ad(2m+7)+2c(2Cm+C)+Cd(5-2m)) \sqrt{c+d \sin(e+fx)} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} \left(\frac{c}{c-d} + \frac{d \sin(e+fx)}{c-d}\right)^{3/2}}{\sqrt{1-\sin(e+fx)}} d \sin(e+fx)}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a} \sqrt{\frac{c+d \sin(e+fx)}{c-d}}}$$

$$\frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{5/2}}{df(2m+7)}$$

↓ 155

$$\frac{\sqrt{2}a^2(c-d) \sqrt{1-\sin(e+fx)} \cos(e+fx) (Ad(2m+7)+2c(2Cm+C)+Cd(5-2m)) (a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)} \text{AppellF1}\left(m+\frac{1}{2}, \frac{1}{2}, -\frac{3}{2}, m+\frac{3}{2}, \frac{1+\sin(e+fx)}{2}, -\frac{d(1+\sin(e+fx))}{c-d}\right)}{f(2m+1)(a-a \sin(e+fx)) \sqrt{\frac{c+d \sin(e+fx)}{c-d}}}$$

$$\frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{5/2}}{df(2m+7)}$$

input `Int[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2)*(A + C*Sin[e + f*x]^2),x]`

output `(-2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(5/2))/(d*f*(7 + 2*m)) + ((Sqrt[2]*a^2*(c - d)*(C*d*(5 - 2*m) + A*d*(7 + 2*m) + 2*c*(C + 2*C*m))*AppellF1[1/2 + m, 1/2, -3/2, 3/2 + m, (1 + Sin[e + f*x])/2, -(d*(1 + Sin[e + f*x]))/(c - d)])*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]])/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]) + (2*Sqrt[2]*a*C*(c - d)*(d*m - c*(1 + m))*AppellF1[3/2 + m, 1/2, -3/2, 5/2 + m, (1 + Sin[e + f*x])/2, -(d*(1 + Sin[e + f*x]))/(c - d)])*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[c + d*Sin[e + f*x]])/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]))/(a*d*(7 + 2*m))`

## 3.11.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



rule 3267 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]])*Sqrt[a - b*Sin[e + f*x]]) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]`

rule 3466 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]`

rule 3525 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1))/(d*f*(m + n + 2)), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + C*(a*d*m - b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]`

### 3.11.4 Maple [F]

$$\int (a + a \sin(fx + e))^m (c + d \sin(fx + e))^{\frac{3}{2}} (A + C(\sin^2(fx + e))) dx$$

input `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x)`

output `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x)`

### 3.11.5 Fricas [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e) + A)(d \sin(fx + e) + c)^{3/2} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x,  
algorithm="fricas")`

output `integral(-(C*c*cos(f*x + e)^2 - (A + C)*c + (C*d*cos(f*x + e)^2 - (A + C)*  
d)*sin(f*x + e))*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m, x)`

### 3.11.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)**2),  
x)`

output `Timed out`

### 3.11.7 Maxima [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e) + A)(d \sin(fx + e) + c)^{3/2} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x,  
algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(d*sin(f*x + e) + c)^(3/2)*(a*sin(f*x + e)  
) + a)^m, x)`

---

3.11.  $\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx$

**3.11.8 Giac [F]**

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e) + A) (d \sin(fx + e) + c)^{3/2} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x,  
algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + A)*(d*sin(f*x + e) + c)^(3/2)*(a*sin(f*x + e)  
) + a)^m, x)`

**3.11.9 Mupad [F(-1)]**

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \int (C \sin(e + fx)^2 + A) (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} dx$$

input `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(3/  
2),x)`

output `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(3/  
2), x)`

### 3.12 $\int (a+a \sin(e+fx))^m \sqrt{c+d \sin(e+fx)}(A+C \sin^2(e$

3.12.1	Optimal result	123
3.12.2	Mathematica [F]	124
3.12.3	Rubi [A] (verified)	124
3.12.4	Maple [F]	128
3.12.5	Fricas [F]	129
3.12.6	Sympy [F]	129
3.12.7	Maxima [F]	129
3.12.8	Giac [F]	130
3.12.9	Mupad [F(-1)]	130

#### 3.12.1 Optimal result

Integrand size = 39, antiderivative size = 375

$$\int (a+a \sin(e+fx))^m \sqrt{c+d \sin(e+fx)}(A+C \sin^2(e+fx)) dx$$

$$= -\frac{2C \cos(e+fx)(a+a \sin(e+fx))^m (c+d \sin(e+fx))^{3/2}}{df(5+2m)}$$

$$+ \frac{\sqrt{2}(2c(C+2Cm)+d(C(3-2m)+A(5+2m))) \operatorname{AppellF1}\left(\frac{1}{2}+m, \frac{1}{2}, -\frac{1}{2}, \frac{3}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right)}{df(1+2m)(5+2m)\sqrt{1-\sin(e+fx)}\sqrt{c+d \sin(e+fx)}}$$

$$+ \frac{2\sqrt{2}C(dm-c(1+m)) \operatorname{AppellF1}\left(\frac{3}{2}+m, \frac{1}{2}, -\frac{1}{2}, \frac{5}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right) \cos(e+fx)}{adf(3+2m)(5+2m)\sqrt{1-\sin(e+fx)}\sqrt{\frac{c+d \sin(e+fx)}{c-d}}}$$

output

```
-2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)/d/f/(5+2*m)+(2*c
*(2*C*m+C)+d*(C*(3-2*m)+A*(5+2*m)))*AppellF1(1/2+m,-1/2,1/2,3/2+m,-d*(1+si
n(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^m*2^(1/2)*
(c+d*sin(f*x+e))^(1/2)/d/f/(1+2*m)/(5+2*m)/(1-sin(f*x+e))^(1/2)/((c+d*sin(
f*x+e))/(c-d))^(1/2)+2*C*(d*m-c*(1+m))*AppellF1(3/2+m,-1/2,1/2,5/2+m,-d*(1
+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)*2
^(1/2)*(c+d*sin(f*x+e))^(1/2)/a/d/f/(3+2*m)/(5+2*m)/(1-sin(f*x+e))^(1/2)/(
(c+d*sin(f*x+e))/(c-d))^(1/2)
```

### 3.12.2 Mathematica [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

input `Integrate[(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]]*(A + C*Sin[e + f*x]^2), x]`

output `Integrate[(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]]*(A + C*Sin[e + f*x]^2), x]`

### 3.12.3 Rubi [A] (verified)

Time = 1.13 (sec) , antiderivative size = 396, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.282$ , Rules used = {3042, 3525, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) \sqrt{c + d \sin(e + fx)} dx$$

↓ 3042

$$\int (a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2) \sqrt{c + d \sin(e + fx)} dx$$

↓ 3525

$$2 \int \frac{\frac{1}{2}(\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} (a(Ad(2m + 5) + C(3d + 2cm)) + 2aC(dm - c(m + 1)) \sin(e + fx))}{ad(2m + 5)}$$

$$\frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}}{df(2m + 5)}$$

↓ 27

---

3.12.  $\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$

$$\frac{\int (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} (a(Ad(2m + 5) + C(3d + 2cm)) + 2aC(dm - c(m + 1)) \sin(e + fx)) dx}{\frac{ad(2m + 5)}{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}} \frac{df(2m + 5)}{df(2m + 5)}} \downarrow 3042$$

$$\frac{\int (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} (a(Ad(2m + 5) + C(3d + 2cm)) + 2aC(dm - c(m + 1)) \sin(e + fx)) dx}{\frac{ad(2m + 5)}{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}} \frac{df(2m + 5)}{df(2m + 5)}} \downarrow 3466$$

$$\frac{a(Ad(2m + 5) + 2c(2Cm + C) + Cd(3 - 2m)) \int (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} dx + 2C(dm - c(m + 1))}{\frac{ad(2m + 5)}{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}} \frac{df(2m + 5)}{df(2m + 5)}} \downarrow 3042$$

$$\frac{a(Ad(2m + 5) + 2c(2Cm + C) + Cd(3 - 2m)) \int (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} dx + 2C(dm - c(m + 1))}{\frac{ad(2m + 5)}{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}} \frac{df(2m + 5)}{df(2m + 5)}} \downarrow 3267$$

$$\frac{a^3 \cos(e + fx)(Ad(2m + 5) + 2c(2Cm + C) + Cd(3 - 2m)) \int \frac{(\sin(e + fx)a + a)^{m - \frac{1}{2}} \sqrt{c + d \sin(e + fx)} d \sin(e + fx)}{f \sqrt{a - a \sin(e + fx)} \sqrt{a \sin(e + fx) + a}} + \frac{2a^2 C(dm - c(m + 1)) \cos(e + fx) \int (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} dx}{f \sqrt{a - a \sin(e + fx)}}}{\frac{ad(2m + 5)}{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}} \frac{df(2m + 5)}{df(2m + 5)}} \downarrow 157$$

$$\frac{a^3 \sqrt{1 - \sin(e + fx)} \cos(e + fx)(Ad(2m + 5) + 2c(2Cm + C) + Cd(3 - 2m)) \int \frac{\sqrt{2}(\sin(e + fx)a + a)^{m - \frac{1}{2}} \sqrt{c + d \sin(e + fx)} d \sin(e + fx)}{\sqrt{2}f(a - a \sin(e + fx)) \sqrt{a \sin(e + fx) + a}} + \frac{\sqrt{2}a^2 C(dm - c(m + 1)) \cos(e + fx) \int (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} dx}{f \sqrt{a - a \sin(e + fx)}}}{\frac{ad(2m + 5)}{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}} \frac{df(2m + 5)}{df(2m + 5)}} \downarrow 27$$

---

3.12.  $\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (Ad(2m+5)+2c(2Cm+C)+Cd(3-2m)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} \sqrt{c+d \sin(e+fx)}}{\sqrt{1-\sin(e+fx)}} d \sin(e+fx)}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} + \frac{2a^2 C(dm-c(m+1)) \sqrt{c+d \sin(e+fx)}}{ad(2m+5)}$$

$$\frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{3/2}}{df(2m+5)}$$

↓ 156

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (Ad(2m+5)+2c(2Cm+C)+Cd(3-2m)) \sqrt{c+d \sin(e+fx)} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} \sqrt{\frac{c}{c-d} + \frac{d \sin(e+fx)}{c-d}}}{\sqrt{1-\sin(e+fx)}} d \sin(e+fx)}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a} \sqrt{\frac{c+d \sin(e+fx)}{c-d}}} + \frac{2a^2 C(dm-c(m+1)) \sqrt{c+d \sin(e+fx)}}{ad(2m+5)}$$

$$\frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{3/2}}{df(2m+5)}$$

↓ 155

$$\frac{\sqrt{2} a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (Ad(2m+5)+2c(2Cm+C)+Cd(3-2m)) (a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)} \text{AppellF1}\left(m+\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, m+\frac{3}{2}, \frac{1}{2}(\sin(e+fx)+1)\right)}{f(2m+1)(a-a \sin(e+fx)) \sqrt{\frac{c+d \sin(e+fx)}{c-d}}}$$

$$\frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{3/2}}{df(2m+5)}$$

input `Int[(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]]*(A + C*Sin[e + f*x]^2),x]`

output `(-2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2))/(d*f*(5 + 2*m)) + ((Sqrt[2]*a^2*(C*d*(3 - 2*m) + A*d*(5 + 2*m) + 2*c*(C + 2*C*m))*AppellF1[1/2 + m, 1/2, -1/2, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]])/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]) + (2*Sqrt[2]*a*C*(d*m - c*(1 + m))*AppellF1[3/2 + m, 1/2, -1/2, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[c + d*Sin[e + f*x]])/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]))/(a*d*(5 + 2*m))`

## 3.12.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



rule 3267 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]])*Sqrt[a - b*Sin[e + f*x]]) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]`

rule 3466 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]`

rule 3525 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1))/(d*f*(m + n + 2)), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + C*(a*d*m - b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]`

### 3.12.4 Maple [F]

$$\int (a + a \sin(fx + e))^m \sqrt{c + d \sin(fx + e)} (A + C(\sin^2(fx + e))) dx$$

input `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x)`

output `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x)`

**3.12.5 Fracas [F]**

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(fx + e) + A) \sqrt{d \sin(fx + e) + c} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x,  
algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - A - C)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x +  
e) + a)^m, x)`

**3.12.6 Sympy [F]**

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \int (a(\sin(e + fx) + 1))^m (A + C \sin^2(e + fx)) \sqrt{c + d \sin(e + fx)} dx$$

input `integrate((a+a*sin(f*x+e))**m*(c+d*sin(f*x+e))**(1/2)*(A+C*sin(f*x+e)**2),  
x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + C*sin(e + f*x)**2)*sqrt(c + d*sin(  
e + f*x)), x)`

**3.12.7 Maxima [F]**

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(fx + e) + A) \sqrt{d \sin(fx + e) + c} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x,  
algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x + e)  
+ a)^m, x)`

### 3.12.8 Giac [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx \\ &= \int (C \sin^2(fx + e) + A) \sqrt{d \sin(fx + e) + c} (a \sin(fx + e) + a)^m dx \end{aligned}$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x,  
algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + A)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x + e)  
+ a)^m, x)`

### 3.12.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx \\ &= \int (C \sin^2(e + fx) + A) (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} dx \end{aligned}$$

input `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(1/  
2),x)`

output `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(1/  
2), x)`

$$3.13 \quad \int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx$$

3.13.1	Optimal result	131
3.13.2	Mathematica [F]	132
3.13.3	Rubi [A] (verified)	132
3.13.4	Maple [F]	136
3.13.5	Fricas [F]	137
3.13.6	Sympy [F]	137
3.13.7	Maxima [F]	137
3.13.8	Giac [F]	138
3.13.9	Mupad [F(-1)]	138

### 3.13.1 Optimal result

Integrand size = 39, antiderivative size = 365

$$\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx$$

$$= -\frac{2C \cos(e+fx)(a+a \sin(e+fx))^m \sqrt{c+d \sin(e+fx)}}{df(3+2m)}$$

$$+ \frac{\sqrt{2}(2c(C+2Cm)+d(C-2Cm+A(3+2m))) \operatorname{AppellF1}\left(\frac{1}{2}+m, \frac{1}{2}, \frac{1}{2}, \frac{3}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right), -\frac{d}{c-d}}{df(1+2m)(3+2m)\sqrt{1-\sin(e+fx)}\sqrt{c+d \sin(e+fx)}}$$

$$- \frac{2\sqrt{2}C(c+cm-dm) \operatorname{AppellF1}\left(\frac{3}{2}+m, \frac{1}{2}, \frac{1}{2}, \frac{5}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right), -\frac{d(1+\sin(e+fx))}{c-d}}{adf(3+2m)^2\sqrt{1-\sin(e+fx)}\sqrt{c+d \sin(e+fx)}}$$

output

```
-2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)/d/f/(3+2*m)+(2*c
*(2*C*m+C)+d*(C-2*C*m+A*(3+2*m)))*AppellF1(1/2+m,1/2,1/2,3/2+m,-d*(1+sin(f
*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^m*2^(1/2)*((c
+d*sin(f*x+e))/(c-d))^(1/2)/d/f/(1+2*m)/(3+2*m)/(1-sin(f*x+e))^(1/2)/(c+d*
sin(f*x+e))^(1/2)-2*C*(c*m-d*m+c)*AppellF1(3/2+m,1/2,1/2,5/2+m,-d*(1+sin(f
*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)*2^(1/2)
*((c+d*sin(f*x+e))/(c-d))^(1/2)/a/d/f/(3+2*m)^2/(1-sin(f*x+e))^(1/2)/(c+d*
sin(f*x+e))^(1/2)
```

### 3.13.2 Mathematica [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

input `Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/Sqrt[c + d*Sin[e + f*x]],x]`

output `Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/Sqrt[c + d*Sin[e + f*x]], x]`

### 3.13.3 Rubi [A] (verified)

Time = 1.16 (sec) , antiderivative size = 395, normalized size of antiderivative = 1.08, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.282$ , Rules used = {3042, 3525, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \sin(e + fx) + a)^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2)}{\sqrt{c + d \sin(e + fx)}} dx$$

$$\downarrow \text{3525}$$

$$2 \int \frac{(\sin(e+fx)a+a)^m (a(Ad(2m+3)+C(d+2cm))+2aC(dm-c(m+1))\sin(e+fx))}{2\sqrt{c+d\sin(e+fx)}} dx$$

$$\frac{ad(2m+3)}{2C \cos(e+fx)(a \sin(e+fx) + a)^m \sqrt{c+d \sin(e+fx)}} -$$

$$\frac{df(2m+3)}{27}$$

---

3.13.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx$

$$\frac{\int \frac{(\sin(e+fx)a+a)^m (a(Ad(2m+3)+C(d+2cm))+2aC(dm-c(m+1)) \sin(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx}{ad(2m+3) \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}} -$$

↓ 3042

$$\frac{\int \frac{(\sin(e+fx)a+a)^m (a(Ad(2m+3)+C(d+2cm))+2aC(dm-c(m+1)) \sin(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx}{ad(2m+3) \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}} -$$

↓ 3466

$$\frac{a(d(A(2m+3)-2Cm+C)+2c(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx + 2C(dm-c(m+1)) \int \frac{(\sin(e+fx)a+a)^{m+1}}{\sqrt{c+d \sin(e+fx)}} dx}{ad(2m+3) \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}} -$$

↓ 3042

$$\frac{a(d(A(2m+3)-2Cm+C)+2c(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx + 2C(dm-c(m+1)) \int \frac{(\sin(e+fx)a+a)^{m+1}}{\sqrt{c+d \sin(e+fx)}} dx}{ad(2m+3) \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}} -$$

↓ 3267

$$\frac{a^3 \cos(e+fx)(d(A(2m+3)-2Cm+C)+2c(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{a-a \sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx) + \frac{2a^2 C(dm-c(m+1)) \cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{a-a \sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx)}{f \sqrt{a-a \sin(e+fx)} \sqrt{a \sin(e+fx)+a}}}{ad(2m+3) \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}} -$$

↓ 157

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)(d(A(2m+3)-2Cm+C)+2c(2Cm+C)) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx) + \frac{\sqrt{2}a^2 C(dm-c(m+1)) \sqrt{1-\sin(e+fx)} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx)}{\sqrt{2}f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}}}{ad(2m+3) \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}} -$$

↓ 27

---

3.13.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx$

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (d(A(2m+3)-2Cm+C)+2c(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}\sqrt{c+d\sin(e+fx)}} d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}} + \frac{2a^2C(dm-c(m+1))\sqrt{1-\sin(e+fx)}}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}}$$


---


$$\frac{2C \cos(e+fx)(a\sin(e+fx)+a)^m \sqrt{c+d\sin(e+fx)}}{df(2m+3)} \quad ad(2m+3)$$

↓ 156

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (d(A(2m+3)-2Cm+C)+2c(2Cm+C)) \sqrt{\frac{c+d\sin(e+fx)}{c-d}} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}\sqrt{\frac{c}{c-d}+\frac{d\sin(e+fx)}{c-d}}} d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}\sqrt{c+d\sin(e+fx)}} + \frac{2a^2C(dm-c(m+1))\sqrt{1-\sin(e+fx)}}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}}$$


---


$$\frac{2C \cos(e+fx)(a\sin(e+fx)+a)^m \sqrt{c+d\sin(e+fx)}}{df(2m+3)} \quad ad(2m+3)$$

↓ 155

$$\frac{\sqrt{2}a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (d(A(2m+3)-2Cm+C)+2c(2Cm+C))(a\sin(e+fx)+a)^m \sqrt{\frac{c+d\sin(e+fx)}{c-d}} \text{AppellF1}\left(m+\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, m+\frac{3}{2}, \frac{1}{2}(\sin(e+fx)+1)\right)}{f(2m+1)(a-a\sin(e+fx))\sqrt{c+d\sin(e+fx)}}$$


---


$$\frac{2C \cos(e+fx)(a\sin(e+fx)+a)^m \sqrt{c+d\sin(e+fx)}}{df(2m+3)}$$

input `Int[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/Sqrt[c + d*Sin[e + f*x]],x]`

output `(-2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]])/(d*f*(3 + 2*m)) + ((Sqrt[2]*a^2*(2*c*(C + 2*C*m) + d*(C - 2*C*m + A*(3 + 2*m)))*AppellF1[1/2 + m, 1/2, 1/2, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[(c + d*Sin[e + f*x])/(c - d)]/(f*(1 + 2*m)*(a - a*Sin[e + f*x]))*Sqrt[c + d*Sin[e + f*x]]) + (2*Sqrt[2]*a*C*(d*m - c*(1 + m))*AppellF1[3/2 + m, 1/2, 1/2, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[(c + d*Sin[e + f*x])/(c - d)]/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]))/(a*d*(3 + 2*m))`

---

3.13.  $\int \frac{(a+a\sin(e+fx))^m (A+C\sin^2(e+fx))}{\sqrt{c+d\sin(e+fx)}} dx$

## 3.13.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p)*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



rule 3267 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]])*Sqrt[a - b*Sin[e + f*x]]) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]`

rule 3466 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]`

rule 3525 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1))/(d*f*(m + n + 2)), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + C*(a*d*m - b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]`

### 3.13.4 Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + C \sin^2(fx + e))}{\sqrt{c + d \sin(fx + e)}} dx$$

input `int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x)`

output `int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x)`

### 3.13.5 Fricas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{\sqrt{d \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x,  
algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - A - C)*(a*sin(f*x + e) + a)^m/sqrt(d*sin(f*x  
+ e) + c), x)`

### 3.13.6 Sympy [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(a(\sin(e + fx) + 1))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

input `integrate((a+a*sin(f*x+e))**m*(A+C*sin(f*x+e)**2)/(c+d*sin(f*x+e))**(1/2),  
x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + C*sin(e + f*x)**2)/sqrt(c + d*sin(  
e + f*x)), x)`

### 3.13.7 Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{\sqrt{d \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x,  
algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/sqrt(d*sin(f*x + e)  
) + c), x)`

### 3.13.8 Giac [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{\sqrt{d \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x,  
algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/sqrt(d*sin(f*x + e)  
) + c), x)`

### 3.13.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin^2(e + fx) + A)(a + a \sin(e + fx))^m}{\sqrt{c + d \sin(e + fx)}} dx$$

input `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c + d*sin(e + f*x))^(  
1/2),x)`

output `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c + d*sin(e + f*x))^(  
1/2), x)`

$$3.14 \quad \int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx$$

3.14.1	Optimal result	139
3.14.2	Mathematica [F]	140
3.14.3	Rubi [A] (verified)	140
3.14.4	Maple [F]	144
3.14.5	Fricas [F]	145
3.14.6	Sympy [F]	145
3.14.7	Maxima [F]	145
3.14.8	Giac [F]	146
3.14.9	Mupad [F(-1)]	146

### 3.14.1 Optimal result

Integrand size = 39, antiderivative size = 413

$$\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx = \frac{2(c^2C + Ad^2) \cos(e+fx)(a+a \sin(e+fx))^m}{d(c^2 - d^2) f \sqrt{c+d \sin(e+fx)}} + \frac{\sqrt{2}(c(A+C)d - d^2(A-C+4Am) - 2c^2(C+2Cm)) \operatorname{AppellF1}\left(\frac{1}{2} + m, \frac{1}{2}, \frac{1}{2}, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e+fx))\right)}{d(c^2 - d^2) f(1+2m)\sqrt{1 - \sin(e+fx)}\sqrt{c+d \sin(e+fx)}} + \frac{\sqrt{2}(2c^2C(1+m) + d^2(A-C+2Am)) \operatorname{AppellF1}\left(\frac{3}{2} + m, \frac{1}{2}, \frac{1}{2}, \frac{5}{2} + m, \frac{1}{2}(1 + \sin(e+fx))\right), -\frac{d(1+\sin(e+fx))}{c-d}}{ad(c^2 - d^2) f(3+2m)\sqrt{1 - \sin(e+fx)}\sqrt{c+d \sin(e+fx)}}$$

output

```
2*(A*d^2+C*c^2)*cos(f*x+e)*(a+a*sin(f*x+e))^m/d/(c^2-d^2)/f/(c+d*sin(f*x+e))^
(1/2)+(c*(A+C)*d-d^2*(4*A*m+A-C)-2*c^2*(2*C*m+C))*AppellF1(1/2+m,1/2,1/
2,3/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*
x+e))^m*2^(1/2)*((c+d*sin(f*x+e))/(c-d))^(1/2)/d/(c^2-d^2)/f/(1+2*m)/(1-si
n(f*x+e))^(1/2)/(c+d*sin(f*x+e))^(1/2)+(2*c^2*C*(1+m)+d^2*(2*A*m+A-C))*App
ellF1(3/2+m,1/2,1/2,5/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(
f*x+e)*(a+a*sin(f*x+e))^(1+m)*2^(1/2)*((c+d*sin(f*x+e))/(c-d))^(1/2)/a/d/(
c^2-d^2)/f/(3+2*m)/(1-sin(f*x+e))^(1/2)/(c+d*sin(f*x+e))^(1/2)
```

### 3.14.2 Mathematica [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx$$

input `Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(3/2), x]`

output `Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(3/2), x]`

### 3.14.3 Rubi [A] (verified)

Time = 1.27 (sec) , antiderivative size = 431, normalized size of antiderivative = 1.04, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.282$ , Rules used = {3042, 3523, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a \sin(e + fx) + a)^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2)}{(c + d \sin(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{3523} \\ & \frac{2(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m}{df (c^2 - d^2) \sqrt{c + d \sin(e + fx)}} - \\ & 2 \int \frac{(\sin(e + fx)a + a)^m (a(cC(d - 2cm) + Ad(c - 2dm)) + a(2C(m + 1)c^2 + d^2(2mA + A - C)) \sin(e + fx))}{2\sqrt{c + d \sin(e + fx)}} dx \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{(\sin(e + fx)a + a)^m (a(cC(d - 2cm) + Ad(c - 2dm)) + a(2C(m + 1)c^2 + d^2(2mA + A - C)) \sin(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx}{ad (c^2 - d^2)} + \\ & \frac{2(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m}{df (c^2 - d^2) \sqrt{c + d \sin(e + fx)}} \end{aligned}$$

---

3.14.  $\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx$

$$\begin{aligned}
& \int \frac{(\sin(e+fx)a+a)^m (a(cC(d-2cm)+Ad(c-2dm))+a(2C(m+1)c^2+d^2(2mA+A-C)) \sin(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx \\
& \quad \frac{ad(c^2-d^2)}{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m} + \\
& \quad \frac{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}{ad(c^2-d^2)} \\
& \quad \downarrow \text{3042} \\
& \frac{a(cd(A+C)-d^2(4Am+A-C)-2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx + (d^2(2Am+A-C)+2c^2C(m+1)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx}{ad(c^2-d^2)} \\
& \quad \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{a(cd(A+C)-d^2(4Am+A-C)-2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx + (d^2(2Am+A-C)+2c^2C(m+1)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx}{ad(c^2-d^2)} \\
& \quad \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}} \\
& \quad \downarrow \text{3267} \\
& \frac{a^3 \cos(e+fx)(cd(A+C)-d^2(4Am+A-C)-2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{a-a \sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx) + a^2 \cos(e+fx)(d^2(2Am+A-C)+2c^2C(m+1)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{a-a \sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx)}{f \sqrt{a-a \sin(e+fx)} \sqrt{a \sin(e+fx)+a}} + \frac{ad(c^2-d^2)}{f \sqrt{a-a \sin(e+fx)}} \\
& \quad \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}} \\
& \quad \downarrow \text{157} \\
& \frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)(cd(A+C)-d^2(4Am+A-C)-2c^2(2Cm+C)) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx) + a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx)(d^2(2Am+A-C)+2c^2C(m+1)) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx)}{\sqrt{2} f (a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} + \frac{ad(c^2-d^2)}{\sqrt{2} f (a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} \\
& \quad \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}} \\
& \quad \downarrow \text{27}
\end{aligned}$$

---

3.14.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx$

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (cd(A+C) - d^2(4Am+A-C) - 2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx)}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} + \frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (cd(A+C) - d^2(4Am+A-C) - 2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx)}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a} \sqrt{c+d \sin(e+fx)}} + \frac{ad(c^2-d^2)}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

$$\frac{2(Ad^2 + c^2C) \cos(e+fx)(a \sin(e+fx) + a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

↓ 156

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (cd(A+C) - d^2(4Am+A-C) - 2c^2(2Cm+C)) \sqrt{\frac{c+d \sin(e+fx)}{c-d}} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{\frac{c}{c-d} + \frac{d \sin(e+fx)}{c-d}}} d \sin(e+fx)}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a} \sqrt{c+d \sin(e+fx)}} + \frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (cd(A+C) - d^2(4Am+A-C) - 2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{\frac{c}{c-d} + \frac{d \sin(e+fx)}{c-d}}} d \sin(e+fx)}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a} \sqrt{c+d \sin(e+fx)}} + \frac{ad(c^2-d^2)}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

$$\frac{2(Ad^2 + c^2C) \cos(e+fx)(a \sin(e+fx) + a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

↓ 155

$$\frac{\sqrt{2}a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (cd(A+C) - d^2(4Am+A-C) - 2c^2(2Cm+C)) (a \sin(e+fx)+a)^m \sqrt{\frac{c+d \sin(e+fx)}{c-d}} \text{AppellF1}\left(m+\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, m+\frac{3}{2}, \frac{1}{2}(\sin(e+fx)+1)\right)}{f(2m+1)(a-a \sin(e+fx)) \sqrt{c+d \sin(e+fx)}}$$

$$\frac{2(Ad^2 + c^2C) \cos(e+fx)(a \sin(e+fx) + a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

input `Int[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(3/2),x]`

output `(2*(c^2*C + A*d^2)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(d*(c^2 - d^2)*f*Sqrt[c + d*Sin[e + f*x]]) + ((Sqrt[2]*a^2*(c*(A + C)*d - d^2*(A - C + 4*A*m) - 2*c^2*(C + 2*C*m))*AppellF1[1/2 + m, 1/2, 1/2, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[(c + d*Sin[e + f*x])/(c - d)]/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]) + (Sqrt[2]*a*(2*c^2*C*(1 + m) + d^2*(A - C + 2*A*m))*AppellF1[3/2 + m, 1/2, 1/2, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[(c + d*Sin[e + f*x])/(c - d)]/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]))/(a*d*(c^2 - d^2))`

---

3.14.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx$

## 3.14.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



rule 3267 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]])*Sqrt[a - b*Sin[e + f*x]]) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]`

rule 3466 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]`

rule 3523 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-(c^2*C + A*d^2))*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(a*d*m + b*c*(n + 1)) + c*C*(a*c*m + b*d*(n + 1)) - b*(A*d^2*(m + n + 2) + C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && (LtQ[n, -1] || EqQ[m + n + 2, 0])`

### 3.14.4 Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + C \sin^2(fx + e))}{(c + d \sin(fx + e))^{\frac{3}{2}}} dx$$

input `int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x)`

output `int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x)`

### 3.14.5 Fricas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{(d \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x,  
algorithm="fricas")`

output `integral((C*cos(f*x + e)^2 - A - C)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x +  
e) + a)^m/(d^2*cos(f*x + e)^2 - 2*c*d*sin(f*x + e) - c^2 - d^2), x)`

### 3.14.6 Sympy [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(a(\sin(e + fx) + 1))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))**m*(A+C*sin(f*x+e)**2)/(c+d*sin(f*x+e))**(3/2),  
x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + C*sin(e + f*x)**2)/(c + d*sin(e +  
f*x))**(3/2), x)`

### 3.14.7 Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{(d \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x,  
algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/(d*sin(f*x + e) +  
c)^(3/2), x)`

---

3.14.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx$

**3.14.8 Giac [F]**

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{(d \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x,  
algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/(d*sin(f*x + e) +  
c)^(3/2), x)`

**3.14.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin^2(e + fx) + A) (a + a \sin(e + fx))^m}{(c + d \sin(e + fx))^{3/2}} dx$$

input `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c + d*sin(e + f*x))^(  
3/2),x)`

output `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c + d*sin(e + f*x))^(  
3/2), x)`

$$3.15 \quad \int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{5/2}} dx$$

3.15.1	Optimal result	147
3.15.2	Mathematica [F]	148
3.15.3	Rubi [A] (verified)	148
3.15.4	Maple [F]	152
3.15.5	Fricas [F]	153
3.15.6	Sympy [F(-1)]	153
3.15.7	Maxima [F]	153
3.15.8	Giac [F]	154
3.15.9	Mupad [F(-1)]	154

### 3.15.1 Optimal result

Integrand size = 39, antiderivative size = 424

$$\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{5/2}} dx = \frac{2(c^2C + Ad^2) \cos(e+fx)(a+a \sin(e+fx))^m}{3d(c^2 - d^2) f(c+d \sin(e+fx))^{3/2}}$$

$$+ \frac{\sqrt{2}(3c(A+C)d + d^2(A+3C-4Am) - 2c^2(C+2Cm)) \operatorname{AppellF1}\left(\frac{1}{2} + m, \frac{1}{2}, \frac{3}{2}, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e+fx))\right)}{3(c-d)^2 d(c+d) f(1+2m) \sqrt{1 - \sin(e+fx)} \sqrt{c+a \sin(e+fx)}}$$

$$+ \frac{\sqrt{2}(2c^2C(1+m) - d^2(A+3C-2Am)) \operatorname{AppellF1}\left(\frac{3}{2} + m, \frac{1}{2}, \frac{3}{2}, \frac{5}{2} + m, \frac{1}{2}(1 + \sin(e+fx))\right), -\frac{d(1+\sin(e+fx))}{c-d}}{3a(c-d)^2 d(c+d) f(3+2m) \sqrt{1 - \sin(e+fx)} \sqrt{c+d \sin(e+fx)}}$$

output

```
2/3*(A*d^2+C*c^2)*cos(f*x+e)*(a+a*sin(f*x+e))^m/d/(c^2-d^2)/f/(c+d*sin(f*x
+e))^(3/2)+1/3*(3*c*(A+C)*d+d^2*(-4*A*m+A+3*C)-2*c^2*(2*C*m+C))*AppellF1(1
/2+m,3/2,1/2,3/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*
(a+a*sin(f*x+e))^m*2^(1/2)*((c+d*sin(f*x+e))/(c-d))^(1/2)/(c-d)^2/d/(c+d)/
f/(1+2*m)/(1-sin(f*x+e))^(1/2)/(c+d*sin(f*x+e))^(1/2)+1/3*(2*c^2*C*(1+m)-d
^2*(-2*A*m+A+3*C))*AppellF1(3/2+m,3/2,1/2,5/2+m,-d*(1+sin(f*x+e))/(c-d),1/
2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)*2^(1/2)*((c+d*sin(f*x+
e))/(c-d))^(1/2)/a/(c-d)^2/d/(c+d)/f/(3+2*m)/(1-sin(f*x+e))^(1/2)/(c+d*sin
(f*x+e))^(1/2)
```

### 3.15.2 Mathematica [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx = \int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx$$

input `Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(5/2), x]`

output `Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(5/2), x]`

### 3.15.3 Rubi [A] (verified)

Time = 1.29 (sec) , antiderivative size = 451, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.282$ , Rules used = {3042, 3523, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a \sin(e + fx) + a)^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2)}{(c + d \sin(e + fx))^{5/2}} dx \\ & \quad \downarrow \text{3523} \\ & \frac{2(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m}{3df(c^2 - d^2)(c + d \sin(e + fx))^{3/2}} - \\ & \frac{2 \int -\frac{(\sin(e + fx)a + a)^m (a(cC(3d - 2cm) + Ad(3c - 2dm)) + a(2c^2C(m + 1) - d^2(-2mA + A + 3C)) \sin(e + fx))}{2(c + d \sin(e + fx))^{3/2}} dx}{3ad(c^2 - d^2)} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{(\sin(e + fx)a + a)^m (a(cC(3d - 2cm) + Ad(3c - 2dm)) + a(2c^2C(m + 1) - d^2(-2mA + A + 3C)) \sin(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx}{3ad(c^2 - d^2)} + \\ & \frac{2(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m}{3df(c^2 - d^2)(c + d \sin(e + fx))^{3/2}} \end{aligned}$$

---

3.15.  $\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx$

$$\int \frac{(\sin(e+fx)a+a)^m (a(cC(3d-2cm)+Ad(3c-2dm))+a(2c^2C(m+1)-d^2(-2mA+A+3C)) \sin(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx + \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{3ad(c^2-d^2) \cdot 3df(c^2-d^2)(c+d \sin(e+fx))^{3/2}}$$

↓ 3042

$$\frac{a(3cd(A+C)+d^2(-4Am+A+3C)-2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{(c+d \sin(e+fx))^{3/2}} dx + (2c^2C(m+1)-d^2(-2Am+A+3C))}{3ad(c^2-d^2)} + \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{3df(c^2-d^2)(c+d \sin(e+fx))^{3/2}}$$

↓ 3466

$$\frac{a(3cd(A+C)+d^2(-4Am+A+3C)-2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{(c+d \sin(e+fx))^{3/2}} dx + (2c^2C(m+1)-d^2(-2Am+A+3C))}{3ad(c^2-d^2)} + \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{3df(c^2-d^2)(c+d \sin(e+fx))^{3/2}}$$

↓ 3042

$$\frac{a(3cd(A+C)+d^2(-4Am+A+3C)-2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{(c+d \sin(e+fx))^{3/2}} dx + (2c^2C(m+1)-d^2(-2Am+A+3C))}{3ad(c^2-d^2)} + \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{3df(c^2-d^2)(c+d \sin(e+fx))^{3/2}}$$

↓ 3267

$$\frac{a^3 \cos(e+fx)(3cd(A+C)+d^2(-4Am+A+3C)-2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{a-a \sin(e+fx)}(c+d \sin(e+fx))^{3/2}} d \sin(e+fx) + a^2 \cos(e+fx)(2c^2C(m+1)-d^2(-2Am+A+3C))}{f \sqrt{a-a \sin(e+fx)} \sqrt{a \sin(e+fx)+a}} + \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{3ad(c^2-d^2) \cdot 3df(c^2-d^2)(c+d \sin(e+fx))^{3/2}}$$

↓ 157

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)(3cd(A+C)+d^2(-4Am+A+3C)-2c^2(2Cm+C)) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}(c+d \sin(e+fx))^{3/2}} d \sin(e+fx) + a^2 \sqrt{1-\sin(e+fx)}(2c^2C(m+1)-d^2(-2Am+A+3C))}{\sqrt{2}f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} + \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{3ad(c^2-d^2) \cdot 3df(c^2-d^2)(c+d \sin(e+fx))^{3/2}}$$

↓ 27

---

3.15.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{5/2}} dx$

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (3cd(A+C)+d^2(-4Am+A+3C)-2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}(c+d\sin(e+fx))^{3/2}} d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}} + \frac{a^2 \sqrt{1-\sin(e+fx)}}{3ad(c^2-d^2)}$$

$$\frac{2(Ad^2+c^2C)\cos(e+fx)(a\sin(e+fx)+a)^m}{3df(c^2-d^2)(c+d\sin(e+fx))^{3/2}}$$

↓ 156

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (3cd(A+C)+d^2(-4Am+A+3C)-2c^2(2Cm+C)) \sqrt{\frac{c+d\sin(e+fx)}{c-d}} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}\left(\frac{c}{c-d}+\frac{d\sin(e+fx)}{c-d}\right)^{3/2}} d\sin(e+fx)}{f(c-d)(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}\sqrt{c+d\sin(e+fx)}}$$

$$\frac{2(Ad^2+c^2C)\cos(e+fx)(a\sin(e+fx)+a)^m}{3df(c^2-d^2)(c+d\sin(e+fx))^{3/2}}$$

↓ 155

$$\frac{\sqrt{2}a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (3cd(A+C)+d^2(-4Am+A+3C)-2c^2(2Cm+C)) (a\sin(e+fx)+a)^m \sqrt{\frac{c+d\sin(e+fx)}{c-d}} \text{AppellF1}\left(m+\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, m+\frac{3}{2}, \frac{1+S\sin(e+fx)}{2}, -\left(\frac{d(1+\sin(e+fx))}{c-d}\right)\right) \text{Cos}[e+fx] \text{Sqrt}[1-S\sin(e+fx)] (a+a\sin(e+fx))^m \text{Sqrt}[\frac{c+d\sin(e+fx)}{c-d}]}{f(2m+1)(c-d)(a-a\sin(e+fx))\sqrt{c+d\sin(e+fx)}}$$

$$\frac{2(Ad^2+c^2C)\cos(e+fx)(a\sin(e+fx)+a)^m}{3df(c^2-d^2)(c+d\sin(e+fx))^{3/2}}$$

input `Int[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(5/2), x]`

output `(2*(c^2*C + A*d^2)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(3*d*(c^2 - d^2)*f*(c + d*Sin[e + f*x])^(3/2)) + ((Sqrt[2]*a^2*(3*c*(A + C)*d + d^2*(A + 3*C - 4*A*m) - 2*c^2*(C + 2*C*m))*AppellF1[1/2 + m, 1/2, 3/2, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[(c + d*Sin[e + f*x])/(c - d)])/((c - d)*f*(1 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]) + (Sqrt[2]*a*(2*c^2*C*(1 + m) - d^2*(A + 3*C - 2*A*m))*AppellF1[3/2 + m, 1/2, 3/2, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[(c + d*Sin[e + f*x])/(c - d)])/((c - d)*f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]))/(3*a*d*(c^2 - d^2))`

---

3.15.  $\int \frac{(a+a\sin(e+fx))^m (A+C\sin^2(e+fx))}{(c+d\sin(e+fx))^{5/2}} dx$

## 3.15.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



rule 3267 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]])*Sqrt[a - b*Sin[e + f*x]]) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]`

rule 3466 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]`

rule 3523 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-(c^2*C + A*d^2))*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(a*d*m + b*c*(n + 1)) + c*C*(a*c*m + b*d*(n + 1)) - b*(A*d^2*(m + n + 2) + C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && (LtQ[n, -1] || EqQ[m + n + 2, 0])`

### 3.15.4 Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + C \sin^2(fx + e))}{(c + d \sin(fx + e))^{\frac{5}{2}}} dx$$

input `int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(5/2),x)`

output `int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(5/2),x)`

---

3.15.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{\frac{5}{2}}} dx$

### 3.15.5 Fracas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{(d \sin(fx + e) + c)^{5/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(5/2),x,  
algorithm="fricas")`

output `integral((C*cos(f*x + e)^2 - A - C)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x +  
e) + a)^m/(3*c*d^2*cos(f*x + e)^2 - c^3 - 3*c*d^2 + (d^3*cos(f*x + e)^2 -  
3*c^2*d - d^3)*sin(f*x + e)), x)`

### 3.15.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(A+C*sin(f*x+e)**2)/(c+d*sin(f*x+e))**(5/2),  
x)`

output `Timed out`

### 3.15.7 Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{(d \sin(fx + e) + c)^{5/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(5/2),x,  
algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/(d*sin(f*x + e) +  
c)^(5/2), x)`

---

3.15.  $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{5/2}} dx$

**3.15.8 Giac [F]**

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{(d \sin(fx + e) + c)^{5/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(5/2),x,  
algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/(d*sin(f*x + e) +  
c)^(5/2), x)`

**3.15.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin^2(e + fx) + A)(a + a \sin(e + fx))^m}{(c + d \sin(e + fx))^{5/2}} dx$$

input `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c + d*sin(e + f*x))^(  
5/2),x)`

output `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c + d*sin(e + f*x))^(  
5/2), x)`

**3.16** 
$$\int \frac{A+B \sin(e+fx)+C \sin^2(e+fx)}{\sqrt{a+a \sin(e+fx)}(c-c \sin(e+fx))^{3/2}} dx$$

3.16.1	Optimal result	155
3.16.2	Mathematica [A] (verified)	155
3.16.3	Rubi [A] (verified)	156
3.16.4	Maple [B] (verified)	159
3.16.5	Fricas [F]	160
3.16.6	Sympy [F]	161
3.16.7	Maxima [F]	161
3.16.8	Giac [A] (verification not implemented)	161
3.16.9	Mupad [F(-1)]	162

**3.16.1 Optimal result**

Integrand size = 50, antiderivative size = 174

$$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \frac{(A + B + C) \cos(e + fx) \sqrt{a + a \sin(e + fx)}}{4af(c - c \sin(e + fx))^{3/2}} - \frac{(A - B - 3C) \cos(e + fx) \log(1 - \sin(e + fx))}{4cf \sqrt{a + a \sin(e + fx)} \sqrt{c - c \sin(e + fx)}} + \frac{(A - B + C) \cos(e + fx) \log(1 + \sin(e + fx))}{4cf \sqrt{a + a \sin(e + fx)} \sqrt{c - c \sin(e + fx)}}$$

output

```
1/4*(A+B+C)*cos(f*x+e)*(a+a*sin(f*x+e))^(1/2)/a/f/(c-c*sin(f*x+e))^(3/2)-1/4*(A-B-3*C)*cos(f*x+e)*ln(1-sin(f*x+e))/c/f/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f*x+e))^(1/2)+1/4*(A-B+C)*cos(f*x+e)*ln(1+sin(f*x+e))/c/f/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f*x+e))^(1/2)
```

**3.16.2 Mathematica [A] (verified)**

Time = 3.13 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.13

$$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \frac{(A + B + C + (-A + B + 3C) \log(\cos(\frac{1}{2}(e + fx))) - \sin(e + fx)) \sqrt{a + a \sin(e + fx)}}{(c - c \sin(e + fx))^{3/2}}$$

input

```
Integrate[(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2)/(Sqrt[a + a*Sin[e + f*x]]*(c - c*Sin[e + f*x])^(3/2)),x]
```

---

3.16. 
$$\int \frac{A+B \sin(e+fx)+C \sin^2(e+fx)}{\sqrt{a+a \sin(e+fx)}(c-c \sin(e+fx))^{3/2}} dx$$

output  $((A + B + C + (-A + B + 3C) \cdot \text{Log}[\text{Cos}[(e + f \cdot x)/2] - \text{Sin}[(e + f \cdot x)/2]]) \cdot (\text{Cos}[(e + f \cdot x)/2] - \text{Sin}[(e + f \cdot x)/2])^2 + (A - B + C) \cdot \text{Log}[\text{Cos}[(e + f \cdot x)/2] + \text{Sin}[(e + f \cdot x)/2]]) \cdot (\text{Cos}[(e + f \cdot x)/2] - \text{Sin}[(e + f \cdot x)/2])^2 \cdot (\text{Cos}[(e + f \cdot x)/2] - \text{Sin}[(e + f \cdot x)/2]) \cdot (\text{Cos}[(e + f \cdot x)/2] + \text{Sin}[(e + f \cdot x)/2])) / (2 \cdot f \cdot \text{Sqrt}[a \cdot (1 + \text{Sin}[e + f \cdot x])] \cdot (c - c \cdot \text{Sin}[e + f \cdot x])^{(3/2)})$

### 3.16.3 Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {3042, 3514, 27, 3042, 3448, 3042, 3216, 3042, 3146, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a \sin(e + fx) + a(c - c \sin(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \sin(e + fx) + C \sin(e + fx)^2}{\sqrt{a \sin(e + fx) + a(c - c \sin(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{3514} \\ & \frac{(A + B + C) \cos(e + fx) \sqrt{a \sin(e + fx) + a}}{4af(c - c \sin(e + fx))^{3/2}} - \frac{\int -\frac{2(a^2(A - B - C) - 2a^2C \sin(e + fx))}{\sqrt{\sin(e + fx)a + a\sqrt{c - c \sin(e + fx)}} dx}{4a^2c} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{a^2(A - B - C) - 2a^2C \sin(e + fx)}{\sqrt{\sin(e + fx)a + a\sqrt{c - c \sin(e + fx)}} dx}{2a^2c} + \frac{(A + B + C) \cos(e + fx) \sqrt{a \sin(e + fx) + a}}{4af(c - c \sin(e + fx))^{3/2}} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{a^2(A - B - C) - 2a^2C \sin(e + fx)}{\sqrt{\sin(e + fx)a + a\sqrt{c - c \sin(e + fx)}} dx}{2a^2c} + \frac{(A + B + C) \cos(e + fx) \sqrt{a \sin(e + fx) + a}}{4af(c - c \sin(e + fx))^{3/2}} \\ & \quad \downarrow \text{3448} \\ & \frac{a^2(A - B + C) \int \frac{\sqrt{c - c \sin(e + fx)}}{\sqrt{\sin(e + fx)a + a}} dx}{2c} + \frac{1}{2}a(A - B - 3C) \int \frac{\sqrt{\sin(e + fx)a + a}}{\sqrt{c - c \sin(e + fx)}} dx}{2a^2c} + \\ & \quad \frac{(A + B + C) \cos(e + fx) \sqrt{a \sin(e + fx) + a}}{4af(c - c \sin(e + fx))^{3/2}} \end{aligned}$$

---

3.16.  $\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)(c - c \sin(e + fx))^{3/2}} dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \frac{a^2(A-B+C) \int \frac{\sqrt{c-c\sin(e+fx)}}{\sqrt{\sin(e+fx)a+a}} dx + \frac{1}{2}a(A-B-3C) \int \frac{\sqrt{\sin(e+fx)a+a}}{\sqrt{c-c\sin(e+fx)}} dx}{2c} + \\
& \frac{2a^2c}{(A+B+C)\cos(e+fx)\sqrt{a\sin(e+fx)+a}} + \\
& \frac{4af(c-c\sin(e+fx))^{3/2}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \downarrow \text{3216} \\
& \frac{a^3(A-B+C)\cos(e+fx) \int \frac{\cos(e+fx)}{\sin(e+fx)a+a} dx + \frac{a^2c(A-B-3C)\cos(e+fx) \int \frac{\cos(e+fx)}{c-c\sin(e+fx)} dx}{2\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \\
& \frac{2a^2c}{(A+B+C)\cos(e+fx)\sqrt{a\sin(e+fx)+a}} + \\
& \frac{4af(c-c\sin(e+fx))^{3/2}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \downarrow \text{3042} \\
& \frac{a^3(A-B+C)\cos(e+fx) \int \frac{\cos(e+fx)}{\sin(e+fx)a+a} dx + \frac{a^2c(A-B-3C)\cos(e+fx) \int \frac{\cos(e+fx)}{c-c\sin(e+fx)} dx}{2\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \\
& \frac{2a^2c}{(A+B+C)\cos(e+fx)\sqrt{a\sin(e+fx)+a}} + \\
& \frac{4af(c-c\sin(e+fx))^{3/2}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \downarrow \text{3146} \\
& \frac{a^2(A-B+C)\cos(e+fx) \int \frac{1}{\sin(e+fx)a+a} d(a\sin(e+fx)) - \frac{a^2(A-B-3C)\cos(e+fx) \int \frac{1}{c-c\sin(e+fx)} d(-c\sin(e+fx))}{2f\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} - \\
& \frac{2a^2c}{(A+B+C)\cos(e+fx)\sqrt{a\sin(e+fx)+a}} + \\
& \frac{4af(c-c\sin(e+fx))^{3/2}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \downarrow \text{16} \\
& \frac{a^2(A-B+C)\cos(e+fx)\log(a\sin(e+fx)+a) - \frac{a^2(A-B-3C)\cos(e+fx)\log(c-c\sin(e+fx))}{2f\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \\
& \frac{2a^2c}{(A+B+C)\cos(e+fx)\sqrt{a\sin(e+fx)+a}} + \\
& \frac{4af(c-c\sin(e+fx))^{3/2}}{4af(c-c\sin(e+fx))^{3/2}}
\end{aligned}$$

input `Int[(A + B*SIN[e + f*x] + C*SIN[e + f*x]^2)/(Sqrt[a + a*SIN[e + f*x]]*(c - c*SIN[e + f*x])^(3/2)),x]`

output `((A + B + C)*Cos[e + f*x]*Sqrt[a + a*SIN[e + f*x]])/(4*a*f*(c - c*SIN[e + f*x])^(3/2)) + ((a^2*(A - B + C)*Cos[e + f*x]*Log[a + a*SIN[e + f*x]])/(2*f*Sqrt[a + a*SIN[e + f*x]]*Sqrt[c - c*SIN[e + f*x]]) - (a^2*(A - B - 3*C)*Cos[e + f*x]*Log[c - c*SIN[e + f*x]])/(2*f*Sqrt[a + a*SIN[e + f*x]]*Sqrt[c - c*SIN[e + f*x]))/(2*a^2*c)`

---

3.16.  $\int \frac{A+B\sin(e+fx)+C\sin^2(e+fx)}{\sqrt{a+a\sin(e+fx)}(c-c\sin(e+fx))^{3/2}} dx$

## 3.16.3.1 Defintions of rubi rules used

- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3146 `Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1)/2, x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])`
- rule 3216 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[a*c*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])) Int[Cos[e + f*x]/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`
- rule 3448 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A*b + a*B)/(2*a*b) Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[(B*c + A*d)/(2*c*d) Int[Sqrt[c + d*Sin[e + f*x]]/Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`

```
rule 3514 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_)
+ (f_)*(x_)]^2), x_Symbol] :> Simp[(a*A - b*B + a*C)*Cos[e + f*x]*(a + b*
Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(2*b*c*f*(2*m + 1))), x] - Si
mp[1/(2*b*c*d*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e +
f*x])^n*Simp[A*(c^2*(m + 1) + d^2*(2*m + n + 2)) - B*c*d*(m - n - 1) - C*(c
^2*m - d^2*(n + 1)) + d*((A*c + B*d)*(m + n + 2) - c*C*(3*m - n))*Sin[e + f
*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && EqQ[b*c +
a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (EqQ[m + n + 2, 0] &&
NeQ[2*m + 1, 0]))
```

### 3.16.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 746 vs.  $2(156) = 312$ .

Time = 3.90 (sec) , antiderivative size = 747, normalized size of antiderivative = 4.29

method	result
default	$\frac{-A \sin(fx+e) - C - A - B - C \sin(fx+e) + A \sin(fx+e) \cos(fx+e) \ln(-\cot(fx+e) + \csc(fx+e) + 1) - A \ln(\csc(fx+e) - \cot(fx+e) - 1)}{\dots}$
parts	Expression too large to display

```
input int((A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2)/(a+a*sin(f*x+e)
)^(1/2),x,method=_RETURNVERBOSE)
```

---

3.16. 
$$\int \frac{A+B \sin(e+fx)+C \sin^2(e+fx)}{\sqrt{a+a \sin(e+fx)}(c-c \sin(e+fx))^{3/2}} dx$$



output `1/2/c/f*(-A*sin(f*x+e)-C*cos(f*x+e)^2*ln(-cot(f*x+e)+csc(f*x+e)+1)-3*C*cos(f*x+e)^2*ln(csc(f*x+e)-cot(f*x+e)-1)+2*C*cos(f*x+e)^2*ln(2/(1+cos(f*x+e)))-C-A-B-C*sin(f*x+e)+A*sin(f*x+e)*cos(f*x+e)*ln(-cot(f*x+e)+csc(f*x+e)+1)-A*ln(csc(f*x+e)-cot(f*x+e)-1)*sin(f*x+e)*cos(f*x+e)-B*sin(f*x+e)*cos(f*x+e)*ln(-cot(f*x+e)+csc(f*x+e)+1)+B*ln(csc(f*x+e)-cot(f*x+e)-1)*sin(f*x+e)*cos(f*x+e)-B*sin(f*x+e)+B*cos(f*x+e)^2+A*cos(f*x+e)^2+C*cos(f*x+e)^2+A*cos(f*x+e)*ln(csc(f*x+e)-cot(f*x+e)-1)-A*sin(f*x+e)*cos(f*x+e)-B*cos(f*x+e)*sin(f*x+e)+C*ln(-cot(f*x+e)+csc(f*x+e)+1)*sin(f*x+e)*cos(f*x+e)+3*C*ln(csc(f*x+e)-cot(f*x+e)-1)*sin(f*x+e)*cos(f*x+e)-2*C*ln(2/(1+cos(f*x+e)))*sin(f*x+e)*cos(f*x+e)-C*sin(f*x+e)*cos(f*x+e)-C*cos(f*x+e)*ln(-cot(f*x+e)+csc(f*x+e)+1)-3*C*cos(f*x+e)*ln(csc(f*x+e)-cot(f*x+e)-1)+2*C*cos(f*x+e)*ln(2/(1+cos(f*x+e)))-B*cos(f*x+e)^2*ln(csc(f*x+e)-cot(f*x+e)-1)+A*cos(f*x+e)^2*ln(csc(f*x+e)-cot(f*x+e)-1)-B*cos(f*x+e)*ln(csc(f*x+e)-cot(f*x+e)-1)-A*cos(f*x+e)*ln(-cot(f*x+e)+csc(f*x+e)+1)+B*cos(f*x+e)*ln(-cot(f*x+e)+csc(f*x+e)+1)+B*cos(f*x+e)^2*ln(-cot(f*x+e)+csc(f*x+e)+1)-A*cos(f*x+e)^2*ln(-cot(f*x+e)+csc(f*x+e)+1))/(-cos(f*x+e)+sin(f*x+e)-1)/(-c*(sin(f*x+e)-1))^(1/2)/(a*(1+sin(f*x+e)))^(1/2)`

### 3.16.5 Fricas [F]

$$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{C \sin(fx + e)^2 + B \sin(fx + e) + A}{\sqrt{a \sin(fx + e) + a}(-c \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2)/(a+a*sin(f*x+e))^(1/2),x, algorithm="fricas")`

output `integral((C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*sqrt(a*sin(f*x + e) + a)*sqrt(-c*sin(f*x + e) + c)/(a*c^2*cos(f*x + e)^2*sin(f*x + e) - a*c^2*cos(f*x + e)^2), x)`

### 3.16.6 Sympy [F]

$$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a(\sin(e + fx) + 1)}(-c(\sin(e + fx) - 1))^{3/2}} dx$$

input `integrate((A+B*sin(f*x+e)+C*sin(f*x+e)**2)/(c-c*sin(f*x+e))**(3/2)/(a+a*sin(f*x+e))**(1/2),x)`

output `Integral((A + B*sin(e + f*x) + C*sin(e + f*x)**2)/(sqrt(a*(sin(e + f*x) + 1))*(-c*(sin(e + f*x) - 1))**(3/2)), x)`

### 3.16.7 Maxima [F]

$$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{C \sin^2(fx + e) + B \sin(fx + e) + A}{\sqrt{a \sin(fx + e) + a}(-c \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2)/(a+a*sin(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)/(sqrt(a*sin(f*x + e) + a))*(-c*sin(f*x + e) + c)^(3/2)), x)`

### 3.16.8 Giac [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.26

$$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \frac{(A\sqrt{a} - B\sqrt{a} - 3C\sqrt{a}) \log\left(-\cos\left(-\frac{1}{4}\pi + \frac{1}{2}fx + \frac{1}{2}e\right)^2 + 1\right)}{ac^{\frac{3}{2}} \operatorname{sgn}\left(\cos\left(-\frac{1}{4}\pi + \frac{1}{2}fx + \frac{1}{2}e\right)\right) \operatorname{sgn}\left(\sin\left(-\frac{1}{4}\pi + \frac{1}{2}fx + \frac{1}{2}e\right)\right)} - \frac{2(A\sqrt{a} - B\sqrt{a})}{ac^{\frac{3}{2}} \operatorname{sgn}\left(\cos\left(-\frac{1}{4}\pi + \frac{1}{2}fx + \frac{1}{2}e\right)\right)}$$

input `integrate((A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2)/(a+a*sin(f*x+e))^(1/2),x, algorithm="giac")`

output `1/4*((A*sqrt(a) - B*sqrt(a) - 3*C*sqrt(a))*log(-cos(-1/4*pi + 1/2*f*x + 1/2*e)^2 + 1)/(a*c^(3/2)*sgn(cos(-1/4*pi + 1/2*f*x + 1/2*e))*sgn(sin(-1/4*pi + 1/2*f*x + 1/2*e))) - 2*(A*sqrt(a) - B*sqrt(a) + C*sqrt(a))*log(abs(cos(-1/4*pi + 1/2*f*x + 1/2*e)))/(a*c^(3/2)*sgn(cos(-1/4*pi + 1/2*f*x + 1/2*e))*sgn(sin(-1/4*pi + 1/2*f*x + 1/2*e))) + (A*sqrt(a) + B*sqrt(a) + C*sqrt(a))/((cos(-1/4*pi + 1/2*f*x + 1/2*e)^2 - 1)*a*c^(3/2)*sgn(cos(-1/4*pi + 1/2*f*x + 1/2*e))*sgn(sin(-1/4*pi + 1/2*f*x + 1/2*e)))/f`

### 3.16.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{C \sin(e + fx)^2 + B \sin(e + fx) + A}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx$$

input `int((A + B*sin(e + f*x) + C*sin(e + f*x)^2)/((a + a*sin(e + f*x))^(1/2)*(c - c*sin(e + f*x))^(3/2)),x)`

output `int((A + B*sin(e + f*x) + C*sin(e + f*x)^2)/((a + a*sin(e + f*x))^(1/2)*(c - c*sin(e + f*x))^(3/2)), x)`

### 3.17 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^n (A + B \sin(e +$

3.17.1	Optimal result	163
3.17.2	Mathematica [F]	164
3.17.3	Rubi [A] (verified)	164
3.17.4	Maple [F]	168
3.17.5	Fricas [F]	168
3.17.6	Sympy [F(-1)]	169
3.17.7	Maxima [F]	169
3.17.8	Giac [F]	169
3.17.9	Mupad [F(-1)]	170

#### 3.17.1 Optimal result

Integrand size = 46, antiderivative size = 269

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \frac{2^{\frac{1}{2}+n} c ((1 + m + n)(C(1 - m + n) + A(2 + m + n)) + (m - n)(C + 2Cm + B(2 + m + n))) \cos(e + fx) - \frac{(C + 2Cm + B(2 + m + n)) \cos(e + fx) (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n}{f(1 + m + n)(2 + m + n)} + \frac{C \cos(e + fx) (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{1+n}}{cf(2 + m + n)}$$

output

```
2^(1/2+n)*c*((1+m+n)*(C*(1-m+n)+A*(2+m+n))+(m-n)*(C+2*C*m+B*(2+m+n)))*cos(
f*x+e)*hypergeom([1/2-n, 1/2+m],[3/2+m],1/2+1/2*sin(f*x+e))*(1-sin(f*x+e))
^(1/2-n)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1+n)/f/(1+2*m)/(1+m+n)/(2+m
+n)-(C+2*C*m+B*(2+m+n))*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n/f
/(1+m+n)/(2+m+n)+C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1+n)/c/
f/(2+m+n)
```

### 3.17.2 Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input `Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x]`

### 3.17.3 Rubi [A] (verified)

Time = 1.20 (sec) , antiderivative size = 318, normalized size of antiderivative = 1.18, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.239$ , Rules used = {3042, 3518, 25, 3042, 3452, 3042, 3224, 3042, 3168, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (a \sin(e + fx) + a)^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin(e + fx)^2) dx$$

$$\downarrow \text{3518}$$

$$\frac{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{n+1}}{cf(m + n + 2)}$$

$$\frac{\int -(\sin(e + fx)a + a)^m (c - c \sin(e + fx))^n (ac(C(-m + n + 1) + A(m + n + 2)) + ac(2mC + C + B(m + n + 2))) dx}{ac(m + n + 2)}$$

$$\downarrow \text{25}$$

$$\frac{\int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^n (ac(C(-m + n + 1) + A(m + n + 2)) + ac(2mC + C + B(m + n + 2))) dx}{ac(m + n + 2)}$$

$$\frac{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{n+1}}{cf(m + n + 2)}$$

---

3.17.  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^m (c - c\sin(e + fx))^n (ac(C(-m + n + 1) + A(m + n + 2)) + ac(2mC + C + B(m + n + 2)))}{ac(m + n + 2)} \\ \frac{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c\sin(e + fx))^{n+1}}{cf(m + n + 2)}$$

↓ 3452

$$\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+(m-n)(B(m+n+2)+2Cm+C)) \int (\sin(e+fx)a+a)^m (c-c\sin(e+fx))^n dx}{m+n+1} - \frac{ac(B(m+n+2)+2Cm+C)}{m+n+1} \\ \frac{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c\sin(e + fx))^{n+1}}{cf(m + n + 2)}$$

↓ 3042

$$\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+(m-n)(B(m+n+2)+2Cm+C)) \int (\sin(e+fx)a+a)^m (c-c\sin(e+fx))^n dx}{m+n+1} - \frac{ac(B(m+n+2)+2Cm+C)}{m+n+1} \\ \frac{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c\sin(e + fx))^{n+1}}{cf(m + n + 2)}$$

↓ 3224

$$\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+(m-n)(B(m+n+2)+2Cm+C)) \cos^{-2m}(e+fx)(a \sin(e+fx)+a)^m (c-c\sin(e+fx))^m \int \cos^{2m}(e+fx)}{m+n+1} \\ \frac{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c\sin(e + fx))^{n+1}}{cf(m + n + 2)} \quad ac(m + n + 2)$$

↓ 3042

$$\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+(m-n)(B(m+n+2)+2Cm+C)) \cos^{-2m}(e+fx)(a \sin(e+fx)+a)^m (c-c\sin(e+fx))^m \int \cos(e+fx)^{2m}}{m+n+1} \\ \frac{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c\sin(e + fx))^{n+1}}{cf(m + n + 2)} \quad ac(m + n + 2)$$

↓ 3168

$$\frac{ac^3 \cos(e+fx)((m+n+1)(A(m+n+2)+C(-m+n+1))+(m-n)(B(m+n+2)+2Cm+C))(a \sin(e+fx)+a)^m (c-c\sin(e+fx))^{\frac{1}{2}(-2m-1)+m} (c \sin(e+fx))^{m-1}}{f(m+n+1)} \\ \frac{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c\sin(e + fx))^{n+1}}{cf(m + n + 2)}$$

↓ 80

---

3.17.  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$

$$\frac{ac^3 2^{n-\frac{1}{2}} \cos(e+fx)((m+n+1)(A(m+n+2)+C(-m+n+1))+(m-n)(B(m+n+2)+2Cm+C))(1-\sin(e+fx))^{\frac{1}{2}-n} (a \sin(e+fx)+a)^m (c \sin(e+fx)+a)^n}{f(m+n+1)}$$

$$\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c \sin(e+fx))^{n+1}}{cf(m+n+2)}$$

↓ 79

$$\frac{ac^2 2^{n+\frac{1}{2}} \cos(e+fx)((m+n+1)(A(m+n+2)+C(-m+n+1))+(m-n)(B(m+n+2)+2Cm+C))(1-\sin(e+fx))^{\frac{1}{2}-n} (a \sin(e+fx)+a)^m (c \sin(e+fx)+a)^n}{f(2m+1)(m+n+1)}$$

$$\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c \sin(e+fx))^{n+1}}{cf(m+n+2)}$$

```
input Int[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]
```

```
output (C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(1 + n))/(c*f*(2 + m + n)) + (-((a*c*(C + 2*C*m + B*(2 + m + n))*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n)/(f*(1 + m + n))) + (2^(1/2 + n)*a*c^2*((1 + m + n)*(C*(1 - m + n) + A*(2 + m + n)) + (m - n)*(C + 2*C*m + B*(2 + m + n)))*Cos[e + f*x]*Hypergeometric2F1[(1 + 2*m)/2, (1 - 2*n)/2, (3 + 2*m)/2, (1 + Sin[e + f*x])/2]*(1 - Sin[e + f*x])^(1/2 - n)*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(-1/2 + (-1 - 2*m)/2 + m + n)*(c + c*Sin[e + f*x])^((-1 - 2*m)/2 + (1 + 2*m)/2))/(f*(1 + 2*m)*(1 + m + n))/(a*c*(2 + m + n))
```

### 3.17.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 79 Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))
```

---

3.17.  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$

- rule 80 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n, x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3168 `Int[(cos[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[a^2*((g*cos[e + f*x])^(p + 1)/(f*g*(a + b*sin[e + f*x])^((p + 1)/2)*(a - b*sin[e + f*x])^((p + 1)/2))) Subst[Int[(a + b*x)^(m + (p - 1)/2)*(a - b*x)^((p - 1)/2), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, e, f, g, m, p}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[m]`
- rule 3224 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*sin[e + f*x])^FracPart[m]*((c + d*sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])`
- rule 3452 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*sin[e + f*x])^m*((c + d*sin[e + f*x])^n/(f*(m + n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(a + b*sin[e + f*x])^m*(c + d*sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 1, 0]`



rule 3518 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + (b*B*d*(m + n + 2) - b*c*C*(2*m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]`

### 3.17.4 Maple [F]

$$\int (a + a \sin(fx + e))^m (c - c \sin(fx + e))^n (A + B \sin(fx + e) + C(\sin^2(fx + e))) dx$$

input `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2), x)`

output `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2), x)`

### 3.17.5 Fricas [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^n dx$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*(a*sin(f*x + e) + a)^m*(-c*sin(f*x + e) + c)^n, x)`

**3.17.6 Sympy [F(-1)]**

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

= Timed out

input `integrate((a+a*sin(f*x+e))**m*(c-c*sin(f*x+e))**n*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output Timed out

**3.17.7 Maxima [F]**

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(fx + e) + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^n dx$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m*(-c*sin(f*x + e) + c)^n, x)`

**3.17.8 Giac [F]**

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(fx + e) + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^n dx$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m*(-c*sin(f*x + e) + c)^n, x)`

**3.17.9 Mupad [F(-1)]**

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (C \sin(e + fx)^2 + B \sin(e + fx) + A) dx$$

input `int((a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^n*(A + B*sin(e + f*x) + C*  
sin(e + f*x)^2),x)`

output `int((a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^n*(A + B*sin(e + f*x) + C*  
sin(e + f*x)^2), x)`

### 3.18 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^{5/2} (A + B \sin(e -$

3.18.1	Optimal result . . . . .	171
3.18.2	Mathematica [C] (verified) . . . . .	172
3.18.3	Rubi [A] (verified) . . . . .	173
3.18.4	Maple [F] . . . . .	176
3.18.5	Fricas [B] (verification not implemented) . . . . .	177
3.18.6	Sympy [F(-1)] . . . . .	178
3.18.7	Maxima [B] (verification not implemented) . . . . .	178
3.18.8	Giac [F] . . . . .	179
3.18.9	Mupad [B] (verification not implemented) . . . . .	180

#### 3.18.1 Optimal result

Integrand size = 48, antiderivative size = 435

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx =$$

$$\frac{64c^3(B(45 - 8m - 4m^2) - C(39 - 16m + 4m^2) - A(63 + 32m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m}{f(5 + 2m)(7 + 2m)(9 + 2m)(3 + 8m + 4m^2) \sqrt{c - c \sin(e + fx)}} -$$

$$\frac{16c^2(B(45 - 8m - 4m^2) - C(39 - 16m + 4m^2) - A(63 + 32m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m}{f(7 + 2m)(9 + 2m)(15 + 16m + 4m^2)} -$$

$$\frac{2c(B(45 - 8m - 4m^2) - C(39 - 16m + 4m^2) - A(63 + 32m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2}}{f(5 + 2m)(7 + 2m)(9 + 2m)} -$$

$$\frac{2(9B + 2C + 2Bm + 4Cm) \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2}}{f(7 + 2m)(9 + 2m)} +$$

$$\frac{2C \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{7/2}}{cf(9 + 2m)}$$

output

```
-2*c*(B*(-4*m^2-8*m+45)-C*(4*m^2-16*m+39)-A*(4*m^2+32*m+63))*cos(f*x+e)*(a
+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)/f/(8*m^3+84*m^2+286*m+315)-2*(2*B*
m+4*C*m+9*B+2*C)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)/f/(4
*m^2+32*m+63)+2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(7/2)/c/f
/(9+2*m)-64*c^3*(B*(-4*m^2-8*m+45)-C*(4*m^2-16*m+39)-A*(4*m^2+32*m+63))*co
s(f*x+e)*(a+a*sin(f*x+e))^m/f/(32*m^5+400*m^4+1840*m^3+3800*m^2+3378*m+945
)/(c-c*sin(f*x+e))^(1/2)-16*c^2*(B*(-4*m^2-8*m+45)-C*(4*m^2-16*m+39)-A*(4*
m^2+32*m+63))*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)/f/(16*m
^4+192*m^3+824*m^2+1488*m+945)
```

---

3.18.  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$

### 3.18.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 13.21 (sec) , antiderivative size = 1029, normalized size of antiderivative = 2.37

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \frac{(a(1 + \sin(e + fx)))^m (c - c \sin(e + fx))^{5/2} \left( \frac{(18900A - 14175B + 12285C + 15648Am - 4140Bm + 648Cm + 5280A^2m - 832B^2m + 1416Cm^2 + 896A^3m - 208B^3m + 224C^3m + 64A^4m - 16B^4m + 16C^4m)}{(1 + 2m)(3 + 2m)(5 + 2m)(7 + 2m)(9 + 2m)} + \frac{(18900A - 14175B + 12285C + 15648Am - 4140Bm + 648Cm + 5280A^2m - 832B^2m + 1416Cm^2 + 896A^3m - 208B^3m + 224C^3m + 64A^4m - 16B^4m + 16C^4m)}{(1 + 2m)(3 + 2m)(5 + 2m)(7 + 2m)(9 + 2m)} + \frac{(3150A - 3465B + 3150C + 2356Am - 1706Bm + 828Cm + 584A^2m - 316B^2m + 200C^2m + 48A^3m - 24B^3m + 16C^3m)}{(3 + 2m)(5 + 2m)(7 + 2m)(9 + 2m)} + \frac{(3150A - 3465B + 3150C + 2356Am - 1706Bm + 828Cm + 584A^2m - 316B^2m + 200C^2m + 48A^3m - 24B^3m + 16C^3m)}{(3 + 2m)(5 + 2m)(7 + 2m)(9 + 2m)} + \frac{(126A - 315B + 378C + 64Am - 124Bm + 88Cm + 8A^2m - 12B^2m + 8C^2m)}{(5 + 2m)(7 + 2m)(9 + 2m)} + \frac{(126A - 315B + 378C + 64Am - 124Bm + 88Cm + 8A^2m - 12B^2m + 8C^2m)}{(5 + 2m)(7 + 2m)(9 + 2m)} + \frac{(-1/8 + I/8) \cos[(5(e + fx))/2] - (1/8 + I/8) \sin[(5(e + fx))/2]}{(5 + 2m)(7 + 2m)(9 + 2m)} + \frac{(-1/8 - I/8) \cos[(5(e + fx))/2] - (1/8 - I/8) \sin[(5(e + fx))/2]}{(5 + 2m)(7 + 2m)(9 + 2m)} \right)}{(1 + 2m)(3 + 2m)(5 + 2m)(7 + 2m)(9 + 2m)}$$

input `Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(5/2)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `((a*(1 + Sin[e + f*x]))^m*(c - c*Sin[e + f*x])^(5/2)*(((18900*A - 14175*B + 12285*C + 15648*A*m - 4140*B*m + 648*C*m + 5280*A*m^2 - 832*B*m^2 + 1416*C*m^2 + 896*A*m^3 - 208*B*m^3 + 224*C*m^3 + 64*A*m^4 - 16*B*m^4 + 16*C*m^4)*((1/8 + I/8)*Cos[(e + f*x)/2] + (1/8 - I/8)*Sin[(e + f*x)/2]))/((1 + 2*m)*(3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((18900*A - 14175*B + 12285*C + 15648*A*m - 4140*B*m + 648*C*m + 5280*A*m^2 - 832*B*m^2 + 1416*C*m^2 + 896*A*m^3 - 208*B*m^3 + 224*C*m^3 + 64*A*m^4 - 16*B*m^4 + 16*C*m^4)*((1/8 - I/8)*Cos[(e + f*x)/2] + (1/8 + I/8)*Sin[(e + f*x)/2]))/((1 + 2*m)*(3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((3150*A - 3465*B + 3150*C + 2356*A*m - 1706*B*m + 828*C*m + 584*A*m^2 - 316*B*m^2 + 200*C*m^2 + 48*A*m^3 - 24*B*m^3 + 16*C*m^3)*((1/8 - I/8)*Cos[(3*(e + f*x))/2] - (1/8 + I/8)*Sin[(3*(e + f*x))/2]))/((3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((3150*A - 3465*B + 3150*C + 2356*A*m - 1706*B*m + 828*C*m + 584*A*m^2 - 316*B*m^2 + 200*C*m^2 + 48*A*m^3 - 24*B*m^3 + 16*C*m^3)*((1/8 + I/8)*Cos[(3*(e + f*x))/2] - (1/8 - I/8)*Sin[(3*(e + f*x))/2]))/((3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((126*A - 315*B + 378*C + 64*A*m - 124*B*m + 88*C*m + 8*A*m^2 - 12*B*m^2 + 8*C*m^2)*((-1/8 + I/8)*Cos[(5*(e + f*x))/2] - (1/8 + I/8)*Sin[(5*(e + f*x))/2]))/((5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((126*A - 315*B + 378*C + 64*A*m - 124*B*m + 88*C*m + 8*A*m^2 - 12*B*m^2 + 8*C*m^2)*((-1/8 - I/8)*Cos[(5*(e + f*x))/2] - (1/8 - I/8)*Sin[(5*(e + f*x))/2]))/((5 + 2*m)*(7 + 2*m)*(9 + 2*m))`

### 3.18.3 Rubi [A] (verified)

Time = 1.49 (sec) , antiderivative size = 336, normalized size of antiderivative = 0.77, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$ , Rules used = {3042, 3518, 27, 3042, 3452, 3042, 3219, 3042, 3219, 3042, 3217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2) dx$$

$$\downarrow \text{3518}$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2} (a \sin(e + fx) + a)^m}{cf(2m + 9)} - \frac{2 \int -\frac{1}{2} (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{5/2} (ac(C(7 - 2m) + A(2m + 9)) + ac(2mB + 9B + 2C + 4Cm) \sin(e + fx)) dx}{ac(2m + 9)}}$$

$$\downarrow \text{27}$$

$$\frac{\int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{5/2} (ac(C(7 - 2m) + A(2m + 9)) + ac(2mB + 9B + 2C + 4Cm) \sin(e + fx)) dx}{ac(2m + 9)} - \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2} (a \sin(e + fx) + a)^m}{cf(2m + 9)}$$

$$\downarrow \text{3042}$$

$$\frac{\int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{5/2} (ac(C(7 - 2m) + A(2m + 9)) + ac(2mB + 9B + 2C + 4Cm) \sin(e + fx)) dx}{ac(2m + 9)} - \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2} (a \sin(e + fx) + a)^m}{cf(2m + 9)}$$

$$\downarrow \text{3452}$$

$$\frac{-\frac{ac(-A(4m^2 + 32m + 63) + B(-4m^2 - 8m + 45) - C(4m^2 - 16m + 39)) \int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{5/2} dx}{2m + 7} - \frac{2ac(2Bm + 9B + 4Cm + 2C) \cos(e + fx)(c - c \sin(e + fx))^{7/2} (a \sin(e + fx) + a)^m}{ac(2m + 9)}}{cf(2m + 9)}$$

$$\downarrow \text{3042}$$

---


$$3.18. \quad \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$\frac{ac(-A(4m^2+32m+63)+B(-4m^2-8m+45)-C(4m^2-16m+39)) \int (\sin(e+fx)a+a)^m (c-c\sin(e+fx))^{5/2} dx}{2m+7} - \frac{2ac(2Bm+9B+4Cm+2C) \cos(e+fx)}{f(2m+5)}$$

$$\frac{2C \cos(e+fx)(c-c\sin(e+fx))^{7/2} (a \sin(e+fx) + a)^m}{cf(2m+9)}$$

↓ 3219

$$\frac{ac(-A(4m^2+32m+63)+B(-4m^2-8m+45)-C(4m^2-16m+39)) \left( \frac{8c \int (\sin(e+fx)a+a)^m (c-c\sin(e+fx))^{3/2} dx}{2m+5} + \frac{2c \cos(e+fx)(c-c\sin(e+fx))^{3/2} (a \sin(e+fx) + a)^m}{f(2m+5)} \right)}{2m+7}$$

$$\frac{2C \cos(e+fx)(c-c\sin(e+fx))^{7/2} (a \sin(e+fx) + a)^m}{cf(2m+9)}$$

↓ 3042

$$\frac{ac(-A(4m^2+32m+63)+B(-4m^2-8m+45)-C(4m^2-16m+39)) \left( \frac{8c \int (\sin(e+fx)a+a)^m (c-c\sin(e+fx))^{3/2} dx}{2m+5} + \frac{2c \cos(e+fx)(c-c\sin(e+fx))^{3/2} (a \sin(e+fx) + a)^m}{f(2m+5)} \right)}{2m+7}$$

$$\frac{2C \cos(e+fx)(c-c\sin(e+fx))^{7/2} (a \sin(e+fx) + a)^m}{cf(2m+9)}$$

↓ 3219

$$\frac{ac(-A(4m^2+32m+63)+B(-4m^2-8m+45)-C(4m^2-16m+39)) \left( \frac{8c \left( \frac{4c \int (\sin(e+fx)a+a)^m \sqrt{c-c\sin(e+fx)} dx}{2m+3} + \frac{2c \cos(e+fx) \sqrt{c-c\sin(e+fx)} (a \sin(e+fx) + a)^m}{f(2m+3)} \right)}{2m+5} \right)}{2m+7}$$

$$\frac{2C \cos(e+fx)(c-c\sin(e+fx))^{7/2} (a \sin(e+fx) + a)^m}{cf(2m+9)}$$

↓ 3042

$$\frac{ac(-A(4m^2+32m+63)+B(-4m^2-8m+45)-C(4m^2-16m+39)) \left( \frac{8c \left( \frac{4c \int (\sin(e+fx)a+a)^m \sqrt{c-c\sin(e+fx)} dx}{2m+3} + \frac{2c \cos(e+fx) \sqrt{c-c\sin(e+fx)} (a \sin(e+fx) + a)^m}{f(2m+3)} \right)}{2m+5} \right)}{2m+7}$$

$$\frac{2C \cos(e+fx)(c-c\sin(e+fx))^{7/2} (a \sin(e+fx) + a)^m}{cf(2m+9)}$$

↓ 3217

---

3.18.  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$

$$\frac{ac(-A(4m^2+32m+63)+B(-4m^2-8m+45)-C(4m^2-16m+39)) \left( \frac{8c \left( \frac{8c^2 \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)(2m+3)\sqrt{c-c \sin(e+fx)}} + \frac{2c \cos(e+fx)\sqrt{c-c \sin(e+fx)}(a \sin(e+fx))}{f(2m+3)} \right)}{2m+5} \right)}{2m+7} \quad ac(2m +$$

$$\frac{2C \cos(e+fx)(c-c \sin(e+fx))^{7/2}(a \sin(e+fx)+a)^m}{cf(2m+9)}$$

input `Int[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(5/2)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `(2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(7/2))/(c*f*(9 + 2*m)) + ((-2*a*c*(9*B + 2*C + 2*B*m + 4*C*m)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(5/2))/(f*(7 + 2*m)) - (a*c*(B*(45 - 8*m - 4*m^2) - C*(39 - 16*m + 4*m^2) - A*(63 + 32*m + 4*m^2))*((2*c*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2))/(f*(5 + 2*m)) + (8*c*((8*c^2*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) + (2*c*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*Sqrt[c - c*Sin[e + f*x]])/(f*(3 + 2*m))))/(5 + 2*m)))/(7 + 2*m))/(a*c*(9 + 2*m))`

### 3.18.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3217 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[n, -2^(-1)]`



rule 3219 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^n/(f*(m + n))), x] + Simp[a*((2*m - 1)/(m + n)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[m - 1/2, 0] && !LtQ[n, -1] && !(IGtQ[n - 1/2, 0] && LtQ[n, m]) && !(ILtQ[m + n, 0] && GtQ[2*m + n + 1, 0])`

rule 3452 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 1, 0]`

rule 3518 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + (b*B*d*(m + n + 2) - b*c*C*(2*m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]`

### 3.18.4 Maple [F]

$$\int (a + a \sin(fx + e))^m (c - c \sin(fx + e))^{\frac{5}{2}} (A + B \sin(fx + e) + C \sin^2(fx + e)) dx$$

input `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

output `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

### 3.18.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 937 vs.  $2(403) = 806$ .

Time = 0.38 (sec) , antiderivative size = 937, normalized size of antiderivative = 2.15

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Too large to display}$$

```
input integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+B*sin(f*x+e)+C*sin(
f*x+e)^2),x, algorithm="fracas")
```

```
output 2*((16*C*c^2*m^4 + 128*C*c^2*m^3 + 344*C*c^2*m^2 + 352*C*c^2*m + 105*C*c^2
)*cos(f*x + e)^5 + 128*(A + B + C)*c^2*m^2 + (16*(B - C)*c^2*m^4 + 16*(9*B
- 14*C)*c^2*m^3 + 8*(52*B - 97*C)*c^2*m^2 + 4*(111*B - 226*C)*c^2*m + 15*
(9*B - 19*C)*c^2)*cos(f*x + e)^4 + 256*(4*A + B - 2*C)*c^2*m - (16*(A - 2*
B + 3*C)*c^2*m^4 + 16*(10*A - 23*B + 32*C)*c^2*m^3 + 8*(65*A - 169*B + 253
*C)*c^2*m^2 + 4*(150*A - 417*B + 656*C)*c^2*m + 3*(63*A - 180*B + 289*C)*c
^2)*cos(f*x + e)^3 + 96*(21*A - 15*B + 13*C)*c^2 + (16*(A - B + C)*c^2*m^4
+ 32*(7*A - 5*B + 7*C)*c^2*m^3 + 8*(133*A - 97*B + 85*C)*c^2*m^2 + 8*(233
*A - 235*B + 233*C)*c^2*m + 3*(231*A - 255*B + 263*C)*c^2)*cos(f*x + e)^2
+ 2*(16*(A - B + C)*c^2*m^4 + 192*(A - B + C)*c^2*m^3 + 8*(107*A - 99*B +
107*C)*c^2*m^2 + 16*(109*A - 89*B + 85*C)*c^2*m + 3*(483*A - 435*B + 419*C
)*c^2)*cos(f*x + e) + (128*(A + B + C)*c^2*m^2 + (16*C*c^2*m^4 + 128*C*c^2
*m^3 + 344*C*c^2*m^2 + 352*C*c^2*m + 105*C*c^2)*cos(f*x + e)^4 + 256*(4*A
+ B - 2*C)*c^2*m - (16*(B - 2*C)*c^2*m^4 + 16*(9*B - 22*C)*c^2*m^3 + 32*(1
3*B - 35*C)*c^2*m^2 + 4*(111*B - 314*C)*c^2*m + 15*(9*B - 26*C)*c^2)*cos(f
*x + e)^3 + 96*(21*A - 15*B + 13*C)*c^2 - (16*(A - B + C)*c^2*m^4 + 32*(5*
A - 7*B + 5*C)*c^2*m^3 + 8*(65*A - 117*B + 113*C)*c^2*m^2 + 24*(25*A - 51*
B + 57*C)*c^2*m + 9*(21*A - 45*B + 53*C)*c^2)*cos(f*x + e)^2 - 2*(16*(A -
B + C)*c^2*m^4 + 192*(A - B + C)*c^2*m^3 + 8*(99*A - 107*B + 99*C)*c^2*m^2
+ 16*(77*A - 97*B + 101*C)*c^2*m + 3*(147*A - 195*B + 211*C)*c^2)*cos(...
```

**3.18.6 Sympy [F(-1)]**

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(c-c*sin(f*x+e))**(5/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output `Timed out`

**3.18.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 1324 vs.  $2(403) = 806$ .

Time = 0.42 (sec) , antiderivative size = 1324, normalized size of antiderivative = 3.04

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output

```

-2*((4*m^2 + 24*m + 43)*a^m*c^(5/2) - (12*m^2 + 40*m - 15)*a^m*c^(5/2)*si
n(f*x + e)/(cos(f*x + e) + 1) + 2*(4*m^2 + 8*m + 35)*a^m*c^(5/2)*sin(f*x +
e)^2/(cos(f*x + e) + 1)^2 + 2*(4*m^2 + 8*m + 35)*a^m*c^(5/2)*sin(f*x + e)
^3/(cos(f*x + e) + 1)^3 - (12*m^2 + 40*m - 15)*a^m*c^(5/2)*sin(f*x + e)^4/
(cos(f*x + e) + 1)^4 + (4*m^2 + 24*m + 43)*a^m*c^(5/2)*sin(f*x + e)^5/(cos
(f*x + e) + 1)^5)*A*e^(2*m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1) - m*lo
g(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/((8*m^3 + 36*m^2 + 46*m + 15)*
(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1)^(5/2)) - 2*((4*m^2 + 40*m + 115)
*a^m*c^(5/2) - 2*(4*m^3 + 40*m^2 + 115*m)*a^m*c^(5/2)*sin(f*x + e)/(cos(f*
x + e) + 1) + 2*(12*m^3 + 76*m^2 + 97*m + 175)*a^m*c^(5/2)*sin(f*x + e)^2/
(cos(f*x + e) + 1)^2 - (16*m^3 + 76*m^2 + 260*m - 175)*a^m*c^(5/2)*sin(f*x
+ e)^3/(cos(f*x + e) + 1)^3 - (16*m^3 + 76*m^2 + 260*m - 175)*a^m*c^(5/2)
*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 2*(12*m^3 + 76*m^2 + 97*m + 175)*a^
m*c^(5/2)*sin(f*x + e)^5/(cos(f*x + e) + 1)^5 - 2*(4*m^3 + 40*m^2 + 115*m)
*a^m*c^(5/2)*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 + (4*m^2 + 40*m + 115)*a^
m*c^(5/2)*sin(f*x + e)^7/(cos(f*x + e) + 1)^7)*B*e^(2*m*log(sin(f*x + e)/(
cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/((
16*m^4 + 128*m^3 + 344*m^2 + 352*m + (16*m^4 + 128*m^3 + 344*m^2 + 352*m
+ 105)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 105)*(sin(f*x + e)^2/(cos(f*x
+ e) + 1)^2 + 1)^(5/2)) + 4*(2*(4*m^2 + 56*m + 219)*a^m*c^(5/2) - 4*(4...

```

### 3.18.8 Giac [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e)^2 + B \sin(fx + e) + A) (-c \sin(fx + e) + c)^{5/2} (a \sin(fx + e) + a)^m dx$$

input

```

integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+B*sin(f*x+e)+C*sin(
f*x+e)^2),x, algorithm="giac")

```

output

```

integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(-c*sin(f*x + e) + c)^(5
/2)*(a*sin(f*x + e) + a)^m, x)

```

### 3.18.9 Mupad [B] (verification not implemented)

Time = 23.24 (sec) , antiderivative size = 1253, normalized size of antiderivative = 2.88

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Too large to display}$$

```
input int((a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^(5/2)*(A + B*sin(e + f*x)
+ C*sin(e + f*x)^2),x)
```

```
output ((c - c*sin(e + f*x))^(1/2)*((C*c^2*(a + a*sin(e + f*x))^m*(m*352i + m^2*3
44i + m^3*128i + m^4*16i + 105i))/(8*f*(m*3378i + m^2*3800i + m^3*1840i +
m^4*400i + m^5*32i + 945i)) + (c^2*exp(e*5i + f*x*5i)*(a + a*sin(e + f*x))
^m*(18900*A - 14175*B + 12285*C + 15648*A*m - 4140*B*m + 648*C*m + 5280*A*
m^2 + 896*A*m^3 + 64*A*m^4 - 832*B*m^2 - 208*B*m^3 - 16*B*m^4 + 1416*C*m^2
+ 224*C*m^3 + 16*C*m^4))/(4*f*(m*3378i + m^2*3800i + m^3*1840i + m^4*400i
+ m^5*32i + 945i)) + (c^2*exp(e*4i + f*x*4i)*(a + a*sin(e + f*x))^m*(A*18
900i - B*14175i + C*12285i + A*m*15648i - B*m*4140i + C*m*648i + A*m^2*528
0i + A*m^3*896i + A*m^4*64i - B*m^2*832i - B*m^3*208i - B*m^4*16i + C*m^2*
1416i + C*m^3*224i + C*m^4*16i))/(4*f*(m*3378i + m^2*3800i + m^3*1840i + m
^4*400i + m^5*32i + 945i)) + (C*c^2*exp(e*9i + f*x*9i)*(a + a*sin(e + f*x)
)^m*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105))/(8*f*(m*3378i + m^2*3800i
+ m^3*1840i + m^4*400i + m^5*32i + 945i)) - (c^2*exp(e*7i + f*x*7i)*(a + a
*sin(e + f*x))^m*(8*m + 4*m^2 + 3)*(126*A - 315*B + 378*C + 64*A*m - 124*B
*m + 88*C*m + 8*A*m^2 - 12*B*m^2 + 8*C*m^2))/(4*f*(m*3378i + m^2*3800i + m
^3*1840i + m^4*400i + m^5*32i + 945i)) - (c^2*exp(e*2i + f*x*2i)*(a + a*si
n(e + f*x))^m*(8*m + 4*m^2 + 3)*(A*126i - B*315i + C*378i + A*m*64i - B*m*
124i + C*m*88i + A*m^2*8i - B*m^2*12i + C*m^2*8i))/(4*f*(m*3378i + m^2*380
0i + m^3*1840i + m^4*400i + m^5*32i + 945i)) + (c^2*exp(e*3i + f*x*3i)*(2*
m + 1)*(a + a*sin(e + f*x))^m*(3150*A - 3465*B + 3150*C + 2356*A*m - 17...
```

### 3.19 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^{3/2} (A + B \sin(e -$

3.19.1	Optimal result	181
3.19.2	Mathematica [A] (verified)	182
3.19.3	Rubi [A] (verified)	182
3.19.4	Maple [F]	185
3.19.5	Fricas [A] (verification not implemented)	186
3.19.6	Sympy [F(-1)]	186
3.19.7	Maxima [B] (verification not implemented)	187
3.19.8	Giac [F]	188
3.19.9	Mupad [B] (verification not implemented)	188

#### 3.19.1 Optimal result

Integrand size = 48, antiderivative size = 322

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx =$$

$$\frac{8c^2(B(21 - 8m - 4m^2) - C(19 - 8m + 4m^2) - A(35 + 24m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m}{f(5 + 2m)(7 + 2m)(3 + 8m + 4m^2) \sqrt{c - c \sin(e + fx)}} +$$

$$\frac{2c(B(21 - 8m - 4m^2) - C(19 - 8m + 4m^2) - A(35 + 24m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m \sqrt{c}}{f(3 + 2m)(5 + 2m)(7 + 2m)} -$$

$$\frac{2(7B + 2C + 2Bm + 4Cm) \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2}}{f(5 + 2m)(7 + 2m)} +$$

$$\frac{2C \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2}}{cf(7 + 2m)}$$

output

```
-2*(2*B*m+4*C*m+7*B+2*C)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)/f/(4*m^2+24*m+35)+2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)/c/f/(7+2*m)-8*c^2*(B*(-4*m^2-8*m+21)-C*(4*m^2-8*m+19)-A*(4*m^2+24*m+35))*cos(f*x+e)*(a+a*sin(f*x+e))^m/f/(7+2*m)/(8*m^3+36*m^2+46*m+15)/(c-c*sin(f*x+e))^(1/2)-2*c*(B*(-4*m^2-8*m+21)-C*(4*m^2-8*m+19)-A*(4*m^2+24*m+35))*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)/f/(8*m^3+60*m^2+142*m+105)
```

### 3.19.2 Mathematica [A] (verified)

Time = 6.44 (sec) , antiderivative size = 306, normalized size of antiderivative = 0.95

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \frac{c(\cos(\frac{1}{2}(e + fx)) + \sin(\frac{1}{2}(e + fx))) (a(1 + \sin(e + fx)))^m \sqrt{c - c \sin(e + fx)} (700A$$

input `Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `(c*(Cos[(e + f*x)/2] + Sin[(e + f*x)/2])*(a*(1 + Sin[e + f*x]))^m*sqrt[c - c*Sin[e + f*x]]*(700*A - 546*B + 494*C + 760*A*m - 380*B*m + 284*C*m + 272*A*m^2 - 120*B*m^2 + 136*C*m^2 + 32*A*m^3 - 16*B*m^3 + 16*C*m^3 + 2*(3 + 8*m + 4*m^2)*(B*(7 + 2*m) - C*(13 + 2*m))*Cos[2*(e + f*x)] - (1 + 2*m)*(4*A*(35 + 24*m + 4*m^2) - 4*B*(63 + 32*m + 4*m^2) + C*(253 + 80*m + 12*m^2))*Sin[e + f*x] + 15*C*Sin[3*(e + f*x)] + 46*C*m*Sin[3*(e + f*x)] + 36*C*m^2*Sin[3*(e + f*x)] + 8*C*m^3*Sin[3*(e + f*x)]))/(2*f*(1 + 2*m)*(3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2]))`

### 3.19.3 Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 279, normalized size of antiderivative = 0.87, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$ , Rules used = {3042, 3518, 27, 3042, 3452, 3042, 3219, 3042, 3217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c - c \sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

↓ 3042

$$\int (c - c \sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2) dx$$

↓ 3518

$$\frac{2C \cos(e+fx)(c-c\sin(e+fx))^{5/2}(a\sin(e+fx)+a)^m}{cf(2m+7)} - \frac{2 \int -\frac{1}{2}(\sin(e+fx)a+a)^m(c-c\sin(e+fx))^{3/2}(ac(C(5-2m)+A(2m+7))+ac(2mB+7B+2C+4Cm)\sin(e+fx))^{3/2} dx}{ac(2m+7)}$$

↓ 27

$$\frac{\int (\sin(e+fx)a+a)^m(c-c\sin(e+fx))^{3/2}(ac(C(5-2m)+A(2m+7))+ac(2mB+7B+2C+4Cm)\sin(e+fx))^{3/2} dx}{ac(2m+7)} - \frac{2C \cos(e+fx)(c-c\sin(e+fx))^{5/2}(a\sin(e+fx)+a)^m}{cf(2m+7)}$$

↓ 3042

$$\frac{\int (\sin(e+fx)a+a)^m(c-c\sin(e+fx))^{3/2}(ac(C(5-2m)+A(2m+7))+ac(2mB+7B+2C+4Cm)\sin(e+fx))^{3/2} dx}{ac(2m+7)} - \frac{2C \cos(e+fx)(c-c\sin(e+fx))^{5/2}(a\sin(e+fx)+a)^m}{cf(2m+7)}$$

↓ 3452

$$\frac{ac(-A(4m^2+24m+35)+B(-4m^2-8m+21)-C(4m^2-8m+19)) \int (\sin(e+fx)a+a)^m(c-c\sin(e+fx))^{3/2} dx}{2m+5} - \frac{2ac(2Bm+7B+4Cm+2C)\cos(e+fx)}{ac(2m+7)}$$

$$\frac{2C \cos(e+fx)(c-c\sin(e+fx))^{5/2}(a\sin(e+fx)+a)^m}{cf(2m+7)}$$

↓ 3042

$$\frac{ac(-A(4m^2+24m+35)+B(-4m^2-8m+21)-C(4m^2-8m+19)) \int (\sin(e+fx)a+a)^m(c-c\sin(e+fx))^{3/2} dx}{2m+5} - \frac{2ac(2Bm+7B+4Cm+2C)\cos(e+fx)}{ac(2m+7)}$$

$$\frac{2C \cos(e+fx)(c-c\sin(e+fx))^{5/2}(a\sin(e+fx)+a)^m}{cf(2m+7)}$$

↓ 3219

$$\frac{ac(-A(4m^2+24m+35)+B(-4m^2-8m+21)-C(4m^2-8m+19)) \left( \frac{4c \int (\sin(e+fx)a+a)^m \sqrt{c-c\sin(e+fx)} dx}{2m+3} + \frac{2c \cos(e+fx) \sqrt{c-c\sin(e+fx)} (a\sin(e+fx)+a)^m}{f(2m+3)} \right)}{2m+5} - \frac{2ac(2Bm+7B+4Cm+2C)\cos(e+fx)}{ac(2m+7)}$$

$$\frac{2C \cos(e+fx)(c-c\sin(e+fx))^{5/2}(a\sin(e+fx)+a)^m}{cf(2m+7)}$$

↓ 3042

---

3.19.  $\int (a+a\sin(e+fx))^m(c-c\sin(e+fx))^{3/2}(A+B\sin(e+fx)+C\sin^2(e+fx)) dx$



$$\frac{ac(-A(4m^2+24m+35)+B(-4m^2-8m+21)-C(4m^2-8m+19))\left(\frac{4c\int(\sin(e+fx)a+a)^m\sqrt{c-c\sin(e+fx)}dx}{2m+3} + \frac{2c\cos(e+fx)\sqrt{c-c\sin(e+fx)}(a\sin(e+fx)+a)^m}{f(2m+3)}\right)}{2m+5} \quad ac(2m+7)$$

$$\frac{2C\cos(e+fx)(c-c\sin(e+fx))^{5/2}(a\sin(e+fx)+a)^m}{cf(2m+7)}$$

↓ 3217

$$\frac{ac(-A(4m^2+24m+35)+B(-4m^2-8m+21)-C(4m^2-8m+19))\left(\frac{8c^2\cos(e+fx)(a\sin(e+fx)+a)^m}{f(2m+1)(2m+3)\sqrt{c-c\sin(e+fx)}} + \frac{2c\cos(e+fx)\sqrt{c-c\sin(e+fx)}(a\sin(e+fx)+a)^m}{f(2m+3)}\right)}{2m+5} \quad ac(2m+7)$$

$$\frac{2C\cos(e+fx)(c-c\sin(e+fx))^{5/2}(a\sin(e+fx)+a)^m}{cf(2m+7)}$$

input `Int[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x]`

output `(2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(5/2))/(c*f*(7 + 2*m)) + ((-2*a*c*(7*B + 2*C + 2*B*m + 4*C*m)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2))/(f*(5 + 2*m)) - (a*c*(B*(21 - 8*m - 4*m^2) - C*(19 - 8*m + 4*m^2) - A*(35 + 24*m + 4*m^2))*((8*c^2*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) + (2*c*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*Sqrt[c - c*Sin[e + f*x]])/(f*(3 + 2*m))))/(5 + 2*m))/(a*c*(7 + 2*m))`

### 3.19.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3217 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[n, -2^(-1)]`

---

3.19.  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$

rule 3219 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^n/(f*(m + n))), x] + Simp[a*((2*m - 1)/(m + n)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[m - 1/2, 0] && !LtQ[n, -1] && !(IGtQ[n - 1/2, 0] && LtQ[n, m]) && !(ILtQ[m + n, 0] && GtQ[2*m + n + 1, 0])`

rule 3452 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 1, 0]`

rule 3518 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + (b*B*d*(m + n + 2) - b*c*C*(2*m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]`

### 3.19.4 Maple [F]

$$\int (a + a \sin(fx + e))^m (c - c \sin(fx + e))^{\frac{3}{2}} (A + B \sin(fx + e) + C \sin^2(fx + e)) dx$$

input `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

output `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

### 3.19.5 Fracas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 563, normalized size of antiderivative = 1.75

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \frac{2((8Ccm^3 + 36Ccm^2 + 46Ccm + 15Cc) \cos(fx + e)^4 - 16(A + B + C)cm^2 - (8(B - C)cm^3 + 4(11$$

```
input integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fracas")
```

```
output -2*((8*C*c*m^3 + 36*C*c*m^2 + 46*C*c*m + 15*C*c)*cos(f*x + e)^4 - 16*(A + B + C)*c*m^2 - (8*(B - C)*c*m^3 + 4*(11*B - 17*C)*c*m^2 + 2*(31*B - 55*C)*c*m + 3*(7*B - 13*C)*c)*cos(f*x + e)^3 - 32*(3*A + B - C)*c*m - (8*(A + C)*c*m^3 + 4*(13*A - 6*B + 5*C)*c*m^2 + 2*(47*A - 48*B + 47*C)*c*m + (35*A - 42*B + 43*C)*c)*cos(f*x + e)^2 - 4*(35*A - 21*B + 19*C)*c - (8*(A - B + C)*c*m^3 + 4*(17*A - 13*B + 17*C)*c*m^2 + 2*(95*A - 63*B + 63*C)*c*m + (175*A - 147*B + 143*C)*c)*cos(f*x + e) - (16*(A + B + C)*c*m^2 + (8*C*c*m^3 + 36*C*c*m^2 + 46*C*c*m + 15*C*c)*cos(f*x + e)^3 + 32*(3*A + B - C)*c*m + (8*B*c*m^3 + 4*(11*B - 8*C)*c*m^2 + 2*(31*B - 32*C)*c*m + 3*(7*B - 8*C)*c)*cos(f*x + e)^2 + 4*(35*A - 21*B + 19*C)*c - (8*(A - B + C)*c*m^3 + 4*(13*A - 17*B + 13*C)*c*m^2 + 2*(47*A - 79*B + 79*C)*c*m + (35*A - 63*B + 67*C)*c)*cos(f*x + e))*sin(f*x + e)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(16*f*m^4 + 128*f*m^3 + 344*f*m^2 + 352*f*m + (16*f*m^4 + 128*f*m^3 + 344*f*m^2 + 352*f*m + 105*f)*cos(f*x + e) - (16*f*m^4 + 128*f*m^3 + 344*f*m^2 + 352*f*m + 105*f)*sin(f*x + e) + 105*f)
```

### 3.19.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Timed out}$$

```
input integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)
```

```
output Timed out
```

---

3.19.  $\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$

### 3.19.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 950 vs.  $2(302) = 604$ .

Time = 0.38 (sec) , antiderivative size = 950, normalized size of antiderivative = 2.95

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Too large to display}$$

```
input integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(
f*x+e)^2),x, algorithm="maxima")
```

```
output -2*((a^m*c^(3/2)*(2*m + 5) - a^m*c^(3/2)*(2*m - 3)*sin(f*x + e)/(cos(f*x +
e) + 1) - a^m*c^(3/2)*(2*m - 3)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + a^m
*c^(3/2)*(2*m + 5)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3)*A*e^(2*m*log(sin(f
*x + e)/(cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^
2 + 1))/(4*m^2 + 8*m + 3)*(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1)^(3/2)
) - 2*(a^m*c^(3/2)*(2*m + 9) - 2*(2*m^2 + 9*m)*a^m*c^(3/2)*sin(f*x + e)/(c
os(f*x + e) + 1) + (4*m^2 + 15)*a^m*c^(3/2)*sin(f*x + e)^2/(cos(f*x + e) +
1)^2 + (4*m^2 + 15)*a^m*c^(3/2)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 - 2*(
2*m^2 + 9*m)*a^m*c^(3/2)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + a^m*c^(3/2)
*(2*m + 9)*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)*B*e^(2*m*log(sin(f*x + e)/
(cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/
((8*m^3 + 36*m^2 + 46*m + (8*m^3 + 36*m^2 + 46*m + 15)*sin(f*x + e)^2/(cos
(f*x + e) + 1)^2 + 15)*(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1)^(3/2)) +
4*(2*a^m*c^(3/2)*(2*m + 13) - 4*(2*m^2 + 13*m)*a^m*c^(3/2)*sin(f*x + e)/(c
os(f*x + e) + 1) + (8*m^3 + 60*m^2 + 66*m + 91)*a^m*c^(3/2)*sin(f*x + e)^2
/(cos(f*x + e) + 1)^2 - (8*m^3 + 20*m^2 + 82*m - 35)*a^m*c^(3/2)*sin(f*x +
e)^3/(cos(f*x + e) + 1)^3 - (8*m^3 + 20*m^2 + 82*m - 35)*a^m*c^(3/2)*sin(
f*x + e)^4/(cos(f*x + e) + 1)^4 + (8*m^3 + 60*m^2 + 66*m + 91)*a^m*c^(3/2)
*sin(f*x + e)^5/(cos(f*x + e) + 1)^5 - 4*(2*m^2 + 13*m)*a^m*c^(3/2)*sin(f*
x + e)^6/(cos(f*x + e) + 1)^6 + 2*a^m*c^(3/2)*(2*m + 13)*sin(f*x + e)^7...
```

**3.19.8 Giac [F]**

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e) + B \sin(fx + e) + A) (-c \sin(fx + e) + c)^{3/2} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(-c*sin(f*x + e) + c)^(3/2)*(a*sin(f*x + e) + a)^m, x)`

**3.19.9 Mupad [B] (verification not implemented)**

Time = 24.21 (sec) , antiderivative size = 790, normalized size of antiderivative = 2.45

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \frac{\sqrt{c - c \sin(e + fx)}}{4 f (16 m^4 + 128 m^3 + 344 m^2 + 352 m + 105)} \left( \frac{C c (a + a \sin(e + fx))^m (m^3 8i + m^2 36i + m 46i + 15i)}{4 f (16 m^4 + 128 m^3 + 344 m^2 + 352 m + 105)} + \frac{c e^{3i + f x 3i} (a + a \sin(e + fx))}{4 f (16 m^4 + 128 m^3 + 344 m^2 + 352 m + 105)} \right)$$

input `int((a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^(3/2)*(A + B*sin(e + f*x) + C*sin(e + f*x)^2),x)`

output

```

((c - c*sin(e + f*x))^(1/2)*((C*c*(a + a*sin(e + f*x))^m*(m*46i + m^2*36i
+ m^3*8i + 15i))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) + (c*exp
(e*3i + f*x*3i)*(a + a*sin(e + f*x))^m*(1260*A - 840*B + 735*C + 1144*A*m
- 128*B*m - 18*C*m + 336*A*m^2 + 32*A*m^3 + 32*B*m^2 + 100*C*m^2 + 8*C*m^3
))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) - (c*exp(e*4i + f*x*4i
)*(a + a*sin(e + f*x))^m*(A*1260i - B*840i + C*735i + A*m*1144i - B*m*128i
- C*m*18i + A*m^2*336i + A*m^3*32i + B*m^2*32i + C*m^2*100i + C*m^3*8i))/
(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) + (c*exp(e*5i + f*x*5i)*(
2*m + 1)*(a + a*sin(e + f*x))^m*(140*A - 210*B + 175*C + 96*A*m - 88*B*m +
16*C*m + 16*A*m^2 - 8*B*m^2 + 4*C*m^2))/(4*f*(352*m + 344*m^2 + 128*m^3 +
16*m^4 + 105)) - (c*exp(e*2i + f*x*2i)*(2*m + 1)*(a + a*sin(e + f*x))^m*(
A*140i - B*210i + C*175i + A*m*96i - B*m*88i + C*m*16i + A*m^2*16i - B*m^2
*8i + C*m^2*4i))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) + (c*exp
(e*1i + f*x*1i)*(a + a*sin(e + f*x))^m*(8*m + 4*m^2 + 3)*(14*B - 21*C + 4*
B*m - 2*C*m))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) - (c*exp(e*
6i + f*x*6i)*(a + a*sin(e + f*x))^m*(8*m + 4*m^2 + 3)*(B*14i - C*21i + B*m
*4i - C*m*2i))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) - (C*c*exp
(e*7i + f*x*7i)*(a + a*sin(e + f*x))^m*(46*m + 36*m^2 + 8*m^3 + 15))/(4*f*
(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)))/(exp(e*4i + f*x*4i) - exp(e
*3i + f*x*3i)*(m*352i + m^2*344i + m^3*128i + m^4*16i + 105i))/(352*m + ...

```

### 3.20 $\int (a+a \sin(e+fx))^m \sqrt{c-c \sin(e+fx)}(A+B \sin(e-$

3.20.1	Optimal result . . . . .	190
3.20.2	Mathematica [A] (verified) . . . . .	191
3.20.3	Rubi [A] (verified) . . . . .	191
3.20.4	Maple [F] . . . . .	194
3.20.5	Fricas [A] (verification not implemented) . . . . .	194
3.20.6	Sympy [F] . . . . .	195
3.20.7	Maxima [B] (verification not implemented) . . . . .	195
3.20.8	Giac [F] . . . . .	196
3.20.9	Mupad [B] (verification not implemented) . . . . .	197

#### 3.20.1 Optimal result

Integrand size = 48, antiderivative size = 197

$$\int (a+a \sin(e+fx))^m \sqrt{c-c \sin(e+fx)}(A+B \sin(e+fx)+C \sin^2(e+fx)) dx$$

$$= \frac{2c(C-6Cm+A(5+2m)-B(5+2m)) \cos(e+fx)(a+a \sin(e+fx))^m}{f(1+2m)(5+2m)\sqrt{c-c \sin(e+fx)}} + \frac{2c(5B+2C+2Bm+4Cm) \cos(e+fx)(a+a \sin(e+fx))^{1+m}}{af(3+2m)(5+2m)\sqrt{c-c \sin(e+fx)}} + \frac{2C \cos(e+fx)(a+a \sin(e+fx))^m(c-c \sin(e+fx))^{3/2}}{cf(5+2m)}$$

output

```
2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)/c/f/(5+2*m)+2*c*(
C-6*C*m+A*(5+2*m)-B*(5+2*m))*cos(f*x+e)*(a+a*sin(f*x+e))^m/f/(4*m^2+12*m+5
)/(c-c*sin(f*x+e))^(1/2)+2*c*(2*B*m+4*C*m+5*B+2*C)*cos(f*x+e)*(a+a*sin(f*x
+e))^(1+m)/a/f/(4*m^2+16*m+15)/(c-c*sin(f*x+e))^(1/2)
```

### 3.20.2 Mathematica [A] (verified)

Time = 2.07 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.90

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \frac{(\cos(\frac{1}{2}(e + fx)) + \sin(\frac{1}{2}(e + fx))) (a(1 + \sin(e + fx)))^m \sqrt{c - c \sin(e + fx)} (30A - 20B + 19C + 32A \sin(e + fx) + C \sin^2(e + fx))}{f(1 + 2m)(3 + 2m)(5 + 2m)}$$

input `Integrate[(a + a*Sin[e + f*x])^m*Sqrt[c - c*Sin[e + f*x]]*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `((Cos[(e + f*x)/2] + Sin[(e + f*x)/2])*(a*(1 + Sin[e + f*x]))^m*Sqrt[c - c*Sin[e + f*x]]*(30*A - 20*B + 19*C + 32*A*m - 8*B*m + 8*C*m + 8*A*m^2 + 4*C*m^2 - C*(3 + 8*m + 4*m^2)*Cos[2*(e + f*x)] + 2*(1 + 2*m)*(5*B - 4*C + 2*B*m)*Sin[e + f*x]))/(f*(1 + 2*m)*(3 + 2*m)*(5 + 2*m)*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2]))`

### 3.20.3 Rubi [A] (verified)

Time = 1.00 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.02, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.146$ , Rules used = {3042, 3518, 27, 3042, 3450, 3042, 3217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{c - c \sin(e + fx)} (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{c - c \sin(e + fx)} (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2) dx$$

$$\downarrow \text{3518}$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{cf(2m + 5)} -$$

$$\frac{2 \int -\frac{1}{2}(\sin(e + fx)a + a)^m \sqrt{c - c \sin(e + fx)} (ac(C(3 - 2m) + A(2m + 5)) + ac(2mB + 5B + 2C + 4Cm) \sin(e + fx)) dx}{ac(2m + 5)}$$

$$\downarrow \text{27}$$

---

3.20.  $\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$



$$\frac{\int (\sin(e + fx)a + a)^m \sqrt{c - c\sin(e + fx)} (ac(C(3 - 2m) + A(2m + 5)) + ac(2mB + 5B + 2C + 4Cm) \sin(e + fx) + a) dx}{\frac{ac(2m + 5)}{cf(2m + 5)} \frac{2C \cos(e + fx)(c - c\sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{cf(2m + 5)}}$$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^m \sqrt{c - c\sin(e + fx)} (ac(C(3 - 2m) + A(2m + 5)) + ac(2mB + 5B + 2C + 4Cm) \sin(e + fx) + a) dx}{\frac{ac(2m + 5)}{cf(2m + 5)} \frac{2C \cos(e + fx)(c - c\sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{cf(2m + 5)}}$$

↓ 3450

$$\frac{ac(A(2m + 5) - B(2m + 5) - 6Cm + C) \int (\sin(e + fx)a + a)^m \sqrt{c - c\sin(e + fx)} dx + c(2Bm + 5B + 4Cm + 2) \int (\sin(e + fx)a + a)^m \sqrt{c - c\sin(e + fx)} dx}{\frac{ac(2m + 5)}{cf(2m + 5)} \frac{2C \cos(e + fx)(c - c\sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{cf(2m + 5)}}$$

↓ 3042

$$\frac{ac(A(2m + 5) - B(2m + 5) - 6Cm + C) \int (\sin(e + fx)a + a)^m \sqrt{c - c\sin(e + fx)} dx + c(2Bm + 5B + 4Cm + 2) \int (\sin(e + fx)a + a)^m \sqrt{c - c\sin(e + fx)} dx}{\frac{ac(2m + 5)}{cf(2m + 5)} \frac{2C \cos(e + fx)(c - c\sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{cf(2m + 5)}}$$

↓ 3217

$$\frac{\frac{2ac^2(A(2m+5)-B(2m+5)-6Cm+C) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}} + \frac{2c^2(2Bm+5B+4Cm+2C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}}{f(2m+3)\sqrt{c-c\sin(e+fx)}}}{\frac{ac(2m + 5)}{cf(2m + 5)} \frac{2C \cos(e + fx)(c - c\sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{cf(2m + 5)}}$$

input `Int[(a + a*Sin[e + f*x])^m*Sqrt[c - c*Sin[e + f*x]]*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `(2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2))/(c*f*(5 + 2*m)) + ((2*a*c^2*(C - 6*C*m + A*(5 + 2*m) - B*(5 + 2*m))*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) + (2*c^2*(5*B + 2*C + 2*B*m + 4*C*m)*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m))/(f*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]])/(a*c*(5 + 2*m))`

---

3.20.  $\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$

## 3.20.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3217 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[n, -2^(-1)]`
- rule 3450 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[B/d Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] - Simp[(B*c - A*d)/d Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]`
- rule 3518 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + (b*B*d*(m + n + 2) - b*c*C*(2*m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]`

### 3.20.4 Maple [F]

$$\int (a + a \sin(fx + e))^m \sqrt{c - c \sin(fx + e)} (A + B \sin(fx + e) + C(\sin^2(fx + e))) dx$$

input `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

output `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

### 3.20.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 310, normalized size of antiderivative = 1.57

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx =$$

$$\frac{2((4Cm^2 + 8Cm + 3C) \cos(fx + e)^3 - 4(A + B + C)m^2 + (4(B + C)m^2 + 12Bm + 5B - C) \cos$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fracas")`

output `-2*((4*C*m^2 + 8*C*m + 3*C)*cos(f*x + e)^3 - 4*(A + B + C)*m^2 + (4*(B + C)*m^2 + 12*B*m + 5*B - C)*cos(f*x + e)^2 - 8*(2*A + B)*m - (4*(A + C)*m^2 + 4*(4*A - B + 2*C)*m + 15*A - 10*B + 11*C)*cos(f*x + e) - (4*(A + B + C)*m^2 - (4*C*m^2 + 8*C*m + 3*C)*cos(f*x + e)^2 + 8*(2*A + B)*m + (4*B*m^2 + 4*(3*B - 2*C)*m + 5*B - 4*C)*cos(f*x + e) + 15*A - 5*B + 7*C)*sin(f*x + e) - 15*A + 5*B - 7*C)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(8*f*m^3 + 36*f*m^2 + 46*f*m + (8*f*m^3 + 36*f*m^2 + 46*f*m + 15*f)*cos(f*x + e) - (8*f*m^3 + 36*f*m^2 + 46*f*m + 15*f)*sin(f*x + e) + 15*f)`

### 3.20.6 Sympy [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a(\sin(e + fx) + 1))^m \sqrt{-c(\sin(e + fx) - 1)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input `integrate((a+a*sin(f*x+e))**m*(c-c*sin(f*x+e))**(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output `Integral((a*(sin(e + f*x) + 1))**m*sqrt(-c*(sin(e + f*x) - 1))*(A + B*sin(e + f*x) + C*sin(e + f*x)**2), x)`

### 3.20.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 644 vs.  $2(187) = 374$ .

Time = 0.38 (sec) , antiderivative size = 644, normalized size of antiderivative = 3.27

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx =$$

$$2 \left( \frac{2 \left( \frac{2 a^m \sqrt{c m} \sin(fx+e)}{\cos(fx+e)+1} + \frac{2 a^m \sqrt{c m} \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - a^m \sqrt{c} - \frac{a^m \sqrt{c} \sin(fx+e)^3}{(\cos(fx+e)+1)^3} \right) Be^{\left( 2 m \log\left( \frac{\sin(fx+e)}{\cos(fx+e)+1} + 1 \right) - m \log\left( \frac{\sin(fx+e)^2}{(\cos(fx+e)+1)^2} + 1 \right) \right)}}{\left( 4 m^2 + 8 m + \frac{(4 m^2 + 8 m + 3) \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + 3 \right) \sqrt{\frac{\sin(fx+e)^2}{(\cos(fx+e)+1)^2} + 1}} \right) -$$

input `integrate((a+a*sin(f*x+e))~m*(c-c*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output

```

-2*(2*(2*a^m*sqrt(c)*m*sin(f*x + e)/(cos(f*x + e) + 1) + 2*a^m*sqrt(c)*m*
sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - a^m*sqrt(c) - a^m*sqrt(c)*sin(f*x + e
)^3/(cos(f*x + e) + 1)^3)*B*e^(2*m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1
) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/((4*m^2 + 8*m + (4*m^2
+ 8*m + 3)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 3)*sqrt(sin(f*x + e)^2/(
cos(f*x + e) + 1)^2 + 1)) - 4*(4*a^m*sqrt(c)*m*sin(f*x + e)/(cos(f*x + e)
+ 1) - (4*m^2 + 4*m + 5)*a^m*sqrt(c)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 -
(4*m^2 + 4*m + 5)*a^m*sqrt(c)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 4*a^m
*sqrt(c)*m*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 2*a^m*sqrt(c) - 2*a^m*sqrt
(c)*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)*C*e^(2*m*log(sin(f*x + e)/(cos(f
*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/((8*m^
3 + 36*m^2 + 46*m + 2*(8*m^3 + 36*m^2 + 46*m + 15)*sin(f*x + e)^2/(cos(f*x
+ e) + 1)^2 + (8*m^3 + 36*m^2 + 46*m + 15)*sin(f*x + e)^4/(cos(f*x + e) +
1)^4 + 15)*sqrt(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1)) + (a^m*sqrt(c)
+ a^m*sqrt(c)*sin(f*x + e)/(cos(f*x + e) + 1))*A*e^(2*m*log(sin(f*x + e)/(
cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/
(2*m + 1)*sqrt(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/f

```

### 3.20.8 Giac [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A) \sqrt{-c \sin(fx + e) + c(a \sin(fx + e) + a)^m} dx$$

input

```

integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(
f*x+e)^2),x, algorithm="giac")

```

output

```

integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*sqrt(-c*sin(f*x + e) + c
)*(a*sin(f*x + e) + a)^m, x)

```

**3.20.9 Mupad [B] (verification not implemented)**

Time = 19.68 (sec) , antiderivative size = 510, normalized size of antiderivative = 2.59

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx =$$

$$\frac{\sqrt{c - c \sin(e + fx)} \left( -\frac{e^{e^{3i+fx^{3i}} (a+a \sin(e+fx))^m (30A-15B+15C+32Am+4Bm+8Am^2+4Bm^2+4Cm^2)}}{f(m^3 8i+m^2 36i+m 46i+15i)} - \frac{e^{e^{2i+fx^{2i}}}}{f(m^3 8i+m^2 36i+m 46i+15i)} \right)}{f(m^3 8i+m^2 36i+m 46i+15i)}$$

```
input int((a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^(1/2)*(A + B*sin(e + f*x)
+ C*sin(e + f*x)^2),x)
```

```
output -((c - c*sin(e + f*x))^(1/2)*((C*(a + a*sin(e + f*x))^m*(m*8i + m^2*4i + 3
i))/(2*f*(m*46i + m^2*36i + m^3*8i + 15i)) - (exp(e*2i + f*x*2i)*(a + a*si
n(e + f*x))^m*(A*30i - B*15i + C*15i + A*m*32i + B*m*4i + A*m^2*8i + B*m^2
*4i + C*m^2*4i))/(f*(m*46i + m^2*36i + m^3*8i + 15i)) - (exp(e*3i + f*x*3i
)*(a + a*sin(e + f*x))^m*(30*A - 15*B + 15*C + 32*A*m + 4*B*m + 8*A*m^2 +
4*B*m^2 + 4*C*m^2))/(f*(m*46i + m^2*36i + m^3*8i + 15i)) + (C*exp(e*5i + f
*x*5i)*(a + a*sin(e + f*x))^m*(8*m + 4*m^2 + 3))/(2*f*(m*46i + m^2*36i + m
^3*8i + 15i)) + (exp(e*1i + f*x*1i)*(2*m + 1)*(a + a*sin(e + f*x))^m*(10*B
- 5*C + 4*B*m + 2*C*m))/(2*f*(m*46i + m^2*36i + m^3*8i + 15i)) + (exp(e*4
i + f*x*4i)*(2*m + 1)*(a + a*sin(e + f*x))^m*(B*10i - C*5i + B*m*4i + C*m*
2i))/(2*f*(m*46i + m^2*36i + m^3*8i + 15i))))/(exp(e*3i + f*x*3i) + (exp(e
*2i + f*x*2i)*(46*m + 36*m^2 + 8*m^3 + 15))/(m*46i + m^2*36i + m^3*8i + 15
i))
```

**3.21** 
$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{\sqrt{c-c \sin(e+fx)}} dx$$

3.21.1	Optimal result	198
3.21.2	Mathematica [A] (verified)	199
3.21.3	Rubi [A] (verified)	199
3.21.4	Maple [F]	202
3.21.5	Fricas [F]	202
3.21.6	Sympy [F]	203
3.21.7	Maxima [F]	203
3.21.8	Giac [F(-1)]	204
3.21.9	Mupad [F(-1)]	204

**3.21.1 Optimal result**

Integrand size = 48, antiderivative size = 170

$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{\sqrt{c-c \sin(e+fx)}} dx$$

$$= -\frac{2B \cos(e+fx)(a+a \sin(e+fx))^m}{f(1+2m)\sqrt{c-c \sin(e+fx)}} + \frac{(A+B+C) \cos(e+fx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}+m, \frac{3}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right) (a+a \sin(e+fx))}{f(1+2m)\sqrt{c-c \sin(e+fx)}} - \frac{2C \cos(e+fx)(a+a \sin(e+fx))^{1+m}}{af(3+2m)\sqrt{c-c \sin(e+fx)}}$$

output

```
-2*B*cos(f*x+e)*(a+a*sin(f*x+e))^m/f/(1+2*m)/(c-c*sin(f*x+e))^(1/2)+(A+B+C)
*cos(f*x+e)*hypergeom([1, 1/2+m],[3/2+m],1/2+1/2*sin(f*x+e))*(a+a*sin(f*x
+e))^m/f/(1+2*m)/(c-c*sin(f*x+e))^(1/2)-2*C*cos(f*x+e)*(a+a*sin(f*x+e))^(1
+m)/a/f/(3+2*m)/(c-c*sin(f*x+e))^(1/2)
```

### 3.21.2 Mathematica [A] (verified)

Time = 31.18 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.94

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= \frac{\cos(e + fx)(a(1 + \sin(e + fx)))^m (2(A + C)(3 + 2m) \text{Hypergeometric2F1}(1, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e + fx)))}{\dots}$$

input `Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/Sqrt[c - c*Sin[e + f*x]],x]`

output `(Cos[e + f*x]*(a*(1 + Sin[e + f*x]))^m*(2*(A + C)*(3 + 2*m)*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2] + B*(1 + 2*m)*Hypergeometric2F1[1, 3/2 + m, 5/2 + m, (1 + Sin[e + f*x])/2]*(1 + Sin[e + f*x]) - 2*(3*B + 2*C + 2*B*m + 4*C*m + 2*C*(1 + 2*m)*Sin[e + f*x]))/(2*f*(1 + 2*m)*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]])`

### 3.21.3 Rubi [A] (verified)

Time = 0.93 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.03, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$ , Rules used = {3042, 3516, 3042, 3452, 3042, 3224, 3042, 3146, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

↓ 3042

$$\int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2)}{\sqrt{c - c \sin(e + fx)}} dx$$

↓ 3516

$$\int \frac{(\sin(e + fx)a + a)^m (A + C + B \sin(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx - \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^{m+1}}{af(2m + 3)\sqrt{c - c \sin(e + fx)}}$$

↓ 3042

---

3.21.  $\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$



$$\begin{aligned}
& \int \frac{(\sin(e+fx)a+a)^m(A+C+B\sin(e+fx))}{\sqrt{c-c\sin(e+fx)}} dx - \frac{2C\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}} \\
& \quad \downarrow \text{3452} \\
& (A+B+C) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx - \frac{2B\cos(e+fx)(a\sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}} - \\
& \quad \frac{2C\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& (A+B+C) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx - \frac{2B\cos(e+fx)(a\sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}} - \\
& \quad \frac{2C\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}} \\
& \quad \downarrow \text{3224} \\
& \frac{(A+B+C)\cos(e+fx) \int \sec(e+fx)(\sin(e+fx)a+a)^{m+\frac{1}{2}} dx}{\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} - \\
& \frac{2B\cos(e+fx)(a\sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}} - \frac{2C\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{(A+B+C)\cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}}{\cos(e+fx)} dx}{\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} - \frac{2B\cos(e+fx)(a\sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}} - \\
& \quad \frac{2C\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}} \\
& \quad \downarrow \text{3146} \\
& \frac{a(A+B+C)\cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{a-a\sin(e+fx)} d(a\sin(e+fx))}{f\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} - \\
& \frac{2B\cos(e+fx)(a\sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}} - \frac{2C\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}} \\
& \quad \downarrow \text{78} \\
& \frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^m \operatorname{Hypergeometric2F1}\left(1, m+\frac{1}{2}, m+\frac{3}{2}, \frac{\sin(e+fx)a+a}{2a}\right)}{f(2m+1)\sqrt{c-c\sin(e+fx)}} - \\
& \frac{2B\cos(e+fx)(a\sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}} - \frac{2C\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}}
\end{aligned}$$

---

3.21.  $\int \frac{(a+a\sin(e+fx))^m(A+B\sin(e+fx)+C\sin^2(e+fx))}{\sqrt{c-c\sin(e+fx)}} dx$

input `Int[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/Sqrt[c - c*Sin[e + f*x]],x]`

output `(-2*B*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) + ((A + B + C)*Cos[e + f*x]*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (a + a*Sin[e + f*x])/(2*a)]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) - (2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m))/(a*f*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]])`

### 3.21.3.1 Defintions of rubi rules used

rule 78 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3146 `Int[cos[(e_) + (f_)*(x_)]^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1)/2], x], x, b*Sin[e + f*x], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])`

rule 3224 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*Sin[e + f*x])^FracPart[m]*((c + d*Sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*Sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])`

rule 3452 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 1, 0]`

rule 3516 `Int[(((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)])^2)/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*C*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(2*m + 3)*Sqrt[c + d*Sin[e + f*x]]), x] + Int[(a + b*Sin[e + f*x])^m*(Simp[A + C + B*Sin[e + f*x], x]/Sqrt[c + d*Sin[e + f*x]]), x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

### 3.21.4 Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + B \sin(fx + e) + C \sin^2(fx + e))}{\sqrt{c - c \sin(fx + e)}} dx$$

input `int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x)`

output `int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x)`

### 3.21.5 Fracas [F]

$$\begin{aligned} & \int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx \\ &= \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m}{\sqrt{-c \sin(fx + e) + c}} dx \end{aligned}$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x, algorithm="fracas")`

---

3.21.  $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{\sqrt{c-c \sin(e+fx)}} dx$

output `integral((C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(c*sin(f*x + e) - c), x)`

### 3.21.6 Sympy [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= \int \frac{(a(\sin(e + fx) + 1))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{-c(\sin(e + fx) - 1)}} dx$$

input `integrate((a+a*sin(f*x+e))**m*(A+B*sin(f*x+e)+C*sin(f*x+e)**2)/(c-c*sin(f*x+e))**(1/2),x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + B*sin(e + f*x) + C*sin(e + f*x)**2)/sqrt(-c*(sin(e + f*x) - 1)), x)`

### 3.21.7 Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m}{\sqrt{-c \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/sqrt(-c*sin(f*x + e) + c), x)`

**3.21.8 Giac [F(-1)]**

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x, algorithm="giac")`

output `Timed out`

**3.21.9 Mupad [F(-1)]**

Timed out.

$$\begin{aligned} & \int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx \\ &= \int \frac{(a + a \sin(e + fx))^m (C \sin(e + fx)^2 + B \sin(e + fx) + A)}{\sqrt{c - c \sin(e + fx)}} dx \end{aligned}$$

input `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c - c*sin(e + f*x))^(1/2),x)`

output `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c - c*sin(e + f*x))^(1/2), x)`

$$3.22 \quad \int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{3/2}} dx$$

3.22.1	Optimal result . . . . .	205
3.22.2	Mathematica [A] (verified) . . . . .	205
3.22.3	Rubi [A] (verified) . . . . .	206
3.22.4	Maple [F] . . . . .	209
3.22.5	Fricas [F] . . . . .	210
3.22.6	Sympy [F] . . . . .	210
3.22.7	Maxima [F] . . . . .	210
3.22.8	Giac [F(-2)] . . . . .	211
3.22.9	Mupad [F(-1)] . . . . .	211

### 3.22.1 Optimal result

Integrand size = 48, antiderivative size = 216

$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{3/2}} dx = \frac{(A+B+C) \cos(e+fx)(a+a \sin(e+fx))}{4af(c-c \sin(e+fx))^{3/2}} + \frac{(A+B+2Am+2Bm+C(9+2m)) \cos(e+fx)(a+a \sin(e+fx))^m}{4cf(1+2m)\sqrt{c-c \sin(e+fx)}} + \frac{(A(1-2m)-B(3+2m)-C(7+2m)) \cos(e+fx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}+m, \frac{3}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right)}{4cf(1+2m)\sqrt{c-c \sin(e+fx)}}$$

output

```
1/4*(A+B+C)*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)/a/f/(c-c*sin(f*x+e))^(3/2)+1/4*(A+B+2*A*m+2*B*m+C*(9+2*m))*cos(f*x+e)*(a+a*sin(f*x+e))^m/c/f/(1+2*m)/(c-c*sin(f*x+e))^(1/2)+1/4*(A*(1-2*m)-B*(3+2*m)-C*(7+2*m))*cos(f*x+e)*hypergeom([1, 1/2+m], [3/2+m], 1/2+1/2*sin(f*x+e))*(a+a*sin(f*x+e))^m/c/f/(1+2*m)/(c-c*sin(f*x+e))^(1/2)
```

### 3.22.2 Mathematica [A] (verified)

Time = 34.09 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.75

$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{3/2}} dx = \frac{\cos(e+fx)(a(1+\sin(e+fx)))^m ((8C+B(3+2m)) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}+m, \frac{3}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right))}{(c-c \sin(e+fx))^{3/2}}$$

---

3.22.  $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{3/2}} dx$

input `Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2)) / (c - c*Sin[e + f*x])^(3/2), x]`

output `-1/4*(Cos[e + f*x]*(a*(1 + Sin[e + f*x]))^m*((8*C + B*(3 + 2*m))*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2]*(-1 + Sin[e + f*x]) + 2*(B + 4*C + 2*B*m - (A + C)*Hypergeometric2F1[2, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2]*(-1 + Sin[e + f*x]) - 4*C*Sin[e + f*x]))/(c*f*(1 + 2*m)*( -1 + Sin[e + f*x])*Sqrt[c - c*Sin[e + f*x]])`

### 3.22.3 Rubi [A] (verified)

Time = 1.17 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.06, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$ , Rules used = {3042, 3514, 27, 3042, 3452, 3042, 3224, 3042, 3146, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2)}{(c - c \sin(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{3514} \\
 & \frac{(A + B + C) \cos(e + fx) (a \sin(e + fx) + a)^{m+1}}{4af(c - c \sin(e + fx))^{3/2}} - \\
 & \frac{\int - \frac{(\sin(e + fx)a + a)^m (a^2(A(3 - 2m) - (B + C)(2m + 5)) - a^2(2mA + A + B + 2Bm + C(2m + 9)) \sin(e + fx))}{2\sqrt{c - c \sin(e + fx)}} dx}{4a^2c} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{(\sin(e + fx)a + a)^m (a^2(A(3 - 2m) - (B + C)(2m + 5)) - a^2(2mA + A + B + 2Bm + C(2m + 9)) \sin(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx}{8a^2c} + \\
 & \frac{(A + B + C) \cos(e + fx) (a \sin(e + fx) + a)^{m+1}}{4af(c - c \sin(e + fx))^{3/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.22.  $\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx$

$$\begin{aligned}
& \frac{\int \frac{(\sin(e+fx)a+a)^m (a^2(A(3-2m)-(B+C)(2m+5))-a^2(2mA+A+B+2Bm+C(2m+9)) \sin(e+fx))}{\sqrt{c-c\sin(e+fx)}} dx}{\frac{8a^2c}{(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} \frac{1}{4af(c-c\sin(e+fx))^{3/2}}} + \\
& \quad \downarrow \text{3452} \\
& \frac{2a^2(A(1-2m)-B(2m+3)-C(2m+7)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx + \frac{2a^2(2Am+A+2Bm+B+C(2m+9)) \cos(e+fx)(a \sin(e+fx))}{f(2m+1)\sqrt{c-c\sin(e+fx)}}}{\frac{8a^2c}{(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} \frac{1}{4af(c-c\sin(e+fx))^{3/2}}} \\
& \quad \downarrow \text{3042} \\
& \frac{2a^2(A(1-2m)-B(2m+3)-C(2m+7)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx + \frac{2a^2(2Am+A+2Bm+B+C(2m+9)) \cos(e+fx)(a \sin(e+fx))}{f(2m+1)\sqrt{c-c\sin(e+fx)}}}{\frac{8a^2c}{(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} \frac{1}{4af(c-c\sin(e+fx))^{3/2}}} \\
& \quad \downarrow \text{3224} \\
& \frac{2a^2(A(1-2m)-B(2m+3)-C(2m+7)) \cos(e+fx) \int \frac{\sec(e+fx)(\sin(e+fx)a+a)^{m+\frac{1}{2}}}{\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} dx + \frac{2a^2(2Am+A+2Bm+B+C(2m+9)) \cos(e+fx)(a \sin(e+fx))}{f(2m+1)\sqrt{c-c\sin(e+fx)}}}{\frac{8a^2c}{(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} \frac{1}{4af(c-c\sin(e+fx))^{3/2}}} \\
& \quad \downarrow \text{3042} \\
& \frac{2a^2(A(1-2m)-B(2m+3)-C(2m+7)) \cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}}{\cos(e+fx)\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} dx + \frac{2a^2(2Am+A+2Bm+B+C(2m+9)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}}}{\frac{8a^2c}{(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} \frac{1}{4af(c-c\sin(e+fx))^{3/2}}} \\
& \quad \downarrow \text{3146} \\
& \frac{2a^3(A(1-2m)-B(2m+3)-C(2m+7)) \cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{a-a\sin(e+fx)} d(a \sin(e+fx)) + \frac{2a^2(2Am+A+2Bm+B+C(2m+9)) \cos(e+fx)(a \sin(e+fx))}{f(2m+1)\sqrt{c-c\sin(e+fx)}}}{\frac{8a^2c}{(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} \frac{1}{4af(c-c\sin(e+fx))^{3/2}}} \\
& \quad \downarrow \text{78} \\
& \frac{3.22. \int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{3/2}} dx}{\frac{8a^2c}{(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}} \frac{1}{4af(c-c\sin(e+fx))^{3/2}}}
\end{aligned}$$



$$\frac{2a^2(A(1-2m)-B(2m+3)-C(2m+7)) \cos(e+fx)(a \sin(e+fx)+a)^m \operatorname{Hypergeometric2F1}\left(1, m+\frac{1}{2}, m+\frac{3}{2}, \frac{\sin(e+fx)a+a}{2a}\right)}{f(2m+1)\sqrt{c-c\sin(e+fx)}} + \frac{2a^2(2Am+A+2Bm+3C)}{f} \\ \frac{(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}}{4af(c-c\sin(e+fx))^{3/2}}$$

input `Int[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/(c - c*Sin[e + f*x])^(3/2), x]`

output `((A + B + C)*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m))/(4*a*f*(c - c*Sin[e + f*x])^(3/2)) + ((2*a^2*(A + B + 2*A*m + 2*B*m + C*(9 + 2*m))*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) + (2*a^2*(A*(1 - 2*m) - B*(3 + 2*m) - C*(7 + 2*m))*Cos[e + f*x]*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (a + a*Sin[e + f*x])/(2*a)]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]])/(8*a^2*c)`

### 3.22.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 78 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3146 `Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1/2), x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])`

```
rule 3224 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*Sin[e + f*x])^FracPart[m]*((c + d*Sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*Sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])
```

```
rule 3452 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 1, 0]
```

```
rule 3514 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(a*A - b*B + a*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(2*b*c*f*(2*m + 1))), x] - Simp[1/(2*b*c*d*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[A*(c^2*(m + 1) + d^2*(2*m + n + 2)) - B*c*d*(m - n - 1) - C*(c^2*m - d^2*(n + 1)) + d*((A*c + B*d)*(m + n + 2) - c*C*(3*m - n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (EqQ[m + n + 2, 0] && NeQ[2*m + 1, 0]))
```

### 3.22.4 Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + B \sin(fx + e) + C(\sin^2(fx + e)))}{(c - c \sin(fx + e))^{\frac{3}{2}}} dx$$

```
input int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x)
```

```
output int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x)
```

---

3.22.  $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{\frac{3}{2}}} dx$

### 3.22.5 Fricas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin(fx + e))^2 + B \sin(fx + e) + A + c}{(-c \sin(fx + e) - c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x, algorithm="fricas")`

output `integral((C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(c^2*cos(f*x + e)^2 + 2*c^2*sin(f*x + e) - 2*c^2), x)`

### 3.22.6 Sympy [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(a(\sin(e + fx) + 1))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(-c(\sin(e + fx) - 1))^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))**m*(A+B*sin(f*x+e)+C*sin(f*x+e)**2)/(c-c*sin(f*x+e))**(3/2),x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + B*sin(e + f*x) + C*sin(e + f*x)**2)/(-c*(sin(e + f*x) - 1))**(3/2), x)`

### 3.22.7 Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin(fx + e))^2 + B \sin(fx + e) + A + c}{(-c \sin(fx + e) - c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/(-c*sin(f*x + e) + c)^(3/2), x)`

---

3.22.  $\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx$

**3.22.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:Unable to divide, perhaps due to rounding error%%{1,[0,1,1,1,0,0,0,0,0,0]}+%%{1,[0,0,1,1,1,0,0,0,0,0]}+%%{1,[`

**3.22.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(a + a \sin(e + fx))^m (C \sin(e + fx) + \dots)}{(c - c \sin(e + fx))^{3/2}}$$

input `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c - c*sin(e + f*x))^(3/2),x)`

output `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c - c*sin(e + f*x))^(3/2), x)`

$$3.23 \quad \int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx$$

3.23.1	Optimal result	212
3.23.2	Mathematica [A] (verified)	212
3.23.3	Rubi [A] (verified)	213
3.23.4	Maple [F]	216
3.23.5	Fricas [F]	217
3.23.6	Sympy [F(-1)]	217
3.23.7	Maxima [F]	217
3.23.8	Giac [F(-2)]	218
3.23.9	Mupad [F(-1)]	218

### 3.23.1 Optimal result

Integrand size = 48, antiderivative size = 230

$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx = \frac{(A+B+C) \cos(e+fx)(a+a \sin(e+fx))}{8af(c-c \sin(e+fx))^{5/2}} + \frac{(A(5-2m)-B(3+2m)-C(11+2m)) \cos(e+fx)(a+a \sin(e+fx))^m}{16cf(c-c \sin(e+fx))^{3/2}} - \frac{(B(5-8m-4m^2)-A(3-8m+4m^2)-C(19+24m+4m^2)) \cos(e+fx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2} + \dots\right)}{32c^2 f(1+2m)\sqrt{c-c \sin(e+fx)}}$$

output

```
1/8*(A+B+C)*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)/a/f/(c-c*sin(f*x+e))^(5/2)+1/16*(A*(5-2*m)-B*(3+2*m)-C*(11+2*m))*cos(f*x+e)*(a+a*sin(f*x+e))^m/c/f/(c-c*sin(f*x+e))^(3/2)-1/32*(B*(-4*m^2-8*m+5)-A*(4*m^2-8*m+3)-C*(4*m^2+24*m+19))*cos(f*x+e)*hypergeom([1, 1/2+m], [3/2+m], 1/2+1/2*sin(f*x+e))*(a+a*sin(f*x+e))^m/c^2/f/(1+2*m)/(c-c*sin(f*x+e))^(1/2)
```

### 3.23.2 Mathematica [A] (verified)

Time = 29.42 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.99

$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx = \frac{\cos(e+fx)(a(1+\sin(e+fx)))^m (8B \dots)}{\dots}$$

input `Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2)) / (c - c*Sin[e + f*x])^(5/2), x]`

output `(Cos[e + f*x]*(a*(1 + Sin[e + f*x]))^m*(8*B*(1 + 2*m) - 2*(16*C + B*(5 + 2*m))*Hypergeometric2F1[2, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2]*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])^4 + 8*(A + C)*Hypergeometric2F1[3, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2]*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])^4 - 16*C*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2]*(-3 + Cos[2*(e + f*x)] + 4*Sin[e + f*x]))/(32*c^2*(f + 2*f*m)*(-1 + Sin[e + f*x])^2*Sqrt[c - c*Sin[e + f*x]])`

### 3.23.3 Rubi [A] (verified)

Time = 1.25 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.07, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$ , Rules used = {3042, 3514, 27, 3042, 3451, 3042, 3224, 3042, 3146, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2)}{(c - c \sin(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{3514} \\
 & \frac{(A + B + C) \cos(e + fx) (a \sin(e + fx) + a)^{m+1}}{8af(c - c \sin(e + fx))^{5/2}} - \\
 & \frac{\int \frac{(\sin(e+fx)a+a)^m ((A(9-2m)-(B+C)(2m+7))a^2 + ((A+B)(1-2m)-C(2m+15)) \sin(e+fx)a^2)}{2(c-c \sin(e+fx))^{3/2}} dx}{8a^2c} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{(\sin(e+fx)a+a)^m ((A(9-2m)-(B+C)(2m+7))a^2 + ((A+B)(1-2m)-C(2m+15)) \sin(e+fx)a^2)}{(c-c \sin(e+fx))^{3/2}} dx}{\frac{16a^2c}{(A + B + C) \cos(e + fx) (a \sin(e + fx) + a)^{m+1}}} + \\
 & \frac{(A + B + C) \cos(e + fx) (a \sin(e + fx) + a)^{m+1}}{8af(c - c \sin(e + fx))^{5/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.23.  $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx$

$$\frac{\int \frac{(\sin(e+fx)a+a)^m ((A(9-2m)-(B+C)(2m+7))a^2 + ((A+B)(1-2m)-C(2m+15)) \sin(e+fx)a^2)}{(c-c\sin(e+fx))^{3/2}} dx}{\frac{16a^2c}{8af(c-c\sin(e+fx))^{5/2}} (A+B+C) \cos(e+fx)(a \sin(e+fx) + a)^{m+1}} +$$

↓ 3451

$$\frac{a^2(A(5-2m)-B(2m+3)-C(2m+11)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} - \frac{a^2(-A(4m^2-8m+3)+B(-4m^2-8m+5)-C(4m^2+24m+19)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx}{2c}$$

$$\frac{(A+B+C) \cos(e+fx)(a \sin(e+fx) + a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}} \frac{16a^2c}{16a^2c}$$

↓ 3042

$$\frac{a^2(A(5-2m)-B(2m+3)-C(2m+11)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} - \frac{a^2(-A(4m^2-8m+3)+B(-4m^2-8m+5)-C(4m^2+24m+19)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx}{2c}$$

$$\frac{(A+B+C) \cos(e+fx)(a \sin(e+fx) + a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}} \frac{16a^2c}{16a^2c}$$

↓ 3224

$$\frac{a^2(A(5-2m)-B(2m+3)-C(2m+11)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} - \frac{a^2(-A(4m^2-8m+3)+B(-4m^2-8m+5)-C(4m^2+24m+19)) \cos(e+fx)}{2c\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}}$$

$$\frac{(A+B+C) \cos(e+fx)(a \sin(e+fx) + a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}} \frac{16a^2c}{16a^2c}$$

↓ 3042

$$\frac{a^2(A(5-2m)-B(2m+3)-C(2m+11)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} - \frac{a^2(-A(4m^2-8m+3)+B(-4m^2-8m+5)-C(4m^2+24m+19)) \cos(e+fx)}{2c\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}}$$

$$\frac{(A+B+C) \cos(e+fx)(a \sin(e+fx) + a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}} \frac{16a^2c}{16a^2c}$$

↓ 3146

$$\frac{a^2(A(5-2m)-B(2m+3)-C(2m+11)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} - \frac{a^3(-A(4m^2-8m+3)+B(-4m^2-8m+5)-C(4m^2+24m+19)) \cos(e+fx)}{2cf\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}}$$

$$\frac{(A+B+C) \cos(e+fx)(a \sin(e+fx) + a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}} \frac{16a^2c}{16a^2c}$$

↓ 78

---

3.23.  $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c\sin(e+fx))^{5/2}} dx$

$$\frac{a^2(A(5-2m)-B(2m+3)-C(2m+11)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} - \frac{a^2(-A(4m^2-8m+3)+B(-4m^2-8m+5)-C(4m^2+24m+19)) \cos(e+fx)}{2cf(2m+1)\sqrt{c-c\sin(e+fx)}}$$

$$\frac{(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}} \quad 16a^2c$$

input `Int[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/(c - c*Sin[e + f*x])^(5/2),x]`

output `((A + B + C)*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m))/(8*a*f*(c - c*Sin[e + f*x])^(5/2)) + ((a^2*(A*(5 - 2*m) - B*(3 + 2*m) - C*(11 + 2*m))*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(c - c*Sin[e + f*x])^(3/2)) - (a^2*(B*(5 - 8*m - 4*m^2) - A*(3 - 8*m + 4*m^2) - C*(19 + 24*m + 4*m^2))*Cos[e + f*x]*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (a + a*Sin[e + f*x])/(2*a)]*(a + a*Sin[e + f*x])^m)/(2*c*f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]))/(16*a^2*c)`

### 3.23.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 78 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3146 `Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1/2), x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])`

---

3.23.  $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx$



```
rule 3224 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*Sin[e + f*x])^FracPart[m]*((c + d*Sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*Sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])
```

```
rule 3451 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(a*B*(m - n) + A*b*(m + n + 1))/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (ILtQ[m + n, 0] && !SumSimplerQ[n, 1])) && NeQ[2*m + 1, 0]
```

```
rule 3514 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(a*A - b*B + a*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(2*b*c*f*(2*m + 1))), x] - Simp[1/(2*b*c*d*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[A*(c^2*(m + 1) + d^2*(2*m + n + 2)) - B*c*d*(m - n - 1) - C*(c^2*m - d^2*(n + 1)) + d*((A*c + B*d)*(m + n + 2) - c*C*(3*m - n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (EqQ[m + n + 2, 0] && NeQ[2*m + 1, 0]))
```

### 3.23.4 Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + B \sin(fx + e) + C(\sin^2(fx + e)))}{(c - c \sin(fx + e))^{\frac{5}{2}}} dx$$

```
input int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x)
```

```
output int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x)
```

---

3.23.  $\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{\frac{5}{2}}} dx$

### 3.23.5 Fracas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin(fx + e))^2 + B \sin(fx + e) + A}{(-c \sin(fx + e) + c)^{5/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x, algorithm="fracas")`

output `integral((C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(3*c^3*cos(f*x + e)^2 - 4*c^3 - (c^3*cos(f*x + e))^2 - 4*c^3)*sin(f*x + e)), x)`

### 3.23.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(A+B*sin(f*x+e)+C*sin(f*x+e)**2)/(c-c*sin(f*x+e))**(5/2),x)`

output `Timed out`

### 3.23.7 Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin(fx + e))^2 + B \sin(fx + e) + A}{(-c \sin(fx + e) + c)^{5/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/(-c*sin(f*x + e) + c)^(5/2), x)`

---

3.23.  $\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx$

**3.23.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \text{Exception raised: TypeError}$$

```
input integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Error index.cc index_gcd Error: Bad Argument Value
```

**3.23.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \int \frac{(a + a \sin(e + fx))^m (C \sin(e + fx) + A + B \sin(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx$$

```
input int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c - c*sin(e + f*x))^(5/2),x)
```

```
output int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c - c*sin(e + f*x))^(5/2), x)
```

### 3.24 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^{-2-m} (A + B \sin(e+fx) + C \sin^2(e+fx)) dx$

3.24.1	Optimal result	219
3.24.2	Mathematica [A] (verified)	220
3.24.3	Rubi [A] (verified)	220
3.24.4	Maple [F]	224
3.24.5	Fricas [F]	224
3.24.6	Sympy [F]	224
3.24.7	Maxima [F]	225
3.24.8	Giac [F]	225
3.24.9	Mupad [F(-1)]	225

#### 3.24.1 Optimal result

Integrand size = 50, antiderivative size = 232

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx =$$

$$\frac{2^{-\frac{1}{2}-m} C \cos^3(e + fx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}(3 + 2m), \frac{1}{2}(3 + 2m), \frac{1}{2}(5 + 2m), \frac{1}{2}(1 + \sin(e + fx))\right) (1 - \sin(e + fx))}{f(3 + 2m)}$$

$$+ \frac{(A + B + C) \cos(e + fx) (a + a \sin(e + fx))^{1+m} (c - c \sin(e + fx))^{-2-m}}{2af(3 + 2m)}$$

$$+ \frac{(A - B + C) \cos(e + fx) (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-1-m}}{2cf(1 + 2m)}$$

output

```
-2^(-1/2-m)*C*cos(f*x+e)^3*hypergeom([3/2+m, 3/2+m],[5/2+m],1/2+1/2*sin(f*x+e))*(1-sin(f*x+e))^(1/2+m)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(-2-m)/f/(3+2*m)+1/2*(A+B+C)*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)*(c-c*sin(f*x+e))^(-2-m)/a/f/(3+2*m)+1/2*(A-B+C)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(-1-m)/c/f/(1+2*m)
```

### 3.24.2 Mathematica [A] (verified)

Time = 1.16 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.84

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx =$$

$$\frac{\sec(e + fx)(1 + \sin(e + fx))^{-m} (a(1 + \sin(e + fx)))^m (c - c \sin(e + fx))^{-m} \left( 2^{\frac{3}{2}+m} C(3 + 2m) \text{Hypergeometric2F1}[-\frac{1}{2} - m, -\frac{1}{2} - m, \frac{1}{2} - m, (1 - \sin(e + fx))/2] * (-1 + \sin(e + fx)) * \sqrt{1 + \sin(e + fx)} + (1 + \sin(e + fx))^{(1 + m)} * (2A - B + 2C + 2A*m + 2C*m - (A + C - 2B*(1 + m)) * \sin(e + fx)) \right)}{c^2 * f * (1 + 2m) * (3 + 2m) * (-1 + \sin(e + fx)) * (1 + \sin(e + fx))^{-m} * (c - c * \sin(e + fx))^{-m}}$$

input `Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(-2 - m)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `-((Sec[e + f*x]*(a*(1 + Sin[e + f*x]))^m*(2^(3/2 + m)*C*(3 + 2*m)*Hypergeometric2F1[-1/2 - m, -1/2 - m, 1/2 - m, (1 - Sin[e + f*x])/2]*(-1 + Sin[e + f*x])*Sqrt[1 + Sin[e + f*x]] + (1 + Sin[e + f*x])^(1 + m)*(2*A - B + 2*C + 2*A*m + 2*C*m - (A + C - 2*B*(1 + m))*Sin[e + f*x]))/(c^2*f*(1 + 2*m)*(3 + 2*m)*(-1 + Sin[e + f*x])*(1 + Sin[e + f*x])^m*(c - c*Sin[e + f*x])^m)`

### 3.24.3 Rubi [A] (verified)

Time = 1.28 (sec) , antiderivative size = 299, normalized size of antiderivative = 1.29, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {3042, 3514, 3042, 3451, 3042, 3224, 3042, 3168, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-2} (A + B \sin(e + fx) + C \sin(e + fx)^2) dx$$

$$\downarrow \text{3514}$$

$$\frac{\int (\sin(e + fx)a + a)^{m+1} (c - c \sin(e + fx))^{-m-2} ((A + B - C)(2m + 1)c^2 + 2C(2m + 1) \sin(e + fx)c^2) dx}{2ac^2(2m + 1)} +$$

$$\frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

3.24.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^{m+1} (c - c \sin(e + fx))^{-m-2} ((A + B - C)(2m + 1)c^2 + 2C(2m + 1) \sin(e + fx)c^2) dx}{2ac^2(2m + 1)} +$$

$$\frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

↓ 3451

$$\frac{\frac{c^2(2m+1)(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}(c-c \sin(e+fx))^{-m-2}}{f(2m+3)} - 2cC(2m+1) \int (\sin(e+fx)a + a)^{m+1}(c - c \sin(e+fx))^{-m-2} dx}{2ac^2(2m+1)}$$

$$\frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

↓ 3042

$$\frac{\frac{c^2(2m+1)(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}(c-c \sin(e+fx))^{-m-2}}{f(2m+3)} - 2cC(2m+1) \int (\sin(e+fx)a + a)^{m+1}(c - c \sin(e+fx))^{-m-2} dx}{2ac^2(2m+1)}$$

$$\frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

↓ 3224

$$\frac{\frac{c^2(2m+1)(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}(c-c \sin(e+fx))^{-m-2}}{f(2m+3)} - 2ac^2C(2m+1) \cos^{-2m}(e+fx)(a \sin(e+fx)+a)^m}{2ac^2(2m+1)}$$

$$\frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

↓ 3042

$$\frac{\frac{c^2(2m+1)(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}(c-c \sin(e+fx))^{-m-2}}{f(2m+3)} - 2ac^2C(2m+1) \cos^{-2m}(e+fx)(a \sin(e+fx)+a)^m}{2ac^2(2m+1)}$$

$$\frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

↓ 3168

$$\frac{\frac{c^2(2m+1)(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}(c-c \sin(e+fx))^{-m-2}}{f(2m+3)} - \frac{2ac^4C(2m+1) \cos^3(e+fx)(a \sin(e+fx)+a)^m (c-c \sin(e+fx))^{\frac{1}{2}}}{2ac^2(2m+1)}}{2ac^2(2m+1)}$$

$$\frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

↓ 80

3.24.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$\frac{c^2(2m+1)(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}(c-c \sin(e+fx))^{-m-2}}{f(2m+3)} - \frac{ac^3C2^{-m-\frac{1}{2}}(2m+1) \cos^3(e+fx)(1-\sin(e+fx))^{m+\frac{1}{2}}(a \sin(e+fx))^{m+1}}{2ac^2(2m+1)}$$


---


$$\frac{(A-B+C) \cos(e+fx)(a \sin(e+fx)+a)^m(c-c \sin(e+fx))^{-m-1}}{2cf(2m+1)}$$

↓ 79

---


$$\frac{c^2(2m+1)(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}(c-c \sin(e+fx))^{-m-2}}{f(2m+3)} - \frac{ac^2C2^{\frac{1}{2}-m}(2m+1) \cos^3(e+fx)(1-\sin(e+fx))^{m+\frac{1}{2}}(a \sin(e+fx))^{m+1}}{2ac^2(2m+1)}$$


---


$$\frac{(A-B+C) \cos(e+fx)(a \sin(e+fx)+a)^m(c-c \sin(e+fx))^{-m-1}}{2cf(2m+1)}$$

input `Int[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(-2 - m)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `((A - B + C)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(-1 - m))/(2*c*f*(1 + 2*m)) + ((c^2*(A + B + C)*(1 + 2*m)*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m)*(c - c*Sin[e + f*x])^(-2 - m))/(f*(3 + 2*m)) - (2^(1/2 - m)*a*c^2*C*(1 + 2*m)*Cos[e + f*x]^3*Hypergeometric2F1[(3 + 2*m)/2, (3 + 2*m)/2, (5 + 2*m)/2, (1 + Sin[e + f*x])/2]*(1 - Sin[e + f*x])^(1/2 + m)*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(-1/2 + (-3 - 2*m)/2)*(c + c*Sin[e + f*x])^((-3 - 2*m)/2 + (3 + 2*m)/2))/(f*(3 + 2*m)))/(2*a*c^2*(1 + 2*m))`

### 3.24.3.1 Defintions of rubi rules used

rule 79 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))`

rule 80 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n, x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3168 `Int[(cos[(e_.) + (f_.)*(x_)]*(g_.))^(p_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Simp[a^2*((g*cos[e + f*x])^(p + 1)/(f*g*(a + b*sin[e + f*x])^((p + 1)/2)*(a - b*sin[e + f*x])^((p + 1)/2))) Subst[Int[(a + b*x)^(m + (p - 1)/2)*(a - b*x)^((p - 1)/2), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, e, f, g, m, p}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[m]`

rule 3224 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*sin[e + f*x])^FracPart[m]*((c + d*sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])`

rule 3451 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_ + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*sin[e + f*x])^m*((c + d*sin[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(a*B*(m - n) + A*b*(m + n + 1))/(a*b*(2*m + 1)) Int[(a + b*sin[e + f*x])^(m + 1)*(c + d*sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (ILtQ[m + n, 0] && !SumSimplerQ[n, 1])) && NeQ[2*m + 1, 0]`

rule 3514 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(a*A - b*B + a*C)*Cos[e + f*x]*(a + b*sin[e + f*x])^m*((c + d*sin[e + f*x])^(n + 1)/(2*b*c*f*(2*m + 1))), x] - Simp[1/(2*b*c*d*(2*m + 1)) Int[(a + b*sin[e + f*x])^(m + 1)*(c + d*sin[e + f*x])^n*Simp[A*(c^2*(m + 1) + d^2*(2*m + n + 2)) - B*c*d*(m - n - 1) - C*(c^2*m - d^2*(n + 1)) + d*((A*c + B*d)*(m + n + 2) - c*C*(3*m - n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (EqQ[m + n + 2, 0] && NeQ[2*m + 1, 0]))`



**3.24.4 Maple [F]**

$$\int (a + a \sin(fx + e))^m (c - c \sin(fx + e))^{-2-m} (A + B \sin(fx + e) + C(\sin^2(fx + e))) dx$$

input `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(2+m)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

output `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(2+m)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

**3.24.5 Fricas [F]**

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx \\ &= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^{-m-2} dx \end{aligned}$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(2+m)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*(a*sin(f*x + e) + a)^m*(-c*sin(f*x + e) + c)^(-m - 2), x)`

**3.24.6 Sympy [F]**

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx \\ &= \int (a(\sin(e + fx) + 1))^m (-c(\sin(e + fx) - 1))^{-m-2} (A + B \sin(e + fx) \\ & \qquad \qquad \qquad + C \sin^2(e + fx)) dx \end{aligned}$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(2+m)*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output `Integral((a*(sin(e + f*x) + 1))^m*(-c*(sin(e + f*x) - 1))^(2+m)*(A + B*sin(e + f*x) + C*sin(e + f*x)**2), x)`

**3.24.7 Maxima [F]**

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^{-m-2} dx$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m*(-c*sin(f*x + e) + c)^(-m - 2), x)`

**3.24.8 Giac [F]**

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^{-m-2} dx$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m*(-c*sin(f*x + e) + c)^(-m - 2), x)`

**3.24.9 Mupad [F(-1)]**

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int \frac{(a + a \sin(e + fx))^m (C \sin(e + fx)^2 + B \sin(e + fx) + A)}{(c - c \sin(e + fx))^{m+2}} dx$$

input `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c - c*sin(e + f*x))^(m + 2),x)`

output `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c - c*sin(e + f*x))^(m + 2), x)`

### 3.25 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^n (A + B \sin(e +$

3.25.1	Optimal result	227
3.25.2	Mathematica [F]	228
3.25.3	Rubi [A] (verified)	228
3.25.4	Maple [F]	232
3.25.5	Fricas [F]	233
3.25.6	Sympy [F(-1)]	233
3.25.7	Maxima [F]	233
3.25.8	Giac [F]	234
3.25.9	Mupad [F(-1)]	234

#### 3.25.1 Optimal result

Integrand size = 45, antiderivative size = 383

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= -\frac{C \cos(e + fx)(a + a \sin(e + fx))^m (c + d \sin(e + fx))^{1+n}}{df(2 + m + n)}$$

$$+ \frac{\sqrt{2}(c(C + 2Cm) + d(C(1 - m + n) + A(2 + m + n) - B(2 + m + n))) \operatorname{AppellF1}\left(\frac{1}{2} + m, \frac{1}{2}, -n, \frac{3}{2} + m\right)}{df(1 + 2m)(2 + m)}$$

$$- \frac{\sqrt{2}(cC(1 + m) - d(Cm + B(2 + m + n))) \operatorname{AppellF1}\left(\frac{3}{2} + m, \frac{1}{2}, -n, \frac{5}{2} + m, \frac{1}{2}(1 + \sin(e + fx)), -\frac{d(1 + \sin(e + fx))}{c + d \sin(e + fx)}\right)}{adf(3 + 2m)(2 + m + n)\sqrt{1 - \sin^2(e + fx)}}$$

output

```
-C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1+n)/d/f/(2+m+n)+(c*(2*C*m+C)+d*(C*(1-m+n)+A*(2+m+n)-B*(2+m+n)))*AppellF1(1/2+m,-n,1/2,3/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*2^(1/2)/d/f/(1+2*m)/(2+m+n)/(((c+d*sin(f*x+e))/(c-d))^n)/(1-sin(f*x+e))^(1/2)-(c*C*(1+m)-d*(C*m+B*(2+m+n)))*AppellF1(3/2+m,-n,1/2,5/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)*(c+d*sin(f*x+e))^n*2^(1/2)/a/d/f/(3+2*m)/(2+m+n)/(((c+d*sin(f*x+e))/(c-d))^n)/(1-sin(f*x+e))^(1/2)
```

### 3.25.2 Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input `Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x]`

### 3.25.3 Rubi [A] (verified)

Time = 1.17 (sec) , antiderivative size = 410, normalized size of antiderivative = 1.07, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {3042, 3524, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx)) (c + d \sin(e + fx))^n dx$$

$$\downarrow \text{3042}$$

$$\int (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2) (c + d \sin(e + fx))^n dx$$

$$\downarrow \text{3524}$$

$$\int \frac{(\sin(e + fx)a + a)^m (c + d \sin(e + fx))^n (a(Ad(m + n + 2) + C(nd + d + cm)) + a(Cdm - cC(m + 1) + Bd(m + n + 2)))}{df(m + n + 2)}$$

$$\downarrow \text{3042}$$

---

3.25.  $\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$

$$\frac{\int (\sin(e+fx)a+a)^m (c+d\sin(e+fx))^n (a(Ad(m+n+2)+C(nd+d+cm))+a(Cdm-cC(m+1)+Bd(m+n+2)))}{C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d\sin(e+fx))^{n+1}} \frac{ad(m+n+2)}{df(m+n+2)}$$

↓ 3466

$$\frac{(Bd(m+n+2)-cC(m+1)+Cdm) \int (\sin(e+fx)a+a)^{m+1} (c+d\sin(e+fx))^n dx - a(-Ad(m+n+2)+Bd(m+n+2))}{C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d\sin(e+fx))^{n+1}} \frac{ad(m+n+2)}{df(m+n+2)}$$

↓ 3042

$$\frac{(Bd(m+n+2)-cC(m+1)+Cdm) \int (\sin(e+fx)a+a)^{m+1} (c+d\sin(e+fx))^n dx - a(-Ad(m+n+2)+Bd(m+n+2))}{C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d\sin(e+fx))^{n+1}} \frac{ad(m+n+2)}{df(m+n+2)}$$

↓ 3267

$$\frac{a^2 \cos(e+fx)(Bd(m+n+2)-cC(m+1)+Cdm) \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}} (c+d\sin(e+fx))^n}{\sqrt{a-a\sin(e+fx)}} d\sin(e+fx)}{f \sqrt{a-a\sin(e+fx)} \sqrt{a\sin(e+fx)+a}} - \frac{a^3 \cos(e+fx)(-Ad(m+n+2)+Bd(m+n+2))}{ad(m+n+2)}$$

↓ 157

$$\frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx)(Bd(m+n+2)-cC(m+1)+Cdm) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m+\frac{1}{2}} (c+d\sin(e+fx))^n}{\sqrt{1-\sin(e+fx)}} d\sin(e+fx)}{\sqrt{2}f(a-a\sin(e+fx)) \sqrt{a\sin(e+fx)+a}} - \frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)}{ad(m+n+2)}$$

↓ 27

$$\frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx)(Bd(m+n+2)-cC(m+1)+Cdm) \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}} (c+d\sin(e+fx))^n}{\sqrt{1-\sin(e+fx)}} d\sin(e+fx)}{f(a-a\sin(e+fx)) \sqrt{a\sin(e+fx)+a}} - \frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)}{ad(m+n+2)}$$

↓ 156

---

3.25.  $\int (a+a\sin(e+fx))^m (c+d\sin(e+fx))^n (A+B\sin(e+fx)+C\sin^2(e+fx)) dx$

$$\frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (Bd(m+n+2) - cC(m+1) + Cdm) (c+d \sin(e+fx))^n \left(\frac{c+d \sin(e+fx)}{c-d}\right)^{-n} \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}} \left(\frac{c}{c-d} + \frac{d \sin(e+fx)}{c-d}\right)^n}{\sqrt{1-\sin(e+fx)}} dx}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}}$$

$$\frac{C \cos(e+fx) (a \sin(e+fx) + a)^m (c+d \sin(e+fx))^{n+1}}{df(m+n+2)}$$

↓ 155

$$\frac{\sqrt{2a} \sqrt{1-\sin(e+fx)} \cos(e+fx) (a \sin(e+fx)+a)^{m+1} (Bd(m+n+2) - cC(m+1) + Cdm) (c+d \sin(e+fx))^n \left(\frac{c+d \sin(e+fx)}{c-d}\right)^{-n} \text{AppellF1}\left(m+\frac{3}{2}, \frac{1}{2}, -\right)}{f(2m+3)(a-a \sin(e+fx))}$$

$$\frac{C \cos(e+fx) (a \sin(e+fx) + a)^m (c+d \sin(e+fx))^{n+1}}{df(m+n+2)}$$

input `Int[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `-((C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(1 + n))/(d*f*(2 + m + n))) + (-((Sqrt[2]*a^2*(C*d*m - c*C*(1 + m) - A*d*(2 + m + n) + B*d*(2 + m + n) - C*(d + c*m + d*n))*AppellF1[1/2 + m, 1/2, -n, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n)/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*((c + d*Sin[e + f*x])/(c - d))^n) + (Sqrt[2]*a*(C*d*m - c*C*(1 + m) + B*d*(2 + m + n))*AppellF1[3/2 + m, 1/2, -n, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*(c + d*Sin[e + f*x])^n)/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*((c + d*Sin[e + f*x])/(c - d))^n))/(a*d*(2 + m + n))`

---

3.25.  $\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$

## 3.25.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



rule 3267 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]])*Sqrt[a - b*Sin[e + f*x]]) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]`

rule 3466 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]`

rule 3524 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !Lt Q[m, -2^(-1)] && NeQ[m + n + 2, 0]`

### 3.25.4 Maple [F]

$$\int (a + a \sin(fx + e))^m (c + d \sin(fx + e))^n (A + B \sin(fx + e) + C \sin^2(fx + e)) dx$$

input `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2), x)`

output `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2), x)`

**3.25.5 Fracas [F]**

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^n dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e) + c)^n, x)`

**3.25.6 Sympy [F(-1)]**

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(c+d*sin(f*x+e))**n*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output `Timed out`

**3.25.7 Maxima [F]**

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^n dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e) + c)^n, x)`

**3.25.8 Giac [F]**

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A) (a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^n dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e) + c)^n, x)`

**3.25.9 Mupad [F(-1)]**

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (C \sin(e + fx)^2 + B \sin(e + fx) + A) dx$$

input `int((a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^n*(A + B*sin(e + f*x) + C*sin(e + f*x)^2),x)`

output `int((a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^n*(A + B*sin(e + f*x) + C*sin(e + f*x)^2), x)`

### 3.26 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{-2-m} (A+B \sin(e+fx) + C \sin^2(e+fx)) dx$

3.26.1	Optimal result	235
3.26.2	Mathematica [F]	236
3.26.3	Rubi [A] (verified)	236
3.26.4	Maple [F]	241
3.26.5	Fricas [F]	241
3.26.6	Sympy [F(-1)]	242
3.26.7	Maxima [F]	242
3.26.8	Giac [F]	242
3.26.9	Mupad [F(-1)]	243

#### 3.26.1 Optimal result

Integrand size = 49, antiderivative size = 410

$$\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{-2-m} (A+B \sin(e+fx) + C \sin^2(e+fx)) dx$$

$$= \frac{(c^2C - Bcd + Ad^2) \cos(e+fx)(a+a \sin(e+fx))^m (c+d \sin(e+fx))^{-1-m}}{d(c^2 - d^2) f(1+m)}$$

$$- \frac{2^{\frac{1}{2}+m} a (cd(A+C+Am+Bm+Cm) - c^2(C+2Cm) - d^2(Am+B(1+m) - C(1+m))) \cos(e+fx)}{a(c-d)df(3+2m)\sqrt{1-\sin(e+fx)}}$$

$$+ \frac{\sqrt{2}C \operatorname{AppellF1}\left(\frac{3}{2}+m, \frac{1}{2}, 1+m, \frac{5}{2}+m, \frac{1}{2}(1+\sin(e+fx)), -\frac{d(1+\sin(e+fx))}{c-d}\right) \cos(e+fx)(a+a \sin(e+fx))}{a(c-d)df(3+2m)\sqrt{1-\sin(e+fx)}}$$

output

```
(A*d^2-B*c*d+C*c^2)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1-m)/
d/(c^2-d^2)/f/(1+m)-2^(1/2+m)*a*(c*d*(A*m+B*m+C*m+A+C)-c^2*(2*C*m+C)-d^2*(
A*m+B*(1+m)-C*(1+m)))*cos(f*x+e)*hypergeom([1/2, 1/2-m],[3/2],1/2*(c-d)*(1
-sin(f*x+e))/(c+d*sin(f*x+e)))*(a+a*sin(f*x+e))^(1-m)*((c+d)*(1+sin(f*x+e
)))/(c+d*sin(f*x+e))^(1/2-m)/(c-d)/d/(c+d)^2/f/(1+m)/((c+d*sin(f*x+e))^m)+
C*AppellF1(3/2+m,1+m,1/2,5/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))
*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)*((c+d*sin(f*x+e))/(c-d))^m*2^(1/2)/a/(c
-d)/d/f/(3+2*m)/((c+d*sin(f*x+e))^m)/(1-sin(f*x+e))^(1/2)
```

### 3.26.2 Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input `Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(-2 - m)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(-2 - m)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x]`

### 3.26.3 Rubi [A] (verified)

Time = 1.33 (sec) , antiderivative size = 441, normalized size of antiderivative = 1.08, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.245$ , Rules used = {3042, 3522, 25, 3042, 3466, 3042, 3267, 142, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx)) (c + d \sin(e + fx))^{-m-2} dx$$

$$\downarrow \text{3042}$$

$$\int (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2) (c + d \sin(e + fx))^{-m-2} dx$$

$$\downarrow \text{3522}$$

$$\frac{\cos(e + fx) (Ad^2 - Bcd + c^2C) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}}{df(m+1)(c^2 - d^2)} -$$

$$\frac{\int -(\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1} (a(Ad(mc + c - dm) + (cC - Bd)(md + d - cm)) + aC(c^2 - d^2))}{ad(m+1)(c^2 - d^2)} dx$$

$$\downarrow \text{25}$$

$$\frac{\int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1} (a(Ad(mc + c - dm) + (cC - Bd)(md + d - cm)) + aC(c^2 - d^2))}{ad(m+1)(c^2 - d^2)} dx$$

$$\frac{\cos(e + fx) (Ad^2 - Bcd + c^2C) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}}{df(m+1)(c^2 - d^2)}$$

3.26.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1} (a(Ad(mc + c - dm) + (cC - Bd)(md + d - cm)) + aC(c^2 - d^2))}{\cos(e + fx) (Ad^2 - Bcd + c^2C) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}} \frac{ad(m+1)(c^2 - d^2)}{df(m+1)(c^2 - d^2)}$$

↓ 3466

$$\frac{a(cd(Am + A + Bm + Cm + C) - d^2(Am + B(m+1) - C(m+1)) - (c^2(2Cm + C))) \int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1}}{\cos(e + fx) (Ad^2 - Bcd + c^2C) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}} \frac{ad(m+1)(c^2 - d^2)}{df(m+1)(c^2 - d^2)}$$

↓ 3042

$$\frac{a(cd(Am + A + Bm + Cm + C) - d^2(Am + B(m+1) - C(m+1)) - (c^2(2Cm + C))) \int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1}}{\cos(e + fx) (Ad^2 - Bcd + c^2C) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}} \frac{ad(m+1)(c^2 - d^2)}{df(m+1)(c^2 - d^2)}$$

↓ 3267

$$\frac{a^3 \cos(e+fx)(cd(Am+A+Bm+Cm+C)-d^2(Am+B(m+1)-C(m+1))-(c^2(2Cm+C))) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}(c+d \sin(e+fx))^{-m-1}}{\sqrt{a-a \sin(e+fx)}} d \sin(e+fx)}{f \sqrt{a-a \sin(e+fx)} \sqrt{a \sin(e+fx)+a}} \frac{ad(m+1)(c^2 - d^2)}{\cos(e + fx) (Ad^2 - Bcd + c^2C) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}} \frac{ad(m+1)(c^2 - d^2)}{df(m+1)(c^2 - d^2)}$$

↓ 142

$$\frac{a^2 C(m+1)(c^2 - d^2) \cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}(c+d \sin(e+fx))^{-m-1}}{\sqrt{a-a \sin(e+fx)}} d \sin(e+fx)}{f \sqrt{a-a \sin(e+fx)} \sqrt{a \sin(e+fx)+a}} - \frac{a^2 2^{m+\frac{1}{2}} \cos(e+fx)(a \sin(e+fx)+a)^{m-1} (cd(Am+A+Bm+Cm+C) - d^2(Am+B(m+1) - C(m+1)) - (c^2(2Cm+C)))}{ad(m+1)(c^2 - d^2)} \frac{ad(m+1)(c^2 - d^2)}{\cos(e + fx) (Ad^2 - Bcd + c^2C) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}} \frac{ad(m+1)(c^2 - d^2)}{df(m+1)(c^2 - d^2)}$$

↓ 157

3.26.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$\frac{a^2 C(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)\int\frac{\sqrt{2}(\sin(e+fx)a+a)^{m+\frac{1}{2}}(c+d\sin(e+fx))^{-m-1}d\sin(e+fx)}{\sqrt{2}f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}}}{a^2 2^{m+\frac{1}{2}}\cos(e+fx)(a\sin(e+fx)+a)^m}$$

$$\frac{\cos(e+fx)(Ad^2-Bcd+c^2C)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

↓ 27

$$\frac{a^2 C(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)\int\frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}(c+d\sin(e+fx))^{-m-1}d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}}}{a^2 2^{m+\frac{1}{2}}\cos(e+fx)(a\sin(e+fx)+a)^m}$$

$$\frac{\cos(e+fx)(Ad^2-Bcd+c^2C)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

↓ 156

$$\frac{a^2 C(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)(c+d\sin(e+fx))^{-m}\left(\frac{c+d\sin(e+fx)}{c-d}\right)^m\int\frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}\left(\frac{c}{c-d}+\frac{d\sin(e+fx)}{c-d}\right)^{-m-1}d\sin(e+fx)}{f(c-d)(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}}}{a^2 2^{m+\frac{1}{2}}\cos(e+fx)(a\sin(e+fx)+a)^m}$$

$$\frac{\cos(e+fx)(Ad^2-Bcd+c^2C)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

↓ 155

$$\frac{\sqrt{2}aC(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)(a\sin(e+fx)+a)^{m+1}(c+d\sin(e+fx))^{-m}\left(\frac{c+d\sin(e+fx)}{c-d}\right)^m\text{AppellF1}\left(m+\frac{3}{2},\frac{1}{2},m+1,m+\frac{5}{2},\frac{1}{2}(\sin(e+fx)+1)\right)}{f(2m+3)(c-d)(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}}$$

$$\frac{\cos(e+fx)(Ad^2-Bcd+c^2C)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

input `Int[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(-2 - m)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

```
output ((c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e
+ f*x])^(-1 - m))/(d*(c^2 - d^2)*f*(1 + m)) + (-((2^(1/2 + m)*a^2*(c*d*(A
+ C + A*m + B*m + C*m) - c^2*(C + 2*C*m) - d^2*(A*m + B*(1 + m) - C*(1 + m
)))*Cos[e + f*x]*Hypergeometric2F1[1/2, 1/2 - m, 3/2, ((c - d)*(1 - Sin[e
+ f*x]))/(2*(c + d*Sin[e + f*x]))]*(a + a*Sin[e + f*x])^(-1 + m)*((c + d)
*(1 + Sin[e + f*x]))/(c + d*Sin[e + f*x])^(1/2 - m))/((c + d)*f*(c + d*Si
n[e + f*x])^m)) + (Sqrt[2]*a*C*(c^2 - d^2)*(1 + m)*AppellF1[3/2 + m, 1/2,
1 + m, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))] *C
os[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*((c + d*Si
n[e + f*x])/(c - d))^m)/((c - d)*f*(3 + 2*m)*(a - a*Sin[e + f*x])*(c + d*S
in[e + f*x])^m))/(a*d*(c^2 - d^2)*(1 + m))
```

### 3.26.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 142 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^(p_), x_] := Simp[((a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/((b*e
- a*f)*(m + 1)))*Hypergeometric2F1[m + 1, -n, m + 2, (-d*e - c*f))*((a +
b*x)/((b*c - a*d)*(e + f*x)))]/((b*e - a*f)*((c + d*x)/((b*c - a*d)*(e +
f*x))))^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[m + n + p + 2,
0] && !IntegerQ[n]
```

```
rule 155 Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Sim
plify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplifierQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplifierQ[e + f*x, a + b*x])
```



rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p])*((b*(e + f*x)/(b*e - a*f))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`

rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n])*((b*(c + d*x)/(b*c - a*d))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]])*Sqrt[a - b*Sin[e + f*x]]) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]`

rule 3466 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]`

rule 3522 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[-(c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(a*d*m + b*c*(n + 1)) + (c*C - B*d)*(a*c*m + b*d*(n + 1)) + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && (LtQ[n, -1] || EqQ[m + n + 2, 0])`

### 3.26.4 Maple [F]

$$\int (a + a \sin(fx + e))^m (c + d \sin(fx + e))^{-2-m} (A + B \sin(fx + e) + C \sin^2(fx + e)) dx$$

input `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

output `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

### 3.26.5 Fricas [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^{-m-2} dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e) + c)^(-m - 2), x)`

**3.26.6 Sympy [F(-1)]**

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

= Timed out

input `integrate((a+a*sin(f*x+e))**m*(c+d*sin(f*x+e))**(-2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output Timed out

**3.26.7 Maxima [F]**

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(fx + e) + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^{-m-2} dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))**(-2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)**2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)**m*(d*sin(f*x + e) + c)**(-m - 2), x)`

**3.26.8 Giac [F]**

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(fx + e) + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^{-m-2} dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))**(-2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)**2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)**m*(d*sin(f*x + e) + c)**(-m - 2), x)`

3.26.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

**3.26.9 Mupad [F(-1)]**

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int \frac{(a + a \sin(e + fx))^m (C \sin(e + fx)^2 + B \sin(e + fx) + A)}{(c + d \sin(e + fx))^{m+2}} dx$$

input `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c + d*sin(e + f*x))^(m + 2),x)`

output `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c + d*sin(e + f*x))^(m + 2), x)`

### 3.27 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{3/2} (A + B \sin(e -$

3.27.1	Optimal result	244
3.27.2	Mathematica [F]	245
3.27.3	Rubi [A] (verified)	245
3.27.4	Maple [F]	249
3.27.5	Fricas [F]	250
3.27.6	Sympy [F(-1)]	250
3.27.7	Maxima [F]	250
3.27.8	Giac [F]	251
3.27.9	Mupad [F(-1)]	251

#### 3.27.1 Optimal result

Integrand size = 47, antiderivative size = 406

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx =$$

$$\frac{2C \cos(e + fx)(a + a \sin(e + fx))^m (c + d \sin(e + fx))^{5/2}}{df(7 + 2m)}$$

$$+ \frac{\sqrt{2}(c - d)(2c(C + 2Cm) - d(7B - 5C + 2Bm + 2Cm - A(7 + 2m))) \operatorname{AppellF1}\left(\frac{1}{2} + m, \frac{1}{2}, -\frac{3}{2}, \frac{3}{2} + m, \frac{1}{2}\right)}{df(1 + 2m)(7 + 2m)\sqrt{1 - \sin(e + fx)}}$$

$$- \frac{\sqrt{2}(c - d)(2cC(1 + m) - d(2Cm + B(7 + 2m))) \operatorname{AppellF1}\left(\frac{3}{2} + m, \frac{1}{2}, -\frac{3}{2}, \frac{5}{2} + m, \frac{1}{2}(1 + \sin(e + fx))\right), -\frac{d}{a}}{adf(3 + 2m)(7 + 2m)\sqrt{1 - \sin(e + fx)}\sqrt{\frac{c+d}{a}}}$$

output

```
-2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(5/2)/d/f/(7+2*m)+(c-d)
)*(2*c*(2*C*m+C)-d*(7*B-5*C+2*B*m+2*C*m-A*(7+2*m)))*AppellF1(1/2+m,-3/2,1/
2,3/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*
x+e))^m*2^(1/2)*(c+d*sin(f*x+e))^(1/2)/d/f/(1+2*m)/(7+2*m)/(1-sin(f*x+e))^
(1/2)/((c+d*sin(f*x+e))/(c-d))^(1/2)-(c-d)*(2*c*C*(1+m)-d*(2*C*m+B*(7+2*m)
))*AppellF1(3/2+m,-3/2,1/2,5/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e
))*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)*2^(1/2)*(c+d*sin(f*x+e))^(1/2)/a/d/f/
(3+2*m)/(7+2*m)/(1-sin(f*x+e))^(1/2)/((c+d*sin(f*x+e))/(c-d))^(1/2)
```

### 3.27.2 Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input `Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x]`

### 3.27.3 Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 425, normalized size of antiderivative = 1.05, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.234$ , Rules used = {3042, 3524, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin(e + fx)^2) dx \\ & \quad \downarrow \text{3524} \\ & \frac{2 \int \frac{1}{2} (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{3/2} (a(Ad(2m + 7) + C(5d + 2cm)) + a(2Cdm - 2cC(m + 1) + Bd(2m - 1))) dx}{ad(2m + 7)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}}{df(2m + 7)} \\ & \quad \downarrow \text{27} \\ & \frac{\int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{3/2} (a(Ad(2m + 7) + C(5d + 2cm)) + a(2Cdm - 2cC(m + 1) + Bd(2m - 1))) dx}{ad(2m + 7)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}}{df(2m + 7)} \end{aligned}$$

---

3.27.  $\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{3/2} (a(Ad(2m + 7) + C(5d + 2cm)) + a(2Cdm - 2cC(m + 1) + Bd(2m - 1)))}{ad(2m + 7)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}$$

↓ 3466

$$\frac{a(2c(2Cm + C) - d(-A(2m + 7) + 2Bm + 7B + 2Cm - 5C)) \int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{3/2} dx + (2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2})}{ad(2m + 7)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}$$

↓ 3042

$$\frac{a(2c(2Cm + C) - d(-A(2m + 7) + 2Bm + 7B + 2Cm - 5C)) \int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{3/2} dx + (2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2})}{ad(2m + 7)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}$$

↓ 3267

$$\frac{a^3 \cos(e + fx)(2c(2Cm + C) - d(-A(2m + 7) + 2Bm + 7B + 2Cm - 5C)) \int \frac{(\sin(e + fx)a + a)^{m - \frac{1}{2}} (c + d \sin(e + fx))^{3/2} d \sin(e + fx)}{\sqrt{a - a \sin(e + fx)} \sqrt{a \sin(e + fx) + a}} + a^2 \cos(e + fx)(Bd(2m - 1))}{ad(2m + 7)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}$$

↓ 157

$$\frac{a^3 \sqrt{1 - \sin(e + fx)} \cos(e + fx)(2c(2Cm + C) - d(-A(2m + 7) + 2Bm + 7B + 2Cm - 5C)) \int \frac{\sqrt{2}(\sin(e + fx)a + a)^{m - \frac{1}{2}} (c + d \sin(e + fx))^{3/2} d \sin(e + fx)}{\sqrt{2f(a - a \sin(e + fx))} \sqrt{a \sin(e + fx) + a}} + a^2 \cos(e + fx)(Bd(2m - 1))}{ad(2m + 7)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}$$

↓ 27

---

3.27.  $\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+7) + 2Bm + 7B + 2Cm - 5C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} (c+d \sin(e+fx))^{3/2} d \sin(e+fx)}{\sqrt{1-\sin(e+fx)}}}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} + \frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+7) + 2Bm + 7B + 2Cm - 5C))}{ad(2m+7)}$$

$$\frac{2C \cos(e+fx) (a \sin(e+fx) + a)^m (c+d \sin(e+fx))^{5/2}}{df(2m+7)}$$

↓ 156

$$\frac{a^3(c-d) \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+7) + 2Bm + 7B + 2Cm - 5C)) \sqrt{c+d \sin(e+fx)} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} \left(\frac{c}{c-d} + \frac{d \sin(e+fx)}{c-d}\right)}{\sqrt{1-\sin(e+fx)}}}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a} \sqrt{\frac{c+d \sin(e+fx)}{c-d}}}$$

$$\frac{2C \cos(e+fx) (a \sin(e+fx) + a)^m (c+d \sin(e+fx))^{5/2}}{df(2m+7)}$$

↓ 155

$$\frac{\sqrt{2} a^2 (c-d) \sqrt{1-\sin(e+fx)} \cos(e+fx) (a \sin(e+fx)+a)^m (2c(2Cm+C) - d(-A(2m+7) + 2Bm + 7B + 2Cm - 5C)) \sqrt{c+d \sin(e+fx)} \text{AppellF1}\left(m+\frac{1}{2}, m+\frac{1}{2}, m+\frac{1}{2}, m+\frac{1}{2}, \frac{c+d \sin(e+fx)}{c-d}, \frac{c+d \sin(e+fx)}{c-d}\right)}{f(2m+1)(a-a \sin(e+fx)) \sqrt{\frac{c+d \sin(e+fx)}{c-d}}}$$

$$\frac{2C \cos(e+fx) (a \sin(e+fx) + a)^m (c+d \sin(e+fx))^{5/2}}{df(2m+7)}$$

input `Int[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `(-2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(5/2))/(d*f*(7 + 2*m)) + ((Sqrt[2]*a^2*(c - d)*(2*c*(C + 2*C*m) - d*(7*B - 5*C + 2*B*m + 2*C*m - A*(7 + 2*m)))*AppellF1[1/2 + m, 1/2, -3/2, 3/2 + m, (1 + Sin[e + f*x])/2, -(d*(1 + Sin[e + f*x]))/(c - d)]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]])/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]) + (Sqrt[2]*a*(c - d)*(2*C*d*m - 2*c*C*(1 + m) + B*d*(7 + 2*m))*AppellF1[3/2 + m, 1/2, -3/2, 5/2 + m, (1 + Sin[e + f*x])/2, -(d*(1 + Sin[e + f*x]))/(c - d)]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[c + d*Sin[e + f*x]])/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]))/(a*d*(7 + 2*m))`



## 3.27.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]])*Sqrt[a - b*Sin[e + f*x]]) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]`

rule 3466 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]`

rule 3524 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !Lt Q[m, -2^(-1)] && NeQ[m + n + 2, 0]`

### 3.27.4 Maple [F]

$$\int (a + a \sin(fx + e))^m (c + d \sin(fx + e))^{\frac{3}{2}} (A + B \sin(fx + e) + C \sin^2(fx + e)) dx$$

input `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

output `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

**3.27.5 Fracas [F]**

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e) + B \sin(fx + e) + A)(d \sin(fx + e) + c)^{3/2} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `integral(-((C*c + B*d)*cos(f*x + e)^2 - (A + C)*c - B*d + (C*d*cos(f*x + e))^2 - B*c - (A + C)*d)*sin(f*x + e)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m, x)`

**3.27.6 Sympy [F(-1)]**

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(c+d*sin(f*x+e))**(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output `Timed out`

**3.27.7 Maxima [F]**

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e) + B \sin(fx + e) + A)(d \sin(fx + e) + c)^{3/2} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(d*sin(f*x + e) + c)^(3/2)*(a*sin(f*x + e) + a)^m, x)`

### 3.27.8 Giac [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e) + B \sin(fx + e) + A) (d \sin(fx + e) + c)^{3/2} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(d*sin(f*x + e) + c)^(3/2)*(a*sin(f*x + e) + a)^m, x)`

### 3.27.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (C \sin^2(e + fx) + B \sin(e + fx) + A) dx$$

input `int((a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(3/2)*(A + B*sin(e + f*x) + C*sin(e + f*x)^2),x)`

output `int((a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(3/2)*(A + B*sin(e + f*x) + C*sin(e + f*x)^2), x)`

### 3.28 $\int (a+a \sin(e+fx))^m \sqrt{c+d \sin(e+fx)}(A+B \sin(e+fx) + C \sin^2(e+fx)) dx$

3.28.1	Optimal result	252
3.28.2	Mathematica [F]	253
3.28.3	Rubi [A] (verified)	253
3.28.4	Maple [F]	257
3.28.5	Fricas [F]	258
3.28.6	Sympy [F]	258
3.28.7	Maxima [F]	258
3.28.8	Giac [F]	259
3.28.9	Mupad [F(-1)]	259

#### 3.28.1 Optimal result

Integrand size = 47, antiderivative size = 396

$$\int (a+a \sin(e+fx))^m \sqrt{c+d \sin(e+fx)}(A+B \sin(e+fx) + C \sin^2(e+fx)) dx$$

$$= -\frac{2C \cos(e+fx)(a+a \sin(e+fx))^m (c+d \sin(e+fx))^{3/2}}{df(5+2m)}$$

$$+ \frac{\sqrt{2}(2c(C+2Cm) - d(5B-3C+2Bm+2Cm - A(5+2m))) \operatorname{AppellF1}\left(\frac{1}{2}+m, \frac{1}{2}, -\frac{1}{2}, \frac{3}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right)}{df(1+2m)(5+2m)\sqrt{1-\sin(e+fx)}}$$

$$- \frac{\sqrt{2}(2cC(1+m) - d(2Cm+B(5+2m))) \operatorname{AppellF1}\left(\frac{3}{2}+m, \frac{1}{2}, -\frac{1}{2}, \frac{5}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right), -\frac{d(1+\sin(e+fx))}{c}}{adf(3+2m)(5+2m)\sqrt{1-\sin(e+fx)}\sqrt{\frac{c+d \sin(e+fx)}{c}}}$$

output

```
-2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)/d/f/(5+2*m)+(2*c
*(2*C*m+C)-d*(5*B-3*C+2*B*m+2*C*m-A*(5+2*m)))*AppellF1(1/2+m,-1/2,1/2,3/2+
m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^
m*2^(1/2)*(c+d*sin(f*x+e))^(1/2)/d/f/(1+2*m)/(5+2*m)/(1-sin(f*x+e))^(1/2)/
((c+d*sin(f*x+e))/(c-d))^(1/2)-(2*c*C*(1+m)-d*(2*C*m+B*(5+2*m)))*AppellF1(
3/2+m,-1/2,1/2,5/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e
)*(a+a*sin(f*x+e))^(1+m)*2^(1/2)*(c+d*sin(f*x+e))^(1/2)/a/d/f/(3+2*m)/(5+2
*m)/(1-sin(f*x+e))^(1/2)/((c+d*sin(f*x+e))/(c-d))^(1/2)
```

### 3.28.2 Mathematica [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input `Integrate[(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]]*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `Integrate[(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]]*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x]`

### 3.28.3 Rubi [A] (verified)

Time = 1.17 (sec) , antiderivative size = 415, normalized size of antiderivative = 1.05, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.234$ , Rules used = {3042, 3524, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (a \sin(e + fx) + a)^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin(e + fx)^2) dx$$

$$\downarrow \text{3524}$$

$$\frac{2 \int \frac{1}{2} (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} (a(Ad(2m + 5) + C(3d + 2cm)) + a(2Cdm - 2cC(m + 1) + Bd(2m + 5)))}{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}} df(2m + 5)}{ad(2m + 5)}$$

$$\downarrow \text{27}$$

---

3.28.  $\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$

$$\frac{\int (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} (a(Ad(2m + 5) + C(3d + 2cm)) + a(2Cdm - 2cC(m + 1) + Bd(2m + 5)))}{ad(2m + 5)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}}{df(2m + 5)}$$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} (a(Ad(2m + 5) + C(3d + 2cm)) + a(2Cdm - 2cC(m + 1) + Bd(2m + 5)))}{ad(2m + 5)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}}{df(2m + 5)}$$

↓ 3466

$$\frac{a(2c(2Cm + C) - d(-A(2m + 5) + 2Bm + 5B + 2Cm - 3C)) \int (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} dx + (Bd(2m + 5))}{ad(2m + 5)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}}{df(2m + 5)}$$

↓ 3042

$$\frac{a(2c(2Cm + C) - d(-A(2m + 5) + 2Bm + 5B + 2Cm - 3C)) \int (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} dx + (Bd(2m + 5))}{ad(2m + 5)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}}{df(2m + 5)}$$

↓ 3267

$$\frac{a^3 \cos(e + fx)(2c(2Cm + C) - d(-A(2m + 5) + 2Bm + 5B + 2Cm - 3C)) \int \frac{(\sin(e + fx)a + a)^{m - \frac{1}{2}} \sqrt{c + d \sin(e + fx)}}{\sqrt{a - a \sin(e + fx)}} d \sin(e + fx)}{f \sqrt{a - a \sin(e + fx)} \sqrt{a \sin(e + fx) + a}} + \frac{a^2 \cos(e + fx)(Bd(2m + 5))}{ad(2m + 5)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}}{df(2m + 5)}$$

↓ 157

$$\frac{a^3 \sqrt{1 - \sin(e + fx)} \cos(e + fx)(2c(2Cm + C) - d(-A(2m + 5) + 2Bm + 5B + 2Cm - 3C)) \int \frac{\sqrt{2}(\sin(e + fx)a + a)^{m - \frac{1}{2}} \sqrt{c + d \sin(e + fx)}}{\sqrt{1 - \sin(e + fx)}} d \sin(e + fx)}{\sqrt{2}f(a - a \sin(e + fx)) \sqrt{a \sin(e + fx) + a}} + \frac{a^2 \sqrt{1 - \sin(e + fx)} (Bd(2m + 5))}{ad(2m + 5)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}}{df(2m + 5)}$$

↓ 27

---

3.28.  $\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+5) + 2Bm + 5B + 2Cm - 3C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} \sqrt{c+d\sin(e+fx)}}{\sqrt{1-\sin(e+fx)}} d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}} + \frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+5) + 2Bm + 5B + 2Cm - 3C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} \sqrt{\frac{c}{c-d} + \frac{d\sin(e+fx)}{c-d}}}{\sqrt{1-\sin(e+fx)}} d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a} \sqrt{\frac{c+d\sin(e+fx)}{c-d}}}$$


---


$$\frac{2C \cos(e+fx)(a\sin(e+fx)+a)^m (c+d\sin(e+fx))^{3/2}}{df(2m+5)} \qquad ad(2m+5)$$

↓ 156

$$\frac{2C \cos(e+fx)(a\sin(e+fx)+a)^m (c+d\sin(e+fx))^{3/2}}{df(2m+5)}$$

↓ 155

$$\frac{\sqrt{2}a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (a\sin(e+fx)+a)^m (2c(2Cm+C) - d(-A(2m+5) + 2Bm + 5B + 2Cm - 3C)) \sqrt{c+d\sin(e+fx)} \operatorname{AppellF1}\left(m+\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\right)}{f(2m+1)(a-a\sin(e+fx))\sqrt{\frac{c+d\sin(e+fx)}{c-d}}}$$


---


$$\frac{2C \cos(e+fx)(a\sin(e+fx)+a)^m (c+d\sin(e+fx))^{3/2}}{df(2m+5)}$$

input `Int[(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]]*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `(-2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2))/(d*f*(5 + 2*m)) + ((Sqrt[2]*a^2*(2*c*(C + 2*C*m) - d*(5*B - 3*C + 2*B*m + 2*C*m - A*(5 + 2*m)))*AppellF1[1/2 + m, 1/2, -1/2, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]])/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]) + (Sqrt[2]*a*(2*C*d*m - 2*C*C*(1 + m) + B*d*(5 + 2*m))*AppellF1[3/2 + m, 1/2, -1/2, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[c + d*Sin[e + f*x]])/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]))/(a*d*(5 + 2*m))`

---

3.28.  $\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$



## 3.28.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]])*Sqrt[a - b*Sin[e + f*x]]) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]`

rule 3466 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]`

rule 3524 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !Lt Q[m, -2^(-1)] && NeQ[m + n + 2, 0]`

### 3.28.4 Maple [F]

$$\int (a + a \sin(fx + e))^m \sqrt{c + d \sin(fx + e)} (A + B \sin(fx + e) + C(\sin^2(fx + e))) dx$$

input `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

output `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

**3.28.5 Fricas [F]**

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A) \sqrt{d \sin(fx + e) + c} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m, x)`

**3.28.6 Sympy [F]**

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a(\sin(e + fx) + 1))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input `integrate((a+a*sin(f*x+e))**m*(c+d*sin(f*x+e))**(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output `Integral((a*(sin(e + f*x) + 1))**m*sqrt(c + d*sin(e + f*x))*(A + B*sin(e + f*x) + C*sin(e + f*x)**2), x)`

**3.28.7 Maxima [F]**

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A) \sqrt{d \sin(fx + e) + c} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*sqrt(d*sin(f*x + e) + c) * (a*sin(f*x + e) + a)^m, x)`

### 3.28.8 Giac [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx \\ &= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A) \sqrt{d \sin(fx + e) + c} (a \sin(fx + e) + a)^m dx \end{aligned}$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*sqrt(d*sin(f*x + e) + c) * (a*sin(f*x + e) + a)^m, x)`

### 3.28.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx \\ &= \int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (C \sin(e + fx)^2 + B \sin(e + fx) + A) dx \end{aligned}$$

input `int((a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(1/2)*(A + B*sin(e + f*x) + C*sin(e + f*x)^2),x)`

output `int((a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(1/2)*(A + B*sin(e + f*x) + C*sin(e + f*x)^2), x)`

$$3.29 \quad \int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx$$

3.29.1	Optimal result	260
3.29.2	Mathematica [F]	261
3.29.3	Rubi [A] (verified)	261
3.29.4	Maple [F]	265
3.29.5	Fricas [F]	266
3.29.6	Sympy [F]	266
3.29.7	Maxima [F]	266
3.29.8	Giac [F]	267
3.29.9	Mupad [F(-1)]	267

### 3.29.1 Optimal result

Integrand size = 47, antiderivative size = 389

$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx$$

$$= -\frac{2C \cos(e+fx)(a+a \sin(e+fx))^m \sqrt{c+d \sin(e+fx)}}{df(3+2m)}$$

$$+ \frac{\sqrt{2}(2c(C+2Cm)-d(3B-C+2Bm+2Cm-A(3+2m))) \operatorname{AppellF1}\left(\frac{1}{2}+m, \frac{1}{2}, \frac{1}{2}, \frac{3}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right)}{df(1+2m)(3+2m)\sqrt{1-\sin(e+fx)}\sqrt{c+d \sin(e+fx)}}$$

$$- \frac{\sqrt{2}(2cC(1+m)-d(2Cm+B(3+2m))) \operatorname{AppellF1}\left(\frac{3}{2}+m, \frac{1}{2}, \frac{1}{2}, \frac{5}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right), -\frac{d(1+\sin(e+fx))}{c-d}}{adf(3+2m)^2\sqrt{1-\sin(e+fx)}\sqrt{c+d \sin(e+fx)}}$$

output

```
-2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)/d/f/(3+2*m)+(2*c
*(2*C*m+C)-d*(3*B-C+2*B*m+2*C*m-A*(3+2*m)))*AppellF1(1/2+m,1/2,1/2,3/2+m,-
d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^m*2
^(1/2)*((c+d*sin(f*x+e))/(c-d))^(1/2)/d/f/(1+2*m)/(3+2*m)/(1-sin(f*x+e))^(
1/2)/(c+d*sin(f*x+e))^(1/2)-(2*c*C*(1+m)-d*(2*C*m+B*(3+2*m)))*AppellF1(3/2
+m,1/2,1/2,5/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a
+a*sin(f*x+e))^(1+m)*2^(1/2)*((c+d*sin(f*x+e))/(c-d))^(1/2)/a/d/f/(3+2*m)^
2/(1-sin(f*x+e))^(1/2)/(c+d*sin(f*x+e))^(1/2)
```

### 3.29.2 Mathematica [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

input `Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/Sqrt[c + d*Sin[e + f*x]],x]`

output `Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/Sqrt[c + d*Sin[e + f*x]], x]`

### 3.29.3 Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 415, normalized size of antiderivative = 1.07, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.234$ , Rules used = {3042, 3524, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

↓ 3042

$$\int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2)}{\sqrt{c + d \sin(e + fx)}} dx$$

↓ 3524

$$2 \int \frac{(\sin(e + fx)a + a)^m (a(Ad(2m + 3) + C(d + 2cm)) + a(2Cdm - 2cC(m + 1) + Bd(2m + 3)) \sin(e + fx))}{2\sqrt{c + d \sin(e + fx)}} dx$$


---


$$\frac{ad(2m + 3)}{2C \cos(e + fx)(a \sin(e + fx) + a)^m \sqrt{c + d \sin(e + fx)}} - \frac{df(2m + 3)}{2C \cos(e + fx)(a \sin(e + fx) + a)^m \sqrt{c + d \sin(e + fx)}}$$

↓ 27

---

3.29.  $\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$

$$\frac{\int \frac{(\sin(e+fx)a+a)^m (a(Ad(2m+3)+C(d+2cm))+a(2Cdm-2cC(m+1)+Bd(2m+3)) \sin(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx}{ad(2m+3)} - \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}$$

↓ 3042

$$\frac{\int \frac{(\sin(e+fx)a+a)^m (a(Ad(2m+3)+C(d+2cm))+a(2Cdm-2cC(m+1)+Bd(2m+3)) \sin(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx}{ad(2m+3)} - \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}$$

↓ 3466

$$\frac{a(2c(2Cm+C) - d(-A(2m+3) + 2Bm + 3B + 2Cm - C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx + (Bd(2m+3) - 2cC(m+1))}{ad(2m+3)} - \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}$$

↓ 3042

$$\frac{a(2c(2Cm+C) - d(-A(2m+3) + 2Bm + 3B + 2Cm - C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx + (Bd(2m+3) - 2cC(m+1))}{ad(2m+3)} - \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}$$

↓ 3267

$$\frac{a^3 \cos(e+fx)(2c(2Cm+C) - d(-A(2m+3) + 2Bm + 3B + 2Cm - C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{a-a \sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx) + \frac{a^2 \cos(e+fx)(Bd(2m+3) - 2cC(m+1))}{f}}{ad(2m+3)} - \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}$$

↓ 157

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)(2c(2Cm+C) - d(-A(2m+3) + 2Bm + 3B + 2Cm - C)) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx) + \frac{a^2 \sqrt{1-\sin(e+fx)}(Bd(2m+3) - 2cC(m+1))}{f}}{ad(2m+3)} - \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}$$

↓ 27

---

3.29.  $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx$

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+3) + 2Bm + 3B + 2Cm - C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}\sqrt{c+d\sin(e+fx)}} d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}} + \frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+3) + 2Bm + 3B + 2Cm - C))}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}}$$

$ad(2m + 3)$

$$\frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m \sqrt{c + d \sin(e + fx)}}{df(2m + 3)}$$

↓ 156

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+3) + 2Bm + 3B + 2Cm - C)) \sqrt{\frac{c+d\sin(e+fx)}{c-d}} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}\sqrt{\frac{c}{c-d} + \frac{d\sin(e+fx)}{c-d}}} d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}\sqrt{c+d\sin(e+fx)}}$$

$ad(2m + 3)$

$$\frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m \sqrt{c + d \sin(e + fx)}}{df(2m + 3)}$$

↓ 155

$$\frac{\sqrt{2}a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (a \sin(e+fx) + a)^m (2c(2Cm+C) - d(-A(2m+3) + 2Bm + 3B + 2Cm - C)) \sqrt{\frac{c+d\sin(e+fx)}{c-d}} \text{AppellF1}\left(m+\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, m\right)}{f(2m+1)(a-a\sin(e+fx))\sqrt{c+d\sin(e+fx)}}$$

$$\frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m \sqrt{c + d \sin(e + fx)}}{df(2m + 3)}$$

```
input Int[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/Sqrt[
c + d*Sin[e + f*x]],x]
```

```
output (-2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]]/(d*f*(
3 + 2*m)) + ((Sqrt[2]*a^2*(2*c*(C + 2*C*m) - d*(3*B - C + 2*B*m + 2*C*m -
A*(3 + 2*m)))*AppellF1[1/2 + m, 1/2, 1/2, 3/2 + m, (1 + Sin[e + f*x])/2, -
((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a +
a*Sin[e + f*x])^m*Sqrt[(c + d*Sin[e + f*x])/(c - d)]/(f*(1 + 2*m)*(a - a
*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x])) + (Sqrt[2]*a*(2*C*d*m - 2*c*C*(1
+ m) + B*d*(3 + 2*m))*AppellF1[3/2 + m, 1/2, 1/2, 5/2 + m, (1 + Sin[e + f
x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f
x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[(c + d*Sin[e + f*x])/(c - d)]/(f*(3
+ 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]))/(a*d*(3 + 2*m))
```



## 3.29.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]])*Sqrt[a - b*Sin[e + f*x]]) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]`

rule 3466 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]`

rule 3524 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !Lt Q[m, -2^(-1)] && NeQ[m + n + 2, 0]`

### 3.29.4 Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + B \sin(fx + e) + C(\sin^2(fx + e)))}{\sqrt{c + d \sin(fx + e)}} dx$$

input `int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x)`

output `int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x)`

## 3.29.5 Fricas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin(fx + e))^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m}{\sqrt{d \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x, algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*(a*sin(f*x + e) + a)^m/sqrt(d*sin(f*x + e) + c), x)`

## 3.29.6 Sympy [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(a(\sin(e + fx) + 1))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

input `integrate((a+a*sin(f*x+e))**m*(A+B*sin(f*x+e)+C*sin(f*x+e)**2)/(c+d*sin(f*x+e))**(1/2),x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + B*sin(e + f*x) + C*sin(e + f*x)**2)/sqrt(c + d*sin(e + f*x)), x)`

## 3.29.7 Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin(fx + e))^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m}{\sqrt{d \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/sqrt(d*sin(f*x + e) + c), x)`

### 3.29.8 Giac [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m}{\sqrt{d \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/sqrt(d*sin(f*x + e) + c), x)`

### 3.29.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(a + a \sin(e + fx))^m (C \sin(e + fx)^2 + B \sin(e + fx) + A)}{\sqrt{c + d \sin(e + fx)}} dx$$

input `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c + d*sin(e + f*x))^(1/2),x)`

output `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c + d*sin(e + f*x))^(1/2), x)`

$$3.30 \quad \int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx$$

3.30.1	Optimal result	268
3.30.2	Mathematica [F]	269
3.30.3	Rubi [A] (verified)	269
3.30.4	Maple [F]	273
3.30.5	Fricas [F]	274
3.30.6	Sympy [F]	274
3.30.7	Maxima [F]	274
3.30.8	Giac [F]	275
3.30.9	Mupad [F(-1)]	275

### 3.30.1 Optimal result

Integrand size = 47, antiderivative size = 433

$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx = \frac{2(c^2C - Bcd + Ad^2) \cos(e+fx)(a+a \sin(e+fx))^{m-1/2}}{d(c^2 - d^2) f \sqrt{c+d \sin(e+fx)}} + \frac{\sqrt{2}(d^2(A+B-C+4Am) - cd(A+B+C+4Bm) + 2c^2(C+2Cm)) \operatorname{AppellF1}\left(\frac{1}{2}+m, \frac{1}{2}, \frac{1}{2}, \frac{3}{2}+m, \frac{1}{2}\right)}{d(c^2 - d^2) f(1+2m) \sqrt{1-\sin(e+fx)}} + \frac{\sqrt{2}(d(Bc - Ad)(1+2m) + C(d^2 - 2c^2(1+m))) \operatorname{AppellF1}\left(\frac{3}{2}+m, \frac{1}{2}, \frac{1}{2}, \frac{5}{2}+m, \frac{1}{2}(1+\sin(e+fx))\right)}{ad(c^2 - d^2) f(3+2m) \sqrt{1-\sin(e+fx)} \sqrt{c+d \sin(e+fx)}} - \frac{d(1-\sin(e+fx))^{m-1/2}}{d(c^2 - d^2) f}$$

output

```
2*(A*d^2-B*c*d+C*c^2)*cos(f*x+e)*(a+a*sin(f*x+e))^m/d/(c^2-d^2)/f/(c+d*sin
(f*x+e))^(1/2)-(d^2*(4*A*m+A+B-C)-c*d*(4*B*m+A+B+C)+2*c^2*(2*C*m+C))*Appel
lF1(1/2+m,1/2,1/2,3/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*
x+e)*(a+a*sin(f*x+e))^m*2^(1/2)*((c+d*sin(f*x+e))/(c-d))^(1/2)/d/(c^2-d^2)
/f/(1+2*m)/(1-sin(f*x+e))^(1/2)/(c+d*sin(f*x+e))^(1/2)-(d*(-A*d+B*c)*(1+2*
m)+C*(d^2-2*c^2*(1+m)))*AppellF1(3/2+m,1/2,1/2,5/2+m,-d*(1+sin(f*x+e))/(c-
d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)*2^(1/2)*((c+d*sin
(f*x+e))/(c-d))^(1/2)/a/d/(c^2-d^2)/f/(3+2*m)/(1-sin(f*x+e))^(1/2)/(c+d*si
n(f*x+e))^(1/2)
```

---


$$3.30. \quad \int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx$$

### 3.30.2 Mathematica [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx$$

input `Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(3/2),x]`

output `Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(3/2), x]`

### 3.30.3 Rubi [A] (verified)

Time = 1.34 (sec) , antiderivative size = 451, normalized size of antiderivative = 1.04, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.234$ , Rules used = {3042, 3522, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2)}{(c + d \sin(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{3522} \\ & \frac{2 \cos(e + fx) (Ad^2 - Bcd + c^2C) (a \sin(e + fx) + a)^m}{df (c^2 - d^2) \sqrt{c + d \sin(e + fx)}} - \\ & 2 \int \frac{(\sin(e + fx)a + a)^m (a((cC - Bd)(d - 2cm) + Ad(c - 2dm)) - a(-2C(m + 1)c^2 + Cd^2 + d(Bc - Ad)(2m + 1) \sin(e + fx)))}{2\sqrt{c + d \sin(e + fx)} ad (c^2 - d^2)} dx \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{(\sin(e + fx)a + a)^m (a((cC - Bd)(d - 2cm) + Ad(c - 2dm)) - a(-2C(m + 1)c^2 + Cd^2 + d(Bc - Ad)(2m + 1) \sin(e + fx)))}{\sqrt{c + d \sin(e + fx)} ad (c^2 - d^2)} dx}{ad (c^2 - d^2)} + \\ & \frac{2 \cos(e + fx) (Ad^2 - Bcd + c^2C) (a \sin(e + fx) + a)^m}{df (c^2 - d^2) \sqrt{c + d \sin(e + fx)}} \end{aligned}$$

---

3.30.  $\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx$

$$\begin{aligned}
& \int \frac{(\sin(e+fx)a+a)^m (a((cC-Bd)(d-2cm)+Ad(c-2dm))-a(-2C(m+1)c^2+Cd^2+d(Bc-Ad)(2m+1)) \sin(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx \\
& \quad + \frac{ad(c^2-d^2)}{2 \cos(e+fx) (Ad^2-Bcd+c^2C) (a \sin(e+fx)+a)^m} \\
& \quad \frac{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& -a(-cd(A+4Bm+B+C)+d^2(4Am+A+B-C)+2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx - (d(2m+1)(Bc- \\
& \quad ad(c^2-d^2)) \\
& \quad \frac{2 \cos(e+fx) (Ad^2-Bcd+c^2C) (a \sin(e+fx)+a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& -a(-cd(A+4Bm+B+C)+d^2(4Am+A+B-C)+2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx - (d(2m+1)(Bc- \\
& \quad ad(c^2-d^2)) \\
& \quad \frac{2 \cos(e+fx) (Ad^2-Bcd+c^2C) (a \sin(e+fx)+a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}} \\
& \quad \downarrow \text{3267} \\
& \frac{a^3 \cos(e+fx)(-cd(A+4Bm+B+C)+d^2(4Am+A+B-C)+2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{a-a \sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx)}{f \sqrt{a-a \sin(e+fx)} \sqrt{a \sin(e+fx)+a}} - \frac{a^2 \cos(e+fx)(d(2m+1)(Bc- \\
& \quad ad(c^2-d^2)) \\
& \quad \frac{2 \cos(e+fx) (Ad^2-Bcd+c^2C) (a \sin(e+fx)+a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}} \\
& \quad \downarrow \text{157} \\
& \frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)(-cd(A+4Bm+B+C)+d^2(4Am+A+B-C)+2c^2(2Cm+C)) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx)}{\sqrt{2}f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} - \frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx)(d(2m+1)(Bc- \\
& \quad ad(c^2-d^2)) \\
& \quad \frac{2 \cos(e+fx) (Ad^2-Bcd+c^2C) (a \sin(e+fx)+a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}} \\
& \quad \downarrow \text{27}
\end{aligned}$$

---

3.30.  $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx$

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (-cd(A+4Bm+B+C)+d^2(4Am+A+B-C)+2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}\sqrt{c+d\sin(e+fx)}} d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}} - \frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (-cd(A+4Bm+B+C)+d^2(4Am+A+B-C)+2c^2(2Cm+C))}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}}$$

$$\frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{df (c^2 - d^2) \sqrt{c + d \sin(e+fx)}} \quad ad (c^2 - d^2)$$

↓ 156

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (-cd(A+4Bm+B+C)+d^2(4Am+A+B-C)+2c^2(2Cm+C)) \sqrt{\frac{c+d\sin(e+fx)}{c-d}} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}\sqrt{\frac{c}{c-d} + \frac{d\sin(e+fx)}{c-d}}} d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}\sqrt{c+d\sin(e+fx)}} - \frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (-cd(A+4Bm+B+C)+d^2(4Am+A+B-C)+2c^2(2Cm+C))}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}\sqrt{c+d\sin(e+fx)}}$$

$$\frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{df (c^2 - d^2) \sqrt{c + d \sin(e+fx)}} \quad ad (c^2 - d^2)$$

↓ 155

$$\frac{\sqrt{2}a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (a \sin(e+fx) + a)^m (-cd(A+4Bm+B+C)+d^2(4Am+A+B-C)+2c^2(2Cm+C)) \sqrt{\frac{c+d\sin(e+fx)}{c-d}} \text{AppellF1}(m, \frac{1}{2}, \frac{1}{2}, \frac{3}{2} + m, \frac{1 + \sin(e+fx)}{2}, -\frac{d(1 + \sin(e+fx))}{c-d})}{f(2m+1)(a-a\sin(e+fx))\sqrt{c+d\sin(e+fx)}} - \frac{\sqrt{2}a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (a \sin(e+fx) + a)^m (-cd(A+4Bm+B+C)+d^2(4Am+A+B-C)+2c^2(2Cm+C))}{f(2m+1)(a-a\sin(e+fx))\sqrt{c+d\sin(e+fx)}}$$

$$\frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{df (c^2 - d^2) \sqrt{c + d \sin(e+fx)}}$$

input `Int[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(3/2),x]`

output `(2*(c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(d*(c^2 - d^2)*f*Sqrt[c + d*Sin[e + f*x]]) + (-((Sqrt[2]*a^2*(d^2*(A + B - C + 4*A*m) - c*d*(A + B + C + 4*B*m) + 2*c^2*(C + 2*C*m))*AppellF1[1/2 + m, 1/2, 1/2, 3/2 + m, (1 + Sin[e + f*x])/2, -(d*(1 + Sin[e + f*x])/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[(c + d*Sin[e + f*x])/(c - d)])/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]) - (Sqrt[2]*a*(C*d^2 - 2*c^2*C*(1 + m) + d*(B*c - A*d)*(1 + 2*m))*AppellF1[3/2 + m, 1/2, 1/2, 5/2 + m, (1 + Sin[e + f*x])/2, -(d*(1 + Sin[e + f*x])/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[(c + d*Sin[e + f*x])/(c - d)])/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]))/(a*d*(c^2 - d^2))`

$$3.30. \quad \int \frac{(a+a\sin(e+fx))^m (A+B\sin(e+fx)+C\sin^2(e+fx))}{(c+d\sin(e+fx))^{3/2}} dx$$



## 3.30.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p)*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]])*Sqrt[a - b*Sin[e + f*x]]) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]`

rule 3466 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]`

rule 3522 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(a*d*m + b*c*(n + 1)) + (c*C - B*d)*(a*c*m + b*d*(n + 1)) + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && (LtQ[n, -1] || EqQ[m + n + 2, 0])`

### 3.30.4 Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + B \sin(fx + e) + C(\sin^2(fx + e)))}{(c + d \sin(fx + e))^{\frac{3}{2}}} dx$$

input `int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x)`

output `int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x)`

### 3.30.5 Fracas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin(fx + e))^2 + B \sin(fx + e) + A}{(d \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x, algorithm="fricas")`

output `integral((C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(d^2*cos(f*x + e)^2 - 2*c*d*sin(f*x + e) - c^2 - d^2), x)`

### 3.30.6 Sympy [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(a(\sin(e + fx) + 1))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))**m*(A+B*sin(f*x+e)+C*sin(f*x+e)**2)/(c+d*sin(f*x+e))**(3/2),x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + B*sin(e + f*x) + C*sin(e + f*x)**2)/(c + d*sin(e + f*x))**(3/2), x)`

### 3.30.7 Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin(fx + e))^2 + B \sin(fx + e) + A}{(d \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/(d*sin(f*x + e) + c)^(3/2), x)`

---

3.30.  $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx$

**3.30.8 Giac [F]**

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin(fx + e))^2 + B \sin(fx + e) + A}{(d \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/(d*sin(f*x + e) + c)^(3/2), x)`

**3.30.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(a + a \sin(e + fx))^m (C \sin(e + fx) + B + A)}{(c + d \sin(e + fx))^{3/2}} dx$$

input `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c + d*sin(e + f*x))^(3/2),x)`

output `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c + d*sin(e + f*x))^(3/2), x)`

**3.31** 
$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{5/2}} dx$$

3.31.1	Optimal result	276
3.31.2	Mathematica [F]	277
3.31.3	Rubi [A] (verified)	277
3.31.4	Maple [F]	281
3.31.5	Fricas [F]	282
3.31.6	Sympy [F(-1)]	282
3.31.7	Maxima [F]	282
3.31.8	Giac [F]	283
3.31.9	Mupad [F(-1)]	283

**3.31.1 Optimal result**

Integrand size = 47, antiderivative size = 451

$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{5/2}} dx = \frac{2(c^2C - Bcd + Ad^2) \cos(e+fx)(a+a \sin(e+fx))^{m-1}}{3d(c^2 - d^2) f(c+d \sin(e+fx))^{3/2}} + \frac{\sqrt{2}(d^2(A - 3B + 3C - 4Am) + cd(3A - B + 3C + 4Bm) - 2c^2(C + 2Cm)) \operatorname{AppellF1}\left(\frac{1}{2} + m, \frac{1}{2}, \frac{3}{2}, \frac{3}{2} + m\right)}{3(c-d)^2 d(c+d) f(1+2m) \sqrt{1 - \sin(e+fx)}} + \frac{\sqrt{2}(Bcd(1 - 2m) + 2c^2C(1 + m) - d^2(A + 3C - 2Am)) \operatorname{AppellF1}\left(\frac{3}{2} + m, \frac{1}{2}, \frac{3}{2}, \frac{5}{2} + m, \frac{1}{2}(1 + \sin(e+fx))\right)}{3a(c-d)^2 d(c+d) f(3+2m) \sqrt{1 - \sin(e+fx)} \sqrt{c+d \sin(e+fx)}}$$

output

```
2/3*(A*d^2-B*c*d+C*c^2)*cos(f*x+e)*(a+a*sin(f*x+e))^m/d/(c^2-d^2)/f/(c+d*sin(f*x+e))^(3/2)+1/3*(d^2*(-4*A*m+A-3*B+3*C)+c*d*(4*B*m+3*A-B+3*C)-2*c^2*(2*C*m+C))*AppellF1(1/2+m,3/2,1/2,3/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^m*2^(1/2)*((c+d*sin(f*x+e))/(c-d))^(1/2)/(c-d)^2/d/(c+d)/f/(1+2*m)/(1-sin(f*x+e))^(1/2)/(c+d*sin(f*x+e))^(1/2)+1/3*(B*c*d*(1-2*m)+2*c^2*C*(1+m)-d^2*(-2*A*m+A+3*C))*AppellF1(3/2+m,3/2,1/2,5/2+m,-d*(1+sin(f*x+e))/(c-d),1/2+1/2*sin(f*x+e))*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)*2^(1/2)*((c+d*sin(f*x+e))/(c-d))^(1/2)/a/(c-d)^2/d/(c+d)/f/(3+2*m)/(1-sin(f*x+e))^(1/2)/(c+d*sin(f*x+e))^(1/2)
```

### 3.31.2 Mathematica [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx = \int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx$$

input `Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(5/2),x]`

output `Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(5/2), x]`

### 3.31.3 Rubi [A] (verified)

Time = 1.41 (sec) , antiderivative size = 478, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.234$ , Rules used = {3042, 3522, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2)}{(c + d \sin(e + fx))^{5/2}} dx \\ & \quad \downarrow \text{3522} \\ & \frac{2 \cos(e + fx) (Ad^2 - Bcd + c^2C) (a \sin(e + fx) + a)^m}{3df (c^2 - d^2) (c + d \sin(e + fx))^{3/2}} - \\ & \frac{2 \int - \frac{(\sin(e+fx)a+a)^m (a((cC-Bd)(3d-2cm)+Ad(3c-2dm))+a(2C(m+1)c^2+Bd(1-2m)c-d^2(-2mA+A+3C)) \sin(e+fx))}{2(c+d \sin(e+fx))^{3/2}} dx}{3ad (c^2 - d^2)} dx \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{(\sin(e+fx)a+a)^m (a((cC-Bd)(3d-2cm)+Ad(3c-2dm))+a(2C(m+1)c^2+Bd(1-2m)c-d^2(-2mA+A+3C)) \sin(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx}{3ad (c^2 - d^2)} + \\ & \frac{2 \cos(e + fx) (Ad^2 - Bcd + c^2C) (a \sin(e + fx) + a)^m}{3df (c^2 - d^2) (c + d \sin(e + fx))^{3/2}} \end{aligned}$$

---

3.31.  $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{5/2}} dx$

↓ 3042

$$\frac{\int \frac{(\sin(e+fx)a+a)^m (a((cC-Bd)(3d-2cm)+Ad(3c-2dm))+a(2C(m+1)c^2+Bd(1-2m)c-d^2(-2mA+A+3C)) \sin(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx}{3ad(c^2-d^2)} + \frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{3df(c^2-d^2)(c+d \sin(e+fx))^{3/2}}$$

↓ 3466

$$\frac{a(cd(3A+4Bm-B+3C)+d^2(-4Am+A-3B+3C)-2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{(c+d \sin(e+fx))^{3/2}} dx + (-d^2(-2Am))}{3ad(c^2-d^2)} + \frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{3df(c^2-d^2)(c+d \sin(e+fx))^{3/2}}$$

↓ 3042

$$\frac{a(cd(3A+4Bm-B+3C)+d^2(-4Am+A-3B+3C)-2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{(c+d \sin(e+fx))^{3/2}} dx + (-d^2(-2Am))}{3ad(c^2-d^2)} + \frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{3df(c^2-d^2)(c+d \sin(e+fx))^{3/2}}$$

↓ 3267

$$\frac{a^3 \cos(e+fx) (cd(3A+4Bm-B+3C)+d^2(-4Am+A-3B+3C)-2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{a-a \sin(e+fx)}(c+d \sin(e+fx))^{3/2}} d \sin(e+fx) + \frac{a^2 \cos(e+fx)}{3ad(c^2-d^2)}}{f \sqrt{a-a \sin(e+fx)} \sqrt{a \sin(e+fx)+a}} + \frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{3df(c^2-d^2)(c+d \sin(e+fx))^{3/2}}$$

↓ 157

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (cd(3A+4Bm-B+3C)+d^2(-4Am+A-3B+3C)-2c^2(2Cm+C)) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}(c+d \sin(e+fx))^{3/2}} d \sin(e+fx) + \frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{3ad(c^2-d^2)}}{\sqrt{2}f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}}$$

↓ 27

---

3.31.  $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{5/2}} dx$

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (cd(3A+4Bm-B+3C)+d^2(-4Am+A-3B+3C)-2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}(c+d \sin(e+fx))^{3/2}} d \sin(e+fx)}{f(a-a \sin(e+fx))\sqrt{a \sin(e+fx)+a}} +$$

$$3ad(c^2-d^2)$$

$$\frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{3df(c^2-d^2)(c+d \sin(e+fx))^{3/2}}$$

↓ 156

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (cd(3A+4Bm-B+3C)+d^2(-4Am+A-3B+3C)-2c^2(2Cm+C)) \sqrt{\frac{c+d \sin(e+fx)}{c-d}} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}\left(\frac{c}{c-d} + \frac{d \sin(e+fx)}{c-d}\right)}}{f(c-d)(a-a \sin(e+fx))\sqrt{a \sin(e+fx)+a}\sqrt{c+d \sin(e+fx)}}$$

$$\frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{3df(c^2-d^2)(c+d \sin(e+fx))^{3/2}}$$

↓ 155

$$\frac{\sqrt{2}a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (a \sin(e+fx)+a)^m (cd(3A+4Bm-B+3C)+d^2(-4Am+A-3B+3C)-2c^2(2Cm+C)) \sqrt{\frac{c+d \sin(e+fx)}{c-d}} \text{AppellF1}\left(r\right)}{f(2m+1)(c-d)(a-a \sin(e+fx))\sqrt{c+d \sin(e+fx)}}$$

$$\frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{3df(c^2-d^2)(c+d \sin(e+fx))^{3/2}}$$

input `Int[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(5/2),x]`

output `(2*(c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(3*d*(c^2 - d^2)*f*(c + d*Sin[e + f*x])^(3/2)) + ((Sqrt[2]*a^2*(d^2*(A - 3*B + 3*C - 4*A*m) + c*d*(3*A - B + 3*C + 4*B*m) - 2*c^2*(C + 2*C*m))*AppellF1[1/2 + m, 1/2, 3/2, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[(c + d*Sin[e + f*x])/(c - d)])/((c - d)*f*(1 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]) + (Sqrt[2]*a*(B*c*d*(1 - 2*m) + 2*c^2*C*(1 + m) - d^2*(A + 3*C - 2*A*m))*AppellF1[3/2 + m, 1/2, 3/2, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[(c + d*Sin[e + f*x])/(c - d)])/((c - d)*f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]))/(3*a*d*(c^2 - d^2))`

---

3.31.  $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{5/2}} dx$



## 3.31.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p)*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

---

3.31. 
$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{5/2}} dx$$

rule 3267 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]])*Sqrt[a - b*Sin[e + f*x]]) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]`

rule 3466 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]`

rule 3522 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(a*d*m + b*c*(n + 1)) + (c*C - B*d)*(a*c*m + b*d*(n + 1)) + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && (LtQ[n, -1] || EqQ[m + n + 2, 0])`

### 3.31.4 Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + B \sin(fx + e) + C(\sin^2(fx + e)))}{(c + d \sin(fx + e))^{\frac{5}{2}}} dx$$

input `int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(5/2),x)`

output `int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(5/2),x)`

**3.31.5 Fricas [F]**

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin(fx + e))^2 + B \sin(fx + e) + A}{(d \sin(fx + e) + c)^{5/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(5/2),x, algorithm="fricas")`

output `integral((C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(3*c*d^2*cos(f*x + e)^2 - c^3 - 3*c*d^2 + (d^3*cos(f*x + e)^2 - 3*c^2*d - d^3)*sin(f*x + e)), x)`

**3.31.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(A+B*sin(f*x+e)+C*sin(f*x+e)**2)/(c+d*sin(f*x+e))**(5/2),x)`

output `Timed out`

**3.31.7 Maxima [F]**

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin(fx + e))^2 + B \sin(fx + e) + A}{(d \sin(fx + e) + c)^{5/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(5/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/(d*sin(f*x + e) + c)^(5/2), x)`

---

3.31.  $\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx$

**3.31.8 Giac [F]**

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin(fx + e))^2 + B \sin(fx + e) + A}{(d \sin(fx + e) + c)^{5/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/(d*sin(f*x + e) + c)^(5/2), x)`

**3.31.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{5/2}} dx = \int \frac{(a + a \sin(e + fx))^m (C \sin(e + fx) + B + A)}{(c + d \sin(e + fx))^{5/2}} dx$$

input `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c + d*sin(e + f*x))^(5/2),x)`

output `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c + d*sin(e + f*x))^(5/2), x)`

### 3.32 $\int (a+b \sin(c+dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)$

3.32.1	Optimal result . . . . .	284
3.32.2	Mathematica [A] (verified) . . . . .	284
3.32.3	Rubi [A] (verified) . . . . .	285
3.32.4	Maple [A] (verified) . . . . .	286
3.32.5	Fricas [A] (verification not implemented) . . . . .	287
3.32.6	Sympy [B] (verification not implemented) . . . . .	287
3.32.7	Maxima [A] (verification not implemented) . . . . .	288
3.32.8	Giac [A] (verification not implemented) . . . . .	288
3.32.9	Mupad [B] (verification not implemented) . . . . .	289

#### 3.32.1 Optimal result

Integrand size = 31, antiderivative size = 81

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

$$= \frac{1}{2}(bB + a(2A + C))x - \frac{(Ab + aB + bC) \cos(c + dx)}{d}$$

$$+ \frac{bC \cos^3(c + dx)}{3d} - \frac{(bB + aC) \cos(c + dx) \sin(c + dx)}{2d}$$

output `1/2*(b*B+a*(2*A+C))*x-(A*b+B*a+C*b)*cos(d*x+c)/d+1/3*b*C*cos(d*x+c)^3/d-1/2*(B*b+C*a)*cos(d*x+c)*sin(d*x+c)/d`

#### 3.32.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.14

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

$$= \frac{6bBc + 6acC + 12aAdx + 6bBdx + 6aCdx - 3(4Ab + 4aB + 3bC) \cos(c + dx) + bC \cos(3(c + dx)) - 3bC \sin(3(c + dx))}{12d}$$

input `Integrate[(a + b*Sin[c + d*x])*(A + B*Sin[c + d*x] + C*Sin[c + d*x]^2),x]`

output `(6*b*B*c + 6*a*c*C + 12*a*A*d*x + 6*b*B*d*x + 6*a*C*d*x - 3*(4*A*b + 4*a*B + 3*b*C)*Cos[c + d*x] + b*C*Cos[3*(c + d*x)] - 3*b*B*Sin[2*(c + d*x)] - 3*a*C*Sin[2*(c + d*x)])/(12*d)`

**3.32.3 Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.46, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.129$ , Rules used = {3042, 3502, 3042, 3213}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin(c + dx)^2) dx$$

$$\downarrow \text{3502}$$

$$\frac{\int (a + b \sin(c + dx))(b(3A + 2C) + (3bB - aC) \sin(c + dx)) dx}{3b} - \frac{C \cos(c + dx)(a + b \sin(c + dx))^2}{3bd}$$

$$\downarrow \text{3042}$$

$$\frac{\int (a + b \sin(c + dx))(b(3A + 2C) + (3bB - aC) \sin(c + dx)) dx}{3b} - \frac{C \cos(c + dx)(a + b \sin(c + dx))^2}{3bd}$$

$$\downarrow \text{3213}$$

$$\frac{-\frac{\cos(c+dx)(a(3bB-aC)+b^2(3A+2C))}{d} + \frac{3}{2}bx(a(2A+C) + bB) - \frac{b(3bB-aC) \sin(c+dx) \cos(c+dx)}{2d}}{\frac{3b}{3bd} C \cos(c + dx)(a + b \sin(c + dx))^2}$$

input `Int[(a + b*Sin[c + d*x])*(A + B*Sin[c + d*x] + C*Sin[c + d*x]^2),x]`

output `-1/3*(C*Cos[c + d*x]*(a + b*Sin[c + d*x])^2)/(b*d) + ((3*b*(b*B + a*(2*A + C))*x)/2 - ((b^2*(3*A + 2*C) + a*(3*b*B - a*C))*Cos[c + d*x])/d - (b*(3*b*B - a*C)*Cos[c + d*x]*Sin[c + d*x])/(2*d))/(3*b)`

### 3.32.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3213 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(2*a*c + b*d)*(x/2), x] + (-Simp[(b*c + a*d)*(Cos[e + f*x]/f), x] - Simp[b*d*Cos[e + f*x]*(Sin[e + f*x]/(2*f)), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*SIN[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*SIN[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

### 3.32.4 Maple [A] (verified)

Time = 1.13 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.99

method	result
parts	$xaA - \frac{(Ab+Ba)\cos(dx+c)}{d} + \frac{(Bb+aC)\left(-\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d} - \frac{Cb(2+\sin^2(dx+c))\cos(dx+c)}{3d}$
parallelrisch	$\frac{(-3Bb-3aC)\sin(2dx+2c)+bC\cos(3dx+3c)+((-12A-9C)b-12Ba)\cos(dx+c)+(6dxB-12A-8C)b+12a(dxA+\frac{1}{2}Cd)}{12d}$
risch	$xaA + \frac{x B b}{2} + \frac{x a C}{2} - \frac{\cos(dx+c)Ab}{d} - \frac{\cos(dx+c)Ba}{d} - \frac{3\cos(dx+c)Cb}{4d} + \frac{bC\cos(3dx+3c)}{12d} - \frac{\sin(2dx+2c)B}{4d}$
derivativedivides	$-\frac{Cb(2+\sin^2(dx+c))\cos(dx+c)}{3} + Bb\left(-\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) + aC\left(-\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) - Ab\cos(dx+c) - \frac{\cos(dx+c)\sin(dx+c)}{2}$
default	$-\frac{Cb(2+\sin^2(dx+c))\cos(dx+c)}{3} + Bb\left(-\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) + aC\left(-\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) - Ab\cos(dx+c) - \frac{\cos(dx+c)\sin(dx+c)}{2}$
norman	$\frac{(aA+\frac{1}{2}Bb+\frac{1}{2}aC)x+(aA+\frac{1}{2}Bb+\frac{1}{2}aC)x\left(\tan^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+(3aA+\frac{3}{2}Bb+\frac{3}{2}aC)x\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+(3aA+\frac{3}{2}Bb+\frac{3}{2}aC)x}{d}$

input `int((a+b*sin(d*x+c))*(A+B*sin(d*x+c)+C*sin(d*x+c)^2),x,method=_RETURNVERBOSE)`

$$3.32. \int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

output `x*a*A-(A*b+B*a)/d*cos(d*x+c)+(B*b+C*a)/d*(-1/2*cos(d*x+c)*sin(d*x+c)+1/2*d*x+1/2*c)-1/3*C*b/d*(2+sin(d*x+c)^2)*cos(d*x+c)`

### 3.32.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.88

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

$$= \frac{2Cb \cos(dx + c)^3 + 3((2A + C)a + Bb)dx - 3(Ca + Bb) \cos(dx + c) \sin(dx + c) - 6(Ba + (A + C)b)}{6d}$$

input `integrate((a+b*sin(d*x+c))*(A+B*sin(d*x+c)+C*sin(d*x+c)^2),x, algorithm="fricas")`

output `1/6*(2*C*b*cos(d*x + c)^3 + 3*((2*A + C)*a + B*b)*d*x - 3*(C*a + B*b)*cos(d*x + c)*sin(d*x + c) - 6*(B*a + (A + C)*b)*cos(d*x + c))/d`

### 3.32.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 189 vs.  $2(73) = 146$ .

Time = 0.14 (sec) , antiderivative size = 189, normalized size of antiderivative = 2.33

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

$$= \begin{cases} Aax - \frac{Ab \cos(c+dx)}{d} - \frac{Ba \cos(c+dx)}{d} + \frac{Bbx \sin^2(c+dx)}{2} + \frac{Bbx \cos^2(c+dx)}{2} - \frac{Bb \sin(c+dx) \cos(c+dx)}{2d} + \frac{Cax \sin^2(c+dx)}{2} + \frac{C}{2} \\ x(a + b \sin(c)) (A + B \sin(c) + C \sin^2(c)) \end{cases}$$

input `integrate((a+b*sin(d*x+c))*(A+B*sin(d*x+c)+C*sin(d*x+c)**2),x)`

output `Piecewise((A*a*x - A*b*cos(c + d*x)/d - B*a*cos(c + d*x)/d + B*b*x*sin(c + d*x)**2/2 + B*b*x*cos(c + d*x)**2/2 - B*b*sin(c + d*x)*cos(c + d*x)/(2*d) + C*a*x*sin(c + d*x)**2/2 + C*a*x*cos(c + d*x)**2/2 - C*a*sin(c + d*x)*cos(c + d*x)/(2*d) - C*b*sin(c + d*x)**2*cos(c + d*x)/d - 2*C*b*cos(c + d*x)**3/(3*d), Ne(d, 0)), (x*(a + b*sin(c))*(A + B*sin(c) + C*sin(c)**2), True))`



**3.32.7 Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.26

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

$$= \frac{12(dx + c)Aa + 3(2dx + 2c - \sin(2dx + 2c))Ca + 3(2dx + 2c - \sin(2dx + 2c))Bb + 4(\cos(dx + c) - \cos(3dx + 3c))Cb + (4Ba + 4Ab + 3Cb)\cos(dx + c) - (Ca + Bb)\sin(2dx + 2c)}{12d}$$

input `integrate((a+b*sin(d*x+c))*(A+B*sin(d*x+c)+C*sin(d*x+c)^2),x, algorithm="maxima")`

output `1/12*(12*(d*x + c)*A*a + 3*(2*d*x + 2*c - sin(2*d*x + 2*c))*C*a + 3*(2*d*x + 2*c - sin(2*d*x + 2*c))*B*b + 4*(cos(d*x + c)^3 - 3*cos(d*x + c))*C*b - 12*B*a*cos(d*x + c) - 12*A*b*cos(d*x + c))/d`

**3.32.8 Giac [A] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.94

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

$$= \frac{1}{2} (2Aa + Ca + Bb)x + \frac{Cb \cos(3dx + 3c)}{12d} - \frac{(4Ba + 4Ab + 3Cb)\cos(dx + c)}{4d} - \frac{(Ca + Bb)\sin(2dx + 2c)}{4d}$$

input `integrate((a+b*sin(d*x+c))*(A+B*sin(d*x+c)+C*sin(d*x+c)^2),x, algorithm="giac")`

output `1/2*(2*A*a + C*a + B*b)*x + 1/12*C*b*cos(3*d*x + 3*c)/d - 1/4*(4*B*a + 4*A*b + 3*C*b)*cos(d*x + c)/d - 1/4*(C*a + B*b)*sin(2*d*x + 2*c)/d`

**3.32.9 Mupad [B] (verification not implemented)**

Time = 14.10 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.15

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx =$$

$$\frac{6Ab \cos(c + dx) + 6Ba \cos(c + dx) + \frac{9Cb \cos(c+dx)}{2} - \frac{Cb \cos(3c+3dx)}{2} + \frac{3Bb \sin(2c+2dx)}{2} + \frac{3Ca \sin(2c+2dx)}{2}}{6d}$$

input `int((a + b*sin(c + d*x))*(A + B*sin(c + d*x) + C*sin(c + d*x)^2),x)`

output `-(6*A*b*cos(c + d*x) + 6*B*a*cos(c + d*x) + (9*C*b*cos(c + d*x))/2 - (C*b*cos(3*c + 3*d*x))/2 + (3*B*b*sin(2*c + 2*d*x))/2 + (3*C*a*sin(2*c + 2*d*x))/2 - 6*A*a*d*x - 3*B*b*d*x - 3*C*a*d*x)/(6*d)`

**3.33** 
$$\int \frac{(a+b \sin(e+fx))(A+B \sin(e+fx)+C \sin^2(e+fx))}{\sin^{\frac{3}{2}}(e+fx)} dx$$

3.33.1 Optimal result . . . . . 290  
 3.33.2 Mathematica [A] (verified) . . . . . 291  
 3.33.3 Rubi [A] (verified) . . . . . 291  
 3.33.4 Maple [B] (verified) . . . . . 294  
 3.33.5 Fracas [C] (verification not implemented) . . . . . 295  
 3.33.6 Sympy [F] . . . . . 296  
 3.33.7 Maxima [F] . . . . . 296  
 3.33.8 Giac [F] . . . . . 296  
 3.33.9 Mupad [B] (verification not implemented) . . . . . 297

**3.33.1 Optimal result**

Integrand size = 41, antiderivative size = 117

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

$$= \frac{2(bB - a(A - C))E(\frac{1}{2}(e - \frac{\pi}{2} + fx) | 2)}{f}$$

$$+ \frac{2(3Ab + 3aB + bC) \text{EllipticF}(\frac{1}{2}(e - \frac{\pi}{2} + fx), 2)}{3f}$$

$$- \frac{2aA \cos(e + fx)}{f \sqrt{\sin(e + fx)}} - \frac{2bC \cos(e + fx) \sqrt{\sin(e + fx)}}{3f}$$

```
output -2*(b*B-a*(A-C))*(sin(1/2*e+1/4*Pi+1/2*f*x)^2)^(1/2)/sin(1/2*e+1/4*Pi+1/2*
f*x)*EllipticE(cos(1/2*e+1/4*Pi+1/2*f*x), 2^(1/2))/f-2/3*(3*A*b+3*B*a+C*b)*
(sin(1/2*e+1/4*Pi+1/2*f*x)^2)^(1/2)/sin(1/2*e+1/4*Pi+1/2*f*x)*EllipticF(co
s(1/2*e+1/4*Pi+1/2*f*x), 2^(1/2))/f-2*a*A*cos(f*x+e)/f/sin(f*x+e)^(1/2)-2/3
*b*C*cos(f*x+e)*sin(f*x+e)^(1/2)/f
```

### 3.33.2 Mathematica [A] (verified)

Time = 1.57 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.83

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx =$$

$$\frac{6(bB + a(-A + C))E\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) + 2(3Ab + 3aB + bC) \operatorname{EllipticF}\left(\frac{1}{4}(-2e + \pi - 2fx), 2\right)}{3f}$$

input `Integrate[((a + b*Sin[e + f*x])*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/Sin[e + f*x]^(3/2), x]`

output `-1/3*(6*(b*B + a*(-A + C))*EllipticE[(-2*e + Pi - 2*f*x)/4, 2] + 2*(3*A*b + 3*a*B + b*C)*EllipticF[(-2*e + Pi - 2*f*x)/4, 2] + (2*Cos[e + f*x]*(3*a*A + b*C*Sin[e + f*x]))/Sqrt[Sin[e + f*x]])/f`

### 3.33.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.268$ , Rules used = {3042, 3510, 27, 3042, 3502, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin(e + fx)^2)}{\sin(e + fx)^{3/2}} dx$$

$$\downarrow \text{3510}$$

$$-2 \int \frac{bC \sin^2(e + fx) + (bB - a(A - C)) \sin(e + fx) + Ab + aB}{2\sqrt{\sin(e + fx)}} dx - \frac{2aA \cos(e + fx)}{f\sqrt{\sin(e + fx)}}$$

$$\downarrow \text{27}$$

$$\int \frac{bC \sin^2(e + fx) + (bB - a(A - C)) \sin(e + fx) + Ab + aB}{\sqrt{\sin(e + fx)}} dx - \frac{2aA \cos(e + fx)}{f\sqrt{\sin(e + fx)}}$$

---

3.33.  $\int \frac{(a+b \sin(e+fx))(A+B \sin(e+fx)+C \sin^2(e+fx))}{\sin^{\frac{3}{2}}(e+fx)} dx$

$$\begin{aligned}
& \int \frac{bC \sin(e+fx)^2 + (bB - a(A-C)) \sin(e+fx) + Ab + aB}{\sqrt{\sin(e+fx)}} dx - \frac{2aA \cos(e+fx)}{f \sqrt{\sin(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{2}{3} \int \frac{3Ab + Cb + 3aB + 3(bB - a(A-C)) \sin(e+fx)}{2\sqrt{\sin(e+fx)}} dx - \frac{2aA \cos(e+fx)}{f \sqrt{\sin(e+fx)}} - \\
& \quad \frac{2bC \sqrt{\sin(e+fx)} \cos(e+fx)}{3f} \\
& \quad \downarrow \text{27} \\
& \frac{1}{3} \int \frac{3Ab + Cb + 3aB + 3(bB - a(A-C)) \sin(e+fx)}{\sqrt{\sin(e+fx)}} dx - \frac{2aA \cos(e+fx)}{f \sqrt{\sin(e+fx)}} - \\
& \quad \frac{2bC \sqrt{\sin(e+fx)} \cos(e+fx)}{3f} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \int \frac{3Ab + Cb + 3aB + 3(bB - a(A-C)) \sin(e+fx)}{\sqrt{\sin(e+fx)}} dx - \frac{2aA \cos(e+fx)}{f \sqrt{\sin(e+fx)}} - \\
& \quad \frac{2bC \sqrt{\sin(e+fx)} \cos(e+fx)}{3f} \\
& \quad \downarrow \text{3227} \\
& \frac{1}{3} \left( (3aB + 3Ab + bC) \int \frac{1}{\sqrt{\sin(e+fx)}} dx + 3(bB - a(A-C)) \int \sqrt{\sin(e+fx)} dx \right) - \\
& \quad \frac{2aA \cos(e+fx)}{f \sqrt{\sin(e+fx)}} - \frac{2bC \sqrt{\sin(e+fx)} \cos(e+fx)}{3f} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left( (3aB + 3Ab + bC) \int \frac{1}{\sqrt{\sin(e+fx)}} dx + 3(bB - a(A-C)) \int \sqrt{\sin(e+fx)} dx \right) - \\
& \quad \frac{2aA \cos(e+fx)}{f \sqrt{\sin(e+fx)}} - \frac{2bC \sqrt{\sin(e+fx)} \cos(e+fx)}{3f} \\
& \quad \downarrow \text{3119} \\
& \frac{1}{3} \left( (3aB + 3Ab + bC) \int \frac{1}{\sqrt{\sin(e+fx)}} dx + \frac{6E\left(\frac{1}{2}(e+fx - \frac{\pi}{2}) \mid 2\right) (bB - a(A-C))}{f} \right) - \\
& \quad \frac{2aA \cos(e+fx)}{f \sqrt{\sin(e+fx)}} - \frac{2bC \sqrt{\sin(e+fx)} \cos(e+fx)}{3f} \\
& \quad \downarrow \text{3120}
\end{aligned}$$

---

3.33.  $\int \frac{(a+b \sin(e+fx))(A+B \sin(e+fx)+C \sin^2(e+fx))}{\sin^{\frac{3}{2}}(e+fx)} dx$

$$\frac{1}{3} \left( \frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(e+fx-\frac{\pi}{2}), 2\right) (3aB+3Ab+bC)}{f} + \frac{6E\left(\frac{1}{2}(e+fx-\frac{\pi}{2})|2\right) (bB-a(A-C))}{f} \right) - \frac{2aA \cos(e+fx)}{f\sqrt{\sin(e+fx)}} - \frac{2bC\sqrt{\sin(e+fx)} \cos(e+fx)}{3f}$$

input `Int[((a + b*Sin[e + f*x])*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/Sin[e + f*x]^(3/2), x]`

output `((6*(b*B - a*(A - C))*EllipticE[(e - Pi/2 + f*x)/2, 2])/f + (2*(3*A*b + 3*a*B + b*C)*EllipticF[(e - Pi/2 + f*x)/2, 2])/f)/3 - (2*a*A*Cos[e + f*x])/(f*Sqrt[Sin[e + f*x]]) - (2*b*C*Cos[e + f*x]*Sqrt[Sin[e + f*x]])/(3*f)`

### 3.33.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

```
rule 3510 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)]*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[(-b*c - a*d)*(A*b^2 - a*b*B + a^2*C)*Cos[
e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b^2*f*(m + 1)*(a^2 - b^2))), x] - S
imp[1/(b^2*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(
m + 1)*((b*B - a*C)*(b*c - a*d) - A*b*(a*c - b*d)) + (b*B*(a^2*d + b^2*d*(m
+ 1) - a*b*c*(m + 2)) + (b*c - a*d)*(A*b^2*(m + 2) + C*(a^2 + b^2*(m + 1)
))*Sin[e + f*x] - b*C*d*(m + 1)*(a^2 - b^2)*Sin[e + f*x]^2, x], x] /; F
reeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2,
0] && LtQ[m, -1]
```

### 3.33.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 395 vs.  $2(169) = 338$ .

Time = 2.80 (sec) , antiderivative size = 396, normalized size of antiderivative = 3.38

method	result
parts	$\frac{(Ab+Ba)\sqrt{1+\sin(fx+e)}\sqrt{2-2\sin(fx+e)}\sqrt{-\sin(fx+e)}F\left(\sqrt{1+\sin(fx+e)},\frac{\sqrt{2}}{2}\right)}{\cos(fx+e)\sqrt{\sin(fx+e)}} - \frac{(Bb+aC)\sqrt{1+\sin(fx+e)}\sqrt{2-2\sin(fx+e)}}{f}$
default	$-A\sqrt{1+\sin(fx+e)}\sqrt{2-2\sin(fx+e)}\sqrt{-\sin(fx+e)}F\left(\sqrt{1+\sin(fx+e)},\frac{\sqrt{2}}{2}\right) + Ab\sqrt{1+\sin(fx+e)}\sqrt{2-2\sin(fx+e)}\sqrt{-\sin(fx+e)}$

```
input int((a+b*sin(f*x+e))*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/sin(f*x+e)^(3/2),x,me
thod=_RETURNVERBOSE)
```

$$3.33. \int \frac{(a+b\sin(e+fx))(A+B\sin(e+fx)+C\sin^2(e+fx))}{\sin^{\frac{3}{2}}(e+fx)} dx$$

```
output (A*b+B*a)*(1+sin(f*x+e))^(1/2)*(2-2*sin(f*x+e))^(1/2)*(-sin(f*x+e))^(1/2)*
EllipticF((1+sin(f*x+e))^(1/2),1/2*2^(1/2))/cos(f*x+e)/sin(f*x+e)^(1/2)/f-
(B*b+C*a)*(1+sin(f*x+e))^(1/2)*(2-2*sin(f*x+e))^(1/2)*(-sin(f*x+e))^(1/2)*
(2*EllipticE((1+sin(f*x+e))^(1/2),1/2*2^(1/2))-EllipticF((1+sin(f*x+e))^(1
/2),1/2*2^(1/2)))/cos(f*x+e)/sin(f*x+e)^(1/2)/f+C*b*(1/3*(1+sin(f*x+e))^(1
/2)*(2-2*sin(f*x+e))^(1/2)*(-sin(f*x+e))^(1/2)*EllipticF((1+sin(f*x+e))^(1
/2),1/2*2^(1/2))-2/3*cos(f*x+e)^2*sin(f*x+e))/cos(f*x+e)/sin(f*x+e)^(1/2)/
f+a*A*(2*(1+sin(f*x+e))^(1/2)*(2-2*sin(f*x+e))^(1/2)*(-sin(f*x+e))^(1/2)*E
llipticE((1+sin(f*x+e))^(1/2),1/2*2^(1/2))-(1+sin(f*x+e))^(1/2)*(2-2*sin(f
*x+e))^(1/2)*(-sin(f*x+e))^(1/2)*EllipticF((1+sin(f*x+e))^(1/2),1/2*2^(1/2
))-2*cos(f*x+e)^2)/cos(f*x+e)/sin(f*x+e)^(1/2)/f
```

### 3.33.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.98

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

$$= \frac{\sqrt{2}\sqrt{-i}(3Ba + (3A + C)b) \sin(fx + e) \operatorname{weierstrassPInverse}(4, 0, \cos(fx + e) + i \sin(fx + e)) + \sqrt{2}\sqrt{i}}$$

```
input integrate((a+b*sin(f*x+e))*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/sin(f*x+e)^(3/2
),x, algorithm="fracas")
```

```
output 1/3*(sqrt(2)*sqrt(-I)*(3*B*a + (3*A + C)*b)*sin(f*x + e)*weierstrassPInver
se(4, 0, cos(f*x + e) + I*sin(f*x + e)) + sqrt(2)*sqrt(I)*(3*B*a + (3*A +
C)*b)*sin(f*x + e)*weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e)
) - 3*sqrt(2)*sqrt(-I)*(I*(A - C)*a - I*B*b)*sin(f*x + e)*weierstrassZeta(
4, 0, weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e))) - 3*sqrt(2
)*sqrt(I)*(-I*(A - C)*a + I*B*b)*sin(f*x + e)*weierstrassZeta(4, 0, weiers
trassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e))) - 2*(C*b*cos(f*x + e)*
sin(f*x + e) + 3*A*a*cos(f*x + e))*sqrt(sin(f*x + e))/(f*sin(f*x + e))
```

---

3.33.  $\int \frac{(a+b \sin(e+fx))(A+B \sin(e+fx)+C \sin^2(e+fx))}{\sin^{\frac{3}{2}}(e+fx)} dx$



**3.33.6 Sympy [F]**

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

$$= \int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

input `integrate((a+b*sin(f*x+e))*(A+B*sin(f*x+e)+C*sin(f*x+e)**2)/sin(f*x+e)**(3/2),x)`

output `Integral((a + b*sin(e + f*x))*(A + B*sin(e + f*x) + C*sin(e + f*x)**2)/sin(e + f*x)**(3/2), x)`

**3.33.7 Maxima [F]**

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

$$= \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A)(b \sin(fx + e) + a)}{\sin(fx + e)^{\frac{3}{2}}} dx$$

input `integrate((a+b*sin(f*x+e))*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/sin(f*x+e)^(3/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(b*sin(f*x + e) + a)/sin(f*x + e)^(3/2), x)`

**3.33.8 Giac [F]**

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

$$= \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A)(b \sin(fx + e) + a)}{\sin(fx + e)^{\frac{3}{2}}} dx$$

---

3.33.  $\int \frac{(a+b \sin(e+fx))(A+B \sin(e+fx)+C \sin^2(e+fx))}{\sin^{\frac{3}{2}}(e+fx)} dx$

```
input integrate((a+b*sin(f*x+e))*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/sin(f*x+e)^(3/2),x, algorithm="giac")
```

```
output integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(b*sin(f*x + e) + a)/sin(f*x + e)^(3/2), x)
```

### 3.33.9 Mupad [B] (verification not implemented)

Time = 15.72 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.44

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

$$= \frac{2 B b E\left(\frac{e}{2} - \frac{\pi}{4} + \frac{fx}{2} \mid 2\right)}{f} - \frac{2 B a F\left(\frac{\pi}{4} - \frac{e}{2} - \frac{fx}{2} \mid 2\right)}{f}$$

$$- \frac{2 A b F\left(\frac{\pi}{4} - \frac{e}{2} - \frac{fx}{2} \mid 2\right)}{f} + \frac{2 C a E\left(\frac{e}{2} - \frac{\pi}{4} + \frac{fx}{2} \mid 2\right)}{f}$$

$$- \frac{A a \cos(e + fx) (\sin(e + fx)^2)^{1/4} {}_2F_1\left(\frac{1}{2}, \frac{5}{4}; \frac{3}{2}; \cos(e + fx)^2\right)}{f \sqrt{\sin(e + fx)}}$$

$$- \frac{C b \cos(e + fx) \sin(e + fx)^{5/2} {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{2}; \cos(e + fx)^2\right)}{f (\sin(e + fx)^2)^{5/4}}$$

```
input int(((a + b*sin(e + f*x))*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/sin(e + f*x)^(3/2),x)
```

```
output (2*B*b*ellipticE(e/2 - pi/4 + (f*x)/2, 2))/f - (2*B*a*ellipticF(pi/4 - e/2 - (f*x)/2, 2))/f - (2*A*b*ellipticF(pi/4 - e/2 - (f*x)/2, 2))/f + (2*C*a*ellipticE(e/2 - pi/4 + (f*x)/2, 2))/f - (A*a*cos(e + f*x)*(sin(e + f*x)^2)^(1/4)*hypergeom([1/2, 5/4], 3/2, cos(e + f*x)^2))/(f*sin(e + f*x)^(1/2)) - (C*b*cos(e + f*x)*sin(e + f*x)^(5/2)*hypergeom([-1/4, 1/2], 3/2, cos(e + f*x)^2))/(f*(sin(e + f*x)^2)^(5/4))
```

### 3.34 $\int (a+b \sin(e+fx))^m (c+d \sin(e+fx))^n (A + B \sin(e +$

3.34.1	Optimal result . . . . .	298
3.34.2	Mathematica [N/A] . . . . .	298
3.34.3	Rubi [N/A] . . . . .	299
3.34.4	Maple [N/A] (verified) . . . . .	300
3.34.5	Fricas [N/A] . . . . .	300
3.34.6	Sympy [F(-1)] . . . . .	300
3.34.7	Maxima [F(-1)] . . . . .	301
3.34.8	Giac [F(-1)] . . . . .	301
3.34.9	Mupad [N/A] . . . . .	302

#### 3.34.1 Optimal result

Integrand size = 45, antiderivative size = 45

$$\int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \text{Int}((a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)), x)$$

output `Unintegrable((a+b*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

#### 3.34.2 Mathematica [N/A]

Not integrable

Time = 24.92 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.04

$$\int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input `Integrate[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `Integrate[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x]`

### 3.34.3 Rubi [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {3042, 3544}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx)) (c + d \sin(e + fx))^n dx$$

↓ 3042

$$\int (a + b \sin(e + fx))^m (A + B \sin(e + fx) + C \sin(e + fx)^2) (c + d \sin(e + fx))^n dx$$

↓ 3544

$$\int (a + b \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx)) (c + d \sin(e + fx))^n dx$$

input `Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `$Aborted`

#### 3.34.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3544 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Unintegrable[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

**3.34.4 Maple [N/A] (verified)**

Not integrable

Time = 2.04 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00

$$\int (a + b \sin(fx + e))^m (c + d \sin(fx + e))^n (A + B \sin(fx + e) + C(\sin^2(fx + e))) dx$$

input `int((a+b*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2), x)`

output `int((a+b*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2), x)`

**3.34.5 Fricas [N/A]**

Not integrable

Time = 1.43 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.20

$$\begin{aligned} & \int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx \\ &= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(b \sin(fx + e) + a)^m (d \sin(fx + e) + c)^n dx \end{aligned}$$

input `integrate((a+b*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*(b*sin(f*x + e) + a)^m*(d*sin(f*x + e) + c)^n, x)`

**3.34.6 Sympy [F(-1)]**

Timed out.

$$\begin{aligned} & \int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx \\ &= \text{Timed out} \end{aligned}$$

input `integrate((a+b*sin(f*x+e))**m*(c+d*sin(f*x+e))**n*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output Timed out

### 3.34.7 Maxima [F(-1)]

Timed out.

$$\int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

= Timed out

input `integrate((a+b*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output Timed out

### 3.34.8 Giac [F(-1)]

Timed out.

$$\int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

= Timed out

input `integrate((a+b*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="giac")`

output Timed out

**3.34.9 Mupad [N/A]**

Not integrable

Time = 73.80 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.04

$$\int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (C \sin(e + fx)^2 + B \sin(e + fx) + A) dx$$

input `int((a + b*sin(e + f*x))^m*(c + d*sin(e + f*x))^n*(A + B*sin(e + f*x) + C*  
sin(e + f*x)^2),x)`

output `int((a + b*sin(e + f*x))^m*(c + d*sin(e + f*x))^n*(A + B*sin(e + f*x) + C*  
sin(e + f*x)^2), x)`

## APPENDIX

4.1 Listing of Grading functions . . . . .	303
--	-----

## 4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

### 4.1.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
```



```

(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A"," "}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A"," "}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal.$
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
      ,
      finalresult={"F","Contains unresolved integral."}
    ]
  ];

  finalresult
]

```

```

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
  If[Head[expn]===Plus || Head[expn]===Times,
    Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
  If[ElementaryFunctionQ[Head[expn]],
    Max[3,ExpnType[expn[[1]]],
  If[SpecialFunctionQ[Head[expn]],
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
  If[HypergeometricFunctionQ[Head[expn]],
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
  If[AppellFunctionQ[Head[expn]],
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
  If[Head[expn]===RootSum,
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
  If[Head[expn]===Integrate || Head[expn]===Int,
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
  9]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

### 4.1.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

```

```

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
  if is_contains_complex(result) then
    if is_contains_complex(optimal) then
      if debug then
        print("both result and optimal complex");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        return "A"," ";
      else
        return "B",cat("Both result and optimal contain complex but leaf count of
                        convert(leaf_count_result,string)," vs. $2 (" ,
                        convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_
        end if
      else #result contains complex but optimal is not
        if debug then
          print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
      fi;
    else # result do not contain complex
      # this assumes optimal do not as well. No check is needed here.
      if debug then
        print("result do not contain complex, this assumes optimal do not as well");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        if debug then
          print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
      else
        if debug then
          print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string)," $ vs. $2(",
                        convert(leaf_count_optimal,string)," )=" ,convert(2*leaf_cou
        fi;
      fi;
    fi;
  fi;

```

```

else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  end if
end if

```

```

elif type(expn, ``~`) then
  if type(op(2,expn), 'integer') then
    ExpnType(op(1,expn))
  elif type(op(2,expn), 'rational') then
    if type(op(1,expn), 'rational') then
      1
    else
      max(2, ExpnType(op(1,expn)))
    end if
  else
    max(3, ExpnType(op(1,expn)), ExpnType(op(2,expn)))
  end if
elif type(expn, ``+`) or type(expn, ``*`) then
  max(ExpnType(op(1,expn)), max(ExpnType(rest(expn))))
elif ElementaryFunctionQ(op(0,expn)) then
  max(3, ExpnType(op(1,expn)))
elif SpecialFunctionQ(op(0,expn)) then
  max(4, apply(max, map(ExpnType, [op(expn)])))
elif HypergeometricFunctionQ(op(0,expn)) then
  max(5, apply(max, map(ExpnType, [op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6, apply(max, map(ExpnType, [op(expn)])))
elif op(0,expn)='int' then
  max(8, apply(max, map(ExpnType, [op(expn)]))) else
  9
end if
end proc:

```

```

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

```

```

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,

```

```

        GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
        EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
    member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
    member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
    if nops(u)=2 then
        op(2,u)
    else
        apply(op(0,u),op(2..nops(u),u))
    end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma] [LeafCount] (u);
end proc:

```

### 4.1.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):

```



```
if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
        return True
    else:
        return False
else:
    return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
```

```

return 1
elif isinstance(expn,list):
    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
        return 1
    else:
        return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
elif isinstance(expn,Pow): #type(expn,``^`)
    if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
        return expnType(expn.args[0]) #ExpnType(op(1,expn))
    elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or type(expn,``*`)
    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

*#main function*

```

def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

    #print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```

#### 4.1.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
```

```

        return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func ," is special_function")
        else:
            print ("func ", func ," is NOT special_function")

    return m

```

```

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric', 'hypergeometric_M', 'hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=", expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception, AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list:  #isinstance(expn,list):

```

```

    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
elif expn.operator() == operator.pow: #isinstance(expn,Pow)
    if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
        return expnType(expn.operands()[0]) #expnType(expn.args[0])
    elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
        if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or inst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

```

```

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)",type(result))
    print("type(optimal)",type(optimal))

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal." + str(leaf
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```



```
return grade, grade_annotation
```