

# Computer Algebra Independent Integration Tests

Summer 2023 edition with Rubi V 4.17.3

4-Trig-functions/4.2-Cosine/84-4.2.1.1-a+b-cos-<sup>n</sup>

Nasser M. Abbasi

December 8, 2023

Compiled on December 8, 2023 at 8:44pm

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>detailed summary tables of results</b>	<b>20</b>
<b>3</b>	<b>Listing of integrals</b>	<b>43</b>
<b>4</b>	<b>Appendix</b>	<b>410</b>

# CHAPTER 1

## INTRODUCTION

1.1	Listing of CAS systems tested . . . . .	3
1.2	Results . . . . .	4
1.3	Time and leaf size Performance . . . . .	7
1.4	Performance based on number of rules Rubi used . . . . .	9
1.5	Performance based on number of steps Rubi used . . . . .	10
1.6	Solved integrals histogram based on leaf size of result . . . . .	11
1.7	Solved integrals histogram based on CPU time used . . . . .	12
1.8	Leaf size vs. CPU time used . . . . .	13
1.9	list of integrals with no known antiderivative . . . . .	14
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	14
1.11	list of integrals solved by CAS but failed verification . . . . .	14
1.12	Timing . . . . .	15
1.13	Verification . . . . .	15
1.14	Important notes about some of the results . . . . .	15
1.15	Design of the test system . . . . .	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [ 62 ]. This is test number [ 84 ].

## 1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 13.3.1 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

## 1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 ( 62 )	0.00 ( 0 )
Mathematica	100.00 ( 62 )	0.00 ( 0 )
Maple	72.58 ( 45 )	27.42 ( 17 )
Fricas	72.58 ( 45 )	27.42 ( 17 )
Giac	62.90 ( 39 )	37.10 ( 23 )
Maxima	62.90 ( 39 )	37.10 ( 23 )
Mupad	56.45 ( 35 )	43.55 ( 27 )
Sympy	51.61 ( 32 )	48.39 ( 30 )

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> <li>1. antiderivative contains a hypergeometric function and the optimal antiderivative does not.</li> <li>2. antiderivative contains a special function and the optimal antiderivative does not.</li> <li>3. antiderivative contains the imaginary unit and the optimal antiderivative does not.</li> </ol>
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	90.323	9.677	0.000	0.000
Maple	62.903	6.452	3.226	27.419
Giac	61.290	1.613	0.000	37.097
Fricas	59.677	3.226	9.677	27.419
Maxima	58.065	4.839	0.000	37.097
Sympy	12.903	19.355	19.355	48.387
Mupad	0.000	56.452	0.000	43.548

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima

and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	17	100.00	0.00	0.00
Maple	17	100.00	0.00	0.00
Giac	23	100.00	0.00	0.00
Maxima	23	100.00	0.00	0.00
Mupad	27	0.00	100.00	0.00
Sympy	30	96.67	3.33	0.00

Table 1.4: Failure statistics for each CAS

## 1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.



System	Mean time (sec)
Fricas	0.24
Mathematica	0.25
Giac	0.29
Rubi	0.33
Maxima	0.76
Maple	1.11
Sympy	1.30
Mupad	10.54

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	65.26	0.85	67.00	0.90
Giac	70.13	0.88	75.00	0.83
Rubi	80.58	0.92	75.50	1.00
Maple	103.73	1.03	71.00	0.77
Mathematica	104.81	1.12	69.00	1.00
Fricas	135.69	1.41	90.00	1.11
Sympy	346.69	3.51	299.50	3.76
Maxima	2655.69	26.49	94.00	1.19

Table 1.6: Leaf size performance for each CAS

## 1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the  $y$  axis is the percentage solved which Rubi itself needed the number of rules given the  $x$  axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

# 1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

## 1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

## 1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

## 1.8 Leaf size vs. CPU time used

The following gives the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time.




Figure 1.5: Leaf size vs. CPU time. Full range

## 1.9 list of integrals with no known antiderivative

{}

## 1.10 List of integrals solved by CAS but has no known antiderivative

**Rubi** {}

**Mathematica** {}

**Maple** {}

**Maxima** {}

**Fricas** {}

**Sympy** {}

**Giac** {}

**Mupad** {}

## 1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

**Rubi** {}

**Mathematica** {}

**Maple** {5}

**Maxima** Verification phase not currently implemented.

**Fricas** Verification phase not currently implemented.

**Sympy** Verification phase not currently implemented.

**Giac** Verification phase not currently implemented.

**Mupad** Verification phase not currently implemented.

## 1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

## 1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

## 1.14 Important notes about some of the results

### 1.14.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.



The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

### 1.14.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

### 1.14.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at [https://ask.sagemath.org/question/57123/could-we-have-a-leaf\\_count-function-in-base-sagemath/](https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/)

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

#### 1.14.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives  $\sin(x)^2/2$

## 1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer, 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer, Leaf size of result.
4. integer, Leaf size of the optimal antiderivative.
5. number, CPU time used to solve this integral. 0 if failed.
6. string, The integral in Latex format
7. string, The input used in CAS own syntax.
8. string, The result (antiderivative) produced by CAS in Latex format
9. string, The optimal antiderivative in Latex format.
10. integer, 0 or 1. Indicates if problem has known antiderivative or not
11. String, The result (antiderivative) in CAS own syntax.
12. String, The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String, Small string description of why the grade was given.
14. integer, 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

*The following fields are present only in Rubi Table file*

15. integer, Number of steps used.
16. integer, Number of rules used.
17. integer, Integrand leaf size.
18. real number, Ratio, Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String, The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi  
June 27, 2013  
Design v0.01

# CHAPTER 2

## DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS . . . . .	21
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	24
2.3	Detailed conclusion table specific for Rubi results . . . . .	40

## 2.1 List of integrals sorted by grade for each CAS

2.1.1	Rubi . . . . .	21
2.1.2	Mma . . . . .	21
2.1.3	Maple . . . . .	22
2.1.4	Fricas . . . . .	22
2.1.5	Maxima . . . . .	22
2.1.6	Giac . . . . .	23
2.1.7	Mupad . . . . .	23
2.1.8	Sympy . . . . .	23

### 2.1.1 Rubi

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62 }

**B grade** { }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

### 2.1.2 Mma

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 38, 39, 40, 42, 43, 44, 46, 47, 48, 50, 51, 52, 53, 54, 55, 57, 58, 59, 60, 62 }

**B grade** { 37, 41, 45, 49, 56, 61 }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

### 2.1.3 Maple

**A grade** { 1, 2, 3, 4, 7, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 54, 55 }

**B grade** { 6, 50, 51, 52 }

**C grade** { 5, 53 }

**F normal fail** { 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 56, 57, 58, 59, 60, 61, 62 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

### 2.1.4 Fricas

**A grade** { 1, 2, 3, 4, 5, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49 }

**B grade** { 6, 7 }

**C grade** { 50, 51, 52, 53, 54, 55 }

**F normal fail** { 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 56, 57, 58, 59, 60, 61, 62 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

### 2.1.5 Maxima

**A grade** { 1, 2, 3, 4, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49 }

**B grade** { 5, 6, 7 }

**C grade** { }

**F normal fail** { 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

### 2.1.6 Giac

**A grade** { 1, 2, 3, 4, 6, 7, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49 }

**B grade** { 5 }

**C grade** { }

**F normal fail** { 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

### 2.1.7 Mupad

**A grade** { }

**B grade** { 4, 5, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 53 }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { 1, 2, 3, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 50, 51, 52, 54, 55, 56, 57, 58, 59, 60, 61, 62 }

**F(-2) exception fail** { }

### 2.1.8 Sympy

**A grade** { 18, 22, 26, 30, 34, 38, 42, 46 }

**B grade** { 35, 36, 37, 39, 40, 41, 43, 44, 45, 47, 48, 49 }

**C grade** { 19, 20, 21, 23, 24, 25, 27, 28, 29, 31, 32, 33 }

**F normal fail** { 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62 }

**F(-1) timedout fail** { 1 }

**F(-2) exception fail** { }



## 2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column N.S. means **normalized size** and is defined as  $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$ . To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F(-1)</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	119	127	83	86	77	75	0	108	0
N.S.	1	1.07	0.70	0.72	0.65	0.63	0.00	0.91	0.00
time (sec)	N/A	0.443	0.273	0.779	0.465	0.247	0.000	0.422	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	93	71	73	60	62	0	84	0
N.S.	1	1.04	0.80	0.82	0.67	0.70	0.00	0.94	0.00
time (sec)	N/A	0.348	0.126	0.808	0.497	0.272	0.000	0.317	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	59	55	58	38	44	0	55	0
N.S.	1	1.00	0.93	0.98	0.64	0.75	0.00	0.93	0.00
time (sec)	N/A	0.255	0.072	0.810	0.469	0.251	0.000	0.278	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	29	43	20	32	0	30	33
N.S.	1	1.00	1.12	1.65	0.77	1.23	0.00	1.15	1.27
time (sec)	N/A	0.180	0.031	0.800	0.436	0.248	0.000	0.265	0.003

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	B	A	<b>F</b>	B	B
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	46	46	40	56	90	126	0	93	45
N.S.	1	1.00	0.87	1.22	1.96	2.74	0.00	2.02	0.98
time (sec)	N/A	0.197	0.016	0.289	0.425	0.277	0.000	0.277	0.002

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	<b>F</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	77	77	63	138	15721	153	0	112	0
N.S.	1	1.00	0.82	1.79	204.17	1.99	0.00	1.45	0.00
time (sec)	N/A	0.273	0.095	1.030	1.034	0.269	0.000	0.275	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	112	65	174	84332	188	0	129	0
N.S.	1	1.05	0.61	1.63	788.15	1.76	0.00	1.21	0.00
time (sec)	N/A	0.374	0.211	1.099	16.466	0.284	0.000	0.332	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	67	67	69	0	0	0	0	0	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.261	0.115	0.000	0.000	0.000	0.000	0.000	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	66	66	69	0	0	0	0	0	0
N.S.	1	1.00	1.05	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.270	0.070	0.000	0.000	0.000	0.000	0.000	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	65	65	69	0	0	0	0	0	0
N.S.	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.266	0.070	0.000	0.000	0.000	0.000	0.000	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	65	65	67	0	0	0	0	0	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.263	0.061	0.000	0.000	0.000	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	65	65	67	0	0	0	0	0	0
N.S.	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.261	0.061	0.000	0.000	0.000	0.000	0.000	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	68	68	69	0	0	0	0	0	0
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.254	0.069	0.000	0.000	0.000	0.000	0.000	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	73	73	74	0	0	0	0	0	0
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.268	0.076	0.000	0.000	0.000	0.000	0.000	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	75	75	75	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.273	0.098	0.000	0.000	0.000	0.000	0.000	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	59	59	77	0	0	0	0	0	0
N.S.	1	1.00	1.31	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.195	0.097	0.000	0.000	0.000	0.000	0.000	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	60	60	74	0	0	0	0	0	0
N.S.	1	1.00	1.23	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.194	0.081	0.000	0.000	0.000	0.000	0.000	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	20	18	24	26	41	30	38
N.S.	1	1.00	0.65	0.58	0.77	0.84	1.32	0.97	1.23
time (sec)	N/A	0.177	0.048	1.033	0.362	0.253	0.373	0.266	14.619

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	56	61	43	46	68	59	187	59	67
N.S.	1	1.09	0.77	0.82	1.21	1.05	3.34	1.05	1.20
time (sec)	N/A	0.249	0.144	0.679	0.331	0.257	0.905	0.278	15.523

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	81	91	56	62	111	90	359	75	96
N.S.	1	1.12	0.69	0.77	1.37	1.11	4.43	0.93	1.19
time (sec)	N/A	0.364	0.251	0.723	0.341	0.271	1.593	0.287	15.510

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	106	121	66	75	151	121	592	88	96
N.S.	1	1.14	0.62	0.71	1.42	1.14	5.58	0.83	0.91
time (sec)	N/A	0.507	0.345	0.834	0.332	0.280	3.071	0.279	15.507

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	20	18	24	26	41	30	38
N.S.	1	1.00	0.61	0.55	0.73	0.79	1.24	0.91	1.15
time (sec)	N/A	0.181	0.050	1.195	0.373	0.267	0.379	0.269	15.494

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	58	63	43	46	69	59	192	61	67
N.S.	1	1.09	0.74	0.79	1.19	1.02	3.31	1.05	1.16
time (sec)	N/A	0.257	0.199	0.687	0.411	0.272	0.824	0.277	15.194

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	83	93	65	64	112	90	364	77	95
N.S.	1	1.12	0.78	0.77	1.35	1.08	4.39	0.93	1.14
time (sec)	N/A	0.377	0.357	0.827	0.374	0.273	1.427	0.283	15.237

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	108	123	66	77	152	121	597	90	97
N.S.	1	1.14	0.61	0.71	1.41	1.12	5.53	0.83	0.90
time (sec)	N/A	0.506	0.351	0.838	0.347	0.271	2.854	0.288	15.630

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	20	18	24	26	42	30	38
N.S.	1	1.00	0.61	0.55	0.73	0.79	1.27	0.91	1.15
time (sec)	N/A	0.185	0.037	0.593	0.346	0.267	0.368	0.268	14.440

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	58	63	43	46	69	59	192	61	67
N.S.	1	1.09	0.74	0.79	1.19	1.02	3.31	1.05	1.16
time (sec)	N/A	0.255	0.011	0.623	0.323	0.255	0.759	0.284	0.002

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	83	93	65	64	112	90	366	77	96
N.S.	1	1.12	0.78	0.77	1.35	1.08	4.41	0.93	1.16
time (sec)	N/A	0.367	0.173	0.794	0.351	0.266	1.341	0.270	14.396

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	108	123	66	77	152	121	597	90	97
N.S.	1	1.14	0.61	0.71	1.41	1.12	5.53	0.83	0.90
time (sec)	N/A	0.492	0.047	0.669	0.401	0.264	2.977	0.301	0.002

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	20	18	24	26	44	30	38
N.S.	1	1.00	0.65	0.58	0.77	0.84	1.42	0.97	1.23
time (sec)	N/A	0.179	0.045	0.558	0.391	0.286	0.371	0.291	14.507

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	56	61	43	46	68	59	190	59	67
N.S.	1	1.09	0.77	0.82	1.21	1.05	3.39	1.05	1.20
time (sec)	N/A	0.259	0.015	0.709	0.379	0.252	0.942	0.268	0.002



Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	81	91	65	62	111	90	362	75	95
N.S.	1	1.12	0.80	0.77	1.37	1.11	4.47	0.93	1.17
time (sec)	N/A	0.366	0.333	0.763	0.366	0.258	1.544	0.274	14.320

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	106	121	66	75	151	121	595	88	96
N.S.	1	1.14	0.62	0.71	1.42	1.14	5.61	0.83	0.91
time (sec)	N/A	0.491	0.014	0.821	0.331	0.264	3.153	0.270	0.003

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	22	65	33	48	46	41	34	17
N.S.	1	0.34	1.00	0.51	0.74	0.71	0.63	0.52	0.26
time (sec)	N/A	0.182	0.047	0.575	0.278	0.272	0.349	0.285	14.418

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	90	48	143	64	91	88	228	62	47
N.S.	1	0.53	1.59	0.71	1.01	0.98	2.53	0.69	0.52
time (sec)	N/A	0.262	0.147	0.632	0.272	0.254	0.663	0.301	14.183

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	115	78	217	83	135	129	474	78	76
N.S.	1	0.68	1.89	0.72	1.17	1.12	4.12	0.68	0.66
time (sec)	N/A	0.381	0.257	0.655	0.272	0.262	1.178	0.292	14.754

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	140	108	296	99	174	170	813	91	102
N.S.	1	0.77	2.11	0.71	1.24	1.21	5.81	0.65	0.73
time (sec)	N/A	0.514	0.311	0.821	0.245	0.262	2.462	0.284	16.708

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	20	63	37	50	46	44	38	17
N.S.	1	0.32	1.00	0.59	0.79	0.73	0.70	0.60	0.27
time (sec)	N/A	0.182	0.048	0.588	0.244	0.266	0.359	0.286	14.674

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	46	139	71	94	88	240	68	47
N.S.	1	0.52	1.58	0.81	1.07	1.00	2.73	0.77	0.53
time (sec)	N/A	0.258	0.203	0.635	0.275	0.260	0.711	0.284	14.320

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	113	76	211	89	137	129	490	84	76
N.S.	1	0.67	1.87	0.79	1.21	1.14	4.34	0.74	0.67
time (sec)	N/A	0.382	0.362	0.702	0.251	0.263	1.188	0.288	14.155

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	138	106	288	105	177	170	831	97	102
N.S.	1	0.77	2.09	0.76	1.28	1.23	6.02	0.70	0.74
time (sec)	N/A	0.500	0.362	0.730	0.237	0.278	2.475	0.286	15.699

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	20	63	37	50	46	44	38	17
N.S.	1	0.32	1.00	0.59	0.79	0.73	0.70	0.60	0.27
time (sec)	N/A	0.175	0.033	0.653	0.246	0.263	0.403	0.285	14.100

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	46	139	71	94	88	240	68	47
N.S.	1	0.52	1.58	0.81	1.07	1.00	2.73	0.77	0.53
time (sec)	N/A	0.254	0.015	0.687	0.273	0.264	0.684	0.275	0.002

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	113	76	211	89	137	129	490	84	75
N.S.	1	0.67	1.87	0.79	1.21	1.14	4.34	0.74	0.66
time (sec)	N/A	0.368	0.197	0.648	0.277	0.252	1.199	0.278	14.393

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	138	106	288	105	177	170	831	97	102
N.S.	1	0.77	2.09	0.76	1.28	1.23	6.02	0.70	0.74
time (sec)	N/A	0.490	0.013	0.712	0.251	0.266	2.484	0.313	0.002

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	22	65	33	48	46	42	34	17
N.S.	1	0.34	1.00	0.51	0.74	0.71	0.65	0.52	0.26
time (sec)	N/A	0.175	0.043	0.551	0.235	0.277	0.377	0.300	13.159

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	90	48	143	64	91	88	231	62	47
N.S.	1	0.53	1.59	0.71	1.01	0.98	2.57	0.69	0.52
time (sec)	N/A	0.255	0.044	0.779	0.261	0.261	0.680	0.302	0.002

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	115	78	217	83	135	129	478	78	75
N.S.	1	0.68	1.89	0.72	1.17	1.12	4.16	0.68	0.65
time (sec)	N/A	0.365	0.362	0.641	0.277	0.263	1.173	0.297	13.840

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	140	108	296	99	174	170	816	91	102
N.S.	1	0.77	2.11	0.71	1.24	1.21	5.83	0.65	0.73
time (sec)	N/A	0.489	0.012	0.682	0.247	0.264	2.442	0.270	0.002

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>C</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	197	203	177	662	0	437	0	0	0
N.S.	1	1.03	0.90	3.36	0.00	2.22	0.00	0.00	0.00
time (sec)	N/A	1.026	0.924	6.606	0.000	0.112	0.000	0.000	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>C</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	157	158	134	450	0	398	0	0	0
N.S.	1	1.01	0.85	2.87	0.00	2.54	0.00	0.00	0.00
time (sec)	N/A	0.755	0.617	4.087	0.000	0.104	0.000	0.000	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	57	57	57	170	0	355	0	0	0
N.S.	1	1.00	1.00	2.98	0.00	6.23	0.00	0.00	0.00
time (sec)	N/A	0.272	0.137	2.917	0.000	0.099	0.000	0.000	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	C	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	57	57	57	78	0	146	0	0	52
N.S.	1	1.00	1.00	1.37	0.00	2.56	0.00	0.00	0.91
time (sec)	N/A	0.278	0.111	0.821	0.000	0.100	0.000	0.000	13.929

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	106	106	83	217	0	482	0	0	0
N.S.	1	1.00	0.78	2.05	0.00	4.55	0.00	0.00	0.00
time (sec)	N/A	0.418	0.278	2.469	0.000	0.104	0.000	0.000	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	221	231	158	489	0	692	0	0	0
N.S.	1	1.05	0.71	2.21	0.00	3.13	0.00	0.00	0.00
time (sec)	N/A	1.062	1.026	4.483	0.000	0.114	0.000	0.000	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	108	108	246	0	0	0	0	0	0
N.S.	1	1.00	2.28	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.251	2.066	0.000	0.000	0.000	0.000	0.000	0.000

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	105	105	118	0	0	0	0	0	0
N.S.	1	1.00	1.12	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.257	0.312	0.000	0.000	0.000	0.000	0.000	0.000

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	105	105	118	0	0	0	0	0	0
N.S.	1	1.00	1.12	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.251	0.273	0.000	0.000	0.000	0.000	0.000	0.000

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	105	105	118	0	0	0	0	0	0
N.S.	1	1.00	1.12	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.251	0.278	0.000	0.000	0.000	0.000	0.000	0.000

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	105	105	116	0	0	0	0	0	0
N.S.	1	1.00	1.10	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.251	0.275	0.000	0.000	0.000	0.000	0.000	0.000

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	110	110	268	0	0	0	0	0	0
N.S.	1	1.00	2.44	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.254	2.268	0.000	0.000	0.000	0.000	0.000	0.000

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	103	103	121	0	0	0	0	0	0
N.S.	1	1.00	1.17	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.245	0.304	0.000	0.000	0.000	0.000	0.000	0.000



## 2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio  $\frac{\text{number of rules}}{\text{integrand size}}$  is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [50] had the largest ratio of [1.0714300000000010]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	8	8	1.07	14	0.571
2	A	6	6	1.04	14	0.429
3	A	4	4	1.00	14	0.286
4	A	2	2	1.00	14	0.143
5	A	4	3	1.00	14	0.214
6	A	6	5	1.00	14	0.357
7	A	8	7	1.05	14	0.500
8	A	4	4	1.00	14	0.286
9	A	4	4	1.00	14	0.286
10	A	4	4	1.00	14	0.286
11	A	4	4	1.00	14	0.286
12	A	4	4	1.00	14	0.286
13	A	4	4	1.00	14	0.286
14	A	4	4	1.00	12	0.333
15	A	4	4	1.00	13	0.308
16	A	2	2	1.00	12	0.167
17	A	2	2	1.00	12	0.167
18	A	2	2	1.00	12	0.167
19	A	5	5	1.09	12	0.417
20	A	8	8	1.12	12	0.667
21	A	11	11	1.14	12	0.917
22	A	2	2	1.00	12	0.167

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
23	A	5	5	1.09	12	0.417
24	A	8	8	1.12	12	0.667
25	A	11	11	1.14	12	0.917
26	A	2	2	1.00	12	0.167
27	A	5	5	1.09	12	0.417
28	A	7	7	1.12	12	0.583
29	A	11	11	1.14	12	0.917
30	A	2	2	1.00	12	0.167
31	A	5	5	1.09	12	0.417
32	A	7	7	1.12	12	0.583
33	A	11	11	1.14	12	0.917
34	A	4	3	0.34	12	0.250
35	A	7	6	0.53	12	0.500
36	A	10	9	0.68	12	0.750
37	A	13	12	0.77	12	1.000
38	A	4	3	0.32	12	0.250
39	A	7	6	0.52	12	0.500
40	A	10	9	0.67	12	0.750
41	A	13	12	0.77	12	1.000
42	A	4	3	0.32	12	0.250
43	A	7	6	0.52	12	0.500
44	A	9	8	0.67	12	0.667
45	A	13	12	0.77	12	1.000
46	A	4	3	0.34	12	0.250
47	A	7	6	0.53	12	0.500
48	A	9	8	0.68	12	0.667
49	A	13	12	0.77	12	1.000
50	A	15	15	1.03	14	1.071
51	A	12	12	1.01	14	0.857
52	A	4	4	1.00	14	0.286
53	A	4	4	1.00	14	0.286
54	A	7	7	1.00	14	0.500
55	A	15	15	1.05	14	1.071
56	A	5	4	1.00	14	0.286

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
57	A	5	4	1.00	14	0.286
58	A	5	4	1.00	14	0.286
59	A	5	4	1.00	14	0.286
60	A	5	4	1.00	14	0.286
61	A	5	4	1.00	14	0.286
62	A	5	4	1.00	12	0.333

# CHAPTER 3

## LISTING OF INTEGRALS

3.1	$\int (a + a \cos(c + dx))^{7/2} dx$ . . . . .	45
3.2	$\int (a + a \cos(c + dx))^{5/2} dx$ . . . . .	51
3.3	$\int (a + a \cos(c + dx))^{3/2} dx$ . . . . .	56
3.4	$\int \sqrt{a + a \cos(c + dx)} dx$ . . . . .	61
3.5	$\int \frac{1}{\sqrt{a+a \cos(c+dx)}} dx$ . . . . .	65
3.6	$\int \frac{1}{(a+a \cos(c+dx))^{3/2}} dx$ . . . . .	70
3.7	$\int \frac{1}{(a+a \cos(c+dx))^{5/2}} dx$ . . . . .	76
3.8	$\int (a + a \cos(c + dx))^{4/3} dx$ . . . . .	82
3.9	$\int (a + a \cos(c + dx))^{2/3} dx$ . . . . .	87
3.10	$\int \sqrt[3]{a + a \cos(c + dx)} dx$ . . . . .	92
3.11	$\int \frac{1}{\sqrt[3]{a + a \cos(c + dx)}} dx$ . . . . .	97
3.12	$\int \frac{1}{(a+a \cos(c+dx))^{2/3}} dx$ . . . . .	102
3.13	$\int \frac{1}{(a+a \cos(c+dx))^{4/3}} dx$ . . . . .	107
3.14	$\int (a + a \cos(c + dx))^n dx$ . . . . .	112
3.15	$\int (a - a \cos(c + dx))^n dx$ . . . . .	117
3.16	$\int (2 + 2 \cos(c + dx))^n dx$ . . . . .	122
3.17	$\int (2 - 2 \cos(c + dx))^n dx$ . . . . .	126
3.18	$\int \frac{1}{5+3 \cos(c+dx)} dx$ . . . . .	130
3.19	$\int \frac{1}{(5+3 \cos(c+dx))^2} dx$ . . . . .	135
3.20	$\int \frac{1}{(5+3 \cos(c+dx))^3} dx$ . . . . .	141
3.21	$\int \frac{1}{(5+3 \cos(c+dx))^4} dx$ . . . . .	147
3.22	$\int \frac{1}{5-3 \cos(c+dx)} dx$ . . . . .	154
3.23	$\int \frac{1}{(5-3 \cos(c+dx))^2} dx$ . . . . .	159
3.24	$\int \frac{1}{(5-3 \cos(c+dx))^3} dx$ . . . . .	164
3.25	$\int \frac{1}{(5-3 \cos(c+dx))^4} dx$ . . . . .	170
3.26	$\int \frac{1}{-5+3 \cos(c+dx)} dx$ . . . . .	177
3.27	$\int \frac{1}{(-5+3 \cos(c+dx))^2} dx$ . . . . .	182

3.28	$\int \frac{1}{(-5+3 \cos(c+dx))^3} dx$	188
3.29	$\int \frac{1}{(-5+3 \cos(c+dx))^4} dx$	194
3.30	$\int \frac{1}{-5-3 \cos(c+dx)} dx$	201
3.31	$\int \frac{1}{(-5-3 \cos(c+dx))^2} dx$	206
3.32	$\int \frac{1}{(-5-3 \cos(c+dx))^3} dx$	212
3.33	$\int \frac{1}{(-5-3 \cos(c+dx))^4} dx$	218
3.34	$\int \frac{1}{3+5 \cos(c+dx)} dx$	225
3.35	$\int \frac{1}{(3+5 \cos(c+dx))^2} dx$	230
3.36	$\int \frac{1}{(3+5 \cos(c+dx))^3} dx$	236
3.37	$\int \frac{1}{(3+5 \cos(c+dx))^4} dx$	244
3.38	$\int \frac{1}{3-5 \cos(c+dx)} dx$	252
3.39	$\int \frac{1}{(3-5 \cos(c+dx))^2} dx$	257
3.40	$\int \frac{1}{(3-5 \cos(c+dx))^3} dx$	263
3.41	$\int \frac{1}{(3-5 \cos(c+dx))^4} dx$	271
3.42	$\int \frac{1}{-3+5 \cos(c+dx)} dx$	279
3.43	$\int \frac{1}{(-3+5 \cos(c+dx))^2} dx$	284
3.44	$\int \frac{1}{(-3+5 \cos(c+dx))^3} dx$	290
3.45	$\int \frac{1}{(-3+5 \cos(c+dx))^4} dx$	298
3.46	$\int \frac{1}{-3-5 \cos(c+dx)} dx$	306
3.47	$\int \frac{1}{(-3-5 \cos(c+dx))^2} dx$	311
3.48	$\int \frac{1}{(-3-5 \cos(c+dx))^3} dx$	317
3.49	$\int \frac{1}{(-3-5 \cos(c+dx))^4} dx$	325
3.50	$\int (a + b \cos(c + dx))^{5/2} dx$	333
3.51	$\int (a + b \cos(c + dx))^{3/2} dx$	342
3.52	$\int \sqrt{a + b \cos(c + dx)} dx$	350
3.53	$\int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx$	355
3.54	$\int \frac{1}{(a+b \cos(c+dx))^{3/2}} dx$	360
3.55	$\int \frac{1}{(a+b \cos(c+dx))^{5/2}} dx$	366
3.56	$\int (a + b \cos(c + dx))^{4/3} dx$	375
3.57	$\int (a + b \cos(c + dx))^{2/3} dx$	380
3.58	$\int \sqrt[3]{a + b \cos(c + dx)} dx$	385
3.59	$\int \frac{1}{\sqrt[3]{a + b \cos(c + dx)}} dx$	390
3.60	$\int \frac{1}{(a+b \cos(c+dx))^{2/3}} dx$	395
3.61	$\int \frac{1}{(a+b \cos(c+dx))^{4/3}} dx$	400
3.62	$\int (a + b \cos(c + dx))^n dx$	405

### 3.1 $\int (a + a \cos(c + dx))^{7/2} dx$

3.1.1	Optimal result . . . . .	45
3.1.2	Mathematica [A] (verified) . . . . .	45
3.1.3	Rubi [A] (verified) . . . . .	46
3.1.4	Maple [A] (verified) . . . . .	48
3.1.5	Fricas [A] (verification not implemented) . . . . .	48
3.1.6	Sympy [F(-1)] . . . . .	49
3.1.7	Maxima [A] (verification not implemented) . . . . .	49
3.1.8	Giac [A] (verification not implemented) . . . . .	49
3.1.9	Mupad [F(-1)] . . . . .	50

#### 3.1.1 Optimal result

Integrand size = 14, antiderivative size = 119

$$\int (a + a \cos(c + dx))^{7/2} dx = \frac{256a^4 \sin(c + dx)}{35d\sqrt{a + a \cos(c + dx)}} + \frac{64a^3 \sqrt{a + a \cos(c + dx)} \sin(c + dx)}{35d} + \frac{24a^2 (a + a \cos(c + dx))^{3/2} \sin(c + dx)}{35d} + \frac{2a(a + a \cos(c + dx))^{5/2} \sin(c + dx)}{7d}$$

output  $24/35*a^2*(a+a*\cos(d*x+c))^(3/2)*\sin(d*x+c)/d+2/7*a*(a+a*\cos(d*x+c))^(5/2)*\sin(d*x+c)/d+256/35*a^4*\sin(d*x+c)/d/(a+a*\cos(d*x+c))^(1/2)+64/35*a^3*\sin(d*x+c)*(a+a*\cos(d*x+c))^(1/2)/d$

#### 3.1.2 Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.70

$$\int (a + a \cos(c + dx))^{7/2} dx = \frac{a^3 \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \left(1225 \sin\left(\frac{1}{2}(c + dx)\right) + 245 \sin\left(\frac{3}{2}(c + dx)\right) + 49 \sin\left(\frac{5}{2}(c + dx)\right)\right)}{140d}$$

input `Integrate[(a + a*Cos[c + d*x])^(7/2),x]`

output  $(a^3 \sqrt{a(1 + \cos(c + dx))} \sec((c + dx)/2) (1225 \sin((c + dx)/2) + 245 \sin((3(c + dx))/2) + 49 \sin((5(c + dx))/2) + 5 \sin((7(c + dx))/2)) / (140d)$

### 3.1.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$ , Rules used = {3042, 3126, 3042, 3126, 3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \cos(c + dx) + a)^{7/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( a \sin\left(c + dx + \frac{\pi}{2}\right) + a \right)^{7/2} dx \\
 & \quad \downarrow \text{3126} \\
 & \frac{12}{7} a \int (\cos(c + dx)a + a)^{5/2} dx + \frac{2a \sin(c + dx)(a \cos(c + dx) + a)^{5/2}}{7d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{12}{7} a \int \left( \sin\left(c + dx + \frac{\pi}{2}\right) a + a \right)^{5/2} dx + \frac{2a \sin(c + dx)(a \cos(c + dx) + a)^{5/2}}{7d} \\
 & \quad \downarrow \text{3126} \\
 & \frac{12}{7} a \left( \frac{8}{5} a \int (\cos(c + dx)a + a)^{3/2} dx + \frac{2a \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \right) + \\
 & \quad \frac{2a \sin(c + dx)(a \cos(c + dx) + a)^{5/2}}{7d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{12}{7} a \left( \frac{8}{5} a \int \left( \sin\left(c + dx + \frac{\pi}{2}\right) a + a \right)^{3/2} dx + \frac{2a \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \right) + \\
 & \quad \frac{2a \sin(c + dx)(a \cos(c + dx) + a)^{5/2}}{7d} \\
 & \quad \downarrow \text{3126}
 \end{aligned}$$

$$\frac{12}{7}a \left( \frac{8}{5}a \left( \frac{4}{3}a \int \sqrt{\cos(c+dx)a+adx} + \frac{2a \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3d} \right) + \frac{2a \sin(c+dx)(a \cos(c+dx)+a)}{5d} \right. \\ \left. \frac{2a \sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7d} \right) \\ \downarrow \text{3042}$$

$$\frac{12}{7}a \left( \frac{8}{5}a \left( \frac{4}{3}a \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx} + \frac{2a \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3d} \right) + \frac{2a \sin(c+dx)(a \cos(c+dx)+a)}{5d} \right. \\ \left. \frac{2a \sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7d} \right) \\ \downarrow \text{3125}$$

$$\frac{12}{7}a \left( \frac{8}{5}a \left( \frac{8a^2 \sin(c+dx)}{3d\sqrt{a \cos(c+dx)+a}} + \frac{2a \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3d} \right) + \frac{2a \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d} \right) \\ \frac{2a \sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7d}$$

input `Int[(a + a*Cos[c + d*x])^(7/2),x]`

output `(2*a*(a + a*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(7*d) + (12*a*((2*a*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*d) + (8*a*((8*a^2*Sin[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]])) + (2*a*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d)))/5)/7`

### 3.1.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`



rule 3126 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n) Int[(a + b*Sin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0]`

### 3.1.4 Maple [A] (verified)

Time = 0.78 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.72

method	result	size
default	$\frac{16a^4 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(5 \cos^6\left(\frac{dx}{2} + \frac{c}{2}\right) + 6 \cos^4\left(\frac{dx}{2} + \frac{c}{2}\right) + 8 \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 16\right) \sqrt{2}}{35 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$	86

input `int((a+cos(d*x+c)*a)^(7/2),x,method=_RETURNVERBOSE)`

output `16/35*a^4*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)*(5*cos(1/2*d*x+1/2*c)^6+6*cos(1/2*d*x+1/2*c)^4+8*cos(1/2*d*x+1/2*c)^2+16)*2^(1/2)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d`

### 3.1.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.63

$$\int (a + a \cos(c + dx))^{7/2} dx = \frac{2 \left( 5 a^3 \cos(dx + c)^3 + 27 a^3 \cos(dx + c)^2 + 71 a^3 \cos(dx + c) + 177 a^3 \right) \sqrt{a \cos(dx + c) + a \sin(dx + c)}}{35 (d \cos(dx + c) + d)}$$

input `integrate((a+a*cos(d*x+c))^(7/2),x, algorithm="fracas")`

output `2/35*(5*a^3*cos(d*x + c)^3 + 27*a^3*cos(d*x + c)^2 + 71*a^3*cos(d*x + c) + 177*a^3)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/(d*cos(d*x + c) + d)`

### 3.1.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{7/2} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(7/2),x)`

output `Timed out`

### 3.1.7 Maxima [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.65

$$\int (a + a \cos(c + dx))^{7/2} dx = \frac{(5\sqrt{2}a^3 \sin(\frac{7}{2}dx + \frac{7}{2}c) + 49\sqrt{2}a^3 \sin(\frac{5}{2}dx + \frac{5}{2}c) + 245\sqrt{2}a^3 \sin(\frac{3}{2}dx + \frac{3}{2}c) + 1225\sqrt{2}a^3 \sin(\frac{1}{2}dx + \frac{1}{2}c)) \sqrt{a}}{140d}$$

input `integrate((a+a*cos(d*x+c))^(7/2),x, algorithm="maxima")`

output `1/140*(5*sqrt(2)*a^3*sin(7/2*d*x + 7/2*c) + 49*sqrt(2)*a^3*sin(5/2*d*x + 5/2*c) + 245*sqrt(2)*a^3*sin(3/2*d*x + 3/2*c) + 1225*sqrt(2)*a^3*sin(1/2*d*x + 1/2*c))*sqrt(a)/d`

### 3.1.8 Giac [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.91

$$\int (a + a \cos(c + dx))^{7/2} dx = \frac{\sqrt{2}(5a^3 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) \sin(\frac{7}{2}dx + \frac{7}{2}c) + 49a^3 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) \sin(\frac{5}{2}dx + \frac{5}{2}c) + 245a^3 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) \sin(\frac{3}{2}dx + \frac{3}{2}c) + 1225a^3 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) \sin(\frac{1}{2}dx + \frac{1}{2}c)) \sqrt{a}}{140d}$$

input `integrate((a+a*cos(d*x+c))^(7/2),x, algorithm="giac")`

output  $1/140*\sqrt{2}*(5*a^3*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(7/2*d*x + 7/2*c) + 49*a^3*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(5/2*d*x + 5/2*c) + 245*a^3*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(3/2*d*x + 3/2*c) + 1225*a^3*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(1/2*d*x + 1/2*c))*\sqrt{a}/d$

### 3.1.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{7/2} dx = \int (a + a \cos(c + dx))^{7/2} dx$$

input `int((a + a*cos(c + d*x))^(7/2),x)`

output `int((a + a*cos(c + d*x))^(7/2), x)`

## 3.2 $\int (a + a \cos(c + dx))^{5/2} dx$

3.2.1	Optimal result . . . . .	51
3.2.2	Mathematica [A] (verified) . . . . .	51
3.2.3	Rubi [A] (verified) . . . . .	52
3.2.4	Maple [A] (verified) . . . . .	53
3.2.5	Fricas [A] (verification not implemented) . . . . .	54
3.2.6	Sympy [F] . . . . .	54
3.2.7	Maxima [A] (verification not implemented) . . . . .	54
3.2.8	Giac [A] (verification not implemented) . . . . .	55
3.2.9	Mupad [F(-1)] . . . . .	55

### 3.2.1 Optimal result

Integrand size = 14, antiderivative size = 89

$$\int (a + a \cos(c + dx))^{5/2} dx = \frac{64a^3 \sin(c + dx)}{15d\sqrt{a + a \cos(c + dx)}} + \frac{16a^2 \sqrt{a + a \cos(c + dx)} \sin(c + dx)}{15d} + \frac{2a(a + a \cos(c + dx))^{3/2} \sin(c + dx)}{5d}$$

```
output 2/5*a*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d+64/15*a^3*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+16/15*a^2*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d
```

### 3.2.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.80

$$\int (a + a \cos(c + dx))^{5/2} dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \left(150 \sin\left(\frac{1}{2}(c + dx)\right) + 25 \sin\left(\frac{3}{2}(c + dx)\right) + 3 \sin\left(\frac{5}{2}(c + dx)\right)\right)}{30d}$$

```
input Integrate[(a + a*Cos[c + d*x])^(5/2),x]
```

```
output (a^2*sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*(150*Sin[(c + d*x)/2] + 25*Sin[(3*(c + d*x))/2] + 3*Sin[(5*(c + d*x))/2]))/(30*d)
```

### 3.2.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.04, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3042, 3126, 3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \cos(c + dx) + a)^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( a \sin\left(c + dx + \frac{\pi}{2}\right) + a \right)^{5/2} dx \\
 & \quad \downarrow \text{3126} \\
 & \frac{8}{5}a \int (\cos(c + dx)a + a)^{3/2} dx + \frac{2a \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{8}{5}a \int \left( \sin\left(c + dx + \frac{\pi}{2}\right) a + a \right)^{3/2} dx + \frac{2a \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \\
 & \quad \downarrow \text{3126} \\
 & \frac{8}{5}a \left( \frac{4}{3}a \int \sqrt{\cos(c + dx)a + adx} + \frac{2a \sin(c + dx)\sqrt{a \cos(c + dx) + a}}{3d} \right) + \\
 & \quad \frac{2a \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{8}{5}a \left( \frac{4}{3}a \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right) a + adx} + \frac{2a \sin(c + dx)\sqrt{a \cos(c + dx) + a}}{3d} \right) + \\
 & \quad \frac{2a \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \\
 & \quad \downarrow \text{3125} \\
 & \frac{8}{5}a \left( \frac{8a^2 \sin(c + dx)}{3d\sqrt{a \cos(c + dx) + a}} + \frac{2a \sin(c + dx)\sqrt{a \cos(c + dx) + a}}{3d} \right) + \\
 & \quad \frac{2a \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d}
 \end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^(5/2), x]`

```
output (2*a*(a + a*cos[c + d*x])^(3/2)*sin[c + d*x])/(5*d) + (8*a*((8*a^2*sin[c +
d*x])/(3*d*Sqrt[a + a*cos[c + d*x]]) + (2*a*Sqrt[a + a*cos[c + d*x]]*sin[
c + d*x])/(3*d)))/5
```

### 3.2.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3125 Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*b*(Cos
[c + d*x]/(d*Sqrt[a + b*sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && Eq
Q[a^2 - b^2, 0]
```

```
rule 3126 Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos
[c + d*x]*((a + b*sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n)
Int[(a + b*sin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[
a^2 - b^2, 0] && IGtQ[n - 1/2, 0]
```

## 3.2.4 Maple [A] (verified)

Time = 0.81 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{8a^3 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(3 \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8\right) \sqrt{2}}{15 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$	73

```
input int((a+cos(d*x+c)*a)^(5/2),x,method=_RETURNVERBOSE)
```

```
output 8/15*a^3*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)*(3*cos(1/2*d*x+1/2*c)^4+4*c
os(1/2*d*x+1/2*c)^2+8)*2^(1/2)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d
```

### 3.2.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.70

$$\int (a + a \cos(c + dx))^{5/2} dx = \frac{2(3a^2 \cos(dx + c)^2 + 14a^2 \cos(dx + c) + 43a^2) \sqrt{a \cos(dx + c) + a \sin(dx + c)}}{15(d \cos(dx + c) + d)}$$

input `integrate((a+a*cos(d*x+c))^(5/2),x, algorithm="fracas")`

output `2/15*(3*a^2*cos(d*x + c)^2 + 14*a^2*cos(d*x + c) + 43*a^2)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/(d*cos(d*x + c) + d)`

### 3.2.6 Sympy [F]

$$\int (a + a \cos(c + dx))^{5/2} dx = \int (a \cos(c + dx) + a)^{\frac{5}{2}} dx$$

input `integrate((a+a*cos(d*x+c))**(5/2),x)`

output `Integral((a*cos(c + d*x) + a)**(5/2), x)`

### 3.2.7 Maxima [A] (verification not implemented)

Time = 0.50 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.67

$$\int (a + a \cos(c + dx))^{5/2} dx = \frac{(3\sqrt{2}a^2 \sin(\frac{5}{2}dx + \frac{5}{2}c) + 25\sqrt{2}a^2 \sin(\frac{3}{2}dx + \frac{3}{2}c) + 150\sqrt{2}a^2 \sin(\frac{1}{2}dx + \frac{1}{2}c))\sqrt{a}}{30d}$$

input `integrate((a+a*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `1/30*(3*sqrt(2)*a^2*sin(5/2*d*x + 5/2*c) + 25*sqrt(2)*a^2*sin(3/2*d*x + 3/2*c) + 150*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c))*sqrt(a)/d`

---

3.2.  $\int (a + a \cos(c + dx))^{5/2} dx$

### 3.2.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.94

$$\int (a + a \cos(c + dx))^{5/2} dx = \frac{\sqrt{2}(3a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \sin(\frac{5}{2} dx + \frac{5}{2} c) + 25a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \sin(\frac{3}{2} dx + \frac{3}{2} c) + 150a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \sin(\frac{1}{2} dx + \frac{1}{2} c)) \sqrt{a}}{30d}$$

input `integrate((a+a*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `1/30*sqrt(2)*(3*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(5/2*d*x + 5/2*c) + 25*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(3/2*d*x + 3/2*c) + 150*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c))*sqrt(a)/d`

### 3.2.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} dx = \int (a + a \cos(c + dx))^{5/2} dx$$

input `int((a + a*cos(c + d*x))^(5/2),x)`

output `int((a + a*cos(c + d*x))^(5/2), x)`



### 3.3 $\int (a + a \cos(c + dx))^{3/2} dx$

3.3.1	Optimal result . . . . .	56
3.3.2	Mathematica [A] (verified) . . . . .	56
3.3.3	Rubi [A] (verified) . . . . .	57
3.3.4	Maple [A] (verified) . . . . .	58
3.3.5	Fricas [A] (verification not implemented) . . . . .	58
3.3.6	Sympy [F] . . . . .	59
3.3.7	Maxima [A] (verification not implemented) . . . . .	59
3.3.8	Giac [A] (verification not implemented) . . . . .	59
3.3.9	Mupad [F(-1)] . . . . .	60

#### 3.3.1 Optimal result

Integrand size = 14, antiderivative size = 59

$$\int (a + a \cos(c + dx))^{3/2} dx = \frac{8a^2 \sin(c + dx)}{3d\sqrt{a + a \cos(c + dx)}} + \frac{2a\sqrt{a + a \cos(c + dx)} \sin(c + dx)}{3d}$$

output `8/3*a^2*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/3*a*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d`

#### 3.3.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.93

$$\int (a + a \cos(c + dx))^{3/2} dx = \frac{a\sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \left(9 \sin\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{3}{2}(c + dx)\right)\right)}{3d}$$

input `Integrate[(a + a*Cos[c + d*x])^(3/2),x]`

output `(a*sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*(9*Sin[(c + d*x)/2] + Sin[3*(c + d*x)/2]))/(3*d)`

### 3.3.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \cos(c + dx) + a)^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( a \sin \left( c + dx + \frac{\pi}{2} \right) + a \right)^{3/2} dx \\
 & \quad \downarrow \text{3126} \\
 & \frac{4}{3} a \int \sqrt{\cos(c + dx)a + adx} + \frac{2a \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{4}{3} a \int \sqrt{\sin \left( c + dx + \frac{\pi}{2} \right) a + adx} + \frac{2a \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \\
 & \quad \downarrow \text{3125} \\
 & \frac{8a^2 \sin(c + dx)}{3d \sqrt{a \cos(c + dx) + a}} + \frac{2a \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d}
 \end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^(3/2), x]`

output `(8*a^2*Sin[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]]) + (2*a*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d)`

### 3.3.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3126 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n) Int[(a + b*Sin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0]`

### 3.3.4 Maple [A] (verified)

Time = 0.81 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.98

method	result	size
default	$\frac{4a^2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right) \sqrt{2}}{3\sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$	58

input `int((a+cos(d*x+c)*a)^(3/2),x,method=_RETURNVERBOSE)`

output  $\frac{4}{3}a^2 \cos\left(\frac{1}{2}d*x + \frac{1}{2}c\right) \sin\left(\frac{1}{2}d*x + \frac{1}{2}c\right) \left(\cos\left(\frac{1}{2}d*x + \frac{1}{2}c\right)^2 + 2\right) 2^{\frac{1}{2}} \left(\frac{1}{2}\right) / \left(a \cos\left(\frac{1}{2}d*x + \frac{1}{2}c\right)^2\right)^{\frac{1}{2}} / d$

### 3.3.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.75

$$\int (a + a \cos(c + dx))^{3/2} dx = \frac{2(a \cos(dx + c) + 5a) \sqrt{a \cos(dx + c) + a \sin(dx + c)}}{3(d \cos(dx + c) + d)}$$

input `integrate((a+a*cos(d*x+c))^(3/2),x, algorithm="fracas")`

output  $2/3*(a*\cos(d*x + c) + 5*a)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)/(d*\cos(d*x + c) + d)$

### 3.3.6 Sympy [F]

$$\int (a + a \cos(c + dx))^{3/2} dx = \int (a \cos(c + dx) + a)^{\frac{3}{2}} dx$$

input `integrate((a+a*cos(d*x+c))**(3/2),x)`

output `Integral((a*cos(c + d*x) + a)**(3/2), x)`

### 3.3.7 Maxima [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.64

$$\int (a + a \cos(c + dx))^{3/2} dx = \frac{(\sqrt{2}a \sin(\frac{3}{2} dx + \frac{3}{2} c) + 9 \sqrt{2}a \sin(\frac{1}{2} dx + \frac{1}{2} c))\sqrt{a}}{3 d}$$

input `integrate((a+a*cos(d*x+c))^(3/2),x, algorithm="maxima")`

output  $1/3*(\sqrt{2}*a*\sin(3/2*d*x + 3/2*c) + 9*\sqrt{2}*a*\sin(1/2*d*x + 1/2*c))*\sqrt{a}/d$

### 3.3.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.93

$$\int (a + a \cos(c + dx))^{3/2} dx = \frac{\sqrt{2}(a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \sin(\frac{3}{2} dx + \frac{3}{2} c) + 9 a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \sin(\frac{1}{2} dx + \frac{1}{2} c))\sqrt{a}}{3 d}$$

input `integrate((a+a*cos(d*x+c))^(3/2),x, algorithm="giac")`

output  $1/3*\sqrt{2}*(a*\operatorname{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(3/2*d*x + 3/2*c) + 9*a*\operatorname{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(1/2*d*x + 1/2*c))*\sqrt{a}/d$

**3.3.9 Mupad [F(-1)]**

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} dx = \int (a + a \cos(c + dx))^{3/2} dx$$

input `int((a + a*cos(c + d*x))^(3/2),x)`output `int((a + a*cos(c + d*x))^(3/2), x)`

### 3.4 $\int \sqrt{a + a \cos(c + dx)} dx$

3.4.1	Optimal result . . . . .	61
3.4.2	Mathematica [A] (verified) . . . . .	61
3.4.3	Rubi [A] (verified) . . . . .	62
3.4.4	Maple [A] (verified) . . . . .	63
3.4.5	Fricas [A] (verification not implemented) . . . . .	63
3.4.6	Sympy [F] . . . . .	63
3.4.7	Maxima [A] (verification not implemented) . . . . .	64
3.4.8	Giac [A] (verification not implemented) . . . . .	64
3.4.9	Mupad [B] (verification not implemented) . . . . .	64

#### 3.4.1 Optimal result

Integrand size = 14, antiderivative size = 26

$$\int \sqrt{a + a \cos(c + dx)} dx = \frac{2a \sin(c + dx)}{d \sqrt{a + a \cos(c + dx)}}$$

output `2*a*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)`

#### 3.4.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.12

$$\int \sqrt{a + a \cos(c + dx)} dx = \frac{2\sqrt{a(1 + \cos(c + dx))} \tan\left(\frac{1}{2}(c + dx)\right)}{d}$$

input `Integrate[Sqrt[a + a*Cos[c + d*x]],x]`

output `(2*Sqrt[a*(1 + Cos[c + d*x])]*Tan[(c + d*x)/2])/d`

### 3.4.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a \cos(c + dx) + a} dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a} dx$$

$$\downarrow \text{3125}$$

$$\frac{2a \sin(c + dx)}{d\sqrt{a \cos(c + dx) + a}}$$

input `Int[Sqrt[a + a*Cos[c + d*x]],x]`

output `(2*a*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])`

#### 3.4.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && Eq Q[a^2 - b^2, 0]`

### 3.4.4 Maple [A] (verified)

Time = 0.80 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.65

method	result	size
default	$\frac{2a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2}}{\sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$	43
risch	$-\frac{i\sqrt{2} \sqrt{a(e^{i(dx+c)}+1)^2 e^{-i(dx+c)} (e^{i(dx+c)}-1)}}{(e^{i(dx+c)}+1)d}$	60

input `int((a+cos(d*x+c)*a)^(1/2),x,method=_RETURNVERBOSE)`

output `2*a*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)*2^(1/2)/(a*cos(1/2*d*x+1/2*c)^2  
^(1/2)/d`

### 3.4.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.23

$$\int \sqrt{a + a \cos(c + dx)} dx = \frac{2 \sqrt{a \cos(dx + c) + a} \sin(dx + c)}{d \cos(dx + c) + d}$$

input `integrate((a+a*cos(d*x+c))^(1/2),x, algorithm="fracas")`

output `2*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/(d*cos(d*x + c) + d)`

### 3.4.6 Sympy [F]

$$\int \sqrt{a + a \cos(c + dx)} dx = \int \sqrt{a \cos(c + dx) + a} dx$$

input `integrate((a+a*cos(d*x+c))**(1/2),x)`

output `Integral(sqrt(a*cos(c + d*x) + a), x)`



**3.4.7 Maxima [A] (verification not implemented)**

Time = 0.44 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \sqrt{a + a \cos(c + dx)} dx = \frac{2\sqrt{2}\sqrt{a} \sin\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{d}$$

input `integrate((a+a*cos(d*x+c))^(1/2),x, algorithm="maxima")`output `2*sqrt(2)*sqrt(a)*sin(1/2*d*x + 1/2*c)/d`**3.4.8 Giac [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.15

$$\int \sqrt{a + a \cos(c + dx)} dx = \frac{2\sqrt{2}\sqrt{a} \operatorname{sgn}\left(\cos\left(\frac{1}{2} dx + \frac{1}{2} c\right)\right) \sin\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{d}$$

input `integrate((a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`output `2*sqrt(2)*sqrt(a)*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)/d`**3.4.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.27

$$\int \sqrt{a + a \cos(c + dx)} dx = \frac{2 \sin(c + dx) \sqrt{a (\cos(c + dx) + 1)}}{d (\cos(c + dx) + 1)}$$

input `int((a + a*cos(c + d*x))^(1/2),x)`output `(2*sin(c + d*x)*(a*(cos(c + d*x) + 1))^(1/2))/(d*(cos(c + d*x) + 1))`

### 3.5 $\int \frac{1}{\sqrt{a+a \cos(c+dx)}} dx$

3.5.1	Optimal result . . . . .	65
3.5.2	Mathematica [A] (verified) . . . . .	65
3.5.3	Rubi [A] (verified) . . . . .	66
3.5.4	Maple [C] (warning: unable to verify) . . . . .	67
3.5.5	Fricas [A] (verification not implemented) . . . . .	67
3.5.6	Sympy [F] . . . . .	68
3.5.7	Maxima [B] (verification not implemented) . . . . .	68
3.5.8	Giac [B] (verification not implemented) . . . . .	69
3.5.9	Mupad [B] (verification not implemented) . . . . .	69

#### 3.5.1 Optimal result

Integrand size = 14, antiderivative size = 46

$$\int \frac{1}{\sqrt{a+a \cos(c+dx)}} dx = \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{a+a \cos(c+dx)}}\right)}{\sqrt{ad}}$$

output `arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)/d/a^(1/2)`

#### 3.5.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.87

$$\int \frac{1}{\sqrt{a+a \cos(c+dx)}} dx = \frac{2 \operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c+dx)\right)\right) \cos\left(\frac{1}{2}(c+dx)\right)}{d \sqrt{a(1+\cos(c+dx))}}$$

input `Integrate[1/Sqrt[a + a*Cos[c + d*x]],x]`

output `(2*ArcTanh[Sin[(c + d*x)/2]]*Cos[(c + d*x)/2])/(d*Sqrt[a*(1 + Cos[c + d*x])])`

### 3.5.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{1}{\sqrt{a \cos(c + dx) + a}} dx \\
 \downarrow \text{3042} \\
 \int \frac{1}{\sqrt{a \sin(c + dx + \frac{\pi}{2}) + a}} dx \\
 \downarrow \text{3128} \\
 \frac{2 \int \frac{1}{2a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d} \\
 \downarrow \text{219} \\
 \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{\sqrt{ad}}
 \end{array}$$

input `Int[1/Sqrt[a + a*Cos[c + d*x]],x]`

output `(Sqrt[2]*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])]/(Sqrt[a]*d)`

#### 3.5.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3128 Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d
Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x]])],
x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]
```

### 3.5.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.29 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.22

method	result	size
default	$\frac{\sqrt{2} \operatorname{am}^{-1}\left(\frac{dx}{2} + \frac{c}{2}   1\right)}{d \sec\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \operatorname{csgn}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}$	56

```
input int(1/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/d*2^(1/2)/sec(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/csgn(cos(1/2
*d*x+1/2*c))*InverseJacobiAM(1/2*d*x+1/2*c,1)
```

### 3.5.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 126, normalized size of antiderivative = 2.74

$$\int \frac{1}{\sqrt{a + a \cos(c + dx)}} dx = \left[ \frac{\sqrt{2} \log \left( -\frac{\cos(dx+c)^2 - 2\sqrt{2}\sqrt{a \cos(dx+c)+a} \sin(dx+c) - 2 \cos(dx+c) - 3}{\sqrt{a} \cos(dx+c)^2 + 2 \cos(dx+c) + 1} \right)}{2\sqrt{ad}}, \right. \\ \left. - \frac{\sqrt{2} \sqrt{-\frac{1}{a}} \arctan \left( \frac{\sqrt{2}\sqrt{a \cos(dx+c)+a} \sqrt{-\frac{1}{a}}}{\sin(dx+c)} \right)}{d} \right]$$

```
input integrate(1/(a+a*cos(d*x+c))^(1/2),x, algorithm="fricas")
```

output `[1/2*sqrt(2)*log(-(cos(d*x + c)^2 - 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(a) - 2*cos(d*x + c) - 3)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1))/(sqrt(a)*d), -sqrt(2)*sqrt(-1/a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(-1/a)/sin(d*x + c))/d]`

### 3.5.6 Sympy [F]

$$\int \frac{1}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{1}{\sqrt{a \cos(c + dx) + a}} dx$$

input `integrate(1/(a+a*cos(d*x+c))**(1/2),x)`

output `Integral(1/sqrt(a*cos(c + d*x) + a), x)`

### 3.5.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 90 vs.  $2(37) = 74$ .

Time = 0.43 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.96

$$\int \frac{1}{\sqrt{a + a \cos(c + dx)}} dx = \frac{\sqrt{2} \log \left( \cos \left( \frac{1}{2} dx + \frac{1}{2} c \right)^2 + \sin \left( \frac{1}{2} dx + \frac{1}{2} c \right)^2 + 2 \sin \left( \frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right) - \sqrt{2} \log \left( \cos \left( \frac{1}{2} dx + \frac{1}{2} c \right)^2 + \sin \left( \frac{1}{2} dx + \frac{1}{2} c \right)^2 - 2 \sin \left( \frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right)}{2 \sqrt{ad}}$$

input `integrate(1/(a+a*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `1/2*(sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c) + 1) - sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 - 2*sin(1/2*d*x + 1/2*c) + 1))/(sqrt(a)*d)`

### 3.5.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 93 vs.  $2(37) = 74$ .

Time = 0.28 (sec) , antiderivative size = 93, normalized size of antiderivative = 2.02

$$\int \frac{1}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{\frac{\sqrt{2} \log\left(\left|\frac{1}{\sin\left(\frac{1}{2} dx + \frac{1}{2} c\right)} + \sin\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 2\right|\right)}{\sqrt{a} \operatorname{sgn}\left(\cos\left(\frac{1}{2} dx + \frac{1}{2} c\right)\right)} - \frac{\sqrt{2} \log\left(\left|\frac{1}{\sin\left(\frac{1}{2} dx + \frac{1}{2} c\right)} + \sin\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 2\right|\right)}{\sqrt{a} \operatorname{sgn}\left(\cos\left(\frac{1}{2} dx + \frac{1}{2} c\right)\right)}}{4d}$$

input `integrate(1/(a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `1/4*(sqrt(2)*log(abs(1/sin(1/2*d*x + 1/2*c) + sin(1/2*d*x + 1/2*c) + 2))/(sqrt(a)*sgn(cos(1/2*d*x + 1/2*c))) - sqrt(2)*log(abs(1/sin(1/2*d*x + 1/2*c) + sin(1/2*d*x + 1/2*c) - 2))/(sqrt(a)*sgn(cos(1/2*d*x + 1/2*c))))/d`

### 3.5.9 Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.98

$$\int \frac{1}{\sqrt{a + a \cos(c + dx)}} dx = \frac{F\left(\frac{c}{2} + \frac{dx}{2} \mid 1\right) \sqrt{\frac{2(a+a \cos(c+dx))}{a}}}{d \sqrt{a + a \cos(c + dx)}}$$

input `int(1/(a + a*cos(c + d*x))^(1/2),x)`

output `(ellipticF(c/2 + (d*x)/2, 1)*((2*(a + a*cos(c + d*x)))/a)^(1/2))/(d*(a + a*cos(c + d*x))^(1/2))`

### 3.6 $\int \frac{1}{(a+a \cos(c+dx))^{3/2}} dx$

3.6.1	Optimal result	70
3.6.2	Mathematica [A] (verified)	70
3.6.3	Rubi [A] (verified)	71
3.6.4	Maple [B] (verified)	72
3.6.5	Fricas [B] (verification not implemented)	73
3.6.6	Sympy [F]	73
3.6.7	Maxima [B] (verification not implemented)	73
3.6.8	Giac [A] (verification not implemented)	74
3.6.9	Mupad [F(-1)]	75

#### 3.6.1 Optimal result

Integrand size = 14, antiderivative size = 77

$$\int \frac{1}{(a + a \cos(c + dx))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{\sin(c + dx)}{2d(a + a \cos(c + dx))^{3/2}}$$

output `1/2*sin(d*x+c)/d/(a+a*cos(d*x+c))^(3/2)+1/4*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(3/2)/d*2^(1/2)`

#### 3.6.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.82

$$\int \frac{1}{(a + a \cos(c + dx))^{3/2}} dx = \frac{\cos^2\left(\frac{1}{2}(c + dx)\right) \left(\operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c + dx)\right)\right) \cos\left(\frac{1}{2}(c + dx)\right) + \tan\left(\frac{1}{2}(c + dx)\right)\right)}{d(a(1 + \cos(c + dx)))^{3/2}}$$

input `Integrate[(a + a*Cos[c + d*x])^(-3/2),x]`

output `(Cos[(c + d*x)/2]^2*(ArcTanh[Sin[(c + d*x)/2]]*Cos[(c + d*x)/2] + Tan[(c + d*x)/2]))/(d*(a*(1 + Cos[c + d*x]))^(3/2))`

### 3.6.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$ , Rules used = {3042, 3129, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \cos(c + dx) + a)^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a \sin(c + dx + \frac{\pi}{2}) + a)^{3/2}} dx \\
 & \quad \downarrow \text{3129} \\
 & \frac{\int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx}{4a} + \frac{\sin(c + dx)}{2d(a \cos(c + dx) + a)^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a} + \frac{\sin(c + dx)}{2d(a \cos(c + dx) + a)^{3/2}} \\
 & \quad \downarrow \text{3128} \\
 & \frac{\sin(c + dx)}{2d(a \cos(c + dx) + a)^{3/2}} - \frac{\int \frac{1}{2a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{2ad} \\
 & \quad \downarrow \text{219} \\
 & \frac{\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{\sin(c + dx)}{2d(a \cos(c + dx) + a)^{3/2}}
 \end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^(-3/2), x]`

output `ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])]/(2*Sqrt[2]*a^(3/2)*d) + Sin[c + d*x]/(2*d*(a + a*Cos[c + d*x])^(3/2))`



## 3.6.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3129 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*Cos[c + d*x]*((a + b*Sin[c + d*x])^n/(a*d*(2*n + 1))), x] + Simp[(n + 1)/(a*(2*n + 1)) Int[(a + b*Sin[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

## 3.6.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 137 vs.  $2(62) = 124$ .

Time = 1.03 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.79

method	result	size
default	$\frac{\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left( \sqrt{2} \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4a}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sqrt{2}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\sqrt{a} \right)}{4a^{\frac{5}{2}} \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$	138

input `int(1/(a+cos(d*x+c)*a)^(3/2),x,method=_RETURNVERBOSE)`

output `1/4/a^(5/2)/cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2^(1/2)*ln(2*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2*c))*a*cos(1/2*d*x+1/2*c)^2+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2))/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d`

### 3.6.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 153 vs. 2(62) = 124.

Time = 0.27 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.99

$$\int \frac{1}{(a + a \cos(c + dx))^{3/2}} dx = \frac{\sqrt{2}(\cos(dx + c)^2 + 2 \cos(dx + c) + 1) \sqrt{a} \log\left(-\frac{a \cos(dx+c)^2 - 2\sqrt{2}\sqrt{a \cos(dx+c)+a}}{\cos(dx+c)^2 + 2}\right)}{8(a^2 d \cos(dx + c)^2 + 2a^2 d \cos(dx + c))}$$

input `integrate(1/(a+a*cos(d*x+c))^(3/2),x, algorithm="fricas")`

output `1/8*(sqrt(2)*(cos(d*x + c)^2 + 2*cos(d*x + c) + 1)*sqrt(a)*log(-(a*cos(d*x + c)^2 - 2*sqrt(2)*sqrt(a*cos(d*x + c) + a))*sqrt(a)*sin(d*x + c) - 2*a*cos(d*x + c) - 3*a)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1)) + 4*sqrt(a*cos(d*x + c) + a)*sin(d*x + c))/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`

### 3.6.6 Sympy [F]

$$\int \frac{1}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{1}{(a \cos(c + dx) + a)^{3/2}} dx$$

input `integrate(1/(a+a*cos(d*x+c))**(3/2),x)`

output `Integral((a*cos(c + d*x) + a)**(-3/2), x)`

### 3.6.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15721 vs. 2(62) = 124.

Time = 1.03 (sec) , antiderivative size = 15721, normalized size of antiderivative = 204.17

$$\int \frac{1}{(a + a \cos(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate(1/(a+a*cos(d*x+c))^(3/2),x, algorithm="maxima")`

output `1/4*(32*(cos(3/2*d*x + 3/2*c)*sin(2*d*x + 2*c) + cos(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c) + cos(d*x + c)*sin(3/2*d*x + 3/2*c) + cos(3/2*d*x + 3/2*c)*sin(d*x + c))*cos(3*d*x + 3*c)^2 + 96*(cos(3/2*d*x + 3/2*c)*sin(3*d*x + 3*c) + 3*cos(3/2*d*x + 3/2*c)*sin(2*d*x + 2*c) - (3*cos(d*x + c) + 1)*sin(3/2*d*x + 3/2*c) - cos(3*d*x + 3*c)*sin(3/2*d*x + 3/2*c) - 3*cos(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c) + 3*cos(3/2*d*x + 3/2*c)*sin(d*x + c))*cos(4/3*arctan2(sin(3/2*d*x + 3/2*c), cos(3/2*d*x + 3/2*c)))^2 + 96*(cos(3/2*d*x + 3/2*c)*sin(3*d*x + 3*c) + 3*cos(3/2*d*x + 3/2*c)*sin(2*d*x + 2*c) - (3*cos(d*x + c) + 1)*sin(3/2*d*x + 3/2*c) - cos(3*d*x + 3*c)*sin(3/2*d*x + 3/2*c) - 3*cos(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c) + 3*cos(3/2*d*x + 3/2*c)*sin(d*x + c))*cos(2/3*arctan2(sin(3/2*d*x + 3/2*c), cos(3/2*d*x + 3/2*c)))^2 - 32*(cos(3/2*d*x + 3/2*c)*sin(2*d*x + 2*c) + cos(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c) + cos(d*x + c)*sin(3/2*d*x + 3/2*c) + cos(3/2*d*x + 3/2*c)*sin(d*x + c))*sin(3*d*x + 3*c)^2 + 32*(6*cos(d*x + c) + 1)*cos(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c) + 96*cos(2*d*x + 2*c)^2*sin(3/2*d*x + 3/2*c) + 96*sin(2*d*x + 2*c)^2*sin(3/2*d*x + 3/2*c) + 96*(cos(3/2*d*x + 3/2*c)*sin(3*d*x + 3*c) + 3*cos(3/2*d*x + 3/2*c)*sin(2*d*x + 2*c) - (3*cos(d*x + c) + 1)*sin(3/2*d*x + 3/2*c) - cos(3*d*x + 3*c)*sin(3/2*d*x + 3/2*c) - 3*cos(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c) + 3*cos(3/2*d*x + 3/2*c)*sin(d*x + c))*sin(4/3*arctan2(sin(3/2*d*x + 3/2*c), cos(3/2*d*x + 3/2*c)))^2 + 96*(cos(3/2*d*x + 3/2*c)...`

### 3.6.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.45

$$\int \frac{1}{(a + a \cos(c + dx))^{3/2}} dx = \frac{\sqrt{2} \left( \frac{\log(\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{\operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{\log(-\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{\operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{2 \sin(\frac{1}{2} dx + \frac{1}{2} c)}{(\sin(\frac{1}{2} dx + \frac{1}{2} c)^2 - 1) \operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} \right)}{8 \sqrt{ad}}$$

input `integrate(1/(a+a*cos(d*x+c))^(3/2),x, algorithm="giac")`

output `1/8*sqrt(2)*(log(sin(1/2*d*x + 1/2*c) + 1)/(a*sgn(cos(1/2*d*x + 1/2*c))) - log(-sin(1/2*d*x + 1/2*c) + 1)/(a*sgn(cos(1/2*d*x + 1/2*c))) - 2*sin(1/2*d*x + 1/2*c)/((sin(1/2*d*x + 1/2*c)^2 - 1)*a*sgn(cos(1/2*d*x + 1/2*c))))/(sqrt(a)*d)`

**3.6.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{1}{(a + a \cos(c + dx))^{3/2}} dx$$

input `int(1/(a + a*cos(c + d*x))^(3/2), x)`output `int(1/(a + a*cos(c + d*x))^(3/2), x)`

$$3.7 \quad \int \frac{1}{(a+a \cos(c+dx))^{5/2}} dx$$

3.7.1	Optimal result . . . . .	76
3.7.2	Mathematica [A] (verified) . . . . .	76
3.7.3	Rubi [A] (verified) . . . . .	77
3.7.4	Maple [A] (verified) . . . . .	79
3.7.5	Fricas [B] (verification not implemented) . . . . .	79
3.7.6	Sympy [F] . . . . .	80
3.7.7	Maxima [B] (verification not implemented) . . . . .	80
3.7.8	Giac [A] (verification not implemented) . . . . .	81
3.7.9	Mupad [F(-1)] . . . . .	81

### 3.7.1 Optimal result

Integrand size = 14, antiderivative size = 107

$$\int \frac{1}{(a+a \cos(c+dx))^{5/2}} dx = \frac{3\arctanh\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d} + \frac{\sin(c+dx)}{4d(a+a \cos(c+dx))^{5/2}} + \frac{3 \sin(c+dx)}{16ad(a+a \cos(c+dx))^{3/2}}$$

output `1/4*sin(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)+3/16*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(3/2)+3/32*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c)))/a^(5/2)/d*2^(1/2)`

### 3.7.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.61

$$\int \frac{1}{(a+a \cos(c+dx))^{5/2}} dx = \frac{24\arctanh\left(\sin\left(\frac{1}{2}(c+dx)\right)\right) \cos^5\left(\frac{1}{2}(c+dx)\right) + 14 \sin(c+dx) + 3 \sin(2(c+dx))}{32d(a(1+\cos(c+dx)))^{5/2}}$$

input `Integrate[(a + a*Cos[c + d*x])^(-5/2), x]`

output `(24*ArcTanh[Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^5 + 14*Sin[c + d*x] + 3*Sin[2*(c + d*x)])/(32*d*(a*(1 + Cos[c + d*x]))^(5/2))`

### 3.7.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 3129, 3042, 3129, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \cos(c + dx) + a)^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2}} dx \\
 & \quad \downarrow \text{3129} \\
 & \frac{3 \int \frac{1}{(\cos(c+dx)a+a)^{3/2}} dx}{8a} + \frac{\sin(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3 \int \frac{1}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a} + \frac{\sin(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
 & \quad \downarrow \text{3129} \\
 & \frac{3 \left( \frac{\int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx}{4a} + \frac{\sin(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a} + \frac{\sin(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3 \left( \frac{\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a} + \frac{\sin(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a} + \frac{\sin(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
 & \quad \downarrow \text{3128} \\
 & \frac{3 \left( \frac{\sin(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} - \frac{\int \frac{1}{2a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d \left( -\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}} \right)}{2ad} \right)}{8a} + \frac{\sin(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
 & \quad \downarrow \text{219}
 \end{aligned}$$

$$\frac{3 \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{\sin(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a} + \frac{\sin(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}}$$

input `Int[(a + a*Cos[c + d*x])^(-5/2), x]`

output `Sin[c + d*x]/(4*d*(a + a*Cos[c + d*x])^(5/2)) + (3*(ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])]/(2*Sqrt[2]*a^(3/2)*d) + Sin[c + d*x]/(2*d*(a + a*Cos[c + d*x])^(3/2)))/(8*a)`

### 3.7.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3129 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*Cos[c + d*x]*((a + b*Sin[c + d*x])^n/(a*d*(2*n + 1))), x] + Simp[(n + 1)/(a*(2*n + 1)) Int[(a + b*Sin[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

### 3.7.4 Maple [A] (verified)

Time = 1.10 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.63

method	result
default	$\frac{\sqrt{a \left( \sin^2 \left( \frac{dx}{2} + \frac{c}{2} \right) \right)} \left( 3\sqrt{2} \ln \left( \frac{4\sqrt{a} \sqrt{a \left( \sin^2 \left( \frac{dx}{2} + \frac{c}{2} \right) \right) + 4a}}{\cos \left( \frac{dx}{2} + \frac{c}{2} \right)} \right) a \left( \cos^4 \left( \frac{dx}{2} + \frac{c}{2} \right) \right) + 3 \left( \cos^2 \left( \frac{dx}{2} + \frac{c}{2} \right) \right) \sqrt{a \left( \sin^2 \left( \frac{dx}{2} + \frac{c}{2} \right) \right)} \sqrt{2} \sqrt{a} + 2\sqrt{2} \sqrt{a} \right)}{32a^{\frac{7}{2}} \cos \left( \frac{dx}{2} + \frac{c}{2} \right)^3 \sin \left( \frac{dx}{2} + \frac{c}{2} \right) \sqrt{a \left( \cos^2 \left( \frac{dx}{2} + \frac{c}{2} \right) \right)}} d$

input `int(1/(a+cos(d*x+c)*a)^(5/2),x,method=_RETURNVERBOSE)`

output `1/32/a^(7/2)/cos(1/2*d*x+1/2*c)^3*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(3*2^(1/2)*ln(2*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2*c))*a*cos(1/2*d*x+1/2*c)^4+3*cos(1/2*d*x+1/2*c)^2*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*2^(1/2)*a^(1/2)+2*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2))/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d`

### 3.7.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 188 vs. 2(88) = 176.

Time = 0.28 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.76

$$\int \frac{1}{(a + a \cos(c + dx))^{5/2}} dx = \frac{3\sqrt{2}(\cos(dx + c)^3 + 3 \cos(dx + c)^2 + 3 \cos(dx + c) + 1) \sqrt{a} \log \left( -\frac{a \cos(dx + c)}{2a \cos(dx + c) + a} \right)}{64 (a^3 d \cos(dx + c)^3 - \dots)}$$

input `integrate(1/(a+a*cos(d*x+c))^(5/2),x, algorithm="fracas")`

output `1/64*(3*sqrt(2)*(cos(d*x + c)^3 + 3*cos(d*x + c)^2 + 3*cos(d*x + c) + 1)*sqrt(a)*log(-a*cos(d*x + c)^2 - 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sin(d*x + c) - 2*a*cos(d*x + c) - 3*a)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1)) + 4*sqrt(a*cos(d*x + c) + a)*(3*cos(d*x + c) + 7)*sin(d*x + c))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`



### 3.7.6 Sympy [F]

$$\int \frac{1}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{1}{(a \cos(c + dx) + a)^{5/2}} dx$$

input `integrate(1/(a+a*cos(d*x+c))**(5/2),x)`

output `Integral((a*cos(c + d*x) + a)**(-5/2), x)`

### 3.7.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 84332 vs. 2(88) = 176.

Time = 16.47 (sec) , antiderivative size = 84332, normalized size of antiderivative = 788.15

$$\int \frac{1}{(a + a \cos(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(1/(a+a*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `1/32*(512*((2*sin(2*d*x + 2*c) + sin(d*x + c))*cos(5/2*d*x + 5/2*c) + cos(5/2*d*x + 5/2*c)*sin(4*d*x + 4*c) + 2*cos(5/2*d*x + 5/2*c)*sin(3*d*x + 3*c) + (2*cos(2*d*x + 2*c) + cos(d*x + c))*sin(5/2*d*x + 5/2*c) + cos(4*d*x + 4*c)*sin(5/2*d*x + 5/2*c) + 2*cos(3*d*x + 3*c)*sin(5/2*d*x + 5/2*c))*cos(5*d*x + 5*c)^2 + 2560*(5*(2*sin(2*d*x + 2*c) + sin(d*x + c))*cos(5/2*d*x + 5/2*c) + cos(5/2*d*x + 5/2*c)*sin(5*d*x + 5*c) + 5*cos(5/2*d*x + 5/2*c)*sin(4*d*x + 4*c) + 10*cos(5/2*d*x + 5/2*c)*sin(3*d*x + 3*c) - (10*cos(2*d*x + 2*c) + 5*cos(d*x + c) + 1)*sin(5/2*d*x + 5/2*c) - cos(5*d*x + 5*c)*sin(5/2*d*x + 5/2*c) - 5*cos(4*d*x + 4*c)*sin(5/2*d*x + 5/2*c) - 10*cos(3*d*x + 3*c)*sin(5/2*d*x + 5/2*c))*cos(8/5*arctan2(sin(5/2*d*x + 5/2*c), cos(5/2*d*x + 5/2*c)))^2 + 10240*(5*(2*sin(2*d*x + 2*c) + sin(d*x + c))*cos(5/2*d*x + 5/2*c) + cos(5/2*d*x + 5/2*c)*sin(5*d*x + 5*c) + 5*cos(5/2*d*x + 5/2*c)*sin(4*d*x + 4*c) + 10*cos(5/2*d*x + 5/2*c)*sin(3*d*x + 3*c) - (10*cos(2*d*x + 2*c) + 5*cos(d*x + c) + 1)*sin(5/2*d*x + 5/2*c) - cos(5*d*x + 5*c)*sin(5/2*d*x + 5/2*c) - 5*cos(4*d*x + 4*c)*sin(5/2*d*x + 5/2*c) - 10*cos(3*d*x + 3*c)*sin(5/2*d*x + 5/2*c))*cos(6/5*arctan2(sin(5/2*d*x + 5/2*c), cos(5/2*d*x + 5/2*c)))^2 + 10240*(5*(2*sin(2*d*x + 2*c) + sin(d*x + c))*cos(5/2*d*x + 5/2*c) + cos(5/2*d*x + 5/2*c)*sin(5*d*x + 5*c) + 5*cos(5/2*d*x + 5/2*c)*sin(4*d*x + 4*c) + 10*cos(5/2*d*x + 5/2*c)*sin(3*d*x + 3*c) - (10*cos(2*d*x + 2*c) + 5*cos(d*x + c) + 1)*sin(5/2*d*x + 5/2*c) - cos(5*d*x ...`

### 3.7.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.21

$$\int \frac{1}{(a + a \cos(c + dx))^{5/2}} dx = \frac{\sqrt{2} \left( \frac{3 \log(\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{3 \log(-\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{2 \left( 3 \sin(\frac{1}{2} dx + \frac{1}{2} c)^3 - 5 \sin(\frac{1}{2} dx + \frac{1}{2} c) \right)}{\left( \sin(\frac{1}{2} dx + \frac{1}{2} c)^2 - 1 \right)^2 a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} \right)}{64 \sqrt{ad}}$$

input `integrate(1/(a+a*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `1/64*sqrt(2)*(3*log(sin(1/2*d*x + 1/2*c) + 1)/(a^2*sgn(cos(1/2*d*x + 1/2*c))) - 3*log(-sin(1/2*d*x + 1/2*c) + 1)/(a^2*sgn(cos(1/2*d*x + 1/2*c))) - 2*(3*sin(1/2*d*x + 1/2*c)^3 - 5*sin(1/2*d*x + 1/2*c))/((sin(1/2*d*x + 1/2*c)^2 - 1)^2*a^2*sgn(cos(1/2*d*x + 1/2*c))))/(sqrt(a)*d)`

### 3.7.9 Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{1}{(a + a \cos(c + dx))^{5/2}} dx$$

input `int(1/(a + a*cos(c + d*x))^(5/2),x)`

output `int(1/(a + a*cos(c + d*x))^(5/2), x)`

### 3.8 $\int (a + a \cos(c + dx))^{4/3} dx$

3.8.1	Optimal result . . . . .	82
3.8.2	Mathematica [A] (verified) . . . . .	82
3.8.3	Rubi [A] (verified) . . . . .	83
3.8.4	Maple [F] . . . . .	84
3.8.5	Fricas [F] . . . . .	84
3.8.6	Sympy [F] . . . . .	85
3.8.7	Maxima [F] . . . . .	85
3.8.8	Giac [F] . . . . .	85
3.8.9	Mupad [F(-1)] . . . . .	86

#### 3.8.1 Optimal result

Integrand size = 14, antiderivative size = 67

$$\int (a + a \cos(c + dx))^{4/3} dx = \frac{2 \cdot 2^{5/6} a^3 \sqrt{a + a \cos(c + dx)} \operatorname{Hypergeometric2F1}\left(-\frac{5}{6}, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx))\right) \sin(c + dx)}{d(1 + \cos(c + dx))^{5/6}}$$

output `2*2^(5/6)*a*(a+a*cos(d*x+c))^(1/3)*hypergeom([-5/6, 1/2], [3/2], 1/2-1/2*cos(d*x+c))*sin(d*x+c)/d/(1+cos(d*x+c))^(5/6)`

#### 3.8.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.03

$$\int (a + a \cos(c + dx))^{4/3} dx = \frac{6(a(1 + \cos(c + dx)))^{4/3} \cot\left(\frac{1}{2}(c + dx)\right) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{11}{6}, \frac{17}{6}, \cos^2\left(\frac{1}{2}(c + dx)\right)\right) \sqrt{\sin^2\left(\frac{1}{2}(c + dx)\right)}}{11d}$$

input `Integrate[(a + a*Cos[c + d*x])^(4/3), x]`

output `(-6*(a*(1 + Cos[c + d*x]))^(4/3)*Cot[(c + d*x)/2]*Hypergeometric2F1[1/2, 1/6, 17/6, Cos[(c + d*x)/2]^2]*Sqrt[Sin[(c + d*x)/2]^2])/(11*d)`

### 3.8.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \cos(c + dx) + a)^{4/3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( a \sin \left( c + dx + \frac{\pi}{2} \right) + a \right)^{4/3} dx \\
 & \quad \downarrow \text{3131} \\
 & \frac{a \sqrt[3]{a \cos(c + dx) + a} \int (\cos(c + dx) + 1)^{4/3} dx}{\sqrt[3]{\cos(c + dx) + 1}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a \sqrt[3]{a \cos(c + dx) + a} \int \left( \sin \left( c + dx + \frac{\pi}{2} \right) + 1 \right)^{4/3} dx}{\sqrt[3]{\cos(c + dx) + 1}} \\
 & \quad \downarrow \text{3130} \\
 & \frac{2 \cdot 2^{5/6} a \sin(c + dx) \sqrt[3]{a \cos(c + dx) + a} \operatorname{Hypergeometric2F1} \left( -\frac{5}{6}, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx)) \right)}{d(\cos(c + dx) + 1)^{5/6}}
 \end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^(4/3),x]`

output `(2*2^(5/6)*a*(a + a*Cos[c + d*x])^(1/3)*Hypergeometric2F1[-5/6, 1/2, 3/2, (1 - Cos[c + d*x])/2]*Sin[c + d*x])/(d*(1 + Cos[c + d*x])^(5/6))`

## 3.8.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]))]*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a))], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

rule 3131 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[n]*((a + b*Sin[c + d*x])^FracPart[n]/(1 + (b/a)*Sin[c + d*x])^FracPart[n]) Int[(1 + (b/a)*Sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && !GtQ[a, 0]`

## 3.8.4 Maple [F]

$$\int (a + \cos(dx + c)) a^{\frac{4}{3}} dx$$

input `int((a+cos(d*x+c))*a^(4/3),x)`

output `int((a+cos(d*x+c))*a^(4/3),x)`

## 3.8.5 Fricas [F]

$$\int (a + a \cos(c + dx))^{4/3} dx = \int (a \cos(dx + c) + a)^{\frac{4}{3}} dx$$

input `integrate((a+a*cos(d*x+c))^(4/3),x, algorithm="fricas")`

output `integral((a*cos(d*x + c) + a)^(4/3), x)`

**3.8.6 Sympy [F]**

$$\int (a + a \cos(c + dx))^{4/3} dx = \int (a \cos(c + dx) + a)^{\frac{4}{3}} dx$$

input `integrate((a+a*cos(d*x+c))**(4/3),x)`

output `Integral((a*cos(c + d*x) + a)**(4/3), x)`

**3.8.7 Maxima [F]**

$$\int (a + a \cos(c + dx))^{4/3} dx = \int (a \cos(dx + c) + a)^{\frac{4}{3}} dx$$

input `integrate((a+a*cos(d*x+c))^(4/3),x, algorithm="maxima")`

output `integrate((a*cos(d*x + c) + a)^(4/3), x)`

**3.8.8 Giac [F]**

$$\int (a + a \cos(c + dx))^{4/3} dx = \int (a \cos(dx + c) + a)^{\frac{4}{3}} dx$$

input `integrate((a+a*cos(d*x+c))^(4/3),x, algorithm="giac")`

output `integrate((a*cos(d*x + c) + a)^(4/3), x)`

**3.8.9 Mupad [F(-1)]**

Timed out.

$$\int (a + a \cos(c + dx))^{4/3} dx = \int (a + a \cos(c + dx))^{4/3} dx$$

input `int((a + a*cos(c + d*x))^(4/3),x)`output `int((a + a*cos(c + d*x))^(4/3), x)`

### 3.9 $\int (a + a \cos(c + dx))^{2/3} dx$

3.9.1	Optimal result . . . . .	87
3.9.2	Mathematica [A] (verified) . . . . .	87
3.9.3	Rubi [A] (verified) . . . . .	88
3.9.4	Maple [F] . . . . .	89
3.9.5	Fricas [F] . . . . .	89
3.9.6	Sympy [F] . . . . .	90
3.9.7	Maxima [F] . . . . .	90
3.9.8	Giac [F] . . . . .	90
3.9.9	Mupad [F(-1)] . . . . .	91

#### 3.9.1 Optimal result

Integrand size = 14, antiderivative size = 66

$$\int (a + a \cos(c + dx))^{2/3} dx = \frac{2\sqrt[6]{2}(a + a \cos(c + dx))^{2/3} \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx))\right) \sin(c + dx)}{d(1 + \cos(c + dx))^{7/6}}$$

output `2*2^(1/6)*(a+a*cos(d*x+c))^(2/3)*hypergeom([-1/6, 1/2], [3/2], 1/2-1/2*cos(d*x+c))*sin(d*x+c)/d/(1+cos(d*x+c))^(7/6)`

#### 3.9.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.05

$$\int (a + a \cos(c + dx))^{2/3} dx = \frac{6(a(1 + \cos(c + dx)))^{2/3} \cot\left(\frac{1}{2}(c + dx)\right) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2\left(\frac{1}{2}(c + dx)\right)\right) \sqrt{\sin^2\left(\frac{1}{2}(c + dx)\right)}}{7d}$$

input `Integrate[(a + a*Cos[c + d*x])^(2/3), x]`

output `(-6*(a*(1 + Cos[c + d*x]))^(2/3)*Cot[(c + d*x)/2]*Hypergeometric2F1[1/2, 7/6, 13/6, Cos[(c + d*x)/2]^2]*Sqrt[Sin[(c + d*x)/2]^2])/(7*d)`



### 3.9.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \cos(c + dx) + a)^{2/3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( a \sin \left( c + dx + \frac{\pi}{2} \right) + a \right)^{2/3} dx \\
 & \quad \downarrow \text{3131} \\
 & \frac{(a \cos(c + dx) + a)^{2/3} \int (\cos(c + dx) + 1)^{2/3} dx}{(\cos(c + dx) + 1)^{2/3}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(a \cos(c + dx) + a)^{2/3} \int \left( \sin \left( c + dx + \frac{\pi}{2} \right) + 1 \right)^{2/3} dx}{(\cos(c + dx) + 1)^{2/3}} \\
 & \quad \downarrow \text{3130} \\
 & \frac{2^{\frac{6}{\sqrt{2}}} \sin(c + dx) (a \cos(c + dx) + a)^{2/3} \text{Hypergeometric2F1} \left( -\frac{1}{6}, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx)) \right)}{d(\cos(c + dx) + 1)^{7/6}}
 \end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^(2/3),x]`

output `(2*2^(1/6)*(a + a*Cos[c + d*x])^(2/3)*Hypergeometric2F1[-1/6, 1/2, 3/2, (1 - Cos[c + d*x])/2]*Sin[c + d*x])/(d*(1 + Cos[c + d*x])^(7/6))`

### 3.9.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]]))*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a))], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

rule 3131 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[n]*((a + b*Sin[c + d*x])^FracPart[n]/(1 + (b/a)*Sin[c + d*x])^FracPart[n]) Int[(1 + (b/a)*Sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && !GtQ[a, 0]`

### 3.9.4 Maple [F]

$$\int (a + \cos(dx + c)a)^{\frac{2}{3}} dx$$

input `int((a+cos(d*x+c)*a)^(2/3),x)`

output `int((a+cos(d*x+c)*a)^(2/3),x)`

### 3.9.5 Fracas [F]

$$\int (a + a \cos(c + dx))^{2/3} dx = \int (a \cos(dx + c) + a)^{\frac{2}{3}} dx$$

input `integrate((a+a*cos(d*x+c))^(2/3),x, algorithm="fracas")`

output `integral((a*cos(d*x + c) + a)^(2/3), x)`

**3.9.6 Sympy [F]**

$$\int (a + a \cos(c + dx))^{2/3} dx = \int (a \cos(c + dx) + a)^{\frac{2}{3}} dx$$

input `integrate((a+a*cos(d*x+c))**(2/3),x)`

output `Integral((a*cos(c + d*x) + a)**(2/3), x)`

**3.9.7 Maxima [F]**

$$\int (a + a \cos(c + dx))^{2/3} dx = \int (a \cos(dx + c) + a)^{\frac{2}{3}} dx$$

input `integrate((a+a*cos(d*x+c))^(2/3),x, algorithm="maxima")`

output `integrate((a*cos(d*x + c) + a)^(2/3), x)`

**3.9.8 Giac [F]**

$$\int (a + a \cos(c + dx))^{2/3} dx = \int (a \cos(dx + c) + a)^{\frac{2}{3}} dx$$

input `integrate((a+a*cos(d*x+c))^(2/3),x, algorithm="giac")`

output `integrate((a*cos(d*x + c) + a)^(2/3), x)`

**3.9.9 Mupad [F(-1)]**

Timed out.

$$\int (a + a \cos(c + dx))^{2/3} dx = \int (a + a \cos(c + dx))^{2/3} dx$$

input `int((a + a*cos(c + d*x))^(2/3),x)`output `int((a + a*cos(c + d*x))^(2/3), x)`

### 3.10 $\int \sqrt[3]{a + a \cos(c + dx)} dx$

3.10.1	Optimal result . . . . .	92
3.10.2	Mathematica [A] (verified) . . . . .	92
3.10.3	Rubi [A] (verified) . . . . .	93
3.10.4	Maple [F] . . . . .	94
3.10.5	Fricas [F] . . . . .	94
3.10.6	Sympy [F] . . . . .	95
3.10.7	Maxima [F] . . . . .	95
3.10.8	Giac [F] . . . . .	95
3.10.9	Mupad [F(-1)] . . . . .	96

#### 3.10.1 Optimal result

Integrand size = 14, antiderivative size = 65

$$\int \sqrt[3]{a + a \cos(c + dx)} dx = \frac{2^{5/6} \sqrt[3]{a + a \cos(c + dx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx))\right) \sin(c + dx)}{d(1 + \cos(c + dx))^{5/6}}$$

output `2^(5/6)*(a+a*cos(d*x+c))^(1/3)*hypergeom([1/6, 1/2],[3/2],1/2-1/2*cos(d*x+c))*sin(d*x+c)/d/(1+cos(d*x+c))^(5/6)`

#### 3.10.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.06

$$\int \sqrt[3]{a + a \cos(c + dx)} dx = \frac{6 \sqrt[3]{a(1 + \cos(c + dx))} \cot\left(\frac{1}{2}(c + dx)\right) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2\left(\frac{1}{2}(c + dx)\right)\right) \sqrt{\sin^2\left(\frac{1}{2}(c + dx)\right)}}{5d}$$

input `Integrate[(a + a*Cos[c + d*x])^(1/3),x]`

output `(-6*(a*(1 + Cos[c + d*x]))^(1/3)*Cot[(c + d*x)/2]*Hypergeometric2F1[1/2, 5/6, 11/6, Cos[(c + d*x)/2]^2]*Sqrt[Sin[(c + d*x)/2]^2])/(5*d)`

### 3.10.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt[3]{a \cos(c + dx) + a} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt[3]{a \sin\left(c + dx + \frac{\pi}{2}\right) + a} dx \\
 & \quad \downarrow \text{3131} \\
 & \frac{\sqrt[3]{a \cos(c + dx) + a} \int \sqrt[3]{\cos(c + dx) + 1} dx}{\sqrt[3]{\cos(c + dx) + 1}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt[3]{a \cos(c + dx) + a} \int \sqrt[3]{\sin\left(c + dx + \frac{\pi}{2}\right) + 1} dx}{\sqrt[3]{\cos(c + dx) + 1}} \\
 & \quad \downarrow \text{3130} \\
 & \frac{2^{5/6} \sin(c + dx) \sqrt[3]{a \cos(c + dx) + a} \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx))\right)}{d(\cos(c + dx) + 1)^{5/6}}
 \end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^(1/3),x]`

output `(2^(5/6)*(a + a*Cos[c + d*x])^(1/3)*Hypergeometric2F1[1/6, 1/2, 3/2, (1 - Cos[c + d*x])/2]*Sin[c + d*x])/(d*(1 + Cos[c + d*x])^(5/6))`

## 3.10.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]]))*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a))], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

rule 3131 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[n]*((a + b*Sin[c + d*x])^FracPart[n]/(1 + (b/a)*Sin[c + d*x])^FracPart[n]) Int[(1 + (b/a)*Sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && !GtQ[a, 0]`

## 3.10.4 Maple [F]

$$\int (a + \cos(dx + c)a)^{\frac{1}{3}} dx$$

input `int((a+cos(d*x+c)*a)^(1/3),x)`

output `int((a+cos(d*x+c)*a)^(1/3),x)`

## 3.10.5 Fracas [F]

$$\int \sqrt[3]{a + a \cos(c + dx)} dx = \int (a \cos(dx + c) + a)^{\frac{1}{3}} dx$$

input `integrate((a+a*cos(d*x+c))^(1/3),x, algorithm="fricas")`

output `integral((a*cos(d*x + c) + a)^(1/3), x)`

**3.10.6 Sympy [F]**

$$\int \sqrt[3]{a + a \cos(c + dx)} dx = \int \sqrt[3]{a \cos(c + dx) + a} dx$$

input `integrate((a+a*cos(d*x+c))**(1/3),x)`

output `Integral((a*cos(c + d*x) + a)**(1/3), x)`

**3.10.7 Maxima [F]**

$$\int \sqrt[3]{a + a \cos(c + dx)} dx = \int (a \cos(dx + c) + a)^{\frac{1}{3}} dx$$

input `integrate((a+a*cos(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate((a*cos(d*x + c) + a)^(1/3), x)`

**3.10.8 Giac [F]**

$$\int \sqrt[3]{a + a \cos(c + dx)} dx = \int (a \cos(dx + c) + a)^{\frac{1}{3}} dx$$

input `integrate((a+a*cos(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate((a*cos(d*x + c) + a)^(1/3), x)`



**3.10.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt[3]{a + a \cos(c + dx)} dx = \int (a + a \cos(c + dx))^{1/3} dx$$

input `int((a + a*cos(c + d*x))^(1/3),x)`output `int((a + a*cos(c + d*x))^(1/3), x)`

### 3.11 $\int \frac{1}{\sqrt[3]{a + a \cos(c + dx)}} dx$

3.11.1	Optimal result	97
3.11.2	Mathematica [A] (verified)	97
3.11.3	Rubi [A] (verified)	98
3.11.4	Maple [F]	99
3.11.5	Fricas [F]	99
3.11.6	Sympy [F]	100
3.11.7	Maxima [F]	100
3.11.8	Giac [F]	100
3.11.9	Mupad [F(-1)]	101

#### 3.11.1 Optimal result

Integrand size = 14, antiderivative size = 65

$$\int \frac{1}{\sqrt[3]{a + a \cos(c + dx)}} dx = \frac{\sqrt[6]{2} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx))\right) \sin(c + dx)}{d \sqrt[6]{1 + \cos(c + dx)} \sqrt[3]{a + a \cos(c + dx)}}$$

```
output 2^(1/6)*hypergeom([1/2, 5/6], [3/2], 1/2-1/2*cos(d*x+c))*sin(d*x+c)/d/(1+cos
(d*x+c))^(1/6)/(a+a*cos(d*x+c))^(1/3)
```

#### 3.11.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.03

$$\int \frac{1}{\sqrt[3]{a + a \cos(c + dx)}} dx = -\frac{6 \cot\left(\frac{1}{2}(c + dx)\right) \text{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2\left(\frac{1}{2}(c + dx)\right)\right) \sqrt{\sin^2\left(\frac{1}{2}(c + dx)\right)}}{d \sqrt[3]{a(1 + \cos(c + dx))}}$$

```
input Integrate[(a + a*Cos[c + d*x])^(-1/3), x]
```

```
output (-6*Cot[(c + d*x)/2]*Hypergeometric2F1[1/6, 1/2, 7/6, Cos[(c + d*x)/2]^2]*
Sqrt[Sin[(c + d*x)/2]^2])/(d*(a*(1 + Cos[c + d*x]))^(1/3))
```

### 3.11.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt[3]{a \cos(c + dx) + a}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt[3]{a \sin\left(c + dx + \frac{\pi}{2}\right) + a}} dx \\
 & \quad \downarrow \text{3131} \\
 & \frac{\sqrt[3]{\cos(c + dx) + 1} \int \frac{1}{\sqrt[3]{\cos(c + dx) + 1}} dx}{\sqrt[3]{a \cos(c + dx) + a}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt[3]{\cos(c + dx) + 1} \int \frac{1}{\sqrt[3]{\sin\left(c + dx + \frac{\pi}{2}\right) + 1}} dx}{\sqrt[3]{a \cos(c + dx) + a}} \\
 & \quad \downarrow \text{3130} \\
 & \frac{\sqrt[6]{2} \sin(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx))\right)}{d \sqrt[6]{\cos(c + dx) + 1} \sqrt[3]{a \cos(c + dx) + a}}
 \end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^(-1/3), x]`

output `(2^(1/6)*Hypergeometric2F1[1/2, 5/6, 3/2, (1 - Cos[c + d*x])/2]*Sin[c + d*x])/(d*(1 + Cos[c + d*x])^(1/6)*(a + a*Cos[c + d*x])^(1/3))`

---

3.11.  $\int \frac{1}{\sqrt[3]{a + a \cos(c + dx)}} dx$

## 3.11.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]]))*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a))], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

rule 3131 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[n]*((a + b*Sin[c + d*x])^FracPart[n]/(1 + (b/a)*Sin[c + d*x])^FracPart[n]) Int[(1 + (b/a)*Sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && !GtQ[a, 0]`

## 3.11.4 Maple [F]

$$\int \frac{1}{(a + \cos(dx + c)a)^{\frac{1}{3}}} dx$$

input `int(1/(a+cos(d*x+c)*a)^(1/3),x)`

output `int(1/(a+cos(d*x+c)*a)^(1/3),x)`

## 3.11.5 Fricas [F]

$$\int \frac{1}{\sqrt[3]{a + a \cos(c + dx)}} dx = \int \frac{1}{(a \cos(dx + c) + a)^{\frac{1}{3}}} dx$$

input `integrate(1/(a+a*cos(d*x+c))^(1/3),x, algorithm="fricas")`

output `integral((a*cos(d*x + c) + a)^(-1/3), x)`

**3.11.6 Sympy [F]**

$$\int \frac{1}{\sqrt[3]{a + a \cos(c + dx)}} dx = \int \frac{1}{\sqrt[3]{a \cos(c + dx) + a}} dx$$

input `integrate(1/(a+a*cos(d*x+c))**(1/3),x)`

output `Integral((a*cos(c + d*x) + a)**(-1/3), x)`

**3.11.7 Maxima [F]**

$$\int \frac{1}{\sqrt[3]{a + a \cos(c + dx)}} dx = \int \frac{1}{(a \cos(dx + c) + a)^{\frac{1}{3}}} dx$$

input `integrate(1/(a+a*cos(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate((a*cos(d*x + c) + a)^(-1/3), x)`

**3.11.8 Giac [F]**

$$\int \frac{1}{\sqrt[3]{a + a \cos(c + dx)}} dx = \int \frac{1}{(a \cos(dx + c) + a)^{\frac{1}{3}}} dx$$

input `integrate(1/(a+a*cos(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate((a*cos(d*x + c) + a)^(-1/3), x)`

**3.11.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt[3]{a + a \cos(c + dx)}} dx = \int \frac{1}{(a + a \cos(c + dx))^{1/3}} dx$$

input `int(1/(a + a*cos(c + d*x))^(1/3), x)`output `int(1/(a + a*cos(c + d*x))^(1/3), x)`

### 3.12 $\int \frac{1}{(a+a \cos(c+dx))^{2/3}} dx$

3.12.1	Optimal result	102
3.12.2	Mathematica [A] (verified)	102
3.12.3	Rubi [A] (verified)	103
3.12.4	Maple [F]	104
3.12.5	Fricas [F]	104
3.12.6	Sympy [F]	105
3.12.7	Maxima [F]	105
3.12.8	Giac [F]	105
3.12.9	Mupad [F(-1)]	106

#### 3.12.1 Optimal result

Integrand size = 14, antiderivative size = 65

$$\int \frac{1}{(a + a \cos(c + dx))^{2/3}} dx = \frac{\sqrt[6]{1 + \cos(c + dx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx))\right) \sin(c + dx)}{\sqrt[6]{2d(a + a \cos(c + dx))^{2/3}}}$$

```
output 1/2*(1+cos(d*x+c))^(1/6)*hypergeom([1/2, 7/6], [3/2], 1/2-1/2*cos(d*x+c))*si
n(d*x+c)*2^(5/6)/d/(a+a*cos(d*x+c))^(2/3)
```

#### 3.12.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.03

$$\int \frac{1}{(a + a \cos(c + dx))^{2/3}} dx = \frac{6 \cot\left(\frac{1}{2}(c + dx)\right) \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2\left(\frac{1}{2}(c + dx)\right)\right) \sqrt{\sin^2\left(\frac{1}{2}(c + dx)\right)}}{d(a(1 + \cos(c + dx)))^{2/3}}$$

```
input Integrate[(a + a*Cos[c + d*x])^(-2/3), x]
```

```
output (6*Cot[(c + d*x)/2]*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[(c + d*x)/2]^2]*
Sqrt[Sin[(c + d*x)/2]^2])/(d*(a*(1 + Cos[c + d*x]))^(2/3))
```

### 3.12.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \cos(c + dx) + a)^{2/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a \sin(c + dx + \frac{\pi}{2}) + a)^{2/3}} dx \\
 & \quad \downarrow \text{3131} \\
 & \frac{(\cos(c + dx) + 1)^{2/3} \int \frac{1}{(\cos(c + dx) + 1)^{2/3}} dx}{(a \cos(c + dx) + a)^{2/3}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(\cos(c + dx) + 1)^{2/3} \int \frac{1}{(\sin(c + dx + \frac{\pi}{2}) + 1)^{2/3}} dx}{(a \cos(c + dx) + a)^{2/3}} \\
 & \quad \downarrow \text{3130} \\
 & \frac{\sin(c + dx) \sqrt[6]{\cos(c + dx) + 1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx))\right)}{\sqrt[6]{2d}(a \cos(c + dx) + a)^{2/3}}
 \end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^(-2/3), x]`

output `((1 + Cos[c + d*x])^(1/6)*Hypergeometric2F1[1/2, 7/6, 3/2, (1 - Cos[c + d*x])/2]*Sin[c + d*x])/(2^(1/6)*d*(a + a*Cos[c + d*x])^(2/3))`



## 3.12.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]))]*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a))], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

rule 3131 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[n]*((a + b*Sin[c + d*x])^FracPart[n]/(1 + (b/a)*Sin[c + d*x])^FracPart[n]) Int[(1 + (b/a)*Sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && !GtQ[a, 0]`

## 3.12.4 Maple [F]

$$\int \frac{1}{(a + \cos(dx + c)a)^{\frac{2}{3}}} dx$$

input `int(1/(a+cos(d*x+c)*a)^(2/3),x)`

output `int(1/(a+cos(d*x+c)*a)^(2/3),x)`

## 3.12.5 Fricas [F]

$$\int \frac{1}{(a + a \cos(c + dx))^{2/3}} dx = \int \frac{1}{(a \cos(dx + c) + a)^{\frac{2}{3}}} dx$$

input `integrate(1/(a+a*cos(d*x+c))^(2/3),x, algorithm="fricas")`

output `integral((a*cos(d*x + c) + a)^(-2/3), x)`

**3.12.6 Sympy [F]**

$$\int \frac{1}{(a + a \cos(c + dx))^{2/3}} dx = \int \frac{1}{(a \cos(c + dx) + a)^{2/3}} dx$$

input `integrate(1/(a+a*cos(d*x+c))**(2/3), x)`

output `Integral((a*cos(c + d*x) + a)**(-2/3), x)`

**3.12.7 Maxima [F]**

$$\int \frac{1}{(a + a \cos(c + dx))^{2/3}} dx = \int \frac{1}{(a \cos(dx + c) + a)^{2/3}} dx$$

input `integrate(1/(a+a*cos(d*x+c))^(2/3), x, algorithm="maxima")`

output `integrate((a*cos(d*x + c) + a)^(-2/3), x)`

**3.12.8 Giac [F]**

$$\int \frac{1}{(a + a \cos(c + dx))^{2/3}} dx = \int \frac{1}{(a \cos(dx + c) + a)^{2/3}} dx$$

input `integrate(1/(a+a*cos(d*x+c))^(2/3), x, algorithm="giac")`

output `integrate((a*cos(d*x + c) + a)^(-2/3), x)`

**3.12.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a + a \cos(c + dx))^{2/3}} dx = \int \frac{1}{(a + a \cos(c + dx))^{2/3}} dx$$

input `int(1/(a + a*cos(c + d*x))^(2/3), x)`output `int(1/(a + a*cos(c + d*x))^(2/3), x)`

### 3.13 $\int \frac{1}{(a+a \cos(c+dx))^{4/3}} dx$

3.13.1	Optimal result	107
3.13.2	Mathematica [A] (verified)	107
3.13.3	Rubi [A] (verified)	108
3.13.4	Maple [F]	109
3.13.5	Fricas [F]	109
3.13.6	Sympy [F]	110
3.13.7	Maxima [F]	110
3.13.8	Giac [F]	110
3.13.9	Mupad [F(-1)]	111

#### 3.13.1 Optimal result

Integrand size = 14, antiderivative size = 68

$$\int \frac{1}{(a + a \cos(c + dx))^{4/3}} dx = \frac{\text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{11}{6}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx))\right) \sin(c + dx)}{2^{5/6} a d \sqrt[6]{1 + \cos(c + dx)} \sqrt[3]{a + a \cos(c + dx)}}$$

output `1/2*hypergeom([1/2, 11/6], [3/2], 1/2-1/2*cos(d*x+c))*sin(d*x+c)*2^(1/6)/a/d / (1+cos(d*x+c))^(1/6)/(a+a*cos(d*x+c))^(1/3)`

#### 3.13.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.01

$$\int \frac{1}{(a + a \cos(c + dx))^{4/3}} dx = \frac{6 \cot\left(\frac{1}{2}(c + dx)\right) \text{Hypergeometric2F1}\left(-\frac{5}{6}, \frac{1}{2}, \frac{1}{6}, \cos^2\left(\frac{1}{2}(c + dx)\right)\right) \sqrt{\sin^2\left(\frac{1}{2}(c + dx)\right)}}{5d(a(1 + \cos(c + dx)))^{4/3}}$$

input `Integrate[(a + a*Cos[c + d*x])^(-4/3), x]`

output `(6*Cot[(c + d*x)/2]*Hypergeometric2F1[-5/6, 1/2, 1/6, Cos[(c + d*x)/2]^2]* Sqrt[Sin[(c + d*x)/2]^2])/(5*d*(a*(1 + Cos[c + d*x]))^(4/3))`

### 3.13.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \cos(c + dx) + a)^{4/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a \sin(c + dx + \frac{\pi}{2}) + a)^{4/3}} dx \\
 & \quad \downarrow \text{3131} \\
 & \frac{\sqrt[3]{\cos(c + dx) + 1} \int \frac{1}{(\cos(c + dx) + 1)^{4/3}} dx}{a \sqrt[3]{a \cos(c + dx) + a}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt[3]{\cos(c + dx) + 1} \int \frac{1}{(\sin(c + dx + \frac{\pi}{2}) + 1)^{4/3}} dx}{a \sqrt[3]{a \cos(c + dx) + a}} \\
 & \quad \downarrow \text{3130} \\
 & \frac{\sin(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{11}{6}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx))\right)}{2^{5/6} a d \sqrt[6]{\cos(c + dx) + 1} \sqrt[3]{a \cos(c + dx) + a}}
 \end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^(-4/3),x]`

output `(Hypergeometric2F1[1/2, 11/6, 3/2, (1 - Cos[c + d*x])/2]*Sin[c + d*x])/(2^(5/6)*a*d*(1 + Cos[c + d*x])^(1/6)*(a + a*Cos[c + d*x])^(1/3))`

## 3.13.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])]*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a)), x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

rule 3131 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[n]*((a + b*Sin[c + d*x])^FracPart[n]/(1 + (b/a)*Sin[c + d*x])^FracPart[n]) Int[(1 + (b/a)*Sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && !GtQ[a, 0]`

## 3.13.4 Maple [F]

$$\int \frac{1}{(a + \cos(dx + c)a)^{\frac{4}{3}}} dx$$

input `int(1/(a+cos(d*x+c)*a)^(4/3),x)`

output `int(1/(a+cos(d*x+c)*a)^(4/3),x)`

## 3.13.5 Fricas [F]

$$\int \frac{1}{(a + a \cos(c + dx))^{4/3}} dx = \int \frac{1}{(a \cos(dx + c) + a)^{\frac{4}{3}}} dx$$

input `integrate(1/(a+a*cos(d*x+c))^(4/3),x, algorithm="fricas")`

output `integral((a*cos(d*x + c) + a)^(2/3)/(a^2*cos(d*x + c)^2 + 2*a^2*cos(d*x + c) + a^2), x)`

**3.13.6 Sympy [F]**

$$\int \frac{1}{(a + a \cos(c + dx))^{4/3}} dx = \int \frac{1}{(a \cos(c + dx) + a)^{4/3}} dx$$

input `integrate(1/(a+a*cos(d*x+c))**(4/3), x)`

output `Integral((a*cos(c + d*x) + a)**(-4/3), x)`

**3.13.7 Maxima [F]**

$$\int \frac{1}{(a + a \cos(c + dx))^{4/3}} dx = \int \frac{1}{(a \cos(dx + c) + a)^{4/3}} dx$$

input `integrate(1/(a+a*cos(d*x+c))^(4/3), x, algorithm="maxima")`

output `integrate((a*cos(d*x + c) + a)^(-4/3), x)`

**3.13.8 Giac [F]**

$$\int \frac{1}{(a + a \cos(c + dx))^{4/3}} dx = \int \frac{1}{(a \cos(dx + c) + a)^{4/3}} dx$$

input `integrate(1/(a+a*cos(d*x+c))^(4/3), x, algorithm="giac")`

output `integrate((a*cos(d*x + c) + a)^(-4/3), x)`

**3.13.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a + a \cos(c + dx))^{4/3}} dx = \int \frac{1}{(a + a \cos(c + dx))^{4/3}} dx$$

input `int(1/(a + a*cos(c + d*x))^(4/3), x)`output `int(1/(a + a*cos(c + d*x))^(4/3), x)`



### 3.14 $\int (a + a \cos(c + dx))^n dx$

3.14.1	Optimal result . . . . .	112
3.14.2	Mathematica [A] (verified) . . . . .	112
3.14.3	Rubi [A] (verified) . . . . .	113
3.14.4	Maple [F] . . . . .	114
3.14.5	Fricas [F] . . . . .	114
3.14.6	Sympy [F] . . . . .	115
3.14.7	Maxima [F] . . . . .	115
3.14.8	Giac [F] . . . . .	115
3.14.9	Mupad [F(-1)] . . . . .	116

#### 3.14.1 Optimal result

Integrand size = 12, antiderivative size = 73

$$\int (a + a \cos(c + dx))^n dx = \frac{2^{\frac{1}{2}+n} (1 + \cos(c + dx))^{-\frac{1}{2}-n} (a + a \cos(c + dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx))\right) \sin(c + dx)}{d}$$

output `2^(1/2+n)*(1+cos(d*x+c))^(1/2-n)*(a+a*cos(d*x+c))^n*hypergeom([1/2, 1/2-n], [3/2], 1/2-1/2*cos(d*x+c))*sin(d*x+c)/d`

#### 3.14.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.01

$$\int (a + a \cos(c + dx))^n dx = \frac{2(a(1 + \cos(c + dx)))^n \cot\left(\frac{1}{2}(c + dx)\right) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} + n, \frac{3}{2} + n, \cos^2\left(\frac{1}{2}(c + dx)\right)\right) \sqrt{\sin^2\left(\frac{1}{2}(c + dx)\right)}}{d + 2dn}$$

input `Integrate[(a + a*Cos[c + d*x])^n,x]`

output `(-2*(a*(1 + Cos[c + d*x]))^n*Cot[(c + d*x)/2]*Hypergeometric2F1[1/2, 1/2 + n, 3/2 + n, Cos[(c + d*x)/2]^2]*Sqrt[Sin[(c + d*x)/2]^2])/(d + 2*d*n)`

### 3.14.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \cos(c + dx) + a)^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( a \sin\left(c + dx + \frac{\pi}{2}\right) + a \right)^n dx \\
 & \quad \downarrow \text{3131} \\
 & (\cos(c + dx) + 1)^{-n} (a \cos(c + dx) + a)^n \int (\cos(c + dx) + 1)^n dx \\
 & \quad \downarrow \text{3042} \\
 & (\cos(c + dx) + 1)^{-n} (a \cos(c + dx) + a)^n \int \left( \sin\left(c + dx + \frac{\pi}{2}\right) + 1 \right)^n dx \\
 & \quad \downarrow \text{3130} \\
 & \frac{2^{n+\frac{1}{2}} \sin(c + dx) (\cos(c + dx) + 1)^{-n-\frac{1}{2}} (a \cos(c + dx) + a)^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx))\right)}{d}
 \end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^n,x]`

output `(2^(1/2 + n)*(1 + Cos[c + d*x])^(-1/2 - n)*(a + a*Cos[c + d*x])^n*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1 - Cos[c + d*x])/2]*Sin[c + d*x])/d`

## 3.14.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]))]*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a)), x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

rule 3131 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[n]*((a + b*Sin[c + d*x])^FracPart[n]/(1 + (b/a)*Sin[c + d*x])^FracPart[n]) Int[(1 + (b/a)*Sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && !GtQ[a, 0]`

## 3.14.4 Maple [F]

$$\int (a + \cos(dx + c)) a^n dx$$

input `int((a+cos(d*x+c))*a^n,x)`

output `int((a+cos(d*x+c))*a^n,x)`

## 3.14.5 Fricas [F]

$$\int (a + a \cos(c + dx))^n dx = \int (a \cos(dx + c) + a)^n dx$$

input `integrate((a+a*cos(d*x+c))^n,x, algorithm="fricas")`

output `integral((a*cos(d*x + c) + a)^n, x)`

**3.14.6 Sympy [F]**

$$\int (a + a \cos(c + dx))^n dx = \int (a \cos(c + dx) + a)^n dx$$

input `integrate((a+a*cos(d*x+c))**n,x)`

output `Integral((a*cos(c + d*x) + a)**n, x)`

**3.14.7 Maxima [F]**

$$\int (a + a \cos(c + dx))^n dx = \int (a \cos(dx + c) + a)^n dx$$

input `integrate((a+a*cos(d*x+c))^n,x, algorithm="maxima")`

output `integrate((a*cos(d*x + c) + a)^n, x)`

**3.14.8 Giac [F]**

$$\int (a + a \cos(c + dx))^n dx = \int (a \cos(dx + c) + a)^n dx$$

input `integrate((a+a*cos(d*x+c))^n,x, algorithm="giac")`

output `integrate((a*cos(d*x + c) + a)^n, x)`

**3.14.9 Mupad [F(-1)]**

Timed out.

$$\int (a + a \cos(c + dx))^n dx = \int (a + a \cos(c + dx))^n dx$$

input `int((a + a*cos(c + d*x))^n,x)`output `int((a + a*cos(c + d*x))^n, x)`

### 3.15 $\int (a - a \cos(c + dx))^n dx$

3.15.1	Optimal result . . . . .	117
3.15.2	Mathematica [A] (verified) . . . . .	117
3.15.3	Rubi [A] (verified) . . . . .	118
3.15.4	Maple [F] . . . . .	119
3.15.5	Fricas [F] . . . . .	119
3.15.6	Sympy [F] . . . . .	120
3.15.7	Maxima [F] . . . . .	120
3.15.8	Giac [F] . . . . .	120
3.15.9	Mupad [F(-1)] . . . . .	121

#### 3.15.1 Optimal result

Integrand size = 13, antiderivative size = 75

$$\int (a - a \cos(c + dx))^n dx = \frac{2^{\frac{1}{2}+n} (1 - \cos(c + dx))^{-\frac{1}{2}-n} (a - a \cos(c + dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(1 + \cos(c + dx))\right)}{d}$$

output `-2^(1/2+n)*(1-cos(d*x+c))^(1/2-n)*(a-a*cos(d*x+c))^n*hypergeom([1/2, 1/2-n], [3/2], 1/2+1/2*cos(d*x+c))*sin(d*x+c)/d`

#### 3.15.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00

$$\int (a - a \cos(c + dx))^n dx = \frac{\sqrt{2} \sqrt{1 + \cos(c + dx)} (a - a \cos(c + dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} + n, \frac{3}{2} + n, \sin^2\left(\frac{1}{2}(c + dx)\right)\right) \tan\left(\frac{1}{2}(c + dx)\right)}{d + 2dn}$$

input `Integrate[(a - a*Cos[c + d*x])^n,x]`

output `(Sqrt[2]*Sqrt[1 + Cos[c + d*x]]*(a - a*Cos[c + d*x])^n*Hypergeometric2F1[1/2, 1/2 + n, 3/2 + n, Sin[(c + d*x)/2]^2]*Tan[(c + d*x)/2])/(d + 2*d*n)`

### 3.15.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a - a \cos(c + dx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( a - a \sin \left( c + dx + \frac{\pi}{2} \right) \right)^n dx \\
 & \quad \downarrow \text{3131} \\
 & (1 - \cos(c + dx))^{-n} (a - a \cos(c + dx))^n \int (1 - \cos(c + dx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & (1 - \cos(c + dx))^{-n} (a - a \cos(c + dx))^n \int \left( 1 - \sin \left( c + dx + \frac{\pi}{2} \right) \right)^n dx \\
 & \quad \downarrow \text{3130} \\
 & \frac{2^{n+\frac{1}{2}} \sin(c + dx) (1 - \cos(c + dx))^{-n-\frac{1}{2}} (a - a \cos(c + dx))^n \operatorname{Hypergeometric2F1} \left( \frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2} (\cos(c + dx) + 1) \right)}{d}
 \end{aligned}$$

input `Int[(a - a*Cos[c + d*x])^n,x]`

output `-((2^(1/2 + n)*(1 - Cos[c + d*x])^(-1/2 - n)*(a - a*Cos[c + d*x])^n*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1 + Cos[c + d*x])/2]*Sin[c + d*x])/d)`

## 3.15.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]))]*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a)), x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

rule 3131 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[n]*((a + b*Sin[c + d*x])^FracPart[n]/(1 + (b/a)*Sin[c + d*x])^FracPart[n]) Int[(1 + (b/a)*Sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && !GtQ[a, 0]`

## 3.15.4 Maple [F]

$$\int (a - \cos(dx + c) a)^n dx$$

input `int((a-cos(d*x+c)*a)^n,x)`

output `int((a-cos(d*x+c)*a)^n,x)`

## 3.15.5 Fracas [F]

$$\int (a - a \cos(c + dx))^n dx = \int (-a \cos(dx + c) + a)^n dx$$

input `integrate((a-a*cos(d*x+c))^n,x, algorithm="fricas")`

output `integral((-a*cos(d*x + c) + a)^n, x)`



**3.15.6 Sympy [F]**

$$\int (a - a \cos(c + dx))^n dx = \int (-a \cos(c + dx) + a)^n dx$$

input `integrate((a-a*cos(d*x+c))**n,x)`

output `Integral((-a*cos(c + d*x) + a)**n, x)`

**3.15.7 Maxima [F]**

$$\int (a - a \cos(c + dx))^n dx = \int (-a \cos(dx + c) + a)^n dx$$

input `integrate((a-a*cos(d*x+c))^n,x, algorithm="maxima")`

output `integrate((-a*cos(d*x + c) + a)^n, x)`

**3.15.8 Giac [F]**

$$\int (a - a \cos(c + dx))^n dx = \int (-a \cos(dx + c) + a)^n dx$$

input `integrate((a-a*cos(d*x+c))^n,x, algorithm="giac")`

output `integrate((-a*cos(d*x + c) + a)^n, x)`

**3.15.9 Mupad [F(-1)]**

Timed out.

$$\int (a - a \cos(c + dx))^n dx = \int (a - a \cos(c + dx))^n dx$$

input `int((a - a*cos(c + d*x))^n,x)`output `int((a - a*cos(c + d*x))^n, x)`

### 3.16 $\int (2 + 2 \cos(c + dx))^n dx$

3.16.1	Optimal result . . . . .	122
3.16.2	Mathematica [A] (verified) . . . . .	122
3.16.3	Rubi [A] (verified) . . . . .	123
3.16.4	Maple [F] . . . . .	124
3.16.5	Fricas [F] . . . . .	124
3.16.6	Sympy [F] . . . . .	124
3.16.7	Maxima [F] . . . . .	125
3.16.8	Giac [F] . . . . .	125
3.16.9	Mupad [F(-1)] . . . . .	125

#### 3.16.1 Optimal result

Integrand size = 12, antiderivative size = 59

$$\int (2 + 2 \cos(c + dx))^n dx = \frac{2^{\frac{1}{2}+2n} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx))\right) \sin(c + dx)}{d \sqrt{1 + \cos(c + dx)}}$$

output `2^(1/2+2*n)*hypergeom([1/2, 1/2-n], [3/2], 1/2-1/2*cos(d*x+c))*sin(d*x+c)/d/(1+cos(d*x+c))^(1/2)`

#### 3.16.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.31

$$\int (2 + 2 \cos(c + dx))^n dx = \frac{2^{1+n} (1 + \cos(c + dx))^n \cot\left(\frac{1}{2}(c + dx)\right) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} + n, \frac{3}{2} + n, \cos^2\left(\frac{1}{2}(c + dx)\right)\right) \sqrt{\sin^2\left(\frac{1}{2}(c + dx)\right)}}{d + 2dn}$$

input `Integrate[(2 + 2*Cos[c + d*x])^n,x]`

output `-((2^(1 + n)*(1 + Cos[c + d*x])^n*Cot[(c + d*x)/2]*Hypergeometric2F1[1/2, 1/2 + n, 3/2 + n, Cos[(c + d*x)/2]^2]*Sqrt[Sin[(c + d*x)/2]^2])/(d + 2*d*n))`

### 3.16.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (2 \cos(c + dx) + 2)^n dx$$

$$\downarrow \text{3042}$$

$$\int \left( 2 \sin \left( c + dx + \frac{\pi}{2} \right) + 2 \right)^n dx$$

$$\downarrow \text{3130}$$

$$\frac{2^{2n+\frac{1}{2}} \sin(c + dx) \operatorname{Hypergeometric2F1} \left( \frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx)) \right)}{d \sqrt{\cos(c + dx) + 1}}$$

input `Int[(2 + 2*Cos[c + d*x])^n,x]`

output `(2^(1/2 + 2*n)*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1 - Cos[c + d*x])/2]*Sin[c + d*x])/(d*Sqrt[1 + Cos[c + d*x]])`

#### 3.16.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :=> Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]]))*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a))], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

**3.16.4 Maple [F]**

$$\int (2 + 2 \cos(dx + c))^n dx$$

input `int((2+2*cos(d*x+c))^n,x)`

output `int((2+2*cos(d*x+c))^n,x)`

**3.16.5 Fricas [F]**

$$\int (2 + 2 \cos(c + dx))^n dx = \int (2 \cos(dx + c) + 2)^n dx$$

input `integrate((2+2*cos(d*x+c))^n,x, algorithm="fricas")`

output `integral((2*cos(d*x + c) + 2)^n, x)`

**3.16.6 Sympy [F]**

$$\int (2 + 2 \cos(c + dx))^n dx = 2^n \int (\cos(c + dx) + 1)^n dx$$

input `integrate((2+2*cos(d*x+c))**n,x)`

output `2**n*Integral((cos(c + d*x) + 1)**n, x)`

**3.16.7 Maxima [F]**

$$\int (2 + 2 \cos(c + dx))^n dx = \int (2 \cos(dx + c) + 2)^n dx$$

input `integrate((2+2*cos(d*x+c))^n,x, algorithm="maxima")`

output `integrate((2*cos(d*x + c) + 2)^n, x)`

**3.16.8 Giac [F]**

$$\int (2 + 2 \cos(c + dx))^n dx = \int (2 \cos(dx + c) + 2)^n dx$$

input `integrate((2+2*cos(d*x+c))^n,x, algorithm="giac")`

output `integrate((2*cos(d*x + c) + 2)^n, x)`

**3.16.9 Mupad [F(-1)]**

Timed out.

$$\int (2 + 2 \cos(c + dx))^n dx = \int \left( 4 \cos \left( \frac{c}{2} + \frac{dx}{2} \right)^2 \right)^n dx$$

input `int((2*cos(c + d*x) + 2)^n,x)`

output `int((4*cos(c/2 + (d*x)/2)^2)^n, x)`

### 3.17 $\int (2 - 2 \cos(c + dx))^n dx$

3.17.1	Optimal result . . . . .	126
3.17.2	Mathematica [A] (verified) . . . . .	126
3.17.3	Rubi [A] (verified) . . . . .	127
3.17.4	Maple [F] . . . . .	128
3.17.5	Fricas [F] . . . . .	128
3.17.6	Sympy [F] . . . . .	128
3.17.7	Maxima [F] . . . . .	129
3.17.8	Giac [F] . . . . .	129
3.17.9	Mupad [F(-1)] . . . . .	129

#### 3.17.1 Optimal result

Integrand size = 12, antiderivative size = 60

$$\int (2 - 2 \cos(c + dx))^n dx = -\frac{2^{\frac{1}{2}+2n} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(1 + \cos(c + dx))\right) \sin(c + dx)}{d\sqrt{1 - \cos(c + dx)}}$$

output `-2^(1/2+2*n)*hypergeom([1/2, 1/2-n], [3/2], 1/2+1/2*cos(d*x+c))*sin(d*x+c)/d/(1-cos(d*x+c))^(1/2)`

#### 3.17.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.23

$$\int (2 - 2 \cos(c + dx))^n dx = \frac{\sqrt{2}(2 - 2 \cos(c + dx))^n \sqrt{1 + \cos(c + dx)} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} + n, \frac{3}{2} + n, \sin^2\left(\frac{1}{2}(c + dx)\right)\right) \tan\left(\frac{1}{2}(c + dx)\right)}{d + 2dn}$$

input `Integrate[(2 - 2*Cos[c + d*x])^n,x]`

output `(Sqrt[2]*(2 - 2*Cos[c + d*x])^n*Sqrt[1 + Cos[c + d*x]]*Hypergeometric2F1[1/2, 1/2 + n, 3/2 + n, Sin[(c + d*x)/2]^2]*Tan[(c + d*x)/2])/(d + 2*d*n)`

### 3.17.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (2 - 2 \cos(c + dx))^n dx$$

↓ 3042

$$\int \left(2 - 2 \sin\left(c + dx + \frac{\pi}{2}\right)\right)^n dx$$

↓ 3130

$$\frac{2^{2n+\frac{1}{2}} \sin(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(\cos(c + dx) + 1)\right)}{d\sqrt{1 - \cos(c + dx)}}$$

input `Int[(2 - 2*Cos[c + d*x])^n,x]`

output `-((2^(1/2 + 2*n)*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1 + Cos[c + d*x])/2]*Sin[c + d*x])/(d*Sqrt[1 - Cos[c + d*x]]))`

#### 3.17.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :=> Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]]))*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a))], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`



**3.17.4 Maple [F]**

$$\int (-2 \cos(dx + c) + 2)^n dx$$

input `int((-2*cos(d*x+c)+2)^n,x)`

output `int((-2*cos(d*x+c)+2)^n,x)`

**3.17.5 Fricas [F]**

$$\int (2 - 2 \cos(c + dx))^n dx = \int (-2 \cos(dx + c) + 2)^n dx$$

input `integrate((2-2*cos(d*x+c))^n,x, algorithm="fricas")`

output `integral((-2*cos(d*x + c) + 2)^n, x)`

**3.17.6 Sympy [F]**

$$\int (2 - 2 \cos(c + dx))^n dx = \int (2 - 2 \cos(c + dx))^n dx$$

input `integrate((2-2*cos(d*x+c))**n,x)`

output `Integral((2 - 2*cos(c + d*x))**n, x)`

**3.17.7 Maxima [F]**

$$\int (2 - 2 \cos(c + dx))^n dx = \int (-2 \cos(dx + c) + 2)^n dx$$

input `integrate((2-2*cos(d*x+c))^n,x, algorithm="maxima")`

output `integrate((-2*cos(d*x + c) + 2)^n, x)`

**3.17.8 Giac [F]**

$$\int (2 - 2 \cos(c + dx))^n dx = \int (-2 \cos(dx + c) + 2)^n dx$$

input `integrate((2-2*cos(d*x+c))^n,x, algorithm="giac")`

output `integrate((-2*cos(d*x + c) + 2)^n, x)`

**3.17.9 Mupad [F(-1)]**

Timed out.

$$\int (2 - 2 \cos(c + dx))^n dx = \int \left( 4 \sin \left( \frac{c}{2} + \frac{dx}{2} \right)^2 \right)^n dx$$

input `int((2 - 2*cos(c + d*x))^n,x)`

output `int((4*sin(c/2 + (d*x)/2)^2)^n, x)`

### 3.18 $\int \frac{1}{5+3 \cos(c+dx)} dx$

3.18.1 Optimal result . . . . .	130
3.18.2 Mathematica [A] (verified) . . . . .	130
3.18.3 Rubi [A] (verified) . . . . .	131
3.18.4 Maple [A] (verified) . . . . .	132
3.18.5 Fricas [A] (verification not implemented) . . . . .	132
3.18.6 Sympy [A] (verification not implemented) . . . . .	132
3.18.7 Maxima [A] (verification not implemented) . . . . .	133
3.18.8 Giac [A] (verification not implemented) . . . . .	133
3.18.9 Mupad [B] (verification not implemented) . . . . .	134

#### 3.18.1 Optimal result

Integrand size = 12, antiderivative size = 31

$$\int \frac{1}{5+3 \cos(c+dx)} dx = \frac{x}{4} - \frac{\arctan\left(\frac{\sin(c+dx)}{3+\cos(c+dx)}\right)}{2d}$$

output `1/4*x-1/2*arctan(sin(d*x+c)/(3+cos(d*x+c)))/d`

#### 3.18.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.65

$$\int \frac{1}{5+3 \cos(c+dx)} dx = -\frac{\arctan\left(2 \cot\left(\frac{1}{2}(c+dx)\right)\right)}{2d}$$

input `Integrate[(5 + 3*Cos[c + d*x])^(-1), x]`

output `-1/2*ArcTan[2*Cot[(c + d*x)/2]]/d`

### 3.18.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{3 \cos(c + dx) + 5} dx$$

↓ 3042

$$\int \frac{1}{3 \sin(c + dx + \frac{\pi}{2}) + 5} dx$$

↓ 3136

$$\frac{x}{4} - \frac{\arctan\left(\frac{\sin(c+dx)}{\cos(c+dx)+3}\right)}{2d}$$

input `Int[(5 + 3*Cos[c + d*x])^(-1),x]`

output `x/4 - ArcTan[Sin[c + d*x]/(3 + Cos[c + d*x])]/(2*d)`

#### 3.18.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3136 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

### 3.18.4 Maple [A] (verified)

Time = 1.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.58

method	result	size
derivativedivides	$\frac{\arctan\left(\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2}\right)}{2d}$	18
default	$\frac{\arctan\left(\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2}\right)}{2d}$	18
parallelrisc	$-\frac{i\left(\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2i\right) - \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2i\right)\right)}{4d}$	36
risch	$-\frac{i\ln\left(e^{i(dx+c)} + \frac{1}{3}\right)}{4d} + \frac{i\ln\left(e^{i(dx+c)} + 3\right)}{4d}$	38

input `int(1/(5+3*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/2/d*arctan(1/2*tan(1/2*d*x+1/2*c))`

### 3.18.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{1}{5 + 3 \cos(c + dx)} dx = -\frac{\arctan\left(\frac{5 \cos(dx+c)+3}{4 \sin(dx+c)}\right)}{4d}$$

input `integrate(1/(5+3*cos(d*x+c)),x, algorithm="fricas")`

output `-1/4*arctan(1/4*(5*cos(d*x + c) + 3)/sin(d*x + c))/d`

### 3.18.6 Sympy [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.32

$$\int \frac{1}{5 + 3 \cos(c + dx)} dx = \begin{cases} \frac{\operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right) + \pi \left\lfloor \frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2} \right\rfloor}{2d} & \text{for } d \neq 0 \\ \frac{x}{3 \cos(c)+5} & \text{otherwise} \end{cases}$$

input `integrate(1/(5+3*cos(d*x+c)),x)`

output `Piecewise(((atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/  
(2*d), Ne(d, 0)), (x/(3*cos(c) + 5), True))`

### 3.18.7 Maxima [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.77

$$\int \frac{1}{5 + 3 \cos(c + dx)} dx = \frac{\arctan\left(\frac{\sin(dx+c)}{2(\cos(dx+c)+1)}\right)}{2d}$$

input `integrate(1/(5+3*cos(d*x+c)),x, algorithm="maxima")`

output `1/2*arctan(1/2*sin(d*x + c)/(cos(d*x + c) + 1))/d`

### 3.18.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.97

$$\int \frac{1}{5 + 3 \cos(c + dx)} dx = \frac{dx + c - 2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+3}\right)}{4d}$$

input `integrate(1/(5+3*cos(d*x+c)),x, algorithm="giac")`

output `1/4*(d*x + c - 2*arctan(sin(d*x + c)/(cos(d*x + c) + 3)))/d`

**3.18.9 Mupad [B] (verification not implemented)**

Time = 14.62 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.23

$$\int \frac{1}{5 + 3 \cos(c + dx)} dx = \frac{\operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{2d} - \frac{\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}}{2d}$$

input `int(1/(3*cos(c + d*x) + 5),x)`

output `atan(tan(c/2 + (d*x)/2)/2)/(2*d) - (atan(tan(c/2 + (d*x)/2)) - (d*x)/2)/(2*d)`

### 3.19 $\int \frac{1}{(5+3 \cos(c+dx))^2} dx$

3.19.1	Optimal result . . . . .	135
3.19.2	Mathematica [A] (verified) . . . . .	135
3.19.3	Rubi [A] (verified) . . . . .	136
3.19.4	Maple [A] (verified) . . . . .	137
3.19.5	Fricas [A] (verification not implemented) . . . . .	138
3.19.6	Sympy [C] (verification not implemented) . . . . .	138
3.19.7	Maxima [A] (verification not implemented) . . . . .	139
3.19.8	Giac [A] (verification not implemented) . . . . .	139
3.19.9	Mupad [B] (verification not implemented) . . . . .	139

#### 3.19.1 Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \frac{1}{(5 + 3 \cos(c + dx))^2} dx = \frac{5x}{64} - \frac{5 \arctan\left(\frac{\sin(c+dx)}{3+\cos(c+dx)}\right)}{32d} - \frac{3 \sin(c + dx)}{16d(5 + 3 \cos(c + dx))}$$

output `5/64*x-5/32*arctan(sin(d*x+c)/(3+cos(d*x+c)))/d-3/16*sin(d*x+c)/d/(5+3*cos(d*x+c))`

#### 3.19.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.77

$$\int \frac{1}{(5 + 3 \cos(c + dx))^2} dx = -\frac{5 \arctan\left(2 \cot\left(\frac{1}{2}(c + dx)\right)\right) + \frac{6 \sin(c+dx)}{5+3 \cos(c+dx)}}{32d}$$

input `Integrate[(5 + 3*Cos[c + d*x])^(-2),x]`

output `-1/32*(5*ArcTan[2*Cot[(c + d*x)/2]] + (6*Sin[c + d*x])/(5 + 3*Cos[c + d*x]))/d`



**3.19.3 Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {3042, 3143, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(3 \cos(c + dx) + 5)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(3 \sin(c + dx + \frac{\pi}{2}) + 5)^2} dx \\
 & \quad \downarrow \text{3143} \\
 & -\frac{1}{16} \int -\frac{5}{3 \cos(c + dx) + 5} dx - \frac{3 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)} \\
 & \quad \downarrow \text{27} \\
 & \frac{5}{16} \int \frac{1}{3 \cos(c + dx) + 5} dx - \frac{3 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{16} \int \frac{1}{3 \sin(c + dx + \frac{\pi}{2}) + 5} dx - \frac{3 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)} \\
 & \quad \downarrow \text{3136} \\
 & \frac{5}{16} \left( \frac{x}{4} - \frac{\arctan\left(\frac{\sin(c+dx)}{\cos(c+dx)+3}\right)}{2d} \right) - \frac{3 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)}
 \end{aligned}$$

input `Int[(5 + 3*Cos[c + d*x])^(-2), x]`

output `(5*(x/4 - ArcTan[Sin[c + d*x]/(3 + Cos[c + d*x])]/(2*d)))/16 - (3*Sin[c + d*x])/(16*d*(5 + 3*Cos[c + d*x]))`

3.19.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

3.19.4 Maple [A] (verified)

Time = 0.68 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$\frac{-\frac{3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{16\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\right)} + \frac{5 \arctan\left(\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2}\right)}{32}}{d}$	46
default	$\frac{-\frac{3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{16\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\right)} + \frac{5 \arctan\left(\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2}\right)}{32}}{d}$	46
parallelrisch	$\frac{(-15i \cos(dx+c) - 25i) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2i\right) + (15i \cos(dx+c) + 25i) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2i\right) - 12 \sin(dx+c)}{192d \cos(dx+c) + 320d}$	78
risch	$-\frac{i(5e^{i(dx+c)} + 3)}{8d(3e^{2i(dx+c)} + 10e^{i(dx+c)} + 3)} - \frac{5i \ln\left(e^{i(dx+c)} + \frac{1}{3}\right)}{64d} + \frac{5i \ln\left(e^{i(dx+c)} + 3\right)}{64d}$	83

input `int(1/(5+3*cos(d*x+c))^2,x,method=_RETURNVERBOSE)`

output  $1/d*(-3/16*\tan(1/2*d*x+1/2*c)/(\tan(1/2*d*x+1/2*c)^2+4)+5/32*\arctan(1/2*\tan(1/2*d*x+1/2*c)))$

### 3.19.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.05

$$\int \frac{1}{(5+3\cos(c+dx))^2} dx = -\frac{5(3\cos(dx+c)+5)\arctan\left(\frac{5\cos(dx+c)+3}{4\sin(dx+c)}\right)+12\sin(dx+c)}{64(3d\cos(dx+c)+5d)}$$

input `integrate(1/(5+3*cos(d*x+c))^2,x, algorithm="fricas")`

output  $-1/64*(5*(3*\cos(d*x+c)+5)*\arctan(1/4*(5*\cos(d*x+c)+3)/\sin(d*x+c))+12*\sin(d*x+c))/(3*d*\cos(d*x+c)+5*d)$

### 3.19.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.91 (sec) , antiderivative size = 187, normalized size of antiderivative = 3.34

$$\int \frac{1}{(5+3\cos(c+dx))^2} dx = \begin{cases} \frac{x}{(5+3\cosh(2\operatorname{atanh}(2)))^2} \\ \frac{x}{(3\cos(c)+5)^2} \\ 5\left(\frac{\operatorname{atan}\left(\frac{\tan\left(\frac{c}{2}+\frac{dx}{2}\right)}{2}\right)+\pi\left\lfloor\frac{\frac{c}{2}+\frac{dx}{2}-\frac{\pi}{2}}{\pi}\right\rfloor\right)\tan^2\left(\frac{c}{2}+\frac{dx}{2}\right)}{32d\tan^2\left(\frac{c}{2}+\frac{dx}{2}\right)+128d} + \frac{20\left(\operatorname{atan}\left(\frac{\tan\left(\frac{c}{2}+\frac{dx}{2}\right)}{2}\right)+\pi\left\lfloor\frac{\frac{c}{2}+\frac{dx}{2}-\frac{\pi}{2}}{\pi}\right\rfloor\right)}{32d\tan^2\left(\frac{c}{2}+\frac{dx}{2}\right)+128d} - \frac{6\tan\left(\frac{c}{2}+\frac{dx}{2}\right)}{32d\tan^2\left(\frac{c}{2}+\frac{dx}{2}\right)+128d} \end{cases}$$

input `integrate(1/(5+3*cos(d*x+c))**2,x)`

output `Piecewise((x/(5+3*cosh(2*atanh(2)))**2, Eq(c, -d*x-2*I*atanh(2)) | Eq(c, -d*x+2*I*atanh(2))), (x/(3*cos(c)+5)**2, Eq(d, 0)), (5*(atan(tan(c/2+d*x/2)/2)+pi*floor((c/2+d*x/2-pi/2)/pi))*tan(c/2+d*x/2)**2/(32*d*tan(c/2+d*x/2)**2+128*d)+20*(atan(tan(c/2+d*x/2)/2)+pi*floor((c/2+d*x/2-pi/2)/pi))/(32*d*tan(c/2+d*x/2)**2+128*d)-6*tan(c/2+d*x/2)/(32*d*tan(c/2+d*x/2)**2+128*d), True))`

---

3.19.  $\int \frac{1}{(5+3\cos(c+dx))^2} dx$

**3.19.7 Maxima [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.21

$$\int \frac{1}{(5 + 3 \cos(c + dx))^2} dx = -\frac{\frac{6 \sin(dx+c)}{\left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 4\right)(\cos(dx+c)+1)}{32 d} - 5 \arctan\left(\frac{\sin(dx+c)}{2(\cos(dx+c)+1)}\right)$$

input `integrate(1/(5+3*cos(d*x+c))^2,x, algorithm="maxima")`output `-1/32*(6*sin(d*x + c)/((sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 4)*(cos(d*x + c) + 1)) - 5*arctan(1/2*sin(d*x + c)/(cos(d*x + c) + 1)))/d`**3.19.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.05

$$\int \frac{1}{(5 + 3 \cos(c + dx))^2} dx = \frac{5 dx + 5 c - \frac{12 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 4} - 10 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+3}\right)}{64 d}$$

input `integrate(1/(5+3*cos(d*x+c))^2,x, algorithm="giac")`output `1/64*(5*d*x + 5*c - 12*tan(1/2*d*x + 1/2*c)/(tan(1/2*d*x + 1/2*c)^2 + 4) - 10*arctan(sin(d*x + c)/(cos(d*x + c) + 3)))/d`**3.19.9 Mupad [B] (verification not implemented)**

Time = 15.52 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.20

$$\int \frac{1}{(5 + 3 \cos(c + dx))^2} dx = \frac{5 \operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{32 d} - \frac{5 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{32 d} - \frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{16 d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 4\right)}$$

input `int(1/(3*cos(c + d*x) + 5)^2,x)`

output `(5*atan(tan(c/2 + (d*x)/2)/2))/(32*d) - (5*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2))/(32*d) - (3*tan(c/2 + (d*x)/2))/(16*d*(tan(c/2 + (d*x)/2)^2 + 4))`

### 3.20 $\int \frac{1}{(5+3 \cos(c+dx))^3} dx$

3.20.1	Optimal result . . . . .	141
3.20.2	Mathematica [A] (verified) . . . . .	141
3.20.3	Rubi [A] (verified) . . . . .	142
3.20.4	Maple [A] (verified) . . . . .	144
3.20.5	Fricas [A] (verification not implemented) . . . . .	144
3.20.6	Sympy [C] (verification not implemented) . . . . .	145
3.20.7	Maxima [A] (verification not implemented) . . . . .	145
3.20.8	Giac [A] (verification not implemented) . . . . .	146
3.20.9	Mupad [B] (verification not implemented) . . . . .	146

#### 3.20.1 Optimal result

Integrand size = 12, antiderivative size = 81

$$\int \frac{1}{(5 + 3 \cos(c + dx))^3} dx = \frac{59x}{2048} - \frac{59 \arctan\left(\frac{\sin(c+dx)}{3+\cos(c+dx)}\right)}{1024d} - \frac{3 \sin(c + dx)}{32d(5 + 3 \cos(c + dx))^2} - \frac{45 \sin(c + dx)}{512d(5 + 3 \cos(c + dx))}$$

output `59/2048*x-59/1024*arctan(sin(d*x+c)/(3+cos(d*x+c)))/d-3/32*sin(d*x+c)/d/(5+3*cos(d*x+c))^2-45/512*sin(d*x+c)/d/(5+3*cos(d*x+c))`

#### 3.20.2 Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.69

$$\int \frac{1}{(5 + 3 \cos(c + dx))^3} dx = -\frac{59 \arctan\left(2 \cot\left(\frac{1}{2}(c + dx)\right)\right) + \frac{3(182 \sin(c+dx)+45 \sin(2(c+dx)))}{(5+3 \cos(c+dx))^2}}{1024d}$$

input `Integrate[(5 + 3*Cos[c + d*x])^(-3), x]`

output `-1/1024*(59*ArcTan[2*Cot[(c + d*x)/2]] + (3*(182*Sin[c + d*x] + 45*Sin[2*(c + d*x)])))/(5 + 3*Cos[c + d*x])^2/d`

**3.20.3 Rubi [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.12, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {3042, 3143, 25, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(3 \cos(c + dx) + 5)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(3 \sin(c + dx + \frac{\pi}{2}) + 5)^3} dx \\
 & \quad \downarrow \text{3143} \\
 & -\frac{1}{32} \int -\frac{10 - 3 \cos(c + dx)}{(3 \cos(c + dx) + 5)^2} dx - \frac{3 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{32} \int \frac{10 - 3 \cos(c + dx)}{(3 \cos(c + dx) + 5)^2} dx - \frac{3 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{32} \int \frac{10 - 3 \sin(c + dx + \frac{\pi}{2})}{(3 \sin(c + dx + \frac{\pi}{2}) + 5)^2} dx - \frac{3 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} \\
 & \quad \downarrow \text{3233} \\
 & \frac{1}{32} \left( -\frac{1}{16} \int -\frac{59}{3 \cos(c + dx) + 5} dx - \frac{45 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)} \right) - \frac{3 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{32} \left( \frac{59}{16} \int \frac{1}{3 \cos(c + dx) + 5} dx - \frac{45 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)} \right) - \frac{3 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{32} \left( \frac{59}{16} \int \frac{1}{3 \sin(c + dx + \frac{\pi}{2}) + 5} dx - \frac{45 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)} \right) - \frac{3 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} \\
 & \quad \downarrow \text{3136} \\
 & \frac{1}{32} \left( \frac{59}{16} \left( \frac{x}{4} - \frac{\arctan\left(\frac{\sin(c+dx)}{\cos(c+dx)+3}\right)}{2d} \right) - \frac{45 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)} \right) - \frac{3 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2}
 \end{aligned}$$

input `Int[(5 + 3*Cos[c + d*x])^(-3),x]`

output `(-3*Sin[c + d*x])/(32*d*(5 + 3*Cos[c + d*x])^2) + ((59*(x/4 - ArcTan[Sin[c + d*x]/(3 + Cos[c + d*x]])/(2*d)))/16 - (45*Sin[c + d*x])/(16*d*(5 + 3*Cos[c + d*x]))) / 32`

### 3.20.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3233 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`



### 3.20.4 Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.77

method	result
derivativedivides	$\frac{-\frac{69(\tan^3(\frac{dx}{2} + \frac{c}{2}))}{128} - \frac{51 \tan(\frac{dx}{2} + \frac{c}{2})}{32} + \frac{59 \arctan\left(\frac{\tan(\frac{dx}{2} + \frac{c}{2})}{2}\right)}{1024}}{4(\tan^2(\frac{dx}{2} + \frac{c}{2}) + 4)^2} + \frac{59 \arctan\left(\frac{\tan(\frac{dx}{2} + \frac{c}{2})}{2}\right)}{1024}$
default	$\frac{-\frac{69(\tan^3(\frac{dx}{2} + \frac{c}{2}))}{128} - \frac{51 \tan(\frac{dx}{2} + \frac{c}{2})}{32} + \frac{59 \arctan\left(\frac{\tan(\frac{dx}{2} + \frac{c}{2})}{2}\right)}{1024}}{4(\tan^2(\frac{dx}{2} + \frac{c}{2}) + 4)^2} + \frac{59 \arctan\left(\frac{\tan(\frac{dx}{2} + \frac{c}{2})}{2}\right)}{1024}$
risch	$-\frac{3i(59e^{3i(dx+c)} + 295e^{2i(dx+c)} + 241e^{i(dx+c)} + 45)}{256d(3e^{2i(dx+c)} + 10e^{i(dx+c)} + 3)^2} - \frac{59i \ln(e^{i(dx+c)} + \frac{1}{3})}{2048d} + \frac{59i \ln(e^{i(dx+c)} + 3)}{2048d}$
parallelrisch	$\frac{59i(-59 - 9 \cos(2dx + 2c) - 60 \cos(dx + c)) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2i\right) + 59i(59 + 9 \cos(2dx + 2c) + 60 \cos(dx + c)) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2i\right)}{2048d(59 + 9 \cos(2dx + 2c) + 60 \cos(dx + c))}$

input `int(1/(5+3*cos(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/d*(1/4*(-69/128*tan(1/2*d*x+1/2*c)^3-51/32*tan(1/2*d*x+1/2*c))/(tan(1/2*d*x+1/2*c)^2+4)^2+59/1024*arctan(1/2*tan(1/2*d*x+1/2*c)))`

### 3.20.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.11

$$\int \frac{1}{(5 + 3 \cos(c + dx))^3} dx =$$

$$-\frac{59(9 \cos(dx + c)^2 + 30 \cos(dx + c) + 25) \arctan\left(\frac{5 \cos(dx + c) + 3}{4 \sin(dx + c)}\right) + 12(45 \cos(dx + c) + 91) \sin(dx + c)}{2048(9d \cos(dx + c)^2 + 30d \cos(dx + c) + 25d)}$$

input `integrate(1/(5+3*cos(d*x+c))^3,x, algorithm="fricas")`

output `-1/2048*(59*(9*cos(d*x + c)^2 + 30*cos(d*x + c) + 25)*arctan(1/4*(5*cos(d*x + c) + 3)/sin(d*x + c)) + 12*(45*cos(d*x + c) + 91)*sin(d*x + c))/(9*d*cos(d*x + c)^2 + 30*d*cos(d*x + c) + 25*d)`

### 3.20.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.59 (sec) , antiderivative size = 359, normalized size of antiderivative = 4.43

$$\int \frac{1}{(5 + 3 \cos(c + dx))^3} dx$$

$$= \begin{cases} \frac{x}{(5+3 \cosh(2 \operatorname{atanh}(2)))^3} \\ \frac{x}{(3 \cos(c)+5)^3} \\ \frac{59 \left( \operatorname{atan} \left( \frac{\tan \left( \frac{c}{2} + \frac{dx}{2} \right)}{2} \right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right) \tan^4 \left( \frac{c}{2} + \frac{dx}{2} \right)}{1024d \tan^4 \left( \frac{c}{2} + \frac{dx}{2} \right) + 8192d \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right) + 16384d} + \frac{472 \left( \operatorname{atan} \left( \frac{\tan \left( \frac{c}{2} + \frac{dx}{2} \right)}{2} \right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right) \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right)}{1024d \tan^4 \left( \frac{c}{2} + \frac{dx}{2} \right) + 8192d \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right) + 16384d} + \frac{944 \left( \operatorname{atan} \left( \frac{\tan \left( \frac{c}{2} + \frac{dx}{2} \right)}{2} \right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{1024d \tan^4 \left( \frac{c}{2} + \frac{dx}{2} \right) + 8192d \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right) + 16384d} \end{cases}$$

input `integrate(1/(5+3*cos(d*x+c))**3,x)`

output `Piecewise((x/(5 + 3*cosh(2*atanh(2)))**3, Eq(c, -d*x - 2*I*atanh(2)) | Eq(c, -d*x + 2*I*atanh(2))), (x/(3*cos(c) + 5)**3, Eq(d, 0)), (59*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4/(1024*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 16384*d) + 472*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**2/(1024*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 16384*d) + 944*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(1024*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 16384*d) - 138*tan(c/2 + d*x/2)**3/(1024*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 16384*d) - 408*tan(c/2 + d*x/2)/(1024*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 16384*d), True))`

### 3.20.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.37

$$\int \frac{1}{(5 + 3 \cos(c + dx))^3} dx = -\frac{6 \left( \frac{68 \sin(dx+c)}{\cos(dx+c)+1} + \frac{23 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{\frac{8 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{\sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 16} - 59 \arctan \left( \frac{\sin(dx+c)}{2(\cos(dx+c)+1)} \right)$$

input `integrate(1/(5+3*cos(d*x+c))^3,x, algorithm="maxima")`

output 
$$\frac{-1/1024*(6*(68*\sin(d*x + c)/(\cos(d*x + c) + 1) + 23*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3)/(8*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2 + \sin(d*x + c)^4/(\cos(d*x + c) + 1)^4 + 16) - 59*\arctan(1/2*\sin(d*x + c)/(\cos(d*x + c) + 1))}{d}$$

### 3.20.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.93

$$\int \frac{1}{(5 + 3 \cos(c + dx))^3} dx = \frac{59 dx + 59 c - \frac{12 \left( 23 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 68 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\left( \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 4 \right)^2} - 118 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+3}\right)}{2048 d}$$

input `integrate(1/(5+3*cos(d*x+c))^3,x, algorithm="giac")`

output 
$$\frac{1/2048*(59*d*x + 59*c - 12*(23*\tan(1/2*d*x + 1/2*c)^3 + 68*\tan(1/2*d*x + 1/2*c)))/(\tan(1/2*d*x + 1/2*c)^2 + 4)^2 - 118*\arctan(\sin(d*x + c)/(\cos(d*x + c) + 3))}{d}$$

### 3.20.9 Mupad [B] (verification not implemented)

Time = 15.51 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.19

$$\int \frac{1}{(5 + 3 \cos(c + dx))^3} dx = \frac{59 \operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{1024 d} - \frac{59 \left( \operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2} \right)}{1024 d} - \frac{\frac{69 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{512} + \frac{51 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{128}}{d \left( \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 8 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 16 \right)}$$

input `int(1/(3*cos(c + d*x) + 5)^3,x)`

output 
$$\frac{(59*\operatorname{atan}(\tan(c/2 + (d*x)/2)/2))/(1024*d) - (59*(\operatorname{atan}(\tan(c/2 + (d*x)/2)) - (d*x)/2))/(1024*d) - ((51*\tan(c/2 + (d*x)/2))/128 + (69*\tan(c/2 + (d*x)/2)^3)/512)/(d*(8*\tan(c/2 + (d*x)/2)^2 + \tan(c/2 + (d*x)/2)^4 + 16))}{d}$$

### 3.21 $\int \frac{1}{(5+3 \cos(c+dx))^4} dx$

3.21.1	Optimal result	147
3.21.2	Mathematica [A] (verified)	147
3.21.3	Rubi [A] (verified)	148
3.21.4	Maple [A] (verified)	150
3.21.5	Fricas [A] (verification not implemented)	151
3.21.6	Sympy [C] (verification not implemented)	151
3.21.7	Maxima [A] (verification not implemented)	152
3.21.8	Giac [A] (verification not implemented)	153
3.21.9	Mupad [B] (verification not implemented)	153

#### 3.21.1 Optimal result

Integrand size = 12, antiderivative size = 106

$$\int \frac{1}{(5+3 \cos(c+dx))^4} dx = \frac{385x}{32768} - \frac{385 \arctan\left(\frac{\sin(c+dx)}{3+\cos(c+dx)}\right)}{16384d} - \frac{\sin(c+dx)}{16d(5+3 \cos(c+dx))^3} - \frac{25 \sin(c+dx)}{512d(5+3 \cos(c+dx))^2} - \frac{311 \sin(c+dx)}{8192d(5+3 \cos(c+dx))}$$

output `385/32768*x-385/16384*arctan(sin(d*x+c)/(3+cos(d*x+c)))/d-1/16*sin(d*x+c)/d/(5+3*cos(d*x+c))^3-25/512*sin(d*x+c)/d/(5+3*cos(d*x+c))^2-311/8192*sin(d*x+c)/d/(5+3*cos(d*x+c))`

#### 3.21.2 Mathematica [A] (verified)

Time = 0.34 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.62

$$\int \frac{1}{(5+3 \cos(c+dx))^4} dx = -\frac{770 \arctan\left(2 \cot\left(\frac{1}{2}(c+dx)\right)\right) + \frac{9(4883 \sin(c+dx)+2340 \sin(2(c+dx))+311 \sin(3(c+dx)))}{(5+3 \cos(c+dx))^3}}{32768d}$$

input `Integrate[(5 + 3*Cos[c + d*x])^(-4), x]`

output `-1/32768*(770*ArcTan[2*Cot[(c + d*x)/2]] + (9*(4883*Sin[c + d*x] + 2340*Sin[2*(c + d*x)] + 311*Sin[3*(c + d*x)]))/(5 + 3*Cos[c + d*x])^3)/d`

**3.21.3 Rubi [A] (verified)**

Time = 0.51 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.14, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$ , Rules used = {3042, 3143, 27, 3042, 3233, 25, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(3 \cos(c + dx) + 5)^4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(3 \sin(c + dx + \frac{\pi}{2}) + 5)^4} dx \\
 & \quad \downarrow \text{3143} \\
 & -\frac{1}{48} \int -\frac{3(5 - 2 \cos(c + dx))}{(3 \cos(c + dx) + 5)^3} dx - \frac{\sin(c + dx)}{16d(3 \cos(c + dx) + 5)^3} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{16} \int \frac{5 - 2 \cos(c + dx)}{(3 \cos(c + dx) + 5)^3} dx - \frac{\sin(c + dx)}{16d(3 \cos(c + dx) + 5)^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{16} \int \frac{5 - 2 \sin(c + dx + \frac{\pi}{2})}{(3 \sin(c + dx + \frac{\pi}{2}) + 5)^3} dx - \frac{\sin(c + dx)}{16d(3 \cos(c + dx) + 5)^3} \\
 & \quad \downarrow \text{3233} \\
 & \frac{1}{16} \left( -\frac{1}{32} \int -\frac{62 - 25 \cos(c + dx)}{(3 \cos(c + dx) + 5)^2} dx - \frac{25 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} \right) - \frac{\sin(c + dx)}{16d(3 \cos(c + dx) + 5)^3} \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{16} \left( \frac{1}{32} \int \frac{62 - 25 \cos(c + dx)}{(3 \cos(c + dx) + 5)^2} dx - \frac{25 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} \right) - \frac{\sin(c + dx)}{16d(3 \cos(c + dx) + 5)^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{16} \left( \frac{1}{32} \int \frac{62 - 25 \sin(c + dx + \frac{\pi}{2})}{(3 \sin(c + dx + \frac{\pi}{2}) + 5)^2} dx - \frac{25 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} \right) - \frac{\sin(c + dx)}{16d(3 \cos(c + dx) + 5)^3} \\
 & \quad \downarrow \text{3233}
 \end{aligned}$$

$$\frac{1}{16} \left( \frac{1}{32} \left( -\frac{1}{16} \int -\frac{385}{3 \cos(c+dx)+5} dx - \frac{311 \sin(c+dx)}{16d(3 \cos(c+dx)+5)} \right) - \frac{25 \sin(c+dx)}{32d(3 \cos(c+dx)+5)^2} \right) - \frac{\sin(c+dx)}{16d(3 \cos(c+dx)+5)^3}$$

↓ 27

$$\frac{1}{16} \left( \frac{1}{32} \left( \frac{385}{16} \int \frac{1}{3 \cos(c+dx)+5} dx - \frac{311 \sin(c+dx)}{16d(3 \cos(c+dx)+5)} \right) - \frac{25 \sin(c+dx)}{32d(3 \cos(c+dx)+5)^2} \right) - \frac{\sin(c+dx)}{16d(3 \cos(c+dx)+5)^3}$$

↓ 3042

$$\frac{1}{16} \left( \frac{1}{32} \left( \frac{385}{16} \int \frac{1}{3 \sin(c+dx+\frac{\pi}{2})+5} dx - \frac{311 \sin(c+dx)}{16d(3 \cos(c+dx)+5)} \right) - \frac{25 \sin(c+dx)}{32d(3 \cos(c+dx)+5)^2} \right) - \frac{\sin(c+dx)}{16d(3 \cos(c+dx)+5)^3}$$

↓ 3136

$$\frac{1}{16} \left( \frac{1}{32} \left( \frac{385}{16} \left( \frac{x}{4} - \frac{\arctan\left(\frac{\sin(c+dx)}{\cos(c+dx)+3}\right)}{2d} \right) - \frac{311 \sin(c+dx)}{16d(3 \cos(c+dx)+5)} \right) - \frac{25 \sin(c+dx)}{32d(3 \cos(c+dx)+5)^2} \right) - \frac{\sin(c+dx)}{16d(3 \cos(c+dx)+5)^3}$$

input `Int[(5 + 3*Cos[c + d*x])^(-4),x]`

output `-1/16*Sin[c + d*x]/(d*(5 + 3*Cos[c + d*x])^3) + ((-25*Sin[c + d*x])/(32*d*(5 + 3*Cos[c + d*x])^2) + ((385*(x/4 - ArcTan[Sin[c + d*x]/(3 + Cos[c + d*x])])/(2*d)))/16 - (311*Sin[c + d*x])/(16*d*(5 + 3*Cos[c + d*x]))/32)/16`

### 3.21.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3136 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + b*Sin[c + d*x]))], x]] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3233 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

### 3.21.4 Maple [A] (verified)

Time = 0.83 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.71

method	result
derivativedivides	$\frac{-\frac{639 \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{1024} - \frac{117 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{32} - \frac{369 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{64} + \frac{385 \arctan\left(\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2}\right)}{16384}}{8 \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\right)^3} d$
default	$\frac{-\frac{639 \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{1024} - \frac{117 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{32} - \frac{369 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{64} + \frac{385 \arctan\left(\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2}\right)}{16384}}{8 \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\right)^3} d$
risch	$-\frac{i(10395 e^{5i(dx+c)} + 86625 e^{4i(dx+c)} + 239470 e^{3i(dx+c)} + 218466 e^{2i(dx+c)} + 73575 e^{i(dx+c)} + 8397)}{12288d(3 e^{2i(dx+c)} + 10 e^{i(dx+c)} + 3)^3} + \frac{385i \ln(e^{i(dx+c)})}{32768d}$
parallelrisc	$\frac{385i(-770 - 27 \cos(3dx+3c) - 981 \cos(dx+c) - 270 \cos(2dx+2c)) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2i\right) + 385i(770 + 27 \cos(3dx+3c) + 981 \cos(dx+c) + 270 \cos(2dx+2c))}{32768d(770 + 27 \cos(3dx+3c) + 981 \cos(dx+c) + 270 \cos(2dx+2c))}$

3.21.  $\int \frac{1}{(5+3 \cos(c+dx))^4} dx$

```
input int(1/(5+3*cos(d*x+c))^4,x,method=_RETURNVERBOSE)
```

```
output 1/d*(1/8*(-639/1024*tan(1/2*d*x+1/2*c)^5-117/32*tan(1/2*d*x+1/2*c)^3-369/6
4*tan(1/2*d*x+1/2*c))/(tan(1/2*d*x+1/2*c)^2+4)^3+385/16384*arctan(1/2*tan(
1/2*d*x+1/2*c))
```

### 3.21.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.14

$$\int \frac{1}{(5 + 3 \cos(c + dx))^4} dx = \frac{385 (27 \cos(dx + c)^3 + 135 \cos(dx + c)^2 + 225 \cos(dx + c) + 125) \arctan\left(\frac{5 \cos(dx+c)+3}{4 \sin(dx+c)}\right) + 36 (311 \cos(dx + c)^2 + 1170 \cos(dx + c) + 1143) \sin(dx + c)}{32768 (27 d \cos(dx + c)^3 + 135 d \cos(dx + c)^2 + 225 d \cos(dx + c) + 125 d)}$$

```
input integrate(1/(5+3*cos(d*x+c))^4,x, algorithm="fricas")
```

```
output -1/32768*(385*(27*cos(d*x + c)^3 + 135*cos(d*x + c)^2 + 225*cos(d*x + c) +
125)*arctan(1/4*(5*cos(d*x + c) + 3)/sin(d*x + c)) + 36*(311*cos(d*x + c)
^2 + 1170*cos(d*x + c) + 1143)*sin(d*x + c))/(27*d*cos(d*x + c)^3 + 135*d*
cos(d*x + c)^2 + 225*d*cos(d*x + c) + 125*d)
```

### 3.21.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 3.07 (sec) , antiderivative size = 592, normalized size of antiderivative = 5.58

$$\int \frac{1}{(5 + 3 \cos(c + dx))^4} dx = \begin{cases} \frac{x}{(5+3 \cosh(2 \operatorname{atanh}(2)))^4} \\ \frac{x}{(3 \cos(c)+5)^4} \\ \frac{385 \left( \operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right) + \pi \left[ \frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2} \right] \right) \tan^6\left(\frac{c}{2} + \frac{dx}{2}\right)}{16384d \tan^6\left(\frac{c}{2} + \frac{dx}{2}\right) + 196608d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) + 786432d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 1048576d} + \frac{4620 \left( \operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right) + \pi \left[ \frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2} \right] \right) \tan^6\left(\frac{c}{2} + \frac{dx}{2}\right)}{16384d \tan^6\left(\frac{c}{2} + \frac{dx}{2}\right) + 196608d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) + 786432d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 1048576d} \end{cases}$$

---

3.21.  $\int \frac{1}{(5+3 \cos(c+dx))^4} dx$



input `integrate(1/(5+3*cos(d*x+c))**4,x)`

output `Piecewise((x/(5 + 3*cosh(2*atanh(2))))**4, Eq(c, -d*x - 2*I*atanh(2)) | Eq(c, -d*x + 2*I*atanh(2))), (x/(3*cos(c) + 5)**4, Eq(d, 0)), (385*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**6/(16384*d*tan(c/2 + d*x/2)**6 + 196608*d*tan(c/2 + d*x/2)**4 + 786432*d*tan(c/2 + d*x/2)**2 + 1048576*d) + 4620*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4/(16384*d*tan(c/2 + d*x/2)**6 + 196608*d*tan(c/2 + d*x/2)**4 + 786432*d*tan(c/2 + d*x/2)**2 + 1048576*d) + 18480*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**2/(16384*d*tan(c/2 + d*x/2)**6 + 196608*d*tan(c/2 + d*x/2)**4 + 786432*d*tan(c/2 + d*x/2)**2 + 1048576*d) + 24640*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(16384*d*tan(c/2 + d*x/2)**6 + 196608*d*tan(c/2 + d*x/2)**4 + 786432*d*tan(c/2 + d*x/2)**2 + 1048576*d) - 1278*tan(c/2 + d*x/2)**5/(16384*d*tan(c/2 + d*x/2)**6 + 196608*d*tan(c/2 + d*x/2)**4 + 786432*d*tan(c/2 + d*x/2)**2 + 1048576*d) - 7488*tan(c/2 + d*x/2)**3/(16384*d*tan(c/2 + d*x/2)**6 + 196608*d*tan(c/2 + d*x/2)**4 + 786432*d*tan(c/2 + d*x/2)**2 + 1048576*d) - 11808*tan(c/2 + d*x/2)/(16384*d*tan(c/2 + d*x/2)**6 + 196608*d*tan(c/2 + d*x/2)**4 + 786432*d*tan(c/2 + d*x/2)**2 + 1048576*d), True))`

### 3.21.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.42

$$\int \frac{1}{(5 + 3 \cos(c + dx))^4} dx$$

$$= -\frac{18 \left( \frac{656 \sin(dx+c)}{\cos(dx+c)+1} + \frac{416 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{71 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right) - 385 \arctan \left( \frac{\sin(dx+c)}{2(\cos(dx+c)+1)} \right)}{\frac{48 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{12 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{\sin(dx+c)^6}{(\cos(dx+c)+1)^6} + 64} 16384 d$$

input `integrate(1/(5+3*cos(d*x+c))^4,x, algorithm="maxima")`

output `-1/16384*(18*(656*sin(d*x + c)/(cos(d*x + c) + 1) + 416*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 71*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/(48*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 12*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + 64) - 385*arctan(1/2*sin(d*x + c)/(cos(d*x + c) + 1)))/d`

---

3.21.  $\int \frac{1}{(5+3 \cos(c+dx))^4} dx$

**3.21.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.83

$$\int \frac{1}{(5 + 3 \cos(c + dx))^4} dx$$

$$= \frac{385 dx + 385 c - \frac{36 \left( 71 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 + 416 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 656 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\left( \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 4 \right)^3} - 770 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+3}\right)}{32768 d}$$

input `integrate(1/(5+3*cos(d*x+c))^4,x, algorithm="giac")`output `1/32768*(385*d*x + 385*c - 36*(71*tan(1/2*d*x + 1/2*c)^5 + 416*tan(1/2*d*x + 1/2*c)^3 + 656*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 4)^3 - 770*arctan(sin(d*x + c)/(cos(d*x + c) + 3))/d`**3.21.9 Mupad [B] (verification not implemented)**

Time = 15.51 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.91

$$\int \frac{1}{(5 + 3 \cos(c + dx))^4} dx = \frac{385 \operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{16384 d} - \frac{385 \left( \operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2} \right)}{16384 d}$$

$$- \frac{\frac{639 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{8192} + \frac{117 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{256} + \frac{369 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{512}}{d \left( \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 4 \right)^3}$$

input `int(1/(3*cos(c + d*x) + 5)^4,x)`output `(385*atan(tan(c/2 + (d*x)/2)/2))/(16384*d) - (385*(atan(tan(c/2 + (d*x)/2) - (d*x)/2))/(16384*d) - ((369*tan(c/2 + (d*x)/2))/512 + (117*tan(c/2 + (d*x)/2)^3)/256 + (639*tan(c/2 + (d*x)/2)^5)/8192)/(d*(tan(c/2 + (d*x)/2)^2 + 4)^3)`

### 3.22 $\int \frac{1}{5-3 \cos(c+dx)} dx$

3.22.1	Optimal result . . . . .	154
3.22.2	Mathematica [A] (verified) . . . . .	154
3.22.3	Rubi [A] (verified) . . . . .	155
3.22.4	Maple [A] (verified) . . . . .	156
3.22.5	Fricas [A] (verification not implemented) . . . . .	156
3.22.6	Sympy [A] (verification not implemented) . . . . .	156
3.22.7	Maxima [A] (verification not implemented) . . . . .	157
3.22.8	Giac [A] (verification not implemented) . . . . .	157
3.22.9	Mupad [B] (verification not implemented) . . . . .	158

#### 3.22.1 Optimal result

Integrand size = 12, antiderivative size = 33

$$\int \frac{1}{5 - 3 \cos(c + dx)} dx = \frac{x}{4} + \frac{\arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{2d}$$

output `1/4*x+1/2*arctan(sin(d*x+c)/(3-cos(d*x+c)))/d`

#### 3.22.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.61

$$\int \frac{1}{5 - 3 \cos(c + dx)} dx = \frac{\arctan\left(2 \tan\left(\frac{1}{2}(c + dx)\right)\right)}{2d}$$

input `Integrate[(5 - 3*Cos[c + d*x])^(-1),x]`

output `ArcTan[2*Tan[(c + d*x)/2]]/(2*d)`

### 3.22.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{5 - 3 \cos(c + dx)} dx$$

↓ 3042

$$\int \frac{1}{5 - 3 \sin\left(c + dx + \frac{\pi}{2}\right)} dx$$

↓ 3136

$$\frac{\arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{2d} + \frac{x}{4}$$

input `Int[(5 - 3*Cos[c + d*x])^(-1),x]`

output `x/4 + ArcTan[Sin[c + d*x]/(3 - Cos[c + d*x])]/(2*d)`

#### 3.22.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3136 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

### 3.22.4 Maple [A] (verified)

Time = 1.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.55

method	result	size
derivativedivides	$\frac{\arctan\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2d}$	18
default	$\frac{\arctan\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2d}$	18
risch	$-\frac{i \ln\left(e^{i(dx+c)} - \frac{1}{3}\right)}{4d} + \frac{i \ln\left(e^{i(dx+c)} - 3\right)}{4d}$	38
parallelrisch	$-\frac{i\left(\ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - i\right) - \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + i\right)\right)}{4d}$	40

input `int(1/(5-3*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/2/d*arctan(2*tan(1/2*d*x+1/2*c))`

### 3.22.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

$$\int \frac{1}{5 - 3 \cos(c + dx)} dx = -\frac{\arctan\left(\frac{5 \cos(dx+c)-3}{4 \sin(dx+c)}\right)}{4d}$$

input `integrate(1/(5-3*cos(d*x+c)),x, algorithm="fricas")`

output `-1/4*arctan(1/4*(5*cos(d*x + c) - 3)/sin(d*x + c))/d`

### 3.22.6 Sympy [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.24

$$\int \frac{1}{5 - 3 \cos(c + dx)} dx = \begin{cases} \frac{\operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor}{2d} & \text{for } d \neq 0 \\ \frac{x}{5-3 \cos(c)} & \text{otherwise} \end{cases}$$

input `integrate(1/(5-3*cos(d*x+c)),x)`

output `Piecewise(((atan(2*tan(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/  
(2*d), Ne(d, 0)), (x/(5 - 3*cos(c)), True))`

### 3.22.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.73

$$\int \frac{1}{5 - 3 \cos(c + dx)} dx = \frac{\arctan\left(\frac{2 \sin(dx+c)}{\cos(dx+c)+1}\right)}{2d}$$

input `integrate(1/(5-3*cos(d*x+c)),x, algorithm="maxima")`

output `1/2*arctan(2*sin(d*x + c)/(cos(d*x + c) + 1))/d`

### 3.22.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.91

$$\int \frac{1}{5 - 3 \cos(c + dx)} dx = \frac{dx + c - 2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)-3}\right)}{4d}$$

input `integrate(1/(5-3*cos(d*x+c)),x, algorithm="giac")`

output `1/4*(d*x + c - 2*arctan(sin(d*x + c)/(cos(d*x + c) - 3)))/d`

**3.22.9 Mupad [B] (verification not implemented)**

Time = 15.49 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.15

$$\int \frac{1}{5 - 3 \cos(c + dx)} dx = \frac{\operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{2d} - \frac{\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}}{2d}$$

input `int(-1/(3*cos(c + d*x) - 5),x)`

output `atan(2*tan(c/2 + (d*x)/2))/(2*d) - (atan(tan(c/2 + (d*x)/2)) - (d*x)/2)/(2*d)`

### 3.23 $\int \frac{1}{(5-3 \cos(c+dx))^2} dx$

3.23.1 Optimal result . . . . .	159
3.23.2 Mathematica [A] (verified) . . . . .	159
3.23.3 Rubi [A] (verified) . . . . .	160
3.23.4 Maple [A] (verified) . . . . .	161
3.23.5 Fricas [A] (verification not implemented) . . . . .	162
3.23.6 Sympy [C] (verification not implemented) . . . . .	162
3.23.7 Maxima [A] (verification not implemented) . . . . .	163
3.23.8 Giac [A] (verification not implemented) . . . . .	163
3.23.9 Mupad [B] (verification not implemented) . . . . .	163

#### 3.23.1 Optimal result

Integrand size = 12, antiderivative size = 58

$$\int \frac{1}{(5-3 \cos(c+dx))^2} dx = \frac{5x}{64} + \frac{5 \arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{32d} + \frac{3 \sin(c+dx)}{16d(5-3 \cos(c+dx))}$$

output `5/64*x+5/32*arctan(sin(d*x+c)/(3-cos(d*x+c)))/d+3/16*sin(d*x+c)/d/(5-3*cos(d*x+c))`

#### 3.23.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.74

$$\int \frac{1}{(5-3 \cos(c+dx))^2} dx = \frac{5 \arctan\left(2 \tan\left(\frac{1}{2}(c+dx)\right)\right) - \frac{6 \sin(c+dx)}{-5+3 \cos(c+dx)}}{32d}$$

input `Integrate[(5 - 3*Cos[c + d*x])^(-2),x]`

output `(5*ArcTan[2*Tan[(c + d*x)/2]] - (6*Sin[c + d*x])/(-5 + 3*Cos[c + d*x]))/(32*d)`



### 3.23.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {3042, 3143, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(5 - 3 \cos(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(5 - 3 \sin(c + dx + \frac{\pi}{2}))^2} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{3 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))} - \frac{1}{16} \int -\frac{5}{5 - 3 \cos(c + dx)} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{5}{16} \int \frac{1}{5 - 3 \cos(c + dx)} dx + \frac{3 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{16} \int \frac{1}{5 - 3 \sin(c + dx + \frac{\pi}{2})} dx + \frac{3 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))} \\
 & \quad \downarrow \text{3136} \\
 & \frac{5}{16} \left( \frac{\arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{2d} + \frac{x}{4} \right) + \frac{3 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))}
 \end{aligned}$$

input `Int[(5 - 3*Cos[c + d*x])^(-2), x]`

output `(5*(x/4 + ArcTan[Sin[c + d*x]/(3 - Cos[c + d*x])]/(2*d)))/16 + (3*Sin[c + d*x])/(16*d*(5 - 3*Cos[c + d*x]))`

## 3.23.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

rule 3143 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

## 3.23.4 Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.79

method	result	size
derivativedivides	$\frac{\frac{3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{64 \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{1}{4}\right)} + \frac{5 \arctan\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{32}}{d}$	46
default	$\frac{\frac{3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{64 \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{1}{4}\right)} + \frac{5 \arctan\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{32}}{d}$	46
parallelrisch	$\frac{(-15i \cos(dx+c) + 25i) \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - i\right) + (15i \cos(dx+c) - 25i) \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + i\right) - 12 \sin(dx+c)}{192d \cos(dx+c) - 320d}$	82
risch	$\frac{i(5e^{i(dx+c)} - 3)}{8d(3e^{2i(dx+c)} - 10e^{i(dx+c)} + 3)} + \frac{5i \ln(e^{i(dx+c)} - 3)}{64d} - \frac{5i \ln(e^{i(dx+c)} - \frac{1}{3})}{64d}$	83

input `int(1/(5-3*cos(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(3/64*tan(1/2*d*x+1/2*c)/(tan(1/2*d*x+1/2*c)^2+1/4)+5/32*arctan(2*tan(1/2*d*x+1/2*c)))`

### 3.23.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.02

$$\int \frac{1}{(5 - 3 \cos(c + dx))^2} dx = -\frac{5(3 \cos(dx + c) - 5) \arctan\left(\frac{5 \cos(dx+c)-3}{4 \sin(dx+c)}\right) + 12 \sin(dx + c)}{64(3d \cos(dx + c) - 5d)}$$

input `integrate(1/(5-3*cos(d*x+c))^2,x, algorithm="fricas")`

output `-1/64*(5*(3*cos(d*x + c) - 5)*arctan(1/4*(5*cos(d*x + c) - 3)/sin(d*x + c)) + 12*sin(d*x + c))/(3*d*cos(d*x + c) - 5*d)`

### 3.23.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.82 (sec) , antiderivative size = 192, normalized size of antiderivative = 3.31

$$\int \frac{1}{(5 - 3 \cos(c + dx))^2} dx = \begin{cases} \frac{x}{(5-3 \cosh(2 \operatorname{atanh}(\frac{1}{2})))^2} \\ \frac{x}{(5-3 \cos(c))^2} \\ \frac{20 \left( \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right) \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{128d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 32d} + \frac{5 \left( \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{128d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 32d} + \frac{6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{128d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 32d} \end{cases}$$

input `integrate(1/(5-3*cos(d*x+c))**2,x)`

output `Piecewise((x/(5 - 3*cosh(2*atanh(1/2)))**2, Eq(c, -d*x - 2*I*atanh(1/2)) | Eq(c, -d*x + 2*I*atanh(1/2))), (x/(5 - 3*cos(c))**2, Eq(d, 0)), (20*(atan(2*tan(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**2/(128*d*tan(c/2 + d*x/2)**2 + 32*d) + 5*(atan(2*tan(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(128*d*tan(c/2 + d*x/2)**2 + 32*d) + 6*tan(c/2 + d*x/2)/(128*d*tan(c/2 + d*x/2)**2 + 32*d), True))`

**3.23.7 Maxima [A] (verification not implemented)**

Time = 0.41 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.19

$$\int \frac{1}{(5 - 3 \cos(c + dx))^2} dx = \frac{\frac{6 \sin(dx+c)}{\left(\frac{4 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 1\right)(\cos(dx+c)+1)} + 5 \arctan\left(\frac{2 \sin(dx+c)}{\cos(dx+c)+1}\right)}{32 d}$$

input `integrate(1/(5-3*cos(d*x+c))^2,x, algorithm="maxima")`output `1/32*(6*sin(d*x + c)/((4*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)*(cos(d*x + c) + 1)) + 5*arctan(2*sin(d*x + c)/(cos(d*x + c) + 1)))/d`**3.23.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.05

$$\int \frac{1}{(5 - 3 \cos(c + dx))^2} dx = \frac{5 dx + 5 c + \frac{12 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{4 \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1} - 10 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)-3}\right)}{64 d}$$

input `integrate(1/(5-3*cos(d*x+c))^2,x, algorithm="giac")`output `1/64*(5*d*x + 5*c + 12*tan(1/2*d*x + 1/2*c)/(4*tan(1/2*d*x + 1/2*c)^2 + 1) - 10*arctan(sin(d*x + c)/(cos(d*x + c) - 3)))/d`**3.23.9 Mupad [B] (verification not implemented)**

Time = 15.19 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.16

$$\int \frac{1}{(5 - 3 \cos(c + dx))^2} dx = \frac{5 \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{32 d} - \frac{5 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{32 d} + \frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{64 d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + \frac{1}{4}\right)}$$

input `int(1/(3*cos(c + d*x) - 5)^2,x)`output `(5*atan(2*tan(c/2 + (d*x)/2)))/(32*d) - (5*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2))/(32*d) + (3*tan(c/2 + (d*x)/2))/(64*d*(tan(c/2 + (d*x)/2)^2 + 1/4))`

### 3.24 $\int \frac{1}{(5-3\cos(c+dx))^3} dx$

3.24.1	Optimal result . . . . .	164
3.24.2	Mathematica [A] (verified) . . . . .	164
3.24.3	Rubi [A] (verified) . . . . .	165
3.24.4	Maple [A] (verified) . . . . .	167
3.24.5	Fricas [A] (verification not implemented) . . . . .	167
3.24.6	Sympy [C] (verification not implemented) . . . . .	168
3.24.7	Maxima [A] (verification not implemented) . . . . .	168
3.24.8	Giac [A] (verification not implemented) . . . . .	169
3.24.9	Mupad [B] (verification not implemented) . . . . .	169

#### 3.24.1 Optimal result

Integrand size = 12, antiderivative size = 83

$$\int \frac{1}{(5-3\cos(c+dx))^3} dx = \frac{59x}{2048} + \frac{59 \arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{1024d} + \frac{3\sin(c+dx)}{32d(5-3\cos(c+dx))^2} + \frac{45\sin(c+dx)}{512d(5-3\cos(c+dx))}$$

output `59/2048*x+59/1024*arctan(sin(d*x+c)/(3-cos(d*x+c)))/d+3/32*sin(d*x+c)/d/(5-3*cos(d*x+c))^2+45/512*sin(d*x+c)/d/(5-3*cos(d*x+c))`

#### 3.24.2 Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.78

$$\int \frac{1}{(5-3\cos(c+dx))^3} dx = \frac{59 \arctan\left(2 \tan\left(\frac{1}{2}(c+dx)\right)\right) (5-3\cos(c+dx))^2 + 546\sin(c+dx) - 135\sin(2(c+dx))}{1024d(5-3\cos(c+dx))^2}$$

input `Integrate[(5 - 3*Cos[c + d*x])^(-3),x]`

output `(59*ArcTan[2*Tan[(c + d*x)/2]]*(5 - 3*Cos[c + d*x])^2 + 546*Sin[c + d*x] - 135*Sin[2*(c + d*x)])/(1024*d*(5 - 3*Cos[c + d*x])^2)`

**3.24.3 Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.12, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {3042, 3143, 25, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(5 - 3 \cos(c + dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(5 - 3 \sin(c + dx + \frac{\pi}{2}))^3} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{3 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} - \frac{1}{32} \int -\frac{3 \cos(c + dx) + 10}{(5 - 3 \cos(c + dx))^2} dx \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{32} \int \frac{3 \cos(c + dx) + 10}{(5 - 3 \cos(c + dx))^2} dx + \frac{3 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{32} \int \frac{3 \sin(c + dx + \frac{\pi}{2}) + 10}{(5 - 3 \sin(c + dx + \frac{\pi}{2}))^2} dx + \frac{3 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} \\
 & \quad \downarrow \text{3233} \\
 & \frac{1}{32} \left( \frac{45 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))} - \frac{1}{16} \int -\frac{59}{5 - 3 \cos(c + dx)} dx \right) + \frac{3 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{32} \left( \frac{59}{16} \int \frac{1}{5 - 3 \cos(c + dx)} dx + \frac{45 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))} \right) + \frac{3 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{32} \left( \frac{59}{16} \int \frac{1}{5 - 3 \sin(c + dx + \frac{\pi}{2})} dx + \frac{45 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))} \right) + \frac{3 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} \\
 & \quad \downarrow \text{3136} \\
 & \frac{1}{32} \left( \frac{59}{16} \left( \frac{\arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{2d} + \frac{x}{4} \right) + \frac{45 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))} \right) + \frac{3 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2}
 \end{aligned}$$

input `Int[(5 - 3*Cos[c + d*x])^(-3),x]`

output `(3*Sin[c + d*x])/(32*d*(5 - 3*Cos[c + d*x])^2) + ((59*(x/4 + ArcTan[Sin[c + d*x]/(3 - Cos[c + d*x]])/(2*d)))/16 + (45*Sin[c + d*x])/(16*d*(5 - 3*Cos[c + d*x]))) / 32`

### 3.24.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3233 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

### 3.24.4 Maple [A] (verified)

Time = 0.83 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.77

method	result
derivativedivides	$\frac{\frac{51 \left( \tan^3 \left( \frac{dx}{2} + \frac{c}{2} \right) \right) + \frac{69 \tan \left( \frac{dx}{2} + \frac{c}{2} \right)}{512}}{\left( 4 \left( \tan^2 \left( \frac{dx}{2} + \frac{c}{2} \right) \right) + 1 \right)^2} + \frac{59 \arctan \left( 2 \tan \left( \frac{dx}{2} + \frac{c}{2} \right) \right)}{1024}}{d}$
default	$\frac{\frac{51 \left( \tan^3 \left( \frac{dx}{2} + \frac{c}{2} \right) \right) + \frac{69 \tan \left( \frac{dx}{2} + \frac{c}{2} \right)}{512}}{\left( 4 \left( \tan^2 \left( \frac{dx}{2} + \frac{c}{2} \right) \right) + 1 \right)^2} + \frac{59 \arctan \left( 2 \tan \left( \frac{dx}{2} + \frac{c}{2} \right) \right)}{1024}}{d}$
risch	$\frac{3i \left( 59 e^{3i(dx+c)} - 295 e^{2i(dx+c)} + 241 e^{i(dx+c)} - 45 \right)}{256d \left( 3 e^{2i(dx+c)} - 10 e^{i(dx+c)} + 3 \right)^2} - \frac{59i \ln \left( e^{i(dx+c)} - \frac{1}{3} \right)}{2048d} + \frac{59i \ln \left( e^{i(dx+c)} - 3 \right)}{2048d}$
parallelrisc	$\frac{59i \left( 59 + 9 \cos(2dx+2c) - 60 \cos(dx+c) \right) \ln \left( 2 \tan \left( \frac{dx}{2} + \frac{c}{2} \right) - i \right) + 59i \left( -9 \cos(2dx+2c) - 59 + 60 \cos(dx+c) \right) \ln \left( 2 \tan \left( \frac{dx}{2} + \frac{c}{2} \right) + i \right)}{2048d \left( -9 \cos(2dx+2c) - 59 + 60 \cos(dx+c) \right)}$

input `int(1/(5-3*cos(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/d*(4*(51/512*tan(1/2*d*x+1/2*c)^3+69/2048*tan(1/2*d*x+1/2*c))/(4*tan(1/2*d*x+1/2*c)^2+1)^2+59/1024*arctan(2*tan(1/2*d*x+1/2*c))`

### 3.24.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.08

$$\int \frac{1}{(5 - 3 \cos(c + dx))^3} dx = \frac{59 \left( 9 \cos(dx + c)^2 - 30 \cos(dx + c) + 25 \right) \arctan \left( \frac{5 \cos(dx+c) - 3}{4 \sin(dx+c)} \right) + 12 \left( 45 \cos(dx + c) - 91 \right) \sin(dx + c)}{2048 \left( 9 d \cos(dx + c)^2 - 30 d \cos(dx + c) + 25 d \right)}$$

input `integrate(1/(5-3*cos(d*x+c))^3,x, algorithm="fricas")`

output `-1/2048*(59*(9*cos(d*x + c)^2 - 30*cos(d*x + c) + 25)*arctan(1/4*(5*cos(d*x + c) - 3)/sin(d*x + c)) + 12*(45*cos(d*x + c) - 91)*sin(d*x + c))/(9*d*cos(d*x + c)^2 - 30*d*cos(d*x + c) + 25*d)`



### 3.24.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.43 (sec) , antiderivative size = 364, normalized size of antiderivative = 4.39

$$\int \frac{1}{(5 - 3 \cos(c + dx))^3} dx$$

$$= \begin{cases} \frac{x}{(5 - 3 \cosh(2 \operatorname{atanh}(\frac{1}{2})))^3} \\ \frac{x}{(5 - 3 \cos(c))^3} \\ \frac{944 \left( \operatorname{atan} \left( 2 \tan \left( \frac{c}{2} + \frac{dx}{2} \right) \right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right) \tan^4 \left( \frac{c}{2} + \frac{dx}{2} \right)}{16384d \tan^4 \left( \frac{c}{2} + \frac{dx}{2} \right) + 8192d \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right) + 1024d} + \frac{472 \left( \operatorname{atan} \left( 2 \tan \left( \frac{c}{2} + \frac{dx}{2} \right) \right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right) \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right)}{16384d \tan^4 \left( \frac{c}{2} + \frac{dx}{2} \right) + 8192d \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right) + 1024d} + \frac{59 \left( \operatorname{atan} \left( 2 \tan \left( \frac{c}{2} + \frac{dx}{2} \right) \right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right) \tan \left( \frac{c}{2} + \frac{dx}{2} \right)}{16384d \tan^4 \left( \frac{c}{2} + \frac{dx}{2} \right) + 8192d \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right) + 1024d} \end{cases}$$

input `integrate(1/(5-3*cos(d*x+c))**3,x)`

output `Piecewise((x/(5 - 3*cosh(2*atanh(1/2)))**3, Eq(c, -d*x - 2*I*atanh(1/2)) | Eq(c, -d*x + 2*I*atanh(1/2))), (x/(5 - 3*cos(c))**3, Eq(d, 0)), (944*(atan(2*tan(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4/(16384*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 1024*d) + 472*(atan(2*tan(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**2/(16384*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 1024*d) + 59*(atan(2*tan(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(16384*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 1024*d) + 408*tan(c/2 + d*x/2)**3/(16384*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 1024*d) + 138*tan(c/2 + d*x/2)/(16384*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 1024*d), True))`

### 3.24.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.35

$$\int \frac{1}{(5 - 3 \cos(c + dx))^3} dx = \frac{6 \left( \frac{23 \sin(dx+c)}{\cos(dx+c)+1} + \frac{68 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right) + 59 \arctan \left( \frac{2 \sin(dx+c)}{\cos(dx+c)+1} \right)}{1024 d}$$

input `integrate(1/(5-3*cos(d*x+c))^3,x, algorithm="maxima")`

output  $1/1024*(6*(23*\sin(d*x + c)/(\cos(d*x + c) + 1) + 68*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3)/(8*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2 + 16*\sin(d*x + c)^4/(\cos(d*x + c) + 1)^4 + 1) + 59*\arctan(2*\sin(d*x + c)/(\cos(d*x + c) + 1)))/d$

### 3.24.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.93

$$\int \frac{1}{(5 - 3 \cos(c + dx))^3} dx$$

$$= \frac{59 dx + 59 c + \frac{12 \left( 68 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 23 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\left( 4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 1 \right)^2} - 118 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)-3}\right)}{2048 d}$$

input `integrate(1/(5-3*cos(d*x+c))^3,x, algorithm="giac")`

output  $1/2048*(59*d*x + 59*c + 12*(68*\tan(1/2*d*x + 1/2*c)^3 + 23*\tan(1/2*d*x + 1/2*c))/(4*\tan(1/2*d*x + 1/2*c)^2 + 1)^2 - 118*\arctan(\sin(d*x + c)/(\cos(d*x + c) - 3)))/d$

### 3.24.9 Mupad [B] (verification not implemented)

Time = 15.24 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.14

$$\int \frac{1}{(5 - 3 \cos(c + dx))^3} dx = \frac{59 \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{1024 d} - \frac{59 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{1024 d}$$

$$+ \frac{\frac{51 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{2048} + \frac{69 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{8192}}{d \left( \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{2} + \frac{1}{16} \right)}$$

input `int(-1/(3*cos(c + d*x) - 5)^3,x)`

output  $(59*\operatorname{atan}(2*\tan(c/2 + (d*x)/2)))/(1024*d) - (59*(\operatorname{atan}(\tan(c/2 + (d*x)/2)) - (d*x)/2))/(1024*d) + ((69*\tan(c/2 + (d*x)/2))/8192 + (51*\tan(c/2 + (d*x)/2)^3)/2048)/(d*(\tan(c/2 + (d*x)/2)^2/2 + \tan(c/2 + (d*x)/2)^4 + 1/16))$

### 3.25 $\int \frac{1}{(5-3 \cos(c+dx))^4} dx$

3.25.1	Optimal result	170
3.25.2	Mathematica [A] (verified)	170
3.25.3	Rubi [A] (verified)	171
3.25.4	Maple [A] (verified)	173
3.25.5	Fricas [A] (verification not implemented)	174
3.25.6	Sympy [C] (verification not implemented)	174
3.25.7	Maxima [A] (verification not implemented)	175
3.25.8	Giac [A] (verification not implemented)	176
3.25.9	Mupad [B] (verification not implemented)	176

#### 3.25.1 Optimal result

Integrand size = 12, antiderivative size = 108

$$\int \frac{1}{(5-3 \cos(c+dx))^4} dx = \frac{385x}{32768} + \frac{385 \arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{16384d} + \frac{\sin(c+dx)}{16d(5-3 \cos(c+dx))^3} + \frac{25 \sin(c+dx)}{512d(5-3 \cos(c+dx))^2} + \frac{311 \sin(c+dx)}{8192d(5-3 \cos(c+dx))}$$

output `385/32768*x+385/16384*arctan(sin(d*x+c)/(3-cos(d*x+c)))/d+1/16*sin(d*x+c)/d/(5-3*cos(d*x+c))^3+25/512*sin(d*x+c)/d/(5-3*cos(d*x+c))^2+311/8192*sin(d*x+c)/d/(5-3*cos(d*x+c))`

#### 3.25.2 Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.61

$$\int \frac{1}{(5-3 \cos(c+dx))^4} dx = \frac{770 \arctan\left(2 \tan\left(\frac{1}{2}(c+dx)\right)\right) - \frac{9(4883 \sin(c+dx) - 2340 \sin(2(c+dx)) + 311 \sin(3(c+dx)))}{(-5+3 \cos(c+dx))^3}}{32768d}$$

input `Integrate[(5 - 3*Cos[c + d*x])^(-4),x]`

output `(770*ArcTan[2*Tan[(c + d*x)/2]] - (9*(4883*Sin[c + d*x] - 2340*Sin[2*(c + d*x)] + 311*Sin[3*(c + d*x)]))/(-5 + 3*Cos[c + d*x])^3/(32768*d)`

### 3.25.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.14, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$ , Rules used = {3042, 3143, 27, 3042, 3233, 25, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(5 - 3 \cos(c + dx))^4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(5 - 3 \sin(c + dx + \frac{\pi}{2}))^4} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{\sin(c + dx)}{16d(5 - 3 \cos(c + dx))^3} - \frac{1}{48} \int -\frac{3(2 \cos(c + dx) + 5)}{(5 - 3 \cos(c + dx))^3} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{16} \int \frac{2 \cos(c + dx) + 5}{(5 - 3 \cos(c + dx))^3} dx + \frac{\sin(c + dx)}{16d(5 - 3 \cos(c + dx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{16} \int \frac{2 \sin(c + dx + \frac{\pi}{2}) + 5}{(5 - 3 \sin(c + dx + \frac{\pi}{2}))^3} dx + \frac{\sin(c + dx)}{16d(5 - 3 \cos(c + dx))^3} \\
 & \quad \downarrow \text{3233} \\
 & \frac{1}{16} \left( \frac{25 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} - \frac{1}{32} \int -\frac{25 \cos(c + dx) + 62}{(5 - 3 \cos(c + dx))^2} dx \right) + \frac{\sin(c + dx)}{16d(5 - 3 \cos(c + dx))^3} \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{16} \left( \frac{1}{32} \int \frac{25 \cos(c + dx) + 62}{(5 - 3 \cos(c + dx))^2} dx + \frac{25 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} \right) + \frac{\sin(c + dx)}{16d(5 - 3 \cos(c + dx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{16} \left( \frac{1}{32} \int \frac{25 \sin(c + dx + \frac{\pi}{2}) + 62}{(5 - 3 \sin(c + dx + \frac{\pi}{2}))^2} dx + \frac{25 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} \right) + \frac{\sin(c + dx)}{16d(5 - 3 \cos(c + dx))^3} \\
 & \quad \downarrow \text{3233}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{16} \left( \frac{1}{32} \left( \frac{311 \sin(c+dx)}{16d(5-3\cos(c+dx))} - \frac{1}{16} \int \frac{385}{5-3\cos(c+dx)} dx \right) + \frac{25 \sin(c+dx)}{32d(5-3\cos(c+dx))^2} \right) + \\
& \quad \frac{\sin(c+dx)}{16d(5-3\cos(c+dx))^3} \\
& \quad \downarrow \text{27} \\
& \frac{1}{16} \left( \frac{1}{32} \left( \frac{385}{16} \int \frac{1}{5-3\cos(c+dx)} dx + \frac{311 \sin(c+dx)}{16d(5-3\cos(c+dx))} \right) + \frac{25 \sin(c+dx)}{32d(5-3\cos(c+dx))^2} \right) + \\
& \quad \frac{\sin(c+dx)}{16d(5-3\cos(c+dx))^3} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{16} \left( \frac{1}{32} \left( \frac{385}{16} \int \frac{1}{5-3\sin(c+dx+\frac{\pi}{2})} dx + \frac{311 \sin(c+dx)}{16d(5-3\cos(c+dx))} \right) + \frac{25 \sin(c+dx)}{32d(5-3\cos(c+dx))^2} \right) + \\
& \quad \frac{\sin(c+dx)}{16d(5-3\cos(c+dx))^3} \\
& \quad \downarrow \text{3136} \\
& \frac{1}{16} \left( \frac{1}{32} \left( \frac{385}{16} \left( \frac{\arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{2d} + \frac{x}{4} \right) + \frac{311 \sin(c+dx)}{16d(5-3\cos(c+dx))} \right) + \frac{25 \sin(c+dx)}{32d(5-3\cos(c+dx))^2} \right) + \\
& \quad \frac{\sin(c+dx)}{16d(5-3\cos(c+dx))^3}
\end{aligned}$$

input `Int[(5 - 3*Cos[c + d*x])^(-4),x]`

output `Sin[c + d*x]/(16*d*(5 - 3*Cos[c + d*x])^3) + ((25*Sin[c + d*x])/(32*d*(5 - 3*Cos[c + d*x])^2) + ((385*(x/4 + ArcTan[Sin[c + d*x]/(3 - Cos[c + d*x]))/(2*d)))/16 + (311*Sin[c + d*x])/(16*d*(5 - 3*Cos[c + d*x])))/32)/16`

### 3.25.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3136 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + b*Sin[c + d*x]))], x]] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3233 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

### 3.25.4 Maple [A] (verified)

Time = 0.84 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.71

method	result
derivativedivides	$\frac{\frac{369 \tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{512} + \frac{117 \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{256} + \frac{639 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8192} + \frac{385 \arctan\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{16384}}{\left(4 \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^3} \frac{d}{d}$
default	$\frac{\frac{369 \tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{512} + \frac{117 \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{256} + \frac{639 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8192} + \frac{385 \arctan\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{16384}}{\left(4 \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^3} \frac{d}{d}$
risch	$\frac{i(10395 e^{5i(dx+c)} - 86625 e^{4i(dx+c)} + 239470 e^{3i(dx+c)} - 218466 e^{2i(dx+c)} + 73575 e^{i(dx+c)} - 8397)}{12288d(3 e^{2i(dx+c)} - 10 e^{i(dx+c)} + 3)^3} - \frac{385i \ln(e^{i(dx+c)} - i)}{32768d}$
parallelrisc	$\frac{385i(770 - 27 \cos(3dx+3c) - 981 \cos(dx+c) + 270 \cos(2dx+2c)) \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - i\right) + 385i(27 \cos(3dx+3c) + 981 \cos(dx+c))}{32768d(27 \cos(3dx+3c) + 981 \cos(dx+c))}$

input `int(1/(5-3*cos(d*x+c))^4,x,method=_RETURNVERBOSE)`

3.25.  $\int \frac{1}{(5-3\cos(c+dx))^4} dx$

output  $1/d*(8*(369/4096*\tan(1/2*d*x+1/2*c)^5+117/2048*\tan(1/2*d*x+1/2*c)^3+639/65$   
 $536*\tan(1/2*d*x+1/2*c))/(4*\tan(1/2*d*x+1/2*c)^2+1)^3+385/16384*\arctan(2*\tan(1/2*d*x+1/2*c))$

### 3.25.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.12

$$\int \frac{1}{(5 - 3 \cos(c + dx))^4} dx = \frac{385 (27 \cos(dx + c)^3 - 135 \cos(dx + c)^2 + 225 \cos(dx + c) - 125) \arctan\left(\frac{5 \cos(dx+c)-3}{4 \sin(dx+c)}\right) + 36 (311 \cos(dx + c)^2 - 1170 \cos(dx + c) + 1143) \sin(dx + c)}{32768 (27 d \cos(dx + c)^3 - 135 d \cos(dx + c)^2 + 225 d \cos(dx + c) - 125 d)}$$

input `integrate(1/(5-3*cos(d*x+c))^4,x, algorithm="fricas")`

output  $-1/32768*(385*(27*\cos(d*x + c)^3 - 135*\cos(d*x + c)^2 + 225*\cos(d*x + c) -$   
 $125)*\arctan(1/4*(5*\cos(d*x + c) - 3)/\sin(d*x + c)) + 36*(311*\cos(d*x + c)$   
 $^2 - 1170*\cos(d*x + c) + 1143)*\sin(d*x + c))/(27*d*\cos(d*x + c)^3 - 135*d*$   
 $\cos(d*x + c)^2 + 225*d*\cos(d*x + c) - 125*d)$

### 3.25.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.85 (sec) , antiderivative size = 597, normalized size of antiderivative = 5.53

$$\int \frac{1}{(5 - 3 \cos(c + dx))^4} dx = \begin{cases} \frac{x}{(5-3 \cosh(2 \operatorname{atanh}(\frac{1}{2})))^4} \\ \frac{x}{(5-3 \cos(c))^4} \\ \frac{24640 \left( \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) + \pi \left\lfloor \frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2} \right\rfloor \right) \tan^6\left(\frac{c}{2} + \frac{dx}{2}\right)}{1048576d \tan^6\left(\frac{c}{2} + \frac{dx}{2}\right) + 786432d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) + 196608d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 16384d} + \frac{18480 \left( \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) + \pi \left\lfloor \frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2} \right\rfloor \right) \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right)}{1048576d \tan^6\left(\frac{c}{2} + \frac{dx}{2}\right) + 786432d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) + 196608d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 16384d} \end{cases}$$

input `integrate(1/(5-3*cos(d*x+c))**4,x)`

```

output Piecewise((x/(5 - 3*cosh(2*atanh(1/2)))**4, Eq(c, -d*x - 2*I*atanh(1/2)) |
Eq(c, -d*x + 2*I*atanh(1/2))), (x/(5 - 3*cos(c))**4, Eq(d, 0)), (24640*(a
tan(2*tan(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x
/2)**6/(1048576*d*tan(c/2 + d*x/2)**6 + 786432*d*tan(c/2 + d*x/2)**4 + 196
608*d*tan(c/2 + d*x/2)**2 + 16384*d) + 18480*(atan(2*tan(c/2 + d*x/2)) + p
i*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4/(1048576*d*tan(c/2 +
d*x/2)**6 + 786432*d*tan(c/2 + d*x/2)**4 + 196608*d*tan(c/2 + d*x/2)**2 +
16384*d) + 4620*(atan(2*tan(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)
/pi))*tan(c/2 + d*x/2)**2/(1048576*d*tan(c/2 + d*x/2)**6 + 786432*d*tan(c/
2 + d*x/2)**4 + 196608*d*tan(c/2 + d*x/2)**2 + 16384*d) + 385*(atan(2*tan(
c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(1048576*d*tan(c/2 + d*
x/2)**6 + 786432*d*tan(c/2 + d*x/2)**4 + 196608*d*tan(c/2 + d*x/2)**2 + 16
384*d) + 11808*tan(c/2 + d*x/2)**5/(1048576*d*tan(c/2 + d*x/2)**6 + 786432
*d*tan(c/2 + d*x/2)**4 + 196608*d*tan(c/2 + d*x/2)**2 + 16384*d) + 7488*ta
n(c/2 + d*x/2)**3/(1048576*d*tan(c/2 + d*x/2)**6 + 786432*d*tan(c/2 + d*x/
2)**4 + 196608*d*tan(c/2 + d*x/2)**2 + 16384*d) + 1278*tan(c/2 + d*x/2)/(1
048576*d*tan(c/2 + d*x/2)**6 + 786432*d*tan(c/2 + d*x/2)**4 + 196608*d*tan
(c/2 + d*x/2)**2 + 16384*d), True))

```

### 3.25.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.41

$$\int \frac{1}{(5 - 3 \cos(c + dx))^4} dx$$

$$= \frac{18 \left( \frac{71 \sin(dx+c)}{\cos(dx+c)+1} + \frac{416 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{656 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right) + 385 \arctan \left( \frac{2 \sin(dx+c)}{\cos(dx+c)+1} \right)}{16384 d}$$

```

input integrate(1/(5-3*cos(d*x+c))^4,x, algorithm="maxima")

```

```

output 1/16384*(18*(71*sin(d*x + c)/(cos(d*x + c) + 1) + 416*sin(d*x + c)^3/(cos(
d*x + c) + 1)^3 + 656*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/(12*sin(d*x + c
)^2/(cos(d*x + c) + 1)^2 + 48*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 64*sin
(d*x + c)^6/(cos(d*x + c) + 1)^6 + 1) + 385*arctan(2*sin(d*x + c)/(cos(d*x
+ c) + 1)))/d

```



**3.25.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.83

$$\int \frac{1}{(5 - 3 \cos(c + dx))^4} dx$$

$$= \frac{385 dx + 385 c + \frac{36 \left( 656 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 + 416 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 71 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\left( 4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 1 \right)^3} - 770 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)-3}\right)}{32768 d}$$

input `integrate(1/(5-3*cos(d*x+c))^4,x, algorithm="giac")`output `1/32768*(385*d*x + 385*c + 36*(656*tan(1/2*d*x + 1/2*c)^5 + 416*tan(1/2*d*x + 1/2*c)^3 + 71*tan(1/2*d*x + 1/2*c))/(4*tan(1/2*d*x + 1/2*c)^2 + 1)^3 - 770*arctan(sin(d*x + c)/(cos(d*x + c) - 3))/d`**3.25.9 Mupad [B] (verification not implemented)**

Time = 15.63 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.90

$$\int \frac{1}{(5 - 3 \cos(c + dx))^4} dx = \frac{385 \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{16384 d} - \frac{385 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{16384 d}$$

$$+ \frac{\frac{369 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{512} + \frac{117 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{256} + \frac{639 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{8192}}{d \left(4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1\right)^3}$$

input `int(1/(3*cos(c + d*x) - 5)^4,x)`output `(385*atan(2*tan(c/2 + (d*x)/2)))/(16384*d) - (385*(atan(tan(c/2 + (d*x)/2) - (d*x)/2)))/(16384*d) + ((639*tan(c/2 + (d*x)/2))/8192 + (117*tan(c/2 + (d*x)/2)^3)/256 + (369*tan(c/2 + (d*x)/2)^5)/512)/(d*(4*tan(c/2 + (d*x)/2)^2 + 1)^3)`

## 3.26 $\int \frac{1}{-5+3\cos(c+dx)} dx$

3.26.1	Optimal result	177
3.26.2	Mathematica [A] (verified)	177
3.26.3	Rubi [A] (verified)	178
3.26.4	Maple [A] (verified)	179
3.26.5	Fricas [A] (verification not implemented)	179
3.26.6	Sympy [A] (verification not implemented)	179
3.26.7	Maxima [A] (verification not implemented)	180
3.26.8	Giac [A] (verification not implemented)	180
3.26.9	Mupad [B] (verification not implemented)	181

### 3.26.1 Optimal result

Integrand size = 12, antiderivative size = 33

$$\int \frac{1}{-5+3\cos(c+dx)} dx = -\frac{x}{4} - \frac{\arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{2d}$$

output `-1/4*x-1/2*arctan(sin(d*x+c)/(3-cos(d*x+c)))/d`

### 3.26.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.61

$$\int \frac{1}{-5+3\cos(c+dx)} dx = -\frac{\arctan\left(2\tan\left(\frac{1}{2}(c+dx)\right)\right)}{2d}$$

input `Integrate[(-5 + 3*Cos[c + d*x])^(-1), x]`

output `-1/2*ArcTan[2*Tan[(c + d*x)/2]]/d`

### 3.26.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3042, 3137}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{3 \cos(c + dx) - 5} dx$$

↓ 3042

$$\int \frac{1}{3 \sin(c + dx + \frac{\pi}{2}) - 5} dx$$

↓ 3137

$$-\frac{\arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{2d} - \frac{x}{4}$$

input `Int[(-5 + 3*Cos[c + d*x])^(-1),x]`

output `-1/4*x - ArcTan[Sin[c + d*x]/(3 - Cos[c + d*x])]/(2*d)`

#### 3.26.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3137 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[-x/q, x] - Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a - q + b*Sin[c + d*x]))], x]] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] & & NegQ[a]`

### 3.26.4 Maple [A] (verified)

Time = 0.59 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.55

method	result	size
derivativedivides	$-\frac{\arctan\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2d}$	18
default	$-\frac{\arctan\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2d}$	18
risch	$-\frac{i \ln(e^{i(dx+c)} - 3)}{4d} + \frac{i \ln(e^{i(dx+c)} - \frac{1}{3})}{4d}$	38
parallelrisch	$\frac{i\left(\ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - i\right) - \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + i\right)\right)}{4d}$	40

input `int(1/(-5+3*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `-1/2/d*arctan(2*tan(1/2*d*x+1/2*c))`

### 3.26.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

$$\int \frac{1}{-5 + 3 \cos(c + dx)} dx = \frac{\arctan\left(\frac{5 \cos(dx+c)-3}{4 \sin(dx+c)}\right)}{4d}$$

input `integrate(1/(-5+3*cos(d*x+c)),x, algorithm="fricas")`

output `1/4*arctan(1/4*(5*cos(d*x + c) - 3)/sin(d*x + c))/d`

### 3.26.6 Sympy [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.27

$$\int \frac{1}{-5 + 3 \cos(c + dx)} dx = \begin{cases} -\frac{\operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) + \pi \left\lfloor \frac{c + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor}{2d} & \text{for } d \neq 0 \\ \frac{x}{3 \cos(c) - 5} & \text{otherwise} \end{cases}$$

input `integrate(1/(-5+3*cos(d*x+c)),x)`

output `Piecewise((-atan(2*tan(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)/pi)) / (2*d), Ne(d, 0)), (x/(3*cos(c) - 5), True))`

### 3.26.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.73

$$\int \frac{1}{-5 + 3 \cos(c + dx)} dx = -\frac{\arctan\left(\frac{2 \sin(dx+c)}{\cos(dx+c)+1}\right)}{2d}$$

input `integrate(1/(-5+3*cos(d*x+c)),x, algorithm="maxima")`

output `-1/2*arctan(2*sin(d*x + c)/(cos(d*x + c) + 1))/d`

### 3.26.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.91

$$\int \frac{1}{-5 + 3 \cos(c + dx)} dx = -\frac{dx + c - 2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)-3}\right)}{4d}$$

input `integrate(1/(-5+3*cos(d*x+c)),x, algorithm="giac")`

output `-1/4*(d*x + c - 2*arctan(sin(d*x + c)/(cos(d*x + c) - 3)))/d`

**3.26.9 Mupad [B] (verification not implemented)**

Time = 14.44 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.15

$$\int \frac{1}{-5 + 3 \cos(c + dx)} dx = \frac{\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}}{2d} - \frac{\operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{2d}$$

input `int(1/(3*cos(c + d*x) - 5),x)`

output `(atan(tan(c/2 + (d*x)/2)) - (d*x)/2)/(2*d) - atan(2*tan(c/2 + (d*x)/2))/(2*d)`

### 3.27 $\int \frac{1}{(-5+3 \cos(c+dx))^2} dx$

3.27.1	Optimal result . . . . .	182
3.27.2	Mathematica [A] (verified) . . . . .	182
3.27.3	Rubi [A] (verified) . . . . .	183
3.27.4	Maple [A] (verified) . . . . .	184
3.27.5	Fricas [A] (verification not implemented) . . . . .	185
3.27.6	Sympy [C] (verification not implemented) . . . . .	185
3.27.7	Maxima [A] (verification not implemented) . . . . .	186
3.27.8	Giac [A] (verification not implemented) . . . . .	186
3.27.9	Mupad [B] (verification not implemented) . . . . .	187

#### 3.27.1 Optimal result

Integrand size = 12, antiderivative size = 58

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^2} dx = \frac{5x}{64} + \frac{5 \arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{32d} + \frac{3 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))}$$

output `5/64*x+5/32*arctan(sin(d*x+c)/(3-cos(d*x+c)))/d+3/16*sin(d*x+c)/d/(5-3*cos(d*x+c))`

#### 3.27.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.74

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^2} dx = \frac{5 \arctan\left(2 \tan\left(\frac{1}{2}(c + dx)\right)\right)}{32d} - \frac{6 \sin(c+dx)}{-5+3 \cos(c+dx)}$$

input `Integrate[(-5 + 3*Cos[c + d*x])^(-2),x]`

output `(5*ArcTan[2*Tan[(c + d*x)/2]] - (6*Sin[c + d*x])/(-5 + 3*Cos[c + d*x]))/(32*d)`

**3.27.3 Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {3042, 3143, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(3 \cos(c + dx) - 5)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(3 \sin(c + dx + \frac{\pi}{2}) - 5)^2} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{3 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))} - \frac{1}{16} \int -\frac{5}{5 - 3 \cos(c + dx)} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{5}{16} \int \frac{1}{5 - 3 \cos(c + dx)} dx + \frac{3 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{16} \int \frac{1}{5 - 3 \sin(c + dx + \frac{\pi}{2})} dx + \frac{3 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))} \\
 & \quad \downarrow \text{3136} \\
 & \frac{5}{16} \left( \frac{\arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{2d} + \frac{x}{4} \right) + \frac{3 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))}
 \end{aligned}$$

input `Int[(-5 + 3*Cos[c + d*x])^(-2), x]`

output `(5*(x/4 + ArcTan[Sin[c + d*x]/(3 - Cos[c + d*x])]/(2*d)))/16 + (3*Sin[c + d*x])/(16*d*(5 - 3*Cos[c + d*x]))`



### 3.27.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

rule 3143 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

### 3.27.4 Maple [A] (verified)

Time = 0.62 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.79

method	result	size
derivativedivides	$\frac{\frac{3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{64 \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{1}{4}\right)} + \frac{5 \arctan\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{32}}{d}$	46
default	$\frac{\frac{3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{64 \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{1}{4}\right)} + \frac{5 \arctan\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{32}}{d}$	46
parallelrisch	$\frac{(-15i \cos(dx+c) + 25i) \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - i\right) + (15i \cos(dx+c) - 25i) \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + i\right) - 12 \sin(dx+c)}{192d \cos(dx+c) - 320d}$	82
risch	$\frac{i(5e^{i(dx+c)} - 3)}{8d(3e^{2i(dx+c)} - 10e^{i(dx+c)} + 3)} + \frac{5i \ln(e^{i(dx+c)} - 3)}{64d} - \frac{5i \ln(e^{i(dx+c)} - \frac{1}{3})}{64d}$	83

input `int(1/(-5+3*cos(d*x+c))^2,x,method=_RETURNVERBOSE)`

3.27.  $\int \frac{1}{(-5+3 \cos(c+dx))^2} dx$

output `1/d*(3/64*tan(1/2*d*x+1/2*c)/(tan(1/2*d*x+1/2*c)^2+1/4)+5/32*arctan(2*tan(1/2*d*x+1/2*c)))`

### 3.27.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.02

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^2} dx$$

$$= -\frac{5(3 \cos(dx + c) - 5) \arctan\left(\frac{5 \cos(dx+c)-3}{4 \sin(dx+c)}\right) + 12 \sin(dx + c)}{64(3d \cos(dx + c) - 5d)}$$

input `integrate(1/(-5+3*cos(d*x+c))^2,x, algorithm="fricas")`

output `-1/64*(5*(3*cos(d*x + c) - 5)*arctan(1/4*(5*cos(d*x + c) - 3)/sin(d*x + c)) + 12*sin(d*x + c))/(3*d*cos(d*x + c) - 5*d)`

### 3.27.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.76 (sec) , antiderivative size = 192, normalized size of antiderivative = 3.31

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^2} dx$$

$$= \begin{cases} \frac{x}{(-5+3 \cosh(2 \operatorname{atanh}(\frac{1}{2})))^2} \\ \frac{x}{(3 \cos(c)-5)^2} \\ \frac{20 \left( \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right) \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{128d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 32d} + \frac{5 \left( \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{128d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 32d} + \frac{6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{128d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 32d} \end{cases}$$

input `integrate(1/(-5+3*cos(d*x+c))**2,x)`

```
output Piecewise((x/(-5 + 3*cosh(2*atanh(1/2)))**2, Eq(c, -d*x - 2*I*atanh(1/2))
| Eq(c, -d*x + 2*I*atanh(1/2))), (x/(3*cos(c) - 5)**2, Eq(d, 0)), (20*(ata
n(2*tan(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2
)**2/(128*d*tan(c/2 + d*x/2)**2 + 32*d) + 5*(atan(2*tan(c/2 + d*x/2)) + pi
*floor((c/2 + d*x/2 - pi/2)/pi))/(128*d*tan(c/2 + d*x/2)**2 + 32*d) + 6*ta
n(c/2 + d*x/2)/(128*d*tan(c/2 + d*x/2)**2 + 32*d), True))
```

### 3.27.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.19

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^2} dx = \frac{\frac{6 \sin(dx+c)}{\left(\frac{4 \sin(dx+c)^2}{(\cos(dx+c)+1)^2+1}\right)(\cos(dx+c)+1)} + 5 \arctan\left(\frac{2 \sin(dx+c)}{\cos(dx+c)+1}\right)}{32 d}$$

```
input integrate(1/(-5+3*cos(d*x+c))^2,x, algorithm="maxima")
```

```
output 1/32*(6*sin(d*x + c)/((4*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)*(cos(d*x
+ c) + 1)) + 5*arctan(2*sin(d*x + c)/(cos(d*x + c) + 1)))/d
```

### 3.27.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.05

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^2} dx = \frac{5 dx + 5 c + \frac{12 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{4 \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1} - 10 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)-3}\right)}{64 d}$$

```
input integrate(1/(-5+3*cos(d*x+c))^2,x, algorithm="giac")
```

```
output 1/64*(5*d*x + 5*c + 12*tan(1/2*d*x + 1/2*c)/(4*tan(1/2*d*x + 1/2*c)^2 + 1)
- 10*arctan(sin(d*x + c)/(cos(d*x + c) - 3)))/d
```

**3.27.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.16

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^2} dx = \frac{5 \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{32 d} - \frac{5 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{32 d} + \frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{64 d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + \frac{1}{4}\right)}$$

input `int(1/(3*cos(c + d*x) - 5)^2,x)`output `(5*atan(2*tan(c/2 + (d*x)/2)))/(32*d) - (5*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2))/(32*d) + (3*tan(c/2 + (d*x)/2))/(64*d*(tan(c/2 + (d*x)/2)^2 + 1/4))`

### 3.28 $\int \frac{1}{(-5+3 \cos(c+dx))^3} dx$

3.28.1	Optimal result . . . . .	188
3.28.2	Mathematica [A] (verified) . . . . .	188
3.28.3	Rubi [A] (verified) . . . . .	189
3.28.4	Maple [A] (verified) . . . . .	191
3.28.5	Fricas [A] (verification not implemented) . . . . .	191
3.28.6	Sympy [C] (verification not implemented) . . . . .	192
3.28.7	Maxima [A] (verification not implemented) . . . . .	192
3.28.8	Giac [A] (verification not implemented) . . . . .	193
3.28.9	Mupad [B] (verification not implemented) . . . . .	193

#### 3.28.1 Optimal result

Integrand size = 12, antiderivative size = 83

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^3} dx = -\frac{59x}{2048} - \frac{59 \arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{1024d} - \frac{3 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} - \frac{45 \sin(c + dx)}{512d(5 - 3 \cos(c + dx))}$$

output `-59/2048*x-59/1024*arctan(sin(d*x+c)/(3-cos(d*x+c)))/d-3/32*sin(d*x+c)/d/(5-3*cos(d*x+c))^2-45/512*sin(d*x+c)/d/(5-3*cos(d*x+c))`

#### 3.28.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.78

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^3} dx = \frac{-59 \arctan\left(2 \tan\left(\frac{1}{2}(c + dx)\right)\right) (5 - 3 \cos(c + dx))^2 - 546 \sin(c + dx) + 135 \sin(2(c + dx))}{1024d(5 - 3 \cos(c + dx))^2}$$

input `Integrate[(-5 + 3*Cos[c + d*x])^(-3),x]`

output `(-59*ArcTan[2*Tan[(c + d*x)/2]]*(5 - 3*Cos[c + d*x])^2 - 546*Sin[c + d*x] + 135*Sin[2*(c + d*x)])/(1024*d*(5 - 3*Cos[c + d*x])^2)`

**3.28.3 Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$ , Rules used = {3042, 3143, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(3 \cos(c + dx) - 5)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(3 \sin(c + dx + \frac{\pi}{2}) - 5)^3} dx \\
 & \quad \downarrow \text{3143} \\
 & -\frac{1}{32} \int \frac{3 \cos(c + dx) + 10}{(5 - 3 \cos(c + dx))^2} dx - \frac{3 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{32} \int \frac{3 \sin(c + dx + \frac{\pi}{2}) + 10}{(5 - 3 \sin(c + dx + \frac{\pi}{2}))^2} dx - \frac{3 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} \\
 & \quad \downarrow \text{3233} \\
 & \frac{1}{32} \left( \frac{1}{16} \int -\frac{59}{5 - 3 \cos(c + dx)} dx - \frac{45 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))} \right) - \frac{3 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{32} \left( -\frac{59}{16} \int \frac{1}{5 - 3 \cos(c + dx)} dx - \frac{45 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))} \right) - \frac{3 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{32} \left( -\frac{59}{16} \int \frac{1}{5 - 3 \sin(c + dx + \frac{\pi}{2})} dx - \frac{45 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))} \right) - \frac{3 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} \\
 & \quad \downarrow \text{3136} \\
 & \frac{1}{32} \left( -\frac{59}{16} \left( \frac{\arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{2d} + \frac{x}{4} \right) - \frac{45 \sin(c + dx)}{16d(5 - 3 \cos(c + dx))} \right) - \frac{3 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2}
 \end{aligned}$$

input `Int[(-5 + 3*Cos[c + d*x])^(-3), x]`

output  $(-3\sin[c + dx]) / (32d(5 - 3\cos[c + dx])^2) + ((-59(x/4 + \arctan[\sin[c + dx]/(3 - \cos[c + dx])]) / (2d)) / 16 - (45\sin[c + dx]) / (16d(5 - 3\cos[c + dx]))) / 32$

### 3.28.3.1 Defintions of rubi rules used

rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_) /; \text{FreeQ}[b, x]]$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3136  $\text{Int}[(a_*) + (b_*)\sin[(c_*) + (d_*)(x_)]^{(-1)}, x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a^2 - b^2, 2]\}, \text{Simp}[x/q, x] + \text{Simp}[(2/(d*q))*\text{ArcTan}[b*(\cos[c + dx])/(a + b*\sin[c + dx])]], x] /; \text{FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{GtQ}[a^2 - b^2, 0] \ \&\& \ \text{PosQ}[a]$

rule 3143  $\text{Int}[(a_*) + (b_*)\sin[(c_*) + (d_*)(x_)]^{(n_)}, x\_Symbol] \rightarrow \text{Simp}[(-b)*\cos[c + dx]*((a + b*\sin[c + dx])^{(n + 1)} / (d*(n + 1)*(a^2 - b^2))), x] + \text{Simp}[1/((n + 1)*(a^2 - b^2)) \text{Int}[(a + b*\sin[c + dx])^{(n + 1)}*\text{Simp}[a*(n + 1) - b*(n + 2)*\sin[c + dx], x], x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

rule 3233  $\text{Int}[(a_*) + (b_*)\sin[(e_*) + (f_*)(x_)]^{(m_)*((c_*) + (d_*)\sin[(e_*) + (f_*)(x_)]), x\_Symbol] \rightarrow \text{Simp}[(-b*c - a*d)*\cos[e + f*x]*((a + b*\sin[e + f*x])^{(m + 1)} / (f*(m + 1)*(a^2 - b^2))), x] + \text{Simp}[1/((m + 1)*(a^2 - b^2)) \text{Int}[(a + b*\sin[e + f*x])^{(m + 1)}*\text{Simp}[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*\sin[e + f*x], x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntegerQ}[2*m]$

### 3.28.4 Maple [A] (verified)

Time = 0.79 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.77

method	result
derivativedivides	$\frac{4 \left( \frac{51 \tan^3 \left( \frac{dx}{2} + \frac{c}{2} \right)}{512} + \frac{69 \tan \left( \frac{dx}{2} + \frac{c}{2} \right)}{2048} \right) - \frac{59 \arctan \left( 2 \tan \left( \frac{dx}{2} + \frac{c}{2} \right) \right)}{1024}}{\left( 4 \tan^2 \left( \frac{dx}{2} + \frac{c}{2} \right) + 1 \right)^2} \frac{1}{d}$
default	$\frac{4 \left( \frac{51 \tan^3 \left( \frac{dx}{2} + \frac{c}{2} \right)}{512} + \frac{69 \tan \left( \frac{dx}{2} + \frac{c}{2} \right)}{2048} \right) - \frac{59 \arctan \left( 2 \tan \left( \frac{dx}{2} + \frac{c}{2} \right) \right)}{1024}}{\left( 4 \tan^2 \left( \frac{dx}{2} + \frac{c}{2} \right) + 1 \right)^2} \frac{1}{d}$
risch	$-\frac{3i(59e^{3i(dx+c)} - 295e^{2i(dx+c)} + 241e^{i(dx+c)} - 45)}{256d(3e^{2i(dx+c)} - 10e^{i(dx+c)} + 3)^2} + \frac{59i \ln(e^{i(dx+c)} - \frac{1}{3})}{2048d} - \frac{59i \ln(e^{i(dx+c)} - 3)}{2048d}$
parallelrisch	$\frac{59i(-9 \cos(2dx+2c) - 59 + 60 \cos(dx+c)) \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - i\right) + 59i(59 + 9 \cos(2dx+2c) - 60 \cos(dx+c)) \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + i\right)}{2048d(-9 \cos(2dx+2c) - 59 + 60 \cos(dx+c))}$

input `int(1/(-5+3*cos(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/d*(-4*(51/512*tan(1/2*d*x+1/2*c)^3+69/2048*tan(1/2*d*x+1/2*c))/(4*tan(1/2*d*x+1/2*c)^2+1)^2-59/1024*arctan(2*tan(1/2*d*x+1/2*c)))`

### 3.28.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.08

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^3} dx$$

$$= \frac{59(9 \cos(dx + c)^2 - 30 \cos(dx + c) + 25) \arctan\left(\frac{5 \cos(dx + c) - 3}{4 \sin(dx + c)}\right) + 12(45 \cos(dx + c) - 91) \sin(dx + c)}{2048(9d \cos(dx + c)^2 - 30d \cos(dx + c) + 25d)}$$

input `integrate(1/(-5+3*cos(d*x+c))^3,x, algorithm="fricas")`

output `1/2048*(59*(9*cos(d*x + c)^2 - 30*cos(d*x + c) + 25)*arctan(1/4*(5*cos(d*x + c) - 3)/sin(d*x + c)) + 12*(45*cos(d*x + c) - 91)*sin(d*x + c))/(9*d*cos(d*x + c)^2 - 30*d*cos(d*x + c) + 25*d)`



### 3.28.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.34 (sec) , antiderivative size = 366, normalized size of antiderivative = 4.41

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^3} dx$$

$$= \begin{cases} \frac{x}{(-5 + 3 \cosh(2 \operatorname{atanh}(\frac{1}{2})))^3} \\ \frac{x}{(3 \cos(c) - 5)^3} \\ -\frac{944 \left( \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right) \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right)}{16384d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) + 8192d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 1024d} - \frac{472 \left( \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right) \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{16384d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) + 8192d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 1024d} - \frac{59 \left( \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{16384d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) + 8192d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 1024d} \end{cases}$$

input `integrate(1/(-5+3*cos(d*x+c))**3,x)`

output `Piecewise((x/(-5 + 3*cosh(2*atanh(1/2)))**3, Eq(c, -d*x - 2*I*atanh(1/2)) | Eq(c, -d*x + 2*I*atanh(1/2))), (x/(3*cos(c) - 5)**3, Eq(d, 0)), (-944*(atan(2*tan(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4/(16384*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 1024*d) - 472*(atan(2*tan(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**2/(16384*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 1024*d) - 59*(atan(2*tan(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(16384*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 1024*d) - 408*tan(c/2 + d*x/2)**3/(16384*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 1024*d) - 138*tan(c/2 + d*x/2)/(16384*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 1024*d), True))`

### 3.28.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.35

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^3} dx = -\frac{6 \left( \frac{23 \sin(dx+c)}{\cos(dx+c)+1} + \frac{68 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{8 \sin(dx+c)^2 + \frac{16 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 1} + 59 \arctan\left(\frac{2 \sin(dx+c)}{\cos(dx+c)+1}\right) \frac{1}{1024 d}$$

input `integrate(1/(-5+3*cos(d*x+c))^3,x, algorithm="maxima")`

output 
$$\frac{-1/1024*(6*(23*\sin(d*x + c)/(\cos(d*x + c) + 1) + 68*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3)/(8*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2 + 16*\sin(d*x + c)^4/(\cos(d*x + c) + 1)^4 + 1) + 59*\arctan(2*\sin(d*x + c)/(\cos(d*x + c) + 1)))/d$$

### 3.28.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.93

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^3} dx$$

$$= \frac{59 dx + 59 c + \frac{12 \left( 68 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 23 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\left( 4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 1 \right)^2} - 118 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)-3}\right)}{2048 d}$$

input `integrate(1/(-5+3*cos(d*x+c))^3,x, algorithm="giac")`

output 
$$\frac{-1/2048*(59*d*x + 59*c + 12*(68*\tan(1/2*d*x + 1/2*c)^3 + 23*\tan(1/2*d*x + 1/2*c)))/(4*\tan(1/2*d*x + 1/2*c)^2 + 1)^2 - 118*\arctan(\sin(d*x + c)/(\cos(d*x + c) - 3)))/d$$

### 3.28.9 Mupad [B] (verification not implemented)

Time = 14.40 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.16

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^3} dx = \frac{59 \left( \operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2} \right)}{1024 d} - \frac{59 \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{1024 d}$$

$$- \frac{\frac{51 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{2048} + \frac{69 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{8192}}{d \left( \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{2} + \frac{1}{16} \right)}$$

input `int(1/(3*cos(c + d*x) - 5)^3,x)`

output 
$$\frac{(59*(\operatorname{atan}(\tan(c/2 + (d*x)/2)) - (d*x)/2))/(1024*d) - (59*\operatorname{atan}(2*\tan(c/2 + (d*x)/2)))/(1024*d) - ((69*\tan(c/2 + (d*x)/2))/8192 + (51*\tan(c/2 + (d*x)/2)^3)/2048)/(d*(\tan(c/2 + (d*x)/2)^2/2 + \tan(c/2 + (d*x)/2)^4 + 1/16))$$

---

3.28. 
$$\int \frac{1}{(-5+3 \cos(c+dx))^3} dx$$

### 3.29 $\int \frac{1}{(-5+3 \cos(c+dx))^4} dx$

3.29.1	Optimal result . . . . .	194
3.29.2	Mathematica [A] (verified) . . . . .	194
3.29.3	Rubi [A] (verified) . . . . .	195
3.29.4	Maple [A] (verified) . . . . .	197
3.29.5	Fricas [A] (verification not implemented) . . . . .	198
3.29.6	Sympy [C] (verification not implemented) . . . . .	198
3.29.7	Maxima [A] (verification not implemented) . . . . .	199
3.29.8	Giac [A] (verification not implemented) . . . . .	200
3.29.9	Mupad [B] (verification not implemented) . . . . .	200

#### 3.29.1 Optimal result

Integrand size = 12, antiderivative size = 108

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^4} dx = \frac{385x}{32768} + \frac{385 \arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{16384d} + \frac{\sin(c + dx)}{16d(5 - 3 \cos(c + dx))^3} + \frac{25 \sin(c + dx)}{512d(5 - 3 \cos(c + dx))^2} + \frac{311 \sin(c + dx)}{8192d(5 - 3 \cos(c + dx))}$$

output `385/32768*x+385/16384*arctan(sin(d*x+c)/(3-cos(d*x+c)))/d+1/16*sin(d*x+c)/d/(5-3*cos(d*x+c))^3+25/512*sin(d*x+c)/d/(5-3*cos(d*x+c))^2+311/8192*sin(d*x+c)/d/(5-3*cos(d*x+c))`

#### 3.29.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.61

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^4} dx = \frac{770 \arctan\left(2 \tan\left(\frac{1}{2}(c + dx)\right)\right) - \frac{9(4883 \sin(c+dx) - 2340 \sin(2(c+dx)) + 311 \sin(3(c+dx)))}{(-5+3 \cos(c+dx))^3}}{32768d}$$

input `Integrate[(-5 + 3*Cos[c + d*x])^(-4), x]`

output `(770*ArcTan[2*Tan[(c + d*x)/2]] - (9*(4883*Sin[c + d*x] - 2340*Sin[2*(c + d*x)] + 311*Sin[3*(c + d*x)]))/(-5 + 3*Cos[c + d*x])^3/(32768*d)`

**3.29.3 Rubi [A] (verified)**

Time = 0.49 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.14, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$ , Rules used = {3042, 3143, 27, 3042, 3233, 25, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(3 \cos(c + dx) - 5)^4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(3 \sin(c + dx + \frac{\pi}{2}) - 5)^4} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{\sin(c + dx)}{16d(5 - 3 \cos(c + dx))^3} - \frac{1}{48} \int -\frac{3(2 \cos(c + dx) + 5)}{(5 - 3 \cos(c + dx))^3} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{16} \int \frac{2 \cos(c + dx) + 5}{(5 - 3 \cos(c + dx))^3} dx + \frac{\sin(c + dx)}{16d(5 - 3 \cos(c + dx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{16} \int \frac{2 \sin(c + dx + \frac{\pi}{2}) + 5}{(5 - 3 \sin(c + dx + \frac{\pi}{2}))^3} dx + \frac{\sin(c + dx)}{16d(5 - 3 \cos(c + dx))^3} \\
 & \quad \downarrow \text{3233} \\
 & \frac{1}{16} \left( \frac{25 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} - \frac{1}{32} \int -\frac{25 \cos(c + dx) + 62}{(5 - 3 \cos(c + dx))^2} dx \right) + \frac{\sin(c + dx)}{16d(5 - 3 \cos(c + dx))^3} \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{16} \left( \frac{1}{32} \int \frac{25 \cos(c + dx) + 62}{(5 - 3 \cos(c + dx))^2} dx + \frac{25 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} \right) + \frac{\sin(c + dx)}{16d(5 - 3 \cos(c + dx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{16} \left( \frac{1}{32} \int \frac{25 \sin(c + dx + \frac{\pi}{2}) + 62}{(5 - 3 \sin(c + dx + \frac{\pi}{2}))^2} dx + \frac{25 \sin(c + dx)}{32d(5 - 3 \cos(c + dx))^2} \right) + \frac{\sin(c + dx)}{16d(5 - 3 \cos(c + dx))^3} \\
 & \quad \downarrow \text{3233}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{16} \left( \frac{1}{32} \left( \frac{311 \sin(c+dx)}{16d(5-3\cos(c+dx))} - \frac{1}{16} \int \frac{385}{5-3\cos(c+dx)} dx \right) + \frac{25 \sin(c+dx)}{32d(5-3\cos(c+dx))^2} \right) + \\
& \quad \frac{\sin(c+dx)}{16d(5-3\cos(c+dx))^3} \\
& \quad \downarrow \text{27} \\
& \frac{1}{16} \left( \frac{1}{32} \left( \frac{385}{16} \int \frac{1}{5-3\cos(c+dx)} dx + \frac{311 \sin(c+dx)}{16d(5-3\cos(c+dx))} \right) + \frac{25 \sin(c+dx)}{32d(5-3\cos(c+dx))^2} \right) + \\
& \quad \frac{\sin(c+dx)}{16d(5-3\cos(c+dx))^3} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{16} \left( \frac{1}{32} \left( \frac{385}{16} \int \frac{1}{5-3\sin(c+dx+\frac{\pi}{2})} dx + \frac{311 \sin(c+dx)}{16d(5-3\cos(c+dx))} \right) + \frac{25 \sin(c+dx)}{32d(5-3\cos(c+dx))^2} \right) + \\
& \quad \frac{\sin(c+dx)}{16d(5-3\cos(c+dx))^3} \\
& \quad \downarrow \text{3136} \\
& \frac{1}{16} \left( \frac{1}{32} \left( \frac{385}{16} \left( \frac{\arctan\left(\frac{\sin(c+dx)}{3-\cos(c+dx)}\right)}{2d} + \frac{x}{4} \right) + \frac{311 \sin(c+dx)}{16d(5-3\cos(c+dx))} \right) + \frac{25 \sin(c+dx)}{32d(5-3\cos(c+dx))^2} \right) + \\
& \quad \frac{\sin(c+dx)}{16d(5-3\cos(c+dx))^3}
\end{aligned}$$

input `Int[(-5 + 3*Cos[c + d*x])^(-4), x]`

output `Sin[c + d*x]/(16*d*(5 - 3*Cos[c + d*x])^3) + ((25*Sin[c + d*x])/(32*d*(5 - 3*Cos[c + d*x])^2) + ((385*(x/4 + ArcTan[Sin[c + d*x]/(3 - Cos[c + d*x]))/(2*d)))/16 + (311*Sin[c + d*x])/(16*d*(5 - 3*Cos[c + d*x])))/32)/16`

### 3.29.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + b*Sin[c + d*x]))], x]] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3233 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

### 3.29.4 Maple [A] (verified)

Time = 0.67 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.71

method	result
derivativedivides	$\frac{\frac{369 \tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{512} + \frac{117 \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{256} + \frac{639 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8192} + \frac{385 \arctan\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{16384}}{\left(4 \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^3} \frac{d}{d}$
default	$\frac{\frac{369 \tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{512} + \frac{117 \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{256} + \frac{639 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8192} + \frac{385 \arctan\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{16384}}{\left(4 \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^3} \frac{d}{d}$
risch	$\frac{i(10395 e^{5i(dx+c)} - 86625 e^{4i(dx+c)} + 239470 e^{3i(dx+c)} - 218466 e^{2i(dx+c)} + 73575 e^{i(dx+c)} - 8397)}{12288d(3 e^{2i(dx+c)} - 10 e^{i(dx+c)} + 3)^3} - \frac{385i \ln(e^{i(dx+c)} - i)}{32768d}$
parallelrisc	$\frac{385i(770 - 27 \cos(3dx+3c) - 981 \cos(dx+c) + 270 \cos(2dx+2c)) \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - i\right) + 385i(27 \cos(3dx+3c) + 981 \cos(dx+c))}{32768d(27 \cos(3dx+3c) + 981 \cos(dx+c))}$

input `int(1/(-5+3*cos(d*x+c))^4,x,method=_RETURNVERBOSE)`

$$3.29. \int \frac{1}{(-5+3 \cos(c+dx))^4} dx$$

output  $1/d*(8*(369/4096*\tan(1/2*d*x+1/2*c)^5+117/2048*\tan(1/2*d*x+1/2*c)^3+639/65536*\tan(1/2*d*x+1/2*c))/(4*\tan(1/2*d*x+1/2*c)^2+1)^3+385/16384*\arctan(2*\tan(1/2*d*x+1/2*c))$

### 3.29.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.12

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^4} dx = \frac{385 (27 \cos(dx + c)^3 - 135 \cos(dx + c)^2 + 225 \cos(dx + c) - 125) \arctan\left(\frac{5 \cos(dx+c)-3}{4 \sin(dx+c)}\right) + 36 (311 \cos(dx + c)^2 - 1170 \cos(dx + c) + 1143) \sin(dx + c)}{32768 (27 d \cos(dx + c)^3 - 135 d \cos(dx + c)^2 + 225 d \cos(dx + c) - 125 d)}$$

input `integrate(1/(-5+3*cos(d*x+c))^4,x, algorithm="fricas")`

output  $-1/32768*(385*(27*\cos(d*x + c)^3 - 135*\cos(d*x + c)^2 + 225*\cos(d*x + c) - 125)*\arctan(1/4*(5*\cos(d*x + c) - 3)/\sin(d*x + c)) + 36*(311*\cos(d*x + c)^2 - 1170*\cos(d*x + c) + 1143)*\sin(d*x + c))/(27*d*\cos(d*x + c)^3 - 135*d*\cos(d*x + c)^2 + 225*d*\cos(d*x + c) - 125*d)$

### 3.29.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.98 (sec) , antiderivative size = 597, normalized size of antiderivative = 5.53

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^4} dx = \begin{cases} \frac{x}{(-5+3 \cosh(2 \operatorname{atanh}(\frac{1}{2})))^4} \\ \frac{x}{(3 \cos(c)-5)^4} \\ \frac{24640 \left( \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right) \tan^6\left(\frac{c}{2} + \frac{dx}{2}\right)}{1048576d \tan^6\left(\frac{c}{2} + \frac{dx}{2}\right) + 786432d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) + 196608d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 16384d} + \frac{18480 \left( \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right) \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right)}{1048576d \tan^6\left(\frac{c}{2} + \frac{dx}{2}\right) + 786432d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) + 196608d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 16384d} \end{cases}$$

input `integrate(1/(-5+3*cos(d*x+c))**4,x)`

---

3.29.  $\int \frac{1}{(-5+3 \cos(c+dx))^4} dx$

```

output Piecewise((x/(-5 + 3*cosh(2*atanh(1/2)))**4, Eq(c, -d*x - 2*I*atanh(1/2))
| Eq(c, -d*x + 2*I*atanh(1/2))), (x/(3*cos(c) - 5)**4, Eq(d, 0)), (24640*(
atan(2*tan(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*
x/2)**6/(1048576*d*tan(c/2 + d*x/2)**6 + 786432*d*tan(c/2 + d*x/2)**4 + 19
6608*d*tan(c/2 + d*x/2)**2 + 16384*d) + 18480*(atan(2*tan(c/2 + d*x/2)) +
pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4/(1048576*d*tan(c/2
+ d*x/2)**6 + 786432*d*tan(c/2 + d*x/2)**4 + 196608*d*tan(c/2 + d*x/2)**2
+ 16384*d) + 4620*(atan(2*tan(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2
)/pi))*tan(c/2 + d*x/2)**2/(1048576*d*tan(c/2 + d*x/2)**6 + 786432*d*tan(c
/2 + d*x/2)**4 + 196608*d*tan(c/2 + d*x/2)**2 + 16384*d) + 385*(atan(2*tan
(c/2 + d*x/2)) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(1048576*d*tan(c/2 + d
*x/2)**6 + 786432*d*tan(c/2 + d*x/2)**4 + 196608*d*tan(c/2 + d*x/2)**2 + 1
6384*d) + 11808*tan(c/2 + d*x/2)**5/(1048576*d*tan(c/2 + d*x/2)**6 + 78643
2*d*tan(c/2 + d*x/2)**4 + 196608*d*tan(c/2 + d*x/2)**2 + 16384*d) + 7488*t
an(c/2 + d*x/2)**3/(1048576*d*tan(c/2 + d*x/2)**6 + 786432*d*tan(c/2 + d*x
/2)**4 + 196608*d*tan(c/2 + d*x/2)**2 + 16384*d) + 1278*tan(c/2 + d*x/2)/(
1048576*d*tan(c/2 + d*x/2)**6 + 786432*d*tan(c/2 + d*x/2)**4 + 196608*d*ta
n(c/2 + d*x/2)**2 + 16384*d), True))

```

### 3.29.7 Maxima [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.41

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^4} dx$$

$$= \frac{18 \left( \frac{71 \sin(dx+c)}{\cos(dx+c)+1} + \frac{416 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{656 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right) + 385 \arctan \left( \frac{2 \sin(dx+c)}{\cos(dx+c)+1} \right)}{16384 d}$$

```

input integrate(1/(-5+3*cos(d*x+c))^4,x, algorithm="maxima")

```

```

output 1/16384*(18*(71*sin(d*x + c)/(cos(d*x + c) + 1) + 416*sin(d*x + c)^3/(cos(
d*x + c) + 1)^3 + 656*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/(12*sin(d*x + c
)^2/(cos(d*x + c) + 1)^2 + 48*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 64*sin
(d*x + c)^6/(cos(d*x + c) + 1)^6 + 1) + 385*arctan(2*sin(d*x + c)/(cos(d*x
+ c) + 1)))/d

```



**3.29.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.83

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^4} dx$$

$$= \frac{385 dx + 385 c + \frac{36 \left( 656 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 + 416 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 71 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\left( 4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 1 \right)^3} - 770 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)-3}\right)}{32768 d}$$

input `integrate(1/(-5+3*cos(d*x+c))^4,x, algorithm="giac")`output `1/32768*(385*d*x + 385*c + 36*(656*tan(1/2*d*x + 1/2*c)^5 + 416*tan(1/2*d*x + 1/2*c)^3 + 71*tan(1/2*d*x + 1/2*c))/(4*tan(1/2*d*x + 1/2*c)^2 + 1)^3 - 770*arctan(sin(d*x + c)/(cos(d*x + c) - 3))/d`**3.29.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.90

$$\int \frac{1}{(-5 + 3 \cos(c + dx))^4} dx = \frac{385 \operatorname{atan}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{16384 d} - \frac{385 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{16384 d}$$

$$+ \frac{\frac{369 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{512} + \frac{117 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{256} + \frac{639 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{8192}}{d \left(4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1\right)^3}$$

input `int(1/(3*cos(c + d*x) - 5)^4,x)`output `(385*atan(2*tan(c/2 + (d*x)/2)))/(16384*d) - (385*(atan(tan(c/2 + (d*x)/2) - (d*x)/2))/(16384*d) + ((639*tan(c/2 + (d*x)/2))/8192 + (117*tan(c/2 + (d*x)/2)^3)/256 + (369*tan(c/2 + (d*x)/2)^5)/512)/(d*(4*tan(c/2 + (d*x)/2)^2 + 1)^3)`

### 3.30 $\int \frac{1}{-5-3\cos(c+dx)} dx$

3.30.1 Optimal result . . . . .	201
3.30.2 Mathematica [A] (verified) . . . . .	201
3.30.3 Rubi [A] (verified) . . . . .	202
3.30.4 Maple [A] (verified) . . . . .	203
3.30.5 Fricas [A] (verification not implemented) . . . . .	203
3.30.6 Sympy [A] (verification not implemented) . . . . .	203
3.30.7 Maxima [A] (verification not implemented) . . . . .	204
3.30.8 Giac [A] (verification not implemented) . . . . .	204
3.30.9 Mupad [B] (verification not implemented) . . . . .	205

#### 3.30.1 Optimal result

Integrand size = 12, antiderivative size = 31

$$\int \frac{1}{-5-3\cos(c+dx)} dx = -\frac{x}{4} + \frac{\arctan\left(\frac{\sin(c+dx)}{3+\cos(c+dx)}\right)}{2d}$$

output `-1/4*x+1/2*arctan(sin(d*x+c)/(3+cos(d*x+c)))/d`

#### 3.30.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.65

$$\int \frac{1}{-5-3\cos(c+dx)} dx = \frac{\arctan\left(2\cot\left(\frac{1}{2}(c+dx)\right)\right)}{2d}$$

input `Integrate[(-5 - 3*Cos[c + d*x])^(-1),x]`

output `ArcTan[2*Cot[(c + d*x)/2]]/(2*d)`

### 3.30.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {3042, 3137}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{-3 \cos(c + dx) - 5} dx$$

↓ 3042

$$\int \frac{1}{-3 \sin\left(c + dx + \frac{\pi}{2}\right) - 5} dx$$

↓ 3137

$$\frac{\arctan\left(\frac{\sin(c+dx)}{\cos(c+dx)+3}\right)}{2d} - \frac{x}{4}$$

input `Int[(-5 - 3*Cos[c + d*x])^(-1),x]`

output `-1/4*x + ArcTan[Sin[c + d*x]/(3 + Cos[c + d*x])]/(2*d)`

#### 3.30.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3137 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[-x/q, x] - Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a - q + b*Sin[c + d*x]))], x]] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] & & NegQ[a]`

**3.30.4 Maple [A] (verified)**

Time = 0.56 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.58

method	result	size
derivativedivides	$-\frac{\arctan\left(\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2}\right)}{2d}$	18
default	$-\frac{\arctan\left(\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2}\right)}{2d}$	18
parallelrisch	$\frac{i\left(\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2i\right) - \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2i\right)\right)}{4d}$	36
risch	$-\frac{i \ln(e^{i(dx+c)}+3)}{4d} + \frac{i \ln(e^{i(dx+c)}+\frac{1}{3})}{4d}$	38

input `int(1/(-5-3*cos(d*x+c)),x,method=_RETURNVERBOSE)`output `-1/2/d*arctan(1/2*tan(1/2*d*x+1/2*c))`**3.30.5 Fracas [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{1}{-5 - 3 \cos(c + dx)} dx = \frac{\arctan\left(\frac{5 \cos(dx+c)+3}{4 \sin(dx+c)}\right)}{4d}$$

input `integrate(1/(-5-3*cos(d*x+c)),x, algorithm="fricas")`output `1/4*arctan(1/4*(5*cos(d*x + c) + 3)/sin(d*x + c))/d`**3.30.6 Sympy [A] (verification not implemented)**

Time = 0.37 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.42

$$\int \frac{1}{-5 - 3 \cos(c + dx)} dx = \begin{cases} \frac{\operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right) + \pi \left\lfloor \frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2} \right\rfloor}{2d} & \text{for } d \neq 0 \\ \frac{x}{-3 \cos(c) - 5} & \text{otherwise} \end{cases}$$

input `integrate(1/(-5-3*cos(d*x+c)),x)`

output `Piecewise((-atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi)) / (2*d), Ne(d, 0)), (x/(-3*cos(c) - 5), True))`

### 3.30.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.77

$$\int \frac{1}{-5 - 3 \cos(c + dx)} dx = -\frac{\arctan\left(\frac{\sin(dx+c)}{2(\cos(dx+c)+1)}\right)}{2d}$$

input `integrate(1/(-5-3*cos(d*x+c)),x, algorithm="maxima")`

output `-1/2*arctan(1/2*sin(d*x + c)/(cos(d*x + c) + 1))/d`

### 3.30.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.97

$$\int \frac{1}{-5 - 3 \cos(c + dx)} dx = -\frac{dx + c - 2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+3}\right)}{4d}$$

input `integrate(1/(-5-3*cos(d*x+c)),x, algorithm="giac")`

output `-1/4*(d*x + c - 2*arctan(sin(d*x + c)/(cos(d*x + c) + 3)))/d`

**3.30.9 Mupad [B] (verification not implemented)**

Time = 14.51 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.23

$$\int \frac{1}{-5 - 3 \cos(c + dx)} dx = \frac{\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}}{2d} - \frac{\operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{2d}$$

input `int(-1/(3*cos(c + d*x) + 5),x)`

output `(atan(tan(c/2 + (d*x)/2)) - (d*x)/2)/(2*d) - atan(tan(c/2 + (d*x)/2)/2)/(2*d)`

### 3.31 $\int \frac{1}{(-5-3\cos(c+dx))^2} dx$

3.31.1	Optimal result . . . . .	206
3.31.2	Mathematica [A] (verified) . . . . .	206
3.31.3	Rubi [A] (verified) . . . . .	207
3.31.4	Maple [A] (verified) . . . . .	208
3.31.5	Fricas [A] (verification not implemented) . . . . .	209
3.31.6	Sympy [C] (verification not implemented) . . . . .	209
3.31.7	Maxima [A] (verification not implemented) . . . . .	210
3.31.8	Giac [A] (verification not implemented) . . . . .	210
3.31.9	Mupad [B] (verification not implemented) . . . . .	211

#### 3.31.1 Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \frac{1}{(-5-3\cos(c+dx))^2} dx = \frac{5x}{64} - \frac{5 \arctan\left(\frac{\sin(c+dx)}{3+\cos(c+dx)}\right)}{32d} - \frac{3 \sin(c+dx)}{16d(5+3\cos(c+dx))}$$

output `5/64*x-5/32*arctan(sin(d*x+c)/(3+cos(d*x+c)))/d-3/16*sin(d*x+c)/d/(5+3*cos(d*x+c))`

#### 3.31.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.77

$$\int \frac{1}{(-5-3\cos(c+dx))^2} dx = -\frac{5 \arctan\left(2 \cot\left(\frac{1}{2}(c+dx)\right)\right) + \frac{6 \sin(c+dx)}{5+3\cos(c+dx)}}{32d}$$

input `Integrate[(-5 - 3*Cos[c + d*x])^(-2), x]`

output `-1/32*(5*ArcTan[2*Cot[(c + d*x)/2]] + (6*Sin[c + d*x])/(5 + 3*Cos[c + d*x]))/d`

**3.31.3 Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {3042, 3143, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(-3 \cos(c + dx) - 5)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(-3 \sin(c + dx + \frac{\pi}{2}) - 5)^2} dx \\
 & \quad \downarrow \text{3143} \\
 & -\frac{1}{16} \int -\frac{5}{3 \cos(c + dx) + 5} dx - \frac{3 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)} \\
 & \quad \downarrow \text{27} \\
 & \frac{5}{16} \int \frac{1}{3 \cos(c + dx) + 5} dx - \frac{3 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{16} \int \frac{1}{3 \sin(c + dx + \frac{\pi}{2}) + 5} dx - \frac{3 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)} \\
 & \quad \downarrow \text{3136} \\
 & \frac{5}{16} \left( \frac{x}{4} - \frac{\arctan\left(\frac{\sin(c+dx)}{\cos(c+dx)+3}\right)}{2d} \right) - \frac{3 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)}
 \end{aligned}$$

input `Int[(-5 - 3*Cos[c + d*x])^(-2),x]`

output `(5*(x/4 - ArcTan[Sin[c + d*x]/(3 + Cos[c + d*x])]/(2*d)))/16 - (3*Sin[c + d*x])/(16*d*(5 + 3*Cos[c + d*x]))`



3.31.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

3.31.4 Maple [A] (verified)

Time = 0.71 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$\frac{-\frac{3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{16\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\right)} + \frac{5 \arctan\left(\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2}\right)}{32}}{d}$	46
default	$\frac{-\frac{3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{16\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\right)} + \frac{5 \arctan\left(\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2}\right)}{32}}{d}$	46
parallelrisch	$\frac{(-15i \cos(dx+c) - 25i) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2i\right) + (15i \cos(dx+c) + 25i) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2i\right) - 12 \sin(dx+c)}{192d \cos(dx+c) + 320d}$	78
risch	$-\frac{i(5e^{i(dx+c)} + 3)}{8d(3e^{2i(dx+c)} + 10e^{i(dx+c)} + 3)} - \frac{5i \ln\left(e^{i(dx+c)} + \frac{1}{3}\right)}{64d} + \frac{5i \ln\left(e^{i(dx+c)} + 3\right)}{64d}$	83

input `int(1/(-5-3*cos(d*x+c))^2,x,method=_RETURNVERBOSE)`

3.31.  $\int \frac{1}{(-5-3\cos(c+dx))^2} dx$

output `1/d*(-3/16*tan(1/2*d*x+1/2*c)/(tan(1/2*d*x+1/2*c)^2+4)+5/32*arctan(1/2*tan(1/2*d*x+1/2*c)))`

### 3.31.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.05

$$\int \frac{1}{(-5 - 3 \cos(c + dx))^2} dx$$

$$= -\frac{5(3 \cos(dx + c) + 5) \arctan\left(\frac{5 \cos(dx+c)+3}{4 \sin(dx+c)}\right) + 12 \sin(dx + c)}{64(3d \cos(dx + c) + 5d)}$$

input `integrate(1/(-5-3*cos(d*x+c))^2,x, algorithm="fracas")`

output `-1/64*(5*(3*cos(d*x + c) + 5)*arctan(1/4*(5*cos(d*x + c) + 3)/sin(d*x + c)) + 12*sin(d*x + c))/(3*d*cos(d*x + c) + 5*d)`

### 3.31.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.94 (sec) , antiderivative size = 190, normalized size of antiderivative = 3.39

$$\int \frac{1}{(-5 - 3 \cos(c + dx))^2} dx$$

$$= \begin{cases} \frac{x}{(-5-3 \cosh(2 \operatorname{atanh}(2)))^2} \\ \frac{x}{(-3 \cos(c)-5)^2} \\ \frac{5 \left( \operatorname{atan} \left( \frac{\tan \left( \frac{c}{2} + \frac{dx}{2} \right)}{2} \right) + \pi \left[ \frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2} \right] \right) \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right)}{32d \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right) + 128d} + \frac{20 \left( \operatorname{atan} \left( \frac{\tan \left( \frac{c}{2} + \frac{dx}{2} \right)}{2} \right) + \pi \left[ \frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2} \right] \right)}{32d \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right) + 128d} - \frac{6 \tan \left( \frac{c}{2} + \frac{dx}{2} \right)}{32d \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right) + 128d} \end{cases}$$

input `integrate(1/(-5-3*cos(d*x+c))**2,x)`

output `Piecewise((x/(-5 - 3*cosh(2*atanh(2)))**2, Eq(c, -d*x - 2*I*atanh(2)) | Eq(c, -d*x + 2*I*atanh(2))), (x/(-3*cos(c) - 5)**2, Eq(d, 0)), (5*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**2/(32*d*tan(c/2 + d*x/2)**2 + 128*d) + 20*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(32*d*tan(c/2 + d*x/2)**2 + 128*d) - 6*tan(c/2 + d*x/2)/(32*d*tan(c/2 + d*x/2)**2 + 128*d), True))`

### 3.31.7 Maxima [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.21

$$\int \frac{1}{(-5 - 3 \cos(c + dx))^2} dx = -\frac{\frac{6 \sin(dx+c)}{\left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2+4}\right)(\cos(dx+c)+1)} - 5 \arctan\left(\frac{\sin(dx+c)}{2(\cos(dx+c)+1)}\right)}{32 d}$$

input `integrate(1/(-5-3*cos(d*x+c))^2,x, algorithm="maxima")`

output `-1/32*(6*sin(d*x + c)/((sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 4)*(cos(d*x + c) + 1)) - 5*arctan(1/2*sin(d*x + c)/(cos(d*x + c) + 1)))/d`

### 3.31.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.05

$$\int \frac{1}{(-5 - 3 \cos(c + dx))^2} dx = \frac{5 dx + 5 c - \frac{12 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 4} - 10 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+3}\right)}{64 d}$$

input `integrate(1/(-5-3*cos(d*x+c))^2,x, algorithm="giac")`

output `1/64*(5*d*x + 5*c - 12*tan(1/2*d*x + 1/2*c)/(tan(1/2*d*x + 1/2*c)^2 + 4) - 10*arctan(sin(d*x + c)/(cos(d*x + c) + 3)))/d`

**3.31.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.20

$$\int \frac{1}{(-5 - 3 \cos(c + dx))^2} dx = \frac{5 \operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{32 d} - \frac{5 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{32 d} - \frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{16 d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 4\right)}$$

input `int(1/(3*cos(c + d*x) + 5)^2,x)`output `(5*atan(tan(c/2 + (d*x)/2)/2))/(32*d) - (5*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2))/(32*d) - (3*tan(c/2 + (d*x)/2))/(16*d*(tan(c/2 + (d*x)/2)^2 + 4))`

### 3.32 $\int \frac{1}{(-5-3\cos(c+dx))^3} dx$

3.32.1	Optimal result . . . . .	212
3.32.2	Mathematica [A] (verified) . . . . .	212
3.32.3	Rubi [A] (verified) . . . . .	213
3.32.4	Maple [A] (verified) . . . . .	215
3.32.5	Fricas [A] (verification not implemented) . . . . .	215
3.32.6	Sympy [C] (verification not implemented) . . . . .	216
3.32.7	Maxima [A] (verification not implemented) . . . . .	216
3.32.8	Giac [A] (verification not implemented) . . . . .	217
3.32.9	Mupad [B] (verification not implemented) . . . . .	217

#### 3.32.1 Optimal result

Integrand size = 12, antiderivative size = 81

$$\int \frac{1}{(-5-3\cos(c+dx))^3} dx = -\frac{59x}{2048} + \frac{59 \arctan\left(\frac{\sin(c+dx)}{3+\cos(c+dx)}\right)}{1024d} + \frac{3\sin(c+dx)}{32d(5+3\cos(c+dx))^2} + \frac{45\sin(c+dx)}{512d(5+3\cos(c+dx))}$$

output `-59/2048*x+59/1024*arctan(sin(d*x+c)/(3+cos(d*x+c)))/d+3/32*sin(d*x+c)/d/(5+3*cos(d*x+c))^2+45/512*sin(d*x+c)/d/(5+3*cos(d*x+c))`

#### 3.32.2 Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.80

$$\int \frac{1}{(-5-3\cos(c+dx))^3} dx = \frac{59 \arctan\left(2 \cot\left(\frac{1}{2}(c+dx)\right)\right) (5+3\cos(c+dx))^2 + 546 \sin(c+dx) + 135 \sin(2(c+dx))}{1024d(5+3\cos(c+dx))^2}$$

input `Integrate[(-5 - 3*Cos[c + d*x])^(-3), x]`

output `(59*ArcTan[2*Cot[(c + d*x)/2]]*(5 + 3*Cos[c + d*x])^2 + 546*Sin[c + d*x] + 135*Sin[2*(c + d*x)])/(1024*d*(5 + 3*Cos[c + d*x])^2)`

**3.32.3 Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$ , Rules used = {3042, 3143, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(-3 \cos(c + dx) - 5)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(-3 \sin(c + dx + \frac{\pi}{2}) - 5)^3} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{3 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} - \frac{1}{32} \int \frac{10 - 3 \cos(c + dx)}{(3 \cos(c + dx) + 5)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{3 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} - \frac{1}{32} \int \frac{10 - 3 \sin(c + dx + \frac{\pi}{2})}{(3 \sin(c + dx + \frac{\pi}{2}) + 5)^2} dx \\
 & \quad \downarrow \text{3233} \\
 & \frac{1}{32} \left( \frac{1}{16} \int -\frac{59}{3 \cos(c + dx) + 5} dx + \frac{45 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)} \right) + \frac{3 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{32} \left( \frac{45 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)} - \frac{59}{16} \int \frac{1}{3 \cos(c + dx) + 5} dx \right) + \frac{3 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{32} \left( \frac{45 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)} - \frac{59}{16} \int \frac{1}{3 \sin(c + dx + \frac{\pi}{2}) + 5} dx \right) + \frac{3 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} \\
 & \quad \downarrow \text{3136} \\
 & \frac{1}{32} \left( \frac{45 \sin(c + dx)}{16d(3 \cos(c + dx) + 5)} - \frac{59}{16} \left( \frac{x}{4} - \frac{\arctan\left(\frac{\sin(c + dx)}{\cos(c + dx) + 3}\right)}{2d} \right) \right) + \frac{3 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2}
 \end{aligned}$$

input `Int[(-5 - 3*Cos[c + d*x])^(-3), x]`

```
output (3*Sin[c + d*x])/(32*d*(5 + 3*Cos[c + d*x])^2) + ((-59*(x/4 - ArcTan[Sin[c
+ d*x]/(3 + Cos[c + d*x]])/(2*d)))/16 + (45*Sin[c + d*x])/(16*d*(5 + 3*Cos
[c + d*x]))) / 32
```

### 3.32.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3136 Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[
a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q
+ b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] &&
PosQ[a]
```

```
rule 3143 Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos
[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp
[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1)
- b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

```
rule 3233 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e +
f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2))
Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(
m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]
```

**3.32.4 Maple [A] (verified)**

Time = 0.76 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.77

method	result
derivativedivides	$\frac{-\frac{69(\tan^3(\frac{dx}{2} + \frac{c}{2}))}{128} - \frac{51 \tan(\frac{dx}{2} + \frac{c}{2})}{32} - \frac{59 \arctan\left(\frac{\tan(\frac{dx}{2} + \frac{c}{2})}{2}\right)}{1024}}{4(\tan^2(\frac{dx}{2} + \frac{c}{2}) + 4)^2} d$
default	$\frac{-\frac{69(\tan^3(\frac{dx}{2} + \frac{c}{2}))}{128} - \frac{51 \tan(\frac{dx}{2} + \frac{c}{2})}{32} - \frac{59 \arctan\left(\frac{\tan(\frac{dx}{2} + \frac{c}{2})}{2}\right)}{1024}}{4(\tan^2(\frac{dx}{2} + \frac{c}{2}) + 4)^2} d$
risch	$\frac{3i(59 e^{3i(dx+c)} + 295 e^{2i(dx+c)} + 241 e^{i(dx+c)} + 45)}{256d(3 e^{2i(dx+c)} + 10 e^{i(dx+c)} + 3)^2} + \frac{59i \ln(e^{i(dx+c)} + \frac{1}{3})}{2048d} - \frac{59i \ln(e^{i(dx+c)} + 3)}{2048d}$
parallelrisch	$\frac{59i(59 + 9 \cos(2dx + 2c) + 60 \cos(dx + c)) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2i\right) + 59i(-59 - 9 \cos(2dx + 2c) - 60 \cos(dx + c)) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2048d(59 + 9 \cos(2dx + 2c) + 60 \cos(dx + c))}$

input `int(1/(-5-3*cos(d*x+c))^3,x,method=_RETURNVERBOSE)`output `1/d*(-1/4*(-69/128*tan(1/2*d*x+1/2*c)^3-51/32*tan(1/2*d*x+1/2*c))/(tan(1/2*d*x+1/2*c)^2+4)^2-59/1024*arctan(1/2*tan(1/2*d*x+1/2*c)))`**3.32.5 Fricas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.11

$$\int \frac{1}{(-5 - 3 \cos(c + dx))^3} dx$$

$$= \frac{59(9 \cos(dx + c)^2 + 30 \cos(dx + c) + 25) \arctan\left(\frac{5 \cos(dx + c) + 3}{4 \sin(dx + c)}\right) + 12(45 \cos(dx + c) + 91) \sin(dx + c)}{2048(9d \cos(dx + c)^2 + 30d \cos(dx + c) + 25d)}$$

input `integrate(1/(-5-3*cos(d*x+c))^3,x, algorithm="fricas")`output `1/2048*(59*(9*cos(d*x + c)^2 + 30*cos(d*x + c) + 25)*arctan(1/4*(5*cos(d*x + c) + 3)/sin(d*x + c)) + 12*(45*cos(d*x + c) + 91)*sin(d*x + c))/(9*d*cos(d*x + c)^2 + 30*d*cos(d*x + c) + 25*d)`



### 3.32.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.54 (sec) , antiderivative size = 362, normalized size of antiderivative = 4.47

$$\int \frac{1}{(-5 - 3 \cos(c + dx))^3} dx$$

$$= \begin{cases} \frac{x}{(-5 - 3 \cosh(2 \operatorname{atanh}(2)))^3} \\ \frac{x}{(-3 \cos(c) - 5)^3} \\ -\frac{59 \left( \operatorname{atan} \left( \frac{\tan \left( \frac{c}{2} + \frac{dx}{2} \right)}{2} \right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right) \tan^4 \left( \frac{c}{2} + \frac{dx}{2} \right)}{1024d \tan^4 \left( \frac{c}{2} + \frac{dx}{2} \right) + 8192d \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right) + 16384d} - \frac{472 \left( \operatorname{atan} \left( \frac{\tan \left( \frac{c}{2} + \frac{dx}{2} \right)}{2} \right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right) \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right)}{1024d \tan^4 \left( \frac{c}{2} + \frac{dx}{2} \right) + 8192d \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right) + 16384d} - \frac{944 \left( \operatorname{atan} \left( \frac{\tan \left( \frac{c}{2} + \frac{dx}{2} \right)}{2} \right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{1024d \tan^4 \left( \frac{c}{2} + \frac{dx}{2} \right) + 8192d \tan^2 \left( \frac{c}{2} + \frac{dx}{2} \right) + 16384d} \end{cases}$$

input `integrate(1/(-5-3*cos(d*x+c))**3,x)`

output `Piecewise((x/(-5 - 3*cosh(2*atanh(2)))**3, Eq(c, -d*x - 2*I*atanh(2)) | Eq(c, -d*x + 2*I*atanh(2))), (x/(-3*cos(c) - 5)**3, Eq(d, 0)), (-59*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4/(1024*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 16384*d) - 472*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**2/(1024*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 16384*d) - 944*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(1024*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 16384*d) + 138*atan(c/2 + d*x/2)**3/(1024*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 16384*d) + 408*tan(c/2 + d*x/2)/(1024*d*tan(c/2 + d*x/2)**4 + 8192*d*tan(c/2 + d*x/2)**2 + 16384*d), True))`

### 3.32.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.37

$$\int \frac{1}{(-5 - 3 \cos(c + dx))^3} dx = \frac{6 \left( \frac{68 \sin(dx+c)}{\cos(dx+c)+1} + \frac{23 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{\frac{8 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{\sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 16} - 59 \arctan \left( \frac{\sin(dx+c)}{2(\cos(dx+c)+1)} \right) \frac{1}{1024 d}$$

input `integrate(1/(-5-3*cos(d*x+c))^3,x, algorithm="maxima")`

output  $\frac{1}{1024} \cdot (6 \cdot (68 \cdot \sin(dx + c) / (\cos(dx + c) + 1) + 23 \cdot \sin(dx + c)^3 / (\cos(dx + c) + 1)^3) / (8 \cdot \sin(dx + c)^2 / (\cos(dx + c) + 1)^2 + \sin(dx + c)^4 / (\cos(dx + c) + 1)^4 + 16) - 59 \cdot \arctan(1/2 \cdot \sin(dx + c) / (\cos(dx + c) + 1))) / d$

### 3.32.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.93

$$\int \frac{1}{(-5 - 3 \cos(c + dx))^3} dx$$

$$= \frac{59 dx + 59 c - \frac{12 \left( 23 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 68 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\left( \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 4 \right)^2} - 118 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+3}\right)}{2048 d}$$

input `integrate(1/(-5-3*cos(d*x+c))^3,x, algorithm="giac")`

output  $\frac{-1/2048 \cdot (59 \cdot dx + 59 \cdot c - 12 \cdot (23 \cdot \tan(1/2 \cdot dx + 1/2 \cdot c)^3 + 68 \cdot \tan(1/2 \cdot dx + 1/2 \cdot c)) / (\tan(1/2 \cdot dx + 1/2 \cdot c)^2 + 4)^2 - 118 \cdot \arctan(\sin(dx + c) / (\cos(dx + c) + 3))) / d$

### 3.32.9 Mupad [B] (verification not implemented)

Time = 14.32 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.17

$$\int \frac{1}{(-5 - 3 \cos(c + dx))^3} dx = \frac{59 \left( \operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2} \right)}{1024 d} - \frac{59 \operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{1024 d}$$

$$+ \frac{\frac{69 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{512} + \frac{51 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{128}}{d \left( \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 8 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 16 \right)}$$

input `int(-1/(3*cos(c + d*x) + 5)^3,x)`

output  $\frac{(59 \cdot (\operatorname{atan}(\tan(c/2 + (dx)/2)) - (dx)/2)) / (1024 \cdot d) - (59 \cdot \operatorname{atan}(\tan(c/2 + (dx)/2) / 2)) / (1024 \cdot d) + ((51 \cdot \tan(c/2 + (dx)/2)) / 128 + (69 \cdot \tan(c/2 + (dx)/2)^3) / 512) / (d \cdot (8 \cdot \tan(c/2 + (dx)/2)^2 + \tan(c/2 + (dx)/2)^4 + 16))$

### 3.33 $\int \frac{1}{(-5-3 \cos(c+dx))^4} dx$

3.33.1	Optimal result . . . . .	218
3.33.2	Mathematica [A] (verified) . . . . .	218
3.33.3	Rubi [A] (verified) . . . . .	219
3.33.4	Maple [A] (verified) . . . . .	221
3.33.5	Fricas [A] (verification not implemented) . . . . .	222
3.33.6	Sympy [C] (verification not implemented) . . . . .	222
3.33.7	Maxima [A] (verification not implemented) . . . . .	223
3.33.8	Giac [A] (verification not implemented) . . . . .	224
3.33.9	Mupad [B] (verification not implemented) . . . . .	224

#### 3.33.1 Optimal result

Integrand size = 12, antiderivative size = 106

$$\int \frac{1}{(-5-3 \cos(c+dx))^4} dx = \frac{385x}{32768} - \frac{385 \arctan\left(\frac{\sin(c+dx)}{3+\cos(c+dx)}\right)}{16384d} - \frac{\sin(c+dx)}{16d(5+3 \cos(c+dx))^3} - \frac{25 \sin(c+dx)}{512d(5+3 \cos(c+dx))^2} - \frac{311 \sin(c+dx)}{8192d(5+3 \cos(c+dx))}$$

```
output 385/32768*x-385/16384*arctan(sin(d*x+c)/(3+cos(d*x+c)))/d-1/16*sin(d*x+c)/
d/(5+3*cos(d*x+c))^3-25/512*sin(d*x+c)/d/(5+3*cos(d*x+c))^2-311/8192*sin(d
*x+c)/d/(5+3*cos(d*x+c))
```

#### 3.33.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.62

$$\int \frac{1}{(-5-3 \cos(c+dx))^4} dx = -\frac{770 \arctan\left(2 \cot\left(\frac{1}{2}(c+dx)\right)\right) + \frac{9(4883 \sin(c+dx)+2340 \sin(2(c+dx))+311 \sin(3(c+dx)))}{(5+3 \cos(c+dx))^3}}{32768d}$$

```
input Integrate[(-5 - 3*Cos[c + d*x])^(-4), x]
```

```
output -1/32768*(770*ArcTan[2*Cot[(c + d*x)/2]] + (9*(4883*Sin[c + d*x] + 2340*Si
n[2*(c + d*x)] + 311*Sin[3*(c + d*x)]))/(5 + 3*Cos[c + d*x])^3/d
```

**3.33.3 Rubi [A] (verified)**

Time = 0.49 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.14, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$ , Rules used = {3042, 3143, 27, 3042, 3233, 25, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(-3 \cos(c + dx) - 5)^4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(-3 \sin(c + dx + \frac{\pi}{2}) - 5)^4} dx \\
 & \quad \downarrow \text{3143} \\
 & -\frac{1}{48} \int -\frac{3(5 - 2 \cos(c + dx))}{(3 \cos(c + dx) + 5)^3} dx - \frac{\sin(c + dx)}{16d(3 \cos(c + dx) + 5)^3} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{16} \int \frac{5 - 2 \cos(c + dx)}{(3 \cos(c + dx) + 5)^3} dx - \frac{\sin(c + dx)}{16d(3 \cos(c + dx) + 5)^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{16} \int \frac{5 - 2 \sin(c + dx + \frac{\pi}{2})}{(3 \sin(c + dx + \frac{\pi}{2}) + 5)^3} dx - \frac{\sin(c + dx)}{16d(3 \cos(c + dx) + 5)^3} \\
 & \quad \downarrow \text{3233} \\
 & \frac{1}{16} \left( -\frac{1}{32} \int -\frac{62 - 25 \cos(c + dx)}{(3 \cos(c + dx) + 5)^2} dx - \frac{25 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} \right) - \frac{\sin(c + dx)}{16d(3 \cos(c + dx) + 5)^3} \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{16} \left( \frac{1}{32} \int \frac{62 - 25 \cos(c + dx)}{(3 \cos(c + dx) + 5)^2} dx - \frac{25 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} \right) - \frac{\sin(c + dx)}{16d(3 \cos(c + dx) + 5)^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{16} \left( \frac{1}{32} \int \frac{62 - 25 \sin(c + dx + \frac{\pi}{2})}{(3 \sin(c + dx + \frac{\pi}{2}) + 5)^2} dx - \frac{25 \sin(c + dx)}{32d(3 \cos(c + dx) + 5)^2} \right) - \frac{\sin(c + dx)}{16d(3 \cos(c + dx) + 5)^3} \\
 & \quad \downarrow \text{3233}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{16} \left( \frac{1}{32} \left( -\frac{1}{16} \int -\frac{385}{3 \cos(c+dx)+5} dx - \frac{311 \sin(c+dx)}{16d(3 \cos(c+dx)+5)} \right) - \frac{25 \sin(c+dx)}{32d(3 \cos(c+dx)+5)^2} \right) - \\
& \quad \frac{\sin(c+dx)}{16d(3 \cos(c+dx)+5)^3} \\
& \quad \downarrow \text{27} \\
& \frac{1}{16} \left( \frac{1}{32} \left( \frac{385}{16} \int \frac{1}{3 \cos(c+dx)+5} dx - \frac{311 \sin(c+dx)}{16d(3 \cos(c+dx)+5)} \right) - \frac{25 \sin(c+dx)}{32d(3 \cos(c+dx)+5)^2} \right) - \\
& \quad \frac{\sin(c+dx)}{16d(3 \cos(c+dx)+5)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{16} \left( \frac{1}{32} \left( \frac{385}{16} \int \frac{1}{3 \sin(c+dx+\frac{\pi}{2})+5} dx - \frac{311 \sin(c+dx)}{16d(3 \cos(c+dx)+5)} \right) - \frac{25 \sin(c+dx)}{32d(3 \cos(c+dx)+5)^2} \right) - \\
& \quad \frac{\sin(c+dx)}{16d(3 \cos(c+dx)+5)^3} \\
& \quad \downarrow \text{3136} \\
& \frac{1}{16} \left( \frac{1}{32} \left( \frac{385}{16} \left( \frac{x}{4} - \frac{\arctan\left(\frac{\sin(c+dx)}{\cos(c+dx)+3}\right)}{2d} \right) - \frac{311 \sin(c+dx)}{16d(3 \cos(c+dx)+5)} \right) - \frac{25 \sin(c+dx)}{32d(3 \cos(c+dx)+5)^2} \right) - \\
& \quad \frac{\sin(c+dx)}{16d(3 \cos(c+dx)+5)^3}
\end{aligned}$$

input `Int[(-5 - 3*Cos[c + d*x])^(-4), x]`

output `-1/16*Sin[c + d*x]/(d*(5 + 3*Cos[c + d*x])^3) + ((-25*Sin[c + d*x])/(32*d*(5 + 3*Cos[c + d*x])^2) + ((385*(x/4 - ArcTan[Sin[c + d*x]/(3 + Cos[c + d*x]))]/(2*d)))/16 - (311*Sin[c + d*x])/(16*d*(5 + 3*Cos[c + d*x]))/32)/16`

### 3.33.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3136 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x]] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3233 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

### 3.33.4 Maple [A] (verified)

Time = 0.82 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.71

method	result
derivativedivides	$\frac{-\frac{639 \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{1024} - \frac{117 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{32} - \frac{369 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{64} + \frac{385 \arctan\left(\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2}\right)}{16384}}{8 \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\right)^3} d$
default	$\frac{-\frac{639 \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{1024} - \frac{117 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{32} - \frac{369 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{64} + \frac{385 \arctan\left(\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2}\right)}{16384}}{8 \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\right)^3} d$
risch	$-\frac{i(10395 e^{5i(dx+c)} + 86625 e^{4i(dx+c)} + 239470 e^{3i(dx+c)} + 218466 e^{2i(dx+c)} + 73575 e^{i(dx+c)} + 8397)}{12288d(3 e^{2i(dx+c)} + 10 e^{i(dx+c)} + 3)^3} + \frac{385i \ln(e^{i(dx+c)})}{32768d}$
parallelrisc	$\frac{385i(-770 - 27 \cos(3dx+3c) - 981 \cos(dx+c) - 270 \cos(2dx+2c)) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2i\right) + 385i(770 + 27 \cos(3dx+3c) + 981 \cos(dx+c) + 270 \cos(2dx+2c))}{32768d(770 + 27 \cos(3dx+3c) + 981 \cos(dx+c) + 270 \cos(2dx+2c))}$

3.33.  $\int \frac{1}{(-5-3\cos(c+dx))^4} dx$

```
input int(1/(-5-3*cos(d*x+c))^4,x,method=_RETURNVERBOSE)
```

```
output 1/d*(1/8*(-639/1024*tan(1/2*d*x+1/2*c)^5-117/32*tan(1/2*d*x+1/2*c)^3-369/6
4*tan(1/2*d*x+1/2*c))/(tan(1/2*d*x+1/2*c)^2+4)^3+385/16384*arctan(1/2*tan(
1/2*d*x+1/2*c))
```

### 3.33.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.14

$$\int \frac{1}{(-5 - 3 \cos(c + dx))^4} dx = \frac{385 (27 \cos(dx + c)^3 + 135 \cos(dx + c)^2 + 225 \cos(dx + c) + 125) \arctan\left(\frac{5 \cos(dx+c)+3}{4 \sin(dx+c)}\right) + 36 (311 \cos(dx + c)^2 + 1170 \cos(dx + c) + 1143) \sin(dx + c)}{32768 (27 d \cos(dx + c)^3 + 135 d \cos(dx + c)^2 + 225 d \cos(dx + c) + 125 d)}$$

```
input integrate(1/(-5-3*cos(d*x+c))^4,x, algorithm="fricas")
```

```
output -1/32768*(385*(27*cos(d*x + c)^3 + 135*cos(d*x + c)^2 + 225*cos(d*x + c) +
125)*arctan(1/4*(5*cos(d*x + c) + 3)/sin(d*x + c)) + 36*(311*cos(d*x + c)
^2 + 1170*cos(d*x + c) + 1143)*sin(d*x + c))/(27*d*cos(d*x + c)^3 + 135*d*
cos(d*x + c)^2 + 225*d*cos(d*x + c) + 125*d)
```

### 3.33.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 3.15 (sec) , antiderivative size = 595, normalized size of antiderivative = 5.61

$$\int \frac{1}{(-5 - 3 \cos(c + dx))^4} dx = \left\{ \begin{array}{l} \frac{x}{(-5-3 \cosh(2 \operatorname{atanh}(2)))^4} \\ \frac{x}{(-3 \cos(c)-5)^4} \\ \frac{385 \left( \operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right) + \pi \left[ \frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2} \right] \right) \tan^6\left(\frac{c}{2} + \frac{dx}{2}\right)}{16384d \tan^6\left(\frac{c}{2} + \frac{dx}{2}\right) + 196608d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) + 786432d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 1048576d} + \frac{4620 \left( \operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right) + \pi \left[ \frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2} \right] \right) \tan^6\left(\frac{c}{2} + \frac{dx}{2}\right)}{16384d \tan^6\left(\frac{c}{2} + \frac{dx}{2}\right) + 196608d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) + 786432d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 1048576d} \end{array} \right.$$

---

3.33.  $\int \frac{1}{(-5-3 \cos(c+dx))^4} dx$

input `integrate(1/(-5-3*cos(d*x+c))**4,x)`

output `Piecewise((x/(-5 - 3*cosh(2*atanh(2))))**4, Eq(c, -d*x - 2*I*atanh(2)) | Eq(c, -d*x + 2*I*atanh(2))), (x/(-3*cos(c) - 5)**4, Eq(d, 0)), (385*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**6/(16384*d*tan(c/2 + d*x/2)**6 + 196608*d*tan(c/2 + d*x/2)**4 + 786432*d*tan(c/2 + d*x/2)**2 + 1048576*d) + 4620*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4/(16384*d*tan(c/2 + d*x/2)**6 + 196608*d*tan(c/2 + d*x/2)**4 + 786432*d*tan(c/2 + d*x/2)**2 + 1048576*d) + 18480*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**2/(16384*d*tan(c/2 + d*x/2)**6 + 196608*d*tan(c/2 + d*x/2)**4 + 786432*d*tan(c/2 + d*x/2)**2 + 1048576*d) + 24640*(atan(tan(c/2 + d*x/2)/2) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(16384*d*tan(c/2 + d*x/2)**6 + 196608*d*tan(c/2 + d*x/2)**4 + 786432*d*tan(c/2 + d*x/2)**2 + 1048576*d) - 1278*tan(c/2 + d*x/2)**5/(16384*d*tan(c/2 + d*x/2)**6 + 196608*d*tan(c/2 + d*x/2)**4 + 786432*d*tan(c/2 + d*x/2)**2 + 1048576*d) - 7488*tan(c/2 + d*x/2)**3/(16384*d*tan(c/2 + d*x/2)**6 + 196608*d*tan(c/2 + d*x/2)**4 + 786432*d*tan(c/2 + d*x/2)**2 + 1048576*d) - 11808*tan(c/2 + d*x/2)/(16384*d*tan(c/2 + d*x/2)**6 + 196608*d*tan(c/2 + d*x/2)**4 + 786432*d*tan(c/2 + d*x/2)**2 + 1048576*d), True))`

### 3.33.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.42

$$\int \frac{1}{(-5 - 3 \cos(c + dx))^4} dx$$

$$= -\frac{18 \left( \frac{656 \sin(dx+c)}{\cos(dx+c)+1} + \frac{416 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{71 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right) - 385 \arctan \left( \frac{\sin(dx+c)}{2(\cos(dx+c)+1)} \right)}{16384 d}$$

input `integrate(1/(-5-3*cos(d*x+c))^4,x, algorithm="maxima")`

output `-1/16384*(18*(656*sin(d*x + c)/(cos(d*x + c) + 1) + 416*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 71*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/(48*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 12*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + 64) - 385*arctan(1/2*sin(d*x + c)/(cos(d*x + c) + 1)))/d`



**3.33.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.83

$$\int \frac{1}{(-5 - 3 \cos(c + dx))^4} dx$$

$$= \frac{385 dx + 385 c - \frac{36 \left( 71 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 + 416 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 656 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\left( \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 4 \right)^3} - 770 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+3}\right)}{32768 d}$$

input `integrate(1/(-5-3*cos(d*x+c))^4,x, algorithm="giac")`output `1/32768*(385*d*x + 385*c - 36*(71*tan(1/2*d*x + 1/2*c)^5 + 416*tan(1/2*d*x + 1/2*c)^3 + 656*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 4)^3 - 770*arctan(sin(d*x + c)/(cos(d*x + c) + 3))/d`**3.33.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.91

$$\int \frac{1}{(-5 - 3 \cos(c + dx))^4} dx = \frac{385 \operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{16384 d} - \frac{385 \left( \operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2} \right)}{16384 d}$$

$$- \frac{\frac{639 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{8192} + \frac{117 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{256} + \frac{369 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{512}}{d \left( \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 4 \right)^3}$$

input `int(1/(3*cos(c + d*x) + 5)^4,x)`output `(385*atan(tan(c/2 + (d*x)/2)/2))/(16384*d) - (385*(atan(tan(c/2 + (d*x)/2) - (d*x)/2))/(16384*d) - ((369*tan(c/2 + (d*x)/2))/512 + (117*tan(c/2 + (d*x)/2)^3)/256 + (639*tan(c/2 + (d*x)/2)^5)/8192)/(d*(tan(c/2 + (d*x)/2)^2 + 4)^3)`

### 3.34 $\int \frac{1}{3+5 \cos(c+dx)} dx$

3.34.1	Optimal result	225
3.34.2	Mathematica [A] (verified)	225
3.34.3	Rubi [A] (verified)	226
3.34.4	Maple [A] (verified)	227
3.34.5	Fricas [A] (verification not implemented)	227
3.34.6	Sympy [A] (verification not implemented)	228
3.34.7	Maxima [A] (verification not implemented)	228
3.34.8	Giac [A] (verification not implemented)	228
3.34.9	Mupad [B] (verification not implemented)	229

#### 3.34.1 Optimal result

Integrand size = 12, antiderivative size = 65

$$\int \frac{1}{3+5 \cos(c+dx)} dx = -\frac{\log \left(2 \cos \left(\frac{1}{2}(c+dx)\right) - \sin \left(\frac{1}{2}(c+dx)\right)\right)}{4d} + \frac{\log \left(2 \cos \left(\frac{1}{2}(c+dx)\right) + \sin \left(\frac{1}{2}(c+dx)\right)\right)}{4d}$$

output `-1/4*ln(2*cos(1/2*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d+1/4*ln(2*cos(1/2*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d`

#### 3.34.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00

$$\int \frac{1}{3+5 \cos(c+dx)} dx = -\frac{\log \left(2 \cos \left(\frac{1}{2}(c+dx)\right) - \sin \left(\frac{1}{2}(c+dx)\right)\right)}{4d} + \frac{\log \left(2 \cos \left(\frac{1}{2}(c+dx)\right) + \sin \left(\frac{1}{2}(c+dx)\right)\right)}{4d}$$

input `Integrate[(3 + 5*Cos[c + d*x])^(-1),x]`

output `-1/4*Log[2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]]/d + Log[2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]/(4*d)`

### 3.34.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.34, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {3042, 3138, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{1}{5 \cos(c + dx) + 3} dx \\
 \downarrow \text{3042} \\
 \int \frac{1}{5 \sin(c + dx + \frac{\pi}{2}) + 3} dx \\
 \downarrow \text{3138} \\
 \frac{2 \int \frac{1}{8 - 2 \tan^2(\frac{1}{2}(c + dx))} d \tan(\frac{1}{2}(c + dx))}{d} \\
 \downarrow \text{219} \\
 \frac{\operatorname{arctanh}(\frac{1}{2} \tan(\frac{1}{2}(c + dx)))}{2d}
 \end{array}$$

input `Int[(3 + 5*Cos[c + d*x])^(-1),x]`

output `ArcTanh[Tan[(c + d*x)/2]/2]/(2*d)`

#### 3.34.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3138 Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{
e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b +
(a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x]
&& NeQ[a^2 - b^2, 0]
```

### 3.34.4 Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.51

method	result	size
parallelrisch	$\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right) - \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{4d}$	33
derivativedivides	$\frac{\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{4} - \frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{4}}{d}$	34
default	$\frac{\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{4} - \frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{4}}{d}$	34
norman	$-\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{4d} + \frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{4d}$	36
risch	$-\frac{\ln\left(e^{i(dx+c)} + \frac{3}{5} - \frac{4i}{5}\right)}{4d} + \frac{\ln\left(e^{i(dx+c)} + \frac{3}{5} + \frac{4i}{5}\right)}{4d}$	40

```
input int(1/(3+5*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/4*(ln(tan(1/2*d*x+1/2*c)+2)-ln(tan(1/2*d*x+1/2*c)-2))/d
```

### 3.34.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.71

$$\int \frac{1}{3 + 5 \cos(c + dx)} dx$$

$$= \frac{\log\left(\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c) + \frac{5}{2}\right) - \log\left(\frac{3}{2} \cos(dx + c) - 2 \sin(dx + c) + \frac{5}{2}\right)}{8d}$$

```
input integrate(1/(3+5*cos(d*x+c)),x, algorithm="fricas")
```

```
output 1/8*(log(3/2*cos(d*x + c) + 2*sin(d*x + c) + 5/2) - log(3/2*cos(d*x + c) -
2*sin(d*x + c) + 5/2))/d
```

---

3.34.  $\int \frac{1}{3+5 \cos(c+dx)} dx$

**3.34.6 Sympy [A] (verification not implemented)**

Time = 0.35 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.63

$$\int \frac{1}{3 + 5 \cos(c + dx)} dx = \begin{cases} -\frac{\log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 2\right)}{4d} + \frac{\log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 2\right)}{4d} & \text{for } d \neq 0 \\ \frac{x}{5 \cos(c) + 3} & \text{otherwise} \end{cases}$$

input `integrate(1/(3+5*cos(d*x+c)),x)`output `Piecewise((-log(tan(c/2 + d*x/2) - 2)/(4*d) + log(tan(c/2 + d*x/2) + 2)/(4*d), Ne(d, 0)), (x/(5*cos(c) + 3), True))`**3.34.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.74

$$\int \frac{1}{3 + 5 \cos(c + dx)} dx = \frac{\log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 2\right) - \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 2\right)}{4d}$$

input `integrate(1/(3+5*cos(d*x+c)),x, algorithm="maxima")`output `1/4*(log(sin(d*x + c)/(cos(d*x + c) + 1) + 2) - log(sin(d*x + c)/(cos(d*x + c) + 1) - 2))/d`**3.34.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.52

$$\int \frac{1}{3 + 5 \cos(c + dx)} dx = \frac{\log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 2\right|\right) - \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 2\right|\right)}{4d}$$

input `integrate(1/(3+5*cos(d*x+c)),x, algorithm="giac")`output `1/4*(log(abs(tan(1/2*d*x + 1/2*c) + 2)) - log(abs(tan(1/2*d*x + 1/2*c) - 2))))/d`

**3.34.9 Mupad [B] (verification not implemented)**

Time = 14.42 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.26

$$\int \frac{1}{3 + 5 \cos(c + dx)} dx = \frac{\operatorname{atanh}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{2d}$$

input `int(1/(5*cos(c + d*x) + 3),x)`

output `atanh(tan(c/2 + (d*x)/2)/2)/(2*d)`

### 3.35 $\int \frac{1}{(3+5 \cos(c+dx))^2} dx$

3.35.1	Optimal result . . . . .	230
3.35.2	Mathematica [A] (verified) . . . . .	230
3.35.3	Rubi [A] (verified) . . . . .	231
3.35.4	Maple [A] (verified) . . . . .	233
3.35.5	Fricas [A] (verification not implemented) . . . . .	233
3.35.6	Sympy [B] (verification not implemented) . . . . .	234
3.35.7	Maxima [A] (verification not implemented) . . . . .	234
3.35.8	Giac [A] (verification not implemented) . . . . .	235
3.35.9	Mupad [B] (verification not implemented) . . . . .	235

#### 3.35.1 Optimal result

Integrand size = 12, antiderivative size = 90

$$\int \frac{1}{(3 + 5 \cos(c + dx))^2} dx = \frac{3 \log(2 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}{64d} - \frac{3 \log(2 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{64d} + \frac{5 \sin(c + dx)}{16d(3 + 5 \cos(c + dx))}$$

```
output 3/64*ln(2*cos(1/2*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d-3/64*ln(2*cos(1/2*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d+5/16*sin(d*x+c)/d/(3+5*cos(d*x+c))
```

#### 3.35.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.59

$$\int \frac{1}{(3 + 5 \cos(c + dx))^2} dx = \frac{9 \log(2 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) + 15 \cos(c + dx) (\log(2 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) - \log(2 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))))}{64d(3 + 5 \cos(c + dx))}$$

```
input Integrate[(3 + 5*Cos[c + d*x])^(-2), x]
```

output  $(9*\text{Log}[2*\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2]] + 15*\text{Cos}[c + d*x]*(\text{Log}[2*\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2]] - \text{Log}[2*\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]]) - 9*\text{Log}[2*\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]] + 20*\text{Sin}[c + d*x])/(64*d*(3 + 5*\text{Cos}[c + d*x]))$

### 3.35.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.53, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 3143, 27, 3042, 3138, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(5 \cos(c + dx) + 3)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(5 \sin(c + dx + \frac{\pi}{2}) + 3)^2} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{1}{16} \int -\frac{3}{5 \cos(c + dx) + 3} dx + \frac{5 \sin(c + dx)}{16d(5 \cos(c + dx) + 3)} \\
 & \quad \downarrow \text{27} \\
 & \frac{5 \sin(c + dx)}{16d(5 \cos(c + dx) + 3)} - \frac{3}{16} \int \frac{1}{5 \cos(c + dx) + 3} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{5 \sin(c + dx)}{16d(5 \cos(c + dx) + 3)} - \frac{3}{16} \int \frac{1}{5 \sin(c + dx + \frac{\pi}{2}) + 3} dx \\
 & \quad \downarrow \text{3138} \\
 & \frac{5 \sin(c + dx)}{16d(5 \cos(c + dx) + 3)} - \frac{3 \int \frac{1}{8 - 2 \tan^2(\frac{1}{2}(c + dx))} d \tan(\frac{1}{2}(c + dx))}{8d} \\
 & \quad \downarrow \text{219} \\
 & \frac{5 \sin(c + dx)}{16d(5 \cos(c + dx) + 3)} - \frac{3 \operatorname{arctanh}(\frac{1}{2} \tan(\frac{1}{2}(c + dx)))}{32d}
 \end{aligned}$$



input `Int[(3 + 5*Cos[c + d*x])^(-2), x]`

output `(-3*ArcTanh[Tan[(c + d*x)/2]/2])/(32*d) + (5*Sin[c + d*x])/(16*d*(3 + 5*Cos[c + d*x]))`

### 3.35.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

### 3.35.4 Maple [A] (verified)

Time = 0.63 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.71

method	result
derivativedivides	$\frac{-\frac{5}{32(\tan(\frac{dx}{2} + \frac{c}{2}) + 2)} - \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 2)}{64}}{d} - \frac{5}{32(\tan(\frac{dx}{2} + \frac{c}{2}) - 2)} + \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 2)}{64}$
default	$\frac{-\frac{5}{32(\tan(\frac{dx}{2} + \frac{c}{2}) + 2)} - \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 2)}{64}}{d} - \frac{5}{32(\tan(\frac{dx}{2} + \frac{c}{2}) - 2)} + \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 2)}{64}$
norman	$-\frac{5 \tan(\frac{dx}{2} + \frac{c}{2})}{16d(\tan^2(\frac{dx}{2} + \frac{c}{2}) - 4)} + \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 2)}{64d} - \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 2)}{64d}$
risch	$\frac{i(3e^{i(dx+c)} + 5)}{8d(5e^{2i(dx+c)} + 6e^{i(dx+c)} + 5)} - \frac{3 \ln(e^{i(dx+c)} + \frac{3}{5} + \frac{4i}{5})}{64d} + \frac{3 \ln(e^{i(dx+c)} + \frac{3}{5} - \frac{4i}{5})}{64d}$
parallelrisch	$\frac{15 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 2) \cos(dx+c) - 15 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 2) \cos(dx+c) + 20 \sin(dx+c) + 9 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 2) - 9 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 2)}{64d(3+5 \cos(dx+c))}$

input `int(1/(3+5*cos(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(-5/32/(tan(1/2*d*x+1/2*c)+2)-3/64*ln(tan(1/2*d*x+1/2*c)+2)-5/32/(tan(1/2*d*x+1/2*c)-2)+3/64*ln(tan(1/2*d*x+1/2*c)-2))`

### 3.35.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.98

$$\int \frac{1}{(3 + 5 \cos(c + dx))^2} dx = \frac{-3(5 \cos(dx + c) + 3) \log(\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c) + \frac{5}{2}) - 3(5 \cos(dx + c) + 3) \log(\frac{3}{2} \cos(dx + c) - 2 \sin(dx + c) + \frac{5}{2})}{128(5d \cos(dx + c) + 3d)}$$

input `integrate(1/(3+5*cos(d*x+c))^2,x, algorithm="fracas")`

output `-1/128*(3*(5*cos(d*x + c) + 3)*log(3/2*cos(d*x + c) + 2*sin(d*x + c) + 5/2) - 3*(5*cos(d*x + c) + 3)*log(3/2*cos(d*x + c) - 2*sin(d*x + c) + 5/2) - 40*sin(d*x + c))/(5*d*cos(d*x + c) + 3*d)`

### 3.35.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 228 vs. 2(78) = 156.

Time = 0.66 (sec) , antiderivative size = 228, normalized size of antiderivative = 2.53

$$\int \frac{1}{(3 + 5 \cos(c + dx))^2} dx$$

$$= \begin{cases} \frac{x}{(5 \cos(2 \operatorname{atan}(2)) + 3)^2} \\ \frac{x}{(5 \cos(c) + 3)^2} \\ \frac{3 \log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 2\right) \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{64d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 256d} - \frac{12 \log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 2\right)}{64d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 256d} - \frac{3 \log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 2\right) \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{64d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 256d} + \frac{12 \log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 2\right)}{64d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 256d} \end{cases}$$

input `integrate(1/(3+5*cos(d*x+c))**2,x)`

output `Piecewise((x/(5*cos(2*atan(2)) + 3)**2, Eq(c, -d*x - 2*atan(2)) | Eq(c, -d*x + 2*atan(2))), (x/(5*cos(c) + 3)**2, Eq(d, 0)), (3*log(tan(c/2 + d*x/2) - 2)*tan(c/2 + d*x/2)**2/(64*d*tan(c/2 + d*x/2)**2 - 256*d) - 12*log(tan(c/2 + d*x/2) - 2)/(64*d*tan(c/2 + d*x/2)**2 - 256*d) - 3*log(tan(c/2 + d*x/2) + 2)*tan(c/2 + d*x/2)**2/(64*d*tan(c/2 + d*x/2)**2 - 256*d) + 12*log(tan(c/2 + d*x/2) + 2)/(64*d*tan(c/2 + d*x/2)**2 - 256*d) - 20*tan(c/2 + d*x/2)/(64*d*tan(c/2 + d*x/2)**2 - 256*d), True))`

### 3.35.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.01

$$\int \frac{1}{(3 + 5 \cos(c + dx))^2} dx$$

$$= -\frac{\frac{20 \sin(dx+c)}{\left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} - 4\right)(\cos(dx+c)+1)} + 3 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 2\right) - 3 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 2\right)}{64d}$$

input `integrate(1/(3+5*cos(d*x+c))^2,x, algorithm="maxima")`

output `-1/64*(20*sin(d*x + c)/((sin(d*x + c)^2/(cos(d*x + c) + 1)^2 - 4)*(cos(d*x + c) + 1)) + 3*log(sin(d*x + c)/(cos(d*x + c) + 1) + 2) - 3*log(sin(d*x + c)/(cos(d*x + c) + 1) - 2))/d`

**3.35.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.69

$$\int \frac{1}{(3 + 5 \cos(c + dx))^2} dx$$

$$= -\frac{\frac{20 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 4} + 3 \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 2|) - 3 \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 2|)}{64 d}$$

input `integrate(1/(3+5*cos(d*x+c))^2,x, algorithm="giac")`output `-1/64*(20*tan(1/2*d*x + 1/2*c)/(tan(1/2*d*x + 1/2*c)^2 - 4) + 3*log(abs(tan(1/2*d*x + 1/2*c) + 2)) - 3*log(abs(tan(1/2*d*x + 1/2*c) - 2)))/d`**3.35.9 Mupad [B] (verification not implemented)**

Time = 14.18 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.52

$$\int \frac{1}{(3 + 5 \cos(c + dx))^2} dx = -\frac{3 \operatorname{atanh}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{32 d} - \frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{16 d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 4\right)}$$

input `int(1/(5*cos(c + d*x) + 3)^2,x)`output `-(3*atanh(tan(c/2 + (d*x)/2)/2))/(32*d) - (5*tan(c/2 + (d*x)/2))/(16*d*(tan(c/2 + (d*x)/2)^2 - 4))`

### 3.36 $\int \frac{1}{(3+5 \cos(c+dx))^3} dx$

3.36.1	Optimal result . . . . .	236
3.36.2	Mathematica [A] (verified) . . . . .	237
3.36.3	Rubi [A] (verified) . . . . .	237
3.36.4	Maple [A] (verified) . . . . .	240
3.36.5	Fricas [A] (verification not implemented) . . . . .	240
3.36.6	Sympy [B] (verification not implemented) . . . . .	241
3.36.7	Maxima [A] (verification not implemented) . . . . .	241
3.36.8	Giac [A] (verification not implemented) . . . . .	242
3.36.9	Mupad [B] (verification not implemented) . . . . .	242

#### 3.36.1 Optimal result

Integrand size = 12, antiderivative size = 115

$$\int \frac{1}{(3+5 \cos(c+dx))^3} dx = -\frac{43 \log(2 \cos(\frac{1}{2}(c+dx)) - \sin(\frac{1}{2}(c+dx)))}{2048d} + \frac{43 \log(2 \cos(\frac{1}{2}(c+dx)) + \sin(\frac{1}{2}(c+dx)))}{2048d} + \frac{5 \sin(c+dx)}{32d(3+5 \cos(c+dx))^2} - \frac{45 \sin(c+dx)}{512d(3+5 \cos(c+dx))}$$

output `-43/2048*ln(2*cos(1/2*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d+43/2048*ln(2*cos(1/2*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d+5/32*sin(d*x+c)/d/(3+5*cos(d*x+c))^2-45/512*sin(d*x+c)/d/(3+5*cos(d*x+c))`

### 3.36.2 Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.89

$$\int \frac{1}{(3 + 5 \cos(c + dx))^3} dx = -\frac{43 \log(2 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}{2048d} + \frac{43 \log(2 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{2048d} + \frac{512d(2 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))^2}{45 \sin(\frac{1}{2}(c + dx))} - \frac{2048d(2 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}{5} - \frac{512d(2 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))^2}{45 \sin(\frac{1}{2}(c + dx))} - \frac{2048d(2 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{5}$$

input `Integrate[(3 + 5*Cos[c + d*x])^(-3), x]`

output `(-43*Log[2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]]/(2048*d) + (43*Log[2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]/(2048*d) + 5/(512*d*(2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^2) - (45*Sin[(c + d*x)/2])/(2048*d*(2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2])) - 5/(512*d*(2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2) - (45*Sin[(c + d*x)/2])/(2048*d*(2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))`

### 3.36.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.68, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {3042, 3143, 25, 3042, 3233, 27, 3042, 3138, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(5 \cos(c + dx) + 3)^3} dx$$

↓ 3042

$$\int \frac{1}{(5 \sin(c + dx + \frac{\pi}{2}) + 3)^3} dx$$

$$\begin{aligned}
& \downarrow \text{3143} \\
& \frac{1}{32} \int -\frac{6-5\cos(c+dx)}{(5\cos(c+dx)+3)^2} dx + \frac{5\sin(c+dx)}{32d(5\cos(c+dx)+3)^2} \\
& \downarrow \text{25} \\
& \frac{5\sin(c+dx)}{32d(5\cos(c+dx)+3)^2} - \frac{1}{32} \int \frac{6-5\cos(c+dx)}{(5\cos(c+dx)+3)^2} dx \\
& \downarrow \text{3042} \\
& \frac{5\sin(c+dx)}{32d(5\cos(c+dx)+3)^2} - \frac{1}{32} \int \frac{6-5\sin(c+dx+\frac{\pi}{2})}{(5\sin(c+dx+\frac{\pi}{2})+3)^2} dx \\
& \downarrow \text{3233} \\
& \frac{1}{32} \left( -\frac{1}{16} \int -\frac{43}{5\cos(c+dx)+3} dx - \frac{45\sin(c+dx)}{16d(5\cos(c+dx)+3)} \right) + \frac{5\sin(c+dx)}{32d(5\cos(c+dx)+3)^2} \\
& \downarrow \text{27} \\
& \frac{1}{32} \left( \frac{43}{16} \int \frac{1}{5\cos(c+dx)+3} dx - \frac{45\sin(c+dx)}{16d(5\cos(c+dx)+3)} \right) + \frac{5\sin(c+dx)}{32d(5\cos(c+dx)+3)^2} \\
& \downarrow \text{3042} \\
& \frac{1}{32} \left( \frac{43}{16} \int \frac{1}{5\sin(c+dx+\frac{\pi}{2})+3} dx - \frac{45\sin(c+dx)}{16d(5\cos(c+dx)+3)} \right) + \frac{5\sin(c+dx)}{32d(5\cos(c+dx)+3)^2} \\
& \downarrow \text{3138} \\
& \frac{1}{32} \left( \frac{43 \int \frac{1}{8-2\tan^2(\frac{1}{2}(c+dx))} d \tan(\frac{1}{2}(c+dx))}{8d} - \frac{45\sin(c+dx)}{16d(5\cos(c+dx)+3)} \right) + \frac{5\sin(c+dx)}{32d(5\cos(c+dx)+3)^2} \\
& \downarrow \text{219} \\
& \frac{1}{32} \left( \frac{43 \operatorname{arctanh}(\frac{1}{2} \tan(\frac{1}{2}(c+dx)))}{32d} - \frac{45\sin(c+dx)}{16d(5\cos(c+dx)+3)} \right) + \frac{5\sin(c+dx)}{32d(5\cos(c+dx)+3)^2}
\end{aligned}$$

input `Int[(3 + 5*Cos[c + d*x])^(-3), x]`

output `(5*Sin[c + d*x])/(32*d*(3 + 5*Cos[c + d*x])^2) + ((43*ArcTanh[Tan[(c + d*x)/2]/2])/(32*d) - (45*Sin[c + d*x])/(16*d*(3 + 5*Cos[c + d*x]))) / 32`

## 3.36.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3143 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`
- rule 3233 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`



### 3.36.4 Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.72

method	result
norman	$-\frac{35 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{85 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{512d}}{(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 4)^2} - \frac{43 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{2048d} + \frac{43 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{2048d}$
derivativedivides	$\frac{\frac{25}{512\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)^2} + \frac{85}{1024\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)} - \frac{43 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{2048} - \frac{25}{512\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)^2} + \frac{85}{1024\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)} + \frac{43 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{2048}}{d}$
default	$\frac{\frac{25}{512\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)^2} + \frac{85}{1024\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)} - \frac{43 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{2048} - \frac{25}{512\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)^2} + \frac{85}{1024\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)} + \frac{43 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{2048}}{d}$
risch	$-\frac{i(215 e^{3i(dx+c)} + 387 e^{2i(dx+c)} + 325 e^{i(dx+c)} + 225)}{256d(5 e^{2i(dx+c)} + 6 e^{i(dx+c)} + 5)^2} - \frac{43 \ln\left(e^{i(dx+c)} + \frac{3}{5} - \frac{4i}{5}\right)}{2048d} + \frac{43 \ln\left(e^{i(dx+c)} + \frac{3}{5} + \frac{4i}{5}\right)}{2048d}$
parallelrisch	$\frac{(-2580 \cos(dx+c) - 1075 \cos(2dx+2c) - 1849) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right) + (2580 \cos(dx+c) + 1075 \cos(2dx+2c) + 1849) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{2048d(43 + 25 \cos(2dx+2c) + 60 \cos(dx+c))}$

input `int(1/(3+5*cos(d*x+c))^3,x,method=_RETURNVERBOSE)`

output 
$$\frac{(-35/128/d*\tan(1/2*d*x+1/2*c)+85/512/d*\tan(1/2*d*x+1/2*c)^3)/(\tan(1/2*d*x+1/2*c)^2-4)^2-43/2048/d*\ln(\tan(1/2*d*x+1/2*c)-2)+43/2048/d*\ln(\tan(1/2*d*x+1/2*c)+2)}$$

### 3.36.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.12

$$\int \frac{1}{(3 + 5 \cos(c + dx))^3} dx$$

$$= \frac{43 (25 \cos(dx + c)^2 + 30 \cos(dx + c) + 9) \log\left(\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c) + \frac{5}{2}\right) - 43 (25 \cos(dx + c)^2 + 30 \cos(dx + c) + 9) \log\left(\frac{3}{2} \cos(dx + c) - 2 \sin(dx + c) + \frac{5}{2}\right) - 40 (45 \cos(dx + c) + 11) \sin(dx + c)}{4096 (25 d \cos(dx + c))^2}$$

input `integrate(1/(3+5*cos(d*x+c))^3,x, algorithm="fricas")`

output 
$$\frac{1/4096*(43*(25*\cos(d*x + c)^2 + 30*\cos(d*x + c) + 9)*\log(3/2*\cos(d*x + c) + 2*\sin(d*x + c) + 5/2) - 43*(25*\cos(d*x + c)^2 + 30*\cos(d*x + c) + 9)*\log(3/2*\cos(d*x + c) - 2*\sin(d*x + c) + 5/2) - 40*(45*\cos(d*x + c) + 11)*\sin(d*x + c)}{(25*d*\cos(d*x + c))^2 + 30*d*\cos(d*x + c) + 9*d}$$

### 3.36.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 474 vs.  $2(102) = 204$ .

Time = 1.18 (sec) , antiderivative size = 474, normalized size of antiderivative = 4.12

$$\int \frac{1}{(3 + 5 \cos(c + dx))^3} dx$$

$$= \begin{cases} \frac{x}{(5 \cos(2 \operatorname{atan}(2)) + 3)^3} \\ \frac{x}{(5 \cos(c) + 3)^3} \\ -\frac{43 \log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 2\right) \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right)}{2048d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) - 16384d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 32768d} + \frac{344 \log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 2\right) \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{2048d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) - 16384d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 32768d} - \frac{688 \log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 2\right)}{2048d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) - 16384d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 32768d} \end{cases}$$

input `integrate(1/(3+5*cos(d*x+c))**3,x)`

output `Piecewise((x/(5*cos(2*atan(2)) + 3)**3, Eq(c, -d*x - 2*atan(2))) | Eq(c, -d*x + 2*atan(2))), (x/(5*cos(c) + 3)**3, Eq(d, 0)), (-43*log(tan(c/2 + d*x/2) - 2)*tan(c/2 + d*x/2)**4/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d) + 344*log(tan(c/2 + d*x/2) - 2)*tan(c/2 + d*x/2)**2/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d) - 688*log(tan(c/2 + d*x/2) - 2)/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d) + 43*log(tan(c/2 + d*x/2) + 2)*tan(c/2 + d*x/2)**4/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d) - 344*log(tan(c/2 + d*x/2) + 2)*tan(c/2 + d*x/2)**2/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d) + 688*log(tan(c/2 + d*x/2) + 2)/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d) + 340*tan(c/2 + d*x/2)**3/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d) - 560*tan(c/2 + d*x/2)/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d), True))`

### 3.36.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.17

$$\int \frac{1}{(3 + 5 \cos(c + dx))^3} dx$$

$$= \frac{20 \left( \frac{28 \sin(dx+c)}{\cos(dx+c)+1} - \frac{17 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{\frac{8 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - \frac{\sin(dx+c)^4}{(\cos(dx+c)+1)^4} - 16} + 43 \log \left( \frac{\sin(dx+c)}{\cos(dx+c)+1} + 2 \right) - 43 \log \left( \frac{\sin(dx+c)}{\cos(dx+c)+1} - 2 \right)$$

$$2048 d$$

---

3.36.  $\int \frac{1}{(3+5 \cos(c+dx))^3} dx$

input `integrate(1/(3+5*cos(d*x+c))^3,x, algorithm="maxima")`

output  $\frac{1}{2048} \cdot (20 \cdot (28 \cdot \sin(dx + c) / (\cos(dx + c) + 1) - 17 \cdot \sin(dx + c)^3 / (\cos(dx + c) + 1)^3) / (8 \cdot \sin(dx + c)^2 / (\cos(dx + c) + 1)^2 - \sin(dx + c)^4 / (\cos(dx + c) + 1)^4 - 16) + 43 \cdot \log(\sin(dx + c) / (\cos(dx + c) + 1) + 2) - 43 \cdot \log(\sin(dx + c) / (\cos(dx + c) + 1) - 2)) / d$

### 3.36.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.68

$$\int \frac{1}{(3 + 5 \cos(c + dx))^3} dx = \frac{20 \left( 17 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 28 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\left( \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 4 \right)^2} + 43 \log\left(\left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 2 \right|\right) - 43 \log\left(\left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 2 \right|\right)$$


---

2048 d

input `integrate(1/(3+5*cos(d*x+c))^3,x, algorithm="giac")`

output  $\frac{1}{2048} \cdot (20 \cdot (17 \cdot \tan(1/2 \cdot dx + 1/2 \cdot c)^3 - 28 \cdot \tan(1/2 \cdot dx + 1/2 \cdot c)) / (\tan(1/2 \cdot dx + 1/2 \cdot c)^2 - 4)^2 + 43 \cdot \log(\text{abs}(\tan(1/2 \cdot dx + 1/2 \cdot c) + 2)) - 43 \cdot \log(\text{abs}(\tan(1/2 \cdot dx + 1/2 \cdot c) - 2))) / d$

### 3.36.9 Mupad [B] (verification not implemented)

Time = 14.75 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.66

$$\int \frac{1}{(3 + 5 \cos(c + dx))^3} dx = \frac{43 \operatorname{atanh}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{1024 d} - \frac{\frac{35 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{128} - \frac{85 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{512}}{d \left( \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 8 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 16 \right)}$$

input `int(1/(5*cos(c + d*x) + 3)^3,x)`

output  $(43*\operatorname{atanh}(\tan(c/2 + (d*x)/2)/2))/(1024*d) - ((35*\tan(c/2 + (d*x)/2))/128 - (85*\tan(c/2 + (d*x)/2)^3)/512)/(d*(\tan(c/2 + (d*x)/2)^4 - 8*\tan(c/2 + (d*x)/2)^2 + 16))$

### 3.37 $\int \frac{1}{(3+5 \cos(c+dx))^4} dx$

3.37.1	Optimal result . . . . .	244
3.37.2	Mathematica [B] (verified) . . . . .	245
3.37.3	Rubi [A] (verified) . . . . .	245
3.37.4	Maple [A] (verified) . . . . .	248
3.37.5	Fricas [A] (verification not implemented) . . . . .	249
3.37.6	Sympy [B] (verification not implemented) . . . . .	249
3.37.7	Maxima [A] (verification not implemented) . . . . .	250
3.37.8	Giac [A] (verification not implemented) . . . . .	251
3.37.9	Mupad [B] (verification not implemented) . . . . .	251

#### 3.37.1 Optimal result

Integrand size = 12, antiderivative size = 140

$$\int \frac{1}{(3 + 5 \cos(c + dx))^4} dx = \frac{279 \log(2 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}{32768d} - \frac{279 \log(2 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{32768d} + \frac{5 \sin(c + dx)}{48d(3 + 5 \cos(c + dx))^3} - \frac{25 \sin(c + dx)}{512d(3 + 5 \cos(c + dx))^2} + \frac{995 \sin(c + dx)}{24576d(3 + 5 \cos(c + dx))}$$

```
output 279/32768*ln(2*cos(1/2*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d-279/32768*ln(2*cos
(1/2*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d+5/48*sin(d*x+c)/d/(3+5*cos(d*x+c))^3
-25/512*sin(d*x+c)/d/(3+5*cos(d*x+c))^2+995/24576*sin(d*x+c)/d/(3+5*cos(d*
x+c))
```

### 3.37.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 296 vs.  $2(140) = 280$ .

Time = 0.31 (sec) , antiderivative size = 296, normalized size of antiderivative = 2.11

$$\int \frac{1}{(3 + 5 \cos(c + dx))^4} dx$$

$$= \frac{467046 \log\left(2 \cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) + 104625 \cos(3(c + dx)) \log\left(2 \cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) - \sin\left(\frac{1}{2}(c + dx)\right)}{(3 + 5 \cos(c + dx))^4}$$

input `Integrate[(3 + 5*Cos[c + d*x])^(-4),x]`

output `(467046*Log[2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 104625*Cos[3*(c + d*x)]*Log[2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 765855*Cos[c + d*x]*(Log[2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - Log[2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]) + 376650*Cos[2*(c + d*x)]*(Log[2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - Log[2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]) - 467046*Log[2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] - 104625*Cos[3*(c + d*x)]*Log[2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + 226140*Sin[c + d*x] + 190800*Sin[2*(c + d*x)] + 99500*Sin[3*(c + d*x)])/(393216*d*(3 + 5*Cos[c + d*x])^3)`

### 3.37.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.77, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {3042, 3143, 25, 3042, 3233, 25, 3042, 3233, 27, 3042, 3138, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(5 \cos(c + dx) + 3)^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(5 \sin(c + dx + \frac{\pi}{2}) + 3)^4} dx$$

$$\downarrow \text{3143}$$

$$\frac{1}{48} \int -\frac{9 - 10 \cos(c + dx)}{(5 \cos(c + dx) + 3)^3} dx + \frac{5 \sin(c + dx)}{48d(5 \cos(c + dx) + 3)^3}$$

$$\begin{aligned}
& \downarrow 25 \\
& \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} - \frac{1}{48} \int \frac{9-10 \cos(c+dx)}{(5 \cos(c+dx)+3)^3} dx \\
& \downarrow 3042 \\
& \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} - \frac{1}{48} \int \frac{9-10 \sin(c+dx+\frac{\pi}{2})}{(5 \sin(c+dx+\frac{\pi}{2})+3)^3} dx \\
& \downarrow 3233 \\
& \frac{1}{48} \left( -\frac{1}{32} \int -\frac{154-75 \cos(c+dx)}{(5 \cos(c+dx)+3)^2} dx - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} \\
& \downarrow 25 \\
& \frac{1}{48} \left( \frac{1}{32} \int \frac{154-75 \cos(c+dx)}{(5 \cos(c+dx)+3)^2} dx - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left( \frac{1}{32} \int \frac{154-75 \sin(c+dx+\frac{\pi}{2})}{(5 \sin(c+dx+\frac{\pi}{2})+3)^2} dx - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} \\
& \downarrow 3233 \\
& \frac{1}{48} \left( \frac{1}{32} \left( \frac{1}{16} \int -\frac{837}{5 \cos(c+dx)+3} dx + \frac{995 \sin(c+dx)}{16d(5 \cos(c+dx)+3)} \right) - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \\
& \quad \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} \\
& \downarrow 27 \\
& \frac{1}{48} \left( \frac{1}{32} \left( \frac{995 \sin(c+dx)}{16d(5 \cos(c+dx)+3)} - \frac{837}{16} \int \frac{1}{5 \cos(c+dx)+3} dx \right) - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \\
& \quad \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left( \frac{1}{32} \left( \frac{995 \sin(c+dx)}{16d(5 \cos(c+dx)+3)} - \frac{837}{16} \int \frac{1}{5 \sin(c+dx+\frac{\pi}{2})+3} dx \right) - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \\
& \quad \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} \\
& \downarrow 3138
\end{aligned}$$

$$\frac{1}{48} \left( \frac{1}{32} \left( \frac{995 \sin(c+dx)}{16d(5 \cos(c+dx)+3)} - \frac{837 \int \frac{1}{8-2 \tan^2(\frac{1}{2}(c+dx))} d \tan(\frac{1}{2}(c+dx))}{8d} \right) - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3}$$

↓ 219

$$\frac{1}{48} \left( \frac{1}{32} \left( \frac{995 \sin(c+dx)}{16d(5 \cos(c+dx)+3)} - \frac{837 \operatorname{arctanh}(\frac{1}{2} \tan(\frac{1}{2}(c+dx)))}{32d} \right) - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3}$$

input `Int[(3 + 5*Cos[c + d*x])^(-4), x]`

output `(5*Sin[c + d*x])/(48*d*(3 + 5*Cos[c + d*x])^3) + ((-75*Sin[c + d*x])/(32*d*(3 + 5*Cos[c + d*x])^2) + ((-837*ArcTanh[Tan[(c + d*x)/2]/2])/(32*d) + (995*Sin[c + d*x])/(16*d*(3 + 5*Cos[c + d*x]))) / 32) / 48`

### 3.37.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`



rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3233 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

### 3.37.4 Maple [A] (verified)

Time = 0.82 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.71

method	result
norman	$-\frac{295 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{265 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{768d} - \frac{745 \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{8192d}}{\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 4\right)^3} + \frac{279 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{32768d} - \frac{279 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{32768d}$
derivativedivides	$\frac{-\frac{125}{6144 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)^3} - \frac{175}{4096 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)^2} - \frac{745}{16384 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)} + \frac{279 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{32768} - \frac{125}{6144 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)^3}}{d}$
default	$\frac{-\frac{125}{6144 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)^3} - \frac{175}{4096 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)^2} - \frac{745}{16384 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)} + \frac{279 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{32768} - \frac{125}{6144 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)^3}}{d}$
risch	$\frac{i(20925 e^{5i(dx+c)} + 62775 e^{4i(dx+c)} + 111042 e^{3i(dx+c)} + 119310 e^{2i(dx+c)} + 68625 e^{i(dx+c)} + 24875)}{12288d(5 e^{2i(dx+c)} + 6 e^{i(dx+c)} + 5)^3} + \frac{279 \ln(e^{i(dx+c)} + 2)}{32768d}$
parallelrisch	$\frac{(765855 \cos(dx+c) + 376650 \cos(2dx+2c) + 104625 \cos(3dx+3c) + 467046) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right) + (-765855 \cos(dx+c) - 391311 \cos(2dx+2c) - 104625 \cos(3dx+3c) - 467046)}{98304d(558 + 125 \cos(3dx+c))}$

input `int(1/(3+5*cos(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `(-295/512/d*tan(1/2*d*x+1/2*c)+265/768/d*tan(1/2*d*x+1/2*c)^3-745/8192/d*tan(1/2*d*x+1/2*c)^5)/(tan(1/2*d*x+1/2*c)^2-4)^3+279/32768/d*ln(tan(1/2*d*x+1/2*c)-2)-279/32768/d*ln(tan(1/2*d*x+1/2*c)+2)`

3.37.  $\int \frac{1}{(3+5 \cos(c+dx))^4} dx$

**3.37.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.21

$$\int \frac{1}{(3 + 5 \cos(c + dx))^4} dx = \frac{837 (125 \cos(dx + c)^3 + 225 \cos(dx + c)^2 + 135 \cos(dx + c) + 27) \log\left(\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c) + 5/2\right) - 837 (125 \cos(dx + c)^3 + 225 \cos(dx + c)^2 + 135 \cos(dx + c) + 27) \log\left(\frac{3}{2} \cos(dx + c) - 2 \sin(dx + c) + 5/2\right) - 40 (4975 \cos(dx + c)^2 + 4770 \cos(dx + c) + 1583) \sin(dx + c)}{(125 d \cos(dx + c)^3 + 225 d \cos(dx + c)^2 + 135 d \cos(dx + c) + 27 d)}$$

input `integrate(1/(3+5*cos(d*x+c))^4,x, algorithm="fracas")`

output `-1/196608*(837*(125*cos(d*x + c)^3 + 225*cos(d*x + c)^2 + 135*cos(d*x + c) + 27)*log(3/2*cos(d*x + c) + 2*sin(d*x + c) + 5/2) - 837*(125*cos(d*x + c)^3 + 225*cos(d*x + c)^2 + 135*cos(d*x + c) + 27)*log(3/2*cos(d*x + c) - 2*sin(d*x + c) + 5/2) - 40*(4975*cos(d*x + c)^2 + 4770*cos(d*x + c) + 1583)*sin(d*x + c))/(125*d*cos(d*x + c)^3 + 225*d*cos(d*x + c)^2 + 135*d*cos(d*x + c) + 27*d)`

**3.37.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 813 vs. 2(126) = 252.

Time = 2.46 (sec) , antiderivative size = 813, normalized size of antiderivative = 5.81

$$\int \frac{1}{(3 + 5 \cos(c + dx))^4} dx = \text{Too large to display}$$

input `integrate(1/(3+5*cos(d*x+c))**4,x)`

```

output Piecewise((x/(5*cos(2*atan(2)) + 3)**4, Eq(c, -d*x - 2*atan(2)) | Eq(c, -d
*x + 2*atan(2))), (x/(5*cos(c) + 3)**4, Eq(d, 0)), (837*log(tan(c/2 + d*x/
2) - 2)*tan(c/2 + d*x/2)**6/(98304*d*tan(c/2 + d*x/2)**6 - 1179648*d*tan(c
/2 + d*x/2)**4 + 4718592*d*tan(c/2 + d*x/2)**2 - 6291456*d) - 10044*log(ta
n(c/2 + d*x/2) - 2)*tan(c/2 + d*x/2)**4/(98304*d*tan(c/2 + d*x/2)**6 - 117
9648*d*tan(c/2 + d*x/2)**4 + 4718592*d*tan(c/2 + d*x/2)**2 - 6291456*d) +
40176*log(tan(c/2 + d*x/2) - 2)*tan(c/2 + d*x/2)**2/(98304*d*tan(c/2 + d*x
/2)**6 - 1179648*d*tan(c/2 + d*x/2)**4 + 4718592*d*tan(c/2 + d*x/2)**2 - 6
291456*d) - 53568*log(tan(c/2 + d*x/2) - 2)/(98304*d*tan(c/2 + d*x/2)**6 -
1179648*d*tan(c/2 + d*x/2)**4 + 4718592*d*tan(c/2 + d*x/2)**2 - 6291456*d
) - 837*log(tan(c/2 + d*x/2) + 2)*tan(c/2 + d*x/2)**6/(98304*d*tan(c/2 + d
*x/2)**6 - 1179648*d*tan(c/2 + d*x/2)**4 + 4718592*d*tan(c/2 + d*x/2)**2 -
6291456*d) + 10044*log(tan(c/2 + d*x/2) + 2)*tan(c/2 + d*x/2)**4/(98304*d
*tan(c/2 + d*x/2)**6 - 1179648*d*tan(c/2 + d*x/2)**4 + 4718592*d*tan(c/2 +
d*x/2)**2 - 6291456*d) - 40176*log(tan(c/2 + d*x/2) + 2)*tan(c/2 + d*x/2)
**2/(98304*d*tan(c/2 + d*x/2)**6 - 1179648*d*tan(c/2 + d*x/2)**4 + 4718592
*d*tan(c/2 + d*x/2)**2 - 6291456*d) + 53568*log(tan(c/2 + d*x/2) + 2)/(983
04*d*tan(c/2 + d*x/2)**6 - 1179648*d*tan(c/2 + d*x/2)**4 + 4718592*d*tan(c
/2 + d*x/2)**2 - 6291456*d) - 8940*tan(c/2 + d*x/2)**5/(98304*d*tan(c/2 +
d*x/2)**6 - 1179648*d*tan(c/2 + d*x/2)**4 + 4718592*d*tan(c/2 + d*x/2)*...

```

### 3.37.7 Maxima [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.24

$$\int \frac{1}{(3 + 5 \cos(c + dx))^4} dx = \frac{20 \left( \frac{2832 \sin(dx+c)}{\cos(dx+c)+1} - \frac{1696 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{447 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right) + 837 \log \left( \frac{\sin(dx+c)}{\cos(dx+c)+1} + 2 \right) - 837 \log \left( \frac{\sin(dx+c)}{\cos(dx+c)+1} - 2 \right)}{\frac{48 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - \frac{12 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{\sin(dx+c)^6}{(\cos(dx+c)+1)^6} - 64} 98304 d$$

```
input integrate(1/(3+5*cos(d*x+c))^4,x, algorithm="maxima")
```

```

output -1/98304*(20*(2832*sin(d*x + c)/(cos(d*x + c) + 1) - 1696*sin(d*x + c)^3/(
cos(d*x + c) + 1)^3 + 447*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/(48*sin(d*x
+ c)^2/(cos(d*x + c) + 1)^2 - 12*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + si
n(d*x + c)^6/(cos(d*x + c) + 1)^6 - 64) + 837*log(sin(d*x + c)/(cos(d*x +
c) + 1) + 2) - 837*log(sin(d*x + c)/(cos(d*x + c) + 1) - 2))/d

```

---

3.37.  $\int \frac{1}{(3+5 \cos(c+dx))^4} dx$

**3.37.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.65

$$\int \frac{1}{(3 + 5 \cos(c + dx))^4} dx = \frac{20 \left( 447 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 1696 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 2832 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\left( \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 4 \right)^3} + 837 \log \left( \left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 2 \right| \right) - 837 \log \left( \left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 2 \right| \right)$$


---

98304 d

input `integrate(1/(3+5*cos(d*x+c))^4,x, algorithm="giac")`output `-1/98304*(20*(447*tan(1/2*d*x + 1/2*c)^5 - 1696*tan(1/2*d*x + 1/2*c)^3 + 2832*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 4)^3 + 837*log(abs(tan(1/2*d*x + 1/2*c) + 2)) - 837*log(abs(tan(1/2*d*x + 1/2*c) - 2)))/d`**3.37.9 Mupad [B] (verification not implemented)**

Time = 16.71 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.73

$$\int \frac{1}{(3 + 5 \cos(c + dx))^4} dx = -\frac{279 \operatorname{atanh}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{16384 d} - \frac{\frac{745 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{8192} - \frac{265 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{768} + \frac{295 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{512}}{d \left( \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - 12 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 48 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 64 \right)}$$

input `int(1/(5*cos(c + d*x) + 3)^4,x)`output `-(279*atanh(tan(c/2 + (d*x)/2)/2))/(16384*d) - ((295*tan(c/2 + (d*x)/2))/512 - (265*tan(c/2 + (d*x)/2)^3)/768 + (745*tan(c/2 + (d*x)/2)^5)/8192)/(d*(48*tan(c/2 + (d*x)/2)^2 - 12*tan(c/2 + (d*x)/2)^4 + tan(c/2 + (d*x)/2)^6 - 64))`

### 3.38 $\int \frac{1}{3-5 \cos(c+dx)} dx$

3.38.1	Optimal result . . . . .	252
3.38.2	Mathematica [A] (verified) . . . . .	252
3.38.3	Rubi [A] (verified) . . . . .	253
3.38.4	Maple [A] (verified) . . . . .	254
3.38.5	Fricas [A] (verification not implemented) . . . . .	254
3.38.6	Sympy [A] (verification not implemented) . . . . .	255
3.38.7	Maxima [A] (verification not implemented) . . . . .	255
3.38.8	Giac [A] (verification not implemented) . . . . .	255
3.38.9	Mupad [B] (verification not implemented) . . . . .	256

#### 3.38.1 Optimal result

Integrand size = 12, antiderivative size = 63

$$\int \frac{1}{3-5 \cos(c+dx)} dx = \frac{\log(\cos(\frac{1}{2}(c+dx)) - 2 \sin(\frac{1}{2}(c+dx)))}{4d} - \frac{\log(\cos(\frac{1}{2}(c+dx)) + 2 \sin(\frac{1}{2}(c+dx)))}{4d}$$

output `1/4*ln(cos(1/2*d*x+1/2*c)-2*sin(1/2*d*x+1/2*c))/d-1/4*ln(cos(1/2*d*x+1/2*c)+2*sin(1/2*d*x+1/2*c))/d`

#### 3.38.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00

$$\int \frac{1}{3-5 \cos(c+dx)} dx = \frac{\log(\cos(\frac{1}{2}(c+dx)) - 2 \sin(\frac{1}{2}(c+dx)))}{4d} - \frac{\log(\cos(\frac{1}{2}(c+dx)) + 2 \sin(\frac{1}{2}(c+dx)))}{4d}$$

input `Integrate[(3 - 5*Cos[c + d*x])^(-1),x]`

output `Log[Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2]]/(4*d) - Log[Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2]]/(4*d)`

### 3.38.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.32, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {3042, 3138, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{1}{3 - 5 \cos(c + dx)} dx \\
 \downarrow \text{3042} \\
 \int \frac{1}{3 - 5 \sin(c + dx + \frac{\pi}{2})} dx \\
 \downarrow \text{3138} \\
 \frac{2 \int \frac{1}{8 \tan^2(\frac{1}{2}(c+dx)) - 2} d \tan(\frac{1}{2}(c + dx))}{d} \\
 \downarrow \text{220} \\
 \frac{\text{arctanh}(2 \tan(\frac{1}{2}(c + dx)))}{2d}
 \end{array}$$

input `Int[(3 - 5*Cos[c + d*x])^(-1),x]`

output `-1/2*ArcTanh[2*Tan[(c + d*x)/2]]/d`

#### 3.38.3.1 Defintions of rubi rules used

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3138 Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{
e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b +
(a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x]
&& NeQ[a^2 - b^2, 0]
```

### 3.38.4 Maple [A] (verified)

Time = 0.59 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.59

method	result	size
parallelrisch	$\frac{-\ln\left(2\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)+\ln\left(2\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}{4d}$	37
derivativedivides	$\frac{-\frac{\ln\left(2\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{4}+\frac{\ln\left(2\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}{4}}{d}$	38
default	$\frac{-\frac{\ln\left(2\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{4}+\frac{\ln\left(2\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}{4}}{d}$	38
norman	$\frac{\ln\left(2\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}{4d}-\frac{\ln\left(2\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{4d}$	40
risch	$\frac{\ln\left(e^{i(dx+c)}-\frac{3}{5}-\frac{4i}{5}\right)}{4d}-\frac{\ln\left(e^{i(dx+c)}-\frac{3}{5}+\frac{4i}{5}\right)}{4d}$	40

```
input int(1/(3-5*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/4*(-ln(2*tan(1/2*d*x+1/2*c)+1)+ln(2*tan(1/2*d*x+1/2*c)-1))/d
```

### 3.38.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.73

$$\int \frac{1}{3-5\cos(c+dx)} dx$$

$$= \frac{\log\left(-\frac{3}{2}\cos(dx+c)+2\sin(dx+c)+\frac{5}{2}\right)-\log\left(-\frac{3}{2}\cos(dx+c)-2\sin(dx+c)+\frac{5}{2}\right)}{8d}$$

```
input integrate(1/(3-5*cos(d*x+c)),x, algorithm="fricas")
```

```
output -1/8*(log(-3/2*cos(d*x + c) + 2*sin(d*x + c) + 5/2) - log(-3/2*cos(d*x + c)
) - 2*sin(d*x + c) + 5/2))/d
```

---

3.38.  $\int \frac{1}{3-5\cos(c+dx)} dx$

**3.38.6 Sympy [A] (verification not implemented)**

Time = 0.36 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.70

$$\int \frac{1}{3-5\cos(c+dx)} dx = \begin{cases} \frac{\log\left(2\tan\left(\frac{c}{2}+\frac{dx}{2}\right)-1\right)}{4d} - \frac{\log\left(2\tan\left(\frac{c}{2}+\frac{dx}{2}\right)+1\right)}{4d} & \text{for } d \neq 0 \\ \frac{x}{3-5\cos(c)} & \text{otherwise} \end{cases}$$

input `integrate(1/(3-5*cos(d*x+c)),x)`output `Piecewise((log(2*tan(c/2 + d*x/2) - 1)/(4*d) - log(2*tan(c/2 + d*x/2) + 1)/(4*d), Ne(d, 0)), (x/(3 - 5*cos(c)), True))`**3.38.7 Maxima [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.79

$$\int \frac{1}{3-5\cos(c+dx)} dx = -\frac{\log\left(\frac{2\sin(dx+c)}{\cos(dx+c)+1} + 1\right) - \log\left(\frac{2\sin(dx+c)}{\cos(dx+c)+1} - 1\right)}{4d}$$

input `integrate(1/(3-5*cos(d*x+c)),x, algorithm="maxima")`output `-1/4*(log(2*sin(d*x + c)/(cos(d*x + c) + 1) + 1) - log(2*sin(d*x + c)/(cos(d*x + c) + 1) - 1))/d`**3.38.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.60

$$\int \frac{1}{3-5\cos(c+dx)} dx = -\frac{\log\left(\left|2\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - \log\left(\left|2\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1\right|\right)}{4d}$$

input `integrate(1/(3-5*cos(d*x+c)),x, algorithm="giac")`output `-1/4*(log(abs(2*tan(1/2*d*x + 1/2*c) + 1)) - log(abs(2*tan(1/2*d*x + 1/2*c) - 1)))/d`



**3.38.9 Mupad [B] (verification not implemented)**

Time = 14.67 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.27

$$\int \frac{1}{3 - 5 \cos(c + dx)} dx = -\frac{\operatorname{atanh}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{2d}$$

input `int(-1/(5*cos(c + d*x) - 3),x)`

output `-atanh(2*tan(c/2 + (d*x)/2))/(2*d)`

### 3.39 $\int \frac{1}{(3-5 \cos(c+dx))^2} dx$

3.39.1	Optimal result . . . . .	257
3.39.2	Mathematica [A] (verified) . . . . .	257
3.39.3	Rubi [A] (verified) . . . . .	258
3.39.4	Maple [A] (verified) . . . . .	260
3.39.5	Fricas [A] (verification not implemented) . . . . .	260
3.39.6	Sympy [B] (verification not implemented) . . . . .	261
3.39.7	Maxima [A] (verification not implemented) . . . . .	261
3.39.8	Giac [A] (verification not implemented) . . . . .	262
3.39.9	Mupad [B] (verification not implemented) . . . . .	262

#### 3.39.1 Optimal result

Integrand size = 12, antiderivative size = 88

$$\int \frac{1}{(3-5 \cos(c+dx))^2} dx = -\frac{3 \log(\cos(\frac{1}{2}(c+dx)) - 2 \sin(\frac{1}{2}(c+dx)))}{64d} + \frac{3 \log(\cos(\frac{1}{2}(c+dx)) + 2 \sin(\frac{1}{2}(c+dx)))}{64d} - \frac{5 \sin(c+dx)}{16d(3-5 \cos(c+dx))}$$

```
output -3/64*ln(cos(1/2*d*x+1/2*c)-2*sin(1/2*d*x+1/2*c))/d+3/64*ln(cos(1/2*d*x+1/2*c)+2*sin(1/2*d*x+1/2*c))/d-5/16*sin(d*x+c)/d/(3-5*cos(d*x+c))
```

#### 3.39.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.58

$$\int \frac{1}{(3-5 \cos(c+dx))^2} dx = \frac{9 \log(\cos(\frac{1}{2}(c+dx)) - 2 \sin(\frac{1}{2}(c+dx))) - 15 \cos(c+dx) (\log(\cos(\frac{1}{2}(c+dx)) - 2 \sin(\frac{1}{2}(c+dx))) - \log(\cos(\frac{1}{2}(c+dx)) + 2 \sin(\frac{1}{2}(c+dx))))}{64d(-3+5 \cos(c+dx))}$$

```
input Integrate[(3 - 5*Cos[c + d*x])^(-2),x]
```

output  $(9*\text{Log}[\text{Cos}[(c + d*x)/2] - 2*\text{Sin}[(c + d*x)/2]] - 15*\text{Cos}[c + d*x]*(\text{Log}[\text{Cos}[(c + d*x)/2] - 2*\text{Sin}[(c + d*x)/2]] - \text{Log}[\text{Cos}[(c + d*x)/2] + 2*\text{Sin}[(c + d*x)/2]]) - 9*\text{Log}[\text{Cos}[(c + d*x)/2] + 2*\text{Sin}[(c + d*x)/2]] + 20*\text{Sin}[c + d*x])/(64*d*(-3 + 5*\text{Cos}[c + d*x]))$

### 3.39.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.52, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 3143, 27, 3042, 3138, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(3 - 5 \cos(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(3 - 5 \sin(c + dx + \frac{\pi}{2}))^2} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{1}{16} \int -\frac{3}{3 - 5 \cos(c + dx)} dx - \frac{5 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))} \\
 & \quad \downarrow \text{27} \\
 & -\frac{3}{16} \int \frac{1}{3 - 5 \cos(c + dx)} dx - \frac{5 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{3}{16} \int \frac{1}{3 - 5 \sin(c + dx + \frac{\pi}{2})} dx - \frac{5 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))} \\
 & \quad \downarrow \text{3138} \\
 & \frac{3 \int \frac{1}{8 \tan^2(\frac{1}{2}(c + dx)) - 2} d \tan(\frac{1}{2}(c + dx))}{8d} - \frac{5 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))} \\
 & \quad \downarrow \text{220} \\
 & \frac{3 \arctanh(2 \tan(\frac{1}{2}(c + dx)))}{32d} - \frac{5 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))}
 \end{aligned}$$

input `Int[(3 - 5*Cos[c + d*x])^(-2), x]`

output `(3*ArcTanh[2*Tan[(c + d*x)/2]]/(32*d) - (5*Sin[c + d*x])/(16*d*(3 - 5*Cos[c + d*x]))`

### 3.39.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3143 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

**3.39.4 Maple [A] (verified)**

Time = 0.64 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.81

method	result
norman	$-\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{16d\left(4\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)} - \frac{3 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{64d} + \frac{3 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{64d}$
derivativedivides	$\frac{-\frac{5}{64\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)} + \frac{3 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{64} - \frac{5}{64\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} - \frac{3 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{64}}{d}$
default	$\frac{-\frac{5}{64\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)} + \frac{3 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{64} - \frac{5}{64\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} - \frac{3 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{64}}{d}$
risch	$-\frac{i\left(3 e^{i(dx+c)} - 5\right)}{8d\left(5 e^{2i(dx+c)} - 6 e^{i(dx+c)} + 5\right)} + \frac{3 \ln\left(e^{i(dx+c)} - \frac{3}{5} + \frac{4i}{5}\right)}{64d} - \frac{3 \ln\left(e^{i(dx+c)} - \frac{3}{5} - \frac{4i}{5}\right)}{64d}$
parallelrisch	$\frac{-15 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) \cos(dx+c) + 15 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) \cos(dx+c) + 20 \sin(dx+c) + 9 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) - 9 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{64d(-3+5 \cos(dx+c))}$

input `int(1/(3-5*cos(d*x+c))^2,x,method=_RETURNVERBOSE)`output `-5/16/d*tan(1/2*d*x+1/2*c)/(4*tan(1/2*d*x+1/2*c)^2-1)-3/64/d*ln(2*tan(1/2*d*x+1/2*c)-1)+3/64/d*ln(2*tan(1/2*d*x+1/2*c)+1)`**3.39.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00

$$\int \frac{1}{(3 - 5 \cos(c + dx))^2} dx$$

$$= \frac{3(5 \cos(dx + c) - 3) \log\left(-\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c) + \frac{5}{2}\right) - 3(5 \cos(dx + c) - 3) \log\left(-\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c) - \frac{5}{2}\right) + 40 \sin(dx + c)}{128(5d \cos(dx + c) - 3d)}$$

input `integrate(1/(3-5*cos(d*x+c))^2,x, algorithm="fracas")`output `1/128*(3*(5*cos(d*x + c) - 3)*log(-3/2*cos(d*x + c) + 2*sin(d*x + c) + 5/2) - 3*(5*cos(d*x + c) - 3)*log(-3/2*cos(d*x + c) - 2*sin(d*x + c) + 5/2) + 40*sin(d*x + c))/(5*d*cos(d*x + c) - 3*d)`

**3.39.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 240 vs.  $2(78) = 156$ .

Time = 0.71 (sec) , antiderivative size = 240, normalized size of antiderivative = 2.73

$$\int \frac{1}{(3 - 5 \cos(c + dx))^2} dx$$

$$= \begin{cases} \frac{x}{(3 - 5 \cos(2 \operatorname{atan}(\frac{1}{2})))^2} \\ \frac{x}{(3 - 5 \cos(c))^2} \\ -\frac{12 \log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 1\right) \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{256d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 64d} + \frac{3 \log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 1\right)}{256d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 64d} + \frac{12 \log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 1\right) \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{256d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 64d} - \frac{3 \log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 1\right)}{256d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 64d} \end{cases}$$

input `integrate(1/(3-5*cos(d*x+c))**2,x)`

output `Piecewise((x/(3 - 5*cos(2*atan(1/2)))**2, Eq(c, -d*x - 2*atan(1/2)) | Eq(c, -d*x + 2*atan(1/2))), (x/(3 - 5*cos(c))**2, Eq(d, 0)), (-12*log(2*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**2/(256*d*tan(c/2 + d*x/2)**2 - 64*d) + 3*log(2*tan(c/2 + d*x/2) - 1)/(256*d*tan(c/2 + d*x/2)**2 - 64*d) + 12*log(2*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)**2/(256*d*tan(c/2 + d*x/2)**2 - 64*d) - 3*log(2*tan(c/2 + d*x/2) + 1)/(256*d*tan(c/2 + d*x/2)**2 - 64*d) - 2*0*tan(c/2 + d*x/2)/(256*d*tan(c/2 + d*x/2)**2 - 64*d), True))`

**3.39.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.07

$$\int \frac{1}{(3 - 5 \cos(c + dx))^2} dx$$

$$= -\frac{\frac{20 \sin(dx+c)}{\left(\frac{4 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - 1\right)(\cos(dx+c)+1)}}{64d} - 3 \log\left(\frac{2 \sin(dx+c)}{\cos(dx+c)+1} + 1\right) + 3 \log\left(\frac{2 \sin(dx+c)}{\cos(dx+c)+1} - 1\right)$$

input `integrate(1/(3-5*cos(d*x+c))^2,x, algorithm="maxima")`

output `-1/64*(20*sin(d*x + c)/((4*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 - 1)*(cos(d*x + c) + 1)) - 3*log(2*sin(d*x + c)/(cos(d*x + c) + 1) + 1) + 3*log(2*sin(d*x + c)/(cos(d*x + c) + 1) - 1))/d`

**3.39.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.77

$$\int \frac{1}{(3 - 5 \cos(c + dx))^2} dx$$

$$= -\frac{\frac{20 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{4 \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 1} - 3 \log(|2 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|) + 3 \log(|2 \tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{64 d}$$

input `integrate(1/(3-5*cos(d*x+c))^2,x, algorithm="giac")`output `-1/64*(20*tan(1/2*d*x + 1/2*c)/(4*tan(1/2*d*x + 1/2*c)^2 - 1) - 3*log(abs(2*tan(1/2*d*x + 1/2*c) + 1)) + 3*log(abs(2*tan(1/2*d*x + 1/2*c) - 1)))/d`**3.39.9 Mupad [B] (verification not implemented)**

Time = 14.32 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.53

$$\int \frac{1}{(3 - 5 \cos(c + dx))^2} dx = \frac{3 \operatorname{atanh}(2 \tan(\frac{c}{2} + \frac{dx}{2}))}{32 d} - \frac{5 \tan(\frac{c}{2} + \frac{dx}{2})}{64 d \left( \tan(\frac{c}{2} + \frac{dx}{2})^2 - \frac{1}{4} \right)}$$

input `int(1/(5*cos(c + d*x) - 3)^2,x)`output `(3*atanh(2*tan(c/2 + (d*x)/2)))/(32*d) - (5*tan(c/2 + (d*x)/2))/(64*d*(tan(c/2 + (d*x)/2)^2 - 1/4))`

### 3.40 $\int \frac{1}{(3-5 \cos(c+dx))^3} dx$

3.40.1	Optimal result . . . . .	263
3.40.2	Mathematica [A] (verified) . . . . .	264
3.40.3	Rubi [A] (verified) . . . . .	264
3.40.4	Maple [A] (verified) . . . . .	267
3.40.5	Fricas [A] (verification not implemented) . . . . .	267
3.40.6	Sympy [B] (verification not implemented) . . . . .	268
3.40.7	Maxima [A] (verification not implemented) . . . . .	268
3.40.8	Giac [A] (verification not implemented) . . . . .	269
3.40.9	Mupad [B] (verification not implemented) . . . . .	269

#### 3.40.1 Optimal result

Integrand size = 12, antiderivative size = 113

$$\int \frac{1}{(3 - 5 \cos(c + dx))^3} dx = \frac{43 \log(\cos(\frac{1}{2}(c + dx)) - 2 \sin(\frac{1}{2}(c + dx)))}{2048d} - \frac{43 \log(\cos(\frac{1}{2}(c + dx)) + 2 \sin(\frac{1}{2}(c + dx)))}{2048d} - \frac{5 \sin(c + dx)}{32d(3 - 5 \cos(c + dx))^2} + \frac{45 \sin(c + dx)}{512d(3 - 5 \cos(c + dx))}$$

```
output 43/2048*ln(cos(1/2*d*x+1/2*c)-2*sin(1/2*d*x+1/2*c))/d-43/2048*ln(cos(1/2*d
*x+1/2*c)+2*sin(1/2*d*x+1/2*c))/d-5/32*sin(d*x+c)/d/(3-5*cos(d*x+c))^2+45/
512*sin(d*x+c)/d/(3-5*cos(d*x+c))
```



### 3.40.2 Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.87

$$\int \frac{1}{(3 - 5 \cos(c + dx))^3} dx = \frac{43 \log(\cos(\frac{1}{2}(c + dx)) - 2 \sin(\frac{1}{2}(c + dx)))}{2048d} - \frac{43 \log(\cos(\frac{1}{2}(c + dx)) + 2 \sin(\frac{1}{2}(c + dx)))}{2048d} - \frac{512d (\cos(\frac{1}{2}(c + dx)) - 2 \sin(\frac{1}{2}(c + dx)))^2}{45 \sin(\frac{1}{2}(c + dx))} - \frac{1024d (\cos(\frac{1}{2}(c + dx)) - 2 \sin(\frac{1}{2}(c + dx)))}{5} + \frac{512d (\cos(\frac{1}{2}(c + dx)) + 2 \sin(\frac{1}{2}(c + dx)))^2}{45 \sin(\frac{1}{2}(c + dx))} - \frac{1024d (\cos(\frac{1}{2}(c + dx)) + 2 \sin(\frac{1}{2}(c + dx)))}{5}$$

input `Integrate[(3 - 5*Cos[c + d*x])^(-3), x]`

output `(43*Log[Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2]])/(2048*d) - (43*Log[Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2]])/(2048*d) - 5/(512*d*(Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2])^2) - (45*Sin[(c + d*x)/2])/(1024*d*(Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2])) + 5/(512*d*(Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2])^2) - (45*Sin[(c + d*x)/2])/(1024*d*(Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2]))`

### 3.40.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.67, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {3042, 3143, 25, 3042, 3233, 27, 3042, 3138, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(3 - 5 \cos(c + dx))^3} dx$$

↓ 3042

$$\int \frac{1}{(3 - 5 \sin(c + dx + \frac{\pi}{2}))^3} dx$$

---

3.40.  $\int \frac{1}{(3 - 5 \cos(c + dx))^3} dx$

$$\begin{aligned}
& \downarrow \text{3143} \\
& \frac{1}{32} \int -\frac{5 \cos(c+dx)+6}{(3-5 \cos(c+dx))^2} dx - \frac{5 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \\
& \downarrow \text{25} \\
& -\frac{1}{32} \int \frac{5 \cos(c+dx)+6}{(3-5 \cos(c+dx))^2} dx - \frac{5 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \\
& \downarrow \text{3042} \\
& -\frac{1}{32} \int \frac{5 \sin(c+dx+\frac{\pi}{2})+6}{(3-5 \sin(c+dx+\frac{\pi}{2}))^2} dx - \frac{5 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \\
& \downarrow \text{3233} \\
& \frac{1}{32} \left( \frac{45 \sin(c+dx)}{16d(3-5 \cos(c+dx))} - \frac{1}{16} \int -\frac{43}{3-5 \cos(c+dx)} dx \right) - \frac{5 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \\
& \downarrow \text{27} \\
& \frac{1}{32} \left( \frac{43}{16} \int \frac{1}{3-5 \cos(c+dx)} dx + \frac{45 \sin(c+dx)}{16d(3-5 \cos(c+dx))} \right) - \frac{5 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \\
& \downarrow \text{3042} \\
& \frac{1}{32} \left( \frac{43}{16} \int \frac{1}{3-5 \sin(c+dx+\frac{\pi}{2})} dx + \frac{45 \sin(c+dx)}{16d(3-5 \cos(c+dx))} \right) - \frac{5 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \\
& \downarrow \text{3138} \\
& \frac{1}{32} \left( \frac{43 \int \frac{1}{8 \tan^2(\frac{1}{2}(c+dx))-2} d \tan(\frac{1}{2}(c+dx))}{8d} + \frac{45 \sin(c+dx)}{16d(3-5 \cos(c+dx))} \right) - \frac{5 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \\
& \downarrow \text{220} \\
& \frac{1}{32} \left( \frac{45 \sin(c+dx)}{16d(3-5 \cos(c+dx))} - \frac{43 \operatorname{arctanh}(2 \tan(\frac{1}{2}(c+dx)))}{32d} \right) - \frac{5 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2}
\end{aligned}$$

input `Int[(3 - 5*Cos[c + d*x])^(-3), x]`

output `(-5*Sin[c + d*x])/(32*d*(3 - 5*Cos[c + d*x])^2) + ((-43*ArcTanh[2*Tan[(c + d*x)/2]])/(32*d) + (45*Sin[c + d*x])/(16*d*(3 - 5*Cos[c + d*x]))) / 32`

## 3.40.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3143 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`
- rule 3233 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

### 3.40.4 Maple [A] (verified)

Time = 0.70 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.79

method	result
norman	$-\frac{85 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 35 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{512d} + \frac{35 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{128d} + \frac{43 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{2048d} - \frac{43 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2048d}$
derivativedivides	$-\frac{25}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} + \frac{35}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} + \frac{43 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{2048} + \frac{25}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^2} + \frac{35}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}$
default	$-\frac{25}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} + \frac{35}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} + \frac{43 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{2048} + \frac{25}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^2} + \frac{35}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}$
risch	$\frac{i(215 e^{3i(dx+c)} - 387 e^{2i(dx+c)} + 325 e^{i(dx+c)} - 225)}{256d(5 e^{2i(dx+c)} - 6 e^{i(dx+c)} + 5)^2} + \frac{43 \ln\left(e^{i(dx+c)} - \frac{3}{5} - \frac{4i}{5}\right)}{2048d} - \frac{43 \ln\left(e^{i(dx+c)} - \frac{3}{5} + \frac{4i}{5}\right)}{2048d}$
parallelrisch	$\frac{(2580 \cos(dx+c) - 1075 \cos(2dx+2c) - 1849) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{1}{2}\right) + (-2580 \cos(dx+c) + 1075 \cos(2dx+2c) + 1849) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{1}{2}\right)}{2048d(-25 \cos(2dx+2c) - 43 + 60 \cos(dx+c))}$

input `int(1/(3-5*cos(d*x+c))^3,x,method=_RETURNVERBOSE)`

output 
$$\frac{(-85/512/d*\tan(1/2*d*x+1/2*c)+35/128/d*\tan(1/2*d*x+1/2*c)^3)/(4*\tan(1/2*d*x+1/2*c)^2-1)^2+43/2048/d*\ln(2*\tan(1/2*d*x+1/2*c)-1)-43/2048/d*\ln(2*\tan(1/2*d*x+1/2*c)+1)}{d}$$

### 3.40.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.14

$$\int \frac{1}{(3-5 \cos(c+dx))^3} dx = \frac{43(25 \cos(dx+c)^2 - 30 \cos(dx+c) + 9) \log\left(-\frac{3}{2} \cos(dx+c) + 2 \sin(dx+c) + \frac{5}{2}\right) - 43(25 \cos(dx+c) - 11) \sin(dx+c)}{4096(25d \cos(dx+c) - 30d + 9)}$$

input `integrate(1/(3-5*cos(d*x+c))^3,x, algorithm="fracas")`

output 
$$\frac{-1/4096*(43*(25*\cos(d*x+c)^2 - 30*\cos(d*x+c) + 9)*\log(-3/2*\cos(d*x+c) + 2*\sin(d*x+c) + 5/2) - 43*(25*\cos(d*x+c) - 11)*\sin(d*x+c)}{(25*d*\cos(d*x+c)^2 - 30*d*\cos(d*x+c) + 9*d)}$$

---

3.40. 
$$\int \frac{1}{(3-5 \cos(c+dx))^3} dx$$

### 3.40.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 490 vs.  $2(102) = 204$ .

Time = 1.19 (sec) , antiderivative size = 490, normalized size of antiderivative = 4.34

$$\int \frac{1}{(3 - 5 \cos(c + dx))^3} dx$$

$$= \begin{cases} \frac{x}{(3 - 5 \cos(2 \operatorname{atan}(\frac{1}{2})))^3} \\ \frac{x}{(3 - 5 \cos(c))^3} \\ \frac{688 \log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 1\right) \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right)}{32768d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) - 16384d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 2048d} - \frac{344 \log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 1\right) \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{32768d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) - 16384d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 2048d} + \frac{43 \log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 1\right)}{32768d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) - 16384d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 2048d} \end{cases}$$

input `integrate(1/(3-5*cos(d*x+c))**3,x)`

output `Piecewise((x/(3 - 5*cos(2*atan(1/2)))**3, Eq(c, -d*x - 2*atan(1/2)) | Eq(c, -d*x + 2*atan(1/2))), (x/(3 - 5*cos(c))**3, Eq(d, 0)), (688*log(2*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**4/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d) - 344*log(2*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**2/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d) + 43*log(2*tan(c/2 + d*x/2) - 1)/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d) - 688*log(2*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)**4/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d) + 344*log(2*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)**2/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d) - 43*log(2*tan(c/2 + d*x/2) + 1)/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d) + 560*tan(c/2 + d*x/2)**3/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d) - 340*tan(c/2 + d*x/2)/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d), True))`

### 3.40.7 Maxima [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.21

$$\int \frac{1}{(3 - 5 \cos(c + dx))^3} dx$$

$$= \frac{20 \left( \frac{17 \sin(dx+c)}{\cos(dx+c)+1} - \frac{28 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{\frac{8 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - \frac{16 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} - 1} - 43 \log \left( \frac{2 \sin(dx+c)}{\cos(dx+c)+1} + 1 \right) + 43 \log \left( \frac{2 \sin(dx+c)}{\cos(dx+c)+1} - 1 \right)$$

2048 d

---

3.40.  $\int \frac{1}{(3-5 \cos(c+dx))^3} dx$

input `integrate(1/(3-5*cos(d*x+c))^3,x, algorithm="maxima")`

output  $\frac{1}{2048} \cdot (20 \cdot (17 \cdot \sin(dx + c) / (\cos(dx + c) + 1) - 28 \cdot \sin(dx + c)^3 / (\cos(dx + c) + 1)^3) / (8 \cdot \sin(dx + c)^2 / (\cos(dx + c) + 1)^2 - 16 \cdot \sin(dx + c)^4 / (\cos(dx + c) + 1)^4 - 1) - 43 \cdot \log(2 \cdot \sin(dx + c) / (\cos(dx + c) + 1) + 1) + 43 \cdot \log(2 \cdot \sin(dx + c) / (\cos(dx + c) + 1) - 1)) / d$

### 3.40.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.74

$$\int \frac{1}{(3 - 5 \cos(c + dx))^3} dx = \frac{20 \left( 28 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 17 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{(4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 1)^2} - 43 \log\left(\left| 2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1 \right|\right) + 43 \log\left(\left| 2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1 \right|\right)$$


---

$2048 d$

input `integrate(1/(3-5*cos(d*x+c))^3,x, algorithm="giac")`

output  $\frac{1}{2048} \cdot (20 \cdot (28 \cdot \tan(1/2 \cdot dx + 1/2 \cdot c)^3 - 17 \cdot \tan(1/2 \cdot dx + 1/2 \cdot c)) / (4 \cdot \tan(1/2 \cdot dx + 1/2 \cdot c)^2 - 1)^2 - 43 \cdot \log(\text{abs}(2 \cdot \tan(1/2 \cdot dx + 1/2 \cdot c) + 1)) + 43 \cdot \log(\text{abs}(2 \cdot \tan(1/2 \cdot dx + 1/2 \cdot c) - 1))) / d$

### 3.40.9 Mupad [B] (verification not implemented)

Time = 14.15 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.67

$$\int \frac{1}{(3 - 5 \cos(c + dx))^3} dx = -\frac{43 \operatorname{atanh}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{1024 d} - \frac{\frac{85 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{8192} - \frac{35 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{2048}}{d \left( \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{2} + \frac{1}{16} \right)}$$

input `int(-1/(5*cos(c + d*x) - 3)^3,x)`

output  $-\frac{43 \operatorname{atanh}\left(2 \tan\left(\frac{c}{2} + \frac{d x}{2}\right)\right)}{1024 d} - \frac{\left(85 \tan\left(\frac{c}{2} + \frac{d x}{2}\right)\right)^2}{819} - \frac{\left(35 \tan\left(\frac{c}{2} + \frac{d x}{2}\right)\right)^3}{2048} / \left(d \left(\tan\left(\frac{c}{2} + \frac{d x}{2}\right)\right)^4 - \tan\left(\frac{c}{2} + \frac{d x}{2}\right)^2 + \frac{1}{16}\right)$

### 3.41 $\int \frac{1}{(3-5 \cos(c+dx))^4} dx$

3.41.1	Optimal result . . . . .	271
3.41.2	Mathematica [B] (verified) . . . . .	272
3.41.3	Rubi [A] (verified) . . . . .	272
3.41.4	Maple [A] (verified) . . . . .	275
3.41.5	Fricas [A] (verification not implemented) . . . . .	276
3.41.6	Sympy [B] (verification not implemented) . . . . .	276
3.41.7	Maxima [A] (verification not implemented) . . . . .	277
3.41.8	Giac [A] (verification not implemented) . . . . .	278
3.41.9	Mupad [B] (verification not implemented) . . . . .	278

#### 3.41.1 Optimal result

Integrand size = 12, antiderivative size = 138

$$\int \frac{1}{(3-5 \cos(c+dx))^4} dx = -\frac{279 \log(\cos(\frac{1}{2}(c+dx)) - 2 \sin(\frac{1}{2}(c+dx)))}{32768d} + \frac{279 \log(\cos(\frac{1}{2}(c+dx)) + 2 \sin(\frac{1}{2}(c+dx)))}{32768d} - \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} + \frac{25 \sin(c+dx)}{512d(3-5 \cos(c+dx))^2} - \frac{995 \sin(c+dx)}{24576d(3-5 \cos(c+dx))}$$

```
output -279/32768*ln(cos(1/2*d*x+1/2*c)-2*sin(1/2*d*x+1/2*c))/d+279/32768*ln(cos(
1/2*d*x+1/2*c)+2*sin(1/2*d*x+1/2*c))/d-5/48*sin(d*x+c)/d/(3-5*cos(d*x+c))^
3+25/512*sin(d*x+c)/d/(3-5*cos(d*x+c))^2-995/24576*sin(d*x+c)/d/(3-5*cos(d
*x+c))
```



### 3.41.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 288 vs.  $2(138) = 276$ .

Time = 0.36 (sec) , antiderivative size = 288, normalized size of antiderivative = 2.09

$$\int \frac{1}{(3 - 5 \cos(c + dx))^4} dx$$

$$= \frac{467046 \log\left(\cos\left(\frac{1}{2}(c + dx)\right) - 2 \sin\left(\frac{1}{2}(c + dx)\right)\right) - 104625 \cos(3(c + dx)) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) - 2 \sin\left(\frac{1}{2}(c + dx)\right)\right) - 2 \sin\left(\frac{1}{2}(c + dx)\right) - 104625 \cos(3(c + dx)) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + 2 \sin\left(\frac{1}{2}(c + dx)\right)\right) + 376650 \cos(2(c + dx)) \left(\log\left(\cos\left(\frac{1}{2}(c + dx)\right) - 2 \sin\left(\frac{1}{2}(c + dx)\right)\right) - \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + 2 \sin\left(\frac{1}{2}(c + dx)\right)\right)\right) - 467046 \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + 2 \sin\left(\frac{1}{2}(c + dx)\right)\right) + 104625 \cos(3(c + dx)) \left(\log\left(\cos\left(\frac{1}{2}(c + dx)\right) - 2 \sin\left(\frac{1}{2}(c + dx)\right)\right) + \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + 2 \sin\left(\frac{1}{2}(c + dx)\right)\right)\right) + 226140 \sin(c + dx) - 190800 \sin(2(c + dx)) + 99500 \sin(3(c + dx))}{(393216 d^3 - 5 \cos(c + dx))^3}$$

input `Integrate[(3 - 5*Cos[c + d*x])^(-4),x]`

output `(467046*Log[Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2]] - 104625*Cos[3*(c + d*x)]*Log[Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2]] - 765855*Cos[c + d*x]*(Log[Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2]] - Log[Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2]]) + 376650*Cos[2*(c + d*x)]*(Log[Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2]] - Log[Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2]]) - 467046*Log[Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2]] + 104625*Cos[3*(c + d*x)]*Log[Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2]] + 226140*Sin[c + d*x] - 190800*Sin[2*(c + d*x)] + 99500*Sin[3*(c + d*x)]/(393216*d*(-3 + 5*Cos[c + d*x])^3)`

### 3.41.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.77, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {3042, 3143, 25, 3042, 3233, 25, 3042, 3233, 27, 3042, 3138, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(3 - 5 \cos(c + dx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(3 - 5 \sin(c + dx + \frac{\pi}{2}))^4} dx$$

$$\downarrow \text{3143}$$

$$\frac{1}{48} \int -\frac{10 \cos(c + dx) + 9}{(3 - 5 \cos(c + dx))^3} dx - \frac{5 \sin(c + dx)}{48d(3 - 5 \cos(c + dx))^3}$$

---

3.41.  $\int \frac{1}{(3 - 5 \cos(c + dx))^4} dx$

$$\begin{aligned}
& \downarrow 25 \\
& -\frac{1}{48} \int \frac{10 \cos(c+dx) + 9}{(3-5 \cos(c+dx))^3} dx - \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 3042 \\
& -\frac{1}{48} \int \frac{10 \sin(c+dx + \frac{\pi}{2}) + 9}{(3-5 \sin(c+dx + \frac{\pi}{2}))^3} dx - \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 3233 \\
& \frac{1}{48} \left( \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} - \frac{1}{32} \int -\frac{75 \cos(c+dx) + 154}{(3-5 \cos(c+dx))^2} dx \right) - \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 25 \\
& \frac{1}{48} \left( \frac{1}{32} \int \frac{75 \cos(c+dx) + 154}{(3-5 \cos(c+dx))^2} dx + \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \right) - \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left( \frac{1}{32} \int \frac{75 \sin(c+dx + \frac{\pi}{2}) + 154}{(3-5 \sin(c+dx + \frac{\pi}{2}))^2} dx + \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \right) - \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 3233 \\
& \frac{1}{48} \left( \frac{1}{32} \left( \frac{1}{16} \int -\frac{837}{3-5 \cos(c+dx)} dx - \frac{995 \sin(c+dx)}{16d(3-5 \cos(c+dx))} \right) + \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \right) - \\
& \quad \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 27 \\
& \frac{1}{48} \left( \frac{1}{32} \left( -\frac{837}{16} \int \frac{1}{3-5 \cos(c+dx)} dx - \frac{995 \sin(c+dx)}{16d(3-5 \cos(c+dx))} \right) + \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \right) - \\
& \quad \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left( \frac{1}{32} \left( -\frac{837}{16} \int \frac{1}{3-5 \sin(c+dx + \frac{\pi}{2})} dx - \frac{995 \sin(c+dx)}{16d(3-5 \cos(c+dx))} \right) + \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \right) - \\
& \quad \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 3138
\end{aligned}$$

---

3.41.  $\int \frac{1}{(3-5 \cos(c+dx))^4} dx$

$$\frac{1}{48} \left( \frac{1}{32} \left( -\frac{837 \int \frac{1}{8 \tan^2(\frac{1}{2}(c+dx))-2} d \tan(\frac{1}{2}(c+dx))}{8d} - \frac{995 \sin(c+dx)}{16d(3-5 \cos(c+dx))} \right) + \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \right) - \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3}$$

↓ 220

$$\frac{1}{48} \left( \frac{1}{32} \left( \frac{837 \operatorname{arctanh}(2 \tan(\frac{1}{2}(c+dx)))}{32d} - \frac{995 \sin(c+dx)}{16d(3-5 \cos(c+dx))} \right) + \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \right) - \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3}$$

input `Int[(3 - 5*Cos[c + d*x])^(-4), x]`

output `(-5*Sin[c + d*x])/(48*d*(3 - 5*Cos[c + d*x])^3) + ((75*Sin[c + d*x])/(32*d*(3 - 5*Cos[c + d*x])^2) + ((837*ArcTanh[2*Tan[(c + d*x)/2]])/(32*d) - (99*5*Sin[c + d*x])/(16*d*(3 - 5*Cos[c + d*x]))) / 32) / 48`

### 3.41.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos [c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp [1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3233 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

### 3.41.4 Maple [A] (verified)

Time = 0.73 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.76

method	result
norman	$\frac{-\frac{745 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8192d} + \frac{265 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{768d} - \frac{295 \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{512d}}{\left(4 \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)^3} - \frac{279 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{32768d} + \frac{279 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{32768d}$
risch	$\frac{i(20925 e^{5i(dx+c)} - 62775 e^{4i(dx+c)} + 111042 e^{3i(dx+c)} - 119310 e^{2i(dx+c)} + 68625 e^{i(dx+c)} - 24875)}{12288d(5 e^{2i(dx+c)} - 6 e^{i(dx+c)} + 5)^3} + \frac{279 \ln(e^{i(dx+c)})}{32768d}$
derivativedivides	$\frac{-\frac{125}{49152 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^3} + \frac{25}{8192 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{295}{32768 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} - \frac{279 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{32768}}{d} - \frac{125}{49152 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^3} + \frac{25}{8192 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{295}{32768 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} - \frac{279 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{32768}}{d}$
default	$\frac{-\frac{125}{49152 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^3} + \frac{25}{8192 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{295}{32768 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} - \frac{279 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{32768}}{d} - \frac{125}{49152 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^3} + \frac{25}{8192 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{295}{32768 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} - \frac{279 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{32768}}{d}$
parallelrisch	$\frac{(-765855 \cos(dx+c) + 376650 \cos(2dx+2c) - 104625 \cos(3dx+3c) + 467046) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{1}{2}\right) + (765855 \cos(dx+c) - 376650 \cos(2dx+2c) + 104625 \cos(3dx+3c) - 467046) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{1}{2}\right)}{98304d(-558 + 125 \cos(3dx+c))}$

input `int(1/(3-5*cos(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `(-745/8192/d*tan(1/2*d*x+1/2*c)+265/768/d*tan(1/2*d*x+1/2*c)^3-295/512/d*tan(1/2*d*x+1/2*c)^5)/(4*tan(1/2*d*x+1/2*c)^2-1)^3-279/32768/d*ln(2*tan(1/2*d*x+1/2*c)-1)+279/32768/d*ln(2*tan(1/2*d*x+1/2*c)+1)`

3.41.  $\int \frac{1}{(3-5 \cos(c+dx))^4} dx$

**3.41.5 Fricas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.23

$$\int \frac{1}{(3 - 5 \cos(c + dx))^4} dx$$

$$= \frac{837 (125 \cos(dx + c)^3 - 225 \cos(dx + c)^2 + 135 \cos(dx + c) - 27) \log\left(-\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c)\right)}{196608 (125 \cos(dx + c)^3 - 225 \cos(dx + c)^2 + 135 \cos(dx + c) - 27) \log\left(-\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c)\right) + 40 (4975 \cos(dx + c)^2 - 4770 \cos(dx + c) + 1583) \sin(dx + c)}$$

input `integrate(1/(3-5*cos(d*x+c))^4,x, algorithm="fricas")`

output `1/196608*(837*(125*cos(d*x + c)^3 - 225*cos(d*x + c)^2 + 135*cos(d*x + c) - 27)*log(-3/2*cos(d*x + c) + 2*sin(d*x + c) + 5/2) - 837*(125*cos(d*x + c)^3 - 225*cos(d*x + c)^2 + 135*cos(d*x + c) - 27)*log(-3/2*cos(d*x + c) - 2*sin(d*x + c) + 5/2) + 40*(4975*cos(d*x + c)^2 - 4770*cos(d*x + c) + 1583)*sin(d*x + c))/(125*d*cos(d*x + c)^3 - 225*d*cos(d*x + c)^2 + 135*d*cos(d*x + c) - 27*d)`

**3.41.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 831 vs. 2(126) = 252.

Time = 2.47 (sec) , antiderivative size = 831, normalized size of antiderivative = 6.02

$$\int \frac{1}{(3 - 5 \cos(c + dx))^4} dx = \text{Too large to display}$$

input `integrate(1/(3-5*cos(d*x+c))**4,x)`

```
output Piecewise((x/(3 - 5*cos(2*atan(1/2))))**4, Eq(c, -d*x - 2*atan(1/2)) | Eq(c,
, -d*x + 2*atan(1/2))), (x/(3 - 5*cos(c))**4, Eq(d, 0)), (-53568*log(2*tan
(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**6/(6291456*d*tan(c/2 + d*x/2)**6 - 47
18592*d*tan(c/2 + d*x/2)**4 + 1179648*d*tan(c/2 + d*x/2)**2 - 98304*d) + 4
0176*log(2*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**4/(6291456*d*tan(c/2 +
d*x/2)**6 - 4718592*d*tan(c/2 + d*x/2)**4 + 1179648*d*tan(c/2 + d*x/2)**2
- 98304*d) - 10044*log(2*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**2/(629145
6*d*tan(c/2 + d*x/2)**6 - 4718592*d*tan(c/2 + d*x/2)**4 + 1179648*d*tan(c/
2 + d*x/2)**2 - 98304*d) + 837*log(2*tan(c/2 + d*x/2) - 1)/(6291456*d*tan(
c/2 + d*x/2)**6 - 4718592*d*tan(c/2 + d*x/2)**4 + 1179648*d*tan(c/2 + d*x/
2)**2 - 98304*d) + 53568*log(2*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)**6/(
6291456*d*tan(c/2 + d*x/2)**6 - 4718592*d*tan(c/2 + d*x/2)**4 + 1179648*d*
tan(c/2 + d*x/2)**2 - 98304*d) - 40176*log(2*tan(c/2 + d*x/2) + 1)*tan(c/2
+ d*x/2)**4/(6291456*d*tan(c/2 + d*x/2)**6 - 4718592*d*tan(c/2 + d*x/2)**
4 + 1179648*d*tan(c/2 + d*x/2)**2 - 98304*d) + 10044*log(2*tan(c/2 + d*x/2
) + 1)*tan(c/2 + d*x/2)**2/(6291456*d*tan(c/2 + d*x/2)**6 - 4718592*d*tan(
c/2 + d*x/2)**4 + 1179648*d*tan(c/2 + d*x/2)**2 - 98304*d) - 837*log(2*tan
(c/2 + d*x/2) + 1)/(6291456*d*tan(c/2 + d*x/2)**6 - 4718592*d*tan(c/2 + d*
x/2)**4 + 1179648*d*tan(c/2 + d*x/2)**2 - 98304*d) - 56640*tan(c/2 + d*x/2
)**5/(6291456*d*tan(c/2 + d*x/2)**6 - 4718592*d*tan(c/2 + d*x/2)**4 + 1...
```

### 3.41.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.28

$$\int \frac{1}{(3 - 5 \cos(c + dx))^4} dx = \frac{20 \left( \frac{447 \sin(dx+c)}{\cos(dx+c)+1} - \frac{1696 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{2832 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right) - \frac{12 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - \frac{48 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{64 \sin(dx+c)^6}{(\cos(dx+c)+1)^6} - 1}{98304 d} - 837 \log \left( \frac{2 \sin(dx+c)}{\cos(dx+c)+1} + 1 \right) + 837 \log \left( \frac{2 \sin(dx+c)}{\cos(dx+c)+1} - 1 \right)$$

```
input integrate(1/(3-5*cos(d*x+c))^4,x, algorithm="maxima")
```

```
output -1/98304*(20*(447*sin(d*x + c)/(cos(d*x + c) + 1) - 1696*sin(d*x + c)^3/(c
os(d*x + c) + 1)^3 + 2832*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/(12*sin(d*x
+ c)^2/(cos(d*x + c) + 1)^2 - 48*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 64
*sin(d*x + c)^6/(cos(d*x + c) + 1)^6 - 1) - 837*log(2*sin(d*x + c)/(cos(d*
x + c) + 1) + 1) + 837*log(2*sin(d*x + c)/(cos(d*x + c) + 1) - 1))/d
```

---

3.41.  $\int \frac{1}{(3-5 \cos(c+dx))^4} dx$

**3.41.8 Giac [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.70

$$\int \frac{1}{(3 - 5 \cos(c + dx))^4} dx = \frac{20 \left( 2832 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 1696 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 447 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\left( 4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 1 \right)^3} - 837 \log \left( \left| 2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1 \right| \right) + 837 \log \left( \left| 2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1 \right| \right)$$


---

98304 d

input `integrate(1/(3-5*cos(d*x+c))^4,x, algorithm="giac")`output `-1/98304*(20*(2832*tan(1/2*d*x + 1/2*c)^5 - 1696*tan(1/2*d*x + 1/2*c)^3 + 447*tan(1/2*d*x + 1/2*c))/(4*tan(1/2*d*x + 1/2*c)^2 - 1)^3 - 837*log(abs(2*tan(1/2*d*x + 1/2*c) + 1)) + 837*log(abs(2*tan(1/2*d*x + 1/2*c) - 1)))/d`**3.41.9 Mupad [B] (verification not implemented)**

Time = 15.70 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.74

$$\int \frac{1}{(3 - 5 \cos(c + dx))^4} dx = \frac{279 \operatorname{atanh}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{16384 d} - \frac{\frac{295 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{32768} - \frac{265 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{49152} + \frac{745 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{524288}}{d \left( \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - \frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{4} + \frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{16} - \frac{1}{64} \right)}$$

input `int(1/(5*cos(c + d*x) - 3)^4,x)`output `(279*atanh(2*tan(c/2 + (d*x)/2)))/(16384*d) - ((745*tan(c/2 + (d*x)/2))/524288 - (265*tan(c/2 + (d*x)/2)^3)/49152 + (295*tan(c/2 + (d*x)/2)^5)/32768)/(d*((3*tan(c/2 + (d*x)/2)^2)/16 - (3*tan(c/2 + (d*x)/2)^4)/4 + tan(c/2 + (d*x)/2)^6 - 1/64))`

### 3.42 $\int \frac{1}{-3+5\cos(c+dx)} dx$

3.42.1	Optimal result . . . . .	279
3.42.2	Mathematica [A] (verified) . . . . .	279
3.42.3	Rubi [A] (verified) . . . . .	280
3.42.4	Maple [A] (verified) . . . . .	281
3.42.5	Fricas [A] (verification not implemented) . . . . .	281
3.42.6	Sympy [A] (verification not implemented) . . . . .	282
3.42.7	Maxima [A] (verification not implemented) . . . . .	282
3.42.8	Giac [A] (verification not implemented) . . . . .	282
3.42.9	Mupad [B] (verification not implemented) . . . . .	283

#### 3.42.1 Optimal result

Integrand size = 12, antiderivative size = 63

$$\int \frac{1}{-3+5\cos(c+dx)} dx = -\frac{\log\left(\cos\left(\frac{1}{2}(c+dx)\right) - 2\sin\left(\frac{1}{2}(c+dx)\right)\right)}{4d} + \frac{\log\left(\cos\left(\frac{1}{2}(c+dx)\right) + 2\sin\left(\frac{1}{2}(c+dx)\right)\right)}{4d}$$

output `-1/4*ln(cos(1/2*d*x+1/2*c)-2*sin(1/2*d*x+1/2*c))/d+1/4*ln(cos(1/2*d*x+1/2*c)+2*sin(1/2*d*x+1/2*c))/d`

#### 3.42.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00

$$\int \frac{1}{-3+5\cos(c+dx)} dx = -\frac{\log\left(\cos\left(\frac{1}{2}(c+dx)\right) - 2\sin\left(\frac{1}{2}(c+dx)\right)\right)}{4d} + \frac{\log\left(\cos\left(\frac{1}{2}(c+dx)\right) + 2\sin\left(\frac{1}{2}(c+dx)\right)\right)}{4d}$$

input `Integrate[(-3 + 5*Cos[c + d*x])^(-1), x]`

output `-1/4*Log[Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2]]/d + Log[Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2]]/(4*d)`



### 3.42.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.32, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {3042, 3138, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{1}{5 \cos(c + dx) - 3} dx \\
 \downarrow \text{3042} \\
 \int \frac{1}{5 \sin(c + dx + \frac{\pi}{2}) - 3} dx \\
 \downarrow \text{3138} \\
 2 \int \frac{1}{2 - 8 \tan^2(\frac{1}{2}(c + dx))} d \tan(\frac{1}{2}(c + dx)) \\
 \downarrow \text{219} \\
 \frac{\operatorname{arctanh}(2 \tan(\frac{1}{2}(c + dx)))}{2d}
 \end{array}$$

input `Int[(-3 + 5*Cos[c + d*x])^(-1), x]`

output `ArcTanh[2*Tan[(c + d*x)/2]]/(2*d)`

#### 3.42.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3138 Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{
e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b +
(a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x]
&& NeQ[a^2 - b^2, 0]
```

### 3.42.4 Maple [A] (verified)

Time = 0.65 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.59

method	result	size
parallelrisch	$\frac{\ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) - \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{4d}$	37
derivativedivides	$\frac{\frac{\ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{4} - \frac{\ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{4}}{d}$	38
default	$\frac{\frac{\ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{4} - \frac{\ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{4}}{d}$	38
norman	$-\frac{\ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{4d} + \frac{\ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{4d}$	40
risch	$-\frac{\ln\left(e^{i(dx+c)} - \frac{3}{5} - \frac{4i}{5}\right)}{4d} + \frac{\ln\left(e^{i(dx+c)} - \frac{3}{5} + \frac{4i}{5}\right)}{4d}$	40

```
input int(1/(-3+5*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/4*(ln(2*tan(1/2*d*x+1/2*c)+1)-ln(2*tan(1/2*d*x+1/2*c)-1))/d
```

### 3.42.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.73

$$\int \frac{1}{-3 + 5 \cos(c + dx)} dx$$

$$= \frac{\log\left(-\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c) + \frac{5}{2}\right) - \log\left(-\frac{3}{2} \cos(dx + c) - 2 \sin(dx + c) + \frac{5}{2}\right)}{8d}$$

```
input integrate(1/(-3+5*cos(d*x+c)),x, algorithm="fracas")
```

```
output 1/8*(log(-3/2*cos(d*x + c) + 2*sin(d*x + c) + 5/2) - log(-3/2*cos(d*x + c)
- 2*sin(d*x + c) + 5/2))/d
```

---

3.42.  $\int \frac{1}{-3+5 \cos(c+dx)} dx$

**3.42.6 Sympy [A] (verification not implemented)**

Time = 0.40 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.70

$$\int \frac{1}{-3 + 5 \cos(c + dx)} dx = \begin{cases} -\frac{\log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 1\right)}{4d} + \frac{\log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 1\right)}{4d} & \text{for } d \neq 0 \\ \frac{x}{5 \cos(c) - 3} & \text{otherwise} \end{cases}$$

input `integrate(1/(-3+5*cos(d*x+c)),x)`output `Piecewise((-log(2*tan(c/2 + d*x/2) - 1)/(4*d) + log(2*tan(c/2 + d*x/2) + 1)/(4*d), Ne(d, 0)), (x/(5*cos(c) - 3), True))`**3.42.7 Maxima [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.79

$$\int \frac{1}{-3 + 5 \cos(c + dx)} dx = \frac{\log\left(\frac{2 \sin(dx+c)}{\cos(dx+c)+1} + 1\right) - \log\left(\frac{2 \sin(dx+c)}{\cos(dx+c)+1} - 1\right)}{4d}$$

input `integrate(1/(-3+5*cos(d*x+c)),x, algorithm="maxima")`output `1/4*(log(2*sin(d*x + c)/(cos(d*x + c) + 1) + 1) - log(2*sin(d*x + c)/(cos(d*x + c) + 1) - 1))/d`**3.42.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.60

$$\int \frac{1}{-3 + 5 \cos(c + dx)} dx = \frac{\log\left(\left|2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right|\right) - \log\left(\left|2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1\right|\right)}{4d}$$

input `integrate(1/(-3+5*cos(d*x+c)),x, algorithm="giac")`output `1/4*(log(abs(2*tan(1/2*d*x + 1/2*c) + 1)) - log(abs(2*tan(1/2*d*x + 1/2*c) - 1)))/d`

**3.42.9 Mupad [B] (verification not implemented)**

Time = 14.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.27

$$\int \frac{1}{-3 + 5 \cos(c + dx)} dx = \frac{\operatorname{atanh}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{2d}$$

input `int(1/(5*cos(c + d*x) - 3),x)`

output `atanh(2*tan(c/2 + (d*x)/2))/(2*d)`

### 3.43 $\int \frac{1}{(-3+5 \cos(c+dx))^2} dx$

3.43.1	Optimal result . . . . .	284
3.43.2	Mathematica [A] (verified) . . . . .	284
3.43.3	Rubi [A] (verified) . . . . .	285
3.43.4	Maple [A] (verified) . . . . .	287
3.43.5	Fricas [A] (verification not implemented) . . . . .	287
3.43.6	Sympy [B] (verification not implemented) . . . . .	288
3.43.7	Maxima [A] (verification not implemented) . . . . .	288
3.43.8	Giac [A] (verification not implemented) . . . . .	289
3.43.9	Mupad [B] (verification not implemented) . . . . .	289

#### 3.43.1 Optimal result

Integrand size = 12, antiderivative size = 88

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^2} dx = -\frac{3 \log(\cos(\frac{1}{2}(c + dx)) - 2 \sin(\frac{1}{2}(c + dx)))}{64d} + \frac{3 \log(\cos(\frac{1}{2}(c + dx)) + 2 \sin(\frac{1}{2}(c + dx)))}{64d} - \frac{5 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))}$$

output `-3/64*ln(cos(1/2*d*x+1/2*c)-2*sin(1/2*d*x+1/2*c))/d+3/64*ln(cos(1/2*d*x+1/2*c)+2*sin(1/2*d*x+1/2*c))/d-5/16*sin(d*x+c)/d/(3-5*cos(d*x+c))`

#### 3.43.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.58

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^2} dx = \frac{9 \log(\cos(\frac{1}{2}(c + dx)) - 2 \sin(\frac{1}{2}(c + dx))) - 15 \cos(c + dx) (\log(\cos(\frac{1}{2}(c + dx)) - 2 \sin(\frac{1}{2}(c + dx))) - \log(\cos(\frac{1}{2}(c + dx)) + 2 \sin(\frac{1}{2}(c + dx))))}{64d(-3 + 5 \cos(c + dx))}$$

input `Integrate[(-3 + 5*Cos[c + d*x])^(-2), x]`

output  $(9*\text{Log}[\text{Cos}[(c + d*x)/2] - 2*\text{Sin}[(c + d*x)/2]] - 15*\text{Cos}[c + d*x]*(\text{Log}[\text{Cos}[(c + d*x)/2] - 2*\text{Sin}[(c + d*x)/2]] - \text{Log}[\text{Cos}[(c + d*x)/2] + 2*\text{Sin}[(c + d*x)/2]]) - 9*\text{Log}[\text{Cos}[(c + d*x)/2] + 2*\text{Sin}[(c + d*x)/2]] + 20*\text{Sin}[c + d*x])/(64*d*(-3 + 5*\text{Cos}[c + d*x]))$

### 3.43.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.52, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 3143, 27, 3042, 3138, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(5 \cos(c + dx) - 3)^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{(5 \sin(c + dx + \frac{\pi}{2}) - 3)^2} dx \\ & \quad \downarrow \text{3143} \\ & \frac{1}{16} \int -\frac{3}{3 - 5 \cos(c + dx)} dx - \frac{5 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))} \\ & \quad \downarrow \text{27} \\ & -\frac{3}{16} \int \frac{1}{3 - 5 \cos(c + dx)} dx - \frac{5 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))} \\ & \quad \downarrow \text{3042} \\ & -\frac{3}{16} \int \frac{1}{3 - 5 \sin(c + dx + \frac{\pi}{2})} dx - \frac{5 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))} \\ & \quad \downarrow \text{3138} \\ & \frac{3 \int \frac{1}{8 \tan^2(\frac{1}{2}(c + dx)) - 2} d \tan(\frac{1}{2}(c + dx))}{8d} - \frac{5 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))} \\ & \quad \downarrow \text{220} \\ & \frac{3 \text{arctanh}(2 \tan(\frac{1}{2}(c + dx)))}{32d} - \frac{5 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))} \end{aligned}$$

input `Int[(-3 + 5*Cos[c + d*x])^(-2),x]`

output `(3*ArcTanh[2*Tan[(c + d*x)/2]]/(32*d) - (5*Sin[c + d*x])/(16*d*(3 - 5*Cos[c + d*x]))`

### 3.43.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3143 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

### 3.43.4 Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.81

method	result
norman	$-\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{16d\left(4\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)} - \frac{3 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{64d} + \frac{3 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{64d}$
derivativedivides	$\frac{-\frac{5}{64\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)} + \frac{3 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{64} - \frac{5}{64\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} - \frac{3 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{64}}{d}$
default	$\frac{-\frac{5}{64\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)} + \frac{3 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{64} - \frac{5}{64\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} - \frac{3 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{64}}{d}$
risch	$-\frac{i\left(3 e^{i(dx+c)} - 5\right)}{8d\left(5 e^{2i(dx+c)} - 6 e^{i(dx+c)} + 5\right)} + \frac{3 \ln\left(e^{i(dx+c)} - \frac{3}{5} + \frac{4i}{5}\right)}{64d} - \frac{3 \ln\left(e^{i(dx+c)} - \frac{3}{5} - \frac{4i}{5}\right)}{64d}$
parallelrisch	$\frac{-15 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) \cos(dx+c) + 15 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) \cos(dx+c) + 20 \sin(dx+c) + 9 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) - 9 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{64d(-3+5 \cos(dx+c))}$

input `int(1/(-3+5*cos(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `-5/16/d*tan(1/2*d*x+1/2*c)/(4*tan(1/2*d*x+1/2*c)^2-1)-3/64/d*ln(2*tan(1/2*d*x+1/2*c)-1)+3/64/d*ln(2*tan(1/2*d*x+1/2*c)+1)`

### 3.43.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^2} dx$$

$$= \frac{3(5 \cos(dx + c) - 3) \log\left(-\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c) + \frac{5}{2}\right) - 3(5 \cos(dx + c) - 3) \log\left(-\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c) - \frac{5}{2}\right) + 40 \sin(dx + c)}{128(5d \cos(dx + c) - 3d)}$$

input `integrate(1/(-3+5*cos(d*x+c))^2,x, algorithm="fracas")`

output `1/128*(3*(5*cos(d*x + c) - 3)*log(-3/2*cos(d*x + c) + 2*sin(d*x + c) + 5/2) - 3*(5*cos(d*x + c) - 3)*log(-3/2*cos(d*x + c) - 2*sin(d*x + c) + 5/2) + 40*sin(d*x + c))/(5*d*cos(d*x + c) - 3*d)`

---

3.43.  $\int \frac{1}{(-3+5 \cos(c+dx))^2} dx$



### 3.43.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 240 vs. 2(78) = 156.

Time = 0.68 (sec) , antiderivative size = 240, normalized size of antiderivative = 2.73

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^2} dx$$

$$= \begin{cases} \frac{x}{(-3 + 5 \cos(2 \operatorname{atan}(\frac{1}{2})))^2} \\ \frac{x}{(5 \cos(c) - 3)^2} \\ -\frac{12 \log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 1\right) \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{256d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 64d} + \frac{3 \log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 1\right)}{256d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 64d} + \frac{12 \log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 1\right) \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{256d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 64d} - \frac{3 \log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 1\right)}{256d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 64d} \end{cases}$$

input `integrate(1/(-3+5*cos(d*x+c))**2,x)`

output `Piecewise((x/(-3 + 5*cos(2*atan(1/2)))**2, Eq(c, -d*x - 2*atan(1/2)) | Eq(c, -d*x + 2*atan(1/2))), (x/(5*cos(c) - 3)**2, Eq(d, 0)), (-12*log(2*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**2/(256*d*tan(c/2 + d*x/2)**2 - 64*d) + 3*log(2*tan(c/2 + d*x/2) - 1)/(256*d*tan(c/2 + d*x/2)**2 - 64*d) + 12*log(2*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)**2/(256*d*tan(c/2 + d*x/2)**2 - 64*d) - 3*log(2*tan(c/2 + d*x/2) + 1)/(256*d*tan(c/2 + d*x/2)**2 - 64*d) - 20*tan(c/2 + d*x/2)/(256*d*tan(c/2 + d*x/2)**2 - 64*d), True))`

### 3.43.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.07

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^2} dx$$

$$= -\frac{\frac{20 \sin(dx+c)}{\left(\frac{4 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - 1\right)(\cos(dx+c)+1)} - 3 \log\left(\frac{2 \sin(dx+c)}{\cos(dx+c)+1} + 1\right) + 3 \log\left(\frac{2 \sin(dx+c)}{\cos(dx+c)+1} - 1\right)}{64d}$$

input `integrate(1/(-3+5*cos(d*x+c))^2,x, algorithm="maxima")`

output `-1/64*(20*sin(d*x + c)/((4*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 - 1)*(cos(d*x + c) + 1)) - 3*log(2*sin(d*x + c)/(cos(d*x + c) + 1) + 1) + 3*log(2*sin(d*x + c)/(cos(d*x + c) + 1) - 1))/d`

**3.43.8 Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.77

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^2} dx$$

$$= -\frac{\frac{20 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{4 \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 1} - 3 \log(|2 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|) + 3 \log(|2 \tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{64 d}$$

input `integrate(1/(-3+5*cos(d*x+c))^2,x, algorithm="giac")`output `-1/64*(20*tan(1/2*d*x + 1/2*c)/(4*tan(1/2*d*x + 1/2*c)^2 - 1) - 3*log(abs(2*tan(1/2*d*x + 1/2*c) + 1)) + 3*log(abs(2*tan(1/2*d*x + 1/2*c) - 1)))/d`**3.43.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.53

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^2} dx = \frac{3 \operatorname{atanh}(2 \tan(\frac{c}{2} + \frac{dx}{2}))}{32 d} - \frac{5 \tan(\frac{c}{2} + \frac{dx}{2})}{64 d \left( \tan(\frac{c}{2} + \frac{dx}{2})^2 - \frac{1}{4} \right)}$$

input `int(1/(5*cos(c + d*x) - 3)^2,x)`output `(3*atanh(2*tan(c/2 + (d*x)/2)))/(32*d) - (5*tan(c/2 + (d*x)/2))/(64*d*(tan(c/2 + (d*x)/2)^2 - 1/4))`

### 3.44 $\int \frac{1}{(-3+5 \cos(c+dx))^3} dx$

3.44.1	Optimal result . . . . .	290
3.44.2	Mathematica [A] (verified) . . . . .	291
3.44.3	Rubi [A] (verified) . . . . .	291
3.44.4	Maple [A] (verified) . . . . .	294
3.44.5	Fricas [A] (verification not implemented) . . . . .	294
3.44.6	Sympy [B] (verification not implemented) . . . . .	295
3.44.7	Maxima [A] (verification not implemented) . . . . .	295
3.44.8	Giac [A] (verification not implemented) . . . . .	296
3.44.9	Mupad [B] (verification not implemented) . . . . .	296

#### 3.44.1 Optimal result

Integrand size = 12, antiderivative size = 113

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^3} dx = -\frac{43 \log(\cos(\frac{1}{2}(c + dx)) - 2 \sin(\frac{1}{2}(c + dx)))}{2048d} + \frac{43 \log(\cos(\frac{1}{2}(c + dx)) + 2 \sin(\frac{1}{2}(c + dx)))}{2048d} + \frac{5 \sin(c + dx)}{32d(3 - 5 \cos(c + dx))^2} - \frac{45 \sin(c + dx)}{512d(3 - 5 \cos(c + dx))}$$

```
output -43/2048*ln(cos(1/2*d*x+1/2*c)-2*sin(1/2*d*x+1/2*c))/d+43/2048*ln(cos(1/2*d*x+1/2*c)+2*sin(1/2*d*x+1/2*c))/d+5/32*sin(d*x+c)/d/(3-5*cos(d*x+c))^2-45/512*sin(d*x+c)/d/(3-5*cos(d*x+c))
```

### 3.44.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.87

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^3} dx = -\frac{43 \log(\cos(\frac{1}{2}(c + dx)) - 2 \sin(\frac{1}{2}(c + dx)))}{2048d}$$

$$+ \frac{43 \log(\cos(\frac{1}{2}(c + dx)) + 2 \sin(\frac{1}{2}(c + dx)))}{2048d}$$

$$+ \frac{512d (\cos(\frac{1}{2}(c + dx)) - 2 \sin(\frac{1}{2}(c + dx)))^2}{45 \sin(\frac{1}{2}(c + dx))}$$

$$+ \frac{1024d (\cos(\frac{1}{2}(c + dx)) - 2 \sin(\frac{1}{2}(c + dx)))}{5}$$

$$- \frac{512d (\cos(\frac{1}{2}(c + dx)) + 2 \sin(\frac{1}{2}(c + dx)))^2}{45 \sin(\frac{1}{2}(c + dx))}$$

$$+ \frac{1024d (\cos(\frac{1}{2}(c + dx)) + 2 \sin(\frac{1}{2}(c + dx)))}{5}$$

input `Integrate[(-3 + 5*Cos[c + d*x])^(-3),x]`

output `(-43*Log[Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2]]/(2048*d) + (43*Log[Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2]]/(2048*d) + 5/(512*d*(Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2])^2) + (45*Sin[(c + d*x)/2])/(1024*d*(Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2])) - 5/(512*d*(Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2])^2) + (45*Sin[(c + d*x)/2])/(1024*d*(Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2]))`

### 3.44.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.67, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {3042, 3143, 3042, 3233, 27, 3042, 3138, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(5 \cos(c + dx) - 3)^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(5 \sin(c + dx + \frac{\pi}{2}) - 3)^3} dx$$

$$\begin{aligned}
& \downarrow \text{3143} \\
& \frac{1}{32} \int \frac{5 \cos(c + dx) + 6}{(3 - 5 \cos(c + dx))^2} dx + \frac{5 \sin(c + dx)}{32d(3 - 5 \cos(c + dx))^2} \\
& \downarrow \text{3042} \\
& \frac{1}{32} \int \frac{5 \sin(c + dx + \frac{\pi}{2}) + 6}{(3 - 5 \sin(c + dx + \frac{\pi}{2}))^2} dx + \frac{5 \sin(c + dx)}{32d(3 - 5 \cos(c + dx))^2} \\
& \downarrow \text{3233} \\
& \frac{1}{32} \left( \frac{1}{16} \int -\frac{43}{3 - 5 \cos(c + dx)} dx - \frac{45 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))} \right) + \frac{5 \sin(c + dx)}{32d(3 - 5 \cos(c + dx))^2} \\
& \downarrow \text{27} \\
& \frac{1}{32} \left( -\frac{43}{16} \int \frac{1}{3 - 5 \cos(c + dx)} dx - \frac{45 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))} \right) + \frac{5 \sin(c + dx)}{32d(3 - 5 \cos(c + dx))^2} \\
& \downarrow \text{3042} \\
& \frac{1}{32} \left( -\frac{43}{16} \int \frac{1}{3 - 5 \sin(c + dx + \frac{\pi}{2})} dx - \frac{45 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))} \right) + \frac{5 \sin(c + dx)}{32d(3 - 5 \cos(c + dx))^2} \\
& \downarrow \text{3138} \\
& \frac{1}{32} \left( -\frac{43 \int \frac{1}{8 \tan^2(\frac{1}{2}(c+dx))-2} d \tan(\frac{1}{2}(c+dx))}{8d} - \frac{45 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))} \right) + \frac{5 \sin(c + dx)}{32d(3 - 5 \cos(c + dx))^2} \\
& \downarrow \text{220} \\
& \frac{1}{32} \left( \frac{43 \operatorname{arctanh}(2 \tan(\frac{1}{2}(c + dx)))}{32d} - \frac{45 \sin(c + dx)}{16d(3 - 5 \cos(c + dx))} \right) + \frac{5 \sin(c + dx)}{32d(3 - 5 \cos(c + dx))^2}
\end{aligned}$$

input `Int[(-3 + 5*Cos[c + d*x])^(-3), x]`

output `(5*Sin[c + d*x])/(32*d*(3 - 5*Cos[c + d*x])^2) + ((43*ArcTanh[2*Tan[(c + d*x)/2]])/(32*d) - (45*Sin[c + d*x])/(16*d*(3 - 5*Cos[c + d*x]))) / 32`

## 3.44.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3143 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`
- rule 3233 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

### 3.44.4 Maple [A] (verified)

Time = 0.65 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.79

method	result
norman	$\frac{85 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{35 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{128d}}{512d} - \frac{43 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{2048d} + \frac{43 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2048d}$
derivativedivides	$\frac{-\frac{25}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^2} - \frac{35}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)} + \frac{43 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2048} + \frac{25}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{35}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}}{d}$
default	$\frac{-\frac{25}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^2} - \frac{35}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)} + \frac{43 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2048} + \frac{25}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{35}{2048 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}}{d}$
risch	$-\frac{i(215 e^{3i(dx+c)} - 387 e^{2i(dx+c)} + 325 e^{i(dx+c)} - 225)}{256d(5 e^{2i(dx+c)} - 6 e^{i(dx+c)} + 5)^2} + \frac{43 \ln\left(e^{i(dx+c)} - \frac{3}{5} + \frac{4i}{5}\right)}{2048d} - \frac{43 \ln\left(e^{i(dx+c)} - \frac{3}{5} - \frac{4i}{5}\right)}{2048d}$
parallelrisc	$\frac{(-2580 \cos(dx+c) + 1075 \cos(2dx+2c) + 1849) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{1}{2}\right) + (2580 \cos(dx+c) - 1075 \cos(2dx+2c) - 1849) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{1}{2}\right)}{2048d(-25 \cos(2dx+2c) - 43 + 60 \cos(dx+c))}$

input `int(1/(-3+5*cos(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `(85/512/d*tan(1/2*d*x+1/2*c)-35/128/d*tan(1/2*d*x+1/2*c)^3)/(4*tan(1/2*d*x+1/2*c)^2-1)^2-43/2048/d*ln(2*tan(1/2*d*x+1/2*c)-1)+43/2048/d*ln(2*tan(1/2*d*x+1/2*c)+1)`

### 3.44.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.14

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^3} dx$$

$$= \frac{43 (25 \cos(dx + c)^2 - 30 \cos(dx + c) + 9) \log\left(-\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c) + \frac{5}{2}\right) - 43 (25 \cos(dx + c) + 2 \sin(dx + c) + \frac{5}{2}) \log\left(-\frac{3}{2} \cos(dx + c) - 2 \sin(dx + c) + \frac{5}{2}\right) + 40 (45 \cos(dx + c) - 11) \operatorname{si}\left(\frac{5}{2} + \frac{3}{2} \cos(dx + c)\right) - 40 (45 \cos(dx + c) - 11) \operatorname{si}\left(\frac{5}{2} - \frac{3}{2} \cos(dx + c)\right)}{4096 (25 d \cos(dx + c) + 9 d)}$$

input `integrate(1/(-3+5*cos(d*x+c))^3,x, algorithm="fricas")`

output `1/4096*(43*(25*cos(d*x + c)^2 - 30*cos(d*x + c) + 9)*log(-3/2*cos(d*x + c) + 2*sin(d*x + c) + 5/2) - 43*(25*cos(d*x + c)^2 - 30*cos(d*x + c) + 9)*log(-3/2*cos(d*x + c) - 2*sin(d*x + c) + 5/2) + 40*(45*cos(d*x + c) - 11)*si(n(d*x + c))/(25*d*cos(d*x + c)^2 - 30*d*cos(d*x + c) + 9*d)`

---

3.44.  $\int \frac{1}{(-3+5 \cos(c+dx))^3} dx$

### 3.44.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 490 vs.  $2(102) = 204$ .

Time = 1.20 (sec) , antiderivative size = 490, normalized size of antiderivative = 4.34

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^3} dx$$

$$= \begin{cases} \frac{x}{(-3 + 5 \cos(2 \operatorname{atan}(\frac{1}{2})))^3} \\ \frac{x}{(5 \cos(c) - 3)^3} \\ -\frac{688 \log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 1\right) \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right)}{32768d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) - 16384d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 2048d} + \frac{344 \log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 1\right) \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{32768d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) - 16384d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 2048d} - \frac{43 \log\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 1\right)}{32768d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) - 16384d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 2048d} \end{cases}$$

input `integrate(1/(-3+5*cos(d*x+c))**3,x)`

output `Piecewise((x/(-3 + 5*cos(2*atan(1/2)))**3, Eq(c, -d*x - 2*atan(1/2)) | Eq(c, -d*x + 2*atan(1/2))), (x/(5*cos(c) - 3)**3, Eq(d, 0)), (-688*log(2*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**4/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d) + 344*log(2*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**2/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d) - 43*log(2*tan(c/2 + d*x/2) - 1)/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d) + 688*log(2*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)**4/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d) - 344*log(2*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)**2/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d) + 43*log(2*tan(c/2 + d*x/2) + 1)/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d) - 560*tan(c/2 + d*x/2)**3/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d) + 340*tan(c/2 + d*x/2)/(32768*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 2048*d), True))`

### 3.44.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.21

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^3} dx$$

$$= -\frac{20 \left( \frac{17 \sin(dx+c)}{\cos(dx+c)+1} - \frac{28 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right) - 43 \log \left( \frac{2 \sin(dx+c)}{\cos(dx+c)+1} + 1 \right) + 43 \log \left( \frac{2 \sin(dx+c)}{\cos(dx+c)+1} - 1 \right)}{2048 d}$$

---

3.44.  $\int \frac{1}{(-3+5 \cos(c+dx))^3} dx$



input `integrate(1/(-3+5*cos(d*x+c))^3,x, algorithm="maxima")`

output 
$$\frac{-1/2048*(20*(17*\sin(d*x + c)/(\cos(d*x + c) + 1) - 28*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3)/(8*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2 - 16*\sin(d*x + c)^4/(\cos(d*x + c) + 1)^4 - 1) - 43*\log(2*\sin(d*x + c)/(\cos(d*x + c) + 1) + 1) + 43*\log(2*\sin(d*x + c)/(\cos(d*x + c) + 1) - 1))/d}{2048 d}$$

### 3.44.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.74

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^3} dx = \frac{20 \left( 28 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 17 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\left( 4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 1 \right)^2} - 43 \log \left( \left| 2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1 \right| \right) + 43 \log \left( \left| 2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1 \right| \right) - 1}{2048 d}$$

input `integrate(1/(-3+5*cos(d*x+c))^3,x, algorithm="giac")`

output 
$$\frac{-1/2048*(20*(28*\tan(1/2*d*x + 1/2*c)^3 - 17*\tan(1/2*d*x + 1/2*c))/(4*\tan(1/2*d*x + 1/2*c)^2 - 1)^2 - 43*\log(\text{abs}(2*\tan(1/2*d*x + 1/2*c) + 1)) + 43*\log(\text{abs}(2*\tan(1/2*d*x + 1/2*c) - 1)))/d}{2048 d}$$

### 3.44.9 Mupad [B] (verification not implemented)

Time = 14.39 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.66

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^3} dx = \frac{43 \operatorname{atanh}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{1024 d} + \frac{\frac{85 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{8192} - \frac{35 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{2048}}{d \left( \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{2} + \frac{1}{16} \right)}$$

input `int(1/(5*cos(c + d*x) - 3)^3,x)`

output  $(43*\operatorname{atanh}(2*\tan(c/2 + (d*x)/2)))/(1024*d) + ((85*\tan(c/2 + (d*x)/2))/8192 - (35*\tan(c/2 + (d*x)/2)^3)/2048)/(d*(\tan(c/2 + (d*x)/2)^4 - \tan(c/2 + (d*x)/2)^2/2 + 1/16))$

### 3.45 $\int \frac{1}{(-3+5 \cos(c+dx))^4} dx$

3.45.1	Optimal result . . . . .	298
3.45.2	Mathematica [B] (verified) . . . . .	299
3.45.3	Rubi [A] (verified) . . . . .	299
3.45.4	Maple [A] (verified) . . . . .	302
3.45.5	Fricas [A] (verification not implemented) . . . . .	303
3.45.6	Sympy [B] (verification not implemented) . . . . .	303
3.45.7	Maxima [A] (verification not implemented) . . . . .	304
3.45.8	Giac [A] (verification not implemented) . . . . .	305
3.45.9	Mupad [B] (verification not implemented) . . . . .	305

#### 3.45.1 Optimal result

Integrand size = 12, antiderivative size = 138

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^4} dx = -\frac{279 \log(\cos(\frac{1}{2}(c + dx)) - 2 \sin(\frac{1}{2}(c + dx)))}{32768d} + \frac{279 \log(\cos(\frac{1}{2}(c + dx)) + 2 \sin(\frac{1}{2}(c + dx)))}{32768d} - \frac{5 \sin(c + dx)}{48d(3 - 5 \cos(c + dx))^3} + \frac{25 \sin(c + dx)}{512d(3 - 5 \cos(c + dx))^2} - \frac{995 \sin(c + dx)}{24576d(3 - 5 \cos(c + dx))}$$

```
output -279/32768*ln(cos(1/2*d*x+1/2*c)-2*sin(1/2*d*x+1/2*c))/d+279/32768*ln(cos(
1/2*d*x+1/2*c)+2*sin(1/2*d*x+1/2*c))/d-5/48*sin(d*x+c)/d/(3-5*cos(d*x+c))^
3+25/512*sin(d*x+c)/d/(3-5*cos(d*x+c))^2-995/24576*sin(d*x+c)/d/(3-5*cos(d
*x+c))
```

### 3.45.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 288 vs.  $2(138) = 276$ .

Time = 0.01 (sec) , antiderivative size = 288, normalized size of antiderivative = 2.09

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^4} dx$$

$$= \frac{467046 \log\left(\cos\left(\frac{1}{2}(c + dx)\right) - 2 \sin\left(\frac{1}{2}(c + dx)\right)\right) - 104625 \cos(3(c + dx)) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) - 2 \sin\left(\frac{1}{2}(c + dx)\right)\right) - 2 \sin\left(\frac{1}{2}(c + dx)\right) - 104625 \cos(3(c + dx)) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + 2 \sin\left(\frac{1}{2}(c + dx)\right)\right) + 376650 \cos(2(c + dx)) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) - 2 \sin\left(\frac{1}{2}(c + dx)\right)\right) - \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + 2 \sin\left(\frac{1}{2}(c + dx)\right)\right) - 467046 \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + 2 \sin\left(\frac{1}{2}(c + dx)\right)\right) + 104625 \cos(3(c + dx)) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + 2 \sin\left(\frac{1}{2}(c + dx)\right)\right) + 226140 \sin(c + dx) - 190800 \sin(2(c + dx)) + 99500 \sin(3(c + dx))}{(393216 d (-3 + 5 \cos(c + dx))^3)}$$

input `Integrate[(-3 + 5*Cos[c + d*x])^(-4), x]`

output `(467046*Log[Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2]] - 104625*Cos[3*(c + d*x)]*Log[Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2]] - 765855*Cos[c + d*x]*(Log[Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2]] - Log[Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2]]) + 376650*Cos[2*(c + d*x)]*(Log[Cos[(c + d*x)/2] - 2*Sin[(c + d*x)/2]] - Log[Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2]]) - 467046*Log[Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2]] + 104625*Cos[3*(c + d*x)]*Log[Cos[(c + d*x)/2] + 2*Sin[(c + d*x)/2]] + 226140*Sin[c + d*x] - 190800*Sin[2*(c + d*x)] + 99500*Sin[3*(c + d*x)]/(393216*d*(-3 + 5*Cos[c + d*x])^3)`

### 3.45.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.77, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {3042, 3143, 25, 3042, 3233, 25, 3042, 3233, 27, 3042, 3138, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(5 \cos(c + dx) - 3)^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(5 \sin(c + dx + \frac{\pi}{2}) - 3)^4} dx$$

$$\downarrow \text{3143}$$

$$\frac{1}{48} \int -\frac{10 \cos(c + dx) + 9}{(3 - 5 \cos(c + dx))^3} dx - \frac{5 \sin(c + dx)}{48d(3 - 5 \cos(c + dx))^3}$$

$$\begin{aligned}
& \downarrow 25 \\
& -\frac{1}{48} \int \frac{10 \cos(c+dx) + 9}{(3-5 \cos(c+dx))^3} dx - \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 3042 \\
& -\frac{1}{48} \int \frac{10 \sin(c+dx + \frac{\pi}{2}) + 9}{(3-5 \sin(c+dx + \frac{\pi}{2}))^3} dx - \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 3233 \\
& \frac{1}{48} \left( \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} - \frac{1}{32} \int -\frac{75 \cos(c+dx) + 154}{(3-5 \cos(c+dx))^2} dx \right) - \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 25 \\
& \frac{1}{48} \left( \frac{1}{32} \int \frac{75 \cos(c+dx) + 154}{(3-5 \cos(c+dx))^2} dx + \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \right) - \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left( \frac{1}{32} \int \frac{75 \sin(c+dx + \frac{\pi}{2}) + 154}{(3-5 \sin(c+dx + \frac{\pi}{2}))^2} dx + \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \right) - \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 3233 \\
& \frac{1}{48} \left( \frac{1}{32} \left( \frac{1}{16} \int -\frac{837}{3-5 \cos(c+dx)} dx - \frac{995 \sin(c+dx)}{16d(3-5 \cos(c+dx))} \right) + \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \right) - \\
& \quad \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 27 \\
& \frac{1}{48} \left( \frac{1}{32} \left( -\frac{837}{16} \int \frac{1}{3-5 \cos(c+dx)} dx - \frac{995 \sin(c+dx)}{16d(3-5 \cos(c+dx))} \right) + \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \right) - \\
& \quad \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left( \frac{1}{32} \left( -\frac{837}{16} \int \frac{1}{3-5 \sin(c+dx + \frac{\pi}{2})} dx - \frac{995 \sin(c+dx)}{16d(3-5 \cos(c+dx))} \right) + \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \right) - \\
& \quad \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3} \\
& \downarrow 3138
\end{aligned}$$

---

3.45.  $\int \frac{1}{(-3+5 \cos(c+dx))^4} dx$

$$\frac{1}{48} \left( \frac{1}{32} \left( -\frac{837 \int \frac{1}{8 \tan^2(\frac{1}{2}(c+dx))-2} d \tan(\frac{1}{2}(c+dx))}{8d} - \frac{995 \sin(c+dx)}{16d(3-5 \cos(c+dx))} \right) + \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \right) - \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3}$$

↓ 220

$$\frac{1}{48} \left( \frac{1}{32} \left( \frac{837 \operatorname{arctanh}(2 \tan(\frac{1}{2}(c+dx)))}{32d} - \frac{995 \sin(c+dx)}{16d(3-5 \cos(c+dx))} \right) + \frac{75 \sin(c+dx)}{32d(3-5 \cos(c+dx))^2} \right) - \frac{5 \sin(c+dx)}{48d(3-5 \cos(c+dx))^3}$$

input `Int[(-3 + 5*Cos[c + d*x])^(-4), x]`

output `(-5*Sin[c + d*x])/(48*d*(3 - 5*Cos[c + d*x])^3) + ((75*Sin[c + d*x])/(32*d*(3 - 5*Cos[c + d*x])^2) + ((837*ArcTanh[2*Tan[(c + d*x)/2]])/(32*d) - (99*5*Sin[c + d*x])/(16*d*(3 - 5*Cos[c + d*x]))) / 32) / 48`

### 3.45.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3233 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

### 3.45.4 Maple [A] (verified)

Time = 0.71 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.76

method	result
norman	$\frac{-\frac{745 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8192d} + \frac{265 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{768d} - \frac{295 \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{512d}}{\left(4 \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)^3} - \frac{279 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{32768d} + \frac{279 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{32768d}$
risch	$\frac{i(20925 e^{5i(dx+c)} - 62775 e^{4i(dx+c)} + 111042 e^{3i(dx+c)} - 119310 e^{2i(dx+c)} + 68625 e^{i(dx+c)} - 24875)}{12288d(5 e^{2i(dx+c)} - 6 e^{i(dx+c)} + 5)^3} + \frac{279 \ln(e^{i(dx+c)})}{32768d}$
derivativedivides	$\frac{-\frac{125}{49152 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^3} + \frac{25}{8192 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{295}{32768 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} - \frac{279 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{32768}}{d} - \frac{125}{49152 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^3} + \frac{25}{8192 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{295}{32768 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} - \frac{279 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{32768}}{d}$
default	$\frac{-\frac{125}{49152 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^3} + \frac{25}{8192 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{295}{32768 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} - \frac{279 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{32768}}{d} - \frac{125}{49152 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^3} + \frac{25}{8192 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{295}{32768 \left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} - \frac{279 \ln\left(2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{32768}}{d}$
parallelrisch	$\frac{(-765855 \cos(dx+c) + 376650 \cos(2dx+2c) - 104625 \cos(3dx+3c) + 467046) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{1}{2}\right) + (765855 \cos(dx+c) - 376650 \cos(2dx+2c) + 104625 \cos(3dx+3c) - 467046) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{1}{2}\right)}{98304d(-558 + 125 \cos(3dx+c))}$

input `int(1/(-3+5*cos(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `(-745/8192/d*tan(1/2*d*x+1/2*c)+265/768/d*tan(1/2*d*x+1/2*c)^3-295/512/d*tan(1/2*d*x+1/2*c)^5)/(4*tan(1/2*d*x+1/2*c)^2-1)^3-279/32768/d*ln(2*tan(1/2*d*x+1/2*c)-1)+279/32768/d*ln(2*tan(1/2*d*x+1/2*c)+1)`

3.45.  $\int \frac{1}{(-3+5 \cos(c+dx))^4} dx$

**3.45.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.23

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^4} dx$$

$$= \frac{837 (125 \cos(dx + c)^3 - 225 \cos(dx + c)^2 + 135 \cos(dx + c) - 27) \log\left(-\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c)\right)}{196608}$$

input `integrate(1/(-3+5*cos(d*x+c))^4,x, algorithm="fracas")`

output `1/196608*(837*(125*cos(d*x + c)^3 - 225*cos(d*x + c)^2 + 135*cos(d*x + c) - 27)*log(-3/2*cos(d*x + c) + 2*sin(d*x + c) + 5/2) - 837*(125*cos(d*x + c)^3 - 225*cos(d*x + c)^2 + 135*cos(d*x + c) - 27)*log(-3/2*cos(d*x + c) - 2*sin(d*x + c) + 5/2) + 40*(4975*cos(d*x + c)^2 - 4770*cos(d*x + c) + 1583)*sin(d*x + c))/(125*d*cos(d*x + c)^3 - 225*d*cos(d*x + c)^2 + 135*d*cos(d*x + c) - 27*d)`

**3.45.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 831 vs.  $2(126) = 252$ .

Time = 2.48 (sec) , antiderivative size = 831, normalized size of antiderivative = 6.02

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^4} dx = \text{Too large to display}$$

input `integrate(1/(-3+5*cos(d*x+c))**4,x)`



output `Piecewise((x/(-3 + 5*cos(2*atan(1/2)))**4, Eq(c, -d*x - 2*atan(1/2)) | Eq(c, -d*x + 2*atan(1/2))), (x/(5*cos(c) - 3)**4, Eq(d, 0)), (-53568*log(2*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**6/(6291456*d*tan(c/2 + d*x/2)**6 - 4718592*d*tan(c/2 + d*x/2)**4 + 1179648*d*tan(c/2 + d*x/2)**2 - 98304*d) + 40176*log(2*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**4/(6291456*d*tan(c/2 + d*x/2)**6 - 4718592*d*tan(c/2 + d*x/2)**4 + 1179648*d*tan(c/2 + d*x/2)**2 - 98304*d) - 10044*log(2*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**2/(6291456*d*tan(c/2 + d*x/2)**6 - 4718592*d*tan(c/2 + d*x/2)**4 + 1179648*d*tan(c/2 + d*x/2)**2 - 98304*d) + 837*log(2*tan(c/2 + d*x/2) - 1)/(6291456*d*tan(c/2 + d*x/2)**6 - 4718592*d*tan(c/2 + d*x/2)**4 + 1179648*d*tan(c/2 + d*x/2)**2 - 98304*d) + 53568*log(2*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)**6/(6291456*d*tan(c/2 + d*x/2)**6 - 4718592*d*tan(c/2 + d*x/2)**4 + 1179648*d*tan(c/2 + d*x/2)**2 - 98304*d) - 40176*log(2*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)**4/(6291456*d*tan(c/2 + d*x/2)**6 - 4718592*d*tan(c/2 + d*x/2)**4 + 1179648*d*tan(c/2 + d*x/2)**2 - 98304*d) + 10044*log(2*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)**2/(6291456*d*tan(c/2 + d*x/2)**6 - 4718592*d*tan(c/2 + d*x/2)**4 + 1179648*d*tan(c/2 + d*x/2)**2 - 98304*d) - 837*log(2*tan(c/2 + d*x/2) + 1)/(6291456*d*tan(c/2 + d*x/2)**6 - 4718592*d*tan(c/2 + d*x/2)**4 + 1179648*d*tan(c/2 + d*x/2)**2 - 98304*d) - 56640*tan(c/2 + d*x/2)**5/(6291456*d*tan(c/2 + d*x/2)**6 - 4718592*d*tan(c/2 + d*x/2)**4 + ...`

### 3.45.7 Maxima [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.28

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^4} dx = \frac{20 \left( \frac{447 \sin(dx+c)}{\cos(dx+c)+1} - \frac{1696 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{2832 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right) - \frac{12 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - \frac{48 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{64 \sin(dx+c)^6}{(\cos(dx+c)+1)^6} - 1}{98304 d} - 837 \log \left( \frac{2 \sin(dx+c)}{\cos(dx+c)+1} + 1 \right) + 837 \log \left( \frac{2 \sin(dx+c)}{\cos(dx+c)+1} - 1 \right)$$

input `integrate(1/(-3+5*cos(d*x+c))^4,x, algorithm="maxima")`

output `-1/98304*(20*(447*sin(d*x + c)/(cos(d*x + c) + 1) - 1696*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 2832*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/(12*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 - 48*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 64*sin(d*x + c)^6/(cos(d*x + c) + 1)^6 - 1) - 837*log(2*sin(d*x + c)/(cos(d*x + c) + 1) + 1) + 837*log(2*sin(d*x + c)/(cos(d*x + c) + 1) - 1))/d`

---

3.45.  $\int \frac{1}{(-3+5 \cos(c+dx))^4} dx$

**3.45.8 Giac [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.70

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^4} dx = \frac{20 \left( 2832 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 1696 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 447 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\left( 4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 1 \right)^3} - 837 \log \left( \left| 2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1 \right| \right) + 837 \log \left( \left| 2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1 \right| \right)$$


---

98304 d

input `integrate(1/(-3+5*cos(d*x+c))^4,x, algorithm="giac")`output `-1/98304*(20*(2832*tan(1/2*d*x + 1/2*c)^5 - 1696*tan(1/2*d*x + 1/2*c)^3 + 447*tan(1/2*d*x + 1/2*c))/(4*tan(1/2*d*x + 1/2*c)^2 - 1)^3 - 837*log(abs(2*tan(1/2*d*x + 1/2*c) + 1)) + 837*log(abs(2*tan(1/2*d*x + 1/2*c) - 1)))/d`**3.45.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.74

$$\int \frac{1}{(-3 + 5 \cos(c + dx))^4} dx = \frac{279 \operatorname{atanh}\left(2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{16384 d} - \frac{\frac{295 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{32768} - \frac{265 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{49152} + \frac{745 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{524288}}{d \left( \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - \frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{4} + \frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{16} - \frac{1}{64} \right)}$$

input `int(1/(5*cos(c + d*x) - 3)^4,x)`output `(279*atanh(2*tan(c/2 + (d*x)/2)))/(16384*d) - ((745*tan(c/2 + (d*x)/2))/524288 - (265*tan(c/2 + (d*x)/2)^3)/49152 + (295*tan(c/2 + (d*x)/2)^5)/32768)/(d*((3*tan(c/2 + (d*x)/2)^2)/16 - (3*tan(c/2 + (d*x)/2)^4)/4 + tan(c/2 + (d*x)/2)^6 - 1/64))`

### 3.46 $\int \frac{1}{-3-5\cos(c+dx)} dx$

3.46.1	Optimal result . . . . .	306
3.46.2	Mathematica [A] (verified) . . . . .	306
3.46.3	Rubi [A] (verified) . . . . .	307
3.46.4	Maple [A] (verified) . . . . .	308
3.46.5	Fricas [A] (verification not implemented) . . . . .	308
3.46.6	Sympy [A] (verification not implemented) . . . . .	309
3.46.7	Maxima [A] (verification not implemented) . . . . .	309
3.46.8	Giac [A] (verification not implemented) . . . . .	309
3.46.9	Mupad [B] (verification not implemented) . . . . .	310

#### 3.46.1 Optimal result

Integrand size = 12, antiderivative size = 65

$$\int \frac{1}{-3-5\cos(c+dx)} dx = \frac{\log\left(2\cos\left(\frac{1}{2}(c+dx)\right) - \sin\left(\frac{1}{2}(c+dx)\right)\right)}{4d} - \frac{\log\left(2\cos\left(\frac{1}{2}(c+dx)\right) + \sin\left(\frac{1}{2}(c+dx)\right)\right)}{4d}$$

output `1/4*ln(2*cos(1/2*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d-1/4*ln(2*cos(1/2*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d`

#### 3.46.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00

$$\int \frac{1}{-3-5\cos(c+dx)} dx = \frac{\log\left(2\cos\left(\frac{1}{2}(c+dx)\right) - \sin\left(\frac{1}{2}(c+dx)\right)\right)}{4d} - \frac{\log\left(2\cos\left(\frac{1}{2}(c+dx)\right) + \sin\left(\frac{1}{2}(c+dx)\right)\right)}{4d}$$

input `Integrate[(-3 - 5*Cos[c + d*x])^(-1),x]`

output `Log[2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]]/(4*d) - Log[2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]/(4*d)`

**3.46.3 Rubi [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.34, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {3042, 3138, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{-5 \cos(c + dx) - 3} dx$$

↓ 3042

$$\int \frac{1}{-5 \sin(c + dx + \frac{\pi}{2}) - 3} dx$$

↓ 3138

$$2 \int \frac{1}{2 \tan^2(\frac{1}{2}(c+dx)) - 8} d \tan(\frac{1}{2}(c + dx))$$

↓ 220

$$\frac{\operatorname{arctanh}(\frac{1}{2} \tan(\frac{1}{2}(c + dx)))}{2d}$$

input `Int[(-3 - 5*Cos[c + d*x])^(-1), x]`

output `-1/2*ArcTanh[Tan[(c + d*x)/2]/2]/d`

**3.46.3.1 Defintions of rubi rules used**

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3138 Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{
e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b +
(a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x]
&& NeQ[a^2 - b^2, 0]
```

### 3.46.4 Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.51

method	result	size
parallelrisc	$\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right) - \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{4d}$	33
derivativedivides	$-\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{4} + \frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{4}$ $d$	34
default	$-\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{4} + \frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{4}$ $d$	34
norman	$\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{4d} - \frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{4d}$	36
risc	$\frac{\ln\left(e^{i(dx+c)} + \frac{3}{5} - \frac{4i}{5}\right)}{4d} - \frac{\ln\left(e^{i(dx+c)} + \frac{3}{5} + \frac{4i}{5}\right)}{4d}$	40

```
input int(1/(-3-5*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/4*(ln(tan(1/2*d*x+1/2*c)-2)-ln(tan(1/2*d*x+1/2*c)+2))/d
```

### 3.46.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.71

$$\int \frac{1}{-3 - 5 \cos(c + dx)} dx$$

$$= -\frac{\log\left(\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c) + \frac{5}{2}\right) - \log\left(\frac{3}{2} \cos(dx + c) - 2 \sin(dx + c) + \frac{5}{2}\right)}{8d}$$

```
input integrate(1/(-3-5*cos(d*x+c)),x, algorithm="fricas")
```

```
output -1/8*(log(3/2*cos(d*x + c) + 2*sin(d*x + c) + 5/2) - log(3/2*cos(d*x + c)
- 2*sin(d*x + c) + 5/2))/d
```

---

3.46.  $\int \frac{1}{-3-5\cos(c+dx)} dx$

**3.46.6 Sympy [A] (verification not implemented)**

Time = 0.38 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.65

$$\int \frac{1}{-3 - 5 \cos(c + dx)} dx = \begin{cases} \frac{\log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 2\right)}{4d} - \frac{\log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 2\right)}{4d} & \text{for } d \neq 0 \\ \frac{x}{-5 \cos(c) - 3} & \text{otherwise} \end{cases}$$

input `integrate(1/(-3-5*cos(d*x+c)),x)`output `Piecewise((log(tan(c/2 + d*x/2) - 2)/(4*d) - log(tan(c/2 + d*x/2) + 2)/(4*d), Ne(d, 0)), (x/(-5*cos(c) - 3), True))`**3.46.7 Maxima [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.74

$$\int \frac{1}{-3 - 5 \cos(c + dx)} dx = -\frac{\log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 2\right) - \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 2\right)}{4d}$$

input `integrate(1/(-3-5*cos(d*x+c)),x, algorithm="maxima")`output `-1/4*(log(sin(d*x + c)/(cos(d*x + c) + 1) + 2) - log(sin(d*x + c)/(cos(d*x + c) + 1) - 2))/d`**3.46.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.52

$$\int \frac{1}{-3 - 5 \cos(c + dx)} dx = -\frac{\log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 2\right|\right) - \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 2\right|\right)}{4d}$$

input `integrate(1/(-3-5*cos(d*x+c)),x, algorithm="giac")`output `-1/4*(log(abs(tan(1/2*d*x + 1/2*c) + 2)) - log(abs(tan(1/2*d*x + 1/2*c) - 2)))/d`

**3.46.9 Mupad [B] (verification not implemented)**

Time = 13.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.26

$$\int \frac{1}{-3 - 5 \cos(c + dx)} dx = -\frac{\operatorname{atanh}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{2d}$$

input `int(-1/(5*cos(c + d*x) + 3),x)`

output `-atanh(tan(c/2 + (d*x)/2)/2)/(2*d)`

### 3.47 $\int \frac{1}{(-3-5 \cos(c+dx))^2} dx$

3.47.1	Optimal result . . . . .	311
3.47.2	Mathematica [A] (verified) . . . . .	311
3.47.3	Rubi [A] (verified) . . . . .	312
3.47.4	Maple [A] (verified) . . . . .	314
3.47.5	Fricas [A] (verification not implemented) . . . . .	314
3.47.6	Sympy [B] (verification not implemented) . . . . .	315
3.47.7	Maxima [A] (verification not implemented) . . . . .	315
3.47.8	Giac [A] (verification not implemented) . . . . .	316
3.47.9	Mupad [B] (verification not implemented) . . . . .	316

#### 3.47.1 Optimal result

Integrand size = 12, antiderivative size = 90

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^2} dx = \frac{3 \log(2 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}{64d} - \frac{3 \log(2 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{64d} + \frac{5 \sin(c + dx)}{16d(3 + 5 \cos(c + dx))}$$

```
output 3/64*ln(2*cos(1/2*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d-3/64*ln(2*cos(1/2*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d+5/16*sin(d*x+c)/d/(3+5*cos(d*x+c))
```

#### 3.47.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.59

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^2} dx = \frac{9 \log(2 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) + 15 \cos(c + dx) (\log(2 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) - \log(2 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))))}{64d(3 + 5 \cos(c + dx))}$$

```
input Integrate[(-3 - 5*Cos[c + d*x])^(-2), x]
```



output  $(9*\text{Log}[2*\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2]] + 15*\text{Cos}[c + d*x]*(\text{Log}[2*\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2]] - \text{Log}[2*\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]]) - 9*\text{Log}[2*\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]] + 20*\text{Sin}[c + d*x])/(64*d*(3 + 5*\text{Cos}[c + d*x]))$

### 3.47.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.53, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 3143, 27, 3042, 3138, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(-5 \cos(c + dx) - 3)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(-5 \sin(c + dx + \frac{\pi}{2}) - 3)^2} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{1}{16} \int -\frac{3}{5 \cos(c + dx) + 3} dx + \frac{5 \sin(c + dx)}{16d(5 \cos(c + dx) + 3)} \\
 & \quad \downarrow \text{27} \\
 & \frac{5 \sin(c + dx)}{16d(5 \cos(c + dx) + 3)} - \frac{3}{16} \int \frac{1}{5 \cos(c + dx) + 3} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{5 \sin(c + dx)}{16d(5 \cos(c + dx) + 3)} - \frac{3}{16} \int \frac{1}{5 \sin(c + dx + \frac{\pi}{2}) + 3} dx \\
 & \quad \downarrow \text{3138} \\
 & \frac{5 \sin(c + dx)}{16d(5 \cos(c + dx) + 3)} - \frac{3 \int \frac{1}{8 - 2 \tan^2(\frac{1}{2}(c + dx))} d \tan(\frac{1}{2}(c + dx))}{8d} \\
 & \quad \downarrow \text{219} \\
 & \frac{5 \sin(c + dx)}{16d(5 \cos(c + dx) + 3)} - \frac{3 \operatorname{arctanh}(\frac{1}{2} \tan(\frac{1}{2}(c + dx)))}{32d}
 \end{aligned}$$

input `Int[(-3 - 5*Cos[c + d*x])^(-2), x]`

output `(-3*ArcTanh[Tan[(c + d*x)/2]/2])/(32*d) + (5*Sin[c + d*x])/(16*d*(3 + 5*Cos[c + d*x]))`

### 3.47.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3143 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

### 3.47.4 Maple [A] (verified)

Time = 0.78 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.71

method	result
derivativedivides	$\frac{-\frac{5}{32(\tan(\frac{dx}{2} + \frac{c}{2}) + 2)} - \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 2)}{64}}{d} - \frac{5}{32(\tan(\frac{dx}{2} + \frac{c}{2}) - 2)} + \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 2)}{64}$
default	$\frac{-\frac{5}{32(\tan(\frac{dx}{2} + \frac{c}{2}) + 2)} - \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 2)}{64}}{d} - \frac{5}{32(\tan(\frac{dx}{2} + \frac{c}{2}) - 2)} + \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 2)}{64}$
norman	$-\frac{5 \tan(\frac{dx}{2} + \frac{c}{2})}{16d(\tan^2(\frac{dx}{2} + \frac{c}{2}) - 4)} + \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 2)}{64d} - \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 2)}{64d}$
risch	$\frac{i(3e^{i(dx+c)} + 5)}{8d(5e^{2i(dx+c)} + 6e^{i(dx+c)} + 5)} - \frac{3 \ln(e^{i(dx+c)} + \frac{3}{5} + \frac{4i}{5})}{64d} + \frac{3 \ln(e^{i(dx+c)} + \frac{3}{5} - \frac{4i}{5})}{64d}$
parallelrisch	$\frac{15 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 2) \cos(dx+c) - 15 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 2) \cos(dx+c) + 20 \sin(dx+c) + 9 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 2) - 9 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 2)}{64d(3+5 \cos(dx+c))}$

input `int(1/(-3-5*cos(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(-5/32/(tan(1/2*d*x+1/2*c)+2)-3/64*ln(tan(1/2*d*x+1/2*c)+2)-5/32/(tan(1/2*d*x+1/2*c)-2)+3/64*ln(tan(1/2*d*x+1/2*c)-2))`

### 3.47.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.98

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^2} dx = \frac{-3(5 \cos(dx + c) + 3) \log\left(\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c) + \frac{5}{2}\right) - 3(5 \cos(dx + c) + 3) \log\left(\frac{3}{2} \cos(dx + c) - 2 \sin(dx + c) + \frac{5}{2}\right) + 40 \sin(dx + c)}{128(5d \cos(dx + c) + 3d)}$$

input `integrate(1/(-3-5*cos(d*x+c))^2,x, algorithm="fracas")`

output `-1/128*(3*(5*cos(d*x + c) + 3)*log(3/2*cos(d*x + c) + 2*sin(d*x + c) + 5/2) - 3*(5*cos(d*x + c) + 3)*log(3/2*cos(d*x + c) - 2*sin(d*x + c) + 5/2) - 40*sin(d*x + c))/(5*d*cos(d*x + c) + 3*d)`

---

3.47.  $\int \frac{1}{(-3-5 \cos(c+dx))^2} dx$

**3.47.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 231 vs. 2(78) = 156.

Time = 0.68 (sec) , antiderivative size = 231, normalized size of antiderivative = 2.57

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^2} dx$$

$$= \begin{cases} \frac{x}{(-3 - 5 \cos(2 \operatorname{atan}(2)))^2} \\ \frac{x}{(-5 \cos(c) - 3)^2} \\ \frac{3 \log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 2\right) \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{64d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 256d} - \frac{12 \log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 2\right)}{64d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 256d} - \frac{3 \log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 2\right) \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{64d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 256d} + \frac{12 \log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 2\right)}{64d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 256d} \end{cases}$$

input `integrate(1/(-3-5*cos(d*x+c))**2,x)`

output `Piecewise((x/(-3 - 5*cos(2*atan(2)))**2, Eq(c, -d*x - 2*atan(2)) | Eq(c, -d*x + 2*atan(2))), (x/(-5*cos(c) - 3)**2, Eq(d, 0)), (3*log(tan(c/2 + d*x/2) - 2)*tan(c/2 + d*x/2)**2/(64*d*tan(c/2 + d*x/2)**2 - 256*d) - 12*log(tan(c/2 + d*x/2) - 2)/(64*d*tan(c/2 + d*x/2)**2 - 256*d) - 3*log(tan(c/2 + d*x/2) + 2)*tan(c/2 + d*x/2)**2/(64*d*tan(c/2 + d*x/2)**2 - 256*d) + 12*log(tan(c/2 + d*x/2) + 2)/(64*d*tan(c/2 + d*x/2)**2 - 256*d) - 20*tan(c/2 + d*x/2)/(64*d*tan(c/2 + d*x/2)**2 - 256*d), True))`

**3.47.7 Maxima [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.01

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^2} dx$$

$$= -\frac{\frac{20 \sin(dx+c)}{\left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} - 4\right)(\cos(dx+c)+1)}}{64d} + 3 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 2\right) - 3 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 2\right)$$

input `integrate(1/(-3-5*cos(d*x+c))^2,x, algorithm="maxima")`

output `-1/64*(20*sin(d*x + c)/((sin(d*x + c)^2/(cos(d*x + c) + 1)^2 - 4)*(cos(d*x + c) + 1)) + 3*log(sin(d*x + c)/(cos(d*x + c) + 1) + 2) - 3*log(sin(d*x + c)/(cos(d*x + c) + 1) - 2))/d`

**3.47.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.69

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^2} dx$$

$$= -\frac{\frac{20 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 4} + 3 \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 2|) - 3 \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 2|)}{64 d}$$

input `integrate(1/(-3-5*cos(d*x+c))^2,x, algorithm="giac")`output `-1/64*(20*tan(1/2*d*x + 1/2*c)/(tan(1/2*d*x + 1/2*c)^2 - 4) + 3*log(abs(tan(1/2*d*x + 1/2*c) + 2)) - 3*log(abs(tan(1/2*d*x + 1/2*c) - 2)))/d`**3.47.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.52

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^2} dx = -\frac{3 \operatorname{atanh}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{32 d} - \frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{16 d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 4\right)}$$

input `int(1/(5*cos(c + d*x) + 3)^2,x)`output `-(3*atanh(tan(c/2 + (d*x)/2)/2))/(32*d) - (5*tan(c/2 + (d*x)/2))/(16*d*(tan(c/2 + (d*x)/2)^2 - 4))`

### 3.48 $\int \frac{1}{(-3-5 \cos(c+dx))^3} dx$

3.48.1	Optimal result . . . . .	317
3.48.2	Mathematica [A] (verified) . . . . .	318
3.48.3	Rubi [A] (verified) . . . . .	318
3.48.4	Maple [A] (verified) . . . . .	321
3.48.5	Fricas [A] (verification not implemented) . . . . .	321
3.48.6	Sympy [B] (verification not implemented) . . . . .	322
3.48.7	Maxima [A] (verification not implemented) . . . . .	322
3.48.8	Giac [A] (verification not implemented) . . . . .	323
3.48.9	Mupad [B] (verification not implemented) . . . . .	323

#### 3.48.1 Optimal result

Integrand size = 12, antiderivative size = 115

$$\int \frac{1}{(-3-5 \cos(c+dx))^3} dx = \frac{43 \log(2 \cos(\frac{1}{2}(c+dx)) - \sin(\frac{1}{2}(c+dx)))}{2048d} - \frac{43 \log(2 \cos(\frac{1}{2}(c+dx)) + \sin(\frac{1}{2}(c+dx)))}{2048d} - \frac{5 \sin(c+dx)}{32d(3+5 \cos(c+dx))^2} + \frac{45 \sin(c+dx)}{512d(3+5 \cos(c+dx))}$$

```
output 43/2048*ln(2*cos(1/2*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d-43/2048*ln(2*cos(1/2
*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d-5/32*sin(d*x+c)/d/(3+5*cos(d*x+c))^2+45/
512*sin(d*x+c)/d/(3+5*cos(d*x+c))
```

### 3.48.2 Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.89

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^3} dx = \frac{43 \log(2 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}{2048d} - \frac{43 \log(2 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{2048d} - \frac{512d(2 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))^2}{45 \sin(\frac{1}{2}(c + dx))} + \frac{2048d(2 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))^2}{5} + \frac{512d(2 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))^2}{45 \sin(\frac{1}{2}(c + dx))} + \frac{2048d(2 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))^2}{5}$$

input `Integrate[(-3 - 5*Cos[c + d*x])^(-3),x]`

output `(43*Log[2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]])/(2048*d) - (43*Log[2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]])/(2048*d) - 5/(512*d*(2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^2) + (45*Sin[(c + d*x)/2])/(2048*d*(2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2])) + 5/(512*d*(2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2) + (45*Sin[(c + d*x)/2])/(2048*d*(2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))`

### 3.48.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.68, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {3042, 3143, 3042, 3233, 27, 3042, 3138, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(-5 \cos(c + dx) - 3)^3} dx$$

↓ 3042

$$\int \frac{1}{(-5 \sin(c + dx + \frac{\pi}{2}) - 3)^3} dx$$

---

3.48.  $\int \frac{1}{(-3 - 5 \cos(c + dx))^3} dx$

$$\begin{aligned}
& \downarrow \text{3143} \\
& \frac{1}{32} \int \frac{6 - 5 \cos(c + dx)}{(5 \cos(c + dx) + 3)^2} dx - \frac{5 \sin(c + dx)}{32d(5 \cos(c + dx) + 3)^2} \\
& \downarrow \text{3042} \\
& \frac{1}{32} \int \frac{6 - 5 \sin(c + dx + \frac{\pi}{2})}{(5 \sin(c + dx + \frac{\pi}{2}) + 3)^2} dx - \frac{5 \sin(c + dx)}{32d(5 \cos(c + dx) + 3)^2} \\
& \downarrow \text{3233} \\
& \frac{1}{32} \left( \frac{1}{16} \int -\frac{43}{5 \cos(c + dx) + 3} dx + \frac{45 \sin(c + dx)}{16d(5 \cos(c + dx) + 3)} \right) - \frac{5 \sin(c + dx)}{32d(5 \cos(c + dx) + 3)^2} \\
& \downarrow \text{27} \\
& \frac{1}{32} \left( \frac{45 \sin(c + dx)}{16d(5 \cos(c + dx) + 3)} - \frac{43}{16} \int \frac{1}{5 \cos(c + dx) + 3} dx \right) - \frac{5 \sin(c + dx)}{32d(5 \cos(c + dx) + 3)^2} \\
& \downarrow \text{3042} \\
& \frac{1}{32} \left( \frac{45 \sin(c + dx)}{16d(5 \cos(c + dx) + 3)} - \frac{43}{16} \int \frac{1}{5 \sin(c + dx + \frac{\pi}{2}) + 3} dx \right) - \frac{5 \sin(c + dx)}{32d(5 \cos(c + dx) + 3)^2} \\
& \downarrow \text{3138} \\
& \frac{1}{32} \left( \frac{45 \sin(c + dx)}{16d(5 \cos(c + dx) + 3)} - \frac{43 \int \frac{1}{8 - 2 \tan^2(\frac{1}{2}(c + dx))} d \tan(\frac{1}{2}(c + dx))}{8d} \right) - \frac{5 \sin(c + dx)}{32d(5 \cos(c + dx) + 3)^2} \\
& \downarrow \text{219} \\
& \frac{1}{32} \left( \frac{45 \sin(c + dx)}{16d(5 \cos(c + dx) + 3)} - \frac{43 \operatorname{arctanh}(\frac{1}{2} \tan(\frac{1}{2}(c + dx)))}{32d} \right) - \frac{5 \sin(c + dx)}{32d(5 \cos(c + dx) + 3)^2}
\end{aligned}$$

input `Int[(-3 - 5*Cos[c + d*x])^(-3),x]`

output `(-5*Sin[c + d*x])/(32*d*(3 + 5*Cos[c + d*x])^2) + ((-43*ArcTanh[Tan[(c + d*x)/2]/2])/(32*d) + (45*Sin[c + d*x])/(16*d*(3 + 5*Cos[c + d*x]))) / 32`



## 3.48.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3143 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`
- rule 3233 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

### 3.48.4 Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.72

method	result
norman	$\frac{35 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 85 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{128d - 512d \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 4\right)} + \frac{43 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{2048d} - \frac{43 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{2048d}$
derivativedivides	$\frac{\frac{25}{512 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)^2} - \frac{85}{1024 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)} - \frac{43 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{2048} - \frac{25}{512 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)^2} - \frac{85}{1024 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)} + \frac{43 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{2048}}{d}$
default	$\frac{\frac{25}{512 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)^2} - \frac{85}{1024 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)} - \frac{43 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{2048} - \frac{25}{512 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)^2} - \frac{85}{1024 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)} + \frac{43 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{2048}}{d}$
risch	$\frac{i(215 e^{3i(dx+c)} - 387 e^{2i(dx+c)} + 325 e^{i(dx+c)} + 225)}{256d(5 e^{2i(dx+c)} + 6 e^{i(dx+c)} + 5)^2} - \frac{43 \ln\left(e^{i(dx+c)} + \frac{3}{5} + \frac{4i}{5}\right)}{2048d} + \frac{43 \ln\left(e^{i(dx+c)} + \frac{3}{5} - \frac{4i}{5}\right)}{2048d}$
parallelrisch	$\frac{(2580 \cos(dx+c) + 1075 \cos(2dx+2c) + 1849) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right) + (-2580 \cos(dx+c) - 1075 \cos(2dx+2c) - 1849) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{2048d(43 + 25 \cos(2dx+2c) + 60 \cos(dx+c))}$

input `int(1/(-3-5*cos(d*x+c))^3,x,method=_RETURNVERBOSE)`

output 
$$\frac{(35/128/d*\tan(1/2*d*x+1/2*c)-85/512/d*\tan(1/2*d*x+1/2*c)^3)/(\tan(1/2*d*x+1/2*c)^2-4)^2+43/2048/d*\ln(\tan(1/2*d*x+1/2*c)-2)-43/2048/d*\ln(\tan(1/2*d*x+1/2*c)+2)}$$

### 3.48.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.12

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^3} dx = \frac{43 (25 \cos(dx + c)^2 + 30 \cos(dx + c) + 9) \log\left(\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c) + \frac{5}{2}\right) - 43 (25 \cos(dx + c)^2 + 30 \cos(dx + c) + 9) \log\left(\frac{3}{2} \cos(dx + c) - 2 \sin(dx + c) + \frac{5}{2}\right) - 40 (45 \cos(dx + c) + 11) \sin(dx + c)}{4096 (25 d \cos(dx + c) + 30 d + 9)}$$

input `integrate(1/(-3-5*cos(d*x+c))^3,x, algorithm="fricas")`

output 
$$\frac{-1/4096*(43*(25*\cos(d*x + c)^2 + 30*\cos(d*x + c) + 9)*\log(3/2*\cos(d*x + c) + 2*\sin(d*x + c) + 5/2) - 43*(25*\cos(d*x + c)^2 + 30*\cos(d*x + c) + 9)*\log(3/2*\cos(d*x + c) - 2*\sin(d*x + c) + 5/2) - 40*(45*\cos(d*x + c) + 11)*\sin(d*x + c)}{(25*d*\cos(d*x + c)^2 + 30*d*\cos(d*x + c) + 9*d)}$$

---

3.48. 
$$\int \frac{1}{(-3-5 \cos(c+dx))^3} dx$$

### 3.48.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 478 vs.  $2(102) = 204$ .

Time = 1.17 (sec) , antiderivative size = 478, normalized size of antiderivative = 4.16

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^3} dx$$

$$= \begin{cases} \frac{x}{(-3 - 5 \cos(2 \operatorname{atan}(2)))^3} \\ \frac{x}{(-5 \cos(c) - 3)^3} \\ \frac{43 \log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 2\right) \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right)}{2048d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) - 16384d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 32768d} - \frac{344 \log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 2\right) \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{2048d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) - 16384d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 32768d} + \frac{688 \log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 2\right)}{2048d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) - 16384d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 32768d} \end{cases}$$

input `integrate(1/(-3-5*cos(d*x+c))**3,x)`

output `Piecewise((x/(-3 - 5*cos(2*atan(2)))**3, Eq(c, -d*x - 2*atan(2)) | Eq(c, -d*x + 2*atan(2))), (x/(-5*cos(c) - 3)**3, Eq(d, 0)), (43*log(tan(c/2 + d*x/2) - 2)*tan(c/2 + d*x/2)**4/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d) - 344*log(tan(c/2 + d*x/2) - 2)*tan(c/2 + d*x/2)**2/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d) + 688*log(tan(c/2 + d*x/2) - 2)/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d) - 43*log(tan(c/2 + d*x/2) + 2)*tan(c/2 + d*x/2)**4/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d) + 344*log(tan(c/2 + d*x/2) + 2)*tan(c/2 + d*x/2)**2/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d) - 688*log(tan(c/2 + d*x/2) + 2)/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d) - 340*tan(c/2 + d*x/2)**3/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d) + 560*tan(c/2 + d*x/2)/(2048*d*tan(c/2 + d*x/2)**4 - 16384*d*tan(c/2 + d*x/2)**2 + 32768*d), True))`

### 3.48.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.17

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^3} dx$$

$$= -\frac{20 \left( \frac{28 \sin(dx+c)}{\cos(dx+c)+1} - \frac{17 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{\frac{8 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - \frac{\sin(dx+c)^4}{(\cos(dx+c)+1)^4} - 16} + 43 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 2\right) - 43 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 2\right)$$

$2048 d$

---

3.48.  $\int \frac{1}{(-3-5 \cos(c+dx))^3} dx$

input `integrate(1/(-3-5*cos(d*x+c))^3,x, algorithm="maxima")`

output `-1/2048*(20*(28*sin(d*x + c)/(cos(d*x + c) + 1) - 17*sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/(8*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 - sin(d*x + c)^4/(cos(d*x + c) + 1)^4 - 16) + 43*log(sin(d*x + c)/(cos(d*x + c) + 1) + 2) - 43*log(sin(d*x + c)/(cos(d*x + c) + 1) - 2))/d`

### 3.48.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.68

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^3} dx = \frac{20 \left( 17 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 28 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\left( \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 4 \right)^2} + 43 \log \left( \left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 2 \right| \right) - 43 \log \left( \left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 2 \right| \right)$$


---

2048 d

input `integrate(1/(-3-5*cos(d*x+c))^3,x, algorithm="giac")`

output `-1/2048*(20*(17*tan(1/2*d*x + 1/2*c)^3 - 28*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 4)^2 + 43*log(abs(tan(1/2*d*x + 1/2*c) + 2)) - 43*log(abs(tan(1/2*d*x + 1/2*c) - 2)))/d`

### 3.48.9 Mupad [B] (verification not implemented)

Time = 13.84 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.65

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^3} dx = \frac{\frac{35 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{128} - \frac{85 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{512}}{d \left( \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 8 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 16 \right)} - \frac{43 \operatorname{atanh}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{1024 d}$$

input `int(-1/(5*cos(c + d*x) + 3)^3,x)`

output  $((35*\tan(c/2 + (d*x)/2))/128 - (85*\tan(c/2 + (d*x)/2)^3)/512)/(d*(\tan(c/2 + (d*x)/2)^4 - 8*\tan(c/2 + (d*x)/2)^2 + 16)) - (43*atanh(\tan(c/2 + (d*x)/2)/2))/(1024*d)$

### 3.49 $\int \frac{1}{(-3-5 \cos(c+dx))^4} dx$

3.49.1	Optimal result . . . . .	325
3.49.2	Mathematica [B] (verified) . . . . .	326
3.49.3	Rubi [A] (verified) . . . . .	326
3.49.4	Maple [A] (verified) . . . . .	329
3.49.5	Fricas [A] (verification not implemented) . . . . .	330
3.49.6	Sympy [B] (verification not implemented) . . . . .	330
3.49.7	Maxima [A] (verification not implemented) . . . . .	331
3.49.8	Giac [A] (verification not implemented) . . . . .	332
3.49.9	Mupad [B] (verification not implemented) . . . . .	332

#### 3.49.1 Optimal result

Integrand size = 12, antiderivative size = 140

$$\int \frac{1}{(-3-5 \cos(c+dx))^4} dx = \frac{279 \log(2 \cos(\frac{1}{2}(c+dx)) - \sin(\frac{1}{2}(c+dx)))}{32768d} - \frac{279 \log(2 \cos(\frac{1}{2}(c+dx)) + \sin(\frac{1}{2}(c+dx)))}{32768d} + \frac{5 \sin(c+dx)}{48d(3+5 \cos(c+dx))^3} - \frac{25 \sin(c+dx)}{512d(3+5 \cos(c+dx))^2} + \frac{995 \sin(c+dx)}{24576d(3+5 \cos(c+dx))}$$

```
output 279/32768*ln(2*cos(1/2*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d-279/32768*ln(2*cos
(1/2*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d+5/48*sin(d*x+c)/d/(3+5*cos(d*x+c))^3
-25/512*sin(d*x+c)/d/(3+5*cos(d*x+c))^2+995/24576*sin(d*x+c)/d/(3+5*cos(d*
x+c))
```

### 3.49.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 296 vs.  $2(140) = 280$ .

Time = 0.01 (sec) , antiderivative size = 296, normalized size of antiderivative = 2.11

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^4} dx$$

$$= \frac{467046 \log\left(2 \cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) + 104625 \cos(3(c + dx)) \log\left(2 \cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) - \sin\left(\frac{1}{2}(c + dx)\right)}{\dots}$$

input `Integrate[(-3 - 5*Cos[c + d*x])^(-4),x]`

output `(467046*Log[2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 104625*Cos[3*(c + d*x)]*Log[2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 765855*Cos[c + d*x]*(Log[2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - Log[2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]) + 376650*Cos[2*(c + d*x)]*(Log[2*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - Log[2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]) - 467046*Log[2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] - 104625*Cos[3*(c + d*x)]*Log[2*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + 226140*Sin[c + d*x] + 190800*Sin[2*(c + d*x)] + 99500*Sin[3*(c + d*x)])/(393216*d*(3 + 5*Cos[c + d*x])^3)`

### 3.49.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.77, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {3042, 3143, 25, 3042, 3233, 25, 3042, 3233, 27, 3042, 3138, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(-5 \cos(c + dx) - 3)^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(-5 \sin(c + dx + \frac{\pi}{2}) - 3)^4} dx$$

$$\downarrow \text{3143}$$

$$\frac{1}{48} \int -\frac{9 - 10 \cos(c + dx)}{(5 \cos(c + dx) + 3)^3} dx + \frac{5 \sin(c + dx)}{48d(5 \cos(c + dx) + 3)^3}$$

$$\begin{aligned}
& \downarrow 25 \\
& \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} - \frac{1}{48} \int \frac{9-10 \cos(c+dx)}{(5 \cos(c+dx)+3)^3} dx \\
& \downarrow 3042 \\
& \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} - \frac{1}{48} \int \frac{9-10 \sin(c+dx+\frac{\pi}{2})}{(5 \sin(c+dx+\frac{\pi}{2})+3)^3} dx \\
& \downarrow 3233 \\
& \frac{1}{48} \left( -\frac{1}{32} \int -\frac{154-75 \cos(c+dx)}{(5 \cos(c+dx)+3)^2} dx - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} \\
& \downarrow 25 \\
& \frac{1}{48} \left( \frac{1}{32} \int \frac{154-75 \cos(c+dx)}{(5 \cos(c+dx)+3)^2} dx - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left( \frac{1}{32} \int \frac{154-75 \sin(c+dx+\frac{\pi}{2})}{(5 \sin(c+dx+\frac{\pi}{2})+3)^2} dx - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} \\
& \downarrow 3233 \\
& \frac{1}{48} \left( \frac{1}{32} \left( \frac{1}{16} \int -\frac{837}{5 \cos(c+dx)+3} dx + \frac{995 \sin(c+dx)}{16d(5 \cos(c+dx)+3)} \right) - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \\
& \quad \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} \\
& \downarrow 27 \\
& \frac{1}{48} \left( \frac{1}{32} \left( \frac{995 \sin(c+dx)}{16d(5 \cos(c+dx)+3)} - \frac{837}{16} \int \frac{1}{5 \cos(c+dx)+3} dx \right) - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \\
& \quad \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left( \frac{1}{32} \left( \frac{995 \sin(c+dx)}{16d(5 \cos(c+dx)+3)} - \frac{837}{16} \int \frac{1}{5 \sin(c+dx+\frac{\pi}{2})+3} dx \right) - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \\
& \quad \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3} \\
& \downarrow 3138
\end{aligned}$$



$$\frac{1}{48} \left( \frac{1}{32} \left( \frac{995 \sin(c+dx)}{16d(5 \cos(c+dx)+3)} - \frac{837 \int \frac{1}{8-2 \tan^2(\frac{1}{2}(c+dx))} d \tan(\frac{1}{2}(c+dx))}{8d} \right) - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3}$$

↓ 219

$$\frac{1}{48} \left( \frac{1}{32} \left( \frac{995 \sin(c+dx)}{16d(5 \cos(c+dx)+3)} - \frac{837 \operatorname{arctanh}(\frac{1}{2} \tan(\frac{1}{2}(c+dx)))}{32d} \right) - \frac{75 \sin(c+dx)}{32d(5 \cos(c+dx)+3)^2} \right) + \frac{5 \sin(c+dx)}{48d(5 \cos(c+dx)+3)^3}$$

input `Int[(-3 - 5*Cos[c + d*x])^(-4), x]`

output `(5*Sin[c + d*x])/(48*d*(3 + 5*Cos[c + d*x])^3) + ((-75*Sin[c + d*x])/(32*d*(3 + 5*Cos[c + d*x])^2) + ((-837*ArcTanh[Tan[(c + d*x)/2]/2])/(32*d) + (995*Sin[c + d*x])/(16*d*(3 + 5*Cos[c + d*x]))) / 32) / 48`

### 3.49.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3233 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

### 3.49.4 Maple [A] (verified)

Time = 0.68 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.71

method	result
norman	$-\frac{295 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{265 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 745 \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{512d} + \frac{279 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{32768d} - \frac{279 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)}{32768d}$
derivativedivides	$\frac{-\frac{125}{6144 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)^3} - \frac{175}{4096 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)^2} - \frac{745}{16384 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)} + \frac{279 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{32768} - \frac{125}{6144 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)^3}}{d}$
default	$\frac{-\frac{125}{6144 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)^3} - \frac{175}{4096 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)^2} - \frac{745}{16384 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)} + \frac{279 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right)}{32768} - \frac{125}{6144 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right)^3}}{d}$
risch	$\frac{i(20925 e^{5i(dx+c)} + 62775 e^{4i(dx+c)} + 111042 e^{3i(dx+c)} + 119310 e^{2i(dx+c)} + 68625 e^{i(dx+c)} + 24875)}{12288d(5 e^{2i(dx+c)} + 6 e^{i(dx+c)} + 5)^3} + \frac{279 \ln(e^{i(dx+c)} + 2)}{32768d}$
parallelrisc	$\frac{(765855 \cos(dx+c) + 376650 \cos(2dx+2c) + 104625 \cos(3dx+3c) + 467046) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\right) + (-765855 \cos(dx+c) - 391311 \cos(2dx+2c) - 104625 \cos(3dx+3c) - 467046)}{98304d(558 + 125 \cos(3dx+c))}$

input `int(1/(-3-5*cos(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `(-295/512/d*tan(1/2*d*x+1/2*c)+265/768/d*tan(1/2*d*x+1/2*c)^3-745/8192/d*tan(1/2*d*x+1/2*c)^5)/(tan(1/2*d*x+1/2*c)^2-4)^3+279/32768/d*ln(tan(1/2*d*x+1/2*c)-2)-279/32768/d*ln(tan(1/2*d*x+1/2*c)+2)`

**3.49.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.21

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^4} dx = \frac{837 (125 \cos(dx + c)^3 + 225 \cos(dx + c)^2 + 135 \cos(dx + c) + 27) \log\left(\frac{3}{2} \cos(dx + c) + 2 \sin(dx + c)\right)}{\dots}$$

input `integrate(1/(-3-5*cos(d*x+c))^4,x, algorithm="fracas")`

output `-1/196608*(837*(125*cos(d*x + c)^3 + 225*cos(d*x + c)^2 + 135*cos(d*x + c) + 27)*log(3/2*cos(d*x + c) + 2*sin(d*x + c) + 5/2) - 837*(125*cos(d*x + c)^3 + 225*cos(d*x + c)^2 + 135*cos(d*x + c) + 27)*log(3/2*cos(d*x + c) - 2*sin(d*x + c) + 5/2) - 40*(4975*cos(d*x + c)^2 + 4770*cos(d*x + c) + 1583)*sin(d*x + c))/(125*d*cos(d*x + c)^3 + 225*d*cos(d*x + c)^2 + 135*d*cos(d*x + c) + 27*d)`

**3.49.6 Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 816 vs.  $2(126) = 252$ .

Time = 2.44 (sec) , antiderivative size = 816, normalized size of antiderivative = 5.83

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^4} dx = \text{Too large to display}$$

input `integrate(1/(-3-5*cos(d*x+c))**4,x)`

```
output Piecewise((x/(-3 - 5*cos(2*atan(2)))**4, Eq(c, -d*x - 2*atan(2)) | Eq(c, -
d*x + 2*atan(2))), (x/(-5*cos(c) - 3)**4, Eq(d, 0)), (837*log(tan(c/2 + d*
x/2) - 2)*tan(c/2 + d*x/2)**6/(98304*d*tan(c/2 + d*x/2)**6 - 1179648*d*tan
(c/2 + d*x/2)**4 + 4718592*d*tan(c/2 + d*x/2)**2 - 6291456*d) - 10044*log(
tan(c/2 + d*x/2) - 2)*tan(c/2 + d*x/2)**4/(98304*d*tan(c/2 + d*x/2)**6 - 1
179648*d*tan(c/2 + d*x/2)**4 + 4718592*d*tan(c/2 + d*x/2)**2 - 6291456*d)
+ 40176*log(tan(c/2 + d*x/2) - 2)*tan(c/2 + d*x/2)**2/(98304*d*tan(c/2 + d
*x/2)**6 - 1179648*d*tan(c/2 + d*x/2)**4 + 4718592*d*tan(c/2 + d*x/2)**2 -
6291456*d) - 53568*log(tan(c/2 + d*x/2) - 2)/(98304*d*tan(c/2 + d*x/2)**6
- 1179648*d*tan(c/2 + d*x/2)**4 + 4718592*d*tan(c/2 + d*x/2)**2 - 6291456
*d) - 837*log(tan(c/2 + d*x/2) + 2)*tan(c/2 + d*x/2)**6/(98304*d*tan(c/2 +
d*x/2)**6 - 1179648*d*tan(c/2 + d*x/2)**4 + 4718592*d*tan(c/2 + d*x/2)**2
- 6291456*d) + 10044*log(tan(c/2 + d*x/2) + 2)*tan(c/2 + d*x/2)**4/(98304
*d*tan(c/2 + d*x/2)**6 - 1179648*d*tan(c/2 + d*x/2)**4 + 4718592*d*tan(c/2
+ d*x/2)**2 - 6291456*d) - 40176*log(tan(c/2 + d*x/2) + 2)*tan(c/2 + d*x/
2)**2/(98304*d*tan(c/2 + d*x/2)**6 - 1179648*d*tan(c/2 + d*x/2)**4 + 47185
92*d*tan(c/2 + d*x/2)**2 - 6291456*d) + 53568*log(tan(c/2 + d*x/2) + 2)/(9
8304*d*tan(c/2 + d*x/2)**6 - 1179648*d*tan(c/2 + d*x/2)**4 + 4718592*d*tan
(c/2 + d*x/2)**2 - 6291456*d) - 8940*tan(c/2 + d*x/2)**5/(98304*d*tan(c/2
+ d*x/2)**6 - 1179648*d*tan(c/2 + d*x/2)**4 + 4718592*d*tan(c/2 + d*x/2...
```

### 3.49.7 Maxima [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.24

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^4} dx = \frac{20 \left( \frac{2832 \sin(dx+c)}{\cos(dx+c)+1} - \frac{1696 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{447 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right) + 837 \log \left( \frac{\sin(dx+c)}{\cos(dx+c)+1} + 2 \right) - 837 \log \left( \frac{\sin(dx+c)}{\cos(dx+c)+1} - 2 \right)}{98304 d} - \frac{\frac{48 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - \frac{12 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{\sin(dx+c)^6}{(\cos(dx+c)+1)^6} - 64}{98304 d}$$

```
input integrate(1/(-3-5*cos(d*x+c))^4,x, algorithm="maxima")
```

```
output -1/98304*(20*(2832*sin(d*x + c)/(cos(d*x + c) + 1) - 1696*sin(d*x + c)^3/(
cos(d*x + c) + 1)^3 + 447*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/(48*sin(d*x
+ c)^2/(cos(d*x + c) + 1)^2 - 12*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + si
n(d*x + c)^6/(cos(d*x + c) + 1)^6 - 64) + 837*log(sin(d*x + c)/(cos(d*x +
c) + 1) + 2) - 837*log(sin(d*x + c)/(cos(d*x + c) + 1) - 2))/d
```

---

3.49.  $\int \frac{1}{(-3-5 \cos(c+dx))^4} dx$

**3.49.8 Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.65

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^4} dx = \frac{20 \left( 447 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 1696 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 2832 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right)}{\left( \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 4 \right)^3} + 837 \log \left( \left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 2 \right| \right) - 837 \log \left( \left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 2 \right| \right)$$


---

98304 d

input `integrate(1/(-3-5*cos(d*x+c))^4,x, algorithm="giac")`output `-1/98304*(20*(447*tan(1/2*d*x + 1/2*c)^5 - 1696*tan(1/2*d*x + 1/2*c)^3 + 2832*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 4)^3 + 837*log(abs(tan(1/2*d*x + 1/2*c) + 2)) - 837*log(abs(tan(1/2*d*x + 1/2*c) - 2)))/d`**3.49.9 Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.73

$$\int \frac{1}{(-3 - 5 \cos(c + dx))^4} dx = -\frac{279 \operatorname{atanh}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2}\right)}{16384 d} - \frac{\frac{745 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{8192} - \frac{265 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{768} + \frac{295 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{512}}{d \left( \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - 12 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 48 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 64 \right)}$$

input `int(1/(5*cos(c + d*x) + 3)^4,x)`output `-(279*atanh(tan(c/2 + (d*x)/2)/2))/(16384*d) - ((295*tan(c/2 + (d*x)/2))/512 - (265*tan(c/2 + (d*x)/2)^3)/768 + (745*tan(c/2 + (d*x)/2)^5)/8192)/(d*(48*tan(c/2 + (d*x)/2)^2 - 12*tan(c/2 + (d*x)/2)^4 + tan(c/2 + (d*x)/2)^6 - 64))`

### 3.50 $\int (a + b \cos(c + dx))^{5/2} dx$

3.50.1	Optimal result . . . . .	333
3.50.2	Mathematica [A] (verified) . . . . .	334
3.50.3	Rubi [A] (verified) . . . . .	334
3.50.4	Maple [B] (verified) . . . . .	338
3.50.5	Fricas [C] (verification not implemented) . . . . .	339
3.50.6	Sympy [F] . . . . .	340
3.50.7	Maxima [F] . . . . .	340
3.50.8	Giac [F] . . . . .	341
3.50.9	Mupad [F(-1)] . . . . .	341

#### 3.50.1 Optimal result

Integrand size = 14, antiderivative size = 197

$$\int (a + b \cos(c + dx))^{5/2} dx = \frac{2(23a^2 + 9b^2) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{15d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{16a(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{15d \sqrt{a + b \cos(c + dx)}} + \frac{16ab \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{15d} + \frac{2b(a + b \cos(c + dx))^{3/2} \sin(c + dx)}{5d}$$

```
output 2/5*b*(a+b*cos(d*x+c))^(3/2)*sin(d*x+c)/d+16/15*a*b*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d+2/15*(23*a^2+9*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/d/((a+b*cos(d*x+c))/(a+b))^(1/2)-16/15*a*(a^2-b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/d/(a+b*cos(d*x+c))^(1/2)
```

### 3.50.2 Mathematica [A] (verified)

Time = 0.92 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.90

$$\int (a + b \cos(c + dx))^{5/2} dx = \frac{2(23a^3 + 23a^2b + 9ab^2 + 9b^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right) - 16a(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right) + b(22a^2 + 3b^2 + 28ab \cos[c + dx] + 3b^2 \cos[2(c + dx)]) \sin[c + dx]}{15d \sqrt{a + b \cos(c + dx)}}$$

input `Integrate[(a + b*Cos[c + d*x])^(5/2), x]`

output `(2*(23*a^3 + 23*a^2*b + 9*a*b^2 + 9*b^3)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticE[(c + d*x)/2, (2*b)/(a + b)] - 16*a*(a^2 - b^2)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)] + b*(22*a^2 + 3*b^2 + 28*a*b*Cos[c + d*x] + 3*b^2*Cos[2*(c + d*x)])*Sin[c + d*x]/(15*d*Sqrt[a + b*Cos[c + d*x]])`

### 3.50.3 Rubi [A] (verified)

Time = 1.03 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.03, number of steps used = 15, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.071$ , Rules used = {3042, 3135, 27, 3042, 3232, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \cos(c + dx))^{5/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \left( a + b \sin\left(c + dx + \frac{\pi}{2}\right) \right)^{5/2} dx \\ & \quad \downarrow \text{3135} \\ & \frac{2}{5} \int \frac{1}{2} \sqrt{a + b \cos(c + dx)} (5a^2 + 8b \cos(c + dx)a + 3b^2) dx + \frac{2b \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \\ & \quad \downarrow \text{27} \\ & \frac{1}{5} \int \sqrt{a + b \cos(c + dx)} (5a^2 + 8b \cos(c + dx)a + 3b^2) dx + \frac{2b \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \end{aligned}$$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \frac{1}{5} \int \frac{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} \left(5a^2 + 8b \sin\left(c + dx + \frac{\pi}{2}\right) a + 3b^2\right) dx +}{2b \sin(c + dx)(a + b \cos(c + dx))^{3/2}} \\
& \downarrow \text{3232} \\
& \frac{1}{5} \left( \frac{2}{3} \int \frac{a(15a^2 + 17b^2) + b(23a^2 + 9b^2) \cos(c + dx)}{2\sqrt{a + b \cos(c + dx)}} dx + \frac{16ab \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \right) + \\
& \quad \frac{2b \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \\
& \downarrow \text{27} \\
& \frac{1}{5} \left( \frac{1}{3} \int \frac{a(15a^2 + 17b^2) + b(23a^2 + 9b^2) \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx + \frac{16ab \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \right) + \\
& \quad \frac{2b \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \\
& \downarrow \text{3042} \\
& \frac{1}{5} \left( \frac{1}{3} \int \frac{a(15a^2 + 17b^2) + b(23a^2 + 9b^2) \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{16ab \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \right) + \\
& \quad \frac{2b \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \\
& \downarrow \text{3231} \\
& \frac{1}{5} \left( \frac{1}{3} \left( (23a^2 + 9b^2) \int \sqrt{a + b \cos(c + dx)} dx - 8a(a^2 - b^2) \int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx \right) + \frac{16ab \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \right) + \\
& \quad \frac{2b \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \\
& \downarrow \text{3042} \\
& \frac{1}{5} \left( \frac{1}{3} \left( (23a^2 + 9b^2) \int \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} dx - 8a(a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx \right) + \frac{16ab \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \right) + \\
& \quad \frac{2b \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \\
& \downarrow \text{3134}
\end{aligned}$$



$$\frac{1}{5} \left( \frac{1}{3} \left( \frac{(23a^2 + 9b^2) \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - 8a(a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) + \frac{2b \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \right) \downarrow \text{3042}$$

$$\frac{1}{5} \left( \frac{1}{3} \left( \frac{(23a^2 + 9b^2) \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - 8a(a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) + \frac{2b \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \right) \downarrow \text{3132}$$

$$\frac{1}{5} \left( \frac{1}{3} \left( \frac{2(23a^2 + 9b^2) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - 8a(a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) + \frac{2b \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \right) \downarrow \text{3142}$$

$$\frac{1}{5} \left( \frac{1}{3} \left( \frac{2(23a^2 + 9b^2) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{8a(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{\sqrt{a + b \cos(c + dx)}} \right) + \frac{2b \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \right) \downarrow \text{3042}$$

$$\frac{1}{5} \left( \frac{1}{3} \left( \frac{2(23a^2 + 9b^2) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{8a(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a + b \cos(c + dx)}} \right) + \frac{2b \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \right) \downarrow \text{3140}$$

$$\frac{1}{5} \left( \frac{1}{3} \left( \frac{2(23a^2 + 9b^2) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{16a(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \right)}{d \sqrt{a + b \cos(c + dx)}} \right) - \frac{2b \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \right)$$

input `Int[(a + b*Cos[c + d*x])^(5/2), x]`

output `(2*b*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*d) + (((2*(23*a^2 + 9*b^2)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (16*a*(a^2 - b^2)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]])))/3 + (16*a*b*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d))/5`

### 3.50.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`



```
input int((a+cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)
```

```
output -2/15*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(24*cos(
1/2*d*x+1/2*c)^7*b^3+56*cos(1/2*d*x+1/2*c)^5*a*b^2-48*cos(1/2*d*x+1/2*c)^5
*b^3+22*cos(1/2*d*x+1/2*c)^3*a^2*b-84*cos(1/2*d*x+1/2*c)^3*a*b^2+30*cos(1/
2*d*x+1/2*c)^3*b^3-8*a^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/
2*c)^2+a-b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+
8*a*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b)
)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+23*(sin(1/2*d*x+1
/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticE(cos(
1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^3-23*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2
*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2
*b/(a-b))^(1/2))*a^2*b+9*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/
2*c)^2+a-b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*
a*b^2-9*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b)
)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*b^3-22*cos(1/2*d*
x+1/2*c)*a^2*b+28*cos(1/2*d*x+1/2*c)*a*b^2-6*cos(1/2*d*x+1/2*c)*b^3)/(-2*s
in(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)
/(-2*b*sin(1/2*d*x+1/2*c)^2+a+b)^(1/2)/d
```

### 3.50.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 437, normalized size of antiderivative = 2.22

$$\int (a + b \cos(c + dx))^{5/2} dx = \frac{\sqrt{2}(i a^3 - 33i ab^2)\sqrt{b}\text{weierstrassPInverse}\left(\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8a^3 - 9ab^2)}{27b^3}, \frac{3b \cos(dx+c) + 3i b \sin(dx+c) + 2a}{3b}\right)}{1}$$

```
input integrate((a+b*cos(d*x+c))^(5/2),x, algorithm="fricas")
```

```
output 1/45*(sqrt(2)*(I*a^3 - 33*I*a*b^2)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2
- 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*s
in(d*x + c) + 2*a)/b) + sqrt(2)*(-I*a^3 + 33*I*a*b^2)*sqrt(b)*weierstrassP
Inverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos
(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*sqrt(2)*(-23*I*a^2*b - 9*I*b^
3)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2
)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2
)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*sqrt(2)*
(23*I*a^2*b + 9*I*b^3)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/
27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/
27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a
)/b)) + 6*(3*b^3*cos(d*x + c) + 11*a*b^2)*sqrt(b*cos(d*x + c) + a)*sin(d*x
+ c))/(b*d)
```

### 3.50.6 Sympy [F]

$$\int (a + b \cos(c + dx))^{5/2} dx = \int (a + b \cos(c + dx))^{5/2} dx$$

```
input integrate((a+b*cos(d*x+c))**(5/2), x)
```

```
output Integral((a + b*cos(c + d*x))**(5/2), x)
```

### 3.50.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{5/2} dx = \int (b \cos(dx + c) + a)^{5/2} dx$$

```
input integrate((a+b*cos(d*x+c))^(5/2), x, algorithm="maxima")
```

```
output integrate((b*cos(d*x + c) + a)^(5/2), x)
```

**3.50.8 Giac [F]**

$$\int (a + b \cos(c + dx))^{5/2} dx = \int (b \cos(dx + c) + a)^{\frac{5}{2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*cos(d*x + c) + a)^(5/2), x)`

**3.50.9 Mupad [F(-1)]**

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} dx = \int (a + b \cos(c + dx))^{5/2} dx$$

input `int((a + b*cos(c + d*x))^(5/2),x)`

output `int((a + b*cos(c + d*x))^(5/2), x)`

### 3.51 $\int (a + b \cos(c + dx))^{3/2} dx$

3.51.1	Optimal result . . . . .	342
3.51.2	Mathematica [A] (verified) . . . . .	343
3.51.3	Rubi [A] (verified) . . . . .	343
3.51.4	Maple [B] (verified) . . . . .	347
3.51.5	Fricas [C] (verification not implemented) . . . . .	347
3.51.6	Sympy [F] . . . . .	348
3.51.7	Maxima [F] . . . . .	348
3.51.8	Giac [F] . . . . .	349
3.51.9	Mupad [F(-1)] . . . . .	349

#### 3.51.1 Optimal result

Integrand size = 14, antiderivative size = 157

$$\int (a + b \cos(c + dx))^{3/2} dx = \frac{8a\sqrt{a + b \cos(c + dx)}E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{3d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2(a^2 - b^2)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{3d\sqrt{a + b \cos(c + dx)}} + \frac{2b\sqrt{a + b \cos(c + dx)} \sin(c + dx)}{3d}$$

```
output 2/3*b*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d+8/3*a*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*
(a+b*cos(d*x+c))^(1/2)/d/((a+b*cos(d*x+c))/(a+b))^(1/2)-2/3*(a^2-b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/d/(a+b*cos(d*x+c))^(1/2)
```

### 3.51.2 Mathematica [A] (verified)

Time = 0.62 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.85

$$\int (a + b \cos(c + dx))^{3/2} dx = \frac{8a(a+b) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right) - 2(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{3d \sqrt{a+b \cos(c+dx)}}$$

input `Integrate[(a + b*Cos[c + d*x])^(3/2), x]`

output `(8*a*(a + b)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticE[(c + d*x)/2, (2*b)/(a + b)] - 2*(a^2 - b^2)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)] + 2*b*(a + b*Cos[c + d*x])*Sin[c + d*x]/(3*d*Sqrt[a + b*Cos[c + d*x]])`

### 3.51.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.01, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$ , Rules used = {3042, 3135, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \cos(c + dx))^{3/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \left( a + b \sin\left(c + dx + \frac{\pi}{2}\right) \right)^{3/2} dx \\ & \quad \downarrow \text{3135} \\ & \frac{2}{3} \int \frac{3a^2 + 4b \cos(c + dx)a + b^2}{2\sqrt{a + b \cos(c + dx)}} dx + \frac{2b \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\ & \quad \downarrow \text{27} \\ & \frac{1}{3} \int \frac{3a^2 + 4b \cos(c + dx)a + b^2}{\sqrt{a + b \cos(c + dx)}} dx + \frac{2b \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\ & \quad \downarrow \text{3042} \end{aligned}$$



$$\begin{aligned}
& \frac{1}{3} \int \frac{3a^2 + 4b \sin(c + dx + \frac{\pi}{2}) a + b^2}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \frac{2b \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \quad \downarrow \text{3231} \\
& \frac{1}{3} \left( 4a \int \sqrt{a + b \cos(c + dx)} dx - (a^2 - b^2) \int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx \right) + \\
& \quad \frac{2b \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left( 4a \int \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} dx - (a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) + \\
& \quad \frac{2b \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \quad \downarrow \text{3134} \\
& \frac{1}{3} \left( \frac{4a \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - (a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) + \\
& \quad \frac{2b \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left( \frac{4a \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - (a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) + \\
& \quad \frac{2b \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \quad \downarrow \text{3132} \\
& \frac{1}{3} \left( \frac{8a \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - (a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) + \\
& \quad \frac{2b \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \quad \downarrow \text{3142}
\end{aligned}$$

$$\frac{1}{3} \left( \frac{8a\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx}{\sqrt{a+b\cos(c+dx)}} \right) +$$

$$\frac{2b\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{3} \left( \frac{8a\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b\cos(c+dx)}} \right) +$$

$$\frac{2b\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3d}$$

↓ 3140

$$\frac{1}{3} \left( \frac{8a\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{2(a^2-b^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}} \right) +$$

$$\frac{2b\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3d}$$

input `Int[(a + b*Cos[c + d*x])^(3/2), x]`

output `((8*a*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*(a^2 - b^2)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]]))/3 + (2*b*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d)`

### 3.51.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3135 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[1/n Int[(a + b*Sin[c + d*x])^(n - 2)*Simp[a^2*n + b^2*(n - 1) + a*b*(2*n - 1)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

### 3.51.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 449 vs.  $2(199) = 398$ .

Time = 4.09 (sec) , antiderivative size = 450, normalized size of antiderivative = 2.87

method	result
default	$-\frac{2\sqrt{\left(2b\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a-b\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{\left(4\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b^2+2\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)ab-6\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b^2-a^2\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)}$

input `int((a+cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)`

output 
$$-\frac{2}{3} * \left( (2*b*\cos(1/2*d*x+1/2*c))^2 + a - b \right) * \sin(1/2*d*x+1/2*c)^2)^{(1/2)} * \left( 4*\cos(1/2*d*x+1/2*c)^5*b^2 + 2*\cos(1/2*d*x+1/2*c)^3*a*b - 6*\cos(1/2*d*x+1/2*c)^3*b^2 - a^2 * (\sin(1/2*d*x+1/2*c)^2)^{(1/2)} * \left( (2*b*\cos(1/2*d*x+1/2*c))^2 + a - b \right) / (a - b) \right)^{(1/2)} * \text{EllipticF}(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{(1/2)}) + b^2 * (\sin(1/2*d*x+1/2*c)^2)^{(1/2)} * \left( (2*b*\cos(1/2*d*x+1/2*c))^2 + a - b \right) / (a - b) \right)^{(1/2)} * \text{EllipticF}(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{(1/2)}) + 4 * (\sin(1/2*d*x+1/2*c)^2)^{(1/2)} * \left( (2*b*\cos(1/2*d*x+1/2*c))^2 + a - b \right) / (a - b) \right)^{(1/2)} * \text{EllipticE}(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{(1/2)}) * a^2 - 4 * (\sin(1/2*d*x+1/2*c)^2)^{(1/2)} * \left( (2*b*\cos(1/2*d*x+1/2*c))^2 + a - b \right) / (a - b) \right)^{(1/2)} * \text{EllipticE}(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{(1/2)}) * a*b - 2*\cos(1/2*d*x+1/2*c)*a*b + 2*\cos(1/2*d*x+1/2*c)*b^2 / (-2*\sin(1/2*d*x+1/2*c)^4*b + (a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)} / \sin(1/2*d*x+1/2*c) / (-2*b*\sin(1/2*d*x+1/2*c)^2 + a + b)^{(1/2)} / d$$

### 3.51.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 398, normalized size of antiderivative = 2.54

$$\int (a + b \cos(c + dx))^{3/2} dx = \frac{12i \sqrt{2} ab^3 \text{weierstrassZeta}\left(\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8a^3 - 9ab^2)}{27b^3}\right), \text{weierstrassPInverse}\left(\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8a^3 - 9ab^2)}{27b^3}\right)}{\dots}$$

input `integrate((a+b*cos(d*x+c))^(3/2),x, algorithm="fracas")`

output `1/9*(12*I*sqrt(2)*a*b^(3/2)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 12*I*sqrt(2)*a*b^(3/2)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)) + 6*sqrt(b*cos(d*x + c) + a)*b^2*sin(d*x + c) + sqrt(2)*(-I*a^2 - 3*I*b^2)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + sqrt(2)*(I*a^2 + 3*I*b^2)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b))/(b*d)`

### 3.51.6 Sympy [F]

$$\int (a + b \cos(c + dx))^{3/2} dx = \int (a + b \cos(c + dx))^{\frac{3}{2}} dx$$

input `integrate((a+b*cos(d*x+c))**(3/2),x)`

output `Integral((a + b*cos(c + d*x))**(3/2), x)`

### 3.51.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{3/2} dx = \int (b \cos(dx + c) + a)^{\frac{3}{2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((b*cos(d*x + c) + a)^(3/2), x)`

**3.51.8 Giac [F]**

$$\int (a + b \cos(c + dx))^{3/2} dx = \int (b \cos(dx + c) + a)^{\frac{3}{2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*cos(d*x + c) + a)^(3/2), x)`

**3.51.9 Mupad [F(-1)]**

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} dx = \int (a + b \cos(c + dx))^{3/2} dx$$

input `int((a + b*cos(c + d*x))^(3/2),x)`

output `int((a + b*cos(c + d*x))^(3/2), x)`

### 3.52 $\int \sqrt{a + b \cos(c + dx)} dx$

3.52.1	Optimal result . . . . .	350
3.52.2	Mathematica [A] (verified) . . . . .	350
3.52.3	Rubi [A] (verified) . . . . .	351
3.52.4	Maple [B] (verified) . . . . .	352
3.52.5	Fricas [C] (verification not implemented) . . . . .	353
3.52.6	Sympy [F] . . . . .	353
3.52.7	Maxima [F] . . . . .	354
3.52.8	Giac [F] . . . . .	354
3.52.9	Mupad [F(-1)] . . . . .	354

#### 3.52.1 Optimal result

Integrand size = 14, antiderivative size = 57

$$\int \sqrt{a + b \cos(c + dx)} dx = \frac{2\sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

output `2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/d/((a+b*cos(d*x+c))/(a+b))^(1/2)`

#### 3.52.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00

$$\int \sqrt{a + b \cos(c + dx)} dx = \frac{2\sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

input `Integrate[Sqrt[a + b*Cos[c + d*x]],x]`

output `(2*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)])`

### 3.52.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3134, 3042, 3132}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a + b \cos(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{3134} \\
 & \frac{\sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
 & \quad \downarrow \text{3132} \\
 & \frac{2\sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}
 \end{aligned}$$

input `Int[Sqrt[a + b*Cos[c + d*x]],x]`

output `(2*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)])`



3.52.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3132 Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a
+ b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a,
b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3134 Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[Sqrt[a +
b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (
b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2
, 0] && !GtQ[a + b, 0]
```

3.52.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 169 vs. 2(82) = 164.

Time = 2.92 (sec) , antiderivative size = 170, normalized size of antiderivative = 2.98

method	result	size
default	$-\frac{2\sqrt{\left(2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b + (a+b)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{\frac{2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b}{a - b}} E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{-\frac{2b}{a - b}}\right) (a - b)$	170
risch	Expression too large to display	104

```
input int((a+cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -2*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*
x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticE(c
os(1/2*d*x+1/2*c), (-2*b/(a-b))^(1/2))*(a-b)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+
b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(-2*b*sin(1/2*d*x+1/2*c)
^2+a+b)^(1/2)/d
```

### 3.52.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 355, normalized size of antiderivative = 6.23

$$\int \sqrt{a + b \cos(c + dx)} dx$$

$$= -i \sqrt{2a} \sqrt{b} \text{weierstrassPInverse} \left( \frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8a^3 - 9ab^2)}{27b^3}, \frac{3b \cos(dx+c) + 3i b \sin(dx+c) + 2a}{3b} \right) + i \sqrt{2a} \sqrt{b} \text{weierstrassPInverse} \left( \frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8a^3 - 9ab^2)}{27b^3}, \frac{3b \cos(dx+c) - 3i b \sin(dx+c) + 2a}{3b} \right) + \frac{1}{b} \sqrt{a + b \cos(c + dx)}$$

```
input integrate((a+b*cos(d*x+c))^(1/2),x, algorithm="fracas")
```

```
output 1/3*(-I*sqrt(2)*a*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + I*sqrt(2)*a*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) + 3*I*sqrt(2)*b^(3/2)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*I*sqrt(2)*b^(3/2)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)))/(b*d)
```

### 3.52.6 Sympy [F]

$$\int \sqrt{a + b \cos(c + dx)} dx = \int \sqrt{a + b \cos(c + dx)} dx$$

```
input integrate((a+b*cos(d*x+c))**(1/2),x)
```

```
output Integral(sqrt(a + b*cos(c + d*x)), x)
```

**3.52.7 Maxima [F]**

$$\int \sqrt{a + b \cos(c + dx)} dx = \int \sqrt{b \cos(dx + c) + a} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*cos(d*x + c) + a), x)`

**3.52.8 Giac [F]**

$$\int \sqrt{a + b \cos(c + dx)} dx = \int \sqrt{b \cos(dx + c) + a} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*cos(d*x + c) + a), x)`

**3.52.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a + b \cos(c + dx)} dx = \int \sqrt{a + b \cos(c + dx)} dx$$

input `int((a + b*cos(c + d*x))^(1/2),x)`

output `int((a + b*cos(c + d*x))^(1/2), x)`

### 3.53 $\int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx$

3.53.1	Optimal result	355
3.53.2	Mathematica [A] (verified)	355
3.53.3	Rubi [A] (verified)	356
3.53.4	Maple [C] (verified)	357
3.53.5	Fricas [C] (verification not implemented)	358
3.53.6	Sympy [F]	358
3.53.7	Maxima [F]	358
3.53.8	Giac [F]	359
3.53.9	Mupad [B] (verification not implemented)	359

#### 3.53.1 Optimal result

Integrand size = 14, antiderivative size = 57

$$\int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx = \frac{2\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b \cos(c+dx)}}$$

output `2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/d/(a+b*cos(d*x+c))^(1/2)`

#### 3.53.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx = \frac{2\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b \cos(c+dx)}}$$

input `Integrate[1/Sqrt[a + b*Cos[c + d*x]],x]`

output `(2*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]])`

### 3.53.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{3142} \\
 & \frac{\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{\sqrt{a + b \cos(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin\left(c+dx + \frac{\pi}{2}\right)}{a+b}}} dx}{\sqrt{a + b \cos(c + dx)}} \\
 & \quad \downarrow \text{3140} \\
 & \frac{2\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d\sqrt{a + b \cos(c + dx)}}
 \end{aligned}$$

input `Int[1/Sqrt[a + b*Cos[c + d*x]],x]`

output `(2*sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*sqrt[a + b*Cos[c + d*x]])`

## 3.53.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3140 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

## 3.53.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.82 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.37

method	result	size
default	$\frac{2\sqrt{-\frac{2b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - a - b}{a + b}} \operatorname{am}^{-1}\left(\frac{dx}{2} + \frac{c}{2} \mid \frac{\sqrt{2}\sqrt{b}}{\sqrt{a+b}}\right)}{d\sqrt{2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b}}$	78

input `int(1/(a+cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

output `2/d/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)^(1/2)*(-(2*b*sin(1/2*d*x+1/2*c)^2-a-b)/(a+b))^(1/2)*InverseJacobiAM(1/2*d*x+1/2*c,2^(1/2)/(a+b)^(1/2)*b^(1/2))`

**3.53.5 Fracas [C] (verification not implemented)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 146, normalized size of antiderivative = 2.56

$$\int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx$$

$$= \frac{-i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}\left(\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8a^3 - 9ab^2)}{27b^3}, \frac{3b \cos(dx+c) + 3i b \sin(dx+c) + 2a}{3b}\right) + i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}\left(\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8a^3 - 9ab^2)}{27b^3}, \frac{3b \cos(dx+c) - 3i b \sin(dx+c) + 2a}{3b}\right)}{bd}$$

input `integrate(1/(a+b*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output `(-I*sqrt(2)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + I*sqrt(2)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b))/(b*d)`

**3.53.6 Sympy [F]**

$$\int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx$$

input `integrate(1/(a+b*cos(d*x+c))**(1/2),x)`

output `Integral(1/sqrt(a + b*cos(c + d*x)), x)`

**3.53.7 Maxima [F]**

$$\int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{1}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate(1/(a+b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(b*cos(d*x + c) + a), x)`

**3.53.8 Giac [F]**

$$\int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{1}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate(1/(a+b*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(b*cos(d*x + c) + a), x)`

**3.53.9 Mupad [B] (verification not implemented)**

Time = 13.93 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.91

$$\int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx = \frac{2 F\left(\frac{c}{2} + \frac{dx}{2} \middle| \frac{2b}{a+b}\right) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{d \sqrt{a + b \cos(c + dx)}}$$

input `int(1/(a + b*cos(c + d*x))^(1/2),x)`

output `(2*ellipticF(c/2 + (d*x)/2, (2*b)/(a + b))*((a + b*cos(c + d*x))/(a + b))^(1/2))/(d*(a + b*cos(c + d*x))^(1/2))`



### 3.54 $\int \frac{1}{(a+b \cos(c+dx))^{3/2}} dx$

3.54.1	Optimal result	360
3.54.2	Mathematica [A] (verified)	360
3.54.3	Rubi [A] (verified)	361
3.54.4	Maple [A] (verified)	363
3.54.5	Fricas [C] (verification not implemented)	363
3.54.6	Sympy [F]	364
3.54.7	Maxima [F]	364
3.54.8	Giac [F]	365
3.54.9	Mupad [F(-1)]	365

#### 3.54.1 Optimal result

Integrand size = 14, antiderivative size = 106

$$\int \frac{1}{(a+b \cos(c+dx))^{3/2}} dx = \frac{2\sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{(a^2-b^2) d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2b \sin(c+dx)}{(a^2-b^2) d \sqrt{a+b \cos(c+dx)}}$$

output `-2*b*sin(d*x+c)/(a^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)+2*(cos(1/2*d*x+1/2*c))^2  
)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/(a^2-b^2)/d/((a+b*cos(d*x+c))/(a+b))^(1/2)`

#### 3.54.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.78

$$\int \frac{1}{(a+b \cos(c+dx))^{3/2}} dx = \frac{2(a+b) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right) - 2b \sin(c+dx)}{(a-b)(a+b) d \sqrt{a+b \cos(c+dx)}}$$

input `Integrate[(a + b*Cos[c + d*x])^(-3/2), x]`

output `(2*(a + b)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticE[(c + d*x)/2, (2*b)/(a + b)] - 2*b*Sin[c + d*x])/((a - b)*(a + b)*d*Sqrt[a + b*Cos[c + d*x]])`

**3.54.3 Rubi [A] (verified)**

Time = 0.42 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 3143, 27, 3042, 3134, 3042, 3132}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + b \cos(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{3143} \\
 & -\frac{2 \int -\frac{1}{2} \sqrt{a + b \cos(c + dx)} dx}{a^2 - b^2} - \frac{2b \sin(c + dx)}{d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \sqrt{a + b \cos(c + dx)} dx}{a^2 - b^2} - \frac{2b \sin(c + dx)}{d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} dx}{a^2 - b^2} - \frac{2b \sin(c + dx)}{d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
 & \quad \downarrow \text{3134} \\
 & \frac{\sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2b \sin(c + dx)}{d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2b \sin(c + dx)}{d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
 & \quad \downarrow \text{3132} \\
 & \frac{2\sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{d(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2b \sin(c + dx)}{d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}}
 \end{aligned}$$

---

3.54.  $\int \frac{1}{(a+b \cos(c+dx))^{3/2}} dx$

input `Int[(a + b*Cos[c + d*x])^(-3/2),x]`

output `(2*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/((a^2 - b^2)*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*b*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]])`

### 3.54.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

### 3.54.4 Maple [A] (verified)

Time = 2.47 (sec) , antiderivative size = 217, normalized size of antiderivative = 2.05

method	result
default	$\frac{2 \left( 2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left( \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) b + E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{-\frac{2b}{a-b}}\right) \sqrt{-\frac{2b \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a-b} + \frac{a+b}{a-b} \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}} a - E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{-\frac{2b \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a-b} + \frac{a+b}{a-b} \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}}\right) \right)}{(a-b)(a+b) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2b \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a+b} d}$

input `int(1/(a+cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)`

output

$$\frac{-2*(2*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^2*b + \text{EllipticE}(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{1/2})) * (-2*b/(a-b)*\sin(1/2*d*x+1/2*c)^2 + (a+b)/(a-b))^{1/2} * (\sin(1/2*d*x+1/2*c)^2)^{1/2} * a - \text{EllipticE}(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{1/2}) * (-2*b/(a-b)*\sin(1/2*d*x+1/2*c)^2 + (a+b)/(a-b))^{1/2} * (\sin(1/2*d*x+1/2*c)^2)^{1/2} * b}{(a-b)/(a+b)/\sin(1/2*d*x+1/2*c)/(-2*b*\sin(1/2*d*x+1/2*c)^2 + a+b)^{1/2}/d}$$

### 3.54.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 482, normalized size of antiderivative = 4.55

$$\int \frac{1}{(a + b \cos(c + dx))^{3/2}} dx = \frac{6 \sqrt{b \cos(dx + c) + ab^2 \sin(dx + c)} + (i \sqrt{2} ab \cos(dx + c) + i \sqrt{2} a^2) \sqrt{b} \text{weierstrassPInverse}\left(\frac{4(4a^2 - 3b^2)}{3b^2}, -\right)}{d}$$

input `integrate(1/(a+b*cos(d*x+c))^(3/2),x, algorithm="fricas")`

output `-1/3*(6*sqrt(b*cos(d*x + c) + a)*b^2*sin(d*x + c) + (I*sqrt(2)*a*b*cos(d*x + c) + I*sqrt(2)*a^2)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + (-I*sqrt(2)*a*b*cos(d*x + c) - I*sqrt(2)*a^2)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*(I*sqrt(2)*b^2*cos(d*x + c) + I*sqrt(2)*a*b)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*(-I*sqrt(2)*b^2*cos(d*x + c) - I*sqrt(2)*a*b)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)))/((a^2*b^2 - b^4)*d*cos(d*x + c) + (a^3*b - a*b^3)*d)`

### 3.54.6 Sympy [F]

$$\int \frac{1}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{1}{(a + b \cos(c + dx))^{\frac{3}{2}}} dx$$

input `integrate(1/(a+b*cos(d*x+c))**(3/2), x)`

output `Integral((a + b*cos(c + d*x))**(-3/2), x)`

### 3.54.7 Maxima [F]

$$\int \frac{1}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{1}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate(1/(a+b*cos(d*x+c))^(3/2), x, algorithm="maxima")`

output `integrate((b*cos(d*x + c) + a)**(-3/2), x)`

**3.54.8 Giac [F]**

$$\int \frac{1}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{1}{(b \cos(dx + c) + a)^{3/2}} dx$$

input `integrate(1/(a+b*cos(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*cos(d*x + c) + a)^(-3/2), x)`

**3.54.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{1}{(a + b \cos(c + dx))^{3/2}} dx$$

input `int(1/(a + b*cos(c + d*x))^(3/2),x)`

output `int(1/(a + b*cos(c + d*x))^(3/2), x)`

### 3.55 $\int \frac{1}{(a+b \cos(c+dx))^{5/2}} dx$

3.55.1	Optimal result	366
3.55.2	Mathematica [A] (verified)	367
3.55.3	Rubi [A] (verified)	367
3.55.4	Maple [A] (verified)	371
3.55.5	Fricas [C] (verification not implemented)	372
3.55.6	Sympy [F]	373
3.55.7	Maxima [F]	373
3.55.8	Giac [F]	374
3.55.9	Mupad [F(-1)]	374

#### 3.55.1 Optimal result

Integrand size = 14, antiderivative size = 221

$$\int \frac{1}{(a+b \cos(c+dx))^{5/2}} dx = \frac{8a\sqrt{a+b \cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{3(a^2-b^2)^2 d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{3(a^2-b^2)d\sqrt{a+b \cos(c+dx)}} - \frac{2b \sin(c+dx)}{3(a^2-b^2)d(a+b \cos(c+dx))^{3/2}} - \frac{8ab \sin(c+dx)}{3(a^2-b^2)^2 d\sqrt{a+b \cos(c+dx)}}$$

output `-2/3*b*sin(d*x+c)/(a^2-b^2)/d/(a+b*cos(d*x+c))^(3/2)-8/3*a*b*sin(d*x+c)/(a^2-b^2)^2/d/(a+b*cos(d*x+c))^(1/2)+8/3*a*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/(a^2-b^2)^2/d/((a+b*cos(d*x+c))/(a+b))^(1/2)-2/3*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/(a^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)`

### 3.55.2 Mathematica [A] (verified)

Time = 1.03 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.71

$$\int \frac{1}{(a + b \cos(c + dx))^{5/2}} dx = \frac{8a(a + b)^2 \left(\frac{a+b \cos(c+dx)}{a+b}\right)^{3/2} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right) - 2(a - b)(a + b)^2 \left(\frac{a+b \cos(c+dx)}{a+b}\right)^{3/2}}{3(a - b)^2(a + b)^2 d}$$

input `Integrate[(a + b*Cos[c + d*x])^(-5/2), x]`

output `(8*a*(a + b)^2*((a + b*Cos[c + d*x])/(a + b))^(3/2)*EllipticE[(c + d*x)/2, (2*b)/(a + b)] - 2*(a - b)*(a + b)^2*((a + b*Cos[c + d*x])/(a + b))^(3/2)*EllipticF[(c + d*x)/2, (2*b)/(a + b)] + 2*b*(-5*a^2 + b^2 - 4*a*b*Cos[c + d*x])*Sin[c + d*x])/(3*(a - b)^2*(a + b)^2*d*(a + b*Cos[c + d*x])^(3/2))`

### 3.55.3 Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.05, number of steps used = 15, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.071$ , Rules used = {3042, 3143, 27, 3042, 3233, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(a + b \cos(c + dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2}} dx \\ & \quad \downarrow \text{3143} \\ & -\frac{2 \int -\frac{3a-b \cos(c+dx)}{2(a+b \cos(c+dx))^{3/2}} dx}{3(a^2 - b^2)} - \frac{2b \sin(c + dx)}{3d(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{3a-b \cos(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx}{3(a^2 - b^2)} - \frac{2b \sin(c + dx)}{3d(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\ & \quad \downarrow \text{3042} \end{aligned}$$

---

3.55.  $\int \frac{1}{(a+b \cos(c+dx))^{5/2}} dx$



$$\begin{aligned}
& \frac{\int \frac{3a-b \sin(c+dx+\frac{\pi}{2})}{(a+b \sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{3(a^2-b^2)} - \frac{2b \sin(c+dx)}{3d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3233} \\
& \frac{2 \int -\frac{3a^2+4b \cos(c+dx)a+b^2}{2\sqrt{a+b \cos(c+dx)}} dx}{3(a^2-b^2)} - \frac{8ab \sin(c+dx)}{d(a^2-b^2)\sqrt{a+b \cos(c+dx)}} - \frac{2b \sin(c+dx)}{3d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{3a^2+4b \cos(c+dx)a+b^2}{\sqrt{a+b \cos(c+dx)}} dx}{3(a^2-b^2)} - \frac{8ab \sin(c+dx)}{d(a^2-b^2)\sqrt{a+b \cos(c+dx)}} - \frac{2b \sin(c+dx)}{3d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{3a^2+4b \sin(c+dx+\frac{\pi}{2})a+b^2}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{3(a^2-b^2)} - \frac{8ab \sin(c+dx)}{d(a^2-b^2)\sqrt{a+b \cos(c+dx)}} - \frac{2b \sin(c+dx)}{3d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3231} \\
& \frac{4a \int \sqrt{a+b \cos(c+dx)} dx - (a^2-b^2) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}{3(a^2-b^2)} - \frac{8ab \sin(c+dx)}{d(a^2-b^2)\sqrt{a+b \cos(c+dx)}} - \\
& \quad \frac{2b \sin(c+dx)}{3d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{4a \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx - (a^2-b^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{3(a^2-b^2)} - \frac{8ab \sin(c+dx)}{d(a^2-b^2)\sqrt{a+b \cos(c+dx)}} - \\
& \quad \frac{2b \sin(c+dx)}{3d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3134} \\
& \frac{4a \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - (a^2-b^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{8ab \sin(c+dx)}{d(a^2-b^2)\sqrt{a+b \cos(c+dx)}} - \\
& \quad \frac{2b \sin(c+dx)}{3d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

---

3.55.  $\int \frac{1}{(a+b \cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \frac{4a\sqrt{a+b\cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}} dx - (a^2-b^2) \int \frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{8ab\sin(c+dx)}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \\
& \frac{3(a^2-b^2)}{2b\sin(c+dx)} \\
& \frac{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3132} \\
& \frac{8a\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right) - (a^2-b^2) \int \frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{8ab\sin(c+dx)}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \\
& \frac{3(a^2-b^2)}{2b\sin(c+dx)} \\
& \frac{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3142} \\
& \frac{8a\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right) - (a^2-b^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{8ab\sin(c+dx)}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \\
& \frac{3(a^2-b^2)}{2b\sin(c+dx)} \\
& \frac{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{8a\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right) - (a^2-b^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{8ab\sin(c+dx)}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \\
& \frac{3(a^2-b^2)}{2b\sin(c+dx)} \\
& \frac{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3140} \\
& \frac{8a\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right) - 2(a^2-b^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{8ab\sin(c+dx)}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \\
& \frac{3(a^2-b^2)}{2b\sin(c+dx)} \\
& \frac{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}
\end{aligned}$$

input `Int[(a + b*Cos[c + d*x])^(-5/2), x]`

$$3.55. \quad \int \frac{1}{(a+b\cos(c+dx))^{5/2}} dx$$

```
output (-2*b*Sin[c + d*x])/(3*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) + (((8*a*
Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[(a
+ b*Cos[c + d*x])/(a + b)]) - (2*(a^2 - b^2)*Sqrt[(a + b*Cos[c + d*x])/(a
+ b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]])
)/(a^2 - b^2) - (8*a*b*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x
]]))/(3*(a^2 - b^2))
```

### 3.55.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3132 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a
+ b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a,
b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3134 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a +
b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (
b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2
, 0] && !GtQ[a + b, 0]
```

```
rule 3140 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*S
qrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[
{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3142 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a
+ b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b)
+ (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && !GtQ[a + b, 0]
```

```
rule 3143 Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos
[c + d*x]*((a + b*SIN[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp
[1/((n + 1)*(a^2 - b^2)) Int[(a + b*SIN[c + d*x])^(n + 1)*Simp[a*(n + 1)
- b*(n + 2)*SIN[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

```
rule 3231 Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (
f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*SIN[e + f*x
]], x], x] + Simp[d/b Int[Sqrt[a + b*SIN[e + f*x]], x], x] /; FreeQ[{a, b
, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]
```

```
rule 3233 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*SIN[e +
f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2))
Int[(a + b*SIN[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(
m + 2)*SIN[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]
```

### 3.55.4 Maple [A] (verified)

Time = 4.48 (sec) , antiderivative size = 489, normalized size of antiderivative = 2.21

method	result
default	$\frac{\sqrt{-(-2b(\cos^2(\frac{dx}{2} + \frac{c}{2})) - a + b)(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}{\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2}))b + (a+b)(\sin^2(\frac{dx}{2} + \frac{c}{2}))}} + \frac{16(\sin^2(\frac{dx}{2} + \frac{c}{2}))}{3b(a-b)(a+b)(\cos^2(\frac{dx}{2} + \frac{c}{2}) + \frac{a-b}{2b})^2} + \frac{16(\sin^2(\frac{dx}{2} + \frac{c}{2}))}{3(a-b)^2(a+b)^2\sqrt{-(-2b(\cos^2(\frac{dx}{2} + \frac{c}{2})) - a + b)(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}$

```
input int(1/(a+cos(d*x+c)*b)^(5/2), x, method=_RETURNVERBOSE)
```

output

$$\begin{aligned}
 & -(-(-2*b*\cos(1/2*d*x+1/2*c)^2-a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(1/3/b/(a-b) \\
 & )/(a+b)*\cos(1/2*d*x+1/2*c)*(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/ \\
 & 2*c)^2)^{(1/2)}/(\cos(1/2*d*x+1/2*c)^2+1/2*(a-b)/b)^2+16/3*\sin(1/2*d*x+1/2*c) \\
 & ^2*b/(a-b)^2/(a+b)^2*\cos(1/2*d*x+1/2*c)*a/(-(-2*b*\cos(1/2*d*x+1/2*c)^2-a+b) \\
 & )*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+2*(3*a-b)/(3*a^3+3*a^2*b-3*a*b^2-3*b^3)*( \sin \\
 & (1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)}/(-2* \\
 & \sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticF(\cos(1/2 \\
 & *d*x+1/2*c), (-2*b/(a-b))^{(1/2)})-8/3*a/(a-b)/(a+b)^2*(\sin(1/2*d*x+1/2*c)^2) \\
 & ^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c) \\
 & ^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(EllipticF(\cos(1/2*d*x+1/2*c), (-2*b \\
 & /a-b))^{(1/2)})-EllipticE(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{(1/2)}))/\sin(1/2* \\
 & d*x+1/2*c)/(-2*b*\sin(1/2*d*x+1/2*c)^2+a+b)^{(1/2)}/d
 \end{aligned}$$

### 3.55.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 692, normalized size of antiderivative = 3.13

$$\int \frac{1}{(a+b\cos(c+dx))^{5/2}} dx = \frac{6(4ab^3\cos(dx+c)+5a^2b^2-b^4)\sqrt{b\cos(dx+c)+a}\sin(dx+c)-(\sqrt{2}(-ia^2b^2-3ib^4)\cos(dx+c)^2-}{$$

input `integrate(1/(a+b*cos(d*x+c))^(5/2),x, algorithm="fricas")`

output

```
-1/9*(6*(4*a*b^3*cos(d*x + c) + 5*a^2*b^2 - b^4)*sqrt(b*cos(d*x + c) + a)*
sin(d*x + c) - (sqrt(2)*(-I*a^2*b^2 - 3*I*b^4)*cos(d*x + c)^2 - 2*sqrt(2)*
(I*a^3*b + 3*I*a*b^3)*cos(d*x + c) + sqrt(2)*(-I*a^4 - 3*I*a^2*b^2))*sqrt(
b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^
3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) - (sqrt(2)*(I*a^2*
b^2 + 3*I*b^4)*cos(d*x + c)^2 - 2*sqrt(2)*(-I*a^3*b - 3*I*a*b^3)*cos(d*x +
c) + sqrt(2)*(I*a^4 + 3*I*a^2*b^2))*sqrt(b)*weierstrassPInverse(4/3*(4*a^
2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b
*sin(d*x + c) + 2*a)/b) + 12*(-I*sqrt(2)*a*b^3*cos(d*x + c)^2 - 2*I*sqrt(2
)*a^2*b^2*cos(d*x + c) - I*sqrt(2)*a^3*b)*sqrt(b)*weierstrassZeta(4/3*(4*a
^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a
^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*
b*sin(d*x + c) + 2*a)/b)) + 12*(I*sqrt(2)*a*b^3*cos(d*x + c)^2 + 2*I*sqrt(
2)*a^2*b^2*cos(d*x + c) + I*sqrt(2)*a^3*b)*sqrt(b)*weierstrassZeta(4/3*(4*
a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*
a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I
*b*sin(d*x + c) + 2*a)/b)))/((a^4*b^3 - 2*a^2*b^5 + b^7)*d*cos(d*x + c)^2
+ 2*(a^5*b^2 - 2*a^3*b^4 + a*b^6)*d*cos(d*x + c) + (a^6*b - 2*a^4*b^3 + a^
2*b^5)*d)
```

### 3.55.6 Sympy [F]

$$\int \frac{1}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{1}{(a + b \cos(c + dx))^{\frac{5}{2}}} dx$$

input `integrate(1/(a+b*cos(d*x+c))**(5/2), x)`

output `Integral((a + b*cos(c + d*x))**(-5/2), x)`

### 3.55.7 Maxima [F]

$$\int \frac{1}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{1}{(b \cos(dx + c) + a)^{\frac{5}{2}}} dx$$

input `integrate(1/(a+b*cos(d*x+c))^(5/2), x, algorithm="maxima")`

output `integrate((b*cos(d*x + c) + a)**(-5/2), x)`

**3.55.8 Giac [F]**

$$\int \frac{1}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{1}{(b \cos(dx + c) + a)^{5/2}} dx$$

input `integrate(1/(a+b*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*cos(d*x + c) + a)^(-5/2), x)`

**3.55.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{1}{(a + b \cos(c + dx))^{5/2}} dx$$

input `int(1/(a + b*cos(c + d*x))^(5/2),x)`

output `int(1/(a + b*cos(c + d*x))^(5/2), x)`

### 3.56 $\int (a + b \cos(c + dx))^{4/3} dx$

3.56.1	Optimal result . . . . .	375
3.56.2	Mathematica [B] (verified) . . . . .	375
3.56.3	Rubi [A] (verified) . . . . .	376
3.56.4	Maple [F] . . . . .	378
3.56.5	Fricas [F] . . . . .	378
3.56.6	Sympy [F] . . . . .	378
3.56.7	Maxima [F] . . . . .	379
3.56.8	Giac [F] . . . . .	379
3.56.9	Mupad [F(-1)] . . . . .	379

#### 3.56.1 Optimal result

Integrand size = 14, antiderivative size = 108

$$\int (a + b \cos(c + dx))^{4/3} dx = \frac{\sqrt{2}(a + b) \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -\frac{4}{3}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx)), \frac{b(1 - \cos(c + dx))}{a + b}\right) \sqrt[3]{a + b \cos(c + dx)} \sin(c + dx)}{d \sqrt{1 + \cos(c + dx)} \sqrt[3]{\frac{a + b \cos(c + dx)}{a + b}}}$$

```
output (a+b)*AppellF1(1/2,-4/3,1/2,3/2,b*(1-cos(d*x+c))/(a+b),1/2-1/2*cos(d*x+c))
*(a+b*cos(d*x+c))^(1/3)*sin(d*x+c)*2^(1/2)/d/((a+b*cos(d*x+c))/(a+b))^(1/3)
)/(1+cos(d*x+c))^(1/2)
```

#### 3.56.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 246 vs. 2(108) = 216.

Time = 2.07 (sec) , antiderivative size = 246, normalized size of antiderivative = 2.28

$$\int (a + b \cos(c + dx))^{4/3} dx = \frac{3 \sqrt[3]{a + b \cos(c + dx)} \csc(c + dx) \left( 4(-a^2 + b^2) \operatorname{AppellF1}\left(\frac{1}{3}, \frac{1}{2}, \frac{1}{2}, \frac{4}{3}, \frac{a + b \cos(c + dx)}{a - b}, \frac{a + b \cos(c + dx)}{a + b}\right) \sqrt{-\frac{b(-1 + \cos(c + dx))}{a + b}} \right)}{d}$$



input `Integrate[(a + b*Cos[c + d*x])^(4/3),x]`

output `(-3*(a + b*Cos[c + d*x])^(1/3)*Csc[c + d*x]*(4*(-a^2 + b^2)*AppellF1[1/3, 1/2, 1/2, 4/3, (a + b*Cos[c + d*x])/(a - b), (a + b*Cos[c + d*x])/(a + b)]*Sqrt[-((b*(-1 + Cos[c + d*x]))/(a + b))]*Sqrt[-((b*(1 + Cos[c + d*x]))/(a - b))] + 5*a*AppellF1[4/3, 1/2, 1/2, 7/3, (a + b*Cos[c + d*x])/(a - b), (a + b*Cos[c + d*x])/(a + b)]*Sqrt[-((b*(-1 + Cos[c + d*x]))/(a + b))]*Sqrt[-((b*(1 + Cos[c + d*x]))/(a - b))]*(a + b*Cos[c + d*x]) - 4*b^2*Sin[c + d*x]^2)/(16*b*d)`

### 3.56.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + b \cos(c + dx))^{4/3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( a + b \sin\left(c + dx + \frac{\pi}{2}\right) \right)^{4/3} dx \\
 & \quad \downarrow \text{3144} \\
 & - \frac{\sin(c + dx) \int \frac{(a + b \cos(c + dx))^{4/3}}{\sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}} d \cos(c + dx)}{d \sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}} \\
 & \quad \downarrow \text{156} \\
 & - \frac{(a + b) \sin(c + dx) \sqrt[3]{a + b \cos(c + dx)} \int \frac{\left(\frac{a}{a+b} + \frac{b \cos(c + dx)}{a+b}\right)^{4/3}}{\sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}} d \cos(c + dx)}{d \sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1} \sqrt[3]{\frac{a + b \cos(c + dx)}{a + b}}} \\
 & \quad \downarrow \text{155} \\
 & \frac{\sqrt{2}(a + b) \sin(c + dx) \sqrt[3]{a + b \cos(c + dx)} \text{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -\frac{4}{3}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx)), \frac{b(1 - \cos(c + dx))}{a + b}\right)}{d \sqrt{\cos(c + dx) + 1} \sqrt[3]{\frac{a + b \cos(c + dx)}{a + b}}}
 \end{aligned}$$

input `Int[(a + b*cos[c + d*x])^(4/3),x]`

output `(Sqrt[2]*(a + b)*AppellF1[1/2, 1/2, -4/3, 3/2, (1 - Cos[c + d*x])/2, (b*(1 - Cos[c + d*x]))/(a + b)]*(a + b*cos[c + d*x])^(1/3)*Sin[c + d*x])/(d*Sqrt[1 + Cos[c + d*x]]*((a + b*cos[c + d*x])/(a + b))^(1/3))`

### 3.56.3.1 Defintions of rubi rules used

rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p)*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`

rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*((b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3144 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Simp[Cos[c + d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]`

**3.56.4 Maple [F]**

$$\int (a + \cos(dx + c)b)^{\frac{4}{3}} dx$$

input `int((a+cos(d*x+c)*b)^(4/3),x)`

output `int((a+cos(d*x+c)*b)^(4/3),x)`

**3.56.5 Fricas [F]**

$$\int (a + b \cos(c + dx))^{\frac{4}{3}} dx = \int (b \cos(dx + c) + a)^{\frac{4}{3}} dx$$

input `integrate((a+b*cos(d*x+c))^(4/3),x, algorithm="fricas")`

output `integral((b*cos(d*x + c) + a)^(4/3), x)`

**3.56.6 Sympy [F]**

$$\int (a + b \cos(c + dx))^{\frac{4}{3}} dx = \int (a + b \cos(c + dx))^{\frac{4}{3}} dx$$

input `integrate((a+b*cos(d*x+c))**(4/3),x)`

output `Integral((a + b*cos(c + d*x))**(4/3), x)`

**3.56.7 Maxima [F]**

$$\int (a + b \cos(c + dx))^{4/3} dx = \int (b \cos(dx + c) + a)^{4/3} dx$$

input `integrate((a+b*cos(d*x+c))^(4/3),x, algorithm="maxima")`

output `integrate((b*cos(d*x + c) + a)^(4/3), x)`

**3.56.8 Giac [F]**

$$\int (a + b \cos(c + dx))^{4/3} dx = \int (b \cos(dx + c) + a)^{4/3} dx$$

input `integrate((a+b*cos(d*x+c))^(4/3),x, algorithm="giac")`

output `integrate((b*cos(d*x + c) + a)^(4/3), x)`

**3.56.9 Mupad [F(-1)]**

Timed out.

$$\int (a + b \cos(c + dx))^{4/3} dx = \int (a + b \cos(c + dx))^{4/3} dx$$

input `int((a + b*cos(c + d*x))^(4/3),x)`

output `int((a + b*cos(c + d*x))^(4/3), x)`

### 3.57 $\int (a + b \cos(c + dx))^{2/3} dx$

3.57.1	Optimal result . . . . .	380
3.57.2	Mathematica [A] (verified) . . . . .	380
3.57.3	Rubi [A] (verified) . . . . .	381
3.57.4	Maple [F] . . . . .	382
3.57.5	Fricas [F] . . . . .	383
3.57.6	Sympy [F] . . . . .	383
3.57.7	Maxima [F] . . . . .	383
3.57.8	Giac [F] . . . . .	384
3.57.9	Mupad [F(-1)] . . . . .	384

#### 3.57.1 Optimal result

Integrand size = 14, antiderivative size = 105

$$\int (a + b \cos(c + dx))^{2/3} dx = \frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -\frac{2}{3}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx)), \frac{b(1 - \cos(c + dx))}{a + b}\right) (a + b \cos(c + dx))^{2/3} \sin(c + dx)}{d \sqrt{1 + \cos(c + dx)} \left(\frac{a + b \cos(c + dx)}{a + b}\right)^{2/3}}$$

```
output AppellF1(1/2, -2/3, 1/2, 3/2, b*(1-cos(d*x+c))/(a+b), 1/2-1/2*cos(d*x+c))*(a+b*cos(d*x+c))^(2/3)*sin(d*x+c)*2^(1/2)/d/((a+b*cos(d*x+c))/(a+b))^(2/3)/(1+cos(d*x+c))^(1/2)
```

#### 3.57.2 Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.12

$$\int (a + b \cos(c + dx))^{2/3} dx = \frac{3 \operatorname{AppellF1}\left(\frac{5}{3}, \frac{1}{2}, \frac{1}{2}, \frac{8}{3}, \frac{a + b \cos(c + dx)}{a - b}, \frac{a + b \cos(c + dx)}{a + b}\right) \sqrt{-\frac{b(-1 + \cos(c + dx))}{a + b}} \sqrt{\frac{b(1 + \cos(c + dx))}{-a + b}} (a + b \cos(c + dx))^{5/3} \csc(c + dx)}{5bd}$$

```
input Integrate[(a + b*Cos[c + d*x])^(2/3), x]
```

output  $(-3*\text{AppellF1}[5/3, 1/2, 1/2, 8/3, (a + b*\text{Cos}[c + d*x])/(a - b), (a + b*\text{Cos}[c + d*x])/(a + b)]*\text{Sqrt}[-((b*(-1 + \text{Cos}[c + d*x]))/(a + b))]*\text{Sqrt}[(b*(1 + \text{Cos}[c + d*x]))/(-a + b)]*(a + b*\text{Cos}[c + d*x])^{(5/3)}*\text{Csc}[c + d*x])/(5*b*d)$

### 3.57.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \cos(c + dx))^{2/3} dx \\ & \quad \downarrow \text{3042} \\ & \int \left( a + b \sin\left(c + dx + \frac{\pi}{2}\right) \right)^{2/3} dx \\ & \quad \downarrow \text{3144} \\ & - \frac{\sin(c + dx) \int \frac{(a + b \cos(c + dx))^{2/3}}{\sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}} d \cos(c + dx)}{d \sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}} \\ & \quad \downarrow \text{156} \\ & - \frac{\sin(c + dx)(a + b \cos(c + dx))^{2/3} \int \frac{\left(\frac{a}{a+b} + \frac{b \cos(c + dx)}{a+b}\right)^{2/3}}{\sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}} d \cos(c + dx)}{d \sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1} \left(\frac{a + b \cos(c + dx)}{a + b}\right)^{2/3}} \\ & \quad \downarrow \text{155} \\ & \frac{\sqrt{2} \sin(c + dx)(a + b \cos(c + dx))^{2/3} \text{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -\frac{2}{3}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx)), \frac{b(1 - \cos(c + dx))}{a + b}\right)}{d \sqrt{\cos(c + dx) + 1} \left(\frac{a + b \cos(c + dx)}{a + b}\right)^{2/3}} \end{aligned}$$

input  $\text{Int}[(a + b*\text{Cos}[c + d*x])^{(2/3)}, x]$

output  $(\text{Sqrt}[2]*\text{AppellF1}[1/2, 1/2, -2/3, 3/2, (1 - \text{Cos}[c + d*x])/2, (b*(1 - \text{Cos}[c + d*x]))/(a + b)]*(a + b*\text{Cos}[c + d*x])^{(2/3)}*\text{Sin}[c + d*x])/(d*\text{Sqrt}[1 + \text{Cos}[c + d*x]]*((a + b*\text{Cos}[c + d*x])/(a + b))^{(2/3)})$

## 3.57.3.1 Defintions of rubi rules used

```
rule 155 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))
^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simp
lify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplerQ[e + f*x, a + b*x])
```

```
rule 156 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))
^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p
]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Si
mp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c,
d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] &
& GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3144 Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x
)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]
```

## 3.57.4 Maple [F]

$$\int (a + \cos(dx + c)b)^{\frac{2}{3}} dx$$

```
input int((a+cos(d*x+c)*b)^(2/3),x)
```

```
output int((a+cos(d*x+c)*b)^(2/3),x)
```

**3.57.5 Fricas [F]**

$$\int (a + b \cos(c + dx))^{2/3} dx = \int (b \cos(dx + c) + a)^{2/3} dx$$

input `integrate((a+b*cos(d*x+c))^(2/3),x, algorithm="fricas")`

output `integral((b*cos(d*x + c) + a)^(2/3), x)`

**3.57.6 Sympy [F]**

$$\int (a + b \cos(c + dx))^{2/3} dx = \int (a + b \cos(c + dx))^{2/3} dx$$

input `integrate((a+b*cos(d*x+c))**(2/3),x)`

output `Integral((a + b*cos(c + d*x))**(2/3), x)`

**3.57.7 Maxima [F]**

$$\int (a + b \cos(c + dx))^{2/3} dx = \int (b \cos(dx + c) + a)^{2/3} dx$$

input `integrate((a+b*cos(d*x+c))^(2/3),x, algorithm="maxima")`

output `integrate((b*cos(d*x + c) + a)^(2/3), x)`



**3.57.8 Giac [F]**

$$\int (a + b \cos(c + dx))^{2/3} dx = \int (b \cos(dx + c) + a)^{2/3} dx$$

input `integrate((a+b*cos(d*x+c))^(2/3),x, algorithm="giac")`

output `integrate((b*cos(d*x + c) + a)^(2/3), x)`

**3.57.9 Mupad [F(-1)]**

Timed out.

$$\int (a + b \cos(c + dx))^{2/3} dx = \int (a + b \cos(c + dx))^{2/3} dx$$

input `int((a + b*cos(c + d*x))^(2/3),x)`

output `int((a + b*cos(c + d*x))^(2/3), x)`

### 3.58 $\int \sqrt[3]{a + b \cos(c + dx)} dx$

3.58.1	Optimal result . . . . .	385
3.58.2	Mathematica [A] (verified) . . . . .	385
3.58.3	Rubi [A] (verified) . . . . .	386
3.58.4	Maple [F] . . . . .	388
3.58.5	Fricas [F] . . . . .	388
3.58.6	Sympy [F] . . . . .	388
3.58.7	Maxima [F] . . . . .	389
3.58.8	Giac [F] . . . . .	389
3.58.9	Mupad [F(-1)] . . . . .	389

#### 3.58.1 Optimal result

Integrand size = 14, antiderivative size = 105

$$\int \sqrt[3]{a + b \cos(c + dx)} dx = \frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -\frac{1}{3}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx)), \frac{b(1 - \cos(c + dx))}{a + b}\right) \sqrt[3]{a + b \cos(c + dx)} \sin(c + dx)}{d \sqrt{1 + \cos(c + dx)} \sqrt[3]{\frac{a + b \cos(c + dx)}{a + b}}}$$

output `AppellF1(1/2, -1/3, 1/2, 3/2, b*(1-cos(d*x+c))/(a+b), 1/2-1/2*cos(d*x+c))*(a+b*cos(d*x+c))^(1/3)*sin(d*x+c)*2^(1/2)/d/((a+b*cos(d*x+c))/(a+b))^(1/3)/(1+cos(d*x+c))^(1/2)`

#### 3.58.2 Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.12

$$\int \sqrt[3]{a + b \cos(c + dx)} dx = \frac{3 \operatorname{AppellF1}\left(\frac{4}{3}, \frac{1}{2}, \frac{1}{2}, \frac{7}{3}, \frac{a + b \cos(c + dx)}{a - b}, \frac{a + b \cos(c + dx)}{a + b}\right) \sqrt{-\frac{b(-1 + \cos(c + dx))}{a + b}} \sqrt{\frac{b(1 + \cos(c + dx))}{-a + b}} (a + b \cos(c + dx))^{4/3}}{4bd}$$

input `Integrate[(a + b*Cos[c + d*x])^(1/3), x]`

output  $(-3*\text{AppellF1}[4/3, 1/2, 1/2, 7/3, (a + b*\text{Cos}[c + d*x])/(a - b), (a + b*\text{Cos}[c + d*x])/(a + b)]*\text{Sqrt}[-(b*(-1 + \text{Cos}[c + d*x]))/(a + b)]*\text{Sqrt}[(b*(1 + \text{Cos}[c + d*x]))/(-a + b)]*(a + b*\text{Cos}[c + d*x])^{(4/3)}*\text{Csc}[c + d*x])/(4*b*d)$

### 3.58.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt[3]{a + b \cos(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt[3]{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} dx \\ & \quad \downarrow \text{3144} \\ & \frac{\sin(c + dx) \int \frac{\sqrt[3]{a + b \cos(c + dx)}}{\sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}} d \cos(c + dx)}{d \sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}} \\ & \quad \downarrow \text{156} \\ & \frac{\sin(c + dx) \sqrt[3]{a + b \cos(c + dx)} \int \frac{\sqrt[3]{\frac{a}{a + b} + \frac{b \cos(c + dx)}{a + b}}}{\sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}} d \cos(c + dx)}{d \sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1} \sqrt[3]{\frac{a + b \cos(c + dx)}{a + b}}} \\ & \quad \downarrow \text{155} \\ & \frac{\sqrt{2} \sin(c + dx) \sqrt[3]{a + b \cos(c + dx)} \text{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -\frac{1}{3}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx)), \frac{b(1 - \cos(c + dx))}{a + b}\right)}{d \sqrt{\cos(c + dx) + 1} \sqrt[3]{\frac{a + b \cos(c + dx)}{a + b}}} \end{aligned}$$

input  $\text{Int}[(a + b*\text{Cos}[c + d*x])^{(1/3)}, x]$

```
output (Sqrt[2]*AppellF1[1/2, 1/2, -1/3, 3/2, (1 - Cos[c + d*x])/2, (b*(1 - Cos[c
+ d*x]))/(a + b)]*(a + b*Cos[c + d*x])^(1/3)*Sin[c + d*x]/(d*Sqrt[1 + Co
s[c + d*x]]*((a + b*Cos[c + d*x])/(a + b))^(1/3))
```

### 3.58.3.1 Defintions of rubi rules used

```
rule 155 Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simp
lify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplerQ[e + f*x, a + b*x])
```

```
rule 156 Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p
]*b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Si
mp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c,
d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] &
& GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3144 Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x
)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]
```

**3.58.4 Maple [F]**

$$\int (a + \cos(dx + c)b)^{\frac{1}{3}} dx$$

input `int((a+cos(d*x+c)*b)^(1/3),x)`

output `int((a+cos(d*x+c)*b)^(1/3),x)`

**3.58.5 Fricas [F]**

$$\int \sqrt[3]{a + b \cos(c + dx)} dx = \int (b \cos(dx + c) + a)^{\frac{1}{3}} dx$$

input `integrate((a+b*cos(d*x+c))^(1/3),x, algorithm="fricas")`

output `integral((b*cos(d*x + c) + a)^(1/3), x)`

**3.58.6 Sympy [F]**

$$\int \sqrt[3]{a + b \cos(c + dx)} dx = \int \sqrt[3]{a + b \cos(c + dx)} dx$$

input `integrate((a+b*cos(d*x+c))**(1/3),x)`

output `Integral((a + b*cos(c + d*x))**(1/3), x)`

**3.58.7 Maxima [F]**

$$\int \sqrt[3]{a + b \cos(c + dx)} dx = \int (b \cos(dx + c) + a)^{\frac{1}{3}} dx$$

input `integrate((a+b*cos(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate((b*cos(d*x + c) + a)^(1/3), x)`

**3.58.8 Giac [F]**

$$\int \sqrt[3]{a + b \cos(c + dx)} dx = \int (b \cos(dx + c) + a)^{\frac{1}{3}} dx$$

input `integrate((a+b*cos(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate((b*cos(d*x + c) + a)^(1/3), x)`

**3.58.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt[3]{a + b \cos(c + dx)} dx = \int (a + b \cos(c + dx))^{1/3} dx$$

input `int((a + b*cos(c + d*x))^(1/3),x)`

output `int((a + b*cos(c + d*x))^(1/3), x)`

**3.59**  $\int \frac{1}{\sqrt[3]{a + b \cos(c + dx)}} dx$

3.59.1 Optimal result . . . . . 390  
 3.59.2 Mathematica [A] (verified) . . . . . 390  
 3.59.3 Rubi [A] (verified) . . . . . 391  
 3.59.4 Maple [F] . . . . . 393  
 3.59.5 Fricas [F] . . . . . 393  
 3.59.6 Sympy [F] . . . . . 393  
 3.59.7 Maxima [F] . . . . . 394  
 3.59.8 Giac [F] . . . . . 394  
 3.59.9 Mupad [F(-1)] . . . . . 394

**3.59.1 Optimal result**

Integrand size = 14, antiderivative size = 105

$$\int \frac{1}{\sqrt[3]{a + b \cos(c + dx)}} dx = \frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{3}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx)), \frac{b(1 - \cos(c + dx))}{a + b}\right) \sqrt[3]{\frac{a + b \cos(c + dx)}{a + b}} \sin(c + dx)}{d \sqrt{1 + \cos(c + dx)} \sqrt[3]{a + b \cos(c + dx)}}$$

output `AppellF1(1/2, 1/3, 1/2, 3/2, b*(1-cos(d*x+c))/(a+b), 1/2-1/2*cos(d*x+c))*((a+b*cos(d*x+c))/(a+b))^(1/3)*sin(d*x+c)*2^(1/2)/d/(a+b*cos(d*x+c))^(1/3)/(1+cos(d*x+c))^(1/2)`

**3.59.2 Mathematica [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.12

$$\int \frac{1}{\sqrt[3]{a + b \cos(c + dx)}} dx = \frac{3 \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{2}, \frac{1}{2}, \frac{5}{3}, \frac{a + b \cos(c + dx)}{a - b}, \frac{a + b \cos(c + dx)}{a + b}\right) \sqrt{-\frac{b(-1 + \cos(c + dx))}{a + b}} \sqrt{\frac{b(1 + \cos(c + dx))}{-a + b}} (a + b \cos(c + dx))^{2/3}}{2bd}$$

input `Integrate[(a + b*Cos[c + d*x])^(-1/3), x]`

---

3.59.  $\int \frac{1}{\sqrt[3]{a + b \cos(c + dx)}} dx$

output  $(-3*\text{AppellF1}[2/3, 1/2, 1/2, 5/3, (a + b*\text{Cos}[c + d*x])/(a - b), (a + b*\text{Cos}[c + d*x])/(a + b)]*\text{Sqrt}[-(b*(-1 + \text{Cos}[c + d*x]))/(a + b)]*\text{Sqrt}[(b*(1 + \text{Cos}[c + d*x]))/(-a + b)]*(a + b*\text{Cos}[c + d*x])^{2/3}*\text{Csc}[c + d*x])/(2*b*d)$

### 3.59.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt[3]{a + b \cos(c + dx)}} dx$$

↓ 3042

$$\int \frac{1}{\sqrt[3]{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx$$

↓ 3144

$$\frac{\sin(c + dx) \int \frac{1}{\sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1} \sqrt[3]{a + b \cos(c + dx)}} d \cos(c + dx)}{d \sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}}$$

↓ 156

$$\frac{\sin(c + dx) \sqrt[3]{\frac{a + b \cos(c + dx)}{a + b}} \int \frac{1}{\sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1} \sqrt[3]{\frac{a}{a + b} + \frac{b \cos(c + dx)}{a + b}}} d \cos(c + dx)}{d \sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1} \sqrt[3]{a + b \cos(c + dx)}}$$

↓ 155

$$\frac{\sqrt{2} \sin(c + dx) \sqrt[3]{\frac{a + b \cos(c + dx)}{a + b}} \text{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{3}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx)), \frac{b(1 - \cos(c + dx))}{a + b}\right)}{d \sqrt{\cos(c + dx) + 1} \sqrt[3]{a + b \cos(c + dx)}}$$

input  $\text{Int}[(a + b*\text{Cos}[c + d*x])^{-1/3}, x]$

---

3.59.  $\int \frac{1}{\sqrt[3]{a + b \cos(c + dx)}} dx$



```
output (Sqrt[2]*AppellF1[1/2, 1/2, 1/3, 3/2, (1 - Cos[c + d*x])/2, (b*(1 - Cos[c
+ d*x]))/(a + b)]*((a + b*Cos[c + d*x])/(a + b))^(1/3)*Sin[c + d*x])/(d*Sq
rt[1 + Cos[c + d*x]]*(a + b*Cos[c + d*x])^(1/3))
```

### 3.59.3.1 Defintions of rubi rules used

```
rule 155 Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simp
lify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplerQ[e + f*x, a + b*x])
```

```
rule 156 Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p
]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Si
mp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c,
d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] &
& GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3144 Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x
)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]
```

**3.59.4 Maple [F]**

$$\int \frac{1}{(a + \cos(dx + c)b)^{\frac{1}{3}}} dx$$

input `int(1/(a+cos(d*x+c)*b)^(1/3),x)`

output `int(1/(a+cos(d*x+c)*b)^(1/3),x)`

**3.59.5 Fricas [F]**

$$\int \frac{1}{\sqrt[3]{a + b \cos(c + dx)}} dx = \int \frac{1}{(b \cos(dx + c) + a)^{\frac{1}{3}}} dx$$

input `integrate(1/(a+b*cos(d*x+c))^(1/3),x, algorithm="fricas")`

output `integral((b*cos(d*x + c) + a)^(-1/3), x)`

**3.59.6 Sympy [F]**

$$\int \frac{1}{\sqrt[3]{a + b \cos(c + dx)}} dx = \int \frac{1}{\sqrt[3]{a + b \cos(c + dx)}} dx$$

input `integrate(1/(a+b*cos(d*x+c))**(1/3),x)`

output `Integral((a + b*cos(c + d*x))**(-1/3), x)`

**3.59.7 Maxima [F]**

$$\int \frac{1}{\sqrt[3]{a + b \cos(c + dx)}} dx = \int \frac{1}{(b \cos(dx + c) + a)^{\frac{1}{3}}} dx$$

input `integrate(1/(a+b*cos(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate((b*cos(d*x + c) + a)^(-1/3), x)`

**3.59.8 Giac [F]**

$$\int \frac{1}{\sqrt[3]{a + b \cos(c + dx)}} dx = \int \frac{1}{(b \cos(dx + c) + a)^{\frac{1}{3}}} dx$$

input `integrate(1/(a+b*cos(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate((b*cos(d*x + c) + a)^(-1/3), x)`

**3.59.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt[3]{a + b \cos(c + dx)}} dx = \int \frac{1}{(a + b \cos(c + dx))^{\frac{1}{3}}} dx$$

input `int(1/(a + b*cos(c + d*x))^(1/3),x)`

output `int(1/(a + b*cos(c + d*x))^(1/3), x)`

### 3.60 $\int \frac{1}{(a+b \cos(c+dx))^{2/3}} dx$

3.60.1	Optimal result	395
3.60.2	Mathematica [A] (verified)	395
3.60.3	Rubi [A] (verified)	396
3.60.4	Maple [F]	397
3.60.5	Fricas [F]	398
3.60.6	Sympy [F]	398
3.60.7	Maxima [F]	398
3.60.8	Giac [F]	399
3.60.9	Mupad [F(-1)]	399

#### 3.60.1 Optimal result

Integrand size = 14, antiderivative size = 105

$$\int \frac{1}{(a+b \cos(c+dx))^{2/3}} dx = \frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, \frac{2}{3}, \frac{3}{2}, \frac{1}{2}(1-\cos(c+dx)), \frac{b(1-\cos(c+dx))}{a+b}\right) \left(\frac{a+b \cos(c+dx)}{a+b}\right)^{2/3}}{d \sqrt{1+\cos(c+dx)} (a+b \cos(c+dx))^{2/3}}$$

output `AppellF1(1/2,2/3,1/2,3/2,b*(1-cos(d*x+c))/(a+b),1/2-1/2*cos(d*x+c))*((a+b*cos(d*x+c))/(a+b))^(2/3)*sin(d*x+c)*2^(1/2)/d/(a+b*cos(d*x+c))^(2/3)/(1+cos(d*x+c))^(1/2)`

#### 3.60.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.10

$$\int \frac{1}{(a+b \cos(c+dx))^{2/3}} dx = \frac{3 \operatorname{AppellF1}\left(\frac{1}{3}, \frac{1}{2}, \frac{1}{2}, \frac{4}{3}, \frac{a+b \cos(c+dx)}{a-b}, \frac{a+b \cos(c+dx)}{a+b}\right) \sqrt{-\frac{b(-1+\cos(c+dx))}{a+b}} \sqrt{\frac{b(1+\cos(c+dx))}{-a+b}} \sqrt[3]{a+b \cos(c+dx)} \operatorname{csc}(c+dx)}{bd}$$

input `Integrate[(a + b*Cos[c + d*x])^(-2/3),x]`

output `(-3*AppellF1[1/3, 1/2, 1/2, 4/3, (a + b*Cos[c + d*x])/(a - b), (a + b*Cos[c + d*x])/(a + b)]*Sqrt[-((b*(-1 + Cos[c + d*x]))/(a + b))]*Sqrt[(b*(1 + Cos[c + d*x]))/(-a + b)]*(a + b*Cos[c + d*x])^(1/3)*Csc[c + d*x])/(b*d)`

### 3.60.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + b \cos(c + dx))^{2/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a + b \sin(c + dx + \frac{\pi}{2}))^{2/3}} dx \\
 & \quad \downarrow \text{3144} \\
 & \frac{\sin(c + dx) \int \frac{1}{\sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1} (a + b \cos(c + dx))^{2/3}} d \cos(c + dx)}{d \sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}} \\
 & \quad \downarrow \text{156} \\
 & \frac{\sin(c + dx) \left(\frac{a + b \cos(c + dx)}{a + b}\right)^{2/3} \int \frac{1}{\sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1} \left(\frac{a}{a + b} + \frac{b \cos(c + dx)}{a + b}\right)^{2/3}} d \cos(c + dx)}{d \sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1} (a + b \cos(c + dx))^{2/3}} \\
 & \quad \downarrow \text{155} \\
 & \frac{\sqrt{2} \sin(c + dx) \left(\frac{a + b \cos(c + dx)}{a + b}\right)^{2/3} \text{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, \frac{2}{3}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx)), \frac{b(1 - \cos(c + dx))}{a + b}\right)}{d \sqrt{\cos(c + dx) + 1} (a + b \cos(c + dx))^{2/3}}
 \end{aligned}$$

input `Int[(a + b*Cos[c + d*x])^(-2/3),x]`

output `(Sqrt[2]*AppellF1[1/2, 1/2, 2/3, 3/2, (1 - Cos[c + d*x])/2, (b*(1 - Cos[c + d*x]))/(a + b)]*((a + b*Cos[c + d*x])/(a + b))^(2/3)*Sin[c + d*x])/(d*Sqrt[1 + Cos[c + d*x]]*(a + b*Cos[c + d*x])^(2/3))`

## 3.60.3.1 Defintions of rubi rules used

```
rule 155 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))
^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simp
lify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplerQ[e + f*x, a + b*x])
```

```
rule 156 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))
^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p
]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Si
mp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c,
d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] &
& GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3144 Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x
)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]
```

## 3.60.4 Maple [F]

$$\int \frac{1}{(a + \cos(dx + c)b)^{\frac{2}{3}}} dx$$

```
input int(1/(a+cos(d*x+c)*b)^(2/3),x)
```

```
output int(1/(a+cos(d*x+c)*b)^(2/3),x)
```

**3.60.5 Fricas [F]**

$$\int \frac{1}{(a + b \cos(c + dx))^{2/3}} dx = \int \frac{1}{(b \cos(dx + c) + a)^{2/3}} dx$$

input `integrate(1/(a+b*cos(d*x+c))^(2/3),x, algorithm="fricas")`

output `integral((b*cos(d*x + c) + a)^(-2/3), x)`

**3.60.6 Sympy [F]**

$$\int \frac{1}{(a + b \cos(c + dx))^{2/3}} dx = \int \frac{1}{(a + b \cos(c + dx))^{2/3}} dx$$

input `integrate(1/(a+b*cos(d*x+c))**(2/3),x)`

output `Integral((a + b*cos(c + d*x))**(-2/3), x)`

**3.60.7 Maxima [F]**

$$\int \frac{1}{(a + b \cos(c + dx))^{2/3}} dx = \int \frac{1}{(b \cos(dx + c) + a)^{2/3}} dx$$

input `integrate(1/(a+b*cos(d*x+c))^(2/3),x, algorithm="maxima")`

output `integrate((b*cos(d*x + c) + a)^(-2/3), x)`

**3.60.8 Giac [F]**

$$\int \frac{1}{(a + b \cos(c + dx))^{2/3}} dx = \int \frac{1}{(b \cos(dx + c) + a)^{2/3}} dx$$

input `integrate(1/(a+b*cos(d*x+c))^(2/3),x, algorithm="giac")`

output `integrate((b*cos(d*x + c) + a)^(-2/3), x)`

**3.60.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a + b \cos(c + dx))^{2/3}} dx = \int \frac{1}{(a + b \cos(c + dx))^{2/3}} dx$$

input `int(1/(a + b*cos(c + d*x))^(2/3),x)`

output `int(1/(a + b*cos(c + d*x))^(2/3), x)`



### 3.61 $\int \frac{1}{(a+b \cos(c+dx))^{4/3}} dx$

3.61.1	Optimal result	400
3.61.2	Mathematica [B] (verified)	400
3.61.3	Rubi [A] (verified)	401
3.61.4	Maple [F]	403
3.61.5	Fricas [F]	403
3.61.6	Sympy [F]	403
3.61.7	Maxima [F]	404
3.61.8	Giac [F]	404
3.61.9	Mupad [F(-1)]	404

#### 3.61.1 Optimal result

Integrand size = 14, antiderivative size = 110

$$\int \frac{1}{(a + b \cos(c + dx))^{4/3}} dx = \frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, \frac{4}{3}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx)), \frac{b(1 - \cos(c + dx))}{a + b}\right) \sqrt[3]{\frac{a + b \cos(c + dx)}{a + b}}}{(a + b)d\sqrt{1 + \cos(c + dx)}\sqrt[3]{a + b \cos(c + dx)}}$$

output `AppellF1(1/2,4/3,1/2,3/2,b*(1-cos(d*x+c))/(a+b),1/2-1/2*cos(d*x+c))*((a+b*cos(d*x+c))/(a+b))^(1/3)*sin(d*x+c)*2^(1/2)/(a+b)/d/(a+b*cos(d*x+c))^(1/3)/(1+cos(d*x+c))^(1/2)`

#### 3.61.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 268 vs. 2(110) = 220.

Time = 2.27 (sec) , antiderivative size = 268, normalized size of antiderivative = 2.44

$$\int \frac{1}{(a + b \cos(c + dx))^{4/3}} dx = \frac{15a \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{2}, \frac{1}{2}, \frac{5}{3}, \frac{a+b \cos(c+dx)}{a-b}, \frac{a+b \cos(c+dx)}{a+b}\right) \sqrt{-\frac{b(-1+\cos(c+dx))}{a+b}} \sqrt{-\frac{b(1+\cos(c+dx))}{a+b}}}{(a + b \cos(c + dx))^{4/3}}$$

input `Integrate[(a + b*cos[c + d*x])^(-4/3), x]`

output  $(15*a*AppellF1[2/3, 1/2, 1/2, 5/3, (a + b*\text{Cos}[c + d*x])/(a - b), (a + b*\text{Cos}[c + d*x])/(a + b)]*\text{Sqrt}[-((b*(-1 + \text{Cos}[c + d*x]))/(a + b))]*\text{Sqrt}[-((b*(1 + \text{Cos}[c + d*x]))/(a - b))]*(a + b*\text{Cos}[c + d*x])*Csc[c + d*x] - 6*(5*b^2 + 2*AppellF1[5/3, 1/2, 1/2, 8/3, (a + b*\text{Cos}[c + d*x])/(a - b), (a + b*\text{Cos}[c + d*x])/(a + b)]*\text{Sqrt}[-((b*(-1 + \text{Cos}[c + d*x]))/(a + b))]*\text{Sqrt}[(b*(1 + \text{Cos}[c + d*x]))/(-a + b)]*(a + b*\text{Cos}[c + d*x])^2*Csc[c + d*x]^2*\text{Sin}[c + d*x])/(10*b*(a^2 - b^2)*d*(a + b*\text{Cos}[c + d*x])^(1/3))$

### 3.61.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + b \cos(c + dx))^{4/3}} dx$$

↓ 3042

$$\int \frac{1}{(a + b \sin(c + dx + \frac{\pi}{2}))^{4/3}} dx$$

↓ 3144

$$-\frac{\sin(c + dx) \int \frac{1}{\sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1} (a + b \cos(c + dx))^{4/3}} d \cos(c + dx)}{d \sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}}$$

↓ 156

$$-\frac{\sin(c + dx) \sqrt[3]{\frac{a + b \cos(c + dx)}{a + b}} \int \frac{1}{\sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1} \left(\frac{a}{a + b} + \frac{b \cos(c + dx)}{a + b}\right)^{4/3}} d \cos(c + dx)}{d(a + b) \sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1} \sqrt[3]{a + b \cos(c + dx)}}$$

↓ 155

$$\frac{\sqrt{2} \sin(c + dx) \sqrt[3]{\frac{a + b \cos(c + dx)}{a + b}} \text{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, \frac{4}{3}, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx)), \frac{b(1 - \cos(c + dx))}{a + b}\right)}{d(a + b) \sqrt{\cos(c + dx) + 1} \sqrt[3]{a + b \cos(c + dx)}}$$

input  $\text{Int}[(a + b*\text{Cos}[c + d*x])^(-4/3), x]$

---

3.61.  $\int \frac{1}{(a + b \cos(c + dx))^{4/3}} dx$

output  $(\text{Sqrt}[2]*\text{AppellF1}[1/2, 1/2, 4/3, 3/2, (1 - \text{Cos}[c + d*x])/2, (b*(1 - \text{Cos}[c + d*x]))/(a + b)]*((a + b*\text{Cos}[c + d*x])/(a + b))^{1/3}*\text{Sin}[c + d*x])/((a + b)*d*\text{Sqrt}[1 + \text{Cos}[c + d*x]]*(a + b*\text{Cos}[c + d*x])^{1/3})$

### 3.61.3.1 Defintions of rubi rules used

rule 155  $\text{Int}[(a + b*x)^m * (c + d*x)^n * (e + f*x)^p, x] \rightarrow \text{Simp}[(a + b*x)^{m+1} / (b*(m+1)*\text{Simplify}[b/(b*c - a*d)]^{n+1} * \text{Simplify}[b/(b*e - a*f)]^p) * \text{AppellF1}[m+1, -n, -p, m+2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /;$   $\text{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x$  &&  $\text{IntegerQ}[m]$  &&  $\text{IntegerQ}[n]$  &&  $\text{IntegerQ}[p]$  &&  $\text{GtQ}[\text{Simplify}[b/(b*c - a*d)], 0]$  &&  $\text{GtQ}[\text{Simplify}[b/(b*e - a*f)], 0]$  &&  $\text{!(GtQ}[\text{Simplify}[d/(d*a - c*b)], 0] \&\& \text{GtQ}[\text{Simplify}[d/(d*e - c*f)], 0] \&\& \text{SimplerQ}[c + d*x, a + b*x]) \&\& \text{!(GtQ}[\text{Simplify}[f/(f*a - e*b)], 0] \&\& \text{GtQ}[\text{Simplify}[f/(f*c - e*d)], 0] \&\& \text{SimplerQ}[e + f*x, a + b*x])$

rule 156  $\text{Int}[(a + b*x)^m * (c + d*x)^n * (e + f*x)^p, x] \rightarrow \text{Simp}[(e + f*x)^{\text{FracPart}[p]} / (\text{Simplify}[b/(b*e - a*f)]^{\text{IntPart}[p]} * (b*((e + f*x)/(b*e - a*f)))^{\text{FracPart}[p]}) * \text{Int}[(a + b*x)^m * (c + d*x)^n * \text{Simp}[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /;$   $\text{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x$  &&  $\text{IntegerQ}[m]$  &&  $\text{IntegerQ}[n]$  &&  $\text{IntegerQ}[p]$  &&  $\text{GtQ}[\text{Simplify}[b/(b*c - a*d)], 0]$  &&  $\text{!(GtQ}[\text{Simplify}[b/(b*e - a*f)], 0]$

rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$   $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3144  $\text{Int}[(a + b*\sin[(c + d*x)])^n, x\_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x] / (d*\text{Sqrt}[1 + \text{Sin}[c + d*x]]*\text{Sqrt}[1 - \text{Sin}[c + d*x]]) * \text{Subst}[\text{Int}[(a + b*x)^n / (\text{Sqrt}[1 + x]*\text{Sqrt}[1 - x]), x], x, \text{Sin}[c + d*x]], x] /;$   $\text{FreeQ}\{a, b, c, d, n\}, x$  &&  $\text{NeQ}[a^2 - b^2, 0]$  &&  $\text{IntegerQ}[2*n]$

**3.61.4 Maple [F]**

$$\int \frac{1}{(a + \cos(dx + c)b)^{\frac{4}{3}}} dx$$

input `int(1/(a+cos(d*x+c)*b)^(4/3),x)`

output `int(1/(a+cos(d*x+c)*b)^(4/3),x)`

**3.61.5 Fricas [F]**

$$\int \frac{1}{(a + b \cos(c + dx))^{4/3}} dx = \int \frac{1}{(b \cos(dx + c) + a)^{\frac{4}{3}}} dx$$

input `integrate(1/(a+b*cos(d*x+c))^(4/3),x, algorithm="fricas")`

output `integral((b*cos(d*x + c) + a)^(2/3)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2), x)`

**3.61.6 Sympy [F]**

$$\int \frac{1}{(a + b \cos(c + dx))^{4/3}} dx = \int \frac{1}{(a + b \cos(c + dx))^{\frac{4}{3}}} dx$$

input `integrate(1/(a+b*cos(d*x+c))**(4/3),x)`

output `Integral((a + b*cos(c + d*x))**(-4/3), x)`

**3.61.7 Maxima [F]**

$$\int \frac{1}{(a + b \cos(c + dx))^{4/3}} dx = \int \frac{1}{(b \cos(dx + c) + a)^{4/3}} dx$$

input `integrate(1/(a+b*cos(d*x+c))^(4/3),x, algorithm="maxima")`

output `integrate((b*cos(d*x + c) + a)^(-4/3), x)`

**3.61.8 Giac [F]**

$$\int \frac{1}{(a + b \cos(c + dx))^{4/3}} dx = \int \frac{1}{(b \cos(dx + c) + a)^{4/3}} dx$$

input `integrate(1/(a+b*cos(d*x+c))^(4/3),x, algorithm="giac")`

output `integrate((b*cos(d*x + c) + a)^(-4/3), x)`

**3.61.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a + b \cos(c + dx))^{4/3}} dx = \int \frac{1}{(a + b \cos(c + dx))^{4/3}} dx$$

input `int(1/(a + b*cos(c + d*x))^(4/3),x)`

output `int(1/(a + b*cos(c + d*x))^(4/3), x)`

### 3.62 $\int (a + b \cos(c + dx))^n dx$

3.62.1	Optimal result . . . . .	405
3.62.2	Mathematica [A] (verified) . . . . .	405
3.62.3	Rubi [A] (verified) . . . . .	406
3.62.4	Maple [F] . . . . .	407
3.62.5	Fricas [F] . . . . .	408
3.62.6	Sympy [F] . . . . .	408
3.62.7	Maxima [F] . . . . .	408
3.62.8	Giac [F] . . . . .	409
3.62.9	Mupad [F(-1)] . . . . .	409

#### 3.62.1 Optimal result

Integrand size = 12, antiderivative size = 103

$$\int (a + b \cos(c + dx))^n dx = \frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -n, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx)), \frac{b(1 - \cos(c + dx))}{a + b}\right) (a + b \cos(c + dx))^n \left(\frac{a + b \cos(c + dx)}{a + b}\right)^{-n} \sin(c + dx)}{d \sqrt{1 + \cos(c + dx)}}$$

```
output AppellF1(1/2, -n, 1/2, 3/2, b*(1-cos(d*x+c))/(a+b), 1/2-1/2*cos(d*x+c))*(a+b*cos(d*x+c))^n*sin(d*x+c)*2^(1/2)/d/(((a+b*cos(d*x+c))/(a+b))^n/(1+cos(d*x+c)))^(1/2)
```

#### 3.62.2 Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.17

$$\int (a + b \cos(c + dx))^n dx = \frac{\operatorname{AppellF1}\left(1 + n, \frac{1}{2}, \frac{1}{2}, 2 + n, \frac{a + b \cos(c + dx)}{a - b}, \frac{a + b \cos(c + dx)}{a + b}\right) \sqrt{-\frac{b(-1 + \cos(c + dx))}{a + b}} \sqrt{\frac{b(1 + \cos(c + dx))}{-a + b}} (a + b \cos(c + dx))^n}{bd(1 + n)}$$

```
input Integrate[(a + b*Cos[c + d*x])^n, x]
```

output `-((AppellF1[1 + n, 1/2, 1/2, 2 + n, (a + b*Cos[c + d*x])/(a - b), (a + b*Cos[c + d*x])/(a + b)]*Sqrt[-((b*(-1 + Cos[c + d*x]))/(a + b))] * Sqrt[(b*(1 + Cos[c + d*x]))/(-a + b)]*(a + b*Cos[c + d*x])^(1 + n)*Csc[c + d*x])/(b*d*(1 + n))`

### 3.62.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + b \cos(c + dx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( a + b \sin \left( c + dx + \frac{\pi}{2} \right) \right)^n dx \\
 & \quad \downarrow \text{3144} \\
 & \frac{\sin(c + dx) \int \frac{(a + b \cos(c + dx))^n}{\sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}} d \cos(c + dx)}{d \sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}} \\
 & \quad \downarrow \text{156} \\
 & \frac{\sin(c + dx) (a + b \cos(c + dx))^n \left( \frac{a + b \cos(c + dx)}{a + b} \right)^{-n} \int \frac{\left( \frac{a}{a + b} + \frac{b \cos(c + dx)}{a + b} \right)^n}{\sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}} d \cos(c + dx)}{d \sqrt{1 - \cos(c + dx)} \sqrt{\cos(c + dx) + 1}} \\
 & \quad \downarrow \text{155} \\
 & \frac{\sqrt{2} \sin(c + dx) (a + b \cos(c + dx))^n \left( \frac{a + b \cos(c + dx)}{a + b} \right)^{-n} \text{AppellF1} \left( \frac{1}{2}, \frac{1}{2}, -n, \frac{3}{2}, \frac{1}{2} (1 - \cos(c + dx)), \frac{b(1 - \cos(c + dx))}{a + b} \right)}{d \sqrt{\cos(c + dx) + 1}}
 \end{aligned}$$

input `Int[(a + b*Cos[c + d*x])^n,x]`

output `(Sqrt[2]*AppellF1[1/2, 1/2, -n, 3/2, (1 - Cos[c + d*x])/2, (b*(1 - Cos[c + d*x]))/(a + b)]*(a + b*Cos[c + d*x])^n*Sin[c + d*x])/(d*Sqrt[1 + Cos[c + d*x]])*((a + b*Cos[c + d*x])/(a + b))^n`

## 3.62.3.1 Defintions of rubi rules used

```
rule 155 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))
^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simp
lify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplerQ[e + f*x, a + b*x])
```

```
rule 156 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))
^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p
]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Si
mp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c,
d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] &
& GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3144 Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x
)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]
```

## 3.62.4 Maple [F]

$$\int (a + \cos(dx + c)b)^n dx$$

```
input int((a+cos(d*x+c)*b)^n,x)
```

```
output int((a+cos(d*x+c)*b)^n,x)
```



**3.62.5 Fricas [F]**

$$\int (a + b \cos(c + dx))^n dx = \int (b \cos(dx + c) + a)^n dx$$

input `integrate((a+b*cos(d*x+c))^n,x, algorithm="fricas")`

output `integral((b*cos(d*x + c) + a)^n, x)`

**3.62.6 Sympy [F]**

$$\int (a + b \cos(c + dx))^n dx = \int (a + b \cos(c + dx))^n dx$$

input `integrate((a+b*cos(d*x+c))**n,x)`

output `Integral((a + b*cos(c + d*x))**n, x)`

**3.62.7 Maxima [F]**

$$\int (a + b \cos(c + dx))^n dx = \int (b \cos(dx + c) + a)^n dx$$

input `integrate((a+b*cos(d*x+c))^n,x, algorithm="maxima")`

output `integrate((b*cos(d*x + c) + a)^n, x)`

**3.62.8 Giac [F]**

$$\int (a + b \cos(c + dx))^n dx = \int (b \cos(dx + c) + a)^n dx$$

input `integrate((a+b*cos(d*x+c))^n,x, algorithm="giac")`

output `integrate((b*cos(d*x + c) + a)^n, x)`

**3.62.9 Mupad [F(-1)]**

Timed out.

$$\int (a + b \cos(c + dx))^n dx = \int (a + b \cos(c + dx))^n dx$$

input `int((a + b*cos(c + d*x))^n,x)`

output `int((a + b*cos(c + d*x))^n, x)`

## APPENDIX

4.1 Listing of Grading functions . . . . .	410
--	-----

## 4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

### 4.1.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
```

```

(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A"," "}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A"," "}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal.$
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
      ,
      finalresult={"F","Contains unresolved integral."}
    ]
  ];

  finalresult
]

```

```

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType,expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]],2]],
            Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3,ExpnType[expn[[1]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
            If[Head[expn]===RootSum,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

### 4.1.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
      return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
            print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
      return "F","Result contains unresolved integral";
fi;

```

```

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
  if is_contains_complex(result) then
    if is_contains_complex(optimal) then
      if debug then
        print("both result and optimal complex");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        return "A"," ";
      else
        return "B",cat("Both result and optimal contain complex but leaf count of
                        convert(leaf_count_result,string)," vs. $2 (" ,
                        convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_
        end if
      else #result contains complex but optimal is not
        if debug then
          print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
      fi;
    else # result do not contain complex
      # this assumes optimal do not as well. No check is needed here.
      if debug then
        print("result do not contain complex, this assumes optimal do not as well");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        if debug then
          print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
      else
        if debug then
          print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string)," $ vs. $2(" ,
                        convert(leaf_count_optimal,string)," )=" ,convert(2*leaf_cou
        fi;
      fi;
    fi;
  fi;

```



```

else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  end if
end proc:

```

```

elif type(expn, ``~`) then
  if type(op(2,expn), 'integer') then
    ExpnType(op(1,expn))
  elif type(op(2,expn), 'rational') then
    if type(op(1,expn), 'rational') then
      1
    else
      max(2, ExpnType(op(1,expn)))
    end if
  else
    max(3, ExpnType(op(1,expn)), ExpnType(op(2,expn)))
  end if
elif type(expn, ``+`) or type(expn, ``*`) then
  max(ExpnType(op(1,expn)), max(ExpnType(rest(expn))))
elif ElementaryFunctionQ(op(0,expn)) then
  max(3, ExpnType(op(1,expn)))
elif SpecialFunctionQ(op(0,expn)) then
  max(4, apply(max, map(ExpnType, [op(expn)])))
elif HypergeometricFunctionQ(op(0,expn)) then
  max(5, apply(max, map(ExpnType, [op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6, apply(max, map(ExpnType, [op(expn)])))
elif op(0,expn)='int' then
  max(8, apply(max, map(ExpnType, [op(expn)]))) else
  9
end if
end proc:

```

```

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

```

```

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,

```

```

        GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
        EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
    member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
    member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
    if nops(u)=2 then
        op(2,u)
    else
        apply(op(0,u),op(2..nops(u),u))
    end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma] [LeafCount] (u);
end proc:

```

### 4.1.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):

```

```

if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
        return True
    else:
        return False
else:
    return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):

```

```

return 1
elif isinstance(expn,list):
    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
        return 1
    else:
        return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
elif isinstance(expn,Pow): #type(expn,``^`)
    if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
        return expnType(expn.args[0]) #ExpnType(op(1,expn))
    elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or type(expn,``*`)
    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

*#main function*

```

def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

    #print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```

#### 4.1.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
```

```

        return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

```



```

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric', 'hypergeometric_M', 'hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=", expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception, AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list:  #isinstance(expn,list):

```

```

    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
elif expn.operator() == operator.pow: #instance(expn,Pow)
    if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)
        return expnType(expn.operands()[0]) #expnType(expn.args[0])
    elif type(expn.operands()[1])==Rational: #instance(expn.args[1],Rational)
        if type(expn.operands()[0])==Rational: #instance(expn.args[0],Rational)
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #instance(expn,Add) or inst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

```

```

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)",type(result))
    print("type(optimal)",type(optimal))

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal." + str(leaf
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```